*The University of Michigan
Institute of Science and Technology Series*

MBL/WHOI

0 0301 0012671 0

THE NATURE OF BIOLOGICAL DIVERSITY

The University of Michigan
Institute of Science and Technology Series

ALLEN • The Molecular Control of Cellular Activity

ALLEN • The Nature of Biological Diversity

LILLER • Space Astrophysics

The Nature of

BIOLOGICAL DIVERSITY

Edited by

John M. Allen

*Associate Professor of Zoology
The University of Michigan*

McGraw-Hill Book Company, Inc.

NEW YORK SAN FRANCISCO TORONTO LONDON

THE NATURE OF BIOLOGICAL DIVERSITY

Copyright © 1963 by the McGraw-Hill Book Company, Inc. All Rights Reserved. Printed in the United States of America. This book, or parts thereof, may not be reproduced in any form without permission of the publishers. *Library of Congress Catalog Card Number* 62-20182

01078

Preface

The papers contained within this volume are the result of a series of lectures held at The University of Michigan in the spring of 1961, under the auspices of its Institute of Science and Technology.

Of the several areas of inquiry that are germane to an understanding of biological phenomena, one of the more challenging is that which attempts to define the factors responsible for the diversification of cellular structure and function. In planning the lectures from which this volume is derived, we felt that examination of the elements leading to the diversification of biological systems from a broad vantage point might yield a degree of synthesis to an area which is fraught with as much variation in its approaches as the systems which it attempts to investigate. This volume deals nearly exclusively with the development of heterogeneity at the cellular and subcellular level, for elucidation of the problems involved will come largely from these areas of investigation which are basically biochemical and molecular in their philosophy.

In pondering the general problem of diversification in biological systems, we arrive at the realization not only that the properties of cells are the result of a long history of organic evolution but also that prebiological evolution has placed basic restrictions upon what cells may become and what they may do. These restrictions, of course, relate to the properties of the molecules available for biological use. Thus, in any general discussion of diversification, an examination of the origin of organic molecules makes a logical starting point. This is the topic which has been chosen to introduce the subject. From this point an examination of ways in which molecules may interact to yield biologically useful energy needed for cell maintenance may be considered. Perhaps the most informative system in this respect is the evolution of the photosynthetic mechanism.

Cell diversification depends upon the variety of catalytic and structural proteins from which the cells are fabricated. These aspects of the problem are examined in three chapters dealing with biochemical

evolution, the origin of specific proteins, and the nature of catalytic proteins. The macromolecular aspects of these processes are examined in chapters dealing with cellular organization and the genetic control of cellular organization, with protozoa as a model system. The final chapters of this book attempt to illustrate the ways in which cellular interactions and cellular products modify the processes of molecular diversification.

Throughout these pages we have tried to dissect the fabric of molecular and cellular interaction which lead to the bewildering array of biological complexity confronting us. We hope this collection of papers will lend perspective to this area. In some small measure we feel we have accomplished this.

I wish to thank the contributors to this volume who gave so freely and graciously of their thoughts and time. Thanks are also due Drs. Wilbur Ackermann and M. J. Coon of The University of Michigan, who spent much time in planning the lecture series from which this book is derived.

John M. Allen

Contents

PREFACE	v
CHAPTER	
1 THE ORIGIN OF ORGANIC MOLECULES <i>Harold C. Urey</i>	1
2 EVOLUTION OF PHOTOSYNTHETIC MECHANISMS <i>Melvin Calvin</i>	15
3 BIOCHEMISTRY AND EVOLUTION <i>Ernest Baldwin</i>	45
4 THE NATURE AND DIVERSITY OF CATALYTIC PRO- TEINS <i>Paul D. Boyer</i>	69
5 THE ORIGIN OF SPECIFIC PROTEINS <i>Clement L. Markert</i>	95
6 DIVERSITY AT THE SUBCELLULAR LEVEL AND ITS SIGNIFICANCE <i>Keith R. Porter</i>	121
7 DOES PREFORMED CELL STRUCTURE PLAY AN ESSENTIAL ROLE IN CELL HEREDITY? <i>Tracy M. Sonneborn</i>	165
8 MICROENVIRONMENTAL INFLUENCES IN CYTO- DIFFERENTIATION <i>Clifford Grobstein</i>	223
9 DIFFERENTIATION AND MORPHOGENESIS IN IN- SECTS <i>Carroll M. Williams</i>	243
10 GROWTH AND DIFFERENTIATION IN THE NERV- OUS SYSTEM <i>Rita Levi-Montalcini</i>	261
INDEX	297

THE NATURE OF BIOLOGICAL DIVERSITY

The Origin of Organic Molecules

Harold C. Urey

School of Science and Engineering
University of California, San Diego
La Jolla, California

Darwin and Wallace proposed the theory of evolution almost simultaneously and quite independently. They and their numerous followers of the past century have demonstrated that, given an organism of the most primitive type, all the species of today and of past geological periods could and did evolve from such cells. We are beginning to understand the processes by which inherited characteristics are carried from generation to generation by means of deoxyribonucleic acids, and also we understand how variations in species of animals and plants occur. Many details remain, but the broad outline has become clearer during recent years and decades. Also, we find marked biochemical similarities between all living things, indicating that they are all part of a single evolutionary process. These fundamental biochemical processes were established in the distant past before the time of the oldest fossils, namely, some 500 million or more years ago.

At least after the development of sexual reproduction, evolution must have occurred by mutation in one individual, it is true, but the

2 The Nature of Biological Diversity

characteristic must have been segregated from a community. It seems likely that precellular evolution also involved the segregation of chemical constituents from many sources.

The complexity of living organisms astounds all students of biology, and the more knowledge the observer has the more astounding do the facts appear to be. Is it not true that the development of a particular minute drop of clear protoplasm, such as a fertilized egg, into a sea urchin and nothing else is one of the greatest wonders of which we have knowledge? Yet the study of the compounds of carbon, nitrogen, oxygen, and hydrogen shows us that a very great variety of these compounds is possible. Two illustrations of the varieties of the molecules can be given. The human egg contains strands of total length of about 1 meter long of deoxyribonucleic acid containing about 10^9 units, each of which can be chosen in four different ways. The number of possible molecules becomes 4^{10^9} , which is an immense number. To specify a human being at each moment of its existence requires less specific information; much redundancy is possible and surely exists. Again the number of possible proteins consisting of, say, 170 amino acids of 20 kinds is very great. If one of each kind were placed in a cubical box, the length of the edge of the box would be 10 raised to nearly the 50th power light-years. Not all the possible protein molecules exist now in the observed universe nor could they have existed during the last 4.5 billion years.

Natural phenomena are very complex in all their various manifestations. We can follow in considerable detail the more simple and isolated examples of these phenomena, for example, many chemical reactions and structures, the structure of atoms and the nuclei of atoms, and the gross structure of stars and galaxies of stars. The details of natural phenomena exceed in complexity the complexity of structure of the human brain, and we cannot follow these except in an approximate way. But we can say something about the comparative complexities of the chemistry of the carbon, nitrogen, oxygen, hydrogen compounds and those of other elements. I shall state my conclusion categorically in this way. There is no known chemistry of other elements which approaches in ordered complexity that of these four elements, and we know enough of the chemistry of all the elements that we can be certain that no other such chemistry exists. Only these elements supply such a complexity of compounds and chemical reactions that we are forced to conclude that only the chemistry of these elements could supply the complexity of structure and behavior which we recognize as those of living things. A system of life based on other

elements would necessarily be so vastly simpler in behavior that we would not recognize it as such.

The liquid in which all terrestrial life exists is water. It is interesting to inquire whether other liquids might be suitable. Only two seem possible candidates, namely, ammonia and hydrocarbons. It is difficult to imagine circumstances that would supply liquid ammonia to a planet without water being present, but its chemical properties might not rule it out. The hydrocarbons might be present on a planet but the generally slow reactions occurring in hydrocarbon solvents look less likely as substitutes for moderately lively reactions of living things.

Living organisms do grow in petroleum products, though these organisms are of the hydrophilic variety and depend for their energy on the oxidation of petroleum by atmospheric oxygen. Free oxygen should be nonexistent on a planet with excess hydrocarbon rather than an excess of water. The sulfur deposits of the world have been produced by the reduction of CaSO_4 by petroleum through the action of bacteria, but again sulfate has been produced on earth by oxidation due to free atmospheric oxygen. These arguments may not exclude the possibility of life on a planet having excess hydrocarbon compounds but no oxidizing atmosphere, and any conclusion to this effect may only indicate a prejudice arising from our terrestrial experience. It seems likely to the writer that only a water solvent could lead to any reasonable approximation to compounds and chemical reactions of sufficient complexity to be regarded as constituting a living organism.

Primitive conditions on the earth

Hydrogen is the most abundant element in the cosmos. In particular the sun contains 80 per cent H_2 , and it can be confidently supposed that the earth and other planets evolved from matter in which hydrogen was equally abundant initially. Also, carbon is reported to be about equally as abundant as oxygen. The abundances of some of the relevant elements in solar and terrestrial material are given in Table 1.

It is evident that solar material is highly reducing, with about 1,100 times as many atoms of hydrogen as of oxygen, and that the relative abundances of these elements are quite different in terrestrial matter at the present time. Meteoritic matter is in general very reducing also, i.e., it contains iron in the elementary and ferrous states. (However, the carbonaceous chondrites contain carbon compounds which are partially oxidized and also highly oxidized sulfur as magnesium sul-

4 The Nature of Biological Diversity

fate.*) The abundances of some elements in the terrestrial surface regions are given also in Table 1. It will be noted that hydrogen is present mostly in oxidized form as water and that carbon and nitrogen are far less abundant than oxygen in surface water and even less abundant relative to oxygen in igneous rocks. Terrestrial surface carbon is present mostly as carbonates, i.e., in the highest oxidation state. If one added a small amount of hydrogen to the earth, one would expect that carbon and nitrogen would be converted in time to methane and ammonia and all sulfates would be reduced to sulfides.

Table 1. Atomic abundances of some elements relative to oxygen

Element	Sun	Earth's atmosphere and oceans	Earth's igneous rocks
H	1.100	2	4.4×10^{-2}
C	0.57	2.1×10^{-2}	9.1×10^{-4}
N	0.105	3.4×10^{-3}	1.1×10^{-4}
O	1	1	1
Mg	2.75×10^{-2}	9.4×10^{-4}	3×10^{-2}
P	2.4×10^{-4}	5.8×10^{-8}	1.3×10^{-3}
S	2.2×10^{-2}	5.0×10^{-4}	5.6×10^{-4}
K	5.5×10^{-5}	1.75×10^{-4}	2.3×10^{-2}
Ca	1.55×10^{-3}	1.8×10^{-4}	3.1×10^{-2}

* The carbon abundance is that of the sedimentary rocks relative to oxygen in the oceans. The abundances of the mineral elements in sedimentary rocks are similar to those in igneous rocks.

SOURCES: Sun: L. Goldberg, E. A. Müller, and L. H. Aller, *Astrophys. J., Suppl. Ser.* 5:1-138 (1960). Earth: K. Rankama and T. G. Sahama, *Geochemistry*, University of Chicago Press, Chicago (1950).

It is evident that two important processes from the standpoint of the present discussion have occurred on the earth and also on Mars and Venus. (1) The more volatile fraction of solar matter has been lost to a high degree, including the elements H, N, C, and O, and the inert gases. (2) The surface regions of these planets have become highly oxidized. The first process is veiled in the obscurity of the earth's origin. Since any discussion of the process would carry considerable doubt and would be very long and involved, it will not be treated here. The second process is well understood. Oxidation of the earth is probably continuing at the present time and is due to the escape of hydrogen from the earth into space. The same process can reasonably be supposed to have occurred in the case of Mars and

* E. Du Fresne, Dissertation, University of Chicago, 1960.

Venus. It is most improbable that hydrogen is now escaping from the major planets. The general course of oxidation of carbon and nitrogen expected is shown in Table 2. The compounds listed are only the most simple examples. Many complex CNOH compounds exist in which the oxidation states of C and N lie between the lowest and highest oxidation states. (The oxidation state is usually defined by counting oxygen as -2 and hydrogen as $+1$ and making the carbon and nitrogen states such that the molecule is neutral. As is well known, the oxidation states are not always uniquely determined.) The general course of

Table 2. Oxidation states

4-	2-	0	2+	4+
CH ₄	CH ₃ OH	CH ₂ O	CHOOH	CO ₂
3-	1-	0	Higher oxidation states	
NH ₃	NH ₂ OH	N ₂		

chemical change on the earth and other terrestrial planets has been from the more reduced to the more oxidized condition, i.e., from left to right in the table. It is evident that a favorable condition for the existence of compounds such as those required for living organisms would be present during this oxidation process.

The escape of hydrogen from the earth and inorganic reactions involved

It is possible to estimate the conditions for the escape of hydrogen from the earth rather plausibly, though hardly with certainty. The escape conditions have been given in considerable detail by Urey (1959). Jeans derived the formula for the escape of planetary atmospheres. Its application to the earth indicated a loss of hydrogen from the earth considerably larger than could be accounted for by the oxidation of surface elements by the oxygen left behind in the process. Harteck and Jensen showed that diffusion was important and suggested a diffusion layer at the tropopause. Subsequent work showed that circulation occurred through this layer and Urey suggested that the diffusion layer was at the top of the atmosphere.* The rate of escape is determined by diffusion in this layer and the rate is proportional to the surface hydrogen pressure, which at present is 10^{-6} atmosphere. The present rate was calculated as 10^7 atoms $\text{sec}^{-1} \text{cm}^{-2}$. This is not a highly precise value. It is probable that the rate of escape

* See Bates and McDowell (1957) for a discussion of the escape problem for He³ and He⁴.

differs with time of day and the latitude. If this rate continued throughout 4.5×10^9 years, the loss would amount to the hydrogen from only 20 g cm^{-2} of water. In order to account for the oxidation of carbon, ammonia, sulfide, and some ferrous oxide, Miller and Urey estimated that probably 1,500 times as much hydrogen must have escaped and hence that the hydrogen pressure must have been some 1.5×10^{-3} atmospheres during much of geologic time and the total water decomposed about $3 \times 10^4 \text{ g cm}^{-2}$ or about one-tenth of the present surface water.* It is reasonable to suppose that hydrogen would be more abundant in an atmosphere that did not contain free oxygen and that it was the appearance of this oxygen which decreased the hydrogen abundance and initiated the present rate of loss.

If hydrogen were present at these comparatively high pressures, it is necessary to ask whether it would interfere with other chemical facts. Carbon dioxide reacts with silicates to give limestone and silicon dioxide, and at the same time hydrogen reacts with carbon dioxide to form methane and water. If the carbon dioxide is destroyed by the second reaction until its pressure becomes lower than that required to precipitate calcium carbonate, the first reaction cannot proceed. Since very ancient limestones are known, this would be impossible. Miller and Urey (1959) showed that these conditions can be met satisfactorily. They also found that nitrogen would be present as NH_4^+ ion if the hydrogen pressure were the assumed value. This study showed that the necessary escape of hydrogen was possible, that calcium carbonate could be formed, and that reduced nitrogen could be present in the oceans. Such a study does not prove that the conditions did exist but only that they are not inconsistent with observations.

The production of carbon compounds

As a first approximation, it is reasonable to assume that the chemical elements will be present in nature in chemical compounds which are thermodynamically stable under the existing conditions. It is well known also that deviations from this condition are so very common as to be almost universal, though such deviations are usually not large. But we may well review briefly some general facts about the stability of carbon-hydrogen-oxygen compounds. The simpler compounds of

* There are indications that deuterium is more abundant in the ocean than in juvenile water by some 4 per cent. This indicates that hydrogen has been lost to a larger extent proportionally than has deuterium. The observations are not inconsistent with the assumptions of Miller and Urey. See N. Kokubu, T. Mayeda, and H. C. Urey (1961), *Geochim. et Cosmochim. Acta*, **21**:247.

this kind are all unstable with respect to the completely oxidized and reduced condition. Thus at 25°C the following reactions all proceed toward the right-hand side.

At least for many other carbon compounds, similar statements are true. Compounds of carbon in intermediate oxidation states such as those common in living organisms are unstable even in the absence of hydrogen or oxygen. It is also true that even small pressures of hydrogen will lead to reactions to produce the completely reduced state of carbon, and that no carbon dioxide would be produced. Also oxygen will burn up all these compounds at ordinary temperatures, given sufficient time with or without suitable catalysts. What we note is that the compounds of which living organisms are composed are unstable. This is necessarily true, for otherwise no metabolism in either fermentative or oxidative organisms would occur. But this emphasizes that during the evolution of the most primitive living organism, as well as during the lives of well-organized living things, a continuous and effective source of energy must be present.

We can estimate the sources of energy available today with some precision, as listed in Table 3. Certain of these can be eliminated

Table 3. Present sources of energy averaged over the earth

Source	Energy, cal cm ⁻² yr ⁻¹
Total radiation from sun	260,000
Ultraviolet light	
λ < 2,500 Å	570
λ < 2,000 Å	85
λ < 1,500 Å	3.5
Electric discharges	4
Cosmic rays	0.0015
Radioactivity (to 1.0 km depth)	0.8
Volcanoes	0.13

immediately from the list of effective supplies. Radioactivity is of no importance now except that it may and probably does produce detrimental mutations as well as an occasional useful mutation. Potassium exists in the bodies of all living organisms but its radioactivity sup-

plies no effective energy for these metabolic processes. The radioactivity of the rocks is unavailable because it is much too violent in character and because it is largely dissipated within the rocks. Cosmic rays are unimportant, as can be seen from the table.

Electrical energy of lightning is too violent to be of direct use, but it does produce some compounds of nitrogen which are useful.

Volcanic heat is of no value today as a source of energy for metabolic process and was probably of minor or in fact negligible importance in the past because of its violent and destructive character and its very sporadic appearance. What is needed for the evolution of life as well as its maintenance is a steady source of energy which will be maintained over billions of years.

Living things require compounds of certain kinds which contain greater quantities of energy than do the degradation products. The particular thermodynamic function that measures the spontaneous character of chemical reactions at constant temperature and pressure is the Gibbs free energy. In any spontaneous reaction this function decreases. Hence, living organisms of the animal or saprophytic plant type must be supplied with chemical compounds of higher free-energy content than the compounds which they discard. In some way free energy must be supplied to these waste products in order that they can be reconverted into the food chemicals again. Such free energy can be supplied by high-energy sources such as sunlight, radioactivity, electric discharges, etc. It could be supplied from heat sources, providing a suitable heat engine operating between high- and low-temperature heat reservoirs is available in accordance with the second law of thermodynamics. Merely high-temperature regions such as water pools alone, supplied by volcanoes or hot springs, are not sufficient to accomplish this. In considering this problem of the energy source for chemical evolution of life, these requirements of thermodynamics should be kept in mind. Though warm water may have accelerated reactions between compounds with suitable free-energy content, warm water alone cannot provide the compounds of high free-energy content.

At present the source of energy is sunlight in the red end of the spectrum, which is absorbed by chlorophyll. However, in prebiological times before the appearance of chlorophyll, ultraviolet light, absorbed in the atmosphere to produce high-energy compounds which in turn were transferred to the oceans, must have furnished most of the free energy required for prebiological evolution. As Table 3 shows, there is only a small fraction of the energy in the solar spectrum below

2,000 Å, but it is larger in amount than all nonradiant sources of energy. It is evident also that the appearance of relatively simple carbon-oxygen-hydrogen compounds would extend the region of absorption into the 2,000 to 3,000 Å region where much larger amounts of energy are available. However, even after 4.5 billion years, living organisms are using only a small fraction of all the energy of sunlight.

Miller's experiments

Miller's synthesis of carbon compounds from simple substances is too well known to require any serious review. The conditions used would not be those used today for the same experiments. It was supposed that the primitive atmosphere contained an appreciable concentration of hydrogen. The high atmosphere of the earth is at some 2000°K or even more, as was shown to be true by Spitzer (1952), and under these conditions elementary hydrogen would be lost very rapidly. Hence today experiments omitting hydrogen as such would appear more appropriate. Also it should be noted that electric discharges were used only because they were easy to provide and not because it was supposed that lightning was the primitive source of energy. It was expected that the electric discharge would activate molecules in a way that would be similar to that produced by ultra-violet light. Table 4 gives the list of substances produced by Miller in one of his experiments. For the first time weighable amounts of substances present in living organisms or readily produced from their

Table 4. *Yields from sparking a mixture of CH₄, NH₃, H₂O, and H₂; 710 mg of carbon was added as CH₄*

Compound	Yield, moles ($\times 10^5$)	Compound	Yield, moles ($\times 10^5$)
Glycine	63.	Succinic acid	4.
Glycolic acid	56.	Aspartic acid	0.4
Sarcosine	5.	Glutamic acid	0.6
Alanine	34.	Iminodiacetic acid	5.5
Lactic acid	31.	Iminoacetic-propionic acid	1.5
N-Methylalanine	1.	Formic acid	233.
α -Amino-n-butyric acid	5.	Acetic acid	15.
α -Aminoisobutyric acid	0.1	Propionic acid	13.
α -Hydroxybutyric acid	5.	Urea	2.0
β -Alanine	15.	N-Methyl urea	1.5

tissues were secured. They were not optically active and a number of unidentified substances were produced.

These experiments were repeated and extended by others, notably by Abelson (1957) in this country. Also the more difficult experiments using ultraviolet light have been done and very similar results obtained. Abelson's experiment showed that various mixtures always gave amino acids if the mixture was reducing, i.e., if excess hydrogen was present in amounts over that required to form water, and that amino acids were not produced if the mixture was on the oxidizing side, i.e., if free oxygen or its equivalent was present. We can conclude that it is most improbable that life would appear in an oxidizing atmosphere. In particular, if all life were destroyed on earth, the oxygen of the atmosphere would probably disappear by combining with carbonaceous compounds. The amount of reduced carbon in living organisms and in soils is very small but the great deposits of shales contain some 10 or 12 times as much carbon as could be oxidized by atmospheric oxygen. This would be brought only very slowly into contact with atmospheric oxygen, and hence the atmosphere would probably remain somewhat oxidizing. Life would probably never evolve again on this planet if it were completely destroyed. Also, if a planet evolved into the oxidized state before well-organized life had appeared, it is probable that no life would evolve on that planet.

Miller's experiments and those of others who have extended this work carry us only a very small way along the long and involved road to the origin of well-organized life. There are many elementary problems which might be mentioned at this point, but possibly one of the most puzzling and also most essential is the introduction of phosphate into effective organic compounds. All inorganic phosphorus compounds are highly unstable except the phosphates, and the calcium, magnesium, and ferrous phosphates are highly insoluble in water (see Table 1). These are mentioned particularly because calcium and magnesium are present in water solution, i.e., in the oceans, as bicarbonates and hence phosphates would be precipitated. In the absence of elementary oxygen, ferrous iron would also be present. Probably organic compounds of some kind which are able to hold phosphate in solution existed in the primitive oceans.

In general, living organisms are excellent chemists and they have found routes for metabolic process which are beautifully adapted to their needs. The production of proteins, chlorophyll, enzymes, and the intricate replicative systems are amazing triumphs of the most skillful of chemists. The mind of man will be able to follow these to some extent in the future.

Development of other planets

It is evident that the processes outlined above could occur on other planets than the earth. But first, nearly all of the solar components of gases must be lost and then conditions must be such that hydrogen can escape at a sufficiently slow rate for the synthesis of very complex carbon compounds to occur. It is possible and indeed probable that this has occurred very generally throughout the entire universe, and it is possible that it has occurred on both Venus and Mars. At present both these planets have carbon in the highly oxidized state of carbon dioxide in their atmospheres, though elementary oxygen has not been detected in either; on the other hand, neither has the highly reduced form of carbon, namely, methane, been detected.

The atmosphere of Venus contains about 1 km atmosphere of carbon dioxide above the cloud layer. Aside from a very small amount of water vapor, no other gas has been detected with certainty in its atmosphere. Several explanations have been offered for the presence of large amounts of carbon dioxide in spite of the affinity of carbon dioxide for silicate rocks which favors its presence as limestone or dolomite. (1) The surface of the planet is arid so that erosion is negligible; hence contact between the gas and solid rocks is ineffective and an equilibrium condition is not approximated. (2) The surface of the planet is covered by oceans and again effective contact is not possible. (3) An excess of carbon dioxide exists such that all surface silicates have been converted to carbonates and the atmospheric carbon dioxide remains in the uncombined state. (4) The temperatures below the atmosphere are so high that the equilibrium pressure is much higher than that appropriate to the earth. The surface temperatures as reported at present are consistent with the first and third alternatives only. It has also been suggested that an excess of hydrocarbons exists and that the surface is covered with oceans of petroleum-like compounds. The temperature in the high atmosphere from infrared measurements is reported to be some -40°C . From the intensities in the rotational spectrum of visible carbon dioxide bands, the temperature is observed to be some 10 or 15°C . But the intensity of the radiation in the centimeter wavelength region indicates temperatures in the region of 300°C . It appears that the first two temperatures must be characteristic of some layers in the atmosphere above the clouds, and the last temperature is generally thought to be that of the solid surface. Sagan (1961) has given reasons for believing that a very effective "greenhouse effect" due to carbon dioxide and water may account for this high temperature. If these high temperatures

do obtain, no life can exist on Venus at the present time. In the past, liquid water may have existed on the planet, the pressure of carbon dioxide may have been low, and hence the greenhouse effect may have been less, the temperature lower, and living organisms may have been present. It would be magnificent had life developed but disappointing if it should have been destroyed.

Mars is a very different planet. It has a thin atmosphere with a pressure at the surface of about one-sixth that of the earth. Carbon dioxide is present, but oxygen has not been detected. The atmosphere is probably mostly nitrogen. It extends to a great height because of the low gravitational field. It is most probable that both hydrogen and oxygen can escape from the planet and hence that the planet may have had extensive oceans in the past. Glaciers would appear to have been probable in this case, but just because the rocks were covered by them carbon dioxide may not have reacted with the rocks effectively and the greenhouse effect may have raised surface temperatures above the very bleak ones that obtain at the present time. Life may have evolved on this planet and in spite of the very arid and cold conditions it may persist to the present time. Carbon dioxide is present as shown by its absorption bands. Water is probably present as indicated by the polar frost caps and the morning haze, but the estimated amounts in the atmosphere are very small. It is probably escaping from the interior because both hydrogen and oxygen should escape at such rates that the observable water would be lost in less than 10^6 years and hence atmospheric water must be continuously replaced.

Life may be present on Mars and its origin would be similar to that outlined for the earth. The changes of color with the seasons, the persistence of the gray areas in spite of dust storms, and, as Sinton (1957) has recently observed, the presence of faint absorption bands, which can be ascribed to the CH band in the light reflected from the gray areas, indicate that life may be present. The proof that this is true would be the most magnificent scientific datum of the twentieth century and would justify all the effort of the space programs of the world. It would substantiate the beliefs of the many students of this subject, namely, that life will evolve when conditions are appropriate in many other places of the universe and that life on earth is not alone in this vast expanse of space that extends for billions of light years in all directions.

References

- Abelson, P. H. (1957), *Ann. N.Y. Acad. Sci.*, **69**:276.
Bates, D. R., and M. R. C. McDowell (1957), *J. Atmospheric & Terrest. Phys.*, **11**: 200.

Miller, S. L., and H. C. Urey (1959), *Science*, **130**:245.

Sagan, C. (1961), *Science*, **133**:849.

Sinton, W. (1957), *Astrophys. J.*, **126**:231.

Spitzer, L. (1952), *Atmospheres of the Earth and Planets*, ed. by G. P. Kuiper, University of Chicago Press, Chicago.

Urey, H. C. (1959), *Handbuch der Physik*, **50**:363ff.

Evolution of Photosynthetic Mechanisms*

Melvin Calvin †

Department of Chemistry and
Lawrence Radiation Laboratory
University of California, Berkeley, California

Introduction

I have found the planning of today's discussion to be particularly difficult, perhaps the most difficult one that I have ever undertaken. The reason for this, I consoled myself, lies in the very nature of the evolutionary process itself. In physical science (and particularly in mathematical sciences) we are accustomed to a single sequence of events, in which each idea is precursor to the next, and one gradually develops a whole pattern of thought—a whole notion from beginning to end—in a single sequence. Those of you who are more familiar with the way biological material has evolved will know that this is not really the way the living organism can be described in its evolutionary

* The preparation of this paper was sponsored by the U.S. Atomic Energy Commission.

† Research Professor of Chemistry, 1960–1961, in the Miller Institute for Basic Research in Science, University of California, Berkeley.

history. The problem with which I am dealing, the problem of photosynthesis, is especially difficult to trace.

It turns out, as I shall try to point out in more specific detail as we go along, that the evolution of photosynthesis entails the fusion of a number of quite independent threads of evolution at some point in time to give rise to the modern process and the modern apparatus as we know it. In trying to describe that sequence of events, I find myself greatly increasing my respect for the novelist who writes historical novels. You know how he does it: He has many chains of events, giving rise to a particular incident at the end, or perhaps at the beginning of the novel, and he is very skillful at starting each of these threads and jumping from one thread to the next, bringing them along so they all come together at the right time and in the right place. I haven't yet been able to do that, and what I am going to have to do is to jump back and forth among the various evolutionary threads that are involved here, which ultimately fuse together to give rise to the very complex process of photosynthesis. The story may appear, therefore, more confused than it really is, since I must jump back and forth between separate evolutionary threads and try to indicate their points of fusion.

Modern Photosynthetic Processes

With this apology, let us begin our study of the evolutionary history of photosynthesis by first describing what we think we know of the modern process toward which we must eventually come. Most of you know that photosynthesis is the process by which living organisms are able to transform electromagnetic energy into chemical energy by inducing the reaction between carbon dioxide and water to evolve molecular oxygen and reduced carbon:

This is the overall process of photosynthesis which has long been recognized as a process for transforming electromagnetic energy, here represented by the quantum, into chemical potential, represented by oxygen in the elementary form and the elements of carbon and hydrogen largely in the oxidation level of carbohydrate.^{1,2,3}

If this were all that we know about the process of photosynthesis, we would be hard pressed to try and predict an evolutionary history which might give rise to this process. Fortunately, in the last decade or two we have learned perhaps more about the process of photo-

synthesis from this point on than in the previous hundred years. This was the stage that was available to us roughly one hundred years ago. Only slow progress was made in increasing the chemical knowledge of photosynthesis until just prior to World War II—beginning in the middle thirties and then going on after the war at an increasingly rapid rate.

What do we know today about the process of photosynthesis? Rather than try to give you a history of how the knowledge has evolved, I am going to (1) put down some of the established things that we know about photosynthesis, represented by the overall reaction, (2) then see which organisms perform this process, (3) then determine what the biological apparatus is within some of the organisms (as far as we can do it), and (4) finally go further on down to the molecular level. You see, the question of the evolution of a process of this sort also raises others: What level shall we deal with? Shall we deal with photosynthesis at the level of the whole organism, the level of the cell, the level of subcellular particles, the level of the macromolecules, or the level of the small substrate molecules that are involved? We should, in fact, deal with all of these, if possible, but this is another complication which makes the organization of such a discussion as this extremely difficult. I am going to try to pick up two aspects of it, the mechanism itself on the substrate, and possibly submolecular level, and the apparatus on the subcellular, or macromolecular level.

Nature of the organisms

I hardly need review for you the nature of the organisms which are capable of performing the process of photosynthesis. Quite obviously, the higher green plants, such as a wheat field or a forest, do this on a grand scale. There is, however, a whole set of other organisms besides the higher green plants which are able to do this, or parts of it, and they represent an important part of the biological scheme of things in the course of our study. These are the marine algae; both the green and the red ones are important in terms of the amount of carbon which is turned over on the surface of the earth per year, as the algae represent the largest single plant family involved in this turnover. Then, there is another group, the blue-green algae, which appear to be more primitive organisms which are capable of doing the entire process of photosynthesis, that is, reducing carbon and evolving oxygen. And, finally, we come to the bacteria, both the green and the red, which are capable of performing part of this conversion

process. The bacteria can transform electromagnetic energy into chemical energy, but not with the evolution of oxygen. They use other reducing agents than water in order to reduce the carbon and therefore they produce other oxidants than oxygen. But the bacteria are able to capture electromagnetic energy from the sun and transform it into chemical potential.

These classifications of organisms really constitute the whole gamut of biological diversity, as far as I am aware of it, which can do all, or some, of this conversion (energy manipulation) process; they all can do the crucial part of it—the quantum absorption and the quantum conversion.

Mechanism of the Photosynthetic Process

A. The path of carbon in photosynthesis

Let us see what we know about the mechanism of the process of photosynthesis itself. Part of this knowledge is a result of the tracer work which was mentioned earlier,^{1,2,3} beginning before the war. My colleague, Sam Ruben, began this work, using radioactive carbon-11. Right after the war in 1945 we took it up again, using carbon-14 labeled carbon dioxide, to examine the sequence of events and determine the sequence of compounds involved in the transformation of CO₂ into carbohydrate. The answer to these questions is now available to us, and we can draw a rather complete road map of the reduction of carbon dioxide. (A simplified version of the carbon reduction cycle is shown in Fig. 1.) The first step in the photosynthetic carbon cycle is the carboxylation of a sugar, ribulose diphosphate, to give phosphoglyceric acid, and this, in turn, can now be reduced to triose phosphate using some kind of reducing agent as well as some pyrophosphate-containing compound. The triose phosphate then goes through a series of rearrangements to produce ribulose diphosphate again, and the carbon cycle can continue.

The light is required to produce these two agents: a reducing agent, here represented by [H] and a particular (pyrophosphate-containing) phosphorus compound (which we shall mention in a moment) to help the reducing agent in the reduction process. This particular phosphorus compound seems to be adenosine triphosphate (ATP) which contains a pyrophosphate linkage. This is of great importance and will be discussed in detail later on.

The major point that I want to introduce at this stage is the idea that the reduction of carbon dioxide through the carbon cycle and

the whole sequence of enzymatic reactions that are involved in this reduction are *dark* reactions. Once we have available the products of the *light* reaction, namely, a reducing agent and some type of "high energy" phosphate, the whole carbon cycle can be operated and car-

FIG. 1. Carbon reduction cycle (simplified version): (1) Ribulose diphosphate reacts with CO_2 to give an unstable 6-carbon compound which splits to give two 3-carbon compounds, one of which is 3-phosphoglyceric acid. The other 3-carbon compounds might be either 3-PGA, as it is known to be in the isolated enzyme system, or some other 3-carbon compound such as a triose phosphate (dashed arrow). (2) PGA is reduced to triose phosphate with ATP and TPNH derived from the light reaction and water. (3) Various condensations and rearrangements convert the triose phosphate to pentose phosphates. (4) Pentose phosphate is phosphorylated with ATP to give ribulose diphosphate. Further carbon reduction occurs via conversion of PGA to phosphoenolpyruvic acid (5) and carboxylation (6), to form a 4-carbon compound (probably oxaloacetic acid). Reactions leading to the formation of some of the secondary intermediates in carbon reduction are also shown.

bon can be taken from CO_2 into a variety of compounds, among them sugar. The sugar can be taken out of the cycle. Every time the cycle turns six times, for example, we can take out a hexose sugar molecule and still have the cycle molecules left. This, indeed, is what happens.

We recognize also that all of the 11 enzymes that are involved in these transformations in the carbon reduction cycle are to be found nearly everywhere widely distributed in the biological world—not limited solely to organisms which are converting solar energy, but also in organisms that have nothing whatever to do with the photosynthetic process. It therefore seems quite clear that at least this sequence, that is, the carbon reduction sequence, undoubtedly evolved in a separate chain of evolutionary events having little or nothing to do in the early stages with the electromagnetic energy conversion process itself.⁴ The electromagnetic energy conversion process appears to produce in a primary act, or very close to it, two materials, a reducing agent and a pyrophosphate linkage, which can then run the carbon reduction cycle.

We can already see the two quite independent evolutionary streams which were joined only very recently in evolutionary history to produce the modern green plant.^{5,6,7} The carbon reduction system was one independent stream. These streams will, of course, break up into finer parts as we go along, but this is our beginning.

B. Quantum conversion in photosynthesis

Let us now return to the photochemical process itself. Having separated out the carbon reduction system as a distinct evolutionary stream, I am going to leave it since there is nothing unique about it for photosynthetic organisms except the combination of the product of the light reaction with a certain collection of enzymes, all of which can be found, either separately or in various combinations, in non-photosynthetic organisms.^{8,9} Therefore, the carbon reduction cycle had a separate evolutionary history until the recent times.

Let us now see what more we can say about the quantum conversion process in photosynthesis. We can say a good deal about it, although not nearly as much as we can about the carbon reduction process. We do not have anywhere near the detailed knowledge of the quantum conversion process that we do of the carbon reduction process. Figure 2 represents the structural formulas of the two molecules which we believe to be essential for running the photosynthetic carbon reduction cycle. (There are undoubtedly others of which we are still unaware required for oxygen evolution as well.) To run the carbon cycle we need the reducing agent, which is a pyridine nucleotide in its reduced form. An adenine and pyridine moiety are tied together by two ribose sugars and a pyrophosphate link to give the molecule known as diphosphopyridine nucleotide. Actually, in photosynthesis

it seems that there is a molecule very similar to this, but involving another phosphate group on one of the ribose molecules, and so I shall actually use the triphosphopyridine nucleotide in its reduced form as the structural formula for the reducing agent which is required to run the carbon reduction cycle.

The possibility exists that still another, and perhaps more specific, reducing agent might be used by photosynthetic organisms in the reductive splitting of the initially produced carboxylation product

TRIPHOSPHOPYRIDINE NUCLEOTIDE
(OXIDIZED FORM) (TPN^+)

ADENOSINE TRIPHOSPHATE (ATP)
IN ADENOSINE DIPHOSPHATE (ADP),
TERMINAL PHOSPHATE IS
REPLACED BY $-\text{OH}$

NICOTINAMIDE PORTION OF
TPNH (REDUCED TPN^+)

FIG. 2. Structural formulas of triphosphopyridine nucleotide and adenosine triphosphate, two of the agents required to run the photosynthetic carbon cycle.

(Step 1, Fig. 1).¹⁰ If so, it is almost certainly as good a reducing agent as TPNH and may or may not be structurally and kinetically related to it. If such a specific photosynthetic reducing agent functions in green plants, it will, in all probability, have been a late addition in the evolutionary development of a higher efficiency, since we already know that the cycle can operate through TPNH.

The other molecule which is essential for running the cycle and which clearly must come somewhere from the *photochemical* reaction is the adenosine triphosphate (ATP). Here, there are two pyrophosphate linkages, and the important one for our purposes is the ter-

minal pyrophosphate link (Fig. 2). These are the two molecules that are required in order to move the cycle around, and clearly these must be manufactured as a result of the photochemical transformation.

How much do we know about how the photochemical transformation manufactures those two substances? Here, we are not so thoroughly informed, but a good deal, nevertheless, is known and some of it is of considerable importance in guiding our thinking as to what the evolutionary relationships between the photosynthetic equipment and other equipment of living organisms might be.

1. Photoinduced Redox System

The principal photochemical reaction we now know is, first, the absorption of light by chlorophyll to produce some kind of an excited chlorophyll, either a molecule or molecular aggregate. (I don't mean this to be a separate chlorophyll molecule in solution, but simply the chlorophyll as it exists in the photosynthetic equipment of the organisms.) This electronically excited molecule must then undergo some kind of transformation; for example, it may react with another molecule or molecules to produce a separation of an oxidant from a reductant.^{11,12} I am using this language first because of a bit of confusion that has arisen in the meaning of these terms. The *oxidant* will eventually become molecular oxygen; the *reductant* will eventually become a reduced compound, pyridine nucleotide. The pyridine nucleotide, together with the ATP, for which we have not yet described a formation mechanism, will then go on to drive the carbon cycle.

Oxidant and reductant are the chemists' terms for what happens after the excited chlorophyll loses its energy to some molecule, or collection of molecules, if any redox system is directly involved. The biologist has been accustomed to writing these two things in different terms. Following van Niel,¹³⁻¹⁶ he has generally associated the term M (Fig. 3) with water and has called the oxidant ([O] in Fig. 3) *hydroxyl*, or [OH], but he has been very careful to put a bracket around it. (Those of you who know what the meaning of a bracket is will understand the significance of this; when you see a biochemist putting a bracket around something of this sort it means that he doesn't really know what he is talking about. It is a general representation, not a chemical formula.) The reductant ([R] in Fig. 3), according to the biologists, has been called [H] (hydrogen), and this had led many to suppose that the primary process of quantum conversion involves the splitting of the water molecule itself. What is meant by the van Niel

theory, at least in chemical terms, is the creation of a reductant of some general character, whose nature we do not know, and of an oxidant, also of some general character whose nature we do not know as yet. These two things are both very closely associated with and must ultimately come from water, as given by the stoichiometry of the primary reaction of photosynthesis in the first place.

In more recent years, still another terminology has entered into this discussion and it comes from quite a different source. The physicist has called the reductant the "electron" and what is left after you

FIG. 3. Simplified photosynthesis scheme. The quantum is first absorbed by the chlorophyll molecule; then something happens (p for primary) to the excited chlorophyll to produce two chemical species, [O] and [R], for example, which can go on, one of them [O] to become molecular oxygen in some way (1), and the other one [R] leading to the reduction of CO_2 to carbohydrate (2). Along these two routes, various other energy-containing species may be created (ATP or $\sim\text{P}$). ATP would be an energy-storage product. This may be created on either, or both, sides. There may be back reaction (3) between the oxidants and reductants, which also could create products of higher energy. The obvious one here is, of course, the pyrophosphate linkage in ATP.

take an electron away from a molecule is called a "hole."¹⁷ These are the physicists' terms for the same phenomenon. You must not become confused about the terminology because all of these—oxidant-reductant, hydroxyl-hydrogen, electron-hole—are different names for essentially the same thing. What we are trying to do now is to discover exactly the best way to describe these things in ultimate and intimate detail.

I introduce the terminology of the physicist because in the last few years we have learned a number of the reactions of excited chlorophyll and one of them is an electron transfer reaction which is observable spectroscopically. An electron is transferred from iron in the divalent state to give iron in the trivalent state,^{18,19,20,21,22} with the

electron located in an as yet unknown place. It is an important recognition that this phenomenon occurs and occurs very quickly after the chlorophyll absorbs the light.* The excited chlorophyll in some way is able to extract an electron from the ferrous iron compound, at present associated with the chlorophyll in modern organisms in the form of cytochrome, to produce the ferricytochrome and an electron in some molecules as yet undesignated.^{19,24} This appears to be an important connection between a molecule that is unique to photosynthetic plants, namely, chlorophyll, and certain kinds of molecules which are not unique to photosynthetic plants, namely, the iron cytochromes (iron hemes). The iron hemes have universal distribution and this is an important fact to remember.

In addition, we now know that electrons must ultimately find their way to pyridine nucleotide, which picks up a proton from the water, making reduced pyridine nucleotide. The oxidized iron, or something close to it, will *eventually* take electrons from water, giving rise to the ferrous iron and molecular oxygen.

At the same time that all these things are happening, somewhere along the line, either on the way from the intermediate oxidant to oxygen (reaction 1, Fig. 3), or on the way from the intermediate reductant to the pyridine nucleotide (reaction 2, Fig. 3), or, perhaps, in a recombination reaction in which the electron falls back into the hole (reaction 3, Fig. 3), we also create adenosine triphosphate. The ATP is designated by $\sim P?$, which represents "high-energy phosphate" linkages. The reactions in Fig. 3 indicate possibilities only, and not knowledge of three different ways (places) in which pyrophosphate could be created: (1) The fall of the intermediate oxidant toward oxygen; (2) the fall of the intermediate reductant (perhaps a sulfhydryl group) to the pyridine nucleotide which would, perhaps, give rise to pyrophosphate; or (3) perhaps the energy of recombination of the hydrogen-hydroxyl (electron-hole), which could also give rise to a number of pyrophosphate linkages.

2. Photoinduced Dehydration

A more profound departure from the basic redox primary photo process is possible, particularly in the light of the recently indicated²⁵

* A recent modification¹⁴ of the van Niel generalization inserts a ferrocyclochrome ahead of the water molecule as the primary electron donor to the excited chlorophyll, but it does not specify the primary fate of the excited electron which must be removed from chlorophyll. The oxidized cytochrome is presumed capable of oxidizing water to oxygen, with the concomitant formation of ATP, a suggestion similar to that of Bassham²³ and corresponding to reaction 1 in Fig. 3.

reversibility of at least some of the steps of oxidative phosphorylation. Thus, there is evidence that in mitochondria it is possible to produce the reaction

If an *independent* (nonredox) method of dehydration could be found for producing ATP according to the reaction,

then *both* ATP and TPNH could be photoproduced without calling upon a photoinduced direct electron transfer reaction.

We already have a precedent for the idea that an optically excited pi-electron system can have an increased affinity for water leading to its hydration by an only *very slowly reversible* process so that energy may be trapped in this manner.²⁶

For example, if the 9-10 enol in chlorophyll were to add orthophosphate (when excited), an enol phosphate could be produced, which presumably would be capable of phosphorylating ADP to make the required ATP.²⁷ Part of this would then be used to reverse the DPNH-cytochrome reduction to produce the ultimately necessary separation of oxidant and reductant (water splitting) required for O_2 production and CO_2 reduction.

The not inconsiderable difficulty with such a plan as this is the necessity for producing a *good many more than one* ATP for each quantum absorbed by chlorophyll. Even if a way of circumventing this difficulty were found, it remains fairly clear that such a device

would be a rather recent evolutionary addition to an already highly developed biosynthetic energy-manipulating system.

Pyrophosphate Linkage in Nonphotosynthetic Processes

The appearance of pyrophosphate linkage in a variety of organisms is well known. In practically all organisms, there are mechanisms for producing ATP which do not involve photosynthetic mechanisms at all. One of them is a reversal of one reaction in which ATP is used in the photosynthetic cycle (triose phosphate dehydrogenase). By running the reaction backward (Step 2, Fig. 1) one can make ATP. A more important source is a reaction which apparently involves iron—the cytochromes, involving also the oxidation and reduction of the pyridine nucleotide. The two reactions together are involved in the creation of ATP in nonphotosynthetic organisms. This process of the oxidation of pyridine nucleotide by the passage of electrons from pyridine nucleotide back to oxygen through the iron cytochromes, with the concomitant formation of ATP, is known as oxidative phosphorylation. It leads to the creation of more ATP than does the substrate oxidation process. The return of a photoexcited electron of chlorophyll through all or part of a similar chain could produce the necessary ATP (see Fig. 3).

Thus the creation of both the reduced pyridine nucleotide and the ATP are not unique to photosynthetic processes. These processes also occur in nonphotosynthetic organisms.⁶ We know something about

how pyridine nucleotide is created, but we know relatively little about how ATP is created in oxidative phosphorylation in which the electrons pass from reduced pyridine nucleotide through iron back to oxygen. This is one of the major problems of energy transformation in all biological organisms.

We have now split up the *photo* process of photosynthesis into two other streams of evolutionary development, the stream which gave rise to pyrophosphate (ATP) and the stream which gave rise to pyridine nucleotide. Neither of these necessarily involves the photo process directly. This leads us to the conclusion that the appearance of the photo reaction, or the coupling of the photo reaction, with the creation of ATP and of reduced pyridine nucleotide was a very late thing in the evolutionary scheme.⁴ You see that we are forced, now, to consider the question of the origin of life in discussing the origin of photosynthesis. We cannot dodge that issue, and we are indeed considering it and doing so in a much more sophisticated way than has been possible up until recent times.

I shall indicate some of the states that we need, in order to try to focus your attention on the separate evolution of mechanisms for making ATP, mechanisms for making the molecules which are involved in the creation of ATP today, mechanisms for creating pyridine nucleotide, and, finally, how the light-capturing molecule, chlorophyll, may have appeared and was coupled to the other energy-transforming processes. This is really the story in principle, and I now want to go through it quickly and try to give you some idea of how I think these things might have occurred.

Evolution of the Photosynthetic Apparatus in the Green Plant

Figure 4 shows the apparatus in the green plant (the chloroplasts) which is responsible for performing the process of photosynthesis. It is perhaps necessary to say a few words here about the relationship of the tangible physical material that performs photosynthesis as it can be seen on the subcellular, but still visible, level. I shall then discuss the macromolecular level (where this apparatus cannot yet be seen), and, finally, go to the substrate level where we can again deal with things in a chemical way.

Three different kinds of chloroplasts are shown in Fig. 4, illustrating the highly ordered array of layers in all of the three types of organisms: a unicellular green alga, a blue-green alga which does not have

FIG. 4. Chloroplasts from a unicellular green alga, from a blue-green alga, and from tobacco, and mitochondria from guinea pig pancreas. (A) Mitochondria from guinea pig pancreas. (B) Chloroplast with several mitochondria. *Chlamydomonas*. (Sager.) (C) Entire cell of *Anabaena* showing lamellae. (Vatter.) (D) Tobacco chloroplasts maintained 24 to 36 hours in dark before fixing with permanganate. (Weier.)

a chloroplast (the layers are still present, however, winding their way in and out through the entire cell), and a chloroplast from a higher plant (tobacco) showing the layering of the green material very clearly. The layers (lamellae) themselves are constructed of arrays of macromolecular subunits which we now think we can see.²⁸ Figure 5 gives a model for chloroplast lamellar structure and Fig. 6 is an electron micrograph of frozen dried spinach chloroplast supernatant.

FIG. 5. Model for the lamellar structure within a spinach chloroplast: (a) Osmium-staining layer of the lamellar structure. Thickness 30 Å in the intergrana regions and 60 Å in the grana regions. (b) Particles forming the granular inner surface of the two layers making up the lamellar structure. The packing of oblate spheres would not be as simple as illustrated, since the central axis of both layers would not be in the same vertical plane shown here.

Figure 4 shows the high degree of order in the chloroplasts, and, furthermore, that this high degree of order exists in other elements in the cell, such as the mitochondria, which perform other functions (formation of ATP by oxidative phosphorylation of pyridine nucleotide).²⁹ The purpose of Fig. 4 is to show the similarity of structure between the photosynthetic apparatus and material which is not photosynthetic, and to show also that it is a highly ordered array in all cases. This highly ordered array must be achieved in some systematic way from molecules which themselves are ordered by virtue of the atoms of which they are made.

FIG. 6. Frozen dried spinach chloroplast sonicate; 880 Å diameter PSL (polystyrene latex) markers.

Chlorophyll structure

The actual detailed structure of the one molecule unequivocally associated with the capture of light and its transformation, i.e., chlorophyll, is shown in Fig. 7. This shows the structure of some of the different kinds of chlorophyll that are known: The first is protochlorophyll, which appears in etiolated plants grown in the dark. When such plants are placed in the light, the protochlorophyll is converted to chlorophyll. The principal difference between protochlorophyll and chlorophyll is the addition of two extra hydrogen atoms at the double bond in ring D. Bacteriochlorophyll is the molecule which is responsible for the capture and conversion of light in the purple and green bacteria; it differs from green plant chlorophyll in having a second, dihydropyrrole ring in it.

We must devise some way of making those ordered chloroplast structures which were seen in Figs. 4, 5, and 6, and we must envisage some way of evolving this particular molecule, chlorophyll, belonging to the general class of tetrapyrrolic substances known as porphyrins. These two things—ordered array within the cells and the development of chlorophyll itself—are two essential features of our evolutionary scheme for the process of photosynthesis.

The structural feature, the appearance of order and structure, is something common to the evolution of all living organisms, and belongs to the general discussion of how ordered structures may be evolved from nonliving materials. This is really part of the problem of the origin of life.

FIG. 7. Structure of protochlorophyll, chlorophyll, and bacteriochlorophyll.

Chemical Evolution

I wish to discuss briefly the beginnings of chemical evolution, starting with the molecules of the primitive atmosphere being subject to a primitive photosynthesis, using the far ultraviolet or radiation from the radioactivity of the earth's crust to transform them. The earliest molecules on the surface of the earth were those shown on the top row of Fig. 8, particularly methane, ammonia, and water. If these molecules are subjected to radiation of energy great enough to break the bonds of carbon-carbon, carbon-hydrogen, hydrogen-hydrogen, nitro-

FIG. 8. Primeval and primitive organic molecules.

gen-hydrogen, hydrogen-oxygen, which can be done by ionizing radiation,³⁰ such as the beta rays of potassium-40 which are plentiful in the earth's crust, or with ultraviolet light of wavelengths shorter than 2,200 Å,³¹ then the atoms which are so formed may reorganize to form more complex molecules, a few of which are shown on the bottom row of Fig. 8. You already recognize these molecules as being the present-day substrate materials (formic acid, acetic acid, succinic acid, and glycine) upon which all living organisms operate. Glycine, shown here, is the only nitrogen-containing compound in the bottom row of Fig. 8, and it is the simplest of the amino acids, of which the proteins are constructed. By exchanging one of the carbon-bound hydrogen atoms of the glycine for any of a group of other atoms, some 20 different amino acids can be built up.

FIG. 9. Radioautograph of paper chromatogram showing compounds which result after irradiation of a mixture of C^{14} , ammonia, and water with 5-Mev electrons.

In the first experiment of this type in 1950, in which we used the cyclotron as a source of ionizing radiation,³⁰ we started with CO₂, hydrogen, and water, and were able to get, by random transformation processes, reduced carbon compounds such as formic acid, acetic acid, and succinic acid. In later experiments, in which ammonia was added to the initial mixture following Miller,³² glycine was obtained. Still more recently (in the last three or four months) we have performed this experiment again, but instead of depending upon ordinary analytical methods to find these randomly occurring compounds, we have used carbon-14 labeled methane in the primitive gas mixture, thus providing radioactive carbon atoms which could be followed around. The discharge from a 5-Mev electron linear accelerator was passed through the mixture of methane, ammonia, and water, and we took the water solution containing the product from this bombardment and spread it out on a piece of filter paper in a systematic way.³³

Figure 9 shows the results of one of these bombardment experiments. It is a photograph of the darkened x-ray film which results when a paper chromatogram containing radioactive products is placed on top of an x-ray film. Wherever there is a black spot on the film a particular compound has been located. We can tell what the nature of the compound is by where it is located on the film with respect to its origin. All the different nonvolatile radioactive compounds which result from one particular bombardment are shown in Fig. 9, and about a dozen compounds have separated out.

We have been able to identify in this way some half dozen compounds,* including adenine, glycine, alanine, and various other amino acids and sugars, some fatty acids and some hydroxy acids—the very things of which today's living matter is composed. One of the compounds, representing about 60 per cent of the total, is urea. We find in neutral and acidic fractions a large number of compounds, including lactic acid and sugars. You can also see that alanine and glycine represent a very small amount of the total. Down in the lower center of the chromatogram are the nucleosides and further up are the bases. There are present in this irradiated mixture adenine, cytosine, guanine, thymine, and perhaps other as yet undetermined bases. Thus, such random processes as these may give rise to all the simple compounds that are needed by present-day living organisms.^{34,35}

Having made these simple compounds (particularly the amino acids) by the random methods, we can build them up into proteins

* HCN was identified in the aqueous solution by a separate procedure.

in various ways. Aside from the more or less laborious and specific methods involving special protective or activating groups, at least two simpler methods, possibly applicable to primitive conditions, have been successfully demonstrated in the laboratory recently. The first involves heating amino acid mixtures in molten glutamic acid together with some polyphosphoric acid to produce a mixed polypeptide resembling protein.³⁶ The second involves heating the amino

FIG. 10. Protein structure. Simple structural principles. Variety of chemical reactivity.

acid in an aqueous ammonia solution to produce a polypeptide of intermediate size.³⁷

The proteins themselves can take on a specific structure which is shown in Fig. 10. The protein is a combination of amino acids, and the helical structure is built in into the linear array of the amino acids because of the particular arrangement of carbon, hydrogen,

nitrogen, and oxygen atoms in such a chain. Figure 11 shows how the helical structure can take on visible order. The upper photograph is an electron micrograph of single filaments of protein which is a component of collagen. When the protein filaments aggregate, as shown in the lower photograph, they do so in a specific ordered array because of the particular arrangement of amino acids in the proteins. Here you can begin to see the appearance of the *visible* order that must be generated to create mitochondria, chloroplasts, and other subcellular particles. This generation of order is, of course, common to all living things and is not unique to photosynthesis. One can gen-

FIG. 11. Electron micrograph of collagen filaments. (a) Filaments of collagen, a protein which is usually found in long fibrils, were dispersed by placing them in dilute acetic acid. This electron micrograph, which enlarges the filament 75,000 times, was made by Jerome Gross of the Harvard Medical School. (b) Fibrils of collagen formed spontaneously out of filaments such as those shown above when 1 per cent of sodium chloride was added to the dilute acetic acid. These long fibrils are identical in appearance with those of collagen before dispersion.

erate order, beginning from the primitive molecules of the early earth's atmosphere (Fig. 8), through proteins (Figs. 10 and 11) into the subcellular material itself (Fig. 4).

Development of Rudimentary Catalysts

Let us now turn to the question of the generation of the porphyrins, which seem to be central not only to the capture of light as represented by chlorophyll but to the appearance of adenosine triphos-

FIG. 12. Evolution of a catalyst (for $\text{H}_2\text{O}_2 \rightarrow \text{H}_2\text{O} + \frac{1}{2} \text{O}_2$). Figures at bottom indicate catalytic activity per milliliter per second at 0°C .

phate in present-day organisms and perhaps to the appearance of ATP in primitive organisms as well.

Figure 12 shows that, starting with the primitive function of iron for the decomposition of hydrogen peroxide, which will be formed in the seas either by ultraviolet radiation or by K^{40} radiation, the iron catalysis can be improved by a factor of 1,000 if it is built into a porphyrin. If we now transform this iron further by encasing the heme into a folded protein and make the molecule of catalase, the catalytic function is improved by another factor of 10 million for this particular peroxide decomposition reaction.³⁵

This fact is of great importance because I believe that peroxide appeared in the primitive seas of the earth at the very earliest stages as a result of both the ultraviolet radiation at the top of the atmosphere and of the potassium-40 radioactivity in the earth's crust. This peroxide can now serve as an evolutionary selection pressure³⁸ to improve the catalytic function of iron from the bare iron to the iron heme to the iron heme-protein combination.

The way in which this can occur is shown by having a look at the way in which hemes are synthesized by modern living organisms (Fig. 13). We start with succinic acid and glycine, which were made by random synthesis from the primitive earth's atmosphere, and by combining these two substances, we make the alpha-amino-beta-keto-adipic acid which then decarboxylates to give the delta-amino-levulinic acid, two of which can combine to form the heterocyclic pyrrole ring. Then there follows a series of oxidation and condensation steps to give rise to the tetrapyrrole ring.³⁹ This reaction is a spontaneous one which involves a number of oxidation steps, several of which are almost certainly catalyzed by iron. The oxidation is achieved either by oxygen or peroxide under the influence of iron and presumably better achieved by iron in a porphyrin than by bare iron. Therefore, once the porphyrin is formed, more of it will be formed because of this autocatalytic self-selection mechanism.^{4,7}

Pyrophosphate formation

This idea is important because the mechanism of the formation of pyrophosphate seems to involve the oxidation of iron. In the last few months, we have been able to demonstrate that one can generate pyrophosphate in aqueous media by simply allowing hydrogen peroxide to oxidize ferrous iron in the presence of orthophosphate.⁴⁰ In

this reaction, a certain amount of orthophosphate is converted into pyrophosphate. The reaction may be written as follows:

* The half circle around the iron symbol is introduced to represent any other coordinated atoms or groups.

I believe this to be evidence of the primitive way in which the highly evolved oxidative phosphorylation which takes place today began. The complexing of phosphate by ferrous iron, followed by the withdrawal of an electron from the ferrous iron to make ferric iron, the elimination of a water molecule to make pyrophosphate, reduction of the ferric iron to ferrous, completes a cycle for the formation and the liberation of the pyrophosphate linkage. This is now demonstrated in a simple system, and I think it will not be long before we will be able to demonstrate it in the highly evolved iron systems that are used in oxidative phosphorylation, both in plants and in animals, and which are also used in photosynthetic phosphorylation probably in a similar manner.

You can see here a driving force which will give rise to the porphyrin molecule. The driving force is the peroxide present in the ocean and the usefulness of transforming orthophosphate to pyrophosphate in aqueous solutions so the pyrophosphate can then be used to assist the combination of amino acids to make proteins. This was the evolutionary sequence which gave rise first to the porphyrin and second to a mechanism for manufacturing pyrophosphate.

Coupling

As yet we have suggested no mechanism for using light to perform these processes. All that would be required in the later stages is to

find a way of removing the electron from the iron, not with hydrogen peroxide but with light, in order to couple the photochemical reaction to what we now know to be nonphotochemical processes.

I think this event happened very late in the evolutionary scheme, and the evidence for it lies in the fact that the chlorophyll molecule is today manufactured by a sequence of reactions almost identical with the sequence of reactions used to manufacture the heme, but just before the iron is put into the heme, a branching occurs, leading not to heme but to the chlorophyll molecule in which magnesium is situated (Fig. 14). I think the reason for that reaction is, first, that the

Fe - PROTOPORPHYRIN NO.9

FIG. 14. Structural relations between heme and chlorophyll.

light-absorbing ability of the heme itself is very poor. Although heme is red, it does not have anywhere near the light-absorbing capacity of chlorophyll, and one of the reasons for the evolutionary selection of magnesium chlorophyll (magnesium chlorin) is the fact that the absorption of light by a magnesium chlorin is several thousand times greater than that of the iron porphyrin. Secondly, something very special about the electronic structure of the magnesium and of the packing together of the chlorophyll molecules in a crystal lattice, leading to the separation of electrons from the chlorophyll,¹¹ is better achieved by the chlorin than it is by the porphyrin. Finally, if the dehydration-phosphate activation idea (by the 9-10 enol of

chlorophyll) turns out to play a role, we would then have a third powerful selective factor favoring the chlorophyll structure.

The mechanism and the detailed chemical and physical reasons for this obvious advantage of the chlorophyll over the porphyrin remain for the future to discover.

References

1. J. A. Bassham and Melvin Calvin, *The Path of Carbon in Photosynthesis*, Prentice-Hall, Inc., Englewood Cliffs, N.J. (1957).
2. J. A. Bassham and Melvin Calvin, *The Photosynthesis of Carbon Compounds*, W. A. Benjamin, Inc., New York (1962).
3. J. A. Bassham, Photosynthesis, *J. Chem. Education*, **36**:548-554 (1959); J. A. Bassham and M. R. Kirk, New aspects of photosynthesis, *J. Chem. Education*, **38**:151-154 (1961).
4. M. Calvin, Evolution of enzymes and the photosynthetic apparatus, *Univ. Calif. Radiation Lab. Rept UCRL-3915* (Aug., 1957). Also, M. Calvin, Evolution of enzymes and the photosynthetic apparatus, *Science*, **130**:1170-1174 (1959).
5. M. Calvin, Chemical evolution and the origin of life, *Idea and Experiment*, vol. 2, no. 4 (June, 1953).
6. M. Calvin, Chemical evolution and the origin of life, *Am. Scientist*, **44**:248-263 (1956).
7. M. Calvin, Round trip from space, *Evolution*, **13**:362-377 (1959).
8. G. Milhaud, J. P. Aubert, and J. Miller, Le métabolisme du carbone dans le chimioautotrophie, *Compt. rend.*, **243**:102 (1956).
9. R. C. Fuller and M. Gibbs, Aberrant patterns of some photosynthetic enzymes, *Plant Physiol.*, **31**, Suppl.: xxi (1956).
10. J. A. Bassham and M. R. Kirk, Dynamics of the photosynthesis of carbon compounds. I. Carboxylation reactions, *Biochim. et Biophys. Acta*, **43**:447-464 (1960).
11. M. Calvin, Some photochemical and photophysical reactions of chlorophyll and its relatives, *Light and Life Symposium*, Johns Hopkins Press, Baltimore (1961), pp. 317-355.
12. M. Calvin, Quantum conversion in photosynthesis, *J. Theoret. Biol.*, **1**:258-287 (1961).
13. C. B. van Niel, Evolution as viewed by the microbiologist, in *The Microbe's Contribution to Biology*, Harvard University Press, Cambridge, Mass. (1956), pp. 155-176.
14. R. Y. Stanier, Photosynthetic mechanisms in bacteria and plants: development of a unitary concept, *Bacteriol. Rev.*, **25**:1-17 (1961).
15. D. I. Arnon, Cell-free photosynthesis and the energy conversion process, *Light and Life Symposium*, Johns Hopkins Press, Baltimore (1961), pp. 489-564.
16. D. I. Arnon, Conversion of light into chemical energy in photosynthesis, *Nature* (London), **184**:10-21 (1959).
17. M. Calvin, The photosynthetic carbon cycle, *J. Chem. Soc.*, **1956**:1895-1915. M. Calvin, Photosynthesis, in *Radiation Biology and Medicine*, ed. by W. D. Claus, Addison-Wesley Publishing Company, Reading, Mass. (1958). M. Calvin, Energy reception and transfer in photosynthesis, *Rev. Mod. Phys.*, **31**:147-156

- (1959). M. Calvin, Free radicals in photosynthetic systems, *Rev. Mod. Phys.*, **31**:157-161 (1959).
18. H. Lundegardh, On the oxidation of cytochrome *f* by light, *Physiol. Plantarum*, **7**:375-382 (1954). H. Lundegardh, Spectrophotometric investigation on enzyme systems in living objects. IV. Kinetics of the steady state, *Biochim. et Biophys. Acta*, **35**:340-353 (1959).
 19. M. D. Kamen, Comments on the function of heme proteins as related to primary photochemical processes in photosynthesis, *Light and Life Symposium*, Johns Hopkins Press, Baltimore (1951), pp. 483-488.
 20. B. Chance and Lucille Smith, Respiratory pigments of *Rhodospirillum rubrum*, *Nature* (London), **175**:803-809 (1959). Lucille Smith, Reactions of cytochrome pigments in photosynthetic bacteria, *Light and Life Symposium*, Johns Hopkins Press, Baltimore (1961), pp. 436-442.
 21. B. Chance and M. Nishimura, On the mechanism of chlorophyll cytochrome interactions: The greater insensitivity of light-induced cytochrome oxidation in *Chromatium*, *Proc. Nat. Acad. Sci. U.S.*, **46**:19-25 (1960).
 22. W. Arnold and R. K. Clayton, The first step in photosynthesis: Evidence for its electronic nature, *Proc. Nat. Acad. Sci. U.S.*, **46**:769-776 (1960).
 23. J. A. Bassham, Energy utilization through coupled systems, *Radiation Research*, Suppl. **2**:497-502 (1960).
 24. M. D. Kamen, in *Enzymes: Units of Biological Structure and Function*, Academic Press, Inc., New York (1956), p. 483.
 25. B. Chance, Energy-linked cytochrome oxidation in mitochondria, *Nature* (London), **189**:719-725 (1961).
 26. D. Shugar and K. L. Wierzchowski, Reversible photolysis of pyrimidine derivatives including trials with nucleic acids, *Biochim. et Biophys. Acta*, **23**:657-658 (1957). D. Shugar and K. L. Wierzchowski, Photochemistry of nucleic acids, nucleic acid derivatives and related compounds, *Postepy Biochem.*, **4**:243-296 (1958).
 27. H. H. Wasserman and David Cohen, 1-Alkoxyvinyl esters of phosphoric acids as phosphorylating agents, *J. Am. Chem. Soc.*, **82**:4435-4436 (1960).
 28. R. B. Park and N. G. Pon, Correlation of structure with function in *Spinacea oleracea* chloroplasts, *J. Mol. Biol.*, **3**:1-10 (1961).
 29. M. Calvin, From microstructure to macrostructure and function in the photosynthetic apparatus, *Brookhaven Symposia in Biology*, **11**:160-180 (1958).
 30. W. M. Garrison, D. C. Morrison, J. G. Hamilton, A. A. Benson, and M. Calvin, Reduction of carbon dioxide in aqueous solutions by ionizing radiation, *Science*, **114**:416-418 (1951).
 31. W. E. Groth and H. v. Weysenhoff, Photochemical formation of organic compounds from mixtures of simple gases, *Planetary Space Sci.*, **2**:79-85 (1960).
 32. S. L. Miller, Production of some organic compounds under possible primitive earth conditions, *J. Am. Chem. Soc.*, **77**:2351-2361 (1955). S. L. Miller and H. C. Urey, Organic compound synthesis on the primitive earth, *Science*, **130**:245-251 (1959).
 33. C. Palm and M. Calvin, Primordial organic chemistry. I. Compounds resulting from the irradiation of C^3H_4 , *J. Am. Chem. Soc.*, **84**:2115 (1962). See also Irradiation of methane, ammonia, hydrogen and water, Univ. Calif. Radiation Lab. Rept. UCRL-9519 (Jan. 31, 1961).
 34. For a more complete discussion of the subject of chemical evolution and the

44 The Nature of Biological Diversity

- origin of life, see M. Calvin, *Chemical Evolution*, Condon Lectures, Oregon State Board of Higher Education, Univ. of Oregon Press, Eugene (1961).
35. M. Calvin, The chemistry of life. 3. How life originated on the earth and in the world beyond, *Chem. Eng. News*, **39**:96-104 (May 22, 1961).
36. S. W. Fox, K. Harada, and A. Vegotsky, Thermal polymerization of amino acids and a theory of biochemical origin, *Experientia*, **15**:81-84 (1959). S. W. Fox, How did life begin? *Science*, **132**:200-208 (1960).
37. J. Oró, Direct synthesis of polypeptides. I. Polymerization of glycine in aqueous ammonia, *Arch. Biochem. Biophys.*, **93**:166-171 (1961).
38. R. Gerschman, Oxygen effects in biological systems, *Proc. 21st. Intern. Physiol. and Pharmacol. Congress* (1959), pp. 222-226.
39. D. Shemin, Biosynthesis of porphyrins, *Harvey Lectures*, **50**:258-284 (1954-1955).
40. J. A. Barltrop, Private communication.

Biochemistry and Evolution

Ernest Baldwin

Department of Biochemistry
University College, University of London

As N. W. Pirie (1937) has written, “‘Life’ and ‘Living’ are clearly words that the scientist has borrowed from the plain man. The loan has worked satisfactorily until comparatively recently. . . . Now, however, systems are being discovered and studied which are neither obviously living nor obviously dead, and it is necessary to define these words or else give up using them and coin others.” With the discovery that viruses are crystallizable nucleoproteins, it began to be necessary for the first time to realize that there is no fundamental gap between what is living and what is not. The effects on biological thinking have been profound and far-reaching. The origin of life, formerly no more than a pseudo problem upon which time, energy, ink, and paper had been most generously—and uselessly—lavished, now became a true and a real problem, worthy at last of serious scientific consideration.

The origin of life has become something to be sought in times far remote from our own—at some early time in terrestrial history. Already many stimulating essays on this important matter have been written, speculation has abounded, and a major symposium has been

held in Moscow (Oparin, 1957) with distinguished scientific contributors from all over the world.

H. C. Urey, among others, has written at length about the origin of the earth and of its early atmosphere, and has concluded that this was originally a reducing rather than the oxidizing medium it is today (Urey, 1959). It has been pointed out that ultraviolet and other forms of radiation, acting upon this primitive atmosphere, must probably have led to the synthesis of large quantities of small-molecular organic compounds and the production of what has been called a "primordial soup."

Miller (1957) has shown—and his observations have been repeatedly confirmed—that among the products of irradiation of gaseous mixtures approximating in composition to that of the earth's primitive atmosphere glycine, formic, acetic, and succinic acids appear in high yields, together with a remarkably assorted collection of other organic materials (Table 1). It can hardly be without significance that even today these are still the starting materials for the biosynthesis of many elaborate compounds—acetate for the synthesis of fatty acids, sterols, and steroids, and glycine and succinate for that of porphyrins.

The gap between this stage in chemical evolution and the eventual emergence of the first organized, self-replicating system is a difficult one to bridge in the present state of knowledge. It has been said that, given enough monkeys, enough typewriters, and enough time, one of the animals would eventually produce a typescript of all Shakespeare's sonnets. There seems to be no reason why this should not be true. Equally, given a large enough number of small molecules as letters of a biochemical alphabet and a few billion years to do it in, there seems to be little reason why random permutations and combinations should not eventually lead to the production of some primitive kind of organized system possessing potentialities for self-replication, survival, and eventual evolution.

No doubt the monkeys would have produced some other interesting documents in the course of their efforts, documents corresponding to other kinds of organized systems; systems that failed to stand up to alterations in a constantly changing external environment, and which subsequently died out and left no trace.

Possibly these random processes were less random than the aimless and totally undirected performance of the monkeys because, as Calvin (1957) has pointed out, autocatalysis and catalysis by simple inorganic compounds or heavy metals may have played a large part, not only in the synthesis of new and more elaborate molecules, but also as a selective and therefore directive agent. There may well have

been still other chemical counterparts of natural selection, so that the production of large-molecular compounds and their organization into a primitive form of living stuff was, in some degree, perhaps a rather less improbable performance than the writing of Shakespeare's sonnets.

Table 1. Products of irradiation of a mixture of H_2 , CH_4 , and NH_3 (spark discharge)

Compound	Yield, moles $\cdot 10^5$		
	Spark. run 1	Silent. run 3	N_2 run, run 6
Glycine	63.(2.1) *	80.(.46) *	14.2 (.48) *
Alanine	34.	9.	1.0
Sarcosine	5.	86.	1.5
β -Alanine	15.	4.	7.0
α -Aminobutyric acid	5.	1.	—
N-Methylalanine	1.	12.5	—
Acids:			
Aspartic	0.4	0.2	0.3
Glutamic	0.6	0.5	0.5
Iminodiacetic	5.5	0.3	3.9
Imino-acetic-propionic	1.5	—	—
Formic	233.	149.	135.
Acetic	15.2	135.	41.
Propionic	12.6	19.	22.
Glycolic	56.	28.	32.
Lactic	31.	4.3	1.5
α -Hydroxybutyric	5.	1.	—
Succinic	3.8	—	22.
Urea	2.	—	2.
Methylurea	1.5	—	0.5
Sum of yields of compounds listed. %	15	3	8

* Per cent yield of glycine based on carbon placed in the apparatus as methane.

SOURCE: Miller (1957).

Supposing, then, that the period of chemical evolution gave rise to some organized system possessing the potentialities for what may be called "life," as we use the word in its everyday connotation, and organic evolution could begin. To our knowledge and understanding of this latter process biochemistry can contribute much; hence the title of this essay, "Biochemistry and Evolution"—a challenging title for any biochemist and above all for one who professes an interest in its comparative aspects.

Nearly 25 years ago the present author wrote a little book (Bald-

win, 1937) whose only claim to lasting distinction was probably the foreword contributed to it by Sir Frederick Gowland Hopkins. In that foreword he gave the following text: "I venture to think that productive thought in biochemistry . . . calls for the widest possible survey of life's manifestations. One of its ultimate tasks is to decide on what, from the chemical standpoint, is essential for these manifestations as distinct from what is secondary and specific." Comparative biochemistry is indeed concerned as much with resemblances as it is with differences, and the further biochemical research progresses, the more does it appear that living organisms closely resemble each other at the molecular level, no matter what their structure, mode of life, or environment. It is as though there is some fundamental structural and metabolic ground plan to which all kinds of living things conform. If we could but strip away everything that is secondary and specific and leave this ground plan revealed and naked, we should very probably have a fair idea of the structural and metabolic make-up of the earliest forms of anything that would be called "alive" by present-day standards. Further back than this, perhaps, we could not go; but that is a problem for the future rather than for this time and place. In writing here about evolutionary diversity the existence of a fundamental ground plan is taken for granted. It is assumed, moreover, that the features of this fundamental plan were laid down very early in the evolution of modern living forms and that they correspond to a common starting point for evolutionary divergence and differentiation.

J. B. S. Haldane (1937) once wrote that: "Our final theory of evolution will see it largely as a biochemical process." Now if, as seems generally to be believed, evolution has proceeded by a long series of individual mutations, and if, as there is every reason to think, individual mutations, or at any rate their consequences, are open to investigation along biochemical lines, we are also bound to believe, again with Haldane, that "future interpretations of genetics will be largely expressed in biochemical terms." Biochemical genetics is now a large and important branch of biochemical inquiry. Moreover, it is well documented (see, for example, Haldane, 1954).

If we accept the notion of a fundamental ground plan—and all modern biochemistry seems to point in this direction—and if we suppose that this ground plan was laid down early in the course of biological time, it would seem probable that mutant forms could only be viable if the underlying mutations were consistent with the maintenance of the primary, basic, and fundamental pattern. Moreover, when at a later stage a number of consistent and specific adaptational super-

structures had been built up on the fundamental plan, further mutations would only be viable if they were consistent with the fundamental plan itself and with the maintenance of everything that had already been superadded to the system. Something of this kind must surely be at the bottom of the fact that the vast majority of random mutations are lethal.

When we come to study spontaneous or artificially induced mutations in modern organisms, ranging from *Neurospora* to man himself, it seems an almost invariable rule that each mutation that turns up results in either the loss or some unfavorable modification of at least one enzyme.

There is an abundance of evidence for this; inherited metabolic disorders such as phenylketonuria, alkaptonuria, hemophilia, and the like have long been known (see Garrod, 1909). Another well-known example on the evolutionary scale is the serial loss of uricolytic enzymes among vertebrate animals (Baldwin, 1949; Florkin, 1949). Fishes in general possess urico-oxidase, allantoinase, and allantoinase, but in some groups and families allantoinase has already disappeared. At the other end of the scale, most mammals possess urico-oxidase but lack allantoinase and allantoinase. Finally, among the primates, even urico-oxidase has been lost.

These last-mentioned enzymes are concerned only with end products of metabolism. If, however, the lost or altered enzyme is one that plays an integral part in intermediary metabolism and normally catalyzes the conversion of *A* into *B*, *A* will tend to accumulate in the cells and the organism thenceforward will be able to survive only in media in which *B* is present and available. A familiar example is found in a mutant of *Escherichia coli*, which lacks the enzyme that converts 4-amino-5-iminazole-carboxamide-ribosyl-5'-phosphate into inosinic acid (Gots, 1953).

This is an important step in the biosynthesis of purines and their nucleotides and was our first clue to the nature of the synthetic mechanism.

A mutation of this kind is likely to result in the restriction of the mutant to certain very circumscribed habitats, habitats in which the necessary metabolite can be found ready made. Further stepwise mutations may result in the loss of ability to produce yet other metabolites and the habitats available become even more restricted. This can lead in the end to the grossest kind of parasitism, especially among bacteria, and it is very possible that symbiosis and frank parasitism may be the end results of similar, serial loss mutations in animals as well.

Among bacteria at any rate, loss of synthetic ability is often a stepwise process and can lead to the most exacting of nutritional requirements; many amino acids and a larger or smaller number of accessory growth factors are often required, and these can sometimes only be found in the tissues or tissue exudates of a specific living host. Such is the case in *Streptococcus haemolyticus* for example, one of the most exacting organisms known. A case that has been studied in much detail is that of coenzyme A synthesis by bacteria. Requirements for the total synthesis of this substance (see Table 2 and Fig. 1) have been established (Snell, 1956).

Table 2. *Biosynthesis of coenzyme A by bacteria*

Organism	Requirement
<i>Corynebacterium diphtheriae</i>	β -alanine
<i>Lactobacillus bulgaricus</i>	pantetheine
<i>Lactobacillus casei</i>	pantothenic acid
<i>Acetobacter suboxydans</i>	pantoic acid
<i>Treponema pallidum</i>	pantetheine-4'-phosphate

SOURCE: Snell (1956).

Yet advantageous mutations, which are probably much less frequent, have clearly taken place over and over again; indeed they have formed the ultimate basis of all adaptation and consequent evolutionary progress. It would seem, however, that among mutations leading to gain, as judged by advantages of survival value, some seem to be more probable than others.

One outstanding example of a gain mutation that seems to be of a rather probable character is the invention of hemoglobin, by far the commonest of the four known respiratory pigments (Lemberg and Legge, 1949). We can say "probable" because, unlike chlorocruorin, hemerythrin, and hemocyanin, hemoglobin has turned up over and over again without the slightest reverence for taxonomy; it crops up

here and there all over the animal kingdom, in many members of many invertebrate groups as well as in vertebrates generally. Keilin suggested many years ago that the formation of hemoglobin is due to modifications (presumably mutational) in the enzymatic machinery ordinarily involved in the production of the cytochromes. Presumably *b* or *c* are the components concerned, since their hemes are very closely related to that of hemoglobin itself. This seems to be confirmed by the comparable fact that chlorocruorin, which occurs only in a small group of so-called chlorhemid worms, carries a heme very similar to and possibly identical with that of cytochrome *a*.

According to all available evidence, different hemoglobins possess a common heme, and the species-specific differences existing between

FIG. 1. Requirements for coenzyme A synthesis by bacteria (see also Table 2).

them reside entirely in the globin part of the molecule. But this may be too narrow a view, for there may well be mutant forms of globin as indeed there are among humans. About two dozen or so human variants are now known (Prankerd, 1961) and what seem to be quite trivial differences from the purely chemical standpoint may have important physiological results: sickle-cell hemoglobin differs from normal in only one of the 300 amino acid units in each half molecule (Hunt and Ingram, 1959). In sickle-cell hemoglobin, valine replaces glutamic acid, while in hemoglobin C the place of glutamic acid is taken by lysine. These and other differences are summarized in Table 3. Small though the chemical differences between the normal and sickle-cell hemoglobins may seem on the surface, individuals carrying two sickle genes commonly die, often in infancy, from the consequent anemia. Here we have what was originally an advanta-

geous mutation operating through a further mutation to a disadvantageous outcome.

In what has just been said it would appear that the invention of hemoglobin is certainly a gain event on the whole, for with its acquisition many animals were enabled to penetrate into regions which they could not otherwise have occupied, and were assured of a more abundant supply of oxygen and a consequently greater potential activity, even in their normal habitats. Subsequent mutations in the human race—and there surely must be others elsewhere among animals—seem to have had only indifferent consequences in some cases (hemoglobin C), while others have been well on the way toward being lethal (hemoglobin S).

Table 3. Chemical differences between normal and abnormal hemoglobins

Hemoglobin	Structural characteristics	Chain in which structures occur
A	Val.his.leu.thr.pro. <i>glu</i> .glu.lys.	N-terminal sequence of β chain
S	Val.his.leu.thr.pro. <i>val</i> .glu.lys.	N-terminal sequence of β chain
C	Val.his.leu.thr.pro. <i>lys</i> .glu.lys.	N-terminal sequence of β chain
G $_{\beta}$	Val.his.leu.thr.pro.glu.lys.lys.	N-terminal sequence of β chain
E	Gly.gly.lys.ala.leu.gly. (Lys. replaces glu. of Hb-A)	Peptide in β chain
I	Try.gly. <i>asp</i> .val.gly. (Asp. replaces lys. of Hb-A)	Peptide in α chain

SOURCE: Prankerd (1961).

Any discussion of the species-specificity of proteins leads one into deep waters, for relatively little is known even today about the detailed chemical structure of these materials. However, insulin is a protein of fairly small molecular size and of which the structure is precisely known through the brilliant work of Sanger and his co-workers (Brown, Sanger, and Kitai, 1955; Harris, Sanger, and Naughton, 1956). They have shown that insulin is species-specific and have discovered the precise nature of the differences between a number of different insulins. Some of their results are summarized in Table 4. All this demonstrates that it is possible to have a number of substances which, though chemically different, nevertheless possess identical physiological or pharmacological activity.

Relatively little is known about the comparative side of enzymology, though important advances have been achieved in elucidating the structure of ribonuclease (Anfinsen & White, 1961). An interesting point of a somewhat different kind is that, as is now well known,

specialized substances produced by the tubercle organisms and the blood group polysaccharides, have much in common. Cellulose occurs widely in plants and occasionally in animals, especially in the tunicates, but we know little about its fine structure. Chitin occurs in many invertebrates, but even its structure is still not precisely known. Galactogen, a major constituent of the eggs and albumin glands of certain snails, is uncommon; indeed it is uncommon if only in that about one in every seven of the constituent galactose units consists of the L- instead of the usual D-isomeride (Bell and Baldwin, 1941). Glycogen, the commonest of animal polysaccharides, somewhat resembles the amylopectin component of the plant starches yet differs somewhat according to its source, but the differences are not chemically very remarkable. Glycogens from various sources give different colors with iodine, ranging from zero to the well-known portwine coloration, probably reflecting differences in chain length and degree of branching. However, it is difficult to know what chain length means in molecules which are continuously changing as new units are constantly added or removed. These are essentially dynamic, ever-changing, and thoroughly recalcitrant molecules.

So far we have mainly considered some variations on some one or other of a few general themes. But it happens from time to time that the comparative biochemist comes across substances that appear at the time of their discovery to be uniquely confined to particular groups or species. Galactogen, for example, is known to be present in the eggs and albumin glands of *Helix pomatia*, *H. aspersa*, and *Limnaea* sp. and might have been thought to be a characteristic feature of gastropod mollusks, but it could not be detected in the eggs of *Aplysia punctata*, a marine gastropod, nor in those of the cephalopod, *Sepia officinalis* (Baldwin, unpublished). Maybe it will turn out to be a feature of pulmonates as opposed to operculates, but too few species of either have as yet been studied.

The trouble here is, as has often been pointed out, that comparative biochemistry has not in the past been comparative enough, and, until it becomes so, it behooves us to be wary of associating this or that substance too closely with some one or other particular group of animals. A recent example of this is the case of homarine. First discovered in lobster muscle by Hoppe-Seyler (1933), it has now been found to be a very widely distributed constituent of invertebrate materials and may, for all we know to the contrary, occur in vertebrate tissues as well. It had indeed already been found in two other phyla, viz., in *Arbacia* sp. by Holtz, Kutscher, and Thielmann (1924)

and in *Arca noae* by Kutscher and Ackermann (1933b), but was wrongly identified as the isomeric trigonelline at the time.

Distribution studies have in the past been one of the more popular kinds of comparative biochemistry, but they have been too few and too far between. Classical examples are those of Kutscher and Ackermann (1933a, 1936). As an illustrative example here we may review work done on the distribution of the phosphagens, with which the author has some personal acquaintance.

Work in this field began with the discovery by Eggleton and Eggleton (1928) that although creatine phosphate occurs widely in vertebrate muscles, it is absent from those of invertebrates. Meyerhof (1928) found that, among invertebrates, arginine phosphate replaces the creatine compound. This pioneer work was soon followed up on a much larger number of species by Needham, Needham, Baldwin, and Yudkin (1932) and gave results which led them to believe that they had important evidence concerning the origin of vertebrates. All the invertebrates examined contained what appeared to be arginine phosphate, with notable exceptions in certain echinoderms (later confirmed by Baldwin and Needham, 1937). In certain echinoids both phosphagens were found to coexist. The same results were found in a hemichordate (*Balanoglossus salmoneus*) and it seemed that the results fitted in with Bateson's theory of the origin of vertebrates, i.e., that they arose through an echinoderm-hemichordate route.

In all this there was a certain element of fortune. If this work had been done not at Roscoff but at Woods Hole, these conclusions could not have been reached because the echinoid and hemichordate studied later at Woods Hole by Baldwin and Yudkin (1950) proved to contain only one phosphagen each—arginine phosphate in *Arbacia pustulosa* and the creatine compound in *Saccoglossus kowalevskii*.

The work of Baldwin and Yudkin (1950) was mainly devoted to a study of the phosphagen of marine annelids and was prompted by the somewhat atypical behavior of the presumptive arginine phosphate of certain annelids studied by Needham, Needham, Baldwin, and Yudkin (1932) and by the work of Arnold and Luck (1933), according to whom several species of marine annelids contain no arginine whatsoever.

The presence of new phosphagens among the annelids was demonstrated and it was shown that none of them is identical with arginine phosphate. One of the new substances was tentatively identified as creatine phosphate, but its positive identification only came later on with the work of Roche and his colleagues (Roche, Thoai, Garcia, and

Robin, 1952; Thoai, Roche, Robin, and Thiem, 1953a, b, c; Thoai and Robin, 1954a, b; see also Ennor and Morrison, 1958). In addition to creatine phosphate, moreover, Roche's group identified two further new phosphagens, one based on glycoeyamine and another on taurocyamine. Yet another new base, lombricin, was subsequently discovered in earthworms and it seems probable that yet another is

FIG. 2. Relationships of guanidine bases of animal phosphagens: (1) Transamidation of glycine; (2) transmethylation of glycocyamine; (3) transamidation of taurine.

present in leeches. The formulas of these bases are shown in Fig. 2. Evidently, then, the annelids are a particularly versatile group.

The discovery of creatine phosphate in worms upset earlier ideas about vertebrate evolution, even though something could still be salvaged from the wreck. But perhaps the most interesting aspect of Roche's series of studies was that glycoeyamine, known for many years as a metabolic intermediate between arginine and creatine, occurs in certain annelids. There seems here to be an evolutionary sequence;

arginine in many marine invertebrates, leading to glycoeyamine in some annelids and thence on to creatine in other annelid species. Perhaps the time has come to reexamine the echinoderms and protochordates in case this intermediate stage has been preserved for us there too.

The discovery of tauroeyamine was interesting inasmuch as transamidation is a fairly rare event. Longest known is the transfer of the amidine group from arginine to glycine, yielding glycoeyamine. It seems likely that tauroeyamine arises by a similar transfer from arginine to taurine, for taurine is widely distributed in the animal kingdom and occurs in remarkably large quantities in certain mollusks, e.g., *Abalone* (Schmidt and Watson, 1918) and, more interesting, in at any rate some annelid worms (Kurtz and Luck, 1935). So, discoveries of this comparative kind can sometimes add something to the vast and ever-growing body of general biochemistry. Another example that comes to mind is the discovery of octopine in cephalopod mollusks (Morizawa, 1927) and in some lamellibranchs and gastropods (Irvin and Wilson, 1939). This substance appeared to give promise of restriction to certain fairly sharply defined groups of animals, but later turned out to be very closely allied to the intermediate compounds formed in the probably universal process of transamination.

As time goes on, then, more and more of the supposedly secondary and specific features of particular groups or species tend to become incorporated into some more general plan, and examples could be multiplied considerably. It may well be, indeed, that as comparative biochemistry becomes more comparative, more and more of the odd, peculiar, and apparently restricted phenomena known today will prove to be widely and in some instances perhaps even universally distributed, perhaps even as parts of the fundamental metabolic ground plan. Even the production of large amounts of steam-volatile fatty acids by parasitic worms (*Ascaris lumbricoides*) (Moyle and Baldwin, 1952), long thought to be unique, at least among animals, finds a close parallel in the production of similar acids by the action of ruminant symbionts upon cellulose (McAnally and Phillipson, 1944), but this does not seem by any means a widespread or fundamental phenomenon.

At this stage two general comments may be made: first, that we are inclined to study too few species and are tempted to draw too far-reaching conclusions from too little evidence. For this we have to blame sometimes ourselves, sometimes shortage of funds, and sometimes the weather, especially at marine stations. Secondly, and this

must be emphasized, no matter what animal or animals we choose to study, there is always a possibility or even a likelihood that we shall stumble across some hitherto unknown phenomenon of general or even fundamental importance.

However, not the whole of comparative biochemistry is in quite this unsatisfactory state. Often facts are scrappy and unsatisfactory because too few species have been studied, but sometimes fairly thorough, systematic studies have been carried out. For example, starting with the work of Städele and Frélich in 1858, evidence has accumulated that the elasmobranch fishes, but not the teleosts or any other group, elaborate urea as their principal nitrogenous excretory end product. This is true also of the holocephalians, which are usually regarded as an aberrant elasmobranch group, so that biochemistry and taxonomy run hand in hand in this case. The function of the urea, long debated, seems now to be certain. It plays a most important role in osmotic regulation in these fishes (Smith, 1936). Here a striking biochemical peculiarity can be correlated with a specific function in a particular kind of environment.

On a smaller scale, urea formation features also in the dipnoan fishes (Smith, 1930). But when the swamps they inhabit dry up in the dry season, the formation of ammonia—important while they have water to live in—is wholly suppressed in favor of urea production. In this case urea formation evidently represents a specific device for the detoxication of ammonia, and without some such mechanism the lungfishes would have little or no chance of survival through their periods of estivation.

A substantially similar case is found in the horned toad, *Xenopus laevis*. This animal, by all taxonomic standards, is a perfectly good amphibian, but it is one of a number of amphibians that have made a secondary and permanent return to the water. In the ordinary way the bulk of its waste nitrogen is excreted as ammonia, but some urea is nevertheless formed as evidence, presumably, of a truly amphibious ancestry (Balinsky and Baldwin, 1961). It is said that *Xenopus* can survive periods of estivation during the dry season, and a number of experiments in which specimens were kept under mildly damp instead of wholly aquatic conditions have proved that this animal, like the lungfish, switches over to large-scale urea production under these conditions (Balinsky, Cragg, and Baldwin, 1961). Like the lungfish emerging from estivation, *Xenopus* excretes large amounts of accumulated urea when returned to water after being kept out of water for some days (Table 5). Here it seems that these are comparative data of real significance, even though the number of species involved

is small. But of course it would be well to have the results either confirmed or contradicted by investigations on other species with similar habits and habitats.

Table 5. Nitrogen excretion of Xenopus in water after 19 days in damp (not wet) conditions.

Days in water	Per cent of total N as	
	Ammonia	Urea
3	17	76
5	22	66
10	56	35
17	82	8

SOURCE: Balinsky, Cragg, and Baldwin (1961).

The results so far mentioned fit in with much other information so precisely that we may perhaps be excused for seeming to jump to conclusions over only one or two species. The basic fact behind all this is that ammonia, familiar enough as a household article, is nevertheless extremely toxic, as was conclusively shown by Sumner many years ago. No animal, probably, could survive for long on dry land without having some means of converting ammonia into a more innocuous material such as urea. Indeed, *Xenopus* apart, all the anuran amphibians so far examined excrete about 80 per cent or more of their waste nitrogen as urea (see Table 6), and the habit

Table 6. Nitrogen excretion of amphibians: Anura

Amphibian	Habitat	Per cent of total N as	
		Ammonia	Urea
<i>Xenopus laevis</i>	FW	65.5	21.4
<i>Rana esculenta</i>	FW/T	7.8	74.8
<i>R. temporaria</i>	FW/T	7.0	78.7
<i>Hyla arborea</i>	T	4.3	83.8
<i>Bufo calamita</i>	T	5.3	86.9

SOURCE: Cragg, Balinsky, and Baldwin (1961).

of urea production has been inherited by the mammals, including even those primitive forms that still lay eggs (Cragg, Balinsky, and Baldwin, 1961). However, the majority of the reptiles, together with

the birds, produce not urea but uric acid as an alternative way of detoxicating ammonia. But from some reptilian groups interesting facts of probable evolutionary significance have emerged (Cragg, Balinsky, and Baldwin, 1961). We know little about the crocodiles and alligators except that ordinarily they live in water and excrete nitrogen mainly in the form of ammonia (Table 7). One would like to know what happens during their overland treks in search of new haunts when the water dries up in the dry season. It should be possible to do "desiccation" experiments similar to those tried on *Xenopus*.

Table 7. Nitrogen excretion in crocodilian reptiles

Reptile	Per cent of total N as		
	Ammonia	Urea	Uric Acid
<i>Alligator mississippiensis</i>	75	6	13
<i>Caiman crocodilus</i>	58	6	15
<i>Crocodylus niloticus</i>	66	5	21

SOURCE: Cragg, Balinsky, and Baldwin (1961).

It is among the tortoises and turtles—perhaps the most primitive reptilian group—that the most interesting results have emerged, though the number of species used was only eight. Three groups have been studied by Moyle (1949), one of which lives an amphibious life and never goes far from the water (Table 8). In these the major end product is urea, inherited presumably from ancestral amphibians. A second group comprises two species that, like *Xenopus*, have made a secondary and apparently permanent return to fresh water. Here

Table 8. Nitrogen excretion in chelonian reptiles

Reptile	Habitat	Per cent of total N as		
		Ammonia	Urea	Uric acid
<i>Kinosternon subrubrum</i>	Aquatic	24.0	22.9	0.7
<i>Pelusios derbianus</i>	Aquatic	18.5	24.4	4.5
<i>Emys orbicularis</i>	Amphibious	14.4	47.1	2.5
<i>Kinixys erosa</i>	Land near water	6.1	61.0	4.2
<i>K. youngii</i>	Land near water	6.0	44.0	5.5
<i>Testudo denticulata</i>	Swamps	6.0	29.1	6.1
<i>T. graeca</i>	Dry land	4.1	22.3	51.9
<i>T. elegans</i>	Very dry land	6.2	8.5	56.1

SOURCE: Moyle (1949).

ammonia sometimes predominates among the nitrogenous excreta, with urea still playing a substantial part. This seems to indicate a reversion toward wholly aquatic habits, but the persistence of urea production gives present evidence of a one-time truly amphibious or terrestrial past.

The third group comprises three species within a single genus, *Testudo*. In the swamp-dwelling *T. denticulata* the picture is similar to that found in the other amphibious species. In *T. graeca*, however, a substantial excretion of uric acid makes its appearance and finally, in the desert-living *T. elegans*, urea no longer appears in more than trivial amounts and uric acid is responsible for practically all the nitrogen accounted for.

In fairness to Moyle, it should be pointed out here that there were a number of analytical difficulties in this work largely because many of these animals have a habit of excreting a good deal of only partially digested food that is not removed by the usual deproteinization procedures, but even so the results do appear to be pretty clear-cut. It certainly seems that in this group of reptiles we have a series of cases which sum up the biochemical steps which were probably entailed in the evolution of the modern snakes, lizards, and birds. Starting from an amphibious stock of urea producers, they took to progressively drier and drier surroundings, making use at first of the mechanisms they already possessed by reason of their ancestry. Eventually the advantages of uric acid became apparent—its extreme insolubility, the fact that it is relatively innocuous and that little water is required for its excretion—and gradually, probably step by step, there came the switchover from the formation of urea to that of uric acid. This, as Joseph Needham (1931) has argued, must also have tied up with the invention of impermeable eggshells and consequential changes in the conditions of embryonic life.

Before leaving these chelonian reptiles, it must be mentioned that there are reports in the literature to the effect that the wholly marine turtles excrete only ammonia and no urea or uric acid worth mentioning. Does this mean, perhaps, that they never had an amphibious phase and so had no use for urea, or that they returned to the water long, long ago and have now altogether forgotten how to make it? Evidently, there is still plenty of room for new work in this field.

From these studies on amphibia and reptiles some recent research has turned toward another problem of no small interest, viz., the nitrogen excretion of fishes. The positive results obtained by the author so far have been published (Baldwin, 1958, 1960), like much of the work already mentioned, but only a few of the negative ones

have so far seen the light of day. However, there are some interesting possibilities from the viewpoint of evolution.

The material used comprised a number of species of elasmobranchs which, as is well known, are large-scale producers of urea, and the primary problem is whether or not they make urea by the same route as that followed in mammals and amphibians. It is now widely though not quite universally believed that urea formation follows the pathways shown in Fig. 3. Carbon dioxide and ammonia in the presence of ATP and acetylglutamate give rise in some way that is still very obscure to carbamyl phosphate. Ornithine is transcarbamylated by carbamyl phosphate to give citrulline, which reacts then with aspartate, giving argininosuccinate. This last product is split into arginine and

Mechanisms of ureogenesis
(as generally accepted)

FIG. 3. Mechanisms of ureogenesis (as generally accepted).

fumarate and the arginine is hydrolyzed by arginase to give urea and regenerate ornithine, which can then be used over again. Most of the enzymes involved have been rather recently more or less purified and studied in considerable detail (see Brown and Cohen, 1960, for references). Their results were published a few years after our own work in this field began.

A number of interesting problems came to mind at the start of this work. In the first place, while arginase is present in the livers of all urea-producing animals, it is normally confined to the liver. In elasmobranchs, however, it is present in every organ in the body; it would be interesting to know whether, in view of its very great osmotic importance, urea might be made not only in the liver as is usually the case but in other organs as well. The whole business presents special problems because one is faced with the determination of small amounts of urea synthesis in the presence of some 2 to 2.5 per cent of

preformed urea, but it has been possible to overcome this, first by the use of well-washed tissue preparations and later (Joseph, unpublished) by the use of ion-exchange resins. A few experiments have been done on urea synthesis in organs other than the liver but have led nowhere so far.

Hitherto it had been usual to think of urea production as a means of detoxicating ammonia, and as something that originated in animals at somewhere about the amphibian level of evolution. This attitude could accommodate not only the modern amphibia but also the lungfishes which, apart from the elasmobranchs, are the only group of fishes in which urea formation is a serious industry. Why, then, and at what stage in their history, did these wholly aquatic elasmobranchs develop the machinery for making urea? It could, of course, be a matter of evolutionary convergence. Alternatively it could be, perhaps, as Romer and others have suggested, that the earliest ancestral fishes lived under conditions similar to those the lungfishes experience today, and that they all used urea as a means of survival. Later on, it could be argued, urea formation became redundant and disappeared from the majority of fishes. The elasmobranchs, it might be thought, had discovered its osmotic advantages in the meantime and retained the synthetic apparatus. If this were true, it might perhaps be anticipated that some vestiges of the one-time synthetic apparatus might be discovered in other kinds of fishes today, and new work is going on to look into this. In the meantime it is known that arginase is present in the livers of a considerable number of teleostean fishes, and that appreciable quantities of urea are often to be found in their excreta and even in their tissues. So this looks like a helpful line to follow up. Finally, there are still a few fishes that, morphologically speaking, have remained practically unchanged since the Jurassic, for example, *Amia* and *Lepidosteus*. *Polypterus* is still with us but has changed little since the Carboniferous or thereabouts. There are, too, the coelocanth. All these and many more would be well worth studying as potential sites of vestigial urea-producing enzymes. So far the only information we have comes from Brown and Cohen (1960), who could find neither carbamylphosphate synthetase nor ornithine transcarbamylase in *Amia* or *Lepidosteus*.

In the earlier part of our own work, we were able without difficulty to show that laboratory-synthesized carbamyl phosphate reacts with ornithine and gives citrulline, and that, in the presence of citrulline, aspartate, and ATP, urea is formed. If any of the reactants was omitted, urea synthesis stopped (Tables 9 to 11). All these results have since been abundantly confirmed. So far there is obviously a close

Table 9. Formation of citrulline from carbamyl phosphate and ornithine

Species	μ mole citrulline/g powder
<i>R. productus</i> (1)	187
<i>R. productus</i> (2)	166
<i>H. californicus</i>	223
<i>M. californicus</i>	205
<i>G. marmorata</i>	166

Reaction mixture: 0.1 ml 0.1 M ornithine; 0.2 ml 0.2 M K-phosphate. buffer (pH 7.4); 2.5 ml 0.02 M dilithium carbamyl phosphate; 0.8 ml water; 0.5 ml extract of Type 1 powder.

Incubation: 30 min at 38°C. Readings at 490 m μ .

SOURCE: Baldwin (1958, 1960).

Table 10. Formation of urea from citrulline and aspartate

Species	μ mole urea/g powder		
	Preformed	After incubation	Synthesized
<i>R. productus</i> (1)	{126	158	27
	{126	178	52
<i>R. productus</i> (2)	{ 95	118	23
	{ 95	121	26
<i>M. californicus</i>	{ 74	108	34
	{ 75	103	28
<i>G. marmorata</i>	{123	145	22
	{125	159	34
<i>U. halleri</i>	{126	158	32
	{127	177	50

Reaction mixture: 0.15 ml 1.0 M L-citrulline; 0.15 ml 1.0 M aspartic acid (K salt); 0.1 ml 1.0 M K-buffer (pH 7.4); 0.1 ml 0.066 M MgSO₄; 0.1 ml 0.025 M ATP (K salt); 0.2 ml 0.1 M K-phosphoglycerate; 0.1 ml arginase (ca. 150 units/ml); 0.1 ml glycolysing enzyme preparation. Incubation: 30 min at 38°C. Readings at 540 m μ .

SOURCE: Baldwin (1958, 1960).

parallelism between urea synthesis in these fishes, in amphibia, in chelonian reptiles, and in mammals. We have since done more detailed work on the stages between citrulline + aspartate and urea, and there is no doubt that the reactions take place in the usual way. But where we have completely failed in more than two years of concentrated work is in the biosynthetic production of carbamyl phosphate by elasmobranchs. We have used more delicate and refined methods for its detection, we have tried all sorts of conditions of temperature and pH, we have tried putting in everything from sodium chloride to the proverbial kitchen sink, but still we have not been able to get a synthesis of carbamyl phosphate. Parallel experiments with rat liver,

*Table 11. Formation of urea from citrulline and aspartate **

(*M. californicus*)

Reactant omitted	μ mole urea/g powder	μ mole urea formed/g powder
All	75	(Preformed)
None	100	25
Citrulline	73	-2
Aspartate	78	+3
ATP	83	+8
Phosphoglycerate	75	0
Arginase	100	25

* 1 ml enzyme and 1 ml reaction mixture as for Table 10. Incubation 30 min at 30°C.

SOURCE: Baldwin (1958, 1960).

frog liver, and slaughterhouse material produce carbamyl phosphate readily enough, but the elasmobranchs systematically refuse to do so. Brown and Cohen (1960) have since reported similar failures.

We shall not try to put any interpretation on these results except to say that there seem to be two possibilities. One is that we may be dealing with a case of evolutionary convergence. The other is that the scheme for urea synthesis on which we are working, and which is widely accepted, may be wrong. Bach, for example, has long maintained that there are more ways than one in which urea may be synthesized and it is even possible that, working on elasmobranch fishes, we may yet find out how urea is really synthesized in mammals.

The author was brought up to believe that the job of biochemistry is to solve biological problems by chemical methods, and he felt in

preparing this essay that the task was, not so much to present new facts as to try to get a certain amount of perspective into the contributions that biochemistry, especially on the comparative side, can make toward the solution of evolutionary problems. Not unnaturally, most space has been spent on matters with which he has had a good deal of personal experience. An attempt has been made to point out some of the shortcomings and some of the dreams, trials, disappointments, and tribulations of a comparative biochemist. But in spite of all these it is still possible to believe that if the recent upward swing of interest in comparative biochemistry can be maintained, it will in the end prove its ability to make valuable contributions to that most biological of all biological theories, the theory of evolution.

References

- Anfinsen, C. B., and F. H. White (1961), The ribonucleases: occurrence, structure and properties in *The Enzymes*, Academic Press, Inc., New York, vol. 5, pp. 95-122.
- Arnold, A., and J. M. Luck (1933), Studies on arginine. III. The arginine content of vertebrate and invertebrate muscle, *J. Biol. Chem.*, **99**:677-691.
- Baldwin, E. (1937), *An Introduction to Comparative Biochemistry*, 1st ed., Cambridge University Press, London and New York.
- (1949), *An Introduction to Comparative Biochemistry*, 3d ed., Cambridge University Press, London and New York, p. 65.
- (1958), Ureogenesis in elasmobranchs, *Nature* (London), **181**:1591-1592.
- (1960), Ureogenesis in elasmobranch fishes. *Comp. Biochem. Physiol.*, **1**:24-37.
- , and D. M. Needham (1937), A contribution to the comparative biochemistry of muscular and electrical tissues, *Proc. Roy. Soc. (London)*, **B**, **122**:197-219.
- , and W. H. Yudkin (1950), The annelid phosphagen: with a note on phosphagen in *Echinodermata and Protochordata*, *Proc. Roy. Soc. (London)*, **B**, **136**:614-631.
- Balinsky, J. B., and E. Baldwin (1961), The mode of excretion of ammonia and urea in *Xenopus laevis*, *J. Exp. Biol.*, **38**:695-705.
- , M. Cragg, and E. Baldwin (1961), The adaptation of waste nitrogen excretion to desiccation, *Comp. Biochem. Physiol.*, **3**:236-244.
- Bell, D. J., and E. Baldwin (1941), The chemistry of galactogen from *Helix pomatia* as a component of a polysaccharide of animal origin. *J. Chem. Soc.*, **1941**:125-132.
- Brown, G. W., and P. P. Cohen (1960), Comparative biochemistry of urea synthesis. 3. Activities of urea-cycle enzymes in various higher and lower vertebrates, *Biochem. J.*, **75**:82-91.
- Brown, H., F. Sanger, and R. Kitai (1955), The structure of pig and sheep insulin, *Biochem. J.*, **60**:556-565.
- Calvin, M. (1957), Chemical evolution and the origin of life, in *The Origin of Life on the Earth*, Rep. Internat. Symp., Publ. House, Acad. Sci. USSR, pp. 338-350.
- Cragg, M., J. B. Balinsky, and E. Baldwin (1961), The comparative study of nitrogen excretion in some amphibia and reptiles, *Comp. Biochem. Physiol.*, **3**:227-235.

- Eggleton, P., and G. P. Eggleton (1928). Further observations on phosphagen, *J. Physiol.*, **65**:15-24.
- Ennor, A. H., and J. F. Morrison (1958). Biochemistry of the phosphagens and related guanidines, *Physiol. Rev.*, **38**:631-674.
- Florkin, M. (1949). *Biochemical Evolution*, Academic Press Inc., New York, pp. 45-47.
- Garrod, A. E. (1909). *Inborn Errors of Metabolism*, Oxford University Press, London and New York.
- Gots, J. S. (1953). Occurrence of 4-amino-5-iminazole carboxamide as a pentose derivative, *Nature* (London), **172**:256-257.
- Haldane, J. B. S. (1937). The biochemistry of the individual, in *Perspectives in Biochemistry*, Cambridge University Press, London and New York, pp. 1-10.
- (1954). *The Biochemistry of Genetics*, George Allen & Unwin, Ltd., London.
- Harris, J. I., F. Sanger, and M. A. Naughton (1956). Species-differences in insulin, *Arch. Biochem. Biophys.*, **65**:427-438.
- Holtz, F., F. Kutscher, and F. Thielmann (1924). Über das Vorkommen des Pflanzenalkaloids Trigonellin in der Tierwelt, *Z. Biol.*, **81**:57-60.
- Hoppe-Seyler, F. A. (1933). Über das homarin, eine bisher unbekannte tierische Base, *Z. physiol. Chem.*, **222**:105-115.
- Hunt, J. A., and V. M. Ingram (1959). A terminal peptide sequence of human haemoglobin, *Nature* (London), **184**:640-641.
- Irvin, J. L., and D. W. Wilson (1939). Studies on octopine. III. The precursor of octopine in scallop muscle, *J. Biol. Chem.*, **127**:575-579.
- Kurtz, A. C., and J. M. Luck (1935). Studies on annelid muscle. I. Taurine in *Audolonia spirabranchus* (Moore), *J. Biol. Chem.*, **111**:577-584.
- Kutscher, F., and D. Ackermann (1933a). Comparative biochemistry of the vertebrates and invertebrates, *Ann. Rev. Biochem.*, **2**:355-376.
- (1933b). Über das Vorkommen von Betainen in der Archenmuschel, *Z. physiol. Chem.*, **221**:33-39.
- (1936). Comparative biochemistry of the vertebrates and invertebrates, *Ann. Rev. Biochem.*, **5**:453-462.
- Lemberg, R., and J. W. Legge (1949). *Haematin Compounds and Bile Pigments: Their Constitution, Metabolism and Function*, Interscience Publishers, Inc., New York, pp. 307, 309.
- McAnally, R. A., and A. T. Phillipson (1944). Digestion in the ruminant, *Biol. Rev.*, **19**:41-54.
- Meyerhof, O. (1928). Ueber die Verbreitung der Argininphosphorsäure, *Arch. Sci. Biol.* (Napoli), **12**:536-548.
- Miller, S. (1957). Formation of organic compounds on the primitive earth, in *The Origin of Life on the Earth*, Rep. Internat. Symp., Publ. House. Acad. Sci. USSR, pp. 73-85.
- Morizawa, K. (1927). The extractive substances in *Octopus octopodia*, *Acta. Schol. Med., Univ. Kioto*, **9**:285-302.
- Moyle, V. (1949). Nitrogenous excretion in chelonian reptiles, *Biochem. J.*, **44**:581-584.
- , and E. Baldwin (1952). Volatile fatty acids of *Ascaris lumbricoides* from the pig, *Biochem. J.*, **51**:504-510.
- Needham, J. (1931). Uricotelic metabolism and the evolution of the terrestrial egg, in *Chemical Embryology*, Cambridge University Press, London and New York, Vol. 2, pp. 1132-1145.

- , D. M. Needham, E. Baldwin, and J. Yudkin (1932), A comparative study of the phosphagens, with some remarks on the origin of vertebrates, *Proc. Roy. Soc. (London)*, **B**, **110**:260-294.
- Oparin, A. (ed.) (1957), *The Origin of Life on the Earth*, Rep. Internat. Symp., Publ. House, Acad. Sci. USSR.
- Pirie, N. W. (1937), The meaninglessness of the terms life and living, in *Perspectives in Biochemistry*, Cambridge University Press, London and New York, pp. 11-22.
- Pranker, T. A. J. (1961), *The Red Cell. An Account of Its Chemical Physiology and Pathology*, Oxford University Press, London and New York, p. 136.
- Roche, J., N. v. Thoai, I. Garcia, and Y. Robin (1952), Sur la nature et la répartition des guanidines monosubstituées dans les tissus des invertébrés. Présence des dérivés guanidiques nouveaux chez des Annélides, *Compt. rend. soc. biol.*, **146**:1902-1905.
- Schmidt, C. L. A., and T. Watson (1918), A method for the preparation of taurine in large quantities, *J. Biol. Chem.*, **33**:499-500.
- Smith, H. W. (1930), Metabolism of the lungfish, *Protopterus aethiopicus*, *J. Biol. Chem.*, **88**:97-130.
- (1936), The retention and physiological role of urea in the Elasmobranchii, *Biol. Rev.*, **11**:49-82.
- Snell, E. E. (1956), Some aspects of vitamin and growth factor research, in *Currents in Biochemical Research*, Interscience Publishers, Inc., New York, pp. 87-114.
- Städeler, G., and F. T. Fréichs (1858), Über das Vorkommen von Harnstoff. Taurin und Scyllit in den Organen der Plagiostomen, *J. prakt. Chem.*, **73**:48-55.
- Thoai N. v., and Y. Robin (1954a), Métabolisme des dérivés guanidylés. II. Isolement de la guanidinotaurine (taurocyamine) et de l'acide guanidinoacétique (glycoyamine) des vers marins, *Biochim. et Biophys. Acta*, **13**:533-536.
- (1945b), Métabolisme des dérivés guanidylés. IV. Sur une nouvelle guanidine monosubstituée biologique, l'ester guanidino-éthylsérylphosphorique (lombri-cine) et le phosphagène correspondant, *Biochim. et Biophys. Acta*, **14**:76-79.
- Thoai, N. v., J. Roche, Y. Robin, and N. v. Thiem (1953a), Sur la présence de la glycoyamine (acide guanidyl acétique), de la taurocyamine (guanidyl taurine) et les phosphagènes correspondants dans les muscles des vers marins, *Biochim. et Biophys. Acta*, **11**:593.
- (1953b), Sur deux nouveaux phosphagènes: la phosphotaurocyamine et la phosphoglycoyamine, *Compt. rend. soc. biol.*, **147**:1241-1243.
- (1953c), Sur le phosphagène de *Lumbricus terrestris* sp., *Compt. rend. soc. biol.*, **147**:1670-1672.
- Urey, H. C. (1959), The atmospheres of the planets, *Handb. Phys.*, **52**:363-418.

The Nature and Diversity of Catalytic Proteins

Paul D. Boyer

**Department of Physiological Chemistry
University of Minnesota, Minneapolis, Minnesota**

Most investigators, in considering the general topic of this lecture series, "The Nature of Biological Diversity," are likely impressed with both the breadth and the importance of the subject matter. I share such a view, and indeed find the subtopic to which my remarks are to be addressed, "The Nature and Diversity of Catalytic Proteins," to be a subject of wide scope. In part our awe of these topics, as scientists, reflects the wealth of experimental material to be considered, together with the recognition that the totality of our present information is meager compared to the scope of the subject, and, further, is to a large extent only descriptive and superficial. We lack the knowledge and the insight to develop the generalized concepts that will give coherence and understanding to the diversity encountered. In this presentation, an attempt will be made to give some important generalizations about enzymes that appear valid at this time, to point out various similarities and diversities which need further understanding, and to suggest some relationships and generalizations which are indicated from present research developments. In illustration of

some concepts or points, considerable use will be made of data from our group at Minnesota because I am most familiar with them and can speak most freely of them, not because these data are necessarily the best to illustrate a particular point.

From the viewpoint of an enzymologist, the most striking characteristic of different living forms is not their diversity but their similarity. Morphological and functional differences among organisms and cells are large compared to chemical differences, and in particular, compared to differences in cellular enzyme chemistry. Some generalizations about the nature of enzyme action, which are so well recognized that their importance may be overlooked, are as follows:

Catalytic proteins (enzymes) are found in all cells. This and the following generalization are absolute in that no known exceptions have been found. Thus biologists would predict with a high degree of confidence that any living form found on earth would have enzymes in its cells.

All cells have in common some basic enzymic reactions. All known types of cells, including many diverse unicellular organisms, have enzyme systems for the formation and use of certain key chemical substances, such as amino acids, particular organic phosphates, and ribonucleic acids.

Various enzymic reactions have important common features of mechanism. The most prominent of these features is that enzymic catalyses occur by specific combination of the substrate molecules with particular regions of the enzyme molecule. Possibilities of some other common features in finer details of mechanism will be examined later.

Enzyme cofactors have similar roles in different cells. This has been an important generalization in the elucidation of metabolic function of vitamins. Biochemists have frequently taken advantage of the knowledge that a cofactor role found with microorganisms will apply to larger, more complex forms in an identical manner or one which is closely related chemically.

Among the implications of the first two generalizations are those related to the origin of life itself, and a few comments of a phylogenetic nature may be in order, particularly in this day of interest in planets other than earth. Existence of living material without the presence of catalysts therein is difficult to visualize, but one may ask if the enzyme protein is the only possible way for a variety of catalysts to be produced from chemical elements. Are the properties of phosphate and phosphate compounds, and of nucleic acids such that

other means of handling cellular energy and reproductive requirements are not possible? The enzymological unity of the forms of life we recognize might find its explanation in one of the following:

1. *A common origin of all living cells.* This is implicit in theories of the origin of life based on the occurrence of at least one highly improbable event.

2. *Given the changing chemical environment to which earth has been exposed, the development of life is a logical consequence.* This implies that the basic chemistry common to various cells is uniquely required for life under our present conditions.

3. *Living forms with different chemical bases have existed on earth, but that form with a chemistry as we know it today has, through adaptability, been the only form to survive.*

Means are not at hand to distinguish among these or other possibilities, and it is not my purpose to dwell further upon such matters but to turn attention to certain questions for which pertinent information is at hand. Some questions about the diversity of enzymes to which the bulk of this lecture will be addressed are:

Do enzymes catalyzing the same or similar reactions in various cells have structural features in common? Features about which meager information is becoming available pertain to size, amino acid composition, amino acid sequences in catalytically important areas, and the relation of degradation and of modification of composition to catalytic activity. In this lecture, some results on amino acids present at catalytic sites of hydrolytic enzymes and on degradation of enzymes in relation to activity will be presented.

Do enzymes from different sources catalyzing the same reaction do so by the same or similar mechanism? At this stage the answer appears to be yes for a number and perhaps for most enzymes, but with some enzymes there is strong suggestion that different mechanisms hold. Attention will be directed toward some results with aldolases, and with several types of enzymes involved in phosphorylation reactions.

Are there additional common features of enzyme mechanisms which are as yet unrecognized? Study of mechanism has many levels, and as finer details of mechanism are revealed, examination for possible similarities in the chemistry of various catalyses is in order. Some of our studies on syntheses coupled to ATP cleavage have given indications of previously unrecognized features of enzymic catalyses, and these implications, together with an anal-

ogy between oxidative phosphorylation and muscle contraction, will be presented.

Some relations of amino acid composition to activity

Amino acid composition and partial or complete amino acid sequence studies have shown only small differences in proteins with a particular biological activity from different animal species, namely, with insulins, hemoglobins, and some pancreatic enzymes. Small but detectable differences in enzymes from one tissue have also been found and the recognition of such variants will likely increase as more discerning methods are applied. Markert has appropriately termed such closely related enzymes "isozymes" (1). The similarities of the protein structure of the same enzyme from different tissues is illustrated by the immunological cross reactions among various mammalian tissue lactate dehydrogenases (2), as well as various lactate dehydrogenases from a single tissue (1). Differences in composition are also shown by differences in catalytic behavior of purified lactate dehydrogenases from different tissues, as documented by Kaplan and co-workers (3). Such differences may have clinical use in identification of the source of an enzyme from its properties.

One aspect of amino acid composition of enzymes of particular recent interest has been the demonstration of the presence of a similar amino acid sequence near the reactive seryl residue of a variety of hydrolytic enzymes. Pertinent findings are summarized in Table 1.

Table 1. Amino acid sequences near reactive seryl residues of some hydrolytic enzymes

Enzyme	Sequence
Trypsin	-Glycyl-aspartyl-seryl-glycyl-
Chymotrypsin	-Glycyl-aspartyl-seryl-glycyl-
Thrombin	-Glycyl-aspartyl-seryl-glycyl-
Elastase	-Glycyl-aspartyl-seryl-glycyl-
Liver aliesterase	-Glycyl-glutamyl-seryl-glycyl-
Pseudocholinesterase	-Glycyl-glutamyl-seryl-glycyl-
Bacterial proteinase	-Threonyl-seryl-methionyl-alanyl-

SOURCES: Original data for the first six enzymes are cited in (4); the sequence for bacterial proteinase is given by Sanger and Shaw (5).

Identification of the peptide containing the active seryl residue has been made by various groups, taking advantage of the basic observation of Jansen, Nutting, and Balls (6), showing that with inhibition by P^{32} -labeled diisopropylphosphorofluoridate (DFP), radioactive phosphorus became attached to an amino acid at the active site of chymotrypsin. The occurrence of the same tetrapeptide sequence, glycyl-aspartyl-(or glutamyl) -seryl-glycyl- in the first six enzymes listed in Table 1 has given rise to the suggestion that such a sequence plays an important role in the activation of the seryl hydroxyl group. That such a sequence is not essential, however, is shown by the recent work of Sanger and Shaw on a bacterial proteinase (5). The sequence they found (Table 1) bears no resemblance to that found in the other enzymes. Clearly more than one means is available for bringing about the participation of a seryl hydroxyl in a catalytic site of a hydrolytic enzyme. The occurrence of a similar sequence in the first group of enzymes may actually reflect some feature other than a basic catalytic requirement.

The probability that only certain regions of enzymes participate in the catalyses raises the question of whether considerable portions of the amino acid chain or chains might not be essential for catalysis, and, further, that much of the observed diversity of composition of enzymes catalyzing a particular reaction might reside in the apparently unessential portions of the protein. Our limited knowledge in this area allows no generalizations as yet. With two enzymes, papain and enolase, retention of activity after considerable degradation has been reported. Hill and Smith (7) removed over one-half of the amino acid residues from papain of mol wt about 21,000 by the action of an aminopeptidase, with retention of full activity. Nylander and Malmström (8) similarly accomplished removal of about 150 amino acid residues of the yeast enolase molecule of mol wt about 66,000 from either the carboxyl terminus or from the amino terminus of the single peptide chain without activity loss. Some enzymes are much more sensitive to degradation, however. Thus bovine pancreatic ribonuclease of mol wt about 13,000 reversibly loses activity with removal of a 20 peptide unit from the amino terminus (9). In contrast to the results with papain and enolase, Drechsler, Boyer, and Kowalsky (10) reported a rather striking loss in catalytic activity and change in specificity of a comparatively large enzyme, rabbit muscle aldolase, of mol wt 147,000, with removal of only three COOH-terminal tyrosyl residues.

The modification of aldolase activity accompanying removal of

tyrosines is accomplished by incubating the crystalline aldolase with carboxypeptidase. The results of a typical digestion are shown in Fig. 1. With release of the tyrosines there is a parallel loss of activity, giving rise to a new species which has about 5 per cent of the original activity toward fructose 1,6-diphosphate as a substrate. This change in the aldolase molecule is not accompanied by any discernible change in its size or physical properties. Aldolase, which catalyzes the cleavage of

FIG. 1. Tyrosine release and activity loss accompanying carboxypeptidase digestion of rabbit muscle aldolase (10).

of fructose 1,6-diphosphate to 2 triose phosphates, will also catalyze cleavage of fructose 1-phosphate to glyceraldehyde and dihydroxyacetonephosphate. Surprisingly, the original weak catalytic activity toward fructose 1-phosphate is actually slightly increased after removal of the tyrosines. These and other aspects of aldolase action are of keen interest, but take us away from the main thread of our topic. A principal reason for giving the findings on aldolase degradation in this lecture is that they have served as the basis for some interesting comparative studies of aldolase by W. J. Rutter at the University of Illinois.

Some comparisons of muscle, liver, and yeast aldolases

The experiments on carboxypeptidase degradation of aldolase have, as indicated above, led to the recognition that muscle and liver aldolase may have considerable structural and catalytic features in common. Some of the characteristics of these two enzymes are summarized in Table 2. The two enzymes are similar in molecular size, number of —SH groups per mole, and ultraviolet absorption spectrum. Carboxypeptidase removes only one tyrosyl residue from liver aldolase, in contrast to the three COOH-terminal residues removed from muscle aldolase. The liver aldolase, *before* carboxypeptidase digestion,

Table 2. Comparative properties of muscle and liver aldolases

Property	Muscle	Liver
Molecular weight	150,000	159,000
—SH per mole	28	29
Tyrosines removed by carboxypeptidase	3	1
Activity $\frac{\text{fructose-di-P}}{\text{fructose-1-P}}$	100/2.1 (Native) 6/2.5 (–3 tyrosines)	100/43 (Native) 40/? (–1 tyrosine)
Ultraviolet spectra	Enzymes have similar spectra	

SOURCES: Refs. 10, 11.

has a catalytic action on fructose 1,6-diphosphate and fructose 1-diphosphate rather similar to that shown by muscle aldolase *after* carboxypeptidase digestion. Thus the principal differences in catalytic behavior between the two enzymes may rest in the presence of the two additional COOH-terminal tyrosines in the muscle aldolase. In addition to their indication of more structural relation between liver and muscle aldolase than previously recognized, these studies are of interest in that they illustrate well the importance of small differences in fine structure of a protein to its biological activity.

In contrast to the similarity of liver and muscle aldolase, yeast aldolase differs markedly from muscle aldolase. Some characteristics of the yeast and muscle enzymes are summarized in Table 3. The metal requirements of the yeast enzyme and the inhibition by chelating agents, not shown by the muscle enzyme, are striking. In addition, the marked difference in the number and behavior of —SH groups, as well

as other differences between the two enzymes, suggests that the enzymes have some important differences in the mechanism of their catalytic reactions. The situation is reminiscent of the familiar biological truism about diversity, namely, "there is more than one way to skin a cat."

A number of other yeast enzymes have not been found to have the marked differences from their mammalian counterparts shown by the aldolases. Thus both liver and yeast alcohol dehydrogenases contain Zn as an essential component (14), and yeast and muscle glyceralde-

Table 3. Some characteristics of yeast and muscle aldolases

Characteristic	Yeast	Muscle
Metal requirement	Zn ⁺⁺ , K ⁺	None
Effect of chelating agents	Inhibit	No effect
—SH groups/mole	2-3	27-28
Substrate effect on <i>p</i> -mercuribenzoate inhibition	Protects	No effect

SOURCES: Refs. 11, 12, 13.

hyde-3-phosphate dehydrogenases appear to have an essential —SH group or groups at the catalytic site (15). A difference between yeast and mammalian lactate dehydrogenases deserves mention however, if only for its social implications. The yeast enzyme uses a flavin-containing cofactor, in contrast to the reaction of the mammalian enzyme, which uses diphosphopyridine nucleotide. Were it not for this difference, lactic acid and not ethyl alcohol would be expected as a principal metabolic product of yeast.

Amino acid analog incorporation and activity

An additional aspect of variation in composition as related to enzyme function comes from studies on the specificity of protein synthesis. Results and implications of recent researches are ably discussed in a review by Vaughan and Steinberg (16); studies of amino acid analog incorporation appear to be of particular pertinence. Various investigators working with bacterial systems have demonstrated that amino acid analogs may be incorporated into tissue proteins in place of naturally occurring amino acids. Recently in our

laboratory Westhead has demonstrated such incorporation in a living animal, namely, that rabbits fed *p*-fluorophenylalanine show extensive incorporation of this amino acid analog into tissue proteins. There are clear indications, particularly with bacterial systems, that synthesis of proteins which are biologically inactive may be one cause of the toxicity of the amino acid analogs. A few highly purified enzymes containing analogs have been isolated and their properties compared to their normal counterparts. Crystalline *Bacillus subtilis* α -amylase containing considerable ethionine in place of methionine retains full activity (17), but replacement of around one-sixth of the phenylalanine residues by *p*-fluorophenylalanine results in 30 per cent loss of activity (18). Similarly, *p*-fluorophenylalanine incorporation into penicillinase gives extensive loss of activity (19). In contrast, Westhead (20) found that rabbit aldolase with replacement of about one-fourth of the phenylalanine residues with *p*-fluorophenylalanine was indistinguishable from the normal enzyme in physical, chromatographic, and catalytic properties. Variable effects of analog incorporation could logically reflect the importance of the residues being replaced to the catalytic site or essential structure of a given enzyme. The occurrence of such variability is of interest because it suggests that even selectivity for normal amino acids may be subject to error, and enzyme microheterogeneity might logically occur without primary genetic variation.

The generality of cofactor function

The similarity of the actions of vitamin-containing cofactors is quite well documented and understood. Also the role of divalent cations as an activator is well recognized. For example, every enzyme system reacting with ATP has been found to require a divalent cation, usually Mg^{++} , for maximal activity. Some years ago, during a brief stay at the Marine Biological Laboratory at Woods Hole, I had occasion to study whether a less well understood cofactor requirement, namely, that of the monovalent cation K^+ , also appeared to be a general requirement for one enzyme from a variety of sources (21). The enzyme studied was pyruvate kinase, catalyzing the transfer of a phosphoryl group from phosphoenolpyruvate to ADP. The results obtained are summarized in Table 4 and show that, in all tissues and species tested, activation by K^+ was demonstrable even without dialysis of the tissue extract. With a fresh-water species, *Anadonta*, which has a characteristically low intracellular salt content, only a weak activation was demonstrable, however, without dialysis. Thus

even in this species, K^+ appears to be required for pyruvate kinase activity but in a considerably lower concentration than for the other tissues.

In addition to the requirement of K^+ for pyruvate kinase from sources listed in Table 4, K^+ has also been shown to activate pyruvate

Table 4. K^+ activation of pyruvate kinase from different sources

Tissue used	Activity *		Muscle source	Activity *	
	No K^+	+ K^+		No K^+	+ K^+
Rabbit					
Heart	15	100	Skate	5	45
Liver	4	18			
Kidney	1	6	Fundulus	2	65
Uterus	3	18			
Brain	14	70	Limulus	41	390
Dogfish			Pecten	8	39
Body muscle	8	230			
Stomach muscle	6	25	Thyone	0.6	6
Brain	7	110	Phascolosoma	0.1	2
Tetrahymena	23	39	Torpedo	0.4	4
			Anadonta †	26	39

* μ moles pyruvate formed per 100 mg tissue per 10 minutes.

† Dialyzed.

SOURCE: Ref. 21.

kinase from yeast (22) and from plants (23). The K^+ activation thus appears to be a rather general phenomenon, suggesting either an irreplaceable role for K^+ in the reaction mechanism or a close phylogenetic relationship of the various organisms.

Mechanistic relations of enzymes catalyzing similar reactions

Enzymologists have quite logically based much of their work on the probability that enzymes catalyzing similar reactions do so by similar mechanisms. Various examples could be cited; a particularly good one comes from the dehydrogenases linked to the cofactor, diphosphopyridine nucleotide. In all instances studied, the reduction of the cofactor occurs with a direct transfer of a hydrogen, probably as a hydride ion, from the substrate to the cofactor (24). Another

example comes from the kinases which catalyze transfer of the terminal phosphoryl group of ATP to various acceptors. Studies in our laboratory (see ref. 4) and that of Mildred Cohn (25) have shown that in various kinase reactions, the catalysis occurs with a cleavage of the bond between the phosphorus atom of the group transferred and the oxygen atom linking the phosphoryl group to the rest of the donor molecule, and that the phosphoryl group is transferred without interchange of its oxygens with oxygens from water or substrates. Recently we have probed further the details of the pyruvate kinase and hexokinase reactions by study of the reaction kinetics and substrate binding (26). These and earlier studies have led to a concept of the catalysis as depicted in Fig. 2 for pyruvate kinase. The pattern given in this

FIG. 2. A schematic illustration of substrate binding and a transition state for pyruvate kinase.

figure likely holds for other kinases as well. Simple but important aspects are that ATP and ADP share a common binding site, and pyruvate and phosphopyruvate another binding site, but that both ATP and phosphoenolpyruvate cannot be bound to the enzyme at the same time because of a common position at the active site for their transferable phosphoryl group.

An important feature of the catalysis by kinases, as indicated for pyruvate kinase in Fig. 2, is the transfer of the phosphoryl group directly from the donor to the acceptor, without formation of any phosphoryl enzyme intermediate. This is in contrast to the intramolecular phosphoryl transfer catalyzed by phosphoglucomutase, in which there is good evidence for a phosphoryl enzyme intermediate (27). Also, speculation has become common that many enzyme re-

actions involve function of discrete, covalently bound intermediates in their catalyses. Such is not the case with the kinases. Of the various experimental results pointing to the absence of a phosphoryl enzyme intermediate in the kinase reactions, results of one recent approach (28) may be of interest because they allow definitive conclusions and because they illustrate a simplicity of experimental design based on the use of comparatively large amounts of a given enzyme. The findings of such an experiment with pyruvate kinase are given in

FIG. 3. Demonstration of lack of phosphoryl enzyme formation by pyruvate kinase.

Fig. 3. The system used is actually a sensitive test for formation of ADP. The initial reaction mixture contains phosphoenolpyruvate, ATP, and reduced diphosphopyridine nucleotide, together with small traces of ADP and pyruvate. Upon addition of catalytic amounts of pyruvate kinase and lactate dehydrogenase, practically all the ADP present is phosphorylated to give ATP through the pyruvate kinase reaction, with maintenance of a high ATP/ADP ratio. The pyruvate formed from phosphoenolpyruvate oxidizes some DPNH, with a resultant slight decrease in absorbancy at 340 millimicrons. The absorbancy as measured on a sensitive recording spectrophotometer

soon becomes stable; the amounts of pyruvate kinase and lactate dehydrogenase present are sufficient to catalyze rapid attainment of equilibrium. At this stage a comparatively large amount of pyruvate kinase is added. If any appreciable phosphoryl enzyme is formed from reaction of the enzyme with ATP, the resultant ADP formation would give an immediate drop in absorbancy at $340\text{ m}\mu$; none such is detected. The continued slow decline represents a side reaction catalyzed by the very large excess of pyruvate kinase (26). The sensitivity of the system to ADP is indicated by the subsequent addition of an amount of ADP equivalent to about one-half of the concentration of active sites of the large amount of pyruvate kinase added. The results demonstrate clearly the lack of formation of phosphoryl enzyme even in the presence of all reactants and under conditions favorable to rapid net reaction.

FIG. 4. The two general patterns for syntheses coupled to ATP cleavage.

The lack of phosphoryl enzyme formation as shown by Fig. 3 could result from the existence of the pyruvate kinase as a stable phosphoryl enzyme. If any such phosphoryl group participated in the catalysis, it would be rapidly replaced by the phosphoryl group from ATP. This possibility is readily checked by the use of radioactive P³²-labeled ATP. Such experiments rule out the formation of a stable phosphoryl enzyme as a catalytic intermediate (28).

Other phosphorylation enzymes which have been of interest are those which catalyze syntheses coupled to cleavage of ATP. The two patterns of ATP cleavage which have been found with this group of enzymes are indicated schematically in Fig. 4. Examples of this important type of reaction are the synthesis of acyl coenzyme A derivatives from fatty acids and coenzyme A, of glutamine from glutamate and ammonia, and of glutathione from its component amino acids. Studies with the stable isotope O¹⁸ have shown that in such syntheses substrates and not water furnish the oxygen necessary for ATP

cleavage, and that of the four possible modes of ATP cleavage leading to release of one or both of the terminal phosphoryl groups of ATP, only the two indicated in Fig. 4 occur (4, 25). Such results give further evidence of the mechanistic similarity of various enzymes catalyzing similar reactions.

One other aspect of the mechanism of enzymes catalyzing similar reactions deserving brief mention pertains to enzyme specificity. An unsettled question is whether enzymes catalyzing a similar reaction, but with marked differences in the degree of specificity, do so by a similar mechanism. Phosphatases are known, for example, with a high degree of specificity, such as a phosphatase which will remove the phosphoryl residue only from the 5' position of nucleotides. Other phosphatases are known with considerably wider specificity, ranging to the extracellular phosphatase from *Escherichia coli*, which Heppel (29) has found to have a remarkable ability to hydrolyze a wide variety of phosphate esters. Aside from their interest as examples of the variability of enzyme specificity, such findings need study and explanation on a mechanistic level.

Possible generalities for enzyme mechanisms

Thus far students of enzyme mechanism have not uncovered any properties of the protein molecule which serve as a general basis for enzyme catalysis, and beyond the implications inherent in the participation of a beautifully specific combination of enzyme and substrate, one rapidly enters an area of speculation. With regard to thermodynamic considerations, perhaps the most important finding is that all enzymes appear to lower the energy of activation for the reaction. Some kinetic findings of possible importance to enzyme mechanisms in general have arisen from our studies of syntheses coupled to ATP cleavage, and I would like to present some aspects of these researches.

In probing at the mechanism of glutamine synthetase, catalyzing

the above reaction, studies of the dynamic reaction rate at equilibrium were undertaken, that is, the rate of interconversion of reactants to products when no net reaction is occurring. Such rates are measurable by adding traces of isotopically labeled substrates to the reaction mixture at dynamic equilibrium. Somewhat to our surprise, we found marked inequalities in the isotopic exchange rates at equilibrium. Thus, with variations in equilibrium conditions, the rate of inter-

change of inorganic phosphate with ATP would be made to be about equal, greater than, or less than the rate of interchange of glutamate with glutamine (30). This led to some theoretical considerations of the factors governing reaction rates at equilibrium. Some clear and useful concepts have emerged (31). With regard to the glutamine synthetase reaction, as depicted in Fig. 5, an explanation for the inequalities of exchange rates is forthcoming if the slow steps in the overall catalysis are not those in which covalent bonds are formed or broken, but those of release of substrates from the enzyme. Inequalities of exchange rates near equilibrium have been demonstrated for the aldolase reaction as well (32), and studies with this and with dehydrogenase reactions are in progress at Minnesota. The experi-

FIG. 5. A scheme for the glutamate synthetase reaction showing substrate binding and release steps.

mental results indicate that the tacit assumption frequently made, that the slow steps in enzymic catalyses are those in which the covalent bonds are broken and formed, is open to serious question.

Recognition that there may be a considerable energy barrier between the free and bound forms of substrates, taken together with the remarkable specificity of enzymic catalyses, suggests that the important structural and electronic changes necessary for catalysis accompany the binding, and that reaction of the bound substrates occurs readily with little further change in their position and properties. Such considerations lend support to the concept of the synthetase and other enzymic reactions as concerted reactions (4, 33), and actually give a simpler picture of the catalysis. This may be illustrated with a typical synthetase reaction, the acetate thiokinase reaction, which also shows inequalities of substrate exchanges at equilibrium (30). In line with the above discussion, the acetate

SCHEMATIC FORMATION OF ACETYL-CoA WITHOUT
ACETYL-AMP AS AN INTERMEDIATE

(Involves only one key transition state, without
shift in reactant binding for catalysis)

FIG. 6. A scheme for the acetate thiokinase reaction with one key transition state.

SCHEMATIC FORMATION OF ENZYME BOUND ACETYL-AMP

(Involves two key transition states with shift of
position of bound reactants)

FIG. 7. A scheme for the acetate thiokinase reaction with two key transition states.

thiokinase reaction can be visualized as occurring with participation of one key transition state, as depicted in Fig. 6. The reactants are bound in an activated form, and as the acetate carboxyl oxygen approaches the P atom of the ATP, the S of coenzyme A approaches the acetate carboxyl carbon, with the formation of the products in one concerted reaction. This is in contrast to the reaction as usually depicted with acetyl adenylate as a discrete intermediate (see ref. 34). Participation of such an intermediate would appear to necessitate rather complicating shifts of the reactants during catalysis. An attempt to depict this difficulty is given in Fig. 7. For discrete formation of acetyl adenylate, the S atom of the bound and activated coenzyme A molecule must somehow be held away from the acetate carboxyl carbon, which would become increasingly susceptible to reaction as its oxygen departed for cleavage of the ATP. Then after acetyl adenylate had formed, a shift would need to occur, to allow the coenzyme A and acetyl adenylate to react.

The manner by which the synthetases and much simpler enzyme systems react is obviously far from settled. The recognition of substrate release as potential slow steps in the catalyses, together with other information, warrants the hypothesis that enzyme reactions in general may proceed through binding of reactants favoring reaction through one key transition state (4).

A similarity of oxidative phosphorylation and muscle contraction

For a number of years our group has been interested in the basic problem of how oxidative enzymic reactions are coupled to formation of ATP, usually referred to as oxidative phosphorylation. Not being content with having one seemingly insoluble problem on our hands, we recently undertook some experiments related to muscle contraction. The divergence is not as wide as it might seem, and one principal reason for our interest in ATP cleavage by myosin and actomyosin was the finding by Levy and Koshland (35) that an unexplained oxygen exchange reaction accompanied the cleavage.

Several years ago in studies on oxidative phosphorylation by mitochondria, the discovery was made that the process was characterized by a rapid exchange of inorganic phosphate with ATP as well as a much more rapid exchange of phosphate oxygens with water. Typical results are given in Table 5. Such rapid exchange of phosphate oxygens is unusual, and one possible explanation may be the rapid formation and hydrolytic cleavage of a phosphorylated intermediate.

Recent studies by Dempsey and Boyer (37) have shown that in the presence of ATP, myosin and actomyosin catalyze an exchange of oxygens of inorganic phosphate with those of water. One experimental result which led to these findings is depicted in Fig. 8. This figure

Table 5. *Exchange reactions catalyzed by mitochondria capable of oxidative phosphorylation*

$\text{P}_i^{32} \rightleftharpoons \text{ATP exchange}$	$\text{O}_{\text{P}_i}^{18} \rightleftharpoons \text{HOH exchange}$
13 μmoles	280 μmoles

Mitochondria with ATP, P_i , Mg^{++} , K^+ , at 23° , 15 minutes. No substrate and no oxygen uptake.

SOURCE: Ref. 36.

shows the time course of appearance of O^{18} from water into phosphate cleaved from ATP by actomyosin. The phosphate appearing in the initial part of the reaction contains only the one water oxygen expected from the hydrolytic reaction. With continued incubation, more than one water oxygen is found in the phosphate, suggesting a concentration dependent on exchange of phosphate oxygens of the medium with water oxygens. The experiments reported in Table 6 provide a con-

Table 6. *P_i oxygen exchange during hydrolysis of ATP*

O ¹⁸ atom, excess, %			
H ₂ O of medium	P _i added	P _i isolated	mM H ₂ O exchanged
Actomyosin hydrolysis			
1.44	0.000	0.099	15.0
0.00	1.005	0.817	15.0
Myosin hydrolysis			
1.44	0.000	0.090	14.0
0.00	1.012	0.854	13.6

SOURCE: Ref. 37.

clusive demonstration that most, if not all, of the oxygen exchange accompanying ATP cleavage results from exchange of phosphate oxygens of the medium. In these studies, the exchange was measured both by incorporation of oxygen from H_2O^{18} into phosphate as well

as from the loss of O^{18} from P_i-O^{18} to water. The results show that, within experimental error, the extent of exchange measured by the two approaches is equal and that therefore the species undergoing exchange is the inorganic phosphate of the medium. As with oxidative phosphorylation, such results suggest the reversible formation and hydrolysis of a phosphoryl derivative. The following points of com-

FIG. 8. The time course of appearance of O^{18} from H_2O^{18} into phosphate cleaved from ATP by actomyosin (37).

parison of oxidative phosphorylation and of ATP cleavage by actomyosin and myosin emerge:

An unexplained exchange of inorganic phosphate oxygens with water oxygens occurs, in both processes, possibly reflecting reversible phosphorylation.

Protein structure changes may accompany both processes. Possibility of structure change accompanying electron transport is clearly indicated by the difference in structure of oxidized and reduced cytochrome *c* as indicated by the difference in proteolytic digestibility (38) and as demonstrated by nuclear magnetic resonance studies of Kowalsky (39) in our laboratory.

In one instance, ATP cleavage may induce structural change; in the other, structural change may induce ATP formation.

Past experience suggests that further work will likely show the last speculation to be invalid, but the possibility that such apparently diverse processes as oxidative phosphorylation and muscle contraction may have an important common mechanistic feature is indeed an intriguing one.

Genetic control of and induced change in the nature of enzymes

A lecture under the heading of the nature and diversity of catalytic proteins would not be complete without some consideration of the factors governing the formation of enzymes. Two exciting areas of research are on the genetic control of the fine structure of enzymes and on the nature of enzymic changes which lead to resistance to antibiotics and anticancer agents.

As an example of genetic control of the fine structure of an enzyme, the excellent researches of Hotchkiss and Evans (40) may be cited. Concepts which have arisen from their work on the genetic modification of a pneumococcal enzyme involved in folic acid synthesis from *p*-aminobenzoic acid, are shown schematically in Fig. 9. Most of the studies of gene-enzyme relationships have concerned all-or-none effects—either a particular enzyme is present or is absent in a given genetic strain. Hotchkiss and Evans have obtained strong genetic evidence that the difference in relative affinities for various analogs of *p*-aminobenzoate noted in their studies results from the variations in the fine structure of a single enzyme. They suggest, as noted in Fig. 9, that DNA controls not only the presence of the enzyme, but that variation in the fine structure of the DNA can result in variation in the enzyme as reflected by the change in inhibitory effects of chemical modifications of *p*-aminobenzoate. As noted by these observers, there will remain some doubt about the interpretation until purification of the enzymes involved can be achieved, but the way seems clear and the authors appropriately state, "It would be a challenging prospect to learn about the linear sequence of amino acids in a series of proteins which have such interesting and characteristic properties in their native folded forms."

Problems of keen fundamental as well as practical importance arise from consideration of the factors responsible for the develop-

ment of resistance to antibiotics and to anticancer agents. Three means of development of resistance appear probable, as follows:

1. Gradual or sudden changes in key properties of an enzyme or enzyme system.
2. Changes in enzyme concentrations leading to altered routes of utilization of former lethal syntheses.
3. Exclusion of the agents from sensitive sites by permeability, binding, or other changes.

Comparative studies on highly purified enzymes from normal and resistant tumor cells or bacterial cells have been extremely limited.

FIG. 9. Genetic control of fine structure of enzyme system using *p*-aminobenzoate (40).

In an important paper some years ago, Kulowitz and Ott (41) found purified lactate dehydrogenase from normal muscle and from the Jensen sarcoma arising from muscle to be closely similar if not identical in catalytic and immunologic properties. With further study of the biochemistry of cancer, an appreciation has developed that cancer cells, like bacterial cells, show a remarkable and deadly ability to adapt to their environment. Recent reviews, such as that of Anderson and Law (42), may be consulted for a summary of various findings. The data point unmistakably to the conclusion that resistance can arise from change in the properties of certain enzymes, although in no instance to our knowledge has proof of such a change been documented with a purified enzyme. Some examples where a change in properties of key enzymes is strongly indicated include the

resistance of *Enterococcus stei* to sulfonamide (43), and of *Escherichia coli* to Aureomycin (44). In the latter case, electron-transport enzymes in cell-free extracts from resistant cells but not from nonresistant cells were demonstrated to be insensitive to added Aureomycin. Resistance of Erlich ascites cells to mercaptopurine and azaguanine has been associated with the loss of enzymes for conversion of these anticancer agents to their ribotides. (45, 46). However, even the decreased ability for ribotide synthesis from the anticancer agents may not be the only basis for resistance. Heidelberger et al. (47) have found that although the rate of conversion of fluorouracil to fluorodeoxyuridylate is decreased, the actual amount of fluorodeoxyuridylate present is about the same in the resistant and normal cells, and the resistance was attributed to a decreased ability of the fluorodeoxyuridylate to inhibit the enzymes involved in thymidine synthesis. Increase in the amount of some enzymes may also be associated with resistance. For example, resistance to folic acid analogs in leukemia has been attributed to increased dihydrofolate reductase activity (48).

Differences in properties of enzymes from a given microorganism or cellular strain may also be encountered in the well-known phenomenon of adaptative enzyme formation. In a valuable paper, Halvorson and his colleagues (49) have shown that a highly purified constitutive yeast β -glucosidase resembles very closely the induced enzyme present in a related species in physical properties and maximal reaction velocity, but that the inducible enzyme has an apparent Michaelis constant one-tenth that of the constitutive enzyme. The data suggest that the two enzymes differ in the fine structure near the catalytic site.

The present information about enzyme change in adaptation and development of resistance, although meager, is sufficient already to outline some fascinating but difficult problems ahead. As in many areas of research, the more important results and controlling factors may not be visualized at present or may be improperly conceived. Obtaining an understanding of the relations between structure change and the properties of a given enzyme promises to be a challenging but rewarding task. The stepwise appearance of resistance in some cancer cells suggests that there may be considerable randomness in the genetic control of fine structure of enzymes in all cell populations, and thus that resistance might result from the continued selection of the better adapted cells. On the practical side, one may even hope that our degree of biochemical erudition will increase to the point where we can, by planned development of resistance of a cancer cell to one agent, actually render it more susceptible to the action of another agent.

Conclusions

The audience is perhaps too well aware at this stage of the validity of a comment made at the beginning of this address, namely, that the topic under discussion is broad. Many other facets than these cited herein could be given for your consideration. In any event it is clear to the lecturer, and I hope to the audience, that the basis for the nature and diversity of catalytic proteins is at best only partly understood. It must be emphasized that the occurrence, properties, and mechanism of enzymes from widely different cells are more striking than the differences. In some instances, however, as with yeast and muscle aldolase, rather wide differences in mechanism do appear possible. With regard to enzyme mechanisms, continued search may reveal more basic and simplifying generalities underlying the behavior of enzymes. Some common mechanistic patterns already recognized have been discussed briefly, and possibilities of further generalizations have been mentioned. The problems sketchily outlined herein call for zealous application of man's ingenuity. It is from the probing at and documenting of the diversities of enzymes and their action that important generalizations about living processes will emerge.

References

1. C. L. Markert and F. Möller, Multiple forms of enzymes: tissue ontogenetic and species specific patterns, *Proc. Nat. Acad. Sci. U.S.*, **45**:753-765 (1959).
2. J. S. Nisselbaum and O. Bodansky, Reactions of human tissue lactic dehydrogenases with antisera to human heart and liver lactic dehydrogenases, *J. Biol. Chem.*, **236**:401-404 (1961).
3. N. O. Kaplan, M. M. Ciotti, M. Hamolsky, and R. E. Bieber, Molecular heterogeneity and evolution of enzymes, *Science*, **131**:392-397 (1960).
4. P. D. Boyer, Mechanism of enzyme action, *Ann. Rev. Biochem.*, **29**:15-44 (1960).
5. F. Sanger and D. C. Shaw, Amino-acid sequence about the reactive serine of a proteolytic enzyme from *Bacillus subtilis*, *Nature (London)*, **187**:872-873 (1960).
6. E. F. Jansen, M. D. F. Nutting, and A. K. Balls, Mode of inhibition of chymotrypsin by diisopropyl fluorophosphate. I. Introduction of phosphorus, *J. Biol. Chem.*, **179**:201-204 (1949).
7. R. L. Hill and E. L. Smith, Hydrolysis of mercuripapain by leucine aminopeptidase without loss of enzymic activity, *J. Biol. Chem.*, **231**:117-134 (1958).
8. O. Nylander and B. Malmström, The degradation of yeast enolase with leucine aminopeptidase and carboxypeptidase without change in enzymic activity: N- and C-terminal residues of the enzyme, *Biochim. et Biophys. Acta*, **34**:196-202 (1959).
9. F. M. Richards and P. J. Vithayathil, The preparation of subtilisin-modified

- ribonuclease and the separation of the peptide and protein components, *J. Biol. Chem.*, **234**:1459-1465 (1959).
10. E. R. Drechsler, P. D. Boyer, and A. G. Kowalsky, The catalytic activity of carboxypeptidase-degraded aldolase, *J. Biol. Chem.*, **234**:2627-2634 (1959).
 11. O. C. Richards and W. J. Rutter, Studies on the properties and mechanism of action of various aldolases, *Federation Proc.*, **19**:86 (1960). W. J. Rutter, private communication.
 12. R. J. Peansky and H. A. Lardy, Bovine liver aldolase. I. Isolation, crystallization, and some general properties, *J. Biol. Chem.*, **233**:365-370 (1958).
 13. R. J. Peansky and H. A. Lardy, Bovine liver aldolase. II. Physical and chemical measurements on the crystalline enzyme, *J. Biol. Chem.*, **233**:371-373 (1958).
 14. B. L. Vallee, Metal and enzyme interactions: correlation of composition, functions and structures, in *The Enzymes*, ed. by Boyer, Lardy, and Myrback, vol. 3, Academic Press, Inc., New York (1960).
 15. P. D. Boyer, Sulfhydryl and disulfide groups of enzymes, in *The Enzymes*, ed. by Boyer, Lardy, and Myrback, vol. 1, Academic Press, Inc., New York (1959).
 16. M. Vaughan and D. Steinberg, The specificity of protein biosynthesis, in *Advances in Protein Chemistry*, vol. 14, Academic Press, Inc., New York (1959), p. 115.
 17. A. Yoshida and M. Yamasaki, Studies on the mechanism of protein synthesis incorporation of ethionine into α -amylase of *Bacillus subtilis*, *Biochim. et Biophys. Acta*, **34**:158-165 (1959).
 18. A. Yoshida, Studies on the mechanism of protein synthesis; incorporation of *p*-fluorophenylalanine into α -amylase of *Bacillus subtilis*, *Biochim. et Biophys. Acta*, **41**:98-103 (1960).
 19. M. H. Richmond, Incorporation of DL- β -(*p*-fluorophenyl)[β - 14 C]alanine into exopenicillinase by *Bacillus cereus* 569/H, *Biochem. J.*, **77**:121-135 (1960).
 20. E. W. Westhead and P. D. Boyer, The incorporation of *p*-fluorophenylalanine into some rabbit enzymes and other proteins, *Biochim. Biophys. Acta*, **54**:145-156 (1961).
 21. P. D. Boyer, The activation by K^+ and occurrence of pyruvic phosphoferase in different species, *J. Cell. and Comp. Physiol.*, **42**:71-78 (1953).
 22. I. F. Seits, Role of potassium and ammonium ions in the transfer of phosphate from phosphopyruvic acid to the adenylic system, *Biokhimiya*, **14**:134-140 (1949); *Chemical Abstracts*, **43**:6281 (1949).
 23. G. Miller and H. J. Evans, The influence of salts on pyruvate kinase from tissues of higher plants, *Plant Physiol.*, **32**:346-354 (1957).
 24. B. Vennesland, Stereospecificity of hydrogen transfer in pyridine nucleotide dehydrogenase reactions, *Federation Proc.*, **17**:1150-1157 (1958).
 25. M. Cohn, Mechanisms of enzymic cleavage of some organic phosphates, *J. Cell. and Comp. Physiol.*, **54**:17-32 (1959).
 26. A. M. Reynard, L. F. Hass, D. D. Jacobson, and P. D. Boyer, The correlation of reaction kinetics and substrate binding with the mechanism of pyruvate kinase, *J. Biol. Chem.*, **236**:2277-2283 (1961).
 27. J. B. Sidbury, Jr. and V. A. Najjar, Further studies on the mechanism of phosphoglucomutase; the phosphoenzyme bond, *J. Biol. Chem.*, **227**:517-522 (1957).
 28. L. F. Hass, P. D. Boyer, and A. M. Reynard, Studies on possible phosphoryl enzyme formation in catalysis by hexokinase, pyruvate kinase, and glucose 6-phosphatase, *J. Biol. Chem.*, **236**:2284-2291 (1961).

29. L. A. Heppel, D. R. Harkness, and R. J. Hilmeo. A Study of the substrate specificity and other properties of the alkaline phosphatase of *Escherichia coli*. *J. Biol. Chem.*, **237**:841-846 (1962).
30. P. D. Boyer, R. C. Mills, and H. J. Fromm, Hypotheses for and some kinetic studies with glutamine synthetase and acetate thiokinase, *Arch. Biochem. Biophys.*, **81**:249-263 (1959).
31. P. D. Boyer. Uses and limitations of measurements of rates of isotopic exchange and incorporation in catalyzed reactions. *Arch. Biochem. Biophys.*, **82**:387-410 (1959).
32. I. A. Rose. The mechanism of action of aldolase and the asymmetric labeling of hexose. *Proc. Nat. Acad. Sci. U.S.*, **44**:10-15 (1958).
33. J. M. Buchanan and S. C. Hartman. Enzymic reactions in the synthesis of the purines. *Advances in Enzymol.*, **21**:199-261 (1959).
34. P. Berg. Role of magnesium in acetyl coenzyme A formation by acetothio kinase. *Science*, **129**:895-896 (1959).
35. H. M. Levy and D. E. Koshland, Jr.. Mechanism of hydrolysis of adenosine-triphosphate by muscle proteins and its relation to muscular contraction. *J. Biol. Chem.*, **234**:1102-1107 (1959).
36. P. D. Boyer, W. W. Luchsinger, and A. B. Falcone. O^{18} and P^{32} exchange reactions of mitochondria in relation to oxidative phosphorylation. *J. Biol. Chem.*, **223**:405-421 (1956).
37. M. E. Dempsey and P. D. Boyer. Catalysis of an inorganic phosphate- H_2O^{18} exchange by actomyosin and myosin. *J. Biol. Chem.*, **236**:PC 6 (1961).
38. M. Nozaki, T. Yamanaka, T. Horio, and K. Okuniki. Cytochrome c. I. Purification procedures and proteinase digestion of bakers' yeast cytochrome c. *J. Biochem. (Tokyo)*, **44**:453-464 (1957).
39. A. Kowalsky, unpublished.
40. R. D. Hotchkiss and A. H. Evans. Fine structure of a genetically modified enzyme as revealed by relative affinities for modified substrate. *Federation Proc.*, **19**:912-925 (1960).
41. F. Kubowitz and P. Ott. Isolierung und Kristallisation eines Gärungsfermentes aus Tumoren. *Biochem. Z.*, **314**:94-117 (1943).
42. E. P. Anderson and L. W. Law. Biochemistry of cancer. *Ann. Rev. Biochem.*, **29**:577-608 (1960).
43. A. Wacker, A. Trebst, and H. Simon. Über den stoffwechsel des Sulfanilamids- S^{35} bei empfindlichen und resistenten Bakterien. *Z. Naturforsch.*, **12b**:315-319 (1957).
44. K. Saz and L. M. Martinez. Enzymatic basis of resistance to Aureomycin. II. Inhibition of electron transport in *Escherichia coli* by Aureomycin. *J. Biol. Chem.*, **233**:1020-1022 (1958).
45. P. Reichard, O. Skold, and G. Klein. Possible enzymic mechanism for the development of resistance against fluorouracil in ascites tumors. *Nature (London)*, **183**:939-941 (1959).
46. R. W. Brockman, L. L. Bennett, Jr., M. S. Simpson, A. R. Wilson, J. R. Thomson, and H. E. Skipper. A mechanism of resistance to 8-azaguanine II. Studies with experimental neoplasms. *Cancer Research*, **19**:856-869 (1959).
47. C. Heidelberger, G. Kaldor, K. L. Mukherjee, and P. B. Danneberg. Studies on fluorinated pyrimidines. XI. In vitro studies on tumor resistance. *Cancer Research*, **20**:903-906 (1960).

48. D. K. Misra, R. H. Stewart, M. Friedkin, A. Goldin, and E. J. Crawford, Increased dihydrofolate reductase activity as a possible basis of drug resistance in leukemia, *Nature* (London), **189**:39-42 (1960).
49. A. S. L. Hu, R. Epstein, H. O. Halvorson, and R. M. Bock, Yeast β -glucosidase: comparison of the physical-chemical properties of purified constitutive and inducible enzyme, *Arch. Biochem. Biophys.*, **91**:210-219 (1960).

The Origin of Specific Proteins

Clement L. Markert

Department of Biology
Johns Hopkins University, Baltimore, Maryland

We are all aware of the fact that every field of biology, including molecular biology, contributes in some degree to our appreciation of biological diversity. Underlying the obvious diversity of living organisms is a fundamental uniqueness in the molecular composition of each individual. This uniqueness is based upon characteristic proportions of small molecules and, even more important, upon the specific structures of numerous macromolecules that compose essential parts of the metabolic machinery. Of all these macromolecules, perhaps the widest range of diversity is found in the proteins. The only other class of molecules of nearly equal significance is the nucleic acids—the raw material of molecular genetics. But even the nucleic acids are, in a fundamental sense, recognized principally through the proteins which they control. Just as nucleic acids have occupied the attention of geneticists, so have the properties and activities of proteins been at the focus of much of the research in biochemistry. The extraordinary advances in each of these two fields, genetics and biochemistry, and the remarkably fruitful collaboration between them in the study of

protein molecules have nearly obscured another area of biology that is larger than either of them and, in fact, encompasses both. This is the area best described as developmental biology. I speak today as a devotee and representative of this area. We developmental biologists share with geneticists an interest in gene function, but in addition we are concerned with the *regulation* of gene function during development. We, like the biochemists, are also concerned with the properties of protein enzymes since the characteristics of differentiating cells are basically a reflection of the enzymes they contain. But going beyond the biochemists, we developmental biologists must also investigate the mechanisms responsible for initiating, regulating, or stopping protein synthesis in accord with the requirements of the developing cell. Developmental biology, even that portion which focuses on specific proteins, is vast and complex, and productive research in this area has not yet matched the size of the problems. Nevertheless, progress has been made and I shall attempt to synthesize and interpret a variety of data collected from diverse fields but all bearing on the problem of the origin or synthesis of specific proteins in an organism. Genetics and biochemistry will necessarily make the major contributions.

The prevailing viewpoint today is that the primary structure of a protein—the linear sequence of amino acids—reflects a corresponding sequence of nucleotides in desoxyribonucleic acid (DNA) and ribonucleic acid (RNA). A growing body of evidence supports this view; in any event since a general correspondence between genes and proteins is well established, no extensive review of this area would be justified now. However, it may be useful by way of introduction to call attention to the evidence relating genes to the protein hemoglobin in human beings. This is a well-known and often-reviewed subject but provides a good foundation for my later remarks.

Hemoglobin synthesis

Hemoglobin is a complex molecule consisting of four polypeptide chains. In the normal adult, two identical α chains combine with a pair of identical β chains to compose the finished molecule, which thus may be written $\alpha^2\beta^2$. Now the α and β chains are each under separate gene control and mutant forms of each of these two genes have been discovered. Individuals with sickle-cell anemia, for example, possess a mutant form of the β gene and synthesize an altered hemoglobin molecule in which valine has replaced glutamic acid at one position in each of the two β chains (for review see Conference on

Hemoglobin, 1958). This is exactly what we have learned to expect from the genetic control of protein synthesis; it tells us only that primary protein structure is related directly to gene structure—presumably to the linear sequence of nucleotides that make up the DNA molecule. Since the synthesis of hemoglobin occurs in the ribosomes of the cytoplasm (Schweet et al., 1958) rather than on the chromosomes in the nucleus, the code for hemoglobin synthesis must have been transferred from the DNA to the ribosome, perhaps by a messenger RNA. Within the ribosome, protein synthesis is probably organized as a cooperative enterprise in which ribonucleic acid and protein both perform essential functions. Hemoglobin synthesis thus sets forth in broad outline the essential features, as we presently understand them, of the genetic control of specific protein synthesis.

Much more interesting and significant from the viewpoint of developmental biology is the fact that the hemoglobin of the fetus and newborn is different from that of the adult human. This fetal hemoglobin, like adult hemoglobin, is composed of four polypeptide chains, but the two β chains of adult hemoglobin have been replaced by two identical γ chains, which differ significantly in amino acid composition from the β chains (Schroeder et al., 1961). This difference in primary protein structure implies that a different nonallelic gene is responsible for the synthesis of the γ polypeptides. Genetic evidence confirms this implication. A fundamental principle of development is thus clearly shown by this example, namely, that the synthesis of a specific protein in the appropriately differentiated cells of a metazoan is dependent upon the activation of the corresponding gene and not simply upon the mere presence of the gene. During the first stages of embryonic development, before the differentiation of erythrocytes, none of the genes for hemoglobin synthesis appear to be active though they are surely present in the cells of the embryo. Then, as erythrocytes begin to develop, the α and γ genes are activated and fetal hemoglobin is synthesized. Later the γ gene is suppressed and the β gene activated, although for a time both function together, producing mixtures of adult and fetal hemoglobin.

With this example of hemoglobin in mind, we may now state the thesis of this lecture. During embryonic development the cells of an organism undergo a progressive differentiation during which they acquire a characteristic repertory of proteins, largely enzymatic, which are the products of genic activity. The appearance of a new protein in a cell is an indication that the gene for that protein has been activated in that cell. By the same token the disappearance of a protein indicates that the corresponding gene has been inhibited.

Thus cell differentiation becomes simply an expression of an antecedent pattern of gene activation or inhibition. The central problem of developmental biology then becomes the problem of the control of gene function.

Some investigators would state the problem differently; they favor the alternative hypothesis that pictures all the genes as functioning continuously. The effect, or consequences, of gene activity in this view would be modified by the specific environment in which the gene was functioning. Thus different cells with identical arrays of functioning genes could be produced. Although this view is logically adequate, it confronts the cell with particularly difficult and complex problems of regulating metabolism. The simplest and most efficient control mechanism should, it seems, be directed toward turning the gene on and off rather than toward controlling the products of gene activity, for these products would be far more numerous and prone to migrate to many different positions in the cell.

Melanogenesis

Whatever the correct hypothesis of cellular differentiation may be, the various developmental and genetic aspects of the problem of cell differentiation are well illustrated during the development of the mammalian melanocyte (Markert and Silvers, 1955). The differentiation of an embryonic cell into a melanocyte requires a highly organized sequence of environmental stimuli. These successive stimuli transform the melanoblast into a cell with distinctive biochemical and morphological characteristics. According to the preferred view, these characteristics soon result in activation of the gene for tyrosinase synthesis. The enzyme tyrosinase is not only characteristic of melanocytes but is indispensable for the terminal stage of differentiation which results in the synthesis of melanin. Thus the origin of this specific protein (tyrosinase) is dependent upon two distinct but related processes—the transformation of an embryonic cell into a mature melanocyte and then the activation of the specific gene for tyrosinase synthesis. These sequential steps in melanocyte differentiation are normally controlled by many different genes. We can recognize several of these steps because mutant genes affecting them have been discovered in a variety of animals, particularly mice. The abnormal effects of these mutant genes enable us to construct a partial picture of the normal activity of several genes involved in the differentiation of mouse melanocytes.

Two quite different kinds of melanocytes—epithelial and dendritic

—occur in mice and in other vertebrates as well. Both types must have arisen from a common precursor cell at some early stage in embryonic development. The progeny of this cell later separated into two groups, each proceeding down its own distinct pathway of embryonic development. One group gave rise to the outer wall of the optic cup, which transforms into the pigmented retina. These are the epithelial melanocytes. The other group gave rise to the neural crest, from which melanoblasts migrate to many different tissues, such as those of the hair follicles. In these terminal tissue sites they receive differentiating stimuli from the surrounding cells and are thus enabled to complete their transformation into mature dendritic melanocytes.

Gene-controlled metabolites from both the embryonic melanoblast and from the cells of the tissue environment are jointly required to bring a melanoblast to full maturity as a pigmented melanocyte. We might reasonably expect, therefore, that some genes affecting melanocyte differentiation would operate primarily within the melanoblast while others would be expressed in the cells of the tissue environment. This expectation is realized.

Since epithelial and dendritic melanocytes reside in quite different tissue environments, they commonly respond differentially to those mutant genes that operate through the tissue environment. For example, in mice carrying the gene for yellow hair, the dendritic melanocytes are induced to synthesize yellow pigment when in the hair follicles, but the epithelial melanocytes of the eye continue to make black pigment. The early steps in the differentiation of the two types of melanocyte can also be controlled by different genes.

Black-eyed white mice carry a gene that prevents the differentiation of all dendritic melanocytes while permitting the differentiation of the pigmented retina. Similarly the genes for eyelessness prevent the appearance of epithelial melanocytes while leaving unaffected the development of dendritic melanocytes. However, many genes affect both types of melanocyte, particularly those genes which control the terminal stages of pigment formation. The albino gene, for example, suppresses melanin pigment formation in all melanocytes by preventing the synthesis of tyrosinase. Likewise the black or brown color of pigmented animals is determined by the same gene equally in both kinds of melanocytes.

Melanin pigment is produced in a complex cell organelle called a melanosome (Moyer, 1961). The development of melanosomes is an indispensable condition for the normal synthesis of the specific protein, tyrosinase. Apparently the protein fibrils of the melanosome are the exclusive sites of tyrosinase activity. Figure 1 shows a sequence

of developmental stages in the synthesis of a melanosome. These fibrillar structures have been found only in melanoblasts and presumably represent the response of the melanoblast to differentiating stimuli. Once the melanosome has been formed, it normally becomes pigmented by the deposition of melanin on the fibrils (Fig. 3) until the protein structure of the granule is completely obscured by melanin. However, the albino gene, which prevents tyrosinase synthesis (and thus melanin formation), does not prevent the appearance of unpigmented melanosomes (Fig. 4). The structural details of the melanosome are also under genetic control. The pink-eye gene brings about a disorganization of the protein fibrils of the melanosome (cf. Figs. 2 and 3) even though tyrosinase synthesis and pigmentation proceed in an essentially normal fashion. These observations on the ultrastructure of melanosomes and the differentiation of melanocytes lead to the general conclusion that the synthesis of a specific protein during development requires a specific gene *and* the proper state of cell differentiation in order for the gene to be activated.

We have strong reasons for believing that the gene for tyrosinase synthesis is present in all the cells of a mouse, not just in the melanocytes. However, the gene reveals itself only in melanocytes by the production of melanin. Recently, however (Peck, 1961), it has been possible to activate the gene for tyrosinase synthesis in cells that normally never give any evidence of possessing this gene in an active form.

The neural retina shares a common ancestry with the pigmented retina, but the cells of the neural retina differentiate into sensory and nerve cells, presumably because of selective environmental stimuli that do not reach the pigmented retina. However, by culturing the cells of the chick neural retina *in vitro* in isolation from one another, they can all be induced to manufacture pigment. This pigment passes the tests for melanin and is presumably synthesized through the catalytic activity of tyrosinase. These results obtained by Peck can best be interpreted as an abnormal activation of the gene for tyrosinase in cells that normally never have this gene in an active form. The word "normally" must be emphasized here, because an inherited disease of human beings known as *retinitis pigmentosa* is characterized by the formation of melanin-like pigment in the neural retina.

FIG. 1. Electron micrograph showing early stages in melanosome formation in the pigmented retina of a pink-eyed black embryonic mouse. Fixed in osmic acid, embedded in methacrylate, stained with uranyl acetate. Numbers indicate successive stages in melanosome development. *R* indicates ribosomes. *Mi* mitochondrion. (Courtesy of Dr. Frank H. Moyer.)

This disease leads to blindness and apparently represents a condition in which the gene for tyrosinase was abnormally provoked to activity.

The chick neural retina, which was cultured *in vitro*, had of course not been exposed to those stimuli that bring about melanosome formation in melanocytes. Nevertheless, pigment granules superficially resembling those of melanocytes, as viewed with the light microscope, were formed. On examination with the electron microscope, however, these granules proved to be quite different in basic structure. Evidently the formation of pigmented melanosomes requires the integrated activity of several genes in addition to the one for tyrosinase. Any one of these genes functioning at the "wrong" time or in the "wrong" cell would probably lead to abnormal cell differentiation. Perhaps the most instructive inference that can be drawn from these experiments is that inactive genes in cells are not lost and may be abnormally aroused to activity by suitable stimuli that lie outside the normal experience of the cell.

Chromosomal differentiation

If we are to understand the nature of cell differentiation, in fact, of all embryonic development, it is essential to elucidate the mechanism by which primary gene function is controlled. The previous observations on the role of genes in melanocyte differentiation and on the synthesis of tyrosinase have served to emphasize the critical dependence of gene activation on the physical-chemical composition of the cell. In view of the crucial importance of this subject, it is surprising that so few investigators have turned their attention to it. The problem is very difficult, of course, but some promising avenues of investigation have been opened. Before discussing these, it is desirable to emphasize that the genes are part of a chromosome in higher organisms—and the chromosomes are the most complicated organelles of the cell. In addition to desoxyribonucleic acid (DNA), the chromosomes contain protein in at least two general varieties, ribonucleic acid (RNA), polyamines, lipids, and perhaps other substances. From studies of viral genes (Cohen, 1957) we infer that DNA can function even though it is not integrated into complicated structures like chromosomes. What then is the function of these accessory substances

FIG. 2. Electron micrograph of pigmented retina of pink-eyed black mouse 13 days *post partum*. Fixed in osmic acid, embedded in EPON-812, stained with uranyl acetate. Note the disorganized arrangement of the fibrils and the irregular shape of the melanosomes. This peculiar melanosome structure is an effect of the gene for pink-eye. (Courtesy of Dr. Frank H. Moyer.)

of the metazoan chromosome? A plausible answer is that their role is primarily to regulate the function of the DNA—a regulation which becomes of critical importance in all organisms composed of many diverse types of cells all equipped with the same set of genes.

FIG. 3. Electron micrograph of pigmented retina of a C57 brown mouse 18.5 days *in utero*. Fixed in osmic acid, embedded in methacrylate, and stained with lead subacetate. Four stages in melanosome development are shown. Note the longitudinal (4) and cross sectional (3) cuts through the melanosomes. (Courtesy of Dr. Frank H. Moyer.)

Direct cytochemical investigations of chromosomes is one promising area of investigation. Recently Bloch and Hew (1960) have shown, in a cytochemical investigation of snail spermatogenesis, that the spermatocyte chromosomes contain histones with staining properties indistinguishable from other somatic tissues. As the differentiation of sperm proceeds, the somatic histone (lysine-rich) is replaced by

an arginine-rich histone and later, in the sperm, by protamine. After fertilization, the protamine of sperm chromosomes disappears and is replaced by faintly basic histones which differ from adult histones in their inability to bind fast green, and from protamines, both by their inability to bind eosin and also by their weakly positive reaction

FIG. 4. Electron micrograph of "pigmented" retina of Swiss albino mouse, 15 days *post partum*. Fixed in osmic acid embedded in EPON-812, stained with uranyl acetate. Note the two nonpigmented melanosomes arrested at successive stages in development. No pigmentation of these melanosomes in albinos has been observed. (Courtesy of Dr. Frank H. Moyer.)

with bromphenol blue. These "cleavage" histones are found in both the male and female pronuclei, the early polar body chromosomes, and the nuclei of the cleaving egg and morula stages. During gastrulation the histone becomes indistinguishable from that of adult somatic cells. Before gastrulation much evidence suggests that the chromosomes are nonfunctional (Moore, 1955; Briggs et al., 1951).

Further studies show that the cytochemical properties of nucleohistone in proliferating tissues differ from those found in physiologically active cells. Both the stainable phosphate groups of DNA and the stainable basic groups of histone decrease in nonreplicating cells, suggesting that these groups are masked by residual protein, which is known to increase in physiologically active cells (Alfert, 1958). In addition to pointing out the changing chemical composition of chromosomes during development, these and other studies also emphasize that chromosome replication is more than just DNA replication. The minimum replicating unit is nucleohistone (Bloch and Godman, 1955), but other constituents of chromosomes are also duplicated during mitosis, though less completely and less consistently. Presumably the variable constituents of chromosomes are the leading candidates for the role of differential gene activators or inhibitors, and thus we must look to protein or RNA. These constituents can be shown to vary in different cells at various stages in differentiation, but at this gross level of observation no correlation with the function of specific genes is possible. Among animals only the chromosomes of *Drosophila* are well enough known to offer hope of identifying chemical changes at known genetic loci. In addition to the characteristic banded pattern of the giant chromosomes of Diptera, it has been known for many years that these chromosomes also exhibit localized areas of enlargement, or puffs (Beermann, 1959). These puffs (Fig. 5) are distributed along the chromosomes in distinctive patterns that are characteristic for the cell type and for the stage of its differentiation (Becker, 1959). They are commonly thought to indicate regions of heightened gene activity, but at any rate they are a clear indication of chromosome differentiation and may be related to differential gene function. Kroeger (1960) in a recent study sought to alter the puffing pattern experimentally. He simply placed nuclei from *Drosophila* salivary gland cells into cytoplasm taken from young embryos of two different ages. After incubation in this younger, less differentiated cytoplasm, the chromosomes of the nuclei were examined for their puffing pattern. His results (Fig. 6) may be summed up succinctly by stating that the puffing patterns changed and were characteristic for each cytoplasm. Some puffs regressed and new ones appeared. Thus the chemical environment of the chromosome elicits changes in chromosome morphology. The effective substances in *Drosophila* embryo cytoplasm are still unknown, but in another organism—the dipteran, *Chironomus tentans*—a specific substance has been shown to change chromosome puffing patterns. Clever and Karlson (1960) injected ecdyson—a hormone obtained from the pro-

FIG. 5. Schematic representation of the end of chromosome arm III in successive stages in development of the third instar larva and prepupa of *Drosophila melanogaster*. Hours before and after the onset of puparium formation are indicated. Note the gradual change in puffing patterns. The puffs are numbered according to the conventional designations of corresponding chromosome regions. (Figure from Becker, 1959.)

thoracic glands of insects—into the last instar larva of a chironomid. The hormone elicited one new puff and induced the disappearance of an old puff on one of the chromosomes. These changes normally occur when the insect metamorphoses. Thus the hormone ecdyson, which stimulated metamorphosis, possibly achieves its effects, at least in part, by inducing chromosome differentiation which may manifest changes in gene function.

FIG. 6. Comparison of regions of salivary-gland chromosomes of *Drosophila busckii* at two stages in development and after incubation for 3 hours in cytoplasm from preblastoderm and blastoderm embryos. Note the changes in puff development in regions 3 and 6. (Figure from Kroeger, 1960.)

We saw previously, during the discussion of tyrosinase synthesis, that the response of a gene to activating stimuli is highly specific for the state of cell differentiation in mice, and probably in all organisms. Apparently substances from the cytoplasm enter the nucleus and induce changes in the structure and function of receptive chromosomes, as shown by the work on dipteran chromosomes. Except for ecdyson, the nature of these substances is completely unknown, but two leading candidates are proteins and ribonucleic acid or some combination between them.

Nuclear differentiation

Chemically induced chromosomal changes also seem to provide an explanation for the results obtained by Moore (1960) in his transplantation of nuclei between two species of frog, *Rana sylvatica* and *R. pipiens*. *Rana sylvatica* eggs were fertilized by *R. pipiens* sperm and then the egg nucleus removed before fusion with the sperm nucleus could occur. Such *R. sylvatica* eggs equipped with haploid *R. pipiens* sperm developed into blastulas, but then development stopped. By the time of developmental arrest, the original sperm nucleus would have divided about 12 to 14 times and the arrested blastula would be composed of approximately 25,000 cells. Nuclei from such arrested blastulas may be transferred again to eggs of the foreign species. The same pattern of cell division and arrest of development at blastula will again occur. Since hybrid amphibian embryos commonly cease development as blastulas, this result is not surprising but it does demonstrate a nucleo-cytoplasmic interdependence. Most suggestive, however, is the discovery that nuclei after a period of residence in foreign cytoplasm were not immediately able to support normal development even when retransplanted to their species-specific cytoplasm. Serial transplantation of these nuclei for six generations produced only arrested blastulas just like those produced by the original hybrid combination. Multiplication of the nuclei in foreign cytoplasm had evidently produced a persistent change, probably in the chromosomes.

An attractive interpretation of these results is that replication of chromosomes in foreign cytoplasm leads to the attachment of foreign substances to the chromosomes, thus preventing their normal differentiation and inhibiting their heterosynthetic functions. It is important to remember that chromosome replication during nuclear division involves more than just the DNA. Many replications might be necessary, therefore, in order for a chromosome to lose a component acquired during its residence in a chemically foreign environment.

These observations on the failure of chromosomes to promote normal development after they had multiplied in foreign cytoplasm prompted us to test the hypothesis that cellular differentiation reflects an antecedent chemical differentiation of the chromosomes. Our experimental design was simple to the point of being naïve. Various macromolecular fractions were prepared from the nuclei of adult frog-liver cells. These fractions were dialyzed against a physiological saline solution and then injected into fertilized eggs of the same

species. Control eggs, injected with the saline solution alone, developed normally except for occasional abnormalities due to physical injuries produced by the injection procedure. However, eggs injected with several of the nuclear fractions exhibited a consistent developmental arrest at the beginning of gastrulation. Cell division stopped in these arrested blastulas, but otherwise they appeared normal. Since

FIG. 7. Diagram of the procedure used in testing the developmental effects of injected protein fractions extracted from adult frog-liver nuclei. Nuclei derived from injected eggs were serially transplanted for seven generations, but each generation arrested at the beginning of gastrulation.

several nuclear fractions produced these effects, we are not sure of the identity of the effective substances, although they appear to be non-dialyzable proteins that are inactivated by prolonged heating. The most potent fraction is still active when diluted, so that its protein content is as little as 0.02 millimicrograms of protein per 0.1 micro-liter—the volume of the injected solution. Further dilution rendered this solution ineffective. During cell division the chromosomes of injected eggs should have been freely accessible to any injected substances. The very small quantity of injected material suggests a highly specific effect. Moreover, when nuclei from arrested blastulas were serially transplanted for seven generations (Fig. 7) to enucleated eggs, the embryos that resulted invariably stopped developing before the onset of gastrulation (Markert and Ursprung, 1962). Since the effect was very characteristic and persistent through many nuclear divisions, it seems reasonable to place the primary responsibility on the chromosomes. The behavior of these nuclei closely resembles that of the transplanted nuclei which multiplied in foreign cytoplasm (Moore, 1960), and indeed the chromosomes of the injected eggs were initially multiplying in “foreign” cytoplasm—a cytoplasm changed by virtue of the injected substances from the adult liver nuclei.

Isozymes

The discussion so far has hopefully revealed the importance of the cytoplasm in activating gene function, an event that is manifested in the synthesis of a specific protein. Is this then an inclusive description of the mechanisms underlying the synthesis of specific proteins? Probably not. Embryologists have often loosely spoken of molecular differentiation when they really meant the differentiation of cells as evidenced by the appearance of new proteins. However, we have recently become aware of a phenomenon that may be a valid example of a true molecular differentiation. Many investigators have shown that single enzymes commonly exist in multiple molecular forms or *isozymes* within the tissues or cells of a single organism (Markert and Möller, 1959). The isozymes of a single enzyme have very similar catalytic properties but can nevertheless be distinguished from one another by their somewhat different physical properties. Usually chromatographic or electrophoretic techniques serve to separate isozymes from one another. These isozymes are not artifacts of analysis but exhibit characteristic patterns of distribution in each tissue (Fig. 8). Moreover, the tissue patterns are species specific (Fig. 9). The characteristically different isozyme patterns of adult tissues must have

FIG. 8. Photograph of zymograms of lactate dehydrogenase isozymes from eight tissues of the mouse. Five isozymes are evident, most of which are found in most tissues. However, the relative concentration of the various isozymes differs markedly in the various tissues.

FIG. 9. Diagrammatic representation of zymograms of lactate dehydrogenase isozymes obtained from heart muscle of four species. Each species exhibits a characteristic pattern of isozymes.

arisen during the course of embryonic development, and direct analysis shows this to be so. The adult pattern is the end product of a long sequence of gradual changes during ontogeny.

The existence of isozymes poses important problems for the origin of specific proteins. Each isozyme may be the responsibility of one or two genes, or the several isozymes may represent modifications of a single gene product. The latter case would represent molecular differentiation. A fully satisfactory explanation for the origin of isozymes can scarcely be made until isozymic differences are related to molecular structure. At least five kinds of molecular changes may be invoked to produce different though closely related isozymes: Small changes in amino acid sequence, amidation of carboxyl groups, conjugation with small molecules, polymerization of different subunits, and changes in conformation achieved by folding the same primary structure in different ways. All these possibilities have served to explain, with varying degrees of experimental support, the existence of numerous enzymes in isozymic forms. Polygenic control will clearly be indicated wherever isozymes are found to differ in primary structure. Other distinctions are more plausibly attributed to modifications imposed secondarily upon a single basic gene product.

The esterase enzymes have been extensively investigated with reference to both ontogenetic (Markert and Hunter, 1959) and genetic control (Wright, 1961). The esterases compose a large family of closely related but distinguishable enzymes. Each tissue of most animals that have been examined contains a characteristic repertory of these esterases in distinctive patterns of concentration. The patterns arise gradually during embryonic development. However, these different esterases are not all isozymes of one another. Wright (1961) has shown in an investigation of *Drosophila* enzymes that an esterase (Fig. 10) is present in position 4 in one strain and in position 5 in another strain. Hybrids between these strains have esterases in both positions. The genetic evidence strongly suggests that each of these esterases is controlled by an allele of a single gene; the heterozygote would then possess both esterases, as is apparent in the offspring from a cross between the two strains (Fig. 10).

In Wright's investigation no isozymes of esterase have yet been identified, although the genetic control of specific protein structure is clearly demonstrated. However, Allen (1960) has shown in *Tetrahymena* that the final structure of the molecule can also be the responsibility of cytoplasmic mechanisms emerging during the course of cellular differentiation. Allen's investigation demonstrated that allelic genes control alternate groups of four esterases in *Tetrahymena*.

The members of each group are distinguished from the members of the alternate group by a difference in charge, which can plausibly be attributed to the substitution of a single amino acid in the primary protein product of the gene. This product would then be modified into four distinguishable esterases that are inherited as a group.

In a similar analysis of alkaline phosphatases in *Escherichia coli*, Bach et al. (1961) found five electrophoretically distinguishable forms

FIG. 10. Zymogram of esterases in *Drosophila*. Homogenates for each insertion were made from five adult male flies less than 24 hours old. Electrophoresis through the starch gel occurred for 6 hours at a voltage drop of 4 v/cm and a pH of 8.7. Starch was made up in Tris buffer, 0.5 M. Alpha-naphthyl acetate was used as substrate and fast blue BB as dye coupler. *Drosophila* strain *Am* is highly inbred and shows an esterase at position 5 but not at position 4. Strain *CH* is not homozygous but contains flies with esterases at positions 4 or 5 or both. When a strain *CH* fly (with an esterase at position 4 but not 5) is crossed to strain *Am*, the F₁ offspring all show esterases at both positions.

of the enzyme. Several mutant strains continued to produce the same number of phosphatases but all the phosphatases were shifted in their electrophoretic mobility to the same extent. These results are consistent with the assumption that all five phosphatases are products of the same gene. Mutations in this gene lead to the production of an altered form of a single basic precursor molecule that is then secondarily modified into the five isozymes of alkaline phosphatase. However, the quantitative proportions in which these isozymes are synthesized are changed by growth in different media, and in effect, therefore, these different patterns represent different states of cell differentiation.

Extensive investigations (Markert and Appella, 1961; Appella and Markert, 1961) on the physicochemical properties of separated and recrystallized LDH isozymes from beef tissues have demonstrated that each of these isozymes is composed of four dissociable polypeptide chains. These chains, like those of hemoglobin, may be encoded by two different genes, thus producing two kinds of polypeptides. Assortment of these two polypeptides in all possible combinations of four would yield the five LDH isozymes which are found in mouse (Fig. 8) and beef tissues. These isozymes would then be under polygenic control although the cytoplasm would doubtless play a role in assembling the finished tetramers in the proportions peculiar to each cell.

However, it is certain that various isozymes may be converted into alternate isozymes by mechanisms resident in the cytoplasm of the cell. Kaji and collaborators (1961) have shown that one of the two major isozymes of yeast hexokinase may be converted into the other by treatment with trypsin or chymotrypsin in the presence of glucose. Presumably this conversion could also occur in the cell, although a fixed ratio of the two isozymes is normally maintained within the yeast cell.

The origin of specific proteins through nucleocytoplasmic mechanisms is also indicated by the work of Keck (1961) on two species of the giant single-celled alga, *Acetabularia* (*Acicularia schenckii* and *Acetabularia mediterranea*). By ingenious transplantation techniques, Keck combined the nuclei and cytoplasm of the two species in various proportions. Each species was shown to have a distinct acid phosphatase which could be readily identified by staining techniques after electrophoresis of a cell homogenate in a starch gel. The electrophoretic mobility of the *med* phosphatase was considerably greater than that of the *acic* enzyme. Mixtures of homogenates from the two species showed two distinct bands after electrophoresis. In the graft combinations shown in Fig. 11, Keck discovered that the *med* enzyme

was, so to speak, dominant. That is, either the nucleus or the cytoplasm of *med* plants sufficed to establish the *med* phosphatase as exclusively characteristic of the hybrid plant. The *acic* enzyme rapidly disappeared from the graft combination at a rate far in excess of its

FIG. 11. Diagram of zymogram patterns of acid phosphatase in two species of *Acetabularia*. Each species contains a single acid phosphatase with a characteristic electrophoretic mobility in starch gel. Strips 1 and 6 show the pattern obtained from mixed homogenates of the two species. Strip 2 depicts the conversion of the *acic* enzyme to the *med* type when a *med* nucleus is substituted for the *acic* nucleus. Strip 3 shows that the *med* enzyme persists even after the *med* nucleus is replaced by an *acic* nucleus. Strip 4 shows that only *med* enzyme is found in hybrid cytoplasm regenerated in the presence of one nucleus from each species. Strip 5 shows the intermediate phosphatase band frequently observed when two *acic* nuclei and one *med* nucleus collaborate in the production of new cytoplasm. This band apparently represents an intermediate stage in the conversion of *acic* enzyme to *med* enzyme.

normal diminution in enucleated *acic* plants. This rapid disappearance suggests that some mechanism in the *med* cytoplasm converts the *acic* enzyme to the *med* type. This mechanism must be initiated by the nucleus because the *med* nucleus, by itself, is sufficient to bring about the conversion of *acic* phosphatase to the *med* type. An intermediate step in this conversion is visible after electrophoresis of graft com-

binations containing two *acic* nuclei and one *med* nucleus joined in a common cytoplasm. In this case, the productivity of the two *acic* nuclei is apparently so great that the *med* cytoplasm cannot complete the conversion of the *acic* enzyme, hence an intermediate stage in the conversion accumulates and becomes visible.

The existence of numerous enzymes in isozymic forms requires us to seek an explanation in terms of physiological utility to the organism. One of the most attractive explanations for isozymes arises out of the work of Stadtman and his collaborators (1961) on the enzyme aspartokinase in *Escherichia coli*. This enzyme catalyzes the phosphorylation of aspartate by ATP and the resulting aspartyl phosphate serves as a precursor in the synthesis of the two amino acids, lysine and threonine. Both of these amino acids inhibit aspartokinase activity in extracts of *E. coli*. The inhibitions are additive and independent, suggesting that two different forms of aspartokinase are present. And indeed, by ammonium sulfate precipitation two different isozymes were separated; one was inhibited by lysine, the other by threonine. Moreover, when *E. coli* is grown in the presence of 10 mM L-lysine, the synthesis of the lysine-sensitive isozyme is completely repressed; thus the amount of the isozyme synthesized is adjusted to the physiological requirements of the organism.

Although both isozymes of aspartokinase carry out the same catalytic activity, the results of Stadtman et al. suggest that the isozymes may be located in different metabolic pathways, one leading to the production of threonine, the other to lysine. The behavior of the isozymes of aspartokinase suggests the following general hypothesis. Isozymes of any given enzyme carry out the same basic catalytic activity, but as parts of different metabolic pathways. These metabolic pathways are either located in different parts of a cell or give rise to different end products. The gradual emergence during embryonic development of specific isozymic patterns would then reflect the origin of distinct though related metabolic pathways.

In summary, the origin of specific proteins during development requires two basic interacting components—the genes and the cytoplasm. Genetic and biochemical evidence clearly demonstrates the role of genes in specifying the primary structure of proteins, but the existence of isozymes also suggests a role for the cytoplasm in modifying at least the finer aspects of protein structure. Both the genes and the cytoplasm are brought to a state of active function by as yet unknown mechanisms of cellular differentiation. The most fundamental expression of this differentiation is the appearance of a new specific protein, which must be regarded as a product involving the collabo-

ration of genes and cytoplasm. During development, genes and cytoplasm interact with one another to produce new patterns of active genes, which lead to the synthesis of new varieties of proteins and thus to an altered cytoplasm. This cyclic interaction makes possible progressive cell differentiation through the synthesis of new proteins and constitutes the foundation of all embryonic development.

References

- Alfert, M. (1958), Variations in cytochemical properties of cell nuclei, *Exp. Cell Research*, Suppl. 6:227-235.
- Allen, Sally Lyman (1960), Inherited variations in the esterases of *Tetrahymena*, *Genetics*, 45:1051-1070.
- Appella, E., and C. L. Markert (1961), Dissociation of lactate dehydrogenase into subunits with guanidine hydrochloride, *Biochem. Biophys. Res. Commun.*, 6:171-176.
- Bach, Marilyn L., E. R. Signer, C. Levinthal, and I. W. Sizer (1961), The electrophoretic patterns of alkaline phosphatase from various *E. coli* mutants, *Federation Proc.*, 20:255.
- Becker, H. J. (1959), Die Puffs der Speicheldrüsenchromosomen von *Drosophila melanogaster*. I. Beobachtungen zum Verhalten des Puffmusters im Normalstamm und bei zwei Mutanten, *giant* und *lethal-giant larvae*, *Chromosoma*, 10: 654-678.
- Beermann, Wolfgang (1959), Chromosomal differentiation in insects, in *Developmental Cytology*, ed. by D. Rudnick, The Ronald Press Company, New York, pp. 83-103.
- Bloch, D. P., and G. C. Godman (1955), A microphotometric study of syntheses of deoxyribonucleic acid and nuclear histone, *J. Biophys. Biochem. Cytol.*, 1:17-28.
- , and H. Y. C. Hew (1960), Changes in nuclear histones during fertilization and early embryonic development in the pulmonate snail, *Helix aspersa*, *J. Biophys. Biochem. Cytol.*, 8:69-81.
- Briggs, Robert, E. U. Green, and T. J. King (1951), An investigation of the capacity for cleavage and differentiation in *Rana pipiens* eggs lacking "functional" chromosomes, *J. Exp. Zool.*, 116:455-500.
- Clever, U., and P. Karlson (1960), Induktion von Puff-Veränderungen in den Speicheldrüsenchromosomen von *Chironomus tentans* durch Ecdyson, *Exp. Cell Research*, 20:623-626.
- Cohen, Seymour S. (1957), The biosynthesis of nucleic acids in some microbial systems, in *The Chemical Basis of Heredity*, ed. by W. D. McElroy and B. Glass, Johns Hopkins Press, Baltimore, pp. 651-685.
- Conference on Hemoglobin (1958), Publication 557, National Academy of Sciences, National Research Council, Washington, D.C.
- Kaji, A., K. A. Trayser, and S. P. Colowick (1961), Multiple forms of yeast hexokinase, *Ann. N. Y. Acad. Sci.*, 94:798-811.
- Keck, Konrad (1961), Nuclear and cytoplasmic factors determining species specificity of enzyme proteins in *Acetabularia*, *Ann. N. Y. Acad. Sci.*, 94:741-752.
- Kroeger, H. (1960), The induction of new puffing patterns by transplantation of

- salivary gland nuclei into egg cytoplasm of *Drosophila*. *Chromosoma*, **11**:129-145.
- Markert, C. L., and E. Appella (1961). Physicochemical nature of isozymes. *Ann. N. Y. Acad. Sci.*, **94**:678-690.
- , and Robert L. Hunter (1959). The distribution of esterases in mouse tissues. *J. Histochem. Cytochem.*, **7**:42-49.
- , and Freddy Møller (1959). Multiple forms of enzymes: Tissue, ontogenetic, and species specific patterns. *Proc. Nat. Acad. Sci. U.S.*, **45**:753-763.
- , and Willys K. Silvers (1955). The effects of genotype and cell environment on melanoblast differentiation in the house mouse. *Genetics*, **41**:429-450.
- , and H. Ursprung (1962). Alteration of zygote chromosomes in *Rana pipiens* by injection of proteins from adult liver nuclei. *Amer. Zool.*, **2**:428.
- Moore, J. A. (1960). Serial back-transfers of nuclei in experiments involving two species of frogs. *Developmental Biology*, **2**:535-550.
- (1955). Abnormal combinations of nuclear and cytoplasmic systems in frogs and toads. *Advances in Genetics*, **7**:139-182.
- Moyer, Frank H. (1961). The developmental genetics of melanocyte differentiation analyzed with the electron microscope. Ph.D. Thesis. Johns Hopkins University.
- Peck, David (1961). The role of tissue organization in the cytodifferentiation and metabolism of the embryonic chick neural retina. Ph.D. Thesis. Johns Hopkins University.
- Schroeder, W. A., Richard T. Jones, J. Roger Shelton, Joan Balog Shelton, Jean Cormick, and Kathleen McCalla (1961). A partial sequence of the amino acid residues in the γ chain of human hemoglobin F. *Proc. Nat. Acad. Sci. U.S.*, **47**:811-818.
- Schweet, R., H. Lamfrom, and E. Allen (1958). The synthesis of hemoglobin in a cell-free system. *Proc. Nat. Acad. Sci. U.S.*, **44**:1029-1035.
- Stadtman, E. R., G. N. Cohen, Gisele LeBras, and Huguette de Robichon-Szulmajster (1961). Feed-back inhibition and repression of aspartokinase activity in *Escherichia coli* and *Saccharomyces cerevisiae*. *J. Biol. Chem.*, **236**:2033-2038.
- Wright, T. R. F. (1961). The genetic control of an esterase in *Drosophila melanogaster*. *Amer. Zool.*, **1**:476.

Diversity at the Subcellular Level and Its Significance

Keith R. Porter

Biological Laboratories
Harvard University, Cambridge, Massachusetts

Diversity in the microscopic anatomy of cells requires no exposition; it is illustrated in numerous histological and cytological treatises. There are in nature essentially as many distinct and separate forms of cells as there are tissues they comprise. Some kinds, especially cells performing similar functions, are remarkably alike in different animals or plants, but enough difference is usually apparent to permit their separation. That equal or greater variations would be apparent at the "subcellular," or "submicroscopic" level, seemed probable but a more thorough investigation of the question was delayed pending the development of modern techniques for electron microscopy. Now these procedures have been quite widely applied and results of recent observations have not been surprising in the degree of diversity they have depicted; it is, in other words, not greater than one might have expected. Diversity at submicroscopic levels naturally expands on the diversity evident at the microscopic because compartments of the cells which formerly appeared structureless now are described in

terms of fibrils, particles, and membranes of diverse organizations and dimensions.

More surprising perhaps than the variations in form revealed by subcellular explorations is the similarity displayed by structures formerly thought to represent diverse units of the cell. Thus it is that the modern cell biologist can relate structures formerly not apparently related. Generalizations encompassing the whole of metazoan cell structure, and including protozoans in some instances, may now be formulated, thus making much easier the task of classifying the new fine-structure information. Simultaneously, there is provided through comparative cytology a clearer understanding of the relationships and functions of several of the newly defined systems.

It will be the purpose of this brief chapter to illustrate the diversity in form displayed by some of the subcellular components of cells, and to point out the pervading similarities and structural associations which relate these components despite their apparent differences.

Diversity in mitochondria

Variation in form at the subcellular level could, of course, be illustrated by comparing representatives of a number of intracellular structures. These might be fibrous, granular, or membranous in their basic structure. But possibly none is better for the purpose than mitochondria. These, as was well known from light microscopy, are universally occurring organelles of the cell. They contain enzymes for the complex series of reactions involved in oxidative phosphorylation in the production of energy-rich ATP. The electron microscope observations of Palade (1953) and of Sjöstrand (1953) about ten years ago on these organelles established the existence of a fairly complicated fine structure, which is uniform among the population in a single cell type but varies in different types (Figs. 1A to 1C). It was apparent, however, from these early studies that a basic architecture was common to all mitochondria (Palade, 1956a). Thus each of these organelles in its fully differentiated form possesses an external limiting membrane; inside this, and generally separated from it by space of about 8 millimicrons, there is an internal membrane which through various infoldings forms shelves, or cristae, as they came to be called. Observations made subsequent to this have not materially changed the picture. What has been revealed is a wide diversity of patterns displayed by the foldings or proliferations of the internal of the two membranes. These achieve a startling complexity in some cells, such as to make structural analysis very difficult (Pappas and Brandt,

1959). The central purpose of these variations seems to be to expose increasing or decreasing areas of membranes to the structureless matrix of the mitochondrion (Figs. 1A to 1C). Recently it has been found by Fernandez-Moran, in collaboration with David E. Green, that the surfaces facing the matrix are constructed of small 100 Å particles. Since biochemical studies of these particles show them to contain all the functional enzymatic components of the electron transport chain, they are regarded as the ultimate unit or "elementary particle" of mitochondrial function (Fernandez-Moran, 1960). It will be valuable to apply similar methods of observation to a variety of mitochondria showing different levels of physiological activity.

Variations in cytoplasmic fine structure

These structural variations in mitochondria—an expression of diversity in the subcellular level—are now well known to students of cell fine structure and deserve this attention only because they serve to illustrate how widely diverse morphologies can, upon analysis, be shown to represent variations on a simpler basic structure. Mitochondria, it must be admitted, present relatively minor problems of interpretation, compared with the whole cytoplasm of the cell. However, even the complexity and apparent diversity of this larger division responds to analysis if one recognizes the existence of separate and distinct systems which occur ubiquitously and can properly be compared.

Some measure of the diversity one encounters in the fine structure of cytoplasm may be gained from a limited survey of representative regions of a few diverse cell types. The range of variation shown is not descriptive of the enormous range one can find in nature, but it is illustrative without being so extreme as to make comparisons and correlations difficult. Figures 2 to 10 which follow are described and compared in the accompanying legends.

FIG. 1A. A mitochondrion and surrounding cytoplasm from the liver cell of a bat (*Myotis lucifugus*). The structure, as is characteristic of liver cell mitochondria, shows few cristae (*cr*), which are paddle-shaped. It is evident that the total membrane surface of the cristae contiguous with the matrix (*mx*) of the mitochondrion, is relatively small. This matrix, which is homogeneous and shows no evidence of patterned fine structure, makes up a large part of the mitochondrial content. The double membrane across the center of the mitochondrion is clearly an extension of the inner of the two membranes limiting the organelle, and may represent a stage in division of the mitochondrion (see Fawcett, 1955, for similar forms). The dense granules evident at various points in the structure are characteristic of many kinds of mitochondria; their function is unknown. Among mitochondria, these of liver cells display one of the simplest morphologies. Magnification: 36,000 $\times$.

FIG. 1B. A group of mitochondria (sarcosomes) as found in cardiac muscle of the bat. It is clear at a glance that these mitochondria differ from those of liver in possessing many more cristae (*cr*) and that the individual crista is much more extensive and more nearly partitions the interior of the organelle. The total membrane surface thus provided is much greater than in the case of liver mitochondria, but the volume of matrix materials is proportionately less. The structural features of these sarcosomes are characteristic of mitochondria associated with systems having large energy requirements. Presumably the production of ATP by such mitochondria is greater than in those of liver. The myofibrils are indicated at *my*, the elements of the meager sarcoplasmic reticulum at *SR*, and Z and M bands of the sarcomere by *Z* and *M*, respectively. The structure at the arrow is unidentified but appears to be continuous with the margins of one or two adjacent mitochondria. This cristae-rich variant on the morphology of the liver mitochondrion is one of the commonest. Magnification: 36,000 $\times$.

FIG. 1C. A more unusual form of mitochondrion as found in *Paramecium aurelia*. The mitochondria here are limited by the typical double membrane, but are distinctive in possessing tubular cristae. These microelements have a diameter approximately equal to the thickness of cristae encountered in other mitochondria. The number of tubular cristae is obviously great and achieves a large contact surface between membrane and matrix. The dense granules in the cytoplasm surrounding the mitochondria represent glycogen. (Courtesy of Giuseppe Millonig.) Magnification: 36,000 $\times$.

FIG. 2. A small region from the cytoplasm of a normal rat liver cell. A section of the margin of the nucleus (*N*) is at the upper left. It is separated from the cytoplasm by the nuclear envelope (*ne*), which in the profile included within the section, shows several pores (arrows). Two structurally dissimilar membrane systems dominate the fine structure of the cytoplasm. The most prominent of these appears in long, slender profiles representing vertical thin sections through large lamellar vesicles 50 to 75 millimicrons thick. The membranes limiting these structures are studded with dense particles (ribosomes) and because of these this form of the endoplasmic reticulum is referred to as the rough ER (*rer*). Continuous with this rough form (see arrow for point of continuity), is another type of reticulum composed for the most part of tubules, interconnected in a three-dimensional lattice. In liver cells this agranular or smooth form of the reticulum (*ser*) is always associated with glycogen. The matrix of the cytoplasm in which these structures are embedded appears homogeneous although in some instances, even in liver cells, there is occasional evidence of bundles of fine, filamentous elements. Where the cisternae of the rough ER are cut obliquely, the distribution of ribosomes on their surfaces is depicted. Beside ribosomes not attached to membranes, the matrix contains resolvable glycogen granules (*gl*), and sometimes ferritin particles. The liver cell, which has become a classical object for cytochemical investigations, has one of the most complex of cytoplasmic, related presumably to the complex physiological role of these cells. Nearly all the organelles and systems found in any cell are represented. It serves as a useful background for the somewhat simpler but diverse morphologies illustrated in the other figures. Magnification: 27,000 $\times$.

N

ne

rer

ql

rer

ser

FIG. 3. The basal portion of a mucous neck cell from a gastric gland of the bat. The nucleus at the lower right contains the nucleolus (*nu*) and is limited by the prominent nuclear envelope (*ne*). A mucous granule is shown at *mg* and mitochondria at *m*. The cytoplasm of this cell is characterized by a prominent development of the rough form of the ER, (*rer*). The individual elements in the section appear as slender line-limited profiles representing sections through lamellar cisternae. The surfaces of these membranes facing the matrix of the cytoplasm are densely coated with ribosomes. This micrograph serves to illustrate a variant of the ER which is common in cells dedicated to the single task of protein synthesis. The total membrane surface for the support of ribosomes is large and the space within the system for the sequestration of the synthesized material is correspondingly extensive. Magnification: 30,000 $\times$.

rer

m

mg

ne

N

nu

FIG. 4. Shows a portion of the cytoplasm of an epithelial cell in the intestine of the mosquito (*Aedes aegypti*). A part of a nucleus is shown at the upper left. The nuclear envelope is not unlike that found in other kinds of cells. The dominant element in the cytoplasm of this cell is again the rough form of the endoplasmic reticulum (*rer*). In its distribution and tendency to appear in clusters of lamellae, it resembles that found in the vertebrate liver cell. The smooth form of the reticulum (*ser*) is also represented in these cells and, as in liver cells, is closely associated with masses of glycogen (*gl*). The Golgi (*G*) is represented by separate stacks of laminate vesicles. Thus in a cell from a very different biological species, one finds a distinctive variant of the ER, but a variant which closely resembles that common to liver cells of other organisms. It will be interesting to learn whether these cells share any of the functions of liver cells. (Courtesy of Thomas F. Roth.) Magnification: 27,000 $\times$.

FIG. 5. Parts of three cells from the sporogenous tissue of the African violet (*Saintpaulia ionantha*). A nucleus (*N*) with a prominent nucleolus (*nu*) is shown at the upper left. Dense primary walls (*cu*) separate the cells and transect the image. The endoplasmic reticulum of these cells is made up characteristically of lamellar and vesicular elements. These are not organized into obvious patterns as in the case of the preceding cells, but seem instead to be more or less randomly distributed. Ribosomes are attached preferentially to the lamellar cisterna, but as is typical of undifferentiated and rapidly proliferating cells, the majority of the ribosomes (*ri*) are free in the cytoplasmic matrix. Mitochondria are indicated at *m*, and smooth ER at *ser*. Here we encounter a morphology that is quite typical of embryonic or meristematic cells: an abundance of free ribosomes and a sparse ER without obvious organization. (Courtesy of Myron C. Ledbetter.) Magnification: 35,000 $\times$.

N

nu

cw

ser

cw

m

rl

FIG. 6. A small part of the cytoplasm of an epithelial cell from a gill filament of *Fundulus heteroclitus*. The cytoplasm is obviously packed with profiles of tubules joined together in a dense three-dimensional lattice. The membranes limiting these structures are free of particles and the whole system is properly interpreted as representing the smooth ER of this cell. In one place, marked by an arrow, there is a short profile of a lamellar vesicle studded with particles. These seem to be extremely rare and are the only representatives of the rough ER found in these cells.

The epithelial cell, of which a part is shown here, has been identified as a salt-secreting unit such as are regularly found in the gills of saltwater fishes. A similar pattern of ER development is common to cells of this type where they have been studied. Chloride-secreting, parietal cells of the stomach are quite similar (Sedar, 1961; Ito, 1961). A part of an adjacent epithelial cell, shown at the lower left, illustrates the diversity which may be evident between two closely associated units. Its cytoplasmic matrix is filled with fibrillar elements (*K*), perhaps keratin in nature. Its mitochondria show fewer cristae than are found in the mitochondria of the salt cell, and the ER is represented by a relatively few profiles. Mitochondria are indicated at *m*, glycogen granules at *gl*. (Courtesy of C. William Philpott.) Magnification: 48,000 $\times$.

k

m

ql

m

ne

N

FIG. 7. A longitudinal section through a small part of a muscle fiber found in the myotome of a green frog (*Rana clamitans*) tadpole. The sarcoplasm between the myofibrils (*my*) is occupied by profiles of membrane-limited structures representing the ER of these cells. A face view of the system is shown at sarcomeres marked X; otherwise images are profiles. It is obviously differentiated with respect to the adjacent sarcomeres of the myofibrils. At the level of the Z band its continuity is interrupted by the interposition of small membrane-limited structures which appear to originate from the plasma membrane of the fiber (Smith, 1961; Porter, 1961c). The intermediate vesicle, plus the two lateral vesicles on opposite sides of the Z band, is the triadic (*tr*) structure of earlier descriptions (Porter and Palade, 1957). A few particles of glycogen are scattered among the membranes. No ribosomes are attached to the membranes, hence they are properly classified as representing the smooth endoplasmic reticulum (*ser*) of the muscle cell.

Though appearing very different from that of other cells, the cytoplasm of the muscle fiber can be interpreted as one in which the fibrous elements of the cytoplasm are prominently developed and collected into large bundles. In spite of this exaggerated development of matrix components, the endoplasmic reticulum is not lost but is similarly crowded into the space between the myofibrils, presumably to perform a function in contraction and relaxation. (Courtesy of Clara Franzini.) Magnification: 50,000 $\times$.

x

tr

tr

M

Z

FIG. 8. Parts of three cells on the epidermis of a larva of *Amblystoma punctatum*. The nucleus with a prominent envelope is shown at the lower right. The cytoplasm shows several profiles of the ER, some with particles attached (*rer*) and some separate, agranular profiles. The dominant feature of these cells is a large number of fine filaments representing the keratin component of the cells. These filaments, in places assembled in bundles (*K*), occupy the continuous or matrix phase of the cytoplasm. The arrow marks a desmosome (*D*), which represents a point of attachment between two adjacent cells; (*is*) indicates intercellular spaces.

This cell, which can be classified as a retaining or nonsecretory type of cell (Mercer, 1961) may ultimately show more and more keratin and less ER, until at the end of its differentiation it is practically free of ER except for the nuclear envelope. Magnification: 22,000 $\times$.

FIG. 9. A longitudinal section through a myelinated nerve fiber in the cortex of a rat. The matrix of the axoplasm in this and similar instances is characterized by the development of a large number of protein filaments (*pf*), which represent a differentiation of the matrix (the keratin or myosin equivalent of this cell). These are arranged with their long axes parallel to the long axis of the fiber. The ER is represented by a very few membrane-limited profiles. The myelin sheath is shown at *ms*, and mitochondria at *m*. The surrounding tissue comprises numerous non-myelinated fibers and dendritic nerve endings. (*Courtesy of Giuseppe Millonig.*) Magnification: 45,000 $\times$.

FIG. 10. Parts of two erythroblasts found in the embryonic liver of the rat. Nuclei are shown at *N*; the image of the one at the lower left includes a distinct envelope (*ne*). It is characteristic of these cells to show very few elements of the ER. The cytoplasmic matrix, on the other hand, is dominated by large numbers of ribosomes and between these an accumulation of material, probably hemoglobin, of lower density. As differentiation proceeds the ribosomes diminish in number, the reticulum disappears, and the matrix becomes uniformly gray. The nuclei are ejected. In this instance a nonfibrous component comes to dominate the cytoplasmic matrix as differentiation proceeds and the ribosomes and ER disappear. Mitochondria are indicated at *M*, the Golgi component at *G*, and centrioles at *C*. (Courtesy of *Wincenty Kilariski*.) Magnification: 20,000 $\times$.

G

C

er

m

N

er

ne

N

It should be evident from this brief review of diversity in cytoplasmic structure that certain components recur, and that they appear as variants of a basic form. One can, for example, find in almost all cells some evidence of fibrous structures in the continuous phase of the cytoplasm; ribosomes of nearly uniform size are present to greater or less extent in most instances; and finally, the cytoplasm shows multiple expressions of membrane-limited tubules and vesicles which seem to comprise a single system. In what follows this will be defined as a unit system of the cytoplasm and considered relative to its structural and functional diversity. There are certainly other components which could be used for the same purpose but which as yet have not been so completely analyzed for variations, occurrence, and function. The majority recur in all cell types and will of course come eventually to be understood. These include structures which are currently known as microbodies, lysosomes, multivesicular bodies, Golgi, and associated microvesicles.

The miscellaneous profiles of tubules and vesicles, which frequently appear to clutter up the cytoplasm, have been shown to represent parts of a continuous or intermittently continuous system, commonly referred to as the endoplasmic reticulum, or ER (Porter and Thompson, 1948; Porter, 1953). As the techniques of electron microscopy have improved over the past ten years, the image of this system has gradually emerged from the optically empty ground substance of the cell, and it is now recognized as a commonly occurring component of the cytoplasm (Hagenau, 1958; Palade and Porter, 1954; Palade, 1955; Porter, 1961a).

The unit structure of this system can be thought of as a vesicular or tubular element which appears in a wide range of forms and dimensions. It is a membrane-enclosed space, usually showing no evidence of internal structure. When these component elements, which appear in thin sections as line or membrane-limited profiles, are followed through serial sections into the depth dimension, they are found to be, in most instances, part of a complex tridimensional reticulum which extends into all parts of the cell.

A further generalization which emerges from electron microscopy of many cell types is that the nuclear envelope is morphologically similar to the unit elements of the ER (Figs. 2, 3, 5), and is found to be structurally continuous with the system at a number of points (Watson, 1955). It is proper, therefore, to include the envelope and the cytoplasmic tubules and vesicles in a single system, the endoplasmic reticulum, or ER. In support of these statements we note that the envelope consists of two membranes and an intervening space.

The outer membrane is studded with ribosomes, like rough cisternae of the ER, and the inner one is in intimate association with the peripheral chromatin of the interphase nucleus. The latter intimacy is such as to suggest an active and important physiologic exchange, or what have you between the envelope membrane and cavities and the genetic material of the nucleus. Perforations or fenestrations in the envelope connect, apparently without structural interception, the matrix of the cytoplasm and less dense nucleoplasm of the nucleus and should provide for a free exchange of fairly large molecules.

In some cells the endoplasmic reticulum is represented almost solely by the nuclear envelope, other parts having disappeared in the later stages of cytoplasmic differentiation (Fig. 10). In another and more commonly encountered structural expression or pattern, there are many lamellar sacs or cisternae in parallel array and this form corresponds to the strongly basophilic component of the cytoplasm long known as ergastoplasm. In other instances the system is represented by a three-dimensional lattice of particle-free tubules (Fig. 6). The diversity of forms ranges widely between the extremes shown, for example, by the cells of the vertebrate pancreas and the interstitial cell of the vertebrate testis. The interesting thing is that the structural pattern of the ER is constant (within limits) for any cell type and it is possible for the experienced observer to identify a cell on the basis of ER morphology alone. These patterns and their functional relationships are discussed more fully below.

From these and other observations on the morphology and variations in this system, it becomes evident that the ER is a complex, finely divided vacuolar system which ramifies and extends to all parts of the cytosome. It effectively creates in the cytoplasm a structureless and sometimes discontinuous internal phase separated by a membrane from the continuous phase of the cytoplasmic matrix. It has been suggested, on the basis of these and related observations, that the ER provides for the channeled diffusion and segregation of metabolites. It is probable that the membrane is electrically polarized. And cytochemical studies support the oft-expressed suggestion that the large surfaces of the system provide for the support and patterned disposition of enzymes within the cytoplasm.

Diversity in ER patterns

One of the most striking features of this complex membrane-limited structure is its tendency to adopt similar patterns in cells of the same type. These patterns, it would seem, are determined by the macro-

molecular composition and arrangement in the system itself rather than by an organization in the matrix, or continuous phase of the cytoplasm. The latter seems to display too much randomness in its behavior to support a patterned structure of this nature. Furthermore, the fragments of the ER which compose the microsome fraction, as isolated in sucrose, retain a form reminiscent of that of the intact system and this in the absence of the cytoplasmic matrix (Palade and Siekevitz, 1956). Hence it is reasoned that the system is to a degree independently structured according to genetic information effective in any particular cell. Except that the expanse of this membranous system is greater than that encountered in mitochondria and chloroplasts, it is not more complicated or more incredible than the internal membrane systems of these smaller organelles. The ER, based on the nuclear envelope, pervades the whole cytoplasm; the cristae mitochondriales, only the cavities of the mitochondria.

Beside being constant for any single type of cell, these patterns are found to repeat to some degree in cells performing similar functions. It was in fact this tendency, brought to light by comparative cytology, which gave the first clues to the functional properties of these structures. Thus it was observed that the ER in cells engaged in the synthesis of protein for export is made up frequently of large, lamellar cisternae which are usually in parallel array (Dalton, 1951; Bernhard et al., 1952). And these cisternae always have dense 150 A particles attached to the outer or matrix surface of the limiting membranes (Palade, 1955b). This has led to the practice of referring to this form of the ER as rough or granular, as opposed to the smooth or agranular forms to be considered in greater detail in a succeeding section.

The rough ER

This form of the ER refers not so much to patterning as it does to a particular characteristic of the component elements. In many instances, it is true, the particle-studded cisternae are arranged in stacks or are regularly spaced in parallel arrays which may occupy a large part of the cytoplasm. This is the ergastoplasm of classical cytology (Garnier, 1899). Thus one expects to find particles associated with this organization and form of the ER vesicle. There are, however, instances where parallel cisternae are particle-free (Porter and Yamada, 1960), as also there are instances where particle-studded elements occur singly (Fig. 6). In general, however, the large flat cisternae of the ER are reserved for particle association and tend to show a degree

of parallel close array. Thus such diversity of pattern as is displayed by this type of ER is in amount and degree of association rather than in the form of individual elements.

The function of this rough form has been more thoroughly explored than any other form of the ER. As is well known, the combined evidence from cytochemistry, comparative cytology, histochemistry, and cytophotometry links the system to protein synthesis. This evidence has been so frequently reviewed that it needs no recital here (see, for example, Lotfield, 1957; Siekevitz and Palade, 1960 a. b). The important aspect of it to note for present purposes is that protein is synthesized in the particle or ribosome and is thence moved through the supporting membrane and segregated, or sequestered from the matrix of the cytoplasm to be transported toward export mechanisms represented usually by the Golgi component (Palade 1956b; Caro, 1961). Thus studies on protein synthesis have established functions for the membrane and cavity of the ER, functions of segregation and transport which might reasonably be extended to other forms of the system.

The smooth or agranular ER

As in the case of the rough ER, this represents a basic form of the system rather than a distinctive pattern for any cell type. Also, as in the case of the particle-studded ER, the smooth appears more frequently in one form, that of slender tubules (50 to 100 millimicrons in diameter) arranged in a three-dimensional lattice. Various departures from this can be found, but the unit elements remain tubular or small vesicular (Fawcett, 1955; Porter and Bruni, 1959; Porter and Yamada, 1960; Christensen and Fawcett, 1961).

The distribution of this type of ER among cells is less constant in its association with any single function than is the case in the rough form. Thus it is not possible to say that the smooth ER is designed for steroid synthesis as opposed to protein synthesis for the rough ER, although it is customary for steroid-producing cells to show this type of ER. Christensen and Fawcett (1961), for example, have described the interstitial cells of the opossum testis as showing only a smooth ER, in a close lattice. This they relate, by virtue of its occurrence in other steroid producers, as common to cells engaged in this activity. Biochemical studies by Lynn and Brown (1958) point in the same direction. The inference is that enzymes active in steroid synthesis are associated with the microsomal membranes. A very similar configuration is found in the pigment cells of the retina (Porter

and Yamada, 1960). Its role here is not so easy to determine, but since these cells are deeply involved in supplying metabolites to the adjacent photosensitive structures, it is probable that the system has some role in the interconversion of vitamin A and retinene (Porter and Yamada, 1960). Several other examples of the involvement of the smooth ER in lipid metabolism could be cited, but for our purposes here it seems more important to point out its association with other activities in other types of cells.

A striking example of the smooth endoplasmic reticulum is found in cells engaged in the concentration and secretion of a specific ion. Thus one finds unusual developments of smooth elements in oxyntic (acid-secreting) cells of the gastric mucosa in a variety of animals (Sedar, 1961). The concentration of these is especially striking near the free surface of these cells, where chloride ion is presumably being discharged into the gastric lumen. "Presumably" is inserted here because a direct demonstration of this phenomenon at the surface of oxyntic cells is still lacking. For present purposes the observation of special patterns and developments of the smooth ER in these acid-secreting cells is all that is stressed. Sedar speculates that substances normally foreign or injurious to the cytoplasmic matrix could be sequestered in the system prior to secretion.

Another chloride-secreting cell is encountered in the gills of salt-water fishes as well as in salt-secreting cells of other forms. In a recent study of these cells as encountered in the gill filaments of *Fundulus*, Philpott (1962) discovered compact reticula of smooth-surfaced ER. The extraordinary character of these is depicted in Fig. 6. The role of this system in chloride secretion is unfortunately not better understood here than in the case of the parietal cell, but the morphological association of the smooth ER with this type of cell is certainly indicative of some involvement (Philpott, 1962 a, b).

Sequestration of small molecular species may be a function common to smooth forms of the ER. This is suggested in the chloride-secreting cells and also by a form of the smooth ER found in liver cells. In these latter, it is common to find a lattice-work of tubules associated with the large rosette granules of glycogen (Fig. 2). The amount of this in a cell seems to vary from time to time in direct proportion to the amount of glycogen. Structural continuities with the rough ER, which is more nearly constant in amount, are perfectly evident and this suggests that the smooth form may have its origins from the margin or edges of the rough form (Figs. 12 and 13). In its association with glycogen, the development of tubular and vesicular elements is

such that the individual glycogen particles are essentially locked or encapsulated by the smooth reticulum (Ashford and Porter, 1961; Porter, 1961b). When glucose secretion is stimulated by glucagon injection of the animal, the glycogen disappears rapidly, and concomitantly the tubules of the smooth ER dilate as though sequestering the glucose. This interpretation is favored because at the same time, vesicles derived from the smooth reticulum accumulate at the basal and lateral cell surfaces and appear to fuse with the plasma membrane, as though to discharge their contents. It is pertinent at least in this regard that glucose-6-phosphatase is localized in membranes of the microsomes. Other enzymes of the glycogenolytic sequence are apparently not associated with the membranes but are soluble or are part of the glycogen granules (Luck, 1961).

The liver cell is not the only instance where smooth and rough forms of the ER intermingle or are at least present in the same cell. Philpott (1962), reports the presence of both in the chloride-secreting cells of *Fundulus*; Christensen and Fawcett (1961) in the interstitial cells; and Ito (1961) in the parietal cells of several forms. The extent to which the two systems are continuous in these latter instances is not indicated, but it is probable that the relationship found in the liver cell will eventually be demonstrated in these other forms.

A discussion of this smooth form of the endoplasmic reticulum would be incomplete without some mention of the system found in striated muscle (Fig. 7). Here a continuous system of tubules and vesicles occupies the sarcoplasm between the myofibrils and differentiates into a pattern of organization which repeats with each sarcomere (Porter and Palade, 1957; Andersson-Cedergren, 1959; Fawcett and Revel, 1961; Revel, 1962). This precise association with the contractile elements has convinced more than one observer that the system must be involved in some phase of muscle physiology. Thus the continuity of the system laterally in the muscle fiber led to the early proposal that it might conduct the excitatory impulse to the myofibrils located centrally within the fiber, sometimes at distances of 50 microns from the sarcolemma. More recently it has been recognized that one component of the system—the middle or intermediary element of the triad (Fig. 7), the T system of Andersson-Cedergren—is a derivative of the plasma membrane and hence a better candidate for lateral conduction (Smith, 1961; Porter, 1961e). This leaves the sarcoplasmic reticulum, or SR—the smooth-surfaced reticulum of the muscle cell—with no assigned function. One might reason from the above-reported observations on the smooth ER of the liver cell that the

systems may sequester glucose and transport it to the triad, or I-band level, for glycolytic breakdown. Fawcett and Revel have made the safe suggestion that the system "takes part in the synthesis of energy-rich compounds." But what role? Actually there is fairly good evidence from cytochemical studies, in which isolated fractions of the smooth ER were examined (Muscatello et al., 1961), that the membranes of the reticulum contain a relaxing factor (the Marsh factor) which inhibits the ATP-ase activity of the myofibril. And there is further evidence (Ebashi, 1961; Revel, 1962) that the triadic element can concentrate Ca^{++} , itself an inhibitor of the relaxing factor. Thus one sees evolving a hypothesis that would involve the sarcoplasmic representative of the smooth reticulum in muscle relaxation.

This idea finds some support from comparative morphology of the SR in muscles showing different rates of contraction and relaxation. It is observed that muscles which contract at extraordinarily high rates of frequency, for example, the toadfish swim bladder (Fawcett and Revel, 1961), the cricothyroid muscle of the bat (Revel, 1962), the synchronous flight muscle of the dragonfly (Smith, 1961), and the extrinsic eye muscles of *Fundulus* (Reger, 1961), all show extraordinarily rich developments of the sarcoplasmic reticulum. Relatively slow muscles, on the other hand, like cardiac muscle, especially in the turtle (Fawcett and Selby, 1958), show a minimal expression of the SR. It appears that the SR achieves its maximal development in muscles where relaxation has to be achieved in a few milliseconds.

The association of the endoplasmic reticulum with intracellular fiber systems is not confined to the myofibril. Fauré-Fremiet et al., (1962) have recently drawn attention to its structural involvement with the contractile bundles of filaments in a *Vorticella* type protozoan. Though the extrapolation is tenuously supported, one can reasonably wonder if many of the motions of cells, especially those which mark the return of the cell to the relaxed or preferred form, may not depend upon the preferred pattern of the ER and the relaxing factors associated with its component elements.

In this article, repeated mention has been made of patterns in the ER. These may consist, as already indicated, partly of ribosome-associated elements and partly of particle-free elements. In speaking of patterns, one refers to the relative amounts of these two forms, and their distribution with respect to one another and to the cell. Thus, although the cells of two salt-secreting organs may look alike in terms of smooth ER, they will differ enough in details of ER structural design or pattern to make a separation of electron microscope images perfectly possible if not easy.

Modulations in ER patterns

When one speaks of these patterns in this membrane system, one refers of course to the preferred pattern in the normal cell. Modulations from this pattern are an expected part of the normal functioning of the cell, and exaggerated expressions of this can be achieved by certain more or less synthetic devices.

It has been known for some time that the secretory cells of the seminal vesicle—along with several other epithelia—are responsive to hormonal control and show, as evidence of this, distinct microscopic changes. For example, under the influence of increased endogenous or exogenous androgen, these secretory cells increase in height and basophilia; whereas after castration both the height and basophilia greatly diminish (Moore, 1939). When these changes are examined for their fine-structural expression, it is discovered that in the castrate the number of vesicles and cisternae of the endoplasmic reticulum is drastically reduced (Figs. 11A and 11B). The profiles representing the system retain an orientation with the long dimension parallel to the nuclear envelope, an organization or pattern which is characteristic of the normal cell. The volume of matrix material is possibly about equal to that in the control. The great difference then is in the volume of the ER and the surface area available for the support of ribosomes. In animals stimulated by exogenous androgens, the ER of the secretory cells shows the opposite response: it obviously proliferates to produce a system with dimensions in excess of normal. We find then a distinct and perhaps direct response in this system to the presence or absence of hormone (Deane and Porter, 1960). The response is not so much in a change of pattern as in area and volume of this functioning system. A morphological hypertrophy accompanies the functional stimulation, and the hypertrophy is limited to the system most clearly associated with the function. The modulation, which appears pronounced, is one of extent rather than design. Just how the hormone achieves this proliferation or where it lodges in the cell is a problem for future investigation.

A somewhat similar though less dramatic response is shown by the liver cells of the common fowl cockerel in response to estrogen stimulation. In this form, the materials for egg-yolk formation are synthesized in the liver. The greater activity of the hen's liver in this regard is expressed in a number of morphological features including a distinctly more extensive development of the endoplasmic reticulum. When a cockerel is injected with estrogenic hormone, lipoproteins and phosphoproteins associated with egg-yolk formation appear in

the blood plasma (Common et al., 1948; Schjeide and Urist, 1956) and the liver cell approximates in its fine structure that of the hen. Among other things, the ER increases approximately two- to threefold (Figs. 12A, 12B). Here again, the pattern of organization remains much the same, with cisternae of the ER associated with mitochondrial surfaces; only the number of cisternae increases, which implies a corresponding increase in volume of the system and the area of its surface.

It is evident from the foregoing that some of the diversity encountered at subcellular levels is controlled by extracellular factors, or factors of extracellular origin. Less dramatic are of course the natural variations that accompany the normal function. When, for example, a muscle fiber contracts, there are well-known alterations in the structure of the fibrils; equally impressive but less well-known changes appear in the fine structure of the sarcoplasmic reticulum. The general pattern of organization shown by the SR of the relaxed fiber persists along with the sarcomeric organization of the fibrils, but the details of fine structure of the system are radically and characteristically altered (Franzini and Porter, unpublished observations).

A better documented example is provided by the liver cell and its response to plasmaphoresis. It has been repeatedly noted (Stenram, 1953; Lagerstedt, 1949; Glinos, 1958) that when an animal is deprived of plasma proteins through repeated bleedings (and return of cells) the distribution of the basophilic material (the ergastoplasm) in the

FIG. 11A. Parts of two secretory cells of the normal mouse seminal vesicle. The basal pole (*bp*) is at the lower right, the apical (*ap*) at the upper left. The cytoplasm is filled with long profiles of the rough ER (*rer*) or ergastoplasm, with their long axes oriented parallel to the long axis of the cell. The larger spherical vesicles toward the apical pole are part of the Golgi complex.

The pattern of organization shown by the endoplasmic reticulum in these cells is characteristic of this tissue in the normal mouse and also in mice given exogenous testosterone. Obviously the cisternae tend to enwrap the nucleus except at the basal pole. The irregular inside dimensions of the cisternae may reflect uneven accumulations of synthesized material. The limiting membranes of these cisternae are typically indistinct and densely packed with ribosomes. The total surface for ribosome association is relatively enormous (Deane and Porter, 1960). (*Courtesy of Helen W. Deane.*) Magnification: 10,200 $\times$.

FIG. 11B. Epithelial cells of the mouse seminal vesicle as they appear two weeks after castration. They have obviously lost height and are not now tall and columnar as in Fig. 11A. Basal and apical poles are indicated. The change in volume and height is reflected in, and is possibly due to a great reduction in the amount of ergastoplasm. The profiles of cisternae are still evident. The pattern of organization shown by these is the same as in the normal animal; only the long dimensions and total surface area have diminished. Magnification: 10,200 $\times$.

cytoplasm is altered. The discrete clumps of densely staining material become less discrete and the components seem to become more diffusely distributed in all cells of the lobule as they are normally in cells bordering the portal areas. One might conclude from this that rather pronounced changes would be found in the form and distribution of the rough form of the endoplasmic reticulum. Actually, when examined (Porter and Bruni, unpublished) in the electron microscope, the phenomenon is found to be referable to the distribution of mitochondria relative to the cisternae of the ER. In the relatively quiescent cell of the normal animal, the cisternae tend to appear in clusters of 8 to 12 units in parallel array (Figs. 13A and 13B). These would reasonably correspond to the discrete basophilic bodies of the light microscope image. In the plasmaphoresed animal, on the other hand, one finds these clusters of cisternae invaded by mitochondria and thus more widely separated (Fig. 13B) which, from an examination of adjacent thick and thin sections, can be seen to account for the diffuse basophilia of the stained cells. Thus in response to demands for increased synthesis of plasma proteins, the structural elements of the ergastoplasm take on a more intimate association with the mitochondria, presumably for the ATP required. Here again it is to be noted that within the limits of this modulation, the pattern of organization exhibited by the ER departs from the normal only to admit the mitochondria to the intercisternal spaces in the otherwise characteristic clumps or clusters of cisternae.

The ER in differentiation

The diverse forms which have been described and illustrated here, and in most published descriptions of this system, are found in fully

FIG. 12A. Portions of two cells from a normal cockerel liver. The picture shows the typical form of the endoplasmic reticulum in the liver cells of this animal. Single cisternae with associated ribosomes are wrapped around the mitochondria. A few vesicles of the agranular type are mixed with the glycogen (*gl*). Micrograph from class project, graduate-student course in cell biology. Magnification: 20,000 $\times$. FIG. 12B. Liver of cockerel 4 days following injection of estrogen. Among other changes induced by this treatment, there is a pronounced increase in the amount of the rough form of the endoplasmic reticulum. The cells come to resemble more closely those of the hen's liver, and the plasma requires lipoproteins and phosphoproteins found normally in the laying hen. These are synthesized for egg-yolk production. Thus exogenous estrogen induces marked proliferation of the rough ER, but the pattern of distribution and mitochondrial association remains similar to that in hen and cockerel. The nucleus is indicated at *N*, the mitochondria at *M*, spaces of Disse at *Di*, rough ER at *rer*. (Complete observations in press.) Magnification: 23,000 $\times$.

or almost fully differentiated cells. In some instances, the most prominent expression of differentiation is to be found in the ER. In other instances this system fades to morphological insignificance as the cell differentiates—as for example, in erythroblasts (Fig. 10) or in cells of the epidermis (Fig. 8) (the retaining cells of Mercer, 1961). Regardless of the degree of its eventual involvement in the fine structure of the differentiated cell, it is a constant structural component of the undifferentiated cell, and it is a constant structural component of the undifferentiated unit. It is represented from the beginning of development by the nuclear envelope, and some evidence of it as a cytoplasmic structure is found as early as the two-cell stage (Sotelo and Porter, 1959).

Since electron microscope studies of differentiating cells are still rare (see Hay, 1958; Salpeter and Singer, 1960; Slaughterback and Fawcett, 1959; Waddington and Perry, 1960; Bellairs, 1959), generalizations are hazardous. One gets the impression, however, that the system exists in embryonic cells as a relatively loose reticulum of tubular and vesicular elements (Fig. 5). Recognizable patterns of organization have so far been described only in the mature functioning cell.

Conclusions

The paper has sought to remind the reader of the wide diversity of forms which may be found at the subcellular level of structure. These are all recognized as variations on a basic architecture found in the

FIG. 13A. Part of a normal rat liver cell, showing a morphology typical for cells around the central vein of the lobule. The parallel cisternae of the rough ER are closely packed in parallel array and in this form coincide with the discrete basophilic bodies of these cells. The mitochondria normally retain a peripheral location relative to such clusters. Adjacent areas, rich in glycogen, also show dense developments of the smooth reticulum. (For symbols, see Fig. 1.) Magnification: 18,000 $\times$. FIG. 13B. Area of liver cell similar to that in Fig. 13A, except that the animal from which this tissue was taken had been deprived of approximately 15 cc of whole plasma within the 24-hour period preceding sacrifice. The change in fine structure of the ER reflects changes in the stained image in which, under the conditions of this experiment, the basophilic material appears in less discrete, more diffuse masses. This electron micrograph reveals that the cisternae in these cells are less closely arrayed and show a larger and less uniform intercisternal space; most significant of all, in accounting for the change in the light microscope image, the mitochondria are here intermingled with the cisternae. Presumably the ER here is more active in the synthesis of plasma proteins, and the changes in fine structure reflect this fact. Despite the mitochondrial invasion, the general ER pattern, characteristic of rat liver cells, persists. (Symbols as for Fig. 1.) Magnification: 18,000 $\times$.

cytoplasm of all cells (excepting bacteria and certain blue-green algae). This includes a nuclear envelope in the form of a thin, membrane-limited vesicle, perforated at numerous points in the surface. Many of the tubular and vesicular elements of the cytoplasm are apparently derivatives of or morphologically continuous with this envelope. These structures, along with mitochondria and other distinctive organelles, are bathed in the cytoplasmic matrix or continuous phase of the cytoplasm, which supports as well such resolvable differentiations as filaments or fibrils, and particulates like ribosomes and ferritin. Diversity in the subcellular appearance of the cytoplasm is largely a reflection of the degree to which one or another of these structures dominates the picture.

Special attention has been directed to the diverse forms adopted by the complex membrane-limited system known as the endoplasmic reticulum. It was emphasized that this newly defined component of cells frequently appears in recognizable patterns in cells—patterns which are constant in differentiated cells of a single tissue type. From evidence available, one may conclude that the system functions in the sequestration and transport of products of synthesis, especially where these are destined for export from the cell. It supports on its surfaces the ribosomes as specific sites of synthesis and one may infer the same surface relationship holds for other enzymic dependent reactions. Thus the ER provides a patterned distribution of functions which extends to all parts of the cytoplasm and is integrated with the nucleus, Golgi, mitochondria, myofibrils, and other structures in support of normal, coordinated cell function. Modulations in the appearance of the system involve not so much a change of pattern as a change in quantity and surface area and minor relationships to associated cell components. The fluid anatomy of the system is evidenced by these modulations. These various properties of the ER make it an excellent candidate for the role of cytoskeleton which Rudolph Peters (1956)

FIG. 14. A hepatoma cell with morphology typical for this transplantable tumor (a rapidly growing hepatoma known as the Dunning). There are many features of fine structure that one expects to see in tumor cells. The surface shows irregularities, in this instance like microvilli; the cytoplasmic matrix contains numerous free ribosomes; and the mitochondria (*m*) are small and abnormal or incompletely differentiated. The features of particular interest for this treatise are the extreme abnormalities of the ER. There are no lamellar cisternae such as one finds in the normal. Instead the system consists of irregular vesicles, with particles unevenly distributed (*er*). There is evidence of unusual extensions from the nuclear envelope (at arrows). The cell fails to store recognizable glycogen; there is no system resembling the smooth ER of the normal cell. The endoplasmic reticulum in general is highly disordered. (*Courtesy of Carlo Bruni.*) Magnification: 15,000 $\times$.

postulated some years ago as required for the integrated behavior of cells—a patterned framework of surfaces and cavities upon which enzymes could be differentially distributed and within which metabolites and products of metabolism could be segregated.

With this thought in mind, it is interesting to look at cells in which at least one expression of normal integration—the capacity to form normal tissues—is lost. These are the cells of tumors.

Carlo Bruni and I have in recent years been comparing the fine structure of rat hepatomas, possessing different growth rates, with one another and with that of the normal liver (Porter and Bruni, 1962). The hepatomas were all induced with chemical carcinogens, have gone through many transplantations, and are cytologically stable. In terms of fine structure, the cells of each tumor show characteristic features which have been constant during the period of study. A number of observations of interest could be mentioned, but in this connection it is important to note only that the structure of the endoplasmic reticulum is very abnormal (Fig. 14) and that the degree of departure from the normal is most pronounced in the more rapidly growing tumors. To what extent the lesions in the ER are involved in other manifestations of malignancy is hard to say, but, if in the normal the endoplasmic reticulum does play a role in the integrated biochemistry and physiology of the cell, the form it shows in these hepatoma cells would not be expected to support that role.

References

- Andersson-Cedergren, E. (1959), Ultrastructure of motor end plate and sarcoplasmic components of mouse skeletal muscle fiber as revealed by three-dimensional reconstructions from serial section, *J. Ultrastruct. Res.*, **2**, Suppl. 1.
- Ashford, T. P., and K. R. Porter (1961), The response of hepatic cell fine structure to isolation perfusion, *Proc. 1st Ann. Meeting Amer. Soc. Cell Biol.*, p. 10.
- Bellairs, R. (1959), The development of the nervous system in chick embryos, studied by electron microscopy, *J. Embryol. Exp. Morphol.*, **7**:94–115.
- Bernhard, W., F. Haguénau, A. Gautier, and C. Oberling (1952), La structure sub-microscopique des éléments basophiles cytoplasmiques dans le foie, le pancréas et les glandes salivaires, *Z. Zellforsch. Mikroskop. Anat., Abt. Histochem.*, **37**: 281–300.
- Caro, L. G. (1961), Electron microscopic radio-autography of thin sections: the Golgi zone as a site of protein concentration in pancreatic acinar cells, *J. Biophys. Biochem. Cytol.*, **10**:37.
- Christensen, A. K., and D. W. Fawcett (1961), The normal fine structure of opossum testicular interstitial cells, *J. Biophys. Biochem. Cytol.*, **9**:653–670.
- Common, R. H., W. Bolton, and W. A. Rutledge (1948), The influence of gonadal hormones on the composition of blood and liver of the domestic fowl, *J. Endocrinol.*, **5**:263–273.

- Dalton, A. J. (1951). Electron micrography of epithelial cells of the gastro-intestinal tract and pancreas, *Am. J. Anat.*, **89**:109-133.
- Deane, H. W., and K. R. Porter (1960). Comparative study of cytoplasmic basophilia and the population density of ribosomes in the secretory cells of mouse seminal vesicle. *Z. Zellforsch. Mikroskop. Anat., Abt. Histochem.*, **52**:697-711.
- Ebashi, S. (1961). Calcium binding activity of vesicular relaxing factor, *J. Biochem. (Tokyo)*, **50**:236-244.
- Fauré-Fremiet, E., P. Favard, and N. Carasso (1962). Etude au microscope électronique des ultrastructures d'*Epistylis anastatica*. *J. Microscopie*, **1**:287-312.
- Fawcett, D. W. (1955). Observations on the cytology and electron microscopy of hepatic cells. *J. Nat. Cancer Inst.*, **15**, suppl.:1475-1507.
- , and J. P. Revel (1961). The sarcoplasmic reticulum of a fast-acting fish muscle, *J. Biophys. Biochem. Cytol.*, **10**:89-108.
- , and C. W. Selby (1958). Observations on the fine structure of the turtle atrium. *J. Biophys. Biochem. Cytol.*, **4**:63-72.
- Fernandez-Moran, H. (1960). Cell membrane ultrastructure. Low temperature electron microscopy and x-ray diffraction studies of lipoprotein components in lamellar systems, *Proc. Ann. Meeting Assoc. Res. Nervous Mental Diseases*, Albert Einstein College of Medicine, New York.
- Franzini, C., and K. R. Porter (unpublished observations).
- Garnier, C. (1899). Contribution à l'étude de la structure et du fonctionnement des cellules glandulaires séreuses. Du rôle de l'ergastoplasme dans la sécrétion. Thesis, Nancy, no. 50.
- Glinos, A. D. (1958). The mechanism of liver growth and regeneration, in *The Chemical Basis of Development*, ed. by W. D. McElroy and Bently Glass, Johns Hopkins Press, Baltimore, pp. 1813-1839.
- Haguenau, F. (1958). The ergastoplasm: its history, ultrastructure and biochemistry, *Intern. Rev. Cytol.*, **7**:425-483.
- Hay, E. D. (1958). The fine structure of blastemia cells of differentiating cartilage in regenerating limbs of *Amblystoma* larvae. *J. Biophys. Biochem. Cytol.*, **4**: 583-592.
- Ito, S. (1961). The endoplasmic reticulum of gastric parietal cells, *J. Biophys. Biochem. Cytol.*, **11**:333-347.
- Lagerstedt, S. (1949). Previous histological and cytochemical investigations on changes in the liver cells in connection with protein assimilation. Section entitled: "Cytological Studies on the Protein Metabolism of the Liver in Rat." *Acta Anat.*, **7**, suppl. 9:1-112.
- Lotfield, Robert B. (1957). The biosynthesis of protein. in *Progress in Biophysics and Biophysical Chemistry*, Pergamon Press, New York, vol. 8, p. 347.
- Luck, D. J. L. (1961). Glycogen synthesis from uridine diphosphate glucose. The distribution of the enzyme in liver cell fractions. *J. Biophys. Biochem. Cytol.*, **10**:195-209.
- Lynn, Jr., W. S., and R. H. Brown (1958). The conversion of progesterone to androgens by testes, *Biochem. J.*, **232**:1015-1030.
- Mercer, E. H. (1961). *Keratin and Keratinization*, Pergamon Press, New York, pp. 1-316.
- Moore, C. R. (1939). Biology of the testes, in *Sex and internal secretions*, 2d ed., ed. by E. Allen, The Williams & Wilkins Company, Baltimore, pp. 353-451.
- Muscattello, U., E. Andersson-Cedergren, G. F. Azzone, and A. van der Decken

- (1961), The sarcotubular system of frog skeletal muscle. A morphological and biochemical study, *J. Biophys. Biochem. Cytol.*, **10**:201-218.
- Palade, G. (1953), An electron microscope study of the mitochondrial structures, *J. Histochem. Cytochem.*, **1**:188-211.
- (1955a), Studies on the ER. II. Simple dispositions in cells *in situ*, *J. Biophys. Biochem. Cytol.*, **1**:567-582.
- (1955b), A small particulate component of the cytoplasm, *J. Biophys. Biochem. Cytol.*, **1**:59-68.
- (1956a), Electron microscopy of mitochondria and other cytoplasmic structures, in *Enzymes: Units of Biological Structure and Function*, Academic Press, Inc., New York.
- (1956b), Intracisternal granules in the exocrine cells of the pancreas, *J. Biophys. Biochem. Cytol.*, **2**:417.
- (1956c), The endoplasmic reticulum, *J. Biophys. Biochem. Cytol.*, **2**, suppl.: 85-98.
- , and K. R. Porter (1954), Studies on the endoplasmic reticulum I. Its identification in cells *in situ*, *J. Exp. Med.*, **100**:641-656.
- , and P. Siekevitz (1956), Liver Microsomes. An integrated morphological and biochemical study, *J. Biophys. Biochem. Cytol.*, **2**:171-200.
- Pappas, G. D., and P. W. Brandt (1958), The fine structure of the contractile vacuole in *Ameba*, *J. Biophys. Biochem. Cytol.*, **4**:485-487.
- Peters, R. (1956), Hormones and the cytoskeleton, *Nature* (London), **177**:426.
- Philpott, C. W. (1962a), Teleost Osmoregulation: A comparative study of the fine structure of chloride cells in three species of *Fundulus*, *Dissertation Abstracts* (in press).
- (1962b), The comparative morphology of the chloride secreting cells of three species of *Fundulus* as revealed by the electron microscope, *Anat. Record*, **142**: 267-268.
- Porter, K. R. (1953), Observations on a submicroscopic basophilic component of cytoplasm, *J. Exp. Med.*, **97**:727-750.
- (1961a), The ground substance; observations from electron microscopy, in *The Cell*, ed. by J. Brachet and A. E. Mirsky. Academic Press, Inc., New York, vol. 2, pp. 621-675.
- (1961b), The endoplasmic reticulum: some current interpretations of its forms and functions, from *Biol. Struct. Function*, **1**:127.
- (1961c), The sarcoplasmic reticulum: its recent history and present status, *J. Biophys. Biochem. Cytol.*, **10**, suppl.:219-226.
- , and C. Bruni (1959), An electron microscope study of the early effects of 3'-Me-DAB on rat liver cells, *Cancer Res.*, **19**:997-1010.
- , and ——— (1962), Comparative fine structure of slow- and fast-growing hepatomas, *Program, 8th Intern. Cancer Congr., Moscow*.
- , and G. E. Palade (1957), Studies on the endoplasmic reticulum. III. Its form and distribution in striated muscle cells, *J. Biophys. Biochem. Cytol.*, **3**:269-300.
- , and H. P. Thompson (1948), A particulate body associated with epithelial cells cultured from mammary carcinomas of a milk factor strain, *J. Exp. Med.*, **88**:15-23.
- , and E. Yamada (1960), Studies on the endoplasmic reticulum: its form and differentiation in pigment epithelial cells of the frog retina, *J. Biophys. Biochem. Cytol.*, **8**:181-205.
- , et al. (unpublished observations).

- Reger, J. F. (1961), The fine structure of neuromuscular junctions and the sarcoplasmic reticulum of extrinsic eye muscles of *Fundulus heteroclitus*, *J. Biophys. Biochem. Cytol.*, **10**:111-121.
- Revel, J. P. (1962), The sarcoplasmic reticulum of the bat cricothyroid muscle, *J. Cell Biol.*, **12**:571-588.
- Salpeter, M. M., and M. Singer (1960), The fine structure of mesenchymatous cells in the regenerating forelimb of the adult newt *Triturus*, *Develop. Biol.*, **2**:516-534.
- Schjeide, O. A., and M. R. Urist (1956), Proteins and calcium in serums of estrogen-treated roosters, *Science*, **124**:1242-1244.
- Sedar, A. W. (1961a), Electron microscopy of the oxyntic cell in the gastric glands of the bullfrog (*Rana catesbiana*). I. The non-acid-secreting gastric mucosa, *J. Biophys. Biochem. Cytol.*, **9**:1-18.
- (1961b), Electron microscopy of the oxyntic cell in the gastric glands of the bullfrog, *Rana catesbiana*. II. The acid-secreting mucosa, *J. Biophys. Biochem. Cytol.*, **10**:47-57.
- Siekevitz, P., and G. E. Palade (1960a), A cytochemical study on the pancreas of the guinea pig. V. *In vivo* incorporation of leucine- ^{14}C into the chymotrypsinogen of various cell fractions, *J. Biophys. Biochem. Cytol.*, **7**:619-630.
- , and ——— (1960b), A cytochemical study on the pancreas of the guinea pig. VI. Release of enzymes and ribonucleic acid from ribonucleoprotein particles, *J. Biophys. Biochem. Cytol.*, **7**:631-644.
- Sjöstrand, F. S. (1953), Electron microscopy of mitochondria and cytoplasmic double membranes, *Nature* (London), **171**:30-31.
- Slautterback, D. B., and D. W. Fawcett (1959), The development of the cnidoblasts of Hydra. An electron microscope study of cell differentiation, *J. Biophys. Biochem. Cytol.*, **5**:441-452.
- Smith, D. S. (1961), The structure of insect fibrillar flight muscle. A study made with special reference to the membrane systems of the fiber, *J. Biophys. Biochem. Cytol.*, **10**, suppl.:123.
- Sotelo, J. R., and K. R. Porter (1959), An electron microscope study of the rat ovum, *J. Biophys. Biochem. Cytol.*, **5**:327-341.
- Stenram, U. (1953), Basophilic rods in the liver cell cytoplasm of rat, *Acta Anat.*, **18**:360.
- Waddington, C. H., and M. M. Perry (1960), The ultra-structure of the developing eye of *Drosophila*, *Proc. Roy. Soc. (London)*, **B**, **153**:155-178.
- Watson, M. L. (1955), Nuclear envelope, *J. Biophys. Biochem. Cytol.*, **1**:257-269.

Does Preformed Cell Structure Play an Essential Role in Cell Heredity?*

Tracy M. Sonneborn †

Department of Zoology
Indiana University, Bloomington, Indiana

I. Introduction

This chapter raises and attempts to answer a simple, fundamental question that is commonly ignored. Does the structure or organization of a cell, aside from that of its chromosomes, play an essential role in the determination of the structure of its cell progeny? This question is a timely one. The underlying simpler and basic questions of the nature of the gene and gene action, it is widely believed, will soon be essentially solved and the time will be ripe for equally concen-

* Contribution No. 716 from the Department of Zoology, Indiana University.

† The author gratefully acknowledges the invaluable collaboration of Dr. Ruth V. Dippell in the researches reported in the first half of this chapter. The investigations were supported by grants from the American Cancer Society (E80 and E81D), the Atomic Energy Commission (AT(11-1)-235), the Rockefeller Foundation, and Indiana University.

trated attacks upon problems at the next higher levels of complexity. Foremost among them are the problems of the nature, development, and inheritance of structure at the higher supramolecular and microscopically visible levels within the cell.

The simplest working hypothesis to guide thought and investigation on these problems, and one which seems to have wide current appeal, is that higher levels of structure within the cell, above that of the polypeptide product of the cistron or gene, are accounted for by three factors: the physicochemical properties of reactants, their random collisions, and the ionic and molecular constitution of the cell "soup" in which the collisions occur. These factors are held to determine how gene products are built up into multipolypeptide enzymes and structural proteins, how multienzyme systems come together in proper sequential arrangement, how enzymes and substrates come together and yield further products, how smaller ribosomes combine to form larger ribosomes, and so on. The "self-assembly" hypothesis in its most extreme form thus ultimately traces the building of *all* cellular structure to molecular contributions from milieu and genes and to random collisions of previously unarranged reactants. While some molecular biologists, at least in oral discussions, seem to have great confidence in the full adequacy of the hypothesis, others adopt it tentatively with the express purpose of seeing how far it can be carried, how much structure can be accounted for without invoking additional factors. That some degree of structure can already be explained in this way seems evident. The question then is: Do we now know, or can we discover, whether the hypothesis is sufficient to account for *all* cell structure?

Doubts have long existed, especially among students of the ciliated protozoa (Sonneborn, 1951, 1960; Wiesz, 1954; Ehret, 1960; Tartar, 1941, 1961), as to the sufficiency of so simple a hypothesis in the light of evidences for the essential role of a fourth factor, a specific preformed organization of the cell or its parts. This organization seems unable to arise *de novo* by gene action in any noncellular milieu or unorganized cell "soup." Moreover, a nucleus from a cell of one species often fails to make its own kind of cell when introduced into an organized cell of a closely related species. Decisive further evidence for an essential role of preformed cell structure in the inheritance of cell structure will be presented in the first half of this chapter. In the second half, some important literature will be considered in relation to our work and an attempt will be made to search for the underlying generalities and the mechanisms involved.

Satisfactory experimental analysis of the genetic function of pre-

existing cell structure requires a cell that possesses conspicuous, constant, normal organizational features. However, it also obviously requires a cell with readily available or producible hereditary variations of its normal features. Finally, the cell should be amenable to standard breeding analysis and other analytic procedures. Few, if any, cell types possess the needed combination of qualifications to the degree exhibited by the ciliated protozoan, *Paramecium aurelia*. Within this species, stock 51 (of syngen 4) and its derivatives are especially suitable because of the availability of adequate nuclear and cytoplasmic markers and an array of varied experimental tricks useful for decisive experimental analysis. This material was therefore used.

Attention was directed to the characteristic pattern of cortical organelles and its experimental modifications. The first and most exhaustively studied modification was the doublet cell, which possesses two complete sets of cortical structures. Doublets and other cortical variations in various ciliates, including *Paramecium*, have been studied from a number of points of view by many workers (see page 198). Some of their conclusions are closely similar to the ones reached in the present study, the unique feature of which is the completeness of the essential genetic analysis and some aspects of the search for generalities and mechanisms. Preliminary abstracts of some of the present work have been published by Sonneborn and Dippell (1960a; 1961a, b, c; 1962).

II. The Pattern of Cortical Structure in *Paramecium aurelia*

A. The cortical pattern in normal singlets

Most of the features of the cortical pattern of *P. aurelia*, which were followed in this investigation, are shown in Fig. 1A of the dorsal (aboral) and Fig. 1B of the ventral (oral) surfaces of a typical normal cell. The most conspicuous features of the dorsal surface are two large dots on the same meridian or not more than a few meridians apart. These are the pores of the contractile vacuoles. Normally, the anterior pore is further from the anterior pole than the posterior pore is from the posterior pole. The most conspicuous features of the ventral surface are all on one meridian, the oral meridian. Near the middle of this meridian is a large oval or comma-shaped depression, the vestibule, at the base of which is the clear, open mouth (technically, the buccal overture). Anterior to the mouth is a long, narrow, clear area—the preoral suture—extending from the mouth forward and left to

the anterior pole. (The directions right and left will always refer to the animal's right and left of the oral meridian which is considered to be the midventral line.) Posterior to the mouth is a long line, the eytopyge or cell anus, which lies on the postoral suture. The latter is the posterior half of the oral meridian, extending from the mouth

FIG. 1. Photographs of *P. aurelia* prepared by the silver impregnation technique. A. Dorsal or aboral surface. B. Ventral or oral surface. Magnification: 730 $\times$. a, anterior pole; al, anterior left kinety field; ar, anterior right kinety field; c, circumoral or vestibular kinety field; c v p, contractile vacuole pore; m, mouth; p, posterior pole; pl, posterior left kinety field; pr, posterior right kinety field; py, eytopyge; s, preoral suture.

posteriorly and to the left toward the posterior pole. Unlike the preoral suture, the postoral suture is not a clear area but only a line of close juncture between two different surface patterns (see below).

Although the contractile vacuole pores and the three parts of the oral meridian (preoral suture, mouth, and eytopyge) are the most conspicuous features of the cortical pattern, the rest of the cortex exhibits a beautiful constancy of finer patterning marked by regularly oriented rows of dots much smaller than those representing the

two contractile vacuole pores. Each dot represents one cilium and its base or kinetosome; each row of dots represents a row of kinetosomes, i.e., a kinety. In order not to interrupt the account of normal cortical pattern, I am going to describe at once the way the kineties are grouped into characteristically patterned areas. Much of this description will be readily followed on Figs. 1*A* and 1*B*; but part will at this point have to be taken on faith until I explain more fully (on pages 170 and 171) some of the details of cortical structure, the cytological technique used in preparing the specimens for study, and just what the dots on the photographs are.

There are six major groupings of kineties into characteristic fields: one dorsal and five ventral. The whole dorsal surface is a great field of longitudinal and parallel kineties, except where they converge near their ends to the anterior and posterior poles (which do not exactly coincide with the ends of the midlongitudinal axis of the body). The following five main ventral fields are on the sides of the oral meridian: (1) The anterior right field is composed of kineties that are nearly parallel to the preoral suture. (2) The anterior left field is composed of kineties that are almost perpendicular to the preoral suture. (3) The circumoral field lies in the vestibule and is composed of arched kineties paralleling the edge of the mouth. In ventral views the vestibule is seen on edge, making its kineties appear so close together as sometimes to give the impression of a heavy ring around the mouth. (4) The posterior right field is composed of kineties that extend parallel to the cytophyge line. (5) The posterior left field includes kineties that meet the cytophyge at an acute angle. Adjacent fields are not sharply set off from one another except where they meet the oral meridian. Thus the two anterior fields grade into the circumoral field and the latter grades into the two posterior fields, especially in the region between the posterior end of the mouth and the anterior end of the cytophyge. Likewise, the distinctive patterns on the right and left of the oral meridian grade into the simple interpolar pattern on the dorsal surface. The five ventral fields and the gullet together will be referred to as the oral segment.

By far the most complex structure of the ventral surface is its internal extension beyond the mouth into the funnel-shaped gullet or cytopharynx (Fig. 2*A*). Figures 2*B* and 2*C* show what appear to be 13 kineties, of which a large part of the gullet is composed. (This appearance may be deceptive in the light of the studies of Ehret and Powers (1959), whose descriptions suggest that the number and orientation of the gullet kineties may be quite different from the superficial appearances.) Parallel to the right margin of the mouth, at the junc-

ture between vestibule and gullet, is a single kinety, the endoral kinety. The left side of the gullet is composed of two groups of four rows of kinetosomes each, the dorsal and ventral peniculi. The dorsal side of the gullet is made up of a group of four rows of kinetosomes, the quadrulus. Between quadrulus and endoral kinety, the right side of the gullet is composed of what is called the ribbed wall (Ehret and Powers, 1959), believed to consist of an unknown number of highly

FIG. 2. The oral apparatus of *P. aurelia*. A. Vestibule (*v*), mouth (*m*), and funnel-shaped gullet (*g*), viewed from right side, in specimen prepared by modified Regaud hematoxylin technique. The positions of the unseen gullet kineties are indicated by the letters used to symbolize them in B and C. B and C. Two focal levels of silver preparation of oral apparatus viewed from its right side; photographed with bright-field objective and phase contrast condenser. Magnification: 2,100 $\times$. B. High focal level showing endoral kinety (*e*) on right side at juncture with vestibule. C. Low focal level showing quadrulus (*q*) of dorsal surface of gullet and the two peniculi (*p*) of the left side of gullet. The two peniculi are so close together as to look like one structure.

modified kineties. The ribbed wall is located to the observer's left of the endoral kinety in Fig. 2B, but it does not regularly show with the technique employed and is missing from this photograph.

The orientation of the kineties, for example, those in the left anterior field, sometimes seems ambiguous in the photographs, yet microscopic examination leaves no doubt. This requires some explanation, which can best be given by some further information on the composition of the cortex and its relation to the dots in the photographs. The cortex is a gelled surface layer 1 to 2 microns thick. (As

used here, cortex is synonymous with ectoplasm. Some authors use the term "pellicle" in the same sense; others use it only for the outermost part of the cortex.) For present purposes, we need only to explain how the orientation of a kinety is recognizable. It is defined by several features. In the first place, trichocysts alternate with one or two kinetosomes in the anteroposterior direction along a kinety. Second, a kinetodesmal fiber passes from the kinetosome to the right and anteriorly. Third, on the right side of each kinetosome (or longitudinally disposed pair of kinetosomes) is another structure known as the parasomal sac (Ehret and Powers, 1959). The kinety, its anteroposterior direction, and its right-left orientation are unambiguously defined by these criteria, as microscopic examination of any short part of a kinety clearly reveals.

The photographs printed here do not show all of these orienting features. The kinetodesmal fibers never appear. The position of the trichocysts is shown only irregularly because the fainter and smaller dots marking their positions (see Fig. 1*A* at edge) are at a slightly different optical level from the larger, darker dots marking the positions of the kinetosomes. When the latter are in sharp focus, the former are either in poor focus or out of focus entirely. However, the photographs often show a blurred, faint, broken line partially connecting the dots in the same kinety and thus showing the direction of the kinety. This serves to remove ambiguity in interpreting kinety direction, as for example in the anterior left field on Fig. 1*B*.

What then are the dots on the photographs? They are silver deposits resulting from the use of a modification of the wet-silver impregnation technique of Chatton and Lwoff (1930). Each of the heavier dots marks the position of one or two cilia (and kinetosomes) and one parasomal sac. With the phase contrast condenser the dot can be resolved into its two or three components; the parasomal sac is always on the right and the one or two kinetosomes are always on the anteroposterior line of the kinety. As Dippell (1962) has shown, these dots are deposited on the body surface; they are not within the cortex. They mark the positions where the cilia and trichocysts meet the body surface.

B. The cortical pattern in doublets

The structural features of the normal single animal are all present in duplicate in doublet animals (Fig. 3). The two sets of structures are in homopolar orientation with 180° separating each pair of homologous structures: the two contractile vacuole meridians, the two oral

meridians, and the two sets of 5 kinety fields adjacent to them. Between the two sets of structures, there is no partition. They appear to converge at a single anterior pole and a single posterior pole near the ends of the longitudinal axis. As one looks to the animal's right from the upper oral meridian in Fig. 3A, one sees first the usual three right kinety fields (anterior, circumoral, posterior) on the extreme

FIG. 3. Doublet animal at two focal levels: *A*, surface at upper focal level; *B*, surface at lower focal level. Silver preparation. Note that on *A* (upper surface) the animal's right is the observer's left; but on *B* (lower surface) the animal's right is the observer's right. *a*, anterior pole; *al*, anterior left kinety field; *ar*, anterior right kinety field; *cyp*, contractile vacuole pore. (The third dot on each contractile vacuole pore meridian is an artifact.) *g*, gullet; *m*, mouth; *ma*, macronucleus; *p*, posterior pole; *pr*, posterior right kinety field; *pl*, posterior left kinety field; *py*, cytophyge; *v*, vestibule with right circumoral kinety field.

left side of the photograph. Continuing to the animal's right, by focusing down on the left from Fig. 3A to 3B, interpolar dorsal kineties come into view and in their midst is the contractile vacuole pore meridian (or few meridians). This wide dorsal kinety field grades into the three left kinety fields (anterior, circumoral, posterior) which border on the second oral meridian seen near the right edge of Fig. 3B. The whole sequence of areas is again repeated in passing further to the right around to the starting oral meridian: the second set of right kinety fields is on the extreme right of Fig. 3B; continuing to

the right now means focusing up to the right edge of Fig. 3A, where the dorsal field and contractile vacuole pores appear; the focus is too deep to show the rest of the dorsal field; but, at the end of the circuit, the anterior left kinety field is in focus beside the starting point, the first oral meridian.

Internally, there is a common endoplasm without partition and usually only one macronucleus. In newly formed doublets, which arise from a fusion of conjugants, there are two macronuclei. After a relatively short time, these give way to a single macronucleus in the later progeny. Normally, the macronucleus of a singlet occupies a position close to the gullet with its major axis extending longitudinally; in doublets, the macronucleus usually lies close to both gullets, making its major axis transverse (Fig. 3A).

III. The Basis of the Hereditary Difference between Singlet and Doublet *P. aurelia*

As has long been known, singlets and doublets reproduce true to type through fissions (with relatively rare exceptions that will be mentioned later). Both kinds of cells also reproduce true to type through repeated autogamies and through repeated conjugations, each with its own kind. Thus, the singlet and doublet are not merely hereditarily diverse; they show a high degree of stability through sexual as well as asexual reproduction.

In order to discover the basis of this hereditary difference, the obvious thing to do is to cross the two types of cells and carry out a typical Mendelian analysis. As is well known (Sonneborn, 1947), conjugation in *P. aurelia* is a process of reciprocal cross-fertilization leading normally to genotypic identity between mates; but there are occasional failures leading instead to double self-fertilization (cytogamy). Therefore, in critical genetic work, the mates must be marked with different alleles in order to know whether reciprocal cross-fertilization actually occurred. Thus in our work on the breeding analysis of doublets and singlets, several unlinked pairs of alleles were always employed. The following account is confined to exconjugant clones in which the phenotypes, with respect to the marker genes, demonstrated that normal cross-fertilization had occurred.

Crosses of doublets by singlets regularly yielded clones of singlets from the singlet conjugants of each mating and a clone of doublets from the doublet conjugant of each mating, in spite of the fact that both kinds of F1 clones had identical genotypes. These results indicate that the difference between singlets and doublets is not due to a

difference in nuclear genotype. However, in order to leave no possibility of reasonable doubt about this conclusion, the breeding analysis was carried further by backcrosses and by crosses between, and autogamy within, F1 clones. All results agreed. No segregation of singlets from doublets occurred in these generations from parents of either type. All exconjugant clones from singlet parents, regardless of genotype, were composed of singlet cells; and all exconjugant clones from doublet parents, regardless of genotype, were composed of doublet cells. Clearly, genotypic differences here had nothing to do with the hereditary difference between singlets and doublets.

The exclusion of genotypic differences suggests a cytoplasmic basis for the hereditary difference between singlets and doublets. A direct test of this possibility involves obtaining and detecting cytoplasmic transfer between conjugants (Sonneborn, 1950, pp. 119, 120, 127). Normally there is no detectable transfer of cytoplasm between mates although it sometimes occurs spontaneously in certain stocks. It is readily inducible, however, by subjecting conjugants for a short time to specific immobilizing antiserum. This establishes a cytoplasmic bridge in the region where gamete nuclei pass across, i.e., just to the right of the mouths. The bridge persists after separation has occurred at all other points, but it may eventually disappear, freeing the mates. The degree of persistence and breadth of the bridge are roughly proportional to the amount of cytoplasm transferred. When bridges are broad, flow of cytoplasm between mates can be directly seen. Transfer of cytoplasm can also be detected indirectly by use of the cytoplasmic marker, kappa, the visible cytoplasmic particle that determines the hereditary killer trait. Each killer cell carries hundreds of kappa particles in its cytoplasm. Cytoplasm which flows from a killer to a sensitive conjugant carries kappa with it; consequently, the cells of the clone from the sensitive mate bear kappa and are killers. If cytoplasm bearing kappa is not introduced in this way into the sensitive mate, the cells of the clone it produces lack kappa and are sensitive.

Free flow of cytoplasm between mates had led to phenotypic identity of the resulting clones with regard to all previously studied hereditary traits that had been shown by breeding analysis not to be due to genotypic differences (Sonneborn, 1943; 1954a; Sonneborn and Lesuer, 1948). Its effect on the inheritance of the singlet versus doublet difference was therefore tested by using kappa and the killer trait as a cytoplasmic marker in addition to gene markers and by inducing cytoplasmic bridges between mates by the antiserum method. As shown in Figs. 4I and 4II, reciprocal crosses were made with respect to which kind of mate bore kappa and was a killer: kappa-bearing

singlet killers were crossed to kappa-free doublet sensitives; and kappa-bearing doublet killers were crossed to kappa-free singlet sensitives.

Again, as in the experiments reported above, singlet mates produced singlet clones and doublet mates produced doublet clones, regardless of whether cytoplasmic transfer had occurred (Figs. 4I β and 4II β) or not (Figs. 4I α and 4II α). Even when the bridge was broad enough to see free cytoplasmic flow through it and to assure thorough mixing of the cytoplasm, eventual separation of the mates, followed by their reproduction, yielded clones showing the same result. Never before in our experience had we encountered results suggesting the paradoxical conclusion that a hereditary difference was neither genotypic nor cytoplasmic in basis. Accordingly, new possibilities had to be considered and subjected to analytical test.

First of all, possibilities of nuclear determination by mechanisms other than genotypic differences were considered. One such possibility was nuclear differentiation. Earlier studies (Sonneborn, 1947, pp. 293, 306-307) on the inheritance of mating types had shown that macronuclei of identical genotype, arising from products of a syncaryon at conjugation or autogamy, could become permanently fixed to control one or the other of two alternative traits, but that this differentiation persists through conjugation and autogamy when new macronuclei arise by "macronuclear regeneration," i.e., by regeneration from a fragment of the prezygotic macronucleus. This knowledge was used to test whether the singlet-doublet alternative was determined by a comparable macronuclear differentiation.

Figure 5 diagrams the design of the experiment. Matings of doublets with singlets were induced to form broad cytoplasmic bridges by exposure to antiserum, as set forth above. This permitted the singlet and doublet mates to exchange some fragments of their prezygotic macronuclei, for these are carried across by the cytoplasm when the bridges are broad. In order to get the fragments to regenerate into macronuclei, all of the conjugants were from stocks that were homozygous for gene *am*. Homozygotes for this gene yield after conjugation a high percentage of first-fission products which lack macronuclei developed from the syncaryon, but which possess fragments of the prezygotic macronuclei (Sonneborn, 1954b; Nobili, 1961a). The fragments in such cells regenerate into macronuclei while segregating at random to the daughter cells formed during the next several fissions. These products of the fissions were isolated and grown into clones. The clonal phenotypes revealed whether the macronuclei were descended from a syncaryon (heterozygous for marker genes) or from

FIG. 4. Cross I: Killer doublet $\times$ sensitive singlets. Cross II: Sensitive doublet $\times$ killer singlets. α , conjugation without transfer of cytoplasm; no cytoplasmic bridges between mates. β , conjugation with transfer of cytoplasm via cytoplasmic bridges connecting mates. A and a , B and b are two pairs of allelic genes. S , sensitive; K , killer; dots represent kappa in cytoplasm.

FIG. 5. Diagram of experiment to test whether macronuclear differentiation is the basis of the singlet-doublet difference. Hollow symbols represent nuclei originally in singlet parents; solid symbols represent nuclei originally in doublet parents; stippled symbols represent hybrid nuclei formed at fertilization and their derivatives. The largest nuclei are macronuclei; the smallest are micronuclei; intermediates are fragments of the prezygotic macronuclei. The diagram is simplified by showing only a few of many fragments of a prezygotic macronucleus; actually the many fragments are not completely segregated until several fissions later.

a, singlet subclone with macronucleus regenerated from fragment of singlet prezygotic macronucleus.

*b**, singlet subclone with macronucleus regenerated from fragment of doublet mate's prezygotic macronucleus.

c, singlet subclone with macronucleus of normal kind, developed from product of fertilization nucleus.

d, doublet subclone with macronucleus regenerated from fragment of doublet's prezygotic macronucleus.

*e**, doublet subclone with macronucleus regenerated from fragment of singlet mate's prezygotic macronucleus.

f, doublet subclone with macronucleus of normal kind, developed from product of fertilization nucleus.

*b** and *e** are the critical cases; their functional nuclei (macronuclei) are derived from their mate's prezygotic functional macronuclei.

All types (*a* through *f*) have hybrid micronuclei, but these do not affect the phenotypes. The latter correspond to the macronuclear genotypes.

a fragment of the prezygotic macronucleus (homozygous for marker genes); and, if the latter, whether from a prezygotic macronucleus of a singlet or doublet conjugant (homozygous for different alleles). This is possible in spite of the fact that all clones have heterozygous micronuclei, because the micronuclear genotype has no detectable effect on the phenotype; the latter corresponds regularly to the genotype of the macronucleus (Sonneborn, 1946).

The experiment was designed to reveal whether the singlet-doublet alternative would prove to be correlated with the source of the macronucleus. The answer should be most clearly provided by two of the six obtainable types of clones, *a* to *f*, shown in Fig. 5. These two, marked with asterisks, are types *b* and *e*. Clones of type *b* are descended from singlet conjugants but possess macronuclei regenerated from a fragment of the prezygotic macronucleus of doublet conjugants. Clones of type *e*, reciprocally, are descended from doublet conjugants but possess macronuclei regenerated from a fragment of the prezygotic macronucleus of a singlet conjugant. Both types *b* and *e* should change character, from singlet to doublet and from doublet to singlet, respectively, if the basis of the difference between singlets and doublets lies in a fixed macronuclear differentiation. That did not happen. Type *b* clones were composed of singlets in spite of possessing macronuclei derived wholly from doublets; type *e* clones were composed of doublets in spite of possessing macronuclei derived wholly from singlets. Macronuclear differentiation thus seems to be excluded as the basis of the singlet-doublet difference, unless macronuclear fragments can be redifferentiated, and that has not been found to occur in any previous work.

The same conclusion is indicated by the character of clones of types *c* and *f* in Fig. 5. These are descended from singlet and doublet conjugants, respectively, that had exchanged cytoplasm (as indicated by transfer of macronuclear fragments, which can pass across only through a broad cytoplasmic bridge), and that had developed new macronuclei from products of syncarya. During this development, macronuclei are known to be subject to differentiation by the action of nuclear products in the cytoplasm (Sonneborn, 1954a). But in the case of clones of types *c* and *f*, the cytoplasm present in the cells of both types were thorough mixtures of the cytoplasm of doublets and singlets. Such mixed cytoplasm, in all previously studied cases, often differentiate the new macronuclei in accord with the newly introduced cytoplasm. But this failed to happen in the present experiment. All clones of types *c* and *f* remained true to their parental conjugant's character; i.e., those from singlet conjugants (*c*) produced singlet

clones and those from doublet conjugants (*f*) produced doublet clones. This evidence thus supports that in the preceding paragraph which excluded macronuclear differentiation as the basis of the singlet-doublet difference.

Another conceivable nuclear basis for the singlet-doublet difference lies in nuclear size. Without making measurements, it is obvious that doublets carry larger macronuclei than singlets. Do the larger nuclei determine larger cells with more cortical structures? If so, is the size of the nucleus an inherent property of the nuclei? The experiments which tested the hypothesis of macronuclear differentiation provided the answer. Mere inspection, without measurement, showed that when a fragment of macronucleus from a doublet regenerated in a singlet (type *b*, Fig. 5) it grew into a macronucleus of normal size, i.e., the size characteristic of singlets; and that, when it regenerated in a doublet (type *d*, Fig. 5), it grew to the large size characteristic of doublets. Conversely, when a macronuclear fragment from a singlet regenerated in a doublet (type *e*, Fig. 5), it grew into a macronucleus of the size characteristic of doublets; but when it regenerated in singlets (type *a*, Fig. 5), it grew to the size characteristic of singlets. The size of the macronucleus is not a cause of, but a response to, the size of the cell, as has also been reported in *Stentor* (Tartar, 1961).

The same conclusion follows from other results in clones homozygous for the *am* gene. As shown in row II of Fig. 6, this gene often yields very unequal macronuclear division at fission (Sonneborn, 1954b; Nobili, 1961a, b). Some cells receive very small pieces of macronucleus (rows II and III, Fig. 6). These small pieces grow into macronuclei of the size characteristic of the type of cell that bears them, regardless of whether the macronuclei were originally derived from singlets or doublets (row IV, Fig. 6). The observations described in this and the preceding paragraph thus effectively exclude macronuclear size as a determinant of the difference between singlets and doublets.

I have not been able to imagine any other possible mechanism of nuclear control of the difference between singlets and doublets. Therefore, attention was redirected back to possible cytoplasmic determination. The test that had been applied (page 174) was one which could reveal an effect of the freely moving cytoplasm, the fluid endoplasm, alone. It could not test an effect of the rigid, nonmobile 1 to 2 micron thick outer layer of ectoplasm, the cortex of the cell, containing nearly all the structures which distinguish singlets from doublets. As the only obvious remaining untested part of the cell, the cortex seems to be or to contain the genetic basis of the difference between

singlets and doublets. However, so potentially important a conclusion should obviously not be left to stand upon mere apparent exclusion of alternatives; it requires the strongest of direct evidence. Such direct evidence might be sought by investigating the genetic consequences of adding and removing cortical parts alone. Would grafted extra cor-

FIG. 6. Diagram showing the cellular determination of nuclear size independently of nuclear origin. *a*, *b*, *d*, and *e* are the same as in Fig. 5; solid macronuclei derived from macronuclei of doublets; hollow macronuclei derived from macronuclei of singlets. I and IV are adult cells before and after a fission, respectively. II shows fission with unequal macronuclear division, an effect of homozygosis for the recessive gene, *am*. III represents the fission products that received much less than half of the dividing macronucleus. The small piece of macronucleus, regardless of its origin (see Fig. 5) from singlets (hollow) or doublets (solid), grows into a normal-sized macronucleus in singlets and into an oversized macronucleus in doublets.

tical parts be reproduced in the progeny? Would deleted cortical parts be regenerated, or would the progeny also lack them? Answers to these questions, together with the results of the breeding analysis set forth above, should leave no doubt as to whether genetic autonomy is possessed by the cell cortex and its parts.

The difficulty in trying to answer these questions in *Paramecium* is that this organism has thus far seemed to be a poor material for the needed sort of operations. Although the body can be cut in various ways, removal of cortical parts alone has met with little or no success and grafting has never succeeded. However, as has so often happened in the past, paramecia themselves frequently accomplish what the experimenter is unable to do. In effect, they themselves performed grafting operations and brought about losses of cortical parts.

In two instances following conjugation, a paramecium grafted onto itself a piece of cortex from its mate. The first case (Fig. 7) was of course the most exciting one. A doublet mating with two singlets was briefly exposed to immobilizing antiserum in order to induce cytoplasmic bridge formation (Fig. 7A). The doublet never separated from one of the singlets; the other singlet, however, separated except for the induced cytoplasmic bridge (Fig. 7B). The singlet remained long united to the doublet, but eventually separated (Fig. 7C). The doublet and its still-attached other singlet mate eventually died, but the free singlet lived. When first observed after separation, it bore a very conspicuous extra piece, as shown in Fig. 7C. Its doublet mate at that time showed a corresponding nick, as if a piece were missing. The singlet, in breaking away from the doublet, had apparently taken along a piece of the doublet's cortex. Instead of growing rapidly and dividing within about 11 hours, as is normal for exconjugants, this singlet failed to grow for two days. During that period, its extra piece flattened out, making the posterior part of the body distinctly wider (Fig. 7D). Then the abnormal animal grew and reproduced (Fig. 7E). From one (but not the other) of the two products of the first fission arose a clone obviously different in form from either singlets or doublets (Fig. 7F). When samples of this clone were observed, after preparation by the silver impregnation technique, we found to our amazement and delight a new hereditary type that went far toward answering our question about the autonomy of cortical parts.

These animals (Fig. 8) were intermediates between singlets and doublets. Like doublets, they had two complete oral segments: two vestibules, mouths, and gullets; two anterior and posterior sutures; usually two cytopyges; and two sets of the five typical associated kinety fields (anterior right and left, circumoral, and posterior right

and left). Unlike doublets, however, these two oral segments were not 180° apart. Instead, the two mouths were 90° or less apart and of course there was much less separation between the left kinety fields of the right segment and the right kinety fields of the left segment. Further, the anterior and posterior sutures of the left oral segment curved to the right instead of to the left, meeting the sutures of the right oral segment at common poles. In many of these cells a *third* cytopygge appeared between the two on the postoral sutures. The

FIG. 7. Diagram of origin of the new type of animal shown in Fig. 8. *A*. Exposure to antiserum of two singlets conjugating with a doublet. *B*. Only one singlet separating, but with cytoplasmic bridge joining it to doublet. *C*. This singlet separated, but bore a hump. The nick in the doublet indicates that the source of the hump was a piece of cortex from doublet mate. † indicates death of doublet and other singlet; these never separated. *D*. The singlet one day later: hump "assimilated"; posterior half of body enlarged. *E*. First fission. *F*. One of the two daughter cells is a new cell type. NOTE: It is not known whether the new type came from the posterior or anterior product of this fission, only that one product produced a clone of the new type and the other did not.

FIG. 8. Silver preparations of cells belonging to new type of clone whose origin was diagrammed in Fig. 7. *A* and *B* are photographs at two optical levels of same animal. Note two complete oral meridians about 90° apart (instead of the 180° separation typical of normal doublets). *C* is another type of cell frequently produced in this clone; it differs from the others only in having three instead of two cytopyges. The animals of this clone have only one dorsal surface and in it only one contractile vacuole pore meridian or narrow segment. *ar*, anterior right kinety field; *al*, anterior left kinety field; *m*, mouth; *pr*, posterior right kinety field; *pl*, posterior left kinety field; *py*, cytopyege; *s*, preoral suture.

structures of the rest of the body were single: there was a single dorsal surface with the usual solitary narrow segment bearing contractile vacuoles. These features of the new cell type were as a rule faithfully reproduced during successive fissions.

This new hereditary cell type is theoretically decisive. A piece of paroral cortex, pulled off of one cell and incorporated on the surface of another cell, has led to the development and inheritance of an entire supernumerary oral segment from pole to pole. In other words, genetic autonomy of a delimited part of the cell cortex has been demonstrated by the consequences of a natural graft of a small piece of cortex.

Further, observations on the cytophyge in the new cell type gave a clue to its developmental and genetic determination. Unlike the normal situation, the third cytophyge is not located on a postoral suture; but like the normal situation, it is located at the juncture of right and left postoral kinety fields. A third juncture of this sort exists in these cells as a result of the positions of the two oral segments: the left postoral field of the right oral segment abuts on the right postoral field of the left oral segment. And there the third cytophyge appears in spite of the absence of a postoral suture and in spite of the reversal of right-left relations. The development and position of the cytophyge are thus correlated with localized structural features, the border between two specific different cortical fields.

The second example of "cortical picking" was the reciprocal of the one just described. The doublet conjugant robbed its singlet mate of a paroral piece of cortex. The singlet failed to grow and died. The doublet developed a third oral segment close to one of its two pre-existing segments. This too was inherited, but reversion to the doublet condition occurred with considerable frequency.

The preceding examples indicate that an interpolar segment of cortex, when integrated into a cell, behaves with a high degree of genetic autonomy. Because growth of *Paramecium* is in the longitudinal direction, the question arises as to whether genetic autonomy is possessed only by one or more entire meridians from pole to pole. The answer to this question is provided by a type of clone which has arisen independently several times and in more than one way. Clones of this type show that some parts of an interpolar segment can be inherited while other parts of the same segment are lacking. In other words, genetic autonomy is not restricted to entire longitudinal segments, but holds also for parts of a segment along the anteroposterior axis.

Animals of this type (Fig. 9), have two oral segments 180° apart, but only one of them is complete. The other oral segment lacks vesti-

bule, mouth, gullet, and the typical circumoral kinety field, but possesses all other structural features of an oral segment, i.e., the preoral suture with its right and left kinety fields and the postoral suture with its cytophyge and right and left kinety fields. Between the complete and incomplete oral segments are two aboral surfaces also

FIG. 9. Doublet lacking vestibule, mouth, and gullet of one oral meridian, but retaining rest of the meridian. Two focal levels of silver preparations of same cell. *A*. Surface with incomplete oral meridian at upper focal level. *B*. Surface with complete oral meridian at lower focal level. (As before, the animal's right and left are the same as the observer's on the lower surface but are reversed on the upper surface.) *al*, anterior left kinety field; *ar*, anterior right kinety field; *m*, mouth; *pl*, posterior left kinety field; *pr*, posterior right kinety field; *py*, cytophyge; *s*, preoral suture; *, area where vestibule, mouth, and gullet are missing.

180° apart, each with its segment bearing contractile vacuole pores. This incomplete doublet, lacking only the middle parts of one oral segment, reproduces true to type. Clearly, the smallest autonomous cortical part is less than an entire normal interpolar segment.

A number of other hereditary cortical variations have been obtained: triplets, quadruplets, a much increased number of contractile

vacuole pores and meridians, and various hereditary disturbances in the development, growth, and relative movement of cortical parts during fission. We shall not go into these in this chapter, although each contributes further evidence of the genetic autonomy of cortical parts and further insight into how cortical structures are reproduced.

The features of cortical reproduction which bear importantly on the present problems will become evident from a comparative examination of how the main components of the highly structured oral segment are reproduced in singlets, doublets, and incomplete doublets. Each of the structures of this segment—the vestibule, mouth, and gullet of the oral apparatus and the sutures, the kintety patterns, and the cytopyge—follows the same rules of reproduction in all three kinds of cells. The differences among the cell types reveal what is essential and what is nonessential for the reproduction of various parts.

We can undertake to give here only some of the salient features of the reproduction of the parts of the oral segment. Actually the story is still not fully known in spite of careful study by many investigators. Among the early signs of reproductive activity, three are outstanding (Fig. 10). (1) A field of new kinetosomes appears near the posterior and right margin of the mouth (Fig. 10*A*). This is the rudiment of the new oral apparatus, destined to pass to the opisthe, the posterior product of fission. (2) Many new kinetosomes appear in the kinetics of the left wall of the vestibule and in a zone extending to the left from the vestibule (Fig. 10*B*). (3) Within this area the kinetics break to form the cleavage line (Fig. 10*B*), which begins between the left posterior edge of the old mouth and the left end of the rudiment of the new oral apparatus. The cleavage line continues around the body

FIG. 10. Formation of the oral rudiment in the fission of singlets, doublets, and one type of incomplete doublet. Silver preparations.

A. Singlet. Rudiment of new oral apparatus forming along the right side and posterior end of the mouth and vestibule of the old oral apparatus.

B. Singlet. Two focal levels showing the new oral rudiment which is beginning to be shifted posteriorly by elongation of vestibular kinetics.

C and *D.* Two focal levels of doublet. *C*, surface at upper focal level; *D*, surface at lower focal level. Both oral meridians and adjacent areas are undergoing the same developments as the one oral meridian and its adjacent areas in the singlet in *B*.

E and *F.* Two focal levels of incomplete doublet lacking one vestibule, mouth, and gullet. *E*, surface at upper focal level showing the oral meridian that lacks vestibule, mouth, and gullet. A cleavage line extends from this meridian almost all the way around the body. *F*, surface at lower focal level with complete oral meridian; new oral rudiment extends slightly posterior to fission line.

Symbols for Fig. 10 are given in the legend for Fig. 12.

and eventually ends in the right wall of the vestibule at the anterior end of the rudiment of the new oral apparatus.

Exactly the same processes occur in doublets, as appears in Figs. 10C and 10D. What happens on the one oral segment of singlets happens on both oral segments of doublets. The same is true for the two oral segments in incomplete doublets possessing two adjacent oral segments but only one dorsal field. Of special significance is what takes place in incomplete doublets that lack only the vestibule, mouth, and gullet on one of their two oral meridians (Figs. 10E and 10F). The complete oral segment behaves as already described for the oral segments of singlets and doublets. The incomplete oral segment does not form a rudiment of a new oral apparatus; but the other two processes—increase in kinetosomes and formation of the cleavage line—occur nevertheless. The cleavage line starts on the left of the defective oral meridian at the place where the missing vestibule, mouth, and gullet should be and passes to the left until it reaches the right edge of the mouth on the complete oral meridian. And the cleavage line that began on the left of that oral meridian passes to the left until it meets the incomplete oral meridian on its right side. Hence the increase of kinetosomes and formation of the cleavage line are independent of the presence of vestibule, mouth, and gullet. And a rudiment of a new oral apparatus appears where and only where an oral apparatus already exists. Thus only one arises in singlets, two in complete doublets, two in incomplete doublets possessing two complete oral meridians, but only one in incomplete doublets possessing an oral apparatus on only one of their two oral meridians.

The main feature of reproduction of the oral segments in the next stages in fission is elongation of the kineties in which new kinetosomes were earlier packed close together. When the cleavage line reaches the right wall of the vestibule, many new kinetosomes appear there. The elongation of the kineties anterior to the cleavage line on both sides of the old oral apparatus has the effect of pushing posteriorly both ends of the rudiment of the new oral apparatus. This places the new rudiment, which is organizing into an oral apparatus, entirely posterior to the old oral apparatus (Fig. 11A). Unequal growth places the new rudiment to the right (Fig. 11B). The same elongation of kineties occurs on both oral meridians of doublets; and wherever there is a rudiment of a new oral apparatus, it is thereby carried posteriorly and to the right. Figures 11C and 11D show this stage on both oral segments of a complete doublet.

The further elongation of kineties occurs differentially in groups, according to a definite pattern which results in the formation of typi-

FIG. 11. Separation of the new oral rudiment from the old oral apparatus in the fission of singlets and doublets. Silver preparations.

A. Singlet. New oral rudiment shifted to position directly posterior to old oral apparatus.

B. Singlet. New oral rudiment posterior and to the right of old oral apparatus.

C and *D.* Two focal levels of doublet animal. *C.* surface at upper focal level. *D.* surface at lower focal level. Note that both oral meridians show nearly the same stage as the one oral meridian of the singlet in *B.*

Symbols for Fig. 11 are given in the legend for Fig. 12.

cal kinety fields. The kineties anterior to the cleavage line begin to elongate before those posterior to the line; in this region, those on the left of the oral meridian elongate more than those on the right. Figure 12*A* shows an early stage of this process and Fig. 12*B* a late stage. The result is to establish the typical posterior right and left kinety fields between the old oral apparatus and the cleavage line. The juncture of the two new kinety fields is the new postoral suture. There the new cytoppyge is destined to be formed, but it is not yet present.

As Fig. 12*A* shows, considerable kinety elongation occurs anterior to the cleavage line before any can be detected posterior to the line. Later, the kineties posterior to the cleavage line begin to elongate (Fig. 12*B*). This takes place differentially, but in reversed relations; i.e., the kineties on the animal's right of the new oral apparatus

FIG. 12. Reconstruction of new kinety fields in the fission of singlets, doublets, and one type of incomplete doublet.

A. Singlet. New oral apparatus still at cleavage line. Between the level of the old oral apparatus and the cleavage line, the kineties have elongated, more on the left than on the right, thus starting the formation of new posterior right and left kinety fields.

B. Singlet. Anterior to the cleavage line, the two new posterior kinety fields have almost been completed by further elongation of kineties, more on the left than on the right. Cytoppyge is not yet developed at the juncture of the right and left fields (postoral suture line). At this stage, the kineties posterior to the cleavage line have also begun to elongate, much more on the right than on the left, beginning the formation of the new anterior right and left kinety fields. The right kineties, by elongating up above that part of the cleavage line which is connected with the anterior part of the new oral apparatus, are converting this part of the cleavage line into the new preoral suture.

C and D. Two focal levels of doublet in a stage of fission between that of the singlets in *A* and *B*. *C*, surface at upper focal level; *D*, surface at lower focal level. The areas on the sides of both oral meridians show the same developments as for the areas adjacent to the single oral meridian in singlets. *C* shows the earliest beginnings of the new preoral suture; i.e., the kineties to its right are just beginning to elongate.

E. Upper surface of an incomplete doublet showing the oral meridian that lacks a vestibule, mouth, and gullet. Same stage of fission as the singlet in *B*; formation of the preoral suture from part of the cleavage line by elongation of right kineties up above part of it.

Symbols for Figs. 10, 11 and 12: *al*, anterior left kinety field; *ar*, anterior right kinety field; *cl*, cleavage line; *cvp*, contractile vacuole pore; *k*, area of increase in number of kinetosomes; *ke*, area of elongating kineties; *m*, mouth; *mn*, mouth of new oral apparatus; *mo*, mouth of old oral apparatus; *pl*, posterior left kinety field; *pr*, posterior right kinety field; *py*, cytoppyge; *pyo*, cytoppyge open (to discharge undigested food); *roa*, rudiment of oral apparatus; *s*, preoral suture; *scl*, preoral suture forming from part of cleavage line.

elongate much more than those on its left. The result (Fig. 12*B*) is to establish the typical anterior right and left kinety fields in an area of new growth between the new oral apparatus and the main part of the cleavage line. I had to say "main part" because, as appears in Fig. 12*B*, part of the cleavage line—the part that was at first immediately on the left of the oral meridian—becomes the new preoral suture. This happens as a consequence of the growth of the right kineties up and to the left, so that they come to lie anterior to the part of the cleavage line which is still connected to the anterior edge of the new oral apparatus.

Exactly the same processes occur in doublets and incomplete doublets wherever there is an oral meridian. Figures 12*C* and 12*D* show them occurring on both oral meridians of a doublet in a stage of fission intermediate between that of the singlets of Figs. 12*A* and 12*B*. Figure 12*C* is a particularly good picture of the beginnings of the elongation of the right kineties posterior to the cleavage line: they are already penetrating into the cleavage line and marking off the earliest trace of the prospective new preoral suture. Figure 12*E* shows that these definite patterns of kinety elongation take place around the oral meridian even when it lacks a vestibule, mouth, and gullet. The new preoral suture line forms posterior to the cleavage plane as a result of the elongation of the right kineties into the cleavage line and around that part of it destined to become the preoral suture.

Thus far only the normal or regular processes in singlets, doublets, and incomplete doublets have been described. They reveal much about how cortical cell heredity is accomplished. More is revealed by observations on exceptions to this course of events, by failures of certain processes to occur, and by the origin of certain structures under conditions not expected from the preceding account of normal events. Only some of the more important exceptions will be mentioned in this paper.

Two principal kinds of failures in the occurrence of expected processes have been noted. On the one hand, certain groups of kineties that normally elongate have failed to do so or have done so to a less than normal extent. In some cases, the vestibular and nearby kineties have failed to elongate and, as a consequence, the rudiment of the new oral apparatus fails to grow apart from the preexisting oral apparatus. This may be dependent upon the prior failure of new kinetosomes to appear in these kineties; but the cause of these failures is still obscure. Often, but not always, correlated with but slight elongation of these kineties is failure of the new gullet to invaginate: its kineties remain up on the body surface more or less posterior to

the preexisting vestibule, mouth, and gullet. We have not previously mentioned this invagination process, which has been noted by Ehret and Powers (1959), and the other details of the transformation of the rudiment into a functional normal oral apparatus because much is still obscure. But invagination of the gullet, the rudiments of which develop in the cortex near the old mouth-vestibule juncture, is part of the story. This invagination may also be correlated with elongation of kineties, possibly, at least in part, those of the rudiment itself.

The other main kind of failure is the failure of the rudiment of vestibule, mouth, and gullet to form and develop even when a pre-existing oral apparatus is present. Two examples can be cited. First, it can happen in complete doublets and it thus results in the origin of incomplete doublets. Significantly, this failure is preceded and accompanied by certain other aberrations. The number of kineties between the oral segment and the contractile vacuole pores to its right is much reduced, from about 25 to about 8 to 15. The preoral suture is much shorter and correspondingly the vestibule, mouth, and gullet are much more anterior; i.e., they are located at its base. In correlation, the postoral suture is much longer than normal. The cytopyge is more variable in size, often being short and much further than normal from the posterior end of the preoral suture. When these relations exist, the vestibule, mouth, and gullet on such an oral meridian are sooner or later destined to be irreversibly lost in the progeny. The dislocation of parts that precedes and accompanies this loss suggests an essential determinative role of localized regional interactions in the formation of the oral apparatus. To this we return later.

The second example is failure of incomplete oral meridians, and of one of two closely placed oral meridians, to persist and be reproduced at autogamy and conjugation. At present this is just a brute fact which has not been studied as to details, even to the extent of knowing whether it invariably happens. No such failure occurs when two complete oral meridians are far apart.

A different type of failure appears in some cells of doublets that have two oral meridians less than 90° apart. At fission, the cleavage line may be inhibited or appear late in the region between the two mouths (Fig. 13A). Correlated with this, the kineties in that region often fail to elongate to the normal extent. Consequently, in the course of successive fissions, the number of kineties between the two oral meridians is progressively reduced, sooner near the poles than near the equator, bringing the polar ends of the anterior and posterior sutures together further and further toward the equator (Fig. 13A). Eventually the two preoral sutures are reduced to one in this way

and the same is true for the two postoral sutures and cytopyges; but the two vestibules, mouths, and gullets may still be present (Fig. 13*B*). In this condition, the latter can still be reproduced (Fig. 13*B*); but eventually even the two vestibules, mouths, and gullets become one in the progeny. This is one way in which cells with two oral

FIG. 13. Two stages (*A* and *B*) in the fusion of oral meridians in a line of descent that had the two oral meridians less than 90° apart.

A. Fission stage: the two new oral apparatuses have formed and separated from the old ones. Each oral meridian has a short preoral suture flanked by right and left kinety fields. Less than halfway to the anterior pole the two preoral sutures meet and proceed to the pole as a single suture flanked by its right and left kinety fields. The two oral meridians have still shorter independent postoral sutures (in the opisthe, of course) before they meet and proceed to the posterior pole as a common suture flanked by a single right and a single left posterior kinety field. There is but one cytopygge; it lies on the common postoral suture. A cleavage line appears to the right of the right oral meridian and (in typical wide form) to the left of the left oral meridian, but it is totally absent between the two oral meridians.

B. A late fission stage of another animal in the same line of descent. The entire suture line (preoral and postoral) is single. The only residue of doubleness appears in the vestibules, mouths, and gullets, both sets of which have reproduced.

al, anterior left kinety field; *ar*, anterior right kinety field; *cl*, cleavage line; *m*, mouth and vestibule; *pl*, posterior left kinety field; *pr*, posterior right kinety field; *py*, cytopygge; *s*, preoral suture; *, area between vestibules where kineties have not cleaved.

meridians can produce progeny with only one. Their reproductive autonomy is guaranteed only when the two oral meridians are 90° or more apart. That is why it has proved impossible to obtain and maintain multiplets of an order higher than quadruplets. It is also evidence that important interactions and interferences occur when corresponding replicate structures are unduly close together, suggesting—as we shall later discuss more fully—that interacting gradients play an important part in cortical morphogenesis and heredity.

In addition to losses in the ways just described, new structures may possibly arise *de novo*. For example, origin of a new oral apparatus at a place where none preexisted possibly occurred as an exception, although this is still clouded with uncertainties. It will be recalled that we twice observed the origin of a whole new oral segment following the “natural grafting” of a piece of cortex from near the oral region of a mate (page 181). Unfortunately, exactly what the grafted piece contained remains unknown. Thus far the event has occurred too rarely to be fully studied in its early stages. But it is clear that no more than a short region of the oral segment, if any, could have been grafted; and that from the graft developed an entire oral segment with all five typical associated kinty fields, as well as vestibule, mouth, and gullet. From the comparison of the behavior at fission of complete and incomplete oral meridians given above (page 186), we may be sure that, in order to do what it did, the graft must have contained an essential component, or all that was necessary for the formation of a new oral apparatus and also for the independent determination of a cleavage line, localized increase in kinetosomes, and localized differential elongation of kineties. That the graft might have lacked many or all of the gross structures of the supernumerary oral apparatus and its associated regions is suggested (but not proved) by two facts. First, one product of the first fission lacked them. Second, in many other ciliates new oral structures arise far from the preexisting ones, indicating a less obvious mechanism of production than one depending upon proximity to the preexisting oral structures themselves. Perhaps something else normally associated with, but separable from, the oral apparatus in *Paramecium* is the basic determiner of formation of vestibule, mouth, and gullet. If so, this—or part of it—but not the oral structures themselves, may have been in the active grafted piece; in which case, the gross structures would have arisen *de novo*, i.e., indirectly, under the influence of a particular, different local cortical pattern.

Such a mode of origin is certainly involved in the formation of the cytopyge. As pointed out, it arises at the juxtaposition of the posterior

right and left kinety fields. Normally, this juxtaposition is on the post-oral suture, but when two oral meridians are not more than 90° apart, a third juxtaposition with reversed right-left relations exists between them, and there a third cytophyge arises, in the absence of a third oral segment or any part of it (page 182). This suggests that cytophyge formation is determined by interaction between these two kinety fields or the cortical regions in which they are located.

This, together with analyses in other ciliates to be discussed later, makes more credible the possibility considered above in relation to the consequences of "natural grafts," of *de novo* origin of the oral apparatus by results of interaction between other juxtaposed diverse kinety fields or cortical regions. Ehret and Powers (1959) have also pointed out this possibility. Indeed, virtually the whole oral meridian from pole to pole, as well as the juncture between vestibule and gullet, is a series of junctures between visibly diverse cortical patterns, which would be more than ample to account in principle for all the determinative actions in morphogenesis and cell heredity correlated with the oral segment and its parts, if indeed such actions are the essential ones involved.

In the light of the preceding account, a preliminary summarization of the main relations may be attempted. In general, new oral segments are formed where and only where oral segments or their decisive correlated cortical pattern junctures preexist. The rudiment of a new oral apparatus likewise arises where and only where an oral apparatus or its cortical determiner preexists. The preexisting oral apparatus and the parts of the oral segment beside and anterior to it pass to the proter, the anterior product of fission; the new rudiment and the parts of the oral segment beside and posterior to it pass to the opisthe, the posterior product of fission. Both proter and opisthe form new half-bodies. Except for the new oral rudiment, formation of these new parts involves: localized and differential (in space and time) increase in the number of kinetosomes in preexisting kineties; formation of the cleavage line by breaks in the kineties starting on the left at the oral meridian and proceeding around the body to the oral meridian; the elongation of the kineties (spacing the originally crowded kinetosomes) according to a definite space-time pattern which creates the new kinety fields; specific elongation of the right circumoral and neighboring kineties of the developing opisthe forward into the cleavage line and to the left, so as to create a new preoral suture in the opisthe from part of the cleavage line; induction of specific new developments at junctures of specific cortical regions—certainly the induction of a cytophyge where new posterior right and left kinety

fields about and possibly induction of vestibule, mouth, and gullet in a comparable way.

In all these normally correlated events, only a few major types of processes seem to be taking place: localized increase in the number of kinetosomes; localized elongation of kinetics in a definite space-time pattern; and progressive differentiation of particular new formations into definite structures. That there is a decisive developmental and genetic role of preformed structure in these processes is evident from the correlations, which have repeatedly been pointed out, between supernumerary and reduced cortical parts and the numbers of new parts that arise during fission. Between the old and the newly formed parts there is no evidence of a simple, direct, template-like, causal relationship. Reproduction of cortical structures is typically very indirect, involving a complex, dynamic series of events. But these events are normally correlated in space with the locus of visible cortical differentiations. Usually, when the localized patterns are present, even in supernumerary and abnormal positions, the corresponding developmental events occur; when they are lacking, the events fail to occur. Remarkable is the fact that such localization of origin and development of parts and even of differential growth occurs within the confines of a single cell.

The major remaining problems are to identify the nature of the particular cortical patterns correlated with and presumably determining the particular pattern changes that lead to the development of new structures, and to discover the nature of the interactions that appear to occur between diverse local regions. Whatever these may be, the end result is two from one, two new cortical patterns normally identical with and essentially dependent upon the original one, both when the original is a typical singlet and when it deviates in any one of several ways from the typical singlet organization. In other words, the processes we have been describing at the gross level of microscopic visibility are the processes underlying cell heredity of the cortex and its highly differentiated parts in *Paramecium aurelia*.

IV. Literature, Discussion, and Conclusion

Are the conclusions reached from our study of one stock of one species of ciliate of limited or general applicability? How should they be modified, if at all, in the light of related work and thought on other organisms? These questions would of course be appropriate in regard to conclusions based on the study of any one organism, perhaps especially when that organism is unicellular and a ciliate, for these organ-

isms differ greatly among themselves and are in some ways unique. We would believe conclusions to be fundamental only if they were found to be applicable to multicellular as well as to unicellular organisms. Such a search for the fundamental is the object of this discussion section. Previous pertinent work on doublets and other cortical variants in ciliates will be considered first; then related work on multicellular organisms; and finally, the general conclusions to be drawn on the role of cortical structure.

A. Doublets and other cortical variants in ciliates

The inheritance and morphogenesis of doublets have been studied in a number of genera of ciliates in the course of the last 40-odd years. Most of this work has been critically and thoughtfully reviewed from several points of view in a series of papers by Fauré-Fremiet (1945; 1948a, b; 1950; 1954). All studies show that doublets reproduce true to type during fissions; but to the question "How long can they do this?" the answer is not so simple. Many workers found that cultures begun with doublets eventually end up with only singlets present (for example, Chatton, 1921, on *Glaucoma*; Sonneborn, 1932, on *Colpidium*; Fauré-Fremiet, 1948a, on *Leucophrys*; Tartar, 1954a, on *Stentor*; Hanson, 1962, on *Paramecium*). Does this mean that the "inheritance" of this cortical variation is limited and that eventually the nuclear genotype prevails by restoring the normal singlet condition? While even limited inheritance would be of some significance, the full genetic import of cortical variations depends in part upon the degree to which they persist during reproduction. What then is the genetic significance of the reversion of cultures of doublets to singlets?

Doublets have been reported to give rise to singlets in several ways. A commonly observed way is for a depression to arise between the two sets of organelles at the anterior end and gradually to deepen in the course of successive fissions until it reaches the fission plane, which then cuts off two singlets and a doublet (Margolin, 1954). There are also some minor variations of this process. Another very different process is asymmetric reduction of the number of kineties and the distance on one side between two corresponding meridians such as the two oral meridians (Fauré-Fremiet, 1948a). This leads to eventual interaction between, and loss of one of, the two sets of structures (see page 193 above). Once this process starts in a subline, it cannot be reversed by selection. A third process is resorption of the

two sets of organelles and replacement by one new set (Fauré-Fremiet, 1945; Tartar, 1954a). Still other processes have been observed.

The question is whether, by one process or another, doublets are destined inevitably to revert to singlets. The answer is "No." Apparent ultimate mass reversion has been shown to be due to one or more of several causes which by no means involve 100 per cent reversion or even a high rate of reversion. Chatton (1921) pointed out that the singlets which arose with low frequency in *Glaucoma* multiplied faster than the doublets and simply overgrew the culture. This has been confirmed in a number of other ciliates, for example, for *Colpidium* by Sonneborn (1932). As he pointed out, the same result would occur even in the absence of differential reproductive rates, but more slowly, because doublets keep producing singlets occasionally, while the latter do not revert to doublets. This one-way change leads to higher and higher proportions of singlets in the culture. Chatton (1921), Margolin (1954), and Uhlig (1960) have further shown that the rate of singlet production by doublets depends upon the cultural conditions; conditions that permit rapid multiplication reduce the frequency of reversion. Uhlig, ascribing Tartar's (1954a) limited maintenance of doublet *Stentors* to suboptimal cultural conditions, succeeded in maintaining them for hundreds of fissions and apparently could do so indefinitely. Indeed, under appropriate conditions, doublets have been maintained indefinitely by culling out the revertant singlets (Dawson, 1920, on *Oxytricha*; Sonneborn, 1932, on *Colpidium*; Uhlig, 1960, on *Stentor*; the present work on *Paramecium*; and by others on other ciliates). Moreover, Sonneborn (1932) showed that selection for morphologically more "perfect" or symmetrical doublets in *Colpidium* could reduce the frequency of reversion from one in seven line-days to zero in 864 line-days. Doublets are not destined inevitably to revert to singlets: aside from occasional production of singlets, they can in general keep reproducing true to type by fissions indefinitely.

However, occasional production of singlets is genetically important in two respects. First, exceptions to doublet self-reproduction are useful, as teratological phenomena usually are, in revealing some of the processes involved in the normal maintenance of type (see page 192). Second, the fact that revertant singlets and their doublet sister lines of descent carry products of division of one and the same ancestral nucleus implies strongly that the difference between the two cell types could hardly be due to a genotypic difference in their nuclei (Sonneborn, 1932).

One report of limited persistence of doublets, not due to reversions at all, requires special mention. Calkins (1925) noted that a doublet clone of *Uroleptus mobilis* lost the capacity to conjugate. The singlets that arose in the clone regained this capacity. Because *Uroleptus* exhibits a clonal life cycle ending eventually in death in the absence of conjugation, the doublet *Uroleptus* was doomed to extinction. No other case of this kind has ever been found. Doublets do conjugate and this has made possible study of the inheritance of the doublet condition through fertilization, especially in *Euplotes* and *Paramecium*.

Both in *Euplotes patella* (Kimball, 1941) and in *Paramecium aurelia* (Sonneborn, 1942, and the present paper) doublets can conjugate with doublets; doublets of *P. aurelia* also undergo autogamy. The doublet condition persists in both the exautogamous and exconjugant progeny. In both genera, doublets were also crossed to singlets (Kimball, 1941; Powers, 1943; Sonneborn, 1942), with the same results as reported in the present paper. Genic markers showed that the doublets had been fertilized by their singlet mates (Powers, 1943). Although genetic analysis of the basis of the difference between singlets and doublets was carried no further than this, the results—as far as they go—are in complete harmony with ours. Both singlet and doublet exconjugants reproduced true to type in spite of having identical genotypes.

That no genotypic difference distinguishes singlets from doublets is also indicated by the modes of origin of doublets. In some genera, the two products of fission of a normal singlet reunite to form a doublet, as was first noted by Chatton (1921) for *Glaucoma*. In this case, the fission is abortive; the two daughter cells remain united by a cytoplasmic bridge and the opisthe moves up alongside of the proter in homopolar orientation. In other genera, such as *Colpidium* (Sonneborn, 1932), the abortive fission may by further replication of parts form a multiple monster from which eventually both homopolar doublets and normal singlets may pinch off. In *Paramecium* (Sonneborn, 1942) doublets arise from a pair of conjugants by the formation of a cytoplasmic bridge which spreads in the main posteriorly. The first fission then yields two singlets from the anterior part of the fused exconjugants and a doublet from the posterior part. The details of origin show minor variations from pair to pair. The two conjugants that yield a doublet are of identical genotypes (as a result of the mating, if not prior to it). Thus, in all three of these modes of doublet origin, the two components of a doublet are genotypically identical

with each other and with singlets of the same clone or clones. The point is even more striking in *Stentor* doublets: they can be produced by experimentally relocating parts of one cell (Tartar, 1961; Uhlig, 1960).

The production of doublets from parts of one cell or by fission or conjugant fusion of two identical cells, the origin of subclones of singlets from doublets, and the limited breeding results from crosses of singlets by doublets, all argued strongly for the absence of pertinent nuclear differences between the two types of cells before our present study proved the point beyond reasonable doubt by exhaustive analysis.

The role of the cortex and its parts in determining the hereditary difference between singlets and doublets, as well as in determining the production and ordering of the diverse localized cortical parts, has also long been studied in many ciliates (Fauré-Fremiet, 1945-1954). Most of these studies are based upon, stimulated by, or related to the kinetosome theory, which can therefore serve as a point of departure for the following discussion.

To Lwoff (1950) we are indebted for a succinct, documented, thoughtful, and influential account of the kinetosome theory. This theory is concerned with the genetic and morphogenetic properties of kinetosomes and kineties. The basic genetic feature of the theory is its attribution of genetic continuity to the kinetosomes. Kinetosomes are held to arise only by division of preexisting kinetosomes. The genetic continuity of kineties is held to be due to the retention of products of kinetosome division in the same row. We shall consider the morphogenetic features of the kinetosome theory after discussing the genetic feature.

The genetic continuity of kinetosomes is accepted by some modern workers, rejected by others. Mazia (1961) argues, as well as can be argued from what is now known, that kinetosomes reproduce themselves from a germinal part. His evidence is largely derived from studies of centrioles; but centrioles are kinetosomes, according to the kinetosome theory, and this is supported by their similar fine structure (De Harven and Bernhard, 1956). On the other hand, Ehret and Powers (1959) contend that definite proof of kinetosomal self-reproduction by division is lacking. The mere appearance of a new kinetosome beside an old one is not proof; and no one has yet demonstrated in a new kinetosome material contributed by a preexisting kinetosome. Ehret and Powers suspect that minute unit elements formed deep in the cell (in *Paramecium*) migrate to the surface and there grow

into a unit of cortical structure, the ciliary corpuscle, which develops kinetosomes and other parts. Definite proof or exclusion of one or the other view is still lacking.

This uncertainty about the self-reproduction of kinetosomes has obvious but limited implications for the genetics of cortical organization. The structures and processes with which we are concerned are at higher levels than that of kinetosomes and the way they arise. They include the ways in which kinetosomes are grouped in more complex structures, as well as the seemingly simpler creation of fields with characteristically positioned and oriented kineties. The problem of the origin, development, and inheritance of structure at this level is independent of the answer to the question of the origin and production of individual kinetosomes.

The morphogenetic aspects of the kinetosome theory are far more important for present purposes. Local differences in cortical structure in other ciliates, as in *Paramecium*, involve variations in the spacing, orientation, differentiation, and relative growth rates of kineties and groups of kinetosomes. It is therefore of basic importance to know whether such variations are inherent in the constituent kinetosomes and kineties or whether they are imposed upon them by other features of the local milieu. On the choice between these alternatives, the kinetosome theory is not firm. Lwoff (1950) recognized both of them and seemed to favor the latter. In other words, he thought it to be more likely that the kinetosomes and kineties were indispensable instruments of cortical differentiation, rather than that they might be the cause of it. Others have come out strongly for the causal interpretation (Wiesz, 1951). Fauré-Fremiet (1945-1954) repeatedly considers the possibilities in varied lights; he sometimes seems to lean toward the one, sometimes toward the other interpretation. Because the choice between these alternatives is fundamental for our theme, the main types of evidence bearing on it will now be considered.

The first type of evidence comes from observations of the normal course of events. Any particular structure normally arises at a particular spot marked either by a certain pattern of kineties or by association with the same kinety. This is well illustrated by the normally constant locus of origin of a new oral apparatus. As the most prominent set of cortical structures, it has been studied most. In many ciliates, such as *Tetrahymena*, it always arises in association with a particular interpolar kinety designated as kinety No. 1 or the "stomatogenic" kinety. In *Paramecium*, the endoral kinety (page 170), by the right edge of the mouth, is held by some to be the stomatogenic kinety (Roque, 1956b; Porter, 1960, at least in part). Their views

imply that this localization of the presumed material source of the new mouth and gullet is the basis for the regular appearance of these structures close to the old one and for the dependence of a new oral apparatus on the presence of a preexisting one. Fauré-Fremiet (1954) refers to this confinement of the stomatogenic kinety within the oral apparatus as the basis for its "autonomization." Here then is one view, based mainly upon the normal course of fission, of why and how new structures are determined by preexisting structures: a particular kinety determines a particular cortical structure.

The second type of evidence comes from experimental interferences with the normal situation and leads in the main to opposite conclusions. The most extensive and intensive work has been done on *Stentor* (Tartar, 1961; Uhlig, 1960). Normally, only the apical end of the new oral apparatus (called the peristomial primordium, the primordium, or the anlage) arises in association with a particular kinety; the basal end extends across several kineties. Thus, the peristomial primordium normally arises in a definite fixed position marked by a particular kinety pattern. In pigmented species of *Stentor*, the kineties regularly alternate with pigmented stripes. Because the latter are more conspicuous, descriptions are commonly given in terms of the stripes. On the midventral surface there is a zone of narrow ramifying stripes flanked on the left by broad interpolar stripes. The apical end of the primordium arises on this border and the basal part within the ramifying zone.

Is this normal position of origin of the primordium due to the inherent properties of one or more kineties in that region? Nearly 60 years ago Stevens (1903) removed the ventral half of the body of *Stentor*, within which lay the ramifying zone, and a considerable region on both sides of it, and found that the peristomial primordium was nevertheless formed. This has subsequently been repeatedly confirmed. Eventually, Tartar (1956) discovered what visible physical conditions were necessary for the formation of the peristomial primordium. The pigment stripes in the ramifying zone are the narrowest on the body. Around the body to the right from this zone the stripes become progressively broader; they are thus broadest on the other side of the ramifying zone. Tartar showed that relocations of body parts yielded primordia only where artificial junctures between wide and narrow stripes were made. For example, when the ventral half of the body is removed, the cut edges of the dorsal half join, making a new juncture between narrow and wide stripes, and there a primordium arises and develops.

Experiments such as this—and many variations of it have been

performed, mainly on *Stentor* (Tartar, 1961; Uhlig, 1960) and on *Blepharisma* (Suzuki, 1957)—show clearly that no kinety or kinety field is alone endowed with the capacity to be the site of formation of the complex peristomial primordium. This can happen anywhere around the whole circumference of the body when a juncture of contrast in stripe (and kinety) widths is set up by cuts and/or grafts. Apparently any kinety or group of kineties can serve as the site of primordium formation. The kineties can be no more than instruments of morphogenesis; they are not its cause.

What then is the cause of specific morphogenesis? Tartar's results seem to prove that the origin and development of the peristomial primordium is determined by interaction between adjacent narrow- and wide-striped cortical areas. As he and Uhlig (1960) note, the visible gradation of stripe widths around the body is the visible expression of a gradient with its extremes meeting on the ventral surface where the apical end of the primordium normally arises. They showed, however, that not all artificially produced junctures of different stripe widths serve to induce primordium formation. If two junctures of unequally diverse stripe contrasts are created, the lesser contrast may fail to induce a primordium, especially if it is close to the greater contrast. Even if only one juncture exists and the stripe contrast is extremely slight, it may fail to induce. There are thus degrees of interaction at junctures correlated with difference in the gradient levels, and degrees of inhibition of primordium formation diffusing in gradient fashion from a major juncture. Another index of gradient action is the delineation during fission of new fine stripes in the fine-stripe ramifying zone: the number of new stripes decreases with distance from the broad-stripe region.

The *Stentor* work shows that the anteroposterior position of the primordium and the differentiation of its parts are determined by another gradient extending from the foot or base toward the apex. This gradient, for example, determines where the primordium bends to the right across the fine-stripe ramifying zone. Further, the differentiation of a mouth at the base of the primordium depends upon the presence of a foot. Supernumerary feet in abnormal positions induce the formation of supernumerary mouths in abnormal positions; absence of a foot results in failure to form a mouth. Altogether, then, the origin and development of the peristomial primordium depend upon the interactions between the two gradients.

Two facts already mentioned give a most important clue to the general nature of operation of the first gradient. Although an adjacent wide stripe zone is essential, all the new formations appear either at

the juncture or within the fine stripe zone. The apical end of the primordium appears at the juncture; its basal portion and the new fine stripes arise within the fine-stripe zone. This strongly suggests that the broad-stripe zone acts as an inductor, its active principle(s) passing into and acting upon the narrow-stripe zone, which behaves as if it were a competent responder to the inductor. We should recall here that supernumerary inductor-response systems are partitioned at fission so that multiplets reproduce true to type indefinitely under favorable cultural conditions (Uhlig, 1960).

Thus far, decisive evidence that the determinative conditions are confined within the cortex has not been given. The cuts and grafts in *Stentor* and *Blepharisma* can hardly be expected to be limited absolutely to the cortex. The cortex merely contains the geographical markers correlated with the observed events. Tartar (1961), however, reports in a preliminary way another kind of operation on *Stentor* which points to a decisive role of the cortex. He made an incision and withdrew all or virtually all of the endoplasm, leaving cortex and nucleus. Such an operation was promptly followed by the restoration of the endoplasm, normal growth, and normal reproduction. Conversely, Tartar also stripped off the entire cortex, leaving endoplasm and nucleus. These cells became spherical, presumably developed a surface membrane (because the remaining cytoplasm was retained), and lived for some time; but they failed to regenerate cortex or to develop further, eventually dying. However, if a piece of cortex was left on the endoplasm, it gradually spread around the latter, reconstituted the visible markers of its gradients, and eventually regenerated and reproduced normally. These experiments indicate that a small piece of cortex possesses or creates at least an essential part of the morphogenetic gradients and that the cortex does not arise in the absence of preexisting cortex. Unfortunately, as Tartar realizes, the experiments fall short of being completely decisive. The cortex might simply be providing certain mechanical or osmotic properties essential for normal cellular functioning, including the production of cortex itself.

The experimental analyses on *Stentor* (and on *Blepharisma*) provide beautiful models of the roles in morphogenesis and cell heredity of gradients, presumably in the cortex, and of inductor-response systems in the cortex. The question is: Are these models generally applicable to ciliates or are they limited to certain genera, for example, those which show an extraordinarily well-developed capacity for regeneration? Uhlig (1960) warns of the possibility of limited applicability and calls for direct evidence from a variety of ciliates.

Although operative analyses of the sort carried out on *Stentor* have not been performed on *Paramecium*, there is nevertheless much to indicate that fundamentally the same models may apply, as will now be set forth.

The existence of cortical gradients in *Paramecium* and morphological evidences of their direction are shown by three lines of evidence. First, the position of lateral spines, a type of abnormality studied by Jennings (1908), shifts with the growth and division of the paramecia. The nearer the spines are to the ends of the body, the less their position shifts per fission cycle; the greatest shifts occur when a spine is near the equator. This indicates that growth is greatest in the equatorial zone and decreases toward the poles. Second, slow growth at the poles is further indicated by the relatively slight restoration of cut ends at a single fission; several fissions are required to complete regeneration of polar parts (Tartar, 1954b and others). Third, the greatest increase of kinetosome number during fission occurs near the equator; there is less and less with increasing distance from the equator. Thus all three lines of evidence point to a growth gradient which is high near the equator and which decreases toward both poles.

Paramecium, like *Stentor*, also possesses sharp juxtapositions between areas exhibiting different kinety patterns. The oral meridian, virtually from pole to pole, is a line of juxtaposition between markedly diverse kinety fields on its right and left (page 169). There may even be differences on the two sides of the line in distance between rows of kinetosomes, those on the left usually being greater than those on the right. But this difference, if extensive measurements confirm it, is much less than the corresponding difference in *Stentor*. Much more striking is the difference on the two sides of the mouth. The vestibular kineties are very closely packed and run parallel to the rim of the mouth; the adjacent kineties forming the gullet not only are differently spaced (some closer, some further apart) but are mostly oriented in a different direction. Ehret and Powers (1959) point out two other differences on the two sides of the vestibule-gullet juncture: hexagonal versus rhomboidal patterns of their "ciliary corpuscles" and one or two versus four kinetosomes per repeating unit.

The important question is whether one or more of these abrupt junctures of visibly diverse cortical patterns has morphogenetic significance. The postoral juncture on the two sides of the cytopyge clearly does, as we have shown (page 184). It will be recalled that the morphogenetic action at the postoral juncture is independent of right-left relations, as was also found to be true for primordium

production at the narrow- and fine-stripe juncture in *Stentor* (Tartar, 1961). These observations appear to indicate that the interactions in both cases are due to an inductive influence from the one region (wide-striped in *Stentor*) which spreads both to the right and to the left. Normally, only the region on one side is competent to respond in both cases; but when competent regions are on both sides at once, both sides respond concurrently. That a competent responding area must be present for the response to occur is shown by the normal limitation of the response to one side. So far as I am aware, present knowledge of the cytotpyge does not permit a choice as to whether the left kinety area is the inductor and the right the competent responder, or the reverse. Perhaps the discovery of induction at the juncture, even when the juncture is not on an oral meridian and when the right-left relations are reversed, will inspire successful efforts to resolve the remaining questions.

With regard to the possible morphogenetic role of the cortical pattern juncture between vestibule and gullet in *Paramecium*, resolution of the alternative interpretations is rendered especially difficult. The position where the anlage of the new oral apparatus is first seen is very close both to the endoral (stomatogenic?) kinety and to the juncture of the strongly marked differences in cortical pattern. Attempts to test which of these structural landmarks, if either, is decisive for the appearance of the anlage of a new oral apparatus could follow either of two paths. First, the earliest beginning of the anlage could be studied to see if it is in fact physically contiguous with the endoral kinety. Roque (1956b) and Porter (1960), using the silver method, claim that it is; Ehret and Powers (1959), using phase and electron microscopy, claim that it arises several microns from this region.

Second, attempts might be made to separate the endoral kinety from the vestibule-gullet juncture. This would not be easy, perhaps not possible, to do surgically; however, repetitions of certain experiments, plus cytological controls, might accomplish the task. One such experiment was Hanson's (1955) destruction of the ability to form a gullet on one oral meridian of a doublet by exposing the area of the right posterior vestibule-gullet juncture on that meridian to a fine beam of ultraviolet irradiation. Cytological study accompanying such an experiment should reveal whether certain structures such as the endoral kinety were destroyed. Hanson did not have cytological controls and did not even know whether the rest of the oral meridian persisted in progeny that lacked one vestibule, mouth, and gullet. Perhaps they were like our incomplete doublets that had this oral segment except for vestibule, mouth, and gullet. Another type of

experiment that might yield the necessary information is induction of "picked" cortex (page 181). Cytological study should reveal whether the effective picked pieces contain an endoral kinety, a vestibule-gullet contrast juncture, only one part of that system, or merely some pattern of cortical structure which is decisively different from that of the area into which it is implanted.

While awaiting decisive evidence as to the possible roles of the endoral kinety and vestibule-gullet juncture, the available facts concerning the general situation point to a tentative choice. Kinetics genetically differentiated for the production of oral apparatuses (or for any other visible structure) do not exist in the decisively analyzed ciliates. Instead, interacting gradients, visibly expressed as juxtaposed different cortical areas, have proved to be determinative. In *Paramecium*, the determination of at least one structure—the cytopyge—occurs in the same way. In the absence of contrary decisive evidence in any ciliate and in the light of supporting evidence for the principle of interacting gradients in all well-analyzed cases (including one in *Paramecium*), tentative rejection of the interpretation of genetically differentiated kineties, including the endoral kinety of *Paramecium*, seems justified. For the same reasons, tentative adoption of the principle of determination of morphogenesis by interacting gradients or fields seems justified, even when they are not correlated with visible pattern junctures, as in the ciliate *Glaucoma* (Frankel, 1960a, b, 1961).

Up to this point, the problem under discussion has been the part played by the cortex in the perpetuation through fissions of its own specific pattern of differentiation. However, the discussion has proceeded upon a tacit assumption which now needs to be made explicit and to be subjected to careful scrutiny. Tacitly it has been assumed that the processes of morphogenesis at fission partake of an all-or-none character. Either a structure develops or it doesn't. And this is determined by the interactions of a few primary gradients at junctures of gradient contrast. So relatively simple a model is a good first approximation, but the impression of its adequacy comes from putting aside a large class of facts which will now have to be examined.

To begin with, the reader will find on page 193 an account of a series of cortical deviations from a complete doublet type which led to eventual complete and irreversible loss of vestibule, mouth, and gullet on one of the two interpolar oral segments of doublets, leaving all other landmarks of this segment. A number of stages in the loss were observed. The process thus occurred gradually over a series of successive fissions. Observations of comparable progressive cortical

losses over a number of fissions were earlier reported by Fauré-Fremiet (1948a) for *Leucophrys patula*: asymmetric reduction in the number of kineties led inevitably to loss of one set of cortical structures from doublets. Likewise, Hanson (1962) has noted progressive changes. Localized irradiation of one oral apparatus of a doublet with a fine beam of ultraviolet inflicts damage on the exposed oral apparatus and on the capacity to produce new ones on that meridian at fission. Both of these kinds of damage, without further irradiation, can independently lead to greater or lesser abnormality over the course of a considerable number of fissions until one or the other of two stable conditions is reached: complete loss of one oral apparatus and the capacity to produce one on that oral meridian, or complete normality.

Observations like these are important in at least two respects. First, they show that morphogenesis and cell heredity of particular cortical structures are not all-or-none phenomena. The initiation of a cortical structure is only the first step in its creation. Initiation alone does not assure complete normal development. For the latter, other factors are involved. Apparently, there is a sequence of such factors, the operation or effect of later factors in the sequence being dependent upon the normal or abnormal results of the operation of earlier factors in the sequence. The nature of these factors and of their operation in ciliates is almost totally unknown, although initial attempts to understand them have been made by Fauré-Fremiet (1948a) and Hanson (1962).

Second, observations of these sorts, by providing an example of progressive cortical changes over a series of successive fissions, may provide a connection in principle between the short-term phenomena of cell heredity and morphogenesis and the long-term progressive changes in morphogenesis and cell heredity, extending over hundreds of successive fissions, that constitute the clonal life cycle of immaturity, maturity, senescence, and death (Maupas, 1888; Calkins, 1926a; Sonneborn, 1954c; Jennings, 1944). Among the events of clonal aging is an increasing frequency of abnormalities of cortical structures and of abnormalities in the processes of production of cortical structures at fission (Sonneborn and Schneller, 1955; Sonneborn and Dippell, 1960b), including final failure to form at fission an oral apparatus (Dippell, 1955). The basic mechanisms underlying this one-way progression away from "normality" (i.e., the condition in young clones) may be similar to those operating in the progression toward complete loss of one set of oral organelles from doublets. Further analysis of either may throw light on both.

Obviously ciliates would long ago have become extinct were there not also reversals of the unidirectional, progressive, degenerative clonal age changes in morphogenetic processes. Return to the starting point is required for survival. This is commonly accomplished in ciliates at the time of the fertilization processes. In *Paramecium*, for example, conjugation and autogamy are the only occasions on which a preexisting oral apparatus involutes, disappears, and is replaced by a new oral apparatus (Hertwig, 1889; Roque, 1956a; Porter, 1960). The new one arises in the same manner as at fission, but it remains in the cell in which it arises. At the same time, the cilia on part of the ventral surface of the two conjugants are lost and later regenerated (Hiwatashi, 1955). Replacement of the corresponding kinetosomes (or ciliary corpuseles) and comparable replacements on the rest of the surface have not yet been reported. In view of the initiation of a new life cycle at fertilization and of the evidence for progressive age changes in the cortex, a deliberate and careful search for more extensive renewal of the cortex at the time of fertilization might prove fruitful. In any case, it is clear that at least some cortical changes form not merely a sequence, but a cycle. Similar cyclic returns to the starting point are associated in some ciliates with encystment and with "physiological regeneration," a disappearance of some old organelles, and their regeneration.

Cyclic return to the starting point is also shown during the asexual reproduction of parasitic ciliates with multiple hosts (Lwoff, 1950). Markedly different cortical patterns are characteristic of the period of association with different hosts. One pattern leads regularly to the next in correlation with a definite and regular sequence of hosts. The sequence of cortical changes and hosts leads back to the starting point. It too is not merely a sequence, but a cycle.

These observations on the cycles of cortical changes in the course of the lives of multihost, parasitic, asexual ciliates and of free-living sexual ciliates are of great significance for any general theory of the role of the cortex. They show that, in ciliates, cortical morphogenesis and heredity are not—as is commonly supposed—normally limited to the regular bipartition and reconstruction of exactly the same cortical pattern fission after fission, but are progressive—in part, very slowly progressive—sequences of changes forming a closed cycle back to the starting point. This larger view of the cortical events in ciliates provides a model of what would have to be involved in multicellular organisms if the cortex played in them a role of importance comparable to the role it plays in ciliates.

B. Multicellular organisms and cortical evolution

Double animals, parabiotic twins, and the like are—as everyone knows—commonplaces of the experimental embryology of Metazoa. But they are not reproduced. This is because the individual cells are not changed and reproduction is from a single cell or a fusion of two cells. Only the pattern of cellular association is altered. However, the cellular association itself functions genetically in the asexual reproduction of certain Metazoa and these can form hereditary doublets as stable as those in ciliates. Sonneborn (1930) obtained two differently oriented doublets of the Rhabdocoel turbellarian, *Stenostomum incaudatum*. Each reproduced true to type indefinitely through asexual reproduction. These cases are particularly instructive not only because they are in Metazoa, but especially because detailed comparison with the doublets in ciliates reveals so much about the basic similarities. In both the ciliates and Metazoa, the gross reproduction true to type involved indirect, dynamic processes quite independent of any template processes of replication per se. In both, the basis lay in the perpetuation of a new combination of the number and the arrangement of parts which were indirectly self-reproducing. In both, the perpetuation depended upon inductor-response systems of morphogenesis with complex development of the hereditary structural parts in each generation. The chief difference lay in the units of organization: cells and tissues in the case of the Metazoa, parts of the cortex of a single cell in the case of the ciliates.

What do the Metazoa have to tell us about the role of the cell cortex? The literature of experimental embryology of higher animals is replete with evidences of the importance of the cell cortex. It plays decisive roles, for example, in cell “recognition” during reaggregation of dissociated cells (Moscona, 1957), in organizer and inductor actions and the response to them by competent cells (Weiss, 1939), in the mosaic of determinative regions of the egg (Daleq and Pasteels, 1937, 1938; Curtis, 1960), and presumably in the whole distinctive molecular organization of each cell type (Weiss, 1962, and his earlier papers on molecular ecology and related matters). The conclusions set forth above as to the existence and roles of gradients, fields, interactions between parts, and movement or differential growth in ciliate morphogenesis and cell heredity have their counterparts in the older and much more extensive experimental embryological studies on Metazoa.

Recently Curtis (1960) has demonstrated the cortical localization

of the determinative parts of the pattern of the mosaic egg of *Xenopus*. In such eggs, the different parts of the egg cortex are segregated into different cells during cleavage, presumably in the absence of growth or further cortical developments. When growth and further development begin, the cells endowed with diverse cortices appear to follow different lines of cortical development (cf. Sonneborn, 1960). Juxtaposition of cells with different cortical patterns leads to organizer and inductor-response reactions. The parallels to regionally diverse cortical parts in ciliates, and the interactions between those that are adjacent, are too obvious to be labored. The main difference between unicellular and multicellular organisms in these respects is in the nature of the differentiated and interacting units: different regions of the cortex of one Protozoan cell correspond in principle to the cortices of different cells in the Metazoa (Fauré-Fremiet, 1954).

Finally, in Metazoa as in Protozoa, the cortical patterns and events undergo a progressive self-directed sequence of changes, during the clonal life cycle in the latter and during the individual life cycle in the former. In both, one change leads to the next and the sequence becomes a cycle by a return to the starting point. In the ciliates, any cell (after immaturity and before advanced senescence) can undergo fertilization (or encystment or physiological regeneration) and return to the cortical starting point. Likewise all cells capable of division may be able to do this in certain plants (Steward, 1961). Only certain cells can do it in the Metazoa. In the Protozoa and the Metazoa, the return may be rapid and normally confined to a cell that has undergone fertilization or one specialized to undergo it. The many parallels suggest that the roles of the cell cortex and the principles of their operation may be fundamentally alike in unicellular and multicellular animals.

This basic similarity implies an evolutionary development of cortical specificities; and the considerable degree of autonomy of cortical parts suggests a corresponding degree of independence in their evolution. This is strongly indicated by the existence of a number of genera of flagellates and ciliates (for example, *Giardia*, *Teutophrys*) in which the animals appear morphologically to be doublets or higher multiples of the animals of other genera (Fauré-Fremiet, 1945). Within this limited group, the evolution of genera clearly runs parallel to the laboratory production of hereditary doublets and multiplerts.

In general, there is no doubt about the existence of cortical evolution in the ciliates. The whole taxonomy of this group hinges upon cortical characteristics (Corliss, 1961). However, nearly all of the gross morphological features of ciliates are cortical features. The

important point is whether the evolutionary changes in the cortex were due to genomic changes, independent cortical changes, or parallel series of independent but selectively correlated changes in genome and cortex. No direct evidence on these alternatives is available or is likely to be obtained. If cortical changes other than mere losses or additions of parts (for example, shifting the position of the oral apparatus from equator to pole, a type of difference that distinguishes certain ciliate taxa) could be produced and shown to be autonomously inherited, then independent cortical evolution would be indicated. Thus far, nothing like this has been accomplished in the laboratory or seems likely to be. I am inclined to favor the possibility of parallel, independent, and selectively correlated evolution of genome and cortex, for it has well-established precedent in the parallel evolution of genome and plastids in *Oenothera* (Stubbe, 1960; Cleland, 1962).

Some of the main trends in cortical evolution are both obvious and instructive. Zooflagellates, from which the ciliates evolved, are an enormously varied group. The progression from simplest to most complex is outstandingly accompanied by corticalization of an increasingly large number and variety of kinetosomes and their associated structures. In the simplest flagellates, these are unitary, simple, and for the most part deep in the endoplasm. The centriole is in them the key kinetosome. While it may be compound in structure, it is associated with both spindle fibers and the one or few flagella. In somewhat less simple flagellates, there are one or few kinetosomes and more complex and more numerous fibrous structures, such as cresta, axostyle, and costa, in addition to spindle fibers and flagella. The kinetosomes, as well as most of the other parts, still lie deep in the endoplasm. In the more complex flagellates, several changes occur: the number of kinetosomes increases greatly, they concentrate more and more upon association with flagella, and they lie prevalingly near the surface of the body. Finally, in the opalinids the cortex of the anterior pole or apex of the cell reaches its fullest development as a sort of organization center, with a ring of apical kinetosomes organizing longitudinal kineties. In the ciliates these longitudinal kineties become independent of the apical region and the cortex of the equatorial region becomes the center of growth and organization.

The original internal location of kinetosomes and other parts in Zooflagellates and their evolutionary migration to the cortex support Ehret and Powers' (1959) view of the internal origin of kinetosomes (or of ciliary corpuseles) in the ontogeny of *Paramecium*. But other aspects of the evolutionary picture do not support their view that the cortex of *Paramecium* is composed solely of packed ciliary corpuseles,

aside from trichocysts. Flagellates have a cortex even when the kinetosomes lie in the endoplasm; it may be little more than a limiting membrane system. Perhaps the outermost membranes of *Paramecium* are homologous with the outer membranes of flagellates. The well-known inducible shedding of the pigment-containing "pellicle" in *Blepharisma* (Nadler, 1929) and *Stentor* (Tartar, 1961) without loss of essential nonreplaceable cortical components, taken in connection with Tartar's earlier cited result on the irreplaceability of the *Stentor* cortex, seems to weigh against that part of the Ehret and Powers (1959) conception which reduces the *entire* cortex to elements having internal origin.

The account just given indicates that new parts were added to the cortex during the evolution of the Protozoa. Knowledge is lacking as to whether certain cortical parts or the whole cortex is the seat of the gradients and physiological diversities that direct the course of morphogenesis and the processes which result in the inheritance of cortical characteristics. In the Metazoa, as we have seen, gross morphological differentiation of the cortex has largely disappeared; but the underlying determinative physiological differentiations of the cortex are present in highly developed form, as Curtis's (1960) experiments demonstrate. This leads to the guess that the basic genetically and morphogenetically important part of the cortex is not the part that is unique to the Protozoa, the part bearing the morphologically visible differentiations.

C. The role of preformed cortical structure

The question with which this chapter began—whether the structure of a cell (aside from that of its chromosomes) plays an essential or nonessential role in the determination of the structure of its cell progeny—has been answered unambiguously by experimental analysis. Such structure may play little or no part in many essential intracellular activities which depend only on the physicochemical properties, amounts and random collisions of newly imbibed food (in the broadest sense), and newly produced direct and indirect products of gene action. On the other hand, our study of *Paramecium*—in general agreement with a number of other studies—shows that preformed cortical differentiations are essential for their own reproduction. Certain visible cortical structures failed to arise *de novo* when they were initially lacking. Experimental modifications of the visible cortical organization were perpetuated during cell reproduction. Thorough breeding analysis, combined with other accessory modes of

analysis, proved that the difference between two hereditary cortical types, singlets and doublets, was not due to any genic, nuclear, or endoplasmic difference. Finally, natural grafts of a piece of cortex showed that it was decisive in morphogenesis and cell heredity. The autonomy and genetic significance of cortical structure was thus fully established.

We then tried to pin down just what cortical structure is genetically autonomous and just how this genetic autonomy is achieved. Conspicuous cortical structures, like kinetosomes, kineties, and groups of kineties, to which a causal role in morphogenesis and cell heredity has often been attributed (and still is, in many quarters), are clearly not genetically fixed as to their developmental and genetic roles. They are one instrument, not the cause, of morphogenesis and cell heredity. They are responsive to, and express visibly, the more basic, underlying causal cortical structure. This finer structure, probably at the molecular level, is the basis of the genetically and morphogenetically decisive interactions we have described as occurring between juxtaposed diverse cortical areas which are the seats of gradients of structure and action.

What are the determinative molecular species, configurations, and modalities in the cortex? At present this remains almost or quite completely unknown, not only in ciliates but in all organisms. Fauré-Fremiet (1950, 1954) has put forth stimulating suggestions based upon the general geometrical parallels between kineties and macromolecules such as polypeptides and polysaccharides. He points out that both exhibit similar symmetry, polarity, and other polymeric features. He suggests that the properties of the visible structures are consequences of the comparable properties of the constituent and/or surrounding cortical macromolecules. This suggestion is attractive, but alone it of course does not account even in principle for the modalities of morphogenetic processes and their genetic consequences.

How can we in general bridge the gap between the (unknown) molecular structure of the cortex and the morphogenetic and genetic processes which surely are linked to it? Speculation about the answer to this question cannot be entirely free; it must be limited to conform with a good deal of pertinent knowledge. In particular, it must conform with what is known about nuclear functions. The cistrons or genes make polypeptides, proteins, enzymes, and indirectly the products of enzymic activities. Some of these doubtless enter into the composition of the cortex of the cell. Differences between allelic cistrons or genes would therefore be expected to result in molecular differences in the cortex and thus to determine at least some cortical differences.

Preer (1959), Maly (1958, 1960), Sonneborn (unpublished), and Hanson (1962) have indeed found in *Paramecium* genic or nuclear differences that determine differences in visible cortical structures or in their morphogenetic processes. These findings have to be accommodated in reasonable speculations about the decisive determinative molecular organization of the cortex and the modalities of its action.

Perhaps the main clue—aside from the demonstrated fact that the cortex is essential and determinative—lies in information bearing on the limits of nuclear action. It should be obvious, but commonly is not recognized as such, that the nucleus alone cannot make its corresponding cell or any cell at all (Sonneborn, 1951). This is not merely because the rest of the cell provides a limiting membrane with the function of regulating ingress and egress of materials, for there are regionally diverse decisive areas in the cortex of a single cell. The decisive contribution of the cell cortex is thus its specific organization. Specificity of cytoplasm is correlated with that of the nucleus, for chromosomes in the cytoplasm of a closely related species often cannot function normally and are injured or destroyed (Levine, 1953; Hennon, 1962). This implies that there must be, for cellular existence, a delicate correlation between genotype and plasmatype which is lethally disrupted by relatively minor changes in either and which slowly co-evolves (see above). This intricate intracellular coordination, the result of ruthless long selection, offers little hope that man could ever devise a noncellular milieu in which a nucleus could operate so as to make its own cell or any cell at all.

These considerations leave but few sorts of speculative possibilities for the mode of operation of the cortex. On the one hand, the diverse parts of the cortex might be the seat of production of certain molecules not directly or indirectly producible by the action of genes (and food) alone. However, there is at present, so far as I am aware, no evidence for the production by cells of specific molecules independently of those taken in as food or formed directly or indirectly by genic action. On the other hand, the diverse parts of the cortex (and other preformed cytoplasmic structures) might be the seat of specific absorption and orientation of molecules derived from the milieu and genic action. This alternative is more in harmony with present knowledge. Preexisting cortical structure would then play its essential part by determining where some gene products go in the cell, how these combine and orient, and what they do (Sonneborn, 1951; Tartar, 1961). Apparently such specific assemblies could confer new properties upon molecular groupings. A contemporary example of such origin of new properties appears to be provided by transfer

RNA's. Their triplets appear to be capable of uniting effectively with their complementary triplets in messenger RNA only after the transfer RNA's have combined with their amino acids. Otherwise the messenger could be clogged with blanks. In like manner, one may suppose that the gene products which unite with molecules of the cortex acquire specific activities as a result of the union. The various developmental and genetic events which are regionally localized in different parts of the cortex may thus be dependent upon specific molecular combinations between newly formed molecules deriving from the genes and preexisting molecular patterns already present in the cortex.

This conception further agrees with the contemporary view that function is intimately connected with molecular structure. Preexisting structure determines processes that lead to different structures and different processes in sequences that are self-determined at every step and that lead back cyclically to the starting point. This dynamic interplay of structure and process contrasts with the static view of an unchanging, persistent, fundamental ground substance or organization of the cytoplasm which always underlies the developmental and regenerative capacities of the cell (Calkins, 1926b; Ephrussi, 1952; Fauré-Fremiet, 1950; 1954; Harrison, 1945). Present knowledge rather indicates a cyclically changing scene dependent at every level on pre-existing structure, from the preformed enzymes and ribosomes that are indispensable for genic replication and action to the organized cortical fields, gradients, and localized inductor-response systems required for morphogenesis and hereditary transmission of cortical structure and action systems.

At this stage in discovery, we can only formulate the systems in such general terms. The much more difficult task for the future is to define and specify in molecular terms the decisive structures, gradients, and inductor-response systems and to reveal how specific absorption, orientation, and activation of migratory molecules leads to visible morphogenesis and genetic stability of cell organization.

References

- Calkins, G. N. (1925), *Uroleptus mobilis*. V. The history of a double organism, *J. Exp. Zool.*, **41**:191-213.
- (1926a), *Biology of the Protozoa*, Lea & Febiger, Philadelphia.
- (1926b), Organization and variation in Protozoa. *Sci. Monthly*, **22**:341-351.
- Chatton, E. (1921), Réversion de la scission chez les Ciliés. Réalisation d'individus distomes et polyenergides de *Glaucoma scintillans* se multipliant indéfiniment par scissiparité. *Compt. Rend. Acad. Sci.*, **173**:393.
- , and A. Lwoff (1930), Imprégnation, par diffusion argentique, de l'infracilia-

- ture des Ciliés marins et d'eau douce, après fixation cytologique et sans desiccation, *Compt. Rend. Soc. Biol.*, **104**:834.
- Cleland, R. E. (1962), Plastid behavior in North American Euoenotheras, *Planta*, **57**:699-712.
- Corliss, J. O. (1961), *The Ciliated Protozoa*, Pergamon Press, New York.
- Curtis, A. S. G. (1960), Cortical grafting in *Xenopus laevis*, *J. Embryol. Exp. Morphol.*, **8**:163-173.
- Dalcq, A., and J. Pasteels (1937), Une conception nouvelle des bases physiologiques de la morphogénèse, *Arch. Biol. (Paris)*, **48**:667-710.
- , and ——— (1938), Potentiel morphogénétique, régulation et "Axialgradients" de Child, *Bull. Acad. Méd. Belg. 6^e Sér.*, **3**:261-308.
- Dawson, J. A. (1920), An experimental study of an amiconucleate Oxytricha. II. The formation of double-animals or twins, *J. Exp. Zool.*, **30**:129-157.
- De Harven, E., and W. Bernhard (1956), Étude au microscope électronique de l'ultrastructure du centriole chez les Vertébrés, *Z. Zellforsch. Mikroskop. Anat., Abt. Histochem.*, **45**:378-398.
- Dippell, Ruth V. (1955), Some cytological aspects of aging in variety 4 of *Paramecium aurelia*, *J. Protozool.*, **2** (Suppl.):7.
- (1962), The site of silver impregnation in *Paramecium aurelia*, *J. Protozool.*, **9** (Suppl.): 24.
- Ehret, C. F. (1960), Organelle systems and biological organization, *Science*, **132**: 115-123.
- , and E. L. Powers (1959), The cell surface of *Paramecium*, *Intern. Rev. Cytol.*, **8**:97-133.
- Ephrussi, B. (1952), *Nucleo-cytoplasmic Relations in Microorganisms*, Oxford University Press, Fair Lawn, N.J.
- Fauré-Fremiet, E. (1945), Symmétrie et polarité chez les Ciliés bi- ou multicomposites, *Bull. Biol. Fr. Belg.*, **79**:106-150.
- (1948a), Doublets homopolaires et régulation morphogénétique chez le Cilié *Leucophrys patula*, *Arch. Anat. Microscop. Morphol. Exp.*, **37**:183-203.
- (1948b), Les mécanismes de la morphogénèse chez les Ciliés, *Folia Bio-theoretica*, **3**:25-58.
- (1950), Problème moléculaire de la morphogénèse, *Ann. Biol.*, **26**:361-379.
- (1954), Les problèmes de la différenciation chez les protistes, *Bull. Soc. Zool. Fr.*, **79**:311-329.
- Frankel, J. (1960a), Morphogenesis in *Glaucoma chattoni*, *J. Protozool.*, **7**:362-376.
- (1960b), Effects of localized damage on morphogenesis and cell division in a ciliate, *Glaucoma chattoni*, *J. Exp. Zool.*, **143**:175-194.
- (1961), Spontaneous astomaty: loss of oral areas in *Glaucoma chattoni*, *J. Protozool.*, **8**:250-256.
- Hanson, E. D. (1955), Inheritance and regeneration of cytoplasmic damage in *Paramecium aurelia*, *Proc. Nat. Acad. Sci. U.S.*, **41**:783-786.
- (1962), Morphogenesis and regeneration of oral structures in *Paramecium aurelia*, *J. Exp. Zool.*, **150**:45-68.
- Harrison, R. G. (1945), Relations of symmetry in the developing embryo. *Trans. Conn. Acad. Arts and Sci.*, **36**:277-330.
- Hemmon, Sally (1962), A cytological and embryological analysis of the nuclear changes occurring in a lethal amphibian hybrid (*Rana sylvatica* female by *R. pipiens* male), Dissertation, Indiana University.

- Hertwig, R. (1899). Über die Conjugation der Infusorien. *Abhandl. k. bayer. Akad. Wiss.*, II Cl., 17:151-233.
- Hiwatashi, K. (1955). Studies on the conjugation of *Paramecium caudatum*. VI. On the nature of the union of conjugation. *Sci. Rep. Tôhoku Univ.*, 4th ser., *Biology*, 21:207-218.
- Jennings, H. S. (1908). Heredity, variation and evolution in Protozoa. I. The fate of new structural characters in *Paramecium*, in connection with the problem of the inheritance of acquired characters in unicellular organisms. *J. Exp. Zool.*, 5:577-632.
- (1944). *Paramecium bursaria*: life history. I. Immaturity, maturity and age. *Biol. Bull.*, 86:131-145.
- Kimball, R. F. (1941). Double animals and amiconucleate animals in *Euplotes patella* with particular reference to their conjugation. *J. Exp. Zool.*, 86:1-33.
- Levine, M. (1953). The interaction of nucleus and cytoplasm in the isolation and evolution of species of *Paramecium*. *Evolution*, 7:366-385.
- Lwoff, A. (1950). *Problems of Morphogenesis in Ciliates*, John Wiley & Sons, Inc., New York.
- Maly, R. (1958). Eine genetische bedingte Störung der Zelltrennung bei *Paramecium aurelia*. *Z. Vererbungslehre*, 89:397-421.
- (1960). Die Normalisierung genetisch bedingter Defekte der Zelltrennung bei *Paramecium aurelia* durch Sauerstoffmangel und Kohlenmonoxyd. *Z. Vererbungslehre*, 91:226-236.
- Margolin, P. (1954). A method of obtaining amacronucleated animals in *Paramecium aurelia*. *J. Protozool.*, 1:174-177.
- Maupas, E. (1888). Recherches expérimentales sur la multiplication des infusoires ciliés. *Arch. Zool. Exp. Gen.* (2), 6:165-277.
- Mazia, D. (1961). Chapter 2, in *The Cell*, ed. by J. Brachet and A. E. Mirsky, Academic Press, Inc., New York, vol. 3, pp. 77-394.
- Moscona, A. (1957). The development *in vitro* of chimeric aggregates of dissociated embryonic chick and mouse cells. *Proc. Nat. Acad. Sci. U.S.*, 43:184-194.
- Nadler, J. E. (1929). Notes on the loss and regeneration of the pellicle in *Blepharisma undulans*. *Biol. Bull.*, 56:327-330.
- Nobili, R. (1961a). L'azione del gene *am* sull'apparato nucleare di *Paramecium aurelia* durante la riproduzione vegetativa e sessuale in relazione all'età del clone ed alla temperatura di allevamento degli animali. *Caryologia*, 14:43-58.
- (1961b). Variazioni volumetriche del macronucleo e loro effetti nella riproduzione vegetativa in *Paramecium aurelia*. *Atti Soc. Toscana Sci. Nat. (Ser. B)*: 217-232.
- Porter, E. D. (1960). The buccal organelles in *Paramecium aurelia* during fission and conjugation, with special reference to the kinetosomes. *J. Protozool.*, 7:211-217.
- Powers, E. L. (1943). The mating types of double animals in *Euplotes patella*. *Am. Midland Naturalist*, 30:175-195.
- Preer, J. R. (1959). Nuclear and cytoplasmic differentiation in the Protozoa. in *Developmental Cytology*, ed. by Dorothea Rudnick. The Ronald Press Company, New York, pp. 3-20.
- Roque, Madeleine (1956a). L'évolution de la ciliature buccale pendant l'autogamie et la conjugaison chez *Paramecium aurelia*. *Compt. Rend. Acad. Sci.*, 242:2592-2595.

- (1956b). La stomatogénèse pendant l'antogamie, la conjugaison et la division chez *Paramecium aurelia*, *Compt. Rend. Acad. Sci.*, **243**:1564-1565.
- Sonneborn, T. M. (1930). Genetic studies on *Stenostomum incaudatum*. II. The effects of lead acetate on the hereditary constitution, *J. Exp. Zool.*, **57**:409-439.
- (1932). Experimental production of chains and its genetic consequences in the ciliate protozoan, *Colpidium campylum* (Stokes), *Biol. Bull.*, **63**(2):187-211.
- (1942). Double animals and multiple simultaneous mating in variety 4 of *Paramecium aurelia* in relation to mating types, *Anat. Record*, **84**(4):29-30.
- (1943). Gene and cytoplasm. I. The determination and inheritance of the killer character in variety 4 of *Paramecium aurelia*, *Proc. Nat. Acad. Sci. U.S.*, **29**:329-338.
- (1946). Inert nuclei: inactivity of micronuclear genes in variety 4 of *Paramecium aurelia*, *Genetics*, **31**:231.
- (1947). Recent advances in the genetics of *Paramecium* and *Euplotes*. *Advan. Genet.*, **1**:263-358.
- (1950). Methods in the general biology and genetics of *Paramecium aurelia*, *J. Exp. Zool.*, **113**:87-148.
- (1951). The role of the genes in cytoplasmic inheritance, in *Genetics in the 20th Century*, ed. by L. C. Dunn, The Macmillan Company, New York, chap. 14.
- (1954a). Patterns of nucleocytoplasmic integration in *Paramecium*, *Caryologia*, vol. suppl.:307-325.
- (1954b). Gene-controlled aberrant nuclear behavior in *P. aurelia*, *Micr. Gen. Bull.*, **11**:24-25.
- (1954c). The relation of autogamy to senescence and rejuvenescence in *Paramecium aurelia*, *J. Protozool.*, **1**:38-53.
- (1960). The gene and cell differentiation, *Proc. Nat. Acad. Sci. U.S.*, **46**:149-165.
- , and Ruth V. Dippell (1960a). The genetic basis of the difference between single and double *Paramecium aurelia*, *J. Protozool.*, **7**(Suppl.):26.
- , and ——— (1960b). Cellular changes with age in *Paramecium*, in *The Biology of Aging*, ed. by B. I. Strehler et al., AIBS, Washington, D.C., p. 285.
- , and ——— (1961a). The modes of replication of cortical organization in *Paramecium aurelia*, syngen 4, *Genetics*, **46**:899-900.
- , and ——— (1961b). Self-reproducing differences in the cortical organization in *Paramecium aurelia*, syngen 4, *Genetics*, **46**:900.
- , and ——— (1961c). The limit of multiplicity of cortical organelle systems in *P. aurelia*, syngen 4, *Am. Zool.*, **1**:390.
- , and ——— (1962). Two new evidences of cortical autonomy in syngen 4 of *Paramecium aurelia*, *J. Protozool.*, **9** (Suppl.): 28.
- , and Arlene Lesuer (1948). Antigenic characters in *Paramecium aurelia* (variety 4): determination, inheritance and induced mutations, *Am. Naturalist*, **82**:69-78.
- , and Myrtle V. Schneller (1955). The basis of aging in variety 4 of *Paramecium aurelia*, *J. Protozool.*, **2** (Suppl.):6.
- Stevens, Nellie M. (1903). Notes on regeneration in *Stentor coerulesus*, *Arch. Entwicklungsmech. Org.*, **16**:461-475.
- Steward, F. C. (1961). Growth induction in explanted cells and tissues: metabolic and morphogenetic manifestations, in *Synthesis of Molecular and Cellular*

- Structure*, ed. by Dorothea Rudnick. The Ronald Press Company, New York, chap. 8, pp. 193-246.
- Stubbe, W. (1960). Untersuchungen zur genetische Analyse des Plastoms von *Oenothera*, *Z. Botau.*, **48**:191-218.
- Suzuki, S. (1957). Morphogenesis in the regeneration of *Blepharisma undulans japonicus* Suzuki, *Bull. Yamagata Univ., Nat. Sci.*, **4**:85-192.
- Tartar, V. (1941). Intracellular patterns: facts and principles concerning patterns exhibited in the morphogenesis and regeneration of ciliate Protozoa, *Growth* **5** (Suppl.): 21-40.
- (1954a), Reactions of *Stentor coeruleus* to homioplasic grafting, *J. Exp. Zool.*, **127**:511-575.
- (1954b), Anomalies in regeneration of Paramecium, *J. Protozool.*, **1**:11-17.
- (1956), Grafting experiments concerning primordium formation in *Stentor coeruleus*, *J. Exp. Zool.*, **131**:75-122.
- (1961), *The Biology of Stentor*, Pergamon Press, New York.
- Uhlig, G. (1960). Entwicklungsphysiologische Untersuchungen zur Morphogenese von *Stentor coeruleus*, *Arch. Protistenk.*, **105**:1-109.
- Weiss, P. (1939). *Principles of Development*, Holt, Rinehart and Winston, Inc., New York.
- Weisz, P. B. (1951). A general mechanism of differentiation based on morphogenetic studies in Ciliates, *Am. Naturalist*, **85**:293-311.
- (1954). Morphogenesis in Protozoa, *Quart. Rev. Biol.*, **29**:207-229.
- (1962). From cell to molecule, in *The Molecular Control of Cellular Activity*, ed. by J. M. Allen, McGraw-Hill Book Company, Inc., New York, chap. 1, pp. 1-72.

Microenvironmental Influences in Cytodifferentiation

Clifford Grobstein

**Department of Biological Sciences
Stanford University, Stanford, California**

My objective is to present some facts which suggest that the materials between, and normally closely associated with, the surfaces of cells may be important in controlling cytodifferentiative behavior. Since cytodifferentiation and organismal diversification are opposite faces of the developmental coin, controls for the one must underlie the regularities of the other. There is a large body of information indicating that cytodifferentiation is sensitive to extrinsic influence. However, my emphasis here is not on extrinsic factors generally, but on those which reside in the immediate vicinity of the cell, in the large-molecular materials which are produced by the cell or its close neighbors and which may continuously interact with their source subsequent to production. It is in these terms that I define what is referred to in this discussion as the microenvironment.

I shall limit myself to consideration of two of the several available

sources of information on the developmental role of microenvironment—embryonic induction and the reaggregation of dissociated cells. The information which I shall summarize raises the interesting possibility that there are important relationships among the mechanisms of aggregation, induction, and cytodifferentiative control.

Embryonic induction long has been recognized as relevant to cytodifferentiative control because it involves developmental dependency—the differentiative course of one group of cells in the embryo is different depending upon whether or not it comes into association with a second group. The possibility of microenvironmental effects is raised because the association between the two tissues normally is extremely close.

In lens induction by optic vesicle, for example, the morphological association is so close, as judged by histological sections and by efforts to separate the two mechanically, that mechanisms have been proposed (Weiss, 1958) which assume direct surface-to-surface contact of the cells involved. These assumptions were supported by the finding that interposition of such materials as cellophane between the lentogenic epithelium and the optic vesicle completely blocks the induction (McKeehan, 1951). In recent years, however, evidence has been accumulating that actual contact is not required in certain inductive systems, providing the interspace between has suitable properties. It appears, nonetheless, that even under these circumstances there is a limit to the separation distance across which the reaction can proceed. I should like, first, to recall some of the evidence which seems to justify these statements.

Let me begin by introducing the experimental system which has so far been most useful in our laboratory. It involves the rudiment of the definitive kidney or metanephros of the mouse. In the 11-day embryo the rudiment consists of two components, the outer metanephrogenic mesenchyme and the inner ureteric bud. It is important to recall that the ureteric bud branches to form the collecting ducts of the kidney, while the outer mesenchyme contributes the glomeruli and secretory tubules. If the intact rudiment is placed under suitable conditions *in vitro*, each component undergoes continued morphogenesis in sufficiently normal fashion to be recognized in the developing culture and in subsequent histological sections (Figs. 1 to 6). If the rudiment is first treated with trypsin, however, and the ureteric bud removed, the metanephrogenic mesenchyme alone entirely fails to undergo kidney-type morphogenesis under the same culture conditions. If the separated mesenchyme is recombined in culture with ureteric bud, or with the dorsal region of embryonic spinal cord,

rudiments of secretory tubules appear (Fig. 7). These facts, together with others previously gathered on kidney development in the intact organism, demonstrate that epithelial tubule formation by nephrogenic mesenchyme is dependent upon an outside contribution. Normally this is assumed to be provided by the ureteric bud; experimentally it can also be supplied by dorsal spinal cord.

For reasons of convenience, it is the combination of dorsal spinal cord and metanephrogenic mesenchyme which has received most attention. When a fragment of cord is included among fragments of metanephrogenic mesenchyme, secretory tubule rudiments appear in the mesenchyme after 30 to 40 hours of incubation. It is important to note that these rudiments do not appear generally in the mesenchyme but only in immediate association with the surface of the spinal cord. I say "immediate association" now; when the observation was first made the relationship suggested "contact." I note this to emphasize that the system under consideration shows the usual characteristic of intimacy in embryonic induction.

Having such an inductive system under reasonable control *in vitro*, the next question is the one of separability of components. It turned out that if dorsal spinal cord and metanephrogenic mesenchyme were cultured "back to back" on a highly porous membrane filter (millipore), secretory tubule rudiments appeared in the metanephrogenic mesenchyme immediately opposite the spinal cord (Fig. 8). I say "immediately opposite" to emphasize again that the tubules do not appear just anywhere in the mesenchyme; they form as close to the cord as they can under the circumstances, separated only by the thickness of the membrane filter. The filter used in the early experiments was approximately 20 microns thick, with average pore diameter in the vicinity of $\frac{1}{2}$ micron. By examining sections with the optical microscope, particularly under phase conditions, it was possible to discern materials penetrating into the filter. Electron microscopy showed that some of this material, at least, was cytoplasmic—so one could not say with certainty with this filter that a 20-micron separation between cell surfaces was achieved. Subsequent experiments, however, with filters of lesser porosity (in the vicinity of 0.1 micron), showed that cytoplasm-free separation of at least 20 microns can be achieved without blocking the induction, though the intensity is reduced.

This last point, the reduction of intensity at 20 microns by membranes of porosity in the vicinity of 0.1 micron, raises a red flag to easy assumptions that separability implies free diffusion and indefinite mobility on the part of the active materials. The caution is

emphasized by measurements made on the transmission distance of the inductive activity, even using higher-porosity filters which allow relatively free entry of cytoplasmic processes into the interspace. Measurements were performed by interposing either single filters of different thickness or multiple layers of filters of the same thickness (Figs. 9 and 10). When the response, expressed either as thickness of the metanephrogenic layer or as occurrence of tubules, was plotted

PLATE I. Developing metanephric rudiment of the 11-day mouse embryo in vitro. ($\times 38.4$.) FIG. 1. Immediately after explantation. FIG. 2. On the second day. FIG. 3. On the fourth day. FIG. 4. On the seventh day.

PLATE II. Metanephros of the 11-day mouse embryo cultured for 8 days on a millipore filter platform. Note glomeruli and epithelial tubules. FIG. 5. Fixed and stained whole mount. FIG. 6. Fixed and stained section.

PLATE III. FIG. 7. Dorsal spinal cord in metanephrogenic mesenchyme. Note induced tubules surrounding cord.

against intervening distance, an "inductive activity" curve was obtained (Fig. 11). This declined sharply beyond about 30 microns and reached the level of nondetectability, by the criteria used, at 60 to 80 microns. The close association of induced structure to inductive source thus appeared not to be based on a requirement for surface-to-surface contact, but on limited transmission distance in active form of the inductive material. Stated in another way, we seem to be dealing with materials which are not of the cell surface itself, but which normally are closely associated with it. In turn, this would imply that such surface-associated, microenvironmental materials are important in controlling differentiative processes.

We would, of course, like to know very much more about these

PLATE IV. Transfilter induction by dorsal spinal cord of epithelial tubules in metanephrogenic mesenchyme. FIGS. 8, 9. Positive effect with two filter layers. FIG. 10. Absence of effect with four filter layers.

8

9

10

materials, how they are produced and how they operate. It seems clear that essential components of the material are large-molecular. This is suggested by the restraining effect, already mentioned, of filter porosities in the 0.1 micron range. Sieving in terms of molecular size is not by any means the only mechanism by which the filter may restrain mobility, but it is noteworthy that 0.1 micron is quite large in terms of molecular dimensions. Moreover, cellophane blocks the inductive effect. This can be shown by simply interposing cellophane

PLATE V. FIG 11. Effect of separation distance (filter thickness) on metanephrogenic mesenchyme thickness and tubule formation. *AA*, *TA*, *TH*, filters of different porosity and thickness. Each point represents an average of four or five cases.

between the interactants, but it is shown more dramatically by making use of the multiple-filter technique, as in the measurement of transmission distance. If a three-layer assembly is used, the middle layer of which consists of cellophane with a small hole (Fig. 12), a localized induction is obtained only immediately over the hole (Fig. 13). This seems to say not only that the cellophane is "opaque" to the activity, but that it comes through the small pinpoint hole to form a "bright" spot on the opposite side. It is interesting that this "shadowing" effect of cellophane is not only what would be expected

FILTER-CELLOPHANE ASSEMBLY

12

13

PLATE VI. FIG. 12. Diagram of cellophane with small hole interposed between inductive interactants. FIG. 13. Fixed and stained whole mount. Note edge of hole in cellophane marked by arrows, epithelial rudiments forming.

of large molecules, but it suggests that mobility of the material is primarily vertical in the filter substance with little lateral component. This is in conformity with what is believed by the manufacturers to be the structure of the filters used. It may be an important characteristic for obtaining the transfilter inductive effect, since it would tend to reduce dilution of materials in the interspace by the general environment.

It also seems clear that protein is a constituent of the interspace material. At an early stage in the study of the transfilter effect, it was noted that the filter interspace between the tissues in sections differed

PLATE VII. Residual spot in filter. FIG. 14. Before exposure to trypsin. FIG. 15. After exposure to trypsin.

slightly in staining and optical properties from the filter lateral to the tissues. This led to studies of spinal cord incubated on filters in the absence of metanephrogenic mesenchyme, in an effort to detect inductive activity or material "secreted" by the spinal cord into the filter. Such preparations, after incubation for 20 to 24 hours, were fixed in alcohol-formalin. The spinal cord was found to be tightly cemented to the filter, so tightly that it could be removed only by scraping and fragmentation. When virtually all of the tissue had been scraped away, a translucent spot was left, clearly different from the surrounding filter (Fig. 14). The spot stained occasionally but variably with the periodic acid-Schiff procedure. When spotted filters

were placed in trypsin, the spot gradually disappeared over a 10-minute interval (Fig. 15). Curiously, the spinal cord could be removed from the filter relatively easily before fixation, and such filters following fixation showed no sign of a spot. In general, the behavior suggested deposition in the filter of a mucoid slime. Neither unspotted filters, however, nor spotted filters following careful washing to remove fixative had any inductive effect on metanephrogenic mesenchyme subsequently added.

The impression that protein is a constituent of the interspace material is strengthened by recent autoradiographic studies in which dorsal spinal cord was labeled by exposure to tritiated amino acids, before being put into culture on the membrane in the presence or absence of metanephrogenic mesenchyme. After varying periods of incubation, the cultures were fixed in alcohol-formalin, sectioned, and processed for autoradiography. The presence of silver grains in the overlying radiation-sensitive film marks the location of the tritium and, because tritium emission has low energy and a short emission path, the resolution of the method is high. Dorsal spinal cord becomes very heavily labeled during incubation for 2 hours in the tritiated amino acid solution. This labeling, presumably primarily of proteins in the tissue, is indicated by marked blackening of the film immediately over the cord (Fig. 16). More interesting, however, is the fact that in cultures incubated for 24 hours there is evident radioactivity in the filter interspace beyond the tissue, as well as in the metanephrogenic mesenchyme on the opposite side. Grain counts over the filter gave values 7 to 16 times background count, and these indications of radioactivity were limited to the general region previously identified as the "spot." It thus appears that administration of labeled protein precursors to the spinal cord deposits label in the filter in a form fixable by alcohol-formalin. It remains to be demonstrated that the tritium-labeled material and the trypsin-sensitive material are one and the same, but it will be more surprising if they are not than if they are.

Using the grain-counting procedure, it is possible to get an estimate of the amount of labeled material with increasing distance from the spinal cord. For this purpose labeled cord was incubated on a thicker filter, 125 to 150 microns, to provide a distance greater than that which inductive activity will cross. After incubation for varying periods such cultures were autoradiographed, and counts were made on successive 13-micron ranks across the thickness of the filter. Omitting details, it is clear that radioactivity falls off sharply in the first 60 microns and approaches background in the vicinity of 100 microns

(Fig. 17), when higher-porosity filters (about 0.5 micron) are used, and declines very much more rapidly when lower-porosity (about 0.1 micron) filters are used. The impression is that there is quite good agreement between the distribution of label and of inductive activity with distance.

It is interesting that the gradient of label exists not only in the filter but in the surrounding clot in which the tissue is embedded (Fig. 18). In fact, as one looks at the autoradiogram, one notes first

PLATE VIII. FIG. 16. Autoradiogram of transfilter culture involving dorsal spinal cord (below), heavily labeled by exposure to tritiated amino acids. Note grains above filter and over metanephrogenic mesenchyme on opposite side of filter.

the dense and reasonably sharply bounded “picture” of the dorsal cord, and then becomes aware—particularly knowing the result of the grain counts—that there is a much less dense “halo” of grains grading off around it.

Recent data obtained by William Koch, a graduate student in our laboratory, indicate that the label observed in the filter is noncellophane-passing. He has repeated the experiment referred to above involving two filter layers with intervening cellophane. He finds label in the filter on the dorsal cord side of the cellophane, but none in the cellophane or in the filter on the opposite side. Under these circum-

stances label is found in the metanephrogenic mesenchyme but at reduced levels in comparison with noncellophane cultures. This label presumably represents nonfixable small molecular material which passes cellophane.

PLATE IX. FIG. 17. Curve of grain counts across thick filter with increasing distance from spinal cord source labeled with tritiated amino acids.

The entire body of data can be explained on the assumption that dorsal spinal cord produces at free surfaces a protein-containing material which moves into the environment over a limited distance. Among its components this material includes one or more which is inductively active. It is important to note that though inductive activity conforms with the distribution of label, labeling does not necessarily indicate

inductive activity. Labeled metanephrogenic mesenchyme, for example, puts label into the filter and across it into unlabeled dorsal cord, though no *known* inductive activity moves in this direction. Moreover, label is present in the filter after incubation times as short

PLATE X. FIG. 18. Curve of grain counts with increasing distance in clot surrounding dorsal spinal cord labeled with tritiated amino acids.

as 3 hours, but definitive changes in the metanephrogenic mesenchyme are not known to occur until 24 to 30 hours. Keeping these reservations in mind, it is safe only to say that inductive activity appears to be associated with a kind of "halo" of large-molecular material, including protein, which exists in the immediate vicinity of dorsal spinal cord.

These results fortify the notion, earlier advanced by Baitsell (1925), Weiss (1933), Huzella (1941), and others, that microenvironmental materials produced by cells have important developmental significance. Additional support is afforded by data from the second major line of investigation to which I referred earlier—the reaggregation of dissociated cells. This phenomenon was studied first, among animals, in sponges and other relatively uncomplicated invertebrates. More recently it has been under investigation in slime molds, in amphibia, and even in higher vertebrates. Observations on reaggregating mouse and chick cells in the last several years, reported by Moscona (1960), are particularly relevant and striking. Moscona finds that at an early stage in the reaggregation of freshly dispersed chick embryonic cells a transparent mucoid material, “evidently an exudate of cellular origin,” makes its appearance. The reaggregating cells appear to move within fine strands of this material, which collects on introduced cotton fibers and on the glass floor of the vessel and appears to be the “immediate substratum or microenvironment” for migration and aggregation. Staining reactions and susceptibility to enzymatic digestion suggest that the material probably is a mucoprotein.

Freshly disaggregated cells produce considerable quantities of mucoid; it can be accumulated in strands or ropes by gently swirling reaggregating cells toward a central vortex. Cells which are grown as monolayer cultures for a period of time undergo change in the quality, and reduction in the quantity, of the mucoid material they produce. Concomitantly, aggregative behavior and cell coherence gradually decline. Thus, dissociated mouse embryonic skin cells, after more than 10 days in monolayer culture, form only small clusters, even when they are swirled into proximity or centrifugally packed to favor aggregation. Such clusters show very limited histogenesis and differentiation. The aggregation and histogenesis are little affected when the cells are intermixed with freshly dissociated chick or mouse precartilaginous cells (heterotypic), but are markedly improved on intermixing with freshly dissociated chick skin cells (homotypic). Under the latter conditions the mouse cells are found grouped together and participating in the formation of follicle-like structures.

The decline in aggregability and histogenesis with culturing described by Moscona recalls and illuminates an observation made some years ago on embryonic mouse salivary rudiments *in vitro*. The salivary rudiment is a two-component structure like the metanephros and, similarly, the components can be separated by trypsin treatment. When the two components are recombined immediately, the inner

epithelial one continues the dichotomous branching characteristic of normal morphogenesis. When, however, the mesenchymal component is cultured over a 20-day period, its capacity to support the dichotomous branching of freshly isolated epithelium gradually declines. Accompanying the decline there is a change in the outgrowth pattern of the mesenchyme, from sheet-like behavior with cohesive boundary to increasingly dispersive behavior with individually migrating fibroblast-like cells at the boundary (Fig. 19). Here a behavioral change, which may well conform with Moscona's observation of decline in aggregability and production of extracellular material, is accompanied by altered morphogenetic effect on the associated dependent epithelium.

Note that in Moscona's experiments aggregative and differentiative behavior are simultaneously altered during the period in monolayer culture. This is, of course, in conformity with many observations that culturing, under conditions which promote cell dispersal, tends to be antidifferentiative (Grobstein, 1959). A well-documented example of this has just been reported by Holtzer et al. (1960). They examined chondrocytes freed from already formed cartilage. Over a period of 10 days of culture in a dispersed state, these cells lost all detectable cartilage properties, as judged by their ability to incorporate S^{35} or to form recognizable matrix when returned to conditions under which freshly dissociated cells undergo chondrogenesis. Although changes in aggregative behavior were not stressed in the study, it is interesting that freshly liberated chondrocytes on a clot spontaneously aggregate within 24 hours and show little spreading tendency. After several passages in culture, while the differentiative properties were declining, the cells failed to aggregate spontaneously, forming a thin, rapidly spreading sheet.

There is, therefore, good reason to suspect a relationship between the mucoid, microenvironmental material and differentiation. Results reported by Wilde (1960) may offer the most direct demonstration to date that such a relationship actually exists. Working with 1 to 15 cells from early amphibian embryos in restricted, microdrop nutrient environments, he found that under particular conditions none of the cells differentiated. When, however, the microdrops containing similar cells also included "the opalescent, slightly viscous material which poured off the cells in the disaggregating fluid," differentiation of the cells was normal and similar to that seen in organotypic cultures of the source tissue. Wilde reports further data, involving combination of cells with extracellular material from heterotypic sources, suggesting that the material may not only be necessary for differentiation but

DIRECT

1ST GENERATION

2ND GENERATION

3RD GENERATION

PLATE XI. FIG. 19. Upper row, freshly isolated salivary epithelium combined with salivary mesenchyme cultured for increasing numbers of culture generations (culture intervals of 5 to 6 days between transfers). Lower row, mesenchymal border of the respective cultures. Note change in epithelial morphogenesis and associated change in mesenchymal cohesiveness.

may contain informational cues as to direction of differentiation. Chemical data on the nature of Wilde's material, particularly in relation to Moscona's and to the conditioned medium of Niu (1956), will be awaited with great interest.

What do these facts add up to? Certainly not to definitive statement about the mechanisms of induction, of aggregation, or of cyto-differentiative control. They point, rather, to continued examination of the couplings among these three phenomena, with emphasis on the possibility that they share a common basis in microenvironmental large-molecular materials. If it be assumed that these materials simultaneously are cell product and cell controls—both for the source cells and their immediate neighbors—they would then provide the basis for “external feedback” and a common matrix underlying the integral character of many multicellular systems. How such materials operate, if they do, in controlling intrinsic cellular activities is not at all clear. In some fashion they must mesh with intracellular control mechanisms radiating outward from the replicative genetic code. A number of models are available from which to choose. They might operate completely superficially, as “gatekeepers” on a permease model or the modified molecular ecology model of Weiss. They might carry “instructions” more deeply into the cell, to interact with the genetic sites themselves, with messenger materials, or with cytoplasmic synthetic sites. Or, an idea which attracts me particularly at the moment, they might operate at or close to the cell boundary, modifying the many subtle factors which influence polymerization and complexing of monomers synthesized deeper in the cell. Whatever is the answer, and most likely there will be more than one, when we have it we shall be one step closer to rationalizing the mutual operation of replicative and regulative factors in the control system of the cell.

References

- Baitsell, G. A. (1925). On the origin of the connective-tissue ground-substance in the chick embryo. *Quart. J. Microscop. Sci.*, **69**:571-590.
- Grobstein, C. (1959). Differentiation of vertebrate cells, in *The Cell*, ed. by J. Brachet and A. E. Mirsky, Academic Press, Inc., New York, vol. 1, pp. 437-496.
- Holtzer, H., J. Abbott, J. Lash, and S. Holtzer (1960). The loss of phenotypic traits by differentiated cells in vitro. I. Dedifferentiation of cartilage cells. *Proc. Nat. Acad. Sci. U.S.*, **46**:1533-1542.
- Huzella, T. (1941). *Die Zwischenzellige Organisation*, Verlag von Gustav Fischer, Jena.
- McKeehan, M. S. (1951). Cytological aspects of embryonic lens induction in the chick. *J. Exp. Zool.*, **117**:31-64.
- Moscona, A. A. (1960). Patterns and mechanisms of tissue reconstruction from dis-

sociated cells, in *Developing Cell Systems and Their Control*, ed. by D. Rudnick, The Ronald Press Company, New York.

Niu, M. C. (1956), New approaches to the problem of embryonic induction, in *Cellular Mechanisms in Differentiation and Growth*, ed. by D. Rudnick, Princeton University Press, Princeton, N.J.

Weiss, P. (1933), Functional adaptation and the role of ground substances in development, *Am. Naturalist*, **67**:322-340.

——— (1958), Cell contact, *Intern. Rev. Cytol.*, **7**:391-423.

Wilde, C. E. (1960), Factors concerning the degree of cellular differentiation in organotypic and disaggregated tissue cultures, *Colloq. intern. sur la Cult. Organo-typique* (in press).

Differentiation and Morphogenesis in Insects

Carroll M. Williams

The Biological Laboratories
Harvard University, Cambridge, Massachusetts

Diversity is introduced into biological systems by the slow-paced events of organic evolution and by the swift unfolding of the individual life. In the diversifications which shape the egg into an organism, what we see is the implementation at the cellular level of species-specific information inherited from the preceding generation. We may think of the genes as supplying a kind of "construction manual" which is translated into action during the course of development. But the construction manual itself is the product of trial-and-error testing in the lives of tens of thousands of individuals over enormous periods of time. Therefore, the two kinds of diversifications, developmental and evolutionary, come together in the management of genetic information—its genesis, implementation, and testing in the careers of the individual and the species.

The construction manual (to continue the analogy) is "read" by the individual cells, which in all but the earliest embryonic stages are numbered in the millions. From a fairly early stage in embryonic development we can separate these cells into a series of cytological types—muscle, nerve, epithelium, and so on. What begins as a unity rapidly becomes a multiplicity of vastly different forms and functions.

All these cells, as far as we know, continue to possess a full set of genes, corresponding to the construction manual for the organism as a whole. And yet, during the course of embryonic development, the construction "crew" resolves itself into clones of diverse cell types, which (in the manner of bricklayers, plumbers, and electricians) make use of only their own specific assignments in the construction manual.

The central dogma

In the interpretation of these happenings we come face-to-face with a series of concepts which everyone will recognize as a central dogma of present-day biology. At the risk of recapitulating what has gone before, let me briefly review this theory.

The hereditary information of the genes is coded in the nucleotide sequence of DNA, presumably in the form of a series of three-letter words made up of the letters A, T, G, and C. At any one time, much and perhaps most of this genetic information is not at the disposal of the synthetic centers of the cell. In one way or another a certain fraction of the DNA is repressed and therefore unexpressed. Under appropriate conditions, specific genes or gene combinations are de-repressed. At these particularly activated sites, the coded information of the DNA is transferred to newly synthesized RNA in the form of a series of three-letter words made up of the letters A, U, G, and C. This so-called "messenger RNA" then passes to the cytoplasm. It enters the ribosomes and each of its three-letter words is used to code the precise position of one of the 20 kinds of amino acids in newly synthesized protein. Once the amino acids have been properly positioned in the peptide chain, the molecules of protein automatically fold into their distinctive so-called "tertiary structure." The protein thereby attains its characteristic enzymatic activity, as dictated by its amino acid sequence and overall shape. And since most, if not all, of the reactions of the cell are catalyzed by enzymes, it is alleged that all happenings in the cell are a sequel to the synthesis of specific enzymes.

Control of genetic information in microbial systems

The central dogma has, more recently, been extrapolated to account for the moment-to-moment control of enzyme synthesis in microbial systems. This extrapolation takes the following form (Jacob and Monod, 1961; Jacob and Wollman, 1961) :

There are certain enzymes which are necessary for the survival of the cell at all times, for example, all the enzymes which lie on the main pathways of intermediate metabolism and energy transformations. These so-called "constitutive enzymes" are continuously synthesized—a state of affairs which signals the activity of the corresponding genes. By contrast, there are a large number of other enzymes which are needed only under certain conditions and circumstances. These "inductive enzymes" are not synthesized until they are needed; therefore, the genes for their synthesis are ordinarily maintained in a state of chronic repression.

The theory tries to account for the ability of specific metabolites to induce the synthesis of specific enzyme systems. It is postulated that the synthetic reactions for groups of enzymes serving a given biochemical pathway are programmed by a collection of genes which are under the control of a single "operator gene," the whole complex being known as an "operon." But under most conditions the operator gene is in combination with a specific "repressor substance," and the corresponding operon is thereby repressed.

Each repressor substance is the synthetic product of a particular "regulator gene." The repressor substance is freely diffusible. If all goes well, it combines with and represses its specific target. But on its way to the operator gene, the repressor substance is subject to inactivation if it encounters and combines with a substrate of the particular enzyme system that is repressed. In this case the operator gene escapes from repression and can go ahead and activate the collection of genes in its operon. So, in this indirect manner, a given substrate induces the synthesis of an appropriate enzyme system.

It is also proposed that certain regulator genes produce inactive repressor substances which can be activated only by combination with specific small molecules. This accounts for the ability of metabolic end products to suppress the on-going synthesis of specific enzyme systems.

In this bird's-eye view of the theory, one cannot fail to be impressed by the increasing attention that centers on mechanisms of control.

Whereas it formerly sufficed to think of genes that program the synthesis of specific enzymes, it is now necessary to postulate several categories of genes whose sole purpose is to control and modulate other genes. If this system of control seems unnecessarily complex, one is well advised to reflect on the metabolic pathways for a miniature molecule such as acetic acid. What if the controlling mechanisms of a cell turn out to be as intricate as the "metabolic maps"?

Microbes versus higher organisms

The central dogma is based on studies of microbes and their viruses. Therefore, the question may well be asked whether it has any bearing on developmental processes in the higher and more pretentious forms of life.

At this point I must confess that I find the concept of regulator genes and operator genes—not to mention the rather mysterious "episomes"—strikingly reminiscent of McClintock's genetic "controllers," including "activators," "modulators," and "dissociators," which were described in 1951 (for review see Brink, 1958). So, the pertinence of these concepts has already been documented in at least one "higher organism," the corn plant.

Despite all this duplication in terminology, I believe that the new point of view has much to tell us about the management of genetic information in animal cells and that a judicious application of its principles can promote a great leap in what may prove to be the forward direction.

As a first and somewhat hesitant step toward this objective, I propose to consider how the new theory, derived largely from studies of bacteria and viruses, may be applied to certain aspects of insect growth and metamorphosis.

Differential gene activity in giant chromosomes

In the banana orchards of Brazil there lives a mosquito-like fly called *Rhynchosciara angelae*. The female of this insect oviposits a cluster of some 150 eggs, all of which hatch simultaneously a few days later. The larvae then assemble into a compact mass and crawl around together as a slug-like object.

The lives of the 150 larvae are wonderfully synchronized. Over a period of a month or so, they grow at the same rate, molt at the same

time, pupate in synchrony, and some days later they all emerge as adult flies within a few minutes of one another. Therefore, as one would hope and as Breuer and Pavan (1955) have confirmed, the activities of their cells and tissues show a degree of synchronization that is unmatched in any other metazoan. Each larva, in effect, is equivalent to the next: so, in principle, one can design experiments in which the same individual is sacrificed 150 times.

As is the practice in most dipteran insects, the growth of larval *Rhynchosciara* takes place solely by cell enlargement, the chromosomes being duplicated again and again without any cell division. Ordinarily, this would lead to a vast increase in chromosome number. However, in *Rhynchosciara* as in most Diptera, the daughter chromatids do not separate following replication. Instead, they align themselves side by side to form compact, ribbon-like or cable-like chromosomes up to 16,000 times larger than ordinary chromosomes.

Within the giant chromosome each chromatid is a much-coiled structure, the tightness of the coil dictating a periodic structure which stains more intensely than do the adjacent loosely coiled regions (Ris and Crouse, 1945). And since the daughter chromatids pair "gene for gene," the net result is that each giant chromosome can be recognized under the compound microscope, not only by its overall size and shape, but also by the pattern of its crossbanding.

In *Rhynchosciara* there are four giant chromosomes in the nucleus of each larval cell. The total number of crossbands is in the thousands: in *Drosophila melanogaster* (where the number has been counted) there are 5,072, corresponding, it is alleged, to the number of gene loci (Bridges, 1942). By taking advantage of the giant size of the chromosomes, one may inquire as to whether a full and complete set of genetic material is distributed to the various cell types during embryonic development.

In *Rhynchosciara*, as in *Chironomus*, the answer to this question is clear-cut. In such diverse cells as those of salivary glands, midgut, rectum, seminal vesicles, and Malpighian tubules, one can identify, not only a full complement of giant chromosomes, but also the characteristic fine structure in homologous parts of homologous chromosomes (Berger, 1940; Slizynsky, 1950; Beerman, 1952; Pavan and Breuer, 1952).

On more detailed cytological examination, one finds that the giant chromosomes of each type of cell are distinguished by the swollen appearance or "puffing" of certain specific bands. This is apparently due to the uncoiling of the chromatids at these localized places—a

process which, more rarely, may proceed to the formation of enormous rosette-like projections called "Balbiani rings" (Beerman, 1952; Mechelke, 1953; Beerman and Bahr, 1954).

Cytochemical studies suggest that the projections from the chromosomes consist of unraveled DNA (Beerman, 1952, 1959; Gall, 1958). Moreover, the individual "puffs" prove to be the sites of intensive synthetic activity, especially of RNA (Beerman, 1952; Pavan and Breuer, 1952; Ficq and Pavan, 1957; Gall, 1958). It is difficult to avoid the conclusion that the puffing phenomenon is, in fact, a caricature of the de-repression of specific gene loci and the coding of messenger RNA.

In general, one can say that the pattern of puffing is uniform throughout a single tissue, but it differs from that in other tissues of the same individual (Beerman, 1952, 1959; Mechelke, 1953; Breuer and Pavan, 1955). Each tissue is, of course, engaged in synthetic operations peculiar to itself—a fact which is apparently reflected in its pattern of puffing.

If attention is centered on a single tissue, say, the salivary glands, it is worth inquiring as to whether the chromosomal puffs undergo any systematic change during larval life. Because of the precise synchronization within each brood of larvae, *Rhynchosciara* is obviously the perfect animal for this type of study (Breuer and Pavan, 1955). However, similar investigations have been carried out with considerable success on the salivary glands of the Chironomidae (Beerman, 1952, 1959; Mechelke, 1953).

Just prior to pupation, the salivary glands show a pronounced change in their synthetic activity, in that the clear secretion which they previously produced is replaced by a brownish fluid which is now synthesized for the first time. In cytological preparations of salivary glands, the new synthetic acts are signaled by striking changes in the pattern of puffing in the giant chromosomes. Most of the puffs that had been prominent during larval life now collapse, the DNA strands apparently being folded back into the chromosome to re-form compact bands (Breuer and Pavan, 1955). Meanwhile, new puffs are formed at a series of other specific loci.

There is a comprehensive body of evidence to show that the change in the secretory activity of the salivary glands is promoted by the same endocrine agents that control the metamorphosis of the animal as a whole. As we shall later consider, pupation is the response to an increasing titer of "ecdysone" and a decreasing titer of "juvenile hormone." Therefore, by injecting ecdysone into immature larvae, one can

establish the endocrine conditions which favor precocious pupation. The question arises as to the behavior of the giant chromosomes under these conditions.

Experiments of this type have been performed on *Chironomus* by Clever and Karlson (1960). Two hours after the injection of ecdyson, the pattern of puffing in the salivary glands shows a changeover from larval to pupal type. Clever and Karlson conclude that the primary effect of ecdyson is to alter the activity of specific genes and that this action is documented in the giant chromosomes.

RNA and the synthesis of silk

In Nigeria and certain other parts of Africa, there occurs a lepidopteran called *Epanaphe moloneyi* whose mode of life bears a certain resemblance to that of *Rhynchosciara*. Here again, the larval stages are gregarious. They move about on the food plant as "processionaries," spinning a thread of silk to guide themselves back to their common nesting place. When the larvae are full grown, the entire group of 100 or more individuals collaborate in spinning a large silken chamber almost the size and shape of a football. Within this chamber, each larva finally spins its own separate cocoon and pupates. After about a month of adult development, the pupae are transformed into moths which escape from the cocoons.

Interest in the *Epanaphe* silkworm stems from the fact that its silk consists essentially of two amino acids, glycine (42.5 moles %) and alanine (53.1 moles %), thereby qualifying as the simplest of naturally occurring proteins (Lucas et al., 1958).

The silk is synthesized and secreted into the lumen of the silk gland by a monolayer of epithelial cells which form the distal regions of the two glands. These cells are extremely rich in ribosomal RNA—a fact which probably accounts for their rapid synthesis of what is essentially a single, diagrammatically simple protein. If the ribosomal RNA is responsible for coding the glycine-alanine sequence in *Epanaphe* silk, this fact should signal itself in certain predictable peculiarities in the ratios of the four bases in RNA: namely, very high ratios of guanine to adenine and of uracil to cytosine (Speyer et al., 1962).

Yčas and Vincent (1960) have tested this proposition by journeying to Africa, collecting large numbers of *Epanaphe* silkworms, and isolating the RNA from the silk-secreting part of the silk gland, as well as from a number of other tissues and organs.

The answer was clear-cut. The base ratios encountered in the silk-

secreting cells showed no correlation with the peculiar features of the synthetic product and, in fact, did not depart from those encountered in the other tissues that were examined.

As Yčas and Vincent (1960) point out, this implies that the "bulk RNA" of the ribosomes does not contain the genetic information which codes the amino acid sequence in silk. This finding is completely in line with that obtained on microbial systems where, as previously mentioned, genetic information is continuously transmitted to the ribosomal "factories" by a certain short-lived messenger RNA which makes up only a tiny fraction of the total RNA.

Metamorphosis of the Cecropia silkworm

In the case of giant chromosomes of *Rhynchosciara*, we considered the evidence for gene repression and de-repression as well as the cytological picture of the synthesis and coding of what appears to be messenger RNA. Then from a study of *Epanaphe* we learned that the ribosomes are "ignorant" synthetic centers to which genetic information is apparently conveyed in the form of messenger RNA. I now propose to consider the harnessing of genetic information to specific morphogenetic acts. And for this purpose I hasten to introduce our final performer.

In fields and forests of eastern North America there occurs a handsome silkworm whose outstanding properties as an experimental object entitle it to rank alongside *Rhynchosciara* and *Epanaphe*. Taxonomists continue to amuse themselves by changing the scientific name of this insect; therefore, let us simply call it the *Cecropia* silkworm.

The eggs are oviposited in early summer on any of a number of common trees and shrubs. Ten days later they hatch into little spinose caterpillars which grow rapidly, punctuating this growth by a series of four larval molts. By transforming leaves into silkworm, they increase their mass some 5,000-fold and finally become mature fifth-instar larvae.

Through the workings of hormonal factors which we shall shortly consider, there comes a day when the behavior of the mature silkworm suddenly changes. It ceases to feed, having already built up within its own tissues a stockpile of molecules to sustain the 10 remaining months of life. After a period of random locomotor activity, it begins to spin a cocoon around itself. Work on the structure continues for 2 days, during which time a silk thread, nearly a mile long, is transformed into a complicated edifice.

The completion of the cocoon signals the beginning of a new and even more remarkable sequence of events. On the third day after the cocoon is finished, a great wave of death and destruction sweeps over the internal organs of the caterpillar. The specialized larval tissues break down. But meanwhile, certain more or less discrete clusters of cells, tucked away here and there within the body, begin to grow rapidly, nourishing themselves on the breakdown products of the dead and dying larval tissues. These are the "imaginal discs" which throughout larval life have been slowly enlarging within the caterpillar. Their spurt of growth now shapes the organism according to a new plan. The specialized structures of the caterpillar are swept away and replaced by new organs arising from the imaginal discs. In addition, some of the less specialized larval tissues, such as the epidermal layer of the abdomen, are transformed directly into pupal tissues. Finally, the old larval cuticle is shed to unmask an essentially new organism, the pupa. All these changes normally take place within the snug confines of the cocoon.

Once the pupa has formed, there is an abrupt halt to the events which during a period of 8 weeks have transformed the egg into a larva and the larva into a pupa. During the months that follow, the insect persists in a kind of suspended development—an obligatory dormancy termed "pupal diapause." The insect overwinters in this state of developmental standstill.

Endocrine control of metamorphosis

The pupal diapause proves to be the result of an endocrine deficiency, namely, a failure of the brain to secrete a "brain hormone" required for the continuation of development (Williams, 1946). The brain hormone is the synthetic product of 26 neurosecretory nerve cells (Williams, 1952a). For reasons that are not understood, the brain's neurosecretory cells are somehow turned off at the time of pupation. In order to get turned on again, they require several months of exposure to low temperatures—a need which is obviously met in the overwintering pupa (Van der Kloot, 1955; Williams, 1956b). When the chilled pupa is returned to a warmer temperature (as normally happens in the spring of the year), the brain secretes its hormone and adult development begins.

By appropriate experiments, it is possible to show that the brain hormone does not exert its effect throughout the insect as a whole. Its primary function is to activate another endocrine organ, the "pro-

thoracic glands" (Williams, 1952b, 1958). And it is this second endocrine organ which then secretes the growth hormone itself—a molecule which has been isolated and crystallized and called "ecdysone." The hormone is a water-soluble, heat-stable, heterocyclic ketone with the provisional formula $C_{18}H_{30}O_4$ (Butenandt and Karlson, 1954; Karlson, 1956).

When exposed to ecdysone, the pupal tissues, after 8 months of developmental standstill, are promptly stimulated to intense morphogenetic activity. The result is a predictable pattern of death and birth at the cellular level as the specialized tissues of the pupa make way for the equally specialized tissues of the adult moth (Williams, 1961a). Spectacular changes occur throughout all parts of the insect: in the head, the formation of compound eyes and feather-like antennae; in the thorax, the molding of legs, wings, and flight muscles; in the abdomen, the shaping of the genitalia and, internally, the exorbitant growth of ovaries and testes. And in the newly formed skin we can witness the strangest behavior of all—the extrusion and transformation of tens of thousands of individual cells into the colorful but lifeless scales so typical of moths and butterflies (Stossberg, 1938).

After 3 weeks of adult development, the process is complete. The full-fledged moth escapes from the cocoon and unfurls its wings.

The genetic construction manual

This summary of the life history of the *Cecropia* silkworm brings us back to the concept of the genetic "construction manual" mentioned at the outset. In the case of the *Cecropia* silkworm, as in all insects that undergo a complete metamorphosis, the construction manual is obviously divided into three distinct chapters.

The first chapter tells how to make the egg into a larva. The second chapter gives directions for the destruction of the larva and the reworking of the same materials into a pupa. The third and final chapter tells how to reshape the pupa into an adult moth. This analogy serves to emphasize that the sequential polymorphism of insect metamorphosis involves the decoding and acting out of what is little short of successive batches of genetic information (Snodgrass, 1954; Wigglesworth, 1961a). Perhaps the day is not far distant when we may speak of "larval operons, pupal operons, and adult operons"—collections of genes which are sequentially brought into play to build and rebuild the insect on the unchanging foundation of the constitutive enzymes prerequisite for life at all stages.

Though these overall happenings are amazing enough, it is well to

recall that the metamorphosis of the animal as a whole is a mosaic of metamorphoses at the level of the individual cells. And here we come face-to-face with the most baffling phenomenon of all. It turns out that the individual cells at an early stage in embryonic development are "determined" or "programmed" for the specific sequence of differentiations which days, weeks, or months later they will act out in the future larva, pupa, and adult. In this sense, the construction manual is more akin to the script of a three-act play in which the individual cells are cast in individual roles.

Metamorphosis of skin implants

The programming of cells is particularly well illustrated in the experiments first performed by Kühn and Piepho (1938, 1940) and subsequently continued in great detail by Piepho and his students at Göttingen (Piepho, 1938a, b; 1943).

If a tiny fragment of skin is cut from a caterpillar and implanted into another caterpillar of the same or even of a different species, the implant not only survives but undergoes a complicated career of growth and metamorphosis in synchrony with that of the host.

The first thing that happens is that the epithelium of the implant grows around to close on itself to form a cyst—a hollow ball with walls only one cell thick. Curiously enough, this growing around occurs in such a manner that the cuticular surface of the epithelium always faces inward to form a lining to the cyst.

Days or weeks later, when the host larva undergoes a larval molt, so does the little hollow ball of cells. The cuticular lining is shed into the lumen and each cell now secretes a new larval cuticle. This shedding and re-forming of larval cuticle is repeated over and over again as long as the host is undergoing larval molts. If the implant is removed prior to the pupation of the host and re-implanted into a sequence of immature larvae, there is apparently no limit to the numbers of larval molts that it can undergo (Piepho, 1943; Piepho and Meyer, 1951). So, as far as its cells are concerned, the larval molts correspond to the use and re-use of the coded instructions for synthesizing and secreting a larval cuticle.

But, in the normal course of events, sooner or later the host metamorphoses to a pupa. So does the implant. The larval cuticle is cast off into the lumen of the cyst and each cell now secretes an overlying island of pupal cuticle. The cyst, in short, has pupated. Manifestly, each cell is now tapping information that had previously been repressed, presumably within its own chromosomes. Still later, when

the pupal host transforms into an adult, the cyst casts off its pupal cuticle and secretes an adult cuticle covered with scales and hairs. Here again, it is necessary to conclude that each epithelial cell is de-repressing a fresh assortment of genetic information.

This experiment teaches us two things. The metamorphosis of the implant shows that each cell possesses a definite repertoire of morphogenetic information—a sequential program which it is able to act out even in the semi-isolated situation. And in the synchronization between the behavior of implant and host, we are reminded once again of the control and coordination which is exercised by hormones.

Endocrine control of genetic information

On the basis of present knowledge, the programming and prepat-terned of the cellular community is little short of incomprehensible. The endocrine control of metamorphosis is, by contrast, a far simpler problem which has begun to clarify itself.

We have seen that the prothoracic gland hormone, ecdyson, is the growth hormone of insects. In its absence, the insect lapses into a state of developmental standstill. When it is again secreted, diapause is terminated and growth is resumed.

When ecdyson is secreted and acts with little or no opposition, it causes the cells to undertake synthetic acts accompanied by the de-repression and utilization of fresh genetic information (Williams, 1961a, b). We have illustrated this action in the case of the giant chromosomes of *Chironomus* (Clever and Karlson, 1960). The larval cells pupate; the pupal cells undergo adult differentiation. So, to revert to an earlier analogy, it is ecdyson that prompts the cells to proceed from one chapter to the next. Therefore, it is necessary to conclude that ecdyson acts directly or indirectly on the nucleus to cause the de-repression of genetic information which can then be implemented by the cytoplasm in new synthetic acts.

At this point I call attention to a minor technicality—the successive instars of the larval life. During the larval period the insect is somehow stabilized as a larva. It grows and molts and grows and molts. The Cecropia silkworm, as we have seen, increases its mass 5,000-fold during the five larval instars. But it fails to metamorphose until the end of the fifth instar.

It is easy to show that ecdyson is the necessary stimulus for larval growth and molting. But somehow there is at work a conservative force which opposes change, which blocks “growing up” without interfering with growth in an unchanging state.

The juvenile hormone

This conservative force proves to be yet another hormone, the so-called "juvenile hormone." As demonstrated by Wigglesworth (1936), the juvenile hormone is secreted by a tiny pair of cephalic glands, the corpora allata.

The action of juvenile hormone is to modify the cellular response to ecdyson—to suppress new synthetic acts without interfering with the use and reuse of the information already at the disposal of the cells (Williams, 1959, 1961a, b). If this "brake" on progressive differentiation is removed by the excision of its source in the corpora allata, then the immature larva reacts to ecdyson by undergoing metamorphosis into a midget pupa and adult (Bounhiol, 1938; Fukuda, 1944; Williams, 1961a, b). What we witness, in short, is the precocious acting out of the life plan.

Under normal conditions and circumstances, pupation becomes possible late in larval life when the corpora allata undergo a progressive loss of their endocrine activity. Ecdyson now acts in the presence of low concentrations of juvenile hormone to cause the de-repression of the fresh genetic information for constructing the pupa (Williams, 1961b). But in many insects one can reapply the "brakes" by implanting active corpora allata obtained from younger larvae. When juvenile hormone is supplied, the mature larva can postpone its metamorphosis and continue to grow as a larva. The net result is that the insect after one or more extra larval instars finally transforms into a giant pupa and adult (Wigglesworth, 1954).

Equally spectacular is the effect of juvenile hormone on the transformation of a pupa into an adult moth, a phase of metamorphosis which normally takes place in the presence of ecdyson and what appears to be the total absence of juvenile hormone. If juvenile hormone is supplied by the implantation of active corpora allata, the pupa is prevented from turning into an adult moth. To varying degrees, as dictated by the concentration of juvenile hormone, the formation of the adult moth is inhibited. In the presence of a sufficiently high titer of juvenile hormone, the pupa responds to ecdyson by molting into a second pupal stage (Williams, 1959).

By the use of the pupal assay, the juvenile hormone has been extracted and extensively purified. It proves to be a water-insoluble oil containing numerous methyl and methylene functions and what appears to be a lactone ring (Williams, 1956a, 1961a). Meanwhile, the German investigator, Schmialek (1961), has been able to mimic the action of juvenile hormone by the injection of the 15-carbon

unsaturated alcohol, farnesol (Wigglesworth, 1961b). The same is true for the 20-carbon unsaturated alcohol, phytol, according to unpublished results of experiments which I have performed in collaboration with John Law. However, in our experience, these well-known chemicals are only about one-thousandth as active as the juvenile hormone itself.

Despite persistent uncertainties as to the chemistry of juvenile hormone, we can state with considerable assurance that the function of the hormone within the living insect is to block the flow of fresh genetic information from nucleus to cytoplasm (Williams, 1961a, b). By analogy with the simpler and more accessible microbial systems, we find no dearth of mechanisms which could account for the conservative action of juvenile hormone. It could affect one or more systems of positive or negative feedback concerned with the repression of specific regulators, operators, or operons; it could interfere with the flow of messenger RNA from nucleus to cytoplasm; it could selectively block the utilization of this information in the synthetic factories.

This much we can say with confidence. Juvenile hormone somehow prevents the cytoplasm from receiving or acting on fresh instructions whose ultimate source must be the coded genetic information of the nucleus. Meanwhile the hormone fails to interfere in any way with the use and reuse of the information already at the disposal of the cytoplasm. In the presence of juvenile hormone a cell can read and reread the same chapter in the construction manual. But it cannot press on to the next chapter.

Insects and microbes

So, in this brief excursion into entomology, we find that the insects obey and in some instances illuminate a number of concepts of developmental biology derived from studies of microbial systems. These concepts, without exception, have to do with the management of genetic information—its de-repression and flow from its source in the hereditary material to its implementation in specific synthetic acts—more particularly, the synthesis of specific enzymes and structural proteins.

In insects as in microbes, it is increasingly clear that cells do not continuously express all the potentialities inherent in their genetic equipment. At any particular cross section of time, most cells are in a highly repressed state.

A bacterial cell synthesizes only those enzymes that are needed and

refrains from synthesizing those which are not needed. The decision not to synthesize a new enzyme is passive in the sense that it involves a continuation of the repressed state of the corresponding gene or genes—a further withholding of the genetic information prerequisite for the synthesis. An affirmative decision to undertake the synthesis of a new enzyme is active in that it involves the de-repression of the corresponding genes and a flow of the appropriate genetic information to the synthetic centers of the cell.

A bacterial cell is equipped, not so much with a “construction manual” as with a “cookbook”—a collection of recipes for the management of any number of energy-rich molecules which it may encounter. Therefore, its genetic equipment is delicately tuned to the exigencies of the environment. In a bacterial cell the countdown on the synthesis of a new enzyme must be counted in milliseconds.

Anyone familiar with the elegant organization of animal cells is entitled to the view that a bacterium is a nondescript object. The cells of a louse are, by comparison, of outstanding beauty. But a far more important distinction is that an animal cell, say, the egg of an insect, has a vastly complicated charge on life. Its method of making another egg is, in the least, a roundabout process.

A central dogma of experimental embryology has been the concept of the totipotent egg—a cell that starts out doing everything and in the course of embryonic development gives rise to a progeny which specializes and differentiates by doing less and less. The microbes have disenchanted us from this point of view. It now appears that the egg is a highly repressed and specialized cell. It is specialized for reproduction, for cloning itself to establish the “critical mass” of the blastoderm (Grobstein, 1959). Even the molecules of yolk seem, for the most part, to have been synthesized elsewhere and “injected” into the ovum prior to fertilization (Bonhag, 1958; Telfer, 1961; Hisaw, 1962).

Unlike the bacterial cell, the egg can scarcely allow itself to be at the disposal of the molecules of the environment. Therefore, it commonly envelops itself with one or more shells and membranes to become a self-contained system. Meanwhile, the genetic plans for the embryo must be speedily programmed.

Here we trip over the phenomenon of “embryonic determination,” which is encountered in its extreme expression in the so-called “mosaic eggs” of the *Diptera* and *Lepidoptera*. The cells of the very early embryo are somehow programmed for a series of future differentiations which they and their progeny will undertake, not only in the embryo, but also during successive phases of metamorphosis. And

in the acting out of these various and sundry programs, the entire community of cells will be paced and coordinated by certain endogenous molecules which, as endocrine agents, will give stop-and-go signals reminiscent of the "controlling elements" in bacterial systems.

It is clear that the programming of cells has to do with sequential gene action—the "taping" of the individual cells for a subsequent "playback." In the concrete case of the higher insects, it is a process that anticipates the future of the individual life and predicts with utmost precision the chemical engineering of larva, pupa, and adult.

References

- Beerman, W. (1952). Chromomerenkonstanz und spezifische Modifikationen der Chromosomenstruktur in der Entwicklung und Organdifferenzierung von *Chironomus tentans*, *Chromosoma*, **5**:139–198.
- (1959). Chromosomal differentiation in insects, in *Developmental Cytology*, ed. by D. Rudnick, The Ronald Press Company, New York, pp. 83–104.
- , and G. F. Bahr (1954). The submicroscopic structure of the Balbiani-ring, *Exp. Cell Research*, **6**:195–201.
- Berger, C. A. (1940). The uniformity of the gene complex in the nuclei of different tissues, *J. Heredity*, **31**:3–4.
- Bonhag, P. F. (1958). Ovarian structure and vitellogenesis in insects, *Ann. Rev. Entomol.*, **3**:137–160.
- Bounhiol, J. J. (1938). Recherches expérimentales sur le déterminisme de la métamorphose chez les Lépidoptères, *Bull. Biol. France et Belg. Suppl.*, **24**:1–199.
- Breuer, M. E., and C. Pavan (1955). Behavior of polytene chromosomes of *Rhynchosciara angelae* at different stages of larval development, *Chromosoma*, **7**:371–385.
- Bridges, P. N. (1942). A new map of the salivary gland 2L-chromosome, *J. Heredity*, **33**:403–408.
- Brink, A. (1958). Mutable loci and development of the organism, *J. Cellular Comp. Physiol.*, **52**, Suppl. 1:169–195.
- Butenandt, A., and P. Karlson (1954). Ueber die Isolierung eines Metamorphose-Hormons der Insekten in kristallisierter Form, *Z. Naturforsch.*, **9b**:389–391.
- Clever, U., and P. Karlson (1960). Induktion von Puff-Veränderungen in den Speicheldrüsenchromosomen von *Chironomus tentans* durch Ecdyson, *Exp. Cell Res.*, **20**:623–626.
- Fisq, A., and C. Pavan (1957). Autoradiography of polytene chromosomes of *Rhynchosciara angelae* at different stages of larval development, *Nature (London)*, **180**:983–984.
- Fukuda, S. (1944). The hormonal mechanism of larval molting and metamorphosis in the silkworm, *J. Fac. Sci., Univ. Tokyo, Sect. IV*, **6**:477–532.
- Gall, J. G. (1958). Chromosomal differentiation, in *McCullum Pratt Symposium on the Chemical Basis of Development*, ed. by W. D. McElroy and Bently Glass, Johns Hopkins Press, Baltimore, pp. 103–135.
- Grobstein, C. (1959). Differentiation of vertebrate cells, in *The Cell*, ed. by J. Brachet and A. E. Mirsky, Academic Press, Inc., New York, vol. 1, pp. 437–496.

- Hisaw, F. L. (1962), The evolution of endocrine adaptations of the ovarian follicle, *Oregon State University Symposium, 1961* (in press).
- Jacob, F., and J. Monod (1961), Genetic regulatory mechanisms in the synthesis of proteins, *J. Mol. Biol.*, **3**:318-356.
- , and E. L. Wollman (1961), *Sexuality and the Genetics of Bacteria*, Academic Press, Inc., New York.
- Karlson, P. (1956), Biochemical studies on insect hormones, *Vitamins and Hormones*, **14**:227-266.
- Kühn, A., and H. Piepho (1938), Die Reactionen der Hypodermis und der Versionschen Drüsen auf das Verpuppungshormone bei *Ephestia kühniello*, *Biol. Zentr.*, **58**:12-51.
- , and ——— (1940), Über die Ausbildung der Schuppen in Hauttransplantaten von Schmetterlingen, *Biol. Zentr.*, **60**:1-22.
- Lucas, F., J. T. B. Shaw, and S. G. Smith (1958), The silk fibroins, *Adv. in Protein Chem.*, **13**:108-243.
- Mechelke, F. (1953), Reversible strukturmodifikationen der Speicheldrüsenchromosomen von *Acricotopus lucidus*, *Chromosoma*, **5**:511-543.
- Pavan, C., and M. E. Breuer (1952), Polytene chromosomes, *J. Heredity*, **43**:150-157.
- Piepho, H. (1938a), Wachstum und totale Metamorphose an Hautimplantaten bei der Wachsmotte *Galleria mellonella* L., *Biol. Zentr.*, **58**:356-366.
- (1938b), Über die Auslösung der Raupenhäutung, Verpuppung und Imagoentwicklung an Hautimplantaten von Schmetterlingen, *Biol. Zentr.*, **58**:481-495.
- (1943), Wirkstoffe in der Metamorphose von Schmetterlingen und anderen Insekten, *Naturwiss.*, **31**:329-335.
- , and H. Meyer (1951), Reaktionen der Schmetterlingshaut auf Häutungshormone, *Biol. Zentr.*, **70**:252-260.
- Ris, H., and H. Crouse (1945), Structure of the salivary gland chromosomes of Diptera, *Proc. Nat. Acad. Sci. U.S.*, **31**:321-327.
- Schmialek, P. (1961), Die Identifizierung zweier im Tenebrio kot und in Hefe vorkommender Substanzen mit Juvenilhormonwirkung, *Z. Naturforsch.*, **16b**:461-464.
- Slizynsky, B. M. (1950), *Chironomus* versus *Drosophila*, *J. Genet.*, **50**:77-79.
- Snodgrass, R. E. (1954), Insect metamorphosis, *Smithsonian Misc. Collections*, **122** (9):1-124.
- Speyer, J. F., P. Lengyel, C. Basilio, and S. Ochoa (1962), Synthetic polypeptides and the amino acid code, H, *Proc. Nat. Acad. Sci. U.S.*, **48**:63-68.
- Stossberg, M. (1938), Die Zellvorgänge bei der Entwicklung der Flügelschuppen von *Ephestia kühniello*, *Z. Morphol. Oecol. Tiere*, **34**:174-206.
- Telfer, W. H. (1961), The route of entry and localization of blood proteins in the oocytes of saturniid moths, *J. Biophys. Biochem. Cytol.*, **9**:747-759.
- Van der Kloot, W. G. (1955), The control of neurosecretion and diapause by physiological changes in the brain of the Cecropia silkworm, *Biol. Bull.*, **109**:276-294.
- Wigglesworth, V. B. (1936), The function of the corpus allatum in the growth and reproduction of *Rhodnius prolixus* (Hemiptera), *Quart. J. Microscop. Sci.*, **79**:91-121.
- (1954), *The Physiology of Insect Metamorphosis*, Monographs in Experimental Biology, no. 1, Cambridge University Press, London and New York, pp. 1-152.

- (1961a), A tentative synthesis, in *Symposium No. 1*, Royal Entomological Society, London, ed. by J. S. Kennedy, pp. 103–113.
- (1961b), Some observations on the juvenile hormone effect of farnesol in *Rhodnius prolixus* Stal (Hemiptera), *J. Insect Physiol.*, **7**:73–78.
- Williams, C. M. (1946), Physiology of insect diapause: The role of the brain in the production and termination of pupal dormancy in the giant silkworm *Platysamia cecropia*, *Biol. Bull.*, **90**:234–243.
- (1952a), Morphogenesis and the metamorphosis of insects, *Harvey Lectures*, **47**:126–155.
- (1952b), Physiology of insect diapause. IV. The brain and prothoracic glands as an endocrine system in the *Cecropia* silkworm, *Biol. Bull.*, **103**:120–138.
- (1956a), The juvenile hormone of insects, *Nature* (London), **178**:212–213.
- (1956b), Physiology of insect diapause. X. An endocrine mechanism for the influence of temperature on the diapausing pupa of the *Cecropia* silkworm, *Biol. Bull.*, **110**:201–218.
- (1958), Hormonal regulation of insect metamorphosis, in *McCollum Pratt Symposium on the Chemical Basis of Development*, ed. by W. D. McElroy and Bently Glass, Johns Hopkins Press, Baltimore, pp. 794–806.
- (1959), The juvenile hormone. I. Endocrine activity of the corpora allata of the adult *Cecropia* silkworm, *Biol. Bull.*, **116**:323–338.
- (1961a), Insect metamorphosis: An approach to the study of growth, in *Growth in Living Systems*, ed. by M. X. Zarrow, Basic Books, New York, pp. 313–320.
- (1961b), The juvenile hormone. II. Its role in the endocrine control of molting, pupation, and adult development in the *Cecropia* silkworm, *Biol. Bull.*, **121**:572–585.
- Yčas, M., and W. S. Vincent (1960), The ribonucleic acid of *Epanaphe moloneyi* Druce, *Exp. Cell Res.*, **21**:513–522.

Growth and Differentiation in the Nervous System*

Rita Levi-Montalcini

**Department of Zoology
Washington University, St. Louis, Missouri**

In the course of divergence of cell lineages and blocking out of different systems from early products of division of the fertilized egg cell, the nervous system is one of the first to assert its differentiation. The thickening of the ectoderm along the middorsal line of the embryo is a barely noticeable event among the many developmental processes which take place at the same time. It is, however, a major event in the history of the nervous system, which has dissociated itself from the

* The work described in this paper was supported in part by grants from the National Institute of Neurological Diseases and Blindness, Public Health Service, Bethesda, Md., and from the National Science Foundation, Washington, D.C., and by a contribution from the American Cancer Society to Washington University, St. Louis, Mo.

The microphotographs were made by Cramer Lewis, Department of Illustration, Washington University Medical School, St. Louis, Mo.

adjacent epithelial tissues and no longer shares their fate. From that moment, two major processes set in, to transform the thin neural plate in that complex structure known as the vertebrate nervous system: The one is the gross shaping of the brain and of the spinal cord which results from intense proliferative activity and from massive cell movements; the other is the elaboration of a multitude of nerve centers and of the wiring which interconnects these centers in the general framework of the central nervous system. Obviously no sharp line can be drawn between these two processes, since they occur simultaneously and the first sets the stage for the second, but they can be singled out and analyzed as two distinct aspects of the same developmental process.

The analysis of the factors involved in the shaping of the brain and the determination of its regional differences has been a favorite object of embryologists ever since Spemann and Mangold discovered the organizing potency of the upper blastoporal lip in amphibians. The molding of the brain and of the spinal cord texture in a myriad of cell aggregates, the formation of fiber tracts, and the outgrowth of peripheral nerves are instead the object of analysis of the neurologist, who explores the developing nerve centers as a clue to the understanding of their ultimate structure and function.

To the biologist who is preoccupied not with the function in the nervous system, but with the more general problems of growth and differentiation, the developing nervous system offers an unequalled field of exploration, for there he finds cell populations differentiating along divergent lines in restricted temporal and spatial dimensions.

To the writer, who shares with the neurologist the interest in structure and function in the nervous system, and with the biologist the curiosity in growth and differentiative processes, the intricacies of the nerve structures represent a never-ending object of wonder and exploration. In the following pages only a few of the manifold aspects of this problem will be considered. The selection is guided by personal experience and interest in some specific developmental aspects rather than by a logical sequential line.

At first we shall consider the developmental mechanisms which operate in the early segregation and differentiation of nerve centers. Then we shall discuss the effects produced on sensory nerve cells by changes inflicted to their peripheral fields of innervation. Finally we shall report on the growth response of sensory and sympathetic nerve cells to some recently discovered nerve growth factors. The similarities and differences between the response of sensory cells to nerve growth factors and to an increased field of innervation will offer an

opportunity of discussing one of the most obscure and yet important aspects of growth and differentiation in the nervous system—the control mechanism exerted by extrinsic factors on these processes.

I. Developmental Mechanisms in the Central Nervous System

The following analysis is based on the study of the developing nervous system of the chick embryo, which has been our favorite object of investigation during the past years. A few observations performed on the nervous system of a mammalian embryo (mouse) show that the developmental processes in birds and mammals follow a similar pattern.

a. First differentiative phase

The band of thickened ectoderm along the middorsal line of the chick embryo has folded into a neural groove and the neural groove has changed in turn into a neural tube, before the cylindrical-shaped cells which form the tube show any sign of their essence. In the head region, differential growth of the walls results in the shaping of the tube in three vesicles. The further elaboration of these vesicles in the main brain regions will not be discussed here. Rather we shall consider the differentiative waves which sweep the neural tube along its rostro-caudal axis and lay down the building blocks of brain and spinal cord nerve centers.

It may be worth mentioning that this first and the following differentiative processes would perhaps have never come to the attention of the neurologist, were it not for the discovery by Cajal, toward the end of the past century, of the unusually high affinity of differentiating nerve cells for silver salts. The silver technique introduced by Cajal offers an invaluable tool to detect the early differentiated nerve cells, which stand out in sharp relief for their dark brown color on the pale background of the undifferentiated cells.

The neural tube has barely formed, when a swarming of neuroblasts takes place from the ventromedial lining of the central canal to the lateral aspect of the tube. In the head region, where this differentiative process first starts, the slender migrating neuroblasts are the forerunner of the diencephalic-mesencephalic main nuclear aggregates and of the motor cephalic nuclei. This process has been described in detail in previous publications (1, 2, 3) and will not be considered here. In the spinal cord, the neuroblasts build up a continuous column in a ventrolateral position. The further differentiation

of this column and the segregation of its different components involve a series of developmental steps which have been analyzed in detail (4) and will be summarized here. In this initial phase of their maturation, the neuroblasts move as isolated units. They cover the short distance between the central canal and the lateral edge of the neural tube by means of amoeboid movements of their bodies, which glide between the parallel and tightly packed rows of undifferentiated cells. As soon as they reach the ventrolateral area of the tube, they assemble together in a column which increases in size as new units join the ones which have already settled. This process continues till the fourth day. A similar migratory movement takes place also from other more dorsal sectors of the ependyma around the central canal. The neuroblasts originating from these segments do not gather in a column but take position as isolated units or small cell groupments along the dorsolateral edge of the neural tube. They give origin to the associative and commissural neurons, which establish links between the two halves of the spinal cord and between rostral and caudal segments of the tube. At the end of the fourth day of incubation, this early differentiative phase is practically completed in the brain centers and in the rostral segments of the spinal cord, while it is still in progress but near completion in the thoracic and lumbosacral segments which mature later, according to the rostrocaudal gradient of differentiation. The subsequent phase, between the end of the fourth and the eleventh day of incubation, represents the most dynamic period in the developmental history of the spinal cord and brain centers of the chick embryo. It is in this period that the apparent homogeneous population of early differentiated nerve cells breaks down at some levels and is swept away in a few hours, while at other levels it splits in morphologically and functionally distinct cell agglomerates; some of the cells continue their differentiation *in situ* while others migrate in compact columns or cell rows to settle in regions at considerable distance from the ones where they first underwent maturation. Numerous instances of such migratory processes were described in different segments of the spinal cord, brain stem, and higher brain centers (4, 5, 6). Here we shall outline only the main developmental patterns of differentiation of nerve centers and present some considerations on the mechanism of these differentiative processes.

b. Second differentiative phase

At the end of the fourth day of incubation, the spinal cord in the chick embryo exhibits a deceptively simple architecture. A compact

column of neuroblasts extends from the cervical to the sacral segment. It is easily identified as the motor column from its position as well as from the direction of the axons which gather in small nerve bundles and emerge from the neural tube as motor roots. The associative and commissural neurons mentioned above give origin to slender fiber tracts; the commissural fiber tract marks the borderline between the compact core of undifferentiated cells which still represent the main component of the neural tube and the differentiated neurons collected in the lateral column. A detailed description of this process is presented in a previous publication (4); here we shall briefly outline the developmental patterns of the motor system at the cervical, brachial, and thoracic levels.

Regional Patterns

Cervical Segment. The cervical lateral motor column undergoes a dramatic and massive degeneration between $4\frac{1}{2}$ and 5 days of incubation; in these few hours about three-fourths of the neuroblasts which form the column disintegrate and are swept away by macrophages which gather in the area (4). The combined use of three different techniques—silver impregnation, vital stains, and standard histological techniques—gave a complete picture of this striking process. At 5 days the column is reduced to about one-fourth of its previous width. The surviving cells form the slender medial column which innervates the trunk muscles at that level (Fig. 1c).

The significance of this extensive and short-lasting disintegrative process in the cervical segment of the spinal cord is obscure. We are not aware of the occurrence of similar mass destruction taking place under normal conditions in other sections of the developing nervous system, although we described two other instances of cell disintegration in the cervical and thoracic spinal ganglia of the chick embryos in stages between 5 and 6 days (7). Considerations presented in the previous publication (4) suggested the possibility that the disintegrating cells at the cervical level might represent an abortive preganglionic center. The experiments performed to test this hypothesis will be presented after discussion of the differentiative pattern in other spinal cord segments.

Brachial and Lumbosacral Segments. In both segments the spinal motor column segregates into a large lateral component which innervates, respectively, the anterior and posterior limbs, and into a medio-lateral column of the same width and position as the column described at the cervical level. This column is also present at the thoracic and

FIG. 1. Diagrammatic illustration of the emergence of regional differences from a morphologically uniform system in the spinal motor column of the chick embryo. (a) 3-day embryo: The motor column is of uniform width from the cervical to the sacral level. (b) 5-day embryo: The majority of the differentiating neuroblasts in the cervical segment of the motor column undergo degeneration. They are represented as solid black circles. In the thoracic and sacral segments the migration of the preganglionic columns is under way. (c) 8-day embryo: The degenerated nerve cells in the cervical segment of the motor column have disappeared. The remaining nerve cells in this segment form the slender medial motor columns. Note the size increase of the brachial and lumbar motor columns innervating the limbs. In the thoracic and sacral segments, the two preganglionic columns have reached their definitive position adjacent to the central canal. The two slender columns in a peripheral position represent the medial motor columns. C, cervical level; B, brachial level; L, lumbar level; PG.S, preganglionic sacral center; PG.TH, preganglionic thoracic center; P.V.SY, paravertebral sympathetic ganglia; S, sacral level; S.C, superior cervical ganglion; TH, thoracic level.

sacral level. In all segments it serves the same function—innervation of the trunk musculature (Fig. 1).

Thoracic and Sacral Segments. In both segments two massive migratory movements take place between the middle of the fourth and the end of the seventh day of incubation. The following description is based on the analysis of this process at the thoracic level. Cell movements in the sacral segment present similar features and therefore will not be discussed.

Before describing the migratory movement, it is of interest to consider the motor column, which has barely formed and which consists at 4 days of neuroblasts of similar size and shape in a random distribution. The duality of this apparently homogeneous population becomes manifest if one inspects the peripheral distribution of their nerve fibers. The majority of these fibers end in contact with the primordia of the sympathetic ganglia at the two sides of the spinal cord, whereas only a small fiber contingent is directed toward the trunk muscles. It is therefore evident that the motor column consists of a mixed population of preganglionic and somatomotor nerve cells. In later developmental stages the two populations are topographically well distinct from each other: The preganglionic column is located in birds in a dorsomedial position adjacent to the central canal while the somatomotor column has a medioventral position and is identified as the medial motor column.

The segregation of the two cell populations is foreshadowed by a change in the texture of the newly formed motor column. At $4\frac{1}{2}$ days about three-fourths of the entire cell population assembles in compact and parallel rows of neuroblasts oriented toward the central canal (Figs. 1 and 7). A migration of these cells starts immediately after, while the remaining cells continue their differentiation *in situ*. They form the medial motor column.

The progression of the migratory cells during the three following days is diagrammatically represented in Fig. 1 and needs no further comment. The intense silver affinity of the migrating cells and the slowness of the entire process, which was studied in close time series of embryos between the fourth and the end of the seventh day, gave the possibility of exploring all the aspects of this movement, which results in the active dislocation of thousands of differentiated nerve cells from their early ventrolateral position to their terminal settlement dorsal to the central canal (Figs. 1, 8, and 9).

At the end of the seventh day the migratory movement is completed, and one can easily identify in the cells which gather in two

distinct nuclei adjacent to the central canal the cells of origin of the preganglionic motor centers (Figs. 1, 10, and 11).

An Experimental Analysis of the Regional Patterns in the Cervical and Thoracic Spinal Cord Segments

The hypothesis (see page 265) that the disintegrating nerve cells in the cervical segment of the spinal cord might represent an abortive visceral column was tested by transplanting the cervical segment of the spinal cord from 2-day chick embryo to the thoracic level of hosts of the same age (8). The thoracic segment of the host was extirpated and the transplant inserted in its place. In performing this operation, Shieh wanted to test the hypothesis mentioned above, by providing the "abortive cells" with the possibility of establishing connections with the sympathetic ganglia. The operation presents technical difficulties and only a few embryos survived till the seventh day. In the few specimens examined, Shieh observed the formation of a preganglionic motor column similar to the one which forms at the thoracic level. He also observed a large number of degenerating neurons. These results suggest that at least a number of the cells that normally undergo degeneration at the cervical level survive in the transplant and build a preganglionic motor column with the same mechanism as normally occurs in the thoracic level. These experiments, if confirmed by others now in progress, will support the hypothesis that an abortive preganglionic center is present in the cervical level of the spinal cord. They would also give evidence of the flexibility of differentiative patterns in the central nervous system.

Mass Migration in Other Sectors of the Central Nervous System

Between the fifth and the eleventh day, the brain stem and the suprasegmental brain structures are the stage of an incessant and yet highly organized movement of differentiated nerve cell populations. We shall consider below the main features of these movements, as they appear in the developing nervous system of the chick embryo.

Migratory Movements in the Brain Stem. The early continuous motor column (see page 265) segregates at first in the primordia of the XII, XI, X, VII, V, IV, and III nuclei. Each of these centers segregates in turn in the following days into two or more cell groupments through more or less complex migratory movements of its components.

At some levels, movements in opposite directions take place at the same time, or two cell populations, moving along slightly divergent

paths, merge and segregate again, each to pursue its specific course. In all instances the migrating cells are well differentiated at the time they start their journey. Their slender, elongated bodies, dark-stained in silver preparations, detach sharply on the pale background of the undifferentiated cells, which still represent the major component of the brain stem at this stage. Each population is easily recognizable from adjacent populations by the different shade of its color in silver-impregnated material. All migrating cells trail their axon behind while moving toward their final location; the direction of the fibers represents in the mature organism a valuable guide to detect the route followed by a nerve center to reach its final destination.

It was the peculiar path traced in the brain stem by the axons of the accessory nucleus of the VIth nerve (Fig. 2) which suggested that its cells of origin have undergone a migratory movement in previous developmental stages. This movement was actually observed in embryos examined between the sixth and the eighth day of incubation.

During this period a considerable number of cells segregate from the main nucleus located near the midline on the floor of the fourth ventricle; they move in a straight line, one after the other in a single row in an oblique direction from their former mediodorsal position to their final ventrolateral location. The neuroblasts cut their way across the dense texture of the brain stem, which consists of closely packed rows of undifferentiated cells with their long axes oriented at 45° with the axes of the migrating cells (Figs. 2b, 2b', 12). Toward the middle of their journey the shifting population crosses and intermingles with the population of the facial nucleus, which is also moving from a laterodorsal to a ventrodorsal position (Fig. 12). The two populations dissociate at the beginning of the seventh day and each reaches its destination about 24 hours later (Figs. 2, 13).

Migratory Movement in the Third Nucleus. This movement, which was first described by Biondi (9), consists of the exchange across the midline of one of the four oculomotor nuclei with its partner of the opposite center. The exchange occurs between the two ventromedial nuclei; it starts at the end of the fifth day and is completed at 8 days. Since the two populations moving across the midline are of a considerable size, the traffic of the neuroblasts moving in the opposite direction and crossing each other is heavy. At 6 days a continuous trail of neuroblasts in different rows bridge the two oculomotor centers (Figs. 3b', 14). When their final settlement is completed at 7 days, the axons trailing behind make a ventral commissure on the midline (Figs. 3c', 15).

Migratory Movements in Suprasegmental Structures. In all supra-

MIGRATION IN THE ABDUCENS

FIG. 2. Diagrammatic frontal and transversal representation of the segregation of the accessory nucleus from the main nucleus of the VIth nerve in the chick embryo. *a*, *b*, *c*: stereographic views of the VIth nuclei at 5, 6, and 11 days. In *a*, the nuclei appear as two columns; in *b*, the arrows indicate the migration of neuroblasts from the main nuclei in a lateral direction; in *c*, the migration is completed and the two accessory columns are formed. *a'*, *b'*, *c'*: the same process as viewed in transversal sections. VI A, main nucleus; VI B, accessory nucleus.

MIGRATION IN THE OCULOMOTOR

FIG. 3. Diagrammatic frontal and transverse representation of the segregation of the four nuclei of the IIIrd nerve and of the migration of the two ventromedial nuclei across the midline. Solid black, left oculomotor column; white, right oculomotor column. *a, b, c*: stereographic views in a frontal plane. *a', b', c'*: the same process as viewed in transverse sections.

segmental structures, migration of differentiated nerve cells plays an important role in the molding of nerve centers. These movements are more difficult to analyze than the ones in the brain stem since the active displacement of cell populations occurs at the same time as gross morphological changes in the shape of the brain. In the following discussion we shall outline only two of these movements, which occur in the cerebellar cortex and in the avian telencephalon.

Migratory Movements in the Cerebellum. The complex changes in size and shape in the Purkinje cells during their late differentiation were first described by Cajal (10) and nothing can be added to his exact analysis of the process. Until recently it had not been settled whence these cells originate and how they appear during the early developmental phases of their differentiation [unpublished observations (11)]. We traced these cells in our series of avian brains in stages between 7 and 8 days of incubation. They appear as elongated neuroblasts in proximity to the deep subependymal germinal layer. Since they react intensely to silver and exhibit the characteristic fusiform shape of cells in a migratory phase, it was easy to explore their position during the following days and trace their active displacement from the deep to the cortical cerebellar layer. During the early phases they are closely packed in dense cellular rows; as they reach the cortical layer they spread in a fan-like fashion and settle in the layer which will be known as the Purkinje layer in the fully developed organism (Figs. 4, 16 to 19).

The origin of the Purkinje cells from the germinal layer was recently described by Uzman (12), who utilized the technique of labeling the migrating cells with tritiated thymidine. His observations establish beyond doubt the migratory movements of these cells. This technique can be very useful to detect the migratory movements in nerve cell populations. In future research both this technique and the silver method, which has the advantage of staining selectively the migrating cells, should be combined.

Migratory Movements in the Cerebral Hemispheres. The mass migration of one of the avian telencephalic centers was described in detail in a previous publication (6). Since it does not substantially differ from other migratory movements of differentiated nerve cells, we shall mention here only its main features.

The mass migration of the nucleus, which we designated as nucleus epibasalis centralis, starts at 8 days. It consists of two dense neuroblast columns moving along the lamina medullaris dorsalis. The cells exhibit the characteristic elongated shape of migrating neuroblasts and the intense silver affinity. One can trace the two moving columns and

MIGRATION IN THE CEREBELLUM

FIG. 4. Diagrammatic representation of the migratory movements of the Purkinje cells in the cerebellum of the chick embryo.

their progression in closely timed series till they have reached their final location in the central epibasal area at 12 days. Then the stationary cells lose their parallel alignment and become multipolar (Fig. 5).

In seeking for an explanation for these migratory movements, which resemble the organized and directed migration of a school of fish in a stream or of a colony of termites, we are faced with several questions: Why should differentiated cells engage in such long routes and settle down in areas so far removed from the ones where they first underwent their early differentiation? Which is the major motive force underlying and directing the migrating cell populations?

Similar questions were raised in connection with other cell movements such as the migratory movements of cells in developmental processes, the disaggregation and reaggregation of cell groupments intermingled at random in tissue culture, and the synergistic movements which play such a prominent role in the developmental cycle of slime molds. We shall consider briefly the results of these investigations since the explanations offered in all three instances may have a bearing on our own problem.

Migratory patterns of chromatophores in Amphibia were investigated by Twitty and by Twitty and Niu (13, 14) *in vivo* and *in vitro*, with refined and precise techniques. The results of the experiments performed *in vitro* suggest that the cells move in response to mutual stimulation or repulsion mediated through the action of substances released by the cells.

Reaggregation patterns of suspensions of freshly dispersed embryonic cells in liquid culture medium were the object of detailed analysis by a number of investigators (15, 16, 17, 18, 19). The early hypothesis by Moscona and Moscona (15), that cell exudates might have an orienting influence on the migration and aggregation of dissociated cells, found support in more recent investigations by Moscona (20). He observed that cells dispersed in a liquid medium frequently "appear to follow each other in clearly discernible rows." Phase-contrast, time-lapse movies of similar cultures reveal that the cells move within fine strands of highly transparent slimy substance, evidently an exudate of cellular origin. The author suggests that the extracellular matrix might act as a cell-integrating system endowed with specific cell-directing activities.

The aggregative and migratory patterns of slime molds (21, 22, 23) present a striking instance of cell movements under the controlling action of diffusible substances. The mechanism of the movement and the chemical nature of the agent acrasin, produced by some of the cells, lent themselves to a more precise exploration than animal cells.

Migration in Telencephalon

FIG. 5. Diagrammatic illustration of the migration of the epibasal central nuclei in the avian telencephalon. In *a*, the black and white arrows indicate the direction of movement of the two cell populations from their early formation at 7 days to the completion of the process at 12 days. In *b*, the two migrating populations are represented. ED, nucleus epibasalis dorsalis; ECM, nucleus epibasalis centralis, pars medialis; ECP, nucleus epibasalis centralis, pars posterior.

PLATE I. Figures 6 to 11 illustrate the formation and migration of the preganglionic thoracic column in the spinal cord of the chick embryo between 3 and 11 days. All sections are from material impregnated with the Cajal-De Castro silver technique. The microphotos in this and in the following plates are not retouched. FIG. 6. 3-day embryo. FIG. 7. 4-day embryo. FIG. 8. 5½-day embryo. FIG. 9. 6-day embryo. FIG. 10. 7-day embryo. FIG. 11. 11-day embryo. See explanation in text.

PLATE II. Fig. 12. Transverse section through the VIth nerve of a 6-day embryo. Arrow points to neuroblasts in the process of migrating from the main nucleus (*a*) of the VIth nerve to the accessory nucleus (*b*) of the same nerve. *c*, neuroblasts of the VIIth nucleus. FIG. 13. Transverse section at the same level as Fig. 12, in an 11-day chick embryo. The migration of the accessory nucleus (*b*) from the main nucleus (*a*) of the VIth nerve is completed. *c*, VIIth nucleus. Fig. 14. Transverse section through the nuclei of the IIIrd nerve at the mesencephalic level of a 6-day embryo. The arrow points to the two populations (*a* and *a'*) as they cross the midline and move in opposite directions to join the nuclei of the contralateral IIIrd nerve. FIG. 15. 15-day embryo. Cross section at the same level as in Fig. 14. The two ventromedial nuclei of the IIIrd nerve (*a* and *a'*) are seen in their definitive contralateral position. Note the segregation of the oculomotor nerve cells in four nuclei.

These last investigations gave definite support to the concept that diffusible agents control cell movements. Since these movements (as in the case of migrating nerve cells) occur simultaneously with other differentiative processes, it is difficult to draw a line between the effects exerted by these agents on cell movements and on other aspects of cell differentiation.

In the case of the migratory movements in the central nervous system, we are under no illusion that the processes might lend themselves to a precise experimental analysis. Even though the hypothesis that the complex locomotion pattern of nerve cells might be directed by some agent released in the matrix by other cells or by the matrix itself is plausible, the demonstration would be extremely more difficult than in other situations where cell movements occur in more suitable environmental conditions. This conclusion should not discourage further attempts to explore the complex migratory patterns of nerve cells, since this exploration will undoubtedly contribute to our knowledge of the developing nervous system. The results of the experimental analysis of the spinal cord by Shieh (see page 268) and results of experiments now in progress on the brain stem show that the migratory patterns can be altered by transposition or ablation of segments of the neural tube. They give evidence of the flexibility of these processes and indicate that environmental factors play a role in the differentiation of the central nervous system.

We shall devote the following section to the analysis of developmental and growth processes of nerve cells which early in life migrate out of the central nervous system and establish themselves in discrete cell aggregates known as sensory and sympathetic ganglia. This position is most favorable for exploring the cells under normal and experimental conditions. The results to be reported below show that these cells are highly receptive to agents present in the medium not only during embryonic life but throughout all their life cycle.

II. Role of Peripheral Effectors and Receptors in the Differentiation and Growth of Nerve Centers

When in 1935 R. G. Harrison presented in the Croonian Lecture the results of his brilliant analysis of the developing nervous system of the amphibian (24), the newly opened field of experimental neuroembryology seemed to offer unlimited possibility to explore the nervous system and to uncover the mechanisms which operate in its differentiation.

Three aspects of this analysis impressed particularly the reader of the Croonian Lecture and encouraged further experimentation along this line: (1) the extraordinary tolerance of the embryo for any kind of surgical performance, ranging from restricted ablations to destruction of most of its nervous structures; (2) the flexibility of the developing nervous system, which readily adapts itself to any new situation and even accepts foreign tissues as peripheral receptors and effectors; and (3) the rapidity of the response of developing nerve centers to changes inflicted to their fields of innervation. The situation in the early thirties seemed to be very similar to the situation which prevailed a few years earlier in the field of experimental embryology, when the discovery of embryonic induction had transformed the static field of descriptive embryology into one of the most dynamic and promising areas of biology. In the years which followed the publication of the Croonian Lecture, all possible avenues of this new field were explored. First the amphibian larvae and then the chick embryos submitted obediently to the glass or steel needle, which performed all kinds of ablation and transplantation experiments. Many times during the past years the results and perspective in this field have been considered (25, 26, 27). We shall examine here only one of the most discussed problems of neuroembryology—the mechanism of control exerted by peripheral effectors and receptors on their associated nerve centers. Since the first experiments were performed by Harrison, Detwiler, and their students (24, 28) a good deal has been learned about the effects elicited by decreasing or increasing the peripheral field of innervation, while practically nothing has been learned about the regulatory mechanisms which operate under normal and experimental conditions. Yet it is the understanding of these mechanisms which is important if we wish to gain information on growth and developmental processes in the nervous system.

In the past the attention has been mainly focused on the differentiation of spinal ganglia. As a result we know today more about the differentiative pattern of sensory nerve cells than about any other nerve cell. Most of the investigation centered on the chick embryo, which provides excellent material for normal and experimental analysis. We shall consider in the following section the results of this analysis, which recently received new impulse from the discovery of specific nerve growth factors well identified in their chemical and biological aspects. A closely related cell, the sympathetic nerve cell, will also be considered since its response to growth factors, which is even more impressive than the response of the sensory cell, sheds light on other facets of the problem under consideration.

PLATE III. FIGS. 16 to 19 show the mass migration and progressive differentiation of the Purkinje cells in the cerebellum of embryos at 8, 11, 14, and 19 days of incubation. See explanation in the text.

***Developmental pattern of sensory ganglia
under normal and experimental conditions***

In 1934 and in 1939 Hamburger performed the first detailed analysis of the growth and differentiation of sensory ganglia in chick embryos under normal conditions and under the experimental conditions of unloading or overloading their peripheral field of innervation (29, 30). The extirpation of the limb bud at the stage of 3 days of incuba-

tion or the implantation of an additional limb at the same stage provided the conditions for this investigation. He observed severe atrophy of the ganglia deprived of their peripheral field and noticeable size increase in the ganglia confronted with a larger than normal field of innervation. His results were confirmed in subsequent investigations (31, 32). In 1943, while performing this analysis, we observed two distinct classes of cells in the sensory ganglia (31). The presence in these ganglia of two populations differing from each other in cell size as well as in their location in the ganglia was first described by Hamburger in his early investigation of normal embryos and embryos deprived of one limb bud (29). We now observed that the two cell populations differ strikingly from each other also in their affinity for silver. The lateroventral population is the first to differentiate and is formed of large nerve cells which react intensively to silver from their first differentiation at 4 to 5 days to the end of the incubation period. The mediodorsal population does not show silver affinity till the end of the incubation period. The two cell groups, which we designated in a previous publication (32) as V-L and M-D from their position, differ therefore from each other in time pattern of differentiation, cell size, position, and silver affinity (Fig. 22). Recently a cytochemical difference between these two types of neurons was shown by Gerebtzoff, who investigated the acetyl-cholinesterase activity in different cells and fiber systems (33). Although the author called attention only to the different content of the enzyme in the fibers of the dorsal funiculi, his photographic documentation shows a clear-cut difference in the amount of the enzyme present in the two cell populations: The V-L but not the M-D cells show the presence of the enzyme (Fig. 23). The structural and chemical differences between the two types of sensory nerve cells come in sharp relief when the ganglia are confronted with changes in their field of innervation. In 1944 (31), we observed that the extirpation of the limb bud at 3 days of incubation results in an almost immediate breakdown and disappearance of most of the V-L cells, whereas the M-D cells persist in atrophic condition till the end of the incubation period. In 1949 (32), we described a differential response of these two cell populations following implantation of an additional limb bud in 3-day-old embryos. Counts of the V-L neurons in ganglia confronted with an enlarged field showed an increase of about 80 per cent of these neurons over the control cell population, whereas no changes were detected in the number of the M-D cells. Results to be presented in the following section gave evidence of a differential and opposite response of the two types of neurons when confronted with neoplastic tissue.

PLATE IV. Figures 20 and 21 represent transverse sections through the thoracic level of 10-day embryos. FIG. 20. Control. FIG. 21. Embryo injected for 3 days with the purified salivary NGF. In both microphotos, arrows point to spinal

A comparison of the effects called forth by extirpation of the limb primordia or by implantation of additional limb buds showed that in both instances all developmental processes of sensory nerve cells seemed to be affected—the mitotic activity, the initial differentiation of sensory nerve cells, as well as their further growth (32).

Two hypotheses were advanced as to the mechanism of this control. Either (1) the peripheral field of innervation controls the growth processes of the associated nerve centers by releasing into the circulation some agent which selectively promotes the growth and differentiation of nerve cells tributary to that field, or (2) the periphery affects only those nerve cells which establish contact through their nerve fibers with the field of innervation. Both alternatives failed to explain satisfactorily all the effects evoked in the sensory ganglia by extirpation or transplantation of an additional limb. The hypothesis listed at (1) was in conflict with the observation that an increase or a decrease of the peripheral field of innervation affects only the sensory and motor centers which supply nerves to that area. The other alternative, which considers the nerves as mediators between the end organs and the associated nerve centers, agrees better with these observations. It does not, however, explain other aspects of the growth response of the sensory ganglia to changes inflicted in their peripheral field of innervation. It was in fact found that the mitotic activity in these ganglia increases or decreases according to the extension of the area they innervate. Since dividing cells are obviously deprived of nerve fibers, these results cannot be accounted for as a direct effect of the periphery on individual nerve cells. The experiments to be reported in the following section suggest a more satisfactory explanation of the above results. They will therefore be reconsidered after presentation of the new findings.

ganglia. Note the increase in size of ganglia of the treated embryo. FIG. 22. Spinal ganglion of a control 11-day embryo. The two cell populations in the spinal ganglion (*MD* and *VL*) show the difference in their affinity for silver. FIG. 23. The acetylcholinesterase activity in the *V-L* nerve cells of the spinal ganglia in 10- to 11-day chick embryos. (*Reproduced with the permission of M. A. Gerebtzoff from Fig. 32 in Cholinesterases, Pergamon Press.*) FIG. 24. Spinal ganglion contributing nerve fibers to a fragment of mouse sarcoma 180 (not apparent in the picture). Same magnification as Fig. 22. Note the size increase of the *MD* population. The *VL* population does not show in this section. FIG. 25. The massive invasion of an implanted fragment of mouse sarcoma 180 by nerve fibers from adjacent sensory and sympathetic ganglia. *T*, tumor; *S*, sensory ganglion; *SY*, sympathetic ganglion.

III. Growth Response of Nerve Cells to a Diffusible Protein Agent

The finding in 1948 by Bueker of a nerve growth stimulating effect of mouse sarcoma 180 on the sensory ganglia of chick embryos (34) may be defined as the outcome of a fortuitous discovery and a calculated search. The results so much exceeded the expectation that the unprepared mind of the observer overlooked the exceptional magnitude of the response in the attempt to make it fit into the previously accepted schemes of growth and differentiation in the nervous system.

In transplanting a fragment of mouse sarcoma 180 in the body wall of 3-day chick embryos, Bueker wanted to test the effect of a rapidly expanding tissue on sensory and motor nerve cells. When he examined the embryos 5 days later he found in a number of cases that the tumor had become established and was invaded by sensory but not by motor nerve fibers. He also observed that the sensory ganglia contributing fibers to the tumor were enlarged in size as compared to the contralateral normal ganglia. No changes were observed in the motor column. The author concluded that the tumor elicited the growth response in the sensory ganglia by providing them with a larger field of innervation than their usual field. Since the effect appeared to be restricted to the ganglia sending nerve fibers to the tumor, the nerve fibers were considered as the mediators of the effect. The results and the conclusions therefore fitted well into the picture presented above of the effect of the periphery on associated nerve centers. A reinvestigation of this effect led to a different interpretation of the phenomenon.

a. Response of sensory embryonic nerve cells to implantation of mouse sarcomas

The response of the sensory ganglia to implantation of mouse sarcoma 180 or 37 was investigated in a close series of chick embryos ranging in age between 4 and 18 days (35). Ganglia adjacent to the transplant provide its neurotization with large nerve bundles, which branch in all directions inside the tumor (Fig. 25). At 11 days the ganglia appear three times larger than controls. The overgrowth is due to an increase in cell size and cell number of the M-D population, whereas no changes are apparent in the V-L population, which does not seem to participate in the innervation of the tumor (Fig. 24). From the eleventh day to the end of the incubation period, the effect

of the tumor on M-D cells gradually decreases in spite of the steady increase in size of the transplant, which at 18 days normally fills the abdominal cavity of the embryo (36, 37).

A series of investigations on the effects of mouse sarcomas 180 and 37 on sensory ganglia of the chick embryo, explanted in vitro with the hanging drop technique, showed that the neoplastic cells elicit a remarkable growth effect from the sensory cells (Figs. 32, 33). The effect consists of the outgrowth of a dense halo of nerve fibers from the ganglia adjacent to but not in contact with the tumor within the first 10 hours of culture in vitro (38). The effect is maximal if the ganglia are explanted from 7- to 9-day chick embryos. It decreases in ganglia explanted from older specimens and is no longer apparent in ganglia from 18-day chick embryo. The effects in vitro show therefore a parallelism with the effects elicited by the tumor in the embryo: In both instances the sensory cells appear to be receptive to the growth agent present in the neoplastic tissue only during a limited period of their growth.

The results of the in vitro experiments showed that the tumor evokes a response from the sensory ganglia even if no contact is established between the two tissues. Experiments to be reported below show that a growth response can be elicited from the sensory ganglia in vivo by simply injecting diffusible agents into the yolk of developing embryos.

It was concluded from the above findings that a direct contact between nerve fibers and the tissue producing the growth agent is not necessary to elicit the increase in size and number of the M-D population in the sensory ganglia.

b. Response of sensory ganglia of chick embryos to diffusible nerve growth factors

It was the substitution of the tissue culture technique for the laborious and time-consuming technique of transplantations in the embryo which made possible an extensive search for nerve growth factors and also offered the possibility of investigating the biochemical aspect of the problem. Two potent nerve growth factors (NGF) discovered in snake venom and mouse submaxillary salivary glands (39, 40, 41, 42) show striking biochemical similarities with the factor isolated from the tumor. In all three instances the agent isolated and purified by Cohen was identified with a protein or a protein-bound particle (41). Here we shall consider the biological effects of the purified extract of

PLATE V. FIG. 26. Whole amounts of the sympathetic thoracic chain ganglia of experimental (*E*) and control (*C*) mice 19 days old. Experimental mouse injected with the NGF salivary factor from birth. *ST*, stellate ganglia. FIG. 27. Transverse section of stellate ganglia (*ST*) in experimental (*E*) and control (*C*) ganglia of Fig. 26. Sections through levels indicated by upper arrows in both chains of Fig. 26. FIG. 28. Transverse section of superior cervical ganglia of two adult mice. *E*, mouse

the mouse salivary glands on the sensory ganglia of chick embryos. Similar results were obtained by injecting the purified agent extracted from snake venom.

The addition to the culture medium of the purified protein extracted from the salivary glands or from the snake venom elicits the same nerve growth response as fragments of tumor or tumor extract. Both the snake venom and the mouse salivary glands harbor this agent in such a large amount that a very small quantity of the purified extracts is sufficient to evoke the *in vitro* effects. It was shown by Cohen that the purified salivary extract is active at a concentration of 0.01 gamma of protein per milliliter (41). This solution, when added to the medium of culture in the proportion of one part to one part of synthetic medium and one part of plasma, elicits a dense halo of nerve fibers from the explanted sensory ganglia. It was designated as one biological unit. A comparison between the potency of the tumor and the salivary gland extract indicates that the latter is about 6,000 times more potent on a dry weight basis than the tumor.

Daily injections of a few gamma of the purified protein isolated from mouse submaxillary salivary gland into the yolk of 6- to 10-day chick embryos call forth a growth response from the M-D population of sensory nerve cells, comparable to the response elicited by intra-embryonic transplants of mouse sarcomas (Figs. 20, 21). The size increase of this population has its counterpart in the hyperneurotization of the exteroceptive embryonic fields. Since no increase in the V-L population was observed and no hyperneurotization of the muscles is apparent, it is tempting to correlate these two populations with different fields of distribution. Further investigations are in progress to elucidate this point and to trace the distribution of nerves from the hypertrophic ganglia to their terminal structures.

injected for 1 week with the NGF salivary factor; C, control ganglion. FIG. 29. Transverse sections through superior cervical ganglia of 4-month-old mice. C, control; E, mouse injected for 5 days after birth with the antiserum to the salivary NGE. FIGS. 30 and 31. Sympathetic nerve trunks of control and experimental chain ganglia of Fig. 26 at higher magnification. Low arrows in Fig. 26 indicate areas enlarged in Figs. 30 and 31. FIG. 32. Spinal ganglion of a 7-day-old chick embryo cultured *in vitro* for 24 hours. FIG. 33. Spinal ganglion as in Fig. 32, combined *in vitro* with a fragment of mouse sarcoma 180. Note the dense fibrillar halo. FIG. 34. Sympathetic ganglion of a human fetus 3 months old cultured for 24 hours in a medium to which the NGF salivary factor was added.

***c. Response of sympathetic nerve cells
of chick embryos to implantation
of mouse sarcomas and to the injection
of other nerve growth factors***

Since the beginning of this investigation in 1950, we realized that the sympathetic nerve cells are far more affected by the tumoral factor than the sensory nerve cells. While the size increase of the sensory ganglia adjacent to the transplant ranges between two and three times that of the controls, the size increase of sympathetic ganglia is four to six times that of the controls (42). Two factors may in part account for these differences: (1) The sympathetic nerve cells at variance with the sensory nerve cells are all receptive to the nerve growth factor; and (2) the effect is not restricted to a given period of their differentiation but continues till the end of the incubation time. As will be shown below, sympathetic nerve cells of mammals remain receptive to the growth factor throughout all life (Fig. 28).

In a number of cases the effects of the tumor are restricted to the sympathetic ganglia which establish contact with the neoplastic tissue through their nerve fibers (35). In the majority of the cases, however (about 85 per cent of the total number of embryos examined, which amounts to many hundreds), the entire sympathetic chain ganglia are strikingly enlarged (36). Nerve fibers emerging from the hypertrophic and hyperplastic ganglia overflow the adjacent viscera, which are not innervated in corresponding developmental stages in control embryos; they also enter and fill large blood vessels with thick nerve bundles. Fragments of tumor transplanted onto the chorioallantoic membrane of 4- to 6-day chick embryos elicit the same effects. In this last group of experiments, the tumor and the host share the circulation, but no direct contact is established between them.

Sympathetic ganglia from 9- to 11-day chick embryos, explanted in vitro in proximity to fragments of mouse sarcomas, produce a dense halo of nerve fibers similar to the halo produced by sensory ganglia (see page 285). The in vitro technique was also used to test the effects of the purified extract of mouse salivary glands and of snake venom on the sympathetic ganglia of chick embryos. The injection of the purified extract of the salivary gland into the yolk of 6- to 10-day chick embryos evokes an overgrowth of the entire sympathetic chain ganglia and hyperneurotization of the viscera (43).

In summary, we may conclude that the M-D sensory neurons and the sympathetic nerve cells of the chick embryos are highly receptive

to a growth factor present in mouse sarcomas, snake venom, and mouse submaxillary salivary glands.

Some of the results are of particular interest for the light they shed on the growth and differentiative processes of embryonic nerve cells. They show that the growth potentialities of these cells far exceed their growth range under normal conditions. Furthermore, they show that the active factor can reach the cells through the nerve fibers or through the circulatory system. The penetration of sympathetic nerve fibers into viscera and blood vessels normally impermeable to nerves, as well as the hyperneurotization of exteroceptive fields with sensory nerve fibers, shows that both the qualitative as well as the quantitative aspects of peripheral nerve distribution are not rigidly fixed in the embryo but that considerable deflections from the normal pattern are tolerated.

d. Response of sympathetic nerve cells of newborn and adult mice to the purified salivary NGF

After the experiments in tissue culture indicated that sympathetic ganglia of mammals (rodents and human fetuses) are receptive to the NGF, we tested its effects in newborn and then in adult mice. Daily injections of this agent in the amount of 2,000 biological units per gram of body weight in mice between the day of birth and the nineteenth day result in a six-time increase of the volume of these ganglia (42). The increase is due to a twofold increase in the cell population and a threefold increase in the size of nerve cells (Figs. 26, 27, 30, 31). A parallel increase in the supply of sympathetic nerve fibers to the blood vessels, viscera, and hair was observed. In the adult, the NGF calls forth increase in the size of individual neurons but no increase in their number.

Distribution of the NGF in the organism

Every time during the past years that we detected a new source of the NGF, we hoped to have at last identified its source in the organism. In no instance, however, was the evidence strong enough to implicate the tissue as the "source" of the NGF, and the search is still in progress at the present time. One interesting and perhaps revealing observation was to find that the NGF is present everywhere in the organism (44). It is harbored in exceptionally high amount in the

mouse salivary glands and in their homologue, the snake venom glands; it is present in lesser amount in the salivary glands of other rodents, which in turn seem to possess it in higher amount in the kidney (unpublished observations). It is present in embryonic tissues of chick embryo (45) and in granuloma tissue experimentally produced in different vertebrates (44). It is also detectable in the sensory and sympathetic nerve cells of embryonic birds and in the sympathetic cells of a variety of mammals, man included. A preliminary exploration indicates its presence in human serum.

The functional significance of this agent in the growth and developmental processes of sympathetic nerve cells can hardly be questioned. We have evidence that in the mouse the sympathetic cells remain receptive to the NGF throughout life and it is conceivable (but still not proved) that this applies also to the same cells in other mammals (Fig. 34). Additional support in favor of the hypothesis that the NGF might play an essential role in the growth and differentiation of the receptive nerve cells came from experiments reported in detail elsewhere (41, 46), which showed that the injection of this antiserum destroys selectively the sympathetic ganglia in newborn mammals (Fig. 29).

A re-evaluation of the role of extrinsic factors as controlling agents of growth and differentiation of nerve cells

Before attempting an evaluation of the results presented in the preceding pages, it may be of interest to consider how past interpretations were abandoned as new facts were discovered and as we gained a more precise picture of the phenomenon. We believe that the present information is still at best only partial and fragmentary; this belief and past experience suggest offering a working hypothesis rather than a conclusion in the following pages.

It will be recalled that at first the growth effects elicited by mouse sarcomas on the sensory ganglia of the chick embryo were compared to the effects elicited by implantation of an additional limb in the embryo (34). At that time, the similarities rather than the differences between the two effects were stressed. A reinvestigation and a more close inspection of the phenomenon resulted in a radical change of position. When it was discovered that the tumor elicits a generalized growth effect on the sensory and sympathetic ganglia of the embryo, the differences rather than the similarities between the effects of the

tumor and the effects of an additional limb were stressed (37). We concluded that the tumor calls forth atypical responses which cannot be considered as merely an intensification of normal developmental processes. The question was then raised whether these exceptional effects were not due to some unique property of neoplastic tissues. This hypothesis, suggested in 1953 (37), was ruled out 3 years later when it was discovered that snake venom harbors a nerve growth factor remarkably similar to the NGF of mouse sarcomas. When, 2 years later, it was found that the mouse submaxillary salivary glands also possess this factor, it was definitely established that even normal structures can harbor the NGF. At the same time, the difference between the effects elicited by the implantation of a limb and the effects elicited by implantation of mouse sarcomas or by injection of the venom or of the salivary NGF seemed to come into an even sharper relief than before.

The results obtained with these last factors established beyond doubt that the specific protein can avail itself of different routes to gain access to the nerve cells. While in experiments of transplantation of additional limbs we had only evidence of the role played by nerve fibers in conveying the message from the periphery to the nerve cells, now we were able to show that the NGF can utilize the nerve fibers (intraembryonic tumor transplantation); it can utilize the circulatory channels (chorioallantoic tumor transplantation or injection of the NGF into the yolk); or it can diffuse in the medium, as proved by tissue culture experiments.

A next step in the investigation was the discovery that the NGF is present in the receptive nerve cells and can also be detected in other structures and body fluids. Of particular interest is the finding that it is present in the serum of mammals, man included, and that it is produced by undifferentiated structures such as embryonic cells (45) and granuloma tissue experimentally produced in adult mammals (44).

The above results raised the question whether the NGF is produced in a special organ and from there spread in the organism, or whether it is normally produced by different types of cells. This question is still undecided, although we believe that the evidence is in favor of the second alternative. Since we have evidence that it is produced in anaplastic tumors and it is present in embryonic tissues, we are again faced with the problem which we considered as definitely settled: Are the growth effects elicited by a transplanted additional limb entirely different from the effects elicited by the NGF discussed above? In order to answer this question, one should have a more precise

notion of the biochemical changes which take place in ganglia confronted with an enlarged peripheral field such as the one provided by an additional limb bud. We know that the mitotic activity increases and as a consequence more nerve cells are produced. The effect is restricted to the ganglia which contribute fibers to the graft. This is in fact the main difference between the effects elicited by a transplanted limb and an implanted tumor. There are, however, instances of moderately active tumors where the effect is restricted to the ganglia which send fibers into the neoplastic tissue. The difference is therefore more of a quantitative than of a qualitative order. The massive and generalized effects elicited by rapidly growing tumors, or by injection of large quantities of the salivary NGF, represent exceptional events with no parallel in the normal development of the embryo. Under normal conditions, it seems conceivable that the peripheral end organs release small quantities of growth factors and that these factors may utilize the nerve fibers as channels of diffusion to the associated nerve centers. This hypothesis would explain the mitotic effects and at the same time would not be in conflict with the observation that the effects are restricted to nerve centers contributing fibers to the implanted organs.

In suggesting that the effects evoked by peripheral structures and by the NGF isolated from tumor, snake venom, and the mouse salivary glands might in ultimate analysis operate in a similar way, we wish to make clear that the growth factors need not be the same in all experimental conditions examined above. It was in fact shown that the effects elicited by the transplantation of an additional limb are mainly apparent on the V-L sensory cells while the effects of the tumor and of the purified NGF are mainly and perhaps entirely restricted to the M-D cells and to the sympathetic ganglia.

A confirmation of the above hypothesis would come from the isolation of growth factors from peripheral structures with a specific effect on other nerve cells, such as the V-L sensory cells and the motor neurons. This is one of the problems we propose to investigate. The realization that it is a very difficult problem makes it more challenging and worth the efforts it will require.

References

1. J. F. Tello (1922), Les diferenciaciones neuronales dans l'embryon du poulet pendant les premiers jours de l'incubation, *Trav. Lab. Rec. Biol., Univ. Madrid, T.*, 21:1-93.
2. W. F. Windle and D. W. Orr (1936), Neurofibrillar development in the central

- nervous system of chick embryos up to 5 days incubation, *J. Comp. Neurol.*, **63**:431, 463.
3. F. Visintini and R. Levi-Montalcini (1939), Relazione tra differenziazione strutturale e funzionale dei centri e delle vie nervose nell'embrione di pollo. *Arch. suisses neurol. et psychiat.*, **43**:1-45.
 4. R. Levi-Montalcini (1950). The origin and development of the visceral system in the spinal cord of the chick embryo. *J. Morph.*, **86**:253-284.
 5. R. Levi-Montalcini (1942). Origine ed evoluzione del nucleo accessorio del nervo abducente nell'embrione di pollo. *Acta Pontif. Acad. Sci.*, **6**:335-345.
 6. A. W. Jones and R. Levi-Montalcini (1958). Patterns of differentiation of the nerve centers and fiber tracts in the avian cerebral hemispheres. *Arch. ital. biol.*, **96**:231-284.
 7. V. Hamburger and R. Levi-Montalcini (1949), Proliferation, differentiation, and degeneration in the spinal ganglia of the chick embryo under normal and experimental conditions. *J. Exp. Zool.*, **111**:467-502.
 8. P. Shieh (1951). The neof ormation of cells of preganglionic type in the cervical spinal cord of the chick embryo following its transplantation to the thoracic level. *J. Exp. Zool.*, **117**:359-396.
 9. C. Biondi (1910), Osservazioni sullo sviluppo dei nuclei d'origine dei nervi oculomotore e trocleare nel pollo. *Riv. ital. neuropatol. Psychiatr.*, **3**:302-312.
 10. S. R. Cajal (1929). La charpente neurofibrillaire dans la cellule de Purkinje jeune, et son evolution, in *Études sur la Neurogenese de Quelques Vertebres*, ed. by Instituto de Estudios científicos, Madrid, pp. 192-211.
 11. R. Levi-Montalcini (unpublished observations).
 12. L. L. Uzman (1960). The histogenesis of the mouse cerebellum as studied by its tritiated thymidine uptake. *J. Comp. Neurol.*, **114**:137-160.
 13. V. C. Twitty (1944). Chromatophore migration as a response to mutual influences on the developing pigment cells. *J. Exp. Zool.*, **95**:259-290.
 14. V. C. Twitty and M. C. Niu (1954). The motivation of cell migration, studied by isolation of embryonic pigment cells singly and in small groups *in vitro*, *J. Exp. Zool.*, **125**:541-573.
 15. A. Moscona and H. Moscona (1952). The dissociation and aggregation of cells from organ rudiments of the early chick embryo. *J. Anat.*, **86**:287-301.
 16. C. Grobstein (1954). Tissue interaction in the morphogenesis of mouse embryonic rudiments *in vitro*, in *Aspects of Synthesis and Order in Growth*, 13th Symposium, Princeton University Press, Princeton, N.J., pp. 233-268.
 17. A. Moscona (1957). The development *in vitro* of chimaeric aggregates of dissociated embryonic chick and mouse cells, *Proc. Nat. Acad. Sci. U.S.*, **43**:184-194.
 18. P. Weiss (1945). Experiments on cell and axon orientation *in vitro*: The role of colloidal exudates in tissue organization, *J. Exp. Zool.*, **100**:353-386.
 19. P. Weiss (1958). Cell contact. *Intern. Rev. Cytol.*, **7**:391-422.
 20. A. Moscona (1959). Patterns and mechanisms of tissue reconstruction from dissociated cells, in *Developing Cell Systems and Their Control*, ed. by Dorothea Rudnick. The Ronald Press Company, New York.
 21. J. T. Bonner (1947). Evidence for the formation of cell aggregates by chemotaxis in the development of the slime mold *Dictyostelium discoideum*, *J. Exp. Zool.*, **106**:1-27.
 22. M. Sussman (1952). An analysis of the aggregation stage in the development of the slime molds. Dictyosteliaceae. II. Aggregative center formation by mixture of *Dictyostelium* wild type and aggregateless variants, *Biol. Bull.*, **103**:446-457.

23. M. Sussman and F. Lee (1955), Interactions among variant and wild type strains of cellular slime molds across thin agar membranes, *Proc. Nat. Acad. Sci. U.S.*, **41**:70-78.
24. R. G. Harrison (1935), The Croonian Lecture on the origin and development of the nervous system studied by the methods of experimental embryology, *Proc. Roy. Soc. (London) B*, **118**:155-196.
25. V. Hamburger and R. Levi-Montalcini (1950), Some aspects of neuroembryology, in *Genetic Neurology*, ed. by P. Weiss, University of Chicago Press, Chicago, pp. 128-160.
26. V. Hamburger (1954), Trends in experimental neuroembryology, in *Biochemistry of the Developing Nervous System*, Academic Press, Inc., New York.
27. P. Weiss (1955), Nervous system, in *Analysis of Development*, ed. by Benjamin H. Willier, Paul A. Weiss, and Viktor Hamburger, W. B. Saunders Company, Philadelphia, vol. 7, pp. 346-401.
28. S. R. Detwiler (1936), *Neuroembryology: An Experimental Study*, The Macmillan Company, New York.
29. V. Hamburger (1934), The effects of wing bud extirpation on the development of the central nervous system in chick embryos, *J. Exp. Zool.*, **68**:449-494.
30. V. Hamburger (1939), Motor and sensory hyperplasia following limb-bud transplantations in chick embryos, *Physiol. Zool.*, **12**:268-284.
31. R. Levi-Montalcini and G. Levi (1944), Correlazioni nello sviluppo tra varie parti del sistema nervoso. *Commentationes Pontif. Acad. Sci.*, **8**:527-568.
32. V. Hamburger and R. Levi-Montalcini (1949), Proliferation, differentiation and degeneration in the spinal ganglia of the chick embryo under normal and experimental conditions, *J. Exp. Zool.*, **111**:457-502.
33. M. A. Gerebtzoff (1959), *Cholinesterases*, Pergamon Press, New York, pp. 1-195.
34. E. D. Bueker (1948), Implantation of tumors in the hind limb field of the embryonic chick and the developmental response of the lumbo-sacral nervous system, *Anat. Record*, **102**:369-390.
35. R. Levi-Montalcini and V. Hamburger (1951), Selective growth stimulating effects of mouse sarcoma on the sensory and sympathetic nervous system of the chick embryo, *J. Exp. Zool.*, **116**:321-362.
36. R. Levi-Montalcini (1952), Effect of mouse tumor transplantation on the nervous system, *Ann. N.Y. Acad. Sci.*, **55**:330-343.
37. R. Levi-Montalcini and V. Hamburger (1953), A diffusible agent of mouse sarcoma, producing hyperplasia of sympathetic ganglia and hyperneurotization of viscera in the chick embryo, *J. Exp. Zool.*, **123**:233-288.
38. R. Levi-Montalcini, H. Meyer, and V. Hamburger (1954), In vitro experiments on the effects of mouse sarcomas 180 and 37 on the spinal and sympathetic ganglia of the chick embryo, *J. Exp. Zool.*, **116**:321-362.
39. R. Levi-Montalcini and S. Cohen (1956), In vitro and in vivo effects of a nerve growth-stimulation agent isolated from snake venom, *Proc. Nat. Acad. Sci. U.S.*, **42**:695-702.
40. S. Cohen and R. Levi-Montalcini (1956), A nerve growth-stimulating factor isolated from snake venom, *Proc. Nat. Acad. Sci. U.S.*, **42**:571-574.
41. S. Cohen (1960), Purification of a nerve-growth promoting protein from the mouse salivary gland and its neuro-cytotoxic antiserum, *Proc. Nat. Acad. Sci. U.S.*, **46**:302-311.
42. R. Levi-Montalcini and B. Booker (1960), Excessive growth of the sympathetic

- ganglia evoked by a protein isolated from mouse salivary glands, *Proc. Nat. Acad. Sci. U.S.*, **46**:373-384.
43. R. Levi-Montalcini (1958), Chemical stimulation of nerve growth, in *The Chemical Basis of Development*, ed. by W. D. McElroy and Bently Glass, Johns Hopkins Press, Baltimore.
44. R. Levi-Montalcini and P. U. Angeletti (in press), *Biological Properties of a Nerve-growth Promoting Protein and Its Antiserum*, Pergamon Press, New York.
45. E. D. Bueker, I. Schenkein, and J. L. Bane (1960), The problem of distribution of a nerve growth factor specific for spinal and sympathetic ganglia, *Cancer Research*, **20**:1220-1228.
46. R. Levi-Montalcini and B. Booker (1960), Destruction of the sympathetic ganglia in mammals by an antiserum to a nerve-growth protein, *Proc. Nat. Acad. Sci. U.S.*, **46**:384-391.
47. R. Levi-Montalcini and S. Cohen (1960), Effects of the extract of the mouse submaxillary salivary glands on the sympathetic system of mammals, *Ann. N.Y. Acad. Sci.*, **85**:324-341.
48. R. Levi-Montalcini and P. U. Angeletti (in press), Growth control of the sympathetic system by a specific protein factor, *Quart. Rev. Biol.*

Index

Page references in **boldface** type indicate illustrations

- Absorption bands, 2
- Acetate thiokinase, 83, **84**
- Acetic acid, 32, 34, 46
- Acids, acetic, 32, 34, 46
 - alpha-amino, beta-ketoadipic, 38
 - amino, 10, 34, 72, **235, 236**
 - isozymes, 113
 - delta-amino-levulinic, 38
 - formic, 32, 34
 - hydroxy, 34
 - succinic, 32, 38, 46
- ADP (adenosine diphosphate), 76, 78, 80
 - and ATP, 78
 - and pyruvate, 80
- Aggregation, 224, 237, 240
- Alanine, 249
- Aldolases, 71, 73, **74**, 76, 77
 - muscle, liver, yeast, **75**, 76
- Algae, photosynthesis, in blue-green, 17
 - (See also Blue-green algae)
 - in marine, 17
- Alleles, 173
- Alpha-amino-beta-ketoadipic acid, 38
- Amino acid sequences, 72
- p-Aminobenzoate, 89
- Ammonia, 3, 32, **33**, 60, 61, 63
 - converted to urea, 59
 - irradiation, 47
 - x-ray identification, 33
- Amphibia, 237
- Androgen, endogenous, 151
 - exogenous, 151
- Anticancer agents, 88
- Antiserum, 175, **182**
- Arginase, 62, 63
- Arginine phosphate, 55
- Asexual reproduction, 173
- Assay, pupal, 255
- ATP (adenosine triphosphate), 18, 38
 - ADP-bound, 78
 - cleavage, 71, 77, 78, **81**, 82, 85
 - phosphate by actosin, **87**
 - synthesis, 81
 - hydrolysis, **86**
 - photoproduced, 25
 - pyrophosphate linkage, 19, **21**, 22, **23**, 24, 26, 27, 29
- Autogamy, 175, 193
- Autonomization, 203
- Autoradiography, 233, **234**
- Avian telencephalon, migration of epi-
basal central nuclei, **275**
- Bacteriochlorophyll, 30
 - structure, **31**
- Balbani rings, 248
- Biochemistry and evolution, 45
- Biology, developmental, 96
- Biosynthesis, of coenzyme A by bacteria, **50**
 - of purines, 48
- Blue-green algae, 159
 - chloroplasts, **28**
 - photosynthesis, 17
- Brain, 263
- Brain differentiation, 262

- Carbon, chemistry of, 2
 hydro-, 13
 methane, 14, 34
 oxidized, 5
 on Venus, Mars, 11
 radioactive-11, 18
 reduction cycle, 18, 19, 20
 terrestrial surface, 4
- Carbon compounds, of other planets, 11
 synthesis of, 9
- Carbon cycle in photosynthesis, 18, 21
- Carbon dioxide, 11, 12, 19
 reduction through carbon cycle, dark reactions, 19
 light reactions, 19
- Carbon-14, 18
- Carbon-hydrogen-oxygen compounds, 6
 in metabolism, 7
- Carbon reduction cycle, 18, 19, 20
 evolutionary history, 20
- Carbonaceous compounds, 9, 10
- Catalysis, 72
 auto-, 38
 iron, 39
 protein, 69, 70, 73
- Catalysts, evolution, 37
 rudimentary development, 36
- Cecropia silkworm, 252
- Cell(s), bacterial, 257
 differentiation, 98
 doublet (see Doublets)
 enlargement, 247
 epithelial, mouse, after castration, 153
 heredity, 165, 205, 208-214
 liver, estrogen stimulation, 151
 movement, 278
 nerve (see Nerve cells)
Paramecium, structure, 167
 programming of, 258
 structure, preformed, 165
 synthesis, 243ff.
- Cellophane, 231, 235
- Centrioles, 201
- Chick embryo, nervous system, 263-265, 268, 277, 278
 spinal column, 266
 preganglionic thoracic column, 276
 segregation, in III nerve, 271
 in VI nerve, 270
 thoracic level, 282
 transverse section of VI nerve, 217
- Chitin, 54
- Chlorophyll, bacterio-, 30
 converted phoro-, 30
 structure, 31
- Chlorophyll, structural relation to heme, 41
 structure, 30, 31
- Chloroplasts, 27, 29
 blue-green alga, 28
 green alga, 28
 guinea pig pancreas, 28
 lamellar structure, 29
 sonicate, 30
 spinach, 29, 30
 tobacco, 28
 visible order, 36
- Chondrocytes, 238
- Chromatid, 247
- Chromosomes, 214, 215
 DNA in, 103
 gene, 246
 giant, 106, 247, 249
 Diptera, 106
 puffing in, 247
 puffing patterns, 106, 107, 108, 248
 RNA, 103
- Ciliates, 196, 201, 210-212
 cortical variants, 198
 doublets, 201
- Citrulline formation, 64
- Clones, 174, 175, 178, 179, 181, 183, 184, 200, 209, 244, 246, 257
 phenotype, 173
- Coenzyme A, synthesis, 51, 81, 84
- Cofactor function, 76, 78
 divalent cation, 76
- Collagen filaments, 36
- Conjugation, 193
- Construction manual, 243, 244, 253
 genetic, 252
- Cortex, 201, 205, 208, 211-213, 215
 inductor response systems, 205
 patterns, 197
- Cortical differentiation, 197
- Cortical evolution, multicellular organisms, 211
- Cortical morphogenesis and heredity, 195
- Cortical picking, 184
- Cortical structure, preformed, 214
- Coupling, 40
- Cyst, pupated, 253
- Cytochrome(s), production, 51
 reduction, 25
- Cytochrome c, 87
- Cytodifferentiation, control, 224, 240
 microenvironmental influences in, 223ff.
- Cytogamy, 173
- Cytological technique, 169

- Cytoplasm, 125, 126, 130, 136, 138, 144, 155, 159, 174, 205, 216, 217, 244
 structure, 157
 Cytoplasmic bridges, 175, 181
 Cytopyge, 182, 184, 186, 194, 195, 207, 208

 Dehydration, nonredoxed, 25
 photoinduced, 24
 Delta-amino-levulinic acid, 38
 Determination, embryonic, 257
 Development, embryonic, 244
 Differentiation, 238
 brain, 202
 cell, 98
 cortical, 197
 ER in, 155
 erythrocytes, 97
 macronuclear, 113
 melanocyte, 98, 103
 molecular, 113
 nervous system, 261, 278
 nuclear, 109
 and segregation, 262
 Diffusion layer, 5
 Diphosphopyridine nucleotide, 20, 76, 78
 cofactor, 78
 Diptera, giant chromosomes, 106
 Diversification, gene, 243
 organismal, 223
 Diversity, in cytoplasmic structure, 144
 in ER, 145, 146
 in mitochondria, 122
 protein, 95
 subcellular, 121
 DNA (deoxyribonucleic acid), 88, 97, 244
 in chromosomes, 103, 104
 replication, 106
 Doublets, 167, 171, 172, 179-181, 184, 185, 188, 193, 198-201
 ciliate, 201
 clone, 200
 conjugants, 178
 cortical variants, 198
 fission, 190
 killer, 176
 kinety fields, 190
 oral apparatus, 209
 oral rudiment, 186, 189
 Paramecium, 175
 sensitive, 176
 (See also Cell(s))
 DPNH-cytochrome reduction, 25

Drosophila, 106, 108
 esterases in, 114

 Ecdyson, 106, 108, 252, 254, 255
 Ectoplasm, 171
 Egg, totipotent, concept, 257
 Egg mosaic, 257
 Elasmobranchs, 62, 63
 Embryo, chick (see Chick embryo)
 Embryonic determination, 257
 Embryonic development, 244
 Endoplasm, 173, 179, 205
 (See also ER)
 Energy, chemical, 20
 electromagnetic, 20
 in bacteria, 18
 Gibbs free, 8
 radioactive, 7
 sources, earth, 7
 , sunlight, 8
 Enolase, 73
 Enzyme(s), 70, 89, 91, 166, 215, 217
 acetate thiokinase, 83, 84
 aldolases, 71, 73, 74, 76, 79
 muscle, liver, yeast, 75, 76
 amino acid sequences, 171
 p-aminobenzoate, 89
 catalyzing, 178
 cofactors, 70
 constitutive, 245
 control, genetic, 88, 89
 dehydrogenases, 78
 in *Drosophila*, 113
 enolase, 73
 esterase, 113
 genetic and ontogenetic control, 113
 glutamine synthesis, 82
 hexokinase, 79
 hydrolytic, 71, 72
 induced change, 88
 inductive, 245
 isozymes (see Isozymes)
 mechanistic relations, 78
 in microbial systems, 245
 papain, 73
 phosphorylation, muscle contraction, 85
 pyruvate kinase, 78-80
 resistance to antibiotics, 88, 90
 specificity, 82
 synthesis, 245
 tyrosinase, 101-103, 108
 Episomes, 246
 Epithelium, 238
 lentogenic, 224
 morphogenesis, 239

- Epithelium, salivary, 239
tubules, 227
- ER (endoplasmic reticulum), 126, 128, 130, 132, 134, 136, 138, 142, 144, 145, 151-153, 155, 159, 160
differentiation, 238
diversity, 145, 146
modulation, 151
rough, 146, 148
smooth, 147
structural continuities, 148
- Erythroblasts, 142
- Erythrocytes, differentiation, 97
- Esterase, 113
Drosophila, 106, 108
evolution and biochemistry, 45
theory of, 1
- Fatty acids, 34
- Fertilization, 200
- Filter, 232, 233
- Fission, 186, 188, 195, 196, 198, 209, 210
abortive, 200
singlets and doublets, kinetic fields, 190
oral rudiment, 185, 186, 189
- Flagellates, 212, 213
zoo-, 213
- p*-Fluorophenylalanine incorporation, 77
- Formic acid, 32, 34
- Galactogen, 54
- Gene(s), 215, 243-245
am, 179
construction manual (see Construction manual)
diversification, developmental, evolutionary, 243
environmental effects, 98
function, 98
regulation of, 96
in giant chromosomes, 246
heterozygous, 175
inhibition or activation, 98
operator, 245, 246
products, 166
regulator, 245
repression, 242
- Genetic information, control in microbial systems, 245
endocrine control, 254
management of, 256
- Genotype, 173, 175, 198, 200, 216
micronuclear, 178
- Glaucoma*, 198, 203-205, 208
- Glomeruli, 227
- Glucagon injection, 149
- Glucolytic breakdown, 150
- Glucose secretion, 149
- Glutamine, 81
- Glutamine synthetase, 82
- Glutathione, 81
- Glycine, 32, 34, 38, 46, 249
x-ray identification, 34
- Glycoeyamine, 56, 57
- Glycogen, 34, 124, 136, 149, 155
- Golgi complex, 130, 142, 144, 153, 159
- Grain-counting procedure, 233, 235, 236
- Greenhouse effect, 12
- Heme, 41
structural relation to chlorophyll, 41
- Hemoglobin, 52, 142
adult, 97
chemical differences, 52
fetal, 97
gain mutation, 50
sickle-cell, 51
C, 52
S, 52
synthesis, 97
- Heterocyclic pyrrole ring, 38
- Histones, 104
cleavage, 105
- Homarine, 54
- Hormone, juvenile, 255, 256
- Hydrogen, 4, 9, 10
chemistry of, 2
escape from earth, 5
irradiation, 47
oxidized, 4
- Hydrogen peroxide, decomposition, 38
- Hydroxy acids, 34
- Imaginal discs, 251
- Induction, 224, 225, 228, 240
- Insects, differentiation, 243
growth, 246
metamorphosis, 246
microbes, 243ff.
morphogenesis, 243
- Insulin, 52
species specificity, 53
- Invagination, 193
- Iron, catalysis, 39
ferrous, 38
function, 38
- Irradiation, ammonia, 47
hydrogen, 47
methane, 47
- Isozymes, 72, 111, 113

- Isozymes, amino acid sequence.** 113
 aspaerokinase, 117
 esterase, 113
 lactate dehydrogenase, **112**
 LDH, 115
 lysine, 117
 molecular changes, 113
 threonine, 117
 transplantation, 115
 (See also Enzymes)
- Killer trait.** 174
Kinetics, interpolar dorsal. 172
Kinetosome(s). 186, 188, 192, 196, 210, 213, 215
 self-reproduction, 202
 theory of, 201
Kinety. 171, 188, 192, 193, 195, 202, 207
 elongation, 188, 192, 215
 endoral, 170, 208
Kinety fields, 195
 doublets, **190**
 fission, **190**
 singlets, **190**
Kinety patterns, 186
- Lactate dehydrogenases.** 72, 76, 39
Lamellar cisternae. **159**
 ribosomes, **132, 142, 144, 153**
Lamellar structure, chloroplasts, **29**
Larva, cuticle. 253
Larval molts, 253
Leucophrys patula, 209
Limestone. 6
Limiting membrane. 216
Lombricin. 56
Lysosomes. 144
- Macromolecules.** 215
Macronucleus. 179
Mars, 12
Matrix. **125, 126, 128, 278**
 cytoplasmic, **134, 136, 140, 142, 151, 159**
Melanin. 101
 in neural retina, 101
 chick, 102
 in pigmented retina, 101, **103**
 tyrosinase synthesis, 101–103, 108
Melanoblasts, 98, 101
Melanocyte, 98
 dendritic, 98
 differentiation, 98, 103
 epithelial, 98
 mammalian, 98
Melanogenesis, 98
- Melanosome(s).** 99, 101
 formation, **101**
 pigmented retina, 102, **104, 105**
 in protein fibrils, 99
Meridian(s). 86, 192
 oral, 190, 194
 fusion, 194
Mesenchyme, metanephrogenic. **224, 226, 230, 232**
 nephrogenic, 225
 salivary, 239
 spinal cord, 225, 228
 ureteric bud, 224, 234
Metabolic maps. 246
Metamorphosis, endocrine control, 251, 254
 insect, 246
 of skin implants, 253
Metanephric rudiment, **226**
Metanephrogenic layer, 226
Metanephrogenic mesenchyme, **224, 226, 230, 232**
Metanephros. **227**
 mouse, 224
Metazoa. 211, 212
Meteoric matter, 3
Methane. 32, **33**
 carbon-14, 34
 irradiation, 47
 x-ray identification, **33**
Microbes, genetic control. 245
 versus higher organisms, 246
 and insects, 243ff.
Microenvironment. 224, 228, 237
Microvesicles. 144
Migration. 237
 nuclei, **275**
Mitochondria. **124, 125, 128, 132, 134, 140, 142, 155, 159**
 diversity, 122
 structural variations, 123
Molecule(s), autocatalysis. 46
 catalysis, by heavy metals, 46
 by organic compounds, 46
 differentiation, 113
 organic origin, 1
 primeval and primitive, **32**
 synthesis, 46
Monolayer culture. 238
Multivesicular bodies. 144
Mutation. 1
 gain, hemoglobin, 50
- Nerve cells, growth response,** 284
 to diffusible protein agent, 285
 migratory patterns, 278

- Nerve cells, sensory, 262
 implantation of mouse sarcomas, 284, 285
 sympathetic, implantation of mouse sarcomas, 288
 response to purified salivary NGF, 289
- Nerve centers, differentiation and growth, 278
 role of peripheral effectors and receptors in, 278
 sensory ganglia, 280, 281
- Nervous system, central, 261
 chick embryo, 263
 developmental mechanisms, 263, 278
 differentiation, 261
 first differentiative phase, 263
 growth of, 261
 mass migration, 268ff.
 Purkinje cells, 280
 regional patterns, 265ff., 268
 second differentiative phase, 264
- Neural tube, 265, 278
- Neuroblasts, 263-265
- NGF, 291, 292
 distribution, 289
 response in sympathetic nerve cells, 289
- Nitrogen, 12
 chemistry of, 2
 excretion, 59, 60
- Nucleohistone, 106
- Nucleotide, diphosphopyridine, 20, 76, 78
 pyridine, 20-27
 sequence, 244
 triphosphopyridine, 21
- Nucleus, 179, 216
 cellular determination of size, 180
 cellular differentiation, 109
 macromolecular fractions, dialyzed, 109, 115
 injected into fertilized eggs, 109
 migration, 275
 multiplication in foreign cytoplasm, 109
- Octopine, 57
- Opalinids, 213
- Operon, 245
 adult, 252
 larval, 252
 pupal, 252
- Opisthe, 196
- Optic vesicle, 224
- Oral apparatus, 196
 doublet, 209
 meridian, 183, 190, 194
 Paramecium, 170, 175, 209
 rudiment, doublet, 186, 189
 singlet, 186, 189
- Organelles, 198, 199, 209
- Organic compounds, phosphate in, 10
- Ornithine, 62
- Orthophosphate, 38
 converted to pyrophosphate, 40
- Oxidation, of carbon, 5
 of nitrogen, 5
 phosphorylation, 25, 26, 72, 86, 87
 of surface elements, 6
- Oxidation state, 5
- Oxygen, atomic abundances relative to, 4
 chemistry of, 2
 free, 6
 molecular, 22
- Paramecium aurelia*, 167, 168, 184, 206-208, 210, 212-214, 216
 cell structure, genetic function, 167
 cortical structure, 167
 hereditary differences in singlets and doublets, 175
 oral apparatus, 170, 175, 209
- Parasomal sac, 171
- Periodic acid-Schiff procedure, 232
- Petroleum, 3
- PGA (phosphoglyceric acid), 19
- Phenotypes, 173
 clonal, 175
- Phosphagens, 55, 56
- Phosphate, carbamyl, 62
 creatine, 55
 in organic compounds, 10
- Phosphorylation, aldolases, 71
 enzymes, 81
 oxidation, 25, 26, 72, 86, 87
 reversibility, 25
- Photosynthesis, 16-18
 in algae, 17
 alkaline, 114, 115
 acid diagram, 116
 apparatus, 17
 ATP, 25
 in bacteria, 17
 carbon cycle, 18, 21
 evolution, 15, 17
 in green plants, 17, 27
 mechanism, 17
 modern, 16
 primitive, 32
 quantum conversion, 20

- Photosynthesis, scheme, **23**
 subcellular, 17
 submolecular, 17
 Plasmaphoresis in liver cells, 153
 Plasmatype, 216
 Polypeptides, synthesis, 97
 Porphyrins, 40
 biosynthesis, **39**
 generation, 36
 Potassium-40, beta rays, 32
 Primordium (anlagen), 203-205
 Protein(s), 233
 catalytic, 69ff., 73
 chemical activity, 35
 diversity, 95
 specific, origin, 95
 structure, 35
 tertiary, 244
 synthesis, 97, 244
 Prothoracic glands, 252
 Puffing phenomenon, de-repression, 248
 in giant chromosomes, 247
 Pupa, assay, 255
 cyst, 253
 diapause, 251
 operon, 252
 Purines, biosynthesis, 48
 coenzyme A, **50**
 Purkinje cells, 280
 migration in chick embryo, **273**
 Pyridine nucleotide, 20, 22, 24, 26, 27
 reduced, 24
 Pyrophosphate, 24
 formation, 38
 linkage, ATP, **19, 21, 22, 23, 24, 26, 27, 29**
 nonphotosynthetic, 26
 Pyruvate kinase activity, **78, 79**
 phosphoryl formation, **80**
- Quadruplets, 185
 Quantum absorption, 18
 Quantum conversion, 18
 in photosynthesis, 20
- Radiation, ionizing, cyclotron source, 34
 K⁴⁰, 38
 Radioactive carbon-11, 18
 Radioactive energy, 7
 Redox system, photoinduced, 22
 Reduction cycle, carbon, 18-20
 carbon dioxide, 19
 DPNA, 25
 phosphate, by ferrous iron, 40
 Regeneration, macronuclear, 175
 Replication, DNA in, 106
- Repressor substance, 245
 Reproduction, asexual, 173
 Reptiles, chelonian, 60, 61
Rhynchosciara angelae, 246ff.
 synchronization, 247, 248
 Ribosome(s), 97, **126, 128, 159, 166, 217, 244, 250**
 lamellar cisterna attached, **132, 142, 144, 153**
 RNA in, 249
 Ribulose diphosphate, 18, **19**
 RNA (ribonucleic acid), 97, 106, 217, 244
 bulk, 250
 in chromosomes, 103
 messenger, 244, 250, 256
 ribosomal, 249
 synthesis, 240, 249
 Rudiment, 196
 oral, 185, **186, 189**
- Sarcolemma, 149
 Sarcoplasm, 149
 Self-selection mechanism, autocatalytic, 38
 Sensory nerve cells, 262, 284, 285
 Seryl residue, **72, 73**
 Singlet(s), 179-**181**, 188, 197, 198, 200, 201
 killer, **176**
 kinety fields, **190**
 oral rudiment, **186, 189**
 organization, 197, 199
 Paramecium, 175
 sensitive, **176**
 Solar material, 3
 Solar matter, 4
 Spinal column, chick embryo, 266
 regional differences, **266**
 Spinal cord, 232, **235, 236**, 263, 264
 mesenchyme, 225, 228
 Spinal ganglia, **283**
 Sponges, 237
 SR (sarcoplasmic reticulum), 149, 152
Stenostomum incaudatum, 211
Stentor, 198, 203-205, 214
 Substrates, 166
 Succinic acid, 32, 38, 46
 Sulfur, 3
 Surface pressure rate, 5
 Sutures, postoral, 194
 preoral, 169
 Sympathetic nerve cells, 288, 289
 Syncaryon, 175
 Synthesis, carbon compounds, 11

- Synthesis, cell, 243ff.
 coenzyme A, 81, 84
 enzyme, 245
 hemoglobin, 97
 mechanism, 49, 50
 molecular, 46
 polypeptides, 97
 protein, 97, 244
 RNA, 240, 249
 tyrosinase, 101-103, 108
 urea, 62, 65
- Taurocyamine, 56, 57
- Teratological phenomena, 199
- TPNH (triphosphopyridine nucleotide),
 19, 21
 photoproduced, 25
- Transfilter inductive effect, 232
- Trichocysts, 171
- Triose phosphate, 18, 19
- Triose phosphate dehydrogenase, 26
- Triplets, 185
- Trypsin, 224, 232, 233
- Tyrosines, 74
 release, 74
- Ultraviolet, far, 32
- Ultraviolet irradiation, 38, 207
- Urea, 58, 61, 63
 ammonia converts to, 59
 formation, 64, 65
 production, 59
 synthesis, 62, 65
 in *Xenopus laevis*, 58
- Ureogenesis, 62
- Ureteric bud, 224, 225
 mesenchyme, 224
- Uric acid, 61
- Vacuole(s), contractile, 193
 meridians, 172
 pores, 185, 186
- Vestibular-gullet juncture, 207
- Water, 3, 12, 32-34
- X-ray identification, ammonia, 33
 glycine, 34
 methane, 33
Xenopus, 212