Mark Rolles

Illustrations
of
British Birds.
by
H. L. Meyer.
Vol. 4.

London, Longman & Co.

596
674
17612
1837
v. 4
SCNHRB

Contents of Vol. 4.

Water Birds.

Bewick's Swan -	Little Auk
Sheldrake	Great Auk
Ruddy Sheldrake	Razor Bill
Shoveller	Puffin
Gadwall	Cormorant
Mallard	Shag
Avotail -	Solan Gannet
Teal	Caspian Tern
Gargany Teal	Sandwich d.
Bimaculated Duck	Common d.
Wigeon	Roseate d.
Summer Duck	Arctic d.
Black Scoter	Lesser d.
Velvet Scoter	Black d.
Surf Scoter	Gull billed d.
Eider Duck	Black Noddy
King Eider	Sabines Gull
Red Headed Pochard	Little Gull
Red Crested Pochard	Brown headed gull
Nyroca Pochard	Pewit Gull
Scaup. Pochard	Laughing Gull
Tufted Pochard	Common Gull
Western Pochard -	Kittiwahe
Long-tailed Hareld	Ivory Gull
Golden-eye	Glaucus Gull
Harlequin Duck	Iceland Gull
Goosander	Herring Gull
Red breasted Merganser	Larger Herring Gull
Hooded Merganser	Great black backed Gull
Smew	Lesser d.
Red necked Grebe	Skua (common)
Red crested Grebe	Pomerine Skua
Slavonian Grebe	Arctic Skua
Eared Grebe	Richardson's Skua
Little Grebe	Fulmar Petrel
Northern Diver	American Petrel -
Young ditto	Manks Petrel
Black throated Diver	Strickland's Petrel
Red throated Diver	Storm Petrel
Foolish Guillemot	Fork tailed Petrel
Black Guillemot	Wilson's Petrel.

BEWICK'S SWAN.

Cygnus Bewickii, (Yarr.)

Length 3 ft 9 inches. Winter visitant. Food, aquatic plants, insects and small fish.

SHELDRAKE.

Anas Belloni. (Steph.)

Length, 20. Natural Size.—Indigenous
to the Arctic Regions. Insects, etc. Eggs, 80.

RUDDY SHELDRAKE.

Tadorna rutila. (Steph.)

Half the Natural Size...rare visitant.
Food, small fry & insects.

SHOVELLER.
Nº1. Male
2. Female.
Spathulæ clypeata, (Flem.)

Half the Natural Size. Indigenous. Food worms, insects and their larvæ. Breeds among the marshes. Eggs 10 to 12. The male changes his plumage towards autumn so as nearly to resemble the female.

GADWALL.

Adult Male.

Chauliodus strepera; (Swainson).

Half the natural size. Rare winter visitant. Food fish insects etc. Breeds in marshes. Eggs 9 or 10. Male and female nearly alike.

MALLARD.
Anas Boschas, (Linn.)

Half the Natural Size. Food insects, grain and water plants. Winter visitant some few remain to breed. Frequent lakes and rivers.
 Nest placed on the ground by the water-side. Eggs 10 or 12.

M. M. (1858)

PINTAIL DUCK.

Anas acuta, (Linn.)

Half the Natural Size. Winter visitant. Food aquatic insects and small fish. Eggs 8 or 9.

TEAL.
Adult Male.
Querquedula crecca; (Stephens.)

Length 14½ inches. Food insects and aquatic plants. Winter visitant. Eggs 10 to 12. The female resembles the male only in the Colour of the speculum all the rest of her plumage is mottled with black and brown.

GARGANY TEAL.
Adult Male.
Querquedula circia; (Steph.)

Half the Natural size. Occasional visitant. Food seeds and aquatic plants. Eggs 10 or 12 deposited on the ground, by the side of lakes or rivers.

BIMACULATED DUCK.

Anas falcata. (Pallas)

About half the Natural Size very rare.

WIGEON.
Male in Winter Plumage
Mareca Penelope, (Steph.)

Half the natural size. Food aquatic insects, small fish etc. Winter visitant. Eggs 8 or 9.

SUMMER DUCK.

Dendrocygna Javanica. (Swamp)

Half the Natural Size, Rare visitant.

BLACK SCOTER.

Oidemia nigra, (Flem.)

Half the Natural Size. Winter visitant. Food small shell-fish. Breeds in the Northern climes

VELVET SCOTER.

Male and Female

Oidemia fusca; (Flem.)

Half the Natural Size. Periodical Visitor. Food, spawn of fish and mollusca. Eggs 8 or 10.

SURF SCOTER.

Oidemia perspicillata. (Flem.)

Half the Natural Size — Rare visitant,
Food, bivalve shell fish & crustacea.

EIDER DUCK (MALE)
Somateria mollissima. (Leach)

about Half the Natural Size.—Indigenous.

Food, shell-fish, etc.—Eggs, 5.

KING EIDER.

Somateria spectabilis (Leach)

One half of the Natural Size, Food,
Shell Fish &c. — Eggs, 4 to 6.

RED-HEADED POCHARD.

Fuligula ferina; (Steph.)

1. Male 2. Female.

Length 19 inches Food marine plants and fish Visits the British coast in Spring and autumn and retires Northward to breed Eggs 12 or 13 deposited among the reeds. The young of the year resemble the female.

RED-CRESTED POCHARD.

Fuligula rufina. (Steph.)

Half the Natural Size. Rare visitant.

NYROCA POCHARI.

Fuligula Nyroca. (Steph.)

One half the Natural Size. — Rare visitant.
Food, aquatic plants and Insects.

No 2.

SCAUP POCHARD.

Fuligula marila; (Steph.)

No 1. Male in immature Plumage.

2 Female.

Half the natural Size Winter visitant Food fish, marine plants and insects. The adult male in perfect plumage has the neck and breast pitch black the belly and flanks pure white, forehead entirely black.

TUFTED POCHARD.

Adult Male.

Fuligula cristata (Steph.)

Half the Natural Size. Periodical Visitor. Food aquatic insects. Breeds in High Northern Latitudes.

WESTERN POCHARD.

Fuligula dispar. (Steph.)

Length 17 inches. — very rare.

LONG-TAILED HARELD.

No. 1. Adult Male in Winter.

No. 2. do do in Summer.

Harelda glacialis, (Leach.)

Half the Natural size. Winter Visitant on the Northern Coast. Food marine insects and mollusca. Breeds near the Sea in the Arctic Regions. No. 2 is the Anas byemalis of Linn.

GOLDEN EYE
N.Y. Male, 2. Female.
Anas Clangula, (Linn.)

Length 19 inches Food small fish, marine plants and insects Winter visitant Returns far north to breed on the borders of lakes or on the sea shore depositing from 12 to 14 eggs. The young resemble the female.

HARLEQUIN DUCK.
Anas histrionica, (Linn.)

Half the natural size. Very rare occasional visitant. Food aquatic insects etc. Eggs 10 to 12.

GOOSANDER.
Mergus Merganser. (Linn.)
Half the Natural Size. Indigenous.

RED-BREASTED MERGANSER.

Mergus serrator, (Linn.)

About half the natural size. Winter visitant. Food fish and reptiles. Eggs 9 deposited on the ground, in a nest made of sticks, lined with the down of the female.

HOODED MERGANSER.

Mergus Cucullatus. (*Linn.*)

Half the Natural Size. Rare visitant.

Food, fish. — Eggs 8 or 10.

No. 1. SNEW. Adult Male.
Mergus Albellus, (Linn.)

Length of the male 17 inches, female 16 inches. Food fish. Winter visitant, frequenting the coast and rivers. Eggs from 10 to 12, deposited on the banks of rivers and lakes. No. 2. Adult Female. The young of the first year resembles the female. No. 3. Male of the 2nd year.

RED-NECKED GREBE.

Podiceps rubricollis. (*Lath.*)

Half the Natural Size.—Periodical Visitor.

ALMiger 1858

GREAT CRESTED GREBE.

Podiceps cristatus, (Lath.)

Half the Natural size. Indigenous food fish and shell-fish. Eggs 4.

SLAVONIAN GREBE.

Nº 1. Adult. Nº 2. Young.

Podiceps cornutus (Lath.)

Two thirds of the Natural Size. Indigenous. Food small fish and marine insects. Eggs 3 or 4.

REDFEETED GREBE.

Podiceps Auritus. (Lath.)

Two thirds of the Natural Size.
winter visitant. Eggs, 3 or 4.

H. Meyer fec.

LITTLE GREBE.
Podiceps minor, (Lath.)

Length 10 inches. Food small fish and aquatic insects. Indigenous. Frequents lakes and rivers where it forms a nest of dried rushes among the flags and reeds. Eggs 5 or 6. In the female the breast and belly are tinged with ash grey.

NORTHERN DIVER: *Nº 1. Adult Male.*
Colymbus glacialis, (Linn.)

Length 41 inches. Food small fish and marine insects. Frequents the northern Scottish isles rarely found in the more southern parts of Britain although the young are frequently met with all along the coast. Eggs 2, deposited on the borders of fresh water lakes. *Nº 2.* represents the young of the year

Nº 2. Young of the year of the Northern Diver.

Egg of the Northern Diver.

BLACK THROATED DIVER.

Colymbus arcticus (Linn)

One third of the Natural Size, Occasional
visitant. Food, Fish & Reptiles, &c. Eggs 2

RED-THROATED DIVER.

Colymbus septentrionalis. (Linn.)

Two fifths of the Natural Size. Periodical
visitant. Food, small fish. Eggs, 2.

FOOLISH GUILLEMOT.

Uria Troile. (Latham)

H. L. Meyer fecit

N° 1. Adult in Summer Plumage. 2. Adult in Winter Plumage.

Length from beak to feet 15 or 16 inches. Locality, the Arctic Seas whence they migrate in great numbers on the approach of winter to the coast of Norway, England &c. remain until the spring to breed. The female deposits her single egg in a hole in rocks on the sea coast. Male & female alike.

BLACK GUILLEMOT.

Uria Grylle; (Lath.)

Length 15½ inches.—Indigenous.

Food, Marine Insects & small Fish.

LITTLE AUK.
Mergulus melanoleucus; (Ray.)
 (Winter Plumage.)

Three fourth of the Natural Size. Periodical Visitant. Food marine insects and shell fish. Egg one deposited upon the ledge of a rock. In summer Plumage the sides of the head and neck and the breast are black.

GREAT AUK.
Alca impennis. (Linn.)

Half the natural size.—Occasional Visitor.
Food, Fish—Larvæ & egg.

RAZOR BILL. *Adult in Summer Plumage.*

Alca Torda; (Linn.)

Length 18 inches. Food small fish and marine insects. They visit the British shores in summer and breed on the highest rocks on the coast depositing one egg. In winter those young birds that have been reared further north visit our Islands in the plumage of Nº 2. Black-billed Auk of *Derrick*. *Alca pica (Linn.)* which nearly resembles that of the adult at that season.

PUFFIN.
Fratrcula Arctica. (*Steph.*)

Threefourths of the Natural Size. Summer Visitant. Food small fish and marine insects. Egg 1, deposited in a burrow which the bird excavates for that purpose.

CORMORANT.

Phalacrocorax Carbo. (Steph.)
MALE & FEMALE.

One third of the Natural size. — Indigenous. Food, Fish. Eggs 3 to 5.

S H A G.

Phalacrocorax cristatus. (Steph.)

One third of the Natural size. Indigenous Food, Fish. Eggs 4 or 5.

SOLAN GANNET.

Adult

Sula Bassana, (Briss.).

About one third of the Natural Size. Indigenous. Food fish, lays one Egg.

CASPIAN TERN.

Summer Plumage.

Sterna Caspia, (Pall.)

Half the Natural Size. Occasional Visitor. Food fish. Eggs 2 or 3.

SANDWICH TERN.

Summer Plumage

Sterna Cantiaea, (Gmel.)

Length 18 inches. Food small fish etc. Summer Visitant. Eggs 2 or 3. In winter the forehead and crown are white.

COMMON TERN.

Sterna Hirundo, (Linn.)

About two thirds of the Natural Size. Food small fish etc. Summer Visitant. Eggs 2 or 3 deposited on the Sand.

41-18-20-22
1-2-3-4

ROSEATE TERN.
Sterna Dougalli. (Mont.)

Length 16½ inches. Periodical visitant. Food small fish and marine insects. Nidification much the same as that of the other terns.

ARCTIC TERN.
Sterna arctica (Temm.)

Two thirds of the Natural Size — Summer Visitant.

Food, small Fish. — Eggs 3 or 4.

LESSER TERN.

Sterna minuta, (Linn.)

Length 9 inches. Food, marine insects. Summer Visitant frequenting the sea coast during the breeding season where it deposits 2 or 3 eggs on the ground. Male and Female alike.

BLACK TERN.

Male in Summer Plumage.

Sterna nigra, (Linn.)

Length about 10 inches. Summer visitant. Food worms and aquatic insects. Eggs 2 to 4, deposited on the ground among sedges. In winter the forehead neck and breast are white.

GULL-BILLED TERN

Sterna Anglica (Mont)

Length 13 inches. - Rare visitant. - Food,
insects. - Eggs 3 or 4

BLACK NODDY.

Sterna stolida. (*Linn.*)

Two thirds of the Natural Size.

SABINE'S GULL.
Xema Sabini. (*Leach.*)
Two thirds of the Natural Size.

LITTLE GULL.

Larus minutus, (Pallas).

Adult in perfect Summer Plumage.

Two thirds of the natural size. A very rare visitant from the Eastern parts of Europe, supposed to breed on the banks of the Wolga and other rivers in Russia. Food worms and insects. Nidification unknown.

BROWN HEADED GULL.

Larus capistratus. (Temm.)

Half the Natural Size.

PEWIT GULL.
Larus ridibundus, (Linn.)
 N^o 1. Adult in Summer Plumage.
 „ 2. Immature

Half the natural size. Food small fish, worms and insects. Indigenous. Eggs 3. The adult in winter has the head and neck entirely white with the exception of a black spot before the eye and a dusky patch on the ear coverts.

LAUGHING GULL.

(in winter plumage.)

Larus Atricilla. (Linn.)

Half the Natural Size.—Occasional visitant.

COMMON GULL.

Larus canus, (Linn.)

No. 1. Adult in Summer Plumage.

2. Young after the first Autumnal Moults.

Length 17 inches Food small fish, worms and marine insects etc. Indigenous. Eggs 3. In winter the adult has the head and nape spotted with dusky.

H. L. Meyer fecit

KITTIAKIE.
N^o 1 Adult in Summer Plumage.
2. Young after the first Autumnal Molt.
Larus tridactylus, (Lath.)

Length 16 inches. Food fish and marine insects. Eggs 3.

IVORY GULL.

Larus eburneus. (Gmel.)

Two thirds of the Natural Size. rare visitant.

GLAUCOUS GULL.

Larus glaucus. (Brunn)

One third of the Natural Size. Periodical
visitant. — Food, fish, and young birds &c.

ICELAND GULL.

Larus islandicus. (*Edmonson*)

One third of the Natural Size. Periodical
visitant. Food, Fish & Carrion.

HERRING GULL.

Larus argentatus.

Length 23 Inches. Food, Fish &c. Eggs 2 or 3.

LARGER HERRING GULL.

Larus argentatoides (Bonap.)

Two fifths of the Natural Size.

GREATER BLACK BACKED GULL.

Adult in perfect Winter Plumage.

Larus marinus, (Linn.)

One third of the Natural Size. Indigenous. Food. Fish etc. Breeds in various islands on the coast. Nest placed on the shelf of a rock. Eggs 3 or 4.

LESSER BLACK-BACKED GULL.

Adult in perfect Summer Plumage

Larus Fuscus, (Linn.)

About one third of the natural size Food Fish, worms &c. Indigenous Nest of dry grass placed upon the ground. Eggs 3 or 4.

COMMON SKUA.

Lestris Cataractes. (Temm.)

Nearly half the Natural Size. — Indigenous.
Food, fish. Eggs, 2.

Wittig 1845

POMARINE SKUA.

Lestris pomarinus. (Temm.)
(young)

Half the Natural Size. — occasional visitant.
Food, fish, carrion, etc. — Eggs, 2 or 3.

ARCTIC SKUA.

Lestris Parasiticus, (Temm.)

Length 21 inches Rare visitant Food principally fish.

RICHARDSON'S SKUA :

Lestris Richardsonii (Swain.)

Length 21 inches. — Periodical visitant —

Food, Fish. — Eggs 2.

FULMAR PETREL.

Fulmarus glacialis (Linn.)

One-half of the Natural Size. Indigenous.
Food. Carrion, Blubber & Offal. — Egg 1

MANX PETREL.
Procellaria Anglarum (Temm.)

Two thirds of the Natural Size. Summer Visitant.

Food, Fish &c. Lays one egg only.

46. 1/2 2/8 1/2

STRICKLANDS PETREL.
Procellaria fuliginosa. (Kuhl.)

Half the Natural Size.—very rare.

STORM PETREL.
Procellaria pelagica, (Linn.)

Natural Size. Food, marine insects. Indigenous. Is found in all the seas around Britain, skimming the surface of the water, in pursuit of its food. Breeds upon the rocky coast. Eggs 1 or 2.

FORK TAILED STORM PETREL.

Procellaria Bullockii; (*Flem.*)

Length $7\frac{1}{2}$ inch; Indigenous. Food marine insects etc.

WILSON'S PETREL
Procellaria Wilsoni. (*Bonap.*)

Natural Size. very rare.

SILVER-WINGED TERN.
Sterna leucopleta.

BULWER'S PETREL.
Thalassidroma Bulweri.

BLACK SWAN.
Cygnus atratus

CINEREOUS PETREL.

Procellaria cinerea. (Kuhl.)

In the second year.

Half the Natural Size.

ANDALUSIAN HEMIPODE.
Turnix tachydroma.

BARBARY PARTRIDGE.
Perdix petrosa

WHITE WAGTAIL.

Motacilla alba

Egg of Magpie

Egg of Bittern

1000. *Strophomena*

1001. *Strophomena*

1002. *Strophomena*

Great White Heron.

Crane.

Spizella monticola

Spizella monticola

Greenish-brown

Purple-brown

Egg of Sandpiper

Egg of Hawk Owl

Egg of the Golden Eagle.

Egg of the White-tailed Eagle.

Turtle Egg

Small Egg

Great Egg

Spotted Plover

Meadow Lark

Crowsbill

Black Woodpecker

Green Egg

Orange Egg

Blue Egg

Pink Egg

f QL
674
.M692
1837
v. 4
SCNHRB

Meyer, H. L.
Illustrations of
British birds
1837

SMITHSONIAN INSTITUTION LIBRARIES

3 9088 01570 9108