When duplicate of this book is required, send us the TITLE of the book, and NUMBER below.

No. 13328

VAN KLEECK, CLARK & Co.,
Blank Book Manufacturers,
AND STATIONERS,
No. 233 BROADWAY, NEW YORK.

A
B
C
D
E
F
G
H
I
J
K
L
M
N
O
P
Q
R
S
T
U
V
W
X
Y

COLLECTING LIST
OF
EDGAR A. MEARNS,
HIGHLAND FALLS,
ORANGE COUNTY, NEW YORK.

A
B
C
D
E
F
G
H
I
J
K
L
M
N
O
P
Q
R
S
T
U
V
W
Y

RECORDING SCHEDULE.

1. Observations on Birds Shot.
2. Color of Eyes, Bill and Feet.
3. State of Plumage.
4. Contents of Stomach.
5. Description of Nests Collected.
6. Record of Trip.
7. Haunts and Habits.
8. Manners and Actions.
9. Pairing.
10. Songs.
11. Migration Notes.
12. Birds Seen not Shot.
13. Absence of Birds Seen Before.
14. Abundance or Scarcity.

The Bird Names in the Index are from the A.O.
U. Check-List, as amended by 1st and 2nd Supplements.
Acanthis linaria. — Specimens (Take index A. O. U. Check-List.)

See Index to Specimens, p. 466.

A
B
C
D
E
F
G
H
I
J
K
L
M
N
O
P
Q
R
S
T
U
V
W
X
Y

C
D
E
F
G
H
I
J
K
L
M
N
O
P
Q
R
S
T
U
V
W
X
Y

Salem, Mass., June 19, 1880.

My Dear Sir:

List of Birds of Hudson Highlands.

pt I. Nos 1 to 11 - vol X Nos. 10, 11, 12 pages 166-179 (May 15, 1879).

pt II. Nos 12 to 30 - vol XI. Nos. 1, 2, 3. pages 43-52 (June 12, 1879).

pt III. Nos 31 to 48 - Vol XII. Nos. 7, 8, 9. pages 154-168 (^{Oct} ~~Nov~~ 29, 1879).

pt IV. Nos. 49 to 71 - Vol. XII. Nos. 10, 11, 12. pages 189-204 (April 26, 1880)

The above contains an answer to your question - the dates are those when received at the rooms of the Institute from the printer + distribution followed immediately

Your Mss. has been received

Yrs &c

H. Wheatland."

1874.

1

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	March 14th, 1879.					
	The following is a complete catalogue of all the Birds collected by me up to date, excepting only my original collection of mounted specimens, and unlabeled skins collected for exchanges with the dealers:					
	<u>Highland Falls, N. Y.</u>					
1.	<i>Trochilus colubris</i> , L.	May 30, 1874.	—	—	—	♀ ad.
2.	<i>Chaetura pelagica</i> .	May 20, 1874.	—	—	—	♂ ad.
3.	do	do	—	—	—	♀ ad.
4.	<i>Antrostomus vociferus</i> .	June 1, 1874.	—	—	—	♂ ad.
5.	<i>Tyrannus carolinensis</i> .	June 6, 1874.	—	—	—	♂ ad.
6.	<i>Myiarchus crinitus</i> .	May 24, 1874.	—	—	—	♂ ad.
7.	^{Poughkeepsie, Westchester County, N. Y.} <i>Anothura troglodytes</i> , <i>Myiarchus</i> .	Apr. 27, 1874.	4.	5.81	—	♀ ad.
8.	<i>Pyrranga rubra</i> .	May 30, 1874.	—	—	—	♂ ad.
9.	do	do	—	—	—	♀ ad.
10.	<i>Regulus calendula</i> .	Apr. 21, 1874.	—	—	—	♂ ad.
11.	" <i>satrapa</i> .	Apr. 20, 1874.	—	—	—	♂ ad.
12.	<i>Goniaphea ludoviciana</i> .	May 11, 1874.	—	—	—	♀ ad.
13.	do	May 12, 1874.	—	—	—	♀ ad.
14.	do	May 11, 1874.	—	—	—	♂ ad.
15.	do	do	—	—	—	♂ ad.
16.	do	do	—	—	—	♂ ad.
17.	<i>Icterus baltimore</i> .	May 15, 1874.	—	—	—	♂ ad.
18.	do	May 11, 1874.	—	—	—	♂ ad.
19.	do	May 20, 1874.	—	—	—	♂ ad.
20.	do	May 19, 1874.	—	—	—	♂ ad.
21.	<i>Agelaius phoeniceus</i> .	Apr. 28, 1874.	—	—	—	♂ ad.
22.	do	May 10, 1874.	—	—	—	♂ ad.
23.	<i>Sialia sialis</i> .	Feb. 20, 1874.	6.90	12.00	—	♂ ad.
24.	<i>Colaptes auratus</i> .	June 8, 1874.	—	—	—	♂ ad.
25.	do	May 20, 1874.	—	—	—	♂ ad.
26.	<i>Coccyzus erythrophthalmus</i> .	do	—	—	—	♂ ad.
27.	do	May 24, 1874.	—	—	—	♀ ad.
28.	<i>Melospiza melodia</i> .	June 5, 1874.	—	—	—	♂ ad.
29.	<i>Pyrranga rubra</i> .	May 20, 1874.	—	—	—	♀ ad.

1874.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	Highland Falls, N. Y.					
30.	<i>Picus pubescens.</i>	March 7.				♂ ad.
31.	<i>Tachycineta bicolor.</i>	May 22.				♂ ad.
32.	do	May 23.				♂ ad.
33.	do	do				♀ ad.
34.	<i>Petrochelidon lunifrons.</i>	do				♂ ad.
35.	<i>Hyriundo erythrogastra.</i>	June 10.				♂
36.	<i>Colaptes auratus.</i>	May 1.				♂ ad.
37.	do	May 2.				♂ ad.
38.	do	May 6.				♂ ad.
39.	do	June 21, 1873.				♀ juv.
40.	do	May 7.				♀ ad.
41.	<i>Buteo pennsylvanicus.</i>	May 4.	14.88	37.38		♀ ad.
42.	<i>Pandion haliaetus.</i>	do	25.00			♀ ad.
43.	do	Nov. 25, 1873.	23.40	65.00		♀ ad.
44.	<i>Sialia sialis.</i>	May 9.				♂ ad.
45.	<i>Ectopistes migratorius.</i>	May 2.				♂ ad.
46.	do	do				♀ ad.
47.	<i>Dendroica virens.</i>	May.				♂ ad.
48.	<i>Sialia sialis.</i>	Feb. 20.				♂ ad.
49.	do	March 3.				♂ ad.
50.	<i>Ceryle alcyon.</i>	April 12.				—
51.	<i>Dendroica virens.</i>	May				♀ ad.
52.	<i>Corvus americanus.</i>	May 2.				♂ juv.
53.	<i>Sturnus motacilla.</i>	May 6.				♀ ad.
54.	do	May 7.				♂ ad.
55.	<i>Sturnus aurocapillus.</i>	May 10.				♂ ad.
56.	<i>Parula americana.</i>	May.				—
57.	<i>Vireo olivaceus.</i>	June 3.				♀ ad.
58.	<i>Tyrannus carolinensis.</i>	June 6.				♂ ad.
59.	<i>Cyanurus cristatus.</i>	May 25.				♂ ad.
60.	do	May 27.				♀ ad.
61.	<i>Molothrus ater.</i>	June 1.				♂ ad.
62.	do	May 21.				♀ ad.
63.	<i>Picus pubescens.</i>	April 12.				♀ ad.
64.	<i>Pipilo erythrophthalmus.</i>	June 10.				♂ ad.
65.	<i>Antrostomus vociferus.</i>	May 12.				—

1874.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
66.	<i>Icterus spurius.</i>	May.				♂ ad.
67.	"	"				♂ juv.
68.	<i>Tringoides macularius.</i>	May 9.				♂ ad.
69.	<i>Spizella pusilla.</i>	Apr. 28.				♂ ad.
70.	<i>Chrysomitris tristis.</i>	May.				♂ ad.
71.	"	"				♂ ad.
72.	<i>Telmatochelys palustris.</i>	May 28.				♀ ad.
73.	<i>Dendroica maculosa.</i>	May.				♂ ad.
74.	"	"				♀ ad.
75.	" <i>aestiva.</i>	"				♀ ad.
76.	" <i>striata.</i>	"				♂ ad.
77.	<i>Mniotilta varia.</i>	"				♂ ad.
78.	<i>Geothlypis trichas.</i>	"				♂ ad.
79.	<i>Icteria virens.</i>	June 2.				♂ ad.
80.	"	May 10.				♀ ad.
81.	<i>Dendroica caerulescens.</i>	May.				♂ ad.
82.	"	"				♂ ad.
83.	" <i>coronata.</i>	"				♂ ad.
84.	" <i>castanea.</i>	"				—
85.	" <i>pennsylvanica.</i>	May 8.				♀ ad.
86.	"	May 11.				♂ ad.
87.	" <i>coronata.</i>	May.				♀ ad.
88.	" <i>pennsylvanica.</i>	May 12.				♂ ad.
89.	"	June 6.				♂ ad.
90.	" <i>palmarum.</i>	Apr. 14.				♂ ad.
91.	<i>Helminthophaga peregrina.</i>	May 14.				♂ ad.
92.	" <i>ruficapilla.</i>	May 18.				—
93.	"	May.				—
94.	"	"				—
95.	<i>Empidonax minimus.</i>	"				♂ ad.
96.	<i>Dendroica blackburniae.</i>	"				♂ ad.
97.	<i>Vireo flavifrons.</i>	May 25.				♀ ad.
98.	"	June 6.				♂ ad.
99.	"	May.				♂ ad.
100.	" <i>olivaceus.</i>	"				♂ ad.
101.	<i>Icterus baltimore.</i>	Sept. 1.				juv.
102.	<i>Harporhynchus rufus.</i>	May 28.				♂ ad.

1874.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
103.	Melospiza melodia.	Feb. 26.				♂ ad.
104.	Ampelis cedrorum.	June 17.				♂ ad.
105.	"	June.				♀ ad.
106.	"	Dec. 24, 1873.				♀ ad.
107.	Helminthorus vermicivorus.	July 23, 1875.	5.44	8.75		juv.
108.	Turdus migratorius.	March 10.				♂ ad.
109.	Icterus baltimore.	May 14.				♂ ad.
110.	Sialia sialis.	March 11.				♂ ad.
111.	"	March 1.				♀ ad.
112.	"	Dec. 1, 1873.				♂ ad.
113.	Passerella iliaca.	March 15.				♀ ad.
114.	"	March 20.				♀ ad.
115.	Spizella socialis.	May 3.				♀ ad.
116.	Turdus pallasi.	Apr. 29.				♀ ad.
117.	Cyanospiza cyanea.	May 13.				♂ ad.
118.	"	"				♂ ad.
119.	Icterus spurius.	May 11.				♂ ad.
120.	" baltimore.	"				♀ ad.
121.	Mimus carolinensis.	May.				♂ ad.
122.	Icterus baltimore.	May 14.				♂ ad.
123.	"	"				♂ ad.
124.	Zonotrichia albicollis.	5.19, 1872.				♂ ad.
125.	Sialia sialis.	March 10.				♂ ad.
126.	Nyctiardea grisea, naevia.	June 4.				♂ ad.
127.	Zonotrichia albicollis.	1872.				♂ ad.
128.	Tyrannus carolinensis.	May 12.				♂ ad.
129.	Myiarchus crinitus.	June 17.				♂ ad.
130.	Nyctiardea grisea, naevia.	June 4.				♀ ad.
131.	"	"				♂ ad.
132.	"	"				♀ ad.
133.	"	"				♂ ad.
134.	"	"				♀ ad.
135.	Myiodiodes mitratus.	July 13.				♀ ad.
136.	Vireo olivaceus.	"				♂ ad.
137.	Ceryle alcyon.	July 14.				♂
138.	Dendroica virens.	July 23, 1875.	4.69	7.06		♀ ad.
139.	Nyctiardea grisea, naevia.	June 26.				♂ juv.

1874.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
140.	Nyctiardea grisea, naevia.	June 26.				♂ ad.
141.	Cyanospiza cyanea.	July 24.				♂ ad.
142.	"	July 25.				♂ ad.
143.	Dendroica virens.	July 24.				♂ ad.
144.	Helminthorus vermicivorus.	July 25.				♂ ad.
145.	Harporhynchus rufus.	July 20.				♀ juv.
146.	Pipilo erythrophthalmus.	July 20.				♂ ad.
147.	"	"				juv.
148.	Cyanospiza cyanea.	July 24.				♂
149.	Certhia familiaris.	Apr. 24.				♀ ad.
150.	Petrochelidon lunifrons.	July 28, 1875.	5.75	12.00		juv.
151.	Ampelis cedrorum.	" 29, "				♂ ad.
152.	Myiarchus crinitus.	" 30, "	8.50	12.62		♂ juv.
153.	"	"	8.62	12.62		♀ juv.
154.	Parula americana.	June 10.				♂ ad.
155.	Dendroica striata.	June.				♂ ad.
156.	Numenius longirostris.	Aug. 18.	24.00	40.00		♂ ad.
157.	Sitta canadensis.	Aug. 31.				♀ juv.
158.	"	Sep. 4.	4.75	8.25		♂
159.	"	Aug. 31.				♂
160.	"	Sep. 2.				
161.	Vireo olivaceus.	Aug. 30.	6.00	10.00		♂
162.	Ampelis cedrorum.	Aug. 31.				♂ juv.
163.	Picus pubescens.	Sept. 3.				♂ juv.
164.	Agelaius phoeniceus.	Sep. 10.	8.75	14.50		♂ juv.
165.	Turdus migratorius.	Aug. 29, 1871.				♂ juv.
166.	"	Aug. 29.				♂ juv.
167.	"	Aug. 29.				♂ juv.
168.	Picus villosus.	Sep. 1.				♂ ad.
169.	Accipiter fuscus.	Sep. 4.	11.50			♂ juv.
170.	Porzana carolina.	Sep. 8.	8.87	14.19		♂
171.	"	Sep. 9.	9.37	14.00		♂
172.	Ceryle alcyon.	Aug. 31.				
173.	"	Aug. 31.				
174.	Helminthorus vermicivorus.	Aug. 4.				
175.	Picus pubescens.	Aug. 26.				♂ juv.
176.	Agelaius phoeniceus.	July 3.				♂

1874.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
177.	<i>Colaptes auratus.</i>	Aug. 17.				♂ juv.
178.	<i>Myiarchus crinitus.</i>	July 30.		8.50	12.25	♂ juv.
179.	<i>Turdus migratorius.</i>	Aug. 22.				♀
180.	<i>Turdus macularius.</i>	Aug. 26.				—
181.	<i>Empidonax minimus.</i>	Aug. 12.				—
182.	<i>Dendroica aestiva.</i>	Aug. 26.				♂ ad.
183.	<i>Totanus solitarius.</i>	Aug. 14.				—
184.	<i>Vireo noveboracensis.</i>	Aug. 11.				juv.
185.	<i>Empidonax minimus.</i>	Aug. 10.				—
186.	<i>Setophaga ruticilla.</i>	Aug. 19.				♂ ad.
187.	<i>Dendroica pensylvanica.</i>	Aug. 20.				♀
188.	<i>Trochilus colubris.</i>	Sep. 29.				♀ ad.
189.	<i>Colaptes auratus.</i>	June 10.				♂ ad.
190.	<i>Icterus baltimore.</i>	May 10.				♀ ad.
191.	<i>Ardea virescens.</i>	Aug. 19.				♀
192.	<i>Sayornis fuscus.</i>	May 20.				♀ ad.
193.	<i>Passerella iliaca.</i>	March 24.				♂ ad.
194.	<i>Cyanospiza cyanea.</i>	May 12.				♂ ad.
195.	<i>Haliaeetus leucocephalus.</i>	Sep. 15.	32.25	84.00		♀ juv.
196.	<i>Ardea herodias.</i>	Aug. 17.				♀ juv.
197.	<i>Pandion haliaetus.</i>	Sep. 8.	21.00	61.75		♂ juv.
198.	<i>Ampelis cedrorum.</i>	Sep. 26.				♂ ad.
199.	<i>Totanus solitarius.</i>	Sep. 16.	8.00	16.19		♂
200.	<i>Agelaius phoeniceus.</i>	Sep. 25.	8.00	14.00		♂ juv.
201.	"	Sep. 25.	8.44	13.38		♂ juv.
202.	"	Sep. 25.	6.75	11.87		♀
203.	<i>Vireo olivaceus.</i>	Sep. 20.	6.06	10.00		♂ ad.
204.	<i>Coccyzus erythrophthalmus.</i>	Sep. 23.	11.63	15.50		♂ ad.
205.	" <i>americanus</i>	Sep. 23.	12.50	17.31		♀ ad.
206.	<i>Pyrranga rubra.</i>	Sep. 23.	7.06	11.75		♂ juv.
207.	<i>Dendroica blackburniae.</i>	Sep. 23.	5.70	9.17		♂
208.	"	Sep. 23.	5.63	8.88		♀
209.	<i>Vireo olivaceus.</i>	Sep. 23.	6.06	10.00		♂ ad.
210.	<i>Melospiza melodia.</i>	Sep. 23.	6.12	8.48		♂ ad.
211.	<i>Cyanurus cristatus.</i>	Sep. 25.	11.75	16.50		♀ ad.
212.	<i>Perisoreus carolina.</i>	Sep. 25.	8.12	12.88		♀
213.	"	Sep. 25.	8.42	13.12		♀

1874.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
214.	<i>Dendroica striata.</i>	Sep. 24.				♀
215.	<i>Agelaius phoeniceus.</i>	Sep. 25.		8.00	14.00	♂ juv.
216.	<i>Flachycineta bicolor.</i>	July 28, 1875.		5.75	12.25	♂ juv.
217.	<i>Totanus solitarius.</i>	Sep. 16.		8.00	16.19	♂
218.	<i>Agelaius phoeniceus.</i>	Sep. 26.		7.75	12.50	♀
219.	<i>Ardea herodias.</i>	July 30.				juv.
220.	<i>Anthus motacilla.</i>	July 21.				"
221.	<i>Ceryle alcyon.</i>	Sep. 29.		12.68	21.12	♀
222.	<i>Cyanurus cristatus.</i>	Sep. 30.				♂ ad.
223.	<i>Pipilo erythrophthalmus.</i>	Sep. 30.		8.37	11.12	♂ ad.
224.	<i>Carpodacus purpureus.</i>	Oct. 6.		6.12	9.60	juv.
225.	"	Oct. 8.				♂ ad.
226.	<i>Sitta carolinensis.</i>	Oct. 7.				♀
227.	"	Oct. 7.				♂ ad.
228.	"	Oct. 8.				♀
229.	<i>Turdus swainsoni, aliciae.</i>	Oct. 7.				♀
230.	"	Oct. 8.				♀
231.	<i>Parula americana.</i>	Oct. 8.				—
232.	<i>Dendroica coronata.</i>	Oct. 7.				♂ juv.
233.	<i>Vireo olivaceus.</i>	Oct. 7.				♂ ad.
234.	<i>Corvus americanus.</i>	Oct. 8.		17.75	39.00	♂ ad.
235.	"	Oct. 7.				♀ ad.
236.	"	Oct. 7.				♀ ad.
237.	"	Sep. 29.				♂ ad.
238.	<i>Sphyrapicus varius.</i>	Oct. 8.				♂ juv.
239.	<i>Zonotrichia albicollis.</i>	Oct. 8.		7.00	9.50	♀
240.	<i>Agelaius phoeniceus.</i>	Oct. 15.				♀
241.	<i>Scops asio (red).</i>	Oct. 23.				♂
242.	<i>Sphyrapicus varius.</i>	Oct. 7.		8.62	15.25	♀ juv.
243.	"	Oct. 15.				♀
244.	<i>Aix sponsa.</i>	Oct. 11.				♂ juv.
245.	<i>Melospiza palustris.</i>	Oct. 16.				♀
246.	<i>Ammodramus caudacutus.</i>	Oct. 16.				—
247.	<i>Carpodacus purpureus.</i>	Oct. 11.		6.25	10.13	♂ ad.
248.	<i>Cyanurus cristatus.</i>	Oct. 17.				♂ ad.
249.	<i>Chrysomitris pinus.</i>	Oct. 16.				♀
250.	<i>Cyanurus cristatus.</i>	Oct. 14.				♂ ad.

1874.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
251.	<i>Scops asio</i> (red).	Oct. 24.				♂
252.	<i>Melospiza palustris</i> .	Oct. 26.				♂
253.	<i>Bonasa umbellus</i> .	Oct. 29.				♂ juv.
254.	<i>Junco hyemalis</i> .	Oct. 20.				♂ juv.
255.	<i>Ceothia familiaris</i> .	Oct. 21.				♀
256.	<i>Junco hyemalis</i> .	Oct. 31.				♀
257.	<i>Ceryle alcyon</i> .	Nov. 2.				♀ ad.
258.	" "	Nov. 4.				♀ ad.
259.	<i>Zonotrichia albicollis</i> .	Oct. 15.		7.00	9.50	♀
260.	<i>Colaptes auratus</i> .	Nov. 4.				♂ juv.
261.	<i>Erimaturus rubida</i> .	Nov. 2.				♀
262.	" "	Nov. 2.				♀
263.	<i>Anorthura troglodytes hyemalis</i> .	Nov. 2.				♂
264.	<i>Bucephala albeola</i> .	Nov. 7.				♂ ad.
265.	<i>Mergus serrator</i> .	Nov. 2.				♂ juv.
266.	<i>Larus argentatus</i> .	Nov. 9.				♀ ad.
267.	<i>Philohela minor</i> .	Nov. 3.				♀ ad.
268.	<i>Larus argentatus</i> .	Nov. 9.				♂ juv.
269.	<i>Bucephala albeola</i> .	Nov. 12.				♀
270.	<i>Bucephala albeola</i> .	Nov. 12.				♂ juv.
271.	<i>Mergus cuculatus</i> .	Nov. 15.				♂ juv.
272.	<i>Bonasa umbellus</i> .	Nov. 20.				juv.
273.	" "	Nov. 11.				♂ ad.
274.	<i>Ampelis cedrorum</i> .	Nov. 25.				♀ ad.
275.	<i>Pinicola enucleator</i> .	Dec. 8				♀
276.	" "	" "				♂
277.	" "	" "				♂
278.	" "	" "	9.25	14.00		♂
279.	" "	" "				♂ juv.
280.	" "	" "	8.69	13.00		♀
281.	" "	" "	8.75	13.56		♀
282.	" "	" "				♀
283.	" "	" "	8.56	13.19		♀
284.	" "	" "				♀
285.	" "	" "				♀
286.	" "	" "				♀
287.	" "	" "				♀

1874-75.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
288.	<i>Regulus satrapa</i> .	Dec. 29.				♂ ad.
289.	<i>Turdus pallasi</i> .	" 9.				♀ ad.
290.	" "	" 28.				♂ ad.
291.	<i>Spizella monticola</i> .	" "				♂ ad.
292.	<i>Chrysomitris tristis</i> .	" 27.				♂
293.	<i>Pinicola enucleator</i> .	" 29.		8.56	13.19	♀
294.	" "	Jan. 5.		9.00	12.63	♀
295.	" "	" "		9.06	13.75	♀
296.	" "	" "				♀
297.	" "	" "				♀
298.	" "	" "		9.00	14.00	♂ juv.
299.	" "	" "				♂ juv.
300.	<i>Egiothus linaria</i> .	Dec. 30.				♂ juv.
301.	" "	" "				♂ juv.
302.	" "	" "				♂ juv.
303.	" "	" "				♀ ad.
304.	<i>Sialia sialis</i> .	Jan. Feb. '74.				♂ ad.
305.	" "	Jan. or Feb. '74.				♂ ad.
306.	" "	do				♂ ad.
307.	" "	do				♂ ad.
308.	" "	do				♂ ad.
309.	" "	do				♂ ad.
310.	" "	do				♂ ad.
311.	" "	do				♂ ad.
312.	" "	do				♂ ad.
313.	<i>Pinicola enucleator</i> .	Dec. 20.				♀
314.	" "	do				♀
315.	" "	do				♀
316.	" "	do				♀
317.	<i>Picus pubescens</i> .	do				♂ ad.
318.	<i>Sialia sialis</i> .	do				♂ ad.
319.	<i>Egiothus linaria</i> .	do				♀ ad.
320.	<i>Pinicola enucleator</i> .	Jan. 1875.				♀
321.	" "	do				♀
322.	" "	do				♀
323.	" "	do				♀
324.	" "	do				♀

1875.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
325.	<i>Pinicola enucleator.</i>	1875. Jan. 5.		—	—	♀
326.	"	"	do	—	—	♀
327.	"	"	do	—	—	♀
328.	"	"	do	—	—	♀
327.	"	"	do	—	—	♀
328.	"	"	do	—	—	♀
329.	"	"	do	—	—	♂ juv.
330.	"	"	do	—	—	♂ juv.
331.	"	"	do	—	—	♀
332.	"	"	do	—	—	♀
333.	<i>Loxia curvirostra, americana.</i>	Jan. 11.		—	—	♂ ad.
334.	"	"	do	—	—	♂ ad.
335.	"	"	do	—	—	♀ ad.
336.	<i>Parus atricapilla.</i>	Jan. 8.		—	—	♂ ad.
337.	<i>Picus pubescens.</i>	" 7.		—	—	♀ ad.
338.	<i>Spizella monticola.</i>	" "	6.44	9.75	—	—
339.	<i>Loxia curvirostra, americana.</i>	" 14.		—	—	♂ ad.
340.	"	"	"	—	—	♂ ad.
341.	"	"	"	—	—	♂ juv.
342.	"	"	"	—	—	♀ ad.
343.	" <i>leucoptera.</i>	"	"	—	—	♂ ad.
344.	"	"	"	—	—	♂ ad.
345.	"	"	"	—	—	♀ ad.
346.	"	"	"	—	—	♀ ad.
347.	"	"	"	—	—	♀ ad.
348.	<i>Picus villosus.</i>	" 16.		—	—	♂ ad.
349.	<i>Melospiza melodia.</i>	"	"	—	—	♂ ad.
350.	<i>Picus pubescens.</i>	" 14.		—	—	♂ ad.
351.	<i>Loxia curvirostra, americana.</i>	" 16.		—	—	♂
352.	<i>Pinicola enucleator.</i>	" 19.	8.94	13.69	—	♀
353.	"	"	"	9.06	14.00	♀
354.	"	"	"	—	—	♀
355.	"	"	"	—	—	♀
356.	<i>Loxia curvirostra, americana</i>	" 16.		—	—	♀
357.	"	" 24.		—	—	♂
358.	<i>Loxia leucoptera.</i>	" 25.		—	—	♂ ad.
359.	"	" 22.		—	—	♂ ad.

1875.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
360.	<i>Loxia leucoptera.</i>	Jan 14.		—	—	♀ ad.
361.	"	"	"	—	—	♀ ad.
362.	"	"	"	—	—	♀ ad.
363.	<i>Melospiza melodia.</i>	" 23.		—	—	—
364.	<i>Egiotthus linaria.</i>	" 20		—	—	♂
365.	"	"	"	—	—	♀
366.	<i>Regulus satrapa.</i>	" 14.		—	—	♂ ad.
367.	"	"	"	—	—	♀ ad.
368.	<i>Certhia familiaris.</i>	"	"	—	—	♀ ad.
369.	"	"	"	—	—	♂ ad.
370.	"	"	"	—	—	♂ ad.
371.	<i>Picus pubescens.</i>	" 20.		—	—	♂ ad.
372.	<i>Spizella monticola.</i>	"	"	—	—	♂ ad.
373.	<i>Sitta carolinensis.</i>	" 23.		—	—	♂ ad.
374.	<i>Anorthura troglodytes, pygmalis.</i>	" 20.		—	—	♂ ad.
375.	<i>Egiotthus linaria.</i>	" 27.		—	—	♂ ad.
376.	"	"	"	—	—	♂ ad.
377.	"	"	"	—	—	♀ ad.
378.	"	"	"	—	—	♂ ad.
379.	"	"	"	—	—	♂ ad.
380.	"	"	"	—	—	♀ ad.
381.	<i>Chrysomitris pinus.</i>	"	"	—	—	♀ ad.
382.	<i>Egiotthus linaria.</i>	"	"	—	—	♂ juv.
383.	<i>Spizella monticola.</i>	"	"	6.06	9.50	♀ ad.
384.	<i>Melospiza melodia.</i>	"	"	—	—	♂ ad.
385.	<i>Egiotthus linaria.</i>	"	"	—	—	♀
386.	"	"	"	—	—	♀
387.	"	"	"	—	—	♀
388.	"	"	"	—	—	♂ juv.
389.	"	"	"	—	—	♀
390.	"	"	"	—	—	♀
391.	"	"	"	—	—	♀
392.	"	"	"	—	—	♀
393.	"	"	"	—	—	♂ juv.
394.	"	"	"	—	—	♀
395.	"	"	"	—	—	♂ juv.
396.	"	"	"	—	—	♂ juv.

1875.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
397.	<i>Egiothus linaria.</i>	Jan. 27.				♂ juv.
398.	"	"	"			♀
399.	"	"	"			♀
400.	<i>Loxia curvirostra, americana</i>	"	"			♀ ad.
401.	<i>Anorthura troglodytes, hyemalis.</i>	"	"			♀ ad.
402.	<i>Carpodacus purpureus.</i>	" 29	"			♂ ad.
403.	"	"	"			♂ ad.
404.	<i>Chrysomitris pinus</i>	" 27.	"			—
405.	"	" 27.	"			♀ ad.
406.	"	"	"			♂ ad.
407.	"	"	"			♂ ad.
408.	"	"	"			♀ ad.
409.	"	"	"			♀ ad.
410.	"	"	"			—
411.	<i>Bonasa umbellus.</i>	" 24.	"			♂ ad.
412.	<i>Pinicola enucleator</i>	" 29.	"	9.06	13.88	♀
413.	"	"	"	9.06	13.88	♀
414.	<i>Loxia curvirostra, americana.</i>	Feb. 1.	"	8.66	13.88	♂ ad.
415.	<i>Sitta carolinensis</i>	"	"	8.86	13.88	♀ ad.
416.	<i>Chrysomitris pinus.</i>	"	"	8.13	13.73	♀ ad.
417.	<i>Egiothus linaria.</i>	"	"			♂ ad.
418.	<i>Junco hyemalis.</i>	"	"			♂ ad.
419.	<i>Melospiza melodia.</i>	"	"			♂ ad.
420.	<i>Spizella monticola.</i>	"	"	86.31	9.44	♀ ad.
421.	<i>Picus pubescens.</i>	"	"			♀ ad.
422.	<i>Loxia curvirostra, americana.</i>	Jan. 29.	"			♀ ad.
423.	"	" 31.	"			♀ ad.
424.	<i>Junco hyemalis.</i>	Feb. 1.	"			♂ ad.
425.	<i>Spizella monticola.</i>	"	"			♂ ad.
426.	<i>Egiothus linaria.</i>	" 19.	"			♂ ad.
427.	"	"	"			♂ ad.
428.	"	"	"			♂ ad.
429.	"	" 1.	"			♂ ad.
430.	"	" 1.	"			♀ ad.
431.	"	" 2.	"			♀ ad.
432.	"	" 20.	"			♀ ad.
433.	<i>Sitta canadensis.</i>	" 19.	"			♀ ad.

1875.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
434.	<i>Sitta canadensis.</i>	Feb. 19.				—
435.	<i>Melospiza melodia.</i>	" 1.	"			♀ ad.
436.	<i>Spizella monticola.</i>	" 1.	"			♀ ad.
437.	<i>Picus villosus.</i>	" 17.	"	9.50	15.81	♂ ad.
438.	<i>Spizella monticola.</i>	" 16.	"			♂ ad.
439.	<i>Plectrophanes nivalis.</i>	" 19.	"			♂ ad.
440.	"	"	"			♀ ad.
441.	"	"	"			♀ ad.
442.	"	"	"			♀ ad.
443.	<i>Spizella monticola.</i>	" 16.	"			♀ ad.
444.	<i>Melospiza melodia.</i>	"	"			♂ ad.
445.	<i>Regulus satrapa.</i>	" 22.	"			♂ ad.
446.	<i>Cecropia familiaris.</i>	" 21.	"			♂ ad.
447.	"	" 26.	"			♀ ad.
448.	"	" 24.	"			♀ ad.
449.	<i>Picus pubescens.</i>	" 26.	"			♀ ad.
450.	<i>Sitta carolinensis.</i>	"	"			♀ ad.
451.	<i>Cyanurus cristatus.</i>	" 28.	"	—	—	♂ ad.
452.	<i>Egiothus linaria.</i>	" 26.	"			♂ ad.
453.	"	" 28.	"			♀
454.	<i>Contopus virens.</i>	(Spring of 1874).				—
455.	<i>Cupidonia cupido</i> (Muller)	Feb. 10.	"			♂ ad.
456.	<i>Junco hyemalis.</i>	" 28.	"			♂ ad.
457.	<i>Spizella monticola.</i>	"	"			♂ ad.
458.	"	"	"	6.25	9.13	♀ ad.
459.	<i>Bonasa umbellus.</i>	March 2.	"			♂
460.	<i>Ampelis cedrorum</i>	" 1.	"			♂ ad.
461.	"	"	"			♀ ad.
462.	<i>Spizella monticola.</i>	" 2.	"			♂ ad.
463.	"	" 2.	"			♀ ad.
464.	<i>Parus atricapillus.</i>	" 2.	"			♂ ad.
465.	<i>Sitta carolinensis.</i>	" 1.	"			♂ ad.
466.	<i>Picus pubescens.</i>	" 1.	"			♂ ad.
467.	<i>Turdus migratorius</i>	" 3.	"			♂ ad.
468.	"	" 4.	"			♀ ad.
469.	<i>Pinicola enucleator.</i>	" 4.	"			♀.
470.	<i>Carpodacus purpureus.</i>	" 4.	"			♀ ad.

1875.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
471.	<i>Carpodacus purpureus.</i>	March 4.				♂ ad.
472.	"	"	"	5.		♂ ad.
473.	"	"	"	5.		♂ ad.
474.	<i>Passer domesticus.</i>	"	"	3.		♂ ad.
475.	<i>Ampelis cedrorum.</i>	"	"			♂ ad.
476.	"	"	"	7.00	11.75	♂ ad.
477.	<i>Spizella monticola.</i>	"	"	6.38	9.63	♂ ad.
478.	<i>Picus pubescens.</i>	"	"			♂ ad.
479.	<i>Egiotthus linaria</i>	"	5			♂ ad.
480.	"	"	"			♂ ad.
481.	<i>Plectrophanes nivalis.</i>	"	"			♀
482.	"	"	"			—
483.	"	"	"			—
484.	<i>Bonasa umbellus.</i>	"	10.			♀ ad.
485.	"	"	"	5.		♂ ad.
486.	<i>Turdus migratorius.</i>	"	9.			♂ ad.
487.	<i>Sialia sialis.</i>	"	"			—
488.	<i>Ampelis cedrorum.</i>	"	5.			♂ ad.
489.	"	"	"			♀ ad.
490.	<i>Toxia curvirostra, americana.</i>	"	10.			♂ ad.
491.	"	"	"			♂ ad.
492.	<i>Passer domesticus.</i>	"	9.			♂ ad.
493.	"	"	"			♀ ad.
494.	<i>Plectrophanes nivalis</i>	"	"			♀ ad.
495.	<i>Haliaetus leucocephalus.</i>	"	12.			♂ ad.
496.	<i>Plectrophanes nivalis.</i>	"	9.			♂
497.	<i>Picus villosus.</i>	"	15.			♀ ad.
498.	<i>Egiotthus linaria.</i>	"	"			♂ ad.
499.	<i>Junco hyemalis.</i>	"	14.			♂ ad.
500.	<i>Plectrophanes nivalis.</i>	"	5.			♂
501.	<i>Pinicola enuleator.</i>	"	14.			♀
502.	"	"	"			♀
503.	"	"	"	17.		♀
504.	"	"	"	"		♀
505.	"	"	"	"		♀
506.	"	"	"	"		♀
507.	"	"	"	"		♂ juv.

1875.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
508.	<i>Agelaius phoeniceus.</i>	March 18.		9.31	15.00	♂ ad.
509.	<i>Holcophagus ferrugineus.</i>	" 17.		9.69	15.25	♂ ad.
510.	<i>Melospiza melodia.</i>	"	"			♀ ad.
511.	<i>Passer domesticus.</i>	" 15.		6.38	9.88	♂ ad.
512.	<i>Melospiza melodia.</i>	"	"			♂ ad.
513.	<i>Sialia sialis.</i>	" 16.				♂ ad.
514.	"	"	"			♂ ad.
515.	<i>Passer domesticus.</i>	"	"	6.13	9.13	♂ juv.
516.	<i>Sialia sialis.</i>	" 17.				♂ ad.
517.	"	"	"			♀ ad.
518.	<i>Passerella iliaca.</i>	"	"	7.44	11.25	♂ ad.
519.	"	"	"			♀ ad.
520.	<i>Melospiza melodia.</i>	"	"			♀ ad.
521.	<i>Turdus migratorius.</i>	"	"			♀ ad.
522.	<i>Ampelis cedrorum.</i>	" 30.		7.44	11.88	♀ ad.
523.	"	"	"			♂ ad.
524.	"	"	"			♂ ad.
525.	"	"	"			♂ ad.
526.	<i>Sitta canadensis.</i>	March 24.				♀ ad.
527.	"	"	"			♂ ad.
528.	<i>Egiotthus linaria.</i>	"	"			♂ ad.
529.	"	"	"			♂ ad.
530.	"	"	"			♂ ad.
531.	<i>Carpodacus purpureus.</i>	" 30.				♀ ad.
532.	<i>Sialia sialis.</i>	"	"			♂ ad.
533.	<i>Melospiza melodia.</i>	"	"			♀ ad.
534.	<i>Sayornis fuscus.</i>	" 29.		7.00	10.81	♀ ad.
535.	<i>Egiotthus linaria.</i>	" 24.				♀ ad.
536.	"	"	"			♀ ad.
537.	<i>Zenaidura carolinensis.</i>	Apr. 2.				♂ ad.
538.	<i>Colaptes auratus.</i>	"	"			♂ ad.
539.	<i>Carpodacus purpureus.</i>	" 1.				♂ ad.
540.	"	"	"			♀ ad.
541.	<i>Passerella iliaca.</i>	"	"			♂ ad.
542.	<i>Sialia sialis.</i>	"	"			♂ ad.
543.	<i>Ampelis cedrorum.</i>	"	"			♂ ad.
544.	<i>Buteo borealis.</i>	"	"			♂ ad.

1875.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
545.	<i>Ampelis cedrorum.</i>	Apr. 11.				♂ ad.
546.	<i>Turdus migratorius.</i>	" 15.				♂ ad.
547.	"	" "				♀ ad.
548.	<i>Passerella iliaca.</i>	" "				♀ ad.
549.	<i>Scolecophagus ferrugineus.</i>	" "		9.56	14.38	♀ ad.
550.	<i>Anas obscura.</i>	" "				♂ ad.
551.	<i>Branta canadensis.</i>	" 9.				♂ ad.
552.	<i>Larus argentatus.</i>	" 4.				♂ ad.
553.	<i>Podiceps cornutus.</i>	" 15.				♀ ad.
554.	<i>Sialia sialis.</i>	" "				♂ ad.
555.	<i>Turdus pallasi.</i>	" 13.				♂ ad.
556.	"	" "				♂ ad.
557.	<i>Passerculus savanna.</i>	" 19.				♂ ad.
558.	<i>Spizella monticola.</i>	" 15.				ad.
559.	<i>Melospiza palustris.</i>	" 19.				♂ ad.
560.	"	" "				ad.
561.	"	" "				ad.
562.	<i>Tringoides macularius.</i>	" 26.		7.81	13.81	♀ ad.
563.	<i>Junco hyemalis.</i>	" "		6.25	9.75	♂ ad.
564.	<i>Poocetes gramineus.</i>	" 27.		6.00	10.12	♂ ad.
565.	"	" "		5.94	9.81	♀ ad.
566.	<i>Regulus satrapa.</i>	" "		4.00	6.94	♂ ad.
567.	"	" "				
568.	<i>Poocetes gramineus.</i>	" "		6.31	10.88	♂ ad.
569.	"	" "		6.19	10.31	♀ ad.
570.	"	" "		6.38	10.94	♂ ad.
571.	"	" "		6.25	10.44	♀ ad.
572.	"	" "		6.25	10.63	♂ ad.
573.	<i>Podiceps cornutus.</i>	" 29.		13.62	24.00	♀
574.	<i>Simus motacilla.</i>	" 30.		6.12	10.56	♂ ad.
575.	<i>Mniotilta varia.</i>	" "		5.25	8.31	♂ ad.
576.	<i>Dendroica coronata.</i>	" "		5.69	9.06	♂ ad.
577.	<i>Regulus calendula.</i>	" "		4.37	7.12	♂ ad.
578.	<i>Molothrus ater.</i>	" 29.		8.06	13.75	♂ ad.
579.	<i>Pipilo erythrophthalmus.</i>	" 30.		8.50	11.12	♂ ad.
580.	<i>Poocetes gramineus.</i>	" 27.		6.00	10.25	♀ ad.
581.	<i>Regulus calendula.</i>	" 30.		4.50	7.25	♂ ad.

1875.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
582.	<i>Scolecophagus ferrugineus.</i>	Apr. 30.		9.00	13.50	♀ ad.
583.	<i>Colaptes auratus.</i>	" "		12.88	20.50	♂ ad.
584.	<i>Spizella socialis.</i>	" "		5.44	8.75	♂ ad.
585.	"	" "		5.19	8.44	♀ ad.
586.	<i>Hirundo horreorum.</i>	May 3.		6.88	13.00	♂ ad.
587.	<i>Dendroica coronata.</i>	" "		5.56	9.19	♂ ad.
588.	"	" "		5.50	8.75	♂ ad.
589.	" <i>pinus.</i>	" "		5.25	8.56	♀ ad.
590.	" <i>palmarum.</i>	" "		5.25	8.12	♂ ad.
591.	<i>Tachycineta bicolor.</i>	" "		5.88	13.00	♂ ad.
592.	<i>Hirundo horreorum.</i>	" 4.		7.44	13.25	♂ ad.
593.	<i>Chaetura pelagica.</i>	" 3.		5.25	12.50	♂ ad.
594.	<i>Petrochelidon lunifrons.</i>	" "		5.75	12.25	♂ ad.
595.	<i>Tachycineta bicolor.</i>	" "		5.88	12.88	♂ ad.
596.	"	" "		5.88	12.63	♂ ad.
597.	<i>Dendroica coronata.</i>	" "		5.56	8.81	♂ ad.
598.	<i>Tachycineta bicolor.</i>	" 4.		6.12	12.63	♂ ad.
599.	"	" "		5.50	12.31	♀ ad.
600.	<i>Hirundo horreorum.</i>	" "		7.00	12.75	♂ ad.
601.	"	" "		6.50	12.62	♀ ad.
602.	<i>Zonotrichia albicollis.</i>	" "		6.50	9.37	♂ ad.
603.	"	" "		7.00	9.75	♂ ad.
604.	"	" "		6.50	9.44	♂ ad.
605.	"	" "		6.69	9.25	♂ ad.
606.	<i>Tachycineta bicolor.</i>	" 3.		5.56	12.19	♂ ad.
607.	<i>Hirundo horreorum.</i>	" 4.		—	—	♂ ad.
608.	"	" "		—	—	♂ ad.
609.	<i>Tachycineta bicolor.</i>	" "		—	—	♂ ad.
610.	"	" "		—	—	♂ ad.
611.	<i>Pipilo erythrophthalmus.</i>	" 5.		8.19	10.56	♀ ad.
612.	<i>Dendroica palmarum.</i>	" 5.		—	—	♀ ad.
613.	<i>Passerculus savanna.</i>	" "		5.62	9.37	♂ ad.
614.	"	" "		5.37	8.69	♀ ad.
615.	"	" "		5.37	8.56	♀ ad.
616.	<i>Troglodytes aëdon.</i>	" 6.		4.69	6.25	♂ ad.
617.	<i>Spizella socialis (albino).</i>	" 8.		5.50	8.56	♂ ad.
618.	<i>Junco hyemalis.</i>	" "		—	—	♂ ad.

1875.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
619.	<i>Spizella pusilla.</i>	May 8.		5.75	8.19	♂ ad.
620.	<i>Mniotilta varia.</i>	" "		5.50	8.88	♂ ad.
621.	<i>Sturnus motacilla</i>	" "		6.50	10.62	♀ ad.
622.	<i>Dendroica palmarum.</i>	" "		—	—	♂ ad.
623.	<i>Hirundo horreorum.</i>	" "		—	—	♂ ad.
624.	<i>Tyrannus carolinensis.</i>	" 10.		8.19	14.06	♀ ad.
625.	<i>Sturnus auroparillus.</i>	" "		—	—	♂ ad.
626.	<i>Geothlypis trichas.</i>	" "		—	—	♂ ad.
627.	" "	" "		5.12	7.19	♂ ad.
628.	<i>Tyrannus carolinensis.</i>	" "		—	—	♂ ad.
629.	<i>Minus carolinensis.</i>	" "		—	—	♀ ad.
630.	<i>Icterus baltimore.</i>	" 11.		—	—	♀ ad.
631.	" "	" "		—	—	♂ juv.
632.	" "	" "		—	—	♂ ad.
633.	<i>Pipilo erythrophthalmus.</i>	" "		—	—	♂ ad.
634.	<i>Chaetura pelagica.</i>	" "		—	—	♂ ad.
635.	" "	" "		—	—	♀ ad.
636.	" "	" "		—	—	♂ ad.
637.	<i>Dendroica maculosa.</i>	" "		5.06	7.75	♂ ad.
638.	" <i>caerulea.</i>	" "		5.12	7.81	♂ ad.
639.	<i>Helminthophaga ruficapilla.</i>	" "		4.31	7.81	♀ ad.
640.	<i>Myiodytes mitratus.</i>	" "		5.25	7.81	♀ ad.
641.	<i>Actostomus vociferus.</i>	" "		9.56	18.56	♀ ad.
642.	<i>Myiodytes mitratus</i>	" 12.		5.44	7.75	♂ ad.
643.	<i>Turdus swainsoni.</i>	" 11.		6.31	12.00	♀ ad.
644.	<i>Tringoides macularius</i>	" 10.		8.00	13.81	♂ ad.
645.	<i>Myiodytes mitratus.</i>	" 12.		5.44	7.75	♂ ad.
646.	<i>Dendroica caerulea.</i>	" 11.		5.12	7.81	♂ ad.
647.	" <i>virens.</i>	" "		5.00	7.56	♂ ad.
648.	" <i>pennsylvanica.</i>	" 12.		5.00	7.88	♂ ad.
649.	<i>Turdus swainsoni.</i>	" "		—	—	♂ ad.
650.	" <i>mustelinus.</i>	" 10.		8.19	13.19	♂ ad.
651.	<i>Dendroica maculosa.</i>	" 12.		4.94	7.50	♂ ad.
652.	<i>Empidonax minimus.</i>	" 11.		5.25	8.25	♂ ad.
653.	" "	" "		4.94	7.50	♀ ad.
654.	<i>Setophaga ruticilla.</i>	" "		5.19	7.88	♂ ad.
655.	<i>Parula americana.</i>	" 12.		4.37	7.37	♂ ad.

1875.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
656.	<i>Myiodytes pusillus.</i>	May 12.		5.00	7.06	♂ ad.
657.	" <i>mitratus.</i>	" "		—	—	♀ ad.
658.	<i>Turdus swainsoni.</i>	" 13.		—	—	♂ ad.
659.	" "	" "		—	—	♀ ad.
660.	" "	" "		—	—	♂ ad.
661.	" "	" "		—	—	♀ ad.
662.	<i>Icterus baltimore.</i>	" 12.		—	—	♂ ad.
663.	" "	" 13.		—	—	♂ ad.
664.	<i>Pyrranga rubra.</i>	" 14.		7.37	12.12	♂ ad.
665.	<i>Ceryle alcyon.</i>	" 11.		13.00	22.50	♂ ad.
666.	<i>Mniotilta varia.</i>	" 14.		4.88	8.44	♂ ad.
667.	<i>Helminthophaga celata.</i>	" 13.		4.69	7.25	♂ ad.
668.	" <i>ruficapilla.</i>	" "		—	—	♂ ad.
669.	" "	" "		—	—	♂ ad.
670.	<i>Parula americana.</i>	" "		4.69	7.00	♀ ad.
671.	<i>Pyrranga rubra.</i>	" 14.		7.12	12.31	♂ ad.
672.	<i>Mniotilta varia.</i>	" 15.		—	—	—
673.	<i>Pyrranga rubra.</i>	" "		—	—	♂ ad.
674.	<i>Dendroica caerulea.</i>	" 17.		5.00	8.06	♂ ad.
675.	<i>Petrochelidon lunifrons.</i>	" "		6.25	10.94	♂ ad.
676.	<i>Dendroica maculosa.</i>	" "		5.00	7.50	♀ ad.
677.	" <i>caerulea.</i>	" "		5.25	8.12	♂ ad.
678.	<i>Trochilus colubris.</i>	" "		3.69	4.12	♂ ad.
679.	<i>Vireo novboracensis.</i>	" 16.		5.12	7.81	♀ ad.
680.	<i>Parula americana.</i>	" 17.		—	—	♂ ad.
681.	<i>Chordeiles virginianus.</i>	" "		10.00	24.00	♀ ad.
682.	<i>Icteria virens.</i>	" "		—	—	♀ ad.
683.	<i>Turdus migratorius.</i>	" "		—	—	♀ ad.
684.	<i>Goniophes ludoviciana.</i>	" 20.		8.25	12.81	♀ ad.
685.	" "	" "		—	—	♂ ad.
686.	<i>Dendroica caerulea.</i>	" "		4.94	7.37	♀ ad.
687.	<i>Coccyzus erythrophthalmus.</i>	" 12.00		19.12	—	♂ ad.
688.	<i>Setophaga ruticilla.</i>	" 17.		—	—	♂ ad.
689.	<i>Helminthophaga ruficapilla.</i>	" 19.		—	—	♂ ad.
690.	<i>Dendroica caerulea.</i>	" 20.		—	—	♂ ad.
691.	" <i>virens.</i>	" "		—	—	♀ ad.
692.	" <i>pennsylvanica.</i>	" "		—	—	♂ ad.

1875.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
693.	<i>Carpodacus purpureus</i> .	May 20.		—	—	♂ ad.
694.	<i>Pyrranga rubra</i> .	" "		—	—	♂ ad.
695.	<i>Parula americana</i> .	" "		—	—	♂ ad.
696.	<i>Vireo gilvus</i> .	" "		—	—	♂ ad.
697.	<i>Trochilus colubris</i> .	" "		—	—	♂ ad.
698.	<i>Dendroica aestiva</i> .	" "		—	—	♂ ad.
699.	" "	" "	" "	5.06	7.62	♀ ad.
700.	" <i>caerulea</i> .	" 21.		—	—	♂ ad.
701.	<i>Myiodytes mitratus</i> .	" "		—	—	♂ ad.
702.	" "	" 20.		—	—	♀ ad.
703.	<i>Helminthophaga peregrina</i> .	" "	" "	5.19	8.12	♂ ad.
704.	<i>Agelaius phoeniceus</i> .	" 21.		—	—	♂ ad.
705.	<i>Dolichonyx oryzivorus</i> .	" "	" "	7.50	12.00	♂ ad.
706.	<i>Zonotrichia albicollis</i> .	" "		—	—	♂ ad.
707.	<i>Dendroica caerulea</i> .	" 22.		—	—	♂ ad.
708.	" "	" 21.		—	—	♂ ad.
709.	<i>Vireo olivaceus</i> .	" "		—	—	♀ ad.
710.	<i>Parula americana</i> .	" "		—	—	♀ ad.
711.	" "	" "		—	—	♀ ad.
712.	<i>Scops asio</i> (nestling).	" 31.		6.00	11.12	♀ juv.
713.	<i>Dolichonyx oryzivorus</i> .	" "	" "	7.50	11.94	♂ ad.
714.	<i>Empidonax flaviventris</i> .	" "	" "	5.69	8.75	♂ ad.
715.	<i>Petrochelidon lunifrons</i> .	" "		—	—	♂ ad.
716.	" "	" "		—	—	♀ ad.
717.	<i>Telmatochelys palustris</i> .	" "	" "	5.12	6.50	♀ ad.
718.	<i>Scops asio</i> (parent).	June 1.		9.37	24.50	♀ ad.
719.	" "	" 2.		9.00	22.12	♂ ad.
720.	<i>Coereba erythrocephala</i> .	May 31.		—	—	♀ ad.
721.	<i>Tyrannus carolinensis</i> .	June 6.		—	—	♀ ad.
722.	<i>Icterus palmarum</i> .	" "		—	—	♂ ad.
723.	<i>Vireo olivaceus</i> .	July 12.		6.44	10.50	♂ ad.
724.	<i>Melospiza melodia</i> .	" "		—	—	♂ juv.
725.	<i>Junco motacilla</i> .	" "	" "	6.25	10.00	♂ juv.
726.	<i>Picus pubescens</i> .	" "	" "	6.62	11.75	♂ juv.
727.	<i>Setophaga ruticilla</i> .	" "		—	—	♂ juv.?
728.	<i>Myiodytes mitratus</i> .	" "	" "	5.56	8.00	♀ ad.
729.	<i>Contopus virens</i> .	" "	" "	6.56	10.50	♂ ad.

1875.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
730.	<i>Turdus migratorius</i> .	July 13.		—	—	juv.
731.	<i>Icterus spurius</i> .	" "	" "	6.50	9.75	♀ juv.
732.	<i>Telmatochelys palustris</i> .	" "	" "	5.12	6.62	♂ ad.
733.	<i>Myiarchus crinitus</i> .	" 30.		8.69	12.50	♂ juv.
734.	" "	" "	" "	8.88	13.12	♂ juv.
735.	<i>Ceryle alcyon</i> .	" "	" "	12.81	21.44	♂ juv.
736.	<i>Mniotilta varia</i> .	" "	" "	5.25	8.50	♂ juv.
737.	<i>Contopus borealis</i> .	Aug. 27.		7.31	12.50	♀
738.	<i>Picus pubescens</i> .	Feb. 1.		—	—	♀ ad.
739.	<i>Ampelis cedrorum</i> .	Aug. 16.		7.06	11.50	♂ juv.
740.	" "	" "	" "	—	—	♀ juv.
741.	<i>Dendroica pinus</i> .	" 31.		5.56	9.06	♀
742.	<i>Vireo solitarius</i> .	May, 1872.		—	—	—
743.	<i>Pyrranga rubra</i> .	Sept. 1, 1874.		—	—	♂ juv.
744.	<i>Sialia sialis</i> .	Sept. 1.		7.00	12.62	♂ juv.
745.	<i>Pyrranga carolina</i> .	" 4.		8.88	14.00	♂ juv.
746.	<i>Picus pubescens</i> .	" 7.		6.75	12.00	♂ juv.
747.	<i>Dendroica virens</i> .	" "	" "	5.06	7.75	♂ juv.
748.	<i>Setophaga ruticilla</i> .	" "	" "	5.06	7.50	♀ ad.
749.	<i>Contopus borealis</i> .	" 18.		7.50	12.88	♀
750.	" "	" "	" "	7.56	12.88	♀
751.	<i>Pyrranga carolina</i> .	" "	" "	9.00	14.69	♀
752.	<i>Dendroica virens</i> .	" "	" "	5.06	7.62	♂ juv.
753.	" <i>maculosa</i> .	" "	" "	5.12	7.50	♂
754.	" <i>striata</i> .	" 19.		5.50	8.06	—
755.	" "	" "	" "	5.56	7.69	♀
756.	" <i>castanea</i> .	" "	" "	5.44	7.44	♂ juv.
757.	<i>Myiarchus crinitus</i> .	" "	" "	8.75	—	♂ juv.
758.	<i>Aldea virescens</i> .	" "		—	—	♀
759.	<i>Dendroica (Perisoreglossa) tigrina</i> .	20.		5.00	8.00	♂ juv.
760.	" <i>blackburniae</i> .	" 20.		4.56	7.50	♂ juv.
761.	<i>Junco aurocapillus</i> .	" "	" "	6.00	9.19	—
762.	<i>Dendroica striata</i> .	" "	" "	5.31	9.12	—
763.	<i>Helminthophaga ruficapilla</i> .	" "	" "	4.44	7.00	♀ juv.
764.	<i>Ampelis cedrorum</i> .	" "	" "	7.31	11.56	♂ ad.
765.	<i>Sialia sialis</i> .	" "	" "	—	—	♂ juv.
766.	<i>Colymbus torquatus</i> .	" 21.		30.50	54.00	♀ juv.

1875.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
767.	<i>Ectopistes migratorius.</i>	Sept. 21.		15.31	24.00	♂ juv.
768.	"	"	"	16.62	24.25	♂
769.	<i>Turdus swainsoni.</i>	" 23.		6.50	11.12	—
770.	<i>Pyrranga rubra.</i>	"	"	7.00	11.12	♂ ad.
771.	<i>Vireo solitarius.</i>	"	"	5.56	9.31	♀?
772.	<i>Parula americana.</i>	"	"	4.62	7.50	—
773.	<i>Ardea virescens.</i>	"	"	—	—	♀
774.	<i>Spizella socialis.</i>	"	"	—	—	—
775.	<i>Ardea virescens.</i>	July 21.		17.75	26.75	♀
776.	<i>Dendroica striata.</i>	Sept. 28.		5.37	8.75	♂
777.	"	"	"	5.50	8.75	—
778.	"	"	"	5.47	8.69	♀
779.	" <i>coronata.</i>	"	"	5.50	9.12	—
780.	" <i>virens.</i>	"	"	4.88	7.56	♀
781.	<i>Empidonax flaviventris.</i>	"	"	5.50	8.75	♂
782.	<i>Podilymbus podiceps.</i>	" 30.		12.50	22.00	♀
783.	<i>Ceryle alcyon.</i>	"	"	13.00	21.81	♀
784.	<i>Dendroica virens.</i>	"	"	5.06	8.00	—
785.	<i>Antrostomus vociferus.</i>	"	"	9.50	19.00	♀
786.	<i>Turdus aliciae.</i>	Oct. 2.		7.44	12.31	♀
787.	<i>Ceryle alcyon.</i>	July 21.		—	—	♀
788.	<i>Dendroica castanea.</i>	May, 1872.		—	—	♂ ad.
789.	<i>Turdus swainsoni</i>	Oct. 6.		7.12	12.00	♂
790.	" <i>aliciae.</i>	"	"	7.50	12.50	♂
791.	<i>Turdus swainsoni, aliciae.</i>	"	"	7.50	12.44	♀
792.	<i>Oedemia fusca, velvetina.</i>	" 7.		22.00	40.00	♂ ad.
793.	<i>Turdus swainsoni, aliciae.</i>	" 6.		7.12	12.12	♂
794.	"	"	"	7.75	13.00	♂
795.	"	"	"	7.44	12.25	♀
796.	<i>Oedemia americana.</i>	" 9.		20.50	33.62	♂ ad.
797.	<i>Carpodacus purpureus.</i>	"	"	6.37	10.44	♀
798.	"	"	"	6.12	10.44	♂ juv.
799.	<i>Dendroica coronata.</i>	" 6.		5.69	9.12	—
800.	<i>Salica americana.</i>	" 11.		14.50	26.00	♂ ad.
801.	<i>Oedemia americana.</i>	"	"	18.37	32.75	♀
802.	" <i>perspicillata.</i>	"	"	18.75	32.75	♀
803.	" <i>americana.</i>	" 13.		18.19	31.88.	♀

1875-76.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
804.	<i>Ceryle alcyon.</i>	Oct. 13.		13.06	22.56	♂
805.	<i>Sphyrapicus varius.</i>	"	"	8.62	15.25	♂ juv.
806.	<i>Harleia glacialis.</i>	" 19.		21.81	30.00	♂ ad.
807.	<i>Scolecophagus ferrugineus.</i>	" 24.		—	—	♂ ad.
808.	<i>Mergus cucullatus.</i>	Nov. 6.		18.37	26.50	♂ juv.
809.	<i>Scops asio.</i>	"	"	9.56	24.00	♀
810.	<i>Turdus migratorius (albinus?)</i>	" 1.		10.00	16.00	♂
811.	<i>Carpodacus purpureus.</i>	"	"	—	—	♂
812.	<i>Certhia familiaris.</i>	" 10.		—	—	♀
813.	<i>Corvus americanus.</i>	" 13.		18.75	30.00	♂
814.	<i>Chrysomitris tristis.</i>	" 25.		5.00	8.88	♂
815.	<i>Nyctea niva. (Long Island).</i>	Dec. 1.		23.50	58.31	♂ ad.
816.	<i>Tringa alpina</i>	"	"	8.81	15.50	♀
817.	<i>Squatarola helvetica.</i>	do	"	12.19	24.69	♂?
818.	<i>Tringa canutus.</i>	do	"	11.06	21.12	♂
819.	<i>Bucephala clangula.</i>	" 8.		17.12	29.00	♀
820.	<i>Chrysomitris tristis.</i>	" 9.		5.06	9.12	♂
821.	"	"	"	5.12	8.91	♂
822.	"	"	"	5.19	8.94	♂
823.	"	"	"	5.06	8.81	♂
824.	<i>Collurio borealis.</i>	" 11.		10.00	14.00	♀ juv.
825.	<i>Parus atricapillus.</i>	"	"	5.06	7.75	♀ ad.
826.	<i>Certhia familiaris.</i>	"	"	5.81	8.06	♀ ad.
827.	<i>Spizella monticola.</i>	" 8.		6.19	9.62	♂ ad.
828.	<i>Certhia familiaris.</i>	" 7.		5.75	8.00	♂ ad.
829.	<i>Anorthus troglodytes, hyemalis.</i>	"	"	4.00	6.12	♂ ad.
830.	<i>Spizella monticola.</i>	" 18.		6.44	9.44	♂ ad.
831.	<i>Chrysomitris tristis.</i>	" 17.		4.88	8.62	—
832.	"	"	"	4.88	8.62	—
833.	"	"	"	4.75	8.69	—
834.	<i>Regulus satrapa.</i>	"	"	4.00	6.81	♀ ad.
835.	<i>Setta carolinensis.</i>	" 16.		5.88	10.56	♂ ad.
836.	<i>Ortyx virginianus.</i>	" 21.		10.12	15.19	♀ ad.
837.	<i>Carpodacus purpureus</i>	" 22.		6.44	10.50	♀ ad.
838.	<i>Certhia familiaris.</i>	" 24.		5.50	7.75	♂ ad.
839.	<i>Regulus satrapa.</i>	" 21		4.25	6.81	♀ ad.
840.	"	"	"	4.12	6.59	♀ ad.

1876.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
841.	<i>Pinicola enucleator.</i>	Jan. 12.		8.75	12.75	♂ imm.
842.	"	"	"	8.25	13.62	♂ imm.
843.	"	"	"	8.31	13.37	♀ ad.
844.	<i>Chrysomitris tristis.</i>	" 11.		4.69	8.75	♂
845.	<i>Regulus satrapa.</i>	" 12.		4.31	6.88	♂ ad.
846.	<i>Chrysomitris tristis.</i>	" 11.		5.00	8.31	♀
847.	"	"	"	5.06	8.94	♂
848.	"	"	"	5.06	8.75	♂
849.	<i>Sitta carolinensis.</i>	" 15.		6.12	11.12	♂ ad.
850.	<i>Junco hyemalis.</i>	" 17.		6.25	9.94	♂ ad.
851.	<i>Parus atricapillus.</i>	" 12.		5.00	7.81	♂ ad.
852.	<i>Picus pubescens.</i>	"	"	6.25	11.75	♂ ad.
853.	<i>Ampelis cedrorum.</i>	" 15.		7.00	11.62	♀ ad.
854.	<i>Sitta carolinensis.</i>	"	"	6.00	11.12	♀ ad.
855.	<i>Ampelis cedrorum.</i>	"	"	7.00	11.75	♀ ad.
856.	"	"	"	7.00	11.56	♀ ad.
857.	<i>Parus atricapillus.</i>	" 17.		5.25	8.00	♀ ad.
858.	<i>Spizella monticola.</i>	" 12.		6.12	9.50	♀ ad.
859.	<i>Ampelis cedrorum.</i>	" 15.		7.00	11.37	♂ ad.
860.	<i>Junco hyemalis.</i>	" 12.		6.12	9.06	♂ ad.
861.	<i>Melospiza melodia.</i>	" 21.		6.19	8.50	♂ ad.
862.	<i>Spizella monticola.</i>	"	"	6.37	9.50	♂ ad.
863.	<i>Parus atricapillus.</i>	" 12.		4.94	7.62	♂ ad.
864.	<i>Regulus satrapa.</i> (Moult)	" 13.		2.75	6.88	♂ ad.
865.	"	" 12.		3.94	6.62	♀ ad.
866.	<i>Picus pubescens.</i>	Feb. 8.		7.00	12.12	♂ ad.
867.	<i>Junco hyemalis.</i>	"	"	6.31	10.00	♂ ad.
868.	"	"	"	6.37	9.69	♂ ad.
869.	"	"	"	6.19	10.12	♀ ad.
870.	"	"	"	6.37	9.81	♂ ad.
871.	"	"	"	6.50	10.06	♂ ad.
872.	<i>Anorthura troglodytes hyemalis</i>	"	"	4.25	6.12	—
873.	<i>Certhia familiaris.</i>	" 22.		5.75	8.06	♀ ad.
874.	"	"	"	5.75	8.06	♀ ad.
875.	<i>Picus pubescens.</i>	"	"	7.12	12.31	♀ ad.
876.	<i>Bonasa umbellus.</i>	"	"	—	—	♀
877.	<i>Picus pubescens.</i>	March 2.		7.00	12.12	♂ ad.

1876.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
878.	<i>Corvus americanus.</i>	March 4.		19.00	37.50	♀ ad.
879.	<i>Picus pubescens.</i>	"	"	6.62	11.94	♀ ad.
880.	"	"	"	6.94	11.88	♂ ad.
881.	<i>Bonasa umbellus.</i>	"	"	18.50	23.50	♂ ad.
882.	<i>Sphyrapicus varius.</i>	" 7.		8.37	15.31	♀ ad.
883.	<i>Sialia sialis.</i>	" 6.		7.12	12.50	♂ ad.
884.	"	"	"	7.00	12.56	♂ ad.
885.	"	"	"	7.00	12.81	♀ ad.
886.	<i>Passerella iliaca.</i>	" 7.		7.25	11.00	♀ ad.
887.	<i>Melospiza melodia.</i>	" 5.		6.62	8.75	♂ ad.
888.	<i>Mareca americana.</i>	" 12.		20.25	34.50	♂ ad.
889.	<i>Sialia sialis.</i>	"	"	7.12	12.50	♂ ad.
890.	<i>Chrysomitris tristis.</i>	" 10.		5.25	8.88	—
891.	<i>Ampelis cedrorum.</i>	" 23.		7.00	11.81	♂ ad.
892.	"	"	"	7.00	12.00	♀ ad.
893.	"	"	"	6.75	11.75	♀ ad.
894.	"	"	"	7.37	12.00	♂ ad.
895.	<i>Junco hyemalis.</i>	" 24.		6.25	9.75	♂ ad.
896.	<i>Melospiza melodia.</i>	"	"	6.44	8.69	♀ ad.
897.	<i>Passerella iliaca.</i>	"	"	7.31	11.50	♂ ad.
898.	"	"	"	7.00	10.75	♀ ad.
899.	<i>Turdus migratorius.</i>	" 28.		9.94	16.37	♂ ad.
900.	<i>Cyanurus cristatus.</i>	"	"	11.75	16.75	♂ ad.
901.	<i>Passerella iliaca.</i>	" 27.		7.69	11.88	♂ ad.
902.	<i>Junco hyemalis.</i>	" 28.		6.12	9.50	♂ ad.
903.	"	"	"	6.37	10.00	♂ ad.
904.	<i>Ampelis cedrorum.</i>	"	"	—	—	♀ ad.
905.	<i>Chrysomitris tristis.</i>	" 31.		5.50	9.06	♂ ad.
906.	<i>Melospiza melodia.</i>	"	"	6.44	8.88	♂ ad.
907.	<i>Ampelis cedrorum.</i>	"	"	7.00	11.75	♂ ad.
908.	<i>Scolecophagus ferrugineus.</i>	Apr. 15.		9.12	13.75	♀ ad.
909.	"	"	"	9.31	14.00	♀ ad.
910.	<i>Passer domesticus.</i>	" 11.		6.31	9.37	♂ ad.
911.	<i>Dendroica palmarum.</i>	" 15.		5.50	8.12	♂ ad.
912.	<i>Molothrus ater.</i>	" 17.		7.88	13.75	♂ ad.
913.	<i>Regulus calendula.</i>	"	"	4.25	7.00	♂ ad.
914.	<i>Spizella pusilla.</i>	"	"	5.56	8.00	♀ ad.

1876.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
915.	<i>Regulus calendula.</i>	Apr. 18.		4.50	7.19	♂ ad.
916.	<i>Dendroica coronata.</i>	" 21.		5.75	9.06	♂ ad.
917.	<i>Melospiza palustris.</i>	" "		5.88	7.94	♂ ad.
918.	<i>Poocetes gramineus.</i>	" "		6.12	10.25	♀ ad.
919.	<i>Sialia sialis.</i>	" "		7.00	12.31	♂ ad.
920.	<i>Sayornis fuscus.</i>	" "		6.69	10.37	♀ ad.
921.	<i>Certhia familiaris.</i>	" "		5.75	8.19	♀ ad.
922.	<i>Regulus calendula.</i>	" "		4.37	7.12	♂ ad.
923.	<i>Turdus pallasi.</i>	" "		7.00	11.37	♀ ad.
924.	<i>Tachycineta bicolor.</i>	" "		6.00	13.06	♂ ad.
925.	<i>Poocetes gramineus.</i>	" "		6.19	10.12	♀ ad.
926.	<i>Regulus calendula.</i>	" "		—	—	♂ ad.
927.	"	"	"	—	—	♂ ad.
928.	"	"	"	—	—	♂ ad.
929.	"	"	"	—	—	♂ ad.
930.	<i>Dendroica pinus.</i>	" 22.		5.75	9.12	♂ ad.
931.	" <i>palmarum.</i>	" "		5.62	8.62	♂ ad.
932.	"	"	"	5.37	8.12	♀ ad.
933.	"	"	"	5.31	8.56	♂ ad.
934.	" <i>coronata.</i>	" "		5.75	9.25	♂ ad.
935.	" <i>palmarum.</i>	" "		5.50	8.31	♀ ad.
936.	<i>Sivrus motacilla.</i>	" "		6.25	10.50	♂ ad.
937.	<i>Dendroica palmarum.</i>	" "		5.31	8.06	♀ ad.
938.	<i>Turdus pallasi.</i>	" "		7.06	11.25	♂ ad.
939.	"	"	"	7.25	11.37	♀ ad.
940.	<i>Certhia familiaris.</i>	" "		5.37	7.88	—
941.	<i>Spizella socialis.</i>	" "		5.25	8.88	♂ ad.
942.	<i>Sayornis fuscus.</i>	" "		7.06	11.12	♂ ad.
943.	<i>Regulus calendula.</i>	" 24.		4.50	7.06	♂ ad.
944.	<i>Junco hyemalis.</i>	" 24.		6.00	9.25	♀ ad.
945.	<i>Dendroica coronata.</i>	" "		5.62	9.00	♂ ad.
946.	<i>Molothrus ater.</i>	" "		7.88	13.50	♂ ad.
947.	<i>Passer domesticus.</i>	" "		6.37	9.88	♂ ad.
948.	<i>Sayornis fuscus.</i>	" "		6.25	10.69	♀ ad.
949.	<i>Sivrus motacilla.</i>	" "		6.19	10.50	♂ ad.
950.	"	"	"	6.19	10.50	♂ ad.
951.	<i>Regulus satrapa.</i>	" 26.		4.19	—	♂ ad.

1876.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
952.	<i>Regulus calendula.</i>	April 26.		4.50	7.25	♂ ad.
953.	"	"	"	4.37	7.14	♂ ad.
954.	"	"	"	4.50	7.25	♂ ad.
955.	<i>Sivrus motacilla.</i>	" 25.		6.06	10.31	♂ ad.
956.	<i>Melospiza palustris.</i>	" "		—	—	—
957.	<i>Picus pubescens.</i>	" 26.		—	—	♂ ad.
958.	<i>Regulus calendula.</i>	" "		—	—	♂ ad.
959.	<i>Sialia sialis.</i>	" "		6.75	12.00	♀ ad.
960.	<i>Fringoides macularius.</i>	" 29.		7.75	13.62	♀ ad.
961.	<i>Passer domesticus.</i>	" 28.		6.44	10.19	♀ ad.
962.	<i>Dendroica palmarum.</i>	" "		5.50	8.69	♂ ad.
963.	<i>Zenaidura carolinensis.</i>	" 29.		11.81	18.25	♂ ad.
964.	<i>Sivrus motacilla.</i>	" "		6.50	10.50	♂ ad.
965.	<i>Molothrus ater.</i>	" "		7.75	13.50	♂ ad.
966.	<i>Melospiza palustris.</i>	" "		6.06	8.31	♂ ad.
967.	"	"	"	6.12	8.25	♂ ad.
968.	<i>Dendroica coronata.</i>	" "		5.75	9.00	♂ ad.
969.	"	"	"	5.69	9.00	♂ ad.
970.	"	"	"	5.81	9.06	♂ ad.
971.	<i>Turdus swainsoni.</i>	May 9.		7.25	12.25	♂ ad.
972.	"	"	"	7.19	12.06	♂ ad.
973.	<i>Fringoides macularius.</i>	" 10.		8.31	14.00	♀ ad.
974.	<i>Helminthophaga ruficapilla.</i>	" "		4.75	7.50	♂ ad.
975.	<i>Dendroica virens.</i>	" "		4.94	7.25	♀ ad.
976.	" <i>pennsylvanica.</i>	" "		5.00	7.81	♀ ad.
977.	<i>Vireo solitarius.</i>	" "		5.50	9.50	♀ ad.
978.	<i>Myiodiocetes mitratus.</i>	" "		5.81	8.31	♀ ad.
979.	<i>Dendroica virens.</i>	" "		5.25	8.06	♂ ad.
980.	" <i>coronata.</i>	" "		5.44	8.44	♀ ad.
981.	<i>Mniotilta varia.</i>	" "		5.31	8.44	♂ ad.
982.	<i>Regulus calendula (mored).</i>	" 11.		4.12	6.88	♀ (ad?)
983.	<i>Myiodiocetes canadensis.</i>	" 22.		5.56	8.06	♂ ad.
984.	<i>Dendroica maculosa.</i>	" 28.		5.25	7.56	♂ ad.
985.	" <i>blackburniae.</i>	" 17.		5.25	8.37	♂ ad.
986.	<i>Myiodiocetes canadensis.</i>	" 22.		5.75	8.06	♂ ad.
987.	"	" 22.		5.56	8.06	♀ ad.
988.	"	" 25		5.69	8.44	♂ ad.

1876.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
989.	<i>Dendroeca caerulea</i> .	May 18.		5.44	8.06	♂ ad.
990.	"	" 16.		5.25	7.94	♀ ad.
991.	<i>Helminthophaga ruficapilla</i> .	12.		4.94	7.75	♂ ad.
992.	<i>Siurus naevius</i> .	" 26.		6.12	9.88	♂ ad.
993.	<i>Myiodioetes mitratus</i> .	" 17.		5.62	8.37	♂ ad.
994.	<i>Helminthus vermivorus</i> .	" "		5.31	8.56	♀ ad.
995.	<i>Dendroeca castanea</i> .	" 26.		5.81	9.06	♂ ad.
996.	<i>Parula americana</i> .	" 18.		4.81	7.65	♂ ad.
997.	<i>Dendroeca maculosa</i> .	" 16.		5.25	7.75	♂ ad.
998.	" <i>caerulea</i> .	" 18.		5.25	7.94	♀ ad.
999.	<i>Geothlypis trichas</i> .	" 27.		5.62	7.44	♂ ad.
1000.	" <i>philadelphia</i> .	" 26.		5.62	8.12	♂ ad.
1001.	<i>Dendroeca pennsylvanica</i> .	" 18.		5.19	8.19	♂ ad.
1002.	<i>Helminthophaga ruficapilla</i> .	" 17.		4.69	7.25	♀ ad.
1003.	<i>Setophaga ruticilla</i> .	" "		5.25	7.62	♂ ad.
1004.	<i>Siurus auricapillus</i> .	" "		6.00	9.37	♀ ad.
1005.	<i>Dendroeca blackburniae</i> .	" "		5.44	8.50	♂ ad.
1006.	"	" "		—	—	♂ ad.
1007.	<i>Myiostila varia</i> .	" 26.		5.25	8.25	♀ ad.
1008.	<i>Dendroeca blackburniae</i> .	" 18.		5.31	8.12	♀ ad.
1009.	"	" 13.		5.12	8.62	♂ ad.
1010.	" <i>castanea</i> .	" 26.		5.56	9.06	♂ ad.
1011.	"	" "		5.25	8.62	♀ ad.
1012.	" <i>maculosa</i> .	" 16.		5.19	7.88	♂ ad.
1013.	" <i>striata</i> .	" 26.		5.75	9.25	♂ ad.
1014.	" <i>maculosa</i> .	" 17.		5.12	7.62	♂ ad.
1015.	" <i>blackburniae</i> .	" "		5.44	8.56	♂ ad.
1016.	<i>Myiodioetes canadensis</i> .	" 23.		—	—	♂ ad.
1017.	<i>Dendroeca maculosa</i> .	" 16.		5.25	7.75	♂ ad.
1018.	" <i>blackburniae</i> .	" "		5.19	8.12	♂ ad.
1019.	<i>Geothlypis trichas</i> .	" 17.		5.37	7.12	♂ ad.
1020.	<i>Dendroeca virens</i> .	" 18.		5.25	8.06	♂ ad.
1021.	<i>Siurus auricapillus</i> .	" 12.		—	—	♀ ad.
1022.	<i>Dendroeca castanea</i> .	" 25.		5.69	9.25	♂ ad.
1023.	<i>Myiodioetes canadensis</i> .	" 26.		5.50	7.94	♂ ad.
1024.	<i>Dendroeca maculosa</i> .	" 17.		5.19	7.62	♂ ad.
1025.	<i>Myiodioetes mitratus</i> .	" 18.		5.75	8.50	♂ ad.

1876.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
1026.	<i>Dendroeca striata</i> .	May 23.		5.81	9.50	♂ ad.
1027.	" <i>castanea</i> .	" "		5.62	8.94	ad.
1028.	"	" 22.		5.62	8.94	♂ ad.
1029.	"	" 26.		5.81	9.06	♂ ad.
1030.	"	" 22.		5.75	9.19	♂ ad.
1031.	"	" 25.		5.62	8.94	♀ ad.
1032.	<i>Icteria virens</i> .	" 23.		7.50	10.19	♂ ad.
1033.	<i>Helminthus vermivorus</i> .	" "		5.62	9.00	♂ ad.
1034.	<i>Myiodioetes canadensis</i> .	" "		5.62	8.00	♂ ad.
1035.	<i>Dendroeca pennsylvanica</i> .	" 22.		5.31	7.88	♀ ad.
1036.	" <i>blackburniae</i> .	" 23.		—	—	♀ ad.
1037.	"	" 18.		5.15	8.31	♂ ad.
1038.	<i>Siurus auricapillus</i> .	" 16.		6.31	10.00	♂ ad.
1039.	<i>Dendroeca pennsylvanica</i> .	" 12.		—	—	♂ ad.
1040.	" <i>striata</i> .	" 22.		5.62	8.81	♀ ad.
1041.	"	" 25.		5.56	9.19	♂ ad.
1042.	" <i>castanea</i> .	" 20.		5.81	9.25	♂ ad.
1043.	" <i>blackburniae</i> .	" 23.		5.37	8.69	♂ ad.
1044.	" <i>pennsylvanica</i> .	" 20.		—	—	♀ ad.
1045.	"	" 23.		5.19	7.75	♀ ad.
1046.	" <i>aestiva</i> .	" 12.		—	—	♂ ad.
1047.	" <i>castanea</i> .	" 25.		—	—	♂ ad.
1048.	"	" "		—	—	♂ ad.
1049.	<i>Myiodioetes canadensis</i> .	" "		5.69	8.44	♂ ad.
1050.	<i>Dendroeca caerulea</i> .	" 20.		—	—	♂ ad.
1051.	" <i>castanea</i> .	" 25.		—	—	♂ ad.
1052.	" <i>striata</i> .	" 23.		5.69	9.00	♀ ad.
1053.	<i>Myiodioetes canadensis</i> .	" 18.		5.62	8.00	♂ ad.
1054.	<i>Dendroeca blackburniae</i> .	" "		5.19	8.44	♂ ad.
1055.	<i>Myiodioetes canadensis</i> .	" 22.		5.69	8.06	♂ ad.
1056.	<i>Setophaga ruticilla</i> .	" 26.		5.56	8.00	♂ ad.
1057.	<i>Dendroeca striata</i> .	" 25.		5.69	9.25	♂ ad.
1058.	" <i>maculosa</i> .	" "		5.00	7.37	♀ ad.
1059.	" <i>blackburniae</i> .	" 23.		5.37	8.75	♂ ad.
1060.	<i>Myiodioetes mitratus</i> .	" 17.		5.62	8.37	♂ ad.
1061.	<i>Icteria virens</i> .	" 23.		7.31	9.81	♂ ad.
1062.	<i>Dendroeca blackburniae</i> .	" 25.		—	—	♀ ad.

1876.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
1063.	<i>Spinus naevius.</i>	May 25.		6.19	9.50	♂ ad.
1064.	<i>Geothlypis trichas.</i>	" 19.		—	—	♀ ad.
1065.	<i>Myiiodioides mitratus.</i>	" 17.		5.56	8.31	♂ ad.
1066.	<i>Spinus naevius.</i>	" 13.		5.81	9.50	♂ ad.
1067.	<i>Dendroica castanea.</i>	" 25.		—	—	♂ ad.
1068.	<i>Spinus motacilla.</i>	" 18.		6.31	10.25	♀ ad.
1069.	<i>Helminthophaga ruficapilla.</i>	" 13.		5.62	9.00	♂ ad.
1070.	<i>Dendroica castanea.</i>	" 26.		5.69	8.94	♀ ad.
1071.	<i>Spinus auricapillus.</i>	" 5.		6.12	9.94	♂ ad.
1072.	<i>Geothlypis trichas.</i>	" 8.		5.50	7.31	♂ ad.
1073.	<i>Mniotilta varia.</i>	" 5.		5.12	8.56	♂ ad.
1074.	<i>Helminthophaga ruficapilla.</i>	" 8.		4.81	7.31	♂ ad.
1075.	<i>Parula americana.</i>	" 8.		4.94	7.69	♂ ad.
1076.	<i>Dendroica blackburniae.</i>	" 9.		5.19	8.37	♂ ad.
1077.	" <i>palmarum.</i>	" 5.		5.31	8.19	♀ ad.
1078.	<i>Setophaga ruticilla.</i>	" 8.		5.50	7.75	♂ ad.
1079.	<i>Dendroica virens.</i>	" 1.		5.12	7.62	♂ ad.
1080.	" <i>coronata.</i>	" 3.		5.56	8.88	♀ ad.
1081.	" "	" 8.		5.75	9.12	♂ ad.
1082.	" <i>palmarum.</i>	" 3.		5.25	8.19	♀ ad.
1083.	<i>Mniotilta varia.</i>	" 4.		5.19	8.75	♂ ad.
1084.	<i>Myiiodioides mitratus.</i>	" 8.		5.81	8.56	♂ ad.
1085.	<i>Helminthophaga ruficapilla.</i>	" 6.		5.00	7.69	♂ ad.
1086.	<i>Dendroica virens.</i>	" "		5.31	8.19	♂ ad.
1087.	" <i>caerulea.</i>	" 9.		5.31	8.19	♂ ad.
1088.	<i>Helminthophaga ruficapilla.</i>	" 8.		4.94	7.75	♂ ad.
1089.	<i>Dendroica maculosa.</i>	" 16.		5.12	7.69	♂ ad.
1090.	<i>Helminthophaga ruficapilla.</i>	" 6.		5.06	7.70	♂ ad.
1091.	<i>Dendroica virens.</i>	" 9.		5.12	7.81	♂ ad.
1092.	<i>Spinus auricapillus.</i>	" 8.		6.37	10.06	♂ ad.
1093.	<i>Dendroica caerulea.</i>	" 9.		5.31	8.25	♂ ad.
1094.	" <i>virens.</i>	" "		5.12	7.81	♂ ad.
1095.	<i>Helminthophaga ruficapilla.</i>	" 12.		5.75	8.88	♂ ad.
1096.	<i>Dendroica caerulea.</i>	" "		5.12	7.69	♀ ad.
1097.	" <i>castanea.</i>	" 16.		5.50	9.00	♂ ad.
1098.	<i>Geothlypis trichas.</i>	" 9.		5.37	7.19	♂ ad.
1099.	<i>Dendroica blackburniae.</i>	" 16.		5.25	8.44	♂ ad.

1876.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
1100.	<i>Dendroica caerulea.</i>	May 16.		5.12	7.50	♀ ad.
1101.	" <i>blackburniae.</i>	" "		5.25	8.44	♂ ad.
1102.	<i>Helminthophaga ruficapilla.</i>	" 9.		5.44	8.62	♂ ad.
1103.	<i>Dendroica blackburniae.</i>	" 16.		5.19	8.19	♂ ad.
1104.	" <i>coronata.</i>	" "		5.44	8.50	♀ ad.
1105.	<i>Geothlypis trichas.</i>	" "		5.12	7.00	♀ ad.
1106.	<i>Helminthophaga ruficapilla.</i>	" "		5.50	8.81	♂ ad.
1107.	<i>Myiiodioides mitratus.</i>	" 13.		5.81	8.62	♂ ad.
1108.	<i>Dendroica blackburniae.</i>	" 11.		5.31	8.56	♂ ad.
1109.	<i>Myiiodioides canadensis.</i>	" 16.		5.75	8.25	♂ ad.
1110.	<i>Spinus motacilla.</i>	" 13.		6.37	10.25	♀ ad.
1111.	<i>Tyrus solitarius.</i>	" 11.		—	—	♀ ad.
1112.	<i>Sayornis fusca.</i>	" 18.		—	—	♂ ad.
1113.	<i>Spizella socialis.</i>	" 8.		5.31	8.81	♀ ad.
1114.	<i>Troglodytes aedon (domesticus).</i>	" 3.		5.06	6.75	♂ ad.
1115.	<i>Sialia sialis.</i>	" 25.		6.00	11.25	♂ juv.
1116.	<i>Spizella pusilla.</i>	" 27.		5.75	8.12	♀ ad.
1117.	<i>Tyrus solitarius.</i>	" 9.		5.50	9.31	♂ ad.
1118.	<i>Spizella socialis.</i>	" 1.		5.37	8.62	♂ ad.
1119.	<i>Tyrus flavifrons.</i>	" 22.		6.00	9.75	♂ ad.
1120.	<i>Chrysomitris tristis.</i>	" 2.		5.31	9.06	♂ ad.
1121.	" "	" "		4.69	8.25	♂ ad.
1122.	<i>Empidonax flaviventris.</i>	" 23.		5.75	8.88	♀ ad.
1123.	<i>Telmatochlamys palustris.</i>	" 18.		—	—	♂ ad.
1124.	<i>Passer domesticus.</i>	" 26.		6.50	9.69	♀ ad.
1125.	<i>Troglodytes aedon (domesticus).</i>	" 3.		5.19	6.81	♂ ad.
1126.	<i>Mimus carolinensis.</i>	" 8.		8.88	11.44	♀ ad.
1127.	<i>Antrostomus vociferus.</i>	" 12.		9.56	18.12	♀ ad.
1128.	<i>Tringoides macularius.</i>	" 23.		7.56	13.31	♂ ad.
1129.	<i>Turdus mustelinus.</i>	" 8.		8.19	13.25	♀ ad.
1130.	" <i>fuscescens.</i>	" "		7.37	11.75	♀ ad.
1131.	<i>Tringoides macularius.</i>	" 30.		7.69	13.25	♂ ad.
1132.	<i>Colaptes auratus.</i>	" 5.		11.81	20.00	♀ ad.
1133.	<i>Pyrranga rubra.</i>	" 23.		6.94	11.94	♀ ad.
1134.	<i>Turdus mustelinus.</i>	" 26.		8.19	13.62	♀ ad.
1135.	<i>Empidonax minimus.</i>	" 23.		5.44	8.19	♀ ad.
1136.	" "	" 22.		5.50	8.00	♀ ad.

1876.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
1137.	<i>Empidonax flaviventris.</i>	May 26.		5.62	8.69	♂ ad.
1138.	"	" 23.		5.69	8.75	♂ ad.
1139.	" <i>minimus.</i>	" 20.		—	—	♂ ad.
1140.	"	" 8.		5.37	8.12	♀ ad.
1141.	"	" 12.		5.56	8.37	♀ ad.
1142.	" <i>traillii.</i>	" 26.		6.00	9.19	♂ ad.
1143.	"	" 27.		6.25	9.50	♂ ad.
1144.	" <i>minimus.</i>	" 8.		5.50	8.25	♂ ad.
1145.	" <i>flaviventris.</i>	" 23.		5.69	8.88	♂ ad.
1146.	<i>Ceryle alcyon.</i>	" 5.		12.81	22.81	♂ ad.
1147.	<i>Pipilo erythrophthalmus.</i>	" 2.		8.00	10.56	♀ ad.
1148.	<i>Harporhynchus rufus.</i>	" 8.		11.50	13.62	♂ ad.
1149.	<i>Cyanurus cristatus.</i>	" 2.		11.81	17.37	♂ ad.
1150.	<i>Coccyzus erythrophthalmus.</i>	" 22.		11.75	15.69	♀ ad.
1151.	<i>Zonotrichia albicollis.</i>	" 20.		—	—	♂ ad.
1152.	<i>Agelaius phoeniceus.</i>	" 2.		8.06	12.88	♀ ad.
1153.	<i>Turdus swainsoni, aliciae.</i>	" 25.		7.62	12.75	♀ ad.
1154.	" <i>pallasi.</i>	" 3.		7.00	11.12	♀ ad.
1155.	<i>Pipilo erythrophthalmus.</i>	" 27.		8.50	11.62	♂ ad.
1156.	"	" 5.		8.06	11.06	♂ ad.
1157.	<i>Agelaius phoeniceus.</i>	" 2.		9.44	15.12	♂ ad.
1158.	<i>Turdus fuscescens.</i>	" 26.		7.31	11.75	♀ ad.
1159.	" <i>aliciae.</i>	" 25.		7.88	13.12	♂ ad.
1160.	<i>Totanus solitarius.</i>	" 19.		8.75	16.37	♂ ad.
1161.	<i>Agelaius phoeniceus.</i>	" 2.		9.50	15.25	♂ ad.
1162.	<i>Turdus swainsoni.</i>	" 11.		7.56	12.94	♀ ad.
1163.	<i>Tyrannus carolinensis.</i>	" 13.		—	—	♂ ad.
1164.	<i>Turdus swainsoni.</i>	"		—	—	♂ ad.
1165.	"	" 25.		7.44	12.31	♂ ad.
1166.	<i>Agelaius phoeniceus.</i>	" 23.		9.50	15.50	♂ ad.
1167.	<i>Carpodacus purpureus.</i>	" 5.		6.19	9.75	♂ ad.
1168.	<i>Zonotrichia albicollis.</i>	" 2.		6.75	9.50	♂ ad.
1169.	<i>Pyrranga rubra.</i>	" 27.		7.31	11.81	♂ ad.
1170.	<i>Goniaphea ludoviciana.</i>	" 12.		8.00	12.50	♂ ad.
1171.	<i>Myiarchus crinitus.</i>	" 13.		9.12	13.50	♂ ad.
1172.	<i>Pyrranga rubra.</i>	" 25.		7.31	11.75	♂ ad.
1173.	"	" 15.		7.19	11.88	♀ ad.

1876.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
1174.	<i>Pyrranga rubra.</i>	May 23.		7.31	11.69	♂ ad.
1175.	<i>Icterus spurius.</i>	"		7.25	10.50	♂ ad.
1176.	<i>Fringoides macularius.</i>	"		8.12	13.88	♀ ad.
1177.	<i>Zonotrichia leucophrys.</i>	"		6.88	9.75	♀ ad.
1178.	<i>Dolichonyx oryzivorus.</i>	"		7.69	12.37	♂ ad.
1179.	<i>Icterus baltimore.</i>	" 26.		7.50	11.31	♀ ad.
1180.	<i>Petrochelidon lunifrons.</i>	" 23.		5.88	12.75	♀ ad.
1181.	<i>Zonotrichia leucophrys.</i>	"		6.88	9.88	♀ ad.
1182.	<i>Pyrranga rubra.</i>	"		—	—	♂ ad.
1183.	"	"		—	—	♂ ad.
1184.	"	"		7.44	12.12	♂ ad.
1185.	"	"		7.37	12.12	♂ ad.
1186.	"	"		7.31	11.81	♂ ad.
1187.	"	"		16. 6.94	11.75	♂ ad.
1188.	"	"		23. 7.25	11.62	♀ ad.
1189.	<i>Turdus swainsoni.</i>	" 18.		7.37	12.69	♂ ad.
1190.	<i>Icterus baltimore.</i>	" 11.		7.25	11.31	♂ ad.
1191.	<i>Pyrranga rubra.</i>	" 23.		7.19	12.00	♂ ad.
1192.	"	"		7.19	12.00	♂ ad.
1193.	"	"		7.06	11.44	♀ ad.
1194.	<i>Turdus swainsoni.</i>	"		7.00	11.31	♂ ad.
1195.	<i>Agelaius phoeniceus.</i>	"		8.00	12.75	♀ ad.
1196.	<i>Vireo flavifrons.</i>	" 27.		—	—	♀ ad.
1197.	<i>Goniaphea ludoviciana.</i>	" 19.		—	—	♂ ad.
1198.	<i>Pyrranga rubra.</i>	" 23.		7.06	11.75	♂ ad.
1199.	"	"		7.31	12.12	♂ ad.
1200.	"	" 18.		—	—	♂ ad.
1201.	<i>Turdus fuscescens.</i>	" 5.		7.56	12.25	♂ ad.
1202.	<i>Goniaphea ludoviciana.</i>	" 19.		—	—	♀ ad.
1203.	"	"		—	—	♂ ad.
1204.	<i>Icterus baltimore.</i>	" 18.		7.75	12.00	♂ ad.
1205.	<i>Cotyle riparia.</i>	" 23.		5.19	10.88	♂ ad.
1206.	<i>Vireo solitarius.</i>	" 18.		5.81	9.69	♂ ad.
1207.	"	" 9.		5.75	9.69	♂ ad.
1208.	"	" 11.		5.69	9.50	♂ ad.
1209.	"	" 9.		5.62	9.62	♀ ad.
1210.	" <i>flavifrons.</i>	" 12.		—	—	♀ ad.

1876.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
1211.	<i>Vireo flavifrons.</i>	May 13.		6.00	9.69	♂ ad.
1212.	" "	" 8.		5.94	10.00	♂ ad.
1213.	" <i>gilvus.</i>	" "		5.94	9.25	♂ ad.
1214.	" "	" 18.		5.75	8.94	♂ ad.
1215.	" "	" "		5.75	9.19	♂ ad.
1216.	" <i>olivaceus.</i>	" 22.		6.44	10.31	♂ ad.
1217.	" <i>novboracensis.</i>	" 17.		5.31	8.00	♀ ad.
1218.	<i>Cyanospiza cyanea.</i>	" 22.		5.69	8.81	♂ ad.
1219.	" "	" 27.		5.75	8.50	♀ ad.
1220.	<i>Parus atricapillus.</i>	" 5.		5.12	7.75	—
1221.	<i>Troglodytes domesticus</i> , Bartram.	3.		4.94	7.00	♂ ad.
1222.	<i>Haliaeetus leucocephalus.</i>	June 2.		35.50	89.25	♀ ad.
1223.	<i>Myiodytes mitratus</i> (parent). "	7.		—	—	♀ ad.
1224.	<i>Buteo pennsylvanica.</i>	May 3.		15.69	36.69	♂ ad.
1225.	" "	" 5.		17.50	38.69	♀ ad.
1226.	<i>Accipiter cooperi.</i>	" 10.		19.75	35.00	♀ ad.
1227.	<i>Mniotilta varia.</i>	July 26.		5.75	8.88	♂ juv.
1228.	<i>Siurus auricapillus.</i>	Aug. 16.		6.44	9.81	♀ ad.
1229.	<i>Buteo pennsylvanicus.</i>	" 17.		16.37	36.75	♂ juv.
1230.	<i>Contopus virens.</i>	" 16.		6.37	10.37	♀ juv.
1231.	<i>Cotyle riparia.</i>	" 17.		5.06	10.75	♂?
1232.	<i>Haliaeetus leucocephalus</i>	" "		33.00	81.50	♀ ad.
1233.	<i>Turdus swainsoni.</i>	Oct. 6, 1875.		7.12	12.00	♂
1234.	<i>Icterus spurius.</i>	Aug. 21, 1875.		7.50	10.56	♀ juv.
1235.	<i>Ampelis cedrorum</i>	" 17.		—	—	♂ juv.
1236.	<i>Contopus virens.</i>	" 30.		6.50	10.50	♂
1237.	" "	" "		6.50	10.56	♂
1238.	" "	" "		—	—	♂
1239.	" "	" "		—	—	♂
1240.	<i>Geothlypis trichas.</i>	" 16.		5.00	6.62	♀
1241.	<i>Vireo olivaceus.</i>	" "		—	—	♂
1242.	<i>Pyrranga rubra.</i>	" 30.		7.37	11.88	♀
1243.	" "	" 31.		7.37	11.81	♀
1244.	<i>Icterus baltimore.</i>	" "		7.50	11.88	♂ juv.
1245.	<i>Icterus spurius.</i>	Aug 21, 1876.		7.37	10.25	♂ juv.
1246.	<i>Dolichonyx oryzivorus.</i>	" "		7.50	11.50	♂ ad.
1247.	" "	" "		7.31	11.75	♂

1876.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
1248.	<i>Dolichonyx oryzivorus.</i>	Aug 21.		7.56	11.94	♂
1249.	" "	" "		7.69	12.00	♂.
1250.	<i>Contopus borealis.</i>	Sep. 5.		7.19	12.25	—
1251.	<i>Siurus naevius.</i>	Aug. 31.		6.00	9.25	♀
1252.	<i>Parus carolinia.</i>	Sep. 6.		9.75	15.25	♂ juv.
1253.	<i>Mimus carolinensis.</i>	" 8.		9.00	11.75	♂ juv.
1254.	<i>Pyrranga rubra.</i>	" 12.		7.37	11.69	♀
1255.	<i>Totanus solitarius.</i>	" 11.		8.12	16.00	♂ juv.
1256.	<i>Contopus virens.</i>	" 14.		6.50	10.50	♀
1257.	<i>Siurus naevius.</i>	" 11.		5.94	9.37	♀
1258.	<i>Hirundo horreorum.</i>	" 12.		6.25	12.56	♂ juv.
1259.	<i>Myiodytes canadensis.</i>	" 11.		5.50	8.00	♀
1260.	<i>Passerculus savanna.</i>	" 12.		5.81	9.19	♀
1261.	<i>Dendroica pennsylvanica.</i>	" 14.		5.06	7.75	♂ juv.
1262.	<i>Fringoides macularius.</i>	" "		7.81	13.75	♀ juv.
1263.	<i>Vireo olivaceus.</i>	" 8.		6.19	10.37	♀
1264.	<i>Dendroica maculosa.</i>	" 11.		5.00	7.56	♀ juv.
1265.	" <i>virens.</i>	" 12.		5.19	7.81	—
1266.	<i>Vireo olivaceus.</i>	" 8.		—	—	♀ juv.
1267.	<i>Parula americana.</i>	" 11.		4.81	7.62	♂ juv.
1268.	<i>Mniotilta varia.</i>	" 15.		5.12	8.12	♀
1269.	<i>Vireo olivaceus.</i>	" 11.		6.12	10.19	♂ juv.
1270.	<i>Chrysomitris tristis.</i>	" 12.		5.00	9.00	♂ ad.
1271.	<i>Dendroica virens.</i>	" 11.		5.00	7.62	♀?
1272.	<i>Parula americana.</i>	" 8.		4.25	7.44	♂ juv.
1273.	<i>Dendroica virens.</i>	" 6.		5.00	7.31	—
1274.	<i>Troglodytes domesticus.</i>	" 14.		5.19	6.69	♂
1275.	<i>Dendroica (Perisoglossa) tigrina.</i>	" 8.		5.22	8.31	—
1276.	" <i>castanea.</i>	" 18.		5.37	8.31	♀ juv.
1277.	<i>Helminthophaga ruficapilla.</i>	" 16.		4.75	7.37	♂
1278.	<i>Telmatodytes palustris.</i>	" 18.		4.75	6.00	♀ juv.
1279.	<i>Helminthophaga peregrina.</i>	" 20.		5.00	7.69	♂ juv.
1280.	<i>Dendroica virens.</i>	" 29.		5.12	7.81	♂ juv.
1281.	" <i>pennsylvanica.</i>	" "		4.94	7.50	juv.
1282.	<i>Sphyrapicus varius.</i>	" "		8.00	14.50	♂ juv.
1283.	<i>Podiceps albicollis.</i>	" "		6.44	9.00	♂
1284.	<i>Parus atricapillus.</i>	" "		5.62	8.25	♂

1876.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
1285.	<i>Vireo solitarius.</i>	Sep. 29.		5.75	9.37	♀ juv.
1286.	<i>Parula americana.</i>	" "		4.94	7.75	♂ juv.
1287.	<i>Dendroica caerulescens</i>	" "		5.44	8.12	♂ ad.
1288.	<i>Spizella pusilla.</i>	" "		5.69	8.12	♂
1289.	<i>Helminthophaga peregrina.</i>	" 30.		4.72	7.25	♀ juv.
1290.	<i>Dendroica blackburniae.</i>	" "		5.00	8.25	♂ juv.
1291.	" "	" "		5.41	8.37	♀ ad.
1292.	<i>Sphyrapicus varius.</i>	" "		8.81	15.50	♂ juv.
1293.	<i>Melospiza palustris.</i>	" "		5.81	7.37	♀ juv.
1294.	" "	" "		6.06	7.88	♂ juv.
1295.	<i>Harporhynchus rufus.</i>	Oct. 3.		11.25	12.69	♂ juv.
1296.	<i>Sphyrapicus varius.</i>	" "		8.62	15.75	♀ ad.
1297.	<i>Turdus swainsoni.</i>	" "		7.12	11.69	♀
1298.	<i>Dendroica maculosa.</i>	" "		5.19	7.56	♀ ad.
1299.	" <i>castanea.</i>	" "		5.69	9.00	♂ juv.
1300.	" <i>caerulescens.</i>	" "		5.25	7.75	♂ ad.
1301.	" <i>virens.</i>	" "		5.06	7.50	♀
1302.	<i>Regulus satrapa.</i>	" "		4.44	7.69	♂ ad.
1303.	" "	" "		4.19	6.62	♀ ad.
1304.	<i>Sphyrapicus varius.</i>	" 4.		8.69	15.50	♀ ad.
1305.	" "	" "		8.44	15.00	♂ juv.
1306.	<i>Junco hyemalis.</i>	" "		6.25	9.88	♂ ad.
1307.	<i>Porzana carolina.</i>	" "		8.75	14.00	♀ juv.
1308.	<i>Anorthura troglodytes, hyemalis.</i>	" "		4.06	6.00	juv.
1309.	<i>Dendroica maculosa.</i>	" 5.		5.06	7.44	♂ juv.
1310.	" <i>caerulescens.</i>	" "		5.25	7.94	♂ ad.
1311.	<i>Certhia familiaris.</i>	" "		5.62	7.75	♀
1312.	<i>Gallinago wilsoni.</i>	" 16.		11.12	17.25	♂ ad.
1313.	" "	" "		10.50	16.12	♂ juv.
1314.	<i>Carduelis purpureus.</i>	" 6.		6.12	10.44	♂ ad.
1315.	<i>Troglodytes domesticus.</i>	" 16.		4.81	6.37	♀
1316.	<i>Dendroica caerulescens.</i>	" 14.		5.31	7.88	♂ ad.
1317.	<i>Mniotilta varia.</i>	" "		5.50	8.56	♀ ad.
1318.	<i>Regulus calendula.</i>	" "		—	—	♂ ad.
1319.	<i>Turdus swainsoni, aliciae.</i>	" 17.		8.25	13.00	♂
1320.	<i>Colaptes auratus.</i>	" 18.		13.00	20.81	♂ ad.
1321.	<i>Sialia sialis.</i>	" "		7.00	12.31	♂ ad.

1876-77.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
1322.	<i>Turdus pallasi.</i>	Oct. 18.		7.19	11.56	♂
1323.	" "	" "		6.94	11.00	♀
1324.	<i>Bubo virginianus.</i>	" 17.		21.88	55.25	♂ juv.
1325.	<i>Gallinago wilsoni.</i>	" 19.		11.00	17.19	♂ ad.
1326.	<i>Ardea sponsa.</i>	" 23.		16.12	29.44	♂ ad.
1327.	<i>Sphyrapicus varius.</i>	" 21.		8.88	15.37	♂ ad.
1328.	<i>Syrnium nebulosum.</i>	" 23.		19.25	44.75	♀ ad.
1329.	<i>Porzana carolina</i> ^{Fort Miller, Washington County, N. Y.}	Sept. 12.		9.50	14.56	♂ ad.
1330.	<i>Picus villosus.</i> ^{Fort Miller, N. Y.}	Nov. 12.		9.69	15.88	♂ ad.
1331.	" <i>pubescens</i> do	" "		7.00	11.88	♂ ad.
1332.	" <i>villosus.</i> do	" "		10.12	16.31	♂ ad.
1333.	" do	" "		9.50	15.62	♀ ad.
1334.	<i>Melanerpes erythrocephalus.</i> ^{Highland Falls, N. Y.}	" 1.		9.44	17.25	♀ juv.
1335.	<i>Turdus pallasi.</i>	Oct. 30.		7.12	11.75	♂ juv.
1336.	<i>Hearelda glacialis.</i>	Nov. 29.		13.94	29.69	♂ ad.
1337.	<i>Ortyx virginianus.</i>	Dec. 21, 1875.		11.00	15.75	♂ ad.
1338.	<i>Scops asio.</i>	" 8,		9.69	24.00	♀
1339.	" "	1877 Dec. 27.		9.19	22.69	♀
1340.	<i>Colurio borealis.</i>	Feb. 12.		10.06	14.00	♀
1341.	<i>Fuligula vallisneria.</i>	Mar. 23.		22.31	35.12	♂ ad.
1342.	" "	" "		22.31	34.75	♂ ad.
1343.	<i>Mergus merganser.</i>	" "		26.65	37.00	♂ ad.
1344.	" "	" "		24.30	34.00	♀ ad.
1345.	<i>Sturnella magna.</i>	Fairfield, Conn. Apr. 7.		10.82	16.43	♂ ad.
1346.	<i>Tachycineta bicolor.</i>	" "		6.25	13.50	♂ ad.
1347.	<i>Junco hyemalis.</i>	Apr. 2.		6.56	10.00	♂ ad.
1348.	<i>Passerella iliaca.</i>	" "		7.25	10.81	♀ ad.
1349.	<i>Accipiter fuscus.</i>	" 28.		11.25	20.75	♂ ad.
1350.	<i>Colaptes auratus.</i>	" 21.		12.73	20.20	♂ ad.
1351.	<i>Vireo solitarius.</i>	" 23.		5.50	9.07	♀ ad.
1352.	<i>Dendroica pinus.</i>	" 24.		5.50	8.90	♂ ad.
1353.	<i>Sturnus motacilla.</i>	" "		6.20	10.28	♂ ad.
1354.	<i>Regulus calendula.</i>	" "		4.50	6.80	♂ ad.
1355.	<i>Falco sparverius.</i>	" 25.		10.42	22.38	♂ ad.
1356.	<i>Carpodacus purpureus.</i>	" 28.		6.25	10.20	♂ ad.
1357.	<i>Mniotilta varia.</i>	May 1.		5.30	8.70	♂ ad.
1358.	<i>Sphyrapicus varius.</i>	" 5.		8.62	15.50	♂ ad.

1877.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
1359.	<i>Mniotilta varia.</i>	May 1.		5.30	8.70	♂ ad.
1360.	<i>Corvus ossifragus.</i>	" 7.		16.00	33.20	♀ ad.
1361.	<i>Tringoides macularius.</i>	" 8.		7.75	13.55	♀ ad.
1362.	" "	" "		7.82	13.85	♀ ad.
1363.	<i>Vireo solitarius.</i>	" "		5.50	9.25	♂ ad.
1364.	<i>Geothlypis trichas.</i>	" "		5.42	7.30	♂ ad.
1365.	<i>Setophaga ruticilla.</i>	" 9.		5.35	8.00	♂ ad.
1366.	<i>Vireo solitarius</i>	" "		5.50	9.62	♂ ad.
1367.	" <i>gilvus.</i>	" "		5.96	9.42	♂ ad.
1368.	<i>Sturnus auricapillus.</i>	" "		5.96	9.54	♂ ad.
1369.	" "	" "		6.25	9.80	♂ ad.
1370.	<i>Zonotrichia albicollis.</i>	" "		7.00	9.60	♂ ad.
1371.	<i>Agelaius phoeniceus.</i>	" 11.		7.62	12.50	♀ ad.
1372.	<i>Mniotilta varia.</i>	" "		5.25	8.00	♀ ad.
1373.	<i>Setophaga ruticilla.</i>	" "		5.37	7.75	♂ ad.
1374.	<i>Pipilo erythrophthalmus.</i>	" 12.		8.50	11.48	♂ ad.
1375.	<i>Hirundo horreorum.</i>	" "		6.94	12.53	♀ ad.
1376.	<i>Myiurus carolinensis.</i>	" 11.		8.94	11.50	♂ ad.
1377.	<i>Setophaga ruticilla.</i>	" 12.		5.30	8.25	♂ ad.
1378.	<i>Chrysomitris tristis.</i>	" 11.		5.25	9.06	♂ ad.
1379.	<i>Sturnus motacilla.</i>	" "		6.31	10.19	♀ ad.
1380.	" <i>auricapillus.</i>	" 12.		6.09	9.75	♂ ad.
1381.	<i>Parula americana.</i>	" "		5.00	7.63	♂ ad.
1382.	<i>Geothlypis trichas.</i>	" "		5.25	7.19	♂ ad.
1383.	<i>Parula americana.</i>	" "		4.81	7.62	♂ ad.
1384.	<i>Antrostomus vociferus.</i>	" 14.		9.91	18.70	♂ ad.
1385.	<i>Picus villosus.</i>	" "		9.00	15.12	♀ ad.
1386.	<i>Harporhynchus rufus.</i>	" "		11.25	12.75	-ad.
1387.	<i>Zonotrichia albicollis.</i>	" "		6.87	9.52	♂ juv.
1388.	" "	" "		6.80	9.00	♀ ad.
1389.	<i>Dendroica caerulescens.</i>	" "		5.25	8.00	♂ ad.
1390.	<i>Totanus solitarius.</i>	" 15.		8.53	16.45	ad.
1391.	<i>Cyanospiza cyanea.</i>	" "		5.50	8.61	♂ ad.
1392.	<i>Sturnus naevius.</i>	" "		6.00	9.75	♂ ad.
1393.	<i>Dendroica palmarum.</i>	Apr. 16.		5.63	8.75	♂ ad.
1394.	" "	" "		5.50	8.77	♂ ad.
1395.	<i>Sayornis fuscus.</i>	March 26.		7.06	11.12	♀ ?

1877.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
1396.	<i>Ectopistes migratorius.</i>	March 26.		16.75	24.31	♂ ad.
1397.	<i>Turdus mustelinus.</i>	May 15.		8.50	14.13	-ad.
1398.	" <i>swainsoni.</i>	" "		7.39	12.39	-ad.
1399.	" <i>fuscescens.</i>	" "		7.75	12.00	♂ ad.
1400.	<i>Harporhynchus rufus.</i>	" "		11.25	13.24	-ad.
1401.	<i>Pipilo erythrophthalmus.</i>	" "		8.50	11.41	♂ ad.
1402.	<i>Turdus fuscescens.</i>	" "		7.60	12.00	♀ ad.
1403.	<i>Pyrranga rubra.</i>	" "		7.38	12.00	♀ ad.
1404.	<i>Dendroica maculosa.</i>	" "		5.11	7.88	♂ ad.
1405.	" <i>blackburniae.</i>	" "		5.38	8.40	♂ ad.
1406.	" <i>pennsylvanica.</i>	" "		5.34	7.81	♂ ad.
1407.	" <i>caerulescens.</i>	" "		5.40	7.99	♂ ad.
1408.	<i>Parula americana.</i>	" "		5.82	7.50	♂ ad.
1409.	<i>Dendroica blackburniae.</i>	" 16.		5.25	8.64	♂ ad.
1410.	" "	" "		5.40	8.48	♂ ad.
1411.	" <i>caerulescens.</i>	" "		5.30	7.82	♀ ad.
1412.	" "	" "		5.39	8.25	♂ ad.
1413.	<i>Vireo olivaceus.</i>	" "		6.39	10.25	♂ ad.
1414.	<i>Icterus baltimore.</i>	" "		7.50	11.96	♂ ad.
1415.	<i>Helminthophaga pinus.</i>	" 17.		4.80	7.24	♀ ad.
1416.	<i>Myiodiocetes pusillus.</i>	" "		5.01	6.88	♂ ad.
1417.	<i>Dendroica aestiva.</i>	" "		5.18	7.87	♂ ad.
1418.	<i>Icterus spurius.</i>	" "		7.38	10.50	♂ ad.
1419.	" "	" "		7.50	10.36	♂ ad.
1420.	<i>Goniaphea ludoviciana.</i>	" "		8.25	13.25	♂ ad.
1421.	" "	" "		8.00	12.39	♂ ad.
1422.	<i>Pyrranga rubra.</i>	" "		7.25	11.88	♂ ad.
1423.	<i>Icterus spurius.</i>	" "		7.14	10.04	♀ ad.
1424.	<i>Dendroica blackburniae.</i>	" "		5.32	8.39	♂ ad.
1425.	" "	" "		5.26	8.78	♂ ad.
1426.	<i>Myiodiocetes mitratus.</i>	" "		5.75	7.96	♀ ad.
1427.	" "	" "		5.70	7.98	♀ ad.
1428.	" "	" "		5.65	8.00	♀ ad.
1429.	<i>Dendroica blackburniae.</i>	" 18.		5.35	8.40	♂ ad.
1430.	<i>Myiodiocetes ^{canadensis} mitratus.</i>	" "		5.52	8.23	♂ ad.
1431.	<i>Pyrranga rubra.</i>	" "		7.33	11.79	♂ ad.
1432.	" "	" "		7.50	11.96	♂ ad.

1877.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
1433.	<i>Dendroica virens.</i>	May	18.	5.14	7.65	♀ ad.
1434.	<i>Myiodioctes canadensis.</i>	"	"	5.62	7.55	♀ ad.
1435.	"	"	"	5.65	8.09	♂ ad.
1436.	<i>Turdus swainsoni, aliciae.</i>	"	19.	7.75	13.00	♂ ad.
1437.	<i>Dendroica pennsylvanica.</i>	"	"	5.04	7.33	♀ ad.
1438.	<i>Myiodioctes canadensis.</i>	"	"	5.38	7.75	♀ ad.
1439.	<i>Dendroica castanea.</i>	"	"	5.78	9.49	♂ ad.
1440.	" <i>striata.</i>	"	"	5.64	9.00	♂ ad.
1441.	<i>Cotyle riparia.</i>	"	21.	5.39	10.88	♀ ad.
1442.	<i>Tyrannus carolinensis.</i>	"	"	8.58	14.96	♂ ad.
1443.	<i>Cyanospiza cyanea.</i>	"	"	5.40	8.12	♀ ad.
1444.	<i>Helmatodytes palustris.</i>	"	"	5.25	6.54	♂ ad.
1445.	<i>Pyrrhula rubra.</i>	"	22.	7.13	11.63	♂ ad.
1446.	<i>Agelaius phoeniceus.</i>	"	21.	7.88	12.60	♀ ad.
1447.	<i>Helmatodytes palustris.</i>	"	"	5.25	6.30	♂ ad.
1448.	<i>Nyctiardea grisea, naevia.</i>	"	23.	27.00	46.10	♂ ad.
1449.	"	"	"	25.68	46.15	♀ ad.
1450.	"	"	"	27.00	46.05	♂ juv.
1451.	<i>Petrochelidon lunifrons.</i>	"	"	6.03	12.54	♂ ad.
1452.	"	"	"	5.85	12.50	♂ ad.
1453.	"	"	"	6.55	12.45	♀ ad.
1454.	"	"	"	6.00	12.02	♂ ad.
1455.	<i>Cotyle riparia.</i>	"	"	5.27	10.82	♂ ad.
1456.	<i>Dolichonyx oryzivorus.</i>	"	"	7.15	11.25	♀ ad.
1457.	<i>Icterus spurius.</i>	"	"	7.30	10.50	♂ ad.
1458.	<i>Melospiza palustris.</i>	"	"	6.00	8.14	♂ ad.
1459.	<i>Geothlypis trichas.</i>	"	"	5.27	7.00	♀ ad.
1460.	<i>Chrysomitris tristis.</i>	"	"	5.26	9.07	♂ ad.
1461.	"	"	"	4.82	8.50	♀ ad.
1462.	<i>Corvus americanus.</i>	"	"	19.20	36.75	♂ ad.
1463.	<i>Buteo pennsylvanicus.</i>	"	25.	18.85	36.70	♀ ad.
1464.	<i>Corvus americanus.</i>	"	23.	18.85	36.70	♀ ad.
1465.	"	"	26.	19.30	36.80	♂ ad.
1466.	<i>Myiarchus cinerascens.</i>	"	25.	8.86	13.50	♂ ad.
1467.	<i>Dendroica castanea.</i>	"	"	5.60	8.60	♀ ad.
1468.	<i>Helminthophila vermivorus.</i>	"	"	5.50	8.75	♂ ad.
1469.	"	"	"	5.52	8.50	♀ ad.

1877.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
1470.	<i>Dendroica striata.</i>	May 26.		5.38	9.00	♂ ad.
1471.	<i>Vireo noveboracensis.</i>	"	"	5.28	7.60	♀ ad.
1472.	<i>Bonasa umbellus.</i>	" 30		—	—	pullus.
1473.	"	"	"	—	—	pullus.
1474.	<i>Vireo olivaceus (parent).</i>	"	"	6.06	9.75	♀ ad.
1475.	<i>Myiodioctes mitratus, do.</i>	" 31.		5.70	8.06	♀ ad.
1476.	<i>Dendroica pennsylvanica.</i>	June 4.		5.15	7.70	♀ ad.
1477.	"	"	"	5.15	7.70	♀ ad.
1478.	<i>Bonasa umbellus.</i>	June 18.		6.00	11.00	♂ juv.
1479.	<i>Dendroica discolor.</i>	" 23.		4.88	7.15	♂ ad.
1480.	<i>Geothlypis trichas.</i>	" 21.		5.55	7.48	♂ ad.
1481.	<i>Cyanospiza cyanea.</i>	" 23.		5.63	8.60	♂ ad.
1482.	<i>Spizella socialis.</i>	Apr. 26.		8.25	15.30	♀ ad.
1483.	<i>Cyanospiza cyanea.</i>	June 21.		5.62	8.70	♂ ad.
1484.	<i>Dendroica virens.</i>	" 23.		5.22	7.91	♂ ad.
1485.	<i>Icteria virens.</i>	" 21.		7.75	10.12	—
1486.	<i>Turdus pallasi.</i>	May 2.		7.09	11.30	♀ ad.
1487.	<i>Agelaius phoeniceus.</i>	June 23.		9.60	15.12	♂ ad.
1488.	<i>Zonotrichia albicollis.</i>	Apr. 2.		6.83	9.65	♂ ad.
1489.	<i>Quercus motacilla.</i>	" 30.		6.30	10.50	♂ ad.
1490.	<i>Goniaphea ludoviciana.</i>	July 28.		8.05	12.85	♂ juv.
1491.	<i>Nyctiardea grisea, naevia.</i>	Aug. 27.		—	—	♂ juv.
1492.	<i>Melanerpes erythrocephalus.</i>	Sept. 25.		9.90	17.70	♂ juv.
1493.	"	"	"	9.36	17.25	♀ juv.
1494.	<i>Dendroica palmarum.</i>	Oct. 12.		5.24	7.88	♀.
1495.	<i>Ammodramus caudatus, nelsoni.</i>	"	"	5.50	7.50	♀??
1496.	<i>Accipiter fuscus.</i>	" 11.		11.55	22.00	♂ ad.
1497.	<i>Dendroica palmarum.</i>	" 13.		5.62	8.63	♀
1498.	<i>Turdus pallasi.</i>	"	"	7.13	11.38	♂ (ad?).
1499.	<i>Oedemia americana.</i>	" 23		—	—	♀ ad.
1500.	" <i>fusca</i>	"	"	—	—	♀ ad.
1501.	"	" 26.		22.40	37.90	♂ ad.
1502.	"	"	"	21.00	37.50	♀ ad.
1503.	"	" 27.		—	—	♂ ad.
1504.	<i>Turdus pallasi.</i>	"	"	—	—	♂ juv.
1505.	<i>Mergus serrator.</i>	Nov. 3.		22.80	33.25	♂ juv.
1506.	"	"	"	22.50	33.00	♀

1877-'78.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
1507.	<i>Passerella iliaca.</i>	Nov. 9.		7.00	10.80	♀ ad.
1508.	<i>Sialia sialis.</i>	" "		7.00	12.60	♂ ad.
1509.	<i>Dendroica coronata.</i>	" 15.		5.50	8.75	♀ ad.
1510.	<i>Regulus satrapa.</i>	" 17.		4.24	6.92	♂ ad.
1511.	" "	" "		4.11	6.56	♀ ad.
1512.	<i>Certhia familiaris.</i>	" "		5.62	8.03	♂ ad.
1513.	<i>Passerella iliaca.</i>	" "		7.52	11.39	♂ ad.
1514.	<i>Passer domesticus.</i>	" "		6.50	9.75	♀ ad.
1515.	<i>Larus philadelphia.</i>	" "		14.70	33.00	♂ ad.
1516.	" "	" "		14.52	33.65	♂ ad.
1517.	<i>Dendroica coronata.</i>	" 24.		5.68	9.18	♂ ad.
1518.	<i>Regulus satrapa.</i>	" "		4.21	6.63	♀ ad.
1519.	" "	" "		4.18	6.60	♀ ad.
1520.	<i>Spizella monticola.</i>	" "		6.58	9.65	♂ ad.
1521.	" "	" "		6.25	9.10	♀ ad.
1522.	<i>Ampelis cedrorum.</i>	Dec. 7.		7.27	11.75	♀ ad.
1523.	<i>Carpodacus purpureus.</i>	" "		6.13	10.12	♀ ad.
1524.	" ^{" Locust Grove, Lewis County, N. Y."}	" "		6.25	10.30	♀ ad.
1525.	<i>Melanerpes erythrocephalus.</i>	Dec. 28.		—	—	♂ juv.
1526.	" "	" "		—	—	♂ juv.
1527.	<i>Picus villosus.</i>	" "		—	—	♂ ad.
1528.	<i>Melanerpes erythrocephalus.</i>	" 29.		—	—	♂ juv.
1529.	<i>Picus pubescens.</i>	1878. " "		—	—	♂ ad.
1530.	<i>Melanerpes erythrocephalus.</i>	Jan. 1.		9.95	17.90	♂ ad.
1531.	" "	" "		9.96	17.80	♂ ad.
1532.	<i>Carpodacus ^{Highland Falls, N. Y.} purpureus.</i>	" 18.		6.26	10.09	♀ ad.
1533.	<i>Chrysomitris tristis.</i>	" 12.		5.15	8.89	♂ ad.
1534.	" "	" "		5.31	9.13	♂
1535.	<i>Sialia sialis.</i>	" 10.		—	—	♀ ad.
1536.	<i>Chrysomitris tristis.</i>	" 12.		—	—	♀ ad.
1537.	<i>Carpodacus purpureus.</i>	" 20.		6.50	10.28	♂ ad.
1538.	<i>Spizella monticola.</i>	" 22.		6.28	9.38	♂ ad.
1539.	<i>Parus atricapillus.</i>	" "		5.39	8.10	♀ ad.
1540.	" "	" "		5.36	8.36	♂ ad.
1541.	<i>Regulus satrapa.</i>	" 24.		—	—	♂ ad.
1542.	" "	" 25.		4.24	6.64	♀ ad.
1543.	" "	" "		4.20	7.04	♂ ad.

1878.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
1544.	<i>Dendroica coronata.</i>	Jan. 25.		5.75	9.08	♂ ad.
1545.	<i>Spizella monticola.</i>	" "		6.41	9.52	♂ ad.
1546.	" "	" "		6.50	9.50	♂ ad.
1547.	" "	" "		6.51	9.50	♂ ad.
1548.	<i>Chrysomitris tristis pinus.</i>	Feb. 6.		5.06	8.76	♀ ad.
1549.	" <i>tristis.</i>	" "		4.93	8.75	♂ ad.
1550.	<i>Dendroica coronata.</i>	" "		5.61	9.81	♂ ad.
1551.	<i>Sialia sialis.</i>	" 5.		7.13	12.34	♂ ad.
1552.	" "	" 9.		7.04	12.82	♂ ad.
1553.	<i>Haliaeetus leucocephalus.</i>	" 15.		32.87	84.10	♂ ad.
1554.	<i>Carpodacus purpureus.</i>	" 21.		6.42	10.52	♂ ad.
1555.	" "	" "		6.13	10.16	♂
1556.	" "	" "		6.06	9.90	♀ ad.
1557.	<i>Picus pubescens.</i>	" "		6.87	12.00	♀ ad.
1558.	<i>Ampelis cedrorum.</i>	" 23.		7.31	11.79	♂ ad.
1559.	" "	" "		7.28	11.86	♂ ad.
1560.	" "	" "		7.49	11.98	♀ ad.
1561.	<i>Parus atricapillus.</i>	" "		5.37	8.00	♂ ad.
1562.	<i>Ampelis cedrorum.</i>	" "		7.37	12.05	♂ ad.
1563.	" "	" "		7.20	11.81	♀ ad.
1564.	<i>Carpodacus purpureus.</i>	" "		6.18	9.99	♀ ad.
1565.	" "	" "		6.38	10.15	♀ ad.
1566.	" "	" "		6.25	10.25	♀ ad.
1567.	<i>Picus pubescens.</i>	" "		6.87	12.10	♀ ad.
1568.	<i>Ampelis cedrorum.</i>	" "		7.04	11.80	♂ (ad.?).
1569.	<i>Passerella iliaca.</i>	" 28.		6.97	10.62	♀ ad.
1570.	<i>Carpodacus purpureus.</i>	March 6.		6.26	10.25	♂ ad.
1571.	" "	" "		6.27	9.92	♂ ad.
1572.	" "	" "		6.19	10.12	♀ ad.
1573.	" "	" "		6.07	9.83	♀ ad.
1574.	<i>Passerella iliaca.</i>	" "		7.15	11.00	♂ ad.
1575.	<i>Passer domesticus.</i>	" "		6.23	9.50	♂ ad.
1576.	" "	" "		6.40	9.70	♀ ad.
1577.	<i>Melospiza melodia.</i>	" "		6.13	8.59	♂ ad.
1578.	<i>Ampelis cedrorum.</i>	" "		7.23	11.40	♂ ad.
1579.	<i>Passer domesticus.</i>	" "		6.65	9.88	♂ ad.
1580.	" "	" "		6.61	10.00	♂ ad.

1878.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
1581.	<i>Ampelis cedrorum.</i>	March 6.		7.28	12.05	♀ ad.
1582.	"	"	"	7.38	11.62	♂ ad.
1583.	<i>Sitta carolinensis.</i>	" 9.	"	6.12	11.18	♂ ad.
1584.	"	"	"	6.00	10.94	♀ ad.
1585.	<i>Chrysomitris tristis.</i>	" 2.	"	5.15	8.88	♂ ad.
1586.	<i>Ampelis cedrorum.</i>	" 13.	"	7.18	12.00	♀ ad.
1587.	"	"	"	7.25	11.81	♀ ad.
1588.	"	"	"	7.02	11.77	♀ ad.
1589.	"	"	"	7.35	11.84	♂ ad.
1590.	"	"	"	7.15	12.04	♂ ad.
1591.	"	"	"	7.10	11.85	♂ ad.
1592.	<i>Carpodacus purpureus.</i>	" 15.	"	6.29	10.39	♂ ad.
1593.	<i>Picus pubescens.</i>	"	"	7.02	12.25	♂ ad.
1594.	<i>Ampelis cedrorum.</i>	" 13.	"	7.25	11.62	♂ ad.
1595.	"	"	"	7.19	11.60	♀ ad.
1596.	"	"	"	7.14	11.75	♀ ad.
1597.	"	"	"	7.25	11.85	—
1598.	"	"	"	—	—	♀
1599.	"	"	"	—	—	♀
1600.	<i>Turdus migratorius.</i>	" 15.	"	10.10	15.96	♂ ad.
1601.	<i>Ampelis cedrorum.</i>	" 11.	"	—	—	♀
1602.	"	"	"	—	—	♀
1603.	<i>Carpodacus purpureus.</i>	" 26.	"	6.24	10.38	♀ ad.
1604.	<i>Sayornis fuscus.</i>	" 25.	"	7.18	11.38	♂ ad.
1605.	"	"	"	7.26	11.80	♂ ad.
1606.	<i>Turdus migratorius.</i>	" 30.	"	9.83	15.50	♀ ad.
1607.	<i>Junco hyemalis.</i>	" 29.	"	6.35	10.11	♂ ad.
1608.	<i>Passerella iliaca.</i>	Apr. 1.	"	7.18	11.25	♂ ad.
1609.	<i>Sayornis fuscus.</i>	"	"	7.24	11.52	♂ ad.
1610.	<i>Carpodacus purpureus.</i>	" 4.	"	6.10	—	♂ ad.
1611.	<i>Sayornis fuscus.</i>	" 3.	"	7.27	11.32	♂ ad.
1612.	<i>Passerella iliaca.</i>	" 4.	"	7.65	11.78	♂ ad.
1613.	<i>Dendroica coronata.</i>	"	"	5.71	9.32	♂ ad.
1614.	<i>Melospiza melodia.</i>	" 10.	"	6.25	8.70	♂ ad.
1615.	<i>Colaptes auratus.</i>	" 11.	"	11.80	20.40	♀ ad.
1616.	<i>Molothrus ater.</i>	"	"	7.75	13.30	♂ ad.
1617.	<i>Turdus migratorius.</i>	" 12.	"	9.70	15.40	♀ ad.

1878.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
1618.	<i>Turdus pallasi.</i>	Apr. 16.		7.30	11.53	♂ ad.
1619.	<i>Spizella socialis.</i>	"	"	5.38	8.80	♂ ad.
1620.	<i>Sturnella magna.</i>	Fishkill, N.Y.	" 18.	10.60	16.45	♂ ad.
1621.	"	"	"	11.12	16.75	♂ ad.
1622.	<i>Agelaius phoeniceus.</i>	do	"	10.13	15.53	♂ ad.
1623.	<i>Sturnella magna.</i>	Putnam Co.	" 20.	9.57	14.86	♀ ad.
1624.	<i>Siurus motacilla.</i>	"	"	—	—	♂ ad.
1625.	<i>Regulus calendula.</i>	"	"	—	—	♂ ad.
1626.	<i>Sphyrapicus varius.</i>	"	"	8.56	15.00	♀ ad.
1627.	<i>Molothrus ater.</i>	"	"	7.38	12.53	♀ ad.
1628.	"	"	"	8.07	13.75	♂ ad.
1629.	"	"	"	7.87	13.50	♂ ad.
1630.	"	"	"	8.12	13.58	♂ ad.
1631.	<i>Junco hyemalis.</i>	N.Y.	" 22.	6.48	10.16	♂ ad.
1632.	<i>Siurus motacilla.</i>	do	" 23.	6.27	10.54	♂ ad.
1633.	<i>Agelaius phoeniceus.</i>	do	" 24.	9.50	15.60	♂ ad.
1634.	<i>Dendroica palmarum.</i>	do	" 24.	5.42	8.60	♂ ad.
1635.	<i>Sphyrapicus varius.</i>	do	" 25.	8.80	15.85	♀ ad.
1636.	<i>Siurus motacilla.</i>	do	"	6.56	10.77	♂ ad.
1637.	"	do	"	6.07	10.40	♂ ad.
1638.	" <i>naevius.</i>	do	" 26.	6.04	9.40	♂ ad.
1639.	<i>Pipilo erythrophthalmus.</i>	"	"	8.40	11.10	♂ ad.
1640.	<i>Hirundo thoracorum.</i>	" 27.	"	7.32	13.10	♂ ad.
1641.	"	"	"	7.06	12.90	♂ (juv?)
1642.	<i>Siurus naevius.</i>	"	"	6.14	9.52	♂ ad.
1643.	<i>Passerculus savanna.</i>	"	"	5.78	9.12	♂ ad.
1644.	"	"	"	5.88	9.40	♂ ad.
1645.	<i>Dendroica palmarum.</i>	"	"	5.27	8.06	♀ ad.
1646.	<i>Setophaga ruticilla.</i>	"	"	5.60	7.92	♂ ad.
1647.	<i>Pipilo erythrophthalmus.</i>	"	"	8.24	10.65	♀ ad.
1648.	<i>Chrysomitris tristis.</i>	" 29.	"	5.19	8.90	♂ ad.
1649.	<i>Parus atricapillus.</i>	"	"	5.30	7.97	♀ ad.
1650.	"	"	"	5.28	8.00	♂ ad.
1651.	<i>Melospiza palustris.</i>	" 30.	"	5.87	7.73	♀ ad.
1652.	" <i>melodia.</i>	"	"	6.46	8.66	♂ ad.
1653.	<i>Passerculus savanna.</i>	"	"	5.83	9.20	♂ ad.
1654.	<i>Melospiza melodia (parent).</i>	"	"	6.30	8.40	♀ ad.

1878.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
1655.	<i>Turdus migratorius.</i>	Apr. 30.		10.10	16.25	♂ ad.
1656.	<i>Harporhynchus rufus.</i>	" "		11.85	13.25	♂ ad.
1657.	<i>Cyanurus cristatus.</i>	" "		11.50	16.40	♀ ad.
1658.	<i>Pipilo erythrophthalmus.</i>	" "		8.50	11.87	♂ ad.
1659.	<i>Molothrus ater.</i>	" "		7.90	13.65	♂ ad.
1660.	<i>Troglodytes domesticus.</i>	May 2.		5.10	6.42	♀ ad.
1661.	<i>Melanerpes erythrocephalus.</i>	" "		9.60	18.00	♀ ad.
1662.	<i>Myiodioctes mitratus.</i>	" 4.		—	—	♂ ad.
1663.	" "	" "		5.67	8.38	♂ juv.
1664.	<i>Helmatodytes palustris.</i>	" "		5.35	6.62	♂ ad.
1665.	<i>Trindoides macularius.</i>	" "		7.95	13.50	♀ ad.
1666.	<i>Geothlypis trichas.</i>	" "		5.30	7.30	♂ ad.
1667.	<i>Coccyzus gramineus.</i>	" 3.		5.30	7.30	♂ ad.
1668.	<i>Molothrus ater.</i>	" 4.		—	—	♂ ad.
1669.	" "	" "		—	—	♀ ad.
1670.	<i>Icterus baltimore</i>	" "		—	—	♂ ad.
1671.	<i>Turdus mustelinus.</i>	" "		8.20	13.75	♂ ad.
1672.	" "	" "		8.50	13.75	♂ ad.
1673.	<i>Miniotilta varia.</i>	" "		5.40	8.53	♀ ad.
1674.	<i>Coccyzus americanus.</i>	" 6.		11.88	16.75	♀ ad.
1675.	" "	" "		12.40	17.35	♀ ad.
1676.	" "	" "		12.00	16.75	♀ ad.
1677.	<i>Siurus motacilla.</i>	" 7.		6.25	10.38	♀ ad.
1678.	<i>Vireo olivaceus.</i>	" "		6.33	10.15	♂ ad.
1679.	<i>Setophaga ruticilla.</i>	" "		5.38	7.90	♂ ad.
1680.	<i>Myioidibetes mitratus</i>	" "		5.75	8.47	♂ ad.
1681.	" "	" "		5.75	8.55	♂ ad.
1682.	" "	" "		5.64	8.40	♂ ad.
1683.	" "	" "		5.62	8.25	♀ ad.
1684.	<i>Sphyrapicus varius.</i>	" "		8.62	15.70	♀ ad.
1685.	<i>Goniaphea ludoviciana.</i>	" 9.		8.15	13.40	♂ ad.
1686.	<i>Helminthophaga chrysoptera</i>	" 11.		5.25	8.35	♂ ad.
1687.	" <i>ruficapilla.</i>	" 10.		—	—	♀ ad.
1688.	<i>Myiodioctes mitratus.</i>	" 10.		—	—	♂ ad.
1689.	" "	" "		—	—	♂ ad.
1690.	<i>Icterus baltimore.</i>	" "		—	—	♂ ad.
1691.	<i>Vireo flavifrons.</i>	" "		—	—	♂ ad.

1878.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
1692.	<i>Dendroica virens.</i>	May 10.		—	—	♂ ad.
1693.	<i>Turdus mustelinus.</i>	" "		—	—	♂ ad.
1694.	" <i>swainsoni.</i>	" "		—	—	♂ ad.
1695.	" <i>mustelinus.</i>	" "		—	—	♂ ad.
1696.	<i>Pyrranga rubra.</i>	" "		—	—	♂ ad.
1697.	<i>Siurus motacilla.</i>	" 11.		—	—	♂ ad.
1698.	<i>Pyrranga rubra.</i>	" 10.		—	—	♂ ad.
1699.	<i>Icterus baltimore.</i>	" "		—	—	♂ juv.
1700.	<i>Agelaius phoeniceus</i>	" 11.		—	—	♂ juv.
1701.	" "	" "		—	—	♂ ad.
1702.	<i>Sialia sialis (parent).</i>	" 13.		—	—	♀ ad.
1703.	<i>Trochilus colubris.</i>	" 15.	3.78	4.12		♂ juv.?
1704.	<i>Myiodioctes mitratus.</i>	" 14.	5.80	8.60		♂ ad.
1705.	" "	" 15.	5.85	8.10		♀ ad.
1706.	<i>Dendroica caerulescens.</i>	" "	5.27	8.16		♂ ad.
1707.	<i>Helminthophaga chrysoptera</i>	" 16.	5.02	8.15		♂ ad.
1708.	<i>Contopus borealis.</i>	" "	7.37	12.88		♂ ad.
1709.	<i>Icteria virens.</i>	" 15.	7.40	9.90		♂ ad.
1710.	<i>Picus pubescens (parent).</i>	" 17.	—	—		♀ ad.
1711.	<i>Vireo flavifrons.</i>	" 16.	5.75	9.80		♂ ad.
1712.	<i>Dendroica virens.</i>	" 15.	—	—		♂ ad.
1713.	<i>Cyanospiza cyanea.</i>	" 16.	—	—		♂ ad.
1714.	" "	" "	—	—		♂ juv.
1715.	<i>Coturniculus passerinus.</i>	" 18.	5.35	8.30		♀ ad.
1716.	" "	" "	5.30	8.60		♂ ad.
1717.	<i>Dendroica aestiva.</i>	" "	5.15	7.80		♂ ad.
1718.	" "	" "	5.10	7.80		♂ ad.
1719.	<i>Nyctiardea grisea, naevia</i>	" "	25.00	45.25		♀ ad.
1720.	<i>Oelichomyx rufirostris</i>	" "	7.65	12.10		♂ ad.
1721.	<i>Icterus baltimore.</i>	" "	7.65	11.85		♀ ad.
1722.	<i>Turdus mustelinus.</i>	" 15.	—	—		♂ ad.
1723.	<i>Contopus virens.</i>	" 20.	5.60	10.95		♂ ad.
1724.	<i>Hirundo horreorum.</i>	" "	7.35	12.80		♂ ad.
1725.	" "	" "	—	—		♂ ad.
1726.	" "	" "	—	—		♂ ad.
1727.	<i>Pyrranga rubra.</i>	" 21.	7.40	12.25		♂ ad.
1728.	<i>Myiarchus crinitus.</i>	" "	9.05	13.45		♂ ad.

1878.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
1729.	<i>Icteria virens.</i>	May 21.		7.50	10.00	♂ ad.
1730.	<i>Myiodiocetes mitratus.</i>	" "		5.75	8.10	♀ ad.
1731.	" "	" "		6.00	8.75	♂ ad.
1732.	<i>Turdus migratorius.</i>	" "		10.00	15.67	♀ ad.
1733.	<i>Icterus spurius.</i>	" "		6.85	10.05	♂ ad.
1734.	<i>Helminthophorus vermicivorus.</i>	" 22.		5.60	8.62	♀ ad.
1735.	<i>Tachycineta bicolor.</i>	" "		—	—	♂ ad.
1736.	<i>Telmatorhynchus palustris.</i>	" "		—	—	♂ ad.
1737.	<i>Dendroica castanea.</i>	" "		5.85	9.30	♂ ad.
1738.	" <i>blackburniae.</i>	" "		5.28	8.35	♂ ad.
1739.	<i>Empidonax flaviventris.</i>	" "		5.45	8.20	♀ ad.
1740.	<i>Dendroica blackburniae.</i>	" "		5.25	8.40	♂ ad.
1741.	<i>Chaetura pelagica.</i>	" 23.		5.35	12.37	♀ ad.
1742.	<i>Icterus spurius.</i>	" "		7.55	10.60	♂ juv.
1743.	<i>Dendroica blackburniae.</i>	" "		5.20	8.40	♂ ad.
1744.	<i>Tyrannus carolinensis.</i>	" "		8.75	14.80	♂ ad.
1745.	<i>Vireo olivaceus.</i>	" 25.		6.40	10.20	♂ ad.
1746.	<i>Mniotilta varia.</i>	" "		5.35	8.55	♀ ad.
1747.	<i>Ampelis cedrorum.</i>	" "		7.30	12.00	♀ ad.
1748.	<i>Agelaius phoeniceus.</i>	" "		9.30	15.00	♂ ad.
1749.	" "	" "		9.75	15.60	♂ ad.
1750.	" "	" 28.		8.12	12.85	♀ ad.
1751.	<i>Mimus carolinensis.</i>	" "		8.95	11.65	♂ ad.
1752.	<i>Pyrranga rubra.</i>	" "		7.30	11.85	♂ ad.
1753.	<i>Turdus swainsoni, alciac.</i>	" "		7.55	12.90	♀ ad.
1754.	<i>Hirundo horreorum.</i>	" 10		—	—	♀ ad.
1755.	" "	" "		—	—	♀ ad.
1756.	<i>Dendroica pennsylvanica.</i>	" 3.		—	—	♂ ad.
1757.	<i>Coccyzus erythrophthalmus.</i>	" 27.		—	—	♂ pullus.
1758.	<i>Pyrranga rubra.</i>	" "		—	—	♂ ad.
1759.	<i>Turdus migratorius.</i>	June 12.		—	—	pullus.
1760.	<i>Icterus baltimore.</i>	" "		—	—	pullus.
1761.	<i>Ampelis cedrorum (parent).</i>	" "		7.50	12.22	♀ ad.
1762.	<i>Myiodiocetes mitratus.</i>	" 15.		5.30	7.80	♀ ad.
1763.	<i>Harporhynchus rufus.</i>	July 6.		—	—	juv.
1764.	<i>Oetophaga ruticilla.</i>	" "		—	—	juv.
1765.	<i>Mniotilta varia.</i>	" "		—	—	♀ ad.

1878.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
1766.	<i>Myiodiocetes mitratus.</i>	July 8.		—	—	juv.
1767.	" "	" "		—	—	juv.
1768.	<i>Sayornis fuscus (parent).</i>	May 3.		—	—	♀ ad.
1769.	<i>Picus pubescens.</i>	Feb. 23.		6.87	12.10	♀ ad.
1770.	<i>Troglodytes aëdon (= domesticus).</i>	Aug. 9.		—	—	juv.
1771.	<i>Tyrannus carolinensis.</i>	" 26.		8.30	10.20	♀ juv.
1772.	<i>Sterna fuliginosa.</i>	Sep. 13.		17.05	34.70	♀ ad.
1773.	<i>Ardea herodias.</i>	" 23.		43.75	69.50	♀ juv.
1774.	<i>Podilymbus podiceps.</i>	" "		13.10	21.40	♀
1775.	<i>Dendroica caerulescens.</i>	" "		5.38	8.25	♂ ad.
1776.	<i>Parula americana.</i>	" "		4.80	7.65	♀
1777.	<i>Chaetura pelagica.</i>	" "		5.38	12.30	—
1778.	<i>Turdus swainsoni, alciac.</i>	" 28.		7.55	12.65	♂
1779.	" "	" "		7.40	11.88	♀
1780.	<i>Amphispiza varius.</i>	" "		8.30	15.15	♀ juv.
1781.	<i>Podiceps griseigena, holbolli.</i>	Oct. 4.		20.75	34.90	♂
1782.	" "	Oct. 7.		22.35	34.25	♂ (ad?)
1783.	<i>Aix sponsa.</i>	Aug. Aug. 21.		18.10	28.75	♀ ad.
1784.	<i>Turdus pallasi.</i>	" 18.		7.73	11.90	♂
1785.	<i>Zonotrichia albicollis.</i>	" "		6.90	9.40	♀ ad.
1786.	<i>Dendroica coronata.</i>	" "		5.55	8.50	—
1787.	<i>Spizella monticola.</i>	" 26.		6.47	9.25	♂ ad.
1788.	<i>Sitta carolinensis.</i>	" "		6.12	11.05	♂ ad.
1789.	" "	" "		6.35	11.20	♀
1790.	<i>Spatula clypeata.</i>	Nov. 9.		20.10	32.00	♂ juv.
1791.	<i>Trismaturus rubida.</i>	" 15.		14.80	21.70	♀ ad.
1792.	<i>Picus pubescens.</i>	" 17.		6.65	11.90	♀ ad.
1793.	<i>Parus atricapillus.</i>	" "		5.45	8.25	♂
1794.	" "	" "		5.60	8.40	♂
1795.	<i>Pinicola enucleator.</i>	" 30.		9.35	14.00	♂ juv.
1796.	" "	" "		9.20	13.75	♀
1797.	" "	" 29.		9.50	14.50	♂ juv.
1798.	" "	" "		8.95	13.45	♀
1799.	" "	" "		9.00	14.05	♂ juv.
1800.	" "	" "		8.62	13.62	♂ juv.
1801.	" "	" "		8.90	13.90	♀
1802.	<i>Nisus fuscus (W. L. Bruce).</i>	" 13.		14.00	25.75	♀ juv.

1878-79.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
1803.	<i>Spizella monticola</i> .	Nov. 29.	Highland Falls, N.Y.	6.55	9.65	♂ ad.
1804.	<i>Cyanurus cristatus</i> .	" 30.	" "	11.50	17.00	♀ ad.
1805.	"	Dec. 27.	" "	11.75	16.62	♀ ad.
1806.	"	"	" "	11.65	16.80	♀ ad.
1807.	<i>Sitta carolinensis</i> .	"	" "	6.25	11.30	♂ ad.
1808.	<i>Bonasa umbellus</i> .	" 30.	" "	17.12	23.35	♂ ad.
1809.	<i>Egiothus linaria</i> .	"	" "	5.52	9.10	♂ ad.
1810.	"	"	" "	5.35	8.65	♂ juv.
1811.	"	"	" "	5.35	8.80	♂ ad.
1812.	"	"	" "	5.50	8.80	♂ ad.
1813.	"	"	" "	5.60	9.35	♂ juv.
1814.	"	"	" "	5.50	9.05	♂ juv.
1815.	"	"	" "	5.60	8.85	♂ juv.
1816.	"	"	" "	5.15	8.45	♀ ad.
1817.	"	"	" "	5.40	8.80	♂ juv.
1818.	"	"	" "	5.12	8.25	♀ ad.
1819.	<i>Mareca penelope</i> .	Jan 25.	Washington Market, N.Y.	19.50	33.60	♂ ad.
1820.	<i>Pediorcetes phasianellus</i> .	Feb'y. 1.	" "	17.75	27.25	♀ ad.
1821.	"	"	" "	—	—	♀ ad.
1822.	<i>Comophila alpestris</i> .	"	Fort Montgomery, N.Y.	8.05	13.75	— ad.
1823.	<i>Picus pubescens</i> .	March 3.	" "	6.70	11.65	♂ ad.
1824.	* <i>Lanius borealis</i> .	Feb. 26.	Putnam Co. N.Y.	10.15	14.15	♀ juv.
1825.	<i>Bonasa umbellus</i> .	March 5.	" "	—	—	♀ juv.
1826.	* <i>Quiscalus purpureus</i> .	" 12.	" "	12.50	17.75	♂ ad.
1827.	<i>Turdus migratorius</i> .	"	" "	10.80	17.10	♀ ad.
1828.	* <i>Melospiza melodia</i> .	" 13.	" "	6.40	8.50	— ad.
1829.	* <i>Picus villosus</i> .	" 14.	" "	9.35	15.25	♂ ad.
1830.	"	"	" "	8.95	15.35	♀ ad.
1831.	<i>Pyrranga rubra</i> .	July 21, 1875.	" "	7.12	11.50	♂ ad.
1832.	<i>Ammodramus haemastica</i> .	Oct. 2, 1874.	" "	—	—	♀ ad.
1833.	<i>Icterus baltimore</i> .	May 18, 1876.	" "	—	—	♂ ad.
1834.	<i>Helminthorus vermivorus</i> .	July 23, 1875.	" "	6.37	8.62	♀ ad.

It is to be understood that when not otherwise stated, all of the above specimens were shot by myself somewhere in the neighborhood of my house; all of

* See p. 91 for its measurements.

1879.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
them in the Hudson Highlands. Some were shot on the opposite bank of the river, in Putnam county; others to the southward, in Rockland. A few were taken higher up the Hudson, in Ulster & Putnam counties; and not a few were obtained upon the islands of Constitution, Cousack, and Jona, in the Hudson River. Those taken at a greater distance are so described in the headings.						

Saturday, March 22, 1879.

Highland Falls, N.Y.

I left the College of Phys. & Surgeons, New York, on the 24th of February last. I found the Hudson completely ice-bound. All of the air-holes were closed tight. A sleigh with one horse was awaiting at Garrison, which brought me and my heavy baggage across the river.

I had hoped to find the Pine Grosbeaks, Crossbills, Lesser Redpolls, Pine Finches and Snow Buntings still here; but I have seen none of those birds since my return.

I walked across the river on the 26th, and walked through Sec'y Fish's place to the public road, opposite to the Osborn's; thence up to the Constitution meadows, Cold Spring, and Constitution Island. The only noteworthy birds, were *Buteo lineatus*, of which I saw several, and *Lanius borealis*. I shot a young female of the latter species, on the meadows, after a long chase, and several unsuccessful shots. Walking down upon the railroad at Mr. Fish's place, I started a *Junco hyem.* from the track a few paces in front of me. A sprike dashed down from the ledge above, in hot pursuit. The Snowbird made every effort to escape, doubling and twisting, and pecking.

1879.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

most piteously as it endeavored to escape by dashing into a clump of hemlocks; but the Shrike followed closely all of its windings till at last the terrified creature flew on top of the ledge, followed closely by its enemy. I could not see the result, but have no doubt that it ended in a tragedy.

On Sunday, March 1st, I took a walk in the deep snow. I saw three Ruffed Grouse, *Bonasa umbellus*, below Mr. Pell's, among the cedars. I saw the first one when at quite a distance. I approached cautiously till within a few paces, where I had an excellent opportunity to observe its actions. At first it stood erect, with neck outstretched but as I remained quiet, it seemed less alarmed, and disposed to get away without taking to flight. Spreading its tail, and with drooping wings, it sallied softly through the snow to a tree or stump. I nearly missed which it took refuge till another favorable opportunity seemed to occur. In the mean time another Grouse was seen to start from a nearer situation, and seeming as little disposed to flight as its predecessor. They skulked behind stones and trees, slowly retreating. Their tails were widely spread, their ruffs erect, and every attitude betokening great solicitude and alarm; also indicated by their scared, clucking notes. At every step their tails were alternately erected and depressed in a jerky manner, while their heads were oscillated spasmodically in a longitudinal direction. Thus they moved away in divergent directions, occasionally encountering a small ledge, which they scaled by means of their wings,

1879.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

much as a hen attains her perch. At length, much to my surprise, a third bird started from a bunch of tall grass, where it had remained concealed, and with a loud clucking note started in a direct line in the same direction in which the others had by this time disappeared. It seemed greatly alarmed, and made little attempt at concealment, but ran clucking after its companions, seeming to prefer this means of escape, as it had been successfully adopted by its comrades.

It was a very pleasant sight to see these handsome birds under such favorable circumstances, and exhibiting so much animation. Such opportunities are not frequent.

Early in March there were heavy rains, and considerable foggy and windy weather. On the morning of the 10th, I heard the Herring Gulls screaming over head. I concluded that the ice must have broken up in the Hudson, and, on going to see, found that I was correct.

No Robins or Bluebirds were here on my first arrival. The first Bluebirds were seen in Putnam county on March 3rd. The first Robin two or three days later, at the Forest-o-Dean mines, where I spent two days fox hunting.

No Purple Finches were seen; but I think I heard one a few days since.

I have not seen or heard a Cedar Bird since my return from New York.

<i>Truscalus purpureus</i> .	First seen	March 12th.
<i>Passerella iliaca</i> .	" "	" "
<i>Sayornis fusca</i> .	" "	" 13th

1879.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

The Phoebe have only uttered their sharp "chip" till this morning (March 22nd) when I heard, for the first time their characteristic note.

The Goshawk, Astur atricapillus, I have seen twice. - March 13th & 20th.

Junco hyemalis. The Snowbirds are twittering everywhere, in great abundance. Occasionally one of them essays a song, which is executed with really good effect, being very pure & sweet. The Fox Sparrow Passerella iliaca has been singing for some days past.

March 24th.

Pandion haliaetus: "Saw the first [Osprey?] to-day." [Fide C. Dingley.]

Amphisp. cedrorum. First seen yesterday.

A flock remained in the trees about the house all of to-day. I observed that they separated into pairs, and they are probably mated already. Two would separate themselves from the mass of their brethren, and edge toward each other, making alternate advances, applying their bills together - doubtless a means of caressing.

Shot two Muskrats on the Consook Marsh. Ducks are quite numerous.

A number of Eagles (Haliaetus leucocephalus) are inhabiting the Garrisons Cove, as Ed, the Expressman, informed me this morning. I intend going up there to-morrow, as there are said to be numbers of Ducks in the Cove.

1879.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

1835. Mergus merganser. March 20 24.00 33.00 ♀ ad. Wing 9.00; tail, 4.35; culmen, 1.92; Tarsus, 2.00. Irides deep brown.

Shot on Consook Island. H. R.

1836. Stelgidopteryx serripennis. May 1872. ♀ ad.

I have neglected to record this, the only specimen of the Rough-winged Swallow that I have ever captured, till now. I took her off her nest, in a bank beside the Parry Pond, along with one of the four eggs that the nest contained. Mr. Fred Fitchett secured the other three eggs.

See Bulletin of Nuttall Club, Vol. III, p. 46, 1878.

Also see my Zoological Journal, No 67, Set 2, p. 34.

Tuesday, March 25th.

Highland Falls, N.Y.

To-day went ducking on the river. Saw a large number of Ducks, and several White-headed Eagles were seen at the Garrisons Cove. One fine fellow sat on the top of a large dead tree, just north of Garrisons. We landed but failed to get him for the same reason that Simple Simon could not shoot wild Ducks.

We got nothing; stayed all day.

Wednesday, March 26th.

H. N. Y.

Went ducking (three guns) down to Lona Island. Visited "Snake Hole" and various places. The following species of Duck were noticed yesterday and to-day:

Anas boschas, H. Mallard.

Fuligula marila Greater Blackhead.

" ferna, americana. Redhead.

" vallisneria Canvas-back.

Bucephala clangula Golden-eye

1879.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

Bucephala albeola. Buff-head.
Harelda glacialis. Long-tailed Duck.
Mergus merganser. Gooseander.
 " *sertator*. Red-breasted Merganser.

A Goose, *Branta canadensis*, was shot by Reed, last Sunday (March 23rd), in front of Haven's Landing.

Larus argentatus is the only Gull seen.

March 26.

Lona Island, H. Riv.

1837. *Turdus migratorius*, L. Robin. 10.04 16.40 ♂ ad.
 Wing, 5.10; tail, 4.15. Iris hazel.

1838. *Ampelis cedrorum*. Cedar Bird. 7.35 11.75 ♀ ad.
 Wing, 3.70; tail, 2.50.

1839. *Picus pubescens*, Linn. Downy W. 7.00 11.95 ♂ ad.
 Wing, 3.65; tail, 2.80.

March 28.

Hudson River, at Cornwall.

1840. *Anas obscura* Dusky Duck 24.50 38.50 ♂ ad.
 Wing, 11.00; tail, 4.35; culmen, 2.33; gape, 2.65; tarsus, 2.00; middle toe, 2.17; its claw, .44.

From Josh Ward. Iris hazel. Bill with base of mandible, bright greenish yellow, shading into olive-green towards tip; nail blackish.

Point of mandible black, like nail of maxilla; resediment of mandible, olive. Legs and toes uniform red; webs blackish red.

1841. *Fuligula marila* Blue-bill. 19.65 33.75 ♂ ad.
 Wing, 8.80; tail, 2.85; culmen, 1.69;

gape, 2.23; tarsus, 1.58; middle toe, 2.50; its claw, .38. Membranes between rami of mandible, purplish. Iris bright orange. Bill, pale blue on maxilla; nail blk. lower mandible black, with edges blue (like

1879.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

maxilla) nearly to point; resediment mottled with purplish blue. Legs & feet, bluish ash; webs, blackish. Shot by Josh Ward, at Cornwall.

April 2.

Highland Falls, N.Y.

Agelaius phoeniceus Red-winged Blackbird.

First seen March 29th.

Ectopistes migratoria Wild Pigeon. A large flock was seen twice, to-day, flying over the woods at the south, though they were not seen to alight there.

1842. *Passerella iliaca*. Fox Sparrow. 7.35 10.80 ♂ ad.
 Wing, 3.25; tail, 2.85. culmen, .50; tarsus, .95.

1843. *Ampelis cedrorum*. Cedar Bird. 7.50 12.00 ♀ ad.
 Wing, 3.77; tail, 2.50; culmen, .39; from nostril, .27; gape, .70; tarsus, .70. Rectrices, reddish at tip.

1844. *Ampelis cedrorum*. Cedar Bird. 6.95 11.25 ♀ juv.
 Wing, 3.55; tail, 2.35; culmen, .40; from nostril, .25; gape, .71; tarsus, .64. Tail just acquiring the yellow tip; greenish, growing bright yellow at tip.

April 3, 1879.

H. Riv.

1845. *Passerella iliaca* (Murren). Fox Sp. 7.50 11.35 ♂ ad.
 Wing, 3.40; tail, 2.90; culmen, .50; tarsus, .95; middle toe, .65; its claw, .31.

1846. *Passerella iliaca* (Murren). Fox Sparrow. 7.08 11.05 ♂ ad.
 Wing, 3.35; tail, 2.90; culmen, .50; tarsus, .94; middle toe, .60; its claw, .30.

1847. *Ectopistes migratoria* (Linn.) W. Pigeon. 15.60 23.85 ♀ ad.
 Wing, 7.75; tail, 7.00; culmen, .74; gape, 1.10; tarsus, 1.08; middle toe, 1.14; its claw, .37. Iris, orange-red. Bare space about eye, purplish blue. Bill, black. Legs & feet, light red.

1879.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
1848	<i>Amphelis cedrorum</i> (Vieill.). Cedar Bird.	April 4, 1879. H. M. J.		7.30	12.05	♂ ad.
	Wing, 3.87; tail, 2.45; culmen, .41; tarsus, .70; middle toe, .61; its claw, .24.					
1849.	<i>Ectopistes migratoria</i> (Linné). Wild Pigeon.			16.75	24.00	♂ ad.
	Wing, 7.80; tail, 7.75; culmen, .75; gape, 1.18; tarsus, 1.10; middle toe, 1.14; its claw, .35. Colors the same as No. 1847.					
1850.	<i>Ectopistes migratoria</i> (Linné). W. Pigeon.			16.35	24.00	♀ ad.
	Wing, 7.75; tail, 7.60; culmen, .72; tarsus, 1.05; middle toe, 1.10; its claw, .35; gape, 1.06. Colors same as No. 1847.					
1851.	<i>Ectopistes migratoria</i> (Linné). W. Pigeon			15.75	24.00	♀ ad.
	Wing, 7.75; tail, 7.15; culmen, .68; gape, 1.05; tarsus, 1.07. Colors the same as No. 1847. Middle toe, 1.06; its claw, .33.					
1852.	<i>Ectopistes migratoria</i> (Linné). Wild Pigeon.			16.00	24.00	♀ ad.
	Wing, 7.80; tail, 7.35; culmen, .68; gape, 1.05; tarsus, 1.09; middle toe, 1.06; its claw, .35. Colors like No. 1847.					

April 5, 1879.

Harrisons, Putnam Co., N. Y.

Molothrus ater, Cow-Bird. First seen.*Sceloporus ferrugineus*, Rusty Snake. First seen.*Sturnella magna*. Meadow Lark. At the Osborn's.1853. *Buteo borealis*. Red-tailed Hawk. 21.90 51.00 ♀ ad.

Wing, 14.75; tail, 9.00; culmen, 1.07; cere, .54; gape, 1.80; tarsus, 3.34; middle toe, 1.60; its claw, .88; hallux, 1.05; its claw, 1.21. Iris, yellow; legs and feet, yellow. Cere, greenish yellow. Bill, blue-black; lead-color at base. Shot with a rifle bullet, from a wagon, and presented to me by Capt. Weyant, of Ferryboat "Highlander".

1879.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
1854.	<i>Mergus serrator</i> , Red-breasted Merganser	April 8, 1879. Cornwall, on the Hudson		23.75	34.25	♂ ad.
	Wing, 9.25; tail, 4.25; culmen, 2.27; gape, 2.87; tarsus, 1.87; middle toe, 2.40; its claw, .43. Iris, bright carmine. Bill, purplish black on ridge of maxilla; remainder, red. Legs and feet, bright red; web, dusky; nails, dusky at base, white at tip.					
	Shot by Josh Ward.					
1855.	<i>Mergus serrator</i> , L. Red-breasted Merganser.			24.25	35.50	♂ ad.
	Wing, 9.50; tail, 4.20; culmen, 2.42; gape, 2.92; tarsus, 1.85; middle toe, 2.35; its claw, .37. Colors the same as No. 1854, above described.					
1856.	<i>Mergus serrator</i> , L. Red-breasted Merganser.			21.60	33.00	♀ ad.
	Wing, 9.05; tail, 3.75; culmen, 2.16; gape, 2.61; tarsus, 1.75; middle toe, 2.21; its claw, .35. Colors the same as No. 1854.					

April 5, 1879. H. M. J.

1857. *Corvus americanus*. Crow. 19.50 38.25 ♂ ad

Wing, 12.05; tail, 7.40; culmen, 1.92; gape, 2.23; tarsus, 2.37; its claw, .52. Iris, hazel.

1858. *Junco hyemalis*. Snow-Bird. 6.03 9.30 ♀ ad.

Wing, 2.80; tail, 2.55; culmen, .44; tarsus, .82. Iris, hazel.

April 9th.

Highland Falls, N. Y.

Ceryle alcyon. Kingfisher. First seen this year.1859. *Podiceps cornutus*. Horned Grebe. 14.00 24.00 ♀ ad.

Wing, 5.40; tail tuft, 1.60; culmen, .88; gape, 1.25; tarsus, 1.77; middle toe, 1.67; its nail, .30; both together, 1.97; length to end of toes, 17.40.

1879.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
(1859.)	Iris, with a narrow inner rim of yellow; outer rim broad, light red. Bill, blackish; tipped white; flesh-color at base of mandible.					
1860.	<i>Podiceps cornutus</i> . Horned Grebe. 14.65 24.75 ♂ ad.					
	Wing, 5.40; tail tuft, 1.50; culmen, .93; gape, 1.37; tarsus, 1.93; middle toe, 1.75; its claw, .32.					
	A bare space, extending from corner of mouth to eye, red. Tarsi and toes, lead color anteriorly; black posteriorly. Rest of parts colored as above, No. 1859.					
	Both of the above were shot by Josh Ward, at Cornwall.					
	April 10th.					
	Harrisons, N. Y.					
	The following species were seen for the first time this season:					
	<i>Colaptes auratus</i> . Yellow-shafted Flicker.					
	<i>Spizella socialis</i> . Chipping Sparrow.					
	<i>Spizella pusilla</i> . Field Sparrow.					
	<i>Regulus calendula</i> . Ruby-crowned Kinglet.					
	<i>Spizella</i> .					
	<i>Pooecetes gramineus</i> . Bay-winged Sparrow.					
	The following species were also observed:					
	<i>Regulus satrapa</i> . Golden-crested Kinglet.					
	<i>Ampelis cedrorum</i> . Cedar Bird.					
	<i>Buteo borealis</i> . Red-tailed Hawk.					
	<i>Nisus fusces</i> . Sharp-shinned Hawk.					
1861.	<i>Sphyrapicus varius</i> . Yellow-bellied W. 8.85 15.80 ♂ ad.					
	Wing, 4.80; tail, 3.10; culmen, .97; gape, 1.13; tarsus, .87. Iris, hazel; bill, black; legs and feet, lead-color.					
1862.	<i>Turdus migratorius</i> . Robin. 10.00 16.25 ♂ ad.					
	Wing, 5.03; tail, 4.35; culmen, .87; from nostril, .56; gape, 1.15; tarsus, 1.30.					
1863.	<i>Passerella iliaca</i> . Fox Sparrow. 6.90 10.80 ♀ ad.					
	Wing, 3.15; tail, 2.60; culmen, .49; gape, .57; tarsus, .95; middle toe, .63; its claw, .28.					

1879.

1864.	<i>Melospiza melodia</i> . Song Sparrow. 6.25 8.70 ♂ ad.					
-------	--	--	--	--	--	--

April 11th.

The following species were seen to-day for the first time this year:

Dendroica palmarum. Yellow-Redpoll Warbler. A number were seen upon the Constitution Meadows, frequenting the burnt spaces. Will Osborn shot two at a shot, amongst the burnt tussocks of grass. I found them about Pell's Pond, after reaching home.

Passerculus savanna. Savannah Sparrow. Will Osborn shot a very handsome one, on the Railroad, at Constitution Island. The yellow stripe on its head was unusually bright.

Other species seen:—

The Eagles were seen about the Cove; most of them immature.

We walked down the track just after several freight trains had passed over the bridge. Will was away ahead, and surprised me by firing at a Merganser (*Mergus merganser* ♀) that rose from beneath the bridge, while he was passing over it. This bird must have been in the same place while the trains were passing right over her head. She got away, unhurt by either of his barrels; and she flew well and vigorously, showing that she had not been previously injured.

1865.	<i>Sphyrapicus varius</i> . Yellow-bellied W. 8.50 15.35 ♀ ad.					
	Wing, 4.85; tail, 3.10; culmen, .90; gape, 1.04; tarsus, .83. Colors the same as No. 1861.					

April 12, th.

Passerculus savanna was abundant on the Cousack Meadows.

1866.	<i>Dendroica palmarum</i> . 5.65 8.75 ♂ ad.					
	Wing, 2.70; tail, 2.33.					

1879.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
1,867.	<i>Certhia familiaris</i> , L. Brown Creeper.	5.75	8.11	♂ ad.		
	Wing, 2.62; tail, 2.70; culmen, .59; from nostril .41; tarsus, .58; middle toe, .40; its claw, .22.					

The following 33 species were observed on March 18th and 19th, 1879, at Brookhaven, Lincoln County, Mississippi, By Wm. Church Osborn.

Turdus migratorius. Robin.
Harporhynchus rufus. Brown Thrasher.
Mimus polyglottus. Mocking-Bird. Common
Sialia sialis. Bluebird.
Poliophtila caerulea. Blue Grey Gnat-catcher.

Parus atricapillus^{carolinensis}. Chickadee. Common.
Tophophanes bicolor. Tufted Titmouse. Common.
Sitta carolinensis. White-bellied Nuthatch.
Sitta pusilla. Brown-headed Nuthatch.
Common. A female shot contained a somewhat developed ovum.

Dendroica coronata. Yellow-rumped Warbler.
Lanius ludovicianus. Loggerhead Shrike.
Progne subis. Purple Martin.
Passerculus savanna. Savannah Sparrow.
Common.

Pooecetes gramineus. Bay-winged Sparrow.
Melospiza palustris. Song Sparrow.
Melospiza melodia. Song Sparrow.
Zonotrichia albicollis. White-crowned Sparrow.
Very common.

Junco hyemalis. Snowbird.
Coturniculus heslowi. Henslow's Bunting.
Common; exceedingly shy and difficult to procure. A pair were preserved.
Pipilo erythrophthalmus. Chewink. Common
Sturnella magna. Meadow-lark.
Scolecophagus ferrugineus. Rusty Grackle.

1879.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	<i>Corvus americanus</i> . Crow.					
	<i>Cyanurus cristatus</i> . Blue Jay.					
	<i>Picus pubescens</i> . Downy Woodpecker.					
	<i>Picus borealis</i> . Red-cockaded Woodpecker.					
	<i>Centurus carolinus</i> . Red-bellied Woodpecker. <u>Common</u> .					
	<i>Colaptes auratus</i> . Yellow-shafted Flicker.					
	<i>Falco sparverius</i> . Sparrow Hawk.					
	<u>Quite numerous; two specimens collected.</u>					
	<i>Zenaidura carolinensis</i> . Carolina Dove.					
	<i>Ortyx virginianus</i> . Quail; Bob White. <u>Common</u> .					
	<i>Bonasa umbellus</i> . Ruffed Grouse.					
	<i>Cathartes aura</i> . Turkey Buzzard. <u>Common</u> .					
	<i>Sayornis fuscus</i> . Phoebe.					

April 13th. H.M.P.
Sphyrapicus varius. Saw one while walking to church.

April 15th. H.M.P.
Buteo borealis. Several observed.
Colaptes auratus. Now common for the first.

April 16th. H.M.P.
 1,868: *Sphyrapicus varius*. Yellow-bellied. 8.58 15.45 ♂ ad.
 Wing, 4.68; tail, 3.20; culmen, .94; gape, 1.04; tarsus, .80.
Swallows. A number of Swallows flew over the woods; but I did not get a good view of them, and so could not tell the species. One that I saw afterward, was, without doubt, the *Petrochelidon lunifrons*. Cliff Swallow. First seen
Owls. Calling in the evening, to enquire about the health of Mr. Chas. P. Smith, I was engaged in conversation with himself and wife, when the servant came, in some excitement, to tell of an Owl that she had seen in the garden. We all repaired to the kitchen, where, to my surprise, I saw a small Owl (probably *Scops asio*, though I could discover no ear-tufts).

1879.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

which I supposed might be Nyctale acadica; it sat upon a japonica-bush close to the window, where it was plainly visible in the twilight. I ran home after my gun, but before I got back, another Owl had come upon the scene, and they had flown away together. I hunted about the grave-yard and various places in vain; but met Miss Wheeler, who told me that they had "seen two Owls about their house for several evenings past."

April 17. H. H. H.

Sat up with Grandfather, who is ill. It began to rain about 2 o'clock, and has continued all day. Blackbirds are about the house.

April 18th. H. H. H.

Rained hard all night; wind north, and cold. Turned into snow this morning. The snow is deep on the mountains. Ground covered, here, with slushy snow, & hail. The Chippies and Field Sparrows seem unhappy; but the Song Sparrows are singing bravely. A number of Bay-winged Sparrows are about the house. A flock of six (6) Pigeons were found in Gatterlee's woods. I returned for a gun; found them in another place; shot one female.

What has happened to the Swallows? Chordeiles virginianus. Nighthawk.

Saw one flying over the garden; the first seen.

April 19. H. H. H.

Furdus pallasi. Hermit Thrush. The one shot is the first I have seen this year. Gallinago wilsoni. Wilson's Snipe.

1879.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

The ground is covered with snow in most places; yet the Spotted Sandpipers, Fringoides macularius, were seen for the first about the Consook creek, where three (3) were seen at once, besides another single one.

I shot a Muskrat, whose skeleton I will preserve. Passerculus savanna. Only saw the one shot. Wild Pigeon, Ectopistes migratoria. The remaining five Pigeons were seen this morning; but they were too shy to be approached. This P. M., I again found them, but so shy that I only procured one, an old male, at long range, after they had taken to flight.

Ceryle alcyon. An old male shot at Echo Lake. Weather. Rained hard all night; snowed a while this morning; wind blowing hard from the north; occasional droppings of rain this P. M. Birds quite numerous. No Woodpeckers seen.

Just received the following Bds. from Henshaw, determined as follows:

1. Eremophila alpestris, ["var.?"]
2. Passerculus savanna, ["alaudinus"].
3. Melospiza melodia, ["fallax?"]

The above specs. were collected at Fort Klammath, Oregon, and sent to Henshaw to determine vars.

Corvus americanus, Audubon.

Found a Crow's nest in a pine-tree (P. rigida), on Consook Island. Too windy to climb to it.

1869. Ectopistes migratoria. Wild Pigeon. 16.55 24.75 ♂ ad.
Wing, 8.00; tail, 7.75; culmen, .69; gape, 1.06; tarsus, 1.13; middle toe, 1.18; claw, .38.

1870. Gallinago wilsoni. Wilson's Snipe. 10.50 17.00 ♂ ad.
Wing, 4.90; tail, 2.22; culmen, 2.43; tarsus, 1.20; middle toe, 1.18; its claw, .28. Iris, hazel. Legs & feet, greenish.

1879.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
1871.	<i>Colaptes auratus</i> . Yellow-shafted Flicker.	13.30	20.80	♂ ad.		
	Wing, 4.6 6.25; tail, 4.90; culmen, 1.47; gape, 1.65; tarsus, 1.12. Bill, blue-black; legs and feet, lead-color.					
1872.	<i>Passerculus savanna</i> . Savannah sp.	5.50	9.25	♂ ad.		
	Wing, 2.67; tail, 2.03; culmen, .39; gape, .47; tarsus, .80.					
1873.	<i>Turdus pallasi</i> . Hermit Thrush. — — —			♂ ad.		
1873a	<i>Ceryle alcyon</i> . Belted Kingfisher.	13.35	22.00	♂ ad.		
	Wing, 6.10; tail, 3.90. Bill from nostril, 1.81; gape, 2.92; tarsus, .41; middle toe, .58; its claw, .40. (Not preserved).					

April 22.

	<i>Hirundo horreorum</i> . Barn Swallow. First seen.					
	<i>Lachyornis bicolor</i> . White-bellied Swallow. do					
	<i>Zenaidura carolinensis</i> . Carolina Dove.					
	Quite numerous about the Constitution Marches.					
	<i>Pandion haliaetus</i> . Fish Hawk. Common.					
1874.	<i>Zenaidura carolinensis</i> . Dove.	11.60	17.00	♀ ad.		
	Wing, 5.25; tail, 5.50; culmen, .50; gape, .72; tarsus, .75; middle toe, .72; its claw, .18.					
	Iris, hazel; bill, black; feet, red.					
1875.	<i>Ceryle alcyon</i> . Belted Kingfisher.	13.35	22.15	♂ ad.		
	Wing, 5.97; tail, 3.65; Bill from nostril, 1.88; gape, 2.92; tarsus, .48; middle toe, .60; claw, .35.					
1876.	<i>Molothrus ater</i> . Cow Bunting.	7.95	13.40	♂ ad.		
	Wing, 4.12; tail, 3.05; culmen, .66; gape, .68; tarsus, 1.00; middle toe, .72; claw, .25.					

April 23.

Zonotrichia albicollis. White-throated Sparrow. The first arrivals were seen to-day; very late.

1879.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	<i>Agelaius phoeniceus</i> . Phoebe.					
	I saw the first nest on the 21st. It was situated just above the door, in the Guard House, at the South Gate, West Point. The parent was sitting upon it.					
	I have seen other nests since, that were also seemingly finished.					
	<i>Accipiter cooperi</i> . Cooper's Hawk. One seen.					
	<i>Buteo pennsylvanicus</i> . Broad-winged Hawk.					

April 26th.

Troglodytes domesticus. First seen, by Will Osborn.
Sialia motacilla. First seen, by Will Osborn, at Harrisburg. None seen on this side of the river, yet.
Mniotilta varia. First seen, at Highland Falls.

April 27th.

Seiurus motacilla. Seen at Highland Falls.
Minus carolinensis. First seen.
Harporhynchus rufus. First seen.
Pipilo erythrophthalmus. First seen.
Accipiter fuscus. Sharp-shinned Hawk.
Accipiter cooperi. Cooper's Hawk.
Anas obscurus. Dusky Duck. A pair are staying at the Poppler Pond, and will probably breed there.

April 29th.

Troglodytes aëdon (= *domesticus*) Seen at N. Y.
Empidonax minimus. Least Flycatcher. First seen.

May 2nd.

Dendroica coronata. Yellow-rumped Warbler.
 First seen since last autumn, though I have watched for them since my return from New York.

1879.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	<i>Setophaga ruticilla</i> .	May 3 rd .	Redstart.			First seen.
	<i>Turdus mustelinus</i> .		Wood Thrush.			First seen.
	<i>Sciurus aurocapillus</i> .		Oven Bird.			First seen.
	<i>Bonasa umbellus</i> .		Ruffed Grouse.			
	<i>Buteo pennsylvanicus</i> .		Broad-winged Hawk.			
1,877.	<i>Sciurus motacilla</i> .		Highland Falls, N.Y.	6.32	10.40	♂ ad.
1,878.	<i>Setophaga ruticilla</i> .		do	5.45	8.10	♂ ad.
1,879.	<i>Minotilta varia</i> .		do	5.45	8.95	♂ ad.
1,880.	<i>Dendroica coronata</i> .		do	5.75	8.95	♀ ad.

May 4th.

Antrostomus vociferus. Whippoorwill. First seen.
Chaetura pelagica. Swift. First seen.

May 6th.

1,881. *Icterus baltimore*. Baltimore Oriole. First seen.
Antrostomus vociferus. Whippoorwill.
 Wing, 6.10; tail, 5.00; bill from nostril, 1.03; 18.60 ♂ ad.
 30; gape, 1.28; tarsus, .63; middle toe, .65; claw, .23.

May 7th.

1,882. *Icteria virens*. Yellow-breasted Chat. 7.53 9.90 ♂ ad.
 Wing, 3.10; tail, 3.25; bill from nostril, .43;
 gape, .77; tarsus, 1.00; middle toe, .65; claw, .27.
 1,883. *Regulus calendula*. Ruby-crowned Kinglet. 4.50 6.90 ♀ ad.
 Wing, 2.15; tail, 1.70; bill from nostril, .25; gape, .48;
 tarsus, .73; middle toe, .36; claw, .16.
 1,884. *Regulus calendula*. Ruby-crowned Kinglet. 4.40 6.95 ♀ ad.
 Wing, 2.12; tail, 1.65; bill from nostril, .25; gape, .50;
 tarsus, .75; middle toe, .37; claw, .16.
 1,885. *Sphyrapicus varius*. Gt. bellied Woodpecker. 8.62 15.60 ♀ ad.
 Wing, 4.92; tail, 3.43; culmen, .92; gape, 1.06;
 tarsus, .68.
 1,886. *Dendroica virens*. Blk-thr. Vireo Warbler. 5.02 7.75 ♂ ad.
 Wing, 2.42; tail, 1.95; bill from nostril, .25;
 tarsus, .67.

1879.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
1,887.	<i>Minotilta varia</i> .	H. N.Y.		5.38	8.65	♂ ad.
1,889.	<i>Sciurus aurocapillus</i> .	do		6.35	10.00	♂ ad.
	Wing, 2.95; tail, 2.05; bill from nostril, .35; tarsus, .93.					
1,890.	<i>Passer domesticus</i>	H. N.Y.		6.37	9.62	♂ ad.
	Wing, 2.90; tail, 2.40; culmen, .53; gape, .64; tar- sus, .77; middle toe, .63; its claw, .21.					
1,891.	<i>Carpodacus purpureus</i>	H. N.Y.		6.20	10.30	♂ ad.
	<i>Turdus migratorius</i>	do		9.75	15.60	♀ ad.
	<i>Bonasa umbellus</i>	do		17.00	23.35	♀ ad.
	Wing, 6.55; tail, 5.40; bill from nostril, .48; gape, 1.05; tarsus, 1.75; middle toe, 1.40; claw, .32.					
	<i>Helminthophaga ruficapilla</i> . First seen.					
	<i>Dendroica virens</i> . First seen.					
	<i>Dendroica caerulescens</i> . First seen.					
	<i>Dendroica pennsylvanica</i> . First seen.					
	<i>Icteria virens</i> . First seen.					

May 8th. H. N.Y.

Helminthophaga ruficapilla.
Helminthophaga vermivorus. First seen.
Dendroica maculosa. First seen.
Geothlypis trichas. First seen.
Parula americana. First seen.
Vireo solitarius. First seen.
Vireo gilvus. First seen.
Pyrrhula rubra. First seen.
Junco hyemalis. One observed.
Spizella monticola. Last observed April 28th.
Icterus spurius. First seen.

1,892. *Pyrrhula rubra*. Scarlet Tanager. 7.20 11.75 ♂ ad.
 1,893. *Dendroica caerulescens*. 5.40 8.15 ♂ ad.
 1,894. *Dendroica virens*. 5.20 8.00 ♂ ad.
 1,895. *Setophaga ruticilla*. Redstart. 5.40 7.85 ♂ ad.

1879.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
1896.	<i>Dendroica coronata</i> .	H. May.		5.80	9.25	♂ ad.
1897.	<i>Dendroica virens</i> .			5.25	8.00	♂ ad.
1898.	<i>Setophaga ruticilla</i> .			5.55	8.00	♂ ad.
1899.	<i>Geothlypis trichas</i> .			5.40	7.50	♂ ad.
1900.	<i>Vireo gilvus</i> .			5.50	8.75	♀ ad.
	<i>Mimus carolinensis</i> .			9.00	11.75	—

May 9th.

Tyrannus carolinensis. First seen.
Contopus borealis. First seen.
Myiarchus crinitus. First seen.
Coccyzus erythrophthalmus. First seen.

May 10th.

Vireo olivaceus. First seen.

1901. *Siurus motacilla* 6.28 10.40 ♂ ad.
 Wing, 3.20; tail, 2.20; bill from nostril, .38;
 gape, .70; tarsus, .93; middle toe, .58; claw, .16.
1902. *Siurus motacilla* 6.25 10.15 ♀ ad.
 Wing, 3.13; tail, 2.13; bill from nostril, .40;
 gape, .70; tarsus, .90; middle toe, .58; claw, .17.
1903. *Chaetura pelagica* 5.40 13.00 ♂ ad.
Siurus motacilla. Large-billed Water Thrush. The female above (No. 1902) contained an egg; full size; and one 1/4 inch in diameter. They were a pair, and doubtless had built their nest.
- Turdus fuscescens*. First seen.
Goniophya ludoviciana First seen

May 12th.

List of Birds seen:
Turdus migratorius. Robin.
Turdus mustelinus. Wood Thrush.

1879.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	<i>Turdus fuscescens</i> .		Wilson's Thrush.			
	<i>Mimus carolinensis</i> .		Cat Bird.			
	<i>Harpophynchus rufus</i> .		Brown Thrasher.			
	<i>Sialia sialis</i> .		Bluebird. Feeding its young.			
	<i>Parus atricapillus</i> .		Chickadee.			
	<i>Sitta carolinensis</i> .		White-bellied Nuthatch.			
	<i>Troglodytes aedon</i> .		House Wren.			
	<i>Minotilta varia</i> .		Black and White Creeping Warbler.			
	<i>Parula americana</i> .		Blue Yellow-backed Warbler.			
	<i>Dendroica pennsylvanica</i> .		Chestnut-sided Warbler.			
	<i>Dendroica coronata</i> .		Yellow-rumped Warbler.			
	<i>Siurus motacilla</i> .		Large-billed Water Thrush.			
	<i>Siurus aurocapillus</i> .		Golden-crowned Thrush.			
	<i>Geothlypis trichas</i> .		Maryland Yellow-throat.			
	<i>Myiodioctes nitratus</i> .		Hooded Warbler.			
			First seen; singing.			
	<i>Myiodioctes canadensis</i> .		Canada Flycatching Warbler. First seen.			
	<i>Setophaga ruticilla</i> .		American Redstart.			
	<i>Pipilo rubra</i> .		Scarlet Tanager.			
	<i>Vireo noveboracensis</i> .		White-eyed Vireo, etc.			
			etc, etc.			
			Visited the Cowcock Meadows & Island.			
			Saw no <i>Marsh Wrens</i> ; seem not to have made their appearance yet. The <i>Savanna Sparrows</i> (<i>Passerculus savanna</i>) were still there, tho I only saw one or two. Have seen no <i>Green Herons</i> (<i>Ardea virescens</i>).			
			not.			
1904.	<i>Siurus aurocapillus</i> .			6.25	9.90	♂ ad.
1905.	<i>Siurus aurocapillus</i> .			5.80	8.80	♀ ad.
1906.	<i>Vireo noveboracensis</i> .			5.37	8.00	♂ ad.
1907.	<i>Pipilo erythrophthalmus</i>			8.62	11.50	♂ ad.

1879.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	<u>May 14th. N. H. N. Y.</u>					
	<u>Turdus swainsoni.</u>					<u>First seen.</u>
	<u>Dendroica castanea.</u>					<u>First seen.</u>
	<u>Dendroica striata.</u>					<u>First seen.</u>
	<u>Dendroica tigrina.</u>					<u>First seen.</u>
	<u>Dendroica blackburniae.</u>					<u>First seen.</u>
	<u>Empidonax flaviventris.</u>					<u>First seen.</u>
	<u>Contopus virens.</u>		<u>Wood River.</u>			<u>First seen.</u>
	<u>May 15th. N. H. N. Y.</u>					
	<u>Helminthophaga chrysoptera.</u>					<u>First seen.</u>
	<u>Ardea virescens.</u>					<u>First seen.</u>
1,908.	<u>Helminthophaga chrysoptera.</u>			5.00	7.90	♂ ad.
	Wing, 2.47; tail, 1.95; bill from nostril, .34; gape, .57; tarsus, .65; middle toe, .42; its claw, .16.					
1,909.	<u>Icteria virens.</u>			7.35	9.75	♂ ad.
	Wing, 2.98; tail, 3.06; bill from nostril, .41; gape, .76; tarsus, 1.05; middle toe, .68; its claw, .24.					
1,910.	<u>Dendroica pennsylvanica.</u>			5.28	7.95	♂ ad.
	Wing, 2.48; tail, 2.05; bill from nostril, .29; gape, .52; tarsus, .72; middle toe, .38; its claw, .17.					
1,911.	<u>Myiodiastes canadensis.</u>			5.60	8.00	♂ ad.
	Wing, 2.42; tail, 2.35; bill from nostril, .29; gape, .55; tarsus, .75; middle toe, .38; its claw, .17.					
1,912.	<u>Myiodiastes nitratus.</u>			5.70	8.25	♂ ad.
	Wing, 2.51; tail, 2.35; bill from nostril, .31; gape, .60; tarsus, .73; middle toe, .42; its claw, .20.					
1,913.	<u>Myiodiastes nitratus.</u>			5.80	8.50	♂ ad.
	Wing, 2.65; tail, 2.45; bill from nostril, .32; gape, .59; tarsus, .75; middle toe, .45; its claw, .19.					
1,914.	<u>Setophaga ruticilla.</u>			5.60	7.78	♂ juv.
	Wing, 2.45; tail, 2.37; bill from nostril, .27; gape, .52; tarsus, .65; middle toe, .40; its claw, .16.					
	<u>Cyanospiza cyanea.</u>					<u>First seen.</u>

	<u>May 14th</u>					
1,915.	<u>Dendroica castanea.</u>			5.75	9.05	♂ ad.
	Wing, 2.86; tail, 1.10; bill from nostril, .32; gape, .60; tarsus, .70; middle toe, .45; its claw, .17.					
1,915a.	<u>Dendroica caerulescens.</u>			5.45	8.12	♂ ad.
	<u>May 17th. N. H. N. Y.</u>					
1,916.	<u>Myiodiastes pusillus.</u>			4.70	6.75	♂ ad.
	Wing, 2.10; tail, 2.00; bill from nostril, .24; gape, .48; tarsus, .70; middle toe, .44; its claw, .18. Iris, hazel; bill, with maxilla, blackish; mandible, yellow; legs & feet, yellow. <u>First seen.</u>					
1,917.	<u>Myiodiastes canadensis.</u>			5.60	8.10	♂ ad.
	Wing, 2.55; tail, 2.20; bill from nostril, .33; gape, .61; tarsus, .74; middle toe, .43; its claw, .19.					
1,918.	<u>Parula americana.</u>			4.70	7.70	♂ ad.
	Wing, 2.40; tail, 1.75; bill from nostril, .32; gape, .54; tarsus, .68; middle toe, .40; its claw, .16.					
1,919.	<u>Myiodiastes canadensis.</u>			5.60	8.10	♂ ad.
	Wing, 2.55; tail, 2.24.					
1,920.	<u>Dendroica castanea.</u>			6.00	9.30	♂ ad.
1,921.	<u>Dendroica striata.</u>			5.75	9.00	♂ ad.
1,922.	<u>Coccyzus erythrophthalmus.</u>			11.50	15.70	♂ ad.

Measurements:

Myiostilta varia. Black and White Creeping Warbler. May 17, N. H. N. Y. 5.12 8.50 ♂ ad.
 wing, 2.58; tail, 1.88; bill from nostril, .36; gape, .57; tarsus, .69; middle toe, .53; claw, .17.
Myiostilta varia. 5.40 8.40 ♂ ad.
 wing, 2.58; tail, 2.00; bill from nostril, .38; gape, .62; tarsus, .67; middle toe, .47; its claw, .18.

May 19th.

Dendroica aestiva. First seen.

1879.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	<u>May 20, 1879. H.F.N.Y.</u>					
1,923.	<u>Dendroica striata.</u>			5.75	9.10	♂ ad.
	Wing, 2.90; tail, 2.15; bill from nostril, .30; gape, .56; tarsus, .75; middle toe, .45; its claw, .17.					
1,924.	<u>Icterus spurius.</u>			7.40	10.60	♂ ad.
	Wing, 3.14; tail, 3.00; bill from nostril, .47; gape, .77; tarsus, .86; middle toe, .62; its claw, .26.					
1,925.	<u>Tyrannus carolinensis.</u>			8.50	14.60	♀ ad.
	Wing, 4.68; tail, 3.60; bill from nostril, .55; gape, 1.00; tarsus, .72; middle toe, .57; its claw, .30.					
	<u>Sturnella magna.</u> One seen on Cousick Island.					
	<u>May 21st.</u>					
1,926.	<u>Dendroica striata.</u>			5.70	9.10	♂ ad.
	Wing, 2.90; tail, 2.00; bill from nostril, .31; gape, .57; tarsus, .76; middle toe, .45; its claw, .17.					
1,927.	<u>Dendroica maculosa.</u>			5.12	7.55	♂ ad.
	Wing, 2.27; tail, 2.00; bill from nostril, .30; gape, .53; tarsus, .71; middle toe, .38; its claw, .17.					
1,928.	<u>Helminthophaga chryoptera.</u>			5.10	7.60	♂ ad.
	Wing, 2.45; tail, 1.85; bill from nostril, .36; gape, .59; tarsus, .68; middle toe, .42; claw, .15.					
1,929.	<u>Geothlypis trichas.</u>			5.15	6.75	♀ ad.

Several specimens of H. chryoptera were seen or heard, near Pell's Pond.

Walking through Jatterlee's place in the evening, a bird flew from a tree over head. I followed it, and discovered that it was an Owl. The nest was discovered in the top of a decayed tree, growing beside the pond; the same in which I captured the parents and young of Scops asio (gray plumage) in

1879.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	<u>May 22. H.F.N.Y.</u>					
1,930.	<u>Pyrranga rubra.</u>			7.37	11.70	♂ ad.
1,931.	<u>Pyrranga rubra.</u>			7.30	11.75	♂ ad.
	Shot on Bear Mountain.					
	<u>May 23. H.F.N.Y.</u>					

1875. On thrusting my hand down into the hole, I found a single young bird, and an addled egg, which I captured and preserved. The birds were doubtless Screech Owls; they snapped their bills about the place, but I did not get a distinct view of them; their cry was very singular, and new to me. Willson visit the place in the day-time and get the birds.

To-day I visited the Owl's nest. I captured the female in her nest, along with the young bird. After a search, I saw the male sitting in the crotch of a white oak-tree, in the midst of a clump of branchlets; his ears were very conspicuous, and his neck extended, as he endeavored to get a better view of me. Both of the parents were red; the young gray, like those I discovered in the same tree in 1875.

1,932. Scops asio (Linne'). Screech Owl (Red). ♀ ad.
Wing, 6.45; tail, 3.10; culmen, .65; 10.15 25.25
gape, 1.00; tarsus (about), 1.60;
middle toe, 1.80; its claw, .48. Iris, yellow.

1,933. Scops asio (Linne'). red. 9.25 23.50 ♂ ad.
Wing, 6.50; tail, 3.00; gape, .95; culmen, .60;
tarsus, 1.48; middle toe, 1.80; its claw, .47.

1,934. Scops asio (Linne') fulvus
The above 3 spec. are a family taken together. An addled egg (set 1^a No. 152) was also taken: see "Coll. Nests & Eggs," p. 77.

1879.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
1935.	<u>Sayornis fuscus.</u>	May 24th.	H. M. N. Y.	6.90	10.20	♀ ad.
	Wing, 3.06; tail, 2.45; bill from nostril, .41; gape, .82; tarsus, .64; middle toe, .40; claw, .82.					

May 27, H. M. N. Y.

To-day, after a long search, I discovered a nest of Siurus motacilla, which contained five nearly fledged young birds.

The following migrants were seen:—
Dendroica striata, Myiodiastes canadensis.

May 30th. H. M. N. Y.

Visited the Rookery of Night Herons, on Constitution Island, in company with Lieut. Wittich and Mr. Rhett. We stopped at the lower falls, on our way up; there we were all somewhat startled by the accidental discharge of a gun, loaded with large shot, the contents of which passed within a few feet of my companions. I was the culprit, and felt correspondingly foolish, but thankful that it resulted so harmlessly. Lieut. Wittich observed, quietly, that "lightning never strikes twice"; while Mr. Rhett philanthropically remarked that a "miss is as good as a mile".

We heard the "barking" of the Herons before we entered the swamp, and were glad to find that we were not going to be disappointed. As we entered a little way, we started the first Heron from its nest quite close to the Railroad, and soon there were numbers of them flying about over the swamp. Their

1879.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	<p>numbers have greatly increased since last season. I succeed in killing three adult birds out of a number of snap shots on the wing; and Lieutenant Wittich made a beautiful wing shot, at very long range, so that we secured four fine specimens. We arrived too late to get any fresh eggs, though I saw a bird flying with a stick in its bill. Lieutenant W. climbed to a nest which contained five eggs that were about to be hatched; he having brought an egg down for trial. I climbed up and examined two nests, one of which contained five, and the other four eggs. The shells of the former were already cracked, while an egg from the latter nest was found on trial to contain a well developed embryo. Shells were found under numerous nests, proving that the eggs were already hatched, and the young birds were heard crying in some nests. The wounded birds were sure to lodge in some tree-top, where it was impossible to dislodge them. After a great deal of effort and flapping of wings, one would manage to sit upright upon its perch; then, when the tree was shaken vigorously, he would lose his hold and topple over, but invariably recovered his hold by means of his bill. With two strong feet, and a bill that could be hooked upon a branch, it was out of the question to dislodge a bird without climbing up the tree, and then the bird was almost certain to escape into a neighboring tree. The bird should always be caught by its bill, for they always strike precisely</p>					

1879

- with that powerful organ, thus endangering
ones eyes. On the ground they often
escape by running swiftly through
the ferns, brakes and bushes that
grow so abundantly in the swamp.
- 1,936. *Nyctiardea grisea*, var. *naevia* (Boddaert).
Wing, 12.50; tail, 5.10 27.50 47.50 ♂ ad.
Bill (culmen), 3.04; gape, 4.25; tarsus, 3.40;
middle toe, 2.96; its claw, .57. Iris, carmine.
Bill, blue-black, except extreme tip of man-
-dible, which is horn-color. Bare space about
face, greenish; about eye, bluish. Legs,
pale yellow.
- 1,937. *Nyctiardea grisea*, var. *naevia*. 28.00 47.50 ♂ ad.
Wing, 12.20; tail, 5.05; culmen, 3.15; gape, 4.35;
tarsus, 3.40; middle toe, 2.95; its claw, .55
- 1,938. *Nyctiardea grisea*, var. *naevia*. 25.00 44.25 ♀ ad.
Wing, 12.00; tail, 4.70; culmen, 3.03; tarsus, 3.20;
gape, 4.12; middle toe, 2.85; its claw, .54.
- 1,939. *Nyctiardea grisea*, var. *naevia*. 27.10 47.75 ♂ ad.
Wing, 12.60; tail, 5.20; culmen, 3.27; gape,
4.40; tarsus, 3.46; middle toe, 3.12;
its claw, .58.

Colors of parts in all the same as No.
1,936. There are, occasionally, a few black
spots on the tarsi and toes of these species.

June 14th.

Visited the Roostery of Night Herons, on
Constitution Island to-day, in company
with Messrs. Rutt, Wittich & Lockwood.
The young were very noisy, but none
had left the nests. I found that
the colony had increased a greatly
and spread itself over the whole of
the Southwestern part of the swamp,
where only a few scattered nests

1879

had been seen before.

June 16th.

A large Heron ("Crane") has been frequently seen
about the pond ("Echo Lake") for a week or two
past. To-day I went to look after it.
I found a large Heron which I think
was a Night Heron.

June 17th.

The Heron was at the pond, where I found
it, in company with Wittich, who made
an unsuccessful shot at it, at which
time I observed it more satisfactorily, and
saw that it was a Night Heron. From
the abundant excretion found under the
tree I supposed that it usually perched
in the same spot. It may have nested
in this vicinity, as it has been seen about
the pond for some weeks, I am told.

June 28th. H. H. H.

- 1,940. *Icteria virens*. 7.20 9.60 ♂ ad.
Wing, 2.90; tail, 3.15; bill from nostril, .39; gape,
.82; tarsus, .95; middle toe, .63; its claw, .22.
- 1,941. *Myiodiastes nigratus*. 5.70 8.10 ♀ ad.
- 1,942. *Colaptes auratus*. — — — ♂ juv.
- 1,943. *Dendroica pennsylvanica*. — — — juv.
- 1,944. *Helminthophaga chrysoptera*. 5.15 8.25 ♂ ad.
Wing, 2.52; tail, 2.10; bill from nostril, .35;
gape, .57; tarsus, .69; middle toe, .42; its claw, .17.
Iris, hazel. Bill, black, with base of mandible, horn-color.
Tarsi and toes, greenish-lead-color.
- 1,945. *Helminthophaga chrysoptera*. — — — ♂ juv.
Testes plainly visible.
- 1,946. *Helminthophaga chrysoptera*. — — — ♂ juv.

1879.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

June 28th (continued).

List of Birds seen:—

Turdus migratorius.*Turdus mustelinus*.*Minus carolinensis*.*Harporhynchus rufus*.*Sialia sialis*. Incubating second brood*Troglodytes domesticus*. do*Amistilla varia*. Quite numerous.*Helminthophaga chrysoptera*.

I first heard the note of the young birds; presently saw an old male, and shot it; could not find the female, but succeeded in getting a good specimen of a young male. Searched carefully & thoroughly, but could not see the mother. I afterwards heard another young one, which I watched carefully for a half-hour, hoping, as it kept up an incessant call, that its mother might come to feed it. The sun was excessively hot and my patience at length became exhausted; so I shot the young one, and came home. It could only make short flights from branch to branch, and must have been reared in the immediate neighborhood where I found it. Its call, was a monotonous, rasping, beetle-like note, which, tho' unlike the singular utterance of the old bird, brought its species to my mind before I had seen any of the birds.

Dendroica aestiva.*Dendroica pennsylvanica*. Common*Sturnus aureicapillus*.*Geothlypis trichas*.*Icteria virens*. Numerous.*Myiodytes mitratus*. do*Setophaga ruticilla*. In young flying.*Arundo horreorum* (= *erythrogaster*).

1879.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

Vireo olivaceus.*Myiarchus crinitus*.*Empidonax (academicus?)* Shot at a Flycatcher which I thought to be of this species - Academic Flycatcher*Trochilus colubris*. Several seen.*Nyctiardea grisea naevia*. Night Herons are constantly seen, flying toward the north-east, in the evening; probably returning from the pond, to the Rookery on Constitution Island.

June 29th.

To-day I called a fine male *Helminthophaga chrysoptera* close up to me, when calling the various species of nesting species from the vicinity of the outlet to Pell's Pond.

June 30th.

1947. *Myiodytes mitratus*. 5:50 8:10 ♂ ad.
Wing, —; tail, —. Iris, dark hazel.1948. *Vireo olivaceus flavifrons*. 6:00 9:65 ♂ ad.
Wing, 2.90; tail, 2.08; bill from nostril, .35; gape, .71; tarsus, .76; middle toe, .47; its claw, .20.

July 7th.

Note. - *Vireo flavifrons* has nested in considerable numbers here this season. I see or hear it very frequently, and in various localities.1949. *Agelaius phoeniceus*. 9:45 15:00 ♂ ad.
Wing, 4.55; tail, 3.85; culmen, .97; gape, 1.00; tarsus, 1.14; middle toe, .80; its claw, .33.1950. *Helminthophaga chrysoptera* 4:85 7:30 ♀ ad.
Wing, 2.21; tail, 1.85; bill from nostril, .32; gape, .55; tarsus, .66; middle toe, .37; its claw, .15.
Iris, dark hazel. Bill brownish-black above, horn-color at base of mandible. Tarsi & toes, brownish-lead-color; ventral surface of toes, yellowish.

1879.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

- July 7th (continued) H. Story.
- 1,951. *Helminthophaga chrysoptera*. — — pullus.
- 1,952. *Helminthophaga chrysoptera*. — — pullus.
- This and the preceding are the young of the female parent, above (No. 1,950 ♀ ad).
- 1,953. *Picus pubescens* (top of head red). — — ♂ juv.
- 1,954. *Geothlypis trichas*. 5:50 7:35 ♂ ad.
- Wing, 2:10; tail, 2:15; bill from nostril, .30; gape, .58; tarsus, .80; middle toe, .48; its claw, .20.
- The gray on forehead less restricted than usual.
- 1,955. *Geothlypis trichas*. 5:25 7:35 ♂ ad.
- Wing, 2:15; tail, 2:06; bill from nostril, .32; gape, .57; tarsus, .78; middle toe, .52; its claw, .20.

July 10th. H. Story.

- 1,956. *Icterus baltimore*. — — ♀ ad.
- This specimen had black feathers on its head, and nape, which lead me to suppose that it was a young ♂. It proved, on dissection, to be an unmistakable female (♀). I showed the ovary to Louis Zerega and lifted out one of the largest ova on the point of my scalpel.
- 1,957. *Myiodytes mitratus*. 5:73 8:45 ♂ ad.
- Wing, 2:63; tail, 2:32; bill from nostril, .32; gape, .57; tarsus, .79; middle toe, .45; its claw, .16.
- 1,958. *Myiodytes mitratus*. 5:60 8:00 ♀ ad.
- Wing, 2:40; tail, 2:40; bill from nostril, .32; gape, .57; ~~tarsus~~ middle toe, .78; middle toe, .44; claw, .19.

Helminthos vermivorus. — I saw a male bird (probably ♂) in very good plumage, which uttered a faint metallic note, and came close up, as I called; afterwards it flew down the hill, and began to sing a pretty, metallic, trill, as it sat upon the end of a dead limb of a large tree, in plain view.

Helminthophaga. — None were observed.

1879.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

- July 13th. H. Story.
- Nyctiardea grisea naevia*. Frequently observed about Echo Lake and Pell's Pond; also flying in a northeasterly or southwesterly direction.
- 1,959. July 19th. H. Story.
- Geothlypis trichas*. — — pullus.
- 1,960. *Spizella socialis*. — — pullus.
- 1,961. *Felmatorytes palustris*. 4:95 6:65 ♂ ad.
- 1,962. *Vireo flavifrons*. 6:00 9:80 ♂ ad.
- Wing, 3:03; tail, 2:23; bill from nostril, .36; gape, .73; tarsus, .76; middle toe, .47; its claw, .20. Iris, hazel.

July 24, 1879. Constitution Island, Huds. Riv.

- 1,963. *Nyctiardea grisea naevia*. — — juvenis
- 1,964. " " " — — ♂ juv.
- 1,965. " " " — — ♂ juv.
- 1,966. *Cotyle riparia*. 5:20 11:00 ♂.
- Wing, 3:90; tail, 2:05; bill from nostril, .17; gape, .50; tarsus, .41; middle toe, .40; its claw, .19; outer toe, .26; its claw, .17.
- 1,967. *Stelgidopteryx serripennis*. 5:62 12:10 ♂ juv.
- Wing, 4:12; tail, 2:05; bill from nostril, .17; gape, .55; tarsus, .43; middle toe, .39; ~~outer~~ its claw, .17; outer toe, .25; its claw, .12.
- Iris, dark hazel. Bill & feet, black.
- 1,968. *Cotyle riparia*. 5:20 10:88 ♂
- 1,969. *Cotyle riparia*. 5:12 10:75 ♀

To-day visited the Heronry on Constitution Island, where we found the Herons numerous, and quite shy. I shot four young birds, and Lieutenant Mattick and Louis Zerega shot three more, making seven, all but one of which were in the first plumage; that one appeared to be in second year's plumage, having the wings marked with a rufous cast, and eyes red, while the eyes of the other six, were yellow.

1879.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

or yellowish-red. Louis & myself prepared five specimens. Six of the birds were cooked, and served up at the table; all of them were eaten, and pronounced very good, though I did not consider it to be much of a delicacy, tho I managed my portion with a very good grace.

The following species were seen in the swamp: Helminthophila vermivorus, Vireo noveboracensis, and Myiarchus crinitus.

On the southeastern point of the Island the various species of Swallows were, as is usually the case at this season, collected in immense numbers, I procured specimens of three different genera at a single shot, with No. 5 shot (Cotyle, Hirundo, & Petrochelidon).

Several Rough-winged Swallows (Stelgidopteryx serripennis) were observed, and one fine specimen was procured, though the immense numbers of the other species (Hirundo, Tachycineta, Cotyle & Petrochelidon) made it difficult to discern the rare species in time for a shot. Time & small shot being limited, we were obliged to leave without thoroughly investigating this army of Swallows.

1970. Picus villosus (Yellowish on head) — juv.
August 10th, 1879. Highland Falls, New York.

August 12th.

Garrisons, Putnam County, N.Y.

1971. Cotyle riparia. 5.35 11.10 ♂ ad.
Wing, 3.85; tail, 1.88; bill from nostril, .20;
gape, .48; tarsus, .45; middle toe, .39; its
claw, .22; outer toe, .27; its claw, .17.

1972. Cotyle riparia. 5.00 10.60 ♂ juv.

1973. Hirundo erythrogastra. 6.90 13.00 ♀ ad.
Wing, 4.60; tail, 3.30; bill from nostril, .24;

1879.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

1974. Cyanospiza cyanea. 5.60 8.50 ♂ ad.
Wing, 2.60; tail, 2.25; culmen, .44; gape, .47;
tarsus, .65; middle toe, .50; its claw, .20.

August 15th.

Rockland County, N.Y.

To-day, Louis Zerega and myself drove down to Stony Point, in the phaeton. We saw the White-headed Eagles about Highland Lake, and nothing else in the bird line that was particularly remarkable. Enjoyed the ride immensely; and shot one Indigo Bird.

1975. Cyanospiza cyanea. 5.50 8.30 ♀ ad.
Wing, 2.48; tail, 2.05; culmen, .42; gape, .47;
tarsus, .70; middle toe, .52; its claw, .22. Shot
it on the telegraph wire.

Capture of a Fish Crow (Corvus ossifragus)
near Seabright, Monmouth County, New Jersey.

April 14th, 1879, Edward Keeler brought me a "Sea Crow", stating that it was shot the day before, in company with many "Land Crows" (probably C. americanus; but C. corax also occurs); all were very shy. It proved, on dissection, to be a male bird; and of remarkably large size. Its dimensions are greater than any I can find on record, viz.: Length, 17.50; ~~the~~ extent, 34.00; wing, 11.50; tail, 7.10; culmen, 1.63; gape, 1.92; tarsus, 1.90; middle toe and claw, 1.96; graduation of tail, .55. Stomach contained two whole shrimps, besides several fragments of the same."

Louis A. Zerega, 23 North Washington Square,
N.Y. City. August 23, 1879.

1976. Buteo pennsylvanicus. 15.50 34.50 ♂ juv.
Wing, 10.25; tail, 6.70; culmen, .68; cere, .45; gape,
1.20; tarsus, 2.35; mid. toe & claw, 1.70; toe alone, 1.18; claw, .58.
Iris, hazel.

1879.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
1,977.	<i>Podilymbus podiceps</i>		Hudson River at Cornwall.	14.25	23.75	♂
	Wing, 5.10; culmen, .84; gape, 1.35; Tarsus, 1.62.	August 25th.	Shot by John Ward.			

September 1st 1879.

Saw two Olive-sided Flycatchers (*Contopus borealis*) sitting on a dead pine-tree, in Rockland County, opposite Peekskill.

September 6, 1879.

Shot specimens of the Black and Yellow Warbler and Black-throated Blue Warbler (*D. D. maculosa* and *caerulescens*) in Putnam Co. near Harrison's.

1,978.	<i>Setophaga ruticilla</i>	5.53	7.85	♂ ad.
	Putnam County, N. Y., near Hamilton			Hick's house.
1,979.	<i>Dendroica virens</i>	4.98	7.75	♀
1,980.	<i>Porzana carolina</i>	9.12	14.40	♂ juv.
	Wing, 4.32; tail, 2.25; culmen, .89; gape, .90; tarsus, 1.40; middle toe, 1.45; its claw, .31.			
1,981.	<i>Corvus americanus</i>			♂ juv.

September 9

Consook Island, Hudson River, N. Y.

The Reed Birds (*Dolichonyx oryzivorus*) were very numerous on the 6th, when hundreds were seen by Miss. Ella Wittich and myself.

Yesterday evening, Misses. Rose & Ella Wittich & myself rowed a little way up the creek, at high tide, in the evening. Large numbers of Rails flew up along the creek, or ran through the grass near the boat.

1,982.	<i>Porzana carolina</i>	9.40	14.50	♂ juv.
	Wing, 4.20; tail, 2.12; culmen, .88; gape, .96; tarsus, 1.39; middle toe, 1.48; its claw, .34.			
	iris, reddish hazel. Bill, greenish yellow.			

1879.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
1,983.	<i>Porzana carolina</i>			8.65	13.65	♀ juv.
	Wing, 4.02; tail, 2.00; culmen, .78; gape, .87; tarsus, 1.27; middle toe, 1.37; its claw, .32.					

Note. -

Selmatodytes palustris. Several seen in the Consook marsh September 10th.

Melanerpes erythrocephalus. A young bird seen on the wild cherry tree, near the house, on the 11th.

Sialia sialis. Large flocks were seen flying, on the evening of the 12th.

September 16th

Highland Falls, N. Y.

1,984.	<u>Turdus mustelinus</u>	—	—	♂ ad.
1,985.	<u>Coccyzus erythrophthalmus</u>	—	—	♀ ad.
1,986.	<u>Dendroica caerulescens</u>	—	—	♂ ad.
1,987.	<u>Chaetura pelagica</u>	5.55	12.50	♂ ad.
	Wing, 4.90; tail, 1.90; bill from nostril, .16; gape, .59; tarsus, .48; middle toe, .27; its claw .21.			
1,988.	<u>Chaetura pelagica</u>	5.45	12.25	♂ ad.
	Wing, 4.90; tail, 1.88; bill from nostril, .14; gape, .60; tarsus, .48; middle toe, .27; its claw .22.			
1,989.	<u>Chaetura pelagica</u>	5.50	12.65	♂ ad.
	Wing, 5.00; tail, 1.90; bill from nostril, .14; gape, .60; tarsus, .47; middle toe, .28; its claw, .22.			

September 18th.

Highland Falls, N. Y.

1,990.	<i>Selmatodytes palustris</i>					♂ ad.
1,991.	<i>Selmatodytes palustris</i>					♂ juv.
1,992.	<i>Layornis fuscus</i>					♀ ad.

Dolichonyx oryzivorus was still quite numerous on the Consook marsh.

Note. - The last Barn Swallows were seen on the Consook marsh September 6th.

1879.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

September 19th. H.F., N.Y.

1993. *Cyanospiza cyanea*. Several seen at the gate.
Pipilo erythrophthalmus. 8.35 11.00 ♀ ad.

Iris, bright red.

1994. *Podilymbus podiceps*. 14.75 23.25 ♂

Shot at Cornwall, on the Hudson, by "Josh" Ward.

Wing, 5.05; tail, 1.50; culmen, .88; bill from nostril, .55; tarsus, 1.66; middle toe and claw, 2.30; toe alone, 2.00; claw, .30.

September 20th. H.F., N.Y.

1995. *Dendroica caerulescens*. 5.30 8.15 ♂ ad.

1996. *Dendroica striata*. 5.70 9.30 ♂ (ad.?)

1997. *Dendroica striata*. 5.60 8.85 ♂ (ad.?)

1998. *Dendroica palmarum hypochrysea* 5.35 8.05 ♀ ad.

September 22nd, Highland Falls, N.Y.

Chaetura pelagica. Several were seen at Echo Lake; one that I shot could not be recovered.

1999. *Vireo solitarius* 5.62 9.15 ♀ ad.

Wing, 2.87; tail, 2.20; bill from nostril, .28; gape, .65; tarsus, .75; middle toe, .45; its claw, .21.

September 23rd. H.F., N.Y.

Chaetura pelagica still present.

Ardea herodias. One seen in the Cousack Creek.

Halibutus leucocephalus. A young one seen - same place.

September 25th. H.F., N.Y., and Cold Spring, N.Y.

Hydrochelidon lariformis. - Mr. Thos. W. Wilson, of Cold Spring, described two Terns which he saw in the "Upper Cove", near Cold Spring, about September 11th, 1879, which were undoubtedly of the present species.

Podilymbus podiceps. - One was chased unsuccessfully in the upper cove; it dived whenever approached, and escaped by retreat-

1879.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

-ing into the edge of the bulrushes; on rising to the surface, after diving, it frequently exposed no more than its head above the surface.

Podiceps cornutus. - A pair was chased in the ~~lower~~ Upper Cove; they kept close together, even when pursued. One that was severely wounded dived to the bottom, where it remained, and the other escaped into the bulrushes.

Fringa minutilla. On firing a shot near the sluice, at the north end of the Lower Cove, I saw a flock of about twelve small Snipes, which arose from the edge of the bulrushes at the side of the cove (it was then low tide), and flew northward in a very crooked and eccentric manner, now high, and again low, and diving from side to side; their pure white ventral surfaces made them appear distinctly against the background of green mountains, so that we could follow their flight for a long distance, when they rose above the horizon line, and were lost to view. I knew that they were birds that I had never seen before alive, and new to this region; so we started toward the Upper Cove, hoping that they had stopped there. After passing from the Railroad as far round the cove as the road, I saw four of the same birds, so far as I could judge, which were fly low about a large pool in the midst of the bulrushes to the south. My friend, Mr. Thos. W. Wilson, who accompanied me, advised me not to attempt to reach them there, but I determined to do so if possible; so I plunged into the reeds, and waded towards the pool, which was several rods distant, my friend having promised to apprise me of any movement on the part of the birds. Having reached the

1879.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

margin of the pool, I anxiously looked around for the birds which seemed nowhere visible. I was already wet to my knees, but the pool was black and muddy, but I plunged in, regardless, until I reached a point where I could view its whole extent. After scanning the border of the rushes carefully for a moment, I discovered the four Snipes which were feeding, deeply immersed, in the slimy vegetable truck close to the reeds. I fired from where I stood nearly waist deep in mud & water. One bird was killed, and another slightly wounded, but it escaped. I reloaded my gun, a single barreled Maynard, and fired a second shot after the retreating flock, and succeeded in wounding the bird that was hurt by my first shot, so that it fell at Mr. Wilson's feet. Then I waded across the pool, with my coat-tails dragging on top of the water, and soon had in my hand the first Least Sandpiper that has been captured in this region.

Ardea herodias. - Four were seen in the Lower Cove at low tide. Mr. Wilson and myself fired several shots from the Maynard rifle, but they were never closer than 300 yards. I fired one shot which splashed water all over the bird, and scared it out of its seven senses.

Eximetus pusillus Cold Spring, New York. Sept. 25th.

2,000. Tringa semipalmata. Semipalmated Sandp. 6.15 12.00 ♂ ad.
Wing, 3.70; tail, 1.58; culmen, .70; gape, .68;
tarsus, .83; middle toe, .60; its claw, .12.

Iris, hazel. Bill, legs & feet, black.

2,001. Tringa semipalmata. Semipalm. Sandp. 5.90 11.65 ♂ ad.
Wing, 3.62; tail, 1.56; culmen, .66; gape, .65;
tarsus, .79; middle toe, .57; its claw, .12. Colors like 2,000.

1879.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

2,002. Totanus solitarius. Highland Falls, N. Y. (Pell's Pond). 8.75 16.75 ♂ ad.
Wing, 5.12; tail, 2.45; culmen, 1.11; gape, 1.30;
tarsus, 1.21; middle toe, .95; its claw, .17.

September 29th. H. J. May.

Chaetura pelagica. - Several (twenty or so) were seen by Miss Wittich and myself, this evening.

Scops asio. - Several heard to-night. I went after one with my gun, but my shot only brought down leaves.

Note. - Lanius borealis ♀ juv. No. 1,824.

Measurements: Length, 10.15; stretch, 14.15; wing, 4.35; tail, 4.65; bill from nostril, .55; gape, 1.10; culmen, .73; tarsus, 1.00; middle toe, .61; its claw, .28.

Iris, hazel. Bill, blue-black; base of mandible, flesh-color. Legs and feet, black.

Luscinia sylvia ♂ ad. No. 1,826.

Measurements: Length, 12.50; stretch, 17.75; wing, 5.55; tail, 5.40; culmen, 1.17; gape, 1.35; tarsus, 1.45; middle toe, 1.00; its claw, .34. Iris, yellow. Bill & feet, black.

Melospiza melodia ad. No. 1,828.

Measurements: Length, 6.40; stretch, 8.50; wing, 2.50; tail, 2.75; culmen, .46; tarsus, .80. Iris, hazel. Bill, flesh-color; maxilla, blackish brown. Legs and feet, light fleshy brown.

Picus villosus ♂ ad. No. 1,829.

Measurements: Length, 9.35; stretch, 15.25; wing, 4.60; tail, 3.40; culmen, 1.31; tarsus, .84. Iris, hazel. Bill, lead-color. Feet, greenish-head-color.

Picus villosus ♀ ad. No. 1,830.

Measurements: Length, 8.95; stretch, 15.35; wing, 4.50; tail, 3.18; culmen, 1.16; tarsus, .75.

Iris, hazel. Bill, lead-color. Feet, greenish-lead-color.

Turdus migratorius ♀ ad. No. 1,827.

Measurements: Length, 10.80; stretch, 17.10; wing, 5.35; tail, 4.30; bill from nostril, .54; gape, 1.13; tarsus, 1.41; middle toe, .87; claw, .30.

1879.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
2,003.	<i>Spizella socialis</i> .	October 1st.	N. Y. City.	5.50	9.00	♂ juv.

October 6th. Hudson River, New York.
Left home for the College of Physicians and Surgeons, New York City, there to remain till Spring. On my way down, on the Steamer "Mary Powell", I saw the following birds:

Larus delawarensis. - Two Gulls which I saw about opposite Tarrytown were probably of this species.

Colymbus torquatus. - One seen flying up the Hudson, not very far from our vessel, near the City.

Note. - *Empidonax traillii*. "J. M. B. No. 1978. Catskill Mountains, N. Y. Dr. James C. Merrill. .72 X .53; .70 X .52. One of these is very nearly an unspotted white." From a separate sent me by Dr. Brewer. "Proc. Nat. Mus. 79 — 1 April 29, 1879." p. 4.

Empidonax acadicus, Baird.
"J. M. B. No. 2735. Staten Island, N. Y. S. D. Osborne. June 5, 1876. .81 X .60; (nearly unspotted) .80 X .59; .79 X .59." From same paper as above, p. 5.

October 18th. Highland Falls, N.Y.
Returned from the Medical College last night. This morning I rose early and enjoyed an excellent swim in the Hudson. The water was rough, but did not seem cold. Gulls were quite numerous. Herring Gulls were the only species noticed.
Scops asio - One screeching near my window,

1879.

early in the morning, just before daybreak.
Warblers. - *D. coronata*. Abundant. The only species seen.

R. R. calendula and *satrappa* were numerous.

I remained at home till Monday, Oct. 20th. during which time very few birds were observed. *Junco hyemalis* was numerous. Miss Wittich & myself took several long walks, but saw, besides the above, only a few Robins, Blue jays, Ruffed Grouse, etc.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
2,004.	<i>Turdus pallasi</i> , Cabanis.	October 21st.	226 W. 15th St., N. Y. City.			♂ juv.

Mrs. Smith just brought me a Hermit Thrush (*Turdus pallasi*) which "Lopsy" (the cat) caught in the back yard. It is evidently a wild bird, and in beautiful plumage.
The bird was fat, and its bones soft, showing that it was a young bird.

October 26th. H.F., N.Y.
Returned from New York last night. To-day I saw the following birds: -
Turdus migratorius. Abundant; Feeding on the berries of the sour gum (*Nyssa multiflora*).
Turdus pallasi. Common; Seen with the Robins.
Regulus satrapa. Very common.
Regulus calendula. do
Sialia sialis.
Ampelis cedrorum. Abundant
Sitta carolinensis.
Parus atricapillus.
Junco hyemalis. Abundant
Spizella socialis. do
Spizella pusilla. do
Casserella iliaca. do
Zonotrichia albicollis. do

1879.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	<u>Corvus americanus</u> .					
	<u>Cyanurus cristatus</u> .					
	<u>Accipiter cooperi</u> . One or two seen.					
	<u>Bonasa umbellus</u> . -					
	<u>Larus argentatus</u> . Abundant on the Hudson.					
	<u>Sayornis busckii</u> .					
	Note - Louis Zerega informs me that there were numbers of Hermit Thrasher (<u>T. pallasi</u>) in the yards at the rear of No. 118 Waverly Place, in this city during last week.					

November 3rd. New York City

Went home Sat. afternoon by 6 o'clock train. On Sunday morning I walked to West Point, and back, before going to church. In the afternoon Rose & I walked down to the river. The following birds were observed.

Turdus migratorius; Lalia sialis; Regulus satrapa; Parus atricapillus; Sitta carolinensis; Certhia familiaris; Dendroica coronata, abundant; Ampelis cedrorum, in large flocks, by the river; Chrysomitris tristis; Junco hyemalis; Passer domesticus; Corvus americanus; Cyanurus cristatus; Picus pubescens; and Larus argentatus.

This morning I returned to New York by the early train. The mountains were white with snow, and the roofs & bare spots along the river were white also, as well as the fields lying back of Harrison. Large flocks of Robins were flying overhead, and some some Yellow-rumps were seen on the way down to the ferry.

I made the following observation on the rapidity of flight in the Long-tailed Duck (Harle glacialis) while coming down on the train: -

Just after leaving Sing Sing (about 8.43), a small flock of Old Wives rose from the water, and flew in a direct course down the Hudson. Their white

1879.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	colors were very distinctly visible against the deep blue of the river. I noticed that they maintained about the same relative position with regard to the train, which had not yet, however, gotten under full headway; but the birds changed their course at intervals, and, also, at times, they would rise in the air, separating from each other at those times, and then they would descend swiftly to near the surface of the water, and					

THE NEW YORK CENTRAL & HUDSON RIVER RAILROAD CO.

HUDSON RIVER AND HARLEM DIVISIONS.

Office of the GENERAL SUPERINTENDENT,

GRAND CENTRAL DEPOT,

J. M. TOUCEY,
Gen'l Superintendent.New York, Dec 20th 1880E. A. Mearns
Sir,

Referring to yours of this date relative to Train stops & distances Nov 3/79 -

The train alluded to left Garrison at 7.52 A.M. & Enig 8.30 & Sing 8.58 and the distance between Sing & Sing is

19 ⁵⁶/₁₀₀ milesJ. M. Toucey
Gen'l Supt.cedrorum, Melospiza melodia, Spizella monticola

As thus required would be. As our train ultimately for my to move indirectly, paying about in the middle of the day, in water. It so happened at which the train the gait at which the they kept their as long as our in we slowed up, Hudson, the Ducks 9.10 o'clock by my according to my the railroad - in guess. The distance, 20 miles.

Time from where weeks ago when I place, was just

y with me at home I made. The follow- migratorius, Lalia atricapillus, upon the rocks olivaceus, Ampelis

1879.

Collecting Number	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
----------------------	-------	-------	-----------	---------	----------	-----------------

Corvus americanus.

Cyanurus cristatus.

Accipiter cooperi. One or two seen.

Bonae umbellus. —

Larus argentatus. Abundant on the Hudson.

Sayornis suscu.

Note - Louis Zerega informs me that there were numbers of Hermit Thrushes (T. pallasi) in the yards at the rear of No. 118 Waverly Place, in this city.

Went home

On Sunday

back, before 9

Rose & I walk

birds were at

Indus miana

Parus stricklandi

lunulicris : ♀

adiposum, adiposum

ad a nam, in
tristis. Dura

Quercus, Pinus
Leaves, Branches

corolla and
tubecous: an

pure class, and
This is a pure

to his mother
travelling. The rest

train. She had
a lot of money

and the rock
white also

while also, as
Harrison &

Harrison's. La
head and

head, and so
with the head.

on the way, do
I will ~~be~~

I made the
list in it

of flight in the

while coming do

Just after tea

flock of Old

1879.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
-----------------------	-------	-------	-----------	---------	----------	-----------------

colors were very distinctly visible against the deep blue of the river. I noticed that they maintained about the same relative position with regard to the train, which had not yet, however, gotten under full headway; but the birds changed their course at intervals, and, also, at times, they would rise in the air, separating from each other at those times, and then they would descend swiftly to near the surface of the water, and then for some time the impetus thus acquired would carry them ^{onward} at an accelerated speed. As our train got fairly under way, and fortunately for my observations, the Ducks ceased to move indirectly, and skimmed along close to the water. ^{flying along in the middle of the river.} It so happened that the regular rate of speed at which the train moved was exactly equal to the ^{rate} gait at which the Ducks flew, and, accordingly they kept their relative position opposite to us as long as our course lay along the river. When we slowed up, before turning back from the Hudson, the Ducks flew ahead. The time was just 9:10 o'clock by my watch. The time given was according to my own piece and not that of the railroad - in fact I had set my watch by guess. The distance according to time-table is just 20 miles.

November 17th

Came down on early train. Time from where the flock of Ducks started, two weeks ago when I timed them, to the turning-off place, was just 27 minutes.

Dr. A. K. Fisher spent Sunday with me at home. No remarkable bird notes were made. The following birds were common. Turdus migratorius, Sialia sialis, Regulus satrapa, Parus atricapillus, Certhia familiaris - Climbing upon the rocks down by the River, Sitta carolinensis, Ampelis cedrorum, Melospiza melodia, Spizella monticola.

1879.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

Junco hyemalis, *Passerella iliaca* - numerous, *Passer domesticus*, *Chrysomitris tristis*, *Corvus americanus*, *Cyanurus cristatus*, *Picus pubescens*, *Accipiter Cooperi*, and *Larus argentatus*.

Note.

"October 31, 1874. - Hudsonian Godwit? Near the mouth of Consook Creek, I saw this bird standing out on the mud-flat, knee-deep in water. When fired at and wounded it flew about in circles, uttering a plaintive cry consisting of three syllables, loudly repeated; at last it fell into the river, but could not be reached."

Nov. 30th N.Y.N.Y.

Spizella socialis - seen yesterday, Nov. 29.

Passerella iliaca - seen Nov. 29th.

Catskill Mountains, New York. Dec. 2d.

2,005. *Astur atricapillus* 23.75 44.25 ♀ juv.
Wing, 13.10; tail, 11.00; culmen, .90; gape, 1.40; tarsus, 2.88; middle toe, 1.81; its claw, .77; inner toe, 1.15; its claw, 1.12; outer toe, 1.41; its claw, .63; hallux, 1.07; its claw, 1.19. Iris, yellow.

December 29, 1879. H.E.N.Y.

2,006. *Sialia sialis*. Bluebird. 7.05 12.30 ♂ ad.
Wing, 3.90; tail, 2.80; bill from nostril, .35; gape, .78; tarsus, .83; middle toe, .69; its claw, .26. Iris, hazel. Bill & feet, black. Inside of mouth, yellow.

Notes on Birds Seen.

Turdus migratorius. - A few near Fort Montgomery.
Sialia sialis. - Numerous near "

1879-'80.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

Regulus satrapa. Singing a sweet and very singular song.

Parus atricapillus. *Picus pubescens*.

Sitta carolinensis. *Certhia familiaris*.

Colaptes auratus. One or two seen near Ft. Montgomery.

Spizella monticola, *Junco hyemalis*.

Chrysomitris tristis. *Bonasa umbellus*.

The above were all seen by me on a walk down to Poppler Creek and back, this afternoon. The only other species of Birds seen since my arrival ^(December 31st) are *Bucephala clangula americana* and *Larus argentatus*.

Note.

January 2. Saw several Bonaparte's Gulls (*Larus philadelphia*), near Peekskill, on my way to New York.

In January I saw a Horned Grebe (*Podiceps cornutus*), near Riverdale, N. Y., and informed Mr. Bicknell of its occurrence there, at a meeting of the Linnaean Society.

2,007. *Larus glaucus*, Bränn. 29.00 67.00 immature.
Wing, 18.00; tail, 7.55; culmen, 2.50; bill from nostril, 1.10; gape, 3.70; tarsus, 2.88; middle toe and claw, 2.90; toe alone, 2.45; claw, .55. Bill, flesh-pink on basal one-half; succeeded by a wide band of blue-black; tip, horn-color. Legs and feet, pinkish white; nails blackish, tipped with horn-color. Irides, white.
I bought this bird of Mr. Robin, at Fulton Market, N. Y. He told me it was shot on Long Island, N. Y., and said it was a very rare bird, that not one Gull in a thousand was white above; that nearly

1880.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

all were saddle-backs. It is the immature bird, described as L. mitchinsii. All white.
Note.

Sayornis fuscus appeared here March 2nd, when Pat Cox saw a pair at the dock. I have seen none yet (March 5).

March 6th.

To-day the thermometer ranged from 60°-70°, and the weather was pleasant. In the morning, I saw turtles basking upon a rock in Pell's pond: both Spotted (Xanemus guttata) and Painted Turtles (Chrysemus picta). In the evening, at 6.30 o'clock, I saw a bat flying about. This is my first at home from the Medical College, and the following birds were seen: Turdus migratorius. Abundant; feeding on Rhus. Sialia sialis. Abundant; singing. Regulus satrapa. Several seen. Certhia familiaris. Carpodacus purpureus. Junco hyemalis. Singing. Spizella monticola. Singing. Melospiza melodia. Abundant; singing. Passer domesticus. About the house. Passerella iliaca. One that I shot was ♀. Corvus americanus. Cyanurus cristatus. Zenaidura carolinus. I shot one on the Cousook Island.

Colaptes auratus. One seen.

2008. Zenaidura carolinus. 12/10 18.85 ♂ ad.
Wing, 6.05; tail, 5.20; culmen, .50; bill from nostril, .36; gape, .80; tarsus, .86; middle toe & claw, 1.05; toe alone, .82; claw alone, .25. Iris, hazel. Skin about

1880.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

eye, blue. Legs and feet, red; nails brownish black. Bill, black. Crop contained triangular-shaped seeds which were preserved with sternum. Shot on Cousook Island.

March 9th.

Yesterday there was a light fall of snow, and somewhat colder weather. The following birds are some of those seen yesterday: Parus atricapillus.

Sitta carolinensis.

Chrysomitris tristis.

Anas obscura. - While I was at Pell's pond, six Dusky Ducks came, and, after circling a few times above the pond, gradually coming lower, they dropped almost vertically down to the middle of the pond, making a loud splash! They made a loud quacking when flying. At first they remained quiet in the middle, then swam swiftly about, flapping their wings and quacking, and at times uttering a curious clanging cry, till they saw me, when they raised with a splash and loud quacking, and left.

To-day the Fox Sparrows are still abundant, and I heard a Purple Finch sing. No birds about Cousook Island, except Song Sparrows and Crows.

March 10th.

Had a severe snow-storm yesterday, with high wind: thermometer 20° at 9 o'clock A.M. Snow 3 inches deep this morning. Flickers (Colaptes auratus) were calling loudly early in the morning, and Robins were abundant.

1880.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

(March 10th)
Snow nearly gone this afternoon.

Note. - Mr. Edward Conway, of 55 Carmine St., New York, showed me a skin of the Red-bellied Woodpecker (Centurus carolinus) which he shot at "Pine Island," (I found a "Pine Bush" near there on the map), on the Wall Kill, back of Newburgh, in Orange County, N.Y. It was one of a number that he shot there himself.

The Red-heads (Melanerpes erythrocephalus) are very abundant at the same place, and it may be that the same causes which prevent that bird from being common here, also keep the Red-bellied Woodpecker from visiting us. Cornwall is the only place near here where it has been taken.

March 11th.

Scolecophagus ferrugineus. - First seen
A flock (8) on Consook Island.
Weather cold. Snowed all day. Snow
4 inches this evening.

March 12th.

Anas obscura. - One seen in Pell's pond,
which was covered with thin ice.
Picus villosus. - Two were seen.

2,009. Picus pubescens. 6.75 11.90 ♀ ad.
Wing, 3.75; tail, 2.55; culmen, .69; gape, .78;
middle toe and claw, .60; tarsus, .61.

2,010. Picus villosus. 9.10 15.90 ♀ ad.
Wing, 4.80; tail, 3.45; culmen, 1.20;
gape, 1.30; tarsus, .83; middle toe & claw, .80.
Iris, hazel. Bill & feet, bluish. Claws,
benish-black.

1880.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

2,011. Corvus americanus, Audubon. March 13th. Consook Id. 18.10 34.50 ♀ ad.
Wing, 10.85; tail, 6.85; culmen, 1.73; gape, 1.98;
tarsus, 2.27; middle toe and claw, 1.80; toe alone,
1.40; claw, .50.

March 14th.

Passerella iliaca. - Singing
Carpodacus purpureus. - Singing.
Anas obscura. - One in Pell's pond.

March 15th.

Turdus migratorius. - First heard singing.
Ducks numerous in Pell's pond.
Mergus merganser. - One at Consook Island.

Note. - A Cow-Bird (Molothrus ater) was reared in
the nest of the European House Sparrow, in Highland
Falls, and lived for some time in the village,
amongst the Sparrows.

March 23.

2,012. Buteo pennsylvanica. - First seen: one pair.
Melospiza melodia. 6.27 8.40 ♂ ad.
Wing, 2.45; tail, 2.42; culmen, .46; gape, .55;
tarsus, .87; middle toe & claw, .85; toe alone,
.67; its claw, .23.
Anas obscura. - Found a flock in a swamp
on Mr. Tracy's place, where the rain has
settled. One was picked up on the salt
marsh, which had a hole eaten in its side;
and ^{through it} one side of the breast was eaten.
Found one in the same place & condition
two years ago.

March 24.

2,013. Passerella iliaca, (Merrem). 7.20 11.30 ♂ ad.
Wing, 3.39; tail, 2.88; culmen, .48; gape, .56; middle
toe and claw, 1.00; toe alone, .71; claw, .32.

1880.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
2014.	<i>Passerella iliaca</i> , (Merrem).	March 26th.	H.F.N.Y.	7.15	11.50	♀ ad.
	Wing, 3.55; 2.92=tail; culmen, .52; gape, .60; tarsus, .98; middle toe and claw, .96.					
2015.	<i>Spizella monticola</i> .	6.27		9.25		♀ ad.
	Wing, 2.88; tail, 2.83; culmen, .41; gape, .47; tarsus, .80; middle toe and claw, .76; toe alone, .56; claw, .24.					
2016.	<i>Cyanurus cristatus</i> .	12.20		16.75		♂ ad.
	Wing, 5.19; tail, 5.50; culmen, 1.08; bill from nostril, .70; gape, 1.19; tarsus, 1.45; middle toe and claw, — (shot off on both feet).					

March 28th.

Agelaius phoeniceus. — First seen.

Note. —

The following specimens were some of those collected in the year 1873. All were mounted, and not numbered or recorded in journals:—

I	<i>Cyanurus cristatus</i> .	Jan. 28, 1873.	12.00	17.00	—?
II	<i>Parus atricapillus</i> .	Jan. 31, 1873.	4.90	7.50	—?
III	<i>Bonasa umbellus</i> .	Feb. 15, 1873.	17.50	22.00	—?
IV	<i>Cyanurus cristatus</i>	Feb. 22, 1873.	11.00	15.70	—?
V	"	Mar. 3, 1873.	11.30	17.00	—?
VI	"	do	11.30	16.40	—?
VII	<i>Turdus migratorius</i> .	Mar. 15, 1873.	9.20	16.00	—?
VIII	<i>Ectopistes migratorius</i> .	Mar. 19, 1873.	16.30	23.70	♀ ad.
IX	<i>Melospiza meloda</i> .	Mar. 15, 1873.	6.60	8.70	—?
X	<i>Junco hyemalis</i> .	Mar. 17, 1873.	6.40	8.90	—?
XI	<i>Melospiza meloda</i> .	do	6.60	8.80	—?
XII	<i>Sialia sialis</i> .	Mar. 19, 1873.	7.20	12.10	♂ ad.
XIII	do	do	7.10	12.00	♀ ad.
XIV	<i>Mergus serrator</i> .	Mar. 24, 1873.	—	—	♀ ad.
XV	<i>Sitta carolinensis</i> .	Mar. 23, 1873.	6.00	10.70	♂ ad.
XVI	"	Mar. 28, 1873.	6.10	11.00	—?
XVII	"	Mar. 29, 1873.	5.60	10.30	—?
XVIII	<i>Scolecophagus ferrugineus</i> .	3, 1873	9.90	15.20	♂ ad.

1880.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
XIX	<i>Scolecophagus ferrugineus</i> .	Mar. 31, 1873.		9.90	15.00	♂ ad.
XX	<i>Lanius borealis</i> .	do		9.50	14.00	— ad.
XXI	<i>Larus argentatus</i>	Apr. 3, 1873.		24.00	59.50	♂ ad.
XXII	<i>Scolecophagus ferrugineus</i> .	Apr. 2, 1873.		9.60	15.20	—?
XXIII	<i>Anas obscura</i> .	Apr. 4, 1873.		34.70	36.20	— ad.
XXIV	<i>Sayornis fuscus</i> .	Apr. 12, 1873.		6.50	10.40	— ad.
XXV	<i>Turdus migratorius</i> .	do		10.40	—	— ad.
XXVI	<i>Trochilus colubris</i> .	May 17, 1873.		3.50	4.00	— ad.
XXVII	<i>Carpodacus purpureus</i> .	Apr. 14, 1873.		6.40	10.60	♂ ad.
XXVIII	<i>Sialia sialis</i> .	do		6.70	12.00	♂ ad.
XXIX	<i>Turdus pallasi</i> .	Apr. 16, 1873.		7.40	12.00	— ad.
XXX	<i>Poocetes gramineus</i> .	Apr. 19, 1873.		6.00	10.20	— ad.
XXXI	<i>Bucephala albeola</i> .	Apr. 23, 1873.		12.60	21.40	♀ ad.
XXXII	<i>Colaptes auratus</i> .	Apr. 24, 1873.		12.40	19.50	♂ ad.
XXXIII	<i>Ampelis cedrorum</i> .	Apr. 28, 1873.		7.00	11.80	♂ ad.
XXXIV	<i>Quiscalus purpureus</i> .	Apr. 29, 1873.		11.20	16.60	♀ ad.
XXXV	<i>Dendroica coronata</i> .	May 1, 1873.		6.00	9.00	♂ ad.
XXXVI	<i>Junco hyemalis</i> .	do		6.20	9.70	♂ ad.
XXXVII	<i>Carpodacus purpureus</i> .	May 3, 1873.		6.30	10.04	♂ ad.
XXXVIII	"	do		6.10	10.40	♀ ad.
XXXIX	<i>Zonotrichia albicollis</i> .	May 5, 1873.		6.50	9.70	♂ ad.
XL	<i>Scolecophagus ferrugineus</i> .	do		9.70	15.00	♂ ad.
XLI	<i>Pipilo erythrophthalmus</i> .	May 8, 1873.		8.00	11.00	♂ ad.
XLII	<i>Empidonax flaviventris</i>	do		5.00	—	—?
XLIII	<i>Setophaga ruticilla</i> .	May 10, 1873.		5.20	8.30	♂ ad.
XLIV	<i>Contopus virens</i> .	May 12, 1873.		5.60	8.50	— ad.
XLV	<i>Icterus baltimore</i> .	do		7.70	11.40	♂ ad.
XLVI	<i>Ampelis cedrorum</i> .	do		6.80	11.50	— ad.
XLVII	<i>Mniotilta varia</i> .	do		5.20	7.70	— ad.
XLVIII	<i>Icterus baltimore</i> .	May 13, 1873.		7.70	11.90	♂ ad.
XLIX	do	do		7.60	11.60	♂ ad.
L	do	do		7.50	11.30	♂ ad.
LI	do	do		7.10	11.00	♀ ad.
LII	<i>Goniaphea ludoviciana</i> .	do		8.00	12.60	♂ ad.
LIII	<i>Cyanospiza cyanea</i> .	do		5.70	8.50	♂ ad.
LIV	<i>Trochilus colubris</i> .	May 15, 1873.		3.50	4.00	♂ ad.
LV	<i>Harporhynchus rufus</i> .	May 17, 1873.		10.90	12.50	— ad.

1880.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
LVI	<i>Icterus spurius</i> .	May 19, 1873.		7.00	9.60	♂ ad.
LVII	<i>Mimus carolinensis</i> .	May 17, 1873.		9.00	12.00	♀ ad.
LVIII	<i>Icterus baltimore</i> .	May 19, 1873.		7.70	11.20	♀ ad.
LIX	<i>Dendroica castanea</i> .	do		5.70	9.00	♂ ad.
LX	<i>Icterus baltimore</i> .	do		7.60	11.00	♂ ad.
LXI	<i>Chaetura pelagica</i> .	May 20, 1873.		5.60	12.00	- ad.
LXII	<i>Pyrranga rubra</i> .	May 22, 1873.		7.20	11.80	♂ ad.
LXIII	<i>Setophaga ruticilla</i> .	do		5.20	8.20	♂ ad.
LXIV	<i>Pyrranga rubra</i> .	do		7.50	12.00	♂ ad.
LXV	<i>Sclerodactylus palustris</i> .	May 29, 1873.		5.10	6.20	♂ ad.
LXVI	<i>Tyrannus carolinensis</i> .	May 24, 1873.		8.10	14.30	♂ ad.
LXVII	<i>Circus olivaceus</i> .	June 2, 1873.		6.50	10.00	- ad.
LXVIII	<i>Ardea virescens</i> .	June 21, 1873.		5.50	5.50	♂ ad. ^{pullus}
LXIX	<i>Tyrannus carolinensis</i> .	June 28, 1873.		8.30	14.50	—
LXX	<i>Picus pubescens</i> .	June 30, 1873.		6.60	11.70	♂ ad.
LXXI	do	do		6.50	11.50	♀ ad.
LXXII	<i>Spizella pusilla</i> .	July 6, 1873.		5.90	7.50	♀ ad.
LXXIII	<i>Cyanospiza cyanea</i> .	do		—	—	♂ ad.
LXXIV	<i>Ardea virescens</i> .	June 10, 1873.		17.30	26.00	♀
LXXV	<i>Philohela minor</i> .	July 11, 1873.		10.50	19.00	♀ ad.
LXXVI	<i>Picus villosus</i> .	July 15, 1873.		8.60	15.60	♀
LXXVII	<i>Pyrranga rubra</i> .	July 14, 1873.		7.40	12.00	—
LXXVIII	<i>Ampelis cedrorum</i> .	do		7.20	11.80	—
LXXIX	<i>Myiarchus crinitus</i> .	July 15, 1873.		9.00	13.50	♂
LXXX	<i>Dolichonyx oryzivorus</i> .	July 25, 1873.		7.00	11.00	—
LXXXI	"	Aug. 18, 1873.		7.50	11.80	—
LXXXII	<i>Totanus solitarius</i> .	Aug. 18, 1873.		—	—	—
LXXXIII	<i>Cyanurus cristatus</i> .	Sep. 20, 1873.		11.50	16.60	—
LXXXIV	<i>Colaptes auratus</i> .	Sep. 24, 1873.		12.10	17.10	♀ ad.
LXXXV	<i>Myiarchus crinitus</i> .	do		7.50	13.00	♂ ad.
LXXXVI	<i>Cyanurus cristatus</i> .	Sep. 27, 1873.		11.50	17.20	♂ ad.
LXXXVII	do	do		11.90	16.40	- ad.
LXXXVIII	<i>Pyrranga rubra</i> .	do		7.00	11.20	♀
LXXXIX	<i>Pandion haliaetus</i> .	Oct. 1, 1873.		23.00	64.80	♂ ad.
LXL	<i>Buteo borealis</i> .	Oct. 11, 1873.		21.00	48.50	♂ ad.
43 LXL	<i>Pandion haliaetus</i> .	Nov. 25, 1873.		23.40	65.00	♀ ad.
LXLI	<i>Bucephala clangula</i> .	Dec. 5, 1873.		19.25	30.88	♀

1880.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
LXLIII.	<i>Fuligula affinis</i> .	Dec. 6, 1873.		16.10	28.30	—
LXLIV.	<i>Fuligula collaris</i> .	Dec. 16, 1873.		17.50	28.00	♂ ad.
LXLV.	<i>Haliaetus leucocephalus</i> .	Nov. 14, 1873.		36.90	80.00	♀ juv.
LXLVI.	<i>Scops asio</i> . (red)	Nov. 19, 1873.		9.40	22.00	♂ ad.

The above is a nearly complete list of all of the birds collected (all mounted) during the year 1873.

At the beginning of the year 1874 I commenced a cabinet of bird-skins as well as of mounted specimens. The following specimens of birds are among those collected during that year; most of the earlier ones were preserved as mounted specimens; most of the later ones as skins:

LXLVII.	<i>Spizella monticola</i> .	Jan. 28, 1874.		6.00	8.60	- ad.
LXLVIII.	<i>Sialia sialis</i> .	do		6.60	12.00	- ad.
LXLIX.	<i>Parus atricapillus</i> .	do		5.30	7.50	- ad.
C.	<i>Spizella monticola</i> .	Feb. 26, 1874.		6.12	9.19	♂ ad.
CI.	do	do		6.06	9.19	♀ ad.
CII.	<i>Melospiza meloda</i> .	do		5.44	8.00	♂ ad.
CIII.	<i>Junco hyemalis</i> .	Mar. 2, 1874.		6.09	9.88	♂ ad.
CIV.	<i>Sialia sialis</i> .	do		7.00	12.19	♂ ad.
CV.	<i>Passer domesticus</i> .	Mar. 5, 1874.		6.25	9.38	—
CVI.	<i>Cotopistes migratoria</i> .	do		16.00	23.25	♀ ad.
CVII.	<i>Sialia sialis</i> .	do		6.62	11.25	♀ ad.
CVIII.	do	do		6.75	12.12	♂ ad.
CIX.	<i>Passerella iliaca</i> .	Mar. 11, 1874.		7.50	11.38	♂ ad.
CX.	do	do		7.12	10.88	♀ ad.
CXI.	<i>Sialia sialis</i> .	do		6.56	12.25	♂ ad.
CXII.	do	do		6.50	12.38	♂ ad.
CXIII.	do	do		6.62	12.43	♂ ad.
CXIV.	do	do		6.56	12.31	♂ ad.
CXV.	do	do		6.56	12.43	♂ ad.
CXVI.	<i>Turdus migratorius</i> .	do		10.00	15.75	♀ ad.
CXVII.	<i>Junco hyemalis</i> .	Mar. 12, 1874.		6.00	10.81	♀ ad.
CXVIII.	<i>Passerella iliaca</i> .	do		7.25	11.00	♀ ad.
CXIX.	do	do		7.50	11.13	♂ ad.
CXX.	<i>Melospiza meloda</i> .	March 13, 1874.		6.06	8.69	- ad.

1880.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
CXXI.	<i>Bubo virginianus</i> .	Mar. 14, 1874.		21.00	52.50	♂ ad.
CXXII.	<i>Turdus migratorius</i> .	Mar. 20, 1874.		10.06	16.25	♀ ad.
CXXIII.	<i>Sialia sialis</i> .	Mar. 19, 1874.		6.75	12.62	♂ ad.
CXXIV.	<i>Junco hyemalis</i> .	Mar. 21, 1874.		6.13	9.44	♂ ad.
CXXV.	<i>Ampelis cedrorum</i> .	Mar. 20, 1874.		6.69	11.50	♀ ad.
CXXVI.	<i>Chrysomitris tristis</i> .	do		5.13	9.06	♂ ad.
CXXVII.	<i>Picus pubescens</i> .	Mar. 21, 1874.		6.88	12.00	♀ ad.
CXXVIII.	<i>Sayornis fuscus</i> .	do		7.00	11.44	— ad.
CXXIX.	<i>Junco hyemalis</i> .	do		6.06	9.25	♂ ad.
CXXX.	do	do		5.56	9.00	♀ ad.
CXXXI.	<i>Picus villosus</i> .	Mar. 25, 1874.		9.00	15.56	♂ ad.
CXXXII.	<i>Turdus migratorius</i> .	do		9.88	15.56	♂ ad.
CXXXIII.	do	do		9.25	16.50	♀ ad.
CXXXIV.	<i>Regulus satrapa</i> .	Apr. 7, 1874.		4.00	6.88	♀ ad.
CXXXV.	do	do		4.25	7.25	♂ ad.
CXXXVI.	<i>Regulus calendula</i> .	do		4.50	6.88	♂ ad.
CXXXVII.	<i>Regulus satrapa</i> .	do		4.25	7.25	♂ ad.
CXXXVIII.	do	do		4.00	6.75	♀ ad.
CXXXIX.	<i>Sayornis fuscus</i> .	Apr. 5, 1874.		7.00	11.13	— ad.
CXL.	<i>Anthus trogl. hyem.</i>	Apr. 27, 1874.		4.00	5.81	♀ ad.
CXLI.	<i>Ceryle alcyon</i> .	do		13.25	29.31	ad.
CXLII.	<i>Spizella monticola</i> .	Apr. 29, 1874.		6.13	9.31	♂ ad.
CXLIII.	<i>Spizella pusilla</i> .	do		5.56	8.38	— ad.
CXLIV.	<i>Spizella socialis</i> .	do		5.13	8.88	— ad.
CXLV.	do	do		5.13	8.81	— ad.
CXLVI.	do	do		5.25	9.00	— ad.
CXLVII.	do	do		5.25	9.06	— ad.
CXLVIII.	<i>Agelaius phoeniceus</i> .	do		9.06	15.00	ad.
CXLIX.	<i>Ectopistes migratoria</i> .	May 4, 1874.		15.50	23.75	♀ ad.
41	<i>Buteo pennsylvanicus</i> .	do		14.25	37.38	♀ ad.
42	<i>Pandion haliaetus</i> .	do		25.00	—	♀ ad.
CLII.	<i>Numenius longirostris</i> .	Aug 24, 1874.		24.00	40.00	♂ ad.
CLIII.	<i>Accipiter fuscus</i> .	Sep. 4, 1874.		11.50	—	♂ juv.
CLIV.	<i>Siurus aurocapillus</i> .	do		6.31	9.44	♂ ad.
CLV.	<i>Chrysomitris tristis</i> .	Sep. 3, 1874.		5.00	8.56	♂ ad.
CLVI.	<i>Sitta canadensis</i> .	Sep. 4, 1874.		4.75	8.25	♂ ad.
CLVII.	<i>Colaptes auratus</i> .	Apr. 14, 1874.		13.06	21.00	♂ ad.

1880.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
CLVIII.	<i>Turdus migratorius</i> .	Apr. 14, 1874.		9.25	14.62	— ad.
CLIX.	<i>Dendroica palmarum</i> .	do		5.56	8.13	♂ ad.
CLX.	<i>Molothrus ater</i> .	Apr. 16, 1874.		7.62	13.56	♂ ad.
CLXI.	<i>Picus pubescens</i> .	Apr. 14, 1874.		6.81	12.00	♂ ad.
CLXII.	do	do		6.81	12.00	♀ ad.
CLXIII.	<i>Regulus calendula</i> .	Apr. 16, 1874.		4.25	6.81	♂ ad.
CLXIV.	<i>Siurus motacilla</i> .	Apr. 15, 1874.		6.13	10.00	♀ ad.
CLXV.	do	do		6.19	10.13	♂ ad.
CLXVI.	do	Apr. 21, 1874.		6.13	10.62	— ad.
CLXVII.	<i>Regulus calendula</i> .	do		4.19	6.81	♂ ad.
CLXVIII.	<i>Siurus motacilla</i> .	Apr. 24, 1874.		6.06	9.94	♂ ad.
CLXIX.	<i>Turdus pallasi</i> .	do		7.25	11.62	♂ ad.
CLXX.	<i>Picus pubescens</i> .	Apr. 22, 1874.		6.75	11.88	♂ ad.
CLXXI.	<i>Zonotrichia albicollis</i> .	May 5, 1874.		6.75	9.75	♂ ad.
CLXXII.	do	do		6.50	9.25	♂ ad.
CLXXIII.	<i>Harporhynchus rufus</i> .	do		10.56	13.13	♂ ad.
CLXXIV.	<i>Coccyzus erythrophthalmus</i> .	Sep. 23, 1874.		11.62	15.50	♂ ad.
CLXXV.	<i>Coccyzus americanus</i> .	do		12.50	17.31	♀ ad.
CLXXVI.	<i>Pyrranga rubra</i> .	do		7.06	11.75	♂ juv.
CLXXVII.	<i>Pandion haliaetus</i> .	Sep. 8, 1874.		21.00	61.75	♂ ad.
CLXXVIII.	<i>Vireo noveboracensis</i> .	Sep. 2, 1874.		5.31	7.62	—
CLXXIX.	<i>Vireo olivaceus</i> .	Sep. 7, 1874.		6.00	10.00	—
CLXXX.	<i>Harporhynchus rufus</i> .	do		13.00	11.44	♂ —
CLXXXI.	<i>Porzana carolina</i> .	Sep. 8, 1874.		8.88	14.19	♂ ad.
CLXXXII.	do	Sep. 9, 1874.		9.38	14.00	♂ juv.
CLXXXIII.	<i>Agelaius phoeniceus</i> .	do		8.75	14.50	♂ —
CLXXXIV.	<i>Haliaeetus leucocephalus</i> .	Sep. 15, 1874.		32.25	84.00	♀ juv.
CLXXXV.	<i>Totanus solitarius</i> .	Sep. 16, 1874.		8.00	16.19	♂ —
CLXXXVI.	<i>Melospiza meloda</i> .	Sep. 23, 1874.		6.13	8.18	♂ ad.
CLXXXVII.	<i>Vireo olivaceus</i> .	do		5.94	9.75	♀ —
CLXXXVIII.	do	do		6.06	10.00	♂ —
CLXXXIX.	<i>Dendroica striata</i> .	do		5.69	9.19	♂ —
CLXL.	do	do		5.62	8.88	♀ —
CLXLI.	<i>Cyanurus cristatus</i> .	Sep. 25, 1874.		11.75	16.50	♀ —
CLXLII.	<i>Porzana carolina</i> .	do		8.12	12.88	♀ juv.
CLXLIII.	do	do		8.44	13.13	♀ juv.
CLXLIV.	<i>Agelaius phoeniceus</i> .	do		8.00	14.00	♂ juv.

1880.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
CLXLIV.	<i>Agelaius phoeniceus</i> .	Sep. 25, 1874.		8.44	13.38	♂ juv.
CLXLV.	do	do		7.62	12.19	♀ -
CLXLVI.	do	do		6.75	11.88	♀ -
CLXLVII.	<i>Melospiza melodia</i> .	do		—	—	—
CLXLVIII.	<i>Corvus americanus</i> .	Sep. 29, 1874.		20.00	37.19	♂ ad.
CLXLIX.	<i>Ceryle alcyon</i> .	do		12.69	21.13	♀ -
CC.	<i>Agelaius phoeniceus</i> .	Sep. 26, 1874.		7.75	12.50	♀ -
CCI.	<i>Spizella pusilla</i> .	Sep. 30, 1874.		5.62	7.69	♂ -
CCII.	<i>Empidonax</i> — ?	do		6.06	9.37	—
CCIII.	<i>Pipilo erythrophthalmus</i> .	do		8.37	11.13	♂ -
CCIV.	<i>Pyrenus pubescens</i> .	do		6.31	11.56	♂ juv.
CCV.	<i>Bonasa umbellus</i> .	Oct. 17, 1874.		16.88	23.00	—
CCVI.	<i>Cyanurus cristatus</i> .	Oct. 14, 1874.		12.00	17.13	♂ -
CCVII.	<i>Sphyrapicus varius</i> .	Oct. 15, 1874.		8.13	14.75	♀ -
CCVIII.	<i>Agelaius phoeniceus</i> .	do		8.06	12.75	♀ -
CCIX.	<i>Fringoides macularius</i> .	do		7.50	13.06	♂ -
CCX.	<i>Agix sponsa</i> .	Oct. 11, 1874.		17.81	28.44	♂ juv.
CCXI.	<i>Carpodacus purpureus</i> .	do		6.25	10.13	♂ ad.
CCXII.	<i>Ammodramus caudacutus</i> .	Oct. 16,		5.82	7.44	—
CCXIII.	<i>Melospiza palustris</i> .	do		5.62	7.56	♀ -

2,017. *Carpodacus purpureus*. April 2nd. H.F., N.Y. 6.35 10.45 ♂ ad.
Wing, 3.25; tail, 2.52; culmen, .45; gape, .55;
tarsus, .73; middle toe & claw, .70. Iris, hazel.

Quiscalus purpureus. April 3rd. H.F., N.Y. — A large flock and several small ones seen. They were the first seen this year.

Spizella pusilla. April 4th. — First seen.
Regulus calendula. — First seen; singing. In company with the Golden-crested (*R. satrapa*).

Spizella socialis. April 5th. — First seen.

1880.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	<i>Ceryle alcyon</i> .	—	First seen.			
	<i>Dendroica coronata</i> .	—	One seen: the first.			
	<i>Dendroica palmarum</i> .	—	First seen; numerous.			
	Iransacked the Consook Marsh, but saw no sparrows but <i>Melospiza melodia</i> .					
2,018.	<i>Dendroica palmarum hypochrysea</i> , Ridgway.					
	Wing, 2.77; tail, 2.24; culmen, .46; .563 8.85 ♂ ad. gape, .57; bill from nostril, .30; tarsus, .77; middle toe, .48; middle toe & its claw, .65. Iris hazel.					
2,019.	<i>Dendroica palmarum hypochrysea</i> , Ridgway.					
	Wing, 2.60; tail, 2.17; culmen, .42; .548 8.62 ♂ ad. gape, .52; bill from nostril, .28; middle toe, .50; middle toe and its claw, .67; tarsus, .78; Iris hazel.					

April 6th. H.F., N.Y.
2,020. *Sturnella magna*. 10.45 16.50 ♂ ad.
Wing, 4.78; tail, 3.00; culmen, 1.30; gape, 1.40;
tarsus, 1.69; middle toe, 1.18; middle toe & claw, 1.50
Iris, hazel.

The Meadowlark flew across to the garden and lit on top of the chestnut-tree at the foot of the garden. I went to the house, got my gun, and shot it. It was the first seen this year.

Ceryle alcyon. Several lit on the place.

April 7th
Constitution Island, Hudson, River, N.Y.
2,021 *Otus vulgaris wilsonianus*, Lesson. Long-eared Owl. Sternum preserved. 14.30 36.00 ♂ ad.
Wing, 11.00; tail, 6.00; culmen (measured from frontal feathers), 1.06; bill from nostril, .54; gape, 1.20; tarsus, 1.55; middle toe, 1.11; middle toe and its claw, 1.65. Irides, yellow.
Captured by Miss Anna B. Warner. She has a fragmentary egg in her cabinet, taken on the Island which probably belongs to this species. Dr. A.K. Fisher has an egg, taken at Sing Sing,

1880.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

N. Y., and identified by Prof. S. F. Baird.

April 8th.

To-night dissected a Fowl (*Gallus domesticus*), which contained an ovarian tumor weighing no less than $13\frac{1}{2}$ oz. It measures inches, and I have it preserved in alcohol; it was encapsuled, and solid. The fowl (♀ ad) weighed 6 lbs. Its ovary was diseased on one side, to which the ^{tumor} was attached by a pedicle; the other organs appeared to be quite normal, tho' the osseous structure was deformed and its body presented a square or flattened appearance in front, caused by the development and pressure outwards of the tumor. The Hen appeared to enjoy good health, and was killed for the table, and sold by the butcher.

My Preceptor, Dr. H. P. Chace also examined the specimen, and supposed that it might be a human uterus diseased, & enquired what it was, and was astonished to learn that such an mis case & other had been taken from a tree. The label slip in alcohol refers to this note.

April 10th.

Molothrus ater. - A small flock spent the day in the garden, and are the first seen this year, tho' the species must have arrived some time since.

April 11th.

Zenaidura carolinensis. - One seen in the evergreen woods, where nest I searched for, but unsuccessfully.

1880.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

Parus atricapillus. - A pair was seen near the river, which appeared to be engaged in building a nest in a very slender, dead sapling which looked much too small to accommodate their nest & young ones.

Ducks. -

I succeeded in crawling close upon a flock of Ducks, in Pell's Pond, most of which were Dusky Ducks (*Anas obscura*); but there was one Mallard (*Anas boschas*).

Turdus pallasi. - First seen - a single bird in a lonely spot in the deep woods.

Sayornis fuscus. - A nest was noticed upon the barn at the foot of the garden which appeared to be nearly or quite finished on yesterday - the 10th. To-day saw them gathering moss upon stones in the Morgan's brook.

April 12th.

2,022.

Colaptes auratus.

11:90 19:40 ♀ ad.

Wing, 5.88; tail, 4.15; culmen, 1.30; bill from nostril, 1.00; gape, 1.45; tarsus, 1.10; middle (3d) toe, .82; middle toe and its claw, 1.30.

Iris brownish-red. Shot in the pine-tree (*P. strobus*) beside the house.

2,023.

Molothrus ater.

7:22 12:05 ♀ ad.

Wing, 3.85; tail, 2.75; culmen, .57; gape, .66; tarsus, .95; middle toe, .65; middle toe & claw, .90.

April 13th.

Regulus satrapa } associated together; both singing.
Regulus calendula }

Zenaidura carolinensis. - Three flew up together from the Consook Marsh; one was shot.

Dendroica palmarum. - The first seen since the 5th. One shot at Pell's Pond.

1880.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

April 13th - continued.

Parus atricapillus. - The pair seen building on the 11th are still at work. One bird's head protruded from the hollow of the tree, while its mate sat in front on a branch and scolded, meanwhile holding its crest erect.

Corvus americanus. - Found a fresh nest on Cousook Island, in the top of a tall, very slender pitch pine tree (*P. rigida*). It was so windy that I feared the tree would not bear my weight, and I was obliged to leave it for another time.

2,024. *Dendroica palmarum*. 5.28 8.40 ♂ ad.

2,025. *Penædura carolinensis*. 11.60 17.50 ♀ ad.

Wing, 5.35; tail, 4.85; culmen, .57; gape, .77; tarsus, .86; middle toe, .75; middle toe and its claw, .92. Shot on Cousook Island.

April 14th.

Yesterday I heard the Yell. Red-bell Wren sing for the first time - a not very musical dittey. Also heard to-day.

Regulus calendula. - Shot a specimen with a ruby crown, which proved, on dissection, to be a female.

2,026. *Regulus calendula* (with red.) 4.60 7.15 ♀ ad.

2,027. *Regulus calendula* do 4.60 7.05 ♂ ad.

Gonotrichia albicollis. - First seen.

April 16th.

Corvus americanus. - Took five eggs from the nest on Cousook Island, and five others from a nest on shore.

Melospiza palustris. - First seen.

Branta canadensis. - A large flock.

1880.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

2,028. *Agelaius phoeniceus*. 9.75 15.40 ♂ ad.
Wing, 4.67; tail, 4.05; culmen, .94; gape, .95; tarsus, 1.16; middle toe, .76; middle toe + its claw, 1.05.

2,029. *Corvus americanus*, Audubon. 18.50 35.50 ♀ ad.
Wing, 12.20; tail, 7.00; culmen, 1.75; bill from nostril, 1.31; gape, 2.05; tarsus, 2.35; middle toe, 1.38; middle toe + its claw, 1.90. Iris, hazel.
Shot as she flew from her nest, on Cousook Island, from which I took 5 eggs - the first this year! Set

April 17th.

Pandion haliaetus carolinensis. - First seen.

April 18th.

Larus argentatus. - Several seen upon the Hudson.

Seiurus motacilla. - First seen. There were two pairs at Morgan's brook singing sweetly.

Corvus americanus. - Took five eggs from a nest in an oak-tree, on Gilbert's place; also found a nest in the top of a tall cedar, which contained but one egg.

Anas boschas. - Two seen in Tracy's swamp.

Buteo pennsylvanicus. - One sat near the road, and paid very little attention to me as I passed.

Otus vulgaris wilsonianus. - one seen this evening.

April 20th.

2,030. *Lurdus pallasi*. 7.35 11.95 ♀ ad.

Wing 3.75; tail, 2.77; bill from nostril, .38; gape, .84; middle toe, .75; middle toe + its claw, .97; tarsus, 1.22.

2,031. *Seiurus motacilla*. 6.28 10.65 ♂ ad.

Wing, 3.27; tail, 2.15; bill from nostril, .45; gape, .79; tarsus, .88; middle toe, .56; middle toe and its claw, .75.

1880.

- 2,032. *Sturnus motacilla*. 6.25 10.40 ♂ ad.
 Wing, 3.20; tail, 2.15; bill from nostril, .41;
 gape, .76; tarsus, .88; middle toe, .56;
 middle toe and its claw, .74. Iris, hazel.
 Legs & feet, flesh-color; before death they have
 a beautiful glow of pink. It was singing
 in a tree-top after seven o'clock P.M.
 They are usually musical ^{at night} late in the evening.
~~Anas~~ *Anas obscura*. - The two Dusky
 Ducks were seen this evening in Tracy's Swamp.
Dendroica palmarum. - ~~seen~~.

April 22d.

Fringoides macularius. - First seen.*Anthostomus vociferus*. - First seen.

April 23d.

- 2,033. *Agelaius phoeniceus*. 8.90 14.60 ♂ ad.
 Wing, 4.55; tail, 3.60; culmen, .85; gape, .90;
 tarsus, 1.10; middle toe, .80; middle toe and
 its claw, 1.05.

- 2,034. *Sturnus motacilla*. 6.20 10.30 ♂ ad.
 Wing, 3.15; tail, 2.05; bill from nostril, .44;
 gape, .75; tarsus, .84; middle toe, .58; middle
 toe and its claw, .75. Iris, hazel.

Bubo virginianus. - A pair were hooting this
 evening, on the mountain to the southeast.*Junco hyemalis*. - One seen.

April 24th.

Harporhynchus rufus. - First seen; singing.*Passerella iliaca*. - Last seen April 8th.*Dendroica virens*. - First seen.*Melospiza palustris*. Numerous.*Mniotilta varia*. - First seen.*Vireo solitaria*. - First seen.*Buteo pennsylvanicus*. Whistling.*Turdus palasi*. - Has not been very
 numerous yet.

1880.

- 2,035. *Melospiza palustris*. 5.90 8.00 ♂ ad.
 Wing, 2.33; tail, 2.20; culmen, .46; gape, .49;
 tarsus, .84; middle toe, .61; middle toe and its
 claw, .85.
- 2,036. *Harporhynchus rufus*. 11.85 13.75 ♂ ad.
 Wing, 4.15; tail, 5.25; bill from nostril, .83;
 gape, 1.40; tarsus, 1.87; middle toe, .91; middle
 toe and its claw, 1.21.
- 2,037. *Vireo solitarius*. 5.65 9.12 ♀ ad.
 Wing, 2.85; tail, 2.27; bill from nostril, .30;
 gape, .60; tarsus, .73; middle toe, .45; middle
 toe and its claw, .63.
- 2,038. *Dendroica virens*. 5.25 7.80 ♂ ad.
- 2,039. *Dendroica virens*. 5.10 7.85 ♂ ad.

April 25th.

- 2,040. *Buteo pennsylvanicus*. 16.00 35.00 ♂ ad.
 Wing, 10.35; tail, 6.80; culmen from cere, .74;
 cere, .48; from frontal feathers to tip of bill, 1.10;
 tarsus, 2.50; middle toe, 1.33; middle toe & its
 claw, 1.84; claw alone, .65. Irides yellow,
 except a trace of the brown of the im-
 mature bird. Cere, greenish-yellow. Tarsi
 and toes, yellow. Bill & claws, blue-black.
 Feet muddy. Beneath its eyelid were some
 long, slender white ^{parasitic} worms, one of which is preserved
 in alcohol. Its stomach contained hair
 of mice and moles, and some coleopterous
 insects. Sternum preserved. Do not
 remember ever to have found remains
 of birds in their stomachs. This bird was
 in very poor, emaciated condition, as
 they usually are at this season.

1880.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

April 26th.

Coccyzus erythrophthalmus. - First seen.*Empidonax minimus*. - First seen.*Hirundo horreorum*. - First seen.*Bubo virginianus*. - At noon it was dark and cloudy, and I heard the Owl screech or hoot then.*Buteo pennsylvanicus*. - One pair flying overhead, uttered two or three screams in rapid succession, and then their usual squealing or whistling note.2,041. *Empidonax minimus*. 5:70 8:50 ♀ ad.

Wing, 2.57; tail, 2.41; bill from nostril, .31; gape, .65; tarsus, .65; middle toe, .32; middle toe and its claw, .49.

April 27th.

To-day took a long walk with Mr. Dr. Wheatley; shot a Winter Wren & White-throated Sparrow.

Hirundo horreorum. - Not numerous yet.*Chaetura pelagica*. - First seen.2,042. *Molothrus ater*. 7:10 12:30 ♀ ad.

Wing, 3.85; tail, 2.70; culmen, .62; gape, .62; tarsus, .94; middle toe, .60; middle toe and its claw, .89.

2,043. *Molothrus ater*. 7:03 12:00 ♀ ad.

Wing, 3.80; tail, 2.60; culmen, .60; gape, .60; tarsus, .90; middle toe, .65; middle toe and its claw, .90.

2,044. *Zonotrichia albicollis*. 6:07 10:00 ♂ ad.

Wing, 3.10; tail, 3.11; culmen, .50; gape, .54; tarsus, .90; middle toe, .66; middle toe and its claw, .88.

2,045. *Geothlypis troglodytes hyemalis*. 3:95 5:70 ♀ ad.

Wing, 1.65; tail, 1.05; bill from nostril, .32; gape, .59; tarsus, .65; middle toe & its claw, .63.

1880.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

April 28th.

Troglodytes domesticus. - First seen.*Pipilo erythrophthalmus*. - First seen.*Sialia sialis*. - A male attacked a Red Squirrel and drove it from the vicinity of his nest. There are young ones in the nest.2,046. *Colaptes auratus*. 12:85 21:00 ♀ ad.

Wing, 6.35; tail, 4.55; culmen, 1.47; bill from nostril, 1.12; gape, 1.58; tarsus, 1.18; middle toe, .92; its claw & toe together, 1.33.

April 30th. N.H., N.Y.

2,047. *Spizella pusilla*. 5:65 8:25 ♂ ad.

Wing, 2.52; tail, 2.70; culmen, .40; gape, .42; tarsus, .74; middle toe, .50; middle toe and its claw, .65. Iris, red.

2,047a. *Hirundo erythrogaster*. 6:62 12:75 ad.

Wing, 4.53; tail, 3.15; bill from nostril, .24; gape, .62; tarsus, .47; middle toe, .48; middle toe and its claw, .68. Not skinned.

2,047b. *Dendroica palmarum*. 5:60 8:40 ad.

Wing, 2.55; tail, 2.18; bill from nostril, .31; gape, .57; tarsus, .82; middle toe, .48; middle toe and its claw, .67. Not skinned.

May 1st.

Turdus mustelinus. - First seen. One was singing as I walked through Kinsley's cornfield from West Point.*Quiscalus purpureus* et var. *amertus*. - Took stage from Newburgh, to Cornwall. Woodna Creek bridge was being rebuilt, so stage went via Vail's Gate. Saw numbers of Grackles of var. *purpureus* et *amertus*. Of all the horribly rickety stages, that one was the worst I have ever ridden in. On top of it were piled some long, bent, heavy iron

1880.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

rods, which rattled dangerously overhead. The horses were lean, - in fact were skeletons, and as the driver whipped 'em up the long hills, I could not help feeling that I was doing violence to my humanity by sitting on the driver's seat instead of getting out and helping the horses pull, as my conscience told me I ought to be doing. The roads and fences were lined with sturdy old cedars, and in these the grackles were ensconced. As we drove past almost brushing the branches, they hopped down close to us and leered at the bony nags and peered into the coach and screamed derisively at us and spread their tails in high glee as they called to their neighbors in advance to join in the merriment at our expense; and they in turn peered us as we passed. I could not help feeling ashamed, and involuntarily looked at the driver to see if he was not blushing at being so ridiculed.

They were scattered over the fields on both sides of the road, and the bronzed variety shone like gold in the sunlight, while the purple ones glittered brilliantly in their metallic plumage. I have never known them to breed in the mountains; but in all the low-lying meadow country along the Hudson they do so abundantly - especially where coniferous trees are abundant.

Tachycineta bicolor. - First seen.

Nyctardea grisea naevia. - Visited their Roosting on Constitution Island, and found them abundant. The flagman said they had been there three weeks.

1880.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

Pandion haliaetus carolinensis. - One seen at the Island (Constitution).

Corvus ossifragus. - One seen at Cornwall. I got out of the stage at Dan Ward's. Found the old gentleman very ill, and his son Dan was out shad-fishing. I wanted to see a Black-headed Gull in their house, but did not like to ask the daughter; so walked around the bay to the Cornwall landing. On my way heard note which I recognized to be that of a Fish Crow. I saw the bird coming towards me, and it passed directly over my head so that I had a distinct view of it. Its note was unmistakable; but I longed for a gun to make assurance doubly sure. My first (only previous) record was also early in May.

Note. [May 1st.]

After dining with Mr. Roe at Cornwall, took steamer "Boardman" to Cold Spring, where I landed and called upon Mr. Butterfast, a barber, who has a small collection of mounted birds which he has made. He took me up stairs, and was extremely polite, and I saw the following very interesting specimens which he took himself at Cold Spring: -

Larus atricilla. - One specimen in his collection shot in Spring, several years ago, on the river.

Another one he mounted a few weeks since for Mr. Clark (boat-builder at Cornwall, where the specimen was shot). He promised to show me the letter which Mr. Clark sent him accompanying the bird, but it was forgotten; so I did not learn the date.

Mr. Butterfast saw another specimen several springs ago, on the

1880.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

Hudson at Cold Spring. It was so tame that he tried to strike it with an oar. It then flew and alighted on the buoy.

Haliaeetus leucocephalus. - I was shown a female that Mr. B. said was 9 ft 2 in in extent of wings; and the bird certainly was an unusually large one.

Mr. B. showed me four eggs which he said were those of this Eagle; he took them from the nest himself the first Sunday in May, 1877. There were no less than five eggs in the nest. He had given one away. He carried a revolver to the nest, but had no occasion to use it for the birds did not come about.

He says that White-headed Eagles build in a cave in Bull Hill, north of Cold Spring.

Centurus Carolinus. - A specimen in his collection, he shot in Mr. Dr. Rham's woods at Cold Spring. He never saw another.

Melospiza palustris. - One about one-half white.

Gallinula chloropus galeata. - Mr. B. has a specimen which he shot beside the river at Cold Spring.

1880.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

May 2nd.

Dendroica palmarum. - Still present.

Regulus calendula. - Still abundant.

Setophaga ruticilla. - First seen; abundant; singing.

Myiodynastes virens. - First seen; singing.

Helminthophaga ruficapilla. - First seen; abundant.

Mimus carolinensis. - First seen; several.

Vireo olivaceus. - First seen.

Vireo gilvus. - First seen.

Larus argentatus. - One seen.

Icterus baltimore. - First seen.

May 3d.

Went shooting & was joined by Dr. Wheatley in the woods, who came in search of me for a tramp. We stayed in the woods till after dark, and heard the Large-billed Accutor (Sialia motacilla) singing.

Chordeiles virginianus. - First seen.

Vireo flavifrons. - First seen.

Dendroica caerulescens. - First seen.

Dendroica pensylvanica. - First seen.

Sialia naevius. - First seen.

Sialia aurocapillus. - First seen.

Icterus spurius. - First seen.

Dendroica palmarum. - One seen.

Regulus calendula. - Still numerous.

2,048. Chaetura pelagica. 5:50 12:20 ♂ ad.

Wing, 4.90; tail, 1.95; bill from nostril, .15; culmen, .21; gape, .58; tarsus, .50; middle toe, .33; middle toe and its claw, .45.

2,049. Dendroica caerulescens. 5:45 8:25 ♂ ad.

Turdus fuscescens. - First seen.

1880.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

2,050. May 4th.
Goniaphea ludoviciana. 8:15-13:25 Bad.
Wing, 4'05; tail, 3'05; culmen, '72; gape, '79;
tarsus, '92; middle toe, '62; middle toe
and its claw, '83.

Regulus calendula. Still present.

Pyranca rubra. - First seen.

Goniaphea ludoviciana. - First seen.

Eagle. - Mr. Edward S. Denton and myself, accompanied by the Rev. Dr. Wheatley, spent a number of hours in making a thorough exploration of the Cro' Nest cliff and valley in search of the nest of the White-headed Eagle which Mr. Butterfass asserted to exist in a certain large hemlock (Abies canadensis) which he pointed out, near the shoulder of the mountain. There is no nest upon that tree nor any other one upon the mountain; but we saw the Eagle, and it doubtless has a nest upon the ledge somewhere along one of the cliffs. We visited the swamp beneath the highest point of the mountain. The Mr. Denton descended at the foot of the cliff, and I at the top, all the way to the river. It was a tough scramble. We set out at 1 o'clock P.M. & reach West Point again at 8 o'clock. Reached home at eleven, having supped at the Point with the others. Bad bruise & loss of integument on shin, and no Eagle's eggs. No Duck Hawks were seen. The Purple Trillium or Birthroot (Trillium erectum), and a Dicentra were flowering abundantly upon the mountain.

1880.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

May 5th.
Icteria virens. - First seen.

May 6th.
Parula americana. - First seen.

May 7th.
Cyanospiza cyanea. - First seen.

May 8th.
Sitta carolinensis. Still present.
Myiarchus crinitus. - First seen.
Tyrannus carolinensis. - First seen.

May 9th.
Lurdus swainsoni. - First seen.
Dendroica aestiva. - First seen.
Dendroica maculosa. - First seen.
Dendroica striata. - First seen.
Myiodiptes canadensis. - First seen.
Helminthophila vermivorus. - First seen.
Geothlypis trichas. - First seen.

May 10th.
Sitta carolinensis. - Heard to-day
Regulus calendula do
Coccyzus americanus. - First seen.

May 13th.
A list of all the birds seen:-
Lurdus migratorius
Lurdus multicolor
Lurdus swainsoni
Lurdus fuscescens
Minus carolinensis
Harporhynchus rufus.

1880.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	<i>Sialia sialis.</i>	<u>May 13th - continued.</u>				
	<i>Parus atricapillus.</i>					
	<i>Sitta carolinensis.</i>	- One is probably breeding near here, as it is seen constantly.				
	<i>Troglodytes domesticus.</i>					
	<i>Mniotilta varia.</i>					
	<i>Parula americana.</i>					
	<i>Helminthophaga vermivorus.</i>	- Quite numerous.				
	<i>Helminthophaga chrysoptera.</i>	- Must be quite numerous, for a number were seen and heard; first seen				
	<i>Helminthophaga ruficapilla.</i>	- Seen in apple-trees.				
	<i>Dendroica virens.</i>					
	<i>Dendroica coerulescens.</i>					
	<i>Dendroica striata.</i>					
	<i>Dendroica maculosa.</i>					
	<i>Dendroica pennsylvanica.</i>					
	<i>Dendroica aestiva.</i>					
	<i>Sinus aurocapillus.</i>					
	<i>Sinus naevius.</i>					
	<i>Sinus motacilla.</i>	- Searched fruitlessly for nest.				
	<i>Geothlypis trichas.</i>					
	<i>Icteria virens.</i>					
	<i>Myiodioetes nigratus.</i>					
	<i>Myiodioetes pusillus.</i>	- Several seen; singing; first seen				
	<i>Myiodioetes canadensis.</i>	- Numerous; singing.				
	<i>Setophaga ruticilla.</i>					
	<i>Pyrrhula rubra.</i>	- Abundant.				
	<i>Hirundo horreorum.</i>					
	<i>Amphisp. cedrorum.</i>					
	<i>Vireo solitarius.</i>					
	<i>Vireo flavifrons.</i>					
	<i>Vireo noveboracensis.</i>	- First seen.				
	<i>Vireo gilvus.</i>					
	<i>Vireo olivaceus.</i>					
	<i>Carpodacus purpureus.</i>					
	<i>Chrysomitris tristis.</i>	- Abundant; singing; in full summer plumage.				

1880.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	<i>Melospiza melodia.</i>					
	<i>Spizella socialis.</i>					
	<i>Spizella pusilla.</i>					
	<i>Zonotrichia albicollis.</i>					
	<i>Passer domesticus.</i>					
	<i>Cyanospiza cyanea.</i>					
	<i>Pipilo erythrophthalmus.</i>					
	<i>Icterus spurius.</i>					
	<i>Icterus baltimore (= galbula).</i>					
	<i>Corvus americanus.</i>					Very busy feeding young, which are nearly out of nest.
	<i>Myiarchus crinitus.</i>					
	<i>Tyrannus carolinensis</i>					
	<i>Sayornis fusca</i>					
	<i>Contopus virens.</i>					First seen.
	<i>Empidonax minimus.</i>					
	<i>Empidonax flaviventris.</i>					First seen.
	<i>Empidonax traillii</i>					First seen.
						Inhabits swampy fields grown up to alders and has curious note.
	<i>Antrostomus vociferus.</i>					
	<i>Chordeiles virginianus.</i>					
	<i>Chaetura pelagica</i>					
	<i>Irochilus colubris.</i>					
	<i>Coccyzus erythrophthalmus.</i>					
	<i>Coccyzus americanus.</i>					
	<i>Ceryle alcyon.</i>					
	<i>Picus pubescens.</i>					
	<i>Colaptes auratus.</i>					
	<i>Fringoides macularius.</i>					
	<i>Ardea virescens.</i>					
2,051.	<i>Empidonax traillii.</i>			6.03	9.40	♂ ad.
	Wing, 2.85; tail, 2.31; bill from nostril, .35; gape, .70; tarsus, .66; middle toe, .42; middle toe and its claw, .58.					
2,052.	<i>Coccyzus erythrophthalmus.</i>			12.25	16.70	♀ ad.
	Wing, 5.62; tail, 6.28; bill from nostril, .74;					

1880.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
				gape, 1.27; tarsus, 1.02; middle toe, .79; middle toe and its claw, 1.06.		
2,053.	<u>Helminthophaga chrysoptera</u>	5.02	7.75	♂ ad.		
	wing, 2.45; tail, 2.02; bill from nostril, .35; gape, .56; tarsus, .66; middle toe, .41; middle toe and its claw, .57.					
2,054.	<u>Helminthosus verrucosus</u>	5.76	9.12	♂ ad.		
	Wing, 2.82; tail, 1.95; bill from nostril, .45; gape, .71; tarsus, .71; middle toe, .53; middle toe and its claw, .67.					
2,055.	<u>Myiodiectes pusillus</u>	4.88	6.57	♀ ad.		
	Wing, 2.02; tail, 1.85; bill from nostril, .25; gape, .50; tarsus, .72; middle toe, .46; middle toe and its claw, .58.					
2,056.	<u>Myiodiectes pusillus</u>	5.00	6.86	♂ ad.		
	Wing, 2.10; tail, 1.88; bill from nostril, .25; gape, .50; tarsus, .74; middle toe, .46; middle toe and its claw, .57.					
2,057.	<u>Dendroeca maculosa</u>	5.02	7.40	♂ ad.		
	Wing, 2.26; tail, 2.05; bill from nostril, .27; gape, .52; tarsus, .71; middle toe, .40; middle toe and claw, .56.					
2,058.	<u>Tyranga rubra</u>			♂ ad.		

May 15th.

Sitta carolinensis. - One seen; doubtless breeding.

May 20th.

Sitta carolinensis. - Have noticed one above my grandfather's stable, where it is probably breeding.

May 25th.

Sitta carolinensis. - One near the house, & frequently seen in various places. They are undoubtedly breeding in this vicinity.

1880.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
2,059.	<u>Anas boschas</u>	May 1st.	?	—	—	—
<p>Head turning, only, preserved. Mr. Francis Butterfass gave me the specimen, & said it was a wild duck which had been wounded, on the Hudson, near Cold Spring; and he had kept it alive in his yard till May 1st, when he killed it for the table.</p>						
		June 5th.				
2,060.	<u>Bonasa umbellus</u> .					pullus.
2060.1.	<u>Measurements of Nyctiardea grisea naevia</u> :-					
	1. <u>Wing</u> , 24.25; <u>tail</u> , 49.50; 24.25 44.50 ♀ ad.					
	<u>Wing</u> , 11.40; <u>tail</u> , 4.50; <u>culmen</u> , 2.85; <u>gape</u> , 3.90; <u>tarsus</u> , 3.23; <u>middle toe</u> , 2.96; <u>its claw</u> , .62; <u>middle toe and its claw</u> , 3.57.					
2060.2.	<u>Nyctiardea grisea, naevia</u> . ♂ ad.					
	2. <u>Wing</u> , 11.60; <u>tail</u> , 4.55; 25.90 45.25 ♂ ad.					
	<u>culmen</u> , 3.10; <u>gape</u> , 4.10; <u>tarsus</u> , 3.20; <u>middle toe</u> , 2.92; <u>its claw</u> , .62; <u>middle toe and its claw</u> , 3.52. Shot on Constitution Id., by Louis A. Perezga, and skinned by him also.					

2061. Trochilus colubris. 3.75 4.12 ♂ ad.
Wing, 1.52; tail, 1.20; culmen, .67; gape, .80;
tarsus, .18; middle toe and its claw, .24.

This Hummingbird killed itself by flying against a telegraph wire.
Sitta carolinensis. - Frequently seen about.

June 22 d.

To-day is the longest of the year. Return-
ing from New York by rail, I saw
the last of the sun as it disappeared
behind the mountain west of Fort
Montgomery, at 7.20 P.M.
Saw a White-headed Eagle (Haliaeetus
leuccephalus) on Long Id., going down to N.Y.

1880.

2062. Otus vulgaris wilsonianus. 14.90 37.75 ♀ ad. ^{June 23d}
 Wing, 11.00; tail, 6.40; culmen measured
 from frontal feathers, 1.15; measured
 from cere, .70; gape, 1.32; tarsus, 1.68;
 middle toe and its claw, 1.85. Iris, yellow.
 To-day rowed, in the afternoon, with
 Dr. Wheatley up to Canatutuon Island.
 We visited the Herons, and found that
 most of the young ~~herons~~ had left
 the nests; but a considerable number
 were still calling in the nests. The
 noise of the young half-grown birds
 is most singular. The adults were
 numerous and very noisy. Young
Ruffed Grouse were numerous; they
 were already on wing June 5th, when
 Louis A. Jerega visited the Herons; it
 procured a specimen, which I skinned.
 We came upon a family of Long-eared
 Owls (Otus vulgaris wilsonianus). They
 appeared to see very well though
 the sun was dazzlingly bright. They
 sat erect and frequently flew a short
 distance. I shot one, and the rest
 all flew away & could not again
 be found.

July 6th.

Sitta carolinensis. - Constantly present.
Dendroica virens. - Saw a pair that, from
 their actions, I have no doubt were near their
 nest or young one. They were in the hemlock
 trees, just south of Davisons.

July 25th.

To-day (Sunday) walked to Highland Lake,
 around the lake, and so home. In the
 old fort to the north of Popple Creek there

1880.

a young Black-throated Green Warbler (Dendroica
 virens); and, while calling it a fine Worm-
 eating Warbler came close to where I stood.

A pine Woodchuck (Arctomys monax) was seen
 close to the shaft of a mine.

During the night, I heard the Night
 Herons (Nyctiardea grisea naevia) several times.

July 17th.

2063. Myiodytes mitratus. 5.65 8.00 ♀ ad.
 Heard Night Herons again last night.
 Wing, 2.50; tail, 2.32; bill from nostril, .31;
 gape, .60; tarsus, .76; middle toe and claw, .60;
 middle toe alone, .44.
 It was picked up dead in the main hall, at
 Cozzens West Point Hotel.

Note.

"Large-billed Water Thrush. - The nest of this
 species I discovered in a marsh under some
 low bushes. It was situated in a tuft of sword
 grass one foot from the ground. The nest is
 somewhat elongated, composed of sword grass
 principally, and lined with fine grass and
 hairs. It was six inches long, and contained
 four eggs of the bird and two Cow Bird's eggs.
 The ~~eggs~~ were so tender that I broke them all
 but one. I judge it to be about .65 by .50 inch.
 W. H. Ballou." < Oölogist, vol.

IV, No. 4, p. 27, 1878.

July 27th

To-day visited Indian Falls with a party of
 Cadies. Coming out of the creek, a Rail
 flew up and I had a distinct view
 of it. It was Rallus virginianus, which
 is probably the species whose nest

1880.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

Miss Anna Warner has, from the same marsh. Helminthophaga ruficapilla. - A very pretty Nashville Warbler was seen in a pear-tree in my garden.

July 28th.

Went this afternoon in search of the Virginia Rail seen yesterday. Waded all through the marsh but did not see so much as a feather of it.

Ardea herodias. - Shot at a fine adult specimen that was dancing about and flapping its wings in the most lively manner amongst a lot of lily-pads. Another bird was seen; for a long time past they have been seen flying towards the mountain at dusk - all summer in fact. The species is resident in summer, but its nest has not been yet detected.

2,064 Nyctiardea grisea naevia. - - - - - ♀ juv.

The Night Herons both young-of-the-year, and adults were quite numerous in the Heronry on Constitution Id. Having shot two of the young, I ascended to a high tree-top and viewed the swamp from my lofty perch, and saw a very interesting sight. The crouching forms of the young birds, so hard to get sight of from the ground were seen distinctly, and in the pine-trees farther off a number of old & young ones were sitting, while other old birds flew in wide circles overhead.

I shot a fine Dove (Zenaidura carolinensis) on the R.R. track, and a pair each of Cave and Bank Swallows. We left the marsh at dusk when the Night Herons were beginning to come out and feed on the flats. They feed there all night on moonlight nights and

1880.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

come to the flats in large numbers just at daylight - especially should the tide be out, leaving their prey exposed. Bats were very numerous; as we rowed across the cove at least 50 could be seen at once. I shot one.

July 29th.

Helminthophaga ruficapilla. - One in a swamp near Cousack Island.

Helminthosus vermivorus. - One in the swamp. Came at call and staid near me some time; flew into a tree top and rattled a bunch of dry leaves, probably searching for food. Its note was sharp and metallic.

August 7th

Went shooting with A. K. Fisher. Shot four specimens of Myiodiodes mitratus, and one Sinus notacilla.

Helminthosus vermivorus is quite numerous.

August 9th

2,065. Pipilo erythrophthalmus. Iris hazel. - - - - - ♂ juv.

2,066. Contopus virens. 6.60 10.90 ♂ ad.

Wing, 3.40; tail, 2.65; culmen, .65; gape, .83; bill from nostril, .42; tarsus, .50; middle toe, .32; middle toe and claw, .48.

Nyctiardea grisea naevia. Saw it at Pell's Pond.

Totanus solitarius. - Saw tracks which were probably its, at Pell's Pond.

Sinus notacilla. One seen at Pell's Pond brook.

August 11th.

2,067. Myiodiodes canadensis. ♀

Porzana carolina. - One shot: the first seen this year. The only one seen, at Cousack Philohela minor. Cousack swamp.

1880.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

August 24th.

Pandion haliaetus carolinensis. First seen since Spring.

August 30th.

Myiiodiodes nitratus. - A fine male seen.Nyctiardea grisea naevia. - Several were seen on August 33d at "Echo Lake", behind my house.

September 1st.

Dendroica palmarum. - One singing its odd little song in a pear-tree in my garden.

September 7th.

Geothlypis colubris. - Seen at Highland Falls, and quite numerous upon the marshes near Constitution Island.Ardea herodias. - About a dozen arose at the report of the first gun we fired in the lower cove. Several lit on trees in plain view of our boat, and one succeeded in establishing itself upon the very summit of a high hemlock tree after flapping and balancing for a long time. I attempted to stalk them, while Mr. Wheatley lay ambushed in the boat drawn up under the low branches of trees that shaded the creek; he thus procured two shots, both of them unsuccessful.

One Great Blue flew out of the Night Heronry on the island just at dusk and flew down upon the flats.

Haliaeetus leucocephalus. - Only one, an adult bird seen in the lower cove.

1880

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

Buteo pennsylvanicus. One flew close over my head, and lit a little way in advance; my gun was loaded with "B" and "dust" shot; I fired the dust at the Hawk as it close to me upon a limb eyeing me curiously. I afterward shot it again; it fell in the trees and could not be recovered.Scolichonox oryzivorus. - Still very plentiful.Ardea virescens. - Numerous.Sitta canadensis. - One bird.Syrnium nebulosum. - One heard.Scops asio. - Abundant; heard at night.Helmatodytes palustris. - Still very plentiful.and very noisy. Found in dry ^{part of} meadows.Antrostomus vociferus. - Heard to night; has been heard regularly of late.

September 8th.

Turdus swainsoni. - One shot - the first seen.Dendroica virens. -Dendroica maculosa. - First seen. One shot.Dendroica pennsylvanica. - One shot.Geothlypis trichas.Siurus naevius. - Several seen at Pell's Pond and along the shore of Cousack Marsh.Setophaga ruticilla. - An adult male in full plumage was seen.Pyrranga rubra.Botaurus minor. - I fired a shot at a bird which I believe to have been a Bittern that was flying before a high wind over Cousack Marsh.Accipiter fuscus. Common now.Buteo pennsylvanica.Pandion haliaetus carolinensis. One seen.Ardea virescens.Ardea herodias. - Plenty of its voice.

1880.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

tracks about Pell's Pond.

Totanus solitarius. - Plenty of tracks, but no birds were about Pell's Pond when we visited first, but, on returning we saw one just alighting; but it flew off the moment I approached; but I succeeded in shooting its mate. The first bird flew very high and crookedly. When migrating they fly very high.

2,068. Totanus solitarius. 8.60 16.00 ♂(ad?).

Wing, 5.03; tail, 2.20; culmen, 1.15; bill from nostril, .96; gape, 1.27; tarsus, 1.25; middle toe, .95; middle toe + its claw, 1.13.

2,069. Vireo olivaceus. 6.12 10.12 ♂(ad?).

2,070. Sturnus naevius. 6.10 9.65 ♂(ad?).

2,071. Sturnus naevius. 5.90 9.50 ♀(ad?).

September 10th.

Sitta canadensis. - One was skipping about after insects in the Norway Spruce tree in front of the house.

Dendroica striata. - Has probably been here before but first seen today.

Dendroica coronata. - do

September 11th.

To-day is my 24th birthday.

Miss Warner sent me a Broad-winged Hawk (Buteo pennsylvanicus), doubtless the same bird I wounded on shore the 7th. In the afternoon I shot another fine specimen north of Pell's Pond.

Totanus solitarius. - One at Pell's Pond, Vireo gilvus. - Singing about the house.

1880.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

2,072. Buteo pennsylvanicus, Wils. 16.15 26.25 ♀ juv. Wing, 11.00; tail, 7.10; culmen measured from base of cere, 1.10; measured from anterior margin of cere, .75; bill from nostril, .70; gape, 1.33; tarsus, 2.40; middle toe, 1.33; middle toe and its claw, 1.80; its claw, .57. Irides, hazel. Stomach contained mole-worms & Arvicolas.

2,073. Buteo pennsylvanicus. 16.75 37.12 ♀ juv. Wing, 11.15; tail, 7.20; bill from base of cere, 1.14; measured from anterior border of cere, .78; bill from nostril, .73; gape, 1.40; tarsus, 2.45; middle toe, 1.32; middle toe and claw, 1.85; claw alone, .62. Iris hazel.

September 10th.

Zonotrichia albicollis. First seen, one shot.

2,074. Myiodiodes mitratus. 5.70 8.10 ♂ ad. Wing, 2.55; tail, 2.20; bill from nostril, .31; gape, .58; tarsus, .77; middle toe, .43; middle toe and its claw, .60.

2,075. Myiodiodes mitratus. 5.75 8.45 ♂ ad. Wing, 2.70; tail, 2.35; bill from nostril, .30; gape, .60; tarsus, .82; middle toe, .47; middle toe and its claw, .66.

2,076. Dendroica maculosa. 5.30 7.65 ♂ juv.

2,077. Setophaga ruticilla. 5.70 8.00 ♂ ad.

September 13th.

Trochilus colubris. - One seen in H. G. this morning. This afternoon one was seen flying about the Constitution Marsh, although it was raining hard.

In the afternoon Mr. Wheatley and myself went to shoot Hons, Rails, etc. about the Lower Cove. It rained very hard before we reached West Point, and we took refuge in the tunnel, and watched the work until

1880.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

The rain ceased; and we hoped for clearing weather; but it soon began to drizzle - while we were rowing across the river - and not a Heron of any sort, nor yet an Eagle or Hawk was to be seen about the Cove. We were completely drenched, and our only consolation was in trying to shoot at the Rails (Porzana carolina), which were numerous about the mouth of Indian Falls Brook; but our guns were wet, and we were seldom able to make a successful wing shot owing to the guns hanging fire; however, we managed before dark, to bag nine, only two of which were adults. The tide was high, and cover was scarce, as only the tops of the reeds were above water, so we often had the Rails in full view, though they were very skillful in taking advantage of the slightest cover. They often came close together, being apparently very affectionate, but perhaps they were one family, led by the parents. At frequent intervals a bird would fly up in the air, and alight at a short distance. They were extremely pretty as they sat, or swam about.

2,078. Porzana carolina. 7.30 14.15 ♂ ad.
Wing, 4.22; tail, 2.30; culmen, .89; bill from nostril, .45; gape, .93; tarsus, 1.35; middle toe, ~~1.37~~ 1.37; middle toe and its claw, 1.67; claw alone, .37. Iris, reddish hazel.

2,079. Porzana carolina. 7.35 14.25 ♂ ad.
Wing, 4.35; tail, 2.30; culmen, .87; bill from nostril, .43; gape, .94; tarsus, 1.32; middle toe, 1.45; middle toe and its claw, 1.75; claw alone, .38. Iris reddish hazel. Bill, greenish-yellow. Legs & feet, yellowish & green.

1880.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

2,080. Porzana carolina. 7.85 14.00 ♀ juv.
No vestige of black on its throat.

I picked out the plumpest Rail of the nine birds shot (a young male, whose throat was slightly marked with black and its breast with drab. It weighed $3\frac{3}{4}$ oz., and presented the following dimensions: 8.75 15.30 (♂ juv.)
Wing, 4.50; tail, 2.25; culmen, .84; bill from nostril, .42; gape, .90; tarsus, 1.40; middle toe, 1.45; middle toe and claw, 1.75; claw, .32.

September 15th.

2,081. Agelaius phoeniceus. — — — ♂ juv.
2,082. — — — — — ♀ juv.
2,083. Geothlypis trichas — — — ♂ juv.

The Wood Pewee (Contopus virens) was last seen on September 11th.

September 16th.

Trochilus colubris - Still present.

Pyrranga rubra - One female seen.

2,084. Dendroica virens. — — — 5.35 8.15 ♂ ad.
2,085. Falco columbarius. 12.60 26.40 ♀ juv.

Wing, 8.25; tail, 5.75; bill measured from base of cere, .72; culmen from cere, .59; bill from nostril, .54; gape, .85; tarsus, 1.57; middle toe, 1.45; middle toe and its claw, 1.80; claw alone, .48. Irides, hazel. Lairs & toes, lemon-yellow; cere, yellow; bill bluish-black, shading into horn-color and then yellow at base.

Shot in the act of carrying off a chicken. Its sternum will be preserved.

1880.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	<u>September 17th.</u>					
2,086.	<i>Dendroica castanea.</i>	5.65	9.05	juv.		
2,087.	<i>Dendroica coerulescens.</i>	5.65	8.50	♂ ad.		
	Wing, 2.70; tail, 2.20; bill from nostril, .28; gape, .53; tarsus, .75; middle toe, .46; middle toe and claw, .58; claw, .17. <u>First seen.</u>					

September 18th.

Chaetura pelagica. - Still plentiful.
Cyanospiza cyanea. - Still present. Appears unusually shy. Inhabits weeds in gardens, and skulks shyly amongst the bulrushes of the salt marshes. No blue males are seen.

2,088.	<i>Turdus swainsoni.</i>	7.25	11.90	♀ ad.		
	Wing, 3.82; tail, 2.87; bill from nostril, .36; gape, .80; tarsus, 1.09; middle toe, .66; middle toe and its claw, .88; claw, .23.					

September 19th.

Carpodacus purpureus. - Has been numerous for a week past, and sings.
Sitta canadensis. - Comes about the house, but have not met it in the woods yet.

Chrysomitris tristis. - Assuming its winter plumage.
Dendroica pinus. - Three were sporting in the white pine-tree in front of the piazza. They were very active, and remarkably gentle. Their note was a short, metallic clicking, bearing a slight resemblance to the Brown Thrasher's. I have not seen the Pine Creeping Warbler before since the spring of 1879, and only once in autumn before, viz: on August 31st, when I shot a female in the soft autumn plumage, that was sitting quietly in the wild cherry in front of the house.

1880.

September 26th.

This afternoon rowed up to the Constitution Marsh, in company with Wheatley, to shoot Rails. Those birds were not very numerous, but I shot three, all of which were immature. I fired into the flock of Blackbirds, and brought three of them to bag; afterwards shot at two Blackbirds flying together and got 'em both. Shot a Kingfisher sitting, with No. 12 shot, and killed another at long range, flying, with the same sized shot.

Rallus. - A Rail, or some such bird, not a Carolina but of considerably larger size rose out of the grass along with a Carolina Rail, and dropped into the bulrushes across the creek. I was pushing the boat and had no gun, and my companion did not shoot it. We could not get it up out of the bulrushes again. It may have been a Virginia Rail, but I think it was a larger species; possibly even a Florida Gallinule.

Telmatochlamys palustris. - Some were seen and heard singing pleasantly in the marsh, but its numbers are greatly reduced, and it will probably be gone in a few days.
Cyanospiza cyanea. - A very few still remain in the marsh, all in brown plumage. I shot one.

Pandion haliaetus. - One alighted in the cove, but flew as we approached.
Trochilus colubris. - One seen yesterday, but not observed about the meadows to-day.

1880.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

Turdus swainsoni alviciæ. - One was brought to me that flew against a telegraph-wire, and died soon after. The only one seen this autumn.

Chaëtura pelagica. - Not seen since the 18th.

September 28th

Cyanospiza cyanea. - Several seen to-day amongst the weeds and blackberries near the gate, and in the corn-fields.

Went to Peekskill on the "River Belle", and saw a Hawk which was probably Falco columbarius, as we passed Patterlee's. Paid them a pleasant visit at the Peekskill Military Academy, and, after taking tea, returned by 7:00 p.m. train.

September 29th

Cyanospiza cyanea. - Still in the blackberries by the gate.

Spizella pusilla. - Still numerous.

Mimus carolinensis. - Several were seen, as Mr. Deuze & myself were going to the river for what may be our last swim of this season; went in yesterday.

Zonotrichia albicollis. - First seen yesterday since the one shot on September 10th.

September 30th.

Pipilo erythrophthalmus. - One seen.

Mimus carolinensis. - Still present.

Ectopistes migratorius. - One flew up from the ground into a tree. It had a good cropful of acorns.

Liriodus mustelinus. - One seen.

1880.

October 3d.

I came down on "Mary Powell" on Friday the first of October, and registered at the College and attended the opening lecture that evening.

This afternoon Morris and I walked all the way through the Central Park from one end to the other. We had a very good afternoon, and saw numerous birds. At first we found a great deal of dust and a great crowd of people. The trees were loaded with dust, and the leaves were fairly gasping for breath; and of course no birds were seen save the ubiquitous Sparrow (Passer domesticus); then came shady walks, fresher air and less people. The first wood bird was a handsome Red-bellied Nuthatch (Sitta canadensis) up on top of a big elanthus limb, tugging and digging into a crack where its prey laid concealed; at last we saw him seize the larva & swallow it, then flit to a new branch, with its highly pitched houk, houk, houk. We saw this Nuthatch in several other places. One was hammering at a dead tree-top; another was catching insects in some bushes bordering a pond; and we saw two prying into the cones of the yellow pine-trees in search of seeds or insects. Then we came upon a pretty flock of Cedar Birds (Ampelis cedrorum) that were feeding upon ~~some~~ ^{the} berries of some black thorn bushes, which occupation they varied by darting in curves after

1880.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	insects; some of them still present the striate breast of the young birds.					
	A Red-shouldered Hawk (<u>Buteo lineatus</u>) circled overhead, and, a short time after we saw a Bittern (<u>Botaurus minor</u>), which must have been disturbed by some rumbler beside some pond. Robins (<u>Turdus migratorius</u>) were numerous in one or two places, and its familiar notes quite frequently heard. We saw a few Song Sparrows (<u>Melospiza fasciata</u>) and one Swamp Sparrow (<u>M. palustris</u>) beside the ponds. A Brown Thrasher's ^(<u>Harporhynchus rufus</u>) note was heard, and the White-throated Sparrow (<u>Zonotrichia albicollis</u>) was quite plentiful in all the bushy places. I watched for Catbird & Towhee Bunting but we saw neither. We heard the notes of the Golden-crested Kinglet ^(<u>Troglodytes aedon</u>) amongst a clump of Norway spruces; and we came upon a beautiful Ruby-crown (<u>Regulus calendula</u>) was in some bushes beside a pond: the first I have seen since last spring. Then we came upon a a wild little ravine covered with underwood and pines. I heard the note of a <u>Suirus</u> , and succeeded in calling it up the ravine from the pond, to where we were standing. Then we identified it as the Small-billed Water Thrush (<u>Suirus naevius</u>). A note was just then heard which was almost new to me. It was that of the Cardinal Redbird (<u>Cardinalis</u>					

1880.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	virginianus). I chirruped until I brought it close up to us. It kept amongst the bushes, and appeared to be very shy. It was unusually dark red, and sat with erect crest. Its note was sharp and metallic like the Water Thrush's. We heard a fine song down beside the pond, which we attributed to the Small-billed Water Thrush, as it did not seem to be like the Cardinal's song. The Yellow-rumped warblers were seen and heard everywhere. A feather was found which came from the left wing of the Great Blue Heron (<u>Ardea herodias</u>), and a Spotted Sandpiper was feeding on the margin of the reservoir. We saw two birds - Warblers - which puzzled us very much for a while; but when I got a near view I saw that they were Black-poll Warblers (<u>Dendroica striata</u>). There was a Night Heron sitting in a tree over my head. When it flew, I expected to see some large Hawk, but it proved to be <u>Nycticorax</u> , and only flapped across the narrow pond and alighted upon a slender bough of an oak tree, flapped a moment, while perfecting a balance, sat with neck extended for a moment, then drew its head close upon its breast, and seemed to repose. White-bellied Swallows were quite numerous; and the Purple Martins (<u>Progne subis</u>) were flocking about some Shanties & empty lots outside the Park. We saw two Gray Rabbits, and some Gray Squirrels. Muskrat signs were about the ponds; and one House Rat was seen beside a pond; it jumped					

1880.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

in a circle around one end of a brick arch. To sum up the birds seen were as follows:—

- Larus immigratorius*. — Quite numerous.
Harporhynchus rufus. — Thought we heard its note.
Regulus satrapa. — Heard in the evergreens.
Regulus calendula. — One beside our pond.
Sitta canadensis. — Several seen.
Dendroica coronata. — Numerous.
Dendroica striata. — One pair.
Siurus naevius. — Two were seen.
Tachycineta bicolor. — Numerous.
Progne purpureus. — One flock.
Parus atricapillus. — Numerous.
Ampelis cedrorum. — Several flocks.
Melospiza fasciata.
Melospiza palustris.
Zonotrichia albicollis. — Abundant.
Junco hyemalis. — One beside reservoir.
Passer domesticus.
Cardinalis virginianus. — One seen—others heard.
Sayornis fuscus. — Beside the pond.
Centurus vociferus. — One sitting on low branch.
Buteo lineatus. — One flying.
Tringoides macularius. — Beside reservoir.
Nyctanassa grisea naevia. — One at a pond.
Botaurus minor. — One flying south.
Ardea herodias. — Found a feather.

October 8th.

On Tuesday Oct. 5th, I took 4:20 P.M. train from Hunter's Point for Sayville, which place I reached after dark, and found some trouble in getting settled for the night; but, finally I was directed to one Captain E. V. Dames, at whose pleasant house I was very comfortably entertained. That night I

1880.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

retired early, after a talk about birds and shooting with the Knid Captain. Early next morning we embarked on board the Captain's little sail boat bound for across the bay to the beach which separates the Atlantic Ocean from Great South Bay. The bay shore is distant from Capt. Dames's house about $\frac{3}{4}$ of a mile, and all along the way we saw the meadow larks and many other birds; many of them were singing sweetly. The Capt. wheeled our table on a barrow, and our traps and ourselves were soon safe in the little skiff, and we pushed out. The little sloop & soon were ready to set sail. A flock of "Sand Sripes" passed by before we could get hold of our guns, but when four more came along, I brought down two of them, one of which was winged and afterwards ran about the skiff, and squealed wherever we tried to catch it. They were Semipalmated Sandpipers (*Ereunetes pusillus*). Crossing the bay we saw only a few "Broadbills" and one loon. The waves were high, & I considered myself fortunate not to be seasick. The loon flew before we could procure a shot. We landed at the narrowest part of the beach—the "Cherry Grove House". We parted at the beach; the Captain went west, and I proceeded eastward. The first bird I put up was a Bittern which I summarily missed with both barrels; another one was seen a little farther along, and it was also missed. A Savannah Sparrow and Ruby-crowned Kinglet were shot, and also a Rail (*Porzana carolina*) which I could not recover. Then I visited the

1880

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	ocean side of the beach, examined the low forests about the higher part of the beach, and then started towards our boats. Pigeon Hawks were numerous, and I missed one that passed overhead, as I likewise did a Short-tailed Tern, and a small Gull, which I thought, perhaps erroneously, were within range. A huge Cormorant flew past close to the water - the craziest looking bird I ever saw. Then I came across some Snipes, and shot about thirty specimens of about five different species, four of which I had never shot before, and three were seen alive for the first time. I was then joined by Capt. Dams, and we procured ten more Snipes, making a total of forty for our first day's bag. We then cooked dinner and rested a while. Then walked up along the bay, but, finding no Snipes went to the ocean side of the beach, and gathered some mollusks, and gradually meandered up along the shore to opposite the Government Life-Saving Station, which we visited, and where we were most hospitably entertained by the Captain of the crew, who showed the whole establishment and all the apparatuses pertaining thereto. I shot a Pigeon Hawk (<i>Falco columbarius</i>), but it was flying & fell amongst a tangled growth of bushes and cat-briars which was impossible to explore. So I lost a good bird. There is a path across the wooded portion of the beach from the Govt. House to the bay side					

1880.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	; crossing it, we saw a Pigeon Hawk sitting upon a dry pine. Capt. Dams tried to procure a shot at it, but failed; I followed it, and was rewarded by getting a shot at a passing Night Heron, which I brought down at long range in fine style with a charge of No. 4 shot. I happened to strike upon a small patch of sand bare of bushes, else I should probably never have recovered it. Walking down the bay shore of the beach, I spied a Hawk sitting upon a dry pine-tree in the middle of the woods. I succeeded in shooting it, but was for a long time unsuccessful in finding it although we knew just where it sat when I fired at; at length I saw a dark object in ^{the} tangle which I thought might be a Hawk's wing, and so it proved, for I at length succeeded in reaching it, and, to my boundless delight found a beautiful Peregrine Falcon or Duck Hawk (<i>Falco communis</i>). This was the crowning success of the day and we returned to our boats with the following birds, all of which were carefully plucked with cotton & wrapped in papers to prevent soiling.					

- 1 Savannah Sparrow
- 1 Ruby-crowned Kinglet
- 1 Bluebird (shot by mistake)
- 36 Semipalmated Sandpipers
- 1 Least Sandpiper.
- 1 Greater Yellow-legs.
- 1 ^{Sonopartia} Red-backed Sandpiper
- 2
- 1 Night Heron
- 1 Peregrine Falcon
- 1 White-bellied Nuthatch
- 1 Phoebe
- 1 Pewee

Total 47 birds

1880.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

We set sail for home and had a good wing, which brought us in just at dark. There was a new moon and a glorious sunset. So ended Wednesday (Oct. 6th).

On Thursday we took an early start, and shot a flying Meadow Lark at long range on our way to the shore. We crossed to Cherry Grove House as before, and on the day saw a few Ducks, Gulls, and one Loon. Only a few flocks and scattered individuals of the Semipalmated Sandpiper were seen, and no other species was observed. On Wednesday we saw a number of Snipes which I did not recognize and could not shoot. The Captain also missed a good shot at two Black-bellied Plovers. We found the smaller sand bird even more numerous than on Wednesday and following & preying upon them were numbers of Pigeon Hawks, and a few Marsh Hawk. Capt. Daves missed a good shot at an adult Duck Hawk that flew overhead. The Pigeon Hawk were constantly dashing past us, and I fired a number of times at them but they are so extremely swift that I did not hit one of them. We saw them constantly striking down small birds, and I saw one miss a flicker several times, while another one struck down one at the first attempt; the poor Woodpecker made a great noise when first caught, but its cries were soon hushed in death. Catbirds, Nuthatches, Alice's Thrushes, Brown Thrashers, Robins, Yellowbirds, Flickers and Yellow-bellied Woodpeckers and especially

1880.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

The Yellow-rumped Warblers were exceedingly abundant upon the beach. We left the beach at noon and sailed for Ducks. We shot two beautiful Horned Grebes. On shore we hunted the meadows for English Snipe, but only shot a Meadow Lark in beautiful autumnal plumage. Just caught the 4:11 P.M. train where I said good-bye to Captain Daves and reached New York in time for dinner, and Dr. A. K. Fisher awaiting me. We unpacked the birds together.

List of Birds Seen:

Turdus migratorius. - Rather abundant on Island & beach.

Turdus swainsoni aliciae. - Very numerous on beach.

Harporhynchus rufus. - Numerous on the beach.

Mimus carolinensis. - On the beach; more abundant than I ever knew it to be anywhere ^{else}.

Sialia sialis. - Only found on the Island (Long?)

Regulus calendula. - Rather abundant on beach.

I shot one thinking it was the Short-billed Marsh Wren, judging from its Wren-like note and from seeing it in the grass.

Regulus satrapa. - In the cedars on shore of?

Parus atricapillus. - Very numerous everywhere on the Island shore and also on the beach.

Sitta carolinensis. - Abundant on the Long Island shore and on the beach, where one was shot.

Sitta canadensis. - Exceedingly abundant upon the beach; also observed on shore in the cedars. Numbers were seen on the beach on Wednesday, but on Thursday there was one constant succession of flights of Red-bellies, which were most abundant in the cedars, but alighted in a abundance upon the ground and in the grass and low bushes.

1880.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

Certhia familiaris. - Numerous on the beach.
Dendroica coronata. - Excessively abundant upon the seaward, sand beaches, grass, bushes, trees, in the air, and in short everywhere both on the beach and on shore.

Dendroica palmarum. - Several seen upon the beach on Thursday

Amphispiza cedrorum. - Three birds seen together on the beach on Thursday

Carpodacus purpureus. - Everywhere.

Chrysomitris pinis. - Its notes were heard all day Thursday upon the beach, and on the following morning a number of scattering ones were seen. I shot one flying singly overhead & killed it with a charge of No. 4 shot. Strange to say, it was ^{but} slightly mutilated and made a good specimen.

Chrysomitris tristis. - Very numerous on shore and especially so on the beach.

Paperculus savanna. - On shore and beach.

Poocetes gramineus. - Abundant on shore.

Melospiza palustris. - Seen on shore.

Melospiza melodia. - Everywhere numerous.

Junco hyemalis. - A few on shore.

Spizella monticola. - Seen on Thursday

October 7th for the first, on shore

Spizella socialis. - Very abundant on shore.

Spizella pusilla. - do do do.

Zonotrichia albicollis. - Abundant.

Paper domesticus. - Abundant in Sayville

Sturnella magna. - Very numerous; two shot on shore, none on the beach.

Corvus americanus. - Everywhere numerous.

Cyanurus cristatus. - Very common on shore.

Sayornis fuscus. - Very common on the beach, and also on shore.

1880.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

Sphyrapicus varius. - Migrating in large numbers along the beach, flying low and frequently alighting.

Colaptes auratus. - Excessively numerous upon the beach; perched upon the dead pines, and ate the berries on bushes.

Constantly moving westward. I saw the Pigeon Hawks strike at them, and one that was caught squealed loudly.

Falco communis. - Two Duck Hawks were seen; one was shot upon a dry pine from the bay side of the beach, and though killed dead, was recovered with great difficulty owing to the dense, matted tangle of cat-briers & bushes surrounding the tree; another flew overhead on the following day (Oct. 7) and Capt. Danes missed a good shot at it. None others were seen.

Falco columbarius. - Very large numbers were seen flying upon the beach. One was shot on wing on the 6th. It came straight down as I fired my second barrel, but could not be recovered in the thick bushes. I shot a several that were flying past like lightning on the following day, but got none. They darted into every thicket after little birds, and were following the immense numbers of small birds that were moving westward over the beach. The rapidity of their flight is marvelous. One captured a Flicker (Colaptes auratus) in my presence. They were so numerous that they were scarcely ever out of sight, though they constantly pass out of view to

1880.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

the westward, not to return.

Circus cyaneus hudsonius - A few seen on the beach trying to capture small birds; one seen on shore.

Squatarola helvetica. - Captain Daves missed two "Black-bellied Plover" that were sitting upon the beach.

Ereunetes pusillus. - One flock flew past as we were embarking for the beach early on the morning of the 6th. Another band of four came along a few moments later, and I brought down two at one shot. On the beach they were numerous in flocks of ^{from} about thirty down to single birds. They were very gentle and always returned to their dead companions after a shot was fired. Their note when flying was agreeable & wild. When wounded ones were caught they squealed loudly. Single individuals and large flocks were seen on shore and on the beach on the 7th also.

Tringa minutilla. - One was found in the lot after I unpacked my birds here in New York.

. - One was shot amongst a flock of Semipalmated Sandpiper feeding upon ^{the} seaweeds on the bay side of the beach. Several sandpipers killed same shot.

. - One shot as it flew along shore of beach uttering a loud note; another was shot at as it stood upon the beach; it never moved, and I fired another shot; then it flew out upon some seaweeds, and I shot it. These two were all that were seen.

1880.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

. - One shot at long range from a passing flock of 4. Another seen on beach also Nyctiardea grisea naevia. - Several young were seen flying & feeding on the beach on the 6th. On that afternoon I shot one of a pair of adult birds that were flying over the beach. None were seen on the 7th.

Botaurus minor. - Two were shot at on the morning of the 6th. None others seen.

Porzana carolina. - One solitary Sora was shot as it flew over the marshy part of the beach. The only one seen.

Anas obscura. - Two lit in a small pond on the beach.

Fuligula marila. - Several flocks seen.

Oedemia fusca. - One or two flocks seen.

Eraulus — ? - One was shot at as it flew low above the water to the westward along the bay shore of the beach.

Larus argentatus. - One or two young and a number of old birds seen.

One or two small Gulls were seen, but not identified.

Hydrochelidon fissipes. - One ^{bird} that I supposed to be of this species was seen on the beach.

Podiceps cornutus. - Two were shot together in the middle of the bay - both in young plumage; they dived several times before killed, and about 7 shots were fired.

Colymbus torquatus. - Several loons seen in the bay; shy.

Note. - A number of land birds doubtless escaped observation, and probably several additional species of Snipes were seen flying, but were not identified. No Owls were seen, save a Screech Owl (Scops asio) as I walked up from the station on the 5th.

1880.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
2,089.	<i>Sturnus melanoleucus</i> .	Oct. 6.		14.15	25.00	♀ ad.
2,090.	<i>Fringa minutilla</i> .	"		6.05	11.25	♀ ad.
2,091.	<i>Sturnella magna</i> .	Oct. 7.		10.15	16.85	♂ ad.
2,092.	<i>Polidna alpina americana</i>	Oct. 6.		8.75	15.10	♂
2,093.	<i>Fringa maculata</i> .	"		9.50	18.00	♂
2,094.	"	"		9.30	17.90	♂
2,095.	<i>Ereunetes pusillus</i> .	"		6.25	12.00	♂
2,096.	"	"		6.40	12.10	♂
2,097.	"	"		6.30	11.90	♂
2,098.	"	"		6.20	11.90	♂
2,099.	"	"		6.30	12.10	—
2,100.	<i>Falco communis</i> .	"		12.40	45.50	♀ juv.

Wing, 14.00; tail, 8.25; culmen from frontal feathers, 1.09; from cere, .85; cere alone, .30; bill from nostril, .81; gape, 1.85; tarsus, 2.23; middle toe and claw, 2.78; middle toe, 2.15; claw, .75.

Irides very dark. Crop & gizz. were packed with fragments of numerous birds — all apparently passerine —, which had been roughly picked before swallowing.

Legs & wings were swallowed entire, the latter with feathers on. I preserved a number of entire legs which apparently belonged to the following species:—

Turdus migratorius.

Turdus swainsoni aliciae.

Minus carolinensis.

Dendroica coronata.

1880.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
2,101.	<i>Nyctiardea grisea naevia</i> .	Oct. 6.		25.50	44.00	♂ ad.
	Iris red.					
2,102.	<i>Passerculus savanna</i> .	"		5.88	9.60	♂ ad.
2,103.	<i>Chrysomitris pinus</i> .	Oct. 7.		5.25	9.20	♂ ad.
2,104.	<i>Podiceps cornutus</i> .	"		—	—	♂ juv.

Sunday, October 10th. Highland Falls, N.Y.
Junco hyemalis. — Now common here.
Spizella monticola. — First seen here.
Spizella socialis. — Very numerous.
Spizella pusilla. — do.
Turdus aliciae. — Very abundant here.

Sunday, October 24th. Highland Falls, N.Y.
Turdus migratorius.
Turdus swainsoni aliciae. — Very typical *Aliciae*.
 Thrushes were quite numerous.
Turdus pallasi.
Sialia sialis. — Has a loud rattling note.
Regulus calendula. — Still abundant, and generally distributed.
Sitta carolinensis.
Parus atricapillus.
Dendroica coronata. — Very abundant.
Chrysomitris tristis.
Spizella socialis. — Several seen.
Junco hyemalis. — Very plentiful.
Melospiza fasciata.
Zonotrichia albicollis.
Passer domesticus. — A cold wind prevailed, and the sparrows as usual at such times frequented the evergreens, in large, noisy flocks.

Corvus americanus. — Mother says the Crows have been migrating in large flocks for a week past. I saw a flock of several hundred flying westward this morning.

1880.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.Lagomys cristatus.Falco columbarius. - One down by the riverAnas obscura. - One in the river.Larus argentatus. - A few on the river.

The above were all the birds noticed.

Note.

A Swan was shot on October 21st (1880) on one of the lakes near Highland Falls. It was purchased by Mr. Charles Butler of West Point, and sent to Newburgh to be mounted. When it returns, I shall be able to see it and determine its species, though there is little doubt of its being C. americanus.

October 28th.

Mr. Morris & myself arrived at Sayville, Long Island, N. Y. by the evening train (October 27th). We took our supper at Captain Dances, after which we went down to the beach and got out our decoys and put them all in the boat. We had about 75 decoys in all, both ducks & brant. Then we dumped the water out of the battery, and took it out to the sailboat, and got everything ready for an early start. We were off by five o'clock on the morning of the 28th. It was very cold and windy. We crawled out of the cabin when we got to the ground, but found the bay at that point too rough to put out the battery; we were obliged to go down to the leeward of Five Island. On the way we had several shots at Loons that were flying over. I wound one, but he did not come down; another one Morris

1880.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

and myself fired at together, one of us broke his leg, but he kept on flying. A Velvet Duck was swimming in front of the boat; I got up by the mast, and fired my gun into the air by accidentally pressing the trigger before I got the gun up, and then missed a wing shot with the other barrel. We did not get the battery out till nine o'clock. Just before I got into the battery, I brought down a Gull (Larus) at long range with "B" shot, at the second shot. I fired 15 shots in the battery, and only got two Gulls (L. L. argentatus and L. L. affinis) and one Duck (Fuligula). Two good flocks of Broadbills came to the stools, and several bands of two or three, and some single birds, but most of them kept pretty well off, & only afforded snap shots, which I make a practice of missing. In the afternoon, Morris shot seven Ducks (five Red-breasted Mergansers and two Big Blackheads. He fired eighteen shots. I chased one of his wounded birds, a Red-breasted Merganser, and shot him over again on the wing. Loons and Red-throated Divers (Colymbus torquatus and C. septentrionalis) were seen; plenty of the former, and one of the latter. Horned Grebes (Podiceps cornutus) were seen a few times. One came within five feet of the battery while Morris was shooting.

The following Sea Birds were noticed:
Branta canadensis. - One bird seen
Anas obscura.

Fuligula macrila. - A bandant. Morris shot two
Fuligula affinis - I shot one of a flock of fifty.

1880.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	<i>Harelda glacialis</i> .	October 28th	continued.			
	<i>Oedenia fusca</i> .		Batter plentiful			
	<i>Oedenia americana</i> .		Abundant			
	<i>Mergus serrator</i> .		Immense numbers.			
	<i>Larus argentatus</i> .		Quite a number of Herring Gulls seen.			
	<i>Larus</i>		Two were shot, and several others seen.			

A White-headed Eagle flew over the battery; it was an adult bird.

105. *Larus delawarensis*. 17.50 42.50 7 juv.
Wing, 12.40; tail, 4.80; culmen, 1.73; bill from nostril, .77; gape, 2.32; tarsus, 2.15; middle toe and claw, 1.73; middle toe, 1.40; the claw, .35. Iris, hazel. Bill black, with a reddish tinge at base of inferior mandible.

106. *Larus delawarensis*. 17.60 43.25 7 juv.
Wing, 12.50; tail, 5.00; culmen, 1.55; bill from nostril, .72; gape, 2.17; tarsus, 1.95; middle toe and its claw, 1.71; toe alone, 1.42; claw, .22. Iris, hazel. Terminal one-half of bill (from tip to nostril), black; residual, flesh-color. Legs + feet, white, slightly brownish in front of tarsi + toes.

Measurements.

2106a *Fuligula marila*. 19.45 32.25 6 22 years.
Wing, 8.65; tail, 2.65; culmen, 1.80; bill from nostril, 1.25; gape, 2.33; tarsus, 1.66; middle toe and its claw, 2.90.

2106b *Fuligula affinis*. 17.25 29.75 3 22 years.
Wing, 8.00; tail, 2.60; culmen, 1.61; bill from nostril, 1.12; gape, 2.03; tarsus, 1.45; middle toe and its claw, 2.50.

The wind kept to northward all day, and consequently the day was bad, and ducks scarce. No Brant were seen.

1880.

(On the evening of the 27th, when we were out on the bay, about ten o'clock, the myriad voices of little migrating birds, and, again before daylight on the 28th we heard them in abundance far out on the middle of the bay.)

October 29th.

To-day left Sayville by 7.31 A.M. train. When we got within two miles of the Garden City depot, Morris called my attention to a Snowy Owl (*Nyctea nyctea*) that the train had started up from near the track. It flew slowly, close to the ground, and finally alighted on a sandy roll amongst some hummocks in a slight rise of ground. I intended to jump off the train which was moving very slowly, but Morris persuaded me to wait and get off at Garden City, which I did, and where I left my valise, first taking out all the carriages. I walked rapidly back two miles to where the bird was seen, but it was not there. The country on both sides of the track was a desolate tract of red bunch-grass, nearly level, and sandy in spots. Here and there a farmhouse, or, rather, a homestead, for the ground is very little productive, except of a stunted growth of oak and cedar-trees and red grass. I walked on another mile, and was repaid by seeing my Snowy Owl sitting in the midst of a large flock of Crows (*Corvus americanus*) upon a sandy spot near the railroad, where I walked behind the track railroad, where

1880.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.

was elevated above the surrounding country, until I came to a culvert underneath it, through which I passed; as I emerged, a large number of the Crows flew off, but many remained, and continued teasing the Owl. I made a hasty calculation as to how many Crows I could kill at one shot, and what were the probabilities of killing the Owl at the same shot, but decided to make a sure thing of it. I approached slowly when the Crows all flew scaring the Owl at the same time. I aimed at it, with my finger on the left trigger, but dared not pull lest I should lose my last chance of killing a Snowy Owl; so I let him fly slowly away close to the ground, with the balance of the flock of Crows chasing him & diving at him. He disappeared behind a neighboring rise of ground; but I knew he had settled into the grass when I saw the Crows turn and alight. I ran as far as the hill, then laid down and crawled towards the Owl, whose head and neck were visible above the grass. The Crows flew off, and then I raised and fired at the Owl a charge of No. 4 shot. He spread his wings with a quiver; and I turned and fired my left barrel's "B's" at the hindmost Crow, which I hit splendidly at long range, and the feathers float away back to where I was catching my Owl. The latter was not killed, but attempted to fly, but I struck him down with my gun, and

1880.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.

then seized both humeri close over the bird's back. It struggled and attempted to bite, arching his neck & snapping his bill and trying to grasp my clothing with his claws. I handed it an empty paper shell which it seized with its bill, and held until the small taste of burnt powder caused it to drop it. Then I poked another at it which it bit fiercely. Its bright yellow eyes were wonderfully fierce and handsome. I finally killed it, after carrying it a good way, and tied it up in my silk muffler. It was the first Snow Owl I ever saw in a wild state. It uttered a curious ^{grating} sound, and ^{squealed} loudly. Passerculus savanna. - This was the only sparrow seen about those desolate meadows, and, though quite evenly distributed. Every four or five rods, one or two would fly up with a whirr and a curious little note. Dendroica coronata. - A few were seen by the railroad in one place. Corvus ossifragus. - A large flock of this crow was seen near Garden City, and I started after my Owl, but I missed the only two that came near enough for a shot. I saw several others straggling about, but could not obtain a shot. The flock first seen contained a hundred or more birds. Sturnella magna. - In the fields of bunch grass, I saw very few, and got one last; but in a large field of stubble & weeds near Garden City, I saw at least fifty meadow larks at one moment, and the adjacent fields abounded with them. I fired at some flying birds with No. 13 but with no result.

1880.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

- 2,107. *Sturnella magna*. 10.95 16.80 ♂ juv.
 2,108. *Nyctea nivra*. Snowy Owl. 22.85 59.00 ♂ ad.
 Wing, 15.75; tail, 9.50; culmen measured from frontal feathers, 1.53; from cere, 1.00; gape, 1.98; tarsus (about), 1.90; claw of the middle toe, 1.18. Irides, yellow.
 Uttered a squealing and also a grunting noise while alive, after being caught.
 Its stomach was absolutely empty.

Mr. Morris saw several Greater Yellow-legged Snipe, near the Hunters Point depot, in some marshes at the oil works.

October 31st.

This afternoon we walked through the Central Park to 100th St. At the little pond opposite the 100th St entrance, I ran across the Cardinal Birds again, perhaps the same we saw there on the 3d of October. There were two of them, male & female, in the thick bushes bordering on the pond. They uttered a rather faint, metallic chirp. When I got near, they both flew across the pond. I found them again on the other side amongst the bushes. I hid amongst the bushes, and watched them. The male flew up on a low tree over the edge of the water and began chirping louder than before, the female always answering him. Then they flew across to the old place where the brook runs in. They were extremely shy, and kept concealed in the bushes, and sometimes descended to the water.

1880.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

- The following birds were seen in the park:
Turdus migratorius. - Everywhere numerous.
Turdus pallasi. - Saw them going into the bushes beside a pond to sleep.
Regulus satrapa. - Quite numerous.
Parus atricapillus. - do
Certhia familiaris. - One was seen creeping on a small tree, but shyly, kept on the other side of the tree.
Ampelis cedrorum. - Several small flocks.
Melospiza fasciata. - Quite numerous.
Junco hyemalis. - Roosted on the ground, amongst the bushes.
Zonotrichia albicollis. - Abundant.
Passer domesticus. -
Passerella iliaca. - One seen.
Cardinalis virginianus. - Male & female seen.
Accipiter cooperi. - One over the lake.
Falco columbarius. - One seen.
 No Gulls have been seen in the park this autumn, yet.

November 2d - Election day.

- Spent the morning in the Central Park, looking at the birds. The following species were noticed:
Turdus migratorius. - Very plentiful.
Turdus pallasi. - do
Sialia sialis
Regulus calendula. - In considerable numbers.
Parus atricapillus. -
Sitta carolinensis. - One seen.
Dendroica coronata. - A few still present.
Ampelis cedrorum. - Some small flocks.
Chrysomitris tristis. - Scattered flocks.
Melospiza palustris. - One beside a pond.
Melospiza fasciata. - Not very numerous.
Junco hyemalis. - Common.

1880.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.Spizella pusilla. - One Field Sparrow seen.Zonotrichia albicollis. - Everywhere abundant, and singing its peculiar ditty; besides it has a metallic chirp.Passer domesticus.Cardinalis virginianus. - I found some Cardinals as usual, both males and female, but not in the old place.

My attention was attracted by a peculiar squealing noise upon the hillside near the path beside the upper, middle lake. There were two male Cardinals upon the ground close together, shaking their feathers and acting altogether very like an amorous pair of English Sparrows. I had them in view for an hour or more during which time they kept constantly side by side, feeding each other, and exhibiting the greatest affection for each other. The female bird that accompanied them flew down in the swamp beside the brook and pond. They kept up a shivering, squealing note everywhere.

Picus pubescens. - One seen.

No Water-fowl or birds of prey seen with the exception of a Night Heron (Nycticorax nycticorax) in immature plumage, which flew from a tree directly over my head, on the Harlem lake.

November 7th Highland Falls, N.Y.
Walked through the woods all the morning, and observed the following birds:

Larus migratorius. - Not abundant.Larus pallasii. - Only one seen.Sialia sialis.Regulus satrapa.Regulus calendula. - Only a few were seen, in com-

1880.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.- Many with R. satrapa. Nearly all seen on theParus atricapillus.Certhia familiaris.Sitta carolinensis. - One was feeding in the nestof the road, in company with Snowbirds (J. hyemalis).Ampelis cedrorum. One large flock.Spizella monticola. Abundant. The only Spizella seen.Junco hyemalis. Abundant.Act. cinis tristis. - Abundant. Full winter plumage.Passerella iliaca. - Abundant. Singing everywhere

in the woods and about the house.

Passer domesticus.Corvus americanus.Cyanurus cristatus. - Abundant.Picus pubescens.Colaptes auratus.November 8th.

Came to New York by first train. Saw several American New Gulls (Larus calurus brachyphyllos?) while crossing on the ferry to Harrisburg, and one Herring Gull (Larus argentatus). The New Gull was recognized on the river for the first time.

November 12th + 13th. Branford, Conn.

Thursday morning it rained hard, but cleared about eleven o'clock. At Robt. Morris's suggestion, I took the one o'clock, P.M. train (on the 11th) to New-Haven; went directly to Morris's house, No. 86 Prospect Street, and got plenty of cartridges and a box of primers. Spent a very pleasant afternoon looking at game-pieces, specimens, drawings etc., and, after tea, took train to Branford. Amongst the birds at Morris's, was an albino mounted specimen of the Bluebird (Sialia sialis), which showed the trace, only, of the blue color.

1880

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

At Linsley's Hotel, at Branford, I spent a comfortable night. Retired at eight and slept till daybreak. Awoke to find a heavy wind blowing from southwest, but, as I was extremely anxious to shoot something, Mr. Linsley went out with me but we found it impossible to get out of the river (Branford River), and were obliged to tie our boats off shore, and walk down to the beach. We saw a good-sized flock of Black Ducks (nas. obscura). Oldsquaws (Harelda glacialis) and Coots (Oedemia - Melanetta velutina) were abundant; and a few Hulls (L. L. argentatus smithsonianus, and brachyrhynchus). One small flock of Snow Buntings (Plectrophanes nivalis) flew off shore to the islands. It was too rough for any boat to lie off shore, and the only bird I shot was a female White-winged Coot. I tried some long range shots at Oldsquaws off Lemphearts Hotel with No. 3 shot. Slipped on a high rock covered with fucus whose slimy pods were exceedingly slippery, and missed and cut my elbow severely, smashed lock stock and barrel of my gun. In front of the hotel the beach and boats were strewn with dead Duck of the three species of Coots, etc., which were thrown away to rot, by the swell gunners who come to shoot for sport. I saw the following species most of which were shot on the 10th.

<u>Melanetta velutina</u> -	Probably 80 dead ones on the shore.
<u>Pelionetta perspicillata</u> -	" 20 on beach (only 2 ♂ ad).
<u>Oedemia americana</u> -	" 30; both sexes all stages.
<u>Harelda glacialis</u> -	" 40. All plumages.
<u>Colymbus torquatus</u> -	1 immature
<u>Podiceps cornutus</u>	1 immature
<u>Corvus lugivorus</u>	5
Total (about)	177 birds

1880.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

One man living near the beach said he thought it a shame to slaughter Ducks and leave them there to rot, and remarked that he was in the habit of supplying his pigs with plenty of fresh Ducks.

Note. - Mr. Sil. Linsley said that the Duck Hawk (Falco peregrinus naevius) usually came around in cold weather. He has had them kill and eat wounded Ducks, and those which he has shot and allowed to drift off a little way.

Towards night I took train to New Haven and was fortunate in getting my gun nicely repaired, so that I could shoot on Saturday. Strolled about New Haven, and met the gunners at Fulsum's gun-store. Got back to Linsley Hotel at 9.30 P. M., and went to bed tired. Next morning at four o'clock A. M. Linsley was one out, and we sat down together to a hearty breakfast, and then set out in our little duck-boats, decked over fore and aft. The wind blew hard, but I was determined to shoot; so we rowed down the river. The sea rolled very strong. At the mouth of the river the waves were so high that the Captain's head often disappeared from view, and often only his black wide-awake hat was visible. The Coots and Skunk-nads (Pelionetta perspicillata) were flying in vast flocks, and floating in large flocks, now rising on the waves and then disappearing from view. Passing over the shoal between two small islands off a point, the waves ran very high and our boats dashed through at a fearful rate. Linsley's boat would entirely disappear from view and then rise up perched on top of a huge wave with the grizzled Captain sitting erect blandly surveying the scene.

1880.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

and taking cognizance of the various flocks of Sea Hawks moving in different directions. He said it was too rough to anchor and we were the only boats afloat. We ran our boats ashore to the leeward of an island and tried one or two long shots at passing Coots and Oldsquaws at long range. The wind blew fiercely. At length we reembarked, and shot a few Coots between the two islands. After while the wind abated somewhat and we were able to cast anchor between the two islands. The Ducks were flying fairly well. Three other boats then came out, and we formed a line of five boats, but the other boats could not stand it, and soon went ashore. We remained till about one o'clock and then went in, and warmed ourselves in a house on the point. I spent the rest of the afternoon shooting Oldsquaws, and hunting the islands for Purple Sandpipers. We were two hours getting back up the river: the toughest row I have had in a long time. Both hands have a row of immense blisters, and they were not tender either. We got in after dark, and had just time to eat a gorgeous meal and pick out a few fine Ducks, eight in all, for Louis Zerega and myself to skin, and then take the train to New York.

While shooting at anchor two flocks of White Snowbirds (*Plectrophenax nivalis*) visited the island near by boat. In the crevices of the bare rock a few streaks of grass were growing; the Snowbirds flew scattering about eating the seeds and alighting beside the seaweed near the breakers. They

1880.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

left the island, visiting each of the shooter's boats on their way to the next island. They flew stragglingly, leaving the rock one after another. A flock of Purple Sandpipers (*Arguata marina*) flew from out at sea and lit on a ridge of rock covered with seaweed (kelp), which was being uncovered gradually by the receding tide, and over which the gulls were occasionally dashing. It was delightful to see the Sandpipers rise in the air to let the former pass over from a giant comber, and then settle down upon the rock and go on feeding. By dint of hard pulling, I managed to get to windward of them, and shot two with my first barrel; they flew scattering, and I missed the single bird I shot; but they soon wheeled and came in again. A third shot brought down three of the survivors, leaving only two of the original flock of seven. I was glad to get my birds and get off the rock without any mishap. I had them all lying on the deck of my little boat along with the Ducks.

Captain Sinsley made some fine shots. Four Oldwives flew past; he killed two with each barrel; two old drakes. Another time he killed two birds, ^(Coots) alone that came over, taking one with each barrel. Another time he killed an Oldsquaw at an immense height, making what the gunners call "one of Sil. Sinsley's tall shots." Often when a flock flew over he would kill one with each barrel.

The Sea Scoter (*Pelionetta perspicillata*) is known only by the name of Skunk-head by the gunners at Brauford and about there. There was one Chipping Sparrow (*Spizella socialis*) and a few Yellow-shafted

1880.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	Clicker (Colaptes auratus) on the beach on the 12th. No other birds of note were seen.					
2109.	<i>Pelionetta perspicillata</i> .	Nov. 10.	20.10	34.00	♂ ad.	
	Wing, 9.15; tail, 3.10; culmen, 1.53; gape, 2.58; tarsus, 1.82; middle toe and claw, 3.00; middle toe alone, 2.67; claw, .40. Iris, white. Stomach filled with clams.					
2110.	<i>Melanetta velutina</i> .	Nov. 13th.	23.00	39.20	♂ ad.	
	Wing, 10.75; tail, 4.00; gape, 2.77; culmen, 1.50; tarsus, 2.03; middle toe and claw, 3.22; middle toe alone, 2.85; claw, .40. Iris, white. Stomach filled with clams.					
2111.	<i>Harelda glacialis</i> .	Nov. 13th.	23.00	31.00	♂ ad.	
	Wing, 8.75; tail, 8.75 (same as wing!); culmen, 1.12; gape, 1.77; tarsus, 1.40; middle toe and claw, 2.20; middle toe, 1.95; its claw, .34. Iris, light hazel.					
2112.	<i>Fuligula marila</i> .	Nov. 13th.	19.50	32.80	♂ juv.	
	Wing, 8.90; tail, 2.70; culmen, 1.85; gape, 2.28; tarsus, 1.65; middle toe and claw, 2.88; claw alone .38; toe alone, 2.58. Iris bright gold-color. Shot at head of flock at long range.					
2113.	<i>Arquatella maritima</i> .	Nov. 13th.	9.35	16.40	♂ ad.	
	Wing, 5.00; tail, 2.45; culmen, 1.37; gape, 1.36; tarsus, .93; middle toe and claw, 1.68; middle toe alone, .91; its claw, .19. Iris, hazel. Bill, yellow at base, shading into greenish-yellow at base in middle; terminal one-third, black. Legs + feet, olive-yellow; nails, black.					
2114.	<i>Arquatella maritima</i> .	Nov. 13th.	8.90	15.60	♂ ad.	
	Wing, 4.87; tail, 2.45; culmen, 1.10; gape, 1.10; tarsus, .90; middle toe and claw, 1.05; middle toe, .86; claw, .21. Iris, bill + feet, same as above. Colors in this & the three succeeding specimens the same as above (No. 2113). Their stomachs contained small univalves and other shells besides sand. Iris, hazel.					

1880.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
2115.	<i>Arquatella maritima</i> .	Nov. 13th.	9.33	16.25	♀ ad.	
	Wing, 5.10; tail, 2.41; culmen, 1.38; gape, 1.36; tarsus, .98; middle toe and its claw, 1.07; middle toe, .10; its claw, .23. Iris, hazel.					
2116.	<i>Arquatella maritima</i> .	Nov. 13th.	9.10	16.00	♀ ad.	
	Wing, 5.00; tail, 2.30; culmen, 1.28; gape, 1.25; tarsus, .95; middle toe and claw, 1.10; middle toe, .90; its claw, .23. Iris, hazel.					
2117.	<i>Arquatella maritima</i> .	Nov. 13th.	8.90	15.80	♀ ad.	
	Wing, 4.85; tail, 2.40; culmen, 1.25; gape, 1.22; tarsus, .93; middle toe + claw, 1.12; middle toe, .92; claw, .23. Iris, hazel.					

November 14th.

This afternoon Morris & I went to Central Park to see the Gulls. Nearly all were sitting on the division between the two parts of the large reservoir. We were not permitted to go out there to see what species there were; but those seen flying were all Herring Gulls. Four Cardinal Redbirds (*Cardinalis virginianus*) were seen, two males and two females, in a part of the park where I have never seen the species before; but Morris has seen it there in spring. Two females and a male were in sight at once. Their notes were usual.

November 17th. New York City.
Three Black Snowbirds (*Junco hyemalis*) are eating crumbs on the roof opposite our window.

November 21 & 22. Highland Falls, N.Y.
Sialia sialis. - Bluebird.
Regulus satrapa. - Golden-crested Kinglet.
Parus atricapillus. - Chickadee.
Sitta carolinensis. - White-bellied Nuthatch.
Certhia familiaris. - Brown Creeper.

1880.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

Astragalinus tristis. - Yellowbird.
Spizella montana. - Tree Sparrow.
Junco hyemalis. - Snow-bird.
Passer domesticus. - European House Sparrow.
Corvus americanus. - Crow.
Bonasa umbellus. - Ruffed Grouse.
Larus argentatus. - Herring Gull.

The above species were all that I saw
 although I took a long tramp on Sunday.
 No Fox Sparrows, Cedar-birds, Robins or
 Yellow-rumped Warblers were seen.

November 19th. Sayville, L. I., N.Y.

The following birds were seen along the
 shore of Great South Bay, at Sayville; the
 weather was too severe to permit us to go
 out in the battery, so we spent the
 morning on the bay shore, and sailed
 across the bay in the afternoon and
 anchored on the shooting grounds near
 the beach and spent the night in very
 close quarters. At three o'clock on the follow-
 ing morning (Nov. 20th) we rose, and, finding
 the wind had again changed to the North
 and was blowing hard, we weighed anchor
 and tacked across the bay to Sayville, in
 time to get breakfast and take the first
 train to New York. The train scared up a flock of
Ortyx virginianus. near Babylon.
Harporhynchus rufus. One seen in the woods
 near the marshes on the bay.

Parus atricapillus.

Carpodacus purpureus. - Cedars on the bay.

Astragalinus tristis. - In enormous flocks.

Chrysomitris pinus. - Flying in small
 companies high in the air.

Melospiza fasciata.

1880.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

Melospiza palustris. - Several were seen on creeks.

Spizella montana. - Very plentiful.

Passerculus savanna. - A few along the bay.

Corvus frugivorus. - Very abundant.

Cyanocitta cristata. - Abundant.

Circus hudsonius. - One on marsh along bay.

Ortyx virginianus.

Branta canadensis. - Not yet abundant on the bay.

Pelionetta perspicillata. - One male shot.

Melanetta velutina. - Very abundant.

Harelda glacialis. - Very abundant.

Fuligula marila.

Colymbus torquatus. - Becoming less abundant
 than a few weeks ago.

Larus argentatus. - The only gull seen.

Falco columbarius. - One flew close past.

Pelionetta perspicillata. 18.85 32.10 ♂ juv.

Wing, 9.75; tail, 3.65; culmen, 1.36; gape, 2.18;
 tarsus, 1.65; middle toe and claw, 2.73;
 middle toe alone, 2.40; claw, .35. Iris, hazel.

This bird was very small, lacked the white patch
 on the top of head, but possessed the glossy
 black plumage of the adult male, and
 had the triangular white patch on the nape
 well developed. The bill lacked the red color
 of the adult, but had a few irregular tawny
 blotches, rather indistinct. I shot it swimming
 along the edge of a marsh on the bay shore.
 Is this the last stage through which the
 male bird passes before arriving at the
 perfect adult plumage? viz: having
 black plumage and white nape complete,
 but lacking red color of bill, white iris
 and white patch on top of the head.

Sciurus carolinensis. - Gray Squirrels were quite
 numerous in the oak woods. I shot a very large
 one, and gave to Louis Beres a specimen.

2,118

1880

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

On November 22d, I took the six o'clock P.M. train from Jersey City for Circleville, Ohio. Slept comfortably until daylight; then I did the curtain & looked out of the car window from bed. We were three or four hours east of Pittsburg, and, although I kept a lookout for birds, only one small bird (probably a Sparrow of some sort) was seen east of that city. From Pittsburg to ~~Columbus~~ ^{Stoubeville}, Ohio, a few flocks of Carolina Doves, Tree Sparrows and Black Snowbirds were all the birds noted.

Stoubeville is situated on the west bank of the Ohio River, which forms the boundary between Pennsylvania and Ohio. From this place westward to Columbus the birds increased constantly in number and variety. The following species were noted: Turdus migratorius. - Three near Cadiz Junction. Lanius borealis. - Several Shrikes, which I noted at the time as Great Northern were seen sitting upon trees at various points along the Pan Handle Road; but Dr. Howard Jones thinks they were Logger-headed Shrikes. Dr. J. M. Wheaton, however, informed me that it does not occur in winter about the city of Columbus, where I saw specimens of L. borealis that were collected there by Mr. Oliver Davis. I think that my observations were correct, and that the Shrikes seen east of Columbus were all Great Northern Shrikes.

Junco hyemalis. - Abundant

Spizella montana. - do

Sturnella magna. - do

Colaptes auratus. - do

Ovis accipitrinus. - Two seen flying about in bright sunshine -

1880.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

Buteo borealis. - Quite numerous.

Buteo lineatus. - do

Buteo pennsylvanicus. - One set close to the railroad as we passed. I think several were seen. Zenaidura carolinensis. - In flocks of six to twenty all along the road from Pittsburg to Columbus.

The first snow was encountered fifty miles east of Pittsburg, Pa., and the weather during the previous night had been severely cold. At Columbus the snow was several inches deep and the temperature several degrees below zero: the same at Circleville, where I arrived at seven o'clock P.M. November 23d. On the following morning, ^{Nov. 24th} Ed. Wittich took me down to Drs. C. M. & Howard Jones's office and introduced me to the old Doctor and his son - Dr. Howard, who writes the text accompanying the illustrations of the work entitled "Illustrations of the Nests & Eggs of Ohio Birds," by Messrs. Jones & Schulze. In answer to my inquiries concerning the winter birds I learned many interesting things, among them - that the Turkey Buzzards go to roost every night in a certain woods upon the Scioto River, six miles north of Circleville. The old doctor brought in a Virginia Rail which his spaniel had caught. It was chewed into a pulp, but I preserved its head which was all right. I formed a very pleasant impression of the Jones's, and promised to call on the following morning.

On Thursday (Nov. 25th) morning I again called upon Dr. Howard Jones, who showed me three plates from Mr. J. J. Audubon's elegant folio edition of Birds of America - the large-called Puffin, the

1880.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

American Avocet, and the Pinnulated Duck, which he had received from Mr. Wade, of Rockville, Connecticut. I was then taken up stairs and presented to Mrs. Dr. Jones (his mother), who was engaged in drawing the nest of *Zenaidura macroura* with its two eggs, upon stone for the engraving. I was also introduced to Miss ^{Nephtis} Jacobs who was coloring one of the plates. She was working slowly and carefully upon the foliage and blossoms of the standing blackberry (*Rubus vellosus*). I saw the recently completed drawing on the stone of the Blue-gray Gnatcatcher (*Polioptila caerulea*) and Ruby-throated Hummingbird (*Trochilus colubris*) both nests drawn on one plate. In a back room were found the greater part of Dr. Jones's collection of nests and eggs of Ohio Birds, each nest accompanied by its set of eggs and resting upon the branch or materials just as it was found, and accompanied by a paper giving a full account of its situation, environments, materials, habits of birds etc., all to be used by Dr. Jones in his descriptive text. Down stairs I was shown a remarkably fine series of the nests and eggs of the Acadian Flycatcher (*Empidonax acadicus*) whose nest Dr. Howard has been especially fortunate in finding. Mrs. Jones was very courteous; and kindly showed me most of the drawings used in their work. Dr. Jones offered to give me some of the nests and eggs figured, and afterwards brought me a nest and set of eggs of the Virginia Cardinal (*Cardinalis virginianus*) taken at Circleville, and

1880.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

promised me the specimen figured in their plate as soon as the text pertaining to that species has been written. The Doctor promised to call for me to go out with him in search of specimens as soon as his sore eyes would permit.

In the afternoon Nov. 25th. Circleville, Ohio. Alph Wittich went out shooting with me. We walked a couple of miles up the canal, without seeing much Save Quail's tracks and some Swamp Sparrows. We were going into a woods beside the canal, when (we saw) short-eared Owls (*Asio accipitrinus*) flying about the prairie to the left. I crept out through the deep snow to the middle of the prairie, scraped away the snow from a small place, and waited for one of the Owls to fly near enough for a shot. Soon one came along and did not notice me, for I was dressed in brown shooting coat & light cap. When he got directly overhead I arose, and, after letting it fly a little way off, shot it. ✓ Meadow Larks (*Sturnella magna*) were buried deep in the snow and flew out with a loud whirr when nearly trodden upon. It was growing dark so, although I shot several, I only succeeded in finding one. Whether they had dropped down into the snow simply to roost for the night, or had allowed them to be snowed under I could not tell.

I made a game-piece of the Lark. Its length was 10.65; stretch, 15.60. ♂ ad

Friday, November 26th Circleville, Ohio.
After dinner Ed Wittich borrowed my

1880.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

and we set out together. We walked up the canal, and then walked across to the Scioto River at a point north of Craus's Island. We crossed the river road, where a large flock of Shore Larks (*Eremophila alpestris*) was feeding in the road. They were the first I ever saw living. I shot four specimens, and also shot a Carolina Dove there. One Dove was so tame that I attempted to catch it in my hands, and actually almost clutched it before it flew. On Craus's Island we saw several Redbirds (*Cardinalis virginianus*), and I shot two males and a female. We also saw a Red-headed Woodpecker & Flickers.

Saturday, Nov. 27th Circleville, Ohio.

Dr. Jones called for me with a sleigh, for a collecting excursion, but did not bring his gun, as his eyes still troubled him. We drove three miles south into a large woods, where a couple of Great-horned Owls were seen. Dr. Jones shot at a Hairy Woodpecker, a Redbird, and at a Tufted Titmouse, but the small charges failed to kill any of them. We put up a large flock of Quail. Tufted Titmice were very numerous. I saw them that morning for the first time and shot four nice specimens besides a male White-bellied Nuthatch. We were only out a couple of hours or so. Three Redbirds were seen in the trees in the city.

Sunday, Nov. 28th Circleville, Ohio.

I received an invitation to dine at the Hulse's, at their farm four miles west of Circleville. The two young ladies and their "aunt Hepais" were very agreeable

1880.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

Kind people; and the brother, Isaiah Hulse is a good fellow. Doves were seen all along the road, picking up the corn which had spilled from the wagons on the previous day. They were in flocks of variable size, also single, and in pairs. As a rule they would allow us to drive close up to them, and then merely flew up on the fence beside the road or upon the nearest bush. Flocks of twenty to thirty birds were not uncommon. Several times we stopped the carriage and looked at those that were sitting on the fences beside the road, within a few feet of us. Shore Larks were seen all along the way in small straggling flocks. Jays, Meadow Larks & Red-headed Woodpeckers were plentiful. One flock of Quail was seen in a perfectly open field, near the road, and only a couple of rods from a house. I saw the first Logger-headed Shrike sitting on a sharp fence-post beside the road. We stopped close to it and took a good look; as we started on it flew to the nearest corn shock and wagged its tail until we were out of sight. This was the only Logger-headed Shrike I saw during the 13 days I spent at Circleville; but Dr. J. M. Wheaton presented me with a nice female shot by himself at Columbus, in April, 1876. Ed. & I accepted an invitation to go out the next day and shoot during the afternoon, which we were glad to do.

Monday Nov. 29th Circleville, Ohio.

We were delayed in starting for Hulse's, but got off in a light buggy at half-past one o'clock P.M. We got to the farm at 2 25.

1880

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

started out soon after. I shot a Dove flying in the orchard, three Rabbits & a Chipmunk in the woods, five Quails and a Bank Mouse (*Arvicola riparia*) on the prairie, and a Red-headed Woodpecker in the big elm-tree beside the house.

We bagged six Rabbits, 13 Quails & 1 Dove, together, for game. Ed says the Red-bird.

Tuesday Nov. 30th. Circleville, Ohio.

In the afternoon Ella Wittich & I went out in a buggy. We tried to get a shot at a flock of Doves upon a board-fence, but they flew before we could tie the horse.

I shot two Doves at a shot, in an apple-tree, besides a Meadow Lark. There were numbers of Meadow Larks and large & small flocks of Shore Larks. We saw one Red-bird (?) and the first Bluebirds I saw in Circleville.

Wednesday, Dec. 1st. Circleville, Ohio

I went up the river alone and shot two Red-birds & two Quails. Missed or failed to kill a Sparrow-Hawk, Great-horned & Barred Owls. Saw Towhee Buntings & large flocks of Red-birds.

Thursday Dec. 2d. Circleville, Ohio.

Ed & I set out early in the morning for Hulse's farm. Jays & Red-headed Woodpeckers were very numerous in the woods about the house. We shot 13 Woodpeckers & 3 Jays in a short time, and killed some rabbits and Quail before dinner. After dinner we drove up to another of Hulse's places, where we got out of the wagon and walked home. Altogether we

1880

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

bagged for game. 5 Rabbits, 8 Quails & one Fox Squirrel. My own bag for the day was. 1 Rabbit, 1 Fox Squirrel, 3 Quails, 10 Red-headed Woodpeckers, 1 Red-bellied Woodpecker, 1 Downy Woodpecker, 1 Towhee Bunting, & 3 Jays. Fox Sparrows were observed about the back orchard.

Friday Dec. 3d. Circleville, Ohio

Spent the day skinning birds & with the Jones's.

Saturday Dec. 4th. Circleville Ohio.

Set out alone for Cram's Island, walking up the river. Where the creek empties into the river I shot the first Great Carolina Wren I ever saw living. It fell up on the edge of the ice where I had to crawl upon all fours in order to reach it without breaking through the ice. I saw and shot one other specimen of this Wren far up a creek to the north of the island. Before I reached the island I killed a male Redbird beside the river. Fox & White-throated Sparrows were quite numerous along the creek & on the island. So were Towhee Buntings. The Great-horned Owls were seen; but no Quail were put up. My bag amounted to:— One Rabbit, three Tufted Titmice, 8 Red-birds, 1 Towhee Bunting, 1 Sparrow Hawk, and 2 Great Carolina Wrens.

Sunday Dec. 5th. Circleville, Ohio.

Stayed in the house all day save an hour's walk about the city in the afternoon. The Tufted Titmouse was seen in the city about the porches of the houses and in the trees. Gray Squirrels are abundant in the "bone orchard" where are also

1880.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

Monday, Dec 6th Circleville, Ohio.
Saw a Cardinal Red-bird in the city in the morning. On our way to the 3:32 P.M. train, Ed. & I saw a Marsh Harrier. I arrived at Columbus at 4:50 P.M., and went directly to Dr. J. M. Wheaton's, and there spent the evening. The old gentleman received me very cordially. He handed me a Duck's Head to diagnose. I said it might be a Black Duck's; but it turned out to be a female Eider - the first record for Ohio, where, however, the King Eider has been taken. Dr. Wheaton then presented me with a copy of his report on "The Food of Birds as Related to Agriculture" published in 1875, after making several additions in pencil and repaging to correspond with the pagination of the Agricultural report in which it was printed. He then presented me with some specimens which he had collected, and showed me his mode of bookkeeping. He keeps one page of his visiting book for his patients and the other for the Birds. We discussed the local fauna; and he alluded to the great difference between that of Columbus & of the Scioto Valley in the region of Circleville where I had been collecting, stating that the fauna of Circleville was markedly more carolinian than that about Columbus. He was surprised to learn that Turkey Buzzards roosted north of Circleville during winter as Dr. Jones had informed me, and inquired specially whether I had

1880.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

seen any Logger-headed Shrikes. He said that Cardinal Red-birds, Tufted Titmice and Red-bellied Woodpeckers wintered at Columbus; and also the Great Carolina Wrens, rarely; but the Shrikes he has never known to do so, and the Turkey Buzzards have not been seen there in winter of late years.

Late at night, on my way to the train, I called on Mr. Oliver Davie, who keeps a Natural History Store at No. 170 North High St., Columbus. He showed me his entire stock of birds, mounted & in skin, and birds' eggs. I bought some specimens, and had a pleasant chat about the local birds, finding Mr. Davie very well posted on the local avi-fauna, as well as general.

Tuesday, Dec 7th. Columbus, O. to Jersey City, N.J.
Saw a few Cardinal Grosbeaks between Steubenville, O. and Pittsburg, Pa. Mostly near the Ohio valley.

I saw no Quail on my journey east, as I passed through the best quail country in the night; but, on my way out they were very abundant in Ohio. I saw six flocks between Columbus and the first station east of there. Some were so tame that they merely flew from the track upon a board-fence at the side, and sat there until the train had passed by.

No Crows were seen until we were 50 miles east of Altuna, Pa., where I saw the first Crows seen since leaving New York on Nov. 22d. Dr. Wheaton informed me that it is rare in winter about the vicinity of Columbus in winter although he gives it a permanent

1880.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

resident in his list.

When at Dr. Wheaton's house, he showed me a book in which he records the occurrence of birds in his little yard in the city. He has found exactly 100 species, only one of which is at all doubtful. Among them are the Green Heron, and Quail; of the latter he says he has sometimes procured a mess for his small family during the autumn migration by firing a couple of shots from his back door.

List of Birds seen at Circleville, Ohio.

November 23d to Dec. 6th

1. Sialia sialis. - Quite numerous; singing.
2. Regulus satrapa.
3. Certhia familiaris.
4. Lophophanes bicolor. - Abundant in all situations; extremely numerous in the woods along the river & creeks. Noisy.
5. Parus atricapillus. - Not very abundant, but possibly var. carolinensis.
6. Setta carolinensis.
7. Setta canadensis.
8. Thryothorus ludovicianus. - Dr. Jones shot one Nov. 27th or 28th, at his house in the city, and skinned it for me. I only saw two, both of which I shot. Isaac Hulse said they were seen constantly about their house, and were only absent when I came after 'em.
9. Ampelis cedrorum.
10. Lanius ludovicianus. - One seen.
11. Eremophila alpestris. - Abundant in different sized flocks; some very large.

1880.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

12. Astragalinus tristis. - Abundant.
13. Melospiza fasciata. - In full song.
14. Melospiza palustris. - Quite abundant.
15. Junco hyemalis.
16. Spizella montana.
17. Zonotrichia albicollis. - Quite numerous.
18. Passerella iliaca. - do
19. Cardinalis virginianus. - Seen near by everywhere: twice in the city. Some flocks contained thirty to fifty specimens. Was singing on the 4th of December.
20. Pipilo erythrophthalmus. - Quite plentiful.
21. Passer domesticus. - Abundant in town.
22. Cyanocitta cristata. - Very abundant.
23. Ceryle alcyon. - Seen December 1st.
24. Picus pubescens.
25. Picus villosus.
26. Centurus carolinus. - Only two were seen both on December 4th. I shot the one I saw. Recognized it at once by its different note, although in a tree top in a woods where Melanerpes were very numerous.
27. Melanerpes erythrocephalus. - Very abundant in the woods everywhere, croaking and rattling & chasing one another.
28. Colaptes auratus. - Abundant everywhere.
29. Asio accipitrinus. - Seen on two days.
30. Syrnium nebulosum. - Two seen.
31. Scops asio. - One seen.
32. Bubo virginianus. - One or two in nearly every large wood.
33. Circus hudsonius. - Seen Nov. 29 & Dec. 1.
34. Falco columbarius.
35. Falco sparverius. - Quite numerous. One was so gentle that we stopped the carriage beneath the tree on which it sat and shouted, before it would fly.

1880.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.36 *Buteo borealis*.37 *Buteo lineatus*.38 *Zenaidura carolinensis*. - Numerous.

In flocks of six to forty; sometimes single.

39. *Oxyechus virginianus*. - Very plentiful. Ishot one with a black mark down the
centre of the white throat. I lost my
first trail. Killed ten altogether.40. *Gallinago wilsoni*. - Dr. Howard Jones
shot two on November 23d.41. *Rallus virginianus*. - Dr. Howard
Jones shot two on Nov. 23d, when
the thermometer was near zero,
and the snow deep on the ground.
Old Dr. Jones brought in one which
his Spanish caught, up the canal,
on Nov. 24th. He saw three, but
would not shoot them. I preserved
the head of this bird, its body having
been mangled by the dog.42. *Branta canadensis*. - In the
corn-fields.Specimens collected at Circleville, O.

- 2,119. *Asio accipitrinus*. Nov. 25. 41.00
2,120. *Falco sparverius*. Dec. 4. 10.12 21.75 ♂ ad.
Wing, 7.00; tail, 4.65; culmen from frontal
feathers, .58; culmen from cere, .50;
cere, .12; gape, .70; tarsus, 1.43; ~~tarsus~~
middle toe and claw, 1.20; middle
toe, .88; its claw, .40. Iris bright hazel.
- 2,121. *Picus pubescens*. Dec. 2. 16.75 10.88 ♂ ad.
Wing, 3.75; tail, 2.53; culmen, .73; gape,
.78; tarsus, .73; middle toe & claw, .70;
middle toe alone, .46; claw, .34. Ella Wittich.

1880.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.2,122. *Centurus carolinus*. Dec. 2. 10.00 16.65 ♂ ad.
Wing, 5.20; tail, 3.45; culmen, 1.27; gape, 1.48;
tarsus, .82; middle toe and claw, 1.12; middle
toe, .75; claw, .45. Iris, bright carmine.2,123. *Melanerpes erythrocephalus*. Nov. 29. 9.70 17.00 ♂ ad.
Wing, 5.08; tail, 5.50; culmen, 1.10; gape, 1.30;
tarsus, .87; middle toe and claw, .97;
middle toe, .70; claw, .35.2,124. *Melanerpes erythrocephalus*. Dec. 2. 9.62 16.87 ♀ ad.
Wing, 5.30; tail, 3.56; culmen, 1.07; gape, 1.25;
tarsus, .92; middle toe & claw, .95; middle
toe alone, .70; claw, .37.2,125. *Melanerpes erythrocephalus*. 9.50 16.75 ♀ ad.
Wing, 5.31; tail, 3.45; culmen, 1.07; gape,
1.27; tarsus, .93; middle toe and claw, .97;
middle toe, .69; claw, .37. Ella Wittich.2,126. *Melanerpes erythrocephalus*. 10.25 17.24 ♂
Wing, 5.18; tail, 3.50; culmen, 1.18; gape, 1.37;
tarsus, .91; middle toe and claw, 1.03;
middle toe, .76; claw, .39.2,127. *Melanerpes erythrocephalus*. 9.50 16.75 ♀ ad.
Wing, 5.00; tail, 3.36; culmen, 1.10; gape, 1.35;
tarsus, .96; middle toe and claw, .99; middle toe,
its claw, .36.2,128. *Melanerpes erythrocephalus*. 10.25 17.24 ♀ juv.
Wing, 5.18; tail, 3.50; culmen, 1.18; gape, 1.37; tarsus,
.91; middle toe and claw, 1.03; middle toe, .76;
its claw, .39.2,129. *Cyanocitta cristata*. Dec. 2. 11.85 16.62 ♂ ad.
Wing, 5.13; tail, 5.25; culmen, 1.12; gape,
1.25; tarsus, 1.57; middle toe and claw, 1.20
middle toe, .93; claw, .36. Ella Wittich.2,130. *Cyanocitta cristata*. Dec. 2. — — ♀ ad.2,131. *Sturnella magna*. Nov. 30. 10.25 16.30 ♂ ad.
Wing, 4.90; tail, 3.20; culmen, 1.32; gape,
1.2; tarsus, 1.66; middle toe and claw, 1.33;
middle toe, 1.07; claw, .31.

1880.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
2,132.	<i>Zenaidura carolinensis</i> .	Nov. 26.	11.75	17.20	♀ ad.	
2,133.	<i>Zenaidura carolinensis</i> .	Nov. 30.	11.75	17.50	♀ ad.	
	Wing, 5.60; tail, 5.40; culmen, .50; gape, .79; tarsus, .80; middle toe and claw, 1.00; middle toe, .77; claw, .25.					
2,134.	<i>Zenaidura carolinensis</i> .	Nov. 30.	11.70	17.85	ad.	
	Wing, 5.65; tail, 5.15; culmen, .52; gape, .79; tarsus .83; middle toe and claw, 1.00; middle toe, .79; claw, .25.					
2,135.	<i>Ortyx virginiana</i> .	Dec. 2.	9.75	15.15	♂ ad.	
	Wing, 4.50; tail, 2.70; culmen, .55; gape, .62; tarsus, 1.31; middle toe and claw, 1.43.					
2,136.	<i>Ortyx virginiana</i> .	Dec. 2.	9.65	14.70	♂ ad.	
	Wing, 4.28; tail, 2.60; culmen, .55; gape, .62; tarsus, 1.25; middle toe and claw, 1.44. ^{irrad} A black stripe down white of throat.					
2,137.	<i>Ortyx virginiana</i> .	Dec. 1.	9.75	15.35	♂ ad.	
	Wing, 4.38; tail, 2.67; culmen, .55; gape, .67; tarsus, 1.29; middle toe and claw, 1.46.					
2,138.	<i>Ortyx virginiana</i> .	Dec. 1.	9.90	15.00	♀ ad.	
	Wing, 4.50; tail, 2.80; culmen, .53; gape, .66; tarsus, 1.20; middle toe and claw, 1.37.					
2,139.	<i>Ortyx virginiana</i> .	Nov. 29.	9.75	15.10	♂ ad.	
	Wing, 4.62; tail, 2.85; culmen, .57; gape, .65; tarsus, 1.30; middle toe and claw, 1.40; claw, .37; toe alone, 1.10.					
2,140.	<i>Ortyx virginiana</i> .	Nov. 29.	10.00	15.30	♀ ad.	
	Wing, 4.63; tail, 2.80; culmen, .57; gape, .67; tarsus, 1.32; middle toe and claw, 1.39; middle toe, 1.09; claw, .32.					
2,141.	<i>Ortyx virginiana</i> .	Nov. 29.	9.80	15.10	♀ ad.	
	Wing, 4.60; tail, 2.88; culmen, .56; gape, .64; tarsus, 1.25; middle toe and claw, 1.40; middle toe, 1.09; claw, .31.					
2,142.	<i>Ortyx virginiana</i> .	Nov. 29.	10.25	15.15	♂ ad.	
	Wing, 4.40; tail, 2.87; culmen, .59; gape, .64; tarsus, 1.37; middle toe and claw, 1.48; middle toe, 1.24; toe alone, .70.					

1880.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
2,143.	<i>Ortyx virginiana</i> .	Nov. 29th.	9.40	15.40	♂ ad.	
	Wing, 4.70; tail, 2.65; culmen, .55; gape, .61; tarsus, 1.38; middle toe and claw, 1.45; middle toe, 1.10; claw, .31. Black dashes on throat.					
2,144.	<i>Ortyx virginianus</i> .	Nov. 29th.	9.90	15.10	♂ ad.	
	Wing, 4.55; tail, 2.80; culmen, .56; gape, .62; tarsus, 1.30; middle toe and claw, 1.38; middle toe, 1.15; claw, .35. Irides, hazel.					
2,145.	<i>Lophophanes bicolor</i> .	Nov. 27th.	6.55	10.25	♂ ad.	
	Wing, 3.10; tail, 2.90; culmen, .48; bill from nostril, .37; gape, .54; tarsus, .82; middle toe and claw, .75; middle toe, .54; claw, .27. Sex carefully determined.					
2,146.	<i>Lophophanes bicolor</i> .	Nov. 27th.	6.85	10.75	♀ ad.	
	Wing, 3.38; tail, 3.20; culmen, .52; bill from nostril, .42; gape, .57; tarsus, .85; middle toe and claw, .75; middle toe, .55; claw, .26. Sex carefully examined.					
2,147.	<i>Lophophanes bicolor</i> .	Nov. 27th.	6.10	10.00	♀ ad.	
	Wing, 3.00; tail, 2.65; culmen, .50; bill from nostril, .37; gape, .55; tarsus, .82; middle toe and claw, .76; middle toe, .54; claw, .29.					
2,148.	<i>Lophophanes bicolor</i> .	Nov. 27th.	6.50	10.50	♂ ad.	
	Wing, 3.15; tail, 2.97; culmen, .47; bill from nostril, .36; gape, .56; tarsus, .82; middle toe and claw, .78; middle toe, .57; claw, .27.					
2,149.	<i>Lophophanes bicolor</i> .	Dec. 4th.	6.00	9.50	♀ (?)	
	Wing, 2.82; tail, 2.65; culmen, .46; bill from nostril, .35; gape, .50; tarsus, .78; middle toe and its claw, .65.					
2,150.	<i>Sitta carolinensis</i> .	Nov. 27th.	6.00	11.40	♂ ad.	
2,151.	<i>Cardinalis virginianus</i> .	Nov. 26.	9.05	12.00	♂ ad.	
	Wing, 3.62; tail, 4.40; culmen, .70; gape, .74; tarsus, 1.02; middle toe and claw, .92; middle toe, .65; claw, .29. Iris, hazel.					
2,152.	<i>Cardinalis virginianus</i> .	Nov. 26.	8.75	11.60	♂ ad.	
	Wing, 3.62; tail, 4.20; culmen, .73; gape, .73; tarsus, .97; middle toe and claw, .95.					

1880.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
2,153.	<i>Cardinalis virginianus</i> .	Nov. 26.		8.75	11.55	♀ ad.
	Wing, 3.50; tail, 4.10; culmen, .72; gape, .72; tarsus, .97; middle toe and claw, .90; middle toe, .70; claw, .22. Iris, hazel.					
2,154.	<i>Cardinalis virginianus</i> .	Dec. 1.		8.90	11.90	♂ ad.
	Wing, 3.65; tail, 4.23; culmen, .70; gape, .73; tarsus, 1.00; middle toe + claw, .96.					
2,155.	<i>Cardinalis virginianus</i> .	Dec. 1.		8.40	11.55	♀ ad.
	Wing, 3.70; tail, 4.05; culmen, .63; gape, .65; tarsus, .92; middle toe + claw, .90.					
2,156.	<i>Cardinalis virginianus</i> .	Dec. 2.		9.10	12.10	♂ ad.
	Wing, 3.80; tail, 4.30; culmen, .71; gape, .72; tarsus, 1.00; middle toe and claw, 1.00.					
2,157.	<i>Cardinalis virginianus</i> .	Dec. 4.		8.70	11.90	♂ ad.
	Wing, 3.80; tail, 4.10; culmen, .75; gape, .75; tarsus, .95; middle toe and claw, .87.					
2,158.	<i>Cardinalis virginianus</i> .	Dec. 4.		8.85	11.90	♂ ad.
	Wing, 3.70; tail, 4.35; culmen, .70; gape, .72; tarsus, .93; middle toe and claw, .95.					
2,159.	<i>Cardinalis virginianus</i> .	Dec. 4.		8.50	11.60	♀ ad.
	Wing, 3.60; tail, 4.25; culmen, .70; gape, .70; tarsus, .95; middle toe + claw, .95.					
2,160.	<i>Cardinalis virginianus</i> .	Dec. 4.		8.50	11.85	♀ ad.
	Wing, 3.65; tail, 4.10; culmen, .67; gape, .68; tarsus, .94; middle toe and claw, .90.					
2,161.	<i>Pipilo erythrophthalmus</i> .	Dec. 2.		8.45	11.20	♂ ad.
	Wing, 3.35; tail, 3.90; culmen, .61; gape, .72; tarsus, 1.07; middle toe + claw, 1.05. Iris, red.					
2,162.	<i>Pipilo erythrophthalmus</i> .	Dec. 4.		8.20	11.40	♂ ad.
	Wing, 3.20; tail, 3.69; culmen, .55; gape, .68; tarsus, 1.07; middle toe + claw, 1.07.					
2,163.	<i>Eremophila alpestris</i> .	Nov. 26.		7.40	14.00	♂ ad.
	Wing, 4.50; tail, 3.00; culmen, .50; gape, .66; tarsus, .91; middle toe and claw, .76; middle toe, .50; claw, .30.					
2,164.	<i>Eremophila alpestris</i> .	Nov. 26.		7.75	14.10	♂ ad.
	Wing, 4.45; tail, 2.80; culmen, .55; gape, .70;					

1880.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	tarsus, .93; middle toe and claw, .82; middle toe, .55; claw, .30.					
2,165.	<i>Eremophila alpestris</i> .	Nov. 26.		7.25	13.00	♂ ad.
	Wing, 4.00; tail, 2.85; culmen, .50; gape, .62; tarsus, .90; middle toe and claw, .77; middle toe, .48; claw, .31.					
2,166.	<i>Eremophila alpestris</i> .	Nov. 26.		7.50	13.40	♀
2,167.	<i>Thryothorus ludovicianus</i> .	Dec. 4.		5.78	7.75	♀
	Wing, 2.28; tail, 1.93; culmen, .66; gape, .83; tarsus, .85; middle toe + claw, .82; middle toe, .65; claw, .21; bill from nostril, .46.					
2,168.	<i>Thryothorus ludovicianus</i> .	Dec. 4.		5.85	7.80	♀
	Wing, 2.27; tail, 2.08; culmen, .72; gape, .88; bill from nostril, .47; tarsus, .75; middle toe and claw, .75; middle toe, .58; claw, .21. Irides, hazel.					

December 6th. New York City.

Rob. L. Morris saw a Fox Sparrow (*Passerella iliaca*) in old Trinity Churchyard on above date.

December 9th + 10th. Highland Falls, N.Y.

A few Herring Gulls were the only water birds seen. *Sialia sialis*. - Singing. - species whose occurrence is worth noting. Did not go about the woods any.

December 12th.

In the Central Park were White-throated Sparrows (*Zonotrichia albicollis*), *Regulus satrapa*, *Parus stricapillus*, *Sitta canadensis*, *Junco hyemalis*, *Melospiza fasciata*, and *Passerella iliaca*. The last species was abundant at various places in the park.

1880

Collecting Number.	NAME.	DATE	LOCALITY.	Length.	Stretch.	Sex and Age.
2169.	<i>Scoops asio</i> (gray plumage).	9.70 24.25	♀ ad.	Wing, 6.75; tail, 3.60; culmen from frontal feathers, .80; from cere, .55; cere, .30; tarsus, 1.35; gape, 1.06; middle toe and claw, 1.25; middle toe alone, .87; claw, .44. Iris, yellow. Dec. 27, 1880.		
2170.	<i>Astur atricapillus</i> .	21.50 41.50	♂ ad.	Wing, 12.50; tail, 9.50; culmen, 1.12; cere, .40; culmen from cere, .83; gape, 1.25; tarsus, 2.88; middle toe and claw, 2.23; middle toe, 1.75; claw, .67; hallux, .97; claw, 1.08; inner toe, 1.00; claw (same as toe), 1.00; outer toe, 1.12; claw, .53. Iris, bright carmine; cere, greenish-yellow; base of both mandibles, blue (except on dorsal surface of maxilla), shading into black rather abruptly; residue of bill, black. Tarsi and toes, yellow; claws, blue-black; corners of mouth, yellow. Shot on Constitution Island, on December 27, 1880. Mounted for Miss Warner.		
2171.	<i>Lanius borealis</i> .	Dec. 27, 1880. 9.90 13.75	♀ ad.	Wing, 4.28; tail, 4.48; culmen, .73; bill measured from cere ^{nostril} , .53; gape, 1.09; tarsus, 1.00; middle toe and claw, .84; middle toe alone, .59; claw, .28. Iris, dark hazel. Shot by Rob. Morris near Harrisons.		
2172.	<i>Ardea herodias</i> .	Dec. 12, 1880. 47.50 74.00	♂ juv.	Wing, 18.55; tail, 7.50; length from tip of bill to end of tail, 63.00; culmen, 5.80; gape, 7.35; bare part of tibia, 4.95; tarsus, 7.30; middle toe and claw, 5.10; middle toe alone, 4.45; its claw, .70. Iris, orange-red. This bird was shot by Smith Mandigo, on the flats of Consook Marsh, on the 12th of December. It was in good condition and evidently a strong bird. It had been		

1880.-'81.

seen about the Consook Marsh for a good while previous to its capture. At the time it was shot, the weather was cold, and a good deal of ice in the Hudson. What it meant by staying here so late is a conundrum. Its back was patched with the darker feathers of the adult plumage. Sternum preserved.

January 1st, 1881. Highland Falls, New York.
The following species have been observed here during the past week:—
Sialia sialis.—Bluebirds have been seen and heard warbling their full song nearly every day.
Regulus satrapa.—Not abundant.
Parus atricapillus.—Not abundant.
Sitta carolinensis.—
Sitta canadensis.—Only one seen.
Certhia familiaris.—Not abundant.
Anorthura troglodytes hyemalis.—One seen Dec. 27th.
Ampelis cedrorum.—One very large flock, only.
Lanius borealis.—Only one seen, and shot.
Melospiza fasciata.—Quite plentiful; singing.
Spizella monticola.—Abundant.
Junco hyemalis.—Abundant.
Passer domesticus.
Corvus americanus.—Not numerous.
Cyanocitta cristata.—One shot. Not very plentiful.
Picus pubescens.—Scarce.
Scoops asio.—One in gray plumage shot.
Circus hudsonius.—Two on Constitution Marsh.
Tinnunculus sparverius.—One on " Island.
Astur atricapillus.—One adult male shot on do.
Larus (? *argentatus*).
Chrysomitris tristis.—One flock.
Agrotis linaria.—Several flocks were seen on both sides of the Hudson. One bird shot.
Haliaeetus leucocephalus.—Several seen.

1881.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
2,173.*	<i>Astragalinus tristis</i> .	January 1st.	H.S.M.Y.	5.15	8.90	♂ ad.
2,174.	<i>Sitta carolinensis</i> .			6.05	10.90	♀ ad.

Only one flock of Ducks (probably *Fuligula marila*) was seen - on December 27th, when the river was still open. The Hudson closed on the 28th, and the thermometer ranged below zero Fahr. for several days together. No Purple Finches, Robins, Yellow-rumps or Fox Sparrows were seen; but Bluebirds warbled loudly even when the weather was coldest.

January 12th. New York City.
 To-day accepted invitation to dine with Rev. Wendell Prime, first spending the afternoon in Central Park and the American Museum. Our special object was to see what birds were in the park, but we also went to have a good look at the obelisk, which is now exactly over the pedestal, mounted like a cannon on a pivot. I was presented to Captain Forging, who promised to let us know when the thing "is over."
 We observed the following bird in the park:-
Bonasa atricapillus. - Quite numerous.
Certhia familiaris. - One heard.
Passer domesticus. - Singing quite a song.
Zonotrichia albicollis. - White-throats several seen, feeding of seeds of White Birch.
Junco hyemalis. - Not abundant.
Cardinalis virginianus. - Was delighted to find one or two Cardinal Grosbeaks in the park on this cold winter's day. I had a strong suspicion that it wintered there, and was glad to verify it. I recognized the note before we saw the bird, and another was heard, after the first, & five males, & one or two

1881.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	<i>A partial Case of Partial Melanism in the Canvas-back Duck (Aythya valisineria).</i>	January 13th.				

On my way from the New York Hospital, I found an adult Canvas-back in full plumage, having a patch of plumbeous-black feathers on the under surface of the body as well as a very long, beginning anteriorly about an inch below the termination of the normal black of the neck and breast, and extending well down on the abdomen. As the bird was badly shot and extremely so, so I simply pulled out a few of the feathers for examination. They are of about the same color as the Coat.

P.S. Color probably due to paint off of boat.

January 14th.
 Went to Roosevelt Hospital (59th-st. & 9th Avenue) at 2:30 P.M., and saw operations for lacerated cervix and perinaeum, and saw sutures removed from a uterus operated upon a week ago, in which perfect union of the parts had resulted. I left the hospital about four o'clock, and walked over to the Hudson River, and then north to about 86th-street, where I came upon a flock of thirty or thereabouts White Snowbirds (*Electrophanes nivalis*). At first they appeared very gentle, but suddenly flew away down the river, uttering their metallic twittering note. They were feeding on the seeds of the hog-weed (*Ambrosia elatior*). Not far from there I saw five Shore Larks (*Eremophila alpestris*), which flew out over the North River.
 I - - - Eastern Snowbird (*Junco hyemalis*)

1881.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	<p>4 few Tree Sparrows (<i>Spizella monticola</i>), and some English Sparrows (<i>Passer domesticus</i>), were the only other Land Birds; but Herring Gulls were seen upon the river, their wings beautifully gilded by the reflected splendor of the glowing sunset.</p> <p>A policeman whom I met said that an occasional Eagle, and numbers of Crows were often seen floating on the ice off the Hudson.</p>					

January 21st

To-night Mr. Bainbridge Bishop called with an introduction from Miss Avery. He resides in Essex County, New York, where he informed us that Ravens (*Corvus corax*) reside permanently there, are are rather numerous about Lake Champlain. I also learned that the Red Squirrel (*Sciurus hudsonius*) destroys large numbers of eggs of the Ruffed Grouse (*Bonasa umbellus*). Mr. Bishop has seen a Squirrel pick each young bird in a nest of Robins through its head till all were destroyed. The Spruce Partridge (*Canace canadensis*) is found in Essex County.

Corvus corax

Mr. Robt. J. Morris has found the Raven on Eagle Creek stillwater, near Fourth Lake, Fulton Chain, during July 1880, where also the Canada Jay (*Perisoreus canadensis*) was found.

February 13th New York City.

Morris, Louis Zerega and myself took a long walk in the Central Park. We saw several fine Gray Squirrels (*Sciurus carolinensis*) and Gray Rabbits (*Lepus sylvaticus*) and

1881.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	<p>plenty of traces of <i>Scalops aquaticus</i> for mammals. Of birds I saw:—</p> <p><i>Turdus migratorius</i>.—One was found on the brook at the head of the Loch, where there were plenty of Sumachs (<i>Rhus typhina</i>) loaded with berries. It was a fine bird, with yellow bill and black head.</p> <p><i>Certhia familiaris</i>.—One.</p> <p><i>Regulus satrapa</i>.</p> <p><i>Tarus atricapillus</i>.</p> <p><i>Corvus americanus</i>.</p> <p><i>Astragalinus tristis</i>.—At Harlem Lake.</p> <p><i>Spizella montana</i>.—A flock at Harlem Lake & others scattered about the park.</p> <p><i>Junco hyemalis</i>.—Not abundant.</p> <p><i>Zonotrichia albicollis</i>.—Abundant; Silent.</p> <p><i>Cardinalis virginianus</i>.—One female at the head of the lake, where Mr. Prime and I saw a male on January 12th.</p> <p><i>Passerella iliaca</i>.—A number were found at the same place as the Red-bird. Silent, except their usual note softly uttered.</p> <p><i>Plectrophanes nivalis</i>.—Upper part, south of Park.</p> <p><i>Melospiza fasciata</i>.—One at head Harlem Lake.</p> <p><i>Passer domesticus</i>. Noisy. Singing.</p>					

March 7th.

Went home on Saturday the 5th, and took a long walk on the following day chiefly in search of the Fox Sparrow (*Passerella iliaca*), but, although a number were seen in Central Park on February 13th, none had yet arrived in the Highlands, so far as I could discover. There were no Robins nor Phoebe, although I was informed that a flock of the former was seen a couple of weeks before. I walked and carried my

1881.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

valice filled chiefly with heavy books, from
Garrisons to Havens, and saw a fine White
headed Eagle. I crossed the river again
Monday morning on the ice, and saw
a splendid Eagle. A Robin was whistling
on top of a big white pine south of Garrison.
I saved a seat for A.K. Fisher, who called
my attention to the large numbers of Ducks -
chiefly Mergansers, and W. bustlers. There
were numbers of Herring Gulls both on
the Hudson and on Harlem River.

March 3th.

Right after lunch took Elevated car to 155th
street, where I walked through the woods
to High Bridge, turning over stones in a
fresh-water brook in search of crawfish
and Salamanders, neither of which, however,
have I found any where on the island.

There were no turtles either. The soil is
clay and sticky, and I imagined they don't
like it; still I should suppose there would
be some Salamanders amongst the stones
in the bed of the brook.

The view from the reservoir at High Bridge
is superb. From there I coasted through
the woods beside Harlem River, where the
Herring Gulls were flying, casting handsome
black shadows upon the water. There
were Song Sparrows, White-bellied Nuthatches,
Chickadees, Golden-crested Kinglets, and plenty of Crows, and Snowbirds.
I saw a fine Acadian Owl (Nyctale
acadiea) where Sherman's Creek breaks
the line of the river-bank, and a
large marsh is situated which looks like
a good place for Snipes. The Owl lit on

1881.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

the stump of a tree on the side of the hill
among the red cedars. The afternoon was
dark and cloudy. It was the first Acadian
Owl I had ever seen wild. There was a
Red Squirrel (Sciurus hudsonius) beside the
Harlem River, the first I have seen on New-
York Island. We crossed to the Kingsbridge
Road, encountering a Hawk and some
very wild country on the way. Then we
went over to the Hudson River side and
walked down through the places, and
through Audubon Park, a lovely place
such as one could not imagine to exist on
this populous island. I crossed and took
the elevated road at 125th street. On the
Hudson River were noticed a number
of Red Squirrels and the same birds
seen as on the Harlem River bank, and
in addition a flock of White-throated
Sparrows, Orioles, Robins and many
other birds had left nests in abun-
dance in all the upper part of the
island. I saw what I suppose was
a Yellow-breasted Chat's nest, and
there were Catbirds and Song Sparrows
in abundance among the patches
of briars, cat-briars, and thickets of hune
mixed with other shrubs.
It was a foggy, dark afternoon not
favorable for bird-life.

March 20th.

This morning, Macdonald, a young artist
staying at 131 W. 23d St., and myself took
the 3d Avenue Elevated railroad to 129th St.,
the terminus of the road, where we got off,
and walked across Harlem bridge into

1881.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

the cedar grove beside Harlem River. There the Song Sparrows were singing and a pair of Bluebirds were found. We crossed the bridge at 155th Street, and walked over to Trinity Churchyard. Macdonald had brought along his sketch-book and artist's stool, intending to make a sketch of Audubon's tomb. The Cemetery is a very pretty place. Large forest trees are growing abundantly, and among them were some splendid examples of our native hemlock (Abies canadensis); among the native flowering trees are some dogwoods (Cornus florida), and tulip-trees (Liriodendron tulipifera). Crossing the bridge over the boulevard that divides the cemetery we strolled down to the river-side where the Audubon vault is situated. The day was cloudy, but the Song Sparrows were singing cheerily, Red Squirrels could be heard chattering in the neighboring Audubon Park, and Nuthatches and Chickadees were making merry; the former among the branches, and the latter upon the bole of some handsome white-woods in which we saw a couple of Baltimore Orioles' nests. Carved in the plain granite above the marble entrance to the vault was simply

AUDUBON.

floral wreath and cross of mountain laurel (Kalmia) lying upon the marble; showing that the dead were not forgotten.

Right behind the vault I picked up a Hair-bird's nest which winter winds had blown from the tree overhead. To the right was described among the branches of a crooked chestnut-tree the hanging nest of the Red-eyed Vireo; and, a little to the left, saddled on a forked branch

1881.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

of a slender oak, there was a Robin's nest. Before and behind the vault stand two graceful hemlocks, and a host of little ones have essayed to grow amongst the wood plants in the pleasantly neglected grass beneath them, in company with sprouting acorns and chestnuts, just beginning their precarious existence. The graveyard slopes gently down to the ^{edge of it} Hudson, along which runs the railroad. Game pigeons were cooing upon the track, and picking up grain which had dropped from the cars. In one corner ~~stand~~ of the cemetery a red cow was tied, and stood patiently, chewing the cud. The leaves on the white-oaks rustled in the wind.

In this quiet spot, beneath the forest trees where the Robin, Red-eyed Vireo, and Baltimore Oriole may be heard singing, lies the dust of the great Audubon, beside the river he loved so well, and a way from the noise and bustle of the city which he preferred to avoid - almost in sight of the home in which he lived for many years.

After making a hasty sketch of the tomb, I walked about the place. The cow in the corner looked friendly, and was glad when I scratched its forehead, and stroked her long, shaggy winter coat of hair. As I looked towards the river a single vessel was passing, her sails flapping idly in the wind. Herring Gulls were passing; and in winter Eagles are sometimes seen floating on the ice. As we left the place, a Red Squirrel scuttled under a tomb-stone at our approach, but quickly showed its head again at my call. Another Robin's nest was noticed in an American elm (ulmus americana) near the entrance.

1881.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

March 28th. Highland Falls, New York.
On Saturday March 26th, I went home. There was no ice in the Hudson. It was thick and strong three weeks before when I last was at home, & walked across the river in both directions. Gulls and Ducks were quite numerous. No Eagles were seen.

On Sunday the 27th I took a long walk in the woods. The day was cold and very windy. Snowbirds were abundant; but no other Sparrows except the Fox Sparrow (*Passerella iliaca*) were seen. They were singing very sweetly, as usual, and were seen at Highland Falls for the first time, as none were there three weeks ago, at my last visit. Robins were abundant everywhere, but none were singing. Bluebirds kept in the cedars beside the river, out of the wind, and were silent. No Phoebe birds were seen.

On March 28th I saw the first Phoebe (*Sayornis Aeneus*). They were calling beside the warm ledges along the river, as I walked down to take ferry at Cozzen. I saw a Fish Hawk (*Pandion haliaetus Carolinensis*) soaring over the bay north of Peekskill, when coming down on the train. It is the only one seen this year - first of the season.

March 29th. New York City.
This afternoon I walked up to the Central Park with Macdonald. About the lower Lake were a hundred or more Crow Blackbirds (*Quiscalus purpureus*). They flew about continually, now alighting beside the water, again flying off to

1881.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

some clump of trees, whence they dropped one or two at a time gracefully down to the ground. They were very noisy.

On the hill near the Lake were confined, in a little pool, six fine Wood Ducks (*Aix sponsa*), two of which were males. The males uttered a note resembling the single note of *Astragalinus tristis*, only louder. They were very active, constantly running out of the water, and then diving in off the bank. A Black Swan and Barnacle Goose were confined in the same wire pen.

Near the Wood Duck's we saw a male and female Cardinal Bird (*Cardinalis Virginianus*). Robins singing.

I saw a Muskrat (*Fiber zibeticus*) swimming in the upper end of the Lake.

A Swan that sat on a rock near me, appeared to be moulting the sheath of its bill; the latter being orange-red, and black in spots. Bought Willughby's Ornithology, Eng. Edition, 1678, at Palmer's, on my walk home.

At April 1st
In a bookstore on Ann Street, I saw to-day a book that I have been looking for:

History of Long Island, Second Edition, 1843. In Vol. II is a list of the Birds of Long Island, by Dr. Jas. E. De Kay. The following is a paragraph extracted from the prefatory remarks preceding the list, which gives some information concerning breeding, etc. by symbols (+ denotes breeders); otherwise there are no annotations:

"In the 3d Vol. of the Medical Repository for the year 1800, there is a list of about

1881.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

40 species, designated by their popular names, but I am unacquainted with any other essay, towards increasing our knowledge of the birds of the district.

The price of the 21 vols. was \$15.00. Saw this work on the way home, after visiting the Fish Exhibition at Fulton Market, where I saw many beautiful species of Salmonidae and other fishes.

Measurements, etc., of Birds shot at Sayville, Long Island, by Robt. J. Morris, April 1st.

2175. Branta bernicla (Linne) Scop. 2540 48.00 ♂ ad. Wing, 13.30; tail, 4.50; culmen, 1.35; bill from nostril, .69; gape, 1.62; nail on bill, .43; nasal orifice, .36; tarsus, 2.43; middle toe and its claw, 2.33; its claw alone, .35; outer toe and claw, 2.16; its claw alone, .29; inner toe and claw, 1.90; its claw alone, .33; hallux, .55; its claw alone, .22. Irides, dark hazel.

Entire bill, legs, feet, webs and toes, black; claws, bluish-black. [My best bird.]

2176. Branta bernicla (Linne) Scop. 2430 46.50 ♂ ad. Wing, 12.50; tail, 4.50; culmen, 1.30; bill from nostril, .70; gape, 1.50; nail on bill, .40; nasal orifice, .33; tarsus, 2.40; middle toe and claw, 2.33; its claw alone, .36; outer toe and claw, 2.17; its claw alone, .30; inner toe and claw, 1.85; its claw alone, .33. Irides, dark hazel. Hallux + claw, .58; claw, .25. Entire bill, legs, toes and webs, black; claws, bluish-black. [My second (immature) bird.] Light margins to wing-coverts as in many flycatchers.

1881.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

2176a Branta bernicla (Linne) Scop. 2520 46.00 ♂ ad. Wing, 13.00; tail, 4.70; culmen, 1.35; bill from nostril, .69; gape, 1.60; nail on bill, .44; nasal orifice, .35; tarsus, 2.47; middle toe and its claw, 2.37; its claw alone, .45; outer toe and claw, 2.25; its claw alone, .36; inner toe and claw, 1.88; its claw alone, .40; hallux and claw, .55; claw, .29. [Louis J. Ziegler's] Colors of bill and feet same as above (No. 224).

2176b Branta bernicla (L.) Scop. 2375 43.50 ♀ ad. Wing, 13.40; tail, 4.40; culmen, 1.23; bill from nostril, .68; gape, 1.52; nail on bill, .43; nasal orifice, .32; ~~largest~~ greatest width of bill, .66; depth, .75; tarsus, 2.31; middle toe and claw, 2.15; its claw, .34; outer toe and claw, 2.00; its claw alone, .28; inner toe and claw, 1.78; its claw alone, .32; hallux and claw, .52; its claw alone, .22. Colors of bill and feet same as above (No. 224). This bird, Morris sent with a pair to "Geo. W. Van Sice, of the New York Bar."

2177. Glaucionetta clangula americana. 20.10 32.75 ♂ juv. Wing, 10.60; tail, 4.25; culmen, .62; bill from nostril, .87; gape, 2.25; nail on bill, .48; nasal orifice, .32; tarsus, 1.60; middle toe and its claw, 2.95; its claw alone, .45; outer toe and claw, 3.03; its claw alone, .35; inner toe and claw, 2.38; its claw alone, .38; hallux and its claw, .92; its claw alone, .27. Bill, leathery; black. Iris, gold. Tarsi and toes, brownish-yellow; webs, blackish; nails, bluish-black.

1881.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

2178. *Halidula marila*. 1920 31.50 ♂ ad.
 Wing, 8.55; tail, 3.10; culmen, 1.72;
 bill from nostril, 1.23; bill from highest
 bare prominence (name of process)
 2.15; nail on bill, .45; nasal orifice, .26;
 gape, 2.22; greatest breadth of bill, 1.05;
 tarsus, 1.55; middle toe and its claw, 2.66;
 its claw alone, .37; outer toe and claw, 2.75;
 its claw alone, .28; inner toe and claw, 2.15;
 its claw alone, .34; hallux and its claw, .85;
 its claw alone, .22.

Bill, blue, with black tip from base of
 nail to extremity, forming a concavity,
 directed backwards on either side of nail.

Posterior aspect of tarsus, under side of
 toes, and all both surfaces of web, black.

Anterior surface of tarsus, and superior
 surface of toes, bluish lead-color; claws,
 all black.

The Geese (*Branta canadensis*) were in
 thousands on the Bay, making up in
 flocks all day, and leaving in the teeth
 of norwester. The only Gulls (*Larus*)
 were of the common species (*Larus argentatus*),
 except one *L. marinus*.
~~Chilodactyls~~ and Whistlers were all paired
 i. e. in pairs. Only one pair of Velvet
 Ducks (*Cedemia* (= *Melanetta*) *melanotos*) seen.
 Robins, Meadow Larks, Song & Fox
 Sparrows all singing; and the Yellow-
 shafted Flickers were all calling.
 Two Ospreys were seen.

April 3d.

Took a long walk in Central Park with
 Louis Gorge. The Robins sang; and
 there were about 40 Herring Gulls

1881.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

on the upper reservoir, and several on
 the lower ones.

April 4th.

Received to-day from Isaiah Hulse, kindness
 of Ed. Wittich, a Red-tailed Hawk, shot at
 Circleville, Ohio, on

April 1st.

2179. *Buteo borealis*. 12.00 50.90 ♀ ad.

Wing, 15.00; tail, 9.75; culmen from base of
 cere, 1.48; cere, .55; culmen from cere, 1.10;
 gape, 1.80; tarsus, 3.20; middle toe and
 claw, 2.30; its claw alone, .84; outer toe and
 claw, 1.75; its claw alone, .70; inner toe
 and claw, 2.03; its claw alone, 1.12;
 hallux and claw, 1.92; its claw alone, 1.25.
 Iris, hazel. Cere greenish-yellow.

Bill, pale-blue, shading gradually into
 blue-black at extremity. Tarsi and toes,
 lemon-yellow. Claws, blue-black. ^{Stomach} ~~empty~~.

April 11th. Highland Falls, N.Y.

Spent Sunday at H.F., and came down on the
 first train as usual. The birds seen
 yesterday ^(both) were as follows:—

Turdus migratorius.—One was singing at
 dusk, as late as 7.10 P.M.

Sialia sialis; *Corvus americanus*, building.

Parus atricapillus; *Passerella iliaca*,

Junco hyemalis, very abundant; *Sayornis*

pusillus, plentiful; *Seiophaga ferruginea*;

Mergus serrator.—One flock in the lower south
 of Consook Island.

Philopiza meloda (= *fasciata*).

Spizopus varius.—Two seen together.

Other species were seen, although I spent a
 good part of the day walking and driving with
 Dr. Chace to Ft. Montgomery Creek to see a patient.

1881.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

April 17th

Morris and I walked in Central Park all of this morning. I picked up a Bank Mouse (*Pericoma riparia*) in the grass. There are abundant traces of its presence there in some numbers. A Mole (*Scalops aquaticus*) was also seen, and two very large Gray Rabbits (*Lepus sylvaticus*).

We entered the Park at 6th Avenue, and walked around the lower lake. Near the bridge a Black Australian Swan was sitting on a large nest of straw on a rock in the middle of neck connecting the two parts of the lower lake.

Robins were everywhere in the Park, and in full spring song. We watched the Wood Ducks and several other species (among which was a large bird which I took to be a Massena Stork, and two Cranes) in some wire enclosures with small ponds in them. The Wood Ducks were singing, and the females talked to themselves as they swam so gracefully about without the slightest apparent effort. While watching them, a fine male red bird (*Cardinalis virginianus*). There was a Brown Creeper (*Certhia familiaris*) in the evergreens beside the lower reservoir. Further north, on the west side of the Park, I saw a very unusually large Rabbit, and followed it up to get a better view, but just as I found it again my attention was called away by a new bird's song. Looking up, I saw a pair of European Goldfinches (*Carduelis elegans*). I called Rob. and we sat down on the rocks and watched the pair, and listened to them singing. They were very gittie, and one of them came close up, and

1881.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

tugged away at a big feather in a bush close by us. Four other Goldfinches flew past, while we were watching this pair. We found two more singing beside the Pool, where the White-throated Sparrow (*Zonotrichia albicollis*) was also singing its full spring song. Junco hyemalis (Black Snowbird) was abundant everywhere, and some singing. The Ruby-crowned Kinglet (*Regulus calendula*) was singing beside the Pool, and the Golden-crested (*E. satrapa*) were about the same spot. A pair (♂ & ♀) of beautiful Cardinal Birds were beside the Pool, and were very tame. Mockers (*Sayornis fuscus*) were seen in all parts of the Park. Near the Loch were seen some Yellow Red-poll Warblers (*Dendroica palmarum*); and a Winter Wren (*Anorthura troglodytes hyemalis*) was found beside the brook. In the northeast corner of the Park we watched six Fish Crows (*Corvus ossifragus*) whose notes, flight and actions were unmistakable. A watering cart came around where they were, and frightened them. They left the flock circling circularly about in the way that Common Crows do not.

Crow Blackbirds (*Luscalus purpureus*) were abundant in all parts of the Park. They sat on the floating boxes in the Lake, and ate corn belonging to the Swans, Wild Geese & Ducks. One of the Swans was sitting on an island out in the Lake. A few Purple Finches (*Carduelis purpureus*) were seen; and Song Sparrows were singing everywhere. There were no Gulls about the Reservoir. One Sparrow Hawk was observed hunting in the Park.

1881.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

April 24th.

This morning I spent in Central Park. Entered at Sixth avenue and, walking around the lower lake, crossed the bridge where the Black Swan was sitting on her nest on a rock out in the pond. A Swamp Sparrow (*Melospiza palustris*) lit on the rock, ran all around the Swan's nest, and looked at her in evident surprise.

Robins were seen in every part of the Park. One pair was busily engaged in building a nest in a white-oak tree, which was nearly completed. They were singing all over the Park.

Several Hermit Thrushes (*Turdus pallasi*) were seen about the Pool and Loch, and were very gentle.

Song Sparrows (*Melospiza fasciata*) were everywhere in abundance and singing; and quite a number of Swamp Sparrows were also seen.

White-throated Sparrows (*Spizella albicollis*) were whistling on every side.

Chipping Sparrows (*Spizella socialis*) were quite numerous in the upper part of the Park, and were the first I have seen this year.

English Sparrows (*Passer domesticus*) are building, but none of the nests contained eggs.

The Snowbirds (*Junco hyemalis*) which were plentiful a week ago had, apparently, all passed on to the North and none were seen.

The Cardinal Red Birds (*Cardinalis virginianus*) were singing about the Lake, and were seen about the Pool and Loch.

The European Goldfinch (*Carduelis elegans*, Stephens) was singing all over in the upper end of the Park, and flocks were frequently seen flying overhead. I do not know whether they wintered in the park, or not.

1881.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

Dendroica palmarum. - The Palm Warbler was singing, chirping ("tsip!"), and wagging its tail about the Pool and Loch, and Harlem Lake.

Dendroica coronata. - The Yellow-rumped Warbler was abundant, and sang in all parts of the Park.

Regulus calendula. - Singing and uttering its Wren-like note.

Cardinalis purpureus. - A fine old male Purple Finch in the best plumage flew and alighted close to where I stood.

Luscinia purpureus. - Scattered all over the park. One flock lives about the Lake; and a pair had settled to housekeeping.

Motodus ater. - Small companies of Cowbirds were seen in different places. The females appeared to be much less numerous than males. Uttered a rattling noise.

Sayornis fusca. - Two seen.

Hirundo erythrogastra. - Curious enough the Barn Swallow was the first of the family to appear. Usually the Purple Martin and White-bellied Swallow arrive first.

The first Barn Swallow was flying alone about the bridge at the Lake. Others were at the north end of the Park.

Sialia sialis. Singing.

Ceryle alcyon. - One Kingfisher, a female, was at the Pool, where, as in all the waters, the Goldfishes (*Carassius auratus*) were swarming, and of all sizes.

There were plenty of Rabbits (*Lepus sylvaticus*) and Gray Squirrels (*Sciurus carolinensis*). Although the day was very warm and the sun very bright, a Carolina Bat (*Vespertilio fuscus*, Beauv. [= *carolinensis*]) was flying about over the lower Reservoir for some time.

1881.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

April 28th. New York City.

Immediately after breakfasting, Rob. Morris and I started for Central Park, but decided to go to the end of the Elevated and have a good walk on the upper part of the Island. We got off at 155th Street, and walked across to Trinity Cemetery. As we entered the Chippies (*Spizella socialis*) were seen on every hand, and were trilling their simple ditty everywhere. A few Robins were singing; and Red Squirrels (*Sciurus hudsonicus*) were heard and seen in all parts of the churchyard on both sides of the boulevard. We heard the first one gnawing at a nut, and had some trouble in seeing it; but, at length Morris discovered it in a thick hemlock. Another was seen immediately afterwards on a low branch, and a third ran across the road. The Song Sparrow (*Melospiza fasciata*) was singing, and Yellow-birds (*Astragalinus tristis*) were there also.

In Audubon Park we saw the first House Wren (*Troglodytes domesticus*) among some leaves and rubbish beside the road. There were two (♂ & ♀) Cow-birds (*Molothrus ater*) near the same spot. The male sat in a tree uttering its characteristic notes and swelling himself up as he did so.

Further up along the Hudson we heard the Song of the American Redstart (*Setophaga ruticilla*) also the first of the season. We soon came up with him, a fine male, and watched him catching countless insects that were swarming in the warm sunshine among the branches of a hickory tree. We also saw a little further on I found a Sparrow Hawk was found in Audubon Park.

1881.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

small brook, and put our pet spotted Juvenile (*Nannus guttata*) into it. I brought it down from Highland Falls several weeks ago.

A Crow, which was seen beside the River was doubtless a Fish Crow (*Corvus ossifragus*) although it maintained perfect silence, its slight size, tail, and manners were proof to me that it was not *C. frugivorus*.

We saw a single Red-headed Woodpecker (*Melanerpes erythrocephalus*) that croaked and rattled amongst some tall oak trees on top of the hill. A Humming-bird (*Trochilus colubris*) and the Golden-winged Woodpecker (*Colaptes auratus*) completed the list of birds seen on the Hudson River side of the Island, with the exception of the Cardinal Grosbeak (*Cardinalis virginianus*) which whistled its beautiful notes clear and loud down in a ravine.

We crossed over to Harlem River above High Bridge, and found one Common Crow (*Corvus frugivorus*). Below High Bridge we saw one fine Brown Thrasher (*Harporhynchus rufus*), a pair of Bluebirds (*Sialia sialis*), and several Hermit Finches (*Turdus pallasi*) besides countless numbers of White-throated Sparrows (*Zonotrichia albicollis*) and a small flock of Eastern Snowbirds (*Junco hyemalis*) which have stayed rather late for them. Then we came to a large swamp as wild as any in the wilderness. In it we found, beside the ubiquitous White-throated Sparrows which whistled from every thicket, a pair of Cardinal Grosbeaks (*Cardinalis virginianus*). Everywhere the Hair-birds (*Spizella socialis*) were billing

1881.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

In the swamp were plenty of Skunk's Cabbage (*Symphlocarpus foetidus*) and American Hellebore (*Viratrum viride*). In Audubon Park, along the Hudson were large beds of Adder's Tongue and violets. Along the Harlem were Dutchman's Breeches (*Dicentra cucullaria*) and Anemones (*Anemone nemorosa*). There are large patches of Prickly Pear (*Opuntia vulgaris*), our only species ofactus.

We came upon a beautiful Wood Thrush (*Turdus mustelinus*) beside the Harlem, and saw a number of Chimney Swifts (*Chaetura pelagica*). There were three European Goldfinches near the 155th Street Station, on the Harlem marsh. Coming down on the "El" we saw the Purple Martin (*Progne subis*). Morris brought home a chrysalis of Prometheus Moth (*Attacus promethus*), which was attached to the Spice Bush (*Benjoin odoriferum*) which were covered with yellow blossoms. This moth almost invariably deposits the chrysalis to the Spicewood bush. Crossing from the Hudson to Harlem we found a pretty flower, the Blueets (*Houstonia coerulesa*) growing in the meadows.

May 1st.

131 W. 23d St., N.Y. City.

This morning, while sitting in nursery, we saw Purple Martins flying over the housetops, and one Sparrow Hawk flew overhead.

This afternoon I visited Central Park, entering it at 104th St. About the Pool were an abundance of Yellow-rumps (*Dendroica coronata*), Yellow Warblers (*Dendroica aestiva*), Chippies (*Spizella socialis*), Catbirds (*Mimus carolinensis*), and Song Sparrows (*Melospiza fasciata*), all heard singing, as

1881.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

as soon as I entered the Park.

The following birds were seen: —

Turdus migratorius. — Some sitting on their eggs; everywhere numerous.

Turdus fuscescens. — First seen. Abundant about Harlem Lake.

Turdus mustelinus. — All over in north end of Park.

Harporhynchus rufus. — Quite numerous.

Mimus carolinensis. — First seen. Abundant all over.

Certhia familiaris. — One seen beside the Pool.

Dendroica aestiva. — First seen.

Very abundant in all parts of the Park, especially beside water; singing loudly; utters a smart chirp.

Dendroica coronata. — Everywhere abundant, singing, and flycatching.

Dendroica discolor. — One alighted close beside me below Harlem Lake, and I watched it for a long time. At first I thought it was a Cape May Warbler, but it dropped down upon the ground where I had a good view of it. It caught insects very rapidly in the trees, then came down to terra firma, and went down beside the water (Harlem Lake) to make friends with a Maryland Yellow-throat, and hopped out on a rock in the lake along with its new acquaintance; then it flew back into a tree.

Dendroica pennsylvanica. — Several seen in the woods near the Loch. There were no Yellow Red-polls.

Merula varia. — Quite numerous. First seen.

Setophaga ruticilla. — Several were seen.

Parula americana. — First seen.

Only one Blue Yellow-backed Warbler

1881

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

was seen, and that beside the Pool. It was in a bush beside the water; and I got within a few feet of it.

Sciurus harrisi. — One — the first seen.

It was on the ground near the Loch.

Sciurus naevius. — First seen.

One on the brook between the Pool and Loch, and several about Harlem Lake.

Geothlypis trichas. — Several males about the margin of Harlem Lake.

Harundo horreorum [erythrogaster]. — There were a number seen, about a pool in a vacant lot, from the Elevated; and there were a number at the Pool and Harlem Lake.

Vireo flavifrons. — First seen. At the Loch.

Vireo solitarius. — First seen.

Everywhere quite plentiful.

Carpodacus purpureus. — As I approached the Wood Duck's pond, a bird was heard singing high up in the air as it descended; it kept flying up in the air over the trees and singing loudly until I reached the spot. It was a female Purple Finch. I stood close to it, and wondered that a female should sing so loud and sweetly.

Melospiza fasciata. — One Song Sparrow beside the Pool is a remarkable singer.

Melospiza palustris. — Quite numerous beside all the waters.

Spizella socialis. — Numerous.

Zonotrichia albicollis. — Still plentiful.

Cardinalis virginianus. — Seen beside Harlem Lake.

Pipilo erythrophthalmus. — Quite numerous about the Loch & Harlem Lake.

Passer domesticus. —

1881.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

Empidonax minimus. — First seen.

Abundant about the Pool, Loch & Harlem Lake.

Luscinia purpurea. — Lots of 'em.

Pandion haliaetus carolinensis. — A beautiful Fish Hawk was soaring over the Park.

May 4th. 131 W. 23, N. Y. City.

This morning, Morris, having got through with his last examination yesterday, spent the morning in Central Park, where he found a large number of Warblers and other birds. Among the warblers were Black-throated Blues (Dendroica caerulescens) in abundance, and some Black and Yellow Warblers (Dendroica maculosa). There was one Baltimore Oriole (Icterus galbula), and a Kingbird (Tyrannus carolinensis).

I passed my examination in Materia Medica in the morning, and Practice of Medicine in the afternoon. I finished the latter at half-past four o'clock, and, on getting to my room, found Dr. Clinton L. Bagg, with whom I went up to 104th St., on the "L", and walked around the Pool, Loch, and Harlem Lake.

The following birds were seen there:—

Icterus migratorius. — Singing. One sitting on its nest.

Turdus mustelinus. — Abundant.

Turdus fuscescens.

Minimus carolinensis. — Singing very sweetly.

Harporhynchus rufus. — Plentiful.

Regulus calendula. — One seen beside the Pool.

Minotilta varia. — Quite abundant.

Dendroica aestiva. — Everywhere singing; in numbers; utters a sharp, loud "chee".

Dendroica coronata. — Abundant; singing.

1881.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	<i>Parula americana</i> .					Singing.
	<i>Geothlypis trichas</i> .					Singing.
	<i>Spinus naevius</i> .		About the Pool.			
	<i>Setophaga ruticilla</i> .					Abundant; singing.
	<i>Dendroica pennsylvanica</i> .					
	<i>Dendroica caerulescens</i> .		We saw one by the brook.			
	<i>Hirundo horreorum</i> (= <i>erythrogaster</i>).					
	<i>Vireo flavifrons</i> .					
	<i>Vireo gilvus</i> .		We only saw one, at the Pool.			
	<i>Vireo norboracensis</i> .		Morris saw one in the morning.			
	<i>Carduelis elegans</i> .		Singing sweetly about the Pool.			
	<i>Astragalinus tristis</i> .					
	<i>Melospiza palustris</i> .		Abundant.			
	<i>Melospiza fasciata</i> .		Singing.			
	<i>Spizella socialis</i> .		Singing.			
	<i>Zonotrichia albicollis</i> .					
	<i>Passer domesticus</i> .					
	<i>Molothrus ater</i> .					
	<i>Quiscalus purpureus</i> .		At the pool.			
	<i>Empidonax minimus</i> .		Plentiful.			
	<i>Tyrannus carolinensis</i> .		Several in the upper part of the Park.			
	<i>Chaetura pelagica</i> .		Abundant.			

May 6th. Highland Falls, N.Y.
 Left New York City by Steamboat "Thomas Cornell", on time at 1 o'clock P.M., for Highland Falls. No birds besides Crows and Kingfishers were seen on the trip. We reached Cozzens at 4 o'clock. In the evening, I walked about the ponds and woods, and noticed the following birds:
Turdus migratorius. - Building & Singing.

1881.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	<i>Turdus mustelinus</i> .					Singing.
	<i>Harporhynchus rufus</i> .					Singing.
	<i>Mimus carolinensis</i> .					Singing.
	<i>Sialia sialis</i> .		Nesting in the birdhouse.			
	<i>Troglodytes domesticus</i> .		Singing about the house.			
	<i>Setophaga ruticilla</i> .					Singing.
	<i>Dendroica coronata</i> .		Very abundant.			
	<i>Spinus motacilla</i> .		About Echo Lake and Pell's L.			
	<i>Spinus aurocapillus</i> .		Singing.			
	<i>Hirundo horreorum</i> (= <i>erythrogaster</i>).					
	<i>Ampelis cedrorum</i> .					
	<i>Carpodacus purpureus</i> .					
	<i>Astragalinus tristis</i> .					
	<i>Melospiza fasciata</i> .					
	<i>Spizella socialis</i> .		Very abundant. Singing.			
	<i>Spizella pusilla</i> .		" "			
	<i>Zonotrichia albicollis</i> .		A few.			
	<i>Pipilo erythrophthalmus</i> .		Numerous.			
	<i>Agelaius phoeniceus</i> .		Singing in the old orchard.			
	<i>Icterus galbula</i> .		Mother saw the first, after I got home, this afternoon.			
	<i>Corvus frugivorus</i> .					
	<i>Sayornis fuscus</i> .		Breeding under the eaves, on the water-pipe.			
	<i>Empidonax minimus</i> .		Abundant.			
	<i>Trochilus colubris</i> .		Two seen. One in the back lot, and one near the house.			
	<i>Ceryle alcyon</i> .		One at Green Point, seen from the Steamboat.			
	<i>Colaptes auratus</i> .		Calling in Roe's woods.			
	<i>Chordeiles virginianus</i> .		First heard.			
	<i>Caprimulgus vociferus</i> .		First heard.			
	<i>Chaetura pelagica</i> .		Abundant.			
	The foliage is several days more advanced in New York City than it is here; and the birds are about as much behind as the vegetation is.					

1881.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

May 8th Highland Falls, N.Y.
 The following birds were seen to-day:-
Turdus migratorius.
Turdus mustelinus.
Turdus fuscescens. - First seen here.
Mimus carolinensis.
Harporhynchus rufus.
Sialia sialis.
Troglodytes domesticus.
Miniotilta varia. - Quite plentiful; singing.
Parula americana. - " " "
Helminthophaga ruficapilla. - Quite a
 number were seen in in the apple
 trees about the house.
Helminthophaga peregrina. - One was
 seen in the same plum-tree where I
 shot the partial albino of this species.
Dendroica aestiva. - Only one was seen,
 and that near a pool beside the Hudson.
Dendroica coronata. - Exceedingly common.
Dendroica pennsylvanica. - Only two seen.
Siurus auricapillus. - Common everywhere.
Siurus motacilla. - Wherever fresh water
 is found, there the Large-Billed
 Cuckoo's song may be heard, loud
 and clear, all through the heat of the
 day. Quite abundant.
Geothlypis trichas. - Only a few were
 seen, and those were all in Consook Marsh.
Icteria virens. - First seen.
Setophaga ruticilla. - Singing everywhere.
Pyrranga rubra. - Just come. First seen.
Hirundo horreorum.
Amphisp. cedrorum. - Small flocks
 are quite common. Their breasts
 are much paler than in autumn.
Carpodacus purpureus. - Singing.

1881.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

Astragalinus tristis. - Scattering flocks.
Melospiza fasciata.
Spizella socialis.
Spizella pusilla.
Zonotrichia albicollis. - Only two seen.
Casser domesticus.
Zamelodia ludoviciana. First seen.
Pipilo erythrophthalmus.
Molothrus ater. - Scattered flock.
Agelaius phoeniceus. - About the house + marsh.
Icterus galbula. - Abundant to-day.
Corvus frugivorus.
Cyanurus cristatus. - Two flocks of Blue
 Jays were seen passing northward.
 On my way up from the river, I threw
 a stone into a thick cedar tree, thinking
 that a nest in it might be newly built,
 for the purpose of starting off the old bird.
 A bird (Blue Jay) did fly out of the
 tree close to the nest. I climbed up,
 shaking the tree and making a good
 deal of noise, as the tree was a very
 compact, dense one. I saw a jays
 bill and tail sticking over the edge of
 the nest, and could have easily captured
 the old bird had I not supposed, from
 her position on the nest, that she was dead.
 She flew away, and did not return. I
 brought the nest and its set of 5 eggs home.
Tyrannus carolinensis. - One pair was
 seen beside Echo Lake; very quiet.
Sayornis fuscus.
Empidonax minimus. - Abundant.
Chordeiles virginianus.
Caprimulgus vociferus.
Ceryle alcyon. - Several seen.
Trochilus colubris. - Quite numerous.

1881.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.*Chaetura pelagica*. - Abundant.*Picus pubescens*.*Colaptes auratus*. - Calling.*Zenaidura carolinensis*.*Fringoides macularius*.

I walked all over Cousook Marsh without seeing any Savanna Sparrows, Swamp Sparrows, or Marsh Wrens.

The Shad-fishermen are on the island.

They told me that both kinds of Sturgeons, fork-tailed Catfish, and Stripped Bass were the only fishes they caught, besides Shad, in their nets.

The Shad blossoms were out, on Cousook Island. While there, I took a bath in the Hudson, the first of the season, and such a cold one that I was glad to get out of the river and into my clothes. The wild columbines are just beginning to come out nicely, and violets are in their prime. I saw, besides the above species, a fine White-headed Eagle (*Haliaeetus leucocephalus*). It was sailing circularly, and descended to the bank of the river.

May 9th. Highland Falls, N.Y.

Turdus migratorius.*Turdus mustelinus*.*Mimus carolinensis*.

Harporhynchus rufus. - This evening I flushed several out of the bushes by the roadside about half-past eight o'clock. They flew off chattering at a great rate.

Sialia sialis. - Have young ones.*Parus atricapillus*. - Two pairs.*Troglodytes domesticus*.*Mniotilta varia*. - Very abundant; singing.

1881.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

Parula americana. - Very abundant; singing.
Dendroica caerulescens. - Abundant. First seen here. Singing.

Dendroica coronata. - Abundant; singing.*Dendroica striata*. - First seen. Only one.*Dendroica pennsylvanica*. - Singing loudly everywhere. Very common.*Dendroica maculosa*. - First seen. Only one.*Sciurus auricapillus*. - Singing and abundant everywhere.*Sciurus naevius*. - First seen. Only one.*Sciurus motacilla*. - Singing wherever there is water. I found a nest within a

few feet of the spot where I found the last one containing young ones. I left

all along under the bank beside the nest without discovering it; at last

as I was leaving the spot - a small brook emptying into a large one a few

rods from the nest - I saw the old bird sitting upon its eggs, which

as near as I could see without disturbing them, were five in number.

I visited the nest just at dusk to-morrow. The old bird left the nest just as I

was putting my hand over it, and ran down the bed of the stream,

uttering a sharp chirping note. Then it came fluttering right into my face,

and passed over my head. It then flew down the brook to a witch hazel branch

where I saw indistinctly its white breast undulating as it bowed up

and down in the moonlight.

Geothlypis trichas. - Has now got up on high ground away from the marshes.

1881.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	<i>Icteria virens</i> .		Chattering everywhere			
	<i>Myiodioctes mitratus</i> .		First seen.			
	Quite abundant, and singing very delightfully along the brooks.					
	<i>Setophaga ruticilla</i> .		Very abundant, and singing quite energetically.			
	<i>Myiodioctes canadensis</i> .		First seen. Abundant. Singing.			
	<i>Tyranga nua</i> .		not yet common.			
	<i>Ampelis cedrorum</i> .					
	<i>Vireo olivaceus</i> .		First seen here			
	<i>Vireo gilvus</i> .		First seen here			
	<i>Vireo flavifrons</i> .		First seen here, and chattering ridiculously.			
	<i>Vireo solitarius</i> .		First seen.			
	<i>Vireo noveboracensis</i> .		First seen.			
	<i>Carpodacus purpureus</i> .		Singing.			
	<i>Astragalinus tristis</i> .					
	<i>Melospiza fasciata</i> .					
	<i>Spizella socialis</i> .					
	<i>Spizella pusilla</i> .					
	<i>Zonotrichia albicollis</i> .					
	<i>Zamelodia ludoviciana</i> .		One seen only.			
	<i>Passer domesticus</i> .					
	<i>Pipilo erythrophthalmus</i> .					
	<i>Molothrus ater</i> .					
	<i>Icterus galbula</i> .		Very abundant. Females.			
	<i>Corvus frugivorus</i> .		Shot one wild old chap.			
	<i>Cyanurus cristatus</i> .					
	<i>Tyrannus carolinensis</i> .		Only one seen, and that one was quiet.			
	<i>Sayornis fuscus</i> .					
	<i>Eupidonax minimus</i> .					
	<i>Chordeiles virginianus</i> .		Flying at dusk.			
	<i>Caprimulgus vociferus</i> .					
	<i>Chaetura pelagica</i> .					
	<i>Oreochilus colubris</i> .					
	<i>Ceryle alcyon</i> .					
	<i>Coccyzus americanus</i> .		First seen.			

1881

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

Coccyzus erythrophthalmus. - First seen.
Picus pubescens.
Colaptes auratus.

I believe these may fifty-four species, several in advance of the number seen yesterday, although a number of the species noticed yesterday were not observed to-day.

I received a notice this evening that my examinations had been satisfactory to the Faculty of the College of Physicians and Surgeons. Then I went to the Water Thrush's nest, and noticed besides that the migrating Brown Thrashers sleep at night in bushes, as well as (or instead of!) on the ground. The Whippoorwills were calling everywhere.

2,180.	<i>Coccyzus americanus</i> .	11.75	16.75	♀ ad.
2,181.	<i>Zamelodia ludoviciana</i> .	8.00	13.35	♂ ad.
2,182.	<i>Myiodioctes canadensis</i> .	5.65	8.15	♂ ad.

May 10th.

Was busy about the place all day, and the only new arrival noticed was *Icterus spurius* (Orchard Oriole), the first seen. This evening I tried to catch the Water Thrush in a net; but it has grown shy, and fluttered off the nest, and swam down the brook very rapidly. This one is the only bird we have that deliberately takes to the water, resembling in this respect the Dippers (*Cinclus mexicanus*).

May 11th.

Turdus swainsoni. - First seen. Quite numerous.
Helminthophaga vermivorus. - First seen.
Helminthophaga chrysoptera. - First seen.
Dendroica pennsylvanica. - One that I shot was in the plumage that I described in the

1881.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	Bulletin of the Nuttall Ornithological Club, viz.: without the black and chestnut markings.					
	<i>Dendroica caerulescens</i> . - Now very abundant.					
	<i>Dendroica maculosa</i> . - Quite plentiful.					
	<i>Geothlypis trichas</i> . - Now abundant.					
	<i>Myiodiodes nigratus</i> . - Abundant, and singing.					
	<i>Myiodiodes Canadensis</i> . - Very abundant. Singing.					
	<i>Icteria virens</i> . - Abundant.					
	<i>Siurus motacilla</i> . - I shot the female as she flew from the nest, and took the nest with its five eggs, which were secured just in time for preservation, as embryos were considerably advanced. The old bird had grown quite shy. She left the nest by running and swimming down the brook, as usual.					
	<i>Siurus naevius</i> . - Only one seen.					
	<i>Melospiza palustris</i> . - One seen near the stable. The first seen here this season. There were none on Cansack Marsh.					
	<i>Junco hyemalis</i> . - A fine old male was seen in the path before the house. A late one!					
	<i>Corvus</i> ?. - A flock of Crows were soaring circularly overhead, and croaking occasionally. I am nearly certain that some of them, at least, were Fish Crows (<i>Corvus ossifragus</i>).					
	<i>Ardea canadensis virescens</i> . - A fine Green Heron was seen at Morgan's brook. The first seen this year.					
2,183.	<i>Siurus motacilla</i> .			6.20	10.20	♀ ad.
	Wing, 3.12; tail, 2.00; bill from nostril, .40; gape, .75; tarsus, .90; middle toe and claw, .73; middle toe alone, .56; its claw, .18.					
	Iris, dark hazel. Maxilla and tip of mandible, blackish; basal one-half of mandible, flesh-color. Legs and feet, flesh-color. Taken with					

1881.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
2,184.	nest and set of 5 eggs (No. 163 ² / ₅).			5.10	8.20	♂ ad.
2,185.	<i>Helminthophaga chrysoptera</i> .			5.50	8.65	♀ ad.
	<i>Helminthophaga vermicivorus</i> .					

May 13th, 1881. New York City.

This morning early, being unable to sleep, I rose and breakfasted, and then started for the upper part of Central Park, taking the Elevated Road to 104th Street as usual. The foliage was well out, and the air was filled with fragrance, and sounds of singing birds. I reached the margin of the Pool, and soon found myself, or rather lost myself in the middle of a thicket, trying to identify a new bird's note. The voice that puzzled me was a sharp and metallic, and its owner proved to be the Kentucky Warbler (*Oporornis formosus*), and a bird never yet seen as far north as the Hudson Highlands. There were two of them; and their sharp call-note differed in pitch and quality from anything else that I have heard. They were shy and not disposed to court the society of naturalists. I soon lost sight of them; but, before I could extricate myself from the tangle into which I had plunged, I had noted species of our family of American Warblers (*Minotiltidae*). There were Yellow-crowned Warblers (*Dendroica coronata*), in full spring dress, accompanied by females in plainer garb, singing a song that, if not remarkable of itself, helps the general chorus. They were hovering over the water, fluttering among the terminal branches above it, or flying out in circles after insects

1881.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

Then there were Maryland Yellow-Throats (*Geothlypis trichas*), Canadian Flycatching Warblers (*Myiodiodes canadensis*), Small-billed Accentors or Water Thrushes (*Siurus naevius*), Summer Yellowbirds (*Dendroica aestiva*), Black and White Creeping Warblers (*Mniotilta varia*), Blue Yellow-backed Warblers (*Parula americana*), and some Black-poll Warblers (*Dendroica striata*).

The Wood Thrushes song heard; and so were the Catbird and Brown Thrasher singing. On a tree beside the Pool a pair of European Goldfinches (*Carduelis elegans*) were singing a sprightly song resembling our Yellowbirds.

As I walked down beside the brook, and the Loch, other Warblers, as the Chestnut-sided (*Dendroica pennsylvanica*), and the Black-throated Blue (*Dendroica caerulescens*), were seen; and the loud song of the Golden-crowned Accentor (*Siurus auricapillus*) was heard on the hillsides above the ravine. A beautiful Alice's Thrush (*Turdus Swainsoni glacie*) was found at the head of Harlem Lake.

There all the birds seemed inspired to produce their finest melody. A pair of Cardinal Grosbeaks were answering each other back and forth across the ravine; and four of the Vireos, viz: the Red-eyed, White-eyed, Warbling, and Yellow-throated (*C. V. V. Vireo olivaceus*, *noveboracensis*, *silvus* and *flavifrons*) were filling the woods with melody. Even the Redstart, emulating the example of better songsters was making music that excited my admiration. The east side of Harlem Lake is a favorite place for birds to

1881.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

congregate; but my time was too short for a thorough exploration. The birds whose presence was evident without making search were Robins, Catbirds and Towhees. The Chippies were trilling on every bush, and a few Song Sparrows were singing; but there was none among them all whose song could match that of the one whose nest is built near eighth avenue, beside the Pool. Small companies of Cow-birds (*Molothrus ater*) were ~~excursing~~ about, and there were some Crow Blackbirds (*Quiscalus purpureus*). A few Kingbirds (*Tyrannus carolinensis*) were seen; and the Least Flycatchers were hunting in pairs at the Loch. As I passed the Pool, on my return, a pair of Large-billed Accentors (*Siurus motacilla*) were noticed on the brink of the Pool. *Chaetura*. I reached the train in time to get down and dress in time for the rehearsal for our College Commencement exercises. In the evening I was decorated with one of the Harsen Prizes (the third and smallest one) for the best report in writing on the medical and surgical clinics held at the New York Hospital during the past winter. I received a Harsen diploma, in addition to one from the College, and a bronze medal, all of which tickled me immensely. While Barton & myself were going home from the Reception at Prof. Markov's house in 30th Street, we heard the Night hawks crying overhead, although it was a dark night and after twelve o'clock.

1881.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

Sunday, May 15th. Highland Falls, N.Y.
Took a long drive with Dr. Chace out to the
Forest o' Dea mines and then to Fort Montgomery
and visited a number of patients.
Near the mines we saw a Robin's nest
built under a bank beside road, whence
dirt had been carted to make the road.
The nest was built under a root, in
such a spot as the Large-billed Acceptor
(*Siurus motacilla*) would build in.
We saw one White-crowned Sparrow
(*Zonotrichia leucophrys*) on a fence
near the mines.

May 17th. H. & M.

Helminthophaga chrysoptera. - One fine male.*Dendroica maculosa*. - Still scarce.*Myiodiodes canadensis*. - Abundant.*Cyanospiza cyanea*. - First seen.

L. 186. *Myiodiodes canadensis*. 6.78 8.35 ♂ ad.
Wing, 2.55; tail, 2.20; bill from nostril, .28;
tarsus, .77.

Note.

On May 12th large numbers of Bank
Swallows (*Cotyle riparia*) were seen flying
about the sandbanks north of Peekskill.

On May 14th a bird was seen swimming
near Jona Island which I suppose to have
a Carolina Grebe (*Podilymbus podiceps*).

On the morning of May 12th, a large
flock of Blue Jays (*Cyanocitta cristata*)
were seen to alight on the bank of the River.
They were evidently migrating; and why
should so many flocks be moving
northward, when those about here
are incubating or have young ones in the nest.

1881.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

May 19th. H. & M.

Turdus swainsoni. - Plentiful*Turdus swainsoni aliciae*. - Quite numerous.*Mniotilta varia*. - Now becoming less numerous.*Parula americana*. - Singing.

Helminthophaga chrysoptera. - Several. They
rarely sing in the bushes, but fly to a tree top,
and remain pretty quiet, so that it is difficult
to discover one, even when it is singing.

Dendroica virens. - Has not been seen yet
this year. Neither has the Blackburnian
(*Dendroica blackburniae*).

Dendroica coerulescens. - Not very numerous.*Dendroica coronata*. - Not very plentiful.*Dendroica pennsylvanicus*. - Mated. Singing.*Dendroica maculosa*. - Rather abundant.*Dendroica striata*. - Quite numerous. Singing.*Dendroica castanea*. - First seen. Abundant. Singing.*Siurus auricapillus*. - Abundant. Mated. Singing.*Siurus motacilla*. - Common. Singing.

Siurus naevius. - I heard the lovely song of
the Small-billed Acceptor to-day for the first
time. It bears little, if any, resemblance to
the song of *Siurus motacilla*, except in its
rich mellow quality. It is superb.

Geothlypis trichas. - Sharp metallic click, a
rattling cry & a sweet song are its notes.

Myiodiodes canadensis. - Still plentiful.*Myiodiodes nitratu*. - Common. Singing.*Setophaga ruticilla*.

Icteria virens. - Less numerous. Settled down
to family matters in swampy thickets.

Pyrranga rubra. - Not abundant.

Chrysomitris (= *Astragalinus*) *tristis*. - Frequent,
as usual at this season, alder-grown swampy
pastures. Full plumage. Singing vivaciously,
and constantly roving about.

1881.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.*Zonotrichia albicollis*. - A few.

Empidonax minimus. - At dusk last evening (7:45 P.M.) a Least Flycatcher flew high up in the air uttering a loud note rapidly repeated, until the little fellow got away up in the air; then he came down uttering his spasmodic two syllabled note in such rapid succession and so fervently as really to produce melody. He evidently thought that he was performing a rapturous love-song.

Chordeiles popetue (= *virginianus*). Heard at nightfall.

Ardea virescens. - At Morgan's brook.

Tingoides macularius. - Heard at dark, flying over some high field near the River.

2,187. *Helminthophaga chrysoptera* 5.10 7.85 ♂ ad.
Wing, 2.48; tail, 1.95; bill from nostril, .35;
culmen, .53; gape, .60; tarsus, .67; middle
toe and its claw, .56; claw alone, .19.

2,188. *Dendroica castanea*. 6.00 9.50 ♂ ad.
Wing, 3.00; tail, 2.15; bill from nostril, .34;
culmen, .52; gape, .61; tarsus, .75; middle
toe and claw, .62; claw alone, .20.

2,189. *Dendroica castanea*. 5.65 8.95 ♀ ad.
Wing, 2.87; tail, 2.12; bill from nostril, .30;
culmen, .47; gape, .59; tarsus, .75; middle
toe and claw, .60; claw alone, .18.

2,190. *Dendroica maculosa*. 5.25 7.85 ♂ ad.
Wing, 2.50; tail, 1.95; bill from nostril, .29;
gape, .53; culmen, .45; tarsus, .72; middle
toe and claw, .55; claw alone, .16.

2,191. *Pyrranga rubra*. — — — ♂ ad.

2,192. *Seiurus naevius*. 6.10 9.62 ♂ ad.

Measurements of *Dendroica maculosa* 5.12 7.80 ♂ ad.
Wing, 2.40; tail, 1.90; bill from nostril, .30;
culmen, .45; gape, .54; tarsus, .70; middle
toe and claw, .55; claw alone, .17.

1881.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

May 21st. H. S. M.

Drove to Moses Clark's with Dr. Chace. Near his house *Helminthophaga chrysoptera* was heard; and an Olive-sided Flycatcher (*Contopus borealis*) was sitting on a fence-post with a big bumblebee in its beak. It was the first seen this season.

June 3d. H. S. M.

To-day went to Newburgh by train. At the mouth of Indian Falls Brook, an White-headed Eagle (*Haliaeetus leucocephalus*) descended and seized some object which was unable to rise with. It was in full view of the train for a distance of half a mile, during which time it kept constantly flapping and trying to rise. It had doubtless grappled with a large fish — perhaps a Sucker, which was too heavy. Two other Eagles were sitting on the shore.

I noted the spot where I shot my first adult Eagle near D. & C. Junction.

At Newburgh, in the gun-store I saw a "Ring-tailed Eagle" (*Aquila chrysaetus*) which was said to have been shot on the Hudson below Newburgh. Its tail was very broadly banded, basally, with pure-white, with a broad blackish terminal band of blackish.

A pair of Great-crested Flycatchers have been about my house for several days, and will probably breed in the vicinity.

Helminthophaga chrysoptera is still here. A pair are evidently breeding below Knisley's house, where I have heard it singing several successive days.

1881.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

June 4th. N.E.M.Y.
Turdus migratorius. - To-day saw a young
one already on wing.

June 16th. N.E.M.Y.
Sitta carolinensis. - A pair were calling in
Cozzens's woods.
Nyctiardea grisea naevia. - One was found
in the edge of the woods, south of the house.

June 20th. N.E.M.Y.
To-day, I saw a black hen of the Spanish breed,
lay an egg. Another hen that was sitting
upon a nest beside the rock where the trumpet
vine grows, was seen to rush from her nest
in, seemingly, great terror, with loud outcrying.
I went to the nest to see the cause of her
alarm, thinking a snake or weasel must
have disturbed her; but found only a hen
engaged in parturition. She showed no fear,
and did not pay attention when I stroked her
feathers the wrong way with a stick, but
seemed to be absorbed in the business on hand.
The labor pains came on rapidly, and, after
several forcible efforts, the egg popped out,
and I was able to see distinctly, what
caused Prof. Pro. C. Dalton so much labor, viz:
that the egg came out small end first. The
egg was large, and considerably rounded
at both ends. It is preserved in my cabinet.

June 23d. N.E.M.Y.
To-day I found the nest of a Purple Finch
(*Carpodacus purpureus*) on a low limb
of an apple-tree, and saw the female
sitting on her eggs. In the afternoon I
loaded two shells, too heavily alas! and repaired

1881.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

to the nest. I drove the female off of the nest,
and made a screeching noise with my lips,
sucking air through them, and finally
succeeded in attracting the male bird; but
it was in the plumage of the female, so that
I was lead to believe that there were two
nests. After a prolonged effort to attract any
red plumaged males, and noticing the devotion
of the two supposed females to each other,
and their evident mutual interest in the
nest, I became convinced that one of them
must be the male parent. Both birds were
then shot, and one showed *prunus fasciae*
evidence of its maternity, while the other
bird, although in complete female dress, exhibited
an absence of the characteristics noted on exam-
ination of the female parent's abdomen. On
dissection it proved to be a male plumage. The
only external character it possessed which would
be a clue to its male sex, was a solitary yellowish-
red feather in the crown.

2,193. *Carpodacus purpureus*. 6:15 10:10 ♀ ad.
Wing, 3:10; tail, 2:35; culmen, .46; gape, .56;
bill from nostril, .34; tarsus, .65; middle toe
and claw, .71; toe alone, .54; claw alone, .20.
Iris, hazel. Bill brownish-black on maxilla and
tip of superior mandible; the basal two-thirds
of the lower mandible, horn-color. Legs and
feet, brown. Female parent of set

2,194. *Carpodacus purpureus*. 5:90 mutilated ♂
Wing, 3:00; tail, 2:35; culmen, .47; gape, .53;
bill from nostril, .35; tarsus, .67; middle
toe and claw, .73; claw alone, .22; toe
alone, .52. Iris, hazel, and other colors the
same as No. 2,193 above except that the
horn-color on the lower bill extends nearly

1881.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

to the tip of the bill, blending gradually into black at the tip. The above two specimens were the parents of the nest and set of eggs,

July 14th. H. S. N.Y.

Myiodiastes nigratus. - Near Forest of Dean mine, I saw a beautiful male in unusually fine plumage, and singing.
Sturnus motacilla. - Several along Popple Creek.
Harporhynchus rufus. - As Dr. Chase & I drove home from Forest of Dean at midday, the day being uncommonly hot, and the sun shining with burning heat upon the dusty road, a number of these birds were seen dusting themselves in the road, where a thermometer placed in the sand would doubtless rise to 115° or more.

Ardea virescens. - One feeding by J. P. Morgan's bridge.

July 15th. H. S. N.Y.

Turdus fuscescens. - Several young ones were seen in the raspberries near the mine.
Helminthophila vermicivora. - One near the mine.
Empidonax minimus. - One seen to fly to its nest. Second brood?

Harporhynchus rufus. - Dozens were seen dusting in the middle of the road in the broiling sunshine.

Pyranga rubra. - Two old males together.

Sturnus motacilla. - Singing on Popple Creek.

August 16th. H. S. N.Y.

To-day Mr. John Redner presented me with a tolerably well-skinned specimen of the American Egret (Herodias alba egretta) (Gmel.) Ridgway, which he shot on August 9th, 1881, at Popple Pond, Orange

1881.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

County, New York, from a flock of eight. They were first seen sitting upon the treetops near the dam, but when the specimen was shot they were feeding upon the mud flat, or shallow water along shore. The waded out till immersed in water to their bodies.

2,195.

~~Ardea~~ Herodias alba egretta (Gmel.) Ridgw.
Popple Pond, Orange Co., N.Y., Sept. 9th, 1881.

September 5th. Sing Sing, N.Y.

Louis Zerega and myself went to see Dr. A. K. Fisher, and spent the afternoon shooting about Sing Sing, where we saw four Ruffed Grouse (Bonasa umbellus), and numerous species of Hawks. In a field containing a number of very large oak-trees, we found the Olive-sided Flycatcher (Contopus borealis) very common, and shot a King bird (Tyrannus carolinensis). Dr. Fisher showed us the ground where Oporornis formosus and Empidonax acadicus breed so commonly every season, but there seems little woods of a similar character about Highland Falls, and I deem it very doubtful if more than an occasional straggler of either species ever occurs about here.

September 6th. H. S. N.Y.

Spent the day at Popple Pond, fishing, in company with Chief Eng'r. Wm. B. Brooks, U. S. N., and his two boys, and Mr. Louis A. Zerega. Louis shot a Woodcock in a cornfield, and I shot at one in an onion field near by. One Great Blue

1881.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

Heron was seen; and a Bittern had been shot at a day or two previously on the Creek. We saw numbers of Kingbirds (*Tyrannus carolinensis*). A fisherman on the Lake (Popple) told me that he had not seen "them White Cranes" in four years when the flock of eight appeared there this summer. He stated that a flock of four of these birds (*Aerodius alba egretta*) were about the pond four summers ago.

2,196. *Amphelis cedrorum*. 7.37 12.00 ♂ ad.
Popple Pond, Sept 6, 1881.

September 8th H.F.M.Y.

To-day rode to Queensborough to see patients. Tied my horse at the bridge, and began casting for trout (*Salmo fontinalis*) of which I saw several nice specimens. While fishing I heard the sharp, metallic note of the Hooded Warbler (*Myiodytes mitratus*), and soon had the pleasure of seeing a fine old male in beautiful plumage taking a bath. It bathed amongst the smooth stones where the water was shallow and kept a perfect fountain of spray going around it all the time.

Night Hawks (*Chordeiles popetue*) were flying in the twilight as I rode home; flew low. Hummingbirds (*Trochilus colubris*) were seen about the cardinal flowers beside the brook. *Tyrannus carolinensis*. - A few still remain.

September 10th H.F.M.Y.

Antrostomus vociferus. - Still common, and heard every night.

1881.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

Sept. 16th, H.F.M.Y.
Trochilus colubris. - Still present.
Empidonax minimus. - Still here.

September 28th H.F.M.Y.

This evening, while riding up from Fort. Montgomery, I heard the note of a Night Heron (*Nyctiardea grisea naevia*), and soon was able to distinguish the form of a solitary heron, flying as swiftly as possible toward the South, uttering its quack at infrequent intervals.

October 6th. Forest of Dean. N.Y.

2,197. *Sturnella magna*. 10.88 16.75 ♂ ad.
Wing, 4.88; tail, 3.30; culmen, 1.42; gape, 1.50; tarsus, 1.73; middle toe, 1.12; middle toe and its claw, 1.55. Iris, hazel.

October 7th. H.F.M.Y.

To day went shooting with Mr. Ridner. We shot a Surf Duck (*Pelecanus purpuricollis*) below Harrisons. It was skulking behind a bush along shore. An immature Eagle (*Haliaeetus leucocephalus*) was flying about Gouverneur's Cove. On Constitution Island the dog breed a Night Heron (*Nyctiardea grisea naevia*) and barked until we came up. Then it flew across the marsh, and I missed it with both barrels.

We saw several Grebes, but we couldn't kill any. A Rail, probably *Rallus virginianus* flew across the creek; we put the dog in the reeds after it, and once we saw it run through the rushes just in front of Dash's nose, but we could not get it to fly.

1881.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

I saw a Semipalmated Sandpiper (Ereunetes pusillus) flying, and watched it until it alighted. Then we rowed up and I shot it on the weeds, when another one arose beside it and I killed that too. I have never seen the species in the Highlands but once before (see entry p. 90).

I saw a male Black-throated Blue Warbler (Dendroica caerulescens) in full plumage. Bonasa Umbellus. - Several seen on Constitution Island, and one on shore.

A Spotted Sandpiper (Tringoides macularius) was seen sitting on a stone on shore, and I got ready to shoot it when it flew; but it had barely gone a ^{rod} yard out over the water when a Pigeon Hawk (Asio columbarius) darted swift as thought in hot pursuit. The terrified Sandpiper instantly dropped to the water and dived. The Hawk reached the spot just as the Snipe got under and my charge struck it just at the same moment; but the shot were No. 12s and the Hawk flew a little way back into the woods and alighted. The Sandpiper rose to the surface, and remained on the water several minutes before it could muster courage to fly away.

2,198. Ereunetes pusillus. 6.62 12.30 ♀ ad.
Wing, 3.75; tail, 1.70; culmen, .81; gape, .80; tarsus, .90; middle toe, .61; middle toe and its claw, .78. Iris, dark hazel.

2,199. Ereunetes pusillus. 6.38 12.25 ♂ ad.
Wing, 3.80; tail, 1.80; culmen, .71; gape, .70; tarsus, .88; middle toe, .63; middle toe and its claw, .80. Iris, dark hazel. Bill, black. Legs & feet, blackish lead-color [one with broken bill].

1881.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

2,200 Pelionetta perspicillata. 17.65 31.00 ♀
Wing, 8.75; tail, 3.05; culmen, 1.40; gape, 2.15; tarsus, 1.70; middle toe and its claw, 2.63; claw, .31.

October 8. H. J. M. Y.

Ardea herodias. - One seen flying over the lower Cove.

October 11th. H. J. M. Y.

Spizella monticola. - The first seen.
Junco hyemalis. - Seen Oct. 6th for the first.
Dendroica virens. - One seen.

October 14th. H. J. M. Y.

The Ruby-crowned Kinglets (Regulus calendula) are now present in full force. So are the White-throated Sparrows (Zonotrichia albicollis). For the past week the Yellow-bellied Woodpeckers (Sphyrapicus varius) has been extremely abundant. You can find them in the apple orchard any time, and also in the woods, where I shot an old male in an elm-tree with my pocket shot-gun.

No Warblers save Dendroica coronata was seen during a long walk; but the Hermit Thrushes (Turdus pallasi) is very abundant. The one recorded below was the first bird shot at with my little gun.

2,201. Turdus pallasi. 7.40 11.65 ♂ ad.
Wing, 3.65; tail, 2.85; bill from nostril, .38; gape, .80; tarsus, 1.20; middle toe and claw, .92; claw alone, .22.

2,202. Sphyrapicus varius. 8.60 15.50 ♂ ad.
Wing, 5.00; tail, 3.30; culmen, .90; bill measured front nostril, .76; gape, 1.05; tarsus, .76; middle (outer) toe and its claw, .80; claw alone, .34; toe without claw, .52. Iris, dark reddish hazel. Bill, dusky bluish-horn-color. Legs & feet, lead-color.

1881.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

October 15th. H. S. N.Y.

As I was fishing in Popple's Creek a beautiful little Blue Yellow-backed Warbler (*Parula americana*) flew into a willow bush and began searching the leaves for insects. It was only a few feet from me, and showed no fear. This is my latest date by one week (Oct. 8, 1876; 8, 1878).

One Catbird was making considerable noise in a thicket by the roadside, so I returned. The only date date given in my list is Oct. 16, 1874.

October 21st. N.Y. City.

To-day Louis and I took the elevated road up to 155th Street, and went through the swamp below High Bridge to see what birds were about, and to try to procure a Red-bird (*Cardinalis virginianus*), but found none, although

Song and White-throated Sparrows were very numerous in the thickets, and we thought we heard the note of a Cardinal Grosbeak. Robins were in great plenty, and were found everywhere feeding upon the berries of the dogwood (*Cornus*). Crows were common, and large flocks of Purple Grackles (*Quiscalus*) flew overhead. A few Catbirds were seen, and a number of Hermit Thrushes and Golden-winged Woodpeckers were feeding on the dogwoods.

October 22nd. N.Y. City.

To-day Louis and I had nothing to do, but enjoy ourselves. So we decided to go on a picnic. We called for the Messrs Davidson in 9th St., and succeeding in obtaining their society, and took the L road from there to 155th St.,

1881.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

We went through Trinity Cemetery, and thence, after visiting Audubon's tomb, strolled through Audubon Park, where I succeeded in bringing down a fine specimen of the Red-headed Woodpecker (*Melanerpes erythrocephalus*) with Louis' big horse pistol, which made a noise like a cannon, kicked furiously, and doubtless aroused the fears of all the inhabitants. We found White-throated Sparrows in plenty, and I shot one for the embellishment of Louis' cabinet. We lunched on the rocks beside the Hudson, just below the West End Hotel, and watched the boys fishing, and netting shrimps. Red-headed Woodpeckers were in great plenty, and we watched one storing up food in the crack of a tree. They were lively, noisy and full of life as ever. Brown Creepers, Robins, White-bellied Nuthatches, Yellow-rumped Warblers, Yellow-birds (*Astragalinus tristis*) and Yellow-shafted Flickers were also quite numerous; and one Swamp Sparrow was seen in the marsh beside the Hudson. Our picnic was a success, and we got home just in time for the 6:30 train which brought me home to the Mearns House.

2, 203. *Melanerpes erythrocephalus*. 9.75 17.75 ♂ juv. Wing, 5.25; tail, 3.25; culmen, 1.08; bill from nostril, .80; gape, 1.25; tarsus, .95; middle toe and its claw, 1.02; claw alone, .37; toe alone, .70; Iris, very dark brown. Bill, bluish horn-color, with ridge of maxilla, terminal half of maxilla and terminal third of mandible, blackish. Legs and feet, lead-color. Shot in Audubon Park, New York (Manhattan) Island, in company with Misses Louisa and Nancy Davidson and Mr. Louis A. Zerega, with an old horse pistol. In memoriam.

1881.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

Mr. Redner brought me a Meadow Lark some two weeks ago, and told me he had shot a Coot (Fulica Americana) on Popoplen's Creek or Pond.

October 25. H. F., N.Y. Popoplen's Pond

Mr. Redner brought me a Grebe
2,207. Podilymbus podiceps. 12.25 20.50 ♀
Length to end of toes, 15.60; wing, 4.87; tarsus, 1.33; mte., 1.90; int., 1.70; claw, .25; culmen, .68; bill from nostril, .46; gape, 1.00.

October 27th. H. F., N.Y.

Regulus satrapa and R. calendula are quite abundant. Nov. 2, 1874, is my latest autumn record for calendula.

Bonasa umbellus. - Quite scarce. One seen to-day.

2,204. Turdus migratorius. 10.00 15.90 ♀

Wing, 5.00; tail, 4.00; culmen, .79; bill from nostril, .52; gape, .79; tarsus, 1.40; middle toe and claw, 1.15; toe alone, .85; claw alone, .30.

Iris, dark hazel; bill, blackish, with extreme base of under mandible, yellow. Tarsi and toes, blackish, with soles of feet, yellow.

October 28th. H. F., N.Y.

Regulus calendula. - Quite a number still here.

Carpodacus purpureus. - In flocks; quite numerous.

Mr. Redner sent me a Greater Yellow-legs.

2,205. Totanus melanoleucus. 14.00 24.25 ♂ ad.

Wing, 7.25; tail, 2.95; culmen, 2.20; gape, 2.50; bare part of tibia, 1.67; tarsus, 2.40; middle toe and claw, 1.53; middle toe alone, 1.30; its claw, .24; outer toe and claw, 1.29; inner toe and claw, 1.25; hallux and its claw, .43; claw alone, .13. Iris, very dark hazel. Bill,

1881.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

bluish-black. Tarsi and toes, greenish yellow; claws, black. Shot at Popoplen's Pond, N.Y.

October 31st. H. F., N.Y.

Drove to Canterbury to-day. Birds seen were Turdus migratorius, Dendroica coronata, Turdus pallasi, Bonasa umbellus, all common besides other ordinary species.

November 1st. H. F., N.Y.

Regulus calendula. - Quite a number are flying about the house, uttering their Wren-like note, and once in a while singing.

November 2nd. H. F., N.Y.

Regulus calendula. - Quite a number were noted in the apple-trees about the house, and in the woods.

Passerella iliaca. - First seen. Quite a good-sized flock was found. They whurred through the brush like small partridges, uttered a loud tweet, and a loud clicking call, borrowed from the Brown Thrasher.

November 4th. H. F., N.Y.

Spent the afternoon in the woods. The following birds were seen: -

Turdus migratorius. - Quite numerous at various localities; upland they sought shelter from the cold wind in the thickets, and down by the river in the red cedars.

Turdus pallasi. - Still quite common.

Sialia sialis. - Collecting in large flocks, preparing to migrate, warbling and chattering constantly.

Regulus satrapa. - Scarce. R. calendula has not been seen since the 2nd.

1881.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	<i>Sitta carolinensis</i> .					Not abundant.
	<i>Dendroica coronata</i> .					Not as abundant as before.
	<i>Ampelis cedrorum</i> .					Large flocks.
	<i>Carpodacus purpureus</i> .					Common. Feeds on seeds of the tulip-tree.
	<i>Astragalinus tristis</i> .					Has lost its summer plumage. The time of change is the month of October.
	<i>Melospiza fasciata</i> .					Not migrating, and only a few permanent ones here.
	<i>Junco hyemalis</i> .					Very abundant.
	<i>Spizella montana</i> .					One seen on Cousook Island.
	<i>Spizella pusilla</i> .					Several seen.
	<i>Passerella iliaca</i> .					Numerous.
	<i>Zonotrichia albicollis</i> .					One large flock.
	<i>Corvus frugivorus</i> .					
	<i>Cyanocitta cristata</i> .					Common.
	<i>Picus pubescens</i> .					
	<i>Fulica americana</i> .					As I walked along Cousook Marsh, a Coot flew up from the edge of the bank and was shot; being wounded severely, it swam out on the mud flats, and cost me a wetting to get it, as I had no boat.
	<i>Bucephala clangula americana</i> .					Several swimming in the middle of the river.

- 2,206. *Fulica americana*. 13.80 24.10 ♀ ad.
 Wing, 7.00; tail, 2.30; culmen, 1.50; bill from nostril, .52; gape, 1.20; Tarsus, 2.00; bare part of tibia, .80; middle toe and claw, 2.98; toe alone, 2.55; claw alone, .45; outer toe and claw, 2.50; toe alone, 2.25; claw, .29; inner toe and claw, 2.40; toe alone, 2.07; claw, .40; hallux, 1.18; toe alone, .90; claw alone, .33. Iris, reddish brown. Bill, bluish flesh-color, with base of maxilla,

1881.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
2,208.	<i>Turdus migratorius</i> .	10.50		16.65		♂ ad.
				Wing, 5.25; tail, 4.15; culmen, .80; bill from nostril, .52; gape, 1.10; tarsus, 1.40; middle toe and claw, 1.14; toe, .89; claw, .31. Iris, hazel. Tarsi, toes & nails, black. Bill with upper surface of maxilla, and extreme tip of mandible, blackish; sides of maxilla, and all save tip of mandible, yellow; inside of mouth, yellow.		
2,209.	<i>Turdus migratorius</i> .	10.00		18.52		♀ ad.
				Wing, 4.87; tail, 3.80; culmen, .88; bill from nostril, .55; gape, 1.15; tarsus, 1.27; middle toe and its claw, 1.11; middle toe, .83; its claw, .29. Colors of parts same as above.		
2,210.	<i>Turdus migratorius</i> .	10.25		19.25		♂ ad.
				Wing, 5.00; tail, 4.00; culmen, .96; bill from nostril, .62; gape, 1.20; tarsus, 1.35; middle toe and its claw, 1.13; middle toe alone, .85; claw, .30. Colors of parts like No. 2,208.		
2,211.	<i>Carpodacus purpureus</i> .	6.37		10.50		♂ juv.
				A few red feathers on chin. Iris, hazel.		

November 5th. H.F., N.Y.

To-day was spent at Poplopen's Pond, fishing and gunning. We reached the pond just in time to witness a gorgeous sunrise, and a lot of migratory Crows (*Corvus frugivorus*). The first flock, we estimated, contained about 150 Crows, and straggling bands kept pouring over the top of a mountain in their wake, until I had counted 322 besides the first flock.

1881

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

(November 5th.)

H. J. M.

While fishing a Carolina Grebe (*Podilymbus podiceps*) swam down the lake towards us. Our boat appeared to excite greatly its curiosity, for, although apparently afraid of us, it kept gradually drawing nearer, by alternate shy advances and scared retreats, until at length it was within range and a fortunate shot killed it dead.

Towards evening Mr. John Redner came over, and asked me to go with him to the head of the lake and look up a pair of Greater Yellow-legs (*Totanus melanoleucus*) which had been staying there for a week or so, and which he had tried to shoot on the day previous. I had already walked half-way around the pond in search of Snipes, and had found only some fine plates of mica, near what is called Singglass Spring. I accepted the invitation, and soon shot a big yellow-legs as it sat beside a pool in the marsh. At the head of the lake we found three more, that flew wildly about from place to place, crying loudly. The place was a large tract (several acres) of partially submerged land, boggy, and mucky, the bottom being perfectly black. I took long shots at the flying birds, as Redner hunted them from place to place, and succeeded in killing two out of the three, at long range, with No. 4 shot.

Mr. Redner gave me a Great Northern Shrike (*Lanius borealis*) which he had shot early in the afternoon.

A Fish Hawk (*Pandion haliaetus carolinensis*) was seen about the

1881.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

pond all day.

2,212.

Podilymbus podiceps.

15:00 24:00 ♂ ad.

Wing, 4.95; tail tuft, 1.73; culmen, .86; bill from nostril, .31; gape, 1.37; tarsus, 1.66; middle toe and its claw, 2.25; middle toe alone, 2.00; claw, .29. Outer toe and claw, 2.40; inner toe & claw, 1.75; hallux and claw, .57; claw alone, .12.

Iris, reddish brown; eyelid encircled by a narrow yellow rim. Bill, with maxilla dusky, lightest on edges; mandible, yellow, dusky on edge and point. Tarsi and toes, bluish olive-green, with dusky shadings; outer aspect of tarsus and bottoms of feet, black; nails, blackish, edged with horn color.

2,213.

Totanus melanoleucus.

14:10 25:35 ♂ ad.

Wing, 7.75; tail, 3.12; length to end of toes, 17.40; culmen, 2.25; bill measured from nostril, 1.88; gape, 2.50; tarsus, 2.60; bare part of tibia, 1.65; middle toe and claw, 1.67; toe alone, 1.40; claw, .28; outer toe and claw, 1.43; inner toe and claw, 1.35; hallux and its claw, .44; claw alone, .15.

Its claws like No. 2,205. Its left tibia, tarsus, foot, and toes had been ^{previously} fractured & deformed.

2,214.

Totanus melanoleucus.

13:85 24:30 ♂ ad.

Wing, 7.50; tail, 3.10; length to end of toes, 17.75; culmen, 2.20; bill measured from nostril, 1.86; gape, 2.46; tarsus, 2.47; bare part of tibia, 1.67; middle toe & claw, 1.60; middle toe alone, 1.38; claw, .25; outer toe and claw, 1.43; inner toe and claw, 1.27; hallux and its claw, .40; claw alone, .12. Colors,

like No. 2,205. Had been previously wounded in both tibiae.

2,215.

Totanus melanoleucus.

14:50 24:75 ♂ ad.

Wing, 7.50; tail, 3.00; culmen, 2.15; gape, 2.43; tarsus, 2.62; bare part of tibia, 1.75; middle toe and claw, 1.70; middle toe alone, 1.45; claw, .28; outer toe and claw, 1.44; inner toe and claw, 1.33; hallux and its claw, .35; claw alone, .14.

over

1881.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

November 5th. H.E.M.Y.
Colors the same as No. 2, 205, only yellow on
lower surface of lower mandible towards base.
2, 216. Lanius borealis. 10.35 14.45 ♂ ad.
Wing, 4.50; tail, 4.50; culmen, .75; bill meas-
ured from nostril, .55; gape, 1.14; tarsus,
1.04; middle toe and its claw, .88; middle
toe alone, .60; claw, .29. Basal half of man-
dible & edge of maxilla, flesh-color; remainder of
bill, black. Tarsi & toes, black. Iris, hazel.

November 6th. H.E.M.Y.
To-day noticed, in the garden, a number
of Chipping and Field Sparrow (Spizella
domestica and S. pusilla).

November 8th. H.E.M.Y.
Spent the afternoon fishing at Poplophen's
Pond. Caught a large mass of Pickerel
& Yellow Perch; and netted a couple of
good-sized Sunfishes (Eupomotus aureus)
and 3 mullet suckers, and some very
large-sized, red-finned Shiners.

Went up to the head of the lake and
found there the one Greater Yellow-leg
(Totanus melanoleucus) that we left on the 5th.
It was shy and flew well out of the way
the first time it was started up, and,
being alone, I did not pursue it.
Spizella pusilla. — A number were seen.

November 9th. H.E.M.Y.
Luscalus purpureus. — One flying south.
Pandion haliaetus carolinensis. — The one
seen last Saturday (5th) was in the same place,
on a hemlock tree, beside Poplophen's
Pond.

1881.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

Being desirous to procure the last of the
flock of Greater Yellow-legs (Totanus
melanoleucus) at Poplophen's Pond, I drove
up there, and got Sam. Redner to get his
gun and help me hunt it. When first started
up the bird flew the whole length of the
pond to the north end, where I shot it
with a number four shot beneath the
wing, a side shot, passing clear through
and out on the other side. Notwithstanding
this mortal wound the snipe flew a long
distance, and alighted beside the pond,
where we found it and had no trouble
in taking it, although it was not dead.
2, 217. Totanus melanoleucus. 13.88 25.10 ♂ ad.
Wing, 7.50; tail, 3.20; Length from end of
bill to end of toes, 17.75; culmen, 2.05;
bill measured from nostril, 1.72; gape, 2.28;
tarsus, 2.55; bare part of tibia, 1.70; middle
toe & its claw, 1.70; middle toe alone, 1.43;
its claw, .25; outer toe and its claw, 1.45;
inner toe and its claw, 1.29; hallux and
its claw, .35; claw alone, .13.
Iris, dark hazel. Tarsi and toes, greenish
yellow; claws, black. This specimen, like
its four companions previously taken, was
a male, apparently adult. My interest with
regard to the lateness of the departure of this
little flock of five unlucky Snipes was exceeded
by the desire for their possession as specimens
but I think would not have been likely to
have remained there many days later.

November 12th. H.E.M.Y.
Last night I slept on the Island, at Poplophen's
Pond, beside large fire built of logs by
Mr. Denna and myself. We reached the pond

1881.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

in the evening (10 o'clock, P.M.), and walked way down the east side of the pond in search of the boat; not finding it we reversed our direction and walked entirely around the pond, to find our boat within a stone's throw of the spot we had reached when we first started out. We set a gill net, and caught during the night several "mulletts," a common sucker, a pickerel, and several perch.

We slept beside a large fire of logs during the rest of the night, and caught about 30 pickerel & some perch during the day with our lines, and some more yellow perch, mullets, and a couple of pickerel in the net. I preserved a "mullet" and sucker in alcohol.

Spizella socialis. - Three were seen near Poplopen's Pond; now extremely scarce. We did not see the Fish Hawk to-day. It rained some during the night and heavily all day.

November 15th. H.E.M.

To-day rode to Fort Montgomery, where I turned my horse into the woods, and came suddenly upon two Ruffed Grouse (Bonasa umbellus) in two small cedars, eating "many berries". Then I rode within a few feet of a Wild Pigeon (Ectopistes migratorius) that was sitting on the ground beneath the thickly growing hemlocks. It flew on to a branch. I told my horse and walking close to it, threw a stone at it, when it flew away.

I saw one Chipping Sparrow (Spizella domestica) in front of my house. It was in the plumage of a bird of the year. Bluebirds in large flocks, feeding on sumach berries.

1881.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

November 25th. H.F., N.Y.

Sialia sialis. - In large flocks, feeding on berries of the sumach (Rhus).

Fundus migratorius. - Heard about house every day.

Dendroica coronata. - Not seen since November 22nd. The ground has been covered with snow for two days.

November 30th. N.Y. City.

This afternoon Louis Zerega and myself were walking through Central Park together when I saw a Mocking-bird (Mimus polyglottus) sitting still upon a low branch of a tree, but was not quite certain about its identity, as it flew into a thicket. There, however, we got close to it, and sat down within a few paces of it and watched it eating some red berries resembling bitter sweet. I gathered some and afterwards showed them to Prof. J. M. Wheaton, M.D., and he told me the name of the plant. It is the Burning-Bush or Waahoo (Euonymus atropurpureus) [Gray's "Botany of the Northern United States," p. 81.]. The shrub or small tree is very handsome when laden with fruit in autumn. I have never seen it in the Hudson Highlands. This is the first wild Mocking-bird I have ever seen alive. The northern locality is interesting, although Mr. Eugene P. Bicknell has taken it a far off up the Hudson River as his residence at Riverdale, N.Y.

December 1st. Baltimore & Ohio Railroad.

Near Deer Park, I saw a Hawk sitting quietly upon a tree near the railroad, which I believe was the Rough-legged Hawk (Archibuteo lagopus sancti-johannis), in very light plumage over.

1881.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

December 1st. - Continued.

Near Oakland I saw the remains of a Bluebird (*Sialia sialis*) which one of the Shrikes (*Lanius*) had impaled upon a thorn tree.

Going west on the Baltimore and Ohio Railroad I first saw Red-Birds (*Cardinalis virginianus*) at Moundsville, W. Va.

December 2nd. Columbus to Circleville, Ohio.

From Columbus to Circleville, Ohio, on the Scioto Valley road, the following birds were observed:—

1. *Junco hyemalis*.
2. *Timonius spawerius*.—A number seen.
3. *Buteo* ———?—Two Buteos seen.
4. *Zenaidura carolinensis*.
5. *Sturnella magna*.—Common.
6. *Colaptes auratus*.—Common.

December Birds of Circleville, Ohio.

On December 5th, went out in the morning on the Scioto River, north, and along Darby Creek.

On December 6th, hunted on the Hulse farms; and on the 7th we drove out to the Flats and spent the day.

On December 10th, I walked up the Scioto, followed the Darby up into the woods, and then crossed over through the corn fields to the Four Mile Red Bridge; went a little farther up the river, and crossed over to the prairie beside the Canal, and walked down to Circleville, Ohio.

Altogether I went shooting only five times. On Dec. 12th, Mr. John Kruehler drove me

1881.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

out to Campbell's swampy prairie, where we spent part of the afternoon in the rain hunting unsuccessfully for Hawks & Owls. Then we drove to Hulse's to take tea, and shot a couple of Gray Squirrels.

The following is a list of the birds seen:—
1. *Sialia sialis*.—A good-sized flock was found in a stumpy field, on Hulse's farm at the Flats. None others were seen.

2. *Regulus satrapa*.—Quite numerous, and found in the largest timber in the deep woods.

3. *Sitta carolinensis*.—Quite abundant, and found in equal numbers in the shrubby thickets and in the deep woods of heavy timber. No Red-bellied Nuthatches were seen.

4. *Lophophanes bicolor*.—Everywhere abundant, noisy, pugnacious and restless.

5. *Parus atricapillus*.—Not abundant. I shot but one, in a flock of Tufted Titmice, on Dec. 6th.

6. *Parus atricapillus carolinensis*.—Quite abundant, and always associated with the Tufted Titmouse. It is found almost everywhere, but perhaps most often in the bushes, weeds and reeds along the rivers and creeks. The sharply-defined black patch of the throat, and its rougher notes serve to distinguish it from the preceding.

7. *Certhia familiaris*.—Not common.

8. *Thryothorus ludovicianus*.—One was heard singing a few sweet notes, on "Aunt Hepsies" farm, on Dec. 6th. We afterwards saw it on the lower rails of a Virginia rail-fence edged with brushwood, but could not capture it.

Another one was heard about a tumble down old house on the Home Farm, at Hulse's; but before I could get to it, a Red-bellied Woodpecker (*Centurus carolinus*)

1881.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	engaged my attention, and I never got back to the spot, after catching my wounded Woodpecker.					
	9. <u>Anorthura troglodytes hyemalis</u> .— Only one seen, on Dec. 10th, amongst some heavy logs and driftwood beside the Scioto River, above the Four Mile Red Bridge.					
	10. <u>Eremophila alpestris</u> .— One good-sized flock of Shore Larks was seen on the 7th, as we were driving out to the flats. They were flying and driving wildly about like a flock of Snow Buntings. The wind was blowing hard, and they flew high most of the time.					
	11. <u>Lanius ludovicianus</u> .— I did not see the Loggerhead Shrike this year, but several have been seen by different persons since I came to Circleville. I saw one here last winter (1880).					
	12. <u>Astragalinus tristis</u> .— The "wild canaries" were occasionally seen or heard.					
	13. <u>Melospiza fasciata</u> .— Abundant every where, but not heard singing.					
	14. <u>Melospiza palustris</u> .— Not very numerous, but common enough in wet places.					
	15. <u>Spizella montana</u> .— Rather abundant, and scattered; but not so plentiful as in the East.					
	16. <u>Junco hyemalis</u> .— Quite abundant.					
	17. <u>Passerella iliaca</u> .— One seen at "Aunt Hepsie's farm" on Dec. 6th; another on Darryl Creek on December 7th.					
	18. <u>Passer domesticus</u> .— Quite a number about Circleville & Columbus.					
	19. <u>Cardinalis virginianus</u> .— Quite widely diffused, and in small companies. They feed on Indian corn, and are very shy.					

1881.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	Not heard singing. They are most commonly met with in weedy places beside the Scioto and its tributaries, but are found every where. I saw one in an apple-tree in the back-yard at Wittich's, in Circleville, on December 13th. They build nests in the City in summer.					
	20. <u>Sturnella magna</u> .— Found in the prairie grass, in corn fields, etc. I shot three specimens in succession as I flushed them out of the prairie in the evening twilight on December 10th.					
	21. <u>Cyanocitta cristata</u> .— Abundant in flocks of 12 to 20 in the woods, and in orchards about farmer's houses.					
	22. <u>Corvus frugivorus</u> .— Crows are rare in Ohio in winter. Dr. Wheaton told me that it is generally believed that there was a disease among Ohio Crows which destroyed them as Chicken Cholera sweeps off the genus <u>Gallus</u> . This is supposed to have destroyed the Crow crop several years since, as they were formerly plentiful in the State during winter, whereas they are now extremely scarce. A flock of about a hundred dozen flew over the Home Farm at Hulse's, on the morning of Dec. 7th; the sight of this flock was thought by Mr. Hulse to be very extraordinary. I have never seen any other crows in the state of Ohio, and they seem rare in all that inland country, gradually becoming more numerous as one approaches the coast coming eastward. Coming east, on Dec. 16th we saw enormous flocks of Crows in north-					

1881.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

-eastern Maryland, and southern Pennsylv-
-vania.

23. Picus pubescens. - Common.

24. Picus villosus. - Not very often seen.

25. Centurus carolinus. - Specimens shot on Dec. 5th and 7th were the only ones seen.

Their loud cry is easily recognized. Dr. J. M. Wheaton told me that Red-bellied Woodpeckers were exceedingly scarce in summer and rather abundant, but not in flocks but usually found singly in winter, in the vicinity of Columbus, O. The same is doubtless true for Circleville, Ohio.

26. Colaptes auratus. - Quite common, but usually seen singly, and widely diffused.

The "Wood Cock" (Hylotornus) has been seen several times of late at the Hulse's woods at the flats, but we could not find it there.

27. Scops asio. - Screech Owls were heard.

28. Asio accipitrinus. - Wherever there are swampy prairies, the "nests" of the Short-eared Owl may be seen in the grass in great plenty. The forms contain excrement and pellets of hair and bones in great plenty abundance. They are gregarious, and can see very well on wet dark days. At such times they do not like to have their plumage bedagged in the long grass, and they betake them to higher places in flocks, where the grass is short, or sit on stumps and rail-fences. Occasionally a Short-eared Owl will be seen flying about on bright days; but, ordinarily, they will not leave their forms in the prairie grass until you are close upon them. Dr. Howard Jones has taken sets of their eggs at

1881.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

Circleville, Ohio, and thinks they are about as common in summer as during winter.

29. Asio americanus. - On Dec. 7th, while hunting ~~in the prairie~~ in a strip of prairie land between two corn-fields, Mr. Campbell, who was with us, shot a Long-eared Owl in the grass. It seems to me very unusual to find a Long-eared Owl in such a situation.

30. Bubo virginianus. - Nearly every woods of any considerable size contains its pair of Great Horned Owls. They are always wild and difficult to approach. They are often seen flying about during daytime but are hard to shoot. I shot a fine old male that was sitting in the top of an enormous sycamore tree on Darby Creek. It sat perfectly motionless with ears erect, and I had to scrutinize it carefully before I was certain that it was not a part of the tree. When I fired a cartridge loaded with No. 3 shot, it made a great fluttering, but succeeded in launching out, and for a little way flew very well; then it began to rise high up in the air, and, finally, fell perfectly dead to the ground from a great height. Three shots were afterwards found to have penetrated its thorax, passing through the os sternum, and one leg was broken.

31. Circus hudsonius. - Common. One that was shot had its crop filled with mice.

32. Sturnella sparreri. - Pretty common everywhere. Seen in pairs sitting on the highest tree-tops.

33. Buteo borealis. - Abundant. We found a pair roosting in Hulse's woods at night and shot at them as they flew from the tree-tops. They made a great clamor.

1881.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

34. Accipiter fuscus. — One seen on Darby Creek, on the 10th of December.

35. Zenaidura carolinensis. — Common in large flocks along the river, and in the corn-fields, where they get most of their food.

36. Ortyx virginiana. — Quails are not numerous this season. A few flocks were seen or heard. The female I shot was seen to alight beside a straight rail fence, without any cover. Nevertheless we passed the bird, ~~and~~ going on either side of the fence, seven or eight times before we could find and put it up.

37. Bonasa umbellus. — We saw about half a dozen "Pheasants" on the Hulse farm at the Flats, one of which Isaiah Hulse shot, and I preserved. All that I saw were red.

38. Botaurus lentiginosus. — One was captured in Circleville on December 3rd, 1881.

39. Querquedula carolinensis. — Two Green-winged Teals were seen on December 10th in a little creek emptying into the Darby. Several flocks of Ducks were seen flying, but the species was not made out.

No Thrushes or Warblers were seen.

Kingfishers were also absent.

Turkey Buzzards (Cathartes aura) were not seen, but have regular roosting-places, at night, in the sycamore trees eight miles up the river, and four miles down the Scioto River. Mr. Isaiah Hulse showed me a sycamore-tree in which a family of Turkey Buzzards is reared every season. Dr. Howard Jones has two sets of eggs taken at Circleville, Ohio.

1881.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

Mammals.

Of Mammalia, Rabbits (Lepus sylvaticus), Gray Squirrels (Sciurus carolinensis), and Bank Voles (Arvicola riparia) were numerous.

One Thirteen-lined Gopher (Spermophilus tridecemlineatus) was seen by Mr. Isaiah Hulse on the 9th of December. I saw plenty of their burrows. They are very abundant here, but seldom seen but in winter.

No Chipmunks (Tamias striatus) were seen.

Specimens.

2,218. Sialia sialis. Dec. 7th. 6.88 12.65 ♂ ad.
Wing, 3.90; tail, 2.72; culmen, .49; bill from nostril, .31; gape, .79; tarsus, .83; middle toe and its claw, .88. Iris, hazel. Bill, legs and feet, black. Inside of mouth, yellow.

2,219. Regulus satrapa. Dec. 6th. 4.10 6.65 ♀ ad.
Wing, 2.18; tail, 1.58; bill measured from nostril, .22; gape, .43; tarsus, .67; middle toe + claw, .50. Iris, hazel. Bill, black. Tarsi, brownish-black; dorsal surface of toes, yellowish-brown; under surface, yellow.

2,220. Lophophanes bicolor. Dec. 6. 6.40 10.00 ♀ ad.
Wing, 13.05; tail, 2.85; culmen, .50; bill from nostril, .37; gape, .55; tarsus, .80; middle toe and its claw, .75. Iris, hazel. Bill, blue-black. Tarsi and toes, blue.

2,221. Parus atricapillus. Dec. 6. 5.37 8.45 — ? ad.
Wing, 2.55; tail, 2.45; bill from nostril, .28; culmen, .38; gape, .45; tarsus, .65; middle toe and claw, .60; Iris, dark hazel. Bill, blue-black. Tarsi and toes, blue.

2,222. Parus carolinensis. Dec. 5. 4.90 8.00 ♂ ad.
Wing, 2.50; tail, 2.28; culmen, .36; bill measured from nostril, .28; gape, .40;

over

1881.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
2,223.	<i>Parus carolinensis</i> .	Dec. 5.		4.90	7.75	♀ ad.
2,224.	<i>Parus carolinensis</i> .	Dec. 5.		4.73	7.50	♀ ad.
2,225.	<i>Cardinalis virginianus</i> .	9.15		12.25		♂ ad.
	Wing, 3.80; tail, 4.35; culmen, .75; bill from nostril, .52; gape, .77; tarsus, 1.01; middle toe and its claw, .94. Iris, hazel. Bill, red. Legs and feet light brown. Shot on Aunt Hepsie's farm Dec. 6th.					
2,226.	<i>Cardinalis virginianus</i> .	8.86		12.20		♂ ad.
	Wing, 3.80; tail, 4.32; culmen, .75; bill from nostril, .50; gape, .73; tarsus, 1.00; middle toe and its claw, .92. Colors of parts same as No. 2,225. December 6th.					
2,227.	<i>Cardinalis virginianus</i> .	9.00		12.25		♂ ad.
	Wing, 3.85; tail, 4.22; culmen, .78; bill from nostril, .50; gape, .75; tarsus, 1.00; middle toe and claw, .95. Iris, hazel. Bill, red. Legs and feet, light brown.					
2,228.	<i>Cardinalis virginianus</i> .	8.55		11.80		♀ ad.
	Wing, 3.73; tail, 4.10; culmen, .75; bill from nostril, .48; gape, .75; tarsus, .96; middle toe + claw, .92. Colors of parts like 2,227.					
2,229.	<i>Cardinalis virginianus</i> .	8.75		11.65		♀ ad.
	Wing, 3.62; tail, 4.03; culmen, .77; bill from nostril, .48; gape, .80; tarsus, 1.04; middle toe + claw, .98. Colors of parts like No. 2,227.					
2,230.	<i>Cardinalis virginianus</i> .	8.70		11.60		♀ ad.
	Wing, 3.55; tail, 3.95; culmen, .74; bill from nostril, .50; gape, .79; tarsus, .98; middle toe and its claw, .94. Colors of parts the same as No. 2,227. Nos. 2,227, 2,228, 2,229 and 2,230 were all shot at the bridge over Darby Creek on Dec. 7th.					
2,231.	<i>Sturnella magna</i> .	Dec. 10.		10.85	16.50	♂ ad.
	Wing, 5.10; tail, 3.40; culmen, 1.45; gape, 1.55; tarsus, 1.65; middle toe + claw, 1.55.					

1881.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
2,232.	<i>Sturnella magna</i> .	Dec. 10.		8.85	13.90	♀ ad.
	Wing, 4.85; tail, 3.90; culmen, 1.14; gape, 1.22; tarsus, 1.43; middle toe and claw, 1.36.					
2,233.	<i>Sturnella magna</i> .	Dec. 10.		9.85	15.35	♂ ad.
	Wing, 4.55; tail, 3.00; culmen, 1.25; gape, 1.34; tarsus, 1.46; middle toe and claw, 1.45.					
2,234.	<i>Cyanocitta cristata</i> .	Dec. 6.		11.15	16.00	♀ ad.
	Wing, 4.90; tail, 4.85; culmen, 1.12; bill from nostril, .80; gape, 1.29; tarsus, 1.35; middle toe and its claw, 1.05. Iris, hazel. Bill and feet, black.					
2,235.	<i>Picus villosus</i> .	Dec. 15.		9.10	15.00	♀ ad.
2,236.	<i>Centurus carolinus</i> .	Dec. 5.		9.90	16.45	♀ ad.
	Wing, 5.12; tail, 3.50; culmen, 1.17; gape, 1.37; tarsus, .80; middle toe with its claw, 1.08. Claw alone, .40. Iris, red. Bill and feet, bluish.					
2,237.	<i>Centurus carolinus</i> .	Dec. 7.		10.35	17.35	♀ ad.
	Wing, 5.40; tail, 3.65; culmen, 1.25; bill from nostril, .98; gape, 1.43; tarsus, .85; middle toe and its claw, 1.05; claw alone, .43; Iris, red. Bill, blue-black. Tarsus and toes, greenish-gray; claws, bluish-black.					
2,238.	<i>Asio americanus</i> .	Dec. 7.		14.75	36.90	♀ ad.
	Wing, 12.00; tail, 6.55; culmen measured from base of cere, 1.13; cere, .56; culmen from cere, .65; gape, 1.15; tarsus, 1.65; middle toe and its claw, 1.72; middle toe alone, 1.40; claw, .73. Iris, yellow.					
2,239.	<i>Bubo virginianus</i> .	Dec. 10.		22.00	52.50	♂ ad.
	Wing, 14.75; tail, 9.40; culmen from base of cere, 1.70; cere, .77; culmen measured from cere, 1.15; gape, 1.80; tarsus, 2.12; middle toe and claw, 2.90; middle toe alone, 2.35; claw, 1.15. Claws, blue-black. Bill, blue-black, also. Iris, yellow. Stomach contained a Rabbit (<i>Lepus sylvaticus</i>).					

1881.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
2,240.	<i>Circus hudsonius</i> .	Dec. 7.	18°00'	41.75	♂ ad.	
	Wing, 13.00; tail, 8.30; base of cere to point of bill, 1.03; cere alone, .46; culmen alone, .63; gape, 1.23; tarsus, 2.92; middle toe & claw, 1.78; middle toe alone, 1.30; claw, .62. Iris, yellow. Bill, horn-color at base, shading into blue within a narrow space, and becoming black on its greater extent. Cere, greenish-yellow. Tarsi and toes, lemon-yellow; claws, black.					
2,241.	<i>Zenaidura carolinensis</i> .	Dec. 10.	11°60'	17.60	♀ ad.	
	Wing, 5.50; tail, 5.20; culmen, .53; gape, .75; tarsus, .82; middle toe and claw, 1.03.					
2,242.	<i>Ortyx virginiana</i> .	Dec. 7.	9°80'	15.30	♀ ad.	
	Wing, 4.50; tail, 2.60; culmen, .54; bill from nostril, .32; gape, .61; tarsus, 1.33; middle toe & claw, 1.40; middle toe alone, 1.13; claw, .33. Iris, hazel. Maxill. and tip of mandible, black; base of mandible, flesh (or horn)-color, shading into black. Legs and feet, purplish fleshy-brown; claws, blackish.					
2,243.	<i>Bonasa umbellus</i> .	Dec. 7.	16°30'	22.65	♀	
	Wing, 6.90; tail, 5.35; bill measured from nostril, .47; gape, 1.05; tarsus, 1.70; middle toe and its claw, 1.80; middle toe alone, 1.40; claw, .46. Iris, hazel. Culmen and extreme tip of mandible, blackish. Extreme point and edge of maxilla at base, and nearly all of mandible, flesh-color. Legs and feet, grayish-blue. Claws, brownish-black.					

1881.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	December 16th. Baltimore and Ohio Railroad.					
	Coming east on the Baltimore and Ohio Railroad we saw a number of birds:—					
	<i>Corvus frugivorus</i> . — In eastern Maryland we first came across the Crows in large flock, up to 500. West of that we had seen a few scattered birds. From there east to Jersey City they were seen in large flocks.					
	<i>Cathartes aura</i> . — In eastern Maryland we saw a Turkey Buzzard flying which I recognized by its flight, although I had never seen one alive before. Shortly afterwards we saw a dead horse lying in a field; and a little later another Turkey Buzzard. After that we saw a number of them before reaching Wilmington, but none east of that point. In one place a flock was seen flying in the air. At another time we saw a number settling down upon some carion not far from the railroad. Some were floating motionless in the air.					
	<i>Molothrus ater</i> . — A number seen in Maryland, Delaware and Pennsylvania.					

December 21st. H. J. N. Y.

Took my wife out shooting for the first time. We drove into the woods below Fort Montgomery, and tied the horse. There were some Purple Finches (*Carpodacus purpureus*) and Goldfinches (*Astragalinus tristis*) feeding in the hemlocks. A number of Yellow-rumped Warblers (*Dendroica coronata*) were seen at different points. Leaving my wife to admire the view from Fort , I strolled a few rods into the edge of the woods. Whilst

1881.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

standing and watching several species of common winter birds, I noticed a bird flying strongly from the North. It alighted in a tree-top near me, and seemed animated and shy; and I thought it was going to fly, so shot it at once. It proved to be a fine young male Louisiana Tanager (*Pyrranga ludoviciana*), in good plumage, and fairly plump and well nourished. See *Forest and Stream*, X, p. 95. In Mass.

- 2,244. *Pyrranga ludoviciana*. 7.45 11.35 ♂ juv.
Wing, 3.62; tail, 3.00; culmen, .65; bill measured from nostril, .48; gape, .75; tarsus, .82; middle toe and its claw, .80; middle toe, .58; claw, .25. Iris, hazel. Bill, with maxilla dusky brown, with yellowish on edges; and mandible yellow. Tarsi and toes, bluish lead-color. This bird was in the plumage of the female, but careful dissection proved that it was a young male.

- 2,245. *Carpodacus purpureus*. 6.35 10.00 ♂ juv.
Wing, 3.05; tail, 2.38; culmen, .45; bill from nostril, .35; gape, .55; tarsus, .70; middle toe and its claw, .75. This specimen, proved by dissection to be a male in female plumage, with no trace of that of the male. On the 23rd of last June [See *antea*, p. 234] I shot a pair of Purple Finches from their nest, the male of which was in female plumage, but had one yellowish-red feather in its crown.

- 2,246. *Asragalinus tristis*. 5.15 9.00 ♂ ad.
Wing, 2.80; tail, 1.90; culmen, .43; bill from nostril, .34; gape, .43; tarsus, .55; middle toe and claw, .62. Crop filled with seeds of hemlock.

1881.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

- 2,247. *Asragalinus tristis*. 5.12 8.90 ♂ ad.
Wing, 2.75; tail, 1.90; culmen, .40; bill from nostril, .30; gape, .42; tarsus, .53; middle toe and its claw, .60. Iris, hazel. Crop filled with seeds of *Abies canadensis*.

December 25th. H. S. M.

While dressing for church a Yellow-bellied Woodpecker (*Sphyrapicus varius*) alighted on the sugar maple just outside the window. It afterwards flew to an apple tree in the orchard. It is the first I have seen this winter. There were flocks of Cedar Birds (*Aspidia cedrorum*) about the house.

December 28th. H. S. M.

- 2,248. *Dendroica coronata*. 5.75 9.05 ♀ ad.
Wing, 2.90; tail, 2.32; bill from nostril, .27; gape, .50; tarsus, .73; middle toe and its claw, .62. Iris, hazel. Bill and feet, black.

December 30th. H. S. M.

To-day Louis Jerega and myself spent hunting Ducks. There were plenty of Oldsquaws (*Harleia glacialis*) and some large black looking ducks, and one or two large flocks of *Mergus merganser*. We wounded two Oldsquaws; and they afterwards kept us busy nearly all day trying to catch them. Gulls (*Larus argentatus*) were abundant, and a pair of White-headed Eagles (*Haliaeetus leucocephalus*) was seen. On Constitution Island we flushed a couple of Grouse (*Bonasa umbellus*), & we saw a few others on shore.

1881-82.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

December 31st, 1881. H. J. N.Y.

Dendroica coronata. - One seen.*Carpodacus purpureus*. - Two small flocks.2,249. *Sialia sialis*. 7.15 12.50 ♂ ad.

Wing, 4.00; tail, 2.85; bill measured from nostril, .36; gape, .79; tarsus, .83; middle toe and its claw, .86; toe alone, .63; claw alone, .21. Iris, dark hazel. Bill, tarsi and toes above, black. Under side

2,250 *Sialia sialis*. 7.10 12.20 ♂ ad.

Wing, 3.95; tail 2.59; bill from nostril, .37; gape, .82; tarsus, .82; middle toe and its claw, .90; middle toe alone, .69; its claw .22. Colors same as No. 2,249.

2,251. *Sialia sialis*. 7.15 12.25 ♂ ad.

Wing, 3.90; tail, 2.68; bill from nostril, .34; gape, .78; tarsus, .79; middle toe & claw, .83; middle toe .64; claw .22.

Colors of parts the same as No. 2,249.

2,252. *Sialia sialis*. 7.15 12.25 ♂ ad.

Wing, 3.95; tail, 2.72; bill measured from nostril, .37; gape, .83; tarsus, .79; middle toe and claw, .84; middle toe alone, .63; claw, .22. Colors like 2,249.

January 2nd, 1882. H. J. N.Y.

2,253. *Sialia sialis*. 7.12 12.45 ♂ ad.

Wing, 4.00; tail, 2.80; bill from nostril, .34; gape, .77; tarsus, .78; middle toe and claw, .83; middle toe alone, .64; claw, .22. Colors like No. 2,249.

2,254. *Cyanocitta cristata*. 11.40 16.55 ad.

Wing, 5.25; tail, 5.12; culmen, 1.03; bill measured from nostril, .68; gape, 1.22; tarsus, 1.33; middle toe and claw, 1.22; middle toe alone, .90; middle toe alone,

1882.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

.90; claw, .37. Iris, hazel. Bill, tarsi and toes, black.

January 4th. H. J. N.Y.

Late this afternoon I saw a flock of Robins (*Turdus migratorius*) flying northward. Later I found a long straggling flight of several hundreds flying from southeast to northwest. They alighted frequently but were moving steadily forwards over the mountain, in a northwesterly direction.*Turdus pallasi*. - One or more seen and heard.

January 5th. H. J. N.Y.

Turdus migratorius. - A large flock seen in Morgan's place. Very wild!
Ampelis cedrorum. - One flock.

January 6th. H. J. N.Y.

Turdus migratorius. - A very large flock was seen under the hill, by the river, amongst the cedars. The snow-storm which commenced this morning ended in rain, and I went out in the drizzle to look for Robins, to get a full series of measurements of winter specimens. I found them in abundance, but very shy. They were singing in numbers at the time I first found them. They were wild and restless, and not inclined to feed.*Colaptes auratus*. - One seen with the Robins.*Carpodacus purpureus*. - Three seen.*Ampelis cedrorum*. - One good-sized flock.*Bonasa umbellus*. - Three were seen; one on Consook Island; two under Mr. Pell's place, among the cedars, where they had sat quietly all the morning in the snow.

1882.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

- January 6th - Continued
- 2,255. *Turdus migratorius*. 10:50 17:00 ♂ ad.
Wing, 5.35; tail, 4.25; bill measured
from nostril, .54; gape, 1.15; tarsus, 1.40;
middle toe and claw, 1.22.
Iris, hazel. Bill with top and tip of maxilla
and terminal $\frac{1}{3}$ of mandible, brownish-black;
basal half of sides of maxilla and basal $\frac{2}{3}$
of mandible, yellow. Tarsi, toes & claws,
brown; toes, blackish-brown; soles of feet,
yellowish flesh-color.
- 2,256. *Turdus migratorius*. 10:50 16:40 ♂ ad.
Wing, 5.25; tail, 4.00; bill from nostril, .57;
gape, 1.13; tarsus, 1.40; middle toe & claw, 1.15.
Colors of bill, feet, irides, etc. the same as No. 2,255.
- 2,257. *Turdus migratorius*. 10:47 16:40 ♀ ad.
Wing, 5.15; tail, 4.05; bill from nostril, .53;
gape, 1.15; tarsus, 1.29; middle toe and
claw, 1.20. Colors of parts same as No. 2,255.

January 7th. H.F., N.Y.

- Turdus migratorius*. - A large number.
- Carpodacus purpureus*. - A number. They
utter a chattering, English Sparrow-like
note while feeding. One or two old males seen.
- Harelda glacialis*. - Shot a female at Cousook Island.
- Bonasa umbellus*. Shot one in the swamp
beside Cousook marsh, and another on
Cousook Island in gray plumage.
- Ampelis cedrorum*. - One large flock.
- Melospiza fasciata*. - One shot on Cousook Id.
- 2,258. *Bonasa umbellus* (gray) 18:25 23:40 ♂ ad.
Wing, 7.30; tail, 6.40; culmen (about), .98;
nostril, .55; gape, 1.15; tarsus, 1.75; middle
toe and claw, 1.95; toe alone, 1.55; claw, .45.
Iris, hazel. Bill, dusky-brown on maxilla
above; sides of maxilla at base and mandible,

1882.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

- flesh-color. Tarsi and toes, bluish-ashy;
claws, dusky. Plumage, gray. Ruffs with
some brown and tail-band decidedly
brownish. Tail band not extending across
central tail-feathers. Crop filled with long
narrow green leaves - not laurel or wintergreen.
- 2,259. *Bonasa umbellus*. 16:30 23:65 ♀
Wing, 7.35; tail, 5.25; bill from nostril, .54;
gape, 1.13; tarsus, 1.65; middle toe and
claw, 1.90; middle toe alone, 1.40; claw, .45.
Parts colored the same as No. 2,258.
- 2,260. *Melospiza fasciata*. 6:85 8:85 ♂ ad.
Wing, 2.85; tail, 2.85; culmen, .53;
nostril, .34; gape, .54; tarsus, .90; mid-
dle toe and claw, .84. Iris, hazel.
Maxilla brownish-black; mandible horn-
color; dusky in part. Tarsi and toes, light
brown. Shot on Cousook Island.

January 8th. H.F., N.Y.

- Sialia sialis*. - Commonly present.
- Carpodacus purpureus*. - One red male.

January 9th. H.F., N.Y.

- To-day we drove to Fort Montgomery.
We saw one large flock of Cedar Birds
(*Ampelis cedrorum*). Shot 7 in two shots.
three had red appendages, one of them three
appendages to rectrices.
- Chrysomitris tristis*. - A large flock of
yellowbirds was feeding in the hemlocks,
on seeds of cones, below Fort Montgomery.
- Sialia sialis*. - Still seen or heard everywhere.
- Carpodacus purpureus*. - A few eating
seeds of hemlock cones.
- Harporhynchus rufus*. - Mr. Arthur Pell shot
one near his house on December 9th, 1881.

1882.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	January 9th. — Continued.					
2,261.	<i>Ampelis cedrorum</i> (Waxen appendages)	7.12	11.50	♀ ad.		
	Wing, 3.70; tail, 2.37; bill from nostril, .27; gape, .75; tarsus, .65; middle toe + claw, .75					
2,262.	<i>Ampelis cedrorum</i> . No appendages	7.25	11.88	♂		
	Wing, 3.75; tail, 2.40; bill measured from nostril, .27; gape, .75; tarsus, .68; middle toe and claw, .77. No wax-like wingtips.					
2,263.	<i>Ampelis cedrorum</i> (No appendages)	7.45	11.95			
	Wing, 3.85; tail, 2.66; bill measured from nostril, .27; gape, .73; tarsus, .68; middle toe and claw, .80. No wax-like appendages.					
2,264.	<i>Ampelis cedrorum</i> (Waxen append.)	7.40	12.30	♀ ad.		
	Wing, 3.90; tail, 2.68; bill from nostril, .29; gape, .79; tarsus, .70; middle toe and claw, .82. This specimen's plumage is interesting because it is a female, and possesses waxen appendages to three of its rectrices. It has met with the same accident as had the Ruffed Grouse (No. 2,258) which I shot a few days since, viz: it must have floundered against a branch or some object and torn through the skin on its neck which was found in both cases to be considerably torn, the edges of the wound having united by plastic adhesion to the trachea of the bird.					
2,265.	<i>Ampelis cedrorum</i> (Waxen append.)	7.20	11.80	♀ ad.		
	Wing, 3.75; tail, 2.25; bill from nostril, .26; gape, .70; tarsus, .68; middle toe and claw, .80.					
2,266.	<i>Ampelis cedrorum</i> (No appendages)	7.22	11.50	♂ juv.		
	Wing, 3.65; tail, 2.30; bill from nostril, .25; gape, .70; tarsus, .69; middle toe and claw, .80.					
2,267.	<i>Ampelis cedrorum</i> (No appendages)	7.22	11.75	—?		
	Wing, 3.75; tail, 2.35; bill from nostril, .27; gape, .75; tarsus, .68; middle toe and claw, .82. Sex indiscernable upon dissection.					

1882.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	January 11th H.C.M.					
	Wife and I went to the top of Crow Nest Mountain this afternoon after dinner. We heard a Bluebird (<i>Sialia sialis</i>) warbling as we stood on the summit.					
	January 12th H.C.M.					
	<i>Sialia sialis</i> . — Drove towards Ft. Montgomery and on the way saw a number of Bluebirds, besides a dozen or more Yellow-rumped Warblers (<i>Dendroica coronata</i>), and a flock of Cedar Birds (<i>Ampelis cedrorum</i>) of which wife shot two specimens in two shots.					
2,268.	<i>Parus atricapillus</i> .	5.37	7.90	♀ ad.		
	Wing, 2.50; tail, 2.40; bill from nostril, .28; culmen, .40; gape, .43; tarsus, .68; middle toe and its claw, .57.					
2,269.	<i>Dendroica coronata</i> .	5.50	8.65	♀ ad.		
	Wing, 2.70; tail, 2.04; bill from nostril, .29; gape, .51; tarsus, .70; middle toe and claw, .60. Iris, hazel. Bill and feet, black. Shot from a flock of a dozen feeding among the cedars. This one had cedar berries (<i>Juniperus virginiana</i>) in its crop.					
2,270.	<i>Ampelis cedrorum</i> (red appendages)	6.98	11.50	♀ ad.		
	Wing, 3.65; tail, 2.20; bill measured from nostril, .28; gape, .73; tarsus, .67; middle toe and claw, .80. Iris, reddish-brown. Bill, Tarsi + toes, black. The first bird my wife ever shot.					
2,271.	<i>Junco hyemalis</i> .	6.50	10.35	♂ ad.		
	Wing, 3.20; tail, 2.80; bill from nostril, .32; gape, .48; tarsus, .86; middle toe and claw, .79. Iris, hazel. Bill, flesh-color, slightly tipped with black. Tarsi and toes, brown; toes, darkest.					
2,272.	<i>Ampelis cedrorum</i> . (No waxen appendages)			♂ juv.		
	Mounted for my wife, who shot it.					

1882.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	January 13th. H.G.M.Y.					
	Three Bluebirds (<i>Sialia sialis</i>) were seen on the maple-tree outside my window, while the snow was lying deep & still falling.					
	January 14th. H.G.M.Y.					
	To-day went shooting in the cutter with my wife. At West Point we saw two male Bluebirds (<i>Sialia sialis</i>) sitting on a telegraph wire. A little way on we saw three females in a chestnut-tree.					
	<i>Astragalinus tristis</i> . - Large flocks seen near the cemetery at West Point.					
	<i>Bonasa umbellus</i> . - One near West Point. Birds very scarce in that direction. We scarcely saw any others driving through W. Point and home by the mountain road.					
	January 16th. H.G.M.Y.					
	Yesterday, Jan. 15th Robins (<i>Turdus migratorius</i>) and Bluebirds (<i>Sialia sialis</i>) were seen.					
	To-day an enormous flock of Cedar Birds (<i>Ampelis cedrorum</i>) was seen (and plenty of Robins (<i>Turdus migratorius</i>) in good-sized flocks flying north.					
	<i>Dendroica coronata</i> . - One flock of nine flying.					
	<i>Sialia sialis</i> . - Quite a number seen.					
	<i>Astragalinus tristis</i> . - One very large flock.					
2,273.	<i>Ampelis cedrorum</i> (red appendages)	7.05	11.85	♀ ad.		
	Wing, 3.90; tail, 2.60; bill from nostril, .25; gape, .72; tarsus, .66; middle toe and claw, .80. Iris, reddish-hazel. Bill and feet, black. The waxen appendix to the smallest remige of one wing was yellowish.					
2,274.	<i>Ampelis cedrorum</i> (red)	7.12	11.65	♀ ad.		
	Wing, 3.65; tail, 2.35; bill from nostril, .25; gape, .73; tarsus, .70; middle toe and claw, .84. Colors like 2,273.					

1882.

2,275.	<i>Ampelis cedrorum</i> (red)	7.35	11.75	♀ ad.		
	Wing, 3.65; tail, 2.42; bill from nostril, .25; gape, .73; tarsus, .67; middle toe and claw, .80. Iris, reddish-brown. Bill and feet, black.					
2,276.	<i>Ampelis cedrorum</i> .	7.15	11.80	♂ juv.		
2,277.	<i>Ampelis cedrorum</i> .	7.05	11.60	♀ juv.		
	This specimen exhibited faintly the light edging to the lesser wing-coverts characteristic of the young. I was probably (doubtless) a bird hatched last summer.					

January 18th. H.G.M.Y.

	To-day drove down to Fort Montgomery, where we found the following birds, all in large flocks: -					
	<i>Dendroica coronata</i> <i>Astragalinus tristis</i> . Yellow-rumped warbler.					
	<i>Ampelis cedrorum</i> . Cedar-bird.					
	<i>Turdus migratorius</i> . Robin.					
	<i>Carpodacus purpureus</i> . Purple Finch.					
	Besides the above a few Bluebirds (<i>Sialia sialis</i>) and Ruffed Grouse (<i>Bonasa umbellus</i>) were seen.					
2,278.	<i>Turdus migratorius</i> .	10.90	16.85	♂ ad.		
	Wing, 5.52; tail, 4.60; bill from nostril, .55; gape, 1.18; tarsus, 1.37; middle toe and claw, 1.20. Extreme tip of maxilla, and of mandible, black; maxilla a little brownish above; residue of bill, yellow. Tarsi and toes, dark brown; soles of feet, yellowish.					
2,279.	<i>Turdus migratorius</i> .	10.62	16.10	♀ ad.		
	Wing, 5.15; tail, 4.18; bill from nostril, .54; gape, 1.15; tarsus, 1.37; middle toe and claw, 1.20. Iris, hazel. Maxilla (except edge) and extremity (say $\frac{2}{5}$) of mandible, brownish-black. Mandible basally and edges of maxilla, yellow. Tarsi and toes, dark brown; soles, yellowish.					
2,280.	<i>Ampelis cedrorum</i> . Red	7.48	11.65	♂ ad.		
	Wing, 3.75; tail, 2.50; bill from nostril, .28; gape, .74; tarsus, .68; middle toe and claw, .80. Iris, reddish-brown. Bill and feet, black. Red appendages on wings and shafts of rectrices through yellow band also red.					

1882.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
2,281.	<i>Ampelis cedrorum</i> .	Red off.		7.40	11.50	♀ ad.
	Wing, 3.60; tail, 2.42; bill from nostril, .25; gape, .70; tarsus, .69; middle toe and claw, .81. Iris, reddish. Bill & feet, black. Only one toe anteriorly; looks as if it was congenital deformity. Throat contained 26 cedar berries.					
2,282.	<i>Ampelis cedrorum</i> .	(No red).		7.38	11.85	♂.
	Wing, 3.65; tail, 2.25; bill from nostril, .29; gape, .78; tarsus, .71; middle toe and claw, .78. Colors, same as above.					
2306.	<i>Ampelis cedrorum</i> .	(No red).		7.40	12.10	juv.
	Wing, 3.75; tail, 2.22; bill from nostril, .27; gape, .77; tarsus, .69; middle toe and claw, .77. Colors same as No. 2,281.					

State

January 19th, H.E.M.
To-day we drove to Fort Montgomery, where in the evening the Robins, Cedar-birds, Yellow-rumps and Snowbirds come to roost. We saw the following birds:—

Turdus migratorius. — Very good-sized flocks.

Sialia sialis. — One large flock.

Ampelis cedrorum.

Carpodacus purpureus.

Bonasa umbellus. — One seen in a hemlock-tree.

Deerocera coronata. — Seen in some numbers, both in the cedars at Mr. Wm. Faure's and also at Fort Montgomery.

Plectrophanes nivalis. — Yesterday Ella saw a flock of about fifty Snow Buntings fly past our west window; and to-day Mother saw nine feeding under the grape arbor while we were out shooting.

2,283.	<i>Sialia sialis</i> .			6.90	12.25	♂ ad.
	Wing, 3.85; tail, 2.50; bill from nostril, .34; gape, .80; tarsus, .82; middle toe and claw, .88. Iris, hazel; bill and feet, black; inside of mouth, yellow.					

1882.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
2,284.	<i>Turdus migratorius</i> .			10.44	16.50	♂ ad.
	Wing, 5.22; tail, 4.50; bill from nostril, .50; gape, 1.10; tarsus, 1.35; middle toe and claw, 2.18. Iris, hazel. Tarsi and toes, dark brown; soles of feet yellowish. Basal half of mandible and edges of maxilla, especially on basal half, yellowish. Inside of bill, brownish-black. Plumage mottled with white on breast.					
2,285.	<i>Turdus migratorius</i> .			10.05	15.80	♀ ad.
	Wing, 4.95; tail, 3.90; bill from nostril, .52; gape, 1.14; tarsus, 1.31; middle toe and claw, 1.22. Colors of parts the same as above. Mottled white and red breast.					
2,286.	<i>Turdus migratorius</i> .			10.00	16.00	♂ ad.
	Wing, 5.15; tail, 4.25; bill from nostril, .56; gape, 1.08; tarsus, 1.31; middle toe and claw, 1.15. Extremity of both mandibles, black; maxilla brownish above; bill otherwise, yellow. Iris, hazel. Tarsi and toes, dark brown; soles of feet, yellowish. This specimen was emaciated and dwarfed, having once had a broken wing at the carpus, allowing of very imperfect motion in flight. I think this accounts for its small size, although it is in excellent plumage.					

January 21st

Spent the day, in company with Mr. Jerome Denna, fishing through the ice on Papopen's Pond. We caught a number of Pickerel & Yellow Perch. I caught 13 Pickerels & 2 Perch for my basket. *Anorthura troglodytes nyemalis* was seen in the brush heaps beside the pond. None others have been seen here this winter, although I saw it on the Scioto River, when in Ohio. It is usually common here in winter, autumn & spring.

January 23rd.

Thermometer 10° above zero in morning; 2° below zero at 9 o'clock P.M. Hudson freezing. Did not go out, except professionally.

1882.

January 24th. N.H.M.
To-day at 9 o'clock a.m. the thermometer stood at 8° F. below zero. The sun was shining on it then, and it got up to 2° below by eleven o'clock a.m.; at 2 P.M. 1° below, and not a cloud in the sky. The Hudson completely closed. Thermometer 5° below zero at 9 P.M.

January 25th. N.H.M.
Thermometer 7° above zero at 8:30, a.m.
A few Snowbirds (*Junco hyemalis*) have returned to the yard, which they deserted during the bitter cold.

January 26th. N.H.M.
Thermometer rises 36° above zero at 9:30 a.m.
I drove down to Fort Montgomery, and took a long walk at several points on the road to see what the birds had been doing during the cold snap, and whether any of the less hardy species had left us. The species seen were as follows:—

Hirundo migratoria.—One flock of about a hundred were feeding in a narrow ravine down which a stream poured, at which, as I saw by tracks they had been drinking in numbers. They were very shy and flew out of the cedars in a scattering manner as I approached, so that I could not shoot a single specimen. They were, as I have constantly noticed to be the case during this month, accompanied by a flock of Cedar Birds (*Ampelis cedrorum*).

Sialia sialis.—Saw a few along the road.

Parus atricapillus.—Several flocks seen.

Ampelis cedrorum.—One flock of an hundred. One shot brought down 15 birds, all young; without waxed appendages, while one was

1882.

specially interesting on account of having no yellow band on tail, except a trace on one feather. The second shot, fired after all the birds were driven out that could be scared away, brought down four adults with finely-developed red appendages, while one had yellow tips to wing-feathers, and another red appendages on rectrices. I have often noticed that these fine birds associate together apart from the ill-developed ones.

Carpodacus purpureus.—Common.

Astragalinus tristis.—Common.

Junco hyemalis.—Abundant. No Tree Sparrows seen.

Bonasa umbellus.—Saw four all sitting in trees when flushed. Shot two of them, one of which carried my shot a fair long and dropped where I chanced to run across it just as it died.

Dendroica coronata.—Dozens of them; four shot.

Corvus frugivorus.—Quite gentle, as the river is closed, and men were crossing at Ft. Montgomery.

Cyanocitta cristata.—A number seen; many seen.

2,287. *Junco hyemalis*. 6.35 9.85 ♂ ad.
Wing, 3.00; tail, 2.55; culmen, .45; bill from nostril, .31; gape, .47; tarsus, .87; middle toe and claw, .75. Bill very slightly tipped black; residue flesh-pink. Tarsi, light brown; feet dark brown. Irides dark brown.

2,288. *Carpodacus purpureus*. 6.40 10.40 ♂ ad.
Wing, 3.40; tail, 2.62; bill from nostril, .36; culmen, .49; gape, .55; tarsus, .67; middle toe and claw, .77.

2,289. *Dendroica coronata*. 6.10 9.12 ♂ ad.
Wing, 2.85; tail, 2.60; bill from nostril, .31; gape, .56; tarsus, .77; middle toe and its claw, .62. Iris, hazel. Bill and feet, black.

1882.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	January 26th. - Continued.					
2,290.	<i>Dendroica coronata</i> .			5.85	9.10	♂ ad.
	Wing, 2.83; tail, 2.28; bill from nostril, .29; gape, .55; tarsus, .77; middle toe and claw, .62. Iris, hazel. Bill and feet, black.					
2,291.	<i>Dendroica coronata</i> .			5.85	9.05	— ad.
	Wing, 2.68; tail, 2.35; bill from nostril, .29; gape, .55; tarsus, .79; middle toe and claw, .64. Iris, hazel. Bill and feet, black.					
2,292.	<i>Dendroica coronata</i> .			5.65	8.80	♀ ad.
	Wing, 2.68; tail, 2.35; bill from nostril, .29; gape, .52; tarsus, .75; middle toe and claw, .61. Iris, hazel. Bill and feet, black.					
2,293.	<i>Carpodacus purpureus</i> .			6.40	10.40	♀ ad.
	Wing, 3.25; tail, 2.50; culmen, .50; bill from nostril, .38; gape, .60; tarsus, .69; middle toe and claw, .78.					
2,294.	<i>Ampelis cedrorum</i> . (red.)			7.25	11.75	♂ ad.
	Wing, 3.65; tail, 2.50; bill from nostril, .24; gape, .67; tarsus, .65; middle toe and claw, .78. Iris, reddish. Bill, legs & feet, black. Three or four primary feathers tipped yellow.					
2,295.	<i>Ampelis cedrorum</i> . (red.)			7.30	11.50	♂ ad.
	Wing, 3.55; tail, 2.52; bill from nostril, .27; gape, .75; tarsus, .68; middle toe and claw, .81. Iris brownish-red. Bill, legs and feet, black. Nearly all of rectrices possess appendages of red.					
2,296.	<i>Ampelis cedrorum</i> . (No red)			7.50	11.55	♀ juv.
	Wing, 3.60; tail, 2.45; bill from nostril, .25; gape, .71; tarsus, .68; middle toe and claw, .80. Iris reddish-brown. Bill & feet, black. Tail lacks terminal yellow band, except where indicated on one feather; probably it has just moulted, and is just regaining its tail, the last development of which may be supposed to be yellow band across the rectrices. No red appendages to wings.					

1882.

2,297. Bonasa umbellus. 17.00 23.25 ♀ ad.
Wing, 7.25; tail, 6.00; culmen, 1.00; bill from nostril, .55; gape, 1.10; tarsus, 1.65; middle toe and claw, 1.80; claw alone, .39. Iris, bright hazel. Feet, bluish-gray.

2,298. Bonasa umbellus. 16.40 22.90 ♀ ad.
Wing, 7.30; tail, 5.70; culmen, .90; bill from nostril, .47; gape, 1.05; tarsus, 1.60; middle toe and claw, 1.85; claw alone, .45. Iris, bright hazel. Feet, bluish-gray.

February 1st, N.J.N.Y.

Turdus migratorius. — Quite a number were found among the cedars. Very shy.

Sialia sialis. — One small flock at home place.

Bonasa umbellus. — Two were started from a patch of green briar, where they were buried beneath a foot of snow. One was shot as they blurred off.

Ampelis cedrorum. — One large flock with the Robins. One shot brought down six, none of which had any red waxen appendages. All were, apparently, young birds.

2,299. Sialia sialis. 7.30 12.75 ♂ ad.
Wing, 4.20; tail, 2.85; bill from nostril, .35; gape, .80; tarsus, .83; middle toe and claw, .88. Iris, hazel. Bill and feet black; soles of feet, yellowish. Inside of mouth yellow.

2,300. Sialia sialis. 7.12 12.60 ♂ ad.
Wing, 4.10; tail, 2.65; bill from nostril, .33; gape, .78; tarsus, .80; middle toe and claw, .87. Colors of bill and feet same as above.

2,301. Sialia sialis. 6.82 12.15 ♂ ad.
Wing, 4.05; tail, 2.52; bill from nostril, .36; gape, .85; tarsus, .83; middle toe and claw, .89. Iris, hazel. Bill & feet, black; soles of feet, yellowish. Inside of mouth, yellow.

1882.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
2,302.	<i>Sialia sialis</i> .			7.25	12.80	♂ ad.
	Wing, 4.00; tail, 2.85; bill from nostril, .37; gape, .82; tarsus, .78; middle toe + claw, .84. Colors, same as above.					
2,303.	<i>Sialia sialis</i> .			7.15	12.60	♂ ad.
	Wing, 3.95; tail, 2.73; bill from nostril, .37; gape, .81; tarsus, .82; middle toe + claw, .85. Colors, same as above.					
2,304.	<i>Sialia sialis</i> .			6.80	11.75	♀ ad.
	Wing, 3.75; tail, 2.53; bill from nostril, .35; gape, .77; tarsus, .77; middle toe and claw, .82. Colors same as above.					
	<i>Bonasa umbellus</i> . - The cat spoiled it. Its ruffs and brown tail-band were brown. An old cock bird.					

February 2nd. H.S.M.

A family of Ruffed Grouse (*Bonasa umbellus*) were found between Fort Clinton and Lee's Point on West Point. The snow is 14 inches deep.

- 2,305 *Rallus virginianus*. — — — ad.
Caught at Circleville, Ohio, on November 24th, 1880, by Dr. N. E. Jones's Spaniel. See *antea* p. 186.

February 7th. H.S.M.

Today walked to West Point with my rifle to shoot an Eagle near Battery Knox. Got an excellent shot at the Eagle at very close range, and missed, or only wounded it. It was a large light gray Eagle (*Haliaeetus leucocephalus*, juv.). The Blue Birds (*Sialia sialis*) were singing as I walked home.

February 10th. H.S.M.

Bluebirds (*Sialia sialis*) have been about every day since the 7th.

1882.

- 2,307. *Junco hyemalis*. 6.40 9.50 ♀ ad.
Wing, 2.93; tail, 2.70; culmen, .44; gape, .48; tarsus, .85; middle toe and claw, .76.
Carpodacus purpureus. - Singing.
During January Bluebirds have been heard almost daily, and yellow-rumped Warblers seen.

February 14th. H.S.M.

- 2,308. *Sialia sialis*. 6.97 12.65 ♂ ad.
Wing, 4.00; tail, 2.60; bill from nostril, .37; gape, .81; tarsus, .83; middle toe and claw, .85.
2,309. *Sialia sialis*. 7.00 12.38 ♂ ad.
Wing, 3.90; tail, 2.75; bill from nostril, .35; gape, .79; tarsus, .80; middle toe and claw, .88.
2,310. *Sialia sialis*. 6.75 12.20 ♀ ad.
Wing, 3.90; tail, 2.58; bill from nostril, .37; gape, .85; tarsus, .73; middle toe and claw, .82.
Haliaeetus leucocephalus. - Rode to West Point, tied my horse behind Ft. Clinton, and went to hunt my Eagle. Got a fairly good shot at the same bird I shot at Feb. 7th - a very large gray bird. My bullet struck the ice exactly underneath the Eagle, and as it arose I saw that my first shot had cut one or two quills from its left wing.
Dendroica coronata. - Seen on the 12th near Fort Montgomery among the cedars & Cedar Birds.

February 16th. H.S.M.

Esalon columbarius. - One seen sitting on a tall maple-tree in Fort Clinton, West Point, and sat still until I was tying my horse preparatory to trying a crack at him with my little rifle, when he recollected an engagement with a flock of Purple Finches (*Carpodacus purpureus*), which he scattered right and left. It was a fine old blue bird.

1882.

- Collecting Number. NAME. DATE. LOCALITY. Length. Stretch. Sex and Age.
- 2, 311. Parus atricapillus. February 17th. H.S.M. 5.40 8.10 ♂ ad.
Wing, 2.50; tail, 2.50; culmen, .37; nostril, .29; gape, .46; tarsus, .67; middle toe and claw, .62. Bill, black. Iris, dark hazel. Tarsi and toes, bluish. Shot by E.W.M.
- 2, 312. Parus atricapillus. 5.35 7.85 ♀ ad.
Wing, 2.45; tail, 2.50; culmen, .36; gape, .43; tarsus, .64; middle toe and claw, .56; bill measured from nostril, .27. Colors of Irides, bill and feet the same as No. 2,311. Shot by E.W.M.

February 18th.

This morning the thermometer was 10° above zero F. at 8 A.M., and I rode over the mountain to Mr. E. P. Roe's residence in Cornwall. Near Cornwall I saw one Robin (Turdus migratorius), a flock of Purple Finches and a number of Bluebirds & one Eagle.

February 19th

This evening, rode over the mountain leaving Mr. Roe's house about five P.M. Near the top of the mountain a dense cloud or fog enveloped everything, and it was impossible to distinguish objects. Still my pony kept on in the road all right until a flock of Cedar Birds (Ampelis cedrorum) whurred up from the road almost under my horse's feet, with their loudest notes which alone served to identify them. The horse was frightened and almost threw me, as I was riding very carelessly. The birds must have gone down to drink in the road, or else have gotten lost in the fog.

1882.

- Collecting Number. NAME. DATE. LOCALITY. Length. Stretch. Sex and Age.
- 2, 313. Sialia sialis. February 27th. H.S.M. 7.10 12.70 ♂ ad.
Wing, 4.00; tail, 2.75; bill, measured from nostril, .37; gape, .88; tarsus, .83; middle toe and claw, .89. Bill, black, with base of mandible and inside of mouth, yellowish. Iris, hazel. Tarsi and dorsal surface of toes, black; bottom of feet and toes, yellow. Shot by E.W.M.
- 2, 314. Picus pubescens. 6.85 11.75 ♀ ad.
Wing, 3.65; tail, 2.60; culmen, .68; gape, .78; tarsus, .60; middle toe and claw, .65. Iris, brownish-red. Bill, blue-black. Tarsi and toes, bluish gray. Shot by E.W.M.
- February 28th. H.S.M.
- 2, 315. Ampelis cedrorum. (Red.) 7.35 11.80 ♂ ad.
Wing, 3.70; tail, 2.75; bill, measured from nostril, .27; gape, .79; tarsus, .65; middle toe and claw, .77. Red appendages on wings, and very faint yellow edging to a few remiges. Iris, reddish-brown. Bill and feet, black.
- 2, 316. Ampelis cedrorum. (Red.) 7.45 11.90 ♀ ad.
Wing, 3.80; tail, 2.65; bill, measured from nostril, .26; gape, .76; tarsus, .69; middle toe and claw, .83. Colors like No. 2,315.
- 2, 317. Ampelis cedrorum. (Red.) 7.50 11.85 ♂ ad.
Wing, 3.77; tail, 2.65; bill, measured from nostril, .29; gape, .77; tarsus, .67; middle toe and claw, .83. Colors the same as No. 2,315.
- 2, 318. Ampelis cedrorum. (Red.) 7.40 11.65 ♂ ad.
Wing, 3.85; tail, 2.60; bill from nostril, .27; gape, .75; tarsus, .67; middle toe and claw, .83. Colors the same as No. 2,315. Yellow on remiges.
- 2, 319. Ampelis cedrorum. (Red.) 7.25 11.90 ♂ ad.
Wing, 3.75; tail, 2.50; bill from nostril, .28; gape, .77; tarsus, .69; middle toe and claw, .80. Colors of Irides, bill and feet the same as No. 2,315.

1882.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
February 28th — Continued.						
2,320.	<i>Ampelis cedrorum</i> . (No. red.)	7.20	11.90	—?		
	Wing, 3.70; tail, 2.25; bill from nostril, .25; gape, .75; tarsus, .67; middle toe and claw, .80. Colors of bill, feet and tides the same as No. 2,315.					
2,321.	<i>Ampelis cedrorum</i> . (No. red.)	7.50	11.90	♂		
	Wing, 3.85; tail, 2.60; bill from nostril, .28; gape, .75; tarsus, .72; middle toe and claw, .81. Colors of tides, bill & feet, the same as No. 2,315.					
2,322.	<i>Ampelis cedrorum</i> . (red.)	7.30	11.75	♀ ad.		
	Wing, 3.68; tail, 2.40; bill from nostril, .26; gape, .79; tarsus, .72; middle toe and claw, .84. Colors of parts the same as No. 2,315.					
2,323.	<i>Ampelis cedrorum</i> . (No. red.)	7.25	11.85	♀		
	Wing, 3.70; tail, 2.38; bill from nostril, .25; gape, .75; tarsus, .65; middle toe and claw, .76. Colors of parts the same as No. 2,315.					
2,324.	<i>Passer domesticus</i> .	6.40	9.90	♂ ad.		
	Wing, 3.00; tail, 1.55; culmen, .53; gape, .65; tarsus, .76; middle toe and claw, .87. Iris, hazel. Bill, black. Tarsi & toes, brown.					
2,325.	<i>Junco hyemalis</i> .	6.40	9.90	♂ ad.		
	Wing, 3.10; tail, 2.70; culmen, .44; gape, .49; tarsus, .82; middle toe and claw, .75. Iris, dark hazel. Bill, pale pink. Tarsi, flesh-color; feet, dusky-brown.					
2,326.	<i>Junco hyemalis</i> . Shot by E.W.M.	6.40	9.95	♂ ad.		
	Wing, 3.15; tail, 2.90; bill (culmen), .44; gape, .48; tarsus, .83; middle toe and claw, .75. Iris, dark hazel. Bill, pale pink. Tarsi, fleshy-brown; feet, brown. Shot by Ella W. Mearns.					
2,327.	<i>Junco hyemalis</i> . Shot by E.W.M.	6.35	10.24	♂ ad.		
	Wing, 3.12; tail, 2.78; culmen, .45; gape, .50; tarsus, .85; middle toe and claw, .83. Iris, dark hazel. Bill pale pink; maxilla slightly tipped with black. Tarsi and toes dusky brown. Shot by Ella W. Mearns.					

1882.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
2,328.	<i>Sialia sialis</i> .	Shot by E.W.M.	6.80	12.00	♀ ad.	
	Wing, 3.65; tail, 2.50; bill, measured from nostril, .35; gape, .80; tarsus, .80; middle toe and claw, .81. Irides, hazel. Bill and feet, black; inside of mouth, heels, and "soles" of feet, yellow. This bird was mated with No. 2,329, and they were hunting up a nesting place.					
2,329.	<i>Sialia sialis</i> .	7.00	12.00	♂ ad.		
	Wing, 3.90; tail, 2.68; bill, measured from nostril, .34; gape, .83; tarsus, .68; middle toe and claw, .80. Colors of parts the same as those of its mate, No. 2,328. The measurements (including tarsus) exact.					

March 2nd. H. S. M.

To-day the weather is still very warm for this season; the snow nearly all melted away by the warm heavy rain, which has filled all of the depressions in the fields with miniature lakes. The Blue Birds (*Sialia sialis*) are looking for breeding places. Several European House Sparrows have been prospecting for nesting sites about the house, ~~had~~ and have paid the penalty of intruding with their skins and lives. Cedar Birds (*Ampelis cedrorum*) have been abundant. I, shooting into a flock, killed six, all immature (?) having no waxen appendages. Most of the flock flew away; but a small number alighted at a short distance. I fired into and killed five of them, all having waxen appendages.

Again my wife shot nine from a pear-tree, all of which were young (?), having no red appendages. The flock then alighted in the edge of the woods, where she fired again and killed six more, four of which had red appendages.

Again, driving home from a shooting excursion we came to a large flock of Cedar Birds, sitting at a distance from the road, I believe that they

1882.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

were the old birds, and, if I had shot into them I would have secured only well-developed specimens having waxen appendages to all of their wings, & some of their tails, and a few with yellow spots on their wings. But nearer the road, feeding in the cedars were large numbers of the birds, which, on being flushed, alighted all together in a large tree-top. I fired into them and killed eleven, without a single red mark on one of them.

Now it has been my constant experience in winter, when shooting Cedar Birds, to kill ^{heavily or quite} all old, or else all young ones at a single shot. The old birds, when feeding, keep apart by themselves, and are the last to leave when frightened or shot at. A great deal of experience has made me perfectly certain of these points.

It is a week or two since I saw the Yellow-rumped Warbler (Dendroica coronata) about. I think that the warm spell has started them all off northward, and, the migration not having commenced as yet, we are left without them. This was the case in the spring of 1878.

Our winter Robins (Turdus migratorius) have also deserted us.

The migrations have not begun even with the Snowbirds (Junco hyemalis) and Tree Sparrows (Spizella montana); and no migrating Song Sparrows (Melospiza fasciata) have been seen. There are only the few winter residents in their usual haunts, and not singing.

Crows (Corvus frugivorus) are migrating in a constant, straggling manner. They

1882.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

are mating also.

No Blackbirds, Fox Sparrows, or Phoebes have yet been seen, although the weather is so warm. The ordinary early butterfly, having buff edging to its chocolate-colored wings was seen to-day for the first time.

The Piping Frogs (Hyla Pickeringii) were also first heard to-day.

I caught a Spotted Turtle (Neuremys guttata) this morning, also the first of the season. Its left fore-foot was gone, and the stump had the appearance of being congenitally deformed, rather than the result of an accident.

The ice in the Hudson, which had been blocked above West Point has been broken up by the heavy rain fall; and Herring Gulls (Larus argentatus) are flying in flocks.

Golden-crested Kinglets (Regulus satrapa) which have been absent (?) or extremely scarce this winter are beginning to come about the house, and are seen in the woods.

White-bellied Nuthatches (Sitta carolinensis) have been rather numerous for some weeks, and a few have been about all winter. No Red-bellied Nuthatches (Sitta canadensis) were seen during last autumn's migration, nor have any been seen this winter past.

The Brown Creeper (Certhia familiaris), usually a common winter bird, has not been once seen this year; and a small number seen near Harrison's, on December 30, 1881, are the only ones I have noticed this winter.

The following species have been abundant all winter:

Turdus migratorius, Sialia sialis, Carpodacus purpureus, Astragalinus tristis, Dendroica coronata, and Empelis cedrorum, besides a

1882.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

March 2nd. — Continued.

Few of the regular birds like the Snowbird & Crow. Tree Sparrows (*Spizella montana*) though not scarce have been much less common than they usually are.

Picus villosus I have not seen this year more than once or twice.

The Winter Wren (*Troglodytes hyemalis*) I have seen but once this year — on Jan 21st at Poplofen's Pond. See p. 277.

The Yellow-bellied Woodpecker has been seen only once this winter — on Dec. 25, 1881.

The Golden-winged Woodpecker (*Colaptes auratus*) has also but one record for the last winter, viz: Jan. 6th, 1882. See p. 269.

Snow Buntings (*Plectrophanes nivalis*) have been seen but twice — Jan. 18th & 19th, 1882.

Eagles (*Haliaeetus leucocephalus*) have been about all winter, probably breeding, but have been rather scarce.

It has been a poor season for winter birds.

2,330* *Astragalinus tristis*. 5.00 8.60 ♀ ad.

Wing, 2.85; tail, 1.90; culmen, .40; gape, .42; tarsus, .53; middle toe and claw, .59.

Iris, hazel. Tarsi and toes, brown anteriorly and superiorly, respectively; flesh-colored posteriorly and inferiorly. Shot by E. W. M.

2,331. *Carpodacus purpureus*. 6.12 10.00 ♀ ad.

Wing, 3.10; tail, 2.35; culmen, .46; gape, .57; tarsus, .68; middle toe and claw, .70. Iris, hazel.

Bill above, dusky; below, horn-color. Tarsi and toes, brown. Shot by E. W. M.

2,332. *Ampelis cedrorum*. (Red.) 7.05 11.70 ♀ ad.

Wing, 3.75; tail, 2.50; bill, measured from nostril, .25; gape, .70; tarsus, .65; middle toe and claw, .75. Iris, brownish-red.

Bill and feet, black. Shot by E. W. M.

1882.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

2,333. *Ampelis cedrorum*. (Red.) 7.35 11.90 ♂ ad.
Wing, 3.80; tail, 2.60; bill, measured from nostril, .25; gape, .74; tarsus, .67; middle toe and claw, .80. Iris, brownish-red. Bill and feet, black. Its throat contained 18 large cedar berries. Shot by E. W. M.

2,334. *Ampelis cedrorum*. (Red.) 7.40 11.90 ♂ ad.
Wing, 3.75; tail, 2.50; bill, measured from nostril, .27; gape, .77; tarsus, .76; middle toe and claw, 1.85. Iris, brownish-red. Bill and feet, black. Shot by E. W. Mearns.

2,335. *Ampelis cedrorum*. (No red.) 7.40 11.90 ♂
Red shafts through the yellow terminal band of tail; but, as stated above, there are no waxen appendages. Shot by Ella W. Mearns.

2,336. *Ampelis cedrorum*. (No red.) 7.15 11.50 ♂
Traces of brown in the blue of the rump, which is a sign of youth. Shot by Ella W. Mearns.

March 3d.

H. S. M.

2,337. *Passer domesticus*. 6.00 10.15 ♂ ad.

Wing, 3.15; tail, 2.60; culmen, .56; gape, .63; tarsus, .77; middle toe and claw, .80. Iris, hazel. Bill, black. Legs and feet, brown. Shot by E. W. M.

2,338. *Passer domesticus*. 6.35 9.50 ♀ ad.

Wing, 2.90; tail, 2.40; culmen, .55; gape, .63; tarsus, .71; middle toe and claw, .78.

Iris, hazel. Bill, ~~black~~ olive-brown; basal half of mandible and edges of maxilla at base, yellow. Tarsi and toes, light brown. Shot with its mate above by Ella W. Mearns.

March 4th.

H. S. M.

Passerella iliaca. — First seen.

One flock of three, of which Ella shot two. Then a single bird, and another flock of four were seen. We could find no more.

1882.

- March 4th — Continued.
 although we hunted over all their favorite ground, as, indeed, we have done for several days past. So I think these were the first arrivals.
Iurdus migratorius. — Have a dozen of migrants were seen, — the first of the season.
Dendroica coronata. — One seen.
- 2,339. Passerella iliaca. 7.65 11.55 ♂ ad.
 Wing, 3.60; tail, 3.15; culmen, .55; gape, .64; tarsus, 1.00; middle toe and claw, .95.
 Iris, hazel. Bill, dusky brown above; sides of maxilla at base and basal one-half of mandible, yellow; terminal half of mandible, horn-color. Legs and feet, brown.
 Shot by Ella W. Mearns.
- 2,340. Passerella iliaca. 7.35 11.10 ♂ ad.
 Wing, 3.45; tail, 2.90; culmen, .55; gape, .62; tarsus, .99; middle toe and claw, 1.00.
 Colors of parts the same as No. 2,339. Shot by Ella W. Mearns.

March 5th. H.G.M.

Walking to church this morning we saw and heard a Song Sparrow, the first of the season. It sat in a small tree in a Blackberry patch. Coming home others were seen. These are the first migrants although in certain spots they have stayed here all winter.

A Yellow-bellied Woodpecker (Sphyrapicus varius) was also seen as we walked home.

March 6th. H.G.M.

Melospiza fasciata. — More abundant.

Singing all the morning in the rain.

Passerella iliaca. — Quite numerous this afternoon, but only its Cedar-bird note heard.

Iurdus migratorius. — A few about.

1882.

- Asas obscurus. — A flock started at Pell's Pond, but could not get a shot.
- 2,341. Passerella iliaca. 7.30 11.00 ♀ ad.
 Wing, 3.45; tail, 2.95; culmen, .56; gape, .59; tarsus, 1.00; middle toe and claw 1.00 (the same as tarsus). Iris, hazel. Bill with maxilla above, dusky-brown; basal half of mandible and sides of maxilla at base, yellow; tip of mandible, dusky, shading through horn-color to the yellow. Tarsi and toes pale brown, with a very perceptible pink color shining through when first shot.
 Shot by E. W. M.
- 2,342. Orcus villosus pubescens. 6.85 12.05 ♂ ad.
 Wing, 3.62; tail, 2.50; culmen, .73; gape, .82; tarsus, .56; middle toe and its claw, .68.
 Shot by E. W. M.

March 8th. H.G.M.

This morning I heard a Purple Finch (Carpodacus purpureus) singing with unusual sweetness. I shot it, and supposed it to be a female, as it was in the plumage of the female. On dissection it proved to be a male.

Mr. Arthur Pell shot a Black-throated Green Warbler (Dendroica virens) in an apple tree near his house, as late as November 23, 1881.

- 2,343. Carpodacus purpureus. 6.35 10.30 ♂.
 Wing, 3.15; tail, 2.35; culmen, .49; gape, .57; tarsus, .69; middle toe and claw, .77. Iris, hazel. Tarsi and toes, dusky brown. In female plumage.

March 9th. H.G.M.

Iurdus migratorius. — A stormy day with snow and hail brought Snowbirds, Tree Sparrows, Song Sparrows, Fox Sparrows,

1882.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

March 9th - Continued.
and Robins about the house.

A Great Horned Owl (*Bubo virginianus*)
was chased for some time in the woods
below the house, but was too shy to be
obtained.

2,344. *Ceuthia familiaris*. 5.50 7.90 ♂ ad.

Wing, 2.58; tail, 2.42; culmen, .65;
bill, measured from nostril, .47; gape, .79;
tarsus, .58; middle toe and claw, .64.
Iris, hazel. Bill, with ^{maxilla} mandible, black;
mandible, flesh-color, -pinkish. Tarsi
and toes, fleshy-brown. Shot by E. W. M.

March 12th. No. 111 E. 72nd St., N.Y. City.

Yesterday my wife and I came to New York
to do some shopping. The tide was low when we
reached the Harlem River, and on the mud were
Herring Gulls (*Larus argentatus*) in large flocks.

To-day Louis Jerega and I spent the
afternoon in Central Park. On the central
wall of the Croton Reservoir were hundreds
of Sea Gulls (*Larus argentatus*) and others
were seen frequently passing two and four
between the river and the Reservoir. At the
north-west end of the Park we found a pair
of them catch goldfishes in the "Pool".
The pond is so small that the Gulls were
somewhat cramped for space in making their
aerial evolutions. It was a very pretty thing
to watch them, as we could easily see every
motion from the brink of the Pool where we
were standing. The goldfishes were seen
lying in shallow water along the edge; but
whenever one approached the surface out in
the pond a Gull would swoop down, and
we found that his chance was about one

1882.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

two. They sometimes chased each other and
tried to take a fish which the other bird had
captured, uttering screams.

Fox Sparrows (*Passerella iliaca*) were numerous
and singing on every hand very sweetly.

The Phoebe (*Sayornis fusca*) was found
in the dusk of evening in a quiet corner of
"the lake" wagging its tail, which attracted
our attention. We got within a few feet of it, but
it was silent still, and sat wagging its tail in
the alders along the shore, apparently enjoying
the shadow effect pictured in the water beneath
it. It is the first seen this year.
Luscalus purpureus was abundant.

March 19th. H. S. M.

Spent the whole of last week in the libraries of
the American Museum. The library is opened at
ten A.M. & closes at five P.M. I always walked
across the Park, and saw a number of wild
birds. Fox Sparrows were always plentiful.
Song Sparrows (*Melospiza fasciata*) sang every day.
A pair of Cardinal Grosbeak (*Cardinalis virginiana*)
was seen on March 13th, and a male on
the 16th. White-throated Sparrows (*Zonotrichia
albicollis*) were first heard singing on the 16th.
The Swamp Sparrow was heard on the 12th.
Crow-Blackbirds (*Quiscalus*) were seen every
time. Robins (*Turdus migratorius*) were
singing on the evening of March 16th.
Snow-birds (*Junco hyemalis*) and Goldfinches
were also singing. A Brown Creeper was
seen on the 15th. Black-capped Chickadees
(*Parus atricapillus*) were sometimes seen, as were
the Bluebirds (*Sialia sialis*). Both Coopers
and Red-shouldered Hawks (*Buteo lineatus*)
were seen on the 12th. In walking across
the park one morning five different Gray

1882.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

March 19th — Continued.

Squirrels (*Sciurus carolinensis*) were seen; and one evening returning after five o'clock with Louie, we saw a Rabbit (*Lepus sylvaticus*) and six Gray Squirrels. Two of the latter were playing like Kittens on the grass, jumping over each other, and frisking entertainingly about.

To-day, mother Ella and myself saw the Bluebirds (*Sialia Sialis*) gathering materials for a nest, and inspecting the various tin cans which I have placed in trees for their use.

A Red Squirrel (*Sciurus hudsonius*) was seen to come and climb up the pole covered with rose vines to the bird-house, where he may have gone to look for birds eggs or to deposit store of food; or possibly to form a nest for himself.

March 20. H.F., N.Y.

2,345. *Lanius ludovicianus*. 9:10 12:90 ♂ ad.

Wing, 3.80; tail, 4.10; bill, measured from nostril, .47; culmen, .70; gape, .98; tarsus, 1.10; middle toe and claw, .93; toe alone, .70; claw, .27. Iris, dark hazel. Bill black; blue at base of mandible. Tarsi and toes, black.

The above specimen (No. 2,345) was shot at Circleville, Ohio, on March 16th, 1882, and was sent to be by mail. Shot on Campbell's meadow.

To-day a pair of Phoebe (*Sayornis fuscus*), evidently the old ones that were in the same place last season, came to Mr. Maloney's barn at the end of our place, where they were heard before we were up in the morning, and began to work at the old nest at once. They are the first seen about here this season. I saw one in Central Park on the 12th, but no others

1882.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

were seen there during the whole of the ensuing week; and my wife says that none were seen here during the last week.

This morning the Wood Frogs were making a great racket in the swamps where they go to breed every spring. I captured a pair in a brook, as I was going to pick up a pair of Fox Sparrows which I had shot. The female was double as large as the male, and brown, while the male was blackish. They were copulating and the male held the female fast while I held the latter by the hind leg for a long time. Then I put both inside of a paper, and when I reached home he still was embracing the female. I put both together in a jar of alcohol, and they are preserved in the same attitude.

The first Painted Turtle (*Chrysemys picta*) was seen and captured this morning. It was an immature one, and a large leech was attached to it. It was preserved with the two specimens of *Rana sylvatica* above mentioned.

The Fox Sparrows (*Passerella iliaca*) were singing, and so were Purple Finches (*Carpodacus purpureus*). My wife, comparing the songs of the two birds, said that there was the difference between them that there is between a clear soprano and a rich contralto voice.

Since writing the above I found the Wood Frogs in large numbers in a shallow pool north of Highland Lake, where they fairly swarmed, and their eggs were in large masses. The Phoebe (*Sayornis fuscus*) was seen by the water-wheel at the old mill as I rode past. There are still tons of ice on the side of the mountain behind the lake, where miniature glaciers formed during winter.

1882.

- Collecting Number. NAME. DATE. LOCALITY. Length. Stretch. Sex and Age.
- 2,346. Passerella iliaca. March 20th - Continued. 7.45 11.30 ♂ ad.
Wing, 3.55; tail, 3.20; culmen, .55; gape, .58; tarsus, .93; middle toe and claw, .97.
- 2,347. Passerella iliaca. 7.20 11.30 ♂ ad.
Wing, 3.53; tail, 2.92; culmen, .58; gape, .62; tarsus, .95; middle toe and claw, .98.

March 21st. H.G.M.Y.
Agelaius phoeniceus. First seen. Several were feeding on the ground in the "old orchard".

March 24th. H.G. N.Y.
Visited Long Pond this evening and heard the ~~tweedle-tweedle-tweedle~~ of the Woodcock (Philohela minor) as it flew up in the air in the neighboring swamp, and then a loud note uttered in a harsh tone as it descended.

A Blackbird (Agelaius phoeniceus) was flying over the lake. The weather was cold. Thermometer about 26° F. Saw one Winter Wren; the first seen since January 21st.

March 25th. H.G.M.Y.
To-day caught five good-sized quackrels in Long Pond, and shot a Woodcock and a Hairy Woodpecker.

- 2,348. Philohela minor. 11.10 17.90 ♂ ad.
Wing, 4.95; tail, 2.70; culmen, 2.60; gape, 2.55; tarsus, 1.22; middle toe and its claw, 1.42. Iris, hazel.

- 2,349. Picus villosus. 9.45 15.60 ♂ ad.
Wing, 4.75; tail, 3.30; culmen, 1.27; gape, 1.35; tarsus, .88; middle toe and claw, .90; toe alone, .61; claw, .39.
Bill, uniform blue-black. Tarsi and toes, bluish-gray.

1882.

- March 26th. H.G.M.Y.
Sayornis fuscus, is now abundant. In the evening they fly high up in the air uttering rapid cries like those made by the Kingbird.
- 2,350. Passerella iliaca. Shot by E.W.M. 7.15 11.20 ♀ ad.
Wing, 3.45; tail, 2.80; culmen, .54; gape, .57; tarsus, .99; middle toe and claw, .95.
- 2,351. Sayornis fuscus. Shot by E.W.M. 7.00 10.90 ♂ ad.
Wing, 3.38; tail, 2.90; bill, measured from nostril, .40; gape, .86; tarsus, .74; middle toe and claw, .62. Iris, hazel. Bill and feet, blackish.
- 2,352. Sayornis fuscus. Shot by E.W.M. 7.55 11.65 ♂ ad.
Wing, 3.60; tail, 3.33; bill, measured from nostril, .44; gape, .85; tarsus, .73; middle toe and claw, .65. Iris, hazel. Bill and feet, black.
- 2,353. Turdus migratorius. Shot by E.W.M. 10.40 16.55 ♂ ad.
Wing, 5.10; tail, 4.25; bill, measured from nostril, .55; gape, 1.13; tarsus, 1.38; middle toe and claw, 1.15. Iris, hazel. Bill, yellow, with tip of both mandibles, yellow. Tarsi and toes, dark brown.
- 2,354. Turdus migratorius. Shot by E.W.M. 10.00 15.65 ♀ ad.
Wing, 4.85; tail, 4.00; bill, measured from nostril, .55; gape, 1.08; tarsus, 1.33; middle toe and claw, 1.15. Bill, all dusky. Tarsi and toes brownish-black. Iris, hazel.

- March 29th. H.G.M.Y.
2,355. Colaptes auratus. 13.10 20.40 ♂ ad.
Flickers are now abundant.

- March 30th. H.G.M.Y.
2,356. Colaptes auratus. 13.00 20.55 ♂ ad.
Wing, 6.25; tail, 5.00; culmen, 1.50; gape, 1.63; tarsus, 1.13; middle toe and claw, 1.30.
- 2,357. Sayornis fuscus. 7.10 11.30 ♂ ad.
Wing, 3.40; tail, 3.00; bill, measured from nostril, .44; gape, .88; tarsus, .70; middle toe and claw, .60.

1882.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

April 4th. H. G. M. Y.

Dendroica coronata. - One alighted in front of the house. It was the first seen for several weeks, although, through the winter, they were rather common. Mr. Arthur Pell told me that he noted a White-throated Sparrow (Zonotrichia albicollis) on March 29th. I have not found it yet.

April 5th. H. G. M. Y.

2,358. Colaptes auratus. 12.80 20.10 ♂ ad.
Wing, 6.00; tail, 4.80; culmen, 1.53; gape, 1.65; tarsus, 1.13; middle toe and claw, 1.27; Iris, dark red. Bill, blue-black. Tarsi and toes, bluish-gray.

April 7th.

Siurus motacilla. - First seen. One was singing very sweetly on Poplopen's Creek late this afternoon.

Ceryle alcyon. - Doubtless they have been about a long time. But was first seen to-day on Poplopen's Creek.

Scolecophagus ferrugineus. - First seen to-day.
Turdus pallasi. - First seen. Only one solitary one, which I shot.

2,359. Astragalinus tristis. 5.05 9.00 ♂ ad.
Wing, 2.87; tail, 2.02; culmen, .39; gape, .40; tarsus, .55; middle toe and claw, .64.

2,360. Turdus pallasi. 7.25 11.70 ♂ ad.
Wing, 3.65; tail, 2.95; bill, measured from nostril, .38; gape, .88; tarsus, 1.23; middle toe and its claw, .92.

2,361. Zenaidura carolinensis 11.90 17.65 ♀ ad.
Wing, 5.75; tail, 5.65; culmen, .56; gape, .81; tarsus, .82; middle toe and claw, .97.

1882.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

April 8th. - H. G. M. Y.

Spizella socialis. - First seen. A single one was trilling in a pear-tree by the house. Pandion haliaetus carolinensis was first seen on April 6th; but has probably been here for some time. I have not seen much of the River lately. Larus argentatus is still seen in small numbers on the River.

A small Gull, probably Bonaparte's or the Laughing Gull, was seen early this morning flying over the River, near Green Point. Ceryle alcyon was seen in some numbers on Poplopen's Creek.

Siurus motacilla was heard again singing on Poplopen's Creek, loud & sweet. On the banks of the Creek were beds of Hepaticas in full bloom, the first I have seen this season. I gathered a large bouquet of them.

Mr. Jerome Denna and myself set a net in Poplopen's creek the night before. In it we caught a number of Common Suckers, One "Wind Fish" a nice mess of good-sized, White, and Yellow Perches, ^{and one sign fish.} We saw one Pickerel in the stream and a number of minnows. The sun fish was the Short-eared species (Eupomotis aureus); but last autumn I took Sepiopus auritus in great numbers high up the Poplopen branch of the stream and in the "Devil's Hole" under the bridge, and in the part of the creek which the tide affects - at the Mill. I also took them at Garrison's drawbridge and Satterell's Point, in the Hudson itself.

2,362. Ceryle alcyon. 13.35 22.00 ♂ ad.
Wing, 6.25; tail, 3.75; culmen, 2.33; gape, 2.88; tarsus, .42; bare part of tibia, anteriorly, .40; middle toe and claw, .88.

1882.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

April 8th - Continued

Mr. Jerome Denna told me that when fishing last summer at Lake Oseawana, in Putnam County, N.Y., he was fishing around the corner of a high rock, when a Kingfisher swallowed his minnow bait, and was reeled in, screaming and fluttering wildly. After examining this strange product of his piscatorial skill, and apologizing to it for their mutual mistake, he gave the poor thing its liberty again.

April 9th.

N.Y.

To-day Mr. Redner brought me a Bonaparte's Gull (Larus philadelphiae) which he shot on Poplopen's Pond on April 6th, from a large flock. Another was shot some miles west of Poplopen's on a small lake. The bird's wing was broken and it was still alive when I got it; but it had refused all food. It snapped and bit viciously but not very severely. After showing it to my wife, I killed it.

2,363.

Larus philadelphiae.

14.25 33.75 ♂ ad.

Wing, 10.80; tail, 4.15; culmen, 1.35; gape, 1.90; bare part of tibia, .77; tarsus, 1.50; middle toe and its claw, 1.55; toe alone, 1.27; claw alone, .28. Iris, bright hazel. Bill, entirely jet black. Bare part of tibiae, tarsi & toes, red; nails, brownish-black. Inside of mouth, bright red. Tongue, red, shading through horn-color to blue and black. This specimen was in transition plumage. Its head showed a mottling of plumbeous and white.

Spizella socialis. - To-day I learned something new: the Chipping Sparrow has

1882.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

a real song, not loud, but sweet and Finch-like in addition to the sweet trill which, later, becomes so decidedly monotonous.

April 10th. N.Y.

This morning the ground is white with snow, and the atmosphere chilly; still the Large-billed Acentor sings along the brooks. A single Rusty Grackle was whistling in the woods below the house. Its notes are peculiar and exciting. Had I not seen the bird and known it to be solitary, I should have supposed, from the sounds that this one bird produced, that at least half a dozen were whistling at once.

2,364.

Sialia motacilla.

6.45 10.60 ♂ ad.

Wing, 3.35; tail, 2.25; bill from nostril, .45; gape, .75; tarsus, .91; middle toe and claw, .75.

2,365.

Turdus pallasi.

7.37 11.75 ♂ ad.

Wing, 3.70; tail, 2.85; bill, measured from nostril, .38; gape, .80; tarsus, 1.18; middle toe and claw, .92.

2,366.

Scolecophagus ferrugineus.

9.90 15.15 ♂ ad.

Wing, 4.65; tail, 4.00; culmen, .92; gape, 1.00; tarsus, 1.22; middle toe and claw, 1.24. Iris, light yellow. Bill and feet, black.

2,367.

Sphyrapicus varius.

8.50 15.45 ♂ ad.

Wing, 4.85; tail, 3.20; culmen, .94; gape, 1.04; tarsus, .78; middle toe and claw, .90.

April 13th. N.Y.

Spizella pusilla. - First seen. All three species of Spizellae are numerous to-day.

Dendroica coronata. - One seen.

2,368.

Spizella pusilla.

5.50 8.05 ♂ ad.

Wing, 2.42; tail, 2.42; culmen, .37; gape, .40; tarsus, .74; middle toe and its claw, .69. Iris hazel.

1882.

- 2, 369. *Melospiza fasciata*. Ap. 13. 6.15 8.80 ♂ ad.
Wing, 2.65; tail, 2.65; culmen, .50; gape, .54; tarsus, .86; middle toe and claw, .88.
Iris, hazel. Shot by Ella at "Echo Lake".

April 15th. H. G. M. Y.

Passerella iliaca. — Shot a single bird, which was alone, and proved on dissection to be a female, but was too badly shot for preservation. This is the only one seen for a week or more.

April 16th. H. G. M. Y.

- 2, 370. *Gunco hyemalis*. Shot by E. W. M. 6.30 9.80 ♂ ad.
Wing, 3.00; tail, 2.80; culmen, .45; gape, .51; tarsus, .83; middle toe and claw, .76.
Corvus frugivorus. — Ella found a nest which contained three eggs. The only other nest I have found was not completed.

April 17th. H. G. M. Y.

- 2, 371. *Poocetes gramineus*. 6.50 10.65 ♂ ad.
Wing, 3.10; tail, 2.70; culmen, .44; gape, .50; tarsus, .84; middle toe and claw, .77.
2, 372. *Melospiza fasciata*. Shot by E. W. M. 6.60 8.70 ♂ ad.
Wing, 2.60; tail, 2.75; culmen, .52; gape, .54; tarsus, .87; middle toe and claw, .85.
Poocetes gramineus. — First seen to-day.
Luscalus purpureus. — First seen to-day.

April 19th. H. G. M. Y.

- 2, 373. *Carpodacus purpureus*. Shot by E. W. M. 6.35 9.90 ♀ ad.
Wing and other measurements not taken. Shot by E.
2, 374. *Sialia sialis*. ♂ ad.
Wing, 4.02; tail, 2.72; bill, measured from nostril, .36; gape, .80; tarsus, .79.

1882.

Collecting Number. NAME. DATE. LOCALITY. Length. Stretch. Sex and Age.

middle toe and claw, .86. One of a pair having a nest in a large tomato can in the pear orchard. The Purple Finch was singing when shot. On dissection it was found to be a female.

April 20th. H. G. M. Y.

Spent the afternoon, with Albert, at Poplophen's pond, where I saw four Ducks, a Great Blue Heron (*Ardea herodias*), and an adult White-headed Eagle (*Haliaeetus leucocephalus*). All of these were shot at in turn with my rifle. A Loon (*Colymbus torquatus*) had spent the previous two weeks on the lake, but although my object in going was to try to shoot it, the bird could not be found. Yellow-rumped Warblers (*Dendroica coronata*) were singing by the lake. White-bellied Swallows (*Iachycineta bicolor*) and a few Barn Swallows (*Hirundo erythrogastra*) were flocking about the lake, and were the first Swallows I have seen this season. *Harporhynchus rufus* was also first seen.

April 21st. H. G. M. Y.

Dendroica coronata. — In a flock; singing.
Harporhynchus rufus. Singing as they were yesterday.

April 22nd. H. G. M. Y.

To-night brought the can containing the Bluebirds nest into the house. The female sat close upon her nest while the old rusty can was broken into from behind. Then she flew out, and was killed. After that to my chagrin and remorse five young ones about 3 days old were found, and I put in alcohol as the best way of ending a specially disagreeable operation. No. 2, 274 is the male of this pair.

Ever since the first week in March a female Bluebird has visited our

1882.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

April 22 - Continued.

bedroom window, fluttered against the glass repeatedly and tried to enter, beating hard against the glass. This was kept up almost without interruption, except once for about three days when she was not noticed. She was here as usual this morning, and may possibly be the parent above spoken of. She would alight on the sash, and peck and fly against the window ~~for~~ at intervals for several hours, quite frequently, in the morning before we were up. This was kept up six weeks. Once I arose and drew down the upper sash. Soon the Bluebird came to the stone ledge below the window flew against the glass and tapped, then flew up to the top of the sashes; as soon as she discovered that the window was open, she flew away precipitately with a scared note.

Sturnus notacilla. - In the dusky twilight last night, I saw several large-billed Water Thrushes fly high up over the tree-tops and descend singing deliciously. The Cow Bird (S. auricapillus) shared this habit. I would like to know if S. naevius, the only remaining species of the genus has this habit.

2,275.

Sialia sialis (Mate of No. 2,274.) 6.88 12.15 ♀ ad.
Wing, 3.80; tail, 2.68; bill, measured from nostril, .32; gape, .77; tarsus, .79; middle toe and claw (much worn down), .80. Nest preserved.

April 23rd.

H. G. M. Y.

Mr. E. S. Denton and myself went botanizing. Saw the first White-throated Sparrow (Zonotrichia albicollis) that I have seen. See p. 300.

1882

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

April 24th.

H. G. M. Y.

2,376.

Colaptes auratus.

12.65 20.25 ♂ ad.

Wing, 6.05; tail, 4.85; culmen, 1.45; gape, 1.60; tarsus, 1.10; middle toe and claw, 1.28. Iris, dark red.

April 25th.

H. G. M. Y.

2,377.

Harporhynchus rufus.

11.50 13.50 ♂ ad.

Wing, 4.05; tail, 5.25; bill, measured from nostril, .82; gape, 1.40; tarsus, 1.30; middle toe and claw, 1.15. Iris, yellow.

2,378.

Sialia sialis.

6.90 12.35 ♂ ad.

Wing, 3.77; tail, 2.80; bill, measured from nostril, .38; gape, .80; tarsus, .80; middle toe and claw, .86.

2,379.

Sialia sialis.

7.00 12.35 ♀ ad.

Wing, 3.75; tail, 2.55; bill, measured from nostril, .39; gape, .84; tarsus, .85; middle toe and claw, .88. This female was shot with its mate on Cousook Island, and bore unmistakable signs of maternity. Its nest could not be found. It measures larger than ♂.

2,380.

Agelaius phoeniceus.

9.40 15.40 ♂ ad.

Wing, 4.75; tail, 3.90; culmen, .88; gape, .88 (same); tarsus, 1.15; middle toe and claw, 1.15 (same). Iris very dark. Bill and feet, black.

April 26th.

H. G. M. Y.

2,381.

Agelaius phoeniceus (Shot by E. M. M.)

9.50 15.35 ♂ ad.

Wing, 4.85; tail, 3.95; culmen, .88; gape, .90; tarsus, 1.17; middle toe and claw, 1.13. Iris very dark brown. Bill and feet, glossy black. Shot by Ella in the marsh above the Lake.

April 27.

H. G. M. Y.

Spent the day botanizing, in company with Mr. Edward S. Denton, of West Point.

1882.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

We found the first wild ginger in bloom, and a large bed of the exquisite white blossoms of the blood-root (Sanguinaria) and numerous other flowers.

On the mountain we saw a White-headed Eagle (Haliaeetus leucocephalus).

Dendroica palmarum. - First seen.

Regulus calendula. - First seen. Both in "Stony Lonesome", where we went to look at the Rhododendrons whose buds were well expanded for the season, although a late one, as seen by the late coming of both birds and flowers. Several Ruffed Grouse (Bonasa umbellus) were seen singly.

April 28th.

H. E. M.

2,382. Ortyx virginiana 9.40 — ♂ ad.

Wing, 4.25; tail, 2.55; culmen, .58; gape, .62; tarsus, 1.21; middle toe and claw, 1.60; toe alone, 1.15; claw (grown long by confinement), .45. This Quail was caught near Poplopen's Pond, and kept in a cage since its capture last fall by Mr. Jerome Denna. It died and he gave it to me. One wing was clipped, and could not be measured.

2,383. Liurus motacilla (Shot by E. W. M.) 6.50 10.55 ♂ ad.
Wing, 3.30; tail, 2.23; bill, measured from nostril, .43; gape, .73; tarsus, .93; middle toe and claw, .50.

2,384. Dendroica coronata (E. W. M.) 5.90 9.30 ♂ ad.
Wing, 3.00; tail, 2.52; bill from nostril, .29; gape, .55; tarsus, .72; middle toe and claw, .65.

2,385. Minotilta varia (E. W. M.) 5.48 8.90 ♂ ad.
Wing, 2.85; tail, 2.15; bill from nostril, .39

1882.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

gape, .65; tarsus, .72; middle toe and claw, .71. The first one seen this season.

2,386. Regulus calendula (E. W. M.) 4.52 7.35 — ad.
Wing, 2.35; tail, 1.80; nostril, .26; gape, .45; tarsus, .76; middle toe and claw, .56.

2,387. Dendroica palmarum (E. W. M.) 5.30 8.40 ♀ ad.
Wing, 2.50; tail, 2.17; bill, measured from nostril, .28; gape, .53; tarsus, .83; middle toe and claw, .64.

2,388. Zonotrichia albicollis (E. W. M.) 6.98 9.60 ♂ ad.
Wing, 2.88; tail, 3.00; culmen, .51; gape, .56; tarsus, .88; middle toe and claw, .93.

2,389. Picus villosus (Shot by E. W. M.) 9.50 16.00 ♂ ad.
Wing, 4.65; tail, 3.10; culmen, 1.23; gape, 1.30; tarsus, .88; middle toe and claw, .97.
Minotilta varia. - First seen.

Pipilo erythrophthalmus. - First seen.

Pandion haliaetus carolinensis. - One alighted in a tree beside the lake, and stayed about there a long while. A Red Salamander caught in Cro' Nest cove.

April 29th. H. E. M.

Pandion haliaetus carolinensis. - Seen at Bell's pond.

Troglodytes domesticus. - First seen.

Chaetura pelagica. - First seen by us; but Mr. Edward S. Denton is positive that he saw a number of them yesterday at West Point.

Tringoides macularius. - First seen. One at Elder Lake.

2,390. Turdus pallasi (E. W. M.) 7.25 11.40 ♀ ad.
Wing, 3.60; tail, 2.90; bill, measured from nostril, .35; gape, .85; tarsus, 1.18; middle toe and claw, .90.

2,391. Astragalinus tristis (E. W. M.) 5.05 9.00 ♂ ad.
Wing, 2.85; tail, 2.05; culmen, .42; gape, .42; tarsus, .52; middle toe and claw, .57. Iris, dark hazel.

1882.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
April 29th — Continued.						
2,392.	<i>Harporhynchus rufus</i> (E.W.M.)	11.50	13.65	♂ ad.		
	Wing, 4.25; tail, 5.35; bill, measured from nostril, .80; gape, 1.30; tarsus, 1.35; middle toe and claw, 1.21. Iris, yellow. Two white feathers in crown.					
2,393.	<i>Harporhynchus rufus</i> (E.W.M.)	11.60	13.70	♂ ad.		
	Wing, 4.15; tail, 5.40; bill from nostril, .83; gape, 1.38; tarsus, 1.40; middle toe and claw, 1.28. Iris, yellow.					
2,394.	<i>Picus villosus</i> (Shot by E.W.M.)	9.35	15.15	♀ ad.		
	Wing, 4.55; tail, 3.20; culmen, 1.10; gape, 1.19; tarsus, .83; middle toe and its claw, .88.					
2,395.	<i>Molothrus ater</i> (Shot by E.W.M.)	8.15	13.75	♂ ad.		
	Wing, 4.40; tail, 3.20; culmen, .72; gape, .75; tarsus, 1.05; middle toe and claw, 1.02. Iris dark hazel.					
2,396.	<i>Pipilo erythrophthalmus</i> (E.W.M.)	8.75	11.62	♂ ad.		
	Wing, 3.50; tail, 4.10; culmen, .55; gape, .72; tarsus, 1.10; middle toe and claw, 1.05. Sides, dark red. Bill, black. Tarsi and toes, pale brown.					

April 30th. H.G.M.

Hirundo erythrogastra. — This species and the White-bellied Swallow were both seen the latter in large numbers, at Poplopau's Pond, on April 20th. To-day a single Barn Swallow was seen on the salt marsh, the first one I have seen since the 20th.

Mimus carolinensis. — First seen. A single one by Consook Marsh.

The season is late. Several Warblers and other birds, which usually come to us in April have not yet arrived. The

1882.

first shad-blossoms were gathered to-day.

May 1st. H.G.M.

Vireo solitarius. — First seen. One bird, which I shot but mutilated so as to be worthless.

The Goldfishes (*Crassius auratus*) made their first appearance along shore in Echo Lake. They will probably soon begin spawning.

Hyla pickerinii is piping about his last. Toads are beginning to spawn.

Herrings are running well in the Hudson, and so are Shad.

2,397. *Zonotrichia albicollis* (E.W.M.) 7.00 9.60 ♂ ad.

Wing, 2.90; tail, 3.00; culmen, .53; gape, .56; tarsus, .90; middle toe and claw, .95. Iris hazel.

Antrostomus vociferus. — First seen, vide E.P.Roe.

May 2d. H.G.M.

Empidonax minimus. — First seen.

Tyrannus carolinensis. — First seen.

Setophaga ruticilla. — First seen.

Vireo solitarius. — First seen.

May 3d. H.G.M.

Lurdus mustelinus. — First seen. On Cro' Nest.

Dendroica aestiva. — First seen. Two on Woodna Creek.

Lurdus migratorius. — While walking with Rev. E. P. Roe over his fruit-farm at Cornwall, we found the first Robin's nest, containing 3 eggs.

A large Mussel as big as a good-sized clam was caught alive in Woodna Creek. Also a big Wood Terripiu (*Glyptemys insculpta*) was caught. Both were brought home for Master Elting Roe to put in his father's brook.

A flock of *Quiscalus purpureus* was seen.

Two Great White Egrets (*Herodias alba egretta*) were seen this morning in Woodna Creek.

1882.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	May 3d — Continued.					
	by Mr. Willets; but we did not find them in the afternoon.					
	A Red-tailed Hawk (<u>Buteo borealis</u>) was seen on Mr. Roe's place.					
2,398.	<u>Dendroica aestiva</u> .			5.05	7.75	♀ ad.
	Wing, 2.38; tail, 1.90; bill, measured from nostril, .33; gape, .56; tarsus, .65; middle toe and claw, .57.					
	May 4th. H.S.M.Y.					
	<u>Icterus galbula</u> . — First seen.					
	<u>Cyanospiza cyanea</u> . — First seen.					
	<u>Dendroica pennsylvanica</u> . — First seen.					
	<u>Vireo noveboracensis</u> . — First seen.					
	<u>Vireo gilvus</u> . — First seen.					
2,399.	<u>Cyanospiza cyanea</u> (E.W.M.)			5.75	8.85	♂ ad.
	Wing, 2.70; tail, 2.23; culmen, .40; gape, .49; tarsus, .70; middle toe and claw, .75.					
2,400.	<u>Vireo gilvus</u> . (Shot by E.W.M.)			5.90	9.25	♂ ad.
	Wing, 2.90; tail, 2.30; bill, measured from nostril, .34; gape, .72; tarsus, .70; middle toe and claw, .57. Iris, hazel. The first one seen.					
2,401.	<u>Vireo noveboracensis</u> (E.W.M.)			5.35	8.25	♂ ad.
	Wing, 2.48; tail, 2.07; bill, measured from nostril, .30; gape, .64; tarsus, .78; middle toe and claw, .57; Iris, white. First one seen.					
2,402.	<u>Dendroica pennsylvanica</u> (E.W.M.)			5.10	7.90	♂ ad.
	Wing, 2.45; tail, 1.95; bill from nostril, .30; gape, .64; tarsus, .67; middle toe and claw, .55. The first one seen this spring.					
2,403.	<u>Pipilo erythrophthalmus</u> (E.W.M.)			8.40	11.45	♂ ad.
	Wing, 3.47; tail, 3.90; culmen, .57; gape, .75; tarsus, 1.13; middle toe and claw, 1.03. Iris, red.					
	<u>Dendroica virens</u> and <u>D. caerulea</u> have not yet been seen, nor <u>D. pinus</u> .					

1882.

May 5th. H.S.M.Y.
To-day Mr. E.S. Denton and myself went up the south ravine to Cro' Nest and down the East ravine to the Hudson. The locality of the long-sought White-headed Eagle's nest (Haliaeetus leucocephalus) was at length discovered, and the young heard scream-ing on the ledge while being fed. There were two of them, and they were able to fly a couple of hundred yards with some difficulty, and return to the nest. The mystery of the Eagle's nest is solved, although we are still in doubt as to the precise spot on the ledge where the nest is placed. We hope to get the eggs next winter. Trochilus colubris. — First seen by my wife on the Japonica bush by the house.

May 6th. H.S.M.Y.
A Fish Hawk (Pandion haliaetus carolinensis) was caught by the fishermen in a herring net. They took the poor fellow out alive and finally let him go free after amusing themselves by tying a string to its legs and allowing it to fly a short distance and then pulling it back. Then they devised another way of amusing themselves. They tied a stone to the cord. The Osprey was barely able to raise this from the ground, and flew slowly along until pulled down and let go by one of the kinder hearted fishermen. We think he will look twice before he dives for caught herring again!

Eliza and I drove to the top of Cro' Nest to-day. There was a Pheasant's

1882.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

May 6th - Continued.

(*Bonasa umbellus*) nest there which Mr. Denton and I found yesterday in this wise; Climbing the last ascent to the summit of Crow Nest I saw a species of *Archid* growing on a damp ledge, and, stooping, called Mr. D. to come and look at the plant. As he started to do so, a hen Ruffed Grouse flew with a terrible whirr past my face and whizzed by Mr. D. like a cannon shot. We were both scared. We found her nest under a pointed projecting stone within a yard of where I was stooping down to look at the *Archid* plant. It was perfectly concealed, and a lot of dry oak-leaves fell upon the eggs as the old bird rushed out, so as to cover them. We counted the eggs. There were eleven. Then we left the nest, but when we visited it ~~to-day~~ this afternoon the mother Grouse had not returned and the eggs were stone cold so we took them all.

Furdus fuscescens. - First seen by E. at the head of the southern ravine, where we went to see the view.

2,404. *Furdus fuscescens* (E.W.M.) 7:80 12:63 ♂ ad. Wing, 4.05; tail, 3.20; bill from nostril, .39; gape, .87; tarsus, 1.15; middle toe and claw, .89. Iris, hazel.

Eupidonax minimus. - Now abundant.

Eupidonax trailii. - First seen.

Buteo pennsylvanicus. - First seen. None seen last winter. All kinds of Hawks have been rare this spring.

May 8th N.Y.

Junco hyemalis. Here yet, but scarce.

1882.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

May 9th. N.Y.

E., myself and Mr. E. S. Denton spent the day botanizing. Found a new Club-moss in the lower ravine of Cro' Nest, and we all heard the Eaglets (*Haliaeetus*).

E. shot *Myiodiocetes nitratus* at the bottom of their cliff. She shot three other birds with the little gun without a miss.

Dendroica virens. - First seen. Singing.

Parula americana. - First seen. Singing.

Dendroica caerulescens. - First seen. Singing.

Sciurus aurocapillus. - First seen. Singing.

Geothlypis trichas. - First seen. Singing.

Myiodiocetes nitratus. - First seen. Singing.

Regulus calendula. - Still here.

2,405. *Dendroica virens* (E.W.M.) 5:10 8:00 ♂ ad. Wing, 2.45; tail, 2.05; bill, measured from nostril, .30; gape, .50; tarsus, .72; middle toe and claw, .55. Iris, hazel. Bill and feet, black. The first one seen.

2,406. *Lanius solitarius* (E.W.M.) 5:70 9:40 ♀ ad. Wing, 2.83; tail, 2.12; bill, measured from nostril, .24; gape, .62; tarsus, .76; middle toe and claw, .65.

Measurements of:-

Sciurus aurocapillus (E.W.M.) 6:25 9:80 ♂ ad. Wing, 2.95; tail, 2.25; bill from nostril, .38; gape, .67; tarsus, .88; middle toe and claw, .75.

May 10th N.Y.

Sciurus naevius. - First seen. Singing.

Icterus spurius. - First seen.

2,407. *Icterus spurius* (E.W.M.) 7:35 10:40 ♂ ad. Wing, 3.17; tail, 3.05; culmen, .74; gape, .77; tarsus, .88; middle toe and claw, .88 (same as tarsus).

1882.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	Iris, hazel. Bill, with maxilla and terminal half of mandible, black; basal half of mandible, blue. Tarsi and toes, grayish blue. The first one seen this season.					
	May 11th. H. S. N.Y.					
	<u>Helminthophaga ruficapilla</u> . - <u>First seen</u> .					
	May 12th. H. S. N.Y.					
2,408.	<u>Helminthophaga ruficapilla</u> .	4.85	7.70	♂ ad.		
	Wing, 2.42; tail, 1.95; bill, measured from nostril, .30; gape, .48; tarsus, .70; middle toe and claw, .60.					
	May 13th. H. S. N.Y.					
	<u>Icteria virens</u> . - <u>First seen</u> .					
	<u>Helminthophaga ruficapilla</u> . - <u>First seen</u> .					
	<u>Dendroica blackburniae</u> . - <u>First seen</u> .					
	To-day I gathered the first specimens of our beautiful Showy Oriole (<u>Oriolus Spectabilis</u>). Four were well out, and ten more were nearly out. Two other species were seen in leaf.					
2,409.	<u>Dendroica blackburniae</u> .	5.15	8.10	♀ ad.		
	Wing, 2.60; tail, 2.12; bill, measured from nostril, .29; gape, .51; tarsus, .66; middle toe and claw, .55.					
2,410.	<u>Helminthophaga ruficapilla</u> .	5.50	9.00	♂ ad.		
	Wing, 2.75; tail, 2.15; bill, measured from nostril, .40; gape, .59; tarsus, .73; middle toe and claw, .67.					
2,411.	<u>Geothlypis trichas</u> .	5.35	7.35	♂ ad.		
	Wing, 2.18; tail, 2.23; bill, measured from nostril, .31; gape, .56; culmen, .80; middle toe and claw, .67.					
2,412.	<u>Melospiza fasciata</u> .	6.55	8.75	♂ ad.		
	Wing, 2.54; tail, 2.82; culmen, .53; gape, .58; tarsus, .84; middle toe and claw, .86.					

1882.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	May 15th. H. S. N.Y.					
2,413.	<u>Setophaga ruticilla</u> . (E.M.M.)	5.50	8.15	♂ ad.		
	Wing, 2.65; tail, 2.40; bill, measured from nostril, .30; gape, .55; tarsus, .68; middle toe and claw, .57.					
2,414.	<u>Vireosylva gilva</u> . (E.M.M.)	5.65	8.90	♀ ad.		
	Wing, 2.68; tail, 2.10; bill, measured from nostril, .31; gape, .67; tarsus, .73; middle toe and claw, .58. Iris, hazel.					
	May 16th. H. S. N.Y.					
2,415.	<u>Helminthophaga ruficapilla</u> (E.)	4.65	7.55	♀ ad.		
	Wing, 2.30; tail, 1.75; bill, measured from nostril, .28; gape, .50; tarsus, .65; middle toe and claw, .58. (Shot by E.M.M.)					
2,416.	<u>Dendroica coronata</u> (E.M.M.)	5.60	8.75	♀ ad.		
	Wing, 2.77; tail, 2.30; bill, measured from nostril, .28; gape, .52; tarsus, .73; middle toe and claw, .60.					
2,417.	<u>Mimus carolinensis</u> (E.M.M.)	9.13	11.75	♂ ad.		
	Wing, 3.55; tail, 4.10; bill, measured from nostril, .45; gape, 1.60; tarsus, 1.18; middle toe and claw, 1.04. Iris, dark hazel.					
2,418.	<u>Agelaius phoeniceus</u> (E.M.M.)	7.95	12.50	♀ ad.		
	Wing, 3.90; tail, 3.20; culmen, .70; gape, .75; tarsus, 1.05; middle toe and claw, 1.60.					
2,419.	<u>Agelaius phoeniceus</u> (E.M.M.)	9.50	15.40	♂ ad.		
	Wing, 4.90; tail, 4.00; culmen, .90; gape, .93; tarsus, 1.15; middle toe and claw, 1.10. Ella shot these two, a mated pair, on the border of Cousook Marsh.					
	<u>Turdus migratorius</u> . - A nest in the silver maple in front of the house has 5 eggs.					
	<u>Mimus carolinensis</u> . - Male accidentally shot of a pair having a nest and one egg beside the Cousook Marsh. One by the house probably has eggs, but not sitting yet.					

1882.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	May 17th. N.Y. N.Y.					
2,420.	<i>Amelodia ludoviciana</i> (E.W.M.)	8.10	13.00	♂ ad.		
	Wing, 4.00; tail, 3.10; culmen, .69; gape, .74; tarsus, .90; middle toe and claw, .97.					
2,421.	<i>Otocoris colubris</i> (E.W.M.)	3.77	4.15	♂ ad.		
	Wing, 1.50; tail, 1.22; culmen to frontal feathers, .62; gape, .80; tarsus, .18; middle toe and claw, .23.					
2,422.	<i>Botaurus minor</i> .	25.00	37.60	♀ ad.		
	Wing, 10.10; tail, 3.45; culmen, 2.80; gape, 3.75; tarsus, 3.40; bare part of tibia, 1.43; middle toe and claw, 3.75; claw alone, .79. Length from extremity of bill to that of outstretched toes, 34.40. Iris, yellow, externally granulated with reddish-brown. Bare part of tibiae, tarsi and toes, yellowish-green. Bill, brownish-black above; residue varied with yellow, and pale green and blue.					
	Mr. Parry stopped on the road, while I was talking to the gardener, and informed me that there was a "Crane" down the road. I was kept for some time, but at length took my gun and hunted up the bird which proved to be the above specimen. I found it in a swamp some distance from where Mr. P. saw it, and brought it down with a "B" shot as he flew from far across the swamp, a charge of 4 shot having previously failed to reach it at that distance. Its wing was broken, but it had received no other injury. When I approached it sprang furiously at me, flattened out the feathers upon its neck, and looked for-					

1882.

-midable. Before I got to it I noticed that it lay upon the ground with its bill pointing straight up in the air. This was the position it assumed after I got it home, where I took it to show to Grandfather Carswell and my wife. This is the first seen this year.
Myiarchus crinitus. — first seen.
Turdus swainsoni. — first seen.

May 18th. N.Y. N.Y.
Zonotrichia leucophrys. — Grandfather Carswell had gone with me to visit Grandfather Mearns (the former gentleman over 80, and the latter more than 95 years old), and we reached our gate, when I saw a specimen of the above species, the first seen this year.

May 20th. N.Y. N.Y.
 Louis A. Zelega came up last night to get some Heron's Eggs and birds. We visited the Heronry to-day, and saw during the day the following species:—
Turdus migratorius.
Hylocichla mustelina.
Hylocichla fuscescens.
Hylocichla viciae. first seen.
Hylocichla ustulata swainsoni.
Galeoscoptes carolinensis.
Harporhynchus rufus.
Sialia sialis.
Parus atricapillus.
Sitta carolinensis.
Troglodytes aedon.

1882.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.Helmatodytes palustris. - First seen.Mniotilta varia. -Parula americana.Helminthophaga chrysoptera. - First seen.Helminthophaga ruficapilla.Dendroica aestiva. - Singing.Dendroica virens. - Singing.Dendroica caerulescens. - Singing.Dendroica coronata. - Still here.Dendroica pennsylvanica.Dendroica castanea. - First seen.Dendroica blackburniae. -Siurus naevius. - Constitution Marsh.Siurus auricapillus.Siurus ludovicianus.Geothlypis trichas.Icteria virens.Myiodynastes nigratus.Setophaga ruticilla.Vireo solitarius.Vireosylva gilva.Ampelis cedrorum.Hirundo erythrogastra.Tachycineta bicolor.Cotyle riparia. - First seen.Petrochelidon lunifrons. - First seen.Astragalinus tristis.Melospiza fasciata.Melospiza palustris. - Constitution Island.Spizella socialis.Spizella pusilla.Zonotrichia leucophrys. - Only one seen; shot upon the ground amongst some currant bushes.Passerina cyanea. - One seen.Pipilo erythrophthalmus.Zamelodia ludoviciana. - Quite a

1882.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

number stay in our cherry-trees, and eat the blossoms.

Passer domesticus.Agelaius phoeniceus.Molothrus ater.Icterus galbula.Icterus spurius.Corvus rugivorus.Cyanocitta cristata.Sayornis fusca.Euphonia minima.Tyrannus carolinensis. First since May 2.Myiarchus crinitus.Trochilus colubris.Chaetura pelagica.Chordeiles popetue. - First seen.Picus pubescens.Colaptes auratus.Ceryle alcyon.Haliaeetus leucocephalus.Bonasa umbellus. - Constitution Island.Zenaidura macroura.Tringoides macularius.

Nyctiardea grisea naevia. - Only a few pairs were about the heronry. Just a week ago (May 13th) a young man by name of Pearson who attends a school in Sing Sing came up to the heronry and took 56 eggs of this species, and a set of 4 eggs of the Green Heron (Ardea virescens). We found two more youngsters in the swamp from the same school, and not an egg could be found in any of the nests. All had been robbed; and I fear that the heronry will be broken up entirely.

1882.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	<u>May 21.</u>	<u>N.Y.</u>				
	<u>Dendroica striata.</u>	<u>First seen.</u>				
	<u>Dendroica maculata.</u>	<u>First seen.</u>				
	<u>Helminthophaga peregrina.</u>	<u>First seen.</u>				
	Two were noticed in an apple-tree by the house.					
	<u>May 22.</u>	<u>N.Y.</u>				
	<u>Myiodytes canadensis.</u>	<u>First seen.</u>				
	<u>Pyrranga rubra.</u>	<u>First seen.</u>	This is very late. There is a notice of its arrival in Central Park in April in "Forest & Stream". None have been seen about here, although carefully sought for. A few <u>Pue Anemones</u> and <u>Hepaticas</u> are still in blossom. <u>Purple Azaleas</u> are just coming in blossom.			
	<u>May 19th.</u>	<u>N.Y.</u>				
2,423.	<u>Zamelodia ludoviciana (E.)</u>	8.20	13.50	♂ ad.		
	Wing, 4.25; tail, 3.10; culmen, .69; gape, .75; tarsus, .90; middle toe and claw, .90.					
	Ella shot two fine males in the cherry trees. Berega took one of them.					
	<u>May 20th.</u>	<u>N.Y.</u>				
2,424.	<u>Zonotrichia leucophrys.</u>	7.25	10.25	♂ ad.		
	Wing, 3.10; tail, 3.30; culmen, .49; gape, .54; tarsus, .95; middle toe and claw, .90.					
	Irides, hazel. Bill, pale reddish-brown; extreme tip of maxilla, black. Tarsi and toes, pinky-brown, darkest on toes. This bird was shot in the same spot where the other one was seen, and after it was shot none others were seen.					
2,425.	<u>Helminthophaga chrysoptera.</u>	5.10	7.95	♂ ad.		
	Wing, 2.50; tail, 2.05; bill, measured from nostril, .34; gape, .57; tarsus, .65; middle					

1882.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
2,426.	<u>Icterus galbula (E.W.M.)</u>	7.75	12.15	♂ ad.		
	Wing, 3.75; tail, 3.12; culmen, .77; gape, .84; tarsus, .90; middle toe and claw, .86. Iris, dark hazel. Bill, with sides of the maxilla, and mandible, pale blue; maxilla above, black. Tarsi and toes, grayish-blue.					
2,427.	<u>Tringoides macularius.</u>	8.00	13.50	♂ ad.		
	Wing, 4.05; tail, 2.10; culmen, .95; gape, 1.10; tarsus, 1.00; bare part of tibia, .55; middle toe and claw, 1.05.					
2,428.	<u>Tringoides macularius.</u>	8.10	14.15	♀ ad.		
	Wing, 4.35; tail, 2.20; culmen, .98; gape, 1.10; tarsus, .99; bare part of tibia, .40; middle toe and claw, .95.					
2,429.	<u>Icterus spurius.</u>	7.15	10.45	♂ ad.		
	Wing, 3.15; tail, 3.00; culmen, .70; gape, .74; tarsus, .85; middle toe and claw, .85.					
	Note. - A few days since, Mr. Louis A. Ber- -ega saw a Sparrow Hawk (<u>Falco sparverius</u>) fly up after a bass-ball that had been thrown in the air, and hover over it a moment when it had reached the highest point.					
	<u>May 23rd.</u>	<u>N.Y.</u>				
2,430.	<u>Sinus auricapillus (E.W.M.)</u>	6.35	9.75	♀ ad.		
	Wing, 2.90; tail, 2.20; bill, measured from nostril, .35; gape, .67; tarsus, .88; middle toe and claw, .72.					
	<u>Ceryle alcyon.</u> - This afternoon drove to a sand-bank on Mr. J. P. Morgan's place with Ella, to dig out a Kingfisher's hole or nest. It ran perpendicularly inward for two feet. I dug downward from above upon					

1882.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

May 23 — Continued.
the first angle of the hole. Found the nest, a globular excavation, at the end of the second reach. It was about four feet from the entrance, and within half that distance of the side of the bank. It contained only four eggs, three of which were got out whole. While digging out the eggs, the old bird returned, flew all about me and perched upon a tree near by. Then it flew away, and did not return.
Ceontopus virens. — First seen.
Lanius flavifrons. — First seen.

May 24. N.E.M.

- 2,431. Dendroica catullescens (E.) 5.15 7.75 ♀ ad.
Wing, 2.38; tail, 1.95; bill, measured from nostril, .28; gape, .54; tarsus, .75; middle toe and claw, .62.
- 2,432. Geothlypis philadelphia. 5.50 8.00 ♂ ad.
Wing, 2.40; tail, 2.08; bill, measured from nostril, .30; gape, .59; tarsus, .81; middle toe and claw, .76. Iris, hazel. Bill above, black; below, flesh-color. Tarsi and toes, brownish flesh-color.
- 2,433. Myiodiotes canadensis (E.W.M.) 5.75 8.55 ♂ ad.
Wing, 2.70; tail, 2.45; bill, measured from nostril, .29; gape, .57; tarsus, .74; middle toe and claw, .60.
- 2,434. Parula americana (E.W.M.) 4.90 7.80 ♂ ad.
Wing, 2.30; tail, 1.95; bill, measured from nostril, .30; gape, .50; tarsus, .68; middle toe and claw, .52.
- 2,435. Geothlypis trichas (E.W.M.) 5.00 7.05 ♂ ad.
Wing, 2.07; tail, 1.85; bill, measured from nostril, .30; gape, .53; tarsus, .80;

1882.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

- 2,436. Geothlypis trichas (E.W.M.) 5.05 6.95 ♂ ad.
Wing, 2.05; tail, 1.95; bill, measured from nostril, .32; gape, .57; tarsus, .80; middle toe and claw, .70. Iris, hazel.
- 2,437. Colaptes auratus. 13.05 20.60 ♂ ad.
Wing, 6.25; tail, 4.75; culmen, 1.54; gape, 1.70; tarsus, 1.20; middle toe and claw, 1.30. Made a good wing-shot.
- 2,438. Coccyzus erythrophthalmus (E.) 11.65 16.00 ♀ ad.
Wing, 5.25; tail, 6.15; culmen, .90; gape, 1.27; tarsus, 1.02; middle toe and claw, .99.
- 2,439. Turdus aliciae (E.) 7.60 12.65 ♂ ad.
Wing, 3.90; tail, 3.00; bill, measured from nostril, .38; gape, .86; tarsus, 1.25; middle toe and claw, .90; iris, hazel. Bill, with terminal $\frac{1}{2}$ of mandible and maxilla brownish-black; basal $\frac{1}{2}$ of mandible, yellow. Tarsi and toes, brown, darkest on feet.

Geothlypis philadelphia. — Only one, the First Seen, in a swampy watercourse north of Pell's pond.

Coccyzus erythrophthalmus. I do not remember having seen or heard this bird before this spring. It must have been late in coming, and scarce.

Bonasa umbellus. — One at Pell's Pond.

May 26th. N.E.M.

Vireosylva olivacea. — First seen.

Empidonax flaviventris. — First seen.

Myiodiotes pusillus. — First seen. One shot behind the stable.

Helminthophaga penins. — First seen. A female behind the stable in birch swamp.

1882.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
May 26th — Continued.						
2,440.	<i>Icteria virens</i> .	6.90	9.50	♀ ad.		
	Wing, 2.90; tail, 2.95; bill, measured from nostril, .40; gape, .73; tarsus, 1.00; middle toe and claw, .82.					
2,441.	<i>Dendroica striata</i> .	5.45	8.85	♂ ad.		
	Wing, 2.85; tail, 2.12; bill, measured from nostril, .29; gape, .54; tarsus, .75; middle toe and claw, .66.					
2,442.	<i>Parula americana</i> .	4.80	7.55	♂ ad.		
	Wing, 2.45; tail, 1.78; bill, measured from nostril, .31; gape, .53; tarsus, .62; middle toe and claw, .54.					
2,443.	<i>Dendroica maculosa</i> (E.)	5.10	7.80	♂ ad.		
	Wing, 2.40; tail, 2.10; bill, measured from nostril, .28; gape, .52; tarsus, .73; middle toe and claw, .56.					
2,444.	<i>Dendroica maculosa</i> (E.)	5.00	7.40	♀ ad.		
	Wing, 2.25; tail, 2.00; bill, measured from nostril, .28; gape, .50; tarsus, .70; middle toe and claw, .53.					
2,445.	<i>Myiodiocetes pusillus</i> .	5.00	6.72	♂ ad.		
	Wing, 2.05; tail, 2.07; nostril, .26; gape, .50; tarsus, .70; middle toe and claw, .55.					
2,446.	<i>Helminthophaga ruficapilla</i> (E.)	5.65	8.75	♀ ad.		
	Wing, 2.65; tail, 2.15; bill, measured from nostril, .40; gape, .65; tarsus, .72; middle toe and claw, .67.					
2,447.	<i>Helminthophaga ruficapilla</i> (E.)	4.75	7.50	♀ ad.		
	Wing, 2.30; tail, 1.80; bill, measured from nostril, .27; gape, .48; tarsus, .65; middle toe and claw, .54.					
2,448.	<i>Myiodiocetes canadensis</i> (E.)	5.50	8.15	♂ ad.		
	Wing, 2.54; tail, 2.25; bill, measured from nostril, .27; gape, .52; tarsus, .76; middle toe and claw, .56.					

1882.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
2,449.	<i>Vireosylva olivacea</i> (E.)	6.10	9.90	♀ ad.		
	Wing, 3.05; tail, 2.15; bill, measured from nostril, .35; gape, .73; tarsus, .74; middle toe and claw, .60.					
2,450.	<i>Lurdus aliciae</i> (E.)	7.85	12.65	♂ ad.		
	Wing, 4.15; tail, 3.10; bill, measured from nostril, .38; gape, .85; tarsus, 1.23; middle toe and claw, .90.					

May 27th. H. & M.
Contopus borealis. — First seen

May 30th. H. & M.
Dendroica castanea. — Still present.
Myiodiocetes canadensis. — Still numerous.
Geothlypis philadelphia. — One seen (♂).
Lamelodia ludoviciana. — Saw one female.
 2,451. *Passerina cyanea*. Long P. 5.60 8.75 ♂ ad.
 Wing, 2.60; tail, 2.17; culmen, .44; gape, .50; tarsus, .69; middle toe and claw, .67.

May 31st, H. & M.
Icterus galbula. — Three or four female Baltimore have been gathering the colored threads & worsteds that I have fastened to shrubs in front of the house. The males came about too, and sang; but none of them took any part in building the nests. I went to one tree and watched the female bird weaving. She dived head first into the nest, and worked with such energy as to shake the whole branch on which the nest was built. The male did not assist by bringing materials or by weaving them when brought. Females from other nests came and tried to

1882.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

steal threads from this one, but were driven off by the angry proprietress.

June 7th, 1882. H.E.M.Y.

Drove around the Eagle Valley road to West Point with Ella. We found a Wood Thrush's nest and several rare plants, among them the Naked Broom-Rope (*Aphyllon uniflorum*) and a number of a handsome Orchid, *Aplectrum hyemale* ("Adam and Eve").

When I reached home Dr. Chace awaited me and wished me to go at once, without supper, to see a woman in labor ten miles back in the mountains beyond Queensboro, which I did. On the way I saw a fine male Snow-bird (*Junco hyemalis*) beside the school-house at Queensboro, the first I have ever seen here in summer. We drove a mile out of the right way and had to come back, making our trip 22 miles, or altogether about 30 miles driving for me, from 4 P.M. to midnight.

June 12th. H.E.M.Y.

Icterus galbula. - Took a nest and set of 5 eggs from a pear-tree in the garden.

Icterus spurius. - Took a set of 4 eggs and nest from a pear-tree in the garden. Both sets were slightly incubated.

Troglodytes domesticus. - A House Wren built a nest in a out-house, entering the door through an aperture. The door was opened and left open. The bird left, but came back and carried

1882.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

off all the sticks. No eggs had been deposited.

June 17th. H.F., N.Y.

Drove to Iona Island. At Fort Moutgomery I saw a Rough-winged Swallow (*Stelgidopteryx serripennis*) sitting upon a telegraph wire. Seeing a couple of holes in a bank, I examined and from one of them another one flew. Both perched together upon the telegraph wire after flying around me a few times; but before I could get hold of my gun both had flown away, and did not soon return. On my return I dug a little way into the bank, and then could easily reach the nest, which was composed entirely of dry grass, with one small piece of paper and one of leaf. It contained four eggs, and the female contained another one, which would shortly have been deposited. I then shot both of the old birds.

2,452 *Stelgidopteryx serripennis* 5:50 12:05 ♂ ad.
Wing, 4:12; tail, 2:18; bill, measured from nostril, .20; gape, .58; tarsus, .42; middle toe and its claw, .60. Iris, hazel. Bill, all black. Left tarsus and toes, black; but the right tarsus was brownish-black; metatarsus, flesh-color; toes and claws black.

2,453 *Stelgidopteryx serripennis* 5:55 11:80 ♀ ad.
Wing, 4:03; tail, 2:10; bill, measured from nostril, .21; gape, .59; tarsus, .45; middle toe and claw, .60. Iris, hazel. Bill & feet, black.

1882.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
		August 5th.	N.Y.			
	In a rocky hillside grown up with young timber we found a number of young and old Whippoorwills (<i>Antrostomus vociferus</i>). We shot two.					
2,454.	<i>Antrostomus vociferus</i>					♀ juv.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
		August 11th.	N.Y.			
2,455.	<i>Ceryle alcyon</i> .			12.65	21.25	♂ ad.
	Wing, 6.00; tail, 4.00; culmen, 2.20; gape, 2.65; tarsus, .46; middle toe and claw, .87; claw alone, .30. Iris, dark hazel. Shot by Ella W. Mearns at Echo Lake.					

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
		August 31st.	N.Y.			
2,456.	<i>Merula erythrogastra</i> .					♀ juv.
	This bird flew against the chimney of our house and broke its wing.					

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
		Sept. 7th.	N.Y.			
2,457.	<i>Colaptes auratus</i> .					♀ ad.
	Moulting. Shot it on wing at very long range, with No. 4 shot, near Pell's Pond.					

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
		Sept. 23rd.	N.Y.			
	Spent the early part of the day in procuring specimens of fishes, at Poplopen's Pond. We took the following:					
	<i>Eupomotis aureus</i> (Common Sun Fish).					
	<i>Catostomus commersonii</i> (Common Sucker).					
	<i>Erismyzon sucetta</i> (Chub Sucker).					
	<i>Perca americana</i> (Common Yellow Perch).					
	<i>Esox reticulatus</i> (Common Eastern Pickerel).					
	<i>Amiurus catus</i> (Small Cat Fish).					
	The day was very stormy, as had been the day previous. Six inches of rain fell in 24 hours. All of the					

1882.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	lakes and streams were enormously swollen. The road beside Long Pond was covered 2½ feet deep with water, the pond itself having risen to that level. All of the boats were washed over the dam at Poplopen's Pond. We had to go back to Clark's, yoke up the oxen to the sled, and haul over Moses Clark's boat. On the way over we put up a flock of Quails (<i>Ortyx virginiana</i>) and I discovered four flowers that were new to me and my wife, viz:					

Aster Novae-Angliae (New England Aster).
Gentiana crinita (Blue-fringed Gentian).
Epiphegus virginiana (Beech-drops).
Diplopappus umbellatus

One of the yellow Perch caught had a duplicate dorsal fin. I skinned and mounted it, and the largest Yellow Perch caught, which weighed about two pounds. A Wood Pewee (*Contopus virens*) came to the house during the storm, and stayed on the plum-tree, beside the kitchen door.

September 27th. N.Y.
Chaetura pelagica. - A few were seen.
Contopus virens has not been seen since the 23d, and has probably departed.

September 29th. N.Y.
 An interesting animal (*Homo sapiens*, ♀ juvenile) was captured in my wife's bedroom this morning, at 9.30 ^{o'clock} A.M. A notice of it will doubtless occur in this journal, further on. Edgar A. Mearns.

1882.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.October 1st. N.Y.

Bonasa umbellus. — To-day, shortly after morning church service, I received a mysterious-looking covered basket, which contained a full-grown Ruffed Grouse alive. It was bird of the year. The letter read as follows: —

"Parsonage. Oct. 1st, 1882.
My Dear Doctor!

Excuse me for sending you this feathered Burglar for confinement or refinement just as you may deem him deserving.

He broke into our house this morning whilst we were all at church.

We have not as yet discovered any other loss than a large pane of glass.

As we have no means of keeping life in him, and do not feel quite injured enough to take the life out of him, we resign him or consign him to your Taxidermical skill.

With congratulations to self, Wife and Daughter. I am

Your Very Truly
P. P. Williams."

The bird in question, after being duly admired, was taken into the orchard, and the basket opened and set upon the ground; but the affrighted creature did not move until I picked it up and tossed it into the air, when it whurred off to the woods with the velocity of a cannon shot.

One frequently hears of its flying

1882.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

against windows. I have known of five or six instances of its having done so, in this place. An explanation of this occurrence has frequently been called for, and a very satisfactory and plausible one is this: The bird, approaching a house, looks towards the window of an otherwise darkened apartment, and sees reflected there an attractive image of the landscape before it, and, mistaking what it sees for the reality, makes towards the supposed spot. If the window-glass yields before its impetuous flight, the bird finds itself a prisoner within the apartment, and, if not, it falls injured to the ground.

2,458. Cyanocitta cristata. — — 3 ad.
Shot Oct. 2nd, 1882. N.H., N.Y.

October 4th. N.H., N.Y.

2,459. Cyanocitta cristata. — — 7 ad.

Note. — Mr. Edward Conway, of 55 Carmine St., N.Y. City, showed me a mounted specimen of the Barn Owl (Aluco flammeus americanus), which was brought to him by Mr. Joseph Wilde, on April 13, 1878, in the flesh. It was shot somewhere near New York City, but just where, I could not ascertain, since Mr. Wilde never came back after his bird, which he said "was killed just outside of New York."

November 6th. N.H., N.Y.

To-day, walked down to see what progress has been made at the rock cut, on the

Parsonage Oct 1st 1882
My dear Doctor:

Excuse me for sending you this feathered Burglar for Confinement or refinement just as you may deem him deserving.

He broke into our house this morning whilst we were all at Church.

We have not as yet discovered any other loss than a large pane of glass.

As we have no means of keeping life in him, and do not feel quite injured enough to take the life out of him, we resign him or consign him to your taxidermic skill.

Best congratulations to Self, Wife and Daughter Sam.

Cours by line
D. J. Williams

1882.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	West Shore Railroad, at Satterlee's Point.					
	Returning home, I flushed a Rabbit and a Woodcock simultaneously, for beneath a pile of brushwood. I saw the Woodcock tumble down amongst some saplings of Ironwood, and watched it sitting there for some time. After dinner I took my gun and walked down there and flushed the bird within a rod of the same spot where I left it. I brought it down, making a good wingshot. I also saw the Rabbit but did not get a shot.					
2,460.	<u>Philohela minor.</u>			10.50	17.65	♂ ad.
	Wing, 4.90; tail, 2.90; culmen, 2.50; gape, 2.30; tarsus, 1.21; middle toe, 1.23; claw, .21; middle toe and claw, 1.42. Iris, hazel. Bill & feet, dusky flesh-color.					

November 7th. N.H.M.

To-day, election-day, was spent in fishing on Poplopen's pond, and in hunting on the mountain west of it.

Mr. J. Deuna accompanied me, and together we caught about 60 Pickeral & Perch (Yellow), besides 8 "Mud noses" and 2 Catfishes. The largest Yellow Perch weighed $1\frac{3}{4}$ pounds, and was preserved as a mounted specimen Ortyx Virginianus. - Mr. Deuna saw a couple of Quads in the lane, near Poplopen's Pond, as we were carrying our new boat home to Mr. Clark's. I got ready, put up the birds, three in number, and shot one with my first barrel, but missed with the second barrel.

2,461. Ortyx virginiana.

Wing, 4.35; Tail, 2.60; culmen, .55; bill,

1882.

measured from nostril, .32; gape, .69; tarsus, 1.24; middle toe & claw, 1.34. Iris, hazel. Probably young of the year.

November 16th. N.H.M.

Loxia leucopterus (White-winged Crossbill). This afternoon, Ella & I walked through Mrs. Stephen R. Roe's place, opposite to ours. In a pine tree right in front of the house we saw a female White-winged Crossbill extracting seeds from the cones of Pinus rigida. I came back and got a gun, leaving Ella to watch the birds. She saw three birds at once, and thought there were four, altogether. I gave her the gun and she shot a female. We both saw it drop a few inches and disappear from sight. We could not get sight of the others afterwards. Ella thinks that all were in female dress, as she saw no red. We watched for half an hour, and then fired another shot hoping to dislodge the dead bird; but failing, we returned home disconsolate. It rained during the night. The next forenoon I walked over, and looked under the tree hoping to find that the bird had fallen during the night. It had not, so I got a big ladder at the stable, and went up to the top of the tree, from whence I saw it clinging to a branch, whence it required considerable shaking to dislodge it. The bird, when dry, turned out to be an excellent specimen. Have not seen it there since the winter 1874-75.

2,462.

Loxia leucopterus. Ella shot it. 6.25 10.35 ♀ ad. Wing, 3.30; tail, 2.45; culmen, .65; gape, on right side, .60; left ~~right~~ side, .64; tarsus, .60; middle toe, .46; claw, .34; middle toe and

1882

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.

claw, .75. Irides, hazel. Bill, blackish on maxilla; ^{most of} mandible and sides of maxilla except near tip, horn-color. ~~Bill~~ Tarsi and toes, dusky brown; claws, blackish. Stomach well filled with seeds of Pinus rigida; fat.

November 23rd. H. S. N.Y.

As I rode through Capt. S. R. Roofs grounds, opposite ours, I found an immature male White-winged Crossbill (Loxia leucoptera) lying in the road, between the pine trees in which Ella shot her female on the 16th instant. Probably she had wounded it when she shot the other specimen; and it had flown a little way and fallen. Its neck was injured; but I was not sure that a cat had not done it. Although a week has elapsed, the temperature has been low for this season, and the bird makes a good skin.

The remarkable thing about it is - that it only had one leg. On skinning it I found no trace of a femur. The acetabulum was a round hole; and the pubic bone was abortive upon that side. The bird was somewhat emaciated; but no signs of inflammatory action were found about the pelvis. The skin showed a slight fold ^{and sac} at the point opposite the acetabulum, but was entirely imperforate.

2,463. Loxia leucoptera. 8:35 10:60 ♂
Wing, 3.35; tail, 2.75; culmen, .65; gape, on right side, .65; tarsus, .63; middle toe and claw, .73; toe alone, .47; claw, .30. Iris, hazel. Bill with maxilla above and tip and sides of mandible, dusky; residue yellowish-horn-color. Tarsi and toes, dusky brown.

1882.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.

The first Pine Grosbeak (Pinicola enucleator) was seen to day. I heard its note before breakfast, and stepped out upon the piazza and saw a single one sitting on the summit of the largest Maple-tree. Ella shot it.

2,464. Pinicola enucleator. 8:80 13:35 ♀ ad.
Wing, 4.35; tail, 3.85; culmen, .85; gape, .62; tarsus, .93; middle toe, .65; its claw, .34; middle toe and claw, .98. Iris, hazel. Bill, dusky; horn-color at base of mandible. Tarsi and toes, black.

November 26th. H. S. N.Y.

Turdus migratorius. - None have been seen near this place for a fortnight or three weeks; but I found a few, on the 25th, in the hemlocks south of Fort Montgomery.

Sialia sialis. - One flock (♂♂ five, ♀ one) seen to-day, and others heard warbling sweetly. The stellate involucre of the asters and golden rods, as well as the cottony matrix of those flowers still left upon the dry stems are especially pretty now. The leaves of Aplectrum hyemale, the Adam and Eve, are now full-grown, and interesting. Yesterday I found the ripe capsules of that rare Orchis coralorrhiza odonorrhiza.

November 28th. H. S. N.Y.

Zonotrichia albicollis. - To-day the ground is deeply covered with snow; and among the birds that came about the house was a fine adult White-throated Sparrow. I have never seen it here later than the middle of November before.

Nov. 1882.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

November 29th. N.Y.

It snowed during the night and is snowing still (11 am.) The Crows (*Corvus frugivorus*) are tamer than I have ever seen them before. Yesterday I drove under a tree in which several were sitting, and they paid no attention to me. Today they sat close to the stable and did not fly away when I drove in, as the usual do.

December 9th. N.Y. & N.Y. City.

Pinicola enucleator. - One was seen in Mr. Henry Cranston's hotel grounds. Louis A. Zerega shot one to-day at Westchester, in the East River, N. Y.

Zonotrichia albicollis. - Numerous in Central Park, New York City.

Passerella iliaca. - Several companies seen in Central Park, uttering their call-note only.

This evening attended the opening meeting of the Linnaean Society of Natural History of New York City. The first volume of the "Transactions" of the Society was laid before the Society, and Mr. Eugene P. Bicknell, ^{President} informed the Society that the cost of publishing an edition of 600 copies had been \$415-⁷⁷/₁₀₀.

I had the pleasure of meeting my old friends, Messrs. Wm. K. Latta of Florida, and Dr. C. Hart Merriam, and A. K. Fisher, besides many other friends, among whom were Dr. C. L. Bagg, Dr. Robt. J. Morns and Newbold J. Lawrence.

1882.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

December 14th. H.N.Y.

Pinicola enucleator. - Walked out with my wife, and saw a number of Pine Grosbeaks.

December 16th. N.Y.N.Y.

Pinicola enucleator. - A flock of a dozen or more were eating the seeds of *Abies excelsa* in the row of handsome trees of that species in front of Mr. Bigelow's place, as I drove to town. After dinner, Ella & I went out and found several flocks, some of them quite wild, others exceedingly gentle. Ella shot three specimens! None of the birds seen was an adult male. One bird was feeding upon ~~all~~ the berries of the Sheep Bush (*Viburnum lentago*) which the people here designate by the name of "Nanny Berry", and permitted my wife and I to stand beside it and watch it eating for many minutes. Others were eating the Red Cedar berries.

- 2,465. *Pinicola enucleator* 8:30 13:00 ♀ (ad.)
Wing, 4.13; tail, 3.45; culmen, .57; gape, .59; tarsus, .88; middle toe and its claw, 1.00; claw alone, .36. Shot by E. W. Mearns.
- 2,466. *Pinicola enucleator* 8:65 13:40 ♀ ad.
Wing, 4.20; tail, 3.60; culmen, .60; gape, .62; tarsus, .89; middle toe and claw, .97; claw alone, .35. Shot by E. W. Mearns.
- 2,467. *Pinicola enucleator*. 9:75 14:00 ♀ ad.
Wing, 4.40; tail, 4.00; culmen, .59; gape, .63; tarsus, .92; middle toe and claw, 1.03; claw alone, .40. Shot by Ella W. Mearns. Red on rump but a ♀.

1882

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

December 18th. N.Y.

Pinicola enucleator. - One came to the house this morning, several were seen in the Norway Spruces (*Abies excelsa*) along Mrs. Bigelow's place, and a flock of about twenty was found in the same sort of trees just inside of Mrs. S.R. Roe's place. I watched this flock for a long time. Every one flew down to the reservoir, where there was a little water and took a thorough bath, notwithstanding the severely cold weather. He splashed the water about and enjoyed themselves immensely. Not a Red one among. I omitted to state that the above specimens (Nos. 2465, 2466 & 2467) had a large number of the ^{small} green buds of *Abies excelsa* as well as the seeds in their crops.

December 19th. N.Y.

Sialia sialis. - Several females seen at West Point.

December 21st. N.Y.

Anorthura troglodytes hyemalis. A Winter Wren was about the house this morning. The first I have seen this winter.

2,468. *Regulus satrapa*. 4.10 6.62 ♀ ad.
Wing, 2.17; tail, 1.70; bill measured from nostril, .20; gape, .44; tarsus, .66; middle toe and its claw, .52; claw alone, .15. Irides, hazel. Bill, black. Tarsi, & toes above, ~~the~~ brownish-black; feet yellow underneath. Shot by Ella W. Mearns.

1882

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.2,469. *Regulus satrapa*. 3.97 6.45 ♀ ad.

Wing, 2.07; tail, 1.70; bill, measured from nostril, .21; gape, .45; tarsus, .66; middle toe and claw, .53; claw alone, .16. Colors of parts the same as No. 2468. Shot by Mrs. E. W. Mearns.

2,470. *Sitta carolinensis*. 6.00 10.75 ♀ ad.

Wing, 3.45; tail, 1.97; bill, measured from nostril, .50; culmen, .73; gape, .88; tarsus, .73; middle toe and claw, .85; claw alone, .25. Shot by Mrs. E. W. Mearns.

2,471. *Sitta carolinensis*. 6.02 11.10 ♂ ad.

Wing, 3.65; tail, 2.02; culmen, .77; bill measured from nostril, .54; gape, .92; tarsus, .75; middle toe and claw, .92; claw alone, .28. Irides, hazel. Bill, blue-black; basal one-half of mandible, pale blue. Tarsi and toes, plumbeous. Shot by E. W. Mearns.

December 23rd. N.Y.

To-day it's warm. Flies are about the house; and I saw a wasp flying across the road. This morning I rode to the hemlock woods below Fort Montgomery in search of Northern birds. I saw none. This afternoon was called to see the Rev. Mr. Campbell back of Ft. Monty. Ella went along. We saw two female Pine Grosbeaks (*Pinicola enucleator*).

December 25th (Xmas). N.Y.

Sialia sialis. - A female Bluebird was seen near the stable to-day.

I rode from the rock cut on Mrs. Roe's place to the spiling across Consook Marsh, upon "Dony", on the newly finished

1882.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

West Shore Railroad track. At the marsh saw
Colaptes auratus. It flew across to the
(Corisook) Island. Have seen no others
this winter.

December 26th. H.S.N.Y.

Received from Lieut. Willis Wittich a young
Loon.

No. 87. Colymbus torquatus

Loon. Shot at Fort Canby, W. T., mouth of
the Columbia River. Dec. 15th, 1882.
Length, 33.00; stretch, 60.50; wing, 14.90;
tail, 3.25; culmen, 3.25; gape, 4.72;
height of bill at middle of nostril,
1.03; width of bill at posterior border of
nostril, .67; tarsus, 3.90; middle toe
and claw, 5.10; its claw, .57.

Although a young bird, its
measurements exceed those of Eastern
specimens of Colymbus torquatus, as
given by Dr. Coues, Bds. N.W., pp. 720, 721.
They are, in fact, intermediate between
the Eastern torquatus, and adamsii.

Its bill has the dorsal surface of max-
illa to the extreme tip, black; sides of
maxilla nearly to tip and whole of
mandible, bluish. Outer side of tarsus,
black. Inferior surface of toes, and
whole of ~~of~~ outer toe except the three
terminal phalanges (which are brown-
dusky at joints), black. Toes above
(excepting the outer), pale greenish-blue,
varying to flesh color in ^{and dusky at joints} places; Webbs,
dusky on sides, flesh-color in the
middle, and slightly bluish above.
Hallux, blackish, bordered with brownish
on the outer side; flesh-color on the

1882.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

inner surface. Nails, horn-color at
base; black at extremity.

Egiothus linari (Linn.) Cab. Common Red-
poll. - To-day a flock of about
thirty was found. They were feeding
upon the seeds of the white birch,
and were very gentle. A single
shot brought down three specimens.
I have not seen any others this year.

2,472. Egiothus linaria (Linn.) Cab. 5.40 8.65 ♂ ad.
Wing, 2.90; tail, 2.40; culmen, .38; gape, .44;
tarsus, .59; middle toe and claw, .55; claw
alone, .25. Irides, hazel. Bill, with a
black line on terminal one-half, both above
(on maxilla) and below (on mandible). Tarsi
and toes, dark-brown; claws black.

2,473. Egiothus linaria (Linn.) Caban. 5.12 8.35 ♀ ad.
Wing, 2.82; tail, 2.48; culmen, .40; gape, .44;
tarsus, .54; middle toe and claw, .58;
claw alone, .26. Colors of parts like No. 2,472.

2,474. Egiothus linaria (Linn.) Cab. 5.10 8.32 ♀ ad.
Wing, 2.70; tail, 2.43; culmen, .39; gape, .44;
tarsus, .55; middle toe and claw, .57;
claw alone, .24. Colors of parts like No. 2,472.

Mr. Wm. Church Osborn found the Snowbird
(Junco hyemalis) at Lake Mohawk about
July 25th, 1877.

December 27th. H.S., N.Y.

Egiothus linaria. - There was a flock
of a dozen at the stable this morning.

2,475 Regulus satrapa. 4.28 7.02 ♂ ad.
Wing, 2.25; tail, 1.75; nostril, .21; gape, .45;
tarsus, .72; middle toe and claw, .54; claw
alone, .16. Irides, hazel. Bill, black. Tarsi

1882

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

and toes, brown - except soles or under surface of feet and toes which are yellow, a yellow shade extending into the brown of the dorsal surface of the toes. Claws, brown. Shot by E. W. Mearns.

Note. - As an interesting fact in ornithological pathology, I will mention that I have preserved in alcohol this day a fatty tumor weighing eleven (11) ounces, which was taken to-day (Dec. 27th, 1882), from the abdomen of a hen (Gallus). I have previously preserved, and noted the fact, a huge ovarian tumor taken from a hen.

December 28th. H. F. N.Y.

Egiothus linaria. - We found the flock this morning again, feeding on white birch seeds behind the stable. Ella shot two females.

Picus villosus. - Saw three; the first seen or heard this winter. I shot a large female as it was flying overhead.

2,476. Picus villosus. 9:35 1500 ♀ ad.

Wing, 4.70; tail, 3.60; culmen, 1.09; gape, 1.19; tarsus, .83; middle toe and claw, .85; claw alone, .38.

Irides, bright hazel. Bill, bluish. Tarsi and toes bluish lead-color.

2,477. Spizella montana (E. W. M.) 6:12 9:25 ♂ ad.

Wing, 3.00; tail, 2.80; culmen, .43; gape, .45; tarsus, .85; middle toe and claw, .79; claw alone, .23; Irides, hazel. Bill,

with maxilla blackish; yellow at base above; mandible, yellow, tipped with black. Tarsi, brown. Feet dusky brown. Claws, black. Shot by E. W. Mearns.

1882

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

2,478. Egiothus linaria (E. W. M.) 5:48 8:75 ♀ ad. Wing, 3.00; tail, 2.54; culmen, .42; gape, .45; tarsus, .58; middle toe and its claw, .56; claw alone, .23. Irides, hazel. Bill, yellow, with a black line above and below. Tarsi & toes brownish-black; claws, black.

2,479. Egiothus linaria (E. W. M.) 5:24 8:50 ♀ ad. Wing, 2.75; tail, 2.42; culmen, .35; gape, .39; tarsus, .55; middle toe and claw, .55; claw alone, .20. Colors of parts like No. 2,478.

December 29th. H. F. N.Y.

Pinicola enucleator. - Four were seen at Capt. Roe's gate, opposite to our house.

I rode to Mountville, west of Jones Island, but saw no noteworthy birds.

2,480. Egiothus linaria. 5:30 8:52 ♀ ad. Wing, 2.94; tail, 2.40; culmen, .39; gape, .43; tarsus, .53; middle toe and claw, .60; claw alone, .25. Colors of parts the same as No. 2,478. Shot by E. W. Mearns.

December 30th. H. F. N.Y.

To-day rowed across the Hudson, and tramped for half a day in Putnam county, where I saw the following species: Regulus satrapa. - Very numerous among the hemlocks.

Parus atricapillus. - Numerous.

Certhia familiaris. - Common.

Sitta capolinensis. - Common. In the middle of the day the sun was very powerful. I heard a sleepy note from this species. Looking up, I saw a male White-bellied Nuthatch picking or rather nodding upon a branch. Found downy in

1882.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

the crotch was a female sound asleep, with her head drawn into her feathers dreaming sweet dreams of buggy tree-trunks. I tossed a branch against the tree and roused her from dream-land. She flew to the next tree.

Anorthura troglodytes hyemalis. - Two were seen near a stream. I saw one on December 21st. No others seen this winter.

Pinicola enucleator. - Heard the note when in a hemlock woods, but did not find the birds.

Melospiza melodia. - Two found in the cat-tails across the river, near Cox's.

Junco hyemalis. - A few.

Spizella monticola. - Scarce.

Astragalinus tristis. - One small flock was feeding upon the seeds of the alder.

Corvus frugivorus. - Common.

Cyanocitta cristata. - Very abundant back of Osborn's house. Three shot.

Picus villosus. - One seen.

Larus argentatus. - Two adults.

Note. The railroad men all say that no "White Snowbirds" (Plectrophaeus nivalis) have been about yet this winter.

Sciurus carolinensis (Gray Squirrel).

I saw two; and nests are very abundant & generally distributed this year, on both sides of the Hudson.

2,481. Anorthura troglodytes hyemalis 4.23 6.25 ♂ ad.

Wing, 1.90; tail, 1.28; bill, measured from nostril, .32; culmen, .47; gape, .65; tarsus, .72; middle toe and claw, .70; claw alone, .18. Irides, hazel. Bill, black above; mandible & edges of maxilla, flesh-color. Tarsi, toes & claws, light brown. Stomach filled with ants and other insects.

1882.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

2,482. Melospiza fasciata. 6.45 8.75 ♀ ad.
Wing, 2.67; tail, 2.75; culmen, .50; gape, .57; tarsus, .85; middle toe and claw, .84; claw alone, .24. Irides, hazel. Bill, black above; dusky at tip of mandible; residue of mandible, horn-color. Tarsi, fleshy-brown; feet, brown; claws, dusky-brown.

2,483. Cyanocitta cristata. 11.75 16.80 ♂ ad.
Wing, 5.40; tail, 5.40; culmen, 1.16; gape, 1.30; tarsus, 1.38; middle toe and claw, 1.15; claw alone, .38. Iris, hazel. Bill, legs & feet, black. Top of head paler than usual.

2,484. Cyanocitta cristata. 12.02 17.00 ♂ ad.
Wing, 5.50; tail, 5.55; culmen, 1.12; gape, 1.28; tarsus, 1.37; middle toe and claw, 1.20; claw alone, .40. Irides, hazel. Bill, legs & feet, black.

2,485. Cyanocitta cristata. 12.05 17.05 ♂ ad.
Wing, 5.60; tail, 5.80; culmen, 1.12; gape, 1.27; tarsus, 1.30; middle toe and claw, 1.15; claw alone, .37. Irides, hazel. Bill, legs & feet, black.

December 31st. H. F., N. Y.

A flock of eight or ten Pine Grosbeaks (Pinicola enucleator) is feeding on the Norway Spruce-tree in front of the house. They are eating the buds; and none were eating seeds while I watched them. A few weeks ago, both the seeds and buds of this tree (Abies excelsa) were found. Although hemlock (Abies canadensis) and pitch pine (Pinus rigida) cones are numerous, I have not known the Pine Grosbeaks to feed, this season, upon other evergreens than Abies excelsa.

1882-'83.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

, nor upon ~~upon~~ the seeds of apples, of which latter they usually seem fond.

Jan. 1, 1883.

At the beginning of the year, I will give a statement of the number of specimens taken each year since 1871, as the specimens collected may be said to fairly represent the amount of field work done.

In 1874, I wrote at the beginning of my Journal, vol. I, a catalogue of the skins then in my collection, which were taken by myself. I did not include in the catalogue a collection of mounted birds which I had made during the years 1873 and 1874, nor the skins which had previously left my cabinet. I had a separate record of a part of these specimens, which I catalogued separately from the balance of my collection, in 1880, in this book, pp. 102 to 108, using Roman numerals, thus making a separate catalogue without exactly duplicating the numbers.

Both collections are given in this statement, only the earlier collection is represented by Roman numerals and then added.

Birds Collected the Year

1871		1.
1872		5.
1873	LXLVI + 4 =	100.
1874	CXVI + 303 =	419.
1875		544.
1876		487.

1883.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

1877		190.
1878		289.
1879		183.
1880		166.
1881		80.
1882		233.

Twelve years. Total, 2,697

January 2nd.

Pinicola enucleator. — Heard whistling on my way down to the dock.

January 7th.

Reached home last night. Rode on Pony to Fort Montgomery, and into the hemlocks, but saw no birds of special interest.

This afternoon my wife walked down to the River. We saw one Herring Gull (Larus argentatus).

The dock cut in front of Mrs. S. R. Roe's place is about finished; and Havens dock covered with racks of ties which will soon be laid.

January 14th. New York City.

This morning took a walk through the whole length of Central Park. We were especially anxious to find the Fox Sparrow (Passerella iliaca) in the Park. We have seen it there late in December, and early in February, and a middle of December record is all that is essential to establish its winter residence in Central Park. There is no record of its occurrence throughout the winter so far north; but Mr. Arthur H. Helme informed me a short time since that it winters regularly on the north shore of Long Island, at his residence. We saw none of them to-day, however.

1883.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

The Muskrat (Fiber zibethicus) was quite numerous. We saw its signs & tracks about nearly all of the waters in the Park.

Gray Squirrels (Sciurus carolinensis leucotis) were abundant. We saw three at one time, all chewing on pine cones of the yellow species. Rob. packed a snowball hard, and was just going to send it after one of them, when a Park policeman shouted in his ear, and completely paralyzed him!

The following birds were seen:

Regulus satrapa. — Abundant.

Sitta canadensis. — One.

Certhia familiaris. — Very numerous.

Zonotrichia albicollis. — Common.

Melospiza fasciata. — Scarce.

Passer domesticus. —

Corvus frugivorus. — Several.

Carduelis elegans. — Flock of 20 at Pool.

Junco hyemalis. A number.

Cardinalis virginianus. — Two were seen, one at the upper and one at the lower part.

January 21st. N.Y., N.Y.

This afternoon Ella and I took a long walk by the river. The Hudson is frozen over and people are walking across it.

We saw a large flock Lesser Redpolls (Aegithus linaria) and a couple of Fox Sparrows (Spizella montana).

In the dusk of evening I saw a shadow without as I sat by the dining-room window, and, looking outside, saw a fine Owl (Scops asio). It sat there for some time. Ella got the big shotgun and filled it with a charge of dust shot. I skinned it for her cabinet.

1883.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

2,486. Scops asio. Red plumage. 9.85 24.80 7.10. Wing, 7.00; tail, 3.65; cere, .33; culmen, measured from cere, .67; from base of cere to tip of bill, .90; tarsus, 1.33; middle toe and claw, 1.45; toe alone, 1.14; claw alone, .50; gape, 1.09. Irides, bright yellow. Mounted in alcohol, given to me by Mr. J. S. Morris, N.Y.

January 22nd. New York City.

Took the 7.42 A.M. train to New York. Just above High Bridge, I saw an immature White-headed Eagle (Haliaeetus leucocephalus), which the train started up from the ice on the river.

January 28th. New York City.

Spent the whole afternoon walking in Central Park with Dr. Rob. J. Morris. We were watching the wild water-fowl in a pool near the American Museum of Natural History, when I heard the note of the bird that attracted us to the Park in this stormy afternoon. It was the Fox Sparrow (Passerella iliaca). It came from a Norway Spruce-tree (Abies excelsa) where a pair of them were seeking shelter from the storm. In the same tree were several Purple Finches (Carpodacus purpureus), one of ^{them} a male and possessed of two white tail-feathers, showing that he was somewhat affected by the disease known as albinism. From the same point we saw a bird sitting upon a telegraph wire on 8th Ave. Soon it was joined by another. They flew past us to a peninsula in the Lake, where we followed and found several others of what proved to be Cardinal Grosbeaks (Cardinalis virginianus), and another Fox Sparrow. We afterwards saw several Fox Sparrows at the upper part of Central Park. We found a number of European Goldfinches.

1883.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

(*Carduelis elegans*), that were eating seeds from the button wood (*Platanus occidentalis*), and singing.

A. Cooper's Hawk (*Accipiter cooperi*) was seen trying to catch a bird, near the lower reservoir (west) and alighted close to us. We watched him. He watched us. He was very bold and handsome. He flew swiftly, when started, and low; made a splendid, swift upward swoop and alighted on a tree-top on the opposite side of the green. It sat there until after we left him.

Gray Squirrels (*Sciurus carolinensis leucotis*) eat the long acorns which grow near the lower side of the large reservoir, in deep, mossy-fringed cups. They also eat the seeds of many of the Pine Cones in Central Park. This afternoon we saw a pair eating the seeds of the large beans which grow in the pods of the thorny locusts. They dispatched them rapidly, sometimes holding them in their arms like a performer on the harp. Rob suggested that a certain Squirrel was able to pry himself off of the limb with the huge pod he was manipulating. When he had extracted all the seeds, he threw the pod down at us. It measures about a foot long and an inch in width. They showed skill in managing their clumsy provisions.

We saw, in addition the following birds:

Certhia familiaris. Brown Creeper.
Zonotrichia albicollis. White-throated Sparrow.
Melospiza fasciata. Song Sparrow.
Passer domesticus. European House Sparrow.
Junco hyemalis. Eastern Snowbird.
Carduelis elegans. European Goldfinch.
Passerella iliaca. Fox Sparrow. I have previously noted the presence of the Fox Sparrow in Central

1883.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

Park, during the months of December and February. Its occurrence is now proven for the remaining winter month (January), and that, too, after a spell of unusually cold winter weather, and in some numbers.

Cardinalis virginianus. Cardinal Grosbeak.
Regulus satrapa. Golden-crested Kinglet.
Picus pubescens. Downy Woodpecker. One seen.
Cyanocitta cristata. - Blue Jay. (Heard once.)

January 31st

Came from New York today. The Hudson is nearly all frozen over; but some Herring Gulls (*Larus argentatus*) were seen at different points on the Hudson, and also upon the Harlem River in New York.

February 1st. Highland Falls, N.Y.

Took a long walk with my wife. We found a large flock of Lesser Redpolls (*Aegithus linaria*) feeding upon the white bird seeds. Ella shot three with her first barrel and two with second. She has got a winged one in a cage; a female.

2,487. *Aegithus linaria*. Shot by E.W.M. 5:35 8:65 ♂ ad. Wing, 2.96; tail, 2.45; culmen, .38; gape, .42; tarsus, .59; middle toe and claw, .57; claw alone, .24. Irides, hazel. Bill, yellow, with extremity of both mandibles, especially the upper tipped with black. Tarsi and toes, black.

2,488. *Aegithus linaria*. Shot by E.W.M. 5:40 8:70 ♂ ad. Wing, 2.87; tail, 2.33; culmen, .36; gape, .41; tarsus, .57; middle toe and claw, .53; claw alone, .21. Colors of parts the same as No. 2,487.

2,489. *Aegithus linaria*. Shot by E.W.M. 5:30 8:45 ♀ ad. Wing, 2.85; tail, 2.42; culmen, .38; gape, .43; tarsus, .60; middle toe and claw, .55; claw alone, .24.

1883.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

February 2nd. N.Y.

A Pine Goldfinch (*Chrysomitris pinus*) was heard calling as it flew over the place. I think I heard them early in the winter.

February 5th. N.Y.

Pinicola enucleator. - A Pine Grosbeak came to the front of the house, and called loudly.

February 6th. N.Y.

This afternoon a Pine Grosbeak alighted near the house.

In this afternoon I drove across down to Fort Montgomery, where I found my Reefed House (*Chama umbellata*) enjoying excellent health. While in the hemlock woods a flock of birds passed overhead just above the tree tops uttering notes like those of the Red Crossbill (*Loxia curvirostris umbrinera*); but as I have not heard that species since '87, I am not quite certain, unless I should see and hear them again.

Empidonax griseus. - I shot a ^{white} *Empidonax* during a couple of ^{white} *Empidonax* on the side of the house in the hemlock to-day. I have never noted seeing a single one last winter; and in '87-88 I procured specimens and saw others during that winter. I have only noted the *Emmit* *Emmit* here on two ^{white} *Empidonax* winters.

2,490. *Hylocichla unalascae pallasii*. 7.12 11:35 ♂ ad. Wing, 3.53; tail, 2.82; bill, measured from nostril, .40; gape, .88; tarsus, 1.18; middle toe and claw, .91; toe alone, .71; claw, .21. Irides, dark hazel. Bill, brownish-black on maxilla and terminal one-half of mandible. Basal one-half of mandible and inside of mouth, yellow. Legs and

1883.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

feet, fleshy-brown. This bird was fat, and remarkably strong and well developed.

February 8th, H.F., New York.

Pinicola enucleator. - A number of Pine Grosbeaks came about the house this afternoon, and ate the buds of the Silver maple, and the apples that still hung upon the trees. All were in female plumage. I tried to make a double shot on two birds that were flying very swiftly but missed with my second barrel. The last bird alighted on the big white oak, and whistled loudly for a long time.

2,491. *Pinicola enucleator*. 8.65 13:40 ♂ juv. Wing, 4.25; tail, 3.50; culmen, .59; gape, .60; tarsus, .87; middle toe and claw, .95; its claw alone, .33. The crop, ~~stomach~~ of this bird was filled with the seeds of the Sugar Maple (*Acer saccharinum*) and the meat of the apples which are upon the trees still. There were no apple seeds in its crop, but only the apple itself, in morsels the size of a split pea.

February 9th. H.F., N.Y.

A flock of fourteen (14) Pine Grosbeaks (*Pinicola enucleator*) came to the sugar maple by the house and began feeding on its buds; they continued all over the tree, and were not at all alarmed when Ella and I went out and stood under the tree and watched them, the lowest being but a few feet above us. They have a pretty, soft note of conversation which sounds when they become excited, tho at times they were perfectly quiet, except

1883.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

for the noise their bills made in biting off the buds, which was quite considerable. After a while a number of them flew to some apple-trees behind the house, and commenced eating again. Two of them descended to the ground and ate those which were down there. Our cutter came up and frightened them two, so that they flew up into the trees; and we drove off leaving the flock still feeding, but all were in female plumage, and none red males. They were gone when we returned.

February 10th. H.E.M.Y.

Elting Roe came down from Cornwall today on his skates. We went out hunting specimens, but found nothing, although we took a long tramp on this side of the Hudson, and then crossed over on the ice to Green Point and after hunting up that region, walked on the Railroad track to a point opposite here and then crossed over. None of the employees on the Railroad remember having seen any Snow Buntings (*Plectrophenax nivalis*) this winter.

Sialia sialis. — Elting saw some at Cornwall, on the Hudson, yesterday.

February 12th. H.E.M.Y.

This morning, early, a large flock of Snow Buntings (*Plectrophenax nivalis*) flew across our place, in front of the house. They are the first I have seen, or heard of their having been seen by others, this winter. They flew at

1883.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

first close to the snow, following the inequalities of the ground, and afterwards arose to a height so as to pass over the tree-tops below our place. They twittered so loud that I saw them coming long before they reached our grounds.

This afternoon several Lesser Redpolls (*Aegithus linaria*) alighted in the neighborhood of our house. Afterwards found a small flock in a hemlock tree (*Abies canadensis*) and a red cedar (*Juniperus virginiana*) that grew near together south of Echo Lake, in a pasture grown up to alder bushes. They hid quite nicely in the evergreens, in which situation I never before saw a Lesser Redpoll. The species is now separating into small flocks and more widely distributed over the region.

2,492. *Aegithus linaria*. 5:50 9:00 ♂ ad.
Wing, 3.05; tail, 2.40; culmen, .40; gape, .45; tarsus, .59; middle toe and its claw, .57; claw alone, .23. Irides, dark hazel. Bill, yellow, with a ridge on culmen and gonys, black. Tarsi, toes and claws, black.

2,493. *Aegithus linaria*. 5:45 8:55 ♂ juv.
Wing, 2.87; tail, 2.37; culmen, .40; gape, .45; tarsus, .57; middle toe and its claw, .58; claw alone, .23. Colors the same as No. 2492, except that the neck & tarsi, toes & claws have a slight brownish tinge.

2,494. *Aegithus linaria*. 5:45 8:50 ♂ juv.
Wing, 2.95; tail, 2.45; culmen, .40; gape, .45; tarsus, .61; middle toe and claw, .59; claw alone, .25. Irides, hazel. Bill, yellow, with a narrow ridge on culmen and gonys of brownish-black. Tarsi, toes & claws, brownish-black.

1883.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

February 13th. N. G. M. Y.

This morning I saw a flock of Pine Grosbeaks in the Norway Spruce trees in front of Mr. Bigelow's place. One of the birds had some red on its rump. As I returned home, later in the morning, they were eating apples in Captain S. R. Roe's orchard.

The Lesser Redpolls (*Aegiothus linaria*), came about the house this afternoon.

February 17th. N. G. M. Y.

Passerella iliaca. — I shot a fine specimen in a grove of cedars near Fort Montgomery. This is the earliest record I have for the Fox Sparrow in the Highlands. My only other February record for this region is Feb. 28th, 1878.

Anorthura troglodytes hiemalis. — One was seen today; and another Winter Wren was heard near the house a couple of days ago.

Pinicola enucleator. — Two good-sized flocks were seen this morning; both were feeding upon the buds of the Sugar Maple. All were females, so far as I could see.

2,495. *Corvus umbellus*. — One flew from a hemlock tree in Paeroti's woods; and there were flocks among the hemlocks at Fort Montgomery.

Passerella iliaca (Merrem) Sw. 7:30 11:10 ♂ ad.
Wing, 3.40; tail, 3.10; culmen, .55; gape, .63; tarsus, 1.00; middle toe and claw, .98; claw alone, .33. Irides, hazel. Bill, with maxilla brownish-black, except laterally on basal one-half, where it is yellowish; mandible, pale yellow at base, shading through bluish and horn color to black at the extremity. This bird was fat, and robust. Its tarsi and feet were brownish flesh-color; claws brownish. Bill not so yellow as in late March specimens.

1883.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

February 18th. N. G. M. Y.

Aegiothus linarium. — The Lesser Redpolls were feeding upon the seeds of rag weeds near our gate this morning. One called "two-e-e-t" like a Yellowbird (*Geothlypis trichas*) as Ella & I passed, on our way to Church.

February 19th. N. G. M. Y.

Corvus frugivorus. — Crows have been only moderately common hitherto, this winter, but now they are appearing in numbers, showing that their northward movement is begun.

Aegiothus linaria. — One flock heard to-day.

February 20th. N. G. M. Y.

Halieetus leucocephalus. — Three were seen soaring together high in the air.

February 21st. N. G. M. Y.

Took a walk after dinner, but not birds of note were seen, with the exception of an immature female Yellow-bellied Woodpecker (*Sphyrapicus varius*), which I shot near our house in the orchard. It was eating apples upon the tree, and its bill was encased in a layer of frozen, decayed apple, as the day is cold.

When I reached home, a message awaited me; and I had to go to Fort Montgomery dock to see a sick child. A flock of Bluebirds (*Sialia sialis*) was seen there. Some alighted on the iron bridge that is being constructed above Poppen's Creek, notwithstanding the fact that an hundred workmen were pounding upon the noisy affair.

I heard a Bluebird on February 17th near the same spot. None others seen since I returned from New York City the last of January.

1883.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

February 21st - Continued.

2,496.

Sphyrapicus varius 8.60 15.40 ♀ ad.
Wing, 4.90; tail, 3.20; culmen, .99; gape,
1.14; tarsus, .75; middle toe and claw, .88;
claw alone, .34. Irides, dark hazel.
Bill olive-brown-black. Tarsi & toes,
grayish-blue; claws, dusky. See p. 359.

February 24th. H. J. N.Y.

Sialia sialis. - The first Bluebird I have seen
on our place since I returned from the City, the
end of January. It was warbling loudly.

February 25th. H. J. N.Y.

Melospiza fasciata. - Saw a Song Sparrow
at Dr. Chase's place this afternoon; and I saw
one at Faurot's brook on the 21st. These are
the first migrants. The few winter residents
frequent other situations, of small
extent, which they do not leave during winter.

February 26th. H. J. N.Y.

Sialia sialis. - Warbling sweetly.

February 28th.

Took a long walk. Crossed the Hudson on
foot. The ice is very thick. The only birds
of note seen were a number of Pine Grosbeaks
Pinicola enucleator seen on Messrs. Osborn
and Fishes' premises. Also a White-headed Eagle
near Garisons, where I walked to mail
some letters, the ferry having broken down.
Melospiza fasciata - Numerous in the
Cox marsh near Fish's. Common Crows
(Corvus frugivorus) are tame, and migra-
ting in numbers.

1883.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

I also noticed a flock of Crows that
were pursuing a large white bird of prey
of some species. It was quite white
below, nearly white above, & tail short.
At first sight I thought it a Snowy
Owl. Nyctea scandiaca; but afterwards
thought it soared more like a Hawk. It
was soaring very high up, and was in view
for some time. It also seemed too small for
a Snowy Owl.

March 1st. H. J. N.Y.

Walked across the River on the ice, which
is slushy on top owing to the warm day, but
is still thick and strong. On Mr. Osborn's
place a few Pine Finches (Chrysomitris pinus)
were seen. They were with some Pine Grosbeaks
(Pinicola enucleator), among some mixed
Norway spruces and larch-trees, and they
also descended to the ground, where some
tall grass was standing in a wet meadow.
The latter uttered a note like "Chee-a" when flying.
I shot two of the Pine Grosbeaks. One of
them was slightly wounded in the wing. I
brought it home alive, & it is scarcely
afraid of me even now, although it bit
viciously when I first caught it and called
loudly. Now it does not try to bite, but raises
the feathers on its head and neck, and
utters a sweet, subdued note.
Among the alder (Alnus) along the brook at
the head of Cox marsh, I found a large
flock of Lesser Redpolls; and while I watched
them, another flock flew from the south
calling. As soon as my flock heard them
they showed some excitement and answered
the strangers in the most cordial manner.

1883.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

, giving them a hearty and vociferous welcome as they settled down upon the Alder bushes among themselves. They were "at home" at once. They were thirsty rather than hungry, and settled down among the weeds beside the glaze (Cut-tails) to drink. I saw one very bright male, and shot into the flock; hoping to get some adult male specimens; but I only killed two in female dress. I left them lying in the shallow water a few moments and got six more from the flock, all of them in female plumage. Returning to my first two dead birds I found their legs literally covered with a mass of leeches which were each about half an inch in length. I shot these birds in the very alders where most of my specimens of the memorable winter of 1874-75 were obtained. I have been wishing to go over and visit the old, familiar spot all winter, but have been too busy for much gunning. Sialia socialis. — Bluebirds common. Singing.

About the first plant that starts growth in spring in this locality is the Skunk's Cabbage (Symplocarpus foetidus). In a small stream in Fish's place, at the end of the saltmarsh, I saw two plants, one of them four or five inches in height, but not yet in flower.

- 2,497. Pinicola enucleator. 8.80 13.30 ♂ juv
Wing, 4.20; tail, 3.70; culmen, .58; gape, .63; tarsus, .90; middle toe and claw, .98; claw alone, .38. Irides, hazel. Bill, black, with an olive tinge on sides of maxilla at base and on base of mandible. Tarsi, toes and claws, black. The bird was fat, and its crop was full of decayed apple.

1883.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

- 2,498. Aegiothus linaria. 5.40 8.85 ♀ ad.
Wing, 2.92; tail, 2.30; culmen, .38; gape, .44; tarsus, .62; middle toe and claw, .56; claw, .22. Iris, hazel. Bill, yellow, with a line of black above and below (shading to dusky towards base above). Tarsi, toes and claws, black. Crop full of alder seeds.
- 2,499. Aegiothus linaria. Breast, rose. 5.12 8.40 ♀ ad.
Wing, 2.75; tail, 2.17; culmen, .38; gape, .41; tarsus, .61; middle toe and claw, .55; claw alone, .23. Colors, etc. as above, only the tarsi are brownish-black instead of black.
- 2,500. Aegiothus linaria. 5.30 8.80 ♀ ad.
Wing, 2.97; tail, 2.35; culmen, .37; gape, .42; tarsus, .62; middle toe and claw, .56; claw, .22. Colors, etc. like No. 2498.
- 2,501. Aegiothus linaria. 5.10 8.20 ♀ ad.
Wing, 2.70; tail, 2.22; culmen, .36; gape, .41; tarsus, .60; middle toe and its claw, .57; claw alone, .24. Colors, etc. like 2498.
- 2,502. Aegiothus linaria. 5.35 8.50 ♀ ad.
Wing, 2.78; tail, 2.43; culmen, .39; gape, .45; tarsus, .58; middle toe and its claw, .58; - same as tarsus; claw alone, .24. Colors like No. 2498.
- 2,503. Aegiothus linaria. 5.05 8.30 ♀ ad.
Wing, 2.79; tail, 2.22; culmen, .37; gape, .43; tarsus, .57; middle toe and its claw, .56; claw alone, .23. Colors of parts, etc. like No. 2498.
- 2,504. Aegiothus linaria. 5.08 8.28 ♀ ad.
Wing, 2.72; tail, 2.10; culmen, .39; gape, .43; tarsus, .62; middle toe and claw, .55; claw alone, .22. Colors of parts, etc. same as No. 2498.
- 2,505. Aegiothus linaria. 5.25 8.43 ♀ ad.
Wing, 2.75; tail, 2.25; culmen, .38; gape, .44; tarsus, .62; middle toe and its claw, .38; claw alone, .20. Colors of parts, etc., the same as in No. 2498.

1883.

- 2,506. Passer domesticus Linn. 6.30 9.45 ♀ ad.
Wing, 2.90; tail, 2.37; culmen, .53; gape, .60; tarsus, .75; middle toe and its claw, .82; claw alone, .21. Irides hazel. Tarsi and toes, brown. Bill, olive-brown; yellow at base of mandible. Shot at Meams House!

March 2nd. H. G. M.
Sialia sialis. - In numbers singing loudly.
Buteo pennsylvanicus - One seen.

March 3rd.
Turdus migratorius. - A large flock was feeding on the ground and among the Sambucus (Rhus typhina) near the stable. They are the first Robins I have seen since the first part of the winter.

Pinicola enucleator. - A flock of Pine Grosbeaks was seen behind the house, eating apples on the trees. They were tame, as usual and all in female dress.

- 2,507. Turdus migratorius Linn. 9.75 15.20 ♀ ad.
Wing, 4.80; tail, 3.90; bill measured from nostril, .52; gape, 1.07; tarsus 1.22; middle toe and its claw, 1.09; claw alone, .25; Irides, hazel. Bill, olive-brown at base of maxilla above, shading into black towards the extremity; terminal one-third of mandible, black; residue of bill, yellow. Tarsi and dorsum of feet, dusky brown; claws, blackish above; under surface of feet, flesh-color.

- 2,508. Pinicola enucleator. 8.90 13.25 ♀ ad.
Wing, 4.35; tail, 3.75; culmen, .60; gape, .59; tarsus, .83; middle toe and claw, .85; claw alone, .28. Irides, hazel. Bill olive on basal one-half of mandible; residue, black. Tarsi,

1883.

toes and claws, black. Top of head, mixed yellow and red; rump, reddish. Crop full of apple and apple-seeds, from which the shells had been neatly removed. I noted that the seeds of these frozen apples had commenced to germinate.

- 2,509. Pinicola enucleator. 9.00 13.60 ♂ juv.
Wing, 4.40; tail, 3.85; culmen, .61; gape, .63; tarsus, .87; middle toe and claw, .95; claw alone, .53. Iris, hazel. Bill black; live green on basal one-half of mandible. Tarsi, toes and claws, black. Contents of crop, apple and apple-seeds.

- 2,510. Pinicola enucleator. 8.90 13.50 ♂ juv.
Wing, 4.25; tail, 3.60; culmen, .58; gape, .60; tarsus, .89; middle toe and claw, .97; claw alone, .36. Colors and contents of stomach same as No. 2,508.

- 2,511. Pinicola enucleator. 9.12 13.60 ♂ juv.
Wing, 4.40; tail, 3.90; culmen, .60; gape, .64; tarsus, .87; middle toe and claw, .95; claw alone, .32. Colors of tail and contents of crop the same as No. 2,508. This specimen, as well as Nos. 2,509 + 2,510 are in female dress, although the breast is quite distinctly rusty. They were carefully dissected and their sex accurately determined.

March 5th. H. G. M.
Passerella iliaca. - heard near the house.
Turdus migratorius. - Numbers of them on the place. Thermometer 14° above zero at 8.30 A.M.
Our Pine Grosbeak (Pinicola enucleator) is perfectly tame. He eats white food close beside him, and whistles so loudly that I hear him when out of doors. I have given him a large room. He calmly, as a rule, sits to the back of a chair upon which he roosts.

1883.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

He likes the seeds of the Norway Spruce (*Abies excelsa*) which I fed him very much. He also eats large quantities of apple.

March 6th. H.S.M.Y.

To-day is stormy. The wind has blown all day and a good deal of snow has fallen. I have not been out; but Robins (*Merula migratoria*) have been numerous about the house, as have the Bluebirds (*Sialia sialis*). A few Fox Sparrows have also been about (*Passerella iliaca*).

March 8th. H.S.M.Y.

Weather still cold. Ground snow-covered. *Egiothus linaria*. - Seen near Echo Lake. *Merula migratoria*. - Robins plenty.

March 9th. H.S.M.Y.

Merula migratoria. - Now quite numerous and generally dispersed over the region. *Passerella iliaca*. - Two seen at Fort

Montgomery, amongst the cedars. *Chrysomitris pinus*. - A small flock was seen feeding upon the berries of the sheep bush (*Viburnum lentago*), at Fort Montgomery. I fired a shot into the bush and killed two good specimens.

2,512. *Chrysomitris pinus*. 5:18 8:90 ♂ ad.

Wing, 2.90; tail, 2.00; culmen, .44; gape, .49; tarsus, .55; middle toe and its claw, .65; claw alone, .21. Irides, dark hazel. Bill, bluish horn-color, tipped with black above and below. Tarsi, toes and claws, dark olive-brown.

The crop of this bird was full of small black seeds, and contained no sheep-berries.

1883.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

2,513. *Chrysomitris pinus*. 5:23 8:90 ♂ ad.

Wing, 2.92; tail, 1.95; culmen, .46; gape, .50; tarsus, .56; middle toe and its claw, .67; claw alone, .21. Colors of parts like 2512. I was, evidently, mistaken in thinking that these birds were feeding upon sheep-berries. The flock was in a sheep bush when they were shot; but the crops & gizzards of both of the specimens contained only a multitude of small seeds, black in color, and resembling grass-seeds. They certainly appeared to be feeding before I fired.

March 12th. H.S.M.Y.

The Hudson is clear from West Point down to the southern point of Consock Island, where the ice is piled 15 feet high, the cakes being over a foot thick. This afternoon saw the first Flicker (*Colaptes auratus*). A single one was on Consock Island.

March 13th. H.S.M.Y.

Junco hyemalis. - To-day is a bright day. Thermometer 42° F. at 11 o'clock, A.M. The Snowbirds are all singing sweetly. I have not heard them ^{singing} before this season. *Accipiter cooperi*. - First seen this spring.

March 15th. H.S.M.Y.

Sayornis fusca. - First seen. One was heard calling Phoebe before I got up; and before breakfast I saw him flying about the house, and sitting on apple trees wagging his tail.

Buteo lineatus. - One was observed to raise a big commotion amongst a flock of Robins, although I did not see him try to

1883.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

strike any of them.

Melospiza fasciata. — First heard singing about the house this morning.

Quiscalus purpureus. — One flew over head.

The first seen this season.

To-day is the first real spring-like day, and it has brought its spring begins. This morning I noted the two first spring arrivals among the birds, excepting the Fox Sparrow, which I saw first on the 17th of February; and this afternoon I saw the first Swamp Sparrow (Melospiza palustris), in the cat-tails beside Consook Island. The flicker was seen in the same place on the Island as when seen there on the 12th inst.

I saw a pair of Chipmunks (Tamias striatus) on a stone-wall. They were not run out before. None were seen since last autumn.

The early spring butterfly, with orange, purplish-brown wings, edged with buff was seen first this afternoon, as was the Yellow Spotted Tortoise (Emydoidea blandingii).

March 16th. N.E.N.Y.

Egiothus linaria. — Three were seen feeding upon the seeds of the alders, below Echo Lake.

Passerella iliaca. — This species is now present in large numbers. They were heard (freezing temp. all day) singing this-morning, although the weather is windy & temperature below freezing point all day, although the sun shone brightly. Yesterday was warm.

1883.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

March 17th. H.E.N.Y.

Egiothus linaria. — This afternoon I met with a large flock of, say forty, Lesser-Redpolls, south of Echo Lake.

2,514. Egiothus linaria. 5.20 8.37 ♀ ad.

Wing, 2.83; tail, 2.33; culmen, .38; gape, .43; tarsus, .56; middle toe and its claw, .50; claw alone, .20.

Iris, hazel. Bill, yellow, with a black line above & below extending to the point. Tarsi toes and claws, blackish-brown.

The "jussy willows" were first seen in bloom, and I brought some home for little Lillian.

March 18th. N.E.N.Y.

Passerella iliaca. — Singing sweetly all about the place. The day warm. Thermometer ranging between 45 and 60° F.

March 19th.

Egiothus linaria. — This morning a female alighted in an apple tree over my head, and then flew away. It uttered a louder note than usual.

Dr. Albert K. Fisher reported at one of the Linnaean Society, N.Y., meetings during the past winter, the capture of the Greater Redpoll at Sing Sing, New York.

To-day is nearly as warm as yesterday. Spotted Tortoises (Emydoidea blandingii) are basking in the sun in the little brook below our place. The ponds are still frozen, and bear one's weight. The Hudson is entirely open, in the channel; but a few coves still have some ice in. Yesterday hundreds of Herring Gulls

1883.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

(*Larus argentatus*) were seen in the Cove above Harrison's, where there is still considerable ice.

Three White-headed Eagles (*Haliaeetus leucocephalus*) were seen in one tree in Cro' Nest ravine yesterday.

To-day Robins, Bluebirds, Tree Sparrows, Fox Sparrows, and Snowbirds are all plentiful about the place. No Robins have been heard to sing yet.

March 20th. H. S. N.Y.

Aegithus linaria. - There was a small flock of Redpolls in the alder-swamp below Echo Lake this afternoon. There were as many males as females. I stood on a rock in the swamp & called them all around me. The Fox Sparrows clicked angrily and the Redpolls uttered their usual note and a loud tweet. Both species erected the feathers on top of their heads.

Sayornis fuscus. - Saw one; the only one seen since the first one seen on March 15th.

Passerella iliaca. - Saw one having white tail feathers and rump.

March 21st. H. S. N.Y.

Zenaidura carolinensis. - Saw the first Mourning Dove to-day. None were seen during the winter.

Sitta carolinensis. - This morning, while seated at breakfast, we saw a White-bellied Nuthatch descend a maple-tree to the ground and then hop about on the lawn for some time. They seldom descend to the ground. Later in the day I saw the same species and the Chickadee, both feeding

1883.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

upon the ground; they were eating the meat upon some refuse bones, which had been thrown upon the ground.

When migrating upon the sea-shore, both Woodpeckers and Nuthatches occasionally alight upon the beach, where vegetation is scarce; but it is unusual for them to do so at other times, except in the case of *Colaptes auratus*, which is quite terrestrial. I am speaking only of the species of the Eastern United States.

March 25th. H. S. N.Y.

To-day is nearly as warm as last Sunday. Early in the morning I took a walk below the Lake. I found one Lesser Redpoll (*Aegithus linaria*) across the lower end of Echo Lake, which I was able to cross upon the ice. It was a female and flew down upon the ground from the top of a tree where I heard it calling "tweet". Afterwards I heard Redpolls again, and saw a flock of three fly down amongst some alder bushes. I went to the spot and lay down upon the ground near a little brook, in which the three Redpolls were bathing. Two of them were handsome adult males. I was unable to distinguish the plumage of the third. They erected the red feathers on their heads while bathing, and uttered the usual twittering note and "tweet".

Zenaidura carolinensis. - Several seen. *Sayornis fuscus* has now become pretty common. This afternoon I killed *Nannemys guttata* are basking in sunshine; and a Wood Frog (*Rana sylvatica*) was heard

1883.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

croaking lustily in the place where the turtles were first seen; and a single "Peeper" (*Hyla pickeringi*) was heard piping near the same spot. I caught and preserved one of the early Spring Butterflies. Neither of the Frogs above mentioned have been heard from before since last season.

Sialia sialis. - Bluebirds are mating or mated. A pair has been suspected a gallon tomato can in a pear-tree all day long, and they may even have begun a nest for I have seen them enter the can many times. One bird goes in & another sits on top of it while its mate is inside.

Corvus frugivorus. - Crows are mating and flying about a great deal as is their custom at this season.

March 26th. N. J. N.Y.

Spizella pusilla. - This morning I shot a Field Sparrow; the first one seen this season.

Sialia sialis. - To-day the Bluebirds began to fly against the windows and flitter and peck against the glass persistently.

March 27th. N. J. N.Y.

The ice still will bear one's weight upon Echo Lake.

Sialia sialis. - A female Bluebird pecked and flew against our window from daylight (shortly after) until breakfast time.

The Garrison Cove is nearly free of ice. Eagles are there in some numbers and also Ducks and Gulls. The following was shot there

1883.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

2,515.

Haliaeetus leucocephalus. 3400 8350 ♂ juv. Wing, 24.60; tail, 13.75; bill, measured from base of cere to end, 2.50; top of cere, .78; culmen from cere, 1.97; gape, 2.95; depth of bill, measured over the end of the cere, 1.33; tarsus, 3.45 (the tarsus was measured from the joint above to the fold of skin which is apparent upon excluding the toes. As nearly as I could get at the exact measurements it is a trifle less: about 3.25.); middle toe and claw, 3.75; toe alone, 2.50; claw, 1.30; hallux, 1.35; its claw, 1.60; inner toe, 1.75; its claw, 1.55; outer toe, 1.77; its claw, 1.10. The last ~~the~~ measurements (toes) were taken from the ~~tarsus~~ metatarso-phalangeal articulation to the end of the toes above.

Colors. - Irides, yellowish, still showing signs of hazel. This specimen leads me to believe that the plumage in which it is, is characteristic of a plumage ^{more} nearly adult than in the dark and uniformly marked specimens, since the irides are nearly as in the adult. This bird shows a great deal of white both above & below, while its head above & neck (above) are very light. It has an albinistic look; but I have seen several of them alive & know that is a regular phase of plumage, though somewhat uncommon. Bill, blue-black, marked with horn-color, yellow and green on the maxilla; but the mandible is greenish-yellow. Tarsi and toes, lemon-yellow. Claws uniformly black, with a faint bluish cast. Stomach & crop empty. Fishy smelling.

1883.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

March 31st H. J. M. Y.

Sialia sialis. - The female Bluebird that has built a nest in the tomato-can near the arbor comes regularly every morning and beats itself furiously against the window panes. If the window is opened half-way down from the top, she taps at the lower part, occasionally alighting upon the top of the sash. Sometimes she makes a short circuit within the room and then flies away.

Yesterday it snowed all day long, and to-day the ground is deeply covered with snow, notwithstanding which, Robins, Turdus migratorius were singing vigorously all the morning until breakfast-time. None have been heard to sing before this season.

On March 29th I saw those insects which inhabit clear water, and which look like Lady Long-legs with lifeboats on their toes () skating about upon Satterlee's reservoir. The first I have seen.

Chrysomitris (= Astragalinus) tristis. - This afternoon I shot an adult male. It was getting a black cap and had two narrow tracts of bright yellow below. Its rump was white, barely tinged with yellow, extending well up upon its back. The bird was moulting. This affords proof positive that this species, in spring, changes color by moulting. It was in full song, and uttered a loud tweet.

April 1st H. J. M. Y.

To-day is sunny. Thermometer 36° F. at 2.30 P.M. The snow had already disappeared from the ground in exposed

1883.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

places. The ice has melted from the whole of the north half of Echo Lake; but is still so solid in the south end that I walked safely across it this afternoon.

Erythraea linaria. - I saw a female near Echo Lake. She was alone and gentle. Uttered the "tweet" and chattering note when flying.

Turtles (Trachemys guttata) are out again. I obtained a good proof of their acuteness of hearing, if one were needed. I walked up to a wall and leaned over, looking into a swamp-hole. As I stood there a Turtle came up. It did not see me. I made a sharp sound, and it instantly turned its head round and looked at me.

Spizella domestica. - One. The first seen.

April 7th. N.Y. City + H. J. M. Y.

Ceryle alcyon. - Several Kingfishers, the first seen this year, were observed along the River. Agelaius phoeniceus. - An adult male was seen just below Peekskill. The first seen.

Yesterday, a number of Crow Blackbirds (Quiscalus purpureus) were seen in Central Park, New York City.

April 8th. H. J. M. Y.

To-day has been quite bright. Thermometer 40° to 50° F. Went to church this morning. This afternoon rode around Highland Lake, and through the hemlocks at Fort Montgomery. The Wood Frogs (Rana sylvatica) and Hyla pickeringii are abundant and vociferous. There is still some ice upon the north slopes of the mountains; but none upon Highland Lake, Echo Lake or Pell's pond.

1883.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

April 8th - Continued.

This morning, near Pell's pond I found a large specimen of the Large Spotted Salamander (Amblystoma punctatum).

This afternoon, I tied my horse near the bridge at Poplopen's Creek, and went to spot on the north, steep bank of that stream sloping to the sun, where I found the first Hepaticas (Hepatica triloba) just a year ago to-day - Saturday, April 8th, 1882. To-day I found numbers of them in the same place, and the white ones were sweet scented, but, disappointing my hopes that they might be H. acutiloba, all were typical of H. triloba. On the steep, rocky roadside leading up the north bank of the stream I also found some Early Saxifrages (Saxifraga virginensis) in flower. Very probably they were the very first ones to open in this region. I brought half of them home along with a large bunch of leaves and flowers of the hepatica. Various species of alders & willows are flowering.

The following birds were seen to-day.

Merula migratoria. - Everywhere common & singing.

Sialia sialis. - Nesting and scattered.

Regulus satrapa. - A number seen.

Parus atricapillus. - Uttering its disyllabic note.

Anorthura troglodytes hyemalis. - One at Highland Falls.

Spizella pusilla. - The first heard to sing this season.

Junco hyemalis. - Abundant. Singing.

Melospiza fasciata. - Singing. Rather common.

Passerella iliaca. - A number seen. Singing.

Corvus frugivorus. - Unusually common in small flocks, and so tame that I rode under the trees in which they sat several times during the afternoon.

1883.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

Sayornis fuscus. - One flew with some moss to the old mill on Poplopen's Creek, where it is building its nest.

Picus pubescens.

Zenaidura carolinensis. - A pair seen near Echo Lake.

Anas obscura. - This morning I flushed a couple of big, strong birds from the brushy margin of Pell's pond. They rose with as mighty splash. Splendid birds! Furns argentatus. - One seen on the Hudson.

April 9th.

H. G. N. Y.

The first Shad was caught here on April 7th. To-day I crossed the Hudson, chiefly for the purpose of gathering specimens of the Skunk's Cabbage (Symplocarpus foetidus). There were plenty of both flowers and leaves; but the flowers were gone from such plants as had large leaves. The Veratrum viride was growing plentifully, tho' all of those seen were small, save a few which were upwards of a foot high. Both plants were found at the lower brook in Gouverneur's Cove, north of Garrison's, where I also saw a few Hepaticas (H. triloba), which were white pinkish or blue like those found yesterday at Poplopen's Creek. One of the latter had eight sepals, but round lobed leaves, and was fragrant. In both places where these first Hepaticas were found, within half a stone's throw lay banks of ice - many tons of it!

To-day two small Brown Butterflies were seen, having black spots.

Near Garrison's I found a new plant in bud, which I have not seen elsewhere.

Quite a number of Gulls and Ducks were

1883.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	seen in Gouverneur's Cove.					
	<i>Larus delawarensis</i> . - One or two adults seen.					
	<i>Larus argentatus</i> . - Quite numerous.					
	<i>Anas obscura</i> . - About one hundred Dusky Ducks were feeding in the cove. They almost always swam in line, single file. I watched them under cover for four hours during which time I was quite near them. All of that time they were always swimming or feeding in lines, as the flock frequently divided up. Their note was exactly like the quack of domestic ducks. I shot one from my Berenof raffles.					
	<i>Clangula glaucium americana</i> . - About twenty were in the cove. They swam in solid bunches and not in lines. Their note was like the croaking of a pool-ful of Wood Frogs (<i>Rana sylvatica</i>); but the note is of shorter duration and sharper quality than the frogs' chorus. Now and then I heard a loud whistle or trumpeted note, which I fancied also came from the Golden-eyes. Perhaps it is from this note that they are called "whistlers" instead of from the sound made by their wings in flying as I have always supposed. I also heard a hollow rasping sound from time to time; but as there were several species of Ducks near-by I could not distinguish which species produced it.					
	<i>Ceryle alcyon</i> . - One in the cove & one at Caustoir.					
	<i>Colaptes auratus</i> . - Producing their laughing cry.					
	<i>Pandion haliaetus carolinensis</i> . - One alighted in a tree very near to me. The first seen.					
	<i>Sciurus carolinensis</i> . - One seen on the ground near the lower falls in the Cove.					

1883.

2, 516.	<i>Anas obscura</i> Gmel.	2440	3780	♂ ad.
Wing, 10.95; tail, 4.50; culmen, 2.02; gape, 2.62; nail on bill, length, .44; breadth, .30; tarsus, 1.88; middle toe and claw, 2.57; claw alone, .42. Irides, hazel. Bill with maxilla greenish-yellow; nail, black; mandible irregularly marked with olive, yellowish and black. Tarsi, toes and their adjacent margins of webs, bright orange; webs and claws and bottom of feet, blackish. Although the flock appeared to be actively engaged in feeding, this bird contained only a quantity of sharp-edged, white, quartz pebbles, which were in its gizzard.				

April 13th. H. S. M. Y.

Mr. Denton drove with me to Highland Lake. At the Poplopen's Creek we found sweet-scented, five-lobed specimens of *Hepatica triloba acutiloba*; and on the north bank of that stream we found a quantity of Saxifrage (*S. virginensis*) and the first specimens of the Rock Cress (*Arabis leptota*). Near Highland Lake we found the Arbutus, the first of the season (*Epigaea repens*).

I looked for the Large-billed Accentor along Poplopen's Creek; but none have yet arrived.

April 14th. H. S. M. Y.

Orchis spectabilis. - Several plants of the Showy Orchis were seen starting, about an inch high.

April 17th. H. S. M. Y.

Sivrus notacilla. - Several Large-billed Accentors were seen on our walk. The first this season. The Marsh Marigolds (*Caltha palustris*) are just about to open their buds. *Hepatica triloba* is now in its prime.

1883.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
2,517.	<i>Corvus frugivorus</i> (Baird)	April 18th.	N. J. N.Y.	19.10	37.35	♂ ad.
	Wing, 11.60; tail, 7.50; culmen, 1.92; gape, 2.20; depth of bill opposite base of culmen, .80; tarsus, 2.45; middle toe and claw 2.07; claw alone, .65. Irides, hazel. Bill, legs & feet, black. Shot while watching for a flock of its feeding comrades.					

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
2,518.	<i>Colaptes auratus</i> .	April 19th.	N. J. N.Y.	12.50	19.80	♀ ad.
	Wing, 5.95; tail, 4.65; culmen, 1.31; gape, 1.53; tarsus, 1.15; middle toe and claw, 1.32; claw alone, .46. Irides, brownish-red. Bill, blue-black. Tarsi and toes bluish lead-color.					

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
2,519.	<i>Sivus motacilla</i> .	April 21st.	N. J. N.Y. & Cornwall.	6.30	10.25	♂ ad.
	Wing, 3.25; tail, 2.23; bill, measured from nostril, .45; gape, .75; tarsus, .93; middle toe & claw, .75; claw alone, .19. Irides, hazel. Bill, dusky brown, flesh-color on basal half of mandible. Tarsi and toes, flesh-color.					

April 21st. N. J. N.Y. & Cornwall.
To-day rode over the mountain to spend Sabbath day with Rev. E. P. Roe. The day was bright. Near the summit of Cro' nest I saw some fine dandelions (*Taraxacum dens-leonis*) in flower. On the summit of the mountain I saw a Hermit Thrush (*Hylocichla ustulata* *pallasi*), and another was seen in the fort on Plum Point above Cornwall. Both the *Taraxacum* and the Thrush were the first seen this season except the single Hermit Thrush (No. 2,490) which I shot on February 6th. The Marsh Marigold (*Caltha palustris*) was also in bloom on top of the mountain, beside a stream, and was also the first seen in blossom.

1883.

Willows of several species scented the air. When I reached Cornwall, I saw the Dutchman's Breeds (*Dicentra cucullaria*), Blood-root (*Sanguinaria canadensis*) and Blue Anemone (*Thalictrum aneminioides*) all blooming, and the first seen this season. A single Winter Wren (*Troglodytes hyemalis*) was seen on the mountain; also the first *Regulus calendula*.

After dinner I played lawn tennis until Elting asked me to go shooting on Plum Point. We found four Fish Hawks (*Pandion haliaetus carolinensis*) fishing in the Bay. A Crow (*Corvus frugivorus*) was seen sitting on her eggs; and the first swallow (*Iachycineta bicolor*) was seen towards evening. It was solitary, and flew over our heads in circles for some time. In the Fort (in which there are sixteen places for cannons) the "scurrying" rush (*Equisetum*) was very rank, making a solid bed of green. Hepaticas and blue anemones grow upon the breastwork.

April 22nd. Cornwall on the Hudson.
This morning heard Lyman Abbott preach. After the church service Mr. Roe, the ladies and myself took a long drive along Moodna Creek. We saw the first Chewink (*Pipilo erythrophthalmus*), a fine male; but he seemed silent and moody. We failed, even by pelting him vigorously with sticks and stones, to elicit a syllable from him. The first White-throated Sparrows (*Zonotrichia albicollis*) were also first seen in the same spot. A fine Osprey (*Pandion haliaetus carolinensis*) sat on a branch of a huge plane tree. We drove to within easy shotgun range when he started and flew, but not away as we

1883.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

expected he would: he came straight towards us until very near, then darted perpendicularly into the stream, and would have disappeared from view beneath its surface, had not the water been as clear as crystal, so that we could distinguish both bird and fish. It soon emerged with its prey—a fine “wind-fish,” and flew down the stream bearing it in its claws.

After dining Mr. Roe and myself walked into the woods. At the back part of his place we found a Woodchuck (*Arctomys monax*) so tame and habituated to the sight of man that he scarcely seemed to fear us; and when we chased him into his den in a rocky cliff beside the garden, his head was seen cropping first out of one crevice and then another, until he finally tumbled clear out and began scratching amongst the leaves. The rocks amongst which bedwells are covered with moist moss and that rare fern, the walking leaf (*Camptosorus rhyzophyllus*), of which I gathered a specimen. Upon the ledges of Storm King we found a pair of Ruffed Grouse (*Bonasa umbellus*).

April 23rd. Cornwall + N.H., N.Y.

This morning I drove over the mountain. It had stormed during the night; and the mountain-top was capped with white. The ice, though nearly melted, still lies in miniature glaciers upon the north side of Storm King. A mossy rock beside the road was covered with the *Camptosorus* or Walking Fern, making three localities in this region in which I have found it growing.

1883.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

The Marsh Mergolds on the mountain are buried in snow.

April 24th. N.H., N.Y.

Nyctardea grisea naevia.—A fine adult seen in Satterlee's woods beside a pool near the Hudson. It merely flapped up to the top of a low pine-tree when I surprised it. The first seen.

Quiscalus purpureus.—A few seen.

Carpodacus purpureus.—A flock of thirty, nearly all of which were males and sang. *Dendroica coronata*. Yellow-rumps first seen. A single.

Summary of the Birds Found During the Winter of 1882-'83.

The winter noted may be said to have commenced with the advent of the White-winged Crossbill on the 16th of November, 1882 and the departure of the Lesser Redpoll on the 1st of April, 1883.

I was in New York a week in December and nearly the whole month of January; and even when at home took little time for studying the birds. Still I made as many observations upon them as circumstances permitted. I took few rides or drives, so that many birds may have escaped me. Fifty-four (54) specimens were procured, and added to my cabinet.

Merula migratoria.—A few were seen amongst the hemlocks at Fort Montgomery until November 25th. None had been seen in this immediate neighborhood for two weeks previous; and none were again encountered until March 3rd, when numbers arrived suddenly.

Hylocichla unalascae pallasi.—A male (No. 2490) was shot on February 6th. It was not again seen until April 21st.

Sialia sialis.—Have been present, but not numerous throughout the winter. The

1883.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	first mated pair I noted commenced building a nest in a tomato-can on March 25th. On the 26th the female parent began to fly against our windows, and flutter and peck against the glass persistently. She continued this for several days, and was, I think either killed by so doing; or, else, a cat caught her upon her nest. Last season a pair of Bluebirds nested in a can near the house; and the female behaved in the same manner. See notes for that time.					
	<u>Regulus satrapa</u> . — Frequent during the winter.					
	<u>Parus atricapillus</u> . — Numerous as usual.					
	<u>Sitta carolinensis</u> . — Common throughout.					
	<u>Certhia familiaris rufa</u> . — As common as usual.					
	<u>Anorthura troglodytes hyemalis</u> . — Occasionally seen during the winter. Noted on April 21st, and others may still be seen.					
	<u>Paricola enucleator</u> . — The first one seen was solitary; shot by Ella in the maple beside the house on November 23rd. Several specimens were shot on March 3rd; but none were seen later. During the winter they were fairly numerous; but no red males were seen, although immature males were not rare. Notes are scattered through the Journal relating to their habits, etc.					
	<u>Carpodacus purpureus</u> . — I mention this species in order to note what is more remarkable than its presence would be, viz. its entire absence during the entire period which this summary includes. In Central Park, N.Y. City, however, I saw several Jan 28th. Here, I saw none till late in April. Other birds whose absence is remarkable are the Cedar Bird (<u>Amphispiza cedrorum</u>) of which I have seen not one during the given period; and the Northern Shrike (<u>Lanius borealis</u>),					

1883.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	The absence of the Yellow-rumped Warbler (<u>Dendroica coronata</u>) may also be noted, although this species is not regular here in winter. They were numerous during the winters of 1877-'78 and 1881-'82.					
	<u>Loxia leucoptera</u> . — Several were seen in some pine trees in front of the residence of Capt. S. R. Roe, where I wife shot a pair on November 16th, 1882. None have since been seen. I thought I heard the notes of a passing flock of Red Crossbill (<u>Loxia curvirostris americana</u>) when in the dense hemlocks at Fort Montgomery, on February 6th; but I may have been wrong.					
	<u>Aegiothrus linaria</u> . — First seen on December 26th, when a flock of 30 were seen near our stable; three of which I procured at a single shot, one of them an adult male. A solitary female, the last seen, was noted on April 1st, the same date on which I last saw them in the spring of 1875. They have been quite common during the winter.					
	<u>Astragalinus tristis</u> . — Very scarce all winter, in fact scarcely noted at all. On March 31st, I shot a male that was getting its black cap and summer plumage. This species was noted a few times only; but it is generally common in winter.					
	<u>Chrysomitris pinus</u> . — I think I heard the call-note early in the winter. I certainly did on February 2nd, when one called "che-a" several times as it flew over our place. I saw a small flock on Mr. Osborn's place across the River, on March 1st; and on the 9th of March, at Fort Montgomery I shot two specimens from a small flock. They have been scarce during the winter.					

1883.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

Plectrophanes nivalis. — On February 12th, a large flock swept over our place. No other Snow Buntings were seen, or heard of, during the winter.

Zonotrichia albicollis. — The last one seen was a fine adult that came about the back of our house, when the ground was deeply covered with snow, on November 28th. None were seen here after that until April 22nd, after which they were abundant. First heard singing their curious song on the 24th. In Central Park, New York City they spent the winter as usual. It is, therefore, strange that they were so long in reaching us in Spring. This, however I have always observed.

Spizella montana. — Generally pretty common during the winter. I find I have neglected to note the date of its arrival last autumn, nor do my notes furnish any record of its last occurrence here this spring. Possibly I may yet see it, as I have records of its staying until April 29th three different years.

Spizella domestica. — I have not noted the date of its last occurrence last autumn; but the first one appeared, this spring, on the 1st of April.

Spizella pusilla. — I have no record of its last occurrence last autumn. The first one seen this spring, was shot on March 26th.

Junco hyemalis. — Has been common, as usual all winter. It is still here (April 24th). On the 7th of June, 1882, I found the Snow Bird at Queensborough, near this place, but in the mountains. It was doubtless, breeding; but I was hastening to a patient, and had no time to stop and investigate. I have

1883.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

I failed to note the date of its arrival last fall. Melospiza fasciata. — A few remained all winter. The first migrants were noted on February 28th. Sang at our house first, March 15th.

Melospiza palustris. — Have no late fall date. The first one seen this spring, at Consook Marsh, was on the 15th of March.

Passerella iliaca. — Have not got the date of its last occurrence last fall. On the 17th of February, 1883, I shot a fine male specimen (No. 2495). My only other February record for this species was February 28th, 1878.

This year, none were seen from February 17th till March 5th, in this region; but I found them throughout the winter in Central Park, N. Y. City, for the first time.

Quiscalus purpureus. — One, the first seen, was noted on March 15. None were seen last fall.

Corvus frugivorus. — Common as usual. Found sitting on its eggs, at Cornwall, on April 21st.

Cyanocitta cristata. — Common as usual.

Sayornis fuscus. — I have no late record for last autumn. This spring one came to our back yard on March 15th, and called "Phoebe" before I was up. She stayed about for some time, and then we saw her fly away. We did not see another until March 20th. By the 25th they were common.

Picus villosus. — Occasionally seen throughout the winter.

Picus pubescens. — Fairly common all winter.

Sphyrapicus varius. — The only one seen during the time specified above, was a young female (No. 2496) shot in an apple tree on this place, February 21st. None have

1883.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

been seen since up to the present time (Apr. 24).
Colaptes auratus. — Saw one at Cousook
Marsh on Xmas day; and think he
stayed there all winter, for when I
next visited the place, on March 12th,
I found one on the Island; and he was
still there when I went again several
days later.

Strix nebulosa. — Mr. Royal Taylor sent a
living specimen to Mr. Edward Denton
from Peekskill. Mr. Denton let it go at West
Point; but it was recaptured by a young
soldier who keeps it still in the basement
of the Band Barracks, where I saw it along
with a whole menagerie of animals,
including a fine Robin, two Catbirds,
an English Sparrow, four Flying Squirrels,
several Red Squirrels and four Gray
Squirrels. The Red Squirrel bites the Gray
through the bars of its cage, and barks savagely.
The Gray Squirrels were so tame as to
run all over Mr. Denton and myself,
and jump from one shoulder to another.
The Owl was chained to a perch, and was
quite tame. The owner of this happy family
is a handsome young German fellow,
whom we routed from his Soldier's couch
to exhibit his Owl, which I wished to identify.

Scops asio. — Ella shot an adult female
(No. 2,486) in red plumage on January
21st. The only one seen.

Accipiter cooperi. — First seen on March 13th.
One was seen in Central Park, N.Y. City on
January 28th; but none were seen here
during the winter.

Tamniacus sparverius. — Only one seen
during the winter. In February.

1883.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

Buteo lineatus. — One seen March 15th: the only one.

Buteo pennsylvanicus. — First seen March 2.
None were noted during the winter.

Haliaeetus leucocephalus. — Occasional all
winter. One specimen (No. 2515 ♂ juv.) was
taken March 27th, in interesting plumage.

Bonasa umbellus. — A few seen, but not
at all abundant. First seen March 21st.
Zenaidura macroura. — First seen March 21st.
Larus argentatus. — Seen throughout the
winter, at intervals.

Besides the above mentioned species which I
have noted in my Journal, a few species
of Ducks, as the Golden-eye, etc. and
Grebes were seen in the Hudson. Other-
wise the above list is a complete enu-
meration of all the Birds seen by me
in the Hudson Highlands from November
16th, 1882 until April 1st, 1883.

Haliaeetus leucocephalus. — On April 9th,
I received from Lieutenant Willis Wittich, U.S.A.,
a skin of a Bald Eagle in adult plumage,
except that the white feathers of the head
were marked in occasional spots with fine lines
of deep brown; and the terminal part of the
shafts of a few of the small feathers of the crown
were also blackish-brown. The specimen was
accompanied by the following information: —

"This bird is a female, and the stretch
is 7 ft., 3 inches. Length, 34½ inches. Weight,
12 pounds. It was shot by Corporal Darnell,
Co. E., 21 Inf., near the fort. The Eagle was
sitting on a dead tree overlooking Baker's
Bay: was shot in the back (at 300 yds.), the
ball passing through the breast some place.
The Eagle flew then about 500 yds., and
lit in the top of a hemlock tree, then lodged

1883.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

somewhere in the branches. The Corporal cut down the tree which was about 18 inches in diameter, and secured the bird, intending to stuff it. He kept it one day, and the longer he thought about it, the larger the job looked; so he gave it away to the Keeper of the Life Saving Station here - Capt. Harris, who was going to get Mr. Allen to help him stuff it for the Life-boat house. They both went hunting and forgot all about it. I accidentally discovered it at the store, and carried it off as a great prize; but there are a lot of them about here, and at Fort Stevens, in the spring."

Dimensions. - Length, 87.00; stretch, 34.50 [Wittich.]; wing, 24.70; tail, 12.50; cere, .75; bill measured from base of cere (including it), 2.75; culmen, 2.25; gape, 3.15; tarsus, depth of bill, measured over the end of cere, 1.42; tarsus, 3.70; middle toe, 4.25; toe alone, 3.00; claw alone, 1.42; hind claw, 1.80. [Ad. Wittich.]

April 25th. N. F., N.Y.

To-day has been cold and windy. The Large-billed Accentor (*Sialia motacilla*) was heard singing sweetly; and a Chewink (*Pipilo erythrophthalmus*), the second one seen this spring and a female, was heard calling.

Beside the runlet proceeding from Bell's pond, I flushed a Ruffed Grouse (*Bonasa umbellus*). Her nest was found under the rock from which she sprang, and contained four (4) eggs, three of which were weather stained from the recent rains, and the fourth appeared to be perfectly fresh. *Corvus sinuatus*. - Five eggs were taken from a nest upon which the old

1883.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

bird was sitting.

Scops asio. - An Owl was flushed from a cedar-tree. As it flew away a Robin started in pursuit, and followed it to the spot where it alighted, and was there joined by its mate. I followed the sound of their scolding voices, and thus found and shot the Screech Owl, which, without the Robins' help I never could have found; but in shooting it, I accidentally killed one of the Robins, which I did not see until I started to leave the spot, when I discovered it lying dead at a little distance beyond the tree in which the Owl had sat.

I walked all over Consook marsh, but saw nothing of interest save a few Swamp Sparrows (*Melospiza palustris*), two of which I shot, and noted a great discrepancy between their sizes.

2,520. *Scops asio*. (Red pl.) 9.15 23.10 ♂ ad. Wing, 6.80; tail, 3.35; from base of cere to point of bill, .87; cere alone, .37; culmen exclusive of cere, .57; gape, 1.07; tarsus (about), 1.40; middle toe and claw, 1.28; toe alone, .98; claw, .48. Irides, bright yellow. Bill, bluish flesh-color. Feet, flesh-color: Basal one-half of claws, flesh-color, shading into black on terminal half.

2,521. *Merula migratoria*. 9.55 15.62 ♂ ad. Wing, 4.90; tail, 3.95; bill, measured from nostril, .50; gape, 1.05; tarsus, 1.29; middle toe and claw, 1.18; claw alone, .29. Irides, hazel. Bill, yellow-saving the extremity of maxilla, which is black. Tarsi + toes, deep brown.

2,522. *Melospiza palustris*. 6.25 8.30 ♂ Wing, 2.53; tail, 2.70; culmen, .52; gape, .55; tarsus, .90; middle toe and claw, .87; claw, .20. Irides, hazel. Bill, with maxilla black; man-

1883.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	-dible, bluish + flesh-color. Tarsi, toes and claws, light-brown.					
	This specimen has the top of its head striped; but is far larger than the following specimen, an adult male (No. 2523), although both are males. This is not the first time I have noted this discrepancy both in size and plumage; and it is a subject well worth investigating.					
2, 523.	<u>Melospiza palustris</u> .	5.77	7.75	♂ ad.		
	Wing, 2.32; tail, 2.37; culmen, .46; gape, .50; tarsus, .83; middle toe and claw, .84; claw alone, .23. Colors of parts as above (No. 2522).					
2, 524.	<u>Dendroica coronata</u> .	5.95	9.30	♂ ad.		
	Wing, 2.90; tail, 2.60; bill, measured from nostril, .30; gape, .54; tarsus, .72; middle toe + claw, .68; claw alone, .18. Irides, hazel. Bill, legs + feet, black.					
2, 525.	<u>Regulus calendula</u> (Crowned).	4.53	6.97	♀ ad.		
	Wing, 2.27; tail, 1.85; bill, measured from nostril, .24; gape, .48; tarsus, .73; middle toe and claw, .52; claw alone, .17. Irides, hazel.					
	April 26th. H.F., N.Y.					
	<u>Junco hyemalis</u> . - Several about the house.					
	<u>Pipilo erythrophthalmus</u> . - One or two only.					
	<u>Seiurus motacilla</u> . - Singing, just before dark, after 7 o'clock, P. M., very sweetly.					
2, 526.	<u>Carpodacus purpureus</u> . E.W.M.	6.30	10.20	♂ ad.		
	Wing, 3.23; tail, 2.43; culmen, .48; gape, .52; tarsus, .69; middle toe and claw, .74; claw alone, .22. Irides, hazel. Bill, pale-brown, with a purplish, fleshy tint, lightest below. Tarsi and toes, brown; claws dark-brown.					
2, 527.	<u>Carpodacus purpureus</u> .	6.25	10.25	♂ ad.		
	Wing, 3.27; tail, 2.40; culmen, .47; gape, .53; tarsus, .70; middle toe + claw, .76; claw alone, .23. Colors of parts like No. 2526. Shot by Ella W. Meams.					

1883.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
2, 528.	<u>Carpodacus purpureus</u> . Shot by E.W.M.	6.15	10.25	♂ ad.		
	Wing, 3.23; tail, 2.40; culmen, .49; gape, .55; tarsus, .68; middle toe and claw, .73; claw alone, .22. Colors of parts the same as No. 2526.					
2, 529.	<u>Carpodacus purpureus</u> . Shot by E.W.M.	6.00	10.03	♂ ad.		
	Wing, 3.10; tail, 2.35; culmen, .47; gape, .54; tarsus, .66; middle toe and claw, .74; claw alone, .20. Colors of parts like No. 2526.					
2, 530.	<u>Spizella pusilla</u> .	5.50	8.15	♂ ad.		
	Wing, 2.43; tail, 2.60; culmen, .40; tarsus, .70.					
	April 27th. H.F., N.Y.					
2, 531.	<u>Carpodacus purpureus</u> .	6.15	10.10	♂ ad.		
	Wing, 3.25; tail, 2.36; culmen, .47; gape, .57; tarsus, .70; gape, middle toe + claw, .78; claw alone, .23. Colors of parts like 2, 526.					
	<u>Pipilo erythrophthalmus</u> . - To-day numbers were seen, and their migration has fairly set in.					
	<u>Zenaidura carolinensis</u> . - There were upwards of a dozen in the pasture-lands south of Echo Lake; and their mournful notes resounded all day.					
	April 28th. H.F., N.Y.					
2, 532.	<u>Dendroica coronata</u> .	5.75	9.00	♂ ad.		
	Mr. Zerega and myself shot three adult males, all of which were moulting. The gray plumage comes in with the spring moult. The above specimen was mixed winter and summer plumage, the latter prevailing.					
2, 533.	<u>Junco hyemalis</u> .	6.15	9.55	♀ ad.		
2, 534.	<u>Seiurus motacilla</u> .	6.25	10.45	♂ ad.		
2, 535.	<u>Harporhynchus rufus</u> .	11.50	13.12	♂ ad.		
	<u>Minioilta varia</u> . - One. The first seen this year. Singing.					
	<u>Caprimulgus vociferus</u> . - We found one in the woods below the house, on the ground. The first seen this year.					

1883.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	<u>Harporhynchus rufus</u> .					The specimen shot was the first one seen this season.
	<u>Troglodytes domesticus</u> .					First noted this season. Singing on the porch.
	<u>Viola cucullata</u>					was first seen in flower to-day.
	<u>Viola canina</u>					was also first seen in flower to-day.
		April 30th.	N.H.M.			
2535.A.	<u>Fringoides maculatus</u> .					First seen this year.
	<u>Corvus ossifragus</u> .					While dredging a creek in Consook marsh, we plainly distinguished the note of a Fish Crow that sat croaking on the top of a pine-tree on Consook Island. Mr. Zerega who accompanied me went in pursuit of the crow, and shot it when soaring very high in the air. It affords the following dimensions:—
				16.45	33.00	♀ ad.
	Wing, 11.00 11.00; tail, 6.95; culmen, 1.53; gape, 1.79; tarsus, 1.76; middle toe and its claw, 1.72; toe alone, 1.25; claw, .52; grad-uation of tail, .77. Sides, hazel. Bill and feet, black.					
2535.B.	Mr. Zerega shot a fine Broad-winged Hawk (<u>Buteo pennsylvanicus</u>), which presented the following dimensions:—			16.60	37.50	♀ ad.
	Wing, 11.25; tail, 7.40; culmen and cere, 1.14; cere alone, .44; culmen alone, .78; gape, 1.28; tarsus, 2.67; middle toe and claw, 1.95; claw alone, .54. Sides hazel (the only sign of immaturity). Cere, greenish-yellow. Bill, blue-black, except the basal one-third of mandible which is pale-blue. Tarsi and toes, greenish-yellow.					
2536.	<u>Fringoides macularius</u> .			7.80	13.50	♀ ad.
	Wing, 4.15; tail, 2.25; culmen, .93; gape, 1.03; tarsus, .93; middle toe and claw, 1.90; claw, .20.					
	<u>Hirundo erythrogastra</u> .					First seen.

1883.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	<u>Siurus auricapillus</u> .					First seen. Singing.
	Mr. Louis A. Zerega and myself spent most of the day in collecting fishes in Consook marsh creek. We took the following species:—					
	<u>Notemigonus chrysolenus</u> (Mitch.) Jor.					Only one small specimen taken.
	<u>Fundulus heteroclitus</u> (L.) Gunther.					Exceedingly numerous. We took a number of specimens of both sexes.
	<u>Fundulus diaphanus</u> (L.S.) Ag.					Quite numerous. We took several.
	<u>Roccus Americanus</u> (Gmel.) J. & G. Common.					
	<u>Lepomis gibbosus</u> (L.) McKay.					Two only.
	<u>Apeltes quadracus</u> (Mitchill.) Brevoort.					
	We preserved 100 specimens, after counting the number of dorsal spines on each. In 79 specimens there were four spines each; in 19 there were five; in two there were but three. It deserves the name " <u>inconstans</u> " more than does that species itself.					
	We caught one small specimen of the Fork-tailed Cat-fish of the Hudson. The Shad fishermen are catching them weighing ten (10) pounds in their nets. The only other species which they take are Shad, the two species of Sturgeon and the Striped Bass. The Cat-fish are called "Lake Cats."					
	<u>Unguilis rostrata</u> (L. Suer) De Kay.					
	Several species of shells and one crustacean were also taken as well as <u>Chrysemys picta</u> , the first Painted Turtle I have seen this year.					
	<u>Naremys guttata</u> was also found copulating in the saltmarsh creek.					

1883.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
2,537.	<u>Luscalus purpureus.</u>	May 1st.	N.Y., N.Y.	12.40	17.80	♂ ad.
	Wing, 5.60; tail, 5.25; culmen, 1.19; gape, 1.37; tarsus, 1.50; middle toe and claw, 1.43; claw alone, .35. Irides, yellow. Bill, legs and feet, black. Shot back of the stable.					
2,538.	<u>Luscalus purpureus.</u>			10.85	15.97	♀ ad.
	Wing, 4.92; tail, 4.50; culmen, 1.08; gape, 1.27; tarsus, 1.32; middle toe and claw, 1.28; claw alone, .32. Irides, bill & feet colored as above.					
	<u>Corvus ossifragus.</u> — To-day near Echo Lake we plainly distinguished the note of a Fish Crow. Soon seven crows flew over our head, one of which croaked repeatedly as they passed by. Mr. Zerega noticed two Crows about the top of a large hemlock tree (<u>Cedrus canadensis</u>) in which he found a nest. From that direction the Crows came whose note proved one at least of their number to be a Fish Crow; but as the birds could not be shot, the nest was not disturbed until this evening, when we went to the tree expecting to kill the parent Crow, in which ^{we} were mistaken, however, for although the eggs were warm no bird was in sight. This was just about dusk after supper. The eggs five in number, were much less in size than average eggs of <u>Corvus frugivorus</u> . They measured as follows: 1.68 x 1.18; 1.68 x 1.16; 1.60 x 1.16; 1.70 x 1.18; 1.65 x 1.18.					
	After noting the Fish Crow in the flock and afterwards found one or two more of the same species near the Hudson, in the direction in which the flock had flown. They were chased and pestered by Common Crows.					

1883.

Buteo pennsylvanicus. — Several seen.

Note.

About April 14th, 1883, Lieutenant Willis Wittich, U.S.A., found a short-tailed Albatross floating upon the seabeach at Fort Canby, Washington Territory. He skinned it nicely and sent it to me, along with the following memorandum: "And now I'll tell you about another 'bird' which I will send you by mail in a few days, viz. nothing more or less than a ~~xxx~~ female Albatross — var. 'short as to his tail'. They are rarely seen about here, though I am informed that 500 or 600 miles off the coast they are abundant. This specimen was washed ashore during a storm in a very emaciated condition, though in a fair state of preservation. Her eye-ball was yet pretty full and good, though her plumage was somewhat bedraggled. Her length was 36 inches, and stretch 7 ft., 5 1/2 inches. She floated about some time and was thought to be a Gull, many of which are found dead after a severe storm, on the weather beach; but as it seemed a pretty large Gull, Mr. Allen, the Post Grader, fished it out and showed it to me. I asked what he would do with it; and he spoke of saving only the head — asked me if I wanted it; I told him that if I had it I would skin it for you. He gave it to me for you, and you will soon get it. Some of the old sailors about here say it is small — that they have seen the 'brown kind' that measured sixteen feet stretch." Lillie says you

1883.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

have two or three; but I do not remember to have seen them in your collection. The bird proves to be the Short-tailed Albatross (Diomedea brachyura Temm.).

Dimensions. - Length, 36.00; stretch, 89.50 ♀ ad. [written]; wing, 20.60; tail, 6.80; culmen, 5.60; distance between nostrils, .67; gape, 6.20; depth of bill at base, 2.00; depth of bill towards extremity, 1.40; depth of bill about the middle, 1.35; bare portion of tibia, 1.30; tarsus, 3.80; middle toe and its claw, 5.25; claw, .68. Upper maxilla, flesh-color, with the extremity for some distance, hatched; under mandible, rose-color; with large process at extremity, bluish-gray. Tarsi and feet, bluish-flesh-color, with toes and webs marked with ~~the~~ dusky. Claws brownish black, shading into horn-color at sides, bases & tips. The specimen reached me about a fortnight after it was shot, while in a soft condition, so that the webs of feet expanded perfectly; but it is probable that the colors of bill and feet had changed somewhat. The description of the dark portions of the plumage given by authors (Lawrence, P.R.R. Rep. Vol. IX, p. 822 and Coues, Key, p. 325) differs from my bird; and I will therefore describe my bird:

"Primaries, tips of secondaries and tertiaries, upper edge of the wing and greater wing coverts and terminal three-fifths of tail dark. This dark area varies in color from plumbeous-gray to brown and black. The outer webs of primaries are black; shafts white, shading to black ^{towards} end; shafts of rectrices the same, except middle pair which are white to end; residue of dark area varying from bluish-gray to brown and dusky. Residue of plumage, pure-white except hind neck buffy-tinted

1883.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

Note. - On April , 1883, I heard the note of a Fish Crow (Corvus ossifragus), and got my gun and went to the place, and found a flock of Crows, all but one of which were probably Corvus rugivorus. They were silent when I got near them; so I picked out the bird I thought looked smallest and shot it; but it got away; and as the flock flew away I again heard the Fish Crow's note. I was unable to find it again.

Note. - On April 25th, 1883, I found, north of Canisook Island, in the Hudson River, a string of periwinkle's "eggs". The ovaries measured 17 inches in length. I have never found the shells here; and never the ovaries till now. This is the species described by De Kay as Pyrrula canaliculata, pl. ~~IX~~, fig. 190, page, 140, 1843.

May 1st - Continued from page 397.

2, 539. Zenaidura carolinensis. 11.90 17.90 ♀ ad. Wing, 5.80; tail, 5.40; culmen, .58; gape, .82; tarsus, .80; middle toe and claw, 1.00; claw alone, .21. Shot near Echo Lake. Junco hyemalis. - Several seen.

May 2nd. H.F., N.Y.

To-day Mr. Ziegler accompanied me to Queensborough to work up the fishes in Poplopen's Creek. We worked the brook from the point where it approaches the ^{old} bridge to the Green St. bridge, going over the ground both ways taking the whole day to do it in. We found the following species of fishes only: - Semotilus corporalis (Mitch.) Putn. - We caught a number, but all were small specimens. Some had a rosy tint; but none had horns on head.

1883.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

Minnilus cornutus (Mitchill) D. & G.

We caught some fair sized Red-fins.

They were exquisitely colored, with crimson fins, and horns (spines) on their heads. They are vigorous biters.

Lepomis auritus (L.) Raf.

I caught a Long-eared Sunfish on a worm bait in the most rapid part of the stream when fishing for Trout. We caught several others.

Esox reticulatus Le Sueur.

The young ones are called "Jacks" They were numerous in still parts of the stream. We snared a couple of little "Jacks" in an arm of the brook, where we also took a number of ^(Cambarus bartoni) Crawfishes, and Desmystylus viridescens. The latter were copulating.

Salvelinus fontinalis (Mitch.) Gill & Por.

We saw several fine ones; but they refused to bite, so we got no specimens.

We found three species of Turtles in the brook, as follows:-

Chelopus insculptus Le C. - Mr. Berega took an unusually large one; but it escaped from the wagon while we ate lunch, and could not be re-captured.

Nanemys guttatus (Schu.) Ag. - The shell markings from specimens taken in the clear brook water were especially clear and bright.

Chrysemys picta (Herm.) Ga. - Several.

A small crustacean was also taken in the brook. Eothenops trichas was first seen.

A pair of Ospreys (Pandion haliaetus carolinensis) were fishing in the Creek.

Buteo pennsylvanicus. - A pair stayed about the brook all day.

Anemone nemorosa. - The Wind Flower was first found in bloom to-day, as was

1883.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

also the Dog-tooth Violet (Erythronium americanum).

The Broom Thrasher (Harporhynchus Rufus) was first heard singing to-day.

May 3rd. H.F., N.Y.

Corvus ossifragus. - Heard this morning.

Buteo pennsylvanicus. - Whistling.

Spinus motacilla. We spent the afternoon in Cro' Nest Ravine where we found several of these lovely birds. They serenaded us until the evening twilight; and the song rang out with unusual clearness and sweetness in the narrow ravine.

In the Ravine we found four species of Orchids growing: Orchis spectabilis of which there were several, Goodyearia pubescens one plant, Aplectrum hyemale perhaps an hundred leaves seen, and the Habenaria viridis, var. bracteata of which we had never before found more than a single specimen was quite numerous in one spot. The handsome leaves of Allium tricoccum, the Wild Leek, were well grown, and beautiful but vile-scented. Nervularia sessilifolia, the Sessile-leaved Bellwort we found in bloom, but only a single one. Ranunculus abortivus and Mitella diphylla (the Bishop's-Cap) were also first found in flower, as was the Sweet-scented White Violet (Viola blanda). "Jack-in-the-pulpit" or the Indian Turnip was also first flowered. A new species of Crustacean was found in the Ravine brook, along with some Salamanders and Crawfishes.

1883.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

May 4th

N. J., N.Y.

Setophaga ruticilla. - First seen to-day.Dendroica blackburniae. - First seen. Mr.

Louis A. Zerega shot a fine male.

Dendroica virens. - First seen. Mr. L. A.

Zerega shot an adult male. Several

others were seen and heard singing.

Empidonax minimus. - First seen.Dendroica junco. - First seen. One.Chactura pelagica. - First seen. Several.Sinus notacilla. - heard singing until fairly dark to-night; long after the Whippoorwills began singing.Bonasa umbellus. - Yesterday I heard them drumming. A day Mr. Zerega visited the Pheasant's nest which I found on the instant. The hen bird hadreturned and laid two more eggs since I shot at her as she flew from the nest, but had subsequently deserted the nest, so we took the six eggs, which were much weather-stained. Semotilus corporalis. -

A male with large horns was seen in brook flowing from Pell's pond.

Rana sylvatica. - Croaking in the swamp still.

Toads were heard yesterday.

Troglodytes domesticus. - Numerous. Singing all day.2540. Regulus calendula. No ruby crown. 4.37 7.00 ♀.

Wing, 2.18; tail, 1.78; bill, measured from nostril, .27; gape, .50; tarsus, .76; middle toe and claw, .54; claw alone, .18.

2541. Troglodytes aëdon. 4.96 6.80 ♂ ad.

Wing, 2.00; tail, 1.84; bill, measured from nostril, .35; gape, .68; tarsus, .67; middle toe and claw, .59; claw alone, .13.

Irides, hazel.

1883.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

2542. Hylocichla unalascae pallasi. 7.35 11.60 ♀ ad.

Wing, 3.55; tail, 2.88; bill, measured from nostril, .42; gape, .93; tarsus, 1.26; middle toe and claw, .98; claw alone, .23.

May 5th.

N. J., N.Y.

Dendroica pennsylvanica. - First seen.Parula americana. - First seen.Helminthophaga ruficapilla. - First seen.

One shot in an apple-tree by the house.

Taloscops carolinensis. - First seen.

One was nesting in the lilac-bushes by the house when I arose; and it was singing sweetly after breakfast.

The Shad-bushes are now in bloom. I got specimens of it (Amelanchier canadensis) to send to the Department, this morning.2543. Dendroica pennsylvanica. 5.07 7.90 ♂ ad.

Wing, 2.47; tail, 2.00; bill, measured from nostril, .29; gape, .55; tarsus, .75; middle toe and claw, .60; claw alone, .16. Iris, hazel. Bill, with maxilla black; mandible bluish horn-color. Tarsi and toes, bluish-gray.

2544. Parula americana. 4.98 7.70 ♂ ad.

Wing, 2.40; tail, 1.95; bill, measured from nostril, .30; gape, .54; tarsus, .66; middle toe and claw, .57; claw alone, .17. Iris, hazel. Bill, with maxilla black, barely edged with yellow; mandible bright yellow. Tarsi, yellowish-brown; feet, brownish-yellow. Claws, brown.

2545. Helminthophaga ruficapilla. 4.65 7.45 ♂ ad.

Wing, 2.32; tail, 1.80; bill, measured from nostril, .29; gape, .50; tarsus, .67; middle toe and its claw, .54; claw alone, .17. Iris, hazel. Bill, with maxilla brownish-black, edged with yellowish; mandible, yellow, with

1883.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
2546.	<i>Dendroica virens</i> .	5.65	7.85	♂ ad.		
	a bluish lateral line on either side. Tarsi and toes, yellowish-brown; claws, pale brown. Wing, 2.47; tail, 2.12; bill, measured from nostril, .29; gape, .53; tarsus, .70; middle toe and claw, .57; claw alone, .17. Irides, hazel. Bill and feet, black. Tarsi, toes and claws, blackish; soles of feet, yellowish.					
2547.	<i>Dendroica coronata</i> .	5.40	8.70	♀ ad.		
	Wing, 2.65; tail, 2.23; bill, measured from nostril, .27; gape, .54; tarsus, .69; middle toe and claw, .64; claw alone, .20. Iris, hazel. Bill, feet and claws, black.					
2548.	<i>Dendroica coronata</i> .	5.65	9.00	♀ ad.		
	Wing, 2.82; tail, 2.45; bill, measured from nostril, .29; gape, .55; tarsus, .73; middle toe and claw, .65; claw alone, .20. Irides, hazel. Bill, feet and claws, black. This specimen is positively feminine, notwithstanding the disparity in size between it and the preceding.					

May 6th.

H. G. M.

Myiodiocetes mitratus. — First seen. Singing
Dendroica palmarum. — First seen.

May 7th.

H. G. M.

To-day found the first specimens we have seen of that rare, northern Violet, *Viola ^{rostrata} ~~albertina~~*. It is a beautiful species.

Lanius solitarius. — First seen.

Hylocichla mustelina. — First seen.

Myiarchus crinitus. — First seen. Its call was such a distinct whistle that I thought a boy was whistling, although familiar with the whistle of this species.

Cistudo clausa. — The first Box Turtle was caught to-day.

1883.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	<i>Bonasa umbellus</i> . — Drumming loudly. Quite a number were flushed at different places. Last winter seems to have been favorable for their preservation. The winter before last they were very scarce, owing, doubtless to some epidemic disease. Now their ranks, happily, are increasing.					
	<i>Quiscalus purpureus</i> . — One flock of perhaps a dozen alighted in the woods. They are rare here, and seldom seen so late in the season. They breed plentifully over the mountains.					
2,549.	<i>Helminthophaga ruficapilla</i> .	4.90	7.75	♂ ad.		
	Wing, 2.38; tail, 1.95; bill, measured from nostril, .30; gape, .54; tarsus, .67; middle toe and claw, .57; claw alone, .18.					
2,550.	<i>Lanius solitarius</i> .	5.75	9.55	♀ ad.		
	Wing, 2.93; tail, 2.28; bill, measured from nostril, .29; gape, .63; tarsus, .76; middle toe and its claw, .63; claw alone, .21.					
2,551.	<i>Dendroica coronata</i> .	5.60	8.85	♀ ad.		
	Wing, 2.75; tail, 2.32; bill, measured from nostril, .24; gape, .52; tarsus, .72; middle toe and its claw, .66; claw alone, .15.					
	<i>Hylocichla mustelina</i> . 8.40 13.60 ♂ ad.					
2,552.	Wing, 4.40; tail, 3.25; bill, measured from nostril, .40; gape, 1.04; tarsus, 1.23; middle toe + claw, 1.03; claw alone, .27. Iris, hazel.					
2,553.	<i>Pipilo erythrophthalmus</i> .	8.12	10.85	♂ ad.		
	Wing, 3.33; tail, 3.75; culmen, .56; gape, .72; tarsus, 1.06; middle toe and its claw, 1.03; claw alone, .28. Irides, dark red. Bill, black. Tarsi, feet + claws, brown, with a fleshy tint on tarsi; feet and claws, successively darker.					
	Shot near Echo Lake, a small pond near the Mearns House.					

1883.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	<u>May 8th.</u>	<u>N. J. N. Y.</u>				
	<u>Dendroica maculosa.</u>					First seen.
	<u>Icteria virens.</u>					First seen.
	<u>Icterus spurius.</u>					First seen.
	<u>Icterus galbula.</u>					First seen. Abundant. One ♀.
	<u>Pyrranga rubra.</u>					First seen.
	<u>Zamelodia ludoviciana.</u>					First seen Several.
	<u>Tyrannus carolinensis.</u>					First seen.
	<u>Vireosylva olivacea.</u>					First seen.
	<u>Helminthophaga chrysophtera.</u>					First seen.
	<u>Tochilus colubris.</u>					First seen.
2,554.	<u>Helminthophaga ruficapilla.</u>	4.70	7.25	♀ ad.		
	Wing, 2.15; tail, 1.75; bill, measured from nostril, .27; gape, .52; tarsus, .70; middle toe and claw, .58; claw alone, .16.					
2,555.	<u>Helminthophaga ruficapilla.</u>	4.85	7.52	♀ ad.		
	Wing, 2.30; tail, 1.85; bill, measured from nostril, .28; gape, .52; tarsus, .68; middle toe and its claw, .57; claw alone, .18.					
2,556.	<u>Dendroica virens.</u>	5.30	8.10	♂ ad.		
	Wing, 2.55; tail, 2.15; bill, measured from nostril, .32; gape, .56; tarsus, .70; middle toe and claw, .59; claw alone, .20.					
2,557.	<u>Dendroica maculosa.</u>	5.05	7.45	♂ ad.		
	Wing, 2.35; tail, 2.40; bill, measured from nostril, .29; gape, .54; tarsus, .73; middle toe and its claw, .73; claw alone, .16.					
2,558.	<u>Taleoscoptes carolinensis.</u>	9.15	12.00	♀ ad.		
	Wing, 3.65; tail, 4.20; bill, measured from nostril, .45; gape, .96; tarsus, 1.06; middle toe and claw, 1.02; claw, .28.					
2,559.	<u>Icterus galbula.</u>	7.40	11.70	♂ ad.		
	Wing, 3.60; tail, 2.95; culmen, .70; gape, .80; tarsus, .84; middle toe and its claw, .85; claw alone, .26.					
2,560.	<u>Pyrranga rubra.</u>	7.10	11.75	♀ ad.		
	Wing, 3.65; tail, 2.80; bill, measured					

1883.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	from nostril, .44; culmen, .65; gape, .74; tarsus, .77; middle toe and claw, .79; claw alone, .24.					
2,561.	<u>Geothlypis trichas.</u>	5.47	7.30	♂ ad.		
	Wing, 2.12; tail, 2.06; bill, measured from nostril, .33; gape, .60; tarsus, .80; middle toe and its claw, .71; claw alone, .22.					
2,562.	<u>Tochilus colubris.</u>	3.70	4.10	♂ ad.		
	Wing, 1.55; tail, 1.20; culmen, .67; gape, .79; tarsus, .15; middle toe and its claw, .27; claw alone, .12.					
	<u>May 9th.</u>	<u>N. J. N. Y.</u>				
	<u>Lanivireo flavifrons.</u>					First seen.
	<u>Vireo noveboracensis.</u>					First seen.
	<u>Dendroica caerulescens.</u>					First seen.
2,563.	<u>Myiodioctes mitratus.</u>	5.45	8.20	♂ ad.		
	Wing, 2.60; tail, 2.25; bill, measured from nostril, .30; gape, .58; tarsus, .76; middle toe and its claw, .65; claw alone, .18. Irides, hazel. Bill, black. Legs and feet, flesh-color.					
2,564.	<u>Dendroica caerulescens.</u>	5.30	8.10	♂ ad.		
	Wing, 2.50; tail, 2.16; bill, measured from nostril, .30; gape, .55; tarsus, .82; middle toe and its claw, .65; claw alone, .18.					
2,565.	<u>Siurus auricapillus.</u>	6.25	9.60	♂ ad.		
	Wing, 2.93; tail, 2.30; bill, measured from nostril, .38; gape, .68; tarsus, .88; middle toe and its claw, .68; claw alone, .15.					
2,566.	<u>Lanivireo flavifrons.</u>	5.75	9.70	♂ ad.		
	Wing, 3.00; tail, 2.12; bill, measured from nostril, .33; gape, .70; tarsus, .75; middle toe and its claw, .67; claw alone, .21. Irides, hazel. Bill, blue black, whitish along commissure towards the end. Tarsi, feet and claws, bluish-gray.					

over

1883.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
2,567.	<u>Vireo noveboracensis.</u>	5.38	7.85	♂ ad.		
	Wing, 2.45; tail, 2.10; bill, measured from nostril, .29; gape, .39; tarsus, .78; middle toe and its claw, .60; claw alone, .20. Irides, white. Bill, blue-black; commissure white towards extremity. Tarsi, toes and claws, bluish-gray.					
2,568.	<u>Helminthophaga ruficapilla.</u>	4.90	7.50	♂ ad.		
2,569.	<u>Cyanocitta cristata.</u>	11.90	16.85	♂ ad.		
2,570.	<u>Tyrannus carolinensis.</u>	8.75	15.10	♂ ad.		
2,571.	<u>Icterus galbula.</u>	7.75	11.70	♂ ad.		
2,572.	<u>Pipilo erythrophthalmus.</u>	8.00	10.20	♀ ad.		

To-day (May 9) I noticed the first Cow-bird (Molothrus ater).

May 10th. N. S. N.Y.

Helminthotherus vermivorus. - First seen.
Siurus naevius. - First seen. Heard singing sweetly in the lower arm of Gouverneur's Cove, where this species usually occurs.
Nyctiardea grisea naevia. - Mr. Denton visited the Heronry with me; but not a single Heron could be found.
Corvus ossifragus. - A pair of these birds are without doubt breeding on Constitution Island. Mr. Denton notes one there some time ago; and to-day we found a pair that seemed very anxious over our movements and investigation of the Island. They soared overhead and croaked at us all the time we spent on the Island; but the Common Crows build on the island in numbers, so that we did not think it worth while to hunt for a nest since the eggs when found could not be

1883.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	positively distinguished from those of the Common Crow, unless a parent could be secured on the nest.					
	At the lower end of Gouverneur's Cove we found a small colony of Yellow Violets, of the round-leaved species (<u>Viola rotundifolia</u>), the first we had either of us seen growing, although the downy yellow Violet (<u>Viola pubescens</u>) grows profusely in this region. We also found some remarkably large and beautiful specimens of trailing Arbutus (<u>Epigaea spens</u>) which in size & habit resemble closely the New-England examples of it.					
	We procured a mutilated, but large, specimen of the Great Sea Lamprey (<u>Petromyzon marinus</u> L.) from Mr. Van Sternberg, who fishes from Constitution Island. He caught it in a fyke, and kept it for me; but something ate its tail off.					
	We called upon the Messrs Warner. Miss Anna went with us and showed us a patch of "Indian Kinnikinnick" (<u>Arctostaphylos Uva-ursi</u> Spreng.). We also procured specimens of <u>Trillium erectum</u> in Gouverneur's Cove.					
2,573.	<u>Myiodytes nitratu.</u>	5.65	8.90	♂ ad.		
	Wing, 2.75; tail, 2.35; bill, measured from nostril, .32; gape, .60; tarsus, .80; middle toe and its claw, .65; claw alone, .19. Irides, hazel.					
2,574.	<u>Helminthotherus vermivorus.</u>	5.70	9.00	♂ ad.		
	Wing, 2.75; tail, 2.05; bill, measured from nostril, .44; gape, .66; tarsus, .67; middle toe and claw, .66; claw alone, .20. Irides, hazel.					
2,575.	<u>Mniotilta varia.</u>	5.35	8.50	♂ ad.		
	Wing, 2.70; tail, 2.07; bill, measured from nostril, .38; gape, .63; tarsus, .70; middle toe & claw, .72; claw alone, .19.					

1883.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
2,576.	<i>Dendroica maculosa</i> .	5.25	7.85	♂ ad.		
	Wing, 2.40; tail, 2.15; bill, measured from nostril, .27; gape, .50; tarsus, .76; middle toe and its claw, .58; claw alone, .19. Irides, hazel.					
2,577.	<i>Dendroica virens</i> .	5.15	7.90	♂ ad.		
2,578.	<i>Hylocichla mustelina</i> .	8.10	13.50	♂ ad.		
2,579.	<i>Hylocichla mustelina</i> .	7.95	13.50	♀ ad.		

May 11th. H. F. N.Y.

Rain nearly all day. This evening walked to Echo Lake, where I found an adult Musk Rat (*Fiber zibethicus*) swimming about and across the lake. Squatting beside the lake where I had liberated some turtles just before, I noticed the strong scent of musk. Looking up, I saw a large Musk Rat swimming about in front of me. After a while it swam up to shore only a few feet from me, and seated itself upon a stone beneath the surface so that only its head was visible. After watching me for some minutes it swam within a yard or two past me and climbed out upon the bank at a short distance.

May 12th, H. F., N.Y.

2,580.	<i>Tyrannus carolinensis</i> .	8.20	14.00	♀ ad.		
	Wing, 4.30; tail, 3.50; bill, measured from nostril, .54; gape, .98; tarsus, .73; middle toe + claw, .85; claw alone, .29. Irides, hazel. Bill, legs, feet + claws black.					
2,581.	<i>Tyrannus carolinensis</i> .	8.62	15.25	♂ ad.		
	Wing, 4.70; tail, 3.75; bill, measured from nostril, .54; gape, 1.00; tarsus, .78; middle toe and claw, .84; claw alone, .25. Irides, hazel. Bill, legs, feet and claws, black. These two birds were a mated pair. The specimen had no right hand, outer toe. Probably a congenital deformity.					

1883.

2,582.	<i>Cyanocitta cristata</i> .	11.60	16.60	♀ ad.		
	Wing, 5.20; tail, 5.30; culmen, 1.00; gape, 1.17; tarsus, 1.42; middle toe and its claw, 1.17; claw alone, .36; Irides, hazel. Bill, tarsi, feet + claws, black.					
2,583.	<i>Pyrranga rubra</i> (Linn.) <i>Summer Tanager</i> .	7.00	11.40	♀ ad.		
	Wing, 3.60; tail, 2.80; culmen, .60; gape, .77; tarsus, .74; middle toe and claw, .75; claw alone, .22. Iris, hazel. Bill, greenish-olive. Legs & feet, bluish-gray; claws, brownish.					
2,584.	<i>Pyrranga rubra</i> . <i>Scarlet Tanager</i> .	7.33	11.95	♂ ad.		
	Wing, 3.85; tail, 3.05; Culmen, .65; gape, .75; tarsus, .77; middle toe and claw, .72; claw alone, .22. Colors of parts like No. 2583.					
2,585.	<i>Pyrranga rubra</i> . <i>Scarlet Tanager</i> .	7.12	11.35	♂ ad.		
	Wing, 3.55; tail, 2.90; culmen, .65; gape, .80; tarsus, .75; middle toe and claw, .76; claw alone, .26. Colors of parts the same as No. 2583.					
2,586.	<i>Trochilus colubris</i> .	3.65	4.15	♂ ad.		
	Wing, 1.62; tail, 1.14; culmen, .67; gape, .78; tarsus, .17; middle toe and claw, .26; claw, .12.					
2,587.	<i>Dendroica maculosa</i> .	5.05	7.55	♂ ad.		
	Wing, 2.35; tail, 2.10; bill, measured from nostril, .28; gape, .53; tarsus, .72; middle toe and its claw, .60; claw alone, .16.					
2,588.	<i>Myiodiocetes pusillus</i> .	4.95	6.80	♂ ad.		
	Wing, 2.13; tail, 2.05; bill, measured from nostril, .25; gape, .50; tarsus, .70; middle toe and claw, .59; claw alone, .16. Irides, hazel. Maxilla and tip of mandible (1/3) black; mandible, yellow, otherwise. Legs & feet, yellow. Claws, brownish.					
2,589.	<i>Hylocichla aliciae</i> .	7.75	12.75	♂ ad.		
	Wing, 4.18; tail, 3.23; bill, measured from nostril, .39; culmen (about), .57; gape, .89; tarsus, 1.22; middle toe and claw, 1.02; claw alone, about .28. Irides, hazel. Bill, blackish, with basal one-half of man-					

1883.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

-dible, yellow. Legs and feet, pale brown.
Claws, blackish. Testicles, blackish.

Myiodioides pusillus. - First seen. Several.

Myiodioides canadensis. - First seen. Numerous.

Dendroica striata. - First seen. One ♂ ad.

Passerina cyanea. - First seen. One ♂ ad.

Hylocichla aliciae. - First seen.

Hylocichla fuscescens. - First seen.

May 13th. N. F., N.Y.

Cyanocitta cristata. - To-day all the woods were filled with Blue Jays that trumpeted and screamed by dozens at a time. I have before noted this late spring movement.

To-day Ella found a Violet New to this region, *Viola prunellaefolia*.

Before breakfast a man brought me my first specimen of the Lampyrell (*Petromyzon nigricans* Le Sueur). Now I have specimens of both species of this genus taken in the Highlands.

May 14th. N. F., N.Y.

2590. *Dendroica blackburniae*. 5.20 8.05 ♂ ad.

2591. *Dendroica pennsylvanica*. 5.10 7.85 ♂ ad.

2592. *Parula americana*. 4.90 7.75 ♂ ad.

2593. *Quiscalus purpureus*. 12.35 17.65 ♂ ad.

Wing, 5.45; tail, 5.15; culmen, 1.24; gape, 1.37; tarsus, 1.43; middle toe and claw, 1.35; claw alone, .34. Sides, yellow. Bill, legs, feet and claws, black.

2594. *Scops asio*. Light red plumage. 9.40 23.75 ♂ ad.
Wing, 6.85; tail, 3.60; cere, .29; culmen, .59; cere and culmen, .82; tarsus (about), 1.35; middle toe & claw, 1.24; claw alone, .47. Sides, yellow. Stomach contained bird & insect debris.

1883.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

2595. *Tachycineta bicolor*.

6.20 9.45 ♂ ad.

2596. *Sinus motacilla*.

6.40 10.10 ♀ ad.

May 17th.

N. F., N.Y.

2597. *Cyanocitta cristata*.

11.50 16.25 ♀ ad.

Wing, 4.95; tail, 5.20; culmen, 1.08; gape, 1.25; tarsus, 1.33; middle toe and claw, 1.10; claw alone, .32. Iris, hazel. Bill, feet and claws, black.

2598. *Cyanocitta cristata*.

11.25 16.10 ♀ ad.

Wing, 5.00; tail, 5.05; culmen, 1.05; gape, 1.21; tarsus, 1.30; middle toe and claw, 1.10; claw alone, .34. Sides, hazel. Bill, legs, feet and claws, black.

2599. *Passerina cyanea*.

5.50 8.60 ♂ ad.

Wing, 2.65; tail, 2.13; culmen, .44; gape, .50; tarsus, .67; middle toe and claw, .69; claw, .19.

2600. *Helminthogaster vermicivorus*. 5.60 8.75 ♀ ad.

Wing, 2.70; tail, 2.15; bill, measured from nostril, .40; gape, .61; tarsus, .73; middle toe and claw, .66; claw alone, .19. Sides, hazel. Bill, with upper mandible, dusky; lower mandible, flesh-color. Legs, feet and claws, flesh-color.

2601. *Zamelodia ludoviciana*. 8.38 13.20 ♂ ad.

Wing, 4.15; tail, 3.25; culmen, .80; gape, .82; tarsus, .87; middle toe and claw, .93; claw alone, .29. Sides, hazel. Bill, flesh-color, slightly brownish above, and dark on tip of maxilla. Legs & feet, bluish-gray; claws, brownish.

2602. *Zamelodia ludoviciana*. 8.00 12.65 ♀ ad.

Wing, 3.85; tail, 3.15; culmen, .70; gape, .78; tarsus, .88; middle toe and claw, .92; claw alone, .27. Sides, hazel.

2603. *Perissoglossa tigrina*. 5.25 8.40 ♂ ad.

Wing, 2.63; tail, 2.00; bill, measured from nostril, .32; gape, .54; tarsus, .73; middle toe and claw, .65; claw alone, .19. Sides, hazel. Bill, legs, feet and claws, black; soles of feet, dusky-yellowish.

1883.

2,604. *Perissoglossa tigrina*. 5.20 8.42 ♂ ad.
 Wing, 2.70; tail, 2.00; bill, measured from
 nostril, .32; gape, .53; tarsus, .70; middle
 toe and claw, .65; claw alone, .18.
 Irides, hazel. Bill, legs, feet and claws, black;
 Soles of feet, dusky.

Cyanocitta cristata. - The woods were filled
 with them to-day; and dozens of them were
 trumpeting in our apple-trees all day.

Perissoglossa tigrina. - Three of these rare
 Warblers were shot in our cherry trees among
 the blossoms, in company with Humming-
 birds, Indigo Birds, Rose-breasted Grosbeaks,
 Orioles and Purple Finches. Its song is
 simple, distinctly uttered, and has the
 peculiar insect-like tone which so many
 Warblers' songs have. Its little ditty is
 constantly repeated. One of the birds shot
 could not be recovered. They were the first
 seen this year.

Helminthophaga vermicorus. - This species
 is now pretty numerous. Found often in
 dry woods, but not far from wet places.

Gamelodia ludoviciana. - The bodies of
 those skinned had a peculiar and strong
 fragrance, resembling the odor of cherry
 blossoms upon which they feed. Their
 bills smelled sweet.

May 18th. H. F. N.Y.

2,605. *Gamelodia ludoviciana*. 8.30 13.35 ♂ ad.
 Wing, 4.10; tail, 3.37; culmen (about, tip shot off), .70;
 gape (about), .75; tarsus, .89; middle toe and claw,
 .93; claw alone, .26. Irides, hazel. Shot
 while picking cherry blossoms. Several
 were seen in the mountains besides.

1883.

May 19th. H. F., N.Y.
Ranunculus recurvatus L. First seen
 in flower.

May 22nd. H. F., N.Y., Sing Sing, N.Y.
 Took 2.30 P.M. train from New York, and
 stopped at Sing Sing for the balance of the
 afternoon. Found *Viola primulaefolia*,
V. rotundifolia, *V. pubescens*, *V. cucullata*
 and *V. canina*. The Primrose-leaved
 Violet was abundant, but not a single
 blanda could be found. The only *Trillium*
 found was *T. erectum*. The *Anemone*
memorosa was also plentiful.

I saw *Empidonax acadicus* (as I
 presume. Have not had time to look it up)
 for the first time. A single bird was
 found in a deep ravine, grown with good
 sized hemlocks (*Abies canadensis*), beside
 a brook. It uttered at occasional
 intervals a loud yelp.

Oporornis formosus was not seen.

Whippoorwill (*Caprimulgus vociferus*)
 were singing about home all night.

2,606. *Empidonax traillii* 6.40 9.65 ♂ ad.
 Wing, 2.97; tail, 2.70; bill, measured from
 nostril, .38; gape, .74; tarsus, .60; middle
 toe and claw, .53; claw alone, .16.

Irides, hazel. Bill, with maxilla, black;
 mandible pale yellow. Tarsi toes and
 claws, brownish-black. Shot at Sing Sing, N.Y.

2,607. *Helminthophaga pinus*. 5.00 7.70 ♂ ad.
 Wing, 2.42; tail, 1.98; bill, measured from
 nostril, .33; gape, .54; tarsus, .70;
 middle toe and its claw, .58; claw alone,
 .18. Irides, hazel. Bill, black. Tarsi, toes
 and claws, dusky yellowish-brown. Sing Sing.

1883.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

May 25th. N. H. M. Y.

This morning Ella called me down stairs to rescue the family of week-old Robins (Merula migratoria) that lived in the Norway Spruce (Abies excelsa) in front of our house. A Red Squirrel (Sciurus hudsonius) had already bitten and thrown two of the young birds to the ground. I saw the old bird dash fearlessly at him it and dash it from the branch to one below upon which it caught, and immediately climbed up toward the nest, when my shot killed it. It proved to be a large, pregnant female Red Squirrel. I have preserved one of the bitten young Robins. Before I got down, the old Robins had once driven the Squirrel down from the nest; but it climbed up again while Ella was looking and threw another little Robin down upon the ground.

2,614.

Merula migratoria.

— — Pullus.

Found a Black Snake (Bascanium constrictor) lying in the road, which ~~weighed~~ measured four feet, ten inches. It was too stale to permit of preservation.

After dinner Ella and I took a walk, but found no birds of interest saving the Yellow-throated Vireo (Lanius flavifrons) one of which Ella shot:

2,615.

Lanius flavifrons.

6:00 9:80 ♀ ad.

Wing, 3.10; tail, 2.15; bill, measured from nostril, .31; gape, .69; tarsus, .75; middle toe and claw, .65; claw alone, .18. Irides, hazel. Bill, with maxilla above & tip, black-black; the residue pale blue, shading to whitish on middle third of bill along gape. Legs and feet, dark bluish-gray. Claws, horn-color,

1883.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

marked with dusky stripes.

In the evening Mr. Denton accompanied me to Crow Nest Ravine, where we found four flowering specimens of Habenaria viridis, var. bracteata. Near the ravine we found buds of Liparis lilifolia about ready to open, and good sized leaves of Habenaria lacera. In the Ravine itself we found a plant of Goodyearia pubescens, scapes of Aplectrum hyemale having buds nearly ready to open, and a few plants of Cypripedium pubescens, one of which had an opening bud. The root leaves of Habenaria tridentata were numerous. Orchis spectabilis grew in fair abundance; and one clump of plants was the handsomest we ever saw. Six large flowering scapes, and two last years seed-scares, together with twice that number of unflowering leaves grew in a compact bunch. We dug the whole thing up with plenty of earth, and shipped it entire to Prof. W. W. Bailey of Providence, R. I.

Thus eight species of Orchids were found in and about the Ravine this afternoon.

The leaves of the Wild Garlic (Allium tricoccum) are, many of them, just beginning to fade, the flowers coming out after the leaves die.

The leaves of the Ginger (Asarum canadense) are satiny and very beautiful now. The seed-pods of the Blood-root (Sanguinaria) are large and filled with seeds. The Blue Cohosh (Caulophyllum thalictroides) has dropped most of its flowers, and the frutage begins to appear. We found Viola cucullata palmata, its most typical form, in the old spot, near the ravine. I found the Purple Oxalis (Oxalis purpurea), the first found by either of us. Miss Warner has, however, found it on Constitution Island, years ago.

1883.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

May 23rd. H.S.N.Y.

To-day drove to Mountville (nearly 10 miles) to see a patient upon whom we operated for lacerated perineum last Sunday.

On the north side of Dunderburg Mountain I found a number of interesting plants. Among them were, *Cypripedium pubescens* quite numerous and scattered, *Arctostaphylos* abundant and large, *Viola primulaefolia*, *V. pubescens*, and *V. rostrata*, the latter quite plentiful: the two first very abundant. The Purple Trillium (*Trillium erectum*) is very plentiful there; likewise *Aplectrum hyemale*.

Near Highland Lake I found fourteen flowering plants of *Cypripedium acaule* growing within an area ten feet square. I procured a double specimen having two flowers + two sets of leaves from one root. Besides the flowering plants there were dozens of plants not in flower - sterile.

On Dunderburg I found a number of Worm-eating Warblers (*Helminthophaga vermivorus*), and saw one bird utter repeatedly a rattling note that suggests the "song" of a *Cecada*, and also resembles somewhat in character the peculiar notes of *Helminthophaga chrysoptera* and *Parula americana*. I have tried for many years to identify the author of this curious sound. I always believed it to be produced by a bird, although several persons to whose attention I directed it supposed that it was produced by some insect.

I saw there an *Euphonia* which possibly may have been *E. acadicus*, but as *E. traillii* is generally numerous here in spring it was probably that species. *Myiodynastes nigratus* was numerous.

1883.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

May 24th. H.S.N.Y.

2,608. *Geothlypis trichas*. Shot by E.W.M. 5:10 6:60 ♀ ad.
Wing, 2.00; tail, 1.98; bill, measured from
nostril, .31; gape, .55; tarsus, .80; middle toe
and claw, .68; claw alone, .18. Irides, hazel.
Bill, blackish. Tarsi, flesh-color; feet, pale brownish.

2,609. *Dendroica striata*. 5:60 9:10 ♂ ad.
Wing, 2.90; tail, 2.25; bill, measured from
nostril, .30; gape, .56; tarsus, .77; middle toe
and its claw, .68; claw alone, .20. Irides, hazel.
Bill, blackish, with basal one-half of mandible
yellowish. Tarsi toes and claws, yellow; the
latter brownish-yellow. E.W.M.

2,610. *Pyrranga rubra*. Shot by E.W.M. 7:30 11:55 ♀ ad.
Wing, 3.65; tail, 3.95; culmen, .67; gape,
.79; tarsus, .75; middle toe and claw, .79;
claw alone, .25. Irides, hazel. Bill, olive-
green. Tarsi and toes, lead-color; claws
brownish. Shot by Ella W. Mearns.

2,611. *Tyrannus carolinensis*. 8:15 14:10 ♀ ad.
Wing, 4.40; tail, 3.38; bill, measured from
nostril, .52; gape, .96; tarsus, .74; middle
toe and claw, .81; claw alone, .26. Irides, hazel.
Bill and feet, black. E.W.M.

2,612. *Carpodacus purpureus*. 6:15 10:05 ♂ ad.*
Wing, 3.05; tail, 2.38; culmen, .43; gape,
.55; tarsus, .65; middle toe and claw, .74;
claw alone, .21. Irides hazel. Shot by E.W. Mearns.

2,613. *Urosylvia olivacea*. 6:40 10:40 ♂ ad.
Wing, 3.30; tail, 2.40; bill, measured from
nostril, .40; gape, .81; tarsus, .73; middle toe
and its claw, .65; claw alone, .19. Irides,
red. Bill, with maxilla, blue-black;
mandible, bluish horn-color. Legs,
feet and claws, bluish gray.
Shot by Ella W. Mearns, near Echo
Lake, Mearns House.

* Plumage entirely feminine; no red or yellow whatever.

1883.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	May 27th.	N. S. N. Y.				
	To-day drove to Mountville to see a patient. Mr. E. S. Denton accompanied me. We afterward explored the ravines of Dunderberg and found <i>Cladonia viridis</i> var. <i>bracteata</i> , <i>Cypripedium parviflorum</i> , and <i>C. pubescens</i> all flowering.					
	We surprised an old hen Pheasant (<i>Bonasa umbellus</i>). She gave a loud Squall, and whirled loudly as she beat her wings upon the ground in front of us. The little chicks, only two or three days old and about fifteen in number, at first piped out most lustily in their first surprise and bewilderment; but soon they recovered, and piled over each others backs pell mell in their efforts to escape and hide. Several of them rushed directly into the water of the brook close beside which we discovered them. One swam clear across. Another was carried a distance down the stream and then crawled under a stone beside the bank. Others hid under stones and debris in the brook; and one fellow was actually drowned in its effort to hide. We found it floating dead upon the water when we returned to the spot some time afterward. The old bird dragged herself over the ground with a great clucking; but when I ran rapidly after her, she took to wing and flew a little way off, and for a few minutes remained silent. We had captured three of the little chicks, and now examined to see what sort of hiding places the balance of the flock had chosen. One was packed like a sardine between two stones in the brook, with its head and a streak					

1883.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	of its back visible; another was wedged tightly between a stone and some herbage growing beside it. Two little fellows were found under shelving stones in the brook, running in imminent risk of drowning. The drops of water were trickling off of one bird's head in rapid succession, and it was found fairly driected. I truly believe that, had I not rescued it from the perilous situation it would surely have died rather than be discovered.					
	We moved a little way from the spot with the three little ones chirruping in our pockets, to witness the mother's return. At first she uttered a loud frog-like note, which we interpreted as a warning to her little ones to "lie low," an admonition which they certainly heeded. Soon she changed the note to one which bore some resemblance to a puppy dog's whine; and, after a while, we occasionally heard a feeble chirping from the chicks. The old bird uttered a great variety of notes, and grew more and more vociferous. At length her maternal anxiety more than counterbalanced her instinctive fear, and she began to approach. At first slowly, until we had her clearly in view; then rapidly and fearlessly she pressed close past us, watching us meanwhile, with her head rapidly extended and then retracted, her tail raised and fully spread, and clucking and calling vehemently at every step. When she reached the spot where the young had scattered, her note changed to a loud, peculiar and difficult note to describe, and when she found the dead bird drowned in the brook, as we supposed, then she seemed fairly frantic leaping about and uttering loud whines, the grief expressed in them was most rending.					

1883.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

Then she slowly succeeded in coaxing the little ones out of their respective hiding places, till we could now and then distinguish a little Partridge running after the old bird. She collected and escorted them to a distance; and we afterwards, were unable to discover them.

We now walked up to the old bird with the little ones in our hands. They answered their mother's call; and then she came intrepidly forward, opening her mouth widely to admit the same grief-stricken sounds which we had heard before when she discovered the drowned bird. We put the birds down upon the ground intending to restore them to her; but she suddenly whirled off to a stone at some distance, and stood upon it watching us for a few minutes, then flew to the top of a stump and eyed us again. Then she wandered away through the herbage. We followed; but she flew off to a distance. Hearing that she would not return to care for the little birds we carried them home along with the drowned bird. In the evening we got one of them to peck at some wheat Indian meal; but, after that, they refused to eat.

2,616. Bonasa umbellus. — — pullus

May 28th H. F., N.Y.

The Partridges (Bonasa) refuse to eat. One of them drowned itself in a mix of water in a salt-cup.

2,617. Bonasa umbellus. — — pullus

May 29th.

Last night we opened the little Partridges' mouths and made them swallow a little food; but both of them were dead this morning.

1883.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

2,618. Bonasa umbellus. — — pullus.
2,619. Bonasa umbellus. — — pullus.

May 31st. H. F., N.Y.

This evening at twilight, I went out for Whippoorwills and had the good fortune to shoot two with the little gun, — a mated pair.

2,620. Caprimulgus vociferus. 10.50 19.30 ♂ ad.

Wing, 6.35; tail, 5.50; bill, measured from nostril, .31; gape, 1.35; tarsus, .70; middle toe and claw, .98; claw alone, .31. Irides, dark hazel. Bill, black. Tarsi, brown anteriorly; fleshy above and posteriorly. Toes, brown, shading into blackish towards tips. Claws, nearly black; the "comb" on inner side of middle toe, fleshy-brown.

2,621. Caprimulgus vociferus. 9.10 17.90 ♀ ad.

Wing, 5.96; tail, 4.40 (Tail moulting.); bill, measured from nostril, .31; gape, 1.25; tarsus, .64; middle toe and claw, .90; claw alone, .29. Colors, just about the same as 2,620.

June 2nd. H. F., N.Y.

2,622. Icteria virens. 7.60 10.25 ♂ ad.

Wing, 3.12; tail, 3.25; bill, measured from nostril, .54; gape, .82; tarsus, 1.10; middle toe and its claw, .94; claw alone, .26. Irides, hazel; bill, blue-black. Legs and feet, dark lead.

2,623. Icteria virens. 7.00 9.60 ♂ ad.

Wing, 2.87; tail, 2.95; bill, measured from nostril, .40; gape, .75; tarsus, 1.00; middle toe and its claw, .83; claw, .19. Irides, ~~toe and its claw,~~ hazel; Bill, blue-black. Tarsi and toes dark bluish-gray.

June 4th H. F., N.Y.

Amphelis cedrorum. — Several small flocks were seen here to-day. They are the first

1883.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

I have seen since last autumn. Although I collected a number of specimens of Birds during the winter and spring just passed, this species was not once noted, although it is usually an abundant species here during the entire year.

This evening Mr. Louis A. Zerega and myself took Edward, the gardener, and drove, after tea, to a Kingfisher's (Ceryle alcyon) burrow, which we dug out. The bank was a slight excavation beside the road in nearly level ground, and the bank only four feet high. We found the nest at about six feet from the opening of the burrow. The burrow was straight for five feet, then turned at a little less than a right angle to the left, where an excavation about 18 (eighteen) inches or more in diameter was discovered. The sand at the bottom of the cavity was loose, and no nest-building materials whatever had been used. The nest contained only three young birds and one nearly hatched egg, whose shell was cracked, and the shells of the eggs which were already hatched, besides the female bird, which I caught by the bill upon the nest, and drew out of the hole. When I first reached her, I could barely touch the tip of her bill, which was slightly open. She did not bite until withdrawn from her burrow; when she took hold of Mr. Zerega whenever occasion offered and bit severely. We were sorry to have opened up the burrow since a set of eggs could not be obtained; but, as it would be cruel to allow the little ones to starve, we resolved to kill the female

1883.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

and young, which were from one to three days old, and looked as if the tips of their upper mandibles had been snipped off with a scissors. We put them in alcohol, and preserved the parent as a cabinet skin.

Empidonax flaviventris. - Mr. Zerega shot a fine female specimen to-day. A late record! 13.00 22.05 ♀ ad.

2,624.

Ceryle alcyon.

Wing, 6.15; tail, 3.90; culmen, 2.00; gape, 2.85; tarsus, .49; middle toe and claw, .92; claw alone, .37. Irides, dark hazel.

I found a No. 6 shot inside of its skin, although we never fired at it. Some one else had shot it.

June 13th.

H. G. M. J.

In the swamp below our stable I found a nest of the Spotted Turtle (Nanemys guttata), which contained three eggs, which were of course white in color, equally round at both ends and measuring, respectively, 1.35 X .70; 1.29 X .68; 1.31 X .68.

The eggs were placed in the mossy turf on top of a stone in a bog in the swamp. A small shallow hole had been excavated by the Turtle, the eggs deposited in it and covered up.

June 14th

H. G. M. J.

Hemithysanota chrysoptera. - Mr. Burnstead and myself found a pair that were doubtless breeding, west of Canby's Swamp. The male sang loudly. We could not find the nest.

Pyrausta rubra. - Several pairs were found, the males singing sweetly. The swamp blade orchid (Liparis lilifolia) is now in full flower, and is fairly common hereabouts.

1883.

Collecting Number. NAME. DATE. LOCALITY. Length. Stretch. Sex and Age.

2,625. *Corvus brachyrovus*. July 7th. H. I. N.Y. — ♀ juv.
Mr. Arthur Taylor shot a young Crow, a bird of the year, and one of a family reared below the house.

Sept. 8th. H. I. N.Y.
To-day Mr. Edward S. Denton sent me a specimen of *Helianthus giganteus* which he got on Hackensack Meadows where the train stopped a few moments. It is the first I have seen.

Sept. 4th. H. I. N.Y.
My boy, Martin, and myself went fishing to Highland Lake, where I shot two fine Ospreys (*Pandion haliaetus carolinensis*). They fed at the Lake every evening, coming in from the Hudson towards Dundown.

2,626. *Pandion haliaetus carolinensis*. 23.75 67.00 ♀ ad.
Wing, 18.80; tail, 9.00; Culmen, measured from base of cere, 1.60; culmen, .35; cere, .35; gape, 1.75; tarsus, 1.20; middle toe, 1.95; its claw, 1.21; inner claw, 1.25; outer claw, 1.27; hind claw, 1.25. Iris, yellow. Bill, blue-black. Feet, pale greenish. Stomach filled with fish.
Caprimulgus vociferus. — Heard.

Sept. 6th.
Caprimulgus vociferus. — Several heard. One shot at Highland Lake.

Sept. 15th.
Went fishing to Highland Lake with Mr. J. Senna. Caught a large Black Bass, which weighed 3 lbs. It was skinned &

1883.

mounted dry. Sex, ♀. Gizzard filled with large Grasshoppers. Length 15 inches, ^{caudal 2 1/2 in.} ~~excl. c.~~
Pandion haliaetus carolinensis. — Several were seen fishing at the Lake and one fine one was shot.

An Orchis (*Corallorrhiza odontorhiza*) which I have hitherto regarded as somewhat scarce, was found in large beds. I counted up to 50 specimens growing within a space one yard square.

A large Gray Squirrel (*Sciurus leucotis carolinensis*) ran across the road beside the Lake, where there were many fresh leaf-nests.

We caught the largest specimen of the Long-eared Sun-fish (*Lepomis auritus*) that I ever saw. I have no balance to weigh it; but its measurements are: length (exclusive of caudal), 6.25; greatest depth, 2.75.

September 26th. American Museum Nat. Hist.
Came to New York to attend and help to found the American Ornithologists' Union at its first meeting in Library of the American Museum. I had the pleasure of meeting many of the ornithologists with whom I had not previously met.

One of the delegates to the Union was Mr. Montague Chamberlain of St. John, New Brunswick, with whom I had long corresponded. The old ornithologist, Mr. Thos. McIlraith of Hamilton, Canada West, was present; and I found him to be extremely agreeable.

Capt. Charles E. Bendire of Fort Klamath, Oregon and Dr. R. W. Shufeldt, U.S.A. from Louisiana were present & had travelled all the way from their posts in order to be here.

1883.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

The other founders of the Union are :-

Prof. J. A. Allen, Cambridge, Mass.

C. Aldrich, Iowa.

Prof. S. F. Baird, Washington, D. C.

H. B. Bailey, New York.

C. F. Batchelder, Cambridge, Mass.

Eugene P. Bicknell, Riverdale, New York.

William Brewster, Cambridge, Mass.

Nathan Clifford Brown, Portland, Me.

Charles B. Cory, Boston, Mass.

Prof. Elliott Coues, M. D., Washington, D. C.

D. G. Elliot, Staten Island, New York.

Dr. A. K. Fisher, Sing Sing, New York.

Dr. J. B. Holder, New York City.

Dr. C. Hart Merriam, Locust Grove, Lewis Co., N. Y.

Dr. D. W. Prentiss, Washington, D. C.

H. A. Purdie, State House, Boston, Mass.

Robert Ridgway, Smithsonian Inst., Washington, D. C.

Dr. J. M. Wheaton, Columbus, Ohio.

It will be noticed that the Medical Profession was represented at the first congress of the American Ornithologists' Union by no less than seven Doctors.

The first Congress lasted three days, and adjourned on September 28th.

A Constitution was adopted and the following officers elected:-

President, Prof. J. A. Allen.

Vice - Presidents, Dr. Elliott Coues and Mr.

Robert Ridgway.

Secretary and Treasurer, Dr. C. Hart Merriam.

Council of the Union, consists of the officers,

ex-officio, and of Prof. Spencer F. Baird,

Mr. Geo. N. Lawrence, Mr. William Brewster,

Mr. H. W. Henshaw, and Mr. Montague

Chamberlain.

1883.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

Six committees were appointed as follows:-

a. The Classification and Nomenclature of North American Birds: Messrs. Ridgway, Allen, Henshaw, Brewster, Coues.

b. The Migration of Birds: Messrs. Merriam, Brown, Purdie, Wheaton, Chamberlain, Grinnell, Henshaw, Cory, Merrill, Fisher, Mearns, McIlhenny.

c. Avian Anatomy: Messrs. Shufeldt, Merriam, Jeffries, Coues.

d. Oölogy: Messrs. Bendire, Bailey, Brewster, Ridgway, Merrill.

e. Faunal Areas: Messrs. Allen, Ridgway, Bicknell, Merriam, Fisher, Mearns.

f. Eligibility or Ineligibility of the English House Sparrow in America: Messrs. Holder, Purdie, Chamberlain, Brown, Bicknell.

October 1st. Am. Mus. Nat. Hist, N.Y.

I came down to the Museum to arrange my collections, which I have arranged to store in in this building now, and in the future. The Museum is to pay me the same salary that I would receive should I accept a proffered contract as acting Assistant Surgeon, U. S. Army. I am engaged to label the European Birds in the collection & select from my collection for the Museum a series of specimens which are lacking in their collection, to make as perfect as possible their collection of mounted birds.

My intention is to use as an authority for nomenclature the "Ibis" List of British Birds compiled by a Committee of the British Ornithologists' Union. London. 1883, as far as it covers the ground, and

1883.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

Dresser's "Birds of Europe" for the rest.

I spoke to Prof. Bickmore last summer of the necessity of buying a copy of Dresser's work for me to have to use in labelling the collection. It has not yet been purchased; neither is there a copy of the Ibis list to be had. I have just sent for a copy of the Ibis list, myself; and have advised them to get a copy of "Dresser" for their library.

October 16th. Highland Falls, N.Y.

To-day I went to Highland Falls, to meet Mr. L. W. Cass, who wished to see our house - "the Weems House" - with reference to renting it. I found a handsome specimen of the Double-crested Cormorant which Mr. "Josh." Ward had shot at Cornwall, on the Hudson, on October 10th. I brought it to N.Y. and had Mr. Conway mount it for the exhibition collection of the American Museum.

2,627. *Phalacrocorax dilophus* (Sw. & Rich.) Nutt. 33.60 53.65
Wing, 12.90; tail, 7.80; culmen, 2.30; gape, 3.60;
tarsus, 2.40; middle toe and claw, 3.20; claw, .47; ~~middle toe & claw, 3.20;~~ outer
toe and claw, 3.90; claw alone, .38; 2nd toe &
claw, 2.33; claw alone, .47; hallux & claw, 1.57;
claw alone, .50. Iris, green. Feet and
claws jet-black. Gular sac, orange.

1883.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

2,628. *Nuncio hyemalis* (Linn.). 6.30 9.90 ♂ ad.
Wing, 3.12; tail, 2.95; culmen, .33; gape, .50;
tarsus, .80; middle toe and its claw, .78;
claw alone, .21. Irides, hazel. Bill,
flesh-color; extreme tip of maxilla, black.
Legs & feet, dusky-brownish. This bird
flew into the Museum and, struck against
a glass, and fell dead. Mr. Benedict,
naturalist of the Fish Commission Steamship,
picked it up and handed it to me,
while still warm, remarking that it
had offered itself, a sacrifice to the
"Study Collection", which I was endeavoring
to found in the American Museum. I
therefore skinned it and will enter it
first upon the register of bird-skins
in the "Study Collection of Birds of
the American Museum of Natural History."

November 18th to 22nd Fort Miller, Washington
County, New York.

During five days which I spent at
my grandfather's house, on "The Ridge" at
Fort Miller, I had two good hunts in
the old woods where I used to shoot Gray
Squirrels and Partridges; and Uncle Will
accompanied me with the same zest as
in bygone days. My cousin, John Blawie,
who was a kid then is now a six-foot
man, and accompanied us to the woods
and there killed his first Grouse in a
hemlock-tree, with an old musket which
uncle Will carried through the war.

Perhaps the pleasantest thing I noted
in "Grandpa's woods", aside from the
big trees which have always been there,

1883.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

was the beautiful carpeting of green club-mosses. When studying the same genus at home in the Highlands - the genus Lycopodium - I had often remembered and spoken to my wife of the beauty of those growing so luxuriantly in my grandfather's old woods; and we wondered if some of the species would not prove to be new to us. Sure enough, when I came to scrutinize closely the beautiful forms of the Lycopodiums under the big hemlocks and amongst the firs and maple saplings, where they grew luxuriantly, I found a species that we never saw in the Highlands of the Hudson, and beside it were each of the four species of the genus which we before knew. Several other interesting plants we noted casually, some of which I will note here, with the Club-mosses: -
Lycopodium lucidulum, Michx. In fruitage.
Lycopodium dendroideum, Michx.
Lycopodium complanatum, Linn. - I have never seen this species growing so luxuriantly, nor have I seen it before with the fruitage attached, and therefore gathered specimens for our collection.
Lycopodium clavatum, Linn. This species, so uncommon in the Highlands was also rather rare here, all of the other species being common.
Lycopodium annotinum, L. - This species was abundant and very beautiful. It was also fruiting in considerable quantities. We have never yet seen the fruitage of L. clavatum or of L. dendroideum, but have specimens of the

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

fruitage of the other three species.
Monotropa hypopitys, Linn. The largest specimens of the Many-flowered Indian Pipe that I ever saw were growing in a ravine in the woods - last summer's plants, of course.
Chimaphila umbellata, Nutt. Abundant. I did not see C. maculata, which is abundant in the Highlands.
Pyrola rotundifolia, Linn.
Pyrola elliptica, Nutt. - Both species were quite plentiful.
Epigaea repens, Linn. - One day I drove to Lakeville with my cousin Carrie, and we found arbutus growing plentifully in places beside the road and in fence-corners - localities in which one would always look in vain in our Highlands, where it prefers north slopes of hills under laurels (Kalmia latifolia) and shrub-oaks.
Mitella diphylla, Linn. Abundant.
Camposorus rhizophyllus, Link. We found a quantity of the walking fern growing on a large rock near the Lake at Lakeville, where I collected a specimen which I afterward lost.
The end of November is not a favorable season for botanizing. I should like to visit the woods in spring and summer. The principal forest trees are oak, birch, hickory, birch, maple, white ash, and sycamore. The conifers are: Tamarack, hemlock, balsam spruce (P. balsamea), and white pine (Pinus strobus). A peculiar feature of the forests there is the constant intermingling, in each bit of woodland, of coniferous and deciduous trees.

1883.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

The following Mammals were noted:—

Lepus sylvaticus.

Lepus americanus.

Arvicola riparia.

Vulpes fulvus.

Mephitis mephitis.

Sciurus carolinensis.—A black specimen

Arctomys monax.—One seen Nov. 29th. I saw only "Grays"

Sciurus hudsonius.—Abundant.

The following Birds were observed:—

Sialia sialis.

Regulus satrapa.

Sitta canadensis.—A number of these merry little birds were found disporting themselves amongst the hemlocks in grandfather's woods. They are great favorites of mine!

Sitta carolinensis.

Pinicola enucleator.—I saw a small flock of Pine Grosbeaks, near some Norway Spruce trees which were covered with cones, a little way south of Albany, beside the Hudson, on Nov. 18th.

Astragalinus tristis.—Noisy, restless, roving flocks of the Goldfinches were seen in the woods, where they were feeding exclusively upon the seeds of a large black species of birch-tree. Out of the woods I saw them eating seeds in the garden, and feeding on the alders beside the kills or creeks.

Spizella montana.—Not numerous.

Junco hyemalis.—Not common.

Melospiza fasciata.—Abundant beside the Hudson at Fort Miller.

Passer domesticus.—These pests were abundant at Fort Edward & at Ft. Miller.

1883.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

Lanius borealis.—One seen near Lakeville.

Circus hudsonius.—Several seen between Albany and Troy.

Buteo borealis.

Buteo lineatus.

Falco columbarius.—One seen near Albany.

Accipiter fuscus.—One seen.

Bonasa umbellus.—Common.

Anas obscura.—A flock below Albany.

Podilymbus podiceps.—We had rare sport with a little fellow that we cornered in shallow water beside the Hudson south of Fort Miller, on the 20th.

Larus delawarensis.—Very numerous upon the ice below Castleton, and noted all the way from Poughkeepsie to Troy on the 18th.

Corvus frugivorus.—Extremely abundant in many places upon the ice.

Larus argentatus.—Abundant as high as Albany and noted above Troy.

Cyanocitta cristata.—Abundant.

Peris atricapillus.—Abundant, and never so jolly and companionable.

Picus villosus.—Only one pair in an orchard.

Picus pubescens.—Only one specimen.

Hylatomys pileatus.—Uncle Will saw one in his orchard a short time ago.

Ceryle alcyon.—Near Albany Nov. 18th.

2629. Sitta canadensis. Nov. 20. 5.00 8.35 ♂ ad.

2630. Picus villosus. do 9.55 15.70 ♀ ad.

2631. Astragalinus tristis. do 5.35 8.95 ♂ ad.

All of the above specimens, including the tough Hairy Woodpecker (No. 2630) were shot with my miniature shot gun, with a barrel only fifteen inches in length, and using dust shot.

1883.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

November 24th. New York City.

To-day, as I walked straight across the Park from 72nd St. + 5th Avenue, I noted a small flock of European Goldfinches (Carduelis elegans), and a pair of Cardinal Grosbeaks (Cardinalis virginianus), and heard both the White-throated and Fox Sparrows singing sweetly. I counted no less than sixteen Gray Squirrels.

November 25th. New York City.

Quiscalus purpureus. - Two Cow Blackbirds were noted in the Park; and the Cardinals were seen near the same spot where before seen.

December 3rd. New York City.

A small flock of five Cow Blackbirds (Molothrus ater), two males and three females were seen near the Lake in Central Park. They were lively, and active. I have never seen them hereso late, although I saw them in some numbers in Maryland, Delaware and ^{middle of} Southern Pennsylvania as late as ^{the} December, 1881.

December 23rd. H. F., N.Y.

One Pine Grosbeak (Pinicola enucleator) was seen and it was heard at an other place. Birds of all kinds are exceedingly scarce this winter.

December 31st. H. F., N.Y.

Louis A. Bergea and myself spent the afternoon in collecting birds, but saw

1883-'84.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

nothing of special interest. Song Sparrows (Melospiza fasciata) were quite numerous along the border of Consook Marsh, where I shot three for the purpose of adding their dimensions to the tables which I am making, with a view to exhibiting the variation in measurements between winter and summer birds of this species. I have prepared similar tables of measurements of the Bluebird (Sialia sialis) and Robin (Turdus migratorius).

2632. Melospiza fasciata 6.27 8.55 ♂ ad.
Wing, 2.62; tail, 2.83; culmen, .50; gape, .57; tarsus, .86; middle toe and claw, .85; claw alone, .22. Consook Marsh. Iris, hazel.

2633. Melospiza fasciata 6.75 8.90 ♂ ad.
Wing, 2.75; tail, 3.00; culmen, .51; gape, .54; tarsus, .85; middle toe and claw, .85; claw alone, .23. Iris hazel.

2634. Sialia sialis 6.88 11.80 ♂ ad.
Wing, 3.75; tail, 2.60; bill measured from nostril, .35; gape, .78; tarsus, .76; middle toe and claw, .85; claw alone, .22. Iris, hazel.

2635. Sialia sialis 6.94 12.25 ♂ ad.
Wing, 3.85; tail, 2.63; bill measured from nostril, .35; gape, .81; tarsus, .76; middle toe and its claw, .85; claw alone, .22. Iris, hazel.

January 20th. H. F., N.Y.

Mr. Edward S. Denton spent the afternoon and evening with us. He brought a collection of plants which Prof. W. W. Bailey of Providence R. I. had sent him, and which he insisted upon our sharing with him in every case where there were duplicates. Among them were a large number of types of genera, which will be a great help to us in working up grasses + sedges.

1884.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

During the afternoon a flock of Pine Grosbeaks (*Pinicola enucleator*) appeared in the Norway Spruce tree (*Abies excelsa*) in front of our house. Ella saw them, and pointed out an old male. I was anxious to procure an adult red male; but when I found my gun and saw that it was empty, and all of my cartridges had been fired, and there was no shot in the house, the prospect of getting this bird did not look favorable; but I found three old shells which belonged to Dr. Merriam's little Dexter Smith shotgun with which I killed an Eagle and Dr. Merriam killed a Bear, and soon extracted the shot and loaded three cartridges for my ten bore. By the time I was ready the birds had left the Norway Spruce, but were found eating the seeds of the cones, in a large White Pine (*Pinus strobus*) on the north side of the house. Ella was looking at them when they left the Spruce-tree, and said "I expected to see the male bird fly first and lead the flock, but the females flew away first."

Before I could see the male bird, the whole flock flew back to the Spruce, where I fired at the red bird as it sat on the topmost branch of the tree. My shot wounded it and killed a young male in female dress. By the feathers which dropped, I was fortunate enough to track and secure the red one, also, for it fell at the end of our place, where by a good fortune, I noted the spot upon the snow where it had fallen, and found it dead. It proved to be an unusually fine adult red male.

1884.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

2636. *Pinicola enucleator* 9.20 14.25 ♂ ad.
Wing, 4.60; tail, 3.95; culmen, .60; gape, .60; tarsus, .86; middle toe and claw, .95; claw alone, .36. Iris, hazel. Bill, blackish, with an olive tinge at the base of the lower mandible. Legs, feet and claws, black.

2637. *Pinicola enucleator* 9.05 13.80 ♂ juv.
Wing, 4.30; tail, 3.85; culmen, .57; gape, .60; tarsus, .89; middle toe and claw, .90; claw alone, .36. Iris, dark hazel. Bill blackish, with an olive tinge on basal half of lower mandible. The bill is not so black as in the adult, and the olive more extended on the under bill. Legs, feet and claws, black. The stomachs of both specimens contained seeds of the White Pine (*P. strobus*), Norway Spruce (*A. excelsa*) and maple (*Acer* ?).

January 25th Central Park, N.Y. City.
Cardinalis virginianus. - A pair seen at the northwest corner of the Lake, near the bridge, where I have seen this pair nearly every day during this winter, but never singing. Their only note is a sharp chip. Yellowbirds (*Astragalinus tristis*) are seen every day, feeding exclusively upon the seeds of birch trees.

Note: - On Dec. 31st, 1883, some Eider Ducks (*Somateria mollissima*) were sent to Fulton Market, New York City. I purchased two females. They were shot in Labrador about Dec. 25th, 1883. They were sent to Mr. Leongay's place to be skinned, and are now in my collection.

2638. *Somateria mollissima* 23.00 37.00 ♀ ad.
2639. *Somateria mollissima* 23.00 36.00 ♀ ad.

1884.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

2640. January 9th. Highland Falls, N.Y.
Archibuteo lagopus sanctijohannis (Omel.).
 Ridgway. American Rough-legged Buzzard. 34.75 55.50 ♀.
 Wing, 17.25; tail, 10.00; culmen, including cere, 1.37; culmen alone, .94; cere, .56; gape, 1.73; tarsus, 2.65; middle toe and claw, 2.25; toe alone, 1.52; claw, .83.
 Irides, pale brownish-yellow. Cere, greenish-yellow. Bill, blue-black; head-color at base except on maxilla in front of cere. Feet, lemon-yellow; claws, black.
 Shot at Forest of Dean, near Highland Falls, by John Redner, who brought it to me.

January 28th. H. F., N.Y.
 Walked across the Hudson from Havens' Dock to Garrisons. While waiting for the train I walked from Garrisons Station down to Mr. Osborn's house and into the premises of Mr. Hamilton Fish, then returned to the station just in time to catch the train. The following birds were noted:
Spizella montana. - A few were found huddled up against the cliffs at Benny Havens'; and the Snow-birds likewise sought shelter there, owing to the bitter cold, for the Hudson is thickly frozen over.
Junco hyemalis. - Was also seen among the hemlocks, near Garrisons.
Astragalinus tristis. - Feeding, in flocks, upon the seeds of the hemlock (A. canadensis).
Melospiza fasciata. - One upon the Railroad, just below Garrisons.
Carpodacus purpureus. - A few were seen on Hamilton Fish's place, where I saw them feeding in the same trees with a flock of Pine Grosbeaks.

1884.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

Pinicola enucleator. - A good-sized flock was seen feeding upon the seeds of the Sugar Maple (Acer saccharinum) and Hemlock (Abies canadensis) on Mr. Osborn's place. An other flock, smaller in size, was found in Mr. Fish's place. They were very gentle; and there were several adult red males. I saw two fine red birds at once; and I think there were more. They were actively engaged in eating the tender buds of a number of handsome Norway Spruces (Abies excelsa); and did not pay any attention, so far as I observed, to the seeds, although the trees were laden with cones in abundance. I watched them for some time. Crows, Blue Jays, and Chickadees were the only other birds seen, with the exception of a large Buteo or Archibuteo. Many ice-nets were set through the ice, but a fisherman told me that their catch was very small, as the fishes were not running. Ice-boats are plying; and teams are crossing.

February 1st. H. F., N.Y.
 My wife shot a Red male Pine Grosbeak!
 2641. Pinicola enucleator. 9.25 13.80 ♂ ad.
 Wing, 4.50; tail, 3.80; culmen, .59; gape, .61; tarsus, .92; middle toe and claw, .94; claw, .31.
 Stomach contained seeds of the Silver Maple (Acer). Bill, blackish; horn-color on basal one-half of lower mandible. Legs, feet and claws, black. Shot by Ella W. Mearns.

New York City.
Passerella iliaca (Merrem). - Saw the first Fox Sparrow of this year in Central Park. It uttered its clicking metallic note & sharp tweet. Mr. Sanders - son Smith saw the first Robins on Staten Island.

1884.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

A List of Lepidoptera, Collected at Highland Falls, N. Y., during the Summer of 1883.

1. Papilio turnus, Linn.
2. " asterias, Fab.
3. " eriphontes, Cramer.
4. Melitaea phacton, Cramer.
5. Vanessa antiopa, Linn.
6. Limenitis ursula, Fab.
7. Pesia buffaloensis.
8. Sphinx cinerea, Harris.
9. Attacus luna, Doury.
10. " polyphemus, Fab.
11. Arctia virgo, Linn.
12. Saturina io, Linn.

February 21st.

Highland Falls, N. Y.

My wife shot two Pine Grosbeaks.

2642. Pinicola enucleator

♂ juv.

2643. Pinicola enucleator

♀ ad.

March 1st.

Highland Falls, N. Y.

Pinicola enucleator. A flock has been seen about our place nearly every day since Jan. 20th, when I shot the first specimens. Ella has shot four others, but they still are tame, and come about the house, eating seeds of maple upon the ground, & buds of Abies excelsa.

1884.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

March 11th.

Highland Falls, N. Y.

The Pine Grosbeaks (Pinicola enucleator) still remain about the place.

We left Highland Falls in the 1.07 P. M. train on the West Shore Railroad en route to Whipple Barracks, Prescott, Arizona Territory. The snow was deep on the ground in the Highlands; and we drove to the depot in a sleigh. We found snow as far as Philadelphia, where we retired and awoke in time for breakfast at Cumberland, Md., from which place to Grafton, W. Va., the mountain scenery was wild and beautiful. The woods were clothed in many places with a heavy growth of the Rhododendron or Rose-Bay (Rhododendron maximum) and Mountain Laurel (Kalmia latifolia). Savage Creek, a tumultuous stream, flows in a deep ravine beside the B. and O. Railroad; its banks clothed with fine hemlocks (Abies canadensis) and Rhododendron shrubs. We saw a handsome Club-moss (Lycopodium) upon the bank, probably L. clavatum, and a laurel-like plant of the Heath Family, which we failed to recognize.

After leaving Scranton we saw only a few Herring Gulls, and a White-headed Eagle, from the train, going down the Hudson for the last time, perhaps for years. We took tea with my aunt, Mrs. M. J. Potter, in Jersey City, and left for the West on the Baltimore and Ohio Railroad at 7.12 P. M.

March 12th. Cumberland, W. Va. to Cincinnati, O.

A large flock of Blackbirds (species not determined) was seen in W. Virginia, and another in Ohio. One Fox Sparrow (Passerella iliaca) seen in Eastern Ohio. Crows (Corvus jugosus) were occasional, and so were Doves (Zenaidura).

1884.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

carolinensis). A Kingfisher (Ceryle alcyon) was observed in the western part of West Virginia; the first I have seen this year.

Besides an occasional Hawk, the above were all the birds noted along the road although I watched carefully for them.

The scenery along the Kanawha Cheat River is very beautiful. We long to see the beautiful Rhododendrons in flower. They were abundant in many localities throughout the mountains of all of West Virginia, but none were seen in Ohio.

The Baltimore and Ohio Railroad from to is very rough. Between the two places there are 23 tunnels and 52 bridges.

We crossed the Ohio at and saw evidences of the great flood of the past month. The Railroad embankment had been washed away on the Ohio side, and, for a week or more the passengers on the B. and O. had to transfer on foot across the breach. At Cincinnati a large section of usually dry land was inundated. We slept from Cincinnati, Ohio until we were nearly to Clay City, Illinois, and so missed the Indiana scenery, which would have been new to me.

March 13th. Clay City, Ill. to St. Louis, Mo. We dressed and looked out of the cars at Clay City. We were passing through a heavy forest, with the ground beneath the trees covered with water as far as the eye could reach. On the top of one of many huge dead trees, standing beside the Railroad, sat a Pilated Woodpecker (Heplatomus pileatus). As the train passed it flew into

1884.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

the forest, its red crest gleaming in the sunlight. Thence to St. Louis we saw & heard many Meadow Larks. Small flocks of Crow Blackbirds were seen frequently, as were Blue Jays, and Woodpeckers of several species (I only recognized specifically one Red-headed Woodpecker (Melanerpes erythrocephalus). A few Larks (Zenaidura carolinensis) were seen, and a large species of Snipe or Plover with snowy white breast. Flocks of Ducks were seen at a distance. Two large Buteos (probably Buteo borealis) were perched in trees quite close to the track, whence they looked unconcernedly at us, without taking to flight. I saw some Robins (Turdus migratorius). English Sparrows (Passer domesticus) were abundant all along the line, and other small passerine birds were seen but not identified all along the line. Crows (Corvus frugivorus) were occasionally seen, but not in large numbers. No Quails (Colinus virginiana) have so far been noted.

We reached St. Louis before noon, and spent the afternoon and evening pleasantly at the "Southern" hotel. ^{Eight Coots (Fulica americana) were seen in the Mississippi under the St. Louis bridge.}

March 14th. St. Louis, Mo. to Halstead, Kansas. We took the 8.25 P.M. train from St. Louis, and reached Springfield, Mo. in time to breakfast there at 6.25 A.M. During the day we crossed the western one-third of Missouri and eastern one-half of Kansas, reaching Wichita, K. before dark, and retiring for the night at Halstead.

In Western Missouri we saw few birds which I was able to identify as they flew. There were a good many Sparrows and little fellows that looked like Titlarks.

1884.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

I thought I saw Lark Finches (*Chondestes grammacus* (Say) Bp., but was not positive of the identification. I saw a few Red-tailed Hawks (*Buteo borealis*), one or two Sparrow Hawks (*Falco sparverius*), plenty of Meadow Larks (*Sturnella* — ?) and a few Marsh Hawks (*Circus hudsonius*). Flocks of Ducks of several species were seen upon some reedy marshes on the stream near Cronogo, Mo., but the only species recognized was the Golden-eye (*Colaptes auratus*). Crows (*C. frugivorus*), which were scarce through Ohio and W. Illinois, were exceedingly abundant in large flocks in the corn-fields. Robins (*Turdus migratorius*) were also plentiful in enormous flocks observed in places; and the species was seen as far west as the Eastern border of Colorado, where I noted them on the 15th. Several were seen at Coolidge. Blue Jays (*Cyanocitta cristata*) were abundant in Mo.; and large flocks of Bluebirds (*Sialia sialis*) were seen, as well as large flocks of Red-winged Blackbirds (*Agelaius phoeniceus*) and Crow Blackbirds (*Zonotrichia querula*). Plovers of some species were frequently seen in ploughed fields and grassy prairie. A large, long-winged bird, which I guessed was Bartram's Upland Plover (*Bartramia longicauda*), a bird which I have never otherwise seen was also occasional. The only Woodpeckers seen were Yellow-shafted Flickers (*Colaptes auratus*), and they were scarce. No signs of mammals save squirrel-nests.

In Kansas as far west as Wachtita the following species were observed:—

1884.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

Meadow Larks (*Sturnella* — ?) were plentiful and singing all the way. A few Robins were seen. One flock of Quail (*Ortyx virginiana*). Near Augusta, Kansas, three Prairie Hens (*Cupidonia cupido*) were seen — the first by me! The one first observed flew up beside the track and around behind the train, soaring with wings set as it disappeared. The other two were squatting close to the ground in a corn-field. A few Blue Jays in the eastern part of the State. I saw a number of Pigeon Hawks (*Falco columbarius*), always in pairs, one Rough-legged Hawk (*Archibuteo lagopus sancti-johannis*) near Fredonia, K., and a few Marsh Hawks (*Circus hudsonius*) both blue and brown; also one Red-tail (*Buteo borealis*). A few Shrikes (*Lanius ludovicianus excubitorides* ?), always solitary, were seen haunting the hedges which are much used in lieu of fencing through Kansas. Larks (*Anthus* or *Eremophila* ?) were plentiful. Although we skirted streams which contained a few ducks, no Kingfishers were seen since the one observed east of the Mississippi.

A solitary Antelope (*Antilocapra americana*, Ord) was seen near Hall River; the first I ever saw, and the only ^{wild} mammal seen in Kansas.

Baltimore Orioles' nests were numerous in wooded places. One Red-wing and a few Crow Blackbirds in E. Kansas, and a few Plovers as far as Wachtita, near which place a number of singing birds were heard and seen, but not identified.

Through Western Missouri some heavy timber is found, which decreases in size and extent as we approach the Kansas line, except

1884.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

along the streams, which were bordered by large sycamores (*Platanus occidentalis*) and other trees. Gradually the Scrub Oaks disappear, and prairies of large extent become more numerous; and when one crosses the Kansas line, trees almost disappear, and when present indicate the presence of a stream whose banks they fringe.

Through the Eastern half of Kansas long stretches of prairie extend as far as the eye can reach and form the horizon boundary on every side. Trees only grow beside streams. The grass is plentiful and apparently good. Towns and ranches abound. The prairie grass was burned in many places over large areas by the inhabitants on account of the benefit to their land from the ashes. Emigrant trains and single wagons were frequent.

We retired after leaving Halstead, having watched the burning grass in the prairie fires which surrounded us, for a long time.

March 15th. Halstead, Col. to Las Vegas, N.M.
Last night we travelled from Halstead to Aubrey, Kansas, and all slept soundly. We awoke upon a level prairie, covered thickly with cattle; and many carcasses of dead beasts from the "hoof and mouth disease" were seen along the road.

At Coolidge, near the Colorado border we broke our fast to the extent of a sandwich and cup of coffee. Between Aubrey and Coolidge we saw the first of the beautiful tufts of sword-grass or Spanish bayonet (*Yucca glauca*) each broad, tough blade armed with a spear

1884

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

or needle at its point. The roots are strong and fibrous. Small plants remind one of the pine-apple. This beautiful bright green plant soon became a common feature, and we found it at the highest altitude of the spur of the Rocky Mountains which we crossed (17000 feet) when coming south into New Mexico, and in New Mexico as far as Lipton, where darkness overtook us.

I first noted the Cotton Wood trees (*Populus fremonti*) near Coolidge, Kansas, and they were the only trees seen afterwards until we came to the low, gnarled cedars of the foot-hills of the Rockies near Hatch, Col., excepting low willows (*Salix*) along streams. The Cotton Woods were quite tall and symmetrical, and grew in broad belts along the Arkansas River and other streams.

The first Cacti were seen near Las Animas, Col. They were covered with long, sharp spines. In general color they varied from reddish to woolly gray. I got a specimen of joints, of course not yet in flower. This cactus (*Opuntia rafinesquiana*) covered the ground in many places as far as the mountains. Although there is much desert land through Colorado, where no vegetation can be seen at this season, a good part of the country is clothed with thin grass, on which many cattle graze. Large areas of sandy sage plains were crossed. The colors of the grass land are bright and beautifully colored with red, brown and yellow in varying shades.

Between Granada and La Junta the first Prairie Dogs (*Cynomys ludovicianus*) were seen. There were several large villages, and the inhabitants did not seem in the least

1884.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

disturbed by the passing train. Some sat up erect and looked towards it, others sat upon the ground eating, and others were sprawled flat across the top of their mounds. As the train passed by they would frequently jump playfully up in the air. In Colorado, the last Prairie Dog village was noted near Earle Junction in the southern part of the State.

A large Jack Rabbit (*Lepus texianus*) was seen near Trinidad, Colorado.

Small burrows were abundant all through Colorado, along the road, and were probably inhabited by Gophers.

Groves of horses were first seen at Limpas. Birds. - At Coolidge, Kansas, a few Robins. Little Larks of some sort were abundant through Colorado. A few flocks of Ducks were seen along the Arkansas River, and a few Crows near La Junta, here Rough-legged and another large Hawk which I think was Swainson's Buzzard (*Buteo swainsoni*) were numerous.

One Sparrow Hawk was seen near Granada. One Bluebird (probably *Sialia arctica*) was seen high up in the mountains of Colorado.

I saw the first Magpie (*Pica rustica hudsonica*) near Earle Junction at a Mexican ranch where beaver had been slaughtered. There was not a tree within sight of the place. A few others were seen between there and Trinidad, always near Mexican dwellings. Their large round nests were numerous about Trinidad and thence over the mountains into New Mexico, where there were trees.

In the mountains the first trees seen were the cedars of the foot-hills. As we ascended we found pines in abundance and

1884.

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

tall spruce trees. On the high mountain slopes were stunted oaks and pinyons.

The Mexican inhabitants lived in dwellings varying from respectable adobe mansions to mere burrows in the ground.

We passed Albuquerque, New Mexico during the night, and were in Western New Mexico when we awoke next morning.

March 16th.

New Mexico to Ash Fork, A.T.

In western New Mexico we noted the beautiful shapes and colors of the sandstone walls of rock, and the volcanic character of the region through which we passed. There were vast lava beds which seemed scarcely to have cooled; mostly blackish in color. We saw the first Ravens (*Corvus corax caninus*) about the red cliffs which walled in the valleys and large flocks of small Ducks, probably teal, where there was water. We crossed the sandy plains of Arizona and the beautiful Bill Williams Mountains which were deeply covered with snow and supported a growth of huge and magnificent pine-trees. In the foot hills there were cedars as usual, then low pines, and towards the higher part were the magnificent pines noted above.

Birds were scarce, however, and only a few small species, some Woodpeckers whose species I could not recognize and a large Hawk which may have been the Duck Hawk (*Falco peregrinus*).

We reached Ash Fork on time at ten o'clock P.M., and put up for the night at the "Palace Hotel", a rough shanty, for the night.

1884.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

March 17th, 1884. Ash Fork, Arizona.

My previously dispatched telegram had not reached Ash Fork when we arrived there, as the wires were down. We, therefore telegraphed to Whipple Barracks for an ambulance immediately upon our arrival, but this compelled us to remain at Ash Fork until the forenoon of the 18th.

Ash Fork is a typical border town, made up of rough board shanties in most of which liquor is sold. It is a rough place, made up of the usual population in such places, viz: freighters, miners, and the usual complement of desperadoes, gamblers and prostitutes. The "Palace Hotel" to which I was compelled to take my wife and little Lillian was indeed a hard place to take a lady unaccustomed to rough living. A few weeks after we left, the guests of this establishment engaged in a quarrel with the proprietors and the whole building was torn down, and the proprietors escorted to the station and put on the first train and shipped. At my second visit a fairly comfortable "hotel" was being kept.

While the A. & P. Railroad was being built "shooting affairs" were of nightly occurrence. A row took place on the morning after we came. A good deal of shooting was done, and that too within a few yards of my wife and baby who were in a car getting some milk from a lady who owned the only cow in Ash Fork. One man was shot through the arm producing a compound comminuted fracture of the humerus. I was called in professionally, but everyone was drunk and carried a cocked pistol, and I was glad to leave the patient as soon as I had taken precautions against hemorrhage and adjusted a bandage.

Later in the day I went out to see the country. In town, about the shanties, were a few light colored

1884.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

, very shy Snowbirds (*Junco*). The first bird I shot in Arizona was the Western Grass Finch (*Poocetes gramineus confinis*). The species was abundant about the town.

Finding a congenial companion, we started off together in a wagon behind a team of bronchos and drove about six miles to Johnston's Cañon. Ash Fork is situated beside the foot-hills surrounding the Bill Williams mountains, at quite an altitude. The region is covered with a scattered growth of red cedars. Antelope, Deer, Prairie Wolves, and Foxes are numerous in the vicinity. There are a few "Mexican Larks" (*Felis concolor*) and plenty of Wild Cats in Johnston's Cañon. Among the cedars I saw a single Savannah Sparrow (*Passerculus savanna*) which I shot on top of a cedar-tree. Shore Larks (*Eremophila alpestris chrysolaema*) were numerous in large flocks. Bluebirds were quite numerous (*Sialia mexicana*), as were the Western Robins (*Merula migratoria propinqua*) one of which latter my friend shot in the Cañon where a good-sized stream then flowed, which he informed me dries up entirely later in the spring.

There were some Ducks in it, some of which we shot. They were *Fulix marila*; but if the characters exhibited by these specimens are constant they certainly should constitute a well-marked variety, which I will name here *Fulix marila arizonae*. I scarcely recognized them as Pease Ducks.

Near the Cañon I saw a number of Ravens one of which I made a good wing shot at, and "collected". *Spyrapicus varius nuchalis* was also quite numerous among the cedars, and I also collected a fine male specimen of that.

"Rock Squirrels" (*Spermophilus grammurus*) of large size were seen in numbers about the rocky walls of the Cañon, as were some little ground

1884

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	Squirrels (<i>Tamias asiaticus dorsalis</i>). Numbers of the Small "Cotton-tail" Rabbits (<i>Lepus sylvaticus arizonae</i>) were seen and shot, as were also seen the large Jack Rabbits (<i>Lepus callotis texianus</i>) by my companion; but I did not see any Jacks myself until I reached Prescott.					
	No flowers are in bloom at this early season; but numbers of interesting plants were seen, among them Aloss, Spanish Bayonets, and cacti in abundance.					
2644.	<i>Corvus corax</i>		<i>carnivorus</i> (Baird.) Ridgw.			♂ ad.
2645.	<i>Spyrapicus</i>		<i>varius</i> <i>michalis</i> , Baird.			♂ ad.

March 18th. Ash Fork to Bangharts, Arizona.

The ambulance got in late last night. We started about 9 o'clock A.M. for Prescott, and reached Bangharts rancho, where we stopped for the night and met Lieut. Anderson, who was on his way to Ft. Grant. Before breakfast I took my gun and took a stroll over a stony, cedar-covered hill near Ash Fork, and collected specimens of *Lophophanes inornatus* (The Plain Titmouse), and *Sturnella neglecta* (Western Meadow Lark). The only other birds seen were Sparrow Hawks (*Accipiter velox*). The Coyotes (*Canis latrans*) were yelling loudly and the dogs in town were all barking back at them.

Old Pease, a veteran driver & mulater drove us, with a team of six mules, and we rattled over the malapai roads, through mud of unknown depth, and over rocks until we were all three nearly pounded into jelly. We crossed grassy plains and sandy wastes of great extent. The country near Ash Fork, however, was forested with the red cedar, with open spaces between the wooded ones, and deep canyons were occasionally crossed. The cacti were the most interesting feature of the scenery to us. A few large "Century plants" (*Cloe*) were seen with dead flower stalks 10 to 15 feet high.

1884

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	Birds were not very numerous. There were two species of Bluebirds in the cedars (<i>Sialia arctica</i> and <i>S. mexicana</i> , of which the latter were much the most numerous. The Western Grass Finch (<i>Poocetes gramineus confinis</i>) was everywhere common. A few Snowbirds were noted (Juneo) and a small Lark was occasionally seen which looked like a Pipit (<i>Anthus</i>). Shore Larks were in great plenty, and associated in large flocks. This species (<i>Eremophila alpestris chrysolaema</i>) outnumbered all the rest. A few Ravens (<i>Corvus corax</i> <i>carnivorus</i>) were passed, as they sat on cedars crouching at us from no great distance.					
	We stopped for dinner at the eating station on Creek, a good-sized stream at this season, where I saw the first Jays. Woodhouse's Jay (<i>Aphelocoma woodhousei</i>) utters a catbird-like note and a series of passionate shrieks. I also saw a single Song-crested Jay (<i>Cyanocitta stelleri macrolopha</i>) at this place. We then crossed a wide belt of dense cedar timber, in which I shot my first specimen, a female, of Gambel's Quail (<i>Lophortyx gambeli</i>); and I secured several more, all females, before we got to Hell's Cañon, where we found a number of flocks of Gambel's Quails and some Song-crested Jays & a Fox. The only other birds noted were <i>Buteos</i> . <i>Sturnella neglecta</i> was seen at intervals along the route.					
2646.	<i>Sturnella neglecta</i>		Andubon.			♂ ad.
	Shot in a cedar-tree at Ash Fork.					
2647.	<i>Lophophanes inornatus</i>		(Gamb.) Cass.			♂ ad.
	Shot among the cedars at Ash Fork.					
2648.	<i>Lophophanes inornatus</i>		(Gamb.) Cass.			♀ ad.
	Shot in a cedar-tree at Ash Fork.					
2649.	<i>Lophortyx gambeli</i>		Nuttall.			♀ ad.
	Hell's Cañon, Yavapai County, Arizona.					
2650.	<i>Lophortyx gambeli</i>		Nuttall			♀ ad.
	Hell's Cañon, Yavapai County, Arizona.					

1884.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
2651.	<i>Lophortyx gambeli</i> , Nuttall		Keell's Canon, Yavapai County, Arizona.			♀ ad.

March 19th From Banghart's to Prescott, Arizona.
We left Banghart's at eight o'clock A.M. and reached Whipple Barracks at 4 o'clock P.M., where I reported to Medical Director B. J. D. Irwin and his Adjutant General, and was assigned to Fort Verde, Arizona, as Post Surgeon, with A. A. Surg. A. P. Frick. Then Post Surgeon for my assistant. We then repaired to the Williams House at Prescott, one mile distant from Whipple Barracks, the Headquarters of the Army in the Department of Arizona.

Before starting, in the morning, I took a long walk along the little stream near Banghart's house. About the corrals, I found enormous flocks of Brewers and one variety of Red-winged Blackbird. I killed five females of the latter (*Agelaius*) at a single shot, and also procured a male Brewer's Blackbird. Along the stream, I found a small species of Sandpiper (*Fringilla*?) and a number of Wilson's Snipe (*Phalaropus lobatus* *Gallinago media wilsoni*). Killdeer were abundant (*Oxyechus vociferus*). Several Sandhill Cranes (*Grus canadensis*) were seen on the plain from Banghart's house early in the morning, before we awoke; but they were gone when I went out to see them. Five Cranes had been seen for several successive days at this place. Ducks were very numerous on this little stream. The only one I succeeded in shooting was the Red-breasted Teal. A pair of Black Pewees (*Sayornis nigricans*) were seen on the stream; the first I ever saw. Western Robins (*Merula migratoria propinqua*), and Mountain Song Sparrows (*Melospiza fasciata fallax*) were abundant here; and so were Western Meadowlarks (*Sturnella neglecta*) and Woodhouse's Jay (*Uphelocoma woodhousei*). The first Red-shafted Flickers

1884.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

(*Colaptes auratus mexicanus*) were seen here. The Shore Larks (*Ammodramus alpestris chrysolaema*) were infinitely abundant all the way from Banghart's to Prescott. A few Western Grass Finches (*Poocetes gramineus*) and Snowbirds (*Junco*) and numbers of Sparrow Hawks were the only other birds, except occasional Meadow Larks seen along the level road from Banghart's until we reached the brush hills near Prescott, where I saw the first Spurred Towhee (*Pipilo maculatus megalonyx*). When we reached the cedars at the Point of Rocks we saw quantities of Bluebirds of both species and of Snowbirds. About Whipple we saw a number of Ravens. Little Lillian petted a "Cottontail" (*Lepus syl. arizonae*) which I shot near Banghart's all the way in and was delighted with the pretty little beast which she calls a "boga".

2652. *Uphelocoma woodhousei* (Baird) Ridgw. — — — — — ♀ ad.
Banghart's Ranch, Yavapai County, Arizona.
2653. *Scolecophagus cyanocephalus* (Wagl.) Lat. — — — — — ♂ ad.
Banghart's Ranch, Yavapai County, Arizona.
2654. *Agelaius phoeniceus phoeniceus*. — — — — — ♀ ad.
Banghart's Ranch, Yavapai County, Arizona.
2655. *Querquedula cyanoptera* (Vieill.) Cass. — — — — — ♂ ad.
Banghart's Ranch, Yavapai County, Arizona.

March 20th Prescott, Arizona.

The day was spent in looking about town, visiting the post at Whipple, skinning birds, and a short ramble in the woods, amongst the tall pines, which we used to read about with envy in the writings of Dr. Elliott Coues. Now we were free to tread where his illustrious feet had preceded us, and much we enjoyed doing so. I took Mrs. O'Hearn and Lillian along, and then returned to the hotel for our gun.

1884.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	We saw and heard the cognacious Long-crested Jay and the Californian Woodpecker (<i>Melanerpes formicivorus</i>) and wandered along Granite Creek, among the rocks, where I saw the first Jack Rabbit (<i>Lepus cal. texianus</i>), a truly wonderful animal.					
	About Prescott and Whipple we heard and saw flocks of those delightful little birds - the House Finches (<i>Carpodacus frontalis</i>). Their song is delicious, their chattering like House Sparrows, and their call-note like the Eastern Goldfinches (<i>Astragalinus tristis</i>).					
	Pygmy Nuthatches were creeping upon the trunks of the huge pines, and were noisy and restless in their habits, as well as gregarious, although single birds were seen.					
	Mexican Bluebirds (<i>Sialia mexicana</i>) were quite numerous, and uttered a note somewhat like the House Finches, and a song which is like the Eastern Bluebirds (<i>S.ialis</i>), but is lacking in the melodious sweetness of our Bluebird. They sat upon the lower dead branches of the pines and flew out after insects.					
2656.	<i>Sialia mexicana</i> , Swainson.			—	—	♂ ad.
2657.	<i>Sitta pygmaea</i> , Vig.			—	—	♂ ad.
2658.	<i>Carpodacus frontalis</i> (Say) Gray.			—	—	♂ ad.

March 21st.

Prescott, Arizona.

I spent the morning in the woods about Prescott, skinned birds, and received calls.

The House Finches were found in large flocks, feeding upon the ground, perhaps on the seeds of the pine cones.

Woodhouse's Jay was quite numerous, and found high up in the tall pines, as well as in the oak-trees, bushes and upon the ground. Wherever found its passionate screams "rent the air" and its whole demeanor is savage and irascible.

The Californian Woodpecker was found in some numbers. When not disturbed or alarmed they

1884.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	are as fond of noise and play as are the Eastern Red-heads (<i>M. erythrocephalus</i>). At times they are quite bold, and will permit you to approach to within a few feet of the spot where it is working. Again it seeks to hide by getting up in the thickest part of the tree and sticking close to the bark and there remaining motionless for a long time.					
	One that I wounded concealed itself so effectually that, although I knew about where it was, I was unable to find it.					
	The Pygmy Owl (<i>Glaucidium gnoma</i>), I owe to the Pygmy Nuthatches, who were scolding and making such a fuss that I went to see what cause our circumstane had brought together such a congregation of these noisy little fellows, and found them attacking & pestering the smallest Owl I ever beheld. The owl flew from one tree to the next with ease, in broad day light.					
	Red-shafted Flickers were numerous, but shy.					
	<i>Regulus calendula</i> . - I shot one of the familiar birds among some willow bushes along Granite Creek. It is the first bird I have seen in Arizona that is exactly like those found in the East.					
	Robins were abundant and singing pleasantly.					
	Snowbirds were excessively common and feed upon the ground in company with <i>Carpodacus frontalis</i> . Seeing up with them into the lower branches of the pines when approached, the House Finches issuing a loud whurring sound like the <i>C. purpureus</i> at home.					
2659.	<i>Apelocoma woodhousei</i> (Baird) Ridgw.			—	—	♀ ad.
2660.	<i>Sialia mexicana</i> , Swainson.			—	—	♀ ad.
2661.	<i>Melanerpes formicivorus</i> (Sw.) Bp.			—	—	♀ ad.
2662.	<i>Melanerpes formicivorus</i> (Sw.) Bp.			—	—	♂ ad.
2663.	<i>Melanerpes formicivorus</i> (Sw.) Bp.			—	—	♂ ad.
2664.	<i>Glaucidium gnoma</i> , Wagl.			—	—	♀ ad.

(over)

1884.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length. in m. m.	Stretch. in m. m.	Sex and Age.
2665.	<i>Zonotrichia gambeli</i> , Nuttall.			—	—	♂ ad.
2666.	<i>Sturnellus sparverius</i> (L.) Vieill.			—	—	♀ ad.
2667.	<i>Cyanocitta stelleri macrolopha</i> (Bd.) Ridgw.			—	—	♂ ad.

March 22nd.

Prescott, Arizona.

Spent the morning in the woods again. In addition to the birds found here before, I found a number of Slender-billed Nuthatches and shot one of a number of Plain Titmice (*Lophophanes inornatus*) that were seen in the low oaks. Western Meadow Larks (*Sturnella neglecta*), Harris's Woodpecker (*Picus villosus harrisi*), and White-throated Wren were seen; also Plumbeous Titmice (*Psittiparus plumbeus*).

I also shot two large fat Rock Squirrels (*Spermophilus grammurus*) which live in the singular huge pyramidal piles of granite boulders around Prescott. I also shot a couple of little Hoard Squirrels (*Tamias asiaticus dorsalis*) in the same place.

2668.	<i>Cyanocitta stelleri macrolopha</i> (Bd.) Ridgw.			—	—	♂ ad.
2669.	<i>Cyanocitta stelleri macrolopha</i> (Bd.) Ridgw.			—	—	♀ ad.
2670.	<i>Aphelocoma woodhousei</i> (Bd.) Ridgw.			—	—	♀ ad.
2671.	<i>Sitta carolinensis aculeata</i> (Cass.) Allen.			—	—	♂ ad.
2672.	<i>Sitta carolinensis aculeata</i> (Cass.) Allen.			—	—	ad.
2673.	<i>Colaptes auratus mexicanus</i> (Sw.) Ridgw.			—	—	♂ ad.
2674.	<i>Picus villosus harrisi</i> (Aud.) Allen.			—	—	♂ ad.
2675.	<i>Picus villosus harrisi</i> (Aud.) Allen.			—	—	♀ ad.
2676.	<i>Carpodacus frontalis</i> (Say) Gray.	161	253			♂ ad.

79; 67; culmen, 10; gape, 11; tarsus, 17; middle toe and claw, 18; claw, 5. Iris, hazel.

2677.	<i>Carpodacus frontalis</i> (Say) Gray.	160	255			♂ ad.
-------	---	-----	-----	--	--	-------

Wing, 80; tail, 70; culmen, 11; gape, 12; tarsus, 17; middle toe and claw, 18; claw, 5. Iris, hazel.

2678.	<i>Catherpes mexicanus conspersus</i> , Ridgw.					♀ ad.
-------	--	--	--	--	--	-------

This specimen was found in a large pile of rocks. It uttered a very singular note which I do not know how to describe, and was so tame that I shot several Squirrels from the rocks about the spot.

1884.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

before attempting to secure it. I was then able to approach to within a few feet of it. This is the only individual seen.

March 23rd.

Prescott, Arizona.

The day was spent in skinning birds and receiving calls from the officers at the Post. I also walked over to Whipple and called on Dr. Hopkins, Dr. McCree, and the Medical Director. About the quarters were flocks of House Finches which were singing sweetly in spite of the execrable weather, for it snowed all last night and to-day the snow is melting and it is sleeting and snowing at intervals. Woodhouse's Jays were numerous in the oaks & cedars around the post; and a few Ravens are making themselves at home about the corral. Sparrow Hawks were sitting on the pine-trees in the garden.

March 24th.

Prescott to Hudson's, Arizona.

We left Whipple at nine o'clock, our trunks having just arrived from Ash Fork the night before. The Woodhouse's Jays were abundant all the way to Hudson's rancho where ~~where~~ we stopped for the night. The Spurred Towhee became common soon after we left the evergreens of the foothills. (At Whipple the principle forests are huge pines (*Pinus*) and there are two species of cedar, the red cedar (*Juniperus*) and the rough-barked Juniper (*Juniperus pachyphloea*). The oaks are of two species—a medium sized tree (*Quercus emoryi*) and the scrub oak (*Quercus*) which is a shrub. There are a few cottonwoods, willows and other deciduous trees along Granite Creek.) Its note is very similar to that of the Catbird. The Shore Larks, a little Lark supposed to be *Anthus* and the Snowbirds were plentiful, and we saw large flocks of Gambel's Quail, *Regulus calendula*.

1884.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

was occasionally observed all along the road to Hudson's ranche, which is on Ash Creek about 19 miles from Fort Verde, and where the species was very common, but not heard singing. Both species of Bluebird (*Sialia arctica*) and *S. mexicana* were noted at various points along the road.

Pipilo fuscus mesoleucus. — A specimen of this pretty Towhee was shot among some broken rocks; but it was afterwards lost out of the ambulance. A few Meadow Larks and Grass Finches were occasionally seen. Some *Margaritas* were the first flowers seen. We reached Hudson's late in the afternoon, but in time to take a ramble about the place.

Woodhouse's Jays were vociferating on every hand, and the call of Gambel's Quail was heard in all directions. The Spurred Towhee was very common and I saw several Red-vented Thrashers (*Harporthynchus crissalis*) for the first time.

2679. *Lophortyx gambeli*, Nuttall. 277 377 ♂ ad.

Hudson's Rancho, on Ash Creek, 19 miles from Ft. Verde, Arizona. Wing, 115; tail, 107; culmen, 11; gape, 13; tarsus, 32; middle toe and claw, 36; claw, 9. Irides, dark hazel. Bill, black. Tarsi and toes, olivaceous bluish-gray; claws, black.

March 25th. Hudson's Rancho to Ft. Verde, Arizona. Besides the birds noted yesterday we saw one or two ladder-backed Woodpeckers (*Picus scalaris*) and a White-rumped Shrike (*Lanius ludovicianus excubitorides*). The Curved-billed, red-vented Thrashers (*Harporthynchus crissalis*) were occasionally noted, and the large Rock Squirrels (*Spermophilus grammurus*) were quite numerous in the vicinity of Ash Creek and again in Copper Cañon where we got our first view of Fort Verde and the Verde River, which, at the time is a rapid, muddy stream of very respectable size. In the level, open spaces along the route we saw a number of pretty flowers, among which were

1884.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

a purple *ternstroemia* plant, quantities of small species of *Geranium*, and a very singular purple-flowering plant with palmate or pinnatifid leaves. In Copper Cañon were brilliant red Indian Pink and many other interesting plants. The yellow *Aquilegia*, though not yet flowering was seen putting forth leaves. The pussy willows were in flower, and the leaves on the cottonwoods were just beginning to unfold. A large *Umbelifera* with purple-green palmate or pinnatifid leaves and large umbels of small yellow flowers was seen in abundance. As we came through the Cañon numbers of Woodhouse's Jays flew screaming up the hillside which are wooded with red cedars, and a small species of pine (*Pinus edulis*).

As we descended into the Verde Valley the temperature rose rapidly. The only birds noted were Shore Larks ^{and *Amphispiza bilineata*, seen for the first time.} As we rounded the corner of an adobe corral large flocks of Brewer's Blackbirds (*Scelopophagus cyanocephalus*) wheeled overhead and a large, whitish Falcon (*Falco mexicanus polyagris*) flew low over the ambulance. Captain Russell, who commands Ft. Verde, was swinging in his hammock, and escorted us to Dr. Frick's house, where we found a pleasant room and fire ready for us; and presently Lieutenants Vogdes and Carrington called, and kindly asked us to join their mess which we were very glad to do temporarily. Lieut. and Mrs. Morgan then called and asked us to come to lunch. We found the table decorated with wild flowers and fragrant peach blossoms. We dined at the mess, while our chinaman, whom we brought from Prescott with us, Dick Ah Moon, accepted tendered courtesies from his countrymen. After leaving Copper Cañon, no vestige of spring foliage was seen. The

1884

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

only deciduous trees visible were cottonwoods which margin the winding course of the Verde River as far as the eye can reach to the north and south. The Valley is, otherwise, destitute of trees, for the red cedars and small pines are only found in scattered patches in the ravines of the lower foot-hills. At the post, a double row of small cottonwoods in front of Officers' Row and a single scattered line of the same trees around the parade-ground were the only trees. The altitude of Fort Verde is 3,500 feet. It is much lower than Whipple Barracks, which is feet above sea level; and, consequently it has a much warmer climate. It is said to take the third rank in point of heat of all the posts in the U. S., Mojave, on the Colorado River ranking first. Verde is built on a little plain in the open valley, about 100 feet above the River, which, upon the Post side has low sandy banks; but on the farther side the bank rises through several irregular terraces crowned by grassy mesas to high steep and rocky walls of limestone rock, which were occupied by Aztec inhabitants, whose cave dwellings may be seen from the Post. In summer the Verde is a good-sized brook, fordable anywhere except where there are quicksands. At this time it was a rushing torrent, tearing the river bed, sweeping away great masses of clay from its perpendicular banks, which the water undermined causing to fall tremendous chunks into the edge of the stream. Cottonwoods ~~woods~~ and willows are being torn up and carried down the River, and the whole river bed shifting from day to day, so that a good crossing one day cannot be relied upon another.

The river bottom is composed of clay and sand, bare in large areas, in others covered with a more or less scattered growth of mesquite bushes and

1884

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

sage brush and other bushes. In the neighborhood of the Post the pretty little Geraniums are the only flowers.

March 26th. Fort Verde, Arizona.

This afternoon I drove up to the Middle Verde, six miles, to see a patient, in company with Dr. Frick. In a Cañon we saw a large flock of Gambel's Quails, which are very plentiful in the Verde Valley.

During the night it snowed hard and the hills and mountains around us are deeply covered with snow, and to-day it is hailing and sleeting in the valley. I was peeping out of the ambulance through a slit in the curtain when a brilliant red bird flew from near us and alighted in a bush beside the Verde. It was the first Vermilion Flycatcher (Tijrocephalus rubinedes mexicanus) that I ever saw. Red-shafted Flickers (Colaptes auratus mexicanus) and Robins (Merula migratoria propinqua) were very common at the ford at Parson Bristow's, where we experienced considerable difficulty in getting the ambulance across the river on account of the high water, which came nearly to the mules' backs. Many small birds were seen, but owing to the stormy weather I could not see distinctly enough to identify them, excepting Picus scalaris + Eremophila.

March 27th. Fort Verde, Arizona.

This afternoon I went shooting with Lieutenant Vogdes. We rode on mules. Mine was a little white fellow with extraordinary development of ears. ^(Muley Crook) We rode down the Valley, and along the edge of the foot-hills. I bagged eight Gambel's Quails and saw several new birds. Among those seen were Oreoscoptes montanus, Regulus calendula, Sialia mexicana, Carpodacus frontalis, Proccetes grammacus confinis, Amphispiza bilineata, Pipilo aberti, Sturnella neglecta, Corvus corax caninus, Sayornis sayi, Picus scalaris, Linumolus sparverius, Buteo calurus, and ^(small)

1884

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

Aquila chrysaetus canadensis, which latter species was seen within a few rods, flying low above my head, but, before I could dismount and slip ^{change off} a large shot in my gun it was out of range. I watched the bird until I saw it joined by another one.

Neurofalcon mexicanus polyagris. - Several of these birds in very pale plumage were seen about the Post, where they have killed a number of chickens in the yards, and pigeons about the corral.

Oxyechus vociferus. - Heard late last night, and occasionally seen flying uttering its loud plaintive cry. Numbers of small birds were seen which were not identified. Rabbits of both species (*L. I. arizonae* and *L. texianus*) were seen in abundance.

2680. *Lanius scalaris*, Wagl. 185 333 ♀ ad.
Wing, 99; tail, 68; culmen, 21; gape, 23; tarsus, 17; middle toe and claw, 21; claw, 8.

2681. *Oreoscoptes montanus* (Townsend) Baird. 222 323 ♂ ad.
Wing, 98; tail, 95; nostril, 11; gape, 25; tarsus, 30; middle toe and claw, 26; claw, 7. Irides, yellow.

March 28th. Fort Verde, Arizona.

Sceloporus cyanocephalus. - Every morning since we have been at Verde, we have awoken to hear the note of this bird, and on looking out have seen the parade-ground black with them. They are very tame, coming upon the gravelled road in front of our quarters, and occasionally even upon our piazza. They flock about the corals, and feed upon the ground amongst the horse, avoiding their hoofs with wonderful dexterity. They spend the day upon the parade ground and adjacent localities, and the ~~near~~ houses & buildings as well as the adobe walls of the corals are resorted to as favorite roosting places by day, but in

1884.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
--------------------	-------	-------	-----------	---------	----------	--------------

the evening they repair to the distant cottonwoods along the River. When retiring for the night they leave the Post in large bands, but never all at one flight, but usually in a dozen or more flocks, keeping close to the ground. In these flights the females fly in separate flocks by themselves; and the sexes often separate even when feeding.

6846 Another capture of the Widgeon (*Anas penelope*) on the Atlantic Coast. - On January 11, 1890, I saw a very fine adult male hanging in a marketman's stall in New York City, near "Washington Market". Upon inquiry, I learned that it had been shipped from Baltimore to New York, and doubtless was shot on Chesapeake Bay. It was hanging in a bunch of Dusky Ducks, with which it was said to have been received. It has been mounted, and placed in the collection of the American Museum of Natural History, New York. Its fresh measurements were as follows: length, 495 mm.; alar expanse, 845 mm.; wing, 260 mm.; tail, 116 mm.; culmen, 34 mm.; tarsus, 40 mm. - Edgar A. Mearns, M. D., Fort Snelling, Minn.

Index to the Birds in this Volume (Specimens).

(Arranged according to the A. C. U. Check-List, amended by the 1st and 2nd Supplements.)

Colymbus holboellii. - Nos. 1781, 1782.

Colymbus auritus. - 553, 573, 1859, 1860, 2104.

Podilymbus podiceps. - 782, 1774, 1977, 1994, ~~2104~~, 2207, 2212.

Tringator imber. - 766.

Larus glaucus. - 2007.

Larus argentatus smithsonianus. - 266, 268, 352, XXI p. 103.

Larus delawarensis. - 2105, 2106.

Larus philadelphia. - 1515, 1516, 2363.

Sterna fuliginosa. - 1772.

Phalacrocorax dilophus. - 2627.

Index to Specimens of Birds.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	<i>Merganser americanus</i> .	- 1343, 1344, 1835.				
	" <i>serrator</i> .	- 265, 1505, 1506, 1854, 1855, 1856, XIV p. 102.				
	<i>Lophodytes cucullatus</i> .	- 272, 808.				
	<i>Anas boschas</i> .	- 2059 (a hybrid or domestic duck. From Butterfuss of C. S. p. 9).				
	" <i>obscura</i> .	- 550, 1840, XXIII p. 103, 2516.				
	" <i>penelope</i> .	- 1819, (p. 467. 6846)				
	" <i>americana</i> .	- 888.				
	" <i>cyanoptera</i> .	- 2655 (Arizona).				
	<i>Spatula clypeata</i> .	- 1790.				
	<i>Aix sponsa</i> .	- 244, 1326, 1783; (CCX p. 108 = 244).				
	<i>Aythya valisineria</i> .	- 1341, 1342.				
	" <i>marila nearctica</i> .	- 1841, 2112, 2178, 2106a, 2106b.				
	" <i>affinis</i> .	- LXLIII p. 106. (2106b.?)				
	" <i>collaris</i> .	- LXLIV p. 105.				
	<i>Glaucochetta clangula americana</i> .	- 819, LXLII p. 104, 2177.				
	<i>Cidemia americana</i> .	- 796, 801, 803, 1499.				
	<i>Cidemia deglandi</i> .	- 792, 1500, 1501, 1502, 1503, 2110.				
	" <i>perspicillata</i> .	- 802, 2109, 2118, 2200.				
	<i>Erismatura rubida</i> .	- 261, 262, 1791.				
	<i>Somateria dresseri</i> ?	- 2638, 2639. From Labrador				
	<i>Branta canadensis</i> .	- 551.				
	" <i>bermuda</i> .	- 2175, 2176 + 2 following without numbers.				
	<i>Charitonetta albeola</i> .	- 264, 270, 271, XXXI p. 103, 2111				
	<i>Clangula hyemalis</i> .	- 806, 1336, 2111.				
	<i>Botaurus lentiginosus</i> .	- 2432.				
	<i>Ardea herodias</i> .	- 196, 219, 1773, 2173.				
	<i>Ardea egretta</i> .	- 2195.				
	<i>Ardea virescens</i> .	- 191, 758, 773, 775, 1838 , LXVIII p. 104, LXXIV p. 104.				
	<i>Nycticorax nycticorax naevius</i> .	- 126, 130, 131, 132, 133, 134, 139, 140, 1448, 1449, 1450, 1491, 1719, 1936, 1937, 1938, 1939, 1963, 1964, 1965, 2060 , 2064, 2101.				
	<i>Rallus virginianus</i> .	- 2305.				
	<i>Porzana carolina</i> .	- 170, 171, 212, 213, 745, 757, 1252, 1307, 1329, 1980, 1982, 1983, CLXXXI p. 107, CLXXXII p. 107, CLXLIV p. 107, 2078, 2079, 2080.				
	<i>Fulica americana</i> .	- 800, 2206.				
	<i>Philohela minor</i> .	- 267, LXXV p. 104, 2348, 2460.				

Index to Specimens of Birds.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	<i>Gallinago delicata</i> .	- 1312, 1313, 1325, 1870				
	<i>Tringa canutus</i> .	- 818.				
	<i>Tringa maritima</i> .	- ^{Did not buy one in market one day with Gains 3.} 2113, 2114, 2115, 2116, 2117.				
	<i>Tringa maculata</i> .	- 2093, 2094.				
	<i>Tringa minutilla</i> .	- 2090.				
	<i>Tringa alpina</i> .	- ^{pacific} 816, 2092.				
	<i>Ereunetes pusillus</i> .	- 2000, 2001, 2095, 2096, 2097, 2098, 2099, 2198, 2199.				
	<i>Totanus melanoleucus</i> .	- 2089, 2205, 2213, 2214, 2215, 2217.				
	<i>Totanus solitarius</i> .	- 180, 199, 217, 1160, 1255, 1390, 2002, LXXXII p. 104, CLXXXIV p. 107, 2068.				
	<i>Actitis macularia</i> .	- 68, 181, 562, 644, 960, 973, 1128, 1131, 1176, 1262, 1361, 1362, 1665, CCIX p. 108, 2427, 2428, 2536.				
	<i>Limosa haemaphysalis</i> .	- 1832.				
	<i>Numenius longirostris</i> .	- 156, CLII p. 106.				
	<i>Charadrius squatarola</i> .	- 817.				
	<i>Colinus virginianus</i> .	- 836, 1337, 2135 to 2144 (10 specimens), 2242, 2382, 2461.				
	<i>Bonasa umbellus</i> .	- 253, 269, 273, 411, 459, 484, 485, 876, 881, 1472, 1473, 1478, 1808, 1825, III p. 102, CCV p. 108, 2060, 2243, 2258, 2297, 2298, 2616, 2617, 2618, 2619.				
	<i>Gymnarchus americanus</i> .	- 455.				
	<i>Pediocetes phasianellus</i> .	- 1820, 1821.				
	<i>Colaptes migratorius</i> .	- 45, 46, 767, 768, 1396, 1847, 1849, 1850, 1851, 1852, 1869, VII p. 102, CVI p. 105, CXLIX p. 106.				
	<i>Zenaidura macroura</i> .	- 537, 963, 1874, 2008, 2025, 2132, 2133, 2134, 2241, 2361, 2539.				
	<i>Circus hudsonius</i> .	- 2240.				
	<i>Accipiter velox</i> .	- CLIII, 169, 1349, 1496, 1802.				
	<i>Accipiter cooperi</i> .	- 1226.				
	<i>Accipiter atricapillus</i> .	- 2005, 2170.				
	<i>Buteo borealis</i> .	- LXL, 544, 1853, 2179.				
	<i>Buteo borealis</i> .	- CL, 41, 1224, 1225, 1229, 1463, 1976, 2040, 2072, 2073, 2535 B., 6506 ^{Quelling} .				
	<i>Archibuteo lagopus sancti-johannis</i> .	- 2640.				
	<i>Haliaeetus leucocephalus</i> .	- LXLV, CLXXXIII, 195, 495, 1222, 1232, 1553, 2515.				
	<i>Falco peregrinus anatum</i> .	- 2100.				

Index to Specimens of Birds.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	<i>Falco columbarius</i> .	-2085.				
	<i>Falco sparverius</i> .	-1355, 2120.				
	<i>Pandion haliaetus carolinensis</i> .	-LXXXIX, LXXI, CLT, CLXXVII,				
		42, 43, 197, 2626,				
	<i>Asio wilsonianus</i> .	-2021, 2062, 2238.				
	<i>Asio accipitrinus</i> .	-2119.				
	<i>Syrnium nebulosum</i> .	-1328.				
	<i>Megascops asio</i> .	- ^{LXLVI} 241, 251, 712, 718, 719, 809, 1338, 1339, 1932,				
		1933, 1934, 2169, 2486, 2520, 2594.				
	<i>Bubo virginianus</i> .	-CXXI, 1324, 2239.				
	<i>Nyctea nyctea</i> .	-815, 2108.				
	<i>Coccyzus americanus</i> .	-CLXXV, 26, 27, 205, 1674, 1675, 1676, 2180.				
	<i>Coccyzus erythrophthalmus</i> .	- ^{CLXXIV} 204, 687, 720, 1150, 1757, 1922, 1985,				
		2052, 2438.				
	<i>Ceryle alcyon</i> .	-CXXI, CLXLIX, 50, 137, 172, 173, 221, 257, 258, 665,				
		735, 783, 787, 804, 1146, 1873.1, 1876, 2362, 2455, 2624.				
	<i>Dryobates villosus</i> .	-LXXVI, CXXXI, 168, 348, 437, 497, 1330, 1332,				
		1333, 1385, 1527, 1829, 1830, 1970, 2010, 2235, 2349, 2389, 2394,				
		2476, 2630.				
	<i>Dryobates pubescens</i> .	-LXX, LXXI, CXXVII, CLXI, CLXXII , CLXX, CCIV,				
		30, 63, 163, 175, 317, 337, 350, 371, 421, 449, 466, 478, 726,				
		738, 746, 853, 866, 875, 877, 879, 880, 957, 1331, 1529, 1557, 1567, 1593,				
		1710, 1769, 1792, 1823, 1839, 1953, 2009, 2121, 2314, 2342.				
	<i>Sphyrapicus varius</i> .	-CCVII, 238, 242, 243, 805, 882, 1282, 1292,				
		1296, 1304, 1305, 1327, 1358, 1482, 1626, 1635, 1684,				
		1780, 1861, 1865, 1868, 1885, 2202, 2367, 2496.				
	<i>Melanerpes erythrocephalus</i> .	-1334, 1492, 1493, 1525, 1526,				
		1528, 1530, 1531, 1661, 2123, 2124, 2125, 2126, 2127,				
		2128, 2203.				
	<i>Melanerpes carolinus</i> .	-2122, 2236, 2237.				
	<i>Colaptes auratus</i> .	-XXXII, LXXXIV, CLVII, 24, 25, 36, 37, 38,				
		39, 40, 177, 189, 260, 538, 583, 1132, 1330, 1350, 1615, 1871,				
		1942, 2022, 2046, 2355, 2356, 2358, 2376, 2437, 2457, 2518.				
	<i>Antrostomus vociferus</i> .	-4, 65, 641, 785, 1127, 1384, 1881, 2454,				
		2620, 2621.				
	<i>Chordeiles virginianus</i> .	-681.				
	<i>Chaetura pelagica</i> .	-LXI, 2, 3, 593, 634, 635, 636, 1741, 1777,				
		1903, 1987, 1988, 1989, 2048.				

Index to Specimens of Birds.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	<i>Trochilus colubris</i> .	-XXVI, LIV, 1, 188, 678, 697, 1703, 2061,				
		2421, 2562, 2586.				
	<i>Tyrannus tyrannus</i> .	-LXVI, LXIX, 5, 58, 128, 624, 628,				
		721, 1163, 1442, 1744, 1771, 1925, 2570, 2580, 2581, 2611.				
	<i>Myiarchus crinitus</i> .	- ^{LXXXIX, LXXXV} 6, 129, 152, 153, 178, 733, 734, 757,				
		1171, 1466, 1728,				
	<i>Sayornis phoebe</i> .	-XXIV, CXXVIII, CXXXIX, 192, 534, 920, 942,				
		948, 1112, 1395, 1604, 1605, 1609, 1611, 1768, 1935, 1992,				
		2351, 2352, 2357.				
	<i>Contopus borealis</i> .	-737, 749, 750, 1250, 1708.				
	<i>Contopus virens</i> .	-XLIV, 454, 729, 1230, 1236, 1237, 1238,				
		1239, 1256, 1723, 2066.				
	<i>Empidonax flaviventris</i> .	-XLII, (CCII?) , 714, 781, 1122,				
		1137, 1138, 1145, 1739.				
	<i>Empidonax pusillus traillii</i> .	-(CCII?), 1142, 1143, 2051,				
		(2606?).				
	<i>Empidonax minimus</i> .	-(CCII?), 95, 182, 185, 652, 653,				
		1135, 1136, 1139, 1140, 1141, 1144, 2041.				
	<i>Otocoris alpestris</i> .	-1822, 2163, 2164, 2165, 2166.				
	<i>Cyanocitta cristata</i> .	-I, IV, V, VI, LXXXIII, LXXXVI, LXXXVII,				
		CL, XL, CCVI, 59, 60, 211, 222, 248, 250, 451, 900, 1149,				
		1657, 1804, 1805, 1806, 2016, 2129, 2130, 2234, 2254, 2458,				
		2459, 2483, 2484, 2485, 2569, 2582, 2597, 2598.				
	<i>Corvus americanus</i> .	-CLXLVIII, 52, ²³⁴ 235, 236, 237, 813, 878,				
		1462, 1464, 1465, 1857, 1981, 2011, 2029, 2517, 2625.				
	<i>Corvus ossifragus</i> .	-1360, 2535.a.				
	<i>Dolichonyx oryzivorus</i> .	-LXXX, LXXXI, 706, 713, 1178, 1246,				
		1247, 1248, 1249, 1456, 1720.				
	<i>Molothrus ater</i> .	-CLX, 61, 62, 578, 912, 946, 965, 1616, 1627, 1628,				
		1629, 1630, 1659, 1668, 1669, 1876, 2023, 2042, 2043, 2395.				
	<i>Agelaius phoeniceus</i> .	-CLXVIII, CLXXXII, CLXLIII, CLXLIV, CLXLV,				
		CLXLVI, CC, CCVIII, 21, 22, 164, 176, 200, 201, 202, 215,				
		218, 240, 508, 704, 1152, 1157, 1161, 1166, 1195, 1371, 1446, 1487,				
		1622, 1633, 1700, 1701, 1748, 1749, 1750, 1949, 2028, 2033,				
		2081, 2082, 2380, 2381, 2418, 2419.				
	<i>Sturnella magna</i> .	-1345, 1620, 1621, 1623, 2020, 2091,				
		2107, 2131, 2197, 2231, 2232, 2233.				

Index to Specimens of Birds.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	<i>Icterus galbula</i> .	XLV, XLVIII, XLIX, L, LI, LVIII, LX, 1190, 1204 , 17, 18, 19, 20, 101, 109, 120, 122, 123, 190, 630, 631, 632, 662, 663, 722, 1179, 1190, 1204, 1244, 1414, 1670, 1690, 1699, 1721, 1760, 1833, 1956, 2426, 2559, 2571				
	<i>Icterus spurius</i> .	LVI, 66, 67, 119, 731, 1175, 1234, 1245, 1418, 1419, 1423, 1457, 1733, 1742, 1924, 2407, 2429.				
	<i>Sceloporus carolinus</i> .	XVIII, XIX, XXI, XL, 509, 549, 582, 807, 908, 909, 2366.				
	<i>Luscalus guscula</i> .	XXXIV, 1826, 2537, 2538, 2593.				
	<i>Pinicola enucleator</i> .	275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 293, 294, 295, 296, 297, 298, 299, 313, 314, 315, 316, 320, 321, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 352, 353, 354, 355, 412, 413, 469, 501, 502, 503, 504, 505, 506, 507, 841, 842, 843, 1795, 1796, 1797, 1798, 1799, 1800, 1801, 2464, 2465, 2466, 2467, 2491, 2497, 2508, 2509, 2510, 2511, 2636, 2637, 2641, 2642, 2643.				
	<i>Carpodacus purpureus</i> .	XXVII, XXXVII, XXXVIII, CCX , 224, 225, 247, 402, 403, 470, 471, 472, 473, 531, 539, 540, 693, 797, 798, 811, 837, 1167, 1314, 1356, 1523, 1524, 1532, 1537, 1554, 1555, 1556, 1564, 1565, 1566, 1570, 1571, 1572, 1573, 1592, 1603, 1610, 1891, 2017, 2193, 2194, 2211, 2245, 2288, 2293, 2331, 2343, 2373, 2526, 2529, 2531, 2612.				
	<i>Loxia curvirostra minor</i> .	333, 334, 335, 339, 340, 341, 342, 351, 356, 357, 400, 414, 422, 423, 490, 491.				
	<i>Loxia leucoptera</i> .	343, 344, 345, 346, 347, 358, 359, 360, 361, 362, 2462, 2463.				
	<i>Acanthis linaria</i> .	300, 301, 302, 303, 319, 364, ³⁶⁵ 375, 376, 377, 378, 379, 380, 382, 385, 386, 387, 388, 389, 390, 391, 392, 393, 394, 395, 396, 397, 398, 399, 417, 426, 427, 428, 429, 430, 431, 432, 2492, 2493, 2494, 2498, 2499, 2500, 2501, 2502, 2503, 2504, 2505, 2514, ^{452, 453, 479, 480, 498, 528, 529, 530, 535, 536, 1809 to 1818,} ^{2472 to 2474, 2478 to 2480, 2487 to 2489,}				
	<i>Pooecetes gramineus</i> .	XXX, 564, 565, 568, 569, 570, 571, 572, 580, 918, 925, 1667, 2371.				
	<i>Ammodramus sandwicensis savanna</i> .	557, 613, 614, 615, 1260, 1643, 1644, 1653, 1872, 2102.				

Index to Specimens of Birds.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	<i>Ammodramus savannarum passerinus</i> .	1715, 1716.				
	<i>Ammodramus caudacutus nelsoni</i> .	CCXII, 246, 1496.				
	<i>Zonotrichia leucophrys</i> .	1177, 1181, 2424.				
	<i>Zonotrichia albicollis</i> .	XXXIX, CLXXI, CLXXII, 124, 127, 239, 259, 602, 603, 604, 605, 705, 1151, 1168, 1283, 1370, 1387, 1388, 1488, 1785, 2044, 2388, 2397.				
	<i>Spizella monticola</i> .	LXLVII, C, CI, CXLII, 291, 338, 372, 383, 420, 425, 436, 438, 443, 457, 458, 462, 463, 477, 558, 828, 830, 858, 862, 1620, 1521, 1538, 1545, 1546, 1547, 1787, 1803, 2015, 2477.				
	<i>Spizella socialis</i> .	CXLIV, CXLV, CXLVI, CXLVII, 115, 584, 585, 617, 774, 941, 1113, 1118, 1619, 1960, 2003.				
	<i>Spizella pusilla</i> .	LXXII, CXLIII, CCI, 69, 619, 914, 1116, 1288, 2047, 2368, 2530.				
	<i>Junco hyemalis</i> .	X, XXXVI, CIII, CXVII, CXXIV, CXXIX, CXXX, 254, 256, 418, 424, 456, 499, 563, 618, 850, 860, 867, 868, 869, 870, 871, 895, 902, 903, 944, 1306, 1347, 1607, 1631, 1858, 2271, 2287, 2307, 2325, 2326, 2327, 2370, 2533, 2628.				
	<i>Melospiza fasciata</i> .	IX, XI, CH , CXX, CLXXX , CLXLVII, 28, 103, 210, 349, 363, 384, 419, 435, 444, 510, 512, 520, 533, 724, 861, 887, 896, 906, 1577, 1614, 1652, 1654, 1828, 1864, 2013, 2260, 2369, 2372, 2412, 2482, 2632, 2633.				
	<i>Melospiza georgiana</i> .	CCXIII, 245, 252, 559, 560, 561, 917, 958, 966, 967, 1293, 1294, 1458, 1651, 2035, 2522, 2523.				
	<i>Passerella iliaca</i> .	CIX, CX, CXVIII, CXIX, 113, 114, 193, 518, 519, 541, 548, 886, 897, 898, 901, 1348, 1507, 1513, 1569, 1574, 1608, 1613, 1842, 1845, 1846, 1863, 2013, 2014, 2339, 2340, 2341, 2346, 2347, 2350, 2495.				
	<i>Pipilo erythrophthalmus</i> .	XLI, CCIII , 64, 146, 147, 223, 579, 611, 633, 1147, 1155, 1156, 1374, 1401, 1639, 1647, 1658, 1907, 1993, 2065, 2161, 2162, 2396, 2403, 2553, 2572.				
	<i>Cardinalis cardinalis</i> .	2151, 2152, 2153, 2154, 2155, 2156, 2157, 2158, 2159, 2160, 2225, 2226, 2227, 2228, 2229, 2230.				
	<i>Heabia ludoviciana</i> .	LII, 12, 13, 14, 15, 16, 684, 685, 1170, 1197, 1202, 1203, 1420, 1421, 1490, 1685, 2050, 2181, 2420, 2423, 2601, 2602, 2605.				

Index to Specimens of Birds.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	<i>Passerina cyanea</i> .	—	LIII, LXXIII, 117, 118, 141, 142, 148, 194, 1218, 1219, 1391, 1443, 1481, 1483, 1713, 1714, 1974, 1975, 2399, 2451, 2599.			
	<i>Spinus tristis</i> .	—	CXXVI, CLV, 70, 71, 292, 814, 820, 821, 822, 823, 831, 832, 833, 844, 846, 847, 848, 890, 905, 1120, 1121, 1270, 1378, 1460, 1461, 1533, 1534, 1536, 1549, 1585, 1648, 2173, 2246, 2247, 2330, 2359, 2391, 2631.			
	<i>Spinus pinus</i> .	—	249, 381, 404, 405, 406, 407, 408, 409, 410, 416, 1548, 2103, 2512, 2513.			
	<i>Plectrophenax nivalis</i> .	—	439, 440, 441, 442, 481, 482, 483, 494, 496, 500.			
	<i>Passer domesticus</i> .	—	CV, 474, 492, 493, 511, 515, 910, 947, 961, 1124, 1514, 1575, 1576, 1579, 1580, 1890, 2324, 2337, 2338, 2506.			
	<i>Piranga ludoviciana</i> .	—	2244.			
	<i>Piranga erythromelas</i> .	—	LXII, LXIV, LXXVII, LXXXVIII, CLXXVI, 80, 9, 29, 206, 664, 671, 673, 694, 743, 770, 1133, 1169, 1172, 1173, 1174, 1182, 1183, 1184, 1185, 1186, 1187, 1188, 1191, 1192, 1193, 1198, 1199, 1200, 1242, 1243, 1254, 1403, 1422, 1431, 1432, 1445, 1696, 1698, 1727, 1752, 1758, 1831, 1892, 1930, 1931, 2058, 2191, 2560, 2584, 2585, 2610.			
	<i>Piranga rubra</i> .	—	2583.			
	<i>Petrochelidon lunifrons</i> .	—	34, 150, 594, 675, 715, 1180, 1451, 1452, 1453, 1454.			
	<i>Chelidon erythrogaster</i> .	—	35, 586, 592, 600, 601, 607, 608, 623, 1258, 1375, 1640, 1641, 1724, 1725, 1726, 1754, 1755, 1973, 2047, a, 2456.			
	<i>Tachycineta bicolor</i> .	—	31, 32, 33, 216, 591, 595, 596, 598, 599, 606, 609, 610, 924, 1346, 1735, 2595.			
	<i>Clivicola riparia</i> .	—	1205, 1231, 1441, 1455, 1966, 1968, 1969, 1971, 1972.			
	<i>Stelgidopteryx serripennis</i> .	—	1836, 1967, 2452, 2453.			
	<i>Ampelis cedrorum</i> .	—	XXXIII, XLVI, LXXVIII, CXXV, 104, 105, 106, 151, 162, 198, 274, 460, 461, 475, 476, 488, 489, 522, 523, 524, 525, 543, 545, 739, 740, 764, 853, 855, 856, 859, 891, 892, 893, 894, 904, 907, 1235, 1522, 1558, 1559, 1560, 1562, 1563, 1568, 1578, 1581, 1582, 1586, 1587, 1588, 1589, 1590, 1591, 1594, 1595, 1596, 1597, 1598, 1599, 1601, 1602, 1747, 1761, 1838, 1843, 1844, 1848, 2196, 2261, 2262, 2263, 2264, 2265, 2266, 2267, 2270, 2272, 2273, 2274, 2275, 2276, 2277, 2280, 2281, 2282,			

Index to Specimens of Birds.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	<i>Ampelis cedrorum</i> — Continued.					
			2294, 2295, 2296, 2306, 2315, 2316, 2317, 2318, 2319, 2320, 2321, 2322, 2323, 2332, 2333, 2334, 2335, 2336.			
	<i>Lanius borealis</i> .	—	XX, 824, 1340, 1824, 2171, 2216.			
	<i>Lanius ludovicianus</i> .	—	2345.			
	<i>Vireo olivaceus</i> .	—	LXVII, CLXXIX, CLXXXVI, CLXXXVII, 57, 100, 136, 161, 203, 209, 233, 709, 723, 1216, 1241, 1263, 1266, 1269, 1413, 1474, 1678, 1745, 2069, 2449, 2613.			
	<i>Vireo gilvus</i> .	—	696, 1213, 1214, 1215, 1367, 1900, 2400, 2414.			
	<i>Vireo flavifrons</i> .	—	97, 98, 99, 1119, 1196, 1210, 1211, 1212, 1691, 1711, 1948, 1962, 2566, 2615.			
	<i>Vireo solitarius</i> .	—	742, 771, 977, 1111, 1117, 1206, 1207, 1208, 1209, 1285, 1351, 1363, 1366, 1999, 2037, 2406, 2550.			
	<i>Vireo noveboracensis</i> .	—	CLXXVIII, 184, 679, 1217, 1471, 1906, 2401, 2567.			
	<i>Mniotilta varia</i> .	—	XLVII, 77, 575, 620, 666, 672, 736, 981, 1007, 1073, 1083, 1227, 1268, 1317, 1357, 1359, 1372, 1673, 1746, 1765, 1879, 1887, 2385, 2575.			
	<i>Helminthosus vermivorus</i> .	—	107, 144, 174, 994, 1033, 1069, 1095, 1102, 1106, 1468, 1469, 1734, 1834, 2054, 2185, 2410, 2446, 2574, 2600.			
	<i>Helminthophila pinus</i> .	—	1415, 2607.			
	<i>Helminthophila chrysoptera</i> .	—	1686, 1707, 1908, 1928, 1944, 1945, 1946, 1950, 1951, 1952, 2053, 2184, 2187, 2425.			
	<i>Helminthophila ruficapilla</i> .	—	92, 93, 94, 639, 668, 669, 689, 763, 974, 991, 1002, 1074, 1085, 1088, 1090, 1277, 1687, 2408, 2415, 2447, 2545, 2549, 2554, 2555, 2568.			
	<i>Helminthophila celata</i> .	—	667.			
	<i>Helminthophila peregrina</i> .	—	91, 703, 1279, 1289.			
	<i>Comptosia americana</i> .	—	56, 154, 231, 655, 670, 680, 695, 710, 711, 772, 996, 1075, 1267, 1272, 1286, 1381, 1408, 1776, 1918, 2434, 2442, 2544, 2592.			
	<i>Dendroica tigrina</i> .	—	759, 1275, 2603, 2604.			
	<i>Dendroica aestiva</i> .	—	75, 183, 698, 699, 1046, 1417, 1717, 1718, 2398.			
	<i>Dendroica caerulescens</i> .	—	81, 82, 638, 646, 677, 686, 690, 700, 707, 708, 989, 990, 998, 1050, 1087, 1093, 1096, 1100, 1287, 1300, 1310, 1316, 1389, 1407, 1411, 1412, 1706, 1775, 1893, 1915, a, 1986, 1995, 2049, 2087, 2431, 2564.			

Index to Specimens of Birds.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	<i>Dendroica coronata</i> .	XXXV, 83, 87, 232, 576, 587, 588, 597, 779, 799, 916, 934, 945, 968, 969, 970, 980, 1080, 1081, 1104, 1509, 1517, 1544, 1550, 1613, 1786, 1880, 1896, 2248, 2269, 2289, 2290, 2291, 2292, 2384, 2416, 2524, 2532, 2547, 2548, 2551.				
	<i>Dendroica maculosa</i> .	73, 74, 637, 661, 676, 753, 984, 997, 1012, 1014, 1017, 1024, 1058, 1089, 1264, 1298, 1309, 1404, 1927, 2057, 2076, 2190, 2443, 2444, 2557, 2576, 2587.				
	<i>Dendroica caerulea</i> .	674.				
	<i>Dendroica pennsylvanica</i> .	85, 86, 88, 89, 187, 648, 692, 976, 1001, 1035, 1039, 1044, 1045, 1261, 1281, 1406, 1437, 1476, 1477, 1756, 1910, 1943, 2402, 2543, 2591.				
	<i>Dendroica castanea</i> .	LIX, 84, 756, 788, 995, 1010, 1011, 1022, 1027, 1028, 1029, 1030, 1031, 1042, 1047, 1048, 1051, 1067, 1070, 1097, 1276, 1299, 1439, 1467, 1737, 1915, 1920, 2086, 2188, 2189.				
	<i>Dendroica striata</i> .	CLXXXVIII, CLXXXIX, 76, 155, 214, 754, 755, 762, 776, 777, 778, 1013, 1026, 1040, 1041, 1052, 1057, 1440, 1470, 1921, 1923, 1926, 1996, 1997, 2441, 2609.				
	<i>Dendroica blackburniae</i> .	96, 207, 208, 760, 985, 1005, 1006, 1008, 1009, 1015, 1018, 1036, 1037, 1043, 1054, 1059, 1062, 1076, 1099, 1101, 1103, 1108, 1290, 1291, 1405, 1409, 1410, 1424, 1425, 1429, 1738, 1740, 1743, 1979, 2409, 2590.				
	<i>Dendroica virens</i> .	47, 57, 138, 143, 647, 691, 747, 752, 780, 784, 975, 979, 1020, 1079, 1086, 1091, 1094, 1265, 1271, 1273, 1280, 1301, 1433, 1484, 1692, 1712, 1886, 1894, 1897, 1979, 2038, 2039, 2084, 2405, 2546, 2556, 2577.				
	<i>Dendroica vigorsii</i> .	589, 741, 930, 1352.				
	<i>Dendroica palmarum hypochrysea</i> .	CLIX, 90, 590, 612, 622, 911, 931, 932, 933, 935, 937, 962, 1077, 1082, 1393, 1394, 1494, 1497, 1634, 1645, 1866, 1998, 2018, 2019, 2024, 2047, 2387.				
	<i>Dendroica discolor</i> .	1479.				
	<i>Seiurus aurocapillus</i> .	CLIV, 55, 625, 761, 1004, 1021, 1038, 1071, 1092, 1228, 1368, 1369, 1380, 1889, 1904, 1905, 2430, 2565.				

Index to Specimens of Birds.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	<i>Seiurus noveboracensis</i> .	992, 1063, 1066, 1251, 1257, 1392, 1638, 1642, 2070, 2071, 2192.				
	<i>Seiurus motacilla</i> .	CLXIV, CLXV, CLXVI, CLXVIII, 53, 54, 220, 574, 621, 725, 936, 949, 950, 955, 964, 1068, 1110, 1353, 1379, 1489, 1624, 1632, 1636, 1637, 1677, 1697, 1877, 1901, 1902, 2031, 2032, 2034, 2183, 2364, 2383, 2519, 2534, 2596.				
	<i>Geothlypis philadelphia</i> .	1000, 2432.				
	<i>Geothlypis trichas</i> .	78, 626, 627, 999, 1019, 1064, 1072, 1098, 1105, 1240, 1364, 1382, 1459, 1480, 1666, 1899, 1929, 1954, 1955, 1959, 2083, 2411, 2435, 2436, 2561, 2608.				
	<i>Icteria virens</i> .	79, 80, 682, 1032, 1061, 1485, 1709, 1729, 1882, 1909, 1940, 2440, 2622, 2623.				
	<i>Sylvania nitrata</i> .	135, 640, 642, 645, 657, 701, 702, 728, 978, 993, 1025, 1060, 1065, 1084, 1107, 1223, 1426, 1427, 1428, 1475, 1662, 1663, 1680, 1681, 1682, 1683, 1688, 1689, 1704, 1705, 1730, 1731, 1762, 1766, 1767, 1912, 1913, 1941, 1947, 1957, 1958, 2063, 2074, 2075, 2563, 2573.				
	<i>Sylvania pusilla</i> .	656, 1416, 1916, 2055, 2056, 2445, 2688.				
	<i>Sylvania canadensis</i> .	983, 986, 987, 988, 1016, 1023, 1034, 1049, 1053, 1055, 1109, 1259, 1430, 1434, 1435, 1438, 1911, 1917, 1919, 2067, 2182, 2186, 2433, 2448.				
	<i>Setophaga ruticilla</i> .	XLIII, LXIII, 186, 654, 688, 727, 748, 1003, 1056, 1078, 1365, 1373, 1377, 1646, 1679, 1764, 1878, 1895, 1898, 1914, 1978, 2077, 2413.				
	<i>Saloscopes carolinensis</i> .	LVII, 121, 629, 1126, 1253, 1376, 1751, 2417, 2558.				
	<i>Geothlypis rufus</i> .	LV, CLXXIII, CLXXX, 102, 145, 1148, 1295, 1386, 1400, 1656, 1763, 2036, 2377, 2392, 2393, 2535.				
	<i>Thryothorus ludovicianus</i> .	2167, 2168.				
	<i>Troglodytes aëdon</i> .	616, 1114, 1125, 1221, 1274, 1315, 1660, 1770, 2541.				
	<i>Troglodytes hiemalis</i> .	(CXL = 7), 263, 374, 401, 829, 872, 1308, 2045, 2481.				

Index to Specimens of Birds.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	<u><i>Cistothorus palustris</i></u> .	— LXV, 72, 717, 1444, 1447, 1664, 1736, 1961, 1990, 1991.		732, 1123, 1278,		
	<u><i>Certhia familiaris americana</i></u> .	— 149, 370, 446, 447, 448, 812, 826, 827, 921, 940, 1311, 1512, 1867, 2344.		255, 368, 838, 873, 874,		369, 874,
	<u><i>Sitta carolinensis</i></u> .	— XV, XVI, XVII, 226, 227, 228, 450, 465, 835, 849, 854, 1583, 1584, 1788, 1789, 1807, 2150, 2174, 2470, 2471.		373, 1807,		
	<u><i>Sitta canadensis</i></u> .	— CLV , 157, 158, 159, 434, 526, 527, 2629.		160, 415, 433,		
	<u><i>Parus bicolor</i></u> .	— 2145, 2146, 2147, 2148, 2149, 2220.				
	<u><i>Parus atricapillus</i></u> .	— II, LXLIX, 857, 863, 1220, 1284, 1539, 1540, 1561, 1649, 1650, 1793, 1794, 2221, 2268, 2311, 2312.		336, 464, 825, 851,		851,
	<u><i>Parus carolinensis</i></u> .	— 2222, 2223, 2224.				
	<u><i>Regulus satrapa</i></u> .	— CXXXIV, CXXXV, CXXXVII, CXXXVIII, 11, 288, 366, 367, 445, 566, 834, 839, 840, 845, 864, 865, 957, 1302, 1303, 1510, 1511, 1518, 1519, 1541, 1542, 1543, 2219, 2468, 2469, 2475.				
	<u><i>Regulus calendula</i></u> .	— CXXXVI, CXLIII, CXLVII, 10, 577, 581, 913, 915, 922, 926, 927, 928, 929, 943, 952, 953, 954, 958, 982, 1318, 1354, 1625, 1883, 1884, 2026, 2027, 2386, 2525, 2540.				
	<u><i>Turdus mustelinus</i></u> .	— 650, 1129, 1134, 1397, 1671, 1672, 1693, 1695, 1722, 1984, 2552, 2578, 2579.				
	<u><i>Turdus fuscescens</i></u> .	— 1130, 1158, 1201, 1399, 1402, 2404.				
	<u><i>Turdus olivaceus</i></u> .	— 229, 230, 786, 790, 791, 793, 794, 795, 1153, 1159, 1319, 1436, 1753, 1778, 1779, 2439, 2450, 2589.				
	<u><i>Turdus ustulatus swainsoni</i></u> .	— 643, 649, 658, 659, 660, 661, 769, 789, 971, 972, 1162, 1164, 1165, 1189, 1194, 1233, 1297, 1398, 1694, 2088.				
	<u><i>Turdus aonalaschkae pallasii</i></u> .	— XXIX, CLXIX, 116, 289, 290, 555, 556, 923, 938, 939, 1154, 1322, 1323, 1335, 1486, 1498, 1504, 1618, 1784, 1873, 2004, 2030, 2201, 2360, 2365, 2390, 2490, 2542.				

Index to Specimens of Birds.

Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.
	<u><i>Merula migratoria</i></u> .	— VII, XXV, CXVI, CLVIII, 108, 165, 166, 167, 179, 467, 468, 486, 521, 546, 547, 683, 730, 810, 899, 1600, 1606, 1617, 1655, 1732, 1759, 1827, 1837, 1862, 2204, 2208, 2209, 2210, 2255, 2256, 2257, 2278, 2279, 2284, 2285, 2286, 2353, 2384, 2507, 2521, 2614.				
	<u><i>Sialia sialis</i></u> .	— XII, XIII, XXVIII, LXLVIII, CIV, CVII, CVIII, CXI, CXII, CXIII, CXIV, CXV, CXXIII, 23, 44, 48, 49, 110, 111, 112, 125, 304, 305, 306, 307, 308, 309, 310, 311, 312, 318, 487, 513, 514, 516, 517, 532, 542, 554, 744, 766, 883, 884, 885, 889, 919, 959, 1115, 1321, 1508, 1535, 1567, 1552, 1702, 2006, 2218, 2249, 2250, 2251, 2252, 2253, 2283, 2299, 2300, 2301, 2302, 2303, 2304, 2308, 2309, 2310, 2313, 2328, 2329, 2374, 2375, 2378, 2379, 2634, 2635.				

Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.Collecting
Number.

NAME.

DATE.

LOCALITY.

Length.

Stretch.

Sex and
Age.

	Sex and Age.	Collecting Number.	NAME.	DATE.	LOCALITY.	Length.	Stretch.	Sex and Age.

8 X 11 1/2

Epinus maximus

Whippoorwill, Woodcock and Field Sparrow
were first seen or heard April 24th, in 1895, at
Ft. Myer, Va. E. A. M.

Edgar A. Kearney, Esq.,
131 West 23rd St.,
New York City.

WRITE THE ADDRESS ON THIS SIDE - THE MESSAGE ON THE OTHER

Stevens Rifle. 15 inch barrel. 5 shots fired.
Distance (measured) 100 yards
Elevation 4 half turns. March 19, 1883.
The other shots struck just below the paper.

Stevens rifle at 160 feet. 9 shots fired.
March 17th, 1883. No elevation.

"See
and
Hudson
in Ph
88
57)

p2.
m. 55,

58.
8.
0.

QUESTIONS RELATING TO BIRDS AND

PLEASE GIVE WHAT INFORMATION YOU CAN IN ANSWER TO
ANY ADDITIONAL ITEMS OF INTEREST REGARDING
MAY BE WRITTEN ON THE BACK OF THIS

Name of species.....

Name of person furnishing the information

Address

1. The average number of eggs in a set or clutch?.....
2. The largest number of eggs in one nest?.....
3. The earliest date of nesting?.....
4. The latest date of nesting?.....
5. Whether this species has one or more broods in a season?.....

Notes on City Birds Outside
the Park.

May 1, 1881, p. 214. Purple Martin & Hawk
May 1, 1881, p. 416. Barn Swallow.

For New York, via Phila. & Reading R.R., 2:45, 9, 11
4:30, 6 p. m. For Keyport, via Phila. & Reading R.R., 11:15
a. m., 2:45, 4:30, 6 p. m. For Vineland, Bridgeton, Atlantic City,
&c., 2:45 p. m.
G. G. HANCOCK, J. R. WOOD, H. H. NIEMAN,
G. P. & T. A., P & R. RR. G. P. A. P. RR. Act'g Supt.

**NEW-YORK CENTRAL AND HUDSON
RIVER RAILROAD.**—Commencing Sept. 17, 1883, through
trains will leave Grand Central Depot:

8 a. m., Western New-York and Northern Express, with drawing,
room cars to Montreal via Central Vermont R.R., and to Rochester;
also to Saratoga.

*8:40 a. m., Fast Limited Chicago Express, with dining cars, stop-
ping at Albany, Utica, Syracuse, Rochester, Buffalo, Erie, Cleveland
and Toledo, arriving at Chicago 10:20 a. m. next day.

10:30 a. m., Chicago Express, drawing-room cars to Canandaigua,
Rochester and Buffalo; connects for Sharon Springs.

11 a. m., to Albany and Troy, connecting for Utica, Saratoga,
Glen's Falls, Lake George and Rutland.

3:30 p. m., Albany and Troy special.

4 p. m., Accommodation to Albany and Troy.

*6 p. m., St. Louis Express, with sleeping cars for Niagara Falls,
Buffalo, Cincinnati (except Saturday), Toledo, Detroit and St.
Louis.

6:30 p. m., Express, with sleeping cars to Syracuse and to Auburn
Road; also to Saratoga and Montreal.

*9 p. m., Pacific Express, with sleeping cars for Rochester, Buffalo,
Cleveland, Toledo, Detroit, Chicago, St. Louis, Cape Vincent (except
Sunday), and Clayton (except Saturday.)

11 p. m., Night Express, with sleeping cars to Albany and Troy.
Connects with the morning trains for the West and North (except
Saturday night.)

Tickets on sale at Grand Central Depot, No. 5 Bowling Green, 252
and 413 Broadway, and at Westcott's Express Offices, 3 Park-place,
and 785 and 942 Broadway, New-York, and 333 Washington and
864 Fulton-st., Brooklyn, and 79 4th-st., Williamsburg.

Baggage called for and checked from residence.

*These trains run daily; all others daily except Sunday.

J. M. TOUCEY, C. B. MEEKER,
Gen. Supt. Gen. Passenger Agent.

PENNSYLVANIA RAILROAD.
On and after Oct. 2, 1883.

GREAT TRUNK LINE

AND UNITED STATES MAIL ROUTE.

Trains leave New-York via Desbrosses and Cortlandt

Stations as follows:

New-York, Pittsburg

"See a list of animal, vegetable
and mineral productions which
Hudson saw on his voyage
in Philosophical Discourses by
Dr Witt Clinton, S. L. D., Note
57, p 117." [Page 270]

2346

2346

Fox Sparrow. March 20, 1882.
7.45; 11.30; 3.55; 3.20; culmen; .55;
gape, .58; tarsus, .93; mte. .97.

Fox Sparrow. 2347

7.20; 11.30; 3.53; 2.92; cul., .58;
gape, .62; tarsus, .95; mte., .98.

Notes on City Birds: C. P.

March 29, 1881.	p.	202
April 3, 1881.	p.	206
April 17, 1881.	p.	208
April 24, 1881.	p.	210
May 1, 1881.	p.	214.
May 4, 1881.	p.	217.
May 13, 1881.	p.	227
November 30, 1881.	p.	253
March 12, 1882	p.	294

[Notes on Birds outside the Park are
not here indexed.]

1020
DeLafield April 12, 1881.

Yellow-bellied Woodpecker
nesting a series
12th St & 6th Avenue!

Fishes of the Hudson River Valley.

Notemigonus chrysolenus (Mitch.) Ford.
No. 128-133. Shiner; Stille; Kream.
Echo Lake, Orange Co. N.Y. 7, 1876.
EDGAR A. MEARNS, M. D.

Stevens rifle at 160 feet. 9 shots fired.
March 17th, 1883. No elevation.

MEARNS HOUSE,

HIGHLAND FALLS, N. Y.

Office of EDGAR A. MEARNS, M. D.

March 16th, 1883.

Dear Sir

Will you kindly send
a copy of vol. 3 of the
"Proceedings of the U.S. National
Museum" and very greatly
oblige

Yours very respectfully
To Prof. Spencer F. Baird
Secy Smithsonian
Washington, D.C.

Stevens Rifle. 15 inch barrel. 58 shots fired.
Distance (measured) 100 yards
Elevation 4 half turns. March 19, 1883.
The other shots struck just below the paper.

WRITE THE ADDRESS ON THIS SIDE—THE MESSAGE ON THE OTHER

Edgar A. Inearns, Esq.,
131 West 23 street,
New York City.

Woodcock in city West 72 Str.,
New York City.
F. & Stream V, 228. Mar. 27, 1881.

In. car. —;

In case you get
out No. 1 of "Early Birds", I
send you these; viz. (from note
book);

Central Park Apr. 27, 1881.

Hirundo horreorum.

Quiscalus purpureus.

May 11. 1881.

Hirundo horreorum.

Quiscalus purpureus.

Chaetura pelagica.

Expect you & Morris
about 3.00 P. M., next Sunday; a
walk in "the Park" & expect you
both to tea & to stand the evening.
Be sure to come, to the
yours, etc.

See birds in Tribune building
Forest and Stream V, 195.
Ed. Mearns.