For Reference

NOT TO BE TAKEN FROM THIS ROOM

A STUDY OF MEIOSIS IN A HAPLOID OF
TRITICUM VULGARE AND ITS PROGENIES

R.C. McGinnis

Department of Plant Science
University of Alberta

Thesis

1961

#28

Ex LIBRIS
UNIVERSITATIS
ALBERTAENSIS

Digitized by the Internet Archive
in 2017 with funding from
University of Alberta Libraries

<https://archive.org/details/studyofmeiosisin00robe>

T H E U N I V E R S I T Y O F A L B E R T A

A STUDY OF MEIOSIS IN A
HAPLOID OF TRITICUM VULGARE AND ITS PROGENIES

A DISSERTATION
SUBMITTED TO THE SCHOOL OF GRADUATE STUDIES
IN PARTIAL FULFILMENT OF THE REQUIREMENTS FOR THE DEGREE
OF MASTER OF SCIENCE

FACULTY OF AGRICULTURE
DEPARTMENT OF PLANT SCIENCE

by

Robert Cameron McGinnis

EDMONTON, ALBERTA

APRIL, 1951

ABSTRACT

A study was conducted on the meiotic behaviour of a 21 chromosome haploid of Triticum vulgare and its progenies.

Metaphase I and anaphase I of meiosis were not clearly differentiated, but chromosome counts were made when the univalents had migrated to the poles. Random distribution to the poles was apparently restricted by the presence of true bivalents and secondary pairing. When no pairing occurred, the distribution of univalents appeared random.

One, two or three bivalents were present in 67.5% of the cells counted. The presence of true bivalents indicated considerable homology between chromosomes not normally considered homologous. No bridges or fragments were observed.

A few cells contained a completely reduplicated set of chromosomes. In these cases, 42 chromosomes were present. It is believed that chromosome doubling with failure of cell wall formation in a sporogenous cell prior to microsporocyte formation, resulted in the duplications observed.

Chains of 3, 4, and 5, were observed in the haploid. These associations appeared as continuations of secondary pairing.

A study was conducted on the effect of the following factors on the rate of reaction of the following reaction:

$$2\text{H}_2\text{O}_2 \rightarrow 2\text{H}_2\text{O} + \text{O}_2$$

The reaction was carried out in a closed system at a constant temperature of 25°C. The rate of reaction was determined by measuring the volume of oxygen gas evolved over a period of 10 minutes. The following factors were varied:

- Concentration of H_2O_2
- Concentration of Fe^{2+} ions
- Temperature
- Catalyst

The results of the experiment are shown in the following table:

Factor	Initial Concentration	Final Concentration	Volume of O_2 (ml)
Concentration of H_2O_2	0.1 M	0.05 M	10
	0.2 M	0.1 M	20
	0.3 M	0.15 M	30
Concentration of Fe^{2+} ions	0.01 M	0.005 M	10
	0.02 M	0.01 M	20
	0.03 M	0.015 M	30
Temperature	25°C	20°C	10
	30°C	25°C	20
	35°C	30°C	30
Catalyst	None	None	10
	None	Fe^{2+}	20
	None	Mn^{2+}	30

From the results of the experiment, it can be concluded that the rate of reaction is affected by all four factors. The rate of reaction increases with increasing concentration of H_2O_2 , increasing concentration of Fe^{2+} ions, increasing temperature, and increasing concentration of catalyst.

The F₂ and F₃ generations were studied cytologically. Although a high frequency of aberrations was observed, a general trend toward a normal complement of chromosomes was indicated in later generations. Univalents, trivalents and quadrivalents were the most frequent aberrations. It is postulated that crossing-over between partially homologous chromosomes in the haploid resulted in reciprocal translocations which were transmitted to succeeding generations. Deficient and duplicated gametes resulting from such translocations probably gave rise to aneuploid plants with various types of associations in later generations.

TABLE OF CONTENTS

	<u>Page</u>
INTRODUCTION	1
LITERATURE REVIEW	3
Occurrence	3
Origin	4
Meiotic Behaviour	6
MATERIALS AND METHODS	12
EXPERIMENTAL RESULTS AND INTERPRETATIONS	14
CYTOLOGICAL DATA FROM THE HAPLOID PLANT	14
Distribution of Univalents	14
Chromosome Associations	16
Paired Chromosomes	16
Chains	20
Cells with Reduplicated Chromosomes	20
CYTOLOGICAL DATA FROM THE F ₂ AND F ₃ GENERATIONS ..	25
CONCLUSIONS	34
ACKNOWLEDGEMENTS	35
REFERENCES	36

A STUDY OF MEIOSIS IN A
HAPLOID OF TRITICUM VULGARE AND ITS PROGENIES

R. C. McGinnis

INTRODUCTION

A haploid plant contains the gametic number or only half the normal chromosome complement. Haploids are usually somewhat dwarfed and less vigorous than their sister diploids, and their reproductive organs are correspondingly reduced in size. They occur rarely in most species under natural conditions, and presumably arise from parthenogenic development of an egg cell or an accompanying synergid.

Because of the aberrant chromosomal behaviour during meiosis, haploids are generally highly sterile. Theoretically however, they need not be completely sterile, for no matter how many chromosomes a species may have, a certain percentage of the eggs and sperms should contain a haploid set of chromosomes.

STATE OF NEW YORK
IN SENATE
JANUARY 1, 1901.

REPORT

OF THE

COMMISSIONERS OF THE LAND OFFICE
IN RESPONSE TO A RESOLUTION PASSED BY THE SENATE
JANUARY 1, 1901.
ALBANY: J. B. LEECH, STATE PRINTER.
1901.

ALBANY: J. B. LEECH, STATE PRINTER.
1901.

Union of such gametes would give rise to a diploid plant which would be completely homozygous for all characters. Diploids could also be obtained from haploids by the union of unreduced gametes. Such diploid plants would be of considerable value for studying the genetics of quantitative characters.

Haploids have been of interest in the study of the phylogeny of a species. Cytological observations of the extent of chromosome pairing tendencies in haploids, have given information concerning basic chromosome numbers and relationships in the species studied.

A practical application of haploidy has recently been demonstrated with the derivation of monosomic lines in common wheat. Monosomics in turn, have been found of value in associating certain genes with specific chromosomes and may also prove to be useful for transferring inherited characteristics by a method of chromosome substitution from one variety to another.

In Triticum vulgare, haploids are primarily of interest in a study of chromosomal associations between genomes, and the meiotic behaviour of unpaired chromosomes. Aberrations involving whole chromosome deficiencies or duplications, and reciprocal translocations observed in the progeny of haploids, make possible a more complete study of linkage and inheritance in this species.

In this paper the results of cytological investigations of chromosome behaviour at meiosis of a haploid of Triticum vulgare and its progenies are presented and discussed.

LITERATURE REVIEW

Occurrence

Work on haploid plants has become prominent only in recent years and records of haploidy are still rare. The first haploid seed plant was described by Blakeslee, et al (1), in 1922. Two such plants appeared in the offspring of normal *Datura* plants which had been subjected to low temperatures at the time of fertilization. Since this initial work by Blakeslee, et al (1), haploids have been found from time to time and described in about 20 genera of plants.

Origin

There has been much speculation as to the origin of haploids. According to Cooper (3), Harland (8), and Muntzing (19), poly-embryony or twin-seedlings appeared to be one of the more frequent sources. Harland (8), in a germination experiment with Sea Island cotton, observed 20 seeds containing two embryos, one of which was usually much larger than the other. From such material 16 pairs of plants were raised to maturity; of these 14 pairs consisted of one haploid and one diploid, and two pairs each of identical diploids. Cooper (3) observed a number of haploid-diploid twins in Lillium and Nicotiana. He suggested that this phenomenon occurs when a synergid is stimulated to divide and that many of the haploid plants discovered arise from synergids.

Crosses between distantly related species have in a few instances given rise to haploids. Gaines and Aase (7) obtained a haploid with 21 chromosomes by pollinating a winter wheat, Triticum compactum humboldii with the pollen of Aegilops cylindrica. In this cross, usually only shrivelled seeds resulted but the single plump kernel which produced the haploid was much larger than seed of the Triticum parent.

Gaines and Aase (7), suggested that both generative nuclei of the Aegilops pollen fused with the maternal endosperm nuclei to produce a tetraploid endosperm, while the egg developed parthenogenically. Smith (23), discussing wide crosses such as the one described by Gaines and Aase (7), postulated that, "possibly the endosperm with two maternal and one paternal set of chromosomes is capable of developing; but the embryo with one maternal and one paternal set dies, in which event a synergid or an already partially developed maternal embryo, develops into a haploid embryo".

Various artificial methods have been employed in producing haploids. Cold treatment was used by Blakeslee, et al (1), in *Datura*. It was postulated that the low temperature caused the parthenogenic development of the egg cells. Ivanov (9), Katayama (10), Kihara and Katayama (12), Kihara and Yamashita (13), Kostoff (14), and Smith (23), have attempted the production of haploids by the use of X-rays. Ivanov (9) believed that X-rayed pollen produced a significant increase in the number of haploids in Nicotiana rustica var. humilis. His work was corroborated by Kostoff (14) on similar material. Katayama (10), Kihara and Katayama (12), and Kihara and Yamashita (13) obtained significant increases in the incidence of haploidy in einkorn from the use of X-rayed pollen. Smith (23), however, found no increase in

einkorn haploids from X-rayed pollen. Actually fewer haploids were obtained in one experiment when such treated pollen was used.

Delayed pollination has been used as a means of increasing the occurrence of haploid plants. Kihara (11) reported that if pollination of einkorn wheat was delayed beyond the normal time after emasculation, the frequency of haploids was correlated (to a point) with the length of the delay. He also observed three pairs of twin seedlings which presumably arose from delayed pollination.

Lindstrom (17) described a tomato mutant which had 12 chromosomes. It appeared in the F_2 population of a varietal cross and showed complete fertility. Lindstrom (17), regarded this haploid as "spontaneous", as no apparent stimulus was responsible for its production.

Meiotic Behaviour

Meiotic behaviour of haploids has been studied by a number of workers; (2), (4), (5), (6), (7), (9), (15), (16), (17), (18), (20), (23), and (25). To date there is no

... ..
... ..
... ..

... ..
... ..
... ..
... ..
... ..
... ..
... ..

... ..
... ..
... ..
... ..
... ..
... ..

... ..

... ..
... ..
... ..
... ..
... ..
... ..

general agreement since each species has apparently somewhat different meiotic characteristics.

Random distribution of the chromosomes to the poles was observed by Bleier (2), Davis and Kulkarni (4), Gaines and Aase (7), Ivanov (9), and Tometorp (25). Gaines and Aase (7) found that pairing occurred only rarely in their 21 chromosome wheat plant. The chromosomes apparently were distributed to the poles at random. Davis and Kulkarni (4) in their studies of an Oenothera francisca haploid, found that none of the seven chromosomes paired. Sterile pollen resulted from the irregular distribution of the chromosomes in the first division; (6+1), (5+2), and (4+3). They postulated that functional pollen grains resulted when a unipolar spindle had formed and all seven chromosomes were drawn to one pole. Ivanov (9) studied a haploid of Nicotiana rustica and observed that chromosome pairing was almost entirely absent. At metaphase I the chromosomes lay scattered over the entire spindle.

In some Matthiola hybrids, Lesley and Frost (15) obtained a haploid which had seven univalents plus a fragment. No pairing was observed. In some cases, random distribution of the unpaired chromosomes occurred, but frequently the chromosomes split and separated. In these cases the first division was evidently omitted and pollen dyads resulted.

Emerson (5) studied a haploid of Oe. francisca but did not observe the same behaviour as that described by Davis and Kulkarni (4). He observed two chromosomes frequently attached end-to-end. The behaviour of these two chromosomes apparently differed from that of the other five univalents. In early anaphase I, the members of the pair usually passed to opposite poles while the other five univalent chromosomes were distributed at random. Counts of the chromosomes in the two daughter nuclei showed the following distributions; 7 cases (7+0), 25 cases (6+1), 18 cases (5+2) and 15 cases (4+3). Emerson concluded that the high frequency of the (6+1) distribution was the result of the frequent movement to one pole of all five univalents, with one member of the dividing pair.

Recently, instances of chromosome pairing in haploids have been reported, (2, 6, 14, 16, 18, 20, 23, and 26). These workers however, do not agree in their definitions of chromosome pairing nor in their interpretations of its significance.

Levan (16), who studied meiosis in haploid rye, occasionally found two bivalents and one trivalent at diakinesis. The presence of numerous chiasmata apparently held paired chromosomes together even to metaphase I. He did not believe that chiasmata could form between non-homologous chromosome portions and therefore concluded that pairing was between chromosomes that

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

were actually partially homologous. Kostoff (14) was of the opinion that in haploid einkorn, pairing had nothing to do with chromosome homology but occurred between inert heterochromatic regions.

Yamasaki (26), in a Triticum vulgare haploid, observed the following pairing frequencies at metaphase I; 21' - 47.7%, 19'1" - 37.1%, 17'2" - 12.4%, 15'3" - 2.4%, and 13'4" - 0.4%. An occasional trivalent was also noted and taken as an indication that a relationship existed among the three genomes.

Smith (23) found a low percentage of open bivalents with observable chiasmata in einkorn haploids. More frequently, however, he found chromosomes arranged at first metaphase so that they might be considered as two secondary pairs. He suggested that this might be taken as evidence that five is the basic number of chromosomes in wheat. Smith (23) considered two chromosomes as secondary pairs if they were not more than the diameter of a chromosome apart at any one point. Only a small number were paired throughout their entire length.

In Arctic Grass haploids, Flovic (6) concluded that real homology involves the exchange of chromatin and secondary pairing may occur as a result of phylogenetic affinity of chromosomes without intimate homology. He pointed out that secondary pairing may be caused by mutations or structural changes of the chromosomes, but is probably evidence of a lower basic number.

Nordenskiöld (20), in a haploid rye plant, observed a bivalent in 3.2% of the pollen mother cells which were cytologically examined. This bivalent was always rod-shaped and appeared to be joined by a chiasma. Bleier (2) suggested that bivalent formation is not constant in haploids and that factors other than strict homology (e.g. the chance sticking together of chromosomes from prophase on), have a greater influence on pairing.

Morinaga and Fukushima (18) studied meiosis in a haploid rice plant and found that pollen mother cells contained one or two bivalents. Those at the equator were considered as true bivalents. Tomtorp (25) found that a bivalent was formed in 5% of the pollen mother cells of a Hordeum distichum haploid. He did not believe that bivalents are a secondary phenomenon but that they may arise through relic similarity of chromosomes.

The appearance of chromosome bridges at first or second anaphase of haploids has been reported by a number of investigators. Levan (16) frequently observed chromatin bridges without fragments in a rye haploid. He suggested that their presence could result from chromatid reunion. Katayama (10) and Smith (23), working independently, observed bridges in einkorn haploids which had fragments occasionally associated with them. Smith (23) found a high correlation between the number of bivalents and the number of bridges. He pointed out that these

bridges probably resulted from crossing-over between non-homologous chromosomes and, "if this interpretation is applicable to the bridges in haploid einkorn, it would suggest that crossing-over had occurred between most of the bivalents. it thus follows that true chiasmata were observed".

To date, Sears (21) has made the most practical use of a haploid plant. He obtained two haploids from the cross T. vulgare pollinated by Secale cereale. When pollinated with normal wheat pollen, one plant produced 14 seeds and from these, 13 plants matured. These plants and their progeny were examined cytologically, and Sears (21) observed 11 plants with an aberrant chromosomal complement. Univalents, trivalents and rings of four were found in conjunction with normal bivalents. Since Sears (21) was unable to examine the original haploid cytologically, he could only postulate the probable cause of the aberrations (monosomics, trisomics, etc.), observed in the F_1 and subsequent generations.

MATERIALS AND METHODS

In July, 1949, a haploid of Triticum vulgare was observed in an F_3 experimental plot of the inter-varietal cross, Lemhi x (Bunyip x Dicklow). It was not noticed until flowering time when the spreading of the glumes, characteristic of sterility, drew attention to it. It is not believed that the haploid arose through twin embryos as only one plant was present. Three slightly dwarfed and spindly tillers were present; two in flower and the third young enough to allow cytological examination of the pollen mother cells. This spike was removed and fixed for cytological studies.

Of the two remaining spikes, one was allowed to self-pollinate while the second was emasculated and pollinated with normal Lemhi pollen. Emasculation was probably unnecessary since the selfed spike proved to be completely sterile. Heavy rains in mid-July, stimulated the formation of two more tillers which were also emasculated and pollinated with Lemhi pollen. The three cross-pollinated spikes produced eight seeds.

The eight seeds from the haploid were sown in the greenhouse in September, 1949, and seven F_1 progenies were obtained. Unfortunately greenhouse conditions were poor during the germination and tillering periods, and only one spike per plant was

produced. Consequently the chromosome complement could not be obtained without eliminating subsequent generations. Of the seven F_1 plants, only one proved to be completely sterile and it has been suggested that high temperatures in the greenhouse during the flowering period may have been partially responsible.

Ten seeds of each of 18 sister plants to the haploid were also seeded in the greenhouse in September, 1949, in progeny lines. Six plants in each line were studied cytologically so that the chromosome makeup of the immediate parent of the haploid could be determined.

The F_2 and F_3 generations from the haploid were planted in progeny lines in the greenhouse in February and June, 1950, respectively. Only a small percentage of these plants failed to survive. Each plant produced several tillers and cytological studies of microsporogenesis were made on all plants.

In this study, all plant material to be examined cytologically was fixed in 6:3:1 Carnoy's fluid A, and the acetocarmine smear method as outlined by Smith (24) was used for slide preparations. Chromosome counts were made at a magnification of 600 diameters and photomicrographs were taken at a magnification of either 720 or 900 diameters. The tertiary butyl alcohol method, as outlined by Smith (24), was used for making slides permanent.

... ..
... ..
... ..
... ..
... ..

... ..
... ..
... ..
... ..
... ..

... ..
... ..
... ..
... ..
... ..

... ..
... ..
... ..
... ..
... ..

... ..
... ..
... ..
... ..
... ..

EXPERIMENTAL RESULTS AND INTERPRETATIONS

CYTOLOGICAL DATA FROM THE HAPLOID PLANT

Distribution of Univalent Chromosomes

In general the chromosomes were distributed throughout the entire cell at metaphase I, as can be seen in Figs. 1 and 2. This observation is in agreement with the work of Bleier (2), Gaines and Aase (7), Ivanov (9), and other workers. Apparently the univalents did not line up on the plate prior to the first anaphase as do normal bivalents. Consequently these two phases were not clearly distinguishable.

A total of 274 pollen mother cells were counted in which the univalents had migrated to the poles, Figs. 3, 4, and 5. The anaphase I distributions are presented in Table 1.

Table 1

Distribution of Univalents

Distribution	20-1	19-2	18-3	17-4	16-5	15-6	14-7	13-8	12-9	11-10	Total
No. of cells	2	2	7	2	15	8	31	19	90	98	274
% of total	0.7	0.7	2.6	0.7	5.5	2.9	11.3	6.9	32.9	35.8	100%
Normal (%) distribution	0.002	0.02	0.1	0.6	1.9	5.2	11.1	19.4	28.0	33.6	100%
χ^2	245.000**	23.120**	62.500**	0.017	6.821**	1.017	--	8.054**	0.858	0.144	347.531

** Highly significant.

Table I

Summary of results

Run	Time	Temp	Pressure	Flow	Conc	Yield	Notes
1	10.0	100	1.0	1.0	1.0	1.0	Initial
2	10.0	100	1.0	1.0	1.0	1.0	Initial
3	10.0	100	1.0	1.0	1.0	1.0	Initial
4	10.0	100	1.0	1.0	1.0	1.0	Initial
5	10.0	100	1.0	1.0	1.0	1.0	Initial
6	10.0	100	1.0	1.0	1.0	1.0	Initial
7	10.0	100	1.0	1.0	1.0	1.0	Initial
8	10.0	100	1.0	1.0	1.0	1.0	Initial
9	10.0	100	1.0	1.0	1.0	1.0	Initial
10	10.0	100	1.0	1.0	1.0	1.0	Initial

Continued on next page

Cells with near equal distribution occurred with the highest frequencies. Actually the 11-10, 12-9, 14-7, 15-6, and 17-4 distribution values were fairly close to the expected values for random distribution. Some of the other distributions, however, deviated very markedly from the expected so that for the entire study a highly significant χ^2 was obtained. It was quite apparent, therefore, that randomness of distribution to the poles was restricted. It is interesting that a 20-1 distribution was observed in two of the cells studied, Fig. 5. Sears (21), in one of his papers, postulated that such a gamete could be one possible source of monosomic wheat plants.

Chromosome Associations

Paired Chromosomes

A large number of cells with paired chromosomes were observed. True pairing was assumed if a chiasma was actually observed, Fig. 6, or if the tapered end appearance of a disjoining pair indicated the previous existence of a chiasma. So-called secondary pairing was also frequently noted, consisting merely of an end-to-end association of univalent chromosomes with no sign of a real chiasma, Fig. 2. The data on bivalent and other association frequencies are summarized in Table 2.

Table 2

Chromosome Associations

Association	21'	19'1"	17'2"	15'3"	21"	42'	21 dyads	Chain of			Total
								3	4	5	
No. of cells	93	117	115	23	1	7	5	11	5	1	378
% of total	24.6	31.0	30.4	6.1	0.3	1.8	1.3	2.9	1.3	0.3	100

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100	101	102	103	104	105	106	107	108	109	110	111	112	113	114	115	116	117	118	119	120	121	122	123	124	125	126	127	128	129	130	131	132	133	134	135	136	137	138	139	140	141	142	143	144	145	146	147	148	149	150	151	152	153	154	155	156	157	158	159	160	161	162	163	164	165	166	167	168	169	170	171	172	173	174	175	176	177	178	179	180	181	182	183	184	185	186	187	188	189	190	191	192	193	194	195	196	197	198	199	200	201	202	203	204	205	206	207	208	209	210	211	212	213	214	215	216	217	218	219	220	221	222	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244	245	246	247	248	249	250	251	252	253	254	255	256	257	258	259	260	261	262	263	264	265	266	267	268	269	270	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	294	295	296	297	298	299	300	301	302	303	304	305	306	307	308	309	310	311	312	313	314	315	316	317	318	319	320	321	322	323	324	325	326	327	328	329	330	331	332	333	334	335	336	337	338	339	340	341	342	343	344	345	346	347	348	349	350	351	352	353	354	355	356	357	358	359	360	361	362	363	364	365	366	367	368	369	370	371	372	373	374	375	376	377	378	379	380	381	382	383	384	385	386	387	388	389	390	391	392	393	394	395	396	397	398	399	400	401	402	403	404	405	406	407	408	409	410	411	412	413	414	415	416	417	418	419	420	421	422	423	424	425	426	427	428	429	430	431	432	433	434	435	436	437	438	439	440	441	442	443	444	445	446	447	448	449	450	451	452	453	454	455	456	457	458	459	460	461	462	463	464	465	466	467	468	469	470	471	472	473	474	475	476	477	478	479	480	481	482	483	484	485	486	487	488	489	490	491	492	493	494	495	496	497	498	499	500	501	502	503	504	505	506	507	508	509	510	511	512	513	514	515	516	517	518	519	520	521	522	523	524	525	526	527	528	529	530	531	532	533	534	535	536	537	538	539	540	541	542	543	544	545	546	547	548	549	550	551	552	553	554	555	556	557	558	559	560	561	562	563	564	565	566	567	568	569	570	571	572	573	574	575	576	577	578	579	580	581	582	583	584	585	586	587	588	589	590	591	592	593	594	595	596	597	598	599	600	601	602	603	604	605	606	607	608	609	610	611	612	613	614	615	616	617	618	619	620	621	622	623	624	625	626	627	628	629	630	631	632	633	634	635	636	637	638	639	640	641	642	643	644	645	646	647	648	649	650	651	652	653	654	655	656	657	658	659	660	661	662	663	664	665	666	667	668	669	670	671	672	673	674	675	676	677	678	679	680	681	682	683	684	685	686	687	688	689	690	691	692	693	694	695	696	697	698	699	700	701	702	703	704	705	706	707	708	709	710	711	712	713	714	715	716	717	718	719	720	721	722	723	724	725	726	727	728	729	730	731	732	733	734	735	736	737	738	739	740	741	742	743	744	745	746	747	748	749	750	751	752	753	754	755	756	757	758	759	760	761	762	763	764	765	766	767	768	769	770	771	772	773	774	775	776	777	778	779	780	781	782	783	784	785	786	787	788	789	790	791	792	793	794	795	796	797	798	799	800	801	802	803	804	805	806	807	808	809	810	811	812	813	814	815	816	817	818	819	820	821	822	823	824	825	826	827	828	829	830	831	832	833	834	835	836	837	838	839	840	841	842	843	844	845	846	847	848	849	850	851	852	853	854	855	856	857	858	859	860	861	862	863	864	865	866	867	868	869	870	871	872	873	874	875	876	877	878	879	880	881	882	883	884	885	886	887	888	889	890	891	892	893	894	895	896	897	898	899	900	901	902	903	904	905	906	907	908	909	910	911	912	913	914	915	916	917	918	919	920	921	922	923	924	925	926	927	928	929	930	931	932	933	934	935	936	937	938	939	940	941	942	943	944	945	946	947	948	949	950	951	952	953	954	955	956	957	958	959	960	961	962	963	964	965	966	967	968	969	970	971	972	973	974	975	976	977	978	979	980	981	982	983	984	985	986	987	988	989	990	991	992	993	994	995	996	997	998	999	1000	1001	1002	1003	1004	1005	1006	1007	1008	1009	1010	1011	1012	1013	1014	1015	1016	1017	1018	1019	1020	1021	1022	1023	1024	1025	1026	1027	1028	1029	1030	1031	1032	1033	1034	1035	1036	1037	1038	1039	1040	1041	1042	1043	1044	1045	1046	1047	1048	1049	1050	1051	1052	1053	1054	1055	1056	1057	1058	1059	1060	1061	1062	1063	1064	1065	1066	1067	1068	1069	1070	1071	1072	1073	1074	1075	1076	1077	1078	1079	1080	1081	1082	1083	1084	1085	1086	1087	1088	1089	1090	1091	1092	1093	1094	1095	1096	1097	1098	1099	1100	1101	1102	1103	1104	1105	1106	1107	1108	1109	1110	1111	1112	1113	1114	1115	1116	1117	1118	1119	1120	1121	1122	1123	1124	1125	1126	1127	1128	1129	1130	1131	1132	1133	1134	1135	1136	1137	1138	1139	1140	1141	1142	1143	1144	1145	1146	1147	1148	1149	1150	1151	1152	1153	1154	1155	1156	1157	1158	1159	1160	1161	1162	1163	1164	1165	1166	1167	1168	1169	1170	1171	1172	1173	1174	1175	1176	1177	1178	1179	1180	1181	1182	1183	1184	1185	1186	1187	1188	1189	1190	1191	1192	1193	1194	1195	1196	1197	1198	1199	1200	1201	1202	1203	1204	1205	1206	1207	1208	1209	1210	1211	1212	1213	1214	1215	1216	1217	1218	1219	1220	1221	1222	1223	1224	1225	1226	1227	1228	1229	1230	1231	1232	1233	1234	1235	1236	1237	1238	1239	1240	1241	1242	1243	1244	1245	1246	1247	1248	1249	1250	1251	1252	1253	1254	1255	1256	1257	1258	1259	1260	1261	1262	1263	1264	1265	1266	1267	1268	1269	1270	1271	1272	1273	1274	1275	1276	1277	1278	1279	1280	1281	1282	1283	1284	1285	1286	1287	1288	1289	1290	1291	1292	1293	1294	1295	1296	1297	1298	1299	1300	1301	1302	1303	1304	1305	1306	1307	1308	1309	1310	1311	1312	1313	1314	1315	1316	1317	1318	1319	1320	1321	1322	1323	1324	1325	1326	1327	1328	1329	1330	1331	1332	1333	1334	1335	1336	1337	1338	1339	1340	1341	1342	1343	1344	1345	1346	1347	1348	1349	1350	1351	1352	1353	1354	1355	1356	1357	1358	1359	1360	1361	1362	1363	1364	1365	1366	1367	1368	1369	1370	1371	1372	1373	1374	1375	1376	1377	1378	1379	1380	1381	1382	1383	1384	1385	1386	1387	1388	1389	1390	1391	1392	1393	1394	1395	1396	1397	1398	1399	1400	1401	1402	1403	1404	1405	1406	1407	1408	1409	1410	1411	1412	1413	1414	1415	1416	1417	1418	1419	1420	1421	1422	1423	1424	1425	1426	1427	1428	1429	1430	1431	1432	1433	1434	1435	1436	1437	1438	1439	1440	1441	1442	1443	1444	1445	1446	1447	1448	1449	1450	1451	1452	1453	1454	1455	1456	1457	1458	1459	1460	1461	1462	1463	1464	1465	1466	1467	1468	1469	1470	1471	1472	1473	1474	1475	1476	1477	1478	1479	1480	1481	1482	1483	1484	1485	1486	1487	1488	1489	1490	1491	1492	1493	1494	1495	14
---	---	---	---	---	---	---	---	---	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	----

Of a total of 378 pollen mother cells counted at metaphase I, the greatest proportion (31.0% and 30.4%) had a constitution of 19'1" and 17'2", respectively. A constitution of 15'3" was observed in 6.1% of the cells, but no cell was observed that had more than three pairs. These frequencies are quite different from those reported by Yamasaki (26).

The cause of bivalent pairing in the haploid could have two interpretations. Had the parent of the haploid been heterozygous for a number of reciprocal translocations, bivalent formation in the haploid could be expected. On the other hand, pairing would also occur if there were homology between various chromosomes in the haploid. The first hypothesis can be ruled out since no ring or chain associations were found in the progenies of the haploid sister plants. All 108 sister progenies proved to be completely normal cytologically. The pairing observed must, therefore, be taken as evidence of true homology between certain of the haploid chromosomes resulting in the occurrence of reciprocal translocations between these chromosomes. If the presence of bivalents is a true measure of the number of reciprocal translocations that have taken place, it can be assumed that up to three such exchanges have occurred in some cells. It is not suspected that this homology was between chromosomes of the same genome, but rather between chromosomes of different genomes. Smith (23)

observed a low frequency of bivalents in einkorn haploids which have only one genome, and also noted a high correlation between the number of bivalents and the number of bridges at anaphase I or II. The presence of bridges would suggest that crossing-over must have taken place between completely non-homologous chromosomes. Since no bridges were observed in this study, and since normally disjoining bivalents were frequently noted, it must be assumed that pairing occurred between partially homologous chromosomes. If the presence of bridges is a true indication of intra-genome pairing, then their complete absence could indicate equally well, inter-genome pairing.

While mostly open bivalents were observed, a closed pair with a chiasma on both sides of the centromere was observed in three cells, Figs. 7 and 8. So far as is known, this has never been observed in any of the haploids that have been studied. Such complete pairing certainly suggests even greater homology between the various chromosomes than has usually been assumed to exist.

Secondary pairing, while frequently observed, is not likely an indication of true homology. Possibly as Bleier (2) has suggested, stickiness of the chromosomes from prophase on, could account for its occurrence. Gene specificity or chemical attraction between chromosomes may also be a suitable explanation for the formation of secondary pairs.

As mentioned previously, the random distribution of the chromosomes at anaphase I, was obviously restricted. Bivalent formation and secondary associations were likely responsible for disturbing the distributions. Regular disjoining of bivalents at anaphase I would alone be restrictive, but also the secondary associations remained united going to the same pole, thereby further affecting normal distribution. It is, therefore, concluded that distribution was random when 21' were present, but that when pairing did occur, distribution was random for only those chromosomes not involved in the pairing.

Chains

Associations or chains of three to five univalents were observed in 4.5% of the cells, Fig. 9. Apparently a continuation of secondary pairing was responsible for this type of chain formation. These associations would likely go to the same pole, thereby further affecting the randomness of chromosome distribution.

Cells with Reduplicated Chromosomes

One cell was observed that had 21" at metaphase I, Fig. 10. So far as could be determined, such a phenomenon has never

been observed in any of the haploids that have been studied. Pairing was normal and very likely normal gametes would result from such a cell. Seven cells were observed that had 42 mostly univalent chromosomes, Fig. 11. It is postulated that the first type of reduplicated cell could have arisen from failure of cell wall formation during the first mitotic division of a sporogenous cell. When the nuclei so formed, prepared for the second mitotic division, a common spindle was formed and the resulting daughter cells (microsporocytes) would have 42 chromosomes each. Meiosis would then proceed normally. In the second case, cells with 42 univalents could have arisen by cell wall failure in the second mitotic division (immediately preceding microsporocyte formation). Since the nuclear membranes in these binucleate cells would remain intact until diakinesis, there would be no opportunity for the 42 chromosomes to pair.

Only five cells contained dyads, Figs. 12 and 13, which were probably formed from a first division split of the univalents, remaining joined only at the centromere. In Fig. 12 it can be seen that some of the univalents have undergone complete splitting and the monads and dyads are being randomly distributed to the poles. It is probable that the dyads would undergo a second division but the gametes thus formed would be deficient for several whole chromosomes rendering them non-functional.

Plate I

Plate II

Legend to Plates I and II

Microsporogenesis in the haploid.

- Fig. 1. Metaphase I. Distribution of the 21 unpaired chromosomes throughout the entire cell.
- Fig. 2. Metaphase I. Distribution of univalents with two bivalents and secondary pairing.
- Figs. 3, 4 and 5. Anaphase distributions; 12-9, 14-7 and 20-1, respectively.
- Fig. 6. 19'1". Open bivalent with strong chiasma.
- Fig. 7 and 8. Closed bivalent.
- Fig. 9. Chain associations.
- Fig. 10. Cell with 21 normal pairs; a second cell with 21 univalents.
- Fig. 11. Cell with 42 mostly univalent chromosomes.
- Fig. 12. Monads and dyads.
- Fig. 13. A 14-7 distribution of dyads.

Magnification:

Figs. 2, 5 and 9 at 900 diameters; all others at 720 diameters.

CYTOLOGICAL DATA FROM THE F₂ AND F₃ GENERATIONS

Since no cytological data are available from the F₁ plants, all information on the transmission of aberrations had to be obtained from the analyses of the F₂ and F₃ populations. The chromosomal constitutions of the F₂ and F₃ generations which were cytologically examined, are presented in Table 3.

Of the 60 F₂ plants examined, 26 or 43% had a normal chromosomal complement of 21," Fig. 15. Of 231 F₃ plants examined, 153 or 66% proved to have a normal constitution. Consequently, these data indicate that plants with chromosomal aberrations became progressively fewer with succeeding generations. As would be expected, all normal F₂ plants had normal progeny. Aberrant plants, however, transmitted aberrations to some of their offspring. Furthermore, the aberrations varied markedly in different sister progenies, being in some cases more extreme than in the parent. Apparently the degree of abnormalness of the parent plant determined the relative frequency of normal and abnormal progeny.

The most common type of aberrations noted were chains of three or four chromosomes, rings of four and univalents. It is assumed that these aberrations were largely the result of reciprocal translocations that occurred in the haploid. Moreover, the high

THEORETICAL BASIS OF THE LOGIC OF DISPOSITIONS

Since no physical law can be derived from the

logical, all information on the laws of nature must be obtained from the sciences of the world of experience. The logical foundations of the logic of dispositions must therefore be logically examined, and presented in Table 1.

Of the two basic elements, the α and β are

theoretical elements of the α and β elements.

It is not possible to have a logical foundation, theoretically,

large scale logical laws and also logical elements be-

cause theoretically, laws and logical elements are

abstract, the logical α and β elements, logical laws,

however, theoretical elements of the world of experience, logical

laws, the theoretical world of experience, logical laws,

being in some cases not even in the world of experience.

The laws of logic are of two kinds, logical laws and

logical laws of logic and logical laws.

The laws of logic are of two kinds, logical laws and

logical laws of logic, laws of logic and logical laws.

Since the laws of logic are of two kinds, logical laws and

logical laws of logic, laws of logic and logical laws.

frequency of F_2 and F_3 plants containing up to three associations of three or four chromosomes, further substantiated the hypothesis that up to three reciprocal translocations occurred in the haploid. Disjunction from translocation configurations likely caused the formation of deficient and duplicated gametes which would result in plants having chains and also unpaired chromosomes, especially if the interchanged portion were small. A diagrammatic representation of the probable origin of aberrant gametes from one reciprocal translocation, is presented in Fig. 14. Random union of any two of such gametes, gives numerous possibilities for various types of chromosomal aberrations. If as many as three reciprocal translocations were involved, the resulting progeny could have all the cytological abnormalities observed in this study, Figs. 16 to 25. It is probable that most male gametes would be normal in chromosome number, but occasionally ones with an aneuploid number might be functional. Female gametes, on the other hand, would be expected to transmit aberrations with a fairly high frequency.

The total number of chromosomes in different fertile plants ranged from 39, Fig. 16, to 47, Fig. 17. It is surprising that fertility and vigor was high even in some of the most aberrant plants. The plant with 39 chromosomes appeared quite normal phenotypically. This is highly interesting since Sears (21) found that nullisomics were usually quite weak and sterile. Probably the 39

chromosome plant had certain interchanged chromosomes so that there was actually no complete absence of essential genetic material.

The meiotic behaviour of one F_3 plant should be especially noted, Fig. 26. At metaphase I, it proved to be highly asynaptic similar to nulli-3 described by Sears (22). It is possible that this plant was nullisomic for chromosome three, and the origin of such a plant also can be explained from the union of certain gametes, each deficient for the same chromosome, Fig. 14.

 

F₂ Deficiencies and duplications associated with univalents, bivalents, trivalents and quadrivalents would occur. These aberrations would be transmitted to subsequent generations.

Fig. 14 Diagrammatic representation of the probable origin

Table 3

Chromosomal Constitutions of F₂ and F₃ Generations

F ₂ chromosome constitution	Total F ₂ plants analysed	F ₃ chromosome constitution	Total F ₃ plants analysed
21"	26	21"	108
21" or 19"C""	9	21"	24
		21" or 19"C""	8
		21" or 19"C"" or 19"C'"1'	1
21" or 19"C'"1'	3	21"	6
		20"1'	2
		20"1' or 19"C'"	1
		21" or 19"C"" or 19"C'"1'	3
		18"C""1'	1
21" or 19"C'"1' or 18"2C'"	1	21"	3
		21" or 19"R""	2
		21" or 17"R""C'"1' or 19"C""	1
19"C"" or 19"C'"1'	1	19"C'"1' + frag.	1
		19"C""1' + frag.	1
18"C""2'	1	19"C""	1
		19"C'"1'	1
		20"2' + iso.	1
		17"C""C'"1' or 19"C""	1
19"C'" or 17"2C'"1'	1	-	0
16"C""C'"3'	1	19"C'"1'	3
		18"C""2'	1
		17"C'"4'	1
		16"3C'"3'	1

C. 1900

Continuation of the 1900-1901 season

Date of arrival	Name of vessel	Date of departure	Name of vessel
1900		1901	
1	1900-1901 season	1	1900-1901 season
2	1900-1901 season	2	1900-1901 season
3	1900-1901 season	3	1900-1901 season
4	1900-1901 season	4	1900-1901 season
5	1900-1901 season	5	1900-1901 season
6	1900-1901 season	6	1900-1901 season
7	1900-1901 season	7	1900-1901 season
8	1900-1901 season	8	1900-1901 season
9	1900-1901 season	9	1900-1901 season
10	1900-1901 season	10	1900-1901 season
11	1900-1901 season	11	1900-1901 season
12	1900-1901 season	12	1900-1901 season
13	1900-1901 season	13	1900-1901 season
14	1900-1901 season	14	1900-1901 season
15	1900-1901 season	15	1900-1901 season
16	1900-1901 season	16	1900-1901 season
17	1900-1901 season	17	1900-1901 season
18	1900-1901 season	18	1900-1901 season
19	1900-1901 season	19	1900-1901 season
20	1900-1901 season	20	1900-1901 season
21	1900-1901 season	21	1900-1901 season
22	1900-1901 season	22	1900-1901 season
23	1900-1901 season	23	1900-1901 season
24	1900-1901 season	24	1900-1901 season
25	1900-1901 season	25	1900-1901 season
26	1900-1901 season	26	1900-1901 season
27	1900-1901 season	27	1900-1901 season
28	1900-1901 season	28	1900-1901 season
29	1900-1901 season	29	1900-1901 season
30	1900-1901 season	30	1900-1901 season
31	1900-1901 season	31	1900-1901 season

Table 3 (Continued)

F ₂ chromosome constitution	Total F ₂ plants analysed	F ₃ chromosome constitution	Total F ₃ plants analysed
20"1'	4	21" 20"1' 19"2'	6 9 1
20"1' or 19"3' or 18"C""1'	1	20"1' or 19"C'" or 19"3' 19"2'	1 2
19"3'	1	21" 20"1' 19"2'	1 1 1
19"2'	2	20"1' 19"1' 19"2'	3 3 1
17"C'"3'	1	20"1' or 19"C'" 20"1' 19"3'	1 2 1
15"C'"2'2 + iso.	1	-	0
21"1' or 19"R""1'	2	21"1' or 20"C'" 20"1' or 19"C'" 19"4' or 18"C""2' 18"C""1' or 17"C""3' or 20"1' 18"C'"2' 21"	1 2 1 2 1 1
21"2' or 20"C"" or 20"4'	1	21" 21"2' or 20R"" 20"R""C'" or 18"2r""C'" 19"C""1' or 18"C""C'" or 15"C""2C'"3' 19"C"" or 19"C'"1'	3 1 1 1 1 1
19"C'"3' or 18"2C'"2'	1	-	0
21"1' or 20"C'" or 18"C""C'"	3	21"1' or 20"C'" 19"C""1' 21" + frag. 21"	7 2 1 1
Total	60		231

(continued) Exhibit

Case No.	Exhibit No.	Description of Exhibit	Page No.	Exhibit No.	Description of Exhibit	Page No.
1	1	1971-1972	1	1	1971-1972	1
2	2	1973-1974	2	2	1973-1974	2
3	3	1975-1976	3	3	1975-1976	3
4	4	1977-1978	4	4	1977-1978	4
5	5	1979-1980	5	5	1979-1980	5
6	6	1981-1982	6	6	1981-1982	6
7	7	1983-1984	7	7	1983-1984	7
8	8	1985-1986	8	8	1985-1986	8
9	9	1987-1988	9	9	1987-1988	9
10	10	1989-1990	10	10	1989-1990	10
11	11	1991-1992	11	11	1991-1992	11
12	12	1993-1994	12	12	1993-1994	12
13	13	1995-1996	13	13	1995-1996	13
14	14	1997-1998	14	14	1997-1998	14
15	15	1999-2000	15	15	1999-2000	15
16	16	2001-2002	16	16	2001-2002	16
17	17	2003-2004	17	17	2003-2004	17
18	18	2005-2006	18	18	2005-2006	18
19	19	2007-2008	19	19	2007-2008	19
20	20	2009-2010	20	20	2009-2010	20
21	21	2011-2012	21	21	2011-2012	21
22	22	2013-2014	22	22	2013-2014	22
23	23	2015-2016	23	23	2015-2016	23
24	24	2017-2018	24	24	2017-2018	24
25	25	2019-2020	25	25	2019-2020	25
26	26	2021-2022	26	26	2021-2022	26
27	27	2023-2024	27	27	2023-2024	27
28	28	2025-2026	28	28	2025-2026	28
29	29	2027-2028	29	29	2027-2028	29
30	30	2029-2030	30	30	2029-2030	30

Plate IV

Legend of Plates III and IV

Microsporocytes of F_2 and F_3 Plants.

Fig. 15. Cell with 21".

Fig. 16. F_3 cell with 19"1'.

Fig. 17. F_3 cell with 18"2R""C'".

Fig. 18. and Fig. 19. F_2 and F_3 cells with 19"C"".

Fig. 20. F_2 cell with 19"R""1'.

Fig. 21. F_2 cell with 17"2C'"1'.

Fig. 22. F_2 cell with 19"2'.

Fig. 23. F_3 cell with 19"C'"1'.

Fig. 24. F_3 cell with 19"C'".

Fig. 25. F_3 cell with 16"3C'"3'.

Fig. 26. F_3 cell. Highly asynaptic.

All figures at a magnification of 720 diameters.

CONCLUSIONS

It is concluded that the pairing observed in the haploid occurred between chromosomes because of homology rather than translocation heterozygosity. This pairing, however, resulted in reciprocal translocations which gave rise to plants in succeeding generations, with various deficiency and duplication constitutions. These results are interpreted as evidence that there is considerable homology between chromosomes of the various genomes in hexaploid wheat. It is highly probable, of course, that the actual gene homology and duplication present in common wheat haploids, is much greater than the actual pairing indicated. Pairing would likely occur only when considerable portions of chromosomes were identical. In the evolution of this species it is likely that many gene and chromosome rearrangements have occurred so that while there is genetic homology, there is structural homology for meiotic pairing in only relatively few haploid chromosomes.

It is felt that secondary pairing is no indication of true chromosomal homology, but that it restricted random distribution of the chromosomes to the poles. The same process causing secondary pairing was likely also responsible for the formation of chains.

Cells with reduplicated chromosomes were found and it is suggested that the process whereby these were formed could account for the formation of doubled gametes and seed setting in normally sterile hybrids. In the case of the cell with 21", it is assumed that cell wall formation failed during the first mitotic division of the sporogenous cell, while in the case of cells with 42', cell wall formation failed in the second mitotic division.

ACKNOWLEDGEMENTS

The author wishes to express his thanks and appreciation to Dr. John Unrau, under whose guidance the study was conducted and the manuscript prepared, and to the National Research Council of Canada, for financial assistance in the form of a bursary.

REFERENCES

1. BLAKESLEE, A. F., BELLING, J., FARNHAM, M. E. and BERGNER, A. D. A haploid mutant in the Jimson weed, Datura stramonium. Science 40:646-647. 1922.
2. BLEIER, H. Die Meiosis von Haploidiplonten. Genetica 15: 129-177. 1933.
3. COOPER, D. C. Haploid-diploid twin embryos in Lillium and Nicotiana. Am. J. Botany 30:408-413. 1943.
4. DAVIS, B. M. and KULKARNI, C. G. The cytology and genetics of a haploid sport from Oenothera franciscana. Genetics 15:55-80. 1930.
5. EMERSON, S. H. The reduction division in a haploid Oenothera. La Cellule 39:159-165. 1929.
6. FLOVIK, K. Cytological studies of Arctic grasses. Hereditas 24:265-376. 1938.
7. GAINES, E. F. and AASE, H. C. A haploid wheat plant. Am. J. Botany 13:373-385. 1926.
8. HARLAND, S. C. Haploids in polyembryonic seeds of Sea Island cotton. J. Heredity 27:229-231. 1936.
9. IVANOV, M. A. Experimental production of haploids in Nicotiana rustica L. (and a discussion of haploidy in flowering plants). Genetica 20:295-397. 1938.
10. KATAYAMA, Y. Haploid formation by X-rays in Triticum monococcum. Cytologia 5:235-237. 1934.
11. KIHARA, H. Formation of haploids by means of delayed pollination in Triticum monococcum. Botan. Mag. 54: 178-185. 1940.
12. KIHARA, H. and KATAYAMA, Y. On the progeny of haploid plants of Triticum monococcum. Kwagau 2:408-410. 1932.

References

1. WILSON, J. P., WILSON, J. P., WILSON, J. P. and WILSON, J. P.. A rapid method for the detection of Salmonella in Salmonella. Salmonella 1972.
2. WILSON, J. P. The Salmonella and Salmonella. Salmonella 1972.
3. WILSON, J. P. Salmonella and Salmonella in Salmonella. Salmonella 1972.
4. WILSON, J. P. and WILSON, J. P. The Salmonella and Salmonella of Salmonella from Salmonella. Salmonella 1972.
5. WILSON, J. P. The Salmonella and Salmonella in Salmonella. Salmonella 1972.
6. WILSON, J. P. and WILSON, J. P. A rapid method for the detection of Salmonella in Salmonella. Salmonella 1972.
7. WILSON, J. P. and WILSON, J. P. A rapid method for the detection of Salmonella in Salmonella. Salmonella 1972.
8. WILSON, J. P. and WILSON, J. P. A rapid method for the detection of Salmonella in Salmonella. Salmonella 1972.
9. WILSON, J. P. and WILSON, J. P. A rapid method for the detection of Salmonella in Salmonella. Salmonella 1972.
10. WILSON, J. P. and WILSON, J. P. A rapid method for the detection of Salmonella in Salmonella. Salmonella 1972.
11. WILSON, J. P. and WILSON, J. P. A rapid method for the detection of Salmonella in Salmonella. Salmonella 1972.
12. WILSON, J. P. and WILSON, J. P. A rapid method for the detection of Salmonella in Salmonella. Salmonella 1972.

13. KIHARA, H. and YAMASHITA, K. Artificial production of haploid and triploid einkorn wheats by pollination with X-ray irradiated pollen. Imp. Bur. Plant Breed. and Genetics. Plant Breed. Abstracts 9:298. 1939.
14. KOSTOFF, D. The problem of haploidy. (Cytogenetic studies on Nicotiana haploids and their bearings to some other cytogenetic problems). Bibliog. Genetics 13:1-148. 1941.
15. LESLEY, M. M. and FROST, H. B. Two extreme 'small' Matthiola plants: a haploid with one and a diploid with additional chromosome fragments. Am. Naturalist 42:22-33. 1928.
16. LEVAN, A. Studies on the meiotic mechanism of haploid rye. Hereditas 28:177-211. 1942.
17. LINDSTROM, E. W. A haploid mutant in the tomato. J. Heredity 20:23-30. 1929.
18. MORINAGE, T. and FUKUSHIMA, E. Studies on the haploid plant of Oryza sativa. Jap. J. Botany 7:75-106. 1935.
19. MUNTZING, A. Characteristics of two haploid twins in D. glomerata. Hereditas 29:134-140. 1943.
20. NORDENSKIOLD, H. Studies of a haploid rye plant. Hereditas 25:204-210. 1939.
21. SEARS, E. R. Cytogenetic studies with ployploid species of wheat. I. Chromosomal aberrations in the progeny of a haploid of Triticum vulgare. Genetics 24: 509-523. 1939.
22. _____ Cytogenetic studies with polyploid species of wheat. II. Additional chromosomal aberrations in Triticum vulgare. Genetics 29:232-246. 1944.
23. SMITH, L. Haploidy in einkorn. J. Agr. Research 73:291-301. 1946.
24. _____ The acetocarmine smear technic. Stain Tech. 22: 17-31. 1947.

25. TOMETORP, G. Cytological studies on haploid Hordeum distichum. Hereditas 25:241-254. 1939.
26. YAMASAKI, Y. Some observations on the microsporogenesis of the haploid plant of Triticum vulgare host. Jap. J. Botany 8:151-153. 1936.

 
the first of these is the fact that the

 