TTA 1522

PHARMACEUTICAL BOTANY

YOUNGKEN

A. Tschirch

Alexander Tschirch, Emeritus Professor of Pharmacognosy in the University of Bern, whose brilliant investigations and numerous contributions in the fields of pharmaceutical botany and pharmacognosy during the present century have added much to our knowledge of drugs.

QK99
.A1
y68
1938

A College
Textbook of

Pharmaceutical Botany

With Pharmaceutical and Medical Applications, especially adapted for the use of Students of Pharmacy and Economic Botany and as a reference for Pharmacists, Chemists and Students in Structural and Systematic Botany.

By

Heber W. Youngken, PH.M., PH.D., SC.D.

Professor of Botany and Pharmacognosy in the Massachusetts College of Pharmacy. Member of the Committee of Revision of the Pharmacopœia of the United States, Botanical Editor of the United States Dispensatory; Author of "A Textbook of Pharmacognosy."

Sixth Edition

Thoroughly Revised

P. Blakiston's Son and Co., Inc.
Philadelphia

MISSOURI BOTANICAL
GARDEN LIBRARY

4207

COPYRIGHT, 1938, BY P. BLAKISTON'S SON & CO., INC.

THE USE IN THIS VOLUME OF CERTAIN PORTIONS OF THE TEXT OF THE UNITED STATES PHARMACOPOEIA, ELEVENTH EDITION, IS BY VIRTUE OF PERMISSION RECEIVED FROM THE BOARD OF TRUSTEES OF THE UNITED STATES PHARMACOPOEIAL CONVENTION. THE SAID BOARD OF TRUSTEES IS NOT RESPONSIBLE FOR ANY INACCURACY OF QUOTATIONS, NOR FOR ANY ERRORS IN THE STATEMENT OF QUANTITIES OR PERCENTAGE OF STRENGTHS.

PERMISSION TO USE FOR COMMENT PARTS OF THE TEXT OF THE NATIONAL FORMULARY, SIXTH EDITION, IN THIS VOLUME HAS BEEN GRANTED BY THE COMMITTEE ON PUBLICATIONS BY AUTHORITY OF THE COUNCIL OF THE AMERICAN PHARMACEUTICAL ASSOCIATION.

PRINTED IN U. S. A.
THE MAPLE PRESS COMPANY, YORK, PA.

TO
MY WIFE
Clara Eastman Youngken

Preface to the Sixth Edition

Since the last edition of "Pharmaceutical Botany" was published, knowledge in plant science has advanced rapidly and a new edition of the United States Pharmacopœia and of the National Formulary have appeared. For these reasons another revision has been called for.

In preparing this revision the writer has endeavored to keep in mind the requirements of varied courses in botany given to students of pharmacy in departments of botany of both the university schools and the independent pharmaceutical institutions. In both types of schools botany should be taught with cultural and professional objectives, and the pharmacy students provided with the proper pharmaceutical slant which need not detract from adequate treatment of the purely academic phases of the subject.

A revision has, accordingly, been prepared which presents a broad general treatment of botany with particular stress upon those phases of this science such as histology, cell contents, taxonomy, and medicinal plants which are believed to be most helpful in preparing students to cope with botanical problems in pharmacognosy and in the practice of pharmacy.

In the course of this revision the writer has not only made changes in the text so as to bring the subject matter up to date but has reorganized some of the chapters, provided some better illustrations and added 121 new figures.

The old chapter on Cytology has been deleted and in its place new chapters have been substituted on "The Living Cell" and on "Non-protoplasmic Cell Contents." Important additions have been made to the subject of the cell and its contents including an amplification of the subject matter on mitosis and

new material on plasmodesma, waxes, plant hormones and vitamins.

A new chapter has been added on "Genetics and Evolution" in keeping with the needs of some of the courses in botany.

The chapters on "The Microscope" and on "Histological Technique" have been transferred from their former positions in the main body of the text to the back of the book where they now appear as Appendix I and Appendix II respectively. New material has been incorporated under both of these captions including modern methods of fixation, staining, imbedding and mounting plant materials for microscopic examination.

The chapter on "Plant Tissues" has been lengthened and improved by the introduction of much new material on the structure and physiology of meristems, endodermis, tracheary tissue, laticiferous tissue, xylem, phloem, the stele, bundles, leaf and branch traces, gaps and nectaries.

The histological and physiological portions of the various chapters dealing with Plant Organs have also been augmented with considerable new text and illustrative material.

The Glossary has been retained and somewhat enlarged as has also the Classified List of Reference Works. A complete Index is also provided, making the data desired by the reader easily found.

Many new drawings, photographs and photomicrographs have been added in this revision which should prove helpful to the student. A considerable number of these are original; many have been borrowed.

The writer desires to express his gratitude to the various publishers and authors from whom permission to use a number of the cuts used in this edition was cheerfully granted. Acknowledgment for these is given under the respective cuts in the text. He is particularly indebted to Dr. A. F. Sievers and the Bureau of Plant Industry for the loan of a number of prints on medicinal plants from which cuts were prepared, to Instructor Richard W. St. Clair of the Massachusetts College of Pharmacy for valued assistance in photographing many of the original figures, to Dr. F. J. Bacon of the Biology Department of Western Reserve University who furnished photographs on *Zamia* plants,

and to Asst. Professor H. L. Reed of the Massachusetts College of Pharmacy who contributed the prints from which Figs. 138, 139 and 140 were prepared.

Finally, the author wishes to express his appreciation to P. Blakiston's Son & Co. Inc. for doing its utmost in producing an attractive and well bound book.

HEBER W. YOUNGKEN.

BOSTON, MASS.,
June, 1938.

Preface to the Fifth Edition

Since the fourth edition of this volume was issued, there have been notable advances in the development of pharmaceutical botany. A new United States Pharmacopœia and a new National Formulary have appeared. The Commonwealth Fund Educational Research Committee has rendered its report on the study of Pharmacy from the functional standpoint and in it the findings of its botanical studies.

These events alone made some additions and alterations of the text necessary. But the writer has gone further. A large part of the text has been rearranged, rewritten and amplified with a view toward improving the clarity of certain portions of the subject matter, and extending its adaptability.

More than a hundred pages of new matter and one hundred and twenty-four new figures comprising original drawings and photographs and borrowed cuts from classic works have been added, and some of the older illustrations have been replaced by better ones.

The botanical findings of the Commonwealth Fund Educational Research Committee have been incorporated, although the text is not limited to these.

Sufficient new information has been added to make the book a distinct improvement over the former editions, and this should make it better for use in professional pharmaceutical colleges and for the courses in botany now prescribed for Pre-medical students in academic colleges, as well as for students in Bachelor of Science courses in colleges of pharmacy.

The table of contents clearly indicates the changes made in the form of the text which now contains twenty-five chapters; an increase of sixteen over the previous edition.

It is believed the increased number of chapters and their new sequence will be found convenient for teachers in assigning work alike for study and collateral reading.

It is hoped that the new edition will also be of value to research students in various fields of botany, since it furnishes in the classified list of reference works a summary of a large amount of literature of the subject.

H. W. Y.

Preface to the Fourth Edition

The value of a thorough systematic training in Botany with emphasis upon the structural and taxonomic aspects of this science is now generally recognized as a prerequisite to the proper comprehension of problems in pharmacognosy and materia medica. But apart from the direct relation of such training to these and other applied fields, Botany *per se* stands out preeminently as a cultural subject and, when presented with laboratory work, is of great value in developing, in students, the power of observation.

In preparing the new edition of this text-book, the author has primarily kept in mind the needs of the pharmaceutical student but he has not been unmindful of its use by students of academic colleges and accordingly has increased the scope of the work.

Fifty-three new pages of subject matter and 25 illustrations have been added. Chapter I on "Fundamental Considerations" has been augmented by the inclusion of a list of standard micro-chemical reagents used in food and drug work together with formulæ for their preparation. Chapter VII on "Plant Organs and Organisms" has been lengthened by 23 pages of new subject matter. Among the outstanding features of its revision may be mentioned the introduction of considerable more plant physiology as well as new drawings and photographs, most of which are original.

There has been an amplification of the subject matter of roots, rhizomes, leaves, seeds and seedlings including data on soil and water relation, diffusion, osmosis, osmosis applied to root hairs, turgor, plasmolysis, essential and non-essential elements absorbed by root hairs, transpiration and the histology of a type of Monocotyl rhizome. New plates illustrating the gross structure of Monocotyl and Dicotyl seeds and of seedlings are introduced for the first time. Chapter IX on Ecology has

been lengthened by 20 pages and several illustrations on Carnivorous plants.

Every part of the work has been carefully revised and brought up to date. As in former editions, the index has been so planned as to make the information contained in this book readily accessible.

To the authors from whom cuts were borrowed, the writer's thanks are due.

H. W. Y.

Preface to the Third Edition

The author has first endeavored to present in a clear, systematic way those fundamental principles of structural and taxonomic botany which serve as a key to the approach of pharmacognic problems. But he has not been unmindful that the use of the work has extended to academic institutions and, so, in this edition, has broadened the scope of the former text. To this end about ninety additional pages of subject matter have been introduced. Several old cuts have been removed. Forty-three new ones have been inserted. Hypothetical discussions have been avoided which saves time for the reader.

The arrangement and plan of the chapters are similar to that of the former edition, in order to adapt the work to several methods of approach. Chapter I on "Fundamental Considerations" has been augmented by treatises on Botanical Nomenclature, Paraffine and Celloidin Imbedding, Sectioning, Staining and Mounting, Microtomes and other information dealing with the preparation of materials for microscopic examination.

Ten pages have been added to Chapter V on Cytology. Under "Protoplasm and its Properties," six pages have been written on the subject of Irritability and Irritable Reactions. Under "Non-Protoplasmic Cell Contents" several additional commercial starches are discussed and two original plates on starch grains added. Additional cuts on Collenchyma, Stone Cells, Sclerenchyma Fibers, Trichomes and Fibrovascular Bundles have been inserted in Chapter VI.

Nine additional pages of subject matter and illustrations have been added to Chapter VII. Original figures of all of the important types of fruits appear here for the first time.

Chapter VIII on "Taxonomy" has been increased by seven pages of new data, and the whole former text carefully revised.

Chapter IX on "Ecology" has been newly introduced as has also a Glossary of Botanical Terms. The index has been so planned as to make the information contained in this book readily accessible.

To the authors of works from which cuts were borrowed the writer's thanks are due.

H. W. Y.

Contents

	PAGES
PREFACE TO THE SIXTH EDITION.	vii
PREFACE TO THE FIFTH EDITION.	xi
PREFACE TO THE FOURTH EDITION	xiii
PREFACE TO THE THIRD EDITION	xv
CHAPTER I	
THE SCOPE AND IMPORTANCE OF BOTANY	
INTRODUCTION.	1-2
DEPARTMENTS OF BOTANICAL INQUIRY.—1. Plant Morphology (Gross Anatomy, Histology, Cytology). 2. Plant Embryology. 3. Plant Physiology. 4. Taxonomy of Plants or Systematic Botany. 5. Plant Ecology. 6. Plant Genetics. 7. Phytopathology. 8. Phytogeography. 9. Phytopalæontology. 10. Cryptogamic Botany. 11. Phanerogamic Botany. 12. Algology. 13. Mycology. 14. Bryology. 15. Economic Botany and its subdivisions	2-4
THE IMPORTANCE OF BOTANY TO MAN	4-5
CHAPTER II	
THE GENERAL CHARACTERISTICS OF PLANTS	
THE PLANT AS A LIVING ENTITY.—Organism, defined. Organ, defined. Vegetative and reproductive processes of plants. Parts of the plant body. Work of the root, stem, green leaves, flowers, fruits and seeds. Tissues. Cells. Protoplasm.	6-12
GROUPS OF PLANTS.—The Thallophytes, Bryophytes, Pteridophytes and Spermatophytes, their etymology, leading subgroups and introduction to their habits. Cryptogams. Phanerogams. Origin and development of a seed plant. Physiological division of labor. Differentiation	12-16
CHAPTER III	
THE LIVING CELL	
THE CELL AS THE FUNDAMENTAL UNIT.—History of the cell and protoplasm. (Hooke, Brown, Schleiden, Schwann, Von Mohl, Corti, Treviranus, Dujardin). The Protoplast. A typical colorless Plant Cell. Typical green cells.	12-37
THE CELL WALL.	23-24

	PAGES
THE AMŒBA.	24-29
THE RELATION OF PLANTS AND ANIMALS	29-30
PROTOPLASM AND ITS PROPERTIES.—Structure. Metabolism. Growth. Irritability. Tropisms, Taxies and Nasties. (Ther- motropism, Chemotropism, Sitotropism, Oxytropism, Hydro- tropism, Rheotropism, Heliotropism, Phototropism, Geotro- pism, Galvanotropism, Thigmotropism). Reproduction. Protoplasmic Movement	30-40
PROTOPLASMIC CELL CONTENTS.—The protoplast. Cyto- plasm; nucleus; nucleolus; plastids (proplastids, leucoplastids, chloroplastids, chromoplastids), cell membranes; chondriosomes. Plasmodesma.	40-44
CELL FORMATION AND REPRODUCTION.—Asexual and sexual cells. Reproduction defined. Asexual Reproduction: Fis- sion; Gemmation; Free Cell Formation; Rejuvenescence or Zoöspore Formation; Vegetative Multiplication. Sexual Repro- duction: Conjugation and Fertilization.	44-47
INDIRECT NUCLEAR DIVISION	47-50
MEIOSIS (Reduction division).	50-52

CHAPTER IV

LIFE HISTORY OF THE MALE FERN [AN INTERMEDIATE TYPE OF PLANT] MEDICINAL USE. PHASES IN ITS LIFE CYCLE	53
HISTORY OF THE SPOROPHYTE OR ASEXUAL GENERATION. —Gross structure of the stem. Gross structure of the frond. Histology of mature stem. Histology of growing stem apex. Histology of mature root. Histology of root apex. Con- tinuity of crude sap flow. Histology of stipe. Histology of a pinnule. Comparative physiology of root, stem and leaf. Gross structure and histology of the sori and sporangia. Rup- ture of sporangium and spore dissemination.	53-68
HISTORY OF THE GAMETOPHYTE OR SEXUAL GENERA- TION.—Formation of protonema and prothallus. Origin of new sporophyte or diploid plant from fertilized egg. Growth of embryo into mature sporophyte. Alternation of generations	68-73

CHAPTER V

NON-PROTOPLASMIC CELL CONTENTS

PROTOPLAST INCLUSIONS.—Sugars. Starch (Assimilation, Tran- sitory and Reserve). Structure and Composition of Starch. Method of Examining Reserve Starches. Characteristics of Important Commercial Starches. Dextrin. Amylodextrin. Inulin. Characteristics of Glucosides. The glucosides Hesperidin, Strophanthin, Salicin, Saponins, Coniferin and Digitoxin. Cyanogenetic glucosides. Pentosides. Alkaloids and
--

CONTENTS

xix
PAGES

their properties. The alkaloids, Strychnine, Brucine, Nicotine, Caffeine, Cocaine, Aconitine, Morphine, Codeine and Colchicine. Gluco-alkaloids. Asparagine. Calcium Oxalate. Micro-crystals. Cystoliths. Silica. Tannins. Proteins. Aleurone Grains. Mucilages and Gums. Fixed Oils and Fats. Waxes. Volatile Oils. Resins. Oleoresins. Gum Resins. Balsams. Pigments (Chlorophyll, Carotin, Xanthophyll, Etiolin, Anthocyanins, etc.) Latex. Enzymes. Classification of Enzymes. Plant Hormones (Auxins). Vitamins 74-107

CELL WALLS.—Their formation and composition. Growth in area and thickness. Pores (Pits). Various kinds of cell walls and behavior of each to micro-chemic reagents 107-110

CHAPTER VI

PLANT TISSUES

PLANT TISSUES.—Tissue defined. The Tissues of Spermatophytes and Pteridophytes: Classification of tissues. Primary and secondary tissues. Simple and complex tissues. Permanent tissues. Primary Meristems. List of Tissues. 111-114

MERISTEM.—Definition. Primordial meristem. Primary meristems (dermatogen, periblem and plerome) secondary meristems (cambium and phellogen). Apical, intercalary and lateral meristems. Their distribution 114-116

PARENCHYMA.—Definition; ordinary parenchyma; chlorenchyma or assimilation parenchyma; conducting parenchyma; reserve parenchyma; their structure, distribution and function 116-117

COLLENCHYMA.—Definition, structure, function and distribution 117-118

SCLERENCHYMA.—Definition; stone cells; sclerenchyma fibers; wood fibers; phloem- or bast-fibers; pericyclic fibers; cortical fibers; their distribution 118-122

EPIDERMIS.—Definition; transpiration and water stomata; epidermal papillæ; trichomes; scales; their distribution and functions. 122-127

ENDODERMIS.—Definition, structure, distribution and functions. Casparian spots. 127-129

CORK.—Definition; its derivation, function and distribution. Rhytidome. Borke. 129-131

PERICYCLE.—Derivation, structure, position and function 131

LATICIFEROUS TISSUE.—The structure, origin and distribution of latex cells, laticiferous vessels and cœnocytic latex cells. Latex. 131-133

SIEVE (LEPTOME) OR CRIBIFORM TISSUE.—Definition; derivation; structure; distribution; function. 133-135

TRACHEARY TISSUE.—Tracheæ: Definition, origin, structure, function and classification; Tracheids: Definition, structure and function. Distinction from tracheæ. 135-139

MEDULLARY RAYS.—Primary and Secondary; their structure, distribution and functions.	139-140
VASCULAR AND FIBRO-VASCULAR BUNDLES.—Description of the various types; the distribution of each type; Xylem and Phloem. Leaf and branch traces.	140-144
THE STELE.—Definition. Prostele. Siphonostele. Dictyostele. Branch Traces. Leaf and Branch Gaps	144-147
SECRETION SACS.—Definition and distribution.	147
INTERCELLULAR AIR SPACES.—Definition. Function. Schizogenous and lysigenous air spaces; their relative size.	147-148
INTERNAL SECRETION GLANDS AND RESERVOIRS.—Definition; structure; internal glands; secretion canals; vittæ. Their origin	148-149
NECTARIES	150
CLASSIFICATION OF TISSUES ACCORDING TO FUNCTION	151

CHAPTER VII

THE ROOT

THE ROOT.—Definition; functions; rootlets; root hairs; piliferous layer; soil and water relation, diffusion, osmosis, osmosis applied to root hairs, turgor, plasmolysis, absorption of nutrient salts, essential elements and their importance, non-essential elements absorbed by root hairs; regions of the root (root cap, embryonic region, growing point, primary meristems or generative tissues, region of elongation, region of maturation and mature region). Differences between root and stem	151-160
CLASSIFICATION OF ROOTS AS TO ORDER.—Primary root; tap root; secondary roots; fibrous and fleshy roots; prop roots; adventitious roots; epiphytic roots; haustoria.	160-161
CLASSIFICATION OF ROOTS AS TO FORM.—Fusiform, napiform, conical fibrous, tortuous, nodose, moniliform, tuberous, fascicled	161-162
ROOT CONSISTENCY OR TEXTURE.—Woody roots. Fleshy roots: single and multiple	162
CLASSIFICATION OF PLANTS ACCORDING TO DURATION OF ROOT.—Annual; Biennial; Perennial	162-163
ROOT HISTOLOGY.—A. <i>Monocotyledons</i> : Histology of Similax root. B. <i>Dicotyledons</i> : Histology of Dicotyl roots: A. young growth, B. transitional growth and development of secondary structure. Histology and Development of the California Privet root. Abnormal structure of Dicotyl roots. Histology of a Dicotyl Tuberous Root (<i>Aconitum</i>). Root branching.	163-176
ROOT TUBERCLES.—Definition; occurrence. Leguminous tubercles. Rotation of crops. <i>Myrica</i> tubercles; their etiology. Nitrogen cycle	176-182

MYCORRHIZA.	183
---------------------	-----

CHAPTER VIII

THE BUD

BUDS.—Definition; plumule, scaly buds, naked buds; Classification of Buds According to Position on Stem: terminal bud; axillary or lateral bud; adventitious bud; accessory bud; dormant bud; sub-petiolar bud. Classification of Buds According to Development: leaf bud; flower bud and mixed bud. Classification of Buds According to their Arrangement on the Stem: alternate; opposite; whorled.	184–185
---	---------

CHAPTER IX

THE STEM

THE STEM.—Definition; direction of growth; functions; size; nodes and internodes; stem elongation; generative tissues; duration of stems; forms of stems; stem modifications; above ground stems; herb, shrub and tree defined; fall of leaves; underground stems (rhizomes, bulbs, corms, tubers, suckers) exogenous and endogenous stems.	186–194
STEM HISTOLOGY.—Histology of Annual Dicotyl Stem, Growth of Perennial Dicotyl Stem and its Histology. Exceptional Types of Dicotyl Stems; Lenticels and Their Formation; Annual Thickening of Stems; Method of formation; “Annual Ring.” Bark: Definition; zones; Cork; Periderm; Phelloderm; Histology of Cascara Sagrada bark. Wood: alburnum; duramen; tyloses; microscopic characteristics of Angiospermous and Coniferous woods. Histology of Typical Herbaceous Monocotyl Stems. Histology of a Typical Monocotyl Rhizome. Histology of a Typical Woody Monocotyl Stem. Histology of Grass Stems. Secondary Growth in Monocotyl Stems.	194–220

CHAPTER X

THE LEAF

THE LEAF.—Definition; the complete leaf and its structure; sessile, petiolate, stipulate and exstipulate leaves; leaf functions: photosynthesis; assimilation; respiration; transpiration.	221–227
TYPES OF LEAVES DEVELOPED IN ANGIOSPERMS.—Cotyledons; Scale leaves; Foliage leaves; Bract leaves: bracts and bracteoles; Sepals; Petals; Microsporophylls; Megasporophylls	227–228
ORIGIN AND DEVELOPMENT OF LEAVES.—Primordial leaf. Its formation	228–229
PHYLLOTAXY.—Definition; spiral law of leaf arrangement; alternate; opposite, decussate; whorled; fascicled; leaf rank.	229–230
VERNATION.—Definition; inflexed or reclinate; conduplicate; convolute; circinate; plicate; involute; revolute.	230

LEAF VENATION.—Furcate; parallel; reticulate; pinnately-veined; palmately-veined; anastomosing, tessellated and impressed veins	230–232
LEAF INSERTION.—Definition; radical; cauline; ramal; perfoliate; amplexicaul; connate-perfoliate; equitant.	232
FORMS OF LEAVES.—Simple and Compound. Forms of the Lamina: (a) General Outline; linear; filiform; ovate, obovate, rhomboidal; lanceolate; elliptical; oblong; equilateral; inequilateral; reniform; orbicular; peltate; oblanceolate; cuneate; spatulate; ensiform; subulate; acerose; falcate, deltoid. (b) Apex: acute; acuminate; tapering; attenuate; obtuse; rounded; truncate; mucronate; cuspidate; aristate; emarginate; retuse; obcordate. (c) Base: cordate; reniform; hastate; auriculate; cuneate; peltate; sagittate; oblique; acute, tapering, attenuate, truncate; obtuse, rounded, decurrent, etc. (d) Margin: entire; revolute; serrate; serrulate; dentate; denticulate; crenate; crenulate; doubly serrate, etc.; spinose, ciliate; sinuses, segments and lobes; repand; sinuate; incised; runcinate; lobed; cleft; parted; divided; pinnatifid; pinnatipartite; pinnatisect; palmatifid; palmatipartite; palmatisect; pedate; interruptedly-pinnate; lyrate. Forms of Compound Leaves	233–240
LEAF MODIFICATIONS.	240–241
LEAF TEXTURE.—Membranous; succulent; coriaceous.	241
LEAF COLOR.—Variations in color	241
LEAF SURFACE.—Glabrous; glaucous; pellucid-punctate; papillose; scabrous; pubescent; appressed-hairy; villose; sericous; hispid; strigose; tomentose; spinose; rugose; verrucose; tuberculate; bullate	241–242
DURATION OF LEAVES.—Persistent or evergreen; deciduous; caducous; fugacious	242
GROSS STRUCTURE AND HISTOLOGY OF THE LEAF PETIOLE.—In Monocotyledons. The ligule and pulvinus. In Dicotyledons. Pulvinus. Aggregation Body. Pericladium. Phyllode.	242–245
STIPULES.—Definition; lateral; free-lateral; lateral-adnate; lateral-connate; lateral interpetiolar. The ligule. Axillary: free axillary and connate. The ochrea. Modified Stipules.	245–246
THE LAMINA.—Definition. Mode of Development of the Lamina of Leaves: Dorsiventral; convergent; centric; bifacial; reversed ob-dorsiventral. A. Dorsiventral: (a) Dorsiventral Umbrophytic; (b) Dorsiventral Mesophytic; (c) Dorsiventral Xerophytic; (d) Dorsiventral Hydrophytic. Gross Structure and Histology of Different Types of Dorsiventral Leaf Blades. Gross Structure and Histology of the following types: B. Convergent; C. Centric; D. Bifacial.	246–251
STRUCTURE AND DEVELOPMENT OF STOMATA	251–254

CONTENTS

xxiii

PAGES

HISTOLOGIC DIFFERENCES BETWEEN LEAVES OF DICOTYLEDONS AND MONOCOTYLEDONS	255
--	-----

CHAPTER XI

THE INFLORESCENCE

INFLORESCENCE.—Definition. Determinate, Indeterminate and Mixed Inflorescences defined. Parts of Inflorescences: peduncle, rachis, pedicel, bracts, involucre, bracteoles. Scape. Spathe.	256-257
INDETERMINATE INFLORESCENCES.—Elongated and flat-topped. Forms of: Solitary indeterminate; raceme; corymb; umbel; compound umbel; spike; compound spike; catkin; head; strobile; spadix; panicle	257-259
DETERMINATE INFLORESCENCES.—Elongated and flat-topped. Forms of: Solitary determinate; cyme; simple and dichasial cymes; scorpioid cyme; glomerule; verticillaster	259-261
MIXED INFLORESCENCES.—Thyrsus; raceme of cymes; corymb of cymes.	261

CHAPTER XII

THE FLOWER

THE FLOWER.—Definition; floral parts; essential organs; accessory organs; complete; perfect; hermaphrodite; regular or actinomorphic; irregular; zygomorphic; symmetrical; imperfect; double; staminate; pistillate; neutral; hypogynous; perigynous; Epigynous. Monœcious, diœcious and polygamous plants.	262-264
PREFLORATION: Definition. Convolute; involute; revolute; plicate; imbricate; valvate; vexillary; contorted.	264-265
CONNATION AND ADNATION.—Definitions and illustrations	265
THE RECEPTACLE, TORUS OR THALAMUS.—Definition; variations in structure; anthophore; gonophore; gynophore; carpophore.	265-266
THE PERIANTH.—Definition; dichlamydeous; monochlamydeous.	266
THE CALYX.—Definition; parts; physical characteristics; connation; kinds and forms; persistence; adnation; sepaline nectaries and spurs; sepaline stipules; the epicalyx; sepaline position	266-269
THE COROLLA.—Definition; parts; physical characteristics; functions. Forms of the Corolla and Perianth. Appendages of the Perigone	269-274
THE ANDRŒCIUM OR STAMEN SYSTEM.—Definition; parts; terms denoting number of stamens in flower; insertion of stamens; proportions of stamens; connation of stamens; stamen color. Gross Structure and Histology of the Filament. Gross Structure and Histology of the Anther. Another Dehiscence.	

	PAGES
Development of the Anther. Attachment of Anther. Pollen; description; forms. Pollinia. Hay Fever pollens.	274-283
THE GYNÆCIUM OR PISTIL SYSTEM.—Definition; Gymnospermous and Angiospermous; parts; the pistil a modified leaf; carpel; dehiscence; apocarpous and syncarpous pistils; terms denoting the number of carpels entering into the formation of the pistil; compound pistils; ovules or megasori as transformed buds; position of ovules in ovary; Gymnospermous and Angiospermous ovules; structure of Angiospermous ovule; forms of ovules.	283-287
THE PLACENTA.—Definition; types of placenta arrangement	287-288
THE STYLE.—Definition; style-arms; relation to carpels forming the gynœcium; variations from typical styler development	288-289
THE STIGMA.—Definition; forms in wind- and animal-pollinated flowers; stigmatic papillæ.	290
POLLINATION.—Definition. Close and Cross Pollination. Protandrous and Protogynous flowers. Terms applied to plants pollinated by various agencies.	290-292
MATURATION OF THE POLLEN GRAIN AND FORMATION OF MALE GAMETOPHYTE (in Angiosperms)	292
MATURATION OF THE EMBRYO SAC AND FORMATION OF THE FEMALE GAMETOPHYTE (in Angiosperms)	292
FERTILIZATION IN ANGIOSPERMS.—Process; formation of embryo and endosperm; double fertilization	292-294
DEVELOPMENT OF THE EMBRYO.—1. In Dicotyledons. 2. In Monocotyledons	295-296

CHAPTER XIII

THE FRUIT

THE FRUIT.—Definition; modifications.	297
FRUIT STRUCTURE.—Pericarp; pseudocarp; anthocarp; epicarp; mesocarp; endocarp; sarcocarp; putamen; sutures; valves; dehiscent and indehiscent fruits; kinds of dehiscence	297-300
CLASSIFICATION OF FRUITS.—Simple; Aggregate; Multiple; dry dehiscent; dry indehiscent; fleshy indehiscent. Forms of Simple Fruits: 1. Capsular (follicle, legume, loment, capsule, pyxis, regma, siliqua, sillicule). 2. Schizocarpic (carcerulus, cremocarp, samara). 3. Achenial (akene, utricle, caryopsis, nut). 4. Baccate (berry, uva, bacca, pepo, hesperidium). 5. Drupaceous (drupe, pome). Aggregate Fruits: Etærio. Forms of Multiple Fruits: strobile or cone; sorosis; syconium; galbulus.	300-307
HISTOLOGY OF A CAPSULE, VANILLA.	307-309
HISTOLOGY OF A MERICARP, FCENICULUM.	309-310

CHAPTER XIV

THE SEED

- THE SEED.—Definition; The Embryo and its parts; vitality of seeds; seed structure; functions; appendages including the coma, awn, caruncle, etc.; aril and arillode; kernel, funiculus; hilum; raphe. 311–314
- MODE OF FORMATION OF DIFFERENT TYPES OF ALBUMEN:
Perispermic; endospermic; perispermic and endospermic.
Perisperm; Endosperm; exalbuminous and albuminous seeds. 314–315
- A MONOCOTYL SEED.—Its gross structure and histology 315–318
- GERMINATION OF THE SEED.—Germination of Indian Corn and structure of the seedling.. . . . 318–319
- A DICOTYL SEED.—Its gross structure and histology; germination of dicotyl seeds.. . . . 319–322

CHAPTER XV

THE CLASSIFICATION AND NAMING OF PLANTS

- PRINCIPLES OF CLASSIFICATION.—The modern classification of plants and its basis. Organic evolution. Natural system. Artificial system. Artificial systems of Theophrastus, Fuchs, Bock, Cæsalpinus, Bauhin, Ray, Tournefort and Linnaeus. The binomial system. Natural systems of De Jussieu, De Candolle, Bentham and Hooker, Eichler, Engler and Prantl, and Hutchinson. The groups of a modern natural system. Complete classification of a medicinal plant, Aconite. 323–328
- OUTLINE OF PLANT GROUPS 328
- BOTANICAL NOMENCLATURE.—Practice of Pre-Linnean botanists. The binomial system of Linnaeus. Rules for the naming of genera, species, varieties, families, orders, classes, sub-classes, divisions and sub-divisions. Names of Cryptogram groups. . 329–332

CHAPTER XVI

THE THALLOPHYTES

- GENERAL CHARACTERISTICS OF THALLOPHYTES 333
- ALGAE.—Characteristics of group 333–334
- CYANOPHYCEÆ (BLUE GREEN ALGÆ).—Their characteristics.
Glœocapsa. Oscillatoria. Nostoc 334–336
- CHLOROPHYCEÆ (GREEN ALGÆ).—Their characteristics. Orders of green algae and life histories of representative forms of each. . 336–348
- BACILLARIEÆ (DIATOMS).—Characteristics, distribution and economic uses 348–351
- PHAEOPHYCEÆ (BROWN ALGÆ).—Their characteristics and economic uses. Life history of a filamentous type, Ectocarpus. The kelps. The rockweeds. Life history of Fucus. 351–357

RHODOPHYCEÆ (RED ALGÆ).—Their characteristics. Chondrus, Agar and Irish Dulse	357-361
--	---------

CHAPTER XVII

THE FUNGI

FUNGI.—Their characteristics and classes	362
SCHIZOMYCETES (BACTERIA).—Definition and characteristics. Classification according to the phenomena they produce. Fermentation, decay and putrefaction. Bacterial cultures. Forms of bacteria. Motility. Nutrition. Nitrification. Den- trification. Growth Products. Reproduction. Morphology due to cleavage. Forms of cell groups after cleavage. Sporula- tion. Rapidity of growth and multiplication. Chemical com- position. Exotoxins and endotoxins.	363-370
MORPHOLOGICAL CLASSIFICATION OF BACTERIA.—The lower bacteria. The higher bacteria. Pathogenic bacteria. Some bacteria producing disease in man, the lower animals and in plants.	370-374
MYXOMYCETES (SLIME MOLDS).—Their characteristics. . . .	374-375
EUMYCETES (TRUE FUNGI).—Their characteristics and sub-classes	375-376
PHYCOMYCETES (ALGA-LIKE FUNGI).—Their characteristics. The orders Zygomycetales and Oömycetales. Cultivation of Molds. Life history of the Black Bread Mold. Related molds.	376-381
ASCOMYCETES (SAC FUNGI).—Their characteristics. Order Protoascales. Family Saccharomycetaceæ. The Yeasts. Life history of a yeast, <i>Saccharomyces cerevisiæ</i> . Top and bottom yeasts. Wine yeast. Other yeasts. The orders Pezizales, Plectascales, Tuberales, Helvellales, Pyrenomycetales. Characteristics of the cup-fungi, blue and green molds, truffles, saddle-fungi and black-fungi. Production of citric acid by molds. Life history of <i>Claviceps purpurea</i>	381-397
BASIDIOMYCETES (BASIDIA FUNGI).—Their characteristics. The Lower Basidiomycetes. 1. The Ustilaginales or Smuts: their characteristics. Life history of corn smut. 2. The Uredinales or Rusts: Characteristics. Life history of wheat rust. The Higher Basidiomycetes: Their characteristics and sub-groups, 1. Hymenomycetes: weeping fungi, exobasidiales, coral-fungi, mushrooms and toadstools including tooth-fungi, pore-fungi, and gill-fungi. Groups of Agaricaceæ based on color of their spores. Spore prints. Life history of the Common Field Mushroom and of the Fly- and Deadly-Agaric. Antidote for poisoning by the Fly Agaric. 2. Gasteromycetes: puff-balls, earth-stars, nest-fungi and carrion-fungi.	397-411
FUNGI IMPERFECTI.—Their characteristics, and forms producing diseases in man	412

CHAPTER XVIII

THE LICHENS

- LICHENS.—Definition. Asco-lichens. Basidio-lichens. Foliaceous, crustaceous and fruticose lichens. Structure of a typical lichen. Reproduction. Economic uses of various lichens 413–417

CHAPTER XIX

THE BRYOPHYTES

- BRYOPHYTES.—Their general characteristics. 418
 HEPATICÆ OR LIVERWORTS.—Their characteristics and orders. . 418–419
 MARCHANTIA (A TYPICAL LIVERWORT).—Life history and structure. 419–423
 MUSCI OR MOSSES.—Their characteristics. Sphagnales or bog mosses. Andraeales or rock mosses. Bryales or true mosses. . 426–430
 LIFE HISTORY OF A TYPICAL TRUE MOSS (POLYTRICUM). 426–430

CHAPTER XX

THE PTERIDOPHYTES

- PTERIDOPHYTES.—Their general characteristics and classes. 431
 THE FERNS.—Their characteristics. Eusporangiatæ and Leptosporangiatæ. True ferns. Water ferns 431–437
 THE HORSETAILS.—Their characteristics. Life history of *Equisetum arvense* 437–439
 THE CLUB MOSSES.—Their characteristics. Life history of *Lycopodium clavatum*. The Little Club Mosses and Quillworts. . . . 439–442

CHAPTER XXI

THE SPERMATOPHYTES

- SPERMATOPHYTES.—Characteristics and subdivisions. 443
 THE GYMNOSPERMS.—Their characteristics and sub-groups. Life history of a Gymnosperm (*Pinus Strobus*). The Cycads or fern palms. Life history of *Zamia*. The Ginkgos. The Conifers. Classification of Conifers. The Pinaceæ. Table of official and unofficial drugs yielded by the Pinaceæ, including official names, parts used, botanical origins and habitats. The Taxaceæ and representatives. The Gnetales. The Gnetaceæ. *Welwitschia*, *Gnetum* and *Ephedra*. The drug *Ephedra* and its uses. . . . 443–471
 THE ANGIOSPERMS.—Their characteristics. Life history of an Angiosperm (*Erythronium*). Resemblances between Gymnosperms and Angiosperms. Fundamental differences between Gymnosperms and Angiosperms. Classes of Angiosperms. . . 471–480

CHAPTER XXII

THE MONOCOTYLEDONS

MONOCOTYLEDONS.—Their characteristics. 481-483

DRUG YIELDING MONOCOTYLEDONS.—The orders and families of monocotyledons yielding drugs with characteristics of the families. Tables of official and unofficial drugs with names of drugs, parts used, botanical origins and habitats. 483-496

CHAPTER XXIII

THE DICOTYLEDONS

DICOTYLEDONS.—Their characteristics. 497-498

DRUG YIELDING DICOTYLEDONS.—The sub-classes, orders and families of Dicotyledons yielding drugs. Tables of official and unofficial drugs under each family description, with the drug names, parts used, botanical origins, and habitats. . . . 499-615

CHAPTER XXIV

ECOLOGY

INTRODUCTION.—Ecology defined. Environmental factors. . . . 616

PLANT COMMUNITIES AND ASSOCIATIONS.—Definitions. Classification based on relation plant associations have assumed in regard to water. Characteristics of Hydrophytes, Helophytes, Halophytes, Xerophytes, Mesophytes and Tropophytes. Plant Succession. 616-621

RELATION OF PLANTS AND ANIMALS.—CARNIVOROUS PLANTS: The Sundews, Venus Fly-trap, Submerged *Dionæa*, the Pitcher Plants, the Bladderworts, the Butterworts and their biological relations. Significance of the carnivorous habit. 621-642

CHAPTER XXV

GENETICS AND EVOLUTION

SCOPE OF GENETICS.—Genetics and Heredity. Characters. Hereditary substance 643

MENDEL'S LAWS OF ALTERNATIVE INHERITANCE.—Mendel and his experiments. Allelomorphs. Principles of Mendel's Laws. Structure of the Chromosome. Male and female chromosomes. Distribution of chromosomes during maturation and fertilization. Distribution of genes. Dominants and recessives. Genotype and Phenotype. Monohybrids. Results of crossing individuals with one pair of contrasting characters. Results of crossing individuals with two pairs of contrasting characters. 643-652

GALTON'S LAW OF ANCESTRAL INHERITANCE 652

VARIATIONS.—Definition. Kinds of variations including mutations. 652-653

PHARMACEUTICAL BOTANY

CHAPTER I

THE SCOPE AND IMPORTANCE OF BOTANY

BOTANY is the science which treats of plants. Plants are living things or organisms belonging to one of the two great kingdoms of living things, called the VEGETABLE or PLANT KINGDOM. The other kingdom of living creatures is that made up of animals and called the ANIMAL KINGDOM. The science which treats of animals is called ZOÖLOGY. Botany and Zoölogy collectively constitute the branch of natural science called BIOLOGY, the science which seeks to inquire into all living things.

The term botany is derived from the Greek word *βοτάνη* meaning pasture, grass, or fodder. Man's first inquiry into plants was undoubtedly prompted by his instinct to seek food and doubtless began with his first appearance on the earth. History records the first pursuit of botany as a science by the ancient Chaldeans, Egyptians and Greeks who cultivated it particularly in relation to sources of food supply.

Later man found that some plants possessed medicinal properties and interest was next focused in the direction of plants as sources of medicines. The earliest pioneers in pharmaceutical botany were the medicine men of primitive peoples including the Indians of America who were acquainted with the virtues of many medicinal plants before the coming of the white settlers. They were followed in Europe by the *rhizotomoi* or root collectors who gathered medicinal herbs and prepared them for the *pharmacopolai* or drug venders. Probably the greatest of the early European pioneers in this field was Theophrastus, a Greek philosopher and pupil of Aristotle, who lived between 372 and 285 B.C. He wrote ten books on the *History of Plants* in which about 500 species of plants used in the treatment of various

diseases are described. Dioscorides, a Greek physician, about 77 A.D. wrote a *Materia Medica* which included a large number of descriptions of medicinal plants and all the medicinal substances then known together with their properties. The early Romans also contributed to the development of Pharmaceutical Botany. Outstanding among these were Pliny the Elder (23–79 A.D.), who in his *Historia Naturalis* crudely described about 1000 plants, many of which possessed medicinal properties, and Galen (131–200 A.D.), a Græco-Roman, who wrote a *Materia Medica*.

Attention was next concentrated upon the classification of plants, and gradually investigations followed on their structure, functions, habits, relationships, diseases, etc., until to-day the scope of botanical study embraces every kind of inquiry about plants.

The field of botany has thus developed many specialties or departments of inquiry.

DEPARTMENTS OF BOTANICAL INQUIRY

1. PLANT MORPHOLOGY treats of the parts, or structure of plants. It is divided into:

(a) MACROMORPHOLOGY or GROSS ANATOMY which deals with the external characters of plants and of their parts or organs; (b) MICROMORPHOLOGY or HISTOLOGY which considers the minute or microscopical structure of plants and plant tissues and (c) CYTOLOGY which treats of plant cells and their contents.

2. PLANT EMBRYOLOGY treats of the stages of growth, differentiation and development of the individual plant body.

3. PLANT PHYSIOLOGY deals with the study of the life processes or functions of plants. It explains how the various parts of plants perform their work of nutrition, growth, reproduction, and the preparation of food for the support of animal life from substances not adapted to that use.

4. TAXONOMY OF PLANTS or SYSTEMATIC BOTANY considers the classification or arrangement of plants in groups in accordance with their relationships to one another. It also is concerned with the identification, the accurate description and the naming of plants. All of the plants found growing in a certain region constitute the *Flora* of that region.

5. PLANT ECOLOGY treats of plants and their parts in relation to their environment, *i.e.* to soil, moisture supply, temperature, light and other plants and animals.

6. PLANT GENETICS deals with the coming into being of new individuals or races of plants. It seeks to account for the resemblances and differences shown by plants related by descent.

7. PLANT PATHOLOGY or PHYTOPATHOLOGY treats of diseases of plants.

8. PLANT GEOGRAPHY or PHYTOGEOGRAPHY treats of the distribution of plants upon the earth. The center of distribution for each species of plant is the *habitat* or the original source from which it spreads, often over widely distant regions. When plants grow in their native countries they are said to be *indigenous* to those regions. When they grow in a locality other than their original home they are said to be *naturalized*.

9. GEOLOGICAL BOTANY or PHYTOPALÆONTOLOGY treats of plants of former ages of the earth's history traceable in their fossil remains.

10. CRYPTOGAMIC BOTANY is the botany of the flowerless plants or cryptogams such as the pond scums, fungi, mosses, ferns and their allies.

11. PHANEROGAMIC BOTANY is the botany of seed plants.

12. ALGOLOGY is the botany of algae.

13. MYCOLOGY is the botany of fungi.

14. BRYOLOGY is the botany of mosses and liverworts.

15. ECONOMIC or APPLIED BOTANY deals with the science from a practical standpoint, showing the special adaptation of the vegetable kingdom to the needs of everyday life. It comprises a number of subdivisions, viz.: PHARMACEUTICAL BOTANY, or the botany of drug, spice, dye and poisonous plants; BACTERIOLOGY, which treats of microorganisms called bacteria; PHARMACOGNOSY which treats of the history, commerce, selection, identification, valuation and preservation of crude drugs and other raw materials of plant and animal origin; AGRICULTURAL BOTANY or AGRONOMY, which treats of crop plants; HORTICULTURE, which treats of the cultivation of garden and orchard plants; FORESTRY which treats of the management of forests and of timber plants; PLANT BREEDING or the application of principles

of selection to the growing of improved races of plants. All of these subdivisions interlock closely with departments of pure botanical inquiry and no training in either of them can be considered sound without the assimilation of the broad underlying principles of morphology, taxonomy and physiology.

PHARMACEUTICAL BOTANY should include a sound foundation in the morphology, taxonomy and elementary physiology of plants before taking up its applied aspects, in order to adequately prepare the student for the more advanced studies of Phar. macognosy, Industrial Microscopy and Drug Plant Cultivation-

THE IMPORTANCE OF BOTANY TO MAN

It is an established fact that without plants there could be no human or lower animal life, for green plants supply the food for all living animals and maintain the oxygen content of the atmosphere. They supply man with timber to build homes, ships and furniture, with hairs and fibers with which to manufacture clothing, papers and books, with fuel to heat our buildings, with dyes to color our garments and ornaments and preparations, with oils and resins to manufacture our soaps, paints and varnishes, with drugs to treat our ills, with raw materials for our explosives, with spices and flavorings to season our food, and with vitamins without which we could not maintain our health, with perfumes, waxes, etc. In fact, for our very existence, comfort and protection we are dependent upon plant life.

Again, many of the fundamental principles of biology and genetics have been first established through observations and experiments with plants.

We look to the plants for a large number of our drugs and to the patterns obtained by the examination of many of their constituents for the chemical synthesis of a large number of important organic chemicals, medicines and dyes. Coal, representing the fossil remains of plants which lived millions of years ago, furnishes the chemist with the raw material from which he has built up hundreds of organic chemicals which are daily used in pharmacy, medicine and the industries.

Without some of the microscopic forms of plants, especially the bacteria, yeasts and molds which represent agencies of decay and fermentation, the earth would be literally covered with the remains of the dead bodies of animals and plants accumulated through the ages. While a number of these produce disease, most are believed to be harmless, and many are of great economic importance to pharmacy, agriculture, and to the dairy, alcohol and tanning industries.

The study of botany unfolds many of *Nature's* secrets, enhances one's appreciation of *Nature's* beauty, develops one's powers of observation, and forms a sound foundation for the understanding of many problems in pharmacy, chemistry, agriculture, industry, and the biological sciences.

CHAPTER II

THE GENERAL CHARACTERISTICS OF PLANTS

If we look about in nature we are confronted at once with many kinds of living objects called plants, but the more familiar ones to the beginner in Botany are the higher green plants such as the common herbs, shrubs and trees of field, meadow or woodland. Each of these is a living entity or ORGANISM which has descended from a parent or parents with an ancestry and which is capable of carrying out all of the life processes which are essential to its existence and to the perpetuation of its species or kind. Each is composed of ORGANS whose life processes are separate but mutually dependent and essential to the healthful maintenance of the individual, and the continuation of its kind.

These organs are the ROOTS, STEMS, LEAVES, FLOWERS, FRUITS and SEEDS. The roots, stems and leaves are so constructed as to carry out the VEGETATIVE PROCESSES, *i.e.*, the activities which have to do with the maintenance and protection of the individual, whereas the flowers, fruits and seeds are structurally adapted to the carrying out of the REPRODUCTIVE PROCESSES or the activities which have to do with the continuance of the species of the plant of which the individual is a representative.

PARTS OF THE PLANT BODY.—The parts of the body of a higher green plant are the AXIS and APPENDAGES. The axis consists of an usually aerial portion or STEM and an underground portion or ROOT. The appendages include the LEAVES, BRANCHES, EMERGENCES such as prickles, and the HAIRS.

The ROOT grows down into the soil and anchors the plant, absorbs water with nutrient salts in solution and stores food to tide the plant over the season when growth activity is suspended. It represents the descending axis. The STEM grows upward and so represents the ascending axis of the plant. Its special processes are those of bearing and supporting leaves, branches

FIG. 1.—Diagram of the body of a seed plant, showing its principal parts. (From Robbins, after Holman and Robbins.)

flowers and fruits, conducting sap containing nutrient materials from roots to leaves and other aerial organs and vice versa. Many stems also store food and green parts of stems manufacture

food. The GREEN LEAVES, representing outgrowths of the stem, are spread out to the sun's rays for carrying out the process of manufacturing complex food from simple inorganic materials absorbed from the soil and the air. This process is called *photosynthesis*. The FLOWERS are equipped with sexual organs called STAMENS (male organs) and PISTILS (female organs) and are so constructed as to bring about fertilization. The FRUITS which follow the flowers function to form and disseminate the seeds. The SEEDS serve to carry a young plant or EMBRYO which has separated from its parent in its dormant stage to the soil and so function to spread the plant.

FIG. 2.—Flower and fruit of Flax. A, floral diagram—*c*, calyx; *co*, corolla; *s*, stamens; *p*, pistil. B, median length-wise section of flower. C, flower with calyx and corolla removed showing the essential organs, stamens and pistil. D, fruit, external view. E, cross section of fruit. (Robbins.)

In addition to the life processes noted, all of these plant organs carry out activities necessary for their own maintenance such as TRANSPORTATION OF FOOD MATERIALS, GROWTH (increase in size), RESPIRATION (the breaking down of carbohydrates accompanied by the taking in of oxygen and giving off of carbon dioxide), and ASSIMILATION (the changing of food materials into living matter).

Just as the plant organism is made up of organs which are visible to the unaided eye, so each of these organs is made up of parts called TISSUES, but the tissues are visible only with the aid of a simple or compound microscope.

Some of the TISSUES in the stem for instance, are the outer protective tissue or EPIDERMIS, the soft region beneath the epidermis and in the pith of the stem or PARENCHYMA, the woody

FIG. 3.—Cross section of a representative portion of the stem of a clover (*Trifolium*) illustrating the tissues and tissue regions found in a stem of an herbaceous dicotyledon. $\times 80$. *ep.*, epidermis or outer protective tissue; *s*, stoma, *ss*, sub-stomal air-space, *co*, primary cortex containing parenchyma, *f*, sclerenchyma fibers, *ph*, phloem containing sieve tissue, *ca*, cambium, and *x*, xylem of an open collateral fibrovascular bundle; *mr*, medullary ray composed of parenchymatous cells, *m*, medulla or pith composed of large parenchyma cells and small angular air-spaces. Note the xylem contains long conducting tubes, here shown in cross section as large open circular areas arranged radiately, which conduct water upward through the plant axis.

portion which gives support to the stem or SCLERENCHYMA (stony tissue), the water tubes which conduct crude sap upward or TRACHEARY TISSUE and the sieve tubes and phloem cells which carry elaborated sap downward or SIEVE TISSUE.

Other plant organs are also differentiated into secondary parts or tissues which have been set aside to perform some definite kind of work in the economy of the organs. The healthful performance of the functions of an organ depends upon the harmonious relation existing between the tissues of which it is composed.

FIG. 4.—Sections of fixed and stained cells from the root tip of the garden pea (*Pisum sativum*). A, young cell before appearance of vacuoles. B, older cell with vacuoles. (Mottier.)

Every tissue, when examined microscopically, is seen to be composed of numerous units of similar form which are in intimate relation one with another. These appear like small compartments or chambers, separated from each other by walls and have been termed **CELLS**. It will be noted that every cell possesses contents.

While these cell contents vary in character depending upon the cell examined, every living cell contains a viscid, often granular, reticular or alveolar, mobile, or jelly-like substance in which life resides. This is called **PROTOPLASM** or living matter. It is this living substance which constructs the cells or funda-

mental units of which the plant body is formed, and carries out all of the life processes of the plant.

The herbs, shrubs and trees discussed in the preceding paragraphs represent but few of the vast array of forms which

FIG. 5.—Two Blue-green Algæ. A, B, C, D, E, *Glæocapsa*; F, *Oscillatoria* showing a dead cell (*d*) which marks a place of separation into segments. (A), *Glæocapsa*, parent cell composed of central protoplast containing scattered chromatin granules, surrounded by cell wall and 3 mucilaginous envelopes; (B), parent cell is shown elongated, the protoplast in process of division to form two daughter protoplasts; (C), daughter protoplasts, each surrounded by two gelatinous envelopes and both within the original parent envelopes; (D) the daughter protoplasts shown in C have just divided to form granddaughter protoplasts which have later separated, each forming envelopes of its own but all four encircled by the parent envelope.

make up the Plant Kingdom. Many additional kinds of plants varying in form, habitat and method of living will be discussed in the chapters to follow, but in order that the student may approach the study of these with better understanding, it seems

necessary, at this point, to make some general introductory comment on the larger groups of plants.

GROUPS OF PLANTS.—There are four great divisions in the Plant Kingdom, the *thallophyta*, *bryophyta*, *pteridophyta* and *spermatophyta*. Each of these divisions is further divided into subdivisions, classes, orders, families, genera and species. The species are made up of individuals.

FIG. 6.—A mold, *Mucor mucedo*, showing the branched filaments or hyphæ which collectively constitute the mycelium of this fungus and the erect spore bearing hyphæ or sporangiophores. (Palladin.)

The lowest of these in scale of organization and the most ancient is the THALLOPHYTA or Thallophytes [Gk. *thallos*, a young branch; *phyton*, plant]. They have been so named because their bodies, each of which is called a *thallus*, show no differentiation into roots, stems and leaves. To this division belong two large groups commonly known as the ALGAE and FUNGI. The Algae possess *chlorophyll*, a complex mixture of green and yellow pigments, while the Fungi are chlorophyllless. The ALGAE include the green and blue-green scums which occur in stagnant pools, ditches and in running streams, the diatoms and the green, brown and red seaweeds. The FUNGI include the bacteria or

minute germs of decay and disease, the slime molds, the yeasts, molds which occur on bread, cheese, fleshy drugs, some pharmaceutical preparations, etc., the water molds, some of which attack the gills of fishes, the mildews, the mushrooms and toadstools, and many forms which cause plant diseases, such as the smuts, rusts, chestnut tree blight, ergot of rye, etc. There is another subdivision of Thallophytes called LICHENS which comprises the leathery-like growths which occur on the bark of trees, on rocks and the soil of woodland. They are peculiar plants in that each represents a living together of an alga and a fungus.

Next to the Thallophytes in order of ascending complexity of structure come the BRYOPHYTA or Bryophytes [Gk. *bryon*, moss; *phyton*, plant]. This division includes the liverworts or HEPATICÆ and the mosses or MUSCI. The liverworts are small, leaf-like plants which occur on moist earth and in water. Some of them are thought to be descendants of forms which were the first plants to take up a land existence. The mosses include pale green forms of swampy ground and bogs called peat mosses, the black mosses of siliceous rock,

and the bright green or common mosses which so conspicuously carpet portions of our forests and rocks. (See Figs. 315 and 318.)

The next highest division is that known as the PTERIDOPHYTA or Pteridophytes [Gk. *pteris*, fern; *phyton*, plant] which have as their representatives the CLUB MOSSES, HORSETAILS or Scouring Rushes and the FERNS. While most of these are low growing terrestrial plants, a few forms are aquatic in habit and some of the tropical ferns assume tree-like forms. (See Figs. 325, 329 and 330.)

The Thallophytes, Bryophytes and Pteridophytes are collectively termed the CRYPTOGAMS [Gk. *cryptos*, hidden; *gamos*,

FIG. 7.—A liverwort (*Lunularia*). Below, portions of the thalloid body, showing the lunar-shaped cupules, with brood-buds, or gemmæ. Above a single gemma, greatly magnified. (Gager.)

marriage or union], since it was formerly believed that the sexual organs of these plants were obscure. They reproduce by spores, and, for the greater number, also by sexual reproduction.

FIG. 8.—A gymnosperm, *Zamia*. Note the large cone which when mature bears naked seeds.

The most highly evolved group of plants is that of the seed plants or SPERMATOPHYTES [Gk. *sperma*, seed; *phyton*, plant] of which the more familiar herbs, bushes, vines and trees are examples. The Spermatophytes are also called the PHANEROGAMS [Gk. *phaneros*, evident; *gamos*, union], because it was formerly believed that they alone showed evident sexual organs. They are characterized by possessing flowers and producing seeds.

The SPERMATOPHYTES include two sub-divisions, namely the GYMNOSPERMS and the ANGIOSPERMS. The Gymnosperms [Gk. *gymnos*, naked; *sperma*, seed] have their seeds uncovered. They include such plants as the pines, spruces, firs, hemlocks,

FIG. 9.—A dicotyledon. The English Oak (*Quercus pedunculata*). A, twig with leaves and flowers. B, staminate flowers. C, two stamens. D, pistillate flower, enlarged. E, fruit, acorns. F, cup (cupula) with nut removed. G, seed in cross section. H, seed in longitudinal section. The perianth of the staminate flower consists of 5–7 segments; stamens 6–12. (After Wossidlo.)

redwoods, larches, yews, junipers, cedars, arbor vitæ, the cycads, ginkgos, gnetums and ephedras.

The ANGIOSPERMS [Gk. *angeion*, vessel; *sperma*, seed] have their seeds enclosed within the seed vessel or fruit wall. They include two sub-groups or classes, namely, the MONOCOTYLEDONS and

the DICOTYLEDONS. The MONOCOTYLEDONS [Gk. *monos*, alone; *kotyledon*, a cup-shaped hollow] are represented by the cat-tails, grasses, sedges, lilies, palms, bananas, gingers, orchids, etc. They have a single seed-leaf or cotyledon, mostly parallel veined leaves and the parts of their flowers are in threes.

The DICOTYLEDONS [Gk. *di*, two; *kotyledon*, a cup-shaped hollow] are represented by such forms as the willows, poplars, maples, oaks, chestnuts, geraniums, violets, cherries, roses, beans, asters, etc. They have two seed-leaves or cotyledons, netted veined leaves, and the parts of their flowers are in fours and fives.

Every seed plant normally comes from an *ovum* (egg cell) which has been fertilized within the ovule of a flower. The fertilized egg or *zygote* divides and redivides to form at first a nearly uniform mass of cells. As these cells continue to multiply a process early becomes manifest. This process is called PHYSIOLOGICAL DIVISION OF LABOR and consists of the apportioning of separate kinds of work to different parts of the same organism. The result of the changes occurring in physiological division of labor is called DIFFERENTIATION. As the cells continue to increase, unequal growth takes place among them and, in time, they become differentiated to form a young seed plant or *embryo*. The embryo lies within the ovule which by this time has ripened to form with the embryo the structure called the seed.

Upon reaching suitable soil the seed absorbs water and its embryo swells and sprouts forth or germinates, forming a young seedling plantlet with roots, green foliage leaves and stem. Eventually, through further growth and differentiation, a mature adult plant is developed which bears flowers in which female sexual cells or *eggs* and male sexual cells or *sperms* are produced. The sperm unites with the egg, fertilizing it, thus starting a new cycle. This process is continued from generation to generation.

CHAPTER III

THE LIVING CELL

THE CELL AS THE FUNDAMENTAL UNIT

Robert Hooke, an Englishman, in 1665, first described and figured cells. He observed compartments regularly arranged in rows in thin sections of cork placed under a microscope and, likening these to the rooms of monasteries, named them cells. In 1831, Robert Brown, an English botanist, discovered the nucleus which he found in a number of the plant cells.

Matthias J. Schleiden, a German botanist, in 1838, showed the cell to be the unit of plant structure. The following year Theodor Schwann, a German zoologist, showed that the bodies of animals are also composed of cell units and announced the *Cell Theory*, based upon Schleiden's earlier work and his own observations. The bodies of all plants and animals are composed of one or more of these fundamental units of structure and function. Each cell consists of a unit mass of living matter to which Hugo Von Mohl gave the name of *protoplasm* in 1851. But protoplasm had been observed (though not so called) for the first time in the cells of some plants by Corti and Treviranus as early as 1772 and in the cells of small animals by Dujardin in 1835. The latter described it as a jelly-like substance, "the primary animal substance," and called it SARCODE. The cell protoplasm may or may not have a cell wall surrounding it. Most plant cells consist of a nucleated mass of protoplasm and a cell wall surrounding it. The term PROTOPLAST is given to the mass of protoplasm found within the cavity of the cell. While most plant cells contain a nucleus and some contain a number of nuclei, the cells of the blue-green algæ and the bacteria have been found to lack definitely organized structures of this kind but rather contain chromatin within their protoplasm in a more or less diffuse or loosely aggregated condition.

A TYPICAL COLORLESS PLANT CELL

If we peel off a portion of the thin colorless skin or *epidermis* from the inner concave surface of an onion bulb scale, mount in water and examine under the microscope, we find it to be composed of a large number of similar cells which are separated from one another by means of lines, the bounding cell walls. Under high power each of these cells will exhibit the following characteristics:

An outer wall, highly refractile in nature and composed of *cellulose* which surrounds the living matter or *protoplasm* (see Fig. 10). This wall is not living itself but is formed by the living matter within the cell. It is sufficiently rigid to give definite shape to the cell and sufficiently porous to allow water with substances in solution to pass through it. Somewhere within the protoplasm will be noted a denser-looking body. This is the *nucleus*. The nucleus is called "the superintendent of the cell" since it is believed to control all the activities of the protoplast. Its removal results in the death of the cell. It is usually spherical in shape but in old epidermal cells of the onion becomes somewhat flattened and pushed up against the outer plasma membrane and cell wall. It is surrounded by a *nuclear membrane* and contains a *nuclear reticulum*, *nuclear sap* and one or more *nucleoli*. The nuclear reticulum can best be seen after the cells are fixed and stained. It consists of a thread-like net of a colorless substance called *linin* adhering to which are granules of a substance which takes the stain of a basic dye called *chromatin*. Chromatin carries the *genes* or hereditary characters from parent to offspring. The nuclear-sap is a watery solution of nourishing substances. Within the nucleus will be seen one or more smaller highly refractile and definitely circumscribed, dense *protoplasmic bodies*, the *nucleolus* or *nucleoli*. During cell division the nucleoli disappear and reappear in the two nuclei of the daughter cells when cell division is completed.

The protoplasm of the cell outside of the nucleus is called the "*cytoplasm*." It will be seen to be clear and granular-looking. Within the cytoplasm will be observed a number of clear spaces. These are *vacuoles* and, because they are filled with *cell sap* (water

with nutrient substances in solution), are called "*sap-vacuoles*." The vacuoles appear as tiny droplets in young cells but grow with the cell, becoming larger as the cell increases in size, then fusing with each other until, in a fully mature old cell, one large vacuole is present in the center of the cell with protoplasm and nucleus pushed up against the cell wall.

FIG. 10.—Portion of the inner epidermis of an Onion bulb scale, showing cells at various stages of maturity. Young cell (1); old cell (3); cell intermediate in age between 1 and 3 (2); cell wall (*c*); outer plasma membrane (*pl*); middle lamella (*ml*); nucleus (*n*¹); nucleolus (*n*²); nuclear membrane (*n*³); cytoplasm (*cy*); vacuole (*v*). Note that the young cell (1) shows numerous small vacuoles and a spheroidal nucleus near center of cytoplasm. In 2 (cell of intermediate age) the cell has enlarged, larger vacuoles have formed through the bursting of films of cytoplasm separating smaller ones, and the nucleus has moved toward the cell wall. In 3, the films have all burst, the cytoplasm and nucleus have been pushed up against the cell wall, the nucleus is flattened out, and a large vacuole appears in the center of the cell.

Protoplasm is in intimate relation to water. The reaction of the cytoplasm to a bounding film of water between it and the cell wall forms the *outer plasma membrane* or *ectoplasm*, a clear, homogeneous, outer band of cytoplasm; the reaction of cytoplasm to the water within the sap vacuoles forms the *vacuolar membranes*; the reaction of the dense protoplasm of the nucleus to the water in the cytoplasm around it forms the *nuclear membrane*. All of these membranes are bounding films of protoplasm belonging to the region in contact with water and are collectively termed *plasma membranes*. Upon mounting another portion of epidermis in iodine solution, removing the excess of stain and adding a drop of sulfuric acid and then examining under high power, we note

that the cell walls of cellulose are stained a deep blue. A yellow line is evident in the middle of each cell wall and separates each cell from its bounding cells. This line represents the fused

FIG. 11.—Elodea, an aquatic plant.

primary walls of adjacent cells and is called the *middle lamella*. It is composed largely of *calcium pectate*.

TYPICAL GREEN CELLS

Excellent material for the microscopic study of green cells of plants is to be found in the leaf of *Elodea*, a fresh water plant, found in ponds and house aquaria. The leaves of this plant are small enough to be mounted beneath an ordinary cover glass on a microscopic slide. They are oblong to oblong-ovate and occur in whorls of 2 to 4 along a slender, elongated stem.

When one of these is detached, mounted in water and examined under low power magnification, it is seen to be two layers of cells thick, except along the margin, where only one layer of cells is to be observed and in the central midrib region, where there occur several layers of narrow cylindrical cells. The

FIG. 12.—Portion of the leaf of *Elodea*, magnified. *v*, midrib; *t*, pointed cells projecting at intervals from the margin.

cells are arranged in rows and vary in size, being shorter and broader toward the midrib and longer and narrower near the margin. Projecting tooth-like cells occur along the margin. The cells of the upper surface appear about twice as broad as those of the lower surface. Each cell is separated from its neighbors by means of transparent *cell walls* of cellulose and pectic substances. Just as in the onion epidermis, what appears to be a single partition wall between two cells is really two cell walls separated by a *middle lamella*.

If some of the larger cells are now examined under high power magnification, each is found to be brick-shaped. Beneath the cell wall will be seen a narrow zone of *cytoplasm* containing numerous, green, ovoid, living bodies, the *chloroplasts*. Each chloroplast is a dense, porous, mass of protoplasm containing the green pigment or rather a mixture of four pigments two green and two yellow and collectively called *chlorophyll*. Chlorophyll is a highly complex and important material. By means of it, energy in the form of light from the sun is absorbed and utilized by the chloroplast for the manufacture of sugar. In some of the

FIG. 13.—A typical green cell of *Elodea*. Cy, cytoplasm; N, nucleus; N', nucleolus; C, chloroplast; Pm, outer plasma membrane; VM, vacuolar membrane; V, sap vacuole; W, cell wall; S, intercellular-air-space. Greatly magnified.

cells the protoplasm will be found to be in motion (*rotation*), carrying the chloroplast along with it. The *nucleus* will be observed as a clear, dense, somewhat hemispherical shaped body embedded in the cytoplasm and surrounded by the *nuclear membrane* which separates it from the cytoplasm. A *nucleolus* will be seen within the nucleus. The center of the cell is occupied by a large *sap vacuole* containing a watery solution of salts and other nourishing substances. A clear *vacuolar membrane* lines the inner surface of the cytoplasm, while another clear membrane, the *outer plasma membrane*, lines the outer surface of the cytoplasm.

If the leaf be boiled in water so as to kill the plasma membranes and make them permeable and then mounted in dilute iodine solution and examined under the microscope, bluish to bluish-black bodies will be observed in the chloroplasts. These

are starch granules. They have been formed in the chloroplast through the agency of chlorophyll and sunlight by the process called *photosynthesis*. This process is discussed more fully in another part of the book.

THE CELL WALL

Nearly all plant cells possess cell walls. The cell wall is composed of non-living substances which are secreted by the

FIG. 14.—Primary, secondary and tertiary walls. A, B, longitudinal and transverse sections of trachea of Linden (*Tilia americana*). C, D, longitudinal and transverse sections of tracheid of a yew (*Taxus brevifolia*). (After Eames and MacDaniels, *Introduction to Plant Anatomy*; McGraw-Hill Book Co. publishers.)

protoplasm. The youngest meristematic cells in the growing apical regions of roots, stems, and leaves are separated from each other by delicate *primary cell walls* made up of carbohydrate material called *pectose*. As these cells grow older each of their protoplasts secretes an individual *secondary cell wall* composed of *cellulose* and some *pectose*. So what often appears to be a single wall separating two adjacent cells consists of the primary wall between 2 layers of secondary wall. The primary wall has

undergone some changes due to the infiltration or addition of other substances and now becomes known as the *middle lamella*. At first the secondary walls are thin but become thickened in many cells owing to the deposition of successive layers of cell wall substances through the intercalation of new particles of cell wall materials between those already present. Some cells like tracheae and tracheids also possess a *tertiary wall* which is deposited by the protoplast in the form of spiral or annular bands on the inner face of the secondary wall.

Thick cell walls are said to be stratified, since they show lamellæ or layers which vary in appearance under the microscope due to the presence of different quantities of water or to their chemical composition. They may also show, when examined in surface view and especially after special staining, fine lines running spirally or obliquely in the wall.

Other substances besides cellulose and pectose may be present in mature cell walls. These include *lignin* found in woody walls, the water-proof substances *cutin* and *suberin* common to walls of protective tissues, and mucilage found in the walls of the epidermis of some seeds. A further consideration of the cell wall will be taken up in Chapter V.

THE AMŒBA

Amœba provides excellent material for the study of protoplasm. It is a microscopic organism rarely more than $\frac{1}{50}$ in. in diameter, classified by most biologists within the phylum, *Protozoa*, a group of unicellular animals. It is found in stagnant water almost universally, where it adheres to weeds, dead leaves, or other submerged objects.

If a drop of Amœba culture be placed on the slide, covered with a cover-glass, and the mount searched under low power, an Amœba should be found.

The beginner in microscopy is very likely to overlook an Amœba altogether, or to think that every little irregular, clear spot in the field of view is one. When a clear object is seen which is never the same in outline in consecutive moments, one is certain of viewing Amœba.

Move the slide if necessary so as to bring the organism into the center of the field of view. Swing on the clips and bring the high-power objective into position. Focus.

Note that *Amœba* is a one-celled organism devoid of a cell wall, hence a naked mass of *protoplasm*. *Protoplasm* is a semifluid, viscid, granular like substance, in which life resides. It is shown by chemical analysis to consist mainly of certain substances known as proteids, bodies of extreme complexity in chemical constitution, examples of which are white of egg or the albumen of blood serum. Besides proteids, the protoplasm of

FIG. 15.—The *Amœba*. A, An *amœba* moving in the direction of the arrows; *n*, nucleus; *v*, contractile vacuole; *f*, food within food vacuole. B, the process of fission in *Amœba*, a nucleus in each half. B, after Jordan, Kellogg and Heath. (From Bergen and Davis *Principles of Botany*.)

Amœba contains fats and small proportions of mineral matters such as salts of potassium, calcium, magnesium and iron. It also contains a large proportion of water, called *water of organization*.

Protoplasm is caused to shink by strong alcohol, which withdraws the water of organization. It is coagulated by heat and dissolved by prolonged treatment with weak acids or alkalies.

Note that the protoplasm of the *Amœba* consists of an outer non-granular portion called the *ectoplasm* and an inner portion called the *endoplasm*. Within the endoplasm is to be found the *nucleus* which in the *Amœba* is difficult to discern without the use of special stains.

Observe the Amœba as it moves and find out whether the direction of movement is in one line or several.

Note that Amœba moves by extending protoplasmic processes called *pseudopodia* (false feet). At a particular point the ectoplasm is pushed out in the form of a small pimple-like elevation; This increases in size, still consisting of ectoplasm only until, at last, granules from the endoplasm stream into it and the projection or pseudopod comes to possess the same structure as the rest of the Amœba. This method of movement is called *amœboid movement*. The same kind of protoplasmic movement is shown by the slime molds and white blood corpuscles.

Note that when the organism comes in contact with a diatom or any other particle of food pseudopodia are extruded and the food is enveloped, so sinking into the cytoplasm. The food is taken in with a small amount of water, which forms a surrounding bubble or *food vacuole*. This constitutes the first step in the process of *nutrition* and is called *ingestion*. If watched sufficiently long, the protoplasm of the diatom or substance of the food particle ingested is slowly dissolved or digested through the agency of some enzyme or enzymes within the endoplasm. As it moves on, the Amœba ejects or expels the cell wall of the diatom or any other insoluble substance.

An *enzyme* is a substance of biologic origin which has the property of causing a change in its surrounding medium without becoming involved itself. Through enzyme action the previously water insoluble (colloidal) protoplasm or food of the Amœba is broken down into water-soluble (crystalloidal) substances which are capable of diffusion. The process whereby water-insoluble foodstuffs are made water-soluble through enzymic action is called *digestion*.

Mineral matters are taken in by the Amœba with its food in the form of a weak, watery solution. The next step is for the organism to rearrange the elements of digested food, mineral salts and water in such a way as to form new particles of living protoplasm. These are deposited among the pre-existing particles and so *assimilated* or converted into the living substance of the Amœba. One effect of this formation of new protoplasm is *growth*. This process of growth by the interposition of new

particles among the old ones is called *intussusception*. All living things, both animals and plants, grow or increase in size by this method.

Minerals or other organized non-living things grow by a deposition of successive layers on their original surfaces. This process of growth is called *accretion*.

FIG. 16.—Stages in the division of *Amœba polypodia*. (From Marshall, after Schulze and Lang.)

Look for a large, clear, pulsating structure or space within the protoplasm of the Amœba. This is the *contractile vacuole*.

Alike with all other living organisms, Amœba is continually undergoing destructive as well as constructive changes. Every time it extrudes pseudopodia or performs work of any kind, a portion of its protoplasm is broken down by oxidation, which is accompanied by a release of energy. The nitrogenous waste in solution, resulting from the oxidation of protoplasm, makes its way to the contractile vacuole and is expelled by its contraction. This function of getting rid of nitrogenous waste is called *excretion*.

In order that the oxidation of protoplasm, referred to in the preceding paragraph, may take place, a constant supply of oxygen is essential. The water in which *Amœba* lives contains this gas in solution. Now, the protoplasm of the *Amœba* (as well as other living organisms) is continually forming carbon dioxide. When two gases are separated from one another by a porous septum, an interchange takes place between them, each diffusing into the space occupied by the other. The process of gaseous interchange is precisely what takes place between the H_2O in the surrounding water and the CO_2 in the interior of the *Amœba*, the protoplasm acting as a porous partition. This gaseous interchange whereby this organism takes in oxygen and gives carbon dioxide off is called *respiration* or *breathing*. The oxygen taken in by the *Amœba* oxidizes the complex carbon compounds within its body with an accompanying release of energy.

In order that the race of *Amœba* may not be exterminated, there must be some provision whereby its numerous casualties may be balanced. In other words, *Amœba* must produce new individuals like itself. This function is called *reproduction*.

Amœba reproduces itself by the simplest method of reproduction, known as *simple division* or *binary fission*. At first the nucleus divides into two; then the whole organism elongates, the two nuclei at the same time traveling away from one another; next a furrow appears across the middle of the elongated body between the nuclei; the furrow deepens until finally the organism separates into two separate *Amœbæ*, which henceforth lead an independent existence.

Under unfavorable circumstances, such as the drying up of the water in pools in which it exists and the approach of cold weather, the *Amœba* draws in its pseudopodia, loses most of its water content, becomes spherical in shape and develops on its outer surface a thick resistant wall called a *cyst*. Inside of the cyst the nucleus undergoes division to form a large number of nuclei. Cells walls then appear dividing up the cell contents into as many little *Amœbas* called *Amœbulæ* as there are nuclei. Upon the approach of favorable conditions, these *Amœbulæ* escape by rupture of the cyst, each developing into an active

Amoeba. This process is called *sporulation* and represents another method of reproduction shown by some species of the Amoeba.

THE RELATION OF PLANTS AND ANIMALS

One can see many striking differences between the highest forms of plants such as the sunflower and orchid and the highest forms of animals such as man and the gorilla, but as one passes to the lower forms of plants and animals these differences gradually disappear. There are a number of low types of organisms such as *Euglena* and *Volvox* which have characteristics common to both plants and animals and which might be placed in either group.

PLANTS

ANIMALS

1. Most plants possess the power of constructing complex organic food substances like sugar and starch from water and carbon dioxide by means of chlorophyll and light.	1. Most animals cannot construct their own foods but are dependent upon green plants for the food essential to their nutrition and existence.
2. The cells of most plants possess rigid cell walls composed of cellulose and pectose which surround their protoplasts.	2. The cells of most animals are devoid of such walls.
3. Most plants are incapable of moving from place to place (locomotion).	3. Most animals are capable of locomotion.
4. In most of the simple plants growth is not localized but in the more complex plants growth in length becomes localized in the tip regions of organs and in these continues indefinitely.	4. In most animals growth in length is not restricted to the ends of body or to the extremities of organs and ceases when the animal attains maturity.
5. Most plants are constructive agents.	5. Most animals are destructive agents.

Still more primitive and less differentiated are the *Amoeba*, the vegetative stages or plasmodia of slime moulds, and the bacteria which, along with many other unicellular organisms of simple structure, have been called "*Protista*." It is the belief of most biologists that plants and animals of the present day have gradually developed from a common origin which, during an early era in the earth's history, gave rise to the *Protista* from which arose two diverging branches of organisms known as the plant and animal kingdoms.

It is impossible to draw sharp lines of distinction which will separate all plants from all animals owing to the similarity of structures, functions and laws of inheritance found in organisms belonging to both groups. It is possible, however, to find points of difference between most plants and most animals. These are given in the table on page 29.

PROTOPLASM AND ITS PROPERTIES

Protoplasm, or living matter, is the viscous cell substance in which life resides. Huxley has defined it as the "physical basis of life."

Protoplasm occurs as a colloidal solution and as an emulsion of colloidal particles. It has well been called a "*polyphase colloidal system*." *Colloids* are substances like glue, agar or gelatin which do not crystallize when solidifying from solution and which will not pass through a semipermeable membrane like pig's bladder or parchment. Substances like salt and sugar which crystallize from a solution upon evaporation of the solvent and which pass through a semipermeable membrane are called crystalloids. When examined under the microscope, protoplasm appears either as a colorless jelly in which are frequently imbedded globules and granules of various shapes and sizes or as a streaming liquid. It occurs both as a *sol* or liquid that flows readily or as a *gel* or jelly-like mass. It is extremely complex and unstable. It may change from the liquid to the jelly state or vice versa under certain conditions. The liquid portion of the protoplasm contains particles of ultramicroscopic size while the jelly-like, granular portion contains particles, granules or globules visible under a compound microscope.

Protoplasm cannot be analyzed chemically. Dead protoplasm, which is the material examined when protoplasm is subjected to reagents in the procedure of chemical analysis, consists of a variety of different chemical compounds including proteins, carbohydrates, fats, salts of sodium, potassium, magnesium, calcium and sometimes iron, etc., also gases like oxygen and carbon dioxide and water.

PROPERTIES OF PROTOPLASM.—The peculiar properties which distinguish protoplasm from non-living matter are as follows:

1. STRUCTURE.—Protoplasm, as it occurs in cells of plants and animals, invariably exhibits structure. No region of it, however small, has been found to be homogeneous. Each advance in microscopical technique reveals new complexities. The protoplasm of a single cell, far from being a single unit, must rather be looked upon as a microcosm.

2. METABOLISM.—Perhaps the most significant peculiarity of living matter is found in its instability and the chemical changes which continually go on within it. It is constantly wasting away, and as constantly being built up. These losses and gains are not upon the exterior surface but throughout its mass. Its growth and renewal are by *intussusception*, or the taking in of new particles and storing them between those already present. Inorganic substances, such as crystals on the other hand, grow by *accretion*, *e.g.* through the addition of similar material in layers on their outer surfaces. A bit of protoplasm may retain its identity while all the matter of which it is composed is changed over and over. It is like a whirlpool or wave in a river which remains the same while the water of which it is composed changes continually. Metabolism has been aptly defined by Huxley as the whirlpool character of the organism. It may also be defined as the series of processes concerned with the building up and breaking down of living matter. To those processes concerned with the building up of protoplasm, such as absorption, photosynthesis, digestion and assimilation, the term *anabolism* is applied. Another term for anabolism is *nutrition*. To those processes which involve the breaking down of protoplasm, such as respiration, excretion and secretion, the term *katabolism* has been given.

3. IRRITABILITY.—This is the property possessed by protoplasm of responding to a stimulus. All living matter responds

to stimulation. When matter fails to be irritable or responsive to stimuli, we declare it to be dead. The stimuli that excite reactions in living matter are of two kinds, viz., intrinsic and extrinsic.

Intrinsic stimuli are inherited stimuli, being carried by the chromosomes from parent to offspring. They determine that the plant shall conform to a particular type, carry on certain activities, pass through a definite life cycle, and detach a portion of its own substance for the formation of new individuals of its kind.

Extrinsic stimuli initiate, inhibit, accelerate or modify the effects of intrinsic stimuli. They comprise agents of the external world such as cold, heat, chemicals, food, water, light, oxygen, electricity, gravity, etc.

TROPISMS, TAXIES AND NASTIES.—The irritable reactions manifested by protoplasm and living things to the effects of these external agents are termed *tropisms*, *taxies* or *nasties* and will now be considered briefly. The term *tropism* is now largely limited to the response of non-motile plants and is usually applied to growth movements of these in response to external stimuli. The response made to a stimulus is usually designated by a prefix name which implies the kind of stimulus, as for example, the response to light is called *phototropism*, to gravity, *geotropism*, etc. A response in the direction of the source of a stimulus is called *positive* while one which is away from the source of the stimulus is designated as *negative*. The term *taxy* (pl. *taxies*) is applied to responses of motile plants and plant cells, as gametes and zoöspores, which involve locomotion. The term *nasty* (pl. *nasties*) is applied to responses of non-motile plants induced by stimuli in which the movements have no particular direction, as for example movements brought about in some flowers as a result of the succession of night and day (*nyctinasty*).

THERMOTROPISM is the response of living substance to the stimulus of temperature. All living substance is influenced by variations in temperature. Freezing disintegrates it while excessive heat causes its coagulation. Active vital phenomena are therefore only evident within these extremes, the limits differing depending upon the endurance of the organism under

examination. The lowest temperature at which the activity of an organism becomes evident is known as the *minimum*, that at which the activities are at their best, the *optimum*, and the highest at which they can be continued, the *maximum*. Some plants are able to endure greater extremes of temperature variation than others because of special adaptations. Thus, certain bacteria produce spores which resist exposure for an hour to the temperature of liquid hydrogen ($-225^{\circ}\text{C}.$) or to that of a hot air oven at $100^{\circ}\text{C}.$ Many higher plants can endure moderately low temperatures by the development of a hairy covering; others which are killed by frost produce seeds which can endure rigid cold, still others adapt themselves to existence through periods of cold by passing through a latent stage in the form of bulbs, like the Squill or the Lily, or rhizomes, as the Blood Root or the Hellebores.

CHEMOTROPISM (or *chemotaxy*) is the response of protoplasm to chemical stimulation. Any substances that possess the property of producing a deleterious effect upon protoplasm are termed *poisons*. Poisons may effect an immediate destructive combination with living substance when they are called *caustics*, or they may have an exciting or depressing effect which may eventually prove destructive without visible structural change, when they are termed *toxins*. Caustics may liquefy the protoplasm, as the alkalies, or coagulate it, as the acids or salts of metals. When well diluted, chemicals may occasion no destructive effects, but may call forth positive or negative responses, known as positive or negative chemotropism or chemotaxis.

Thus, Pfeffer, working with the motile sperms of ferns, found that if a capillary tube, containing a solution of malic acid, be introduced into water containing them, the sperms moved toward it and entered. It is now generally believed that the motile male sexual cells of all flowerless plants are attracted to the appropriate female sexual cells by means of positive chemotaxic influences. Among flowering plants, it has been observed that pollen grains brought by various agencies from anthers to stigmas of certain plants of different species will not germinate, but when they are carried from one plant to another of the same species or variety, they readily send their pollen tubes through

the stigma and style to the ovule below. In the former instance, negative chemotropism is illustrated, while in the latter, positive chemotropism is shown.

SITOTROPISM is the reaction of living matter to the influence of food. When the response is by motile organisms or motile cells, it is called *sitotaxy*. Hertwig found that if a fine capillary tube be filled with a 1 per cent solution of asparagin or beef extract and held in contact with a drop of water containing certain bacteria, a mass of these soon plugged the mouth of the tube. His experiment shows that these organisms moved from a poorer to a richer nutrient medium in response to a positive sitotaxic influence.

OXYTROPISM is the response to the stimulating influence of oxygen. We see evidence of this everywhere in nature. No living thing can continue to exist without this element. A mistaken idea is often prevalent regarding obligate anaërobic bacteria. Like all other bacteria or organisms, these plants require oxygen but can only assimilate it in its combined form. The tetanus bacillus is a good example. Aërobic bacteria, on the other hand, require free (uncombined) oxygen for assimilative purposes. Thus the tetanus organism (an *anaërobe*) grows in the depth of culture media, whereas the tubercle bacterium (an *aërobe*) grows only on the surface.

HYDROTROPISM is the response of protoplasm to the stimulus of water. This reaction is seen in both positive and negative phases in the slime molds. The vegetative stage of these lowly plants is characterized by a naked, many nucleated mass of protoplasm, confining itself to the moist crevices of rotten logs, etc. until the surface of the substratum becomes wet, when and only when it will emerge (positive reaction). As soon, however, as its fruiting stage begins, the whole protoplasmic mass wells up from the substratum, away from moisture (negative reaction). The roots of young seedling plants show positive hydrotropism by growing toward moisture in the soil.

RHEOTROPISM is the response made by roots and slime moulds to currents of water.

HELIOTROPISM is the response of living substance to the stimulus of sunlight.

PHOTOTROPISM is the response of living things to light of any source. The stems of higher plants tend to grow toward the light and are, therefore, positively phototropic, whereas the roots grow away from the source of light and are so negatively phototropic.

FIG. 17.—Venus fly-traps, *Dionaea muscipula*, growing in field near Wilmington, N. C.
(Photograph by Steckbeck and Palmer.)

GEOTROPISM is the response of protoplasm to the stimulus of gravity. Roots of Pteridophytes and seed plants invariably grow downward toward the center of gravity and so are positively geotropic. The fruiting organs of the fungi and the main stems of higher plants tend to grow perpendicular to the earth's

surface and so are negatively geotropic. Branches of stems that assume a relation parallel to the earth's surface are *diageotropic*. The Lima Bean, Sarsaparilla, Poison Ivy, and other plants whose stems twine about supports exhibit *lateral geotropism* in their horizontal curvatures.

GALVANOTROPISM is the reaction of protoplasm to electrical stimuli. In this connection it may be said that the degree of response bears a definite relation to the intensity of the stimulus. No visible external electrotropic reactions have been observed in higher plants, although when their cells are examined microscopically, the reaction becomes manifest. Kühne has shown that when an electric current is passed through the hairs of the Spiderwort, the cytoplasm becomes gathered into small globular masses.

THIGMOTROPISM is the response of living matter to mechanical stimulation. Examples of this form of irritability appear to be far less common among plants than among animals. Certain species of *Mimosa* (Sensitive Plants), *Oxalis*, *Drosera* (Sundew), *Desmodium* and *Dionaea muscipula* or the Venus Fly-trap exhibit this phenomenon to a marked degree. A few instances only will be considered. When the tendrils of climbing plants come into contact with the uneven surface of solid bodies they are induced to coil. When the tentacles on a modified leaf of the Sundew (*Drosera*) are stimulated mechanically by an insect or artificially they are induced to curve over. If a good plant of the Venus Fly-trap (*Dionaea*) is selected, it will be seen to possess leaves, the terminal portions of which are modified as traps for catching insects (Fig. 17). Hairs will be seen projecting from the upper surface of each valve of the hinged blade. If one of these hairs is touched with a pencil no reaction will be evident but if, after a lapse of twenty seconds, the hair is touched again, the 2 valves close. If the stamens of *Berberis* (Barberry) be touched near the base during their pollen shedding stage they will be observed to curve toward the stigma.

The most highly specialized form of thigmotropism observed in plants appears to be found in *Mimosa Spegazzini*, a member of the Bean family. According to Steckbeck "when a mechanical stimulus, such as a forceps pinch, is applied to one of the terminal

secondary leaflets, after a latent period of less than $\frac{1}{4}$ second, the leaflet stimulated rises and its partner almost at the same time.

FIG. 18.—Leaves of *Mimosa pudica*. *A*, normal position; *B*, after stimulation. *p*, pulvinus or leaf cushion at base of petiole. Note the compound leaf consists of a long petiole ending in a blade of 4 leaflets, each of which possesses a secondary petiole with a pulvinus at its base, and a midrib which bears a large number of small secondary leaflets. A stimulus applied to the large pulvinus at the base of the main petiole will cause the latter to fall and the secondary leaflets to become erect with the upper surfaces of each pair of opposite leaflets against each other (*B*). (After Pfeffer.)

The stimulus is then carried down the midrib, the pairs of secondary leaflets closing in order; in 9 seconds all the secondary leaflets have closed, the midribs converge, followed in 3 seconds

by a drop of the entire leaf. The stimulus moves up the other leaflet with the result that the secondary leaflets close in order. In 20 seconds after the stimulus has been applied all of the secondary leaflets, are closed. The stimulus is propagated through the stem to other leaves."¹ (Cf. Figs. 18 and 19.)

FIG. 19.—*Mimosa Spegazzini*. After the application of a stimulus. Compare with Fig. 18. (After Steckbeck.)

4. REPRODUCTION.—Protoplasm also shows a very remarkable ability to increase and to give off detached portions which retain the infinitely complex peculiarities and properties of the original. The process, moreover, may be continued indefinitely. These detached portions may be of the nature of sexual cells which unite to form a cell called a *zygote* or *oöspore* from which a mature individual develops or of the nature of asexual cells detached singly or enmasse which eventually develop into mature individuals.

¹ "The comparative histology and irritability of sensitive plants" by D. W. Steckbeck in Contributions from The Botanical Laboratory of the U. of Pa., vol. IV, No. 2, p. 217, 1919.

Other physiological characteristics, as motility and conductivity, might be added, but the four just discussed are the most satisfactory criteria by which living may be distinguished from non-living matter.

PROTOPLASMIC MOVEMENT

Living matter in its active state of existence exhibits motion. The several kinds of movement of protoplasm which may be observed within the closed cells or in naked cells are *amoeboid movement*, *ciliary movement*, *cytoplasmic rotation* and *cytoplasmic circulation*. Cold retards and heat accelerates movement within certain ranges.

1. *Amoeboid movement* is a creeping or flowing movement such as is typical of the plasmodium or vegetative body of slime molds and of the animal *Amoeba*.

2. *Ciliary movement* can be observed in the motile bacteria and in the sex cells of the lower and some of the higher plants which possess slender protoplasmic outgrowths called *cilia* which lash the water.

3. *Cytoplasmic rotation* is well illustrated in *Elodea* and *Nitella*, two common aquarium plants. In rotation, the protoplasm moves down one side of the cell then across the end and returning along the other side and end. In *Elodea* the chloroplasts found

FIG. 20.—A, embryonic cells from onion root tip; *d*, outer plasma membrane; *c*, cytoplasm; *a*, nuclear membrane enclosing the thread-like nuclear reticulum *b*, nucleolus; *e*, plastids (black dots scattered about). B, older cells farther back from the root tip. The cytoplasm is becoming vacuolate; *f*, vacuole. C, a cell from the epidermis of the mid-rib of *Tradescantia zebrina*, in its natural condition on the right, and plasmolyzed by a salt solution on the left; *g*, space left by the recedence of the cytoplasm from the wall; the plasma membrane can now be seen as a delicate membrane bounding the shrunken protoplast. All highly magnified. (Stevens.)

in the inner layer of cytoplasm are carried along by the moving cytoplasm. In some plants like *Nitella* the nucleus is carried along with the current. (See Fig. 21.)

4. *Cytoplasmic circulation* or *streaming* may be seen in the cells of the stamen hairs of the Spiderwort, in some epidermal cells of the onion and in the epidermal hairs of the squash. Here the cytoplasm will be seen in several strands traversing the cell cavity as well as lining the inner face of the cell wall. The

FIG. 21.—Protoplasmic movement. 1 and 2, nuclei from the tapetal plasmodium of the spore case of *Botrychium*, 1, in the usual form, and 2, flattened while entering a crevice between spore-groups; 3, cell from *Nitella* showing cytoplasmic rotation as indicated by the arrows; 4, cell from stamen-hair of the Spiderwort (*Tradescantia*), showing circulation of the cytoplasm as indicated by the arrows. (After Stevens.)

cytoplasm moves to and from the nucleus and cell wall and may after streaming in one direction for awhile, reverse itself.

PROTOPLASMIC CELL CONTENTS

The name "PROTOPLAST" has been given to the entire living or protoplasmic matter within a single cell. The protoplast consists of six well-differentiated portions, viz.:

(a) THE CYTOPLASM, or the foamy, often granular matrix of protoplasm outside of the nucleus. Embedded in the cytoplasm are non-protoplasmic bodies called *inclusions*. The most prominent of these are the *vacuoles* which are spaces within the cytoplasm containing either a watery solution of nourishing materials (*cell sap*) or, in old cells, air.

(b) THE NUCLEUS, a denser, definitely circumscribed region of protoplasm containing nucleoplasm which consists of gels of varying consistency containing *chromatin*, a substance staining heavily with certain basic dyes, adhering in the form of granules to a network of *linin*. NUCLEAR SAP, consisting of a watery solution of nourishing substance is also found in the nucleus. The nucleus may be regarded as the superintendent of the cell

FIG. 22.—Diagram of Plant Cells: *a*, cell wall; *b*, cytoplasm; *c*, cytoplasmic strand; *d*, nucleus; *e*, plastid; *f*, plasmodesma, passing through pit in cell wall; *g*, sap vacuole. (Drawing by H. McCarthy, from Stanford's "Economic Plants," D. Appleton-Century Co. Publishers.)

since it controls the structural changes and physiological activities that take place within the living cell. It is usually spherical in shape but in some plant cells is ovate, ellipsoidal, stellate, discoidal or elongated. The chromatin is the substance which carries those characters or factors from parent to offspring which determine that the offspring shall resemble its parent or parents and past ancestors.

(c) THE NUCLEOLUS, a small body of dense protoplasm within the nucleus. Sometimes two or more nucleoli may occur within a nucleus. The nucleoli are believed to represent reserve food which is employed in the metabolic processes of the nucleus.

(d) THE PLASTIDS, composed of differentiated protoplasm called plastic plasm. They are small, discoidal, spheroidal, ellipsoidal, stellate, ribbon-shaped, or variously shaped, dense, porous, protoplasmic bodies scattered about in the cytoplasm. The plastids may become modified and specialized for definite functions depending upon the position of the cells containing them in the plant and the external conditions to which the cells are exposed. Sometimes, as in the cells of lower plants like the *Spirogyra*, plastids are large and are then called CHROMATOPHORES.

In the meristem cells of root and stem tips of plants the plastids are very small, colorless and usually found around the nucleus. These minute plastids are termed *primordia* or *proplastids*. As the meristematic cells enlarge and undergo differentiation, the proplastids increase in size and develop into *leucoplastids*, *chromoplastids* and *chloroplastids*.

According to the position of the cells in which plastids occur and the work they perform, they differ in color, viz.:

LEUCOPLASTIDS are colorless plastids found in the underground portions of a plant and also in seeds, and other regions given up to the storage of starch. Their function is to build up reserve foods like reserve starch and oils from sugar and other carbohydrates. Leucoplastids may become transformed into chloroplastids upon exposure of cells containing them to light through the formation within them of chlorophyll and related pigments. They are only evident after properly fixing and staining cells containing them. When leucoplastids build up and store starch they are sometimes called *amyloplastids*; when they build up and store oils they are called *elaioplastids*.

CHLOROPLASTIDS are plastids found in cells exposed to light and contain the green pigments *chlorophyll a* and *chlorophyll b* together with two yellow pigments known as *carotin* and *xanthophyll*. All green parts of plants contain these bodies. Chloroplastids may develop into chromoplastids, as in the ripening of many green fruits. The fully developed chloroplastid is capable of division by constriction.

CHROMOPLASTIDS are plastids found in cells independent of their relation to light or darkness and contain a yellow, orange

or red pigment called *chromophyll*. The chromoplastids may be circular, oval, needle-shaped, angular or fan-shaped and are averagely smaller than chloroplastids. They are responsible for the yellow and orange colors of many petals and sepals of flowers and of ripened fruits. The root of the carrot contains large numbers of them.

(e) CELL MEMBRANES.—These are of three kinds, viz.: (1) *The outer plasma membrane or ectoplasm*, found as a transparent, colorless, outer bounding film of cytoplasm enveloping the

FIG. 23.—A, cell from the epidermis of the upper surface of the calyx of the Nasturtium (*Tropaeolum majus*) with crystalline chromoplasts; B, cells from the petal of *Lupinus luteus*, with yellow chromoplasts; C, cell showing numerous chloroplasts scattered through the cytoplasm. A, after Strasburger; B, after Frank. (Stevens.)

granular, inner cytoplasm, the latter sometimes termed “*endoplasm*”; (2) *the vacuolar membranes*, clear bounding films of cytoplasm surrounding sap vacuoles; and (3) *the nuclear membrane*, a clear, transparent film of protoplasm forming the outer bounding zone of the nucleus. All of these membranes are *semipermeable membranes*, since, in their living condition, they will permit the passage through them of water and certain crystalloidal (water-soluble) substances in solution while they will not admit the passage of other crystalloidal substances in solution.

(f) CHONDRIOSOMES or MITOCHONDRIA.—These are small granular or rod-shaped protoplasmic bodies found scattered in

the cytoplasm of many cells. They take a deep stain with basic dyes, like chromatin, but their significance is not understood.

PLASMODESMA.—The protoplasts of adjacent cells in many plant organs especially those possessing thick cell walls are connected by delicate threads of cytoplasm which pass through tiny holes (*pits*) in the walls between the cells. These protoplasmic connections are called *plasmodesma*. They are probably more common between cells with non-lignified walls than is generally realized owing to their frequent extreme fineness and the need of a special method of fixing and staining of preparations to bring them out. They occur in meristem, medullary rays, storage tissue, phloem, etc. The author has found large and prominent plasmodesma in the endosperm of species of *Plantago* yielding *Psyllium* seeds. They have been found in the red algæ, in stems of mistletoe and pine, in nearly every part of the mosses, ferns, and gymnosperms, especially in the young condition of the cells, and in the thick walled endosperm of seeds of *nux vomica*, *ignatia*, date, persimmon, etc. (See Fig. 22.)

CELL FORMATION AND REPRODUCTION

The cells of plants have all been derived from preëxisting cells. In the bacteria and many other low forms of plant life, the division of the cell always results in reproduction; in higher forms, however, it merely increases the size of the individual and so is a phenomenon of growth.

There are two kinds of cells formed by plants, viz.: *asexual* and *sexual*. Both of these are endowed with the possibilities of reproduction, although the former are frequently limited to the process of growth.

Reproduction is the power possessed by an organism of giving rise to new individuals. This may take place through the agency of either asexual or sexual cells and is accordingly asexual or sexual in character. Whenever a union of cells or their protoplasmic contents takes place the process is called "*sexual reproduction*"; if, however, there is a mere separation of a cell or cells from an individual which later form a new organism, the process is termed "*asexual or vegetative reproduction*."

There are five chief modes of ASEXUAL REPRODUCTION, viz.: *Fission* or *Binary Division*, *Gemmation*, *Free Cell Formation*, *Rejuvenescence* and *Vegetative Multiplication*.

FISSION.—This is the separation of a cell into two equal halves, each of which may grow to the size of the original parent cell from which it was derived. Fission is seen in the reproduction of bacteria, growth of many algæ and the formation of tissues of higher plants.

In *fission* or *binary division*, as the process is frequently called, there may occur (1) a division of the cell by means of a cell plate of the nature of a plasma membrane formed by the shrinkage of the spindle fibers to the equator of the cell during mitosis. This cell plate splits, forming two plasma membranes, between which the cell walls separating the daughter cell are laid down, or (2) a division by constriction. In the latter method, as seen in some of the algæ, the cytoplasm becomes constricted or pushed away from the wall of the cell at a point midway between the ends of the cell and at this constriction the cytoplasm lays down a transverse wall of ring-like form. As the constriction deepens, the new transverse wall broadens until, when division is completed, it forms a completely closed circular disk.

GEMMATION or **BUDDING.**—This is the method of reproduction common among yeasts. (See Fig. 284.) The cell forms a protuberance called a bud which increases in size until it equals the size of the cell which formed it and then becomes detached, although frequently not until it has developed other buds and these still others.

FREE CELL FORMATION.—This is a type of reproduction which consists of the cutting out of cells within the common cytoplasm of a cell by the activity of spindle fibers attached to the nuclei. These curve around the nuclei and form plasma membranes each of which lays down a cell wall. Seen in the formation of ascospores within the ascus of Ascomycetes.

REJUVENESCENCE or **ZOÖSPORE FORMATION.**—In this mode of reproduction the protoplasm of the cell becomes rounded out, escapes by rupture of the cell wall, forms cilia and moves about as a *zoöspore*. Later it loses its cilia, develops a cell wall and passes into a resting condition. Under favorable circumstances

it grows into a new organism. It is found in *Edogonium*, *Ectocarpus*, and many other algae. (See Fig. 257z.)

VEGETATIVE MULTIPLICATION.—This form of asexual reproduction involves the giving off of multicellular portions or vegetative outgrowths of the parent plant, such as bulbs, tubers and runners, which become detached and develop into new plants. It occurs frequently in nature but is also accomplished artificially through the removal by man of parts of the parent plant, such as stem or root cuttings or leaves, or by layering, *i.e.* bending branches over until they touch the soil and subsequently give

FIG. 24.—Propagation of Strawberry (*Fragaria*) by runners (stolons). The two plants in the small pots developed from the tips of the runners produced by the older plant in the central pot. (Gager.)

off roots and buds, when they are cut away from the parent plant. (See Fig. 24.)

There are two kinds of **SEXUAL REPRODUCTION**, viz.: *Conjugation* and *Fertilization*. In both of these the sexual cells, called *gametes*, or their nuclei come together and their protoplasm blends to form a new cell. Fertilization is the common method seen in higher plants.

CONJUGATION.—A union of two gametes, alike in physical character, the product being a *zygote* or *zygospore*. This method of reproduction is seen in the molds, *Spirogyra*, *Zygnema*, etc.

FERTILIZATION.—A union of two unlike gametes or their nuclei, the product being an *oöspore* (or *zygote*). One gamete, the male sexual cell, is smaller and active, while the other, the

female sexual cell, is larger and passive. This process is seen among the higher and many of the lower plants.

INDIRECT NUCLEAR DIVISION (MITOSIS OR KARYOKINESIS)

This is the general method of division seen in the formation of tissues of higher plants.

The process begins in the nucleus and ends with the formation of a cell wall dividing the new-formed cells.

When we examine a cell in its *resting stage*, we find the nucleus more or less spherical in shape, surrounded by a *nuclear membrane* and containing a *nuclear network*, *nuclear sap* and one or more *nucleoli*. The nuclear network consists of a colorless network of *linin* adhering to which are numerous minute granules called *chromatin* which take the stain of a basic dye. Surrounding the nucleus is the *cytoplasm*.

As the cell commences to divide, the nucleus elongates and the linin threads of the nuclear reticulum shorten, drawing the chromatin granules together into a thickened, twisted, chromatic thread called the *spireme*. This thread splits transversely and thus becomes divided into a number of rods termed *chromosomes*. Each of these then splits into two longitudinal halves that may be termed the *daughter-chromosomes*. They lie within the nuclear cavity united by delicate threads. There now begins a phenomenon concerned with the cytoplasm which is primarily a process of spindle formation. The granular cytoplasm accumulates at the poles of the elongated nucleus, forming the *cytoplasmic caps*. Presently it begins to show a fibrillar structure, the threads extending outward around the periphery of the nucleus forming an umbrella-like arrangement of fibers from both cytoplasmic caps. With the formation of fibers comes a breaking down of the nuclear membrane and, in consequence, the fibers enter the nuclear cavity and organize the *spindle*. Some of the fibers become attached to the split chromosomes and push, draw or pull them to the *equatorial plate*, halfway between the poles. Meanwhile the nucleolus disappears. As the chromosomes line up at the *equatorial plate* their daughter halves are drawn apart in V-shaped fashion. The split extends and eventually one daughter-chromosome is drawn to one pole and the remaining half to the other. At the respective poles the

FIG. 25.—Semi-diagrammatic representation of nuclear and cell-division. *a*, resting cell ready to begin division; *b*, the nuclear reticulum is assuming the form of a thickened thread, and the cytoplasm at opposite poles is becoming thread-like to form the spindle fibers; *c*, the nuclear thread has divided longitudinally through the middle, and the spindle fibers have become more definite; *d*, the nuclear membrane and the nucleolus have disappeared, and the nuclear thread has become segmented into chromosomes which are assembling at the equator of the cell. All of the phases of division thus far are called *prophases*. *e*, the *metaphase*, where the longitudinal halves of the chromosomes are being drawn apart preparatory to their journey toward the opposite poles; *f*, the *anaphase*, or movement of the chromosomes toward the poles, is about completed, connecting fibers extend from pole to pole; *g*, *telophase* where the chromosomes have begun to spin out in the form of a nuclear reticulum. The connecting fibers have begun to thicken in the equatorial plane; *h*, the connecting fibers have spread out and come into contact with the wall of the mother cell in the equatorial plane, and the thickening of the fibers throughout this plane has made a complete cell plate within which the dividing wall will be produced; *i*, a nuclear membrane has been formed about each daughter nucleus, and the dividing cell-wall is completed. The two daughter cells are now ready to grow to the size of the parent cell in *a*, when the daughter nuclei will appear as does the nucleus there. All highly magnified. (Stevens.)

daughter chromosomes form a dense compact knot. A cell membrane (*cell plate*), composed of material contributed largely through the shrinking of the spindle fibers, is now formed through the middle of the spindle. This soon splits to form a thin vacuole lying between the two membranes (the *plasma membranes*). Presently there appears within the vacuole and between the membranes a substance composed chiefly of pectose (and representing the *primary cell wall*) which later, by union with mineral substances, chiefly calcium, becomes the *middle lamella*. On either side of this deposit the plasma membranes form a *cellulose membrane*. The flattened vacuole extends toward the periphery and ultimately a complete cell wall is formed.

The dense compact knots of chromosomes, at the poles of the spindle, that are to form the daughter-nuclei, now begin to expand and clear mesh-like spaces to appear between the expanding threads. As this process advances the chromosome substance becomes distributed through the nuclear cavity. It is soon possible to distinguish the chromatin from the linin. Eventually an irregular *network* of *linin* carrying *chromatin granules* is formed through the area of the nucleus. A *nuclear membrane* also is formed and the *nucleolus* reappears. The *spindle fibers* disappear. The *daughter-nuclei* increase in size and each *daughter-cell* formed by this process now assumes the resting stage.

FIG. 26.—Enlarged views of spindle fibers and cell-plate formation. A, swellings appearing on the fibers. B, lateral union of the swellings, forming a continuous cell plate. C, splitting of the cell plate into two plasma membranes. D, a cell wall is secreted between the new plasma membranes of the daughter cells. (From "A Textbook of General Botany" by Smith, Overton et al., Macmillan Co. Publishers.)

The process of mitosis may be divided into four stages or phases and each of these briefly summarized, as follows:

1. PROPHASE.—Construction of the chromatic thread or spireme. Transverse segmentation of spireme into chromosomes and their longitudinal splitting into double chromosomes. Disappearance of the nucleolus or nucleoli. Formation of the cytoplasmic caps and spindle. Disappearance of nuclear membrane.

2. METAPHASE.—Arrangement of the double chromosomes at the equatorial plate with their daughter halves drawn apart in V-shaped fashion. Attachment of spindle fibers from poles of spindle to the halves of the chromosomes.

3. ANAPHASE.—Separation of the daughter halves of each double chromosome. Movement of one-half of each double chromosome to each of the poles of the spindle. Formation of two dense knots or masses of daughter chromosomes, one at each pole.

4. TELOPHASE.—Expansion of the masses of daughter chromosomes at the poles, and their division into linin-carrying chromatin granules, and the complete organization of the daughter nuclei. Appearance of a nuclear membrane around each daughter nucleus. Appearance of a nucleolus or of nucleoli in each daughter nucleus. Construction of a cell plate between the two daughter cells. Splitting of the cell plate to form plasma membranes. Formation of the middle lamella between the plasma membranes. Formation of a cell wall by each plasma membrane.

NUMBER OF CHROMOSOMES.—The number of chromosomes found in the cell nucleus varies with the kind of cells considered in each individual and with the species. Every species of plant or animal has a definite number of chromosomes that appear when its body cells undergo mitosis. In all sexually-reproducing plants or animals two kinds of cells occur, the body- or *somatic-cells* and the *germ cells*. Each of the body cells contains twice as many chromosomes as each of the germ cells. Thus, the *Stramonium* contains 24 chromosomes in each of its body cells and 12 in each egg and sperm cell. The number of chromosomes in the body cells is constant, is called the *diploid* or species number, and is expressed as $2x$. The number in the germ cells is called the *haploid* number and is expressed as x .

MEIOSIS OR MATURATION

The process whereby the diploid number becomes haploid is called *meiosis* or *maturation*.

Each of the spore mother cells which give rise to pollen grains and to eggs in plants are formed by the division of body cells and

like them contain $2x$ chromosomes, of which x chromosomes are contributed by the egg nucleus and x chromosomes by the sperm nucleus. It is apparent that if each uniting egg and sperm had $2x$ chromosomes, the zygote resulting would contain twice the number of chromosomes as found in the body cells of the parents. Moreover, the body cells of the sporophyte resulting from the

FIG. 27.—Diagram showing various stages in the reduction division of a spore-mother-cell of a plant. A, resting stage of the mother-cell nucleus; B, the nuclear skein or spirem during synapsis; C, the spirem after synapsis, showing its double (diploid) nature; the dotted line indicates the segmentation of the spirem into 2 diploid chromosomes, each of which has split longitudinally in D; E, the diploid chromosomes on the equator of the spindle of the first (heterotypic) division; F, late anaphase; G, metaphase of the second or homotypic division; H, late anaphase of same, two haploid chromosomes approaching the poles of each spindle; I, the four daughter cells (spores) of the tetrad. (After Gager.)

division and redivision of the oöspore would contain $4x$ chromosomes. This does not occur, for at a certain stage in the life history of every sexually reproducing plant between each zygote formation and the next, a reduction division takes place whereby the $2x$ chromosomes are reduced to x chromosomes. This reduction in the chromosome number takes place in the two nuclear divisions of spore-mother cells to form pollen grains and embryo sacs with the result that each of the starting points of the gametophyte generation have their chromosomes reduced to the x or *haploid* number.

In the first division called *heterotypic mitosis*, the resultant cells contain, in their two daughter nuclei, only one-half the number of chromosomes found in the nuclei of the body cells. These nuclei do not pass into the resting condition but soon undergo a second division.

In the second division called *homotypic mitosis*, each of the two nuclei divide again to form an aggregation of four daughter cells in all, each with a nucleus containing the x number of chromosomes. The x number of chromosomes continues through the whole gametophyte generation, so that the uniting egg nucleus and sperm nucleus possess this number. In the act of fertilization the x chromosomes from the egg nucleus are combined with the x chromosomes from the sperm nucleus to form $2x$ (diploid) chromosomes in the zygote. When the zygote divides, each of its daughter cells contains $2x$ (*diploid*) chromosomes and their number is continued throughout the sporophyte generation.

CHAPTER IV

LIFE HISTORY OF THE MALE FERN [AN INTERMEDIATE TYPE OF PLANT]

The Male Fern (*Dryopteris Filix-mas*) has long been known to the pharmaceutical and medical professions as the source of the drug ASPIDIUM, a most valuable remedy for the expulsion of tapeworm. The parts of the plant employed are the rhizome and stipes which are collected in autumn, freed of the roots and dead portions and dried at a temperature not exceeding 70° C.

Alike with other ferns, the Male Fern shows two phases or generations in its life cycle. The first of these is that which is represented by the more conspicuous plant which we call "the fern." It is in this phase that the spores or asexual cells are produced, hence the plant producing them is called the SPOROPHYTE (spore-bearing plant). The second phase is that which is represented in the development of an inconspicuous, tiny heart-shaped body which occurs on the surface of woodland soil and which bears the sexual organs in which sexual cells or gametes are produced. This phase is called the GAMETOPHYTE or sexual generation.

HISTORY OF THE SPOROPHYTE OR ASEXUAL GENERATION

GROSS STRUCTURE OF STEM.—The main axis of *Dryopteris Filix-mas* is the creeping underground stem or RHIZOME which is oblique or ascending in habit. It gives off numerous ROOTS from its lower and posterior portions and LEAVES from its upper and anterior portions. The leaves of a fern are called FRONDS. Behind the fronds of the present year are to be noted the persistent stalk bases of fronds of previous seasons. The roots are slender, brown and fibrous with semi-transparent tips. They are inserted on the bases of the fronds, close to their junction with the stem. The growing end of the rhizome is called the ANTERIOR EXTREMITY and is marked by the presence of an APICAL

FIG. 28.—*Dryopteris* (*Aspidium*) *Filix-mas*. $\frac{2}{5}$ nat. size. A, Sorus and portion of pinnule bearing same, in cross section, $\times 20$. (After Kny.) B, Pinnule with young sori, each covered with an indusium. C, the same, showing a later stage of the sori, with shrivelled indusia. Both slightly enlarged. (From Strasburger, Noll, Schenck and Schimper's *Lehrbuch der Botanik*.)

BUD overarched by young fronds. The opposite end is known as the POSTERIOR EXTREMITY and in the living plant is constantly decaying, as the anterior portion elongates.

GROSS STRUCTURE OF THE FROND.—Each frond consists of (1) a stem like portion called the STIPE or leaf stalk and (2) a green, expanded, leafy portion called the LAMINA or blade. The lamina is further divided into a MIDRIB bearing its first divisions or lobes which are called PINNÆ. Each PINNA (*sing.* for pinnæ) is further divided into secondary lobes which are called PINNULES. If a pinnule be examined carefully it will be seen to possess VEINS which are arranged in forked fashion. The under side of some of the pinnules will be seen to possess “fruit dots” or SORI.

HISTOLOGY OF STEM (RHIZOME).—Passing from periphery toward the center, the following structures are to be observed, when cross sections are examined under a microscope (see Fig. 30):

1. EPIDERMIS, a protective outer covering tissue, composed of a single layer of brownish cells from which are given off scales.

2. OUTER CORTEX (*hypodermis*), a zone of several layers of thick-walled, lignified, fibrous cells separating the epidermis from the softer underlying tissue. Its main function is to support the epidermis.

3. INNER CORTEX of several layers of more or less isodiametric cells (cells of nearly the same length, breadth and thickness) with thin cellulose walls and containing stored starch surrounded by a protoplasmic investment. Intercellular-air-spaces occur between many of the cells. This tissue is called *parenchyma* or *fundamental tissue*. Its cells conduct sap by osmosis and store food. Between the cells are to be noted intercellular-air-spaces, many of which contain internal glandular hairs with greenish contents.

4. PITH, of parenchyma, resembling the last in aspect and function.

5. VASCULAR BUNDLES.—These are elongated strands of water tubes and other tissues which appear in cross section as isolated spot-like areas. If the rhizome be cut lengthwise they will be observed to join each other forming an open network. Branch bundles come off of this network, some of which course

into the roots and others into the leaves. Upon reaching the blade of the leaf, they branch in such a way as to form the characteristic veins of that organ. These are of two kinds, viz.: *stem bundles* and *leaf-trace bundles*. The leaf trace bundles are the smaller of the two kinds seen. Both are of elliptical outline, as seen in cross section, and are embedded in the parenchyma forming the broad central matrix. The stem bundles are

FIG. 29.—A, Cylindrical network of vascular bundles in the stem of *Dryopteris Filix-mas*. B, A portion of the same more highly magnified. At L are the interstices over which the leaves are inserted; at G are branches (leaf trace bundles) passing into the leaves from the main vascular bundles. (Sayre.)

comparatively broad and, as viewed in longitudinal sections, form a continuous network with good-sized meshes, each mesh being opposite the point of insertion of one of the leaves (see Fig. 29). In transverse section these bundles are seen to be usually ten in number and arranged in an interrupted circle within the fundamental tissue. The leaf-trace bundles are comparatively narrow and are observed to come off of the stem-bundles and pass out through the cortex into the leaves (fronds). When each "bundle" is examined under a high-power magnification it is seen to be composed of: (a) an *endodermis* or *bundle sheath*, a single layer of cells with yellowish walls and granular contents enveloping the bundle; (b) a *pericycle* (*pericambium*) or phloem sheath of one to three layers of delicate thin-walled cells, rich in protoplasm; (c) a *phloem*, a broad zone of tissue formed of *phloem* cells, with thin cellulose walls and protoplasmic contents, which convey sugar in solution from the leaves to the roots, and broader *sieve tubes* which appear polygonal in transverse section and whose function is that of conveying soluble proteins in the same direction; (d) a broad central zone, the *xylem* (wood) formed of thin-walled *xylem cells* which conduct sap slowly and store food, and *scalariform tubes* or *tracheids* which conduct crude sap (water with mineral salts in solution) rapidly from the roots to the leaves (fronds). The scalariform tubes in lengthwise view have ladder-like thickenings in their walls which alternate with thin areas.

FIG. 30.—*Dryopteris Filix-mas*. A, Transverse section of the rhizome, $\times 3$. B, Transverse section of the stipe base, $\times 5$. C, Longitudinal section through portion of the parenchyma region showing intercellular air-spaces (*lc*) into which oleoresin containing glandular hairs project, $\times 65$. D, Transverse section, through a concentric bundle and adjacent parenchyma, $\times 65$. E, Longitudinal section through the same, $\times 65$. *a*, cortex; *c* pith; *i*, parenchyma; *y*, xylocentric bundles; *lc*, intercellular air-spaces; *l* internal glandular hairs; *sp*, scalariform tracheids. (After Berg.)

Since the xylem is surrounded by the phloem, the vascular bundle is of the *xylocentric* type. Strictly speaking, the endodermis and pericambium are accessory regions, surrounding, but not part of the bundle proper.

HISTOLOGY OF GROWING APEX.—When the bases of the leaves of the current year, the young rolled up leaves of the following year and the large mass of brown scales have been removed from around the apical bud of a well-grown plant, the following structures may readily be observed with a hand lens:

1. The **APICAL CONE** (*punctum vegetationis*), a rounded papilla, which occupies a terminal position in the apical region.

FIG. 31.—Apical cell of a fern rhizome in vertical longitudinal section. *a.c.*, apical cell; *h*, hair; *m*, meristem. (After Hofmeister.) *Sedgwick & Wilson's General Biology*, Henry Holt & Co.

2. The **YOUNG FRONDS**, arranged around the apical cone.

Upon removing the extreme apex of the apical cone with a sharp razor, mouting in dilute glycerine or water and examining under low power, it will be noted that a large pyramidal cell occupies the center of the apical cone. This is the *apical cell* (Fig. 31). The cells surrounding it have been derived by segmentation (cell-division) from it, by means of walls parallel to its three sides; they are termed *segment cells* and in turn undergo further division and redivision to originate the entire stem tissue and leaf tissue. Step by step the tissue cells become modified into epidermal, cortical, bundle and fundamental cells.

HISTOLOGY OF MATURE ROOT.—Transverse sections cut some distance above the root apex will present the following structures for examination: (See Fig. 32).

1. EPIDERMIS, of epidermal cells whose outer walls are brown. Some of these cells have grown out as root hairs which surround soil particles and absorb water with mineral salts in solution from the soil.

FIG. 32.—Cross section of a fern root. Photomicrograph, showing the central, radial diarch bundle and surrounding tissues. Scleriform tracheids of xylem patches (x^1 and x^2); phloem patches (p^1 and p^2); pericambium (pe); endodermis (en); cortex (co). $\times 300$.

2. CORTEX, of many layers of cortical parenchyma cells with brown walls. The outer layers of cells of this region are thin-walled, while the extreme inner ones are lignified and form a *sclerenchymatous ring* which surrounds the endodermis.

3. ENDODERMIS, a single layer of cells tangentially-elongated, whose radial walls are lenticularly thickened.

4. PERICYCLE (Pericambium), usually of one or two layers of thin-walled cells containing protoplasm and large nuclei. This region surrounds the single bundle.

5. RADIAL VASCULAR BUNDLE, consisting of two phloem patches of phloem cells and sieve tubes on either side of two radial xylem arms of scalariform tubes (tracheids) and a few xylem cells.

6. LATERAL ROOTLETS, which take origin in the pericambium.

HISTOLOGY OF ROOT APEX.—Microscopic examination shows this region to be composed of soft, pale, growing cells ending in the triangular *apex-cell* of the root. From the free base of the apex cell *segment cells* are cut off as *calyptrogen* cells. These, by dividing, form the *root cap*. The root cap or *calyptra* consists of a mass of loosely attached cells which forms a protective covering around the tip of the root.

From the inner sloping sides of the apex cell the segment cells divide to form three primary meristems. These are (1) the *dermatogen* which, by repeated division of its cells, originates the *epidermis* (outer protective covering of the root), (2) the *periblem*, originating *cortex*, and (3) the *plerome* originating the *bundle* and *related tissue*.

CONTINUITY OF CRUDE SAP FLOW.—The *crude sap* (water with mineral salts in solution) penetrates the thin walls of the root hairs by osmosis and passes into the interior of hairs, thence into the root xylem and through this to stem xylem, thence through the stem xylem into the leaves.

HISTOLOGY OF STIPE (PETIOLE).—This, in transverse section, passing from periphery toward the center, presents the following structural characteristics (See Fig. 33):

1. EPIDERMIS, a single layer of epidermal cells with dark brown outer walls.

2. OUTER CORTEX (*hypodermis*), a wide band of small cells with lignified walls.

3. INNER CORTEX, similar to inner cortex of stem but devoid of leaf-trace bundles.

4. PITH, of fundamental parenchyma, similar to same region of the rhizome, in the outer region of which are embedded a number of *xylocentric vascular bundles* arranged in an interrupted

circle. Each of these shows a central xylem mass surrounded by an outer phloem mass. Each bundle is enveloped by a *pericambium* and an *endodermis* or bundle sheath.

HISTOLOGY OF A PINNULE.—When transverse and surface sections of the *pinnule* are studied microscopically, the following structural details will be observed (See Fig. 34):

FIG. 33.—Transverse section of stipe of *Dryopteris Filix-mas* showing epidermis (*e*); hypodermis (*h*); inner cortex (*ic*); concentric fibrovascular bundles, one of which is shown at (*b*); endodermis (*en*); pericambium (*p*); xylem (*x*); and phloem (*p*). (Photomicrograph.) $\times 50.4$.

1. **UPPER EPIDERMIS**, of wavy-walled, slightly chlorophylloid, flat upper epidermal cells, devoid of stomata (breathing pores), but with rather thick cuticle.

2. **MESOPHYLL**, a region between the upper and lower epidermis consisting largely of chlorenchyma tissue which is traversed by vascular bundles. The chlorenchyma contains irregular shaped, chlorophylloid cells, with abundant chloroplasts. Intercellular-air-spaces are found between various cells which are larger in the lower than in the upper region.

Internal glandular-hairs with greenish oleoresin contents are frequently to be discerned projecting into many of these spaces.

3. XYLOCENTRIC VASCULAR BUNDLES or laminar veins, that distribute sap to, and carry sap from the mesophyll. These are

FIG. 34.—Structure of the Fern Pinnule. (*Dryopteris Filix-mas*). A, lower epidermis in surface section. B, upper epidermis in surface section. C, cross section of mid-vein and adjacent interneural regions of lamina. D, cross section of marginal region. *v*, epidermal cells below a vein; *st*, stoma; *gc*, guard cells; *n*, nucleus; *c*, chloroplastid; *sv*, sap vacuole; *chl*, chlorenchyma; *col*, collenchyma; *gl*, internal glandular hair; *a*, intercellular-air-space; *si*, sieve tissue (phloem); *xy*, xylem; *pe*, pericycle; *en*, endodermis; *uep*, upper epidermis; *mes*, mesophyll; *lep*, lower epidermis.

seen to be embedded in the chlorenchyma of the mesophyll. The xylem portion of each bundle is nearest, in end regions of veins, to the upper surface of the leaf and so the bundles there approach the collateral type.

4. ENDODERMIS, a continuous layer of cells whose radial and inner walls are more thickened than the outer walls and which surrounds each bundle and binds it in place.

Between the endodermis and the vascular bundle will be noted one or more layers of thin-walled PERICYCLE cells. In the regions of larger bundles, *collenchyma* tissue composed of thick-angled cells occurs between the endodermis and the two epidermal layers.

5. LOWER EPIDERMIS of wavy-walled, flattened, chlorophylloid cells with thin cuticle and many *stomata* (breathing pores). Each STOMA is surrounded by a pair of crescent-shaped *guard cells* which regulate its opening and closing. The upper and lower epidermis are continuous around the laminar margin.

COMPARATIVE PHYSIOLOGY OF ROOT, STEM AND LEAF (FROND).—The primary function of the roots of the Male Fern is that of ABSORPTION of water with mineral salts in solution. The secondary function is that of SUPPORT for the stem, the tertiary, that of STORING FOODSTUFFS to tide the plant over the season when vegetative activities are lessened. Water is the most essential of all materials absorbed by vegetable organisms. It is found in the soil surrounding the soil particles with certain mineral salts dissolved in it. The delicate root-hairs with thin cellulose walls, protoplasmic lining and sap denser than the soil water, are firmly adherent to these particles. The soil water diffuses through these walls by osmosis and comes into relation with the *outer plasma membrane* or *ectoplasm*, a delicate protoplasmic membrane, which has the power of selecting what it wants and rejecting what it does not need. In this way only such solutes as are of value to the plant are admitted. The water with mineral salts in solution, once within the root-hair protoplast, is called "*crude sap*." This passes through the hair into the cortical parenchyma cells which are in contact with the scalariform tracheids. It passes from one cortex cell to another by osmosis and, under considerable root pressure, is forced into the scalariform tubes of the xylem. Therein it is conveyed upward by root pressure through the tracheids of the stem bundles into those of the leaves and finally osmoses into the leaf parenchyma cells (mesophyll).

Carbon dioxide (CO_2), from the air, enters the leaf through the stomata. From the stomata it moves through the inter-

cellular-air-spaces to the mesophyll cells which line these, whence it is absorbed. Within the mesophyll cells are found small chloroplasts composed of protoplasm and chlorophyll. The kinetic energy of the sun's rays is absorbed by the chlorophyll, which renders the energy of the absorbed light effective in breaking up the CO_2 and H_2O into their component elements C, H and O, and in rearranging them in such a way as to ultimately form sugar. This process is called PHOTOSYNTHESIS. The stroma of

FIG. 35.—Demonstration of the production of starch in illuminated parts of foliage leaves. A, leaf with piece of cork attached which cut off the direct rays of the sun. B, the same leaf decolorized and treated with iodine after removal of cork 24 hours later to show that starch was removed from the shaded spot and not replenished. The illuminated portion turned blue on addition of iodine, an indication of an abundance of starch. C, the entire leaf was shaded except the part indicated by letters CHO. (*Mottier.*)

the chloroplast has the power of manufacturing starch from the sugar formed within it.

According to von Baeyer, CO_2 is split into C and O_2 , the C being retained, the O_2 given off. The nascent C is linked with H_2O to form CH_2O (formic-aldehyde); six molecules of this are then united to form grape sugar or glucose ($\text{C}_6\text{H}_{12}\text{O}_6$). The Baeyer theory, however, has been criticized by Willstatter and others on the ground that formaldehyde is employed with great difficulty for the synthesis of carbohydrates and so cannot be considered as an intermediate product of carbon dioxide assimila-

tion. According to the theory of Willstatter which is not as yet generally accepted, CO_2 is first united with chlorophyll to form a compound, the atoms of this compound become rearranged to form a peroxide which later is decomposed by catalytic enzymes with the liberation of free oxygen. The formation of starch may be expressed by the following equation: $6\text{CO}_2 + 5\text{H}_2\text{O} = \text{C}_6\text{H}_{10}\text{O}_5 + 6\text{O}_2$. The molecules of glucose and starch contain stored-up energy. A portion of the grape sugar is removed from solution by the chloroplast and converted into starch which is stored up within it; another portion is used to nourish the protoplasm of the cell. But the greater portion of sugar manufactured descends in solution through the phloem cells of the bundles of the veins, mid-rib and stipe to the stem or roots, where it is removed from solution by the action of the leucoplasts which convert it into reserve starch.

Sugar and starch, however, are not the only food materials manufactured in the leaf. Proteins are likewise formed. These are composed of carbon, hydrogen, oxygen, nitrogen, sulfur and sometimes phosphorus. They are formed from grape sugar with the addition of nitrogen and the other elements by the living protoplasm. The source of nitrogen, sulfur and phosphorus is the mineral salts which are found in the crude sap. These proteins descend through the sieve tubes of the veins, *midrib* and *petiole* to the stem and roots, nourishing all of these parts with protein material. The surplus protein material is stored in various cells for future use.

Not all of the water carried to the green cells of the leaves by the tracheids is used in the manufacture of sugar and starch. Some of it is employed in chemical reactions taking place within the cells and, some by the protoplasm itself of these cells for the purpose of repair. Whenever there is a surplus of water in the cells of the leaf, some of it escapes into the intercellular-air-spaces as water-vapor. When the humidity of the external atmosphere is less than that of the air-spaces, the water vapor passes out through the stomata. The passage of water vapor through the stomata is called TRANSPIRATION.

All of the living cells of root, stem, leaf as well as other organs of the fern require *oxygen*. This gas enters the plant through the

stomata and passes into the intercellular-air-spaces of all of the organs which communicate one with another throughout the entire plant. It diffuses through the cell walls lining these spaces into the cells. Here it unites with the food materials present, oxidizing a portion of them. Oxidation of glucose, etc. is a breaking down process, similar to the combustion of fuel in a furnace, but less intensive and not accompanied by light. The chief products of oxidation are *carbon dioxide* and *water*.

The oxidation of food materials within the cells of organisms is called **RESPIRATION**. The carbon dioxide resulting from this process diffuses through the walls of the cells into the intercellular-air-spaces and finally passes out through the stomata. When food materials such as grape sugar undergo oxidation, their stored-up energy (potential energy) is set free as energy in motion (kinetic energy). This energy is utilized by the protoplasm of the cell in carrying on its various kinds of works.

Respiration may be expressed by the following equation:

Respiration differs from photosynthesis in the following particulars:

RESPIRATION	PHOTOSYNTHESIS
1. It occurs in all living cells.	1. It occurs only in cells containing chlorophyll.
2. It takes place during the day and the night.	2. It takes place only during the day under natural conditions.
3. Oxygen is absorbed and carbon dioxide is set free.	3. Carbon dioxide is absorbed and oxygen is set free.
4. Carbohydrates are combined with oxygen and oxidized.	4. Carbohydrates are formed.
5. Potential energy is released from the food molecules as kinetic energy.	5. Kinetic energy is stored in the food molecules as potential energy.
6. It is a destructive process.	6. It is a constructive process.

There is another process which goes on in the various cells of these organs. It will be recalled that the surplus glucose is stored up in the form of starch by the leucoplasts and that surplus proteins are also stored. These stored foods are not in the form whereby they can be used by the protoplasm when needed for

repair and growth. In order to be serviceable for these purposes, they must be changed into water-soluble products. This is accomplished through the activity of substances called *enzymes* which are manufactured by the cell protoplasm. The process whereby water-insoluble foods are converted into water-soluble products is called **DIGESTION**.

GROSS STRUCTURE AND HISTOLOGY OF THE SORI AND SPORANGIA.—The *sporangia* or spore cases are found clustered together in circular groups on the under surface of the *pinnules* nearer the mid-vein than the margin. Each group of sporangia is covered

FIG. 36.—Sporangia of an undetermined species of fern; *li*, lip-cells; *an*, annulus; *st*, stalk; *sp*, mature spores. Each of the four nuclei in the upper cells of the stalk is in the terminal cell of one of the four rows of cells that compose the stalk. (Gager.)

with a membranous expansion of the epidermis called the *indusium*. The whole is called a *sorus* (Fig. 28A) (pl. sori) and contains many sporangia. Each *sporangium* is composed of: (a) the *stalk* of considerable length and usually comprising three rows of cells, outgrowths of the epidermis of the *pinnule*; and (b) the *head*, sub-globular and hollow, consisting for the most part of a covering wall of thin-walled, flattened cells, within which will be noted a marginal ring of cells, with walls having U-shaped thickenings, and called the *annulus*. Beneath the wall of an immature head of a spore case occurs a *tapetum* of a single layer of cells and in the center spore mother cells. The spore mother cells undergo two successive divisions to form *tetrads* or groups of four spores which later separate as mature *spores*. During the first of these divisions reduction in the number of chromosomes takes place from the number characteristic of the species (*diploid* number) to the *haploid* number or half the species or

sporophyte number, so that each spore contains only half the number of chromosomes found in the sporophyte generation and thus belongs to the gametophyte generation.

Each spore is a single cell composed of an outer brown wall with band-like markings called an *exosporium*, an inner thinner wall or *endosporium*, and within this a mass of protoplasm containing a nucleus.

The sporophyte generation ends with division of the spore mother cell to form a tetrad of young spores.

HISTORY OF THE GAMETOPHYTE OR SEXUAL GENERATION

The gametophyte generation begins with the tetrad of young spores within the sporangium.

RUPTURE OF SPORANGIUM AND SPORE DISSEMINATION.—As was previously indicated, each sporangial head had a row of cells with U-shaped thickenings around the margin called an annulus. As the sporangium matures, the water escapes from the cells, pulling them together and holding the annulus like a bent spring. The thinner walled cells at the side of the spore case opposite the annulus, unable to stand the strain, are consequently torn; the annulus then straightens and a wide rent is made in the sporangium. The annulus then recoils and hurls the spores out of the sporangium.

GERMINATION.—The fern spore, falling upon a moist surface, germinates, producing a delicate, green, septate filament called a *protonema*. One end of this structure shows larger cells, which, by the formation of oblique walls, cut out an apical cell of somewhat triangular shape. This is the growing point of what eventually becomes a mature, green, heart-shaped body called the "*prothallium*" or "*prothallus*." The prothallium, about the size of an infant's finger nail, develops on its under surface *antheridia*, or male sexual organs, *archegonia*, or female sexual organs, and *rhizoids* or hair-like absorptive structures. The antheridia appear three to five weeks after spore germination. They are hemispherical in shape and are situated among the rhizoids toward the posterior end. Each antheridium consists of a three-celled wall which completely surrounds the *spermatocytes* or mother-cells of the spermatozoids. Within each *spermatocyte*

the protoplasm arranges itself in a spiral fashion forming a *spermatozoid*, a spiral, many ciliated, male sexual cell. From two to four weeks after the maturation of the antheridia, the *archegonia* makes their appearance toward the indented apex of the lower prothallial surface as outgrowths of the *prothallial cushion*. Since they appear later than the antheridia they are not likely to be fertilized by spermatozoids from the antheridia of the same

FIG. 37.—Group of fern prothallia growing on the surface of soil. Young sporophytes are developing on some of the prothallia. (Gager.)

prothallium. Each *archegonium* is composed of a *venter*, *neck*, *neck canal-cells*, *ventral canal-cell*, and *ovum* or egg-cell (see Fig. 38B). The neck is composed of cells arranged in four rows, forming a cylinder, one layer of cells thick. This protrudes from the surface of the prothallium and encloses the neck canal-cells and ventral canal-cell. The ovum is embedded in the prothallial cushion just beneath the ventral canal-cell. Upon the maturation of the archegonium, the canal cells are transformed into a

mucilaginous substance which fills a canal extending from the outside opening (mouth) to the ovum.

FIG. 38.—A, under surface of a fern prothallium showing archegonia (*f*), antheridia (*m*) and rhizoids (*r*); B, immature archegonium showing binucleate neck canal cell (*n.c.c.*), ventral canal cell (*v.c.c.*), and egg (*e*); C, mature archegonium showing sperms (*s.p.*) moving through neck canal (*n.c.*) toward ovum (*e*); and venter (*v*). All highly magnified.

During wet weather the mature antheridial wall bursts open and the many ciliated *spermatozoids* escape into the water. These, moving in the water, are drawn by the chemotactic

malic acid to the mouths of the archegonia of another prothallus, and, passing down the canal of each of these, gather around the ovum. One, probably the best adapted, fuses with the ovum and fertilizes it forming a zygote (*oöspore*) or fertilized egg. This completes the gametophyte generation.

ORIGIN OF NEW SPOROPHYTE PLANT FROM FERTILIZED EGG

The fertilized egg now divides into hemispheres, then into quadrants and again to form eight cells arranged in a spherical mass and called *octant cells*. The octant cells divide to form a many celled *embryo* which shows the following parts: *foot rudiment*, *root rudiment*, *stem or rhizome rudiment*, *hair rudiments*, and *first leaf*.

GROWTH OF EMBRYO INTO MATURE SPOROPHYTE.—The foot rudiment develops into the *foot* which obtains nourishment from

FIG. 39.—A, median longitudinal section through immature antheridium, and cell of prothallium showing prothallial cell (*p*), and antheridial wall surrounding a number of spermatocytes; B, similar section through mature antheridium and cell of prothallium showing fully developed spermatozooids (*s.p.*) enclosed by wall of antheridium (*an*). Both highly magnified.

the prothallium, upon which the young sporophyte is for a time parasitic. The root rudiment becomes the *first root* which grows downward into the soil. The stem rudiment becomes a *rhizome* which, with the first leaf, turns upward. In a few weeks the prothallus decays and the sporophyte is established as an independent plant. More leaves (fronds) and roots are developed and ere long continued growth results in the formation of a mature sporophyte which presents for examination: (1) a *subterranean stem* or *rhizome*, (2) *roots*; and (3) *aerial fronds*, each of which consists of a *stipe* or *petiole* and a *lamina* or *blade*, the

latter divisible into *pinnæ* or *lobes* and *pinnules*, upon which last *sori* are developed.

ALTERNATION OF GENERATIONS.—It will be observed that in the life cycle of the Male Fern there occur two distinct genera-

FIG. 40.—*Pteris* (The brake fern). A. Early stage in development of embryo from oöspore showing division of octant cells to form: *f*, foot rudiment; *s*, stem rudiment; *b*, first leaf rudiment; *w*, root rudiment. B. Later stage in the development of embryo; *f*, foot embedded in archegonium (*aw*) of prothallus (*pr*); *s*, young stem; *b*, young first leaf; *w*, young root. A, after Kienitz-Gerloff; B, after Hofmeister. (From Strasburger's *Lehrbuch der Botanik*.)

FIG. 41.—Diagram of life-cycle of a fern. 1, Mature sporophyte; 2, sporangium; 3, spore mother cell; 4, tetrad; 5, spores; 6, germinating spore; 7, prothallus; 8, archegonium; 8a, antheridium; 9, ovum; 9a, antherozoid; 10, fertilization; 11, zygote. (Gager.)

tions, one, a sporophyte or asexual generation which begins with the zygote (or oöspore) and ends with the division of the spore mother cells to form tetrads of spores; a second, the

gametophyte or sexual generation, beginning with the spore and ending with the fertilization of the egg to form the zygote or oöspore. The sporophyte gives rise to the gametophyte which in turn gives origin to the sporophyte. This is called "*alternation of generations.*"

During this alternation of generations, a series of nuclear changes takes place in which the *diploid* chromosomes or full quota of the sporophyte generation are reduced to the *haploid* number in the gametophyte generation. These changes occur in the mitotic divisions of the spore mother cells to form spores which contain only half the number of chromosomes found in the spore mother cell and the rest of the sporophyte generation. At the time of fertilization the two gametes, antherozoid and ovum, fuse and restore the full quota or *diploid* number of chromosomes to the zygote.

CHAPTER V

NON-PROTOPLASMIC CELL CONTENTS

The non-protoplasmic cell contents comprise many kinds of substance of non-living character sometimes referred to as inclusions in the protoplast. They include inorganic and organic substances which are deposited in the cell as a result of its physiological activities. Many of them, when separated from plants, constitute valuable drugs and foods. Some of them like glucose and other sugars, starch and inulin are products of photosynthesis, others like proteids, fixed oils and fats are formed from the elements of glucose, still others like enzymes are direct products of protoplasm, and a number of mineral compounds such as calcium oxalate, calcium carbonate and silica bodies are formed as deposits as a result of chemical processes within the cell.

1. SUGARS.—*Sugars* comprise a group of crystalline substances found in the cell sap of many plants either free or in combination with glucosides. They may be divided into two main groups: monosaccharoses and disaccharoses. The MONOSACCHAROSES are simple sugars containing two to nine atoms of carbon, which are known respectively as bioses, trioses, tetroses, pentoses, hexoses, etc. Of these the hexoses ($C_6H_{12}O_6$) are the most important and of wide distribution. Examples of the hexoses found in drug plants are: (a) DEXTROSE (grape sugar), found in the leaves, stems, fruits, sprouting grains and nectaries of flowers of nearly all plants; (b) FRUCTOSE (levulose or fruit-sugar), commonly associated with dextrose; (c) D-MANNOSE, found in the saccharine exudation of the Manna Ash (*Fraxinus Ornus*); and (d) SORBINOSE, found in ripe Mountain Ash berries. Upon evaporating the sap or treating the parts containing these principles with alcohol they can be crystallized out.

Flückiger's Micro-chemic Test for the determination of different kinds of sugars: Dissolve a small portion of copper tartrate in a drop of sodium hydrate on a glass

slide; in this place the section and put on the cover slip. If fructose is present cuprous oxide crystals will at once be formed without warming. If grape sugar is also present, a gentle warming will produce another crop of reddish-yellow crystals. If dextrin be present, continued heating will still further augment the number of crystals. Cane sugar and mannite, on the other hand, will respond negatively to this test.

The DISACCHAROSES, having the chemical formula of $C_{12}H_{22}O_{11}$, include sucrose, maltose, trehalose, melibiose, touranose and agavose. Of these SUCROSE is the most important. It is found in the stems of sugar cane, sorghum, corn and Mexican grass; in many fleshy roots notably the sugar beet; in the sap of the sugar maple and various palms; in various fruits, as apples, cherries, figs, etc., in the nectaries of certain flowers; in honey; and in a number of seeds. It crystallizes in monoclinic prisms or pyramids.

When sections of plant parts containing cane sugar are placed for a few seconds in a saturated solution of copper sulphate, then quickly rinsed in water, transferred to a solution of 1 part KOH in 1 part of water, and heated to boiling, the cells containing the sugar take on a sky-blue color.

Invertase, an enzyme of the yeast, reduces cane sugar to dextrose and levulose and zymase, another enzyme of the same plant, ferments these forming carbon dioxide and alcohol.

MALTOSE is found in the germinating grains of barley and other cereals as a product of the action of the ferment diastase on starch. It reduces Fehling's solution, forming cuprous oxide, but one-third less with equal weights.

TREHALOSE or mucose is found in ergot, *Boletus edulis*, the Oriental Trehala and various other fungi.

MELIBIOSE is formed with fructose upon hydrolyzing the trisaccharose melitose which occurs in the molasses of sugar manufacture and in Australian manna.

TOURANOSE is produced upon hydrolyzing melizitose, a trisaccharose which occurs in Persian manna.

AGAVOSE is found in the cell sap of the American Century Plant, *Agave americana*.

2. STARCH.—Starch is a carbohydrate having the chemical formula of $(C_6H_{10}O_5)_n$ which is generally found as the first visible product of photosynthesis in most green plants. It is

found in or more rarely adhering to the chloroplasts of green parts of plants in the form of minute granules. This kind of starch is known as ASSIMILATION STARCH.

Assimilation starch is dissolved during darkness within the chloroplasts by the action of a starch-splitting ferment called *amylase* and passes into water-solution as a glucose which is conveyed downward to those parts of the plant requiring food. In its descent, some of the surplus glucose not used by the cells in

FIG. 42.—Cell of *Pellionia Daveauana*, showing reserve starch-grains. The black, crescent-shaped body on the end of each grain is the *leucoplast*. Greatly enlarged. (Gager.)

their nutrition, is stored up by leucoplasts in medullary-ray cells, and in various parts of the xylem, phloem, pith and cortex in the form of transitory starch grains. Considerable of it, however, is carried to the underground parts, such as rhizomes, tubers, corms, bulbs or roots, or into aerial storage tissues, as the storage cells of seeds, where the leucoplasts store it in the form of larger-sized grains called RESERVE STARCH. This type of starch is generally characteristic for the plant in which it is found. It constitutes stored-up food for the plant during that period of the year when the vegetative processes are more or less dormant.

STRUCTURE AND COMPOSITION OF STARCH.—Starch grains vary in shape from spheroidal to oval to conchoidal to polygonal. They are composed of layers of soluble carbohydrate material and probably other substances called "*lamellæ*," separated from each other by a *colloidal substance* resembling a mucilage in its behavior toward aniline dyes. They contain a more or less distinct highly refractile point of origin or growth called the "*hilum*," which also takes the stain of an aniline dye. The layers of carbohydrate material stain variously, blue, indigo, purple, etc., with different strengths of iodine solutions. Each grain is covered with a stainable *elastic membrane*.

Starch grains may be grouped, according to the condition in which they are found in the cells of storage regions into three kinds, viz.: *simple starch grains*, *compound starch grains* and *fill starch grains*.

Simple starch grains are such as occur singly. Compound starch grains occur in groups of two, three, four, five, six or more and are designated as two-, three-, four-, five-, six-, etc., compound, according to the number of grains making up the group. *Fill starch grains* are small grains filling up the spaces between the larger grains in storage cells. These are common in commercial starches.

METHOD OF EXAMINING RESERVE STARCHES.—Many of the reserve starches are used commercially, such as potato, corn, rice, maranta, oat, wheat, sago, tapioca, etc., and it frequently becomes necessary for the microscopist to determine their purity or their presence in a sample of food or drug. The following characteristics should be noted in determining the identity or source of the starch.

1. The shape of the grain.
2. Whether simple or compound or both; if compound, the number or range in numbers of grains composing it.
3. The size of the grain in microns.
4. The position of the hilum, if distinct; whether central or excentric (outside of the center).
5. The shape of the hilum and the degree to which it is often fissured.
6. The nature of the lamellæ, whether distinct or indistinct; if distinct, whether concentric (surrounding the hilum) or eccen-

FIG. 43.—A, wheat starch grains; B, rye starch; C, barley starch; D, potato starch; E, Maranta starch; F, Sago starch. Explanation in text.

tric (apparently ending in the margin of the grain and not surrounding the hilum), or both, as in potato starch.

7. The color of the grains when stained with dilute iodine solutions; whether indigo, blue, purple, red or yellowish-red, etc.

8. The appearance under polarized light.

9. The temperature at which the paste is formed.

10. The consistency of the paste.

11. Whether the grains are unaltered or altered.*

CHARACTERISTICS OF IMPORTANT COMMERCIAL STARCHES

POTATO STARCH (*Solanum tuberosum*)

Mostly simple, conchoidal or ellipsoidal, with occasional spheroidal and two- to three-compound grains.

Size: 5 to 125 μ .

Hilum: circular, at smaller end of grain.

Lamellæ: concentric and eccentric.

Polarization cross very distinct; beautiful play of colors with selenite plate.

MARANTA STARCH (*Maranta arundinacea*)

Ellipsoidal to ovoid.

Simple.

Size: mostly 10 to 65 μ .

Hilum: a transverse or crescent-shaped cleft in center or near broad end of grain, or circular.

Lamellæ: usually faint, concentric and eccentric.

Polarization cross very distinct; fine play of colors with selenite plate.

CORN STARCH (*Zea Mays*)

Polygonal to rounded.

Up to 35 μ . Most grains over 15 μ in diameter.

Simple.

Hilum: circular or a two- to five-rayed cleft.

Lamellæ: indistinct.

Polarization cross distinct but no marked play of color with selenite plate.

RICE STARCH (*Oryza sativa*)

Polygonal.

2 to 10 μ in diameter.

Simple or two- to many-compound.

Hilum: usually indistinct, occasionally a central cleft.

Lamellæ: indistinct.

Polarization cross distinct but no play of colors with selenite plate.

WHEAT STARCH (*Triticum sativum*)

Circular grains appearing lenticular shaped on edge view; simple.

Large grains 28 to 45 μ in diameter.

Hilum: central, appearing as dot, but usually indistinct.

Lamellæ: generally indistinct, when present in partially germinated wheat, concentric.

Polarization cross indistinct; no play of colors with selenite plate.

RYE STARCH (*Secale cereale*)

Grains having a similar shape to those of wheat starch but many larger; simple.

Large grains 25 to 64 μ in diameter.

Hilum: a star-shaped central cleft or indistinct in most grains.

Lamellæ: indistinct; concentric in partly germinated rye.

Polarization cross distinct.

* Starch grains may be altered by heat, pressure, acids and enzymes. *Soluble starch* is starch altered by boiling. It gives a sky blue color with iodine water. See also dextrin, p. 82 and corrosion starch, p. 102.

G

H

I

J

K

L

FIG. 44.—G, corn starch; H, rice starch; I, corn dextrin; J, pea starch; K, cassava starch; L, bean starch. Explanation in text.

BARLEY STARCH (*Hordeum distichon*)

Grains having a similar shape to those of wheat starch but frequently tending to bulge on one side and so appear sub-reniform; large grains smaller; simple.

Grains appear elliptical to lemon shaped in edge view. Large grains usually 18 to 25 μ , occasionally up to 30 μ in length.

Hilum: centric or circular or cleft, often indistinct.

Lamellæ: concentric in partly germinated barley; usually indistinct.

Polarization cross distinct.

BUCKWHEAT STARCH (*Fagopyrum esculentum*)

Grains simple and compound.

Simple grains polygonal or rounded-polygonal.

BUCKWHEAT STARCH

Compound grains more or less rod-shaped.

2 to 15 μ in diameter.

Hilum: central.

Lamellæ: generally indistinct.

Polarization cross distinct.

CASSAVA STARCH (*Manihot utilissima*)

Grains rounded, truncated on one side.

Simple or two- to three- or four- to eight-compound.

6 to 35 μ in diameter.

Hilum: central, circular or triangular with radiating clefts frequently.

Lamellæ: indistinct.

Polarization cross prominent.

FIG. 45.—Potato starch, polarized as viewed by light passing through crossed nicol prisms of a petrographic microscope. (Schneider.)

FIG. 46.—Maranta or Arrowroot starch under polarized light. Note that the two dark bands cross at the hilum of each starch grain. (Schneider.)

BEAN STARCH (<i>Phaseolus vulgaris</i>)	CANNA STARCH (<i>Canna edulis</i> and other species of <i>Canna</i>)
Ovoid, ellipsoidal or reniform shaped simple grains, occasionally obscurely 3- or 4-sided.	Broadly elliptical, flattened, with beak or obtuse angle at one end.
Up to 60μ in length. Generally from $30-35\mu$.	Most grains 50 to 135μ in length.
Hilum: central, elongated with branching clefts.	Hilum: excentric near narrower end.
Lamellæ: distinct, concentric. In some indistinct.	Lamellæ: concentric and eccentric.
Polarization crosses shaped thus, χ .	Polarization cross very distinct; fine play of colors with selenite plate.
PEA STARCH (<i>Pisum sativum</i>)	SAGO STARCH (<i>Metroxylon Sagu</i> et al. spp., <i>Cycas</i> spp.)
Oval-oblong, elliptical or sub-reniform.	Ovoid, muller-shaped, or irregularly 3 or 4 sided with rounded angles.
Up to 51μ in length. Generally from $20-40\mu$.	Some more or less gelatinised.
Hilum: similar to that of bean starch but less cleft or simply elongated.	Simple or 2, 3 or 4-compound.
Lamellæ: distinct, concentric.	Larger grains $30-60\mu$ long.
Polarization crosses similar to bean starch.	Hilum: excentric, often altered by gelatinisation.
	Lamellæ: eccentric and concentric.
	Polarization cross distinct.

3. DEXTRIN.—Dextrin is a carbohydrate made from starch (chiefly from corn or potato starch) by the application of heat (yellow dextrin) or by treatment with both heat and acids (white dextrin). It forms a paste with water, the yellow variety tending to swell up and dissolve much more readily than the white. When examined microscopically in alcohol mounts, the grains, while conforming in general outline to those of the type of starch from which the dextrin was prepared, nevertheless show more conspicuous striations and clefts. Corn dextrin shows distinct striations, whereas striations in corn starch are absent. The grains take on a red coloration with iodine solutions.

4. AMYLODEXTRIN.—This is a carbohydrate intermediate in properties between starch and dextrin. It occurs in the form of small, irregularly shaped granules, in Mace, that take on a reddish brown to reddish violet color with iodine solutions.

5. INULIN.—Inulin is a carbohydrate isomeric with starch which has the chemical formula of $C_{12}H_{20}O_{10}$. It is found dissolved in the cell sap of many plants, especially those of the *Compositæ*. If pieces of a plant part containing this substance be placed directly in alcohol for at least a week, then sectioned and mounted in alcohol, sphæro-crystals of inulin will be seen

applied to the walls of the cells. When these sections are treated with a 10 per cent. solution of alpha naphthol in alcohol and 2 or 3 drops of strong H_2SO_4 , the sphæro-crystals will dissolve with a violet color. Fehling's solution is not reduced by inulin.

6. GLUCOSIDES (Glycosides).—These substances are very numerous in the plant kingdom. They arise in the cell sap of plants containing them as products of constructive metabolism (anabolism) and are thought by many to have the function of protecting plants against the ravages of animals. Some are known to serve as reserve food. All glucosides

are characterized by the property of being split up into glucose and other substances when acted upon by an enzyme, dilute acids or alkalies.

FIG. 47.—Inulin, in the form of sphæro-crystals from the tuber of *Dahlia*. A, precipitated from an aqueous solution; B, precipitated within the cells by long standing in alcohol. (From Stevens, after Sachs.)

EXAMPLES OF GLUCOSIDES

HESPERIDIN.—Hesperidin is a glucoside having the chemical formula of $\text{C}_{21}\text{H}_{26}\text{O}_{12}$. Like inulin it occurs in solution within the cell sap. It is found in abundance in the Rutaceæ family but occurs in many other plants. (See Fig. 85h.)

If sections of alcoholic material containing this substance, such as Buchu leaves or unripe orange peel, are mounted in alcohol and examined, sphæro-crystals will be seen. If these are then treated with a drop of alpha naphthol solution and 2 or 3 drops of strong H_2SO_4 , they dissolve with a yellow color. The same coloration is evident when 5 per cent. solution of KOH is substituted for the alpha naphthol and H_2SO_4 .

STROPHANTHIN.—This is a glucoside occurring in the cell sap of the endosperm of *Strophanthus Kombe*, *S. hispidus* and other species of *Strophanthus*. If sections of fresh *Strophanthus* seeds are mounted in a drop of water and then transferred to a drop of concentrated H_2SO_4 , the cells containing strophanthin will assume a bright green color.

SALICIN.—Salicin is a glucoside occurring in the cell sap of the bark and leaves of the Willows and Poplars. Sections of these

mounted in concentrated H_2SO_4 will show a red coloration in the cells containing this substance. If water be added, a red powder is thrown down.

SAPONINS are glucosides which have the property of frothing with water. They are found in Soap Bark, Senega, Saponaria, Yucca and other drugs and take a red color with strong H_2SO_4 .

CONIFERIN is a glucoside, occurring in the cell sap of the spruce, pine, and other plants of the *Coniferae*. If sections containing it are first treated with a solution of phenol and then with sulfuric acid, the cells containing it take on a deep blue color.

DIGITOXIN, a glucoside found in the leaves of *Digitalis purpurea*, is colored green with hydrochloric acid.

CYANOGENETIC GLUCOSIDES are common in the seeds of the Rose and Legume families. They yield hydrocyanic acid when hydrolyzed by enzymes or other agents.

7. PENTOSIDES.—These are substances related chemically to glucosides, differing from the latter in yielding pentose as one of the products, when decomposed (hydrolyzed) by enzymes, dilute acids, etc. The aloins found in the juices of various species of *Aloe* are examples of these principles. One of these, "barbaloin," found in the official *Aloe* gives a greenish fluorescence in aqueous solutions, when the latter are saturated with borax.

8. ALKALOIDS.—Chemically, these are basic carbonaceous amines which like glucosides are products of metabolism. Their method of formation in plants is uncertain. Some hold that they are katabolic products, resulting from the breaking down of tissues, while others believe them anabolic in character. They undoubtedly serve as defensive agents in plants containing them on account of their bitter taste and poisonous properties.

PROPERTIES OF ALKALOIDS

Alkaloids are invariably found in combination with acids forming salts which dissolve in water or alcohol. They are composed of carbon, hydrogen and nitrogen. Some contain oxygen. They are precipitated from saline solutions by the addition of alkalies. They are mostly colorless and crystallizable. They can be precipitated by one or more of the following alka-

loidal reagents: tannic acid, gold chloride, phospho-molybdic acid, picric acid and potassio-mercuric iodide (Maver's reagent).

EXAMPLES OF ALKALOIDS

STRYCHNINE.—This alkaloid, with a chemical formula of $C_{21}H_{22}N_2O_2$, occurs in the seeds of *Strychnos Nux-vomica*, *Strychnos Ignatii* and other species of *Strychnos*. When sections of strychnine containing seeds, previously defatted with petroleum ether and absolute alcohol, are mounted in a solution of 1 Gm. ammonium vanadate in 100 cc. of sulfuric acid, they take on a violet-red color which later changes to brown. Strychnine itself produces with sulfuric acid containing 1 per cent. ammonium vanadate a deep violet-blue color changing to a deep purple and finally a cherry red. It crystallizes from alcoholic or aqueous solutions, forming rhombohedral crystals.

BRUCINE.—This is another alkaloid obtained from *Strychnos* seeds having the formula of $C_{23}H_{26}O_4N_2$. It crystallizes from watery solutions in the form of monoclinic prisms. Nitric acid imparts to it a red color.

NICOTINE.—This is a volatile alkaloid having the formula of $C_{10}H_{14}N_2$ which is found in the *Nicotiana* genus of the Nightshade family. Sections of tobacco leaves or stems mounted in dilute Lugol's solution will show first a carmine-red color and then a reddish-brown precipitate which in time loses its color.

CAFFEINE.—This alkaloid, with a formula of $C_8H_{10}N_4O_2 + H_2O$, occurs in Tea, Coffee, Maté or Paraguay Tea, Cassine, leaves of *Ilex vomitoria* or Yaupon, and seeds of Kola, Cocoa, etc. If thin sections containing it are placed on a glass slide in 2 or 3 drops of concentrated hydrochloric acid and gently heated and then 2 or 3 drops of gold chloride solution are added, the sections then pushed to the side and the liquid allowed to evaporate, slender, yellowish, branching needles of caffeine gold chloride will be seen to separate.

COCAINE.—This narcotic alkaloid, having the formula $C_{17}H_{21}NO_4$, is found in the leaves of *Erythroxylon Coca* and *E. Truxillense*. If sections of these leaves are prepared in the same manner as indicated for those containing Caffeine, but platinum chloride solution substituted for that of gold chloride, large

feathers or fern frond crystals of cocaine-chloro-platinate will be seen separating.

ACONITINE ($C_{34}H_{47}O_{11}N$) is found in various parts of *Aconitum Napellus*. It is particularly abundant in the tuberous root of this

FIG. 48.—Cocaine chloroplatinate crystals, $\times 100$.

FIG. 49.—Morphine. Orthorhombic prisms from alcoholic solution of this alkaloid. $\times 120$.

plant. If sections of aconite root are treated on a glass slide with solution of potassium permanganate, a red precipitate of aconitine permanganate will appear in the cells containing this alkaloid.

MORPHINE ($C_{17}H_{19}O_3N$), the chief active principle of Opium, crystallizes from alcoholic solutions in orthorhombic prisms and

acicular crystals. Sulfuric acid containing 5 mg. of selenous acid in each cc. gives with morphine sulfate a blue color changing to green and then to brown.

CODEINE ($C_{18}H_{21}O_3N \cdot H_2O$), another alkaloid occurring in Opium, produces with sulfuric acid containing 5 cc. of selenous acid in each cc. a green color which changes rapidly to blue, then slowly back to green.

FIG. 50.—Codeine phosphate crystals from aqueous solution. $\times 120$.

COLCHICINE ($C_{22}H_{25}O_6N$).—This alkaloid occurs in the corm and seeds of *Colchicum autumnale*. It is very abundant in the cells surrounding the fibro-vascular bundles of the corm. If a section of either corm or seed be treated with a mixture of 1 part of H_2SO_4 and 3 parts of H_2O , the cells containing colchicine will be colored yellow. If a crystal of KNO_3 then be added, the color will change to a brownish-violet.

9. GLUCO-ALKALOIDS (Alkaloidal glucosides).—These are compounds intermediate in nature between alkaloids and glucosides, having characteristics of each. To this group belongs *solanine*, a poisonous alkaloidal glucoside with a saponin-like action, which is found in the potato (*Solanum tuberosum*), in *Solanum nigrum*, *Solanum Dulcamara*, *Solanum carolinense* and other species of the *Solanaceæ*. When sections of those plant parts

which contain this constituent are mounted in a solution of 1 part of ammonium vanadate in 1000 parts of a mixture of 49 parts of sulfuric acid with 18 parts of water, the cells containing solanin take on a yellow color which changes successively to orange,

FIG. 51.—Various forms of calcium oxalate crystals. A, stylolids from the bark of *Quillaja saponaria*; B, rosette aggregate from rhizome of *Rheum officinale*; C, raphides from the bulb of *Urginea maritima*; D, crystal fiber as seen in longitudinal section in either the xylem or phloem regions of *Glycyrrhiza*; E, micro-crystals (crystal sand) isolated from the parenchyma of *Belladonna* root; F, monoclinic prisms; and G, twin-crystals from leaves of *Hyoscyamus niger*. All highly magnified.

various shades of red, blue-violet, grayish-blue and then disappears.

10. ASPARAGINE ($C_4H_8N_2 + H_2O$).—This is an amino compound of crystalline nature which occurs widely in the plant kingdom. It has been found in certain of the slime molds and fungi, in the roots of *Althæa officinalis* and *Atropa belladonna*, in young shoots of *Asparagus*, in the seeds of *Castanea dentata*, in the

tubers of *Solanum tuberosum* and varieties of *Dahlia*, and is known to play an important part in metabolism. Stevens claims that proteids are reduced for the most part to asparagine during seed germination.¹ If thick sections are cut from a plant part containing this substance and mounted in alcohol, rhombohedral crystals of asparagin in the form of plates will be deposited upon the evaporation of the alcohol. If to these a few drops of a saturated solution of asparagine are added the crystals already formed will increase in size. To get satisfactory results the saturated solution must be of the same temperature as the mount.

11. CALCIUM OXALATE.—This substance occurs in many plants always in the form of crystals. It is formed by the reaction of salts of calcium, which have found their way into the cell sap from the soil, with oxalic acid which is manufactured by the plant as a by-product of protein synthesis and other metabolic processes. Calcium oxalate crystals dissolve readily in mineral acids without effervescence. They are insoluble in acetic acid or water. These crystals are classified according to form and belong either to the monoclinic or tetragonal system. (See Fig. 51.)

Crystals belonging to the monoclinic system possess 3 axes of unequal length, two of which are obliquely inclined to each other, the other axis forming right angles with these two. They appear to be more frequently found in plants than those of the tetragonal system.

Crystals belonging to the tetragonal system have 3 axes at right angles to each other; two of the axes are equal in length, the third being of a different length.

CRYSTALS BELONGING TO THE MONOCLINIC SYSTEM AND
EXAMPLES OF DRUGS CONTAINING THEM:

1. Solitary—Hyoscyamus, Acer Spicatum, Viburnum Prunifolium.

2. Rosette Aggregates (Druses)—Althæa, Gossypii Radicis Cortex, Stramonium, Granatum, Rheum, Fœniculum, Viburnum, Eucalyptus.

3. Columnar (Styloids)—Quillaja, Iris.

4. Raphides—Convallaria, Sarsaparilla, Veratrum Viride, Scilla, Phytolacca, Ipecacuanha, Hydrangea.

¹ Stevens' Plant Anatomy, 3d Edit., p. 189.

5. Micro-crystals (Crystal sand)—*Belladonnæ Radix*, *Cinchona*, *Hyoscyamus*, *Stramonium*, *Phytolacca*, *Capsicum*. This type is frequent in the *Solanaceæ* family.

6. Crystal Fibers—*Cascara Sagrada*, *Prunus Virginiana*, *Glycyrrhiza*, *Aspidosperma*, *Senna*, *Viburnum*, *Unpeeled Calamus*.

7. Membrane Crystals—*Aurantii Dulcis Cortex*, *Limonis Cortex*, other Citrus peels, *Condurango*, *Wafer Ash bark*.

FIG. 52.—Cystoliths from the leaf of the Fig (*Ficus carica*). A, Cystolith within cystolith sac; B, cystolith from which the calcium carbonate has been removed for use in other parts of the plant. B is from leaf that has fallen off in autumn. (From Stevens after Haberlandt.)

Solitary crystals, usually in the form of rhombohedra, occasionally in twin crystals, occur as sharp angular bodies, each one often completely filling up the lumen of a cell. *Twin crystals* occur in elm bark, *hyoscyamus*, etc. They are single prisms which have the appearance of two prisms united by their flat surfaces.

Rosette aggregates or *druses* consist of numerous small prisms or pyramids, or hemihedral crystals arranged around a central axis, appearing like a rosette or star. They often contain an organic center.

Columnar crystals or *styloids* are elongated prisms.

Raphides or acicular or needle-shaped crystals, which frequently occur in bundles in long, thin-walled cells containing mucilage called *raphide sacs*. They are more frequently found in Monocotyledons than in any other plant group.

Crystal fibers are longitudinal rows of superimposed parenchyma cells each of which contains a single monoclinic prism or rosette aggregate. Crystal fibers are found adjacent to sclerenchyma fibers such as bast or woody fibers.

Membrane crystals are monoclinic prisms, each of which is surrounded by a wall or membrane. In the process of formation, a crystal first is formed in the cell sap and then numerous oil globules make their appearance in the protoplasm surrounding it; later some of the walls of the cell grow around the crystal and completely envelop it. (See Fig. 85k.)

12. MICRO-CRYSTALS (sphenoidal micro-crystals or crystal sand) are minute arrow-shaped or deltoid forms completely filling the parenchyma cells in which they occur and giving these a grayish-black appearance. While sometimes classified with the calcium oxalate crystals, they are thought to represent some other compound as yet undetermined.

13. CYSTOLITHS.—Cystoliths are clustered bodies formed by the thickening of the cell wall at a certain point and subsequent ingrowth which latter forms a cellulose skeleton consisting of a stalk and body. Silica is subsequently deposited on the stalk while calcium carbonate is piled up on the body in layers, forming an irregular spheroidal or ellipsoidal deposit. These structures are abundantly found in the plants of the Nettle, Mulberry and Acanthus families and constitute a leading peculiarity of the same (see Fig. 52). They also occur in the Combretum and Borage families, in species of *Phlox*, *Klugia*, etc. They are probably excretory products. Excellent materials for the study of cystoliths are the leaves of the rubber plant (*Ficus elastica*) and the fig (*F. carica*). Here, the cystoliths appear as stalked mulberry-like structures suspended from the outer walls of specialized epidermal cells called "cystolith sacs."

Hair cystoliths differ from the average type in that they are devoid of a stalk. Such are seen in the non-glandular hairs of *Cannabis sativa*.

The calcium carbonate incrustation of a cystolith dissolves with effervescence on the addition of a mineral or organic acid.

14. SILICA.—Silica (SiO_2) occurs in a number of plants either as an incrustation in the cell wall, as in Diatoms, the *Equisetineæ* and *Gramineæ* or more rarely in the form of "silica bodies," such as are found in certain Palms and Orchids. It is insoluble in all the acids except hydrofluoric. It may be obtained in pure form by placing tissue containing it in a drop or two of concentrated sulfuric acid and after a time treating with successively stronger solutions of chromic acid (starting with 25 per cent.) and then washing with water and alcohol.

In a number of members of the *Palmæ*, *Orchidaceæ*, *Scitamineæ*, etc. the strands of sclerenchyma fibers are accompanied by rows of small, tongue-shaped cells called *stigmata*.

In these cells the walls adjacent to the fibers are thickened while the opposite walls bulge outwards. Each *stegma* possesses a mass of silica which usually fills the cell lumen.

15. TANNINS.—Tannins are amorphous substances occurring in plants having an astringent taste, and turning dark blue or green with iron salts. They occur in greatest quantity in the bark of exogens, and in gall formations. They are soluble in water, alcohol, glycerine, and a mixture of alcohol and ether. They are almost insoluble in absolute ether and chloroform. They give insoluble precipitates with organic bases such as alkaloids and with most of the salts of the heavy metals.

According to their behavior with solution of iron chloride or other soluble iron salts, two kinds of tannic acid are recognized: (*a*) a form of tannic acid giving a bluish-black color, as that which is found in *Rhus*, *Castanea*, *Granatum*, *Galla*, etc.; (*b*) another tannic acid producing a dark green coloration, as that found in *Krameria*, *Kino*, Mangrove bark, *Quercus*, *Catechu*, etc.

If sections are placed in a 7 per cent. solution of copper acetate for a week or more, then placed on a slide in 0.5 per cent. aqueous solution of ferric chloride, and after a while washed with water and mounted in glycerin, an insoluble brownish precipitate will be produced in those cells containing tannin.

Braemer's reagent (Sodium tungstate 1 Gm. and Sodium acetate 2 Gm. dissolved in 10 cc. of water) is superior to solutions of iron salts in the detection of tannin, since some other plant constituents give a dark-green or bluish-black color with iron. It gives a yellowish-brown precipitate.

16. PROTEINS.—Proteins are complex nitrogenous substances forming the most important of the reserve foods of plants. They are found in the cytoplasm, nucleus and plastids of all the living and many of the dead cells of plants, although most abundant in seeds. They occur as crystalline or amorphous bodies or in colloidal solution. Protoplasm, itself, is composed largely of these substances. They all contain carbon, hydrogen, oxygen, nitrogen and sulfur, and many contain in addition phosphorus. They are formed by the addition of nitrogen, sulfur and frequently phosphorus to elements of grape sugar. The nitrogen, sulfur and phosphorus elements are obtained from nitrates, sulfates and phosphates which are dissolved in the water taken in through the roots. All living plant cells are capable of synthesizing proteins from carbohydrates. The process of protein

synthesis appears to involve the following steps: (1) the union of molecules of dextrose or some other simple sugar to nitrogen to form *amino acids*; (2) the combining of amino acids to form *polypeptides*; (3) the building up of polypeptides into *simple proteins*; (4) the union of phosphorous or sulfur to simple proteins to form *complex proteins*. The names of proteins recorded may be found by the hundreds. These are grouped into chemical classes, the most important of which from the standpoint of their occurrence in plants are the *globulins*, *albumens*, *glutelins*, *nucleins*, and *gliadins*. Of these the *globulins* are found most extensively. Globulins are insoluble in water but soluble in sodium chloride solutions. They do not coagulate upon the application of heat.

Albumens are soluble in water and coagulate with heat.

Glutelins are insoluble in water, sodium chloride solution and strong alcohol.

Gliadins are nearly or wholly insoluble in water but soluble in 70 to 90 per cent. alcohol.

Nucleins are insoluble in water but soluble in alkaline solutions.

The following tests are of value in determining the presence of proteins:

Lugol's solution (made by dissolving 5 Gm. of iodine and 10 Gm. of potassium iodide in enough water to make product weigh 100 Gm.) stains proteins yellow or brown

Concentrated nitric acid stains proteins yellow. This color becomes deeper upon the addition of ammonia water.

Milon's reagent (made by dissolving 3 cc. of mercury in 27 cc. of fuming nitric acid without heat and adding an equal volume of distilled water) stains proteins a brick-red.

Concentrated solution of nickel sulfate colors proteins yellow or blue.

If sections are placed for an hour or two in a solution of 1 Gm. of sodium phospho-molybdate in 90 Gm. of distilled water and 5 Gm. of nitric acid, the proteid substances appear as yellowish granules.

The *globulins* (phytoglobulins) frequently occur in bodies called "aleurone grains."

ALEURONE GRAINS

Aleurone grains are small bodies found in seeds particularly those containing oil, and like starch grains often are characteristic

of the genus or species. They are produced by the drying up of protein-containing vacuoles. Each aleurone grain consists of a ground substance (composed of amorphous proteid matter soluble in water, dilute alkali or acid), in which are usually embedded one or more *crystalloids* or *phyto-globulins* (insoluble in cold water, but soluble in less than 1 per cent. solution of an alkali, in dilute HCl and acetic acid), one or more transparent, globular *globoids* composed of Ca and Mg phosphate (insoluble in water and dilute potash solution but soluble in 1 per cent. acetic acid solution), and frequently a crystal of *calcium oxalate*, the whole being enclosed by a *protoplasmic membrane* (soluble in water). (Fig. 53B.)

FIG. 53.—To show Aleurone Grains. A, cells from cotyledon of seed of garden bean; n, aleurone grain; m, starch; B, cell from endosperm of castor bean; a, aleurone grain; l, ground substance; k, phytoglobulin; i, globoid. (A, After Sachs; B, After Frank.)

The proteins insoluble in the cell-sap water are made soluble for translocation by means of *proteolytic enzymes* which change them into proteoses and peptones.

17. MUCILAGES AND GUMS are those substances occurring in plants which are soluble in water or swell in it, and which are precipitated by alcohol.

Mucilage is formed in plants in several ways, viz.; either as a product of the protoplasm, as a disorganization product of some of the carbohydrates, as a secondary thickening or addition to the cell wall, or as a metamorphosis of it. In the first two cases the mucilage is called *cell-content mucilage*; in the last two, *membrane mucilage*.

Mucilage is stored as reserve food in the tubers of Salep and many other Orchids and also in the seeds of some species of the *Leguminosæ* as Fenugreek, etc.

CELL-CONTENT MUCILAGE has been found in the leaves of *Alcæ*, the rhizomes of *Triticum*, the bulb scales of *Squill* and *Onion* and in certain cells of many other Monocotyledons, especially those containing raphides.

MEMBRANE MUCILAGE has been observed in *Buchu* leaves, *Elm* bark, *Psyllium* seeds, in *Althæa*, *Linum*, *Astragalus*, and *Acacia* species, in the Blue-green Algæ, and many of the Brown and Red Algæ.

GUM.—When mucilage is collected in the form of an exudate from shrubs and trees it constitutes what is termed a *gum*. Chemically gums consist of glucosidal acids combined either partly or completely with calcium, potassium and magnesium. Many gums are used in pharmacy, medicine, and the arts. The four most important from a pharmaceutical standpoint are: ACACIA, yielded by *Acacia Senegal* and other species of *Acacia*; TRAGACANTH, yielded by *Astragalus gummifer* and other Asiatic species of *Astragalus*; CHERRY GUM, obtained from *Prunus Cerasus* and its varieties; and RED GUM obtained from *Eucalyptus rostrata* and some other species of *Eucalyptus*.

Mucilage may be demonstrated in plant tissues containing it by placing sections of these in a deep blue solution of methylene-blue in equal parts of alcohol, glycerin and water on a glass slide, allowing them to remain in the solution for several minutes, then draining off the stain and mounting in glycerin. Those cells containing mucilage will exhibit bluish contents of deeper aspect than other elements.

18. FIXED OIL AND FATS.—These are fatty acid-esters of glycerin which are found in the vacuoles of cells or formed with the cell walls from which they may be liberated as globules upon treating sections with chloral hydrate or sulfuric acid or heating them. The fats differ from the fixed oils only in that they tend to solidify and crystallize at ordinary temperatures. They are quite soluble in ether, chloroform, benzol, acetone and volatile oils but insoluble in water, and, with the exception of castor oil, insoluble in alcohol. They are readily distinguished from the volatile oils in that they leave a greasy stain upon paper which does not disappear. Fixed oils and fats take a brownish to black color with osmic acid, a red color with alkannin or Sudan III and a blue color with cyanin. In *Vaucheria*, the *Diatoms* and a few of the other *Thallophytes*, fixed oil is formed

in the chromatophores instead of starch as the first visible product of photosynthesis. In higher plants it is generally found in storage regions, such as the parenchyma of seeds, fruits and the medullary-ray cells and the parenchyma of barks, roots and rhizomes.

19. WAXES.—These are esters of fatty acids which differ from fixed oils and fats in not forming glycerin when saponified, but rather certain of the higher monatomic alcohols which are insoluble in water.

The characteristic bloom on the epidermis of fruits of junipers, plums and grapes and the glaucous surfaces of many leaves are due to waxy coatings. Three types of waxy covering occur on plants. The most common type consists of densely crowded granules as seen on the leaves and stems of many lilies, grasses and irises. On the nodes of the sugar cane the covering assumes the form of a layer of vertical rods, rolled or hooked at their distal ends [Fig. 68(3)]. On the leaves of the houseleek (*Sempervivum*) and the wax palms it occurs as a crust interspersed with cracks. On the fruits of the wax myrtle and bayberry this wax crust is secreted by glandular emergences.

Wax serves as an effective control over transpiration by the epidermis. It also gives rigidity to the epidermis, lessening the compression and distortion of subjacent parenchyma which is apt to occur from its contraction during loss of water.

The more important vegetable waxes of commerce are Carnauba Wax, from the leaf buds and young leaves of *Copernicia cerifera*, the Brazilian Wax Palm, Myrtle or Bayberry Wax from the fruits of *Myrica cerifera*, *M. carolinensis* and *M. xalapensis*, and Japan Wax from the fruits of *Rhus succedana*, *R. vernicifera* and *R. sylvestris*.

20. VOLATILE OILS.—These are volatile odoriferous principles found in various parts of numerous plants which arise either as a direct product of the protoplasm or through a decomposition of a layer of the cell wall which Tschirch designates a "resinogenous layer." They are readily distilled from plants, together with watery vapor, are slightly soluble in water, but very soluble in fixed oils, ether, chloroform, glacial acetic acid, naphtha, alcohol, benzin and benzol. They leave a spot on paper which,

however, soon disappears. They respond to osmic acid, alkanin, Sudan III, and cyanin stains similar to the fixed oils and fats.

Volatile oils may be grouped into four classes:

A. PINENES or TERPENES, containing carbon and hydrogen and having the formula of $C_{10}H_{16}$. Examples: Oil of Turpentine and various other volatile oils occurring in coniferous plants.

B. OXYGENATED OILS, containing carbon, hydrogen and oxygen. Examples: Oil of cassia and other cinnamons.

C. NITROGENATED OILS, containing carbon, hydrogen and oxygen with nitrogen (from HCN). Example: Oil of Bitter Almonds.

D. SULFURATED OILS, containing carbon, hydrogen and sulfur. Example: Volatile oil of mustard.

21. RESINS, OLEORESINS, GUM RESINS, AND BALSAMS.—These substances represent products of metabolism in many plants which are formed either normally, as Turpentine, Asafoetida, Mastiche, etc., or as a result of pathological processes through injury to the plant tissues, as Storax, Benzoin, Balsam of Tolu and Peru, etc. They occur usually in special cavities such as secretion cells, glands, or secretion reservoirs.

RESINS are amorphous, solid or semisolid, fusible substances insoluble in water but mostly soluble in alcohol and other organic solvents. They combine with alkalies to form soap. Many of them are oxidized oils of plants. Examples: Guaiac, Rosin, Mastic.

OLEORESINS are secretions composed of resins dissolved in volatile oil. Examples: Turpentine, Copaiba, Balsam of Fir and Venice Turpentine.

GUM RESINS are natural compounds of resin, gum and oil. Examples: Asafoetida, Myrrh, Gamboge.

BALSAMS are mixtures of resins with cinnamic or benzoic acid or their esters and frequently both, the esters being generally called a volatile oil. Examples: Balsam of Tolu, Storax, Balsam of Peru.

If sections of a resin-containing plant part are placed in a saturated aqueous solution of copper acetate for a week or two and mounted in dilute glycerin, the resin will be stained an emerald green.

22. PIGMENTS.—These are substances which give color to various plant parts in which they are found. They occur either

in special protoplasmic structures, as chloroplasts, chromoplasts or chromatophores, or dissolved in the cell sap. Of the pigments named, the following will be considered: Chlorophyll, Xanthophyll, Chromophyll, Carotin, Etiolin, Anthocyanin, Phycocyanin, Phycophæin, and Phycoerythrin.

FIG. 54.—Absorption spectra of five different concentrations of chlorophyll *a*. The strongest is represented by the lowest spectrum. (From Palladin after Willstätter and Stoll.)

CHLOROPHYLL is a collective name given to the two green pigments found in the chloroplastids or chromatophores of leaves or other green parts of plants. Its composition is not definitely known although it yields products similar to the hæmoglobin of the blood when decomposed. Iron is known to be essential to its formation and Magnesium, Carbon, Hydrogen, Oxygen and Nitrogen are components of it. If green leaves be boiled in water to kill the plasma membranes, the water poured off, strong alcohol added and the leaf material allowed to macerate in the alcohol for some time, an alcoholic extract of the green leaves can be obtained. If a portion of this alcoholic extract of green leaf be diluted with water to an 85 per cent. alcoholic solution and xylene or benzine be gradually added and the mixture shaken, the pigments in solution will break up

into a yellowish and a greenish portion. The greenish portion dissolves in the xylene, which rises, forming the upper stratum, while the yellowish portion dissolves in the alcohol and water forming the lower stratum. By repeatedly shaking and adding alcohol to the greenish fraction it can be entirely separated from the yellowish fraction. The isolated greenish portion of alcoholic leaf extract has been found to consist of two green pigments called *chlorophyll a* and *chlorophyll b*. When separated and dissolved in alcohol, *chlorophyll a* appears blue-green and *chlorophyll*

FIG. 55.—Absorption spectrum of five different concentrations of chlorophyll *b*. The strongest is represented by the lowest spectrum. (From Palladin after Willstätter and Stoll.)

b, yellowish-green in transmitted light. By reflected light, the former is blood-red, the latter brownish-red. *Chlorophyll a* constitutes 72 per cent. by weight of the green pigments and has, as so far known, the formula of $C_{55}H_{72}O_5N_4Mg$. *Chlorophyll b* represents the remaining 28 per cent. and has, as far as is known, the formula of $C_{55}H_{70}O_6N_4Mg$. The isolated yellowish portion of the leaf extract consists of two pigments, *carotin* and *xanthophyll*.

Alcoholic solution of chlorophyll, when examined spectroscopically in several concentrations, shows absorption bands in the red, orange, yellow and green with a diffuse band in the

violet-blue zone of the spectrum, the sharpest band being in the red.

CHROMOPHYLL is a broad term applied to either the yellow or orange pigments found in chromoplastids. Sulfuric acid forms a blue color with chromophyll.

CAROTIN or carotene is an orange to orange-red pigment which has the chemical formula of $C_{40}H_{56}$. It is found accompanying the chlorophyll within chloroplasts, but exists separately in chromoplastids of leaves, flowers, fruits and seeds, as well as in subterranean parts of higher plants. It is especially abundant in the roots of carrots. It is readily soluble in chloroform, ether and carbon bisulfide, slightly soluble in hot alcohol and insoluble in water. When examined with a spectroscope, it shows two dark bands in the green-blue half of the spectrum. Carotin is a source of Vitamin A, the growth-promoting factor. It is changed into Vitamin A in the body.

XANTHOPHYLL is a yellow pigment which accompanies the chlorophyll in the chloroplast. Chemically it represents an oxidation product of carotin, having the formula of $C_{40}H_{56}O_2$.

ETIOLIN is a pale yellow pigment which appears when green plants are kept for some time in darkness. It is probably identical with xanthophyll.

ANTHOCYANINS are applied to the blue, purple and red pigments which occur in the cell sap of floral organs, stems, leaves, roots and fruits. The character of the color is claimed to be due to the alkalinity or acidity of the cell sap.

PHYCOCYANIN is the blue pigment found in the blue-green algæ, associated with chlorophyll. It is soluble in water and glycerin but insoluble in ether and alcohol.

PHYCOPHÆIN or FUCOXANTHIN is the brown pigment found in the brown algæ. It is readily soluble in water.

PHYCOERYTHRIN is the red pigment found in many of the red algæ and some of the blue-green algæ. It is readily soluble in water but insoluble in ether and alcohol.

The last two pigments are always associated with chlorophyll but frequently conceal it.

23. LATEX.—This is an emulsion of varying composition and color found in special passages, as latex cells and laticiferous

vessels, of many plants. It may contain starch, sugar, proteid, oil, enzymes, bitter principles, tannins, alkaloids, gum, resins, caoutchouc and mineral salts. The color may be absent, as in *Oleander*; whitish, as in *Asclepias*, *Papaver*, *Hevea*, and *Apocynum*; yellowish to orange, as in *Celandine*, or red, as in *Sanguinaria*.

Chlor-zinc-iodine solution imparts to latex a wine-red color.

The latex of the following plants is of value to pharmacy and the arts:

Papaver somniferum and its variety *album* which yields OPIUM. That from the unripe capsules is alone used for this drug.

Palaquium Gutta and *Payena* spp. which yields GUTTA PERCHA.

Hevea species, *Ficus elastica*, *Landolphia* species, *Castilloa elastica*, *Hancornia speciosa*, *Forsteronia* species, *Funtumia elastica* and *F. africana*, *Manihot Glaziovii*, *Clitandra* species and various species of *Urceola* and *Euphorbia* furnish most of the RUBBER of commerce.

Lactuca virosa and other species of *Lactuca* yield the drug LACTUCARIUM.

Euphorbia resinifera, contains a white resinous latex which is gathered as the drug EUPHORBIIUM.

24. ENZYMES.—An *enzyme* or soluble *ferment* or unorganized ferment (according to Hepburn) is a soluble organic compound of biologic origin functioning as a thermolabile (sensitive to heat) catalyst in solution. Ostwald has defined a catalyst as an agent which alters the rate of a reaction without itself entering into the final product, or which does not appear to take any immediate part in the reaction, remains unaltered at the end of the reaction and can be recovered again from the reaction product unaltered in quantity and quality. The biologic catalysts (enzymes) differ from the inorganic catalysts in that they are sensitive to heat and light. They are destroyed quickly at 100°C. and most of them cannot be heated safely above 60°C. The optimum temperature range for this activity usually is from 40° to 45°C. Enzymes are soluble in water, dilute alcohol, glycerin or dilute saline solutions and are precipitated by concentrated alcohol. They are colloidal and non-diffusible. They are stimulated to activity by substances known as “*activators*” and their activity is checked by other substances called

“paralyzers.” Frequently the paralyzers consist of products of enzyme action. Cold inhibits and warmth accelerates enzyme action. Moisture must always be present for enzymic activity.

CLASSIFICATION OF ENZYMES

A. ACCORDING TO DIFFUSIBILITY THROUGH CELL WALL.

ENDOCELLULAR: Those that cannot diffuse out of the cell.
Example: Zymase of Yeast.

EXTRACELLULAR: Those that can diffuse out of the cell.
Example: Invertase of Yeast.

B. ACCORDING TO KIND OF SUBSTANCES ACTED UPON AND TRANSFORMED.

1. CARBOHYDRATE ENZYMES:

DIASTASE, found in the germinating seeds of barley and other grains and in *Aspergillus oryzae*, etc., converts starch to maltose and dextrin. As observed under the microscope, starch grains partly transformed by diastase (*amylase*) show depressions, later deepening to form irregular tunnels. Such grains are called *corrosion starch*.

INVERTASE, secreted by yeasts, and found in younger parts of higher plants, transforms cane sugar, producing dextrose and levulose.

MALTASE, found in malt, in green leaves and *Saccharomyces octosporus*, transforms maltose to dextrose.

TREHALASE, found in *Polyporus*, hydrolyzes trehalose to dextrose.

CYTASE, found in *Nux Vomica* and *Ignatia* seeds, in barley, dates, etc., decomposes hemicellulose and cellulose to galactose and mannose.

LACTASE, found in Kephir grains, hydrolyzes lactose to dextrose and galactose.

PROTOPECTINASE splits protopectin of the primary cell wall to soluble pectin.

PECTASE hydrolyzes the methyl alcohol from soluble pectin to produce pectic acid.

PECTINASE, found in fruits and other plant organs, transforms the soluble pectin, pectic acid or pectates of the middle lamella into galactose, arabinose and galacturonic acid.

INULASE, found in plants of the Daisy and Bluebell families, etc., transforms inulin to levulose.

ZYMASE, found in yeast, hydrolyzes glucose (dextrose and levulose) to alcohol and carbon dioxide according to the following equation: $C_6H_{12}O_6 = 2C_2H_5OH + 2CO_2$.

2. FAT AND OIL ENZYME:

LIPASE splits up fats and oils into fatty acids and glycerin. It is found in various mildews, molds and numerous oily seeds and other fatty-oil storage regions of higher plants.

3. PROTEINACEOUS ENZYMES:

PEPSIN converts proteids into proteoses and peptones.

TRYPSIN, found in yeast and other fungi, as *Boletus edulis*, *Amanita* spp., etc., resolves proteins to peptones and amino-acids.

BROMELIN, found in the fruit of the Pineapple, and PAPAYIN (Papain), found in the latex of the fruit of the Papaw, act similarly to trypsin.

NEPENTHIN, found in the pitchers of *Nepenthes* species, acts similarly to pepsin.

4. GLUCOSIDE ENZYMES:

EMULSIN (synaptase), found in the seeds of the Bitter Almond, Cherry Laurel leaves, in the barks of the Wild Black Cherry and Choke Cherry and in other Rosaceous plant parts, in *Manihot utilissima*, *Polygala* species, etc., hydrolyzes the glucoside present (either amygdalin or l-mandelonitrile glucoside) to hydrocyanic acid, benzaldehyde and glucose.

MYROSIN (myronase), found in the seeds of the black mustard (*Brassica nigra*) and other members of the *Cruciferae*, converts the glucoside, sinigrin, into allyl-iso-sulphocyanide and glucose.

RHAMNASE, found in Alder Buckthorn (*Rhamnus Frangula*) and probably other species of *Rhamnus*, hydrolyzes the anthra-glucoside frangulin to rhamnose and emodin.

GAULTHERASE, found in Wintergreen (*Gaultheria procumbens*) and other Ericaceous plants, resolves the glucoside, gaultherin, to methyl-salicylate and glucose.

5. ORGANIC ACID ENZYMES:

REDUCTASE of yeast decomposes lactic acid to pyrotartaric acid and hydrogen. It also decomposes formic acid into carbon dioxide and hydrogen.

CARBOXYLASE, found in higher plants, splits pyrotartaric acid into acetic aldehyde and carbon dioxide.

C. OXIDIZING ENZYMES.

These are enzymes which absorb molecular oxygen from the air and combine it with plant constituents.

PEROXIDASE occurs in many plants including apples, many fleshy fungi, potatoes, etc.

PEROXIDASES are common in plants and are responsible for discolorations seen in the tissues of many barks and fruits when they are bruised or broken and exposed to the air. The juices and extracts of plants containing oxidases turn brown upon exposure to air. This is owing to the formation of organic peroxides by the action of atmospheric oxygen on phenolic substances present and the presence of a peroxidase which, with an oxygen acceptor, brings about oxidation with a resulting colored oxidation product.

25. HORMONES.—The term “hormone” means chemical messenger and is appropriately named because it produces its effect at a distance from its point of origin. Hormones are substances which govern the proper development or functioning of organisms and organs. The hormones of animals, sometimes termed “internal secretions,” are formed by the endocrine glands as the pituitary, thyroid, parathyroid, adrenal, pineal, pancreas, ovaries, testes, and are secreted directly into the blood, lymph and cerebro-spinal fluid.

The hormones of plants, while anticipated for some time, remained unknown until 1928, when F. W. Went demonstrated the existence of a plant growth hormone controlling the enlargement of the coleoptile of the oat seedling. In 1933, F. Kögl and coworkers isolated 3 plant-growth hormones in crystalline form, *auxin a* ($C_{18}H_{32}O_5$) which they obtained from malt, maize germ oil and human urine, *auxin b* ($C_{18}H_{30}O_4$) and *hetero-auxin* from vegetable material. Hetero-auxin has been identified as indole-3-acetic acid. Their formation and distribution has been found to be influenced by light. They regulate growth in plants.

When seedlings are illuminated on one side, the growth hormone moves from the illuminated side to the shaded side.

The result is that the cells elongate more rapidly on the shaded side, bending the plants toward the light. It has been recently shown that though growth-hormones are produced in the light, they function better in the dark.

26. VITAMINS.—Vitamins are essential food principles for the most part manufactured by plants without which animals cannot maintain their health. They are designated alphabetically as A, B₁, B₂ or G, B₃, B₄, B₅, C, D, E, and K and by names indicative of the deficiency disease in animals they correct.

VITAMIN A, also known as Fat-soluble A, Anti-xerophthalmic, and *Anti-infective Vitamin*, was discovered by McCollom and Davis in 1915. It occurs in conspicuous amounts in carrots, green leafy vegetables, in a number of fruits, as tomatoes, prunes, pineapple and banana, in sweet potatoes, in wheat germ, yellow corn meal, whole milk and in cod and halibut liver oils. In its chemical and physiological properties, Vitamin A is closely related to carotene which occurs in carrots and many other plants. Its presence in oils can be detected by the addition of a few drops of a chloroform solution of antimony trichloride to the oil when a blue color appears which deepens and then fades. This vitamin protects epithelial tissues against bacterial infection, especially that of the respiratory mucosa, and is essential to the proper growth of the bodies of young animals. Animals deprived of it in their diet lose weight and develop a hemorrhagic condition in their eyes known as xerophthalmia. The U.S.P. unit of vitamin A is the amount of vitamin contained in $\frac{1}{3000}$ of a grain of the U.S.P. Reference Oil taken as a standard.

VITAMIN B₁ or *Anti-neuritic Vitamin* is a water-soluble, thermolabile factor which has been isolated in pure crystalline form from rice polishings. Its absence in the diet has been found to be the cause of beri-beri, or polyneuritis, a disease of the nervous system characterized by muscular incoordination followed by paralysis of certain sets of muscles. Its presence in the animal's diet promotes growth by stimulating the metabolic processes, protects it from nerve disease and stimulates appetite and digestion. It is also essential for normal gestation. This vitamin occurs abundantly in yeast, rice polishings and wheat

germ and to a lesser extent in whole rice, rye, wheat, barley, milk, eggs and a number of vegetables and fruits.

VITAMIN B₂ or G, sometimes called the *Anti-pellagric factor* of the Vitamin B complex, prevents pellagra, a disease characterized by soreness of the mouth and tongue, diarrhea, skin eruptions and disturbances of digestion and mentality. It is water-soluble and little affected by heat. It occurs in yeast, in spinach, kale and other green vegetables, in potato, banana, fresh carrots, cabbage, beets, tomatoes, lettuce, wheat bran, meat, etc.

VITAMINS B₃, B₄, and B₅ are all growth-promoting principles which have been found to occur in yeast. They are water-soluble and thermolabile.

VITAMIN C or *Antiscorbutic Vitamin* is the factor which prevents scurvy. It is water-soluble and destroyed by heat. It has been found to be ascorbic (cevitamic) acid, C₆H₈O₆ and has been made synthetically. This vitamin is required for the proper metabolism of bones and the formation and nutrition of teeth. It occurs abundantly in the juice of oranges, lemons and limes, in lettuce, spinach, cabbage and tomatoes and to a less extent in pineapple, raspberry, strawberry, peach, beans, corn, and some other fruits and vegetables.

VITAMIN D or *Anti-rhachitic Vitamin* is essential for the prevention of rickets, since it regulates the absorption, retention and deposition of calcium and phosphorous by the bones and teeth. It has been found to be irradiated ergosterol. It is formed in animals by the action of ultra-violet light upon cholesterol present in the skin. This vitamin has not been found in plants. It is soluble in oil and relatively stable to heat. It occurs abundantly in cod-liver and halibut liver oils and in smaller amounts in salmon, egg-yolk, oysters, clams, etc. The U.S.P. unit employed in the biological standardization for vitamin D is that amount of vitamin contained in $\frac{1}{95}$ of a gram of the Reference Oil taken as a standard.

VITAMIN E or *Anti-sterility Vitamin* is a fat-soluble principle which has been found essential for the normal maturation of the germ cells of the male animal and for the maintenance of the placenta in the female. Its most abundant source is wheat

germ oil. It also occurs in lettuce, water cress, whole cereals, nuts, in cocoanut, corn and cottonseed oils, meat, etc.

VITAMIN K or *Anti-hemorrhagic Vitamin* has been found to represent the substance necessary in the diet of chickens (and possibly other animals) for the prevention of multiple hemorrhages. It has been found in the fat of hog liver, in tomatoes, kale and hemp seed.

CELL WALLS

The cell walls of plants make up the plant skeleton. They are all formed by the living contents of the cells (protoplasts) during cell-divisions. In most plants the cell wall when first formed (primary cell wall) consists of pectose or a related carbohydrate. Later the daughter protoplasts secrete two layers of a mixture of *cellulose*, $(C_6H_{10}O_5)_n$, and protopectin, one on either side of a middle lamella, the latter composed of calcium pectate. Each of these layers on either side of the middle lamella represents a *secondary cell wall*. A *tertiary wall* may be laid down by the protoplast beneath the secondary wall. (See Fig. 14.)

The cell walls may remain of such composition or become modified to meet certain functions required of them. Thus, in the case of outer covering cells as epidermis and cork, whose function is that of protecting the underlying plant units, the walls become infiltrated with *cutin* (in the case of epidermal cells), *suberin* (in the case of cork cells), waxy-like substances, which make them impermeable to water and gases, as well as protect them against easy crushing. Again, in the case of stone cells and sclerenchyma fibers whose function is that of giving strength and support to the regions wherein found, the walls become infiltrated with *lignin* which increases their strength, hardness, and in the case of sclerenchyma fibers, their elasticity also. Moreover, in the case of the cells comprising the testa or outer seed coat of the pumpkin, squash, mustard, flax, psyllium, etc., whose function is that of imbibing quantities of water, the walls undergo a *mucilaginous modification*. These, upon the imbibition of water, swell up and form layers of mucilage within the cell cavities.

The sub-epidermal cells beneath the upper epidermis of Buchu leaves exhibit a striking example of *mucilaginous modi-*

fication of cellulose as to their walls. If transverse sections of either fresh or dried Buchu leaf-blades are first mounted in strong alcohol and observed under the microscope, the walls of the sub-epidermal cells appear no different from ordinary cellulose walls; if, however, the alcohol be gradually withdrawn from beneath the cover slip and gradually replaced with water, these walls will be seen to swell and form layers of mucilage in the cell lumina. The mucilage formed absorbs water so rapidly that, in a relatively short time, the pressure (turgor) within the cells becomes so great as to cause a rupture of the less elastic vertical walls of the sub-epidermis with a consequent separation of the upper epidermal cells from those of the palisade layer beneath and the appearance of a broad rent in the section.

In pathological conditions, as a result of natural or induced injury to the tissues, the walls of cellulose and pectose may be completely transformed to mucilage. This condition is called *gummosis*. The gum drugs Acacia and Tragacanth are produced in this way.

The walls of some bacteria, other fungi and some algae are infiltrated with *chitin*. Those of the outer layers of stems of grasses, sedges and horsetails contain silica deposits, while many plant walls contain calcium oxalate and calcium carbonate.

GROWTH IN AREA AND THICKNESS.—The cell wall when first formed is limited in both extent and thickness. As the protoplast within enlarges, new particles are placed within the wall by the process called *intussusception*. This increases its area. New particles, also, are deposited on its surface which gradually increases its thickness. The latter process is known as *growth by apposition*.

PORES (PITS).—Pores or pits are small thin spots or holes in the cell wall. The following terms are used to describe the character of these structures:

Beaded. Circular pores arranged close together in a single row.

Bordered. Pores partially covered on either side of the middle lamella by a thickened, overhanging projection of the cell wall.

Branching. Pore canals which divide and sometimes subdivide.

VARIOUS KINDS OF CELL WALLS AND BEHAVIOR OF EACH TO MICROCHEMIC REAGENTS

<i>Nature of wall</i>	<i>Where found</i>	<i>Reagent and behavior toward same</i>
Cellulose	Parenchyma cells, trichomes such as cotton, etc.	Schweitzer's reagent dissolves it. Chlorzinciodine solution imparts a blue or violet color. Iodine solution followed by sulfuric acid colors it blue.
Lignocellulose (Lignified wall).	Woody parts of plants, such as stem cells, bast fibers, wood fibers, etc.	Phloroglucin with HCl imparts a red color except to bast fibers of flax, mezereon, etc. Corallin-soda solution imparts a pink color. Aniline sulphate with H ₂ SO ₄ colors it a golden-yellow. Chlorzinc-iodine imparts a yellow color.
Reserve cellulose.	Found in certain seeds such as nux vomica, ignatia, ivory nut, date, coffee, etc.	As for cellulose.
Mucilaginous modification of cellulose.	In various parts of plants.	Alcoholic or glycerin solution of methylene-blue imparts a blue color.
Suberized walls . .	In cork, wounded areas of plants, endodermis.	Alcoholic extract of chlorophyll, in the dark, imparts a green color. Alcannin and Sudan III impart a red coloration. Converted into yellowish droplets and granular masses upon heating with a strong solution of KOH. Sulfuric acid is resisted.
Cutinized walls . .	Forming outer walls of many epidermal cells.	As for suberized walls.
Callus of sieve plates.	Plates of sieve tubes.	Corallin-soda solution imparts pink color.
Silicified walls . . .	Epidermis of Horsetails, Grasses, etc.; Diatoms.	Soluble in hydrofluoric acid.

PORES (continued)

Elliptical. Somewhat elongated pores with rounded ends.

Oblique. Slit-like pores in a slanting position in the cell wall.

Simple. Unbranched or non-bordered pores.

Slit-like. Very narrow, elongated pores.

Transverse. Slit-like pores extending in a direction perpendicular to the lateral cell walls.

CHAPTER VI

PLANT TISSUES

A TISSUE is an aggregation of cells of common source, structure and function in intimate union.

THE TISSUES OF SPERMATOPHYTES AND PTERIDOPHYTES

The tissues of seed plants and pteridophytes are all derived from a fertilized egg (zygote) which has undergone repeated divisions. At first either an apical cell arises (pteridophytes) or a mass of cells is formed (spermatophytes) at the tips of growing regions which are essentially alike and constitute the *primordial meristem*; but gradually we find that a division of labor has become operative setting aside many different groups of cells, each group of which has its particular role to perform in the economy of the whole. Each group of cells similar in source, structure and function is called a tissue. The tissues found in higher plants range from those whose component cells are more or less rounded, in a rapid state of division, and whose thin, cellulose, cell walls enclose a mass of protoplasm, devoid of vacuoles, or with exceedingly small ones to those whose cells through various physical and chemical factors become compressed, elongated, and highly modified in respect to their contents and walls.

CLASSIFICATION OF TISSUES.—Tissues may be classified according to method of development, form, structure, function, and whether simple or complex.

According to their *method of development*, tissues are either primary or secondary. PRIMARY TISSUES are those which develop at the growing points of different organs by cell division. They arise from apical and intercalary meristems. They include *parenchyma*, *epidermis*, *endodermis*, *pericycle*, *primary phloem*, *primary xylem*, and *cambium*.

SECONDARY TISSUES are those which take their origin in a cambium or cork cambium. Examples of secondary tissues

FIG. 56.—Diagram showing the evolution of tissues in a stem from the primordial meristem down to the beginning of cambial activity. In the longitudinal diagram, at the bottom, the initial C of the word cambium stands directly beneath this tissue which is radially but one cell in thickness. (After Stevens.)

are *secondary xylem* and *secondary phloem* which arise from a *primary cambium*, and *cork* and *secondary cortex* which arise from *cork cambium*. These will be discussed more fully in a later part of this chapter.

SIMPLE TISSUES are those which are composed of one kind of cell, as *ordinary parenchyma*, *collenchyma* and *sclerenchyma*, whereas COMPLEX TISSUES are made up of two or more kinds of cells as *xylem* and *phloem*.

The adult or permanent tissues represent those which have been laid down by the cell-division of meristems.

PRIMARY MERISTEMS.—As was shown by Hanstein,¹ the embryo of Angiosperms, while still constituted of only a few cells in the process of division, becomes differentiated into three layers of cells which differ in their arrangement and direction of division; these were called by him *Dermatogen*, *Periblem* and *Plerome*. In roots a fourth layer of cells is sometimes evident at the apex. This was termed by Janczewski² the *Calyptrogen* layer. These primary layers or groups of cells are called *primary meristems* or *generative tissues*. They are composed of more or less rounded cells having delicate cell walls of cellulose which enclose protoplasm and nucleus, and wherever found in living embryos are in a rapid state of division.

The primary meristems or generative tissues are found in the growing apices of plant organs, such as root, stem and leaf apex. By the division and redivisions of their cells they give rise to the mature or adult tissues of plants.

1. DERMATOGEN OR PROTODERM originates epidermal tissue and derivative structures such as stomata, non-glandular and glandular hairs, glands, and sometimes cork cambium.

2. PERIBLEM OR GROUND MERISTEM originates primary cortex tissue, chlorophylloid cells (*chlorenchyma*), colloid cells (*collenchyma*), strengthening cells (*sclerenchyma*), crystal cells (*raphiderchyma*), latex cells (*lacterchyma*), pericycle, endodermis, sometimes cork cambium and pith.

3. PLEROME OR PROCAMBIUM originates the primary or first fibrovascular bundles, the cambium and sometimes the pith.

According to the structure of the component cells and their location, the following tissues are found in various forms of higher plants:

¹ Hanstein, "Die Scheitelzellgruppe im Vegetationspunkt der Phanerogamen," Bonn, 1868.

² Am. Sci. Nat. 5 série, tom. xx.

1. MERISTEM	8. CORK
2. PARENCHYMA	9. LATICIFEROUS TISSUE
3. COLLENCHYMA	10. SIEVE OR CRIBIFORM TISSUE
4. SCLERENCHYMA	11. TRACHEARY TISSUE
5. EPIDERMIS	12. MEDULLARY RAYS
6. ENDODERMIS	13. GLANDS AND SECRETION RESERVOIRS
7. PERICYCLE	14. NECTARIES

MERISTEM

MERISTEM, frequently called embryonic tissue, is undifferentiated tissue composed of cells in the state of rapid division. The first kind of meristem to appear arises from the division of apical cells at or near the apices of roots, stems and leaves and is in these regions called PRIMORDIAL MERISTEM. The primordial meristem gives rise through cell divisions and slight differentiation to the PRIMARY MERISTEMS (*plerome*, *periblem* and *dermatogen*) which in turn develop the primary permanent tissues of plants. Primary meristems retain the power of independent growth and capacity for division as long as the plant part survives which contains them. Meristem is also found in other regions of plant organs such as the cambium and cork cambium, and is there called SECONDARY MERISTEM. Secondary meristem is a derivative of primary meristem. It gives rise to the secondary permanent tissues. It loses with its transition to the permanent tissues the power of division and independent growth.

CAMBIUM is a secondary meristem lying between the phloem and xylem in certain collateral bundles (intrafascicular cambium) or between the phloem and xylem portions of the medullary rays (interfascicular cambium). It is cylindrical and extends longitudinally through the plant body. The cambium is found in roots and stems of gymnosperms and dicotyledons where it divides to form secondary xylem and secondary phloem. These formations are responsible for the increase in diameter of these organs. (Cf. Fig. 76.)

CORK CAMBIUM or *phellogen* originates in the pericycle of roots and in the epidermis or outermost layer of the cortex of stems. Through division of its cells it cuts off cork on its outer face and frequently secondary cortex on its inner face. (See Fig. 70.)

According to the position of the meristems in the plant, they may be classified as *apical*, *intercalary* and *lateral*.

APICAL MERISTEMS consist of growth cells which occur at the tips of roots and stems. They produce *growth in length* by giving

FIG. 57.—The cambium and medullary rays in tangential-longitudinal section. $\times 50$. The section has been cut through the cambium zone adjacent to the phloem or inner bark of the stem of the tulip poplar, *Liriodendron tulipifera*, and shows besides the elongated taper-ended cambial cells, phloem parenchyma which has been cut off by the transverse divisions of cambial cells, also medullary rays. *ca*, cambial cell, in the upper portion of which two phloem cells have been cut off; *ca'*, cambial cell at left of which is a row of phloem parenchyma; *r*, medullary ray.

rise to *primary tissues*. In the bryophytes and pteridophytes the apical cell represents apical meristem, while in spermatophytes a group of apical cells makes up this tissue.

INTERCALARY MERISTEMS represent parts of apical meristems which have been separated from the rest of apical meristem by

permanent tissues and left with the latter as the apical meristem advances in growth. They are found just above or below the nodes of many stems and at the bases of some leaves, especially those of monocotyls.

LATERAL MERISTEMS are those which occur lateral in an organ, as the cambium and cork cambium.

PARENCHYMA

PARENCHYMA or FUNDAMENTAL TISSUE is the soft, primitive tissue of plants, consisting of cells nearly equal in length, breadth and thickness (isodiametric) with usually thin, cellulose cell walls enclosing cytoplasm and a nucleus and frequently substances of a non-protoplasmic nature. It is capable of cell division after maturity. There are four generally recognized types of parenchyma, viz.:

ORDINARY PARENCHYMA (SOFT GROUND TISSUE, FUNDAMENTAL TISSUE).—Next to the meristem this is the least modified of all plant tissues. It is generally composed of thin-walled cells, commonly polyhedral or spheroidal in form and often of approximately the same length, breadth, and thickness (isodiametric); the cell walls are composed of cellulose which is usually unmodified. Intercellular air-spaces occur in the angles of adjoining cells. Occasionally the outline of the cells is star-shaped or stellate, as in the stems of Wood Rush or Pickerel Weed or the cells may be several times as long as wide, as in the stems of Geranium, etc. Moreover, markings may occur on the walls. These may be of the nature of pores, as in the parenchyma cells of the pith of the Elder or Sassafras, annular thickenings, as in the Mistletoe, or spiral thickenings, as in certain Orchids. Protoplasm and a nucleus are always present as well as colorless plastids, but in old cells are only seen as a thin layer pushed up against the cell wall. Ordinary Parenchyma may be seen composing the soft tissues of roots, stems, and barks. (See Fig. 69.)

ASSIMILATION PARENCHYMA (CHLOROPHYLL PARENCHYMA, CHLORENCHYMA). This form of parenchyma tissue is found in foliage leaves, floral leaves, in the outer region of young green stems and fruits. Its cells are thin-walled and vary in shape from

more or less isodiametric to irregular and elongated forms. The cells always contain chloroplasts or plastids, in whose pores may be found some other coloring substance. Good examples of assimilation tissue are seen in the green column-shaped cells below the upper epidermis of foliage leaves which, because of their shape, are termed "*palisade parenchyma cells*" and in the green rounded to irregular-shaped cells beneath the palisade parenchyma of foliage leaves which, because they are more loosely arranged around air-spaces, are termed "*spongy parenchyma cells*." (See Fig. 166.)

CONDUCTING PARENCHYMA.—

This type of parenchyma functions in the rapid translocation of food materials to distant regions in the plant. It includes the *wood parenchyma cells* of the *xylem* which convey a portion of the crude sap (water with mineral salts in solution) and the *phloem parenchyma* (soft bast) which transports the elaborated sap (carbohydrate and proteid material in solution). Conducting parenchyma cells differ from those of ordinary parenchyma in being usually more elongated and in conducting soluble food materials with greater celerity.

RESERVE PARENCHYMA.—This resembles ordinary parenchyma in many particulars of structure but differs from it mainly by its cells being filled with starch, protein crystals, or oil globules. It is usually found in seeds, fleshy roots, or underground stems such as tubers, corms, and bulbs.

COLLENCHYMA.—This form of tissue is characterized by its cells being prismatic, more elongated than ordinary parenchyma, with soft, plastic walls of cellulose thickened in areas, usually in their angles, but sometimes on the tangential walls or on the sides abutting air-spaces, with a colloidal substance. The thickened areas are in the form of longitudinal strips. The

FIG. 58.—Transverse section of part of leaf-stalk of a begonia. *e*, Epidermis; *c*, cuticle; *B*, collenchyma, with walls thickened at the angles, *v*; *chl*, chloroplasts. (Sayre after Vines.)

cells, like those of parenchyma tissue, contain cytoplasm and a nucleus, and frequently chloroplasts. It remains alive after maturity (Fig. 58). Collenchyma is generally found underneath the epidermis in the elongating parts of many stems, less often in the midribs and petioles of leaves, and gives support to these regions. It is the first strengthening tissue to develop in stems. It is frequently observed forming the "ribs" of stems and fruits of the Parsley Family and "ribs" of stems of the Mint

FIG. 59.—Stone cells from different sources. 1, From coffee; 2, 3, and 4 from stem of clove; 5 and 6, from tea leaf; 7, 8 and 9, from powdered star-anise seed. (Stevens, after Moeller.)

Family. In many leaves it has been found as the supporting and strengthening tissue between the stronger veins and the epidermis.

SCLERENCHYMA or stony tissue comprises a variety of supporting elements having thickened cell walls usually composed of lignocellulose. When first formed these cells resemble those of ordinary parenchyma in having walls of pure cellulose, but later *lignin*, a hard, woody substance, becomes deposited on the inner surface of the walls in one or more layers. (Occasionally, as in the bark region of stems of Flax and Mezereon, no lignin is

deposited on the walls of the sclerenchyma fibers.) When sclerenchyma is composed of cells which are more or less isodiametric or moderately elongated, with thickened, lignified

FIG. 60.—Stone cells from various sources. 1, From olive pit; 2, from cocoanut endocarp; 3, from flesh of pear; 4, from aconite root; 5, from capsicum; 6, from hazelnut; 7, from allspice. (Drawing by Hoffstein.)

walls and conspicuous pores, its elements are called **STONE CELLS**. Stone cells are distributed in fruits, seeds and barks of many plants, rarely in woods. They occur singly or in small groups or large masses. They have been found forming the

gritty particles in the "flesh" of certain fruits as the Pear, the endocarp or stone region of drupaceous fruits as the Olive, Peach, Cubeb, Pepper, etc., the hard portions of seed coats as in *Physostigma*, Walnuts, etc. Each stone cell presents for examination a cell wall of cellulose with one or several layers of lignin on its inner surface which surround a *central lumen*.

FIG. 61.—Sclerenchyma fibers from different sources. 1, From powdered cinnamon bark; 2, end of bast fiber of flax stem showing transverse markings (*b*); 3, middle portion of flax fiber showing characteristic cross markings at *b*; 4, bast fiber from cinchona bark; 5, branched bast fiber from choke cherry bark; 6, above, end, and below, medium portion of bast fiber of jute. All highly magnified.

The latter is in communication with radial *pore canals* leading outward to the middle lamella. Longitudinal pore canals are also evident.

FIBERS.—When sclerenchyma is composed of cells which are greatly elongated and more or less obtusely or taper ended, its component elements are termed **SCLERENCHYMA FIBERS** or **FIBERS**. These fibers are frequently spindle-shaped. When young they contain protoplasts but when mature their lumina usually contain air. They exhibit pits of the nature of oblique

slits or rounded pores in their walls. They are either polygonal, rectangular or somewhat rounded in transverse section. They

FIG. 62.—Transverse section of a portion of the wood of the Black Ash. *A*, summer wood; *B*, autumn wood; *C*, spring wood. *wf*, wood fibers; *mr*, medullary ray; *tr*, tracheae. $\times 170$. Note the polygonal to rounded-polygonal shape of the wood fibers in cross section.

occur in various parts of roots, stems, leaves, fruits and seeds as supporting elements. When sclerenchyma fibers occur in the xylem region of fibro-vascular bundles, they are termed *Xylem*

Fibers, or *Wood Fibers*; when they appear in the phloem region, *Phloem Fibers*, *Bast Fibers* or *Liber Fibers*; when in the pericycle, *Pericyclic Fibers*; when in the cortex, *Cortical Fibers*. Wood fibers are usually cut off by the cambium only and so are generally

FIG. 63.—Tangential-longitudinal section of wood of the Black Ash. $\times 170$. *wf*, wood fibers; *mr*, medullary rays. Note how the tapered ends of the wood fibers are spliced over each other.

seen in the secondary xylem. They differ from the fibers of the phloem, pericycle and cortex in that they possess bordered pits although in many instances these are so reduced by the thickening of the wall as to be simple. Phloem, pericyclic and cortical fibers possess simple pits.

EPIDERMIS

EPIDERMIS is the outer covering tissue of a plant and is protective in function. It provides against mechanical injury and

loss of water. In old stems and roots of secondary growth it is ultimately replaced by cork. Its cells are living, containing a thin marginal layer of cytoplasm, a nucleus and a large central vacuole and frequently leucoplastids. Non-protoplasmic inclusions as crystals, mucilage, etc., may occur in the lumen. They may be brick-shaped, tabular or polygonal, the vertical wall being equilateral or wavy in outline. Their outer walls are frequently *cutinized* (infiltrated with a waxy-like, waterproof substance called cutin). The radial walls also are sometimes cutinized.

FIG. 64.—Upper epidermis of Sweet Fern (*Comptonia asplenifolia*) leaf (surface view) showing epidermal cells and two non-glandular trichomes.

STOMATA.—Among the epidermal cells of leaves and young green stems may be found numerous pores or *stomata* (sing. *stoma*) surrounded by pairs of crescent-shaped cells, called guard cells. The guard cells contain chloroplasts and are capable of undergoing changes in size and shape dependent upon changes in their turgor. Increased turgor causes them to open. A decrease in turgor causes them to close. The stomata are in direct communication with the sub-stomal air chambers beneath them which in turn are in communication with intercellular space systems of the tissues of the entire plant. Through the stomata an exchange of gases between the tissues of the plant and the external air takes place. The function of the stomata is

to give off watery vapor (transpiration) and take in or give off carbon dioxide, and oxygen. In addition to stomata some leaves possess groups of *water stomata* which differ from the ordinary or *transpiration stomata* in that they always remain open, are circular in outline, give off water in droplets directly, and lie over an area of small, secreting, glandular cells (*epithem*) which is in connection with a bundle of tracheids that springs

FIG. 65.—Radial-longitudinal section through a hydathode from the leaf margin of *Primula sinensis*. *i*, upper, and *j*, lower epidermis; *h*, palisade cell; *e*, thin walled parenchyma called epithem; *g*, intercellular space; *f*, guard cell of a water stoma; *k*, tracheids. (From Stevens, after Haberlandt.)

from the end of one or more fibrovascular bundles. An air chamber usually occurs directly beneath the water pore. Haberlandt gave the name of *hydatode* to this and related types of structures which secrete water in the form of droplets. Examples: Leaves of *Primula*, *Corn*, various grasses, *Potato*, *Elm*, *Larkspur*, *Crassula*, *Saxifraga* and *Ficus*.

EPIDERMAL APPENDAGES.—The epidermis of leaves, stems, fruits, and seeds of many plants frequently gives rise to out-growths in the form of *papillæ*, *hairs* and *scales*.

EPIDERMAL PAPILLÆ are short protuberances of epidermal cells. They may be seen to advantage on the upper epidermis of the ligulate corolla of various species of *Chrysanthemum*, on the lower epidermis of the foliage leaves of species of *Erythroxylon* and upon the upper epidermis of the petals of the Pansy (*Viola tricolor*). **EPIDERMAL HAIRS** or **TRICHOMES** are more elongated outgrowths of one or more epidermal cells. They may be unicellular, as those of the seed of cotton, or multicellular, non-glandular (simple) or glandular. The *non-glandular* hairs may be of various shapes, viz.: clavate (club-shaped) as on *Rhus glabra* fruits; stellate (or star-shaped) as on *Deutzia* leaves; candelabra-shaped, as on Mullein leaves; filiform, as on Hyoscyamus, Belladonna and Digitalis leaves; hooked, as on the stems of the Scarlet Runner or Hops; barbed, as on the stems of *Loasa* species; or tufted, as found on the leaves of Horehound (*Marru-*

FIG. 66.—Trichomes from different sources. 1, Unicellular non-glandular trichomes as seen growing out of epidermal cells of *Senna*; 2, uniseriate nonglandular trichomes of *Digitalis*; 3, unicellular stellate trichomes from *Deutzia scabra*; 4, trichomes of *Digitalis*; 5, clavate non-glandular twisted trichomes from lower epidermis of *Eriodictyon*; 6, 2-glandular trichomes from scraping of epidermis of the fruits of *Rhus glabra*; 7, branched trichomes of *Hygoscymus muticus*, a substitute for Henbane; 8, multicellular trichome of *Marrubium*; 9, glandular trichomes from leaves of *Digitalis purpurea*; 10, *Humulus* (Lupulin) 9, glandular trichomes from leaves of *Digitalis purpurea*; 10, aggregate, non-glandular trichomes of *Kamala*; 11, lateral view (to left) and vertical view (to right) of glandular trichomes of *Kamala*; 12, vertical view (above) and profile view (below) of 8-celled glandular hair from *Mentha piperita*. All highly magnified.

bium vulgare). They may be simple as in Cotton, etc., or branched as in *Hyoscyamus muticus*.

Nonglandular hairs are said to be *uniseriate* when they consist of a single row of cells, as those of *Digitalis* or *Stramonium*, etc.,

FIG. 67.

FIG. 68.

FIG. 67.—1, Epidermis of oak leaf; 2, epidermis of Iris leaf, both viewed from the surface; 3, group of cells from petal of *Viola tricolor*, showing conical papillæ; 4, two epidermal cells in cross-section showing thickened outer wall differentiated into three layers, namely, an outer cuticle, cutinized layer (shaded), and an inner cellulose layer; 5 and 6, epidermal outgrowths in the form of scales and hairs. (1, 2, 6 after Stevens, 3 after Strasburger, 4 after Sachs, and 5 after de Bary.)

FIG. 68.—Different forms of epidermal outgrowths. 1, Hooked hair from *Phaseolus multiflorus*; 2, climbing hair from stem of *Humulus Lupulus*; 3, rod-like wax coating from the stem of *Saccharum officinarum*; 4, climbing hair of *Loasa hispida*; 5, stinging hair of *Urtica urens*. (Fig. 3. after de Bary; the remainder from Haberlandt.)

multiseriate, when they consist of two or more rows of cells, as those of the corolla of *Calendula*.

The *glandular hairs* comprise those whose terminal cell or cells are modified into a more or less globular gland for gummy, resinous or oily deposits. They are generally composed of a stalk and a head region although rarely the stalk may be absent. The stalk may be unicellular, bicellular, uniseriate (consisting of a single series of superimposed cells) or multiseriate (consisting

of two or more rows of superimposed cells). The head varies from a one- to many-celled structure. The drug *Lupulin* consists of the glandular hairs separated from the strobiles of the Hop (*Humulus lupulus*).

SCALES are flat outgrowths of the epidermis composed of one or several layers of cells. They occur attached to the stipes of *Aspidium*, *Osmunda* and other ferns, where they are called "chaff scales." They are also found on a number of higher plants.

FUNCTIONS OF HAIRS.—Plant hairs are adapted to many different purposes. They may absorb nourishment in the form of moisture and mineral matter in solution, *e.g.*, root hairs. Those which serve as a protection to the plant may be barbed and silicified, rendering them unfit for animal food, or, as in the nettle, charged with an irritating fluid, penetrating the skin when touched, injecting the poison into the wound. A dense covering of hairs also prevents the ravages of insects and the clogging of the stomata by an accumulation of dust. They fill an important office in the dispersion of seeds and fruits, as with their aid such seeds as those of the milkweed and *Apocynum* are readily scattered by the wind.

The reproductive organs of many Cryptogams are modified hairs, as the sporangia of Ferns.

ENDODERMIS

ENDODERMIS is the layer of cells, constituting the innermost layer of the cortex. It forms a cylindrical sheath around the stele in roots and stems of pteridophytes and spermatophytes. It also occurs around the vascular bundles of leaves. In Angiospermous stems it usually resembles the other parenchyma layers of cortex as to structural characteristics, save that it frequently contains more starch. In fern stems, roots of Monocotyledons and of Dicotyledons of primary growth, however, its cells are clearly distinguished from the other cells of the primary cortex by their vertically elongated form and cutinized or suberized (occasionally lignified) radial and end walls.

The most outstanding characteristics of typical endodermal cells are to be found in their walls. These may be thin or thick

but the radial and end walls are usually thickened with deposits of a waxy, waterproof material representing *cutin* or *suberin*. The thin-walled endodermal cells possess strips or bands of cutin which extend around the cells on the inner surface of the radial and end walls. These strips or bands are called *Casparian*

FIG. 69.—Cross section of the stele and a portion of the epidermis and cortex of the monocotyledonous root of the corn. The outgrowths of the epidermis are root hairs. (From “*A Textbook of General Botany*” by Smith, Overton et al., Macmillan Co., publishers.)

strips. In cross sections the Casparian strips often appear like dots or lenticular shaped areas in the radial walls and are frequently termed “*Casparian dots or spots*.” In the roots of Mexican Sarsaparilla the inner as well as the radial walls are suberized; in those of the Honduras variety, inner, radial and outer walls all show suberization. Endodermal tissue is devoid of intercellular-air-spaces. Its cells contain protoplasm and

nucleus; starch grains are frequently but not always present and sometimes mucilage, tannin and crystals occur. Its functions seem to be to give protection to the *stele* (tissues within it), to reduce permeability between primary cortex and stele

FIG. 70.—Transverse section of the outer regions of the stem of the elder (*Sambucus canadensis*), showing cork, cork cambium, etc. (From Strasburger, Macmillan Co.)

and to serve as an air-dam, preventing the clogging with air of the water-conducting elements.

CORK

CORK or suberous tissue is a protective tissue composed of cells of tabular shape, whose walls possess a layer of water-proof substance called *suberin*. It is formed by the division of the cells of the *phellogen* or cork cambium which cuts off cells outwardly. Its cells, when first formed, contain cytoplasm and nucleus but, as the cork cambium forms new layers of cork on its exterior, the older cork cells lose their protoplasmic contents, die, become filled with air or a yellowish or brownish substance, and gradually sluff off. Under the microscope, dead cork cells usually appear black due to air filling their cavities. Thin sections of cork, when mounted in suitable reagents, will show

that the walls of the cork cells are composed of three layers or lamellæ. The lamella next to the cell cavity is cellulose. Outside of this is a middle lamella which may be lignified and a lamella of suberin. Cork tissue is usually devoid of inter-cellular-air-spaces. It forms a protective covering to the roots of secondary growth, perennial stems (after the first season)

FIG. 71.—Cross section of the herbaceous dicotyl stem of *Aconitum Napellus* var. *roseum*. $\times 50$. *ep*, epidermis; *co*, primary cortex; *per*, pericycle filled with pericyclic fibers; *ph*, phloem; *ca*, cambium (intrafascicular); *x*, xylem; *mr*, medullary ray; *m*, pith, composed of ordinary parenchyma.

of Dicotyledons and Gymnosperms, and wounds of stems and branches. It protects the delicate underlying tissues from invasion by fungal parasites, from sudden changes in temperature, and from loss of water.

The suberized walls of cork cells resist the action of concentrated sulfuric acid. They are colored green, when in contact with alcoholic extract of chlorophyll for several days in the dark.

RHYTIDOME sometimes called "scale bark" or "shell bark" represents the outer crust of alternating cork and cortical or pericyclic and phloem tissues resulting when cork cambia arise

in deeper regions of the bark and form cork or periderm. (See Fig. 131.)

BORKE is a term which has been applied by German botanists to designate the dead tissues which sluff off when cork cambium arises and forms layers of cork in deep regions of the cortex, pericycle and phloem. This occurs in the oak, birch, grape vine, elm, viburnum, hickory, cherries, etc.

PERICYCLE

The PERICYCLE or PERICAMBIUM, as it was formerly called, is a derivative of the *ground meristem*. It is a narrow cylinder of tissue which lies between the endodermis and the outer margin of the primary phloem of the fibrovascular bundles of stems or between the endodermis and the radial fibrovascular bundle of roots. In many stems it is composed entirely of thin-walled parenchyma cells but in others, whether woody or herbaceous and notably climbing or twining stems, it is composed of two kinds of tissues, thin-walled parenchyma and sclerenchyma. Frequently stone cells as well as sclerenchyma fibers occur in this region, as in stems of Cinnamon, etc. The sclerenchyma fibers occurring in the pericycle are called "*pericyclic fibers*." Normally these fibers are lignified but may consist of cellulose, as in Flax and Mezereon stems. The parenchyma cells sometimes contain chloroplasts and function in the slow conduction of sap and the storage of starch. The sclerenchyma fibers give strength. In roots, the pericycle consists of one or two rows of thin-walled cells which undergo division in places to form meristems which give rise to lateral roots. In roots of primary growth of Exogens the pericycle divides into two layers, the outer layer becoming a *phellogen* or cork cambium. The pericycle also may give rise to meristems forming adventitious roots and stems.

LATICIFEROUS TISSUE

This form of tissue comprises either *latex cells*, *laticiferous vessels*, or *coenocytic latex cells*, differing from each other in origin and method of development. (1) LATEX CELLS (*latex sacs*) are uninucleate cells which take their origin from minute

meristematic cells of the embryo. They are arranged in longitudinal rows and often occur in all of the organs. Each latex cell is a single unit and does not anastomose with other latex

FIG. 72.—Laticiferous vessels from the cortex of root of *Scorozonora hispanica*. A, As seen under low power, and B, a smaller portion under high power. (Stevens, after Sachs.)

cells. Such cells are abundant in the following families: *Apocynaceæ*, *Papaveraceæ*, *Urticaceæ*, *Sapotaceæ* and *Convolvulaceæ*.

(2) LATICIFEROUS VESSELS are long branching tubes, which owe their origin to chains of superimposed cells whose transverse walls have early become absorbed, the lumina of the cells then becoming filled with latex. They are found in various parts of

roots, stems, and leaves. The branches connect with those of other tubes forming an anastomosing network. These vessels occur in the following families: *Compositæ*, *Papaveraceæ*, *Campanulaceæ*, *Convolvulaceæ*, *Euphorbiaceæ*, *Araceæ*, *Oleaceæ*, *Geraniaceæ*, and *Musaceæ*.

(3) CÆNOCYTIC LATEX CELLS are elongated, multinucleate, laticiferous cells which occur in the milkweed family (*Asclepiadaceæ*) and in *Euphorbia* species. They arise, in each instance, from a single cell in the embryo which elongates at an equal rate with the growth of the plant, extending itself between other cells like a filamentous hypha of a mold, branching freely, but unlike laticiferous vessels the branches do not anastomose.

All of the latex elements are living, possessing thin layers of protoplasm just within their walls and containing one or more nuclei depending upon the type, also at times, *leucoplastids*, *elaioplastids* and *proteinoplastids*. The leucoplastids form starch grains, the elaioplastids build up oil globules and the proteinoplastids organize protein granules.

The exact role of latex elements remains undiscovered. Some botanists believe them to be food reservoirs, others excretory reservoirs, while still others regard them as representing a conducting system for the movement of starch and proteins.

All laticiferous elements contain a colorless, milky-white, or otherwise colored emulsion of gum-resins, fat, wax, caoutchouc and, in some cases, alkaloids, tannins, salts, ferments, etc. This emulsion is called "*latex*." For further details see under "Latex" in Chapter V.

SIEVE (LEPTOME OR CRIBIFORM) TISSUE

This tissue, found in the phloem region of fibro-vascular bundles, is a product first of the procambium and later of the cambium. It consists of superimposed, elongated, tubular cells whose longitudinal walls are thin and composed of cellulose and whose oblique or transverse walls, called "*sieve plates*," are thickened and perforated, permitting of the passage of proteids, amino-acids and amides from one cell to another. Frequently sieve plates are formed on the longitudinal walls, as in *Cascara Sagrada*. The mature sieve tube has a thin layer of cytoplasm

lining its walls and a large central vacuole but no nucleus. The tubes are arranged in longitudinal rows and the protoplasts of each series are connected by means of protoplasmic strands which extend from one sieve tube to another through perforations in the sieve plates. The cell sap found in the vacuole of sieve tubes contains protein substances often of slimy character. When fixed

FIG. 73.

FIG. 73.—Stages in the development of sieve tubes, companion cells, and phloem parenchyma. *A.* *a* and *b*, Two rows of plerome cells; in *c* and *d*, *a* has divided longitudinally and *c* is to become companion cells; *d*, a sieve tube, and *b*, phloem parenchyma. *B.* *c*, Companion cells, and *d*, a beginning sieve tube from *c* and *d*, respectively in *A*. The cross-walls in *d* are pitted; *b*, phloem parenchyma grown larger than in *A*. *C.* The same as *B* with the pits in the cross-walls of the sieve tubes become perforations, and the nuclei gone from the cells composing the tube. (From Stevens.)

FIG. 74.

FIG. 74.—Vascular elements. *A*, annular tracheal tube; *B*, spiral tracheal tube; *C*, reticulated tracheal tube; *D*, pitted tracheal tube; *E*, cross-section through plate of sieve tube, and adjoining companion cell; *F*, length-wise section of sieve tube; *G*, portions of two companion cells. (*A, B, C, D, Robbins; E, F, and G, after Strasburger.*)

and stained sections are examined, coagulated masses of protein will often be seen filling the pores of the sieve plates. These are called *slime plugs*.

Sieve tubes are usually accompanied by *companion cells* excepting in Pteridophytes and many Gymnosperms. Both

companion cells and sieve tubes arise by the division of the same mother-cell of the *procambium*. The *companion cells* occur in the phloem of Angiosperms only. In cross sections they appear as small, rounded or angular cells at the corner or lateral to a larger sieve tube. In longitudinal section they are seen to be elongated, slender cells. They may be distinguished from the mature sieve tubes by their abundant granular, cytoplasmic contents, their small vacuoles, and also by the fact that they retain their nuclei after complete maturation. Their walls are pitted where they are in contact with sieve tubes with which they are associated, phloem parenchyma and medullary rays. Their work is to take over food materials from the sieve tubes and pass them on to other tissues needing them for growth, repair or storage.

Besides sieve tubes and companion cells, phloem fibers also called bast fibers, and parenchyma cells called "*phloem-parenchyma*" or "*sieve parenchyma*" are often found in the phloem. The phloem-parenchyma cells transmit soluble carbohydrates, amino-acids and amides slowly downward and, through their communication with medullary rays, deliver food materials to them for radial distribution and storage.

TRACHEARY TISSUE

The tracheary tissue of plants comprises two kinds of elements, the *tracheæ* (ducts or vessels) and *tracheids*. They are formed first by the *procambium* or *plerome* and later by the *cambium*. Both of these conduct *crude sap* (water with mineral salts in solution) upward through the roots and stems into the leaves. The *tracheæ* are very long tubes of a cylindrical or prismatic shape which are formed by the breaking down of the transverse walls between vertical groups of superimposed cells, during the growth of the plant. Thousands of these tracheæ are arranged one above the other from the smallest branches of the root to the highest part of the stem and out into the farthest reaches of the leaves. Cross or end walls which are perforated with openings separate each trachea from its neighbors. In this way each line of tracheæ serves as a continuous tube for the conduction of water rapidly from root to leaf. Most tracheæ have two perforations, one in either end but as many as four have been

found, each opening into a different cell. The cross or end wall separating one trachea from another usually possesses a single, large, rounded or elliptical perforation called a *simple perforation*, but some end walls possess two or more smaller perforations forming an opening divided by transverse bars and called a *scalariform perforation*. The longitudinal or vertical walls of these tubes are of varying thickness, usually, however, thinner than those of woody fibers. The thickness is due to an infiltration of lignin upon the original cellulose wall. The walls show characteristic thickenings on their inner surfaces. These thickenings are for the purpose of giving strength to the tube while the thin parts of the wall are for the passage into and out of the tube of water with nutrient materials in solution.

TRACHEÆ are classified according to their markings as follows:

Annular, with ring-like thickenings.

Spiral, with spiral thickenings.

Reticulate, with reticulate thickenings.

Porous or *pitted*, with spherical or oblique slit pores.

Bordered pored, with rows of pores or slits, each being surrounded by a thickening of variable shape (circular, ellipsoidal and prismatic are the usual shapes).

Annulo-spiral, with both ring and spiral thickenings.

Scalariform, with ladder-like thickenings.

TRACHEIDS are primitive conducting xylem cells which have thickenings similar to tracheæ but are usually, though not always, smaller in size. Their ends are usually tapered, sometimes chisel-like, but not sharply pointed. They may be distinguished from tracheæ by the absence of perforations in their end walls. Their end walls have, however, thin places or pits which permit the more rapid passage of sap from cell to cell. Like tracheæ their walls usually give the characteristic lignin reaction with phloroglucin and HCl. The tracheids of conifers (see Fig. 75E, D.) such as the pine, spruce, cedar, etc., usually show one row of good-sized *bordered pores* (bordered pits) on their longitudinal walls. Each bordered pore exhibits a wall surrounding the pore which forms a dome-shaped protrusion into the cell. Like tracheæ, also, tracheids convey water with mineral salts in solution upward through the plant axis. Tracheids with bordered

FIG. 75.—Types of tracheae and tracheids. *A*, annulo-spiral trachea; *B*, spiral trachea; *C*, tracheid with several rows of bordered pores; *H*, pitted vessel; *E*, portion of a coniferous tracheid with a single row of bordered pits; *D*, tangential-longitudinal section of a bordered pit and portion of adjacent wall shown in *E*, enlarged; *p*, pore or orifice pit; *b*, protruding border; *cm*, limiting membrane which bears at its center a thickened pod or torus (*t*); *c*, cavity of pit; *F*, reticulate trachea in longitudinal section; *G*, longitudinal section of spiral trachea.

FIG. 76.—Stages in the development of the elements of the xylem. *A*, progressive steps in the development of a tracheal tube. 1, Row of plerome or cambial cells that are to take part in the formation of a tube; 2, the same at a later stage enlarged in all dimensions; 3, the cells in 2 have grown larger, their cross-walls have been dissolved out, and the wall has become thickened and pitted; 4, the walls in 3 have become more thickened, the pits have an overhanging border, the walls have become lignified as indicated by the stippling, and finally the protoplasts have disappeared, and the tube is mature and dead. *B*, stages in the formation of tracheids from plerome or cambial cells. The steps are the same as in *A*, excepting that the cross-walls remain and become pitted. *C*, steps in the development of wood fibers from cambial cells. 1, Cambial cells; 2, the same growth larger in all dimensions with cells shoving past each other as they elongate; 3, a later stage

pits and medullary-rays make up the wood of Conifers and most of the other orders of Gymnosperms.

The functions of tracheids are longitudinal conduction of crude sap and support to the part containing them.

MEDULLARY RAYS

These are bands of parenchyma cells which extend radially from the cortex to the pith (*primary medullary rays*) or from a part of the xylem to a part of the phloem (*secondary medullary rays* or “vascular rays”). Secondary medullary rays occur only as a result of the activity of the cambium during secondary growth of the plant axis. The xylem portion of a medullary ray is termed a *xylem ray* or *wood ray* while the phloem portion is termed a *phloem ray*.

In tangential-longitudinal section medullary rays usually appear spindle-shaped, (Fig. 63), while in radial-longitudinal sections they are seen crossing the other elements. Their primary function is to supply

FIG. 77.—Portion of radial-longitudinal section through wood of White Pine stem. *t*, tracheid; *p*, bordered pore of same; *m*, medullary ray. Magnified.

the cambium and wood with elaborated sap formed in the leaves and conveyed away by the sieve tubes and phloem parenchyma and to supply the cambium and phloem with crude sap which passes up mainly through the tracheæ and tracheids from the absorptive regions of the roots. They distribute food materials radially. They furthermore

with cells longer and more pointed and walls becoming thickened and pitted; 4, complete wood fibers with walls more thickened than in the previous stage and lignified, as shown by the stippling. The protoplasts in this last stage have disappeared and the fibers are dead. *D*, steps in the formation of wood parenchyma from cambial or procambial cells. 1, Group of cambial or procambial cells; 2, the same enlarged in all dimensions; 3, the same with walls thickened and pitted; 4 and 5 show the same stages as 2 and 3, but here the cells have enlarged radially or tangentially more than they have vertically. The walls of these cells are apt to become lignified, but the cells are longer lived than the wood fibers. (From Stevens.)

serve as storage places for starch, alkaloids, resins, and other substances. (Cf. Figs. 62, 127, 139.)

VASCULAR AND FIBROVASCULAR BUNDLES

VASCULAR BUNDLES are strands of conducting elements consisting of elongated cells specialized for the conduction of sap. The conducting strands are of two kinds: the PHLOEM which carries elaborated food materials (*elaborated sap*) from the leaves and green parts of stems to all parts of the plant, and the XYLEM which carries soil water with dissolved mineral salts (*crude sap*) from the roots to the leaves.

FIG. 78.—Diagram showing the transport of food through the sieve tubes, medullary-rays and tracheal tubes (tracheæ), and its storage in the parenchyma cells of the wood and bark. The black bodies in the cells indicate stored food. (Stevens.)

FIBROVASCULAR BUNDLES are groups of fibers, vessels and cells coursing through the various organs of a plant and serving for conduction and support. The term "Vascular Bundle" is employed to indicate bundles whose xylem and phloem contain no fibers and which are devoid of enveloping fibrous sheaths. According to the relative structural arrangement of their xylem and phloem masses, they may be classed as follows:

I. *Collateral*, consisting of a xylem and a phloem strand extending side by side.

II. *Closed collateral*, consisting of a strand of xylem lying alongside of a strand of phloem, in intimate contact, with no cambium zone between them. Stems and leaves of most Monocotyledons and Horsetails.

III. *Open collateral*, consisting of a strand of xylem and a strand of phloem separated from each other by a cambium.

Stems and leaves of Dicotyledons and roots of Dicotyls and Gymnosperms of secondary growth. In stems and roots the phloem faces toward the exterior and the xylem toward the center of the axis.

IV. *Bicollateral*, characterized by a xylem strand being between an inner and an outer phloem mass. Cambium may

FIG. 79.—Closed collateral bundle of stem of *Zea mays*. VG, Bundle sheath; L, intercellular space; A, ring from an annular tracheal tube; SP, spiral tracheal tube; M, pitted vessels; V, sieve tubes; S, companion cells; CP, crushed primary sieve tubes; F, thin-walled parenchyma of the ground or fundamental tissue. (From Sayre after Strasburger.)

be present or absent on either outer or inner face of the xylem or on both faces. Seen chiefly in stems and leaves of the *Cucurbitaceæ*, *Combretaceæ*, *Loganiaceæ*, *Apocynaceæ*, *Asclepiadaceæ*, *Oleaceæ*, *Convolvulaceæ*, *Gentianaceæ*, *Myrtaceæ* and *Solanaceæ*.

V. *Concentric*, characterized by a central xylem mass surrounded by a phloem mass or *vice versa*. No cambium present.

(a) *Concentric*, with xylem central in bundle. (*Xylocentric bundle*.) Seen in stems and leaves of nearly all ferns and the *Lycopodiaceæ*.

(b) *Concentric*, with phloem central in bundle. (*Phlocentric bundle*.) Seen in stems and leaves of some *Monocotyledons*. Examples: *Calamus* and *Convallaria* rhizomes.

VI. *Radial*, characterized by two or more xylem and phloem masses arranged in alternate manner usually around a central pith. The terms diarch, triarch, tetrarch, pentarch, hexarch, heptarch and polyarch are respectively employed to designate radial bundles having 2, 3, 4, 5, 6, 7 or many xylem strands and

FIG. 80.—Vascular bundles. A, the concentric-monocotyl or phlocentric type, with xylem, k, surrounding the phloem, h. B, the collateral type, with phloem, h, standing in front of the xylem, k. C, a portion of the radial type (undergoing transition) shown complete in D, where the part outlined at a corresponds to C. c, xylem, b, phloem, f, cambium ring; e, pericycle; d, endodermis. Pericycle and endodermis are not parts of the bundle. (From Stevens after Haberlandt.)

a like number of phloem strands. Seen in the primary roots of all *Spermatophytes* and *Pteridophytes*.

XYLEM¹ is that part of a conducting or vascular bundle that usually contains wood parenchyma cells, tracheæ and tracheids. It frequently contains, in addition, wood fibers. The tracheid is the fundamental cell type in the xylem. The xylem of most *Gymnosperms* is devoid of tracheæ.

¹ The conducting elements of the xylem comprising tracheæ, tracheids and wood parenchyma constitute the *hadrome*.

PRIMARY XYLEM (PROTOXYLEM) is that xylem in all conducting bundles which directly develops from the plerome. In exogenous stems of secondary growth, the primary xylem is in small strands adjacent to the pith, and in roots of secondary growth it appears as a small, several rayed strand in the center.

FIG. 81.—Bicollateral bundle of the stem of Squash (*Cucurbita*). $\times 70$. P' , external phloem; P^2 , internal or intraxylary phloem; x , xylem; ca , cambium; si , a sieve tube.

SECONDARY XYLEM (METAXYLEM) is xylem layed down by the cambium.

PHLOEM¹ is that part of a vascular bundle that contains sieve tubes, phloem parenchyma cells, and often bast fibers. The sieve tubes are the most important structural and functional elements of the phloem. In Pteridophytes and many Gymnosperms these and phloem parenchyma are alone present in the

¹ The conducting elements of the phloem which comprise the sieve tubes, companion cells and phloem parenchyma constitute the *leptome*.

phloem while in Angiosperms sieve tubes, companion cells, phloem parenchyma, bast fibers, secretion cells and stone cells may occur in this region.

PRIMARY PHLOEM (PROTOPHLOEM) is the phloem formed by the plerome and occurs directly beneath the pericycle.

FIG. 82.—A Polyarch Radial Bundle. Cross-section through a portion of a root of *Acorus calamus*, a monocotyledon. A, Cortical parenchyma; B, endodermis; C, pericycle; E, phloem; F, xylem. At Y, Y, are large tracheal tubes, which were formed last, the narrow tubes near the periphery of the xylem being formed first. At the center of the root, within the circle of the polyarch, radial vascular bundle, occur thin-walled, parenchymatous pith cells. (From Sayre after Frank.)

SECONDARY PHLOEM (METAPHLOEM) is phloem formed by the cambium.

LEAF AND BRANCH TRACES.—These are prolongations of the vascular bundles from the stele of the stem which extend into leaves and branches. (See Fig. 84.)

THE STELE

The STELE represents the central cylinder of a stem or root and includes all of the tissues interior to the cortex.

A PROSTELE is a primitive form of stele in which the vascular tissues form a solid cylinder, the phloem surrounding the xylem. It occurs in young parts of stems of club-mosses, in the stems of some ferns and in the roots of secondary growth of most plants. (See Fig. 83A.)

A SIPHONOSTELE resembles the prostele except that a pith occurs in its centre. There are two kinds of siphonosteles, viz., the *ectophloic* in which the phloem is external to the xylem, as

FIG. 83.—Diagrams showing the types of arrangement of vascular tissues in steles. A, protosteles. B, siphonostele. C, dictyostele. (From "An Introduction to Plant Anatomy" by Eames and Mac Daniels, McGraw-Hill Book Co., Inc. Publishers.)

in the stems of gymnosperms and angiosperms, and (2) the *amphiphloic* in which there is an external phloem outside the xylem and an internal phloem on the inner face of the xylem, as in the stems of some ferns and certain families of dicotyledons. (See Fig. 83B.)

A DICTYOSTELE is a siphonostele whose vascular mass is broken up into a number of longitudinal strands or vascular bundles. This type is common to most monocotyl stems, the bundles being sometimes called *meristeles*. (See Fig. 83C.)

BRANCH TRACES.—These are strands of vascular tissue which arise from the vascular bundles of the stele of the stem and run out into the lateral branches. Usually two traces or bundles arise for each branch, but sometimes only one arises. When

FIG. 84. Diagrams illustrating leaf traces, branch traces and gaps. A, longitudinal section of node through leaf trace and gap. B, similar to A, but with branch trace and gap also present. C, view of vascular cylinder showing departure of leaf and branch traces, and the gaps associated with each. D, E, F, cross sections of stem illustrated in A at levels *a-a*, *b-b*, and *c-c*, respectively. G, face view of outside of cylinder shown in C, the leaf and branch traces cut away at the surface of the cylinder. H, transverse section of G at *a-a*. (In diagrams A-H the vascular tissue is not differentiated as xylem and phloem; the traces are doubly cross-hatched. In I, more detailed structure is shown, protoxylem, metaxylem (secondary xylem) and pith being indicated. (From "An Introduction to Plant Anatomy" by Eames and Mac Daniels, McGraw-Hill Book Co., Inc., publishers.)

the branch supply consists of two bundles, these soon unite on their way into the branch to form a cylindrical stele which appears circular as viewed in cross section. In instances where only one trace or bundle strand courses into the branch, this appears in cross section as a horseshoe or crescentic structure with an opening. The opening is closed as the trace passes into the branch so that in the branch a complete cylindrical stele is formed.

LEAF AND BRANCH GAPS.—These are breaks in the continuity of the vascular cylinder of stems around and above the places where the leaf or branch traces leave the cylinder. The break or opening is designated as a *leaf gap* or *branch gap* according as to whether a leaf trace or branch trace is involved in its formation. It contains parenchyma and through it the cortex and pith are confluent. (See Fig. 84.)

SECRETION SACS (SECRETION CELLS)

These were formerly parenchyma cells which sooner or later lost their protoplasm and nucleus and became receptacles for oil, resin, oleoresin, mucilage or some other secretory substance. They are generally found in parenchyma regions of stems, roots, leaves, flower or fruit parts and frequently possess suberized walls. Good illustrations of these structures may be seen in Ginger and Calamus.

INTERCELLULAR AIR SPACES

Intercellular air spaces are cavities filled with air found between cells or groups of cells throughout the bodies of higher plants. Their function is to permit of the rapid movement of atmospheric gases through the entire plant body. They are formed either by the breaking down of the middle lamella of the cell walls, where several cells come together, and a later separation of the cells at these places (*Schizogenous intercellular-air-spaces*), or by a breaking down and disappearance of cell walls common to groups of cells (*Lysigenous intercellular-air-spaces*). In terrestrial plants which live in middle regions (*mesophytes*) and in desert plants (*xerophytes*) the intercellular-air-spaces are averagely small and more or less angular. In plants of swamp

or marsh habit they are medium-sized, while in those which live entirely in the water (*hydrophytes*) they are of large size and more or less rounded.

INTERNAL SECRETION GLANDS AND RESERVOIRS

These structures are either found as globular cavities as in Orange and Lemon Peel and Eucalyptus leaves, containing oil

FIG. 85.—Cross section through a portion of Orange Peel showing the cavity of an internal, schizogenous globular gland at *g*; crystals of hesperidin at *h*; membrane crystals of calcium oxalate at *k*. (After Tschisch and Oesterle.)

FIG. 86.—Lysigenous gland in the leaf of *Dictamnus fraxinella*, *B*, young glands, with cells beginning to secrete oil; *C*, mature gland where the secreting cells have broken down and left their secretion in the cavity thus formed; *o*, large drop of secreted oil. (Stevens after Sachs.)

or oil and resin, when they are called *internal glands*, or, as tube-like spaces filled with hydrocarbon principles, such as are found in Pine leaves and stems, when they sometimes receive the name of *secretion canals* or *reservoirs*. In the Parsley family these canals are found in the fruits, where they are called *vittæ*. Occasionally they are named according to the nature of their contents—resin or oil canal, reservoir, etc. The *internal glands* usually arise by the division of a cell or group of cells of the ground

meristem but sometimes the dermatogen gives rise to them, as in Clove, Myrtle, Asarum, etc., or both the ground meristem and dermatogen may contribute to the formation of the same gland. Internal glands and secretion reservoirs are termed *schizogenous* when the secreting cells split apart at the center of the group and draw away from the line of separation, as in the

FIG. 87.—Schizogenous secretion reservoir (resin duct) as seen in cross section in the young stem of English Ivy. A, early and E, later stage in the formation of the duct. g, the mature duct; c, cambium; wh, phloem; b, bast fibers, rp, parenchyma. (Stevens, after Sachs.)

stems and leaves of the pines; when they break down completely, leaving their secretion in the resulting cavity they are termed *lysigenous*, as in the rinds of the orange, lemon and other Citrus fruits. The *secretion reservoirs*, also called *internal tubular glands*, are formed by a long vertical series of cells forming a space schizogenously or lysigenously. The schizogenous type of gland and reservoir is lined with a layer of cells, usually more or less flattened, which are characterized by possessing large nuclei. To this layer has been assigned the name "*secretory epithelium*."

NECTARIES

Nectaries are nectar-secreting tissues, consisting either of groups of modified cells or of specialized glands, tubes or cavities found on insect-pollinated plants. The nectar secreting tissue occurs usually on parts of the flower but may occur on petioles, stipules, bracts or other structures. *Nectar* is a sugary secretion which is exuded through the outer walls of the cells of the

FIG. 88.—Resin duct (secretion reservoir) in leaf of *Pinus sylvestris*, in cross-section at *A*, and in longitudinal section at *B*; *h*, cavity surrounded by the secreting cells; *f, f*, sclerenchyma fibers surrounding and protecting the duct. (*Stevens after Haberlandt.*)

nectary. When the nectar secreting tissue is superficial the modified epidermal cells are often palisade-like or papillose in form, but may consist of dense groups of club-shaped hairs. In numerous plants as more elaborate structures such as sunken pouches, as in the buttercups, or elevated glandular structures lined with nectar secreting structures, as on the petioles of *Viburnum Opulus*, etc.

CLASSIFICATION OF TISSUES ACCORDING TO FUNCTION.—According to their particular function, tissues may be classified as follows:

I. CONDUCTING TISSUES	{	Parenchyma (fundamental tissue)
		Medullary rays
		Xylem cells (wood parenchyma)
		Tracheæ (vessels) and tracheids
		Phloem cells (phloem parenchyma)
		Sieve tubes
		Companion cells
II. PROTECTIVE TISSUES	{	Epidermis (outer cell walls cutinized)
		Cork (suberized tissue)
		Phloem fibers
		Wood fibers
III. MECHANICAL TISSUES	{	Sclerenchyma fibers
		Pericyclic fibers
		Stone cells
		Collenchyma
		Internal glands
IV. SECRETING TISSUES	{	Secretion reservoirs
		External glands (glandular hairs)
		Nectaries
		Parenchyma
		Reserve parenchyma
V. STORAGE TISSUES	{	Medullary-rays
		Wood parenchyma
		Phloem parenchyma
		Laticiferous tissue
		Primordial meristem
		Dermatogen (protoderm)
VI. MERISTEMATIC TISSUES	{	Periblem (ground meristem)
		Plerome (procambium)
		Cork cambium
		Cambium

CHAPTER VII

THE ROOT

The **ROOT** is that part of the plant that grows into or toward the soil, that never develops leaves, rather rarely produces buds, and whose growing apex is covered by a cap.

The chief functions of a root are absorption, storage and support. Its principal function is the absorption of nutriment and to this end it generally has branches called **ROOTLETS** that are covered with **ROOT-HAIRS** which largely increase the absorb-

FIG. 89.—Cross-section of rootlet in the region of the root-hairs. (*From Stevens.*)

ing surface. These **ROOT-HAIRS** are of minute and simple structure, being merely elongations of the epidermis of the root, for a short distance back of the root cap, into slender tubes with thin walls. Like other living cells, each shows cytoplasm, sap vacuoles containing cell sap, a nucleus with nuclear membrane, outer plasma membrane against the covering cell wall of cellulose and vacuolar membranes around sap vacuoles. That portion of the epidermis which bears root-hairs is called the **PILIFEROUS LAYER**.

SOIL AND WATER RELATION.—Under conditions favorable to the growth of land plants, the soil is open and porous, the particles of soil being separated by spaces containing air, while every soil particle is enveloped by a film of water. The root-hairs are in close contact with these soil particles (Fig. 92) and

soil water from the films surrounding them soaks through their cell walls into the root-hairs. This soil water is a solution of various mineral salts. In order to understand how this solution passes into the root-hairs, it is essential to first understand the processes of diffusion and osmosis.

DIFFUSION.—This is the process whereby the particles of substances making up a solution tend to be uniformly intermingled. It may be observed by introducing a crystal of copper sulfate into a tumbler of water. The crystal slowly dissolves and the particles of which it is composed in time diffuse themselves equally throughout the water, so that the water is colored uniformly blue. The movement of the particles of the dissolved copper sulfate from the region of greater density or concentration to that of less implies a pressure which is termed diffusion tension.

OSMOSIS.—If the denser copper sulfate solution in the bottom of the tumbler were separated from the less dense water above by a semipermeable membrane such as parchment or bladder, diffusion would take place through the membrane and the water above would become colored as proof of this diffusion. Moreover, the water above the membrane would pass through it in the opposite direction and more rapidly than the solution of the copper salt. It would continue to do so until the solution was of the same density on both sides of the membrane. *Diffusion through a porous membrane is known as osmosis.* It is a well-known law of physics that when two liquids or gases of different densities are separated by a porous (osmotic) membrane, diffusion through the membrane will take place until the density of the fluids or gases becomes the same on each side. The diffusion will be more rapid from the less dense to the more dense region.

FIG. 90.—Osmosis. The more rapid inflow of liquid from A to B has elevated the liquid surface within from b to a. (From Brownell's General Science.)

OSMOSIS APPLIED TO ROOT-HAIRS.—Normally the cell sap of the root-hairs is a denser liquid than the soil water outside of

FIG. 91.—Successive stages of plasmolysis. *N*, nucleus; *V*, vacuole; *p*, protoplasm; *e*, area formerly occupied by protoplast now replaced by plasmolyzing solution. (*Palladin after de Vries.*)

the hairs. The outer plasma membrane and the vacuolar membranes represent porous osmotic membranes separating the

FIG. 92.—Root-hairs, with soil-particles adhering. (*Gager, after Sachs.*)

denser solution within the hair from the less dense soil water without. The soil water imbibed by the cell wall passes by osmosis through the outer plasma membrane, diffuses through the cytoplasm to the vacuolar membranes through which it passes by osmosis into the sap vacuoles and there becomes a portion of the cell sap.

The kind of osmosis just indicated is known as *endosmosis* or osmosis from without in. A reverse process takes place when small traces of CO_2 , acid

and other substances are excreted, *e.g.*, passed out of the root-hair. This is called *exosmosis*.

TURGOR.—If the cell sap of the root-hair or any other plant cell becomes concentrated above that of the soil solution, the cell is caused to bulge. This bulging is a manifestation of osmotic pressure within and the condition resulting is termed *turgor*. Cells of plants exhibiting this phenomenon are said to be in the state of turgescence or rigidity. Since turgor is coexistent with growth, conditions affecting turgescence affect at the same time all growth processes.

PLASMOLYSIS.—If root-hairs or other plant cells are placed in solutions of a greater density than that of their cell sap content, as for instance 5 to 10 per cent. salt or sugar solution, exosmosis becomes more rapid than endosmosis. The cell sap is extracted more rapidly than the fluid enters from without, with a result that the protoplasm is loosened from the cell wall and caused to collapse. This condition is called *plasmolysis* and the cell is said to be *plasmolyzed*. Plasmolysis throughout a tissue or organ results in wilting. Many plants wilt on account of too high a concentration of soil solutes.

ABSORPTION OF NUTRIENT SALTS.—The plasma membranes of all living plant cells are permeable to certain solutes (substances in solution) and impermeable to others. These membranes in the root-hairs absorb each solute particle separately and according to the need or attraction for that substance; this is called *selective absorption*.

The higher green plants vary greatly in respect to their mineral requirements. Of thirty-one elements found in the ash of plants, only eleven occur regularly, *e.g.*—sulfur, chlorine, phosphorus, silicon, potassium, sodium, calcium, magnesium, iron, aluminum (a trace) and manganese (a trace). Ten elements are absolutely required in the form of water-soluble salts for the normal growth of higher green plants. These ten are termed *essential elements* and comprise the following: *nitrogen, sulfur, phosphorus, calcium, potassium, magnesium, iron, manganese, boron* and *zinc*. Traces of copper have also been found essential for the growth of some plants.

IMPORTANCE OF ESSENTIAL ELEMENTS.—*Nitrogen, sulfur* and *phosphorus* are constituents of proteids and so of protoplasm and are essential to their formation. Without sufficient nitrogen,

growth is retarded. Without sufficient sulfur, cell-division and fruit formation are checked. Deficiency in phosphorus retards the maturation of the plant.

FIG. 93.—Median longitudinal section through the root tip of the barley. *k*, calyptrogen; *d*, dermatogen, *c*, the thick outer wall of the same; *pr*, periblem; *pl*, plerome; *en*, endodermis; *i*, intercellular-air-spaces; *a*, cell rows which will develop the central vessels; *r*, detached cells of root cap; *s*, starch grains in cells of root cap. $\times 180$. (After Strasburger and Koernicke in *Lehrbuch der Botanik*.)

Calcium not only neutralizes many harmful substances but facilitates the absorption of other salts and is necessary for normal leaf development. It is the chief element in the ash of leaves.

Potassium accompanies carbohydrates during their translocation and is thought to play a part in their formation.

Magnesium is a constituent of chlorophyll and accompanies proteins. It is necessary for the formation of fats. Deficiency in this element results in chlorosis and poor fruit formation.

Iron is essential to the formation of chlorophyll although it is not a component of that substance. When deprived of iron, green plants become pale and chlorotic, even though they are grown in light.

While not absorbed as mineral salts, *carbon*, *hydrogen* and *oxygen* are likewise indispensable elements. The very life of all plants depends upon their availability. All three enter into the formation of proteins, carbohydrates and protoplasm and the first two are essential to the formation of fats and oils. Water, which makes up the greatest part of protoplasm and forms the solvent for soil salts, consists of a combination of two parts of hydrogen to one part of oxygen.

In determining which elements are essential for the normal growth and health of plants, it is customary to grow them in water solutions of various salts or in sterile sand or ground pumice containing a water solution of the salts.

NON-ESSENTIAL ELEMENTS ABSORBED BY ROOT-HAIRS.—These elements include those which are also absorbed as water-soluble salts but which are not required for the normal growth of all plants. In many specific instances, however, they may be beneficial.

Silicon is an abundant element in many plants, notably the grasses, sedges and scouring rushes. It occurs in the ash of these plants as silicon dioxide (SiO_2) and in most soils as silicic acid (H_2SiO_3). It is deposited mainly in cell walls of the peripheral parts of stems, leaves and seeds and affords protection from penetration by the hyphæ and parasitic fungi as well as from animal attack such as plant lice and scale insects.

Chlorine, another inessential element, probably has an influence on the translocation of carbohydrates from leaves to other organs under natural conditions. Most plants can attain complete development without it. The question of its rôle still remains unsettled.

Aluminum is only occasionally found in plants. It is known to influence the color of flowers in *Hydrangea hortensis*. When

grown in forest or moor soil, this species has reddish flowers but when grown in soil containing soluble aluminum compounds, its flowers are blue.

REGIONS OF THE ROOT.—There are five regions of a root beginning with the extreme tip and passing backward. These are the *root cap*, the *embryonic region*, the *region of elongation* or growth in length, the *region of maturation* or cell differentiation and the *mature region* (see Fig. 94). These phases of growth can be seen in fixed and stained longitudinal sections of root tips.

ROOT CAP.—The tip of each root or rootlet is protected by a smooth, sheath- or scale-like covering known as the *root cap* or *calyptra*. This is a mass of cells which not only protects the delicate cells of the embryonic region but serves as a mechanical aid as the root pushes its way through the soil.

EMBRYONIC REGION.—This region is composed of small, densely arranged, somewhat polyhedral or quadrangular cells which undergo repeated division. This region is about a millimeter in length.

GROWING POINT.—The growing point of the root of a seed plant consists of a group of actively dividing cells directly beneath the root cap. These cells make up the tissue called the *primordial meristem*.

PRIMARY MERISTEMS.—Just in back of the primordial meristem, the cells have begun to show differentiation into three groups of tissues called *primary meristems* or *generative tissues*. In the posterior part of this region the cells gradually cease their rapid division and become larger due to the absorption of soil water.

GENERATIVE TISSUES.—The generative tissues directly back of the growing point are, passing from the center toward the outside: *plerome* or *procambium*, producing fibro-vascular tissue (xylem and phloem) and pith; *periblem* or *ground meristem*, producing cortex including endodermis, and pericycle; and *dermatogen* or *protoderm*, producing epidermis. The name *calyptrogen* has been given to those cells of the primordial meristem which form the root cap. (Cf. Fig. 93.)

REGION OF ELONGATION.—This region occurs just behind the embryonic region where the cells enlarge chiefly in length. Root

FIG. 94.—Lengthwise section of a root tip (diagrammatic), showing its various regions. (From "A Textbook of General Botany" by Smith, Overton et al., Macmillan Co., Publishers.)

hairs begin to make their appearance in this region. Differences in size and shape of the cells also begin to occur here.

REGION OF MATURATION.—In this region more differentiation takes place, the cells become larger and frequently their walls become thicker. Root hairs are usually evident throughout this region.

MATURE REGION.—This region is the part of the root between the region of maturation and the base of the stem. In its lowest part the primary tissues have been laid down by the primary meristems. These, in order of their arrangement from the periphery to the center, are the *epidermis*, *primary cortex* (composed of parenchyma), *endodermis*, *pericycle*, *radial vascular bundle* composed of alternating strands of *primary xylem* and *primary phloem*, and *pith* (composed of parenchyma). *Root hairs* usually occur in this lower region as outgrowths of epidermal cells but gradually disappear in most roots as we pass further upward.

DIFFERENCES BETWEEN ROOT AND STEM

THE ROOT	THE STEM
1. Descending axis of plant.	1. Ascending axis of plant.
2. Growing point sub-apical, being covered by the root cap.	2. Growing point apical, but covered by protective bud scales or young foliage leaves.
3. Contains no chlorophyll.	3. Chlorophyll sometimes present.
4. Does not bear leaves or leaf rudiments.	4. Bears leaves and modifications.
5. Branches arranged irregularly, and arising from deeper tissue.	5. Branches with mathematical regularity, the branches arising at surface of stem, in axils of leaves and from nodes.
6. No division into nodes and internodes.	6. Division into nodes and internodes.
7. Internal structure different.	7. Internal structure different.

CLASSIFICATION OF ROOTS AS TO ORDER.—1. *Primary* or *first root*, a direct downward growth of the radicle (from embryo within seed), which, if greatly in excess of the lateral roots, is called the *main* or *tap root*.

2. *Secondary roots* are lateral branches of the primary root. They may, in turn, give rise to finer branches, or rootlets. Both primary and secondary roots may be either *fibrous* or *fleshy*.

3. *Adventitious roots* are such as arise from any part of the plant other than the seed embryo or normal root system. Examples: Roots developing on *Bryophyllum* and *Begonia* leaves when

placed in moist sand; roots arising from stems of climbing plants or from stem cuttings.

4. *Prop roots* or *brace roots* are such as grow out of the stem a short distance above the soil and extend diagonally into the ground, serving as supports to the stem. Examples: Indian Corn and Pandanus.

5. *Epiphytic roots* are the roots of epiphytes or air-plants, many of which are common to tropical forests. In some instances, as in *Vanilla* and other epiphytic Orchids, these roots, arising from aerial stems, hang down free in the air and absorb water from rain by means of their several layered epidermis called a *velamen*. In others, as in the English Ivy, several short roots grow out of the stem at various intervals and adhere by their tips to walls, thus serving as supports for climbing.

The Banyan (*Ficus religiosa*) of India can extend itself over large areas by means of its aerial roots. These extend down vertically from horizontal branches and form trunks, upon anchoring themselves in the soil.

FIG. 95.—*Vanilla planifolia*, an epiphyte, growing on a tree in Jamaica, W. I. (Gager, after Maxon, U. S. Dept. Agr.)

6. The roots of parasitic plants are known as *Haustoria*. These penetrate the bark of plants upon which they find lodgment, known as hosts, and absorb nutritious juices from them. The *Mistletoe*, *Dodder* and *Gerardia* are typical parasitic higher plants.

CLASSIFICATION OF ROOTS AS TO FORM.—*Fusiform*, or *spindle-shaped*, like that of the radish or parsnip.

Napiform, or *turnip-shaped*, somewhat globular and becoming abruptly slender then terminating in a conical tip, as the roots of the turnip and some radishes.

Conical, having the largest diameter at the base then tapering, as in the Maple and Aconite.

Fibrous, when very slender or fiber-like, as in the grasses and cereals.

Tortuous, when irregularly bent or twisted, as Pareira.

Nodose, when swollen at intervals in its course and fibrous otherwise, as the roots of the Dropwort.

Moniliform, when the swellings occur at very close intervals giving the root the appearance of a close chain of beads.

Tuberous, when fleshy, swollen, especially in the central portion, and resembling tubers in shape, as in Jalap and the Sweet Potato.

Fascicled, when a number of thickened and similarly looking roots occur in a cluster, as in the Dahlia.

ROOT CONSISTENCY OR TEXTURE.—In this particular, roots are termed *woody*, when the

FIG. 96.—Dodder (*Cuscuta* sp.) in flower. Parasitic on a golden rod (*Solidago ulmifolia*). (Gager, photo by Elsie M. Kittredge.)

hard, lignified tissue predominates and *fleshy*, when the soft cellular tissues predominate. Fleshy roots may be (1) simple or single or (2) multiple, when a number arise from the base of the stem in a cluster, as in the Dahlia.

DURATION OF ROOT.—Plants are classified according to the duration of the root, as follows:

1. *Annual plants* are *herbs* with roots containing no nourishment for future use. They complete their growth, producing

flowers, fruit and seed in a single season, then die. Examples: Stramonium, Lobelia, etc.

2. *Biennial* plants develop but one set of aerial organs the first year, *e.g.*, the leaves, and, as *Digitalis*, *Conium*, etc., a large amount of reserve food material is stored in the roots for the

FIG. 97.—Photomicrograph of a transverse section of the stem of a dicotyledonous host-plant infested with the parasite, Dodder. Note the haustoria extending from the Dodder (D, D') into the cortex of the host (H). Greatly enlarged. (Gager.)

support of the plant the following season, when it flowers, fruits and dies.

3. *Perennial plants* are those whose roots or underground stems and roots live indefinitely, as trees, shrubs and perennial herbs.

ROOT HISTOLOGY.—The histology of roots varies, depending upon character of the surrounding water relations or soil in which

the plants are accustomed to live which may be aquatic, mesophytic, halophytic or xerophytic, etc.

Aquatic plants or *hydrophytes* are those which live in water or wet soil as the water-lilies, calamus, cat-tails, etc.

FIG. 98.—Root systems. Fleshy tap root of Carrot (1); napiform root of a Radish (2); multiple, tuberous roots of a Dahlia (3); nodose roots of the Dropwort (4); fibrous roots of a Grass (5); fibrous-tuberous root system of an Orchid (6).

Mesophytic plants or *mesophytes* are those which thrive with a moderate supply of moisture, as most of the land plants of our fields and meadows.

Halophytic plants or *halophytes* are those which thrive in salt marshes, on saline flats near the sea coast and on the alkali flats of the interior, as the salt marsh samphire, the mangroves, chenopodium, etc.

Xerophytic plants or *xerophytes* are those which thrive in very dry soil, as the cactus, century plant, hemlock, spruce, etc.

HISTOLOGY OF MONOCOTYL ROOT

In this connection we will discuss only the type of greatest pharmacognic importance, *i.e.*, the mesophytic type as seen in its most typical form in the transverse section of a *Smilax* root.

Examining such a section from periphery toward the center, one notes the following:

1. *Epidermis* (derived from dermatogen) of a single layer of cells many of which give rise to root-hairs.

2. *Hypodermis* (derived from periblem) of one, two or three layers of cells whose walls are extremely thickened.

3. *Cortex* (derived from periblem) consisting of a broad zone of parenchyma cells many of which contain starch grains.

4. *Endodermis* (derived from periblem) of one layer of endodermal cells whose walls are extremely thickened through the infiltration of suberin and lignin.

5. *Pericycle*, also called pericambium (derived from periblem), of one or more layers of meristematic cells whose walls are extremely thin.

6. A *radial fibro-vascular bundle* (derived from plerome) of many alternating xylem and phloem strands and hence *polyarch*. The phloem tissue consists of phloem cells and sieve tubes arranged in small ovate areas beneath the pericycle and between the outer reaches of the xylem strands. The xylem is composed of xylem cells, tracheæ and wood fibers.

7. *Medulla* or *pith* (derived from plerome) composed of parenchyma cells containing starch and often showing xylem patches cut off and enclosed within it.

The tissues within the endodermis collectively constitute the STELE.

The *Smilax* root resembles most monocotyl roots in showing no secondary increase in thickness which is sometimes called secondary growth. *Secondary increase in thickness* is the rule in roots of most dicotyledons and gymnosperms and is due to the formation in those roots of a cambium. No such tissue arises in most monocotyl roots. However, some difference in diameter will often be noted between sections cut through the same root at different levels above the root cap, those cut nearer the stem end

FIG. 99.—Cross section through a representative portion of a monocotyl root of a species of *Smilax*. ($\times 70$.) *ep*, epidermis; *hy*, hypodermis with lignified radial and outer walls; *co*, cortex; *par*, parenchyma; *en*, endodermis whose radial and inner walls are thicker and more lignified than the outer walls and which show a V-shaped lumen; *pc*, pericycle; *ph*, strand of phloem, the end of the leader being on a sieve tube; *x*, xylem; *wf*, wood fibers in xylem; *tr*, trachea in xylem; *m*, pith. Note the numerous xylem and phloem strands collectively constituting a polyarch radial fibrovascular bundle.

being broader. The explanation is that the cells in the broader sections have become larger through the process of growth, rather than having been augmented in number.

HISTOLOGY OF DICOTYL ROOTS

The typical dicotyl root is a tetrarch one, which is a root whose radial bundle exhibits four xylem patches alternating with four phloem patches. These roots usually have an unlimited power of growth. The early growth is called *primary growth* which means first growth and is represented in sections cut a little above the root cap in the region of the root hairs. Above the region of primary growth is the *region of transition* where changes are taking place in the root which lead to the appearance of secondary structure.

A transverse section of a dicotyl root in its YOUNG GROWTH shows the following structure from periphery toward center:

1. *Epidermis* (from dermatogen) with cutinized outer walls, the cells often elongating to form root-hairs.

2. *Primary cortex* (from periblem) of parenchyma cells with usually small intercellular spaces. Whenever the outermost layer or layers of cells of this region have more thickened walls than those beneath, the term *hypodermis* is employed for this outer portion of the cortex.

3. *Endodermis* (from periblem) or innermost layer of cells of the cortex, with lenticularly thickened radial walls.

4. *Pericycle* (*Pericambium*) (from periblem) of one to two layers of actively growing cells which may produce side rootlets.

5. *Radial fibro-vascular bundle* (from plerome) of four, less frequently of two, three, five or six *primary phloem* (protophloem)

FIG. 100.—Cross-section of a young root of *Phaseolus multiflorus*. A, pr, cortex; m, pith; x, stele or central cylinder—all tissue within the pericycle, inclusive; g, primary xylem bundles; b, primary phloem bundles. B, cross-section of older portion of root; lettered as in A; b', secondary phloem; k, cork. (Stevens, after Vines.)

patches alternating with as many *primary xylem* (protoxylem) arms. It is not uncommon to find bast fibers along the outer face of each phloem patch. Xylem has spiral tracheæ, internal

FIG. 101.—Diagram of a longitudinal section of a root, showing secondary thickening and the disappearance of the primary cortex. Endodermis then becomes surface layer beneath which is formed the cork by the pericycle. p^1, p^2, p^3 , phloem formed during first, second and third years; xy^1, xy^2, xy^3 , xylem formed during first, second and third years. (Mottier.)

to these a few pitted vessels, then, as the root ages, more pitted vessels, also xylem cells and wood fibers make their appearance.

6. *Pith* (from *plerome*), a small zone of parenchyma cells (not present in all dicotyl roots of primary growth).

HISTOLOGY OF TRANSITIONAL GROWTH AND DEVELOPMENT OF SECONDARY STRUCTURE.

At about six weeks of growth, one notes cells dividing by tangential walls in the inner curve of each phloem patch. This is *intrafascicular cambium*. Its cells in each location start to cut off on their inner side a quantity of *secondary xylem* and add a little *secondary phloem* on their outer side. The protophloem tracts become pushed out and the protoxylem tracts in. The pith (if originally present) disappears as the protoxylem encroaches upon it. Secondary xylem finally fills up the patches between the arms. The pericambium has a tendency to start division into an outer and an inner layer. The outer layer becomes a *cork cambium* (phellogen) surrounding the bundle inside of the endodermis. It cuts off cork tissue on its outer face, hence all liquid material is prevented from filtering through and cortex including endodermis, as well as the epidermis, shrivel, dry up and separate off at the age of two to three months. The cork cambium (phellogen) lays down *secondary cortex* internal to itself and external to the phloem.

Patches of cells of the *inner layer* of *pericambium* divide rapidly and are called *interfascicular cambium*. These join the *intrafascicular cambium* to form a continuous *cambium* ring which then cuts off additional secondary xylem on its inner face and additional secondary phloem on its outer face pushing inward the first-formed or protoxylem and outward the first-formed or protophloem. Medullary-rays are formed by the cambium as it cuts off secondary xylem and secondary phloem elements.

Thus, in a transverse section made through a portion of a Dicotyl root showing *secondary growth*, the following regions are noted passing from periphery to center:

1. Cork
2. Cork cambium (phellogen)
3. Secondary cortex
- Fibro-vascular tissue.

{
 4. Protophloem
 5. Secondary phloem
 6. Cambium
 7. Secondary xylem
 8. Protoxylem

9. Strands of cells extending radially from the cortex to the center of the section separating each open fibro-vascular bundle from its neighbors. These are called medullary-rays.

The xylem or wood portion of each fibrovascular bundle appears as a wedge placed between the xylem portions of two medullary-rays and so is sometimes called a WOOD WEDGE. The xylem portions of the medullary-rays between the wood wedges are termed WOOD-RAYS.

FIG. 102.—Photomicrograph of a transverse section of a California Privet root of primary growth showing epidermis (*ep*); hypodermis (*h*); primary cortex (*pc*); endodermis (*en*); pericambium (*pca*); a xylem arm of the radial bundle (*x*); a phloem arm (*ph*); and pith (*m*). $\times 40$.

HISTOLOGY AND DEVELOPMENT OF A TYPICAL DICOTYL ROOT (CALIFORNIA PRIVET).—A. Make a permanent mount of a cross section of the root of the California Privet (*Ligustrum Californicum*) cut just above the root cap, and note the following structures, passing from periphery toward the center (see Fig. 102):

1. *Epidermis*, composed of a layer of epidermal cells whose outer walls have been infiltrated with a substance called *cutin*.

2. *Hypodermis*, a layer of somewhat thick-walled cells just beneath the epidermis.

3. *Cortex*, composed of cortical parenchyma cells with small, angular, intercellular-air-spaces.

4. *Endodermis*, or innermost layer of cells of the cortex, whose radial walls are lenticularly thickened.

5. *Pericambium* (*Pericycle*), of a layer of actively growing meristematic cells, which has the power of producing lateral rootlets.

FIG. 103.—Photomicrograph of a transverse section of a California Privet root made about $1\frac{1}{2}$ inches above the root tip and showing transition structure. The epidermis (*ep*), primary cortex (*pc*) and endodermis are in the process of sluffing off, since cork (*ck*) has been laid down by the cork cambium (*pn*) directly beneath the endodermis. The cork cambium has also formed several layers of secondary cortex (*sc*) on its inner face. The protophloem (*pp*) represented largely by hard bast has been pushed out, while a small amount of secondary phloem (*sp*) represented by soft bast has been deposited beneath it by the cambium (*c*) which now is nearly circular in aspect. The protoxylem (*px*) has been pushed into the center by the encroaching secondary xylem (*sx*) which has been laid down by the cambium on its inner face. *mr*, medullary ray. Highly magnified.

6. *Radial fibro-vascular bundle* of five or six primary xylem arms alternating with as many primary phloem patches. Note the narrow spiral tracheæ in the xylem patches.

7. *Pith*, a small central zone of parenchyma.

The section you have just studied illustrated in general the appearance of any Dicotyl root of primary growth.

B. Mount permanently another cross section cut through the same root a short distance above the first.

Note that this is somewhat larger in diameter. Observe the *root hairs* starting from the *epidermis*; a broad *cortex*; a large clear

FIG. 104.—Transverse section of California Privet root ($\times 35$) made a short distance above the section shown in Fig. 103 and showing further secondary development. More secondary xylem has been laid down on the inner face of the cambium and there has been a slight increase in the secondary phloem. *ep*, epidermis; *h*, hypodermis; *pc*, primary cortex; *ck*, cork; *pn*, phellogen or cork cambium; *sc*, secondary cortex; *pp*, protophloem; *sp*, secondary phloem; *c*, cambium; *sx*, secondary xylem; *px*, protoxylem.

and open looking *endodermis*; then *pericambium*; next, a central patch of *xylem* showing a faint pentarch relation. Pushed out are five *phloem* tracts. Each of these is composed of a mass of *protophloem* (first formed phloem). On the inner face of each phloem mass may be seen *intrafascicular cambium*. At the outer end of each xylem tract there has been cut off a patch of fine cambial cells (*interfascicular cambium*) which becomes joined to the *intrafascicular cambium* to develop *secondary phloem* on the outer surface and *secondary xylem* on the inner face.

C. Mount permanently a third T. S. cut through the same root a short distance above the second. Note that this is still larger in diameter than the second. The *pericambium* has already divided into an inner and an outer layer. The outer layer has become the *cork cambium*, cutting off the *cork* on its outer face

FIG. 105.—Transverse section of California Privet root made about an inch and a half above the section shown in Fig. 104 and showing early secondary structure. Note that epidermis, primary cortex and endodermis have completely disappeared. Cork (*ck*); phellogen (*ph*); secondary cortex (*sc*); protophloem (*p'*); secondary phloem (*p*²); cambium (*c*); secondary xylem (*x*²) and protoxylem (*x'*). (Photomicrograph, $\times 40$.)

beneath the endodermis. Cork, being an impermeable barrier to water, has prevented the nourishing sap from percolating through to the endodermis, cortex and epidermis. These regions have consequently begun to sluff off. Note that the cambium has begun to spread out into the form of a ring. More *secondary xylem* has been formed on its inner face and additional *secondary phloem* has appeared on its outer face (Fig. 104).

D. Make a permanent mount of a fourth cross section cut through the same root some distance above the third. Note that the *epidermis*, *primary cortex* and *endodermis* have completely peeled off. *Cork* is found as the external bounding layer and underneath it, *cork cambium*. This cork cambium has developed *secondary*

FIG. 106.—Photomicrograph of a transverse section of an older portion of California Privet root, showing further secondary development. $\times 35$. Note the prominent medullary rays (*mr*); cork (*ck*); phellogen (*ph*); secondary cortex (between *ph* and *p'*); protophloem (*p'*); secondary phloem (*p*²); cambium (*c*); secondary xylem (*x*²), the leader pointing to a trachea; wood fibers between tracheæ; and protoxylem (*x'*).

cortex on its inner face. The *cambium* has assumed a circular aspect. Just beneath the secondary cortex will be found flattened patches of *protophloem*, and beneath these *secondary phloem* masses have been formed through the activity of the cambium. The cambium has developed new or *secondary xylem* on its inner face which has pushed the first formed or *protoxylem* toward the center of the root. The pith has disappeared and its place is taken by *protoxylem* (Figs. 105 and 106).

ABNORMAL STRUCTURE OF DICOTYL ROOTS.—In certain Dicotyl roots as the Beet, Jalap, Pareira, and Poke, after the normal bundle system has been formed, there then develop successive circles of cambiums in the cortex outside of these bundles.

FIG. 107.—Cross section of the daughter tuberous root of *Aconitum Napellus*, Spark's variety, cut slightly below mid-region. Photomicrograph $\times 25$. *ep*, epidermis; *pc*, primary cortex; *en*, endodermis; *si*, sieve strands in inner bark; *c*, cambium; *wp*, wood parenchyma; *xy*, tracheary tissue of xylem; *m*, pith.

Each of these cuts off phloem on its outer face and xylem on its inner face, thus producing concentric series of open collateral bundles.

HISTOLOGY OF A DICOTYL TUBEROUS ROOT (ACONITUM).—Transverse sections made through the tuberous roots of *Aconitum Napellus* vary in appearance depending upon the level at which

they are cut and upon whether cut from the daughter or parent tuberous root. In general, the commercial root shows the following internal structure: An *outer layer* consisting either of an *epidermis* or, in older parts, of one or more rows of *suberized cells* with blackish-brown walls, beneath which in younger portions of the root occur a *primary cortex* of starch containing parenchyma in which occur scattered *stone cells* of numerous shapes and a *modified endodermis*, all or a portion of these regions being suberized in older parts, forming a brown tissue called *metaderm* or entirely exfoliated; an *inner bark* of numerous rows of *starch parenchyma* and scattered *islets of sieve tissue*. Stone cells may occur in this zone directly under the endodermis. Next, a five- to eight-angled *cambium*, in each angle of which is a 2- to 4-rayed collateral vascular bundle. In some sections of Aconite roots smaller bundles may be found between the angles and along the cambium line. In the center is found a broad, five- to eight-rayed *pith* composed of large parenchyma cells. The parenchyma cells of the cortical regions and pith contain single or two- to five-compound starch grains.

ROOT BRANCHING.—The branches of roots are formed by the pericycle. The cells of this region adjacent to the protoxylem undergo division by laying down tangential walls forming a mass of meristem. This meristem develops a root tip which pushes its way out laterally through the tissues of the primary root (Fig. 108). In its passage outward, enzymes are secreted by its cells which dissolve the tissues in its path, the root growing into the cavity thereby formed. The first branches of a root are termed *secondary roots*. These may give rise to branch roots and those to further branches by the same method.

ROOT TUBERCLES

The roots of plants of the Bean and Bayberry families as well as some species of the Birthwort family and of the Alders, Cycads, Spruces, etc. are characterized by the appearance upon them of nodule-like swellings called *root tubercles*.

LEGUMINOUS TUBERCLES.—In the case of the members of the Bean family (*Leguminosæ*), the causative factor is a species of bacteria named *Pseudomonas radiculicola*. This is a motile rod-

FIG. 108.—Cross section of root of lupine, showing a secondary root breaking through the cortex. (From Curtis.)

FIG. 109.—Root system of a legume showing tubercles. (Marshall.)

shaped organism which appears widely distributed in soils. It is apparently attracted to the root-hairs of leguminous plants by a chemotaxic influence probably due to the secretions poured out by these structures. A number of these organisms penetrate the walls of the root-hairs by enzymic action. Upon entering the hairs they form *bacterial strands* which branch and rebranch and

FIG. 110.—*Pseudomonas radicicola*. 1, From *Melilotus alba*; 2 and 3, from *Medicago sativa*; 4 and 5, from *Vicia villosa*. (Marshall, after Harrison and Barlow from Lipman.)

extend into the middle cortex cells. Within the cortex cells the organisms multiply rapidly producing nest-like aggregations. Their presence here stimulates the rapid multiplication of the parenchyma cells with the consequent formation of nodules or tubercles. Under oil-immersion magnification these bacteria are often found to exhibit variously shaped forms called involution forms. They remain within the cells of the middle cortex region breaking down the starch and sugar present into simpler compounds. In doing this the *Pseudomonas* obtains the energy required to “fix” the free atmospheric nitrogen. This process is called “*fixation of atmospheric nitrogen*” and consists of uniting the free nitrogen of the air with some other elements to form a nitrogenous compound. This is assimilated into its own body.

The bacteria gradually swell up into *zoöglæa* masses, until finally their bodies break down into soluble nitrogenous substances which are partly absorbed and assimilated and partly stored as reserve nitrogenous food by the green leguminous plant.

ROTATION OF CROPS.—Upon the decay of the leguminous roots and their bacteria in the soil, the nitrogenous compounds are set free and are ready to be decomposed by various soil organisms into the various decomposition products, including amino-acids and ammonia.

In the modern *rotation of crops*, plant growers plough under the leguminous crops or their nodule-producing roots which decay and enrich the soil with ample nitrogenous material to supply the next season's crop of nitrogen-consuming plants, such as cereals, cabbage, potatoes, etc.

Many strains of *Pseudomonas radicola* exist, one for every kind of leguminous crop. Cultures of many of these, such as "Nitro-Germ," etc. are now obtainable from the drug trade.

FIG. 111.—Tubercular clusters on underground stem and roots of a Bayberry; *Myrica Macfarlanei*, observed by the author at North Wildwood, N. J., Jan. 31, 1915.

MYRICA TUBERCLES.—The writer has found tubercles on *Myrica cerifera*, *Myrica Caroliniensis* and *Myrica Macfarlanei* seedling primary roots of 5 to 6 months' growth, and from thence onward on the secondary roots inserted on the hypocotyl axis, on nearly all the adventitious roots of subterranean branches and on the subterranean branches of *Myrica cerifera*, *M. Caroliniensis*, *M. Gale*, *M. Macfarlanei*, and *Comptonia asplenifolia*. The inciting organism has been isolated by him in pure culture, according to Koch's postulates, and named *Actinomyces Myricarum* Youngken.

The tubercles occur either singly, as is frequently the case on subterranean branches, in small groups the size of a pea, or in

FIG. 112.—Nitrogen cycle in the life of alfalfa, a leguminous plant. (From Brownell's *General Science*.)

larger coralloid loose or compact clusters which frequently attain the size of a black walnut. Each tubercle is a short, cylindrical, blunt-ended, root-like structure which branches di- or trichot-

FIG. 113.—Nitrogen cycle. (Marshall.)

FIG. 114.—Ectotrophic mycorrhiza of the beech (*Fagus*); *a*, humus particles; *b*, strands of fungus hyphæ penetrating the soil. (Palladin.)

omously after attaining a certain length. The branches frequently rebranch at their tips which grow out into long thread-like structures from 1–3 cm. in length that may also branch and become entwined about the roots of other plants. The color of the youngest tubercles is a pinkish-gray brown. As the tubercles become older their color changes to brown, dark-brown and even black.

THE NITROGEN CYCLE

Living plants make use of nitrogen in the form of nitrates found in the soil and water. The nitrates are absorbed by the

FIG. 115.—Endotrophic mycorrhiza in epidermal cells of the root of a member of the heath family, *Andromeda polifolia*, the root shown in cross section. (Palladin.)

plant and split up within its cells into the component elements, the nitrogen of which is linked with some other simple elements there present including carbon, hydrogen, oxygen, sulfur and sometimes phosphorus to form proteins. Animals eat the plants and part of the proteins are digested by them and excreted as urea, uric acid and hippuric acid. These products are later decomposed by bacteria (*micrococcus ureæ*, etc.) to form ammonia. The remainder of the proteins are assimilated and stored in the growing animal cells. When the animals die their bodies decay and the protein or nitrogenous compounds are split into decomposition products (peptones, amino-acids, ammonia) of which ammonia is the final end product.

The soil contains NITRIFYING BACTERIA. Of these the *Nitrosomonas* and *Nitrosococcus* secrete enzymes which oxidize ammonia to nitrous acid. Some of the nitrous acid is neutralized by alkaline substances found in the soil to form nitrites. Another nitrifying organism, the *Nitrobacter* secretes enzymes which oxidize nitrous acid and nitrites forming nitric acid; free nitric acid then combines with other substances (bases) in the soil forming nitrates, thus completing the cycle.

MYCORRHIZA

The roots of many plants, especially those growing in humus, are associated with fungal hyphæ which form a weft over their surface, as in the case of the beech, etc., when this weft is called an *ectotrophic mycorrhiza*, or the surrounding hyphæ may penetrate into the cells of the roots, as in many plants of the Heath and Orchid families, when the association is termed an *endotrophic mycorrhiza*.

There are a number of plants, particularly non-green seed plants, which cannot thrive without the mycorrhizæ with which their roots are normally associated and die when transplanted to soil not containing them. Mycorrhizæ assist the roots to absorb nutrient materials from the soil.

A number of mushrooms form ectotrophic mycorrhizas on the roots of trees.

CHAPTER VIII

THE BUD

BUDS are short, young, undeveloped stems or shoots with rudimentary leaves compactly arranged upon them.

The *plumule* represents the first bud on the initial stem or caulicle.

Scaly buds are such as have their outer leaf rudiments transformed into scales; these are often coated with a waxy or resinous substance without and a downy lining within to protect them from sudden changes in climate. Buds of this character are common among shrubs and trees of temperate regions. The official Poplar Bud is an example.

Naked buds are those which are devoid of protective scales. They are common to herbaceous plants. The rose buds, head of cabbage or brussels sprouts are good examples.

CLASSIFICATION OF BUDS ACCORDING TO DEVELOPMENT.—1. A *leaf bud* is a young shortened shoot bearing a number of small rudiments of leaves. It is capable of elongating into a branch which bears leaves. The leaf buds of *Populus balsamifera* and *P. candicans* are official in the National Formulary.

2. A *flower bud* is a rudimentary shoot bearing one or more concealed and unexpanded young flowers. The drug Clove is an example of a flower bud which, if allowed to expand, would form a single flower.

3. A *mixed bud* is an undeveloped stem or branch bearing concealed, unexpanded leaves and flowers. The buds which give rise to flowers and leaves on the apple trees are mixed buds.

CLASSIFICATION OF BUDS ACCORDING TO POSITION ON THE STEM.—1. A *terminal bud* is one which is located on the end of a stem (shoot). It is capable of elongating into a shoot which bears leaves or both leaves and flowers.

2. An *axillary* or *lateral bud* is one which arises in the leaf axil. It is capable of giving rise to a side branch or to a flower. Occa-

sionally axillary buds do not develop and are then called *dormant buds*.

3. An *adventitious bud* is one which occurs on some position of the plant other than at the apex of the stem or in the axil of a leaf. Such buds may be seen developing along the veins of a Begonia leaf or in the notches along the margin of a Bryophyllum leaf after these have been planted in moist soil for several days. The roots of the poplars, willows, cherries and many other trees develop adventitious buds underground which develop *suckers* which may eventually become mature plants that lead an independent existence.

4. An *accessory bud* is an extra bud which forms in or near the leaf axil. These may be seen on the young stems of the Red Maple.

5. A *sub-petiolar bud* is one arising beneath the basal part of the petiole or leaf stalk, so that when the leaf falls off the bud becomes exposed, as in the Buttonwoods.

CLASSIFICATION OF BUDS ACCORDING TO THEIR ARRANGEMENT ON THE STEM.

1. When a single bud is found at each joint or *node* of a stem, the buds are said to be *alternate*.

2. When two buds are found at a node they are *opposite*.

3. When several buds occur at a node they are *whorled*.

FIG. 116.—Cottonwood twig, two years old. (From Robbins after Longyear.)

CHAPTER IX

THE STEM

The STEM is that part of the plant axis which bears leaves or modifications of leaves and its branches are usually arranged with mathematical regularity.

Stems usually grow toward the light and so are positively heliotropic.

The principal functions of a stem are to bear and support leaves, branches and reproductive organs, connect roots with leaves, conduct water with minerals in solution from roots to leaves, to transport elaborated plant foods from the leaves to the roots and to store reserve foods.

When the stem rises above ground and is apparent, the plant is said to be *caulescent*, as the various trees, shrubs, geraniums, etc.

When no stem is visible, but only flower or leaf stalks, the plant is said to be *acaulescent*, as the Bloodroot and Violet.

Stems vary in size from scarcely $\frac{1}{25}$ inch in length, as in certain mosses, to a remarkable height of 400 feet or more. The giant Sequoia of California attains the height of 420 feet and has a trunk diameter at its base of between 25 and 30 feet. Some of the Eucalyptus trees of Australia and Tasmania are reported to attain the height of 500 feet.

NODES AND INTERNODES.—The nodes are the joints of stems. They represent the parts of the stem from which leaves or branches arise. Internodes are the parts of stems between nodes.

DIRECTION OF STEM GROWTH.—Generally the growth of the stem is upright or *erect*. Very frequently it may be:

Ascending, or rising obliquely upward. Example: Saw Palmetto.

Reclining, or at first erect but afterward bending over and trailing upon the ground. Examples: Raspberry and Blackberry.

Procumbent, lying wholly upon the ground. Example: Pipsissewa.

Decumbent, when the stem trails and the apex curves upward. Examples: Vines of the *Cucurbitaceæ* or Gourd family.

Repent, creeping upon the ground and rooting at the nodes, as the Strawberry.

STEM ELONGATION.—At the tip of the stem there is found a group of very actively dividing cells (*primordial meristem*) which is the growing point of the stem. All the tissues of the stem are derived from the cells of the growing point whose activity gives rise in time to three generative regions which are from without, inward:

1. *Dermatogen* or *protoderm* forming epidermis;
2. *Periblem* or *ground meristem* forming the cortex, pericycle, and pith.
3. *Plerome* or *procambium* forming the fibro-vascular elements.

DURATION OF STEMS.

Annual, the stem of an herb whose life terminates with the season. Example: Corn.

Perennial, when the stem lives for many years. Example: Oak.

FORMS OF STEMS.—The following adjectives appear in pharmacognostic and botanical literature in relation to the form of stems and in some instances other plant organs: *Cylindrical* or cylinder-shaped; *subcylindrical* or somewhat cylinder-shaped; *terete* or circular in cross section; *compressed* or pressed together above and below or laterally; *triquetrous* or three-angled, *quadrangular*, or four-angled; *alate* or winged; *exfoliating* or

FIG. 117.—End portion of a branch of the horse-chestnut (*Aesculus hippocastanum*) in the winter condition. (Mottier.)

shedding cork in layers; *obconical*, or having the shape of an inverted cone; *flexuous* or bent; *tortuous* or twisted; *truncate*, or ending abruptly, as if cut across at the summit.

STEM MODIFICATIONS.

Twining stems are those which coil spirally around a support, as the stems of the Hop, Bittersweet, Dodder, etc.

Tendriform stems are those which undergo thread-like modification and become sensitive to contact of a side branch or other

object, coiling around it, as in Passion flower, Grape, Squash, etc.

Spines are formed by the checking and hardening of a branch that may then become defensive, as in hawthorn, honey locust, etc.

Aerial tuberous stems are those in which one or more internodes enlarge above ground and store reserve food, as in pseudobulbs of orchids.

Subterranean tuberous stems are those in which an underground stem or branch enlarges

as a food-storing center: (a) annual type, tuber, as in potato, etc., corm, as in crocus, etc.; (b) perennial type, bulbs as in lily (scaly) and onion or hyacinth (tunicated).

Phylloid stems or *phylloclades* are green, flat, leaf-like stems in which flattening branch expansion has occurred, as in *Asparagus*, *Ruscus*, etc.

Cactoid stems are those in which reduced, condensed branches or stems become swollen for water (and food) storage, as in Cacti, some *Euphorbia* species, etc.

ABOVE-GROUND OR AERIAL STEMS.—A *twining* stem winds around a support, as the stem of a bean or Morning Glory.

A *culm* is a jointed stem of the Grasses and Sedges.

A *climbing* or *scandent* stem grows upward by attaching itself to some support by means of aerial rootlets, tendrils or petioles. Examples: Ivy, Grape, etc.

FIG. 118.—Diagram of a longitudinal section through the tip of a stem. 1, 2, 3, 4, successively older beginnings (primordia) of leaves. (Mottier.)

The *scape* is a stem rising from the ground and bearing flowers but no leaves, as the dandelion, violet, or blood root.

A *tendrils* is a modification of some special organ, as of a leaf stipule, leaflet or branch, capable of coiling spirally and used by a plant in climbing. The tendrils of the Grape Vine are modified inflorescence branches, those of Sarsaparilla are modified stipules and those of the Pea are modified leaflets. The ends of the tendrils of the Japan Ivy become swollen and flattened, forming adhesive discs which cling to objects with which they come into contact.

A *spine* or *thorn* is the indurated termination of a stem tapering to a point, as the thorns of the Honey Locust.

Prickles are outgrowths of the epidermis and cortex and are seen on the stems of the roses, green-briers, etc.

A *stolon* or *runner* is a prostrate or reclining branch, the end of which, on coming in contact with the soil, takes root, so giving rise to a new plant. Examples: Currant and Raspberry. Short stolons like those of the Houseleek are called *offsets*.

A *bulbel* or *bulbil* is a small, bulb-shaped, young aerial shoot which serves as an organ of vegetative multiplication by falling off the stem and developing into a new plant. The Garlic and Tiger Lily are good examples of plants which produce bulbels.

An *herbaceous stem* is one which is soft in texture and readily broken. Example: Lily-of-the-Valley.

An *undershrub* or *suffruticose* stem is a stem of small size and woody only at the base. Examples: Bittersweet, Thyme, etc.

FIG. 119.—Tendrils of Bryonia (*Bryonia dioica*). (Palladin.)

A *shrubby* or *fruticose* stem is a woody stem larger than the preceding and freely branching near the ground. Example: Lilac, etc.

A *trunk* is the woody main stem of a tree.

A *twig* is that portion of a woody branch which has developed during the last growing season.

HERB, SHRUB AND TREE

An **HERB** is a plant whose stem does not become woody and permanent, but usually dies, at least down to the ground, after

FIG. 120.—An elm tree in summer and winter. To illustrate alike the deliquescent habit of branching and the deciduous nature of this tree (dropping its leaves in autumn and developing new ones in the spring). (Atwood.)

flowering. Herbs are either annual, biennial or perennial (see pp. 162–163). When a perennial herb possesses an aerial stem which develops leaves that persist on the stem over winter, as the Partridge Berry, it is called an *evergreen herb*.

A **TREE** is a perennial woody plant of considerable size, attaining a height of 15 or more feet, and having as the above-ground parts a trunk and a crown of leafy branches.

There are two plans of branching in trees. When the trunk, or main stem, extends vertically upward to the tip, as it does in the pines, junipers, spruces and other conical trees, the type of branching is called *excurrent*; when it divides into several more or less equal divisions as in the elm and other spreading trees, it is said to be *deliquescent*. The deliquescent plan is the more common one among our deciduous trees.

FIG. 121.—A, tunicate bulb of Onion (*Allium Cepa*), in longitudinal section; B, Scaly bulb of Canada Lily (*Lilium canadense*); C, Same as B, but in longitudinal section; D, Bulb of Garlic (*Allium sativum*) with portion of the scales cut away so as to enclose the young bulbs within; E, Portion of stem of Tiger Lily (*Lilium tigrinum*), showing bulbils in the axils of the leaves; G, Corm of Crocus showing a terminal and one lateral bud. The horizontal lines represent the scars of the few scales that have been removed. At the base is the remnant of last year's corm. The various buds will develop into new corms. F, Vertical section of a familiar corm. H, flat corm of Cyclamen; I, Bulblets at the apex of a flower stalk of the Onion. (After Gager.)

A **SHRUB** is a perennial woody plant of smaller stature than a tree whose stem is relatively short and usually branches near the ground. Examples: Viburnum, Elder, Buchu, Krameria, etc.

FALL OF LEAVES.—The fall of leaves in autumn from our deciduous woody dicotyledons and gymnosperms is caused by the formation beneath the leaf stalks of an abscission or cut-off layer or band of delicate cells. The cellulose walls of these break down into pectic acid and pectin which forms a mucilaginous substance, the middle lamellae between the cells disintegrates, the veins become ruptured and the leaf falls of its own weight. A protec-

tive layer of cork is developed beneath the abscission layer either before or after the fall of the leaf which seals the wound and forms the leaf scar. The shape of many of the leaf scars are diagnostic for the stems bearing them.

UNDERGROUND OR SUBTERRANEAN STEMS.—A **RHIZOME** is a creeping underground stem, more or less scaly, sending off roots from its lower surface and stems or leaves from its upper. The rhizome grows horizontally, vertically or obliquely, bearing a

FIG. 122.—Rhizome of Solomon's Seal (*Polygonatum biflorum*). 1, scar of last year's overground stem; 2, base of this year's aerial stem; 3, terminal bud of next year's aerial stem. *sc*, scale leaves; *s*, scars (annulations) of old scales; *r*, rootlets. $\frac{5}{8}$ natural size. (Gager, after Gapp.)

terminal bud at its tip. Its upper surface is marked with the scars of the bases of aerial stems or leaves of previous years. Examples: Triticum, Rhubarb, etc.

The **TUBER** is a short and excessively thickened underground stem, borne usually at the end of a slender, creeping branch, and having numerous "eyes" or axillary buds. Examples: Tubers of the Potato and Jerusalem Artichoke.

The **CORM** is an underground stem excessively thickened and solid and characterized by the production of buds from the center of the upper surface and rootlets from the lower surface. Examples: Colchicum, Jack-in-the-Pulpit, Crocus, etc. (See Fig. 127.)

A **BULB** is a very short underground stem invested by fleshy scales which represent storage leaves. It produces rootlets from its lower face and leaves and flower from its upper.

Tunicated bulbs are completely covered by broad scales which form concentric coatings. Examples: Onion, Squill, Daffodil.

Scaly bulbs have narrow imbricated scales, the outer ones not enclosing the inner. Example: Lily. (See Fig. 127.)

A **SUCKER** is a slender underground branch that arises either on a root or a stem. *Root suckers* are produced by roses, poplars, elms, etc. and arise as adventitious buds on the roots of these plants. *Stem suckers* are produced by peppermint and other mints, etc., arising from the axils of scaly leaves on the rhizome,

FIG. 123.—Portion of a sprouting potato tuber. (After Robbins.)

or they may arise from wound tissue which has developed on sawed-off branches or from old wood near the base of old trees.

EXOGENOUS AND ENDOGENOUS STEMS.—*Exogenous stems* are typical of Gymnosperms and Dicotyledons and can increase materially in thickness due to the presence of a cambium. Such stems show differentiation into an outer bark cylinder and an inner or wood cylinder. Between the bark and the wood cylinders is a cambium.

Endogenous stems are typical of most Monocotyledons and cannot increase materially in thickness due to the absence of cambium. The limited increase in diameter that does take place is due to the enlargement of the cells of the primary tissues. Such stems show no differentiation into bark and wood regions.

FIG. 124.—Sunflower stem. Representative portion of a transverse section cut through upper, young part of stem. *ep*, epidermis bearing trichomes (*t*); *hy*, hypodermis; *co*, primary cortex; *pf*, pericyclic fibers; *p*, protophloem of primary fibrovascular bundle; *c*, intrafascicular cambium of bundle; *x*, protoxylem of bundle; *m*, pith or medulla. All of the tissue between each two bundles is a primary medullary ray. Greatly enlarged.

HISTOLOGY OF ANNUAL DICOTYL STEM.—Excellent stem material of an annual dicotyledon for the study of the stems of this type is to be found in the sunflower. Transverse sections cut through the upper, thin, younger portion of this stem, present the following structures: (In both annual and perennial dicotyledonous stems, endodermis and pericycle are rarely dis-

tinguishable from the cortex, except for the knowledge of their relative position, since each has become so similar to the cortex through passage of food, etc.)

1. EPIDERMIS, a single layer of somewhat flattened cells whose outer walls are cutinized. Some of the cells show outgrowths as hairs. Stomata also occur.

FIG. 125.—Diagram of a representative portion of a transverse section cut through an old part of the Sunflower stem. *ep*, epidermis; *col*, collenchyma; *co*, cortex; *p.f.*, pericyclic fibers in pericycle; *ph*, protophloem; *ph'*, secondary phloem; *cam*, intrafascicular cambium; *cam'*, interfascicular cambium; *x*, protoxylem; *x'*, secondary xylem; *mr*, primary medullary ray; *mr'*, secondary medullary ray; *m*, pith. The three longest bundles represent primary bundles which have been broadened by the cambium adding secondary phloem outwardly and secondary xylem inwardly. The smaller bundles represent secondary bundles, each consisting of secondary phloem, cambium and secondary xylem.

2. PRIMARY CORTEX, composed of two zones: an outer region of several layers of narrow, thick-walled cells (hypodermis) and a broad zone of larger, thin-walled cortical parenchyma cells, many of which contain chloroplasts.

3. The innermost layer of cells of the cortex is called the ENDODERMIS. (Not generally distinguishable.)

4. PERICYCLE (Pericambium). Not generally distinguishable in most dicotyl stems but showing early formation of groups of fibers in sunflower stems.

5. PRIMARY FIBRO-VASCULAR BUNDLES of open collateral type arranged in a circle with primary medullary rays between the bundles. Each bundle shows an outer *protophloem*, a *cambium* called *intrafascicular cambium*, because it is within the bundle, and an inner *protoxylem*.

6. PITH, composed of parenchyma cells.

If a series of additional cross sections are now cut at different descending levels along the Sunflower stem until the old part of

FIG. 126.—Portion of a longitudinal section of a sunflower (*Helianthus*) stem. (From "A Text Book of General Botany" by Smith, Overton et al. The Macmillan Co. Publishers.)

the stem near the ground is reached, and these be examined microscopically, it will be noted that some secondary changes have taken place which explain the gradual increase in thickness of this stem as the base is approached. One of the first of these changes incident to secondary growth is that observed in the cells of the primary medullary rays between the *intrafascicular cambia* of the various bundles. These lay down tangential walls to form *interfascicular cambia* or *cambiums* between the bundles. When this process is completed, a continuous *cambium ring* is formed.

The cambium ring adds more xylem elements called *secondary xylem* on its inner face and more phloem elements called *secondary*

FIG. 127.—A diagram to show the character of the tissues and their disposition in a young stem of the typical dicotyledon type. (From Stevens.)

phloem on its outer face. As it cuts off sieve tubes and companion cells outwardly and tracheæ and wood fibers inwardly all along the line it also cuts off here and there a medullary ray cell on

each of its faces. It will be noted that more secondary xylem is added than secondary phloem. The result of the activity of the cambium ring is the deepening of the primary bundles and the

FIG. 128.—Diagram similar to the preceding but representing a later stage and showing the tissues formed by the cambium. (From Stevens.)

primary medullary rays, the appearance of new or *secondary bundles* (which are smaller than the primary bundles) in the primary medullary rays and the appearance of narrow secondary medullary rays between the secondary bundles as well as inside

of the deepened primary bundles. The secondary medullary rays within the bundles are also called "vascular rays."

As secondary increase progresses, it will be noted that certain cells in widely separated regions of the pericycle begin to divide rapidly producing groups of thick-walled *pericyclic fibers* here and there within the pericycle but chiefly outside of the protophloem of the larger or primary bundles.

FIG. 129.—Photomicrograph of cross-section of stem of *Aristolochia siphon*, where cambial activity is just beginning. *a*, Epidermis; *b*, collenchyma; *c*, thin-walled parenchyma of the cortex, the innermost cell layer of which is the starch sheath or endodermis; *d*, sclerenchyma ring of the pericycle; *e*, thin-walled parenchyma of the pericycle; *f*, primary medullary-ray; *g*, phloem; *h*, xylem; *i*, interfascicular cambium; *j*, medulla or pith. $\times 20$. (From Stevens.)

It will be noted that cross sections through the young terminal portion of the sunflower stem show an interrupted cylinder of separate bundles whereas similar sections cut through the older lower portion of the stem show a continuous cylinder of bundles, the continuity being due to the laying down by the interfascicular cambia of secondary bundles between the primary bundles during secondary growth. The stele or central cylinder of the younger part of this stem is known as a *dictyostele* or *dissected siphonostele* whereas that of the older portion is a *siphonostele* (see Fig. 83).

GROWTH OF PERENNIAL DICOTYL STEM AND ITS HISTOLOGY.—A perennial dicotyl stem, in the first year of its growth does not

differ in structure from an annual. By the close of the year a cork cambium (phellogen) has originated beside the epidermis. In the origin of the *cork cambium*—one of two methods: (a) either the epidermis may divide into an outer layer of cells that remains

FIG. 130.—Portion of cross-section of four-year-old stem of *Aristolochia sipho*, as shown by the rings of growth in the wood. The letters are the same as in Fig. 129 but new tissues have been added by the activity of the cambium; and a cork cambium has arisen from the outermost collenchyma cells and given rise to cork. The new tissues are: *l*, cork cambium; *k*, cork; *g*, secondary phloem from the cambium, and just outside this is older crushed phloem; *n*, secondary xylem produced by the cambium; *m*, secondary medullary ray made by the cambium (notice that this does not extend to the pith). Half of the pith is shown. Notice how it has been crushed almost out of existence. Compare Figs. 129 and 130, tissue for tissue, to find out what changes the primary tissues undergo with age, and to what extent new tissues are added. Photomicrograph $\times 20$. (From Stevens.)

epidermis and an inner layer of cells that becomes cork cambium, or, (b) the outermost layer of cortex cells underneath the epidermis becomes active after being passive for one year, and lays down walls, the inner layer becoming cork cambium, the outer becoming a layer of cork. The cork cuts off water and food supplies from the epidermis outside, and so the epidermis sepa-

rates and falls off as a stringy layer. The cork cambium produces cork on its outer face and secondary cortex on its inner.

Between the bundles, certain cells of the primary medullary-rays become very active and form *interfascicular cambium* which

FIG. 131.—Diagrams, based on the red oak, *Quercus rubra*, showing the position and extent of successively formed periderm or corky layers in a typically woody stem. *A*, a one-year-old twig, the first periderm layer, a complete cylinder, formed beneath the epidermis. *B*, a two-year-old twig, the epidermis and first periderm ruptured; new, shell-shaped layers formed deeper in the cortex. *C*, a three-year-old stem, the outer tissues weathered away and more periderm layers formed still more deeply in the stem, invading the secondary phloem. *D*, a four-year-old stem, the cortex and outer secondary phloem with their periderm layers weathered away, the new cork layers invading the younger phloem. *E*, the outer tissues of an old tree-trunk, showing the narrow band of young, living secondary phloem, and the thick, deeply fissured layer of older, dead phloem with its many shell-shaped periderm layers; a considerable amount of similar tissue has exfoliated. (From Eames and MacDaniels' "Introduction to Plant Anatomy," McGraw-Hill Book Co., Inc., Publishers.)

joins the cambium of the first-formed bundles (*intrafascicular cambium*) to form a complete *cambium ring*. By the rapid multiplication of these cambial cells, new (secondary) xylem is cut off internally and new (secondary) phloem externally, pushing inward the first-formed, or *protoxylem*, and outward the first-

formed, or *protophloem*, thus increasing the diameter of the stem. The primary medullary-rays are deepened. Cambium may also give rise to secondary medullary-rays.

Sometimes, as in Grape Vines, Honeysuckles, Hickories, Cherries, Viburnums, and Oaks, etc., instead of cork cambium arising only from the epidermis or the outer layer of cortex cells, it may arise at any point in the cortex or even as far inward as the pericycle or outer layers of phloem, and frequently several cork cambia appear successively. As each new cork cambium arises and cuts off cork cells on its outer face, all the living tissues outside of the cork are deprived of nutrient sap and die. It is the origin of cork cambia at varying depths that causes extensive sheets of tissue to separate off. This dead tissue has been termed *borke* by the German botanists (see Fig. 131). That is what gives the stringy appearance to the stems of climbers.

At close of first year in a *Perennial Dicotyl Stem* we note:

1. Epidermis—development of dermatogen—in process of peeling off, later on entirely absent.
2. Cork tissue.
3. Cork cambium or phellogen.
4. Sometimes a zone of thin-walled cells containing chloroplasts cut off by cork cambium on inner face and known as *phelloderm* or secondary cortex.
5. Primary cortex—in perennials, stem cells of cortex may undergo modification into mucilage cells, into tannin receptacles, crystal cells, spiral cells, etc.
6. Endodermis, or innermost layer of cells of cortex, usually similar to other cortex layers in character of cells.
7. Pericycle, a parenchymatous region usually appearing similar to the cortex, but sometimes containing sclerenchyma fibers or stone cells or both.
8. Vascular bundles of open collateral type which are now arranged in a compact circle, and between which are found primary- and often secondary medullary rays. The longest bundles represent Primary Bundles which have been deepened by the cutting off of secondary xylem and phloem by the cambium.

From without inward the following tissues make up the bundles:

Protophloem	{	Hard Bast—long tenacious bast fibers (sometimes absent).
Secondary Phloem		Soft Bast—phloem cells, companion cells and sieve tubes.

Cambium—active layer giving rise to secondary phloem on outer and secondary xylem or inner face, and adding to depth of med. rays.

Secondary xylem—wood fibers, pitted vessels, tracheids.

Protoxylem—spiral tracheæ.

9. Pith.

EXCEPTIONAL TYPES OF DICOTYL STEMS

In a number of Dicotyledons and Gymnosperms, the secondary growth in thickness of the stem and frequently of the root differs from that which is found in the vast majority of species and so is called exceptional or anomalous.

In stems of *Phytolacca*, etc., there first arises a ring of primary bundles with broad, loose medullary-rays. Then the stem

FIG. 132.—Transverse section of the herbaceous dicotyledonous stem of *Solanum tuberosum* (Fam. Solanaceæ) showing bicollateral bundles with internal phloem. $\times 75$. *ep*, epidermis; *co*, cortex; *ph*, external phloem; *ca*, cambium; *x*, xylem; *ip*, internal phloem; *si*, sieve tissue; *m*, medulla. Both phloem regions contain primary phloem.

cambium ceases its activity and, outside the bast of the bundles already formed, in the pericycle or tissue developing from it, a new cambium starts to lay down another ring of bundles in rather irregular fashion. Then, after developing a wavy ring of bundles and connecting tissue, the cambium closes up. Still another cambium ring arises without this which lays down another circle of bundles and, in a single season quite a number of these are found successively arranged in concentric fashion.

In *Gelsemium*, species of *Solanaceæ*, *Combretaceæ*, *Cucurbitaceæ*, etc., there arises a cambium on the inner face of the xylem which

FIG. 133.—White birch (*Betula populifolia*). Portion of a branch showing the prominent lenticels. (Gager.)

forms *internal phloem* (or *intraxylary phloem*), thus giving rise to bicollateral bundles. (See Fig. 132.)

In *Strychnos Nux Vomica*, internal phloem, exactly as in *Gelsemium*, etc. appears, but in addition *interxylary phloem* is developed. In the wood region of this plant axis the cambium starts at a certain age to lay down patches of phloem which become wedged in between xylem tissue as *interxylary phloem*.

LENTICELS AND THEIR FORMATION.—*Lenticels* are openings in the periderm which provide a means of communication between the external air and the living cells of the bark beneath. The epidermis in a great majority of cases

produces stomata, apertures, surrounded by a pair of guard cells, which function as passages for gases and watery vapor from and to the active cells of the cortex beneath.

FIG. 134.—Cross-section through a lenticel of *Sambucus nigra*. *E*, Epidermis; *PH*, phellogen; *L*, loosely disposed cells of the lenticel; *PL*, cambium of the lenticel; *PS*, phelloderm; *C*, cortical parenchyma containing chlorophyll. (From Sayre after Strasburger.)

There very early originate in the region beneath some of the stomata loosely arranged cells called "*complimentary cells*" which

are cut off by the cork cambium in certain places instead of cork. These swell up during rain and pressing outward, form fissures in the epidermis, called lenticels.

The function of lenticels is similar to that of stomata, namely, to permit of aëration of delicate cells of the cortex beneath.

FIG. 135.—Representative portion of a transverse section of a Linden stem, 3 years old.—1x, 2x, 3x, Successive annual rings of wood. *ph*, Phloem, with bast fibers showing as lighter parts and sieve-tubes as darker parts. *c*, Cortex containing chloroplasts and rosette crystals of CaO . *m.r.*, spreading end of primary medullary ray. *pe*, periderm. (From Small, after Kny.)

While common to many plants, they are very conspicuously prominent in the birches and cherries.

ANNUAL THICKENING.—In all woody exogenous stems, such as trees and shrubs, the persistent cambium gives rise to secondary xylem thickening every spring, summer and autumn. Soon a great cylinder of xylem arises which constitutes the wood of the trunk and branches. In the spring, growth is more active, and large ducts with little woody fiber are produced while in summer and autumn growth is lessened and small ducts and much mechanical woody fiber are formed. Thus the open, loosely

arranged product of the spring growth abuts on the densely arranged product of the last summer and autumn growth and the sharp contrast marks the periods of growth. To the spring, summer and autumn regions of growth of each year is given the term of "ANNUAL RING." By counting the number of these rings it is possible to estimate the age of the tree or branch.

FIG. 136.—Stem of Linden, showing detailed structure of a small square area of Fig. 135, magnified 200 times. *S*, sieve tubes; the dark spaces at one corner of the sieve-tubes are companion cells. *V*, pitted tracheæ. The summer wood consists of wood fibers interspersed with parenchyma. (From F. O. Bowers' "Plants and Man." By permission of the Macmillan Co., Publishers.)

BARK.—Bark is a term applied to all that portion of a woody exogenous plant axis outside of the cambium line.

In pharmacognic work, bark is divided into three zones, these from without inward being:

1. *Outer Bark or Cork region (sometimes with epidermis).*
2. *Middle Bark or Cortex region.*
3. *Inner Bark or Phloem region (sometimes with pericycle attached).*

CORK.—Cork or *phellem* is all that tissue produced by the cork cambium on its outer face. Its walls become impregnated with a waterproof substance of a waxy character called *suberin*.

FIG. 137.—Part of a cross-section through branch of *Cytisus laburnum*. (The branch was cut from the tree at the end of October.) From A to E the last annual ring of wood; from A to B the spring growth with large tracheal tubes (T, T, T); between B and C and D and D are wood-fibers; between C and D and D and E, wood parenchyma; from E to F, cambium; F to G, phloem portion; G to H, cortical parenchyma; M, medullary ray. Below A the last wood-fibers and wood parenchyma formed the previous year. (From Sayre after Haberlandt.)

PERIDERM.—Periderm is a name applied to all the tissue produced by the cork cambium (*Phellogen*) externally and inter-

nally and includes the cork cambium itself. This term appears often in pharmacognic and materia medica texts.

PHELLODERM.—Phelloderm or secondary cortex is all that tissue produced by the cork cambium on its inner face. Its cells frequently contain chloroplasts.

FIG. 138.—Cascara Sagrada Bark. Transverse section. *K*, cork; *kk*, cork cambium; *ph*, phelloderm; *st*, group of stone cells; *C*, primary cortex; *rcr*, rosette aggregate of calcium oxalate; *P*, phloem; *bf*, bast fibers; *crf*, crystal fibers more or less surrounding a group of bast fibers; *mr*, medullary ray; *mp*, monoclinic prism of CaO .

HISTOLOGY OF A TYPICAL BARK, CASCARA SAGRADA.—*In transverse section* passing from outer to inner surface, the following structural characteristics are evident:

1. **CORK**, or outer bark, composed of several layers of rectangular cork cells. The most external layers are dead and appear black because they are filled with air. The inner layers of this region are living, and contain brownish, brownish red or purple contents.

2. CORK CAMBIUM (phellogen), a layer of delicate cells in the process of division with protoplasmic contents.

FIG. 139.—Cascara Sagrada Bark. Radial-longitudinal section. *k*, cork, *kk*, cork cambium; *ph*, phelloderm of collenchymatous cells; *c*, primary cortex; *st*, group of stone cells; *mp*, monoclinic prism; *rcr*, rosette crystal of calcium oxalate; *mr*, medullary ray; *bf*, bast fibers; *s*, sieve tube; *p*, phloem; *crf*, crystal fibers adhering to bast fibers.

3. CORTEX, or middle bark, consisting of two regions, viz.: an outer zone of two or three rows of brownish collenchyma cells, and an inner broader zone of tangentially elongated, cortical parenchyma cells. Imbedded within this zone will be noted numerous groups of stone cells. Rosette crystals and monoclinic

prisms of calcium oxalate occur in parenchyma cells of this region.

4. PHLOEM, or inner bark, a very broad zone composed of irregular-shaped, elongated phloem masses separated from each other by medullary-rays which converge in the outer phloem region. Each phloem mass consists of numerous sieve tubes,

FIG. 140.—Cascara Sagrada Bark. Tangential-longitudinal section through outer phloem region. *p*, phloem parenchyma; *st*, group of stone cells; *rcr*, rosette aggregates of calcium oxalate; *mr*, medullary ray; *si*, sieve tubes; *crf*, crystal fibers containing monoclinic prisms of calcium oxalate; *mp*, monoclinic prism.

companion cells and phloem cells, some of which latter contain spheroidal starch grains while others contain either monoclinic prisms or rosette aggregates of calcium oxalate. Embedded within the phloem masses, in tier-like fashion, will be noted groups of bast or phloem fibers, each group of which is surrounded by a row of crystal fibers, individual cells of which can only be made out in this kind of a section. Each of these contains a monoclinic prism of calcium oxalate. The medullary-

rays possess brownish contents which take a red color with an alkaline solution.

In radial longitudinal section, a lengthwise view of the tissues will be seen. The medullary-rays appear 15 to 25 cells in height and crossing at right angles to the other elements. The crystal

FIG. 141.—Transverse section of representative portion of the stem of the second year's growth of spruce (*Picea* sp.), a conifer. ($\times 80$.) *ep*, epidermis; *ck*, cork; *pn*, phellogen; *co*, cortex; *sr*, secretion reservoir; *ph*, phloem; *ca*, cambium; *xr*, xylem ray; *td*¹, tracheids of autumn wood of first year; *td*², tracheids of wood of second year.

fibers here will be seen to be composed of vertical rows of superimposed thin-walled cells each of which contains a monoclinic prism of calcium oxalate. The phloem (bast) fibers appear elongated and taper-ended and are associated with crystal fibers. The sieve tubes show sieve plates on their longitudinal as well as their transverse walls in this bark.

In a tangential longitudinal section which has been cut through the phloem, the exact range in width of the medullary-rays may

FIG. 142.—Diagrammatic representation of a block of pine wood, highly magnified. *a*, Early growth; *b*, late growth; *c*, intercellular space; *d*, bordered pit in tangential wall of late growth; *m*, *f* and *e*, bordered pit in radial wall of early growth from different points of view; *h*, row of medullary cells for carrying food; *g*, row of medullary ray cells for carrying water; *k*, thin plate in radial wall of ray cells that carry food. (From Stevens.)

be ascertained. In this bark the medullary-rays are spindle-shaped in tangential view and one to four cells in width.

WOOD.—From a practical pharmacognic standpoint as well as that of the lumber trade, wood is all that portion of woody exogenous plant axis inside of the cambium line. In Dicotyl

and Gymnosperm stems it therefore includes the xylem regions of the bundles, the xylem portions of the medullary-rays and the pith, while in the roots of secondary growth of these plants it comprises the xylem portions of the bundles and the xylem medullary-rays. From the pure botanical viewpoint, wood is synonymous with xylem and so represents that portion of a vascular bundle that contains tracheæ or tracheids or both.

As the cambium year after year adds new layers of wood to that already present on its inner face, the conveying of sap and storing of starch, etc. is gradually relegated to the outer wood layers, since the inner layers, step by step, lose their protoplasmic contents and power of conducting sap and become filled with extractive, resinous and coloring matters. The outer whitish layers of wood which contain living cells, functioning in the vegetative processes of the plant, constitute the *ALBURNUM* or *SAP-WOOD*. The drug *Quassia* is a good example of this kind of wood. The inner dead colored layers constitute the *DURAMEN* or *HEART-WOOD*. Important examples of this kind of wood used in pharmacy are *Lignum Guaiaci*, *Hæmatoxylon* and *Santalum Album*.

TYLOSES.—These are bladder-like enlargements of cell walls usually of wood parenchyma or medullary ray cells which extend into the lumina of adjacent tracheæ or tracheids through pit cavities in the walls of the latter. They are formed by the surface growth of the pit membranes of the one-sided bordered pits which push their way into the lumina of the vessels or tracheids. In its living state the tylosis contains cytoplasm, cell sap, and often a nucleus, the latter having passed into it from the parent cell. In its fully developed state it may contain starch, gum, resin and crystals, and its wall may be wrinkled. Sometimes a single parenchyma cell may produce several tyloses. Tyloses occur both in angiospermous and coniferous woods although are probably more common in the former. They are more abundant in heartwood than in sapwood and are of constant occurrence in some species. Excellent examples of tyloses can be seen in the rhizomes of *Asarum* and *Serpentaria* and in the stems of *Sassafras*, *Juglans*, the locust (*Robinia*), *Catalpa*, *Poplar*, *Sumachs*, *Oaks* and *White Pine*.

MICROSCOPIC CHARACTERISTICS OF ANGIOSPERMOUS AND CONIFEROUS WOODS.—The wood of Angiosperms is characterized by the presence of tracheæ (vessels) with various markings on their walls, particularly by small pits in the walls of some of

FIG. 143.—A, diagram of the course of vascular bundles in a palm stem (monocotyledon); 1m, 2m, 3m, bundles from the median portions of the leaves. B, diagram of vascular bundles in external view and in cross section of the dicotyledonous stem of *Cerastium*. The leaves are shown cut off at 1, 2, 2' and 3. (After Vines.)

the tracheæ, together with wood fibers, fiber tracheids,¹ wood parenchyma cells and medullary-rays.

The wood of Conifers is made up for the larger part of tracheids with bordered pits which latter are characterized in radial longitudinal section by the presence of two rings, one within the other. A single row of these is seen on the tracheid wall. Medullary-rays, frequently diagnostic for different species, and wood parenchyma cells (in larches and Douglas Fir, etc.), are also found.

¹ Forms intermediate between typical tracheids and wood fibers.

HISTOLOGY OF TYPICAL HERBACEOUS MONOCOTYL STEMS (ENDOGENOUS).—Passing from exterior toward the center the following structures are seen:

1. *Epidermis* whose cells are cutinized in their outer walls.
2. *Hypodermis*, generally collenchymatic in young, terminal portions, becoming later sclerenchymatous.

FIG. 144.

FIG. 144.—Photomicrograph of cross-section of very young cornstalk (a herbaceous monocotyl stem), where certain pith strands have just gone over into vascular bundles. For comparison with Fig. 145. (Stevens.)

FIG. 145.

FIG. 145.—Photomicrograph of cross-section of older cornstalk stem; *a*, epidermis; *b*, cortex and *c*, ground tissue of pith. (After Stevens.)

3. *Cortex*, of cortical parenchyma cells. Sometimes vascular bundles of the closed collateral or the concentric type occur in this region.

4. *Endodermis* or innermost layer of cortex and Pericycle (frequently appearing like and often indistinguishable from cortical parenchyma cells).

5. *Stele* or *Pith*, a large central zone of parenchyma matrix in which are found scattered fibro-vascular bundles of the closed collateral or more rarely of the concentric monocotyl (phlocentric) type. In this latter type, which is typical of old monocotyl stems and also seen in *Calamus*, *Convallaria*, etc., the xylem grows completely around the phloem so that phloem is found in the center and xylem without and surrounding it.

HISTOLOGY OF A TYPICAL MONOCOTYL RHIZOME.—The rhizome of *Convallaria majalis* (Lily-of-the-Valley) shows typical herbaceous monocotyl rhizome structure. A transverse section

(see Fig. 147) presents the following structural details, passing from periphery toward the center:

1. *Epidermis* of a layer of epidermal cells whose outer walls are covered with a waxy cutin deposit.

2. *Cortex* of about 20 rows of ordinary parenchyma cells, some of which contain more or less spheroidal starch grains, others raphides of calcium oxalate.

FIG. 146.—Transverse section through a representative portion of the monocotyledonous stem of the corn (*Zea Mays*) ($\times 50$). *ep*, epidermis; *h*, hypodermis containing lignified fibers; *co*, cortex (poorly defined in corn); *ph*, primary phloem and *x*, primary xylem of a closed collateral bundle surrounded by a sclerenchyma sheath of lignified fibers (*scl*); *par*, parenchyma of pith in which the bundles are scattered; *si*, sieve tissue; *tr*, tracheae; *a*, a ring of an annular trachea bordering a large air-space; *scl*, sclerenchyma fibers. The two large lateral tracheae in each bundle are pitted tracheae.

3. *Endodermis* of usually 2 layers of endodermal cells, the radial and inner walls of which are strongly indurated with lignin.

4. *Stele*, a broad central region consisting of a matrix of starch- and crystal-bearing parenchyma and through which region course closed collateral and concentric bundles.

The closed collateral bundles are arranged in an interrupted circle just beneath the endodermis. The concentric phlocentric bundles are scattered through the center of the stele.

Longitudinal-radial sections show that the tracheæ are of the spiral, scalariform, and pitted types.

HISTOLOGY OF A TYPICAL WOODY MONOCOTYL STEM.—The stem of the Greenbrier, a woody monocotyl, will here be considered. In transverse section passing from periphery toward the center the following structural details will be noted:

FIG. 147.—Transverse section of the rhizome of *Convallaria majalis*, a monocotyledon. $\times 40$. *ep*, epidermis; *co*, cortex; *end*, endodermis; *p*, phloem and *x*, xylem of a closed collateral fibrovascular bundle; *p'*, phloem and *x'*, xylem of a concentric bundle; *m*, pith, in which the concentric bundles are imbedded.

1. *Epidermis*, of a single layer of epidermal cells whose outer walls are strongly cutinized. *Cutin* is a wax-like substance which forms a protective coat to the epidermis, preventing the evaporation of water, the ingress of destructive parasites, and injury from insects.

2. A *cortex*, composed of about ten or twelve layers of thick-walled parenchyma cells, the outer two or three layers of which are termed hypodermis.

3. An *endodermis*, wavy in character and composed of endodermal cells whose brownish walls are strongly suberized.

4. A sclerenchymatous *cylinder sheath* composed of somewhat separated masses of sclerenchyma (pericyclic) fibers and undeveloped fibro-vascular bundles of the closed collateral type.

FIG. 148.—Photomicrograph of a representative portion of Greenbrier stem showing epidermis (*e.p.*), cortex (*c*), endodermis (*e.n.d.*), cylinder sheath (*c.s.*), sclerenchyma fibers of closed collateral bundle (*b*), fundamental parenchyma (*f.p.*), trachea (*t*). $\times 22$.

5. A *pith* or *central matrix* of strongly thickened, more or less lignified, parenchyma cells in which are scattered, irregularly, numerous closed collateral bundles. Small starch grains will be found in the parenchyma cells. Examine a representative bundle, and note the two very large tracheæ and several smaller ones in the xylem portion of the bundle which faces toward the center of the section. In the outer or phloem portion of the

bundle will be seen an area of soft, small-celled sieve tubes, companion cells and phloem parenchyma. The entire bundle is enclosed by a several layered ring of *sclerenchyma fibers*.

HISTOLOGY OF STEMS OF GRASSES (MONOCOTYLEDONOUS).—The stems of grasses are solid in their terminal and nodal regions;

FIG. 149.—Structure of a Grass Stem. Transverse section of the underground stem (rhizome) of the couch grass, *Agropyron repens*, $\times 35$. *ep*, epidermis; *h*, hypodermis; *co*, cortex; *b*, concentric bundle; *en*, endodermis; *scl*, sclerenchymatous ring of several rows of sclerenchyma fibers in which are imbedded an irregular, interrupted circle of undeveloped vascular bundles; *tr*, trachea and *ph*, a phloem strand of a closed collateral bundle; *cb*, closed collateral bundle; *m*, pith. Note the hollow space in the center due to the disintegration of the pith in centrifugal fashion.

but as we pass backward from the terminal internode into the older internodes the central pith is seen to disintegrate in centrifugal fashion, leaving a hollow cavity in the central region bordered by parenchyma cells. A distinctly differentiated endodermis may or may not be present. The vascular bundles may be entirely closed collateral, as in corn, but are frequently of two kinds, namely, small, phlocentric bundles in the cortex and

larger closed collateral bundles in the pith. The pith bundles may be more or less uniformly scattered, as in the stems of corn and bermuda grass, or arranged in one or more interrupted circles, as in *Triticum* species, etc. The epidermal cells usually possess cutinized outer walls and frequently all of their walls are lignified. The hypodermis contains lignified fibers, and there is usually present a sclerenchyma ring directly beneath the endodermis or the cortex in which bundles may be imbedded or to which they may be attached. A sclerenchyma sheath usually surrounds each bundle.

SECONDARY GROWTH IN MONOCOTYL STEMS.—Secondary thickening in stems of monocotyledons is exceedingly rare and the exception rather than the rule. It is seen, however, in some of the more woody and tree-like genera of the Lily family as *Aloë*, *Yucca*, *Cordyline* and *Dracaena* and in the tuberous *Dioscoreaceæ*. In these forms a cambium arises in the inner region of the cortex or the pericycle and lays down procambial strands and secondary parenchyma on its inner face. As it retreats outward, the procambial strands, by repeated division and differentiation, develop into secondary collateral (rarely phlocentric) bundles which lie imbedded in secondary parenchyma.

CHAPTER X

THE LEAF

The leaf is a usually flattened, rarely semi-centric, or centric-lateral expanse developed by the stem or by branches and in whose axil one or more branches arise.

FIG. 150.—Complete leaf of Slippery Elm (*Ulmus fulva*). *l*, lamina; *p*, petiole; *s*, stipules; *mr*, midrib; *v*, secondary vein; *st*, branch to which leaf is attached. The margin of this leaf is doubly serrate.

Leaves seldom develop buds over their surface or along their margin and in connection therewith roots. The capacity for bud development by the leaf is restricted to three families, viz.: *Crasulaceæ*, *Begoniaceæ* and *Gesneraceæ*.

THE COMPLETE LEAF.—The leaf when complete consists of three parts, *lamina*, *petiole*, and *stipules*. The LAMINA or BLADE

is the largest, most prominent and essential part. Other parts may be wanting. It consists of upper and lower surfaces provided with a layer of protective *epidermal cells* which may give rise to outgrowths in the form of hairs or papillæ. Either or both of these surfaces show openings called *stomata* (sing. *stoma*) which are protected by guard cells that regulate their opening and closing. The stomata serve for the exchange of carbon

FIG. 151.—Stereogram of leaf structure. Part of a veinlet is shown on the right. Intercellular spaces are shaded. (From Stevens.)

dioxide and oxygen and the escape of water vapor. Between the epidermal layers, in the region called the *mesophyll*, is a soft green tissue called *chlorenchyma* which is made up of leaf parenchyma cells containing *chloroplasts* and intercellular-air-spaces. Coursing through the mesophyll are the *veins* or *fibro-vascular bundles* which are in continuity with the fibro-vascular bundles of the stem. These veins branch and rebranch and so make up the frame-work of the lamina. The outer walls of the epidermal cells are provided with a cuticle which is generally thicker on the upper than on the lower surface. In most leaves the mesophyll is differentiated into upper *palisade parenchyma*, lower

spongy-parenchyma and *border parenchyma* regions, and stomata are only found in the lower epidermis.

The PETIOLE is the leaf stalk which supports the lamina in a position where it will receive the most light for photosynthesis. The STIPULES are leaf-like appendages appearing at the base of the petiole.

The leaf of the Slippery Elm (*Ulmus fulva*) affords a good example of a *Complete Leaf* (see Fig. 150).

FIG. 152.—Camera-lucida drawing of a tangential section of a leaf (section parallel to leaf surface) showing *f*, intercellular-air-spaces between the palisade parenchyma cells (*e*); *g*, border parenchyma; *h*, tracheal elements of a vein. (Stevens.)

Sometimes the lamina or blade is attached directly to the stem by its base and the leaf is then said to be *sessile*. If the petiole is present, the leaf is *petiolate*.

When leaf stipules are absent, the leaf is said to be *exstipulate*; when present, *stipulate*.

The petiole is seldom cylindrical in form, but usually channelled on the upper side, flattened, or compressed. The stipules are always in pairs and closely resemble the leaf in structure.

LEAF FUNCTIONS.—The most essential function of green plants is the conversion of inorganic into organic matter; this takes place ordinarily in the green parts, containing chlorophyll, and in these when exposed to sunlight. Foliage is an adaptation for increasing the extent of green surface.

The chief functions of a leaf are *photosynthesis*, *assimilation*, *respiration* and *transpiration*.

PHOTOSYNTHESIS is the process possessed by the cells of all green leaves or other green parts of plants of building up sugar, starch or other complex organic (carbon containing) substances by means of chlorophyll and sunlight. This process takes place in nature, only during sunlight but will also occur in artificial light. In this process CO_2 is taken into the leaf through the

FIG. 153.—The end of a very small veinlet as seen in a section of the spongy parenchyma cut parallel to the leaf surface. The walls of the conducting cells of the veinlet have spiral thickenings. The border parenchyma cells surrounding the veinlets serve to conduct the sugar from the mesophyll to the phloem cells of the vein. (From Mottier, after Strasburger, Macmillan Co.)

stomata and passes into the intercellular-air-spaces between the chlorenchyma cells. It diffuses through the cell walls of these into the protoplasts where it passes into solution and enters the chloroplastids along with H_2O from the soil. The chloroplasts contain *chlorophyll* which absorbs the light energy. This light energy is then used either by the chlorophyll or, as seems most likely, by the whole chloroplast to unite the CO_2 with the H_2O to form carbon compounds. These contain stored or potential energy. The kinetic energy of the sun's rays thus becomes stored as potential energy and O_2 is given off. The plant gains

in weight and so the process is constructive in character. In photosynthesis, sunlight is the power, the chloroplasts the working machinery or energy transformers, the chlorenchyma cells the factories, CO_2 and H_2O the raw materials and glucose, starch or oil the finished products, while oxygen is a waste

FIG. 154.—Diagram of photosynthesis. (After Atwood.)

product. The production of glucose may be represented by the equation:

ASSIMILATION is the process of converting food material into protoplasm. The chlorenchyma cells of the leaf synthesize proteins from the carbohydrates. In this process molecules of glucose derivatives containing CHO are united with nitrogen absorbed from the soil as nitrates to form *amino acids*. The amino acids combine to form *polypeptides* which are later built into *simple proteins*. The simple proteins are joined with sulfur or phosphorous or both to form *complex proteins*. The latter enter into the formation of protoplasm.

RESPIRATION is the process which takes place in all living plant and animal cells whereby complex carbon compounds are oxidized with an accompanying release of energy. In higher green plants oxygen is taken in through the stomata of leaves, lenticels of stems, and root hairs. It passes into the intercellular-air-spaces which communicate with each other

throughout the entire plant and diffuses through the various cell-walls of cells, in contact with these, and passes into the interior of the cells. Here it oxidizes some of the carbohydrates breaking these down into carbon dioxide and watery vapor.

FIG. 155.—Demonstration of the liberation of oxygen during photosynthesis. A, release of bubbles of oxygen gas from the freshly cut end of the stem of *Elodea* in sunlight. B, collection of the gas, mainly oxygen, from *Elodea* in sunlight during photosynthesis. If an aquatic plant be placed in a glass vessel of water under an inverted funnel over whose stem is suspended an inverted test tube filled with water, the gas given off by the plant will collect in the upper end of the tube and displace the water. Upon removing the tube and thrusting into it a glowing splinter, the latter will increase in brightness, indicating the presence of oxygen. (After Mottier.)

These diffuse out of the cells into the intercellular-air-spaces and pass out of the plant through the same channels as oxygen entered. Respiration goes on both in light and darkness, organic compounds are broken down, O is absorbed and CO₂ set free, and potential energy is transformed into kinetic energy. The plant loses weight and, accordingly, this process is destructive in character.

TRANSPIRATION is the action of giving off watery vapor. The greater portion of the crude sap consisting very largely of water is conveyed upward as a *transpiration stream* through the tracheæ and tracheids of the roots and stems into those of the leaves. The latter pervade the soft, green leaf parenchyma and end in proximity to air spaces between the green cells. A portion of the crude sap diffuses into the leaf parenchyma cells and is utilized in the nutritive processes such as photosynthesis, digestion and assimilation occurring there. The remainder, which normally constitutes the larger part, passes through the intercellular-air-spaces and out of the leaves as watery vapor. About 90 per cent. of the water thus transpired escapes through the stomata, the remainder through the epidermal surface. It has been estimated that a large oak tree

transpires about 180 gallons of water per day and that during a growing season of five months it would give off water enough to cover the ground shaded by it to the depth of 20 feet.

TYPES OF LEAVES DEVELOPED IN ANGIOSPERMS.—These may be tabulated as follows:

1. Cotyledons (the primitive or seed leaves).
2. Scale leaves.
3. Foliage leaves.

FIG. 156.—Semi-diagrammatic cross section of a DORSIVENTRAL LEAF of a mesophyte showing by arrows how the water passes from the tracheal elements of a vein into the border parenchyma cells, and thence into the palisade and spongy parenchyma, from which it evaporates into the intercellular spaces and passes from the leaf through the stomata. *a*, upper epidermis; *b*, lower epidermis; *c*, palisade parenchyma; *g*, spongy parenchyma; *d*, border parenchyma; *e*, tracheal elements; and the stippled cells below *e*, the phloem cells. (After Stevens.)

4. Bract leaves; (*a*) primary at base of inflorescence; (*b*) bracteolar leaves (bracteoles) at the base of individual flowers.

5. Sepals.

6. Petals.

7. Microsporophylls (stamens).

8. Megasporophylls (carpels).

COTYLEDONS.—Cotyledons are the first leaves to appear upon the ascending axis and are single in Monocotyledons, double in Dicotyledons. Occasionally, as in certain Maples, there may be three cotyledons shown. This is due to a splitting of one of the cotyledons. There exist no true cases of polycotyledony (development of many cotyledons) among Angiosperms, as in

Gymnosperms. In Monocotyledons the single cotyledon is a terminal structure and truly axial in relation to the hypocotyl and radicle. From a primitively Monocotyl-like ancestry Dicotyledons develop a second cotyledon on the epicotyledonary node. Later, by a suppression of the second node the second cotyledon is brought to the level of the first.

SCALE LEAVES.—Scale leaves are reduced foliage leaves. They are found on certain rhizomes, above ground stems, such as Dodder, Indian Pipe, etc., on bulbs, and forming the protective scales of scaly buds.

FIG. 157.—The growing end of the stem of the aquatic plant, *Elodea canadensis*, showing the origin of leaves. The larger over-lapping leaves are omitted. A, surface view; B, in section. (Courtesy of Standard Scientific Corp., formerly Scientific Equipment Dept. of Kny-Scheerer Corp.)

FOLIAGE LEAVES.—These are the common green leaves so familiar to all.

BRACT LEAVES are modified leaves appearing on inflorescence axes.

SEPALS, PETALS, MICROSPOROPHYLLS and MEGASPOROPHYLLS are floral leaves and will be treated at length under the subject of the flower.

ORIGIN AND DEVELOPMENT OF LEAVES.—Leaves arise around the growing apex region of a stem or branch as lateral outgrowths, each consisting at first of a mass of cells called the *primordial leaf*. Through continued cell-division and differentiation of these cells

in time the mature leaf is developed. The primordial leaf is formed by a portion of the dermatogen of the growing stem apex, which becomes epidermis, a portion of the periblem, producing mesophyll or leaf parenchyma which grows into this, and a part of the plerome, which becomes vascular tissue within the mesophyll.

In the sub-divisions of cells around the growing stem-apex, the primordial leaves (*primordia*) do not arise exactly at the same time. There is a tendency toward spiral arrangement.

PHYLLOTAXY.—Phyllotaxy is the study of *leaf arrangement* upon the stem or branch, and this may be either alternate, opposite whorled, or verticillate, or fascicled. It is a general law in the arrangement of leaves and of all other plant appendages that they are spirally disposed, or on a line which winds around the axis like the thread of a screw. The spiral line is formed by the union of two motions, the circular and the longitudinal, and its most common modification is the circle.

Foliage leaves are usually arranged in separated fashion along the stem, one occurring at each node. This is the form of spiral arrangement called the *alternate* arrangement, as shown by most plants.

The alternate arrangement is said to be two-ranked, when the third leaf is over the first, as in all Grasses; three-ranked, when the fourth is over the first. Example: Sedges. The five-ranked arrangement is the most common, and in this the sixth leaf is directly over the first (counting the leaf from which the start is made), two turns being made around the stem to reach it. Examples: Cherry, Apple, Peach, Oak and Willow, etc. As the distance between any two leaves in this arrangement is two-fifths of the circumference of the stem or 144° , the five-ranked arrangement is expressed by the fraction $\frac{2}{5}$. In the eight-ranked arrangement, the ninth leaf stands over the first, and three turns are required to reach it, hence the fraction $\frac{3}{8}$ expresses it. Of the series of fractions thus obtained, the numerator represents the number of turns to complete a cycle, or to reach the leaf which is directly over the first; the denominator, the number of perpendicular rows on the stem, or the number of leaves, counting along the spiral, from any one to the one directly above it.

The one which is chosen as the starting point is marked and taken as 0.

Other forms of leaf arrangement are:

Opposite, when a pair of leaves is developed at each node, on opposite sides of the stem. Examples: Mints, Lilac.

Decussate, when the leaves are arranged in pairs successively along the stem, at right angles to each other. Example: Thoroughwort.

Whorled or *Verticillate*, when three or more form a circle about the stem. Examples: Canada Lily and Culver's Root.

Fascicled or *Tufted*, when a cluster of leaves is borne from a single node, as in the Larch and Pine.

VERNATION.—*Prefoliation* or *Vernation* relates to the way in which leaves are disposed in the bud. A study of the individual

FIG. 158.—Three principal types of vernation. (Robbins.)

leaf enables us to distinguish the following forms. When the apex is bent inward toward the base, as in the leaf of the Tulip Tree, it is said to be *inflexed* or *reclinate vernation*; if doubled on the midrib so that the two halves are brought together, as in the Oak or

Peach, it is *conduplicate*; when rolled inward from one margin to the other, as in the Wild Cherry, it is *convolute*; when rolled from apex to base, as in Ferns, it is *circinate*; when folded or plaited, like a fan, as in Ricinus, Maples, Aralias, etc., it is *plicate*; if rolled inward from each margin toward the midrib on the upper side, as the leaves of the Apple or Violet, *involute*; when rolled outward from each margin as Dock or Willow leaves, *revolute*. The inner surface is always that which will form the upper surface when expanded.

LEAF VENATION.—This refers to the arrangement of the veins within the leaf. It is dependent upon the course pursued by the fibrovascular bundles after they enter the leaf blade.

The central vein in the leaf is termed the *midrib* or *primary*. The branches coming off of the midrib are called *secondary veins* or *secondaries*; those arising from secondary veins are called *tertiaries*.

Strong and thick veins are called *ribs* and less prominent ones, especially if lateral to the midrib, *nerves*.

FURCATE- or FORKED-VENATION is characteristic of leaves of the Ferns.

FIG. 159.—Leaf venation. 1. Furcate (*Botrychium*, a fern). 2. Parallel (*Convallaria*). 3. Pinnately-parallel (*Musa*). 4. Pinnate-reticulate (*Castanea*). 5. Palmate-reticulate (*Acer*). 6. Pinnate-reticulate with marginal veins anastomosing (*Eucalyptus*). (Drawing by *Canis*.)

PARALLEL VENATION is typical of the leaves of *Monocotyledons*, as Palms, Lilies, Grasses, etc. In this type of veining the veins visible to the unaided eye extend in somewhat parallel fashion through the leaf, converging near the summit. In pinnately-parallel veined leaves, as the Banana and Calla Lily, parallel veins extend from midrib to margin.

RETICULATE or NETTED VEINS characterize the leaves of *Dicotyledons*, as Matico and the Poplar or Oak. The primary veins in these are generally pinnate or palmate while the secondary ones and their branches are arranged in netted fashion.

PINNATELY-VEINED or FEATHERED-VEINED leaves consist of a midrib with lateral veins, extending from midrib to margin at frequent intervals and in a regular manner. Examples: Beech and Chestnut leaves.

PALMATELY VEINED leaves consist of a number of veins of nearly the same size, radiating from petiole to margin. Examples: Maple and Castor Oil leaves.

In palmately-veined leaves, the central vein is termed the *midrib* or *middle primary* and those veins lateral to it, the *lateral primaries*.

Most pinnately-veined leaves and palmately-veined leaves show net-veins combined with the basic pinnate or palmate type and so are said to exhibit *pinnate-reticulate* or *palmate-reticulate* venation respectively.

Veins are said to be ANASTOMOSING when they subdivide and join each other, as the veins near the margin of *Eucalyptus* leaves; TESSELLATED, when they form square meshes, as in Matico leaves; and IMPRESSED when sunken below the surface.

LEAF INSERTION.—The point of attachment of the leaf to the stem is called the insertion. A leaf is:

Radical, when inserted upon an underground stem or a root.

Cauline, when upon an aerial stem.

Ramal, when attached directly to a branch.

When the base of a sessile leaf is extended completely around the stem it is *perfoliate*, the stem appearing to pass through the blade. Example: *Uvularia perfoliata* or Mealy Bellwort.

When a sessile leaf surrounds the stem more or less at the base, it is called *clasping* or *amplexicaul*. Example: Poppy (*Papaver somniferum*).

When the bases of two opposite leaves are so united as to form one piece, the leaf is called *connate-perfoliate*, as *Eupatorium perfoliatum* or Boneset.

Leaves are called *equitant* when they are all radical and successively folded on each other toward their bases, as in *Iris* spp.

THE FORMS OF LEAVES.—SIMPLE LEAVES are those having a single blade, either sessile or petiolate.

COMPOUND LEAVES are those whose blades are divided into two or more distinct subdivisions called *leaflets*. The leaflets may

FIG. 160.—Leaf outline: Linear (1); linear-lanceolate (2); lanceolate (3); ovate (4); elliptical (5); orbicular (6); oblanceolate (7); cuneate (8); spatulate (9); falcate (10); ensiform (11). (Drawing by Canis.)

be either sessile or petiolate. If the compound leaf has leaflets arising from the midrib, it is called *Pinnately Compound*. The midrib of a compound leaf is sometimes called the *rachis*. If the leaflets of a compound leaf come off of the summit of the

petiole, the compound leaf is termed **PALMATELY-** or **DIGITATELY-COMPOUND**. The leaf stalk of a leaflet is called the *petiolule*.

A compound leaf develops from a single leaf bud as does a simple leaf. Even though a group of small leaves may appear as

FIG. 161.—Leaf bases (12–17); leaf apices (18–26); compound leaves (27–31). Cordate (12); auriculate (13); connate-perfoliate (14); sagittate (15); hastate (16); peltate (17). Acuminate (18); acute (19); obtusate (20); truncate (21); retuse (22); emarginate (23); cuspidate (24); mucronate (25); aristate (26). Imparipinnate (27); paripinnate (28); bi-pinnate (29); decompound (30); palmately 5-foliate (31).

a compound leaf, they can be distinguished as simple leaves by the presence of a bud in each of their axils.

FORMS OF LAMINA.—The lamina of simple leaves and the leaflets of compound leaves are described as to general outline, apex, base, marginal indentations, surface and texture.

(a) GENERAL OUTLINE (form viewed as a whole without regard to indentations of margin).

Linear, long, narrow, ribbon-like of nearly equal breadth throughout, as *Linaria*.

Filiform, thread-like, as the *Asparagus* and one of the *Sundews*.

Acerose or *acicular*, needle-like, as in the *Pines*.

Subulate, awl-shaped, as in the *Arbor Vitæ*.

Ovate, egg-shaped, with its length not more than two or three times its breadth, as *Boldo*.

Obovate, reversely ovate as some leaves of *Short Buchu*.

Rhomboidal, angularly-ovate, as the *Salt Bush*.

Lanceolate, like the linear with the exception that the broadest part is a little below the center. Example: *Long Buchu*.

Elliptical, somewhat longer than wide, with rounded ends and sides. Example: *Leaf of Pear*.

Oblong, when longer than broad, margins parallel. Example *Matico*.

Equilateral, margin of the same length on both sides.

Inequilateral, margin longer on one side than the other, as the *Hamamelis*, *Elm* and *Linden*.

Orbicular, the blade circular in shape. Example: *Nasturtium*.

Reniform, or kidney-shaped, as the leaves of *Canada Snakeroot*.

Oblanceolate, reversely lanceolate. Example: *Chimaphila*.

Spatulate, like a spatula, with narrow base and broad rounded apex. Example: Some leaves of *Uva Ursi*.

Ensiform, when shaped like a sword. Example: *Calamus*.

Falcate, scythe- or sickle-shaped, as leaves on older branches of the *Eucalyptus*.

Deltoid, when the shape of the Greek letter Δ , as *Chenopodium*.

(b) APEX OF LEAF.—*Acute*, when the margins form an acute angle at the tip of the leaf giving the tip the appearance of a sharp point. Examples: *Eriodictyon*, *Digitalis*.

Acuminate, when the point is longer and more tapering than the acute. Examples: *Pellitory*, *Coffee*.

Tapering, if the summit of the leaf be narrowed very gradually to a point, as in leaves of many grasses.

Attenuate, if the summit is still more drawn out than the tapering type.

Obtuse, blunt or rounded. Example: Long Buchu.

Rounded, when extremely obtuse, as in some leaves of Foxglove.

Truncate, abruptly obtuse, as if cut square off. Example: Melilotus leaflets.

Mucronate, terminating in a short, soft point. Example: some Senna leaflets.

Cuspidate, like the last, except that the point is long and rigid.

Aristate, with the apex terminating in a bristle.

Emarginate, notched. Example: Pilocarpus.

Retuse, with a broad, shallow sinus at the apex. Example: Petal of *Rose gallica*.

Obcordate, inversely heart-shaped. Example: Oxalis.

(c) BASE OF LEAF.—*Cordate*, heart-shaped. Examples: Catnep, Violet and Coltsfoot leaves.

Reniform, kidney-shaped. Examples: Ground Ivy, Asarum.

Hastate, or halberd-shaped, when the lobes point outward from the petiole. Example: *Aristolochia Serpentaria*.

Auriculate, having ear-like appendages at the base. Example: Philodendron.

Sagittate, arrow-shaped. Examples: Bindweed and Sagittaria.

Cuneate, wedge-shaped. Examples: Short Buchu and Uva Ursi.

Peltate, or shield-shaped, having the petiole inserted at the center of the lower surface of the lamina. Example: Podophyllum.

Oblique, when the base of the lamina is attached lower on one side of the midrib than the other, as in the Stramonium.

The bases of many leaves extend downward along the stem beyond the main point of insertion when they are called *decurrent*. If the base of the lamina is prolonged downward along the sides of the petiole to which it is joined, the leaf is said to have a *winged petiole*, as in Digitalis.

The terms *acute*, *tapering*, *attenuate*, *truncate*, *obtuse* and *rounded* which have been defined under "Apex of Leaf" are also applied to bases of like character.

(d) MARGIN OF LEAF.—*Entire*, when the margin is an even line. Example: Belladonna.

Revolute, margin rolled backward, as in Thyme.

Involute or *incurved*, margin rolled inward.

Serrate, with sharp teeth which incline forward like the teeth of a hand-saw. Examples: Peppermint, Yerba Santa, Buchu.

Serrulate, when finely saw-toothed, as in Rose leaflets.

Dentate, or *toothed*, with outwardly projecting teeth. Example: Damiana.

Denticulate, when finely toothed.

Crenate, or *Scalloped*, similar to the preceding forms, but with the teeth much rounded. Examples: Digitalis, Catnip.

Crenulate, when finely scalloped, as in Sage.

In the serrate, dentate and crenate forms, when the teeth show divisions into smaller or secondary teeth, the word "*doubly*" is used as a prefix to the term expressing the character of the primary tooth, thus, *Doubly-serrate*, etc.

Spinose, when the teeth are extended as spines, as in the American Holly.

Ciliate, when the margin is fringed with hairs, as in Drosera.

The *sinuses* are the indentations of the margin. If sufficiently deep, they separate the leaf into *segments* or *lobes*.

Repand, or *Undulate*, margin—a wavy line. Example: Hamamelis.

Sinuate, when the margin is more distinctly sinuous than the last. (*Stramonium*.)

Incised, cut by sharp, irregular incisions. Example: Hawthorn.

Runcinate, the peculiar form of pinnately-incised leaf observed in the Dandelion and some other *Compositæ* in which the teeth are recurved.

A *Lobed* leaf is one in which the indentations extend toward the mid-rib, or the apex of the petiole, the segments or sinuses, or both, being rounded. Example: Sassafras.

Cleft is the same as lobed, except that the sinuses are deeper, and commonly acute. Example: Dandelion.

A *Parted* leaf is one in which the incisions extend nearly to the mid-rib or the petiole. Example: *Geranium maculatum*.

In the *Divided* leaf the incisions extend to the mid-rib, or the petiole, but the segments are not stalked nor are they in the form of leaflets. Example: Watercress.

If the venation is pinnate, the preceding forms may be described as pinnately-cleft (incised), -lobed, -parted, or -divided. If the venation is (radiate) palmate, then the terms palmately-lobed, -incised, etc., are employed.

FIG. 162.—Leaf margins: Pinnately-lobed (32); pinnately-cleft (33); pinnately-parted (34); pinnately-divided (35); palmately tri-lobed (36); palmately tri-cleft (37); palmately 3-parted (38); palmately 3-divided (39); crenate (40); serrate (41); dentate (42); repand or undulate (43); sinuate-dentate (44).

Pinnatifid is employed by some authors for describing a pinnately-cleft leaf, *pinnatipartite* for a pinnately-parted one, and *pinnatisect* for one that is pinnately-divided. Likewise *palmatifid*, *palmatipartite* and *palmatisect* are sometimes employed in place

of palmately-cleft, palmately-parted and palmately-divided respectively.

FIG. 163.—Forms of Leaves. Runcinate leaf of Dandelion (1); lyrate leaf of Turnip (2); interruptedly-pinnate leaf of Potato (3); palmate leaf of Horsechestnut (4); inequilateral leaf of Witch Hazel (5); pedate leaf of Hellebore (6). Modified leaves of insectivorous plants 7–9. Leaf of a Sundew bearing tentacles (7); leaf of Venus Trap (8); pitched leaf of *Sarracenia purpurea* (9); leaf of Heartsease showing free lateral stipules (10).

A *Pedate* leaf is one which is palmately-parted or divided but which has its lateral lobes in turn divided in more or less linear fashion.

When the lobes or divisions of a pinnately-divided, -parted or cleft leaf are alternately large and small, the leaf is *interruptedly pinnate*, as the Potato leaf. When the terminal lobe is the largest, and the remaining ones diminish in size toward the base the form is known as *lyrate*, illustrated in the leaf of the Turnip.

COMPOUND LEAVES.—The transition from Simple to Compound Leaves is a very gradual one, so that in many instances it is difficult to determine whether a given form is to be regarded as simple or compound. The number and arrangement of the parts of a compound leaf correspond with the mode of venation, and the same descriptive terms are applied to outline, margin, etc., as in simple leaves.

Leaves are either *pinnately* or *palmately compounded*. The term *pinnate* is frequently given to the former while that of *palmate* is often assigned to the latter. They are said to be abruptly pinnate or *paripinnate* when the leaf is terminated by a pair of leaflets; odd pinnate or *imparipinnate* when it terminates with a single leaflet.

Palmately compound leaves have the leaflets attached to the apex of the petiole. When these are two in number the leaf is *bifoliate*, or *binate*; if three in number, *trifoliate*, or *ternate*, as in *Menyanthes*; when four in number, *quadrifoliate*, as in four-leaved clover, etc. If each of the leaflets of a palmately compound leaf divides into three, the leaf is called *biterminate*; if this form again divides a *triternate* leaf results. Beyond this point the leaf is known as *decompound*. In the case of pinnately-compound leaves, when division progresses so as to separate what would be a leaflet into two or more, the leaf becomes *bipinnate*, as the compound leaves of *Acacia Senegal* or on the new wood of the Honey Locust (*Gleditschia*); if these become again divided, as in many *Acacia* species, the leaf is termed *tripinnate*. Beyond this point the leaf is known as *decompound*. Examples of pinnately-decompound leaves are seen in *Cimicifuga* and Parsley.

LEAF MODIFICATIONS.—The leaves of a number of plants have become modified in one or more parts for the purpose of carrying out special functions coincident with habits acquired by the plants possessing them. Thus, in the common Garden Pea, the

upper leaflets are transformed into tendrils for climbing purposes; in the Barberry some of the leaves have become transformed into spines for the defense of the plant from browsing animals; in Squill and Garlic the leaves of the bulbs have become succulent scales for food storage; in the Bladderworts certain submerged leaves have become modified as bladder-like traps for capturing crustaceans; in Sundews, Venus Fly Trap and the Pitcher Plants (*Heliamphora*, *Darlingtonia*, *Sarracenia*, *Nepenthes* and *Cephalotus*) the leaves are variously modified as traps for alluring, capturing and devouring insects. (See Carnivorous Plants in Chapter on Ecology.)

LEAF TEXTURE.—Leaves are described as:

Membranous, when thin and pliable, as Coca.

Succulent, when thick and fleshy, as Aloe, and Live Forever.

Coriaceous, when thick and leathery, as Eucalyptus, Uva Ursi and Magnolia.

LEAF COLOR.—*Petaloid*, when of some brilliant color different from the usual green, as the Coleus and Begonia, and other plants which are prized for the beauty of their foliage rather than their blossoms.

LEAF SURFACE.—Any plant surface is:

Glabrous, when perfectly smooth and free from hairs or protuberances. Example: Tulip.

Glaucous, when covered with bloom, as the Cabbage leaf.

Punctate, when dotted as if by punctures, as in the leaf of the Wax Myrtle.

Pellucid-punctate, when dotted with projections formed by subjacent oil glands, as the leaves of the many members of the Orange family.

Papillose, when covered with minute, teat-shaped projections, as the lower surface of Coca leaves.

Scabrous leaves have a rough surface with minute, elevated, hard papillæ or warts, as those of Slippery-elm and many composites.

Pubescent, covered with short, soft hairs. Example: Digitalis.

Appressed-hairy, when the hairs are curved so as to appear somewhat pressed against the surface, as in Senna leaflets.

Villose, covered with long and shaggy hairs. Example: Forget-me-not.

Sericeous, when covered with hairs which give a silky appearance. Example: Silverleaf.

Hispid, when covered with short, stiff hairs. Example: Borage.

Strigose, when covered with appressed, stiff and sharp hairs.

Tomentose, densely pubescent and felt-like, as the Mullein leaf.

Spinose, beset with spines, as in the Thistle.

Rugose, when roughened with closely set wrinkles. Example: Sage.

Verrucose, covered with protuberances or warts, as the calyx of *Chenopodium*.

Tuberculate, when covered with minute wart-like projections.

Bullate, when blister-like projections appear between the veins.

DURATION OF LEAVES.—Leaves vary as to their period of duration. They are: *Persistent*, or *evergreen*, if they remain green on the tree for a year or more.

Deciduous, if unfolding in spring and falling in autumn.

Caducous, or *fugacious*, if falling early in the season.

PARTS OF TYPICAL LEAF.—The parts of a typical leaf are *petiole* or leaf stalk, *lamina* or blade, and *stipules*.

THE PETIOLE

The PETIOLE IN MONOCOTYLEDONS is usually a broadened, sheathing basal structure which connects the lamina to the stem. In the Grasses and Bananas it forms a long tubular sheath which surrounds the stem. At the point of union of the sheathing petiole and the lamina of a grass is a membranous or leathery ring or flap called the **LIGULE**. The ligule is not a part of the petiole but is a formation due to the upgrowth and union of a pair of stipules. Into the Grass petiole a set of closed collateral vascular bundles of the stem extend, these showing xylem uppermost and phloem beneath. In the Palms, Aroids, Yams, Bananas, the petiole in part or throughout may be much thickened, strengthened and developed as a semi-cylindric or cylindric structure frequently showing, as in Palms, generally, two sets of

bundles. In all of these the petiole shows distinct closed collateral to concentric bundles embedded in parenchyma which in turn is surrounded by epidermis. In the Monocotyl genus *Maranta* of the Arrowroot family, a special swelling is found at the apex of the petiole which is termed a *Pulvinus*.

IN DICOTYLEDONS, the PETIOLE attains its most perfect development and here usually shows differentiation into a short basal part called the PULVINUS or *leaf cushion* and a longer part, the STALK portion. (See Fig. 18.)

The pulvinus is sensitive to environal stimuli and in some groups, as *Oxalidaceæ* and *Leguminosæ*, a gradual increase in sensitivity up to a perfect response can be traced. Moreover, in these, if we start with the simpler less sensitive pulvini and pass by stages to the most complex, we note that a special substance known as the *aggregation body* develops in the pulvinar cortex cells and that this substance undergoes rapid molecular change on stimulation of the leaf.

The stalk portion of the petiole in Dicotyledons is usually plano-convex or nearly to quite circular in outline; rarely in certain families does it resemble Monocotyledons in becoming abruptly or gradually thinned or flattened or widened out so as to sheath round the stem. The most striking example of this is seen in the Parsley family (*Umbelliferae*) where the flattened sheathing leaf stalk is known as the PERICLADIUM. Such a structure is not peculiar to the *Umbelliferae*, for in many members of the Crowfoot family (*Ranunculaceæ*), etc., a similar sheathing development is observed. The stalk may bear the laminar tissue on its extremity. This is most commonly the rule, but when the

FIG. 164.—Barley, (a member of the Grass Family). A, portion of leaf at juncture of blade and decurrent sheath; B, stem cut in median longitudinal section. $\times 2\frac{1}{2}$. (Robbins.)

plant is exposed to xerophytic conditions, as the Acacias of Africa and Australia, the stalk, instead of being cylindric or sub-cylindric, becomes flattened from side to side, until there is produced a bifacial, vertically placed petiole, with a large green surface that wholly takes the place of the lamina. Such a structure is known as a PHYLLODE or PHYLLODIUM. (Fig. 165.)

FIG. 165.—A Phyllode, or Blade-like Petiole, from *Acacia melanoxylon*, and its Cross-Section. Drawing by H. McCarthy. (After Stanford, *General and Economic Botany*, D. Appleton-Century Co. publishers.)

The INTERNAL STRUCTURE OF THE PETIOLE in primitive types of Dicotyledons resembles that seen in Monocotyledons except that the bundles are more condensed side by side. In these the petiole is somewhat flattened and shows an external epidermis, a flattened cortex and a single row of vascular bundles, each with xylem uppermost and phloem below. From this we can pass to another group of these plants in which the bundles form three-fourths of a circle and in which the upper bundles show incurving

orientation, to still another in which, as in *Nepenthes*, all of the bundles form nearly a cylinder. Finally, in *Ficus*, *Geranium*, *Podophyllum* and other plants showing completely formed cylindrical petioles, the bundles form a continuous ring enclosing pith and surrounded by cortex and epidermis, as in Dicotyl stems.

STIPULES

STIPULES are lateral leafy or membranous outgrowths from the base of the petiole at its junction with the stem. They may be divided into two groups, viz.: *lateral* and *axillary*. The *lateral group* includes four types, namely, free lateral, lateral adnate, lateral connate and lateral interpetiolar.

FREE LATERAL STIPULES are seen in *Leguminosæ*, *Rosaceæ*, *Beeches*, etc. They are free on either side of the petiole and supplied by vascular tissue from the petiolar bundle mass. In appearance and duration they may be either green, foliaceous and persistent or membranous to leathery, scale-like and caducous. *Caducous scaly stipules* only function as bud scales, forming a protective covering to the bud through the winter and fall in spring as the buds expand. Cf. Fig. 163 (10).

LATERAL ADNATE STIPULES are such as fuse with and are carried up with the petiole as wing-like appendages. This type is seen in the Roses, in Clovers, etc.

LATERAL CONNATE STIPULES are such as join and run up with the petiole to form a structure which is called a *LIGULE*. This structure is common to the *Gramineæ* or Grass family. (See Fig. 164.)

LATERAL INTERPETIOLAR STIPULES are common to many species of the *Rubiaceæ*. This family has opposite leaves and in some of the genera, as *Cinchona*, one stipule from each leaf on either side of the stem unites by its margin with the margin of the other stipule to form an interpetiolar stipule. In this way a stipule is formed on either side of the stem between the opposite leaves.

The AXILLARY GROUP represents stipules which stand in the axil of the leaf with the stem. Such may be FREE AXILLARY STIPULES, arising as distinct processes, or CONNATE, when the two

stipules unite at their margins, grow up and sheathe the stem, forming a legging-like sheath, as in many species of the *Polygonaceæ* such as Buckwheat, Rhubarb, Yellow Dock, Knot Weeds, etc. The sheath formed is called an OCHREA.

MODIFIED STIPULES.—In some plants such as the Locust and several other trees and shrubs of the Legume family, the stipules become modified for defensive purposes as spines or prickles. In the Sarsaparilla-yielding plants and other species of the genus *Smilax* they undergo modification into tendrils which are useful in climbing. (See Fig. 366.)

THE LAMINA

This, as was previously indicated, represents an expansion of the tissues of the petiole, but in sessile leaves is directly attached to the stem and so is a direct stem outgrowth.

MODE OF DEVELOPMENT OF THE LAMINA OF LEAVES.—The lamina of leaves develops in one of six ways.

1. Normal or Dorsiventral.
2. Convergent.
3. Centric.
4. Bifacial.
5. Reversed.
6. Ob-dorsi-ventral.

The first four will only be considered.

A. DORSIVENTRAL (the commonest).

(a) DORSIVENTRAL UMBROPHYTIC.—Flattened from above downward. Plants with such leaf blades tend to grow in the shade.

(b) DORSIVENTRAL MESOPHYTIC.—Similar to the former, but plants usually grow directly in the open and exposed to sunlight and winds.

(c) DORSIVENTRAL XEROPHYTIC.—Similar to former, but plants not only grow exposed, but exposed to hot desert conditions or to cold vigorous conditions.

(d) DORSIVENTRAL HYDROPHYTIC.—All transitions occur between typical mesophytic forms to those of marshy places, to swamps and borders of streams and finally with leaves wholly emersed, the last a completely hydrophytic type.

GROSS STRUCTURE AND HISTOLOGY OF DIFFERENT TYPES OF DORSIVENTRAL LEAF BLADES.—1. DORSIVENTRAL UMBROPHYTIC. Characterized by leaves mostly undivided and having the largest and most continuous leaf expanse. Usually the deepest green leaves we have, to enable the leaves to absorb scattered and reduced rays that pass in through high trees and shrub overhead.

FIG. 166.—Transverse section of the dorsiventral leaf blade of the Lilac, a mesophytic shrub. *s*, stoma. (Mottier.)

Their texture is usually thin and soft. In microscopic structure they are covered with a cutinized epidermis which has all the stomata on the lower surface. The chlorenchyma is fairly spongy, the spongy parenchyma having decided intercellular spaces. Examples: Hydrastis, Asarum, Maranta.

2. DORSIVENTRAL MESOPHYTIC.—In this type of dorsiventral leaf there is a tendency to subdivision, either to slight or

moderate lobing, seldom to complete subdivision in pinnate or tripinnate fashion. Examples: Belladonna, Hyoscyamus, Dandelion, Grasses, etc. In microscopic structure, they consist of an upper and lower epidermis, the upper epidermis being the thicker of the two, the lower epidermis having wavy vertical walls. The stomata are wholly or are mainly on the lower epidermis. In mesophytic trees and shrubs they are restricted generally to the lower surface while in herbs of mesophytes they may occur on both surfaces although mostly on the lower surface. In mesophytic dicotyledons, the palisade parenchyma is toward the upper epidermis and consists of one to three layers of cells. The spongy parenchyma is toward the lower and consists of irregular, loosely arranged cells with prominent intercellular-air-spaces. In a number of mesophytic monocotyledons like the grasses and sedges the chlorenchyma is devoid of palisade cells, being compact and uniform.

3. DORSIVENTRAL XEROPHYTIC.—Leaves characterized by a thick upper and lower cuticle and by having their numerous, small stomata restricted to the lower surface or present more or less equally on both surfaces, where they are sunken in depressions or pits. They may be either firm, leathery, tough, or fibrous, or may become swollen up in their mesophyll, chiefly in their spongy parenchyma cells, and store considerable mucilage. The object of mucilage is to retain water. Very frequently they develop a water-storage tissue above the mesophyll. Their palisade tissue is usually well developed, occurring in the cacti in many rows. Examples: Boldo, Eriodictyon, *Ficus*.

4. DORSIVENTRAL HYDROPHYTIC.—All gradations are seen. In pond plants, such as the Water Lily, the leaves have long split petioles which bring the blade up to the surface of the water. The stomata are entirely on the upper surface. In *Ranunculus*, the lower leaves are cut up into filiform segments. These are devoid of stomata. Their mesophyll is soft, open, and spongy showing large air-spaces. The epidermis is quite thin. The upper leaves are floating, trilobed, and have stomata only on their upper surface. In *Utricularia*, some of the filiform submerged leaves are modified into bladders which trap insect larvæ and smaller Crustaceæ (see Fig. 481).

B. CONVERGENT.—This type is mostly seen in monocotyledons. In *Phormium tenax* or New Zealand Flax the base of the blade is sheathing; it then converges and opens out above. In the various species of *Iris* the petiole is sheathing, the sides of the upper part being fused and functioning as a lamina.

FIG. 167.—Transverse section through portion of dorsiventral leaf blade of horehound (*Marrubium vulgare*). Upper epidermis devoid of stomata (*up. ep.*); lower epidermis which possesses stomata (*l. ep.*); palisade parenchyma (*pal.*); spongy parenchyma (*sp. p.*); xylem (*x*) and phloem (*ph*) regions of fibrovascular tissue of stronger vein; long-pointed non-glandular trichome (*t*); branched trichomes (*ft*¹, *ft*², *ft*³); several types of glandular trichomes (*gt*¹, *gt*², *gt*³, *gt*⁴).

C. CENTRIC.—This type of lamina is semi-centric to centric in shape. It is nearly always associated with Xerophytes. Centric laminae are produced gradually by an encroachment of the under on the upper surface, and the swelling of the whole. There are two kinds of centric laminae, the *succulent-centric* and the *xerophytic centric*.

SUCCULENT.—In a completely centric leaf of the succulent kind, like that of *Sedum murale* the difference between the upper and lower surface is lost. Stomata are found scattered over the entire epidermis. The bundles are arranged in a circle, the mid-rib being in the center. A great deal of mucilage is found stored in the central cells.

XEROPHYTIC.—In a typical Xerophytic Centric leaf, like that of the Pine (see Figs. 168 and 169) or *Sansevieria cylindrica*, the

FIG. 168.—Transverse section of the xerophytic-centric leaf of the Austrian Pine (*Pinus nigra*), magnified. *ep*, epidermis; *s*, deeply sunken stomata; *hf*, hypodermal fibers; *ip*, infolded parenchyma; *o.r.*, schizogenous oil reservoirs; *f*, fibers; *n*, bundle sheath (endodermis) surrounding the stele in which will be noted two vascular bundles with xylem (*x*) directed toward the flat face of the leaf; *ph*, phloem of vascular bundle; *mr*, medullary ray; *scl*, sclerenchymatous cells; *p*, small celled parenchyma of pith.

epidermis shows a thick cuticle; the stomata are sunken in cavities of the epidermis; the epidermis and leaf tissue are strengthened by scleroid bands in the outer portion of the centric mesophyll.

D. BIFACIAL.—Leaves with laminæ which stand edge on in relation to the sun's rays. The best illustrations are seen among dicotyledons, such as *Eucalyptus*, *Callistemon*, and other genera of

the *Myrtaceæ*. The leaflets of *Cassia Senna* and *Cassia angustifolia* which constitute the drug Senna, show a marked bifacial structure. Both surfaces are similar, having stomata about equal in number. The mesophyll (chlorenchyma) is differentiated into a central spongy parenchyma containing bundles, and a zone of palisade cells on either side facing the epidermises (see Fig. 170).

STRUCTURE AND DEVELOPMENT OF STOMATA.—Stomata or stomates are slit-like openings in the epidermis of leaves or young green stems surrounded by a pair of cells, called guard cells, whose sides opposite one another are concave. They form a

FIG. 169.—Portion of a transverse section of the White Pine leaf showing xerophytic-centric structure. *ep*, Epidermis; *s*, stoma; *gc*, guard cells; *s'*, substomal air-space; *h*, hypodermis, consisting of strengthening fibers which form a scleroid band beneath epidermis; *p*, infolded parenchyma cell of mesophyll. Greatly enlarged.

communication between the intercellular-air-space (respiratory cavity) beneath them and the exterior. The slit-like openings, taken with the guard cells, constitutes what is known as the *stomatal apparatus*. The stomatal apparatus is frequently called in practice, "stoma" or "stomate."

The epidermal cells which abut on the stomatal apparatus are called *neighboring cells* or *subsidiary cells*. These, in many cases, as in species of *Helleborus*, *Sambucus*, *Hyacinthus*, *Pæonia*, Ferns, etc., are very similar to the other epidermal cells, but in a large number of plants they differ in size, arrangement and shape from the other cells of the epidermis which do not abut upon the stomatal apparatus. In *Senna* they are mostly two in number, one larger than the other and arranged parallel to the guard cells of the

stoma; in *Coca* a similar arrangement occurs but the cells are more even in size, nevertheless they lack the characteristic papillæ found on the other epidermal cells; in *Pilocarpus* they are usually four in number but quite narrow and more or less crescent-shaped; in *Uva Ursi* their number is usually seven to eight and their arrangement radial around the stomatal apparatus.

FIG. 170.—Photomicrograph of a transverse section of a bifacial leaf of *Eucalyptus globulus* showing epidermis (*ep*), palisade parenchyma (*pal*), toward both surfaces, spongy parenchyma (*sp*), vein (*v*), and oil reservoir (*o.r.*) lined with secretory epithelium. (Highly magnified.)

On all dorsiventral leaves, the stomata arise more abundantly on the lower epidermis, less abundantly on the upper. Exceptions to this rule are due to the peculiar readaptation of the leaf to its surroundings. Thus, in the reversed types of leaves (twisted in a half circle), the stomata, formerly on the lower surface, have migrated to the upper surface which now has become the physiological lower surface.

In Umbrophytic (shade) plants the stomata are either wholly on the lower surface or partly so with a number on the upper surface. Where the plants are Mesophytic and exposed to dense sunlight and the leaves remain dorsiventral, the stomata are on the lower surface; these stomata are large, if the surroundings are damp. If such plants live in dry soil and dry air, the stomata are of small size and numerous; if they dwell in dry soil in hot surroundings and dense light they are very small and frequently sunken. If the plants are Xerophytic and the leaves dorsiventral, the stomata are quite abundant, small, with narrow slit, and depressed below the level of the epidermis.

There are five types of stomatal development, viz.:

FIRST TYPE.—Each primitive epidermal cell (or the majority, or only certain ones of the epidermis) at the close of the dermatogen stage, gradually lengthens and then cuts off a smaller from a larger cell. The smaller one is equilateral, has a very large nucleus, and is termed the *Stoma Mother-cell*; the larger, quadrangular, and called the *Epidermal Daughter-cell*. The latter, upon maturing, becomes a normal epidermal cell. A partition is laid down lengthwise through the *Stoma Mother-cell* dividing it into two stomatal daughter-cells. The wall laid down lengthwise splits and thus forms the orifice of the stoma; the cells on either side of the orifice are called *Guard Cells*. These, while at first flat and inoperative, soon become bulged and crescent-shaped. This mode of development is seen in *Squill*, *Hyacinth*, *Daffodil*, *Sambucus*, *Silene*, etc.

SECOND TYPE.—After the cutting off of the stomal mother-cell, there are cut off, on either side, portions of neighboring epidermal cells which form subsidiary cells to the stoma. This condition is seen in *Gramineæ*, *Cyperaceæ*, *Juncaceæ*, in various species of *Aloe*, *Musa* and *Proteaceæ*.

THIRD TYPE.—Instead of two parallel subsidiary cells, four are cut off, as in *Heliconia*, in species of *Tradescantia*, *Araucaria*, or four to five, as in *Ficus elastica*, or four to five or more, as in the *Coniferæ* and *Cycads*.

FOURTH TYPE.—Instead of only four subsidiary cells, each of these again subdivides by parallel walls, more rarely by radial

walls, into eight radiating subsidiary cells, as in *Maranta bicolor*, *Pothos argyræa*, some of *Proteaceæ*, etc.

FIFTH TYPE.—The “stomal mother-cell” divides once or several times before becoming the true mother-cell of the stoma. As a result of the divisions there are also formed one or more subsidiary cells. This mode of development is seen in the

FIG. 171.—Types of stomatal apparatuses and neighboring cells from different sources. In A, a portion of the lower epidermis of Easter Lily leaf. The stomatal apparatus is surrounded by neighboring cells that are similar to other epidermal cells adjacent to them; in B, lower epidermis of Senna leaflet, note the two neighboring cells parallel to the guard cells, one being larger than the other; C, lower epidermis of Coca leaf showing two neighboring cells, parallel to the guard cells but nearly equal in size as well as papillated regular epidermal cells; D, lower epidermis of Pilocarpus showing rounded stomatal apparatus and four crescent-shaped neighboring cells; E, lower epidermis of Uva Ursi, showing eight neighboring cells arranged radiately around stomatal apparatus; F, lower epidermis of Stramonium.

Labiataæ, *Papilionaceæ*, *Cruciferaæ*, *Solanaceæ*, *Crassulaceæ*, *Cactaceæ*, and *Begoniaceæ*, also in a number of ferns.

HISTOLOGIC DIFFERENCES BETWEEN LEAVES OF DICOTYLEDONS AND MONOCOTYLEDONS.—The following may be cited as broad comparative histologic differences between Dicotyl and Monocotyl leaves:

DICOTYL LEAVES

1. Epidermal cells usually iso-diametric or sinuous in surface sections.
2. The stomata are on the whole more numerous but smaller.
3. Non-glandular and glandular hairs frequent on upper but more frequent on lower surface, or both.
4. Leaf glands which excrete varied products are rather abundant.
5. Water stomata over the upper surface, more rarely over the lower surface, are frequent, especially along margins of leaves.
6. Palisade and spongy parenchyma in dicotyledons are more distinct and palisade parenchyma is denser.
7. The vascular bundles, in their intrinsic elements, are more indurated but the accessory fibrous sheath is feebly developed.
8. A greater variety of accessory products of assimilation are developed.

MONOCOTYL LEAVES

1. Epidermal cells usually elongate with rectilinear vertical walls, in surface sections.
2. Stomata larger.
3. Hairs rare in Monocotyls.
4. Leaf glands rare and only seen as a rule on the sepals.
5. Water stomata absent or very rare. Present in some *Araceæ*.
6. Palisade and spongy parenchyma are less distinct and dense.
7. The vascular bundles, in their intrinsic elements, are less indurated. The fibrous sheath is strongly developed.
8. A comparatively small variety of accessory products of assimilation are developed.

CHAPTER XI

THE INFLORESCENCE

INFLORESCENCE OF ANTHOTAXY.—The terms Inflorescence and Anthotaxy are applied to the arrangement of the flowers and their position on the stem, both of which are governed by the same law which determines the arrangement of leaves.

Indeterminate, racemose, ascending, or centripetal inflorescence is that form in which the terminal bud of the flower cluster continues to develop and increase the length of the stem indefinitely. As the stem elongates new flower buds arise behind the terminal bud in the axils of reduced foliage leaves called bracts, so that the oldest bud is at the base of the flower cluster and the youngest one is nearest the apex. Example: *Digitalis*.

Determinate, cymose, descending, or centrifugal inflorescence is that form in which the opening of the terminal bud as a flower determines or completes the growth of the stem of the flower cluster. Example: *Ricinus communis*.

Mixed inflorescence is a combination of the other two forms. Example: Horse Chestnut.

The flower stalk is known as the *peduncle*. Its prolongation beyond the point where flowers or branches begin to appear is called the *rachis*, or axis of the inflorescence.

The flower stalk of a single flower of an inflorescence is called a *pedicel*. When borne without such support the flower is *sessile*.

A peduncle rising from the ground is called a *scape*, previously mentioned under the subject of stems.

The modified leaves found on the peduncle or its extension, the rachis, are termed *bracts*. These vary much the same as leaf forms, are described in a similar manner, and may be either green or colored. When collected in a whorl at the base of the peduncle they form an *involucre*, the parts of which are sometimes imbricated or overlapping, like shingles. This is generally green, but sometimes petaloid, as in the Dogwoods. The modified leaves found on pedicels are called *bracteolar leaves* or bracteoles.

The *Spathe* is a large bract more or less enveloping the inflorescence and often colored, as in the Calla, or membranous, as in the Daffodil.

INDETERMINATE INFLORESCENCES.—In the indeterminate or centripetal anthotaxy, the flowers are either produced from base

FIG. 172.—Inflorescences. 1, indeterminate; 2, determinate; 3, mixed. (Drawing by Canis.)

to apex, those blossoming first which are lowest down on the rachis or the flowers open from margin to center. Indeterminate inflorescences thus fall into two groups, viz., the elongated or ascending group and the flat-topped or centripetal group. The principal forms of indeterminate inflorescence are:

SOLITARY INDETERMINATE.—The simplest form of inflorescence in which a single flower springs from the axil of a leaf. A number of these are generally developed on the same stem. Example: Periwinkle.

RACEME, or simple flower-cluster in which the flowers, on pedicels of nearly equal length, are arranged along a lengthened rachis, the first flower to open being at the base of the rachis and the last to open at the summit. Examples: *Convallaria*, *Cimicifuga*, and Wild Cherry.

FIG. 173.—Types of indeterminate or centripetal inflorescence. *a*, raceme; *b*, spike; *c*, panicle; *d*, corymb; *e*, umbel; *f*, head (capitulum) of *Compositæ* type; *g*, head of clover type. (Gager.)

CORYMB, a short, broad cluster, differing from the raceme mainly in its shorter axis and longer lower pedicels, which give the cluster a flat appearance by bringing the individual florets to nearly the same level. Example: Hawthorn.

UMBEL, which resembles the raceme, but has a very short axis, and the nearly equal pedicels radiate from it like the rays of an umbrella. When the pedicels of an umbel put forth branches, these branches are termed secondary pedicels. Such, then, bear flowers at their tips and make up with other parts of the inflorescence a **COMPOUND UMBEL**. Many examples of this mode of inflorescence are seen in the family *Umbelliferae*, as indicated by the name, including Anise, Caraway, Fennel and other drug-yielding official plants.

A SPIKE is a cluster of flowers, sessile or nearly so, borne in the axils of bracts on an elongated rachis. The Mullein and common Plantain afford good illustrations. If branches arise from the rachis of a spike and bear flowers, a COMPOUND SPIKE results. In this type of inflorescence which is seen in Wheat, Rye, Barley and many other members of the Grass Family, the branch is called the *rachilla* and the secondary inflorescences are termed *spikelets*.

The CATKIN or AMENT resembles the *Spike*, but differs in that it is deciduous and usually bears only one kind of flowers, staminate or pistillate, as in the Oak, Hazel, Willow, *Comptonia*, etc.

The HEAD or CAPITULUM is like a spike, except that it has the rachis shortened, so as to form a compact cluster of sessile flowers, as in the Dandelion, Marigold, Clover, and Burdock.

The STROBILE is a compact spike of unisexual flowers with overlapping scales concealing the flowers, as the inflorescence of the Hop.

The SPADIX is a thick, fleshy rachis with flowers closely sessile or embedded on it. It is usually more or less enclosed by a spathe or sheathing bract. Examples: *Calla*, *Calamus*, *Arum triphyllum*.

The raceme, corymb, and umbel, are frequently compounded through the branching of their pedicels. The compound raceme, or raceme with branched pedicels, is called a PANICLE. Examples: *Brayera*, *Yucca* and the Oat.

DETERMINATE INFLORESCENCES.—*Determinate Anthotaxy* is one in which the first flower that opens is the terminal one on the axis, the others appearing in succession from apex to base or from

FIG. 174.—Photomicrograph of longitudinal section through a staminate catkin of *Comptonia asplenifolia* $\times 10$, showing catkin axis (*ax*), anther-lobe (*a*) and bract (*b*).

center to margin. If the floral axis is elongated, the order of flower development is *descending*, or from apex to base; if more or less "flat topped," the order is *centrifugal*, or from center to margin. The principal varieties are:

The SOLITARY DETERMINATE, in which there is a single flower borne on the scape, as in the *Anemone*, or Windflower, the Tulip, and *Hydrastis*.

The CYME, a flower cluster resembling a corymb, except that the flower buds develop and open from center to circumference.

FIG. 175.—Cymose inflorescences. *F*, A terminal flower; *G*, a simple cyme; *H*, a compound cyme. (From Hamaker.)

The simplest kind of cyme is seen in the flower cluster of the Linden tree, where a peduncle (arising from a large bract) is elongated into a rachis which ends in a flower bud. At the junction of the rachis with the peduncle, 2 bracteoles arise. In the axil of each of these bracts a bud is formed which elongates in each instance into a pedicel bearing a flower bud at its end. This type of cyme is called a SIMPLE CYME. If each of the pedicels give rise to a pair of bracts from the axils of which secondary pedicels arise bearing flowers, a type of *compound cyme* arises called a DICHASIAL CYME.

A SCORPIOID CYME imitates a raceme, having the flowers pedicelled and arranged along alternate sides of a lengthened axis. These open in right-left hand fashion from summit to base. Example: Sundew.

A GLOMERULE is a cymose inflorescence of any sort which is condensed into a head, as the so-called head of the Dogwoods (*Cornus*).

A VERTICILLASTER is a compact, cymose flower cluster which resembles a whorl, but really consists of two glomerules situated

FIG. 176.—Flowering-branch of a Dogwood showing glomerules. Note the four notched petaloid bracts subtending each glomerule or determinate head of flowers.

in the axils of opposite leaves. Clusters of this kind are seen in Catnip, Horehound, Peppermint and other plants of the *Labiatae*.

MIXED INFLORESCENCES are those in which the determinate and indeterminate plans are combined, and illustrations of this are of frequent occurrence.

A THYRSUS is a mixed inflorescence more or less pyramidal or oblong in shape consisting of a panicle of cymes, as in the Privet, Lilac, Grape or Horsechestnut. Racemes of cymes and corymbs of cymes are other examples of mixed inflorescences.

CHAPTER XII

THE FLOWER

The FLOWER is a shoot which has undergone a series of changes so as to serve as a means for sexual reproduction. It represents a highly modified stem or branch which bears appendages.

A TYPICAL OR COMPLETE FLOWER possesses four circles or whorls of floral leaves arranged upon a more or less shortened and frequently swollen upper end of a stem axis called a RECEPTACLE, TORUS or THALAMUS. These whorls passing from periphery toward the center are: CALYX, composed of parts called sepals; COROLLA, composed of parts termed petals; ANDRÆCIUM, composed of parts called stamens or microsporophylls; and GYNÆCIUM, composed of one or more parts termed carpels, pistils, or megasporophylls. In a complete flower the calyx and corolla constitute the *perianth* or floral envelope.

The stem bearing the flower is called the flower stem or PEDUNCLE.

The stamens and carpels constitute the ESSENTIAL ORGANS, and a flower is said to be PERFECT when these are present and functional. An IMPERFECT FLOWER shows one set of *essential* organs wanting. The calyx, corolla and receptacle constitute the ACCESSORY ORGANS of the flower.

A HERMAPHRODITE FLOWER is one which possesses both stamens and carpels which may or may not be functionally active. In some cases the stamens may alone be functional while in others the carpels only may function.

A REGULAR FLOWER possesses parts of each whorl of the same shape and size, as the flower of the buttercup. This kind of flower possesses radial symmetry and is also called for that reason ACTINOMORPHIC.

An IRREGULAR FLOWER is one in which the parts of one or more whorls differ in shape and size. If it possesses bilateral symmetry, as the flower of the violet or sweet pea, it is also called

ZYGOMORPHIC. Irregularity may result either from variation in form or size of the parts of a whorl, from appendaging, as in the calyx of the Larkspur where one of the 5 sepals bears a long spur, or from abortion.

A SYMMETRICAL FLOWER is one in which the parts of each whorl are of the same number, or multiples of the same number, as the lily or rose.

FIG. 177.—Diagrams of floral structures. A, Shows the relations of the floral parts in a hypogynous flower; B, the same in a perigynous flower; C, the same in an epigynous flower; D, a stamen; E, a simple pistil in longitudinal section; F, the same in cross-section; G, transitional forms between true petals (left) and true stamens (right); H, slight union of two carpels to form a compound pistil; I and J, union of carpels more complete; K and L, cross-sections of compound pistils, of three carpels. In B: *a*, stamen; *b*, petal; *c*, sepal; *d*, pistil; *e*, receptacle; *f*, pedicel. In D: *a*, anther cell; *b*, connective; *c*, filament. In E: *a*, stigma; *b*, style; *c*, ovules; *d*, ovary. (From Hamaker.)

When either petals or sepals, or both, are present in more than the usual number, the flower is said to be “double,” as the cultivated Rose and Carnation. The doubling of flowers is brought about through cultivation and is due either to the transformation of stamens (as in cases cited), and occasionally of carpels into petals, to a division of the petals, or to the formation of a new series of petals.

If one or more pistils are present and stamens wanting, the flower is called PISTILLATE, or female; if it possesses one or more stamens but no pistil, it is described as STAMINATE, or male; if both are absent, NEUTRAL, as marginal flowers of *Viburnum*

and Hydrangea. When staminate and pistillate flowers are borne separately they are called *diclinous* flowers. Flowers containing stamens and pistils are *monoclinous*. Some plants, as the Begonias, Corn, and Castor oil, bear both staminate and pistillate flowers on the same individual, and are called *Monœcious*. When the staminate and pistillate flowers are borne on different plants of the same species, the plants are termed *Diœcious*, as the Sassafras and Willow. When staminate, pistillate and hermaphrodite flowers are all borne on one plant, as on certain species of Maple and Buckwheat, they are *polygamous*.

A **HYPOGYNOUS FLOWER** is one in which the sepals, petals and stamens are attached near the base of the receptacle while the pistil or the carpels are arranged over the upper portion of the receptacle, as in the magnolia and strawberry. The receptacle of this kind of flower is generally conical, dome-shaped or elongated like a spire. The ovaries of this type of a flower are *superior*.

A **PERIGYNOUS FLOWER** is one in which the sepals, petals and stamens are inserted on the edge or summit of a saucer or cup-shaped receptacle while the pistil or carpels are inserted at the bottom of the receptacle, as in the flowers of the cherry, rose and almond. The ovary of this type of flower is *superior*.

An **EPIGYNOUS FLOWER** is one like that of the apple or quince where the receptacle is greatly hollowed out in vase-shaped manner, its summit overarching the top of the ovary and bearing the sepals, petals and stamens above the ovary. The ovary of such a flower is *inferior*.

PREFLORATION.—By prefloration is meant the arrangement of the floral envelopes in the bud. It is to the flower bud what vernalion is to the leaf bud, the same descriptive terms being largely employed, as convolute, involute, revolute, plicate, imbricate, etc.

In addition to those already defined under *vernalion*, the following are important.

Valvate Prefloration, in which the margins of the parts meet but do not overlap. Of this variety the induplicate has its two margins rolled inward, as in Clematis. In the reduplicate, they are turned outward, as the sepals of Althæa.

Vexillary, the variety shown in the corolla of the Pea, where the two lower petals are overlapped by two lateral ones, and the four in turn overlapped by the larger upper one.

Contorted, where one margin is invariably exterior and the other interior, giving the bud a twisted appearance, as in the Oleander and Phlox.

CONNATION AND ADNATION.—In the development of the flowers of primitive species of flowering plants, the parts of each whorl are disjoined or separate from each other. In many higher types, however, the parts of the same whorl frequently

FIG. 178.

FIG. 179.

FIG. 178.—Flower of the Ash (*Fraxinus*); both calyx and corolla absent (*achlamydeous*).

FIG. 179.—Flower of the Goosefoot (*Chenopodium*) with a calyx but no corolla (*monochlamydeous*). (After Small.)

become partly or completely united laterally. This condition is termed *connation*, *coalescence*, *cohesion* or *syngensis*. Illustrations of this may be seen in Belladonna, Stramonium, and Uva Ursi flowers, where the petals have joined laterally to form gamopetalous corollas. When the one or more parts of different whorls are united, as of stamens with petals (*Rhamnus*) or stamens with carpels (*Apocynum*), the union is called *adnation* or *adhesion*.

THE RECEPTACLE.—The *Receptacle*, *Torus*, or *Thalamus* is a more or less shortened axis or reduced branch which bears the floral leaves. It is usually flat or convex, but may be conical and fleshy as in the Strawberry, concave as in the Rose and Fig, or show a disc-like modification, as in the Orange. Like other stems it has nodes and internodes. The internodes of the receptacle in many species lengthen and separate various whorls. When the lengthening of the internode occurs between calyx and

corolla, as in *Lychnis*, the structure resulting is called an ANTHOPHORE; if between corolla and andrœcium, as in *Passiflora*, a GONOPHORE; if between andrœcium and gynœcium, as in *Geum*, a GYNOPHORE. In the flowers of the *Umbelliferae*, the receptacle elongates between the carpels producing the structure called a CARPOPHORE.

THE PERIGONE.—The perigone or perianth is the floral envelope consisting of calyx and corolla (when present).

When both whorls, *i.e.*, calyx and corolla, are present the flower is said to be *dichlamydeous*; if only calyx is present, *monochlamydeous*.

FIG. 180.

FIG. 181.

FIG. 182.

FIG. 180.—Chorisepalous and hypogynous calyx of the Pimpernel (*Anagallis*).

FIG. 181.—Gamosepalous and bilabiate calyx of the Deadnettle (*Lamium*).

FIG. 182.—Urceolate calyx of the Henbane (*Hyoscyamus*). (After Small.)

THE CALYX.—The Calyx is the outer whorl of modified leaves. Its parts are called SEPALS. These may be distinct or separated from one another when the calyx is called CHORISEPALOUS, APOSEPALOUS or POLYSEPALOUS. If the sepals are more or less united at their margins, the calyx is called GAMOSEPALOUS or SYNSEPALOUS.

The sepals are usually green—foliaceous or leaf-like—but may be brilliantly colored, hence the term *petaloid* (like the petals) is applied. Examples of flowers having a petaloid calyx: Tulip, Larkspur, Columbine and Aconite.

In the Daisy Family, etc., the calyx is modified in the form of teeth, bristles, hairs or awns and is then called the PAPPUS.

In a *gamosepalous calyx*, when the union of sepals is incomplete, the united portion is called the *tube*, the free portion, the *limb*, the orifice of the tube, the *throat*.

In form, the calyx may be regular or irregular; regular, if its parts are evenly developed, and irregular if its parts differ in size and shape. The more common forms are *tubular*, resembling a tube; *rotate*, or wheel-shape; *campanulate*, or bell-shaped; *urceolate*

FIG. 183.

FIG. 184.

FIG. 185.

FIG. 183.—Campanulate calyx, surrounded by an epicalyx, of the Rose of Sharon (*Hibiscus*).

FIG. 184.—Perigynous and chorisepalous calyx (*cf*) of Rose with concave receptacular tube (*ct*) with stamens (*e*) inserted perigynously on its edge.

FIG. 185.—Persistente pigynous calyx (*c*) of the Quince (*Cydonia*), showing the receptacular tube completely enveloping the matured carpels. (After Small.)

FIG. 186.

FIG. 187.

FIG. 188.

FIG. 186.—Caducous hairy calyx of the Poppy (*Papaver*).

FIG. 187.—Globose persistent calyx of Winter Cherry (*Physalis*).

FIG. 188.—Perigynous chorisepalous calyx of Indian Cress (*Tropaeolum*) with spurred receptacular cup. (After Small.)

or urn-shape; *hypocrateriform*, or salver-shape; *bilabiate*, or two-lipped; corresponding to the different forms of corolla, under which examples illustrating each will be given.

The calyx usually remains after the corolla and stamens have fallen, sometimes even until the fruit matures, as in the Night-

shade family; in either case it is said to be *persistent*. If it falls with the corolla and stamens, it is *deciduous*, and if when the flower expands, *caducous*, as in the calyx of Poppy and May-apple. If the calyx falls very early, it is called *fugacious*.

SEPALINE NECTARIES AND SPURS.—Occasionally some or all of the sepals may become pouched and at length spurred as nectar receptacles or as receptacles for other parts that are nectariferous. Thus, in the *Cruciferae* or Mustard Family, we occasionally see a

FIG. 189.—Strawberry (*Fragaria chiloensis*). Median lengthwise section of flower. (After Robbins.)

slight pouching of the two lateral sepals. These act as nectar pouches for the nectar secreted by nectar glands on the knobs or girdles surrounding the short lateral stamens. These become deep pouches in *Lunaria* while in others the pouches become elongated spurs. Again, in the Larkspurs, the posterior sepal forms an elongated spur into which pass the two spurred nectariferous petals. In *Aconitum* the same sepal, instead of being spurred, forms an enlarged hood-like body called the *galea* or *hood* which arches over the flower like a helmet; into this pass the

two hammer-shaped nectariferous petals. In the Wild Columbine (*Aquilegia*) each of the 5 sepals develops a spur.

SEPALINE STIPULES.—These structures are well developed and easily traceable in the more primitive herbaceous members of the Rose family. Thus in *Potentilla*, *Fragaria*, *Geum*, etc., in addition to the normal calyx of five sepals, there is a supplementary whorl of green sepal-like lobes. This is outside of the normal calyx, and is called the EPICALYX. The five lobes of the epicalyx may be as large or larger than the sepals or smaller up to the disappearing point. Upon examining a few flowers of *Potentilla* or *Fragaria*, it will be observed that not infrequently one, sometimes two lobes of the epicalyx are bifid, or deeply cleft, or separated completely into two parts. The explanation is that the five sepals, after evolving in the flower bud, form at their bases two lateral swellings or *sepaline stipules*, which, as they grow, fuse in adjacent pairs, one stipule of one sepal joining with the adjacent stipule of another sepal to form five lobes.

SEPALINE POSITION.—The most simple and primitive position for the sepals in relation to the floral parts is HYPOGYNOUS, in which the sepals are inserted directly into the enlarged floral axis (receptacle) below the petals, stamens and carpels. But in the more primitive herbaceous *Rosaceæ*, *Leguminosæ*, etc., the floral axis forms a saucer-like, transverse expansion which pushes out the sepals, petals and stamens on its edge. Thus originates the PERIGYNOUS insertion of the sepals. In not a few higher *Rosaceæ*, *Saxifragaceæ*, *Crassulaceæ*; etc., the saucer-like floral axis becomes deepened and contracted into a cup-shaped structure (Cherry, Peach, Almond, Plum, etc.), and on the edge of this cup the sepals as well as the petals and stamens are inserted at different levels. Finally, in the Apple, Pear, Quince, etc., the greatly hollowed-out receptacle assumes a vase-shaped form and closes over the top of the ovary, at the same time lifting the sepals, petals, and stamens above the ovary. Here the sepals are EPIGYNOUS.

THE COROLLA.—The Corolla is the inner floral envelope, usually delicate in texture, and showing more or less brilliant colors and combinations of color. Its parts are called PETALS, and when the calyx closely resembles the corolla in structure and

coloring they are together called the *perianth*. The purpose of these envelopes is to protect the reproductive organs within, and also to aid in the fertilization of the flower, as their bright colors, fragrance and saccharine secretions serve to attract pollen-carrying insects.

When the petals are not united with each other, the corolla is said to be CHORIPETALOUS, APOPETALOUS or POLYPETALOUS.

FIG. 190.—Sour Cherry (*Prunus cerasus*). Median longitudinal section of flower. Note the perigynous insertion of the sepals, petals and stamens. The ovary is superior. (After Robbins.)

When more or less united into a tubular structure, it is GAMOPETALOUS, often called SYNPETALOUS.

FORMS OF THE COROLLA AND PERIANTH.—When the distinct petals are four in number, and arranged in the form of a cross, the corolla is called *Cruciform*. Example: Mustard and other plants belonging to the family Cruciferae.

The *Papilionaceous corolla* is so called because of a fancied resemblance to a butterfly. The irregularity in this form is very

striking, and the petals bear special names: the largest one is the *vexillum*, or standard; the two beneath it the *alæ*, or wings; the two anterior, the *carina* or keel. Examples: Locust, Pea, and Clover. [Fig. 193 (9).]

Orchidaceous flowers are of peculiar irregularity, combining calyx and corolla. The petal in front of stamen and stigma, which differs from the others in form and secretes nectar, is called the *Labellum* (Fig. 192). Examples: *Cypripedium* and other Orchids.

FIG. 191.—Apple (*Malus sylvestris*). Median lengthwise section of flower. Note the epigynous insertion of sepals, petals and stamens. The ovary is inferior. (After Robbins.)

When calyx and corolla each consist of three parts closely resembling each other in form and color, as in the Tulip and Lily, the flower is called *Liliaceous*.

The *Ligulate* or *Strap-shaped corolla* is nearly confined to the family *Compositæ*. It is usually tubular at the base, the remainder resembling a single petal. Examples: Marigold and Arnica Flowers.

Labiate, or *Bilabiate*, having two lips, the upper composed of two petals, the lower one of three. This form of corolla gives the name to the *Labiatae*, while in the family *Leguminosæ* this arrange-

ment is sometimes reversed. The corolla may be either *ringent*, or gaping, as in Rosemary and Sage, or *personate*, when the throat is nearly closed by a projection of the lower lip, as in Snapdragon and Linaria.

FIG. 192.—Flower of an orchid (*Orchis militaris*). A, a, bract; b, ovary; c, the outer, and d, the two anterior perianth leaves; e, the labellum or lip with spur, f, (nectary); g, the gynostemium the union of stamens and pistil. B, flower after removal of all the perianth leaves with the exception of the upper part of the labellum; h, stigma; l, rostellum, a beak-like extension from the upper edge of the stigma; k, tooth-like prolongation of rostellum; m, anther; n, connective; o, pollinium; q, viscid disk; p, staminodium. C, a pollinium removed and enlarged; r, its stalk (caudicle); s, pollen mass. D, ovary in cross section. (After Strasburger, Macmillan Co., publishers.)

apex. Examples: Canterbury Bell, Harebell.

Urceolate, or *Urn-shaped*, when the tube is globose in shape and the limb at right angles to its axis, as in the official *Uva Ursi*, *Chimaphila* and *Gaultheria*.

Rotate or *Wheel-shaped*, when the tube is short and the divisions of the limb radiate from it like the spokes of a wheel. Example: The Potato blossom.

Crateriform or *Saucer-shaped*, like the last except that the margin is turned upward or cupped. Example: *Kalmia latifolia* (Mt. Laurel).

Hypocrateriform or *Salver-shaped* (more correctly, hypocraterimorphous), when the tube is long and slender, as in Phlox or Trailing Arbutus and abruptly expands into a flat limb. The name is derived from that of the ancient Salver, or hypocraterium with the stem or handle beneath.

When of nearly cylindrical form, the corolla is *Tubular*, as in the Honeysuckle, and Stramonium.

Funnel-form (Infundibuliform), such as the corolla of the common Morning Glory, a tube gradually enlarging from the base upward into an expanded border or limb.

Campanulate, or *Bell-shaped*, a tube whose length is not more than twice the breadth, and which expands gradually from base to

Caryophyllaceous, when the corolla consists of five petals, each with a long slender claw expanding abruptly at its summit into a

FIG. 193.—Illustrating various forms of the corolla. 1, Personate bilabiate corolla of *Linaria*; 2, cruciform corolla of *Rocket*; 3, campanulate corolla of *Harebell*; 4, infundibuliform corolla of *Bindweed*; 5, ringent bilabiate corolla of *Rosemary*; 6, spurred calyx of *Larkspur*; 7, ligulate corolla of *Chrysanthemum*; 8, rotate corolla of *Pimpernel*; 9, papilionaceous corolla of *Irish Broom*; 10, urceolate corolla of *Heath*.

broad limb. Examples: *Carnation* and other members of the *Pink* family.

APPENDAGES OF THE PERIGONE.—In addition to the *spurs* already discussed, the following outgrowths of the perianth may also be encountered:

The AURICLE, an ear-shaped appendage occurring on the calyx of *Lobelia*. This term is also applied to similarly shaped lobes of foliage leaves.

The AWN is a sharp, rigid outgrowth from the tip or dorsal surface of a perigone lobe.

The HORN or CORNUA is a hollow, horn-like appendage.

The CAUDA is a long, drawn out, tail-like appendage, as seen in the petals of the Cacao plant.

The PALATE is a sac-like upgrowth of the lower lip of a personate, bilabiate corolla.

The KEEL is a ridge resembling the keel of a boat.

The CORONA OR CROWN is a circular upgrowth from the throat of the perianth, as seen in *Narcissus*. It may represent the upgrowth and coalescence of stipules of perianth segments or stamens.

THE ANDRÆCIUM OR STAMEN SYSTEM.—The *andræcium* is the single or double whorl of male organs situated within or above the corolla. It is composed of STAMENS or MICROSPOROPHYLLS.

A complete stamen (Fig. 177D) consists of a more or less slender stalk portion called a FILAMENT and a terminal appendage called the ANTHER or MICROSORUS. The anther is generally vertically halved by an upgrowth of the filament, called the CONNECTIVE, dividing the anther into two lobes.

NUMBER OF STAMENS.—When few in number, stamens are said to be *definite*; when very numerous, and not readily counted, they are *indefinite*. The following terms are in common use to express their number:

Monandrous, for a flower with but one stamen.

Diandrous, with two stamens.

Triandrous, with three.

Tetrandrous, with four.

Pentandrous, having five.

Hexandrous, six.

Polyandrous, an indefinite number.

The most primitive flowers have numerous stamens, but passing from these to those of more evolved families there occurs a

gradual reduction from many to ten, as in *Caryophyllaceæ*, *Leguminosæ* and some *Aceraceæ*, these being in two circles. In *Malvaceæ*, *Umbelliferæ* and other Apopetalous families as well as many Synpetalæ, the number five is typical. But in *Scrophulariaceæ*, while five are developed and fertile in *Verbascum*, four with a fifth staminode (sterile stamen) are found in the allied genus *Celsia*. In *Pentstemon* there are four didynamous fertile stamens and an equally long staminode. In *Scrophularia* the fifth staminode is reduced to a petaloid flap in the posterior part of the flower. In *Linaria* this exists only as a small knob at the base of the back part of the corolla and there secretes nectar. In most *Scrophulariaceæ* the fifth stamen is entirely absent and the four stamens left are didynamous; but in *Calceolaria* two of these are rudimentary and thread-like, the other two alone being well-developed and fertile. In *Veronica* three stamens are entirely absorbed and two only are left as fertile representatives.

INSERTION OF STAMENS.—As to insertion the stamens may be:

HYPOGYNOUS, when inserted upon the receptacle below the base of the pistil (see Fig. 177A).

PERIGYNOUS, when inserted on the calyx or corolla above the base of and lateral to the pistil (see Fig. 177B).

EPIGYNOUS, when inserted above the ovary (see Fig. 177C).

GYNANDROUS, when inserted upon the pistil, as in Orchids and *Aristolochia* (see Fig. 194 No. 8).

PROPORTIONS OF THE STAMENS.—The stamens may be of equal length; unequal, or of different length.

DIDYNAMOUS, when there are two pairs, one longer than the other. Examples: Snapdragons, *Digitalis*, Mints. [Fig. 194 (4).]

TETRADYNAMOUS, three pairs, two of the same length, the third shorter. Example: Mustard. [Fig. 194 (6).]

CONNATION OF STAMENS.—Terms denoting connection between stamens are:

MONADELPHOUS (in one brotherhood), coalescence of the filaments into a tube. Examples: *Lobelia* and *Malvaceæ* [Fig. 194 (5)].

DIADELPHOUS (in two brotherhoods), coalescence into two sets. Example: *Glycyrrhiza*. [Fig. 194 (2).]

FIG. 194.—The Stamen System. 1, Stamen of *Hepatica* showing adnate anther, the 2 lobes separated by connective; 2, diadelphous stamens of a leguminous plant; 3, polyadelphous stamens of Orange with calyx at base; 4, flower of *Teucrium* showing didynamous stamens; 5, monadelphous stamens of *Malva*; 6, tetradynamous stamens of a Crucifer; 7, flower of *Hypericum* cut vertically to show triadelphous stamens; 8, androecium and gynoecium of *Aristolochia* showing gynandrous stamens; 9, stamen of a Labiate showing branched connective with one anther-sac aborted and absent; 10, adnate anther of *Campanula*; 11, stamen of *Malva* with confluent anther, which is 1-celled; 12, stamen of Lily with versatile anther; 13, stamen of False Rue Anemone with innate anther; 14, stamens of Sunflower showing syngeneious anthers.

TRIADELPHOUS, with filaments united into three sets. Example: St. John's Wort. [Fig. 194 (7).]

POLYADELPHOUS, when there are several sets or branched bundles. Example: Orange.¹

SYNGENESIOUS, when the anthers cohere. Example: *Compositæ*.

COLOR OF STAMENS.—In most species the color of these organs is seldom pronounced owing to their delicate structure. It varies from greenish-yellow to yellow to white, through pink, pinkish-red, red, purple, purple-blue to blue. It is yellow, for instance, in *Sassafras*, *Cucumber* and *Golden Club*; greenish-yellow, yellow to red, in *Maples*; yellow-pink to pink and pinkish-red, in some *Mallows*; in *Azalea amena* the filaments are crimson-purple and the anthers, purple-blue; in the genus *Scilla* both filaments and anthers are blue.

GROSS STRUCTURE AND HISTOLOGY OF THE FILAMENT.—The filament may be cylindric as in the *Rose*, awl-shaped as in *Tulip*, flat and with a dilated base as in the *Harebell*, three-toothed as in *Garlic*, appendiculate, when it bears an appendage as in *Chaetostoma*, *Alyssum*, etc. The filament is covered with a protective epidermis containing stomata. Beneath this is a soft, loose, cellular tissue, the mesophyll, and in the center a small vascular bundle, the pathway of food from the floral axis to the anther. In some cases the single bundle may split into two or three bundle parts.

GROSS STRUCTURE AND HISTOLOGY OF THE ANTHER.—Each staminal leaf (MICROSPOROPHYLL) bears a special development or appendage, as a rule, on its extremity which is the ANTHER or MICROSORUS. This consists, fundamentally, of a median prolongation of the filament equal to the connective or placenta. This develops on either side a quantity of indusial tissue that grows out to form a covering substance that protects and carries two MICROSPORANGIA on either side. An *anther*, therefore, consists of a median *connective* or placenta, producing on either side two *anther lobes* or indusial expansions. Each anther lobe encloses two pollen-sacs or *microsporangia*, which, in some cases, remain distinct up to the dehiscence (splitting open) of the anther. Thus, in *Butomus*, the anthers show four pollen chambers up to the

¹ In the *Orange* flower each set of stamens consists of a filament bearing in its distal region phalanges of stamens or staminal leaflets.

time of dehiscence. Again, in various species of *Lauraceæ*, the anthers remain four lobed and dehisce by four recurved lids. But in the great majority of Angiosperms each pair of pollen sacs fuse before dehiscence, owing to the breaking down of the partition between them, and so, at that time, show two-celled anthers. Still more rarely the anthers may be two-celled in their young state and by the breaking down of the partition become one-celled, *e.g.*, *Malvaceæ*.

FIG. 195.—Cross section of immature anther of Johnson grass (*Holcus halepensis* L.). The pollen mother cells are contained within four microsporangia. *ex*, exothecium. (Mottier.)

Externally the mature anther is bounded by an EXOTHECIUM or epidermis, often swollen, where lines of dehiscence occur, which may develop stomata, also hairs. Within it is a combined layer or set of one to often two or three, sometimes five or six cell layers (*Agave*, etc.) of indusial and sporangial cells, the ENDOTHECIUM. The outermost one to three layers of this become spirally, annularly or stellately thickened to form the elastic tissue of the anther, which, by pressure against the delicate epidermis or EXOTHECIUM, causes ultimate rupture of the anther wall. Within the innermost endothelial layer, bounding each sporangium, is the TAPETUM, a single-celled layer. This, near the time of dehiscence, undergoes breaking down or absorption

by developing POLLEN or microspore cells. Filling the cavities of the four sporangia are the mature pollen grains. The connective shows in or near its center a vascular bundle with xylem uppermost and phloem downward, surrounded by thin-walled cellular tissue, from which the indusial and sporangial substance has matured by extension.

ANTHER DEHISCENCE.—This is the breaking open of the anther to discharge the pollen.

When fully ripe the dividing partition between each pair of sporangia usually becomes thinned, flattened and ultimately breaks down, while the elastic and resistant endothecium, steadily pushing against the more delicate and now shrinking exothecium causes rupture where endothecium is absent, namely along opposite lines of the anther wall. Thus arises a line of anther dehiscence called *longitudinal anther dehiscence* on either side of the anther sacs. In the division *Solaneæ* of the family *Solanaceæ* which includes Belladonna, in some of the *Ericaceæ* as *Rhododendron* and *Azalea*, etc., the anthers dehisce by small apical pores from which the pollen is shed. This kind of dehiscence is called *apical porous dehiscence*. Again, in *Lauraceæ* and *Berberidaceæ*, the anthers dehisce by recurved valves (see Fig. 202). This is called *valvular dehiscence*.

Moreover, in *Malvaceæ*, the originally longitudinal anther is divided internally by a partition. It gradually swings on the filament so that eventually the anther is transverse and the partition becomes absorbed, thus becoming a one-celled anther with *transverse dehiscence* in its mature state [see Fig. 194 (11)].

DEVELOPMENT OF THE ANTHER.—Each stamen originates as a knob-like swelling from the receptacle between the petals and carpels. This swelling represents mainly future soral (anther) tissue. The filament develops later. When such a young sorus or anther is cut across and examined microscopically, it shows a mass of nearly similar cellular tissue in which the first observable changes are the following:

The surface dermatogen cells become somewhat flattened and regular to form the future epidermis or exothecium of the anther. About the same time, some cells, by more rapid division in the middle of the anther substance, give rise to the elements of the

vascular bundle in the connective. Then, along four longitudinal tracts, rows of cells remain undivided or only divided slowly as they increase in size and around them cells divide and redivide to form the future endothecial and covering tissue to the four sporangia. Next, the four sporangial tracts of undivided cells cut off from their outer surfaces a layer of enveloping cells, the *tapetum*. This consists of richly protoplasmic cells that form a covering to the *spore mother-cells* within. Each spore mother-cell undergoes division and redivision into four spore daughter-cells, at the same time that reduction in the chromatin substance takes place in

FIG. 196.—Cross-section of a mature lily anther. The pairs of pollen chambers unite to form two pollen sacs, filled with pollen grains; *s*, modified epidermal cells at line of splitting. (From a *Text-book of Botany* by Coulter, Barnes, and Cowles. Copyright by the American Book Company, Publishers.)

these cells. Thus originate *tetrads* (groups of four) of *spore daughter-cells* inside the spore mother-cell wall. These continue to enlarge, press against the mother-cell wall, which becomes converted into mucilage, and each of the tetrad cells becomes in time a mature *microspore* or *pollen grain*.

During this time the entire anther is growing in size, the cells of the endothecium in one or more layers becomes thickened by lignin deposits to form a *mechanical endothecium*; the *tapetum* gradually breaks down and appears only, at length, as an irregular layer around the maturing pollen cells. When the anther is finally ripe the partition between each pair of microsporangia becomes narrowed, flattened and ruptured and thus numerous

microspores or pollen grains fill two cavities, one on either side of the connective. The *microspores* or *pollen grains* at first show

FIG. 197.—Various forms of pollen grains. Pollen from *Typha latifolia* (A), *Zea mays* (B), *Ambrosia elatior* (C), *Lilium philadelphicum* (D), *Pinus* (E), *Ranunculus bulbosus* (F), *Carpinus caroliniana* (G), *Althaea rosea* (H), *Oenothera biennis* (I). All highly magnified. Drawing by Hogstad.

only a thin, clear, cellulose layer, but from this, by a differentiation of the exterior film, the *exospore* layer or *exine* becomes cut off.

This becomes cuticular. The cellulose inner layer (*endospore* or *intine*), remains unaltered. In the development of the exospore, one to several deficiencies are usually left in it through which the endospore may protrude later as the rudiment of the pollen tube. These are called *germ pores*.

ATTACHMENT OF ANTHER.—The attachment of the anther to the filament may be in one of several ways, as follows:

Innate, attached at its base to the apex of the filament.

Adnate, adherent throughout its length.

FIG. 198.

FIG. 198.—Orchid Pollinia (*p*), composed of masses of pollen grains, with caudicle (*c*), and disc or retinaculum (*r*). (After Small.)

FIG. 199.

FIG. 199.—Upper part of Orchid flower, showing the retinacula (*a*) in place within the rostellum or sterile stigma. (After Small.)

Versatile, when the anther is attached near its center to the top of the filament, so that it swings freely. The adnate and versatile types are *introrse* when they face inward, *extrorse* when they face outward.

POLLEN.—The pollen grains or microspores vary in form for different species and varieties and while they are averagely constant for these, nevertheless, many exceptions have been recorded. The following are some of the commoner forms:

Four Spore Daughter-cells, hanging together, as in the Cat Tail (*Typha*), forming a pollen grain.

Elongated, simple pollen grains, as in *Zostera*.

Dumb-bell shaped, as the pollen of the Pines.

Triangular, as in the Evening Primrose (*Oenothera*) and Clove.

Echinate, as in the *Malvaceæ*.

Spherical, as in *Geranium*, Cinnamon and Sassafras.

Lens-shaped, as in the Lily.

Spinose, as in the *Compositæ*.

Barrel-shaped, as in *Polygala*.

Under the microscope, the immature pollen grain generally consists of two membranes, an outer firmer one called the *exospore*, which may be variously marked and which possesses deficiencies in the form of "pores" or "clefts," and an inner delicate cellulose membrane called the *endospore*, which surrounds a protoplasmic interior in which are imbedded a nucleus, oil droplets and frequently starch or protein.

POLLINIA.—These are agglutinated pollen masses which are common to the *Orchidaceæ* and *Asclepiadaceæ*.

HAY FEVER POLLENS.—The pollen of many plants, notably certain species of *Compositæ*, *Chenopodiaceæ*, *Gramineæ*, *Amaranthaceæ*, *Fagaceæ*, *Ulmaceæ*, *Salicaceæ* and *Juglandaceæ*, has been shown to be responsible for "Rose Cold" and "Hay Fever." The chief hay-fever pollens are those of the Ragweeds (*Ambrosia* spp.), Pigweeds (*Amaranthus* spp.), Docks (*Rumex*), Wormwoods (*Artemisia*), Russian Thistle (*Salsola*), various grasses and cereals including timothy, corn, wheat, oats and Buffalo grass, and a number of trees including the oaks, elms, poplars, black walnut, and red cedar. From the investigations of Walker and others, it appears that the symptoms of "Rose Cold" and "Hay Fever" which are seasonable forms of bronchial asthma are induced by the sensitivity of individuals to the proteins of certain pollens. At the present time pollen extracts, also called pollen allergens, pollen antigens and pollen atophans, are manufactured from various pollens to be used in the diagnosis, prevention and relief of this disease.

THE GYNÆCIUM OR PISTIL SYSTEM.—This is the female system of organs of flowering plants. It may consist of one or more modified leaves called carpels. Each *carpel* or *megasporophyll* is a female organ of reproduction. In the Spruce, Pine, etc., it consists of an open leaf or scale which bears but does not

FIG. 200.—Pollinia of a Milkweed (*Asclepias*). *b*. Free. *p*. Adhering to the head of the style, *s*. (After Small.)

enclose the *ovules*. In angiosperms it forms a closed sac which envelops and protects the *ovules*, and when complete is composed of three parts, the ovary or hollow portion at the base enclosing the *ovules* or rudimentary seeds, the *stigma* or apical portion which receives the pollen grains, and the *style*, or connective which

FIG. 201.

FIG. 202.

FIG. 203.

FIG. 204.

FIG. 205.

FORMS OF GYNÆCIUM

FIG. 201.—Monocarpellary pistil of Broom (*Scoparius*). *o*, ovary; *s*, style; *t*, stigma. FIG. 202.—Barberry (*Berberis*). *o*, one-celled or *unilocular* ovary; *ov*, ovules attached to marginal placenta (*p*); *st*, sessile stigma. One stamen dehiscing by recurved valves is shown. FIG. 203.—Bicarpellary, syncarpous gynoecium of Tobacco (*Nicotiana Tabacum*). *o*, ovary; *s*, style; *g*, capitate stigma. FIG. 204.—Syncarpous gynoecium of Primrose (*Primula*) with simple style and capitate stigma. FIG. 205.—Median lengthwise section through flower of Pæony (*Pæonia*). *ds*, dorsal suture of carpel; *vs*, ventral suture. (After Small.)

unites these two. The last is nonessential and when wanting the stigma is called sessile, as in the Poppy. The carpel clearly shows its relations to the leaf, though greatly changed in form. The lower portion of a leaf, when folded lengthwise with the margins incurved, represents the *ovary*; the infolded surface upon which the ovules are borne is the *placenta*; a prolongation of the tip of the leaf, the *stigma*, and the narrow intermediate

portion, the *style*. A leaf thus transformed into an ovule-bearing organ is called a *carpel* or *pistil*. The pistils of the Columbine and Pea are made up of single carpels. In the latter the young peas occupy a double row along one of the *sutures* (seams) of the pod. This portion corresponds to the infolded edge of the leaf, and the pod splits open along this line, called the *ventral suture*.

Dehiscence, or the natural opening of the carpel to let free the contained seeds, takes place also along the line which corresponds to the mid-rib of the leaf, the *dorsal suture*.

The gynœcium may consist of a number of separate carpels, as in the buttercup or Water Lily flowers, when it is said to be *apocarpous*, or the carpels composing it may be united together to form a single structure, as in the flowers of Belladonna and Orange, when it is called *syncarpous*.

If the pistil is composed of one carpel, it is called *monocarpellary*; if two carpels enter into its formation, it is said to be *bicarpellary*; if three, *tricarpellary*; if many, *polycarpellary*.

Compound Pistils are composed of carpels which have united to form them, and therefore their ovaries will usually have just as many cells (*locules*) as carpels. When each simple ovary has its placenta, or seed-bearing tissue, at the inner angle, the resulting compound ovary has as many axile or central placentæ as there are carpels, but all more or less consolidated into one. The partitions are called *dissepiments* and form parts of the walls of the ovary. If, however, the carpels are joined by their edges, like the petals of a gamopetalous corolla, there will be but one cell, and the placenta will be parietal, or on the wall of the compound ovary.

OVULES.—The OVULES or MEGASORI are transformed buds, destined to become seeds in the mature fruit. Their number varies from one to hundreds. In position, they are *erect*, growing upward from the base of the ovary, as in the Compositæ; *ascending*, turning upward from the side of the ovary or cell; *pendulous*, like the last except that they turn downward; *horizontal*, when directed straight outward; *suspended*, hanging perpendicularly from the top of the ovary.

In Gymnosperms the ovules are naked; in Angiosperms they are enclosed in a seed vessel.

FIG. 206.

FIG. 207.

FIG. 208.

FIG. 209.

FIG. 210.

FIG. 211.

FIG. 212.

FIG. 213.

FIG. 214.

FIG. 215.

FORMS OF GYNŒCIUM

FIG. 206.—Cone of apocarpous carpels (*c*) of Tulip-tree (*Liriodendron tulipifera*). *a*, receptacle; *e*, stamens. FIG. 207.—Apocarpous carpels of Strawberry, raised upon a swollen, hemispherical carpophore. FIG. 208.—Ripened apocarpous carpels of Raspberry on conical receptacle (*l*). FIG. 209.—Apocarpous carpels of Pheasant's Eye (*Adonis*). FIG. 210.—*Zanthoxylum* with gynophore (*g*) and apocarpous ovary (*o*) free styles but united stigmas (*s*). FIG. 211.—Two bi-lobed carpels and gynobasic style (*s*) of Horehound (*Marrubium vulgare*). FIG. 212.—Similar but more distinctly divided gynœcium of Forget-me-not (*Myosotis*). FIG. 213.—Flower of Rue (*Ruta graveolens*) to show carpels (*ov*) united only at base. FIG. 214.—*a*, Diagram of three apocarpous carpels; *b*, cross-section of the ovaries of the same. FIG. 215.—*a*, Diagram compound pistil of three carpels with free styles and united ovaries; *b*, cross-section of the ovaries of the same. (After Small.)

A complete angiospermous seed ovule which has not undergone maturation consists of a **NUCELLUS** or body; two coats, the outer and inner **INTEGUMENTS**; and a **FUNICULUS**, or stalk. Within the nucellus is found the **EMBRYO SAC** or **MEGASPORE** containing protoplasm and a nucleus. (See Fig. 221.)

The coats do not completely envelop the nucellus, but an opening at the apex, called the **FORAMEN** or **MICROPYLE** admits the pollen tube. The vascular plexus near the point where the coats are attached to each other and to the nucellus is called the **CHALAZA**. The **HILUM** marks the point where the funiculus is joined to the ovule, and if attached to the ovule through a part of its length, the adherent portion of the funiculus is called the **RAPHE**.

FIG. 216.—Forms of Ovules. A, orthotropous. B, campylotropous. C, amphitropous. D, anatropous. (From Gager, redrawn from Gray.)

FORMS OF OVULES.—The shape of the ovule may be **ORTHOTROPOUS**, or straight; **CAMPYLOTROPOUS**, bent or curved; **AMPHITROPOUS**, partly inverted; and **ANATROPOUS**, inverted. The last two forms are most common. A campylotropous ovule is one whose body is bent so that the hilum and micropyle are approximated, as the ovule of mustard.

THE PLACENTA.—The placenta is the nutritive tissue connecting the ovules with the wall of the ovary. The various types of placenta arrangement (placentation) are grouped according to their relative complexity as follows: (1) Basilar, (2) Sutural, (3) Parietal, (4) Central, (5) Free Central.

BASILAR PLACENTATION is well illustrated in the *Polygonaceæ* (Smart Weed, Rhubarb, etc.), in *Piper* and *Juglans*. Here, at the apex of the axis and in the center of the ovarian base, arises a single ovule from a small area of placental tissue.

SUTURAL PLACENTATION is seen in the *Leguminosæ* (Pea, Bean, etc.). Here each carpel has prolonged along its fused edges two cord-like placental twigs, from which start the funiculi or ovule stalks.

PARIETAL PLACENTATION is seen in *Gloxinia*, *Gesneria*, *Papaver*, etc. Here we find two or more carpels joined and placental

tissue running up along edges of the fused carpels bearing the ovules.

CENTRAL OR AXILE PLACENTATION is seen in *Campanulaceæ* (Lobelia, etc.), *Solanaceæ*, etc., where the two, three, or more carpels have folded inward until they meet in the center and in the process have carried the originally parietal placenta with them. This then may form a central swelling bearing the ovules over the surface.

FREE CENTRAL PLACENTATION occurs perfectly in the *Primulaceæ*, *Plantaginaceæ* and a few other families. In this the carpels simply cover over or roof in a central placental pillar around which the ovules are scattered.

STYLE.—The style is the portion of the carpel which connects the stigma with the ovary. It is usually thread-like but may also be considerably thickened. It frequently divides into branches in its upper part. These are called **STYLE ARMS**. As many style arms as carpels may be present. In the one-carpelled pistil of some *Leguminosæ*, the usually bent-up style is the tapered prolongation of a single flower. Again, in the apocarpous carpels of many flowers of the *Ranunculaceæ*, each carpel bears a short to long stylar prolongation. When the carpels, however, are syncarpous, the common

FIG. 217.—Heterostyly, a means of insuring cross pollination. Two forms of flowers of bluets (*Houstonia*) opened out and seen from the inside. A, short stamens and long style. B, long stamens and short style. An insect visiting A, will receive pollen upon the front part of its body; upon visiting B this pollen will rub against the stigmas, and the pollen from this flower will be received by the under part of the body; upon visiting another flower like A, the pollen from B will be brushed against the stigma of A. (From Darwin's "Cross Pollination." Courtesy of D. Appleton & Co.)

styles tend to become more or less fused but usually show lobes, clefts or style arms at their extremities that indicate the number of carpels in each case which form the gynœcium.

In some plants remarkable variations from the typical stylar development may occur. Thus, in *Viola*, the end of the style is a swollen knob on the under surface of which is a concave stigma with a flap or peg. In the genus *Canna*, the style is an elongate, blade-like, flattened body with a sub-terminal stigma. In forms of the *Campanulaceæ*, the style is closely covered with so-called collecting hairs. On these the anthers deposit their pollen at an early period before the flowers have opened. Later, when the flowers open, insects remove the pollen after which the collecting hairs wither. The stigmas then curl apart to expose their viscid stigmatic hairs. In this instance there are two distinct and at separate times functioning hairs on the stylar prolongation, viz.: (a) *collecting stylar hairs*, functioning for pollen collection and distribution; and (b) *stigmatic hairs* for pollen reception from another flower. In *Vinca*, the style swells near its extremity into a broad circular stigma and then is prolonged into a short *column* bearing a tuft of hairs that prevents the entrance of insect thieves into the flower. In the genus *Iris*, the common style breaks up at the insertion of the perianth into three, wide, petaloid, style arms. Each of these bifurcates at its extremity. On the lower or outer face of this is a transverse flap that bears the stigmatic papillæ. In *Physostigma*, the style enlarges at its extremity into a flap-like swelling which bears a narrow stigmatic surface. Finally, in *Sarracenia*, the single style of the five-carpelled pistil enlarges above into a huge umbrella-like portion with five radiating ribs. At the extremity of each bifid end of each rib is a minute, peg-like, stigmatic surface.

FIG. 218.—*Lilium Martagon*. Longitudinal section of the stigma and upper part of the style. The pollen grains, caught on the papillæ of the stigma, have germinated, and the pollen-tubes are growing down along the walls of the style canal. (From Gager after Dodel-Port.)

THE STIGMA.—This is usually a viscid papillose surface of greater or less expanse functioning for pollen reception. If no style intervenes between the ovary and stigma, the stigma is said to be **SESSILE**, as in the poppy.

FIG. 219.—Trimorphism and heterostyly, a means of insuring cross pollination. The three forms of flowers in the loosestrife (*Lythrum salicaria*). The corollas and calyces are omitted in order to show the different lengths of stamens and styles. The functional crosses are indicated by the arrows. (From Darwin's "Cross Pollination," Courtesy of D. Appleton & Co.)

of these are *capitate*, referring to a spheroidal stigma, *bifid*, or two-lobed, *truncate*, when cut square off at the tip, *annular*, or ringed below the summit, *penicillate* or brush-like, *barbellate*, having minute barbs, etc.

POLLINATION

POLLINATION is the transfer of pollen from anther to stigma and the consequent germination thereon. It is a necessary step to fertilization.

In wind-pollinated flowers such as the grasses, the stigmas are the numerous feathery hairs which cover the ends of the styles and are intended to catch flying pollen grains. Such stigmas are called *plumose*. In animal-pollinated flowers, the stigmas are usually small restricted knobs, lines or depressions. The *stigmatic papillæ* vary in size in different plants and even may vary on different individuals of the same species. Thus, in the long styles of *Primula*, the stigmatic papillæ are elongated, columnar, hair-like structures, whereas in the short styles of short-styled flowers the papillæ are small knob-like cellular swellings.

Various terms are ascribed to stigmas to denote their shape, lobing, markings, appendages, etc. Among the more common

When the pollen is transferred to the stigma of its own flower the process is called *Close* or *Self-pollination*; if to a stigma of another flower, *Cross-pollination*. If fertilization follows, these processes are termed respectively, *Close* or *Self-fertilization* and *Cross-fertilization*. Close-fertilization means in time ruination to the race and happily is prevented in many cases by (*a*) the stamens and pistils standing in extraordinary relation to each other, (*b*) by the anthers and stigmas maturing at different times, (*c*) by the pollen in many cases germinating better on the stigma of another flower than its own and (*d*) by staminate and pistillate flowers being borne on different plants of the same species.

In flowers like the geranium, clerodendron and chamomile, the stamens mature and shed their pollen before the stigmas are receptive. Such flowers are called *protandrous*. In the plantains, figworts, etc., the stigma matures before the stamens. Such flowers are *protogynous*.

The agents which are responsible for cross-pollination are the wind, insects, water currents, small animals, and man.

Wind-pollinated flowering plants are called ANEMOPHILOUS; their pollen is dry and powdery, flowers inconspicuous and inodorous, as in the Pines, Wheat, Walnut, Oak, Birch, Beech, Wormwood, Hop, etc.

Insect-pollinated plants are called ENTOMOPHILOUS. These, being dependent upon the visits of insects for fertilization, possess brilliantly colored corollas, have fragrant odors, and secrete *nectar*, a sweet liquid very attractive to insects, which are adapted to this work through the possession of a pollen-carrying apparatus. Examples: Orchids, Irises, Foxglove, Aconites, etc.

Plants pollinated through the agency of water currents are known as HYDROPHILOUS. To this class belong such plants as live under water and which produce flowers at or near the surface of the same. Some of these as *Ruppia* disperse floating pollen grains, others like *Zostera* produce pollen of the same specific gravity as water while *Vallisneria*, the common eelgrass, liberates floating male flowers.

Bird-pollinated plants are called ORNITHOPHILOUS. Some plants like the Trumpet Creeper, Nasturtium and Honeysuckle are pollinated by humming birds. Other plants, as *Aspidistra*,

are pollinated by snails and slugs when they are called MALACOPHILOUS. These have dark colored flowers.

Before the pollen grain has been deposited upon the stigma and during its germination thereon, a series of events affecting both the pollen grain and the embryo sac occur which result in the ultimate formation of the male and female gametophytes.

MATURATION AND FERTILIZATION (IN ANGIOSPERMS)

MATURATION OF THE POLLEN GRAIN AND FORMATION OF THE MALE GAMETOPHYTE.—The substance of the microspore (pollen grain) divides into two cells, the *mother* and *tube cells* of the future male gametophyte. The nucleus of the mother-cell divides to form two sperm nuclei. Within the tube cell is found a tube nucleus embedded in protoplasm. Upon germinating the partition disappears and the thin endospore (*intine*), carrying within it the protoplasm in which are embedded the tube nucleus and two sperm nuclei, penetrates through a deficiency of the exospore (*extine*). The contents of the pollen grain at this stage is called the *male gametophyte*.

MATURATION OF THE EMBRYO SAC AND FORMATION OF THE FEMALE GAMETOPHYTE.—The *nucleus* of the megaspore or embryo sac undergoes division until eight *daughter-nuclei* are produced which are separated into the following groups:

(a) Three of these nuclei occupy a position at the apex, the lower nucleus of the group being the egg or ovum, the other two nuclei being the *synergids* or *assisting nuclei*.

(b) At the opposite end of the sac are three nuclei known as the *antipodals* which apparently have no special function.

(c) The two remaining nuclei (*polar nuclei*) form a group lying near the center of the embryo sac which unite to form a single *endosperm nucleus* from which, after fertilization, the endosperm or nourishing material is derived. This stage of the embryo sac constitutes the *female gametophyte*.

FERTILIZATION IN ANGIOSPERMS.—After the pollen grain reaches the stigma, the viscid moisture of the stigma excites the outgrowth of the male gametophyte which bursts through the

coats of the pollen grain forming a pollen tube. The pollen tube, carrying within its walls two sperm nuclei and one tube nucleus, penetrates through the loose cells of the style until it reaches the micropyle of the ovule, then piercing the nucellus, it enters the embryo sac. The tip of the tube breaks and one of the sperm nuclei unites with the egg nucleus to form the *oöspore*. The second sperm nucleus is absorbed as food. The *oöspore* develops at once into an embryo or plantlet which lies passive

FIG. 220.—Stages in the germination of the pollen grain and development of the pollen tube. (From Robbins, after Holman and Robbins, after Bonnier and Sablon.)

until the seed undergoes germination. In a number of plants including corn, some lilies, etc., the second sperm nucleus unites with the previously fused polar nuclei to form the endosperm nucleus. The union of one sperm nucleus with the egg nucleus and the other sperm nucleus with the two polar nuclei is called *double fertilization*.

FORMATION OF ENDOSPERM.—The endosperm nucleus within the embryo sac soon undergoes rapid division into a large number of nuclei that become scattered about through the protoplasm of the embryo sac. These accumulate protoplasm about them, cell walls are laid down, **ENDOSPERM** resulting.

FIG. 221.—Development of the female gametophyte in an angiosperm. *A*, lengthwise section through a pistil showing a single inverted ovule within the locule of the ovary. The embryo sac is in the center of the ovule surrounded by the nucellus (*N*) and the two integuments (*I*) better shown in *C*. Note how the pollen tube reaches and enters the micropyle of the ovule. *B* shows the development of the eight nuclei in the embryo sac. *C* shows the mature female gametophyte and the tip of the male gametophyte; the former containing eight nuclei and the latter containing only three. *AP*, antipodal nuclei; *E*, egg nucleus; *EN*, two polar nuclei fusing to form endosperm nucleus; *I*, integuments; *N*, nucellus; *S*, sperm nuclei. (After Atwood.)

FIG. 222.—Development of the embryo. This figure is a continuation of the preceding. *D* shows the antipodal cells (*AP*) disappearing, and the fusion of the two sperm nuclei with the endosperm nucleus (*EN*) and the egg nucleus (*E*). The remains of the pollen tube (*P*) are seen in the micropyle (*M*). *E* shows the developing embryo with its suspensor (*S*) and the endosperm (*END*). *F* shows the young embryo and its parts. *C*, cotyledons; *PL*, plumule; *R*, radicle. Note that the embryo is surrounded by four coats. *END*, endosperm; *N*, nucellus; *II* and *OI*, inner and outer integuments. *H* is the hilum. (After Atwood.)

DEVELOPMENT OF THE EMBRYO (DICOTYLEDONS).—In the development of the embryo of a typical Dicotyledonous type, as seen in *Brassica* (the mustard genus), the oöspore, lying within the embryo sac, undergoes division and redivision to form a several celled filament. At the end of the filament farthest away from the micropyle of the ovule, unequal growth and differentia-

FIG. 223.—Successive stages, 1, 2, 3, 4, 6, 8, in the development of the embryo of *Brassica*. 1. Proembryo. 2. Octant stage. *x*, Octant. *i*, *i*, Suspensor. 3. Formation of dermatogen. 4. Differentiation of periblem and plerome. 6. Last cell or hypophysis, (*v*), of suspensor completes the dermatogen by division. 8. Inner part of *v*. *d*¹, *d*², Completes periblem. *v*, Region of stem apex. *h*, *h*, Bases of cotyledons. *D*, Dermatogen. *Pe*, Periblem. *Pl*, Plerome. (From Small, after Kny.)

tion takes place which eventually produces the body or main part of the embryo. Just below this region, the cells of the filament develop into a suspensor whose basal cell is larger than the rest and directed toward the micropylar end of the embryo sac. The body of the embryo is formed as follows: The upper end cell of the axis of the young developing embryo divides to form 2 rows of four cells each. The upper row of 4 cells by unequal growth and repeated division form the terminal epicotyl and 2

lateral cotyledons. The 4 cells of the lower row similarly form the hypocotyl.

MONOCOTYLEDONS.—In the average Monocotyledon, the oöspore divides to form a short row of 3 cells. The basal cell which is the largest of the three does not divide. The middle cell of the row by repeated division and unequal growth of the cells, develops into a suspensor, a hypocotyl and a lateral epicotyl. The terminal cell, by repeated divisions and unequal growth forms the single cotyledon which is terminal on the axis of the embryo.

CHAPTER XIII

THE FRUIT

The FRUIT consists of the matured pistil (carpel or carpels) and contents, or ovarian portion thereof, but may include other organs of the flower which frequently are adnate to and ripen with it. Thus, in the Apples, Pears, and Quinces, the receptacle becomes thick and succulent, surrounds the carpels during the ripening process and forms the edible portion of these fruits. In Dandelion, Arnica, and many other members of the *Compositæ*, the modified calyx or *pappus* adheres to the ovary during its maturation into the fruit and renders the fruit buoyant. In Wintergreen, the calyx becomes fleshy, surrounds the ovary, reddens, and forms the edible part of the fruit. In Ground Cherry (*Physalis*), the calyx enlarges considerably and encloses the ovary in an inflated colored bladder. Involucres frequently persist around and mature with the fruits. These may be membranous as in *Anthemis*, *Matricaria* and other *Compositæ*, leathery and prickly as in the Chestnuts, scaly woody cups (cupules) as in the Oaks, or foliaceous cups as in the Filberts. Occasionally, as in the Fig, Osage Orange, Mulberry, etc., the fruit may consist of the ripened flower cluster or inflorescence.

FRUIT STRUCTURE

THE PERICARP.—The *Pericarp*, or *seed vessel*, is the ripened wall of the ovary. In general the structure of the fruit wall resembles that of the ovary, but undergoes numerous modifications in the course of development.

The number of cells of the ovary may increase or decrease, the external surface may change from soft and hairy in the flower to hard, and become covered with sharp, stiff prickles, as in the *Datura Stramonium* or Jamestown weed. Transformations in consistence may take place, and the texture of the wall of the ovary may become hard and bony as in the Filbert, leathery, as

the rind of the Orange, or assume the forms seen in the Gourd, Peach, Grape, etc.

Frequently, as in the Apple, Pear, Quince, Wintergreen etc., the pericarp consists for the most part of other elements than the ripened ovarian wall and is then termed a *pseudocarp* or *anthocarp*.

The pericarp consists of three layers of different texture, viz.:

FIG. 224.—Fruit of Cherry (a drupe) as seen in median lengthwise section. (After Robbins.)

epicarp, *mesocarp* and *endocarp*. The epicarp is the outer layer. The mesocarp the middle, and the endocarp the inner layer. When the mesocarp is fleshy, as in the Peach, it is called the *sarcocarp*.

When the endocarp within the *sarcocarp* is hard, forming a shell or stone, this is termed a *putamen*.

SUTURES.—The *ventral suture* is a line formed by the coherent edges of a carpellary leaf. The *dorsal suture* is the mid-rib of the carpel. *Parietal sutures* are lines or furrows frequently visible on the walls of fruits, formed by the ripening of a compound ovary. They occur between its dorsal sutures and indicate the places of union between adjacent septa or of two parietal placentæ.

VALVES.—These are the parts into which the mature fruit separates to permit the escape of the seeds. Depending upon the number of these the fruit is said to be *univalved*, *bivalved*, *trivalved*, etc.

DEHISCENCE.—This is the opening of the pericarp to allow the seeds to escape.

Fruits are either *Dehiscent* or *Indehiscent* according as to whether they open to discharge their seeds spontaneously when ripe (DEHISCENT), or decay, thus freeing the seeds, or retain their seeds, the embryo piercing the pericarp in germination (INDEHISCENT). Dehiscent fruits open regularly, or normally, when the

pericarp splits vertically through the whole or a part of its length, along sutures or lines of coalescence of adjacent carpels. Legumes usually dehisce by both ventral and dorsal sutures. Irregular or abnormal dehiscence has no reference to normal sutures, as where it is TRANSVERSE or CIRCUMSCISSILE, extending

FIG. 225.—A. The aggregate fruit of the Strawberry as seen in median longitudinal section, $\times 2\frac{1}{2}$. B, A single achene. The greater bulk of this fruit consists of a fleshy receptacle. Partially imbedded in the surface of this receptacle are numerous seed-like bodies which represent true fruits or achenes. (After Robbins.)

around the capsule either entirely or forming a hinged lid, as in *Hyoscyamus*, or detached.

DEHISCENCE is called POROUS or APICAL when the seeds escape through pores at the apex, as in the Poppy; VALVULAR, when valve-like orifices form in the wall of the capsule.

VALVULAR DEHISCENCE.—There are three kinds of valvular dehiscence, viz.: septicidal, loculicidal and septifragal. SEP-

TICIDAL DEHISCENCE is that form in which the constituent carpels of a fruit become disjoined, and then open along their ventral suture, as in *Colchicum* and Marsh St. John's Wort capsules.

LOCULICIDAL DEHISCENCE, or dehiscence into loculi or cells, is the form which is seen when each component carpel splits down its dorsal suture, and the dissepiments or partition walls separating the ripened carpels remain intact, as in *Cardamom* and *Iris*.

SEPTIFRAGAL DEHISCENCE is that form in which there occurs a breaking away of the valves from the septa or partitions, as in *Stramonium* and *Morning Glory* fruits.

CLASSIFICATION OF FRUITS (according to structure).—**SIMPLE FRUITS** result from the ripening of a single pistil in a flower, as the *Tomato* or *Banana*.

AGGREGATE FRUITS are the product of all the carpel ripenings in one flower, the cluster of carpels being crowded on the ripened receptacle forming one mass, as in the *Raspberry*, *Blackberry*, *Dewberry* and *Strawberry*.

MULTIPLE FRUITS are those which are the product of the ripening of a flower cluster instead of a single flower, as the *Fig* and *Pineapple*.

Simple and Compound fruits are either **Dry** or **Fleshy**. The first may be divided into **Dehiscent**, those which split open when ripe; and **Indehiscent**, those which do not.

SIMPLE FRUITS:

Dry	{	I. Capsular (dehiscing).
		II. Schizocarpic (splitting).
		III. Achenial (indehiscent).
Succulent	{	IV. Baccate (berries).
		V. Drupaceous (stone fruits).

The **CAPSULAR** fruits include all of those, whether formed of one or more carpels, which burst open to let their seeds escape.

SCHIZOCARPIC or splitting fruits are those in which each carpel or each half carpel (in *Labiatae*) splits asunder from its neighbor and then falls to the ground. The split portion is one-seeded.

ACHENIAL fruits are dry, one-celled, one-seeded and indehiscent at the time of final ripening.

BACCATE fruits are such in which the endocarp always and the mesocarp usually becomes succulent and so the seeds lie in the

pulp formed by the endocarp or endocarp and mesocarp combined.

DRUPACEOUS fruits are those in which the endocarp is always fibrous or stony in consistence, while the mesocarp is more or less succulent. The endocarp may become cuticularized as in the Apples. The mesocarp may form stone cells lying in the midst of soft parenchyma cells as in Pears; it may become hardened and thickened by lignin deposits to form fibers as in the Cocoanut, or it may become swollen and soft-succulent as in Peaches, Cherries, etc.

I. CAPSULAR FRUITS.—These may be simple, when composed of one carpel as the follicle and legume, or compound, when composed of two or more carpels as the capsule, pyxis, regma, siliqua or silicule.

The FOLLICLE or pod [Fig. 226 (1)] is a dry, simple, capsular fruit formed of a single carpel which dehisces by one suture. This is usually the ventral suture, as in *Aconite*, *Staphisagria*, *Larkspur* and some other *Ranunculaceæ*, but may be the dorsal suture as in *Magnolia* and *Illicium* (Star anise).

A LEGUME is a dry, simple, capsular fruit formed of a single carpel and dehiscent by both ventral and dorsal sutures. Examples: Peas, Beans, etc. The legume is typical of most *Leguminosæ*, Fig. 226 (8).

A LOMENTUM or LOMENT is a modified, usually jointed legume that is separable transversely into one-seeded portions. Seen in *Acacia*, Peanut, Tamarind, *Entada scandens*, *Cathartocarpus Fistula*, *Desmodium*, etc. of *Leguminosæ*, Fig. 227 (5).

A CAPSULE is a fruit formed of two or more carpels composing a compound pistil which dehisce longitudinally or by apical teeth or valves. Examples: *Cardamom*, Poppy, *Iris*, etc., Fig. 226 (2 and 3).

A PYXIS or PYXIDIUM is a capsular fruit formed of two or more united carpels that dehisce transversely. Examples: *Hyoscyamus*, *Plantago*, *Portulaca*. The upper portion forms a lid which fits upon the lower pot-like portion, Fig. 226 (4).

A REGMA is a capsular fruit of two or more united carpels that first splits into separate carpel parts and then each of these dehisces. This type of fruit is typical of *Hura crepitans* (Sandbox), *Pelargonium* (House Geranium) and *Geranium*, Fig. 226 (5).

A **SILIQUE** is a long, slender, one or two-celled capsule, often with a spurious membranous septum (when two-celled) and two persistent parietal placentæ, the valves opening from below upward. Examples: Mustard, *Chelidonium* and Wallflower, Fig. 226 (6).

FIG. 226.—Types of capsular fruits. 1, Pod of Aconite; 2, Capsule of Colchicum showing septicidal dehiscence; 3, capsule of poppy, having porous dehiscence; 4, pyxis of Henbane; 5, dehiscing regma of Geranium; 6, siliqua of Celandine showing valves opening from below upward; 7, silicule of Cochlearia; 8, Legume of Pea.

A **SILICULE** is a short siliqua in which the length is never much greater than the breadth. Examples: *Cochlearia*, Fig. 226 (7); Shepherd's Purse, etc.

II. **SCHIZOCARPIC FRUITS**.—A **CARCERULUS** or **NUTLET** is the typical fruit of the *Labiata*, but is also seen in the *Boraginaceæ*, The ovary that has become four-celled at the time of flowering matures into four little pieces which split asunder lengthwise.

Each split part is composed of one-half of a ripened carpel, Fig. 227 (2).

A CREMOCARP is the characteristic splitting fruit of the *Umbelliferae* family. It consists of two inferior akenes or mericarps separated from each other by a forked stalk called a *carpopore*. These mericarps usually cling to the forks of the

FIG. 227.—Schizocarpic fruits, and loment. 1, Cremocarp of Fennel, composed of two mericarps (*m*) and a split carpophore (*c*); 2, carcerulus of the Bugle; 3, one-carpelled samara of the Ash; 4, double samara of the maple; 5, loment of purging cassia, a portion of the pericarp being removed to show chambers, each containing a single seed.

carpopore for a time after the cremocarp splits, but sooner or later fall, Fig. 227 (1).

A SAMARA is a winged splitting fruit. It may be one-carpelled as in the Elm, Ash, Tulip Poplar and Wafer Ash or two-carpelled as in the Maples. See Fig. 227 (3 and 4).

III. ACHENIAL FRUITS (all indehiscent).—The AKENE is a dry, one-chambered, indehiscent fruit, in which the pericarp is firm and not united with the seed, the style remaining in many cases as an agent of dissemination, Fig. 228 (1). The latter may be long and feathery as in *Clematis* or be hooked. Exam-

ples of akenes: Fruits of the *Compositæ*, Sedges, Buttercup, *Anemone*, etc. The HIP of the Roses consists of a number of akenes in a ripened concave receptacle.

The UTRICLE is like the akene, except that the enveloping calyx is loose and bladder-like. Examples: *Chenopodium*, Fig. 228 (3).

A CARYOPSIS or GRAIN is a dry, one-celled indehiscent fruit whose pericarp is always fused with the seed coat. This fruit is more likely than any other to be mistaken for a seed. Examples:

FIG. 228.—Achenial fruits. 1, Akene of *Pulsatilla* cut vertically, showing adherent feathery style (*st*), pericarp (*p*), testa (*t*), endosperm (*e*), hypocotyl (*h*) and cotyledons (*cot*) the last two structures making up the embryo; 2, caryopsis of wheat showing beard of hairs above and position of embryo of seed below; 3, utricle of *Chenopodium* cut vertically to show calyx (*c*), pericarp (*p*) and seed (*s*) regions; 4, nut of an oak consisting of a glans (*g*) and cupule (*cu*).

Wheat, Corn, Barley, Oats and other members of the *Gramineæ* or Grass Family. Fig. 228 (2).

A NUT or GLANS is a one-celled, one-seeded fruit with a leathery or stony pericarp. Examples: Oaks, Beeches, Chestnuts, Alders and Palms, Fig. 228 (4).

IV. BACCATE FRUITS (Succulent fruits in which the endocarp is always succulent and the mesocarp sometimes).—The BERRY is a small fleshy fruit with a thin membranous epicarp and a succulent interior in which the seeds are imbedded. Examples: *Capsicum*, Tomato, Belladonna, Grape, Currant, etc.

An UVA is a berry from a superior ovary. Examples: Belladonna, Egg-plant, Tomato, Fig. 229 (1).

A **BACCA** is a berry from an inferior ovary. Examples: Gooseberry, Honeysuckle, Currant.

The **PEPO** or **GOURD FRUIT** is a baccate fruit of large size which has developed from an inferior ovary. It is fleshy internally and has a tough or very hard rind. Examples: Fruits of the *Cucurbitaceæ* and the Banana, Fig. 229 (2).

The **HESPERIDIUM** is a large, thick-skinned, succulent fruit with seeds embedded in the pulp but from a superior ovary. Examples: Orange, Grape-fruit, Lemon, etc. In each of these there is to be noted a glandular leathery epicarp, a sub-leathery mesocarp and an endocarp in the form of the membranous cover-

FIG. 229.—Baccate fruits. 1, Berry (uva) of Belladonna with adherent calyx; 2, Pumpkin, cut transversely illustrating a pepo fruit; (h), a locule; 3, hesperidium fruit of the Orange cut transversely showing epicarp (e), mesocarp (m), endocarp (en), pulp (p), and seed (s).

ings of separate carpels. From the endocarp hairs grow inward into the carpellary cavities and become filled with succulence. The seeds lie amid the hair cells, Fig. 229 (3).

V. **DRUPACEOUS FRUITS** (succulent fruits in which the mesocarp is more or less succulent, but the endocarp leathery or stony).—A **DRUPE** is a one-celled, generally one-seeded, drupaceous fruit such as the fruit of the Plum, Peach, Prune, Sabal, Rhus, Piper, Cherry, etc., whose endocarp or putamen is composed wholly of stone cells or stone cells and sclerenchyma fibers, Fig. 230 (1).

The **POME** is a fleshy, drupaceous fruit, usually 5-carpelled, with a tough epicarp, succulent or fleshy mesocarp, and leathery or stony endocarp, the chief bulk of which consists of the adherent torus. Quince, Apple and Pear are examples. The carpels

constitute the core, and the fleshy part is developed from the torus, Fig. 230 (2) and 231.

FIG. 230.—1, Drupe of cocoanut cut vertically, showing epicarp (*e*), mesocarp (*m*), stony endocarp (*d*), seed coat (*s*), endosperm (*end*), and embryo sac cavity (*e.s.*) which in the mature seed contains a nutritive fluid. 2, Pome of an apple cut vertically to show core composed of 5 ripened carpels and flesh of matured torus. 3, Etærio of raspberry. 4, Same, cut vertically to show arrangement of the little drupes on fleshy receptacle.

AGGREGATE FRUITS.—An ETÆRIO consists of a collection of little drupes or of achenes on a ripened torus of a single flower.

FIG. 231.—Pome of apple. *A*, median lengthwise section; *B*, cross section of fruit. (Robbins.)

Examples: Raspberry, Blackberry, Strawberry, etc., Fig. 230 (3 and 4).

MULTIPLE FRUITS.—The **SYCONIUM** is a multiple fruit consisting of a succulent, hollow torus enclosed within which are akene-like bodies, products of many flowers. Example: Fig. 232 (1).

The **SOROSIS** is represented by the Mulberry, Osage Orange, etc., the grains of which are not the ovaries of a single flower, as in the Blackberry, but belong to as many separate flowers. In the Pineapple all the parts are blended into a fleshy, juicy, seedless mass, and the plant is propagated by cuttings.

FIG. 232.—Multiple fruits. 1, Syconium of Fig cut vertically to show hollowed out receptacle (*r*) of ripened flower cluster; 2, strobile of the Hop; 3, galbulus of Juniper.

The **STROBILE** or **CONE** is a scaly, multiple fruit consisting of a scale-bearing axis, each scale enclosing one or more seeds. The name is applied to the fruit of the Hop, Fig. 232 (2), and also to the fruit of the *Coniferae* in which the naked seeds are borne on the upper surface of the woody scales.

A **GALBULUS** is a more or less globular multiple fruit formed of three fleshy connate scales each bearing a seed on its upper or inner surface, as in Juniper, Fig. 232 (3).

HISTOLOGY OF A CAPSULE, VANILLA.—The vanilla fruit is a one-celled capsule formed by the union of three carpellary leaves and dehiscing by two unequal longitudinal valves.

MICROSCOPIC APPEARANCE OF A TRANSVERSE SECTION.—Passing from periphery toward the center, the following structures present themselves:

1. EPICARP or *epidermis* consisting of a layer of thick-walled epidermal cells whose outer walls show the presence of a thin yellow cuticle. Stomata are present in this tissue. The epidermal cells contain protoplasm and brownish bodies. Some also contain small prisms of calcium oxalate and a few vanillin crystals.

FIG. 233.—Vanilla. Structure of the capsular fruit. *c*, cuticle; *ep*, epidermis; *hy*, hypodermis; *mes*, mesocarp of parenchyma in which region fibrovascular bundles (*b*) course; *tr*, tracheae, *si*, sieve tissue and *f*, sclerenchyma fibers of a bundle; *end*, endocarp; *pl*, placenta; *h*, glandular hairs; *sd*, seed; *s*, suture; *cr'*, prismatic crystals and *cr*², raphides of calcium oxalate.

2. MESOCARP, a broad region divisible into a narrow, outer hypodermis and broad inner parenchyma zone. The *hypodermis* is composed of one to several layers of collenchymatic cells with dark-colored contents. Its cells are somewhat larger than those of the epidermis and thicker-walled. The inner parenchyma zone is composed of somewhat loosely arranged, large, thin-walled parenchyma cells becoming smaller in the inner part of this region. Most of these cells contain brownish contents but some possess long raphides of calcium oxalate. If the section be mounted in phloroglucin solution (5 per cent.) and

a drop of strong sulfuric acid is added, a carmine-red color will be observed showing the presence of vanillin in this region. A number of closed collateral bundles will be seen coursing through the mesocarp.

3. ENDOCARP, an irregular line of inner epidermal cells which is differentiated into two regions, the *interplacental region* and the *placental region*. The interplacental (inner) epidermis shows its cells elongated into numerous thin-walled glandular hairs which contain an abundance of balsam; the placental region covers the six bifid *placentæ* which extend into the cavity of the capsule. Its (inner) epidermis is composed of mucilaginous cells.

FIG. 234.—Photomicrograph of a transverse section of a mericarp of *Fœniculum vulgare*, showing epicarp (A), mesocarp (B), endocarp (F), vitta (C), endosperm of seed (D), carpophore (G) and fibro-vascular bundle in primary rib (E). (Highly magnified.)

4. SEEDS.—These are minute blackish bodies attached to the placental twigs of the placentæ. Some of them may have been torn off in cutting the section.

HISTOLOGY OF A TYPICAL MERICARP, FŒNICULUM.—This five-angled fruit, in transverse section, shows a concave *commissural* and convex *dorsal surface*. Passing from the surface toward the center we note:

1. *Epicarp*, or outer covering tissue, composed of colorless epidermal cells and small stomata. The epidermal cells in cross-section appear rectangular, while in surface view they are both polygonal and rectangular.

2. *MESOCARP*, of several layers of thin-walled, colorless, isodiametric cells, beneath which are two to several additional

layers of thicker-walled cells with brownish walls. Through the angles or *rib* portions of the mesocarp extend the *fibro-vascular bundles*. Each of these consists of a few spiral, annular and simple pored tracheids and tracheae and many sclerenchyma fibers. Between each fibro-vascular bundle and the tip of each rib will be found a zone of collenchyma cells. In the mesocarp between each two ribs on the dorsal side occurs a single *VITTA* or *oil tube* which is lined with a layer of brownish polygonal cells. These vittæ contain the official oil of fennel. Two vittæ generally occur in the mesocarp of the commissural side although four are reported to have been found in this region of some varieties of fennel.

3. *Endocarp*, a narrow layer of cells, transversely elongated, except over the regions of bundles where they may be seen elongated in several directions.

4. *Spermoderm or Seed Coat*, consisting of a layer of somewhat broadened epidermal cells attached to the endocarp and several layers of collapsed cells which are only well defined in the region of the *raphe*.

5. *Endosperm*, a central, somewhat reniform mass of more or less polygonal cells containing aleurone grains and fixed oil globules. Each aleurone grain generally contains a rosette aggregate of calcium oxalate and one or two globoids.

6. *Embryo*, embedded in the endosperm of the upper region of the seed, with its hypocotyl directed upward.

CHAPTER XIV

THE SEED

A SEED is a fertilized and matured ovule containing an embryo.

The EMBRYO is a young plant in a dormant state of existence. It is capable of remaining alive for a long period under a broad range of conditions of temperature and moisture. When fully developed within the seed it consists of four principal parts: the *caulicle* or embryonic stem, the *plumule* or embryonic bud, the *radicle* or embryonic root and one, two or more seed leaves called *cotyledons*.

The plumule is at the summit of the caulicle and consists of the young growing apex of the caulicle with delicate rudimentary leaves enveloping it. The radicle is a continuation of the lower end of the caulicle. It develops, upon the sprouting of the seed, the primary or first root of the seedling. The cotyledons frequently are carried above the soil, upon germination, and there function to build up food for the seedling plantlet until the latter has developed foliage leaves. In some seeds the cotyledons contain stored food which is used by the embryo as it develops into the seedling plantlet.

All that portion of an EMBRYO PLANTLET above the cotyledons is termed the *epicotyl* and all that part below the cotyledons, the *hypocotyl*.

The purpose of the seed is to insure the continuation and distribution of the species. It is therefore an organ of reproduction. Like all other organs of the plant, seeds exhibit irritability, nutrition, respiration and growth. These, in the dry seed, are in the state of reduced activity. When properly stored, seeds may live for one year, as those of Aconite and Larkspur, or one to three years, as those of the Grass family, or two to five years, as those of the Rose and Mallow families, up to 20 years, as those of Tobacco or rarely 5 to 87 years as those of the Bean and Pea family.

SEED STRUCTURE.—The average seed shows one or two **SEED COATS** that enclose an **EMBRYO** which fills the seed cavity; the seed is then termed *exalbuminous*, as in beans, peas and other leguminous seeds; or between the seed coat and embryo reserve food may be stored, and the seed is then termed *albuminous*, as in the cereals, castor bean, etc. The stored food in or around the embryo may consist of starch and protein (pea, bean), or starch,

FIG. 235.—Castor-oil seed (*Ricinus communis*), an albuminous type. A, view of seed halved flatwise, showing surface of cotyledon and hypocotyl. B, longitudinal section at right angles to the surface of the papery cotyledons. (Mottier.)

protein and oil (peanut), or oil and protein and active medicinal principle (castor and croton), or of hard reserve cellulose and protein (date, nux vomica, coffee, ivory-nut, palm-nuts). The seed coats, corresponding to those of the ovule, are one or two in number. If but one seed coat is present it is termed the **SPERMODERM**. If two are present, the outer one is called the **TESTA**, and the inner one the **TEGMEN**. The testa, or outer seed shell, differs greatly in form and texture. If thick and hard, it is

crustaceous; if smooth and glossy, it is polished, if roughened, it may be pitted, furrowed, hairy, reticulate, etc.

The testa may often present outgrowths or seed appendages whose functions are to make the seeds buoyant, whereby they may be disseminated by wind currents. Examples of these are seen in the Milkweed, which has a tuft of hairs at one end of the seed called a COMA, in STROPHANTHUS, which has a long bristle-like appendage attached to one end of the seed and called an

FIG. 236.—Appendages of seeds. A, coma (c) on milkweed (*Asclepias*); B, strophiole (st) on bean (*Phaseolus*) h, hilum, m, micropyle; C, strophiole on *Corydalis aurea*; D, aril on white water-lily (*Castalia*); E, projecting raphe (r) on pitcher plant (*Sarracenia purpurea*); F, wing on trumpet creeper (*Tecoma radicans*); G, arillode ("mace") on nutmeg. A, B, and G from nature; C-F, redrawn from Gray. (After Gager.)

AWN, and in the seeds of the Trumpet Creeper and Pines which have *wings*. The excrescence or slightly developed aril at or near the hilum of some seeds, as in Castor and Colchicum Seed, is called the CARUNCLE or STROPHIOLE.

The tegmen or inner coat surrounds the nucellus closely and is generally soft and delicate.

A third integument, or accessory seed coat covering the outside of the testa, is occasionally present and is called the ARIL or arillus. Examples: *Evonymus* (succulent) and Cardamom (dry).

When such an integument arises from the dilatation of the micropyle of the seed, as in the Nutmeg, it is known as an ARILLODE.

The KERNEL consists of all the seed substance within the seed coats. ALBUMEN is the name of the nutritive matter stored in the seed. The FUNICULUS or seed stalk is absent in most of the official seeds. The scar left by its separation is called the HILUM. When the funiculus is continued along the outer seed coat, it is called the RAPHE.

MODE OF FORMATION OF DIFFERENT TYPES OF ALBUMEN

If the fertilized egg-cell within the embryo sac segments and grows into the embryo and, stretching, fills up the cavity without food material laid down around it, it happens that the nutritive material lingers in the cells of the nucellus, pressing around the embryo. This nutritive material is called *Perispermic albumen*. The tissue of the seed containing it is called the PERISPERM and represents the ripened nucellus of the ovule. Seen in the *Polygonaceæ* or Buckwheat family, etc.

In by far the greater number of Angiosperms, the endosperm nucleus, divides and redivides, giving rise to numerous nuclei imbedded in the protoplasm of the embryo sac, outside of the developing embryo. Gathering protoplasm about themselves and laying down cell walls they form the ENDOSPERM tissue outside of the embryo. Into this tissue food is passed constituting the *Endospermic albumen*.

In the Arrowroot family (*Marantaceæ*), the Pepper family (*Piperaceæ*), etc., nutritive material is passed into the nucellar cells causing them to swell up, while to one side a small patch of endosperm tissue accommodates a moderate amount of nourishing substance, thus resulting in the formation of abundant PERISPERM and a small reduced ENDOSPERM.

EXALBUMINOUS SEEDS are those in which the albumen is stored in the embryo during the growth of the seed. Such seeds show the fleshy embryo taking up all or nearly all the room within the seed coat. Examples: PHYSOSTIGMA, AMYGDALUS, etc.

ALBUMINOUS SEEDS are those in which the nourishment is not stored in the embryo until germination takes place. Such

seeds show a larger nourishing tissue region and a smaller embryo region. Examples: NUX VOMICA, MYRISTICA, COLCHICUM, CASTOR, etc.

GROSS STRUCTURE OF A MONOCOTYL SEED (WITH FRUIT WALL ATTACHED), INDIAN CORN.—The ripened seed of Indian Corn is surrounded by a thin, tough pericarp which is firmly adherent to and inseparable from the *spermoderm* or seed coat. Because of this fact, while in reality a fruit called a caryopsis or grain, this structure is sometimes erroneously termed a seed.

The mature grain of most varieties of Indian Corn is flattened and somewhat triangular in outline, the summit being broad and the base comparatively narrow. The summit is indented and often marked by a small point which represents a vestige of the style. The base or "tip" region marks the part of the grain which was inserted into the cob. Upon it may be found papery chaff, representing parts of the pistillate spikelet. The groove noted on the broader surface indicates the position of the embryo.

HISTOLOGY OF THE INDIAN CORN SEED (WITH FRUIT WALL ATTACHED).—If a longitudinal section be cut through the lesser diameter of a soaked grain, the following histologic characteristics will be observed:

1. The PERICARP or ripened ovarian wall which, alike with all other grains, adheres firmly to the wall of the seed forming a portion of the skin of the grain. The pericarp comprises an outer *epicarp* of elongated cells with thin cuticle, a *mesocarp* of thicker walled cells without, becoming thinner within, and a layer of *tube cells*.

2. The SPERMODERM or seed coat, a single layer of delicate elongated cells.

3. The PERISPERM, another layer directly underneath the Spermoderm, difficult to distinguish without special treatment, and representing the ripened nucellar tissue of the ovule.

4. The ENDOSPERM or nourishing tissue, consisting of: (a) The *Aleurone Layer*, for the most part a single row of cells, containing aleurone grains. Some of the cells may be seen to be divided by tangential partitions. (b) *Starch Parenchyma*, consisting of two regions: an outer horny zone composed of cells containing for the

FIG. 237.—Indian Corn grain and seedling. 1, Median longitudinal section through lesser diameter of grain of corn showing fruit wall (pericarp) and adherent seed coat (spermoderm) $p + s$; horny endosperm (end^1); starchy endosperm (end^2); palisade layer (pal) scutellum (s) with vascular tracts (v); cotyledonary sheath (cot); radicle (r) and plumule (pl). 2. Transverse section of a representative portion of outer region of corn grain showing pericarp (p) and spermoderm (s) with tube cells ($t.c.$); endosperm (end) with an outer layer of aleurone cells containing aleurone grains (al) and an inner starch parenchyma zone (st). 3. Young seedling showing first leaves (l); primary root (r) bearing secondary roots (sr) with root hairs (h); cotyledonary sheath (cs) and seed coat (s).

most part polygonal starch grains and oil droplets; and an inner mealy zone of cells with mostly rounded starch grains.

FIG. 238.—Median longitudinal section through the lesser diameter of a grain of corn (*Zea Mays*). *p*, pericarp which is attached to subjacent seed coat; *al*, aleurone layer of endosperm; *end*, horny endosperm; *fe*, floury endosperm; *e*, epithelium of scutellum; *sc*, scutellum; *cr*, coleorhiza; *r*, radicle; *pl*, plumule. $\times 8$.

5. The EMBRYO, consisting of a single shield-shaped *cotyledon* adjoining the endosperm, the *plumule* or rudimentary bud at the end of the *caulicle* or rudimentary stem and the *radicle* or rudimentary root, with its tip covered by a root cap. The cotyledon

or seed leaf consists of two parts: the *scutellum* which lies next to the endosperm, and is an organ of absorption, and the *sheathing* portion which surrounds and protects the rest of the embryo. That portion of the cotyledonary sheath which surrounds the radicle is called the *coleorhiza*; that portion surrounding the plumule, the *coleoptile*.

The embryo contains oil and proteids, but no starch.

If a similar longitudinal section of a soaked grain be mounted in dilute iodine solution, the contents of the aleurone cells will be colored yellow indicating their proteid nature, while the starch grains will take on a blue to violet coloration. The endosperm will be observed taking up most of the room within the seed coat. The contents of its cells are not baled out to the embryo until after germination begins. Indian Corn is therefore an *albuminous seed*.

GERMINATION OF THE SEED

When any viable seed is planted in suitable soil, and furnished with oxygen and water and a certain degree of heat, *germination* takes place. The first step in germination is the *imbibition* or taking in of water. In the presence of moisture, etc., the seed imbibes water and the embryo and endosperm (when present) swells. The ferments present within the cells of the embryo and endosperm then change the insoluble proteid, starch, and oil to soluble materials, which, in the case of Indian Corn, are absorbed in solution by the scutellum which bales this nourishment out to other parts of the growing embryo, there to be used in part in constructing new tissues, and in part to be consumed by oxidation or respiration. The process of respiration or breathing takes place when the plant takes in oxygen and gives off carbon dioxide. The oxygen oxidizes the food materials within the tissues with an accompanying release of energy, which is necessary to life and growth.

The combined pericarp and spermoderm bursts opposite the tip of the radicle, and the radicle, piercing through the cotyledonary sheath, protrudes. The cleft in the coat lengthens to the point opposite the tip of the plumule, which also protrudes after bursting through the cotyledonary sheath. The radicle, next,

grows downward into the soil forming the *primary root*, and develops upon itself *secondary* or lateral roots, all of which give rise to root-hairs just above their root caps. Additional lateral roots emerge above the scutellar region which ere long attain the size of the first or primary root. The caulicle, carrying upon its tip the plumule, elongates and forms the *stem*; the leaves of the plumule spread out and turn green to function as *foliage leaves*. The perforated cotyledonary sheath grows out surrounding both the root and the stem for a portion of their length. By this time

FIG. 239.—Different ways in which seedlings emerge from the soil. *A*, garden pea; *B*, corn; *C*, Lima bean; *g*, ground level. (Mottler.)

all or nearly all of the nourishment stored in the endosperm has been absorbed and assimilated by the young seedling and the coat and scutellum, left behind, gradually decay and disappear. The root-hairs absorb nourishment from the soil, the green leaves build up carbohydrates, *prop-roots* make their appearance at the first node (joint) above ground, and the seedling grows larger.

GROSS STRUCTURE OF A DICOTYL SEED, PHASEOLUS LUNATUS (Lima Bean).—The Lima Bean Seed shows a flattened-ovate to somewhat reniform outline. Externally it exhibits a polished seed coat which is perforated on its basal side by a minute pore

FIG. 240.—Lima Bean. A, seed showing scar (hilum); B, embryo after removal of seed coat showing one cotyledon (*cot*) and caulicle (*c*); C, embryo after removal of one of the cotyledons showing the second cotyledon (*cot*), plumule (*p*), caulicle (*c*), and radicle (*r*); D, young seedling showing epicotyl (*ep*) bearing first pair of green leaves (*l*) and a terminal stem bud (*b*); cotyledons (*c*), hypocotyl (*h*) with first root (*r*) and rootlets (*rt*); E, cross section of representative portion of seed showing cuticle (*c*), palisade cells (*p*), column cells (*h*) and spongy parenchyma (*sp*) of spermoderm; epidermis (*ep*) and parenchyma (*pa*) of mesophyll of cotyledon (magnified).

called the *micropyle* or *foramen*. Just below this pore will be noted the *hilum* or scar which represents the point of detachment from the *funiculus* or stalk, which connected the seed during its growth with the wall of the fruit. Upon soaking the seed in water it is possible to remove the *seed coat* or *spermoderm*. This done, the embryo will be exposed. The *two fleshy cotyledons* are first seen. Upon spreading these out, convex sides down, the rest of the embryo, consisting of a thin leafy structure surrounding a bud

FIG. 241.—Stages in the germination of the common bean (*Phaseolus vulgaris*). Showing nutation and hyponasty. (After Atkinson.)

and called the *plumule*, the *caulicle* or rudimentary stem, and in line with the latter, the *radicle*, or rudimentary root, will be seen.

HISTOLOGY OF THE LIMA BEAN SEED.—In transverse sections, the following microscopic structure will be evident:

1. SPERMODERM of three regions, viz.; Palisade cells, Column cells, and Spongy Parenchyma. The *palisade cell layer* is composed of vertically elongated, thick-walled cells which are covered on their outer faces by a clear glistening cuticle. The *column cells*, found forming a layer directly beneath the palisade zone, are hour-glass-shaped.

The *Spongy parenchyma* forms a zone of several layers of thin walled parenchyma cells, the cells of the outer and inner layers being considerably smaller than the middle layers.

2. EMBRYO, the two cotyledons of which make up the greatest bulk. These are composed of an *epidermis* covering over a region of *mesophyll*. The mesophyll is constituted of moderately thick-walled cells which contain ellipsoidal and kidney-shaped starch grains up to 65μ in length. A conspicuous branching cleft will be seen in the larger grains.

In the Lima Bean, the nourishment is stored in the embryo during the growth of the seed. It is, therefore, *exalbuminous*.

GERMINATION OF DICOTYL SEEDS.—Following the imbibition of water and swelling of the seed contents, in dicotyledonous plants the *hypocotyl* is generally the first part of the embryo to elongate and emerge from the seed coat. Responding positively to the stimulus of gravity, it grows directly toward the center of the earth and enters the soil, becoming the *primary root*. The primary root continues to grow in the soil and ere long develops *secondary roots*. In the seeds of the Lima and other beans, the cotyledons and epicotyl are pulled above the soil by the elongation of the hypocotyl which makes a crook. In seeds of the pea and corn, etc., in which the cotyledons or cotyledon remain within the seed coat under the ground, the elongation of the epicotyl brings the aerial organs above the soil surface.

CHAPTER XV

THE CLASSIFICATION AND NAMING OF PLANTS

The classification and naming of plants belongs to that department of Botany called Taxonomy, a name which was derived from two Greek words, *taxis*, arrangement, and *nomos*, law.

PRINCIPLES OF CLASSIFICATION

The modern classification of plants is an attempt to arrange plants in groups based upon their natural relationship or genetic affinities. The basis of all modern classification alike of plants and animals is the theory of organic evolution.

Organic evolution presumes that all organisms living to-day have descended from preexisting organisms either by direct descent or by common ancestry. Sufficient evidence has been brought forth during the past century to substantiate plant evolution. The sources of this evidence have been found in part through the study of comparative embryology, anatomy and histology of plants, the study of fossil plants found in the rocks of past ages of the world's history in comparison with living forms of the present age, the comparison of life histories of present day plants, and through the more recent sero-diagnostic tests of Carl Mez and his pupils.

A system of classification based upon natural resemblances of plants related by descent is called a NATURAL SYSTEM. Any system of classification which does not take into account the natural kinships and evolutionary tendencies existing between plants is called an ARTIFICIAL SYSTEM.

A number of artificial systems were derived and used by botanists of the past. Most of them were based upon some arbitrarily chosen character such as woody stems, petals, stamens, etc.

Theophrastus (370-285 B.C.), a pupil of Aristotle, and probably the earliest of pioneers in attempting a scientific grouping of plants, classified them structurally into trees, shrubs, undershrubs,

and herbs. He was the first to distinguish between certain natural families.

Leonhard Fuchs (1501–1566), a German physician and botanist, in his *Stirpium Historia* (1542), an Herbal, chose to arrange plants alphabetically by their Greek names.

Hieronymus Bock (1498–1554) published an Herbal in 1546 called *Kreuter Buch* in which he carefully described from nature many plants which he grouped under herbs, shrubs and trees.

FIG. 242.—Linnaeus, the father of modern systematic botany. (After Gager.)

Caesalpinus (1519–1603) whose real name was Andrea Cesalpino, an Italian physician and botanist, was dissatisfied with the alphabetical arrangement and the narrowness of the earlier artificial systems which were largely restricted to medicinal plants. He attempted in his *De Plantis* (1583) a classification of all plants without regard to their economic importance and based it upon their fruit and seed characters.

Gaspard Bauhin (1560–1624), a Swiss botanist, grouped plants into seaweeds, grasses, lilies, trees and shrubs.

John Ray (1627–1705), an English botanist, in 1682 introduced a method of classification which distinguished between flowerless and flowering plants. He arranged plants into two chief divisions, viz.; woody plants and herbs, and made cotyledons a basis for subdividing flowering herbs into *monocotyledons* and *dicotyledons*.

Joseph P. De Tournefort (1656–1708), an eminent French botanist, chose petals as his basis of grouping and subdivided the flowering plants into the *Apetali* or the plants without petals and the *Petalodes* or those possessing petals, distinguishing herbs and undershrubs, trees and herbs. He was the first botanist to describe genera.

The greatest of these artificial systems was that of the Sexual System of Carl von Linné, popularly called Linnaeus (1707–1778), a Swedish naturalist, who through his great works, *Systema Naturæ*, published in 1735 and *Species Plantarum*, published in 1753, did more to establish Botany on a scientific foundation than any previous investigator. Linnaeus divided plants into sexual and asexual, the former of which he designated as *Phanerogamia* or flowering plants and the latter as *Cryptogamia* or flowerless plants. His classification of the flowering or seed plants was based primarily on the number and position of the stamens, each of the larger groups being subdivided according to the number and character of the carpels. In his *Species Plantarum*, he systematically described a great list of plants giving to each species a binomial name consisting of two Latin words. This system of naming plants, called the “BINOMIAL SYSTEM,” has been followed by all scientific botanists since Linnaeus and is still in use at the present day.

The first natural system of classification was that devised by Antoine Laurent de Jussieu, a French botanist, who in 1789, issued his *Genera Plantarum*. This work formed the basis of all future natural systems. Jussieu was the first to group the genera into families, and many of his concepts on families are used with modifications to-day.

In 1813, Augustin P. De Candolle, a follower of the teachings of Jussieu, issued his *Theorie elementaire de la botanique* in which he demonstrated that the affinities of plants are to be found

through the study of the comparative embryology and structure of plant organs. He later published his famous *Prodromus* which was continued by his son Alphonse de Candolle with the aid of eminent collaborators. This work gives descriptions of genera and species of Dicotyledons and represents a modification and elaboration on the system of Jussieu.

From 1862 to 1883, George Bentham and Joseph D. Hooker, two eminent British botanists, issued their *Genera Plantarum*. This work contains a systematic arrangement of the Phanerogams with a description of all of the genera then known.

This system was followed by the systems of A. W. Eichler (Berlin 1883) and of Engler and Prantl. The latter, called "*Die natürlichen Pflanzenfamilien*," has since its appearance become the most popular guide of systematists. It is based on the belief that the absence of a perianth is a primitive condition and that Monocotyledons precede Dicotyledons.

In 1926, J. Hutchinson, of the Royal Botanical Gardens at Kew, issued the first volume of his work, "*The Families of Flowering Plants*." This is a new natural system on the seed plants and is based on the assumption that plants with sepals and petals associated with other floral and anatomical characters regarded as primitive are more ancient than plants without sepals or petals. The Monocotyledons are placed after the Dicotyledons from which they are claimed to have been derived.

In every modern natural system, every individual belongs to a species or unit of classification.

A SPECIES is a mental concept of a group of individual plants having their more stable characters in common. While the individual plants making up the group called a species may differ in the size and number of their various organs and even show small differences in the shape of their leaves or fruits, the parts of their flowers will usually be quite similar in form, number and arrangement. These more stable characters enable one to recognize any members of a given species and to distinguish it from members of other species. When an individual of a species reproduces, its offspring will show the same essential or more stable characters as its parent or parents. Thus, all the Oak trees which we commonly recognize as white oak constitute the species *Quercus alba*.

Moreover, every *species* belongs to a *genus*, every genus to a *family*, every family to an *order* or cohort, every order to a *class*, every class to a *subdivision* and every subdivision to a *division* or phylum of the Plant Kingdom.

A GENUS is a group of related species. Thus, the plants we recognize as red oak, scarlet oak, pin oak and white oak belong to the genus *QUERCUS*. The chief structure upon which the concept of a genus is based is the flower, but other organs also play a part.

A FAMILY is a group of similar and related genera. Thus, the oak genus (*Quercus*), the chestnut genus (*Castanea*) and the beech genus (*Fagus*) constitute the family called *Fagaceæ*.

An ORDER is a group of related families. Thus, the Birch Family (*Betulaceæ*), the Hazel Family (*Corylaceæ*) and the Oak Family (*Fagaceæ*) constitute the order known as *Fagales*.

It will be noted that the common family ending is *ACEÆ*, more rarely *Æ* and the order ending *ALES*.

A CLASS is a still higher group composed of related orders. Thus, the *Fagales*, *Ranales* and a large number of other orders are placed in the Class *Dicotyledoneæ*, because all of them have several outstanding characteristics in common, such as netted-veined leaves, two cotyledons in embryo, parts of their flowers in fours and fives, and open collateral bundles.

A SUBDIVISION is a group of related classes. Thus, the Class *Dicotyledoneæ* and the Class *Monocotyledoneæ*, while differing in several characters, resemble each other in possessing covered ovules and seeds and so are placed in the subdivision *Angiospermæ*. A DIVISION or PHYLUM is composed of related subdivisions. Thus, the Subdivision *Angiospermæ* or related plants having covered seeds and the Subdivision *Gymnospermæ* or plants with naked seeds agree in possessing seeds and so are placed in the Division *Spermatophyta*.

The Divisions *Spermatophyta* (seed plants), *Pteridophyta* (ferns and their allies), *Bryophyta* (liverworts and mosses) and *Thallophyta* (seaweeds, fungi, etc.) make up the KINGDOM OF PLANTS.

In some instances species are subdivided into smaller groups called VARIETIES and these in turn into SUB-VARIETIES, FORMS and STRAINS. Classes may be subdivided into SUB-CLASSES. Thus, the Class *Dicotyledoneæ* contains two subclasses, the *ARCHI-*

Division-Spermatophyta (seed plants).

Subdivision-Angiospermæ (seeds enclosed).

Class-Dicotyledoneæ (seeds with two cotyledons).

Sub-class-Archichlamydeæ (petals disjoined).

Order-Ranales (carpels disjoined, etc.).

Family-Ranunculaceæ (herbs with many hypogynous stamens).

Sub-family-Helleboreæ (petaloid sepals hooded).

Species-Aconitum Napellus (carpels, 3 or 4; etc.).

OUTLINE OF PLANT GROUPS

DIVISION OF PLANT KINGDOM			
THALLOPHYTA	Algæ	Cyanophyceæ	Blue-green algæ
		Chlorophyceæ	Green algæ
		Bacillarieæ	Diatoms
		Phaeophyceæ	Brown algæ
		Rhodophyceæ	Red algæ
	Fungi	Schizomycetes	Bacteria
		Myxomycetes	Slime molds
		Phycomycetes	Alga-like fungi
		Ascomycetes	Sac-fungi
		Basidiomycetes	Club fungi
BRYOPHYTA	Lichens	Deuteromycetes	Fungi imperfecti
		Ascolichens	Lichens
		Basidiolichens	Lichens
	Hepaticæ	Ricciales	Riccia, etc.
		Marchantiales	Marchantia, etc.
		Jungermanniales	Leafy liverworts
	Musci	Anthocerotales	Anthoceros, etc.
		Andreaëales	Black mosses
		Bryales	True mosses
PTERIDOPHYTA	Lycopodineæ	Sphagnales	Peat mosses
		Lycopodiales	Club mosses
		Selaginellales	Little club mosses
	Equisetineæ	Isoetales	Quillworts
		Equisetales	Horsetails
	Filicineæ	Ophioglossales	Adder's tongue ferns
		Filicales	Ferns
		Hydropterales	Water ferns
SPERMATOPHYTA	Gymnospermæ	Cycadales	Sago palms
		Ginkgoales	Ginkgos
		Coniferales	Conifers
		Gnetales	Gnetums and ephedras
	Angiospermæ	Monocotyledoneæ	Grasses, lilies, etc.
		Dicotyledoneæ	Willows, cherries, etc.

ALGÆ - BACILLARIEÆ
 PHAEOPHYCEÆ
 RHODOPHYCEÆ
 SCHIZOMYCETES
 MYXOMYCETES
 PHYCOMYCETES
 ASCOMYCETES
 BASIDIOMYCETES
 DEUTEROMYCETES
 ASCOLICHENS
 BASIDIOLICHENS
 RICCIALES
 MARCHANTIALES
 JUNGERMANNIALES
 ANTHOCEROTALES
 ANDREAËALES
 BRYALES
 SPHAGNALES
 LYCOPODIALES
 SELAGINELLALES
 ISOETALES
 EQUISETALES
 OPHIOGLOSSALES
 FILICALES
 HYDROPTERALES
 CYCADALES
 GINKGOALES
 CONIFERALES
 GNETALES
 MONOCOTYLEDONEÆ
 DICOTYLEDONEÆ

CHLAMYDEÆ or that group in which the petals are either distinct and separate or entirely wanting and the METACHLAMYDEÆ or SYNPETALÆ in which the petals are more or less joined to form a tubular corolla. Large families may be subdivided into SUBFAMILIES or tribes which in some works are given full family rank.

A HYBRID is a cross breed of two varieties or species, rarely of two genera.

A complete classification of the medicinal plant, Aconite, will show the principal subordinate groups of one of the divisions. (See p. 328.)

BOTANICAL NOMENCLATURE

Before Carl von Linné (Linnaeus), the great Swedish naturalist, brought forth the binomial system of nomenclature, no uniformity existed in the assignment of plant names. Among the pre-Linnean botanists there were some who designated plants by single names, others who employed sentences in naming them, some of which were quite lengthy, and a number who adhered to the practice of naming them in their own modern tongue. The result was quite obvious, a number of systems were employed and confusion prevailed among students.

SPECIFIC NAME.—According to the binomial system which has been universally adopted, every plant belongs to a species which is given two Latin names which together represent its SPECIFIC NAME. The first name is the name of the genus or GENERIC NAME, the second, the name of the species or SPECIFIC EPITHET. The generic name corresponds, in the naming of persons, to the surname or family name, while the specific epithet is analagous to the given name. Thus, the Wild Cherry SPECIES is named *Prunus serotina*, *Prunus* representing the name of the genus, *serotina* the specific epithet or kind of *Prunus*.

The application of the specific name is determined by the type specimen, *i.e.*, the specimen or more than one with which the author of the species worked.

GENERIC NAME.—The name of the genus (pl. genera) is always a noun in the singular number and must not be applied to more than one genus. Its spelling should begin with a capital letter. Genera names may be taken from any source

whatever. Some, like *Fagus* for the Beech genus, and *Acer* for the Maple, are of Latin origin. Others have been latinized from other languages. Some have been named after some therapeutic property, their roots, leaves, flowers or seeds were thought to possess; for example, *Jateorhiza*, a latinized compound of two greek words, *ιάτεια*, healing, + *ρίζα*, root, because of the healing virtues of the root. A number have had names ascribed to them because of some peculiarity of structure, color, taste, odor, behavior, habit or appearance of the plant or portion thereof.

Thus, *Eriodictyon* (from Gr. *ἔριον*, wool + *δίκτυον*, net) alludes to its woolly, netted veined leaves; *Melaleuca* (from Gr. *μέλας*, black, + *λευκός*, white) alludes to the black bark of the trunk and white bark of the branches; *Marrubium* (from Hebrew *marrob*, bitter) refers to its bitter sap; *Barosma* (from Gr. *βαρύς*, heavy + *οσμή*, odor) in allusion to its strong smell; *Epifagus* (from Gr. *ἐπί*, upon + Lat. *Fagus*, the beech) alludes to its growth on the roots of that tree; *Impatiens* (from Lat. *in*, not and *patiens*, enduring) refers to the sudden bursting of the capsules of this genus when touched; *Lycopodium* (from Gr. *λύκος*, a wolf, + *πούς*, a foot) pertains to the appearance of the shoots of this genus. Many have been named in honor of eminent naturalists or friends of these, or other noted persons. For example, *Collinsonia* was named in honor of Peter Collinson, an English botanist of the 18th century; *Dioscorea* in honor of Dioscorides, the Greek naturalist; *Paullinia* after Paullini, a German botanist of the 17th century; *Cinchona* in honor of the countess of Chinchon, who brought the bark to Europe in 1640, and *Jeffersonia*, in honor of Thomas Jefferson.

SPECIFIC EPITHET.—The specific epithets are for the most part adjectives which agree with the names of genera to which they belong in case, gender, etc. They may, however, be nouns and in a few instances consist of two nouns or a noun and an adjective. If an adjective, it should begin with a small letter, as in *Rhus glabra* and *Evonymus atropurpureus*. This holds if it is drawn from a geographical name, e.g. *Rubus allegheniensis*. When the specific epithet is a noun, it may either be a proper noun in the genitive case when it should begin with a capital, as *Garcinia Hanburyi*; or it may be a common noun in the genitive, when it should begin with a small letter, as *Grindelia cam-*

porum; or the noun may be in apposition to the generic name and so in the same case, as *Cytisus scoparius*. Names that had formerly been used for genera but since reduced to species are usually capitalized, whether originally proper nouns or not, as *Aristolochia Serpentaria* and *Anacyclus Pyrethrum*. In cases where two nouns make up the specific epithet, the first of these is in the nominative case, the second in the genitive, the two names being connected by a hyphen, as *Capsella bursa-pastoris*. The botanical name of the species yielding the drug, *Aspidosperma*, (*Aspidosperma Quebracho-blanco*) will serve as an example of the specific portion of the names being composed of a noun and an adjective. Specific epithets taken from names of persons begin with a capital.¹

VARIETY EPITHET.—Names of varieties are applied in two different ways. Either the name of the species is given and followed by the prefix *var.* before the varietal epithet, as *Chenopodium ambrosioides var. anthelminticum*; or the varietal epithet may be appended to the name of the species, as *Chenopodium ambrosioides anthelminticum*.

It frequently happens that a botanist is careless in naming a species and, without ascertaining whether the same name has been assigned to another species, applies it to his, thus causing duplication. For example, there have been two distinct species of plants named *Eupatorium laevigatum*, one a form of Joe Pye Weed, the other a species of the American tropics. In this case the name *Eupatorium laevigatum* does not tell us which species the writer or speaker refers to; it might be the tropical plant called *Eupatorium laevigatum* by Lamarck or the Joe Pye Weed named *Eupatorium laevigatum* by Torrey at a later date. Accordingly, it is necessary to add to the name of the species the author's name. In this connection it is customary to abbreviate the name of the author thus, *Lam.* for Lamarck, *Torr.* for Torrey, *L.* for Linné, *Mill.* for Miller, *Ait.* for Aiton, *Loisel.* for Loiseleur-Deslongchamps, or *Tourn.* for Tournefort.

Whenever a plant is transferred from one genus to another, it must retain its original specific epithet, unless the genus to which it is transferred already possesses a species with that epithet, in

¹ There is a tendency amongst American botanists to use the small letter in beginning the spelling of all specific names irrespective of their source.

which case a new specific name must be given it. Moreover, the name of the botanist who assigned the original specific epithet but placed it under a different genus must be placed in parenthesis between the specific epithet and the name of the botanist who later connects it with another genus. For example, we read as the definition for Purging Cassia in the National Formulary V—"The dried fruit of *Cathartocarpus Fistula* (Linné) Persoon." The significance of the name Linné in parenthesis is that he had previously given the specific epithet *Fistula* to the plant indicated but placed it under a different genus, which genus happened to be *Cassia*.

FAMILY NAME.—The names of families are designated by the name of one of their principal genera or ancient generic names with the ending *aceæ*, *e.g.*, *Ranunculaceæ* from *Ranunculus*, *Malvaceæ* from *Malva*, etc. The following names, because of long usage, are exceptions to the rule: *Palmæ*, *Gramineæ*, *Guttiferæ*, *Umbelliferæ*, *Labiataæ*, and *Compositæ*.

ORDER NAME.—Orders are generally designated by the name of one of their principal families, with the ending *ales*, *e.g.*, *Rhamnales* from *Rhamnaceæ*, *Rosales* from *Rosaceæ*. Suborders are likewise designated, but with the ending—*ineæ*, *e.g.*, *Malvineæ* from *Malvaceæ*. Other older endings may, however, be retained for these names providing they do not lead to confusion or error.

CLASS NAME.—Names of classes, subclasses, divisions and subdivisions are designated from one of their characters by words of Latin or Greek origin, some similarity of form and ending being given to those that designate groups of the same nature, as *Mono-cotyledoneæ*, *Dicotyledoneæ*; *Archichlamydeæ*, *Metachlamydeæ*; *Thallophyta*, *Spermatophyta*; *Gymnospermæ*, *Angiospermæ*.

In the case of Cryptogams, the use of old family names as *Algæ*, *Fungi*, *Lichens*, *Musci*, etc. is permissible for designating groups above the rank of family.

For further details on botanical nomenclature, the student is referred to the following articles:

International rules of nomenclature revised by the International Botanical Congress of Cambridge, 1930, 3 ed., Gustav Fisher, Jena, 1935.

An abridged version of The International rules, inaccurate in a few details, was published as a supplement to volume 72 of the [English] Journal of Botany (1934).

CHAPTER XVI

THE THALLOPHYTES

The Thallophyta or thallophytes constitute Division I of the Vegetable Kingdom. As a group they represent the most simple and most primitive plants. Many of their living representatives like the bacteria, flagellates and slime molds possess characteristics of both plant and animal so that they are believed to be near the first forms of life that existed on the earth. The greater number of the members of this division have bodies consisting of a *thallus*, a structure undifferentiated into roots, stems or leaves. Under the microscope the thallus shows neither tracheids nor tracheæ. Some of the higher forms possess structures superficially resembling and functioning like the roots, stems and leaves of higher groups. They reproduce asexually or asexually and sexually.

They include the *Schizophyta* or fission plants represented by the bacteria and blue-green algae, the *Flagellata* or flagellates, the *Myxomycetes* or slime moulds, the *Zygophyceæ* represented by the diatoms, pend silks and desmids, the *Chlorophyceæ* or green algæ, the *Charophyta* or stoneworts, the *Phaeophyceæ* or brown algæ, the *Rhodophyceæ* or red algae, the *Hyphomycetes* or fungi and the *Lichens*.

The evolutionary or phylogenetic relationship of the above groups is so incompletely understood that the forms now to be considered will be discussed under the long established physiological groups of *Algae*, *Fungi* and *Lichens*.

THE ALGÆ

The algæ (*sing.* alga) are forms of thallophytes found on land and in fresh, brackish or salt water. The lower forms consist for the most part of single cells or rows of single cells joined end to end to form filaments. The higher forms, however, possess structures which, while appearing like stems and leaves of higher

plants, are merely sheets or cylinders of many celled filaments, closely arranged and connected by protoplasmic connections. They contain chlorophyll, and so can use the CO_2 and H_2O in the same manner as higher plants, *e.g.*, in building up food and providing for their own nutrition. In addition to chlorophyll, some contain one or more additional pigments. Asexual repro-

FIG. 243.—Blue-Green Algae. A–C. *Gloeocapsa polydermatica*. A, single cell in which division has begun; B, two daughter cells resulting from fission; the cell on the left is dividing; C, colony of four granddaughter cells; D, fixed and stained cell of *Rivularia pisum*. The net-like central body is the nucleus. $\times 3000$. E, F., *Oscillatoria princeps*. E, terminal portion of a filament; F, portion of a filament showing a dead cell. $\times 540$. G, as maller species of *Oscillatoria*. H–K, *Nostoc*. H, portion of filament; h, heterocyst; I, portion of a filament with spores (sp); J, spore; K, short filament produced by germinating spore. $\times 650$. (A to C, after Strasburger; D, after Haupt; H, I, J, K, after Bornet.)

duction is accomplished by motile or non-motile spores. Sexual reproduction occurs in all but the simpler forms. The gametes may be motile or non-motile. The female gamete is lodged within an oögonium or carpogonium both of which are unicellular female sexual organs. Archegonia are absent in this group.

CLASS I.—CYANOPHYCEÆ, THE BLUE-GREEN ALGÆ

Plants which are sometimes termed *blue-green algæ*. They contain chlorophyll, a green pigment, and phycocyanin, a blue pigment, a combination giving a blue-green aspect to most of the

plants of this group. Some of them also possess *phycoerythrin*, a red pigment. Their bodies are usually covered with a gelatinous envelope which may be in layers. In many of them the protoplast is differentiated into a peripheral pigment containing region and a central colorless region. The unicellular forms reproduce by binary fission while many of the filamentous forms reproduce by the separation of one or more segments of the filament from the rest of the body which move away and develop into new individuals. These segments are called *hormogonia*. They are found everywhere in fresh and salt water and also on damp logs, rocks, bark of trees, stone walls, in the roots of Cycads, etc. Examples: *Oscillatoria*, *Glæocapsa*, *Nostoc*, *Anabæna*, *Rivularia*, and *Spirulina*.

GLÆOCAPSA.—This blue-green alga is commonly found on old, damp flower pots in greenhouses and on damp rocks and walls near springs, where it forms slimy masses. Under the microscope a mount of *Glæocapsa* will be seen to consist of isolated protoplasts and groups of protoplasts, surrounded by concentric gelatinous envelopes. Each protoplast consists of a protoplasmic mass which contains blue and green pigments. No definitely organized nucleus is apparent but chromatin in the form of granules is scattered through the protoplasm. The whole is surrounded by a cell wall which undergoes mucilaginous modification producing thus the soft gelatinous envelopes which encircle parent-, daughter-, grand-daughter- and even great-grand-daughter-cells.

OSCILLATORIA.—*Oscillatoria* is a blue-green, filamentous organism found abundantly on the surface of the mud of drains and ditches as well as in ponds where the water is foul. The filament is slender and composed of compactly arranged disc-shaped cells which are all alike, excepting the terminal ones which appear rounded off. Each filament is surrounded by a gelatinous sheath. The filaments tend to be agglomerated in thick felts or gelatinous masses and each possesses peculiar oscillating and forward creeping movements. At the time of reproduction the filament breaks up transversely into short segments or *hormogonia* each of which, by fission occurring among its cells, grows into a new filament.

NOSTOC.—Nostoc occurs on the damp ground bordering streams or in slow bodies of water as greenish or brownish, tough, gelatinous balls varying in size from a pea to a hen's egg. When one of these masses is dissected and examined microscopically, it is seen to contain, imbedded in a gelatinous matrix, numerous serpentine filaments, composed of spherical or elliptical cells loosely attached to each other in chain-like fashion. Most of

FIG. 244.—Blue-Green Algae. A, *Anabaena azollae*; B, *Anabaena inaequalis*; C, *Cylandrospermum catenatum*; h, heterocyst; s, spore; D–G, *Rivularia bullata*: D, Single filament; E, F, hormogonia; G, formation of hormogonia; H, *Merismopedia*. (A, C, from Campbell's "University Text-Book of Botany." By permission of The Macmillan Co., Publishers; B, from "A Treatise on The British Freshwater Algae" by the late G. S. West; revised by F. E. Fritsch, Cambridge Univ. Press, 1927; D–G, after Bornet, from Engler and Prantl.)

the cells are of the blue-green vegetative kind, but there occur at intervals larger cells, often devoid of protoplasm which are termed *heterocysts*. Frequently the filaments break apart on either side of the heterocyst, setting free segments of cells (*hormogonia*) which grow into new filaments.

CLASS II.—CHLOROPHYCEÆ, THE GREEN ALGÆ

Green algæ are unicellular (sometimes motile), filamentous, colonial, or sheet-like water plants characterized by the presence of solitary, or numerous chloroplasts in the cells which compose

the thallus. These chloroplasts vary considerably in form, being in some cases spiral bands, in others star-shaped, in others like a napkin ring, and in others granular. In the chloroplasts of most green algæ are *pyrenoids*, which consist of a central crystalline portion of protein (aleurone-like) surrounded by a starchy envelope of variable magnitude. These are called starch centers and the starch is frequently in the form of rounded, or angular grains. The nutrition of these algæ is autotrophic. There is a definite nucleus present, but in the cœnocytic forms the nuclei may be many within the confines of the cell wall. The motile cells have two (except in *Oedogoniaceæ* and by fusion in the *Vaucheriaceæ*) to many cilia, as likewise some of the reproductive cells. No cilia occur in the conjugate algæ. Reproduction is by cell division, the formation of zoöspores (motile cells), by zygospores produced by conjugation, by egg cell and sperm cell union (oöspores) oögamous reproduction. Green algæ live mostly in fresh water. Some live in sea water and a few in brackish water. Some are associated with fungi to form lichens.

1. ORDER PROTOCOCALES OR ONE-CELLED GREEN ALGÆ.—This order contains nearly all of the one-celled green algæ excepting the diatoms and desmids. It also contains a variety of many celled forms, the cells of which are devoid of cilia.

FAMILY PLEUROCOCCACEÆ.—*Pleurococcus vulgaris* [Fig. 245 (2)] is a one-celled green alga, millions of which, living together in colonial fashion, constitute the so-called "green stain" that is common on the north sides of tree trunks, stone walls and fences. Each organism consists of a protoplast surrounded by a cell wall of cellulose. The protoplast contains a *chromatophore* (capable of division into 2 or more chromatophores), cytoplasm and nucleus. Reproduction takes place by the protoplast dividing into two equal parts and laying down a cell wall forming two daughter-protoplasts. These may again divide to form four granddaughter-protoplasts. Still another division may occur as a result of which eight great-granddaughter-protoplasts are formed which frequently adhere to one another forming colonies.

2. ORDER VOLVOCALES.—This order comprises free-swimming aquatic forms whose vegetative cells are bi-ciliated, green, more or less spherical or compressed. Some of the organisms

like *Sphærella* [Fig. 245 (3)] and *Chlamydomonas* consist of single cells bearing a pair of cilia, while others like *Pandorina*, *Eudorina*

FIG. 245.—Various types of Green Algæ. *Ulothrix* (1); *Pleurococcus* (2); *Sphærella* (3); *Cosmarium* (4); *Rhaphidium* (5); *Vaucheria* (6); *Rhizoclonium* (7); *Hydrodictyon* (8); *Oedogonium* (9); *Botrydium* (10); *Zygnema* (11); *Draparnaldia* (12); *Zygnema* during conjugation (13).

and *Volvox* show varying degrees of colony formation. Reproduction sexual or asexual.

The organisms of this order are sometimes classified with the *Flagellates*, a group having characteristics partly like plants and partly like animals and represented by *Euglena* and many other unicellular, aquatic forms possessing flagella or cilia.

Volvox globator, a typical representative of this order, is found in fresh water pools as a tiny, hollow, spherical, green colony about $\frac{1}{50}$ to $\frac{1}{30}$ of an inch in diameter. When examined under the microscope (Fig. 246), it is found to consist of hundreds of green, more or less spherical cells, united by fine strands of cytoplasm (protoplasmic bridges), the whole being enveloped by a gelatinous sheath. The peripheral cells are provided with cilia, in order that the colony may rotate and roll through the water. In a young colony, all of the cells are alike, each consisting of a mucilaginous-like cell-wall enclosing cytoplasm, a nucleus, a chloroplastid and often a red pigment spot. In a mature colony, however, throughout the greater part of its existence, two kinds of cells may be discerned: small, sterile, vegetative cells that do not divide and from 10 to 12 larger vegetative ones that divide to form new colonies. The latter slip inward below the level of the smaller cells and through repeated divisions form a number of ciliated cells joined by cytoplasmic threads, which in reality is a miniature colony. This then escapes to the exterior through the rupturing of the gelatinous wall of the old colony.

During autumn of the year, certain of the ordinary cells undergo differentiation, some to form sperm cells, others, egg-cells. When about three times the size of the ordinary sterile cells, the sperm cells divide repeatedly to form a cluster of elongated secondary cells [Fig. 246 (1a, a² and 5)] each of which contains an orange pigment, a red corpuscle, has an elongated beak and bears a pair of *flagella* (lash-like processes). The cluster in time separates into motile *antherozoids* [Fig. 246 (6, 7)] which finally escape into the cavity of the volvox sphere through rupture of the investing wall. The flask-shaped *egg cells* (1b, b) increase greatly in size without dividing. Each shows vacuoles, then becomes filled with a dark green pigment, becomes spherical and acquires a gelatinous envelope. It then passes

FIG. 246.—*Volvox globator*. Mature colony in center (1); sexual cells (2a); endochrome of primary cell has resolved itself into a cluster of secondary cells (1a, a^2 and 5); antherozoids (6, 7); bundle separated into component antherozoids in cavity of primary cell ($1a^3$); breaking of wall of primary cell showing escape of antherozoids into cavity of volvox sphere ($1a^4$); egg cells (1b, b); flask shaped germ (egg) cells with large vacuoles in protoplasm ($1b^2$, b^2); globular egg cells prepared to pass into cavity of volvox sphere (b^3); antherozoids collected about egg cell (3); oöspore (4). (Carpenter.)

into the cavity of the sphere where it is surrounded by numerous antherozoids (3) and is finally fertilized.

The product of this fertilization is an *oöspore* or *zygote* (4) which ere long becomes covered with an internal smooth membrane and a thicker external spinose coat. The chlorophyll within then disappears and starch and a reddish- or orange-colored oil make their appearance. Up to 40 of these oöspores have been observed in a single volvox sphere. Not long after the formation of these oöspores the whole parent colony breaks up and the oöspores fall to the bottom of the pool to pass the winter season. As early as February each oöspore germinates to form another volvox colony, which repeats the life cycle described.

FIG. 247.—*Vaucheria terrestris*, a siphon alga. *anth*, antheridium (empty); *o*, oögonia. (Gager.)

3. ORDER CONFERVALES.—In this order are included a variety of green filamentous and membranous forms some of which show sexual reproduction.

FAMILY ULOTHRICEÆ.—*Ulothrix zonata*, [Fig. 245 (1)] a typical representative of this family, is a filamentous organism found growing on stones around ponds, on rocks along the shores of lakes, in slow-moving streams, etc. Each filament is unbranched and consists of a row of short cells; one of the terminal cells, called the *rhizoid cell* is elongated and serves as an attachment structure. Each cell consists of a cell wall of cellulose enclosing cytoplasm, a nucleus and a wide, band-shaped, green chromatophore, more or less cylindrical in shape. The *chromatophore* (chloroplast) lies next to the cell wall and contains *pyrenoids* or starch-forming centers. The filament

grows in length by the fission of its various component cells. After attaining a certain size it reproduces either asexually or sexually. *Asexual reproduction* takes place by certain cells becoming altered in their protoplasmic contents, through division, to form rounded or pear-shaped *zoöspores*. Each zoöspore contains a red pigment spot and bears four cilia (protoplasmic outgrowths). The zoöspores escape into the water by lateral openings in the walls of cells containing them. They swim rapidly about, propelled by their cilia, and ere long attach themselves to various objects and grow into *Ulothrix* filaments. The *sexual method of reproduction* is effected through the production of many *gametes*, in cells of the filament. These resemble the zoöspores in shape but differ from them in being smaller and possessing but two cilia. These escape into the water and, after swimming about for a short time come together in pairs and fuse with one another. The product of the fusion of each pair of these like gametes is termed a *zygospore*. The zygospore swims about but finally comes to rest, remaining quiescent for a considerable length of time. It then enlarges and its protoplasmic content divides to form several zoöspores which, escaping from the cell, swim about for a while and finally, attaching themselves to objects, grow into filamentous *Ulothrix* organisms.

Closely related to *Ulothrix* is the *sea-lettuce*, *Ulva lactuca* (Fam. *Ulvaceæ*), a membranous, leaf-like form found attached to rocks along the shores exposed to the action of the waves.

4. ORDER CONJUGALES.—To this order belong the desmids and pond scums which are distinguished from other green algæ by presenting no motile stages in their life histories. No zoöspores are, therefore, produced by members of this order. They are all of fresh-water habit and reproduce by conjugation.

FAMILY DESMIDACEÆ.—The desmid family includes a number of genera of unicellular as well as filamentous green plants that present a variety of shapes. Each unicellular desmid [see Figs. 245 (4, 5)] is characterized by being composed of two like halves frequently separated from each other by a constriction called the *isthmus*. In each half there is a *chromatophore* containing *pyrenoids*. The *nucleus* is found in the isthmus. Reproduc-

tion is accomplished either asexually by fission or sexually by conjugation.

FAMILY ZYGNEMACEÆ.—This is a family of pond scums including the well-known genera, *Spirogyra* and *Zygnema* [Fig. 245 (11, 13)].

FIG. 248.—*Spirogyra* sp. A, terminal portion of vegetative filament; B, stages of scalariform conjugation; C, preparation for lateral conjugation; D, zygospores formed by lateral conjugation. (Gager.)

Spirogyra or *Brooksilk* is a filamentous organism found suspended or floating in masses in quiet water. Each filament, when examined microscopically, will be found to consist of more or less elongated cylindrical cells arranged end to end, the terminal cells having rounded extremities. Each cell has a cell wall

of cellulose within which is to be found a thin film of *ectoplasm*. One or more spirally shaped *chromatophores* will be seen directly within this area. Each chromatophore contains *chlorophyll* and a number of *pyrenoids*. In the center of the cell the *nucleus* is found. Fine strands of protoplasm hold it in place and run out to the ectoplasm.

Under favorable circumstances the cells of *Spirogyra* increase rather rapidly in length. Abnormally long cells are not seen, however, because the elongating cells speedily divide, forming two daughter-cells. Under the best of conditions, division may

FIG. 249.—*Spirogyra longata*. Portion of a monoecious filament showing two adjacent cells in the early stages of conjugation. This type of conjugation is called *lateral*. Ladder-like or *scalariform conjugation* of dioecious filaments is shown in figure 248. Note the pulsating or contractile vacuoles. (From Gager, after Lloyd.)

occur every night. In this way the filaments are rapidly made longer. Sooner or later they break and in this way the plant multiplies.

Spirogyra has also a process of sexual reproduction known as conjugation. This process occurs normally from March to June and July, but can be induced in the laboratory by allowing the water in the vessel in which it is growing to slowly evaporate. Two filaments arrange themselves side by side, and the cells lying opposite each other undergo internal changes so as to form *gametes* or sexual cells. Each gamete-containing cell is called a *gametangium*. Each gametangium protrudes a process or lateral branch. The tips of these fuse upon meeting each other and form a *conjugation tube* or *connecting tube*. Then the protoplast (gamete) of one cell passes over and coalesces with that in the cell

opposite. The result of the process is a new cell called a *zygospore* or *zygote*. This method is called *scalariform conjugation*. The zygospore forms a thick cell wall about itself. It is set free by decay of the walls of the old cell and falls to the bottom of the

FIG. 250.—*Vaucheria sessilis*. A, Terminal portion of a filament showing the appearance of a partition wall separating the newly formed sporangium from the rest of the filament. B, Same, after escape of zoospore (Z). C, A portion of the margin of the zoospore. D, Young plant developed from a zoospore with rhizoids (r). (From Strasburger, Noll, Schenk and Schimper; A, B, Z, after Götz; D, after Sachs from Altmann's *Algæ*; C, after Strasburger.)

water, there to undergo a resting stage until favorable conditions for growth arise.

Professor F. E. Lloyd has shown that in *Spirogyra longata*, conjugation takes place between adjacent cells of the same filament.

This second type of conjugation is called *lateral conjugation*. In this case a protrusion of the cell wall of the filament occurs opposite the partition wall of adjacent gametes forming a channel of communication between the gametes. Pulsating or contractile vacuoles, such as have been described in *Amoeba* appear in the protoplast and upon contraction propel the gamete in one cell cavity (*supplying gamete*) over into the cell cavity of the other gamete (*receiving gamete*). The fusion of these gametes results in the formation of a *zygospore*.

5. ORDER SIPHONALES (SIPHON ALGÆ).—This group is characterized by the peculiarity that the organisms constituting it possess protoplasm containing myriads of nuclei within a common filament or cell cavity not segmented by cell walls. The term *cænocyte* has been given to such structures which consist of a many-nucleated mass of protoplasm surrounded by a cell wall. Some of the siphon algæ reproduce by zoöspore formation, others by conjugation as well as zoöspore formation while *Vaucheria*, the “green felt,” stands out alone in reproducing both by the formation of a single compound zoöspore and by the production also of *oögonia* and *antheridia* with resultant fertilization (see Figs. 245 (6) and 247).

The various species of *Vaucheria* occur in water or moist soil where they form dense mats of tubular filaments. Each filament is unicellular until the time of reproduction, containing cytoplasm in which are imbedded thousands of nuclei as well as numerous chloroplasts devoid of pyrenoids, and small drops of oil. At the time of asexual reproduction a transverse wall is laid down near the tip of the filament cutting off a terminal cell or zoösporangium in which a many ciliated compound zoöspore is formed. This escapes by a break in the end of the sporangium, swims about for a brief time and then germinates, sending out one or more *Vaucheria* filaments. In some species a colorless simple or branched rhizoidal cell or organ of attachment is developed by the filament where it contacts a solid substance.

6. ORDER CHARALES (THE STONEWORTS).—FAMILY CHARACEÆ.—The highest group of green algæ, possessing forms which are differentiated into stem-like and leaf-like structures and rhizoids. Examples: *Chara* and *Nitella*.

Chara, a type of this family, is a submerged fresh-water plant which fastens itself to the muddy bottom of ponds, ditches and slow streams by means of slender filaments called rhizoids. Its plant body consists of a many noded (jointed) axis or so-called "stem" which bears whorls of slender, green, dwarf branches,

FIG. 251.—*Chara fragilis*. Habit and structure of certain vegetative parts. A, portion of plant showing long branches (a), dwarf branches (b), and rhizoids. B, longitudinal section through growing tip. C, cross section of internode, showing internodal cell and surrounding cortex. (From Mottier, adapted from Campbell.)

sometimes called leaves, at its nodes. The dwarf branches, like the main axis, possesses nodes and internodes. The nodes of these produce short abortive "leaves." Filamentous rhizoids arise from the lower node. Long branches are also found issuing from the axils of some of the dwarf branches at the nodes which repeat the structure of the main axis. Reproduction is either

asexual or sexual. Asexual reproduction is accomplished by means of tuber-like bodies borne on submerged parts or by special branches which form *rhizoids* on their lower nodes and later become separated from the parent plant. Sexual reproduction is effected through the formation of *oögonia* (female sex organs) and *antheridia* (male sex organs). These in some species are borne on the same plant; in others, on different plants. In

FIG. 252.—A, Portion of the axis of *Chara fragilis*, a stonewort. *s.* Oogonium (one-celled female sexual organ). *a.* Antheridium. *b.* Internode. *B.* Abortive "leaves." *B', B''.* Sterile "leaves." *c.* Crown of oögonium. *B.* Young stage of the same. *br.* Cortical cell. *a.* Antheridium. *sk.* Oögonium. (From Small, after Sachs.)

all cases the sexual organs are produced at the nodes of dwarf branches. The oval oögonium is borne above the spherical antheridium. The oögonium develops within itself a large *ovum* or egg. The antheridium produces within its wall numerous motile sperms. Upon the maturation of the antheridium the sperms are liberated into the water, and, propelled by their cilia, find their way to the oögonia which they enter, the one best adapted fusing with the egg in each case and fertilizing it. The resultant cell is called the *oöspore* or *zygote*. The zygote undergoes a resting stage within the oögonium. The oögonium with the enclosed zygote falls from

the plant and sinks to the bottom of the water. The zygote later germinates as a *proembryo* or juvenile form of *Chara* plant. The proembryo consists of a small shoot of two nodes and a long rhizoid. From the upper node of this proembryo, the adult form of *Chara* arises as a side shoot, and from the lower node rhizoids are formed.

CLASS III.—BACILLARIEÆ OR DIATOMS

This class comprises several thousand species of unicellular plants called *Diatoms* which are found in fresh, brackish and salt water, forming much of the diet of small animals. While

unicellular, they frequently are united in colonies. They all possess chromatophores containing chlorophyll but this green pigment is often obscured by the presence also of a brown pigment.

The most striking peculiarity of the group is the structure of the enclosing cell wall. This is in the form of a siliceous case consisting of two *valves* which fit into each other like the halves of

FIG. 253.—Two views of the diatom *Pinnularia*. A, valve view; B, girdle view, showing cell contents. (After Strasburger's *Text Book of Botany*, Macmillan Co. publishers.)

a pill box. The valves, which are beautifully sculptured with intricate patterns of lacework or of beading, are similar except that one is slightly larger than the other so as to fit over it. Diatoms vary in form, being either circular, linear, elliptical, cylindrical, rhomboidal, triangular, stellate, wedge-shaped, or fan-shaped, etc. Some are borne on the ends of stalks, while others are held in gelatinous masses. Their siliceous skeleta are deposited constantly on the floor of ponds, rivers, lakes and

seas, often in such abundance as to form Diatomaceous earths or Kieselguhrs (Siliceous Earths). Huge geological deposits of this material have been found in various parts of the world. One of the most remarkable for extent as well as for the number and beauty of the species contained in it is that at Richmond, Virginia. It is in many places 24 to 40 feet in depth and extends for many miles. Another deposit at Lompoo, California, is

claimed to cover an area of 12 square miles with an average depth of over 100 feet. Many of the diatomaceous earths are useful as absorbent and polishing powders.

Terra Silicea Purificata (Purified Siliceous Earth) is a powder consisting of the frustules and fragments of diatoms which has been purified by boiling with diluted hydrochloric acid, washed and calcined.

Many diatoms exhibit a gilding or jerky motion. This is stated to be caused by the streaming movement of their cytoplasm.

Diatoms exhibit two modes of reproduction, viz., fission and formation of an *auxospore*.

The more common method is that of fission but this is peculiar for these plants. The cell-contents within the siliceous case separate into two distinct masses and the valves separate slightly from each other. As the two daughter-masses become more and more developed, the valves of the parent-cell are pushed more widely apart. Each of the two masses secretes for itself a new valve on the side opposite to the original valve. When the process is completed the girdle of the parent-diatom separates and the two daughter-diatoms thus become independent plants. Each of these possesses one of the

FIG. 254.—Two species of Diatoms. To left, *Diatoma vulgare*, a, side view of frustule; b, frustule undergoing division. To right, *Grammatophora serpentina*: a, front and side views of single frustule; b, b, front and end views of divided frustule; c, frustule about to undergo division; d, frustule completely divided. (After Carpenter.)

parent valves and a second, which it has formed itself, more or less parallel to the first.

In a number of species, repeated fission results in the formation of succeeding smaller and weaker individuals. This process, however, goes on only for a certain number of generations until the decrease of size has reached a limit for the species, when the plant is rejuvenated by the formation of an *auxospore*. This may be formed with or without the conjugation of two parent-protoplasts. In either case the auxospore resulting undergoes a resting stage after which it develops new valves. The newly formed diatom is then several times the size of the individual or individuals which contributed to its formation and is endowed with renewed vigor for growth and division.

FIG. 255.—*Licmophora flabellata*, a diatom with wedge-shaped frustules borne on the ends of stalks, producing a fan-like arrangement. (After Carpenter.)

CLASS IV.—PHÆOPHYCEÆ, THE BROWN ALGÆ

Mostly marine forms showing great diversity in the form of their vegetative bodies. They occur for the most part in salt water between the high and low tide marks. Some of them are always submerged. Their bodies are usually fixed to some support in the water by means of a holdfast, and are often highly differentiated both as to form and tissues. Some reach a hundred feet or more in length as, for example, *Macrocystis*, a giant kelp, which grows in the Pacific Ocean off the coast of California. They all contain the brown pigment called *phycophæin* or *fucoxanthin* and the green pigments, *chlorophyll*, both of which are present in their chromatophores. A yellowish pigment called *phycoxanthin* has also been isolated from some of the species. Many of the kelps and rock-

weeds belonging to this class have long been sources of iodine, potash and sodium.

There are four orders of brown algae, viz., the *Phæosporales*, *Laminariales*, *Dictyotales* and the *Fucales*. Types of three of these will be considered.

A FILAMENTOUS BROWN ALGA, *ECTOCARPUS SILICULOSUS*.—*Ectocarpus*, a representative of the Order *Phæosporales*, occurs as

FIG. 256.—Fossil diatoms: *a, a, a*, *Gaillonella procera* and *G. granulata*; *b, b*, *Surirella plicata*; *c*, *Surirella craticula*; *d, d, d*, *Gaillonella (Melosira) biseriata* (side view); *e*, *Gomphonema gracile*; *f*, *Cocconema fusidium*; *g*, *Tabellaria vulgaris*; *h*, *Pinnularia dactylus*; *i*, *Pinnularia nobilis*; *k*, *Surirella caledonica*; *l*, *Synedra ulna*. (After Carpenter.)

tufts of branching filaments, each of which is many celled. These tufts are found on eelgrass or other algæ as well as attached to pilings of wharves in salt water. It is a striking illustration of the simplest form of brown algæ and serves to show the beginning of a more complex form of reproduction than that observed in the forms studied up to this time. On examination of a filament we find it to consist of many cells joined end to end. A single cell has a cell wall of cellulose. Just within the cell wall there is a layer of protoplasm. Going toward the center we find an irregular chromatophore containing a brown pigment called *phyco-phæin*. From certain cells of the filament spherical to elliptical

sporangia (spore cases) arise, which are unicellular. They contain numerous biciliate *zoöspores*, which escape into the sea water, move about and later develop into new *Ectocarpus* plants. Along the filaments several branches will be seen. Some of these have undergone division into several cells and these again into still smaller cells until many-celled chambers have resulted, which are called *plurilocular gametangia*. Each cell of a plurilocular

FIG. 257.—*Ectocarpus siliculosus*, a filamentous brown alga. A. Unicellular sporangia, one containing zoöspores, the other empty; z, a zoöspore. B. Plurilocular gametangia, the larger mature, the smaller still showing the outlines of the original cells in the branch from which it arose. C, Union of two gametes to form the zygospore.—C, after Oltmanns. (From Bergen and Davis *Principles of Botany*.)

gametangium contains a *gamete* or sexual cell, which resembles in many details a zoöspore. When the gametangium matures these gametes are discharged into the salt water. They fuse together in pairs and form *zygospores*. Each zygospore undergoes a resting stage and upon the advent of favorable conditions develops into a new *Ectocarpus* filament.

THE KELPS.—To this group (Order *Laminariales*) belong the largest living representatives of the Algæ. They are represented

by the Devil's Aprons or *Laminaria* and the Sea Palms (*Postelsia*) which grow on rocks along the North Atlantic coast and by the giant kelps *Macrocystis* (Fig. 258) and *Nereocystis* (Sea Otter's Cabbage) found along the Pacific coast of the United States.

Laminaria digitata is an abundant kelp off the coast of New England where it grows attached to rocks by a holdfast consisting of a whorl of rootlet-like structures. From this arises a solid cylindrical stipe bearing on its end an oval or lance-shaped

FIG. 258.—A giant Kelp (*Macrocystis*), one of the Brown Algæ. (From U.S.D.A. Bur. Soils.)

blade, the latter becoming split into finger-like segments at its summit. *Postelsia* or the Sea Palm has a holdfast of several whorls of stubby rootlet-like structures and a rigid stipe about a foot in length, the latter terminating in a number of lanceolate blades.

Because of their rich content of potassium salts, the kelps are used extensively as fertilizer. They also contain a fair amount of iodine and represent one of the chief sources of this drug.

THE ROCKWEEDS.—To this group (Order *Fucales*) belong the brown seaweeds which cover many of the rocks along our North

Atlantic shores and mussel beds of our inlets. They all possess floats or bladder-like structures containing carbon dioxide and oxygen which aid in buoying up the plants in the salt water when submerged. Common forms occurring along the eastern coast of North America are *Fucus*, *Ascophyllum* and *Sargassum*. The last named is called the Gulf Weed because it is often found floating freely in the gulf stream. Large numbers of these plants often accumulate in quieter areas of the ocean where they form the so called "Sargasso Sea."

FUCUS VESICULOSUS (THE BLADDER WRACK).—This form, a brown alga, occurs as a flat thallus, which forks repeatedly, a type of branching called dichotomous. It grows near the surface of sea water, attached to rocks or to mussels along banks by means of a basal, *disc-shaped holdfast*. In the upper branches of the thallus are to be found *air-bladders* which are more or less spherical and usually in pairs. The tips of old branches become swollen and are termed *receptacles*. They are dotted over with minute warts, each having an aperture at its summit which leads into an enclosed cavity within the receptacle. This cavity is called the *conceptacle*. Within the conceptacles of the male plants of the species the *antheridia*, or male sexual organs, are formed while in the conceptacles of female plants the *oögonia* or female sexual organs, are produced. The conceptacles also con-

FIG. 259.—Kelps from the North Atlantic. A, The simple type of *Laminaria*, some of whose species grow to be thirty or more feet long. B, The digitate type (*Laminaria digitata*) which is never very long, but is broad at the base. (From Bergen and Davis *Principles of Botany*.)

tain numerous, branching, protective filaments called *paraphyses*, which arise from the cells lining the cavities. The antheridia are found as outgrowths of some of these paraphyses and produce

FIG. 260.—*Fucus vesiculosus*. A, portion of plant showing habit. B, diagram of cross section of thallus; r, outer rind; m, loose inner pith or medulla. C, cross section of a fruiting receptacle. D, longitudinal section of a growing point taken at right angles to the flat surface, showing the pit at the bottom of which is the apical cell. (A, B, C, after Mottier; D, from King's chart.)

sperms or male sexual cells. The *oögonium* is a large, globular, stalked cell and produces eight *eggs*, each of which is a female sexual cell. The eggs and sperms escape into the sea water. The eggs float and are surrounded by myriads of sperms. One sperm, only, gains an entrance, after which its nucleus fuses with

that of the egg to form an oöspore. The oöspore at once develops into a new *Fucus* plant.

The drug *FUCUS* consists of the dried thallus of *Fucus vesiculosus*, *Fucus serratus*, *Ascophyllum nodosum* or *Halidrys siliquosa* (Fam. *Fucaceæ*). *Fucus* has been used in the treatment of obesity and goiter.

CLASS V.—RHODOPHYCEÆ, THE RED ALGÆ

A greatly diversified group comprising the majority of marine algæ but represented also by some fresh-water forms of which *Batrachospermum* and *Lemanea* are examples. The marine red algæ are generally found at or just beyond the low water mark. Their vegetative bodies vary from simple branching filaments through all gradations to forms differentiated into holdfasts and branching stem- and leaf-like structures. It has been observed that many of the higher types are composed of numerous filaments which are arranged so closely and connected so intimately by protoplasmic processes as to superficially resemble the tissues of plants of higher domain. Their color may be red, purple, violet, or reddish-brown or even green. The red color of most of them is due to the presence of *phycoerythrin*, a red pigment, which is found in the chromatophores with, but frequently masking, the chlorophyll. *Phycocyanin*, a blue pigment occurs in the chromatophores of some red algae. In most of them several chromatophores occur in each cell. These are usually small and oval in shape. Starch is not formed but rather another carbohydrate staining red with iodine.

Reproduction is asexual by spores or sexual by spermatia functioning like sperms and by eggs. The spores and gametes are devoid of cilia.

SUB-CLASS FLORIDEÆ.—To this group belong plants which consist of branching filaments or filaments closely bound by an intercellular substance and plasmodesma into dichotomously branched, cylindrical or ribbon like thalli. Some of them have their bodies composed of cells all alike while others show differentiation into an outer chlorenchymatous and an inner conduc-

FIG. 261.—*Fucus vesiculosus*. Receptacle cut transversely. *c*, conceptacle, *T*, cellular thickness. (Magnified.) (After Thuret.)

tive tissue. Several rounded or flattened chromatophores occur in each cell.

Reproduction may be accomplished vegetatively by fragmentation and by budding. Asexual reproduction is commonly by means of *tetraspores* which are formed in *tetrasporangia* that occur on the ends of lateral filaments or in subepidermal cells.

FIG. 262.—*Fucus vesiculosus*. Transverse section through outer region of a portion of the receptacle of male plant showing a male or antheridial conceptacle and adjacent tissues. $\times 50$. *an*, antheridium; *os*, ostium or opening of conceptacle through which paraphyses are seen protruding; *par*, paraphyses protruding from another conceptacle, the latter concealed by surrounding tissue; *b*, compact pseudoparenchyma; *ps*, loose pseudoparenchyma with mucilaginous cells.

Sexual reproduction is by union of the nuclei of *spermata* and *ova*. The female sexual organ is called a *carpogonium*, the male sexual organ an *antheridium*. The carpogonium consists of a basal cell which contains the ovum and a terminal filament which ends in a hair-like portion called the *trichogyne*. The contents of each mature antheridium as it emerges from the organ is termed a *spermatium*. It is a spherical, non-motile male cell. The spermata cling to the trichogyne and the nucleus of a spermatium

enters it and passes to the ovum in the basal cell where fertilization is effected. The zygote forms filaments called *gonimoblasts* which give rise to *carpospores* or fruits called *cystocarps* which contain carpospores.

Chondrus crispus and *Gigartina mamillosa* (Fam. *Gigartinaceæ*) when dried and bleached, constitute the official drug CHONDRUS,

FIG. 263.—*Fucus vesiculosus*. Transverse section through the outer region of a portion of the receptacle of a female plant showing an oögonial conceptacle and adjacent tissues. $\times 50$. o, oögonium within a conceptacle; p, paraphyses; ep, epidermis; co, cortex; ps, pseudoparenchyma of irregular cells having mucilaginous walls.

Irish Moss or Carragheen. Both are purplish-red in color. Each consists of a dichotomously branched thallus the lower portion of which is differentiated as a stipe or stalk; the basal portion of which, called the holdfast, clings to the rock. The upper part is several times forked and its terminal segments appear notched or bilobed. Scattered here and there over the segments of the thallus will be noted sporangia which, when mature, contain *tetraspores*. In *Chondrus crispus* the sporangia are

elliptical and embedded in the thallus near its surface, whereas in *Gigartina* they are ovate and borne on minute stalks which

FIG. 264.—*Fucus vesiculosus*. Fertilization. A, egg approached by biciliated sperms; B, sperms attached to egg and surrounding it prior to fertilization (highly magnified). (After Thuret.)

project outward from the surface of the segments. Upon the ripening of these structures the spores are discharged into the sea

FIG. 265.—*Chondrus crispus*. (After Strasburger.)

water. These sooner or later germinate into new *Chondrus* or *Gigartina* organisms.

The dried mucilaginous substance extracted from *Gelidium corneum* or other species of *Gelidium* (Fam. *Gelidiaceæ*) and closely related red algæ growing in the sea along the eastern coast of Asia and along the coast of California constitutes the drug, AGAR.

FIG. 266.—*Gigartina mamillosa*. s, fruiting body consisting of short projection bearing tiny sporangia. (After Strasburger.)

Agar is a most valuable ingredient in culture media as well as a laxative.

Rhodymenia palmata or IRISH DULSE is a purplish-red, flat, membranous, palmately cleft or dichotomous red alga growing on the tissues of other algæ along northern shores of the Atlantic between the low- and high-tide marks.

CHAPTER XVII

THE FUNGI

The fungi or *hyphomycetes* constitute a great assemblage of thallophytes which are characterized by the total absence of chlorophyll. The members of this subdivision, therefore, possess no independent power of manufacturing food materials such as starches, sugars, etc. from CO_2 and H_2O . Consequently they are either parasites, depending for their nourishment upon other living plants or animals, called *hosts*; or saprophytes, depending upon decaying animal or vegetable matter in solution. Some forms are able to live either as saprophytes or parasites while others are restricted to either the parasitic or saprophytic habit. Some of the fungi like the bacteria and the yeasts are unicellular while others like the molds and mildews, etc. have bodies in the form of thread-like filaments called *hyphæ* (sing. *hypha*) which are frequently branched. A mass of intertangled hyphæ is called a mycelium. The hypha may be unicellular or multicellular. Fungal hyphæ either ramify through decaying matter or invade the tissues of living organisms and derive nourishment therefrom. In the cases of parasitic fungi like the rusts, etc., absorbing connections are present as hyphal outgrowths which penetrate into the cells of the host plant. These are called haustoria. In the higher forms the hyphæ become consolidated into false tissues, and assume definite shapes according to the species. Of this character are the fruiting organs which constitute the above ground or exposed parts of Puff Balls, Cup Fungi, Bract Fungi, Mushrooms, etc.

There are three classes of fungi, viz.:

Class I Schizomycetes (bacteria)

Class II Myxomycetes (slime moulds)

Class III Eumycetes (true fungi)

CLASS I.—SCHIZOMYCETES (BACTERIA)

The bacteria, also called “schizomycetes” or “fission fungi,” are now regarded by leading systematists as the lowest types of plant life extant. Among them are to be found the smallest living organisms known to science.

BACTERIA are minute, unicellular, colorless, rarely weakly red or green colored, vegetable organisms destitute of chlorophyll. They serve as agents of decay and fermentation and are frequently employed in industrial processes. The curing of certain

FIG. 267.—The organic cycle. (After Gager.)

drugs and tobacco, the tanning of leather, the manufacture of vinegar, buttermilk, cheese and indigo, the making of ensilage and sauerkraut, and the retting of flax and some other textile fibers are all dependent upon fermentative processes induced by bacteria. Their form varies from spherical, rod-shaped, spiral to filamentous, the filamentous types being frequently surrounded by a capsule or sheath.

The bodies of most bacteria possess nitrogenous cell walls representing outer plasma membranes, although hemicellulose has been found in the walls of many and mucilage occurs as a

in medicine
pin
gelatinous sheath surrounding the primary wall in some forms. Cytoplasm occurs within the cell as well as chromatin, but the latter is usually diffused in fine particles through the cell protoplasm and never has been demonstrated to be condensed into a definitely organized body called a nucleus. A common method of asexual reproduction is possessed by these plants whereby the cell cleaves or splits into two parts, each of which becomes a separate and independent organism. Sexual reproduction has not been conclusively proven, although suspected by some investigators. Bacteria occur everywhere but are relatively few on high mountain peaks and beneath 3 meters of natural soil. They have even been found in the hot geyser springs where no other form of life can exist (Thermophilic bacteria).

According to the various phenomena they produce, they may be classified as follows: (a) *Zymogens* producing fermentation; (b) *Aerogens* producing gas; (c) *Photogens* producing light; (d) *Chromogens* producing color; (e) *Saprogens*, producing putrefaction; (f) *Pathogens*, producing disease; (g) *Thermogens*, producing heat.

FERMENTATION, DECAY AND PUTREFACTION.—*Fermentation* is the process which consists of the breaking down of organic substances through the action of enzymes produced by living organisms. The organic substances fermented are various, including carbohydrates, proteins and fats. *Decay* is the fermentation of proteins in the presence of oxygen. PUTREFACTION, although frequently used in the sense of the decomposition of proteins, is more properly applied to the anærobic fermentation of proteins, many of the products of which possess an offensive odor. When carbohydrates undergo fermentation, the products are useful. Bacteria cause considerable spoilage of foods and drugs owing to their power of producing enzymes.

study in medicine
BACTERIAL CULTURES.—Because of their minute size (a space the size of a pinhead may hold eight billion of them), the student commences his study of bacterial growths in colonies or cultures, each kind possessing characteristics by which they may be distinguished and differentiated. These cultures are obtained by growing the microorganisms on sterile artificial culture media usually in an incubator, the temperature of which is carefully regulated to suit the requirements of the organisms grown. The

culture media employed are both solid and liquid. Beef broth or nutrient bouillon forms the basis of most of the culture media. To this either agar agar or gelatin are added to solidify the medium (solid media) and various chemical substances and sugars for the purpose of studying the physiological characteristics of the various forms. Formulas for the preparation of various culture media and directions for the isolation of bacteria will be found in the National Formulary VI and in works of bacteriology and mycology.

FORMS OF BACTERIA.—Bacteria occur as spheres, straight rods or cork-screw like (bent rods) forms (lower bacteria) or as filaments (higher bacteria). The spherical shaped form is

FIG. 268.

FIG. 268.—Monotrichous bacteria. *Microspira comma*. (After Migula from Schmidt and Weiss and Marshall.)

FIG. 269.

FIG. 269.—Lophotrichous bacteria. *Pseudomonas synchyanea*. (After Migula from Marshall.)

termed a *micrococcus* or *coccus* (plural-*micrococci* or *cocci*). The straight rod form is called a *bacillus* (plural-*bacilli*). The cork-screw shaped form is known as a *spirillum* (plural-*spirilla*).

MOTILITY.—A large number of bacteria possess the power of independent movement when observed in a fluid medium which will permit of their locomotion. This true movement should be distinguished from the so-called "*Brownian movement*," a quivering or oscillating phenomenon shown by minute particles or bodies when freely suspended in a suitable fluid. *True movement* is due to the activity of the living cell, whereas *Brownian movement* is a purely physical phenomenon due to the bombardment of particles, dead bacteria or non-motile bacteria, by the molecules of the menstruum. In true movement the individual form moves from one part of the microscopic field to another, whereas in

Brownian movement the individual forms do not change their relative positions to any extent.

The power of locomotion which certain bacteria possess is due to the possession by these forms of fine protoplasmic extensions from their bodies which are thought to pass from the cytoplasm through the cell wall into the surrounding liquid medium. These are called flagella or cilia. They are endowed with the power of contractility, and through their vibratory motion the organism is enabled to propel itself through the fluid medium. The great majority of bacteria possessing flagella are the bacilli and spirilla. These flagella are rarely seen without special methods of treatment and staining.

FIG. 270.

FIG. 270.—Lophotrichous bacteria. *Spirillum rubrum*. (After Migula from Schmidt and Weiss and Marshall.)

FIG. 271.

FIG. 271.—Peritrichous bacteria. *Eberthella typhi*. (*Bacillus typhosus*.) (After Migula from Schmidt and Weiss and Marshall.)

The distribution of flagella varies in different forms of bacteria. Some possess a single flagellum at one pole and are called *monotrichous*. When a flagellum occurs at each pole, they are termed *amphitrichous*; when a tuft occurs at one pole, *lophotrichous*; and when they occur over the whole cell, as in the case of the typhoid bacillus, *peritrichous*.

The motility or non-motility of bacteria is best determined through the examination of a hanging drop.

NUTRITION.—Most bacteria derive their sustenance from organic substances. Those which obtain their nourishment from dead organic substances are termed saprophytic bacteria and those which obtain their nourishment from living organic material, parasitic bacteria. A number of the saprophytic forms will thrive both on living and dead organic matter and so are called

faculative parasites, while many others, like the majority of the common soil and water bacteria, are restricted to dead organic matter and so are obligate saprophytes. Most of the parasitic bacteria are faculative saprophytes, since they can be grown upon artificial culture media. The leprosy bacillus is one of the few examples of parasitic bacteria which has resisted attempts to grow it upon artificial culture media and so is an obligate parasite.

NITRIFICATION.—There occur a group of soil bacteria called nitrifying bacteria which thrive in the presence of simple inorganic salts and in the entire absence of organic matter. Some of these (Nitrosomonas and Nitrosococcus) oxidize the ammonia of the soil to nitrous acid while others (Nitrobacter) convert the nitrous acid to nitric acid. The nitric acids reacts with alkaline soil constituents to form nitrates which pass into solution in the soil water and are utilized by higher plants in their growth. This process is called nitrification.

Bacteria
+
NH₃
↓
Nitrous
↓
Nitric

DENITRIFICATION.—Denitrification is a process carried on by denitrifying bacteria including Pseudomonas fluorescens, Ps. pyocyanea, and the colon and typhoid bacilli. Many of the denitrifying organisms flourish in "sour" soil, in the absence of air, and obtain oxygen for their respiration by reducing the nitrates and nitrites to gaseous oxides of nitrogen or to free nitrogen, liberating the gas into the atmosphere.

GROWTH PRODUCTS.—The products resulting from bacterial growth include pigments, enzymes, acid and alkali, putrefactive substances, ptomaines and toxins.

REPRODUCTION.—Bacteria multiply and reproduce themselves by cleavage or fission. A young individual increases in size up to the limits of the adult form, when by simple cleavage at right angles to the long axis, the cell divides into two individuals.

MORPHOLOGY DUE TO CLEAVAGE.—According to limitations imposed by cleavage directors, the spherical forms called micrococci or cocci assume a chain appearance, or a grape cluster-like appearance, or an arrangement in packets or cubes having three diameters. These aggregations are the basis for the following genera of spherical-shaped bacteria:

Diplococcus (plural, diplococci), occurring in twos, as organisms producing meningitis, pneumonia and gonorrhea.

Staphylococcus (plural, *staphylococci*), from a Greek word referring to the shape of a bunch of grapes, as the organism producing golden pus.

Streptococcus (plural, *streptococci*) from a Greek word meaning chain-shaped, as the scarlet fever organism.

Sarcina, package-shaped or cubical, as the *Sarcina lutea*, a common saprophyte.

FORM OF CELL GROUPS AFTER CLEAVAGE.—The individual bacteria after cleavage may separate, or cohere. The amount of cohesion, together with the plane of cleavage, determines the various forms of the cell groups. Thus, among the cocci, diplo- or double forms may result giving rise to distinguishing morpho-

FIG. 272.—Division forms of micrococci. *a*, Diplococcus, perfect form with flattened opposed surfaces (*gonococcus*), lanceolate form (*pneumococcus*); *b*, streptococcus; *c*, consecutive fission yielding a tetrad; *d*, *sarcina* form resulting from division of a tetrad; *e*, staphylococcus. (From Marshall after Novy.)

logical characteristics. Similarly, among the bacilli, characteristic forms result as single individuals and others which form chains of various lengths.

SPORULATION.—A large number of bacteria possess the power of developing into a resting stage by a process known as sporulation or spore formation. Sporulation is regarded as a method of resisting unfavorable environment. This is illustrated by the anthrax bacilli which are readily killed in twenty minutes by a 10 per cent. solution of carbolic acid, and able, when in the spore condition, to resist the same disinfectant for a long period in a concentration of 50 per cent. While the vegetative forms of anthrax show little more resistance against moist heat than the vegetative form of other bacteria, the spores will withstand the action of live steam for as long as ten to twelve minutes or more.

Whenever the spores are brought into favorable condition for bacterial growth, as to temperature, moisture and nutrition, they

return to the vegetative form and then are capable of multiplication by fission in the ordinary way.

Two kinds of spores occur amongst bacteria, *arthrospores* and *endospores*. Arthrospores represent entire bacterial vegetative cells which lose some of their water and thicken their walls and thus become resting cells. Endospores or true spores of bacteria are formed within the cell. They have the power of resisting a greater heat than the arthrospores.

RAPIDITY OF GROWTH AND MULTIPLICATION.—The rapidity with which bacteria grow and multiply is dependent upon species and environment. The rapidity of the growth is surprising. Under favorable conditions they may elongate and divide every

FIG. 273.—Spores and their location in bacterial cells. (Marshall after Frost and McCampbell.)

twenty or thirty minutes. If they should continue to reproduce at this rate for twenty-four hours, a single individual would have 17 million descendants. If each of these should continue to grow at the same rate, each would have in twenty-four hours more, 17 million offspring, and then the numbers would develop beyond conception. However, such multiplication is not possible under natural or even artificial conditions, both on account of lack of nutritive material and because of the inhibition of the growth of the bacteria by their own products. If they did multiply at this rate, in a few days there would be no room in the world but for bacteria.

CHEMICAL COMPOSITION OF BACTERIA.—The quantitative chemical composition of bacteria is subject to wide variations, dependent upon the nutritive material furnished them. About

80 to 85 per cent. of the bacterial body is water; proteid substances constitute about 50 to 80 per cent. of the dry residue. When these are extracted, there remain fats, in some cases wax, in some bacteria traces of cellulose appear, and the remainder consists of 1 to 2 per cent. ash.

FIG. 274.—Spore germination. *a*, direct conversion of a spore into a bacillus without the shedding of a spore wall (*B. leptosporus*); *b*, polar germination of *Bact. anthracis*; *c*, equatorial germination of *B. subtilis* (hay bacillus); *d*, same of *B. megaterium*; *e*, same with “horse-shoe” presentation. (From Marshall after Novy.)

The proteids consist partly of nucleo-proteids, globulins, and protein substances differing materially from ordinary proteids. Toxic substances known as *endotoxins*, to distinguish them from the *exotoxins* or bacterial poisons secreted by certain bacteria like

FIG. 275.—Types of bacilli. (After Williams.)

the tetanus and diphtheria organisms during the process of growth, also occur.

MORPHOLOGICAL CLASSIFICATION OF BACTERIA*

THE LOWER BACTERIA

The lower bacteria comprise most of the forms which produce disease and the majority of the forms which are used in the arts

* No strictly morphological system of classification for the bacteria will subserve the purpose of properly identifying species of this group owing to a deficiency of structural characteristics for the separation of so many species. For the main divisions and some of the smaller details, the application of a morphological key will be sufficient, but cultural tests for their physiological phenomena and staining reactions are essential in working out the identity of a specific bacterial organism.

and industries. They are grouped under the first 3 families, viz., *Coccaceæ*, *Bacteriaceæ* and *Spirillaceæ*.

FIG. 276.—Types of spirilla. (After Williams.)

FAMILY I.—COCCACEÆ.—Cells in their free condition globular, becoming but slightly elongated before division. Cell-division in one, two or three directions of space.

A. Cells without Flagella.

1. Division only in one direction of space forming an aggregation resembling a chain of beads—*Streptococcus*.

2. Division in two directions of space forming an aggregation resembling a cluster of grapes—*Staphylococcus*.

3. Division in three directions of space forming a package-shaped or cubical aggregation—*Sarcina*.

B. Cells with Flagella.

1. Division in two directions of space—*Planococcus*.

2. Division in three directions of space—*Planosarcina*.

FAMILY II.—BACTERIACEÆ. Cells longer than broad, generally two to six times, straight or only with an angular bend, never curved or spiral, division only at right angles to axis or rod; with or without flagella and endospores.

FIG. 277.—*Crenothrix polyspora*, an iron bacterium, which oxidizes iron causing a deposit of iron hydroxide to accumulate in water pipes where it may cause trouble. (After Migula from Schmidt and Weiss and Marshall.)

1. Flagella and endospores absent—*Bacterium*.

2. Flagella and endospores present.

(a) Flagella over entire surface—*Bacillus*.

(b) Flagella at one end—*Pseudomonas*.

FAMILY III.—SPIRILLACEÆ.—Cells curved or spirally bent, generally motile through polar flagella.

1. Cells stiff, not flexible.

(a) Cells without flagella—*Spirosoma*.

(b) Cells with one, very rarely with two polar flagella—*Microspira* (*Vibrio*).

(c) Cells with a bundle of polar flagella—*Spirillum*.

THE HIGHER BACTERIA

FAMILY IV.—MYCOBACTERIACEÆ.—Cells short or long, cylindrical or clavate-cuneate in form, without a sheath surrounding the chains of individuals, without endospores, with true dichotomous branching.

FIG. 278.—Sulfur bacteria (*Beggiatoa alba*). (After Winogradsky from Schmidt and Weiss and Marshall.)

A. In cultures possessing the characters of true bacteria. Growth on solid media smooth, flat, spreading. Rod with swollen ends, or cuneate or clavate forms—*Corynebacterium*.

B. Cultures on solid media raised, folded or warty. Generally short slender rods, rarely short branched. Take the tubercle stain—*Mycobacterium*.

FAMILY V.—CHLAMYDOBACTERIACEÆ.—Thread-like, composed of individual cells, surrounded by a sheath. Simple or with true branching. Ordinary vegetative growth by division in only one direction of space, *i.e.*, at right angles to the longer axis.

A. Cell contents without sulfur granules.

1. Filaments unbranched.

(a) Cell-division only in one direction of space.

(b) Cell-division in gonidial formation in three directions of space—*Actinomyces* (*Streptothrix*).

*Marine forms with cells surrounded by a very delicate hardly discernible sheath—*Phragmidiothrix*.

**Fresh-water forms with easily discernible sheath—
Crenothrix. (*Iron bacteria*.)

2. Filaments branched.

B. Cell contents with sulfur granules—*Thiothrix*.

SOME BACTERIA PRODUCING DISEASE IN MAN OR THE LOWER ANIMALS

ORGANISM	DISEASE
<i>Alcaligenes melitensis</i>	Malta fever
<i>Actinomyces bovis</i>	Actinomycosis in cattle
<i>Bacillus anthracis</i>	Anthrax
<i>Clostridium chauvei</i>	"Blackleg" of cattle
<i>Clostridium tetani</i>	Tetanus or Lock-jaw
<i>Clostridium welchii</i>	Gas-gangrene
<i>Corynebacterium diphtheriæ</i>	Diphtheria
<i>Diplococcus pneumoniae</i>	Lobar pneumonia
<i>Eberthella typhi</i>	Typhoid fever
<i>Hemophilus pertussis</i>	Whooping cough
<i>Hemophilus influenzae</i>	Influenza
<i>Mycobacterium lepræ</i>	Leprosy
<i>Mycobacterium tuberculosis</i>	Tuberculosis
<i>Neisseria catarrhalis</i>	Catarrh
<i>Neisseria intracellularis</i>	Cerebrospinal meningitis
<i>Neisseria gonorrhoeæ</i>	Gonorrhoea
<i>Pasteurella pestis</i>	Bubonic plague
<i>Pfeifferella mallei</i>	Glanders
<i>Rickettsia prowazeki</i>	Typhus
<i>Staphylococcus aureus</i>	Boils, abscesses, carbuncles
<i>Streptococcus scarlatinæ</i>	Scarlet fever
<i>Streptococcus erysipelatis</i>	Erysipelas
<i>Salmonella paratyphi</i>	Paratyphoid fever (A)
<i>Salmonella schottmulleri</i>	Paratyphoid fever (B)
<i>Vibrio comma</i>	Cholera

(Spirochaetales, intermediate forms between bacteria and protozoa)

<i>Borrelia recurrentis</i>	Relapsing fever (European)
<i>Borrelia duttoni</i>	Relapsing fever (African)
<i>Borrelia vincenti</i>	Vincent's angina
<i>Treponema pallidum</i>	Syphilis or lues

Some Bacteria Producing Disease in Plants

ORGANISM	DISEASE
<i>Actinomyces Myricarum</i>	Tubercles upon and lesions within <i>Myrica</i> and <i>Comptonia</i>
<i>Bacterium tumefaciens</i>	Crown gall
<i>Bacterium savastanoi</i>	Olive knot
<i>Bacillus amylovorus</i>	Pear blight
<i>Bacillus Solanacearum</i>	Wilt of Solanaceae
<i>Bacillus tracheiphilus</i>	Wilt of Cucurbits
<i>Pseudomonas juglandis</i>	Walnut blight
<i>Pseudomonas Stewarti</i>	Wilt of Sweet Corn.

The *iron bacteria* utilize iron salts forming iron oxide which is deposited in a sheath around their walls. They are responsible for the rusty appearance of some water.

FAMILY VI.—BEGGIATOACEÆ (Sulfur Bacteria).—Thread-like, without a capsule, but with an undulating membrane. Cell contents show sulfur granules.

Threads apparently not separated, septa only faintly visible with iodine staining. Colorless or faintly rose-colored—*Beggiatoa*.

The *sulfur bacteria* oxidize the hydrogen sulfide of sulfur compounds, liberating sulfur in the form of granules within the cell.

PATHOGENIC BACTERIA.—Bacteria which produce diseases in living organisms are called *pathogenic bacteria* or *pathogens*. A list of some of the more important of these and the names of the disease caused are shown on p. 373.

CLASS II.—MYXOMYCETES, OR SLIME MOLDS

Terrestrial or aquatic organisms, frequently classified as belonging to the animal kingdom and found commonly on decaying wood, leaves, or humous soil in forests. Their vegetative body consists of a naked, multinucleated mass of protoplasm called the *plasmodium*, which has a creeping and rolling amœboid motion, putting out and retracting regions of its body called *pseudopodia*. The size of the plasmodium varies from a ten-cent piece to several square feet of surface. It is net-like, the net being of irregular dimensions. Like the amœba the outer portion of the plasmodium is clear and watery and known as the *ectoplasm*, the inner portion is granular and called the *endoplasm*. Like the amœba and unlike other plants, this slimy body engulfs solid food by means of its pseudopodia instead of admitting it in solution. It is extremely sensitive to light being negatively heliotropic, *i.e.*, turning away from the sun's rays. At the time of reproduction, the plasmodium creeps to the surface. The whole plasmodium then forms one or more fructifications. These fructifications vary from cushion-like masses (*æthallia*) to more elevated bodies in which the net-like structure of the plasmodium is preserved (*plasmodiocarps*) to stalked *sporangia*

(spore cases). All of the fructifications, however, produce *spores*. During wet weather amoeboid protoplasts (*swarm spores*) escape from the spores, each developing a single *cilium* and moving actively about. In time the cilia disappear and these swarm spores coalesce in smaller then larger groups to form a plasmodium.

FIG. 279.—Slime molds. A, B, *Comatricha nigra*. A, Sporangium, natural size; B, capillitium, 20/1; C, E, *Stemonitis fusca*; C, sporangium, natural size; D, and E, capillitia, 5/1, 20/1; F, H, *Enerthema papillatum*, F, unripe; G, mature sporangium, 10/1; H, capillitium, 20/1. (C, D, after nature. A, F, G, H, after Rostafinski; B, E, after de Bary in *Die natürlichen Pflanzenfamilien* I. 1, p. 26.)

CLASS III.—EUMYCETES (TRUE FUNGI)

The True Fungi represent a large class of chlorophyllless plants in which are included plants ordinary called fungi, such as the molds, water-molds, yeasts, blights, mildews, rusts, smuts, toadstools, puffballs, etc. In nutrition, they are either saprophytes or parasites or may be adapted to both modes of existence. Most of them have a vegetative body called a mycelium consisting of filamentous hyphæ. This class is subdivided into the following sub-classes:

Sub-Class I. Phycomycetes or Alga-like Fungi—molds, water-molds, blights, mildews, rots.

Sub-Class II. Ascomycetes or Sac-Fungi—yeasts, blue and green molds, powdery mildews, cup-fungi, morels, ergot, etc.

Sub-Class III. Basidiomycetes or Club-Fungi—smuts, rusts, mushrooms, puff-balls, earth stars, etc.

Sub-Class IV. Deuteromycetes or Imperfect Fungi—fungi whose life histories are incompletely known, as celery-blight, trichophyton, monilia, oidium, etc.

THE PHYCOMYCETES, OR ALGA-LIKE FUNGI

The Phycomycetes represent a small group of fungi showing close affinity with the green algæ. Their mycelium is composed of hyphæ which are usually unicellular up until the time of reproduction and many-nucleated. This suggests a close relation with the Spirogyra group of green algæ. Their sexual organs are likewise similar in structure. Transverse septa appear in the hyphæ upon the formation of reproductive organs which separate these structures from the vegetative hyphæ. They reproduce by spores as well as by sexual reproduction.

Those Phycomycetes in which sexual reproduction is brought about through the union of like gametes belong to the order called *Zygomycetales* which includes the black bread molds and their kin; those in which sexual reproduction takes place through the union of unlike gametes belong to the *Oömycetales* or order of the water molds and their allies.

CULTIVATION OF MOLDS.—Molds and other fungi will grow on most bacteriological culture media but do better if some sugar is added. They may also be grown on moistened slices of bread, on slices of potato or carrot, or on decoctions of prunes, potatoes, manures, peas, beans, etc. Beerwort is also a good substratum for their growth. The liquid media may be made solid by the addition of agar. An excellent medium for the growth of molds and most other fungi in the laboratory is a modification of Czapek's medium, the formula for which follows:

Sucrose.....	30.0	Gm.
Sodium Nitrate.....	2.0	
Dibasic Potassium Phosphate.....	1.0	
Magnesium Sulfate Crystals.....	0.5	
Potassium Chloride.....	0.5	
Ferrous Sulfate.....	0.01	
Water.....	1000.0	cc.
For solid medium, add Agar.....	15.0	Gm.

Slide cultures are of great value in the study of molds since they permit the study of the aerial and submerged hyphæ

which are often tangled in mounts prepared by transferring the mycelium with a forceps or needle. These may be prepared by the Henrici method as follows: Clean large rectangular cover glasses and deposit a drop of de Khotinsky cement or melted sealing wax on each end; spreading it out with a small hot iron file to form a layer 5 mm. wide. Heat a slide in a flame and place the cover glass on it with the cement side down. The slide should be just hot enough to soften the cement rather than to liquefy it. The culture chamber thus prepared has a space between the cover slip and slide about 1 mm. deep. Melt a tube of Czapek's agar, cool and inoculate with spores of the mold to be examined. With a sterile pipette transfer some of the inoculated agar to the edge of the cover glass and allow it to run under until part of the space has been filled. The slides are then placed in a clean glass staining jar containing moistened filter paper in the bottom and the lid sealed with vaseline, and the slides incubated until growth has occurred.

ORDER ZYGOMYCETALES

THE BLACK BREAD MOLD.—RHIZOPUS NIGRICANS (*Mucor stolonifer*), commonly known as "Black Mold" or "Black Bread Mold," is frequently found on bread, jellies, syrups, acetic pharmaceutical extracts and other substrata, where it forms a dense thready mycelium bearing numerous black, tiny spore cases. The source of this mold is the spores, which are found in the air or water with which the attacked substratum is in contact. Each of these, upon germinating, sprouts out and forms a vegetative mycelium consisting of three kinds of hyphæ, viz.: *rhizoidal* or *submerged hyphæ*, *sporangiohores* or *aerial hyphæ* and *stoloniferous hyphæ*. The branching rhizoidal hyphæ penetrate the substratum and secrete a diastatic ferment that changes the water insoluble carbohydrate materials into a soluble sugar which passes into solution and is absorbed by their walls. This, upon entering the hyphæ, is converted into protoplasm, and so the mold increases in size. Sporangiohores or aerial hyphæ arise vertically or obliquely from a bulged-out common base of the rhizoidal hyphæ. Each of these when mature bears upon its summit a spheroidal *sporangium* containing numerous small, brownish, multinucleate

spores called *endospores*. The wall of the sporangium is beset with asperites of calcium oxalate. Springing from the base of the sporangiophores or aerial hyphæ, one or more stoloniferous hyphæ traverse a portion of the surface of the substratum, and their tips, coming in contact with the substratum, swell up forming an adhesive organ or *appressorium* which branches out below into a cluster of spreading submerged hyphæ and above into several aerial hyphæ bearing sporangia. This method of growth proceeds until the entire surface of the nutritive medium is covered with a dense fluffy mycelium.

FIG. 280.—Black mold (*Rhizopus nigricans*). A, older plant; *myc*, mycelia; *sph*, sporangiophore; *sp*, sporangium; *st*, stoloniferous hypha produced by A, and giving rise at its tip to a new plant, B. Greatly enlarged. (Gager.)

Rhizopus reproduces by two methods. The most common one is that of *division by cleavage furrows*. In this **ASEXUAL METHOD**, a transverse wall is laid down in the sporangiophore near its tip. The terminal cell thus formed swells up, becoming globular in shape and its multinucleate protoplasmic contents, by progressive cleavage, become divided to form numerous spores within the wall of the *sporangium* or enlarged terminal cell of the sporangiophore. The partition wall, separating the lumen of the sporangium from that of the sporangiophore, bulges into the sporangium as a dome-shaped structure, which is termed the *columella*. Upon the ripening of the spores the wall of the spore case bursts, liberating them. These, falling upon moist nutrient substrata, germinate and ultimately form new *Rhizopus* plants.

The Cedar and Apple Rust produces aecia on the leaves of the apple and hawthorn and telia on the red cedar (*Juniperus virginiana*). The mycelium of this fungus growing in the leaves of the juniper produces the large brown galls known as "cedar apples."

ORDER 3.—AURICULARIALES.—The so-called "ear fungi" which occur on the bark of many plants, on wooden fences, etc., as auriculate growths which when young are jelly-like and brilliantly colored, when old, hard, grayish and considerably wrinkled. The earlike fruiting body is known as the *sporophore*. Its internal surface is lined with a *hymenium* or fruiting membrane consisting of numerous four-celled basidia, each of which cuts off at its tip a basidiospore. A common example of this group is the Jew's ear fungus.

ORDER 4.—TREMELLALES.—Saprophytes which live on decaying wood as moist, soft, quivering gelatinous growths becoming later dry and horny.

THE HIGHER BASIDIOMYCETES

Mostly fleshy forms characterized by one-celled basidia with generally four, occasionally six, eight or two sterigmata, each of which cuts off a basidiospore at its tip.

DIVISION A.—HYMENOMYCETES

(Hymenium or spore-bearing surface exposed)

This division of higher basidiomycete fungi comprises the following orders: Dacromycetales, Exobasidiales, Thelephorales, Clavariales and Agaricales.

ORDER 1.—DACROMYCETALES.—This order includes the "weeping fungi." One of the most common is *Dacromyces deliquescens* which occurs as a gelatinous body of bright red color on dead wood. The basidiospores are formed during a wet period and the fungus swells up in the water forming a slimy mass. In addition to basidiospores the mycelium may break up into oidiospores, if the wet period is prolonged. In consisting of slimy gelatinous masses the "weeping fungi" approach the *Tremellaceæ* but are distinguished from them in the basidium being undivided in the former and divided in the latter.

ORDER 2.—EXOBASIDIALES.—This group is found growing parasitically on shrubs especially those of the heath family. The mycelium lives in the tissues of the stems, leaves, sepals and petals and produces spongy, fleshy, yellowish or brownish galls which are popularly called "Azalea apples." The galls are edible. They are covered with a hymenium.

ORDER 3.—THELEPHORALES. These possess fruit-bodies appearing on tree trunks as flattened leathery or crust-like

FIG. 298.—Coral-like fruit-bodies of *Clavaria flava*. (Harshberger, from Photo by W. H. Walmsley.)

or bracket-like outgrowths often overlapping each other. Example: *Corticium*.

ORDER 4.—CLAVARIALES, the coral or fairy club fungi. Fleshy coral or club-shaped forms, all of which are saprophytes found in woods growing in bunches out of leaf mold. They are all edible and of a white, yellow, lavender or some other brilliant color. (See Fig. 298.)

ORDER 5.—AGARICALES, the mushroom or toadstool alliance. Alike with the other members of the Basidiomycetes, the plant body consists of the mycelium, ramifying through the sub-

stratum, but the part which rises above the surface (the *sporophore* or fruiting body) is in most cases differentiated into a stalk-like body called a *stipe* bearing upon its summit a cap or *pileus*, the latter having special surfaces for the *hymenium*.

FAMILY 1.—HYDNACEÆ, or tooth fungi. This group is characterized by the hymenium being placed over purplish, spiny or long finger-like projections of the pileus. Many of the species of the genus *Hydnum* are edible.

FIG. 299.—*Boletus felleus* in three stages of development. (After Patterson, *Flora W. and Charles, Vera K.*, Bull. 175, U. S. Dept. Agric., pl. xxxi, Apr. 29, 1915.)

FAMILY II.—POLYPORACEÆ, or pore fungi. The sporophores or fruiting bodies of these fungi are various. They may be entirely supinate with pores or shallow depressions on their upper surfaces (*Merulius*), or mushroom-like (*Boletus*), or of the nature of woody or corky (*Fomes*) or fleshy (*Fistulina*) brackets. In all cases the hymenium or basidial layer lines the inner surface of tubes which open on the exterior as pores.

The sporophore or fruit body of *Polyporus officinalis*, when deprived of its outer rind and dried, constitutes the drug AGARI-

CUS. This species grows abundantly on various species of pines, spruces and larches.

FAMILY III.—AGARICACEÆ, the gill family, in which the hymenium covers blade-like plates of the pileus, called *gills*, generally occurring on the under surface of the same. Accord-

FIG. 300.—Meadow mushroom (*Agaricus campestris* L.). A, view showing under side of pileus; *g*, gills; *a*, annulus, or remains of the veil attached to the stipe. B, side view; *s*, stipe; *a*, annulus; *p*, margin of pileus, showing at intervals the remains of the veil. (Gager, after W. A. Merrill.)

ing to the color of their spores the members of this family are divided into 5 groups:

1. Leucosporæ or white-spored, as *Amanita*.
2. Rhodosporæ or red-spores, as *Pluteus*.
3. Ochrosporæ or brown-spored, as *Pholiota*.
4. Porphyrosporæ or purple-brown spored, as *Agaricus*.
5. Melanosporæ or black-spored as *Coprinus*.

The color of the spores is best determined by cutting off the pileus, placing it gill-surface downward on a piece of smooth paper and covering it with a bell jar. Within a day or less the spores, falling off the gills, leave an imprint of the gill arrangement which is called the *spore print*.

AGARICUS CAMPESTRIS (COMMON MUSHROOM).—This plant, also referred to as *Psalliota campestris* (L.) Fr. by some authors is a common edible gill fungus which grows in open, grassy

Yeast plants grow in dilute saccharine solutions containing dissolved nitrogenous substances such as beerwort, Pasteur's solution, grape juice, etc. Here they are constantly wasting away and as constantly being built up. The question may well arise: "How do they obtain the material necessary for growth and repair?" The answer, in a general way, is not difficult. The fluid in which they live is a solution of sugars and of nitrog-

FIG. 283.—Yeast, *Saccharomyces cerevisiae*, the variety known as brewers' bottom yeast; *a*, spore formation; *b*, elongated cells. (After Schneider, *Pharmaceutical Bacteriology*.)

enous and other matters. The cell walls are readily permeable. Food substances diffuse through it into the cell, and by a series of changes (which, indeed, it is no easy matter to understand) are converted into new living substance. The waste products likewise diffuse readily outward. This method of nutrition is called saprophytic, and the yeast plant is said to be a *saprophyte*.

A striking fact must be briefly mentioned in connection with the metabolism of yeast. Many organisms exercise a profound

effect on the medium in which they live. Yeast causes a wholesale destruction of sugar in the surrounding fluid. One of the decomposition products of sugar is alcohol. The alcohol of commerce is produced by the activity of this plant.

The protoplasm of the yeast plant manufactures two kinds of enzymes, viz., an extracellular enzyme called *invertase* which passes out of the yeast cell and hydrolyzes cane sugar to dextrose (invert sugar) and an endocellular enzyme called *zymase* which

FIG. 284.—*Saccharomyces cerevisiae*. The form or variety known as brewers' top yeast. (Oberhefe.)

remains within the yeast cell and breaks down dextrose into alcohol and carbon dioxide.

Saccharomyces has its times of danger and stress when the cells perish in great numbers from cold, starvation, poisons, etc. If not too suddenly exposed, however, they are able to meet adverse conditions by eliminating most of their water, suspending physiological processes, and becoming dormant. Sometimes they enter the resting condition after a process of division, when each cell divides into four parts, each of which becomes nearly dry and is surrounded by a thick wall. Such cells are

called *ascospores*, and their production serves both as a method of multiplying the plant and of tiding over adverse conditions. They can survive for a long time without food or water, and can endure higher temperatures than the active cells and almost any degree of cold.

The dried cells and spores float in the air as dust and so accomplish a dispersal of the organism. Doubtless most of them never again meet suitable environment and so sooner or

FIG. 285.—*Saccharomyces ellipsoideus*. A common yeast found on grapes, jams, jellies, etc. Budding process is shown in many of the cells as also the vacuoles. (Schneider, *Pharmaceutical Bacteriology*.)

later perish. But some will fall into favorable conditions and be able to multiply enormously again, and so the species is continued.

The general method of reproduction in *Saccharomyces* is that of *gemination* or *budding*. A small protuberance of the cell wall commences to form on the parent-cell. This grows larger and a portion of the cytoplasm and nuclear material pass into it. Eventually a daughter-bud, which may assume the size of the parent-cell, is formed. This generally adheres to the parent-cell and produces one or more granddaughter-buds which in turn may produce great-granddaughter-buds before separation from the parent-cell takes place.

There are two varieties of brewer's yeast, viz.: TOP YEASTS and BOTTOM YEASTS. Top yeasts grow on or near the surface of the liquid and produce rapid fermentation at summer temperatures causing great quantities of carbon dioxide to arise to the surface and thus forming the froth which is characteristic of ale, stout and porter. They tend to remain attached in clusters. Most of the yeasts used for compressed yeast and bread making are top yeasts.

FIG. 286.—Saucer-shaped fruit-bodies of *Peziza repanda*. (Harshberger, from Photo by W. H. Walmsley.)

Bottom yeasts grow at about 4°C. at or near the bottom of the vat. They are used in the manufacture of lager beers.

Compressed yeast (Cerevisiæ Fermentum Compressum) or "yeast cake" consists of the moist, living cells of *Saccharomyces cerevisiæ* or of other species of *Saccharomyces*, combined with a starchy or absorbent base.

Yeast contains the antineuritic vitamin B, and the pellagra preventive vitamin B₂ or G.

OTHER YEASTS.—*Saccharomyces ellipsoideus* (Fig. 285) is a wild species, several varieties of which are found growing on grapes especially in districts where wine is produced. It is termed the true wine yeast to distinguish it from other wild species found in grape juice, like *S. apiculatus* and *S. membranifaciens*

which exert a deleterious effect in wine production. Its cells are ellipsoidal, 6μ long, occurring singly or in rows of several generations, which are rather loosely joined.

FIG. 287.—Cup Fungi. A, B, *Lachnea scutellata*. A, Habit; B, ascus with paraphysis; C, D, *Lachnea hemisphaerica*; C, habit; D, ascus with paraphysis; E, *Sarcosphaera arenosa* habit; F, G, *Sarcosphaera coronaria*; F, ascus; G, habit; H, *Sarcosphaera arenicola* ascus with paraphysis. (See *Die natürlichen Pflanzenfamilien* I, 1, p. 181.) (Harshberger.)

S. pastorianus is a common contaminant in brewer's yeast and in yeast cakes. It forms large sausage-shaped cells. It is a cause of disagreeable flavors in beer.

ORDER 2.—PEZIZALES or cup fungi. Examples: *Peziza*, *Lachnea* and *Ascobolus*.

Parasitic or saprophytic plants, whose vegetative bodies consist of a mycelium ramifying through the substratum and whose above ground fruiting bodies are sessile or stalked, cup or saucer-

shaped structures termed *apothecia* (sing. *apothecium*), in which a fruiting membrane (*ascigeral layer*) lines the concave upper surface. The asci are usually eight-spored and separated from each other by filamentous structures called *paraphyses*. (Figs. 286 and 287.)

FIG. 288.—Three aerial hyphae showing the characteristic brush-like branching and spore formation of *Penicillium glaucum*. This fungus is a true saprophyte and is never found on living fruits or vegetables. *a*, Conidiophore branching above into secondary conidiophores; *b*, sterigmata; *c*, conidiospores. (Schneider.)

ORDER 3.—PLECTASCALES, the blue and green molds. Examples: *Aspergillus* and *Penicillium*.

PENICILLIUM GLAUCUM (GREEN MOLD OR MILDEW), a type of mildew, belonging to the Ascomycetes class of Fungi, forms sage-green crusts on bread, jellies, old boots, gloves, and various pharmaceutical preparations. It consists of a felt-like mass of interlaced tubular hyphae called a *mycelium*. From the mycelium

numerous hyphæ project into the air and bear a green powder, the spores. These hyphæ are called *aerial hyphæ*. Other hyphæ grow down into the substratum and are called *submerged hyphæ*.

When a small portion of the mycelium is mounted in 10 per cent. alcohol and observed under the high-power objective, it will be noted that each hypha has a transparent wall and protoplasmic contents and is divided by transverse septa into a number

FIG. 289.—*Penicillium Roqueforti*. *a*, Part of a conidiophore; *b*, *c*, other types of branching; *d*, young conidiophore, just branching, *e*, *f*, conidiiferous cells; *g*, *j*, diagrams of types of fructification. *k*, *l*, *m*, *n*, germinating spores. (After Thom.)

of cells. Each cell contains protoplasm, which is differentiated into cytoplasm (cell protoplasm) and several nuclei. In the cytoplasm will be seen several large clear spaces. These are vacuoles and contain water with nutritive substances in solution, called *cell sap*. Each hypha with its branches is clearly distinct from every other one.

The aerial hyphæ bear brush-like branches, which become constricted on their ends into a moniliform aggregation of rounded spores appearing like a row of beads. Each aerial

hypha is composed of a vertical septate branch of the mycelium called the *conidiophore*, branches of this, which are called *secondary conidiophores*, and chains of spores at the tips of *sterigmata* (cells

FIG. 290.—*Penicillium Camemberti*. *a*, Conidiophore with common type of branching with conidiospores; (*b*) a common less-branched form; *c*, *d*, *f*, diagrams of large fructifications; *g*, *i*, *j*, germinating conidiospores. (From Bull. 82, Bureau of Animal Industry, also after Thom.)

bearing conidia) which are called *conidia* or *conidiospores*. The conidia form the loose green powder characteristic of *Penicillium*.

A number of species of *Penicillium* are useful in the arts. *Penicillium roqueforti* is the principal ripening agent of Roquefort, Gorgonzola and Stilton cheeses. It possesses blue-green globular conidia 4 to 5 μ in diameter.

Penicillium camemberti is the principal agent in the ripening of Camembert cheese. It possesses ellipsoidal bluish-green conidia 4.5 to 5.5 μ in diameter.

Penicillium brevicaulis grows on old moist paper and has been used to detect the presence of arsenic, for when grown in media containing this element, it develops the compound, diethylarsine. It is yellowish-brown in color and its conidia are rough and spiny.

Penicillium expansum is often found on decaying apples where it produces brownish coremia.

CITRIC ACID is produced by fermenting solutions of carbohydrates (mono- or poly-saccharides) by means of molds of the genera *Penicillium*, *Citromyces*, *Aspergillus* and *Sterigmatocystis*. The unfermented carbohydrate is removed by an alcoholic fermentation using yeast.

ASPERGILLUS HERBARIORUM.—This green mold also named *Aspergillus glaucus* and *Eurotium Aspergillus glaucus* is frequently found on fleshy drugs which have not been properly dried. It has also been observed on dried herbarium material, old extracts, on jams, jellies, tobacco, cotton-seed meal, old leather, stale black bread, etc. Like *Penicillium* its vegetative body consists of a mycelium consisting of aerial and submerged hyphæ. It differs from *Penicillium*, however, mainly in not possessing septated conidiophores and by the upper portion of the conidiophores being globular. Upon the globular extremity of the conidiophores are placed numerous elongated sterigmata which bear chains of grayish-green conidia. These are spherical and prickly and range from 7 to 30 μ in diameter. Under certain conditions closed brownish fruit bodies called *perithecia* are produced. These arise on the surface of the substratum from spirally coiled hyphæ and when mature possess numerous *asci*, each of which contains five to eight ellipsoidal *ascospores*.

Aspergillus oryzae is a yellowish-green to brown mold which secretes diastase, a valuable digestive ferment, having the power of converting starch into sugar and dextrin. For centuries the Japanese have employed this species in the preparation of rice mash for Saké, as well as in manufacture of Miso and Soja sauce. The spherical conidiospores are 6 to 7 μ in diameter and of a yellowish-green color.

Aspergillus fumigatus is a pathogenic species which produces a disease in birds, horses, cattle and even, though rarely, in man that is called aspergillosis. The organ most prone to infection

FIG. 291.—*Aspergillus oryzae* associated with yeasts in the making of the Japanese beverage Saké. Vegetative hyphæ (a) and spore-forming hyphæ (b, c, d) are shown. (Schneider, *Pharmaceutical Bacteriology*.)

by this organism is the lung, although the skin, cornea, ears and other parts are also subject to its parasitic influence. It produces short conidiophores with sterigmata bearing long chains of rounded, colorless conidia 2.5 to 3 μ in diameter. Harsh-

berger¹ cites the presence of perithecia in this organism which are nut-brown, globular, 250 to 350 μ in diameter, and inclose

FIG. 292.—*Aspergillus niger* showing conidiophores and conidiospores formation with stages in germination of spores. (Harshberger, after Henri Coupin.)

oval, thin-skinned asci with eight, red, lenticular ascospores each of which has a diameter of 4 to 5 μ .

Aspergillus niger develops dark brown mycelial masses in which are to be noted tall conidiophores terminating in large, black,

¹ "Mycology and Plant Pathology," p. 147.

globular spore heads. The globular vesicle of the spore head bears radially arranged, branched sterigmata that cut off from their tips chains of rounded black to dark brown conidia 3.5 to 5 μ in diameter. This fungus has been found to produce suppurative inflammation of the external and middle portion of the human ear. It is also a cause of cork disease, so often imparting a disagreeable taste to bottled beverages.

This mold, however, is of considerable economic importance. While a number of molds produce citric and oxalic acids from

FIG. 293.—The morel, *Morchella esculenta*. (Gager, from photo by W. A. Murrill.)

sugars, *Aspergillus niger* is especially active in this direction and selected strains of it are used commercially in the production of citric acid on a large scale.

ORDER 4.—TUBERALES, the truffles. Fungi whose septate mycelium is often connected with the roots of trees forming the structure known as *mycorrhiza*. Several species of the genus *Tuber* growing in woods of France, Germany and Italy produce tuberous subterranean bodies called Truffles, which are highly prized as a table delicacy by the inhabitants of these countries.

ORDER 5.—HELVELLALES, the saddle fungi. Fleshy fungi entirely saprophytic, living attached to leaf mold or growing in humus soil or, in a few cases, on decaying wood. The fleshy fruiting bodies (ascocarps) are divided into stalk (*stipe*) and cap

(*pileus*) portions. The external surface of the cap is covered with a layer of asci and paraphyses which together constitute the ascigeral layer. To this group belong the Morels and the Earth Tongues.

One of the Morels, *Morchella esculenta*, is frequently found in fire-swept woods. Its fruiting body consists of a hollow, externally ridged stipe, bearing upon its summit a fleshy pileus whose outer surface is honeycombed with ridges and depressions. The depressions are covered with an ascigeral layer composed of asci and paraphyses. This species is edible. (See Fig. 293.)

ORDER 6.—PYRENOMYCETALES, the mildews and black fungi, common as superficial parasites on various parts of plants. To the Family *Hypocreaceæ* of the black fungi division of this order the Ergot fungus, *Claviceps purpurea*, belongs.

LIFE HISTORY OF CLAVICEPS PURPUREA.—Through the agency of winds or insects the spores (ascospores or conidia) of this organism are brought to the young ovaries of the rye (*Secale cereale*). They germinate into long filaments called hyphæ, which, becoming entangled to form a *mycelium*, spread over the ovary, enter it superficially, secrete a ferment, and cause decomposition of its tissue and the resultant formation of a yellow, mucous substance called honey dew, which surrounds chains of moniliform reproductive bodies known as conidia. The honey dew attracts certain insects which disseminate the disease to other heads of grain.

The mycelial threads penetrate deeper and deeper into the ovary and soon form a dense tissue which gradually consumes the entire substance of the ovary and hardens into a purple, somewhat curved body called a *sclerotium*, or official ergot—the resting stage of the fungus, *Claviceps*.

The ergot falls to the ground and in the following spring sprouts into several, long stalked, globular heads called *stromata* or *ascocarps*. Each (fruiting) head or ascocarp has imbedded in its surface numerous flask-shaped invaginations called *perithecia*, from the bases of which several sacs or *asci* develop from the ends of hyphæ. Within each ascus are developed eight filiform spores (*ascospores*) which, when the ascus ruptures, are

FIG. 294.—A, *Balansia claviceps* on ear of *Paspalum*; B–L, *Claviceps purpurea*; E, sclerotium; C, sclerotium with *Sphaecelia*; D, cross-section of sphaecelial layer; E, sprouting sclerotium; F, head of stroma from sclerotium; G, section of same; H, section of perithecium; J, ascus; K, germinating ascospore; C, conidiospores produced on mycelium. (See *Die natürlichen Pflanzenfamilien* I, 1, p. 371.)

discharged and are carried by the wind to other fields of grain, there to begin over a new life cycle.

The drug *ergot* contains a number of physiologically active constituents chief among which are the alkaloids *ergonovine*, *ergotoxine* and *ergotamine* and the amines *histamine* and *tyramine*. It is used in medicine chiefly because of its power to contract the pregnant uterus and to check post partum hemorrhage.

THE BASIDIOMYCETES OR BASIDIA FUNGI

This large class of fungi, including the smuts, rusts, mushrooms, gill and tooth fungi, etc., is characterized by the occurrence of a basidium in the life history. A *basidium* is the swollen end of a hypha consisting of one or more cells and giving rise to branches called sterigmata, each of which cuts off at its tip a spore, called a *basidiospore*. In addition to the basidiospores, some forms also produce one or more kinds of other types of spores. All of the members have a mycelium composed of septated hyphæ.

THE LOWER BASIDIOMYCETES

This group comprises the smuts and rusts which are parasites living within the bodies of seed plants.

ORDER 1.—USTILAGINALES, the smuts. Destructive parasites which attack the flowers of various cereals, occasionally other parts of these plants. Examples: *Ustilago Maydis*, the corn smut. The basidiospores in this group are borne on a three- to five-celled basidium called a *promycelium*.

USTILAGO MAYDIS (USTILAGO ZEÆ) (CORN SMUT).—Corn smut is a destructive parasite which for a long time was supposed to be confined to the Indian Corn, but which now is known to occur on Mexican Grass. It is the only smut useful in medicine. The mycelium of the fungus extends through all parts of the infected organ of the host through the intercellular-air-spaces, sending short absorbing branches into its cells. In due course of time, it produces large tumor-like swellings on the ears, tassels, husk, leaves, root and stem. Each swelling is at first made up of masses of hyphæ, the whole being covered with a tightly appressed membrane which has a whitish appearance like

German silver. Later the cells of the hyphæ lay down walls becoming divided into smaller cells, each of which becomes rounded off as a spore. This spore is known as a *chlamydospore* or *brand spore*. The spores are at first a dark olive-green, but on

FIG. 295.—Smut boil of *Ustilago zeæ* on ear of corn, developed from one infected kernel. (After Jackson, F. S., Bull. 83, Del. Coll. Agric. Exper. Stat., December, 1908.)

maturity are dark brown. They are sub-spherical and show prominent but tiny spines. These spores fall to the ground and pass the winter. In the spring each germinates into a three-to five-celled filament called a *promycelium*, from the cells of which

basidiospores arise. (Fig. 296.) The basidiospore, upon coming in contact with young growing tissue or a wound upon the host plant, starts a new infection and develops a mycelium which penetrates the infected organ.

ORDER 2.—UREDINALES, the rusts. Obligate parasites possessing a septated, branched mycelium which ramifies

FIG. 296.—Germination of the chlamydospores of corn smut (*Ustilago zeæ*); 1, Various stages in germination from corn 3 days after being placed in water; 2, spores germinated in contact with air; 3, several days after spores were placed in $\frac{1}{20}$ per cent. acetic acid, formation of infection threads. *a*, Spores; *b*, promycelia; *c*, basidiospores; *d*, infection threads; *e*, detached pieces of mycelia. (After Bull. 57, Univ. Ill. Agric. Exper. Stat., March, 1900.)

through the intercellular-air-spaces of the host and sends haustoria into the cell cavities. The different stages of their life cycle are either restricted to one host or distributed between two or more hosts. An outline of the life history of the wheat rust will give an idea of the peculiarities of the group.

THE WHEAT RUST (*PUCCINIA GRAMINIS*).—If we examine the wheat plant just before harvest we will find on the stems and

leaves some rust-red lines. The presence of the mycelium of the fungus in the intercellular spaces of the host does not kill the host directly or appear to stunt its growth, but the effect of the parasite on the host is seen when the grains mature. The grains are small and mushy, due to the fact that the nutrition of the host had been disturbed and the formation of starch in the grains inhibited. The mycelium is localized and gives rise underneath the epidermis to rounded egg-shaped spores attached

FIG. 297.—Spore forms of wheat rust, *Puccinia graminis*. A, Section through barberry leaf showing pycnia (spermagonia) on upper surface and aecia on lower; B, two urediniospores; C, germinating urediniospore; D, teliosorus showing several teliospores; E, single two-celled teliospore; F, germinating teliospore with four-celled basidium and two basidiospores; G, basidiospore growing on barberry leaf. (Harshberger, adapted from deBary.)

to it by short pedicels. The spores are produced in such numbers that the space beneath is too confined. As the long epidermal cells of grasses run longitudinally, the pressure of the spore masses from within causes the epidermis to crack and its edges become turned back. Through the resultant cleft the *summer spores* or *uredospores* (urediniospores) are thrust out. These uredospores are orange-brown in color and covered with minute spines. The mass of them has been called a *uredinium*. These spores are detached from the pedicels and blown by the wind to healthy plants. After summer is over and dry weather

comes on, an examination of stubble in the field (blades of grass and stems of wheat left carelessly), shows that these rust-red lines are replaced by brownish-black spores called *teleutospores* (teliospores). A mass of these is known as a telium.

The summer stage on wheat is known as *Uredo linearis*.

The autumn stage on wheat is known as *Puccinia graminis*.

The teleutospores are two-celled and have thick walls and persistent pedicels. They remain attached to the stubble until the following spring and then either one or both cells composing them produce an outgrowth known as a *promycelium* (nothing but a basidium divided transversely into four cells). Each cell of the basidium is capable of producing a branch, at the tip of which a *basidiospore* is formed. These basidiospores are blown to the Barberry (*Berberis*) and infect the leaves of this plant. The mycelium runs in the intercellular-air-spaces and causes the appearance of a number of small depressions on the upper side of the leaf. These in section are a rich chocolate brown and known as *spermagonia*. In the center of a spermagonium are produced hyphæ, which project out to its orifice and obstruct off minute spores called *spermacia*. On the opposite side of the leaf, cup-shaped depressions are formed, each with a limiting membrane (*peridium*). Within the cup-shaped depression thousands of spores are formed in chains closely packed together. These are the *æcidiospores* (*æciospores*). The cluster cup is called an *Æcidium* (*Æcium*). These æciospores are conveyed to wheat and cause infection, thus completing the life cycle. It has been observed that in America the uredospores or summer spores may winter over and infect healthy plants, so that the Barberry phase is completely eliminated from the life cycle.

SOME OTHER RUSTS.—The White Pine Blister Rust, *Cronartium ribicola*, causes great damage yearly to the *Pinus strobus* and other 5-needle pines. It has its aecial and spermatial stages on these pines and the uredospore and teleutospore stages on currants (*Ribes* spp.). The Hollyhock Rust, *Puccinia malvacearum* occurs on the hollyhock (*Malva rosea*) and other species of *Malva*, notably the drug-yielding *Malva rotundifolia* and *M. sylvestris*. It passes through two stages, the teleutospore and basidiospore, on the same host, only a single host being involved.

The Cedar and Apple Rust produces aecia on the leaves of the apple and hawthorn and telia on the red cedar (*Juniperus virginiana*). The mycelium of this fungus growing in the leaves of the juniper produces the large brown galls known as "cedar apples."

ORDER 3.—AURICULARIALES.—The so-called "ear fungi" which occur on the bark of many plants, on wooden fences, etc., as auriculate growths which when young are jelly-like and brilliantly colored, when old, hard, grayish and considerably wrinkled. The earlike fruiting body is known as the *sporophore*. Its internal surface is lined with a *hymenium* or fruiting membrane consisting of numerous four-celled basidia, each of which cuts off at its tip a basidiospore. A common example of this group is the Jew's ear fungus.

ORDER 4.—TREMELLALES.—Saprophytes which live on decaying wood as moist, soft, quivering gelatinous growths becoming later dry and horny.

THE HIGHER BASIDIOMYCETES

Mostly fleshy forms characterized by one-celled basidia with generally four, occasionally six, eight or two sterigmata, each of which cuts off a basidiospore at its tip.

DIVISION A.—HYMENOMYCETES

(Hymenium or spore-bearing surface exposed)

This division of higher basidiomycete fungi comprises the following orders: Dacromycetales, Exobasidiales, Thelephorales, Clavariales and Agaricales.

ORDER 1.—DACROMYCETALES.—This order includes the "weeping fungi." One of the most common is *Dacromyces deliquescens* which occurs as a gelatinous body of bright red color on dead wood. The basidiospores are formed during a wet period and the fungus swells up in the water forming a slimy mass. In addition to basidiospores the mycelium may break up into oidiospores, if the wet period is prolonged. In consisting of slimy gelatinous masses the "weeping fungi" approach the *Tremellaceæ* but are distinguished from them in the basidium being undivided in the former and divided in the latter.

ORDER 2.—EXOBASIDIALES.—This group is found growing parasitically on shrubs especially those of the heath family. The mycelium lives in the tissues of the stems, leaves, sepals and petals and produces spongy, fleshy, yellowish or brownish galls which are popularly called "Azalea apples." The galls are edible. They are covered with a hymenium.

ORDER 3.—THELEPHORALES. These possess fruit-bodies appearing on tree trunks as flattened leathery or crust-like

FIG. 298.—Coral-like fruit-bodies of *Clavaria flava*. (Harshberger, from Photo by W. H. Walmsley.)

or bracket-like outgrowths often overlapping each other. Example: *Corticium*.

ORDER 4.—CLAVARIALES, the coral or fairy club fungi. Fleshy coral or club-shaped forms, all of which are saprophytes found in woods growing in bunches out of leaf mold. They are all edible and of a white, yellow, lavender or some other brilliant color. (See Fig. 298.)

ORDER 5.—AGARICALES, the mushroom or toadstool alliance. Alike with the other members of the Basidiomycetes, the plant body consists of the mycelium, ramifying through the sub-

stratum, but the part which rises above the surface (the *sporophore* or fruiting body) is in most cases differentiated into a stalk-like body called a *stipe* bearing upon its summit a cap or *pileus*, the latter having special surfaces for the *hymenium*.

FAMILY 1.—HYDNACEÆ, or tooth fungi. This group is characterized by the hymenium being placed over purplish, spiny or long finger-like projections of the pileus. Many of the species of the genus *Hydnum* are edible.

FIG. 299.—*Boletus felleus* in three stages of development. (After Patterson, *Flora W. and Charles, Vera K.*, Bull. 175, U. S. Dept. Agric., pl. xxxi, Apr. 29, 1915.)

FAMILY II.—POLYPORACEÆ, or pore fungi. The sporophores or fruiting bodies of these fungi are various. They may be entirely supinate with pores or shallow depressions on their upper surfaces (*Merulius*), or mushroom-like (*Boletus*), or of the nature of woody or corky (*Fomes*) or fleshy (*Fistulina*) brackets. In all cases the hymenium or basidial layer lines the inner surface of tubes which open on the exterior as pores.

The sporophore or fruit body of *Polyporus officinalis*, when deprived of its outer rind and dried, constitutes the drug AGARI-

OLD BULB

Removed 5/3/07

CUS. This species grows abundantly on various species of pines, spruces and larches.

FAMILY III.—AGARICACEÆ, the gill family, in which the hymenium covers blade-like plates of the pileus, called *gills*, generally occurring on the under surface of the same. Accord-

FIG. 300.—Meadow mushroom (*Agaricus campestris* L.). A, view showing under side of pileus; *g*, gills; *a*, annulus, or remains of the veil attached to the stipe; B, side view; *s*, stipe; *a*, annulus; *p*, margin of pileus, showing at intervals the remains of the veil. (Gager, after W. A. Murrill.)

ing to the color of their spores the members of this family are divided into 5 groups:

1. Leucosporæ or white-spored, as *Amanita*.
2. Rhodosporæ or red-spores, as *Pluteus*.
3. Ochrosporæ or brown-spored, as *Pholiota*.
4. Porphyrosporæ or purple-brown spored, as *Agaricus*.
5. Melanosporæ or black-spored as *Coprinus*.

The color of the spores is best determined by cutting off the pileus, placing it gill-surface downward on a piece of smooth paper and covering it with a bell jar. Within a day or less the spores, falling off the gills, leave an imprint of the gill arrangement which is called the *spore print*.

AGARICUS CAMPESTRIS (COMMON MUSHROOM).—This plant, also referred to as *Psalliota campestris* (L.) Fr. by some authors is a common edible gill fungus which grows in open, grassy

fields during late summer and early autumn. It is never found in the forest or on trees or fallen trunks, seldom in the mountains. The cultivated form grows in specially constructed houses made of boards. A corridor runs through these houses so that the mushroom beds can be easily reached. In the growth of mushrooms tons of horse manure are used. This is covered with loamy soil $1\frac{1}{2}$ inches thick. The whole mass is compacted together. Into the resultant beds is introduced English-grown spawn, which comes in flat brick-shaped masses (horse manure through which mycelium has grown). Pieces of these "bricks"

FIG. 301.—Fruiting surface (hymenium) of a mushroom (*Agaricus*). *m*, hyphae of the trama and sub-hymenium; *b*, basidium; *st*, sterigma; *sp*, spores. Diagrammatic. (Gager.)

are put in the horse manure bed only after the heat has first disappeared. The beds are then watered well and in a short time the *sporophores* or fruiting bodies of the fungus spring up.

The mycelium or vegetative body of *Agaricus* which develops in the soil from spores (basidiospores) is white and thready. On this mycelium develop little white buttons, first about the size of a pin head, becoming later pea size and then assuming a pear-shaped form. At this stage the sporophore consists of a cylindrical solid stipe or stalk and a pileus or cap. The border of the pileus is joined to the stipe by means of a "*partial veil*." Within this veil is found a circular cavity, into which the *gills* grow. At first the stipe grows faster than the rest of the fruiting

body. The pileus expands transversely and the gills keep pace. After a while the veil ruptures, leaving a portion attached to the stipe. This constitutes the *annulus* or *ring* (*true annulus*). The hyphæ in the pileus form the *Tela contexta*. If we make a section through a gill, the hyphæ are seen to run longitudinally. The central part is called the *trama*; next and outside trama is the *sub-hymenium*; next, *hymenium*, consisting of *basidia* (hence a basidial layer). Each basidium bears two or four little points

FIG. 302.—Deadly amanita (*Amanita muscaria*) showing volva at base of stipe and frill-like stipe ring. (After Chestnut, V. K., Bull. 175, U. S. Dept. Agric., pl. i, Apr. 29, 1915.)

known as sterigmata. Each sterigma bears a purplish-brown basidiospore. The basidiospores falling to the ground germinate into hyphæ and these become interlaced to form a mycelium.

In the wild mushroom the gills are at first white in the button stage, later pink; in the cultivated, fawn-colored. Ultimately in the wild form the gills turn blackish brown. The spores are purplish-brown. The color of the stipe is white; that of the upper surface of the pileus varies from satiny white to pallid brown.

THE AMANITAS (POISONOUS FUNGI).—*Amanita muscaria* and *Amanita phalloides*, commonly known as the “fly agaric” and

the "deadly agaric," respectively, are very poisonous forms. *Amanita muscaria* is common in coniferous forests, although it may be found in cultivated soil along roadsides, in open woods and wood margins. It occurs singly or in patches. Its chief poisonous constituent is muscarine for which atropine is an antidote. An emetic should first be given. *Amanita phalloides*

FIG. 303.—The deadly amanita, *Amanita phalloides*. Note the cup at the base of the stipe. (Gager, from photo by E. M. Kittredge.)

is found in woods and borders of fields and occurs singly and not in groups.

Each of these have fruiting bodies (sporophores), which begin at the surface of the ground as a button similar to that of the edible mushroom. This enlarges and assumes a dumbbell shape. The whole button is covered by an outer veil, known as the *velum universale*, which encloses the pileus, gills and stipe. As the stipe lengthens more rapidly than the pileus, the upper part of the veil is stretched and finally breaks in its middle

portion. The lower part remains as a cup, out of which the stipe grows. The upper part is carried up as shreds adhering to the margin of the pileus. The lower part is called the *volva* or death cup. The annulus present is a *false annulus*, for it represents a peeling down of the upper part of the stipe. Both have chalk-white gills, an usually white stipe with a bulbous base, and white spores.

The pileus of *Amanita muscaria* varies from orange-red or blood-red to orange or yellow. The surface is covered with

FIG. 304.—A colony of Puff Balls, *Lycoperdon*, growing saprophytically upon a portion of a rotten log. (Photograph by author.)

prominent warty scales, the latter being remnants of the veil which are readily removed. Its stipe is white or pale sulfur colored, at first stuffed, later hollow. The annulus is large, white and ragged.

The pileus of *Amanita phalloides* varies from pure white to dull yellow to olive or umber. It does not possess the warty scales found in the *Amanita muscaria*, but occasionally has a few membranous patches. Its stipe is generally smooth, usually white, solid and bulbous below but hollow upwards.

Amanita phalloides is the most poisonous of all species of toadstools. Its chief poisonous constituents are the glucoside amanita-hemolysin and the toxin called amanita-toxin. Amanitahemolysin is a blood laking principle. No antidote is known which will neutralize the deadly effects of this fungus.

DIVISION B.—GASTEROMYCETES

(Hymenium inclosed)

ORDER 1.—LYCOPERDALES, or puff ball alliance. This order includes a number of interesting parasites and saprophytes, the most common of which are the earth stars belonging to the genus *Geaster* and the puff balls, the most common form being *Lycoperdon*. In these, the fruiting sporophore consists for the most part of a shell-like covering called the *peridium*, composed

FIG. 305.—Bird's Nest Fungus (*Crucibulum*). Note the open sporophores appearing like nests with eggs. The "eggs" are the peridiola or special bodies in which the basidiospores are produced. (Mottier.)

of an outer layer or *exoperidium* and an inner layer of *endoperidium*. The peridium in the unripe condition of the sporophore covers a mass of soft cellular tissue called the *gleba*. Upon the ripening of this mass, the interior is seen to be divided into many-branched compartments that are separated from each other by walls made up of branched hyphæ. These walls are lined with a hymenium composed of many basidia, each of which constricts off usually four basidiospores. The earth stars differ from the puff balls in possessing an outer wall or *exoperidium* which splits in star-shaped fashion.

ORDER 2.—NIDULARIALES, the nest fungi. A group of Gasteromycetes whose sporophores are crucible- or crater-like.

These arise from a subterranean mycelium and show an outer and inner peridial layer. The outer peridium is roughened at its base. The inner peridium is leathery and may or may not be continued over the top. When mature the crucible-like body shows black seed-like bodies inside which resemble eggs in a bird's nest. Each one of these is connected with the inner peridium by a cord which resembles the umbilical cord of an animal. These inner bodies are called *peridiola* (sing. *peridiolum*). Each peridiolum consists of a hard glistening outer layer and a spongy inner layer surrounding a cavity into which basidia and basidiospores project. These fungi are found in stiff clayey soil.

ORDER 3.—PHALLALES, the carrion or stink-horn fungi. This group consists of highly and characteristically colored forms which, when mature, emit most vile and penetrating odors. The fruiting body, in each instance, begins as an egg-shaped structure which starts its growth from a widely spread underground mycelium of chalky-white color. As the "eggs" enlarge they push above the surface of the ground. The central

portion, elongating, then breaks through the outer or peridial portion, which remains as a cup or *völva* at the base of the mature fruit body. Upon the summit of the central stalk rests the cup-like, many-chambered gleba. The basidiospores are imbedded in a greenish fetid slime formed by a mucilaginous disintegration of the substance of the hymenium. This fetid green material is attractive to carrion flies which visit the plants and remove the material with its embedded spores. The latter will not germinate until after passing through the alimentary canal of these flies.

FIG. 306.—Mature stink-horn, *Dictyophora duplicata*. (Harshberger, from photo. by W. H. Walmsley.)

THE FUNGI IMPERFECTI

An assemblage of varied forms, the life histories of most of which are imperfectly understood. Only one kind of spore, the conidiospore, is known to occur in this group. In this group are included numerous parasites which produce diseases in crop plants and some in man and the lower animals.

Among the fungi of this group producing diseases in man are *Trichophyton* species which produce a skin affection called "ring-worm," and *Monilia albicans* which produces an infection of the mucous membrane of the mouth known as "thrush."

CHAPTER XVIII

THE LICHENS

Lichens are variously colored, usually dry and leathery plants, consisting of symbioses of algæ and fungi. In each case the fungus derives its food from materials manufactured by the alga and in return extracts water from the substratum and shares it with the alga. The association is therefore mutually benefi-

FIG. 307.—A foliaceous lichen, *Physcia stellaris* (L.) Nyb., growing on a rock. The cup-shaped structures are the fruiting bodies (apothecia). At the left are seen two very young plants. (Gager.)

cial. While Blue-green and Protococcus forms of Green Algæ and Ascomycete Fungi are for the most part concerned in lichen formation, in some instances Basidiomycete Fungi are also involved with the algæ. Thus we have two distinct classes of Lichens depending upon the type of fungus associated with the alga, viz.: *Asco-lichens* and *Basidiolichens*.

Lichens are found on the bark of trees, on rocks, logs, old fences, etc.

According to the manner of growth of the thallus and nature of attachment to the substratum, three different sub-groups of lichens may be distinguished, viz.: (1) Foliaceous, where the thallus is flat, leathery and leaf-like and attached to the substratum at different points. To this group belong *Physcia*, *Sticta* and *Parmelia*. (2) Crustaceous, where the thallus closely adheres to rocks and bark of trees. To this group belong *Graphis* and *Pertusaria*. (3) Fruticose, where the thallus is upright and branching. To the last group belong *Cetraria islandica*, species of *Cladonia*, and *Usnea*.

FIG. 308.—*Cetraria islandica*, or Iceland Moss, a medicinal lichen. (Sayre.)

STRUCTURE.—The body of a typical lichen shows a differentiation into the following regions:

- (1) A *tegumentary layer* of several rows of hyphal cells.
- (2) An *upper cortex* of densely arranged hyphal cells.
- (3) An *algal layer* containing green algal cells, often called *gonidia*.

(4) A *medullary layer* of loosely interwoven fungal hyphæ, with large air-spaces. This is usually the broadest layer and sometimes also contains algal cells.

(5) A *lower cortex* of hyphal cells, less dense than the upper cortex. From the surface cells of this layer rhizoids are developed which absorb water and nutrient salts from the substratum.

FIG. 309.—A lichen, *Parmelia perlata* (L.) Ach. 1. Plant, slightly reduced in size; *a*, apothecia; *b*, lobe of thallus; *c*, soredial patches. The soredia are vegetative reproductive bodies composed of both algal and fungal elements, and therefore able to reproduce the lichen; the ascospores, alone, cannot do this. 2. Longitudinal section of apothecium; *a*, thecium; *b* and *c*, the two layers of the hypothecium; *d*, upper algal layer; *e*, colonies of algae distributed through the medullary layer; *f*, lower algal layer; *g*, lower cortical layer. 3. Cross-section of vegetative portion of thallus. 4. Paraphyses (sterile fungal filaments), and spore-sac (ascus), containing ascospores. 5. Ascospores. 6. Algal cells, surrounded by fungal hyphae with haustoria (absorbing branches). (Gager.)

REPRODUCTION.—Lichens reproduce either by producing spores or by detaching scale-like portions of their bodies. In the Asco-lichens (which make up the majority) either saucer-shaped structures called *apothecia* (sing. *apothecium*) occur on the upper surface or dark-colored, spherical structures called *perithecia* somewhat embedded in the upper surface, arise. In either type of structure *asci* and sterile hyphæ (*paraphyses*) arise. Each ascus usually contains eight spores. The spores upon maturity are liberated, each germinating in the presence of moisture into a

FIG. 310.—*Roccella tinctoria*, a dye-yielding lichen. (From Strasburger after Wiesner.)

hypha. The hypha, upon coming into contact with an alga cell, grows around it to form a lichen.

Scales or *soredia* are found on many lichens. These consist of a network of hyphæ enclosing algal cells. By becoming detached from the parent plant, they develop new lichens and so constitute a means of vegetative propagation.

USES.—To the pharmacist and chemist lichens are chiefly of interest because of the coloring principles which they contain. Species of *Lecanora* and *Roccella tinctoria* yield, when subjected to fermentation, the dyes orcein and litmus. Litmus is one of the best indicators in volumetric analysis. Cudbear, a purplish-red powder, used extensively for coloring pharmaceutical preparations in the form of tincture, is prepared by treating species of *Roccella*, *Lecanora* or other lichens with ammonia water. Other

lichens, such as *Cetraria islandica*, various species of *Parmelia*, *Usnea* and *Alectoria*, have been used in medicine because of demulcent principles which they contain.

One of the most conspicuous lichens in northern latitudes is the Reindeer Moss (*Cladonia rangiferina*). It is a slender, branched, fruticose form which grows to the height of about a foot and bears its brownish apothecia at the ends of branches. In Arctic regions it occurs in dense patches covering large areas and forms the only kind of fodder available in many sections for reindeer and caribou.

Another lichen, *Lecanora esculenta*, is thought to represent the source of Scriptural Manna. It has been employed as a food in periods of famine by the natives of southwestern Asia.

CHAPTER XIX

THE BRYOPHYTES

This interesting group, comprising the second division of the Vegetable Kingdom, is represented by the liverworts and the mosses. These plants show a beginning of definite alternation of generations, *i.e.*, gametophyte (sexual phase) alternating with sporophyte (asexual phase of development) in their life history, the two phases being combined in one plant. The sporophyte is attached to the gametophyte throughout its existence, being parasitic upon it. Of the two phases, the gametophyte is always the larger and more conspicuous structure. The female sexual cell is always lodged in an archegonium (a multicellular female sexual organ).

There are two classes of Bryophytes: CLASS I, HEPATICÆ or liverworts and CLASS II, MUSCI or mosses.

HEPATICÆ OR LIVERWORTS

The liverworts are a class of aquatic or terrestrial plants occurring in tropical and temperate regions. While most of them live in moist land situations, some are found submerged or floating in water, some are *epiphytic*, and others occur in localities where long dry seasons prevail. The body of a liverwort either consists of (1) a small, flat, dichotomously branching thalloid body showing differentiation into a green upper or dorsal region and a nearly colorless under or ventral part, as in *Riccia* or *Marchantia*, or (2) of a branching stem which bears two rows of dorsal leaves containing chloroplasts and one row of pale ventral leaves, as in *Porella* and other higher liverworts. Reproduction is by fertilization or by vegetative multiplication. Some species develop both kinds of sexual organs (antheridia and archegonia) on the same gametophyte and so are monœcious, while others develop antheridia on one gametophyte and archegonia on another gametophyte and are therefore diœcious.

There are three orders of liverworts, viz.:

ORDER 1.—MARCHANTIALES, including *Marchantia* and *Riccia*.

ORDER 2.—JUNGERMANNIALES, the leafy liverworts, including *Porella*.

FIG. 311.—*Marchantia polymorpha*. Male gametophyte. Lower figure: Thalloid body with cupules containing gemmæ and male receptacles (antheridiophores) in various stages of development, *a*, *b*, *c*, *d*. Upper figure: antheridia in depressions of upper surface of disk of male receptacle, in successive stages of development, *f*, *e*, *d*, *c*, *a*. *b*, depression with the antheridium taken out. (From Small after Kny.)

ORDER 3.—ANTHOCEROTALES, having the most complex sporophyte generations among liverworts, including *Anthoceros* and *Megaceros*.

MARCHANTIA (A TYPICAL LIVERWORT)

One of the most common and widely distributed liverworts is *Marchantia polymorpha*. It occurs in dense patches on the moist

surface of soil in flower pots of greenhouses and on the damp rocks and humous soil of woodlands.

GAMETOPHYTE GENERATION.—The gametophyte or sexual plant consists of a dichotomously branched *thalloid body* with a distinct midrib. With a hand lens it will be noted that its green upper surface exhibits numerous pale green lines which divide it into small rhomboidal-shaped areas. In the center of each of these is a pore which leads into an air-chamber within the body. Its lower surface is nearly colorless and exhibits *rhizoids* and delicate *scales*.

FIG. 312.—*Marchantia polymorpha*. Microscopical structure of the thallus which shows dorsioventral differentiation. A, vertical section showing one large air-cavity beneath the epidermis. B, C, D, portions of each of three kinds of rhizoids, the last two with inner thickenings of the wall. (Mottier.)

A cross section of the thalloid body, when examined microscopically, will disclose an upper epidermis of epidermal cells containing chloroplasts and showing *stomata*, each stoma surrounded by four vertical rows of cells. Directly beneath this is a row of *air-chambers* separated by partitions. In each air-chamber branched rows of densely *chlorophylloid cells* grow upward from the cells which constitute its floor. These do most of the work of photosynthesis. Below the air-chambers are several rows of clear parenchyma cells containing only a few *chloroplasts*. From the lower surface of these are to be noted outgrowths as slender rhizoids and scales.

Some of the gametophytes will show cup-like structures on their upper surface. These are called *cupules*. They contain

special buds called *gemmae* which, upon being detached, may give rise directly to a new gametophyte. This is the vegetative method of reproduction.

Marchantia is *diœcious* for it produces two kinds of gametophytes, one bearing antheridia (male) and another archegonia

FIG. 313.—*Marchantia polymorpha*. 1, Vertical section of one lobe of the disk of the archegonial receptacle, showing archegonia on lower surface, the group being surrounded by a laciniated perichaetium. 2a, 2a₂, 3, 4, 5, 9, 10, Successive stages in the development of the archegonium. 6, 7, 8, Transverse sections of the same. 11, Mature archegonium with antherozoids attracted to mouth. 12, Transverse section of venter of mature archegonium. 13, Transverse section of neck of same, showing six cells. 14, Tuberculate rhizoid. 15, Transverse section of same. (From Small after Kny.)

(female). There is no way of distinguishing one from the other until stalked structures or *receptacles* arise from the upper surface of the gametophyte.

The *antheridial receptacle* consists of an erect stalk surmounted by a disk with an eight lobed edge. Embedded in numerous separate cavities of the upper region of the disk are the *antheridia*.

The *archegonial receptacles* consist of a stalk surmounted by an umbrella-shaped disk with about nine somewhat pendant rays. On the lower surface of the disk the *archegonia* arise as flask-shaped structures which hang downward.

The *antheridia* are somewhat oval and consist of a wall of *sterile cells* surrounding a mass of *sperm mother cells*. The latter give rise to the *sperms* or *antherozoids* which are biciliate.

The *archegonia* are flask-shaped structures, each of which possesses a basal bulbous part called the *venter* and an elongated tubular part termed the *neck*. The flask has an outer layer of *sterile cells* and a central axial row of cells which from the base of the venter upward are the *egg cell*, *ventral canal cell* and *neck canal cells*. As the archegonium matures, the horizontal walls of the canal cells are dissolved by an enzyme and the contents of the cells are transformed into a mucilaginous mass. During wet weather the upper portion of the wall of the mature antheridium bursts open and the biciliate sperms are liberated in a mucilaginous mass. Attracted by the substances then issuing from the mouths of the archegonia, they swim toward them and usually a number pass down the neck of the canal of each archegonium and gather around the egg. One of them unites with the egg fertilizing it and forming an *oospore*. The act of fertilization concludes the gametophyte generation.

THE SPOROPHYTE.—The *oospore* enlarges and then divides to form a spherical mass of cells which in time become differentiated to form a sporophyte or asexual plant which consists of (1) a basal *foot* which attaches the sporophyte to the base of the archegonium and absorbs nutrient substances in solution from it, (2) a short *stalk* which gradually lengthens and (3) a *capsule* composed of an outer layer of sterile cells enclosing *spores* and slender, elongated cells called *elaters*.

Upon the rupture of the wall of the capsule the spores are discharged, being aided in their dispersal by the elaters found among the spores which, being hygroscopic and with spirally thickened walls, bend and twist, giving force to the process of spore distribution. Each spore, upon absorbing water, may develop into a gametophyte.

ALTERATION OF GENERATIONS.—In the life history of *Marchantia* there are two distinct generations. The gametophytes bear the sex organs in which gametes (sperm and egg) are produced. The sperm unites with the egg forming an oospore.

FIG. 314.—*Marchantia polymorpha*. 1. Thalloid body of Female Gametophyte with archegonial receptacles (archegoniophores) in successive stages of development, a_1, a_2, b, c, d, e, f . 2, 3, Upper and under views of archegoniophore when the sporophytes are ripe. 4, 5, Stages in the development of the sporophyte. 6, Young sporophyte enclosed by a perigynium. 7, Mature sporophyte shedding spores. 8, Elater and spores. 9, Surface view of cells from wall of capsule. (From Small after Kny.)

The oospore develops into a sporophyte which forms spores. Each spore may develop into a gametophyte.

MUSCI OR MOSSES

Plants found on the ground, on rocks, trees and in running water. Their life histories consist of two generations,

FIG. 315.—*Sphagnum acutifolium*, Ehrb. A, prothallus (*pr*), with a young leafy branch just developing from it; B, portion of a leafy plant; *a*, male cones; *ch*, female branches; C, male branch or cone, enlarged, with a portion of the vegetative branch adhering to its base; D, the same, with a portion of the leaves removed so as to disclose the antheridia; E, antheridium discharging spores; F, a single sperm; G, longitudinal section of a female branch, showing the archegonia (*ar*); H, longitudinal section through a sporogonium; *sg*¹, the foot; *ps*, pseudopodium; *c*, calyptra; *sg*, sporogonium, with dome of sporogenous tissue; *ar*, old neck of the archegonium; I, *Sphagnum squarrosum* Pers.; *d*, operculum; *c*, remains of calyptra; *qs*, mature pseudopodium; *ch*, perichætium. (Gager, from Schimper.)

gametophyte and sporophyte similar to the liverworts but differ from liverworts, generally, by the ever-present differentiation of the gametophyte body into distinct stem and simple leaves, and

the formation of the sexual organs at the end of an axis of a shoot. They are either *monœcious*, when both kinds of sexual organs are borne on the same plant, or *diœcious*, in which case the antheridia and archegonia arise on different plants.

ORDER 1.—SPHAGNALES, or Bog Mosses, including one family, *Sphagnaceæ* with the single genus, *Sphagnum* which has many species. Pale mosses of swampy habit whose gametophytes grow indefinitely at the apices of the stems repeating their growth periodically while their lower portions die away gradually and form peat, hence their frequent name of Peat Mosses. Peat in a dry state is used as fuel. The peat mosses possess a protonema which differs from that of other mosses in being thallose and bearing a resemblance to the prothallus of a fern. The leafy stem ends in a three-angled apical cell and bears a branch at every fourth leaf. The branches divide to form secondary branches. The *antheridia* arise on small lateral branches in the leaf axils, each one consisting of a slender stalk and a globular capsule. The *archegonia* arise on the leafy tips of stems. The zygote gives rise to the *Sporophyte* which possesses a large foot, a small stalk and a spherical sporangium the end of which splits off as a lid to allow the spores to escape.

A number of species of *Sphagnum* have been employed in surgery as absorbents in place of gauze. For this purpose they must be thoroughly cleaned and sterilized.

The power which bog mosses have to absorb water depends on the presence of large chlorophyll-less cells the walls of which have pores and are supported by spiral or ring-like thickenings. These cells absorb water by capillarity. Between

FIG. 316.—Surface view of portion of a leaf of *Sphagnum palustre* showing reservoir cell (*r*); transverse bands that prevent cells from collapsing (*h*); pores in reservoir cells (*p*); and green cells containing chlorophyll (*g*).

these empty cells are elongated narrow cells containing chloroplasts.

ORDER 2.—ANDREÆALES or Rock Mosses consisting of one family, *Andreæaceæ* with the single genus *Andreæa*, of xerophytic habit, occurring on siliceous rock.

ORDER 3.—BRYALES, or true mosses, comprising the most highly evolved types of bryophytes. Examples: *Polytrichum*, *Funaria*, *Hypnum*, and *Mnium*.

LIFE HISTORY OF POLYTRICHUM COMMUNE (A TYPICAL TRUE MOSS).—*Polytrichum commune* is quite common in woods, forming

FIG. 317.—Protonema of a Moss. A, Early stages of germinating spores: *v*, vacuole; *w*, rhizoid; *s*, spore wall or exosporium. B, Developed protonema about three weeks after germination; *h*, primary filament; *k*, bud and *w*, rhizoid. The bud develops into a gametophytic plant. (After Sachs from Small.)

a carpet-like covering on the ground beneath tall tree canopies. It is dioecious, the plants being of two kinds, male and female.

Beginning with a spore which has fallen to the damp soil, we note its beginning of growth (germination) as a green filamentous body composed of chloroplast-containing cells arranged end to end. This is called a *protonema*. This protonema soon becomes branched, giving rise to hair-like outgrowths from its lower portion called *rhizoids* and *lateral buds* above these which grow into leafy stems bearing rhizoids at their bases and which are commonly known as "moss plants." At the tips of some of these leafy stems *antheridia* (male sexual organs) are formed while on others *archegonia* (female sexual organs) are developed. The leafy

tip of the male plant can readily be distinguished from that of the female plant by its cup-like shape and orange color. These

FIG. 318.—Hair-cap moss (*Polytrichum commune*). *A*, male plant; *B*, same, proliferating; *C*, female plant, bearing sporogonium; *D*, same; *g*, gametophyte; *s*, seta; *c*, capsule; *o*, operculum; *a*, calyptra; *E*, top view of male plant. (Gager.)

organs are surrounded at the tips by delicate hairy processes called *paraphyses* as well as leaves for protection. The antheridia bear the *antherozoids*, the archegonia, the eggs or *ova*, as in the liverworts. When an abundance of moisture is present, the

antherozoids are liberated from the antheridia, swim through the water to an archegonium and descend the neck canal, one fertilizing the egg by uniting with it. This completes the sexual or gametophyte generation. The fertilized egg now undergoes

FIG. 319.

FIG. 320.

FIG. 321.

FIG. 319.—Vertical section through summit of male gametophyte of a Moss. *Mnium hornum* ($\times 40$) an, Antheridia; par, Paraphyses.

FIG. 320.—A, Antherozoid of Moss ($\times 1,200$); B, Antherozoid of fern ($\times 700$).

FIG. 321.—*Funaria*.—A, Apex of female gametophyte with archegonia. B, Neck of archegonium, showing mode of separation of the cap cells. C, Immature archegonium. (From Small after Sachs.)

division until an elongated stalk, bearing upon its summit a capsule, is finally produced, this entire structure being known as the *sporogonium*. The base of the stalk remains imbedded in the basal portion of the archegonium, at the tip of the leafy stalk, and forms a *foot* or absorbing process. In growing upward the sporogonium ruptures the neck of the archegonium and

carries it upward as the covering of the capsule, or *calyptra*. The calyptra is thrown off before the spores are matured within

FIG. 322.

FIG. 323.

FIG. 322.—*Funaria hygrometrica*. A, Young sporophyte (*c*) attached to gametophyte (*g*). B, Adult sporophyte with seta (*s*) and capsule (*f*); gametophyte (*g*), part of which forms the calyptra (*o*). C, Vertical section of capsule ($\times 80$). Operculum (*d*); Peristome (*p*); Annulus (*a*); Air-space with trabeculae or filaments (*t*); Sporogenous layer (*s*). (From Small, after Prantl.)

FIG. 323.—Surface view of peristome of *Funaria*, after removal of operculum. *p*, peristome tooth. (After Small.)

the capsule. The upper part of the capsule becomes converted into a lid or *operculum* at the margin of which an *annulus* or ring of thick-walled cells forms. Attached to the annulus by their bases

FIG. 324.—Diagram of the life history of a moss. (After Atwood.)

are a number of long triangular teeth called collectively the *peristome*. The cells of the annulus are hygroscopic and expand

at maturity, throwing off the lid. The *peristome teeth* are very hygroscopic. During rainy weather or when the atmosphere is very humid they bend inward preventing the escape of the spores. During dry weather they straighten out and in this process lift out some of the spores and allow others to sift out from the cavity of the capsule. This completes the asexual or sporophyte generation. The spores falling to the damp soil germinate into protonemata, thus completing the life cycle in which is seen an alteration of generations, the two phases, gametophyte alternating with sporophyte.

CHAPTER XX

THE PTERIDOPHYTES

The Pteridophytes or Vascular Cryptogams constitute the third great division of the Plant Kingdom. They include the plants known as ferns, horsetails or scouring rushes, club mosses and quillworts. These are the most highly developed cryptogams, showing not only a distinct alternation of generations in their life histories, but a considerable structural advance over the Thallophytes and Bryophytes. They differ from the Bryophytes (1) by the sporophyte becoming an independent plant which shows differentiation into true root or roots, stem, and leaf or leaves each of which possesses a well organized vascular bundle system, (2) by the gametophyte body or prothallus nourishing the young sporophyte only until the latter has produced its own root and leaf after which the gametophyte perishes, and (3) by presenting large, conspicuous sporophytes and very small gametophytes.

Pteridophytes formed the chief vegetation of the earth during the Carboniferous period.

The Pteridophytes are grouped into three classes, viz.:

CLASS I. FILICINEÆ OR FERNS.

CLASS II. EQUISETINEÆ OR HORSETAILS.

CLASS III. LYCOPODINEÆ, THE CLUB MOSSES AND QUILLWORTS.

THE FERNS

The ferns, constituting the class *Filicineæ*, represent the largest group of the vascular cryptogams.

Ferns occur in a variety of situations in tropical and temperate countries. Most of them prefer damp, shady places. Some grow in forests and fields, others in the crevices of rocks, still others in water. In the tropics, some ferns become tree-like in form, possessing long erect trunks bearing at their summit a crown of compound leaves called *fronds*. In temperate regions,

the land ferns have underground stems or rhizomes from which fronds arise.

FIG. 325.—*Dryopteris marginalis*, a fern whose rhizome and stipes constitute the drug, American Aspidium.

The main axis of the majority of the typical land-growing ferns is a creeping underground stem or *rhizome* which at its various nodes bears *rootlets* below and *fronds* above. These fronds are highly developed, each being provided with a petiole-

like portion called a *stipe* which is extended into a *lamina* usually showing a forked venation. Some ferns possess *laminæ* which are lobed, each lobe being called a *pinna*. If a pinna be further divided, its divisions are called *pinnules*. In some ferns the pinnules are further subdivided into *secondary pinnules*, as in the "brake" or "bracken," *Pteris aquilina*. The unfolding of a frond is *circinate* and it increases in length by apical growth. On the under surface or on the margin of the *laminæ*, *pinnae*, or *pinnules* may be seen small brown patches each of which is called a *sorus*, and usually covered by a protective membrane called the *indusium*. Each *sorus* consists of a number of *sporangia* (spore cases) developed from epidermal cells. In some ferns the entire leaf becomes a spore-bearing organ (*sporophyll*). Many *sporangia* have a row of cells around the margin, the whole being called the *annulus*. Each cell of the annulus has a U-shaped, thickened cell wall. Water is present within these cells and when it evaporates it pulls the cell walls together, straightening the ring and tearing open the weak side. The annulus then recoils and hurls the spores out of the sporangium. Upon coming into contact with damp earth, each spore germinates, producing a green septate filament called a *protonema*. This later becomes a green heart-shaped body called a *prothallus*. It develops on its under surface *antheridia* or male organs and *archegonia* or female organs as well as numerous *rhizoids*. Within the antheridia are developed motile, many ciliated *sperms*, while *ova* are produced within the archegonia. The many ciliate sperms escape from the antheridia of one prothallus during a wet season and, moving through the water, are drawn by a chemotaxic influence to the archegonia of another prothallus, pass down the neck canals of these and fuse with the ova, fertilizing them. The fertilized egg or oöspore divides and redivides and soon becomes differentiated into a *stem*, *first leaf*, *root*, and *foot*. The foot obtains nourishment from the prothallus until the root grows into the soil, when it atrophies, and the sporophyte becomes independent. Unequal growth and division of labor continue until a highly differentiated sporophyte results, the mature "fern plant."

In some species of ferns as the Cinnamon Fern (*Osmunda cinnamomea*) and the Sensitive Fern (*Onoclea sensibilis*), two kinds

of fronds are produced, *sterile fronds* which represent green foliage leaves and *fertile fronds* which contain little or no chlorophyll and bear clusters of sporangia.

There are two sub-classes of Filicineæ, viz.:

SUB-CLASS EUSPORANGIATÆ, including ferns like *Ophioglossum* or adder's tongue and *Botrychium* or moonwort in which the sporangium originates from a group of epidermal cells and

FIG. 326.—Fern leaves, showing various degrees of subdivision or branching of the blade. A, *Phyllitis*; B, *Polypodium*; C, *Pteris*; D, *Adiantum*. (From Gager.)

contains similar spores, the latter giving rise to prothallia which bear both antheridia and archegonia.

SUB-CLASS LEPTOSPORANGIATÆ which include most of the common ferns and the water ferns. In these the sporangium develops from a single epidermal cell. Most of them have sporangia containing similar spores but the water ferns are heterosporous.

ORDER 1.—FILICALES OR TRUE FERNS (HOMOSPOROUS)

FAMILY CYATHEACEÆ.—Mostly tree-like ferns of the tropics and subtropics possessing sporangia with a complete annulus.

The long, silky, yellow or brownish hairs obtained from the bases of *Cibotium glaucum*, *C. Barometz*, *Alsophila lurida*, and other members of this family growing in Java and Sumatra have been successfully used in checking external hemorrhages under the names of "*Pengawar Djambi*" or "*Golden Moss*."

FAMILY POLYPODIACEÆ.—In this family the capsule of the sporangium possesses a vertical, incomplete annulus which splits

FIG. 327.—Cross section of the main axis or rhizome of the common brake, *Pteris aquilina*. $\times 8$. *ep*, epidermis; *scl*, sclerenchyma fibers in hypodermis; *b*, xylocentric vascular bundles; *sclb*, sclerenchyma band composed of lignified fibers; *r*, lateral ridges; *pa*, parenchyma; *x*, xylem; *si*, sieve tissue (phloem); *end*, endodermis; *ph*, phloem of a vascular bundle.

the mature sporangium transversely. It includes most of the more common land ferns of temperate regions. Outstanding genera of this family are *Pteris* or brake, *Polypodium* or polypody, *Adiantum* or maiden hair, *Asplenium*, or spleenwort, *Dryopteris* or wood fern, *Camptosorus* or walking leaf, *Polystichum* or christmas fern, *Pellaea* or cliff brake and *Phegopteris* or beech fern.

The rhizomes and stipes of *Dryopteris Filix-mas* and *Dryopteris marginalis* are valuable tape worm remedies. Those of the former species are official under "Aspidium" in the U.S.P. The vascular bundles of the stems and leaves of these and other

ferns are *concentric* in type but differ from the concentric vascular bundles of the stems and leaves of some monocotyledons in that xylem is innermost and phloem surrounds the xylem. This kind of concentric bundle is called a *xylocentric bundle*. For further details on fern structure see Chapter IV. The dried fronds of the native *Adiantum pedatum* and of the European *A. Capillus-Veneris* are used in domestic medicine as a demulcent expectorant in the form of syrup or infusion.

FAMILY OSMUNDACEÆ.—This family is characterized by having short sporangia, the annulus of which consists of a group of

FIG. 328.—*Cyrtomium falcatum*. Under (dorsal) surface of a portion of a sporophyll, showing the numerous sori on the pinnae. (Gager.)

thick-walled cells near the apex on either side of the wall of the sporangium, the rest of the wall becoming gradually thinner walled. The mature sporangium splits open longitudinally. It is represented in North America by the genus *Osmunda*, three species of which, *O. regalis*, the royal fern, *O. cinnamomea*, the cinnamon fern and *O. claytoniana*, the interrupted fern, occur locally. The rhizomes and stipe bases of the last named have frequently been found as substitutes and adulterants of *Aspidium*.

ORDER 2.—HYDROPTERALES OR WATER FERNS (HETEROSPOROUS)

FAMILY SALVINIACEÆ.—Floating ferns with broad floating leaves and submerged dissected leaves which bear fruit bodies called sporocarps containing microsporangia and megasporangia. The microsporangia contain microspores which give rise to male prothallia. The megasporangia contain megaspores which

develop female prothallia. Examples: *Salvinia*, *Marsilea* or the Water Shamrock, and *Azolla*.

THE HORSETAILS

The *Equisetineæ*, commonly known as the Horsetails or Scouring Rushes, are perennial herbs with hollow, cylindrical, jointed and fluted stems, sheath-like whorls of united leaves and terminal cone-like fructifications. Their bodies contain large amounts of silicon, hence the name scouring rushes. Only 25 species exist today and all of these belong to the genus *Equisetum*. These are believed to represent degenerate descendants of tree-like Horsetails, fossils of which have been found in coal-bearing rocks of the Carboniferous period.

In some species of *Equisetum* the fruiting cone is borne on the ordinary aerial stem, in others on a special stem of slightly different form. The spores are provided with ribbon-like appendages called *elaters*, which, being hygroscopic, coil and uncoil with increase or decrease in the amount of moisture present, thus aiding in the entanglement of the spores into small masses, and so preventing isolation, an advantage on account of the dioecious character of the prothallia which they originate.

LIFE HISTORY OF *EQUISETUM ARVENSE*, A HORSETAIL

Equisetum arvense, a frequently occurring species along railroad embankments but also in fields, may be taken as a type form of the group. Like other horsetails its life history involves an alternation of sporophyte and gametophyte generations of which the sporophyte is the more conspicuous.

The sporophyte consists of a subterranean, horizontally-branched, food-storing *rhizome* (main axis) bearing on its lower surface water- and mineral nutrient absorbing-roots and on its upper surface 2 kinds of jointed, hollow shoots—*sterile* and *fertile*. The *sterile shoots* are green and bear whorls of shortened, green branches from their upper nodes and scale-like leaves from their lower nodes. Theirs is the work of manufacturing food for the plant. The *fertile shoots* are nearly or completely devoid of chlorophyll and bear whorls of scale-like leaves at their nodes and terminate in a *cone*. The central stalk of this cone is a prolonga-

tion of the shoot axis and bears numerous whorls of *sporophylls*. Each sporophyll consists of a short stalk and a disk; the latter is arranged in peltate fashion on the end of the stalk and bears a number of *sporangia* on its lower surface. Within these sporangia are produced numerous *spores* which are liberated upon the

FIG. 329.—*Equisetum arvense*. *P*, sterile branch; *P'*, fertile branch with strobilus, or cone; *R*, rhizome (underground); *T*, cross-section of cone, showing insertion of sporophylls in a whorl; *N*, *N'*, sporophylls with pendant sporangia; *S*, *S'*, *S''*, spores with coiled elaters (*el*). (Gager.)

ripening of the sporangia. While alike as to size, some of them give rise upon germination to *male prothallia*, others to *female prothallia*. These prothallia represent the sexual or gametophyte plants. The male prothallia produce *antheridia* along the margins or tips of their lobes in which, when mature, numerous many-ciliated *antherozoids* are found. The female prothallia produce

flask-shaped *archegonia* on the prothallial cushion each of which contains an *egg*. Both kinds of prothallia bear absorptive *rhizoids* on their lower surface. The antherozoids, liberated from the antheridia of male prothallia, swim to the archegonia of neighboring female prothallia, enter their mouths, pass down their neck canals and gather about the eggs. In each instance the best adapted one fertilizes the egg. The fertilized egg (*oöspore*) develops into an *embryo* which through continued growth becomes a *mature sporophyte* plant.

THE CLUB MOSSES

The Class *Lycopodineæ* or club mosses consists of small perennial, vascular, dichotomously-branched, trailing, evergreen herbs with stems thickly covered with awl-shaped leaves. They are represented, today, by over 900 extant species. These are relatives of giant tree-like species of club mosses such as *Lepidodendron* and *Sigillaria* which with the giant horsetails and ferns formed the chief flora during the Coal Age. They differ from ferns and horsetails in possessing a *suspensor*, like seed plants, in their early sporophyte stage and in having *biciliate sperms*. Except in a few instances the sporangia are borne on *sporophylls* crowded together and forming cones or spikes (*strobili*) at the ends of erect branches or as lateral branches. The spores found within the sporangia are either alike or unlike in size, hence the plants are either *homosporous* or *heterosporous*. In heterospory the small spores give rise to male gametophytes and the large spores to female gametophytes.

FAMILY I.—LYCOPODIACEÆ, including the single genus *Lycopodium* with widely distributed species. The group is homosporous. The spores of *Lycopodium clavatum* are official.

LIFE HISTORY OF LYCOPODIUM CLAVATUM.—This species of club-moss thrives in temperate and sub-tropical forests of both hemispheres and yields a light-yellow, mobile powder consisting of spores. This powder is of value in pharmacy to prevent the adhering of suppositories, pills, etc., and in medicine as a protective to tender surfaces. When examined under the microscope the spores appear tetrahedral with one convex and three plane surfaces. The outer wall (*exosporium*) of each is reticulated,

FIG. 330.—*Lycopodium clavatum* L. A, portion of sporophyte plant showing habit (natural size); a, foliage leaf; b, sporophyll, with kidney-shaped sporangium at base (magnified); c, d, e, spores viewed from different aspects ($\times 580$); B, prothallus of another species. When the sporophylls are mature, the spikes are cut and dried, and the spores separated by shaking. They are freed from debris by sieving through flour-sieves. (After Luerssen.)

showing meshes that are surrounded by polygonal ridges with straight sides. In nature, these spores, falling out of the ripe spore cases, germinate on moist soil, each producing a fleshy *prothallus* which is submerged in its lower portion and exposed above ground in its upper part. This structure has *antheridia*

FIG. 331.—*Selaginella Martensii*. *a*, vegetative branch; *b*, portion of the stem, bearing cones (*x*); *c*, longitudinal section of a cone, showing microsporangia (*mic. sp.*) in the axils of microsporophylls, and megasporangia in the axils of megasporophylls; *d*, microsporangium with microsporophyll; *e*, microspores; *f*, portion of wall of sporangium, greatly magnified; *g*, megaspore; *h*, microsporangium opened, and most of the microspores scattered; *i*, megasporangium, with megasporophyll; *k*, same, opened, showing the four megaspores. (Gager.)

and *archegonia* imbedded in its upper end and shows elongations of its epidermal cells as *rhizoids* in its tuberous, sunken, lower part which absorb nutriment from the soil. The mature antheridia burst open during wet weather and liberate biciliate *antherozoids*. Each mature archegonium produces an *ovum* in its basal cavity. Antherozoids (*sperms*) swim to the mouth of the archegonium, pass down its canal and gather about the ovum.

One of these fertilizes the ovum forming an *oöspore* (*zygote*). The *oöspore* begins to divide and ere long develops into an embryo which soon becomes differentiated into *foot* and *shoot* regions. The foot absorbs nourishment from the prothallus, for a short time, until the first root commences to function. The *shoot* elongates carrying the first leaves above the soil and giving rise to the first root or *radicle* at its base. Through continued growth and differentiation a mature *sporophyte* is developed which consists of a creeping, prostrate, dichotomously branching stem covered with crowded, light green, linear, awl-shaped, and bristle-tipped leaves and bearing several ascending fertile branches which terminate in one or two *spikes*. Each of these spikes consists of a slender *axis* bearing numerous more or less deltoid, bristle-tipped *scales*, each one having a kidney-shaped *sporangium* at the base of its upper surface containing numerous *spores*.

FAMILY II.—SELAGINELLACEÆ (Little Club Mosses), including the single genus *Selaginella* with species for the greater part tropical. Plants similar in habit to the *Lycopodiaceæ* but producing microspores and megaspores, hence showing heterospory. The microspores are formed in microsporangia which are borne on microsporophylls, the megaspores in megasporangia borne on megasporophylls. (See Fig. 331.) The microspores produce male prothalli and the megaspores, female prothalli or gametophytes. The selaginellas are beautiful moss-like plants cultivated for ornament. One of them, *Selaginella lepidophylla*, a native of Mexico, is a "resurrection plant," curling up if allowed to dry and, when apparently dead, capable of being revived in water.

FAMILY III.—ISOETACEÆ (Quillworts) including the single genus *Isoetes* whose species are plants with short and tuberous stems giving rise to a tuft of branching roots below and a thick rosette of long, stiff, awl-shaped leaves above. They produce microspores and megaspores and so show heterospory.

CHAPTER XXI

THE SPERMATOPHYTES

This is the highest evolved division of the vegetable kingdom. It includes all plants which produce real flowers and seeds. The older name for the group is Phanerogamia. In this group the sporophyte attains its greatest development while the gametophytes are reduced to minute structures parasitic on the sporophyte. The Spermatophytes are subdivided into two subdivisions based primarily upon whether their seeds are borne naked or enclosed within a seed vessel or pericarp.

Subdivision I. Gymnospermæ (Gymnosperms)—plants with naked seeds.

Subdivision II. Angiospermæ (Angiosperms)—plants with covered seeds.

THE GYMNOSPERMS

The Gymnosperms comprise an ancient and historic group of seed plants which were more numerous in the Triassic and Carboniferous periods than now. They differ from the Angiosperms in several respects, viz.: they bear naked ovules on the edges or flat surfaces of leaves called carpels, while Angiosperms bear covered ones; each megaspore produces within itself a bulky prothallus, in the upper portion of which originate one or more archegonia, while in Angiosperms no recognizable prothallus has been proven to exist; the stored food tissue within their seeds is prothallial tissue loaded with starch, etc., while that in Angiosperm seeds (endosperm) is developed from the endosperm nucleus; the mode of growth of their stems is always indefinite while that of Angiosperms is either indefinite or definite.

The groups still extant are the Cycads or Fern Palms, the Gnetums, the Ephedras, the Ginkgos and the Conifers. Of these the Conifers comprising over 300 species are the most numerous. Many of them yield valuable products to pharmacy and the arts.

The Conifers include the pines, spruces, hemlocks, cedars, firs, redwoods, yews, arbor vitæ, chamæcyparis, and larches. All of their number are evergreen except the European larch, which drops its foliage upon the advent of winter.

FIG. 332.—The White Pine (*Pinus Strobus*).

LIFE HISTORY OF A GYMNOSPERM (*PINUS STROBUS*)

The White Pine frequently called the Weymouth Pine (*Pinus Strobus*), one of the principal timber trees of the Northern States and Canada, is also of value in pharmacy and medicine. The inner bark of its trunk and branches is used because of its valuable expectorant properties and is official in the National Formulary.

DESCRIPTION OF THE WHITE PINE TREE (MATURE SPOROPHYTE)

From an underground spreading root system there arises an erect aerial *trunk* or stem that extends from the ground to the apex of the tree, ending in a terminal bud. The trunk rarely exceeds 3 feet in diameter and 125 feet in height and is averagely $1\frac{1}{2}$ to

FIG. 333.—Transverse section of white pine stem of four years' growth, showing cork (*a*), cortex (*b*), phloem (*c*), cambium (*d*), xylem (*e*), secretion reservoir (*f*), pith (*g*) and medullary-ray (*h*). (Photomicrograph.) $\times 16$.

3 feet in diameter and 50 to 90 feet high. At a varying distance above the soil, depending upon environal conditions as well as the age of plant, whorls of lateral branches (three to seven in a whorl) are seen emanating from the trunk, in horizontal fashion, at various levels up to near the apex. These become, under conditions prevalent when the tree is grown in the open, gradually shorter until the summit is reached, giving to the *crown* or upper part of the tree the appearance of a pyramid. These

branches give rise to other branches which agree with the lateral branches in ending in terminal buds. Another kind of branch, however, is found which is always shorter than the former branches. This type of branch is called a "*spur shoot*" and arises from the former branches. The spur shoots bear the *needles* or

FIG. 334.—Transverse section of the needle (leaf) of the white pine. Highly magnified. *ep*, epidermis; *s*, stomata; *m*, infolded parenchyma cells of mesophyll; *c*, cuticle; *r*, oil reservoir; *p*, colorless central parenchyma corresponding to the pith of the stem; *n*, endodermis; *x*, xylem and *ph*, phloem of collateral vascular bundles. The xylem is composed of tracheids with bordered pores. The leaf shows xerophytic-centric structure.

foliage leaves which are light-green, when young, and bluish-green, soft, flexible, $2\frac{1}{2}$ to 5 inches long, when mature. The "needles" occur in tufts (*fascicles*) of five, are triangular in cross-section, have finely *serrate* (*saw-toothed*) edges and are surrounded at the base by a deciduous sheath. These foliage leaves persist until the end of their second year, when they are shed with the spur shoot which bears them.

The white pine, like most of its allies among the Coniferæ, bears cones. These structures are of two kinds, viz.: staminate and carpellate. Both kinds are produced on the same tree.

STAMINATE CONES.—The yellow, ovate, staminate cones appear about May and are clustered at the base of the new growth of the current season. Each consists of a main axis (modified branch) which bears spirals of scales called *microsporophylls* or stamens. On the under surface of each scale are the

FIG. 335.—Staminate cones of the Austrian pine (*Pinus austriaca*). Below, before shedding pollen; above, after shedding. (Gager.)

spore-cases (*microsporangia*), which develop the *microspores* (*pollen grains*). A section of the microsporangium shows it to consist of a wall beneath which is a layer of cells called the *tapetum* and in the interior a large number of spheroidal *spore mother-cells*. Each of the spore mother cells divides into four spores. During this process, reduction division takes place and the chromosomes are reduced to one-half the number, so that the spores are *haploid* as those of the fern. Each pollen grain when mature consists of a central fertile cell and a pair of air-sacs or wings, one on either side of the fertile cell. The purpose of the latter is to give greater buoyancy in the air to the microspore.

CARPELLATE CONES.—The young carpellate cones appear in May or early June as pinkish-purple structures arranged in solitary fashion or in small groups, lateral along the new growth. Each terminates a lateral axillary branch. A carpellate cone is composed of a main axis which bears spirals of scales, termed

FIG. 336.—Scotch pine (*Pinus sylvestris*). A-D, stages in the development of the carpellate cone, and its carpotropic movements. E, very young carpellate cone much enlarged; F, ventral, G, dorsal views of a scale from E; 1, ovuliferous scale, 2, ovule (in longitudinal section); 3, pollen chamber and micropyle leading to the apex of the nucleus (megasporangium); 4, integument of the ovule; G, 1, tip of ovuliferous scale; 5, bract; 4, integument; H, longitudinal section at right angles to the surface of the ovuliferous scale (diagrammatic); 6, megaspore; 7, pollen chamber; I, longitudinal section of a mature cone; 6, ovule; J, scale from a mature cone; 6, seed; w, wing of seed; K, dissection of mature seed; h, hard seed coat; c, dry membrane remains of the nucellus, here folded back to show the endosperm and embryo; e, embryo; p, remains of nucellus; L, embryo; c, cotyledons; e, hypocotyl; r, root-end. (Gager.)

megasporophylls or *carpels*. Each *megasporophyll* is composed of an ovuliferous scale bearing two *ovules* or *megasori* and a bract. Each megasorus (ovule) is attached to the surface of the megasporophyll. It contains a *nucellus* or *megasporangium* which is surrounded by an integument, except at the apex where an opening, the *micropyle*, is evident. The micropyle is the gateway to the *pollen chamber* which lies below it. Within the nucellus a single spore

mother cell is formed. This divides to form four spores but only one persists. This is called the *megaspore* or *embryo sac*. It contains haploid chromosomes.

DESCRIPTION OF THE GAMETOPHYTE GENERATION

The Gametophyte generation of the White Pine begins with the development of the male and female gametophytes and terminates with the fertilization of the egg.

FIG. 337.—Mature carpellate cones of white pine showing separated scales.

THE MALE GAMETOPHYTE.—The male gametophyte commences to form in the mature pollen grain before the pollen is shed. A series of three nuclear divisions takes place which cut off two small prothallial cells (traces of one of which may be seen pushed up against the wall of the fertile cell of the pollen grain), a tube nucleus and a generative cell. (See Fig. 338.) At this stage the pollen is shed and some of it is carried by air currents to the carpellate cones where it sifts in between the ovule-bearing scales and accumulates at the scale bases. A number of the pollen grains are drawn close to the nucellus of the ovule by the drying up of the viscid fluid which fills the *pollen chamber*. In this

fluid they germinate forming pollen tubes. The transfer of pollen grains from the pollen sac to the pollen chamber and consequent germination therein is called *pollination*. The contents of a mature pollen-grain constitutes the male gametophyte.

FIG. 338.—Pine Pollen and development of pollen tube. In *A* the pollen grain is shown containing prothallial cells (*p*). They have been cut off before the grain was shed. The tube nucleus (*T*) has been divided off from the generative nucleus (*G*).

In *B* the tube has been put forth and the tube nucleus has moved down into it. Between the stages represented in *B* and *C* the generative nucleus has divided to form a stalk nucleus (*ST*) and a body nucleus (*B*).

In *C* four nuclei are present in the pollen tube,—the tube nucleus (*T*), the stalk nucleus (*ST*), and the two sperm nuclei (*S*). The sperm nuclei have been produced by a division of the body cell, the latter having disappeared. (After Atwood.)

THE FEMALE GAMETOPHYTE.—If the embryo-sac be examined at about the time of pollination, it will be found to consist of a single cell containing a single nucleus surrounded by cytoplasm. Very shortly afterward, however, the nucleus divides repeatedly to form a large number of nuclei which are scattered throughout the cytoplasm. Each nucleus accumulates around itself a por-

tion of the cytoplasm and ultimately cell walls are laid down and the entire embryo-sac contains *endosperm* (prothallial) tissue. Toward the micropylar end of the endosperm (*prothallus*) originate several archegonia.

FIG. 339.—Longitudinal section of one-year old ovule of Austrian Pine showing internal structure. (Mottier.)

Each archegonium consists of a much-reduced neck of four cells and an egg (*ovum*) which lies embedded in the prothallus which forms a narrow layer of cells around it called the *jacket*. The contents of the mature embryo-sac constitutes the female gametophyte.

After pollination the carpellate cone gradually bends on its axis so as to ultimately reverse its position and hang downward.

FERTILIZATION.—About a year after pollination the pollen tubes, lying within the pollen chamber show signs of renewed activity. The tube nucleus passes to the tip of the tube. The

FIG. 340.—White pine (*Pinus Strobus*). Vertical section through the upper part of an ovule, shortly before fertilization. *s.n.*, sperm-nuceli; *st. c.*, stalk-cell; *t.n.*, tube-nucleus; *arch*, archegonium; *e.n.*, egg-nucleus. (Gager, after Margaret C. Ferguson.)

which it is attached. The embryo is nourished by a portion of the prothallus but the greater part of the prothallus forms the endosperm tissue of the seed surrounding the embryo. The thin nucellus persists as an endosperm covering. The integument becomes modified to form the hard protective seed coat. A portion of the scale of the cone directly above and adjacent to the ovule forms a membranous wing which separates from the scale as part of the seed.

generative-cell divides to form a body and a stalk-cell which pass into the tube. The body-cell later forms two sperm nuclei. While these changes are taking place the tube is penetrating the nucellus and growing toward the embryo sac with its contained female gametophyte. It finally enters it, passing between the neck-cells of the archegonium. The tip of the tube then breaks and the entire tube contents is emptied into the egg. One of the sperm nuclei fuses with the egg nucleus and fertilizes it forming an oöspore or zygote. The oöspore contains the diploid number of chromosomes.

SEED FORMATION AND DISTRIBUTION.—The oöspore undergoes repeated divisions and forms the *embryo* or young sporophyte plant and a *suspensor* to

By this time (about two years after pollination) the scales of the cone, now quite woody, separate, the seeds are shaken out, and many are carried for a considerable distance by winds.

FIG. 341.—Austrian Pine (*Pinus nigra*). Longitudinal section of the upper end of the endosperm showing two archegonia. The one at the right is mature and ready for fertilization; that at the left is younger; the ventral canal cell is being cut off. (Mottier.)

GERMINATION OF THE SEED.—Under favorable conditions, the seeds absorb water and germinate in the spring following their dispersal. The hypocotyl of the embryo appears first, arching upward and downward, and, straightening out, draws the green cotyledons with it, which spread out toward the light, while it

OUTLINE OF LIFE HISTORY OF PINUS STROBUS

1. Embryo is in a *seed structure* composed of a seed coat (developed from the integument of the ovule, "megasorus") enclosing perisperm (developed from megasporangial tissue, the nucellus). The perisperm surrounds endosperm (prothallial tissue developed in embryo sac) and within the endosperm is the embryo which was developed from a fertilized ovule within one of the archegonia. The seed wings are developed from a portion of the scale leaf (ovuliferous scale or megasporophyll).

2. Mature Sporophyte consists of a root system, trunk, lateral branches, spur shoots bearing foliage leaves in fascicles. Staminate cones are borne at base of new growth. Carpellate cones are borne at end of a lateral axillary branch.

3. Megasorus or ovule is composed of an integument surrounding the megasporangium (nucellus) except at apex where the micropyle (opening) leads to pollen chamber.

4. Male Gametophyte—consists of the mature pollen grain containing two prothallial cells, a tube nucleus and a generative cell.

5. Female Gametophyte consists of prothallial tissue (endosperm) with several archegonia, each containing an ovum. It is the mature "embryo sac."

grows into the soil to form the tap root and in time the remainder of the root system. Thus the seedling sporophyte is formed which in time develops into the mature White Pine tree.

FIG. 342.—Diagram of life-cycle of a pine. (From Gager after Schaffner.)

CLASSIFICATION OF THE GYMNOSPERMS

The gymnosperms are grouped into 4 orders, as follows:

- Order 1. Cycadales, the Cycads.
Order 2. Ginkgoales, the Ginkgos.
Order 3. Coniferales, the Conifers.
Order 4. Gnetales, the Gnetums and Ephedras.

ORDER CYCADALES

The Cycads or fern palms, the most primitive of the seed plants, are represented by about 100 species of fern-like plants that are believed to have been descended from the *Cycadofilices*, a group of fossil plants which possessed the leaf characters of ferns and ovules and seeds like the pines. Most of them have plant bodies consisting of a thick cylindrical stem covered by the bases of spirally arranged leaves and bearing upon its summit leathery, pinnately-compound leaves and scale leaves. The male and female cones are borne on different sporophytes. They are represented by the genera *Cycas*, *Zamia*, and *Dioon*.

LIFE HISTORY OF ZAMIA

Zamia, a representative of the Cycads, possesses a plant body or sporophyte which is short with a cylindrical erect stem con-

FIG. 343.—*Cycas revoluta*. The trunk of this cycad stores an enormous amount of starch. It is one of the sources of Sago starch of commerce. Note the crown of pinnately-compound leaves.

taining a large central pith which is surrounded by a circle of open collateral bundles and by a broad cortex in whose cells great amounts of starch are stored. The summit of the stem

is crowned by a cluster of pinnately-compound leaves of a leathery texture.

STAMINATE CONE.—The staminate cone of *Zamia* arises on the ends of short branches emanating from the end of the stem of

FIG. 344.—Mature sporophytes of *Zamia* bearing staminate strobili (cones). $\times \frac{2}{3}$.
(Photograph by F. J. Bacon.)

the male plant (*sporophyte*). Each consists of a central axis bearing compactly arranged *microsporophylls*. Each microsporophyll bears on its lower surface a large number of *microsporangia*. Within each microsporangium *spore mother cells* are developed, each of which by division form 4 *microspores* (*pollen grains*).

CARPELLATE CONE.—The carpellate cone arises on a short branch arising slightly from the end of the main stem of the female plant. It consists of a fleshy central axis bearing peltate *megasporophylls*. Each megasporophyll bears on the inner surface of its expanded outer portion two *megasori* or *ovules*. Each megasorus when mature consists of a *megasporangium* or *nucellus* surrounded by a single *integument*, the latter being perforated by a

narrow tubular opening or *micropyle* leading to a small cavity or *pollen chamber* at the end of the nucellus. (See Fig. 348.)

Within the nucellus is developed one *megaspore mother cell* which by division gives rise to 4 *megaspores* or *embryo sacs*. Only one of these persists, the other three perish.

GAMETOPHYTES.—The *male gametophyte* begins to develop from the young pollen grain (*microspore*) while the latter is still within the *microsporangium*. The pollen grain divides to form 2 cells, a smaller one or *prothallial cell* and a *larger cell*. The larger cell then divides to form a *generative cell* and a *tube cell*. There results a 3-celled, undeveloped *male gametophyte* or *mature pollen grain*.

FIG. 345.—A, staminate cone of *Zamia*. B, microsporophylls bearing on their lower surfaces many microsporangia. (From "A Textbook of General Botany" by Smith, Overton et al. Macmillan Co., Publishers.)

These are liberated by the bursting of the wall of the microsporangium of the now ripened male cone with its separated microsporophylls, and are carried by winds. Some are carried to the ripened carpellate cones of female *Zamias*. Some of the pollen grains sift between the megasporophylls and come to rest about the ovules. Some of them become caught in a mucilaginous fluid at the outer end of the micropyle and are sucked into the pollen chamber. Upon reaching the pollen chamber they resume their development. Each pollen grain germinates

FIG. 346.—*Zamia floridana* bearing an immature carpellate strobilus. $\frac{1}{6}$ actual size. The starch obtained from the stem of this plant is known as Florida arrow-root. (Photograph by F. J. Bacon.)

through the elongation of its tube cell to form a *pollen tube* which grows into the nucellus (megasporangium). Then, the generative cell divides into a *stalk cell* and a *body cell*. The body cell then divides to form 2 top-shaped antherozoids, each of which bears many cilia along a spiral groove. *The mature pollen tube and its contents represents the male gametophyte of Zamia.*

FIG. 347.—A, mature carpellate strobilus (cone) of *Zamia*. B, megasporophyll bearing two megasporangia. C, lengthwise section of a young megasporophyll, showing an early stage in the development of female sexual plants or megagametophytes. D, similar section of a later stage; megagametophytes fully matured. (From "A Textbook of General Botany" by Smith, Overton & Co. by permission from Macmillan Co., publishers.)

The *female gametophyte* develops within the *megaspore* (embryo sac) of the nucellus of the ovule by a series of nuclear divisions of the remaining nucleus within the embryo sac, resulting in many scattered nuclei within the cytoplasm. Cell walls are laid down and many cells result, and the embryo sac increases in size forming a *prothallus*. The nucellus is encroached upon by the enlarged developing gametophyte or prothallus in whose cells nourishment is stored. An *archegonial chamber* appears between the end region of the prothallus and nucellus, in line with the

pollen chamber and micropyle. Two to six archegonia are developed from the prothallial tissue just beneath the archegonial chamber. Each *archegonium* consists of two neck cells protruded into the archegonial chamber and a large *ovum* embedded in the prothallus. *The contents of the mature embryo sac, therefore, represents the female gametophyte.*

FERTILIZATION.—It will be recalled that at the time pollination was effected, the male gametophyte¹ had assumed the char-

FIG. 348.—*Zamia*. Stages in the development of a megasporangium and a macrogametophyte (megagametophyte). A, at the time when the macrospores (megaspores) are formed. B, after the germination of one macrospore to form a macrogametophyte. C, at the time of gametic union. (From "A Textbook of General Botany" by Smith, Overton et al., the Macmillan Co., Publishers.)

acter of a pollen tube (see Fig. 348) containing a prothallial cell, stalk cell, and two antherozoids which lay in the pollen chamber. Several of these are usually present. Now, when the eggs are mature, the tube grows through the nucellus to the archegonial chamber and the end of the tube bursts. The antherozoids of the several tubes move about in the fluid of the chamber and some of these pass through the necks of the archegonia, one uniting with each egg. The sperm nucleus then unites with the egg nucleus forming a *zygote*.

SEED DEVELOPMENT.—The *zygote*, through repeated division, forms a *young sporophyte* or *embryo Zamia* in the upper end of the female gametophyte (*prothallus*) of the ovule. This, by the elongation of several cells, becomes differentiated into a slender

¹ *Microgametophyte* is another name for male gametophyte, and *macrogametophyte* a synonym for female gametophyte

suspensor attached to the anlage of the embryo which has been pushed through the basal membrane of the egg deep into the prothallus by the elongating cells of the suspensor. The embryo lives as a parasite on the tissue of the female gametophyte. After differentiation has proceeded in the body of the embryo to a certain point, *i.e.* where two *cotyledons* or seed leaves and a *central axis* consisting of a *hypocotyl* (part below the attachment of the cotyledons) and an *epicotyl* (part above the cotyledons) have been formed, growth of the embryo becomes suspended.

Nourishment is stored in the cells of the prothallus around the embryo forming the *endosperm*. The integument undergoes changes in its cells to form a firm *seed coat* consisting of a strong and a fleshy layer surrounding the nucellus. The *micropyle* persists as a tubular opening in the seed coat.

GERMINATION OF THE SEED.—The mature seeds drop from the megasporophyll of the mature cone to the ground and some of them germinate. In this process the seed absorbs water. The enzymes within its cells change the reserve or stored food within the endosperm into water soluble substances which are converted into new substances or protoplasm, as a result of which the

OUTLINE OF LIFE HISTORY OF ZAMIA

embryo grows. The hypocotyl, elongating, forces its way out through the micropyle and arches downward into the soil, most of it becoming the primary or main root. The upper part becomes the stem. The cotyledons remain within the seed. The epicotyl pushes its way upward and develops the rest of the stem, also the green leaves of the seedling sporophyte. As

FIG. 349.—*Ginkgo biloba*, the Maiden-hair tree.

soon as the green leaves and the root with its absorbing root hairs are developed, the seedling becomes an independent organism developing either into a mature male or female sporophyte.

ORDER GINKGOALES

The Ginkgoales is represented by the single species, *Ginkgo biloba*, the maidenhair tree, a native of western China and grown extensively in China and Japan and the United States as an ornamental tree. It possesses a straight trunk from which

spring long shoots bearing scattered leaves and short shoots with crowded leaves. Its leaves are fan-shaped, more or less 2-lobed, with furcately-branching veins and deciduous. It bears two kinds of cones (*strobili*) each kind on a different tree. The *staminate strobili* occur in clusters on the short shoots, each strobilus consisting of a central axis bearing a loosely arranged series of stamens (*microsporophylls*). Each stamen consists of a stalk ending in a knob and bearing on its underside two or more

FIG. 350.—Inflorescences of the pine. 1, Terminal twig; 2, ovulate cone; 3, staminate cone; 4, two-year-old cone. (Hamaker.)

microsporangia. The *ovulate (female) strobili* arise from the axils of leaves on the dwarf shoots of female trees. Each consists of an elongated stalk bearing near its end two *ovules* each of which has a small, cup-like *megasporophyll* at its base. The mature, *naked seed* resembles a drupe, the seed coat having an outer fleshy and inner stony layer.

ORDER CONIFERALES

Trees with a single upright stem which develops side branches that spread out horizontally and taper to a point at the summit, giving the crown of the tree the appearance of a huge cone, rarely

shrubs. This order includes two families, the *Pinaceæ* or Pine Family and the *Taxaceæ* or Yew Family.

Pinaceæ (*Coniferæ*) or *Pine Family*.—Trees or shrubs with resinous juice whose wood is characterized by being composed largely of tracheids with bordered pits. Leaves entire, awl- or needle-shaped, frequently fascicled, exstipulate, usually evergreen. Flowers, monoecious (Pines, etc.) or rarely dioecious (Juniper), achlamydeous, in cones. Staminate cone of a large number of microsporophylls (stamens) closely packed together and arranged spirally around a central axis, each stamen bearing usually two pollen sacs. Carpellate cone composed of spirally arranged scales, each of which bears a pair of naked ovules (megasori) near the base of its upper face. Fruit a cone with woody or fleshy scales (*Pinus*, *Thuja*, *Abies*, *Picea*, etc.) or a galbulus (*Juniperus*). Seeds albuminous. Embryo with two or more cotyledons.

The following genera of the pine family are of especial pharmaceutic and general economic interest:

GENUS		COMMON NAME	
Pinus		Pine	
Picea		Spruce	
Abies		Fir	
Larix		Larch	
Tsuga		Hemlock	
Pseudotsuga		Douglass Fir	
Thuja		Arbor vitae	
Juniperus		Juniper	
Sequoia		Redwood	
Cedrus		Cedar	
OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Terebinthina (Turpentine)	Concrete oleoresin	Pinus palustris and other species of Pinus	Southern United States
Resina (Rosin)	Resin	Pinus palustris and other species of Pinus	Southern United States
Oleum Terebinthinæ	Volatile oil	Pinus palustris and other species of Pinus	Southern United States
Pix Pini (Pine Tar)	Product of destructive distillation of wood	Pinus palustris and other species of Pinus	Southern United States
Oleum Picis Rectificatum	Volatile oil from pine tar		
Pinus Alba	Inner bark	Pinus Strobilus	United States and Canada

OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Oleum Pini Pumilionis	Volatile oil	<i>Pinus montana</i>	Tyrolese Alps
Juniperus	Fruit	<i>Juniperus communis</i>	} North America, Europe and Asia
Oleum Juniperi	Volatile oil	<i>Juniperus communis</i>	
Pix Juniperi	Empyreumatic oil	<i>Juniperus Oxycedrus</i>	So. Europe
UNOFFICIAL DRUG			
Sabina	Tops	<i>Juniperus Sabina</i>	Europe
Pix Burgundica	Resinous exudate	<i>Picea excelsa</i>	Europe
Terebinthina Canadensis	Liquid oleoresin	<i>Abies balsamea</i>	Northern United States and Canada
Terebinthina Laricis	Oleoresin	<i>Larix europæa</i>	Alps and Carpathians
Thuja	Leafy young twigs	<i>Thuja occidentalis</i>	United States and Canada
Oleum Abietis (Oil of Siberian Fir)	Vol. oil from fresh leaves	<i>Abies sibirica</i>	Russia and Siberia
Tamarack Bark	Stem bark	<i>Larix laricina</i>	Cold swamps of North America
Sandaraca	Resinous exudate	<i>Callitris quadrivalvis</i>	Africa
Dammar	Resinous exudate	<i>Agathis loranthifolia</i>	E. India
Kauri Gum	Resin	<i>Agathis australis</i>	New Zealand
Succinum (Amber)	Fossil resin	<i>Pinites succinifer</i>	Basin of Baltic
Bordeaux Turpentine	Concrete oleoresin	<i>Pinus maritima</i>	France
Pix Canadensis	Oleoresin	<i>Tsuga canadensis</i>	North America
Oregon Balsam	Oleoresin	<i>Pseudotsuga taxifolia</i>	Western United States and British Columbia
Spruce Gum	Gum	<i>Picea canadensis</i> <i>Picea mariana</i> <i>Picea rubra</i>	} Canada and New England
Oil of Cedarwood	Oil from wood	<i>Juniperus Virginiana</i>	
Hemlock	Bark	<i>Tsuga canadensis</i>	North America

TAXACEÆ OR YEW FAMILY.—Trees or shrubs with dark green, evergreen, exstipulate, linear leaves without resin ducts and with hard, close-grained wood of a reddish-color. The flowers are usually dioecious; the staminate globular with 4 to 8 stamens each having their anthers under a shield-shaped connective; the

pistillate consisting of a single terminal ovule with several bracts at its base. The fruit consists of an erect bony seed situated in a fleshy, reddish, cup-shaped disk, representing an aril.

FIG. 351.—*Ephedra* a member of the order Gnetales. A, branches with ovulate strobili. B, branches with staminate strobili. C, ovulate strobilus. D, an ovulate flower. E, staminate strobilus with staminate flowers in the axils of bracts. (From Watson.)

The leaves, seeds and bark of *Taxus canadensis* (American Yew) and *T. baccata* (European Yew) contain the alkaloid, taxin, which is poisonous to horses, cattle, sheep and pigs. The fleshy disk (aril) surrounding the seed is not poisonous.

FIG. 352.—*Ephedra sinica*. A, entire plant. B, portion of a green aerial stem from A, enlarged; n, node bearing two opposite scale-leaves (l); i, internode. C, fruiting branch showing terminal cone (f) and leaves (l). This species yields the commercial variety of Ephedra known in commerce as Tsaopen Mahuang and is gathered in Northern China.

ORDER GNETALES

The Gnetales comprise the single family, *Gnetaceæ*, which has three genera which in certain respects show a closer resemblance to the angiosperms than do other gymnosperms.

FIG. 353.—Cross section of a lower internode of the green stem of *Ephedra sinica*, a source of the drug Mahuang. *r*, ridge; note many of the ridges are papillated; *c*, cuticle of epidermis; *co*, cortex; *mf*, mesocortical fibers; *hf*, hypodermal fibers; *x*, xylem of an older (primary) bundle which has undergone secondary thickening; *p*, pericyclic fibers just beneath which is phloem, *ca*, cambium; *sx*, secondary xylem; *px*, end of protoxylem; *ph*, phloem of a secondary bundle; *td*, tracheids with bordered pits; *tr*, trachea; *m*, pith. Note the narrow medullary rays separating the open collateral bundles.

Family Gnetaceæ. The Gnetum family includes the genera *Ephedra*, *Welwitschia* and *Gnetum*, each genus representing a different type of plant. The *Ephedras* include about 30 species of straggling shrubs or low bushy plants of warm dry regions of the northern hemisphere, a number of species of which are used medicinally because of their ephedrine content. *Welwitschia*

contains the single species *Welwitschia mirabilis*, a low growing plant of arid regions in southwest Africa. The genus *Gnetum* contains about 15 species of woody climbers native to the tropics of both hemispheres. The principal features of the *Gnetum* family are, as follows:

Stems either unbranched and tuberous (*Welwitschia*) or branched and shrubby (*Ephedra*) or often climbing (*Gnetum*) and

FIG. 354.—Transverse section through the outer region of a representative portion of the stem of *Ephedra sinica*. $\times 400$. *c*, cuticle; *ep*, epidermis; *h*, hypodermal fibers; *ch*, chlorenchyma of cortex, the outer layers of which are palisade-like; *f*, mesocortical fibers; *td*, tracheids and *tr*, trachea of secondary xylem; *st*, stoma.

possessing open-collateral fibrovascular bundles. Resin canals are absent while true vessels (tracheæ) are present in the secondary xylem (distinction from conifers). Tracheids also occur in the xylem. Internodes and nodes are conspicuous in *Ephedra* and *Gnetum*. Leaves usually small, often reduced to scale-like structures and arranged in opposite or verticillate fashion. Flowers nearly always dioecious and exhibiting a membranous perianth of 2- to 4-bracts. Perianth of male flowers surrounding 2 to 8 stamens. Female flowers showing an erect ovule with 1 to

2 integuments and a bract-like perianth. Fruit in *Ephedra* a reddish, somewhat globular, galbulus-like structure (syncarp). Seed albuminous, the embryo with two cotyledons imbedded in endosperm.

The drug EPHEDRA consists of the green stems of *Ephedra sinica*, *Ephedra equisetina*, *E. distachya*, *E. Gerardiana* or other species of *Ephedra* containing the alkaloid *ephedrine*. These species are largely Asiatic in distribution. The United States Pharmacopoeia recognizes the alkaloid ephedrine and two of its salts, ephedrine sulfate and ephedrine hydrochloride. Preparations of these are official in the National Formulary. Ephedrine and its salts are used locally to shrink mucous membranes in colds and as a mydriatic. They are given internally for bronchial asthma, and certain circulatory conditions.

THE ANGIOSPERMS

The angiosperms comprise a large assemblage of seed plants which are more highly developed than those of the preceding group. Like the gymnosperms they possess flowers with sporophylls (stamens and carpels), but the floral axis has been shortened so that the sporophylls and the accessory floral leaves (sepals and petals), when present, are brought closer together on the axis, instead of being separated at strikingly different levels along the axis. The ovules, instead of being borne naked on a scale, are enclosed within the ovary and the seeds are covered by the ripened wall of the ovary which is called the pericarp or seed vessel.

LIFE HISTORY OF AN ANGIOSPERM (ERYTHRONIUM)

This attractive little plant, commonly called the Dog's Tooth Violet but related to the Lily, is found in the hollows of woods and may be seen in flower during the month of April in the Middle Atlantic States. It consists of an underground *stem* bearing *scales* (modified leaves) which is termed a *bulb*. From the lower surface of the bulb are given off numerous slender *rootlets* which penetrate the soil and from the upper surface, a pair of oblong, lance-shaped *leaves* of pale green color mottled with purple and white, and later, a *flower stalk* (*scape*), which bears upon its sum-

mit a single, yellow, nodding flower, which is often marked with purple stripes. The flower consists of a *torus* or *receptacle* which will be observed as the upper swollen end of the flower stalk (scape). Inserted upon it, passing from periphery toward the center, will be noted four whorls of floral leaves which, in order, are *calyx*, *corolla*, *androecium* and *gynæcium*. The calyx is composed of three lance-shaped and recurved, yellow parts called *sepals*; the corolla of three similarly looking parts called *petals* which

FIG. 355.—Dog's tooth violet (*Erythronium americanum*). Stages of development from the seed. 1–5 show the stage of development in each of five successive years. Full explanation in the text. 6, Bulb showing a surface bud (the sprout has been destroyed). (Gager after F. H. Blodgett.)

alternate in position with the sepals. Both of these whorls are collectively called the *perianth* or floral envelope. The androecium or male system of organs is composed of two whorls or circles of structures called *microsporophylls* or *stamens*. There are three *stamens* in each whorl. The outer whorl of stamens will be found opposite the sepals while the inner will be observed opposite the petals. Each stamen (microsporophyll) consists of an awl-shaped stalk or *filament* bearing upon its summit an oblong-linear body called an *anther*. The anther consists of two

lobes called *microsori*. Each lobe or *microsorus* contains two anther sacs or *microsporangia* in which, when mature, are to be found *microspores* or pollen grains. In the center of the flower will be noted the gynœcium or female system of organs. This, upon dissection, will be found to consist of three fused carpellary leaves termed *megasporophylls* (carpels) forming a somewhat flask-shaped structure called a *pistil*. The swollen basal portion of the

FIG. 356.—Cross section of the ovary of a lily formed by the coalescence of the lower portions of three carpels. $\times 70$. *s*, dorsal sutures; *ep*, epidermis; *c*, cuticle of epidermis; *b* bundle in mesophyll; *l*, locule; *ov*, ovule (anatropous); *pl*, infolded edges of adjacent carpellary leaves forming the axile placentas from which the ovules arise.

pistil is called the *ovary*; the stalk which arises from it is called the *style* and the knob-like viscid summit of the style is termed the *stigma*.

Microscopical examination of sections of the ovary will reveal it to be composed of three chambers called *locules*, within each of which are to be noted several inverted *ovules*. Each of these ovules is developed upon a nourishing tissue termed "*placenta*" which connects the ovules to the inner angle of the wall of the locule. The ovule is composed of a central prominent *mega-*

sporangium or *nucellus* which is almost completely invested by two upgrown *integuments* or coverings. The opening between the tips of the inner integument is called the *micropyle* (little gate). This is the gateway for the entrance of the pollen tube on its way to the nucellus. It is also the exit door for the hypocotyl of the embryo after the fertilized and ripened ovule becomes a seed. Within the nucellus, if the sections examined have been properly fixed, will be found a *megaspore* or *embryo sac*.

The *megasporangium* or nucellus is formed from a mass of meristem on the placenta. While in the state of differentiation, one of its cells is set apart as a *megaspore mother cell*. Its nucleus (containing $2x$ chromosomes) undergoes two successive divisions during the first of which a reduction in the number of chromosomes from the *diploid* ($2x$) to the *haploid* (x) number occurs. The x number persists in the ovum or egg.

DEVELOPMENT OF THE FEMALE GAMETOPHYTE THROUGH THE MATURATION OF THE EMBRYO SAC.—In its immature condition the *embryo sac* (*megaspore*) contains a mass of protoplasm surrounding a nucleus. This nucleus undergoes three divisions forming, as a result, eight nuclei which ultimately arrange themselves within the protoplasm of the embryo sac as follows: three of them occupy a position at the apex, the lower nucleus of the group being that of the egg or *ovum*, the other two nuclei being the *synergids* or assisting nuclei; at the opposite end of the sac three nuclei known as *antipodals* take their position; the two remaining nuclei called *polar nuclei* take up a position near the center of the embryo sac. In this condition the contents of the embryo sac constitutes the *female gametophyte*. [See Fig. 358 (1–8).]

MATURATION OF THE POLLEN GRAIN AND FORMATION OF THE MALE GAMETOPHYTE

The pollen grains (microspores), within the anther sacs, all arise from a number of *tetrads* (groups of four) which are formed by the division and redivision of *pollen mother-cells* preceding them. The pollen mother cells like the rest of the sporophyte generation contain the diploid or $2x$ number of chromosomes. During their division to form tetrads a reduction division occurs which reduces the chromosomes to the *haploid* or x number. This

number continues into the sperm nuclei. Each pollen grain, after the tetrads have separated into their components, consists of an outer firm wall or *exosporium*, an inner wall or *endosporium*, within which will be found the region called the *fovilla*, which is nothing other than a mass of protoplasm containing a nucleus. Before the pollen is shed from the anther, its protoplasmic contents undergo a series of changes leading up to the development of the male gametophyte. The nucleus and protoplasm enveloping it divides to form two cells, one a generative-cell

FIG. 357.—A, cross section of dehiscid anther of a lily (*Butomus*). The layer of cells beneath the epidermis (exothecium) acts as mechanical tissue to open the pollen sacs. (After Sachs.) B, Section of a pollen grain, diagrammatic. (Mottier.)

containing a generative nucleus, the other a tube cell containing a tube nucleus. The generative nucleus then divides to form two sperm nuclei and the partition wall between the two cells disappears. In this condition the protoplasmic contents of the pollen grain constitute the *male gametophyte*. (See Fig. 220.)

POLLINATION AND FERTILIZATION

The mature pollen grains are discharged from the ripened anther through the splitting open of its wall. They are transferred to the stigma of the pistil of another *Erythronium* flower through the agency of insects. Here they germinate, each put-

ting forth a tube (pollen tube). The pollen tubes, each carrying within it two sperm nuclei and a tube nucleus embedded in protoplasm, penetrate through the style canal until they reach the micropyles of various ovules. Each enters and passes through a micropyle, then piercing the nucellus, grows toward the embryo sac. The tip of the tube fuses with the end of the embryo sac and the two sperm nuclei are discharged into the sac. One

FIG. 358.—At the left, diagram of the anatomy of an angiospermous flower shortly after pollination; *anth.*, anther; *fil.*, filament; *st.*, stamen; *stig.*, stigma; *p.g.*, pollen grains germinating; *sty.*, style; *pt.*, pollen tube; *o.w.*, ovary wall; *o.* ovule, containing embryo-sac; *pet.*, petal; *sep.*, sepal. 1-8, Stages in the development of the female gametophyte (embryo-sac); *meg.sp.*, megaspore-mother-cell; *i.i.*, inner integument; *o.i.*, outer integument; *fun.*, funiculus; *chal.*, chalaza; *nu.*, nucellus (megasporangium); *emb.*, embryo-sac. All diagrammatic. (Gager.)

of these sperm nuclei (containing x chromosomes) passes between the synergids and fuses with the nucleus of the egg (containing x chromosomes) to form an *oöspore* or *zygote* in which the $2x$ number of chromosomes found in the sporophyte of the species is restored. By this time the tube nucleus has disintegrated. The *oöspore* by repeated divisions develops into as many as four *embryos* or young sporophyte plants. Only one of these, however, persists. The polar nuclei fuse to form the *endosperm nucleus* which soon

undergoes rapid division into a large number of nuclei scattered about through the protoplasm of the embryo sac. Later cell walls are laid down and *endosperm* is formed. The endosperm cells soon become filled with abundant starch which is later to be utilized by the embryo during germination.

RIPENING OF THE OVULE TO FORM THE SEED AND OF THE OVARY TO FORM THE FRUIT

When the embryo and endosperm are being formed, the ovule enlarges and its integuments become modified to form a hard, horny, seed coat which encloses the endosperm surrounding the embryo. The ovary, containing the ovules, has by this time ripened to form a three-valved, loculicidal capsule enclosing the seeds.

GERMINATION OF THE SEED AND DEVELOPMENT OF THE MATURE SPOROPHYTE

The seeds are fully developed by June or July when the capsule or fruit splits open to discharge them. They fall to the ground and lie dormant until the following spring when they germinate or commence to grow. Each seed absorbs water from the ground which stimulates the ferment amylase, contained in the endosperm cells, to break up the insoluble starch into soluble sugar which passes into solution and diffuses into the cells of the embryo, where the protoplasm changes it into additional protoplasm and so the embryo increases in size, therefore, *grows*. The pressure of the swollen endosperm and growing embryo becomes so great that the seed coat bursts; the hypocotyl emerges first, dragging the cylindrical cotyledon out of the seed coat and epicotyl with it. The hypocotyl elongates and extends itself into the soil where it develops a root near its tip. The tip enlarges through the storage of starch, manufactured by the green cotyledon and becomes a bulb. The bulb soon develops within it a plumule, the cotyledon withers, and the young plant (seedling) passes the following winter in this condition.

During the next spring the plumule develops into a foliage leaf and the bulb gives rise from its base to several slender elongated *runners*, which, at their tips, develop *runner bulbs*.

These runner bulbs, the third year, give origin to another set of runners similar to those formed during the second year which also develop runner bulbs at their tips. A foliage leaf is also formed by each. The following spring (spring of fifth year after formation and shedding of seed) one of these bulbs develops into a mature sporophyte plant, bearing a single flower at the summit of its elongated scape. (See Fig. 355.)

RESEMBLANCES BETWEEN GYMNOSPERMS AND ANGIOSPERMS

1. In both are developed flowers.
2. In both the flowers develop at least two sets of leaves (either on one or two plants of the same species) called sporophyll leaves, the stamens and carpels. The stamens or staminal leaves are also termed microsporophylls. The carpels or carpellate leaves are also known as megasporophylls.
3. Both groups produce microspores or pollen grains and megaspores or embryo sacs.
4. In both are developed on the evident generation, the plant or sporophyte and the gametophyte, the latter concealed within and parasitic upon the megaspore of the sporophyte.
5. Both develop seeds with one or two seed coats.
6. In both groups there is developed from the fertilized egg an embryo which lies within the cavity of the megaspore.
7. In both there exists a root and a stem pericycle.
8. Both produce collateral vascular bundles. Occasionally we meet with phlocentric bundles in the stem or leaf of Angiosperms.

FUNDAMENTAL DIFFERENCES BETWEEN GYMNOSPERMS AND ANGIOSPERMS

1. The flowers of Gymnosperms are often monœcious or dioœcious but very rarely hermaphrodite (as in *Welwitschia*), whereas those of Angiosperms are usually hermaphrodite, rather rarely monœcious, still more rarely dioœcious.
2. In the Gymnosperms the sporophylls are usually inserted either spirally or in whorls around a distinctly elongated axis, whereas in Angiosperms the sporophylls are condensed to short

whorls or spirals set around a shortened axis, the floral axis or receptacle, torus or thalamus, or, as in the more modified Angiosperms, the floral axis may even become hollow.

3. In Gymnosperms the microsporophylls or stamens are usually sessile, whereas in Angiosperms the microsporophylls are nearly always stalked. Rarely do we find sessile anthers among Angiosperms, an instance of this being seen in Mistletoe (*Viscum*) where the anthers are set on the staminal leaf.

4. In Gymnosperms there is a traceable prothallus or gametophyte plant that later becomes the so-called "endosperm" of the gymnosperm, whereas in Angiosperms no recognizable prothallus has been proven to exist.

5. The stored food tissue in Gymnosperm seeds is prothallial tissue loaded with starch, etc., whereas in Angiosperm seeds the stored food tissue (endosperm) is a special formation after fertilization.

6. Gymnosperms bear naked ovules and seeds while Angiosperms bear covered ones.

7. In Gymnosperms there are distinct recognizable archegonia formed on or imbedded in the prothallus, whereas in Angiosperms there are no distinct archegonia, only an isolated egg or eggs.

8. In Gymnosperms there are not infrequently found several embryos from one fertilized egg. This condition is called *polyembryony*. Polyembryony is unknown in Angiosperms, only a *false polyembryony* being noticed.

9. In Gymnosperms the secondary xylem (wood) tissue of roots, stems and leaves consists either of punctated or scalariform cells, whereas in Angiosperms the secondary wood tissue may be varied in structural aspect.

CLASSES OF ANGIOSPERMS

The Angiosperms are subdivided into two classes, the Monocotyledoneæ or Monocotyledons, and the Dicotyledoneæ or Dicotyledons.

The Monocotyledons include such plants as the cat-tails, grasses, sedges, palms, lilies, daffodils, irises, bananas, gingers and orchids. The Dicotyledons are represented by a vast

array of plants which include nearly all the familiar trees and shrubs except the conifers and the majority of weeds, garden plants, and ornamental herbs which have netted veined leaves.

The monocotyledons appear from studies in comparative morphology and serum-diagnosis tests to have originated from the Archichlamydeæ of the Dicotyledons.

CHAPTER XXII

THE MONOCOTYLEDONS

This is a class of Angiosperms characterized by the following peculiarities:

FIG. 359.—Morphology of the typical monocotyledonous plant. *A*, leaf, parallel-veined; *B*, portion of stem, showing irregular distribution of vascular bundles; *C*, ground plan of flower (the parts in 3's); *D*, top view of flower; *E*, seed showing monocotyledonous embryo. (*Gager.*)

(1) One cotyledon or seed leaf in the embryo, terminal in position.

(2) Stems usually endogenous with closed collateral or concentric fibro-vascular bundles, which are scattered. ,

(3) Leaves alternate in arrangement, with an entire margin and generally parallel-veined.

FIG. 360.—Wheat plant showing the general habit of grasses. (*Robbins.*)

(4) Flowers usually pentacyclic and trimerous (having the parts of each whorl in threes or a multiple thereof).

(5) Secondary growth in roots generally absent, the polyarch radial bundle persisting from its formation.

- (6) Medullary-rays generally absent.
- (7) Cambium generally absent.

FIG. 361.—Pistillate and staminate inflorescences of corn (*Zea mays*). Pistillate on left, staminate on right. (Robbins.)

DRUG YIELDING GROUPS OF MONOCOTYLEDONS

The Monocotyledons comprise 45 families which contain about 25,000 species of described plants. Only those groups which yield drugs are considered in this chapter.

ORDER GRAMINALES

This is the most important economic order of Monocotyledons. It contains two families, the *Gramineæ* or family of grasses and cereals and the *Cyperaceæ* or family of sedges. The

flowers of both of these families are devoid of sepals and petals but possess scale-like structures called glumes and palets which enclose one or more flowers. The grasses and cereals have stems with solid nodes and hollow internodes whereas the sedges have stems which are solid throughout. The above ground stem of a sedge is usually 3-sided, unlike any grass stem. The fruit of a grass is a caryopsis whereas that of a sedge is an akene.

GRAMINEÆ OR GRASS FAMILY.—Mostly herbaceous (*Triticum*, *Sorghum*, *Saccharum*, cereals, etc.), rarely shrubby (*Arundinaria*, etc.), more rarely arborescent (Bamboo, etc.) plants, varying in height from 2 in. to 100 ft.

FIG. 362.—Diagram of a typical spikelet. *nY* lower glume; *φY* upper glume; *nI* outer palet; *φI*, inner palet; *l*, lodicule; *st*, stamens; *I-I*, rachilla; 2, lateral axes. (Somewhat modified after Warming.)

Stems, leaves and even parts of flowers often abounding in silica. The leaves are alternate, with long split sheaths usually open on the side opposite the lamina, and a ligule. Flowers small, generally hermaphroditic, occasionally monœcious (Corn and Rice Grass) and usually borne in *spikelets* arranged along a spike-axis called a rachis, but sometimes in panicles (Oat). The spikelet is the unit of a grass inflorescence. A typical spikelet consists of a shortened zigzag axis (*rachilla*) bearing floral leaves. The two lowest floral leaves in each spikelet are barren and termed *glumes*. The succeeding floral leaves are of bract-like character, each supporting one flower in its axil, and are called *outer palets* or *lemmas*. Each flower has a bracteole which is placed on the inside, opposite and with its back to the rachilla and known as the *inner palet*. Immediately following the inner palet are 2 delicate scales called *lodicules*, 3 *stamens* (rarely 1, 2 or 6) with versatile anthers that are usually notched at each end and a single *pistil*, having a unilocular, superior ovary with one ovule and usually 2 styles, with simple or spirally branched (plumose) stigmas. Fruit, a caryopsis or grain. Seeds albuminous with starchy endosperm and a small embryo at the base and opposite the hilum side. Seed coat fused with fruit coat to

form one layer. Embryo with a *scutellum*. The scutellum represents a portion of the single cotyledon of these plants, which bales out the nourishment from the endosperm to the rest of the embryo, during the germination of the grain.

OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Triticum	Rhizome and roots	Agropyron repens	Europe and Asia
Sucrosum	Refined sugar	Saccharum officinarum and other sources	Tropics
Glucosum	Product obtained by incomplete hydrolysis of starch		
Dextrosum	Sugar obtained by hydrolysis of starch		
Amylum	Starch	Zea Mays	Mexico
Zea	Styles and stigmas	Zea Mays	Mexico
Avena	Grain	Avena sativa	Eurasia
Oleum Maydis	Fixed oil	Zea Mays	Mexico
UNOFFICIAL DRUG			
Oleum Citronellæ	Volatile oil	{ Cymbopogon Nardus Cymbopogon Winterianus	{ Asia
Job's Tears	Fruits	Coix Lacryma-Jobi	E. Indies
Fructus Canariense	Fruits	Phalaris canariensis	Canary Islands
Diastasum	Enzyme	Hordeum spp.	Asia
Vetiver	Rhizome and roots	Andropogon squarrosus	E. Indies
Maltum	Grain partially germinated and dried	Var. of Hordeum vulgare	Asia

ORDER PRINCIPES

PALMÆ OR PALM FAMILY.—Tropical or subtropical shrubs, rarely trees, having unbranched trunks which are terminated by a crown of leaves, in the axils of which the flowers are produced. The leaves are well developed with pinnate or palmate blades and fibrous, sheathed, clasping petiole. The flowers are small, of one or two sexes, and crowded on a spike or spadix, which is subtended by a large bract, or spathe which may become woody, as in the Cocoanut Palm. The perianth consists of 6 parts in 2 whorls (3 sepals and 3 petals) or it may be inconspicuous or absent. The stamens are 6 in number, rarely 3, inserted below the ovary. The ovary is superior, of 3 cells, with central placenta. The fruit is either a nut, with leathery epicarp, fibrous or cellular mesocarp and thin membranous

endocarp, or a drupe (Cocoanut) with leathery epicarp, broadly fibrous mesocarp and stony endocarp, or a berry as in the Date

FIG. 363.—Cocoanut Palm (*Cocos nucifera*). Note pinnate leaf-blades. (From Bull. Agricole du Congo Belge. 1910.)

Palm, *Phoenix*, with membranous epicarp, succulent mesocarp and soft succulent endocarp. The seeds are albuminous with

the reserve food frequently in the form of hard (reserve) cellulose (ivory-nut-palm) or ruminant albumen (*Areca*).

OFFICIAL DRUG	PART USED	BOTANICAL NAME	HABITAT
Sabal (Saw Palmetto)	Fruit (drupe)	Serenoa serrulata	South Carolina to Florida and Texas
Areca Arecolinæ Hydrobromi- dum	Seed Hydrobromide of alkaloid from seed	{ Areca Catechu	{ Asia and Malayan Islands.
UNOFFICIAL DRUG			
Dragon's Blood	Inspissated juice	Dæmonorops Draco	East Indies
Cocoanut oil	Fixed oil	Cocos nucifera	Tropics
Carnauba wax	Wax from leaves	Copernicia cerifera	Brazil
Palm oil	Fixed oil	Elæis guineensis	West Africa

ORDER ARALES

ARACEÆ OR ARUM FAMILY.—Perennial herbs with fleshy rhizomes or corms, long petioled leaves, and acrid or pungent juice. Leaves mostly radical; when cauline, alternate, entire to divided, often hastate or sagittate, with a membranous sheath at the base. Flowers small, usually imperfect, rarely hermaphroditic, usually monoecious, the pistillate usually below, the staminate above, crowded on a spadix, which springs from the axil of a spathe. Fruit an indehiscent, 1 to several celled, 1 to many seeded berry occurring in a cluster on the fleshy fruiting spadix. Seeds filled with large, curved, fleshy embryo or mostly with copious endosperm with embryo in center.

This family includes many ornamental plants among which are the calla lily (*Richardia æthiopica*), the ceriman (*Monstera deliciosa*) whose large leaves have many holes, the Anthuriums with colored showy bracts and the Elephant's ear (*Colocasia esculenta*).

OFFICIAL DRUG	PART USED	BOTANICAL NAME	HABITAT
Calamus	Peeled rhizome	<i>Acorus Calamus</i>	Europe, Asia, North America
UNOFFICIAL DRUG			
Skunk cabbage	Rhizome	<i>Symplocarpus fœtidus</i>	North America
Indian turnip	Corm	<i>Arisæma triphyllum</i>	North America
Tonga	Rhizome	<i>Epipremnum pin- natum</i> (in part)	Fiji Islands

FIG. 364.—*Acorus Calamus*, the Sweet flag, a member of the *Araceæ*. (Sayre.)

FIG. 365.—Diagram of *A*, lily flower, and *B*, grass flower, showing homologous structures. *A*, *f*, bract; *ax*, axis; *op*, outer perianth; *ip*, inner perianth; *s*, stamens; (*c*) tricarpellary ovary. *B*, shaded structures are aborted; *l*, lemma (bract); *ax*, axis; *p* and *p'*, palea (outer perianth); *l* and *l'*, lodicules (inner perianth); *s* and *s'*, two whorls of stamens; *c*, tricarpellary ovary. (*A*, Robbins. *B*, after Shuster.)

ORDER LILIALES

LILIACEÆ OR LILY FAMILY.—Herbs (*Lilium*), shrubs (*Yucca*), or trees (*Dracena Draco* or Dragon's Blood), with regular and

FIG. 366.—*Smilax officinalis*—Portion of vine and root system. Note the 3-nerved leaves with stipules modified as tendrils and the berry fruits. (Sayre.)

symmetrical almost always six-anded flowers. Stem either short, creeping under ground (*Convallaria*), or, swelling up and forming bulbs (*Hyacinth*), or corms (*Colchicum*), or, stem may

elongate above ground and become wiry and herbaceous or semi-shrubby as *Smilax*, or the stem may remain short, giving rise to thick, fleshy and sap-storing leaves, as in *Aloe*. Leaves linear to lanceolate, ovate rarely wider, divisible into sheathing base, narrow petiole and expanded blade. Venation, parallel, becoming in some ovate leaves parallel with oblique connections, reticulate or highly reticulate, as in the Sarsaparilla plants and other species of *Smilax*. The perianth is parted into six segments, the calyx and corolla being alike in color. Anthers introrse. Ovary three-locular with a single style. Fruit a three-locular, loculicidally dehiscent capsule (*Lilium*, *Colchicum*, etc.) or rarely a berry (*Asparagus*, Lily-of-the-valley, etc.). Seeds usually numerous with fleshy endosperm.

OFFICIAL DRUG	PART USED	BOTANICAL NAME	HABITAT
Sarsaparilla	Root	<i>Smilax medica</i>	Mexico
		<i>Smilax ornata</i>	Costa Rica
		<i>Smilax officinalis</i> or	Guatemala,
		an undetermined species	Honduras and Nicaragua
Veratrum Viride	Rhizome and roots	<i>Veratrum viride</i>	United States
Colchici Cormus	Corm	<i>Colchicum autumnale</i>	Mediterranean Basin
Colchici Semen	Seed	<i>Colchicum autumnale</i>	Mediterranean Basin
Aloe	Inspissated juice of leaves	<i>Aloe vera</i>	Dutch West Indies
		<i>Aloe Perryi</i>	Socotra and East Africa
Aloin	Pentoside		
Scilla	Inner scales of bulb	<i>Urginea maritima</i> (<i>Scilla maritima</i>)	Mediterranean Basin
Convallariæ Radix	Rhizome and roots	<i>Convallaria majalis</i>	Europe, Asia, United States
Trillium	Rhizome and roots	<i>Trillium erectum</i>	North America
Aletris	Rhizome and roots	<i>Aletris farinosa</i>	Eastern United States
Helonias	Rhizome and roots	<i>Chamælorium luteum</i>	Eastern United States
UNOFFICIAL DRUG			
Cape Aloe	Inspissated juice	<i>Aloe ferox</i>	S. Africa
Sabadilla	Seed	<i>Asagraea officinalis</i>	Mexico and Central America
Red Squill	Bulb	<i>Urginea maritima</i> (red variety)	Mediterranean Basin

UNOFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Allium	Bulb (fresh)	Allium sativum	Southern Europe
Convallariæ Flores	Inflorescence	Convallaria majalis	Europe, Asia, United States
Veratrina	Mixture of alkaloids	Asagrea officinalis	Mexico and Central America

DIOSCOREACEÆ OR YAM FAMILY.—Twining perennial herbs or undershrubs arising from large tuberous roots or knotted rootstocks. Leaves alternate, petioled, simple, palmately-ribbed and netted-veined. Flowers small, dioecious, regular, having a six-cleft, calyx-like perianth, six stamens and a three-celled inferior ovary. Fruit usually a membranous, three-angled or winged, three-celled capsule.

OFFICIAL DRUG	PART USED
Dioscorea	Rhizome
BOTANICAL NAME	HABITAT
Dioscorea villosa	United States

IRIDACEÆ OR IRIS FAMILY.—Perennial herbs with underground rhizomes (Iris) corms (Crocus) or bulbs (Gladiolus) sword shaped or linear, equitant, two-ranked leaves and usually showy, perfect, regular or irregular flowers which emerge from a spathe of 2 or more bracts. Perianth of 3 sepals (often petaloid), and 3 petals arising from the summit of a tubular receptacle above the ovary; stamens 3; ovary inferior, 3-celled, style single, stigmas 3, sometimes expanded and colored, as in *Iris*. Fruit a three-celled, loculicidal, many-seeded capsule. Seeds with fleshy or horny endosperm. Rootstocks, tubers, or corms mostly acrid. Fibrovascular bundles in stems and leaves phlocentric.

FIG. 367.—*Iris versicolor* (Blue Flag). Note the fleshy rhizome bearing equitant, sword-shaped leaves and long stalked flower clusters. The flowers are violet-blue, the outer segments of the perianth limb being spatulate, variegated with yellow, green or white on the claw and marked with purple veins. (Reproduced from U. S. Dept. Agric. Miscellaneous Publ. 77.)

OFFICIAL DRUG	PART USED	BOTANICAL NAME	HABITAT
Iris (Orris)	Peeled rhizome	{ Iris florentina Iris pallida Iris germanica	{ Mediterranean Region
Iris Versicolor (Blue Flag)	Rhizome	{ Iris versicolor Iris caroliniana	Eastern North America
Crocus (Saffron)	Stigma	Crocus sativus	Southern Europe and Asia

ORDER SCITAMINALES

This order includes herbs with asymmetrical flowers and pinnately-veined leaves. Some of the members, notably the bananas, are gigantic in size. It contains the *Musaceæ* or Banana Family, *Cannaceæ* or Canna Family, *Zingiberaceæ* or Ginger Family and *Marantaceæ* or Arrowroot Family.

ZINGIBERACEÆ OR GINGER FAMILY.—Tropical plants, perennial herbs, usually with fleshy rhizomes and large, elliptical, pinnately-veined leaves. The leaf sheaths are folded tightly around each other so as to give the appearance of a stem. Flowers, very irregular, trimerous; sepals three, short, often green; petals three, elongate, often fused below; stamens three to four abortive, petaloid, one often absent, a sixth alone fertile and stamen-like. The petaloid stamens constitute the most attractive parts of the flower. Pistil, inferior, tricarpellary. Fruit, a loculicidally dehiscent, usually 3-celled capsule. Seeds albuminous with both perispermic and endospermic albumen.

OFFICIAL DRUG	PART USED	BOTANICAL NAME	HABITAT
Zingiber	Rhizome	Zingiber officinale	S. Asia
Cardamomi Semen	Seeds	Elettaria Cardamomum	Indo-China
Oleum Cardamomi	Volatile oil from seeds	Elettaria Cardamomum	Indo-China
UNOFFICIAL DRUG			
Galanga	Rhizome	Alpinia officinarum	China
Zedoaria	Rhizome	Curcuma Zedoaria	India
Curcuma	Prepared rhizome	Curcuma longa	S. Asia and E. Indies
Grains of Paradise	Seeds	Amomum melegueta	Asia

MARANTACEÆ OR ARROWROOT FAMILY.—Perennial herbs chiefly of tropical climes with fleshy rhizomes or tubers rich in

starch content. The leaves are radical and cauline, usually long petioled, with a sheathing base and a pulvinus at the summit of the petiole. The inflorescence is a panicle or raceme

FIG. 368.—*Zingiber officinale*, the Ginger plant. (Sayre.)

bearing irregular trimerous flowers with a one-celled ovary (by abortion).

UNOFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Amylum	Starch	Maranta	{ West Indies Northern S. America
Marantæ		arundinacea	

ORDER ORCHIDALES

ORCHIDACEÆ OR ORCHID FAMILY.—Perennial herbs of terrestrial or terrestrial saprophytic or epiphytic growth, having grotesque flowers. Roots fibrous or tuberous, often saprophytic

FIG. 369.—Floral organs of an orchid (*Catleya* sp.). A, the entire flower; *sep*, sepal; *pet*, petal; B, column, showing *s*, stigma and *r*, the rostellum (beak), with the small glands at the tip; to the glands are attached the four strap-shaped caudicles of the pollinia; C, pollinia, with the four caudicles; below, the gland; D, longitudinal section of the column; *p*, pollinium; E, the same, enlarged. (Gager.)

in relation, or aerial and with velamen. Stems and branches upright, in epiphytic types, often forming pseudobulbs. Leaves alternate, entire, parallel-veined, sheathing at base, rarely reduced to yellowish or pale scales in saprophytes. Flowers irregular, usually attractive, entomophilous, but sometimes

small, greenish, and inconspicuous, arranged in elongated spikes or racemes, trimerous. Sepals 3, usually similar; petals 3 of which two often resemble sepals, third is variously, often greatly modified and fused with two outer petaloid stamens as a labellum or lighting-board for insects. Third stamen of outer whorl

FIG. 370.—*Vanilla planifolia*—Branch showing leaves and flowers. (Sayre.)

fertile (*Orchideæ*) or a barren knob (*Cypripediæ*); pollen of fertile anther agglutinate and forming a *pollinium*. Three stamens of inner circle barren and petaloid or one absent (*Cypripediæ*). Stamens all epigynous and often three are fused with the style as a gynandrium. Carpels three, syncarpous, with inferior, 3-, rarely 4-, 1 (usually)-celled ovary. Fruit a capsule, 3-valved and 1-celled. Seeds minute, abundant and wind disseminated.

OFFICIAL DRUG	PART USED	BOTANICAL NAME	HABITAT
Vanilla	Fruit (cured, full-grown, unripe)	Vanilla planifolia	Mexico
UNOFFICIAL DRUG			
Salep	Tubers	{ Orchis mascula Orchis morio	{ Europe
Corallorhiza	Rhizome	Corallorhiza odontorhiza	Eastern U. S.
Cypripedium	Rhizome and roots	{ Cypripedium bulbosum Cypripedium parviflorum Cypripedium pubescens	{ United States

CHAPTER XXIII

THE DICOTYLEDONS

The Dicotyledons are a class of angiosperms having the following characteristics:

(1) Two-seed leaves (cotyledons) in embryo, these being lateral on the embryonic axis.

FIG. 371.—Morphology of a typical dicotyledonous plant. *A*, leaf, pinnately-netted veined; *B*, portion of stem, showing concentric layers of wood; *C*, ground-plan of flower (the parts in 5's); *D*, perspective of flower; *E*, longitudinal section of seed, showing dicotyledonous embryo. (*Gager.*)

(2) Foliage leaves with pinnate-reticulate or palmate-reticulate veins.

(3) Leaf blades varying from simple to lobed to compound in pinnate or palmate fashion depending upon character of venation.

(4) Stems, leaves, and roots of secondary growth with open collateral or (excepting roots) bi-collateral fibro-vascular bundles; in stems radially arranged about a central pith.

(5) Exogenous stems.

(6) Medullary-rays present in stems and in roots of secondary growth.

(7) Cambium present in the plant axis.

FIG. 372.—At left, fructing branch of Black Pepper (*Piper nigrum*). Portion of shoot with inflorescences. Terminal portion of an inflorescence (enlarged). (After Wossildo.)

At right, the Cubeb (*Piper Cubeba*). *a*, cluster of fruit; *b*, staminate flower; *c*, pistillate flower in longitudinal section; *d*, fruit in longitudinal section. (After Berg and Schmidt.) Both from Gager after Strasburger, Noll, Schenck and Schimper.

(8) Roots developing secondary structure.

(9) Flowers usually pentacyclic and tetramerous or pentamerous (parts of each whorl, four or five or multiple thereof), with usually a reduced number of parts in the gynœcium.

This is the largest class of angiosperms containing more than 100,000 species of described plants which have been grouped into 240 families. We shall consider in this chapter only those groups of dicotyledons which yield drugs.

The Dicotyledons are subdivided into two sub-classes; viz.:

Sub-class 1. Archichlamydeæ.

Sub-class 2. Sympetalæ or Metachlamydeæ.

DRUG YIELDING DICOTYLEDONS

SUB-CLASS ARCHICHLAMYDEÆ

Those dicotyledonous plants in which the petals are distinct and separate from one another or are entirely wanting. That group of the Archichlamydeæ whose flowers show the absence of petals and frequently of sepals is called the *Apetalæ*. The group whose plants have flowers showing the parts of their corolla (petals) separate and distinct is called the *Choripetalæ* or *Polypetalæ*.

ORDER PIPERALES

PIPERACEÆ OR PEPPER FAMILY.—A family of apetalous aromatic herbs and shrubs with jointed stems, opposite, verticillate, or sometimes alternate leaves without stipules, and spiked, inconspicuous, wind-pollinated flowers. The characteristic fruit is a drupe enclosing a single, upright, albuminous seed with abundant perisperm (from megasporangial tissue) and reduced endosperm (from matured embryo sac).

OFFICIAL DRUG	PART USED	BOTANICAL NAME	HABITAT
Cubeba	Unripe fruit	Piper Cubeba	Borneo, Java Sumatra
UNOFFICIAL DRUG			
Piper (Pepper)	Unripe fruit	Piper nigrum	Cochin-China, India
Piper Longum	Spike of immature fruit	{ Piper longum Piper officinarum	{ Asia
Kava	Rhizome and roots	Piper methysticum	Sandwich Islands
Matico	Leaves	Piper angustifolium	Peru, Bolivia

ORDER SALICALES

SALICACEÆ OR WILLOW FAMILY.—Shrubs or trees of temperate or cold regions, with upright woody stems, rarely herbs (*Salix retusa*). Bark often containing bitter principles (Salicin, etc.).

Leaves alternate, simple, entire, stipulate; stipules rarely green, persistent, usually functioning as winter bud-scales and falling in spring.

Inflorescences dioecious spikes, so on separate plants. Staminate spikes forming deciduous catkins of yellowish flowers, pistillate as persistent spikes of green flowers, at length maturing fruit.

Flowers of catkins numerous, each of two to five (Willow) or six to fifteen (Poplar); stamens in axil of a small bract leaf, sometimes with small nectar knob or girdle at base; pollen abundant, hence plants anemophilous, rarely entomophilous. Pistillate flowers green, each of a bicarpellate pistil in axil of bract, ovary one-celled with parietal placentation, style simple, stigma bilobed.

Fruit a capsule dehiscing longitudinally. Seeds small, exalbuminous, surrounded by a tuft of hairs for dissemination.

OFFICIAL DRUG	PART USED	BOTANICAL NAME	HABITAT
Salicinum	Glucoside	Several species of Salix and Populus	Europe, North America
Populi Gemma	Closed winter leaf bud	<div> <div> Populus balsamifera Populus candicans </div> </div>	<div> </div> North America
UNOFFICIAL DRUG			
Salix	Bark	Salix alba	Europe
Salix Nigra	Bark	Salix nigra	North America
Poplar bark	Bark	Populus tremuloides	North America

ORDER MYRICALES

MYRICACEÆ OR BAYBERRY FAMILY.—Dioecious or sometimes monœcious, aromatic shrubs or trees with watery juice and possessing underground branches which arch downward then upward producing many suckers. Roots fibrous and bearing many short rootlets upon which are frequently found coralloid clusters of tubercles containing the *Actinomyces Myricarum* Youngken. Leaves alternate, revolute in vernation, serrate, irregularly dentate, lobed or entire, rarely pinnatifid, pinnately and reticulately veined, pellucid-punctate, evergreen or deciduous, generally exstipulate, rarely stipulate. Flowers naked, unisexual, monœcious or dioecious, in the axils of unisexual or

androgynous aments from scaly buds formed in the summer in the axils of the leaves of the year, remaining covered during the winter and opening in March or April before or with the unfolding of the leaves of the year.

Staminate flowers in elongated catkins, each consisting of two to eight stamens inserted on the torus-like base of the oval or oval-lanceolate bracts of the catkin, usually subtended by two or four or rarely by numerous bracteoles; filaments short or elongated, filiform, free or connate at the base into a short stipe; anthers ovoid, erect, two-celled, extrorse, showing longitudinal dehiscence. Pistillate flowers in ovoid or ovoid-globular catkins. Gynœcium of two united carpels on a bract. Ovary sessile, unicellular, subtended by two lateral bracteoles which persist under the fruit, or by eight linear-subulate bracteoles, accrescent, and forming a laciniate involucre inclosing the fruit; styles short and dividing into two elongated style arms which bear stigmatic surfaces on their inner face; ovule orthotropous, solitary, with a basilar placenta and superior micropyle. Fruit an akene or ceriferous nut. Pericarp covered with glandular emergences which secrete wax or fleshy emergences, smooth and lustrous or smooth, glandular. Seed erect, exalbuminous, covered with a thin testa. Embryo straight, cotyledons thick, plano-convex; radicle short, superior.

There are two distinct genera of this family, *e.g.*, *Myrica* and *Comptonia*.

FIG. 373.—Flowers and fruit of willow (*Salix viminalis*). A, twig with staminate catkins. B, staminate flower with subtending bract, enlarged. C, pistillate catkin. D, E, pistillate flowers, enlarged. F, fruit, natural size. G, the same enlarged. H, seed enlarged. (From Strasburger, Macmillan Co.)

UNOFFICIAL DRUG	PART USED	BOTANICAL NAME	HABITAT
Myrica	Bark of root	{ Myrica cerifera Myrica carolinensis	{ Eastern North America
Comptonia	Leaf	Comptonia asplenifolia	Eastern North America
Bayberry Wax	Wax from fruits	Myrica cerifera Myrica carolinensis	{ E. North America

FIG. 374.—Fructiferous branches of *Myrica caroliniensis* (to left) and *Myrica cerifera* (to right) as they appear in early winter. The former species is deciduous while the latter is evergreen. Collected by author at Wildwood, N. J., in mid December.

ORDER JUGLANDALES

JUGLANDACEÆ OR WALNUT FAMILY.—Tall, branching, monœcious trees or shrubs with branches and leaf-stalks usually beset with glandular-resinous hairs. Leaves alternate, exstipu-

late and odd-pinnate which, upon falling, leave usually horse-shoe shaped scars on branches, and in whose axils develop frequently one axillary bud and several ascending accessory buds. Inflorescence of staminate catkins and pistillate spikes.

FIG. 375.—Fructiferous branch of *Myrica Gale*, showing mature pistillate catkins and nature of leaves. Collected at the south end of Peak's Island in Casco Bay, Maine, September, 1913.

The staminate catkins occur on the old wood of the previous year. The pistillate spikes are borne at the extremity or on the sides of the new wood. The staminate flower is composed of an elongated floral disk bearing a terminal bract, a perianth of 6 to rarely 4 parts and 40 to 4 stamens. The pistillate flower is composed of an upgrown, hollowed-out receptacle bearing on

its margin or extremity 5 to 4 perianth segments and containing a pistil consisting of an inferior ovary with a single ovule and 2 styles having radiating penicillate stigmas. Fruit an aromatic

FIG. 376.—The sweet- or black birch, *Betula lenta*. Above, branch with staminate catkins. Below, branch with pistillate catkins. The tree occurs in rich woodlands from Newfoundland to Florida. (Reproduced from U. S. Dept. Agric. Miscell. Publ. 77.)

drupe composed of fused receptacular tube and ovarian wall and exhibiting a leathery to rarely membranous epicarp, more or less succulent mesocarp, and stony endocarp enclosing a seed with a double layered seed coat surrounding an enlarged embryo with irregular, semi-ruminate cotyledons, into and between which the fruit and seed layers grow.

The family embraces six genera, of which *Carya* (Hickories) and *Juglans* (Walnuts) are represented in the United States. The seeds of *Carya Pecan* (Pecan) and *C. ovata* (Shagbark Hickory) are valued as food. The wood of the walnuts is used for furniture and the seeds of *Juglans cinerea* (Butternut), *J. nigra* (Black Walnut) and *J. regia* (English Walnut) are prized as foods. Black walnut hulls are used as a coloring agent.

UNOFFICIAL DRUG	PART USED	BOTANICAL NAME	HABITAT
Juglans	Inner root bark	<i>Juglans cinerea</i>	United States
Black Walnut	{ Leaves, bark, hulls, kernels	{ <i>Juglans nigra</i>	United States

ORDER FAGALES

BETULACEÆ OR BIRCH FAMILY.—A family of aromatic trees or shrubs distinguished by monœcious flowers with scaly bracts and astringent resinous bark. Differs from *Fagaceæ* by superior ovary and absence of cupule. To this family belong the hazelnuts (*Corylus*), birches (*Betula*), alders (*Alnus*), the hop hornbeam, ironwood (*Ostrya*) and the American hornbeam (*Carpinus*).

The seeds of the hazelnuts or filberts are prized as food. The bark and oil of the sweet- or black-birch are employed as flavoring agents and antirheumatics.

OFFICIAL DRUG	PART USED	BOTANICAL NAME	HABITAT
(Oleum Betulæ) Methylis Salicylas	Volatile oil	<i>Betula lenta</i>	North America
Oleum Betulæ Empyreu- maticum Rectificatum	Pyroligneous oil	<i>Betula alba</i>	Eurasia
UNOFFICIAL DRUG			
Birch Bark	Bark of stem	<i>Betula lenta</i>	North America

FIG. 377.—*Quercus infectoria*—Branch and nutgall. (Sayre.)

FAGACEÆ OR BEECH FAMILY (CUPULIFERÆ).—*Beeches*, *Chestnuts*, *Oaks*, the trees of this family, are found in the temperate forests of the eastern and western hemispheres and comprise about 368 species. North America has over 50 species of oaks; 2 species of Chestnuts; 1 species of beech and 1 species of golden-leaved chestnut. The most important American oaks used for building, for furniture, for cooperage, for wagons, for tanning

FIG. 378.—*Ulmus fulva*, the Slippery Elm. Flowering branch bearing bell-shaped flowers, leaf and leaf and fruiting branch. The fruits are circular, single samaras. The inner bark of this tree is rich in mucilage. (Reproduced from U. S. Dept. Agric. Miscell. Publ. 77.)

leather etc. are white oak, *Quercus alba*; chestnut oak, *Q. prinus*; black oak, *Q. velutina*; live oak, *Q. virginiana*; swamp white oak, *Q. platanoides*; cow oak, *Q. Michauxii*, and the two Pacific coast oaks, *Q. chrysolepis* and post oak, *Q. garryana*. The uses of the fast disappearing American Chestnut, *Castanea dentata*, are well known. The wood of the beech, *Fagus grandifolia*, is used for chairs, tool handles, plane stocks, shoe lasts and for fuel. The nuts (mast) fatten hogs and feed wild animals and birds. The cork of commerce is obtained from the bark of *Quercus Suber* and *Quercus occidentalis*, plants indigenous to Spain and France.

The above trees are all monœcious, that is the staminate (male) and pistillate (female) flowers are distinct from each other, but borne on the same tree. Most of the species are trees, a few oaks are shrubs. The leaves are simple, netted-veined and alternate. A pair of deciduous stipules are found at the base of the leaf-stalk (petiole). The margins of the chestnut and beech leaves are sharply cut with large teeth. The leaves of the oaks divide the genus into two groups, viz.—the *white oaks* with rounded lobed leaves and annual acorn production, and the *black oaks* with sharp, bristle-tipped lobes and biennial acorn production. The male flowers are in dangling heads

(beech), or in catkins (chestnut and oaks). The male flowers have a united perianth, which is 4–6 parted and encloses an indefinite number of undivided stamens. The female flowers have a superior 6-parted perianth; the pistil consisting of 3 carpels with a corresponding number of stigmas. The ovary is 3–6 celled and each cell has 2 pendulous ovules. The fruit is a one-seeded nut. The cup, or cupule, in the beech is 4-sided and covered externally with weak spines and encloses two 3-sided seeds. The cupule in the chestnut forms the spiny bur, which splits into 4 valves at maturity, enclosing 3 nuts. The cupule in the oak is saucer-, or cup-shaped, and encloses a single rounded nut, or acorn. The seeds are exalbuminous and the cotyledons are thick and fleshy, edible in the beech, chestnut and a few of the oaks.

OFFICIAL DRUG	PART USED	BOTANICAL NAME	HABITAT
Galla	Excrescence	<i>Quercus infectoria</i>	Europe and W. Asia
Castanea	Leaf	<i>Castanea dentata</i>	North America
Acidum Tannicum	Tannin from gall	<i>Quercus infectoria</i>	Europe
Acidum Gallicum	Product made from tannin		
Pyrogallol	Product made from gallic acid or extract of galls		
UNOFFICIAL DRUG			
Quercus	Inner bark	<i>Quercus alba</i>	North America

ORDER URTICALES

ULMACEÆ OR ELM FAMILY.—Forest trees indigenous to the temperate and tropical zones, characterized by being woody plants, with simple, often inequilateral, pinnately-veined leaves having caducous stipules and without milky juice. Their flowers are unisexual or hermaphroditic with six or four parts to the calyx. Fruit a single winged samara (Elms) or a drupe (Hackberry).

OFFICIAL DRUG	PART USED	BOTANICAL NAME	HABITAT
Ulmus	Inner bark	<i>Ulmus fulva</i>	United States and Canada

MORACEÆ OR MULBERRY FAMILY.—Mostly apetalous shrubs or trees, as the mulberries (*Morus*), figs (*Ficus*), osage orange (*Maclura*), etc., rarely herbs as hop and hemp, many of them

containing a milky juice in latex tubes with small, axillary, clustered or solitary, unisexual flowers with sepaloid perianth, variously colored; leaves either simple, ovate with serrate margin

FIG. 379.—Hop (*Humulus lupulus*), a member of the *Moraceæ*. A, portion of plant showing pistillate inflorescences; B, staminate inflorescence; C, rachis of pistillate inflorescence ("hop"). (Robbins.)

or palmately lobed (fig) to palmately compound (hemp) and having caducous stipules; ovules mostly amphitropous; fruit an akene enclosed by the persistent perianth, as in hop and can-

FIG. 380.—*Cannabis sativa* or Indian Hemp (Fam. *Moraceæ*). At left, staminate plant, bearing axillary panicles of staminate flowers. At right, young pistillate plant whose panicles constitute the drug Cannabis. Note the palmately compound leaves, the leaflets of which are lanceolate with serrate margins.

nabis, a sorosis, as in the mulberry and osage orange or a syconium, as in the fig; embryo usually curved.

OFFICIAL DRUG	PART USED	BOTANICAL NAME	HABITAT
Cannabis	Flowering tops of pistillate plant	$\left\{ \right.$ Cannabis sativa	Asia

OFFICIAL DRUG	PART USED	BOTANICAL NAME	HABITAT
Humulus	Strobile	Humulus lupulus	} Europe, Asia } North America
Lupulinum	Glandular trichome	Humulus lupulus	
UNOFFICIAL DRUG			
Fructus Cannabis (Hemp seed)	Fruit	Cannabis sativa	Asia
Ficus	Fruit	Ficus Carica	Persia

ORDER SANTALALES

SANTALACEÆ OR SANDALWOOD FAMILY.—Herbs, shrubs or trees having entire exstipulate leaves, greenish flowers, nut-like or drupaceous fruits and oily seeds. Many are parasitic on the roots of other plants or on trees. The woods and volatile oils of a number of species are of economic importance.

OFFICIAL DRUG	PART USED	BOTANICAL NAME	HABITAT
Oleum Santali	Volatile oil	} Santalum album	India and Malaya
Santalum Album	Heartwood		
UNOFFICIAL DRUG			
Oleum Santali	Volatile oil	Eucarya spicata	W. Australia
Australiensis			

LORANTHACEÆ OR MISTLETOE FAMILY.—Most of the members of this family are woody shrubs, parasitic on trees; leaves opposite or whorled, entire, simple, exstipulate, sometimes reduced to scales; flowers hermaphroditic or dioecious, the calyx annular or cup-like, adnate to the ovary; petals disjoined or united into a tube, often split down one side, stamens equal in number to the petals and inserted on their base, the anthers dehiscent by pores or terminal slits: nectar disk present or absent; ovary inferior with simple style or sessile stigma; fruit a berry or drupe with a solitary seed devoid of testa and containing copious endosperm and a straight embryo.

UNOFFICIAL DRUG	PART USED	BOTANICAL NAME	HABITAT
Phoradendron (American Mistletoe)	Entire plant	Phoradendron flavescens	United States
Viscum (European Mistletoe)	Entire plant	Viscum album	Europe and Asia

ORDER ARISTOLOCHIALES

ARISTOLOCHIACEÆ OR BIRTHWORT FAMILY.—Herbs or twining semi-woody or woody plants, having more or less swollen

nodes from which spring alternate, cordate, reniform or ovate leaves. Flowers regular (*Asarum*, etc.) or irregular (*Aristolochia*), often offensively smelling. Sepals varying from 6 to rarely 5 to more commonly 4 or 3; in *Asarum*, etc. forming a regular symmetrical cup with 3 teeth or lobes, in *Aristolochia* divisible into a bowl, a tube and a limb. Stamens from 24 to 3, the fila-

FIG. 381.—*Aristolochia Serpentaria*. (Sayre.)

ments distinct or slightly fused with styles, in *Asarum*, or completely fused, in *Aristolochia*, to form a gynostemium, the anthers separate. Pistil of as many carpels as stamens, the stylar portions more or less distinct from stamens in *Asarum*, fused in *Aristolochia*, the ovary semi-inferior to inferior in *Asarum*, completely inferior in *Aristolochia* and as many celled as there are carpels. Fruit a capsule. Seeds with copious albumen and minute embryo.

OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Serpentaria	Rhizome and roots	<div style="display: inline-block; vertical-align: middle;"> <div style="display: inline-block; vertical-align: middle;">Aristolochia</div><div style="display: inline-block; vertical-align: middle;">Serpentaria</div><div style="display: inline-block; vertical-align: middle;">Aristolochia reticulata</div> </div>	United States
Asarum	Rhizome and roots	Asarum canadensis	United States

ORDER POLYGONALES

POLYGONACEÆ FAMILY.—Usually herbs [Knot-grasses (*Polygonum*), rhubarb (*Rheum*), docks (*Rumex*), buckwheat (*Fagopy-*

FIG. 382.—Serpentaria—Cross-section of rhizome. (25 diam.) A, parenchyma of cortex; B, medullary ray; C, xylem; D, phloem; E, medulla. (Sayre.)

rum), etc.] rarely trees (*Coccoloba uvifera* or Sea-Grape) or shrubs (*Muehlenbeckia* or Ribbon Bush) having strong vertical tap roots and spreading secondary roots more or less provided with tannin compounds. Stems elongate, green, to woody, rarely flattened, leathery, phylloidal (*Muehlenbeckia platyclada*), still more rarely tendriform (*Antigonon leptopus* or Mountain Rose of Tropical America). Leaves alternate, rarely opposite or whorled (*Eriogonum*), entire, rarely lobed (*Rheum palmatum*, *Rumex acetosella*), petiolate, rarely sessile, and stipulate. Stipules fused and forming a greenish membranous upgrowth (*ocrea*) which sheaths the stem. Inflorescence racemose with many dense scorpioid or helicoid cymes, which in some forms condense into single flowers. Flowers regular, pentamerous, with simple calyx, becoming trimerous with two whorls of three sepals each. Stamens varying from fifteen or twelve to nine or six, more

rarely to five, four, three to one (*Kænigia*), hypogynous, more rarely by enlargement of receptacle and slight fusion of sepals, perigynous. Pistil tri- to bi-carpellate, often three- to two-sided; ovary one-celled with one ovule. Styles three, rarely two, radiating penicillate in wind-pollinated, inconspicuous flowers, becoming condensed, knob-like in conspicuous insect-pollinated flowers. Fruit a triangular or biconvex akene often crowned by persistent styles and surrounded by persistent closely

FIG. 383.—Rhubarb (*Rheum*) flower, external view, median lengthwise section, and with perianth and stamens removed. (Robbins, after Luerksen.)

applied sepals. Seeds solitary, albuminous, with straight embryo, or in *Rumex*, curved embryo.

OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Rheum	Rhizome and roots, deprived of periderm tissues	<div> <div>Rheum officinale</div> <div>Rheum palmatum and other species (excepting <i>R. rhaponticum</i>) or hybrids of Rheum, grown in China and Thibet</div> </div>	China and Thibet
UNOFFICIAL DRUG			
Rumex	Root	<div> <div>Rumex crispus</div> <div>Rumex obtusifolius</div> </div>	Europe
Bistorta	Rhizome	Polygonum Bistorta	Europe and Asia

ORDER CHENOPODIALES OR CENTROSPERMÆ

This order contains seven families, all of which have seeds with a curved or coiled embryo. These families are the Goose-

foot Family (*Chenopodiaceæ*), the Pokeweed Family (*Phytolaccaceæ*); the Amaranth Family (*Amaranthaceæ*) which includes the ornamental Cock's Comb; the Four-o'clock Family (*Nyctaginaceæ*) to which belong the Marvel of Peru or Four-o'clock and the beautiful Bougainvilleas; the *Aizoaceæ* containing the Carpet Weed (*Mollugo verticillata*) and the Sea Purslane (*Sesuvium maritimum*), etc.; the Portulaca Family (*Portulacaceæ*) which is represented by the Spring Beauty (*Claytonia virginica*) and the Purslane (*Portulaca*); the Pink Family (*Caryophyllaceæ*) which includes the Carnations, Corn Cockle, Bouncing Bet, etc.

CHENOPODIACEÆ OR GOOSEFOOT FAMILY.—Usually herbaceous halophytes or shore growers, growing also in any alkaline soil, more rarely shrubs (*Atriplex*) or low trees (*Haloxylon*).

Among them are several garden vegetables (*Spinach*, *Beets*, *Mangels*) and a number of weeds. Leaves alternate to opposite sometimes reduced to teeth, entire or lobed. Inflorescence spikes or short racemes of condensed cymes. Flowers regular, usually small and greenish. They are either perfect (*Beta*), monœcious (*Chenopodium*), diœcious (*Atriplex* sp.), or polygamous (*Kochia* or *Summer Cypress*). Fruit a utricle. Seed albuminous. Embryo curved, enclosing central albumen.

OFFICIAL DRUG	PART USED	BOTANICAL NAME	HABITAT
Oleum Chenopodii	Volatile oil	<i>Chenopodium ambrosioides</i> var. <i>anthelminticum</i>	United States
Sucrosum	Refined sugar	{ <i>Beta vulgaris</i> <i>Beta maritima</i>	Europe
UNOFFICIAL DRUG			
<i>Chenopodium</i>	Fruit	<i>Chenopodium ambrosioides</i> var. <i>anthelminticum</i>	United States

PHYTOLACCACEÆ OR POKE FAMILY.—A family of apetalous trees, shrubs, and herbs with alternate, entire leaves, and flowers resembling those of the goosefoot family (*Chenopodiaceæ*), but differing in having the several-celled ovary composed of carpels united in a ring, and forming a berry in fruit. It embraces 21 genera, and 55 species, mostly tropical and sub-tropical. Only the genus *Phytolacca* occurs in the United States.

OFFICIAL DRUG	PART USED	BOTANICAL NAME	HABITAT
<i>Phytolacca</i> (Poke Root)	Root	<i>Phytolacca americana</i>	North America

ORDER RANALES

This order includes the following families: *Nymphaeaceæ* or Water Lily Family, *Magnoliaceæ* or Magnolia Family, *Ranunculaceæ* or Crowfoot Family, *Berberidaceæ* or Barberry Family, *Menispermaceæ* or Moonseed Family, *Anonaceæ* or Custard Apple

FIG. 384.—*Phytolacca americana* (in foreground), growing in damp woodland.

Family, *Myristicaceæ* or Nutmeg Family, *Monomiaceæ* or Boldo Family, and *Lauraceæ* or Laurel Family.

MAGNOLIACEÆ OR MAGNOLIA FAMILY.—Trees and shrubs having alternate leaves with large deciduous stipules that enclose the young buds and single large flowers with several series of sepals and petals which are often colored alike. Sepals and petals deciduous. Carpels and stamens numerous, spirally

arranged and free. Bark aromatic and bitter. Fruit a collection of follicles dehiscing dorsally. Seeds large, albuminous and oily.

FIG. 385.—*Myristica fragrans*—Branch and fruit. (Sayre.)

To this family belong the stately Tulip Poplar, *Liriodendron Tulipifera*, prized for its wood and beautiful flowers, and the showy Magnolias of our gardens.

OFFICIAL DRUG	PART USED	BOTANICAL NAME	HABITAT
Oleum Anisi	Volatile oil	<i>Illicium verum</i>	South China
UNOFFICIAL DRUG			
Winter's Bark	Bark	<i>Drimys Winteri</i>	South America
<i>Illicium</i> (Star Anise)	Fruit	<i>Illicium verum</i>	S. China

MYRISTICACEÆ OR NUTMEG FAMILY.—A group of apetalous trees comprising the genus *Myristica*, of about 80 species.

MYRISTICA.—A large tropical genus of fragrant, apetalous trees—the nutmegs—coextensive with the nutmeg family, having.

FIG. 386.—*Hydrastis canadensis* (Fam. *Ranunculaceæ*) $\times \frac{1}{2}$. At left, the apocarpous aggregate fruit. (After Strasburger, Noll, Schenck and Schimper.)

alternate, entire, often pellucid punctate leaves, small dioecious, regular flowers, and a succulent, two-valved, one-celled fruit with a solitary seed containing ruminant albumen and covered by a fleshy arillode.

M. fragrans, a handsome tree, 20 to 30 feet high, indigenous to the Molucca Islands in the East Indian archipelago but now cultivated extensively in the West Indies, supplies the nutmegs and mace of commerce.

OFFICIAL DRUG	PART USED	BOTANICAL NAME	HABITAT
Myristica	Kernel of seed	Myristica fragrans	} Molucca Islands
Oleum Myristicæ	Volatile oil	Myristica fragrans	
UNOFFICIAL DRUG			
Macis	Arillode	Myristica fragrans	

RANUNCULACEÆ OR BUTTERCUP FAMILY.—Herbs, rarely shrubs (*Clematis*) with acrid, poisonous, watery juices and with alternate, rarely opposite, simple, rarely compound, exstipulate leaves. Flowers pentamerous, regular to irregular, incomplete to complete, aposepalous and apopetalous. Sepals five—rarely more or less—green to petaloid, regular, passing to irregular (*Larkspur*, *Monkshood*). Petals none or five, regular to rarely irregular, often nectariferous and with nectariferous petals often variously transformed. Stamens indefinite, hypogynous. Pistil of many to few apocarpous carpels, each carpel with one to several ovules. Fruit a collection of achenes as in *Buttercup* (*Ranunculus*), *Pasque Flower* (*Pulsatilla*), or a collection of follicles (*Columbine*, *Larkspur*, *Peony*, *Aconite*) or rarely a berry as in *Baneberry* (*Actæa*) or head of crimson berries as in *Golden Seal* (*Hydrastis*). Seeds albuminous with large oil-containing endosperm and small embryo.

FIG. 387.—Above ground portion of *Aconitum Napellus* (*Monkshood*) showing palmately-divided leaves and hooded flowers.

OFFICIAL DRUG	PARTS USED	BOTANICAL NAME	HABITAT
Hydrastis (Golden Seal)	Rhizome and roots	<i>Hydrastis canadensis</i>	Eastern United States and Canada

OFFICIAL DRUG	PARTS USED	BOTANICAL ORIGIN	HABITAT
Aconitum	Tuberous root	Aconitum Napellus	Europe, Asia
Cimicifuga	Rhizome and roots	Cimicifuga racemosa	E. North America
Pulsatilla	Entire herb	{ Pulsatilla vulgaris Pulsatilla pratensis Pulsatilla patens	{ Europe and Asia W. North America
Adonis	Overground portion	Adonis vernalis	Europe and Asia
Delphinium	Seed	Delphinium Ajacis	Europe
UNOFFICIAL DRUG			
Aconiti Folia	Leaves and flower tops	Aconitum Napellus	Europe and Asia
Coptis (Goldthread)	Entire herb	Coptis trifolia	United States and Canada
Helleborus	Rhizome and roots	Helleborus niger	Alps
Nigella (Black Caraway)	Seed	{ Nigella sativa Nigella damascena	{ Europe
Staphisagria	Seed	Delphinium Staphisagria	{ South Europe, Asia Minor
Western Coptis	Entire herb	Coptis occidentalis	Western United States
Bear's Foot Root	Rhizome and roots	Helleborus foetidus	Europe
Xanthorhiza	Rhizome	Zanthorhiza apiifolia	United States

BERBERIDACEÆ OR BARBERRY FAMILY.—Herbs and woody plants with watery juices and alternate or radical, simple or pinnately-compound, mostly exstipulate leaves, either stems or leaves of *Berberis* (Barberry) species often bearing spines or barbs, which give them a barbed appearance. Rhizomes and roots often of yellow color internally, due to berberine content. Flowers either in racemes of cymes, cymes, racemes or solitary. Sepals and petals usually in 2 series of 3 each, similar, hypogynous; stamens 9 to 4, hypogynous; pistil single. Fruit a berry or capsule. Seeds exalbuminous with a straight large embryo in axis of albumen.

OFFICIAL DRUG	PARTS USED	BOTANICAL ORIGIN	HABITAT
Berberis (Barberry)	Rhizome and roots	Berberis species of the Sect. Mahonia	Western North America
Podophyllum (May Apple)	Rhizome and roots	Podophyllum peltatum	Eastern North America
Caulophyllum	Rhizome and roots	Caulophyllum thalictroides	Eastern North America
UNOFFICIAL DRUG			
Podophyllum Indicum B.P.	Rhizome and roots	Podophyllum emodi	Tibet and Afghanistan
Barberry Bark	Bark of root and stem	Berberis vulgaris	Europe

FIG. 388.—*Cimicifuga racemosa*—Plant and rhizome. Note the pinnately compound leaf, the wand-like racemes which bear white flowers, and the knotted rhizome. This perennial herb grows in rich woods. It is commonly known as Black Snakeroot and Black Cohosh and belongs to the *Ranunculaceæ*. (Sayre.)

MENISPERMACEÆ, OR MOONSEED FAMILY.—Choripetalous, woody, climbing, mostly tropical plants with alternate, exstipulate, simple often peltate leaves. Flowers green to white. Fruit

FIG. 389.—*Podophyllum peltatum*. Entire plant, above ground portion, and fruit. Note the flowering stems bearing in each instance two one-sided leaves and a nodding flower from the forks. This plant also sends up from its rhizome flowerless stems each of which terminates in a 7-9 lobed peltate leaf.

a one-seeded succulent drupe. Seeds albuminous. They usually contain tonic, narcotic or poisonous bitter principles.

OFFICIAL DRUG	PARTS USED	BOTANICAL NAME	HABITAT
Calumba	Root	Jateorrhiza palmata	East Africa
Cocculus	Fruit	Anamirta Cocculus	Asia
UNOFFICIAL DRUG			
Menispermum (Yellow Parilla)	Rhizome and roots	Menispermum canadense	United States and Canada
Pareira	Root	Chondodendron tomentosum	Brazil and Peru

MONOMIACEÆ OR BOLDO FAMILY.—Tropical trees and shrubs with opposite or rarely alternate, entire or serrate, pellucid-punctate leaves devoid of stipules, but containing cystoliths.

Flowers hermaphrodite or unisexual in cymes or racemes; calyx of many to 4 lobes or teeth in 2 or more series, stamens numerous to few, hypogynous, carpels several to one, each with

FIG. 390.—Blue Cohosh (*Caulophyllum thalictroides*). Top of plant. Note the trternately compound leaf bearing 2- to 4-lobed leaflets and the panicle of yellowish-green flowers. The fruits which ripen in August are dark blue berries. (Reproduced from U. S. Dept. Agric. Misc. Publ. 77.)

a single ovule. Fruit a drupe (*Peumus*), nut or achene. Seeds with membranous testa and fleshy endosperm.

OFFICIAL DRUG	PART USED	BOTANICAL NAME	HABITAT
Boldus	Leaf	<i>Peumus Boldus</i>	Chile

LAURACEÆ OR SPICEBUSH FAMILY.—Shrubs or trees of subtropical or tropical, rarely of temperate regions. Bark, wood and leaves rich in spicy, aromatic, hydrocarbon oils. Leaves alternate, simple, entire, often leathery and shining and exstipulate. Inflorescences usually cymose clusters. Flowers small, green, yellow or rarely whitish, hermaphrodite or more or less dioecious; regular calyx alone present as a floral whorl, of three to six

small sepals. Stamens four to twelve in several rows of three to four each; anthers continuous with the filaments, each anther opening by recurved valves, the valves opening from the base upward. Pistil monocarpellary, ovary superior, one-celled with

FIG. 391.—*Jateorhiza palmata*, the source of Calumba—Portion of vine. Note tendril for winding about a support as this plant climbs. (Sayre.)

solitary pendulous ovule, style simple with usually rounded stigma. Fruit a succulent berry (Spicebush and “*Bacca laurea*”) or a succulent drupe (Sassafras). Seeds usually richly albuminous.

OFFICIAL DRUG	PART USED	BOTANICAL NAME	HABITAT
Camphora	Ketone	Cinnamomum Camphora	Eastern Asia
Sassafras	Bark of root	Sassafras variifolium	North America

OFFICIAL DRUG	PART USED	BOTANICAL NAME	HABITAT
Oleum Sassafras	Volatile oil from root	Sassafras variifolium	North America
Cinnamomum	Bark	Cinnamomum Loureirii	Cochin China
Oleum Cinnamomi	Volatile oil	Cinnamomum Cassia	China
UNOFFICIAL DRUGS			
Cinnamomum Zeylanicum	Bark	Cinnamomum Zeylanicum	Ceylon
Coto	Bark	Nectandra Coto	Bolivia
Laurus	Leaves	Laurus nobilis	Europe
Fagot cassia	Bark	Cinnamomum Burmanii	North America
Clove Bark	Bark	Dicypellium caryophyllatum	Asia
Cassia Buds	Immature fruit	Cinnamomum Loureirii	Brazil
Cassia Bark	Bark	Cinnamomum Cassia	Cochin China
Sassafras Medulla	Pith	Sassafras variifolium	China
			North America

ORDER RHÆADALES (PAPAVERALES)

PAPAVERACEÆ OR POPPY FAMILY.—Herbs or low shrubs. Root only, in some cases root and stem, and in still others the entire plant, traversed by anastomosing latex tubes that contain a milky juice, varying from white (*Papaver* or Poppy) or pale-yellow, as in Celandine (*Chelidonium*), to red (*Bloodroot*). Latex tubes in the axis wholly confined to cortex or phloem but occasionally sending branches into the medullary rays, rarely into the pith of the plant. Leaves alternate, varying from simple (*Platystemon*) to pinnatifid, pinnatipartite or even pinnately-compound. Inflorescence varying from a loose raceme or panicle of cymes to a raceme of cymes or condensing until, as in *Papaver*, only one large terminal flower is left. Flowers regular tetramerous or dimerous. Sepals typically two, more rarely three. Petals typically four, more rarely six. Stamens indefinite, in most forms hypogynous, except in California Poppy (*Eschscholtzia*) where hypogyny is modified into perigyny. Pistil of sixteen to four, rarely two carpels generally fused together. Placenta parietal. Ovules numerous, anatropous. Fruit a

FIG. 392.—*Cinnamomum Zeylanicum*, the source of Ceylon cinnamon bark.
A member of the *Lauraceæ*. (Sayre.)

FIG. 393.—Flower of Ceylon Cinnamon (*Cinnamomum Zeylanicum*). Note the uplifted lids of the anther locules in dehiscence, the calyx of six sepals (in *a*), the perigynous stamens and 1-celled ovary with pendulous ovule, in *b*. (Agter Baillon.)

FIG. 394.—Bloodroot (*Sanguinaria canadensis*), a perennial herb native to rich open woods of eastern North America. The waxy white flowers and grayish green 5-9-lobed, kidney-shaped leaves make their appearance only in spring. The rhizome containing a red latex constitutes the drug. It should be gathered in early summer, deprived of its rootlets and carefully dried. The drug possesses expectorant and emetic properties. (Reproduced from U. S. Dept. Agric. Misc. Pub. 77.)

FIG. 395.—*Papaver somniferum*—Flowering branch and fruit. (Sayre.)

capsule (Papaver) or 2-valved pod (Bloodroot, etc.). Seeds richly albuminous.

OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Opium	Air-dried milky exudate from unripe capsules	<i>Papaver somniferum</i> and its <i>var. album</i>	Eastern Mediter- ranean countries
Morphinæ Sulfas	Sulfate of alkaloid		
Morphinæ Hydrochloridum	Hydrochloride of alkaloid		
Codeina	Alkaloid		
Codeinæ Phosphas	Phosphate of alkaloid		
Codeinæ Sulfas	Sulfate of alkaloid		
Papaverinæ Hydrochloridum	Hydrochloride of alkaloid		

OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Sanguinaria	Rhizome	Sanguinaria canadensis	United States and Canada
UNOFFICIAL DRUG			
Chelidonium	Entire flowering plant	Chelidonium majus	United States and Canada
Maw seed	Seeds	Papaver somniferum var. nigrum	Eastern Mediterranean countries
Papaveris Fructus	Unripe capsules	Papaver somniferum	Eastern Mediterranean countries
Papaver Rhœas	Fresh petals	Papaver Rhœas	Europe

FIG. 396.

FIG. 397.

FIG. 398.

FIG. 396.—Transverse section of flower of Poppy. (Sayre.)

FIG. 397.—Gynœcium of Poppy, with one stamen remaining. (Sayre.)

FIG. 398.—Transverse section of ovary of Poppy. (Sayre.)

FUMARIACEÆ OR FUMITORY FAMILY.—Delicate herbs, rarely shrubs containing milky-watery to watery latex containing alkaloids. Leaves radical, alternate and more or less compound. Inflorescence a raceme or spike. Flowers irregular, hermaphrodite, zygomorphic; sepals 2, petals 4, the two outer often saccate or with a spur at the base; stamens 4 or 6; ovary superior, 1-celled with 2 parietal placentas. Fruit a one-chambered capsule. Seeds albuminous. Idioblasts common.

OFFICIAL DRUG	PART USED	BOTANICAL NAME	HABITAT
Corydalis	Tubers	Dicentra canadensis Dicentra Cucullaria	{ United States and Canada

CRUCIFERÆ OR MUSTARD FAMILY.—Herbs, rarely shrubs, mostly of temperate regions. Many species of this family contain glucosides which on hydrolysis yield pungent volatile oils. Stem and branches upright or diffuse spreading (*Arabis*). Leaves alternate, simple, rarely compound, exstipulate, entire or toothed, often more or less hairy. Inflorescence at first

a corymb or shortened raceme, later elongating into a loose raceme. Bracts at base rarely reduced, usually absent. Flowers regular, rarely irregular (Candytuft), tetramerous. Sepals four, green, equal, or two laterals at times pouched as nectar receptacles. Petals four, yellow, white, pink or purple, cruciform, often divisible into claw and blade. Stamens six, four being long and anteroposterior in position, two being short and lateral,

FIG. 399.—The Black Mustard, *Brassica nigra*—Branch. (Sayre.)

hence termed tetradynamous, insertion hypogynous. Pistil syncarpous, bicarpellate, superior, carpels lateral. Ovary one-celled but falsely two-celled by a placental replum (a spurious membranous partition dividing the ovary into 2 locules); style simple; stigma rounded or bifid or bilobed. Ovules several, rarely few, attached to marginal placenta. Fruit a siliqua, as in the mustards or silicule, as in the shepherd's purse (*Capsella*) rarely indehiscent, bursting lengthwise by two valves. Seeds exalbuminous with folded cotyledons. (See Fig. 400.)

OFFICIAL DRUG	PART USED	BOTANICAL NAME	HABITAT
Oleum Sinapis	Volatile oil	{ <i>Brassica nigra</i>	Europe and Asia
Volatile		{ <i>Brassica juncea</i>	Asia
Sinapis Nigra	Seed	{ <i>Brassica nigra</i>	Europe and Asia
		{ <i>Brassica juncea</i>	Asia
UNOFFICIAL DRUG			
Semen Rapæ	Seed	<i>Brassica napus</i>	Europe and Asia
Armoracia	Root	<i>Cochlearia Armoracia</i>	Europe
Sinapis Alba	Seed	<i>Sinapis alba</i>	Europe and Asia
Capsella	Aerial plant	<i>Capsella bursa-pastoris</i>	Europe

ORDER SARRACENIALES

This order includes several families of plants whose leaves are modified for the purpose of catching and digesting insects.

The principal families are the *Sarraceniaceæ* and *Nepenthaceæ* both of which have their leaves modified as pitchers, and the *Droseraceæ* which includes the Fly-Traps and Sundews. (See pp. 622-638.)

FIG. 400.—Cruciferae. Floral diagram above; flower in median longitudinal section below. (Robbins.)

DROSERACEÆ OR SUNDEW FAMILY. Herbs (*Drosera*, *Dionaea*, etc.) rarely shrubs (*Roridula* of South Africa), growing in bogs or swamps or purely aquatic in habit (*Aldrovanda*). Leaves, either rosettes or more or less scattered in alternate fashion over stem, usually glandular-hairy or sensitive-hairy and insectivorous. Inflorescence a loose raceme or cymose umbel. Flowers regular, pentamerous; sepals five, aposepalous, green; petals apopetalous, varying from white to whitish-pink, pink, scarlet, purple to bluish-purple; stamens varying from fifteen to five, hypogynous; pistil syncarpous of five to three, rarely fewer carpels. Fruit a capsule. Seeds albuminous.

UNOFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
<i>Drosera</i> (Sundew)	Entire plant	{ <i>Drosera rotundifolia</i> <i>Drosera anglica</i> <i>Drosera longifolia</i>	{ Eastern and west- ern Hemispheres

ORDER ROSALES

SAXIFRAGACEÆ OR SAXIFRAGE FAMILY.—Herbs (Mitrewort (*Mitella*), Saxifrage (*Saxifraga*), Alum-root (*Heuchera*), etc.) or shrubs (*Hydrangea*, *Philadelphus* or Mock-orange, etc.) rich in tannin content, with opposite or alternate leaves usually devoid of stipules. Flowers regular, hermaphrodite; sepals 5 usually,

FIG. 401.—*Drosera rotundifolia*. Production of a young plant from the leaf of an older plant. (Gager.)

petals 5 to 10, sometimes absent, petals and stamens perigynous to epigynous; ovary 3 to 1 celled, free or adnate to upgrown receptacle. Fruit a follicle, capsule, or berry. Seeds small with copious albumen and tiny embryo.

OFFICIAL DRUG	PARTS USED	BOTANICAL NAME	HABITAT
Hydrangea	Rhizome and roots	Hydrangea arborescens	United States
UNOFFICIAL DRUG			
Heuchera	Root	Heuchera americana	United States

HAMAMELIDACEÆ OR WITCH HAZEL FAMILY.—Shrubs or small or large trees. Leaves simple, alternate and usually pinnately veined; stipules deciduous to caducous, paired and slightly fused

at the bases of petioles. Flowers frequently yellow to yellowish-white, in axillary clusters or heads or spikes, hermaphrodite or monœcious; sepals and petals five to four, rarely indefinite, superior (petals absent in *Fothergilla*); stamens twice as many as the petals, but the outer row alone fertile, the inner row being more or less barren, scale-like; gynœcium of two carpels united

FIG. 402.—*Liquidambar orientalis*, the Sweet Gum—Branch. (Sayre.)

below. Fruit a two-beaked, two-celled, woody capsule dehiscing at the summit, with a bony seed in each cell, or several, only one or two of them ripening.

OFFICIAL DRUG	PART USED	BOTANICAL NAME	HABITAT
Styrax (Storax)	Balsam from wood and inner bark	Liquidambar orientalis	The Levant
		Liquidambar Styraciflua	Central America
Hamamelidis Folium	Leaf	Hamamelis virginiana	United States and Canada
UNOFFICIAL DRUG			
Hamamelidis Cortex	Bark	Hamamelis virginiana	United States and Canada

FIG. 403.—*Hamamelis virginiana*. Upper figure shows this shrub as it appears in autumn after the leaves have fallen. Note that the plant is in blossom. Lower figure shows a flowering branch from the same plant. The bright yellow flowers occur in axillary clusters appearing at the same time as the ripening of fruits from blossoms of the previous year. A leaf of the plant, commonly known as Witch Hazel, is shown on p. 239. The recently cut and partially dried twigs and leaves are used in the preparation of Witch-hazel water or Aqua Hamamelidis.

ROSACEÆ OR ROSE FAMILY.—Herbs, shrubs, or trees mostly of temperate regions. Stem and branches upright or creeping (Strawberry, Cinquefoil), herbaceous to woody. Leaves alternate, stipulate (stipules green, persistent to scaly, deciduous), compound, condensing to “simple.” Flowers regular, pen-

FIG. 404.—Branch of *Quillaja Saponaria*, a Chilean tree from whose trunk and branches Soap Tree Bark is obtained. (Sayre.)

tamerous, either solitary, as in the Peach (*Prunus persica*) or Apricot (*Prunus armeniaca*), in racemes, as in the Wild Black Cherry (*Prunus serotina*), in cymes, as in Apple (*Malus*), or in umbels as in Bird Cherry (*Prunus Avium*) and Sour Cherry (*Prunus Cerasus*), sepals and petals 5, rarely 4, inferior to ovary, becoming by stages superior to it. Sepals green, at times with

epicalyx (Strawberry, Cinquefoil, Kousso, etc.), persistent round fruit. Petals usually yellow to white or pink, crimson, rarely purple, rosaceous, deciduous. Stamens indefinite, perigynous (Strawberry, etc.), to semi-epigynous (Rose, Peach, etc.), and epigynous (Apple, Pear, Quince). Pistil apocarpous with many (Strawberry, Rose) carpels or fewer to five (Apple), or two to one (Plum, Cherry), becoming falsely fused by union with upgrowing receptacle (Hawthorn, Quince, Pear, Apple). Fruit a collection of achenes on dry (Cinquefoil) or succulent receptacle (Strawberry), or dry follicles (Bridal Wreath), or drupels (Blackberry), or a drupe (Peach, Plum, Cherry), or a pome (Apple, Quince, Pear). Seeds exalbuminous, embryo filling seed cavity.

Some botanists divide this family into three sub-families, each of which, in the former American System, had been given the rank of a family. These according to this system of classification are: *Rosaceæ*, *Malaceæ* and *Amygdalaceæ*. *Rosaceæ* include the genera *Geum*, *Agrimonia*, *Duchesnia*, *Fragaria*, *Potentilla*, *Waldsteinia*, *Cercocarpus*, *Spiræa*, *Rubus*, *Dalibarda* and *Rosa*.

Malaceæ include the genera *Sorbus* (Mt. Ash), *Malus* (Crab Apple and Apple), *Amelanchier* (Service Berry and Shad Bush), *Pyrus* (Pear), *Aronia* (Choke Berry), *Cratægus* (Hawthorn) and *Cotoneaster* (Christ's Thorn).

Amygdalaceæ comprise the genera *Prunus* (Plum group), *Padus* (Wild Cherry group) and *Amygdalus* (Almond and Peach group).

OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Prunus Virginiana	Bark	Prunus serotina	United States and Canada
Quillaja	Inner bark	Quillaja Saponaria	Chile and Peru
Rosa	Petals	Rosa gallica	W. Asia and S. Europe
Oleum Rosæ	Vol. oil	Rosa gallica	{ W. Asia and S. Europe Balkans
		Rosa damascena	
Rubus Idæus	Fresh ripe fruit	Rubus Idæus and Rubus strigosus	Europe and Asia
Prunus Cerasus	Fresh ripe fruit	Prunus Cerasus	Asia Minor
Oleum Amygdalæ Amaræ	Vol. oil from kernels	Amygdalus communis, Prunus persica, etc.	W. Asia
Oleum Amygdalæ Expressum	Fixed oil	Amygdalus communis (var's.)	W. Asia

UNOFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Amygdala Dulcis	Seed	Amygdalus communis var. dulcis	W. Asia
Amygdala Amara	Seed	Amygdalus communis	W. Asia
Folia Laurocerasi	Leaves	Prunus Laurocerasus	Asia Minor, Persia
Cýdonium	Seed	Cydonia vulgaris	Cultivated widely
Rosa Centifolia	Petals	Rosa centifolia	Western Asia
Rosa Canina	Spurious Fruit	Rosa canina	Europe
Tormentilla	Rhizome	Potentilla silvestris	Europe and Asia
Rubus	Bark of rhizome and roots	{ Sect. Eubatus of genus Rubus	United States
Prunum	Fruit		
Brayera	Panicles of pistillate flowers	Hagenia abyssinica	Abyssinia
Succus Pomorum	Fresh juice of fruit	Malus (cult. var's.)	Europe

FIG. 405.—*Prunus domestica*, or Damson plum, the source of prunes. Fruiting branch and flowering branch. (Sayre.)

LEGUMINOSÆ OR PEA FAMILY (FABACEÆ).—Herbs, shrubs or trees of all regions, with tubercled roots. Stem usually erect, rarely creeping (*Trifolium repens*). Leaves alternate, compound, rarely simple, stipulate, sometimes tendriform, as in Peas, or reduced to phylloid petioles (some *Acacia spp.*). Inflorescence a

FIG. 406.—*Glycyrrhiza glabra*—Branch. (Sayre.) The source of Licorice. The varieties yielding the drug are perennial herbs with alternate, imparipinnate leaves and axillary spikes of violet-colored papilionaceous flowers.

FIG. 407.—Flower of Papilionaceæ. A, floral diagram of *Vicia faba*; B, sweet pea flower, dissected, diagrammatic. (From Robbins, A after Eichler, B after Bergen and Caldwell.)

raceme, at times, condensed almost to a head or capitulum (Spp. of clover, *Mimosa*, etc.). Flowers pentamerous (rarely tetramerous as in *Krameria*), regular (*Mimosaceæ*), to irregular (*Cæsalpinaceæ*) to very irregular with corolla butterfly-shaped (*Papilionaceæ*). Sepals five united, green; petals five (rarely four) variously related; in *Papilionaceæ* the one superior, external

FIG. 408.—*Cassia Senna*—Branch showing flower and fruit. (Sayre.)

and posterior is called the standard or *vexillum*, the two lateral form the wings or *alæ*, the two inferior, internal and anterior, slightly adherent, form the *keel*. Stamens ten to four, free, or in *Papilionaceæ* united by filaments into a monadelphous (ten) or a diadelphous (nine to one) tube, inserted perigynously. Pistil typically monocarpellary, ovary with sutural placentation, style simple. Fruit a legume, more rarely becoming transformed into a loment by transverse constrictions (*Entada scandens*). Seeds exalbuminous.

There are three natural subdivisions of the *Leguminosæ*: (1) The *Mimosaceæ* or *Mimosa* alliance is characterized by possessing regular flowers, often in capitula, the petals of which do not overlap or are they overlapped, and by often furnishing mucilage. To this group belong the plants yielding Acacia and Catechu. (2) The *Cæsalpinaceæ* which have slightly irregular flowers, the petals of which are imbricate in aestivation, the odd posterior petal in the flower being overlapped by the 2 side petals. To this group belong the plants yielding Copaiba, Hæmatoxylon, Senna, Cassia Fistula, Brazil Wood, Peach Wood, Sappan and Tamarindus. (3) The *Papilionaceæ* which have very irregular flowers, the petals of which are imbricate in aestivation, but the odd petal here called the standard, overlaps the 2 side petals and the two side petals overlap the keel petals. To this group belong the plants yielding Tolu, Balsamum Peruvianum, Baptisia, Glycyrrhiza, Goa Powder, Foenugreek, Galega, Melilotus, Trifolium, Tragacantha, Tonka, Santalum Rubrum, Scoparius, Physostigma, etc.

OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Acacia	Gummy exudation	Acacia Senegal and some other African species of Acacia	Africa
Cascara Amarga	Bark	Sweetia panamensis	S. Mexico to Panama
Tragacantha	Gummy exudation	Astragalus gummifer and other Asiatic species	Asia
Balsamum Peruvianum	Balsam	Toluifera Pereiræ (Myroxylon Pereiræ)	Central America
Balsamum Tolutanum	Balsam	Toluifera Balsamum	Northern South America
Stantalum Rubrum	Heartwood	Pterocarpus santalinus	Indo-China and S. India
Glycyrrhiza	Rhizome and roots	{ Glycyrrhiza glabra var. typica Glycyrrhiza glabra var. glandulifera and other var's. of G. glabra yielding a yellow and sweet wood	{ S. Europe, W. and C. Asia
Senna	Leaflets	{ Cassia Senna Cassia angustifolia	{ Egypt and the Soudan Arabia and Africa

OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Copaiba	Oleoresin	Copaiba species	South America
Chrysarobinum	Neutral principle	Vouacapoua Araroba	Brazil
Kino	Inspissated juice	Pterocarpus Marsupium	India
Trifolium	Inflorescence	Trifolium pratense	Europe
Krameria	Root	{ Krameria triandra	Peru and Bolivia
		{ Krameria argentea	Brazil
Physostigminæ Salicylas	Salicylate of alka- loid from seed	Physostigma veneno- sum	Africa
UNOFFICIAL DRUG			
Baptisia	Root	Baptisia tinctoria	Eastern North America
Hæmatoxylon	Heartwood	Hæmatoxylon campechianum	Central America
Tamarindus	Pulp of fruit	Tamarindus indica	Tropical Africa
Cassia Fistula	Fruit	Cathartocarpus Fistula	India
Scoparius	Tops	Cytisus Scoparius	Europe
Galega	Flowering herb	Galega officinalis	S. Europe and W. Asia
Sappan	Heartwood	Caesalpinia Sappan	India, Malaya
Brazil Wood	Heartwood	Caesalpinia brasiliense	South America
Peachwood	Heartwood	Caesalpinia echinata	South America
Derris (Tuba Root)	Rhizome and root	{ Derris elliptica	{ Malaya
		{ Derris malaccensis	{ E. Indies
Cube Root	Root	Lonchocarpus Spp.	South America
Physostigma	Seed	Physostigma venenosum	Africa
Melilotus	Leaf and flower- ing top	Melilotus officinalis	Europe
Mesquite Gum	Gum	Prosopis juliflora	Mexico
Fenugreek	Seed	Trigonella Fœnum-græcum	Mediterranean region
Piscidia	Bark	Piscidia Erythrina	West Indies
Indigo	Coloring matter	Indigofera tinctoria	India
Soy Bean	Seed	Soja max	China and Japan
Abrus (Jequirity)	Seed	Abrus precatorius	Tropics and sub- tropics
Catechu	Extract	Acacia Catechu	India
Mucuna (Cowhage)	Hairs from fruits	Mucuna pruriens	East and West Indies

ORDER GERANIALES OR GRUINALES

GERANIACEÆ OR GERANIUM FAMILY.—Herbaceous, rarely semi-succulent, sub-shrubby plants. Stems cylindrical, often hairy or glandular hairy. Leaves alternate to opposite, stipulate;

venation from pinnate to palmate, so leaf shape from ovate to pinnatifid to pinnatipartite to sub-palmatifid to palmatipartite to compound palmate. Inflorescence either a dichasial or scorpioid cyme. Flowers regular, pentamerous (*Geranium*) to irregular pentamerous (*Pelargonium*); sepals five, aposepalous; petals five,

FIG. 409.—*Physostigma venenosum*—Portion of plant and fruit. (Sayre.)

apopetalous, varying in color from greenish-white or pink-red to scarlet, scarlet-crimson to crimson-purple; stamens with anthers ten or five, hypogynous or inserted into slightly developed hypogynous disk; pistil pentacarpellary, ovary five-celled with two, rarely one ovule in each cell, styles elongate, fused round a stylar column of receptacle then continued as a stylar tip which splits into five stigmatic surfaces. Fruit a regma, rarely a simple capsule. Regma splits into five recurved carpels, each then dehiscing to set free two or one seeds. Seeds exalbuminous.

UNOFFICIAL DRUG	PART USED	BOTANICAL NAME	HABITAT
Geranium	Rhizome	<i>Geranium maculatum</i>	North America
Oil of Rose } Geranium }	Volatile oil	<i>Pelargonium</i>	Mediterranean regions
		<i>odoratissimum</i>	
		<i>Pelargonium Radula</i>	
		<i>Pelargonium capitatum</i>	

LINACEÆ OR FLAX FAMILY.—Herbs with slender stems and alternate, simple, narrow leaves. Inflorescence cymose with regular pentamerous flowers; pistil five-carpelled with a five-celled ovary containing two ovules in each cavity and having a single

FIG. 410.

FIG. 410.—Cranesbill (*Geranium maculatum*), a perennial herb with palmately-parted leaves, pale purple or rose-purple flowers, a tuberculated rhizome and regma fruits. The latter, appearing like the bill of a crane, are seen to arise from slender stalks forming a fork near the center of the picture. (Reproduced from U. S. Dept. Agriculture Misc. Publ. 77.)

FIG. 411.

FIG. 411.—*Linum usitatissimum*, an annual herb with linear-lanceolate leaves, pale blue flowers and capsular fruits, the source of Flaxseed, Linseed oil and Linen fiber. (Sayre.)

style with a knob-like stigma. While the flower is still in bud condition or soon after, there commences an ingrowth of the mid-rib of each carpel which proceeds until, when plant is in

fruit, there are formed 10 cavities each enclosing a seed. Seeds, anatropous, mucilaginous, flattened, containing a large embryo and small enveloping endosperm.

OFFICIAL DRUG	PART USED	BOTANICAL NAME	HABITAT
Linum	Seeds	Linum usitatissimum	Unknown. Cult. in Temperate regions
Oleum Lini	Fixed oil	Linum usitatissimum	

ERYTHROXYLACEÆ OR COCA FAMILY.—Shrubs (*Erythroxylon*) or trees with alternate, simple, entire, glabrous and pinnately

FIG. 412.—*Guajacum sanctum*—Flowering branch. (Sayre.)

veined leaves, the cells of the lower epidermis of which show papillæ. Flowers regular, hermaphroditic, each with five sepals, five hypogynous petals, ten stamens and a two- to three-celled ovary subtending three styles, each with a capitate stigma. Fruit an ovoid, angular, one celled drupe containing a single seed.

OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Cocaina	Alkaloid	{ Erythroxylon Coca and other species of Erythroxylon }	{ Peru and Bolivia, South America }
Cocainæ	Hydrochloride of		
Hydrochloridum	alkaloid		
UNOFFICIAL DRUG			
Coca	Leaves	Erythroxylon Coca and E. truxillense	Peru and Bolivia

ZYGOPHYLLACEÆ OR CALTROP FAMILY.—Herbs, shrubs, or trees (*Guajacum* or *Guaiac*) having jointed, often divaricate

FIG. 413.—*Citrus Aurantium*, the bitter or Curaçao orange, a small tree bearing evergreen leaves with winged petioles, white flowers and a spherical fruit with bitter, acid pulp. The source of bitter orange peel and volatile oils of the peel and flowers. (Sayre.)

branches. Leaves usually opposite, stipulate and compound. Flowers regular or irregular, pentamerous, white, yellow, red or blue (*G. officinale*). Fruit a capsule.

OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Guaiacum	Resin of wood	{ Guajacum officinale Guajacum sanctum }	{ W. Indies and S. America }

UNOFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Guaiaci Lignum	Heartwood	G. officinale and G. sanctum	{ W. Indies and S. America

RUTACEÆ OR RUE FAMILY.—Herbs (*Rue*), shrubs (*Jaborandi* and *Buchu*) or trees (*Orange*, *Lemon*, *Grape Fruit*, etc.). Stems upright, often wiry and xerophytic in the Rue sub-family or *Ruteæ*, elongated and spiny in the Prickly Ash sub-family or *Zanthoxy-leæ*, woody and green in the Citrus sub-family or *Aurantieæ*. Leaves alternate or opposite, simple (*Buchu*), rarely whorled (*Pilocarpus*) or pin-natifid, as in the Rue plant (*Ruta graveolens*), or pinnate, as lower parts of *Ruta graveolens*, becoming reduced pinnate in the Orange (*Citrus Aurantium*). Leaves exstipulate or with spiny stipules (*Zanthoxylum*). Stems and leaves abound in more or less sunken glands. Flowers pentamerous, varying in color from yellow in *Ruta* to white in *Citrus* to pink (*Barosma betulina*) or pink crim-son, as in some *Barosma* and *Diosma* species, rarely to pinkish-purple (*Pilocarpus*); sepals five, aposepalous, becoming in *Citrus* more or less synsepalous; petals five, apopetalous, becoming more or less synpetalous and tubular (*Correa grandiflora*); stamens five, simple or with expanded bases, lobed, or lobes developed as staminal stipules and more or less split (*Citrus*); pistil of ten, five, three or two carpels, ovary as many-celled, with frequently 2 to 1 ovules in each cell. Fruit a capsule (*Dittany*, *Zanthoxylum*), a hesperidium (*Citrus*) or rarely a samara, as in the Wafer Ash (*Ptelea*). Seeds albuminous or exalbuminous. Many of the plants contain volatile oils in their secretory cavities.

FIG. 414.—*Barosma betulina*, the source of Short Buchu, a S. African shrub. Branch and flower. (Sayre.)

OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Aurantii Dulcis Cortex	Fresh outer rind of ripe fruit	Citrus Aurantium var. sinensis	Northern India
Oleum Aurantii	Vol. oil from fresh peel of ripe fruit	Citrus Aurantium var. sinensis	Northern India
Aurantii Amari Cortex	Rind of fruit	Citrus Aurantium	Northern India
Limonis Cortex	Outer rind of ripe fruit	Citrus medica var. Limon	Northern India
Oleum Limonis	Vol. oil from fresh fruit peel	Citrus medica var. Limon	Northern India
Pilocarpinæ Nitras	Nitrate of alka- loid from leaflets	{ Pilocarpus Jaborandi Pilocarpus microphyllus }	Brazil
Pilocarpinæ Hydrochloridum	Hydrochloride of alkaloid from leaflets		
Buchu	Leaves	{ Barosma betulina, B. crenulata and B. serratifolia }	S. Africa
Xanthoxylum (Prickly Ash)	Bark	{ Zanthoxylum americanum Zanthoxylum Clava-Herculis }	Northern United States Southern United States
Oleum Bergamottæ	Volatile oil from fresh fruit rind	Citrus Aurantium var. Bergamia	France, Italy
Oleum Aurantii Florum	Volatile oil from fresh flowers	Citrus Aurantium	Northern India
Oleum Aurantii Amari	Volatile oil from fresh peel	Citrus Aurantium	Northern India
Xanthoxyli Fructus	Capsules	{ Zanthoxylum americanum Zanthoxylum Clava-Herculis }	Northern United States Southern United States
UNOFFICIAL DRUG			
Ruta (Rue)	Leaves	Ruta graveolens	Southern Europe
Ptelea (Wafer Ash)	Bark of root	Peltea trifoliata	North America
Belæ Fructus	Unripe fruit	Ægle Marmelos	Malabar, India
Pilocarpus	Leaflets	{ Pilocarpus Jaborandi Pilocarpus microphyllus }	Brazil
Cusparia Bark	Bark	Galipea officinalis	
			Venezuela

SIMARUBACEÆ OR AILANTHUS FAMILY.—A family of chiefly tropical shrubs or trees containing bitter principles. The leaves are mostly alternate and pinnate. The flowers are unisexual or polygamous and arranged in axillary panicles (*Picrasma excelsa*) or in terminal racemes (*Quassia amara*). The fruit is a drupe

FIG. 415.—Branch of a Quassia yielding plant, *Picrasma excelsa*. Note the imparipinnate leaves with winged petioles and axillary flower clusters. (After Sayre.)

(*Picrasma*) or pod (*Quassia*). The plants are distinguished from those of the *Rutaceæ* by the absence of secretory cavities.

OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Quassia	Wood	{ Picrasma excelsa Quassia amara	West Indies Surinam
UNOFFICIAL DRUG			
Simaruba	Bark of root	{ Simaruba officinalis Simaruba amara	South America Central America Bahamas and Florida

BURSERACEÆ OR MYRRH FAMILY.—Shrubs and trees of tropical climes having secretion reservoirs in their bark which

contain gum-resin or oil. Leaves alternate, exstipulate and compound (trifoliate in *Commiphora Myrrha*). Flowers small,

regular and hermaphrodite, arranged in racemes or panicles. Fruit a drupe.

FIG. 416.—*Commiphora myrrha*—Branch.
(Sayre.)

OFFICIAL DRUG	PART USED
Myrrha	Gum resin
BOTANICAL ORIGIN	HABITAT
<i>Commiphora</i>	N. E. Africa and
<i>abyssinica</i> et al. spp.	Arabia
UNOFFICIAL DRUG	PART USED
Olibanum	Gum resin
BOTANICAL ORIGIN	HABITAT
<i>Boswellia carterii</i>	East Africa and
	Arabia

MELIACEÆ OR MAHOGANY FAMILY.—Tropical trees or shrubs with wood often hard, colored and odoriferous. Leaves alternate, exstipulate, pinnately-compound, rarely simple and entire. Inflorescence a terminal or axillary raceme. Flowers hermaphrodite or rarely polygamo-dioecious, regular; sepals five to four, small; petals usually five to four, hypogynous; stamens generally ten to eight, rarely

five, very rarely twenty to sixteen, inserted outside the base of the hypogynous disc; filaments united into a tube; carpels usually five to three; style simple; ovary free, usually five- to three-celled. Fruit a drupe (*Melia*), berry (*Vavaea*), or capsule (*Cedrela*). Seeds exalbuminous or with fleshy albumen.

UNOFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Azedarach	Bark of root	<i>Melia</i> Azedarach	Asia
Cocillana	Bark	<i>Guarea</i> Rusbyi	Bolivia

POLYGALACEÆ OR MILKWORT FAMILY.—Herbs or shrubs with upright, herbaceous to woody stems often branching pro-

fusely, the branches occasionally becoming geotropic or subterranean and bearing cleistogamous flowers. Leaves simple, often lanceolate or linear, exstipulate, alternate. Inflorescence a raceme, spike (*Polygala Senega*) or head (*P. lutea*). Flowers

FIG. 417.—*Polygala senega*—Plant and rhizome. (Sayre.)

irregular, hermaphroditic with 5 distinct sepals, the 2 lateral ones being large and petaloid, 5 petals of which the two lateral are wanting or rudimentary and the anterior large and boat-shaped, eight stamens, and a bicarpellate pistil. Fruit a two-celled capsule (*P. Senega*), rarely a drupe or samara. Pollen grains barrel-shaped.

OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Senega	Root	Polygala Senega	United States and Canada

EUPHORBIACEÆ OR SPURGE FAMILY.—Often herbaceous, more rarely shrubby, rather seldom arborescent plants. Stem, leaves and other parts in several genera traversed by latex canals

FIG. 418.—The Castor Oil plant (*Ricinus communis*). (From Strasburger after Baillon.)

that are either ramifying cells (*Euphorbia*) or laticiferous vessels (*Manihot*, *Hevea*, etc.) or rows of laticiferous sacs (*Micrandra*) and contain a white latex with acrid often poisonous contents or alkaloid or hydrocarbon, at times, rubber contents. Leaves alternate, exstipulate to stipulate, simple to pinnate or palmate. Inflorescence cymose. Flowers usually, as in *Ricinus*, etc.,

pentamerous, monœcious; sepals five, green, aposepalous, becoming rudimentary or absent in *Euphorbia*. Petals none or five more or less petaloid; stamens numerous to ten to five or one (*Euphorbia*); pistil in pistillate flowers rarely of twenty to ten apocarpous or loosely syncarpous carpels (Sandbox tree), commonly of three syncarpous carpels with distinct radiate styles; ovary as many-celled as carpels with two to one ovule in each cell. Fruit a tricoccoid regma or capsule, rarely winged, indehiscent, nutlike. Seeds with oily endosperm. Flowers at times surrounded and subtended by more or less petaloid and expanded bracts and bracteoles.

OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Euphorbia Pilulifera	Herb	Euphorbia pilulifera	India
Stillingia	Root	Stillingia sylvatica	Southern United States
Kamala	Hairs of capsule	Mallotus philippinensis	Asia
Oleum Ricini	Fixed oil	Ricinus communis	India
Oleum Tiglii	Fixed oil	Croton Tiglium	Asia
UNOFFICIAL DRUG			
Tapioca	Starch	Manihot utilissima	South America
Elastica (Rubber)	Prepared latex	Hevea brasiliensis and other species	Brazil
Cascarilla	Bark	Croton Eluteria	West Indies
Euphorbium	Resinous latex	Euphorbia resinifera	Morocco
Tung Oil	Fixed oil	Aleurites cordata	China

ORDER SAPINDALES

ANACARDIACEÆ OR SUMAC FAMILY.—Shrubs or trees producing in stems and leaves secretion contents that are either acrid-watery or acrid-opalescent or white-viscid, viscid-acrid and poisonous. Leaves alternate, rarely opposite, simple (*Rhus Cotinus*), three foliate (*Rhus toxicodendron*) or pinnate (*Rhus glabra*, *R. venenata*, etc.). Inflorescence frequently terminal and composed of racemes of cymes, often reduced to a simple raceme. Flowers small, clustered, green, greenish-white to greenish-yellow; sepals five, rarely six or four green, small; petals five, smaller than sepals; stamens equal in number to the petals and alternate, rarely fewer, sometimes double in number, rarely

indefinite, inserted hypogynously or upon an enlarged disc that surrounds or swells up between stamens and pistil; pistil monocarpellary (*Rhus*), more rarely bicarpellary, very rarely, as in *Spondiæ*, of ten to five carpels, ovary one-celled with single ovule. Fruit a drupe. Seeds exalbuminous with large embryo filling seed cavity.

OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Mastiche	Concrete resinous exudate	<i>Pistacia Lentiscus</i>	Grecian Archipelago
UNOFFICIAL DRUG			
Chinese Galls	Excrescences	<i>Rhus semialata</i>	China
Japanese Galls	Excrescences	<i>Rhus japonica</i>	Japan
<i>Rhus Glabra</i>	Ripe fruits	<i>Rhus glabra</i>	Canada and United States
<i>Rhus Typhina</i>	Ripe fruits	<i>Rhus typhina</i>	United States
Acajou Gum	Gum	<i>Anacardium occidentale</i>	West Indies
Pistachio	Seeds	<i>Pistacia vera</i>	Western Asia
<i>Anacardium</i>	Fruit	<i>Anacardium occidentale</i>	West Indies
<i>Rhus Toxicodendron</i>	Fresh leaflets	<i>Rhus toxicodendron</i>	United States

AQUIFOLIACEÆ (ILICACEÆ) OR HOLLY FAMILY.—Trees or shrubs, chiefly evergreen; leaves alternate, petiolate, simple with entire to spiny-toothed margin (*Ilex opaca* or American Holly). Flowers regular, small, greenish to white, hermaphrodite or polygamous or dioecious. Fruit drupaceous and containing 4 to 6 nutlets. Caffeine has been obtained from Maté and Cassena.

UNOFFICIAL DRUG	PART USED	BOTANICAL NAME	HABITAT
Maté	Leaves	<i>Ilex paraguariensis</i>	South America
Cassine (Cassena)	Leaves	<i>Ilex vomitoria</i>	Southeastern United States
Black Alder (Prinos)	Bark	<i>Ilex verticillata</i>	North America

CELASTRACEÆ OR STAFF TREE FAMILY.—Shrubs (*Evonymus* or Burning Bush), or shrubby climbers (*Celastrus* or Climbing Bittersweet). Leaves rarely alternate (*Celastrus*), usually opposite (*Evonymus*, *Pachystima*), simple, entire or toothed. Inflorescence of axillary cymes of terminal racemes. Flowers perfect (*Evonymus*, *Pachystima*) or polygamo-dioecious (*Celastrus*), greenish (*Celastrus*), greenish or yellowish-white (*Evonymus Europæus*),

greenish-purple (*Evonymus americanus*) to dark purple (*Evonymus atropurpureus*); calyx four- to five-lobed; corolla of four to five petals; stamens four to five, perigynous, on a disc, which fills the base of the calyx and sometimes covers the ovary; ovary three- to five-celled. Fruit a two- to five-celled capsule. Seeds albumi-

FIG. 419.—The Burning Bush or Wahoo, *Evonymus atropurpureus*. Flowering branch to left; fruiting branch to right.

nous with fleshy reddish aril (*Evonymus*, *Celastrus*) or white membranous aril (*Pachystima*).

OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Euonymus	Bark of root	Evonymus atropurpureus	United States

SAPINDACEÆ OR SOAPWORT FAMILY.—Trees (*Sapindus*), shrubs (*Paullinia*), undershrubs or rarely herbaceous vines of tropical

climes (*Cardiospermum*) containing the glucoside saponin. Stem erect or climbing (*Paullinia*) often provided with tendrils. Leaves commonly alternate and compound. Flowers in racemes or panicles (*Paullinia*) or corymbs (*Cardiospermum*), perfect or polygamodioecious, yellowish in *Paullinia Cupana*. Fruit a capsule (*P. Cupana*), samara, drupe or berry (*Sapindus*). Seeds exalbuminous.

FIG. 420.—Cascara Sagrada (*Rhamnus purshiana*). Leaf and fruiting branch. The cured bark constitutes a valuable tonic-laxative. (Reproduced from U. S. Dept. Agric. Misc. Publ. 77.)

OFFICIAL DRUG	PART USED
Guarana	Dried Paste, made from crushed seed
BOTANICAL ORIGIN	HABITAT
<i>Paullinia Cupana</i>	Brazil

ACERACEÆ OR MAPLE FAMILY.—Chiefly trees, occasionally shrubs, of temperate regions with watery sap. Leaves opposite, simple and palmately lobed or cleft (*Acer*) or pinnate (*Negundo*). Inflorescence a raceme, condensing in some species to a capitulum of cymes. Flowers small, regular, polygamous or dioecious; sepals five to four, green; petals none or five,

variously colored; stamens usually eight, hypogynous or perigynous; nectar disc around stamens or between them and pistil; pistil bicarpellary with two-celled ovary. Fruit a samara. Seeds green, exalbuminous with coiled or folded embryo and long thin cotyledons.

UNOFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
<i>Acer Spicatum</i>	Bark	<i>Acer spicatum</i>	United States

ORDER RHAMNALES

RHAMNACEÆ OR BUCKTHORN FAMILY.—Shrubs or low trees usually of branching or spreading habit. Branches either cylindric, long, green or hardened, checked back and spinescent, occasionally, especially flowering branches, developing

tendrils for support. Leaves simple, usually alternate and stipulate. Inflorescence cymose. Flowers hermaphrodite or more or less diclinous, pentamerous to tetramerous, greenish to greenish-yellow to yellowish-white; sepals five to four; petals five to four alternating with sepals; stamens five opposite the

FIG. 421.—Alder Buckthorn or *Rhamnus Frangula*—Branch. (Sayre.)

petals, perigynous; pistil either free in center of receptacular cup or more or less fused with it and so semi-inferior, ovary typically three-celled, becoming rarely four-celled with two to one atropous ovules in each cavity. Fruit of three indehiscent cocci, each enclosing a single albuminous seed with straight embryo imbedded in albumen.

OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Cascara Sagrada	Bark	<i>Rhamnus Purshiana</i>	Northern California, Oregon, Washington, and southwestern British America
Frangula	Bark	<i>Rhamnus Frangula</i>	Europe
<i>Rhamnus Cathartica</i>	Fruit	<i>Rhamnus cathartica</i>	Asia and Africa
UNOFFICIAL DRUG			
Ceanothus	Root	<i>Ceanothus americanus</i>	North America
Zizyphus	Fruit	<i>Zizyphus sativa</i>	Mediterranean region

VITACEÆ OR GRAPE FAMILY.—Rarely tall, herbaceous, usually shrubby and climbing, more rarely shrubby, upright

FIG. 422.—Northern Fox Grape (*Vitis Labrusca*). 1, portion of twig, with leaf, inflorescence and tendril, about natural size; 2, median longitudinal section of a perfect flower, enlarged; 3, perfect flower (enlarged) with petals removed; 4, longitudinal section of fruit (berry); 5, seeds, natural size and enlarged. (1-4 after Gray; 5, after Engelmann, from C. E. Bessey, *Nebraska Hort. Soc. Ann. Rept.* 1895.) (Gager.)

plants. Stems rarely short, more usually elongate, feeble, rather brittle, climbing by tendrils which represent modified inflorescence shoots. Leaves alternate, simple to lobed (either pinnately or more often palmately) to compound-pinnate or palmate. Perfect graded series of lanceolate leaves with pinnate veining to palmate veining; from pinnately veined to compound-pinnate; from palmately veined to compound-palmate. Stipules greenish to membranous or none. Flowers in racemes of compressed cymes, hermaphrodite or diclinous, nearly always small, clustered, green to greenish-yellow or greenish-white, rarely otherwise; sepals five, rarely four, small to minute (mere rim of receptacle) more or less persistent. Petals five, deciduous to caducous, typically distinct (in *Vitis*), united by their tips into calypetroform corolla, so in June, as Grape Vine flowers expand, corolla splits at base into five lobes that separate below, being attached at tips, while whole becomes tumbled off by lengthening stamens. Pistil bicarpellate. Ovary two-celled, superior or at most sub-inferior. Ovules two to one, erect. Style short, often more or less thickened, with terminal, capitate, slightly two-lobed stigmas. Stamens equal to petals or sepals and opposite petals. Receptacle internal to stamens, often expanded into nectariferous girdle or, in *Vitis*, into receptacular knobs alternating with stamens. Fruit a berry rarely six- to three-celled, typically two-celled and with two to one seeds in each cavity. Seeds like ovules, erect with bony testa. Embryo small, imbedded at base of cartilaginous albumen.

Brandy or Spiritus Vini *Vitis* is an alcoholic liquid obtained by the distillation of the fermented juice of sound, ripe grapes and containing from 48 to 54 per cent. of ethyl alcohol.

ORDER MALVALES

STERCULIACEÆ OR COLA FAMILY.—Rarely herbs, usually shrubs or tall, often heavy trees with soft wood and broad annual rings. The cambium, in developing bast, produces one, two, three, four, or five alternating layers of hard and soft bast which in some species of this as well as the *Tiliaceæ* or Linden family form long finger-like processes pushing out into the cortex. Leaves alternate, sometimes simple and pinnately-veined or passing to palmately-veined or palmately-compound. Flowers

hermaphrodite; sepals five, sometimes surrounded by bracteoles forming an epicalyx; petals usually five; stamens typically five, hypogynous, opposite petals, distinct or slightly fused in mon-

FIG. 423.—*Theobroma Cacao*, the Chocolate plant. 1. Trunk and branches bearing fruit. 2. Branch bearing flowers. 3. Single flower. 4. Staminal ring. 5. Stamen. (After Strasburger, Noll, Schenck and Schimper.)

adelphous fashion (*Melochia*, *Waltheria*) or, stamens subdivided above into few or numerous staminal leaflets, anthers two-celled; pistil many- to ten- to five- or four-carpelled; carpels apocarpous or more usually partially or completely united. Fruit either

follicles, or fused to form a capsule of ten or more, frequently five dehiscent carpels or, carpels splitting asunder into cocci or, becoming a woody capsular nut (*Theobroma*) or, rarely the fruit may become succulent. Seeds globose or subglobose and often provided with wings, arils or similar appendages; embryo straight, large and surrounded by scanty albumen.

OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Oleum Theobromatis	Fixed oil	Theobroma Cacao	Tropical America
Cacao Præpara- tum	Prepared powder from roasted, cured kernels of seeds	Theobroma Cacao and other species	Tropical America
Kola	Cotyledons	Cola nitida and other species	Africa

TILIACEÆ OR LINDEN FAMILY.—Shrubs or trees, rarely herbs, having stellate hairs on both stems and leaves. Leaves alternate, pinnately- more rarely palmately-veined, stipulate. Inflorescence cymose. Flowers hermaphrodite, more rarely, by absorption, more or less diclinous; sepals and petals five each, more rarely four, sepals deciduous; stamens five opposite the petals or, as in *Sterculiaceæ*, five phalanges of stamens representing subdivided stamens (*Tilia*), pistil of ten to five or two syncarpous carpels; ovary superior. Fruit either a nut-like drupe or drupe, rarely baccate.

FIG. 424.—*Tilia americana*, The American Linden or Basswood. Leaf and flowering branch. The leafy, tongue-shaped bract and simple cyme springing from near the center of its midrib constitutes the drug "Linden Flowers." (Reproduced from U. S. Dept. Agric. Misc. Pub. 77.)

UNOFFICIAL DRUG	PART USED
<i>Tilia</i> (Linden flowers)	Inflorescence
BOTANICAL ORIGIN	HABITAT
<i>Tilia</i> species	United States and Europe

MALVACEÆ OR MALLOW FAMILY.—Herbs in temperate regions (as the Mallow, *Malva rotundifolia*, and Marshmallow,

Althæa officinalis, etc.), occasionally shrubs in temperate regions (*Hibiscus Syriacus* or Rose of Sharon, etc.), frequently shrubs or tall trees in the tropics. Stems, as in *Sterculiaceæ* and *Tiliaceæ*, sometimes forming numerous layers of hard and soft bast. Leaves alternate and stipulate, ovate, ovate-cordate, orbicular or palmately-compound; venation pinnate or palmate. Stems, roots and leaves contain mucilage cells. Inflorescence a raceme

FIG. 425.—Median lengthwise section through flower of American Upland Cotton (*Gossypium hirsutum*) $\times 2$. (After Robbins.)

or fascicle of cymes. Flowers regular, pentamerous; calyx green, of five aposepalous sepals but frequently surrounded outside by an epicalyx. Both calyx and epicalyx are persistent. Corolla of five petals, varying in color, which are more or less fused with stamens at their bases, stamens monadelphous and forming an upright column (*staminal column*) enclosing the styles; anthers one-celled, dehiscing transversely; pollen grains echinate; pistil loosely or strongly syncarpous, rarely sub-apocarpous of thirty to five carpels. Fruit either a circular set of cocci (*Malva*),

FIG. 426.—The Asiatic- or Levant Upland Cotton, *Gossypium herbaceum*—Branch.
(Sayre.)

follicles, or a capsule (*Gossypium* or Cotton). Seeds albuminous with oily and mucilaginous albumen.

OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Althæa	Root (peeled)	Althæa officinalis	Europe and Asia
		(Cultivated varieties of:	
		Gossypium herbaceum	Arabia, United
Gossypii Radicis	Bark of root	Gossypium	States, East Indies
Cortex	(recently gathered)	Barbadense	West Indies, Egypt, and India
		Gossypium hirsutum	
Gossypium Purificatum	Hairs of seed	Cultivated varieties of Gossypium herbaceum	Arabia, East Indies, United States, and India

OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Oleum Gossypii Seminis	Fixed oil from Seeds	{ Cultivated varieties of G. herbaceum and other Gossypium species	{ United States, Asia, Africa and W. Indies
UNOFFICIAL DRUG			
Malvæ Folia	Leaves	{ Malva sylvestris Malva rotundifolia	{ Europe
Althæa Folia	Leaves	Althæa officinalis	Europe and Asia
Althæa Flores (larger mallow flowers)	Flowers (dark colored)	Althæa rosea	Europe
Malvæ Flores (mallow flowers)	Flowers	{ Malva sylvestris Malva rotundifolia	{ Europe
Ambrette Seed	Seed	Hibiscus Abelmoschus	India

FIG. 427.—Upland cotton (*Gossypium hirsutum*). A, mature boll opened out; B, cross-section of young boll; C, single seed with hairs; D, young boll. (Robbins.)

ORDER PARIETALES

This order includes a large variety of plants most of which have parietal placentation, separate sepals and numerous stamens. Belonging to this order are the Violets, Begonias, St. John's Worts, Damiana, Passion Flowers, Rock Roses, Tea and Camellias, Canella, etc.

THEACEÆ OR TEA FAMILY (TERNSTRÆMIACEÆ, CAMMELIACEÆ).—Evergreen shrubs or low branching, or tall, often heavy trees with watery juice. Leaves for the most part alternate, evergreen, often leathery, sometimes membranous; stipules either bud scales and caducous or often absent; leaf margins sinuate or serrate (*Thea*). Inflorescence a raceme becoming by condensation terminal, one-flowered. Flowers regular, perfect, pentamerous; sepals five, rarely four to three, deciduous, occasionally subtended by bracteolar scales; petals five, brittle and succulent, varying from greenish-white or greenish-yellow through yellow to white or, whitish pink to pink, scarlet, crimson, very rarely a tendency toward purple; stamens typically five but, as they grow, they subdivide into staminal leaflets, so that in their mature condition they are apparently indefinite and mono- to polyadelphous; stamens inserted hypogynously or perigynously and opposite the petals; pistil of typically five syncarpous carpels but reduced in some species to four to three or two. Fruit usually a capsule (*Thea*), five- to three-celled, dehiscent longitudinally, more rarely a fleshy, semi-baccate, semi-drupaceous, indehiscent fruit. Seeds with scanty or no albumen and often attached to the inner angle of cells by projecting spongy placentæ.

OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Caffeina	Feebly basic substance	<i>Thea sinensis</i> varieties, etc.	Eastern Asia
UNOFFICIAL DRUG			
<i>Thea</i> (Tea)	Leaves	<i>Thea sinensis</i> var. <i>Bohea</i> and <i>viridis</i>	Eastern Asia

GUTTIFERÆ OR GAMBOGE FAMILY.—Tropical trees (Gamboge and Mangosteen), rarely shrubs, containing resinous principles in schizogenous resin canals found in cortex and pith. Leaves opposite, coriaceous. Flowers dioecious, generally pentamerous or tetramerous with usually five stamens which are subdivided. Fruit a berry (*Garcinia Hanburyi*), drupe or capsule. Seeds generally large; embryo large to huge, often with enlarged radicle and reduced or absorbed cotyledons.

OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Cambogia	Gum resin	<i>Garcinia Hanburyi</i>	{ Siam, Cochin China and Annam

HYPERICACEÆ OR ST. JOHN'S WORT FAMILY.—Herbs or shrubs of temperate climes with opposite (*Hypericum perforatum*), rarely whorled branches and balsamic, resinous juices, which, in the herbaceous species, are secreted by black or pellucid glands found in the leaf parenchyma. Leaves entire, opposite, usually

FIG. 428.—*Garcinia Hanburyi*—Branch of Gamboge plant. (Sayre.)

sessile, exstipulate, and dotted. Flowers, regular, hypogynous and arranged in panicles or forked cymes. Petals usually 5. Stamens usually indefinite, rarely definite, often in 3–5 sets, more rarely monadelphous or free. Pistil of usually 3–5 carpels with 3–5 celled, compound ovary and as many filiform styles as carpels. Fruit a capsule with usually septicidal dehiscence

(*Hypericum*) or a berry. Seeds small, numerous, anatropous and exalbuminous.

UNOFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Hypericum	Entire plant	<i>Hypericum perforatum</i>	Europe

CANELLACEÆ FAMILY.—Trees the bark of which contains aromatic principles. Leaves alternate, pellucid-punctate. Flowers regular, golden-yellow, and arranged in terminal or axillary cymes. Fruit a berry containing two to many seeds with oily and fleshy albumen.

OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Canella	Inner bark	<i>Canella Winterana</i>	{ Florida and West Indies

BIXACEÆ FAMILY.—Tropical shrubs or trees. Leaves alternate, simple with minute or no stipules. Flowers hermaphrodite or unisexual, regular; stamens hypogynous, mostly indefinite with anthers opening by slits, rarely by one or two apical pores (*Bixa*). Fruit fleshy or dry. Seeds with fleshy albumen and sometimes covered with a fleshy arilus (*Bixa Orellana*).

UNOFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Annatto	Coloring matter	<i>Bixa Orellana</i>	{ Tropical America and Madagascar

FLACOURTIACEÆ OR FLACOURTIA FAMILY.—A family of chiefly tropical trees and shrubs with simple, alternate leaves, staminate, pistillate and hermaphrodite flowers, frequently dioecious or polygamous, with a one-celled ovary. The fruits are either berries (*Taraktogenous Kurzii* or Leprosy plant), drupes or rarely capsules. The seeds are albuminous.

OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Oleum	Fixed oil from seeds	<i>Taraktogenos Kurzii</i>	Burma
Chaulmoogræ		<i>Hydnocarpus</i>	India
		<i>Wightiana</i>	
		<i>Hydnocarpus anthelmintica</i>	Cochin China

VIOLACEÆ OR VIOLET FAMILY.—Herbs or shrubs. Stems upright, rarely creeping, spreading or acaulescent. Leaves either cauline or radical, stipulate, alternate, simple to pinnatifid or palmate.

Flowers pentamerous, regular or irregular (*Viola*) and sometimes cleistogamous; sepals 5; petals 5, unequal, the lowermost often spurred at the base; stamens 5, hypogynous, with adnate, introrse anthers and usually closely arranged or coherent around the ovary, in *Violets* the two lower stamens bearing spurs which extend into the spur of the corolla; ovary, sessile, one-celled with 3 to 5 parietal placentas; style usually club-shaped. Fruit a loculicidally dehiscent capsule (*Viola*), rarely baccate. Seeds albuminous.

UNOFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Pansy	Entire herb	<i>Viola tricolor</i>	Temperate regions
<i>Viola Odorata</i>	Fresh flowers and root	<i>Viola odorata</i>	Europe

TURNACEÆ OR DAMIANA FAMILY.—Tropical herbs, shrubs or trees. Leaves alternate, simple, petioled, exstipulate. Flowers perfect, regular, axillary, pentamerous with one-celled ovary. Fruit a capsule with three valves. Seeds strophiolate with albuminous embryo.

OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Damiana	Leaves	<i>Turnera diffusa</i>	California, Lower
		<i>Turnera</i>	California, Mexico, S.
		<i>aphrodisiaca</i>	America, West Indies

PASSIFLORACEÆ OR PASSION FLOWER FAMILY.—Herbaceous or woody vines climbing by tendrils. Leaves alternate, simple, entire, lobed or compound. Flowers perfect or imperfect, solitary; peduncles jointed at the flower; perianth petaloid with urceolate or tubular tube and four to five or eight to ten partite and two-seriate limb, the throat usually crowned by one or more series of subulate filaments which are frequently colored; gynophore elongating, supporting the stamens and pistil. Fruit a one-celled berry (*Passiflora*) or a three- to five-valved, dehiscent capsule containing numerous seeds.

UNOFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
<i>Passiflora</i>	Flowering and fruiting top	<i>Passiflora incarnata</i>	United States

CARICACEÆ OR PAPAW FAMILY.—A family of latex-containing trees composed of two genera indigenous to tropical America. Of chief pharmaceutic interest is the species *Carica Papaya*, the

Papaw or Melon tree, the fruit of which yields Papain, a valuable digestive ferment. This plant is a tree about 20 feet high which bears at its summit a cluster of deeply lobed, petiolate leaves and dioecious flowers. The fruit is a berry, the size of one's head and contains an acrid milky juice from which papain can be precipitated by the addition of alcohol.

CISTACEÆ OR ROCK ROSE FAMILY.—Herbs or shrubs whose stem and branches are often glandular, pubescent or tomentose, with simple or stellate trichomes. Leaves simple, entire, the lower ones opposite, upper alternate. Flowers perfect, regular, terminal, and solitary or in cymes or unilateral racemes; sepals five, the two external ones often bractiform or wanting; petals five, as in *Helianthemum* or Rock Rose and *Hudsonia*, rarely three or none, as in *Lechea* or Pinweeds; stamens hypogynous, indefinite; carpels three to five, ovary free, one-celled. Fruit a one-celled, three- to five-valved capsule.

FIG. 429.—Passion Flower or May-pop (*Passiflora incarnata*), a climbing, perennial herb used as a motor depressant in neuralgia, etc. Note the palmately 3-cleft leaves and axillary tendrils. The flowers possess a lavender or pink and purple corona.

UNOFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
<i>Helianthemum</i>	Herb	<i>Helianthemum canadense</i>	Eastern United States

ORDER OPUNTIALES.—CACTACEÆ OR CACTUS FAMILY.—Herbaceous, rarely arborescent (*Cereus giganteus* or Giant Cactus), more or less succulent plants living in warm, dry (*Pereskia*), usually desert situations, rarely becoming epiphytic and correspondingly modified. Stems, accordingly, varying from elongate, slightly enlarged, green (*Pereskia*), to flattened and ribbed (*Selenicereus*), to flattened and shortened as in the Prickly Pear (*Opuntia*), to condensed (*Echinocactus*, *Echinocereus*,

etc.), to greatly condensed (*Mamillaria*). Leaves alternate, stipulate or exstipulate, enlarged and more or less fleshy (*Pereskia*), becoming reduced, green and semicircular (*Opuntia*), or modified into spines, or wholly absorbed. Flowers regular, solitary, or fascicled in axils of leaves; sepals five; petals similar to sepals, petaloid, small to much enlarged, in color varying from yellow to white or from yellow to yellowish-pink, pink, scarlet or crimson; stamens indefinite, inserted at varying levels in the throat

FIG. 430.—The peyote cactus, *Lophophora Williamsii*, a perennial herb whose dried tops yield the drug known as peyote or mescal buttons. This drug has been used by the Aztec and Kiowa Indians as an intoxicant in their religious ceremonies. *A*, side view; *B*, view from above; *t*, *t*², tubercle; *r* rib; *t'*, scar of tubercle of older growth. $\times \frac{1}{2}$.

of a greatly expanded upgrown receptacle; pistil generally tricarpeal; ovary inferior, often deeply sunk in upgrown receptacular part; style thread-like, divided above into as many stigmas as carpels. Fruit a receptacular berry enclosing numerous small seeds. Seeds exalbuminous.

OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Cactus	Fresh succulent	{ <i>Selenicereus grandiflorus</i> }	Jamaica
Grandiflorus	stem		
UNOFFICIAL DRUG			
Peyote	Crown of plant	<i>Lophophora Williamsii</i>	Mexico

ORDER MYRTALES OR MYRTIFLORÆ

This order includes trees, shrubs and herbs, most of which possess tetramerous flowers with an inferior ovary. It contains

in addition to the families here considered the *Rhizophoraceæ* or Mangrove Family, the *Onagraceæ* or Evening-Primrose Family which is represented by the *Oenotheras* or Evening Primroses, the Willow Herbs (*Epilobium*) and the cultivated Fuchsia, the *Elæagnaceæ*, represented by the Buffalo-Berry, the *Trapaceæ*, con-

FIG. 431.—*Daphne Mezereum* (Fam. *Thymelæaceæ*)—Fruiting branch and flowers. (Sayre.)

taining the Water-Chestnut or *Trapa natans* and the *Lecythidaceæ* or Monkey-Pot Family which contains the Brazil-Nut whose seeds are edible.

THYMELÆACEÆ OR MEZEREUM FAMILY.—Shrubs (*Daphne Mezereum*) or low trees, usually of branching habit, the stems developing long tenacious bast fibers. Leaves alternate, rarely

opposite, exstipulate, coriaceous, simple, varying from lanceolate to ovate. Inflorescence a condensed raceme or spike. Flowers perfect, polygamous or diœcious, small, with calyx alone of the perianth parts developed. This is crimson-purple in *Daphne Mezereum*. Sepals usually fused to form a tube or cup-shaped perianth. Stamens usually eight in two rows of four longer and four shorter (*Daphne Mezereum*) inserted on the calyx tube. Pistil monocarpellary; ovary superior, mostly one-celled with a single pendulous ovule. Fruit a nut, drupe or berry (*Daphne*).

OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Mezereum	Bark	<div style="display: inline-block; vertical-align: middle;"> <div style="display: inline-block; vertical-align: middle;">{</div> <div style="display: inline-block; vertical-align: middle;"> <i>Daphne Mezereum</i> <i>Daphne Gnidium</i> <i>Daphne Laureola</i> </div> </div>	Europe and Asia

LYTHRACEÆ OR LOOSESTRIFE FAMILY.—Herbs, as Swamp Loosestrife (*Decodon*), Cigar plant (*Cuphea*) etc., shrubs (*Lawsonia*) or low trees containing astringent principles. Leaves either alternate, opposite or whorled, simple, exstipulate, usually lanceolate to ovate, entire, often glandular, viscous and containing idioblasts. Stems and leaves possess bicollateral bundles. Inflorescence a raceme, spike, or condensed cyme. Flowers perfect, often dimorphic or trimorphic, as in the spiked or purple loosestrife (*Lythrum Salicaria*), usually regular, but pass more or less to irregular, sometimes very irregular as in genus *Cuphea*; Calyx tubular, 3 to many toothed, the tube ribbed; sepals seven to four, petals equal in number and alternate with the inner teeth of the calyx, inserted on the mouth of the calyx tube, rarely absent; stamens fifteen, ten, or five in alternate rows of five each, inserted hypogynously or perigynously; pistil six-, five-, four-, two-, to one-carpeled with as many cavities in the ovary and numerous small ovules; style elongate with knobbed or 2-lobed stigma. Fruit a capsule. Seeds anatropous, exalbuminous. Calcium oxalate occurs in rosette aggregates and monoclinic prisms.

UNOFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Lawsonia (Henna)	Leaves	<i>Lawsonia inermis</i>	N. Africa to E. India

GRANATACEÆ (PUNICACEÆ) OR POMEGRANATE FAMILY.—A small family of shrubs and trees all of which are included in the

single genus *Punica*. Leaves opposite to whorled, entire, glabrous and extipulate. Flowers hermaphrodite, scarlet in the Pomegranate (*Punica Granatum*) solitary or aggregate, terminal, calyx petaloid; corolla of 7 to 5 petals inserted to calyx throat; stamens numerous; in many series, inserted below the petals; ovary inferior and imbedded in upgrown receptacle, its upper portion 7 to 5 celled with parietal placentation, its lower portion 3 celled with central placentation; style filiform; stigma capitate; ovules

FIG. 432.—The Pomegranate, *Punica Granatum*—Branch with flowers. (Sayre.)

numerous, anatropous. Fruit a spherical, baccate capsule crowned by the persistent calyx, its locules separated by membranous septa. Seeds numerous, exalbuminous.

OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Pelletierinæ Tannas	Mixture of tannates of the alkaloids	<i>Punica Granatum</i>	India
UNOFFICIAL DRUG			
Granati Fructus	Rind of fruit	{ <i>Punica Granatum</i>	India
Cortex Granatum	Bark of stem and root		

MYRTACEÆ OR MYRTLE FAMILY.—Rarely herbs (*Careya*) mostly shrubs or trees, some being the tallest trees known (*Eucalyptus*). Stems often tend to develop cork in flakes which separate

much as in the Buttonwoods. Leaves rarely alternate, nearly always opposite, entire, often glistening, subcoriaceous to coria-

FIG. 433.— The Blue Gum Tree, *Eucalyptus Globulus*,—Branch. (Sayre.)

ceous (*Eucalyptus*, *Pimenta*, etc.), frequently edge-on in position upon branches. Inflorescence cymose, at times forming scor-

poid cymes, becoming condensed into small fascicles, or each cyme condensing into a solitary flower.

Flowers regular or very rarely irregular from the lop-sided development of the stamens. Symmetry rarely hexamerous, typically pentamerous, not infrequently reduced to tetramerous

FIG. 434.—*Caryophyllus aromaticus*. (Sayre.)

(*Clove*); sepals five, six or four, aposepalous, or synsepalous at base, superior, and inserted around the edge of an expanded, upgrown, receptacular disc, varying from green and more or less expanded to short, thick, fleshy (*Clove*) or reduced to teeth (*Eucalyptus*); petals equal in number to the sepals, more or less petaloid and enlarged, rarely reduced and wanting, varying in color from green through greenish-yellow to white (*Caryophyllus*) or from

whitish to pink, scarlet, crimson, purple and blue, petals sometimes synpetalous and cup-like, detaching as the flower opens; stamens usually indefinite and epigynous, varying in the color of their filaments as do the petals; pistil rarely of ten to six carpels usually of five, not infrequently, as in Clove, of four carpels; ovary inferior or semi-inferior, as many-celled as there are carpels and with central placentation; style elongate; stigma undivided. Fruit either a hard, woody indehiscent nut (Brazil Nut), a capsule dehiscing at apex (*Eucalyptus*) or berry (*Caryophyllus*). Seeds exalbuminous.

OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Oleum Eucalypti	Volatile oil	Eucalyptus Globulus and some other Eucalyptus sp.	Australia and Tasmania
Eucalyptol	Organic oxide	Eucalyptus Globulus	
Caryophyllus	Flower buds	Caryophyllus aromaticus	Molucca Islands
Oleum Caryophylli	Volatile oil	Caryophyllus aromaticus	
Eugenol	Aromatic phenol	Caryophyllus aromaticus	
Oleum Pimentæ	Vol. oil	Pimenta officinalis	West Indies Central America Mexico
Oleum Myrciæ	Volatile oil	Pimenta acris	West Indies
UNOFFICIAL DRUG			
Bay Leaves	Leaves	Pimenta acris	West Indies
Eucalypti Gummi	Dried gummy exudate	Eucalyptus rostrata and other sp. of Eucalyptus	Australia
Eucalyptus	Sythe-shaped leaf	Eucalyptus Globulus	Australia
Oleum Cajuputi	Volatile oil from leaves and twigs	{ Var's. of Melaleuca Leucadendron	East Indies
Pimenta	Fruit	Pimenta officinalis	West Indies

COMBRETACEÆ OR MYROBALANS FAMILY.—Mostly tropical shrubs and trees containing considerable tannin. Leaves exstipulate, alternate or opposite, simple, pinnately veined, entire or toothed. Inflorescence a raceme, spike or head. Flowers regular, perfect or imperfect. Fruit a drupe, frequently longitudinally winged, containing a single seed.

UNOFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Combretum	Leaves	Combretum sundaicum	Sumatra

ORDER UMBELLALES OR UMBELLIFLORÆ

This order includes the Ginseng Family (*Araliaceæ*) represented by Ginseng, Spikenard, and Virginian Sarsaparilla, the *Umbelliferae* or Parsley Family which contains Fennel, Anise, Coriander, Caraway, Poison Hemlock, Water Hemlock and other medicinal plants, and the Dogwood Family (*Cornaceæ*) which includes the Dogwoods and Sour Gum. All of these families agree in usually possessing inflorescences consisting of umbels and flowers with an inferior ovary and a single whorl of stamens.

ARALIACEÆ OR GINSENG FAMILY.—Herbs, as Ginseng (*Panax quinquefolium*), English Ivy (*Hedera Helix*), Wild or Virginian Sarsaparilla (*Aralia nudicaulis*), etc., undershrubs (*Aralia hispida*, etc.), shrubs, as Bristly Sarsaparilla (*Fatsia horrida*), or trees, as Hercules Club (*Aralia spinosa*), with stems which are more or less hollow along internodes and solid at nodes. Leaves alternate, varying from simple to trifoliate or to multipinnate (tropical Aralias) or passing by telescoping into compound-palmate. Leaves serrate-margined and along with stem they develop volatile oil, resin and gum contents in secretion reservoirs. Inflorescence varying from a raceme of umbels to a raceme and even to condensed racemose umbels. Flowers regular, generally pentamerous, small, generally inconspicuous, green, greenish-yellow to rarely white, usually hermaphrodite but sometimes polygamous or dioecious; sepals five, rarely four; petals five, rarely four, often greenish to greenish-yellow, occasionally white, seldom pink in color; stamens varying from indefinite to ten to commonly five, opposite sepals, and, like sepals, epigynous in insertion; anthers versatile; pistil occasionally fifteen- to ten-, usually five-carpellate; ovary as many celled with one or rarely two pendulous ovules in each cavity; styles distinct ending in knob-shaped stigmas. Fruit a berry. Seeds albuminous, with a single seed coat.

OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Aralia (Spikenard)	Rhizome and roots	<i>Aralia racemosa</i>	Eastern United States and Canada

UNOFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Aralia nudicaulis	Rhizome	Aralia nudicaulis	Eastern United States and Canada
Aralia spinosa	Bark	Aralia spinosa	Eastern United States
American Ginseng	Root	Panax quinquefolium	North America
Chinese Ginseng	Root	Panax Ginseng	E. Asia
Panax Repens	Rhizome	Panax repens	Japan

UMBELLIFERÆ OR PARSLEY FAMILY.—Herbs (Anise, Fennel, Sweet Cicely, Poison Hemlock, etc.), rarely shrubs, often of rapid growth, and with upright, fistular (hollow at internodes, solid at nodes), often grooved and ridged stems. Leaves alternate, compound and usually much divided, exstipulate, but with expanded sheathing and flattened leaf base (*pericladium*), that ensheathes the stem. Inflorescence a simple or often compound umbel surrounded by an involucre of bracts or of bracteoles. Flowers small, pentamerous, with inferior ovary and superior floral parts. Sepals minute, tooth-like, inserted above inferior ovary, or absorbed. Petals small, usually yellow to white, rarely pink to purple, distinct, each with inflexed tip. Stamens five, epigynous, inserted below a nectariferous, epigynous disc, incurved in bud. Carpels two, fused into bicarpellate pistil. Ovary two-celled, with one pendulous ovule in each cell, ovarian wall traversed by oleoresin canals; styles two, distinct above the nectar disc or stylopod. Fruit a dry, splitting fruit or cremocarp, that splits lengthwise into two mericarps which hang for a time by a forked carpophore. Seeds single in each mericarp, albuminous, each with a single seed coat.

OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Anisum	Ripe fruit	Pimpinella Anisum	Asia Minor, Egypt and Greece
Oleum Anisi	Volatile oil	Pimpinella Anisum	
Anethol	{ Methyl ether of para-pro- penyl phenol	Pimpinella Anisum	
		Foeniculum vulgare	
Foeniculum	Ripe Fruit	Foeniculum vulgare	Mediterranean region
Oleum Fœniculi	Volatile oil	Foeniculum vulgare	
Sumbul	{ Rhizome and roots	Ferula Sumbul or other closely related spp. of Ferula having a musk-like odor	Asia

OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Carum	Fruit	Carum Carvi	Europe, Asia
Oleum Cari	Vol. oil	Carum Carvi	
Asafoetida	Gum resin	Ferula foetida, F. Assa-foetida, F. rubricaulis, etc.	Persia and Afghanistan
Coriandrum	Ripe fruit	Coriandrum sativum	{ Mediterranean and Caucasus regions
Oleum Coriandri	Volatile oil	Coriandrum sativum	
Apii Fructus	Ripe fruit	Apium graveolens	Southern Europe
UNOFFICIAL DRUG			
Angelicae Fructus	Ripe Fruit	Angelica Archangel- ica and other species of Angelica	Northern Europe and Siberia
Angelicae Radix	Rhizome and roots	Angelica Arch- angelica and other species of Angelica	Northern Europe and Siberia
Conium	Unripe fruit	Conium maculatum	Europe
Ammoniacum	Gum-resin	Dorema Ammoniacum	Persia
Galbanum	Gum-resin	Ferula galbaniflua	Persia and Afghanistan
Imperatoria (Masterwort)	Root	Imperatoria Ostruthium	Europe
Levisticum (Lovage)	Root	Levisticum officinale	Europe
Pimpinella	Roots	{ Pimpinella Saxifraga } { Pimpinella magna }	{ Central Europe
Petroselinum	Ripe fruit	{ Petroselinum sativum }	
Petroselini Radix	Root		
Apiol	Oleoresin from root		

CORNACEÆ OR DOGWOOD FAMILY.—Herbs, as Bunchberry (*Cornus canadensis*), etc., shrubs, as Blood-Twig Dogwood (*Cornus sanguinea*), etc. or trees, as Flowering Dogwood (*Cornus florida*) and Sour Gum (*Nyssa sylvatica*), etc. Leaves simple, alternate (Sour Gum), or opposite (Dogwoods). Inflorescence an umbel or determinate head (*glomerule*), the whole being surrounded by an enlarged and more or less petaloid involucre. Flowers regular, rarely pentamerous, more frequently tetramerous; sepals usually four, small, tooth-like or absorbed; petals usually four, small, greenish to yellowish to white (*Cornus florida*), rarely pink or crimson; stamens four or five, alternate to the petals and inserted with the sepals and petals epigynously around and

between the nectar disc; pistil syncarpous, bicarpellate, rarely tricarpetate; ovary as many celled with one pendulous ovule in

FIG. 435.—*Conium maculatum* L., the poison hemlock. Note spotted stems, decompound leaves and compound umbels of white flowers. The poisonous juice of this plant was employed as a potion by the ancient Greeks which was given to criminals as a means of capital punishment.

each cavity; style usually simple, ending in rounded or slightly bilobed stigma. Fruit a two-seeded drupe. Seeds albuminous with single seed coat.

UNOFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Cornus	Bark of root	Cornus florida	Eastern United States and Canada

SUB-CLASS—SYMPETALÆ (GAMOPETALÆ OR METACHLAMYDEÆ)

A division of dicotyledonous plants in which the flowers possess both calyx and corolla, the latter with petals more or less united into one piece.

ORDER ERICALES

ERICACEÆ OR HEATH FAMILY.—Sub-herbaceous, as Pipsissewa (*Chimaphila*), suffruticose, as the Heather (*Calluna*),

FIG. 436.

FIG. 437.

FIG. 436.—Wintergreen (*Gaultheria procumbens*), an aromatic, evergreen, perennial with ovate leaves, nodding, white, urn-shaped, axillary flowers and red, berry-like fruits, each of the latter representing a 5-celled capsule enveloped by a fleshy calyx. Native to sandy soils in cool, damp woods of Eastern North America. (Reproduced from U. S. Dept. Agric. Miscellaneous Publication 77.)

FIG. 437.—Pipsissewa (*Chimaphila umbellata*) on left. The two plants on right are *Chimaphila maculata*, the spotted pipsissewa or "spotted wintergreen." (Reproduced from U. S. Dept. Agric. Misc. Publ. 77.)

fruticose (Azaleas, Mountain Laurel, etc.), rarely sub-arborescent (*Arbutus unedo* or Strawberry Tree) plants. Roots fibrous, often saprophytically associated, rarely tuberous or more or less enlarged. Stem upright, ascending or creeping, more or less

woody, rarely through a saprophytic or parasitic connection becoming soft, annual and white above ground, as in Indian Pipe (*Monotropa uniflora*). Leaves alternate, simple, entire, exstipulate, rarely soft, delicate, herbaceous (*Azaleas*), usually leathery to wiry and evergreen, more rarely (*Pterospora* (Pine Drops), *Monotropa*, etc.) becoming greenish-blue, bluish-yellow, yellowish-white to white and correspondingly saprophytic. Inflorescence typically a raceme (*Pyrola*, *Andromeda*, *Gaylussacia*, *Erica*, *Arctostaphylos Uva Ursi*, etc.) but raceme condensed into a racemose umbel (*Azalea*, etc.) or further reduced to a few flowers or, in the degraded saprophytic condition, to one flower (*Mono-*

FIG. 438.—Flower of a Blueberry (*Vaccinium*). A, median lengthwise section; B, external view. (Robbins.)

tropa uniflora). Flowers regular, passing to irregular (*Rhododendron*), pentamerous or tetramerous; sepals five to four, rarely fewer, apo- to synsepalous, usually green, sometimes brightly petaloid; petals five, more rarely four, slightly to deeply synpetalous, cup-shaped, as in Mountain Laurel (*Kalmia*) to urceolate (*Arctostaphylos*, *Andromeda*, etc.), yellow to white or through yellow-pink to scarlet to crimson to crimson-purple; stamens ten to eight in two circles of five to four each, becoming by absorption of inner circle, five to four only, hypogynous, epipetalous or epigynous; anthers two-celled, dehiscing by apical pores, as in Bearberry (*Arctostaphylos*) or apical slits; pollen sometimes agglutinated into long viscous threads; pistil five- to four-

rarely six- to eight-carpeled, superior, rarely semi-inferior to inferior, as in Blueberries (*Vaccineæ*); ovary as many celled as there are carpels; style elongated, filiform, usually five- to four-lobed. Fruit a capsule, as in *Epigæa repens* or *Trailing Arbutus*, a berry, as in Blueberries and Cranberries (*Vaccinium*) and

FIG. 439.—*Arctostaphylos Uva ursi*. 1. Twig bearing spatulate leaves and urn-shaped flowers. 2. Flower in longitudinal section. 3. Pollen tetrad. 4. Fruit. 5. Cross section of fruit. $\times 2-5$. (From Strasburger, Noll, Schenck and Schimper after Berg and Schmidt.)

Huckleberries (*Gaylussacia*) or false drupe, as in the Wintergreen (*Gaultheria procumbens*). Seeds small, anatropous.

OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Chimaphila	Leaf	Chimaphila umbellata	United States, Canada, Northern Europe and Asia
Uva Ursi	Leaf	Arctostaphylos Uva Ursi	
Methylis Salicylas	Volatile oil	Gaultheria procumbens (in part)	Northern United States and Canada, Europe and Asia
UNOFFICIAL DRUG			
Gaultheria	Leaf	Gaultheria procumbens	United States and Canada
Vaccinium	Leaf	{ Vaccinium corymbosum et al. spp. Vaccinium	United States
Neomyrtillin	Glucoside		

ORDER EBENALES

SAPOTACEÆ OR STAR APPLE FAMILY.—Tropical shrubs or trees (*Palaquium*) characterized by the presence of laticiferous sacs

in the pith and cortex of the stems and adjoining the veins of the leaves. Leaves alternate, exstipulate, evergreen and coriaceous.

FIG. 440.—The manzanita (*Arctostaphylos manzanita*), an ericaceous shrub growing in western United States and common on hillsides near the Pacific coast. (After Stanford, *General and Economic Botany*, D. Appleton-Century Co., publishers.)

Flowers perfect, large and axillary. Fruit a berry (*Palaquium*) rarely a capsule (*Ponteria*).

UNOFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Gutta Percha	Purified coagulated milky exudate	Various species of <i>Palaquium</i> and <i>Payena</i>	Malay Archipelago
Chicle	Dried latex	<i>Achras Sapota</i>	Mexico and Central America
Balata	Dried latex	<i>Mimusops globosa</i>	West Indies and S. America

STYRACEÆ OR BENZOIN FAMILY.—Shrubs or low trees. Leaves alternate to opposite, entire, often acuminate and exstipulate. Flowers perfect, regular, rarely sub-irregular, either in condensed fascicles or solitary in the axils of the leaves; sepals and petals typically five each; corolla often white, rarely pinkish or yellowish; stamens many to four to two, perigynous or sub-hypogynous; pistil bicarpellary or four to five carpellate. Fruit either drupe-like or dry, often winged and rarely as many-celled as there are carpels.

OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Benzoinum	Balsamic resin	Styrax Benzoin	East Indies
		Styrax tonkinensis	Siam
		or other species of Styrax	

ORDER GENTIANALES

OLEACEÆ OR OLIVE FAMILY.—Shrubs, as Lilacs (*Syringa*), Golden Bell (*Forsythia*) or trees (as the Ash (*Fraxinus*), Olive (*Olea*), etc.) with stems possessing close white wood, and slightly swollen or enlarged nodes. Leaves opposite, decussate, simple, rarely pinnately-compound (Ash). Inflorescence dichasial or scorpioid cymes but tending constantly toward condensation and, so in the Lilac, the inflorescence becomes a clustered raceme of cymes (thyrsus). Flowers regular, pentamerous or tetramerous; sepals small, green, rarely petaloid, synsepalous; petals synpetalous, elongated into a narrow tube, expanding above into a stellate limb; stamens very rarely five, rarely four to three, nearly always two, epipetalous and high set on corolla tube; pistil bicarpellate, rarely of three to four carpels; ovary two-celled with two to one pendulous ovules in each cavity. Fruit either a capsule (Lilac), drupe (Olive), berry (Privet) or a winged indehiscent akene (Ash). Seeds with moderate to scanty albumen, becoming occasionally exalbuminous.

OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Manna	Dried exudation	Fraxinus Ornus	Southern Europe
Oleum Olivæ	Fixed oil	Olea europæa	Southern Europe, Algeria, Asia
Chionanthus	Bark of root	Chionanthus virginica	Southern United States
UNOFFICIAL DRUG			
Fraxinus	Bark	Fraxinus americana	Northern United States and Canada

LOGANIACEÆ OR LOGANIA FAMILY.—Herbs (*Spigelia* or Pink Root, etc.), woody vines (*Gelsemium* or Yellow Jasmine, etc.) or trees (*Strychnos Nux Vomica*, etc.) with a bitter juice usually containing alkaloids. Stem, rarely herbaceous, usually woody, often long climbing and rope-like (*Gelsemium*), usually with a bicollateral bundle system. Leaves opposite, stipulate or exstipulate. Inflorescence racemose or cymose (*Spigelia*) (scor-

pioid cyme in *Strychnos*), sometimes condensed into solitary, axillary flowers. Flowers perfect, usually regular; calyx gamosepalous; corolla gamopetalous, hypogynous, rotate, campanulate

FIG. 441.—*Strychnos Nux-vomica*—Flowering branch and seeds. (Sayre.)

or infundibuliform; stamens inserted on the corolla tube or throat and with thread-like filaments; ovary superior, two-celled; style elongate with bifid stigma; ovules numerous. Fruit usually a capsule, septicidally dehiscent (*Gelsemium sempervirens*), or loculicidally dehiscent (*Spigelia marilandica*), sometimes a berry

FIG. 442.—Pinkroot (*Spigelia marilandica*) a perennial herb of the *Loganiaceæ*. Note the opposite leaves and one-sided spike of tubular flowers. The corollas of these are red outside and yellow within. The rhizome and roots are used as a worm remedy or anthelmintic. (Reproduced from U. S. Dept. Agric. Misc. Pub. 77.)

(*Strychnos Nux-Vomica*) or drupe. Seeds numerous or solitary, sometimes winged.

OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Nux Vomica	Seeds	<i>Strychnos Nux-Vomica</i>	India
Brucinæ Sulfas	Sulfate of alkaloid from seed		
Strychnina	Alkaloid from seed also sulfate, nitrate and phosphate of strychnine	{ <i>Strychnos Nux-Vomica</i> and other <i>Loganiaceæ</i>	{ Eastern Hemi- sphere
Gelsemium	Rhizome and roots		
		<i>Gelsemium sempervirens</i>	Southern United States
UNOFFICIAL DRUG			
Curare	Extract from bark	<i>Strychnos toxifera</i> et al. spp.	S. America
Ignatia	Seed	<i>Strychnos Ignatii</i>	Philippine Islands
Spigelia	Rhizome and roots	<i>Spigelia marilandica</i>	Southern United States

GENTIANACEÆ OR GENTIAN FAMILY.—Herbs, often low-growing. Roots and short stems sometimes more or less thickened (*Gentiana lutea*). Leaves opposite, decussate, entire, exstipulate. Inflorescence cymose (*Gentiana lutea*) or condensing to a single

solitary, terminal flower (*Gentiana verna*, *G. acaulis*, etc.). Flowers regular, perfect, pentamerous or tetramerous, sepals five to four, green, more or less synsepalous, not infrequently everted or reflexed, corolla of five, rarely four petals, more or less

FIG. 443.—The yellow gentian or bitter root, *Gentiana lutea*,—Flowering head and dissected flower. (Sayre.)

synpetalous, in shape passing from open-stellate, as in *Gentiana lutea*, through many stages of connation to long-tubed, as in *Gentiana acaulis*; stamens five, epipetalous; pistil bicarpellate; ovary one-celled or incompletely two-celled; style more or less elongated with bilobed to divided stigma. Fruit a capsule. Seeds albuminous. Plants contain bitter glucosides.

OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Gentiana	Rhizome and root	<i>Gentiana lutea</i>	Europe
UNOFFICIAL DRUG			
Chirata	Entire plant	<i>Swertia Chirayita</i>	Northern India
Centaurium	Flowering plant	<i>Erythræa Centaurium</i>	Europe
Menyanthes	Leaves	<i>Menyanthes trifoliata</i>	Europe and Asia
Sabbatia	Herb	<i>Sabbatia angularis</i>	Eastern United States and Canada

FIG. 444.

FIG. 444.—*Menyanthes trifoliata*, the buckbean or bogbean, whose leaves are employed as a simple bitter. This perennial herb occurs in bogs of Europe, Asia and North America. Note the long petioled, trifoliate leaves whose bases sheathe the rhizome and the long peduncle bearing a raceme of tubular flowers which are pinkish-purple or white. (Reproduced from U. S. Dept. Agric. Misc. Pub. 77.)

FIG. 445.

FIG. 445.—*Apocynum cannabinum* or Canadian hemp, a perennial herb whose rhizome and roots contain cardiac stimulant glucosides. Note the opposite, ovate-lanceolate leaves, the central terminal cyme of greenish white flowers and the fruit of paired follicles (on right). (Reproduced from U. S. Dept. Agric. Misc. Publ. 77.)

APOCYNACEÆ OR DOG BANE FAMILY.—Herbs, rarely shrubs, not infrequently clambering or climbing in habit (*Allamanda*). Stem and branches show bicollateral bundles. Stem, leaves and flowers have latex tubes which ramify through the cortex and mesophyll tissues. Leaves alternate, opposite or verticillate,

simple, entire, deciduous or evergreen. Inflorescence cymose. Flowers regular, pentamerous, rarely tetramerous; sepals five, gamosepalous, green, rarely subpetaloid to petaloid; petals five, slightly to deeply gamopetalous, in shape varying from open tubular, stellate, to elongate tubular to elongate funnel-shaped,

FIG. 446.—*Strophanthus hispidus*—Branch and seed with comose awn. (Sayre.)

in color varying from greenish-yellow to white or from yellow-red to crimson to crimson-purple to nearly purple-blue; stamens five, epipetalous; pistil usually bicarpellate; ovary two-celled with central placentation; style more or less elongate with terminal brush of hairs, knobbed or multifid; stigma a circular band or circular spur beneath terminal style swelling. Fruit two follicles (*Apocynum*, etc.), a berry, drupe, or capsule. Seeds flattened, frequently hairy, albuminous.

OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Strophanthus	Seed (deprived of awn)	{ Strophanthus Kombe Strophanthus hispidus }	{ Africa
Strophanthin	Glucoside	Strophanthus Kombé	Africa
Apocynum	Rhizome and roots	Apocynum cannabinum	United States and Canada
UNOFFICIAL DRUG			
Aspidosperma	Bark	Aspidosperma Quebracho blanco	Argentine Republic

ASCLEPIADACEÆ OR MILKWEED FAMILY.—Herbs or shrubs containing a milky juice, many species yielding rubber. Leaves entire, more or less fleshy, sometimes verticillate. Inflorescence usually a dichasial or scorpioid cyme. Flowers regular, pentamerous; sepals woolly, small, synsepalous; petals five, rarely four, synpetalous, elongated into awls; the corolla varying in shape from stellate to campanulate and in color from pale green to yellow, to greenish-brown, chocolate, or from white to yellow, to scarlet, to crimson, to purple, to blue; stamens five, epipetalous, fused in relation, forming a cylindrical swollen mass around the central pistil; filaments flattened and furnished with a crown having various appendages; anthers two-celled, each cell containing a pollen mass (*pollinium*), adhering to the glandular prominences of the stigma; pistil bicarpellate, superior. Fruit typically two dry follicles (*Asclepias*), rarely becoming succulent or bladdery. Seeds numerous, compressed, imbricate, with a comose appendage.

FIG. 447.—*Asclepias tuberosa*, the Pleurisy-root or Orange Milkweed. The medicinal root, to left, is orange brown and devoid of latex. Note the alternate leaves and terminal cymes of umbels, which bear orange-colored flowers. To right, stem bearing follicles two of which are discharging their seeds. (Reproduced from U. S. Dept. Agric. Miscellaneous Publication 77.)

DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Asclepias	Root	<i>Asclepias tuberosa</i>	United States
Condurango, N.F.	Bark	<i>Marsdenia</i> Condurango	Peru and Ecuador

ORDER TUBIFLORÆ OR POLEMONIALES

CONVOLVULACEÆ OR MORNING GLORY FAMILY.—Frequently herbaceous, more rarely sub-woody or woody, perennial climbing plants with underground parts sometimes swollen into tuberous roots (Jalap, Sweet Potato, Wild Man of the Earth). Stems rarely short, upright or tufted, usually elongate and cir-

FIG. 448.—*Convolvulus scammonia*—Branch. (Sayre.)

cummutating in action. Vascular bundles frequently bicollateral. Leaves alternate, simple, exstipulate, varying from cordate to cordate-sagittal, to broad-reniform to reniform, palmately lobed to palmatifid to palmately-compound (*Ipomæa* shows all these transitions). Stem and leaves frequently contain a dull, viscous, watery to milky-white, resinous juice. Inflorescence a scorpioid cyme, becoming reduced in some forms to

a solitary flower. Flowers pentamerous; sepals five, green, gamosepalous; corolla varying in shape from rotate to funnel-like with expanded mouth, in color from greenish-yellow to white or through yellowish-pink to scarlet, crimson, purple or blue; stamens five, often with the bases of the filaments expanded; pistil bicarpellate; ovary two celled, superior, often surrounded by a nectar girdle; style filiform with bilobed or bifid stigma. Fruit usually a capsule (*Exogonium*, *Convolvulus*, etc.), dehiscent septifragally, rarely a berry. Seeds scantily albuminous to exalbuminous.

OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Jalapa	Tuberous root	<i>Exogonium</i> Jalapa	Mexico
<i>Ipomœa</i>	Root	<i>Ipomœa orizabensis</i>	Mexico
UNOFFICIAL DRUG			
Tampico Jalap	Root	<i>Ipomœa simulans</i>	Mexico
Wild Jalap	Root	<i>Ipomœa pandurata</i>	United States
Turpeth Root	Root	<i>Operculina Turpethum</i>	East Indies
Scammonia Radix	Root	<i>Convolvulus</i> <i>Scammonia</i>	Asia Minor, Greece, Syria

HYDROPHYLLACEÆ OR WATER LEAF FAMILY.—Annual, herbaceous, rarely perennial woody plants whose stems, branches, leaves and sepals are often viscous and glandular hairy. Leaves alternate, exstipulate, from simple, linear to pinnatipartite to pinnate. Inflorescence rarely expanded, usually scorpioid cymes. Flowers small to large, funnel-form in Yerba Santa (*Eriodictyon californicum*); sepals five, green; petals five, regular; corolla varying from small stellate with slightly fused petals to large rotate, campanulate or tubular, in color varying from greenish-white or yellow to rarely white, often pink, purple or blue; stamens five, rarely with alternate staminodes; pistil bicarpellate. Fruit a two-celled capsule dehiscing usually septicidally.

OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
<i>Eriodictyon</i>	Leaves	<i>Eriodictyon californicum</i>	California and New Mexico

BORAGINACEÆ OR BORAGE FAMILY.—Herbaceous (*Boragineæ* sub-family) or shrubby (*Heliotropeæ* sub-family) plants, forming a primary root and a single or often branched shoots. Leaves often divisible into expanded, sometimes large basal and alter-

nate, scattered cauline leaves. Each of these simple, exstipulate, often hairy, rarely glabrous. Inflorescence a raceme of dichasial or scorpioid cymes, at times condensed into a dichasium of scorpioids or a simple scorpioid cyme. Flowers pentamerous, regular, passing to slight or marked irregularity as in Blue-weed (*Echium*); sepals five, green, slightly or deeply gamosepalous, often hairy; petals five, the corolla varying in shape from rotate

FIG. 449.—Yerba Santa (*Eriodictyon californicum*), an evergreen shrub found along the coastal ranges from central California to Oregon. The leaves are used in expectorant preparations. (Reproduced from U. S. Dept. Agric. Miscellaneous Publ. 77.)

FIG. 450.—*Symphytum officinale* or Comfrey, a hairy perennial herb with a mucilaginous root and purplish, blue, or pinkish-white tubular flowers borne in pendulous cymes. (Reproduced from U. S. Dept. Agric. Misc. Publ. 77.)

with shallow tube, as in Forget-me-not (*Myosotis*) and *Borage*, to tubular, as in Comfrey (*Symphytum*), to funnel-shaped in most species; in color, all transitions, frequently purple-blue to blue; stamens five; pistil bicarpellate, syncarpous, embryologically two-celled with two ovules in each cavity, but dorsal ingrowths divide ovary by time of flowering into four cells with one ovule in each cavity; style gynobasic. Fruit typically four-nutlets. Seeds solitary in each cavity and either scantily albuminous (*Heliotropææ*) or exalbuminous (*Boragineæ*).

UNOFFICIAL DRUG	PART USED *	BOTANICAL ORIGIN	HABITAT
Symphytum (Comfrey)	Root	Symphytum officinale	Europe and United States
Cynoglossum (Hound's tongue)	Herb and root	Cynoglossum officinale	United States
Alkanet	Root	Alkanna tinctoria	So. Europe and Asia
Lungwort	Leaf	Pulmonaria officinalis	Europe

VERBENACEÆ OR VERVAIN FAMILY.—Herbs, as the Vervains (*Verbena*), shrubs, as the Glory-bower (*Clerodendron*), rarely trees, as *Premna arborea*, one of the sources of the drug Tonga, and *Tectona* or Teakwood, whose stems and branches are usually quadrangular and rarely scented. Leaves generally opposite, exstipulate, simple or compound. Inflorescence a terminal panicle of spikes (*Verbena hastata*), a cyme (Beauty-berry or *Callicarpa*) or head (*Lippia lanceolata*). Flowers white, pink or blue (*Verbena hastata*), irregular, more or less 2-lipped; calyx gamosepalous, tubular; corolla gamopetalous, hypogynous with a 4-5 fid limb; stamens generally 4, didynamous and inserted on the corolla tube or throat; pistil of 2-4 carpels, a terminal style and undivided stigma. Fruit a drupe or 2 to 4 celled berry, usually splitting into as many nutlets. Seeds exalbuminous.

FIG. 451.—The Blue Vervain, *Verbena hastata*. Note the lanceolate leaves with serrate margins and the flowers which are usually blue, and arranged in panicles of narrow spikes. (Reproduced from U. S. Dept. Agric. Misc. Publ. 77.)

UNOFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Verbena	Overground portion	Verbena hastata	United States
Tonga	Bark	Premna arborea (in part)	Fiji Islands

LABIATÆ (LAMIACEÆ) OR MINT FAMILY.—Herbs producing creeping runners that spread out and root at the nodes. Stems

quadrangular, rarely cylindrical in outline. Leaves opposite, decussate, mainly petiolate; leaf margin nearly always serrate, dentate or crenate. Stems and leaves further characterized by the presence of glandular hairs containing aromatic volatile

FIG. 452.—Peppermint (*Mentha piperita*), a perennial herb naturalized from Europe and found in meadows of eastern and central North America. Note the opposite, petiolate, ovate-oblong leaves with serrate margins and the flowers which are purple and borne on blunt spikes of verticillasters terminating the square stems. (Reproduced from U. S. Dept. Agric. Misc. Publ. 77.)

oil. These hairs, for the greater part, consist of a short one-celled stalk and a head (gland) of six or eight cells. Inflorescence a raceme or spike of verticillasters (double dichasial cymes) or, as in the Ground Ivy, a reduced verticillaster. Flowers typically pentamerous, rarely tetramerous; sepals five, synsepalous, ribbed and forming a tubular, regular or irregular bilabiate (Scullcap, etc.) calyx whose upper lip is bifid and lower trifid; corolla of five to four gamopetalous petals, hypogynous, frequently two-lipped, the upper lip bifid, the lower trifid; stamens four, didynamous, rarely one pair alone fertile and the other pair reduced, in some cases almost or quite to disappearing point, as in the Sage (*Salvia*) and the Horsemint (*Monarda*); pistil bicarpellate, embryologically two-celled with two ovules in each cavity, becoming, at time of flowering, four-celled with one ovule in each cavity. Style embryologically terminal, but, upon opening of flower, deeply gynobasic, elongate, slender, with two stigmatic surfaces. Fruit four nutlets enclosing as many exalbuminous seeds.

OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
<i>Mentha Piperita</i> (Peppermint)	Leaves and tops	<i>Mentha piperita</i>	Europe
<i>Oleum Menthae Piperitæ</i>	Volatile oil from fresh aerial parts of flowering plant	<i>Mentha piperita</i>	Europe

OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Menthol	Alcohol separated from vol. oil	<i>Mentha arvensis</i> var. <i>piperascens</i>	Europe
<i>Mentha Viridis</i> (Spearmint)	Leaves and tops	<i>Mentha spicata</i>	Europe
<i>Oleum Menthæ Viridis</i>	Volatile oil from fresh aerial parts of flowering plants	<i>Mentha spicata</i>	Europe
<i>Scutellaria</i> (Scullycap)	Overground portion	<i>Scutellaria lateriflora</i>	United States and Canada
<i>Thymus</i> (Thyme)	Leaves and flowering tops	{ <i>Thymus vulgaris</i>	{ Southern Europe
<i>Oleum Thymi</i>	Volatile oil from flowering plant		
<i>Oleum Rosmarini</i>	Volatile oil from fresh flowering tops	<i>Rosmarinus officinalis</i>	Mediterranean Basin
<i>Oleum Lavandulæ</i>	Volatile oil from fresh flowering tops	<i>Lavandula Spica</i>	Southern Europe

FIG. 453.—Catnip (*Nepeta Cataria*). Note the cordate leaves with crenate margins and the spikes of whorled flowers which are white and spotted with purple. (Reproduced from U. S. Dept. Agric. Misc. Publ. 77.)

Cataria (Catnep)	Leaves and flowering tops	<i>Nepeta Cataria</i>	Europe and Asia
<i>Salvia</i> (Sage)	Leaves	<i>Salvia officinalis</i>	Southern Europe
UNOFFICIAL DRUG			
<i>Marrubium</i> (Horehound)	Leaves and flowering tops	<i>Marrubium vulgare</i>	Europe and Asia
<i>Hedeoma</i> (Pennyroyal)	Leaves and flowering tops	<i>Hedeoma pulegioides</i>	United States and Canada

UNOFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Herba Majoranæ (Sweet Marjoram)	Leaves and flowering tops	Origanum Majorana	Mediterranean regions
Collinsonia (Stone Root)	Rhizome and roots	Collinsonia canadense	United States
Serpyllum (Wild Thyme)	Leaves and flowering tops	Thymus Serpyllum	Europe and Asia
Melissa (Balm)	Leaves and flowering tops	Melissa officinalis	Southern Europe, Asia Minor
Monarda (Horsemint)	Leaves and flowering tops	Monarda punctata	United States
Origanum	Leaves and flowering tops	Origanum vulgare	Europe, Asia and North Africa
Hyssopus	Leaves and flowering tops	Hyssopus officinalis	Southern Europe
Summer Savory	Leaves	Satureja hortensis	Southern Europe
Mountain Mint	Leaves	Pycnanthemum montanum	United States
Sweet Basil	Leaves	Ocimum Basilicum	Asia and Africa
Motherwort	Leaves and flowering tops	Leonurus Cardiaca	Europe
Lavender	Flowers	Lavandula Spica	S. Europe
Rosmarinus	Leaves	Rosmarinus officinalis	} Mediterranean Basin
Greek Tea	Herb	Sideritis spp.	

SOLANACEÆ OR NIGHTSHADE FAMILY.—Herbs (potato, tomato, horsenettle), shrubs (*Lycium* spp.), vines (*Solanum dulcamara* or bittersweet), rarely trees as in some tropical *Daturas*. Stems and leaves possess bicollateral bundles. Many of the plants as belladonna, stramonium, hyoscyamus and scopola contain mydriatic alkaloids. Leaves alternate, exstipulate, entire or more or less lobed, rarely compound; often glandular-hairy. Flowers in cymes or solitary; regular or rarely irregular (*Petunia*, *Tobacco* spp.), pentamerous, perfect, synphyllous; sepals green (rarely petaloid), rotate to tubular, usually persistent and accrescent; petals rotate (*Solanum*), to tubular (*Atropa*), to funnel-shaped (*Tobacco*), and so (1) open to all comers, or (2) to bees or wasps, or (3) to butterflies, moths; color, greenish-yellow, or greenish-white, to white, to pink, crimson, purple, rarely blue; stamens five, epipetalous, hypogynous, along with style usually forming nectar glands. Filaments short to long, anthers dehiscing longitudinally or by apical pores; pistil bicarpellate, syncarpous, with or without

nectar girdle; superior ovary, two-celled with central placentation, ovules numerous, style more or less elongate with bilobed or bifid stigma. Fruit, a capsule dehiscing longitudinally

FIG. 454.—The Henbane (*Hyoscyamus niger* Linné.), a member of the Nightshade family, whose leaves, with or without the tops, constitute the official drug *Hyoscyamus* and are a source of the valuable medicinal alkaloids, hyoscyamine and scopolamine.

(Tobacco, Stramonium) or transversely (Henbane); or a berry (potato, egg-plant, tomato, red pepper). Seeds albuminous.

OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Belladonnæ Folium	Leaves and tops	<i>Atropa Belladonna</i>	Central and Southern Europe, Asia Minor and Persia
Belladonnæ Radix	Root	<i>Atropa Belladonna</i>	
Atropina	Alkaloid	<i>Atropa Belladonna</i>	
Stramonium	Leaves	<i>Datura Stramonium</i>	Asia and Tropical America

OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Hyoscyamus	Leaves and tops	Hyoscyamus niger	Europe, Asia
Capsicum	Fruit	Capsicum frutescens	Tropical America
UNOFFICIAL DRUG			
Dulcamara	Twigs and stems	Solanum Dulcamara	Europe and Asia
Manaca	Root	Brunfelsia Hopeana	Tropical America
Duboisia	Leaves	Duboisia myoporoides	Australia.
Tabacum	Leaves	Nicotiana tabacum	Tropical America
Scopola	Rhizome	Scopola carniolica	Alps and Carpathian Mts.
Paprika	Fruit	Capsicum annum var. longum	Tropical America; cultivated
Pimiento	Fruit	Variety of Capsicum annuum	Spain
Miré	Roots, stems and leaves	Brunfelsia hydrangæformis	South America
Solanum	Ripe fruit	Solanum carolinense	United States

FIG. 455.—The Deadly Nightshade, *Atropa Belladonna*,—Branch. (Sayre.)

SCROPHULARIACEÆ OR FIGWORT FAMILY.—Herbs (*Linaria*, *Verbascum*, *Gerardia*, *Digitalis*, etc.), shrubs (shrubby *Veronicas*, etc.), rarely trees (*Paulownia imperialis*). Stem, branches and leaves usually green and independently vegetating, but in *Pedicularis*, *Gerardia*, *Euphrasia*, *Buchnera*, *Rhinanthus*, etc., the stem, leaves, and branches are condensed from the development of a parasitic root habit. Stems cylindrical to frequently quadrangular, especially when leaves are opposite. Leaves alternate to opposite and decussate, simple, exstipulate, often

FIG. 456.—Bittersweet (*Solanum Dulcamara*). Note transition from simple to lobed to compound leaves and flowers with rotate corollas. (After Gager.)

hairy, but becoming by drought or parasiticism reduced to scales or almost absorbed. Inflorescence a raceme of cymes (*Paulownia*) or a simple raceme (*Foxglove*, *Linaria*, etc.) or spike (*Verbascum Thapsus*) or, if leaves are opposite, often a whorl of axillary flowers or solitary axillary flowers. Flowers rarely regular, mostly irregular; calyx of five sepals, condensed in *Veronica* to four through absorption of one sepal by fusion of two sepals; corolla of five to four petals, deeply synpetalous, varying from rotate (*Verbascum Blattaria*, etc.) to irregular tubular to elongate, irregular bilabiate to funnel-shaped. In color, corolla varies from greenish to greenish-yellow or white (*Scrophularia*)

to pure white or from red to purple to blue (*Veronica*). Stamens five, fertile, equal in length in a few *Verbascum* (mullein) species or unequal in other *Verbascum* species to stamens four with a long sterile staminode (*Pentstemon*, beard-tongue) to four didynamous stamens with a short petaloid staminode (*Scrophularia*, figwort) to four didynamous stamens with a minute often nectariferous staminode (*Linaria*, toadflax), to frequently four didynamous stamens only, the two lateral or two anterior stamens stronger and longer (*Antirrhinum*, snapdragon), to two perfect stamens and two minute staminodes (*Calceolaria*), to two stamens alone developed (*Veronica*, speedwell). Pistil bicarpellate, ovary two-celled with

FIG. 457.—Culver's Root or Tall Speedwell (*Veronica virginica*). Note the rhizome and roots and portions of the stem bearing whorls of lanceolate, serrate leaves and terminal spikes of usually white, often pink or blue flowers. (Reproduced from U. S. Dept. Agric. Misc. Publ. 77.)

central placentation; style terminal with bilobed stigma; ovules numerous, small. Fruit a two-celled and usually many-seeded capsule. Seeds richly albuminous, anatropous or amphitropous.

OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Digitalis	Leaf	{ Digitalis purpurea Veronica virginica	Europe
Digitalis Pulverata	Standardized powd. leaf		
Leptandra (Culver's Root)	Rhizome and roots	Veronica virginica	United States and Canada
UNOFFICIAL DRUG			
Verbasci Flores	Corollas with stamens	{ Verbascum phlomoides Verbascum thapsiforme	{ Europe and Asia
Euphrasia (Eye-bright)	Plant	Euphrasia officinalis	Europe
Verbasci Folia	Leaves	Verbascum Thapsus	Europe and Asia
Balmony	Leaves	Chelone glabra	North America

PEDALIACEÆ OR SESAME FAMILY.—Tropical herbs often thickly covered with viscous hairs. Leaves soft, usually alter-

FIG. 458.—The foxglove, *Digitalis purpurea* var. *gloxinæflora*. Second year of growth.

nate, more rarely opposite, exstipulate. Flowers irregular, pentamerous. Fruit a capsule (*Sesamum*, etc.), drupe, or rarely a one-seeded, indehiscent nut. Seeds exalbuminous usually.

OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Oleum Sesami (Benne Oil)	Fixed oil	Sesamum indicum (cultivated varieties)	Asia and Africa

ACANTHACEÆ OR ACANTHUS FAMILY.—Usually herbaceous (*Ruellia*), rarely sub-woody or woody plants, occasionally bushy

FIG. 459.—*Plantago Psyllium*, the source of an excellent variety of Psyllium seed.

in habit, containing cystoliths in the mesophyll or epidermal cells of the leaves and in the parenchyma of the roots and stems. Leaves opposite, more rarely whorled, entire, exstipulate. Inflorescence a raceme of condensed cymes, becoming a simple raceme or spike, rarely condensed into a solitary

terminal inflorescence. Flowers hermaphrodite, usually irregular; calyx five-cleft; corolla hypogynous, gamopetalous, more or less bilabiate, funnel-form and composed of five sepals; stamens usually four (*Ruellia*, etc.), occasionally reduced to two, as in genus *Dianthera*, didynamous or diandrous, epipetalous; pistil bicarpellate; ovary two-celled, superior, with numerous campylotropal ovules; style terminal, filiform. Fruit a capsule containing numerous curved seeds. The family is of pharmaceutical interest mainly because of *Ruellia ciliosa*, a pubescent perennial herb growing in the Eastern United States, whose rhizome and roots have frequently been admixed with or substituted for *Spigelia*.

PLANTAGINACEÆ OR PLANTAIN FAMILY.—Annual or perennial herbs of caulescent (*Plantago Psyllium*, *P. arenaria*, etc.) or more usually acaulescent habit (*Plantago lanceolata*), rarely sub-shrubs. Leaves all radical in most species, or in a rosette, alternate or opposite to whorled, simple, nerved, flat, entire or toothed. Flowers usually hermaphroditic, regular, tetramerous, and arranged on spikes; calyx of 4 sepals; corolla gamopetalous, hypogynous, tubular with a 4-lobed limb; stamens 4, epipetalous; carpels 2 or 1, the ovary 1 to 4 celled, ovules 1 to 8 in each cell, peltate on the middle of the septum in many ovuled cells or at the bottom in 1-ovuled cells; style filiform with 2 lines of stigmatic papillæ. Fruit a pyxis in the Plantains, rarely a bony nucule. Seeds albuminous with a straight (*Plantago*) to curved (*Bougueria*) embryo. The seed coat of many species abounds in mucilage, accounting for their laxative properties.

OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Plantaginis Semen	Seed	{ <i>Plantago Psyllium</i> }	Mediterranean countries
		{ <i>Plantago arenaria</i> }	
		{ <i>Plantago ovata</i> }	Asia, S. Europe, N. Africa

ORDER RUBIALES

RUBIACEÆ OR MADDER FAMILY.—Herbs, as Cleavers (*Galium*) and Partridge Berry (*Mitchella*), etc., shrubs as the Ipecac (*Cephaëlis*) and the Buttonbush (*Cephalanthus*), or trees (*Cinchona* species) with fibrous roots, sometimes, as in *Cephaëlis Ipecacuanha*, annularly enlarged. Roots, stems and to a less extent leaves

rich in varied alkaloids, some of medicinal value. Leaves opposite, entire, stipulate and interpetiolate. Inflorescence a raceme of ditchesial cymes, occasionally condensing to scorpioids. Flowers perfect, often dimorphic, pentamerous or tetramerous;

FIG. 460.—*Cephaelis Ipecacuanha*—Plant and dried root. (Sayre.)

sepals five (*Cinchona*, etc.) but four in *Galium*, small, green, subtended with other flowers by one or two or more enlarged petaloid bracts; petals five (*Cinchona*, etc.) to four in *Galium*, stellate, varying from shallow-rotate to elongate-tubular or funnel-shaped with stellate limbs; stamens five to four, epipetalous; pistil nearly always bicarpellate, rarely of five to four carpels; ovary inferior, two-celled with central placentation; styles either

distinct with knob-shaped stigmas or style elongate, filiform, ending in bilobed stigmas. Fruit varied, a capsule in *Cinchona*, a berry in *Coffea*, a drupe, or frequently, as in *Galium*, dry and splitting into nutlets; seeds albuminous, each with a curved embryo.

OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Caffeina	Feebly basic principle	{ <i>Coffea arabica</i> <i>Coffea liberica</i>	Eastern Africa Liberia
Cinchona	Bark	{ <i>Cinchona succirubra</i> or its hybrids	South America
Quinina	Alkaloid	{ <i>Cinchona Ledgeri-</i>	
Quinidinæ Sulfas	Sulfate of Alkaloid	{ ana, <i>C. Calisaya</i>	
Cinchoninæ Sulfas	Sulfate of Alkaloid	{ and hybrids of	
Cinchonidinæ Sulfas	Sulfate of Alkaloid	{ these with other <i>Cinchona</i> species	
Gambir	{ Dried aqueous extract prepared from leaves and twigs	{ <i>Ourouparia Gambir</i>	Malay Archipelago
Ipecacuanha	Rhizome and root	{ <i>Cephaëlis</i> <i>Ipecacuanha</i> <i>Cephaëlis acuminata</i>	Brazil United States of Columbia
Emetinæ Hydrochloridum	Hydrochloride of alkaloid emetine obtained from ipecac.		
Mitchella	Entire plant	<i>Mitchella repens</i>	North America
Æthylhydrocupreinæ	Hydrochloridum, a synthetic compound made from cuprea bark or quinine.		
UNOFFICIAL DRUG			
Coffea Tosta	Roasted seeds	{ <i>Coffea arabica</i> <i>Coffea liberica</i>	{ Africa
Yohimbe	Bark	<i>Corynanthe Johimbi</i>	Africa
Cuprea Bark	Bark	{ <i>Remijia pedunculata</i> <i>Remijia purdieana</i>	{ Brazil
Loxa Bark	Bark	<i>Cinchona officinalis</i>	Ecuador and Peru
Ethylhydrocupreine	Alkaloidal compound prepared from cupreine found in bark of <i>Remijia pedunculata</i>		
Cleavers	Herb	<i>Galium aperiene</i>	Europe and Asia
Madder	Root	<i>Rubia tinctorum</i>	Mediterranean region
Sweet Woodruff	Entire plant	<i>Asperula odorata</i>	Europe, W. Asia

CAPRIFOLIACEÆ OR HONEY SUCKLE FAMILY.—Shrubs (Elder, Viburnum, Snowberry, etc.) or rarely herbs (Twin Flower). Leaves entire, opposite, exstipulate or with delicate, attenuate or filiform stipules. Inflorescence varying from a raceme of shortened cymes to a capitulum. Flowers varying from regular and small (*Sambucus*, *Viburnum*, etc.) to increasingly large, slightly irregular and ultimately very irregular, in some Honeysuckles

FIG. 461.—Two medicinal Viburnums. A, *Viburnum Opulus* var. *americanum*, the source of cramp bark. B, *Viburnum prunifolium*, the black haw, whose root bark is official.

(*Lonicera*) and in a few *Weigelas* and allies; calyx pentamerous, superior; corolla superior, gamopetalous, limb pentafid, small in *Viburnum* and *Sambucus* to elongate-tubular or irregular-infundibuliform in Honeysuckles; stamens five, inserted on tube of corolla and alternating with corolla segments; filaments equal or didynamous (in irregular flowers); ovary inferior, rarely five- to three-celled, usually three- or frequently two-celled; style terminal. Fruit a berry (*Honeysuckle*, *Elder*, etc.) from an inferior ovary, several celled, occasionally becoming one-celled with several to rarely one seed and drupaceous (*Viburnum*), or fruit a capsule (*Diervilla*, *Weigelia*). Seeds albuminous.

OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Sambucus	Flower (air-dried)	{ <i>Sambucus canadensis</i> <i>Sambucus nigra</i>	{ United States Europe
Viburnum Prunifolium	Bark of root	{ <i>Viburnum</i> prunifolium	{ Eastern and central United States
Viburnum Opulus	Bark	<i>Viburnum Opulus</i> var. <i>americanum</i>	United States and Canada

ORDER CUCURBITALES

CUCURBITACEÆ OR GOURD FAMILY.—Herbaceous, succulent, very often annual (*Colocynth*, *Pumpkin*, etc.), more rarely perennial (*Bryonia*, etc.), rarely shrubby plants, the perennial and shrubby forms perennating by swollen roots, some of which are heavy and tuberous. Stems very usually grooved and ridged, often provided with roughened and barbed hairs. Tendrils are frequently produced in the axils of leaves from tendril axillary buds (*Pumpkin*, *Colocynth*, *Watermelon*, *Cucumber*, *Bryony*, etc.). Leaves varying from entire, simple, usually deltoid to triangular through stages of trilobate, pentalobate, deeply palmatifid to palmatipartite to seldom approaching compound (*Colocynth*). Venation in nearly all cases palmate. Leaves thin, herbaceous, much expanded, often hairy. Vascular bundles of petioles, branches and stems bicollateral. Inflorescence either of loose cymes or more frequently racemes or spikes or entire axillary inflorescence may become solitary axillary. Flowers pentamerous, very rarely tetramerous, monœcious (*Bryonia alba*) or diœcious (*Bryonia dioica*); sepals 5, gamosepalous, adnate to ovary; corolla of 5, rarely 4 gamopetalous petals varying in size and shape from small to large campanulate (*Pumpkin*) or broadly cup-shaped (*Cucumber*),

FIG. 462.—*Colocynth*—Portion of vine and whole fruit. (Sayre.)

and in color from greenish-yellow to greenish-white to pure yellow to yellowish-white to white; stamens typically 5, epigynous, with anthers either joined by pairs or synantherous; carpels usually 3, rarely 4 or 5, fused to form a compound pistil with a short thick style and lobed stigma; ovary inferior, one- to three-celled. Fruit a pepo (a berry from an inferior ovary with thick skin). Seeds flat and exalbuminous.

OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Bryonia	Root	{ Bryonia alba Bryonia dioica	{ Europe
Colocynthis	Pulp of unripe but full grown fruit	Citrullus Colocynthis	Africa and Asia
UNOFFICIAL DRUG			
Pepo	Seed	Cucurbita Pepo (cultivated varieties)	Probably Tropical America
Elaterinum	Principle from elaterium	Ecballium Elaterium	Mediterranean region
Watermelon Seed	Seeds	Citrullus vulgaris	Southern Asia
Momordica (Balsam apple)	Fruit	Momordica Balsamina	East Indies

ORDER VALERIANALES

This order includes the *Valerianaceæ* or Valerian Family and the *Dipsaceæ* or Teasel Family. Both are represented by herbs with flowers having an inferior ovary containing a single ovule.

VALERIANACEÆ OR VALERIAN FAMILY.—Herbaceous often low succulent plants with creeping rhizomes, frequently strongly scented and possessing stimulating properties. Leaves frequently dimorphic; radical fascicled; cauline opposite; petiole dilated, exstipulate. Inflorescence a raceme of dichasial or scorpioid cymes. Flowers more or less irregular; calyx absent as such, but represented by a series of teeth that are incurved in the bud and flower and which expand later into a pappose crown and act in the fruit as a pappose disseminator; corolla pentamerous, gamopetalous, varying from rotate-synpetalous to irregular-tubular with petals diversely united, in color varying from greenish-white to white or pink (*Valeriana officinalis*) to crimson; stamens three to two or one (*Valerian*), epipetalous; pistil syncarpous; ovary usually one-celled, inferior; style filiform with three stigmatic surfaces. Fruit an akene from inferior

ovary crowned by a persistent, expanded, pappose calyx rudiment. Seeds anatropous, exalbuminous.

OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Valeriana	Rhizome and roots	<i>Valeriana officinalis</i>	Europe and Asia

FIG. 463.—*Valeriana officinalis*—Plant and rhizome. (Sayre.)

ORDER CAMPANALES (CAMPANULALES)

This order comprises the Bluebell Family (*Campanulaceæ*), the Lobelia Family (*Lobeliaceæ*) and the Daisy Family (*Compositæ*). Its outstanding characters are the epigynous corolla,

usually 5 stamens, an inferior ovary and a three- to two-cleft stigma.

CAMPANULACEÆ OR BLUEBELL FAMILY.—Herbs of annual or more commonly perennial growth rarely sub-shrubby or sub-woody in habit, frequently with laticiferous tubes containing a milky juice. Stem upright or feeble and spreading. Leaves alternate, simple, exstipulate. Inflorescence primitively a racemose cyme, condensing into a raceme, to a sub-capitulum and ultimately to a capitulum. Flowers regular, campanulate to campanulate-elongate to elongate and deeply cleft in petals; sepals five, only slightly synsepalous, epigynous; petals five, campanulate to campanulate-tubular to tubular-elongate to tubular and deeply cleft; corolla varying in color from greenish-yellow to yellowish-white to white or again, from yellowish-purple (rarely through yellowish-pink or red) to purple to pure blue; stamens five, epigynous, usually free from corolla; nectary epigynous; pistil usually tricarpellary; ovary as many celled as number of carpels and with central placenta; style single, elongate; stigmas as many as carpels. Fruit a capsule. Seeds albuminous. The plants contain inulin.

LOBELIACEÆ OR LOBELIA FAMILY.—Herbs, with inulin and latex contents, corresponding with *Campanulaceæ* in their vegetative parts, but differing from that group by having irregular flowers (pale blue in *Lobelia inflata*), anthers always united into a tube (synantherous) and pistil always bicarpellate with two-celled ovary and bilobed or bilabiate stigma.

OFFICIAL DRUG	PARTS USED	BOTANICAL ORIGIN	HABITAT
Lobelia	Leaves and tops	<i>Lobelia inflata</i>	United States and Canada

COMPOSITÆ (ASTERACEÆ) OR DAISY FAMILY.—Herbs of annual or perennial habit, rarely shrubs or trees, and with watery or milky juice. Inulin is present in cell sap of parenchyma. Leaves alternate, rarely opposite, simple to compound, exstipulate. Inflorescence a capitulum or a raceme of capitula, each capitulum surrounded by an involucre or protective whorl of bracts, and composed of numerous small flowers called florets that may be: (a) wholly regular, tubular and hermaphrodite (Thistle, etc.); or (b) central florets as in (a), but marginals

strap-shaped or ligulate and usually pistillate (Daisy, Dahlia, etc.); or (*c*) florets all ligulate and hermaphrodite (Dandelion, Chicory, etc.); or (*d*) florets in part or in whole bilabiate (Mutisia, etc.). Flowers small (florets), closely crowded, pentamerous, shaped as above, with ovary inferior and other floral parts superior. Sepals rudimentary, tooth-like (Sunflower), or reduced

FIG. 464.—Capitulum of a composite. Jerusalem artichoke (*Helianthus tuberosus*). A, lengthwise section of capitulum, $\times 1$; B, ray flower, $\times 6$; C, disk flower, cut lengthwise, $\times 6$. (A after Baillon, B and C, Robbins.)

to a pappose or hairy rudiment above the ovary that is functionless during flowering, but that expands in fruit as a hairy fruit disseminator (Dandelion, Thistle, etc.); or sepals wholly absorbed (Daisy). Petals synpetalous, tubular, ligulate or rarely bilabiate, greenish-yellow to white, or through pink-crimson and purple to blue (Chicory). Stamens five, epipetalous, filaments distinct, anthers united into an upright anther-box (so synantherous) into which pollen is shed before or

FIG. 465.—*Arctium minus* or Common Burdock, a source of the drug Lappa or Burdock Root. (Reproduced from U. S. Dept. Agric. Miscel. Publ. 77.)

FIG. 466.—*Grindelia squarrosa* or Gum Plant, a native of the Plains and Prairies from Saskatchewan to Mexico. (Reproduced from U. S. Dept. Agric. Miscel. Publ. 77.)

FIG. 467.—*Brauneria angustifolia*, or Narrow-leaved Purple Cone-flower. (Reproduced from U. S. Dept. Agric. Miscel. Publ. 77.)

FIG. 468.—*Tussilago Farfara* or Coltsfoot. (Reproduced from U. S. Dept. Agric. Miscel. Publ. 77.)

during opening of each floret. Carpels two, syncarpous; ovary inferior, one-celled with single ovule; style simple, at first short, later elongating and by collecting hairs sweeping pollen to top of anther box, then dividing into two stigmatic surfaces with

FIG. 469.—*Matricaria Chamomilla*, the German Chamomile. Branch and dissected flowers. (Sayre.)

stigmatic hairs for pollen reception. Fruit an indehiscent akene, often (Dandelion, Thistle) crowned by the pappose, calyx rudiment. Seed single, exalbuminous.

The Compositæ is subdivided into two sub-groups or series, namely the *Tubulifloræ* and the *Ligulifloræ*. The TUBULIFLORÆ include all those Composites whose flower heads contain either

tubular florets only, as the Eupatoriums and Burdocks or marginal ligulate florets and central tubular florets, as in the Sunflower, Chamomiles, etc. The *LIGULIFLORÆ* include those Composites whose flower heads possess ligulate florets only, as

FIG. 470.—Chicory (*Cichorium Intybus*). A, portion of flowering branch; B, basal leaf (runcinate-pinnatifid); C, median longitudinal section through a head showing the insertion of the flowers; D, individual flower; E, fruit (ripened ovary), showing the persistent pappus (calyx) of short scales. (Gager.)

the Dandelions, Hawkweeds, Chicory, Lettuces, etc. They contain a bitter or acrid milk juice.

OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Arnica	Flower head	<i>Arnica montana</i>	Europe
Matricaria	Flower head	<i>Matricaria</i>	Europe and
		<i>Chamomilla</i>	western Asia
Calendula	Ligulate florets	<i>Calendula officinalis</i>	Mediterranean
			basin
Eupatorium	Leaves and flowering tops	<i>Eupatorium perfoliatum</i>	North America

OFFICIAL DRUG	PART USED	BOTANICAL ORIGIN	HABITAT
Grindelia	Leaves and flowering tops	{ Grindelia camporum Grindelia cuneifolia Grindelia squarrosa	{ Western North America
Taraxacum	Rhizome and roots	Leontodon Taraxacum	Europe
Echinacea	Rhizome and roots	{ Brauneria pallida Brauneria angustifolia	{ Central United States
Lappa	Root of first year plant	Arctium Lappa and Arctium minus	Europe and Asia
Santoninum	Inner anhydride of santoninic acid	Artemisia Cina	Turkestan
UNOFFICIAL DRUG			
Anthemis	Flower head	Anthemis nobilis	Europe
Arnicae Radix	Rhizome and roots	Arnica montana	Europe
Pyrethri Flores	Flower head (unexpanded or partly expanded)	{ Chrysanthemum cinerariæfolium Chrysanthemum roseum Chrysanthemum Marschallii	{ Dalmatia Herzegovina Western Asia
Pyrethrum	Root	Anacyclus Pyrethrum	Northern Africa and southern Europe
Inula	Rhizome and roots	Inula Helenium	Europe and Asia
Senecio	Entire plant	Senecio aureus	United States
Farfara	Leaves	Tussilago Farfara	Europe
Absinthium (Wormwood)	Leaves and flowering tops	Artemisia Absinthium	Europe, Asia and Africa
Lactucarium	Dried milk juice	Lactuca virosa	Europe
Santonica	Unexpanded flower head	Artemisia Cina	Turkestan
Carthamus (Safflower)	Tubular florets	Carthamus tinctorius	India
Achillea (Yarrow)	Leaves and flowering tops	Achillea millefolium	Europe and Asia
Tanacetum (Tansy)	Leaves and flowering tops	Tanacetum vulgare	Europe
Gnaphalium (Everlasting)	Leaves and flowering tops	{ Gnaphalium polycephalum Anaphalis margaritacea	{ North America Asia
Cichorium (Chicory)	Rhizome and roots	Cichorium Intybus	Europe
Oleum Erigerontis	Volatile oil	Erigeron canadensis	North America
Flores Cyani	Florets	Centaurea Cyanus	Europe
Solidago	Herb	Solidago odora	E. United States
Silphium	Root	Silphium laciniatum	United States
Vanilla Leaf	Leaves	Trilisa odoratissima	S. United States
Swamproot	Root	Eupatorium purpureum	United States
Blessed Thistle	Herb	Cnicus benedictus	Europe

CHAPTER XXIV

ECOLOGY

ECOLOGY is that department of biology which deals with the relations of plants and animals of various habitats to their environmental conditions.

Ecology may be divided into two branches, *autecology* and *synecology*. **AUTECOLOGY** deals with the structure and behavior of the individual organism and its parts as related to environment, whereas *synecology* deals with plant communities as related to soil, light, climate and other environmental factors.

Every living thing is a creature of circumstance, dominated and controlled by heredity and environment. Its fundamental structural and functional characteristics are determined by the genes in its protoplasm but its characteristics become modified by external factors of environment. In order to exist and keep healthy it must adapt itself to the various factors of its surroundings. The environmental factors having to do with the existence and health of plants include soils and soil constituents, air, moisture, light, range in temperature, gravity, and surrounding animals and plants of other kinds. Various aspects of most of them have been considered in preceding chapters.

PLANT COMMUNITIES AND ASSOCIATIONS

An entire group of plants of similar habits occurring in a common habitat constitutes what is termed a *plant community*. A *plant association* is a plant community having a definite composition i.e., made up of a definite aggregation of plants developing under similar conditions. Thus, a desert where the cacti and euphorbias form the dominant plants with a definite assemblage of other plants growing with them would represent a *cactus-euphorbia desert association*. Plant communities may be classified either from the point of view of their order of development, as based upon the principle of succession, or upon their

water relation. The latter method appears to be simpler, and will now be considered.

According, therefore, to the relation plant communities have assumed in regard to water, they may be grouped as follows:

1. *Hydrophytes* or water plants.
2. *Helophytes* or marsh plants.
3. *Halophytes* or salt plants.
4. *Xerophytes* or desert plants.
5. *Mesophytes* or intermediate plants.
6. *Tropophytes* or alternate plants.

HYDROPHYTES.—The effect of an aquatic environment on the structure of water plants is most striking. The root systems are reduced both in length and number of branches. The root hairs of those immersed in the water are absent. The supportive action of the water is such that the fibrovascular elements of the stems, which usually function both for support and for the conduction of crude sap, are greatly reduced in size and strength. The leaves, stems and roots possess large air-spaces. The mesophyll of the leaves is spongy and the chloroplasts motile. Stomata are entirely absent from leaves that are submerged and only present on the upper surface of floating ones, where they are nearly always open. Some of these plants have broad floating leaves and dissected submerged ones, often with thread-like divisions. The submerged parts are devoid of special protective walls, *e.g.*, those containing cutin or suberin. The cell sap has a low osmotic pressure. The submerged leaves often absorb more water than the roots. The free floating microscopic plants (blue-green algæ, bacteria, diatoms, desmids, etc.) form the *plankton* of our ponds, rivers and lakes. The free-swimming higher plants (the *pleuston*) comprise certain liverworts like *Riccia* and *Ricciocarpus*, water-ferns and such seed plants as the water-lettuce and water-hyacinth. The aquatic plants, including the algæ, mosses, and flowering plants which live attached to rocks comprise the *lithophilous benthos*. Another class of aquatic plants (*benthos*) include those with true roots, which attach the plant to the substratum, and at most possess floating leaves. This type includes the water-lilies, the water-chestnut, the splatter docks, the floating-heart and the pondweeds.

HELOPHYTES.—To this group belong plants typical to marshes. A **MARSH** is an area with wet soil, wholly or partially covered with water and with annual or perennial herbs (never shrubs and trees) which are adjusted structurally to a mucky soil, lacking the usual supplies of oxygen. These plants likewise show an adjustment to a partial or periodical submergence. Like hydrophytes, marsh plants are for the most part perennial. They produce adventitious roots and possess horizontal rhizomes, or runners, and frequently have air chambers in roots, stems and leaves, so that they are adapted to meet the scarcity of air in wet soils. They also show a striking development of erect chlorophyll-bearing organs in the shape of leaves, in the flags, and stems, in the rushes.

The taller seed-like plants of the marsh-land, such as seed-grass (*Phragmites*), the bur-reed (*Sparganium*), the cat-tails (*Typha*), the blue-flags (*Iris*), the sweet flag (*Acorus calamus*) and the papyrus (*Papyrus*) form associations known as *fresh-water marshes*, *reed-marshes* or *fens*. The channels or pools of water in among these amphibious plants are filled with true aquatic plants.

HALOPHYTES.—The plants of this group live in a soil which is rich in soluble salt, usually common salt (NaCl), and, on account of the fact that the osmotic force of the root is nearly inadequate to overcome that of the concentrated solution of the soil, the soil to such plants is physiologically dry. A halophyte, in fact, is one form of xerophyte. The most striking feature among halophytes is that they are nearly all succulent plants. The leaves of such plants, for example, are thick, fleshy and more or less translucent. They are rich in concentrated cell sap by which they are able to counteract the osmotic pull of the concentrated saline solution of the soils in which they live. Anatomically, they are poor in chlorophyll; the intercellular-air-spaces are small and the palisade tissue is more abundant. Coatings of wax are found and a hairy covering, although infrequent, sometimes occurs. Coriaceous and glossy leaves, especially in tropical halophytes, are noteworthy, while in many salt-loving plants the stomata are sunken. Halophytes are found in our coastal salt marshes and on saline tidal flats in temperate and tropical countries and on the alkali flats of the interior of continents. Notable exam-

ples of these plants are the Salt Marsh Samphire, *Salicornia ambigua*, the Mangroves (*Rhizophora*), the Goosefoots (*Chenopodium*), and the Bald Cypress (*Taxodium*).

XEROPHYTES.—The plants of this group, like the halophytes, are adjusted to live in a soil which is physiologically dry. The soil may owe this condition to its physical nature, such as porosity (sand), or to the presence of humic acids, or by chemical action, which inhibits the absorption of water. They are adapted to meet the conditions of strongest transpiration and most precarious water supply. To meet such conditions of physiological drought, the plants show various structural adaptations. In deserts, where the atmospheric precipitations are less than a certain limit, the plants acquire a xerophytic structure, such as succulency, water storage tissue, associated frequently with mucilage, lignified tissues, thick cuticle to the leaves, depressed stomata (frequently in pits), reduced transpiration surfaces and thorns. Mechanical tissues like wood and bast fibers attain their highest development in these plants. Cacti, conifers, lichens, and the century plants (*Agave*) are types of xerophytes, while many bog plants like the cranberry and Labrador tea, with leathery leaves, are xerophytic.

MESOPHYTES.—These are plants that grow in soil of an intermediate character which is neither specially acid, cold or saline, nor excessively wet or dry, but is sufficiently well supplied with water and rich in the elements required for plant growth. Plants which grow under such conditions do not have structures by which transpiration is closely controlled. They have large leaves frequently toothed and incised, with numerous stomata on lower or both surfaces, a thin, moderately cutinized epidermis, and small intercellular-air-spaces. The leaves and stems are usually of a fresh green color. Typical of the mesophytes are the grasses, the crop plants of field and garden, and most of the annual and biennial herbs of temperate regions.

TROPOPHYTES.—This term was first introduced by Schimper in 1898 for land plants which have deciduous leaves and whose conditions of life are, according to the season of the year, alternately those of mesophytes and xerophytes. The mesophytic condition is found in summer, when the trees, shrubs and peren-

nial herbs, included in this group, are in full leafage, and when, owing to the regular supply of rain during the growing season, the soil is plentifully supplied with water to meet the demands of these plants during the period of active transpiration. During the winter they are xerophytes. The cold of winter freezes the

FIG. 471.—A colony of the Golden Seal (*Hydrastis canadensis*), a mesophytic shade plant or umbrophyte, growing in sparsely shaded woodland.

water in the soil so that the transpiration is reduced to a minimum, and this is associated with the fall of the leaves of the trees and shrubs and the death of the overground parts of the perennial herbs which spring up each year from their underground parts. The vegetation of cold temperate regions is mainly tropophytic.

The deciduous trees and shrubs, also known as the broad-leaved plants and the summer-green plants, form the principal

trophophytes. The deciduous forests, which include the oaks, the beeches, the ashes, the maples, the walnuts, the chestnuts, cover a part of eastern and western China, central Europe (England, France, Belgium, Germany) and eastern Australia, and are coincident with the countries occupied by the most civilized races of man, such as the Americans, Europeans, Chinese and Japanese. The cold temperate climatic conditions which have determined the distribution of the forest trees have been influential also in the development of the energetic races of mankind.

PLANT SUCCESSION.—In the development of vegetation, the same region becomes successively occupied by different communities of plants. This process is called *plant succession*. There are two kinds of plant succession, (1) the *hydrarch* which begins in water and (2) the *xerarch* which may begin on bare rock, rocky talus slopes, or other location where the soil is extremely dry. As the scope of this book forbids little more than mention of this topic, the student interested in it is referred to works on plant ecology.

RELATION OF PLANTS TO ANIMALS

The great and fundamental rôle of green plants in the world's economy is that of constructing highly complex organic compounds for food out of simple, inorganic, raw materials. This food, part and parcel of their very bodies, is eaten by animals whose chief and contrasting rôle is that of breaking down the plant food into its simplest elements.

The raw materials which constitute the ordinary diet of green plants represent salts of various metals, such as nitrates, phosphates, sulfates, etc., the water which they absorb from the soil, and carbon dioxide which they inhale from the air during sunlight.

CARNIVOROUS PLANTS.—There exist, however, about 500 species of green plants which, in addition to the common habits of nutrition possessed by their relatives in the vegetable world, have acquired the luxurious appetite for the flesh and blood of animals. These exhibit a variety of devices for the allurements, capture, imprisonment, digestion and absorption of their prey.

Chief among these carnivorous plants are the sundews, fly-traps, pitcher-plants, bladderworts and butterworts. In each it is the leaves which have become modified for the purposes indicated.

FIG. 472.—The carbon cycle. (After Gager.)

THE SUNDEWS

The sundews are curious, small, bog plants that are mainly members of the genus *Drosera* which comprises 84 species, occurring in tropical, warm temperate and cool temperate regions of both hemispheres. Each of these possesses a rosette of leaves that arise from a greatly reduced root system. The shape of the leaves varies depending upon the species, although it is frequently rotund, spatulate or filiform. The whole upper surface of the leaf-blade is covered with glandular tentacles often of a wine-red color. The glandular heads of these are covered with a viscid secretion that glistens in the sunlight like dewdrops, a phenomenon that accounts for the common name assigned to the members of this group. The tentacles in the central part of the leaf-blade are short and erect and their stalks are green. Toward the margin they become longer and more inclined outward and their stalks are of a wine-red to purple color.

When examined under the microscope each tentacle shows a slender pedicel terminating in a gland. A conducting bundle containing spiral vessels and simple vascular cells is observed to come off of a fibrovascular bundle in the leaf-blade and run

FIG. 473.—*Drosera rotundifolia*, the Round-leaved Sundew. (From Jenkins' "Interesting Neighbors.")

through the center of the pedicel to the gland. Upon entering the gland it is enlarged and spread out into a number of tracheids. These centrally placed tracheids are surrounded by a protective sheath (endodermis) outside of which are two layers of secretory (epidermal) cells with wine-red, granular contents, the outer layer of which is palisade-like. These glands are commonly oval excepting the extreme marginal ones which are considerably

elongated. They secrete, absorb and are acted upon by various stimulants.

Darwin¹ has shown that when a small object is placed on the tentacles in the center of the leaf these transmit an impulse to the marginal tentacles. The nearer ones first respond and slowly bend toward the center and then those farther away until finally all become bent over the object. The time essential varies from 10 seconds to 5 or more hours depending upon the nature, contents and size of the object, upon temperature and also upon the age of the leaf. Again, if the glands are repeatedly touched or brushed or if chemical substances are placed on these, the marginal tentacles curve inward. The bending part of each tentacle is confined to a limited space near the base. Not only the tentacles but the blade of the leaf becomes much incurved when any strongly exciting substance is placed on the blade.

The time during which the tentacles and blade remain curved over the object varies according to the temperature, character of object, age, etc. Dr. Nitschke² found that during cold weather both the blades and tentacles re-expand within a shorter period than when the weather is warm. Darwin³ found that the tentacles remain clasped for a much longer period over objects which yield soluble nitrogenous matter than over those yielding no such matter. After a period varying from 1 to 7 days the tentacles and blade re-expand and are then ready to again respond. It has been shown by the same authority that as soon as the tentacles become inflected over an object yielding soluble nitrogenous matter their glands pour out an increased amount of secretion which becomes acid in nature. These glands, moreover, continue to secrete as long as the tentacles remain closely inflected.

In nature small insects catch sight of the glittering drops on the tips of the reddish tentacles and, mistaking these for honey, alight upon the leaf. They become instantly entangled by the viscid glandular secretion. They try to stroke the viscid fluid off of their legs but only besmear themselves more. Soon they

¹ Darwin: *Insectivorous Plants*, p. 9, 1884.

² Darwin: *Insectivorous Plants*, p. 13, 1884.

³ Darwin: *Insectivorous Plants*, p. 13, 1884.

become covered with the sticky substance which occludes the orifices of their breathing tubes (tracheæ) and they perish in a short time (one-quarter of an hour according to Dr. Nitschke) from suffocation. The surrounding tentacles bend over the insect's body and clasp it on all sides. A digestive juice containing a proteolytic enzyme of peptic character and an acid is now poured out of the glands. This digests the flesh and blood of the insects caught. The water-insoluble chitinous parts of the body are left on the surface of the blade. If the insect be large, the bending of the tentacles is augmented by the inflexion of the whole surface of the leaf-blade which assumes a concave shape. With the tentacles also curved over, the whole leaf simulates a closed fist. The glands then absorb the soluble nitrogenous material which is assimilated by the plant.

Among the insects caught by the sundews are gnats, flies, ants, beetles, small butterflies and dragon flies. The last named are captured by the cooperation of 2 or 3 adjacent leaves.

THE FLY TRAPS

The fly traps comprise two species, viz.: the Venus Fly Trap (*Dionæa muscipula*) and the Submerged *Dionæa* (*Aldrovanda vesiculosa*).

The Venus Fly Trap is restricted to damp localities of a coastal plain strip extending for about 55 miles north and 45 miles south of Wilmington, North Carolina, and nowhere over 15 to 20 miles in width.

Each plant shows a rosette of modified leaves which together are rarely more than 6 or 7 inches across. From the center of these arises a flower stalk bearing a cyme of white flowers which open from April to June.

Each leaf consists of a winged petiole that is studded on both surfaces with small, brown, stellate hairs. The petiole is truncated in front and contracted to the midrib which suddenly broadens out into a blade composed of 2 symmetrical halves that can fold together along the line of the contractile and irritable midrib region. Along the margin of each half of the blade are long, stiff, non-irritable bristles which, on closure of the blade, interlock with each other. On the center of the upper surface

of each half of the blade are three spine-like, sensitive hairs disposed as angles of a triangle. Each of these sensitive hairs is composed of elongated cells whose protoplasmic contents show

FIG. 474.—*Dionaea muscipula*, The Venus Fly Trap. (From Jenkins' "Interesting Neighbors.")

movement. At the base of each is a cylindrical mass of small cells which permits the stiff hair to be bent over. These sensitive hairs are highly irritable and capable of receiving and transmitting a stimulus. Further, over the entire upper surface of

the blade are numerous, closely set, sessile, glandular hairs which, after repeated irritation, secrete an acid digestive juice. Each of these, as observed under a microscope in surface view, consists of a rosette-arranged set of cells composed of 4 to 12 cell radii filled with a crimson-claret pigment.

Each sensitive hair consists of a somewhat elongated structure composed of 3 parts, viz., base, highly sensitive joint and insensitive shaft. The base consists of large epidermal cells enclosing a prolongation of mesophyll cells within, that can receive and propagate a stimulus from the sensitive joint to the leaf interior. The joint consists of elongated columnar cells that enclose similar columnar mesophyll cells with soft, elastic walls. The shaft cells are elongated, thick-walled and almost insensitive.

When a stimulus is applied to any sensitive hair, this affects the joint cells and causes upsetting of turgidity and exudation of liquids with a consequent contraction of the elastic cells. The action is propagated to the midrib region where the cells, by contraction along the upper surface and expansion along the lower, cause closure of the halves.

Macfarlane¹ has shown that when an insect alights on a leaf of *Dionaea*, one hair must neither be stimulated twice or 2 hairs on the same leaf at slight intervals apart in order to cause closure. The caught insect produces a summation of stimuli, and gradual tightening of the lamina occurs under the repeated stimuli until the halves become closely locked. The glandular hairs now pour out an acid digestive secretion which digests the flesh and blood of the insect's body. The soluble nitrogenous substances are then absorbed by the glands and assimilated.

Dr. J. S. Hepburn² found that the secretion of the leaves of *Dionaea* contained a protease which was active in the presence of 0.2 per cent. hydrochloric acid. This enzyme, therefore, resembles pepsin of the gastric juice of man, which also acts in a 0.2 per cent. hydrochloric acid medium.

Dr. J. M. Macfarlane, who with Dr. Canby and Charles Darwin have separately investigated the structure and physiological activities of this plant, reports that 2 touches one-fourth

¹ Macfarlane: Contrib. Bot. Lab. U. of Pa., 1: 7-44, 1892.

² Hepburn: Jour. Franklin Inst., 194: 780, 1922.

to one-third of a second apart will not produce closure of the halves, though a second stimulus that is from 1 to 120 seconds apart from the first will effect closure, also that it is not essential for the stimulus to be on one hair, that a prick sets off the trap at once due to liquid escaping from turgid cells and, further, that all muscle stimulants, mineral acids and ammonia effect closure of the trap.

Among the insects caught by *Dionæa* are earwigs, millipedes, flies, ants, wood-lice and dragonflies. The length of time required for the digestion of their softer parts and the absorption of the soluble products of digestion varies. During this time the trap remains closed. The power of digestion possessed by *Dionæa* is more limited than that of *Drosera*, which has been known to capture and digest many insects in a shorter period of time.

Aldrovanda vesiculosa, which might well be called a "Submerged *Dionæa*," is a relative of the Venus Fly-Trap. It is a floating aquatic plant found widely distributed in shallow ditches and ponds over the old world from Europe to Australia but nowhere abundant. This plant is entirely devoid of roots, but possesses a slender stem which bears whorls of modified leaves at its nodes. Like the Venus Fly Trap, each leaf shows differentiation into a winged petiole ending in 5 narrow processes that connect with a terminal, rounded, incurved blade, divided into equal halves by a sensitive midrib, but the midrib projects beyond the summit of the lamina as a bristle. Long, rigid, spiny bristles extend from the petiole and are thought to prevent the approach of animals unsuitable as prey. The margins of each half of the blade are bent inward and their rims are studded with short conical teeth. Projecting from the upper surface of the midrib and along a line describing the inner third of the upper blade surface are a number of sensitive hairs and short-stalked, disc-shaped glands, while over the outer portion of the surface are to be noted a number of scattered stellate hairs.

Larvæ of aquatic insects and small species of Crustaceans such as *Cyclops*, *Daphnia* and *Cypris*, swimming by, occasionally brush against the sensitive hairs and the 2 halves of the blade close together, just as in *Dionæa*, and the animals are entrapped.

When they attempt to escape through the place where the margins of the blade meet, they find the conical teeth prevent their egress. They die in the trap, and when the latter are forced open and examined a couple of weeks later they only contain the chitinous skeleta.

THE PITCHER PLANTS

The pitcher plants are for the most part found in bogs. Their main representatives belong to the genera *Heliamphora*, *Sarracenia*, *Darlingtonia*, *Nepenthes* and *Cephalotus*. The pitcher itself represents the hollowed out midrib of the leaf in the first four named genera and an inpouching of the leaf in the last named genus.

*Heliamphora*¹ is represented by a single species (*H. nutans*) which is only found on and around the base of Mount Roraima, between British Guiana and Venezuela. Here it flourishes in widespreading dense tufts in wet places where the grass is short. It has a rosette of red-veined pitcher-leaves and delicate white flowers raised high on red-tinted stems. Along the entire length of the pitcher are 2 broad wings. The pitcher itself represent a hollowed out midrib which is tubular in shape, becoming gradually broader from base to mouth and ending in a small lid. Over the entire outside of the leaf are nectar glands and upward directed hairs. The nectar glands secrete a sweet fluid which entices insects so that this surface can be termed the alluring surface. Nectar glands also occur on the inner surface of the lid. Next, the upper one-third to one-half of the inner surface of the pitcher is covered with downwardly projecting hairs and nectar glands. This surface can be called the attractive and conducting surface. A smooth surface follows, which in turn is succeeded by a detentive surface in the lower part of the pitcher, composed of mostly smooth-walled cells, a few of which bear short hairs.

*Sarracenia*¹ is represented by 7 species all of which are confined to Eastern North America except *Sarracenia purpurea*, which extends west to Western Minnesota and West Central Canada. Their pitched leaves are of varying form and color design,

¹ Macfarlane: *Sarraceniaceæ* in Engler, *Das Pflanzenreich*, 34 Heft, (iv, 110) 39 pp., 1908.

depending upon the species. Frequently they are long funnel-shaped, tubular or vase-shaped. The color of every species varies with its age as well as extent to which it is exposed to

FIG. 475.—*Sarracenia purpurea*, a pitcher plant from a northern bog. (From Jenkins' "Interesting Neighbors.")

bright sunshine. In some respects the form of pitchers resembles *Heliamphora*, but there is only a single wing present and further, the lid is considerably larger.

In all we find the pitchers showing an alluring outer surface covered with nectar glands, but devoid of the upwardly directed

hairs seen in *Heliampora*, an attractive inner lid surface, marked with numerous downwardly directed hairs and nectar glands which lead the insect toward the next or conducting surface. This is either nearly smooth (*S. purpurea*) or presenting delicate, downwardly projecting processes and on which the insect loses its foothold, also (excepting *S. purpurea*, which next shows an intervening glandular zone) a lower dententive zone, in the bottom of

FIG. 476.—A colony of *Sarracenia purpurea* growing along the border of a pond in Burlington County, New Jersey. *Chamaecyparis* in background.

the pitcher, which contains a watery fluid and whose surface, save its lowest portion, is marked by the presence of many elongated, downwardly directed, thick-walled hairs, which prevent the ascent of insects that have fallen into this zone from above. Glands are also present on the upper part of the detentive surface.

*Darlingtonia*¹ is represented by a single species, *Darlingtonia californica*, which grows in mountain swamps and borders of small streams at altitudes of 1000 to 6000 feet from Plumas county in the Sierras of California northward to Jackson and

¹ Macfarlane: "Sarraceniaceæ," in Engler, *Das Pflanzenreich*, Heft (iv, 110), 39 pp., 1908.

Josephine counties in Oregon. Its underground rhizome gives rise to pitcher leaves which are from $1\frac{1}{2}$ to nearly 3 feet in height and spirally twisted in about a half revolution. Each

FIG. 477.—*Sarracenia flava*, a pitcher plant from a North Carolina bog. (From Jenkins' "Interesting Neighbors.")

pitcher expands near the summit into an inflated hood which exhibits a circular opening up to an inch in diameter on the under side. The dome of the hood is spotted with large, thin, translucent areas. A wing extends along the pitcher from

rhizome to its orifice. At the upper and outer edge of the opening is a moustache-shaped appendage which possesses stiff hairs all of which point toward the pitcher opening. The color of the pitcher is green, blotched with red and yellow. Within and about the opening and on the bi-lobed appendage are alluring glands which secrete a nectar attractive to insects. The nectar is also secreted by glands along the wing. An insect creeping over the exterior of the pitcher is enticed to the nectar along the wing which it follows up to the orifice where the honey is sweetest. Flying insects are attracted to the pitcher by its mottled and colored features. They alight on one of the flaps of the appendage and follow its curve which narrows to the orifice. Sharp bristles in the path pointing toward the orifice make it the natural direction for the insect to travel. Upon reaching the end of the path, it is tempted farther by honey glands within the opening of the pitcher which it next visits. When satisfied and ready to leave, the translucent areas on the hood above, like illuminated windows, entice it from the opening by which it entered. The insect sees no means of escape, nothing but hairs on the inner surface of the pitcher pointing downward and which direct it to the dark detentive pit below, where it drowns in the watery secretion.

FIG. 478.—Darlington's Pitcher Plant (*Darlingtonia Californica*) from mountains in California. (From Jenkins' "Interesting Neighbors.")

By far the most beautiful and gracefully shaped pitchers are those belonging to the genus *Nepenthes* which is represented by about 68 species that are mainly indigenous to the East Indies with headquarters in North Borneo and with Mount Kina Balu as a center. Many of the plants are continuously exposed to a moist dripping atmosphere.

FIG. 479.—*Nepenthes Rajah*. (From St. John's "Life in the Forests of the Far East.")

Each of the modified, pitched leaves shows a petiole (usually winged) which widens into an expanded blade, the midrib of which is continued as an elongated tendril which expands into a terminal winged pitcher. The pitchers show a large variety of shapes and color designs depending upon the species. Frequently they are tubular, goblet or cornucopia shaped. Each pitcher has a hinged lid which varies from small elliptic to large heart- or kidney-shaped. The tendril is sensitive to contact stimuli and often winds about a limb of a tree. The pitchers, partly filled with a viscid watery secretion, either hang suspended in the air from the tendril or rest on the ground.

One of the largest pitchers found in this group is that of *Nepenthes Rajah*, a species found on Mount Kina Balu along the north coast of Borneo at an altitude of 5000 feet. It has leaves which, exclusive of petioles, are 18 inches long, and the pitched portion is 6 inches in diameter and 12 inches in length with a circumference of 19 inches. The lid is 10 inches long and 8 inches broad. The pitchers rest on the ground in a circle. The color of the pitchers is deep purple. The color of the lid portion is lavender, shading to green at the edges. One pitcher noted by Spenser St. John¹ held 4 pint bottles.

The lower surface of the petiole, lamina and the whole external surface of the tendril and pitcher are covered with alluring nectar glands. These alluring glands also occur along the stem. Their secretion entices insects to the pitcher mouth or lid. Around the mouths of the pitchers of nearly all of the species is a corrugated rim on which are parallel ridges, which are usually extended as teeth that are inclined downward into the pitcher cavity. The surface of this rim is smooth and polished, so affording a poor foothold for insects. Around this rim, in the region of the teeth, are the openings of deeply sunken, marginal glands which pour forth a very sweet nectar, attractive to insects. The inner surface of the lid is usually dotted with attractive glands.

In many species of *Nepenthes*, the entire inner surface of the pitcher cavity constitutes the detentive surface. It is lined with a smooth epidermis that is uniformly studded with glands which secrete a digestive juice containing a proteolytic ferment and absorb the soluble products of digestion. In many others the upper surface of the cavity is smooth, forming a conducting surface, while the lower one-half or one-third is alone detentive. Insects lose their foothold on the conducting surface and fall into the watery secretion at the bottom of the pitcher.

Cephalotus has but one species, *C. follicularis*, which is native to swamps of South West Australia. It has short creeping rhizomes which produce annually one set of 5 foliage leaves and later a set of pitcher leaves which rest on damp soil. The pitchers externally are equipped with winged ridges which provide a handy

¹ St. John, S.: Life in the Forests of the Far East, 1: 334, 1863.

means of ascent to the mouth for creeping animals, while their lids are half-closed and mottled with white areas and purple veins and are often mistaken for flowers by high flying insects. In minute details of structure the pitchers are surprisingly like those of *Sarracenia* and *Nepenthes*. The exterior of the stalk is studded with alluring glands. The margin of the pitcher is corrugated and inflexed, and all over this area and exterior of corrugation as well as on the inner lid surface are an abundance of honey glands (attractive surface). Just inside of the margin is a circular shelf-like ingrowth consisting of downward-directed cells which resemble the conductive surface of *Sarracenia*. The remainder of the inner surface constitutes the detentive surface. At the bottom of the pitcher on either side are two beautiful crimson to crimson-claret patches. These have sunk in them huge, many-celled glands which have a fibrovascular bundle terminating in their base. The glands secrete an acid digestive juice which partly dissolves the animal captives. A viscid secretion partly fills the pitcher cavity.

As to relations of the pitchers of these several types of curious plants to animals, it may be said that insects, spiders, etc., are attracted to the plants partly by the brilliant colors of their pitchers and partly by the nectar drops exuded by honey glands. They make their way to the exterior of the lid (or appendage in *Darlingtonia*) where the secretion along the outer margin is plentiful. From here they move to the inner surface of the lid and sip the honey. Tempted further by the sight of nectar ahead and often by hairs pointing downward, they step inside the pitcher on to the conducting surface. Upon reaching this surface they waver, slip off and fall into the lower part of the pitcher which is the water holding region. They make numerous attempts to escape but find exit impossible on account of the downward-projecting hairs. They drown in the liquid and their bodies are either decomposed or digested by a digestive juice secreted by glands lining the lower portion of the pitcher cavity.

The animals caught by these pitcher plants are various. While insects constitute their chief prey, slugs, spiders and rodents have been recorded by observers as having been captured by a

number of them. Burbidge¹ observed several *Nepenthes* in North Borneo visited by a small rodent, which, while perched on the margins of the pitchers bends its head and neck and scoops out the caught insects and devours them. The same writer states that if it attempted such action with *N. bicalcarata*, the

FIG. 480.—*Nepenthes Edwardsiana*. (From St. John's "Life in the Forests of the Far East.")

two sharp spurs with which the pitcher of this species is provided catch it by the neck and tumble it into the pitcher.

Macfarlane² has carefully observed the relations of *Nepenthes* to animals. He states in part that "running insects such as ants and cockroaches are their principal prey. Cockroaches run up the stem and may pause to sip nectar from the alluring stem glands. Reaching the base of the leaf, they may pass along it,

¹ Burbidge: Gard. of Sun, pp. 40-344, 1880.

² Macfarlane: Nepenthaceæ, in Engler, Das Pflanzenreich, 36 Heft., 88 pp., 1908.

attracted by the presence of honey drops there. They almost invariably run along its under side to shelter themselves from enemies and the hot sun. Moving on restlessly and sipping from nectar glands as they advance, they reach the tendril. The ventral wings and the areas between are more beset with alluring glands than is the outer part of the pitcher and along this they often run till they reach the orifice or lid. The lid glands of the inner surface prove a greater attraction but their secretion does not compare with the marginal glands. Straining to reach the orifices of the glands, the insect visitors often overreach themselves after a few efforts and tumble into the pitcher cavity, and in rare cases is escape again possible."

Mr. Low who accompanied Spenser St. John on his travels in North Borneo found a drowned rat in one of the pitchers of *Nepenthes Rajah*.¹

The character of the enzymes found in the pitcher secretions has been investigated by Hepburn.² This author found a tryptic enzyme in *Nepenthes* which was active in a slightly acid medium. No proteolytic enzyme was found by him in the secretions of *Darlingtonia* while the *Sarracenias* he investigated were found to possess proteolytic enzymes. The protease in the pitcher liquor of *Sarracenia flava* and *S. minor* acts best in 0.2 per cent. hydrochloric acid while that present in *S. purpurea*, *S. rubra*, *S. Sledgei* and *S. Drummondii* acts best in an alkaline solution containing 0.5 per cent. or less of sodium carbonate. Dakin³ found a protease in the pitcher liquor of *Cephalotus*.

THE BLADDERWORTS

The bladderworts belong to the genus *Utricularia*. They are aquatic plants which are found in pools and bogs of various parts of the world with their centers in S. America and the East Indies.

The plants are rootless and according to the season of the year sink to the bottom or rise to near the water surface. They have finely dissected submerged leaves, some of which are specialized

¹ St. John, S.: Life in the Forests of the Far East, 1: 335-336, 1863.

² Hepburn, St. John and Jones, Jour. Frank. Inst., 189: 152, 1920.

³ Dakin: Jour. and Proc. Roy. Soc. W. Australia, 4: 37-53, 1919.

as bladder-like traps which allow the entrance but not the escape of small animals. Each of the traps represents a greatly modified, inflated leaf of vesicular form and usually about $\frac{1}{6}$ to $\frac{1}{5}$ inch in diameter. From the margin of its opening project

FIG. 481.—A Bladderwort (*Utricularia Grafiana*). Note the submerged stems bearing modified, bladder-like leaves. Small animals become entrapped in these bladders where they are decomposed and the water-soluble portions of their remains are absorbed as nourishment by the plant. In the lower right-hand corner is a bladder-like leaf of another species (*Utricularia neglecta*) shown in section. The entrance to the bladder is protected by numerous hairs and by a trap door (t). Drawing based upon Kerner. (Gager.)

stiff bristles. There is an upper and lower lip present. The lower lip is greatly thickened and provided with a cushion-like process that extends into the bladder. From the upper lip is suspended a thin elastic valve, the free edge of which rests upon the inner surface of the lower lip cushion and closes the opening.

Small animals, including crustaceans like *Cyclops* and *Daphnia*, as well as larvae of small insects and worms, enter the bladders either because they seek a sheltered harbor for a time from larger prey or because they expect to find food within the bladder. In entering they have to press upon the valve and push it back. As soon as they stop pressing upon the valve, the elastic character of the valve causes it to close by bringing it back on the under lip cushion. The entrapped animals struggle to escape by pressing themselves against the valve, but their efforts are in vain, for it is impossible for the prisoners to force the valve outwardly over the cushion, and they die in time either from starvation or suffocation. Their bodies decompose and the water-soluble products of decay are absorbed by quadrifid cells lining the entire inner wall of the bladder.

THE BUTTERWORTS

The Butterworts comprise an interesting group of flesh-eating plants which are placed in the genus *Pinguicula*. About 40 species exist of which probably the best known is *Pinguicula vulgaris*, a member of the *Utriculariaceæ* which occurs in bogs or other damp places generally in mountainous districts of northern North America and Europe. Each plant exhibits a root system of from about 5 to 16 short, submerged, unbranched roots from which arises a rosette of oblong-ovate, yellowish-green leaves, the younger central ones being concave and more or less erect, the older marginal ones being flat or convex with their lower surfaces resting upon the moist ground. From the center of the leaf rosette there arises a slender scape bearing a single flower of violet blue hue that is spurred in its corolla portion.

Alike with other investigated species of *Pinguicula*, the leaves of this plant have somewhat upturned margins which give them the form of a broad trough, the upper surface of which is covered with a viscid secretion. Microscopical examination of this surface discloses 2 types of glands (excepting along the margin), the first being of toadstool shape and consisting of a cylindrical stalk bearing a disc-shaped head of 8 to 16 cells arranged radially, the second consisting of a shorter stalk bearing an 8-celled head.

Darwin¹ and others have experimented with the leaves on living plants of several Butterworts and found that drops of water, gum or sugar, and scratching of the surface of these leaves produced no response, that insoluble solid bodies such as grains of sand and powdered glass allowed to remain for some time caused slight incurving of the margins and a slight increase in the quantity of mucilage secreted by the glands, but that when nitrogenous bodies such as bits of cartilage, meat, clotted blood, etc., are placed near the margin in contact with the glands they are excited and pour forth abundant mucilage and an acid secretion. Moreover, prolonged contact with these bodies caused a marked incurving of the margin which pushed them slowly toward the middle of the leaf where more glands are present and so increased secretion. This acid secretion completely dissolved these substances which were later absorbed by the leaf.

When small insects alight from the air upon the leaf of the butterwort, they become fastened by the mucilage and in struggling to extricate themselves only become deeper enmeshed by it. Within a short time the acid juice, secreted by the glands as a result of the stimuli, digests their bodies and all excepting their hard, indigestible outer skeleta (exoskeleta) is absorbed.

If small creeping insects come upon the margin of the leaf they stimulate the edge to curl over and the marginal glands to secrete. While the insect is enmeshed within the adhesive mucilage, it is slowly pushed by the curling up of the edges into the middle of the leaf where the acid secretion glands are most abundant. In about 24 hours the softer parts of its body are digested and absorbed and the leaf expands to its normal condition.

FIG. 482.—*Pinguicula vulgaris*, a butterwort. An extract of this plant containing a proteolytic enzyme is used in certain sedative expectorant compounds. (After Le-Maout and Decaisne.)

¹ Darwin: Insectiv. Plants, pp. 371–390, 1899.

SIGNIFICANCE OF THE CARNIVOROUS HABIT

In all of the animal-eating plants which have been investigated structurally and chemically, sufficient evidence has been recorded to warrant the conclusion that these forms have developed their peculiar habit and correlated structures with the object of providing adequately for their nutrition.

CHAPTER XXV

GENETICS AND EVOLUTION

GENETICS is the science which deals with the coming into being of organisms. It deals with heredity, variation, sex determination, and the origin of individual plants and animals.

HEREDITY is the faithful transmission of characters from parent to offspring. While similar to genetics it is not entirely the same, since the heredity complex, representing what the offspring receives at the time of fertilization, may be changed by environment. Genetics deals with both the parental complex and the environmental complex. Modern genetics is based upon the law that the chromatin material of germ plasm is inherited.

All organisms are described in the literature in terms of characters. These are the units employed to distinguish each individual, variety or species from another. Characters are evident in the outlines of leaves, color of leaves, size of seeds or plants, etc., and are the result of hereditary material acquired from the parent, modified by environment. They are mainly hereditary. Hereditary substance is chromatin material in the nucleus of the cell.

MENDEL'S LAWS OF ALTERNATIVE INHERITANCE

The methods of heredity which have been confirmed in the main by the experiments and observations of many biologists were discovered by Gregor Johann Mendel (1822-1884), an Austro-Silesian monk of Br $\ddot{u}$ nn, Austria.¹ Mendel was the first to determine that the genes or "*unit characters*," as he termed them, can separate from one another and recombine in such a way that the characters which they represent appear to act as separate units that can often be removed from one strain and introduced into another.

¹ Although Mendel's Laws were published in 1866, they were disregarded until 1900 when DeVries unearthed them and brought them to the attention of biologists.

Mendel experimented with the hereditary characters of various kinds of garden peas because these were the best materials he could find to test out his ideas and obtain light on the problems of heredity and variation in which he was so keenly interested. He studied seven pairs of constant differentiating characters in the peas he worked with, namely, tall vs. dwarf stems; round (smooth) vs. wrinkled (angular) seeds; yellow vs. green seeds; colored flowers and seed coats vs. white flowers and seed

FIG. 483.—Gregor Johann Mendel at the age of forty. (Gager.)

coats; parchmented vs. non-parchmented pods; green vs. bright canary yellow, unripe pods; and normal round stems with axillary flowers vs. fasciated (flattened) stems with terminal or umbellate flowers. The members of such pairs of contrasting (alternative) characters were called *allelomorphs* by Bateson. Each member of two different strains of plants which can be crossed, as for example, a tall stemmed pea with a dwarf stemmed pea, is said to be *allelomorphic* to the other.

Mendel's Laws embody the following principles:

1. **THE PRINCIPLE OF UNIT CHARACTERS.**—The inheritance of any organism may be analyzed into a number of independent characters (unit characters) which are inherited in toto and are not further divisible. Thus, every organism is a complex of a vast number of independent heritable units.

2. **THE PRINCIPLE OF DOMINANCE.**—When contrasting unit characters are present in the parents they usually do not blend in the offspring, but one is *dominant* while the other is *recessive* and temporarily not visible.

3. **THE PRINCIPLE OF SEGREGATION OF UNIT CHARACTERS, OR THE PURITY OF THE GERM CELLS (GAMETES).**—Every germ cell is pure with respect to any given unit character. In the germ cells

of hybrids there occurs a separation of the determiners of contrasting characters resulting in the production of different kinds of germ cells, each of which is pure in respect to any given character. This principle was later called "the splitting of hybrids" by DeVries.

Probably the most outstanding principle in the Mendelian Law of heredity is the fact that the heritable characteristics of organisms are independent one of another and may be inherited independently. The determining substances of these characteristics must, accordingly, occur in the germ cells as independent, self-perpetuating particles. Such substances are called *genes*, or *factors* or *unit characters*.

STRUCTURE OF THE CHROMOSOME.—In the resting nucleus the nuclear reticulum consists of chromatin threads. These threads as often observed are longitudinally double. Each unit of the double thread appears as a chain of beads and consists of a linear series of granules called the *chromomeres* and these in turn of one or more smaller particles called *chromioles*. The chromomeres lie opposite each other forming pairs. The determiners within the germ cells are believed to occupy the same arrangement, so that chromomeres or their components may represent genes.

MALE AND FEMALE CHROMOSOMES.—Every sexually produced organism is a double being. Every cell of a plant or animal is double, one half having been derived from the male sexual cell and the other half from the female sexual cell. Each of these cells contains two sets of chromosomes, one set from the spermatozoon and one set from the egg. These two sets of chromosomes contain the genes derived from both parents.

During the maturation of the germ cells the chromosomes become unravelled. Each paternal chromosome is matched by a homologous chromosome of maternal origin. During maturation homologous chromosomes and also their chromomeres lie side by side in pairs. At the time of reduction of the chromosomes each pair of chromosomes (maternal and paternal) appear as two parallel strings of beads, the beads representing chromomeres and containing the genes.

DISTRIBUTION OF CHROMOSOMES DURING MATURATION AND FERTILIZATION.—Maturation involves the reduction in the

FIG. 484.—Diagrams illustrating the behavior of chromosomes during the reduction division which occurs at the time of spore formation in ferns and in most other plants which have sexual reproduction. The maternal chromosomes are shown in black, the paternal in white. Homologous chromosomes (corresponding maternal and paternal chromosomes) are of corresponding length. As a result of the process here shown, the spore mother cell (in the resting condition in 1) gives rise to a tetrad or group of 4 spores as shown in 10. The spore mother cell nucleus (1) has the diploid chromosome number (in this case 6) which is characteristic of the cells of the sporophyte. These six consist of 3 pairs of homologous chromosomes, one of each pair having come from the male gamete and one from the female gamete in the last fusion of gametes. Note that in 2 the three pairs with the chromosomes of each pair in close contact are lying within the nuclear membrane. In 3 the paired chromosomes have shortened and have become closely “knotted” whereas in 4 they have become “unraveled” and have undergone further shortening. In 5 the members of each pair have separated but are still close to one another. In 6 the nuclear membrane and the nucleolus have disappeared and the six chromosomes, each showing evidence of a later lengthwise splitting, have been separated into two groups of three. *This is the actual reduction division.* Note that in 7, only one chromosome of each homologous pair is present in each of the resulting nuclei which are therefore different in the chromosome “stock.” Each of these nuclei now undergoes a second or homotypic division (8) with a splitting of each of the chromosomes, so that there are two kinds of nuclei in the tetrads (shown in 9), two having the same chromosomes as the upper nucleus in 7 and two having the same chromosomes as the lower nucleus shown in 7. (From “*A Text Book of General Botany*” by Holman and Robbins by permission of the Publishers, J. Wiley & Sons, Inc.)

number of chromosomes to one-half that of the somatic cells. In the primary spermatocyte and primary oöcyte of animals at synapsis the chromosomes unite in pairs. One member of each pair was derived from the mother, the other from the father. When division occurs, the chromosomes do not divide, but entire chromosomes separate and are distributed in pairs to two daughter cells. At the division resulting in spermatids and mature egg, the pairs separate and each daughter cell receives univalent chromosomes. A similar condition prevails during the division of the mother cells of pollen grains and embryo sacs in plants. Each spermatozoon and ovum is thus provided with one-half the number of chromosomes found in the somatic cells or primordial germ cells.

DISTRIBUTION OF GENES DURING MATURATION AND FERTILIZATION.—It has been shown that homologous chromosomes consist of homologous chromomeres. Since the homologous chromosomes carry the genes and the genes are also paired, the genes will be distributed as were the chromosomes.

ALLELOMORPHS.—Genes that occupy corresponding positions in homologous chromosomes are called *allelomorphs*. The members of a pair of *allelomorphs* may be the same or they may differ; if they are the same, the individual bearing them is said to be *homozygous* as far as the character is concerned that is controlled by the particular genes; if they are different, the individual is said to be *heterozygous*.

DOMINANTS AND RECESSIVES.—In a heterozygote the genes are of two kinds, *dominants* and *recessives*. A gene is said to be dominant when the character it represents appears in the heterozygote; its *allelomorph* in such a case is said to be recessive. In peas, tallness is dominant, dwarfness recessive; in the sunflower, the branched habit is dominant and the unbranched habit is recessive. The result of dominance and recessiveness in *allelomorphic* genes is that heterozygotic individuals do not exhibit all that they inherit from their parents, since some of the genes are recessive.

GENOTYPE AND PHENOTYPE.—The term *genotype* is used to express the entire genetic make-up of an organism. Individuals having a similar germinal constitution are said to be of the same

genotype. The term *phenotype* is used to describe that part of the inheritance that appears in the individual. Forms alike in body characteristics irrespective of their genetic constitution are of the same *phenotype*.

MONOHYBRIDS.—A monohybrid is an individual whose parents differ with respect to one character.

FIG. 485.—Diagram illustrating the results obtained from crossing a variety of garden pea breeding true to smooth cotyledons with one breeding true to wrinkled cotyledons. (Gager.)

RESULTS OF CROSSING INDIVIDUALS WITH ONE PAIR OF CONTRASTING CHARACTERS

Mendel first made sure the characters with which he was to work were constant for certain varieties and then began crossing one variety with another which differed from each other in some of the previously mentioned characters, studying the offspring of several successive generations with respect to the behavior of the contrasting characters.

In every instance he found that the plants that developed from the cross involving a single pair of alternative characters

showed only one of the characters occurring in the parental plants. The obvious character he termed "*dominant*," the hidden or latent character, "*recessive*." In one of his experiments, he crossed tall pea plants with dwarf pea plants by transferring the pollen from the anthers of the tall plant to the stigma of a dwarf plant; the seeds resulting grew to be tall plants like the tall parent. These offspring plants represented the first hybrid generation called the *first filial generation* and are represented by the symbol F_1 . Accordingly, Mendel designated tallness the dominant character and dwarfness the recessive character. He next permitted the tall F_1 hybrids to be self fertilized and their offspring called the *second filial generation* (F_2) were found to be tall and dwarf in the ratio of three tall to one dwarf. With seed obtained from self fertilized F_2 plants he grew an F_3 generation. He found that all the dwarfs (recessives) of the F_2 when so inbred, gave only dwarfs which bred true indefinitely, while all the tall plants of the F_2 generation, when inbred, proved to be of two kinds, one-third pure dominant talls which bred true indefinitely, and two-thirds hybrids similar in number of kinds and ratio to that obtained by growing seed of F_1 plants, *i.e.*, three dominants to one recessive. These results may be diagrammatically represented as follows:

When there is only one pair of contrasting characters usually but two types of offspring occur in the F_2 generation, dominants and recessives.

RESULTS OF CROSSING INDIVIDUALS WITH TWO PAIRS OF CONTRASTING CHARACTERS

Dihybrids are individuals whose parents differ in respect to two pairs of contrasting characters. When Mendel set out to prove the phenomenon of a 3 to 1 ratio in the number of domi-

nant characters and recessive characters appearing in the progeny of crosses between plants examined for one set of contrasting characters such as tallness and dwarfness, he found that by the same method experiments could be carried on in which two sets of contrasting characters would be involved. His variety of Pea plants that differed in one set of contrasting characters, when crossed, produced monohybrids or factors represented by a suitable symbol such as Dd (Tallness D , Dwarfness d). Two pairs of contrasting characters such as a variety of Pea plant which Mendel found to produce yellow, wrinkled seeds and another plant producing the characters for smooth, green seeds, when cross-pollinated, developed spores which gave rise to four kinds of sperm and four kinds of eggs and not the characteristic two kinds of sperm and eggs as of the monohybrids. This condition of a progeny in which there are two separate contrasted characters is called a *Dihybrid*. In the cross of a variety of Pea plants having green color, smooth surface seeds with another variety having yellow color, wrinkled surface seeds the following progeny appears in the first filial generation: Factors for smooth (indicated by S) and yellow (indicated by Y) are dominant; whereas factors for wrinkled (indicated by s) and green (indicated by y) are recessive. The F_1 diagrammatically is:

$$\begin{array}{c} S y \times s Y \\ \swarrow \searrow \\ SsYy \text{ (all smooth and yellow)} \end{array}$$

In the second filial generation or F_2 , after self-fertilizing the progeny of the first filial generation or F_1 , it is then noted that there are four possible kinds of male gametes and four possible kinds of female gametes. During mitosis, the male gametes (indicated by σ) unite with the female gametes (indicated by φ) and there is a union of four different male characters (indicated by S, s, Y, y) with four different female characters (indicated by S, s, Y, y). The multiple of such a union of gametes is sixteen, and indicates that there will arise an F_2 generation of zygotes with sixteen dihybrid characters in a definite proportion. The actual composition of the F_2 zygotes of this dihybrid may be represented by the following checkerboard diagram:

		♂	♀				
		$SsYy \times SsYy$					
		Male gametes—		SY	Sy	sY	sy
Female gametes	SY	$SSYY$	$SSYy$	$SsYY$	$SsYy$	$ssYY$	$ssYy$
	Sy	$SSYy$	$SSyy$	$SsYy$	$Ssyy$	$ssYy$	$ssyy$
	sY	$SsYY$	$SsYy$	$ssYY$	$ssYy$	$ssyy$	$ssyy$
	sy	$SsYy$	$Ssyy$	$ssYy$	$ssyy$	$ssyy$	$ssyy$

F₂ Generation (Showing progeny with a 9-3-3-1 ratio)

In the checkerboard diagram there are evident homozygous (like) characters and heterozygous (unlike) characters. Any zygote with one or both characters for dominant (SS , YY or Ss , Yy) will have the appearance of smooth surface and yellow color. Zygotes with one dominant character in the dihybrid but with 2 or 3 other recessives ($Ssyy$ or $SSyy$) will appear with smooth surface and green color. Likewise, zygotes homozygous for one contrasting character, if that character is recessive, will appear recessive for that particular character regardless of how the second contrasting character appears. For example, if a zygote has the factors for $ssYY$, the character ss is homozygous for wrinkled surface, a recessive character, and homozygous for yellow color, a dominant character; the dihybrid is, then, yellow and wrinkled. If the dihybrid appears with characters $ssYy$, the character is still homozygous for wrinkled surface but heterozygous for yellow color; it will then appear also yellow and wrinkled. Although the result seems to be the same as far as external appearance is concerned, the second contrasting character of Yy , being heterozygous, will not breed true for yellow in succeeding generations. In the like manner, if the first pair of contrasting characters were Ss instead of ss , again there would be a heterozygous character which would not breed true should that zygote be self-fertilized.

With this fact as a basis, the F₂ generation of the cross between varieties of Pea plants having yellow color, wrinkled-surface seeds and plants having green color, smooth-surface seeds bear zygotes in the ratio of 9 plants with dominant characters for both

yellow color and smooth surface; 3 with homozygous characters for the recessive factor of green color but either homozygous or heterozygous for the dominant factor of smooth surface and thus all appearing with green color and smooth surface; 3 with homozygous characters for the recessive factors of wrinkled surface but either homozygous or heterozygous for yellow color; and 1 with both sets of characters homozygous for the recessive green color and wrinkled surface which breed true when self-fertilized. The four classes when grouped as to the appearance of seeds are then as follows:

- 9 Yellow color, smooth surface
- 3 Yellow color, wrinkled surface
- 3 Green color, smooth surface
- 1 Green color, wrinkled surface

GALTON'S LAW OF ANCESTRAL INHERITANCE.—Francis Galton was the founder of the scientific study of inheritance. He concluded from a study of several selected traits of certain families that the two parents contribute between them on the average one-half of each inherited faculty, each of them contributing one-quarter of it. The four grandparents contribute between them one-quarter, or each of them one-sixteenth; and so on, the sum of the series $\frac{1}{2}$ plus $\frac{1}{4}$ plus $\frac{1}{8}$ plus $\frac{1}{16}$. . . being equal to 1.

VARIATIONS

VARIATIONS represent the differences which exist between the individuals of a species. They are those changes in organisms which make them different from their parents or from their species. No two organisms of any species are exactly alike. This is very apparent from the examination of children in the same family or plants reared from seeds of the same fruit. While having similar characteristics in the main, differences may readily be made out whereby we can distinguish the individuals. Examples of kinds of variations are variations in size, symmetry, outline of organs as leaves, the appearance of a new pigment, etc.

Other kinds of variations are (1) *modifications due to differences in environment to which organisms are exposed*; these are

called *fluctuations*. (2) *Variations due to new combinations* of ancestral characters which in some cases were latent in the parents but present in the grandparents or in more distant ancestors. Such instances are called "*reversions*." (3) *Mutations* or sudden and frequently large differences which appear among individuals not of hybrid origin. It is believed that many varieties of economic plants had their origin in seed mutants or "*sports*." The mutations seen in plants reared from seed sports are due to hereditary causes in which there probably occur a combination of parental characters. *Bud mutations* or bud sports sometimes occur on normal plants as in some varieties of fruit trees especially in certain citrus fruits. The Boston Fern, *Nephrolepis exaltata bostoniensis* is a bud sport of *Nephrolepis exaltata*. This form of mutation is due either to the alternation of a simple gene in a certain chromosome or to abnormalities in the distribution of the chromosomes during mitosis. *Continuous variations* are small additions or diminutions of certain parental characters. *Discontinuous variations* are sudden marked variations which arise without the appearance of transitional stages, as for example where variations occur in the normal number of parts, as in the four-leaved clover. De Vries called these kinds of variations *mutations*.

EVOLUTION

Organic evolution is the established fact that all forms of organisms existing today have been derived from others previously existing either by direct descent or common ancestry. It did not originate with Charles Darwin. It was first conceived by Anaximander, Empedocles and other ancient Greeks. The more important theories as to the method of evolution will now be discussed.

THE LAMARCKIAN THEORY OF USE AND DISUSE.—Jean Baptiste de Lamarck (1744–1829) a French naturalist, propounded his theory on the origin of species in 1809 in his "*Philosophie Zoologique*." He assumed that the environment changes, that use and disuse may alter the character of the individual, that characteristics acquired by an organism in its lifetime are inherited by its offspring. The persistent use of a part of the

body, as the muscles of the arm of a blacksmith, results in greater development of that organ whereas the continued disuse of an organ results in its partial atrophy or degeneration. While these facts about the environment and the use and disuse of organs have been recognized in the lives of many individuals, Lamarck assumed that the changes brought about in an organism through use and disuse of organs were passed on to the descend-

FIG. 486.—Jean Baptiste de Lamarck (1744–1829) who proposed the theory that characters acquired by an individual in its lifetime are inherited by its offspring and so effect evolutionary changes. (Atwood.)

ants. As an illustration of his theory he used the neck of the giraffe. He assumed the early ancestors of the giraffe were grazing animals and that a change in the environment caused them to feed on the leaves of trees. The present long neck was acquired, according to Lamarck, as the cumulative result of the inheritance of slightly greater neck-length brought about in each generation by reaching up persistently to browse upon higher vegetation.

His theory of the inheritance of acquired characters is largely rejected by biologists of the present day because of the lack of convincing evidence.

THE DARWINIAN THEORY OF NATURAL SELECTION.—Charles Darwin (1809–1882) was an English naturalist, who made many observations on the British exploring expedition that circumnavigated the globe on H.M.S. Beagle from 1831–1836. As an outcome of this experience and other later observations he worked out a theory of evolution which he published in 1859 in a book entitled “The Origin of Species by Means of Natural Selection,” or “The Preservation of Favored Races in the Struggle for Life.”

Darwin’s Theory is based on the following fundamental propositions:

1. **PRODIGALITY OF NATURE.**—Living organisms are enormously fertile, yet the total number is approximately stationary. It seems well established that many more offspring are produced by plants and animals than can actually survive. A

very small proportion of the seeds and spores of plants actually germinate and averagely few of the seedlings attain maturity.

2. STRUGGLE FOR EXISTENCE.—Owing to the production of individuals far beyond the earth's facilities to support them, there is a struggle for existence. An intense competition is con-

FIG. 487.—Charles Darwin. (*Gager.*)

stantly taking place in nature for food, water and light among individual plants and animals. The struggle is unequal because of the variability in structures, activities and close relations.

3. ORGANISMS AND SPECIES DIFFER IN RESPECT TO THEIR FITNESS FOR A GIVEN ENVIRONMENT.—Obviously, a water plant is unfitted to live in dry soil and a dry soil plant unfitted to live in the water.

4. VARIATION AND HEREDITY.—The individuals of a species are not all alike and among the offspring of the same parents there are always at least slight variations. These variations are continuous or fluctuating in character.

5. NATURAL SELECTION.—Through the struggle for existence there is a natural selection which results in the survival of the fittest. Darwin argued that in this biological competition some organisms and species possess advantageous variations and are

FIG. 488.—August Weissmann.
(Atwood.)

consequently preserved while the less fortunate perish. The favorable characters are handed down to the descendant, generation after generation, so long as they continue to be advantageous. Thus, the weak and poorly adapted organisms are eliminated while the forms most in harmony with their environment survive.

6. ORIGIN OF SPECIES.—According to Darwin, if certain members of a given group adjust themselves to the conditions of life through variation in one direction and others in other

directions, then divergence among the forms through the continued action of Natural Selection generation after generation must in the course of time be sufficient to make them rank as separate species.

WEISSMANN'S GERM PLASM THEORY.—August Weissmann (1834–1914), a German biologist, developed a theory of heredity based upon the principle of the continuity of the germ plasm. While this principle was recognized earlier by Owen (1849), Galton, Nussbaum and others, it remained for Weissman through a series of brilliant essays and a book on the Germ Plasm (1892) to bring it forcibly to the attention of scientific world.

Weissmann vigorously denied the inheritance of all forms of acquired characters. He believed (1) that the cells of an

organism are of two kinds, the body or somatic cells containing somatoplasm and the germ cells containing germ plasm, (2) that the germ cells reside in the body but are not part of the body or soma, (3) that the germ cells arise as direct descendants of the germ cells of the previous generation, (4) that the germ plasm continues from the beginning of life and the soma protects it, (5) that variations arise from combination of different characters in the germ plasm of each parent, (6) that the determiners originate solely in the germ plasm and migrate from the germ cells out into the various parts of the developing body and that differentiation of the organism is produced in this way, (7) that a new type of organism arises only in consequence of a changed type of germ cell.

Weissmann's ideas about variation were just the reverse of those held by Lamarck and Darwin who believed that variations first originate in the body and are thence passed on to the germ cells. He stressed the importance of heredity in evolution but almost neglected environment.

FIG. 489.—Hugo De Vries, author of the theory of mutation as a method of evolution. (After Walton and Foss.)

DE VRIES MUTATION THEORY.—Professor Hugo De Vries, a Dutch botanist, in 1901, advanced the theory of evolution through sudden abrupt variations. He asserted that new species arise suddenly and become fully established from a parent form which may continue to live side by side with the new form. He called these sudden variations, *mutations*. De Vries reached his conclusions from the striking variations he discovered in successive generations of the Evening Primrose (*Oenothera lamarckiana*), a species he found growing wild, in 1886, in a waste field about Hilversum in Holland. In experimenting on the Evening Primrose in his Amsterdam gardens, he found the mutants when self fertilized proved stable, and when crossed behaved according to the Mendelian laws. De Vries' principal

conclusions have been verified by many subsequent investigators and his methods extended to the discovery of mutants among many species of plants and animals.

CHROMOSOME MUTATIONS.—Since De Vries' discovery, cytological studies have been applied to the study of mutations. These have shown that many mutants possess new chromosome numbers throughout their structure. For instance, *Oenothera lamarckiana* possesses 14 chromosomes while one of its mutants, *Oenothera lata*, has 15.

POLYPLOIDY.—The normal somatic cells of the bodies of higher plants and animals have two sets of chromosomes (diploid). *Polyploids* are forms having multiples higher than two, *e.g.* three or more chromosome sets. Polyploids have been found to originate by hybridization followed by chromosome doubling. Extreme temperatures are claimed by Sax and others to produce doubling of the chromosomes. Polyploidy has been found to account for many mutants in wild plants as the chrysanthemums, maples, roses, apples, cherries, hawthorns, erigeron and rumex and in cultivated plants including wheat, oats, tobacco, sugar cane, dahlias, etc.

THE MODERN THEORY OF THE GENE.—The current theory had its origin in the work of Thomas H. Morgan, an American geneticist, who discovered that the fruit fly, *Drosophila melanogaster*, was in a mutating condition. About 400 mutant characters have been identified in this insect. Morgan showed that the genes can be linked into four groups that correspond to four pairs of homologous chromosomes in the germ cells. The location of the characters or their determiners or genes on the chromosomes has been worked out by Morgan and his associates, and their experiments have shown that genes may cross over from one homologous chromosome to another. They have further shown that genes displaying linkage with each other are located in the same pair of chromosomes, that the substance of the chromosomes binds the genes to each other and causes one factor of a specific character to be inherited along with another gene on the same chromosome.

Several explanations have been advanced concerning the causes of gene mutations or the appearance of new hereditary

characters. Gates suggests they may be environmental, as for example the lowering of the temperature during periods of germ cell maturation. Others have suggested internal causes, such as the weakness of attraction between homologous chromosomes and the consequent failure of synapsis.

The rate of mutations has been slightly increased in *Drosophila* by Muller through raising the temperature of cultures of this insect. In 1926 he showed that the X-ray had the power to alter and rearrange the genes. By exposing adult *Drosophila* flies to powerful X-rays he greatly accelerated the rate of the mutations, increasing these by up to 150 per cent.

It is now thought probable by many geneticists that genes located in the chromosomes represent the sole vehicle of inheritance and that heritable variations can come only from changes in the genes. These changes represent mutations.

Appendix I

THE MICROSCOPE

A MICROSCOPE is an optical instrument, consisting of a lens, or combination of lenses, for making an enlarged image of an object which is too minute to be viewed by the naked eye.

Microscopes are of 3 principal kinds, viz.: simple, dissecting and compound.

THE SIMPLE MICROSCOPE

This consists simply of a convex lens or several combined into a system and appropriately mounted. A good example of a sim-

FIG. 490.—A dissecting microscope. Description in text.

ple microscope is a reading glass. This type of simple microscope is valuable in field work, in the examination of dried herbarium material or the external characters of crude drugs, where only a low magnification of the object is required.

THE DISSECTING MICROSCOPE

But when flowers or other plant parts are to be dissected, it is necessary to have both hands free. To meet this need various

forms of stands have been devised which have been combined with an arm and lens to constitute what are known as "Dissecting Microscopes." One of the simpler forms of these is shown

FIG. 491.—Zeiss stereoscopic dissecting microscope with built in illuminating equipment.

in Fig. 490. It consists of a metal stand with convenient hand rests; a glass stage plate, full size, on which the object to be dissected is placed; a double faced mirror adjustable for reflecting light from various angles, the concave face for concentrating all of the light at the plane of the object. A jointed lens arm is carried

on a rack-and-pinion focusing adjustment having ample range to suit a variety of thicknesses of objects. A black-and-white metal stage plate serves for use as a contrast background. A more elaborate type of dissecting microscope of the twin objective binocular type is shown in Fig. 491. This gives erect images and a stereoscopic effect.

THE COMPOUND MICROSCOPE

A. ITS CONSTRUCTION:

The principal parts of a compound microscope are:

1. The *base*, generally horseshoe shaped, which rests on the table.

2. The *pillar*, an upright bar, which is attached to the base below, supports the rest of the instrument and incorporates the inclination joint.

3. The *stage*, a horizontal shelf upon which is placed the preparation or slide to be examined. The stage is perforated in the center for transmitting light reflected up by the mirror. On the stage are two clips for holding the glass slide.

4. The *mirror*, situated below the stage, by which the light is reflected upward through the opening in the stage. The mirror generally has two faces; the one is plane for initial light intensity, the other concave for concentration of light on the object.

5. The *diaphragm*, inserted in the opening of the stage or attached to its lower face, and used to regulate the amount of light reflected by the mirror.

6. The *body tube*, a cylinder which holds the *draw tube* and *lenses* and moves up and down perpendicularly above the opening in the stage. The tube is raised or lowered either by sliding it back and forth with a twisting movement or by a *rack and pinion* mechanism. The latter is called the *coarse adjustment* and is used for finding the focus.

7. The *fine adjustment*, a micrometer screw back of the tube, which, on being turned, produces a very slow motion of the entire framework which holds the body tube. It permits exact focusing of the higher power lenses.

8. The *oculars* or *eyepieces* which slip into the upper end of the *draw tube*. Each of these consist of two plano-convex lenses, the

lower one being the larger and known as the *collective* or *field lens* because it increases the field of vision. The upper or smaller lens is called the *eye lens*. It magnifies the image formed by the objective. Midway between the field and eye lens is a *perforated diaphragm*, the object of which is to cut out edge rays from the image. It determines the size of the field of view.

FIG. 492.—Illustrating the parts of a compound microscope.

According to the system adopted by the maker, oculars are designated by numbers, as 1, 2, 3, 4, etc., or by figures which represent focal lengths, and by magnification numbers, as $7\times$, $10\times$, etc.

9. The *objectives*, which screw into the bottom of the body tube or nose piece. They consist of a system of two, three or more lenses, some of which are simple, others compounded of a

convex crown lens and a concave flint lens. Objectives like oculars are usually designated by fractions as $\frac{1}{12}$, $\frac{1}{6}$, $\frac{2}{3}$, etc., or in millimeters, as 2 mm., 4 mm., 16 mm., which represent focal

FIG. 493.—Diagram illustrating optics of a compound microscope in use. F_1 , Upper focal plane of objective; F_2 , Lower focal plane of eyepiece; Δ , Optical tube length = distance between F_1 and F_2 ; O_1 , object; O_2 , real image in F_2 , transposed by the collective lens, to O_3 , real image in eyepiece diaphragm; O_4 , virtual image formed at the projection distance C , 250 mm. from EP , eyepoint; CD , condenser diaphragm; L , mechanical tube length (160 mm.); 1, 2, 3, three pencils of parallel light coming from different points of a distant illuminant, for instance, a white cloud, which illuminate three different points of the object. (Courtesy of Bausch and Lomb Optical Co.)

lengths or numbers, as 97, 45, 10, etc. which indicate the magnification of the objective alone.

The smaller the number in millimeters or fraction representing the focal length of an objective, the greater is its magnifying power.

The function of an objective is to take in the divergent light from the object and change it into convergent rays that meet in the upper end of the microscope tube in the front focal plane of the ocular forming there a magnified image of the object. This image is further magnified by the lens of the human eye.

FIG. 494.—Sectional view showing component elements of an apochromatic, high power objective.

OBJECTIVES are either *dry* lenses or *immersion* lenses. If an air space be present between the tip of the objective and the object, the lens is called a *dry* one; if a liquid is present between the tip of the objective and the object, the lens is called an *immersion* lens. If this liquid be oil, the objective is called an *oil immersion objective*; if water, a *water immersion objective*.

Dry objectives are of low, medium and high powers. Dry medium and dry high power objectives are the best types to use on objects in air.

When examining objects in water, glycerin or balsam with the dry objective, the cover glass should be slightly under 0.17 mm. in thickness for best results.

The important properties of the microscope objective are: (1) *Numerical aperture (N.A.)* which measures the amount of light received from the object and represents the index of resolving power. (2) *Resolving power* is that by which two small elements in the structure of an object and which are only a short distance apart are distinctly separated.

The higher the N.A. the greater the resolving power of the objective and the finer the detail it can reveal.

N.A. equal $N \sin u$
wherein

n = the lowest refractive index that appears between the objective and the front of the objective.

u = half the angular aperture of the objective.

FIG. 495.—A new type of objective in a correction mount. Turning the correction collar varies the distance between the two upper lenses and the two lower lenses fixed in the body of the mount.

The degree of angle of illumination directly affects resolving power up to a maximum resulting from filling the aperture of the objective with light.

FIG. 496.—A modern monocular research microscope with revolving nose piece, side fine adjustment and substage.

A 4 mm., 0.85 N.A. objective will resolve lines separated by distance ranging between 0.00062 and 0.00031, dependent upon the aperture employed. For a 4 mm., 0.65 N.A. objective the limiting values are 0.00081 and 0.000405.

Two objectives of the same equivalent focal length (E.F.) and the same N.A. should show the same illuminated area in the back lens, when viewed without an eyepiece and illuminated with the widest cone of light they can take in.

It also is evident that an objective cannot show its full efficiency if it is not used with a condenser of an N.A. large enough to fill the back of the objective with light.

DEPTH OF FOCUS.—Depth of focus (known also as depth of sharpness of penetration) depends on the N.A. and the magnification and is inversely proportional to both. The higher the N.A. and the higher the magnification, the less the depth of focus. An increase of the depth of focus, for instance, by insert-

FIG. 497.—Aplanatic condenser (Zeiss)
N.A. 1.4. Focal length 10.5 mm.

ing diaphragms above the back lens of the objective must necessarily decrease the effective diameter of the back lens and thus decrease the N.A., thereby lowering the efficiency of the objective.

To comply with different requirements in this respect the makers offer two 4 mm. objectives, whose depths of focus, when used with the same magnification, are in the ratio of 3:4.

It should be borne in mind that the finer the definition of an objective the more sensitive it is to incorrect focusing and to slight changes of the fine adjustment screw. This means less depth of focus. Objectives that show greater depth of focus than others of the same N.A. cannot be well corrected.

Working distance of an objective is sometimes confused with the equivalent focus. It varies for different powers being disproportionately less for high powers, and is measured from the front of the objective to the upper surface of the cover glass. For instance, the high dry objective (4 mm. e.f. 0.65 N.A.) has a working distance of 0.6 mm.

Some microscopes are fitted up with a *nose-piece*, capable of carrying two, three, or four objectives, which may be revolved into place at the lower end of the body tube. Others have a *condenser* which is employed to concentrate the light upon the

object examined. When using the condenser, employ only the *plane mirror*.

B. ITS USE:

1. Place the microscope on the table with the pillar nearest you.
2. Screw the objectives into the nose piece and slip an ocular into the upper end, if not already on instrument. Turn the lowest power objective into position.
3. Find the light by looking into the ocular (eye piece) and at the same time turning the mirror at such an angle that it

FIG. 498.—Microscope lamp, useful in illuminating opaque objects.

reflects light from the window or lamp up through the opening in the stage to the objective. When opaque objects are to be illuminated, a stronger illumination is required than that usually afforded by an ordinary laboratory lamp or by the light from a window. For this purpose a microscope lamp, such as the Bausch and Lomb No. 4578 (see Fig. 498) is very satisfactory. Mirrors have two faces, a plane and a concave. Use the concave unless employing the condenser, when the plane mirror should always be used.

4. Regulate the quantity of light by the iris diaphragm. If too bright it must be cut off somewhat. The higher powers

require brighter light than the lower; the oil immersion full opening.

5. Place the slide on the stage in a horizontal position with the object over the middle of the opening through which light is thrown from the mirror.

FIG. 499.—Binocular microscope. This is a compound microscope with a binocular body designed to relieve eyestrain for those doing microscopical research work or otherwise engaged for prolonged periods with the microscope. Note the parallel position of the eyepiece tubes which are adjustable for interpupillary distance by means of a compound slide, with the distance indicated on a numbered millimeter scale for future guidance. One tube has a spiral adjustment for correcting any difference in vision existing between the two eyes. The instrument shown is that designated CAE of the Bausch and Lomb Optical Co.

6. With the lower power in position quite close to the object, move the coarse adjustment upward until either the object or small solid particles on the slide appear distinctly,

which means that the lenses are *in focus*. (If two objectives only accompany your instrument, the low power is the shorter one.) The object, if not under the lens, may now be brought into the field by moving the slide back and forth very slowly while looking through the ocular. To improve the focus, slowly turn the fine adjustment screw.

7. To focus with the high-power objective, first find the object with the low power and arrange in the center of the field. Put clips on slide without moving it. Raise the body tube by means of the coarse adjustment. Then turn the high-power objective into position. Lower the body tube carrying the objective until the objective front lens nearly touches the cover glass. A slight movement of the fine adjustment should show the object clearly. Never focus down with the high-power objective while looking through the ocular because of the danger of pressing it into the cover glass and the risk of ruining the delicately mounted lenses.

Accustom yourself to use both eyes indifferently and always keep both eyes open. If right handed, observe with the left eye, as it is more convenient in making drawings.

9. When the *oil immersion objective* is to be used, a small drop of immersion oil (slightly evaporated cedar oil) should be placed on the cover glass directly above the object, and the body tube should be run down with the coarse adjustment until the front lens of the immersion objective enters the drop and comes almost into contact with the cover glass. This should be done while watching the objective. Then look through the ocular and draw the objective up with the fine adjustment until the object comes into focus.

RULES FOR THE CARE OF THE MICROSCOPE

1. In carrying the microscope to or from your table, grasp it firmly by the pillar and hold in an erect position, so that the ocular which is fitted loosely into the draw tube may not fall out and its lenses become damaged.

2. Never allow the dry objective to touch the cover glass or the liquid in which the object is mounted.

3. Never touch the objective or ocular lenses with fingers or cloths.

4. Never change from lower to higher power objective without first ascertaining that the body tube has been raised sufficiently to allow the high-power objective to be slipped into place without injury to the objective or mounts.

5. Never clean the microscope lenses or stand with cloths that have been used for removing surplus of alkali, acid or other reagent from slides.

6. Note whether the front lens of the objective is clean before attempting to use it. If soiled, breathe on the lens and gently

FIG. 500.—Ocular micrometer.

wipe with an old, clean, soft handkerchief or lens paper. If the lens be soiled with balsam or some other sticky substance, moisten the handkerchief or lens paper with a drop of xylol, taking care to wipe it perfectly dry as soon as possible.

7. Do not let the objective remain long near corrosive liquids, such as strong solutions of iodine, corrosive sublimate, or mineral acids. Never examine objects lying in such fluids without putting on a cover glass.

8. Never lift the slide from the stage, but, after raising the objective, slide it off the stage without upward movement.

9. Never allow the *stand* (microscope without lenses) to be wetted with such substances as alcohol, soap, etc., which dissolve lacquer.

10. Keep the microscope covered when not in use.

MICROMETRY

The unit of length used in microscopic measurement is the *micron* (μ) which is one-thousandth part of a millimeter (0.001 mm.) or one twenty-five thousandth part of an inch.

In measuring microscopic objects it is necessary to make use of a micrometer of some kind. That pretty generally used is the *ocular micrometer* (Fig. 500). It is a circle of glass suitable for insertion within the ocular with a scale etched on its surface. The scale is divided to tenths of a millimeter (0.1 mm.), or the entire

surface of the glass may be etched with squares (0.5 mm.), the *net micrometer* (Fig. 501).

STANDARDIZATION OF OCULAR MICROMETER

The value of each division of the *ocular micrometer scale* must be ascertained for each optical combination (ocular, objective, and tube length) by the aid of a stage micrometer.

The *stage micrometer* (Fig. 502) is a slide with a scale engraved on it divided to hundredths of a millimeter (0.01 mm.), in some cases, also, to tenths of a millimeter (0.1 mm.), every tenth line being made longer than intervening ones, to facilitate counting.

METHOD:

1. Insert the ocular micrometer within the tube of the ocular by placing it on the diaphragm of the ocular, and adjust the stage micrometer by placing it on the stage of the microscope.
2. Focus the scale of the stage micrometer accurately so that the lines of the two micrometers will appear in the same plane. Make the lines on the two micrometers parallel each other.

FIG. 501.—Net micrometer, about $1\frac{1}{2} \times$ actual size. The engraved scale is 10 mm. square and is divided into small squares with 0.5 mm. sides.

FIG. 502.—Stage micrometer used to standardize the ocular micrometer. The mm. readings etched on this micrometer indicate the width of the larger and smaller spaces of the stage micrometer scale seen in the center of the stage micrometer.

This can often be done by turning the ocular to the right or left, while looking into the microscope.

3. Make two of the lines on the ocular micrometer coincide with two on the stage micrometer. Note the number of included divisions.

4. Note the known value for each division of the stage micrometer scale which may either be etched on the stage

micrometer or indicated on a label found pasted upon it. If the value indicated is 0.01 mm. ($\frac{1}{100}$ mm.), then each division of the stage micrometer scale has a value of 10 microns; if 0.1 mm. ($\frac{1}{10}$ mm.), 100 microns.

FIG. 503.—Scale of Stage micrometer, magnified about 24 $\times$. Each small space of actual scale has a value of 10 microns, each large space between the longest lines, 100 microns. The length of the entire scale is 1 mm.

5. Multiply the number of included divisions of the stage micrometer scale by the value in microns given for each division and divide the result by the number of included divisions of the ocular micrometer scale. The quotient represents the value of each division of the ocular micrometer scale.

6. Note the optical combination (number of ocular, objective and tube length) used and keep a record of it with the calculated micrometer value. Repeat for each of the combinations.

To measure an object by this method, read off the number of divisions it occupies of the ocular micrometer scale, and express the result in microns by looking up the recorded value for the optical combination used.

Appendix II

HISTOLOGICAL TECHNIQUE

In this chapter the various methods which are commonly employed in the preparation of plant materials for microscopic examination and the reagents used in the examination of the same will be discussed.

MAKING OF SECTIONS

FREE-HAND SECTIONING.—Free-hand sections are usually satisfactory for the general examination of roots, stems, leaves, barks and many fruits and seeds. Material which is fresh may be sectioned at once, but dry material should be well soaked in warm water before using. Fresh material should always be kept moistened in water on account of the danger of cell shrinkage through loss of water. Dried membranous or subleathery leaves should be kept in a moist chamber for at least 24 hours or until sufficiently supple before they are sectioned. Dried woody roots and stems can usually be best prepared for sectioning by soaking segments of these in equal parts of alcohol, glycerin and water for several weeks prior to using. Very hard material like heartwoods, the shells of nuts and seeds, may be softened in solution of caustic potash or ammonia water and then washed free of alkali before sectioning. Fine shavings can then be whittled off of these with a sharp knife. In some cases alternately boiling and cooling the material will suffice.

The object to be sectioned is held between the thumb and finger of the left hand. If tender and flexible, such as a fresh leaf, it must be placed between the two flat surfaces of elder pith before sectioning. A segment of pith about an inch long is halved lengthwise with a sharp knife and a portion of the leaf is held between the halves of pith while the section is cut through pith and leaf. The pith is later separated from the leaf section. Sections through other delicate parts of plants, as buds, slender

roots, etc., may be made in the same way, only a groove should be made in the pith of such size as is necessary to hold the material firmly enough without crushing it. In certain instances, when, because of the smallness of the object and its resistance to cutting, good sections can not readily be made with the aid of pith, a small sized cork stopper can be used with better results. A hole just large enough to prevent the object from slipping is made in the center of the smaller end and the object inserted preparatory to sectioning. The upper surface of the razor is wetted with 50 per cent. alcohol. The razor, which should be real sharp, is held in the right hand and is drawn across the object with the edge toward the student and the blade sliding on the forefinger of the left hand. *The sections should be cut as thin as possible.* Small bits of representative parts of entire sections yield better results when examined under the microscope than complete sections which have been cut too thick. As soon as a number of sections have been cut, they can be transferred to a vessel of water with a camel's hair brush, before they become dry. If the sections are cut from fresh material and are to be stained, they should be placed in 95 per cent. alcohol for at least several minutes prior to staining. If from material preserved in formalin solutions, they should be washed in water.

Surface sections of leaves may occasionally be prepared by stripping off the leaf epidermis. In most instances, however, the leaf epidermis adheres firmly to the subjacent tissue and it becomes necessary to employ different technique. A good practice is to bend the leaf over the index finger of the left hand and hold it firmly between the index finger and the third finger on the one end and with the thumb on the other. Another method is the following: Place a representative portion of the leaf not over $\frac{1}{4}$ inch square in a watch crystal with 10 per cent. to 25 per cent. chloral hydrate solution, warm gently for several minutes, according to the texture of the leaf, transfer to a clean slide, add a drop or two of chloral hydrate solution, cover with cover slip and exert pressure on the cover slip with a rotary movement, describing the figure 8. The epidermis will thereby usually separate from the underlying tissues. It sometimes becomes necessary to study large areas of the epidermis of

leathery leaves. Such leaves should be prepared by boiling in 10 to 20 per cent. aqueous solution of either sodium or potassium hydroxide until the epidermis puckers up in the form of blisters when the epidermis can be lifted off by means of fine forceps or dissecting needles and floated on the slide.

SECTIONING IN PARAFFINE OR CELLOIDIN. When it is necessary to study the microscopic structure of very delicate plant parts, superior results can generally be obtained by imbedding the material in paraffine or celloidin (see pp. 692 and 699), which is subsequently hardened, and sectioned by means of a sliding or rotary microtome.

KINDS OF SECTIONS

1. A *transverse* or *cross-section* is one made horizontally through the object, hence its plane lies at right angles to the long axis.

2. A *radial-longitudinal* section is one which is made parallel to the long axis of the object in such a way that it lies in the plane of the radius.

3. A *tangential-longitudinal* section is one made parallel to a plane tangent to the cylinder. This type of section is therefore prepared by cutting parallel to the outer long surface.

4. A *surface* section is one made by cutting or stripping off the outer protective covering of the plant organ.

FIG. 504.—Showing the planes in which sections are cut, A, transversely; B, longitudinal radially; C, longitudinal tangentially. (After Stevens.)

MICROTOMES

Microtomes are instruments employed to facilitate the cutting of sections of organic tissues. The three most commonly used types are the hand, sliding and rotary microtomes.

HAND MICROTOME.—This type is shown in Fig. 505. If the object is sufficiently hard to bear the strain, it is placed directly in a clamp of the upper end of the tube that is tightened by the

screw seen on the side of the tube, or it may first be inclosed in elder pith or cork and then clamped in. The object to be sectioned is raised a little at a time through the hole in the glass plate at the top by turning the finely graduated feed near the base of the tube. The section razor is then laid flat on the glass plate and pulled across the object with a long sliding motion.

FIG. 505.—Hand microtome.
Description in text.

The upper surface of the razor blade is kept wet with 50 per cent. alcohol and after several sections have been cut they can be swept by the finger or camel's hair pencil to a dish of water. Each division of the feed represents 10 microns, so that the thickness of sections desired can be regulated by moving the feed, accordingly, just before each stroke of the razor.

SLIDING MICROTOME.—This type of microtome (see Fig. 506) is adapted for cutting all kinds of sections. It consists of an iron supporting frame of horizontal and upright portions. The horizontal base rests on the table and is hollowed out to accommodate a drip pan that can readily be removed and cleaned.

The front of the upright portion exhibits a frame which accommodates a sliding feed mechanism to which is attached the object carrier. The top of the upright portion shows a flat bed which carries a solid iron block which can be readily slid along the bed when the latter is lubricated with paraffin oil. The upper surface of the block is grooved to accommodate the thumb screw. The microtome knife consists of a blade portion that is flat on its lower and hollow ground on its upper face. The latter is placed in the clamp on the top of the microtome, and its position adjusted. Sections of woody material can be cut directly on this microtome and placed in dilute alcohol. When paraffine sections are cut, the cutting edge of the knife should be parallel to the motion; but when celloidin sections are desired the knife must be set at an

oblique angle to the frame and drawn across the block with a long sliding motion. The knife and the top of the celloidin block must be constantly kept wet with 80 per cent. alcohol.

FIG. 506.—Spencer sliding microtome.

FIG. 507.—Rotary microtome. The feed mechanism is covered to protect the wearing parts from dust. (Courtesy Spencer Lens Co.)

The object is placed in the object carrier and clamped in. By means of the graduated disk at the base of the feed mechanism the thickness, in terms of microns, is regulated after each stroke of the razor.

ROTARY MICROTOME.—When paraffin ribbons are desired, especially for the study of serial sections of material, the rotary microtome surpasses by far the efficiency of the sliding type of instrument. The Spencer Rotary Microtome No. 820 is shown in Fig. 507.

REAGENTS AND STAINS AND METHODS OF PREPARATION OF MATERIALS

The following reagents will be found of value in the microscopical examination of plants, powdered vegetable drugs and foods: (additional reagents and formulas for their preparation are presented elsewhere in the text in connection with methods of examination).

ACETIC ACID.—This reagent should contain not less than 36 per cent. nor more than 37 per cent. of pure acetic acid. It is used for various operations such as distinguishing between calcium carbonate which dissolves in it with effervescence and calcium oxalate which is insoluble in it. A 2 to 3 per cent. solution is occasionally employed as a mounting medium for microscopic plants.

ACID ALCOHOL.—Mix $\frac{1}{10}$ cc. hydrochloric acid with 100 cc. of 70 per cent. alcohol. This is employed in removing the excess of stain from overstained material. When used, it should be thoroughly washed out with diluted alcohol or water.

ALCOHOL.—Useful in the form of various percentage solutions for dehydrating, preserving, hardening and solvent purposes. Absolute alcohol (Dehydrated alcohol) should contain not less than 99 per cent. by weight of ethyl hydroxide and must be kept in well stoppered bottles because of its property of rapidly absorbing moisture from the air and so becoming reduced in strength. It should always be used before cedar oil or xylol in making a balsam mount, but if clove oil is used for clearing instead of cedar oil or xylol, 95 per cent. alcohol may be employed.

ALCANNA TINCTURE.—Macerate 20 Gm. of alkanet root for a week in 100 cc. of 90 per cent. alcohol, boil for several minutes and cool. Dilute with an equal volume of water just before

using. This reagent imparts a red color to fixed oils but may stain other contents as well.

AMMONIA WATER.—An aqueous solution of gaseous ammonia containing in each 100 cc. not less than 9 Gm. and not more than 10 Gm. of NH_3 . Useful in clearing highly colored vegetable powders, such as ground roasted coffee, when mixed with equal parts of peroxide of hydrogen. Material should be macerated in this mixture within a tightly corked vial over night or longer and washed with water before examination.

AMMONIA WATER, STRONGER.—This is an aqueous solution of NH_3 containing not less than 27 per cent. nor more than 29 per cent. by weight of ammonia gas. It is employed in the preparation of ammonia water and cuoxam.

ANILINE CHLORIDE.—A saturated aqueous solution acidified with hydrochloric acid is useful in staining lignified walls which are colored a golden yellow.

ANILINE SULFATE.—A saturated aqueous solution is employed as a test for lignified walls. Sections are mounted in this reagent and a drop of sulfuric acid added. A yellow color is imparted to lignified walls.

ANILINE BLUE.—A saturated aqueous solution is useful in staining sieve tubes. Sections should be placed in this solution for 24 hours and then washed to remove excess of stain. A one per cent. solution in 90 per cent. alcohol is used in double-staining with safranin. It stains cellulose walls blue.

BISMARCK BROWN (ANILINE BROWN).—A saturated aqueous solution is useful in double staining with gentian violet. A dilute aqueous solution made by dissolving 0.2 Gm. of Bismark Brown in 100 cc. of distilled water is of value in staining transparent tissues.

CHLORAL HYDRATE.—A solution prepared by dissolving 25 Gm. in 10 cc. of water is an excellent clearing agent. It dissolves starch, resin, protein, chlorophyll, etc., and causes more or less expansion of shrunken cells.

CHLORAL IODINE.—Saturate the previously prepared chloral hydrate solution with iodine by adding crystals of resublimed iodine to it and shaking. Useful for the detection of minute traces of starch or small starch grains, which are colored blue.

CHLORZINCIODINE SOLUTION.—Dissolve 25 Gm. of anhydrous zinc chloride and 8 Gm. of potassium iodide in 8.5 Gm. of water and add iodine crystals to saturation. Keep in dark colored containers. Useful in distinguishing between cellulose and lignified walls. Cellulose walls are colored blue or violet and lignified walls yellow when mounted in this solution. It causes starch grains to swell and colors them blue.

CHROMIC ACID SOLUTION.—Dissolve 5 Gm. of chromic acid in 45 cc. of diluted sulfuric acid (U.S.P.). A valuable reagent for separating the component cells in sections of plant organs. The sections are placed in this solution in a watch-glass and removed one by one, as required, in the course of 10 to 15 minutes to a slide, washed with a few drops of water and subjected to pressure with a glass rod.

CORALLIN SODA SOLUTION.—Dissolve 15 Gm. of carbonate of soda in 35 Gm. of distilled water and add sufficient corallin to produce a pale bright pink color. This reagent stains the callus plates of sieve tubes pink. It should be freshly prepared as needed.

EOSIN.—An aqueous solution prepared by dissolving 1 Gm. of eosin in sufficient water to make 100 cc. is useful in the staining of cytoplasm, aleurone grains and other cell contents. If permanent mounts are desired, transfer to 1 per cent. acetic acid for a minute, wash with water, then to glycerin or through glycerin to glycerin jelly.

A saturated alcoholic solution of eosin (using absolute alcohol as solvent) is employed when eosin-stained material is to be mounted in balsam. Acetic acid is omitted in this case and the sections passed from alcoholic eosin to a clearing agent as cedar oil or xylol, thence into balsam.

ETHER.—Useful in defatting sections or powders of oil seeds.

FAST GREEN.—A one per cent. solution in 95 per cent. alcohol is useful in staining tissue with cellulose walls green.

FEHLING'S SOLUTION. (Alkaline Cupric Tartrate T.S.).

Copper Solution.—Dissolve 34.66 Gm. of small, uneffloresced crystals of cupric sulfate in sufficient distilled water to make the mixture measure 500 cc. Keep in well stoppered bottles.

Alkaline Tartrate Solution.—Dissolve 173 Gm. of crystallized potassium and sodium tartrate and 50 Gm. of sodium hydroxide in sufficient distilled water to make the solution measure 500 cc. Keep in rubber-stoppered bottles.

Mix exactly equal volumes of the two solutions when required. This solution yields a precipitate of cuprous oxide with reducing sugars.

FERRIC CHLORIDE SOLUTION.—A 1 per cent. solution of ferric chloride in distilled water is used to detect the presence of tannin. It yields a dark green to bluish-black precipitate with tannin.

FUCHSIN (Basic).—Dissolve 1 Gm. of fuchsin in 100 cc. of 95 per cent. alcohol and add an equal volume of water. This solution will impart different tints of red to the various tissues. It has been found especially useful in staining sections which are to be photomicrographed.

FUCHSIN, ACID.—A 0.2 per cent. aqueous solution is useful in staining plastids and the phytoglobulins of aleurone grains. A 1 per cent. solution in 50 per cent. or 70 per cent. alcohol is occasionally employed in conjunction with methyl green or fast green in the double staining of tissues of plants.

GENTIAN VIOLET (METHYL VIOLET).—An aqueous solution, made by dissolving sufficient of the dye in 1 per cent. acetic acid solution to impart a deep violet color, is useful in fixing and staining pollen grains, spores and other fresh plant material. A 1 per cent. solution in distilled water is used in conjunction with Safranin and Orange G. solutions in staining material used for the study of nuclear divisions.

GLYCERIN, DILUTE.—Equal parts of pure glycerin and distilled water. Used as a mounting medium.

GLYCERINE GUM.—Dissolve 20 Gm. of gum arabic and 20 Gm. of glycerine in 15 cc. of water. Useful for fixing small fruits or seeds on pith before sectioning.

HÆMATOXYLIN, DELAFIELD'S.—Used for double-staining, see p. 701.

HÆMATOXYLIN, HEIDENHAIN'S.—This consists of two solutions, a mordant and a stain. The mordant (Solution I) is prepared

by dissolving 3 Gm. of Ferric Ammonium Sulfate in sufficient distilled water to make 100 cc. The stain (Solution II) is prepared by dissolving 0.5 Gm. of hematoxylin in 100 cc. of distilled water. The mordant and the stain are used separately on the sections or tissues to be treated.

HYDROCHLORIC ACID.—The concentrated C.P. acid is used with an equal volume of Phloroglucin Solution in determining lignification. Dilute solutions are employed for neutralizing alkaline solutions, etc.

IODINE WATER.—Add as much iodine to distilled water as it will dissolve. Stains starch pale blue to purplish-blue.

IODINE AND POTASSIUM IODIDE SOLUTION.—Dissolve 1 Gm. of iodine and 3 Gm. of potassium iodide in 50 cc. of water. Keep in glass stoppered bottles. This reagent stains cellulose, lignified and corky walls and proteins yellow and starch blue to bluish-black.

LABARRAQUE'S SOLUTION.—Triturate 50 Gm. of fresh chlorinated lime with 250 cc. of water so as to form a uniform mixture. Dissolve 35 Gm. of monohydrated sodium carbonate in 250 cc. of hot water, and add this solution to the chlorinated lime mixture in a suitable vessel. Stir or shake thoroughly. (Warm if solution gelatinizes.) Transfer the mixture to a wetted muslin strainer, returning the first portion until the liquid passes through clear, and when no more liquid drains from it, wash the precipitate with sufficient water to make the product weigh 500 Gm. Keep in amber colored, glass stoppered bottles in cool place. It gradually loses in strength on standing. This reagent is used for bleaching highly colored sections and powders. The materials bleached should always be washed with water before mounting.

LIGHT GREEN.—This is used as a combination stain with safranin. Dissolve 1 Gm. of light green in 100 cc. of clove oil; 25 cc. of absolute alcohol and 75 cc. of clove oil may be used in place of pure clove oil, if the light green is found to precipitate out when xylol is subsequently employed as a clearing agent.

NAPHTHOL SOLUTION.—Dissolve 5 Gm. of alpha-naphthol in 50 cc. of alcohol. Used in conjunction with concentrated sulfuric acid for the detection of inulin. A violet coloration is

produced if sections containing inulin are first treated with a drop or two of this reagent and then, after a minute, a drop or two of sulfuric acid be added.

PETROLEUM ETHER.—Useful in clearing sections or powdered material of oil.

PHLOROGLUCIN SOLUTION.—Dissolve 1 Gm. of phloroglucin in 50 cc. of 95 per cent. alcohol. This solution gradually darkens and loses strength with age. It is unfit for use after three or four months. The lignified walls of elements first treated with several drops of this solution and subsequently with a drop or two of concentrated hydrochloric acid are colored red.

POTASH SOLUTION.—Several percentage solutions are employed in histological technique for varying purposes. The most commonly used one is the 5 per cent. aqueous solution which is used for clearing sections of starch, protein and tannin. Ten to 20 per cent. aqueous solutions are used for the purpose of separating the epidermis of leathery leaves. The leaf material is boiled in the strong potash solution until the epidermis puckers up in the form of blisters, when it should be removed to a slide and the epidermis lifted off by means of a fine forceps or dissecting-needles. *A 2 per cent. solution is used for the isolation of laticiferous vessels from plant organs.* Pieces of stems, barks or roots containing these elements are placed in a test tube and digested on a water-bath for a half hour or longer until the parenchymatous tissues are sufficiently soft. The potash solution is then poured off and the material washed with water, transferred to a slide and the woody or other undesirable parts removed with the aid of dissecting needles. The softer parts are then stained with iodine and potassium iodide solution; a cover slip is adjusted, the slide placed on stage, and the laticiferous tissues are identified by the presence of branching and anastomosing tubes stained yellow to yellowish-brown with granular contents. A 1.5 per cent. solution is used for the digestion of ground cereals or wheat middlings for the purpose of clarification preparatory to microscopical examination of these materials.

POTASSIUM FERROCYANIDE.—A 10 per cent. solution is employed in studying the localization of alkaloids in plant materials. Sections of fresh material are fixed, in order to pre-

vent postmortem filtration of the alkaloids, by placing them in this solution for 5 minutes, and the excess of this reagent washed out with water. The $K_4Fe(CN)_6$ solution precipitates the alkaloids in the cells which can then be colored by adding a drop or two of $FeCl_3$ solution. Clearing of the sections may then be effected with chloral solution.

RUTHENIUM RED AND LEAD ACETATE SOLUTION.—This is an excellent reagent for the detection of mucilage, to which it imparts a *pink color*. Add sufficient ruthenium red to a 10 per cent. solution of lead acetate until it shows a wine-red color. The lead acetate solution should be prepared with distilled water and the ruthenium red added to quantities of it as needed, as the reagent is unstable. In pharmacognosy, it is especially useful in detecting powdered cacao shells in powdered cocoa and chocolate, since the cacao shells are the only portions of the cacao seed which contain mucilage.

SAFRANIN.—Several kinds of safranin solutions are in use, viz., Alcoholic Safranin, Anilin Safranin and Aqueous Safranin. *Alcoholic Safranin* is prepared by making a saturated solution of safranin (alcohol soluble) in 95 per cent. alcohol and then adding an equal volume of water. *Anilin Safranin* is made by mixing equal parts of saturated alcoholic solution of safranin with aniline water. *Aqueous Safranin* represents a 1 per cent. solution of safranin (water-soluble) in water. Safranin solutions per se stain both protoplasmic contents and cell walls red. When properly used in contrast staining they stain all suberized walls and lignified walls red.

SCHWEITZER'S REAGENT (Ammoniated Cupric Oxide Test Solution).—This is a valuable reagent for dissolving cellulose. It is prepared by dissolving 10 Gm. of copper sulfate in 100 cc. of distilled water, adding sufficient 20 per cent. solution of sodium hydroxide to precipitate the copper hydroxide which is collected on a filter and washing the precipitated copper hydroxide free from sulfate with cold distilled water. The precipitate, which must be kept wet during the entire process is then dissolved in the minimum quantity of ammonia water necessary for complete solution.

SULFURIC ACID.—Concentrated sulfuric acid, containing from 93 to 95 per cent. of H_2SO_4 is useful in the detection of suberized or corky walls. These resist its action, while lignified and cellulose walls are completely dissolved by it. This acid as well as dilutions of it with water are also employed *per se* or with various chemicals in a number of microchemic tests.

Additional reagents and stains are discussed under appropriate headings in this and other chapters of the text.

THE TECHNIQUE OF MAKING A TEMPORARY MOUNT

1. Place a drop or two of water (or reagent) in the center of a clean glass slide.

2. With the aid of a forceps take the section or very small quantity of the material to be examined and spread it on the drop of water.

3. Place a clean cover-glass over the material. In placing the cover-glass, do not drop it flat upon the drop of water, but place one side of it down first and allow it to squeeze the water along under it.

4. Keep the top of the cover-glass dry.

When filamentous algæ or molds are to be examined, the material tends to cling together and must be carefully separated, in the drop of water, with dissection needles before the cover-glass is placed over the material. In case a coarse ground drug is to be mounted, the coarser particles should be first crushed in the water on the slide and subsequently teased apart with dissection needles.

Care should always be taken to see that the water or mounting medium used is not contaminated with foreign substances. This can best be practiced by examining the mounting medium under the microscope before the material to be examined is placed in it.

THE TECHNIQUE OF MAKING PERMANENT MOUNTS

1. **THE MOUNTING MEDIUM.**—When a microscopic object is to be preserved permanently it must be kept from decaying and the fluid in which it is placed must be kept from evaporating. These conditions can be met by adding an antiseptic (2 per cent.

acetic acid, or formaldehyde) to the water used in mounting and carefully sealing the cover glass with asphaltum or zinc white. As a rule, a better way is to use a mounting medium that will not evaporate, *e.g.*, glycerine, glycerin gelatin or Canada balsam. These fluids have a high refractive index and so render the objects penetrated by them more transparent. This quality is generally an advantage, but for objects already almost transparent it is quite the reverse. Glycerine has the disadvantage of always remaining soft, so that the mount may at any time be spoiled by careless handling. Glycerin-gelatin has the advantage of mixing readily with 50 per cent. glycerin in which the object should be placed before being mounted in this medium. It should be warmed on a water bath before using and the cover glass applied quickly after it is placed on the specimen. It cools rapidly and constitutes the quickest and simplest means of effecting a durable permanent mount. Its disadvantage is due mainly to its jelly like consistency which is frequently responsible for damaged mounts when the cover-glass above the preparation is too greatly strained. Canada balsam slowly becomes solid, so that the mount is exposed to no accident short of actual breakage. Balsam has the disadvantage of being non-miscible with water, so that before it can be used the object must be carefully dehydrated. Even after this is done, and the object lying in absolute alcohol, a clearing oil such as cedar oil, clove oil or xylol must be used as an intermediate agent between alcohol and balsam.

2. STAINING.—For two reasons it is generally better to stain plant tissues before mounting. Transparent tissues may become almost invisible in glycerine, glycerin-gelatin, or balsam, and different tissues take a stain differently. This being the case it becomes possible to stain one tissue and not another, or one tissue with one stain and another in the same section with a different stain (*double staining*), so that the different parts may be brought out like areas on a colored map. The most common stains are hæmatoxylin derived from logwood, and various anilin stains—safranin, fuchsin, eosin, crystal violet, fast green, methyl violet, light green, iodine green, methyl-green, malachite green, etc.

A number of useful methods of staining and mounting plant materials will now be discussed. For additional methods the

reader is referred to "Stain Technology," published by the Commission on Standardization of Biological Stains, Geneva, N. Y.

METHODS FOR THE PREPARATION OF CANADA BALSAM MOUNTS
No. 1. SAFRANIN AND FAST GREEN

1. Safranin. By means of an eye-dropper put 10 to 15 drops of a strong alcoholic solution of safranin into a watch glass of distilled water. Allow the sections to lie in this mixture until deep red.

2. Transfer the sections to another watch glass containing tap water and rinse off excess of stain.

3. Dehydrate and differentiate in 50 per cent. alcohol.

4. Dehydrate in 95 per cent. and absolute alcohol (1 minute in each).

5. Fast Green. Place 5 drops of a solution of fast green (fast green, 1 Gm., absolute alcohol 25 cc., clove oil, 75 cc.) in a watch glass of absolute alcohol. Allow the sections to remain in this solution until the green color stains the softer tissues properly.

6. Clear sections in xylol.

7. Mount in Canada balsam dissolved in xylol.

8. Label slide, indicating subject and stain combination.

This method has given excellent results in staining sections of a wide variety of fresh and macerated roots, stems and leaves which have been cut free hand or on the sliding microtome.

No. 2. SAFRANIN AND LIGHT GREEN

1. Stain object with either alcoholic safranin or anilin safranin (1 per cent. solution in 50 per cent. alcohol) for 1 to 24 hours or by gently warming object in the safranin solution until vapors arise.

2. Wash out excess of stain with 50 per cent. alcohol.

3. Dehydrate with 95 per cent. alcohol for two minutes.

4. Further dehydrate by placing material in absolute alcohol for 20 to 30 seconds.

5. Stain with a saturated solution of light green in clove oil for from 3 to 30 minutes.

6. Clear in xylol for 1 minute. Blot up excess from around edge of section.

7. Mount in Canada balsam.

8. Label slide.

The safranin will impart a red color to the suberized and lignified walls, while the light green will stain cellulose walls green.

Should the light green precipitate upon clearing in the xylol, add 3 cc. of absolute alcohol to each 100 cc. of xylol.

Should air-bubbles be detected in the balsam shortly after mounting, heat a dissection needle in a flame and touch each with its tip, when they will be found to disappear.

If too much Canada balsam has been used, some of it usually spreads beyond the edge of the cover-glass, or on its surface. In this event wait until the balsam hardens, when it can be scratched off with a knife, and the surface of the glass cleaned with a rag moistened with turpentine oil or xylol.

Should the Canada balsam become too thick, it can be thinned down with either xylol or benzol.

NO. 3. SAFRANIN AND ANILIN BLUE

1. Stain sections with either alcoholic or anilin safranin (1 per cent. sol. in 50 per cent. alcohol) for from 1 to 24 hours or by heating them in the staining fluid until vapors arise.

2. Wash out excess of stain with 50 per cent. alcohol. Be careful not to remove too much of the safranin. Washing should stop as soon as the stain is reduced to a pink in cellulose walls.

3. Stain with Anilin Blue (1 per cent. sol. in 90 per cent. alcohol) for from 3 to 10 minutes.

4. Wash in 95 per cent. alcohol for a few seconds.

5. Rinse with slightly acidulated 95 per cent. alcohol for a few seconds.

6. Wash for 1 to 2 minutes in 95 per cent. alcohol containing a trace of an alkali to remove the acid.

7. Dehydrate with absolute alcohol for two minutes.

8. Clear in cedar oil for one minute. Blot up excess of cedar oil from around section. Clove oil or xylol may be substituted for the cedar oil.

9. Mount in balsam.

10. Label slide.

If the sections are properly stained, the lignified and suberized walls will be stained red and the cellulose walls a brilliant blue.

No. 4. SAFRANIN AND METHYL GREEN

1. Stain sections with either alcoholic safranin or anilin safranin for from 1 to 24 hours or by warming them in the safranin solution until vapors arise.

2. Remove excess of stain by moving sections about in a vessel of 50 per cent. alcohol until the cellulose walls are pink in color.

3. Stain with 0.5 to 1 per cent. solution of methyl green for from 30 seconds to 5 or 10 minutes. Good results with this stain requires experience with the materials to be stained. The average hand cut section requires 10 minutes.

4. Remove excess of stain and dehydrate in 70 per cent. alcohol for several seconds.

5. Dehydrate with 95 per cent. alcohol for 10 seconds.

6. Further dehydrate by placing material in absolute alcohol for one minute.

7. Clear in cedar oil for one minute. Blot up excess of oil from around edge of section.

8. Mount in Canada balsam.

9. Label slide.

If the sections are properly stained, the lignified and suberized walls will be stained red and the cellulose walls green.

METHOD FOR THE PREPARATION OF A GLYCERIN-GELATIN MOUNT

1. Stain object with an aqueous solution of eosin.

2. Transfer to 1 per cent. acetic acid for a minute.

3. Wash out acid and excess of stain by moving the section about in a dish of water.

4. Transfer object to weak glycerin (glycerin 10 parts, water 90 parts) for 3 to 5 minutes.

5. Transfer object to 50 per cent. glycerin for 3 to 5 minutes.

6. Transfer object to concentrated glycerin for 5 minutes.

7. Remove excess of glycerin around object and mount in glycerin-gelatin. The slide and cover slip should be warmed before the glycerin-gelatin is dropped over the object and the cover slip quickly lowered. The preparation of Glycerin-gelatin is as follows: Macerate 14 grams of gelatin in 84 cc. of water for 2 hours, add 76 cc. of glycerin and warm; add 2 cc. of liquefied phenol, warm and stir for 15 minutes until clear. Filter

while hot through glass-wool or filter paper and collect the filtrate in a wide mouthed bottle. Keep well stoppered so as to exclude dust.

Glycerin-gelatin becomes solid when cool. For use warm the bottle in a water bath after first removing the stopper. A glass rod sufficiently long to reach to the bottom of the bottle can be inserted in the cork and used for transferring the material to the slide.

8. Ring mount with zinc white, balsam or asphaltum at the edge of cover slip. If the cover slip is circular, this can best be done by means of a centering turn-table. A camel's hair brush is dipped into the zinc white, balsam or asphaltum and held to the margin of the cover slip while the slide, fastened with clips to the turn table, is rotated with it.

9. Label slide.

If the objects or sections are such as not to be liable to shrink they can be transferred from water or weak glycerin directly to glycerin-gelatin.

TECHNIQUE OF FIXING, DEHYDRATING, HARDENING AND IMBEDDING IN PARAFFIN

When the intention is to study the protoplasts in their natural form or the processes of cell division, the fresh material must be put through the various stages of fixation, hardening and imbedding before it is sectioned. The steps will now be considered in the order in which they must be carried out.

FIXATION.—This is the process of killing and coagulating the protoplast. The essence of good fixation is in rapid killing. It should be simultaneous with coagulation or hardening so that the protoplast will not be modified by later treatment. Fixing fluids are always substances unknown to protoplasm, *e.g.* poisons. The coagulation of protoplasmic structures is due to the fact that these are alkaline in reaction whereas the fixing fluid is acid. Fixing fluids must be judged not only as to killing and hardening but also as to reaction of tissues to stains afterward. Fluids that are mixtures make the best fixing agents. Among the fixing agents employed are the following: *Osmic acid* (OsO_4) comes in sealed glass tubes containing 0.5 gm. or 1 gm. It has a very powerful odor and is easily affected by organic materials.

It is used in 1 to 2 per cent. solutions and should be made up in distilled water. It fixes cytoplasm well but the nucleus not as good. Its disadvantage lies in its inability to penetrate rapidly.

Chromic acid (CrO_3) in 0.5 to 1 per cent. aqueous solution is very favorable for nuclear structure but like osmic acid penetrates rather slowly.

Picric acid $\text{C}_6\text{H}_2(\text{OH})(\text{NO}_2)_3$ is one of the most penetrating fixing fluids but has very little hardening power. It is employed in saturated aqueous solution.

Corrosive sublimate (HgCl_2) in 0.2 per cent. aqueous or alcoholic solution penetrates and hardens rapidly but does not give as sharp optical differentiation as the others considered.

Absolute alcohol can be employed for very small objects that are dry. If the objects are moist, shrinkage will follow.

Carnoy fluid, consisting of 6 parts absolute alcohol, 3 parts formaldehyde and 1 part of glacial acetic acid, can also be used for fixing small objects. It has the advantage of fixing these in about 10 minutes. Moreover, the objects can be carried directly to absolute alcohol, thence to hot melted paraffin and imbedded.

Chrome-acetic-formalin fluid (*Craf Fluid*). This comparatively new fluid has been found to give very favorable results in fixing materials for chromosome studies. The chromosomes are better defined after staining with Heidenhain's hematoxylin and give a good contrast with the cytoplasm. It is prepared by mixing equal volumes of the following solutions designated as Solution A and Solution B just before using. Solution A contains chromic acid, 1 Gm., glacial acetic acid, 7 cc. and distilled water 92 cc. Solution B contains neutral formalin (40 per cent. formaldehyde solution) 30 cc. and distilled water 70 cc. The material to be fixed should be placed in this fluid for from 12 to 24 hours after which it can be directly transferred to 75 per cent. ethyl alcohol, changing the solution every 15 mi. for 3 or 4 changes, thence to 85 per cent. alcohol (1 hr.) to 65 cc. of 80 per cent. ethyl alcohol and 35 cc. n-butyl alcohol (1 hr.) to 45 cc. of 90 per cent. ethyl alcohol and 55 cc. of n-butyl alcohol (1 hr.) to 25 cc. absolute ethyl alcohol and 75 cc. n-butyl alcohol (1 hr.) to n-butyl alcohol for 2 or 3 hours with hourly changes. The material in butyl alcohol can be infiltrated with paraffin by adding melted

paraffin (m.p. 54° to 55°C.) directly to the beaker or other receptacle containing it and placing the beaker in a paraffin oven at 56° for several hours. Then replace the butyl-alcohol-paraffin mixture with pure melted paraffin for 3 hours and embed.

Formalin-alcohol, consisting of from 2 to 6 per cent. of formalin in 50 to 70 per cent. alcohol, is a good fixing and preserving agent for materials which are not of very delicate texture. It should not be used for unicellular or filamentous algæ and fungi. For these 2 to 6 per cent. aqueous solutions of formalin have given good results. Material can be left in the formalin solutions until required for use, provided care be exercised in storing it in air-tight containers.

For most materials the *Flemming fluids* have proven very satisfactory and are the most generally employed. They are of two classes, viz.; 1. Those that simply involve chromic acid and acetic acid (the chrome-acetic fluids) and 2. Those that involve chromic acid, acetic acid and osmic acid (the Chrome-Osmium-Acetic Fluids). The formulæ follow:

CHROME-ACETIC FLUIDS

Strong	{ 1 per cent. Chromic acid solution	100 cc.
	{ Glacial acetic acid	1 cc.
Medium	{ 1 per cent. Chromic acid solution	70 cc.
	{ 1 per cent. Glacial acetic acid	1 cc.
	{ Distilled water	29 cc.
Weak	{ 1 per cent. Chromic acid solution	25 cc.
	{ 1 per cent. Glacial acetic acid	10 cc.
	{ Distilled water	65 cc.

CHROME-OSMIUM-ACETIC FLUIDS

Strong	{ 1 per cent. Chromic acid solution	75 cc.
	{ 2 per cent. Osmic acid solution	20 cc.
	{ Glacial acetic acid	5 cc.
Weak	{ 1 per cent. Chromic acid solution	25 cc.
	{ 1 per cent. Osmic acid solution	10 cc.
	{ 1 per cent. Acetic acid solution	10 cc.
	{ Distilled water	55 cc.

The acetic acid in all of the Flemming fluids is of great advantage since it penetrates very rapidly, carrying the chromic acid or chromic and osmic acids into the tissue depths, thus insuring complete fixation.

The material to be fixed should be cut into small pieces not longer than 5 mm. nor broader than 2 or 3 mm. The amount of fixative to be used should not be less than 15 times the bulk of the material to be fixed. For delicate material, the weak fixing fluid is used, while for the slightly woody the medium strength fluid and for the more woody, the strong fluid is indicated. The material should be placed in the fixing fluid immediately after it is gathered. One or two drachm homeopathic phials are convenient for the process. The material is kept in the fixing fluid for from 12 to 24 hours and then washed in small cheese cloth bags which are placed in running tap water for from 6 to 12 hours or over night.

DEHYDRATING AND HARDENING OF SOFTER PLANT MATERIALS. After washing, the material, still kept in the bags, is placed in 10 per cent. alcohol for 1 hour and is then carried through a series of alcohols. Each of the series 10 per cent. stronger than the one before it, remaining in each grade for $1\frac{1}{2}$ to 2 hours until 70 per cent. alcohol is reached. Take out of bag and place in phial in 70 per cent. alcohol. If the material is not to be imbedded in paraffine immediately, it can remain in 70 per cent. or 85 per cent. alcohol (if very delicate) until needed. It is not safe to leave very valuable material in a grade below 70 per cent. over night. From the 70 per cent. alcohol it is carried to 85 per cent. to 95 per cent. to absolute alcohol, remaining in each at least 6 hours with 2 or 3 changes of the last.

CLEARING AND IMBEDDING OF SOFTER PLANT MATERIALS.—In order to get the material from absolute alcohol into paraffine, some medium must be used which mixes with absolute alcohol and which also dissolves paraffine. Either oils such as cedar, clove or bergamot or substances like xylol, chloroform or benzol satisfy this requirement. To clear with xylol—transfer material from absolute alcohol to a mixture of $\frac{3}{4}$ absolute alcohol and $\frac{1}{4}$ xylol for 12 hours, then to mixture of equal parts of absolute alcohol and xylol for 12 hours, then to $\frac{3}{4}$ xylol and $\frac{1}{4}$ absolute

alcohol for 12 hours to pure xylol for 12 hours. To phial containing material in pure xylol add paraffine in small pieces and put on top of paraffine bath sufficiently long until paraffine is melted. Then add more paraffine and put phial in paraffine bath at 56°C. over night. Pour fluid off and add pure melted paraffine and repeat 2 or 3 times until rid of all trace of xylol. A tray is then prepared by taking a piece of paper and folding up its edges all around to the height of about a half inch. Half fill this on a cool surface with melted paraffine. Heat two dissection needles in bunsen flame and with these dispose pieces of material in orderly fashion over the crust which has by this time formed at the bottom of the tray. Blow upon the surface of the paraffine to harden it more quickly and as soon as the surface crust will bear it, plunge the tray into cold water. The material can now be left imbedded in paraffine until required for sectioning.

If cutting is to be done in a cool room, softer grades of paraffine with melting points between 40° and 50°C. should be used for imbedding. If, on the other hand, cutting is to be done at summer temperatures, the harder grades melting at between 55° and 70°C. should be employed.

USE OF N-BUTYL ALCOHOL IN IMBEDDING WOODY PLANT ORGANS

This method, originally devised by Dr. Conway Zirkle, is designed to eliminate the hardening and shrinking of tissues induced by the ethyl alcohol and xylol used in the common procedure of paraffin embedding. While the common method, heretofore described, is suitable for the softer plant materials, it is unsuited for plant materials containing abundant lignified elements. The higher concentrations of ethyl alcohol and the xylol so harden the woody elements that they are almost impossible to cut, but break and chip the microtome knife.

Woody plant organs or tissues may be dehydrated and cleared with a mixture of ethyl and N-butyl alcohols, eliminating the use of the higher ethyl alcohols and xylol. N-butyl alcohol dissolves sparingly in water but in all proportions in paraffine. A mixture however of equal parts of ethyl and butyl alcohols is completely miscible with water. By substituting butyl alcohol for xylol, the processes of dehydration and paraffine infiltration

are simplified while hardening and shrinking in many materials are avoided.

DEHYDRATION.—For small cubes of woody material, keep one hour in each of the following solutions (over night in No. 6):

	1	2	3	4	5	6	7	8	9	10	11
Water.....	90	85	75	65	45	25	15	5			
95% E. Alc.....	5	10	20	30	35	50	45	40	25		
Abs. butyl.....	..	..	..	..	10	20	35	55	75	100	100

For Root Tips:

Water.....	90	85	75	65	45	25	15	15	5		
95% E. Alc.....	5	10	20	30	50	70	80	50	40		
Abs. E. Alc.....	..	..	..	..	..	..	..	..	..	25	
Abs. Butyl.....	..	..	..	..	..	..	..	35	50	75	100

Material should remain in pure butyl alcohol until all water is removed. This usually requires several hours.

INFILTRATION.—Pour paraffin into vial (one-half or two-thirds full) and allow to cool. Place material in butyl alcohol on top of paraffin and put vial into oven. After material has stood in the melted mixture for an hour or so, pour off mixture and add pure melted paraffin. Repeat after another hour. Two or more such changes are necessary, depending on the size and nature of the pieces of material. All the butyl alcohol should be removed before imbedding, although traces of it do not interfere with sectioning as do similar amounts of xylol.

TECHNIQUE OF SECTIONING AND MOUNTING MATERIAL IMBEDDED IN PARAFFINE

Strip off the paper tray from the imbedded material and cut out a block of paraffine containing the object which is to be sectioned, taking care to include at least 2 or 3 mm. of paraffine on all sides beyond the specimen. Take a segment of pine wood about an inch long and with a surface at one end about $\frac{3}{8}$ in. square, and coat the square area with melted paraffine. Warm the paraffine on the piece of pine wood and quickly press the paraffine block containing the specimen into this melted paraffine in the desired position for cutting. Heat a dissecting needle and apply this all around the base so that the paraffine block is firmly sealed to the wood. Dip paraffine block in cold water to

harden. Now trim the paraffine block with a sharp scalpel so that the faces form right angles with each other. Adjust the wood in the clamp of the microtome and the microtome blade so that the top of the paraffine block just touches the near surface of the microtome knife. Make certain that the knife edge and the two opposite faces of the paraffine block are perfectly parallel. Now trim the remaining two sides of the block close to the object. Adjust the automatic feed of the rotary microtome by moving dial to number on scale representing thickness in microns desired of sections and turn wheel of microtome. It will be observed that the carrier moves up and down and with each downward movement slightly forward, causing the knife to cut sections which adhere in ribbons.

Transfer the ribbons by means of a camel's hair pencil or dissecting needle to a piece of dust free paper with the side downward which was next to the knife. The ribbons are now ready to be mounted on slides.

The slides to be used should only be those which are devoid of grease or dirt of any kind particularly on the surface upon which the ribbons are to be mounted. A very good plan is to keep a number of slides intended for this purpose submerged in a saturated solution of potassium dichromate in concentrated sulfuric acid. These can be taken out as needed and thoroughly rinsed with water.

With a clean cloth stretched over the forefinger vigorously rub one surface of each slide until perfectly dry and free of lint. Then place a small drop of Mayer's albumin fixative on clean surface and rub over the surface. (The formula for *Mayer's Albumin Fixative* is as follows: Egg white and Glycerin, equal parts, Salicylate of Soda, 1 Gm. to each 100 cc. Mix thoroughly and filter.) Now flood the surface with water and cut the ribbons into segments of the desired length and arrange in rows on slide, being careful to have the segments somewhat shorter than the length of the cover slip because of tendency of paraffine to stretch when warmed. Warm slide gently by holding high above a bunsen flame or flame of an alcohol lamp until ribbons stretch out in smooth fashion. Absorb superfluous water from beneath ribbons with blotting paper held to their edges and at same time

push the sections into even rows. Then leave the sections to dry for several hours or over night.

METHOD FOR THE STAINING AND MOUNTING OF MATERIAL IN PARAFFINE RIBBONS AFFIXED TO SLIDE

1. Gently heat the dry slides with paraffine ribbons adhering to the fixative, high above the Bunsen flame (with the ribbon side up).
2. Place the slide upright in a well of xylol or turpentine. The xylol or turpentine will dissolve the melted paraffine in a minute or two.
3. Take the slide out of the well, wipe off the under side and allow a stream of 95 per cent. alcohol to run over the upper side from a pipette.
4. Place the slide upright in a well of safranin for from four to twenty-four hours.
5. Take the slide out of the safranin well and extract excess of stain with 50 per cent. alcohol. If impossible to extract excess of safranin by this procedure, use acid alcohol and remove acid by washing with 50 per cent. alcohol.
6. Place the slide in a well of gentian violet or fast-green for a second or more. The time varies for different objects and can only be determined by trial.
7. Rinse slide with 70 per cent. alcohol from pipette.
8. Pour absolute alcohol over sections, follow with a few drops of clove oil, replace clove oil with cedar oil.
9. Mount in balsam.
10. Label slide.

IMBEDDING IN CELLOIDIN

Whenever material is unsuited for free hand sectioning and will not give good results when imbedded in paraffine on account of size, hardness, or brittleness, celloidin may be resorted to as an imbedding medium.

The technique employed is similar to that of the paraffine method so far as the preliminary fixing, hardening and dehydrating are concerned up to and including the 95 per cent. alcohol

stage. From this point the various succeeding steps in the procedure are as follows:

1. Place material in equal parts of 95 per cent. alcohol and ether (known as ether-alcohol) for several hours.

2. Transfer to a 2 per cent. solution of celloidin in ether-alcohol, for 2-5 days.

3. Transfer to a 6 per cent. solution of celloidin in ether-alcohol, for 2-5 days.

4. Transfer to a 12 per cent. solution of celloidin in ether-alcohol, for 3-10 days.

5. Prepare a pine block sufficiently large in cross section to support the material and otherwise adapted to its being clamped in the object carrier of the microtome. Soak one end of this block in ether-alcohol for a while and then dip it in the 2 per cent. celloidin solution.

6. Take the material from the thick celloidin and set it in proper position, for cutting the sections desired, on the prepared end of the block and allow the celloidin to thicken for a few seconds only.

7. Dip the celloidin end into the thick solution; remove and hold upright so that the new coating may spread out over the end of the block and solidify the union.

8. As soon as the celloidin has hardened a little to form a surface film, drop the preparation into a vessel of chloroform and allow to remain here 1 day.

9. Transfer preparation to a vessel containing equal parts of glycerin and 95 per cent. alcohol until required for sectioning.

SECTIONING CELLOIDIN MATERIAL

Clamp the block in the sliding microtome and set the knife obliquely so that the sections can be cut with a long sliding stroke. Keep the knife and top of the block wet with the alcohol-glycerin mixture and as soon as the sections are cut, sweep them with a camel's hair pencil into a dish of 70 per cent. alcohol. The sections can be attached to a slide by placing the slide in a closed chamber over ether. The ether vapor dissolves the celloidin and causes the sections to adhere to the slide.

STAINING AND MOUNTING CELLOIDIN SECTIONS

1. Place sections in safranin solution for 1 day. This safranin solution should be made by dissolving as much safranin in 95 per cent. alcohol as it will take up and then diluting with an equal quantity of water.
2. Rinse sections in 50 per cent. alcohol to remove excess of stain.
3. Transfer them to Delafield's hæmatoxylin for 10 minutes (Delafield's hæmatoxylin is made by dissolving 1 gm. of hæmatoxylin in 6 cc. of absolute alcohol and adding this gradually to 100 cc. of a saturated aqueous solution of ammonia alum. This is left exposed for a week, filtered, 25 cc. each of methyl alcohol and glycerin added, allowed to stand 6 hours, again filtered, and ripened about 2 months before using).
4. Rinse sections thoroughly first in water, then in 35 per cent. alcohol, then in 50 per cent. alcohol.
5. Put them quickly through acid alcohol (1 drop of HCl in 50 cc. of 70 per cent. alcohol).
6. Transfer to 70 per cent. alcohol for about 2 minutes.
7. Transfer to 85 per cent. alcohol for about 2 minutes.
8. Transfer to 95 per cent. alcohol for about 2 minutes.
9. Transfer to absolute alcohol for about 2 minutes.
10. Clear sections in a mixture of equal parts of cedar oil and phenol for at least 2 minutes.
11. Remove excess of clearing solution and mount in balsam.
12. Label slide.

DESILICIFICATION OF HARD WOODY MATERIALS

It frequently happens, even after prolonged maceration or boiling in alkaline solutions, that thin sections of hard roots, stems, woods or fruits are difficult or impossible to procure. This is due to the presence of deposits of silica and other mineral substances that usually occur in woody tissues. Therefore, it is of prime importance that these substances be removed as thoroughly as possible. For this purpose a 10 per cent. aqueous solution of commercial Hydrofluoric Acid (or stronger solutions up to the pure acid for very hard materials) is most useful. Small fruits

or short segments of other hard material are placed in this acid (which should be kept in a bottle coated internally with a thick layer of paraffine) for from 3 days to a week, depending on the size of the objects, with one or two changes of the acid. The acid is then washed out thoroughly with running water for 2 to 5 hours. This treatment completely frees the tissues of all mineral deposits without affecting the organic structure.

SCHULZE'S MACERATION PROCESS

This method is employed for the separation of cells. Radial-longitudinal sections, that may be cut with a pen knife, are placed in a beaker or test tube containing 50 cc. of nitric acid of specific gravity 1.3 (about 2 volumes of nitric acid and 1 volume of water will serve the purpose). To this add 1 gm. of chlorate of potash crystals and heat gently until the reddish color which first appears in the tissues has disappeared. Stop the action by pouring the whole of the contents into a vessel containing water and wash well with water. The cells can now be readily separated with dissection needles and mounted in water for examination. Do not mount in glycerine, for it makes the already bleached elements too transparent.

Classified List of Reference Works

GENERAL BOTANY

- Bergen, J. Y., and B. M. Davis. *Principles of Botany*. Ginn and Co., Boston, 1906.
- Coulter, J. M., C. R. Barnes, and H. C. Cowles. *A text-book of botany for colleges and universities*. 3v. American Book Co., New York, 1931.
- Eyster, W. H. *College Botany*. R. Long and R. R. Smith, Inc., New York, 1932.
- Gager, C. S. *General Botany*. P. Blakiston's Son and Co., Phila., 1926.
- Ganong, W. F. *A text-book of botany for colleges*. Parts 1 and 2. Macmillan Co., New York, 1919-20.
- Holman, R. M., and W. W. Robbins. *A text-book of general botany for colleges and universities*. 3 ed. John Wiley and Son's, New York, 1934.
- Kerner von Marilaun, A. *The natural history of plants. Their forms, growth, reproduction and distribution*. Transl. by F. W. Oliver. 2v. H. Holt and Co., New York, 1895.
- Mottier, D. B. *A text book of botany for college students*. P. Blakiston's Son & Co., Inc., Phila., 1932.
- Robbins, W. J. and H. W. Rickett. *Botany*. 2 ed. D. Van Nostrand Co., Inc., New York, 1934.
- Sinnot, E. W. *Botany: principles and problems*. McGraw-Hill Book Co., New York, 1923.
- Smith, G. M., J. B. Overton and others. *A text-book of general botany*. Revised ed. Macmillan Co., New York, 1928.
- Strasburger, E. *A Textbook of Botany*. Rewritten by H. Fitting, H. Seip, R. Harder and G. Karsten. 6th English ed. Macmillan Co., N. Y., 1930.

PLANT MORPHOLOGY

- De Bary, H. A. *Comparative morphology and biology of the fungi, mycetoza and bacteria*. Eng. transl. by H. E. F. Garnsey, rev. by I. B. Balfour. Clarendon Press, Oxford, 1887.
- De Bary, H. A. *Comparative anatomy of the vegetative organs of the phanerogams and ferns*. Transl. and annotated by F. O. Bower and D. H. Scott. Clarendon Press, Oxford, 1884.
- Eames, A. J. *Morphology of vascular plants (lower groups)*. McGraw-Hill Book Co., N. Y., 1936.
- Eames, A. J., and L. H. MacDaniels. *An introduction to plant anatomy*. McGraw-Hill Book Co., New York, 1925.
- Goebel, K. *Organography of plants. Part I: General organography*. Eng. ed. by I. B. Balfour. Clarendon Press, Oxford, 1900.
- Goebel, K. *Outlines of the classification and special morphology of plants*. Eng. transl. by H. E. Garnsey, rev. by I. B. Balfour. Clarendon Press, Oxford, 1887.

- Haberlandt, G. *Physiological plant anatomy*. Transl. by M. Drummond. Macmillan and Co., London, 1914.
- Hitchcock, A. S. *The genera of grasses of the United States*. U. S. D. A. Bul. No. 772, 1920.
- Jeffrey, E. C. *The anatomy of woody plants*. Univ. of Chicago Press, Chicago, 1917.
- Molisch, Hans. *Anatomie der Pflanze*. G. Fischer, Jena, 1922.
- Scott, D. H. and F. T. Brooks. *An introduction to structural botany*. Part 1. Flowering plants. 11ed. A. and C. Black, Ltd., London, 1927.
- Scott, D. H. and F. T. Brooks. *An introduction to structural botany*. Part 2. Flowerless plants. 9ed. A. and C. Black, Ltd., London, 1924.
- Solereider, H. *Systematic Anatomy of the Dicotyledons*. Transl. by L. A. Boodle and F. E. Fritsch. Rev. by D. H. Scott. 2v. Clarendon Press, Oxford, 1908.
- Stevens, W. C. *Plant Anatomy from the standpoint of the development of the tissues and handbook of microtechnic*. 4ed. P. Blakiston's Son and Co., Phila., 1924.
- Rusby, H. H. *A manual of structural botany*. Lea and Febiger, Phila., 1911.

PLANT HISTOLOGY AND MICROSCOPY

- Ballard, C. W. *Elements of vegetable histology*. J. Wiley and Sons, New York, 1921.
- Chamot, E. M. *Elementary Chemical Microscopy*. 2ed. New York, 1921.
- Gathercoal, E. N. and E. H. Wirth. *Pharmacognosy*. Lea and Febiger, Phila., 1937.
- Greenish, H. C. *The microscopical examination of foods and drugs*. 3ed. J. and A. Churchill, London, 1923.
- Hanausek, T. F. and A. L. Winton. *The microscopy of technical products*. John Wiley and Sons, New York, 1907.
- Mansfield, W. *Histology of medicinal plants*. John Wiley and Sons, New York, 1916.
- Matthews, J. M. *The textile fibers*. New York, 1924.
- Rusby, H. H., and S. E. Jelliffe. *Morphology and Histology of plants*. New York, 1899.
- Wallis, T. E. *Analytical Microscopy, its aims and methods*. E. Arnold and Co., London, 1923.
- Whipple, G. C. *Microscopy of drinking water*. 3ed. John Wiley and Sons, New York, 1914.
- Winton, A. L. and J. Moeller. *Microscopy of vegetable foods*. 2ed. J. Wiley and Sons, New York, 1916.
- Youngken, H. W. *A textbook of Pharmacognosy*. 4ed. P. Blakiston's Son and Co., Phila., 1936.

MICROSCOPICAL TECHNIQUE

- Chamberlain, C. J. *Methods in plant histology*. 4th ed. University of Chicago Press, Chicago, 1924.
- Conn, H. J. *Biological Stains*. 3ed. Geneva, N. Y., 1936.
- Bailey, I. W. *Microtechnique for woody structures*. Botanical Gazette 49: 57-58, 1910.

- Molisch, H. *Mikrochemie der Pflanze*. 2te Aufl. F. Fischer, Jena, 1921.
- Schneider, H. and A. Zimmermann. *Die botanische Mikrotechnik, ein Handbuch der mikroskopischen Arbeitsverfahren*. 2te Aufl. G. Fischer, Jena, 1922.
- Strasburger, E. *Das botanische Prakticum*. 7te ed. Rev. by Max Koernicke. G. Fischer, Jena, 1923.
- Zimmermann, A. *Botanical microtechnique*. Transl. by J. E. Humphrey. H. Holt and Co., New York, 1893.

PLANT TAXONOMY

- Abrams, L. *An illustrated flora of the Pacific States in 3 volumes*. Vol. I, Ferns to Birthworts. Stanford University Press, 1923.
- American Joint Committee on Horticultural Nomenclature. *Standardized plant names*. Salem, 1923.
- Beal, W. J. *Michigan Flora*. Mich. Agr. Col., 1905.
- Britton, N. L. and Addison Brown. *An illustrated flora of the northern United States, Canada and the British possessions*. 2nd ed. C. Scribner's Sons, New York, 1913.
- Coulter, J. M., and A. Nelson. *New manual of botany of the central Rocky Mountain region*. (Vascular plants.) American Book Co., New York, 1909.
- DeDalla Torre, C. G., and H. Harms. *Genera siphonogamarum ad systema Englerianum conscripta*. W. Engelmann, Leipzig, 1900-1907.
- Engler, A. and K. Prantl. *Die Natürlichen Pflanzenfamilien nebst ihren Gattungen u. wichtigeren Arten, insbesond. der Naturpflanzen*. 23v. W. Engelmann, Leipzig, 1900-1925; 2ed., 1930- .
- Engler, A., and E. Gilg. *Syllabus der Pflanzenfamilien*. 4te and 10te Aufl. Gebrüder Borntraeger, Berlin, 1924.
- Engler, A. *Das Pflanzenreich regni vegetabilis conspectus*. (Issued serially). Heft 1-90, 1900-1927. W. Engelmann, Leipzig, 1900- .
- Gerth Van Wijk, H. L. *A dictionary of plant names*. 2v. M. Nijhoff, 1916.
- Gray, Asa. *Gray's new manual of botany. Handbook of the flowering plants and ferns of central and northeastern United States and adjacent Canada*. 7th ed. Revised by B. L. Robinson and M. L. Fernald. American Book Co., New York, 1908.
- Gray, Asa. *Synoptical flora of North America*. 2ed. Ivison, Blakeman, Taylor & Co., 1886.
- Hitchcock, A. S. *Methods of descriptive systematic botany*. John Wiley and Sons, New York, 1925.
- Hutchinson, J. *Families of flowering plants*. I. Dicotyledons. II. Monocotyledons. Macmillan Co., London, 1926; 1934.
- Index Kewensis*. 4v, 1893-1895 and 8 supplements, 1911-1936. Clarendon Press, Oxford.
- Jackson, B. D. *Guide to the literature of botany*. Longmans Green and Co., London, 1881.
- Jepson, W. L. *A manual of the flowering plants of California*. Associated Student's Store, San Francisco, 1925.
- Le Maout, Em. and J. Decaisne. *General system of botany, descriptive and analytical*. Transl. and ed. by J. D. Hooker. London, 1872.

- Lyons, A. B. Plant names, scientific and popular. Nelson Baker, Detroit, 1900.
- Pfeiffer, L. Nomenclator botanicus. 2v. T. Fischer, Kassel, 1873-1874.
- Pool, R. J. Flowers and flowering plants. 1ed. McGraw-Hill Book Co., N. Y., 1929.
- Pritzel, G. A. Thesaurus literaturae botanicae . . . 2ed. F. O. Brockhaus, Leipsig, 1872.
- Rydberg, P. Flora of the Rocky Mountains and adjacent plains. 2nd ed. New York, 1922.
- Small, J. K. Flora of the Southeastern United States. New York, 1903.
- Swingle, D. B. A textbook of systematic botany. 1ed. McGraw Hill Book Co. New York, 1928.
- Wettstein, R. Handbuch der Systematischen Botanik. 3te Aufl. F. Deuticke, Leipsig und Wien, 1924.
- Wiegand, K. M. and A. J. Eames. The flora of the Cayuga Lake Basin, New York. Cornell Univ. Ag. Exp. Sta. Memoir 92, 1925.

CYTOLOGY AND GENETICS

- Babcock, E. B. and R. E. Clausen. Genetics in relation to agriculture. 2nd ed. McGraw-Hill Co., New York, 1927.
- Castle, W. E. Genetics and eugenics. 4ed. Harvard Univ. Press, 1931.
- Meyer, A. Morphologische und Physiologische Analyse d. Zelle d. Pflanzen u. Tiere. G. Fischer, Jena, 1920.
- Newman, H. H. Evolution, Genetics and Eugenics. 3ed. Univ. of Chicago Press, 1932.
- Sharp, L. W. An introduction to cytology. 2nd ed. Rev. McGraw-Hill Book Co., New York, 1926.

BACTERIOLOGY AND MICROBIOLOGY

- Bergey, D. H. Manual of Determinative Bacteriology. 4th ed. Williams and Wilkins Co., Baltimore, 1934.
- Conn, H. W. and H. J. Conn. Bacteriology. 2nd ed. Williams and Wilkins Co., Baltimore, 1924.
- Jordan, E. O. A text-book of general bacteriology. 11th ed. W. B. Saunder's Co., Phila., 1935.
- Marshall, C. E. Microbiology. A text-book of microorganisms general and applied. 2ed. Rev. P. Blakistons Son and Co., Phila., 1917.
- Park, W. H., A. W. Williams and C. Krumwiede. Pathogenic microorganisms. 7th ed. Rev. Lea and Febiger, Phila., 1920.
- Rice, T. B. Text Book of Bacteriology. W. B. Saunders Co., Phila., 1935.
- Schneider, A. The microbiology and microanalysis of foods. P. Blakiston's Son and Co., Phila., 1920.
- Zinsser, H. A text-book of bacteriology. 5th ed. D. Appleton and Co., New York, 1925.

DRUG PLANT CULTIVATION

- Corbett, L. C. The propagation of plants. Farmers' Bulletin No. 157, U. S. D. A. Government Printing Office, Washington, 1917.
- Corbett, L. C. Pruning. Farmer's Bul. No. 181, U. S. D. A., 1915.

- Dillman, A. C. The production of seed flax. U. S. D. A. Farmer's Bul. No. 1328, 1924.
- Henkel, Alice. Weeds used in medicine. U. S. D. A. Farmer's Bul. No. 188, 1917.
- Henkel, Alice. American root drugs. U. S. D. A. Bur. Pl. Ind. Bul. 107, 1907.
- Hood, S. C. Commercial production of thymol from horsemint (*Monarda punctata*). Bul. No. 372, U. S. D. A.
- Kraemer, H. Cultivation of medicinal plants, pp. 727-748, in *Applied and Economic Botany*, Phila., 1914.
- Kilmer, F. B. The cultivation of medicinal plants. *Am. Jour. Pharm.*, 87: 343-359 and 421-435, 1915. (This article includes an extensive citation of the literature on the subject.)
- Rabak, F. The production of volatile oil and perfumery plants in the United States. U. S. D. A. Bur. Pl. Ind. Bul. No. 195, 1910.
- Rixford, G. P. Smyrna Fig culture. U. S. D. A. Bul. No. 732, 1918.
- Russell, G. A. Drying crude drugs. U. S. D. A. Farmer's Bul. No. 1231, 1921.
- Stansel, R. H. and R. H. Wyche. Fig culture in the Gulf Coast region of Texas. *Tex. Agr. Exp. Sta. Bul.* no. 466, 1932.
- Stockberger, W. W. Drug plants under cultivation. *Farmer's Bul.* No. 663, U. S. D. A., 1915.
- Stockberger, W. W. Ginseng culture. U. S. D. A. Farmer's Bul. No. 1184, 1928.
- Stockberger, W. W. Growing and curing hops. U. S. D. A. Farmer's Bul. No. 304, 1928.
- Shrader, J. H. The castor oil industry. Bul. No. 867, U. S. D. A., 1920.
- Sievers, A. D. Some effects of selection on the production of alkaloids in belladonna. Bul. No. 306, U. S. D. A., 1915.
- Sievers, A. F. American medicinal plants of commercial importance. U. S. D. A. misc. publ. 77, 1930.
- Sievers, A. F. Peppermint and Spearmint as farm crops. U. S. D. A. Farmer's Bul. No. 1555, 1929.
- Teetgen, A. B. Profitable herb growing and collecting. *Country Life, London*, 1919.
- Tschirch, A. *Handbuch der Pharmakognosie*. 2ed. Chr. H. Tauchnitz, Leipsig, 1930.
- Van Fleet, W. Goldenseal under cultivation. U. S. D. A. Farmer's Bul. No. 613, 1922.
- Wood, M. N. Almond varieties in the U. S. U. S. D. A. Dept. Bul. No. 1282, 1924.

PLANT ECOLOGY AND GEOGRAPHY

- Campbell, D. H. An outline of plant geography. Macmillan and Co., New York, 1926.
- Clements, F. E. Research methods in ecology. University Publishing Co., Lincoln, Nebraska, 1905.
- Committee on the Preservation of Natural Conditions of the Ecological Society of America. Naturalists' guide to the Americas. Edited by V. E. Shelford and others. Williams and Wilkins Co., Baltimore, 1926.
- Engler, A., and O. Drude. *Die Vegetation der Erde*. 15v. W. Engelmann, Leipsig, 1896-1923.

- Fernald, M. L. Persistence of plants in unglaciated areas of Boreal America. *Memoirs of the American Academy of Arts and Sciences.* 15: No. 3, 239-342, 1925.
- Grisebach, A. *Die Vegetation der Erde nach ihrer Klimatischen anordnung.* 2v. 2te. Aufl. W. Engelmann, Leipsig, 1884.
- Hilgard, E. W. *Soils, their formation, properties, composition, and relation to climate and plant growth.* Macmillan and Co., New York, 1906.
- Knuth, P. *Handbook of flower pollination.* Transl. by J. R. A. Davis. 3v. Clarendon Press, Oxford, 1906-1909.
- Scharff, R. F. *Distribution and origin of life in America.* Macmillan Co., New York, 1912.
- Schimper, A. F. W. *Pflanzen-Geographie auf Physiologischer Grundlage.* G. Fischer, Jena, 1898.
- Warming, E. and M. Vahl. *Ecology of plants. An introduction to the study of plant communities.* Transl. by P. Groom and I. B. Balfour. Clarendon Press, Oxford, 1909.
- Weaver, J. E. and F. E. Clements. *Plant Ecology.* McGraw-Hill Book Co., Inc., New York, 1929.

PLANT PHYSIOLOGY AND PLANT CHEMISTRY

- Detmer, W. *Practical plant physiology.* Transl. by S. A. Moor. Swan, Sonnenschein and Co., London and Macmillan Co., New York, 1909.
- Duggar, B. M. *Plant physiology with special reference to plant production.* Macmillan Co., New York, 1911.
- Green, J. R. *An introduction to vegetable physiology.* 3rd ed. J. and A. Churchill, London. P. Blakiston's Son and Co., Phila., 1911.
- Haberlandt, G. F. J. *Physiologie und Ökologie.* B. J. Teubner, Leipsig, 1917.
- Hass, P., and T. G. Hill. *An introduction to the chemistry of plant products.* 3rd ed. 2v. Longmans, Green and Co., 1921 and 1922.
- Henry, T. A. *The plant alkaloids.* 2nd ed. J. and A. Churchill, London, 1924.
- Jost, L. *Lectures on plant physiology.* Transl. by R. J. H. Gibson and supplement. Clarendon Press, Oxford, 1907 and 1913.
- Maximov, N. A. *Textbook of Plant Physiology.* McGraw-Hill Book Co., Inc., New York, 1930.
- Oppenheimer, C. *Die Fermente und ihre Wirkung.* 5te Aufl. Issued serially. Georg Thieme, Leipsig, 1924- .
- Palladin, V. I., and B. L. Livingston. *Palladin's Plant Physiology.* 3rd Amer. ed. P. Blakiston's Son and Co., Phila., 1926.
- Pfeffer, W. *The physiology of plants. A treatise upon the metabolism and sources of energy in plants.* Transl. by A. J. Ewart. 3v. Clarendon Press, Oxford, 1900 and 1906.
- Tummann, O. and L. Rosenthaler. *Pflanzenmikrochemie.* Gebr. Bornträger, Berlin, 1931.

HORTICULTURE AND AGRONOMY

- Bailey, L. H. *Standard Cyclopedia of Horticulture.* 6v. Macmillan Co., New York, 1900-1917.
- Bailey, L. H. *A manual of cultivated plants.* Macmillan Co., New York, 1924.

- Harshberger, J. W. Textbook of pastoral and agricultural botany. P. Blakiston's Son and Co., Phila., 1920.
- Meehan, T. The native flowers and ferns of the United States. 2v. Robson, Phila., 1880.
- Rehder, A. Manual of the cultivated trees and shrubs hardy in North America. Macmillan Co., New York, 1927.
- Robbins, W. W. Botany of crop plants. 2ed. P. Blakiston's Son and Co., Phila., 1924.

MYCOLOGY

- Atkinson, G. F. Mushrooms, edible, poisonous, etc. Andrus and Church, Ithaca, 1900.
- Buller, A. H. R. Researches on Fungi. 3v. Vol. 1, 1909, Vol. 2, 1922, Vol. 3, 1924. Longmans Green and Co., London.
- Duggar, B. M. Fungous diseases of plants. Ginn and Co., Boston, 1909.
- Harshberger, J. W. A text-book of mycology and plant pathology. P. Blakiston's Son and Co., Phila., 1917.
- Henrici, A. T. Molds, yeasts and actinomycetes. J. Wiley Sons, Inc., N. Y., 1930.
- Heald, F. D. Manual of plant diseases. 2ed. McGraw-Hill Co., N. Y., 1932.
- McIlvaine, C. One thousand American fungi. Bowen-Merrill Co., Indianapolis, 1900.

MEDICINAL AND POISONOUS PLANTS

- Bentley, R. and H. Trimen. Medicinal plants. 4v. J. and A. Churchill, London, 1880.
- Bernhard-Smith, A. Poisonous plants of all countries. Bailliere, Tindall and Cox, London, 1923.
- Long, H. C. Plants poisonous to live stock. Univ. Press, Cambridge, 1917.
- Millspaugh, C. F. American medicinal plants. 2v. Boericke, New York, 1887.
- Nees von Esenbeck, T. F. L. Plantæ medicinales. 3v. Arnz, Dusseldorf, 1828.
- Pammell, L. H. A manual of poisonous plants. The Torch Press, Cedar Rapids, Ia. Part I, 1910, Part II, 1911.
- Wood, H. C., C. H. LaWall, H. W. Youngken, I. Griffith, A. Osol and L. Gershenfeld. The United States Dispensatory. 22nd ed. J. B. Lippincott Co., Phila., 1937.
- Woodville, W. Medical botany. 5v. 3ed. Bohn, London, 1832.
- Youngken, H. W. A text-book of pharmacognosy. 4th ed. P. Blakiston's Son and Co., Phila., 1936.

HISTORY OF BOTANY

- Harvey-Gibson, R. J. Outlines of the history of botany. A. and C. Black, London, 1919.
- Green, J. R. History of botany, 1860-1900. Clarendon Press, Oxford, 1909.
- Green, E. L. Landmarks of botanical history. Smithsonian Institution, Washington, 1909.
- von Sachs, J. History of botany, 1530-1860. Rev. by I. B. Balfour. Clarendon Press, Oxford, 1906.

Glossary

- ABOR'TION.—The imperfect or non-development of an organ.
- ACAULES'CENT.—Without an obvious aerial stem.
- ACHENE' (akene).—A small, dry, one-celled, indehiscent fruit in which the seed coat and pericarp (fruit wall) are not firmly attached.
- ACHLAMY'DEOUS.—Destitute of calyx and corolla.
- ACIC'ULAR.—Applied to crystals of calcium oxalate, etc., that are needle-shaped.
- ACIDS (Plant Acids).—A widely diverse group of cell constituents having the property of uniting with radicals to form salts or esters.
- ACROP'ETAL.—Development from outside (below) toward the inside (above).
- ACTINOMOR'PHIC.—Applied to flowers like those of the Mustard, etc. which are capable of division into equal halves in more than one direction. Cf. zygomorphic.
- ACU'MINATE.—Tapering gradually to a long point.
- ACUTE'.—Sharp-pointed, the point being less than a right angle.
- AD'NATE.—Applied to the growing together of unlike parts.
- ADVENTI'TIOUS.—Applied to roots and buds that are out of their ordinary position.
- AESTIVA'TION.—Arrangement of the parts of the flower in the bud.
- ALBU'MEN.—Nutritive material stored in the embryo, endosperm, or perisperm.
- ALBUR'NUM.—Sapwood.
- ALEU'NONE.—Applied to small protein bodies in which are usually embedded one or more crystalloids, one or more globoids and at times a small rosette of calcium oxalate.
- ALKALOIDS.—Nitrogenous organic bases.
- ALLELO'MORPHS.—Alternative forms of a gene which are located at the same point on each one of a pair of chromosomes.
- AL'TERNATE.—Applied to leaves, buds, etc., that are arranged singly (one after another) at the nodes.
- AM'ENT.—A scaly, spike-like inflorescence. Another name for catkin.
- AMOR'PHOUS.—Without definite shape.
- AMPHIT'ROPOUS (ovules and seeds).—Half-inverted and straight, with the hilum about the middle, and micropyle terminal.
- AMPLEX'ICAUL.—Clasping the stem.
- ANAL'OGY.—Resemblance in function.
- ANASTOMO'SING.—Applied to veins that are connected with others by cross veins, so forming a network, as with the marginal veins of Eucalyptus.
- ANAT'ROPOUS.—Inverted ovules or seeds with micropyle adjacent to hilum.
- ANDRÆ'CIUM.—The male system of organs in a flower.
- ANDROG'YNOUS.—Applied to inflorescences composed of both staminate and pistillate flowers.
- ANEMOPHI'LOUS.—Wind pollinated.
- ANGIOSPERM'OUS.—Having ovules and seeds borne within a box-like covering, the pericarp.

- AN'NUAL.—Producing flowers, fruit and seed within a year from the time the seed germinated and then dying completely.
- AN'NULAR.—Ring-like.
- ANTE'RIOR.—The front region.
- AN'THER.—That portion of a stamen which bears the pollen.
- AN'THERIDIUM.—Male sexual organ of Thallophytes, Bryophytes and Pteridophytes.
- AN'THEROZOID.—A male sexual cell formed within an antheridium.
- AN'THOPHORE.—A lengthened internode of the receptacle between calyx and corolla.
- APET'ALOUS.—Without petals, as in the oaks, etc.
- APOCAR'POUS.—Carpels separate and distinct.
- APOG'AMY.—The direct development of a sporophyte from a gametophyte without the functioning of sexual organs or cells.
- APOPET'ALOUS.—Petals separate and distinct.
- APOSEP'ALOUS.—Sepals separate and distinct.
- APOTHE'CIUM.—A disk like or cup-shaped fruiting organ of a lichen and some cup-fungi.
- ARCHEGO'NIUM.—A multicellular female sexual organ found in the bryophytes, pteridophytes and gymnosperms.
- AR'IL.—An accessory seed covering outside of the testa and arising at or about the hilum, as in *Evonymus*.
- AR'ILLODE.—A fake accessory seed covering outside of the testa, as in Nutmeg, and arising from the dilatation of the micropyle.
- ARIS'TATE.—Having a stiff bristle-like termination.
- ASCEND'ING.—Growing obliquely upward.
- AS'CUS.—A sac-like spore case of an Ascomycete fungus.
- AT'AVISM.—Reversion to ancestral type. The reappearance of a character after a lapse of one or more generations.
- AURIC'ULATE.—Ear-like.
- AWN.—A bristle-like structure that branches along its axis.
- AX'IL.—Upper angle formed where the leaf joins the stem.
- AX'ILLARY.—In the axil.
- BAC'CATE.—Berry-like.
- BAL'SAM.—A mixture of resin with cinnamic or benzoic acid or both and a volatile oil.
- BÄRB.—A short bristle usually bent back.
- BARK.—All that portion of an exogenous plant axis outside of the cambium line.
- BÄST.—Applied to the phloem region but mainly to the fibrous portion thereof.
- BEAD'ED.—Applied to a thickened, jointed cell wall showing in sectional view somewhat circular areas in the wall like a row of beads, due to a close proximity of the pores in the wall.
- BEARD'ED.—Furnished with long hairs.
- BER'RY.—A fleshy fruit whose mesocarp and endocarp are fleshy and frequently succulent throughout, and with seeds imbedded therein, as tomato, capsicum, belladonna, etc.
- BI.—A prefix of the Latin language indicating two, twice or doubly.
- BI'COLLAT'ERAL BUN'DLE.—A collateral bundle with the xylem strand enclosed between two phloem strands, cambium being present or absent on either outer or inner face of the xylem or on both faces.

- BIEN'NIAL.—Applied to plants that live for more than one year but not longer than two years.
- BILA'BIATE.—Two lipped.
- BLĀDE.—Expanded part of a leaf.
- BLOOM.—The whitish and waxy secretion of epidermal cells, as in the stems of Sugar Cane or the leaves of Cabbage.
- BORKE.—The dead bark tissues which arise when secondary phellogen forms cork in deep regions of the bark.
- BRĀCT.—A modified leaf, frequently scale-like, appearing on inflorescence axes.
- BRAC'TEOLE (bracteolar leaf).—A modified leaf found on pedicels.
- BŮD.—A rudimentary stem.
- BULB.—A very short scaly underground stem.
- BUL'BILS.—Small underground bulbs, as in garlic.
- BUL'BLETS.—Small above ground bulbs, as in the tree onions.
- BUN'DLE.—See Vascular Bundle. Also applied to a more or less cylindrical group of any parallel, elongated cells or other objects.
- BUN'DLE SHEATH.—An endodermis or a border parenchyma enclosing a single bundle.
- CADU'COUS.—Falling with the opening of the flower, as the calyx of *Papaver*.
- CA'LYX.—The outermost whorl of floral leaves.
- CAM'BIUM.—A secondary meristem lying between the phloem and xylem in certain collateral bundles (fascicular or intrafascicular cambium) or between the phloem and xylem portions of medullary rays (interfascicular cambium).
- CAMPAN'ULĀTE.—Bell shaped.
- CAMPYLO'TROPOUS.—Applied to ovules or seeds that are curved so as to bring the apex and base near together.
- CANES'CENT.—White or gray from a coating of fine hairs.
- CAOUT'CHOUC.—A tenacious, elastic hydrocarbon found in the latex of many plants.
- CAPILLIT'IUM.—A network of filaments among spores, as in the sporangium of slime molds, the fruit body of puff balls, etc.
- CAP'ITATE.—Shaped like a head.
- CAP'RIFICATION.—The process of pollinating figs artificially.
- CAP'SULE.—A dry, dehiscent fruit of a compound pistil consisting of two or more carpels.
- CARBO'HY'DRATE.—A substance consisting of carbon and of hydrogen and oxygen in the proportion of two parts of hydrogen to one part of oxygen.
- CAR'PEL.—A transformed leaf bearing one or more ovules; a simple pistil; a part of a compound pistil.
- CARPOGON'IUM.—A unicellular, female sexual organ of certain red algæ and fungi consisting of a cell enlarged at its base, its terminal part ending in a long filament called a *trichogyne*.
- CAR'POPHORE.—A slender stalk, the prolongation of the receptacle, to which the inferior akenes (mericarps) of the *Umbelliferae* are attached.
- CARUN'CLE.—A warty excrescence at or about the hilum of certain seeds, as that of castor-oil seed, etc.
- CARYOP'SIS.—A dry, indehiscent, one-seeded fruit of the grasses or cereals in which the fruit wall (pericarp) and seed coat firmly adhere.
- CAT'KIN.—A scaly spike of flowers.

- CAU'DATE.—Tailed.
- CAULES'CENT.—With an obvious aerial stem.
- CAU'LINE.—Pertaining to the stem.
- CELL.—The structural unit of plants and animals consisting of a protoplast which in most plants is surrounded by a cell wall secreted by the protoplast.
- CENTRIF'UGAL.—Applied to a flower cluster in which the terminal or central flower blossoms first.
- CENTRIP'ETAL.—Applied to a flower cluster in which the lower or outer flowers bloom first.
- CHAFF.—The glumes and paleas of grains; the scaly hairs on the stipes of ferns; the bracts subtending each floret in some heads of *Compositæ*.
- CHALA'ZA.—That portion of the ovule marked by the junction of the integuments with the nucellus.
- CHASMO'GAMOUS.—Pertaining to flowers that regularly open.
- CHLAMY'DOSPORE.—Thick walled spore formed within the hyphæ of smuts.
- CHLO'ROPHYLL.—The mixed pigments forming the green color of all green plants and found in the chloroplastids.
- CHLOROPLAS'TID.—A protoplasmic body in the cells of green parts of plants containing chlorophyll.
- CHRO'MATIN.—That portion of the nucleus which is readily colored by a basic dye. The substance that carries the hereditary characters from parent to offspring.
- CHROMO'MERE.—One of the chromatin granules, arranged linearly like the beads on a string, comprising the chromosome.
- CHROMOPLAS'TID.—A protoplasmic body in the cells of certain parts of plants containing a pigment other than chlorophyll.
- CHRO'MOSOME.—One of the bodies into which the chromatin of the nucleus is resolved during indirect nuclear division. The chromosomes carry the genes, linearly arranged, which control the development of Mendelian characters.
- CIL'IA.—Vibratory, hair-like, protoplasmic outgrowths of zoöspores, bacteria, gametes, etc.
- CIRCUMNUTA'TION.—The repeated bending in different directions of the growing tips of stems of climbing plants.
- CIR'CINATE.—Rolled inward from apex toward base, as the young leaves of ferns.
- CIRCUMSCIS'SILE.—Applied to the splitting open of capsules transversely into lid and pot portions.
- CLAD'ODE.—The flattened branch which somewhat resembles a leaf.
- CLAW.—The narrowed base of some petals, as those of the Pink Family.
- CLEISTOG'AMOUS.—Applied to flowers that never open but are self fertilized, as in some *Polygalas* and *Violets*.
- COCH'LEA.—A spirally coiled legume.
- COE'NOCYTE.—A multinucleate cell.
- COHE'SION.—The union of parts of the same whorl.
- CO'HORT.—A group of natural orders.
- COLEORHI'ZA.—A root sheath through which the radicle bursts in germination.
- COLLAT'ERAL.—Applied to fibrovascular bundles in which the phloem and xylem masses are arranged side by side.
- COLLEN'CHYMA.—Tissue composed of elongated prismatic cells thickened at their angles with a colloidal substance.

- COLUMEL'LA.—The end cell wall of an aerial hypha that bulges into the sporangium; also applied to the axis of a capsule.
- COL'UMN.—The united stamens and carpels in Orchids.
- COL'UMN CELLS.—Hour-glass shaped cells forming a layer beneath the palisade-epidermal layer of leguminous seeds.
- CO'MA.—A tuft of hairs, as found on the seeds of Milkweeds.
- COM'MISSURE.—The contiguous surfaces of two carpels as in the flowers and fruits of the Parsley Family.
- COMPAN'ION CELLS.—Parenchyma cells associated with sieve tubes and having a narrow cavity, abundant protoplasm, large nuclei and the wall adjoining the sieve tube marked by transversely elongated pits.
- CONCEN'TRIC.—Applied to several circles or whorls one within the other. Concentric vascular bundles are those in which the xylem mass surrounds the phloem mass or vice versa.
- CONCEP'TACLE.—A sac bearing the fruiting organs in certain Algæ and Fungi.
- CON'DUCT'ING BUN'DLE.—(See Vascular Bundle.)
- CON'DUCT'ING PAREN'CHYMA.—Somewhat elongated parenchyma cells serving primarily for the conduction of fluids.
- CONDU'PLICATE.—Folded together lengthwise, as for example the bud leaves of the oak or peach.
- CONID'IA.—Asexual spores cut off from the ends of hyphæ or sterigmata by *Penicillium*, *Aspergillus*, *Peronospora*, *Claviceps*, etc.
- CONID'IOPHORE.—A hypha bearing conidia.
- CONJUGA'TION.—One of the sexual methods of reproduction where two like sexual cells unite to form a zygospore.
- CON'NATE.—Applied to parts that have grown together, as the bases of two opposite leaves.
- CONNECT'IVE.—The continuation of the filament of the stamen that connects the two lobes of the anther.
- CONNI'VENT.—Brought close together; converging.
- CON'VOLUTE.—Rolled lengthwise from one edge, as the leaves in the buds of the Wild Cherry and Plum.
- COR'DATE.—Heart-shaped.
- CORIA'CEOUS.—Leathery in texture.
- CORK (SU'BEROUS TISSUE).—A protective tissue derived from the phellogen and with cell walls waterproofed with suberin.
- CORK CAMB'IUM.—(See Phellogen.)
- CORM.—A solid, swollen, fleshy underground stem.
- COROL'LA.—The inner whorl of floral envelopes composed of petals.
- CORO'NA.—A crown like appendage in the throat of the corolla, as in the flowers of *Narcissus* and *Silene*.
- COR'TEX.—That region in dicotyl and gymnosperm roots of primary growth and in roots and stems of monocotyledons between the outer protective tissue or epidermis and the pericycle; in dicotyl and gymnosperm roots of secondary growth or in barks between phellogen and phloem or pericycle. The PRIMARY CORTEX is the region lying between the pericycle and the epidermis or the secondary tissues produced by the phellogen. The SECONDARY CORTEX is all that tissue laid down by the phellogen on its inner face.
- COR'TICAL PAREN'CHYMA.—The parenchyma of the cortex.

- COR'YMB.—A flat topped or convex centripetal inflorescence with the lowermost pedicels the longest.
- COS'TA.—A rib.
- COTYLE'DON.—A seed-leaf of the embryo.
- CREM'OCARP.—The peculiar fruit of *Umbelliferae*, consisting of two inferior akenes (mericarps) separated from each other by a carpophore.
- CRE'NATE.—Applied to leaf margins having rounded teeth.
- CREN'ULATE.—The margin with fine rounded teeth.
- CRIB'RIFORM.—Sieve-like.
- CRU'CIFORM.—Applied to the corolla or the calyx of flowers, the parts of which are arranged in the form of a cross.
- CRUSTA'CEOUS.—Applied to the thallus of a lichen that closely adheres to the substratum.
- CRYP'TOGAM.—A plant belonging to one of the divisions of the vegetable kingdom below the Spermatophytes.
- CRYS'TALS.—The solid geometrical forms assumed by many chemically homogeneous bodies.
- CRYS'TAL SAND.—See Microcrystals.
- CRYS'TALLOID.—A protein body found in the aleurone grains of seed or underground parts.
- CULM.—A jointed stem of a grass or sedge.
- CU'NEATE.—Wedge-shaped.
- CU'PULE.—Applied to the concave involucre enclosing the glans of an acorn but also to other cup-shaped parts of plants.
- CU'TICLE.—A thin covering of a waxy substance called cutin on the outer wall of epidermal cells.
- CU'TIN.—A fatty cell-wall constituent serving to waterproof the wall.
- CUTINI'ZED.—Applied to a cell-wall infiltrated or covered with cutin.
- CUS'PIDATE.—Tipped with a sharp rigid point.
- CYME.—A more or less flat topped determinate inflorescence.
- CY'MOSE.—Cyme-like.
- CYS'TOLITHS.—Cell bodies consisting of deposits of calcium carbonate or silica or both on an extension of the cell wall into the cell.
- CY'TASE.—An enzyme which digests cellulose.
- CYTOL'OGY.—The study of cells and their contents.
- CY'TOPLASM.—The cell protoplasm outside of the nucleus.
- DECAN'DROUS.—Having ten stamens.
- DECID'UOUS.—Applied to leaves which fall in autumn, to plants bearing such leaves and to the calyx and corolla which fall shortly after blossoming, before the development of the fruit.
- DEC'LINATE.—Curved or bent downward.
- DECOMPOUND'.—Several times compounded, as the leaf-blades of *Cimicifuga*.
- DECUM'BENT.—Erect at base, then lying on the ground, with the end rising.
- DECUS'SATE.—Applied to opposite leaves when the pairs stand at right angles to each other along the stem.
- DEHIS'CENCE.—Splitting open.
- DELIQUES'CENT.—Applied to a tree whose trunk or main stem is lost in branches.
- DEL'TOID.—Having the shape of the Greek letter Δ.

- DEN'TATE.—Having broad, acute, marginal teeth pointing outward.
- DENTIC'ULATE.—Finely dentate.
- DERMAT'OGEN.—The generative tissue that gives rise to epidermis.
- DETER'MINATE.—Applied to inflorescences on which flowering begins with the terminal bud, thus ending the elongation of the stem bearing the flowers.
- DIADEL'PHOUS.—Applied to stamens whose filaments are united at their edges into two sets.
- DIAGEOT'ROPIC.—Applied to a plant organ that assumes a horizontal position.
- DIAN'DROUS.—Possessing two stamens.
- DI'ASTASE.—A ferment found in germinating seeds and fungal hyphæ which changes starch into maltose.
- DICHLAMYD'EUS.—Pertaining to flowers that possess both calyx and corolla.
- DICHOG'AMY.—The maturation of one set of sexual organs before the other.
- DICHOT'OMOUS.—Forked.
- DIC'LINOUS.—Pertaining to the stamens and carpels being found in separate flowers.
- DICOT'YLE'DON.—A plant whose embryo possesses two seed leaves or cotyledons.
- DIG'ITATE.—Referring to a compound leaf whose leaflets come off at the end of the petiole.
- DIMOR'PHISM.—Having two forms of flowers, one with long styles and short stamens, the other with short styles and long stamens; the occurrence of two distinct forms.
- DICE'CIOUS.—Applied to species having two kinds of individuals, male and female.
- DISSECT'ED.—Cut deeply into numerous divisions.
- DISSEPI'MENT.—A partition separating cavities in a compound ovary or fruit.
- DIS'TICHOUS.—Pertaining to the arrangement of leaves in two rows.
- DIVI'DED.—Segmented to the mid-rib or base.
- DORSIVEN'TRAL.—Having distinct upper and lower surfaces.
- DOR'SUM.—The back of an organ. The lower surface of a foliage or floral leaf.
- DOWN'Y.—Covered densely with soft hairs.
- DRUPE.—A one-celled, one-seeded fruit whose endocarp is stony.
- DRUPE'LET.—A small drupe.
- DUCT.—A tubular element found in the xylem region of a vascular bundle. See trachea.
- DURA'MEN.—Heartwood.
- E- OR EX-, a prefix meaning devoid of, outside of, or away from.
- ECCEN'TRIC.—Deviating from the center. Applied to the hila of starch grains which are outside of the center, also to woody plants which develop more rapidly on one side than on the other.
- ECHIN'ULATE.—Beset with small prickles or spines.
- ECH'INATE.—Beset with prickles or spines.
- EC'TOPLASM (Outer Plasma Membrane).—The specialized, non-granular, outer limiting layer of the cytoplasm.
- EGG-APPARA'TUS.—The ovum and two synergids at the micropylar end of the embryo-sac.
- ELABORATED SAP.—The nutrient solution of grape sugar, etc. manufactured in the green cells of leaves which passes downward through the phloem region to be distributed to parts needing it.
- ELAIO'PLASTID.—A plastid which stores oil or fat.

- EL'ATER.—An elastic, spiral filament attached to the spores of some Liverworts and Horsetails and aiding in their dispersal when mature.
- EMAR'GINATE.—Notched at the apex.
- EM'BRYO.—A rudimentary plant found within the seed.
- EMBRYOL'OGY.—The study of the embryo and its development.
- EM'BRYO-SAC.—A large cell within the nucellus of the ovule in which the embryo is formed after fertilization.
- EN'DOCARP.—The inner layer of the pericarp. The inner epidermis of the fruit wall.
- ENDO'DERMIS.—A layer of cells forming the innermost boundary of the cortex and surrounding the central cylinder. Also applied to the bundle sheath surrounding concentric bundles in the stems and leaves of pteridophytes and some monocotyledons.
- EN'DOGEN.—A Monocotyledon.
- ENDOG'ENOUS.—Applied to the axes of Monocotyl plants that do not increase materially in diameter.
- EN'DOPHYTE.—A plant which grows within the tissues of another.
- EN'DOSPERM.—A storage tissue of the seed, consisting of the matured embryo sac around the embryo, and formed from the endosperm nucleus.
- EN'DOSPORE.—The inner wall of a spore.
- ENDOTHE'CIUM.—A zone of one or more layers within the exothecium of an anther.
- EN'SIFORM.—Sword-shaped.
- ENTOMOPH'ILOUS.—Insect pollinated.
- EN'TOPHYTE.—See Endophyte.
- EN'ZYMES.—Nitrogenous, colloidal bodies closely related to the proteins and concerned with the chemical changes occurring in all living cells.
- EPHEM'ERAL.—Lasting for a brief period (a day or so).
- EPICA'LYX.—A whorl of bracts resembling the calyx but below it.
- EPI'CARP.—The outer layer of the pericarp or the outer epidermis of fruits.
- EPICOT'YL.—The portion of the embryo-axis above the cotyledon or cotyledons.
- EPIDER'MIS.—The primary outer covering layer of cells of plants, sometimes later replaced by cork.
- EPIG'YNOUS.—Applied to floral leaves that appear to be inserted upon the ovary.
- EPIPET'ALOUS.—Upon the corolla.
- EP'IPHYTE.—An air plant. A plant growing on another plant but not necessarily nourished by it.
- EPITHE'LIUM.—A delicate layer of cells lining an internal cavity.
- EQ'UITANT.—Applied to leaves, as in Iris, when they all spring from a rhizome and are successively folded on each other toward their bases.
- ERYTH'ROPHYLL.—The red coloring matter of leaves.
- ESTIVA'TION (Aestivátion).—The arrangements of the floral organs in the flower bud.
- ETÆ'RIO.—An aggregate fruit like the Raspberry or Blackberry, the product of a single flower, consisting of an aggregation of drupelets on a receptacle.
- E'TIOLATION.—The bleaching of green parts of plants when kept in the dark for some time.
- EVOLU'TION.—The presumable theory that all forms of living things existing today have been derived from others previously existing, either by direct descent or by common ancestry.

EXALBU'MINOUS.—Applied to a seed in which the nourishment is stored in the embryo during the growth of seed from the ovule stage.

EXCEN'TRIC.—See Eccentric.

EXCRES'CENCE.—A morbid outgrowth.

EXCRE'TION.—Getting rid of nitrogenous waste.

EXCUR'RENT.—Applied to trees, the main stems of which do not disappear in branches but grow erect to the summit, ending in a terminal bud. The opposite of Deliquescent.

EXFO'LIATE.—To shed layers of bark. To cast off layers of tissue.

EX'INE.—The outer wall of a pollen grain.

EX'OCARP.—The outer layer of the pericarp.

EXOG'ENOUS.—Applied to the axes of Gymnosperms and Dicotyledons which increase materially in diameter.

EX'OGENS.—Plants with exogenous axes.

EXOSPOR'IUM.—The outer wall of a spore.

EXSERT'ED.—Applied to stamens that protrude from the throat of the corolla.

EX'TINE.—The outer wall of a pollen grain.

EXTORSE'.—Applied to anthers which face outward, away from the gynœcium.

FACE.—The free surface of an organ.

FAL'CATE.—Scythe or sickle-shaped.

FAM'ILY.—A sub-division of an order.

FARINA'CEOUS.—Starchy or mealy.

FAS'CICLE.—A bundle or cluster.

FASCIC'ULAR.—Belonging to a bundle.

FASCIC'ULAR CAMBIUM.—See Cambium.

FASCIC'ULATE.—Clustered.

FATS.—Glyceryl esters of fatty acids that tend to solidify and crystallize at ordinary temperatures.

FEC'ULA.—The nutritive part of a cereal.

FER'TILE.—Producing fruit or reproductive organs. Applied to flowers which contain functionally active stamens and carpels.

FERTILIZA'TION.—That method of reproduction characterized by the union of two dissimilar gametes.

FI'BER.—A cell that is much elongated, thick-walled, usually lignified, tapering more or less to a point at either end and primarily functioning mechanically.

FI'BROUS.—Fiber-like. Referring to root systems composed of many slender rootlets.

FIBROVAS'ULAR BUN'DLE.—A stringy group of fibers, vessels or tracheids and cells coursing through the various organs of the higher plants and serving for support and conduction of sap.

FIL'AMENT.—The stalk of a stamen; a thread-like structure.

FILAMEN'TOUS.—Thread-like.

FIL'IFORM.—Thread-like.

FIM'BRIATED.—Fringed.

FIS'SION.—A form of division in which the cell separates into two equal or nearly equal parts.

FIS'TULAR.—Hollow and cylindrical.

FIXED OILS.—Glyceryl esters of fatty acids that tend to remain liquid at ordinary temperatures.

FLAGEL'LUM.—A whip-like protoplasmic outgrowth of certain organisms, as Flagellates or of zoöspores, serving as an organ of locomotion.

FOLIA'CEOUS.—Leaf-like.

FOL'LICLE.—A one-chambered, dry fruit that dehisces along one suture only.

FOVIL'LA.—The contents of a pollen grain.

FRAC'TURE.—The manner in which a root or other plant part breaks when subjected to sufficient pressure.

FROND.—The leaf of a fern. The leaf-like, branching thallus of marine algæ.

FRUIT.—A matured pistil, or ovarian portion thereof together with any closely adhering part.

FRU'TICOSE.—Shrubby.

FUCO'XAN'THIN.—The brown pigment of the brown algæ (synonymous with phycophæin).

FUGA'CIOUS.—Falling off early.

FUNDAMEN'TAL TIS'SUE.—Ground-tissue. The tissue of plants through which the fibrovascular bundles course.

FUNIC'ULUS.—The stalk of an ovule.

FUR'CATE.—Forked.

FU'SIFORM.—Enlarged in the middle and tapering toward either end.

GAL'BULUS.—A berry-like cone, as in *Juniperus*, formed by the coalescence of fleshy scales and carpels.

GA'LEATE.—Helmet shaped.

GAMETAN'GIUM.—The cell or organ lodging the gamete.

GAM'ETE.—A sexual cell.

GAM'ETOPHYTE.—The sexual generation.

GAMOPET'ALOUS.—Applied to a flower whose corolla is composed of petals which are more or less united at their edges.

GAMOSEP'ALOUS.—Having the sepals more or less united at their margins.

GEM'MA.—An asexual bud-like structure found in the cupules of Liverworts.

GEMMA'TION.—The process of budding as seen in the yeasts.

GENE.—The unit of inheritance which is found in the chromosomes of the germ cells.

GEN'ERA.—Plural of genus.

GENIC'ULATE.—Kneed.

GEOT'ROPISM.—Response to the stimulus of gravity.

GERMINA'TION.—The sprouting of a spore or seed.

GERM CELL.—A reproductive cell as distinguished from a somatic or body cell.

GILLS.—The spore-bearing plates of a toadstool.

GLA'BROUS.—Smooth.

GLAND.—A cell or group of cells whose principal function is secretion.

GLAND'ULAR.—Pertaining to or bearing glands.

GLANS.—A nut.

GLAU'COUS.—Covered with a bloom.

GLO'BODS.—Small spherical crystals of calcium-magnesium-phosphate found in aleurone grains.

GLOB'ULAR.—Spherical.

GLOM'ERULE.—A head-like cyme. A determinate head.

GLU'COSIDE (Gly'coside).—A cell constituent which under the influence of an enzyme or of heat, dilute acids or alkalies splits up into glucose and some other substances.

- GLUME.—A floral bract of the grasses and sedges.
- GLU'TEN.—The proteid matter of cereals.
- GONID'IUM.—Applied to the algal cell in lichens as well as to many forms of asexual reproductive bodies in flowerless plants.
- GON'OPHORE.—An upgrowth of the receptacle between the corolla and stamens, as in *Passiflora*.
- GUARD CELLS.—The pair of special epidermal cells which surround the pore of the stoma.
- GYN'ÆCI'UM.—The female sexual system of a flower.
- GYN'OPHORE.—An upgrowth of the receptacle between gynoecium and androecium, as in *Geum*.
- GYNOSTE'MIUM.—The united stamens and style. The column of orchids.
- HAB'ITAT.—The original home of a plant.
- HA'DROME.—The conducting tissue of the xylem, comprising tracheæ, tracheids and wood parenchyma.
- HAS'TATE.—Shaped like the head of a halberd, the basal lobes diverging.
- HEAD.—An indeterminate form of inflorescence, as seen in the Daisy family, in which the flowers are in a dense cluster on the receptacle.
- HEART'WOOD.—The inner portion of the wood, often containing resinous and coloring matters and which no longer functions in the vegetative processes of the plant.
- HELIOT'ROPISM.—Response to the stimulus of sunlight.
- HERBA'RIVM.—A classified collection of dried plant specimens.
- HERED'ITY.—The faithful transmission of characters from parent or parents to offspring.
- HERMAPH'RODITE.—Applied to flowers which contain both sets of essential organs, not necessarily functionally active.
- HESPERID'IUM.—A large, thick-skinned, succulent fruit like the orange, lemon or grape-fruit.
- HET'EROCYST.—A cell of different size and content from other cells in the filaments of Nostoc and some other blue-green algæ.
- HETEROPHYL'LOUS.—Having more than one kind of foliage-leaves on the same plant.
- HETEROS'POROUS.—Producing asexual spores of more than one kind, as in *Selaginella* and the rusts.
- HETEROTYP'IC.—Applied to the first cell division in meiosis which involves the reduction of the chromosomes from the diploid to the haploid member.
- HETEROZY'GOTE.—An organism to which its two parents have contributed unlike genes with respect to any given allelomorphic pair governing contrasted characters, and which in turn produces two kinds of germ cells with respect to the characters.
- HEX.—A prefix of Greek origin meaning six.
- HEXAG'YNOUS.—Having six carpels or styles.
- HEXAM'EROUS.—Having the parts of the flower in 6's.
- HEXAN'DROUS.—Having six stamens.
- HIBERNA'TION.—Passing the winter in a dormant state of existence.
- HI'LUM.—The scar of a seed, after the stalk of the ovule has fallen off. Also applied to the point of origin or growth of a starch grain.

- HIP.—The fruit of a Rose, consisting of a number of akenes surrounded by a ripened concave receptacle.
- HIRSUTE'.—Covered with numerous long coarse hairs.
- HISPID.—Beset with erect stiff hairs, as Borage.
- HISTOL'OGY.—The study of tissues with the aid of the microscope.
- HOLDFAST.—The base of the stipe or of the plant body of a thallophyte which is differentiated for attaching the plant to its substratum.
- HOMOL'OGOUS.—Having the same structural nature.
- HOMOS'POROUS.—Producing asexual spores of only one kind.
- HOMOZY'GOTE.—An organism whose parents contributed to it a similar member of any given pair of genes and whose germ cells are all alike with respect to the genes for that character.
- HORMOGON'IUM.—A segment of a filament of a blue-green alga which may detach itself from the rest of the filament becoming a new organism.
- HYBRID.—A cross between two varieties or species, rarely between two genera of the same family.
- HYDATH'ODES (Water Pores).—Organs for the exudation of water from plants under conditions of low transpiration and abundant soil moisture.
- HYDROPH'ILOUS.—Applied to flowers that are pollinated through the agency of water currents.
- HY'DROPHYTE.—A water-plant.
- HYDROT'ROPISM.—The response of a plant organ to the stimulus of moisture.
- HYGROSCOP'IC.—The property possessed by certain cells or substances of absorbing moisture with avidity.
- HYME'NIUM.—A spore-bearing membrane of a fungus.
- HY'PHA.—A filament of the mycelium of a fungus.
- HYPO.—A prefix of Greek origin meaning under.
- HY'POCOTYL.—That part of an embryo plantlet below the cotyledon or cotyledons.
- HYPOCRATER'IFORM.—Applied to a calyx or corolla when the tube is long and slender and abruptly expands into a flat limb.
- HYPODER'MIS.—That portion of a plant organ directly beneath the epidermis.
- HYPOGE'OUS.—Beneath the surface of the soil.
- HYPOTHE'CIUM.—That portion of a thallus of a lichen directly beneath or around the apothecium.
- HYPOG'YNOUS.—Applied to the insertion of various floral parts on the receptacle and beneath the pistil.
- Id'IOBLAST.—A cell which differs materially in form, size, character of cell wall, or contents from its neighbors in a tissue, frequently applied to colorless stone cells.
- IMBIBI'TION.—The taking in of water by organic bodies in such a manner as to cause them to swell up.
- IM'BRICATE.—Overlapping like shingles.
- IMMERSED'.—Growing entirely under water.
- IMPARIPIN'NATE.—Applied to a pinnately-compound leaf terminating with a single leaflet.
- INDEF'INITE.—Applied to stamens and other organs of the flower, when too numerous to be conveniently counted.
- INDEHIS'CENT.—Not splitting open in a definite manner when ripe.
- INDIG'ENOUS.—Native.

- INDU'SIUM.—An outgrowth of the lower epidermis of many ferns that covers the cluster of sporangia.
- INEQUILAT'ERAL.—Having unequal sides.
- INFER'IOR.—Applied to an ovary when the other floral parts are inserted above it.
- INFLORES'CENCE.—The arrangement of the flowers on the stem of a plant.
- INFUNDIB'ULIFORM.—Funnel shaped.
- INNATE'.—Applied to anthers that are attached by their base to the summit of the filament.
- INTEG'UMENT.—A covering.
- INTERCEL'LULAR.—Between the cells.
- INTERFASCIC'ULAR CAM'BIUM.—Applied to a cambium layer which extends from one fibrovascular bundle to another in the stems of Dicotyledons and Gymnosperms.
- INTER'NAL GLANDS.—More or less globular or elliptical shaped secretion reservoirs.
- INTER'NAL PHLOEM (Intraxylary Phloem).—A strand of phloem on the inner face of the xylem, as found in bicollateral bundles.
- IN'TERNODE.—That portion of the stem between two nodes.
- INTERRUP'TEDLY-PIN'NATE.—Applied to a pinnate leaf that has either smaller or larger leaflets between those of usual size.
- IN'TINE.—The inner coat of the pollen grain.
- IN'TRA.—A prefix meaning within.
- INTRAPET'IOLAR.—Applied to stipules that are between the petiole and the stem; also to buds that are beneath or inside of the base of the petiole.
- INTRORSE'.—Applied to anthers that face toward the gynœcium.
- INTUSSUSCEP'TION.—The formation of additional particles of protoplasm between those already present.
- IN'ULIN.—A carbohydrate isomeric with starch found in the cell sap of the Compositæ, Lobeliaceæ and some other families, from which it crystallizes in lumps (in air-dried material) or in sphaerocrystals (in alcohol-preserved material).
- IN'VOLUCRE.—A whorl (or whorls) of bracts subtending a flower or flower cluster.
- INVOL'UCEL.—A secondary involucre.
- IN'VOLUTE.—Applied to the arrangement of leaves within a bud when they are rolled inward from both sides.
- IRRITABIL'ITY.—That property of living matter whereby it responds to a stimulus.
- ISO'DIAMET'RIC.—Applied to cell having nearly the same length, breadth and thickness.
- ISOG'AMY.—The union of sexual cells of similar form.
- ISOM'EROUS.—Having the same number of parts in each whorl.
- ISOSTEM'ONOUS.—Having the stamens and petals each in one whorl and of the same number.
- ISTH'MUS.—Applied to the constricted portion between the two half cells in certain desmids.
- KARYOKINE'SIS.—Indirect nuclear division.
- KATAB'OLISM.—Destructive metabolism.
- KEEL.—Applied to a longitudinal ridge or elevation of cortical tissue of Senega root which extends from the crown downward. Also applied to the two inferior petals of a papilionaceous corolla which are more or less united into a body resembling the keel of a boat.

KNEE.—A form of knot which projects upward into the air from the roots of certain trees that grow in wet soil, notably the bald cypress.

LABEL'LUM.—The large, lip-like, lower petal in the flower of an orchid.

LA'BIATE.—Two lipped.

LA'BIUM.—The lower lip of a labiate flower.

LACIN'iate.—Applied to the margins of leaves which are deeply cut into irregular narrow lobes.

LAMEL'LÆ.—Little plates or layers. Applied to the layers of carbohydrate material in a starch grain which surround the growing point; also to the gills of a toad-stool as well as to the more or less regular markings found in cell walls formed by delicate layers of material of different water content or of different composition.

LAM'INA.—The blade or expanded part of any leaf.

LA'NATE.—Covered with long, curled, wool-like hairs.

LAN'CEOLATE.—Lance-shaped.

LA'TEX.—The milk juice of a plant.

LATICIF'EROUS.—Applied to the latex carrying tissue of a plant.

LATIFO'LIATE.—Possessing broad leaves.

LEAF.—An expansion of the stem or branch in whose axil one or more branches arise.

LEAF'LET.—A division of a compound leaf.

LEAF-TRACE.—A vascular bundle while on its way from the stem bundle to the leaf.

LEG'UME.—A dry, simple, capsular fruit formed of a single carpel and dehiscent by both ventral and dorsal sutures.

LEN'TICELS.—Fissures in the cork or epidermis of stems formed by the swelling up and rupture of subjacent cells which were cut off by the phellogen in places where cork was not laid down.

LENTIC'ULAR.—Having the shape of a double convex lens.

LEP'TOME.—The conducting tissue of the phloem, comprising sieve tubes, phloem parenchyma and sometimes companion cells.

LEU'COPLAST.—A colorless plastid found in the cells of plants not exposed to light.

LI'ANE.—A woody climber or twiner of tropical forests.

LI'BER.—The inner bark or phloem region of Gymnosperms and Dicotyledons.

LI'BRIFORM-CELLS.—Those cells of the xylem that are thick-walled and resemble bast fibers.

LIG'NEOUS.—Woody.

LIG'NIFIED.—Covered with deposits of lignin.

LIG'NIN.—A mixture of substances that adheres to the cellulose walls of certain cells and which is characterized by taking on a reddish coloration with phloroglucin and hydrochloric acid reagent.

LIG'ULATE.—Strap-shaped.

LIG'ULE.—A membranous appendage at the summit of the leaf-sheath in many grasses and cereals; a strap shaped corolla of a Composite.

LIGULIFLO'ROUS.—Applied to *Compositæ* flower heads, as those of Dandelion and Chicory, which contain ligulate florets only.

LIMB.—The spreading portion of a gamosepalous calyx or a gamopetalous corolla.

LINE.—One-twelfth of an inch.

LIN'EAR.—Many times longer than broad and with nearly parallel margins.

- LOBE.—A division of a leaf or other flattened organ which is larger than a tooth but which is not a leaflet.
- LOC'ULAR.—Having a cavity or cavities.
- LOCULICI'DAL.—Applied to the dehiscence of a capsule when it splits open along the dorsal suture.
- LOC'ULUS.—A cell or cavity of an anther, ovary, or fruit.
- LO'MENT.—A modified, jointed or multilocular legume that breaks open transversely into segments when mature.
- LU'CID.—Clear.
- LU'MEN.—The cell cavity. The open space bounded by the walls of a cell.
- LU'NIFORM.—Half-moon or crescent shaped.
- LU'RID.—Dingy-brown.
- LUTES'CENT.—Yellowish.
- LY'RATE.—Applied to a pinnatifid leaf, as that of the Turnip, in which the terminal lobe is the largest and the rest decreasing in size toward the base.
- LYSIG'ENOUS.—Applied to the formation of a type of intercellular-air-space which originates through the breaking down of cell walls common to a group of cells; also applied to secretion reservoirs which are formed by the complete breaking down of contiguous secretory cells which leave their products in the cavity formed.
- MACRO.—A prefix of Greek origin meaning large.
- MACROSPORAN'GIUM.—A spore case containing one or more macrospores. (The nucellus in Spermatophytes.)
- MAC'ROSPORES.—The larger of the two different kinds of spores produced by some of the higher Pteridophytes and the Spermatophytes. (The embryo-sac in Spermatophytes.)
- MACROSPO'ROPHYLL.—The leaf bearing the macrosporangium. (The carpel in Spermatophytes.)
- MAC'ULATE.—Spotted.
- MA'MILLATE.—Bearing teat-like protuberances.
- MARCES'CENT.—Withering but not falling, dropping off.
- MARINE'.—Applied to plants which grow in the sea or ocean.
- MEDUL'LA.—Pith.
- MED'ULLARY.—Pertaining to the pith.
- MED'ULLARY RAYS.—Vertical, radial plates of parenchyma connecting the cortex with the pith or a portion of the xylem with a portion of the phloem.
- MED'ULLARY SHEATH.—The inner ring of xylem abutting on the pith.
- MEGASO'RUS.—The ovule.
- MEGASPORAN'GIUM.—See macrosporangium.
- MEG'ASPORE.—See macrospore.
- MEGASPO'ROPHYLL.—The carpel of seed plants. A leaf bearing one or more megasporangia.
- MEIO'SIS.—The reduction divisions during the maturation of the gametes, in which the members of each chromosome pair separate, and the chromosome number in the resulting daughter cells is reduced to half the somatic number.
- MEM'BRANE CRYSTALS.—Crystals enclosed within a membrane developed within the cell.
- MEM'BRANOUS.—Thin, soft and flexible.

- MER'ICARP.—One of the two inferior akenes which are found with the carpophore making up the cremocarp in *Umbelliferae*.
- MER'ISTEM.—Formative or undifferentiated tissue consisting of cells which in the living plant are in an active state of division.
- MERISTEMAT'IC.—Consisting of generative cells or meristem.
- MES'OCARP.—The middle region of the fruit wall or pericarp, lying between the epicarp and endocarp.
- MES'OPHYLL.—The region of leaf parenchyma within the epidermis or, in flat leaves, between the two epidermises.
- MES'TOME.—The conducting portion of a fibrovascular bundle.
- METAB'OLISM.—The sum total of all the chemical changes which take place in a living plant.
- METAGEN'ESIS.—Alternation of generations. The production of sexual individuals by asexual means and asexual or neutral individuals by sexual means.
- METAMOR'PHOSIS.—A change in the form or function of an organ or organism.
- METAPHLO'EM.—Phloem formed by the cambium. Secondary phloem.
- METAXY'LEM.—The secondary xylem, formed by the cambium on its inner face.
- MICRO.—A prefix of Greek origin meaning small.
- MI'CROBE.—A minute vegetable or animal organism.
- MI'CROCrys'TALS (Sphenoidal Crystals).—Very minute, arrowhead- or deltoid-shaped crystals frequently packed in great numbers in a single cell.
- MI'CROPYLE.—The opening between the coats of an ovule through which the pollen tube enters. The orifice or foramen in the seed coat through which the hypocotyl passes during germination.
- MICROSO'MES.—Applied by Strasburger to minute particles in the protoplasm which have a high degree of refringency.
- MICROSO'RUS.—A lobe of the anther.
- MICROSPORAN'GIUM.—A spore case containing microspores. An anther sac.
- MI'CROSPORE.—A small spore found in a microsporangium. The pollen grain of a seed plant.
- MICROSPO'ROPHYLL.—A leaf bearing microsporangia. The stamen of seed plants.
- MID'DLE LAMEL'LA.—A layer of calcium pectate or of some other base in combination with pectic acid, appearing within the vacuole between two plasma membranes during the last stage of cell division and appearing in suitably mounted matured tissues as a dividing line between the cellulose walls of adjacent protoplasts.
- MID'RIB.—The large main central vein of a pinnately-veined leaf which is continuous with the leaf stalk.
- MITO'SIS.—Indirect nuclear division.
- MONADEL'PHOUS.—Applied to stamens which are united by their filaments into one set, as in the *Malvaceae*.
- MONAN'DROUS.—Possessing only one stamen.
- MONAN'THOUS.—Having only a single flower on the peduncle.
- MONIL'IFORM.—Resembling a chain of beads.
- MONO.—A prefix of Greek origin, meaning one or single.
- MONOCAR'PELLARY.—Of one carpel.
- MONOCHLAMYD'EUS.—Possessing but one perianth whorl.
- MONOCLIN'IC PRISMS (Monoclinic Crystals).—Crystals possessing three axes of unequal length, two of which are obliquely inclined to each other and the third forming right angles to these two.

- MONOC'LINOUS.—Having both androecium and gynoecium.
- MONOCOTYLED'ONOUS.—Having only one cotyledon or seed leaf.
- MONŒ'CIOUS.—Having separate staminate and pistillate flowers on the same plant.
- MONOHY'BRID.—The offspring of two parents differing in only one mendelian characteristic.
- MONOLOC'ULAR.—One chambered.
- MONOM'EROUS.—Applied to flowers having one part running through each whorl.
- MONOPO'DIUM.—A plant axis which elongates at the apex and sends off lateral branches in acropetal sequence.
- MONOS'TICHOUS.—Arranged in one vertical row.
- MU'CRONATE.—Terminating abruptly in a small soft point.
- MULTI.—A prefix of Latin origin meaning many.
- MULTICEL'LULAR.—Consisting of many cells.
- MULTICIP'ITAL.—Many-headed; applied to a rhizome or root from which numerous stems arise.
- MULTIFA'RIOUS.—Composed of many diverse parts.
- MULTILOC'ULAR.—Many celled or chambered.
- MUL'TIPLE FRUIT.—A fruit composed of many small fruits, each the product of a separate flower, as in the Fig or Hop.
- MUTA'TION.—A sudden variation that is later passed on through inheritance and that results from changes in a gene or genes.
- MYCE'LIUM.—The vegetative body of a fungus consisting of intertangled hyphæ.
- MYCOL'OGY.—That branch of Botany that treats of the Fungi.
- MYCORRHI'ZA.—An association between the roots of certain plants and the mycelium of certain fungi which forms an investment about their tips.
- NA'PIFORM.—Turnip-shaped. Somewhat globular, becoming abruptly slender and then terminating in a conical tap root.
- NAT'URALIZED.—Applied to plants that have been introduced from another country.
- NAVIC'ULAR.—Boat-shaped.
- NEC'TAR.—A sweet secretion by the flower.
- NEC'TARY.—The part of the flower which secretes nectar.
- NEIGH'BORING CELLS (Subsidiary Cells).—The epidermal cells which abut upon the guard cells of the stoma.
- NERVA'TION.—The arrangement of veins in a leaf.
- NETTED.—See Reticulate.
- NEU'TRAL.—Said of flowers which possess neither stamens nor carpels. Also applied to the asexual generation of plants.
- NIV'EOUS.—Snow-white.
- NODE.—The place on the stem which normally shows outgrowths of a leaf, whorl of leaves or leaf modifications.
- NODOSE'.—Having swollen joints or knobs.
- NOD'ULE.—A small rounded body as a root tubercle.
- NOR'MAL.—Usual.
- NON.—Not.
- NUCEL'LUS.—The body of an ovule.
- NUCIF'EROUS.—Nut-bearing.
- NU'CLEUS.—A dense, definitely circumscribed region of protoplasm within the cell containing linin, chromatin and nuclear sap.

NUCLE'OLUS.—A small body of dense protoplasm within the nucleus.

NUT.—A dry, indehiscent, 1-celled, 1-seeded fruit with a stony or leathery pericarp.

NUT'LET.—A small nut. The characteristic fruit of the *Labiatae*.

NUTRI'TION.—That branch of Physiology which includes the absorption, distribution and assimilation of food stuffs.

OB.—A prefix of Latin origin signifying inversion.

OBCON'ICAL.—Inversely cone-shaped.

OBCOR'DATE.—Inversely heart-shaped.

OBLAN'CEOLATE.—Lance-shaped with the broadest part toward the summit.

OBLATE'.—Flattened at the ends or poles.

OB'LIGATE.—Necessary, indispensable.

OBLIQUE'.—Taking a position between erect and horizontal, as in the case of many stems. More developed on one side than on the other, as in certain leaf blades.

OB'LONG.—Longer than broad with nearly parallel sides.

OBO'VATE.—Ovate with the attachment at the narrower end.

OBTUSE'.—Having a blunt or rounded end.

O'CHREA (o'crea).—A sheathing stipule.

OILS.—See Fixed Oils; Volatile Oils.

OLEORES'INS.—Secretions composed of resins dissolved in volatile oil.

ONTOG'ENY.—The history of the development of an individual.

O'ÖSPORE.—The fertilized egg.

OPEN COLLATERAL BUNDLE.—A collateral bundle in which a cambium zone lies between the xylem and phloem strands.

OPER'CULUM.—The transversely dehiscent lid or cover of a moss capsule.

ORBIC'ULAR.—Circular.

OR'DER.—A division of a class containing one or more families.

ORTHOT'ROPOUS.—Applied to ovules or seeds which are erect, with the micropyle at the apex and the hilum coinciding with the chalaza.

O'VARY.—The lower part of a pistil or carpel containing the ovules.

O'VATE.—Shaped like a lengthwise section of a hen's egg and having the attachment at the broader end.

O'VULE.—A transformed bud destined to become a seed after fertilization.

O'VUM.—The female sexual cell.

PAL'ATE.—A convex projection on the base of the lower lip of a personate corolla.

PA'LEA (*Pal'et*).—An inner bract of a Grass inflorescence which with the lemma incloses the flower.

PALEA'CEOUS.—Chaffy.

PAL'ISADE CELLS.—Radially-elongated cells with more or less indurated walls found in certain seed coats.

PAL'ISADE PAREN'CHYMA (*Palisade Tissue*).—Elongated mesophyll cells lying perpendicular to the surface of the leaf.

PAL'LID.—Pale.

PAL'MATE.—Divided or lobed in radiate fashion.

PALMAT'IFID.—Palmately-cleft.

PANDU'RIFORM.—Fiddle-shaped.

PAN'ICLE.—A compound raceme.

- PAPILLIONA'CEOUS.—Having butterfly-shaped flowers, as in the sub-family *Papilionaceæ* of the Leguminosæ.
- PAPIL'LAE.—Small, short, nipple-shaped protuberances of the epidermal cells.
- PAP'ILLOSE.—Bearing small, nipple-shaped protuberances.
- PAP'PUS.—The calyx of a Composite flower.
- PAPYRA'CEOUS.—Papery.
- PARAPH'YSIS.—A sterile filament found among reproductive organs in certain plants.
- PARASIT'IC.—Growing upon or within and deriving sustenance from another living organism.
- PAREN'CHYMA (Fundamental Tissue).—The soft, filling-in tissue of plants consisting of more or less isodiametric cells and usually with cellulose walls enclosing living contents.
- PARENCHY'MATOUS.—Referring to, consisting of or resembling parenchyma.
- PARI'ETAL.—Situating on or pertaining to the wall of an ovary or pericarp.
- PART'ED.—Incised nearly to the mid-rib or base.
- PARTHENOGEN'ESIS.—The production of an embryo from an unfertilized egg.
- PATHOL'OGY.—The study of diseases.
- PEC'TOSE.—Carbohydrate cell wall material related to the hemi-celluloses and to mucilage.
- PED'ATE.—Palmately parted or divided with two lateral lobes or divisions from each of which more or less linear divisions arise.
- PED'ICEL.—A branch of an inflorescence axis supporting a single flower.
- PEDUN'CLE.—The main stalk of an inflorescence. The flower stalk.
- PELLU'CID.—Transparent, clear.
- PEL'TATE.—Shield-shaped and attached by its lower surface to the support.
- PEN'DULOUS.—Hanging nearly vertically downward, as in the case of some ovules that hang from the sides of a locule.
- PENTAM'EROUS.—Applied to flowers having the number five or a multiple thereof running throughout each whorl.
- PENTAN'DROUS.—Having five stamens.
- PE'PO.—A fruit of a Cucurbit; a gourd.
- PEREN'NIAL.—Living more than two years.
- PER'FECTION.—Applied to flowers that contain both stamens and carpels that are functionally active.
- PERFO'LIATE.—Applied to leaves which are united around the stem at their base.
- PER'IANTH.—The floral envelopes, calyx and corolla or calyx alone when corolla is absent.
- PER'IBLEM.—A region of meristem lying between the dermatogen and plerome in the growing end of a root or stem. The meristem which gives rise to cortex.
- PERICAM'BIUM.—A zone of meristematic tissue lying just within the endodermis.
- PER'ICARP.—The wall of a ripened ovary or fruit surrounding the seed or seeds.
- PERICLA'DIUM.—A sheathing petiole.
- PER'ICYCLE.—A region lying between the endodermis and the conducting bundle or bundles.
- PER'IDERM.—All of the tissues produced by the phellogen, both externally and internally, and including the phellogen itself.
- PERID'IUM.—The outer covering of certain fungus fructifications as puff-balls.
- PER'IGONE.—See perianth.

- PERIG'YNOUS.—Applied to stamens and petals when they are adherent to the calyx throat, and so borne around the gynœcium.
- PER'ISPERM.—The nourishing tissue of some seeds outside of the embryo sac and representing the nucellus of the ovule, which during maturation has become laden with nutriment.
- PER'ISTOME.—The teeth around the mouth of the capsule in mosses.
- PERITHE'CIUM.—The receptacle containing asci in certain Ascomycetes.
- PERSIST'ENT.—Applied to parts of the flower which remain until the fruit ripens or to leaves which remain on the plant over winter.
- PER'SONATE.—Applied to a bilabiate corolla which has its throat closed by a convex projection on the base of the lower lip.
- PET'AL.—One of the floral leaves of the corolla.
- PET'ALOID.—Of some other color than green. Having the color of a petal.
- PET'IOLE.—A leaf stalk.
- PET'IOLOLE.—The stalk of a leaflet.
- PHEL'LODERM.—Secondary cortex containing chloroplasts and formed by the cork cambium or phellogen on its inner face.
- PHEL'LOGEN.—The meristem which gives rise to cork and frequently secondary cortex; cork cambium.
- PHLO'EM.—That part of a vascular bundle which contains sieve tissue and frequently bast (phloem) fibers.
- PHLOROGLU'CIN.—A white crystalline substance having the formula of $C_6H_6O_3$, obtained by the decomposition of phloretin and from certain gummy extracts and used with hydrochloric acid as a test for lignin.
- PHYCOCY'ANIN.—The blue pigment found in the Cyanophyceæ (Blue Green Algæ).
- PHYCOERYTH'RIN.—The red pigment occurring in the Rhodophyceæ (Red Algæ) and Cyanophyceæ (Blue-green Algæ).
- PHYCOPHÆ'IN.—The brown pigment found in the Phæophyceæ (Brown Algæ).
- PHYCOXAN'THIN.—A yellowish pigment occurring in some Algæ.
- PHYLLOCLA'DE.—A flattened branch which resembles a leaf, as in *Ruscus*.
- PHYL'LODE.—A dilated petiole which functions as a leaf blade.
- PHYL'LOTAXY.—The arrangement of leaves on stems.
- PHYLLOXAN'THIN.—See xanthophyll.
- PHYLOG'ENY.—The history of the race.
- PHYS'IOLOGY.—The science which treats of the functions of living organisms.
- PHYTO'GLOBULINS (Crystalloids).—Crystal-like bodies of protein composition found in aleurone grains.
- PHY'TON.—A term given by Gaudichaud to an internode with a node at its upper extremity which bears one or more leaves, in the axils of which buds may appear.
- PIG'MENTS.—Coloring substances found in cells.
- PI'LEUS.—The cap of a toadstool.
- PILIF'EROUS.—Bearing hairs.
- PILOSE'.—Covered with long, straight and scattered hairs.
- PIN'NATE.—Applied to compound leaves when the leaflets are arranged along the mid-rib.
- PINNAT'IFID.—Pinnately-cleft.
- PINNATIPAR'TITE.—Pinnately-parted.
- PINNAT'ISECT.—Pinnately-divided.

- PIN'NULE.—A secondary pinna.
- PI'SIFORM.—Pea-shaped.
- PIS'TIL.—The central female organ of a flower consisting of one or more united carpels.
- PIS'TILLATE.—Applied to flowers that possess one or more carpels but no fertile stamens.
- PITH (Medulla).—The core of soft parenchymatous tissue at the center of stems or roots.
- PLACEN'TA.—The nourishing tissue which connects the ovules with the wall of the ovary.
- PLACENTA'TION.—The arrangement of the placenta within the ovary or the pericarp.
- PLASMA MEM'BRANE (Outer Plasma Membrane).—See Ectoplasm.
- PLASMO'DIUM.—A multinucleated, naked mass of protoplasm having amoeboid movement. The vegetative body of a Slime Mold.
- PLASMOL'YSIS.—A contraction of the protoplasm of a cell due to the extraction of contained water under the influence of reagents of greater density than the protoplasmic sap.
- PLAS'TIDS.—Dense, porous protoplasmic bodies of various shapes scattered about in the cytoplasm.
- PLE'ROME.—A meristem found in the apical regions of plant axes which gives rise to fibrovascular tissue.
- PLI'CATE.—Folded like a fan.
- PLUMOSE'.—Feathery.
- PLU'MULE.—The rudimentary bud between the cotyledons.
- PLURILOC'ULAR.—Having more than one chamber or cell.
- PO'LAR BODY.—A portion of a gamete budded off before fertilization.
- POL'LEN.—The fertilizing dust composed of cells produced in the anthers of flowers.
- POLLEN GRAIN (Microspore).—A cell developed within the anther-sac and which carries the male fertilizing element.
- POLLINA'TION.—The transfer of pollen from anther to stigma and subsequent germination thereon.
- POLLIN'IUM.—A coherent mass of pollen grains in Orchids and Milkweeds, arranged as to be carried by insects.
- POLY.—A prefix of Greek origin meaning many.
- POLYADEL'PHOUS.—Applied to stamens which are united by their filaments into many sets.
- POLYAN'DROUS.—Having many stamens.
- POLYAN'THOUS.—Many flowered.
- POLYARCH'.—Said of a radial vascular bundle having many xylem and phloem rays.
- POLYCAR'PELLARY.—Composed of 3 or more carpels.
- POLYCAR'PIC.—Fruiting successively.
- POLYCEPHAL'IC.—Bearing many heads.
- POLYCOTYLED'ON.—A plant such as a Conifer which possesses more than 2 cotyledons or seed leaves.
- POLYEM'BRYONY.—Producing more than one embryo within a seed.
- POLYG'AMOUS.—Applied to species in which staminate, pistillate and hermaphrodite flowers are borne on the same plant.
- POLYG'ONAL.—Having several or many angles.

- POLYMOR'PHOUS.—Having several to many different forms.
- POLYPET'ALOUS.—Having distinct, disjoined petals.
- POLYPH'YLLOUS.—Many-leaved.
- POLYSEP'ALOUS.—Having distinct, disjoined sepals.
- POLYS'TACHOUS.—Having many spikes.
- POLYSTEM'ONOUS.—Possessing many more stamens than petals.
- POME.—A fleshy indehiscent fruit, two or more carpelled, with fibrous, cartilaginous or stony endocarp, the chief bulk of which consists of an adherent torus.
- PORE CANALS.—Pores through much thickened, usually lignified walls.
- PORES (Pits).—Small thin spots or small holes in the cell wall.
- PREFLORA'TION.—See Aestivation.
- PREFOLIA'TION.—See Vernation.
- PRICK'LE.—A sharp, rigid outgrowth from the epidermis and cortex.
- PRIMOR'DIAL.—First formed.
- PRIMOR'DIAL U'TRICLE.—The outer plasma membrane. The outer layer of protoplasm adjacent to the cell wall.
- PROCAM'BIUM.—The first formed fibrovascular tissue of any organ before differentiation has taken place into xylem and phloem.
- PROCUM'BENT.—Lying flat on the ground.
- PROEM'BRYO.—The primary stage in the development of *Chara* consisting of a single filament and a long rhizoidal cell. The suspensor in flowering plants.
- PROMYCE'LIUM.—A short hyphal growth from resting spores of smuts or rusts upon which basidiospores are borne.
- PROSEN'CHYMA.—Tissue composed of elongated, taper-pointed cells.
- PROTAN'DROUS.—A condition of hermaphrodite flowers in which the stamens mature before the carpels.
- PROTOG'YNOUS.—Applied to hermaphrodite flowers in which the carpels are mature before the stamens.
- PROTHAL'LUS (Prothal'lium).—A thalloid body bearing antheridia and archegonia, produced by the germination of a spore of a Pteridophyte into a protonema which later undergoes differentiation.
- PROTONE'MA.—A simple or branched green filament formed by the germination of a spore of a moss or fern.
- PROTOPHLO'EM.—The first-formed phloem elements in a fibrovascular bundle.
- PRO'TOPLASM.—Living matter.
- PRO'TOPLAST.—A term applied by Hanstein to the smallest body of protoplasm capable of individual action, either with or without a cell-wall, and either associated with other like units in a tissue or independent.
- PROTOXY'LEM.—The first formed elements of xylem in a fibrovascular bundle.
- PROX'IMAL.—Applied to the basal extremity. The attached end of an organ as opposed to the free or distal end.
- PSEUDO.—A prefix of Greek origin indicating spurious or false.
- PSEUDO-BULB.—The fleshy bulb-like internode of an epiphytic Orchid.
- PSEU'DOCARP.—A fruit which represents the product of the ripening of a single ovary as well as one or more accessory parts.
- PSEUDOPAREN'CHYMA (Plectenchyma).—A tissue found in the fruiting bodies of many fungi and consisting of compact hyphae somewhat resembling parenchyma in cross section.
- PUBER'ULENT.—Covered with a fine, soft, hairy coating.

- PUBES'CENT.—Covered with soft, short hairs.
- PULVI'NUS.—An enlargement at the base of the petiole or petiolule of some leaves or leaflets, as in numerous *Leguminosæ*.
- PUNC'TATE.—Dotted with small spots or minute pits.
- PUS'TULAR.—Applied to surfaces having blister- or pimple-like elevations.
- PUTA'MEN.—The stony endocarp of a drupe.
- PYRE'NOIDS.—Small, rounded, colorless, refractile granules embedded in the chromatophores of numerous *Algæ* and thought to be starch forming centers.
- PYX'IS.—A capsule which dehisces transversely into pot and lid portions.
- QUAD- or QUADRI.—A prefix of Latin origin signifying four.
- QUADRAN'GULAR.—Four-angled.
- QUADRIFO'LIATE.—Applied to palmate leaves which have four leaflets arising from the summit of the petiole.
- QUINQUEFOL'iate.—Applied to any compound leaf that has five leaflets.
- RACEME'.—An indeterminate inflorescence having pedicelled flowers arranged along a lengthened axis.
- RAC'EMOSE.—Arranged in racemes.
- RA'CHIS.—The extended portion of a peduncle.
- RAD'ICAL.—Arising from the root or base of the stem.
- RAD'ICLE.—The rudimentary root of an embryo plantlet.
- RA'MAL.—Pertaining to a branch.
- RA'MUS.—A branch.
- RAMOSE'.—Branching.
- RANK.—A row of leaves or other organs arranged vertically on a stem.
- RA'PHE (RHA'PHE).—The adherent portion of the ovule stalk in inverted and half inverted ovules and seeds.
- RAPH'IDES.—Bundles of needle-shaped crystals.
- RECEP'TACLE.—The shortened stem upon which the whorls of floral leaves are inserted.
- RECEPTAC'ULAR.—Pertaining to the receptacle.
- REC'LINATE.—Bent downward.
- RECLIN'ING.—See Reclinate.
- RECURVED'.—Curved outward or backward to a moderate extent.
- REFLEXED'.—Turned outward or backward more abruptly than Recurved.
- REG'MA.—A capsular fruit of 2 or more carpels that first splits into separate parts and then each of these dehisces.
- REJUVENES'CENCE.—Applied to a mode of reproduction in which the protoplasm of the cell becomes rounded out, escapes by rupture of the cell wall, forms cilia and moves about, in time developing into a new plant.
- REN'IFORM.—Kidney-shaped.
- REPAND'.—Having a slightly undulating margin.
- RE'PENT.—Creeping.
- RE'PLUM.—A spurious membranous septum seen in Cruciferous fruits that persists after the valves have fallen away.
- RETIC'ULATE.—Applied to markings or veins which are in the form of a network.
- RETUSE'.—Having a broad, shallow sinus at the apex.
- REV'OLUTE.—Said of leaves in the bud when their margins are rolled backward.

- RHA'PHE.—See Raphe.
- RHI'ZIDS.—Absorption organs of certain plants that are analogous with roots of higher plants but which consist of outgrowths of epidermal cells.
- RHIZOME'.—A creeping underground stem.
- RHI'ZOMORPHS.—Root-like structures composed of united hyphæ and seen in certain fungi.
- RIB.—A prominent vein or ridge.
- RIN'GENT.—Applied to the corolla of a bilabiate type whose throat is open and lips separated.
- RIPA'RIOUS.—Growing along the banks of rivers or other water-courses.
- ROSETTE'.—A cluster of leaves or other organs.
- ROS'TRATE.—Beaked.
- ROSTEL'LUM.—A process found on the column of orchid flowers which supports the disks of the pollinia.
- RO'TATE.—Wheel-shaped.
- ROTUND'.—Rounded in outline.
- RU'FOUS.—Brownish-red.
- RUGOSE'.—Wrinkled.
- RU'MINATE.—Applied to the albumen of certain seeds when the perisperm is found coursing through the endosperm in irregular fashion.
- RUN'CINATE.—Applied to a pinnately-cleft leaf whose lobes are directed backward as in the Dandelion.
- RUN'NER.—A stem or branch which roots at intervals as it trails along the ground.
- SAC'CATE.—Pouch-like.
- SAG'ITTATE.—Arrow-shaped.
- SAMA'RA.—A winged fruit.
- SAP'ROPHYTE.—An organism that lives upon decaying or dead organic matter.
- SAR'COCARP.—The fleshy portion of a drupe or other fruit.
- SCA'BROUS.—Said of leaves, etc., that are rough or harsh to the touch.
- SCALAR'IFORM.—Applied to tracheæ or tracheids whose walls show transversely arranged bars, resembling the rungs of a ladder.
- SCAN'DENT.—Climbing.
- SCAPE.—A naked peduncle arising from a root or underground stem.
- SCA'RIOUS.—Dry and membranous.
- SCHIZ'OCARP.—A fruit that separates when mature into 2 or more indehiscent parts.
- SCHIZOG'ENOUS.—Said of intercellular-air-spaces or of reservoirs that are formed by the breaking down of the middle lamellæ of cells where several come together and the later separation of the cells at these places.
- SCI'ON.—A shoot intended for grafting.
- SCLEREN'CHYMA.—A tissue composed of stone cells, fibers or both.
- SCLERO'TIUM.—A hardened mass of mycelium.
- SCOR'PIOID.—Applied to certain cymes whose flowers are situated on alternate sides of the floral axis.
- SCUTEL'LUM.—A shield-shaped portion of the cotyledon of *Gramineæ*, which absorbs nourishment from the endosperm during germination and bales it out to the rest of the embryo.
- SEC'UNDINE.—The outer coat of the ovule.
- SEED.—A fertilized and matured ovule containing an embryo.

- SE'PAL.—A leaf of the calyx.
- SEP'TATE.—Possessing one or more partitions.
- SEPTICI'DAL.—A mode of dehiscence in which the opening occurs along the line of junction of the carpels.
- SEPTIFRA'GAL.—A method of dehiscence in which the valves of a capsular fruit break away from the partitions or septa.
- SEP'TUM.—A partition between cavities in an ovary or fruit or between cells in a tissue.
- SERIC'EUS.—Silky. Having a covering of fine, soft, appressed, silky, hairs.
- SER'RATE.—Toothed with teeth projecting toward the apex.
- SER'RULATE.—Finely serrate.
- SES'SILE.—Without a stalk.
- SE'TA.—A bristle-like structure.
- SETIG'EROUS.—Bristle bearing.
- SIL'ICULE.—A short silique.
- SIL'IQUE.—The characteristic fruit of the *Cruciferae*, consisting of a capsule of 2 valves which separate from the replum in dehiscence.
- SIN'UATE.—Wavy margined.
- SOFT BAST.—The unligified portion of the phloem.
- SOMAT'IC CELLS.—The body cells of an individual, in distinction from reproductive cells.
- SORE'DIUM.—A scale-like structure found on many lichens and consisting of a group of algæ cells surrounded by a network of hyphæ. When detached from the parent-plant it has the power of developing vegetatively into a mature lichen.
- SORO'SIS.—A multiple fruit, as represented by the Mulberry and Osage Orange, consisting of a swollen up, condensed and mature spike.
- SO'RUS.—An aggregation of sporangia.
- SPA'DIX.—A fleshy spike more or less surrounded by a bract called a spathe.
- SPATHE.—A large bract that encloses or subtends an inflorescence.
- SPAT'ULATE.—Said of flat leaves that are narrow at the base and become gradually broader toward the summit, which is rounded.
- SPERMA'TOPHYTE.—A seed plant.
- SPERMATOZO'ID.—A male sexual cell. See Antherozoid.
- SPERMATOZO'ON.—Another name for Spermatozoid or Antherozoid.
- SPER'MODERM.—The covering of the seed.
- SPHACE'LIA.—The conidia stage of *Claviceps*.
- SPIC'ATE.—Arranged in a spike.
- SPIC'ULE.—A small pointed outgrowth. A needle-shaped crystal.
- SPIKE.—An indeterminate inflorescence consisting of sessile florets arranged along a lengthened axis.
- SPIKE'LET.—A secondary spike.
- SPINE.—A sharp, rigid termination of a branch as in the Honey Locust. A thorn.
- SPINES'CENT.—Spiny in structure.
- SPORAD'IC.—Scattered.
- SPORAN'GIOPHORE.—The stalk or support of a sporangium.
- SPORAN'GIUM.—A spore case.
- SPORE.—An asexual or sexual reproductive cell usually with a highly resistant cell wall.
- SPOROGO'NIUM.—The asexual generation in Bryophytes and Pteridophytes.

SPO'ROPHYLL.—A spore bearing leaf.

SPUR.—A tubular or saccate appendage of some part of the flower, usually containing nectar.

SQUAMOSE'.—Scale-like.

STA'MEN.—A male organ of the flower producing pollen.

STAM'INODE.—An abortive and sterile stamen, or any body without an anther occupying the normal place of a stamen.

STEG'MA (pl. stegma'ta).—A cell containing a warty silicious body adherent to the surface of a sclerenchyma fiber, and occurring in various palms, orchids, the *Scitamineæ*, etc.

STE'LE.—All the structures within the usual type of endodermis.

STEL'LATE.—Star-shaped.

STEM.—The ascending axis of a plant bearing leaves or leaf modifications.

STE'REOME.—The supporting elements of a fibrovascular bundle.

STER'ILE.—1. Unproductive, as a stamen without anther, flower without pistil, or pericarp without seeds. 2. Devoid of living organisms.

STERILIZA'TION.—The process of ridng an object of all living organisms.

STIG'MA.—That part of a pistil or carpel which receives the pollen.

STIPE.—The stem of a moss; the stalk of a fern frond; the stalk of a toadstool or other fungus.

STIP'ULATE.—Possessing stipules.

STIP'ULE.—A modified leaf, usually blade-like and situated at the base of the leaf-stalk.

STO'LON.—A slender running branch above or below the surface of the soil, either capable of taking root or bearing a bulb at its end.

STOLONIF'EROUS.—Bearing stolons.

STO'MA (Stomate).—A minute opening for the passage of gases and water vapor in the epidermis, surrounded by two guard cells, which usually have the power to enlarge and reduce the opening. Also employed for the opening in the epidermis with the guard and accessory cells.

STOM'ATA.—Plural for stoma.

STOMAT'AL APPARA'TUS.—The stoma with its surrounding guard cells and neighboring cells.

STOMAT'AL CHAM'BER.—The intercellular-air-space directly beneath the stoma.

STRAND.—Applied to any tissue or group of tissues forming a narrow, more or less cylindrical mass extending longitudinally through the plant or plant organ.

STRI'ATE.—Marked with fine longitudinal lines or grooves.

STRIGOSE'.—Covered with sharp and rigid appressed hairs.

STROB'ILE.—A scaly multiple fruit consisting of a scale-bearing axis, each scale of which encloses one or more seeds. A cone.

STONE CELLS (Sclerenchyma Cells, Sclereids).—Isodiametric or moderately elongated parenchyma cells modified by thickenings, lignification of the cell wall and loss of protoplasmic contents.

STYLE.—That portion of a pistil connecting the ovary with the stigma.

STYLOPO'DIUM.—The fleshy disk directly above the ovarian portion of an Umbelliferous fruit, formed by the expansion of the bases of the two styles.

SUB.—A prefix of Latin origin meaning under, below, subordinate, nearly or partially.

SU'BER.—Cork tissue.

- SUBERIN.—A fatty cell wall constituent serving to water-proof the wall.
- SUBERI'ZED (Corky).—Applied to a cell wall infiltrated with suberin.
- SUBTERRA'NEAN.—Beneath the surface of the soil.
- SU'BULATE.—Narrow and tapering to an acute end.
- SUC'ULENT.—Soft and juicy or fleshy.
- SUC'KER.—A shoot from the root or lower part of the stem or underground stem.
- SUFFRU'TICOSE.—Applied to stems or plants that are woody at their base and herbaceous above.
- SUL'CATE.—Having longitudinal grooves.
- SUPE'RIOR.—Said of an ovary that is not adherent to and above the calyx; also applied to a calyx which is situated on the upgrown receptacle above the ovary or to a tubular calyx whose limb appears to spring from the top of the ovary.
- SUSPEN'SOR.—A row of cells, representing the first development of the fertilized egg of a seed plant, upon the end of which an embryo is formed.
- SU'TURE.—The line of union of two carpels. The line of dehiscence.
- SWARM SPORE.—A spore which possesses one or more cilia for movement.
- SYCON'IUM.—The characteristic multiple fruit of the Fig, which consists of a fleshy, invaginated receptacle bearing numerous akenes.
- SYMBIO'SIS.—The living together of two individuals having a communion of life interests.
- SYMMET'RICAL.—Said of flowers when the parts of each whorl are of the same number or multiples of the same number.
- SYMPET'ALOUS.—See Gamopetalous.
- SYM'PHYSIS.—A union of parts.
- SYNAPSIS.—The union in pairs of homologous chromosomes, respectively of maternal and paternal origin, to form bivalents.
- SYNCAR'POUS.—Said of fruits and gynoecia when they are formed of two or more united carpels.
- SYNER'GIDS.—Two nuclei in the upper region of the embryo sac above the egg nucleus.
- SYNGENE'SIOUS.—Said of stamens when their anthers are united.
- SYN'ONYM.—Another name for the same thing.
- SYNPHYLL'OUS.—Applied to sepals or petals which are united by their margins.
- SYNSEP'ALOUS.—See Gamosepalous.
- TAB'ULAR.—Flattened from above downward.
- TANGENT'IAL.—Applied to cell walls more or less parallel to a tangential plane of the plant organ. The walls which are at right angles with the radial walls.
- TAPE'TUM.—A layer of cells lining the cavity of an anther sac.
- TAP-ROOT.—The main root coursing directly downward.
- TAXON'OMY.—The science of classification.
- TEG'MEN.—The inner seed coat.
- TELEU'TOSPORE.—A spore produced by the Rusts toward the close of the season which forms a promycelium the next year.
- TEN'DRIL.—A modified stem, stipule, leaf, or leaflet which has taken on the form of a slender appendage that is capable of coiling spirally around some object.
- TERATOL'OGY.—The study of monstrosities.
- TERETE'.—More or less cylindrical.

- TER'MINAL.—Pertaining to the end or apex.
- TER'NATE.—In threes.
- TERRES'TRIAL.—Growing on land.
- TES'SELLATED.—Marked like a checkerboard.
- TES'TA.—The outer seed coat.
- TETRA.—A prefix of Greek origin signifying four.
- TET'RAD.—A group of four immature pollen grains cut off by pollen mother cells.
- TETRACAR'PELLARY.—Having four carpels.
- TETRADYN'AMOUS.—Having six stamens, four of which are longer than the other two.
- TETRAG'ONAL.—Four-angled.
- TETRAM'EROUS.—Said of flowers that have the number four or multiple thereof running through their various whorls.
- TETRAN'DROUS.—Having four stamens.
- TETRAPET'ALOUS.—Having four petals.
- TETRASEP'ALOUS.—Having four sepals.
- TE'TRARCH.—Said of a radial vascular bundle having 4 xylem and 4 phloem arms alternating with one another.
- TET'RASPORES.—Applied to the asexually produced spores of the *Florideæ* group of Red Algæ on account of being formed in groups of four in the mother cell.
- TETRAS'TICHOUS.—Said of leaves when they are arranged in four vertical rows upon a stem.
- THAL'AMUS.—Another name for receptacle.
- THAL'LUS.—A plant body showing no differentiation into root, stem, or leaf.
- THERMOT'ROPISM.—Response of living matter to the stimulus of heat or cold.
- THORN.—See Spine.
- THROAT.—The opening into the tube of a gamosepalous calyx or gamopetalous corolla.
- THYR'SUS.—A compact panicle of flowers like the Lilac or Sumac.
- TIS'SUE.—An aggregation of cells of similar source, structure and function in intimate union.
- TO'MENTOSE.—Covered with dense, matted, wooly hairs.
- TOR'TUOUS.—Bent or twisted irregularly.
- TO'RUS.—Another name for receptacle.
- TRA'CHEA.—An elongated cylindrical or prismatic tube with open ends, found in the xylem of the fibrovascular system and serving for the conduction of crude sap.
- TRA'CHEID.—An elongated, taper ended, tubular cell with closed ends and pitted walls found in the xylem and serving for water conduction and support.
- TRANSPIRA'TION.—The giving off of watery vapor by the plant.
- TRI.—Three.
- TRIADEL'PHOUS.—Having the filaments in 3 sets.
- TRIAN'DROUS.—Possessing three stamens.
- TRI'ARCH.—Applied to a radial fibrovascular bundle having three xylem and three phloem arms alternating with one another.
- TRICAR'PELLARY.—Possessing three carpels.
- TRICH'OBLAST.—An internal hair, like those projecting into the intercellular-air-spaces of the stems of certain Water Lilies.
- TRICH'OGYNE.—A slender appendage to the carpogonium.

- TRICH'OME.—A plant hair.
- TRICHOT'OMOUS.—Three-branched or forked.
- TRIFO'LIATE.—Said of a compound leaf having three leaflets.
- TRIMOR'PHOUS.—Possessing three kinds of hermaphrodite flowers in the same species, differing in the relative length of their stamens and carpels.
- TRIS'TICHOUS.—Three ranked.
- TRITER'NATE.—Applied to a compound leaf whose petiole divides into three secondary petioles, each of which again divides into three tertiary petioles, each division bearing 3 leaflets.
- TRUN'CATE.—Ending abruptly as if cut off or flattened at the summit.
- TU'BER.—A short excessively thickened end of an underground stem.
- TU'BERCLE.—A small wart-like outgrowth upon the rootlets, roots or subterranean stems of various plants.
- TU'BEROUS.—Bearing or resembling tubers.
- TU'NICATED.—Covered with successively overlapping coats as the bulb of an Onion.
- TUR'BINATE.—Top-shaped.
- TURGES'CENT.—Swelling.
- TYLO'SIS.—A protrusion of the wall of a cell through the pit in the wall of an adjacent vessel and appearing in the cavity of the latter.
- TYPE.—An individual possessing the essential characteristics of the group to which it belongs.
- UM'BEL.—The typical inflorescence of the family *Umbelliferae*. A more or less flat-topped indeterminate inflorescence in which the pedicels spread like the stays of an umbrella.
- UN'CIFORM.—Hook-shaped.
- UN'DERSHRUB.—A low shrub-like plant whose base is woody and upper portion herbaceous.
- UN'DULATE.—Having a wavy margin.
- UNI.—A prefix of Latin origin meaning one.
- UNILAT'ERAL.—One-sided.
- UNILOC'ULAR.—One-celled.
- UNISE'RIATE.—Arranged in a single row, as the cells of some plant hairs.
- UR'CEOLATE.—Urn-shaped.
- URE'DOSPORE.—A one-celled spore produced during the life history of a Rust.
- U'TRICLE.—An akene with a bladdery pericarp or calyx as *Chenopodium* fruit.
- VAC'UOLE.—A cavity within the protoplasm of a cell usually containing cell sap.
- VALV'ATE.—Applied to the leaves of a flower in the bud stage when their margins meet but do not overlap.
- VALVE.—One of the halves of a diatom. One of the parts of a pericarp that splits open when ripe.
- VARIA'TION.—The occurrence of differences among the individuals of a species or variety.
- VARI'ETY.—A group of individuals within a species that differ in some minor respect from the rest of the species.
- VAS'CULAR BUN'DLE.—A group of tubular cells frequently combined with fibers or other kinds of cells coursing through the plant and serving primarily for conduction and secondarily for mechanical support, storage, etc.

- VAS'CULUM.—A collecting case used by botanists.
- VEG'ETABLE.—A plant.
- VEIN.—A strand of fibrovascular tissue in a leaf.
- VELA'MEN.—An absorptive tissue composed of several layers of dead cells covering the roots of some tropical epiphytic orchids and aroids.
- VENA'TION.—The arrangement of veins in a leaf.
- VEN'TER.—The enlarged basal portion of an archegonium.
- VEN'TRAL CANAL' CELL.—A cell beneath the entrance of the neck portion of an archegonium.
- VERMIC'ULAR.—Worm-shaped.
- VERNA'TION.—The manner in which leaves are disposed in the bud.
- VER'RUCOSE.—Wart-like.
- VERTICILLAS'TER.—A pair of dense cymes in the axils of opposite leaves.
- VERTIC'ILLATE.—Whorled.
- VES'SEL.—See trachea.
- VILLOSE'.—Covered with soft, thin, rather straight hairs.
- VIRIDES'CENT.—Greenish.
- VIS'CID.—Sticky.
- VITTA.—An oil tube in the fruit of an Umbelliferous plant.
- VOLATILE OIL.—An odorous, liquid hydrocarbon not leaving a greasy stain on filter paper.
- VOL'VA.—The covering extending from the margin of the cap to the base of the stipe in some toadstools, which is ruptured during growth and adheres around the bulbous base.
- WATER PORES.—See Hydathodes.
- WAXES.—Esters of the fatty acids, other than glyceryl esters, usually found in plants as a layer over the cuticle of various stems, leaves and fruits.
- WOOD.—All that portion of a woody exogenous plant axis lying inside of the cambium line; also applied to the xylem region of conducting bundles.
- WOOD CIRCLE.—In exogenous stems and roots of secondary structure, all the region within the cambium line except the pith and internal phloem if present, and consisting of the xylem and the xylem portion of medullary rays.
- WOOD FIBERS.—Fibers found in the xylem.
- WOOD RAY (Xylem Ray).—All that portion of a medullary ray between two wood wedges.
- WOOD WEDGES (Xylem Wedges).—That portion of the xylem region between two wood medullary rays.
- XY'LEM.—That portion of a vascular bundle which contains tracheæ or tracheids or both and often wood parenchyma and fibers.
- ZOÖGLOE'A.—A gelatinous mass of bacteria.
- ZO'ÖSPORE.—A ciliated spore having the power of movement.
- ZYGOMOR'PHIC.—Applied to irregular flowers like those of the Broom and Pea which can only be divided into similar halves in one direction. cf. Actinomorphic.
- ZYG'OSPORE.—A spore resulting from the union of two like gametes.
- ZY'GOTE.—A cell resulting from the union of two sexual cells or their nuclei.
- ZY'MOGEN.—A microorganism capable of producing fermentation.

Index

- Abies*, 465
Abies balsamea, 466
Abies sibirica, 466
Abrus precatorius, 540
Absinthium, 615
Absorption of nutrient salts, 155
 selective, 155
Acacia, 301, 536, **539**
Acacia Catechu, 540
Acacia Senegal, 95, **539**
Acajou Gum, 552
Acanthaceæ, 602
Acanthus Family, 91, 602
Accretion, defined, **27**
Acer, 554
Acer spicatum, 554
Aceraceæ, 554
Acetic acid, 680
Achillea, 615
Achillea millefolium, 615
Achras Sapota, 582
Acid alcohol, 680
Acidum Gallicum, 507
Acidum Tannicum, 507
Aconite, 162, **518**
Aconiti Folia, 519
Aconitine, micro-chemic test for, 86
Aconitum, 518
 calyx of, 266
 fruit of, 301
 galea of, 268
 histology of, 175
Aconitum Napellus, 86, 518, **519**
Aconitum Napellus var. *roseum*, Fig. of
 cross sect. of stem, 130
Aconitum U.S.P., 519
Acorus Calamus, **487**, 618
 Fig. of, 488
 Fig. of cross section of root, 144
Actæa, 518
Actinomyces, 372
Actinomyces bovis, 373
Actinomyces Myricarum, **179**, 373, 500
Adder's Tongue, 434
Adhesion, 268
Adiantum, 434, **435**
Adiantum Capillus-Veneris, 436
Adiantum pedatum, 436
Adnation, 265
Adonis N. F., 519
Adonis vernalis, 519
Æcidiospores, 401
Æcidium, 401
Æcium, 401
Ægle Marmelos, 546
Aerobe, 34
Aerogens, 364
Æthallia, 374
Agar, 361
Agaricaceæ, 405
Agaricales, 403
Agaricus, 404
Agaricus campestris, 405
Agathis australis, 466
Agathis loranthifolia, 466
Agave, 619
Agave americana, 75
Agavose, **75**
Aggregation-body, 243
Agricultural Botany, defined, 3
Agrimonia, 535
Agronomy, defined, 3
Agropyron repens, 485
 Fig. of cross section of stem, 219
Ailanthus Family, 546
Aizoaceæ, 514
Akene, 303
Albumen, defined, 314
 endospermic, 314
 mode of formation, 314
 perispermic, 314
Albumens, 93
Alburnum, 213
Alcaligenes melitensis, 373

- Alcanna tincture, 680
 Alcohol, 680
 N-butyl, 696
 Alder, 504
 Black, 552
 Alder Buckthorn, 103, **555**
 Aldrovanda, 530
 vesiculosa, 628
 Alectoria, 417
 Aletris, 490
 Aletris *farinosa*, 490
 Aleurites *cordata*, 551
 Aleurone grains, 93
 Alga-like Fungi, 376
 Algæ, **333**
 blue green, **334**, 413
 brown, 351
 green, **336**, 413
 red, 357
 siphon, 346
 Alkaloidal reagents, 84, 85
 Alkaloids, 84-87, 397
 Alkanet, 593
 Alkanna *tinctoria*, 593
 Allamanda, 587
 Allelomorphs, 644, 647
 Allium, 491
 Allium sativum, 491
 Fig. of bulb of, 191
 Almond, 535
 Alnus, 504
 Aloe, 490
 Aloe, 84, 220, 241
 ferox, 490
 Perryi, 490
 vera, 490
 Aloin, 84, **490**
 Alpinia *officinarum*, 492
 Alternation of generations, 71, 423
 Althæa, 561
 Flores, 562
 Folia, 562
 mucilage in, 95
 Althæa *officinalis*, 88
 rosea, 562
 Alum root, 531
 Alyssum, 277
 Amanita, 407
 deadly, 408
 Amanita, toxin, 409
 Amanita muscaria, 407, **408**, **409**
 phalloides, 408, **409**
 Amanitahemolysin, 409
 Amaranthaceæ, 514
 pollen of, 283
 Amber, 466
 Ambrette seed, 562
 Ambrosia, 283
 Amelanchier, 535
 Ament, 259
 Amino-acids, 93
 Ammonia water, 681
 Ammoniacum, 577
 Amœba, **24**
 Figs. of, 25, 27
 Amœboid movement, **26**
 Amomum *melegueta*, 492
 Amygdala Amara, 536
 Amygdala Dulcis, 536
 Amygdalaceæ, 535
 Amygdalin, 103
 Amygdalus, 535
 communis, **535**, **536**
 Amylase, 76
 Amylodextrin, 82
 Amyloplastids, 42
 Amylum, 485
 Amylum Marantæ, 493
 Anabæna, 335, **336**
 Anabolism, defined, **31**
 Anacardiaceæ, 551
 Anacardium, 552
 Anacardium *occidentale*, 552
 Anacyclus *Pyrethrum*, 615
 Anærobe, **34**
 Anamirta *Cocculus*, 521
 Anaphase, 48, **50**
 Anaphalis *margaritacea*, 615
 Andræa, 426
 Andræales, 426
 Androcæcium, 262, **274**
 Andromeda, 580
 Andropogon *squarrosus*, 485
 Anemone, 260, 304
 Anethol, 576
 Angelica *Archangelica*, 577
 Angelicæ Fructus, 577
 Angelicæ Radix, 577

- Angiosperm, development of female gametophyte, 292, 474
 development of male gametophyte, 292, 474
 life history of an, 471
 Angiospermæ, 443
 Angiosperms, 15, 443, **471**
 classes of, 479
 distinctions from gymnosperms, 478
 fertilization in, 292
 resemblances between and gymnosperms, 478
 Aniline, blue, 681
 brown, 681
 chloride, 681
 sulphate, 681
 Anise, **258, 576**
 Anisum, 576
 Annatto, 565
 Annual plants, 162
 Annual rings, 205, **206**
 thickening, 205
 Annulus, **67, 407, 433**
 false, 409
 true, 407
 Anonaceæ, 515
 Anthemis, 615
Anthemis nobilis, 615
 Anther, 274, **277**
 adnate, **276, 282**
 attachment of, 282
 confluent, 276
 dehiscence, **279, 475**
 development of, 279
 extrorse, 282
 Figs. of cross sect. of, 278, 280
 gross structure of, 277
 histology of, 277
 innate, 276
 introrse, 282
 lobes, 277
 syngenesious, 276
 versatile, 276, **282**
 Antheridia, 346, 348, 355, 426
 Antheridium, 358
 Antherozoids, 339, 441
 Anthocarp, 298
Anthoceros, 419
Anthocerotales, 419
 Anthocyanins, 100
 Anthophore, 266
 Anthotaxy, 256
Antigonon leptopus, 512
 Antipodals, 292
 Apetalæ, 499
 Apii Fructus, 577
 Apiol, 577
Apium graveolens, 577
Apocynaceæ, 587
 Apocynum, 589
Apocynum cannabinum, **587, 589**
 Apothecia, 388
 Apothecium, 388, **416**
 Appendages, epidermal, 124
 Apple, 305, 534
 Fig. of flower, 271
 Fig. of fruit, 306
 Rust, 402
 Appressorium, 378
 Apricot, 534
Aquifoliaceæ, 552
Aquilegia, 269
Arabis, 528
Araceæ, 487
 Arales, 487
 Aralia, 575
Aralia hispida, 575
 nudicaulis, 575, **576**
 racemosa, 575
 *spinos*a, 575, **576**
Araliaceæ, 575
 Arbor vitæ, 465
Arbutus unedo, 579
 Archegonia, 422, 426
 of pine, 451, 452, 453
 Archegonium, 69, 418
 Archichlamydeæ, 499
Arctium Lappa, 615
 minus, **612, 615**
Arctostaphylos, 580
 manzanita, 582
Uva Ursi, 580, **581**
 Fig. of, 581
Areca Catechu, 487
 Areca, 487
 Arecolinæ Hydrobromidum, 487
 Aril, 313

- Arillode, 314
Arisæma triphyllum, 487
Aristolochia, 275, 276, 511
 reticulata, 512
 Serpentaria, 512
 sipho, 199, 200
Aristolochiaceæ, 510
 Armoracia, 530
 Arnica, 614
Arnica montana, **614**, **615**
Arnicae Radix, 615
Aronia, 535
 Arrowroot Family, 314, **492**
Artemisia Absinthium, 615
 Cina, 615
Artemisia, pollen of, 283
 Arthrospores, 369
 Arum Family, 487
Arum triphyllum, 487
Arundinaria, 484
Asafœtida, 577
Asagræa officinalis, 490, 491
Asarum, N. F., 512
Asarum, 149, 511
 canadensis, 512
Asarum, flower of, 511
 tyloses of, 213
Asclepiadaceæ, 589
Asclepias, 589
Asclepias
 latex cells in, 133
 pollinia of, 283
Asclepias tuberosa, 589
Ascocarp, 394, 395
 Asco-lichens, 413
 Ascomycetes, 375, **381**
Ascophyllum, 355
 Ascospores, 381, 385
 Ascus, 381
 Fig. of, 387
 Ash, 303, 583
 Black, 121, 122
 Mountain, 74, 535
 Asparagine, 88
Asparagus, 88, 188, **490**
Aspergillus, 388, 391
 differences from *Pencillium*, 391
 fumigatus, 392
 glauca, 391
Aspergillus herbariorum, 391
 niger, **393**, 394
 oryza, 102, **391**, 392
Asperula odorata, 605
 Aspidistra, 291
 Aspidium, **53**, 435
 American, 434
 Aspidosperma, 589
Aspidosperma Quebracho-blanco, 589
Asplenium, 435
 Assimilation, defined, 225
 parenchyma, 116
Asteraceæ, 610
Astragalus, 95, 539
Astragalus gummifer, 95, **539**
Atriplex, 514
Atropa, 596
Atropa Belladonna, 88, **597**, **598**
 Atropina, 597
Aurantieæ, 545
Aurantii Dulcis Cortex, 546
 Amari Cortex, 546
 Auricle, 274
 Auriculariales, 402
 Autecology, 616
 Auxin, 104
 Auxospore, 350, **351**
 Avena, N. F., 485
Avena sativa, 485
 Awn, 274, 313
 Azalea, 579, 580
 apples, 403
 Azedarach, 548
Azolla, 437

 Bacca, 305
 Bacillariæ, 348
 Bacilli, anthrax, 368
 types of, 370
 Bacillus, 365, **372**
 Bacteria, 362, **363**
 aërobic, 34
 amphitrichous, 366
 anaërobic, 34
 chemical composition of, 369
 classification of, accd. to phenomena, 364
 morphological, 370
 denitrifying, 367

- Bacteria, disease-producing, 373
 forms of, 355
 higher, 372
 iron, 373, **374**
 lophotrichous, 366
 lower, 370
 monotrichous, 366
 morphology due to cleavage, 367
 motility of, 365
 nitrifying, **182**, 367
 nutrition of, 366
 parasitic, 366
 pathogenic, 374
 peritrichous, 366
 products of, 367
 rapidity of multiplication, 369
 reproduction of, 367
 saprophytic, 366
 sporulation, 368
 sulfur, 372, **374**
 thermophilic, 364
Bacteriaceæ, 371
 Bacterial cultures, 364
 Bacteriology, defined, 3
Bacterium, 372
 Balata, 582
 Bald Cypress, 619
 Balm, 596
 Balmony, 600
 Balsam Apple, 608
 Balsams, 97, **98**
 Balsamum Peruvianum, 539
 Balsamum Tolutanum, 539
 Bamboo, 484
 Banana, 305
 Banana Family, 492
 Baneberry, 518
 Banyan, 161
 Baptisia, 540
Baptisia tinctoria, 540
 Barbaloin, 84
 Barberry, 519
 rust on, 401
 Barberry Family, 515, **519**
 Bark, defined, 206
 Figs. of sections of stem-bark of
 Cascara Sagrada, 208, 209,
 210
 histology of a typical, 208
 Bark, scale, 130
 shell, 130
 Barley, 75, 259
 Fig. of leaf and stem, 243
 Fig. of longitudinal section through
 root-tip of, 156
 fruit, 304
 starch of, 78, **81**
Barosma, 330, **545**
 betulina, **545**, **546**
 crenulata, 546
 serratifolia, 546
 Basidiomycetes, 375, **397**
 Basidiospore, 397
 Basidium, defined, 397
 Basswood, 559
 Bast, fibers, 122
 hard, 202
 Bast, soft, 202
Batrachospermum, 357
 Bauhin, G., 324
 Bay Leaves, 574
 Bayberry Family, 176, **500**
 Bayberry wax, **96**, 502
 Bean Family, 176
 Bean, Fig. of, 313
 starch of, 82
 Bearberry, 580
 Bear's Foot Root, 519
 Beauty-berry, 593
 Beech, 245, 304
 Beech Family, 506
 Beets, 514
Beggiatoa, 374
Beggiatoaceæ, 374
 Begonia, adventitious buds of, 185
 roots of, 160
 Fig. of cross sect. of leaf stalk, 117
Begoniaceæ, 221
 Belæ Fructus, 546
 Belladonna, fruit of, 304
 Belladonnæ Folium, 597
 Belladonnæ Radix, 597
 Bentham, G., 326
 Benthos, 617
 Benzoin, 97
 Benzoin Family, 582
 Benzoinum, 583
Berberidaceæ, 279, 515, **519**

- Berberis, response of stamens to stimulation, 36
 Berberis N. F., 519
 Berry, defined, 304
 Beta, 514
 maritima, 514
 vulgaris, 514
 Betula, 504
 Betula alba, 505
 Betula lenta, 504, 505
 Betula populifolia, 204
 Betulaceæ, 504
 Biennial plants, 163
 Bifacial leaf blade, 250, 251
 Binary division, 45
 Binary fission, 28
 Binomial system, 325
 Biology, defined, 1
 Birch Bark, 505
 Birch Family, 504
 Birthwort Family, 510
 Bismark Brown, 681
 Bistorta, 513
 Bitter Almond, 103
 Bittersweet, 188, 189, 596, 599
 Climbing, 552
 Bixa Orellana, 565
 Bixaceæ, 565
 Black Alder, 552
 Black Caraway, 519
 Black Fungi, 395
 Black Mold, 377
 Black Mustard, 529
 Blackberry, 186, 300, 535
 Bladder Wrack, 355
 Bladderworts, 638, 639
 Blakeslee, 357
 Blessed Thistle, 615
 Bloodroot, 186, 189, 524, 526
 Blue Cohosh, 522
 Blue Flag, 492
 Blue Green Algæ, 334
 mucilage of, 95
 Blue Vervain, 593
 Bluebell Family, 103, 609, 610
 Blueberry, 580, 581
 Blue-weed, 592
 Bock, H., 324
 Bog Mosses, 425
 Bogbean, 587
 Boldo, 248, 522
 Boldo Family, 522
 Boldus, 522
 Boletus, 404
 edulis, 75, 103
 felleus, Fig. of, 404
 Borage, 592
 Borage Family, 91, 591
 Boraginaceæ, 591
 Boragineæ, 591, 592
 Borke, 131
 Borrelia, 373
 Boswellia carterii, 548
 Botanical nomenclature, 329
 Botany, agricultural, 3
 cryptogamic, 3
 defined, 1
 departments of inquiry, 2
 derivation of, 1
 economic, 3
 geological, 3
 importance to man, 4
 phanerogamic, 3
 pharmaceutical, 3
 Botrychium, 434
 Botrydium, 338
 Bougainvillea, 514
 Bract, 256
 Bract leaves, 228
 Bracteolar leaf, 227, 256
 Bracteoles, 227, 256
 Bræmer's reagent, 92
 Brake fern, 72, 433, 435
 Branch Traces, 145
 diagrams of, 146
 Branching, deliquescent, 191
 dichotomous, 355
 excurrent, 191
 of roots, 176
 Brand spores, 398
 Brandy, 556
 Brassica, embryo of, 295
 Brassica juncea, 530
 Brassica napus, 530
 Brassica nigra, 103, 529, 530
 Brauneria angustifolia, 612, 615
 pallida, 615
 Brayera, 259, 536

- Brazil nut, 569
 Brazil wood, 539, **540**
 Bread mold, 377
 Brewer's yeast, 382
 Bridal wreath, 535
 Bromelin, 103
 Brooksilk, 343
 Brown Algæ, 95, **351**
 Brown, Robert, 17
 Brownian movement, 365
 Brucinæ Sulfas, 585
 Brucine, 85
Brunfelsia Hopeana, 598
 hydrangæformis, 598
 Bryales, 426
 Bryology, 3
 Bryonia, 608
Bryonia alba, 607, 608
 dioica, 607, 608
 Bryony, 607
 tendrils of, 189
Bryophyllum, 185
 Bryophyta, 13, **418**
 Bryophytes, 13, **418**
 Buckbean, 587
Buchnera, 599
 Buchu, 95, **546**
 subepidermal cells of, 108
 Buckthorn Family, 554
 Buckwheat, 512
 family, 314, **512**
 starch of, 81
 Bud, defined, 184
 Bud sports, 653
 Budding, **45**
 Buds, 184, 185
 Buffalo Berry, 569
 Bulb, defined, 192
 scaly, 192
 tunicated, 192
 Bulbel, 189
 Bulbil, 189
 Bulbs, Figs. of, 191
 Bunchberry, 577
 Bundles, **140**, 713
 bicollateral, **141**, 203
 closed collateral, 140
 collateral, 140
 concentric, 141
 Bundles, fibrovascular, 140
 open collateral, 140
 phlocentric, 141
 vascular, 140
 xylocentric, 141
 Burdock, 612
 Burning Bush, 552
 Bur-reed, 618
Burseraceæ, 547
Butomus, anthers of, 277
 Buttercup Family, 518
 Butterworts, 640
 Buttonbush, 603
 Buttonwoods, buds of, 185

 Cacao Præparatum, 559
Cactaceæ, 567
 Cactus Family, 567
 Cactus Grandiflorus, 568
Cæsalpinia brasiliense, 538, **540**
 echinata, 538, **540**
 Sappan, 540
Cæsalpiniaceæ, 538, **539**
 Cæsalpinus, 324
 Caffèina, **563**, **605**
 Caffèine, **85**, 552
 micro-chemic test for, 85
 Calamus, 147, **487**
 inflorescence of, 259
 Calamus Draco (see *Dæmonorops Draco*)
Calceolaria, 275, **600**
 Calcium, importance of, 156
 Calcium oxalate, 89
 Fig. of various forms of crystals, 88
 how formed, 89
 tests for, 89
 types of crystals, 89, 90
 Calcium pectate, 20
 Calendula, 614
Calendula officinalis, 614
 California Poppy, 524
 California Privet, 170
 Calla, 231, 257
 Calla Lily, 487
Callicarpa, 593
Callitris quadrivalvis, 466
Calluna, 579
 Caltrop Family, 544
 Calumba, 521

- Calyptra of moss, 429
 of root, 158
 Calyptrogen, 113, 156
 Calyx, 262, **266**
 aposepalous, 266
 bilabiate, 266
 caducous, 268
 Fig. of, 267
 campanulate, 267
 chorisepalous, 266
 deciduous, 268
 defined, 266
 epigynous, 269
 Fig. of, 267
 fugacious, 268
 gamosepalous, 266
 hypocrateriform, 267
 hypogynous, 266
 irregular, 267
 perigynous, 269
 Fig. of, 266
 persistent, 268
 petaloid, 266
 polysepalous, 266
 regular, 267
 rotate, 267
 synsepalous, 266
 tubular, 267
 urceolate, 267
 Cambium, 114
 cork, 114
 interfascicular, 169, 172, 196, 201
 intrafascicular, 169, 172, 196, 201
 origin of, 113
 ring, **169, 196**, 197
 Cambogia, 563
Camelliaceæ, 563
 Campanales, 609
Campanulaceæ, 609, **610**
 placentation in, 288
 styles of, 289
 Campanulales, 609
 Camphora, 523
Camptosorus, 435
 Candytuft, 529
 Cane sugar, micro-chemic test for, 75
 Canella, 565
Canella Winterana, 565
Canellaceæ, 565
 Canna Family, 492
 starch, 82
 style of, 289
 Cannabis, 509
Cannabis sativa, **91, 509, 510**
 Cape Aloe, 490
 Capillitium, 375
 Capitulum, 259
 Fig. of, 611
Caprifoliaceæ, 606
Capsella, 529
Capsella bursa-pastoris, 530
 Capsicum, 304, **598**
Capsicum annuum, 598
 frutescens, 598
 Capsule, 301
 histology of a, 307
 Caraway, 258
 Carbon cycle, 622
 Carboxylase, 104
 Carcerulus, 302
 Cardamomi Semen, 492
 Cardamom, 300, 313
Cardiospermum, 554
Carica Papaya, 566
Caricaceæ, 566
 Carina, 271
 Carnation, 273, **514**
 Carnauba wax, 96, **487**
 Carnivorous plants, 621
 Carnoy fluid, 693
 Carotene, 100
 Carotin, 42, 99, **100**
 Carpel, defined, 285
 Carpet Weed, 514
Carpinus, 504
 Carpogonium, 358
 Carpophore, **266, 286, 303**
 Carpospores, 359
 Carragheen, 359
 Carrion fungi, 411
 Carrot, 164
 Carthamus, 615
Carthamus tinctorius, 615
 Carum, 577
Carum Carvi, 577
 Caruncle, 313
 Carya, 504
Caryophyllaceæ, 514

- Caryophyllus, 574
Caryophyllus aromaticus, 573, 574
 Caryopsis, 304
 Cascara Amarga, 539
 Sagrada, 90, **208, 554, 556**
 Cascarilla, 551
 Casparian spots, 128
 Casparian strips, 128
 Cassava, starch of, 81
Cassia angustifolia, 251
 structure of leaflets, 251
 Cassia Bark, 524
 Buds, 524
 Fagot, 524
 Cassia Fistula, 539, **540**
Cassia Senna, 251, 539
 Fig. of, 538
 Cassine, 552
 Castanea, 507
Castanea dentata, 88, **506, 507**
Castilloa elastica, 101
 Castor-oil plant, 550
 Castor-oil seed, 312
 Cataria, 595
 Catechu, 539, **540**
Cathartocarpus fistula, 301, **540**
 Catkin, 259
Catleya, 494
 Catnip, 261, 595
 Cat-tails, 618
 Cauda, 274
 Caulicle, 311
 Caulophyllum, 519
Caulophyllum thalictroides, **519, 522**
 Caustics, **33**
 Ceanothus, 556
Ceanothus americanus, 556
 Cedar, 465
Cedrus, 465
 Celandine, 524
 latex of, 101
Celastraceæ, 552
Celastrus, 552
 Cell, as a fundamental unit, 17
 body, 452, 460
 formation and reproduction, **44**
 inclusions, **40**
 living, 17
 membranes, 18, 19, **43**
 Cell, mother, 292
 non-protoplasmic contents, 74
 plate, 49
 rhizoid, 341
 sap, 18, **40**
 stalk, 452, 460
 tube, 292
 typical colorless, 18
 typical green, 20
 Cell walls, 23, 107
 behavior of to micro-chemic re-
 agents, 109
 cutinized, 109, **123**
 growth in area and thickness, 108
 primary, 23
 secondary, 23, 107
 silicified, 109
 suberized, 109
 tertiary, 24, 107
 various kinds of and behavior of
 each to micro-chemic re-
 agents, 109
 Celloidin sections, 700, 701
 Cells, 10
 asexual, **44**
 column, 321
 companion, 134, 135
 complementary, 204
 egg, 69, 339, 356
 epidermal-daughter, 253
 germ, 50
 guard, 123, 253
 neck canal, 69, 422
 neighboring, 251
 palisade, 321
 palisade parenchyma, 117
 parenchyma, 116
 secretion, 147
 somatic, 50
 sexual, **44**
 sperm, 339, 356
 sperm mother, 422
 sterile, 422
 stoma-mother, 253
 stone, 119
 subsidiary, 251
 typical green, 20
 ventral canal, 69, 422
 wood parenchyma, 117

- Cellulose, 107, 109
 enzyme hydrolyzing, 102
 mucilaginous modification of, 107
 reserve, 109
Centaurea Cyanus, 615
Centaurium, 587
Centrospermæ, 513
 Century Plant, American, 75
Cephalis acuminata, 605
 Ipecacuanha, 603, 606
Cephalanthus, 603
Cephalotus, 629, 635, 638
 follicularis, 635
Cercoarpus, 535
Cereus giganteus, 567
Cerevisiæ Fermentum Compressum, 386
 Ceriman, 487
Cetraria islandica, 414, 417
 Chalaza, defined, 287
Chamælirium luteum, 490
 Chamomiles, 614
Chara, 346
Chara fragilis, life history, 347
 Figs. of, 347, 348
Characeæ, 346
 Charales, 346
 Charophyta, 333
 Chaulmoogra oil, 565
Chelidonium, 302, 524, 528
 majus, 528
Chelone glabra, 600
 Chemotaxy, 33
 Chemotropism, 33
Chenopodiaceæ, 514
 pollen of, 283
 Chenopodiales, 513
Chenopodium, 304, 514, 619
 ambrosioides anthelminticum, 514
 Fig. of flower, 265
 volatile oil of, 514
 Cherry, 305, 535
 Bird, 534
 Fig. of flower, 270, 534
 Fig. of fruit of, 298
 Sour, 534
 Cherry, Wild Black, 534
 Cherry laurel, emulsin in leaves of, 103
 Chestnut, 506
 Chicle, 582
 Chicory, 611, 615
 Fig. of, 614
Chimaphila, 581
Chimaphila, 579
 maculata, 579
 umbellata, 579, 581
 Chinese Galls, 552
Chionanthus, 583
Chionanthus virginica, 583
 Chirata, 587
 Chlamydobacteriaceæ, 349
Chlamydomonas, 338
 Chlamydospore, 398
 Chloral Hydrate, 681
 Chloral Iodine, 681
 Chlorenchyma, 116
 Chlorine, role of, 157
 Chlorophyceæ, 336
 Chlorophyll, 22, 42, 98
 parenchyma, 116
 Chloroplast, 22
 Chloroplastids, 42
 Chlor-zinc-iodine solution, 682
 Choke Berry, 535
 Choke Cherry, enzyme in bark of, 103
Chondodendron tomentosum, 521
 Chondriosomes, 43
 Chondrus, 359
Chondrus crispus, 359, 360
 Choripetalæ, 499
 Christ's Thorn, 535
 Chromatin, 18
 Chromatophore, 42, 337
 Chrome-acetic fluids, 694
 Chrome-acetic-formalin fluid, 693
 Chrome-osmium-acetic fluids, 694
 Chromic acid, fixation, 693
 solution of, 682
 Chromioles, 645
 Chromogens, 364
 Chromomeres, 645
 Chromophyll, 100
 Chromoplastids, 42
 Chromosomes, 47, 48
 behavior of during reduction division, 646
 distribution during maturation and fertilization, 645

- Chromosomes, number of, 50
 structure of, 645
Chrysanthemum cinerariæfolium, 615
Marschallii, 615
Chrysanthemum, papillae on epidermis of,
 124
roseum, 615
 Chrysarobinum, 540
 Cibotium, 435
 Cichorium, 615
Cichorium Intybus, 615
 Cilia, 366
 Ciliary movement, **39**
 Cimicituga, 258, **519**
Cimicifuga racemosa, **519, 520**
Cinchona, 330, **604, 605**
Calisaya, 605
Ledgeriana, 605
succirubra, 605
Cinchona U.S.P., 605
 Cinchonidinæ Sulfas, 605
 Cinchoninæ Sulfas, 605
Cinnamomum, Burmanii, 524
 Camphora, 523
 Cassia, 524
 Loureirii, 524
 Saigonicum, 524
 Zeylanicum, 524
 Fig. of, 525
Cinnamomum U.S.P., 524
 Cinnamon Fern, 433
 Cinquefoil, 534, 535
Cistaceæ, 567
 Citric acid, 391
Citromyces, 391
Citrullus colocynthis, 608
vulgaris, 608
Citrus, 545
Aurantium, **544, 545, 546**
 var. *Bergamia*, 546
 var. *sinensis*, 546
medica var. *Limon*, 546
Cladonia, 414
rangiferina, 417
 Class, 327
 naming of, 332
 Classification, principles of, 323
Clavaria flava, Fig. of, 403
 Clavariales, 403
Claviceps purpurea, Fig. of, 396
 life history of, 395
Claytonia virginica, 514
 Clearing, 695
 Cleavers, 603, **605**
 Clematis, 518
 Clove, 149, 282, **574**
 Clove Bark, 524
 Clover, 538
 Club Fungi, 375
 Club Mosses, 431, **439**
Cnicus benedictus, 615
 Coalescence, 265
 Coca, 241, **544**
 Coca Family, 543
 Cocaina, 544
 Cocaine, 85
 micro-chemic test for, 85
Coccaceæ, 371
Coccoloba uvifera, 512
 Cocculus, 521
 Coccus, 365
Cochlearia Armoracia, 530
 Cocillana, 548
 Cocoa, 85
 Coconut, 486
 Fig. of fruit, cut vertically, 306
 Palm, 485, **486**
 Coconut oil, 487
Cocos nucifera, 487
 Codeine, **87**, 527
 Cœnocyte, 325
 Cœnocytic latex cells, 133
Coffea arabica, 605
liberica, 605
Coffea Tosta, 605
 Coffee, 85, 605
 Cohesion, 265
Coix Lacryma-Jobi, 485
 Cola Family, 557
Cola nitida, 559
 Colchici Cormus, 490
 Colchici Semen, 490
 Colchicine, micro-chemic test for, 87
 Colchicum, 192, 300
 seed, 313
Colchicum autumnale, 87, **490**
 Coleoptile, 318
 Coleorhiza, 318

- Collenchyma, 63, **117**
 Collinsonia, 596
Collinsonia canadense, 596
 Colloids, **30**
Colocasia esculenta, 487
 Colocynth, 607
 Colocynthis, 608
 Coltsfoot, 612
 Columbine, 266, 518
 Columella, 378
 Coma, 313
Combretaceæ, 574
 Combretum, 574
Combretum sundaicum, 574
 Comfrey, **592, 593**
Commiphora abyssinica, 548
 myrrha, 548
 Companion cells, **134**
Compositæ, 609
 Compressed yeast, 386
Comptonia, 501, **502**
Comptonia asplenifolia, 123, **502**
 photomicrograph of long. -section
 through staminate catkin of,
 259
 tubercles on, 179
 Conceptacles of *Fucus*, 355
 Figs. of, 356, 358, 359
 Condurango, 589
 Cone, defined, 307
 Figs. of, 447, 448, 449
 Confervales, 341
 Conidia, 390
 Conidiophore, 390
 Conidiospores, 390
Coniferae, 84, 253, 465
 Coniferales, 455, **464**
 Coniferin, micro-chemic test for, 84
 Conifers, 443, **444**
 Wood of, 139
 Conium, 163, **577**
Conium maculatum, **577, 578**
 Conjugales, 342
 Conjugation, **46, 342**
 lateral, 346
 scalariform, 345
 Connation, 265
 Connective, **274**
 Convallaria, 258
Convallaria majalis, 490, 491
 histology of rhizome, 215
 Convallariæ Radix, 490
 Flores, 491
Convolvulaceæ, 590
Convolvulus, 591
Convolvulus Scammonia, **590, 591**
 Copaiba, 540
Copernicia cerifera, 96, **487**
 Copper acetate test for tannins, 92
 test for resins, 98
 Coptis, 519
Coptis occidentalis, 519
Coptis trifolia, 519
 Coral fungi, 403
 Corallin-soda solution, 682
Corallorhiza odontorhiza, 496
 Coriandrum, 577
Coriandrum sativum, 577
 Cork, **129, 207**
 commercial, 506
 microchemic tests for, 109
 Cork cambium, 114, 129
 origin in roots, 169
 origin in stems, 200, 202
 Cork formation in roots, 169
 in stems, 200, 202
 Corm, defined, 192
 Fig. of, 191
 Corn, 484
 fruit, 304
 histology of seed, 315
 histology of stem, 215, **216**
 inflorescences, 483
 leaves, 124
 root, Fig. of cross section, 128
 seed, 315
 seedling, 316, 319
 smut, 397
 Fig. of, 398
 starch, 79, 80
 Corn Cockle, 514
Cornaceæ, 577
 Cornua, defined, 274
 Cornus, 579
Cornus canadensis, 577
 florida, 577, 579
 sanguinea, 577

- Corolla, 262
 apopetalous, 270
 bilabiate, 271
 campanulate, 272
 caryophyllaceous, 273
 choripetalous, 270
 crateriform, 272
 cruciform, 270
 defined, 269
 forms of, 273
 gamopetalous, 270
 hypocrateriform, 272
 infundibuliform, 272
 labiate, 271
 ligulate, 271
 papilionaceous, 270
 personate, 272
 polypetalous, 270
 ringent, 272
 rotate, 272
 synpetalous, 270
 tubular, 272
 urceolate, 272
 Corona, 274
 Cortex, 206, 215
 primary, 167, 195
 secondary, 169, 202
Corticium, 403
Corydalis, 528
Corylus, 504
 Corymb, 258
Corynanthe Johimbi, 605
Corynebacterium, **372**
 diphtheriæ, 373
Cosmarium, Fig. of, 338
 Coto, 524
 Cotoneaster, 535
 Cotton, 126, **561**
 Cottonwood, Fig. of twig of, 185
 Cotyledons, **227, 311, 312**
 function of, 311
 Couch grass, structure of stem of, 219
 Cowhage, 540
 Craef fluid, 693
 Cranberries, 581, 619
 Cranesbill, 542
Crassulaceæ, 221
Crategus, 535
 Cremocarp, 303
Crenothrix, 373
 Fig. of, 371
 Cribiform tissue, 133
Crocus, 191, 192, 491, 492
Crocus sativus, 492
Cronartium ribicola, 401
Croton Eluteria, 551
Croton Tiglium, 551
 Crowfoot Family, 243, **515**
 Crown, 274
Cruciferae, 103, 268, **528, 530**
 Crude sap, 117
 Cryptogamia, 325
 Cryptogams, defined, 13
 vascular, 431
 Crystal fibers, 90
 Crystal sand, 91
 Crystalloids, 94
 Crystals, columnar, 89
 membrane, 90
 micro-, 90, 91
 raphides, 89
 rosette aggregate, 89
 solitary, 89
 Cube Root, 540
 Cubeba, 499
 Cucumber, 607
Cucurbita Pepo, 608
Cucurbitaceæ, 607
 Cucurbitales, 607
 Cudbear, 416
 Culm, 188
 Culver's Root, 600
 Cup Fungi, 387
Cuphea, 570
 Cupule, **297, 304, 507**
 Cupules, of liverwort, 420
Cupuliferae, 506
 Cuprea Bark, 605
 Curare, 585
 Curcuma, 492
Curcuma longa, 492
 Zedoaria, 492
 Currant, 189, 304
 Cusparia Bark, 546
 Custard Apple Family, 515
 Cutin, 24, 107, 128
 Cyanogenetic glucosides, 84
 Cyanophyceæ, 334

- Cyatheaceae*, 434
 Cycadales, 455
 Cycadofilices, 455
 Cycads, 253, 455
Cycas revoluta,
 Fig. of, 456
Cydonia vulgaris, 536
 Fig. of fruit, 267
 Cydonium, 536
Cylindrospermum, 336
Cymbopogon Nardus, 485
Cymbopogon Winterianus, 485
 Cyme, 260
 Dichasial, 260
 Scorpioid, 260
 Simple, 260
Cynoglossum officinale, 593
Cyperaceae, 483
Cypripediæ, 495
 Cypripedium, 496
Cypripedium bulbosum, 496
 parviflorum, 496
 pubescens, 496
 Cyst, 28
 Cystocarps, 359
 Cystolith, 90, 91, 602
 hair, 91
 Cytase, 102
Cytisus laburnum, Fig. of cross sect.
 of branch, 207
 Scoparius, 540
 Cytology, defined, 2
 Cytoplasm, 18, 40
 Cytoplasmic caps, 47
 Cytoplasmic rotation, 39
 Czapeck's medium, 376

Dacromyces deliquescens, 402
Dacromycetales, 402
Dæmonorops Draco, 487
 Daffodil, 192, 257
 Dahlia, 162, 611
 Daisy, 611
 Daisy Family, 609, 610
 Damiana, 566
 Dammar, 466
 Dandelion, 189, 611
Daphne Gnidium, 570
 Laureola, 570
 Daphne Mezereum, 569, 570
Darlingtonia, 631
 Californica, 631, 633
 Darwin, C., 624, 627, 654, 655
 Darwin's Theory, 654
 Date palm, 486
Datura Stramonium, 597
 fruit of, 297
 De Candolle, A. P., 325
 De Candolle, Alphonse, 326
 De Jussieu, A. L., 325
 De Lamarck, J. B., 653, 654
 De Tournefort, J. P., 325
 Deadly agaric, 408
 Deadly Nightshade, 598
 Decay, defined, 364
Decodon, 570
 Dehiscence, apical in fruits, 299
 apical porous in anthers, 279
 circumscissile, 299
 in anthers, 279
 in fruits, 298
 loculicidal, 300
 longitudinal anther, 279
 septicidal, 299, 300
 septifragal, 300
 transverse anther, 279
 valvular-fruit, 299
 Dehydrating, 695
 Delafield's hæmatoxylin, 701
 Delphinium, 519
Delphinium Ajacis, 519
 Staphisagria, 519
 Denitrification, 367
 Dermatogen, 113
 Derris, 540
Derris elliptica, 540
Derris malaccensis, 540
 Desilicification of woody materials, 701
 Desmid, 342
Desmidaceæ, 342
Desmodium, 301
 Determinate inflorescences, 256
 Deuteromycetes, 376
 Deutzia, trichomes on leaves of, 125
 Devil's Aprons, 354
 Dewberry, 300
 Dextrin, 82
 corn, 80, 82

- Dextrin, white, 82
 yellow, 82
 Dextrose, **74, 485**
 Diageotropic, 36
Dianthera, 603
 Diastase, **102, 485**
Diatoma vulgare, 350
 Diatomaceous Earth, 350
 Diatoms, 91, **348**
 Fossil, 352
Dicentra canadensis, 528
 Cucullaria, 528
 Dicotyl Plant, morphology of type, 497
 Dicotyl Root, histology of, 167
 of transitional and secondary growth, 169
 Dicotyl Roots, abnormal structure of, 175
 Dicotyl Seed, 319
 histology of, 320, **321**
 Dicotyl Stems, exceptional types of, 203
 structure of, 194-204
 Dicotyledoneæ, **479, 497**
 Dicotyledons, **16, 479, 497**
 development of embryo in, 295
 drug yielding, 499
 leaves of, 232, 255
 petiole of, 243
 roots of, 167
 stems of, 194, 199
 Dictyostele, **145, 199**
 Dictyotales, 353
Dicypellium caryophyllatum, 524
Diervilla, 606
 Differentiation, **16**
 Diffusion, 153
 Digestion, defined, **26, 67**
 Digitalis, 163, 256, 275, 599, **600**
 Digitalis Pulverata, 600
Digitalis purpurea, 84, 600
 trichomes of, 124, **125**
 var. *gloxinæflora*, Fig. of, 601
 Digitoxin, micro-chemic test for, 84
 Dihybrids, 649
 Dioecious, 264, 418
Dionæa, 35, 530
 Submerged, 625
 • *Dionæa muscipula*, 35, **36, 625**
Dioscorea villosa, 491
 Dioscoreaceæ, 491
 Dioscorides, **2, 330**
Diosma, 545
Diplococcus, **367, 368**
Dipsacæ, 608
 Disaccharoses, 74, **75**
 Dissepiments, 285
 Division, by cleavage furrows, 378
 simple, defined, **28**
 Division of labor, **16**
 Docks, 283, **512**
 Dodder, 161, 188, 228
 Figs. of, 162
 Dog Bane Family, 587
 Dog's Tooth Violet, life history of, 471
 Dogwood, 256, 261
 Blood Twig, 577
 Fig. of flowering branch of, 261
 Flowering, 577
 Dogwood Family, 577
Dorema Ammoniacum, 577
 Douglas Fir, 214, **465**
Dracæna, stem of, 220
Dracæna Draco, 489
 Dragon's Blood, 487, 489
Draparnaldia, 338
Drimys Winteri, 516
 Drosera, 530
Drosera, 36, **622**
 anglica, 530
 longifolia, 530
 rotundifolia, **530, 623**
Droseraceæ, 530
 Drupe, 305
 Fig. of, 306
Dryopteris, 435
Dryopteris Filix-mas, **53, 435**
 alternation of generations in, 71
 comparative physiology of root, stem and leaf, 63
 diagram of life cycle, 72
 Fig. of, 54
 gross structure of frond, 55
 sori and sporangia, 66
 stem, 53
 histology of growing apex, 58
 mature root, 58
 pinnule, 61, 62
 root apex, 60

- Dryopteris Filix-mas*, histology of growing apex, sori and sporangia, 66
 stem, 55
 stipe, 60
 Figs. of trans.-sect., 61
 history of gametophyte generation, 67
 sporophyte generation, 53
 life history of, 53
 origin of new sporophyte or diploid plant from fertilized egg, 71
Dryopteris marginalis, **432, 435**
 Duboisia, 598
Duboisia myoporoides, 598
 Dujardin, 17
 Dulcamara, 598
 Duramen, 213
- Earth Stars, 410
 Ebenales, 581
Eberthella typhi, 373
Ecballium Elaterium, 608
 Echinacea, 615
Echinocactus, 567
Echinocereus, 567
Echium, 592
 Ecology, 3, **616**
 Economic Botany, defined, 3
Ectocarpus siliculosus, 352
 Ectoplasm, 25, **43**, 374
 Egg Plant, 304, **597**
 Eichler, A. W., 326
Elæagnaceæ, 569
 Elaioplastids, **42**, 133
Elæis guineensis, 487
 Elastica, 551
 Elaterium, 608
 Elaters, 422, 437
 Elder, 606
Elettaria Cardamomum, 492
 Elm, 303
 Slippery, 506
 Elm Family, 507
Elodea, 20
 Embryo, 8, 295, **311**, 452
 Figs. showing development of, 294, 295
 parts of, 311
- Embryology, defined, 2
 Embryo-sac, 287, 449, 474
 maturation of, 292, 474
 Emetinæ Hydrochloridum, 605
 Emulsin, 103
 Endocarp, 298
 Endodermis, 127
 Endoperidium, 410
 Endoplasm, 25, **43**, 374
 Endosmosis, 154
 Endosperm, 293, 314
 Endospore, 282, 292
 Endospores, 369, 378, 381
 Endosporium, **68**, 475
 Endothecium, 278, 280
 Endotoxins, 370
 Engler and Prantl, 326
 English ivy, 575
 roots of, 161
Entada scandens, 301
 Enzyme, defined, 26, **101**
 Enzymes, 101
 classification of, 102
 proteolytic, 94, **103**
 Eosin, 682
Ephedra, 467, 468, 469, **470**
Ephedra distachya, 471
Ephedra equisetina, 471
Ephedra Gerardiana, 471
Ephedra sinica, 468, 470, 471
 Ephedrine, 471
 Epicalyx, **269**, 535
 Epicarp, 298
 Epicotyl, 311
 Epidermal outgrowths, 124
 hairs, 124
 papillæ, 124
 scales, 124
 Epidermis, 122
Epigæa repens, 581
Epilobium, 569
 Epiphytes, 161
Epipremnum pinnatum, 487
 Epithem, 124
 Equisetineæ, 91, 431, **437**
Equisetum, 437
 arvense, 438
 life history of, 43
 Ergonovine, 397

- Ergot, 395, 397
 Ergot fungus, 395
Erica, 580
Ericaceæ, 279, **579**
 Ericales, 579
Erigeron canadensis, 615
 Eriodictyon, 591
Eriodictyon californicum, **591, 592**
 trichomes of, 125
Eriogonum, 512
Erythræa Centaurium, 587
Erythronium Americanum, life history of,
 471
Erythroxylaceæ, 543
Erythroxylon, 543
 papillæ on epidermis, 132
Erythroxylon Coca, 544
 Truxillense, 544
Eschscholtzia, 524
 Essential elements, 155
 Etærio, 306
 Ether, use of, 682
 Ethylhydrocupreine, 605
 Etiolin, 100
Eubatus, 536
 Eucalypti Gummi, 574
 Eucalyptol, 574
 Eucalyptus, 574
Eucalyptus, 186, 573
 globulus, 572
 lamina, 250
 leaves, internal glands of, 148
 photomicrograph of trans. sect. of
 leaf-blade, 252
 rostrata, 95
Eucarya spicata, 510
Eudorina, 338
 Eugenol, 574
 Eumycetes, 375
 Euonymus, 553
 Eupatorium, 614
Eupatorium perfoliatum, 232, **614**
Eupatorium purpureum, 615
Euphorbia, 133, 188, 550, 551
 pilulifera, 551
 resinifera, 551
Euphorbiaceæ, 550
 Euphorbium, 101, **551**
 Euphrasia, 599, **600**
Euphrasia officinalis, 600
Eurotium, 391
 Evening Primrose, 282, 569, 657
 Everlasting, 615
 Evolution, 653
Evonymus, 313, **552**
Evonymus Americanus, 553
 atropurpureus, 553
 Europæus, 552
 Eye-bright, 600
 Excretion, defined, 27
 Exine, 281, 292
Exoascus, 381
 Exobasidiales, 403
Exogonium Jalapa, 591
 Exoperidium, 410
 Exosmosis, 154
 Exospore, 281
 Exosporium, **68, 439, 475**
 Exothecium, 278
 Exotoxins, 370

Fabaceæ, 536
Fagaceæ, 506
Fagales, 504
Fagopyrum, 512
 Fagot Cassia, 524
Fagus, 330
Fagus grandifolia, 506
 Families, naming of, 332
 Family, defined, 327
 naming of, 332
 Farfara, 615
 Fast Green, 682
 Fats, 95
Fatsia horrida, 575
 Fehling's solution, 682
 Fennel, 258, 576
 Fenugreek, 540
 Ferment, 101
 Fermentation, **364**
 Fern, antheridium, 68, 71
 apical cell, 58
 apical cone, 58
 archegonium, 69, **69**
 Beech, 435
 Boston, 653
 Brake, 433, 435
 Christmas, 435

- Fern, Cinnamon, 433
 foot of, 71, 433
 frond of, 55
 Interrupted, 436
 leaves of, 53, 434
 Male, 53
 prothallium, 68, 70
 Royal, 436
 Sensitive, 433
 sorus, 67
 spermatozoids of, 68, 70
 Walking-leaf, 435
 Wood, 435
 Fern Palms, 443, 455
 Ferns, 431
 True, 434
 Water, 436
 Ferric chloride, solution of, 683
 Fertilization, cross, 291
 defined, 46
 double, 293
 in angiosperms, 292
 in White Pine, 452
 in *Zamia*, 461
 self, 291
Ferula Assafœtida, 577
 fœtida, 577
 galbaniflua, 577
 Sumbul, 576
 Fibers, 120
 Fibro-vascular bundles, 140
 bicollateral, 141
 closed collateral, 140
 collateral, 140
 concentric, 141
 open collateral, 140
 phlocentric, 141
 primary, 196
 radial, 142
 secondary, 198
 xylocentric, 141
 Ficus, 510
Ficus, 248, 507
 Carica, 510
 elastica, 101
 petiole of, 245
 religiosa, 161
 Fig, 507, 509
 Figwort Family, 599
 Filament, 274
 gross structure of, 277
 histology of, 277
 Filbert, 297
 Filicales, 434
 Filicineæ, 431
 Fir, 465
 Fission, 45, 367
 of *Amœba*, 28
Fistulina, 404
 Fixation, 692
 of atmospheric nitrogen, 178
 Fixed oils, 95
 Fixing agents, 692
 Flacourtia Family, 565
Flacourtiaceæ, 565
 Flagella, 339, 366
 Flagellata, 333
 Flagellates, 339
 Flax, 542
 bast fibers of, 118
 New Zealand, 249
 Flax Family, 542
 Flemming fluids, 694
 Flora, defined, 2
 Flores Cyani, 615
 Florideæ, 357
 Flower, accessory organs of, 262
 actinomorphic, 262
 complete, 262
 defined, 262
 diagram of, 476
 diandrous, 274
 dichlamydeous, 266
 diclinous, 264
 double, 263
 epigynous, 264
 essential organs of, 262
 hermaphrodite, 262
 hexandrous, 274
 hypogynous, 264
 imperfect, 262
 irregular, 262
 liliaceous, 271
 monandrous, 274
 monochlamydeous, 266
 monoclinous, 264
 neutral, 263
 orchidaceous, 271

- Flower, pentandrous, 274
 perfect, 262
 perigynous, 264
 pistillate, 263
 polyandrous, 274
 protandrous, 291
 protogynous, 291
 regular, 262
 staminate, 263
 symmetrical, 263
 tetrandrous, 274
 triandrous, 274
 typical, 262
 zygomorphic, 263
 Fly agaric, 407
 Fly-traps, 530, **625**
 Fœniculum, 576
 histology of mericarp of, 309
Fœniculum vulgare, 576
 Folia Laurocerasi, 536
 Foliage leaves, 228
 Follicle, (See pod.), 301
 Fomes, 404
 Foot, **71**, 428
 Foramen, 287
 Forestry, defined, 3
 Forget-me-not, 592
 Formalin-alcohol, 694
Forsythia, 583
Fothergilla, 532
 Four-o'clock Family, 514
 Fovilla, 475
 Foxglove, 599, 601
Fragaria, 269, **535**
Fragaria chiloensis, Fig. of flower, 268
 Frangula, 556
 Frangulin, 103
 Fraxinus, 583
Fraxinus Americana, 583
 Ornus, 583
 Free cell formation, **45**
 Frond, **55**, 431
 Fronds, **53**, 431, 432
 fertile, 434
 sterile, 434
 Fructose, 74
 Fructus Cannabis, 510
 Fruit, 8, **297**
 definition, 297
 Fruit, dehiscence, 298
 indehiscent, 298
 structure, 297
 sutures, 298
 valves, 298
 Fruits, achenial, 300
 aggregate, 300, **305**
 Fig. of, 299
 baccate, 300, **304**
 capsular, 300, **301**
 classification of, 300
 compound, 300, **301**
 dehiscent, 298, 300
 drupaceous, 300, **301**
 indehiscent, 298, 300
 multiple, **300**, 306
 schizocarpic, 300
 simple, 300
 Fucales, 352, **354**
 Fuchs, L., 324
 Fuchsia, 569
 Fuchsin, acid, 683
 basic, 683
 Fucoxanthin, **100**, 351
 Fucus, drug, 357
 Fucus, 355
 Fucus vesiculosus, 355
 Fumariaceæ, 528
 Fumitory Family, 528
 Funaria, 426
 Fig. of, 429
 Fundamental tissue, 116
 Fungi, 12, **362**
 alga-like, 376
 basidia, 397
 black, 395
 carrion, 411
 classes of, 362
 coral, 403
 cup, 387
 fairy club, 403
 fission, 363
 gill, 405
 nest, 410
 poisonous, 407
 pore, 404
 sac, 381
 saddle, 394
 stink horn, 411

- Fungi, tooth, 404
 true, 362, **375**
 weeping, 402
 Fungi Imperfecti, 412
 Funiculus, 287
Funtumia africana, 101
 elastica, 101

Gaillonella, 352
 Galanga, 492
 Galbanum, 577
 Galbulus, defined, 307
 Galea, 268
 Galega, 540
Galega officinalis, 540
 Galen, 2
Galipea officinalis, 546
Galium, **603**, 604
Galium aperine, 605
 Galla, 92, **507**
 Galls, Chinese, 552
 Japanese, 552
 Galton's Law, 652
 Galvanotropism, 36
 Gambir, 605
 Gamboge, 563
 Gamboge family, 563
 Gametangia, 353
 Gametangium, 344, 379
 Gamete, **46**, 344, 353, 379
 Gametophyte, 68, 418, 420, 458
 female, 292, 294, 450, **451**
 male, 292, 449, **450**, 458
 Gamopetalæ, 579
Garcinia Hanburyi, 563
 Garlic, 189, 277
 Fig. of bulb, 191
 Gasteromycetes, 410
 Gaultherase, 103
 Gaultheria, 581
Gaultheria, enzyme in, 103
Gaultheria procumbens, **579**, **581**
 Gaultherin, 103
Gaylussacia, 580, 581
Geaster, 410
Gelidiaceæ, 361
Gelidium, 361
Gelsemium, 585
 Gelsemium, 583
 sempervirens, **584**, **585**
 stem, 204
 Gemmæ, 421
 Gemmation, **45**, 385
 Genera Plantarum, 326
 Generative tissues, 113
 Generic names, 329
 Gene, modern theory of, 658
 Genes, 18, **645**
 Genetics, 3, **643**
 Genotype, 647
 Gentian violet, solutions of, 683
 Gentian Family, 585
 Gentiana, 587
Gentiana acaulis, 586
 lutea, 585, **586**, 587
 verna, 586
Gentianaceæ, 585
 Gentianales, 583
 Generic name, 329
 Genus, defined, 327
 Geological Botany, defined, 3
 Geotropism, 35
 dia-, 36
 lateral, 36
Geraniaceæ, 540
 Geraniales, 540
 Geranium, 542
Geranium, 541
 fruit of, 301
 maculatum, 542
 petiolar structure, 245
 Geranium Family, 540
Gerardia, 161, **599**
 Germination, seed, 318, 453
Geum, 266, 269, **535**
Gigartina mamilliosa, 359, 360
 Gills of mushrooms, 405
 Ginger, 147
 Family, 492
 Fig. of plant, 493
Ginkgo biloba, 463
 Ginkgoales, 455, **463**
 Ginseng, 575, **576**
 American, 576
 Chinese, 576
 Ginseng Family, 575
 Gladiolus, 491

- Glands, external, 151
 internal, 148
 Glans, defined, 304
 Gleba, 410
 Gliadins, 93
 Globoid, 94
 Globulins, 93
Glæocapsa, 335
 Fig. of, 334
 Glomerule, **261**, 577
 Glory-bower, 593
 Gluco-alkaloids, 87
 Glucoside enzymes, 103
 Glucosides, characteristics of, 83
 Glucosum, 485
 Glumes, 484
 Glutelins, 93
 Glycerine, dilute, 683
 gum, 683
 Glycerin-Gelatin, 691
 Glycine hispida, 482
 Glycosides, 83
Glycyrrhiza, 275, **539**
 glabra, **537**, **539**
 glandulifera, 539
 typica, 539
Gnaphalium polycephalum, 615
Gnetaceæ, 469
 Gnetales, 469
Gnetum, 469, 470
 Golden Bell, 583
 Golden Seal, **518**, 620
 Gonidia, 414
 Gonophore, 266
 Gooseberry, 305
 Goosefoot, Fig. of flower, 265
 Goosefoots, 619
 Goosefoot Family, 514
Gossypii Radicis Cortex, 561
Gossypium, 561
 Barbadense, 561
 herbaceum, 560, **561**, **562**
Gossypium hirsutum, 560, **561**, **562**
Gossypium Purificatum, 561
 Gourd, defined, 305
 Gourd Family, 607
 Grain, defined, 304
 Grains of Paradise, 492
 Graminales, 483
Gramineæ, 91, 304, **484**
 pollen of, 267
 silica in, 91
 Gram's method, 351
Granataceæ, 570
Granati Fructus Cortex, 571
Granatum, 92, **571**
 Grape, 188, 261, 304, **557**
 Grape Family, 556
 Grape Fruit, 305, 545
 Grape Sugar, 57
 Grape Vine, 202, 556
 tendrils of, 189
Graphis, 414
 Grass, diagram of flower, 562
 petiole of, 242
 Grass Family, 259, 304, **484**
 Greek Tea, 596
 Green Algæ, 336
 Felt, 346
 Mold, 388
 Greenbrier, histology of stem, 217
 Fig. of T. S. stem, 218
 Grindelia, 615
Grindelia camporum, 615
 cuneifolia, 615
 squarrosa, **612**, **615**
 Gross anatomy, defined, 2
 Ground Cherry, 297
 Ground Ivy, 594
 Ground meristem, 113
 Growth, 26
 of perennial dicotyl stem, 199
 Gruinales, 540
 Guaiac, 544
Guaiaci Lignum, 545
Guaiacum, 544
Guajacum officinale, **544**, **545**
 sanctum, **543**, **544**, **545**
 Guarana, 554
 Guard cells, 253
Guarea Rusbyi, 548
 Gulf Weed, 355
 Gum, 95
 Cherry, 95
 Kauri, 466
 Red, 95
 Spruce, 466

- Gum resins, 97
 defined, 98
 Gums, 94, 95
 Gutta Percha, 101, **582**
Guttiferae, 563
 Gymnosperm, life history of a, 444
Gymnospermæ, 443
 Gymnosperms, 15, **443**
 classification of, 455
 differences from angiosperms, 438
 Gynæcium, 262, 283
 Figs. of forms of, 286
 Gynophore, 266
 Fig. of, 286

 Haberlandt, 123
 Habitat, defined, 3
 Hackberry, 507
 Hadrome, 142
 Hæmatoxylin stain, 683
 Hæmatoxylon, 539, **540**
Hæmatoxylon campechianum, 540
Hagenia abyssinica, 536
 Hair cystoliths, 91
 Hairs, aggregate, 125
 barbed, 124
 branched, 125, **126**
 candelabra-shaped, 124
 clavate, 124
 climbing, 126
 epidermal, 124
 functions of, 127
 glandular, 126
 hooked, 124, 126
 multicellular, 124
 multiseriate, 126
 non-glandular, **124**, 126
 root, 127, **152**, 154
 stellate, 124, 125
 stinging, 126
 tufted, 124
 unicellular, 125, 126
 uniseriate, 125, **126**
Halidrys siliquosa, 357
 Halophytes, 164, 617, **618**
Haloxylon, 514
Hamamelidaceæ, 531
 Hamamelidis Cortex, 532
 Hamamelidis Folium, 532

Hamamelis virginiana, 532
 Fig. of, 533
Hancornia, latex in, 101
 Hard Bast, 202
 Hardening, of soft materials, 695
 Harebell, 272, 277
 Haustoria, 161
 of fungi, 362
 Hawkweeds, 614
 Hawthorn, 258, 535
 Hay Fever, pollens responsible for, 283
 Hazelnuts, 504
 Head, defined, 259
 Heart-wood, 213
 Heath, 273
 Heath Family, 579
 Heather, 579
 Hedeoma, 595
Hedeoma pulegioides, 595
Hedera Helix, 575
Heliamphora, 629
 nutans, 629
 Helianthemum, 567
Helianthemum canadense, 567
Heliconia, 253
Heliotropeæ, 591, 592
 Heliotropism, **34**
 Helleborus, 519
Helleborus fœtidus, 519
Helleborus niger, 519
 Helonias, 490
 Helophytes, 617, **618**
 Helvellales, 394
 Hemlock, 465
Hemophilus, 373
 Hemp, 507, 508
 seed, 510
 Henbane, 597
 Fig. of calyx, 266
 Henna, 570
 Hepaticæ, 13, **418**
 Hepburn, 101, 627
 Herb, defined, 190
 evergreen, 190
 Herba Majoranæ, 596
 Hercules Club, 575
 Heredity, 643
 Hesperidin, 83
 Hesperidium, 305

- Heteroauxin, 104
 Heterocysts, 336
Heuchera, 531
Hevea, 550
 latex of, 101
Hevea braziliensis, 551
Hibiscus, flower of, 267
Hibiscus abelmoschus, 562
Hibiscus Syriacus, 560
 Hickories, 504
 stems of, 202
 Hilum,
 of ovule, 287
 of seed, 321
 of starch grain, 77
 Hip, defined, 304
 Histological technique, 675
 Histology, defined, 2
 of dicotyl roots, 167
 of Aconitum, 175
 of annual dicotyl stem, 194
 of anther, 277
 of Aristolochia Siphon stem, 199
 of bark, 208
 of California Privet root, 170
 of Cascara Sagrada, 208
 of Convallaria rhizome, 215, 217
 Corn stem, 216
 of dicotyl roots, of primary growth, 167
 of secondary growth, 169
 of dicotyl stems, 194
 of dicotyl tuberous root, 175
 of *Ephedra sinica* stem, 469, 470
 of fern rhizome apex, 58
 pinnule, 61
 root, 58
 sori and sporangia, 67
 stem (rhizome), 55
 stipe, 60
 of filament, 277
 of fruits, 307
 of grass stems, 219
 of Greenbrier stem, 217
 of herbaceous monocotyl stem, 215
 of leaves, 247
 of Lilac leaf blade, 247
 of mericarp, 309
 of monocotyl rhizome, 215
 Histology, of monocotyl roots, 165
 of monocotyl stems, herbaceous, 215
 woody, 217
 of perennial dicotyl stem, 199, 202
 of petioles, 242, 244
 of seeds, 315
 of sunflower stem, 194
 of typical monocotyl rhizome, 215
 of Vanilla capsule, 307
 Holdfast, 355
 Holly, 552
 Hollyhock, 401
 Honey dew, 395
 Honey Locust, 189
 Honeysuckle, 272, 291, 305, **606**
 Honeysuckle Family, 606
 Hooke, Robert, 17
 Hooker, J. D., 326
 Hop, 127, 188, **259**, 291, 507
 Fig. of, 508
Hordeum vulgare, 485
 Horehound, 261, **595**
 Hormogonia, 336
 Hormones, 104
 Hornbeam, 504
 Horse chestnut, 187, 256
 Horsemint, 594, **596**
 Horsenettle, 596
 Horsetails, 431, 437
 Horticulture, defined, 3
 Hound's Tongue, 593
 Houseleek, 96, 189
 Huckelberries, 581
Hudsonia, 567
 Humulus, 510
 glandular trichomes of, 125, 127
Humulus Lupulus, 510
 Hutchinson, J., 326
 Huxley, **30, 31**
 Hyacinth, 489
 Hybrid, defined, 329
 Hybrids, "Splitting of," 645
 Hydathode, 124
Hydnaceae, 404
Hydnocarpus anthelmintica, 565
Hydnocarpus Wightiana, 565
Hydnum, 404

- Hydrangea*, 264, 531
 arborescens, 531
Hydrastis, 260, **518**
Hydrastis canadensis, 518
 Fig. of, 517
 Fig. of colony of, 620
 Hydrochloric acid, 684
Hydrodictyon, 338
Hydrophyllaceæ, 591
 Hydrophytes, 164, **617**
 Hydropterales, 436
 Hydrotropism, **34**
 Hymenium, 402
 Fig. of, 406
 Hymenomycetes, 402
Hyoscyamus U.S.P., 598
 fruit of, 301
Hyoscyamus muticus, **125**, 126
 niger, 598
 Fig. of calyx, 266
Hypericaceæ, 564
Hypericum, 276, **565**
 perforatum, **564**, **565**
 Hypha, defined, 362
 Hyphæ, aerial, 377, 389
 rhizoidal, 377
 stoloniferous, 377
 submerged, 377, 389
 Hyphomycetes, 333, **362**
 Hypocotyl, 311
Hypocreaceæ, 395
 Hyssopus, 596
Hyssopus officinalis, 596

 Iceland Moss, 414
 Ignatia, 585
Ilex opaca, 552
 paraguariensis, 552
 verticillata, 552
 vomitaria, 85
Illicaceæ, 552
Illicium verum, 516
 Imbedding, celloidin, 699
 of woody plant organs, 696
 paraffine, 695, 699
Impatiens, 330
 Imperatoria, 477
Imperatoria Ostruthium, 477
 Inclusions, 74
 Indehiscent fruit, 298, 300
 Indeterminate inflorescence, 256
 Indian Corn, 315, 397
 Figs. of grain and seedling, 316, 317
 histology of seed, 315
 Indian Cress, 267
 Indian Hemp, 509
 Indian Pipe, 580
 Indian Turnip, 487
 Indigenous, defined, 3
 Indigo, 540
Indigofera tinctoria, 540
 Indirect nuclear division, 47
 Indusium, **67**, 433
 Inflorescence, ascending, 256
 centrifugal, 256
 centripetal, 256, 257
 cymose, 256
 defined, 256
 descending, 256
 determinate, 256, **259**
 indeterminate, 256, 257
 mixed, 256
 racemose, 256
 solitary determinate, 260
 solitary indeterminate, 258
 Ingestion, **26**
 Inheritance, 643, 652
 Integuments of angiospermous ovule,
 287
 Intercellular-air-spaces, lysigenous, 147
 schizogenous, 147
 Internal glands, 148
 lysigenous, 149
 schizogenous, 149
 Internal phloem, 204
 Internal secretion reservoirs, 148
 Internode, defined, 186
 Intine, 282, 292
 Intussusception, **27**, 31, 108
 Inula, 615
Inula Helenium, 615
 Inulase, 103
 Inulin, 82
 Invertase, **75**, 102
 Involucre, 256
 Involucres, persistent, 297
 Iodine, a source of, 354

- Iodine and potassium iodide, solution of, 684
 Iodine water, 684
 Ipecac, 603
 Ipecacuanha, 605
 Ipomœa, 591
Ipomœa orizabensis, 591
 pandurata, 591
 simulans, 591
 Iridaceæ, 491
 Iris, 300, 491
 caroliniana, 492
 Family, 492
 florentina, 492
 germanica, 492
 pallida, 492
 petiole of, 249
 style of, 289
 versicolor, 492
 Irish Dulse, 361
 Moss, 359
 Iron, as an essential element, 157
 Ironwood, 504
 Irritability, **31**
 Isoetaceæ, 442
Isoetes, 442
 Ivy, 188
 English, 575

 Jacket, 451
 Jack-in-the-pulpit, 192
 Jalap, 590
 abnormal root structure, 175
 Tampico, 591
 Wild, 591
 Jalapa, 591
 Jamestown weed, 297
 Japan ivy, 189
 Japan wax, 96
 Japanese Galls, 552
Jateorhiza palmata, 521
 Fig. of, 523
 Jequirity, 540
 Jerusalem Artichoke, 192, 611
 Jew's ear fungus, 402
 Job's Tears, 485
 Juglandaceæ, 502
 pollen of, 283
 Juglandales, 502
 Juglans, 504
Juglans cinerea, 504
 nigra, 504
 regia, 504
 Jungermanniales, 419
 Juniper, 465
 Juniperus, 466
Juniperus, 465
 communis, 466
 Oxycedrus, 466
 Sabina, 466
 Virginiana, 402, **466**
 Jussieu, A. L., 325
 Jute, bast fiber of, 120

Kalmia, 580
 latifolia, corolla of, 272
 Kamala, 551
 trichomes of, 125
 Karyokinesis, **47**
 Katabolism, **31**
 Kava, 499
 Keel, 274, 538
 Kelps, 353
 Figs. of, 354, 355
 Kernel, 314
 Kieselguhr, 350
 Kino, 92, 540
 Knot-grasses, 512
Kochia, 514
 Kola, 85, **559**
 Krameria, 92, **540**
Krameria argentea, 540
 triandra, 540
 Kühne, 36

 Labarraque's solution, 684
 Labellum, 271
Labiata, 261, **593**
 Labrador tea, 619
Lachnea, 387
 Lactase, 102
 Lactose, 102
Lactuca virosa, 615
 Lactucarium, 101, **615**
 Lamarck, J. B., 653, 654
 Lamarckian Theory, 653
 Lamiaceæ, 593

- Lamina, defined, 55, 246
 forms of, 234
 mode of development of, 246
Laminaria, 354
 Fig. of, 355
Laminaria digitata, 355
 Laminariales, 352, **353**
Landolphia, latex of, 101
 Lappa, 612, **615**
 Larch, 465
Larix, 465
Larix Europæa, 466
Larix laricina, 466
 Larkspur, 518
 calyx of, 263, 266
 spur of, 268
 Latex, **100**, **133**
 cells, 131
 micro-chemic test for, 101
 Laticiferous tissue, 131
 vessels, 132
Lauraceæ, 515, **522**
 anthers of, 278, 279
 Laurel Family, 515, **522**
Laurus, 524
Laurus nobilis, 524
Lavandula Spica, 595, 596
 Lavender, 596
Lawsonia, 570
Lawsonia inermis, 570
 Leaf, 221
 Leaf apex, 234, **235**
 Leaf arrangement, 229
 Leaf base, 234, **236**
 Leaf blade, 221, 247, 249
 Leaf, cleft, 237
 color, 241
 complete, 221
 compound, 233
 cushion, 243
 decompound, 240
 definition of, 221
 divided, 237
 exstipulate, 223
 functions, 223
 imparipinnate, 240
 lobed, 237
 modifications, 240
 palmate, 240
 Leaf, palmately-compound, 234, 240
 palmatifid, 238
 palmatipartite, 238
 palmatisect, 238
 paripinnate, 240
 parted, 237
 parts of typical, 242
 pedate, 239
 petaloid, 241
 petiolate, 223
 pinnate, 240
 pinnately-compound, 233, 240
 pinnatifid, 238
 pinnatipartite, 238
 pinnatisect, 238
 primordial, 228
 sessile, 223
 simple, 233
 sinuses, 237
 stipulate, 223
 Leaf duration, 242
 Leaf fall, 191
 Leaf gaps, 146, **147**
 Leaf insertion, 232
 Leaf margin, **236**, 238
 dentate, 223
 Leaf outline, 234, **235**
 Leaf rank, 229
 Leaf sinuses, 237
 Leaf surface, 241
 Leaf texture, 241
 Leaf traces, 144
 diagrams of, 146
 Leaf venation, 230
 Leaflets, 233
 Staminal, 277
 Leaves, bifacial, **250**, 251
 binate, 240
 bipinnate, 240
 biternate, 240
 bract, 228
 bracteolar, 227
 centric, 249
 succulent, 250
 xerophytic, 250
 compound, 233, **240**
 palmately, **234**, 240
 pinnately, **233**, 240
 convergent, 249

- Leaves, coriaceous, 241
 decompound, 240
 development of lamina of, 246
 dorsiventral, 246
 -hydrophytic, 248
 -mesophytic, 247
 Fig. of cross sect. of, 247
 -umbrophytic, 247
 -xerophytic, 248
 foliage, 228
 forms of, 233
 imparipinnate, 240
 interruptedly pinnate, 240
 lyrate, 240
 origin and development of, 228
 palmate, 240
 paripinnate, 240
 pellucid-punctate, 241
 pinnate, 240
 primordial, 228
 runcinate, 237
 scale, 228
 simple, 233
 stomata of, 251
 ternate, 240
 trifoliate, 240
 types of Angiospermous, 227
 veins of, 230
Lecanora, 416
Lecanora esculenta, 417
Lechea, 509
Lecythidaceæ, 569
Legume, 301
Leguminosæ, 536
 placentation in, 288
 root tubercles of, 176
 stipules of, 245
Lemma, 484
Lemon, 305, **545**
Lenticels, 204
Leontodon Taraxacum, 615
Leonurus cardiaca, 596
Leprosy Plant, 565
Leptandra, 600
Leptome, 133, 143
Leptosporangiatæ, 434
Lettuces, 614
Leucoplastids, **42**, 133
Levisticum, 577
Levisticum officinale, 577
Levulose, 74
Liber fibers, 122
Lichens, 13, **413**, 619
Licmophora, 351
 Life history, of an angiosperm, 471
 of *Aspidium*, 53
 of black mold, 377
 of *Chara*, 347
 of *Claviceps purpurea*, 395
 of *Erythronium*, 471
 of *Ectocarpus*, 352
 of *Equisetum*, 437
 of fern, 53
 of *Fucus*, 355
 of gymnosperm, 444
 of *Lycopodium*, 439
 of a moss, 426
 of mushrooms, 405-409
 of smut, 397
 of *Spirogyra*, 343
 of wheat rust, 399
 of white pine, 444
 of yeast, 382
 of *Zamia*, 456
Light Green, 684
Lignin, 108, **118**
Lignocellulose, 109
Ligule, 242, **245**, 484
Ligulifloræ, **613**, **614**
Ligustrum, histology of root, 170
Lilac, 261, 583
Liliaceæ, 489
Liliales, 489
Lilium, 489, 490
Lily, 192
 Calla, 487
 Canada, 191
 diagram of flower, 488
 Fig. of T. S. anther, 280
 ovary, 473
 pollen of, 281, 283
 Tiger, 189, 191
Lily Family, 489
Lima bean, Figs. of, 320
 histology of, 321
Limb, 266
Limonis Cortex, 546
Linaceæ, 542

- Linaria*, 599, 600
 corolla of, 272
 Fig. of, 273
 stamens of, 275
 Linden, American, 559
 Linden Family, 559
 Flowers, 559
 Linden stem,
 Fig. of T. S. of, 205, 206
 Linin, 18
 Linnæus, Carl, 325
 Linum, 543
 mucilage in, 95
Linum usitatissimum, **542, 543**
 Lipase, 103
Lippia lanceolata, 593
Liquidambar orientalis, 532
 Styraciflua, 532
Liriodendron Tulipifera, 516
 carpels of, 286
 Litmus, 416
 Little Club Mosses, 442
 Liverworts, 13, **418**
 l-mandelonitrile glucoside, 103
Loasa hispida, hairs of, 126
 Lobelia, 163, 274, 275, **610**
Lobelia inflata, 610
Lobeliaceæ, **609, 610**
 Locule, 473
 Locust, corolla of, 271
 modified stipules of, 246
 Lodicule, 484
Loganiaceæ, 583
 Loment, 301
 Fig. of, 303
 Lomentum, 301
Lonchocarpus, 540
Lonicera, 606
 Loosestrife Family, 570
 heterostyly in, 290
 purple, 570
Lophophora Williamsii, 568
Loranthaceæ, 510
 Lovage, 577
 Loxa Bark, 605
 Lugol's solution, 93
Lunaria, 268
 Lungwort, 593
Lunularia, Fig. of, 13
 Lupulin, 125, **127**
 Lupulinum, 510
Lychnis, 266
 Lycoperdales, 410
Lycoperdon, 409, **410**
Lycopodiaceæ, 439
Lycopodineæ, 431, **439**
Lycopodium, 439
Lycopodium clavatum, **439, 440**
Lythraceæ, 570
Lythrum, 290
Lythrum Salicaria, 570
 Mace, 82
 Macfarlane, 627, 629
 Macis, 518
Maclura, 507
Macrocystis, 351, **354**
 Macromorphology, defined, 2
 Madder, 605
 Madder family, 603
 Magnesium as an essential element, 157
 Magnolia, 301, 516
 Magnolia Family, 515
Magnoliaceæ, 515
 Mahogany Family, 548
Mahonia, 519
 Male Fern, 53
 life history of, 53
Mallotus philippinensis, 551
 Mallow Family, 559
 Mallow Flowers, 562
 Maltase, 102
 Maltose, **75**
 Maltum, 485
Malus, **534**, 536
Malus sylvestris, Fig. of flower, 271
Malva, 560
 Malvæ Flores, 562
 Malvæ Folia, 562
Malva rotundifolia, 401, **559, 562**
 sylvestris, 401, **562**
Malvaceæ, 559
 anthers of, 278, 279
 Malvales, 557
 Manaca, 598
 Mangrove bark, 92
 Mangrove Family, 569
 Mangroves, 619

- Manihot*, 550
Manihot Glaziovii, 101
 utilissima, 103, **551**
 Manna, 583
 Scriptural, 417
 Manna Ash, 74
 Manzanita, 582
 Maple, 162, 303
 Red, 185
 Maple Family, 554
Maranta arundinacea, 493
 starch, **79**, 81
Maranta, petiole of, 243
Marantaceæ, 314, **492**
Marchantia, 418
Marchantia polymorpha, 419
 life history of, 419
 Marchantiales, 419
 Marrubium, 595
Marrubium vulgare, 595
 Fig. of cross sect. of leaf blade, 249
 hairs of, 124, **125**
Marsdenia Condurango, 589
 Marsh, defined, 618
 Marshmallow, 559
Marsilea, 437
 Masterwort, 577
 Mastiche, 552
 Maté, 85, **552**
 Matico, 232, **499**
 Matricaria, 614
Matricaria Chamomilla, 613, **614**
 Maturation of pollen grain and formation of male gametophyte in Angiosperms, 292
 in Erythronium, 474
 in White Pine, 449
 of embryo sac and formation of female gametophyte in Angiosperms, 292
 Maw seed, 528
 May Apple, calyx of, 268
 Mayer's albumin fixative, 698
 Mechanical tissues, 151
 Medulla, **165**
 Medullary rays, 139
Megaceros, 419
 Megasori, 285, 448
 Megasporangia, 381
 Megasporangium, 448, 458
 Megaspore, **287**, 448, **449**, 460, 474
 Megasporophyll, defined, 283
 Megasporophylls, 448, 473
 Meiosis, 50
Melaleuca Leucadendron, 574
Melia Azedarach, 548
Meliaceæ, 548
 Melibiose, 75
 Melilotus, 540
Melilotus officinalis, 540
 Melissa, 596
Melissa officinalis, 596
Melochia, 558
 Melon tree, 567
 Membrane crystals, 90
 Membrane mucilage, 94, 95
 Membranes, 19
 cell, **43**
 plasma, 19
 semipermeable, **43**
 vacuolar, 19, **43**
 Mendel, G. J., 643, 644
 Mendel's Laws, 643
Menispermaceæ, 521
 Menispermum, 521
Menispermum canadense, 521
Mentha arvensis, 595
Mentha piperita, 594
 Fig. of glandular trichomes, 125
Mentha Piperita, U.S.P., 594
Mentha spicata, 595
Mentha Viridis U.S.P., 595
Mentha viridis, 595
 Menyanthes, 587
Menyanthes trifoliata, 587
 Mericarp, 303
 histology of a, 309
Merismopedia, 336
 Meristeles, 145
 Meristem, 114
 primordial, 114, 187
 Meristems, 114
 apical, 115
 intercalary, 115
 lateral, 116
 primary, 114
 secondary, 114
Merulius, 404

- Mesocarp, 298
 Mesophyll, **61, 222**
 Mesophytes, 164, 617, **619**
 Mesquite Gum, 540
 Metabolism, **31**
 Metachlamydeæ, 579
 Metaphase, 50
 Metaxylem, 143
 Methyl Salicylate, 103, 505
 Methylis Salicylas, 505, 581
Metroxylon Sagu, 82
 Mez., Carl, 323
 Mezereum, 570
Micrandra, 550
Micrococcus, 365
 division forms of, 368
 ureæ, 182
 Micro-crystals, 90, **91**
 Microgametophyte, 461
 Micrometer, net, 673
 ocular, 672
 stage, 673, 674
 Micrometry, 672
 Micromorphology, defined, 2
 Micron, defined, 672
 Micropyle, **287**, 448, 474
 Microscope, binocular, 670
 care of, 671
 compound, 663, 664
 use of, 669
 defined, 661
 diagram showing optics of, 665
 dissecting, 661, 662
 research, 667
 simple, 661
 Microscopic measurement, 672
 Microsorus, **274, 473**
Microspira, 372
 Microsporangia, 277, 381, 447, 457, 473
 Microspore, 280, 447
 Microsporophylls, **274**, 277, 447, 457
 Microtomes, 677
 hand, 677
 rotary, 679, **680**
 sliding, 678, 679
 Middle lamella, 20, 21, **24, 49**
 Midrib, 230
 Mildew, 388
 Milkweed, 295, **313**
 pollinia of, 283
 Milkweed Family, 589
 Milkwort Family, 548
 Milon's reagent, formula for, 93
Mimosa, **36, 538**
 pudica, 37
 conduction of stimuli by leaves, **37**
 Spegazzini, 36, **38**
 thigmotropic reactions in, **36, 37**
Mimosaceæ, 538, **539**
Mimusops globosa, 582
 Mint Family, 118, **593**
 Miré, 598
 Mistletoe, **161**, 479, **510**
 Mistletoe Family, 510
 Mitchella, 605
Mitchella repens, 605
Mitella, 531
 Mitochondria, **43**
 Mitosis, 47
 heterotypic, 52
 homotypic, 52
 Mitrewort, 531
Mnium, 426, 427
 Mock-orange, 531
 Mold, black, 377
 blue, 388
 bread, 377
 green, 388
 Molds, cultivation of, 376
Mollugo, 514
Momordica Balsamina, 608
Monarda, 594, 596
Monarda punctata, 596
Monilia, 412
Monimiaceæ, 515, **522**
 Monkshood, 518
 Monoclinic prisms, **88**, 89
 Monocotyl
 leaves, 242, **255**
 rhizome, 215
 root, 165
 seed, 315
 seedling, 316
 stems, 215-220
 secondary growth in, 220

- Monocotyledon, morphology of typical
 plant of, 481
- Monocotyledoneæ, 479
- Monocotyledons, 15, 16, **481**
 development of embryo in, 296
 drug yielding, 483
 histology of leaves of, 255
 histology of roots of, 165
 histology of stems of, 215
 petiole in, 242
- Monœcious, **264**, 418, 425
- Monohybrids, 648
- Monosaccharoses, 74
- Monotropa*, 580
 uniflora, 580
- Monstera*, 487
- Moonseed Family, 515, **521**
- Moonwort, 434
- Moraceæ*, 507
- Morchella esculenta*, 395
 Fig. of, 394
- Morels, 395
- Morgan, T. H., 658
- Morning Glory, 188, 300
- Morning Glory Family, 590
- Morphine, **86**, **527**
 test for, 87
- Morphology, defined, 2
- Moss, diagram of life history of, 429
 Iceland, 414
 Golden, 435
 Reindeer, 417
- Mosses, 418, **424**
 bog, 425
 peat, 425
 rock, 426
 true, 426
- Motherwort, 596
- Mountain Ash, 74, **535**
- Mountain Laurel, 272, **579**, **580**
- Mountain Mint, 596
- Mountain Rose, 512
- Mounts, Canada balsam, 689
 glycerin-gelatin, 691
 permanent, 687
 temporary, 687
- Movement, 39
 amœboid, 26, **39**
 Brownian, 365
- Movement, ciliary, **39**
 true, 365
- Mucilage, 94
 cell content, 94, **95**
 membrane, 94, **95**
 micro-chemic test for, 95
- Mucor mucedo*, 380
 stolonifer, 377
- Mucuna pruriens*, 540
- Muehlenbeckia*, 512
- Mulberry, 307, **507**, **509**
- Mulberry Family, 91, **507**
- Mullein, 259
- Musa*, 231
- Musaceæ*, 492
- Musci, 13, **418**, **424**
- Mushroom, common, 405
- Mustard, corolla of, 270
 fruit of, 302
 stamens of, 275
- Mustard Family, 268, **528**
- Mutation Theory, 657
- Mutations, 653, 657, 659
 Chromosome, 658
- Mutisia*, 611
- Mycelium, **362**, 377, 388, 397
- Mycobacteriaceæ*, 372
- Mycobacterium*, **372**, 373
- Mycology, 3
- Mycorrhiza, 181, 182, **183**, 394
- Myosotis*, 592
- Myrica, 502
- Myrica caroliniensis*, 96, **179**, **502**
 cerifera, 96, **179**, **502**
 Gale, 179
 Fig. of, 503
 Macfarlanei, 179
 tubercles of, 179
 xalapensis, 96
- Myricaceæ*, characters of family, 501
 root tubercles of, 179
- Myristica*, 517
 fragrans, 518
 Fig. of, 516
- Myristica, U.S.P., 518
- Myristicaceæ*, 515, **517**
- Myrobalans Family, 574
- Myronase, 103
- Myrosin, 103

- Myrrh Family, 547
 Myrrha, 548
Myrtaceæ, 251, **571**
 Myrtales, 568
 Myrtifloræ, 568
 Myrtle Family, 571
 Myxomycetes, 362, **374**
- Naphthol solution, 684
 Narcissus, 274
 Nasturtium, 291
 Nasty, **32**
 Natural Selection, 654, 656
 Natural system, 323
 Naturalized, defined, 3
Nectandra Coto, 524
 Nectar, 150
 Nectaries, 150
 sepaline, 268
Negundo, 554
 Neighboring-cells, 251
Neisseria gonorrhoeæ, 373
 intracellularis, 373
Nepenthaceæ, 530
Nepenthes, 241, 245, **634**, 638
 bicalcarata, 637
 Edwardsiana, 637
 enzyme in, 103
 petiolar structure in, 245
 Rajah, 634, **635**, 638
 Nepenthin, 103
Nepeta Cataria, 595
 Nephrodium. See *Dryopteris*
Nereocystis, 354
 Nest fungi, 410
 Nettle family, cystoliths in, 91
 Nettle, hairs of, 127
Nicotiana, 85
Nicotiana tabacum, 598
 Nicotine, 85
 micro-chemic test for, 85
 Nidulariales, 410
Nigella, 519
 Nightshade, Fig. of, 535
 Family, 533
Nitella, 40, **346**
 Nitrification, 367
 Nitrifying bacteria, **182**, 367
Nitrobacter, 182
 Nitrogen cycle, 182
 Nitrogen fixation, 178
Nitrosococcus, 182
Nitrosomonas, 182
 Node, defined, 186
 Nomenclature, 329
 Binomial system of, 325
 Non-protoplasmic cell contents, 74
Nostoc, 336
 Fig. of, 334
Nucellus, **287**, 448, 474
 Nuclear membrane, 19, **43**
 Nuclear sap, **41**
 Nuclei, assisting, 292
 polar, 292
 Nucleins, 93
 Nucleolus, 18, **41**
 Nucleoplasm, **41**
 Nucleus, 18, **41**
 Numerical aperture, 666
 Nut, 304
 Nutgall, Fig. of, 505
 Nutlet, 302
 Nutmeg, 314
 Nutmeg Family, 515
 Nutrition, defined, **31**
 of *Amœba*, 26
 of bacteria, 366
 saprophytic, 383
Nux Vomica, 585
Nyctaginaceæ, 514
 Nyctinasty, **32**
Nympheaceæ, 515
Nyssa sylvatica, 577
- Oak, 259
 Oaks, 297, 304, **506**
 Oat, 484
 inflorescence of, 259
 Objectives, microscope, **664**, **666**
 numerical aperture of, 666
 working distance of, 666
Ochrea, **246**, 512
Ocimum Basilicum, 596
 Octant cells, 71
 Ocular micrometer, 672
 standardization of, 673
Cedogonium, 338
Oenothera, 282, **569**

- Oenothera lamarckiana*, 657, 658
Oenothera lata, 658
 Offsets, 189
Oidium, 376
 Oil, benne, 602
 bitter almond, 97
 cassia, 97
 cedarwood, 466
 chaulmoogra, 565
 cocoanut, 487
 enzyme, 103
 linseed, 542
 mustard, 97
 palm, 487
 rose geranium, 542
 Siberian fir, 466
 tung, 551
 turpentine, 97
 Oils, fixed, 95
 nitrogenated, 97
 oxygenated, 97
 sulphurated, 97
 terpene, 97
 volatile, 96
Olea, 583
Olea europæa, 583
Oleaceæ, 583
 Oleander, latex of, 101
 Oleoresins, 97
 defined, 98
 Oleum abietis, 466
 amygdalæ amaræ, 535
 amygdalæ expressum, 535
 anisi, 516, 577
 aurantii, 546
 aurantii amari, 546
 aurantii florum, 546
 bergamottæ, 546
 betulæ, 505
 betulæ empyreumaticum rectifi-
 catum, 505
 cajuputi, 574
 cardamomi, 492
 cari, 577
 caryophylli, 574
 chaulmoogræ, 565
 chenopodii, 514
 cinnamomi, 524
 citronellæ, 485
 Oleum coriandri, 577
 erigerontis, 615
 eucalypti, 574
 foeniculi, 577
 gossypii seminis, 562
 juniperi, 466
 lavandulæ, 595
 limonis, 546
 lini, 543
 maydis, 485
 menthæ piperitæ, 594
 menthæ viridis, 594
 myrciæ, 574
 myristicæ, 518
 olivæ, 583
 picis rectificatum, 465
 pimentæ, 574
 pini pumilionis, 466
 ricini, 551
 rosæ, 535
 rosmarini, 595
 santali, 510
 santali australiensis, 510
 sassafras, 524
 sesami, 602
 sinapis volatile, 530
 terebinthinæ, 465
 theobromatis, 559
 thymi, 595
 tiglii, 551
 Olibanum, 548
 Olive, 583
 Olive Family, 583
Onagraceæ, 569
 Onion, 188, 191
 epidermis of bulb scale, 18
Onoclea sensibilis, 433
 Oögonia, 346, 348
 Oögonium, 348
 Oömycetales, 376
 Oöspore, 46, 293, 341, 476
Operculina Turpethum, 591
 Operculum, 429
Ophioglossum, 434
 Opium, 86, 87, 101, 527
Opuntia, 567, 568
 Opuntiales, 567
 Orange, 545
 flower, 277

- Orange, fruit of, 305
 peel, cross section, 148
 stamens of, 277
 Orcein, 416
 Orchid family, 494
Orchidaceæ, 494
 Orchidales, 494
Orchideæ, 495
 Orchids, Fig. of pollinia, 282
 floral organs of, 494
 parenchyma of, 116
 stamens of, 275
Orchis mascula, 496
Orchis militaris, Fig. of flower, 272
Orchis morio, 496
 Order, defined, 327
 naming of, 332
 Ordinary Parenchyma, 116
 Oregon balsam, 466
 Organ, defined, 6
 Organic cycle, 363
 Organic-acid enzymes, 103
 Organism, defined, 6
 Organs, accessory of flower, 262
 essential of flower, 262
 plant, 6
 reproductive, 6
 vegetative, 6
 Origanum, 596
Origanum Majorana, 596
 vulgare, 596
 Orris, 492
 Osage Orange, 297, **307**, **507**
Oscillatoria, 335
 Fig. of, 11, 334
 Osmic acid, 692
 Osmosis, 153
 applied to root hairs, 154
Osmunda cinnamomea, 433, 436
 Claytoniana, 436
 regalis, 436
Ostraya, 504
 Ostwald, 101
Ourouparia Gambir, 605
 Outline of plant groups, 328
 Ovary, **284**, **473**
 of Poppy, 528
 Ovule, amphitropous, 287
 anatropous, 287, 473
 Ovule, campylotropous, 287
 forms of, 287
 orthotropous, 287
 Ovules, **285**, 448, **473**
 Ovum, 69, 450, **474**
 Oxalic acid, production by molds, 394
 Oxidizing enzymes, 104
 Oxytropism, **34**

Pachystima, 552, 553
Padus, 535
Palaquium, 581, 582
Palaquium gutta, 101
 Palate, 274
 Palet, 484
 Palm, Brazilian Wax, 96
 Cocoanut, 485, 486
 Date, 486
 Family, 485
 Fern, 455
 oil, 487
 petiole, 242
 Sea, 354
 stem, 214
Palmae, 485
 petiole of, 242
 stigmata in, 91
Panax Ginseng, 576
Panax quinquefolium, 576
 repens, 576
Pandanus, roots of, 161
Pandorina, 338
 Panicle, 258, **259**
 Pansy, 566
 Papain, 103, 567
Papaver, 524
 calyx of, 267
 latex of, 101
 placentation in, 288
 Papaver Rhœas, 528
Papaver somniferum, 101, **527**, **528**
Papaveraceæ, 524
 Papaverales, 524
 Papaverinæ Hydrochloridum, 527
 Papaveris Fructus, 528
 Papaw, 567
 Papaw Family, 566
 Papayin, 103
Papilionaceæ, 537, **538**

- Papillæ, 124
 Pappus, 266, 297
 Paprika, 598
Papyrus, 618
 Paraguay Tea, 85
 Paraphyses, **356**, 387, 388, 427
 Parasite, 362
 Parasites, facultative, 367
 obligate, 367
 Pareira, 162, **521**
 Parenchyma, 116
 assimilation, 116
 border, 223, 224
 conducting, 117
 ordinary, 116
 palisade, 117, 222
 phloem, 117, 135
 reserve, 117
 sieve, 135
 spongy, 117, 223
 wood, 117
 Parietales, 562
Parmelia, 414
Parmelia perlata, Fig. of, 415
 Parsley Family, 118, 148, **243**, **576**
 Partridge Berry, 190, **605**
 Pasque Flower, 518
Passiflora, 266, **566**
 incarnata, **566**, **567**
Passifloraceæ, 566
 Passion Flower, 188, **567**
 Passion Flower Family, 566
 Pasteur's Solution, 382
 Pathogens, 364
 Pathology, 3
Paullinia, 553, 554
Paullinia Cupana, 554
Paulownia, 599
Paulownia imperialis, 599
Payena, 101, 582
 Pea, 82, **240**
 tendrils of, 189
 Pea Family, 536
 Peach, 301, 305, **534**, 535
 Peachwood, 540
 Peanut, 301
 Pear, 535
 fruit of, 305
 pericarp of, 298
 Pear, stone cells of, 119, 120, 301
 Peat mosses, 425
 Pecan, 504
 Pectase, 102
 Pectinase, 102
 Pectose, 23
Pedaliaceæ, 601
 Pedicel, 256
Pedicularis, 599
 Peduncle, 256, 262
Pelargonium, 301, 541
 capitatum, 542
 odoratissimum, 542
 Radula, 542
Pellæa, 435
 Pelletierinæ Tannas, 571
 Pengawar Djambi, 435
Penicillium, 388
 brevicaule, 391
 camemberti, 390, **391**
 expansum, 391
 glaucum, 388
 roqueforti, 389, **390**
 Pentosides, 84
Pentstemon, 600
 stamens of, 275
 Peony, 518
 Fig. of flower, 284
 Pepo, 305, **608**
 Pepper, Black, 499
 Red, 597
 Pepper Family, 499
 Peppermint, 594
 Pepsin, 103
 Perennial plants, 163
Pereskia, 567, 568
 Perianth, 262, **266**
 forms of, 270
 outgrowths of, 274
 Periblem, 113
 Pericambium, 56, **131**
 Pericarp, 297
 Pericladium, 243, 576
 Pericycle, 56, 63, **131**
 Pericyclic fibers, 122, 131
 Periderm, defined, 207
 Peridiolum, 411
 Peridium, 401, 410

- Perigone, 266
 appendages of, 274
 Perisperm, 314
 Peristome, 429
 Perithecia, 391, 395, 416
 Periwinkle, 258
 Peroxidase, 104
Pertusaria, 414
 Petals, 269
 Petiole, 223, 242
 gross structure and histology of, 242
 Petiolule, 234
 Petroleum Ether, 685
 Petroselin Radix, 577
 Petroselinum, 577
Petroselinum sativum, 577
 Petunia, 596
Peumus Boldus, 522
 Peyote, 568
Peziza, 387
 repanda, Fig. of, 386
 Pezizales, 387
 Pfeffer, 33
 Phæophyceæ, 351
 Phæosporales, 352
Phalaris canariensis, 485
 Phallales, 411
 Phanerogamia, 325, 443
 Phanerogams, 14
 Pharmaceutical Botany, defined, 3
 Pharmacognosy, defined, 3
Phaseolus lunatus, structure of seed of, 319
Phaseolus multiflorus, cross-sections of
 root of, 167
 hooked hair from, 126
Phegopteris, 435
 Phellem, 207
 Phelloderm, defined, 208
 Phellogen, 114, 129, 131
 Phenotype, 647, **648**
Philadelphus, 531
 Phloem, 140
 defined, 143
 internal, 204
 interxylary, 204
 intraxylary, 204
 parenchyma, **117, 135**
 primary, 143
 ray, 139
 Phloem, secondary, 143
 Phloroglucin, solution of, 685
 Phlox, corolla of, 272
Phoradendron flavescens, 510
 Photogens, 364
 Photosynthesis, **64, 66**
 defined, 224
 Phototropism, 35
Phragmidiothrix, 372
Phragmites, 618
 Phycocyanin, **100, 334, 357**
 Phycoerythrin, **100, 335, 357**
 Phycophæin, 100
 Phycomycetes, 375, **376**
 Phycophæin, 351, 352
 Phycoxanthin, 351
 Phylloclades, 188
 Phyllode, 244
 Phyllotaxy, 229
Physalis, 297
 Fig. of calyx, 267
Physica, 414
Physica stellaris, Fig. of, 413
 Physiological division of labor, 16
 Physiology, defined, 2
 Physostigma, 539, **540**
 style of, 289
Physostigma venenosum, **540, 541**
 Physostigminæ Salicylas, 540
 Phytogeography, defined, 3
 Phytopalæontology, defined, 3
 Phytoglobulins, 94
Phytolacca, 514
 stems of, 203
Phytolacca americana, 514, 515
Phytolacca, N.F., 514
Phytolaccaceæ, 514
 Phytopalæontology, defined, 3
 Phytopathology, defined, 3
Picea, 465
 canadensis, 466
 excelsa, 466
 mariana, 466
 rubra, 466
 Pickerel weed, 116
Picrasma excelsa, 546, **547**
 Picric acid, 39
 Pigments, 98
 Pigweed, pollen of, 283

- Pileus, 404
 Piliferous layer, 152
 Pilocarpinæ Hydrochloridum, 546
 Nitrates, 546
 Pilocarpus, 546
Pilocarpus, 545
Pilocarpus Jaborandi, 546
 microphyllus, 546
 neighboring-cells of leaf epidermis,
 252
 Pimenta, 574
Pimenta acris, 574
 officinalis, 574
 Pimiento, 598
 Pimpinella, 577
Pimpinella Anisum, 576
 magna, 577
 Saxifraga, 577
 Pinaceæ, 465
 Pine, 465
 Austrian, **250, 447, 451**, 453
 Drops, 580
 Family, 465
 inflorescences of, 464
 leaf of, 446
 pollen of, 282
 Scotch, 448
 stem, 445
 tar, 465
 White, 444
 wood of, 212
 Pineapple, enzyme in, 103
 fruit of, 307
 Pinenes, 97
Pinguicula, 640
 vulgaris, **640, 641**
Pinites succinifer, 466
 Pink Family, **273, 514**
 Pink Root, 583, **585**
 Pinna, **55**, 433
Pinnularia, Figs. of, 349, 352
 Pinnule, **55**, 433
Pinus, 465
 austriaca, 447
 maritima, 466
 montana, 466
 nigra, leaf of, 250
 palustris, 465
 Strobilus, 401, **444, 465**
Pinus, sylvestris, 448
Pinus Alba, N.F., 465
 Pinweeds, 509
 Piper, 305, **499**
Piper angustifolium, 499
 Cubeba, 498, **499**
 longum, 499
 methysticum, 499
 nigrum, 498, **499**
 officinarum, 499
Piperaceæ, 499
 formation of endosperm and peri-
 sperm in, 314
 Piperales, 499
 Pipsissewa, 579
 Piscidia, 540
Piscidia Erythrina, 540
 Pistachio, 552
Pistacia vera, 552
 Lentiscus, 552
 Pistil, 285, 473
 bicarpellary, 285
 compound, 285
 monocarpellary, 285
 polycarpellary, 285
 system, 283
 tricarpellary, 285
 Pitcher plants, 629
 Pith, 165, 166
 Pits, 108
 bordered, 136
 Pix Burgundica, 466
 Canadensis, 466
 Juniperi, 466
 Pini, 465
 Placenta, defined, 287
 Placentation, 287, 288
 Plankton, 617
Planococcus, 371
Planosarcina, 371
 Plant, acaulescent, 186
 anemophilous, 291
 annual, 162
 aquatic, 164
 associations, 616
 axis, 6
 biennial, 163
 breeding, 3
 caulescent, 186

- Plant, cell, 41
 cigar, 570
 communities, 616
 dioecious, 264
 entomophilous, 291
 genetics, 3
 geography, 3
 hairs, 124
 halophytic, 164
 homozygous, 647
 heterozygous, 647
 hydrophilous, 291
 indigenous, 3
 malacophilous, 292
 mesophytic, 164
 monœcious, 264
 naturalized, 3
 organs, 6
 ornithophilous, 291
 parts of body of, 6
 perennial, 163
 pitcher, 629
 polygamous, 264
 seed, 7
 tissues, 111
 xerophytic, 164
- Plant Succession, 621
- Plantaginaceæ*, placentation in, 288
- Plantaginis Semen, 603
- Plantago*, 44, 301, **603**
- Plantago arenaria*, 603
- Plantago lanceolata*, 603
- Plantago ovata*, 603
- Plantago Psyllium*, 602, 603
- Plantain, 259, 603
- Plants, aquatic, 164, 617
 carnivorous, 621
 classification of, 323
 desert, 617
 general characteristics of, 6
 groups of, **12, 328**
 intermediate, 617
 marsh, 618
 naming of, 323, **329**
 relation of, and animals, **29**
 relation to animals, 621
 salt, 617
 water, 617
- Plasma membranes, **19, 22**
- Plasmodesma, **44**
- Plasmodiocarp, 374
- Plasmodium, 374
- Plasmolysis, 155
 illustrating, successive stages of, 154
- Plastids, **42**
- Platystemon*, 524
- Plectascales, 388
- Plerome, 113
- Pleurococcaceæ*, 337
- Pleurococcus*, 337
 Fig. of, 338
- Pleuston, 617
- Pliny, 2
- Plum, 305, 535
- Plumule, 311
- Pod, 301
- Podophyllum, 519
- Podophyllum emodi*, 519
 peltatum, 519
 Fig. of, 521
 petiolar structure of, 245
- Podophyllum Indicum*, 519
- Poison Hemlock, 516
 Fig. of, 578
- Poisons, 33
- Poke, Family, 514
 Root, **175, 514**
- Polemoniales, 590
- Pollen, 278
 chamber, 449, 458
 grain, **280**
 mother cells, 278, **474**
 stages in germination, 293
- Pollen grains, 282, 457
 Fig. of, 281, 450
 pine, 281, **450**
- Pollens, Hay-fever, 283
- Pollination, 290
 cross, 291
 in erythronium, 475
 in pine, 450
 self, 290
- Pollinia, 282, **283, 589**
- Polycotyledony, 227
- Polyembryony, 479
- Polygala*, 103, 283
 lutea, 549
 pollen of, 283

- Polygala Senega*, 549, 550
Polygalaceæ, 548
Polygonaceæ, 246, 314, 512
 Polygonales, 512
Polygonum, 512
 Bistorta, 513
 Polypeptides, 93
 Polypetalæ, 499
 Polyploidy, 658
Polypodiaceæ, 435
Polypodium, 434, 435
Polyporaceæ, 404
Polyporus officinalis, 404
Polystichum, 435
Polytrichum commune, Fig. of, 427
 life history of, 426
 Pome, 305
 of apple, Fig. of, 306
 Pomegranate Family, 570
Ponteria, 582
 Poplar, 500
 Poppy, calyx of, 267, 268
 Family, 524
 fruit of, 299, 302
 leaf of, 232
 Populi Gemma, 500
Populus balsamifera, 500
 candicans, 500
 tremuloides, 500
 Pore Fungi, 404
Porella, 418, 419
 Pores, 108, 110
 bordered, 136
 germ, 282
Portulaca, 301, 514
Postelsia, 354
 Potash, solutions of, 685
 Potassium ferrocyanide solution, 685
 Potato, 87, 192, 272, 596
 starch of, 79, 81
Potentilla, 269, 535
 silvestris, 536
 Prefloration, 264
 Prefoliation, 230
Premna arborea, 593
 Prickles, 189
 Prickly Ash, 546
 Prickly Pear, 567
 Primary Meristems, 113
 of root, 157
 Primordial meristem, 111
 of leaf, 228
Primulaceæ, placentation in, 288
 Principes, 485
Prinos, 552
 Privet, 261, 583
 California, 170
 Procambium, 113
 Prodromus, 326
 Proembryo, 348
 Promycelium, 397, 398, 401
 Prophases, in nuclear division, 48, 49
 Proplastids, 42
Prosopis juliflora, 540
 Prostele, 145
Proteaceæ, 254
 Protective tissues, 151
 Proteinaceous enzymes, 103
 Proteinoplastids, 133
 Proteins, 92
 formation of, 65
 tests for, 93
 Prothallia, female, 438
 Fig. of fern, 69
 male, 438
 Prothallial cushion, 69
 Prothallium, 68
 Fig. of fern, 70
 Prothallus, 68, 451, 460
 Protista, 30
 Protoascales, 381
 Protococcales, 337
 Protoderm, 113
 Protonema, 68, 426
 Protopectin, 107
 Protopectinase, 102
 Protophloem, 144
 Protoplasm, 10, 25, 30
 properties of, 30, 31
 Protoplasmic cell contents, 40
 Protoplasmic movement, 39, 40
 Protoplast, 17
 Protoxylem, 143, 168, 169, 196
 Protozoa, 24
 Prune, 305
 Prunum, 536

- Prunus armeniaca*, 534
 Avium, 534
 Cerasus, 95, 535
 Fig. of flower, 270
 domestica, 536
 Laurocerasus, 536
 serotina, 534
Prunus Virginiana, 535
Psalliota campestris, 405
Pseudobulbs, 188
Pseudocarp, 298
Pseudomonas, 372
Pseudomonas radicicola, 176
 Fig. of, 178
Pseudopodia, 26, 374
Pseudotsuga, 465
Pseudotsuga taxifolia, 466
Psyllium seeds, 95, 602
 testa of, 107
Ptelea, 545, **546**
Ptelea trifoliata, 546
Pteridophyta, 13, 431
Pteridophytes, 13, **431**
Pteris aquilina, 433
 cross section of rhizome, 435
Pteris, Fig. of leaf of, 434
 Fig. showing development of embryo, 72
Pterocarpus marsupium, 540
 santalinus, 539
Pterospora, 580
Puccinia Graminis, 399
 Fig. of, 400
Puccinia malvacearum, 401
Puff Balls, 409
Pulmonaria officinalis, 593
Pulsatilla, 518
Pulsatilla patens, 519
 pratensis, 519
 vulgaris, 519
Pulvinus, 243
Pumpkin, 607
Punica, 571
 Granatum, 571
Punicaceæ, 571
Purging Cassia, 332
Purified Siliceous Earth, 350
Purple Cone-flower, 612
Putamen, 298
Putrefaction, 364
Pycnanthemum montanum, 596
Pycnia, 401
Pyrenoid, 337, 341
Pyrenomycetales, 395
Pyrethri Flores, 615
Pyrethrum, 615
Pyrogallol, 507
Pyrola, 580
Pyrus, 535
Pyxidium, 301
Pyxis, 301

Quassia, 547
Quassia amara, 546, **547**
Quercus, 507
Quercus alba, 506, 507
 chrysolepis, 506
 garryana, 506
 infectoria, 507
 Fig. of, 505
 Michauxii, 506
 occidentalis, 506
 pedunculata, 15
 platanoides, 506
 prinus, 506
 Suber, 506
 velutina, 506
 virginiana, 506
Quillaja, 535
Quillaja Saponaria, 88, **534**, **535**
Quillworts, 431, **442**
Quince, 535
 calyx of, 267
 fruit of, 305
 pericarp of, 298
 receptacle of, 297
Quinidinæ Sulfas, 605
Quinina, 605

Raceme, 258
Rachilla, **259**, **484**
Rachis, 233, 256
Radial bundles, 142
Radicle, **311**, 317, 321
Radish, 161, 162
Ragweeds, pollen of, 283
Ranales, 515
Ranunculaceæ, 243, 515, **518**

- Ranunculus*, 248, 518
 Raphe, 287
 Raphides, 89, **90**
 Raspberry, 189
 fruit of, 300, **306**
 stems of, 186
 Ray, J., 325
 Rays, medullary, 139
 vascular, 139
 Reagents, 680
 Receptacle, 262, **265**
 Receptacles of *Fucus*, **355**, 356, **357**
 of *Marchantia*, 421
 Red Algæ, 95, **357**
 Red Gum, 90
 Red Squill, 490
 Reductase, 103
 Reduction division, 51
 Redwood, 465
 Regma, 301
 Reindeer Moss, 417
 Rejuvenescence, 45
Remijia pedunculata, 605
 purdicana, 605
 Reproduction, asexual, **44**, **45**, **378**
 defined, **38**, **44**
 sexual, **44**
 Reserve cellulose, 109, 487
 parenchyma, 117
 starch, 76
 Resin duct, Fig. of, 150
 Resina, 465
 Resinogenous layer, 96
 Resins, 97, **98**
 Respiration, defined, 66, **225**
 distinction from photosynthesis, 66
Rhamnaceæ, 554
Rhamnales, 554
 Rhamnase, 103
 Rhamnose, 103
Rhamnus cathartica, 556
Rhamnus Frangula, 103, **555**, **556**
Rhamnus Purshiana, **554**, **556**
 Rheotropism, **34**
Rheum, 512
 officinale, 513
 palmatum, 512, 513
Rhinanthus, 599
 Rhizoclonium, Fig. of, 317
 Rhizoids, **65**, 347, 420, **426**
 Rhizome, defined, 192
 histology of monocotyl, 215
Rhizophora, 619
Rhizophoraceæ, 569
Rhizopus, 378
 nigricans, 377
 strains of, 380
 Rhododendron, **580**
 Rhodophyceæ, 357
Rhodymenia palmata, 361
 Rhœadales, 524
 Rhubarb, 513
Rhus, 92, 305, 552
 Cotinus, 551
 glabra, 551, 552
 hairs of, 124, 125
 japonica, 552
 semialata, 552
 succedeana, 96
 sylvestris, 96
 toxicodendron, 551, 552
 typhina, 552
 venenata, 551
 vernifera, 96
 Rhytidome, 130
 Ribbon Bush, 512
Riccia, **418**, 419, 617
Ricciocarpus, 617
 Rice starch, **79**, 80
Ricinus communis, 256, 312, **551**
Rivularia, 335, **336**
Robinia, 213
Rocella tinctoria, 416
 Rock Rose Family, 567
 Rockweeds, 354
 Root, adventitious, 160
 branching, 176, 177
 California Privet, 170
 cap, 158
 classification as to form, 161
 classification as to order, 160
 conical, 162
 consistency, 162
 defined, 152
 distinction from stem, 160
 duration of, 162
 epiphytic, 161
 fibrous, 160, **162**

- Root, functions, 6, 152
 fusiform, 161
 generative tissues of, 158
 growing point of, 158
 hairs, **152**, 154
 histology, 163-176
 lateral, 160
 main, 160
 moniliform, 162
 napiform, 162
 Phaseolus, 167
 primary, 142, **160**, 319
 regions of, 158
 secondary, **160**, 177, 319
 Smilax, 165
 soil and water relation of, 152
 systems, 164, 177
 tap, 160
 texture, 162
 tortuous, 162
 tubercles, 176
 Rootlet, 152
 Roots, 160
 abnormal structure in, 175
 brace, 161
 Dicotyledon, 167
 fascicled, 162
 fleshy, 162
 Monocotyledon, 165
 multiple, 162
 napiform, 162
 nodose, 162
 nodule producing, 176
 of parasitic plants, 161
 of primary growth, 167
 of secondary growth, 169
 prop, **161**, 319
 secondary, 160
 tuberous, 162
 woody, 162
Roridula, 530
 Rosa, 535
Rosa canina, 536
 centifolia, 536
 damascena, 535
 gallica, 535
Rosaceæ, 245, **534**
 Rosales, 531
 Rose, 535
 Family, 534
 fruit of, 304
 Rosemary, corolla of, 272
 Rose of Sharon, 560
 Rosette aggregates, 89, **90**
 Rosin, 465
Rosmarinus officinalis, 595, 596
 Rotation, crop, 179
 cytoplasmic, **39**
 of crops, 174
 protoplasmic, 22
 Rubber, **101**, 551
Rubiaceæ, 245, **603**
 Rubiales, 603
Rubia tinctorum, 605
 Rubus, 536
 Rubus Idæus, 535
Rubus Idæus, 535
 strigosus, 536
 Rue, 286, 545, **546**
 Rue Family, 545
Ruellia, 602, 603
Ruellia ciliosa, 603
Rumex, **512**, **513**
 acetosella, 512
 crispus, 513
 obtusifolius, 513
 pollen of, 283
 Runner, defined, 189
Ruppia, 291
 Ruscus, 188
 Russian Thistle, pollen of, 283
 Rust, 399
 Cedar, 402
 Hollyhock, 401
 Wheat, 399
 White Pine Blister, 401
 Ruta, 545
Ruta graveolens, **545**, **546**
Rutaceæ, 545
Ruteæ, 545
 Ruthenium Red and Lead Acetate
 Solution, 686
 Rye, 259
 starch of, 78, **79**
 Sabadilla, 490
 Sabal, 305, **487**

- Sabbatia, 587
Sabbatia angularis, 587
 Sabina, 466
 Sac Fungi, 381
Saccharomyces, 381
 apiculatus, 386
 cerevisiæ, 382
 ellipsoideus, 385, **386**
 membranifaciens, 386
 pastorianus, 387
Saccharomycetaceæ, 381, **382**
Saccharum, 484
 officinarum, 485
 Saddle Fungi, 394
 Safflower, 615
 Saffron, 492
 Safranin, solutions of, 686
 and Aniline Blue, 690
 and Fast Green, 689
 and Light Green, 689
 and Methyl Green, 691
 Sage, 594, **595**
 Sage, corolla of, 272
 Sago starch, 456
 Salep, 94, **496**
Salicaceæ, 499
 Salicales, 499
 Salicin, 83
 micro-chemic test for, 83
 Salicinum, 500
Salicornia ambigua, 619
 Salix, 500
Salix alba, 500
 nigra, 500
 retusa, 499
 viminalis, 501
Salmonella, 373
Salsola, 283
 Salvia, 595
Salvia officinalis, 595
Salvinia, 437
Salviniaceæ, 436
 Samara, 303
 Sambucus, 606, **607**
Sambucus canadensis, 607
 nigra, 607
 Samphire, 619
 Sandalwood Family, 510
 Sandaraca, 466
 Sandbox, 551
 Sanguinaria, 528
 latex of, 101
Sanguinaria canadensis, **526, 528**
Sansevieria cylindrica, leaf of, 250
Santalaceæ, 510
 Santalales, 510
Santalum album, 510
Santalum Rubrum, 539
 Santonica, 615
 Santonium, 615
 Sap, cell, 18
 crude, 117
 elaborated, 117
 nuclear, 18
 vacuoles, 19
Sapindaceæ, 553
 Sapindales, 551
 Sapindus, 553
Saponaria, 84
 Saponins, 84
 micro-chemic test for, 84
 Sapotaceæ Family, 581
 Sappan, 540
 Saprogens, 364
 Saprophyte, **362, 383**
 facultative, 367
 obligate, 367
 Sap-wood, 213
Sarcina, 368, **371**
 Sarcode, 17
 Sarcocarp, 298
Sargassum, 355
Sarracenia, 289, **629, 638**
 Drummondii, 638
 flava, Fig. of, **632, 638**
 purpurea, **629, 630, 631, 638**
 rubra, 638
 Sledgei, 638
 style of, 289
Sarraceniaceæ, 530
 Sarraceniales, 530
 Sarsaparilla, 490
 Bristly, 575
 tendrils of, 189
 Virginian, 575
 Wild, 575
 Sassafras, 523
 Medulla, 524

- Sassafras variifolium*, 523, 524
Satureia hortensis, 596
Saxifragaceæ, 531
 Saxifrage, 531
 Family, 531
 Scale bark, 130
 Scale leaves, 228
 Scales, 127
Scammoniae Radix, 591
 Scape, 189, 256
 Schimper, 619
Schizomycetes, 362, **363**
Schizophyta, 333
 Schleiden, M. J., 17
 Schulze's process of maceration, 702
 Schwann, 17
 Schweitzer's reagent, 686
 Scilla, 490
Scilla maritima, 490
Scitaminales, 492
Scitamineæ, 91
Sclerenchyma, 118
 fibers, 120
Sclerotium, 395
Scoparius, 539, **540**
Scopola, 598
Scopola Carniolica, 598
 Scorpioid cyme, 260
 Scouring Rushes, 437
Scrophularia, 275, **599**, **600**
Scrophulariaceæ, 275, **599**
 Scullcap, 594, 595
Scutellaria, 595
Scutellaria lateriflora, 595
 Scutellum, **318**, 485
 Sea Grape, 512
 Sea Lettuce, 342
 Sea Otter's Cabbage, 354
 Sea Palms, 354
 Sea Purslane, 514
Secale cereale, 395
 Secretion canals, 148, 149
 cells, 147
 reservoirs, 148, **149**
 sacs, 147
 Secretory epithelium, 149
 Sectioning, 675
 free hand, 675
 in celloidin, 700
 Sectioning, in paraffine, 677, **697**
 in pith, 675
 Sections, celloidin, 700, 701
 making of, 675
 paraffine, 697
 radial-longitudinal, 677
 surface, 677
 tangential-longitudinal, 677
 transverse, 677
 Sedges, 304, **483**, **484**
Sedum, leaf of, 250
 Seed, 311
 albuminous, 314
 appendages, 313
 Castor-oil, 312
 coats, 312
 defined, 311
 Dicotyl, 319
 exalbuminous, 314
 functions of, 8, 311
 germination, 318, 322, 453
 histology, 315
 Indian Corn, 315
 Monocotyl, 315
 structure, 312
 vessel, 297
 vitality of, 311
 Seed-grass, 618
 Seedling, 319
 of Indian corn, **316**, 319
 of lima bean, 319, **320**
 of pea, 319
Selaginella, 442
 Martensii, Fig. of, 441
Selaginellaceæ, 442
Selenicereus, 567
 grandiflorus, 568
Semen Rapæ, 530
Sempervivum, 96
Senecio, 615
Senecio aureus, 615
Senega, 550
Senna, 251, **539**
 neighboring cells of, 254
 stomatal apparatus of, 251
 Sensitive Fern, 433
 Sepaline position, 269
 spurs, 268
 stipules, 269

- Sepals, **266**
 epigynous, 269
 hypogynous, 269
 perigynous, 269
Sequoia, 465
 height of, 186
Serenoa serrulata, 487
 Serpentaria, 512
 Serpyllum, 596
 Service Berry, 535
 Sesame Family, 601
Sesamum indicum, 602
Sesuvium maritimum, 514
 Shad Bush, 535
 Shell bark, 130
 Shepherd's Purse, 302
 Shoots, 180, 247
 fertile, 437
 spur, 446
 sterile, 437
 Shrub, **191**, 582
Sideritis, 596
 Sieve, 133
 callus of plates, 109
 plates, 133
 Sieve tubes, 133, 134
 stages in development of, 134
 Silica, 91
 bodies, 91
 Siliceous Earths, 350
 Silicule, 302
 Siliqua, 302
Silphium, 615
 Simaruba, 547
Simaruba amara, 547
 officinalis, 547
Simarubaceæ, 546
Sinapis alba, 530
Sinapis Nigra U.S.P., 530
 Sinigrin, 103
 Siphon Algæ, 346
 Siphonales, 346, 376
 Siphonostele, **145**, 199
 Sitotaxy, **34**
 Sitotropism, **34**
 Skunk Cabbage, 487
 Slime Molds, 362, **374**
 Fig. of, 353
 Response to water, **34**
 Slime plugs, 134
 Slippery Elm, leaf of, **221**, 241
Smilax, 490
 histology of root, 165, 166
 medica, 490
 modified stipules of, 246
 officinalis, 489, **490**
 ornata, 490
 Smuts, 397
 Snapdragon, 272, **600**
 stamens of, 275
 Snowberry, 606
 Soap Bark, 84
 Soapwort Family, 553
 Soft bast, 202
Soja max, 540
Solanaceæ, 596
 development of stomata in, 254
 placentation in, 288
Solaneæ, anther dehiscence in, 279
 Solanine, 87
 Solanum, 598
Solanum, 596
Solanum carolinense, 87, **598**
 Dulcamara, 87, 596, **598**, **599**
 nigrum, 87
 tuberosum, 87, 203
Solidago odora, 615
 Solomon's Seal, 192
 Sorbinose, 74
Sorbus, 535
 Soredia, 416
 Sorghum, 484
 Sorosis, defined, 307
 Sorus, 433
 Sour Gum, 575, **577**
 Soy Bean, 540
 Spadix, 259
 Sparganium, 618
 Spathe, 259
 Spearmint, 595
 Species, defined, 326
 Species Plantarum, 325
 Specific epithet, 330
 Specific name, 329
 Spermacia, 401
 Spermagonia, 401
 Spermatium, 358
 Spermatocyte, 68

- Spermatophyta, 14, **443**
 Spermatophytes, 14, 15, **443**
 Spermatozoid of fern, 68
 Spermoderm, 312
 Sperms, 16
Sphagnaceæ, 425
 Sphagnales, 425
Sphagnum, 425
 acutifolium, 424
 palustre, 425
 squarrosum, 424
Sphaerella, 338
 Spicebush, 523
 Family, 522
 Spiderwort, 36, 40
 Spigelia, 583, **585**
Spigelia marilandica, 584, 585
 Spike, 259
 Spikelet, diagram of typical, 484
 Spikelets, **259**, 484
 Spikenard, 575
 Spinach, 514
 Spine, defined, 189
Spiræa, 535
 Spireme, 47
 Spirilla, types of, 371
Spirillaceæ, 372
Spirillum, 365, 372
Spirochæta, 348
Spirogyra, 343, 344
Spirosoma, 372
Spirulina, 335
Spondiæ, 552
 Spongy parenchyma, 117, **223**
 Sporangiphores, 377
 Sporangium, 67, 353, 377, 379
 Spore, 68
 bacterial, 368, **369**
 brand, 398
 daughter-cells, 280
 germination of, 346
 mother-cells, 67, 280, 447, 457
 print, 405
 swarm, 375
 Sporogonium, 428
 Sporophore, 402, **404**, 406
 Sporophyll, 433
 Sporophyte, **53**, **418**, 431
 Sporulation, 368
 Spring Beauty, 514
 Spruce, 211, **465**
 Spruce Gum, 466
 Spur shoot, 446
 Spurge Family, 550
 Spurs, sepaline, 268
 Squash, 188
 Squill, 192, 490
 St. John's Wort, 277
 St. John's Wort Family, 564
 Staff Tree Family, 552
 Stain,
 Delafield's hæmatoxylin, 701
 Fast Green, 682
 Gentian violet, 683
 Heidenhain's hæmatoxylin, 683
 Ruthenium Red and Lead Acetate,
 686
 Safranin, 686
 Staining, 688
 double, 688, 689, 690, 691
 of celloidin sections, 701
 of material in paraffin ribbons, 699
 Stains, 680
 Stamen System, 274
 Figs. of, 276
 Stamens, color of, 277
 connation of, 275
 definite, 274
 diadelphous, 275
 didynamous, 275
 epigynous, 275
 gynandrous, 275
 histology of parts, 277
 hypogynous, 275
 indefinite, 274
 insertion of, 275
 monadelphous, 275
 monandrous, 274
 number of, 274
 perigynous, 275
 polyadelphous, 277
 proportions of, 275
 tetradynamous, 275
 triadelphous, 277
 Staminode, 275
 Staphisagria, 301, **519**
Staphylococcus, 368, **371**
 aureus, 373

- Star Anise, 516
 Star Apple Family, 581
 Starch, **75**
 assimilation, **76**
 Barley, 78, **81**
 Bean, 80, **82**
 Buckwheat, 81
 Canna, 82
 Cassava, 80, **81**
 characteristics of commercial kinds,
 79-82
 compound grains, 77
 Corn, **79**, 80
 corrosion, 102
 fill, 77
 formation of, 64
 Maranta, 78, **79**
 Pea, 80, **82**
 Potato, 78, **79**
 reserve, **76**, 77
 Rice, **79**, 80
 Rye, 78, **79**
 Sago, 78, **82**, 456
 simple grains, 77
 structure and composition of, 77
 transitory, 76
 Wheat, 78, **79**
 Steckbeck, 36
 Stegmata, 91
 Stele, 144
 Stem, defined, 186
 direction of growth, 186
 distinction from root, 160
 elongation, 187
 functions, 6, **186**
 generative tissues of, 187
 modification of, 188
 Sunflower, 194
Stemonitis fusca, 375
 Stems, above-ground, 188
 aerial, 188
 aerial-tuberous, 188
 alate, 187
 annual, 187
 ascending, 186
 cactoid, 188
 climbing, 188
 compressed, 187
 cylindraceous, 187
 Stems, decumbent, 187
 duration of, 187
 elongation of, 187
 endogenous, **193**, 194
 erect, 186
 exceptional types of dicotyl, 203
 exfoliating, 187
 exogenous, 193
 flexuous, 188
 forms of, 187
 fruticose, 190
 herbaceous, 130, **189**
 histology of aconite, 130
 of annual dicotyl, 194
 of herbaceous dicotyledon, 9,
 130, 203
 of herbaceous monocotyl, 215
 of perennial dicotyl, 199
 of woody monocotyl, 217
 obconical, 188
 perennial, 187
 phylloid, 188
 procumbent, 186
 quadrangular, 187
 reclining, 186
 repent, 187
 scandent, 188
 shrubby, 190
 size of, 186
 spiny, 188
 subcylindrical, 187
 subterranean, 192
 subterranean tuberous, 188
 suffruticose, 189
 tendriform, 188
 terete, 187
 tortuous, 188
 triquetrous, 187
 truncate, 188
 twining, 188
 underground, 192
Sterculiaceæ, 557
 Sterigmata, 390, 397
Sterigmatocystis, 391
 Stevens, 104
Sticta, 414
 Stigma, **290**, 473
 Stillingia, 551
Stillingia sylvatica, 551

- Stimuli, extrinsic, **32**
 intrinsic, **32**
 Stink horn fungi, 411
 Stipe, 55, 404
 Stipule, defined, 223
 Stipules, 223, **245**
 axillary, 245
 caducous scaly, 245
 connate, 245, 246
 free axillary, 245
 free lateral, 245
 lateral, 245
 lateral-adnate, 245
 lateral-connate, 245
 lateral-interpetiolar, 245
 modified, 246
 sepaline, 269
 Stolon, defined, 189
 Stoma, 63, 251
 Stoma Mother-cell, 253
 Stomata, 123, 222
 structure and development of, 251
 transpiration, 124
 water, 124
 Stomatal apparatus, 251
 Fig. of types of, 254
 Stomatal development, 251
 Stone cells, 119
 Stone Root, 596
 Stoneworts, 346
 Storax, 532
 Stramonium, 50, 163, **597**
 Strawberry, 300, **534**, 535
 Fig. of flower, 268
 Fig. of fruit, 299
 Fig. of plant showing propagation,
 86
 stem of, 186
 tree, 579
Streptococcus, 386, **371**
 erysipelatis, 373
 scarlatinae, 373
Streptothrix, 372
 Strobile, **259**, **307**
 Strobili, 457, 464
 Stromata, 395
 Strophanthin, **83**, **589**
 micro-chemic test for, 83
 Strophanthus, 313, **589**
Strophanthus hispidus, 83, **588**, **589**
 Kombé, 83, **589**
 Strophiole, 313
 Strychnina, 585
 Strychnine, 85
 micro-chemic test for, 85
Strychnos Ignatii, 85, **585**
 Nux Vomica, 85, 204, 583, **585**
 Fig. of, 584
 toxifera, 585
 Style, **288**, **473**
 arms, **288**, 289
 collecting hairs of, 289
 column, 289
 Styloids, 89, **90**
Styracæa, 582
 Styrax, 532
Styrax Benzoin, 583
 tonkinensis, 583
 Subdivision, 327
 Suberin, 24, 107
 Succinum, 466
 Succus Pomorum, 536
 Suckers, 185, **193**
 Sucrose, **75**
 Sucrosum, **485**, **514**
 Sugar cane, 75
 Sugars, 74
 Sulfuric acid, as reagent, 687
 Sumac Family, 551
 Sumbul, 576
 Summer Cypress, 514
 Summer Savory, 596
 Sundew, 530
 Sundews, 260, 530, **622**
 Sunflower, 614
 Figs. of cross sections of stem, 194,
 195
 histology of stem of, 194
Surirella, 352
 Suspensor, 439, 452
 Sutures, 298
 Swamproot, 615
 Swarm spores, 375
 Sweet Basil, 596
 Sweet Birch, 504, 505
 Sweet Cicely, 576
 Sweet Flag, 618
 Sweet Marjoram, 596

- Sweet Potato, 162, 590
 Sweet Woodruff, 605
Sweetia panamensis, 539
Swertia Chirayita, 587
 Syconium, 307
 Fig. of, 290
 Sympetalae, 579
 Symphytum, 593
Symphytum officinale, 593
Symplocarpus fœtidus, 487
 Syncarp, 471
Synedra ulna, Fig. of, 352
 Synergids, 292, 474
 Syngensis, 265
 Systema Naturæ, 325
 Systematic Botany, 323
 defined, 2

 Tabacum, 598
Tabellaria vulgaris, Fig. of, 352
 Tamarack bark, 466
 Tamarind, 301
 Tamarindus, 539, **540**
Tamarindus indica, 540
 Tanacetum, 615
Tanacetum vulgare, 615
 Tannins, 92
 Tansy, 615
 Tapetum, 67, **278**, 280
 Tapioca, 551
Taraktogenos Kurzii, 565
 Taraxacum, 615
Taxaceæ, 465, **466**
 Taxies, 32
Taxodium, 619
 Taxonomy, 323
 defined, 2
Taxus baccata, 467
 canadensis, 467
 Tea, 85, **563**
 Tea Family, 563
 Teak-wood, 593
 Teasel Family, 608
Tecoma radicans, Fig. of seed, 313
Tectona, 593
 Tegmen, 312
 Tela contexta, 407
 Teleutospores, 401
 Teliospores, 400, 401

 Telium, 401
 Telophase, 50
 Temporary mount, technique of making a, 687
 Tendril, defined, 189
 Tendrils, 189
 Terebinthina, 456
 Canadensis, 466
 Laricis, 466
Ternstræmiaceæ, 563
 Terpenes, 97
 Terra Silicea Purificata, 350
 Testa, 312
 Tetrads, 67, 280, 474
 Tetrasporangia, 358
 Tetraspores, **358**, 359
 Thalamus, 262, **265**
 Thallophyta, 12, **333**
 Thallophytes, 12, 333
 Thallus, defined, 12
Thamnidium, 381
 Thea, 563
Thea sinensis, 563
Theaceæ, 563
 Thelephorales, 403
Theobroma, 559
Theobroma Cacao, 559
 Fig. of, 558
 Theophrastus, 1, 323
 Thermogens, 364
 Thermotropism, **32**
 Thigmotropism, 36
Thiothrix, 373
 Thistle, 610, 611
 Thorn, defined, 189
 Throat, 266
Thuja, 465, **466**
 occidentalis, 466
 Thyme, 189, **595**
Thymelæaceæ, 569
Thymus Serpyllum, 596
 vulgaris, 595
 Thyrsus, 261
Tilia, 559
 americana, 559
Tiliaceæ, 559
 Tissue, 111
 assimilation parenchyma, 116
 cambium, 114

- Tissue, chlorenchyma, 116
 collenchyma, 113, 117
 conducting, 151
 conducting parenchyma, 117
 cork, 129
 cribriform, 133
 embryonic, 114
 endodermal, 127
 epidermal, 122
 fundamental, 116
 generative, 113
 ground, 116
 laticiferous, 131
 mechanical, 151
 medullary ray, 139
 meristematic, 113, **114**, 151
 ordinary parenchyma, 113, **116**
 parenchyma, 116
 pericycle, 131
 protective, 151
 reserve parenchyma, 116
 sclerenchyma, 113, **118**
 secreting, 151
 sieve, 133
 stony, 118
 storage, 151
 suberous, 129, 130
 tracheary, 135
- Tissues, 8, **111**
 adult, 113
 classifications of, 111, 151
 complex, 113
 primary, 111, 115
 secondary, 112
 simple, 113
- Tobacco, 596, 597
- Tolu, 539
- Toluiifera Balsamum*, 539
Pereira, 539
- Tomato, 304, 596
- Tonga, **487**, **593**
- Tooth Fungi, 404
- Tormentilla, 536
- Torulaceæ*, 381
- Torus, 262, **265**
- Touranose, 75
- Tournefort, 325
- Toxins, **33**
- Tracheæ, **135**
- Tracheary tissue, 135
- Tracheids, 135, **136**, 139
 Fiber, 214
- Tradescantia*, 82, 253
- Tragacanth, 95
- Tragacantha, 539
- Trailing arbutus, 272, **581**
- Trama, 407
- Transpiration, 65, **226**
 Stomata, 124
 Stream, 226
- Trapa natans*, 569
- Trapaceæ*, 569
- Tree, defined, 190
- Trees, branching in, 191
- Trehalase, 102
- Trehalose, **75**, 102
- Tremellaceæ*, 402
- Tremellales, 402
- Treponema*, 373
- Treviranus, 17
- Tribes, 329
- Trichogyne, 358
- Trichomes, 124
- Trichophyton*, 412
- Trifolium, 540
Trifolium pratense, 540
repens, 536
- Trigonella Fœnum-græcum*, 540
- Trilisa odoratissima*, 615
- Trillium, 490
Trillium erectum, 490
- Trimorphism, 289
- Triticum, 485
Triticum sativum, starch of, 79
- Tropæolum*, 267
- Tropisms, **32**
- Tropophytes, 617, **619**
- True Fungi, 375
- True Mosses, 426
- Truffles, 394
- Trumpet Creeper, **291**, **313**
- Trunk, defined, 190
- Trypsin, 103
- Tschirch, 96
- Tsuga*, 465
Tsuga canadensis, 466
- Tuber, defined, 192
Tuber, genus, 394

- Tuberales, 394
 Tubercles, 176, 179
 Tubifloræ, 590
Tubulifloræ, 613
 Tulip, 260, 266
 Tulip Poplar, 516
 Tung Oil, 551
 Turgor, 155
Turnera aphrodisiaca, 566
 diffusa, 566
Turneraceæ, 566
 Turnip, 162, 240
 Turpentine, 465
 Bordeaux, 466
 Turpeth Root, 591
Tussilago Farfara, **612, 615**
 Twig, 190
 Twin crystals, 90
 Tyloses, 213
 Typha, 282

Ulmaceæ, 507
 Ulmus, 507
Ulmus fulva, 221, 223, **506, 507**
Ulothrix zonata, 341
 Fig. of, 338
Ulva lactuca, 342
 Umbel, 258
 compound, 258
 Umbellales, 575
Umbelliferæ, 243, 266, 303, **576**
Umbellifloræ, 575
 Unit characters, 643, 644
 Uredinales, 399
 Uredinium, 400
Uredo linearis, 401
 Uredospore, 400
Urginea maritima, 490
Urtica urens, stinging hair of, 126
 Urticales, 507
Usnea, 414, 417
 Ustilaginales, 397
Ustilago Maydis, 397
Ustilago Zeæ, 397, 398
 Utricle, defined, 304
Utricularia, 248, **638, 639**
Utriculariaceæ, 640
 Uva, defined, 304

 Uva Ursi, 581
 corolla of, 273

Vaccineæ, 581
 Vaccinium, 581
Vaccinium corymbosum, 581
 Vacuole, contractile, 27
 food, 26
 Vacuoles, sap, **19, 40**
 Valerian Family, 608
 Valeriana, 609
Valeriana officinalis, 608, **609**
Valerianaceæ, 608
 Valerianales, 608
Vallisneria, 291
 Valves of fruits, 298
 Vanilla, 496
Vanilla planifolia, 496
 Figs. of plant, 495
 histology of fruit, 307
 roots of, 161
 Vanilla Leaf, 615
 Variations, 652
 Varieties, 327
 naming of, 331
 Variety epithet, 331
 Vascular bundles, 140
 Vascular rays, 199
Vaucheria, 95, **346**
Vaucheria sessilis, Fig. of, 345
Vaucheria terrestris, Fig. of, 341
 Vegetable Kingdom, 1
 Vegetative Multiplication, **46**
 Veil, partial, 406
 universal, 408
 Veins, **222, 230**
 Velamen, 161
 Venation, leaf, 230
 Venter, **69, 70**
 Venus fly-trap, 36, 625
 Figs. of, 35, 626
 Veratrina, 491
Veratrum viride, 490
 Verbasci Flores, 600
 Verbasci Folia, 600
Verbascum, 275, 599
 Blattaria, 599
 phlomoides, 600

- Verbascum thapsiforme*, 600
Thapsus, 599, 600
 Verbena, 593
Verbena hastata, 593
Verbenaceæ, 593
 Vernation, 230
Veronica, 275, 599, 600
 virginica, 600
 Verticillaster, 261
 Vervain Family, 593
 Vessels, 135
 laticiferous, 132
 Vexillum, 271
Vibrio, 372, 373
Viburnum, 202, 263, 606
Viburnum Opulus, 150, 606, 607
 Prunifolium, 606, 607
Vinca, style of, 289
Viola, 289, 566
Viola odorata, 566
Viola tricolor, 124, 126, 566
Violaceæ, 565
 Violet, 189, 566
 Violet Family, 565
Viscum, 479, 510
Viscum album, 510
Vitaceæ, 556
 Vitamins, 105
Vitis, 556
Vitis Labrusca, 557
 Vittæ, 148
 Volatile oils, 96
 Volva, 409
 Volvocales, 337
Volvox, 29
Volvox globator, life history of, 339
 Fig. of, 340
 Von Mohl, H., 17
Vouacapoua Araroba, 540

 Wafer Ash, 303, 545, 546
 Wahoo, 553
Waldsteinia, 535
 Walnut, Black, 504
 Walnut Family, 502
 Water Chestnut, 569, 617
 Water Ferns, 436
 Water Hyacinth, 617
 Water Leaf Family, 591
 Water Lily, 248, 617
 Family, 515
 Water Molds, 376
 Water Net. See *Hydrodictyon*.
 Water of organization, 25
 Water Shamrock, 437
 Water stomata, 124
 Watermelon, 607
 seed, 608
 Waxes, 96
Weigelia, 606
 Weissmann, A., 656
 Weissmann's Theory, 656
Welwitschia, 469
 Weymouth Pine, 444
 Wheat, 259, 304
 Fig. of fruit, 304
 Fig. of plant, 482
 rust, 399
 starch, 79
 White Birch, Fig. of branch showing
 lenticels, 204
 White Pine, life history of, 444
 outline of life cycle, 454
 Wild Black Cherry, 103, 534
 Wild Cherry, 258
 Wild Columbine, 269
 Wild Jalap, 591
 Willow, 259, 500
 Willow Family, 499
 Willow Herb, 569
 Windflower, 260
 Winter Cherry, 267
 Wintergreen, 297, 579
 Winter's Bark, 516
 Witchhazel Family, 531
 Wood, Angiospermous, 214
 Coniferous, 214
 defined, 212
 fibers, 122
 heart-, 213
 microscopic characters of coniferous
 and angiospermous, 214
 parenchyma, 117
 pine, 212
 ray, 139
 sap-, 213

- Wood-rays, 170
 Wood-wedge, 170
 Wormwood, 615

 Xanthophyll, 99, **100**
Xanthorhiza, 519
 Xanthoxyli Fructus, 546
 Xanthoxylum, 546
 Xerophytes, 164, 617, **619**
 Xylem, 140
 defined, 142
 primary, 143
 ray, 139
 secondary, 143
 stages in development of elements
 of, 138

 Yam Family, 491
 Yarrow, 615
 Yaupon, 25
 Yeast, Brewer's, 382
 Compressed, 386
 vitamins in, 105, 106, **386**
 Wine, 386
 Yeasts, bottom, 383, **386**
 classification of, 382
 defined, 381
 top, 384, **386**
 Yellow Jasmine, 583
 Yellow Parilla, 521
 Yerba Santa, 592
 Yew, 467
 Fig. of tracheid of, 23
 Yew Family, 465, **466**
 Yohimbe, 605
Yucca, 220, 259

Zamia, 455
 life history of, 456
 outline of life cycle, 462
Zamia floridana, Fig. of, 459
Zanthorhiza apiifolia, 519
Zanthoxyleæ, 545
Zanthoxylum, 545
 Americanum, 546
Zanthoxylum Clava Herculis, 546
 Zea N. F., 485
Zea Mays, 485
 Fig. of, 483
 starch of, **79**, 80
 structure of stem of, 215, 216
 Zedoaria, 492
 Zingiber, 492
Zingiber officinale, 492
 Fig. of, 493
Zingiberaceæ, 492
 Zizyphus, 556
Zizyphus sativa, 556
 Zöoglœa, 178
 Zöology, defined, 1
 Zoöspore, **45**, 342, **345**, 353
 Figs. of, 353
 Zoöspore formation, 45
Zostera, 291
 Zygnema, 343
 Fig. of, 338
Zygnemaceæ, 343
 Zygomycetales, 376, **377**
 Zygomycetæ, 333
 Zygomycetaceæ, 544
 Zygosporangium, **46**, 342, 345, 353, 379
 Zygote, 16, **46**, 345, 348, 380, **461**
 Zymase, 102, **103**
 Zymogens, 364