SF
994
W33
[v.2]
Birds

F-2

W33
[v.2]
Birds

FINAL REPORT

A SEROLOGICAL AND ECTOPARASITE SURVEY OF MIGRATORY BIRDS IN NORTHEAST AFRICA

Paleartic Migratory Bird Survey

Appendix I

SF
994
W53
[17.2]
Birds

Watson, George Elder

FINAL REPORT

A SEROLOGICAL AND ECTOPARASITE SURVEY OF MIGRATORY BIRDS IN NORTHEAST AFRICA, Palearctic Migratory Bird Survey

Appendix I

Included in this Appendix are distribution maps of the major species of birds, ticks and virus strains studied during the Palearctic Migratory Bird Survey (PMS), as well as lists of banded birds recovered in the Near East. The bird distribution maps (pages 1-137) show Egyptian (star), Cyprus (black circle), Israel (square), and Lebanon (open circle) recoveries. The recovery lists, pages 157-220 should be consulted for details. The tick maps (pages 138-148) were prepared by Dr. Kaiser on the basis of preliminary data and although subject to later revision are the best approximations of distribution currently available. The virus maps (pages 149-156) show only the locations of primary isolations of the various strains. Pages 157-199 present banding details of Near Eastern (Egypt, Cyprus, Lebanon and Israel) birds both those banded in the area, and recovered elsewhere, and foreign bands recovered in the Near East. The recoveries are the result of PMS activities (pages 157-192). Additional Cyprus recoveries (pages 200-220) were previously reported in the *Bulletin of the Cyprus Ornithological Society*. Most of these recoveries are shown on the bird maps. Recovery information on birds banded at Azraq by the British Trust for Ornithology and International Jordan Expedition was provided by Mr. I. J. Ferguson-Lees (pages 221-222). These recoveries are not included on the maps. We have not yet received banding information for a number of recoveries of foreign bands found in Egypt (pages 223-232). Many of these are a number of years old and, in particular, we have had some difficulty getting banding information from the U.S.S.R.

TABLE OF CONTENTS

I. Distribution Maps

A. Birds (in taxonomic sequence with computer codes for

Appendix II)

1 - 137

Species	Code	Page
<i>Podiceps nigricollis</i>	PO NI	1
<i>Ixobrychus minutus</i>	IX MI	2
<i>Ardeola ralloides</i>	AD RA	3
<i>Bubulcus ibis</i>	BL IB	4
<i>Ardea purpurea</i>	AD PU	
<i>Egretta garzetta</i>		
<i>Egretta alba</i>		
<i>Nycticorax nycticorax</i>		
<i>Ciconia ciconia</i>		5
<i>Platalea leucorodia</i>		
<i>Plegadis falcinellus</i>		
<i>Anas crecca</i>	AS CR	
" <i>penelope</i>	AS PE	
" <i>acuta</i>	AS AC	
<i>Nyroca ferina</i>	NY FE	
<i>Milvus migrans</i>	MI MI	
<i>Accipiter brevipes</i>	AP BR	6
" <i>nisus</i>	AP NI	7
" <i>brevipes/nisus</i>	AP BN	
<i>Buteo buteo</i>	BU BU	8
<i>Circaetus gallicus</i>	CQ GA	
<i>Circus macrourus</i>	CC MA	
<i>Pandion haliaetus</i>		
<i>Falco eleonora</i>		
" <i>naumanni</i>	FA NA	9
" <i>subbuteo</i>	FA SU	10
" <i>tinnunculus</i>	FA TI	11
" <i>verpertinus</i>	FA VE	12
<i>Coturnix coturnix</i>	CO CO	13
<i>Rallus aquaticus</i>	RA AQ	14
<i>Porzana pusilla</i>	PZ PU	
" <i>porzana</i>	PZ PO	15
<i>Crex crex</i>	CX CX	16
<i>Gallinula chloropus</i>	GL CH	17
<i>Porphyrio porphyrio</i>		
" <i>madagascariensis</i>	PO MA	
<i>Fulica atra</i>	FU AT	18

Species	Code	Page
<i>Charadrius hiaticula</i>	CH HI	19
" <i>dubius</i>	CH DU	20
" <i>alexandrinus</i>	CH AL	21
<i>Vanellus vanellus</i>	VA VA	
" <i>spinosus</i>	VA SP	22
<i>Arenaria interpres</i>	AR IN	23
<i>Calidris minuta</i>	CA MI	24
" <i>temminckii</i>	CA TE	
" <i>alpina</i>	CA AP	25
" <i>ferruginea</i>	CA FE	
" <i>canuta</i>	CA CA	
" <i>alba</i>	CA AL	
<i>Philomachus pugnax</i>	PL PU	26
<i>Tringa erythropus</i>	TR ER	27
" <i>totanus</i>	TR TO	28
" <i>nebularia</i>	TR NE	29
" <i>ochropus</i>	TR OC	30
" <i>glareola</i>	TR GL	31
" <i>hypoleucos</i>	TR HY	32
<i>Scolopax rusticola</i>	SC RU	
<i>Capella gallinago</i>	CP GA, GA GA	33
<i>Lymnocyptes minima</i>	LY MI	34
<i>Burhinus oedicephalus</i>	BR OE	
" <i>senegalensis</i>	BR SE	
<i>Cursorius cursor</i>	CZ CZ	
<i>Stercorarius parasiticus</i>		
<i>Larus fuscus</i>	LR FU	35
" <i>genei</i>		36
" <i>ridibundus</i>		
<i>Chlidonias niger</i>	CS NI	37
" <i>leucopterus</i>	CS LE	38
" <i>hybrida</i>	CS HY	39
<i>Hydroprogne caspia</i>	HY TS	40
<i>Streptopelia senegalensis</i>	ST SE	
" <i>turtur</i>	ST TU	41
<i>Cuculus canorus</i>	CU CA	42
<i>Clamator glandarius</i>	CL GL	
<i>Centropus senegalensis</i>	CN SE	
<i>Tyto alba</i>	TY AL	
<i>Otus scops</i>	OT SC	43
<i>Athene noctua</i>	AH NO	
<i>Caprimulgus nubicus</i>	CM NU	
" <i>europaeus</i>	CM EU	44
<i>Apus pallidus</i>	AP PA	
" <i>apus</i>	AP AP	
<i>Ceryle rudis</i>	CY RU	
<i>Halcyon smyrnensis</i>	HA SM	
<i>Alcedo atthis</i>	AL AT	45
<i>Merops orientalis</i>	ME OR	46
" <i>apiaster</i>	ME AP	47
" <i>superciliosus</i>	ME SU	

Species	Code	Page
<i>Coracias garrulus</i>	CR GA	48
<i>Upupa epops</i>	UP EP	49
<i>Jynx torquilla</i>	JY TO, JU TO	50
<i>Ammomanes cincturus</i>	AM CI	
<i>Ammomanes deserti</i>	AM DE	
<i>Alamon alaudipes</i>	AL AL	
<i>Calandrella cinerea</i>	CL CI, CL BR	51
<i>Melanocorypha calandra</i>	ML CA	52
<i>Galerida cristata</i>	GA CR	
<i>Lullula arborea</i>	LL AR	53
<i>Alauda arvensis</i>	AU AR	
<i>Riparia riparia</i>	RI RI	54
<i>Hirundo rupestris</i>	HR RU	55
" <i>rustica</i>	HR RS	56
" <i>daurica</i>	HR DA	57
<i>Delichon urbica</i>	DE UR	58
<i>Anthus campestris</i>	AN CA	59
" <i>trivialis</i>	AN TR	60
" <i>pratensis</i>	AN PR	61
" <i>cervinus</i>	AN CE	62
" <i>spinoletta</i>	AN SP	63
<i>Motacilla flava</i>	MO FL	64
" <i>flava feldegg</i>	MO FF	
" <i>flava flavissima</i>	MO FV	
" <i>flava cinereocapilla</i>	MO FC	
" <i>cinerea</i>	MO CI	65
" <i>alba</i>	MO AL	66
<i>Pycnonotus barbatus</i>	PY BA	
<i>Lanius collurio</i>	LA CR, LA CO	67
" <i>nubicus</i>	LA NU	68
" <i>senator</i>	LA SE	69
" <i>minor</i>	LA MI	70
" <i>excubitor</i>	LA EX	71
<i>Prunella modularis</i>	PU MO	72
<i>Cettia cetti</i>	CT CE	73
<i>Locustella luscinioides</i>	LO LU	74
" <i>fluviatilis</i>	LO FL	75
<i>Acrocephalus melanopogon</i>	AC ME	76
" <i>schoenobaenus</i>	AC SH	77
" <i>palustris</i>	AC PA	78
" <i>scirpaceus</i>	AC SC	79
" <i>scirpaceus/palustris</i>	AC SP	
" <i>arundinaceus</i>	AC AR	80
" <i>stentoreus</i>	AC ST	81
<i>Hippolais icterina</i>	HI IC	82
" <i>olivetorum</i>	HI OL	83
" <i>pallida</i>	HI PA	84
<i>Sylvia nisoria</i>	SY NI	85
" <i>hortensis</i>	SY HO	86

Species	Code	Page
<i>Sylvia borin</i>	SY BO	87
" <i>atricapilla</i>	SY AT	88
" <i>communis</i>	SY CO	89
" <i>curruca</i>	SY CU	90
" <i>nana</i>		91
" <i>rueppelli</i>	SY RU	92
" <i>melanocephala</i>	SY ME	93
" <i>melanothorax</i>	SY MX	
" <i>cantillans</i>	SY CA	94
" <i>conspicillata</i>		95
<i>Phylloscopus trochilus</i>	PH TR	96
" <i>collybita</i>	PH CO	97
" <i>trochilus/collybita</i>	PH TC	
" <i>bonelli</i>	PH BO	98
" <i>sibilatrix</i>	PH SI	99
" <i>inornatus</i>	PH IN	
<i>Cisticola juncidis</i>	CG JU	100
<i>Prinia gracilis</i>	PR GR	
<i>Ficedula hypoleuca</i>	FI HY	101
" <i>albicollis</i>	FI AL	102
" <i>semitorquata</i>	FI SE	103
" <i>albicollis/hypoleuca</i>	FI AH	
" <i>parva</i>	FI PA, MU PA	104
<i>Muscicapa striata</i>	MU ST	105
<i>Saxicola rubetra</i>	SA RU	106
" <i>torquata</i>	SA TO	107
<i>Oenanthe oenanthe</i>	OE OE	108
" <i>pleschanka</i>	OE PL	109
" <i>hispanica</i>	OE HI	110
" <i>lugens</i>	OE LU	
" <i>deserti</i>	OE DE	
" <i>finschii</i>		111
" <i>isabellina</i>	OE IS	112
" <i>leucura</i>	OE LC	
<i>Cercomela melanura</i>	CE ME	
<i>Certotrichas galactotes</i>	CC GA	113
<i>Monticola saxatilis</i>	MN SA	114
" <i>solitarius</i>	MN SO	
<i>Phoenicurus ochruros</i>	PN OC	115
" <i>phoenicurus</i>	PN PH	116
<i>Erithacus rubecula</i>	ER RU	117
<i>Luscinia megarhynchos</i>	LU ME	118
" <i>luscinia</i>	LU LU	119
<i>Cyanosylvia svecica</i>	LU SV	
<i>Turdus pilaris</i>	TU PI	
" <i>torquatus</i>	TU TO	120
" <i>merula</i>	TU ME	121
" <i>iliacus</i>	TU IL	122
" <i>philomelos</i>	TU PH	123
" <i>viscivorus</i>	TU VI	
<i>Turdoides squamiceps</i>	TD SQ	
<i>Parus major</i>	PA MA	

Species	Code	Page
<i>Cinnyris osea</i>	CI OS	
<i>Emberiza calandra</i>	EM CL	124
" <i>hortulana</i>	EM HO	125
" <i>caesia</i>	EM CA	126
" <i>melanocephala</i>	EM ME	127
<i>Fringilla coelebs</i>	FR CO	128
" <i>montifringilla</i>		
<i>Serinus serinus</i>	SE SE, SE CA	129
" <i>citrinella</i> (?)	SE CI	
<i>Carduelis chloris</i>	CR CH	130
" <i>spinus</i>	CR SP	131
" <i>carduelis</i>	CR CA	132
<i>Acanthis cannabina</i>	AT CA	133
<i>Carpodacus erythrurus</i>	CD ER	
<i>Coccothraustes coccothraustes</i>	CK CK	
<i>Passer domesticus</i>	PS DM, PS DO	134
" <i>hispaniolensis</i>	PS HI	135
" <i>hispaniolensis/domesticus</i>	PS HD	
" <i>moabiticus</i>	PS MO	136
<i>Sturnus vulgaris</i>	SU VU	
<i>Oriolus oriolus</i>	OR OR	137
<i>Corvus corone</i>	CV CO	

B. Ticks

Species	138-148
<i>Amblyomma lepidum</i>	138
" <i>nuttalli</i>	139
" <i>variegatum</i>	140
<i>Haemaphysalis concinna</i>	141
" <i>parva</i> (<i>otaphila</i>)	142
" <i>punctata</i>	143
<i>Hyalomma m. marginatum</i>	144
" " <i>rufipes</i>	145
<i>Ixodes frontalis</i>	146
" <i>redikorzevi</i> subsp.	147
" <i>ricinus</i>	148

C. Virus

149-156

Strain	
Bahig and EG AN 1047-61	149
EG AN 890-3	150
Ingwavuma	151
Kemerovo	152
Quaranfil	153
Sindbis	154
Unkuniemi	155
West Nile	156

II. Band Recoveries

A. Near Eastern Recoveries
Mostly the Result of PMS Activities	157-192
B. Near Eastern Recoveries
Addendum - Mostly Literature Records of White Storks	193-199
C. Cyprus Recoveries
Previously Reported to Cyprus Ornithological Society	200-220
D. Jordan Recoveries
Birds Banded at Azraq Oasis (31°50' N, 36°47' E),
Jordan by British Trust for Ornithology, International
Jordan Expedition	221-222
E. Band Recoveries
Information Incomplete	223-232

Black-necked Grebe *Podiceps nigricollis*

Only one subspecies with no geographic variation is found in the Palearctic.

Habitat: Fresh, brackish or saline marshes, lagoons, ponds, lakes and streams.

Little Bittern *Ixobrychus minutus*

Monotypic in the Palearctic with no geographic variation.

Habitat: Dense reed beds and mixed willows and reeds along the shores of ponds, lakes, and slow moving streams; also densely wooded swamps.

Squacco Heron

Ardeola ralloides

Palearctic populations winter in tropical Africa. Monotypic, no geographic variation.

Habitat: Reeds and thickets in swamps, lagoons and river banks; wet grasslands with bushes and flooded cultivation.

Cattle Egret

Bubulcus ibis

May also breed in southern France and the Volga delta. Mostly resident and breeding in northern Egypt. After breeding may wander erratically. There is no morphological difference between North African and southwest Asian birds but those from Pakistan east are a different subspecies.

Habitat: Pastures, open grasslands and cultivation near large hooved stock for feeding, but usually nests in marshes and wet thickets near fresh water. In Cairo near Zoo, nests in Plane Trees along city streets.

White Stork

Ciconia ciconia

The western populations (all but isolated one in Turkestan) show no geographic variation but differ in migration routes. Birds breeding east of eastern Germany head south east to the Bosphorus, cross Nile Valley to eastern Africa. West European birds cross the Mediterranean at Gibraltar and spread out in crossing the Sahara on a broad front.

Habitat: Wet meadows or marshes, river valleys and cultivation generally near human settlements.

Levantine Sparrowhawk *Accipiter brevipes*

Monotypic. Closely related to the African and Southeast Asian Shikra, *A. badius*, but overlapping its range in the Middle East.

Habitat: Open forests and plains.

Sparrowhawk *Accipiter nisus*

A moderate west to east cline exists, but only the western subspecies, *nisus*, which breeds east to the Yenisei, winters in northeast Africa. Much of this population is resident but migrants may move south as far as the line in East Africa, southern India, and southeast Asia.

Habitat: Forests and open woodland, both mountains and lowlands

Buzzard

Buteo buteo

European and Siberian subspecies differ but intergrade in eastern Europe. Both, plus intergrades, pass through the eastern Mediterranean to winter in southern and East Africa.

Habitat: Varies from rocky coasts to high plateau in woodlands and open pastureland and steppe. Absent from dense forests.

Lesser Kestrel *Falco naumanni*

Monotypic, no geographic variation.

Habitat: Open country, especially farmland, towns and villages. Breeds colonially in cliffs and buildings.

Hobby

Falco subbuteo

The clinal tendencies of geographic variation are slight and largely obscured by individual variation. Only the extreme eastern population can be recognized on morphological grounds.

Habitat: Open woodland and fields with scattered trees.

Kestrel

Falco tinnunculus

The Eurasian populations, which are partly migratory, winter as far south as the northern Belgian Congo, Tanzania, Arabia, India and Indochina. Banding recoveries in Tunisia show a strong eastward tendency in spring. Presumably birds wintering in or passing through Egypt on migration breed in western Siberia.

Habitat: Open woodland and forested steppes, river valleys, cultivated fields, pastureland, seacoasts and even towns and villages.

Red-footed Falcon *Falco vespertinus*

No variation in northern and western Eurasian population but isolated oriental population is distinct. Both winter in Africa but oriental population does not pass through Mediterranean region.

Habitat: Forested steppes, plains and meadows with scattered groves of trees and bushes; cultivated areas and river valleys. Occasionally colonial.

Quail

Coturnix coturnix

Other races breed in eastern and southern Africa. No variation in birds of continental Europe and western Asia that winter in Africa. Overlaps another species *C. japonica* in northern Mongolia and, on migration, in India.

Habitat: Open grain fields, meadows, pastures and grassy steppes; scrubby areas in semidesert; and cropland in desert oases.

Water Rail

Rallus aquaticus

Variation is slight, and all Continental birds in North Africa, Europe, and Siberia belong to the subspecies that breeds and occurs on passage in Eastern Mediterranean.

Habitat: Dense vegetation in swamps, marshes, and banks of rivers, ponds, and lakes.

Spotted Crane *Porzana porzana*

Monotypic with no geographic variation.

Habitat: Sedge growth in shallow swamps, bogs, very wet meadows, and swampy edges of ponds, lakes and rivers.

Corn Crane

Crex crex

Monotypic but shows slight west to east clinal variation.

Habitat: Usually in lowland meadows, pastures and grain crops with lush growth, occasionally on mountain slopes.

Moorhen *Callinula chloropus*

Monotypic in the Palearctic with no recognizable geographic variation.

Habitat: Dense aquatic vegetation on the banks of fresh water marshes, lakes, and slow moving streams. Generally stays hidden in cover.

Coot *Fulica atra*

Monotypic with no variation in Palearctic.

Habitat: Aquatic vegetation on the banks of fresh water marshes, lakes, and slow-moving streams preferring larger more open stretches than moorhen; on migration on brackish water.

Ringed Plover *Charadrius hiaticula*

Western Palearctic (east of Pechora River) population which migrates to Africa shows no geographic variation.

Habitat: Sandy or pebble beaches, short grass meadows, and tundra along the seacoast and less frequently inland.

Little Ringed Plover *Charadrius dubius*

Monotypic in Palearctic with no variation.

Habitat: Sandy or gravel shores and banks of streams, rivers, lakes; generally frequents fresh water but occasionally found in estuaries.

Kentish Plover *Charadrius alexandrinus*

Only a single subspecies, *alexandrinus*, breeds in western Eurasia and migrates to Africa.

Habitat: Sand, mud or shingle beaches or bars or flats on salt or fresh water.

Spur-winged Plover *Vanellus spinosus*

Resident in Africa, but a migrant to Greece and the Near East. Monotypic with no geographic variation.

Habitat: Open fresh water wetlands and irrigated fields.

Ruddy Turnstone *Arenaria interpres*

Highly migratory; adults outside breeding season and some first-year birds all year, can be found almost anywhere in southern Europe, Africa and southern Asia on shores provided there are pebbles, stones or shells. There is no variation in Palearctic populations.

Habitat: Barren stony or pebble shores and dry stony tundra in breeding season.

Little Stint *Calidris minuta*

Monotypic, no geographic variation.

Habitat: Breeds in wet, dry or forest tundra; sea coasts on migration and in winter.

Dunlin

Calidris alpina

Variation is clinal but it is impossible to differentiate subspecies on migration or in winter quarters.

Habitat: Breeds on wet arctic and mountain tundra, grassy moors, flats and salt marshes. On beaches and flats in winter.

Ruff

Philomachus pugnax

Monotypic with no geographic variation.

Habitat: Wet grassy or forested tundra; shores of ponds, lakes, rivers and marshes in steppes; grassy marshes, ferns, bogs, swampy meadows and coastal marshes.

Spotted Red Shank *Tringa erythropus*

Monotypic, with no geographic variation.

Habitat: Wetlands on forested or bushy tundra and northern edge of taiga.

Greenshank

Tringa nebularia

Monotypic with no geographic variation.

Habitat: Marshes, bogs, swampy clearings or river meadows in the taiga; birch forest; dry ground near marshes, high wet moors and lowland flats.

Green Sandpiper *Tringa ochropus*

Monotypic with no geographic variation.

Habitat: Along quiet brooks, pools, marshes,
and alder swamps in forest or wooded steppes.

Wood Sandpiper

Tringa glareola

Monotypic, no geographic variation.

Habitat: A variety of wetlands including boggy moors, heaths, meadows, and shores of ponds, lakes and rivers in forest, willow zone above timberline and forested tundra.

Common Sandpiper *Tringa hypoleucos*

Monotypic, with no geographic variation.

Habitat: Banks of clear streams, ponds and lakes with sand, gravel, or rocky shores at all altitudes.

Jacksnipe

Lymnocyptes minima

Monotypic, with no geographic variation.

Habitat: Grassy wetlands or low reeds in forested tundra, northern birch or alder woods, and coniferous forest.

Lesser Black-backed Gull *Larus fuscus*

Only the well marked Scandinavian sub-species (*fuscus*) is found on migration and in winter in the eastern Mediterranean.

Habitat: Seacoasts and inland lakes.

Slender-billed Gull *Larus genei*

Monotypic with no geographic variation.

Habitat: Sandy flats on small islands,
on shallow coasts, lakes, lagoons,
and marshes in river estuaries.

Black Tern

Chlidonias niger

Only a single subspecies which shows no geographic variation breeds in the Palearctic and migrates to Africa.

Habitat: Shallow fresh inland waters such as marshes, swamps, lakes and rivers during breeding but occurs at sea in winter.

White-winged Black Tern *Chlidonias leucopterus*

Monotypic with no geographic variation.

Habitat: Shallow fresh inland waters such as marshes, swamps, lakes and rivers.

Whiskered Tern *Chlidonias hybrida*

No variation in western Palearctic subspecies that migrates to Africa.

Habitat: Shallow fresh inland waters such as marshes, swamps, lakes and rivers.

Turtle Dove *Streptopelia turtur*

Variation is clinal in Eurasia and examples of both western (*turtur*) and eastern (*arenicola*) subspecies have been taken in eastern Mediterranean; but complex in birds breeding in Sahara oases and Egypt.

Habitat: Open woodland, cultivation, parkland and gardens with trees, orchards and olive groves, steppeland with trees or bushes, and desert oases.

Cuckoo

Cuculus canorus

Although there is slight clinal variation and several poorly marked subspecies are recognized, migrants cannot be assigned to subspecies on the basis of plumage characters.

Habitat: Open wood or parkland, locally in the far north above the treeline.

Scops Owl

Otus scops

Populations on Iberian peninsula, Mediterranean islands and Near East are distinct and probably resident. Birds breeding on continent from France and North Africa to western Siberia belong to a single race (*scops*) that migrates through the Mediterranean to winter in Africa. East Siberian, Turkestan and Middle East populations probably do not reach Mediterranean on passage while Indian and oriental populations may be a different species.

Habitat: In all sorts of forest in north and east and more open woodland, parks, orchards, and gardens in the south and west.

Nightjar

Caprimulgus europaeus

Clinal with four recognizable subspecies, one in northern Europe and Asia; another in southern Europe, North Africa and the Near East; a third in the Middle East and Central Asia; and the fourth in the extreme eastern part of the range. The first three occur regularly on migration in the eastern Mediterranean.

Habitat: Varies from all types of forest clearings to bushy hillsides and semi-desert.

Kingfisher *Alcedo atthis*

Part of Palearctic population is migratory wintering in southern part of breeding range and regularly in Sinai, Egypt, the Sudan and Arabia. Variation is slight and clinal so that birds in winter quarters are not assignable to subspecies.

Habitat: Fresh water streams and lakes, also on coast in winter.

Little Green Bee-eater *Merops orientalis*

Largely resident, but Egyptian population moves south to Sudan in winter.

Habitat: In Africa, semi-arid or arid areas from open savannah and acacia or thorn scrubs to virtually bare desert.

Roller

Coracias garrulus

Birds on passage in eastern Mediterranean to winter in Africa come predominately, if not exclusively, from eastern Europe, Asia Minor, and northern Turkestan. Iranian and southern Turkestan birds winter in Arabia.

Habitat: Open plains, pastures and cultivated fields with large trees, occasionally in open woods and orchards.

Hoopoe

Upupa epops

Variation is stepped with one subspecies (*epops*) breeding in Northwest Africa, Europe and western Siberia and migrating to tropical Africa, Arabia, the Middle East, and India; another subspecies is resident in Egypt, a third (*saturatus*) is migratory in the eastern Palearctic and several others occur in Africa and southeast Asia.

Habitat: Steppes, open areas in forested steppes and woodland, meadows, pastureland, parks, cultivation, orchards, vineyards, gardens and desert oases.

Wryneck

Jynx torquilla

Variation is clinal but only Western Eurasian (east to the Urals, but excluding Italy) nominate subspecies occurs in the Eastern Mediterranean.

Habitat: Open deciduous and, to a lesser extent, conifer forest, old trees in parkland, brushy plains, hedgerows, orchards and gardens.

Short-toed Lark *Calandrella cinerea*

Other Ethiopian races breed in Southern Arabia, eastern and southern Africa. Variation is clinal and slight so that migrating and winter birds can often not be identified to subspecies. Birds passing through eastern Mediterranean breed in eastern Europe and Near East.

Habitat: Barren scrub and wasteland, dry steppe, pastures and cultivation.

Calandra Lark *Melanocorypha calandra*

Variation is clinal, with southeastern and western extremes recognized as sub-species. Migrants in Egypt probably come from steppes of USSR.

Habitat: Arid grassy and stony fields and steppes up to mountain plateaus.

Sand Martin

Riparia riparia

Both resident and Eurasian migratory birds occur in Egypt. Migrants may come from as far east as Baikal. Birds breeding in Egypt migrate south to Sudan and Eritrea.

Habitat: Open country near water.

Crag Martin *Hirundo rupestris*

Monotypic with a slight clinal tendency to larger size in the easternmost populations.

Habitat: Rocky cliffs and gorges in mountains and seacoast.

Swallow

Hirundo rustica

Variation is clinal with one widespread subspecies (*rustica*) in western Eurasia to the Yerisei south to western North Africa and the Himalayas, which migrates to Africa; another (*transitiva*) in the Near East, which occurs in winter in Egypt; a third (*savignii*) is resident in Egypt; others in the Far East.

Habitat: Open and cultivated country and human settlements.

Red-rumped Swallow *Hirundo daurica*

Precise African winter quarters of European and southwest Asian (east to Kashmir) subspecies (*rufula*) unknown, but there is a regular migration through the Nile Valley. Five subspecies breed in Africa but are not shown on the map.

Habitat: Open country, forest clearings, and human settlements.

House Martin

Delichon urbica

Birds from the western Palearctic east to western Siberia and West Pakistan winter in Africa. Although there is some clinal variation all belong to a single subspecies which also winters in India.

Habitat: Open country of all sorts from rocky cliffs in high mountains through cultivation to towns and cities.

Tree Pipit *Anthus trivialis*

Variation is mostly clinal with European and western Siberian birds wintering in Africa.

Habitat: Meadowland, hillsides and open woodland. Frequently perches in bushes and trees.

Red-throated Pipit *Anthus cervinus*

Monotypic with no geographic variation.

Habitat: Breeds on moist tundra.

Water Pipit

Anthus spinoletta

Southern European and southwest Asian mountain birds are difficult to differentiate in winter plumage. Both probably occur in Eastern Mediterranean and Egypt.

Habitat: Alpine meadows and tundra especially near streams, occurring at lower altitudes and on seacoast in winter. Other, non-Mediterranean, subspecies breed on seacoast in northern Europe.

Yellow Wagtail

Motacilla flava

This highly variable species breaks up into a number of well marked subspecies that can be identified by bands. Isolated individuals of exotic forms may be "sports" and have little significance but general breeding localities can be assigned to male migrants. The following subspecies are common in the eastern Mediterranean and Egypt on passage: *flava* (Southern Scandinavia, central and eastern Europe and Russia to the Urals), *thunbergi* (northern Scandinavia), and *feldegg* (southeast Europe, Ukraine, and Near East to Afghanistan). The last winters sparingly in east Africa and more commonly in India.

Habitat: Meadows, moors, tundra, steppes, marshland and cultivated fields usually near water.

Grey Wagtail *Motacilla cinerea*

Variation in continental populations is clinal, but so slight that migrants cannot be assigned easily to subspecies.

Habitat: Clear running streams or lakes in mountains or hills.

White Wagtail *Motacilla alba*

Variation is complex with a number of well marked geographic subspecies, but only one (*alba* breeds continental Europe to Urals and Asia Minor) migrating through eastern Mediterranean to Africa.

Habitat: Highly variable open country from tundra and farmland to desert, often near water.

Red-backed Shrike *Lanius collurio*

Variation is clinal with all but extreme eastern subspecies wintering in Africa. In addition to northern and western Eurasian birds (east to Urals), which are probably commonest on passage in the eastern Mediterranean, Caucasian and Iranian subspecies (*kobylini*) have been recorded.

Habitat: Isolated small trees and bushes in open country and edges of woods. Occurs in plains and mountains under mesic to near desert conditions.

Masked Shrike *Lanius nubicus*

Monotypic with no geographic variation.

Habitat: Olive groves, gardens, lightly wooded country or high scrub.

Wood-chat Shrike *Lanius senator*

Variation is clinal with both European and Asiatic birds recorded on migration in eastern Mediterranean.

Habitat: Open ground with scattered trees, bushes and scrub, olive groves, gardens and edges of woods.

Lesser Grey Shrike *Lanius minor*

Variation is clinal with no clear-cut separation between the two poorly differentiated subspecies.

Habitat: Grasslands, cultivation and gardens with scattered trees and bushes as well as forested steppes.

Great Grey Shrike *Lanius excubitor*

Two groups of subspecies: a northern one of partly migratory subspecies and a southern one of largely resident forms. Only the resident has been recorded regularly in Egypt.

Habitat: In Egypt in desert scrub.

Dunnock

Prunella modularis

Migrants in eastern Mediterranean belong both to northern continental subspecies (*modularis*) and probably to lesser extent to the Asia Minor and Near Eastern subspecies (*euxina* and *obscura*).

Habitat: Variable, from bushy scrub and garden shrubs to undergrowth in open deciduous and coniferous woods. Occurs from sealevel to mountains.

Cetti's Warbler *Cettia cetti*

Variation is clinal. Migrants in Cyprus appear to be East European and Asia Minor birds.

Habitat: Heavy thickets and reed beds near water.

Savi's Warbler *Locustella luscinioides*

Eastern and western populations can be distinguished; both migrate through the eastern Mediterranean.

Habitat: Open reedbeds with mixed undergrowth of bushes and sedges.

River Warbler *Locustella fluviatilis*

Monotypic, with no geographic variation.

Habitat: Bushes along streams and thick undergrowth in woods.

Moustached Warbler *Acrocephalus melanopogon*

Variation is clinal but only European subspecies, *melanopogon*, winters in Africa.

Habitat: Extensive reedbeds and lake borders.

Sedge Warbler *Acrocephalus schoenobaenus*

Also breeds sparingly on Lake Eber, Turkey.
Monotypic, with no geographic variation.

Habitat: Osier and reedbeds, heavy marshy
undergrowth and wet cropland.

Marsh Warbler *Acrocephalus palustris*

Very slight clinal variation with populations not identifiable on migration. Also breeds sparingly in southern Turkey.

Habitat: Osier beds and heavy undergrowth beside streams and lakes as well as cultivated crop or grassland and steppes.

Reed Warbler *Acrocephalus scirpaceus*

Two subspecies, both passing through north-east Africa.

Habitat: Reed and osierbeds, as well as bushes on stream and lake banks.

Great Reed Warbler *Acrocephalus arundinaceus*

Variation is clinal. All the western Eurasian populations, including that of Mesopotamia, winter in southern Africa. All PMS specimens have been western subspecies *arundinaceus*.

Habitat: Reed beds.

Icterine Warbler *Hippolais icterina*

Monotypic with no variation, but replaced from Spain, France and northwestern Africa to Italy by the closely related Melodious Warbler, *Hippolais polyglotta*.

Habitat: Gardens with trees and bushes, woods with undergrowth, and scrub often in damp areas.

Olive-tree Warbler *Hippolais olivetorum*

Highly local within range shown. Monotypic with no geographic variation.

Habitat: Gardens, olive groves and open oak woods.

Olivaceous Warbler *Hippolais pallida*

Southeast European and Asian subspecies (*elaeica*) passes through Eastern Mediterranean and Egypt on migration; Egyptian subspecies (*pallida*) also migrates south.

Habitat: Small trees in orchards, olive, pine and oak groves, thickets and scrub, often near water.

Barred Warbler *Sylvia nisoria*

Variation is clinal but slight. Majority of birds in eastern Mediterranean probably come from eastern Europe and Russia. Turk-
 estan and Mongolian birds winter-
 ing in Arabia and Kenya, do not
 apparently enter Egypt.

Habitat: Bushes, thickets, hedge-
 rows and scrub and undergrowth
 in clearings in deciduous or mixed
 woods.

Orphean Warbler

Sylvia hortensis

Variation is strongly clinal from west to east with four well-marked subspecies. Only that breeding from Yugoslavia and Cyrenaica east to Transcaucasia has been recorded on passage in the eastern Mediterranean.

Habitat: Large bushes and small trees in open woods with undergrowth, orchards, olive, cork and oak groves, gardens and high maquis.

Garden Warbler *Sylvia borin*

Variation is clinal but slight with eastern and western extremes recognized as subspecies. Both pass through eastern Mediterranean.

Habitat: Deciduous and mixed woodlands with heavy undergrowth, also hedgerows, shrubbery and bushes in gardens.

Blackcap *Sylvia atricapilla*

Variation is clinal, but not pronounced, in continental populations. Most migrants in eastern Mediterranean come from central Europe and probably western Russia.

Habitat: Open deciduous and conifer woodland with thick undergrowth, also gardens.

Whitethroat

Sylvia communis

Variation is clinal with extremes in west, northeast, and southeast recognized as subspecies, all three of which have been recorded on migration in the eastern Mediterranean.

Habitat: Open pastureland and meadows, rough hillsides and gardens with bushes, thickets, hedges, and brambles.

Lesser Whitethroat *Sylvia curruca*

Variation is highly complex and still needs further work. Only European birds have been identified during PMS in eastern Mediterranean, but Siberian, and southwest Asian birds, have been recorded in northeast Africa by others.

Habitat: Similar to that of Whitethroat but prefers areas with more trees and higher bushes.

Desert Warbler *Sylvia nana*

Different subspecies breed western Sahara and central Asia with the latter reaching northeast Africa on migration.

Habitat: Sparse bushes such as acacia, saxaul, or tamarisk in sandy or rocky desert, semidesert, and steppes.

Rüppell's Warbler *Sylvia rueppelli*

Monotypic with no geographic variation.

Habitat: Thick maquis scrub in gullies
and rocky slopes.

Sardinian Warbler *Sylvia melanocephala*

Variation is imperfectly known but may be highly localized in the Near East. At least two populations have been recorded on migration in the eastern Mediterranean—a larger darker continental form (*melanocephala*) and a smaller paler Near Eastern form (*momus*). Mostly resident but some migrate to Cyprus and Egypt and occur irregularly in northern Iraq, Sudan and Aden.

Habitat: Low scrub thickets and bushes in dry areas.

Subalpine Warbler *Sylvia cantillans*

Three slightly differentiated subspecies: in western Europe to Italy (*cantillans*) North Africa (*inornata*) Balkans to Turkey (*albistriata*). Only the last occurs in eastern Mediterranean and migrates through Egypt.

Habitat: Similar low maquis scrub as that of Sardinian Warbler but often in less arid and thicker undergrowth in open woodland.

Spectacled Warbler *Sylvia conspicillata*

Only a single subspecies is found in continental Europe, North Africa and the Near East although another occurs on Atlantic islands. Distribution is poorly documented.

Habitat: Thickets, bushes, shrubs or low scrub maquis in arid localities.

Willow Warbler *Phylloscopus trochilus*

Varies clinally. Migration through Mediterranean region appears to be largely straight north-south while eastern Siberian birds may miss northeast Africa entirely on passage.

Habitat: Open woodlands and open country with scattered trees, conifer and birch forest, willow scrub and tundra.

Chiffchaff

Phylloscopus collybita

Markedly clinal in northern Palearctic populations that are most highly migratory. Only western Palearctic subspecies *collybita* and *abietinus* (England and France to Russia and Caucasus) migrate to eastern Mediterranean.

Habitat: Undergrowth in copses, woods, and heavy forest both deciduous and conifers as well as scrub beyond the tree limit.

Bonelli's Warbler *Phylloscopus bonelli*

Two well marked subspecies, with only the eastern migrating through eastern Mediterranean.

Habitat: Oak and pine forest in hills and mountains.

Wood Warbler *Phylloscopus sibilatrix*

Monotypic with no geographic variation.

Habitat: Upper story in oak, beech, mixed
and coniferous forests.

Fan-tailed Warbler *Cisticola juncidis*

Resident showing no known dispersal movements.

Habitat: Damp grass or wastelands.

Pied Flycatcher *Ficedula hypoleuca*

Variation is clinal with the extremes recognizable. Both northern European and Siberian subspecies migrate through the eastern Mediterranean.

Habitat: Open oak and birch forest and old woods and less frequently coniferous forest; also wooded gardens, orchards and villages.

Collared Flycatcher *Ficedula albicollis*

Monotypic with no geographic variation
but see *F. semitorquata*.

Habitat: Open forest and old woods of
oak, birch, and mixed conifers; wooded
gardens, orchards, and villages.

Half-collared Flycatcher *Ficedula semitorquata*

Morphologically mid-way between Pied and Collared Flycatcher but considered closer to and probably conspecific with the latter.

Habitat: As in Collared Flycatcher.

Red-breasted Flycatcher *Ficedula parva*

Individuals of the European population irregularly pass through the eastern Mediterranean. Isolated Himalayan population may be a separate species (*F. subrubra*).

Habitat: Undergrowth in mixed woodland and less frequently conifer forest.

Whinchat

Saxicola rubetra

Possibly also breeds sparingly in most of North Africa. Monotypic with slight clinal trends that are of no use for differentiating birds on passage.

Habitat: Grassland with bushes, lowland meadows, marshes, upland pastures and hillsides.

Stonechat

Saxicola torquata

Several subspecies, not shown on map, breed in most of Africa. European, Near East, Asia Minor, Caucasus and Iranian birds all winter in Northeast Africa.

Habitat: Bushy grassland, lowland meadows, rocky upland pastures, wasteland, hillsides and mountains, also sea coast.

Wheatear

Oenanthe oenanthe

Continental Eurasian populations do not vary markedly nor constantly enough to be recognized on passage or in winter quarters. Northwest African and Somaliland populations are resident and distinct.

Habitat: Stony barren wastes, with or without scrub, tundra, steppes, deserts, rocky outcrops in forest, pastureland from sealevel to high plateaus and mountaintops.

Pied Wheatear *Oenanthe pleschanka*

Both isolated endemic Cypriot population and Eurasian continental birds occur on migration in Egypt.

Habitat: Stony slopes and broken ground with sparse vegetation especially in steppes and low mountains; also in gardens, cultivation, dry grassland and very open woods.

Black-eared Wheatear *Oenanthe hispanica*

Populations east from Yugoslavia, which are distant subspecies, migrate through the Mediterranean.

Habitat: Open rocky scrub in arid areas, vineyards, pastures and rough cultivation from sealevel to low mountains.

Finsch's Wheatear *Oenanthe finschii*

Two subspecies; birds wintering in eastern Mediterranean come from Asia Minor and Palestine east to the Caucasus; birds breeding from Iran eastward winter from Iraq to India.

Habitat: Arid rocky terrain and rough cultivation, especially in mountainous areas.

Isabelline Wheatear *Oenanthe isabellina*

Morotypic; geographic variation in color and size has not yet been fully documented.

Habitat: Steppes, stony deserts, arid plains and mountains, adopting unused rodent burrows for nesting.

Rufous Bush Chat
or Warbler *Cercotrichas galactotes*

Variation is clinal in Eurasia but subspecies are easily recognizable. In winter and on passage subspecies from southeast Europe, Near East and Middle East are recorded in northeast Africa.

Habitat: Bushes and thickets in dry or arid areas.

Rock Thrush *Monticola saxatilis*

Monotypic, with a slightly developed clinal variation in the far eastern part of range.

Habitat: Open rocky regions in hills or high mountains.

Black Redstart *Phoenicurus ochruros*

European birds are distinctly different from those in Asia with an intermediate population breeding in Asia Minor. European subspecies (*gibraltariensis*) and occasionally Iranian and Himalayan subspecies (*phoenicuroides*) have been taken in Egypt; neareastern subspecies (*semirufus*) occurs in Cyprus in winter.

Habitat: Open rocky ground, cliffs, steep banks, ruins, gardens, orchards, vineyards and even in cities from sealevel to high mountains.

Redstart

Phoenicurus phoenicurus

Variation is clinal with two well marked subspecies, one widespread in western and northern Eurasia and the other in the Middle East, with an area of intergradation in Turkey. Both subspecies are found in the eastern Mediterranean and Egypt but the more widespread predominates.

Habitat: Deciduous woods, parks and gardens with bushes and old trees, occasionally in pine forests and ruins.

Robin

Erithacus rubecula

Highly variable with distinct clinal trends. On color characters birds migrating to Cyprus and northeast Africa belong to north European and to a lesser extent Balkan populations.

Habitat: Undergrowth and scrub in gardens, open woodland, and heavy mixed and conifer woodland.

Nightingale *Luscinia megarhynchos*

Variation is clinal, and both European and Turkestan birds have been recorded in eastern Mediterranean on passage.

Habitat: Undergrowth of deciduous woods, hedgerows, bushes, and thickets.

Thrush Nightingale *Luscinia luscinia*

Monotypic, with no geographic variation.

Habitat: Thick undergrowth in damp deciduous woods; also swampy or water-side thickets.

Ring Ouzel *Turdus torquatus*

Variation is generally clinal with South-east European and less commonly Scandinavian and Caucasian subspecies having been recorded in eastern Mediterranean, Sinai and Egypt.

Habitat: Open moorland, rocky ravines, conifer forest fringe in mountains.

Blackbird

Turdus merula

Variation is clinal with several subspecies recognized. There is a considerable influx of migrants presumably from the northern part of the range, into the Mediterranean in winter while local birds are resident. Birds wintering in Egypt have been identified as belonging to central European subspecies (*merula*).

Habitat: Undergrowth in woods, gardens and scrub from sealevel to treeline.

Redwing *Turdus iliacus*

Monotypic on the continent with no variation.

Habitat: Birch, alder, and open conifer forest and scrub willows and birches in tundra.

Song Thrush *Turdus philomelos*

Variation is clinal with a single subspecies breeding from Scandinavia and central Germany eastward and wintering in the eastern Mediterranean.

Habitat: Woodland undergrowth, hedgerows, thickets, garden shrubbery, and scrub.

Corn Bunting *Emberiza calandra*

Mostly resident; migrants found in Egypt in winter are presumably from northern parts of range. South-eastern birds from Near and Middle East which belong to different subspecies, have only been recorded south to Sinai in winter.

Habitat: Grasslands and cultivation with hedgerows and bushes, pastures, and scrubby wasteland.

Ortolan Bunting *Emberiza hortulana*

Monotypic, with no variation in western Palearctic populations.

Habitat: A variety of open hilly or flat country with scattered trees and or scrub, in southern part of breeding range in mountains.

Cretschmar's Bunting *Emberiza caesia*

Monotypic, no geographic variation.

Habitat: Arid and barren rocky slopes
with scrubby vegetation.

Blackheaded Bunting *Emberiza melanocephala*

Monotypic with no geographic variation.
In southern Caspian region meets and
hybridizes with Red-headed Bunting,
Emberiza bruniceps.

Habitat: Scrubby plains or foothills,
open woodland with undergrowth, or-
chards, vineyards and gardens.

Chaffinch *Fringilla coelebs*

Variation is clinal but slight in European and northern Asian populations that migrate into Mediterranean and northeast Africa. Southern populations are resident.

Habitat: Gardens, hedgerows and thickets; bushy farmland; and open woodland, both conifer and deciduous.

Serin *Serinus serinus*

Monotypic with no geographic variation. The species has spread rapidly north and west in the past 150 years. Closely related to the Syrian Serin, *Serinus syriacus*, of the mountain cedars of Syria and Lebanon that winters in Iraq, Palestine, Sinai and occasionally Egypt.

Habitat: Open woodland, cultivation, gardens, parks, vineyards, orchards and hedges from sealevel to tree limit.

Greenfinch *Carduelis chloris*

Chiefly sedentary, but northern birds may move to southern part of range as well as Cyprus and Sinai in winter. Variation is clinal, with only southern European, Asia Minor, Palestinian birds occurring in the eastern Mediterranean.

Habitat: Open woodland, orchards and gardens.

Siskin *Carduelis spinus*

Monotypic, no geographic variation.

Habitat: Breeds in coniferous and occasionally mixed woodland.

Goldfinch *Carduelis carduelis*

Both sedentary and partly migratory; northern populations winter south to the Iberian Peninsula, Mediterranean Islands, Near East and southern Iran. Variation is clinal in the Western Palearctic (west of the Yenesei and Caspian; a totally different subspecies group occurs in east) with only resident Balkan, Near and Middle East birds (*niediecki*) recorded in Egypt in winter.

Habitat: Open cultivated areas, gardens, pastures and brush land with scattered trees.

Linnet *Acanthis cannabina*

Variation is clinal, both eastern and western Palearctic subspecies winter in Northeast Africa.

Habitat: Open country and gardens with low scrub growth including low mountains in southern range.

House Sparrow *Passer domesticus*

Resident, with two distinct groups of subspecies. One inhabits most of northern Palearctic, southern one occurs in Sudan, Arabia, Iran, southern Turkestan, India and Burma. Within each geographic variation is correlated with environmental factors. Hybridizes with migratory Spanish Sparrow in Italy, some oases of north Africa, and Aegean Islands.

Habitat: Commensal with man inhabiting cities, towns, villages and farm yards and feeding in nearby cultivated land.

Spanish Sparrow *Passer hispaniolensis*

Partly migratory. Variation is clinal with two poorly marked subspecies. Most of the birds migrating into Egypt in winter are probably from Balkans and nearby Near East. Hybridizes with resident House Sparrow in Italy, some oases of North Africa and Aegean Islands.

Habitat: River bottoms, open woods, agricultured fields, and palm groves.

Dead Sea Sparrow *Passer moabiticus*

Isolated populations in Southeast Turkey, northern Syria (small circles), Jordan valley, Iraq, and Seistan.

Habitat: Near water in bushes or deep tamarisk or poplar scrub. In winter feeds in date palms.

Golden Oriole *Oriolus oriolus*

All western and northern Palearctic birds winter in Africa and are not individually distinguishable. Birds from eastern Russian and Chinese Turkestan south, winter in India.

Habitat: Deciduous, mixed and less commonly conifer woods, copses, and second growth forest; forested steppes; orchards; high windbreaks in cultivation and along rivers and parks and gardens with trees.

AMBLYOMMA LEPIDUM

AMBLYOMMA NUTTALLI

AMBLYOMMA VARIEGATUM

HAEMAPHYSALIS CONCINNA

(Also breeds in Japan,
Korean Peninsula, eastern
Manchuria.)

HAEMAPHYSALIS PARVA (OTOPHYLA)

HAEMAPHYSALIS PUNCTATA

HYALOMMA MARGINATUM MARGINATUM

HYALOMMA MARGINATUM RUFIPES

IXODES FRONTALIS

IXODES REDIKORZEVI subsp.

IXODES RICINUS

BAHIG and EG AN 1047-61

EG AN 890-3

QUARANTIFIL

UNKUNIEMI

SPECIES

Band number	<u>Age and Sex</u> Recovery remarks	<u>Locality banded</u> Locality recovered	<u>Date banded</u> Date recovered
ARDEA CINEREA			
B 55534 Moskwa	Juvenile	Mouth of Volga River, Astrakhan Reserve, USSR 46.00 N, 48.33 E	16-27 May, 1961
	Recovered	Port Said, Egypt 31.17 N, 32.18 E	November, 1965
B 55 918	Juvenile	Astrakhan Reserve, Volga Delta, USSR 46.14 N, 49.02 E	25 May, 1962
	Shot by hunter	Saiyala, ca. 80 km S of Aswan, Egypt, UAR	Date unknown (Reported 1970)
ARDEA PURPUREA			
D 440160 Moskwa		Krasnodar Region, Primorsk-Akhtorsk 46.03 N, 38.09 E	2 July 1959
		Lake Burullus, Egypt 31.30 N, 30.50 E	19 March 1969
C-88996 Moskwa	Pullus	Near Primorsko-Akhtarsk (Krasnodar Region) USSR 46.03 N, 38.09 E	11 July 1960
	Shot by hunter	Mersa Matruh, Egypt, UAR 31.21 N, 27.14 E	September, 1968
D 495955 Moskwa	Juvenile	Near Primorsko-Akhtarsk, (Krasnodar Region) USSR 46.03 N, 38.09 E	13 June, 1961
	Recovered	Port Said, Egypt 31.17 N, 32.18 E	Late Fall of 1965 or winter, 1965-66
EGRETTA ALBA (See Platalea leucorodia)			

EGRETTA GARZETTA

D 943 Tel-Aviv Univ.		Tel-Aviv, Israel	31 July 1964
	Shot by hunter	New Kalabsha, Egypt	1965-66
C-85373 Moskwa	Pullus	Chernomorski Reserve Kherson Region, USSR	26 June 1967
	Shot by hunter	46.27 N, 31.55 E Mersa Matruh, Egypt	24 October 1969

NYCTICORAX NYCTICORAX

24775 Budapest	Juvenile	Retszilas, Hungary	26 June 1957
		46.50 N, 18.35 E Salum, Egypt	25 September 1957
C 230378 Zagreb	Pullus	Tikves, Bilje, Hrvatska, Jugoslaviya	23 June 1969
	Shot by hunter	45.37 N, 18.43 E Mersa Matruh, Egypt, UAR	4 October 1969
		31.21 N, 27.14 E	

PLATALEA LEUCORODIA or EGRETTA ALBA [sic]

B 1420 Radolfzell	Nestling	Near Neusiedl am See Burgenland, Austria	7 or 8 June 1950
		47.57 N, 16.51 E) El Manzala, Egypt	30 May 1951
		(Banding records incomplete)	(Reported: 14 June 1951; 6 February 1970)

PLEGADIS FALCINELLUS

127322 Budapest	Juvenile	Kisbalaton, Hungary	16 June 1938
		46.40 N, 17.15 E Lake Burullus, Egypt	May 1946
		31.30 N, 30.50 E	

CICONIA CICONIA

MVD 1628T Denmark	Pullus	Farup, Denmark	11 July 1919
	Shot by hunter	55.21 N, 08.44 E Old Kalabsha, c. 50 km S. of El Shallal, Egypt	Before 1964-65
		23.33 N, 32.52 E	

CICONIA CICONIA

207 876 Zool. Station Helgoland	Nestling	Moorburg, Bez. Hamburg, Germany 53.29 N, 09.55 E	2 July 1932
	Shot	Old Kalabsha, near El Shallal, Egypt 23.26 N, 32.50 E	Before 1964-65 (Reported 6 March 1970)
S 12085 Sofia Bulgaria	Shot by hunter	Bulgaria Old Dihmit, ca. 45 km S El Shallal, Egypt	ca. 1934 Date unknown (Reported 1970)
214 799 Helgoland	Nestling	Seidlitz, Kr. Oppelm, Schlesian, Germany ca. 50.36 N, 17.43 E	30 June 1934
	Shot	El Ambarkab, Egypt 23.38 N, 32.51 E	Reported: 28 January 1970
S 19825 Royal Museum Sofia	Pullet	Kapitan, Andreyevo, Svilengrad, Bulgaria 41.42 N, 26.19 E	23 June 1936
		Suez, Egypt 29.55 N, 32.35 E	27 May 1937
B 500.835 Polonia- Varsovia	Pullus	Niezdow, arr. pualwy Poland 51.09 N, 21.56 E	16 June 1937
	Shot by hunter	Old Kalabsha, Egypt 23.33 N, 32.52 E	Before 1964-65
222913 Helgoland	Pullus	Wathlingen, b. Celle Germany 52.33 N, 10.10 E	8 July 1937
	Killed	Xylophagou, Larnaca, Cyprus 34.58 N, 33.40 E	28 September 1937
B 503.488 Polonia- Varsovia	Shot by hunter	Jaskowice, arr. Krzemieniec, Poland Old Kalabsha, Egypt 23.33 N, 32.52 E	7 July 1938 Before 1964-65
B 1827 Polonia- Varsovia	Shot by hunter	Podhajce, Poland 48.34 N, 25.11 E Old Kalabsha, Egypt 23.33 N, 32.52 E	15 July 1938 Before 1964-65

CICONIA CICONIA

D 30078	Zagreb	Pullus	Nard, Slavonija Hrvatska, Jugoslavija 45.40 N, 18.29 E	29 June 1939
			Sinai, Egypt	1 May 1940
225 942	Helgoland	Nestling	Trippigleben, Kr. Klotze, Bez. Magdeburg, Germany 52.32 N, 11.09 E	2 July 1939
		Shot	Old Kalabsha, near El Shallal, Egypt 23.26 N, 32.50 E	Before 1964-65 (Reported: 6 March 1970)
225241	Helgoland	Pullus	Elsfleth, Germany 54.14 N, 8.27 E	10 June 1939
		Killed	Avgolidha, Famagusta, Cyprus 35.21 N, 33.58 E	19 November 1939 (ca.)
S 34402	Sofia Bulfaria		Bulgaria	ca. 1940-1941
		Shot by hunter	Old Kalabsha, ca. 50 km S El Shallal, Egypt	Before 1964-65 (Reported 1970)
228 825	Helgoland	Nestling	Bergenhusen, Kr. Schleswig, Schleswig-Holstein, Germany 54.22 N, 09.20 E	7 July 1941
		Shot	Old Kalabsha, near El Shallal, Egypt 23.26 N, 32.50 E	Before 1964-65 (Reported 6 March 1970)
229 238	Helgoland	Nestling	Tiddische, Kr. Helmstedt, Niedersachsen, Germany 52.31 N, 10.47 E	1 July 1944
		Shot	Old Kalabsha, 50 km S El Shallal, Egypt 23.26 N, 32.50 E	Before 1964-65 (Reported 6 March 1970)
231 065	Helgoland	Nestling	Kasterf, Kr. Helmstedt, Niedersachsen, Germany 52.32 N, 10.31 E	24 June 1950
		Shot	Old El Umbarakab, near El Shallal, Egypt 23.40 N, 32.52 E	Before 1964-65 (Reported 6 March 1970)
B 1 820	Radolfzell	Nestling	Wichmannsdorf, Kr. Templin, Bez. Neubranden- burg, Germany 53.14 N, 13.39 E	2 July 1950
		Shot	Kalabsha, Egypt 23.33 N, 32.51 E	Before 1964-65 (Reported 6 March 1970)

CICONIA CICONIA

233 645 Helgoland	Nestling	Gamsen, Kr. Gifhorn, Niedersachsen, Germany 52.31 N, 10.32 E	16 June 1951
	Shot	Old El Umbarakab, Egypt 23.40 N, 32.52 E	Before 1964-65 (Reported 6 March 1970)
234 385 Helgoland	Nestling	Holstenniendorf, Kr. Rendsburg, Schleswig- Holstein, Germany 54.03 N, 09.21 E	5 July 1951
	Shot	Old Kalabsha, near El Shallal, Egypt 23.26 N, 32.50 E	Before 1964-65 (Reported 6 March 1970)
224 279 Helgoland	Nestling	Bleckede, Kr. Luneburg, Niedersachsen, Germany 53.18 N, 10.44 E	20 June 1953
	Shot	Old Kalabsha, near El Shallal, Egypt 23.26 N, 32.50 E	1962/63
ZMC 311 Denmark	Pullus	Kasturaa, Denmark 55.18 N, 09.21 E	24 June 1954
	Shot by hunter	El Umbarakab, 47 km S of El Shallal, Egypt 23.40 N, 32.53 E	In the 1950's
BB 7 208 Radolfzell	Nestling(3)	Teicha, Kr. Bautzen, Bez. Dresden, Germany 51.19 N, 14.28 E	26 June 1954
		Sorsok, Egypt 31.05 N, 31.20 E	10 June 1955 (Reported 6 Feb. 1970)
235 535 Helgoland	Nestling	Hamburg-Kirchwerder, Bez. Hamburg, Germany 53.26 N, 10.12 E	1 July 1954
	Shot	Old El Umbarakab, Egypt 23.40 N, 32.52 E	Before 1964-65 (Reported 6 March 1970)
235 642 Helgoland	Nestling	Holstenniendorf, Kr. Rednsburg, Schleswig- Holstein, Germany 54.03 N, 09.21 E	4 July 1954
	Shot	Old Kalabsha, near El Shallal, Egypt 23.26 N, 32.50 E	Before 1964-65 (Reported 6 March 1970)

CICONIA CICONIA

BB 5470 Radolfzell	Pullus	Wechingen, Germany 48.54 N, 10.37 E	16 July 1955
		Dhromolaxia, Larnaca, Cyprus 34.52 N, 33.35 E	13 September 1955
212 245 Helgoland	Just fledged	Lasronne, Kr. Harburg, Niedersachsen, Germany 53.24 N, 10.15 E	3 July 1956
	Shot	Old Kalabsha, near El Shallal, Egypt 23.26 N, 32.50 E	Before 1964-65 (Reported 6 March 1970)
D 103118 Zagreb	Pullus	Ecka, Vojvodina, Srbija, Jugoslavia 45.20 N, 20.26 E	7 July 1956
		El Umbarkab, 40 km S of Aswan, Egypt	Recovery, date un- known (Reported 1970)
BB 5 813 Radolfzell	Nestling	Emerkingen, Kr. Ehingen, Sudwurttemberg, Germany 48.12 N, 09.40 E	18 June 1957
	Shot	Kalabsha, Egypt 23.33 N, 32.51 E	Before 1964-65 (Reported 6 March 1970)
B-51 117 Moskwa	Pullus	Brest Region, Belovezh- skaya, Pushcha Reserve, USSR. 52.40N, 24.00 E	21 June 1957
	Shot by hunter	Old Tushka, 250 km SW of Kalabsha, Egypt	Recovery date un- known (Reported 1970)
B-51 148 Moskwa	Pullus	Brest Region, Belovezh- skaya, Pushcha Reserve, USSR. 52.40 N, 24.00 E	28 June 1957
	Shot by hunter	Old Kalabsha, 50 km S of El Shallal, Egypt	Recovery date un- known (Reported 1970)
241 983 Helgoland	Just fledged	Winsen, Kr. Harburg, Niedersachsen, Germany 53.22 N, 10.13 E	1 July 1957
	Shot	El Shalla, near Aswan, Egypt. 24.00 N, 32.52 E	1964
B-530.344 Polonia- Varsovia	Juvenile	Glezno, arr. Choszczno, Poland. 53.11 N, 15.26 E	7 July 1957
	Shot by hunter	Khor Teify, Egypt, UAR (ca. 40 km from Aswan)	ca. 1960

CICONIA CICONIA

BB 8 855 Radolfzell	Nestling	Eickstadt, Kr. Prenzlär, Bez. Neubrandenburg, Germany. 53.18 N, 14.03 E	15 June 1958
	Shot	Kalabsha, Egypt 23.33 N, 32.51 E	Before 1964-65 (Reported 6 March 1970)
238 066 Helgoland	Nestling	Hamburg-Neuengamme, Bez. Hamburg, Germany 53.27 N, 10.16 E	17 June 1958
	Shot	Suez, Egypt 29.58 N, 32.33 E	10 March 1959
BB 6 954 Radolfzell	Nestling	Sand. Kr. Hasfurt, Unterfranken, Germany 49.59 N, 10.35 E	17 June 1958
	Shot	New Kalabsha, near Kom Ombo, Egypt 24.26 N, 32.57 E	1969
238 731 Helgoland	Nestling	Strohkirchen, b. Hagenow, Bez. Mecklenburg, Germany 53.24 N, 11.19 E	29 June 1958
	Shot	El Ambarkab, Egypt 23.40 N, 32.53 E	ca. 1959
2404 Helgoland	Nestling	Hohenlepte, Kr. Zerbst, Bez. Sachsen Anhalt, Germany. 51.57 N, 12.01 E	1 July 1958
	Shot	Old Kalabsha, near El Shalla, Egypt 23.26 N, 32.50 E	Before 1964-65 (Reported 6 March 1970)
D 104548 Zagreb	Pullus	Idvor, Vojvodina, Srbija, Jugoslavia 45.11 N, 20.33 E	18 July 1958
	Shot by hunter	Dihmit, 25 km S of Aswan Egypt, UAR	Recovery date un- known (Reported 1970)
BB 10 687 Radolfzell	Nestling	Altzauche, Kr. Lubben, Bez. Cottbus, Germany 51.55 N, 14.01 E	28 June 1959
	Shot	bei El-Umbarakab. Egypt 23.40 N, 32.54 E	5 March 1963

CICONIA CICONIA

B 4 221 Radolfzell	Nestling	Pfatter, Kr. Regensburg, Oberpfalz, Germany 48.58 N, 12.23 E	18 June 1960
	Shot	El Umbarakab, Egypt 23.38 N, 32.54 E	Before 1964-65 (Reported 1970)
B-527.533 Polonia- Varsovia	Pullus	Ruda Sulowska, arr. Milicz, Poland 51.09 N, 21.56 E	5 July 1960
	Shot by hunter	Old Kalabsha, Egypt 23.33 N, 32.52 E	Before 1964-65
B-528.945 Polonia- Varsovia		Chycina, arr. Miedzyrzec, Poland. ca. 52.27 N, 15.35 E	27 June 1961
	Shot by hunter	Old El Umbarakab, Egypt 23.40 N, 32.53 E	Before 1964-65
236 578 Helgoland	Nestling	Wittbeck, Kr. Husum, Schleswig-Holstein Germany. 54.28 N, 09.12 E	30 June 1961
	Shot	Old Kalabsha, near El Shallal, Egypt 23.26 N, 32.50 E	Before 1964-65 (Reported 6 March 1970)
B-528.147 Polonia- Varsovia	Pullet	Zelezniki, arr. Milicz, Poland	30 June 1961
	Shot by hunter	El Ambarkab, 25 km S of Aswan, Egypt 23.40 N, 32.53 E	Date unknown (Reported Dec. 1969)
B-527.247 Polonia- Varsovia	Pullus	Stoleczno, arr. Czyluchow, Poland ca. 53.41 N, 17.22 E	14 July 1961
	Shot by hunter	ca. Khor Teify, Egypt, UAR (40 km from Aswan Dam)	(Reported 28 January 1970) Date unknown
244 534 Helgoland	Nestling	Moorhusen, Kr. Steinburg, Schleswig-Holstein, Germany 53.57 N, 09.24 E	12 June 1962
	Shot	Old Tushka, near El Shallal, Egypt 22.30 N, 31.53 E	Before 1964-65 (Reported 6 March 1970)
H 9801 Helgoland	Nestling	Sandel, Kr. Friesland Niedersachsen, Germany 53.33 N, 07.51 E	17 June 1963
	Shot	Old Tushka, near El Shallal, Egypt 22.30 N, 31.53 E	Before 1964-65 (Reported 6 March 1970)

CICONIA CICONIA

B-528.440 Polonia- Varsovia	Shot by hunter	Radziadz, arr. Milicz, Poland. 51.31 N, 16.58 E Old Tushka, Egypt 22.30 N, 31.53 E	1 July 1963 Before 1964-65
245 488 Helgoland	Nestling	Horsten, Kr. Wittmund, Niedersachsen, Germany 53.27 N, 07.56 E	14 July 1963
	Shot	Old Tushka, near El Shallal, Egypt 22.30 N, 31.53 E	Before 1964-65 (Reported 28 Jan 1970)
H 7055 Helgoland	Nestling	Mecklenhorst, Kr. Neustadt, Niedersachsen, Germany 52.30 N, 09.30 E	27 June, 1964
	Shot	Old Tushka, near El Shallal, Egypt 22.30 N, 31.53 E	Before 1964-65 (Reported 6 March 1970)
H 5585 Helgoland	Nestling	Erfde, Kr. Rendsburg, Schleswig-Holstein, Germany 54.18 N, 09.20 E	26 June 1966
	Shot	New Tushka, near Aswan, Egypt. 24.29 N, 32.33 E	1967
D 108129 Zagreb	Pullus	Melenci, Vojvodina, Srbija, Jugoslavija 45.30 N, 20.17 E	22 June 1967
	Shot by hunter	ca. 50 km S of El Shallal, Egypt, UAR	Recovery date un- known (Reported 1970)
B 1774 Romania	Juvenile	Dumbravioara, Mures, Transylvania, Romania 46.38 N, 24.34 E	10 July 1967
	Shot by hunter	Old Tushka, 250 km SW of Kalabasha, Egypt	Before 1964-65

ANAS CRECCA

E2210 Tel-Aviv Univ.	Female	Ma'agan Michael, Israel Wing: 170 mm (32.33 N, 34.55 E)	13 February 1968
	Shot by hunter	Damietta, Egypt 31.25 N, 31.48 E	17 February 1970

ANAS ACUTA

D-264 507 Moskwa	Adult Female	Astrakhan Region, Astrakhan Reserve, Blinov Island, USSR 46.25 N, 49.05 E	12 August 1964
	Killed by hunter	Lake Menzalla, Egypt UAR 31.15 N, 32.00 E	December 1968

PANDION HALIAETUS

E-1.073 Helsinki Museum	Pullet	Takajarvi, Hattula (Hame), Finland 61.00 N, 24.10 E	4 July 1954
	Shot	Old El Umbarakab, Egypt, UAR. 23.40 N, 32.53 E	Unknown

FALCO ELEONORAE

310.4491 British Museum	Pullus	Akrotiri, Cyprus 34.35 N, 32.57 E	15 September 1967
	Shot	Mersa Matruh, Egypt 31.21 N, 27.15 E	September 1969

FALCO NAUMANNI

060-04561 Cairo Zoo		Abassia, Cairo (NAMRU 3) Egypt, UAR	20 April 1967
	Shot - Made into skin for collection of Mr. Hassan Yousef el Hawi; male by des- cription "gray head"	Al Mu'allaqah Zahle, Lebanon. 33.50 N, 35.54 E	23 March 1970

COTURNIX COTURNIX

060-00022 Cairo Zoo	Female	Bahig, Egypt, UAR 30.56 N, 29.35 E	6 April 1967
		Makharadze, Georgian SSR 41.57 N, 42.02 E	22 December 1967
050-0173 Cairo Zoo		Maadi, Cairo, Egypt, UAR 29.58 E, 31.15 E	2 October 1968
	Shot by hunter	Galal, Egypt	4 October 1969

COTURNIX COTURNIX

050-0225	Cairo Zoo		Maadi, Cairo, Egypt, UAR 29.58 N, 31.15 E	4 October 1968
		Shot by hunter	Galal, Egypt	6 October 1969
050-0909	Cairo Zoo	Female	Maadi, Cairo, Egypt, UAR 29.58 N, 31.15 E	9 October 1968
		None given	Bakhchisarai, Crimean Region, USSR 44.45 N, 33.53 E	21 September 1969
050-0742	Cairo Zoo		Maadi, Cairo, Egypt, UAR 29.58 N, 31.15 E	25 October 1968
			Liuljakovo, Burgas, Bulgaria 42.52 N, 27.02 E	10 August 1969
050-4960	Cairo Zoo		Maadi, Cairo, Egypt, UAR 29.58 N, 31.15 E	25 October 1968
			Gurguljat, Sofia, Bulgaria 42.48 N, 22.57 E	24 August 1969
050-4967	Cairo Zoo	Female	Cairo, Egypt 29.58 N, 31.15 E	25 October 1968
			Faiyum, Egypt 29.19 N, 30.48 E	19 February 1969
060-00343	Cairo Zoo	Female	Ikingi Mariut, Egypt, UAR 31.00 N, 29.45 E	14 September 1969
		Shot by hunter	Safita, Syria 34.49 N, 36.07 E	20 April 1970
060-00402	Cairo Zoo	Male	Ikingi Mariut, Egypt, UAR 31.00 N, 29.45 E	16 September 1969
		Shot by hunter	Safita, Tartous Province, Syria. 34.49 N, 36.07 E	4 April 1970
060-00868	Giza Zoo Cairo	Male	Ikingi Mariut, Egypt, UAR 31.00 N, 29.45 E	20 September 1969
		Capture Shot	8 km NW Burg el Arab, Egypt Krivnja, Bulgaria Coordinance unknown	13 September 1970
060-00877	Cairo Zoo	Male	Ikingi Mariut, Egypt, UAR 31.00 N, 29.45 E	20 September 1969
		Shot by hunter	Ras el Hikma (formerly Ras el Kanayis), Egypt, UAR 31.08 N, 27.50 E	24 September 1970
060-00870		Male	Ikingi Mariut, Egypt, UAR 31.00 N, 29.45 E	20 September 1969
		Shot by hunter	10 km from Damascus, Syria 33.30 N, 36.18 E	10 April 1970

COTURNIX COTURNIX

060-01104 Giza Zoo Cairo	Male	Ikingi Mariut, Egypt, UAR 31.00 N, 29.45 E	22 September 1969
	Capture Shot	8 km NW Burg el Arab, EGYPT Samokov, Bulgaria 42.20 N, 23.33 E	29 August 1970
060-01357	Male	Ikingi, Mariut, Egypt, UAR 31.00 N, 29.45 E	26 September 1969
	Shot by hunter	El Homira, Akor, Lebanon	11 April 1970
060-01497 Cairo Zoo	Female	Ikingi Mariut, Egypt, UAR 31.00 N, 29.45 E	29 September 1969
	Shot by hunter	Vakif, Turkey	ca. September 1970
060-01573 Giza Zoo Cairo	Female	Ikingi Mariut, Egypt, UAR 31.00 N, 29.45 E	30 September 1969
	Capture: 8 km NW Burg el Arab, Egypt; weight-121.1 Shot	Peschtera, Bulgaria 42.02 N, 24.18 E	26 September 1970
060-02036 Giza Zoo Cairo	Female	Mersa Matruh, Egypt, UAR 31.21 N, 27.15 E	1 October 1969
	Capture: 82.6, weight Shot	Galal, Egypt 31.02 N, 28.10 E Tsar Asen, Bulgaria	15 August 1970
060-02038 Giza Zoo Cairo	Female	Mersa Matruh, Egypt, UAR 31.21 N, 27.15 E	1 October 1969
	Shot by hunter weight at band- ing, 106.4	El Abiat, Egypt, 18 km W of Mersa Matruh	11 October 1969
060-02055 Giza Zoo Cairo	Female	Mersa Matruh, Egypt, UAR 31.21 N, 27.15 E	2 October 1969
	Capture: Galal, Egypt; weight 91.2 Shot	Liuben Karavelovo, Bulgaria 43.21 N, 27.47 E	16 August 1970
060-01985 Cairo Zoo		Ikingi Mariut, Egypt, UAR 31.00 N, 29.45 E	3 October 1969
	Shot by hunter	Nabtia, Lebanon 33.23 N, 35.29 E	28 March 1970

COTURNIX COTURNIX

060-02074 Cairo Zoo	Male	Mersa Matruh, Egypt, UAR 31.21 N, 27.15 E	4 October 1969
	Killed	Alexandria Bird Market Alexandria, Egypt, UAR 31.13 N, 29.55 E	15 October 1969
060-05771 Giza Zoo Cairo	Female	Ikingi Mariut, Egypt, UAR 31.00 N, 29.45 E	8 October 1969
	Capture: 8 km NW Burg el Arab Shot	Trankovo, Bulgaria 42.17 N, 25.49 E	13 September 1970
060-02578 Giza Zoo Cairo	Male	Mersa Matruh, Egypt, UAR	10 October 1969
	Capture: Galal, Egypt Shot	Dobroplodno, Bulgaria 43.24 N, 27.22 E	9 August 1970
060-02746 Giza Zoo Cairo	Female	Mersa Matruh, Egypt, UAR 31.21 N, 27.15 E	11 October 1969
	Capture: Galal, Egypt Shot	Ognijanovo, Bulgaria	9 August 1970
060-06501 Cairo Zoo	Male	Ikingi Mariut, Egypt, UAR 31.00 N, 29.45 E	18 October 1969
	Shot by hunter	2 km W of Byblos, Lebanon 34.00 N, 35.40 E	8 April 1970
060-03347 Cairo Zoo	Female	Mersa Matruh, Egypt, UAR 31.21 N, 27.15 E	20 October 1969
	(Captured: Galal, Egypt; released Mersa Matruh) Shot by hunter	Lago di Patria, Napoli, Italy	25 September 1970
060-06760 Giza Zoo Cairo	Male	Ikingi Mariut, Egypt, UAR 31.00 N, 29.45 E	23 October 1969
	Capture: 8 km NW Burg el Arab, Egypt; weight 94.7 Shot	Tscherni Brijag, Bulgaria 43.07 N, 26.05 E	Reported in 1970
060-03710 Cairo Zoo	Male	Mersa Matruh, Egypt, UAR 31.21 N, 27.15 E	27 October 1969
	Shot by hunter	Meshta Hassan, Akor, Lebanon 34.37 N, 36.19 E	15 April 1970
060-07162 Cairo Zoo	Male	Ikingi Mariut, Egypt, UAR 31.00 N, 29.45 E	30 October 1969
	Shot by hunter	Banior, Tartous Province Syria. 34.50 N, 35.55 E	7 April 1970

LARUS FUSCUS

461145 Copenhagen		Christiansø, Denmark 55.19 N, 15.12 E El Morra, Egypt	30 June 1951 4 January 1955
H-65.001 Helsinki Museum	Pullus(2)	Koyra, Kangasala, (Häme), Finland 61.33 N, 24.14 E	4 July 1965
	Killed by hunter	Port Said, Egypt, UAR 31.17 N, 32.18 E	November 1966
H-102.031 Helsinki Museum	Pullus	Uotila, Saaksmaki, (Häme) Finland 61.11 N, 24.13 E	12 August 1968
	Found	ca. 52 km W of Alexandria, Egypt. 31.00 N, 29.35 E	September 1969
H-72.804 Helsinki Museum	Pullus	Isokalliosalo, Sahalahti (Häme), Finland 61.32 N, 24.25 E	26 June 1969
	Shot	Salum, Egypt, UAR 31.34 N, 25.09 E	ca. 1-7 Nov 1969
H-47-763 Helsinki Museum	Pullus	Ganlakarleby, (Vaza) Finland. 63.50 N, 23.05 E	15 July 1969
	Shot	Salum, Egypt, UAR 31.34 N, 25.09 E	10 October 1969

LARUS GENEI

E 105372 Moskwa	Pullus	Orlov Island, USSR 46.17 N, 31.45 E	10 October 1947
	Killed	Limassol, Cyprus 34.40 N, 33.00 E	December 1961
E 170336 Moskwa	Juvenile	Orlov Island, USSR 46.17 N, 31.45 E	3 July 1949
		Xeros, Nicosia, Cyprus 35.48 N, 32.50 E	10 October 1952
E 149423 Moskwa	Juvenile	Orlov Island, USSR 46.17 N, 31.45 E	3 July 1949
		Zyggi, Larnaca, Cyprus 34.43 N, 33.20 E	18 October 1949
E 161861 Moskwa	Juvenile	Orlov Island, USSR 46.17 N, 31.45 E	26 June 1950
		Salamis, Famagusta, Cyprus 35.10 N, 33.54 E	29 October 1950

CALIDRIS ALPINA

B1200 Tel-Aviv Univ.

Shot by hunter

Ma'agan Michael, Israel
Wing: 117 mm; Bill: 31 mm;
Weight: 46 gr; moulting.
El Manzalla, Egypt
30.20 N, 31.15 E

16 September 1968

1 January 1970

CALIDRIS ALBA

BE 78.134 Cyprus

Post Juvenile

Killed

Akrotiri, Cyprus
34.35 N, 32.57 E
near Sousse, Tunisia
35.50 N, 10.38 E

26 September 1967

14 May 1969

PHILOMACHUS PUGNAX

53145 Univ. Bologna-
Oss. Pisa

Killed by hunter

San Marco, Pisa, Italia
43.41 N, 10.23 E
Lago Menzalla, Ezbet el
Burg, Egypt, UAR

10 April 1968

February 1969

TRINGA GLAREOLA

4.046.587 Stockholm

Juvenile

Shot by hunter

Kvismaren, Narke, Sweden
59.11 N, 15.24 E
El Sinaniya, Egypt
31.24 N, 31.47 E

14 July 1968

6 April 1970

STERCORARIUS PARASITICUS

S-035.292 Helsinki
Museum

Pullet (2)

Shot

Ranni, Pygamaa (Turku-Porki) Finland. 60.55 N, 21.12 E
Damrou, Sidi Salim (Kafr el Shiekh), Egypt, UAR
31.16 N, 30.47 E

5 July 1969

15 October 1969

LARUS FUSCUS

PSV K 11871 Denmark

Pullus

/?/

Christiansø, Denmark
55.19 N, 15.12 E
Dahshur, Egypt
29.45 N, 25.48 E

June, 1933

11 May 1940

PSV K 11631 Denmark

Pullus

Christiansø, Denmark
55.19 N, 15.12 E
Maadia, Egypt
31.34 N, 25.48 E

26 June 1933

31 January 1934

ZMC M 9835 Denmark

Pullus

/?/

Christiansø, Denmark
55.19 N, 15.12 E
Bughaz, Egypt
31.29 N, 30.22 E (?)

26 June 1940

31 October 1940

ZMC M 9830 Denmark

Pullus

/?/

Christiansø, Denmark
55.19 N, 15.12 E
Port Said, Egypt
31.16 N, 32.18 E

26 June 1940

30 December 1940

LARUS GENEI

E 200241 Moskwa	Juvenile	Orlov Island, USSR 46.17 N, 31.45 E Engomi, Famagusta, Cyprus 35.09 N, 33.52 E	26 June 1950 20 February 1951
E 205357 Moskwa	Juvenile	Orlov Island, USSR 46.17 N, 31.45 E Larnaca, Cyprus 34.55 N, 33.39 E	16-18 June 1952 November 1952
E 205063 Moskwa	Juvenile	Orlov Island, USSR 46.17 N, 31.45 E Limassol, Cyprus 34.41 N, 33.05 E	16-18 June 1952 6 January 1953
E 228623 Moskwa	Juvenile	Orlov Island, USSR 46.17 N, 31.45 E Larnaca, Cyprus 34.55 N, 33.39 E	16-18 June 1952 27 December 1952
E 228386 Moskwa	Juvenile	Orlov Island, USSR 46.17 N, 31.45 E Limassol, Cyprus 34.41 N, 33.05 E	16-18 June 1952 14 January 1953
E-266 743 Moskwa	Pullus	Kherson Region, Chernomorsk Reserve, Orlov Island, USSR 46.17 N, 31.45 E	24 June 1953
	Killed by hunter	Shatta, Egypt, UAR	10 January 1970
E 334469 Moskwa	Juvenile	Orlov Island, USSR 46.17 N, 31.45 E Syrianokhori, Nicosia, 35.13 N, 32.57 E Cyprus	22 June 1954 24 March 1955
E 334319 Moskwa	Juvenile	Orlov Island, USSR 46.17 N, 31.45 E Ayia Irini, Kyrenia, Cyprus. 35.18 N, 32.58 E	22 June 1954 24 November 1954
E 331615 Moskwa	Juvenile	Orlov Island, USSR 46.17 N, 31.45 E Larnaca, Cyprus 34.55 N, 33.39 E	22 June 1954 19 November 1954

LARUS GENEI

E 273208 Moskwa	Juvenile	Orlov Island, USSR 46.17 N, 31.45 E Xerovounos, Nicosia, Cyprus. 35.08 N, 32.44 E	24 June 1953 16 November 1953
E 275646 Moskwa	Juvenile	Orlov Island, USSR 46.17 N, 31.45 E Xeros, Nicosia, Cyprus 35.08 N, 32.50 E	22 June 1955 11 December 1955
E-380431 Moskwa	Juvenile	Orlov Island, USSR 46.17 N, 31.45 E Port Said, Egypt 31.16 N, 32.18 E	22 June 1955 1960
E 380862 Moskwa	Juvenile	Orlov Island, USSR 46.17 N, 31.45 E Larnaca, Cyprus 34.55 N, 33.39 E	22 June 1955 5 January 1956
E-420212 Moskwa	Juvenile	Orlov Island, USSR 46.17 N, 31.45 E Port Said, Egypt 31.16 N, 32.18 E	21 June 1956 1960
E-413349 Moskwa	Juvenile	Orlov Island, USSR 46.17 N, 31.45 E Port Said, Egypt 31.16 N, 32.18 E	21 June 1956 1960
E-413652 Moskwa	Juvenile	Orlov Island, USSR 46.17 N, 31.45 E Port Said, Egypt 31.16 N, 32.18 E	21 June 1956 1960
E-422634 Moskwa	Juvenile	Orlov Island, USSR 46.17 N, 31.45 E Port Said, Egypt 31.16 N, 32.18 E	21 June 1956 1960
E-420512 Moskwa	Juvenile	Orlov Island, USSR 46.17 N, 31.45 E Port Said, Egypt 31.16 N, 32.18 E	21 June 1956 1960
E-420701 Moskwa	Juvenile	Orlov Island, USSR 46.17 N, 31.45 E Port Said, Egypt 31.16 N, 32.18 E	21 June 1956 1960

LARUS GENEI

E-420248 Moskwa	Juvenile	Orlov Island, USSR 46.17 N, 31.45 E Port Said, Egypt 31.16 N, 32.18 E	21 June 1956 1960
E-420744 Moskwa	Juvenile	Orlov Island, USSR 46.17 N, 31.45 E Port Said, Egypt 31.16 N, 32.18 E	21 June 1956 1960
E-420057 Moskwa	Juvenile	Orlov Island, USSR 46.17 N, 31.45 E Port Said, Egypt 31.16 N, 32.18 E	21 June 1956 1960
E-380951 Moskwa	Juvenile	Orlov Island, USSR 46.17 N, 31.45 E Port Said, Egypt 31.16 N, 32.18 E	22 June 1955 1960
E-421 124 Moskwa	Pullus	Kherson Region, Chernomorsk Reserve Orlov Island, USSR 46.17 N, 31.45 E	30 June 1955
	Killed by hunter	Shatta, Egypt, UAR	10 January 1970
E-410255 Moskwa	Juvenile	Orlov Island, USSR 46.17 N, 31.45 E Port Said, Egypt 31.16 N, 32.18 E	22 June 1956 1960
E-414?24 Moskwa	Juvenile	Orlov Island, USSR 46.17 N, 31.45 E Port Said, Egypt 31.16 N, 32.18 E	22 June 1956 1960
E-415503 Moskwa	Juvenile	Orlov Island, USSR 46.17 N, 31.45 E Port Said, Egypt 31.16 N, 32.18 E	22 June 1956 1960
E-415491 Moskwa	Juvenile	Orlov Island, USSR 46.17 N, 31.45 E Port Said, Egypt 31.16 N, 32.18 E	22 June 1956 1960

LARUS GENET

E-410249 Moskwa	Juvenile	Orlov Island, USSR 46.17 N, 31.45 E Port Said, Egypt 31.16 N, 32.18 E	22 June 1956 1960
E-511360 Moskwa	Juvenile	Orlov Island, USSR 46.17 N, 31.45 E Port Said, Egypt 31.16 N, 32.18 E	27 June 1958 1960
E-512528	Juvenile	Orlov Island, USSR 46.17 N, 31.45 E Port Said, Egypt 31.16 N, 32.18 E	27 June 1958 1960
E-512914 Moskwa	Juvenile	Orlov Island, USSR 46.17 N, 31.45 E Port Said, Egypt 31.16 N, 32.18 E	27 June 1958 1960
E-511146 Moskwa	Juvenile	Orlov Island, USSR 46.17 N, 31.45 E Port Said, Egypt 31.16 N, 32.18 E	27 June 1958 1960
D-401531 Moskwa	Juvenile	Orlov Island, Tendra Bay Black Sea, USSR 46.17 N, 31.45 E Port Said, Egypt 31.16 N, 32.18 E	17 June 1959 1960
D-402419 Moskwa	Juvenile	Smaleny Island, Tendra Bay, Black Sea, USSR 46.15 N, 32.00 E Port Said, Egypt 31.16 N, 32.18 E	17 June 1959 1960
D-406931 Moskwa	Juvenile	Orlov Island, USSR 46.17 N, 31.45 E Port Said, Egypt 31.16 N, 32.18 E	17 June 1959 1960
D-403155 Moskwa	Juvenile	Smaleny Island, Tendra Bay, Black Sea, USSR 46.15 N, 32.00 E Port Said, Egypt 31.16 N, 32.00 E	19 June 1959 1960

LARUS GENEI

D-403934 Moskwa	Juvenile	Smaleny Island, Tendra Bay, Black Sea, USSR 46.15 N, 32.00 E Port Said, Egypt 31.16 N, 32.18 E	19 June 1959 1960
HYDROPROGNE CASPIA			
H-12.104 Helsinki Museum	Pullet	Tratten, Kyrkslatt (Nyland), Finland 59.59 N, 24.35 E Lake Manzala, Egypt, UAR 31.15 N, 32.00 E	11 July 1939 Before April 1941
H-12.113 Helsinki Museum	Pullet	Tratten, Kyrkslatt (Nyland), Finland 59.59 N, 24.35 E Lake Burullus, Egypt, UAR 31.30 N, 30.50 E	11 July 1939 8 May 1941
H-12.299 Helsinki Museum	Pullet	Tratten, Kyrkslatt (Nyland), Finland 59.59 N, 24.35 E Lake Manzala, Egypt, UAR 31.15 N, 32.00 E	28 June 1940 January 1943
H-12.158 Helsinki Museum	Pullet	Tirgrund, Kyrkslatt (Nyland), Finland 59.58 N, 24.34 E Suez, Egypt, UAR 29.58 N, 32.22 E	5 July 1940 January 1941
U 25.310 Stockholm	Pullet	Skvolken, off Lulea, Norrbotten, Sweden 65.22 N, 22.28 E	26 July 1947
	Recovered	Lake Burullus, Egypt c 31.25 N, 30.45 E	26 September 1954
U 40.756 Riksmuseum Stockholm	Pullet	Havringe, Sodermanland Sweden. 58.36 N, 17.19 E	6 July 1949
	Recovered	Lake Burullus, Egypt c 31.25 N, 30.45 E	29 November 1954
U 25.163 Stockholm	Pullet	Avlaggaren, off Lulea 65.26 N, 22.15 E	8 July 1948
	Recovered	Lake Burullus, Egypt c 31.25 N, 30.45 E	6 April 1955

HYDROPROGNE CASPIA

U 31.351 Stockholm	Pullet	Kudoxa Mellgrund, Uppland, Sverige, Sweden 59.37 N, 19.09 E	24 June 1952
	Recovered	Lake Burullus, Egypt, UAR c 31.25 N, 30.45 E	6 October 1957
40003, 40020 Sv. Jagareförbundet	Pullet	Sandgrynnan, off Lulea, Norrbotten, Sweden 65.23 N, 22.23 E	4 July 1953
	Recovered	Sidi Salim, Egypt 31.16 N, 30.47 E	10 October 1959
U 61.652 Stockholm	Pullet	Lacka Trutbada, Vasterljug Sodermanland, Sweden 58.44 N, 17.36 E	2 July 1954
	Recovered	Lake Manzala, Egypt, UAR c 31.15 N, 32.00 E	5 April 1957
H-19.828 Helsinki Museum	Pullet	Lokhall, Kyrklatt (Nyland), Finland 60.02 N, 24.41 E	20 June 1955
		Lake Burullus, Egypt, UAR 31.30 N, 30.50 E	5 November 1957
H-19.895 Helsinki Museum	Pullet	Lokhall, Kyrklatt (Nyland), Finland 60.02 N, 24.41 E	12 July 1955
		Lake Burullus, Egypt, UAR 31.30 N, 30.50 E	24 September 1959
C-602 Helsinki Museum	Pullet	Tratten, Kyrklatt (Nyland), Finland 59.59 N, 24.35 E	22 July 1956
		Lake Burullus, Egypt, UAR 31.30 N, 30.50 E	24 September 1959
U 71.513 Stockholm	Pullet	Bredskar, Oxelosund, Sodermanland, Sweden 58.38 N, 17.13 E	18 June 1957
	Recovered	Lake Burullus, Egypt c 31.25 N, 30.45 E	10 October 1957
H-23.897 Helsinki H-23.898 Museum	Pullet	Tratten, Kyrkelatt (Hyland), Finland 59.59 N, 24.35 E	22 June 1957
	Shot by hunter	Mersa Matruh, Egypt, UAR 31.21 N, 27.15 E	circca Fall 1959

HYDROPROGNE CASPIA

7.019.776 Stockholm	Pullet	Algsbadarna, Vestrum, Smaland, H, Sweden 57.41 N, 16.48 E	24 June 1961
	Killed by hunter	Ras el Barr, Egypt, UAR 31.32 N, 31.50 E	End of December 1969
H-46.343 Helsinki Museum	Pullet	Midfjardsbadan, Eckero (Aland), Finland 60.12 N, 19.25 E	24 June 1962
	Killed by hunter	Ras Al Barr, Egypt, UAR 31.31 N, 31.01 E	5 January 1970
7.015.873 Stockholm	Pullus	Fagelsundet, Hallnas Uppland, C. Sweden 60.36 N, 17.56 E	6 July 1964
	Killed by hunter	Ras el Barr, Egypt, UAR 31.32 N, 31.50 E	End of December 1969
7.016.044 Stockholm	Nestling	Ledskär, Österlövsta Uppland, Sweden 60.13 N, 17.43 E	7 June 1965
	Caught "seabird"	Egypt Lake Manzala, Port Said / 31.15 N, 32.15 E	6 September 1966
H-87.634 Helsinki Museum	Pullet	Tunnan, Kirkkonummi, (Uusimaa), Finland 59.59 N, 24.35 E	25 June 1967
	Killed by hunter Reported as <u>Larus</u> ?	Lake Menzalla, Egypt, UAR 31.15 N, 32.00 E	December 1969
H-67143 Helsinki Zoo		Gadden, Brando Finland 60.40 N, 20.56 E	27 June 1967
	Shot	Brolos Lake, UAR	19 March 1969
7.042.137 Stockholm	Pullus	Algabäddarna, Vestrum Smaland, Sweden 57.41 N, 16.48 E	30 June 1967-
	Shot by hunter	Salum, Egypt, UAR 31.34 N, 25.09 E	15 September 1969

LARUS RIDIBUNDUS

E-523135 Moskwa	Pullus	Lobnja (25 km N of Moscow), Russia 56.01 N, 37.30 E Port Said, Egypt 31.16 N, 32.18 E	14 June 1955 1960
-----------------	--------	---	------------------------------

LARUS RIDIBUNDUS

B 412958 Yugoslavia	Pullet	Ludas Lake, Subotica Vojvodina, Yugoslavia 46.04 N, 19.39 E	13 June 1965
	Recovered	Port Said area of Egypt 31.17 N, 32.18 E	Late Fall/1965 or Winter/1965-66

STREPTOPELIA TURTUR

DS 51003	Juvenile	Akrotiri, Cyprus 34.35 N, 32.57 E	20 September 1967
	Killed	Near Gulyay Pole (Zaporozh'ye) Ukraine, SSR 47.39 N, 36.15 E	11 August 1968
060-04591 Cairo Zoo		Cairo, Maadi, Egypt	11 September 1968
	Shot by hunter	16 km W. Alexandria, Egypt	19 September 1969

OTUS SCOPS

060-00025 Cairo Zoo		Bahig, Egypt	9 April 1967
	Trapped	Bahig Egypt	September 1967
	Previous recapture and release:		
	10 Apr 1967--Bahig--78.5 gm		
	11 Apr 1967--Bahig--74.3 gm		

CAPRIMULGUS EUROPAEUS

CS 71066	Adult Male	Akrotiri, Cyprus 34.35 N, 32.57 E	25 September 1967
	Trapped & killed	Mkolani, near Iringa Tanzania. 07.45 S, 35.40 E	6 March 1969
050-05073 Giza Zoo Cairo	F. (plumage)	Mersa Matruh, Egypt, UAR 31.21 N, 27.14 E	28 April 1970
	Shot by hunter	Bugbug, 40 km E of Salum, Egypt, UAR 31.31 N, 25.09 E	30 April 1970

APUS PALLIDUS

040-00168 Cairo Zoo		Ikingi Mariut, Egypt, UAR 31.00 N, 29.45 E	29 May 1969
	Killed-broken wing	Bahig, Egypt, UAR 30.56 N, 29.35 E	20 May 1970

UPUPA EPOPS

CS 71766 Cyprus	Post-Juvenile Taken	Paralimni, Cyprus Ayia Napa, near Famagusta Cyprus	23 March 1968 31 March 1968
-----------------	------------------------	--	--------------------------------

RIPARIA RIPARIA

HR 35.621	Full-grown	Malamfatori, Lake Chad (Bornu), Nigeria 13.33 N, 13.23 E	1 April 1968
	Lined	Paralimni, Cyprus 35.02 N, 34.00 E	September 1969

GALERIDA CRISTATA

030-01992 Cairo Zoo		Ikingi Mariut, Egypt, UAR 31.00 N, 29.45 E	28 May 1969
	Released Ikingi	Bahig, Egypt, UAR, (Khamis) 30.56 N, 29.35 E	28 May 1970

ANTHUS PRATENSIS

El1462 British Museum	First Year Found dead	RAF Nicosia, Cyprus RAF Nicosia, Cyprus	7 January 1959 8 March 1960
--------------------------	--------------------------	--	--------------------------------

MOTACILLA ALBA

ZG 5.949 Stockholm	Pullet	Gallas, Bergstena Vastergotland, Sweden 58.02 N, 12.39 E	9 June 1940
	Recovered	Giza, Egypt 30.01 N, 31.12 E	2 December 1940
A 25693 Goteborg	Pullet	Baktsjaur, Abborrtrask, Lapland, Pite lappmark, Sweden. 65.21 N, 19.16 E	25 June 1941
	Recovered	Baga Sohag, Egypt 26.33 N, 31.42 E	24 December 1941
921.503 Stavanger		Hadland, Time, County Rogaland, SW Norway 58.43 N, 5.47 E	29 June 1952
	Shot	Nagi Khade, Markaz el Koussia, Assiut Province, Upper Egypt (Giza Zool. Gardens)	7 February 1953
2.004.464 Stockholm	Juvenile	Ledskar, Uppland, Sweden Tanta, Egypt 30.48 N, 31.00 E	6 August 1960 25 January 1961

MOTACILLA ALBA

121.137 Ornis Box	Full-grown	Ottenby, Oland, Sweden	7 October 1956
	Recovered	56.12 N, 16.24 E El Alaqma, Egypt 30.38 N, 31.38 E	23 January 1957

MOTACILLA ALBA ALBA

HK 64014 Cyprus	Adult	Paralimmi, Cyprus	3 April 1968
	Caught by cat	35.01 N, 33.59 E Roznov pod Gadh. near Vsetin (Gottwaldov) Czechosl. 49.28 N, 18.09 E	April 1968

MOTACILLA ALBA

K-490.052 Helsinki Museum	Male, after first year	Valassaaret, Bjorkoby (Vaasa), Finland 63.25 N, 21.05 E	28 May 1969
	Killed by hunter	Bird Market, Alexandria 31.13 N, 29.55 E	24 February 1970
2.186.881 Stockholm	Pullet (5)	Stendammen, Vasterlanda Bohuslan, O, Sweden 58.02 N, 12.06 E	14 June 1969
	Killed	Alexandria (Bird Market) Egypt. 31.13 N, 29.55 E	10-17 December 1969
K-575.298 Helsinki Museum	After first year	Savilahti, Pirkkala (Hame), Finland 61.26 N, 23.33 E	15 July 1969
	Shot by hunter	Faiyum, Egypt 29.19 N, 30.50 E	1 January 1970
K-631.560 Helsinki Museum	First Year	Laajalahti, Espoo, Uusimaa, Finland 60.12 N, 24.49 E	11 August 1969
	Shot by hunter	Bourj el Brajeni, Ayn Sekka, Lebanon. ca. 34.10 N, 36.03 E	Fall of 1969
K-544.344 Helsinki Museum	First year	Niemi, Lahti, Hame, Finland 61.00 N, 25.40 E	29 August 1968
	Shot by hunter	Bourj el Brajeni, Ayn Sekka, Lebanon. ca. 34.10 N, 36.03 E	Fall, 1969
K-399.961 Helsinki Museum	Pullet	Holmola, Ruokilshti, (Kymi) Finland. 61.26 N, 29.01 E	18 June 1969
	Killed in Ataba Bird Market, Cairo	Kafr el Zayat, Egypt, UAR 30.50 N, 30.49 E	20 November 1969

MOTACILLA ALBA

K-623.040 Helsinki Museum	Pullet	Claresund, Narpes (Vasa) Finland. 62.30 N, 21.30 E	27 June 1969
	Killed	Alexandria Bird Market Alexandria, Egypt, UAR 31.13 N, 29.55 E	10-17 December 1969
K-626.763 Helsinki Museum	After first year	Mietoinen, Pyharabta (Turku-Pori), Finland 60.38 N, 21.52 E	31 July 1969
	Killed	Alexandria Bird Market Alexandria, Egypt, UAR 31.13 N, 29.55 E	10-17 December 1969
K-603.952 Helsinki Museum	First year	Suomenoja, Espoo (Uusimaa), Finland 60.09 N, 24.44 E	1 September 1969
	Killed	Alexandria Bird Market Alexandria, Egypt, UAR 31.13 N, 29.55 E	10-17 December 1969
K-610.917 Helsinki Museum	First year	Koivisto, Kangasala (Hame), Finland 61.27 N, 24.03 E	3 September 1969
	Caught & killed	Alexandria Bird Market Alexandria, Egypt, UAR 31.13 N, 29.55 E	26 December 1969

LANIUS COLLURIO

518567 sm Stockholm	Adult-male	Ottenby Bird Station Oland Island, Sweden 56.12 N, 16.24 E	24 May 1958
		El Amriya, 15 km W of Alexandria, Egypt 31.10 N, 29.50 E	11 September 1958
P80382 Helsinki Museum	Tammio,	Finland. 60.25 N, 27.25 E	31 August 1962
		Burg el Arab, Egypt 30.49 N, 29.20 E	27 September 1965
G 282066 Radolfzell	Post-Juvenile	Degenfeld, Schwabisch Gmund (Nord-Wurttemberg) Germany 48.44 N, 9.53 E	17 July 1965
	Killed (bird- lime)	Paralimni, Cyprus 35.01 N, 33.59 E	September 1967
P193435		Tali Helsinki, Finland 60.40 N, 24.52 E	22 August 1967
		Ikingi Mariut, UAR 31.13 N, 29.55 E	September 1968

LANIUS COLLURIO

4.061.507 Stockholm	Adult-male	Urakur, Gruso, Uppland Sweden. 60.31 N, 18.24 E	26 June 1968
	Captured on "lime- stick", prepared as study skin.	Bahig, Egypt 30.56 N, 29.35 E	5 September 1969

CALANDRELLA CINEREA

020-05445 Cairo Zoo		Ikingi Mariut, Egypt, UAR 31.00 N, 29.45 E	21 March 1969
	Bird in market May 1, 1970 for sale	Alexandria Bird Market Alexandria, Egypt, UAR 32.12 N, 29.54 E	ca. 25 April 1970

ACROCEPHALUS PALUSTRIS

4B 6471 Bruxelles	Pullus	Ieper, Belgium 50.51 N, 02.53 E	21 June 1952
	Found dead	Famagusta, Cyprus 35.09 N, 33.59 E	October 1952
801 27136 Vogelwarte Helgoland	Adult	Moischt near Marburg (Hessen), Germany 50.46 N, 8.49 E	9 July 1961
	Shot and killed	Cyprus (locality not known)	ca. 30 October 1961
H 806811 Radolfzell	Full-grown	Bretzenheim nr. Bad. Krenznach (Rheinland Pfalz) Germany. 49.53 N, 7.54 E	29 July 1962
	Lined	Famagusta, Cyprus 35.05 N, 33.57 E	10 January or February 1966
H 146062 Warsaw	First year	Zagarow (Slupka), Poland 54.50 N, 17.00 E	22 July 1964
		Famagusta, Cyprus 35.02 N, 34.00 E	Last week of August 1964

ACROCEPHALUS SCIRPACEUS

K578168 Germany	First year	Neusiedl am See, Burgenland, Osterreich 47.57 N, 16.51 E	2 August 1967
	Captured and killed	Bahig, Egypt 30.56 N, 29.35 E	8 August 1968

SYLVIA HORTENSIS

BE 78676 Cyprus	Adult-male	Paralimni, Cyprus 35.02 N, 34.00 E Kfar Akka, El Koura, Lebanon	13 April 1968 31 March 1969
-----------------	------------	---	--------------------------------

SYLVIA ATRICAPILLA

2,194,316 Rik Museum	Juvenile-male	Enskär, Bålinge, Södermanland, Sweden. 58.41 N, 17.29 E	8 Sept 1966
	Caught on lime- stick & killed	Paralimni, Cyprus 35.01 N, 33.59 E	March 1968
S 209897 USSR	Adult-female	Near Rybachii, Kaliningrad Region, USSR 55.11 N, 20.49 E	29 August 1965
	Found	Paralimni, Cyprus 35.02 N, 33.59 E	September 1967
2.276.883 Sweden		Lake Landsjon, Skarstad, Smaland, Sweden 57.52 N, 14.19 E	2 September 1967
	Shot (On band identified as male, upon re- capture, female.	Paralimni, Cyprus 35.01 N, 33.59 E	8 June 1968
HK 63.989 Cyprus	Adult-male	Paralimni, Cyprus 35.02 N, 34.00 E	2 April 1968
	Caught & caged	Beirut, Lebanon 33.52 N, 35.28 E	28 August 1968
A-6747 Tel-Aviv	Male	Elot, Israel 29.33 N, 34.57 E	23 April 1968
	Caught on lime- stick & killed	Paralimni, Cyprus 35.02 N, 34.00 E	October 1969
	Previous recapture: Elot, Israel 24 or 25 April 1968		
HK 64.686 British Museum	Full-grown female	Paralimni, Cyprus 35.02 N, 34.00 E	24 April 1968
	Shot by hunter	Ayn Sekka, Share Jdid, 35 kms NE of Baalbek, Lebanon ca. 34.20 N, 36.30 E	Autumn 1969
HA 51 464 Radolfzell	After first year	Marchegg, Bez. Ganserndorf Niederösterreich, Germany 48.17 N, 16.55 E	28 April 1968
	Caught & killed	Paralimni, Cyprus 35.02 N, 34.00 E	Fall 1969

SYLVIA ATRICAPILLA

K 134618 Radolfzell	Adult-male	Waldsassen, Kr. Tirschenreuth, Oberpfalz, Germany. 50.00 N, 12.18 E	18 May 1968
	Presumably caught on a limestick	Paralimni, Cyprus 35.02 N, 34.00 E	September 1968

SYLVIA COMMUNIS

HJ 32480 Nigeria	Male	Malamfatori, Bornu, Nigeria ca. 13.00 N, 14.00 E	13 April 1967
	Trapped	Bahig, Egypt	23 September 1968
564.243 Museum Paris	First Year female	Oued el Abid (Cap Bon) Republique Tunisienne 36.54 N, 10.44 E	3 May 1968
	Killed	El Dabaa (Matruh Governorate), Egypt UAR 31.03 N, 28.31 E	17 September 1969
2092297 Stockholm	Adult female	Algvikssjon, N. Nynashamn Sodermanland, Sweden 58.57 N, 17.55 E	30 August 1968
	Caught - Bird also banded Giza Zoo 020-09562, at King Mariut Province, 37 km W of Alexandria, UAR.	Bahig (c 60 km W Alexandria) Mariut Province, UAR 30.56 N, 29.35 E	7 May 1969

SYLVIA CURRUCA

P 41780 Chichester, R.G.	Full grown	Chichester (Sussex) England. 50.48 N, 0.48 W	15 August 1964
	Trapped w/lime- stick & released	Paralimni (Famagusta) Cyprus. 35.01 N, 33.59 E	3 April 1968
300815 Paris Museum		Arraincourt Mogelle, France 48.58 N, 6.32 E	16 May 1965
	Shot by hunter	Mersa Matruh, Egypt, UAR 31.21 N, 27.15 E	29 August 1968
M 523186 N. Museum Praha	Full grown	Zehuby, District of Kutna Hora, Czechoslovakia 49.52 N, 15.29 E	23 April 1967
	Trapped & killed	Paralimni, Famagusta area, Cyprus. 35.02 N, 34.00 E	20 March 1968
2.274.981 Stockholm	Adult	Orskar, Oregrund, Uppland, Sweden. 60.31 N, 18.24 E	30 May 1967
	Caught & killed	Paralimni, Cyprus 35.02 N, 34.00 E	Autumn 1969
90130267 Hiddensee	Adult	Serrahn, Krs. Neustrelitz, 53.20 N, 13.14 E E. Germany	14 June 1967
	Caught in net & killed by Bedouins. Fat-medium; male; adult; wing-64mm	8 km NW Burg el Arab, UAR	25 September 1969
HJ 63.634 British Museum	Post juvenile	Bloxham, Banbury (Oxford- shire), England 52.02 N, 01.22 W	11 August 1967
	Shot	Ikingi, Lake Mariut, Egypt ca. 31.05 N, 29.50 E	September 1969
K360405 Muszhki	After first year	Jurmo, Korppoo, (Turku-Pori) Finland. 59.50 N, 21.37 E	30 May 1968
	Obtained from Bedouins (dead) Skull-partial ossification; fat-none; wing-65mm	Bahig, Egypt 30.56 N, 29.35 E	9 October 1968
HK 63.970 Cyprus	Post juvenile	Paralimni, Cyprus 35.02 N, 34.00 E	1 April 1968
	Taken	Schio (Vicenza), Italy 45.43 N, 11.21 E	28 August 1968
HA 71 018 Radolfzell	First year	Lachen, Kr. Newstadt/ Weinstrabe, Pfalz, Germany 49.19 N, 08.12 E	8 August 1968
	Killed	Alexandria, Egypt 31.13 N, 29.55 E	10 October 1969

SYLVIA CURRUCA

12A 79565		Jupille (Lg.), Belgium 50.13 N, 05.32 E	3 September 1968
	Killed	Alexandria, Egypt 31.13 N, 29.55 E	10 October 1969
020-04625 Cairo Zoo		Maadi, Cairo, Egypt, UAR 29.58 N, 31.15 E	24 September 1968
	Shot by hunter	Railroad Station, Burg el Arab, Egypt, UAR 30.55 N, 29.32 E	Sept-Oct 1969
020-04996 Cairo Zoo		Maadi, Cairo, Egypt, UAR 29.58 N, 31.15 E	30 September 1968
	None given	Jambol, Bulgaria 42.28 N, 26.30 E	6 April 1969
020-03600 Cairo Zoo		Maadi, Cairo, UAR 29.58 N, 31.15 E	18 October 1968
	Killed by hunter	Galal, Egypt, UAR 31.02 N, 28.10 E	Nov-Dec 1969
1.419.915 Stockholm	Adult	Enskar, Balinge, Sodermanland, D. Sweden 58.41 N, 17.29 E	28 May 1969
	Caught & killed "Sylvia atricapilla"	Paralimni, Cyprus 35.02 N, 34.00 E	Autumn 1969
S 337.414 Arnhem	Full-grown	Oude Waal bij Nijmegen (Gelderland) Nederland 51.51 N, 05.54 E	25 July 1969
	Release same day, Ikingi Mariut, Egypt 31.00 N, 29.45 E Weight-12.4 gm; fat-light; wing-61.5 mm.	Burg el Arab, Egypt (8 km NW) 30.55 N, 29.32 E	14 October 1969
M 589 606 Praha	Juvenile	Rychory, District of Trutnov Czechoslovakia 50.40 N, 15.52 E	5 August 1969
	Controlled, re- leased at Ikingi Mariut, Egypt 30 October 1969	Hamam (23 km NW of Burg el Arab), Egypt	29 October 1969
A 416228		Tohogne (Luxembourg), Belgium. 50.23 N, 05.29 E	20 August 1969
	Killed	Alexandria, Egypt 31.13 N, 29.55 E	16 October 1969

SYLVIA CURRUCA

HX 55.323 British Museum	Adult	Seaford Head (Sussex), England. 50.46 N, 0.06 E	23 August 1969
	Shot	Ikingi, Lake Mariut, Egypt ca. 31.05 N, 29.50 E	10 October 1969 =

PHYLLOSCOPUS TROCHILUS

K450838 Muszhki Finland	Full-grown	Tauvo, Siikajoki (Oulu) Finland. 64.49 N, 24.35 E	25 August 1967
	Obtained from Bedouins; dead Skull-ossified; fat-medium; wing-65.5 mm The band was found on the tibia and could not slip over the tibia.	Bahig, Egypt (UAR) 30.56 N, 29.35 E	13 October 1968
1.347.979 Stockholm	Full-grown	Algvikssjon, N. Nynashamn Sodermanland, Sweden 58.57 N, 17.55 E	30 August 1968
	Killed by hunter	Mersa Matruh, Egypt 31.22 N, 27.15 E	24 September 1968
S4321263 Moskwa	Juvenile	Pape, Latvian, SSR, USSR 56.09 N, 21.02 E	7 September 1968
	Netted & killed by Bedouins. Wing feathers had been removed and identification is based on presumption of identity of other migrants captured with this individual.	Bahig, Egypt, UAR 30.56 N, 29.35 E	23 September 1969
S-467487 Moskwa	Juvenile	Near Rybachii - Kalingrad Region, USSR 55.11 N, 20.49 E	15 July 1969
	Shot by Bedouin	5 km NW Ikingi Mariut, Egypt, UAR 31.00 N, 29.45 E	25 September 1969
R-75.484 Helsinki Museum	First year	Kuitula, Pori (Turku-Pori) Finland. 61.32 N, 21.40 E	11 August 1969
	Shot by hunter	4 km SE Mersa Matruh Egypt, UAR. 31.21 N, 27.14 E	November 1969
R-65.386 Helsinki Museum	After first year	Polla, Hailucto, (Oulu), Finland. 64.58 N, 24.42 E	28 August 1969
	Killed	Bird Market, Alexandria Egypt, UAR. 31.13 N, 29.55 E	13 October 1969

PHYLLOSCOPUS TROCHILUS

R-65.487 Helsinki Museum	After first year	Polla, Hailucto, (Oulu), Finland. 64158 N, 24.42 E	30 August 1969
	Killed	Alexandria Bird Market Alexandria, Egypt, UAR 31.13 N, 29.55 E	13 October 1969
1.432.507 Stockholm	Fledgling	Enskar, Bulinge, Sodermanland, Sweden 58.41 N, 17.29 E	30 August 1969
	Netted by Bedouins; caught by Hassan Nagi; released at Ikingi wing, 66 mm.	Bahig, Egypt 30.56 N, 29.35 E Mariut, UAR: Fat, light; weight 8.5 gms;	20 September 1969
010-06246 Cairo Zoo		Ikingi Mariut, Egypt, UAR 31.00 N, 29.45 E	10 September 1969
	Found dead (Victim of traffic)	Matinlompolo, Pello, Finland 66.44 N, 24.19 E	27 June 1970

PHYLLOSCOPUS COLLYBITA

HC 27525 Jordan	Post-juvenile, Male	Azraq, Jordan 31.50 N, 36.50 E	24 April 1966
	Killed (lime- stick)	Paralimni, Cyprus 35.01 N, 33.59 E	8 April 1968

MUSCICAPA HYPOLEUCA

493.421 Paris Museum		Luz-Saint-Sauveur (Eastern Pyrenees), France 42.52 N, 0.00	17 August 1968
	"Flycatcher" (pre- sumed Ficedula hypoleuca), Wing- 79, weight 10.9; released at Ikingi Mariut, Cairo Zoo Band 010-03833. Female plumage.	Bahig, Egypt, UAR 30.56 N, 29.35 E	13 May 1969

MUSCICAPA STRIATA

020-02488 Cairo Zoo		Bahig, Egypt, UAR 30.56 N, 29.35 E	3 May 1967
	Shot by hunter	Allis Merias, Volos, Greece 39.22 N, 22/57 E	20 November 1969

OENANTHE OENANTHE

020-17565 Cairo Zoo	Female	Mersa Matruh, Egypt, UAR 31.21 N, 27.14 E	31 March 1970
	Fat-N, Wing-94 Shot by hunter	Bir Ramla, 15 km W of Salum Egypt. 31.31 N, 25.09 E	16 April 1970

OENANTHE HISPANICA

020-01225	Cairo Zoo		Bahig, Egypt, UAR 30.56 N, 29.35 E	6 April 1967
		Trapped	Bahig, Egypt, UAR 30.56 N, 29.35 E	September 1967

PHOENICURUS OCHRURUS

020-05125	Cairo Zoo	Female	Ikingi, Mariut, Egypt, UAR 31.00 N, 29.45 E	13 March 1969
		Released Ikingi Mariut, Egypt.	Bahig, Egypt, UAR 30.56 N, 29.35 E	3 March 1970
		Fat, light; wing, 81 mm; weight, 15.0 gm; female. Recaptured: Ikingi Mariut, Egypt. Killed by hunter		

PHOENICURUS PHOENICURUS

010-00632	Cairo Zoo		Bahig, Egypt, UAR 30.56 N, 29.35 E	5 April 1967
		Fat, light; wing- 79 mm; weight 13.7, Male-adult Released Ikingi Mariut, Egypt, UAR	Bahig, Egypt, UAR 30.56 N, 29.35 E	5 April 1970

010-00673	Cairo Zoo	Male-plumage	Bahig, Egypt, UAR 30.56 N, 29.35 E	6 April 1967
		Reported/Letter to Mohamed Abdel Rahim, 6/2/68	Zlitan, Lybia 32.28 N, 14.34 E	?

A28768	Tel-Aviv	Female	Elat, Israel 29.33 N, 34.57 E	30 April 1970
		Caught on lime- stick & killed	Paralimmi, Cyprus 35.02 N, 34.00 E	14 May 1970

LUSCINIA MEGARHYNCHOS

020-02059	Cairo Zoo		Bahig, Egypt, UAR 30.56 N, 29.35 E	19 April 1967
		Trapped	Bahig, Egypt, UAR	September 1967
		Previous recap- ture & release	30.56 N, 29.35 E Bahig, Egypt, UAR	19 April 1967
HK 62.824	Cyprus	Post-juvenile	Akrotiri, Cyprus 34.35 N, 32.57 E	13 October 1968
		Controlled	Eilat, Gulf of Aquaba, Israel. 29.34 N, 34.57 E	24 March 1969

LUSCINIA LUSCINIA

ZAA 8318 Riksmuseum Stockholm	Pullet	Kungshälla, Malmö, Skåne, Sweden 55.36 N, 13.05 E	21 June 1950
	Shot	Qalyubiya Province Egypt. 30.18 N, 31.18 E	27 March 1953
3.151.964 Stockholm	Adult	Siglefs, N or , Gotland Sweden. 57.06 N, 18.14 E	27 June 1968
	Shot by hunter	Burg el Arab, Egypt, UAR 30.54 N, 29.32 E	Sept-October 1969

TURDUS PHILOMELOS

050-0046 Cairo Zoo	Female	Bahig, Egypt, UAR 30.56 N, 29.35 E	26 March 1967
	Shot by hunter (Fat:light; wing: 117.5, faint buff tips second covert, ovary small, slightly granular.	Bahig, Egypt, UAR (Abdel Aziz). 30.56 N, 29.35 E	10 March 1970

CARDUELIS CHLORIS

BE 78268 Cyprus	First Year	Akrotiri, Cyprus 34.36 N, 32.57 E	8 November 1967
	Trapped & caged	Limassol, Cyprus 34.40 N, 33.02 E	11 February 1968
BE 78348 Cyprus	First Year	Akrotiri, Cyprus 34.36 N, 32.57 E	10 November 1967
	Taken on lime- stick.	Maroni Village (Larnaca), Cyprus. 34.55 N, 33.38 E	16 January 1968

FICEDULA HYPOLEUCA

26642 Jos Museum	Full-grown	Malamfatori, Bornu, Nigeria. ca. 13.00 N, 14.00 E	9 October 1966
	Killed, Lime stick	Paralimni, Cyprus 35.01 N, 33.59 E	ca. 1967

FICEDULA Sp.

010-01126 Cairo Zoo	Male-immature	Bahig, Egypt, UAR 30.56 N, 29.35 E	22 April 1967
	Trapped. Band given to NAMRU & returned to SI.	Bahig, Egypt, UAR 30.56 N, 29.35 E	13 October 1967
	Previous recapture:	Bahig, Egypt, UAR	22 April 1967

FICEDULA SP.

010-03384 Cairo Zoo	Maadi, Cairo, Egypt, UAR	27 April 1969
	29.58 N, 31.15 E	
	Reggio Calabria, Italy	20 October 1969

PASSER HISPANIOLENSIS

030-00071 Cairo Zoo	Male-plumage	Bahig, Egypt, UAR	13 April 1967
		30.56 N, 29.35 E	
		Sicas Savae, Camariotissa,	Fall 1967 or
		Samothraci, Greece	Spring 1968
B-4932 Tel-Aviv Univ.	Female	Elat, Israel	25 February 1969
		29.33 N, 34.57 E	
	Shot by hunter	Near Cairo, Egypt, UAR	Before July 1970
		30.03 N, 31.15 E	

ORIOLOUS ORIOLOUS

050-0078 Cairo Zoo	Female	Bahig, Egypt, UAR	6 May 1967
		30.56 N, 29.35 E	
	Shot	Darya El Shoef, Lebanon	25 April 1969
050-4518 Giza Zoo	Female-adult	Maadi, Cairo, Egypt, UAR	9 September 1968
		29.58 N, 31.15 E	
	Shot	Maadi, Cairo, Egypt, UAR	11 September 1968
		29.58 N, 31.15 E	

NEAR EASTERN RECOVERIES

Addendum - Mostly literature records of White Storks

SPECIES

Band number	Age and Sex Recovery remarks	Locality banded Locality recovered	Date banded Date recovered
ANAS CLYPEATA	Male Adult	Astrakhan Reserve, USSR	24 July, 1934
E-23482 Moskwa	(Works Bur. Bird- Banding, USSR)	46.22 N, 48.55 E Damietta, Egypt	January, 1937
E-108916 Moskwa	Male	31.27 N, 31.49 E	
	(Works Bur. Bird- Banding, USSR)	Lake Matveevo, USSR	21 July, 1949
		54.23 N, 77.21 E	
		nr. El Faiyum, Egypt	4 February, 1951
		30.08 N, 31.20 E	
E-222491 Moskwa	Male Adult	Astrakhan Reserve, USSR	2 August, 1952
	(Works Bur. Bird- Banding, USSR)	46.22 N, 48.55 E	
		Lake Mariut, Egypt	28 December, 1952
		31.11 N, 29.55 E	
CICONIA CICONIA			
B23 526 0		Lichtenfeld, Germany	16 July, 1930
	(Vogelzug, 1937, p. 199)	54.20 N, 20.15 E	
		40 km S of Cairo, Egypt	March, 1931
		30.2 N, 31.15 E	
B 24891 0		Baumgarth, West Prussia	19 July, 1931
	(Vogelzug, 1937, p. 204)	53.58 N, 19.23 E	
		Giza, Egypt	30 August, 1935
		29.58 N, 31.12 E	
208 178 0		Militsch, Germany	24 July, 1931
	Found (Vogelzug, 1937, p. 204)	51.32 N, 17.17 E	
		Talga, Egypt	10 June, 1937
		31.3 N, 31.31 E	
B 26 968 0		Langhofel, East Prussia	26 June, 1932
	Shot (Vogelzug, 1937, p. 203)	54.37 N, 20.52 E	
		Lake Burullus, Egypt	August, 1933
		31. E on the Coast	

CICONIA CICONIA

B 42 984 0		Szierandzen, Germany 54.49 N, 21.55 E	13 July, 1933
	Shot (<u>Vogelzug</u> , 1937, p. 202)	Dershai Delengat, Egypt 30.50 N, 30.30 E	16 May, 1934
B 33 548 0		Bruchhoff, Germany 53.32 N, 16.01 E	29 June, 1933
	Shot (<u>Vogelzug</u> , 1937, p. 203)	Sidi Salem, Egypt (Nile Delta)	mid-May, 1934
B 41 991 0		Liepz, Germany 53.44 N, 15.40 E	26 June, 1933
	Recovered (<u>Vogel-</u> <u>zug</u> , 1937, p. 202)	Nazlet el Dib, Egypt 29.6 N, 31.7 E	26 April, 1934
B 42 547 0		Felstow, Germany 54.32 N, 17.44 E	18 July, 1933
	(<u>Vogelzug</u> , 1937 p. 202)	Mansuria, Egypt 30.9 N, 15 km NW of Cairo	4 May, 1934
B 44 316 0		Wirtkallen, Germany near Insterburg	10 July, 1933
	Shot, (<u>Vogelzug</u> , 1937, p. 202)	Teh el Baroud, Egypt 30.54 N, 31.40 E	6 May, 1934
B 42 929 0		Thieslauken, Germany near Insterburg	22 June, 1933
	Recovered (<u>Vo-</u> <u>gelzug</u> , 1937, p. 198)	near Teefa, Egypt 23.40 N, 32.55	7 May, 1937
BB 1461 0		Lascheninken, Germany Insterburg district	6 July, 1934
	Shot (<u>Vogelzug</u> 1937, p. 203)	Barnoogi, Egypt app. 31 N, 31.30 E	26 May, 1935
BB 319 0		Tannowischken, Germany Insterburg district	2 July, 1934
	Found (<u>Vogelzug</u> , 1937, p. 203)	between Malaha & Esbet Egypt 31.12 N, 29.53 E	1 May, 1935
B 50 536 0		Wilkowe, Germany 51.32 N, 17.17 E	30 June, 1934
	Found (<u>Vogelzug</u> , 1937, p. 204)	near Tor, Sinai 28.15 N, 33.36 E	Oct., 1934

CICONIA CICONIA

211 282 0		Dorf Grabkow, Germany 51.47 N, 14.58 E	1 July, 1934
	Found dead (<u>Vogelzug</u> , 1937 p. 204)	Sinai-Kloster, Egypt 28.36 N, 33.58 E	Oct., 1934
BB 39 94 0		Lomnitz, Germany 50/50 N, 18.32 E	14 June, 1935
	Found (<u>Vogelzug</u> , 1937 p. 205)	West Sinai 30 N, 34 E.	14 Sept., 1935
BB 3965 0		Birkenfeld, Germany Kreuzberg district	13 June, 1935
	Found, (<u>Vogel-</u> <u>zug</u> , 1937 p. 204)	Wadi Gharandal, Sinai 29.20 N, 33.3 E	end March, 1936
BB 2773 0		Scheppetschen, East Prussia	23 June, 1935
	(<u>Vogelzug</u> , 1937	Mataana, hr.'s train fr. Luxor, Upper Egypt 25.43 N, 32.37 E	15 May, 1937
BB 1524 0		Kl. Albrechtsdorf, East Prussia	11 July, 1935
	(<u>Vogelzug</u> , 1937 p. 204	54.25 N, 23. 1 E Meet-el Amel, Egypt app. 30 N, 31.30 E	9 June, 1936
BB 28 52 0		Triaken, East Prussia Insterburg district	25 June, 1937
	Dead, (<u>Vogelzug</u> , 1937, p. 202)	Shanfash Markaz, Egypt Dakahlia Province	22 June, 1937
B 60771		Domane Cantnitz, Germany 53.30 N, 13.18 E	30 June, 1935
	Found dead (?) (<u>Vogelzug</u> , 1937 p. 201)	Mazraet El Orz, Egypt 25.26 N, 30.40 E	27 July, 1937
BB 2946 0		Drutschlauken, East Prussia, Germany	26 June, 1935
	Recovered, (<u>Vogelzug</u> , 1937 p. 201)	Tanta, Egypt 30.47 N, 30.59 E	16 June, 1937
BB 4071 0		Moesen, Schlesien, Germany	14 June, 1935
	Killed in troop of 4 storks, (<u>Vogelzug</u> , 1937 p. 200	50.26 N, 17.5 E Kostamna-Shellal, Egypt 24.5 N, 32.57 E	17 August, 1936

CICONIA CICONIA

B 54 928 0		Piastental, Germany 50.52 N, 17.28 E	9 July, 1935
	(<u>Vogelzug</u> , 1937 p. 199)	Gurta (?) Upper Egypt	app. 25 Aug., 1935
B 45 948 0		Klein-Methling, Germany 53.58 N, 12.43 E	16 June, 1936
	Recovered (<u>Vo- gelzug</u> , 1937, p. 201)	near Aswan, Egypt 24.5 N, 32.57 E	26 June, 1937
B 62 834 0		Prutzke, Germany 52.22 N, 12.38 E	15 June, 1936
	Found freshly dead (<u>Vogelzug</u> , 1937, p. 202)	Ezbet Mazhar Pacha (?) Dakahlia Province, Egypt	11 August, 1937
B 53662 0		Erdmannen, East Prussia 53.38 N, 21.48 E	23 June, 1936
	(<u>Vogelzug</u> , 1937 p. 204)	On Bogma, Sinai 28.20 N, 33.40 E	27 August, 1936
BB 63 10 0		Gravenort, East Prussia Insterburg district	3 July, 1936
	Dead, (<u>Vogelzug</u> 1937, p. 203)	Ras el Khalig, Egypt 31.15 N, 31.37 E	13 June, 1937
B 63 223 0		Gut Wall, Rugen, Germany 54.28 N, 13.17 E	7 July, 1936
	(<u>Vogelzug</u> , 1937, p. 203)	35 km S of Port Said on Suez Canal, Egypt 30.56 N, 32.20 E	1 June, 1937
B 61 982 0		Heiligenbeil-Rosenberg, East Prussia 54.28 N, 19.57 E	28 June, 1936
	Found, (<u>Vogelzug</u> , 1937, p. 202)	Manshiet Battash, Egypt 31.52 N, 31.27 E	6 July, 1937
B 48 281 0		nr. Tauroggen, Germany 55.16 N, 22.17 E	14 July, 1936
	(<u>Vogelzug</u> , 1937 p. 202)	Tanta, Egypt 30.47 N, 30.59 E	17 July 1937
B 48 236 0		Rudienen, Germany 55.21 N, 21.29 E	11 July, 1936
	Found, (<u>Vogel- zug</u> , 1937, p. 202)	On Dama, Egypt 26.30 N, 31.40 E	13 May, 1937

CICONIA CICONIA

BB 6567 0

Found, (Vogelzug,
1937, p. 202Schacken, East Prussia
Germany [=Latvia]
Qina Province, Egypt
26.10 N, 32.44 E

June, 1936

8 May, 1937

COTURNIX COTURNIX

C 2416 Tel Aviv
Univ. IsraelAlive in Alex.
Bird Mkt., but
scheduled to be
killed.Elat, Israel
29.33 N, 34.57 E
Alexandria, Egypt
31.12 N, 29.54 E

9 April, 1969

Autumn, 1970

060-03346
Cairo Zoo

Killed

Mersa Matruh, Egypt
31.21 N, 27.14 E
Obthcharovo, Bulgaria
43.13 N, 27.08 E

20 Oct., 1969

5 Sept., 1970

LARUS FUSCUS PULLUS

H-61.459
Helsinki Mus.

Found dead

Rodharun, Finland
22.15 E
Max, Alexandria,
Egypt, UAR
31.12 N, 29.54 E

16 June, 1965

10 May, 1967

MOTACILLA ALBA

ZOD 1252 Stock-
holm

Adult

Caught & released
(COS Bull.)
no ringOttenby, Sweden
56.12 N, 16.24 E
At sea btween
Famagusta & Pt. Said,
Egypt

1 Sept., 1950

22 Oct., 1950

1.494.704
StockholmKilled before
3 Nov. 1970Klippan, Sweden
56.08 N, 13.07 E
Alexandria, Egypt
31.12 N, 29.54 E

15 Sept., 1970

Autumn, 1970

LANIUS COLLURIO

H 85 437

(Vogelzug, 1936,
p. 127Enkheim, Germany
50.9 N, 8.46 E
El-Agami, Egypt
31.12 N, 29.53 E

18 June, 1929

end Sept., 1929

LANIUS COLLURIO

H 71886		Ober-Baumgarten, Germany 50.53 N, 16.10 E	7 July, 1930 return 23 July, 1931
	(Vogelzug, 1936 p. 126)	Gabbary nr. Alex., Egypt 31.12 N, 20.53 E	26 August, 1932
H 861 449 A		Heuhaldensleben, Germany 52.17 N, 11.25 E	17 July, 1932
	(Vogelzug, 1936, p. 126)	Rashid-Rosetta, Egypt 31.24 N, 30.24 E	6 Sept., 1933
R 202 901 G		Insterburg, Germany [=Latvia]	9 July, 1933
	(Vogelzug, 1936 p. 127)	Nr. Lake Burullus- Rosetta, Egypt 31.22 N, 30.26 E	24 Sept., 1933
H 741 709		Kockwitz-Halle, Germany 51.28 N, 11.58 E	25 April, 1933
	(Vogelzug, 1936 p. 126)	In Mariout desert, 60 km SW of Alex., Egypt	8 Sept., 1935
H 71-854		Ober-Baumgarten, Germany 50.53 N, 10.13 E	21 August, 1933
	(Vogelzug, 1936, p. 127)	Rosetta, Egypt 31.24 N, 30.24 E	16 Sept., 1933
H 851 137		Wetzlar-Hermannstein, Germany	11 Sept., 1933
	(Vogelzug, 1936 p. 127)	Burg-el-Arab, near Alexandria, Egypt 31.12 N, 29.53 E	14 Oct. 1933
R 179 145 F		Langebruck-Dresden, Germany 51.8 N, 13.51 E	24 June, 1934
	(Vogelzug, 1936 p. 126)	Mariout, west of Alex, Egypt 31.7 N, 29.24 E	8 Sept., 1934
BJ 51.066		Kuscenneti, Turkey 40.14 N, 28.03 E	11 August, 1969
British Mus.		Alexandria, Egypt 31.13 N, 29.55 E	14 August, 1970
	Killed (Cambridge Exped.)		

SYLVIA ATRICAPILLA

C 674 Cairo	Adult male	Fayid, Egypt	8 April, 1951
	(Pub. COS Bull.)	30.17 N, 32.19 E	
		Famagusta, Cyprus	11 Oct., 1951
		35.03 N, 33.57 E	

SYLVIA CURRUCA

HV 95.410		Holkham, England	9 May, 1970
British Museum	Adult	52.58 N, 0.48 E	
	Dead	nr. Burg el Arab,	Autumn, 1970
		Egypt	
		30.55 N, 29.32 E	

PHYLLOSCOPUS TROCHILUS

N-09.828		Norr fjarden, Finland	4 August, 1970
Helsinki Museum	after 1st yr.	62.17 N, 21.24 E	
	Found dead	Alexandria, Egypt	befr. 3 Nov., 1970
		31.12 N, 29.54 E	

PHYLLOSCOPUS COLLYBITA

1.543.533	fullgrown	Sundsvall, Sweden	28 August, 1970
Stockholm		62.23 N, 17.18 E	
	Killed	Alexandria, Egypt	Autumn, 1970
	Alex. Bird Mkt.	31.12 N, 29.54 E	

PHYLLOSCOPUS BONELLI ORIENTALIS

Z 1440 Tel Aviv		Maagen Michael, Israel	21 March, 1970
Univ. Israel		32.33 N, 34.55 E	
		near Kom Ombo, Egypt	Rep. 15 June,
		24.28 N, 32.57 E	1970

CYPRUS RECOVERIES

PREVIOUSLY REPORTED TO CYPRUS ORNITHOLOGICAL SOCIETY

SPECIES

Band number	Age and Sex Recovery remarks	Locality banded Locality recovered	Date banded Date recovered
NYCTICORAX NYCTICORAX	Pullus	Kisbalaton, Hungary 46.40 N. 17.15 E	14 June 1929
54164 Budapest	Dead (COS Bull.)	Nicosia, Cyprus 35.10 N. 33.23 E	18 May 1930
170438 Budapest	Pullus	Kisbalaton, Hungary 46.40 N, 17.15 E	10 June 1953
	Killed (COS Bull.)	"Cyprus"	13 September 1953
G 39357 Radolfzell	Pullus	Marchegg Sanseindorf. 48.17 N, 16.55 E	6 June 1963
	Found dead (COS Bull.)	Cape Andreas, Cyprus 35.40 N, 34.35 E	9 April 1967
PLEGADIS FALCINELLUS			
D 51458 Moskwa	Pullus	Kizil Agach Reserve Lenkoran (Azerbaydahn)	3 June 1954
	(COS Bull.)	USSR. 39.00 N, 48.50 E	
	Recovery details not known	Morphou Bay, Cyprus 35.10 N, 32.55 E	30 March 1961
ANAS CRECCA			
E 75556 Moskwa	Adult	Astrakhan Reserve, USSR 46.12 N, 48.55 E	28 July 1948
	(COS Bull.)	Livadhia (Larnaca), Cyprus. 34.57 N, 33.38 E	February 1948
2032002 British Museum	Full-grown Male	near Akrotiri, Cyprus 34.36 N, 32.56 E	22 November 1959
	(COS Bull.)	Ardabil, Iran 38.15 N, 48.22 E	6 March 1960
2032008 British Museum	1st Winter Male	Akrotiri Sewage Outfall Cyprus.	5 December 1959
	Killed (COS Bull.)	Akrotire Sewage Outfall Cyprus.	5 December 1959
2032005 British Museum	1st Winter	Akrotiri, Cyprus 40.28 N, 25.28 E	22 November 1959
	Killed (COS Bull.)	Akrotiri, Cyprus 40.28 N, 25.28 E	Autumn/Winter 1962

ANAS ACUTA

E 133007 Moskwa	Adult	Astrakhan Reserve, USSR. 46.12 N, 48.55 E	4 August 1949
	Killed (COS Bull.)	Ktima (Paphos), Cyprus 34.46 N, 32.25 E	17 March 1951

SPATULA CLYPEATA

E 74278 Moskwa	Adult Male	Astrakhan Reserve, USSR. 46.12 N, 48.55 E	8 August 1940
	Killed (COS Bull.)	Kalokhorio (Larnaca) Cyprus 34.55 N, 33.32 E	17 March 1951

PERNIS APIVORUS

(Written label)		Aseevka (dist. Kuzetsk), USSR. Prob. 53.11 N, 46.25 E	12 September 1929
	Killed (COS Bull.) (Ring 2:85-86, 1958)	Karavas (Kyrenia), Cyprus 35.20 N, 33.11 E	10 October 1929

MILVUS MIGRANS

C 63775 Rossitten	Juvenile	Grabow (Germany), Poland 53.00 N, 14.23 E	11 June 1938
	Killed (COS Bull.)	Famagusta, Cyprus 35.09 N, 33.59 E	21 September 1938

FALCO ELEONORAE

3062516 British Museum	Pullus	Akrotiri (Limassol), Cyprus. 34.36 N, 32.56 E	17 September 1961
	(COS Bull.)	Zakaki (5 m from Akrotiri, Limassol), Cyprus	12 September 1962

GYPS. FULVUS

z17485 British Museum	Pullus	Episkopi, Cyprus 34.40 N, 32.55 E	15 May 1966
	Dead (COS Bull.)	Episkopi, Cyprus 34.40 N, 32.55 E	13 July 1966
z17483 British Museum	Pullus	Episkopi, Cyprus 34.40 N, 32.55 E	16 May 1965
	Injured (COS Bull.)	Athna (Famagusta), Cyprus. 35.07 N, 33.56 E	25 September 1966

FALCO ELEONORAE

306.2515 British Museum	Pullus	Akrotiri, Cyprus 40.28 N, 25.28 E	17 September 1961
	(COS Bull.)	Larnaca, Cyprus 34.55 N, 33.38 E	16 May 1962

FALCO ELEONORAE

309 1453 British Museum	Pullus	Akrotiri, Cyprus 40.28 N, 25.28 E Ankazobe (Tananarive), Madagascar. 18.20 S, 47.07 E	22 September 1962
	(COS Bull.)		30 November 1962
3091466 British Museum	Pullus	Akrotiri, Cyprus 40.28 N, 25.28 E	22 September 1963
	Shot (COS Bull.)	Akrotiri, Cyprus 40.28 N, 25.28 E	5 May 1966
3091485 British Museum	Pullus/2	Akrotiri, Cyprus 40.28 N, 25.28 E	20 September 1964
	Shot (COS Bull.)	West of Limassol, Cyprus	9 September 1966

FALCO TINNUNCULUS

3091490 British Museum	Pullus/5	Akrotiri, Cyprus 40.28 N, 25.28 E	3 June 1965
	Shot (COS Bull.)	Akrotiri, Cyprus 40.28 N, 25.28 E	26 December 1966
3091489 British Museum	Pullus/5	Akrotiri, Cyprus 40.28 N, 25.28 E	3 June 1965
	Shot (COS Bull.)	Sotira (Famagusta), Cyprus 35.07 N, 33.56 E	6 September 1965

CHARADRIUS HIATICULA

V23036 British Museum	1st Winter	Akrotiri, Cyprus 40.28 N, 25.28 E	6 November 1959
	Killed (COS Bull.)	Ahezitou Res., Famagusta Cyprus. 35.07 N, 33.56 E	25 December 1960

CALIDRIS MINUTA

V 23938 British Museum	Adult	Akrotiri Salt Lake, Cyprus 40.28 N, 25.28 E	6 November 1959
	Killed (COS Bull.)	Near Limassol, Cyprus 34.41 N, 33.05 E	2 February 1960

CALIDRIS ALPINA

V 23055 British Museum	Full-grown	Akrotiri Salt Lake, Cyprus 40.28 N, 25.28 E	21 November 1959
	Killed (COS Bull.)	Akrotiri Salt Lake, Cyprus 40.28 N, 25.28 E	28 January 1960

CROCETHIA ALBA

V 23053 British Museum	Full-grown	Akrotiri Salt Lake, Cyprus 40.28 N, 25.28 E	21 November 1959
	Killed (COS Bull.)	Akrotiri Salt Lake, Cyprus 40.28 N, 25.28 E	29 November 1959

PHILOMACHUS PUGNAX

V23121 British Museum	Full-grown	Akrotiri (Limassol), Cyprus 40.28 N, 25.28 E	1 September 1962
	Killed (COS Bull.)	Akrotiri, Cyprus 40.28 N, 25.28 E	7 September 1962

LARUS MELANOCEPHALUS

E 127972 Moskwa	Juvenile	Orlov Island, USSR 46.17 N, 31.45 E	3 July 1948
	(COS Bull.)	Larnaca, Cyprus	1 August 1948
E 190262 Moska	Juvenile	Orlov Island, USSR 46.17 N, 31.45 E	29 June 1950
	(COS Bull.)	No. Coast of Cyprus 34.30 E	25 March 1951

LARUS RIDIBUNDUS

E 133993 Moskwa	Pullus	Lake Kievo, USSR 56.02 N, 37.30 E	6 June 1948
	(COS Bull.)	Zakaki (Limassol), Cyprus 34.39 N, 33.00 E	29 December 1948
N 46172 Moscow	Juvenile	Ivanou Region, Sokolskoe, USSR. 59.10 N, 43.10 E	12 June 1959
	(COS Bull.)	Cyprus	Autumn, 1959

STERNA SANDVICENSIS

F 133144	Pullus	Orlov Island, USSR 46.17 N, 31.45 E	23 June 1952
	(COS Bull.)	"Cyprus"	February 1953
F 163244 Moskwa	Pullus?	Orlov Island, USSR 46.17 N, 31.45 E	5 July 1952
	(COS Bull.)	Larnaca, Cyprus 34.55 N, 33.39 E	19 February 1953

MEROPS APIASTER

D 8771 Sofia	Adult	Sindel, Bulgaria 43.07 N, 27.36 E	14 June 1933
	Caught, or caught & released w/o ring (COS Bull.)	Larnaca, Cyprus 34.55 N, 33.39 E	24 September 1934

MEROPS APIASTER

F 14988 Moskwa	Juvenile	Yorki, USSR 48.53 N, 30.43 E	21 July 1934
	(COS Bull.)	Larnaca, Cyprus 34.55 N, 33.39 E	25 September 1934
E 22929 Sofia	Pullus	Kuzina, Bulgaria 43.17 N, 25.38 E	9 July 1935
	Caught, or caught & released w/o ring (COS Bull.)	Pakhna (Limassol), Cyprus 34.46 N, 32.38 E	21 September 1935
F 20719 Moskwa		Kanev, USSR 40.44 N, 31.02 E	3 August 1940
	(COS Bull.)	Ormidhia (Larnaca), Cyprus 34.59 N, 33.47 E.	26 September 1940

ORACIAS GARRULUS

112152 Riga	Pullus	Nauleni (Latvia), USSR 55.58 N, 27.40 E	16 July 1938
	(COS Bull.)	Ormidhia (Larnaca), Cyprus 34.59 N, 33.19 E	26 September 1938

ALERIDA CRISTATA

E 11358 British Museum	Full-grown	Nicosia, Cyprus 35.10 N, 33.23 E	11 January 1959
	Killed (COS Bull.)	Nicosia, Cyprus 35.10 N, 33.23 E	31 January 1960
E 11370 British Museum	Full-grown	RAF Nicosia, Cyprus 35.13 N, 33.05 E	13 January 1959
	Killed (COS Bull.)	Nicosia, Cyprus 35.10 N, 33.23 E	26 October 1960

LULLULA ARBOREA

141632 Budapest	Adult	Budaors, Hungary 47.26 N, 18.59 E	2 May 1948
	Killed (COS Bull.)	Lefkara (Larnaca), Cyprus 34.51 N, 33.19 E	27 February 1957

HIRUNDO RUSTICA

E 11.139 British Museum	Caught & released with ring (COS Bull.)	Morphou Bay, Cyprus 35.10 N, 32.50 E	14 August 1957
		Morphou Bay, Cyprus 35.10 N, 32.50 E	21 May 1962

HIRUNDO RUSTICA

E 11133 British Museum	Full-grown	Pendalia, Cyprus 34.41 N, 32.27 E	10 April 1957
	Caught (COS Bull.)	W. Tsadha, Cyprus 34.50 N, 32.28 E	10 April 1959
E 11597 British Museum	Juvenile	Nicosia, Cyprus 35.13 N, 33.22 E	13 July 1961
	Found dying (COS Bull.)	Nicosia, Cyprus 35.13 N, 33.22 E	28 June 1964
AE 55567 British Museum	Juvenile	Akrotiri, Cyprus 40.28 N, 25.28 E	30 July 1962
	? (COS Bull.)	Limassol, Cyprus 34.41 N, 33.05 E	14 June 1963

DELICHON URBICA

E 11021 British Museum	Adult	Pano Platres, Cyprus 34.53 N, 32.52 E	1 May 1957
	Killed (COS Bull.)	Pano Platres, Cyprus 34.53 N, 32.52 E	

ANTHUS PRATENSIS

E11462 British Museum	First year	RAF Nicosia, Cyprus 35.13 N, 33.22 E	7 January 1959
	Killed (COS Bull.)	RAF Nicosia, Cyprus 35.12 N, 33.22 E	8 March 1960

MOTACILLA FLAVA

AE 55768 British Museum	First winter	Episkopi, Cyprus 34.40 N, 32.55 E	7 October 1963
	(COS Bull.)	Mersin, Silifko, Turkey 36.22 N, 33.56 E	28 September 1964

MOTACILLA CINEREA

AE 55784 British Museum	Full-grown	Akrotiri, Cyprus 40.28 N, 25.28 E	22 October 1963
	Killed (COS Bull.)	Akrotiri, Cyprus 40.28 N, 25.28 E	5 November 1963

MOTACILLA ALBA

ZO 1073 Stockholm	Adult	Ottenby (Oland Id.) Sweden 56.12 N, 16.24 E	9 September 1938
	Killed (COS Bull.)	Larnaca, Cyprus 34.55 N, 33.39 E	3 November 1938

MOTACILLA ALBA

ZOD 1252 Stockholm	Adult	Ottenby (Oland Id.) Sweden 56.12 N, 16.24 E	1 September 1950
	Caught or caught & released w/o ring. (COS Bull.)	At sea between Famagusta and Port Said	22 October 1950
ZAH 6133 Stockholm	Pullus	Voa, Sweden 61.49 N, 17.23 E	9 July 1952
	(COS Bull.)	Limassol, Cyprus 34.41 N, 33.05 E	12 April 1953
2.025.467 Stockholm	Juvenile	Stallarolmon, Malaron, Sweden.	1 July 1961
	(COS Bull.)	? ? Cyprus	27 February 1963

LANIUS COLLURIO

F 359777 Rossitten	Pullus	Barzdorf (Germany) Poland 51.00 N, 16.24 E	Summer 1939
	(COS Bull.)	Tarvros, Famagusta, Cyprus 35.23 N, 34.05 E	14 September 1939
2000 Ljubljana		Svetje Medvode, Yugoslavia 46.08 N, 14.22 E	16 July 1939
	(COS Bull.)	Paralimni, Famagusta, Cyprus 35.01 N, 33.59 E	ca 3 May 1940

LANIUS NUBICUS

41688 R British Museum	Full-grown	Troodos, Cyprus 34.55 N, 32.53 E	30 June 1963
	Caught or caught & released w/o ring.	Akbiyeh, 12 m SSW Sidon, Lebanon. ca 33.30 N, 35.16 E	13 April 1964

ACROCEPHALUS PALUSTRIS

4B 6471 Bruxelles	Pullus	Ieper, Belgium 50.51 N, 2.53 E	21 June 1952
	Found dead (COS Bull.)	Famagusta, Cyprus 35.09 N, 33.59 E	October 1952
801 27136 Helgoland	Adult	Moischt nr. Marburg (Hessen) Germany. 50.46 N, 8.49 E	9 July 1961
	Killed (COS Bull.)	Cyprus (locality unknown)	ca 30 Oct. 1961
H 806811 Radolfzell	Full-grown	Bretzenheim nr. Bad Krenznach(Rheinland Pfalz) Germany. 49.53 N, 07.54 E	28 July 1962
	Lined (COS Bull.)	Famagusta, Cyprus 35.07 N, 33.57 E	10 Jan or 10 Feb 1966

ACROCEPHALUS PALUSTRIS

H 146062 Warsaw	Juvenile	Zagarow (Slupka) Poland 54.50 N, 17.00 E	22 July 1964
	(COS Bull.)	Famagusta, Cyprus 35.02 N, 34.00 E	Last week/Aug '64

HIPPOLAIS PALLIDA

J 14301 British Museum	Adult	Troodos (ca 6000 ft above sea level) Cyprus 35.16 E	21 May 1963
	(COS Bull.) Controlled	Troodos, Cyprus 34.55 N, 35.16 E	21 June 1964

SYLVIA BORIN

8085637	Pullus	Burg, Germany 52.17 N, 11.52 E	30 May 1934
	Caught or caught & released w/o ring. (COS Bull.)	Famagusta, Cyprus 35.09 N, 22.59 E	16 October 1954 (sic)
898140 A Helgoland	Juvenile	Dalbersdorf (Germany) Poland 51.12 N, 17.44 E	27 June 1933
	Killed (COS Bull.)	Larnaca, Cyprus 34.55 N, 33.39 E	26 August 1933

SYLVIA ATRICAPILLA

49434 Budapest	Pullus	Nagylozs, Hungary 47.33 N, 16.43 E	3 June 1929
	Killed (COS Bull.)	Hadjichtori (Limassol) Cyprus. 34.54 N, 32.56 E	18 September 1930
C 3871 Stockholm	Juvenile	Ulriksdal, Sweden 59.23 N, 18.00 E	1 August 1931
	(COS Bull.) Caught or caught & released w/o ring.	Ayios Theodoros (Larnaca) Cyprus. 34.48 N, 33.24 E	5 November 1932
869093 A Helgoland	Adult	Gimmel (Germany) Poland 51.13 N, 17.35 E	30 August 1932
	Killed (COS Bull.)	Paralimni (Famagusta) Cyprus. 35.01 N, 33.59 E	10 September 1932
H 3929 Lotos	Juvenile	Troppau, Czechoslovakia 49.57 N, 17.54 E	27 June 1933
	(COS Bull.)	Nicosia, Cyprus 35.10 N, 33.23 E	10 October 1934

SYLVIA ATRICAPILLA

8022101 Helgoland	Pullus	Barzdorf (Germany) Poland 51.00 N, 16.28 E	6 June 1933
	Caught or caught & released w/o ring. (COS Bull.)	Athna (Famagusta) Cyprus 35.44 N, 33.07 E	before 22 Sept 1934
9032860 Helgoland	Adult	Helgoland, Germany 54.11 N, 7.55 E	14 September 1934
	Killed (COS Bull.)	Ayios Theodoros (Famagusta) Cyprus. 35.21 N, 34.02 E	1 October 1937
G 237308 Rossitten	Juvenile	Ulmenhorst (Rossitten) USSR 55.09 N, 20.52 E	10 October 1933
	(COS Bull.)	Khirokitia (Larnaca), Cyprus. 34.47 N, 33.22 E	10 October 1934
E 8873 Praha	Adult	Pikovice near Saz, Czechoslovakia. 49.53 N, 14.26 E	10 September 1936
	Found dead (COS Bull.)	Maroni (Larnaca) Cyprus 34.45 N, 33.21 E	18 October 1938
G 404492 Rossitten	Juvenile	Ulmenhorst (Rossitten) USSR 55.09 N, 20.52 E	12 September 1937
	Killed (COS Bull.)	Dhali (Nicosia) Cyprus 35.01 N, 33.25 E	ca 19 Oct 1937
G 404709 Rossitten	Adult	Ulmenhorst (Rossitten) USSR 55.09 N, 20.52 E	19 September 1937
	Killed (COS Bull.)	Ayios Theodoros (Larnaca) Cyprus. 34.48 N, 33.24 E	6 November 1937
A 13322 Goteborg	Juvenile	Ekeby, Sweden 56.01 N, 12.58 E	10 June 1937
	Caught or caught & released w/o ring. (COS Bull.)	Kophinou (Larnaca) Cyprus 34.50 N, 33.24 E	4 October 1937
M20071 Praha	Juvenile	Dubi Bustehrad, Czechoslovakia 50.10 N, 14.09 E	21 June 1941
	Killed (COS Bull.)	Anglisidhes (Larnaca), Cyprus 34.50 N, 33.28 E	31 October 1941
M 33261 Praha	Pullus	Vrapice, Czechoslovakia 50.10 N, 14.10 E	2 June 1942
	Killed (COS Bull.)	Ayios Theodoros (Larnaca) Cyprus. 34.48 N, 33.24 E	6 October 1943
M 64929 Praha	Adult	Turnov, Czechoslovakia 50.36 N, 15.10 E	27 July 1943
	Killed (COS Bull.)	Anaphotia (Larnaca) Cyprus 34.48 N, 33.28 E	1 October 1946

SYLVIA sp. (ATRICAPILLA?)

134391 Riga		Near Kuldiga, Latvia	12 July 1943
		56.57 N, 22.00 E	
	(COS Bull.)	Pyrgha (Larnaca) Cyprus	9 November 1943
		34.50 N, 33.25 E	

SYLVIA ATRICAPILLA

A 44375 Goteborg	Juvenile Caught or caught & released w/o ring. (COS Bull.)	Henriksdorp, Sweden 55.58 N, 13.38 E Kornos (Larnaca) Cyprus 34.56 N, 33.24 E	6 June 1945 25 October 1945
M 143374 Praha	Pullus (COS Bull.)	Zlin, Czechoslovakia 49.13 N, 17.46 E Mazotos (Larnaca) Cyprus 34.48 N, 33.29 E	31 July 1948 ca 1 October 1948
M 183206 Praha	Adult (COS Bull.)	Caslav, Czechoslovakia 49.56 N, 15.20 E Limassol, Cyprus 34.41 N, 33.05 E	17 August 1951 ca 22 October 1951
C 674 Cairo	Adult male (Ringing date orig. reported as 7/8/51 from Egypt; corrected to 8/4/51 by Col. Brownlow from original records) (COS Bull.)	Fayid, Egypt 30.17 N, 32.19 E Dherynia (Famagusta) Cyprus 35.03 N, 33.57 E	8 April 1951 11 October 1951
Z 84636 Praha		Bela Rohovladova Czechoslovakia. 50.06 N, 15.36 E Near Famagusta, Cyprus 35.09 N, 33.59 E	26 July 1952 October 1952
216596 Moskwa	Pullus (COS Bull.)	Moscow, USSR 55.45 N, 37.42 E Famagusta, Cyprus 35.09 N, 33.59 E	23 June 1952 October 1952
M 215929 Praha	Pullus (COS Bull.)	Plzen, Czechoslovakia 49.45 N, 13.23 E Near Famagusta, Cyprus 35.09 N, 33.59 E	2 June 1952 October 1952
M 170567 Praha		Litvinov (Most) Czechoslovakia. 50.35 N, 13.36 E Perocherie, Nicosia, Cyprus 35.10 N, 33.20 E	10 August 1953 1 October 1953

SYLVIA ATRICAPILLA

G 55 1358 Varsovia	Pullus Caught or caught & released w/o ring. (COS Bull.)	Rydzyzna, Poland 51.48 N, 16.11 E Livadhia (Larnaca), Cyprus 34.57 N, 33.38 E	9 June 1957 ca 16 Sept 1957
66560 Madartani		Janoshegy, near Budapest Hungary. Kato Paphos, Cyprus 43.42 N, 32.24 E	3 June 1957 14 April 1959
M 286299 Praha		Plzen, Czechoslovakia 49.45 N, 13.23 E Yialousa (Famagusta dist.) Cyprus. 35.31 N, 34.12 E	28 August 1959 5 May 1960
908622 Moskwa	Juvenile Killed. (COS Bull.)	Rybatschi (Kaliningrad) USSR. 55.11 N, 20.44 E Paralimni (Famagusta dist.) Cyprus. 35.02 N, 34.00 E	17 July 1960 5 October 1960
M 399114 Praha	Pullus (COS Bull.)	Zleby (Caslav) Czechoslovakia 49.54 N, 15.30 E Near Famagusta, Cyprus 37.07 N, 33.57 E	2 July 1961 ca 30 October 1961
H32547 Varsovia	Full-grown Female Lined (COS Bull.)	Mierzigu Wislana (Gdansk) Poland. 54.19 N, 19.14 E Khirokitia, Near Larnaca Cyprus. 34.47 N, 32.21 E	21 August 1962 1 September 1964
P93329 Helsinki	Adult male (COS Bull.)	Jurmo, Korpo, Finland 59.50 N, 21.37 E Nicosia, Cyprus 35.10 N, 33.22 E	20 May 1963 January 1966
VIK.G.R.MUGRUM 2.117.955	Juvenile Male Killed (COS Bull.)	Ottenby, Sweden 56.12 N, 16.24 E Paralimni, Cyprus 35.09 N, 22.59 E	17 September 1963 1963 early November /
2121435 Stockholm	Juvenile Lined (COS Bull.)	Ytterby (Bohusland), Sweden 57.52 N, 11.55 E Ayios Amvrosis, Kividhes, Cyprus. 34.46 N, 32.51 E	10 August 1963 ca 10 Jan 1964
P110336 Helsinki	Adult Female Lined (COS Bull.)	Jurmo, Korpo, Finland 59.50 N, 21.37 E Near Paralimni, Cyprus 35.02 N, 34.00 E	29 May 1964 24 October 1964

SYLVIA ATRICAPILLA

AN26676 British Museum	Adult Female	Happy Valley, Episkopi Cyprus. 34.40 N, 32.55 E	22 March 1964
	Controlled	Happy Valley, Episkopi Cyprus. 34.40 N, 32.55 E	16 January 1965
8001-7344 Hiddensee	Pullus	Leutzsch, Leipzig, Germany 51.20 N, 12.18 E	18 June 1964
	Limed	Paralimni, Cyprus 35.02 N, 34.00 E	24 October 1964
216.7462 Stockholm	Adult Male	Kvismare, Norrlyas (Narhe) Sweden. 59.11 N, 15.24 E	4 August 1964
	Limed	Alaminos Village, Larnaca Cyprus. 34.55 N, 33.40 E	17 October 1965
2199381 Stockholm	Full-grown Male	Skanor (Skane), Sweden 55.25 N, 12.50 E	11 September 1965
	Limed	Near Larnaca, Cyprus 34.55 N, 33.39 E	15 November 1965
H 137343 Warsaw	Post-juvenile Male	Nowy Pasleka, near Olsztyn Poland. 54.26 N, 19.45 E	20 August 1966
	Trapped and Killed	Alaminos, near Larnaca Cyprus. 34.48 N, 33.26 E	3 October 1966
H.137343 Warsaw	Post-juvenile Male	Nowy Pasleka, near Olsztyn Poland. 54.26 N, 19.45 E	20 August 1966
	Trapped and Killed	Alaminos, near Larnaca Cyprus. 34.48 N, 33.26 E	3 October 1966

SYLVIA COMMUNIS

134391 Riga	Juvenile	Kuldiga, Latvia 56.58 N, 21.58 E	12 July 1943
	Recovered	Pyrgha, Larnaca, Cyprus 34.55 N, 33.25 E	9 November 1943

SYLVIA CURRUCA

985500 Copenhagen	Juvenile	Amager, Denmark 55.33 N, 12.36 E	10 August 1953
		Sotira, Famagusta, Cyprus 35.02 N, 33.57 E	ca 22 Sept 1953
CSR N 59486 Praha		Albrechtice, Prov. Lansktoun Czechoslovakia. 49.55 N, 16.39 E	3 June 1960
		Maroni District, Larnaca, Cyprus. 34.54 N, 33.39 E	3 October 1960

SYLVIA CURRUCA

K 93631 Helsinki	Adult	Rukkoila Hauho, Finland 61.13 N, 24.37 E	10 July 1961
	Unknown	Cyprus (locality unknown)	ca 18 Sept 1961
AE 55766 British Museum	Full-grown	Akrotiri, Cyprus 40.28 N, 25.28 E	5 October 1963
	Killed	Zakaki, Limassol, Cyprus 34.41 N, 33.05 E	23 October 1963
1.144.646 Stockholm	Post-juvenile	Ottenby (Oland Id.), Sweden 56.12 N, 16.24 E	21 August 1963
	Caught on limestick and killed	Paralimni, Famagusta Cyprus. 35.01 N, 33.59 E	30 September 1967
1.144 646 Stockholm	Post-juvenile	Ottenby, Oland, Sweden 56.12 N, 16.24 E	21 August 1963
	Caught on Limestick and killed	Paralimni, Famagusta, Cyprus. 35.01 N, 33.59 E	30 September 1967

SYLVIA MELANOCEPHALA

AE 55722 British Museum	1st Winter Female	Akrotiri, Cyprus 40.28 N, 25.28 E	12 December 1962
	Controlled	Akrotiri, Cyprus 40.28 N, 25.28 E	1 January 1964
AN 26776 British Museum	Adult Male	Akrotiri, Cyprus 40.28 N, 25.28 E	29 November 1964
	Controlled	Limassol, Cyprus 34.41 N, 33.05 E	2 January 1965
J 14710 British Museum	Full-grown Female	Happy Valley, Episkopi Cyprus. 34.40 N, 32.55 E	9 February 1964
	Controlled	Happy Valley, Episkopi Cyprus. 34.40 N, 32.55 E	16 January 1965

SYLVIA CONSPICILLATA

J 14247 British Museum	Adult Female	RAF Nicosia, Cyprus 35.13 N, 33.22 E	12 April 1961
	Killed	Makeodonatissa area, ca 2 m SW of RAF Nicosia Cyprus. 35.13 N, 32.55 E	28 May 1961

PHYLLOSCOPUS TROCHILUS

AN 26730 British Museum	Adult	Akrotiri, Cyprus 40.28 N, 25.28 E	16 September 1964
	Dead	Limassol (ca 7 m NE) Cyprus. 34.41 N, 33.05 E	18 September 1964

PHYLLOSCOPUS COLLYBITA

G 354566 Rossitten	Juvenile	Munsterburg (Germany) 50.37 N, 17.02 E Kophinou (Larnaca) Cyprus 34.50 N, 33.24 E	15 July 1935 4 December 1935
AE 55703 British Museum	Adult	Famagusta, Cyprus 35.07 N, 33.56 E	7 December 1962
	Killed	Famagusta, Cyprus 35.07 N, 33.56 E	ca 25 March 1963
J 14614 British Museum	Full-grown	Happy Valley Episkopi Cyprus. 34.40 N, 32.55 E	16 February 1964
	Lined	Apollo's Temple (5m E) Cyprus.	26 December 1964

MUSCICAPA STRIATA

9837132 Helgoland		Island of Helgoland Germany. 54.11 N, 7.55 E	17 September 1959
		Episkopi, Cyprus 34.40 N, 32.55 E	30 October 1959

SAXICOLA TORQUATA

J 14064 British Museum	Full-grown Female	Nicosia, Cyprus 35.13 N, 33.22 E	27 January 1961
	Killed	Nicosia, Cyprus 35.13 N, 33.22 E	ca 25 Nov 1963

OENANTHE PLESCHANKA

AE 55953 British Museum	Pullus/4	Episkopi, Limassol Cyprus. 34.40 N, 32.23 E	27 June 1963
	Found dead.	Episkopi, Limassol Cyprus. 34.40 N, 32.23 E	3 April 1964

PHOENICURUS OCHRUIROS

G 453846 Rossitten		Lossen (Germany) Poland 50.48 N, 17.35 E	2 October 1936
		Moni, Limassol, Cyprus 34.44 N, 33.12 E	5 November 1937
G 59485 Varsovia	Juvenile	Dublany (Poland) USSR 49.54 N, 24.05 E	8 August 1937
	Killed	Gefnyeli, Nicosia, Cyprus 35.12 N, 33.18 E	22 November 1937

PHOENICURUS OCHRUIROS

2.316.737 Praha	Adult	Halda near Vyprachtice (Pardubice (Pardubice) Czechslovakia 49.58 N 16.38 E	8 October 1964
	Lined	Leonarisso (Karpas) Cyprus 35.27 N, 34.09 E	8 February 1965
J 14747 British Museum	Adult Female	Happy Valley, Episkopi Cyprus. 34.40 N, 32.55 E	11 February 1964
	Controlled	Happy Valley, Episkopi Cyprus. 34.40 N, 32.55 E	13 February 1965
H 141945 Varsovia	Full-grown Female	Wapnica, Wolin, Poland 53.52 N, 14.26 E	19 September 1965
	Found dead	Kyrenia, Cyprus 35.20 N, 33.20 E	early January 1966

PHOENICURUS PHOENICURUS

019.4775 Helgoland	Juvenile	Mecklenburg, Germany 53.47 N, 10.52 E	14 July 1964
	Caught w/broken leg & released	Episkopi, Cyprus 34.47 N, 32.31 E	January 1965

ERITHACUS RUBECULA

ZOD 4681 Stockholm	Adult	Ottenby (Oland Island) Sweden. 56.12 N, 16.24 E	5 October 1950
	Caught & released w/ring.	Rizokarpaso (Famagusta) Cyprus. 35.36 N, 34.24 E	19 December 1950
E 11938 British Museum	Full-grown	Nicosia, Cyprus 35.13 N, 33.22 E	12 November 1961
	Killed	Nicosia, Cyprus 35.13 N, 33.22 E	5 December 1961
K 131760 Helsinki	Full-grown	Kirkkoummi, Porkkala Finland. 59.56 N, 24.23 E	8 October 1961
		Kato Lefkara, Larnaca, Cyprus. 34.50 N, 33.20 E	December 1961
J 14174 British Museum	Full-grown	RAF Nicosia, Cyprus 35.13 N, 33.22 E.	24 February 1961
	Caught & released w/ring.	RAF Nicosia, Cyprus 35.13 N, 33.22 E	3 December 1961
AE 55691 British Museum	Full-grown	Akrotiri, Cyprus 40.28 N, 25.28 E	1 December 1962
	Caught & released w/ring.	Akrotiri, Cyprus 40.28 N, 25.28 E	23 November 1963

ERITHACUS RUBECULA

AE 55739 British Museum	Full-grown	Akrotiri, Cyprus 40.28 N, 25.28 E	19 January 1963
	Caught on limestick and killed.	Near Limassol (15 m NE) Cyprus. 34.41 N, 33.05 E	12 March 1965
AN 26619 British Museum	Full-grown	Happy Valley, Episkopi Cyprus. 34.40 N, 32.55 E	1 March 1964
	Controlled	Happy Valley, Episkopi Cyprus. 34.40 N, 32.55 E	16 January 1965
J 14746 British Museum	Full-grown	Happy Valley, Episkopi Cyprus. 34.40 N, 32.55 E	11 February 1964
	Controlled	Happy Valley, Episkopi Cyprus. 34.40 N, 32.55 E	10 January 1965
J 14476 British Museum	Full-grown	Happy Valley, Episkopi Cyprus. 34.40 N, 32.55 E	28 January 1964
	Caught on limestick and killed.	Apollo's Temple (5m E) Cyprus.	26 December 1964
J 14506 British Museum	Full-grown	Akrotiri, Cyprus 40.28 N, 25.28 E	8 March 1964
	Caught alive-Being held.	Ayia Trias, Yialousa (100 m NE), Cyprus 35.32 N, 34.11 E	1 December 1964
AE 55973 British Museum	Full-grown	Happy Valley, Episkopi Cyprus. 34.40 N, 32.55 E	24 January 1964
	Caught on limestick and killed.	Apollo's Temple (5 m E) Cyprus.	26 December 1964

TURDUS PILARIS

F 178211 Moskwa		Voronezh, USSR 51.40 N, 39.13 E	7 June 1953
	Recovery details Unknown	Limassol, Cyprus 34.40 N, 33.03 E	February 1961
G 177269 Radolfzell		Mannheim Sandheifen (Baden) Germany. 49.33 N, 8.28 E	31 December 1960
	Killed	Near Lefkara, Cyprus 34.50 N, 33.20 E	7 January 1962

TURDUS MERULA

F 63745 Moskwa	Adult	Azov Sivash Reserve USSR 46.17 N, 35.08 E	10 October 1954
		Kakopetria (Nicosia) Cyprus. 34.59 N, 32.54 E	2 December 1955

TURDUS MERULA

F 230166 Moskwa	Female	Biruchii Island, Sea of Azov, USSR	14 April 1957
		Lythrophonda (Nicosia) Cyprus. 34.42 N, 33.18 E	November 1958

TURDUS MUSICUS

PR 300 London	Juvenile	Gibraltar Point, Great Britain. 53.06 N, 0.12 E	1 November 1952
	Found dead	Kyrenia, Cyprus 35.20 N, 33.20 E	13 February 1954

TURDUS ERICETORUM

F 63729 Moskwa		Azov Sivash Reserve, USSR 46.07 N, 35.08 E	7 October 1954
		Apostolos Andreas (Famagusta) Cyprus. 35.39 N, 34.35 E	9 January 1955
V 23504 British Museum	1st Winter	RAF Nicosia, Cyprus 35.13 N, 33.22 E	8 February 1961
	Found dead	RAF Nicosia, Cyprus 35.13 N, 33.22	9 February 1961
V 23505 British Museum	1st Winter	Nicosia, Cyprus 35.10 N, 35.22 E	8 February 1961
	Killed	Near Nicosia, Cyprus 35.10 N, 35.22 E	February 1962
V 23515 British Museum	1st Winter	RAF Nicosia, Cyprus 35.10 N, 35.22 E	7 March 1961
		Nisou, ca 12 m S of Nicosia Cyprus.	25 February 1962
V 23508 British Museum	1st Winter	Nicosia, Cyprus 35.10 N, 35.22 E	12 February 1961
	Killed	Near Delphi, Greece 38.29 N, 22.30 E	1 January 1962
V 23.514 British Museum	Full-grown	RAF Nicosia, Cyprus 35.13 N, 33.22 E	24 February 1961
	Killed	Nicosia, Cyprus 35.10 N, 33.23 E	2 March 1961
TURDUS PHILOMELOS			
V 23132 British Museum	Full-grown	Akrotiri, Cyprus 40.28 N, 25.28 E	29 December 1962
	Killed	Akrotiri, Cyprus 40.28 N, 25.28 E	20 January 1963

TURDUS PHILOMELOS

4166 OR British Museum	Full-grown	Akrotiri, Cyprus 40.28 N, 25.28 E	21 January 1963
	Caught	Khalassa (11 m NE) Cyprus	8 January 1964
V 23135 British Museum	Full-grown	Akrotiri, Cyprus 40.28 N, 25.28 E	13 January 1963
	Killed	Akrotiri, Cyprus 40.28 N, 25.28 E	11 February 1963
V 23141 British Museum	Full-grown	Akrotiri, Cyprus 40.28 N, 25.28 E	18 January 1963
	Killed	Akrotiri, Cyprus 40.28 N, 25.28 E	5 February 1963

PARUS ATER

E 11036 British Museum	Pullus	Pano Platres, Cyprus 34.53 N, 32.52 E	1 May 1957
	Killed	Pano Platres, Cyprus 34.53 N, 32.52 E	30 October 1960

EMBERIZA CALANDRA

V 23017 British Museum	First winter	RAF Nicosia, Cyprus 35.13 N, 33.22 E	6 January 1959
	Caught by a cat and killed	RAF Nicosia, Cyprus 35.13 N, 33.22 E	19 March 1959

EMBERIZA CAESIA

E 11 490 British Museum	Adult	Mitsero, Cyprus 35.02 N, 32.50 E	5 May 1958
	Not known	Kato Moni, Cyprus 35.03 N, 33.06 E	March 1960

EMBERIZA MELANOCEPHALA

3930 Zagreb	Juvenile	Gata, Yugoslavia 43.28 N, 16.24 E	23 June 1956
	Caught or caught & released w/o ring.	Famagusta, Cyprus 35.09 N, 33.59 E	20 April 1957

FRINGILLA COELEBS

E 11512 British Museum	Adult	RAF Nicosia, Cyprus 35.13 N, 32.22 E	29 January 1959
	Caught & released w/ring.	Vulkaneshty (Madavia) USSR 45.41 N, 28.23 E	2 April 1960

SERINUS SERINUS

J 14611 British Museum	Adult	Episkopi, Cyprus 34.40 N, 32.55 E	16 February 1964
	Shot	Ayia Philaxis (Limassol) Cyprus. 34.41 N, 33.05 E	10 January 1967
J 14729 British Museum	Female	Episkopi, Cyprus 34.40 N, 32.55 E	11 February 1964
	Shot	Limassol (15 m. E) Cyprus 34.41 N, 33.05 E	24 November 1964

SERINUS CANARIUS

E 11338 British Museum	1st Winter	Nicosia, Cyprus 35.10 N, 33.23 E	28 December 1958
	Killed	Nicosia, Cyprus 35.10 N, 33.23 E	23 January 1960

CARDUELIS CHLORIS

HH 41006 British Museum	Post-juvenile Female	Akrotiri, Cyprus 40.28 N, 25.28 E	8 September 1966
	Killed by Slingshot	Akrotiri, Cyprus 40.28 N, 25.28 E	13 July 1967

CARDUELIS SPINUS

51166 Madartani		Harmashatarhagy near Budapest, Hungary	8 October 1955
	Unknown	Spilia, Cyprus	10 October 1959
S201301 Moskwa	Juvenile	Rybatschi, Kaliningrad USSR. 55.11 N, 20.49 E	30 June 1965
	Killed	Kamaria, near Troodos Cyprus. 34.58 N, 32.48 E	30 October 1965

CARDUELIS CARDUELIS

A 135727 Zagreb	Juvenile	Split, Yugoslavia 43.30 N, 16.27 E	12 October 1948
	Caught and released w/ring.	Limassol, Cyprus 34.41 N, 33.05 E	28 January 1949
E 11402 British Museum	Juvenile	Kykko (Nicosia) Cyprus 35.13 N, 33.22 E	31 August 1958
	Killed	Nicosia, Cyprus 35.13 N, 33.22 E	20 April 1959

CARDEULIS CARDUELIS

E 11321 British Museum	Adult	RAF Nicosia, Cyprus 35.13 N, 33.22 E	27 December 1958
	Found dead	RAF Nicosia, Cyprus 35.13 N, 33.22 E	15 February 1959
J 14262 British Museum	Adult	Akrotiri, Cyprus 40.28 N, 25.28 E	12 April 1963
	Recovered dead	Akrotiri, Cyprus 40.28 N, 25.28 E	24 May 1965
AN 26723 British Museum	Juvenile	Happy Valley, Episkopi Cyprus. 34.40 N, 32.55 E	24 August 1964
	Controlled (as 1st winter female)	Happy Valley, Episkopi Cyprus. 34.40 N, 32.55 E	16 February 1965
HH 41043 British Museum	1st year female	Akrotiri, Cyprus 40.28 N, 25.28 E	12 September 1966
	Caught	Limassol, Cyprus 34.41 N, 33.05 E	19 February 1967
670.696 British Museum	Juvenile	Polemidthia, Cyprus 34.41 N, 33.00 E	17 July 1967
	Caught alive, being held.	Polemidthia, Cyprus 34.41 N, 33.00 E	12 October 1967
670.696 British Museum	Juvenile	Polemidthia, Cyprus 34.41 N, 33.00 E	17 July 1967
	Caught alive, being held	Polemidthia, Cyprus 34.41 N, 33.00 E	12 October 1967

STURNUS VULGARIS

F 75870 Moskwa	Juvenile	Oksky Reserve, USSR 54.45 N, 40.50 E	31 May 1949
		Yeroskibos (Paphos) Cyprus 34.45 N, 32.27 E	29 November 1949
F 75814 Moskwa	Juvenile	Oksky Reserve, USSR 54.45 N, 40.50 E	28 May 1949
		Pergamos (Larnaca) Cyprus 35.02 N, 33.43 E	6 December 1949
F 80248 Moskwa	Juvenile	Oksky Reserve, USSR 54.45 N, 40.50 E	27 May 1950
		Mandria (Paphos) Cyprus 34.42 N, 32.32 E	6 November 1950

STURNUS VULGARIS

F 129647 Moskwa	Juvenile	Oksky Reserve, USSR 54.45 N, 40.50 E	25 May 1953
	(COS Bull. [In- correctly reported in <u>Birds of Cyprus</u> as re- covered in 1956])	Ktima (Paphos), Cyprus 34.46 N, 32.25 E	9 November 1953
F 353949 Moskwa	Juvenile	Zakharov (Ryazan) USSR 54.23 N, 29.17 E	3 June 1957
	Recovery details not known	Ayia Kibir, Cyprus 35.25 N, 34.00 E	18 December 1960
F 244937 Moskwa		Alushta (Crimea), USSR 44.40 N, 34.25 E	June 1960
	Recovery details not known	Kiti (Larnaca), Cyprus 35.50 N, 33.35 E	10 December 1960
F 715709 Moskwa	Pullus	Kamenka (Penza) USSR 53.10 N, 44.05 E	28 May 1961
		Paphos, Cyprus 34.45 N, 32.25 E	12 November 1961

JORDAN RECOVERIES

Birds Banded at Azraq Oasis (31°50' N 36°47' E), Jordan
by British Trust or Ornithology, International Jordan
Expedition

SPECIES

Band number	Age and Sex Recovery remarks	Locality banded Locality recovered	Date banded Date recovered
GALLINAGO GALLINAGO			
	Post-juv.	Azraq	25 April, 1966
CV53.517	shot	Skopin, (Ryazan'sk) USSR 53.50' N, 39.34 E	14 Sept., 1969
STREPTOPELIA TURTUR			
2082951	Post-juv.	Azraq	10 Sept., 1963
	shot	Krasnoarmeyskoye: (Ukraine) USSR 48.19 N, 37.10 E	20 Aug., 1966
JYNX TORQUILLA			
BC11.018	Post-juv. retrapped	Azraq Azraq	18 April, 1965 23 April, 1965
	Found dead	Kozelets (Chernigov) USSR 50.56 N, 31.08 E	9 May, 1965
HIRUNDO RUSTICA			
HC28.354	Post-juv. caught	Azraq Yusta (Kalmytskaya) USSR 47.06 N, 46.19 E	28 April, 1966 May, 1966
HC27.785	Post-juv. Found dead	Azraq Magdalinovka (Ukraine) 48.53 N, 34.47 E	26 April, 1966 10 May, 1966
HC27606	Post-juv. Caught	Azraq Kosciuszki (Kielce) 51.26 N, 21.26 E	24 April, 1966 27 July, 1968
SC17.763	Ad. Male Found Dead	Azraq Maykop (Krasnodarsk) 44.35 N, 40.07 E	14 Sept., 1963 18 June, 1964

REPORT

ON THE PROGRESS OF THE WORK DURING THE YEAR 1900

1901

NAME	AGE	SEX	RELATION	DATE
JOHN DOE	25	M	Son	1900
JANE DOE	22	F	Daughter	1900
WILLIAM DOE	20	M	Son	1900
MARY DOE	18	F	Daughter	1900
EDWARD DOE	15	M	Son	1900
ANNE DOE	12	F	Daughter	1900
CHARLES DOE	10	M	Son	1900
ELIZABETH DOE	8	F	Daughter	1900
FRANK DOE	6	M	Son	1900
MARGARET DOE	4	F	Daughter	1900
ALICE DOE	3	F	Daughter	1900
JOHN DOE	2	M	Son	1900
JANE DOE	1	F	Daughter	1900

RIPARIA RIPARIA

HC28.698	Adult cat	Azraq Tikhoretsk (Krasnodar) USSR	1 May, 1966 25 Sept., 1966
HC27824	Post-juv. Killed	Azraq Yeysk (Krasnodar) USSR 46.41 N, 38.15 E	26 April, 1966 20 Sept., 1969

MOTACILLA ALBA

HC28695	Post-juv. Shot	Azraq Quteifa: Damascus, Syria 33.44 N, 36.36 E	1 May, 1966 3 March, 1967
---------	-------------------	--	------------------------------

MOTACILLA FLAVA ? DOMBROWSKII

HC2737	Post-juv M Killed	Azraq Damascus, Syria 33.30 N, 36.19 E	23 April, 1966 23 Sept., 1966
--------	----------------------	--	----------------------------------

SYLVIA ATRICAPILLA

HR75.653	Male Adult Killed	Azraq Amyun, Lebanon 34.19 N, 35.48 E	5 May, 1968 11 May, 1968
AH50.003	Post-juv. M Shot	Abdulla Bridge Jericho 31.38 N, 35.33 E Dhour El-Shuwair Lebanon 33.53 N, 35.45 E	3 Sept., 1963 1 Sept., 1965

SYLVIA CURRUCA

HC28.686	Post-juv. Shot	Azraq Sidon, Lebanon 33.32 N, 35.33 E	1 May, 1966 19 Sept., 1966
----------	-------------------	---	-------------------------------

PHYLLOSCOPUS COLLYBITA

HC27525	Post. juv. Lined	Azraq Paralimni, Cyprus 35.01 N, 33.59 E	24 April, 1966 8 April, 1968
---------	---------------------	--	---------------------------------

BAND RECOVERIES
INFORMATION INCOMPLETE

COUNTRY	BAND #	RECOVERY DATE	RECOVERY SITE	SPECIES
Czechoslovakia	A 532 Praha C SR	20 April, 1939	Lake Burrullus, Egypt 31.35N, 31.05E	
	B 2519 MUS Praha Bohemia	1 Aug., 1944	El Dilingat, Egypt 30.50N, 30.32E	
	M 24125 Praha O-SR	3 Feb., 1957	Imbaba, Giza, Egypt 30.04N, 31.14E	
	R 6379 ZOO Praha Bohemia	11 Sept., 1957	Rosetta, Egypt 31.24N, 30.25E	
England	HV 95410 Brit. Museum London	Fall 1970	5 km NW Burg el Arab, Egypt 30.55N, 29.32E	
Finland	H-102031 Mus. Zool. Hels.	Sept. 1969	52 km W of Alexandria, Egypt	
	H 61459 Mus. Zool. Hels.	10 May 1967	Max, Alexandria, Egypt 31.21N, 29.54E	
	NO 9828 Mus. Zool. Hels.	Fall 1970	Alexandria Bird Mkt., Alexandria, Egypt 31.21N, 29.54E	
Germany (West)	220952 Helgoland	3 July, 1940	Disuq, Kafrel Sheikh, Egypt	
	223505 Helgoland	20 June, 1937	Maghagha, Egypt 28.39N, 30.50E	
	849223 Helgoland	14 Sept., 1942	El Burullus, Egypt 31.35N, 31.05E	
	247025 Helgoland	1966	Hishani River, Lebanon	Pelican?
	238318 Helgoland	April, 1968	Kom Ombo, Egypt 24.28N, 32.57E	<i>Ciconia</i> <i>ciconia</i>
	E0085 Helgoland	1962	Old Tushka, Egypt 22.30N, 31.53E	

COUNTRY	BAND #	RECOVERY DATE	RECOVERY SITE	SPECIES
Germany (West) cont.	240660 Helgoland	May, 1969	New Tushka, Egypt 24.29N, 32.33E	
	B 59767 Helgoland	May, 1969	New Tushka, Egypt 24.29N, 32.33E	
	244214 Helgoland	1945	Old Kalabsha, Egypt 23.33N, 32.52E	
	H 6524 (brown)	1945	Old Kalabsha, Egypt 23.33N, 32.52E	
	236455 Helgoland	15 June, 1964	Diork, Daqahliya, Egypt 30.55N, 31.30E	
	H 9179 Helgoland	29 May, 1964	Kafr el Sheikh, Egypt	
	233938 Helgoland	26 April, 1952	Rah, North Sinai, Egypt	
	211714 Helgoland	15 June, 1940	Nag Hammadi, Qena, Egypt	
	220952 Helgoland		Disuq, Kafr el Sheikh, Egypt 30.00N, 31.15E	
	223505 Helgoland	20 June, 1937	Maghagha, Egypt 28.39N, 30.50E	
	849223 Helgoland	14 Sept., 1942	El Burullus, Egypt 31.35N, 31.05E	
	BB 20082 BB 20083 Rossettin (Two Bands on same bird)	18 May, 1945	Damanhur, Egypt	
	A 3024 Rossettin	3 May, 1942	Manshiya, Egypt	

BB 14671 Rossettin	25 June, 1941	El Wasta, Beni Suef, Egypt 29.20N, 31.12E
-----------------------	---------------	---

B 78865 Rossettin	29 May, 1942	Damanhur, Egypt
----------------------	--------------	-----------------

COUNTRY	BAND #	RECOVERY DATE	RECOVERY SITE	SPECIES
Germany (West) (cont.)	471366 Rossettin	28 Sept., 1940	Damietta, Egypt 31.25N, 31.49E	
	D 70885 Rossettin	6 June, 1942	Madaba, Egypt	
	511856 Rossettin	1 Sept., 1940	Damietta, Egypt 31.25N, 31.49E	
	B 77387 Rossettin	Before 10 May, 1941	(None available)	
	G 430796 Rossettin	10 Feb., 1941	Damietta, Egypt 31.25N, 31.49E	
	BB 4268 Rossettin	Winter 1939	El Manzala, Egypt 30.20N, 31.15E	
	BB 2199 Rossettin	15 April, 1940	East of Haifa, Palestine	
	B 72032 Rossettin	22 April, 1940	El Salum, Egypt 31.34N, 25.09E	
	B 71599 Rossettin	1 May, 1940	Abu Rauwash, Egypt 30.02N, 31.06E	
	BB 13639 Rossettin	June 1940	Saft el Khirsa, Egypt 28.49N, 30.50E	
	215763 Rossettin	18 April, 1936	Idfu, Egypt 24.58N, 32.52E	
	A 1119 Rossettin	11 Oct., 1935		
	B 57663 Rossettin	Before 1964-65	Old El Umbarakab, Egypt 23.40N, 32.53E	
	B 58163 Rossettin	1940	Old El Umbarakab, Egypt 23.40N, 32.53E	
	BB 2926 Rossettin	1944	Old El Umbarakab, Egypt 23.40 N, 32.53E	
	B 70617 Rossettin	Before 1964-65	Old Kalabsha, Egypt 23.33N, 32.52E	

COUNTRY	BAND #	RECOVERY DATE	RECOVERY SITE	SPECIES
Germany (West) (cont.)	B 73689 Rossettin	Before 1964-65	Old Kalabsha, Egypt 23.33N,32.52E	
	B 79989 Rossettin	Before 1964-65	Old Kalabsha, Egypt 23.33N,32.52E	
	BB 400	Before 1964-65	Old Kalabsha, Egypt 23.33N,32.52E	
	BB 5604	Before 1964-65	Old Kalabsha, Egypt 23.33N,32.52E	
	BB 12587	Before 1964-65	Old Kalabsha, Egypt 23.33N,32.52E	
	B 27974	Before 1964-65	Old Kalabsha, Egypt 23.33N,32.52E	
	B 43120(?)	Before 1964-65	Old Kalabsha, Egypt 23.33N,32.52E	
Hungary	24706 Budapest	Before 1964-65	Old Kalabsha, Egypt 23.33N,32.52E	
Israel	Z 1440 Tel Aviv Univ. Israel	Reported 15 June, 1970	Near Kom Ombo, Egypt 24.28N,32.57E	
	C 2416 Tel Aviv Univ. Israel	Fall, 1970	Alexandria Bird Mkt., Alexandria, Egypt 31.12N,29.54E	<i>Coturnix</i> <i>Coturnix</i>
Poland	B 1400661 St. Ornith., Varsovia	17 May, 1967	Port Said, Egypt 31.16N,32.18E	
	5619 Mus. Zool. Varsovia	1 Oct., 1940	Port Said, Egypt 31.16N,32.18E	
	509731 Mus. Zool. Varsovia	24 March, 1940	Marten Baghush, Egypt	
	G 402127 Mus. Zool. Varsovia	Before Jan., 1942	Damietta, Egypt 31.25N,31.49E	
	J 709240 St. Ornith. Polonia Var.	Sept.-Oct., 1969	Burg el Arab, Egypt 30.55N,29.32E	

Date	Description	Particulars	Amount	Total
1890	Jan 1	Balance	100.00	100.00
	Feb 1	Interest	5.00	105.00
	Mar 1	Interest	5.00	110.00
	Apr 1	Interest	5.00	115.00
	May 1	Interest	5.00	120.00
	Jun 1	Interest	5.00	125.00
	Jul 1	Interest	5.00	130.00
	Aug 1	Interest	5.00	135.00
	Sep 1	Interest	5.00	140.00
	Oct 1	Interest	5.00	145.00
	Nov 1	Interest	5.00	150.00
	Dec 1	Interest	5.00	155.00
1891	Jan 1	Balance	155.00	155.00
	Feb 1	Interest	5.00	160.00
	Mar 1	Interest	5.00	165.00
	Apr 1	Interest	5.00	170.00
	May 1	Interest	5.00	175.00
	Jun 1	Interest	5.00	180.00
	Jul 1	Interest	5.00	185.00
	Aug 1	Interest	5.00	190.00
	Sep 1	Interest	5.00	195.00
	Oct 1	Interest	5.00	200.00
	Nov 1	Interest	5.00	205.00
	Dec 1	Interest	5.00	210.00
1892	Jan 1	Balance	210.00	210.00
	Feb 1	Interest	5.00	215.00
	Mar 1	Interest	5.00	220.00
	Apr 1	Interest	5.00	225.00
	May 1	Interest	5.00	230.00
	Jun 1	Interest	5.00	235.00
	Jul 1	Interest	5.00	240.00
	Aug 1	Interest	5.00	245.00
	Sep 1	Interest	5.00	250.00
	Oct 1	Interest	5.00	255.00
	Nov 1	Interest	5.00	260.00
	Dec 1	Interest	5.00	265.00

COUNTRY	BAND #	RECOVERY DATE	RECOVERY SITE	SPECIES
Romania	B 82 Bucarest	1 June, 1947	Zagazig, Egypt 30.35N,31.31E	
	30 O.V.R.S. Romania	2 May, 1935	El Mahalla, Egypt 30.58N,31.10E	"I" Pelican
	R 12820 C. Ornitol- Romania	25 Sept., 1969	Galal, Egypt 31.02N,28.10E	<i>Luscinia</i> ?
Sweden	1494704 Riks Mus. Stockholm	Fall, 1970 (Before 3 Nov.)	Alexandria Bird Mkt., Alexandria, Egypt 31.12N,29.54E	
	1543533 Riks Mus. Stockholm	Fall, 1970 (Before 3 Nov.)	Alexandria Bird Mkt., 31.12N,29.54E	
Turkey	Journal Hurriyet 5 Istanbul Turkey 31	Nov. 1969	Kom Ombo, Egypt 24.28N,32.57E	Spoonbill
U.S.S.R.	C 85373 Moskwa	24 Oct., 1969	Mersa Matruh, Egypt 31.21N,27.14E	<i>Bubulcus</i> <i>ibis</i>
	C 88996 Moskwa	Sept., 1968	Mersa Matruh, Egypt 31.21N,27.14E	<i>Egretta</i> <i>alba</i>
	F 790615 Moskwa	5 Oct., 1969	Ghbaleh, Casa Kistrwam, Lebanon 34.03N,35.43E	
	F 788306 Moskwa	5 Oct., 1969	Ghbaleh, Casa Kistrwam, 34.03N,35.43E	
	D 403300	?	Lake Mariut, Egypt 31.08N,29.56E	
	E 415000 Moskwa	Date unknown (10 Feb., 1960 reported to Moscow)	Port Said, Egypt 31.16N,32.18E	
	E 517860 Moskwa	Date unknown (Same as above)	Port Said, Egypt 31.16N,32.18E	
	D 479106 Moskwa	Date unknown (Same as above)	Port Said, Egypt 31.16N,32.18E

COUNTRY	BAND #	RECOVERY DATE	RECOVERY SITE	SPECIES
U.S.S.R. (cont.)	D 402427 Moskwa	Date unknown (10 Feb., 1960 reported to Moscow)	Port Said, Egypt 31.16N,32.18E	
	M 27824 Moskwa	Date unknown (same as above)	Port Said, Egypt 31.16N,32.18E	
	E 269997 Moskwa	Date unknown (23 June, 1960 reported to Moscow)	Port Said, Egypt 31.16N,32.18E	
	E 517871 Moskwa	Date unknown (same as above)	Port Said, Egypt 31.16N,32.18E	
	E 245552 Moskwa	Date unknown (same as above)	Port Said, Egypt 31.16N,32.18E	
	E 415183 Moskwa	Date unknown (same as above)	Port Said, Egypt 31.16N,32.18E	
	E 272937 Moskwa	Date unknown (same as above)	Port Said, Egypt 31.16N,32.18E	
	E 250407 Moskwa	Date unknown (same as above)	Port Said, Egypt 31.16N,32.18E	
	E 268988 Moskwa	Date unknown (same as above)	Port Said, Egypt 31.16N,32.18E	
	E 422265 Moskwa	Date unknown (same as above)	Port Said, Egypt 31.16N,32.18E	
	E 420550 Moskwa	Date unknown (same as above)	Port Said, Egypt 31.16N,32.18E	
	E 332177 Moskwa	Date unknown (same as above)	Port Said, Egypt 31.16N,32.18E	
	E 267288 Moskwa	Date unknown (same above)	Port Said, Egypt 31.16N,32.18E	
	E 516952 Moskwa	Date unknown (same above)	Port Said, Egypt 31.16N,32.18E	
	E 243486 Moskwa	Date unknown (same as above)	Port Said, Egypt 31.16N,32.18E	
	E 334086 Moskwa	Date unknown (same as above)	Port Said, Egypt 31.16N,32.18E	

COUNTRY	BAND #	RECOVERY DATE	RECOVERY SITE	SPECIES
U.S.S.R. (cont.)	E 140014	Date unknown (23 June, 1960 reported to Moscow)	Port Said, Egypt 31.16N,32.18E	
	E 272856	Date unknown (same as above)	Port Said, Egypt 31.16N,32.18E	
	E 306078	Date unknown (same as above)	Port Said, Egypt 31.16N,32.18E	
	E 511784	Date unknown (same as above)	Port Said, Egypt 31.16N,32.18E	
	E 517478	Date unknown (same as above)	Port Said, Egypt 31.16N,32.18E	
	E 381335	Date unknown (same as above)	Port Said, Egypt 31.16N,32.18E	
	E 375536	Date unknown (same as above)	Port Said, Egypt 31.16N,32.18E	
	E 409592	Date unknown (same as above)	Port Said, Egypt 31.16N,32.18E	
	E 271532	Date unknown (same as above)	Port Said, Egypt 31.16N,32.18E	
	E 420279	Date unknown (same as above)	Port Said, Egypt 31.16N,32.18E	
	E 516917	Date unknown (same as above)	Port Said, Egypt 31.16N,32.18E	
	E 299927	Date unknown (same as above)	Port Said, Egypt 31.16N,32.18E	
	E 516535	Date unknown (same as above)	Port Said, Egypt 31.16N,32.18E	
	E 512836	Date unknown (same as above)	Port Said, Egypt 31.16N,32.18E	
	E 327578	Date unknown (same as above)	Port Said, Egypt 31.16N,32.18E	
	E 215989	Date unknown (same as above)	Port Said, Egypt 31.16N,32.18E	

COUNTRY	BAND #	RECOVERY DATE	RECOVERY SITE	SPECIES
U.S.S.R (cont.)	E244782	Date unknown (23 June, 1960 reported to Moscow)	Port Said, Egypt 31.16N,32.18E	
	E 437792	Date unknown (same as above)	Port Said, Egypt 31.16N,32.18E	
	E 334499	Date unknown (same as above)	Port Said, Egypt 31.16N,32.18E	
	E 381847	Date unknown (same as above)	Port Said, Egypt 31.16N,32.18E	
	? 326355	Date unknown (same as above)	Port Said, Egypt 31.16N,32.18E	
	D 403062	Date unknown (same as above)	Port Said, Egypt 31.16N,32.18E	
	D 213513	Date unknown (same as above)	Port Said, Egypt 31.16N,32.18E	
	D 406309	Date unknown (same as above)	Port Said, Egypt 31.16N,32.18E	
	D 403486	Date unknown (same as above)	Port Said, Egypt 31.16N,32.18E	
	D 403630	Date unknown (same as above)	Port Said, Egypt 31.16N,32.18E	
	D 208842	Date unknown (same as above)	Port Said, Egypt 31.16N,32.18E	
	D 209249	Date unknown (same as above)	Port Said, Egypt 31.16N,32.18E	
	D 403581	Date unknown (same as above)	Port Said, Egypt 31.16N,32.18E	
	M 41947	Date unknown (same as above)	Port Said, Egypt 31.16N,32.18E	
	P 20264	Date unknown (same as above)	Port Said, Egypt 31.16N,32.18E	
	P 20833	Date unknown (same as above)	Port Said, Egypt 31.16N,32.18E	

COUNTRY	BAND #	RECOVERY DATE	RECOVERY SITE	SPECIES
U.S.S.R. (cont.)	B 406 Musee Zool. Kaunas Lithaunie	1962-63	Old Kalabsha, Egypt 23.33N,32.52E	
	101622 Ornithol. Centr. Riga	Before 1964-65	Old Kalabsha, Egypt 23.33N,32.52E	
	E 204077 Moskwa	19 Jan., 1970	Shatta, Egypt 31.25N,31.52E	
	C 88669 Moskwa	25 Dec., 1969	Port Said, Egypt 31.16N,32.18E	
	E 177276 Moskwa	Jan., 1970	Ezbet el Burg, Egypt	
	114355 Riga	13 Jan., 1941	Beni Suef, Egypt	
	3226 Kaunas Univ.	24 Mar., 1940	Maaten Baggush, Egypt	
	116997 Riga	4 Sept., 1940	Damietta, Egypt 31.25N,31.49E	
	36924F Moskwa	12 Nov., 1969	Ataba Bird Market, Cairo, Egypt 30.03N,31.15E	<i>Sturnus vulgaris</i>
	413728 Moskwa	1967	Port Said, Egypt 31.16N,32.18E	
	42907B Moskwa	19 Sept., 1969	Salum, Egypt 31.34N,25.09E	
	C 86675 Moskwa	ca. 1951	Lake Mariut, Egypt 31.08N,29.56E	<i>Larus ?</i>
	E 327548 Moskwa	ca. 1951	Lake Mariut, Egypt 31.08N,29.56E	<i>Larus ?</i>
	E 192522 Moskwa	ca. 1951	Lake Mariut, Egypt 31.08N,29.56E	<i>Larus?</i>
	S 547063 Moskwa	Fall, 1970 (before 3 Nov.)	Alexandria, Egypt 31.12N,29.54E	

In addition, 193 Russian bands collected by the Giza Zoo Cairo and PMS remain outstanding.

COUNTRY	BAND #	RECOVERY DATE	RECOVERY SITE	SPECIES
Yugoslavia	100006 Ljubljana	July, 1966	Burdein, Egypt 30.30N,31.32E	
Mystery Rings	49887 63-105-DF2- T980		Port Said, Egypt 31.16N,32.18E	
	America Boston College	1934	Egypt	Duck
	CRPS 50- 482	?	El Ambarkab, Egypt 23.40N,32.53E	
	CRPS 51- 673	1969	El Ambarkab, Egypt 23.40N,32.53E	
	CRPS 59- 478	1969	El Ambarkab, Egypt 23.40N,32.53E	
	CRYOVAC	Jan., 1967	Lake Menzalla, Egypt 30.20N,31.15E	

SMITHSONIAN INSTITUTION LIBRARIES

3 9088 00170564 9

nhbird SF994.W33

v. 2 A serological and ectoparasite su