OCT 24 1921

ILLUSTRATED CATALOGUE

57,883

OF THE

MUSEUM OF COMPARATIVE ZOOLOGY,
LIBRARY.
THE MUSEUM OF COMPARATIVE ZOOLOGY
CAMBRIDGE, MASS.

AT HARVARD COLLEGE.

No. VI.

SUPPLEMENT TO THE OPHIURIDÆ AND ASTROPHYTIDÆ.

BY

THEODORE LYMAN.

CAMBRIDGE:

PRINTED FOR THE MUSEUM OF COMPARATIVE ZOÖLOGY.

1871.

S

2001
2002
2003 2004 2005 2006 2007 2008 2009 2010 2011 2012 2013 2014 2015 2016 2017 2018 2019 2020 2021 2022 2023 2024 2025 2026 2027 2028 2029 2030 2031 2032 2033 2034 2035 2036 2037 2038 2039 2040 2041 2042 2043 2044 2045 2046 2047 2048 2049 2050 2051 2052 2053 2054 2055 2056 2057 2058 2059 2060 2061 2062 2063 2064 2065 2066 2067 2068 2069 2070 2071 2072 2073 2074 2075 2076 2077 2078 2079 2080 2081 2082 2083 2084 2085 2086 2087 2088 2089 2090 2091 2092 2093 2094 2095 2096 2097 2098 2099 2100 2101 2102 2103 2104 2105 2106 2107 2108 2109 2110 2111 2112 2113 2114 2115 2116 2117 2118 2119 2120 2121 2122 2123 2124 2125 2126 2127 2128 2129 2130 2131 2132 2133 2134 2135 2136 2137 2138 2139 2140 2141 2142 2143 2144 2145 2146 2147 2148 2149 2150 2151 2152 2153 2154 2155 2156 2157 2158 2159 2160 2161 2162 2163 2164 2165 2166 2167 2168 2169 2170 2171 2172 2173 2174 2175 2176 2177 2178 2179 2180 2181 2182 2183 2184 2185 2186 2187 2188 2189 2190 2191 2192 2193 2194 2195 2196 2197 2198 2199 2200 2201 2202 2203 2204 2205 2206 2207 2208 2209 2210 2211 2212 2213 2214 2215 2216 2217 2218 2219 2220 2221 2222 2223 2224 2225 2226 2227 2228 2229 2230 2231 2232 2233 2234 2235 2236 2237 2238 2239 2240 2241 2242 2243 2244 2245 2246 2247 2248 2249 2250 2251 2252 2253 2254 2255 2256 2257 2258 2259 2260 2261 2262 2263 2264 2265 2266 2267 2268 2269 2270 2271 2272 2273 2274 2275 2276 2277 2278 2279 2280 2281 2282 2283 2284 2285 2286 2287 2288 2289 2290 2291 2292 2293 2294 2295 2296 2297 2298 2299 2300 2301 2302 2303 2304 2305 2306 2307 2308 2309 2310 2311 2312 2313 2314 2315 2316 2317 2318 2319 2320 2321 2322 2323 2324 2325 2326 2327 2328 2329 2330 2331 2332 2333 2334 2335 2336 2337 2338 2339 2340 2341 2342 2343 2344 2345 2346 2347 2348 2349 2350 2351 2352 2353 2354 2355 2356 2357 2358 2359 2360 2361 2362 2363 2364 2365 2366 2367 2368 2369 2370 2371 2372 2373 2374 2375 2376 2377 2378 2379 2380 2381 2382 2383 2384 2385 2386 2387 2388 2389 2390 2391 2392 2393 2394 2395 2396 2397 2398 2399 2400 2401 2402 2403 2404 2405 2406 2407 2408 2409 2410 2411 2412 2413 2414 2415 2416 2417 2418 2419 2420 2421 2422 2423 2424 2425 2426 2427 2428 2429 2430 2431 2432 2433 2434 2435 2436 2437 2438 2439 2440 2441 2442 2443 2444 2445 2446 2447 2448 2449 2450 2451 2452 2453 2454 2455 2456 2457 2458 2459 2460 2461 2462 2463 2464 2465 2466 2467 2468 2469 2470 2471 2472 2473 2474 2475 2476 2477 2478 2479 2480 2481 2482 2483 2484 2485 2486 2487 2488 2489 2490 2491 2492 2493 2494 2495 2496 2497 2498 2499 2500 2501 2502 2503 2504 2505 2506 2507 2508 2509 2510 2511 2512 2513 2514 2515 2516 2517 2518 2519 2520 2521 2522 2523 2524 2525 2526 2527 2528 2529 2530 2531 2532 2533 2534 2535 2536 2537 2538 2539 2540 2541 2542 2543 2544 2545 2546 2547 2548 2549 2550 2551 2552 2553 2554 2555 2556 2557 2558 2559 2560 2561 2562 2563 2564 2565 2566 2567 2568 2569 2570 2571 2572 2573 2574 2575 2576 2577 2578 2579 2580 2581 2582 2583 2584 2585 2586 2587 2588 2589 2590 2591 2592 2593 2594 2595 2596 2597 2598 2599 2600 2601 2602 2603 2604 2605 2606 2607 2608 2609 2610 2611 2612 2613 2614 2615 2616 2617 2618 2619 2620 2621 2622 2623 2624 2625 2626 2627 2628 2629 2630 2631 2632 2633 2634 2635 2636 2637 2638 2639 2640 2641 2642 2643 2644 2645 2646 2647 2648 2649 2650 2651 2652 2653 2654 2655 2656 2657 2658 2659 2660 2661 2662 2663 2664 2665 2666 2667 2668 2669 2670 2671 2672 2673 2674 2675 2676 2677 2678 2679 2680 2681 2682 2683 2684 2685 2686 2687 2688 2689 2690 2691 2692 2693 2694 2695 2696 2697 2698 2699 2700 2701 2702 2703 2704 2705 2706 2707 2708 2709 2710 2711 2712 2713 2714 2715 2716 2717 2718 2719 2720 2721 2722 2723 2724 2725 2726 2727 2728 2729 2730 2731 2732 2733 2734 2735 2736 2737 2738 2739 2740 2741 2742 2743 2744 2745 2746 2747 2748 2749 2750 2751 2752 2753 2754 2755 2756 2757 2758 2759 2760 2761 2762 2763 2764 2765 2766 2767 2768 2769 2770 2771 2772 2773 2774 2775 2776 2777 2778 2779 2780 2781 2782 2783 2784 2785 2786 2787 2788 2789 2790 2791 2792 2793 2794 2795 2796 2797 2798 2799 2800 2801 2802 2803 2804 2805 2806 2807 2808 2809 2810 2811 2812 2813 2814 2815 2816 2817 2818 28

HARVARD UNIVERSITY.

LIBRARY

OF THE

MUSEUM OF COMPARATIVE ZOÖLOGY
54-55

57,883

GIFT OF

Thyrium.

October 24, 1921.

OCT 24 1921

ILLUSTRATED CATALOGUE

OF THE

MUSEUM OF COMPARATIVE ZOÖLOGY,

AT HARVARD COLLEGE.

No. VI.

SUPPLEMENT TO THE OPHIURIDÆ AND ASTROPHYTIDÆ.

BY

THEODORE LYMAN.

CAMBRIDGE:

PRINTED FOR THE MUSEUM OF COMPARATIVE ZOÖLOGY.

1871.

UNIVERSITY PRESS: WELCH, BIGELOW, & CO.,
CAMBRIDGE.

E R R A T A .

I N N o . I .

Page	9,	5th line from bottom,	<i>for</i>	Berthelet	<i>read</i>	Berthelot.
"	10,	13th " " "	"	abyssicola	"	abyssicolum.
"	12,	5th " " top,	"	isocanthum	"	isacanthum.
"	14,	22d " " "	"	Asterochema	"	Asteroschema.
"	105,	15th " " "	"	Five arms	"	Five or six arms.
"	118,	12th, 14th lines from top,	"	innermost	"	outermost.
"	197,	5th line from top, right column	"	Asterochema	"	Asteroschema.
"	"	7th " " " " "	"	Asteromyx	"	Asteronyx.
"	198,	36th, 37th line from top, left column	"	52	"	152.

I N N o . V I .

Page	7,	48th line from top	<i>for</i>	grandisquania	<i>read</i>	grandisquama.
"	8,	10th " " "	"	bevispina	"	brevispina.
"	8,	30th " " "	"	Michlin	"	Michelin.
"	(18)	Description Plates, figs. 1 - 6	"	Ophyoglypha	"	Ophioglypha.

IN the six years that have passed since the publication of the first number of this Catalogue, which treated of Ophiuridæ and Astrophytidæ, our knowledge of these families has considerably increased. Many new species have been discovered, among which the singular forms, brought up by the deep-sea dredgings off Florida, and described in Bulletin No. 10, Vol. I., are of particular interest. In the present Supplement to the Catalogue, figures are given of the most important of these deep-sea Ophiurans; and there are added descriptions of one new genus and of five new species from shallow water. There are given, also, lists of the principal authors, and of the new species described by them, since 1865, making a continuation of similar lists published in Catalogue No. I.

LOUIS AGASSIZ.

October 10, 1871.

PRINCIPAL AUTHORS

SINCE THE PUBLICATION OF CATALOGUE No. I, IN MARCH, 1865.

To which are added some publications not before included. The new species are mentioned with each author.

BRADY & ROBERTSON. Ann. and Magazine of Nat. Hist., III. p. 355.

A notice of *Ophianoplus annulosus*, Sars, from W. Coast of Ireland.

DUCHASSAING, P. *Animaux Radiaires des Antilles*. 1850, p. 4.

Ophioderma variegata. (= *O. brevicauda* Ltk. ?)

" saxatilis. (= Young of *O. cinerea* ?)

Ophiolepis annulosa. M. T. (No such thing there.)

" trisquamosa.

" vicina.

" albida.

" Taneredi.

Ophiocoma scolopendrina. M. T. (No such thing there.)

" serpentaria. M. T. (= *O. echinata* Agass.)

" crassispina. M. T. (= " " ")

" punctata (= *O. pumila* Ltk. ?)

Ophiarachna gorgonia. M. T. (No such thing there.)

Ophiothrix fragilis. M. T. (No such thing there.)

" quinquefissa. (= *O. orstedii* Ltk. ?)

Ophiura hexactinia. Lmk. (No such thing there.)

Euryale costosum. } (= *Astrophyton costosum* Seba. ?)

" muricatum. }

Trichaster Isidis (= *Asteropora*, Ltk. ? vel *Astrogomphus* Lym. ?)

I never saw this little pamphlet till recently, and it seems also to have escaped Dr. Lütken's notice. It antedates his descriptions of West Indian Ophiurans by several years; but the short Latin descriptions are so incomprehensible and so vague that I do not think they can be accepted as a basis for priority.

DEJARDIN ET HUPÉ. *Histoire Naturelle des Zoöphytes*. 1862.

Astroporpa dasycladia. (= *Astroporpa annulata*.)

Astroschema affinis. (Good sp. ?)

GRUBE, A. E. Über einige seltene oder neue Ophiuriden. Verhand. der Schlesischen Gesellschaft. 1867, p. 22.

Astrophyton kevipelle. (= *A. asperum* ?)

Ophiolepis adspersa.

Ophiothrix melanostica.

" roseo-cærulans.

" striolata.

Zeitschrift für die Gesamten Naturwissenschaften, XXX. 1867.

Die Insel Lussin und ihre Meeresfauna. 1864.

HELLER, C. Zoophyten und Echinodermen des Adriatischen Meeres. 1868.

Pectinura Forbesii. Lissa, p. 57.

Ophioglypha Grubei. (= *O. albida* ?) Lesina, p. 58.

Ophiothrix alopecurus. M. T. He considers different from *O. fragilis*. p. 63.

HERKLOTS, J. A. Echinodermes de Kuhl, van Hasselt et Sal. Müller. Bydragen tot de Dierkunde. 1869. Colored drawings of *Ophiomastix annulosa*, *Ophiulepis annulosa*, *Ophioplocus imbricatus*, *Ophiocoma picta*, *Ophiarachna incrassata*, *Ophiethrix longipeda*, *Ophiethrix serrata*, K. & v. Has. Good colors, but no details.

- LIUNGMAN, AXEL.** Om några nya arter af Ophiurider. Öfversigt af Kongl. Vetenskaps-Akademiens Förhandlingar. 1866. No. 6.
- Ophiogymna.** Gen. nov.
- Ophiogymna elegans.** Singapore, Hong Kong. p. 163.
- Ophiothrix clypeata.** Singapore. p. 163.
- Ophiactis megellanica.** Straits of Magellan. p. 164.
- “ *fragilis.* Hawai. p. 164.
- Ophiophragmus Loveni.** Rio Janeiro. p. 165.
- Amphipholis.** Gen. nov. (Amphiura pars.) p. 165. See Bulletin Mus. Comp. Zool., No. 10, Vol. I. p. 335.
- Amphipholis Januarii.** Rio Janeiro. p. 165.
- Ophioglypha Kimbergi.** Sidney, N. H. p. 166.
- Ophiuroidea** Viventia huc usque cognita. Öfversigt af Kongl. Vetenskaps-Akademiens Förhandlingar. 1866. No. 9.
- Ophiarachna stellata.** Singapore. p. 305.
- “ *spinosa.* (= *Ophiopeza fallax* ? See Ltk. Addit. III. 36.) Fona Is. p. 305.
- Ophioglypha multispina.** Sidney. p. 307.
- “ *Tenori.* Mediterranean (*Ophiura*, Delle Chiaje). p. 308.
- Ophiopus.** Gen. nov.
- “ *aretiens.* Spitzbergen. p. 309.
- Ophionereis squamata.** Honolulu. p. 310.
- “ *crassispina.* Honolulu. p. 311. (See Bulletin Mus. Comp. Zool., Vol. I., No. 10, p. 312, note.)
- Amphipholis** (vel *Amphiura*) *depressa.* Deep sea between Batavia and Singapore. p. 312.
- “ “ “ *integra.* Port Natal, S. Af. p. 313.
- “ “ “ *hastata.* Mozambique. p. 313.
- “ “ “ *grisea.* Gulf of Guayaquil. p. 313.
- “ “ “ *impressa.* Deep sea between Batavia and Singapore. p. 314.
- “ “ “ *albida.* (= *A. Januarii* ?) Rio Janeiro. p. 314.
- “ “ “ *subtilis.* (= *A. Januarii* ?) Rio Janeiro. p. 314.
- “ “ “ *lobata.* Near Sidney, N. H. p. 314.
- Ophiophragmus antarcticus.** “ *In fretu Magalhaënsi.*” p. 315.
- “ *gibbosus.* Port Natal, S. Af. p. 316.
- “ *echinatus.* Sea between Batavia and Singapore. p. 316.
- Amphiura** *divaricata.* Deep sea between Batavia and Singapore. p. 318.
- “ *Eugenie.* Atlantic, opposite river La Plata. p. 318.
- “ *candida.* Near Mozambique. p. 318.
- “ *complanata.* Atlantic, off Rio Janeiro. p. 319.
- “ *crassipes.* “ “ “ p. 319.
- “ *flexuosa.* Near Brazil. p. 319.
- “ *verticillata.* Galapagos Islands. p. 320.
- “ *capensis.* Near Port Natal and Cape of Good Hope. p. 320.
- “ *magellanica.* Straits of Magellan. p. 320.
- “ *latispina.* Atlantic, off river La Plata. p. 320.
- “ *atlantica.* Near St. Helena. p. 321.
- Ophiocentrus.** Gen. nov.
- “ *aculeatus.* Deep sea between Batavia and Singapore. p. 321.
- Amphilepis.** Gen. nov.
- “ *norvegica.* Norway. p. 322.
- Hemipholis affinis** (*H. gracilis* Vill.) Guayaquil. p. 322.
- Ophiactis** (?) *Savignyi* (*Ophiolepis* M. T.) Egypt. p. 323.
- “ *carnea.* Port Natal, S. Af. p. 324.
- Ophiacantha indica.** Deep sea between Batavia and Singapore. p. 326.
- Ophioscolex purpurea.** Düben and Koren, Norway. p. 327.
- Ophiocnemis obscura.** Straits of Malacca. p. 333.
- Tillägg till kännedomen af Skandinavians Ophiurider. Öfvers. af K. Vet.-Akad. Förh. 1864. No. 7, with a plate.
- Ophiocten sericeum** Ljn. (= *Ophiura sericea* Forbes. = *Ophiocten* Kroyeri Ltk.) p. 360.
- Amphiura tenuispina.** (= *Amphiura squamata*, var. ?) p. 360.
- “ *norvegica* Norway. p. 363.

- LJUNGMAN, AXEL. *Ophiactis clavigera*. S. Norway. p. 365.
Om tvänne nya arter Ophiurider. Öfversigt af Kongl. Vetenskaps Akademiens Förhandlingar, 1870. No. 5.
- Ophiacantha vivipara*. Atlantic, near E. Patagonia. p. 471.
Ophioglypha Lymani. " " " " p. 472.
- LORENZ, J. R. *Physikalische Verhältnisse*, etc. der Organismen in quarnerischen Golfe.
- LÜTKEN, C. F. *Additamenta ad Historiam Ophiuridarum*. Part III. 1869, from the Vidensk. Selsk. Skr.
- Ophionephthys*. Gen. nov.
Ophionephthys limicola. St. Thomas. p. 98.
Ophionema. Gen. nov.
Ophionema intricata. St. Thomas. p. 98.
Amphipholis fissa. Amur. p. 98.
Ophioconis. Gen. nov. (= *Pectinura* Heller.) (*O. Forbesii*.) p. 98.
Ophiopsammus. Gen. nov. (= *Ohiopceza* Yoldii.) p. 98.
Ophiochæta. Gen. nov.
Ophiochæta setosa. Feejee Islands. p. 98.
Ophiarachna affinis. " " " " p. 98.
Ophiomastix mixta. Samoa Islands. p. 99.
" *asperula*. Feejee Islands. p. 99.
" *earyophyllata*. Feejee Islands. p. 99.
Ophiomyxa australis. Between Australia and Tasmania. p. 99.
Ophiocoma canaliculata. " " " " p. 99.
Ophiacantha pentaerinus. (= *O. meridionalis* Lym.) W. Indies, deep sea. p. 99.
Ophiothrix striolata. Grube. New Guinea and China Sea. p. 99.
" *elegans*. China Sea. p. 100.
" *trilineata*. Samoa Island. p. 100.
" *capensis*. Cape of Good Hope. p. 100.
- Asteromorpha*. (Gen. nov.)
- Asteromorpha Steenstrupii*. Unknown. p. 100.
- LYMAN, THEODORE. Preliminary Report on Ophiuridæ and Astrophytidæ dredged in deep water between Cuba and the Florida Reef, by L. F. de Pourtales. Bulletin Mus. Comp. Zool., Vol. I., No. 13. Nov., 1869.
- Ophioglypha acervata*. (Illustr. Cat., No. VI., Pl. I., fig. 6.) p. 316.
" *falcifera*. (" " " " Pl. I., figs. 3, 4, 5.) p. 319.
Ophiocten depressum. (" " " " Pl. II., fig. 10.) p. 320.
Ophiomusium. Gen. nov.
Ophiomusium eburneum. (Illustr. Cat., No. VI., Pl. II., figs. 1, 2, 3.) p. 322.
Ophiacantha pentaerinus Ltk. (= *O. meridionalis* Lym.) p. 324.
Ophiomitra. Gen. nov.
Ophiomitra valida. (Illustr. Cat., No. VI., Pl. II., figs. 4, 5, 6.) p. 325.
" *sertata*. p. 326.
Ophiochondrus. Gen. nov.
Ophiochondrus convolutus. (Illustr. Cat., No. VI., Pl. II., fig. 7.) p. 328.
Ophiactis humilis. p. 329.
" *plana*. p. 330.
" *loricata*. p. 331.
Amphiura semiermis. p. 332.
" *grandisquania*. p. 334.
" *pulebella*. p. 337.
Ophiocnida olivacea. (Illustr. Cat., No. VI., Pl. I., fig. 7, 8.) p. 340.
Ophiothamnus. Gen. nov.
Ophiothamnus vicarius. (Illustr. Cat., No. VI., Pl. II., figs. 8, 9.) p. 342.
Ophiomyces. Gen. nov.
Ophiomyces mirabilis. p. 343.
" *frutectosus*. (Illustr. Cat., No. VI., Pl. II., figs. 11, 12, 13.) p. 345.
Ophiocreas. Gen. nov.
Ophiocreas lumbricus. (Illustr. Cat., No. VI., Pl. I., figs. 19, 20, 21.) p. 347.
Astrophyton mucronatum. p. 348.
Astrogomphus. Gen. nov.
Astrogomphus vallatus. (Illustr. Cat., No. VI., Pl. I., figs. 16, 17, 18.) p. 350.
- Illustrated Catalogue of the Museum of Comparative Zoölogy. No. VI. Supplement to the Ophiuridæ and Astrophytidæ. 1871. Two plates.
- Ophiomaza*. Gen. nov.
Ophiomaza cacaotica. Zanzibar.

- LYMAN, THEODORE. *Ophiothela tigris*. Unknown.
Ophiocnida Putnami. Hong Kong (?).
Ophioglypha sinensis. " " "
Ophiomastix janualis. Mexillones, Bolivia.

The two plates give also figures of the chief species from the deep-sea Florida dredgings.

- MARTENS, E. v. Die Ophiuriden des indischen Oceans. Wiegmann's Archiv für Naturg. XXXVI. p. 244. 1870.

- Ophiactis* maculosa. South China Sea. p. 248.
 " incisa. Singapore, Java, China Sea. p. 249.
Ophiomyxa bevispina. Amboina? p. 249.
Ophiocoma alternans. S. W. Java. p. 252.
 " ternispina. Larentuka, Flores Is. p. 253.
Ophiothrix carinata. Singapore. p. 255.
 " punctolimbata. Java. p. 257.
 " rotata. Mindanao. p. 259.
 " cataphracta. Singapore. p. 260.
 " triloba. Red Sea. p. 260.

Ueber Ostasiatische Echinodermen. Wiegmann's Archiv für Naturg. XXXII. 1866. p. 87.

Trichaster flagellifer. (= *T. palmiferus* ?)

Monatsbericht der Kön. Akad. Berlin. 1867. p. 345 and 481.

[*Ophiocoma* ocellata.] (= *Ophiarachna* incrassata M. T.) Cape York, N. Australia. p. 345.

- Ophiothrix* purpurea. Amboina. p. 346.
 " viridiabilis. China Sea. p. 347.

Amphiura planispina. Rio Janeiro. p. 347.

Hemieuryale. Gen. nov.

Hemieuryale pustulata. West Indies? p. 484.

MECZNIKOFF. Bullet. Acad. Imper. St. Petersburg. XIII. p. 285; also, Mem. Acad. Imper. St. Petersburg. XIV. No. 8. On the anatomy and metamorphosis of *Amphiura* squamata.

MICHLIN, H. In Annexe to Notes sur l'île de la Réunion. Astroschema Rousseaui.

NORDMAN. Notiz über die Gattung Asterophyton vorkommend in der Littoralfauna Russlands. Ofversigt af finska Vetenskaps-Societets forhandlingar. IV. 1856-57.

NORMAN, A. M. On the Morphology of Ophiurans. Rep. British Assoc. 1863.

PACKARD, A. Memoires Boston Soc. Nat. Hist. Vol. I., Part 2, 1867. See also Canadian Naturalist and Geologist. 1863.

SARS, M. Over det dyriske Livs Umbredning in Havets Dybder. Vidensk.-Selsk. Forhandlinger. 1868. p. 8.

Om Artiske Dyreformer in Christianiafjord. Christiania Videnskab. Selsk. Forhandl. 1865.

SARS, G. O. Nye Echinodermter fra den Norske Kyst. Vidensk.-Selsk. 1871. p. 6.

- Ophiacantha* abyssicola. Lofoten Is., 120-300 fathoms.
 " spectabilis. Bodoe (Norway), 80-100 fath.
 " anomala. Sultenfjord, near Bodoe, 150-200 fath.

Ophiopeltis borealis. Lofoten Is. and Bodoe, 80-300 fath.

Ophioglypha gracilis. " " " 150-300 fath. Bay of Christiania. 100-120.

SCHWALBE. Archiv. für mikrosk. Anatomie. V. Notice of the muscles of *Ophiotrix* fragilis.

TSCHERNIAWSKY. Protocollen der Moskauer Naturf. Versammlung. 1869. *Amphiura* Stepanovii. Black Sea. The only echinoderm found in those waters.

VERRILL, A. E. On the Polyps and Echinoderms of New England. Proceed. Boston Soc. Nat. Hist., X., p. 333. 1865.

Descriptions of Star-fishes and Ophiurans from the Atlantic coasts of America and Africa. Amer. Journ. Sc. Vol. II. 1871.

Amphipholis abdita (specimen with no disk). Near New Haven. p. 4.

Notes on Radiata. Trans. Connecticut Academy Arts and Sc., Vol. I., p. 251. March, 1867.

July, 1867, pp. 325, 341. Feb., 1868, p. 366. April, 1868, p. 372. Jan., 1871, p. 572.

Astrophyton panamense. Panama and Pearl Is. p. 251.

Ophiura Daniana. La Union, San Salvador. p. 254.

Hemipholis gracilis. (= *H. affinis* Ljn.) Panama. p. 262.

Ophiothela. Gen. (subgen.) nov.

Ophiothela mirabilis. Pearl Is., Panama Bay. p. 268.

On new and imperfectly known Echinoderms. Proceed. Boston Soc. Nat. Hist., XII., p. 381. 1869, p. 387.

Astrophyton Stimpsonii. Ochotsk. p. 388.

Ophiarachna maculata. New Zealand. p. 389.

Ophiothela Danae. Feejee Islands. p. 391.

DESCRIPTION OF NEW SPECIES.

OPIHIOMAZA.* GEN. NOV.

Teeth. Tooth-papillæ numerous, and arranged in a close vertical oval, as in *Ophiothrix*. No mouth-papillæ. Disk below naked, but covered above with large, swollen radial-shields and plates. Arm-spines stout, nearly smooth, as in *Ophiocoma*.

This curious genus has the dentition of *Ophiothrix*, the disk of *Ophiocnemis*, and the arms of *Ophiocoma*.

Ophiomaza cacaotica. SP. NOV.

Plate I., fig. 15.

Special Marks. — Five, sometimes six arm-spines, the two or three upper ones considerably largest. No tentacle-scale. Interbrachial plates on back of disk elongated and irregularly broken.

Description of a Specimen. — Diameter of disk, 11^{mm}; from outer side of mouth-shield to outer corner of opposite mouth-slit, 4.2^{mm}; distance from outer side of mouth-shield to inner points of tooth-papillæ, compared to that between outer corners of mouth-slits, 2.5 : 2. Width of arm, 2^{mm}; length of arm about 30^{mm}; tooth-papillæ small and crowded, about 26 to each angle of the mouth; four strong, square-edged teeth. Under arm-plates, beyond edge of disk, broader than long; their laterals and inner side slightly re-enteringly curved; outer side slightly curved; length to breadth (6th plate), 9 : 1.2. Within the disk there are six plates, of which the one next the mouth-slit is small and oval; those outside of it grow gradually broader and are shaped like those beyond the disk; side arm-plates stout, but not much projecting; upper arm-plates broader than long, somewhat irregular, broader without than within, outer side curved; length to breadth (8th plate), .9 : 1.6. The first two or three plates are narrow and wedge-shaped and fit in a notch in the disk; beyond the margin they grow wider, and have their

* *Οφίς, serpent; μάζα, a cake.

full width at the 7th or 8th plate; disk naked below and covered by thick, wrinkled skin; radial shields very large, pear-seed shape, length, to breadth, 4 : 2.5, separated by a band of three or four elongated, narrow, broken, and irregular plates, which come to the edge of the disk, and are continuous with the upper arm-plates. Interbrachial spaces covered by plates similar in form to those between radial-shields, but wider; in the centre, a rosette of small, very irregular pieces, one in the middle, and five close around it. The edge of the disk has a row of stout, swollen plates, three to five in each interbrachial space, of which one is opposite the radial-shield on either side, and from one to three are opposite the interbrachial plates. Arm-spines usually five sometimes six; the two or three upper ones much the longest; all stout, tapering, blunt, cylindrical; lengths, to that of the upper arm-plate (7th joint), 2.2, 2, 1.6, 1, 1, .5 : .9. Near end of arm, three or four spines, of which the under one is much swollen at the base, and slightly hooked at the end. Color, in alcohol: deep chocolate. Zanzibar, collected by Mr. Cooke.

Ophiothela tigris. SP. NOV.

Plate I., figs. 10, 11, 12.

Special Marks. — Back of disk completely covered by the radial-shields, which are quite smooth, as are also the interbrachial spaces below. Five arms.

Description of a Specimen. — Diameter of disk, 7^{mm}; length of arm (which had been repaired), 12^{mm}; from outer corner of mouth-slit to outer edge of opposite mouth-shield, 3^{mm}; width of arm, including side arm-plates, 2^{mm}; distance from outer corner of mouth-shield to points of tooth-papillæ, to that between outer corners of mouth-slits, 1.6 : 1.5. Tooth-papillæ arranged in a close, vertical, oval clump; about fifteen in number; four stout, flattened, squarish teeth, which with the tooth-papillæ form a dentition like that in *Ophiothrix* (a generic character); mouth-shields irregular, sometimes in form only of a small swelling; but, when clearly defined, having an oval heart-shape, with outer side straight, and with length to breadth, .5 : 1. Side mouth-shields large and conspicuous; their free lateral sides re-enteringly curved; their inner laterals joined along their whole length; their inner sides making together a wide curve; under arm-plates covered with thick skin and seen indistinctly, except when dry. They have an irregular triangular form, with a peak within and a lumpy surface. They cover only a part of the arm, and differ in figure one from another. Side arm-plates

standing out independently, as pad-like ridges (a generic character), which are stout and swelled; upper arm-plates represented by a double row of irregular elongated warts, which just at the base of the arm are increased in number, so as to form a clump of different-sized pieces. At the tapering end of the arm there are still the two warts at the base of each joint, and between these a double row of fine grains. Disk wholly covered above by the ten great radial shields, which are regular, slightly swelled, and smooth; at the insertion of the arm and in the interbrachial space they form a slight notch or re-entering angle. Interbrachial spaces below covered with thick skin, without spines or scales; genital-slits with a large prominent scale at their outer end, whose edge can be seen from above. Arm-spines like little stumps, thorny at their ends; towards the end of the arm they become more flattened, and the three lowest ones are armed with three or four flattened hooks. The first joint outside the mouth-shield has two or three spines on each side; the second three, and the rest four. Color, in alcohol: upper surface lined and marbled with dark green and light yellow, interbrachial spaces below dark green; under surface light greenish.

Another specimen had a disk of 5.5^{mm}. and arms of 18^{mm}. The mouth-shields were all regular, except the madreporic, which was swelled. The back of the disk had a central pentagon of marbled bluish and yellow, surrounded by a band of yellowish. The animal doubtless has the power of *moving* the side arm-plates, which in this specimen were folded close on the arm.

Locality unknown, but supposed to have been brought from the Pacific by the Exploring Expedition.

Ophiocnida Putnami.* SP. NOV.

Plate I., fig. 9.

Special Marks.—Nine or ten flat arm-spines, the upper ones with broad ends; they closely cover the arm, as if with grains of rice regularly set in rows. Disk beset with similar, but smaller spines. Two swelled papillæ at the apex of the mouth-angle. Two tentacle-scales.

Description of the Specimen.—Diameter of disk, 16^{mm}. Width of arm, 2.5^{mm}; length of arm about 72^{mm}. From outer side of mouth-shield to outer corner of opposite mouth-slit, 4.8^{mm}; distance from outer corner

* Named after the late Captain W. H. A. Putnam, a man of a fine taste in natural history, and who enriched this Museum with remarkable collections made during his sea voyages.

of mouth-slit to inner point of mouth-papillæ, to that between outer corners of mouth-slits, 2.2 : 2.4 ; mouth-shields small, rounded, narrower without, length to breadth, 1.1 : .8 ; side mouth-shields small and not meeting within ; two large swelled mouth-papillæ at the inner apex of the angle ; they are pressed close together and present a re-entering grinding surface ; above, and partly covered by them, a swollen irregular tooth, followed by three more which are squeezed together and more or less thickened ; above these again are four flat, square-edged teeth separated from each other, the uppermost are longest. Outside each mouth-papilla stands the thick, club-like tentacle-scale of the first mouth-tentacle ; above which, and nearly hidden in the mouth-slit, is the tooth-like scale of the second mouth-tentacle. Under arm-plates small, nearly square, with slightly re-entering sides ; length to breadth (9th plate), .8 : .8. Upper arm-plates small, nearly round, but farther out on the arm, the inner side is straight ; length to breadth (12th plate), .8 : .9. Arm-spines 9 (near base of arm 10), white, flattened, upper ones broadened towards the end, like a narrow spatula ; lowest one broadest and stoutest, next two or three above it narrower than the rest ; length of uppermost, .8^{mm} ; 7th, 1.2^{mm} ; lowest, 1^{mm}. towards point of arm. 5 or 6 spines which are flat and taper to a blunt point. No tentacle-scales ; tentacles large and long. Disk puffy and lobed, covered above and below with fine, regular scales, about 60 to a square ^{mm} ; beset above and below with numerous stout, flattened, spines, .8^{mm} long, and resembling the arm-spines. Radial-shields small, inclining to crescent-shape, separated and standing just over base of arm ; length to breadth, 2.6 : 1. Color, in alcohol : disk gray ; arms straw color.

Collected, almost without doubt, at Hong Kong, by Captain W. H. A. Putnam.

This handsome species is an *Amphipura* with a thorny disk ; and I therefore place it in *Ophiocnida*, as an approximation to its natural position, without asserting that its affinities are thus exactly expressed. See Bulletin of the Mus. Comp. Zool., Vol. I. p. 335.

Ophioglypha sinensis. Sp. nov.

Plate I., figs. 1, 2.

Special Marks. — A pit or depression between the side arm-plates, on the under side of the joints within the disk (as in *O. lucertosa*). Only one tentacle-scale on most of the pores beyond the disk. An arm-comb along the edge of the radial-scale, but none on the arm itself.

Description of a Specimen. — Diameter of disk, 9^{mm} ; from outer corner of mouth-slit to outer side of opposite mouth-shield, 4.2^{mm} ; distance from outer side of mouth-shield to point of innermost mouth-papilla, to that between outer corners of mouth-slits, $2:2$; width of arm, 1.4^{mm} . Mouth-papillæ, six to each angle, whereof the two outer ones on each side are thin and triangular, and the innermost is stouter and more elongated, approaching the form of the teeth, which are four in number, stout, and formed like a blunt spear-head. Mouth-shields large, "lyre-shaped," that is to say, bounded by a curve without, an angle within, and by re-entering curves on the sides; so that the inner corners, at the heads of the genital-slits, make little peaks; length to breadth, $1.7:1$. Side mouth-shields very narrow, meeting within, and running along inner angle of mouth-shield to its corner, where they run across the head of the genital-slit to the side arm-plate. Under arm-plates small, much wider than long, with an angle within and a curve without. This is their general form, except that the first two or three are squeezed laterally; length to breadth (8th plate), $.2:.5$, a little way out they become very minute. Side arm-plates large, meeting below along a line longer than the under arm-plate; those within the disk are indented at their juncture, making a little pit. They make a re-entering curve where the tentacle-scale is fixed, and at little peak between it and the under arm-plate. Upper arm-plates strongly arched, wider without than within; bounded by straight lines on the sides and by a curve without; length to breadth (8th plate), $.6:.8$. Disk covered with larger and smaller irregular scales, among which may be seen the larger, round, primary plates. Radial-shields pear-seed shape, nearly or quite touching without; length to breadth, $1.3:1$. Radial-scales carrying a close comb of rounded tapering papillæ, in form like the arm-spines, which grow smaller as they pass below the disk and end in a row of fine grains, midway the genital-slit. Arm-spines, three, slender, rounded, regularly tapering, not quite so long as the joint; length to that of under arm-plate (8th joint), $.8.7.7:2$. Tentacle-scales thin and nearly round. On the first pair of great pores, three or four on each side; on the second and third, three on one side and two on the other; on the fourth, two on one side and one on the other. The next four or five pores have one large scale, and a very small one, which last stands on the side arm-plate. Color, in alcohol: straw yellow.

Collected, almost without doubt, at Hong Kong, by Captain W. H. A. Putnam.

Variations. — All the specimens had broken arms, but by their character they had probably about the proportions of *O. Sarsii*, say a length of three or four times the diameter of the disk. There were two specimens in which the side mouth-shields were broader; the pits on

the under side of the base of the arm, little marked or wanting, and no little peak in the under arm-plate next the tentacle-scale. Perhaps another species, perhaps only a variety.

O. sinensis looks a good deal like *O. lucertosa* of Europe, but is distinguished by wanting the additional comb of papillæ on the base of the arm, and in having fewer tentacle-scales.

Ophiomastix janualis. Sp. nov.

Plate I., figs. 13, 14.

Special Marks.—Disk-scales conspicuous below, but obscured by thick skin on the upper surface of disk, which is beset with sharp tapering spines; but none below. Three round tapering arm-spines, the uppermost longest.

Diameter of disk, 5.5^{mm} ; arm nearly cylindrical, 1.2^{mm} wide; very slowly tapering to a length of 24^{mm} , where it is broken off. From outer corner of mouth-slit to outer side of opposite mouth-shield, 2.5^{mm} ; distance from outer side of mouth-shield to inner points of tooth-papillæ, to that between outer corners of mouth-slits, $1.2 : 1.5$. Tooth-papillæ four, arranged in a close clump, directly below the teeth, which are as in *Ophiocoma*. Mouth-papillæ, 8, arranged in a close half-circle round the mouth-angle; outer ones flat, rounded, as long as broad; inner ones narrower and smaller. Mouth-shields longer than broad, narrower within than without, angles rounded; length to breadth, $.9 : .6$. Side mouth-shields triangular, lying along the sides of the mouth-shields, but not meeting within. Under arm-plates wider without than within, with a little peak without, and their sides a little re-enteringly curved; length to breadth (6th joint), $.6 : .5$. Farther out, the plates become more elongated, with a clean curve without, re-entering sides, and a truncated angle within. Side arm-plates projecting but slightly; at base of arm they meet neither above nor below, but, at the 11th upper arm-plate, they meet above. Upper arm-plates wedge-shaped, with a curved outer side, and laterals re-enteringly curved; length to breadth (6th plate), $.7 : .6$. Disk finely scaled in the interbrachial spaces below, each about $\frac{1}{8}^{\text{mm}}$ long; above the scaling is covered except near the edge, round the radial-shields, and at the base of the arm, to which it extends. Scattered over the upper surface are a few sharp spines, 1.5^{mm} long, which are mounted on slight elevations and stand over the places of the primary plates. The radial-shields project close to the arm as little, irregular, triangular, swellings about $.7^{\text{mm}}$ long. Arm-spines three, rounded, slightly tapering, blunt;

under a lens, finely grooved; lengths (5th joint). 2.3, 2, 1.2. Tentacle-scales usually two, of which the outer is nearly oval, the inner small, narrow, and tooth-like. Color, in alcohol: chocolate brown, dappled with yellowish.

Mexillones. Bolivia; Captain Putnam.

As to the genus of this specimen, it may be called a spiny *Ophiocoma*, or an *Ophiomastix* wanting the club-shaped arm-spine. Müller & Troschel, in characterizing *Ophiomastix*, say (Syst. Asterid., 107), "Disk beset with scattered spines. . . . On the arms, above the spines, club-shaped bodies running out at their ends into several projections"; and their original, *O. annulosa*, has a fine imbricated scaling, from which stand out little naked radial-shields. Lütken (Addit. ad Hist. Oph., Pt. III. p. 26) has shown that the club-shaped spine is not a constant feature in the genus, which therefore is hardly to be set off from *Ophiocoma*, and that, moreover, the disk may be beset either with spines of various sorts, or with spines and grains. With the addition of this species, his view of the genus would stand thus:—

A. Disk smooth or with very few short spines,	<i>O. venosa</i> Pet.
B. Disk closely beset with grains and moderately long spines,	<i>O. mixta</i> Ltk.
C. Disk densely beset with very short spines,	<i>O. asperula</i> Ltk.
D. Disk on both sides beset with long thin spines,	{ <i>O. annulosa</i> M. T.
	{ <i>O. caryophyllata</i> Ltk.
E. Disk beset on its upper surface, only, with slender spines; arm-spines all slender and tapering,	{ <i>O. janualis</i> Lym.

The species under A, B, C, and D all have certain of the arm-spines either club-shaped or forked at the end, but that under E has no thickening of these spines.

NOTE ON NOMENCLATURE AND CLASSIFICATION.

Dr. Lütken, in a foot-note on page 68 of his Additamenta (Part III.), remarks: 1. That names prior to Linnæus have no place in present nomenclature. 2. That, when an author has described a species, his name must always be put after that species as the authority, no matter if the *generic* name be *changed*. As to the first of these propositions, I may say, that Linnæus first contrived what is called binomial nomenclature, in which each animal has two names, the generic and the specific. Consistency is the first duty of a naturalist, therefore it was the first duty of the followers of Linnæus to respect and to adopt such binomial names as may be found in ante-Linnaean authors, of whom some were among the most celebrated of zoölogists. When, then, I find the name *Astrophyton costosum* distinctly used by Seba (III. Pl. IX., Fig. 1, p. 16, 1758), and moreover an excellent figure given, I shall certainly apply that name to the West Indian species to which it belongs. Did not

Seba, more than a century ago, publish a fine folio, with figures that are better than some of those we see nowadays, and shall we ignore his names when they are such as may properly be taken? In putting such a name as *Ophiopholis bellis* for *Ophiopholis aculeata*, I think I have gone too far, because the name *bellis* of Linck is used as part of a diagnosis and not as a name. I was, however, encouraged in the selection by the usage of Johnson and of Forbes.

As to the second point, Dr. Lütken expresses great astonishment, "after all that has been written," that anybody should change the *authority* with each change in the *combination* of a name. For instance, we have, in 1854, *Ophiura nodosa* Ltk. I afterwards attempted to show that *Ophiura* was preoccupied, and made a name, *Ophioglypha*, to take its place; and the question now is, Shall it be written *Ophioglypha nodosa* Ltk. or *Ophioglypha nodosa* Lym.? Dr. Lütken has no cause for astonishment. There are two parties to this question. That to which he belongs insists on considering *credit* or *honor* the real reason for using names of authors; and always speaks of the "injustice" done when an author's name is lost sight of. The party whose views I hold maintains that nomenclature is *a system of exact registration*, and that, with the present enormous mass and confusion of titles, no other guide is possible; and further, that the credit of a zoölogist does not rest on his monogram, but on something better. Will the reputation of John E. Gray be greater because his name thus appears often; or will that of Lacaze-Duthiers be less because his appears seldom? After what I have said in the Bulletin (Vol. I. p. 336, note), I can add nothing more to the point than a quotation from Alexander Agassiz (American Naturalist, Vol. V. p. 354):—

"The history of the present confused condition of nomenclature is an interesting one; it is the attempt to show by the binomial system, not only the correct name of any animal, but, at the same time, give a short historical sketch of the changes the name has undergone. The name of an animal or plant is that binomial combination which it has first received, let us say *A b* from Linnaeus; [*A* (generic) *b* (specific)]. Subsequent changes, such as the transfer of this to a different genus, *B*, by Cuvier, are simple matters of registration, a part of the *history of the science*, showing what Cuvier thought of the affinities of the species named *A b* by Linnaeus. When, then, we speak of this species as *B b* Cuvier, we are recording his views as an investigator, and this does not lessen whatever credit there may be in the original description of *A b* by Linnaeus. If afterwards Blainville comes and says that Cuvier should have referred *A b* to the genus *C* of Latreille, and quotes this species hereafter as *C b* Blainville, he is only recording his opinion, and so on through indefinite time. *Changes which the progress of science render necessary in the position of A b of Linnaeus are or should be quoted under the*

authority of the author who proposes them as expressing the actual condition of our knowledge of the affinities of the species A b of Linnaeus."

At any rate, let the decision be what it may, Dr. Lütken need have no fear that his own reputation rests on any such unstable basis. To sum up, *Astrophyton costosum* Seba, and *Ophioglyphus nodosa* Lyman, mean just what they should mean, and nothing more; to wit, that in the writings of these two persons will be found these names definitely, used for the *first time*. They do not distinguish these persons as good or as poor zoölogists, nor do they show that such names are the correct ones; but they are simply the "trade-marks" of workmen.

As to the "Synopsis" of *Ophiuridae* presented by Dr. Lütken (l. c. p. 87), I have no objection to it, provided it be understood as a movement towards a true classification, or as a really convenient tabulation of genera. But inasmuch as it rests on the structure of the mouth parts it cannot be expected to be an exposition of nature, any more than can a classification of fishes based on their scales; of mammalia, on their nervous centres; or of molluska, on the character of their shells. Any one who will examine the single original genus *Amphiuva* will be convinced how impossible it is to arrange it only on the *number* of the mouth-papillæ. In fact, the species of *Ophiuridae* are now so numerous, that some one man must examine them all with his own eyes, before we can hope for a good arrangement. And especially do the genera *Ophiocoma* and *Ophiothrix* need severely to be weeded.

DESCRIPTION OF THE PLATES.

These are all alcoholic specimens; but, in drawing them, the spirit was allowed to evaporate, so that the details could be properly seen.

PLATE I.

- Fig. 1. *Ophioglypha sinensis*.
 " 2. " " under side of sixth arm-joint, to show the position of the tentacle-scales.
 Figs. 3, 4. *Ophioglypha falcifera*; arm-spines of the basal joints.
 Fig. 5. " " base of the arm in profile, with edge of disk and comb of papillæ.
 " 6. *Ophioglypha acervata*; under side of an arm-joint. (The lobe on the outer side is not made distinct enough.)
 " 7. *Ophiocnida olivacea*.
 " 8. " " under side of an arm-joint showing the spine-like tentacle-scale.
 " 9. *Ophiocnida Putnami*. The little organs at the outer corners of the mouth-slits are the mouth-tentacles.
 " 10. *Ophiothela tigris*.
 " 11. " " enlarged view of the parts near the mouth, and of the first and second under arm-plates, and the first set of arm-spines.
 " 12. *Ophiothela tigris*. One of the pad-like side arm-plates bearing its arm-spines.
 " 13. *Ophiomastix janualis*.
 " 14. " " under side of an arm-joint, showing the small and great tentacle-scales.
 " 15. *Ophiomaza caecotica*.
 " 16. *Astrogomphus vallatus*.
 " 17. " " first sets of arm-spines magnified; *a*, head of the genital-slit.
 " 18. " " a piece near the tip of the arm, highly magnified, showing the granulations on the arm-joints, and the ridges of larger granules, between the joints, on which stand the hooks.
 " 19. *Ophiocreas lumbricus*.
 " 20. " " an angle of the mouth seen from within, showing the column of teeth, and on each side a bunch of granules or small papillæ, beyond which appear the two pairs of mouth-tentacles.
 " 21. *Ophiocreas lumbricus*; an arm-joint showing the thorny arm-spines.

PLATE II.

- Figs. 1, 2. *Ophiomusium eburneum*, from above and below.
 Fig. 3. " " arm from the side.
 " 4. *Ophiomitra valida*.
 " 5. " " an arm-spine.
 " 6. " " thorny stumps from the disk.
 " 7. *Ophiochondras convolutus*.
 " 8. *Ophiothamnus vicarius*.
 " 9. " " an arm-joint from below.
 " 10. *Ophiocten depressum*.
 Figs. 11, 12. *Ophiomyces frutescens*, from above and below.
 Fig. 13. " " arm-joint in section showing the retracted tentacles and the arrangement of the arm-spines.

Harvard MCZ Library

3 2044 066 300 658