THE
BOTANIST'S POCKET BOOK

W. R. HAYWARD

1st ed.

RB6/400

Library
of the
University of Toronto

1st edition of a book that has
been through 19 or more editions
and was in print in 1939, or ² later.
Not in Cat. B.19, nor in B.19. (Nat. Hist.)

THE
BOTANIST'S POCKET-BOOK.

CONTAINING IN A TABULATED FORM THE
CHIEF CHARACTERISTICS OF
BRITISH PLANTS.

WITH
THE BOTANICAL NAME, COMMON NAME, SOIL OR SITUATION,
COLOUR, GROWTH, AND TIME OF FLOWERING OF EVERY
PLANT, ARRANGED UNDER ITS OWN ORDER.

BY W. R. HAYWARD.

LONDON:
BELL & DALDY, YORK STREET, COVENT GARDEN.
1872.

WILLIAM CLOWES & SONS

PRINTED BY WILLIAM CLOWES & SONS, STAMFORD STREET
AND CHARING CROSS.

LONDON:

PRINTED BY WILLIAM CLOWES AND SONS, STAMFORD STREET
AND CHARING CROSS.

WILLIAM CLOWES & SONS

PRINTED BY WILLIAM CLOWES & SONS, STAMFORD STREET
AND CHARING CROSS.

INTRODUCTION.

HAVING in my botanical rambles frequently found the want of a book of reference to determine some doubtful point, I have endeavoured in the following pages, to compress into the smallest compass such information, as will enable any one tolerably conversant with British flowering plants, to determine their species and varieties.

I venture to hope that my little book will prove useful where a larger book would be an incumbrance, by affording information to the tyro, and also by refreshing the memory of the more advanced botanist, who, by examining on the spot any doubtful plant, may be saved the trouble of carrying home specimens of little value. It is not intended as a book for the study, nor as a rival to the many excellent and complete manuals of our leading botanists ; but I trust it will be accepted for what it is, viz., "A Botanist's Pocket Book."

Although the book is chiefly intended for the more advanced botanist, a list of the characteristic features of the different orders has been added for the benefit of the beginner. I have not attempted to tabulate them, but only to give a short description of the general appearance of the plants, their habit of growth, the usual characteristics of their flowers, and the marks by which they may most generally be distinguished from those of other orders.

I have tabulated the Genera for the use of beginners, and have therefore used the more prominent parts of the plants, such as the

leaves, petals, &c., and omitted, except where absolutely necessary, the more intricate differences of the fruit, &c.

In describing the individual plants, I have used as many abbreviations as could satisfactorily be made, and I believe they are such as the reader will quickly become acquainted with.

In the first column of the left-hand page I have given the scientific name of each plant, with its varieties. These I have adopted from the "London Catalogue," and, with few exceptions, I have arranged them in the order in which they are there classified. Several varieties there mentioned I have omitted, because on reference to our leading Floras, I have either failed to find them described, or their differences are so small as not to warrant their insertion.

In the next column will be found the English names—next their place of growth, colour, general height, duration, and time of flowering. In the English names I have followed those given in "Hooker and Arnott's Flora."

The colour I have given is that of the general or prevailing colour of the corolla. The height and time of flowering naturally vary with the soil, situation, and climate.

On the right-hand page will be found the characters which distinguish the species and varieties from their type. In some cases I have divided the species into sub-species, which may be recognized by the prefixes *a*, *b*, &c. These again are sometimes divided into varieties.

The varieties of several plants I have vainly endeavoured to compress into such limits as are consistent with the character of the book. These belong to the Genera *Rubus*, *Rosa*, *Hieracium*, and *Salix*. They are a study of themselves, and the tyro must not expect to name his specimens by a cursory examination, but must diligently study them by the aid of such books as have treated of them more fully.

I would refer the student of the Rubi to Professor Babington's "British Rubi." The Genus *Rosa* has been ably systematized by

---

Mr. J. G. Baker in the "Linnæan Journal," xi. 197. Mr. Backhouse's "Monograph of British Hieracia" may be advantageously consulted ; and to the Salices, Symes' "English Botany" will be a safe guide.

I have added a short list of plants found in several parts of the kingdom, which have no claim to be indigenous, or even as yet to be naturalized. They are in most cases escapes from gardens, or have been introduced in seed or ballast.

I trust that my little book will supply a want which the various Floras already published do not provide for, and that the botanist reserving deeper study for his library may, when in the field or on the mountain, find it a useful and portable companion.

EXPLANATION OF ABBREVIATIONS.

! following the habitat of a plant denotes that it is very rare.
When a plant is placed in brackets [ ], it denotes that it is a naturalized plant.

LEFT-HAND PAGE.

Third Column—*Soil or Situation.*

Alp.— <i>Alpine.</i>	dp.— <i>damp.</i>	lak.— <i>lakes.</i>	san.— <i>sandy.</i>
bar.— <i>barren.</i>	E.— <i>England.</i>	mar.— <i>marshes.</i>	Sc.— <i>Scotland.</i>
bks.— <i>banks.</i>	fie.— <i>fields.</i>	mea.— <i>meadows.</i>	sh.— <i>shores.</i>
bus.— <i>bushy.</i>	grav.— <i>gravelly.</i>	moi.— <i>moist.</i>	sha.— <i>shady.</i>
calc.— <i>calcareous.</i>	hea.— <i>heaths.</i>	mnt. } — <i>mountain.</i>	thick.— <i>thickets.</i>
ch.— <i>chalky.</i>	hed.— <i>hedges.</i>	mount. }	Tr.— <i>trees.</i>
cl.— <i>cliffs.</i>	hil.— { <i>hills.</i>	mud.— <i>muddy.</i>	was.— <i>waste.</i>
co.— <i>coast.</i>	} <i>hilly.</i>	past.— <i>pastures.</i>	wds.— <i>woods.</i>
cop.— <i>copses.</i>	I.— <i>Ireland.</i>	riv.— <i>rivers.</i>	wo.— <i>woods.</i>
dit.— <i>ditches.</i>	Ir.— <i>Irish.</i>	roc.— { <i>rocks.</i>	
		} <i>rocky.</i>	

Fourth Column—*Colour.*

B.— <i>blue.</i>	Gr.— <i>green.</i>	R.— <i>red.</i>	W.— <i>white.</i>
Br.— <i>brown.</i>	Li.— <i>lilac.</i>	Ro.— <i>rose.</i>	Y.— <i>yellow.</i>
Cm.— <i>cream.</i>	O.— <i>orange.</i>	Sc.— <i>scarlet.</i>	d.— <i>dark.</i>
Cr.— <i>crimson.</i>	Pk.— <i>pink.</i>	Vi.— <i>violet.</i>	p.— <i>pale.</i>
Fl.— <i>flesh.</i>	Pu.— <i>purple.</i>		

Small letters preceding a capital denote that the predominating colour of the flower is that denoted by the capital, tinged, striped, &c., with the colour designated by the small letter; as—pk W. means pinkish white; gr Y. means greenish yellow, &c.

Fifth Column—*Growth.*

Cl.— <i>climbing.</i>	ft.— <i>feet.</i>
Cr.— <i>creeping.</i>	in.— <i>inches.</i>
flo.— <i>floating.</i>	

Where a height is given to a prostrate plant, it refers to the average length of its stem.

Sixth Column—*Duration.*

A.— <i>annual.</i>	Shb.— <i>shrub.</i>
B.— <i>biennial.</i>	T.— <i>tree.</i>
P.— <i>perennial.</i>	

Seventh Column—*Time of flowering.*

- 1—*January.*
2—*February, &c.*

RIGHT-HAND PAGE.

ac.— <i>acute</i> .	cylind.— <i>cylindrical</i> .	interr.— <i>interrupted</i> .	retic.— <i>reticulate</i> .
ach.— <i>achene</i> .	decid.— <i>deciduous</i> .	invol.— <i>involute</i> .	revol.— <i>revolute</i> .
acum.— <i>acuminate</i> .	decumb.— <i>decumbent</i> .	irreg.— <i>irregular</i> .	rhomb.— <i>rhomboid</i> .
aggre.— <i>aggregate</i> .	decurr.— <i>decurrent</i> .	L., l., ls.— $\left\{ \begin{array}{l} \text{leaf.} \\ \text{leaves.} \end{array} \right.$	rostr.— <i>rostrate</i> .
alt.— <i>alternate</i> .	delt.— <i>deltoid</i> .	lacin.— <i>lacinate</i> .	rt.— <i>root</i> .
amplex.— <i>amplexical</i> .	dent.— <i>dentate</i> .	lanc.— <i>lanceolate</i> .	runc.— <i>runcinate</i> .
ang.— <i>angular</i> .	dentic.— <i>denticulate</i> .	lat.— <i>lateral</i> .	sagit.— <i>sagittate</i> .
ann.— <i>annual</i> .	dichot.— <i>dichotomous</i> .	lfts.— <i>leaflets</i> .	scab.— <i>scabrous</i> .
anth.— <i>anther</i> .	diff.— <i>diffuse</i> .	lin.— <i>linear</i> .	seg.— <i>segment</i> .
apic.— <i>apiculate</i> .	digit.— <i>digitate</i> .	lyr.— <i>lyrate</i> .	sep.— <i>sepal</i> .
approx.— <i>approximate</i> .	dist.— <i>distant</i> .	marg.— <i>margins</i> .	serr.— <i>serrate</i> .
arc.— <i>arcuate</i> .	divar.— <i>divaricate</i> .	memb.— <i>membra- neous</i> .	sess.— <i>sessile</i> .
aris.— <i>aristate</i> .	ellip.— <i>elliptical</i> .	mucro.— <i>mucronate</i> .	simp.— <i>simple</i> .
asc.— <i>ascending</i> .	elong.— <i>elongate</i> .	obt.— <i>obtuse</i> .	sinu.— <i>sinuate</i> .
atten.— <i>attenuate</i> .	emarg.— <i>emarginate</i> .	opp.— <i>opposite</i> .	sol.— <i>solitary</i> .
aur.— <i>auricle</i> .	ensi.— <i>ensiform</i> .	ov.— <i>oval</i> .	sp.— <i>species</i> .
axill.— <i>axillary</i> .	ent.— <i>entire</i> .	palm.— <i>palmate</i> .	spath.— <i>spathulate</i> .
bar.— <i>barren</i> .	falc.— <i>falcate</i> .	pan.— <i>panicled</i> .	spher.— <i>spherical</i> .
cal.— <i>calyx</i> .	fert.— <i>fertile</i> .	part.— <i>partite</i> .	spic.— <i>spicate</i> .
campan.— <i>campanulate</i> .	fibr.— <i>fibrous</i> .	pect.— <i>pectinate</i> .	spin.— <i>spinulose</i> .
capill.— $\left\{ \begin{array}{l} \text{capillary.} \\ \text{capillaceo.} \end{array} \right.$	fila.— <i>filament</i> .	pedi.— <i>pedicel</i> .	spread.— <i>spreading</i> .
capit.— <i>capitate</i> .	fili.— <i>filiform</i> .	pedunc.— <i>peduncle</i> .	st.— <i>stem</i> .
caps.— <i>capsule</i> .	Fl., fls.— $\left\{ \begin{array}{l} \text{flower.} \\ \text{flowers.} \end{array} \right.$	pelt.— <i>peltate</i> .	sta.— <i>stamen</i> .
carp.— <i>carpel</i> .	flex.— <i>flexuous</i> .	pend.— <i>pendulous</i> .	stig.— <i>stigma</i> .
catk.— <i>catkin</i> .	flo.— <i>floating</i> .	per.— <i>perennial</i> .	stip.— $\left\{ \begin{array}{l} \text{stipule.} \\ \text{stipuled.} \end{array} \right.$
cil.— <i>ciliate</i> .	foll.— <i>follicle</i> .	perfol.— <i>perfoliate</i> .	subu.— <i>subulate</i> .
cli.— <i>climbing</i> .	fr.— <i>fruit</i> .	peri.— <i>perianth</i> .	succ.— <i>succulent</i> .
con.— $\left\{ \begin{array}{l} \text{conical.} \\ \text{conic.} \end{array} \right.$	fro.— <i>frond</i> .	persis.— <i>persistent</i> .	tend.— $\left\{ \begin{array}{l} \text{tendrill.} \\ \text{tendrilled.} \end{array} \right.$
conn.— <i>connate</i> .	fusi.— <i>fusiform</i> .	pet.— <i>petal</i> .	term.— <i>terminal</i> .
contig.— <i>contiguous</i> .	gl.— <i>glume</i> .	pinn.— <i>pinnate</i> .	tern.— <i>ternate</i> .
contin.— <i>continuous</i> .	glab.— <i>glabrous</i> .	pinnfid.— <i>pinnatifid</i> .	triang.— <i>triangular</i> .
cor.— <i>corolla</i> .	gland.— <i>glandular</i> .	procumb.— <i>procumbent</i> .	trique.— <i>triquetrous</i> .
cord.— <i>cordate</i> .	glauc.— <i>glaucous</i> .	prost.— <i>prostrate</i> .	trunc.— <i>truncate</i> .
coria.— <i>coriaceous</i> .	glob.— <i>globose</i> .	pubesc.— <i>pubescent</i> .	tub.— <i>tuberous</i> .
corym.— $\left\{ \begin{array}{l} \text{corymb.} \\ \text{corym-} \\ \text{bose.} \end{array} \right.$	glu.— <i>glume</i> .	pyram.— <i>pyramidal</i> .	tuberc.— <i>tubercu- late</i> .
cr.— <i>creeping</i> .	hemis.— <i>hemispheri- cal</i> .	rac.— $\left\{ \begin{array}{l} \text{raceme.} \\ \text{racemose.} \end{array} \right.$	turb.— <i>turbinate</i> .
cun.— <i>cuneate</i> .	herb.— <i>herbaceous</i> .	rad.— <i>radical</i> .	umbell.— <i>umbellate</i> .
	imbric.— <i>imbricate</i> .	reflex.— <i>reflexed</i> .	unilat.— <i>unilateral</i> .
	infl.— <i>inflated</i> .	reni.— <i>reniform</i> .	urc.— <i>urceolate</i> .
	inflex.— <i>inflexed</i> .		vertic.— <i>verticillate</i> .

SYNOPSIS OF THE NATURAL ORDERS.

Class I.—Dicotyledons.

Plants with leaves usually with netted veins—Parts of the corolla 4-5 or their multiples.

DIV. I —PLANTS WITH BOTH CALYX AND COROLLA.

* POLYPETALÆ.—*Corolla of free, separate petals.*

A. Stamens hypogynous.

1. RANUNCULACEÆ.—Sta. 12 or more—Pet. 5 or more—Sep. mostly 5, distinct—Fr. of many distinct carpels, or one-seeded achenes—*Herbs mostly with alt. leaves and regular fls.*

2. BERBERIDACEÆ.—Sta. 6—Pet. and sep. 6—Fruit a berry—*A shrub with 3 partite spines, alt. leaves, and pendulous flowers.*

3. NYMPHÆACEÆ.—Sta. many, gradually passing into petals, and these into sepals—Ovary many celled, many seeded—*Aquatic plants with floating leaves and sol. fls.*

4. PAPAVERACEÆ.—Sta. indefinite—Pet. 4—Sep. 2, deciduous—Ovary 1 celled with membranous divisions—*Herbs with milky juice, alternate, exstipulate leaves and regular flowers, usually nodding in bud.*

5. FUMARIACEÆ.—Sta. 6 in two bundles—Pet. 4 irregular—Sep. 0. or 2, deciduous—Ovary 1 celled—*Herbs with usually much divided exstipulate leaves, and racemed irregular flowers (usually purplish).*

6. CRUCIFERÆ.—Sta. 6 (4 long and 2 short)—Pet. 4, cruciate—Sep. 4—Ovary 1-2 celled—Fruit a pouch or pod—*Herbs with alternate, exstipulate leaves, and racemed, regular fls. (usually yellow or white).*

7. RESEDACEÆ.—Sta. many (10-24)—Pet. 4-7 irregular—Sep. 4-5 persistent—Ovary 1-2 lobed, 1 celled—*Herbs (rarely shrubs) with alternate, exstipulate leaves and spiked, irregular, greenish flowers.*

8. CISTACEÆ.—Sta. many—Pet. 5, twisted in bud—Sep. 5, 2 outer smaller—Ovary 3 partite, 1 celled—*Herbs or low shrubs with entire, opposite leaves and mostly racemed, regular flowers (usually yellow).*

9. VIOLACEÆ.—Sta. 5—Pet. 5 unequal, lower spurred—Sep. 5, persistent—Ovary 3 partite, 1 celled—*Small herbs with alternate, stipuled leaves and axillary, irregular flowers.*

10. DROSERACEÆ.—Sta. 5 or 10—Pet. 5—Sep. 5 imbricate—Ovary 1 celled—*Low marsh herbs with (generally) radical, glandular leaves and cymose, regular flowers whitish).*

11. POLYGALACEÆ.—Sta. 8, in 2 bundles—Pet. 3-5 unequal—Sep. 5 inner petaloid—Capsule 2 celled—*Herbs with alternate, exstipulate, simple leaves and racemose, irregular flowers.*

12. FRANKENIACEÆ.—Sta. 4-6—Pet. 4-6, clawed—Sep. 4-6, combined into a tube at base—Ovary 1 celled, 2-4 valved—*A low branched herb with opposite, exstipulate leaves and small, axillary, red flowers.*

13. ELATINACEÆ.—Sta., pet., and sep. 2-5—Capsule 3-5 valved—*Small aquatic herbs with spatulate, stipulate leaves and minute, axillary flowers.*

14. CARYOPHYLLACEÆ.—Sta. 8-10—Pet. 4-5—Sep. 4-5—Capsule 1 celled, opening at top with teeth—*Herbs mostly with jointed stems, opposite ls., and regular flowers (white or red).*

15. LINACEÆ.—Sta. 4-5—Pet. 4-5, twisted in bud, clawed, fugacious—Sep. 4-5 persistent—Capsule 3-5 celled—*Herbs with entire, exstipulate leaves and mostly cymose, regular flowers (generally blue).*

16. MALVACEÆ.—Sta. many, united into a tube—Pet 5, twisted in bud—Sep. 5—Ovary many celled—*Herbs with alternate, stipuled leaves and axillary flowers (red).*

17. TILIACEÆ.—Sta. many—Pet. and sep. 4-5—Ovary 2-10 celled—*Trees with alternate, stipuled leaves and cymose, inconspicuous, regular flowers.*

18. HYPERICACEÆ.—Sta. many, united in bundles—Pet. 4-5, twisted in bud—Sep. 4-5, with glandular dots—Capsule 3-5 celled—*Shrubs or mostly herbs with opposite, exstipulate, often glandular, dotted leaves and mostly cymose, regular flowers (yellow).*

19. ACERACEÆ.—Sta. 8, inserted on disk—Pet. 4-9, imbricate in bud—Sep. 4-9—Fruit a Samara—*Trees with opposite leaves and small, green regular flowers.*

20. GERANIACEÆ.—Sta. 5-10—Pet. 5 clawed—Sep. 5, persistent—Carpels 5 surrounding a long beak—*Herbs with stipuled, lobed leaves and handsome, regular fls. (mostly red).*

21. BALSAMINACEÆ.—Sta. 5—Pet. 5, irreg., united in pairs—Sep. 5 or 3 by abortion—Capsule bursting with elastic valves—*Succulent herbs with simple, alternate leaves and axillary, irregular yellow flowers.*

22. OXALIDACEÆ.—Sta. 10—Pet 5, cohering at base—Sep. 5—Carpel 5 celled, many seeded—*Low herbs with radical, generally trifoliate leaves and axillary, regular flowers.*

b. Stamens perigynous or epigynous.

23. CELASTRACEÆ.—Sta. 4-5, alternating with petals—Sep. 4-5—Capsule fleshy—*Trees or shrubs with simple, opposite leaves and small, regular flowers in axillary cymes.*

24. RHAMNACEÆ.—Sta. 4-5, opposite the petals—Sep. 4-5—Ovary superior, 3 celled—*Shrubs with simple leaves and small, greenish, regular flowers.*

25. LEGUMINOSÆ.—Sta. generally 10—Pet. 5, irregular, papilionaceous—Sep. 4-5, combined—Fruit a pod—Ovary superior—*Herbs or shrubs with alternate, stipuled leaves, mostly ternate or pinnate, sometimes tendrilled, and irregular flowers.*

26. ROSACEÆ.—Sta. indefinite—Pet. 5, equal—Sep. 4-5—Ovary superior, 2 celled—*Shrubs or herbs with stipuled, alternate leaves and showy (mostly red or yellow) flowers.*

27. ONAGRACEÆ.—Sta. 2-4 or 8—Pet. 2-4, twisted in bud—Sep. 2-4—Ovary inferior, 1-4 celled, many seeded—*Herbs with mostly entire leaves and regular, showy flowers (generally red).*

28. HALORAGIACEÆ.—Sta. 1-2-4 or 8—Pet. generally 0—Sep. 2-4—Ovary inferior, 1-4 celled—*Aquatic herbs with whorled leaves and minute flowers.*

29. LYTHRACEÆ.—Sta. as many or twice as many as pet.—Pet. 3-6, crumpled in bud—Sep. 3-6—Ovary superior—*Herbs with opposite or whorled, entire leaves and showy, regular flowers.*

30. TAMARISCACEÆ.—Sta. 4 or more—Pet. 4-5—Sep. 4-5—*Shrub with minute scale-like leaves and small, regular flowers in lateral spikes.*

31. CUCURBITACEÆ.—Sta. 3.—Pet. 5.—Sep. 5-cleft—Ovary inferior—Fr. fleshy—*A climbing herb, with tendrilled leaves, and flowers in axillary racemes.*

32. PORTULACACEÆ.—Sta. 3-5.—Pet. 5.—Sep. 2.—Ovary superior, 1 celled—*A low succulent herb, with opposite leaves and small flowers.*

33. ILLECEBRACEÆ.—Sta. 1-5.—Pet. 4-5, or 0. in Gen. I.—Sep. 4-5.—Ovary superior—*Small branching herbs, with entire sessile leaves and small flowers.*

34. SCLERANTHACEÆ.—Sta. 1-10.—Pet. 0.—Cal. 4-5 lobed—Ovary superior—*Small, low, tufted herbs, with opposite, connate leaves, and small, green, axillary flowers.*

35. GROSSULARIACEÆ.—Sta. 4-5, alternate with pet.—Sep. 4-5.—Ovary inferior—Fruit a pulpy berry—*Shrubs with lobed leaves and small greenish flowers.*

36. CRASSULACEÆ.—Sta. 3-20, inserted on base of sep.—Pet. and sep. 4-12—Ovaries superior, as many as pet.—*Mostly low, succulent herbs, with small, starry, regular flowers.*

37. SAXIFRAGACEÆ.—Sta. 5-10.—Pet. 4-5.—Sep. 4-5.—Ovary superior—*Herbs (mostly mountainous) with various leaves and regular flowers.*

38. ARALIACEÆ.—Sta. 5.—Pet. 5.—Cal. 3-5 cleft—Ovary inferior—Fruit a berry—*A low herb and a climbing shrub, with green flowers.*

39. CORNACEÆ.—Sta. 4-5.—Pet. 4-5.—Cal. 0. or 5 cleft—Ovary inferior—*Herbs or shrubs, with opposite leaves and small flowers.*

40. UMBELLIFERÆ.—Sta. 5.—Pet. 5.—Sep. 5 or 0.—Ovary inferior—Fruit of 2 adhering carpels—*Herbs with mostly compound leaves, and small, white, umbelled flowers.*

41. LORANTHACEÆ.—Sta. 4.—Pet. 4.—Sep. 0.—Ovary inferior—Fruit a pulpy berry—*A parasitic shrub, with entire fleshy leaves and inconspicuous flowers.*

Plants with petals apparently more or less united at base occur in Ord. v., Ord. xi., Ord. xxx., Ord. xxxi., Ord. xxxii., and Ord. xxxvi. (Gen. Cotyledon).

* * MONOPETALÆ.—*Corolla monopetalous.*

† *Ovary inferior.*

a. *Stamens on the corolla.*

(See also Ord. xlvii., Gen. Lobelia.)

42. CAPRIFOLIACEÆ.—Sta. 4-5, alternate with lobes of cor.—Cal. 5 cleft—Fruit a berry—*Mostly shrubs, with opposite leaves and showy (sometimes irregular) flowers.*

43. RUBIACEÆ.—Sta. 3-5, alternate with lobes of cor.—Cal. 4-5 cleft—Carpels 2 celled—*Herbs with whorled (sometimes prickly) leaves and small, regular flowers.*

44. VALERIANACEÆ.—Sta. 1 or 3.—Cor. 5 cleft—Cal. toothed—Ovary 1 celled—*Herbs with opposite leaves and small (sometimes irregular) flowers.*

45. DIPSACEÆ.—Sta. 4.—Cor. 4-5 lobed—Cal. enclosed in a scarious involucre.—Anthers free—Seeds pendulous—*Herbs with capitate involucre heads (mostly blue).*

46. COMPOSITÆ.—Sta. 4-5.—Cor. tubular or ligulate—Cal. 0.—Anthers mostly connate—Seeds erect—*Herbs with various leaves, and flowers mostly white or yellow, collected into compound heads.*

b. *Stamens on the ovary.*

47. CAMPANULACEÆ.—Sta. 5.—Cor. and cal. mostly 5 lobed—Ovary 2-8 celled—*Herbs with milky juice; alternate, entire leaves, and showy (mostly blue) regular flowers.*

48. ERICACEÆ.—(Gen. *Vaccinium*)—Sta. 8-10.—Cor. and cal. 4-5 lobed—Fruit a fleshy berry, 4-5 celled—*Low shrubs (mostly mountainous), with alternate ls. and sol. regular fls. (mostly white or red).*

† † Ovary superior.

a. Stamens free from corolla.

48. ERICACEÆ (except *Vaccinium*).—Sta. 5-8-10—Cor. and cal. 4-5 cleft—Fruit 3-5 celled—*Shrubs with opposite or whorled evergreen leaves, and small, often showy, regular flowers.*

b. Stamens on the corolla.

(See Ords. lxiii. and lxiv., Gen. *Littorella*.)

49. AQUIFOLIACEÆ.—Sta. 4-5, alternate with lobes of cor.—Cal. 4-5 cleft—Fruit a berry—*Shrub with coriaceous leaves and small, greenish regular flowers.*

50. OLEACEÆ.—Sta. 2—Cor. and cal. 4 cleft, sometimes 0—Fruit a berry or samara—*Trees or shrubs, with opposite leaves and small, regular flowers.*

51. APOCYNACEÆ.—Sta. 5—Cor. and cal. 4-5 cleft—Fruit of 2 follicles—*Shrubs with opposite, entire leaves, and showy, regular flowers (purple).*

52. GENTIANACEÆ.—Sta. 4-10, alternate with lobes of cor.—Cal. 4-10 lobed—Fruit a capsule—Ovary 1-2 celled—*Herbs with opposite, entire leaves and solitary, regular, mostly showy, flowers.*

53. POLEMONIACEÆ.—Sta. 5—Cor. and cal. 5 lobed—Fruit a capsule—Ovary 3 celled—*Herbs with pinnate leaves and showy, regular flowers.*

54. CONVULVULACEÆ.—Sta. 4-5—Cor. 4-5 lobed—Cal. of 5 sepals—Fruit a capsule—Ovary 2-4 celled—*Herbs (generally climbing) with alternate leaves (o. in *Cuscuta*), and mostly showy, regular flowers.*

55. SOLANACEÆ.—Sta. 4-5—Cor. and cal. 5 cleft—Fruit a berry—Ovary 2 celled—*Herbs with alternate leaves, and axillary cymes of regular flowers.*

56. SCHROPHULARIACEÆ.—Sta. 4; 2 longer than others—Cor. 4-5 lobed, mostly 2 lipped—Cal. 4-5 lobed—Fruit a capsule—Ovary 2 celled—*Herbs with various leaves and irregular, generally showy, flowers.*

57. OROBANCHACEÆ.—Sta. 4; 2 longer than others—Cor. 4-5 cleft, gaping—Cal. 4-5 cleft—Ovary 1 celled—*Leafless, scaly herbs, with mostly brownish flowers.*

58. VERBENACEÆ.—Sta. 4; 2 longer than others—Cor. tubular, usually 2 lipped—Cal. tubular—Fruit a capsule of 4 nutlets—Ovary 4 celled—*A herb with opposite leaves and small, irregular flowers.*

59. LABIATÆ.—Sta. mostly 4 (rarely 2); 2 longer than others—Cor. mostly 2 lipped—Cal. 2 lipped, or 5 cleft—Fruit of 4 nutlets—Ovary 4 celled—*Herbs mostly with square stems and opposite leaves, and irregular (generally small) flowers, in whorls or cymes.*

60. BORAGINACEÆ.—Sta. 5—Cor. and cal. 5 lobed—Fruit of 4 nutlets—Ovary 4 celled—*Herbs with alternate (usually scabrous) leaves, and spikes of showy, regular flowers.*

61. LENTIBULARIACEÆ.—Sta. 2—Cor. 2 lipped—Cal. 5 part or 2 lipped—Fruit a many-seeded capsule—Ovary 1 celled—*Marsh herbs with radical, entire leaves, or multipartite, floating leaves with bladders, and irregular, rather showy, flowers.*

62. PRIMULACEÆ.—Sta. 4-7, generally opposite segments of cor.—Cor. and cal. 4-7 cleft—Fruit a many-seeded capsule—Ovary 1 celled—*Herbs with various (generally radical) leaves, and mostly showy, regular flowers.*

63. PLUMBAGINACEÆ.—Sta. 5, opposite petals, hypogynous—Cor. of 5 petals united at base—Cal. tubular, scarious—Ovary 1 celled, 1 seeded—*Herbs mostly maritime, with radical or alternate leaves, and mostly blue, regular flowers.*

64. PLANTAGINACEÆ.—Sta. 4, hypogynous in Gen. II.—Cor. 4 lobed, scarious—Cal. 4 cleft—Ovary 2-4 celled (1 celled in Gen. II.), many seeded—*Herbs with generally radical, entire leaves, and spikes of small, greenish flowers.*

(The petals are free, or apparently so, in Ord. xlviii., Genera *Pyrola* and *Monotropa*, and in Ord. lxiii.)

[Plants belonging to the orders in Division I., but which are, or appear to be, wanting in either calyx or corolla, occur in the following orders:—

- Ord. i.—Genera *Clematis*, *Thalictrum*, *Anemone*, and *Caltha*.
- Ord. vi.—Genera *Cardamine* (Sp. 4), *Senebiera* (Sp. 2), and *Lepidium* (Sp. 5).
- Ord. xiv.—Genera *Sagina* (Sp. 1), and *Cherleria*.
- Ord. xxvi.—Genera *Alchemilla* and *Poterium*.
- Ord. xxvii.—Gen. *Isnardia*.
- Ord. xxviii.—All the genera.
- Ord. xxix.—Gen. *Peplis*.
- Ord. xxxiii.—Gen. *Herniaria*.
- Ord. xxxiv.
- Ord. xxxvii.—Gen. *Chrysosplenium*.]

DIV. II.—PLANTS WITH EITHER COROLLA, CALYX, OR BOTH ABSENT.

A. Flowers with the corolla absent, mostly with stamens and pistils in same flower.

* *Ovary superior.*

65. AMARANTHACEÆ.—Flowers 1 sexual—Sta. 3-5, opposite sep.—Fruit indehiscent—*Herbs with exstipulate leaves, and inconspicuous, green flowers.*

66. CHENOPODIACEÆ.—Flowers 1-2 sexual—Sta. mostly 5 (1 or 2 in *Salsola*), opposite sepals—Fruit indehiscent—*Herbs with exstipulate leaves, or sometimes with leafless, jointed stems, and small flowers.*

67. POLYGONACEÆ.—Flowers mostly 2 sexual—Sta. 5-8—Sep. 3-6—Fruit indehiscent, usually enclosed in the sepals—*Herbs with sheathing stipules, alternate leaves and small flowers.*

68. ELEAGNACEÆ.—Flowers 1 sexual (male flowers in catkins)—Sta. 4-8—Sep. of male flc. 3-4—Fruit indehiscent—*A shrub with silvery scales, alternate, entire, exstipulate leaves and small flowers.*

69. THYMELACEÆ.—Flowers 2 sexual—Sta. 8—Cal. 4 lobed—Fruit fleshy, indehiscent—*Shrubs, with entire, exstipulate leaves, and conspicuous, sweet-scented flowers.*

* * *Ovary inferior.*

70. SANTALACEÆ.—Sta. 3-5, opposite segments of calyx—Ovary 1 celled—*Parasitic herbs with entire, alternate leaves, and inconspicuous flowers.*

71. ARISTOLOCHIACEÆ.—Sta. 6-12, epigynous—Calyx 3 lobed or lipped—Ovary 4-6 celled—*Herbs, sometimes climbing, with alternate leaves.*

B. Plants with corolla and calyx absent, and with stamens and pistils in separate flowers.
(*Calyx present in Ord. lxiii., Genera Buxus and Mercurialis.*)

a. Flowers not in catkins.

72. EMPETRACEÆ.—Diœcious—Sta. 3—Perianth of scales, but apparently of 3 pet. and 3 sep.—Ovary 3-9 celled—*Small shrub, with evergreen, alternate leaves, and inconspicuous flowers.*

73. EUPHORBIACEÆ.—Diœcious—Sta. 1 or many—Perianth 3-4 part or 0.—Fruit separating into carpels elastically—*Herbs with entire leaves, often milky juice, and small flowers sometimes enclosed in a calyx like involucre.*

74. URTICACEÆ.—Flowers 1 sexual—Sta. 4-5, opposite sepals—Fruit indehiscent—Ovary 1 celled—*Herbs with often stinging leaves, and clustered small green flowers.*

75. ULMACEÆ.—Flowers 2 sexual—Sta. 4-8—Perianth 4-8 part—Fruit a samara—Ovary 2 celled—*Trees with alternate, distichous leaves.*

b. Flowers in catkins.

76. CUPULIFERÆ.—Sta. 5-20—Perianth 5-6 lobed—Fruit indehiscent, 1 seeded, enclosed in a coriaceous involucre or cup—*Trees with alternate stipulate leaves and small green flowers.*

77. BETULACEÆ.—Sta. 1 or more, opposite each sepal—Fruit small, indehiscent, 2 seeded, not enclosed in a cup—*Trees or shrubs with alternate leaves and small flowers.*

78. SALICACEÆ.—Sta. 1 or more to each scale—Perianth 0—Fruit many seeded, not in a cup—*Trees with alternate, simple leaves and flowers, which mostly precede the leaves.*

79. MYRICACEÆ.—Sta. 4-8—Fruit a drupe—*A small aromatic shrub, with alternate leaves and inconspicuous flowers.*

80. CONIFERÆ.—Male flowers in deciduous catkins—Female in cones—*Shrubs or trees with rigid, evergreen, linear leaves and resinous juices.*

Class II.—Monocotyledons.

Plants with leaves usually with parallel veins—Parts of the flowers generally 3 or 6.

DIV. I.—PETALOIDÆ.—Perianth petaloid (sometimes absent) coloured, not glumaceous.

* Ovary inferior.

a. Ls. with parallel veins.

81. ORCHIDACEÆ.—Flowers 2 sexual, irregular—Sta. 1-2, united to style—Fruit capsular—Ovary 1 celled—*Herbs mostly with tuberous roots and handsome flowers in spikes or racemes.*

82. IRIDACEÆ.—Flowers 2 sexual—Sta. 3—Perianth 6 part—Fruit capsular, 3 valved—Ovary 3 celled—*Herbs with narrow leaves and handsome flowers.*

83. AMARYLLIDACEÆ.—Flowers 2 sexual—Sta. 6—Perianth 6 part—Fruit capsular, 3 valved—Ovary 3 celled—*Herbs with bulbous roots, narrow leaves, and handsome flowers.*

88. HYDROCHARIDACEÆ.—Flowers 1 sexual—Sta. 3 or 6—Fruit a berry—Ovary 1 or 3-6 celled—*Aquatic herbs with floating or submerged leaves, and conspicuous flowers.*

b. Ls. with netted veins.

86. DIOSCOREACEÆ.—Fls. 1 sexual—Sta. 6—Perianth 6 part—Fruit a berry—Ovary 3 celled—*A climbing herb, with broad leaves and small flowers.*

* * Ovary superior.

84. LILIACEÆ.—Flowers 2 sexual—Sta. 6—Perianth 6 partite—Fruit a berry or capsule, mostly 3 celled—Ovary 3 celled—*Herbs with mostly narrow leaves, and showy flowers.*

85. TRILLIACEÆ.—Flowers 2 sexual—Sta. 6-10—Pet. 3-5, filiform—Sep. 3-5—Fruit a berry—*A herb with 4-5 whorled, ovate, netted-veined leaves.*

87. MELANTHIACEÆ.—Flowers 2 sexual—Sta. 6—Perianth petaloid, 6 part—Fruit capsular, 3 valved—*Herbs with linear leaves.*

89. ALISMACEÆ.—Flowers 2 sexual—Sta. 6—Perianth 6 part—Fruit of many carpels—*Aquatic plants with radical, netted-veined leaves, and mostly showy (white) flowers.*

90. NAIDACEÆ.—Flowers 1-2 sexual—Perianth of scale like sepals or 0—Sta. as many as sepals—Fruit of 1-4 carpels—*Aquatic herbs with various, floating or submerged leaves, and inconspicuous flowers.*

91. LEMNACEÆ.—Sta. 1-2—Perianth 0—Minute floating plants, with green, cellular fronds, rarely flowering.

92. ARACEÆ.—Sta. indefinite—Perianth 0—Fruit a berry—Herbs with netted-veined leaves, and flowers on a spadix enclosed in a leafy sheath.

93. TYPHACEÆ.—Sta. many—Perianth 0—Fruit drupaceous—Marsh plants with ensiform leaves, and small flowers in dense, conspicuous heads.

94. RESTIACEÆ.—Sta. 2-3—Perianth 2-6 partite, colourless—Fruit capsular—Aquatic plant, with subulate leaves, and flowers in an involucrate head.

95. JUNCACEÆ.—Sta. usually 6—Perianth 6 partite; segments scarious—Fruit capsular, 3 valved—Herbs (rushes) with narrow leaves and small, brown flowers.

DIV. II.—GLUMIFERÆ.—Flowers without perianth, glumaceous.

96. CYPERACEÆ.—Flowers 1-2 sexual—Sta. 1-3—Grassy herbs, with solid stems, entire sheaths, and flowers in spikelets.

97. GRAMINEÆ.—Flowers generally 2 sexual—Sta. usually 3—Grassy herbs, with hollow stems and split sheaths.

Class III.—Acotyledons, or Cryptogamia.

Plants without flowers, reproduced by minute spores.

98. FILICES.—Leafy plants, with the capsules on the edge, or under-side of the leaves.

99. LYCOPODIACEÆ.—Leafy plants, with the capsules axillary, or in the bracts of a cone.

100. MARSILEACEÆ.—A creeping plant, with slender leaves, and capsules in a globose receptacle at the base of the leaves.

101. EQUISETACEÆ.—Leafless plants, with whorled branches, with sheaths at the joints, and capsules in terminal spikes.

ARTIFICIAL KEYS TO THE GENERA.

ORD. I.—RANUNCULACEÆ.—(p. 2.)

A. Flowers regular.

* Corolla absent.

- Sep.* 4-6—*St. climbing* . . . . . GEN. I. CLEMATIS.
Sep. 4-5, *imbric.* . . . . . GEN. II. THALICTRUM.
Sep. 5-14, *with a 3-partite involucre some distance from*
flower . . . . . GEN. III. ANEMONE.
Sep. 5, *petaloid, coloured* . . . . . GEN. VII. CALTHA.

* * Corolla present.

- Pet.* 5-10, *scarlet—Sep.* 5, *petaloid* . . . . . GEN. IV. ADONIS.
Pet. 5—*Sep.* 5—*Flowers spicate* . . . . . GEN. V. MYOSORUS.
Pet. 5—*Sep.* 5—*Carpels in an obl. or globular head* . . . . GEN. VI. RANUNCULUS.
Pet. 5-15, *small, linear—Sep.* 5 . . . . . GEN. VIII. TROLLIUS.
Pet. 5, *small, tubular, bilabiate—Sep.* 5, *deciduous* . . . . GEN. IX. ERANTHIS.
Pet. 8-10, *small, tubular, bilabiate—Sep.* 5, *persistent* . . . GEN. X. HELLEBORUS.
Pet. 5, *spurred—Sep.* 5 . . . . . GEN. XI. AQUILEGIA.
Pet. 4—*Sep.* 4, *caducous* . . . . . GEN. XIV. ACTÆA.

B. Flowers irregular.

- Sep.* 5, *upper ones spurred—Pet.* 4, *2 upper enclosed in spur* GEN. XII. DELPHINIUM
Sep. 5, *upper ones hooded—Pet.* 2-5, *2 upper enclosed in hood* GEN. XIII. ACONITUM.

ORD. II.—BERBERACEÆ.—(p. 6.)

- Pet., sep., and sta.* 6 . . . . . GEN. I. BERBERIS.

ORD. III.—NYMPHÆACEÆ.—(p. 6.)

- Sep.* 4, *greenish—Pet. inserted on base of ovary.* . . . . GEN. I. NYMPHÆA.
Sep. 5-6, *yellow—Pet. inserted on receptacle.* . . . . GEN. II. NUPHAR.

ORD. IV.—PAPAVERACEÆ.—(p. 6.)

- Fruit ovate—Stigma* 4-20 *rayed—Style sessile* . . . . . GEN. I. PAPAVER.
Fruit oblong—Stigma 5-6 *rayed—Style short* . . . . . GEN. II. MECONOPSIS.
Fruit linear, 1-celled—Flowers small . . . . . GEN. III. CHELIDONIUM
Fruit linear, 2-celled—Flowers large . . . . . GEN. IV. GLAUCIUM.

ORD. V.—FUMARIACEÆ.—(p. 8.)

- Fruit round, 1-seeded* . . . . . GEN. I. FUMARIA.
Fruit long, many-seeded . . . . . GEN. II. CORYDALIS.

ORD. VI.—CRUCIFERÆ.—(p. 8.)

A. Fruit a pouch.

* Pouch 1 or 2 jointed, each 1-seeded.

- Pouch ang. of 2 1-seeded joints; upper fertile, sessile, ensiform, deciduous; lower obconic, pendulous.* . . . GEN. I. CAKILE.
Pouch of 2 joints; upper globose, 1-seeded, deciduous; lower abortive, stalk-like . . . GEN. II. CRAMBE.
Pouch broader than long, wrinkled, notched at base and apex; 2-jointed, each 1-seeded . . . GEN. III. CORONOPUS.
Pouch flat, oblong, with keeled valves, 1-celled, 1-seeded . . GEN. IV. ISATIS.

* * Pouch 2 valved.

a. Laterally compressed.

- Pouch obovate, notched, with winged valves; many-seeded* . GEN. V. THLASPI.
Pouch triangular, cuneate, valves keeled, not winged; many-seeded . . . GEN. VI. CAPSELLA.
Pouch elliptic, entire; valves keeled; cells 2-seeded . . . GEN. VII. HUTCHINSIA.
Pouch obcordate, emarginate; valves boat-shaped, keeled below, slightly winged above . . . GEN. VIII. TEESDALIA.
Pouch globose, notched; valves winged; cells 1-seeded—Petals unequal . . . GEN. IX. IBERIS.
Pouch ovate, notched; valves compressed, keeled; cells 1-seeded . . . GEN. X. LEPIDIUM.

b. Pouch turgid, or dorsally compressed.

- Pouch globose, turgid; valves veined; cells many-seeded* . GEN. XI. COCHLEARIA.
Pouch oval, turgid; valves boat-shaped—Styles deciduous—Cells 4-seeded . . . GEN. XII. SUBULARIA.
Pouch ovate; valves slightly convex; cells many-seeded . . GEN. XIII. DRABA.
Pouch sub-oval; valves inflated, 1-nerved; cells many-seeded GEN. XIV. CAMELINA.
Pouch compressed, pointed—Cells 1-seeded—Pet. white . . GEN. XV. KONIGA.
Pouch compressed—Cells 2-4 seeded—Pet. yellow . . . GEN. XVI. ALYSSUM.

B. Fruit a long pod.

* Pod neither beaked nor jointed.

a. Calyx equal at base.

- Pod lanc., compressed, tapering; valves flat, veinless, opening elastically; seed stalks winged* . . . GEN. XVII. DENTARIA.
Pod linear, compressed; valves veinless, opening elastically; seed stalks filiform . . . GEN. XVIII. CARDAMINE.
Pod linear, compressed; valves flat, with prominent, swelling veins—Seeds in 1 row (2 rows in Sp. 6.) . . . GEN. XIX. ARABIS.
Pod linear, terete; valves with a prominent rib—Seeds in 1 row . . . GEN. XX. BARBAREA.
Pod linear, short; valves convex, nerveless—Seeds in 2 irregular rows . . . GEN. XXI. NASTURTIIUM.
Pod linear, terete; valves convex, with 2 veins (1-veined in Sp. 4)—Seeds in 1 row . . . GEN. XXII. SISYMBRIUM.
Pod linear, quadrangular; valves with a prominent nerve—Seeds in 1 row . . . GEN. XXIII. ERYSIMUM.

b. Calyx succate at base.

- Pod compressed, or 2-edged, with a rib on each valve—Stigma 2 lobed* . . . GEN. XXIV. CHEIRANTHUS.

*Pod linear, rounded or compressed—Stigma 2 lobed, gibbous,
or horned at back* . . . . . GEN. XXV. MATTHIOLA.

*Pod quadrangular, valves nerved—Stigma 2 lobed; lobes
erect, ellip., obtuse* . . . . . GEN. XXVI. HESPERIS.

* * Pods beaked.

Pod linear, angular—Sep. erect . . . . . GEN. XXVII. BRASSICA.

Pod linear, angular—Sep. spreading—Seeds in 1 row . . . GEN. XXVIII. SINAPIS.

Pod compressed—Sep. spreading—Seeds in 2 rows . . . GEN. XXIX. DIPLTAXIS.

* * * Pods jointed.

Pods tapering, divided into 1-seeded joints . . . . . GEN. XXX. RAPHANUS.

ORD. VII.—RESEDACEÆ.—(p. 16.)

Pet. many part.—Sep. divided . . . . . GEN. I. RESEDA.

ORD. VIII.—CISTACEÆ.—(p. 16.)

Pet. 5, deciduous—Sep. 5, 2 outer smaller . . . . . GEN. I. HELIANTHEMUM.

ORD. IX.—VIOLACEÆ.—(p. 16.)

Pet. 5, unequal, lower spurred . . . . . GEN. I. VIOLA.

ORD. X.—DROSERACEÆ.—(p. 18.)

Pet., sep., and sta. 5—Ls. glandular . . . . . GEN. I. DROSERA.

ORD. XI.—POLYGALACEÆ.—(p. 18.)

Cor. irregular—Sep. 5; 2 inner petaloid . . . . . GEN. I. POLYGALA.

ORD. XII.—FRANKENIACEÆ.—(p. 18.)

Style trifid . . . . . GEN. I. FRANKENIA.

ORD. XIII.—ELATINACEÆ.—(p. 20.)

Pet., sep., and sta. 3-4 . . . . . GEN. I. ELATINE.

ORD. XIV.—CARYOPHYLLACEÆ.—(p. 20.)

A. Sepals united in a tube—Sta. 10.

Styles 2—Calyx with two opposite bracts at base . . . . . GEN. I. DIANTHUS.

Styles 2—Calyx without bracts at base . . . . . GEN. II. SAPONARIA.

Styles 3-4—Capsule 3-celled, opening at apex with 6 teeth . . . GEN. III. SILENE.

Styles 5—Petals with a scale at base . . . . . GEN. IV. LYCHNIS.

(Without a scale in Sp. 6.)

B. Sepals distinct—Sta. 4-10.

* Leaves stipuled.

Styles 5 . . . . . GEN. VII. SPERGULA.

Styles 3 . . . . . GEN. IX. SPERGULARIA.

* * Leaves not stipuled.

Sta., pet., and sep. 4—Styles 3 . . . . . GEN. V. MÖENCHIA.

Sta. 4-10—Pet. 4-5 (or 6), shorter than sep.—Styles 4-5 . . . GEN. VI. SAGINA.

Sta. 10—Pet. entire, longer than sep.—Styles 3-5 . . . . . GEN. VIII. HONKENEYA.

Sta. 8-10—Pet. entire, conspicuous—Sep. 5—Styles 2-3 . . . GEN. X. ARENARIA.

- Sta.* 3-5—*Pet.* jagged at end—*Styles* 3 . . . . . GEN. XI. HOLOSTEUM.
Sta. 10—*Pet.* bifid—*Styles* 3 . . . . . GEN. XII. STELLARIA.
Sta. 4-10—*Pet.* bifid—*Styles* 4-5 . . . . . GEN. XIII. CERASTIUM
Sta. 10—*Pet.* entire, very minute, or 0—*Styles* 3 . . . . . GEN. XIV. CHERLERIA.

ORD. XV.—LINACEÆ.—(p. 26.)

- Pet., sep., and sta.* 5 . . . . . GEN. I. LINUM.
Pet., sep., and sta. 4 . . . . . GEN. II. RADIOLA.

ORD. XVI.—MALVACEÆ.—(p. 26.)

- Involute of 3 distinct leaves* . . . . . GEN. I. MALVA.
Involute 6-9 lobed . . . . . GEN. II. ALTHÆA.
Involute 3 lobed . . . . . GEN. III. LAVATERA.

ORD. XVII.—TILIACEÆ.—(p. 26.)

- Pet. and sep.* 5—*Sta.* many . . . . . GEN. I. TILIA.

ORD. XVIII.—HYPERICACEÆ.—(p. 28.)

- Pet. and sep.* 5—*Styles* 3-5 . . . . . GEN. I. HYPERICUM.

ORD. XIX.—ACERACEÆ.—(p. 28.)

- Pet. and sep.* 5—*Sta.* 8 . . . . . GEN. I. ACER.

ORD. XX.—GERANIACEÆ.—(p. 28.)

- Carpels with a glabrous, recurved awn* . . . . . GEN. I. GERANIUM.
Carpels with a spiral awn, bearded inside . . . . . GEN. II. ERODIUM.

ORD. XXI.—BALSAMINACEÆ.—(p. 30.)

- Flowers spurred* . . . . . GEN. I. IMPATIENS.

ORD. XXII.—OXALIDACEÆ.—(p. 30.)

- Sta.* 10—*Styles* 5 . . . . . GEN. I. OXALIS.

ORD. XXIII.—CELASTRACEÆ.—(p. 30.)

- Fls. umbell.—Fruit rose-coloured* . . . . . GEN. I. EUONYMUS.

ORD. XXIV.—RHAMNACEÆ.—(p. 32.)

- Calyx 4-5 cleft, urceolate* . . . . . GEN. I. RHAMNUS.

ORD. XXV.—LEGUMINOSÆ.—(p. 32.)

A. Stamens all united—Pods continuous.

* Leaves simple.

- Cal. deeply 2-lipped, with bracts at base, nearly as long as cor.* . . . . . GEN. II. ULEX.

- Cal. slightly 2-lipped, without bracts, much shorter than cor.* . . . . . GEN. III. GENISTA.

* * Leaves ternate (or upper simple).

- Cal. 2-lipped; upper lip bifid; lower trifid* . . . . . GEN. I. SAROMANTHUS

- Cal. 5 cleft; seg. lin.* . . . . . GEN. IV. ONONIS.

- * * * Leaves pinnate. . . . . GEN. V. ANTHYLLIS.
Cal. inflated, 5 cleft . . . . .
 B. Upper stamen free, the rest united.
 * Leaves trifoliolate—Pod continuous.
Flowers in short racemes—Pod spiral (rarely falcate), many-seeded (1-seeded in Sp. 3) . . . . . GEN. VI. MEDICAGO.
Flowers in long, one-sided racemes—Pod nearly straight, 1-4 seeded . . . . . GEN. VII. MELITOTUS.
Flowers in short racemes—Pod slightly curved, many-seeded, longer than corolla . . . . . GEN. VIII. TRIGONELLA.
Flowers in capitate heads—Pod short, nearly straight, 1-4 seeded—Calyx teeth unequal . . . . . GEN. IX. TRIFOLIUM.
Flowers umbellate—Pod nearly straight, lin., many-seeded—Calyx teeth equal . . . . . GEN. X. LOTUS.
 * * Leaves pinnate, with a terminal leaflet; not tendrilled.
Pod continuous, imperfectly 2-celled; keel obtuse—Pod with the lower suture inflexed . . . . . GEN. XI. ASTRAGALUS.
Keel with a narrow tip—Pod with the upper suture inflexed . . . . . GEN. XII. OXYTROPIS.
 * * * Leaves pinnate, with a terminal leaflet; not tendrilled—Pods jointed.
Flowers umbellate, with a pinnate leaf at apex of pedunc.—Pod compressed, curved, many-jointed . . . . . GEN. XIII. ORNITHOPUS.
Flowers umbellate, without a pinnate leaf—Pod compressed, many-jointed, but scarcely contracted at joints . . . . . GEN. XIV. ARTHROLOBIUM.
Flowers umbellate, not bracteate—Pod of many crescent-shaped joints . . . . . GEN. XV. HIPPOCREPIS.
Flowers racemed—Pod flat, 1-jointed . . . . . GEN. XVI. ONOBRYCHIS.
 * * * * Leaves pinnate, without a terminal leaflet (*leaves apparently simple in Gen. XVIII., Sp. 1 and 2*); mostly tendrilled—Pod continuous.
Leaves all pinnate, tendrilled—Style filiform, hairy all round, or bearded below . . . . . GEN. XVII. VICIA.
Leaves pinnate or simple, mostly tendrilled—Style flattened upwards, hairy only on upper side . . . . . GEN. XVIII. LATHYRUS.

ORD. XXVI.—ROSACEÆ.—(p. 38.)

A. Petals distinct.

* HERBS.

a. Sepals equal.

- Pet. 5—Cal. 5 cleft—Fruit of 3-12 nuts, inserted on cal. tube* . . . . . GEN. II. SPIRÆA.
Pet. 8—Cal. 8-10 cleft—Sta. many—Fruit of many small nuts, each with a feathery awn . . . . . GEN. IV. DRYAS.
Pet. 5—Cal. 5 cleft—Nuts 2 . . . . . GEN. XI. AGRIMONIA.
 b. Sepals with alternate, smaller segments.
Pet. 5—Sta. many—Fruit of many small nuts, tipped with the long, persistent style, on a long, dry receptacle . . . . . GEN. III. GEUM.
Pet. 5, small—Sta. 5—Fruit of 5-10 small nuts on a concave, dry receptacle . . . . . GEN. V. SIBBALDIA.
Pet. 4-5—Sta. many—Fruit of many small nuts on a flat, dry receptacle . . . . . GEN. VI. POTENTILLA.

- Pet.* 5, smaller than sepals—*Sta.* many—Fruit on a conical, spongy receptacle . . . . . GEN. VII. COMARUM.
- Pet.* 5—*Sta.* many—Fruit of many small nuts, sunk in a fleshy, deciduous receptacle . . . . . GEN. VIII. FRAGARIA.
- * * SHRUBS OR TREES.
- a. Styles many.
- Pet.* 5—Fruit of many, succulent drupes, on a spongy, conical receptacle . . . . . GEN. IX. RUBUS.
- Pet.* 4-5—Fruit of many, dry nuts, enclosed in the fleshy tube of calyx . . . . . GEN. X. ROSA.
- b. Styles 1-5:
- Pet.* 5—Calyx 5 cleft, deciduous—Fruit a fleshy drupe . . GEN. I. PRUNUS.
- Pet.* 5, large—*Sep.* large, leafy—Fruit turbinate, the upper ends of the bony cells exposed . . . . . GEN. XV. MESPILUS.
- Cal.* segments acute—Fruit oval, concealing the 1-5 bony carpels . . . . . GEN. XVI. CRATÆGUS.
- Fruit a drupe, with 3-5 half-exserted seeds, not cohering at centre . . . . . GEN. XVII. COTONEASTER.
- Fruit a fleshy drupe, with 5 2-seeded cells, with cartilaginous walls . . . . . GEN. XVIII. PYRUS.
- B. Petals 0.
- Sta.* 4—*Cal.* 4 cleft, with 2-3 scales at base . . . . . GEN. XII. SANGUISORBA.
- Stamens* many . . . . . GEN. XIII. POTERIUM.
- Sta.* 1-4—*Cal.* double, 8 cleft . . . . . GEN. XIV. ALCHEMILLA.

ORD. XXVII.—ONAGRACEÆ.—(p. 50.)

- Sta.* 8—*Cal.* 4 cleft; segments free . . . . . GEN. I. EPILOBIUM.
- Sta.* 8—*Cal.* 4 cleft, combined below . . . . . GEN. II. CENOTHERA.
- Sta.* 4—*Cal.* 4 cleft . . . . . GEN. III. ISNARDIA.
- Sta.* 2—*Pet.* and *sep.* 2 . . . . . GEN. IV. CIRCÆA.

ORD. XXVIII.—HALORAGIACEÆ.—(p. 50.)

- Ls.* entire, whorled—*Sta.* 1 . . . . . GEN. I. HIPPURIS.
- Ls.* cut, whorled—*Sta.* 8 . . . . . GEN. II. MYRIOPHYLLUM.
- Ls.* ent., opposite—*Sta.* 1 . . . . . GEN. III. CALLITRICE.
- Ls.* forked, whorled—*Anth.* 12-20, sessile . . . . . GEN. IV. CERATOPHYLLUM.

ORD. XXIX.—LYTHRACEÆ.—(p. 52.)

- Cal.* cylind., tubular—Style long, filiform . . . . . GEN. I. LYTHRUM.
- Cal.* campan.—Style short . . . . . GEN. II. PEPLIS.

ORD. XXX.—TAMARISCACEÆ.—(p. 52.)

- Pet.* and *sep.* 4-5—Styles 5 . . . . . GEN. I. TAMARIX.

ORD. XXXI.—CUCURBITACEÆ.—(p. 52.)

- Diœcious*—*Cor.* and *cal.* 5 cleft—*Sta.* 5, in three bundles . GEN. I. BRYONIA.

ORD. XXXII.—PORTULACACEÆ.—(p. 52.)

- Pet.* 5, irregular—*Sep.* 2—*Sta.* 3 . . . . . GEN. I. MONTIA.

ORD. XXXIII.—ILLICEBRACEÆ.—(p. 54.)

* Capsule 1-seeded.

- Ls. opp.*—*Pet.* 0, or only like small scales . . . . . GEN. I. ILLECEBRUM.
Ls. alt.—*Pet. obl.*, as long as sepals . . . . . GEN. II. CORRIGIOLA.
Ls. opp.—*Pet. filiform*—*Stigmas* 2, sessile . . . . . GEN. III. HERNIARIA.

* * Capsule many-seeded.

- Ls. opp.*, lower in fours—*Pet.* 5, emarginate . . . . . GEN. IV. POLYCARPON.

ORD. XXXIV.—SCLERANTHACEÆ.—(p. 54.)

- Pet.* 0—*Sep.* 5—*Sta.* 10 . . . . . GEN. I. SCLERANTHUS.

ORD. XXXV.—GROSSULARIACEÆ.—(p. 54.)

- Pet.* 4-5—*Sep.* 5—*Berry* many-seeded . . . . . GEN. I. RIBES.

ORD. XXXVI.—CRASSULACEÆ.—(p. 54.)

- Pet.*, *sep.*, and *sta.* 3-4 . . . . . GEN. I. TILLÆA.
Pet. 5, distinct (4 in *Sp.* 1)—*Sta.* 10-12 . . . . . GEN. II. SEDUM.
Pet. 10-12—*Sta.* many . . . . . GEN. III. SEMPERVIVUM.
Pet. 5, united at base—*Sta.* 10 . . . . . GEN. IV. COTYLEDON.

ORD. XXXVII.—SAXIFRAGACEÆ.—(p. 56.)

- Pet.* 5—*Sta.* 10—*Caps.* 2-celled . . . . . GEN. I. SAXIFRAGA.
Pet. 0—*Caps.* 1-celled . . . . . GEN. II. CHRYSOSPENIUM.
Pet. 5—*Sta.* 5—*Caps.* 1-celled . . . . . GEN. III. PARNASSIA.

ORD. XXXVIII.—ARALIACEÆ.—(p. 58.)

- St. herbaceous*—*Corolla* monopetalous . . . . . GEN. I. ADOXA.
St. woody—*Pet.* 5 . . . . . GEN. II. HEDERA.

ORD. XXXIX.—CORNACEÆ.—(p. 60.)

- Pet.*, *sep.*, and *sta.* 4 . . . . . GEN. I. CORNUS.

ORD. XL.—UMBELLIFERÆ.—(p. 60.)

SEC. I.—Leaves simple.

- Ls. peltate*—*Stem* creeping . . . . . GEN. I. HYDROCOTYLE.
Ls. linear or *ovate*—*Stem* erect . . . . . GEN. XIX. BUPLEURUM.

SEC. II.—Leaves palmate.

- Ls. palmate*—*Fruit* round, prickly . . . . . GEN. II. SANICULA.
Ls. spiny—*Umbels* dense—*Fruit* round, scaly . . . . . GEN. IV. ERYNGIUM.
Ls. palmate—*Involucral* bracts large, coloured—*Fruit*
round, with dentate ribs . . . . . GEN. III. ASTRANTIA.

SEC. III.—Lower leaves pinnate or ternate.

a. Leaves all pinnate.

- Universal* and *partial* *involucral* bracts many . . . . . GEN. XVIII. SIUM.
Involucral bracts 0 . . . . . GEN. XVII. PIMPINELLA (*Sp.* 2).

b. Upper leaves lobed or pinnfid.

- Fruit* dorsally compressed, winged; wings with a thick,
knotted border . . . . . GEN. XXXII. TORDYLIUM.
Fruit dorsally compressed, with even wings . . . . . GEN. XXXI. HERACLEUM.

- Fruit laterally compressed, with blunt ridges—Vittæ solitary, clavate* . . . . . GEN. XIII. SISON.
- Fruit laterally compressed, with blunt ridges—Vittæ many* . . . . . GEN. XVII. PIMPINELLA (Sp. 1).
- Fruit laterally compressed, with prominent ridges—Vittæ solitary* . . . . . GEN. XII. HELOSCIADIUM (Sp. 1).
- c. Ls. ternate, or ternately compound.
- Calyx o—Fruit orbicular—Carpels winged, with 3 prominent ridges—Vittæ many* . . . . . GEN. VII. SMYRNIUM.
- Calyx teeth leafy—Fruit with obscure ridges—Vittæ solitary* . . . . . GEN. VIII. CICUTA.
- Calyx o—Fruit with 5 prominent ridges—Vittæ solitary* . . . . . GEN. IX. APIUM.
- Calyx o—Fruit orbicular, with 5 slender ridges—Vittæ o* . . . . . GEN. XIV. ÆGOPodium.
- SEC. IV.—Leaves compound.
- A. Fruit beaked.
- Beak long* . . . . . GEN. XXXVI. SCANDIX.
- Beak slender, shorter than fruit* . . . . . GEN. XXXVII. ANTHRISCUS.
- B. Fruit narrow, linear, not beaked.
- Carpels obtusely ribbed—Vittæ 1* . . . . . GEN. XXXVIII. CHÆROPHYLLUM.
- Carpels acutely ribbed—Vittæ o* . . . . . GEN. XXXIX. MYRRHIS.
- C. Fruit prickly.
- Fruit laterally compressed, not ribbed, with straight bristles* . . . . . GEN. XXXV. TORILIS.
- Fruit laterally compressed, with prominent ribs and hooked bristles* . . . . . GEN. XXXIV. CAUCALIS.
- Fruit dorsally compressed, with prominent ridges* . . . . . GEN. XXXIII. DAUCUS.
- D. Fruit ovate or oblong; glabrous or hairy.
- * Fruit dorsally compressed.
- a. Lateral wing contiguous.
- Pet. entire, acute—Lateral ridges near the outer edge of the wings* . . . . . GEN. XXX. PASTINACA.
- Pet. obcord.—Lateral ridges at the base of the wings* . . . . . GEN. XXIX. PEUCEDANUM.
- b. Lateral wings separate . . . . . GEN. XXVIII. ANGELICA.
- * * Fruit ovate, roundish; not compressed either way.
- a. Vittæ o.
- Fruit globose, with obscure ridges—Carpels scarcely separating* . . . . . GEN. XL. CORIANDRUM.
- b. Vittæ solitary.
- Fruit corky, with obscure ridges and a long, erect style* . . . . . GEN. XX. CENANTHE.
- Fruit globose, with acute ribs and a short style—Pet. obcordate* . . . . . GEN. XXI. ÆTHUSA.
- Fruit oblong, with prominent, obtuse ribs—Style short—Pet. entire* . . . . . GEN. XXII. FENICULUM.
- Fruit hairy, with thick ribs and long, reflexed styles* . . . . . GEN. XXIII. SESELI.
- c. Vittæ 2 or more.
- Fruit ellip., with sharp, almost winged ridges—Pet. obcord., white—Lfts. ovate* . . . . . GEN. XXIV. LIGUSTICUM.
- Fruit ellip., almost winged—Pet. emarginate, yellow, appendaged at base—Lfts. lanc.* . . . . . GEN. XXV. SILAUS.
- Fruit as Silaus—Pet. entire, ellip., with an incurved point—Lfts. capillary* . . . . . GEN. XXVI. MEUM.

- Fruit with triangular, sharp ridges—Lfts. fleshy* . . . GEN. XXVII. CRITHMUM.
 * * * Fruit laterally compressed.
 a. Petals entire.
Lfts. ov. or lanc.—Involucral bracts many—Fls. yellow . . . GEN. X. PETROSELINUM.
Lfts. linear—Involucral bracts few or 0—Fls. dioecious ;
Pet. yellow . . . . . GEN. XI. TRINIA.
Lfts. ov.-lanc, or submerged, capillary—Fls. white . . . GEN. XII. HELOSCIADIUM (Sp. 2).
 b. Petals emarginate.
Vittæ 0—Style long, reflexed . . . . . GEN. XIV. ÆGOPIDIUM.
Vittæ solitary—Base of style depressed . . . . . GEN. XV. CARUM.
Vittæ several—Base of style conical . . . . . GEN. XVI. BUNIUM.
 * * * * Fruit short, turgid, slightly laterally compressed.
Carpels with 5 thick, waved ridges—Vittæ 0 . . . . . GEN. V. CONIUM.
Carpels round, with slender ridges—Vittæ solitary . . . GEN. VI. PHYSOSPERNUM.

ORD. XLI.—LORANTHACEÆ.—(p. 66.)

- Dioecious—Pet. 4, fleshy* . . . . . GEN. I. VISCUM.

ORD. XLII.—CAPRIFOLIACEÆ.—(p. 68.)

* Stigmas 3, sessile.

- Leaves pinnate* . . . . . GEN. I. SAMBUCUS.
Leaves simple . . . . . GEN. II. VIBURNUM.
 * * Stigma 1, filiform.
Sta. 5—Cor. irregular . . . . . GEN. III. LONICERA.
Sta. 4—Cor. regular . . . . . GEN. IV. LINNÆA.

ORD. XLIII.—RUBIACEÆ.—(p. 68.)

- Cal. 0—Cor. rotate, 5 cleft—Fruit fleshy* . . . . . GEN. I. RUBIA.
Cal. 0—Cor. rotate, 4 cleft—Fruit dry . . . . . GEN. II. GALIUM.
Cal. 0—Cor. bell-shaped—Fruit dry . . . . . GEN. III. ASPERULA.
Calyx teeth 4-6—Cor. funnel-shaped—Fruit dry, crowned
with calyx . . . . . GEN. IV. SHERARDIA.

ORD. XLIV.—VALERIANACEÆ.—(p. 70.)

- Sta. 1—Cor. spurred—Fruit crowned with feathery pappus* . . . GEN. I. CENTRANTHUS.
Sta. 3—Cor. gibbous—Fruit crowned with feathery pappus . . . GEN. II. VALERIANA.
Sta. 3—Fruit crowned with unequal teeth . . . . . GEN. III. VALERIANELLA.

ORD. XLV.—DIPSACEÆ.—(p. 72.)

- Receptacle with spiny scales—Calyx limb entire* . . . . . GEN. I. DIPSACUS.
Receptacle scaly—Calyx limb of 4-5 bristles . . . . . GEN. II. SCABIOSA.
Receptacle hairy—Calyx limb of 8-15 bristles . . . . . GEN. III. KNAUTIA.

ORD. XLVI.—COMPOSITÆ.—(p. 72.)

SUB-ORDER I.—FLOWERS ALL LIGULATE AND PERFECT.

* Pappus 0.

GEN. XIV. LAPSANA.

* * Pappus crown-like or of erect scales.

Flowers terminal, yellow . . . . . GEN. XIII. ARNOSERIS.

Flowers axillary, blue . . . . . GEN. XV. CICHORIUM.

* * * Pappus plumose, especially of the central florets.

a. Receptacle chaffy.

GEN. V. HYPOCHÆRIS.

b. Receptacle naked.

Involucre simple, or of 8-10 connected scales . . . . . GEN. I. TRAGOPOGON.

Involucre simple, surrounded by leafy bracts . . . . . GEN. II. HELMINTHIA.

Involucre simple, with unequal, linear bracts at base . . . . . GEN. III. PICRIS.

Involucre imbricate . . . . . GEN. IV. LEONTODON.

* * * * Pappus pilose, filiform.

a. Hairs of pappus silvery (except *Crepis paludosa*).

† Fruit beaked.

Stem leafy—Fruit compressed . . . . . GEN. VI. LACTUCA.

Stem leafless—Fruit suddenly contracted and muricate above . . . . . GEN. XII. TARAXACUM.

Stem leafy—Fruit terete . . . . . GEN. XI. BARKHAUSIA.

† † Fruit not beaked.

Fruit compressed . . . . . GEN. VII. SONCHUS.

Fruit terete . . . . . GEN. IX. CREPIS.

b. Hairs of pappus rigid, turning brown.

Fls. blue—Fruit compressed . . . . . GEN. VIII. MULGEDIUM.

Fls. yellow—Fruit terete . . . . . GEN. X. HIERACIUM.

SUB-ORDER II.—FLOWERS ALL TUBULAR—STYLE THICKENED BELOW ITS BRANCHES.

* Leaves spiny, or fringed with small prickles.

All the involucre bracts spinous—Fruit compressed . . . . . GEN. XIX. CARDUUS.

All the involucre bracts spinous—Fruit angled, or 4-ribbed . . . . . GEN. XX. ONOPORDIUM.

Outer involucre bracts spinous; inner coloured, spreading . . . . . GEN. XXI. CARLINA.

* * Leaves not spiny.

a. Involucre bracts with hooked points.

GEN. XVI. ARCTIUM.

b. Involucre bracts with straight points.

Pappus longer than fruit—Anthers with bristles at base . . . . . GEN. XVII. SAUSSUREA.

Pappus longer than fruit—Anthers simple at base . . . . . GEN. XVIII. SERRATULA.

Pappus shorter than fruit . . . . . GEN. XXII. CENTAUREA.

SUB-ORDER III.—FLOWERS TUBULAR, OUTER OFTEN RADIANT—STYLE NOT THICKENED BELOW ITS BRANCHES.

A. Heads with a ray of different colour to disk (except *Chrys. segetum*).

* Pappus o.

Receptacle conical, naked—Scapes 1-flowered . . . . . GEN. XLI. BELLIS.

Receptacle flat or convex, naked—Involucre bracts with scarious margins . . . . . GEN. XLII. CHRYSANTHEMUM.

Receptacle conical or convex—Involucre bracts without scarious margins . . . . . GEN. XLIII. MATRICARIA.

Receptacle scaly . . . . . GEN. XLIV. ANTHEMIS.

- * * Pappus pilose.
Ray florets in several rows . . . . . GEN. XXXIV. ERIGERON.
Ray florets in 1 row . . . . . GEN. XXXV. ASTER.
 B. Heads with a ray of same colour as disk, discoid; or with the ray revolute.
 * Pappus of 2-5 rigid bristles. GEN. XXIII. BIDENS.
 * * Pappus of broad ciliate scales. GEN. XLVI. GALINSOGA.
 * * * Pappus feathery or pilose.
 a. Leaves all radical, appearing after flowers.
Heads many, racemed . . . . . GEN. XXXII. PETASITES.
Heads solitary . . . . . GEN. XXXIII. TUSSILAGO.
 b. Stems leafy.
 † Heads discoid—Involucral bracts herbaceous.
Flowers reddish purple . . . . . GEN. XXIV. EUPATORIUM.
Flowers yellow—Ls. linear . . . . . GEN. XXV. CHRYSOCOMA.
Flowers yellow—Ls. pinnfid. . . . . . GEN. XXXVII. SENECIO (in part).
 † † Heads discoid—Involucral bracts dry, scarious.
Receptacle flat, naked—Heads diæcious . . . . . GEN. XXIX. ANTENNARIA.
Receptacle flat, naked—Heads all similar . . . . . GEN. XXX. GNAPHALIUM.
Receptacle conical, scaly at margin . . . . . GEN. XXXI. FILAGO.
 † † † Heads radiant.
Involucral bracts imbricated—Anthers simple . . . . . GEN. XXXVI. SOLIDAGO.
Involucral bracts in 1 row, with scales at base . . . . . GEN. XXXVII. SENECIO.
Involucral bracts in 2-3 rows, equal . . . . . GEN. XXXIX. DORONICUM.
Involucral bracts imbricated—Anthers with bristles at base GEN. XL. INULA.
 * * * * Pappus o.
Receptacle not scaly—Fruit angular, with a large disk . GEN. XXVII. TANACETUM.
Receptacle not scaly—Fruit obovate, with a small disk . GEN. XXVIII. ARTEMISIA.
Receptacle scaly—Heads rayed . . . . . GEN. XLV. ACHILLEA.
Receptacle scaly—Heads discoid . . . . . GEN. XXVI. DIOITIS.

ORD. XLVII.—CAMPANULACEÆ.—(p. 88.)

- Corolla campan.; segments broad* . . . . . GEN. I. CAMPANULA.
Corolla rotate; segments linear—Anthers distinct . . . GEN. II. PHYTEUMA.
Corolla rotate; segments linear—Anthers cohering at base . GEN. III. JASIONE.
Corolla irregular—Anthers cohering in a tube . . . . . GEN. IV. LOBELIA.

ORD. XLVIII.—ERICACEÆ.—(p. 90.)

Div. I.—Ovary inferior.

GEN. VIII. VACCINIUM.

Div. II.—Ovary superior.

* Shrubs.

a. Fruit dry.

- Corolla persistent—Calyx simple, 4 part.* . . . . . GEN. I. ERICA.
Corolla persistent—Calyx double . . . . . GEN. II. CALLUNA.
Corolla deciduous, ventricose—Cal. 4-5 part.—Sta. 10 . . . GEN. III. MENZIESIA.
Corolla deciduous, campanulate—Sta. 5 . . . . . GEN. IV. LOISELEURIA.
Corolla deciduous, ovate—Sta. 10 . . . . . GEN. V. ANDROMEDA.

3. Fruit fleshy.

- Cells many-seeded* . . . . . GEN. VI. ARBUTUS.
Cells 1-seeded . . . . . GEN. VII. ARCTOSTAPHYLOS.

* * Herbs.

- Calyx green—Style 5 lobed* . . . . . GEN. IX. PYROLA.
Calyx coloured—Style simple, peltate . . . . . GEN. X. MONOTROPA.

ORD. XLIX.—AQUIFOLIACEÆ.—(p. 92.)

- Cor. rotate—Cal. 4-5 cleft* . . . . . GEN. I. ILEX.

ORD. L.—OLEACEÆ.—(p. 92.)

- Shrub—Fruit a fleshy berry* . . . . . GEN. I. LIGUSTRUM.
Tree—Fruit winged, dry . . . . . GEN. II. FRAXINUS.

ORD. LI.—APOCYNACEÆ.—(p. 94.)

- Cor. salver-shaped; tube long—Cal. 5 partite* . . . . . GEN. I. VINCA.

ORD. LII.—GENTIANACEÆ.—(p. 94.)

A. Leaves opposite.

- Corolla salver-shaped—Style persistent* . . . . . GEN. I. GENTIANA.
Corolla funnel-shaped—Sta. 4—Style deciduous—Stigma 1
entire . . . . . GEN. II. CICENDIA.
Corolla funnel-shaped—Sta. 5—Style deciduous—Stigmas 2 . . . . . GEN. III. ERYTHRÆA.
Corolla rotate—Sta. 8 . . . . . GEN. IV. CHLORA.

B. Leaves alternate.

- Leaves simple, orbicular* . . . . . GEN. V. VILLARSIA.
Leaves trifoliate . . . . . GEN. VI. MENYANTHES.

ORD. LIII.—POLEMONIACEÆ.—(p. 96.)

- Cor. rotate; tube short—Cal. 5 cleft* . . . . . GEN. I. POLEMONIUM.

ORD. LIV.—CONVOLVULACEÆ.—(p. 96.)

- Leafy plants* . . . . . GEN. I. CONVULVULUS.
Leafless parasites . . . . . GEN. II. CUSCUTA.

ORD. LV.—SOLANACEÆ.—(p. 96.)

- Cor. funnel-shaped—Capsule 2-celled, dry, opening by a lid* . . . . . GEN. I. HYOSCYAMUS.
Cor. rotate; segments reflexed—Anthers opening by terminal
pores . . . . . GEN. II. SOLANUM.
Cor. campanulate—Anthers not opening by pores . . . . . GEN. III. ATROPA.

ORD. LVI.—SCROPHULARIACEÆ.—(p. 98.)

A. Stamens 2.

- . . . . . GEN. II. VERONICA.

B. Stamens 4.

* Calyx 4-cleft.

- Upper lip of cor. entire, not flattened* . . . . . GEN. III. BARTSIA.
Upper lip of cor. bifid, not flattened . . . . . GEN. IV. EUPHRASIA.
Upper lip of cor. flattened—Calyx inflated . . . . . GEN. V. RHINANTHUS.
Upper lip of cor. flattened—Calyx tubular . . . . . GEN. VI. MELAMPYRUM.

* * Calyx 5 cleft.

a. Calyx inflated.

GEN. VII. PEDICULARIS.

b. Calyx tubular.

Corolla globose . . . . . GEN. VIII. SCROPHULARIA.

Corolla bell-shaped, unequal . . . . . GEN. IX. DIGITALIS.

Corolla personate, gibbous at base, not spurred . . . . . GEN. X. ANTIRRHINUM.

Corolla personate, spurred at base . . . . . GEN. XI. LINARIA.

Corolla campanulate, 5 cleft, equal . . . . . GEN. XII. LIMOSELLA.

Cor. rotate, equal . . . . . GEN. XIII. SIPHTHORIA.

Cor. ringent, upper lip with reflexed sides . . . . . GEN. XIV. MIMULUS.

C. Stamens 5,

Corolla rotate . . . . . GEN. I. VERBASCUM.

ORD. LVII.—OROBANCHACEÆ.—(p. 102.)

Cor. 4-5 cleft—Sep. 2 . . . . . GEN. I. OROBANCHE.

Cor. 2-lipped—Cal. 4 cleft . . . . . GEN. II. LATHRÆA.

ORD. LVIII.—VERBENACEÆ.—(p. 104.)

Cor. irregular—Cal. 5 part. . . . . GEN. I. VERBENA.

ORD. LIX.—LABIATÆ.—(p. 104.)

A. Stamens 2,

Corolla and calyx 2-lipped . . . . . GEN. I. SALVIA.

Corolla equally 4 cleft—Cal. 5 cleft . . . . . GEN. II. LYCOPUS.

B. Sta. 4, almost equal,

Corolla 4-5 cleft, nearly equal . . . . . GEN. III. MENTHA.

Corolla 2-lipped—Flowers capitate or in whorls . . . . . GEN. IV. THYMUS.

Corolla 2-lipped—Flowers in terminal heads . . . . . GEN. V. ORIGANUM.

C. Stamens 4, didynamous.

* Upper lip of corolla minute or o.

Upper lip of cor. deeply cleft . . . . . GEN. IX. TEUCRIUM.

Upper lip of cor. entire, or slightly notched . . . . . GEN. X. AJUGA.

* * Upper lip of corolla longer than the stamens.

a. Calyx obviously 2-lipped.

† Stamens remote at base, conniving under the upper lip.

Upper lip of cor. flat; its tube straight . . . . . GEN. VI. CALAMINTHA.

Upper lip of cor. concave; its tube curved . . . . . GEN. VII. MELISSA.

† † Stamens parallel.

Calyx campanulate, open in fruit . . . . . GEN. VIII. MELITTIS.

Calyx ovate, close in fruit; upper lip trifid, lower bifid . . . . . GEN. XVIII. PRUNELLA.

Calyx ovate, closed in fruit; lips entire . . . . . GEN. XIX. SCUTELLARIA.

b. Calyx 5-toothed.

† Lower stamens the longest.

Calyx 10-ribbed, funnel-shaped—Lower lip of cor. trifid; middle segment largest . . . . . GEN. XI. BALLOTA.

Calyx tubular—Upper lip of corolla flat; lower with 3 blunt lobes . . . . . GEN. XII. LEONURUS.

Calyx 10-ribbed, campanulate—Lower lip of cor. entire, with 1 or 2 small teeth on each side . . . . . GEN. XIII. LAMIAM.

- Calyx campanulate*—Lower lip of corolla with 3 unequal lobes and 2 teeth on its upper side . . . . . GEN. XIV. GALEOPSIS.
Calyx bell-shaped, 10-ribbed—Lower lip of corolla with the lateral lobes reflexed. . . . . GEN. XV. STACHYS.
     † † Two upper stamens the longest. . . . . GEN. XVI. NEPETA.
     c. Calyx 10-toothed. . . . .
Two lower stamens the longest . . . . . GEN. XVII. MARRUBIUM.

ORD. LX.—BORAGINACEÆ.—(p. 110.)

A. Throat of corolla naked.

- Cor. irregular* . . . . . GEN. XI. ECHIUM.
Cor. regular—Cal. tubular, 5-fid—Sta. included . . . . . GEN. X. PULMONARIA.
Cor. regular—Cal. short, in 5 deep segments—Sta. included . . . . . GEN. II. LITHOSPERMUM.
Cor. regular—Sta. protruded . . . . . GEN. III. MERTENSIA.

B. Throat of the corolla closed with scales.

* Stamens included.

a. Nuts attached to a disk below style.

- Cor. salver-shaped, twisted in æstivation* . . . . . GEN. I. MYOSOTIS.
Cor. funnel-shaped; its tube straight . . . . . GEN. VII. ANCHUSA.
Cor. funnel-shaped; its tube curved . . . . . GEN. VI. LYCOPSIS.
Cor. campanulate; its tube short, straight . . . . . GEN. IV. SYMPHYTUM.

b. Nuts attached to the elevated, persistent base of style.

- Cal. 5 cleft, with intermediate teeth* . . . . . GEN. VIII. ASPERUGO.
Cal. equally 5 cleft . . . . . GEN. IX. CYNOGLOSSUM.

* * Stamens exserted.

- Corolla rotate* . . . . . GEN. V. BORAGO.

ORD. LXI.—LENTIBULARIACEÆ.—(p. 112.)

- Terrestrial—Leaves entire* . . . . . GEN. I. PINGUICULA.
Aquatic—Leaves multifid . . . . . GEN. II. UTRICULARIA.

ORD. LXII.—PRIMULACEÆ.—(p. 114.)

- Leaves all radical—Cor. salver-shaped, with spreading lobes* . . . . . GEN. I. PRIMULA.
Leaves all radical—Limb of corolla reflexed . . . . . GEN. II. CYCLAMEN.
Corolla rotate—Calyx 7 partite . . . . . GEN. III. TRIENTALIS.
Aquatic—Leaves pectinate—Cor. salver-shaped . . . . . GEN. IV. HOTTONIA.
Cor. rotate, yellow—Cal. 5 partite—Capsule opening at summit . . . . . GEN. V. LYSIMACHIA.
Cor. rotate; tube 0—Calyx 5 part.—Sta. hairy—Capsule opening transversely . . . . . GEN. VI. ANAGALLIS.
Cor. with a sub-globose tube—Calyx 4 partite—Capsule opening transversely . . . . . GEN. VII. CENTUNCULUS.
Cor. salver-shaped, tube short—Cal. 5 part.—Capsule half inferior opening with reflexed teeth . . . . . GEN. VIII. SAMOLUS.
Cor. 0—Calyx tubular, coloured . . . . . GEN. IX. GLAUX.

ORD. LXIII. PLUMBAGINACEÆ.—(p. 116.)

- Flowers capitate* . . . . . GEN. I. ARMERIA.
Flowers paniced . . . . . GEN. II. STATICE.

ORD. LXIV.—PLANTAGINACEÆ.—(p. 118.)

- Flowers in spikes (perfect)* . . . . . GEN. I. PLANTAGO.
Flowers solitary (monœcious) . . . . . GEN. II. LITTORELLA.

ORD. LXV.—AMARANTHACEÆ.—(p. 118.)

- Monœcious—Perianth 3-5 partite* . . . . . GEN. I. AMARANTHUS.

ORD. LXVI.—CHENOPODIACEÆ.—(p. 118.)

A. Flowers perfect.

* St. not jointed.

† Leaves flat.

Pericarp free . . . . . GEN. I. CHENOPODIUM.

Pericarp adhering to perianth . . . . . GEN. IV. BETA.

† † Leaves semi-cylindrical.

Leaves spinous . . . . . GEN. V. SALSOLA.

Leaves not spinous . . . . . GEN. VI. SUCEDA.

* * Stem jointed.

. . . . . GEN. VII. SALICORNIA.

B. Flowers imperfect, monœcious.

Pericarp free . . . . . GEN. III. ATRIPLEX.

Pericarp adhering to perianth . . . . . GEN. II. OBIONE.

ORD. LXVII.—POLYGONACEÆ.—(p. 122.)

Perianth 5 part.; segments equal . . . . . GEN. I. POLYGONUM.

Perianth 6 part.; 3 inner segments large . . . . . GEN. II. RUMEX.

Perianth 4 part.; 2 inner segments large . . . . . GEN. III. OXYRIA.

ORD. LXVIII.—ELEAGNACEÆ.—(p. 124.)

Diœcious—Perianth of 2 leaves adhering at apex . . . . . GEN. I. HIPPOPHAË.

ORD. LXIX.—THYMELACEÆ.—(p. 124.)

Perianth 4 part., deciduous—Sta. 8 . . . . . GEN. I. DAPHNE.

ORD. LXX.—SANTALACEÆ.—(p. 124.)

Cor. funnel-shaped—Sta. 5 . . . . . GEN. I. THESIUM.

ORD. LXXI.—ARISTOLOCHIACEÆ.—(p. 124.)

A low herb—Sta. 12 . . . . . GEN. I. ASARUM.

Climbing—Sta. 6 . . . . . GEN. II. ARISTOLOCHIA.

ORD. LXXII.—EMPETRACEÆ.—(p. 126.)

Pet. 3 part.—Cal. 3 part.—Sta. 3 . . . . . GEN. I. EMPETRUM.

ORD. LXXIII.—EUPHORBIACEÆ.—(p. 126.)

Stamens many, with 1 pistil enclosed in a calyx-like bract . . . . . GEN. I. EUPHORBIA.

Evergreen shrub—Pet. and sep. present—Sta. 4 . . . . . GEN. II. BUXUS.

Perianth 3 partite—Fertile and barren fls. separate . . . . . GEN. III. MERCURIALIS.

ORD. LXXIV.—URTICACEÆ.—(p. 128.)

- Sta.* 4—Flowers (monœcious or diœcious) in racemes . . . GEN. I. URTICA.
Sta. 4—Flowers (polygamous) in axillary clusters, sur-
 rounded by involucre. . . . . GEN. II. PARIETARIA.
Sta. 5—Climbing—Flowers (diœcious) in a catkin . . . GEN. III. HUMULUS.

ORD. LXXV.—ULMACEÆ.—(p. 128.)

- Perianth* 4-5 part.—*Sta.* 5 . . . . . GEN. I. ULMUS.

ORD. LXXVI.—CUPULIFERÆ.—(p. 128.)

- A. Barren flowers in a globose catkin.
 . . . . . GEN. II. FAGUS.
 B. Barren flowers in a long catkin.
 * Stig. 3.

- . . . . . GEN. I. QUERCUS.
 * * Stig. 2.

- Scales of barren catkin roundish* . . . . . GEN. II. CORYLUS.
Scales of barren catkin 3 cleft . . . . . GEN. III. CARPINUS.

ORD. LXXVII.—BETULACEÆ.—(p. 130.)

- Sta.* 4—Achenes not winged . . . . . GEN. I. ALNUS.
Sta. 8-12—Achenes winged . . . . . GEN. II. BETULA.

ORD. LXXVIII.—SALICACEÆ.—(p. 130.)

- Sta.* 4-30—Scales of catkin jagged . . . . . GEN. I. POPULUS.
Sta. 1-5—Scales of catkin entire . . . . . GEN. II. SALIX.

ORD. LXXIX.—MYRICACEÆ.—(p. 134.)

- Sta.* 4-8—Scales of catkin concave—Fr. drupaceous . . . GEN. I. MYRICA.

ORD. LXXX.—CONIFERÆ.—(p. 134.)

- Fruit* a dry, many-seeded cone . . . . . GEN. I. PINUS.
Fruit a fleshy, few-seeded cone . . . . . GEN. II. JUNIPERUS.
Fruit a fleshy, cup-shaped, 1-seeded berry . . . . . GEN. III. TAXUS.

ORD. LXXXI.—ORCHIDACEÆ.—(p. 134.)

A. Leafless.

- Flowers* greyish brown—*Spur* 0 . . . . . GEN. III. NEOTTIA.
Flowers pink or purplish—*Spur* small . . . . . GEN. XIII. NEOTINEA.
Flowers pale yellow—*Spur* inflated . . . . . GEN. VII. EPIPOGIUM.
Flowers yellowish green; lip white, with purple blotches—
Spur 0 . . . . . GEN. VIII. CORALLORHIZA.

B. Plant leafy.

* Anthers 2.

- Lip* slipper-shaped. . . . . GEN. XVIII. CYPREPIDIUM.

* * Anther 1.

a. Lip spurred.

- Pollen glands* in a pouch . . . . . GEN. IX. ORCHIS.

- Pollen glands free, contiguous—Flowers rose* . . . . . GEN. X. GYMNADENIA.
Pollen glands free, distant—Flowers white . . . . . GEN. XI. HABENARIA.
     b. Lip not spurred.
     † Lip entire at apex.
Lip inferior, saccate at base, free from column—2 lateral
     sepals herbaceous, spreading . . . . . GEN. I. GOODYERA.
Lip inferior, with fleshy projections at base—2 lateral
     sepals, coloured, erect . . . . . GEN. II. SPIRANTHES.
Lip superior . . . . . GEN. XVI. MALAXIS.
Lip inferior—Pet. and sep. spreading . . . . . GEN. XVII. LIPARIS.
     †† Lip lobed or notched at apex.
     § Lip divided transversely.
Flowers drooping; ovary straight . . . . . GEN. V. EPIPACTIS.
Flowers erect; ovary twisted . . . . . GEN. VI. CEPHALANTHERA.
     §§ Lip not divided transversely.
Pet. and sep. spreading; lip linear, 2 lobed at apex—Pollen
     glands glandular. . . . . GEN. IV. LISTERA.
Pet. and sep. forming a hood; lip 3 lobed—Pollen glands in
     a common pouch . . . . . GEN. XII. ACERAS.
Pet. and sep. spreading; lip petaloid—Pollen glands naked. GEN. XIV. HERMINIUM.
Pet. and sep. spreading; lip variously lobed, mostly tumid—
     Pollen glands in 2 distinct pouches . . . . . GEN. XV. OPHRYS.

ORD. LXXXII.—IRIDACEÆ.—(p. 138.)

A. Segments of perianth unequal.

- Alternate segments of perianth reflexed* . . . . . GEN. I. IRIS.
Perianth almost labiate . . . . . GEN. II. GLADIOLUS.

B. Segments of perianth equal.

- Perianth tube short—Styles 3, filiform, entire* . . . . . GEN. III. SISYRINCHIUM.
Perianth tube short—Styles 3, bifid; lobes slender . . . . . GEN. IV. TRICHONEMA.
Perianth tube long . . . . . GEN. V. CROCUS.

ORD. LXXXIII.—AMARYLLIDACEÆ.—(p. 140.)

- Perianth with a bell-shaped crown within* . . . . . GEN. I. NARCISSUS.
Perianth without a crown; segments unequal . . . . . GEN. II. GALANTHIS.
Perianth without a crown; segments equal . . . . . GEN. III. LEUCOJUM.

ORD. LXXXIV.—LILIACEÆ.—(p. 140.)

A. Fruit dry.

* Stems leafy.

- Flowers drooping, paniced, flesh-coloured, with purple spots* GEN. I. LILIUM.
Flowers erect, solitary, yellow . . . . . GEN. II. TULIPA.
Flowers drooping, solitary, large, dull red, chequered . . . GEN. III. FRITILLARIA.
Flowers erect, solitary, white, veined with purple . . . . . GEN. X. LLOYDIA.

** Flowers on a leafless scape.

a. Segments of perianth free.

† Flowers capitate or umbellate.

- Fls. enclosed in a 1-2 leaved spathe* . . . . . GEN. IV. ALLIUM.
Fls. (yellow) enclosed in 1-2 linear leaves . . . . . GEN. V. GAGEA.

† † Flowers racemed or panicled.

Fls. white . . . . . GEN. VI. ORNITHOGALLUM.

Fls. blue or pink . . . . . GEN. VII. SCILLA.

Fls. jointed on their pedicels; white inside, purple without . GEN. XI. SIMETHIS.

b. Segments of perianth cohering at base.

Peri. campanulate, reflexed at tips . . . . . GEN. VIII. HYACINTHUS.

Peri. globose, contracted at mouth . . . . . GEN. IX. MUSCARI.

B. Fruit a succulent berry. . . . .

* Stem branched. . . . .

St. herbaceous—Flowers axillary . . . . . GEN. XII. ASPARAGUS.

*St. shrubby, with spinous ls. and fls. (apparently) on
centre of leaf* . . . . . GEN. XIII. RUSCUS.

* * Stem simple. . . . .

St. with 2 leaves—Peri. 4 partite . . . . . GEN. XIV. MAIANTHEMUM.

St. leafless—Peri. bell-shaped . . . . . GEN. XV. CONVALLARIA.

St. many-leaved—Peri. cylindrical . . . . . GEN. XVI. POLYGONATUM.

ORD. LXXXV.—TRILLIACEÆ.—(p. 144.)

Peri. in 2 whorls; outer herbaceous, inner coloured . . . GEN. I. PARIS.

ORD. LXXXVI.—DIOSCOREACEÆ.—(p. 144.)

Diœious—Perianth bell-shaped . . . . . GEN. I. TAMUS.

ORD. LXXXVII.—MELANTHIACEÆ.—(p. 144.)

Peri. tubular, very long—Scape leafless . . . . . GEN. I. COLCHICUM.

Peri. 6 partite—Scape leafy . . . . . GEN. II. TOFIELDIA.

ORD. LXXXVIII.—HYDROCHARIDACEÆ.—(p. 146.)

Leaves reniform, floating . . . . . GEN. I. HYDROCHARIS.

Leaves linear, submerged . . . . . GEN. II. ELODEA.

Leaves ensiform, submerged . . . . . GEN. III. STRATIOTES.

ORD. LXXXIX.—ALISMACEÆ.—(p. 146.)

* Leaves flat—Flowers large.

Sta. 6—Carpels many, free . . . . . GEN. I. ALISMA.

*Sta. 6—Carpels connate at base, spreading in a stellate
manner* . . . . . GEN. II. ACTINOCARPUS.

Sta. many—Carpels free—Ls. arrow-shaped . . . . . GEN. III. SAGGITTARIA.

* * Leaves triquetrous.

Fls. rose red . . . . . GEN. IV. BUTOMUS.

* * * Leaves linear.

Segs. of perianth erect . . . . . GEN. V. TRIGLOCHIN.

Segs. of perianth reflexed . . . . . GEN. VI. SCHENCHZERIA.

ORD. XC.—NAIADACEÆ.—(p. 148.)

* Flowers perfect.

Peri. segments 4 . . . . . GEN. I. POTAMOGETON.

Peri. 0 . . . . . GEN. II. RUPPIA.

* * Flowers imperfect.

- Fls. axill., sol.—Stigma 1, peltate.* . . . . . GEN. III. ZANNICHELLIA.
Fls. (apparently) in a slit in the leaf. . . . . . GEN. IV. ZOSTERA.
Fls. axillary, solitary—Stigmas 2-4, persistent. . . . . GEN. V. NAIAS.

ORD. XCI.—LEMNACEÆ.—(p. 150.)

- Frond with hair-like roots.* . . . . . GEN. I. LEMNA.
Frond rootless. . . . . . GEN. II. WOLFFIA.

ORD. XCII.—ARACEÆ.—(p. 150.)

- Leaves hastate.* . . . . . GEN. I. ARUM.
Leaves ensiform. . . . . . GEN. II. ACORUS.

ORD. XCIII.—TYPHACEÆ.—(p. 152.)

- Fls. in globular heads.* . . . . . GEN. I. SPARGANIUM.
Fls. in dense, cylindrical spikes. . . . . . GEN. II. TYPHA.

ORD. XCIV.—RESTIACEÆ.—(p. 152.)

- Peri. 4-6 part., inner segments united to summit.* . . . . GEN. I. ERIOCAULON.

ORD. XCV.—JUNCACEÆ.—(p. 152.)

- Peri. coloured (yellow).* . . . . . GEN. I. NARTHECIUM.
Peri. glumaceous—Capsule 3-celled, many-seeded. . . . . GEN. II. JUNCUS.
Peri. glumaceous—Capsule 1-celled, 3-seeded. . . . . GEN. III. LUZULA.

ORD. XCVI.—CYPERACEÆ.—(p. 154.)

A. Flowers perfect.

* Glumes 2 ranked.

- Spikelets many-flowered—Glumes all fertile.* . . . . . GEN. I. CYPERUS.
Spikelets few-flowered—Lower glume barren. . . . . . GEN. III. SCHÆNUS.

* * Glumes imbricated.

a. Bristles 0.

- Spikelets in many, crowded, compound cymes—Lower glumes smaller, empty.* . . . . . GEN. II. CLADIUM.
Spikelets 1-3, terminal, or collected into globular heads—Lower glume equal to upper. . . . . . GEN. VII. ISOLEPIS.

b. Bristles included in glumes.

- Spikelets in an ovate head, enclosed in a leafy bract.* . . . . GEN. IV. RHYNOSPORA.
Spikelets alternate, forming a bracteate spike. . . . . . GEN. V. BLYSMUS.
Spikelet 1, terminal. . . . . . GEN. VIII. ELEOCHARIS.
Spikelets fascicled, lateral. . . . . . GEN. VI. SCIRPUS.

c. Bristles much longer than glumes.

- Bristles at length silky.* . . . . . GEN. IX. ERIOPHORUM.

B. Flowers imperfect.

- Fruit not enclosed in the perigone.* . . . . . GEN. X. KOBRESIA.
Fruit enclosed in the perigone. . . . . . GEN. XI. CAREX.

ORD. XCVII.—GRAMINEÆ.—(p. 164.)

SECTION I.—SPIKELETS STALKED, FREE OF THE RACHIS; MOSTLY IN AN OPEN PANICLE, SOMETIMES IN A CYLINDRICAL SPIKE.

A. Spikelets with 2 or more perfect florets.

* Florets awned.

† Lower palea bifid.

- Glumes equalling lower floret—Awn geniculate* . . . . . GEN. XXII. AIRA.
Glumes equalling lower floret—Awn twisted . . . . . GEN. XXIV. AVENA.
Glumes shorter than lower floret—Awn straight . . . . . GEN. XXXIX. BROMUS.

†† Lower palea entire.

- Spikelets enveloped in silky hairs—Fruit free of palea* . . . . . GEN. XIX. ARUNDO.
Spikelets in an interrupted spike—Fruit free of palea . . . . . GEN. XXVIII. KOELERIA.
Spikelets crowded, subsecund—Fruit free of palea . . . . . GEN. XXXVII. DACTYLIS.
Spikelets mostly in a 1-sided panicle—Fruit adhering to palea . . . . . GEN. XXXVIII. FESTUCA.

** Florets not awned.

a. Spikelets with an apparent involucre.

- Spikelets with a pectinate bract at base* . . . . . GEN. XXXVI. CYNOSURUS
Spikelets with a ciliate bract at base . . . . . GEN. XXI. SESLERIA.

b. Spikelets without an apparent involucre.

† Fruit adhering to palea.

- Glumes boat-shaped, saccate, obtuse* . . . . . GEN. XXXV. BRIZA.

†† Fruit not adhering to palea.

- Glumes unequal, acute, as long as the lower palea—Lower palea emarginate, with an intermediate tooth* . . . . . GEN. XXVII. TRIODIA.
Glumes equal, veined, nearly as long as the ovate spikelet—Spikelets of 2 perfect florets and a club-like rudiment . . . . . GEN. XXIX. MELICA.
Glumes unequal, not veined, shorter than the lanc. spikelet—Palea hardening on the fruit . . . . . GEN. XXX. MOLINIA.
Glumes and outer palea eroso-truncate . . . . . GEN. XXXI. CATABROSA.
Glumes obtuse, unequal—Lower palea with 5-7 prominent, parallel veins; convex, obtuse. . . . . . GEN. XXXII. GLYCERIA
Glumes acute, unequal—Outer palea with 5 obscure, parallel veins; cylindrical below . . . . . GEN. XXXIII. SCHLEROCHLOA
Glumes acute, unequal—Outer palea with 3-5 parallel veins, compressed, keeled . . . . . GEN. XXXIV. POA.

B. Spikelets with 1 perfect floret and 1 or more rudimentary or neuter ones.

* Spikelets in a dense spike.

- Spike cylindrical—Palea with hairs at base.* . . . . . GEN. XVIII. AMMOPHILA.
Spike ovate—Outer palea bifid, long-awned. . . . . . GEN. XXIII. LAGURUS.
Spike ovate—Glu. equal, boat-shaped, keeled . . . . . GEN. VII. PHALARIS (Sp. 2).
Spike cylindrical, interrupted below—Palea with an involucre of bristles . . . . . GEN. VI. SETARIA.

** Spikelets in an interrupted, spike-like panicle.

- Sta. 2—Glu. unequal.* . . . . . GEN. VIII. ANTHOXANTHUM.
Sta. 3—Glu. equal, boat-shaped, keeled. . . . . . GEN. VII. PHALARIS (Sp. 1).

*** Spikelets in an open panicle.

- Upper floret with sta. only, awned; lower perfect, unawned.* GEN. XXVI. HOLCUS.

- Upper floret perfect, with short, straight awn; lower barren, with long awn* . . . . . GEN. XXV. ARRHENATHERUM.
- Upper floret perfect; 2 lower with sta. only, all unawned* . GEN. IX. HIEROCLOE.
- C. Spikelets with 1 perfect floret and no rudimentary or barren ones.
- * Spikelets in a dense, cylindrical spike.
- Paleæ awned* . . . . . GEN. XI. ALOPECURUS.
- Paleæ unawned* . . . . . GEN. X. PHLEUM.
- ** Spikelets in a spike-like panicle.
- Glumes acute, united and swollen at base* . . . . . GEN. XIII. GASTRIDIMUM.
- Glumes emarginate, with a long slender awn* . . . . . GEN. XIV. POLYPOGON.
- Paleæ surrounded by silky hairs* . . . . . GEN. XX. CALAMAGROSTIS (Sp. 1).
- *** Spikelets in a lax panicle.
- Glumes 0* . . . . . GEN. I. LEERSIA.
- Glumes equal—Paleæ unawned.* . . . . . GEN. XV. MILIUM.
- Glumes unequal; lower the smaller—Outer paleæ awned* . GEN. XVI. APERA.
- Glumes unequal; upper the smaller; outer paleæ unawned* . GEN. XVII. AGROSTIS.
- Glumes equal—Paleæ surrounded by silky hairs* . . . . GEN. XX. CALAMAGROSTIS.

SECTION II.—SPIKELETS SESSILE, ON A TOOTHED OR JOINTED RACHIS IN A TRUE SPIKE.

A. Spike compound.

- Spikes upright* . . . . . GEN. II. SPARTINA.
- Spikes spreading—Spikelets laterally compressed—Glumes equal* . . . . . GEN. III. CYNODON.
- Spikes spreading—Spikelets dorsally compressed—Glumes unequal.* . . . . . GEN. IV. DIGITARIA.
- Spikes spreading—Spikelets dorsally compressed, with bristles at base—Glumes unequal* . . . . . GEN. V. PANICUM.

B. Spike simple.

- Spikelets sol., unilat., 1-flowered—Paleæ scariose, shaggy, truncate.* . . . . . GEN. XII. KNAPPIA.
- Spikelets sol., sub-sess., broadside to rachis—Glume unequal, smaller than the outer paleæ* . . . . . GEN. XL. BRACHYPODIUM.
- Spikelets sol., broadside to rachis, many-flowered—Glumes equal.* . . . . . GEN. XLI. TRITICUM.
- Spikelets sol., edgewise to rachis, many-flowered—Glume 1, or inner smaller* . . . . . GEN. XLII. LOLIUM.
- Spikelets 2 or 3 together, 2-7 flowered* . . . . . GEN. XLIII. ELYMUS.
- Spikelets in threes, 1-flowered—Paleæ with long awns* . . GEN. XLIV. HORDEUM.
- Spikelets sol., in a 2-rowed, unilateral spike—Glumes 0* . . GEN. XLV. NARDUS.
- Spikelets sol., bilateral, 1-flowered—Glumes cartilaginous* . GEN. XLVI. LEPTURUS.

ORD. XCVIII.—FILICES.

SECTION I.—SPORE CLUSTERS ON BACK OF FROND.

A. Clusters with an indusium.

* Clusters dorsal, roundish.

- Indusia laciniate, attached beneath the sori* . . . . . GEN. III. WOODSIA.
- Indusia hooded, attached by their base* . . . . . GEN. VI. CYSTOPTERIS.
- Indusia circular, attached by their centre* . . . . . GEN. VII. POLYSTICHUM.

- Indusia reniform*, attached by their sides . . . . . GEN. VIII. LASTRÆA.
Indusia "crescentic, attached across the vein by their inner
edge" (Newm.) . . . . . GEN. IX. PSEUDATHYRIUM.
* * Clusters dorsal, linear.
Clusters single—Margins of *indusium laciniate* . . . . . GEN. X. ATHYRIUM.
Clusters single—Margins of *indusium entire* . . . . . GEN. XI. ASPLENIUM.
Clusters double, parallel . . . . . GEN. XII. SCOLOPENDRIUM.
* * * Clusters marginal—*Indusium* apparently consisting of the
reflexed margin of the frond.
Fronds all similar—Clusters short, oblong . . . . . GEN. XV. ADIANTUM.
Fronds all similar—Clusters continuous . . . . . GEN. XIV. PTERIS.
Fertile fronds dissimilar—Clusters oblong . . . . . GEN. V. ALLOSORUS.
Fertile fronds dissimilar—Clusters continuous . . . . . GEN. XIII. BLECHUM.
B. Clusters without an *indusium*.
Clusters linear, covered with chaffy scales . . . . . GEN. I. CETERACH.
Clusters round . . . . . GEN. IV. POLYPODIUM.
Clusters linear . . . . . GEN. II. GYMNOGRAMMA.

SECTION II.—SPORE CLUSTERS AT MARGIN OF FROND, TERMINATING A SINUS.

- Indusium cup-shaped* . . . . . GEN. XVI. TRICHOMANES.
Indusium 2-valved. . . . . GEN. XVII. HYMENOPHYLLUM.

SECTION III.—SPORE CLUSTERS IN SPIKES.

- Ls. bipinn.*, fertile at top. . . . . GEN. XVIII. OSMUNDA.
Ls. entire; spikes paniced . . . . . GEN. XIX. BOTRYCHIUM.
Ls. entire; spikes simple. . . . . GEN. XX. OPHIOGLOSSUM.

ORD. XCIX.¹—LYCOPODIACEÆ.

- Stem long*—Capsules all similar . . . . . GEN. I. LYCOPodium.
Stem long—Capsules of two kinds . . . . . GEN. II. SELAGINELLA.
Stem o—*Ls. filiform* . . . . . GEN. III. ISOETES.

ORD. C.—MARSILEACEÆ.

- Spore clusters sessile, globose* . . . . . GEN. I. PILULARIA.

ORD. CI.—EQUISETACEÆ.

- St. leafless, branched with toothed sheaths* . . . . . GEN. I. EQUISETUM.

TABULAR CLASSIFICATION.

ORD. I.—RANUNCULACEÆ.

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
Gen. I. Clematis, L.	<i>Traveller's Joy.</i>					
i. <i>C. vitalba, L.</i>	<i>common T.</i>	Hedges .	W.	Cl.	Shb.	7-8
Gen. II. Thalictrum, L.	<i>Meadow-Rue.</i>					
i. <i>T. alpinum, L.</i>	<i>Alpine M.</i>	Alp. bogs	Y.	4-8 in.	P.	7-8
ii. <i>T. minus, L.</i>	<i>lesser M.</i>	Cha. past.	g Y.	½-4 ft.	P.	7-8
v. 1. <i>maritimum</i>	. . . .	San. sh.	. . .	. . .	. . .	. . .
v. 2. <i>montanum</i>	. . . .	Stony pas.	. . .	. . .	. . .	. . .
v. 3. <i>flexuosum, Bernh.</i>	. . . .	N. of E.	. . .	. . .	. . .	. . .
v. 4. <i>Kochii, Fries</i>	. . . .	Lake dist.	. . .	. . .	. . .	. . .
v. 5. <i>Saxatile, Schl.</i>	. . . .	. . . .	. . .	. . .	. . .	. . .
iii. <i>T. flavum, L.</i>	<i>common M.</i>	Riv. banks	p Y.	2-4 ft.	P.	7-8
v. 1. <i>særocarpum, Sy.</i>	. . . .	. . . .	. . .	. . .	. . .	. . .
v. 2. <i>riparium, Sy.</i>	. . . .	. . . .	. . .	. . .	. . .	. . .
v. 3. <i>Morisonii, Sy.</i>	. . . .	. . . .	. . .	. . .	. . .	. . .
Gen. III. Anemone, L.	<i>Anemone.</i>					
i. <i>A. pulsatilla, L.</i>	<i>Pasque fl.</i>	Cha. hills	Pu.	4-8 in.	P.	5-6
ii. <i>A. nemorosa, L.</i>	<i>Wood A.</i>	Woods .	W.	4-8 in.	P.	4-5
iii. <i>A. apennina, L.</i>	<i>blue Mount A.</i>	Woods! .	B.	4-8 in.	P.	4-5
Gen. IV. Adonis, L.	<i>Pheasant's-eye.</i>					
i. <i>A. autumnalis, L.</i>	<i>Corn P.</i>	Corn fie.	Cr.	6-10 in.	A.	5-7
Gen. V. Myosorus, L.	<i>Mousetail.</i>					
i. <i>M. minimus, L.</i>	<i>common M.</i>	Corn fie.	y G.	2-6 in.	A.	4-6
Gen. VI. Ranunculus, L.	<i>Buttercup, Crow-foot, Spearwort, &c.</i>					
SEC. I.						
i. <i>R. aquatilis, L.</i>	<i>Water C.</i>	Pds., dit. & riv.	W.	flo.	P.	5-8
* A.						
v. 1. <i>heterophyllus, Fries</i>	. . . .	Ponds .	. . .	. . .	. . .	. . .
v. 2. <i>pseudo-fluitans, Sy.</i>	. . . .	Streams .	. . .	. . .	. . .	. . .
* *						
v. 3. <i>peltatus, Fries</i>	. . . .	Ponds .	. . .	. . .	. . .	. . .
v. 4. <i>floribundus, Bab.</i>	. . . .	Ponds .	. . .	. . .	. . .	. . .
v. 5. <i>confusus, Godr.</i>	. . . .	Salt. mar.	. . .	. . .	. . .	. . .
v. 6. <i>Baudotii, Godr.</i>	. . . .	Ditches .	. . .	. . .	. . .	. . .
B.						
v. 7. <i>fluitans, Lamk.</i>	. . . .	Rivers. .	. . .	. . .	. . .	. . .
v. 8. <i>Bachii, Wirtg.</i>	. . . .	Rivers. .	. . .	. . .	. . .	. . .
C.						
v. 9. <i>Drouettii, Schu.</i>	. . . .	Ponds .	. . .	. . .	. . .	. . .
v. 10. <i>trichophyllus, Cha.</i>	. . . .	Ponds .	. . .	. . .	. . .	. . .
v. 11. <i>circinatus, Sib.</i>	. . . .	Ponds .	. . .	. . .	. . .	. . .
D.						
v. 12. <i>tripartitus, DC.</i>	. . . .	Ditches!	. . .	. . .	. . .	. . .
ii. <i>R. Lenormandi, Sch.</i>	<i>Mud C.</i>	Ditches .	W.	Cr.	P.	6-8

ORD. I.—RANUNCULACEÆ.

CHARACTERISTICS OF SPECIES AND VARIETIES.

Gen. I.

- i. St. climbing—Ls. pinn. ; lfts. (3-5) cord.

Gen. II.

- i. St. simp., naked—Ls. bitern. glauc. beneath—Rac. terni.
 ii. St. leafy—Ls. tri-pinn. ; lfts. trifid—Pan. lax ; fls. drooping.
 v. 1. *Leafless at base—Aur. spread—Pan. leafless, broad ; branches spreading.*
 v. 2. *Leafless at base—Aur. spread—Pan. leafless, narrow ; branches sub-erect.*
 v. 3. *Leafy at base—Lfts. very large—Aur. reflex—Pan. leafy.*
 v. 4. *Leafy at base—Lfts. small—Aur. horizontal—Pan. leafless.*
 v. 5. *Leafy at base—Aur. reflex—St. ang.—Fls. erect—Ach. not gibbous.*
 iii. St. furrowed—Ls. bipinn. ; lfts. trifid—Pan. compact ; fls. erect.
 v. 1. *Pan. contracted—Ach. glob.*
 v. 2. *Pan. lax—Ach. oval.*
 v. 3. *Pan. interrupted—Ach. obl.*

Gen. III.

- i. Fls. erect—Sep. silky—Ls. bipinn. ; seg. lin.—Ach. feathery.
 ii. Fls. drooping—Ls. tern.—Sep. 6 obl.
 iii. Fls. erect, blue—Ls. tritern.—Sep. 10-14 narrow.

Gen. IV.

- i. Ls. decomposed ; seg. lin.—Pet. 5-to concave, emarginate.

Gen. V.

- i. Ls. lin., fleshy—Sp ke slender.

Gen. VI.

SEC. I.—AQUATIC PLANTS WITH WHITE FLOWERS.

- i. Flo. ls. trifid ; submerged ls. capil.-multifid. very variable.

A. Fls. large—flo. ls. (usually) present.

* Seg. of ls. collapsing when taken from water.

- v. 1. *Recept. con. —Pedunc. as long as ls.*
 v. 2. *Recept. spher. —Pedunc. much longer than ls. —Flo. ls. sometimes absent.*

* * Seg. of ls. spreading when taken from water.

- v. 3. *Recept. spher. —Stig. club-shaped—Carp. blunt—Pedunc. tapering.*
 v. 4. *Recept. spher. —Stig. tongue-shaped—Carp. blunt—Pedunc. equal.*
 v. 5. *Recept. ov.-con. —Sta. longer than pist.—Carp. narrowed upwards.*
 v. 6. *Recept. long-con. —Sta. shorter than pist.—Carp. infl. upwards.*

B. Fls. large—Flo. ls. absent.

- v. 7. *Seg. of ls. long, rigid, and parallel—Pedunc. shorter than ls.*
 v. 8. *Similar, but ls. smaller, narrower, and sessile—Pedunc. longer than ls.*

C. Fls. small—Flo. ls. absent.

- v. 9. *Seg. of ls. collapsing when taken from water.*
 v. 10. *Seg. of ls. spreading but not in one plane.*
 v. 11. *Seg. of ls. spreading in one flat rigid plane.*

D. Fls. small—submerged ls. absent.

- v. 12. *Ls. nearly round, long stalked, divided nearly to base.*

- ii. Ls. reni. ; lobes narrowest at base—Pet. twice as long as sep.

ORD. I.—RANUNCULACEÆ—cont.

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Duration.	Time of flowering.
iii. <i>R. hederaceus</i> , <i>L.</i> . . . SEC. II. <i>A.</i>	<i>Ivy leaved C.</i> . . .	Ditches .	W.	Cr.	P.	6-8
iv. <i>R. ficaria</i> , <i>L.</i>	<i>Pilewort</i>	Fie. . .	Y.	3-6 in.	P.	4-5
v. 1. <i>divergens</i> , <i>Schu.</i> . . .	. . . . .	. . . . .	. . .	. . .	. . .	. . .
v. 2. <i>incumbens</i> , <i>Schu.</i> . . .	. . . . .	. . . . .	. . .	. . .	. . .	. . .
v. <i>R. flammula</i> , <i>L.</i>	<i>lesser S.</i>	Wet pl. .	Y.	4-12 in.	P.	6-9
v. 2. <i>reptans</i> , <i>L.</i>	. . . . .	San. sh. of lakes!	. . .	. . .	. . .	. . .
vi. <i>R. lingua</i> , <i>L.</i>	<i>great S.</i>	Bogs . .	Y.	3-5 ft.	P.	6-9
vii. <i>R. ophioglossifolius</i> , <i>Vill.</i> <i>B.</i> *	<i>Adder's Tongue S.</i>	Bogs, Jersey!	Y.	6-10 in.	A.	6-8
viii. <i>R. auricomus</i> , <i>L.</i> . . .	<i>Goldilocks</i>	Thick. .	Y.	6-10 in.	P.	4-8
ix. <i>R. acris</i> , <i>L.</i>	<i>upright B.</i>	Fie. . .	Y.	1-3 ft.	P.	6-7
a. <i>Steiveni</i> , <i>Reich.</i>	. . . . .	. . . . .	. . .	. . .	. . .	. . .
b. <i>vulgatus</i> , <i>Jord.</i>	. . . . .	. . . . .	. . .	. . .	. . .	. . .
c. <i>rectus</i> , <i>Bor.</i>	. . . . .	. . . . .	. . .	. . .	. . .	. . .
x. <i>R. repens</i> , <i>L.</i>	<i>creeping C.</i>	Past. . .	Y.	6-12 in.	P.	5-8
xi. <i>R. bulbosus</i> , <i>L.</i>	<i>bulbous C.</i>	Past. . .	Y.	1-2 ft.	P.	5-8
xii. <i>R. sceleratus</i> , <i>L.</i> . . . * *	<i>Celery-leaved C.</i>	Dit. . .	Y.	1-2 ft.	A.	5-9
xiii. <i>R. hirsutus</i> , <i>Curt.</i> . . .	<i>pale hairy C.</i> . . .	Past. . .	Y.	6-12 in.	A.	6-10
xiv. <i>R. parviflorus</i> , <i>L.</i> . . .	<i>small flowered C.</i>	Corn fie.	Y.	6-18 in.	A.	5-6
xv. <i>R. arvensis</i> , <i>L.</i> . . .	<i>Corn C.</i>	Corn fie.	Y.	6-18 in.	A.	5-7
Gen. VII. <i>Caltha</i> , <i>L.</i>	<i>Marsh Marygold.</i>					
i. <i>C. palustris</i> , <i>L.</i>	<i>common M.</i>	Mar. . .	Y.	8-18 in.	P.	3-6
a. <i>vulgaris</i>	. . . . .	. . . . .	. . .	. . .	. . .	. . .
v. 2. <i>Guerangerii</i>	. . . . .	. . . . .	. . .	. . .	. . .	. . .
v. 3. <i>minor</i>	. . . . .	. . . . .	. . .	. . .	. . .	. . .
b. <i>radicans</i> , <i>Forst.</i>	. . . . .	Scot.! . .	. . .	. . .	. . .	. . .
Gen. VIII. <i>Trollius</i> , <i>L.</i>	<i>Globe flower.</i>					
i. <i>T. Europæus</i> , <i>L.</i> . . .	<i>Mountain G.</i> . . .	Moun.pas.	Y.	6-18 in.	P.	5-6
Gen. IX. <i>Eranthis</i> , <i>Salisb.</i>	<i>Winter Aconite.</i>					
[i. <i>E. hyemalis</i> , <i>Salisb.</i> . .	<i>common W.</i>	Parks . .	Y.	6-12 in.	P.	2-4
Gen. X. <i>Helleborus</i> , <i>L.</i>	<i>Hellebore.</i>					
i. <i>H. viridis</i> , <i>L.</i>	<i>green H.</i>	Woods .	G.	12-18 in.	P.	4-5
ii. <i>H. foetidus</i> , <i>L.</i>	<i>stinking H.</i>	Woods .	G.	1-2 ft.	P.	4-5
Gen. XI. <i>Aquilegia</i> , <i>L.</i>	<i>Columbine.</i>					
i. <i>A. vulgaris</i> , <i>L.</i>	<i>common C.</i>	Thick. .	Pu.	1-2 ft.	P.	5-6
Gen. XII. <i>Delphinium</i> , <i>L.</i>	<i>Larkspur.</i>					
i. <i>I. Ajacis</i> , <i>Reich.</i> . . .	<i>field L.</i>	Corn fie.	B.	10-18 in.	A.	6-7

ORD. I.—RANUNCULACEÆ—*cont.*

CHARACTERISTICS OF SPECIES AND VARIETIES.

- iii. Ls. reni. ; lobes broadest at base—Pet. not longer than sep.

SEC. II.—TERRESTRIAL PLANTS WITH YELLOW FLOWERS.

A. Leaves entire.

- iv. St. one flowered—Ls. cord., angled—Rt. tuberous.
 v. 1. Lobes of lower ls. not overlapping at base.
 v. 2. Lobes of lower ls. overlapping at base.
 v. St. decumb. at base, not rooting at joints—Ls. ov.-lanc., stalked.
 v. 2. St. filiform, rooting at joints—Ls. ellip.-lanc.
 vi. St. erect—Ls. lanc. sess.—Fls. large.
 vii. St. erect—Ls. ov.-lanc.; lower ones cord., long stalked—Pet. scarcely longer than sep.

B. Leaves divided.* *Achenes smooth.*

- viii. Rad. ls. reni., long stalked ; st. ls. digit. ; seg. lin.—Cal. spread., coloured.
 ix. Cal. spreading—Pedunc. cylind., smooth—Rt. fibr.
 a. St. slightly hairy at base—Seg. of rad. ls. not overlapping (*common*).
 b. St. densely hairy at base—Seg. of rad. ls. overlapping (*common*).
 c. St. slightly hairy with adpressed hairs (*rare, Lochnagar*).
 x. Cal. spread.—Pedunc. furrowed—Rt. with cr. scions.
 xi. Cal. reflex.—Pedunc. furrowed—Rt. swollen, almost bulbous.
 xii. Cal. reflex. hairy—Ls. tri-pinn. ; upper lobes lin.—Fls. small—St. succulent.
 * * *Achenes rough.*
 xiii. Ach. tubercled at margins—St. erect hairy, many fld.—Cal. reflexed.
 xiv. Ach. with hooked tubercles—Ls. orb., three-lobed—St. decumb.—Pedunc. lat.
 xv. Ach. covered with hooked spines—St. erect—Cal. spread—Fls. pale yellow.

Gen. VII.

- i. Ls. orb., rad.—St. erect—Sep. imbricate.
 a. Joints not rooting—Rad. ls. orb.
 v. 2. *Smaller—Sep. remote when expanded.*
 v. 2. *Procumb.—Fls. sol., smaller—Sep. remote.*
 b. Rooting at joints—Rad. ls. deltoid, toothed.

Gen. VIII.

- i. Ls. five part. ; seg. cun. lobed—Fls. glob ; sep. coloured.

Gen. IX.

- i. Rad. ls. orb., 3-5 part., long stalked—Fls. sol.—Invol. leafy.]

Gen. X.

- i. Rad. ls. digit. decid.—St. few-flowered—Cal. spread.
 ii. Rad. ls. pedate, evergreen—St. many-flowered—Cal. erect.

Gen. XI.

- i. Ls. 2-3 tern. ; seg. three-lobed, stalked—Pet. spurred—Fls. pan.

Gen. XII.

- i. St. erect, branched—Ls. multifid ; seg. lin.—Rac. of fls. long—Carp. and foll. downy.
 (*D. Consolida, also found in Jersey, has shorter racemes, and the carp. and foll. glabrous.*)

ORD. I.—RANUNCULACEÆ—*cont.*

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
Gen. XIII. Aconitum, L.	<i>Monkshood.</i>					
i. <i>A. napellus, L.</i>	<i>common M.</i>	Riv. bnks.!	d B.	1-2 ft.	P.	7-8
Gen. XIV. Actæa, L.	<i>Bane-berry.</i>					
i. <i>A. spicata, L.</i>	<i>B. or Herb Chris- topher.</i>	Moi. cops.!	W.	1-2 ft.	P.	5-6

ORD. II.—BERBERACEÆ.

Gen. I. Berberis, L.	<i>Barberry.</i>					
i. <i>B. vulgaris, L.</i>	<i>common B.</i>	Hedges .	p Y.	4-6 ft.	Shb.	5-6

ORD. III.—NYMPHÆACEÆ.

Gen. I. Nymphæa, L.	<i>Water-lily.</i>					
i. <i>N. alba, L.</i>	<i>white W.</i>	Lakes, &c.	W.	flo.	P.	7-8
Gen. II. Nuphar, L.	<i>Water-lily.</i>					
i. <i>N. lutea, L.</i>	<i>yellow W.</i>	Lakes, &c.	Y.	flo.	P.	7-8
v. 2. <i>minor</i>						
ii. <i>N. pumila, Sm.</i>	<i>least W.</i>	Alp. lakes!	Y.	flo.	P.	7-8

ORD. IV.—PAPAVERACEÆ.

Gen. I. Papaver, L.	<i>Poppy.</i>					
* i. <i>P. hybridum, L.</i>	<i>round prickly- headed P.</i>	Chal. fie.	Cr.	10-18 in.	A.	5-7
ii. <i>P. argemone, L.</i>	<i>long prickly- headed P.</i>	Corn fie.	R.	6-10 in.	A.	5-7
* * iii. <i>P. dubium, L.</i>	<i>long smooth- headed P.</i>	Corn fie.	Sc.	1-2 ft.	A.	5-7
α. <i>Lamottei, Bor.</i>						
β. <i>Lecoqii, Lam.</i>						
iv. <i>P. Rhæas, L.</i>	<i>common red P.</i> . .	Corn fie.	Sc.	1-2 ft.	A.	5-7
v. 2. <i>strigosum, Boenn.</i> . .						
v. <i>P. somniferum, L.</i> . . .	<i>white P.</i>	Corn fie.	W.	1-2 ft.	A.	6-8
Gen. II. Meconopsis, Vig.	<i>Welch Poppy.</i>					
i. <i>M. cambrica, Vig.</i> . . .	<i>common W.</i>	Rocky pl.!	Y.	1-2 ft.	P.	6-7

ORD. I.—RANUNCULACEÆ—*cont.*

CHARACTERISTICS OF SPECIES AND VARIETIES.

Gen. XIII.

- i. St. erect, simp.—Ls. 3-5 part., seg. deeply cut—Rac. simp.

Gen. XIV.

- i. Ls. bitern, stalked ; lfts. trifid, deeply cut—Rac. simp., elong.

ORD. II.—BERBERACEÆ.

Gen. I.

- i. Branches with three part. spines—Ls. ob.-ov. serr.—Fls. in a pend. rac.

ORD. III.—NYMPHÆACEÆ.

Gen. I.

- i. Ls. cord. ent.—Pet. obl., white.

Gen. II.

- i. Ls. heart-shaped ; lobes at base contiguous—Stig. ent. with 15-20 rays.
 v. 2. *Fls. smaller (1½ in. diam.)—Stig. slightly waved with 10-14 rays (Chartness lock)*
 ii. Ls. heart-shaped ; lobes at base distant—Stig. lobed with 8-10 rays.

ORD. IV.—PAPAVERACEÆ.

Gen. I.

* *Fruit hispid.*

- i. Fr. glob., furrowed.
 ii. Fr. club-shaped.
 * * *Fruit smooth.*
 iii. Fr. obl.—Hairs of pedunc. adpr.

- a. Leaf lobes short, suddenly acute—Caps, broadest at top.
 b. Leaf lobes long, gradually acute—Caps, broadest at one-third from top.
 iv. Fr. glob.—Hairs of pedunc. spread.
 v. 2. *Hairs of pedunc. adpr.*
 v. Fr. glob., glab.—Ls. lobed, amplex., glauc.

Gen. II.

- i Fr. obl., glab.—Ls. long-stalked, pinn. ; lfts. pinn.

ORD. IV.—PAPAVERACEÆ—*cont*

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
Gen. III. <i>Chelidonium</i>, L.	<i>Celandine</i> .					
i. <i>C. majus</i> , L.	<i>Celandine</i>	Sh. pl. .	Y.	1-2 ft.	P.	5-8
Gen. IV. <i>Glaucium</i>, L.	<i>Horned Poppy</i> .					
i. <i>G. luteum</i> , Scop.	<i>yellow H.</i>	San. sh.	Y.	1-2 ft.	B.	6-9
ii. <i>G. violaceum</i> , Fuss.	<i>violet H.</i>	Ch. fie.! .	Vi.	1-2 ft.	P.	5-6

ORD. V.—FUMARIACEÆ.

Gen. I. <i>Fumaria</i>, L.	<i>Fumitory</i> .					
i. <i>F. capreolata</i> , L.	<i>rampant F.</i>	Corn fie.	var.	Cl.	A.	6-9
v. 2. <i>pallidiflora</i> , Ford.	. . . . .	S.W. of E.	C.	. .	. .	. .
v. 3. <i>Boræi</i> , Ford.	. . . . .	. . . . .	p Pu.	. .	. .	. .
v. 4. <i>confusa</i> , Ford.	. . . . .	. . . . .	Pk.	. .	. .	. .
v. 5. <i>muralis</i> , Sond.	. . . . .	. . . . .	Ro.	. .	. .	. .
iv. <i>F. officinalis</i> , L.	<i>common F.</i>	Corn fie.	p Pu.	6-12 in.	A.	5-9
iii. <i>F. micrantha</i> , Lag.	<i>small fld. F.</i> . . .	Corn fie.	p Pu.	6-10 in.	A.	5-9
iv. <i>F. tenuisecta</i> , Sy.	<i>least fld. F.</i>	Corn fie.	p Pu.	6-10 in.	A.	6-9
a. <i>parviflora</i> , Lam.	. . . . .	. . . . .	. .	. .	. .	. .
b. <i>Vaillantii</i> , Lois.	. . . . .	. . . . .	. .	. .	. .	. .
Gen. II. <i>Corydalis</i>, DC.	<i>Corydalis</i> .					
i. <i>C. claviculata</i> , DC.	<i>white climbing C.</i>	Thick. .	y W.	Cl.	A.	6-9
ii. <i>C. lutea</i> , DC.	<i>yellow C</i>	Old walls	Y.	6-10 in.	P.	5-8
iii. <i>C. solida</i> , Hook.	<i>solid rooted C.</i> . .	Thick.! .	Pu.	6-10 in.	P.	4-5

ORD. VI.—CRUCIFERÆ.

Gen. I. <i>Cakile</i>, Gært.	<i>Sea-Rocket</i> .					
i. <i>C. maritima</i> , L.	<i>purple S. R.</i>	San. sh. .	Li.	1-2 ft.	A.	6-7
Gen. II. <i>Crambe</i>, L.	<i>Kale</i> .					
i. <i>C. maritima</i> , L.	<i>Sea K.</i>	Sea sh. .	W.	1-2 ft.	P.	6
Gen. III. <i>Coronopus</i>, Sm.	<i>Wart-Cress</i> .					
i. <i>C. didyma</i> , Sm.	<i>lesser W.</i>	Was. pl. .	W.	Cr.	A.	7-9
ii. <i>C. Ruellii</i> , Gært.	<i>common W.</i>	Was. pl..	W.	Cr.	A.	6-9
Gen. IV. <i>Isatis</i>, L.	<i>Woad</i> .					
i. <i>I. Tinctoria</i> , L.	<i>Dyers' W.</i>	Fie.! . .	Y.	1-3 ft.	P.	7-8

ORD. IV.—PAPAVERACEÆ—*cont.*

CHARACTERISTICS OF SPECIES AND VARIETIES.

Gen. III.

- i. Ls. pinnfid.—Fls. in long stalked umbels.

Gen. IV.

- i. St. branched, glauc.—Ls. amplex., sinuate—Fls. yellow.
 ii. Ls. bipinnfid. ; seg. lin., scabr.—Pet. violet.

ORD. V.—FUMARIACEÆ.

Gen. I.

- i. St. climbing—Petioles twisted—Ls. bipinn.—Sep. as broad as tube of cor.
 y. 2. *Sep. one-half to two-thirds as long as tube of cor.—Fr. longer than broad—Pet. cream-coloured tipped with purple.*
 y. 3. *Sep. one-third as long as and broader than tube of cor.—Fr. broader than long—Pet. pale purple.*
 y. 4. *Sep. one-third as long as and rather narrower than tube of cor.—Fr. longer than broad—Pet. pink tipped with purple.*
 y. 5. *Sep. and fr. as Var. 4, but fls. smaller, rose tipped with purple.*
 ii. St. erect, branched—Ls. bi-tripinn.—Sep. half as broad as tube of cor.
 iii. St. diff.—Ls. bipinn. ; seg. small—Sep. round, dent. half as long and twice as broad as tube of cor.
 iv. St. diff.—Ls. bipinn. ; seg. lin., glauc.—Sep. minute, lin.
 a. Seg. of ls. channelled—Rec. dense—Pedi. as long as bracts.
 b. Seg. of ls. flat—Rac. lax—Pedi. twice as long as bracts.

Gen. II.

- i. St. cli.—Peti. tendrilled—Ls. pinn. ; lfts. 3-5 nate. ; seg. ov.
 ii. St. erect, branched, ang.—Ls. bipinn. ; lfts. lobed—Rac. lat.—Rt. fibr.]
 iii. St. erect, simp.—Ls. 3-4 ternate, glauc.—Rac. term.—Rt. tuberous.]

ORD. VI.—CRUCIFERÆ.

Gen. I.

- i. Branches zigzag—Ls. fleshy, pinnfid.

Gen. II.

- i. Ls. glauc., broad, waved—Root fleshy.

Gen. III.

- i. Rac. dense—Pouch notched, granulated—Sty. short.
 ii. Rac. lax—Pouch ent., crested with sharp points—Sty. prominent.

Gen. IV.

- i. Rad. ls. obl., cren. : St. ls. sagitt.

ORD. VI.—CRUCIFERÆ—*cont.*

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
Gen. V. <i>Thlaspi</i>, <i>L.</i>	<i>Penny-Cress.</i>					
i. <i>T. arvense</i> , <i>L.</i>	<i>Field P.</i>	Fie. . .	W.	1-2 ft.	A.	5-7
ii. <i>T. perfoliatum</i> , <i>L.</i> . . .	<i>perfoliate P.</i> . . .	Lim. dis.	W.	4-6 in.	A.	4-5
iii. <i>T. alpestre</i> , <i>L.</i>	<i>Alpine P.</i>	Moun. pas.	W.	6-10 in.	B.	5-7
a. <i>sylvestre</i> , <i>Ford.</i>	. . . . .	. . . .	. . .	. . .	. . .	. . .
b. <i>occitanum</i> , <i>Ford.</i>	. . . . .	. . . .	. . .	. . .	. . .	. . .
c. <i>virens</i> , <i>Ford.</i>	. . . . .	. . . .	. . .	. . .	. . .	. . .
Gen. VI. <i>Capsella</i>, <i>Vent.</i>	<i>Shepherd's Purse.</i>					
i. <i>C. Bursa-pastoris</i> , <i>DC.</i>	<i>common S.</i>	Every- where.	W.	6-18 in.	A.	2-10
Gen. VII. <i>Hutchinsia</i>, <i>Br.</i>	<i>Hutchinsia.</i>					
i. <i>H. petræa</i> , <i>Br.</i>	<i>Rock H.</i>	Lim. ro..	W.	2-4 in.	A.	3-5
Gen. VIII. <i>Teesdalia</i>, <i>Br.</i>	<i>Teesdalia.</i>					
i. <i>T. nudicaulis</i> , <i>Br.</i> . . .	<i>naked-stalked T.</i>	Grav. pl.	W.	2-4 in.	A.	4-6
Gen. IX. <i>Iberis</i>, <i>L.</i>	<i>Candy-tuft.</i>					
i. <i>I. amara</i> , <i>L.</i>	<i>bitter C.</i>	Chal. fi..	W.	6-9 in.	A.	7-8
Gen. X. <i>Lepidium</i>, <i>L.</i>	<i>Pepper-wort.</i>					
i. <i>L. latifolium</i> , <i>L.</i>	<i>broad-leaved P.</i> . .	Salt ma.!	W.	2-4 ft.	P.	7-8
[ii. <i>L. Draba</i> , <i>L.</i>	<i>Whitlow P.</i>	Banks .	W.	1-2 ft.	P.	5-6
iii. <i>L. Smithii</i> , <i>Hook.</i> . . .	<i>smooth Fie. P.</i> . .	Fie. . .	W.	6-12 in.	P.	4-8
iv. <i>L. campestre</i> , <i>Br.</i> . . .	<i>common P.</i>	Fie. . .	W.	10-12 in.	A.	5-8
v. <i>L. ruderale</i> , <i>L.</i>	<i>narrow-leaved P.</i>	Salt ma.!	W.	6-12 in.	A.	5-6
Gen. XI. <i>Cochlearia</i>, <i>L.</i>	<i>Scurvy Grass.</i>					
i. <i>C. officinalis</i> , <i>L.</i>	<i>common S.</i>	Sea sh. .	W.	4-10 in.	A.	5-8
v. 2. <i>alpina</i> , <i>Wats.</i>	. . . . .	Scot. mo.!	. . .	. . .	. . .	. . .
v. 3. <i>danica</i> , <i>L.</i>	. . . . .	Sea sh. .	. . .	. . .	. . .	. . .
ii. <i>C. anglica</i> , <i>L.</i>	<i>English S.</i>	Riv. bks.	W.	10-18 in.	A.	5-8
[iii. <i>C. armoracia</i> , <i>L.</i>	<i>Horse-Radish</i> . .	Was. pl..	W.	2-3 ft.	P.	5-6
Gen. XII. <i>Subularia</i>, <i>L.</i>	<i>Awlwort.</i>					
i. <i>S. aquatica</i> , <i>L.</i>	<i>Water A.</i>	Marg. of lakes.	W.	3-4 in.	P.	7
Gen. XIII. <i>Draba</i>, <i>L.</i>	<i>Whitlow Grass.</i>					
i. <i>D. aizoides</i> , <i>L.</i>	<i>yellow Alpine W.</i>	Wales! .	Y.	2-5 in.	P.	3-5
ii. <i>D. rupestris</i> , <i>Br.</i>	<i>Rock W.</i>	Alp. roc.!	W.	1-2 in.	P.	5-6
iii. <i>D. incana</i> , <i>L.</i>	<i>twisted-podded W.</i>	Alp. roc.!	W.	6-12 in.	P.	5-6
iv. <i>D. muralis</i> , <i>L.</i>	<i>speedwell-leaved W.</i>	Lim. roc.!	W.	6-12 in.	A.	4-5
v. <i>D. verna</i> , <i>L.</i>	<i>common W.</i>	Dry bks.	W.	1-6 in.	A.	4-5
v. 2. <i>brachycarpa</i> , <i>Ford.</i>	. . . . .	Yorks. .	. . .	. . .	. . .	. . .
v. 3. <i>inflata</i> , <i>Hook.</i>	. . . . .	Ben Lawers.	. . .	. . .	. . .	. . .

ORD. VI.—CRUCIFERÆ—*cont.*

CHARACTERISTICS OF SPECIES AND VARIETIES.

Gen. V.

- i. Pod large, orb. with a broad wing—St. ls. obl., sagitt. toothed, glab.
- ii. Pod ob.-cord. broadly winged—St. ls. sagitt. cord—Sty. very short.
- iii. Pod ob.-ov. retuse—St. simple—Sty. as long as notch.
  - a. Rac. long—Sty. not longer than notch.
  - b. Rac. obl.—Sty. longer than notch—Pod triang.
  - c. Rac. oval—Sty. much longer than notch—Pod ob.-oval.

Gen. VI.

- i. Pod triang., ob.-cord.—Rad. ls. pinnfid.—Very variable.

Gen. VII.

- i. L. pinnfid., seg. ent.—Sep. as long as pet.

Gen. VIII.

- i. Ls. all rad., lyr. pinnfid.—Pet. unequal.

Gen. IX.

- i. St. branched—Ls. obl., lanc., acu.—Fls. spiked.

Gen. X.

- i. Ls. ov. lanc., ent., toothed—Pod oval, downy—Sty. minute.
- ii. Ls. obl., ent.—St. ls. amplex—Pod cord.—Sty. as long as pod.]
- iii. St. branched from base—Sty. twice as long as notch—Anth. violet.
- iv. St. erect, simp., branched at top—Sty. not longer than notch—Anth. yellow.
- v. Rad. ls. pinnfid.—St. ls. lin. ent.—Fls. apetalous—Sty. minute.

Gen. XI.

- i. Rad. ls. cord—St. ls. sess. ; semi-amplex angled—Pod globose.
- v. 2. *Smaller—Ls. fleshy—Pod ob.-ov.*
- v. 3. *Rad. ls. cord—St. ls. stalked, delt.—Pod ovate. veiny.*
- ii. Rad. ls. ovate, obl.—St. ls. lanc., toothed, semi-amplex—Pod ellip. strongly veined.
- iii. Rad. ls. obl. ; cren. long stalked—St. ls. lanc., sess.—Root cylind.]

Gen. XII.

- i. Ls. few, rad., subulate—Roots of long white fibres.

Gen. XIII.

- i. St. naked—Ls. lanc., rigid, fringed—Pet. twice as long as sep.
- ii. St. naked or one-leaved—Ls. lanc., hairy, soft—Pet. slightly notched.
- iii. St. leafy—Ls. lanc., toothed, pubes—Pods twisted.
- iv. St. branched flex.—Ls. cord. toothed, hairy—Pods flat, obt.
- v. St. naked—Ls. lanc., toothed, hairy—Pet. deeply notched—Pod. compr. ob.-ov., twice as long as broad.
- v. 2. *Pods compressed, orb.—1½ times as long as broad.*
- v. 3. *Pods inflated.*

ORD. VI.—CRUCIFERÆ—*cont.*

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
Gen. XIV. Camelina, <i>Crantz.</i>	<i>Gold-of-Pleasure.</i>					
i. <i>C. sativa, Cr.</i>	<i>common G.</i>	Flax fie. .	Y.	2-3 ft.	A.	6-7
Gen. XV. Koniga, Br.	<i>Koniga.</i>					
[i. <i>K. maritima, Br.</i> . . .	<i>Sea-side K.</i>	Sea co. .	W.	4-8 in.	P.	8-9
Gen. XVI. Alyssum, L.	<i>Alyssum.</i>					
[i. <i>A. calycinum, L.</i> . . .	<i>Alyssum</i>	Sea co. .	Y.	3-8 in.	A.	6-8
Gen. XVII. Dentaria, L.	<i>Coral-wort.</i>					
i. <i>D. bulbifera, L.</i>	<i>bulbiferous C.</i> . . .	Sha. pl.!	Li.	1-2 ft.	P.	4-5
Gen. XVIII. Cardamine, L.	<i>Bitter-cress.</i>					
i. <i>C. amara, L.</i>	<i>large fld. B.</i>	Wat. pl. .	W.	1-2 ft.	P.	4-5
ii. <i>C. pratensis, L.</i>	<i>Cuckoo flower</i> . . .	Mead. .	Li.	1-2 ft.	P.	4-6
iii. <i>C. hirsuta, L.</i>	<i>hairy B.</i>	Wat. pl. .	W.	6-12 in.	A.	4-8
v. 2. <i>sylvatica, Link.</i> . . .	. . . . .	. . . . .	. . .	. . .	. . .	. . .
iv. <i>C. impatiens, L.</i>	<i>narrow-leaved B.</i>	Roc. pl.!	W.	1-2 ft.	A.	6-7
Gen. XIX. Arabis, L.	<i>Rock Cress.</i>					
i. <i>A. petræa, Lam.</i>	<i>Alpine R.</i>	Alp. roc.!	W.	3-6 in.	P.	7-8
ii. <i>A. stricta, Huds.</i>	<i>Bristol R.</i>	Rocks! .	W.	5-10 in.	P.	4-5
iii. <i>A. ciliata, Br.</i>	<i>fringed R.</i>	Ir. sea roc.!	W.	4-6 in.	B.	7-8
v. 2. <i>hispida, Sy.</i>	. . . . .	S. Wales	W.	. . .	. . .	. . .
iv. <i>A. hirsuta, Br.</i>	<i>hairy R.</i>	Dry bks.	W.	1-2 ft.	B.	6-8
v. 2. <i>glabrata, Sy.</i>	. . . . .	Arran.!	. . .	. . .	. . .	. . .
v. A. <i>perfoliata, Lamk.</i> . . .	<i>smooth R. C., or</i> <i>Tower Mustard.</i>	Dry bks.!	y W.	2-3 ft.	B.	5-6
[vi. <i>A. Turrita, L.</i>	<i>Tower R.</i>	Walls! .	p Y.	1-2 ft.	B.	5
Gen. XX. Barbarea, Br.	<i>Winter Cress.</i>					
i. <i>B. vulgaris, Br.</i>	<i>yellow Rocket.</i> . . .	Dp. pl. .	Y.	1-2 ft.	B.	5-8
v. 2. <i>arcuata, Reich.</i>	. . . . .	I.!	. . .	. . .	. . .	. . .
v. 3. <i>stricta, And.</i>	. . . . .	N. of E.!	. . .	. . .	. . .	. . .
v. 4. <i>intermedia, Bor.</i>	. . . . .	. . . . .	. . .	. . .	. . .	. . .
ii. <i>B. præcox, Br.</i>	<i>early W.</i>	Was. pl.	Y.	1-2 ft.	B.	5-7
Gen. XXI. Nasturtium, Br.	<i>Water-Cress or</i> <i>Yellow Cress.</i>					
i. <i>N. officinale, Br.</i>	<i>common W.</i>	Brooks. .	W.	2-3 ft.	P.	6-7
v. 2. <i>sifolium, Rei.</i>	. . . . .	. . . . .	. . .	. . .	. . .	. . .
ii. <i>palustre, DC.</i>	<i>Marsh Y.</i>	Wat. pl. .	Y.	1-2 ft.	P.	6-8
iii. <i>N. sylvestre, Br.</i>	<i>creeping Y.</i>	Wet pl. .	Y.	9-12 in.	P.	6-9

CHARACTERISTICS OF SPECIES AND VARIETIES.

Gen. XIV.

- i. Rad. ls. stalked—St. ls. lanc., arrow shaped—Pods large, long stalked.

Gen. XV.

- i. St. procumb.—Ls. lin. lanc., acute—Pods glab.]

Gen. XVI.

- i. St. stellutely hoary—Ls. ob.-ov., lan.—Pods pubes.

Gen. XVII.

- i. St. simple—Lower ls. pinn. ; upper simple.

Gen. XVIII.

- i. Lfts. of st. ls. toothed, ang.—Sta. as long as pet.—Anth. violet.
 ii. Lfts. of st. ls. lanc., ent.—Stam. half as long as pet.—Anth. yellow.
 iii. Lfts. of lower ls. roundish obl. ; cut, petioled ; upper ones narrower—Pet. small, ob.-lanc.—Stam. four—Pods and pedi. erect.
 v. 2. *Perennial*—*Stam. six*—*Pedi. patent*—*Pods erect*.
 iv. Lfts. lanc., narrow, deeply cut—Petio. with sagitt. ciliated step—Pet. lin. or ov.

Gen. XIX.

- i. Rad. ls. lyr. pinnfid. stalked—St. ls. ent.—Pods spr.
 ii. Rad. ls., lyr., toothed, obt., hisped—St. ls. semi-amplex—Pods erect.
 iii. St. simp.—Rad. ls. ob.-ov., ciliate—St. ls. sess., ellip., pods erect.
 v. 2. *Leaves hisped*.
 iv. Ls. all hisped, toothed—St. ls. semi-amplex—Pods adpr.
 v. 2. *St. and ls. glab*.
 v. St. very erect—Rad. ls. toothed ; st. ls. glab., glauc. ent., amplex, auricled—Pods many, crowded, erect.
 vi. Rad. ls. ellip., stalked ; st. ls. narrow, amplex—Pods recurved ; flat with a thickened margin.]

Gen. XX.

- i. Rad. ls. lyr.—Upper ls. toothed ; term. lfts. roundish—Pod short, thicker than pedi.
 v. 2. *Pods in a lax. elong. rac.*—*Pet. twice as long as sep. ; pedi. spreading*.
 v. 3. *Pods in a dense rac.*—*Term. lft. large*—*Pet. half as long again as sep.*—*Pedi. adpr.*—*Seeds twice as long as broad*.
 v. 4. *Term. lft. smaller*—*Seg. of ls. more numerous*—*Pet. twice as long as sep.*—*Seeds as long as broad*.
 ii. Rad. ls. lyr.—Upper pinnfid. ; seg. narrow—Pod long, not thicker than pedi.

Gen. XXI.

- i. Aquatic—Fls. white—Ls. pinn. ; term. lfts. larger.
 v. 2. *St. tall, erect, thick*—*Term. lfts. not larger*.
 ii. Rt. fibr.—Ls. pinnfid. seg. unequally cut—Pet. not longer than sep.—Pods turgid, obl.
 iii. Rt. cr.—Ls. pinn. ; lfts. lanc., cut—Pet. twice as long as sep.—Pods lin.

ORD. VI.—CRUCIFERÆ—*cont.*

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
iv. <i>N. amphibium</i> , <i>Br.</i> .	<i>amphibious Y.</i> .	Wet. pl.	Y.	2-4 ft.	P.	6-9
Gen. XXII. <i>Sisymbrium</i>, <i>L.</i>	<i>Hedge Mustard.</i>					
i. <i>S. officinale</i> , <i>Scop.</i> . .	<i>common H.</i> . . .	Was. pl..	Y.	1-2 ft.	A.	6-7
ii. <i>S. Irio</i> , <i>L.</i>	<i>London Rocket.</i> .	Old walls!	Y.	1-2 ft.	A.	6-7
iii. <i>S. Sophia</i> , <i>L.</i>	<i>Flix Weed.</i> . . .	Was. pl..	Y.	1-2 ft.	A.	6-8
iv. <i>S. thalianum</i> , <i>Hook.</i> .	<i>Thale Cress.</i> . .	Dry pl. .	W.	6-10 in.	A.	5-9
v. <i>S. alliaria</i> , <i>Sc.</i> . . .	<i>Garlic Mustard.</i> .	Banks . .	W.	2-3 ft.	B.	5-6
vi. <i>S. polyceratum</i> , <i>L.</i> .	. . . .	Bury St. Edmunds.	Y.	1-2 ft.	A.	7-8
Gen. XXIII. <i>Erysimum</i>, <i>L.</i>	<i>Treacle Mustard.</i>					
i. <i>E. cheiranthoides</i> , <i>L.</i>	<i>Worm-seed T.</i> .	Was. pl..	W.	1-2 ft.	B.	6-8
ii. <i>E. orientale</i> , <i>Br.</i> . . .	<i>Hare's-ear T.</i> . .	Cliffs! .	W.	1-2 ft.	A.	5-7
Gen. XXIV. <i>Cheiranthus</i>, <i>L.</i>	<i>Wallflower.</i>					
i. <i>C. cheiri</i> , <i>L.</i>	<i>Common W.</i> . .	Walls . .	Y.	6-12 in.	P.	4-5
Gen. XXV. <i>Matthiola</i>, <i>Br.</i>	<i>Stock.</i>					
i. <i>M. incana</i> , <i>Br.</i> . . .	<i>hoary shrubby S.</i>	Cliffs! .	Pu.	1-2 ft.	P.	5-6
ii. <i>M. sinuata</i> , <i>Br.</i> . . .	<i>great sea S.</i> . . .	Sand. sh.!	Pu.	1-2 ft.	B.	6-8
Gen. XXVI. <i>Hesperis</i>, <i>L.</i>	<i>Dame's Violet.</i>					
i. <i>H. matronalis</i> , <i>L.</i> . .	<i>Common D.</i> . . .	Hilly pas.!	Li.	2-3 ft.	B.	5-6
Gen. XXVII. <i>Brassica</i>, <i>L.</i>	<i>Cabbage, &c.</i>					
i. <i>B. oleracea</i> , <i>L.</i> . . .	<i>Wild Cabbage.</i> .	Cliffs . .	Y.	1-2 ft.	B.	6-8
ii. <i>B. campestris</i> , <i>L.</i> . .	<i>Wild Navev.</i> . .	Riv. banks	Y.	1-2 ft.	A. or B.	6-7
iii. <i>B. napus</i> , <i>L.</i>	<i>Rape.</i>	Was. gr.	Y.	1-2 ft.	A.	5-6
iv. <i>B. rapa</i> , <i>L.</i>	<i>Turnip.</i>	Was. pl..	Y.	1-2 ft.	B.	4-7
v. <i>B. monensis</i> , <i>Br.</i> . . .	<i>I. of Man C.</i> . . .	Sand. sh.!	Y.	6-18 in.	B.	6-8
v. 2. <i>cheiranthus</i> , <i>Vill.</i> .	. . . . .	Jersey! .	. .	. .	. .	. .
Gen. XXVIII. <i>Sinapis</i>, <i>L.</i>	<i>Mustard.</i>					
i. <i>S. arvensis</i> , <i>L.</i>	<i>Charlock.</i>	Corn fie..	Y.	1-2 ft.	A.	5-8
ii. <i>S. alba</i> , <i>L.</i>	<i>white M.</i>	Corn fie..	Y.	1-2 ft.	A.	6-7
iii. <i>S. nigra</i> , <i>L.</i>	<i>common M.</i>	Corn fie..	Y.	1-2 ft.	A.	6-8
iv. <i>S. incana</i> , <i>L.</i>	<i>hoary M.</i>	Jersey! .	Y.	1-2 ft.	B.	7-8
Gen. XXIX. <i>Diplotaxis</i>, <i>DC.</i>	<i>Rocket.</i>					
i. <i>D. tenuifolia</i> , <i>DC.</i> . .	<i>Wall R.</i>	Old walls	Y.	1-3 ft.	P.	7-9
ii. <i>D. muralis</i> , <i>DC.</i> . . .	<i>Sand R.</i>	Was. pl..	Y.	6-9 in.	A.	8-9
v. 2. <i>Babingtonii</i> , <i>Bab.</i> .	. . . . .	. . . . .	. .	. .	. .	. .

CHARACTERISTICS OF SPECIES AND VARIETIES.

- iv. Ls. obl., lanc., serr. or pinnfid.—Pet. twice as long as sep.—Pod obl. shorter than the stalk.

Gen. XXII.

- i. St. hairy—Ls. runc., hairy—Pods subulate adpr.
 ii. St. glab.—Ls. runc., glab.—Pods terete, erect.
 iii. St. slightly hairy—Ls. bi.-pinnfid; seg. lin.
 iv. Ls. nearly all rad., obl., lan., toothed, pubes—Pods asc. with four obscure angles.
 v. Ls. large, cord stalked, much veined—Pods erect, long, on short stout pedi.
 vi. St. prost.—Ls. runc. pinnfid., glab.—Rods three together, sess., axill., subu.]

Gen. XXIII.

- i. Ls. lanc., ent. or slightly lobed, hairy.
 ii. Ls. ellip., cord., ent., amplex, glauc.; rad. ls. obt.]

Gen. XXIV.

- i. St. shrubby—Ls. lanc., ent.

Gen. XXV.

- i. St. shrubby, branched, erect—Ls. lanc., ent.—Pod cylin.
 ii. St. herb., spread.—Ls. lin., downy; lower sinuate—Pod compr., muricate.

Gen. XXVI.

- i. St. erect—Ls. oval, lanc., toothed—Pet. obt.

Gen. XXVII.

- i. Rt. fleshy, cylind.—Ls. glauc., lobed, waved.
 ii. Rt. tuber.—St. slender—Lower ls. lyr.—somewhat hisped; upper cord, amplex, glauc.—Pods erect.
 iii. Rt. fusiform—Ls. all glab., glauc., lower lyr.—Pods spread.
 iv. Rt. tub., fleshy—Lower ls. scabrous; upper ls. glab.
 v. St. prost., leafless—Ls. deeply pinnfid., glab.
 2. *St. erect, branched, leafy, hisped below.*

Gen. XXVIII.

- i. Ls. ovate, lyr.—Pods spread., knotted, longer than beak.
 ii. Ls. lyr., pinnfid.—Pods spread., knotted, shorter than beak.
 iii. Upper ls. lin.-lanc., ent. glab.—Pods adpr., four angled; beak sterile.
 iv. St. much branched—Upper ls. lin.-lanc. hisped—Pods adpr. cylind.; beak one-seeded.

Gen. XXIX.

- i. St. leafy, branched, glabr.—Ls. lanc., pinnfid. glauc.
 ii. St. simple, leafless, hisped—Ls. lyr., sinu., glab.
 2. *Biennial—St. ls. several.*

ORD. VII.—RESEDACEÆ.ORD. VIII.—CISTACEÆ.ORD. IX.—VIOLACEÆ.

Gen. I. <i>Viola</i> , <i>L.</i>		<i>Violet, Pansy,</i> <i>Heartsease.</i>					
	<i>*</i>						
i.	<i>V. Palustris, L. . . .</i>	<i>marsh V. . . . .</i>	Mar. . .	Li.	Cr.	P.	4-7
ii.	<i>V. odorata, L. . . . .</i>	<i>sweet V. . . . .</i>	Banks. .	var.	Cr.	P.	3-5
iii.	<i>V. hirta, L. . . . .</i>	<i>hairy V. . . . .</i>	Chal. pas.	♂ B.	Cr.	P.	4-6
v. 2.	<i>calcareæ, Bab. . . . .</i>	. . . . .	Gogmagog Hills.	. .	. .	. .	. .
v 3.	<i>sepincola, Ford. . . .</i>	. . . . .	. . . . .	. .	. .	. .	. .
	<i>* *</i>						
iv.	<i>V. canina, L. . . . .</i>	<i>Dog V. . . . .</i>	Woods .	var.	Cr.	P.	4-6
	<i>A.</i>						
a.	<i>V. sylvatica, Fries. . .</i>	. . . . .	. . . . .	. .	. .	. .	. .
v. 1.	<i>Riviniiana, Reich. . .</i>	. . . . .	. . . . .	. .	. .	. .	. .
v. 2.	<i>Reichenbachiana,</i> <i>Reich.</i>	. . . . .	. . . . .	. .	. .	. .	. .

ORD. VI.—CRUCIFERÆ—*cont.*

CHARACTERISTICS OF SPECIES AND VARIETIES.

Gen. XXX.

- i. Ls. simply lyr.—Sty. twice as long as last joint of pod.
- ii. Rad. ls. interruptedly pinn.—Sty. scarcely longer than last joint of pod.

ORD. VII.—RESEDACEÆ.

Gen. I.

- i. Ls. lanc., ent.—Sep. and pet. four.
- ii. Ls. lower pinn.: upper trifid—Sep. and pet. six unequal.
- iii. Ls. all pin., waved, glauc.; seg. lin.—Sep. and pet. five equal.]
- v. 2. *Sep. and pet. six.*

ORD. VIII.—CISTACEÆ.

Gen. I.

- i. St. shrubby, procumb.—Ls. stip. opp., ov., flat, green above, hoary beneath.
- ii. St. shrubby, procumb.—Ls. sti., opp., obl., revol. at margin, hoary on both sides.
- iii. St. shrubby, procumb.—Ls. not stip., opp., ov.-obl., stalked, flat, hairy above, hoary beneath.
- v. 2. *Ls. glab. above.*
- iv. St. ann. erect—Ls. obl., lanc., lower opp. upper alt.—Rac. without tracts.
- v. 2. *Rac. bracteate.*

ORD. IX.—VIOLACEÆ.

Gen. I.

- * *Stem short, creeping.*
- i. Ls. reni., cord., glab.—Stig. flat—Fr. st. erect.
- ii. Ls. cord., slightly hairy—Stig. hooked—Fr. st. prost.—Runners long—Spur. straight—Fls. violet, lilac or white.
- iii. Ls. cord. very hairy—Runners o. or short.—Fr. st. prost.—Spur. hooked.
- v. 2. *Pet. narrower—Pedunc. longer than ls.*
- v. 3. *much hairier—With longer runners.*
- * * *Stem evident.*
- iv. L. obl. cord. long stalked, smooth, varying—Fr. st. erect—Spur. blunt.
A. *Main. st. flowerless—Fl. stems axill.*
- a. L. orb., cord., acum.—Spur. long—Caps. glab.
- v. 1. *Spur. pale furrowed—Fls. large.*
- v. 2. *Spur. purple, not furrowed—Fls. smaller and later.*

ORD. IX.—VIOACEÆ—*cont.*

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
<i>b. arenaria</i> , DC.	. . . . .	Teesdale !	. .	. .	. .	. .
<i>B.</i>						
<i>c. canina</i> , Bab.	. . . . .	Hea. . .	. .	. .	. .	. .
<i>v. 2. flavicornis</i> , Sm.	. . . . .	Hea. . .	. .	. .	. .	. .
<i>d. lactea</i> , Sm.	. . . . .	Hea. ! . .	. .	. .	. .	. .
<i>e. stagnina</i> , Kit.	. . . . .	Bogs. ! .	. .	. .	. .	. .
<i>v. V. tricolor</i> , L.	<i>Pansy</i>	Pas. . .	P. & Y.	4-10 in.	A.	5-9
<i>v. 2. arvensis</i> , Murr.	. . . . .	Fie. . . .	<i>w</i> Y.	. .	. .	. .
<i>vi. V. Curtisii</i> , Forst.	. . . . .	San. sh. !	B. & Y.	3-10 in.	P.	5-9
<i>v. 1. Forsteri</i>	. . . . .	. . . .	. .	. .	. .	. .
<i>v. 2. Mackaili</i>	. . . . .	. . . .	. .	. .	. .	. .
<i>v. 3. Symei</i>	. . . . .	. . . .	. .	. .	. .	. .
<i>vii. V. lutea</i> , Huds.	<i>yellow P.</i>	Hill. pl. !	Y.	6-12 in.	P.	5-9
<i>v. 2. amœna</i>	. . . . .	. . . .	. .	. .	. .	. .

ORD. X.—DROSERACEÆ.

Gen. I. Drosera , L.	<i>Sundew.</i>					
<i>i. D. rotundifolia</i> , L.	<i>round-leaved S.</i>	Bogs . .	W.	3-6 in.	P.	7-8
<i>ii. D. intermedia</i> , L.	<i>long-leaved S.</i>	Bogs . .	W.	2-4 in.	P.	7-8
<i>iii. D. Anglica</i> , Huds.	<i>great English S.</i>	Bogs . .	W.	4-8 in.	P.	7-8

ORD. XI.—POLYGALACEÆ.

Gen. I. Polygala , L.	<i>Milkwort.</i>					
<i>i. P. vulgaris</i> , L.	<i>common M.</i>	Hea. . .	W.B. or Li.	2-10 in.	P.	6-8
<i>v. 2. grandiflora</i> , Sy.	. . . . .	Benbulbin !	. .	. .	. .	. .
<i>v. 3. oxyptera</i> , Reich.	. . . . .	San. sh. .	. .	. .	. .	. .
<i>v. 4. ciliata</i>	. . . . .	Gogmagog Hills.	. .	. .	. .	. .
<i>v. 5. depressa</i> , Wen.	. . . . .	Hea. . .	. .	. .	. .	. .
<i>ii. P. calcarea</i> , Sch.	. . . . .	Chalky downs.	W.B. or Li.	2-5 in.	P.	5-6
<i>iii. P. uliginosa</i> , Frie.	. . . . .	Teesdale and Kent. !	<i>d</i> Pk.	2-4 in.	P.	6-8

ORD. XII.—FRANKENIACEÆ.

Gen. I. Frankenia , L.	<i>Sea-Heath.</i>					
<i>i. F. lævis</i> , L.	<i>smooth S.</i>	Salt mar. !	Ro.	Cr.	P.	8-9

ORD. IX.—VIOLACEÆ—*cont.*

CHARACTERISTICS OF SPECIES AND VARIETIES

- b.* L. broadly cord., obt.—Sp. short—Caps. pubesc.
B. Main. st. long, branched, flowering.
- c.* L. narrow, cord. at base, roundedly acute—Spur. obt.—Caps. ovoid.
- v.* 2. *Smaller—Ls. more cord.—Spur. paler.*
- d.* L. lanc., base rounded—Pet. narrow, grey—Caps. sub-glob.
- e.* Rootstock creeping—Ls. obl. lanc.; base trunc.—Pet. oval, obl., pale—Caps. three-sided.
- v.* Rootstock o.—Ls. obl., cren.—Stip. leafy—Pet. longer than sep.—Caps. ovoid.
- v.* 2. *Pet. not longer than sep.—Caps. glob.*
- vi.* Rootstock woody, creeping; branches stout—St. ang. rough—Pet. scarcely longer than sep.
- v.* 1. }
v. 2. } *Vars. Forsteri and Mackain differ only in the colour of the fls.*
- v.* 3. *Fls. larger, yellow, one in. long.—Stip. larger—(Mullaghmore, I.).*
- vii.* Rootstock slender, underground—Pet. yellow, much longer than sep.
- v.* 2. *Fls. purple.*

ORD. X.—DROSERACEÆ.

Gen. I.

- i.* Ls. orb., spread.—Peti. hairy.
- ii.* Ls. obl., spath., erect—Peti. glab.—Seeds with a compact rough coat.
- iii.* Ls. ob.-ov., lanc., erect—Peti. glab.—Seeds with a loose chaffy coat.

ORD. XI.—POLYGALACEÆ.

Gen. I.

- i.* St. woody—Lower ls. ov., crowded; upper lanc., alt.—Cal. wings ob.-ov. broader than fr.—Pedi. glab.—Rac. term.
- v.* 2. *Upper ls. large—Cal. wings ellip. ac.—Fls. dark blue.*
- v.* 3. *Fls. smaller—Cal. wings ellip., cun. below, narrower than fr.—Pedi. glab.*
- v.* 4. *Cal. wings ob.-ov. broader than fr.—Pedi. pubesc.—Sep., bracts, and caps. ciliate.*
- v.* 5. *St. fili.—Lower ls. spath., scattered; upper ellip. nearly opp., distichous—Rac. term. and lat.*
- ii.* St. branched; each branch with a rosette of ellip. ls., from which the fl. shoots spring—St. ls. narrower—Cal. wings ob.-ov. broader and longer than fr.
- iii.* Ls. spath. in a rad. rosette—Upper ls. narrow—Cal. wings obl. narrower than fr.

ORD. XII.—FRANKENIACEÆ.

Gen. I.

- i.* Ls. lin., revolute at marg., glab., ciliate at base.

ORD. XIII.—ELATINACEÆ.

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
Gen. I. Elatine, L.	<i>Waterwort.</i>					
i. <i>E. hexandra, DC.</i> . . .	<i>hexandrous W.</i> . .	Marg. of lakes!	Ro.	Cr.	A.	8-9
ii. <i>E. Hydropiper, L.</i> . . .	<i>small octandrous W.</i>	Marg. of lakes!	Ro.	Cr.	A.	8-9

ORD. XIV.—CARYOPHYLLACEÆ.

Gen. I. Dianthus, L.	<i>Pink, Carnation.</i>					
*						
i. <i>D. prolifer, L.</i>	<i>proliferous P.</i> . . .	Grav. pl.!	Ro.	6-12 in.	A.	7
ii. <i>D. Armeria, L.</i>	<i>Deptford P.</i>	Dry banks	Ro.	1-1½ ft.	A.	7-8
***						
[iii. <i>D. plumarius, L.</i> . . .	<i>wild P.</i>	Old walls!	Pk.	1 ft.	P.	6-7
[iv. <i>D. caryophyllus, L.</i> . .	<i>Carnation</i>	Old walls!	Pk.	1-1½ ft.	P.	7-8
v. <i>D. cœsius, Sm.</i>	<i>Cheddar P.</i>	Lim. rocs.!	Pk.	4-12 in.	P.	6-7
vi. <i>D. deltoides, L.</i>	<i>Maiden P.</i>	Hilly pas.!	Ro.	6-12 in.	P.	6-9
v. 2. <i>glaucus, L.</i>	. . . . .	Edinbro'.!	. .	. .	. .	. .
Gen. II. Saponaria, L.	<i>Soapwort.</i>					
i. <i>S. officinalis, L.</i>	<i>common S.</i>	Hedges .	Ro.	1-1½ ft.	P.	8
v. 2. <i>hybrida, Mil.</i>	. . . . .	Waysides	. .	. .	. .	. .
Gen. III. Silene, L.	<i>Catchfly, Campion.</i>					
*						
i. <i>S. inflata, Sm.</i>	<i>Bladder Cam.</i> . . .	Past. . .	W.	2-3 ft.	P.	6-7
v. 2. <i>puberula, Ford.</i>	. . . . .	!	. .	. .	. .	. .
ii. <i>S. unaritima, With.</i> . . .	<i>Sea Cam.</i>	Sea sh. .	W.	3-6 in.	P.	6-8
***						
iii. <i>S. Otites, Sm.</i>	<i>Spanish Catch.</i> . .	San. fie.!	y G.	1-3 ft.	P.	6
iv. <i>S. Anglica, L.</i>	<i>English Catch.</i> . .	San. fie. .	pkW.	1-2 ft.	A.	6-9
v. 2. <i>quinquevulnera, L.</i> . . .	. . . . .	Jersey .	W.	. .	. .	. .
v. <i>S. nutans, L.</i>	<i>Nottingham Catchfly.</i>	Lim. rocs.!	W.	2-3 ft.	P.	6-7
v. 3. <i>paradoxa, Sm.</i>	. . . . .	Dover .	. .	. .	. .	. .
vi. <i>S. noctiflora, L.</i>	<i>night flowering C.</i>	San. fie.!	p R.	1-2 ft.	A.	7-8
vii. <i>S. conica, L.</i>	<i>striated Corn. C.</i>	San. fie.!	Ro.	6-12 in.	A.	5-6
viii. <i>S. acaulis, L.</i>	<i>Moss. Camp.</i> . . .	Alp. rocks	Pk.	2-3 in.	P.	7-8
Gen. IV. Lychnis, L.	<i>Campion, Catch- fly.</i>					
i. <i>L. alpina, L.</i>	<i>red Alpine Catch.</i>	Sc. mount.!	Ro.	5-6 in.	P.	6-7
ii. <i>L. viscaria, L.</i>	<i>red German Catch.</i>	Lim. rocs.!	Ro.	6-10 in.	P.	6-7
iii. <i>L. flos-cuculi, L.</i>	<i>ragged Robin.</i> . . .	Moi. mea.	Ro.	1-2 ft.	P.	5-6
iv. <i>L. diurna, Siph.</i>	<i>red Campion.</i> . . .	Dp. banks	R.	1-2 ft.	P.	6-7

ORD. XIII.—ELATINACEÆ.

CHARACTERISTICS OF SPECIES AND VARIETIES

Gen. I.

- i. Fls. stalked ; pet. three ob.-ov.—Sta. six—Seeds straight.
- ii. Fls. sess. ; pet. four ov.—Sta. eight—Seeds bent.

ORD. XIV.—CARYOPHYLLACEÆ.

Gen. I.

* *Flowers clustered.*

- i. Fls. capitt.—Ls. lin. lanc. ; margins rough—Cal. scales ov., blunt., membr.
- ii. Fls. fascicled—Ls. lin., opp., conn.—Cal. scales lanc., downy.

* * *Flowers solitary.*

- iii. Ls. lin., acute, one-nerved, rough at margin—Fls. sol. in a lax. cyme ; pet multifid.]
- iv. Ls. grooved, glauc., smooth at margin—Fls. in a lax. cyme ; pet. toothed.
- v. St. one flowered—Ls. lin., glauc., rough at marg.—Cal. scales orb., very short.
- vi. Ls. flat, downy—Pet. toothed—Cal. scales two lanc., acute.
- v. 2. *Ls. glauc.—Cal. scales four—Fls. white.*

Gen. II.

- i. Ls. ovate. lanc., glab.—Calyx cylind.
- v. 2. *Ls. conn.—Corolla monopetalous.*

Gen. III.

* *Calyx inflated.*

- i. St. erect—Fls. pan.—Ls. glab.—Pet. deeply cleft.
- v. 2. *Ls. downy.*
- ii. St. decumb.—Fls. sol.—Pet. shortly cleft.

* * *Calyx cylindrical.*

- iii. St. erect, nearly simp.—Ls. spath.—Fls. small whorled—Pet. lin., ent.
- iv. St. erect, viscid.—Ls. lanc.—Fls. sol., axill. alt.—Pet. bifid.
- v. 2. *Pet. entire, white, with a pink spot.*
- v. St. erect, viscid and hairy—Ls. lanc.—Fls. pan., drooping—Pet. deeply cut into lin. segm.
- v. 2. *"Broader" Bab. "not distinguishable" Syme.*
- vi. St. erect, dichot.—Ls. obl.-lanc., acute—Fls. in a forked panicle—Pet. bifid, obt., crowned.
- vii. St. erect, forked, hairy—Ls. lin., lanc.—Fls. term. and in the forks—Cal. coni., many-furrowed.
- viii. St. tufted, one fld.—Ls. lin., ciliate at base.

Gen. IV.

- i. St. glab.—Ls. lin., lanc.—Fls. capit., corymb. ; pet. bifid
- ii. St. viscid—Ls. lanc.—Fls. pan. ; pet. slightly notched.
- iii. St. viscid—Ls. lanc.—Fls. pan. ; pet. four cleft.
- iv. St. hairy—Ls. ovate.-lanc.—Fls. diæcious ; cal. teeth triang.—Caps. glob. ; its teeth recurved.

ORD. XIV.—CARYOPHYLLACEÆ—*cont.*

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
v. <i>L. vespertina</i> , <i>Sipth.</i> . .	<i>white Champion.</i> . .	Fie.. . .	W.	1-2 ft.	P.	6-7
vi. <i>L. Gigatho</i> , <i>Lam.</i> . .	<i>Corn-cockle.</i> . . .	Corn fie..	p Pu.	1-2 ft.	A.	7-8
Gen. V. <i>Mœnchia</i>, Ehrh.	<i>Mœnchia.</i>					
i. <i>M. erecta</i> , <i>Sm.</i> . . .	<i>erect M.</i>	Dry pl. .	W.	2-6 in.	A.	5-6
Gen. VI. <i>Sagina</i>, L.	<i>Pearl-wort.</i>					
i. <i>S. apetala</i> , <i>L.</i>	<i>annual small fld.</i> <i>P.</i>	Dry pl. .	W.	4-10 in.	A.	6-9
ii. <i>S. ciliata</i> , <i>Fr.</i>	<i>ciliated P.</i> . . .	Dry pl. .	W.	Cr.	A.	5-6
iii. <i>S. maritima</i> , <i>Don.</i> . .	<i>Sea P.</i>	Sea co. .	r W.	Cr.	A.	5-9
v. 2. <i>debilis</i> , <i>Ford.</i> . . .	. . . . .	. . . . .	. .	. .	. .	. .
v. 3. <i>densa</i> , <i>Ford.</i>	. . . . .	E. of E.!	. .	. .	. .	. .
v. 4. <i>alpina</i> , <i>Syme.</i>	. . . . .	BenNevis!	. .	. .	. .	. .
iv. <i>S. procumbens</i> , <i>L.</i> . . .	<i>procumbent P.</i> . .	Dry pl. .	W.	Cr.	P.	5-9
v. <i>S. saxatilis</i> , <i>Wimm.</i> . .	<i>Alpine P.</i>	Sc. moun.!	W.	Cr.	P.	6-8
vi. <i>S. subulata</i> , <i>Wimm.</i> . .	<i>Awl-shaped P.</i> . .	Dry pl. .	W.	Cr.	P.	6-8
vii. <i>S. nivalis</i> , <i>Fr.</i>	. . . . .	Sc. moun.!	W.	Cr.	P.	6-8
viii. <i>S. nodosa</i> , <i>L.</i>	<i>knotted P.</i> . . .	Damp. pl.	W.	3-4 in.	P	7-8
Gen. VII. <i>Spergula</i>, L.	<i>Spurrey</i>					
i. <i>S. arvensis</i> , <i>L.</i>	<i>Corn-sp.</i>	Corn fie..	W.	Cr.	A.	6-8
Gen. VIII. <i>Honckenia</i>, Ehrh.	<i>Sea-purslane.</i>					
i. <i>H. peplodes</i> , <i>Ehrh.</i> . .	<i>Sea-purslane.</i> . .	Sea co. .	W.	Cr.	P.	5-8
Gen. IX. <i>Spergularia</i>, Pers.	<i>Sandwort- Spurrey.</i>					
i. <i>S. marina</i>	<i>Sea-side S.</i>	Sea co. .	W.	Cr.	A. or P.	6-9
a. <i>neglecta</i> , <i>Sy.</i>	. . . . .	. . . . .	. .	. .	A.	. .
v. 2. <i>salina</i>	. . . . .	. . . . .	. .	. .	. .	. .
v. 3. <i>media</i>	. . . . .	. . . . .	. .	. .	. .	. .
b. <i>marginata</i> , <i>Sy.</i>	. . . . .	. . . . .	. .	. .	P.	. .
c. <i>rupestris</i> , <i>Lebel.</i> . . .	. . . . .	Sea co.!	. .	. .	P.	. .
ii. <i>S. rubra</i> , <i>Fen.</i>	<i>Field S.</i>	Dry pl. .	W.	Cr.	A.	6-9
Gen. X. <i>Arenaria</i>, L.	<i>Sandwort.</i>					
i. <i>A. ciliata</i> , <i>L.</i>	<i>fringed S.</i> . . .	Ir. moun.!	W.	Cr.	P.	5-6
ii. <i>A. Norvegica</i> , <i>Gunn.</i>	<i>Norwegian S.</i> . .	Sc. moun.!	W.	Cr.	P.	5-6
iii. <i>A. serpyllifolia</i> , <i>L.</i> . .	<i>Thyme-leaved S.</i>	Dry pl. .	W.	Cr.	A.	5-6
v. 1. <i>spærocarpa</i> , <i>Te.</i> . . .	. . . . .	. . . . .	. .	. .	. .	. .
v. 2. <i>Lloydii</i> , <i>Ford.</i>	. . . . .	. . . . .	. .	. .	. .	. .
v. 3. <i>leptoclados</i> , <i>Gus.</i> . .	. . . . .	. . . . .	. .	. .	. .	. .

ORD. XIV.—CARYOPHYLLACEÆ—*cont.*

CHARACTERISTICS OF SPECIES AND VARIETIES.

- v. Like No. 4, but fls. white—Cal. teeth lin. lanc.—Caps. con. its teeth erect.
 vi. St. erect, branched—Fls. large sol. ; cal. teeth woolly, longer than cor.

Gen. V.

- i. L. lin.-lanc., rigid, glauc.—Sep. with membr. marg.

Gen. VI.

- i. Slightly pubesc.—St. ascending—Ls. aristate, fringed—Sep. spread. longer than caps.
 ii. Gland-pubesc.—St. decumb.—Ls. mucro., cil.—Sep. adpr. to caps.
 iii. Glab.—St. slender erect or decumb., with long internodes—Ls. lin., fleshy, blunt—Pedunc. erect—Pet. small or none—Caps. equal to pet.
 v. 2. *Procumb., slend.—Internodes long—Caps. shorter than sep.*
 v. 3. *Erect—St. slender, branched with short internodes—Caps. as Var. 2.*
 v. 4. *Erect—St. stout with short internodes—Caps. as Var. 2.*
 iv. Glab.—St. procumb.; central very short, flowerless; branches long, rooting—Ls. lin., awned—Pedunc. recurved after flowering; sep. spread. in fr.
 v. Closely resembling Sp. 4, but sep. adpressed in fruit, pet. longer, and stamens 10.
 vi. Tufted, gland.-hairy—Ls. lin., awned, ciliate.—Pedunc. long, curved after flowering.
 vii. Densely tufted, glab.—Ls. acum.—Pedunc. always erect.
 (The three last species might well be combined.)
 viii. St. procumb.—Ls. subu. glab., conn. with leaf buds in their axils—Fls. 2-3 term.—Pet. much longer than sep.

Gen. VII.

- i. Ls. whorled, lin. fleshy—Fr. stalks reflexed—Seeds rough.

Gen. VIII.

- i. Ls. ovate, acute, fleshy—Fls. sol., sess. ; pet. not longer than sep.

Gen. IX.

- i. Per.—Ls. lin. fleshy, convex below, blunt—Caps. longer than sep.—Bracts. delt. ovate.
 a. Ann.—Bracts., scarious, shorter than pedi.—Seeds smooth.
 v. 2. *Bracts leafy, as long as pedi.—Seeds winged.*
 v. 3. *Bracts leafy, longer than pedi.—Seeds winged.*
 b. Perenn. larger and quite glab.—Fls. paler—Stip. dull white.
 c. Perenn. resembling Sub.-sp. marginata—But pedicels sep. and stip. gland.-pubesc. and stip. silvery white.
 ii. L. lin., flat, pointed—Caps. not longer than sep.—Stip. lanc., toothed.

Gen. X.

- i. St. procumb., branched—Ls. spath., ciliate, fls. sol.—Sep. half as long as pet. prominently ribbed.
 ii. Like Sp. 1., but ls. fleshy, glab.—Sep. broader and obscurely ribbed.
 iii. St. decumb.—Ls. ovate., acute, sess., rigid—Sep. longer than pet., hairy.
 v. 1. *Rigid—Caps. ovoid., pedicel erect.*
 v. 2. *Stout—Caps. swollen at base (I. of W.).*
 v. 3. *Weak—Caps. narrow, pedicel spreading.*

ORD. XIV.—CARYOPHYLLACEÆ—cont.

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
iv. <i>A. tenuifolia</i> , L. . . .	<i>fine-leaved S.</i> . . .	Sandy fie.	W.	4-6 in.	A.	5-6
v. 2. <i>laxa</i> , Ford.	. . . . .	. . . . .	. . .	. . .	. . .	. . .
v. 3. <i>viscosa</i> , Bab.	. . . . .	. . . . .	. . .	. . .	. . .	. . .
v. <i>A. verna</i> , L.	<i>vernal S.</i>	Dry rocks	W.	2-4 in.	P.	5-9
v. 2. <i>Gerardi</i> , Will. . . .	. . . . .	. . . . .	. . .	. . .	. . .	. . .
vi. <i>A. rubella</i> , Hook. . . .	<i>Alpine S.</i>	Sc. moun.!	W.	1-2 in.	P.	7-8
vii. <i>A. uliginosa</i> , Schl. . .	<i>Bog S.</i>	Teesdale !	W.	2-3 in.	P.	6-7
viii. <i>A. trinervis</i> , L. . . .	<i>three-nerved S.</i> . . .	Damp pl.	W.	6-12 in.	A.	5-6
Gen. XI. <i>Holosteum</i> , L.	<i>Jagged Chickweed.</i>					
i. <i>H. umbellatum</i> , L. . . .	<i>umbelliferous</i> <i>?</i> . .	Old walls !	W.	4-8 in.	A.	4-5
Gen. XII. <i>Stellaria</i> , L.	<i>Stitchwort.</i>					
i. <i>S. nemorum</i> , L.	<i>Wood S.</i>	Dp. wds.	W.	1-2 ft.	P.	5-6
ii. <i>S. media</i> , With.	<i>Chickweed.</i>	every- where.	W.	6-18 in.	A.	3-9
v. 2. <i>Borœana</i> , Ford. . . .	. . . . .	. . . . .	. . .	. . .	. . .	. . .
v. 3. <i>neglecta</i> , Wei.	. . . . .	. . . . .	. . .	. . .	. . .	. . .
v. 4. <i>umbrosa</i> , Opi.	. . . . .	. . . . .	. . .	. . .	. . .	. . .
iii. <i>S. holostea</i> , L.	<i>greater S.</i>	Hedges .	W.	1-2 ft.	P.	4-6
iv. <i>S. glauca</i> , With.	<i>glaucous S.</i>	Mar. . .	W.	1-2 ft.	P.	5-7
v. <i>S. graminea</i> , L.	<i>lesser S.</i>	Dry pl. .	W.	1-2 ft.	P.	5-8
vi. <i>S. uliginosa</i> , Murr. . . .	<i>Bog S.</i>	Bogs . .	W.	3-12 in.	A.	5-6
vii. <i>S. cerastoides</i> , L. . . .	<i>Alpine S.</i>	Sc. moun.!	W.	4-6 in.	P.	7-8
Gen. XIII. <i>Cerastium</i> , L.	<i>Mouse-ear Chick- weed.</i>					
i. <i>C. aquaticum</i> , L.	<i>Water M.</i>	Wet pl. .	W.	1-2 ft.	P.	7-8
ii. <i>C. glomeratum</i> , Thuil.	<i>broad-leaved M.</i> . .	Dry pl. .	W.	6-10 in.	A.	4-9
iii. <i>C. triviale</i> , Link. . . .	<i>narrow-leaved M.</i>	Dry pl. .	W.	6-10 in.	A. or B.	4-9
v. 2. <i>holosteoides</i> , Fri. . . .	. . . . .	N. of E..	. . .	. . .	. . .	. . .
v. 3. <i>pentandrum</i>	. . . . .	Dry walls	. . .	. . .	. . .	. . .
iv. <i>C. semidecandrum</i> , L.	<i>little M.</i>	Sea sh. .	W.	1-8 in.	A.	4-5
v. 2. <i>pumilum</i> , Curt.	. . . . .	!	. . .	. . .	. . .	. . .
v. <i>C. tetrandrum</i> , Curt. . .	<i>Four-cleft M.</i> . . .	San. pl. .	W.	6-12 in.	A.	5-7
vi. <i>C. arvense</i> , L.	<i>Field M.</i>	Dry pl. .	W.	6-10 in.	P.	4-8
v. 2. <i>Andrewsii</i> , Sy.	. . . . .	Arran.!	. . .	. . .	. . .	. . .
vii. <i>C. alpinum</i> , L.	<i>hairy Alpine M.</i>	Alp. rocs.!	W.	3-5 in.	P.	6-8
v. 2. <i>pubescens</i>	. . . . .	. . . . .	. . .	. . .	. . .	. . .
viii. <i>C. latifolium</i> , L. . . .	<i>broad-leaved Alp. M.</i>	Alp. rocs.!	W.	3-5 in.	P.	5-8

CHARACTERISTICS OF SPECIES AND VARIETIES.

- iv. St. erect—slender, branched—Ls. lin., lanc., glab.—Sep. three-nerved, as long as pet.—Sta. 10.
- v. 2. Sta. 5—*Sep. gland.*, shorter than caps. (*Great Wilbraham*).
- v. 3. Sta. 8–10—*Pedunc. and sep. gland.-hairy*—*Sep. as long as caps* (*Thetford*).
- v. St. tufted—Ls. lin., subu. acute, three-veined—Sep. three-nerved, shorter than the obl. pet.
- v. 2. *Ls. blunt; lower ones adpressed.*
- vi. St. densely tufted—Ls. subu. obtuse, three-veined—Fls. sol.—Sep. acute, longer than the lanc. pet.
- vii. St. loosely tufted—Ls. subu., veinless—Pedunc. very long, 1–3-flowered—Sep. ov.-acute, as long as the ob.-ov. pet.
- viii. St. diffuse, hairy—Ls. ov., acute, peti. cili.—Fls. axill., sol., long-stalked.

Gen. XI.

- i. Ls. ellip., ovate—Fls. umbell.—Pedunc. viscid (E. of Eng.).

Gen. XII.

- i. Lower ls. cord., stalked; upper ov. sess.—Pan. dichot.
- ii. St. decumb., with a hairy, alternately lateral line—Ls. ovate.—Sta. 5.
- v. 2. *Petals absent*—Sta. 3.
- v. 3. *St. suberect*—Ls. larger—Sta. 10.
- v. 4. *St. erect, slender*—*Pedi. glab.*—*Sep. glab. with raised points.*
- iii. St. erect, ang.—Ls. lanc., serr., acum.—Sep. nerveless, half as long as pet.
- iv. St. erect, angled—Ls. lin. lanc., ent., glauc.—Fls. sol., axill.—Sep. three-nerved, shorter than pet.
- v. St. diffuse—Ls. lin., lanc., ent., glab.—Sep. three-nerved, as long as pet.
- vi. St. diffuse—Ls. ovate, lanc., callous at tip—Sep. longer than pet.
- vii. St. decumb., with an alt., hairy line, two-flowered—Ls. obl., obt.—Sep. one-nerved, downy.

Gen. XIII.

- i. St. long, straggling—Ls. large cord. sess.—Fls. axill., sol.
* *Annual*—*Sep. as long as pet.*
- ii. St. erect, viscid—Ls. oval, pale green—Pet. equal to sep.; pedi. shorter than sep.
- iii. St. diffuse, viscid, hairy—Ls. lanc., deep green; pedi. longer than hairy sep.
- v. 2. *St. with two lines of hairs*—*Sep. glab*—Sta. 10.
- v. 3. *Annual*—*Sep. hairy*—Sta. 5—*Caps. shorter.*
- iv. St. sub-erect—Cymes few flowered—Pet. notched; sta. 5; bracts half membranous—Caps. shorter than sep.
- v. 2. *Bracts with a narrow memb. margin*—*Caps. longer.*
- v. St. diffuse—Ls. ovt.—Sep. 4—Fr. stalks three or four times as long as sep.
* * *Perennial*—*Sep. shorter than pet.*
- vi. St. hairy—Ls. lin. lanc.—Pet. twice as long as sep.
- v. 2. *Ls. rigid, glab.*—*Fls. solitary.*
- vii. St. ascending—Ls. ov. covered with long white hairs—Pan. dichot, few-flowered—Sep. with a narrow memb. margin.
- v. 2. *Covered with short hairs.*
- viii. St. prost., tufted—Ls. ov., covered with short yellowish hairs—Fls. sol.—Sep. with a broad memb. margin.

ORD. XIV.—CARYOPHYLLACEÆ—*cont.*

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
<i>v. 2. compactum, Sy. . .</i>	. . . .	Braemar !	. .	. .	. .	. .
<i>v. 3. nigrescens, Sy. . .</i>	. . .	Unst. !	. .	. .	. .	. .
Gen. XIV. Cherleria, L.	<i>Cyphel.</i>					
<i>i. C. sedoides, L. . . .</i>	<i>Mossy C. . . . .</i>	Sc. mount. !	γ G.	1-2 in.	P.	6-8

ORD. XV.—LINACEÆ.

Gen. I. Linum, L.	<i>Flax.</i>					
<i>i. L. usitatissimum, L. . .</i>	<i>common F. . . .</i>	Corn fie.	B.	1-1½ ft.	A.	7
<i>ii. L. perenne, L. . . .</i>	<i>perennial F. . . .</i>	Cha. hills !	B.	1-2 ft.	P.	6-7
<i>iii. L. angustifolium, Huds.</i>	<i>narrow-leaved F.</i>	Cha. hills	ϕ B.	1-2 ft.	P.	7-8
<i>iv. L. catharticum, L. . .</i>	<i>purging F. . . .</i>	Heaths .	W.	2-10 in.	A.	6-9
Gen. II. Radiola, Gme.	<i>Flax seed.</i>					
<i>i. R. millegrana, Sm. .</i>	<i>Thyme-leaved F.</i>	Damp pl.	W.	1-4 in.	A.	7-8

ORD. XVI.—MALVACEÆ.

Gen. I. Malva, L.	<i>Mallow.</i>					
<i>i. M. moschata, L. . . .</i>	<i>musk M. . . . .</i>	Grav. pl.	Ro.	2-3 ft.	P.	7-8
<i>ii. M. sylvestris, L. . . .</i>	<i>common M. . . . .</i>	Was. pl. .	ϕ Pu.	2-3 ft.	P.	6-9
<i>iii. M. rotundifolia, L. .</i>	<i>dwarf M. . . . .</i>	Was. pl. .	Li.	6-12 in.	P.	6-9
Gen. II. Althæa, L.	<i>Marsh Mallow.</i>					
<i>i. A. officinalis, L. . . .</i>	<i>common M. . . . .</i>	Mar. . .	Ro.	2-3 ft.	P.	8-9
<i>ii. A. hirsuta, L. . . . .</i>	<i>hispid M. . . . .</i>	Cuxton !.	Pu.	6-18 in.	A.	6-7
Gen. III. Lavatera, L.	<i>Tree Mallow.</i>					
<i>i. L. arborea, L. . . . .</i>	<i>Sea T. . . . .</i>	Sea rocs.	Pu.	3-5 ft.	B.	7-9

ORD. XVII.—TILIACEÆ.

Gen. I. Tilia, L.	<i>Lime.</i>					
<i>i. T. parviflora, Ehrh. .</i>	<i>small-leaved L. .</i>	Woods .	γ G.	. .	T.	7-8
<i>ii. T. intermedia, DC. .</i>	<i>common L. . . .</i>	Woods .	γ G.	. .	T.	6-7
<i>iii. T. grandiflora, Ehrh.</i>	<i>broad-leaved L. .</i>	Woods .	γ G.	. .	T.	6-7
<i>v. 2. corallina, Sm. . .</i>	. . . .	. . .	. .	. .	. .	. .

ORD. XIV.—CARYOPHYLLACEÆ—*cont.*

CHARACTERISTICS OF SPECIES AND VARIETIES.

- v. 2. *Densely tufted—Fl. st. short—Sep. gland.*
 v. 3. *Deep green—St. ls. broader—Hairs all gland.*

Gen. XIV.

- i. *Densely tufted—Ls. crowded, channelled—Fls. sol. sess.*

ORD. XV.—LINACEÆ.

Gen. I.

- i. *Ann.—St. sol.—Ls. alt., lin.-lan.—Sep. ov., acute three-nerved, ciliate.*
 ii. *Peren.—Sts. numerous—Ls. alt., lin.-acute—Sep. ob.-ov., five-ribbed, obt., glab.*
 iii. *Peren.—Sts. numerous—Ls. alt., lin.-lanc.—Sep. ellip., three-ribbed, mucro.*
 iv. *Ann.—St. much branched—Ls. opp., obl.—Sep. acumi. one-nerved.*

Gen. II.

- i. *St. repeatedly forked—Ls. ov. ent.—Fls. axill. and term, sol.*

ORD. XVI.—MALVACEÆ.

Gen. I.

- i. *St. erect—Ls. reni. 5-7 cleft, lobes pinnfid.—Fr. hairy.*
 ii. *St. ascend.—Ls. reni. 5-7 cleft, lobes cren.—Fr. glab.*
 iii. *St. prost.—Ls. cord. 5-7 cleft, lobes cren.—Fr. hairy.*

Gen. II.

- i. *Ls. ent., five lobed, velvety—Fls. cymose, axill.*
 ii. *St. hispid—Ls. long stalked, palmately lobed—Fls. solit.*

Gen. III.

- i. *St. woody—Ls. seven-angled, plaited, downy—Pedunc. axill., one-flowered.*

ORD. XVII.—TILIACEÆ.

Gen. I.

- i. *Ls. small, glauc. beneath, with a woolly tuft in the axils of the nerves—Fr. ang., thin and brittle.*
 ii. *Ls. glab. beneath, with a woolly tuft in the axils of the nerves—Fr. coriaceous, hairy, not ribbed.*
 iii. *Ls. large, downy beneath—Pedunc. 3 fld.—Fr. downy, with 3-5 prominent ribs.*
 v. 2. *Bark of young twigs red.*

ORD. XVIII.—HYPERICACEÆ.

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
Gen. I. Hypericum, L. *	<i>St. John's Wort,</i> <i>&c.</i>					
i. <i>H. Androsænum, L.</i> .	<i>common Tutsan.</i>	Thick. .	Y.	1-2 ft.	P.	6-8
ii. <i>H. calycinum, L.</i> . .	<i>large-flowered St.</i> <i>Ƴ.</i>	Thick.! .	Y.	10-15 in.	P.	7-9
A.						
iii. <i>H. perforatum, L.</i> . .	<i>common St. Ƴ.</i> .	Thick. .	Y.	1-2 ft.	P.	7-9
v. 2. <i>angustifolium.</i> . .						
iv. <i>H. quadrangulum, L.</i>	<i>square-stalked St.</i> <i>Ƴ.</i>	Moi. pl. .	Y.	1-2 ft.	P.	7-9
a. <i>dubium, Leers.</i> . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
v. 2. <i>maculatum, Bab.</i> .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
b. <i>tetrapterum, Fries.</i> .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
c. <i>boeticum, Boiss.</i> . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
v. <i>H. humifusum, L.</i> . .	<i>trailing St. Ƴ.</i> . .	Commons	Y.	4-10 in.	P.	7-8
B.						
vi. <i>H. linariifolium, Vahl.</i>	<i>linear-leaved St. Ƴ.</i>	Corn. and Jersey!	Y.	6-12 in.	P.	7-8
vii. <i>H. pulchrum, L.</i> . . .	<i>small St. Ƴ.</i> . .	Commons	Y.	1-2 ft.	P.	7-8
viii. <i>H. hirsutum, L.</i> . . .	<i>hairy St. Ƴ.</i> . . .	Thick. .	Y.	1-3 ft.	P.	7-8
ix. <i>H. montanum, L.</i> . . .	<i>bearded St. Ƴ.</i> . .	Thick. .	Y.	1-2 ft.	P.	7-8
x. <i>H. elodes, L.</i>	<i>Marsh St. Ƴ.</i> . .	Bogs . .	Y.	3-12 in.	P.	7-8

ORD. XIX.—ACERACEÆ.

Gen. I. Acer, L.	<i>Maple.</i>					
i. <i>A. campestre, L.</i> . .	<i>common M.</i>	Hedges .	G.	10-20 ft.	T.	5-6
ii. <i>A. pseudo-platanus, L.</i>	<i>Sycamore</i>	Woods .	G.	40-50 ft.	T.	5-6

ORD. XX.—GERANIACEÆ.

Gen. I. Geranium, L. *	<i>Crane's-bill.</i>					
i. <i>G. sanguineum, L.</i> . .	<i>bloody C.</i>	Rocky pl.!	Cr.	1-2 ft.	P.	7-8
v. 2. <i>Lancastriense, With.</i> .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
* *						
A.						
ii. <i>G. phœum, L.</i>	<i>dusky C.</i>	Thick. .	Pu.	1-2 ft.	P.	6-7
iii. <i>G. sylvaticum, L.</i> . .	<i>Wood C.</i>	Woods .	b Pu.	1-2 ft.	P.	6-7

ORD. XVIII.—HYPERICACEÆ.

CHARACTERISTICS OF SPECIES AND VARIETIES.

Gen. I.

* *Stamens in five sets.*

i. St. erect, shrubby—Ls. large, ov., blunt, sess.—Fls. in a term cyme.—Sty. three, recurved.

ii. St. creeping, shrubby—Ls. obl. sess.—Fls. sol., large—Sty. five, straight.

* * *Stamens in three sets.**A. Sepals with entire or toothed margins without glands.*

iii. St. two-edged—Ls. obl., obt., with pellucid dots and opaque veins—Sep. lanc., ent., erect.

v. 2. *Ls. strap-shaped.*

iv. St. four-angled—Ls. obl., with pellucid dots.

a. St. with four elevated lines—Ls. with pellucid veins—Pet. broad, ent.—Sep. spreading, recurved.

v. 2. *Ls. narrower—Sep. narrow, denticulate.*

b. St. with four wings—Fls. in compact many small flowered cymes—Sep. lanc., erect, acute.

c. Ls. with black glands beneath—Sep. ov., lanc., crenate (*Cornwall*).

v. St. weak, prost.—Ls. obl., glab.—Sep. unequal.

B. Sepals with glandular margins.

vi. St. erect, terete—Ls. lin., margins revolute—Sep. lanc., acute.

vii. St. erect, round, glab.—Ls. cord., amplex—Sep. ov.-obt.—Fls. tipped with red in bud.

viii. St. erect, hairy—Ls. ov.-obl., shortly stalked—Sep. lanc., acute.

ix. St. erect, glab.—Ls. ov., sess.—Sep. lanc. acute, with black glands—Fls. in dense cymes.

x. St. prost., rooting—Ls. orb., sess., hairy—Fls. few, pale, panicled.

ORD. XIX.—ACERACEÆ.

Gen. I.

i. Ls. five-lobed, crenate—Racemes erect.

ii. Ls. five-lobed, unequally serr.—Racemes pendulous.

ORD. XX.—GERANIACEÆ.

Gen. I.

* *Pedunc., one-flowered.*

i. St. sub-erect—Ls. orb. 5-7 part, with trifid lobes.

v. 2. *St. decumb., tufted—Fls. pink (Walney Island).** * *Pedunc. two-flowered.**A. Root perennial.*

ii. Ls. orb. 3-7 lobed, lobes acute, cut, serr.—Pet. waved, dingy purple.

iii. St. erect, corymb.—Ls. sub-pelt, seven-part, lobes cut and serr.—Pet. slightly notched.

ORD. XX.—GERANIACEÆ—*cont.*

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
iv. <i>G. pratense</i> , L. . . .	<i>blue Meadow C.</i> . .	Thick. .	Pu.	1-3 ft.	P.	6-7
v. <i>G. pyrenaicum</i> , L. . .	<i>Mountain C.</i> . .	Past. . .	r Pu.	1-2 ft.	P.	6-8
<i>B.</i>						
vi. <i>G. rotundifolium</i> , L. .	<i>round-leaved C.</i> . .	Was. pl..	Pk.	6-12 in.	A.	6-7
vii. <i>G. pusillum</i> , L. . . .	<i>small-flowered C.</i> . .	Was. pl..	Ro.	6-18 in.	A.	6-9
viii. <i>C. molle</i> , L.	<i>Dove's-foot C.</i> . . .	Was. pl..	Pk.	8-12 in.	A.	4-8
ix. <i>G. dissectum</i> , L. . . .	<i>jagged-leaved C.</i> . .	Was. pl..	R.	1-2 ft.	A.	5-8
x. <i>G. columbinum</i> , L. . .	<i>long-stalked C.</i> . .	Dry pl. .	Ro.	1-2 ft.	A.	6-7
xii. <i>G. Robertianum</i> , L. .	<i>Herb Robert</i> . . .	Was. pl..	Pk.	1-2 ft.	A.	5-9
v. 2. <i>modestum</i> , Ford. . .	. . . . .	. . . . .	. .	. .	. .	. .
v. 3. <i>purpureum</i> , Forst..	. . . . .	. . . . .	. .	. .	. .	. .
xii. <i>G. lucidum</i> , L. . . .	<i>skining C.</i>	Rocks and Walls.	Ro.	6-18 in.	A.	5-9
Gen. II. <i>Erodium</i>, L'Herit. <i>Stork's-bill.</i>						
i. <i>E. maritimum</i> , Sm. . .	<i>Sea S.</i>	San. Sh.!	Pk.	6-18 in.	B.	5-9
ii. <i>E. cicutarium</i> , Sm. . .	<i>Hemlock S.</i>	Was. pl..	Ro.	6-18 in.	B.	6-9
v. 2. <i>chærophylloides</i> , . .	. . . . .	. . . . .	. .	. .	. .	. .
iii. <i>E. moschatum</i> , Sm. . .	<i>musky S.</i>	Was. pl..	Pu.	1-2 ft.	A.	6-7

ORD. XXI.—BALSAMINACEÆ.

Gen. I. <i>Impatiens</i>, L. <i>Balsam.</i>						
i. <i>I. fulva</i> , Nutt.	. . . . .	Riv. bks.!	Y.	2-3 ft.	A.	8-9
ii. <i>I. noli-me-tangere</i> , L.	<i>yellow B.</i>	Moi. Wds.!	Y.	1-2 ft.	A.	7-9

ORD. XXII.—OXALIDACEÆ.

Gen. I. <i>Oxalis</i>, L. <i>Wood Sorrel.</i>						
i. <i>O. acetosella</i> , L. . . .	<i>common W.</i>	Woods .	W.	3-6 in.	P.	5-6
ii. <i>O. corniculata</i> , L. . .	<i>yellow W.</i>	Waste pl., S. of Eng.!	Y.	6-18 in.	A.	6-9
iii. <i>O. stricta</i> , L.	. . . . .	Waste pl.	Y.	6-18 in.	A.	6-9

ORD. XXIII.—CELASTRACEÆ.

Gen. I. <i>Euonymus</i>, L. <i>Spindle tree.</i>						
i. <i>E. Europæus</i> , L. . . .	<i>common S.</i>	Hedges .	g W.	5-20 ft.	T.	5-6

ORD. XX.—GERANIACEÆ—*cont.*

CHARACTERISTICS OF SPECIES AND VARIETIES.

- iv. St. erect—Ls. orb. 5-7 part; lobes lacin.; seg. acute—Pet. slightly notched—Fr. stalk reflexed.
 v. St. straggling—Ls. reni., 5-7 part, lobes obt., trifid—Pet. deeply notched—Root fusi.

B. Root annual.

- vi. Pet. entire—Caps. keeled, smooth, hairs spreading—Seeds dotted.
 vii. Pet. notched—Caps. keeled, hairs adpressed—Seeds smooth.
 viii. Pet. notched—Caps. wrinkled, glab.—Seeds smooth.
 ix. Pet. notched—Caps. even, hairy—Seeds pitted—Seg. of ls. lin.—Fls. shortly stlkd.
 x. Pet. entire—Caps. glab.—Seeds dotted—Fls. with long stalks.
 xi. Pet. entire—Caps. transversely wrinkled, downy—Seeds smooth—Ls. tern. or quin., with pinnfid. lfts.
 v. 2. *Glabrous—Sep. and pedi. glandular hairy.*
 v. 3. *Like Var. 2, but leaves finely divided.*
 xii. Pet. ent., ob.-ov.—Caps. reticulate—Seeds smooth—St. and ls. glab., spreading, tinged with red.

Gen. II.

- i. Pedunc. 1-2 flowered—Ls. ent. ov.-cord—Pet. small or o.
 ii. Pedunc. many-flowered—Ls. pinn., lfts pinnfid.—Stip. lanc.
 v. 2. *Lfts. with more divided and more acute seg.—Fls. larger and more numerous. (A very variable plant.)*
 iii. Larger than last—Lfts. less deeply and irregularly cut—Stip. ovate.

ORD. XXI.—BALSAMINACEÆ.

Gen. I.

- i. Ls. ac.—tip of closely recurved spur notched.
 ii. Ls. obt.—tip. of loosely recurved spur ent.

ORD. XXII.—OXALIDACEÆ.

Gen. I.

- i. Stemless—Ls. radi., trifol., long-stalked—Pedunc. one-flowered.
 ii. St. branched, hairy; branches procumb.—Pedunc. two-flowered, shorter than ls.
 iii. St. erect, with runners at base—Pedunc. many-flowered, longer than ls.—Stip. o.

ORD. XXIII.—CELASTRACEÆ.

Gen. I.

- i. Ls. ovate, lanc., serr.—Fls. four cleft, umbellate.

ORD. XXIV.—RHAMNACEÆ.

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth	Dura- tion.	Time of flowering.
Gen. I. Rhamnus, L.	<i>Buckthorn.</i>					
i. <i>R. catharticus, L.</i> . . .	<i>common B.</i> . . .	Thick. .	y G.	5-10 ft.	Shb.	5-7
ii. <i>R. Frangula, L.</i> . . .	<i>Alder B.</i> . . .	Thick. .	g W.	5-10 ft.	Shb.	5-7

ORD. XXV.—LEGUMINOSÆ.

Gen. I. Saromanthus,	<i>Broom.</i>					
<i>Wimm.</i>						
i. <i>S. scoparius, Koch.</i> . .	<i>common B.</i> . . .	Heaths .	Y.	2-6 ft.	Shb.	5-6
Gen. II. Ulex, L.	<i>Furze, Gorse.</i>					
i. <i>U. Europæus, L.</i> . . .	<i>common F. or G.</i>	Heaths .	Y.	2-5 ft.	Shb.	2-7
v. 2. <i>strictus, Mack.</i> . . .	<i>Irish F.</i>	I.	..	..	..	..
ii. <i>U. nanus, Forst.</i> . . .	<i>dwarf F.</i>	Heaths, S. of E.	Y.	1-2 ft.	Shb.	7-11
v. 2. <i>Gallii, Planch.</i> . . .	. . . . .	Heaths, W. of E.	..	..	..	..
Gen. III. Genista, L.	<i>Green Weed.</i>					
i. <i>G. tinctoria, L.</i>	<i>Dyer's G.</i>	Thick. .	Y.	1-2 ft.	Shb.	7-9
v. 2. <i>humifusa, Syme.</i> . . .	. . . . .		..	..	..	..
ii. <i>G. pilosa, L.</i>	<i>hairy G.</i>	Heaths! .	Y.	6-12 in.	Shb.	5-9
iii. <i>G. anglica, L.</i>	<i>Needle G. or Petty Whin.</i>	Heaths .	Y.	1-2 ft.	Shb.	5-6
Gen. IV. Ononis, L.	<i>Rest Harrow.</i>					
i. <i>O. arvensis, L.</i>	<i>common R.</i>	Heaths .	Ro.	1-2 ft.	P.	6-9
ii. <i>O. spinosa, L.</i>	<i>spiny R.</i>	Heaths .	Ro.	6-12 in.	P.	6-9
iii. <i>O. reclinata, L.</i>	<i>small spreading R.</i>	Sea cl.! .	Ro.	2-3 ft.	A.	6-7
Gen. V. Anthyllis, L.	<i>Kidney Vetch.</i>					
i. <i>A. vulneraria, L.</i> . . .	<i>common K., or Ladies' fingers.</i>	Dry pas.	Y.	6-16 in.	P.	6-8
v. 2. <i>Dillenii, Schultz.</i> . .	. . . . .	S. of E.! .	R.	..	..	..
Gen. VI. Medicago, L.	<i>Medick.</i>					
*						
[i. <i>M. sativa, L.</i>	<i>purple M., or Lu- cerne.</i>	Fields .	Pu.	1-2 ft.	P.	6-7
ii. <i>M. falcata, L.</i>	<i>yellow M.</i>	Dry bks.! .	Y.	6-24 in.	P.	6-7
v. 2. <i>sylvestris, Fries.</i> . . .	. . . . .		..	..	..	..
iii. <i>M. lupulina, L.</i>	<i>black M., or Non- such.</i>	Was. pl..	Y.	6-24 in.	A.	5-8
iv. <i>M. maculata, Sibt.</i> . . .	<i>spotted M.</i>	Past. . .	Y.	6-24 in.	A.	5-8
v. <i>M. denticulata, Willd.</i>	<i>reticulated M.</i> .	Fie.! E. & S. of E.	Y.	6-24 in.	A.	5-8

ORD. XXIV.—RHAMNACEÆ.

CHARACTERISTICS OF SPECIES AND VARIETIES.

Gen. I.

- i. Branches spinous—Ls. ov., serr.—Fls. four-cleft.
- ii. Branches unarmed—Ls. ob.-ov., ent.—Fls. five-cleft.

ORD. XXV.—LEGUMINOSÆ.

Gen. I.

- i. St. branched, ang., green—Ls. tern. ; lfts. ob.-ov. silky—Fls. large, on short pedicels.

Gen. II.

- i. St. erect—Cal. with lax ov. bracts, and spr. hairs—Wings longer than keel—Spines rigid.
- v. 2. *Branches very erect—Spines soft (Co. Down, I.).*
- ii. St. procumb.—Cal. with minute bracts and adpr. hairs—Wings shorter than keel—Spines weak, deflexed.
- v. 2. *More erect—Wings rather longer than keel, curved—Primary spines strong.*

Gen. III.

- i. St. erect, glab.—Ls. lanc.—Rac. slender—Pods glab.
- v. 2. *St. procumb.—Ls. ov.—Cal. and pods hairy (Kynance Cove, Cornwall).*
- ii. St. procumb., woody—Ls. lanc., obt.—Rac. short, leafy—Pods hairy.
- iii. St. ascend, spinous—Ls. ov.-lanc.—Fls. sol., axill.—Pods glab.

Gen. IV.

- i. St. procumb., unarmed, hairy all round—Pods erect, ov. shorter than cal.
- ii. St. erect, spinous, with two rows of hairs—Pods erect, ov. longer than cal.
- iii. St. ann., spreading, hairy—Pods reflexed, cylind.—(*Alderney, Galloway*).

Gen. V.

- i. St. ascend.—Ls. pinn. ; lfts. unequal—Fls. capit.
- v. 2. *Smaller—Fls. red—Involucral ls. longer.*

Gen. VI.

* *Pods glabrous.*

- i. St. erect—Fls. rac.—Pods twisted two or three times.]
- ii. St. decumb., slightly hairy—Fls. rac.—Pods falcate.
- v. 2. *Fls. turning to dark green—Pods semicircular.*
- iii. St. procumb.—Fls. in a dense ovate spike—Pods reni., veiny, black when ripe.
* * *Pods bristly.*
- iv. St. procumb.—Lfts. ob.-cord., often spotted—Pods compactly spiral, with four ridges and a central furrow—Stip. dent.
- v. St. prost.—Pods loosely spiral, deeply reticu. and spin.—Stip. lacin.

ORD. XXV.—LEGUMINOSÆ—cont.

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
v. 2. <i>apiculata</i> , Willd. . .						
vi. <i>M. minima</i> , L. . . .	<i>little Bur-weed</i> .	San. fie.!	Y.	6-10 in.	A.	5-7
Gen. VII. Melilotus,						
<i>Tourn.</i>						
i. <i>M. officinalis</i> , L. . .	<i>common M.</i> . . .	Fields. .	Y.	2-3 ft.	B.	6-8
ii. <i>M. arvensis</i> , Wallr. .	<i>Field M.</i>	Fields. .	Y.	2-3 ft.	B.	6-8
iii. <i>M. alba</i> , Lam. . . .	<i>white M.</i>	Was. pl. .	W.	2-3 ft.	B.	7-8
Gen. VIII. Trigonella, L.						
<i>Fennugreek.</i>						
i. <i>T. ornithopodioides</i> , DC.	. . . . .	San. pas.	W. or Pk.	2-8 in.	A.	6-7
Gen. IX. Trifolium, L.						
<i>A.</i>						
i. <i>T. subterraneum</i> , L. .	<i>subterranean T.</i> .	Dry. pas.	C.	6-18 in.	A.	5-6
ii. <i>T. glomeratum</i> , L. .	<i>smooth round- headed T.</i>	Dry. pas. E. & S. of Eng.	Pu.	6-12 in.	A.	6-7
iii. <i>T. strictum</i> , L. . .	<i>upright round- headed T.</i>	Roc. bks.! Corn. & Jer.	Ro.	2-6 in.	A.	6-7
iv. <i>T. suffocatum</i> , L. . .	<i>suffocated T.</i> . .	San. sh.! E. & S. of Eng.	W.	1-3 in.	A.	6-7
v. <i>T. repens</i> , L.	<i>Dutch C.</i>	Past. . .	W.	3-18 in.	P.	5-9
[vi. <i>T. hybridum</i> , L. . . .	. . . . .	. . . . .	. .	. .	. .	. .
v. 2. <i>elegans</i> , Savi. . . .	. . . . .	. . . . .	. .	. .	. .	. .
vii. <i>T. fragiferum</i> , L. . .	<i>Strawberry T.</i> .	Fields. .	Ro.	6-12 in.	P.	7-8
<i>B.</i>						
*						
viii. <i>T. arvense</i> , L. . . .	<i>Hare's-foot T.</i> . .	Dry pas.	✓ Pk.	6-12 in.	A.	7-9
ix. <i>T. Bocconi</i> , Savi. . .	<i>Boccone's T.</i> . . .	Dry pl.! Corn.	W.	2-4 in.	A.	7-8
[x. <i>T. incarnatum</i> , L. . .	<i>crimson C.</i>	Fie. . . .	Sc.	6-18 in.	A.	6-7
v. 2. <i>Molinerii</i> , Balb. . .	. . . . .	. . . . .	✓ Y.	3-6 in.	. .	. .
**						
xi. <i>T. ochroleucum</i> , L. .	<i>sulphur-coloured T.</i>	Dry pas. E. of Eng.!	Y.	6-18 in.	P.	6-8
xii. <i>T. pratense</i> , L. . . .	<i>common purple C.</i>	Mead. .	Pu.	6-18 in.	P.	5-9
xiii. <i>T. medium</i> , L. . . .	<i>zigzag T.</i>	Mead. .	Pu.	6-18 in.	P.	5-9
xiv. <i>T. maritimum</i> , Huds.	<i>Teasel-headed T.</i>	Salt mar.!	✓ R.	6-18 in.	A.	6-7
xv. <i>T. striatum</i> , L. . . .	<i>soft-knotted T.</i> . .	Dry pas.	Ro.	4-12 in.	A.	6-7
xvi. <i>T. scabrum</i> , L. . . .	<i>rough-rigid T.</i> . .	San. fie.!	W.	4-10 in.	A.	5-7

CHARACTERISTICS OF SPECIES AND VARIETIES.

v. 2. *Spines of pod short, straight.*

vi. Downy—Pods compactly spiral, glob., spinous; the four spires narrowly keeled—Stip. half cord., almost ent.

Gen. VII.

i. Fls. yell.—Pet. all equal—Pods acute, hairy.

ii. Fls. yell.—Wings and standard longer than keel—Pods ribbed, obt., glab.]

iii. Fls. W.—Standard longer than keel and wings—Pods acute, glab.

Gen. VIII.

i. St. decumb., glab.—Lfts. ob.-cord., toothed—Pedunc. axill., three-flowered.

Gen. IX.

A. Throat of the cal. naked—Fls. white or red.

i. St. most underground—Heads 2-5 fld. hairy, stalked in fr., with abortive palm calices which fold over fr.

ii. St. procumb.—Heads glob., sess., term. and axill.—Cal. ten-veined, teeth ov., reflexed, glab., shorter than coroll. and contracted at base.

iii. St. ascending—Heads glob. term. and axill. shortly stalked—Cal. teeth subulate spread.—Lfts. lin.-lanc.

iv. St. underground—Heads dense, sess., axill.—Cal. membr., hairy, scarcely striate; its teeth lanc., recurved, longer than cor.

v. St. creeping—Heads glob., axill., long-stalked, umbell. at length deflex.—Pods four-seeded.

vi. Similar, but has an erect st. and a two-seeded pod (not native).

v. 2. *St. weak, decumb.—Heads smaller.]*

vii. St. creeping—Heads axill., long-stalked—Cal. much infl. after flowering.

B. Throat of calyx hairy.

* Heads cylindric in flower or seed.

viii. St. branched, sub-erect—Heads term., stalked, very hairy—Cal. teeth plumose, longer than cor.

ix. St. erect—Heads term. and axill., sess., usually in pairs—Cal. teeth erect, as long as cor.

x. St. erect—Heads sol., term., stalked—Cal. teeth shorter than cor.—Fls. scarlet.]

v. 2. *Fls. pale yellow (Lizard and Kynance Cove, Corn.; native).*

** Heads ovate.

xi. St. erect—Heads term., sol., stalked—Lower tooth of cal. very long—Fls. sulphur-coloured.

xii. St. sub-erect—Heads term., dense, sess.—Cal. hairy, lower tooth longer—Stip. ovate.

xiii. St. straggling, zigzag—Heads term., sol., stalked—Cal. glab., teeth hairy—Stip. lanc.

xiv. St. rigid, sub-erect—Heads term., shortly stalked—Cal. teeth rigid, subu., leafy in fr., lower one longer and three-nerved—Stip. subu., very long.

xv. St. procumb.—Heads term. and axill., sess.—Cal. furrowed, densely hairy, infla. in fr.; its teeth straight.

xvi. St. procumb.—Heads term. and axill., sess.—Cal. cylind. in fr.: its teeth unequal, rigid, recurved.

ORD. XXV.--LEGUMINOSÆ--cont.

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
xvii. <i>T. stellatum</i> , L. . . .	<i>Starry-headed T.</i>	Sea sh. Shoreham.	C.	4-12 in.	A.	6-8
<i>C.</i>						
xviii. <i>T. procumbens</i> , L. . .	<i>Hop T.</i>	Past. . .	Y.	6-18 in.	A.	6-8
xix. <i>T. minus</i> , Sm.	<i>lesser yell. T.</i> . . .	Past. . .	Y.	10-20 in.	A.	6-8
xx. <i>T. filiforme</i> , L.	<i>slender yell. T.</i> . .	Dry pas.!	Y.	4-8 in.	A.	6-7
Gen. X. <i>Lotus</i> , L.	<i>Bird's-foot Trefoil.</i>					
i. <i>L. corniculatus</i> , L. . .	<i>common B.</i>	Dry pl. .	Y.	4-18 in.	P.	7-8
v. 2. <i>crassifolius</i>	. . . . .	. . . . .	. . .	. . .	. . .	. . .
v. 3. <i>villosus</i>	. . . . .	. . . . .	. . .	. . .	. . .	. . .
v. 4. <i>tenuis</i> , Kit.	. . . . .	. . . . .	. . .	. . .	. . .	. . .
ii. <i>L. major</i> , Scop.	<i>narrow-leaved B.</i>	Moi. med.	Y.	6-24 in.	P.	7-8
iii. <i>L. angustissimus</i> , L. .	<i>slender B.</i>	Dry banks S. of Eng.!	Y.	2-6 in.	A.	6-8
<i>α. diffusus</i> , Sm.	. . . . .	. . . . .	. . .	. . .	. . .	. . .
<i>β. hispidus</i> , Desf.	. . . . .	. . . . .	. . .	. . .	. . .	. . .
Gen. XI. <i>Astragalus</i> , L.	<i>Milk-vetch.</i>					
i. <i>A. glycyphyllos</i> , L. . . .	<i>sweet M.</i>	Thick. .	<i>ρ</i> Y.	2-3 ft.	P.	6-9
ii. <i>A. hypoglottis</i> , L. . . .	<i>purple Mount M.</i>	Cha. pl. E. & N. of Eng.!	Pu.	2-6 in.	P.	6-7
iii. <i>A. alpinus</i> , L.	<i>Alpine M.</i>	Sc.mount.!	W.	2-4 in.	P.	7
Gen. XII. <i>Oxytropis</i> , L.	<i>Oxytropis.</i>					
i. <i>O. Uralensis</i> , DC. . . .	<i>hairy Mount O.</i> . .	Sc.mount.!	Pu.	4-6 in.	P.	6-7
ii. <i>O. campestris</i> , DC. . .	<i>yell Mount O.</i> . . .	Sc.mount.!	Y.	4-8 in.	P.	7-8
Gen. XIII. <i>Ornithopus</i> , L.	<i>Bird's-foot.</i>					
i. <i>O. perpusillus</i> , L. . . .	<i>common B.</i>	Dry pl. .	<i>ρ</i> R.	6-18 in.	A.	5-7
Gen. XIV. <i>Arthrolobium</i> , Desv.	<i>Joint-vetch.</i>					
i. <i>A. ebracteatum</i> , DC. . .	<i>Sand f.</i>	Sandy gro. C. Isl.!	Y.	6-18 in.	A.	6-8
Gen. XV. <i>Hippocrepis</i> , L.	<i>Horseshoe-vetch.</i>					
i. <i>H. comosa</i> , L.	<i>tufted H.</i>	Chal. pas.	Y.	6-18 in.	P.	5-8
Gen. XVI. <i>Onobrychis</i> , Tourn.	<i>Sain-foin.</i>					
i. <i>O. sativa</i> , Lam.	<i>common S.</i>	Chal. pas.	Ro.	1-2 ft.	P.	6-7
Gen. XVII. <i>Vicia</i> , L.	<i>Vetch, Tare.</i>					
*						
i. <i>V. Orobus</i> , DC.	<i>Wood bitter V.</i> . .	Roc. pl. N. of Eng.!	<i>pu</i> W.	1-2 ft.	P.	5-6

CHARACTERISTICS OF SPECIES AND VARIETIES.

- xvii. St. procumb.—Heads term., stalked—Cal. hairy ; its teeth long, leafy, spread in a stellate manner.]
C. Throat of calyx naked—Heads long-stalked—Fls. bright yellow.
 xviii. Heads large, dense, many-flowered—Standard furrowed when dry.
 xix. Heads 10-15 flowered on rigid pedunc.—Standard smooth, trunc.
 xx. Heads 2-6 flowered on capill. pedunc.—Standard deeply notched.

Gen. X.

- i. St. decumb.—Ls. obov., glab.—Cal. teeth erect in bud ; two upper converging—Claw of standard ob.-ov.
 v. 2. *St. tufted—Lfts. fleshy (sea-side).*
 v. 3. *Ls. and stem very hairy.*
 v. 4. *St. filiform—Lfts. lin.-lanc.—Fls. small.*
 ii. St. erect—Ls. ob.-ov.—Cal. teeth spreading in bud, two upper diverging—Claw of standard linear.
 iii. St. decumb.—Ls. hairy or silky, ov.-lanc.—Cal.-teeth erect in bud, all subu.
 a. Heads 1-2 flowered—Pods six times longer than cal.
 b. Heads 3-4 flowered—Pods rugose, twice as long as calyx.

Gen. XI.

- i. St. zigzag, glab.—Ls. longer than pedunc.—Fls. yell.—Pods lin., glab.
 ii. St. prost., hairy—Ls. shorter than pedunc.—Fls. purple—Pods ov., erect, hairy.
 iii. St. slender, ascend. pubesc.—Ls. as long as pedunc.—Fls. white, tipped purple—Pods obl., pend., hairy.

Gen. XII.

- i. Stemless—Ls. shorter than pedunc.—Fls. purple.
 ii. St. very short, prost.—Ls. equally pedunc.—Fls. yellow.

Gen. XIII.

- i. St. prost.—Ls. pinn.; lfts. ellip., downy—Fls. small white veined crims.

Gen. XIV.

- i. St. prost., fili.—Ls. pinn. ; lfts. ellip.-obl., glauc.—Fls. yell., veined crims.

Gen. XV.

- i. St. prost., glab.—Ls. pinn. ; lfts. ob.-ov., ellip.—Fls. umbell.

Gen. XVI.

- i. St. erect, stout—Lfts. ellip.-obl., mucro.—Fls. in dense rac.

Gen. XVII.

- * *Lfts. many—Pedunc. long, many-flowered—Calyx gibbous at base.*
 i. St. asc.—Lfts. ov.-obl., acute, hairy—Tend. o.—Stip. semi-sagitt., slightly toothed.

ORD. XXV.—LEGUMINOSÆ—*cont.*

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
ii. <i>V. sylvatica</i> , <i>L.</i> . . .	<i>Wood V.</i>	Roc. pl. .	W.	Cli.	P.	6-8
iii. <i>V. Cracca</i> , <i>L.</i> . . . . * *	<i>tufted V.</i>	Bushy pl.	B.	Cli.	P.	6-8
iv. <i>V. bithynica</i> , <i>L.</i> . . .	<i>rough-podded pur- ple V.</i>	Bushy pl.!	ϕ P.	1-2 ft.	P.	7-8
v. 2. <i>angustifolia</i> , <i>Syme.</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
v. <i>V. lathyroides</i> , <i>L.</i> . . . * * *	<i>Spring V.</i>	Dry pas.	Pu.	6-8 in.	A.	4-6
vi. <i>V. sativa</i> , <i>L.</i>	<i>common V.</i>	Fields. .	ϕ Pu.	6-18 in.	A.	5-6
v. 2. <i>angustifolia</i> , <i>Roth.</i> .	. . . . .	Dry pl. .	Pu.	4-6 in.	. . . . .	. . . . .
v. 3. <i>Bobartii</i> , <i>Forst.</i> . . .	. . . . .	Dry pl. .	Pu.	4-6 in.	. . . . .	. . . . .
vii. <i>V. lutea</i> , <i>L.</i>	<i>rough-podded yell. V.</i>	Sea side Rocks!	Y.	6-18 in.	A.	6-8
viii. <i>V. sepium</i> , <i>L.</i> . . . . * * * *	<i>Bush V.</i>	Hedges	ϕ Pu.	2-3 ft.	P.	5-8
ix. <i>V. hirsuta</i> , <i>Koch.</i> . . .	<i>hairy T.</i>	Was. pl. .	ϕ B.	1-2 ft.	A.	6-8
x. <i>V. tetrasperma</i> , <i>Moench.</i>	<i>slender T.</i>	Was. pl. .	ϕ B.	1-2 ft.	A.	6-8
v. 2. <i>gracilis</i> , <i>Lois.</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
Gen. XVIII. <i>Lathyrus</i> , <i>L.</i> *	<i>Everlasting Pea, Bitter Vetch.</i>					
i. <i>L. Aphaca</i> , <i>L.</i>	<i>yell. Vetch</i>	San. fie.!	Y.	1-3 ft.	A.	6-7
ii. <i>L. Nissolia</i> , <i>L.</i> . . . . * *	<i>crimson V., or Grass-Vetch.</i>	Bushy pl.	Cr.	1-2 ft.	A.	5-6
iii. <i>L. hirsutus</i> , <i>L.</i>	<i>rough-podded V.</i> .	Fields! .	Pu.	2-4 ft.	A.	6-7
iv. <i>L. pratensis</i> , <i>L.</i>	<i>Meadow V.</i>	Mead. .	Y.	1-2 ft.	P.	6-9
v. <i>L. sylvestris</i> , <i>L.</i>	<i>narrow-leaved E.</i>	Thick.!	g Pu.	3-6 ft.	P.	6-8
[vi. <i>L. tuberosus</i> , <i>L.</i> . . . . * * *	<i>Earth-nut Pea</i> . .	Corn fie.	Cr.	2-4 ft.	P.	6-8
vii. <i>L. palustris</i> , <i>L.</i>	<i>blue Marsh V.</i> . .	Bog. pl.!	ϕ Pu.	2-4 ft.	P.	6-8
viii. <i>L. maritimus</i> , <i>Bigel.</i> .	<i>Sea-side E.</i>	Sea sh. .	Pu.	1-3 ft.	P.	7-8
v. 2. <i>acutifolius</i> , <i>Bab.</i> . . . * * * *	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
ix. <i>L. macrorrhizus</i> , <i>Wim.</i>	<i>tuberous B.</i>	Thick. .	g Pu.	6-12 in.	P.	5-7
v. 2. <i>tenuifolius</i> , <i>Roth.</i> . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
x. <i>L. niger</i> , <i>L.</i>	<i>black B.</i>	Sco. rocs.	Pu.	1-2 ft.	P.	6-7

ORD. XXVI.—ROSACEÆ.

Gen. I. <i>Prunus</i> , <i>L.</i>	<i>Plum, Cherry, &c.</i>					
i. <i>P. communis</i> , <i>L.</i>	. . . . .	Copses & hedges.	W.	3-8 ft.	T.	3-4
a. <i>spinosa</i> , <i>L.</i>	<i>Sloe, Blackthorn</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
b. <i>institia</i> , <i>L.</i>	<i>Bullace</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .

CHARACTERISTICS OF SPECIES AND VARIETIES.

- ii. St. climb.—Lfts. ellip.-obl., mucro., glab.—tend. branched—Stip. lunate, deeply toothed.
- iii. St. climb.—Lfts. lanc., pubes.—tend. branched—Stip. semi-sagitt. ent.
* * *Lfts. few—Calyx equal at base.*
- iv. St. climb., glab.—Lfts. 1-2 pair, lanc.—Tend. branched—Pods rough.
- v. 2. *Lfts. lin. lanc.*
- v. St. spread, hairy—Lfts. 2-3 pair, retuse—Tend o.—Pods glab.
* * * *Lfts. many—Pedunc. short, few flowered—Calyx gibbous at base.*
- vi. St. trail—Lfts. ellip.-lanc., retuse—Fls. (1-2) axill., sess.—Stip. toothed, spotted—Pods spreading.
- v. 2. *St. slender—Lfts of upper ls. lin. lanc.—Fls. in pairs.*
- v. 3. *St. slender, prost.—Lfts of upper ls. lin.—Fls. sol.*
- vii. St. tufted, prost.—Lfts. ellip.-lanc.—Fls. sol., sess., cal. teeth unequal—Pods reflexed, hairy.
- viii. St. climb.—Fls. in sess. racemes—Pods erect, glab.
* * * * *Lfts. many—Pedunc. long, few-flowered.*
- ix. Lfts. 6-8 pair—Pedunc. 1-6 flowered—Pods two seeded, hairy.
- x. Lfts. 3-6 pair—Pedunc. 1-2 flowered as long as ls.—Pods four-seeded glab.
- v. 2. *Lfts. 3-4 pair acute—Pedunc. 1-4 flowered, longer than ls.—Pods 5-8 seeded.*

Gen. XVIII.

- * *Lfts. none.*
- i. St. climb., tendrilled—Stip. large, leafy, cord-sagitt.
- ii. St. erect, very slender—Petio. grass like.
* * *Lfts. one pair.*
- iii. St. winged—Lfts. lin.-lanc.—Pedunc. two-flowered—Pods hairy (Surrey and Essex).
- iv. St. angled—Lfts. lanc.—Pedunc. many-flowered—Stip. large sagitt.
- v. St. climb., winged—Lfts. large, ensi.—Pedunc. many-flowered—Stip. semi-sagitt. narrow.
(*L. Latifolius with ellip. lfts. and broader stip. is the Everlasting Pea of gardens, and escaped from cultivation.*)
- vi. St. angled—Lfts. ob.-ov.—Pedunc. 2-5 flowered, very long—Stip. large semi-sagitt.—Pods glab. (Ongar, Essex.)
* * * *Lfts. many—Ls. tendrilled.*
- vii. St. winged—Lfts. lin. lanc., acute—Stip. semi-sagitt., lanc.
- viii. St. angled—Lfts. oval—Stip. oval, cord.-hast., large.
- v. 2. *St. slend., straggling—Lfts. ellip.-lanc. (Burra Firth, Unst, Shetland).*
* * * * *Lfts. many—Ls. not tendrilled.*
- ix. St. winged, simp.—Lfts. 2-4 pair ellip.-obl., glab.—Stip. semi-sagitt.
- v. 2. *Lfts. lin. lanc.*
- x. St. angled, branched—Lfts. 3-6 pair, lanc.—Stip. lin. lanc., acute.

ORD. XXVI.—ROSACEÆ.

Gen. I.

- i. Fls. one or two on short pedi.—Ls. ellip.
- a. Pedi. sol., glab.—Ls. glab.—Branches spin.—Fr. glob., erect.
- b. Pedi. in pairs, downy—Ls. downy beneath—Branches slightly spin.—Fr. glob., drooping.

ORD. XXVI.—ROSACEÆ—*cont.*

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
<i>c. domestica</i> , <i>L.</i>	<i>wild Plum</i>	. . . .	. .	. .	. .	. .
ii. <i>P. Padus</i> , <i>L.</i>	<i>Bird Cherry</i> . . .	Woods .	W.	10-15ft.	T.	5-6
iii. <i>P. cerasus</i> , <i>L.</i>	<i>Cherry</i>	Woods .	W.	3-8 ft.	Shb.	5-6
iv. <i>P. avium</i> , <i>L.</i>	<i>wild C. or Gean.</i>	Woods .	W.	10-20ft.	T.	5-6
Gen. II. Spiræa , <i>L.</i>	<i>Dropwort, Mea- dow-Sweet.</i>					
i. <i>S. ulmaria</i> , <i>L.</i>	<i>Meadow-Sweet</i> .	Mead. &c.	W.	3 ft.	P.	6-8
ii. <i>S. filipendula</i> , <i>L.</i> . .	<i>Dropwort</i>	Downs .	W.	12-18in.	P.	6-7
[iii. <i>S. salicifolia</i> , <i>L.</i> . . .	<i>Willow-leaved Spiræa.</i>	Dp. wds.!	Pk.	3-4 ft.	P.	6-7
Gen. III. Geum , <i>L.</i>	<i>Avens.</i>					
i. <i>G. urbanum</i> , <i>L.</i>	<i>Wood A.</i>	Hedgebks.	Y.	1-2 ft.	P.	6-8
ii. <i>G. rivale</i> , <i>L.</i>	<i>Water A.</i>	Dp. woods	Y.	1-2 ft.	P.	6-7
Gen. IV. Dryas , <i>L.</i>	<i>Dryas.</i>					
i. <i>D. octopetala</i> , <i>L.</i> . .	<i>white D. or Moun- tain Avens.</i>	Mount.!	W.	1-3 in.	P.	6-7
Gen. V. Sibbaldia , <i>L.</i>	<i>Sibbaldia.</i>					
i. <i>S. procumbens</i> , <i>L.</i> . .	<i>procumbent S.</i> . .	Sc. moun.!	Y.	3-5 in.	P.	7
Gen. VI. Potentilla , <i>L.</i>	<i>Cinque-foil.</i>					
*						
i. <i>P. fruticosa</i> , <i>L.</i>	<i>shrubby C.</i>	Rocky pl.!	Y.	2-4 ft.	Shb.	6-7
ii. <i>P. rupestris</i> , <i>L.</i>	<i>strawberry-leaved C.</i>	Rocks ! .	W.	1-2 ft.	P.	5-6
iii. <i>P. anserina</i> , <i>L.</i> . . .	<i>Silver-weed</i>	Roadsides	Y.	6-12 in.	P.	6-7
* *						
iv. <i>P. argentea</i> , <i>L.</i>	<i>hoary C.</i>	Roadsides	Y.	6-18 in.	P.	6-7
v. <i>P. verna</i> , <i>L.</i>	<i>Spring C.</i>	Hil. pas.	Y.	3-6 in.	P.	4-6
vi. <i>P. alpestris</i> , <i>Hall, fil.</i>	<i>Alpine C.</i>	Mount. .	Or.	4-10 in.	P.	6-7
vii. <i>P. reptans</i> , <i>L.</i>	<i>creeping C.</i>	Roadsides	Y.	6-18 in.	P.	6-9
viii. <i>P. tormentilla</i> , <i>Sipt.</i> .	<i>Tormentil</i>	Heaths .	Y.	6-10 in.	P.	6-8
<i>a. officinalis</i>	. . . . .	. . . .	. .	. .	. .	. .
<i>b. procumbens</i> , <i>Sipt.</i> . .	. . . . .	. . . .	. .	. .	. .	. .
ix. <i>P. fragariastrum</i> , <i>Ehrh.</i>	<i>Strawberry-leaved C.</i>	Banks .	W.	1-6 in.	P.	3-5
Gen. VII. Comarum , <i>L.</i>	<i>Marsh Cinque-foil.</i>					
i. <i>C. palustre</i> , <i>L.</i>	<i>purple M.</i>	Bogs . .	pu B.	6-18 in.	P.	5-7

CHARACTERISTICS OF SPECIES AND VARIETIES.

- c. Pedi. sol. or in pairs, glab.—Ls. downy on the ribs beneath—Branches unarmed—Fr. obl.
- ii. Fls. in a pend. raceme—Ls. ob.-ov., doubly serr., glab.
- iii. Fls. in almost sess. umbels—Ls. ov.-obl., glab., firm, erect—Calyx tube not contracted.
- iv. Fls. as last—Ls. flaccid, drooping, downy beneath—Calyx tube contracted at mouth.

Gen. II.

- i. Ls. pinn., downy beneath: lfts. ent., term one large, lobed.
- ii. Ls. pinn., mostly rad.; lfts. deeply cut, serr., glab.
- iii. Shrubby—Ls. obl.-lanc., serr., glab.—Rac. term.]

Gen. III.

- i. Fls. erect—Heds of fr. sess.; upper joint of awn. short glab.
- ii. Fls. drooping—Heds of fr. stalked; upper joint of awn. long hairy.
(*G. Intermedium, probably a hybrid, partakes of the characters of both—It has the fls. erect or drooping and the upper joint of the awn. hairy with a glabrous tip.*)

Gen. IV.

- i. St. procumb.—Ls. simp., obt., serr.—Pet. eight.

Gen. V.

- i. St. woody—Ls. tern.; lfts. wedge-shaped, three-toothed.

Gen. VI.

* *Ls. pinnate.*

- i. St. shrubby—Lfts. five, ent., margins revolute.
- ii. St. erect, dichot—Lfts. cuneate, serr.—Fls. white.
- iii. St. creeping—Lfts. deeply serr., silky beneath—Fls. sol.
* * *Ls. digitate—Fls. mostly yellow.*
- iv. St. erect—Ls. quin.; lfts. cuneate, white beneath.
- v. St. prost., woody—Rad. ls. quin.; lfts. ob.-ov., trunc.—Lower stip. lin., acute.
- vi. St. ascend—Ls. as *P. verna*. but not trunc.—Fls. larger—Stip. ovate., obt.
- vii. St. prost., fili., rooting—Ls. quin., long-stalked—Fls. sol.; pet. five.—Ach. granulate.
- viii. St. prost., rarely rooting—Ls. tern., stem ls. sess.—Fls. cymose; pet. four—Ach. wrinkled.
- a. St. erect, not rooting at joints—Ls. alt., sess.—St. ls. tern.; fls. in a term. cyme.
- b. St. longer and procumb., rooting at joints—Ls. larger, stalked; st. ls. quin.—Fls. sol.

P. Mixta is probably a hybrid between Sp. 7 and 8.

- ix. St. procumb.—Ls. tern., lfts. hairy, serrate—Fls. white.

Gen. VII.

- i. St. erect—Ls. pinn.; lfts. lanc., serr.—Fls. dingy purple.

ORD. XXVI.—ROSACEÆ—*cont.*

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
Gen. VIII. <i>Fragaria</i>, L.	<i>Strawberry.</i>					
i. <i>F. vesca</i> , L.	<i>Wood S.</i>	Woods .	W.	1-6 in.	P.	5-7
[ii. <i>F. elatior</i> , Ehrh. . .	<i>Hautboy S.</i>	Woods .	W.	3-10 in.	P.	5-7
Gen. IX. <i>Rubus</i>, L.	<i>Blackberry, Rasp- berry, &c.</i>					
SEC. I.						
i. <i>R. chamæmorus</i> , L. .	<i>Cloudberry</i> . . .	Alpine turfy bogs.	W.	8-10 in.	P.	6-7
ii. <i>R. saxatilis</i> , L. . . .	<i>Stone B.</i>	Stony pl. N. of E.	g Y.	6-12 in.	P.	6-8
SEC. II.						
iii. <i>R. idæus</i> , L.	<i>Raspberry.</i> . . .	Woods .	W.	3-5 ft.	P.	6-8
b. <i>Leesii</i> , Bab.	. . . . .	. . . . .	. . .	. . .	. . .	. . .
SEC. III.						
iv. <i>R. fruticosus</i> , L. . .	<i>Blackberry</i> . . .	Woods, hedges, &c.	W. or Pk.	3-10 ft.	P.	7-8
A.						
i. <i>R. suberectus</i> , Anders.	. . . . .	Boggy woods.	W.	. . .	. . .	. . .
ii. <i>R. plicatus</i> , W. & N.	. . . . .	Heaths .	W.	. . .	. . .	. . .
iii. <i>R. fissus</i> , Lindl. . . .	. . . . .	Wet heaths.	W.	. . .	. . .	. . .
iv. <i>R. affinis</i> , W. & N. .	. . . . .	Heaths .	. . .	. . .	. . .	. . .
B.						
v. <i>R. Lindleianus</i> , Lees.	. . . . .	Thickets	W.	. . .	. . .	. . .
vi. <i>R. rhamnifolius</i> , W. & N.	. . . . .	Hedges .	W.	. . .	. . .	. . .
vii. <i>R. incurvatus</i> , Bab. .	. . . . .	Heaths .	Pk.	. . .	. . .	. . .
viii. <i>R. imbricatus</i> , Hort. .	. . . . .	Woods .	W.	. . .	. . .	. . .
ix. <i>R. latifolius</i> , Bab. . .	. . . . .	Woods .	W.	. . .	. . .	. . .
C.						
*						
x. <i>R. discolor</i> , W. & N.	. . . . .	Hedges .	Pk.	. . .	. . .	. . .
xi. <i>R. thyrsoides</i> , Wimm.	. . . . .	Hedges .	W.	. . .	. . .	. . .
**						
a.						
xii. <i>R. leucostachys</i> , Sm. .	. . . . .	Hedges .	Pk.	. . .	. . .	. . .
xiii. <i>R. Grabowskii</i> , Wei. .	. . . . .	Hedges .	. . .	. . .	. . .	. . .

CHARACTERISTICS OF SPECIES AND VARIETIES.

Gen. VIII.

- i. Hairs of the pedi. adpr.
- ii. Larger—Hairs of the pedi. spread.]

Gen. IX.SEC. I.—*Stem herbaceous.*

- i. St. erect—Ls. simp., lobed, plaited—Fls. sol., term.

- ii. St. prost. ; fl. shoots erect—Ls. tern.—Fls. in term. corymbs.

SEC. II.—*Stem shrubby with suckers.*

- iii. St. round, very prickly—Ls. pinn. ; lfts. ovate, white beneath.
- b. Ls. all tern., lfts. crowded—Prickles with bulbous bases.

SEC. III.—*Stem shrubby without suckers.*

- iv. St. prickly, mostly rooting—Ls. tern. or pinn. very variable—Fls. in term. panicles.

A. Suberecti.—*Stem nearly erect, glabrous, with equal prickles.*

- i. St. ang. ; prickles few, confined to the angles—Lfts. flexible, those of the fl. shoots narrowed below—Panicle thyrsoid—Fr.-cal. reflex.
- ii. St. bluntly ang. ; prickles strong, deflexed, confined to the angles—Lfts. coriaceous ; those of the fl.-shoots gibbous below—Pan. racemose, hairy—Fr.-cal. reflexed.
- iii. St. slightly ang. ; prickles many, slender, not confined to the angles—Lfts. coriaceous—Pan. corymbose—Fr.-cal. erecto-patent.
- iv. St. ang. ; prickles large, deflexed—Lfts. slightly wavy, dull green above, pilose beneath—Pan. broad, corymb. with ascending branches, lower ones widely spreading.

B. Nitidi.—*St. arched, rooting at end, slightly hairy, without setæ or stellate down—Prickles mostly on the angles of the stem.*

- v. St. smooth, shining ; prickles few, strong, declining—Lfts. stalked, green and shining above ; paler and pilose beneath—Pan. corymb., hairy, leafy, with divaricate branches.
- vi. St. ang., furrowed ; prickles strong—Lfts. coriaceous, flat, tomentose beneath—Pan. pyram., compact above with distant branches below.
- vii. St. ang., prickles strong, straight—Lfts. coriaceous, with incurved wavy margins, downy beneath—Pan. narrow ; branches short, patent.
- viii. St. decurved, furrowed, much branched ; prickles small, strong—Lfts. opaque above, term. one round, cord : basal. imbricate, convex—Pan. long narrow ; branches long, racemose, hairy at top.
- ix. St. decumb., furrowed ; prickles short, slender, declining—Lfts. broad, large, dull green above, paler beneath, coarsely serr. ; term. one cord-acum. ; basal. ones sess., imbrica.—Pan. short leafy below.

C. Villicaules.—*St. pilose, or stellately downy with few glands or setæ—Prickles on the angles of stem—Basal. lfts. stalked.** *Prickles strong equal—Hairs adpressed—Lfts. felted beneath.*

- x. St. furrowed, stellately downy—Lfts. coriaceous, slightly convex, closely tomentose beneath—Pan. long narrow ; pet. pink, styles purple.
- xi. St. furrowed, sub-glab.—Lfts. coriaceous, slightly concave, hairy and tomentose beneath—Pan. long, lax ; pet. white, styles green.

* * *Prickles moderate, equal—Hairs patent—Lfts. not felted beneath.**a. St. ang., hairy—Ls. grey or white beneath.*

- xii. St. with copious stellate down—Lfts. softly tomentose beneath—Pan. long, narrow, its prickles slender straight.
- xiii. St. nearly glab. ; prickles many—Lfts. opaque above, ashy beneath—Pan. long compound, prickly ; branches ascending, corymbose.

ORD. XXVI.—ROSACEÆ—*cont.*

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
xiv. <i>R. carpinifolius</i> , <i>W. & N.</i> <i>b.</i>	. . . .	Hills . .	Pk.	. .	. .	. .
xv. <i>R. Salteri</i> , <i>Bab.</i> .	. . . .	Hedges .	Pk.	. .	. .	. .
xvi. <i>R. villicaulis</i> , <i>W. & N.</i>	. . . .	Woods .	Pk.	. .	. .	. .
xvii. <i>R. macrophyllus</i> , <i>Weihe.</i> * * *	. . . .	Woods .	W.	. .	. .	. .
xviii. <i>R. mucronulatus</i> , <i>Bor.</i>	. . . .	Hedges .	<i>p</i> Pk.	. .	. .	. .
xix. <i>R. Sprengelii</i> , <i>Weihe.</i> <i>D.</i> *	. . . .	Heaths .	Pk.	. .	. .	. .
xx. <i>R. Bloxamii</i> , <i>Lees.</i>	. . . .	Woods .	W.	. .	. .	. .
xxi. <i>R. scaber</i> , <i>Weihe.</i>	. . . .	. . . .	. .	. .	. .	. .
xxii. <i>R. hystrix</i> , <i>Weihe.</i>	. . . .	Hedges .	Pk.	. .	. .	. .
xxiii. <i>R. rosaceus</i> , <i>Weihe.</i>	. . . .	. . . .	. .	. .	. .	. .
xxiv. <i>R. rudis</i> , <i>Weihe.</i> .	. . . .	Hedges .	W.	. .	. .	. .
xxv. <i>R. radula</i> , <i>Weihe.</i> * *	. . . .	Hedges .	Pk.	. .	. .	. .
xxvi. <i>R. Koehleri</i> , <i>Weihe.</i>	. . . .	Hedges .	P.	. .	. .	. .
xxvii. <i>R. fusco-ater</i> , <i>Weihe.</i>	. . . .	Hedges .	Pk.	. .	. .	. .
xxviii. <i>R. diversifolius</i> , <i>Lind.</i>	. . . .	Hedges .	W.	. .	. .	. .
xxix. <i>R. Lejeunii</i> , <i>Weihe.</i> * * *	. . . .	Hedges .	. .	. .	. .	. .
xxx. <i>R. pyramidalis</i> , <i>Bab.</i>	. . . .	Woods .	<i>g</i> W.	. .	. .	. .
xxxi. <i>R. Guntheri</i> , <i>Weihe.</i>	. . . .	Sha. pl. .	<i>p</i> Pk.	. .	. .	. .
xxxii. <i>R. humifusus</i> , <i>Weihe.</i>	. . . .	Woods .	W.	. .	. .	. .
xxxiii. <i>R. foliosus</i> , <i>Weihe.</i>	. . . .	Heaths .	W.	. .	. .	. .
xxxiv. <i>R. glandulosus</i> , <i>Bell.</i> <i>E.</i>	. . . .	Woods .	W.	. .	. .	. .
xxxv. <i>R. Balfourianus</i> , <i>Blox.</i>	. . . .	Hedges .	<i>p</i> Pk.	. .	. .	. .
xxxvi. <i>R. corylifolius</i> , <i>Sm.</i>	. . . .	Hedges .	W.	. .	. .	. .
xxxvii. <i>R. althæifolius</i> , <i>Host.</i>	. . . .	Hedges .	W.	. .	. .	. .

CHARACTERISTICS OF SPECIES AND VARIETIES.

- xiv. St. with sol. and clustered hairs, no stellate down—Lfts. hairy beneath—Pan. narrow, its prickles decurved.
b. St. ang.—Ls. green beneath.
- xv. St. furrowed, sub-glab.—Lfts. thin, rough beneath, coarsely and doubly serr.—Pan. long, lax, branches short.
- xvi. St. with spreading hairs—Lfts. thick, pale green shining, velvety, with reddish veins beneath.
- xvii. St. with scattered hairs—Lfts. thin, pilose beneath—Pan. elong., its branches ascending.
* * * Prickles unequal—St. round—Ls. green beneath.
- xviii. St. arched; prickles few, small, slender, straight—Lfts. rough and pilose on both sides, finely serrate.
- xix. St. prost.; prickles many, unequal, deflexed—Lfts. green above, pilose beneath.
D. Glandulosi.—Stem covered with many glands, setæ, and aciculi.
* Basal. lfts. stalked—Prickles passing suddenly into setæ and aciculi.
- xx. St. prost.—Lfts. green beneath, term. one broadly ob.-ovate, cusp.—Pan. long leafy to top.
- xxi. A form of *R. Bloxamii*, with larger prickles, and more lax panicle.
- xxii. St. arched, slightly furrowed—Lfts. green beneath—Term. one obl., ob.-ov. acum.—Pan. long, leafy, upper part leafless.
- xxiii. A prost. form of *R. hystrix*, with a coral red stem, and the prickles more unequal.
- xxiv. St. arched, ang.—Lfts. whitish green beneath, coarsely, doubly serr.—Sep. leaf-pointed, reflexed in fruit.
- xxv. St. arched, ang.—Lfts. white beneath, unequally and finely serr.—Sep. not leaf-pointed, reflexed in fruit.
* * Ls. quinate—Prickles passing gradually into setæ and aciculi.
- xxvi. St. trailing, terete; prickles many, irregular—Lfts. glab. above, pale beneath; basal. ones stalked—Pan. open, broad topped, with spreading branches.
- xxvii. St. decumb., ang.—Prickles many, unequal, declining—Lfts. rugose above, green beneath, term. one cord, ov.; basal. ones imbric.—Pan. long, narrow.
- xxviii. St. prost., ang.; prickles unequal, slender—Lfts. broad, rugose above, pale, pilose beneath—Pan. long, leafy above.
- xxix. St. prost., ang.; prickles unequal, small—Lfts. narrow, dull; paler beneath, with silky hairs on both sides—Pan. lax, axillary branches long.
* * * Ls. ternate, rarely quinate—Prickles passing abruptly into setæ, &c.
- xxx. St. decumb.; prickles many, short, strong—Lfts. with recurved edges—Pan. large, very pyramidal; base of sep. adpressed to fr.
- xxxi. St. decumb.; prickles weak, unequal—Lfts. large, flat, doubly and unequally serr.—Pan. long, leafy, rachis wavy—Fr.-sep. reflexed.
- xxxii. St. prost.; prickles weak, unequal—Lfts. doubly and finely dent.—Pan. racemose, naked at top; branches few, short, distant.
- xxxiii. A form of *R. Koehleri* with st. more ang.; term lft. cord-acum.—Pan. leafy to top.
- xxxiv. St. trailing; prickles slender, small, unequal glandular hairs many, equal—Pan. racemose, open; branches spreading at right angles.
E. Cæsii.—Stem prost. roundish without hairs, &c., but covered with glaucous bloom.
- xxxv. Prickles nearly equal—Lfts. green on both sides—Sep. ov. with a long point, erect in fr.
- xxxvi. Prickles nearly equal—Lfts. pale and soft beneath—Sep. ov. reflexed in fr.
- xxxvii. Prickles very unequal—Term lfts. round, cord. pointed—Sep. ov.-acum.

ORD. XXVI.—ROSACEÆ—*cont.*

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
xxxviii. <i>R. tuberculatus</i> ,	. . . .	Hedges .	Pk.	. .	. .	. .
xxxix. <i>R. cæsius</i> , <i>L.</i> . .	. . . .	Heaths .	W.	. .	. .	. .
Gen. X. <i>Rosa</i>, <i>L.</i>	<i>Rose</i> , <i>Sweet-briar</i> .					
<i>A.</i>						
* i. <i>R. spinosissima</i> , <i>L.</i> .	<i>Burnet-leaved R.</i>	Heaths .	p Pk.	1-3 ft.	Sh.	5-6
v. 2. <i>rubella</i>	. . . .	. . . .	. .	. .	. .	. .
ii. <i>R. Hibernica</i> , <i>Sm.</i> . .	<i>Irish R.</i>	Ireland .	Pk.	1-3 ft.	Sh.	5-7
v. 2. <i>glabra</i> , <i>Baker</i> . . .	. . . .	. . . .	. .	. .	. .	. .
v. 3. <i>cordifolia</i> , <i>Baker</i> . .	. . . .	. . . .	. .	. .	. .	. .
* *						
iii. <i>R. Sabini</i> , <i>Woods.</i> . .	<i>Sabine's R.</i>	N. of E..	Pk.	1-5 ft.	Sh.	5-7
v. 2. <i>involuta</i> , <i>Sm.</i> . . .	. . . .	. . . .	. .	. .	. .	. .
v. 3. <i>Robertsonii</i> , <i>Baker.</i>	. . . .	N. of E..	. .	. .	. .	. .
v. 4. <i>Wilsoni</i> , <i>Bor.</i> . . .	. . . .	Menai. .	. .	. .	. .	. .
<i>B.</i>						
*						
† v. <i>R. mollissima</i> , <i>Fr.</i> . .	<i>villous R.</i>	Hedges, N. of E.	Ro.	3-6 ft.	Sh.	6-7
vi. <i>R. tomentosa</i> , <i>Woods.</i>	<i>downy-leaved R.</i>	Hedges .	Ro.	3-6 ft.	Sh.	6-7
† † vii. <i>R. inodora</i> , <i>Fr.</i> . . .	<i>slightly-scented S.</i>	Thick., S. of E.	R.	3-6 ft.	Sh.	6-7
viii. <i>R. micrantha</i> , <i>Sm.</i> .	<i>small-flowered S.</i>	Heaths .	R.	3-6 ft.	Sh.	6-7
† † † ix. <i>R. rubiginosa</i> , <i>L.</i> . .	<i>true S.</i>	Ch. hths.	R.	3-6 ft.	Sh.	6-7
x. <i>R. sepium</i> , <i>Lindl.</i> . .	<i>small-leaved S.</i> .	Heaths !.	R.	3-6 ft.	Sh.	6-7
* *						
† xi. <i>R. canina</i> , <i>L.</i>	<i>common Dog R.</i>	Hedges .	Pk.	3-6 ft.	Sh.	6-7
† †						
xii. <i>R. systyla</i> , <i>Woods.</i> .	<i>close-styled R.</i> . .	Hedges .	Pk.	2-4 ft.	Sh.	6-7
xiii. <i>R. arvensis</i> , <i>Huds.</i> .	<i>trailing R.</i>	Hedges .	W.	2-6 ft.	Sh.	6-7
Gen. XI. <i>Agrimonia</i>, <i>L.</i>	<i>Agrimony.</i>					
i. <i>A. Eupatoria</i> , <i>L.</i> . .	<i>common A.</i>	Banks .	Y.	1½-2 ft.	P.	6-7
v. 2. <i>odorata</i> , <i>Mill.</i> . . .	<i>fragrant A.</i> . . .	S. of E. .	. .	. .	. .	. .

CHARACTERISTICS OF SPECIES AND VARIETIES.

xxxviii. A very doubtful plant which has the prickles with tubercular bases.

xxxix. Prickles very unequal—Term lfts. ovate or three-lobed—Sep. ov.-lanc.—Fr. glaucous.

Gen. X.

A. Shoots setigerous.

* *Ls. simply serrate.*

i. Lfts. small, glab. above—Prickles many, crowded, unequal—Sep. simple—Fr. glob., erect, glab., black.

v. 2. *Prickles few, more equal—Fr. red, obl., pendulous.*

ii. Lfts. larger, hairy beneath—Prickles few unequal, the larger curved—Sep. pinnfid, glab.—Fr. sub-glob., glab., red.

v. 2. *Peti. and ls. not hairy; pedunc. naked.*

v. 3. *Pedunc. glandular (Northumb.).*

* * *Ls. doubly serrate.*

iii. Lfts., densely hairy and slightly glandular beneath—Prickles scattered, unequal, straight—Sep. somewhat pinn.—Fr. glob., setose.

Vars. Doniana (Woods) and gracilis (Woods) are barely distinguishable.

v. 2. *St. dwarf—Lfts. hairy and gland. beneath; prickles crowded, unequal. straight—Sep. simple, almost as long as cor.—Fr. urceolate, glob., setose.*

v. 3. *Teeth of ls. sharper—Sep. pinn.—Old ls. glab. above.*

v. 4. *Tinged with purple—Ls. simply serr. hairy beneath—Fr. ov.-lanc.*

B. Shoots not setigerous.

* *Ls. glandulose.*

† *Prickles uniform, straight.*

v. Root shoots erect—Sep. persist, simp. connivent in fr.—Fr. scarlet, sub-glob., ripening early.

vi. Root shoots arched—Sep. falling before winter, very pinn., spreading in fr.—Fr. obl., ripening late.

(Several varieties. Subglobosa (Sm.)—Farinosa (Raw.)—Scabrinscula (Sm.)—Sylvestris (Woods)—Fundzilliana (Bak.)—Obovata (Baker), connect this and the last, and they are probably one species.)

† † *Prickles uniform, uncinatæ.*

vii. Ls. doubly serr., downy—Sep. doubly pinn., decid.—Fr. ellip.—Ramuli without setæ.

viii. Ls. doubly serr., downy—Sep. simply pinn.; seg. elongate, decid.—Fr. small, ovate—Ramuli with few setæ.

† † † *Prickles various—Larger uncinatæ, smaller subulatæ.*

ix. Lfts. doubly serr. rounded at base—Fr. pear-shaped—Sep. decid.

x. Lfts. doubly serr. acute at base—Fr. ov. rounded at base—Sep. persist.

* * *Ls. not glandulose.*

† *Styles distinct.*

xi. Lfts. naked, variable—Prickles equal, hooked—Sep. pinn. decid.

(The different forms of this most variable plant have been ably described and systematized by Mr. F. G. Baker, who enumerates some thirty varieties.)

† † *Styles united in a column.*

xii. Root shoots erect, arched—Prickles equal—Sty. in a conic head.

xiii. Root shoots trailing—Prickles unequal, those on the ramuli subul.—Sty. in a round head.

Gen. XI.

i. St. ls. pinn.; lfts. deeply serr.—Tube of cal. obconic, deeply furrowed in fr.—Lower spines spreading.

v. 2. *Tube of cal. bell-shaped, not furrowed in fr.; lower spines reflexed.*

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
Gen. XII. Sanguisorba, <i>L.</i>	<i>Burnet.</i>					
i. <i>S. officinalis, L.</i>	<i>great B.</i>	Dp. mead.	Pu.	1-2 ft.	P.	6-8
Gen. XIII. Poterium, L.	<i>Burnet.</i>					
i. <i>P. Sanguisorba, L.</i>	<i>lesser B.</i>	Cha. hths.	Pu.	6-18 in.	P.	6-8
ii. <i>P. muricatum, Spach.</i>	. . . .	Cha. hths.	Pu.	1-2 ft.	P.	6-8
Gen. XIV. Alchemilla, L.	<i>Lady's Mantle.</i>					
i. <i>A. vulgaris, L.</i>	<i>common L.</i>	Hilly past. N. of E.	y G.	6-18 in.	P.	6-8
v. 2. <i>montana, Willd.</i>	. . . .	. . . .	. . . .	. . . .	. . . .	. . . .
ii. <i>A. alpina, L.</i>	<i>Alpine L.</i>	Sc. mount.	y G.	3-9 in.	P.	6-7
v. 2. <i>conjuncta, Bab.</i>	. . . .	. . . .	. . . .	. . . .	. . . .	. . . .
iii. <i>A. arvensis, Scop.</i>	<i>Field L., or Parsley-Piert.</i>	Dry bks.	y G.	2-8 in.	A.	5-8
Gen. XV. Mespilus, L.	<i>Medlar.</i>					
i. <i>M. germanica, L.</i>	<i>common M.</i>	Hedges!	W.	. .	T.	5-6
Gen. XVI. Cratægus, L.	<i>Hawthorn, May.</i>					
i. <i>C. Oxyacantha, L.</i>	. . . .	Woods & hedges.	W.	. .	T.	6-7
v. 2. <i>monogyna, Jac.</i>	. . . .	. . . .	. .	. .	. .	. .
Gen. XVII. Cotoneaster, <i>Lindl.</i>	<i>Cotoneaster.</i>					
i. <i>C. vulgaris, Lindl.</i>	<i>common C.</i>	Ormes- head!	Pk.	1-2 ft.	Shb.	5-6
Gen. XVIII. Pyrus, L.	<i>Pear, Apple, Service, &c.</i>					
i. <i>P. communis, L.</i>	<i>Pear</i>	Woods .	W.	. .	T.	4-5
a. <i>Pyraster, Bor.</i>	. . . . .	. . . .	. .	. .	. .	. .
b. <i>Achras, Bor.</i>	. . . . .	. . . .	. .	. .	. .	. .
ii. <i>P. malus, L.</i>	<i>Crab p ple</i>	Hedges & Woods.	Pk.	. .	T.	5-6
a. <i>acerba, DC.</i>	. . . . .	. . . .	. .	. .	. .	. .
b. <i>mitis</i>	. . . . .	. . . .	. .	. .	. .	. .
iii. <i>P. torminalis, Sm.</i>	<i>wild S.</i>	Woods .	W.	. .	T.	4-5
iv. <i>P. Aria, Sm.</i>	<i>white Beam-tree</i> .	Woods .	W.	. .	T.	5
a. <i>aria, Sy.</i>	. . . . .	. . . .	. .	. .	. .	. .
b. <i>rupicola, Sy.</i>	. . . . .	. . . .	. .	. .	. .	. .
c. <i>scandica, Bab.</i>	. . . . .	. . . .	. .	. .	. .	. .
d. <i>fennica, L.</i>	. . . . .	. . . .	. .	. .	. .	. .
v. <i>P. aucuparia, Gärtn.</i>	<i>Mountain-Ash</i> .	Woods .	W.	. .	T.	5-6

CHARACTERISTICS OF SPECIES AND VARIETIES.

Gen. XII.

- i. Ls. pinn.; lfts. about thirteen, stalked, cord.—Spikes ov.-obl.; sta. four, not longer than cal.

Gen. XIII.

- i. St. ang.—Ls. pinn.; lfts. many, small, sess.—Sta. many, exserted; cal. of fr. quadrangular.
 ii. Larger—Cal. of fr. winged, pitted and reticulated.

Gen. XIV.

- i. Ls. ent., reni., plaited, lobed—Fls. in lax, term. clusters.
 v. 2. *Ls. and petioles very hairy.*
 ii. Ls. digitate, serr., silky beneath—Fls. in interrupted spikes.
 v. 2. *Lfts. joined to one-third of their length (Faroe Islands).*
 iii. Ls. trifid, lobes cut, pubescent—Fls. sess., axill.

Gen. XV.

- i. Thorny—Ls. lanc.—Fls. sol., sess., term.

Gen. XVI.

- i. Spinous—Ls. deeply pinnfid., obt., glab.—Fls. corymb.; cal. and pedi. glab.
 v. 2. *Cal. and pedi. pubesc. (the commonest form).*

Gen. XVII.

- i. Ls. ov., densely hairy beneath—Pedunc. downy.

Gen. XVIII.

- i. Ls. ellip., serr.—Fls. in corymbs—Sty. distinct—Fr. pyriform.
 a. Base of fr. obconic.
 b. Base of fr. rounded.
 ii. Ls. ov.-obl.—Fls. in sess. umbels—Sty. united—Fr. glab.
 a. Pedicels and undersides of ls. glab.—Fr. drooping.
 b. Pedicels and undersides of ls. toment.—Fr. erect.
 iii. Ls. ov., cord., glab., lobed; lobes triang., serr., lower ones spreading—Fls. corymb.—Fr. small, spotted.
 iv. Ls. ovate, serr., lobed, white and downy beneath—Fls. corymb.—Fr. glob., red.
 a. Ls. ov.; ent. below, serr. above—Lat. veins about twelve.
 b. Ls. ob.-ovate, most deeply lobed above the middle—Lat. veins six or eight.
 c. Ls. with broad toothed lobes, lobes deepest near the middle—Lat. veins 5-8.
 d. Ls. doubly serr. near apex, pinnfid below.
 v. Ls. pinn.; lfts. serr., downy beneath—Fls. corymb.—Fr. glab., scarlet.

ORD. XXVII.—ONAGRACEÆ.

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
Gen. I. Epilobium, L.	<i>Willow herb.</i>					
i. <i>E. angustifolium, L.</i>	<i>Rose bay W.</i> . . .	Copses .	Ro.	4-6 ft.	P.	7-8
v. 2. <i>brachycarpum, Leight.</i>	.. .. .	.. .. .	.. .. .	.. .. .	.. .. .	.. .. .
ii. <i>E. hirsutum, L.</i> . . .	<i>great hairy W.</i> . .	Wat. pl..	Ro.	4-5 ft.	P.	7-8
iii. <i>E. parviflorum, Schreb.</i>	<i>small-flowered W.</i>	Wat. pl..	Ro.	1-1½ ft.	P.	7-8
v. 2. <i>rivulare, Wahl.</i> . . .	.. .. .	.. .. .	.. .. .	.. .. .	.. .. .	.. .. .
v. 3. <i>intermedium, Merat.</i>	.. .. .	.. .. .	.. .. .	.. .. .	.. .. .	.. .. .
iv. <i>E. montanum, L.</i> . . .	<i>broad smooth-leaved W.</i>	Dry bks.	♂ Pu.	1-2 ft.	P.	6-7
v. <i>E. roseum, Schreb.</i> . . .	<i>pale smooth-leaved W.</i>	Copses .	Ro.	1-2 ft.	P.	7-8
vi. <i>E. lanceolatum, Seb.</i> . .	<i>spear-leaved W.</i> . .	Sto. pl.!	♂ Ro.	1-3 ft.	P.	7-9
vii. <i>E. palustre, L.</i>	<i>narrow-leaved W.</i>	Bogs . .	Ro.	6-18 in.	P.	7-8
v. 2. <i>ligulatum, Baker.</i> . . .	.. .. .	.. .. .	.. .. .	.. .. .	.. .. .	.. .. .
viii. <i>E. tetragonum, L.</i> . . .	<i>square-stalked W.</i>	Bogs . .	Ro.	1-2 ft.	P.	7-8
v. 2. <i>obscurum, Schreb.</i> . . .	.. .. .	.. .. .	.. .. .	.. .. .	.. .. .	.. .. .
ix. <i>E. alpinum, L.</i>	<i>Alpine W.</i>	Alp. rills	Ro.	2-4 in.	P.	7-8
v. 2. <i>anagallidifolium, Lam.</i>	.. .. .	.. .. .	.. .. .	.. .. .	.. .. .	.. .. .
x. <i>E. alsinifolium, Vill.</i> . .	<i>Chickweed-leaved W.</i>	Alp. rills	Ro.	4-12 in.	P.	7-8
Gen. II. Oenothera, L.	<i>Evening-primrose.</i>					
i. <i>O. biennis, L.</i>	<i>common E.</i>	San. sh. (Lpool.)	Y.	2-3 ft.	P.	7-9
Gen. III. Isnardia, L.	<i>Isnardia.</i>					
i. <i>I. palustris, L.</i>	<i>Isnardia</i>	Boggy pools.	G.	6-10 in.	P.	6-7
Gen. IV. Circea, L.	<i>Enchanter's Nightshade.</i>					
i. <i>C. lutetiana, L.</i>	<i>common E.</i>	Dp. wds.	♂ W.	1-2 ft.	P.	6-8
ii. <i>C. alpina, L.</i>	<i>Alpine E.</i>	Nor. wds.	♂ W.	1-2 ft.	P.	6-8

ORD. XXVIII.—HALORAGIACEÆ.

Gen. I. Hippuris, L.	<i>Mare's-tail.</i>					
i. <i>H. vulgaris, L.</i>	<i>common M.</i>	Ditches .	G.	6-18 in.	P.	6-7
Gen. II. Myriophyllum, L.	<i>Water Milfoil.</i>					
i. <i>M. verticillatum, L.</i> . . .	<i>whorled-W.</i>	Ponds. .	W.	flo.	P.	7-8
v. 2. <i>pectinatum, DC.</i>	.. .. .	.. .. .	.. .. .	.. .. .	.. .. .	.. .. .
ii. <i>M. spicatum, L.</i>	<i>spiked-W.</i>	Ponds. .	W.	flo.	P.	6-7
iii. <i>M. alternifolium, DC.</i>	<i>alternate-flowered W.</i>	Ponds. .	W.	flo.	P.	6-8
Gen. III. Callitriche, L.	<i>Water Star-wort.</i>					
i. <i>C. verna, L.</i>	<i>vernal W.</i>	Ditches .	.. ..	flo.	A.	4-9

ORD. XXVII.—ONAGRACEÆ.

CHARACTERISTICS OF SPECIES AND VARIETIES.

Gen. I.

- i. St. round, erect—Ls. ellip.-lanc., alt.—Fls. many, in a long rac.
- v. 2. *Fls. smaller—Caps. erect (a garden escape).*
- ii. St. branched, cylind.—Ls. obl.-lanc., opp., hairy, amplex—Rt. cr.
- iii. St. cylind., woolly, almost simple—Ls. lanc., sess., alt.—Rt. fibr.
- v. 2. *Almost glabrous.*
- v. 3. *All the ls. alt.*
- iv. St. slender, mostly simp.—Ls. ov., toothed, stalked, opp., glab.
- v. St. obscurely four-angled—Ls. ellip.-lanc., toothed, stalked, glab., opp.
- vi. St. obscurely angled, pubesc.—Ls. ob.-lanc., toothed, alt., stalked, finely pubesc.
- vii. St. round, terete—Ls. sess., lanc. with a wedge-shaped base.
- v. 2. *Ls. lanc. faintly toothed.*
- viii. St. square—Ls. lanc., sess., finely toothed, shining above.
- v. 2. *Ls. ov.-lanc., opp., not shining above.*
- ix. St. decumb. at base—Ls. ellip.-lanc., obt., ent., slightly hairy.
- v. 2. *Stolons longer—Fls. pale.*
- x. St. tufted, asc.—Ls. ovate, acute, stalked, toothed, glab.—Root creeping.

Gen. II.

- i. St. erect, hairy—Ls. ov.-lanc.—Fls. large, sess.; pet. ob-cord.

Gen. III.

- i. St. glab., square, procumb. or flo., rooting at joints—Ls. opp., ov., acute—Fls. minute, sess., axill.

Gen. IV.

- i. St. erect—Ls. ov., slightly toothed, downy.
- ii. Smaller—Ls. cord., dent., shining—(*Hybrids occur between these two*).

ORD. XXVIII.—HALORAGIACEÆ.

Gen. I.

- i. St. simple—Ls. lin., whorled—Fls. axill.

Gen. II.

- i. Fls. in axill. whorls with pect. bracts.—Ls. five in a whorl.
- v. 2. *Bracts scarcely longer than fls.*
- ii. Fls. in a leafless spike, erect in bud—Ls. four in a whorl—Bracts ent.
- iii. Fls. in a leafless spike, nodding in bud—Ls. three in a whorl—Bracts pect.

Gen. III.

- i. Bracts falc.—Lobes of fr. bluntly keeled at back.

ORD. XXVIII.—HALORAGIACEÆ—*cont.*

CHARACTERISTICS OF SPECIES AND VARIETIES.

- v. 2. *Lobes of fr. slightly winged at back.*
 ii. Bracts o.—Lobes of fr. parallel, bluntly keeled.
 a. Fr. sess.
 b. Fr. stalked—Ls. lin.
 iii. Bracts. o.—Lobes of fr. spreading acutely, winged at back.

Gen. IV.

- i. Fr. armed with two spines—Seg. of cal. notched at tip.
 v. 2. *Fr. smooth—Seg. of cal. ent.*

ORD. XXIX.—LYTHRACEÆ.

Gen. I.

- i. Ls. opp., lanc., cord—Fls. in whorled spikes.
 ii. Ls. alt., lin.-lanc.—Fls. sol., axill.

Gen. II.

- i. Ls. opp., ob.-ov.—Fls. sol., axill.; pet. often absent.

ORD. XXX.—TAMARISCACEÆ.

Gen. I.

- i. Ls. minute, acute, adpr.—Fls. in slender, lat., spikes.

ORD. XXXI.—CUCURBITACEÆ.

Gen. I.

- i. St. climbing—Ls. palm—Fls. in short axill., rac.

ORD. XXXII.—PORTULACACEÆ.

Gen. I.

- i. St. tufted, succulent—Ls. opp., ent., spath.—Fls. at first drooping.
 a. St. short, rigid—Ped. axill. and term.
 b. St. long, flaccid—Ped. all axill.

ORD. XXXIII.—ILLECEBRACEÆ.

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
Gen. I. Illecebrum, L.	<i>Knot-grass.</i>					
i. <i>I. verticillatum, L.</i> . . .	<i>whorled-K.</i> . . .	Bogs! S.W. of E.	W.	4-8 in.	P.	7-8
Gen. II. Corrigiola, L.	<i>Strap-wort.</i>					
i. <i>C. littoralis, L.</i> . . .	<i>Sand S.</i>	San. pl.!	W.	4-8 in.	A.	7-8
Gen. III. Herniaria, L.	<i>Rupture-wort.</i>					
i. <i>H. glabra, L.</i>	<i>Rupture-wort</i> . .	San. pl.!	G.	4-6 in.	P.	7-8
v. 2. <i>ciliata, Bab.</i>	. . . . .	. . . . .	. .	. .	. .	. .
Gen. IV. Polycarpon, L.	<i>Allseed.</i>					
i. <i>tetraphyllum, L.</i>	<i>four-leaved A.</i> .	San. sh.! S.W. of E.	g W.	3-6 in.	A.	6-7

ORD. XXXIV.—SCLERANTHACEÆ.

Gen. I. Scleranthus, L.	<i>Knawel.</i>					
i. <i>S. annuus, L.</i>	<i>annual K.</i>	Fie. . .	G.	2-8 in.	A.	6-9
v. 2. <i>biennis, Syme.</i>	. . . . .	. . . . .	. .	. .	. .	. .
ii. <i>S. perennis, L.</i>	<i>perennial K.</i> . .	San. fie., E. of E.	G.	2-8 in.	P.	6-9

ORD. XXXV.—GROSSULARIACEÆ.

Gen. I. Ribes, L.	<i>Currant, Goose- berry.</i>					
i. <i>R. nigrum, L.</i>	<i>black C.</i>	Woods .	y G.	3-5 ft.	Shb.	4-5
ii. <i>R. rubrum, L.</i>	<i>red C.</i>	Woods .	y G.	3-5 ft.	Shb.	4-5
a. <i>sylvestre.</i>	. . . . .	. . . . .	. .	. .	. .	. .
b. <i>petraeum, Sm.</i>	. . . . .	. . . . .	. .	. .	. .	. .
c. <i>spicatum, Robs.</i>	. . . . .	. . . . .	. .	. .	. .	. .
iii. <i>R. alpinum, L.</i>	<i>Mountain C.</i> . .	Woods, N. of E.	y G.	2-4 ft.	Shb.	4-5
iv. <i>R. grossularia, L.</i> . .	<i>wild C.</i>	Hedges .	G.	2-4 ft.	Shb.	4-5
a. <i>glandulosum</i>	. . . . .	. . . . .	. .	. .	. .	. .
b. <i>uva-crispa</i>	. . . . .	. . . . .	. .	. .	. .	. .

ORD. XXXVI.—CRASSULACEÆ.

Gen. I. Tillæa, L.	<i>Tillæa.</i>					
i. <i>T. muscosa, L.</i>	<i>Mossy T.</i>	San. hths.	W.	1-2 in.	A.	6-7

ORD. XXXIII.—ILLECEBRACEÆ.

CHARACTERISTICS OF SPECIES AND VARIETIES.

Gen. I.

- i. St. fili., prost.—Ls. ovate—Fls. small, in axill. whorls.

Gen. II.

- i. Sts. many, prost.—Ls. lin.-lanc.—Fls. in small clusters.

Gen. III.

- i. St. prost.—Ls. obl.—Fls. sess., in axill. clusters forming a spike.
 v. 2. *Ls. broader, orb., ciliate—Fls. in distinct clusters.*

Gen. IV.

- i. Lower ls. four in a whorl., ob.-ov. ; upper ones opp.—Fls. in crowded clusters.

ORD. XXXIV.—SCLERANTHACEÆ.

Gen. I.

- i. St. much branched—Seg. of cal. acute, spreading in fr.
 v. 2. *St. shorter, biennial—Ls. and fls. smaller.*
 ii. St. nearly simp.—Seg. of cal. obt., connivent in fr.

ORD. XXXV.—GROSSULARIACEÆ.

Gen. I.

- i. Cal. pubesc.—Rac. of fls. pend., lax, downy—Bracts subu., shorter than fls.
 ii. Cal. downy—Rac. of fls. (mostly pend.), glab.—Bracts ov. shorter than fls.
 a. Rac. of fls. drooping in fl. and in fr.
 b. Rac. erect in fl., pend. in fr.
 c. Rac. erect in fl. and fr.
 iii. Cal. glab.—Rac. erect in fl. and fr.—Bracts longer than fls.
 iv. St. spinous—Pedunc. 1-3 fld. short, hairy—Bracts minute.
 a. Fr. hairy.
 b. Fr. smooth.

ORD. XXXVI.—CRASSULACEÆ.

Gen. I.

- i. St. procumb., tufted—Ls. obl., opp.—Fls. sess., axill., three cleft.

ORD. XXXVI.—CRASSULACEÆ—cont.

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
Gen. II. Sedum, L.	<i>Stonecrop, Orpine.</i>					
* i. <i>S. Rhodiola, DC.</i> . . .	<i>Rose-root S.</i> . . .	Wet rocs.	Y.	6-12 in.	P.	6-7
ii. <i>S. Telephium, L.</i> . . .	<i>O., or Livelong.</i> . .	Waysides	Pu.	1-2 ft.	P.	7-8
a. <i>purpurascens, Koch.</i> . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
b. <i>fabaria, Koch.</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
* * A.						
iii. <i>S. villosum, L.</i>	<i>hairy S.</i>	Alp. bogs	puW.	3-4 in.	A.	6-7
iv. <i>S. dasphyllum, L.</i> . . .	<i>thick-leaved S.</i> . . .	Old walls, W. of E.!	ro W.	2-3 in.	P.	6-7
v. <i>S. anglicum, Huds.</i> . . .	<i>English S.</i>	Roc. pl. .	W.	2-3 in.	A.	6-8
vi. <i>S. album, L.</i>	<i>white S.</i>	Rocks & walls.	W.	3-5 in.	P.	7-8
a. <i>teretifolium, Haw.</i> . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
b. <i>micranthum, Bast.</i> . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
* vii. <i>S. acre, L.</i>	<i>biting S., or Wall- pepper.</i>	Walls & roofs.	Y.	3-8 in.	P.	6-7
[viii. <i>S. sexangulare, L.</i> . .	<i>tasteless Mountain S.</i>	Old walls!	Y.	3-6 in.	P.	7-8
[ix. <i>S. reflexum, L.</i>	<i>crooked yellow S.</i>	Walls & roofs.	Y.	6-8 in.	P.	7-8
v. 2. <i>albescens, Haw.</i> . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
x. <i>S. rupestre, Huds.</i> . . .	<i>St. Vincent's Rock S.</i>	Rocks! W. of E.	Y.	6-10 in.	P.	6-7
v. 2. <i>minus.</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
xi. <i>S. Forsterianum, Sm.</i>	<i>Welch Rock S.</i> . .	Rocks, W. of E.!	Y.	6-10 in.	P.	6-7
v. 2. <i>virescens</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
Gen. III. Sempervivum, L.	<i>House-leek.</i>					
i. <i>S. tectorum, L.</i>	<i>common H.</i>	Roofs . .	Pu.	1-2 ft.	P.	7-8
Gen. IV. Cotyledon, L.	<i>Penny-wort.</i>					
i. <i>C. umbilicus, L.</i>	<i>Wall P.</i>	Rocks . .	y G.	6-12 in.	P.	6-8
v. 2. <i>foliosa</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .

ORD. XXXVII.—SAXIFRAGACEÆ.

Gen. I. Saxifraga, L.	<i>Saxifrage.</i>					
i. <i>S. Geum, L.</i>	<i>Kidney-shaped S.</i>	Ir. mount.	W.	6-12 in.	P.	6-7
a. <i>crenata</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
b. <i>serrata</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
c. <i>elegans, Mack.</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
d. <i>gracilis, Mack.</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
ii. <i>S. hirsuta, L.</i>	. . . . .	Ir. mount.	Y.	6-12 in.	P.	6-7

CHARACTERISTICS OF SPECIES AND VARIETIES.

Gen. II.

* *Ls. flat.*

- i. Root woody—Ls. ob.-ov., fleshy, glaucous—Fls. yell. four partite.
- ii. Ls. oval, obl., serr.—Fls. purp. five partite.
  - a. Upper ls. rounded at base.
  - b. Upper ls. cuneate at base.

* * *Ls. subterete.**A. Flowers white or purplish.*

- iii. St. erect—Ls. lin., alt., flat above—St. and ls. gland.-pubesc.
- iv. Flowering st. weak, procumb.—Ls. opp., ovate, fleshy, glaucous.
  - v. Flowering st. ascending—Ls. alt., gibb., ovate, glab.—Pan. bifid—Pet. very pointed.
  - vi. Flowering st. erect—Ls. lin., obt.—Pan. much branched—Pet. lanc.
    - a. Ls. flattened above—Pet. obt.
    - b. Ls. flattened above and below—Pet. acute.
- * *Flowers yellow.*
- vii. Ls. alt., gibb., ov., imbricate on the barren shoots—Pan. trifid. leafy—Sep. obt., glab.
- viii. Ls. lin., obt., spreading—Sep. acute not gibbous.]
- ix. Ls. subu., cylind., lower ones recurved—Pet. obt. ; sep. ov., acute.]
- v. 2. *Lower ls. glauc. not recurved.*
- x. Ls. lin.-lanc., flattened, glauc. ; those of the barren shoots adpr.—Fls. in a flat-topped cyme—Sep. obl., obt. (Cheddar).
- v. 2. *Smaller and more slender (W. of E.).*
- xi. Ls. lanc., not glauc. ; those of the barren shoots spreading—Fls. in a compact, round-topped cyme—Sep. obt.
- v. 2. *More slender.*

Gen. III.

- i. Ls. ov.-lan., cili., in a rose-like tuft—Offsets spreading.

Gen. IV.

- i. Ls. pelt., cren.—St. simple with rac. of pend. fls.—Bracts. ent.
- v. 2. *Stem leafy.*

ORD. XXXVII.—SAXIFRAGACEÆ.

Gen. I.

- i. Ls. orb., reni., toothed, hairy—St. slender, peti. channelled.
  - a. Ls. bluntly toothed ; base deeply cord.
  - b. Ls. less cord., dent.-serr.
  - c. Ls. acutely toothed, sharply narrowed into the peti.
  - d. Ls. orb., hairy on both sides.
- ii. Ls. oval heart-shaped, blunt, with cartilaginous notches, sharply toothed, very hairy.

ORD. XXXVII.—SAXIFRAGACEÆ—*cont.*

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
iii. <i>S. umbrosa</i> , <i>L.</i> . . .	<i>London Pride, or None-so-pretty</i>	Ir. mount.	W.	6-12 in.	P.	6-7
<i>a. genuina</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
<i>b. punctata</i> , <i>Haw.</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
<i>c. serratifolia</i> , <i>Mack.</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
<i>d. duplicato-serrata</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
iv. <i>S. stellaris</i> , <i>L.</i>	<i>starry S.</i>	Alp. rocks	W.	3-8 in.	P.	7-8
v. <i>S. nivalis</i> , <i>L.</i>	<i>Alpine-clustered S.</i>	Alp. rocks	W.	3-6 in.	P.	7-8
vi. <i>S. Hirculus</i> , <i>L.</i>	<i>y. marsh S.</i>	Wet moors!	Y.	4-8 in.	P.	8
vii. <i>S. aizoides</i> , <i>L.</i>	<i>y. mount S.</i>	Alp. rills	Y.	5-7 in.	P.	6-9
viii. <i>S. oppositifolia</i> , <i>L.</i>	<i>pu. mount S.</i>	Alp. roc.!	Pu.	6-8 in.	P.	4-5
ix. <i>S. granulata</i> , <i>L.</i>	<i>W. Mead S.</i>	Mead. . .	W.	6-8 in.	P.	5-6
x. <i>S. ceruna</i> , <i>L.</i>	<i>drooping bulbous S.</i>	Ben Lawers	W.	3-5 in.	P.	8
xi. <i>S. rivularis</i> , <i>L.</i>	<i>Alp. brook S.</i>	Alp. roc.	W.	1-4 in.	P.	8
xii. <i>S. tridactylites</i> , <i>L.</i>	<i>Rue-leaved S.</i>	Walls . .	W.	2-4 in.	A.	4-7
xiii. <i>S. hypnoides</i> , <i>L.</i>	<i>mossy S.</i>	Rocks. . .	W.	3-12 in.	P.	5-7
<i>a. gemmifera</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
<i>b. platypetala</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
<i>c. decipiens</i> , <i>Ehrh.</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
<i>d. hirta</i> , <i>Don.</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
xiv. <i>S. cæspitosa</i> , <i>L.</i>	<i>tufted-Alp. S.</i>	Sc. mnt.!	W.	2-4 in.	P.	5-7
Gen. II. <i>Chrysosplenium</i> , <i>L.</i>	<i>Golden Saxifrage.</i>					
i. <i>C. oppositifolium</i> , <i>L.</i>	<i>common G.</i>	Mar. . .	Y.	2-4 in.	P.	4-7
ii. <i>C. alternifolium</i> , <i>L.</i>	<i>alternate-leaved G.</i>	Mar. . .	Y.	2-4 in.	P.	4-7
Gen. III. <i>Parnassia</i> , <i>L.</i>	<i>Grass of Pär- nassus.</i>					
i. <i>P. palustris</i> , <i>L.</i>	<i>G. of P.</i>	Bogs . .	W.	1-8 in.	P.	8-9

ORD. XXXVIII.—ARALIACEÆ.

Gen. I. <i>Adoxa</i> , <i>L.</i>	<i>Moschatel.</i>					
i. <i>A. Moschatellina</i> , <i>L.</i>	<i>tuberous M.</i>	Sha. pl. .	y G.	4-6 in.	P.	4-5
Gen. II. <i>Hedera</i> , <i>L.</i>	<i>Ivy.</i>					
i. <i>H. Helix</i> , <i>L.</i>	<i>common I.</i>	Walls . .	p G.	Climb.	Shb.	10-11

ORD. XXXVII.—SAXIFRAGACEÆ—*cont.*

CHARACTERISTICS OF SPECIES AND VARIETIES.

- iii. Ls. ob.-ovate, blunt, suddenly narrowed into a broad flat peti.
- a. Ls. obl.-orb., cren., in a compact rosette.
 b. Ls. orb., serr., in a loose rosette—Peti. longer.
 c. Ls. ob.-ov. sharply-toothed, erect.
 d. Ls. doubly serrate.
- iv. Ls. cun.-lan., sess., in a rosette—Cyme lax.
 v. Ls. ob.-ov., toothed, shortly stalked—Cyme dense.
 * * *Ls. entire—Stems leafy.*
- vi. St. erect—Ls. lanc., alt., ent.; rad. ls. lanc. or spath., rosulate—Sep. reflex.
- vii. St. decumb.—St. ls. lin., alt., cili., crowded at base—Sep. conn., erect.
- viii. St. tufted—Ls. opp., ovate, obt. imbri.; upper ones cili.
 * * * *Ls. divided.*
- ix. St. erect, paniced; bulbiferous at base—Ls. reni., lobed.
- x. St. erect, bulbiferous in axils—Ls. palmately lobed.
- xi. St. decumb.—Ls. palm., lobed, long-stalked—Cal. lobes acute.
- xii. St. erect—Ls. alt., cun., three or five fid., upper ent.—Cal. lobes obt.
- xiii. Flowering st. erect.; sterile st. decumb.—Rad. ls. three cleft; upper ones three cleft or ent.—Cal. lobes acute (a very variable plant).
 a. Ls. of sterile shoots undivided, with axill. buds.
 b. Ls. of sterile shoots three cleft, without axill. buds.
(These two varieties are connected by various forms, as leptophylla (Haw.) elongella (Sin.), condensata (Gmel.), denudata (Sin.), pygmæa (Don).)
 c. Ls. all rosulate; 3-5 fid.: cal. lobes obt.
 d. Sterile shoots, short—Ls. trifid; lobes acute—Pet. flat—Sep. triang.-ov.—Very hairy.
Of this there are two other doubtful varieties, affinis (Don), and incurvifolia (Don).
- xiv. Rad. ls. densely tufted, palm., hairy, 3-5 cleft; lobes obt.—Sep. obt.—St. 2-3 flowered.

Gen. II.

- i. St. branched, decumb.—Ls. opp., lower shortly stalked.
 ii. St. simp., erect—Ls. alt., lower long stalked.

Gen. III.

- i. Rad. ls. ov., cord., long-stalked—St.-leaf sess.—Fls. sol., term.

ORD. XXXVIII.—ARALIACEÆ.

Gen. I.

- i. Rad. ls. tri.-tern. long-stalked; st. ls. tern.—Fls. in a term. five-flowered head.

Gen. II.

- i. Ls. five-lobed, cord., variable—Fls. umbell.

ORD. XXXIX.—CORNACEÆ.

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
Gen. I. Cornus, L.	<i>Cornel, Dogwood.</i>					
i. <i>C. sanguinea, L.</i>	<i>Wild C. or D.</i>	Woods .	W.	5-6 ft.	Shb.	6-7
ii. <i>C. Suecica, L.</i>	<i>dwarf C.</i>	Alp. pas.	pkW.	6-8 in.	P.	7-8

ORD. XL.—UMBELLIFERÆ.

Gen. I. Hydrocotyle, L.	<i>White-rot.</i>					
i. <i>H. vulgaris, L.</i>	<i>White-rot, or Marsh penny-wort.</i>	Bogs . .	W.	Cr.	P.	5-8
Gen. II. Sanicula, L. . .	<i>Sanicle.</i>					
i. <i>S. Europæa, L.</i>	<i>Wood S.</i>	Woods .	W.	1-2 ft.	P.	6-7
Gen. III. Astrantia, L.						
[i. <i>A. major, L.</i>	. . . . .	Woods !.	pkW.	1-2 ft.	P.	6-7
Gen. IV. Eryngium, L.	<i>Eryngo.</i>					
i. <i>E. maritimum, L.</i>	<i>Sea Holly.</i>	Sand. sh.	B.	1-2 ft.	P.	7-8
ii. <i>E. campestre, L.</i>	<i>Field E.</i>	Sea sh. !.	B.	1-2 ft.	P.	7-8
Gen. V. Conium, L.	<i>Hemlock.</i>					
i. <i>C. maculatum, L.</i>	<i>common H.</i>	Was. pl.	W.	2-4 ft.	B.	6-7
Gen. VI. Physospermum	<i>Bladder-seed.</i>					
<i>Cuss.</i>						
i. <i>P. cornubiense, Hook.</i>	<i>Cornish B.</i>	Field ! Cornwall.	W.	1½-2 ft.	P.	7-8
Gen. VII. Smyrnum, L.	<i>Alexanders</i>					
i. <i>S. olusatrum, L.</i>	<i>common A.</i>	Was. pl.	y G.	3-4 ft.	B.	4-6
Gen. VIII. Cicutula, L.	<i>Cow-bane.</i>					
i. <i>C. virosa, L.</i>	<i>common C.</i>	Ditches !	W.	3-4 ft.	P.	6-8
Gen. IX. Apium, L.	<i>Celery.</i>					
i. <i>A. graveolens, L.</i>	<i>wild C.</i>	Salt mar.	g W.	1-2 ft.	B.	6-8
Gen. X. Petroselinum,	<i>Parsley.</i>					
<i>Hoffm.</i>						
[i. <i>P. sativum, Hoffm.</i>	<i>common P.</i>	Old walls !	Y.	1-2 ft.	B.	6-8
ii. <i>P. segetum, Koch.</i>	<i>Corn P.</i>	Banks .	W.	1-1½ ft.	A.	8-9
Gen. XI. Trinia, Hoffm.	<i>Hene-wort.</i>					
i. <i>T. vulgaris, DC.</i>	. . . . .	Li. rocks !	Y.	3-6 in.	P.	5-6

ORD. XXXIX.—CORNACEÆ.

CHARACTERISTICS OF SPECIES AND VARIETIES.

Gen. I.

- i. Shrub—Fls. in term., dense-flowered cymes.
- ii. Herb—Fls. in term., few-flowered umbels.

ORD. XL.—UMBELLIFERÆ.

Gen. I.

- i. Ls. peltate, cren.—Umbels five-flowered.

Gen. II.

- i. St. simple, leafless—Ls. all radical, palmate—Fls. sessile.

Gen. III.

- i. Rad. ls. palmate 5-7 partite ; lobes ov.-lanc., serr.—Invol. ent.]

Gen. IV.

- i. Rad. ls. spin., round, plaited, stalked, very glauc. ; upper ls. palm.
- ii. Rad. ls. tern. ; lobes pinnfid., spiny, pale green, upper ls. bi-pinnfid., amplex.

Gen. V.

- i. St. glab., spotted with purple—Partial invol. three-leaved, unilat.

Gen. VI.

- i. St. erect, paniced above—Lfts. cun., deeply cut—Umbels long-stalked.

Gen. VII.

- i. St. terete, stout—Rad. ls. yellow green with sheathing peti.—Umbels dense rounded.

Gen. VIII.

- i. St. furrowed, hollow—Ls. large ; rad. ones pinn. ; lfts. lanc., doubly serr., decurr.—Umbels large, flat.

Gen. IX.

- i. St. furrowed—Lfts. of rad. ls. round, lobed ; those of upper ls. cuneately notched—Umbels sess. or shortly stalked.

Gen. X.

- i. Ls. delt., bipinn., shining ; seg. of lower ls. trifiid—Fls. yellow.]
- ii. Ls. pinn., lower lfts. sess., lobed—Umbels drooping, irreg.—Fls. white.

Gen. XI.

- i. Root woody—Lfts. lin., filiform, glauc.—Fls. diæcious.

ORD. XL.—UMBELLIFERÆ—*cont.*

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
Gen. XII. Helosciadium, Koch.	<i>Marsh-wort.</i>					
i. <i>H. nodiflorum, Koch.</i>	<i>procumbent M.</i>	Bogs . . .	W.	Cr.	P.	7-8
v. 2. <i>repens, Koch.</i>			W.	Fl.	P.	6-7
ii. <i>H. inundatum, Koch.</i>	<i>least M.</i>	Pools . .	W.	Fl.	P.	6-7
Gen. XIII. Sison, L.	<i>Bastard Stone- parsley.</i>					
i. <i>S. Amonum, L.</i>		Waysides	W.	2-3 ft.	B.	8-9
Gen. XIV. Ægopodium, L.	<i>Gout-weed.</i>					
i. <i>Æ. Podagraria, L.</i>	<i>G. or Bishop-weed</i>	Damp pl.	g W.	1-2 ft.	P.	6-8
Gen. XV. Carum, L.	<i>Caraway.</i>					
i. <i>C. Carui, L.</i>	<i>common C.</i>	Was. pl.	W.	1-2 ft.	B.	6-7
ii. <i>C. verticillatum, Koch.</i>	<i>whorled-C.</i>	Past.!	pkW.	1-2 ft.	P.	7-8
iii. <i>C. Bulbocastanum, Koch.</i>	<i>tuberous C.</i>	Ch. fie.!	W.	1-2 ft.	P.	6-7
Gen. XVI. Bunium, Koch.	<i>Earthnut.</i>					
i. <i>B. flexuosum, With.</i>	<i>common E.</i>	Past. . .	W.	2-3 ft.	P.	5-6
Gen. XVII. Pimpinella, L.	<i>Burnet Saxifrage.</i>					
i. <i>P. Saxifraga, L.</i>	<i>common B.</i>	Dry pas.	W.	1-2 ft.	P.	7-9
ii. <i>P. magna, L.</i>	<i>large B.</i>	Bushy pl.	W.	3-4 ft.	P.	7-8
Gen. XVIII. Sium, L.	<i>Water Parsnep.</i>					
i. <i>S. latifolium, L.</i>	<i>broad-leaved W.</i>	Wet pl.	W.	3-4 ft.	P.	7-8
ii. <i>S. angustifolium, L.</i>	<i>narrow-leaved W.</i>	Ditches .	W.	1-3 ft.	P.	7-8
Gen. XIX. Bupleurum, L.	<i>Hare's-ear.</i>					
i. <i>B. tenuissimum, L.</i>	<i>slender H.</i>	Salt mar.!	Y.	6-12 in.	A.	8-9
ii. <i>B. aristatum, Bartl.</i>	<i>narrow-leaved H.</i>	Rocky pl.!	Y.	2-8 in.	A.	7-8
iii. <i>B. falcatum, L.</i>	<i>falcate-leaved H.</i>	Ongar Heath!	Y.	2-4 ft.	P.	8-9
iv. <i>B. rotundifolium, L.</i>	<i>common H.</i>	Corn fie..	Y.	1-1½ ft.	A.	6-7
Gen. XX. Œnanthe, L.	<i>Water-dropwort.</i>					
i. <i>Œ. fistulosa, L.</i>	<i>common W.</i>	Ditches .	W.	2-3 ft.	P.	7-9
ii. <i>Œ. pimpinelloides, L.</i>	<i>callous-fruited W.</i>	Past.!	W.	1-3 ft.	P.	6-8
iii. <i>Œ. Lachenalii, Gmel.</i>	<i>Parsley W.</i>	Salt mar.	W.	1-3 ft.	P.	6-8
iv. <i>Œ. silaifolia, Bieb.</i>	<i>Sulphur-wort</i>	Marshes!	W.	1-3 ft.	P.	6-7
v. <i>Œ. crocata, L.</i>	<i>Hemlock W.</i>	Marshes	W.	2-5 ft.	P.	7-8
vi. <i>Œ. Phellandrium, Lam.</i>	<i>fine-leaved W.</i>	Ditches .	W.	1-4 ft.	B.	7-9
vii. <i>Œ. fluviatilis, Colem.</i>	<i>River W.</i>	Streams .	W.	1-4 ft.	B.	7-9

CHARACTERISTICS OF SPECIES AND VARIETIES.

Gen. XII.

- i. Lfts. ov.-lanc., bluntly serr.—Umbels sub-sess., opp. the ls.
- v. 2. *Smaller—Lfts. acutely serr.—Umbels stalked.*
- ii. St. creeping or submerged—Lower ls. capillaceo-multifid; upper or flo. ls. pinn.—Umbels mostly of two rays.

Gen. XIII.

- i. Lower ls. pinn.; lfts. obl., serr.—Upper ls. smaller—Umbel irregular.

Gen. XIV.

- i. St. grooved, fistular—Ls. bi., or tri.-tern.; lfts. sharply serr., ov., acum.

Gen. XV.

- i. St. branched—Root fusiform—Ls. narrow-triang., bipinn.; lfts. with lin. seg.
- ii. St. slightly branched—Root fascicled—Ls. narrow; lfts. whorled, seg. capill.
- iii. St. erect, much branched—Root tuberous—Ls. broadly triang.; lfts. lin., not whorled.

Gen. XVI.

- i. St. slender, flex, leafless at base—Root tuberous—Lfts. pinnfid.; seg. lin.

Gen. XVII.

- i. Lfts. of lower ls. roundish, serr., variable; those of upper ls. with lin. seg.
- ii. Lfts. all ovate, serr.; term. one three-lobed.

Gen. XVIII.

- i. Lfts. obl.-lanc. equally serr.—Umbels term.
- ii. Lfts. ov., unequally lobed and serr.—Umbels lateral.

Gen. XIX.

- i. Ls. lin.-lanc.—Umbels term. and axill., sess. very small—Bracts. subulate.
- ii. Ls. lanc., striate, rigid, yellow—Umbels term.—Bracts. lanc., cusp.
- iii. Lower ls. ellip., long-stalked; upper ls. lin.-lanc.—Bracts. five lanc.]
- iv. Ls. oval, acute, perfoliate—Bracts. oval, leafy, connate.

Gen. XX.

- i. Root stoloniferous—St. ls. pinn., fili., fistulous—Invol. none.
- ii. Root fibres tuberous—Rad. ls. bipinn.; lfts. cuneate, acutely cut, upper lfts. lin., ent.—Invol. many-leaved, linear—Fr. with a swollen base.
- iii. Root fibres fleshy—Lfts. of lower ls. obl., ent., bluntly lobed; upper lfts. lin., acu.—Fr. narrowed below.
- iv. Root of ellip., obl., knots—Lfts. all lin., acute—Invol. o.
- v. Root of large fusiform tubers—Lfts. stalked cuneate, variously cut; upper narrower—Fr. ent., obl., cylind.
- vi. Aquatic—Rt. fibr.—St. erect—Segm. of submerged ls. with multifid diverging segments.
- vii. Aquatic—Rt. fibr.—St. floating—Segm. of submerged ls. with parallel segments.

ORD. XL.—UMBELLIFERÆ—*cont.*

Botanical Name.	Common Name.	Soil or Situation.	Colour	Growth.	Dura- tion.	Time of flowering.
Gen. XXI. <i>Æthusa</i>, L.	<i>Fool's-Parsley.</i>					
i. <i>Æ. Cynapium</i> , L. . .	<i>common F.</i> . . .	Waste pl.	W.	1-1½ ft.	A.	7-8
Gen. XXII. <i>Foeniculum</i>, <i>Hoffm.</i>	<i>Fennel.</i>					
i. <i>F. vulgare</i> , <i>Gaert.</i> . .	<i>Fennel</i>	Sea cliffs	Y.	3-5 ft.	P.	7-8
Gen. XXIII. <i>Seseli</i>, L.	<i>Meadow Saxi- frage.</i>					
i. <i>S. Libanotis</i> , <i>Koch.</i> .	<i>Mountain M.</i> . .	Ch. pas.!	W.	1-2 ft.	P.	7-8
Gen. XXIV. <i>Ligusticum</i>, L.	<i>Lovage.</i>					
i. <i>L. Scoticum</i> , L.	<i>Scottish L.</i> . . .	Sea rocks!	pkW.	1-3 ft.	P.	7
Gen. XXV. <i>Silaus</i>, Bess.	<i>Pepper-Saxifrage.</i>					
i. <i>S. pratensis</i> , Bess. . .	<i>common P.</i> . . .	Past. . .	Y.	1-3 ft.	P.	7-9
Gen. XXVI. <i>Meum</i>, <i>Tourn.</i>	<i>Spignel.</i>					
i. <i>M. athamanticum</i> , <i>Jacq.</i>	<i>S., Bald-money, or Meu.</i>	Alp. past.	y W.	6-18 in.	P.	6-7
Gen. XXVII. <i>Crithmum</i>, L.	<i>Samphire.</i>					
i. <i>C. maritimum</i> , L. . .	<i>Sea S.</i>	Sea rocks	W.	6-10 in.	P.	5-8
Gen. XXVIII. <i>Angelica</i>, L.	<i>Angelica.</i>					
i. <i>A. sylvestris</i> , L. . . .	<i>wild A.</i>	Moi. wds.	W.	2-3 ft.	P.	7-8
Gen. XXIX. <i>Peucedanum</i>, L.	<i>Hog's-fennel.</i>					
i. <i>P. officinale</i> , L. . . .	<i>Sea H., or Sea Sulphur Weed.</i>	Salt mar.!	Y.	2-3 ft.	P.	7-9
ii. <i>P. palustre</i> , <i>Mæench.</i> .	<i>Marsh H., or Milk Parsley.</i>	Bogs! . .	W.	4-5 ft.	P.	7-8
iii. <i>P. Ostruthium</i> , <i>Koch.</i>	<i>broad-leaved H., or Master-wort.</i>	Moist mead.!	W.	2-3 ft.	P.	6-7
Gen. XXX. <i>Pastinaca</i>, L.	<i>Parsnep.</i>					
i. <i>P. sativa</i> , L.	<i>wild P.</i>	Was. pl. .	Y.	2-3 ft.	B.	7-8
Gen. XXXI. <i>Heracleum</i>, L.	<i>Cow-Parsnep.</i>					
i. <i>H. Spondylium</i> , L. . .	<i>C., or Hog-weed..</i>	Was. pl. .	r W.	4-5 ft.	B.	7-8

ORD. XL.—UMBELLIFERÆ--*cont.*

CHARACTERISTICS OF SPECIES AND VARIETIES.

Gen. XXI.

- i. St. branched, leafy—Lfts. cuneate, lanc., decurr.—Invol. bracts. three, long, pendulous.

Gen. XXII.

- i. Lfts. capill., flaccid ; upper ones long—Fls. yellow.

Gen. XXIII.

- i. Ls. sess., hairy ; lfts. cut, seg. lanc., mucro.—Invol. bracts many.

Gen. XXIV.

- i. Ls. mostly radical ; lfts. dull green, sub-rhomb., dent.-serr.—Invol. of 5-7 ls.

Gen. XXV.

- i. Lfts. opp., lin.-lanc., ent. or three-lobed—Invol. of 1-2 s.

Gen. XXVI.

- i. Lfts. multi-partite ; segm. bristle-shaped, aromatic.

Gen. XXVII.

- i. Lfts. fleshy, lanc., lax—Invol. of many lanc. ls.

Gen. XXVIII.

- i. Lfts. equal, serr.—Teeth of cal. obsolete.

Gen. XXIX.

- i. Ls. five times partite ; segm. lin., flaccid—Umbel large ; fls. yellow—Bracts few, decid.
 ii. Ls. tripinn. ; lfts. pinnfid., seg. lin.-lanc.—Fls. white—Bracts many, persistent.
 iii. Ls. bi.-tern. ; lfts. ovate, serrate.

Gen. XXX.

- i. Lfts. ovate, downy beneath, cut and serr.—Fls. yellow—Invol. bracts o.

Gen. XXXI.

- i. Lfts. few, pinnfid., or lobed, serr., rough, hairy--Pet. unequal.

ORD. XL.—UMBELLIFERÆ—*cont.*

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
Gen. XXXII. Tordylium, L.	<i>Hart-wort.</i>					
i. <i>T. maximum, L.</i> . . .	<i>great H.</i>	Waysides!	r W.	1-2 ft.	A.	6-7
Gen. XXXIII. Daucus, L.	<i>Carrot.</i>					
i. <i>D. carota, L.</i>	<i>wild Carrot.</i> . . .	Past. . .	W.	1-2 ft.	B.	6-8
v. 2. <i>maritimus, With.</i> .	<i>Sea Carrot.</i> . . .	Sea sh.	. .	. .	. .	. .
Gen. XXXIV. Caulalis, L.	<i>Bur-Parsley.</i>					
i. <i>C. daucoides, L.</i> . .	<i>small B.</i>	Corn fie..	dkW.	6-18 in.	A.	6-7
[ii. <i>C. latifolia, L.</i> . . .	<i>great B.</i>	Ch. fie.!	Ro.	6-18 in.	A.	6-7
Gen. XXXV. Torilis, L.	<i>Hedge Parsley.</i>					
i. <i>T. Anthriscus, Gaert.</i>	<i>upright H.</i> . . .	Waste pl.	W.	2-3 ft.	A.	7-9
ii. <i>T. infesta, Spr.</i> . . .	<i>spreading H.</i> . .	Corn fie..	W.	4-12 in.	A.	6-9
iii. <i>T. nodosa, Gaert.</i> . .	<i>knotted H.</i> . . .	Waste pl.	r W.	6-18 in.	A.	5-7
Gen. XXXVI. Scandix, L.	<i>Shepherd's-Needle.</i>					
i. <i>S. Pecten-Veneris, L.</i>	<i>S., or Venus's Comb.</i>	Corn fie..	W.	4-12 in.	A.	6-9
Gen. XXXVII. Anthriscus, Hoffm.	<i>Beaked Parsley.</i>					
i. <i>A. vulgaris, Pers.</i> . .	<i>common B.</i> . . .	Waysides	W.	2-3 ft.	B.	5-6
ii. <i>A. sylvestris, Hoffm.</i> .	<i>wild B.</i>	Waysides	W.	3-4 ft.	P.	4-6
[iii. <i>A. Cerefolium, Hoffm.</i>	<i>Garden B., or Chervil.</i>	Hedges .	W.	1-1½ ft.	A.	5-7
Gen. XXXVIII. Chærophyllum, L.	<i>Chervil.</i>					
i. <i>C. temulum, L.</i> . . .	<i>rough Chervil.</i> . .	Waysides	W.	1-3 ft.	B.	6-7
Gen. XXXIX. Myrrhis, Tourn.	<i>Cicely.</i>					
i. <i>M. odorata, Scop.</i> . .	<i>sweet C.</i>	Northern past.	W.	2-3 ft.	P.	5-6
Gen. XL. Coriandrum, L.	<i>Coriander.</i>					
[i. <i>C. sativum, L.</i>	. . . .	Was. pl..	pkW.	1-2 ft.	A.	6-7

ORD. XLI.—LORANTHACEÆ.

Gen. I. Viscum, L.	<i>Mistletoe.</i>					
i. <i>V. album, L.</i>	<i>common M.</i> . . .	Apple trees.	G.	1-3 ft.	Shb.	3-5

ORD. XL.—UMBELLIFERÆ—*cont.*

CHARACTERISTICS OF SPECIES AND VARIETIES.

Gen. XXXII.

- i. St. hispid, with reflexed hairs—Lfts. lanc., deeply serr.—Umbels small, crowded.

Gen. XXXIII.

- i. Lfts. pinnfid. ; seg. acute—Central fl. purple—Umbel in fr. concave.
 v. 2. *Lfts. more fleshy—Umbels in fr. convex.*

Gen. XXXIV.

- i. St. furrowed, hairy at joints—Ls. bipinn. ; lfts. lin. acute—Umbels trifid ; branched three-seeded.
 ii. St. scabrous—Ls. pinn. ; lfts. lanc., serr.—Umbels trifid ; branches five-seeded.]

Gen. XXXV.

- i. St. erect—Umbels stalked, term.—Invol. of many ls.—Fr. bristles not hooked.
 ii. St. erect, much branched—Umbels stalked, term.—Invol. of one leaf—Fr. scabrous with hooked bristles.
 iii. St. prostr., spreading—Umbels lat., sess.—Fr. with hooked bristles.

Gen. XXXVI.

- i. St. erect—Lfts. many partite ; seg. lin.—Bracts. long, ent. or bifid.

Gen. XXXVII.

- i. Umbels lat., stalked—Fr. ovate, hispid.
 ii. Umbels term., stalked—Fr. lin., glab.—Beak short.
 iii. Umbels lat., sess.—Fr. lin., glab.—Beak long.

Gen. XXXVIII.

- i. Joints of st. swollen—Ls. rough, spotted ; lfts. ovate, cut ; seg. obt., mucro.

Gen. XXXIX.

- i. Ls. tripinn., whitish beneath—Invol. bracts. lanc., acum.

Gen. XL.

- i. St. fistular, slender—Ls. deltoid, bipinn. ; upper lfts. lin., fleshy, lax.]

ORD. XLI.—LORANTHACEÆ.

Gen. I.

- i. St. much branched—Ls. ob.-ov., lanc.—Fls. in axill. clusters.

ORD. XLII.—CAPRIFOLIACEÆ.

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
Gen. I. Sambucus, L.	<i>Elder.</i>					
i. <i>S. nigra, L.</i>	<i>common E.</i>	Woods .	W.	10-12 ft.	T.	6
ii. <i>S. Ebulus, L.</i>	<i>dwarf E., or Dane-wort.</i>	Was. pl..	puW.	2-3 ft.	P.	7-8
Gen. II. Viburnum, L.	<i>Gullder-rose.</i>					
i. <i>V. Opulus, L.</i>	<i>common G.</i>	Woods .	W.	6-8 ft.	Shb.	6-7
ii. <i>V. Lantana, L.</i>	<i>mealy G., or Way- faring tree.</i>	Woods .	W.	10-20 ft.	Shb.	5-6
Gen. III. Lonicera, L.	<i>Honey-suckle.</i>					
i. <i>L. Periclymenum, L.</i> . .	<i>H., or Woodbine.</i>	Hedges .	R.	Cli.	Shb.	6-9
[ii. <i>L. caprifolium, L.</i> . . .	<i>perfoliate H.</i> . . .	Woods! .	R.	Cli.	Shb.	5-6
[iii. <i>L. Xylosteum, L.</i> . . .	<i>upright Fly H.</i> . .	Thick! .	Y.	3-5 ft.	Shb.	5-6
Gen. IV. Linnæa, Gronov.	<i>Linnaea.</i>					
i. <i>L. borealis, Gro.</i>	<i>two-flowered L.</i> . .	Sco. woods	Pk.	3-8 in.	P.	7

ORD. XLIII.—RUBIACEÆ.

Gen. I. Rubia, L.	<i>Madder.</i>					
i. <i>R. peregrina, L.</i>	<i>wild M.</i>	Sto. pl.! SW. of E.	Y.	1-2 ft.	P.	6-8
Gen. II. Galium, L.	<i>Bed-straw.</i>					
*						
i. <i>G. verum, L.</i>	<i>yellow B.</i>	Downs	Y.	1-2 ft.	P.	6-9
v. 2. <i>ochroleucum</i>	.....	.....	Y.	.....	.....	.....
ii. <i>G. cruciatum, With.</i> . .	<i>Cross-wort B., or Mug-wort.</i>	Waysides	Y.	6-18 in.	P.	4-6
A.						
iii. <i>G. palustre, L.</i>	<i>white Water B.</i> . .	Mar. . .	W.	1-3 ft.	P.	7-8
v. 2. <i>elongatum, Presl.</i> . .	.....	.....	.....	.....	.....	.....
v. 3. <i>Witheringii, Sm.</i> . .	.....	.....	.....	.....	.....	.....
iv. <i>G. uliginosum, L.</i> . . .	<i>rough Water B.</i> . .	Mar. . .	W.	2-5 ft.	P.	7-8
B.						
v. <i>G. saxatile, L.</i>	<i>smooth Heath B.</i>	Downs .	W.	4-6 in.	P.	6-8
vi. <i>G. erectum, Huds.</i> . . .	<i>upright B.</i>	Downs .	W.	1-2 ft.	P.	6-8
v. 2. <i>aristatum, Sm.</i> . . .	.....	.....	.....	.....	.....	.....
vii. <i>G. mollugo, L.</i>	<i>great Hedge B.</i> . .	Hedges .	W.	1-4 ft.	P.	7-8
v. 2. <i>scabrum, With.</i> . . .	.....	.....	.....	.....	.....	.....
v. 3. <i>insubricum, Gaud.</i> . .	.....	.....	.....	.....	.....	.....

ORD. XLII.—CAPRIFOLIACEÆ.

CHARACTERISTICS OF SPECIES AND VARIETIES.

Gen. I.

- i. Tree—Cymes with five branches—Stip. o.
- ii. Herb—Cymes three fid.—Stip. leafy.

Gen. II.

- i. Ls. three-lobed, acum., glab.
- ii. Ls. ellip., cord., downy beneath—Young shoots downy.

Gen. III.

- i. St. climb.—Ls. oval, sess.—Fls. in term., stalked heads—Fr. red.
- ii. St. climb.—Upper ls. connate, perfol.—Fls. in term., sess., whorls—Fr. orange.]
- iii. St. erect—Ls. oval, downy—Fls. axill. in pairs—Fr. scarlet.]

Gen. IV.

- i. St. creeping ; pedunc. erect, two-flowered—Ls. ovate, opp.—Cor. campan.

ORD. XLIII.—RUBIACEÆ.

Gen. I.

- i. Ls. obl. 4-6 in a whorl, evergreen, rough—Fls. in term. and axill. cymes.

Gen. II.

* *Flowers yellow.*

- i. Ls. eight, lin., rough—Fls. term. and axill. in dense pan.
- v. 2. *Fls. pale yellow (a hybrid between this and the next). (Deal.)*
- ii. Ls. four, ovate, hairy—Fls. axill., whorled.

* * *Fls. white.*

A. Root perennial—St. with reflexed bristles.

- iii. Ls. 4-6 obl.-lanc., obt., slightly cili.—St. diffuse, rough—Fr. smooth on spreading pedi.

v. 2. *Much larger—St. thicker, less branched—Ls. longer.*

v. 3. *Like var. 2, but margins of ls. with more prickles—Ls. narrower and recurved—Pan. narrow.*

- iv. St. slender, weak—Ls. 6-8 lin.-lanc., rigid, mucro.—Fr. granulated on erect stalks

B. Root perennial—St. without reflexed bristles.

v. St. prost., tufted—Ls. six, ob.-ov., mucro.—Fr. stalks erect ; fr. granulated.

- vi. St. erect—Ls. (6-8) lin.-lanc., glauc., with prickles pointing forward—Pan. with slender, erect branches—Fr. smooth.

v. 2. *Leaves larger and narrower.*

- vii. St. long and straggling, thick at joints—Ls. eight, ellip., obt., mucro., margins rough, with prickles pointing forward or erect—Pan. with spreading branches—Fr. glab.

v. 2. *Stem rough and hairy.*

v. 3. *Ls. six, ob.-ov.—Branches of pan. few-flowered.*

ORD. XLIII.—RUBIACEÆ—cont.

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
v. 4. <i>Bakeri</i> , Syme. . . .						
viii. <i>G. sylvestre</i> , Poll. . .	mount B.	Lim. hills!	W.	3-6 in.	P.	7-8
v. 2. <i>nitidulum</i> , Thuil. .						
ix. <i>G. boreale</i> , L.	cross-leaved B. .	Moi. rocks N. in the N.	W.	1-2 ft.	P.	6-8
	C.					
x. <i>G. Aparine</i> , L.	Goose-grass, or Cleavers.	Hedges .	W.	1-5 ft.	A.	6-7
v. 2. <i>Vaillantii</i> , DC.						
xi. <i>G. tricorne</i> , With. . .	rough-fruited Corn B.	Cha. fie..	W.	1-3 ft.	A.	6-10
xii. <i>G. anglicum</i> , Huds. .	Wall B.	San. pl. SE. of E.	W.	6-12 in.	A.	6-7
Gen. III. <i>Sherardia</i> , L.	Field Madder.					
i. <i>S. arvensis</i> , L.	blue S. or F. . .	Corn fie..	Li.	6-12 in.	A.	4-10
Gen. IV. <i>Asperula</i> , L.	Woodruff.					
i. <i>A. odorata</i> , L.	sweet W.	Sh. woods	W.	6-8 in.	A.	5-6
ii. <i>A. cynanchica</i> , L. . .	small W., or Squinancy-wort.	Downs .	Pk.	6-10 in.	P.	6-7

ORD. XLIV.—VALERIANACEÆ.

Gen. I. <i>Centranthus</i> , DC.	Spur-Valerian.					
[i. <i>C. ruber</i> , DC.	red S.	Ch. pits .	Ro.	1-2 ft.	P.	6-9
Gen. II. <i>Valeriana</i> , L.	Valerian.					
i. <i>V. dioica</i> , L.	small marsh V. .	Mar. . .	p Ro.	6-8 in.	P.	5-6
ii. <i>V. officinalis</i> , L. . .	great wild V. . .	Moi. wds.	Fl.	2-4 ft.	P.	6-8
a. <i>Mikanii</i>	. . . . .	. . . . .	. . .	. . .	. . .	. . .
b. <i>sambucifolia</i> , Mik. . .	. . . . .	. . . . .	. . .	. . .	. . .	. . .
[iii. <i>V. pyrenaica</i> , L. . .	heart-leaved V. .	Woods, Sc.	. . .	. . .	. . .	. . .
Gen. III. <i>Valerianella</i> , <i>Tournef.</i>	Corn-salad.					
*						
i. <i>V. olitoria</i> , Mæench. .	common C., or Lamb's-lettuce.	Corn fie..	p B.	3-12 in.	A.	4-6
ii. <i>V. carinata</i> , Lois. . .	carinate C. . . .	Banks! .	p Li.	6-12 in.	A.	4-6
iii. <i>V. Auricula</i> , DC. . .	sharp-fruited C. .	Corn fie..	p B.	6-12 in.	A.	6-8
iv. <i>V. dentata</i> , Koch. . .	smooth C.	Corn fie..	Fl.	6-12 in.	A.	6-8

ORD. XLIII.—RUBIACEÆ—*cont.*

CHARACTERISTICS OF SPECIES AND VARIETIES.

v. 4. *Ls.* 6-8 *lin.*

viii. *St.* diffuse—*Ls.* 6-8 *lin.-lanc.*, cili., bristle-pointed—*Fr.* stalks spreading; fr. minutely granulate.

v. 2. *Ls.* narrow, margins revolute—*Pan.* compact.

ix. *St.* erect—*Ls.* four, *lanc.*, glab.—*Fr.* covered with hooked-bristles.

C. Root annual.

x. *St.* straggling—*Ls.* *lanc.*, keeled, rough with reflexed prickles—*Fr.* covered with hooked bristles.

v. 2. *Fls.* very small, greenish yellow (*Saffron-Walden*).

xi. *St.* procumb., spreading—*Ls.* 6-8 *lin.-lanc.*, rough with reflexed prickles—*Cymes* three-flowered—*Fr.* large, granulated, drooping.

xii. *St.* slender, spreading—*Ls.* six, *lin.-lanc.*, mucro., with prickles pointing forward—*Fr.* minute, tubercled.

Gen. III.

i. *Ls.* 4-6 *obl.-lanc.*, acute—*Fls.* in term., sess., umbels.

Gen. IV.

i. *Ls.* 6-8 *obl.-lanc.* all equal—*Fls.* white—*Fr.* hispid.

ii. *Ls.* four, *lin.*, very unequal—*Fls.* pink—*Fr.* rugose.

ORD. XLIV.—VALERIANACEÆ.

Gen. I.

i. *Ls.* ov.-*lanc.*—*Fls.* in term., pan., cymes—*Spur* slender.]

Gen. II.

i. *Rad.* *ls.* ov. ; *st.* *ls.* pinnfid.

ii. *Ls.* all pinn. ; *lfts.* *lanc.*

a. *Lfts.* 6-10 pairs.

b. *Lfts.* 4-6 pairs.

iii. *Ls.* large, cord., deeply serr. ; upper with one or two pair of small basal *lfts.*]

Gen. III.

* Fertile cell corky.

i. *Fls.* in dense cymes—*Fr.* compressed, crowned with three minute, reflexed teeth.

** Fertile cell not corky.

ii. *Fls.* in dense cymes—*Fr.* obl. crowned with one straight tooth—*Barren* cells contiguous, equal to fertile cell.

iii. *Fls.* in lax cymes—*Fr.* sub-glob., turgid, crowned with one erect tooth—*Barren* cells contiguous, larger than fertile cell.

iv. *Fls.* in lax cymes—*Fr.* ov., flattish, crowned with a four-toothed lobe—*Barren* cells remote, very small.

ORD. XLV.—DIPSACEÆ.

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
Gen. I. Dipsacus, L.	<i>Teasel.</i>					
i. <i>D. sylvestris, L.</i>	<i>wild T.</i>	Was. pl..	<i>pu</i> Pu.	4-5 ft.	P.	8-9
i. <i>D. pilosus, L.</i>	<i>small T.</i>	Moi. wds.	W.	2-4 ft.	P.	8-9
Gen. II. Scabiosa, L.	<i>Scabious.</i>					
i. <i>S. succisa, L.</i>	<i>Devil's-bit S.</i> . .	Pas. . .	<i>pu</i> B.	1-2 ft.	P.	7-10
ii. <i>S. columbaria, L.</i> . .	<i>small S.</i>	Dry hths.	Li.	1-2 ft.	P.	7-8
Gen. III. Knautia, L.	<i>Knautia.</i>					
i. <i>K. arvensis, Coult.</i> . .	<i>Field K.</i>	Corn fie..	Li.	2-5 ft.	P.	6-8

ORD. XLVI.—COMPOSITÆ.

Gen. I. Tragopagon, L.	<i>Goat's-beard.</i>					
i. <i>T. pratensis, L.</i> . . .	<i>yellow G.</i>	Was. pl..	Y.	1-2 ft.	B.	6-7
v. 2. <i>minor, Fries.</i>	. . . .	. . . .	. .	. .	. .	. .
v. 3. <i>grandiflorus, Sy.</i> . .	. . . .	. . . .	. .	. .	. .	. .
ii. <i>T. porrifolius, L.</i> . .	<i>purple G. or Salsify.</i>	Moi. mead.	Pu.	1-2 ft.	P.	5-6
v. 2. <i>parviflorus, Sy.</i> . .	. . . .	. . . .	. .	. .	. .	. .
Gen. II. Helminthia, Fuss.	<i>Ox-tongue.</i>					
i. <i>echioides, Gaert.</i> . .	<i>bristly O.</i>	Waysides	Y.	2-3 ft.	A.	6-10
Gen. III. Picris, L.	<i>Picris.</i>					
i. <i>P. hieracioides, L.</i> . .	<i>Hawkweed P.</i> . .	Waysides	Y.	2-3 ft.	B.	6-10
v. 2. <i>arvalis</i>	. . . .	. . . .	. .	. .	. .	. .
Gen. IV. Leontodon, L.	<i>Hawkbit.</i>					
i. <i>L. hirtus, L.</i>	<i>hairy H.</i>	Moors .	Y.	4-8 in.	P.	7-8
ii. <i>L. hispidus, L.</i> . . .	<i>rough H.</i>	Past. .	Y.	6-18 in.	P.	6-9
iii. <i>L. autumnalis, L.</i> . .	<i>Autumnal H.</i> . .	Past. .	Y.	6-18 in.	P.	8-9
v. 2. <i>pratensis, Koch.</i> . .	. . . .	Mount. .	. .	. .	. .	. .
Gen. V. Hypochaeris, L.	<i>Cat's-ear.</i>					
i. <i>H. glabra, L.</i>	<i>smooth C.</i>	Dry. pl. .	Y.	6-12 in.	A.	6-8
v. 2. <i>Balbisii, Lois.</i> . . .	. . . .	. . . .	. .	. .	. .	. .
ii. <i>H. maculata, L.</i> . . .	<i>spotted C.</i>	Ch. past.!	Y.	6-12 in.	P.	7-8
iii. <i>H. radicata, L.</i> . . .	<i>long-rooted C.</i> . .	Waysides	Y.	6-18 in.	P.	7-8
Gen. VI. Lactuca, L.	<i>Lettuce.</i>					
i. <i>L. virosa, L.</i>	<i>strong-scented L.</i>	Was. pl..	Y.	3-5 ft.	B.	7-8
ii. <i>L. scariola, L.</i> . . .	<i>prickly L.</i>	Was. pl.!	Y.	3-5 ft.	B.	7-8

ORD. XLV.—DIPSACEÆ.

CHARACTERISTICS OF SPECIES AND VARIETIES.

Gen. I.

- i. Ls. simp., opp., sess.
- ii. Ls. stalked, with a basal pair of lfts.—Heads small, hairy.

Gen. II.

- i. Ls. ent.—Cor. four cleft.
- ii. Ls. pinnfid.—Cor. five cleft.

Gen. III.

- i. Lower ls. sinn. ; upper pinnfid.

ORD. XLVI.—COMPOSITÆ.

Gen. I.

- i. Ls. glab., acum., channelled—Pedunc. slightly thickened—Invol. equalling the florets—Fls. yellow.
- v. 2. *Invol. longer than the florets.*
- v. 3. *Invol. half as long as florets.*
- ii. Ls. as Sp. 1—Pedunc. much thickened above—Fls. purple.
- v. 2. *Florets half as long as the bracts.*

Gen. II.

- i. St. stout, hispid, with rigid, hooked bristles from tubercular bases—Lower ls. aur. ; upper cord., amplex.

Gen. III.

- i. St. rough with hooked bristles—Ls. lanc. toothed—Fls. corymb.
- v. 2. *Branches erect—Fls. umbellate.*

Gen. IV.

- i. Ls. lanc., hisp.—Scapes one-flowered—Invol. glab.
- ii. Ls. runc., hisp.—Scapes one-flowered—Invol. hairy.
- iii. Ls. lanc., toothed, glab.—Scapes branched—Invol. glab.
- v. 2. *Invol. shaggy, with black hairs.*

Gen. V.

- i. St. branched, glab.—Ls. obl., dent., sinu.—Invol. as long as florets—Inner fruits beaked.
- v. 2. *All the fruits beaked.*
- ii. St. sol., glab.—Ls. spath., toothed, spotted, scab.
- iii. St. branched, glab.—Ls. runc., obt.—Invol. shorter than the florets.

Gen. VI.

- i. Scab.—Ls. spreading, ob.-ov., toothed—Beak equalling the black fr.
- ii. Scab.—Ls. erect, lanc., sinu., sagitt.—Beak equalling the grey fr.

ORD. XLVI.—COMPOSITÆ—*cont.*

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
iii. <i>L. saligna</i> , <i>L.</i>	<i>least L.</i>	Ch. pas.! near sea.	Y.	12-18 in.	B.	7-8
iv. <i>L. muralis</i> , <i>L.</i>	<i>Ivy-leaved L.</i> . . .	Walls & woods.	Y.	1-3 ft.	A.	6-8
Gen. VII. <i>Sonchus</i>, <i>L.</i>	<i>Sow-thistle.</i>					
i. <i>S. palustris</i> , <i>L.</i>	<i>Marsh S.</i>	Mar.! . .	Y.	5-7 ft.	P.	7-8
ii. <i>S. arvensis</i> , <i>L.</i>	<i>corn S.</i>	Corn fie..	Y.	3-4 ft.	P.	8-9
iii. <i>S. asper</i> , <i>Hoffm.</i>	<i>common S.</i>	Was. pl..	Y.	2-3 ft.	A.	6-8
iv. <i>S. oleraceus</i> , <i>L.</i>	<i>sharp-fringed S.</i> . .	Was. pl..	Y.	2-3 ft.	A.	6-8
Gen. VIII. <i>Mulgedium</i>, <i>Cass.</i>	<i>Blue Sow-thistle.</i>					
i. <i>M. alpinum</i> , <i>Less.</i>	<i>Alpine B.</i>	Alp. rocs.!	B.	3-4 ft.	P.	7-8
Gen. IX. <i>Crepis</i>, <i>L.</i>	<i>Hawk's-beard.</i>					
i. <i>C. virens</i> , <i>L.</i>	<i>smooth H.</i>	Was. pl..	Y.	1-3 ft.	P.	7-8
ii. <i>C. biennis</i> , <i>L.</i>	<i>rough H.</i>	Ch. pas. .	Y.	1-4 ft.	B.	6-7
iii. <i>C. succisifolia</i> , <i>Tausch.</i>	<i>Succory-leaved H.</i>	Woods, N. of E.	Y.	1-2 ft.	P.	7-8
iv. <i>C. paludosa</i> , <i>Mærch.</i> . .	<i>Marsh H.</i>	Moi. wds.	Y.	1-3 ft.	P.	7-9
Gen. X. <i>Hieracium</i>, <i>L.</i>	<i>Hawkweed.</i>					
*						
i. <i>H. Pilosella</i> , <i>L.</i>	<i>Mouse-ear H.</i> . . .	Dry banks	Y.	2-10 in.	P.	5-8
v. 2. <i>villosissimum</i>	. . . . .	. . . . .	. . .	. . .	. . .	. . .
[ii. <i>H. aurantiacum</i> , <i>L.</i> . .	<i>Orange H.</i>	Copses N. of E.	O.	1-2 ft.	P.	6-8
* *						
<i>A.</i>						
iii. <i>H. alpinum</i> , <i>L.</i>	<i>Alpine H.</i>	Sc. mount.	Y.	4-12 in.	P.	7-8
iv. <i>H. holosericeum</i> , <i>Back.</i>	. . . . .	Cumb. .	. .	. .	. .	. .
v. <i>H. eximium</i> , <i>Back.</i> . . .	. . . . .	Sc. mount.	. .	6-15 in.	. .	. .
v. 2. <i>tenellum</i>	. . . . .	. . . . .	. .	. .	. .	. .
vi. <i>H. calenduliflorum</i> , <i>Back.</i>	. . . . .	Sc. mount.	. .	. .	. .	. .
<i>B.</i>						
vii. <i>H. gracilentum</i> , <i>Back.</i>	. . . . .	Sc. mount.	. .	6-10 in.	. .	. .
viii. <i>H. globosum</i> , <i>Back.</i> .	. . . . .	. . . . .	. .	. .	. .	. .

CHARACTERISTICS OF SPECIES AND VARIETIES.

- iii. Glab.—Ls. lin., ent., acum., sagitt.—Beak twice as long as fr.
- iv. Glab.—Ls. lyr., pinnfid. ; term. lobe angled—Beak much shorter than fr.

Gen. VII.

- i. Perenn.—Rootstock branched—Lower ls. runc., minutely toothed ; upper ls. ent., sagitt.
- ii. Perenn.—Rootstock creeping—Lower ls. runc., sharply toothed ; upper ls. ent., cord.
- iii. Ann.—Ls. ent. or pinnfid. ; auricles rounded—Fr. ribbed, not transversely wrinkled.
- iv. Ann.—Ls. ent. or pinnfid. ; auricles arrow-shaped—Fr. ribbed and transversely wrinkled.

Gen. VIII.

- i. Ls. lyr., sagitt. ; term. lobe large, delt.—Fls. blue.

Gen. IX.

- i. Glab.—Rad. ls. lyr. ; st. ls. lin., sagitt.—Inner bracts of invol. glab. within—Fr. shorter than pappus.
- ii. Hesp.—Rad. ls. runc.-lyr. ; st. ls. lanc., sess.—Inner bracts of invol. downy within—Fr. longer than pappus.
- iii. Glab. or hairy—Rad. ls. spath. nearly glab. ; st. ls. obl. half-amplex.—Outer bracts of invol. short, adpressed.
- iv. Glab.—Rad. ls. obl.-lanc. ; st. ls. oval-lanc., amplex.—Invol. gland. hairy.

Gen. X.

* *Plants with running scions.*

- i. St. slender, leafless—Ls. ent. ellip.-lanc., downy—Fls. sol.
- v. 2. Fls. larger—Plant with long silky hairs—Root scions short.
- ii. St. leafy—Ls. ov.-lanc., hairy—Fls. corymb.]
- * * *Plants producing radical rosettes of ls. in autumn.*
- A. St. generally one-headed—Invol. shaggy or silky ; bracts lax—Florets extremely hairy.
- iii. St. hairy, one-headed—Rad. ls. ov., spath. almost glab. above ; st. ls. lin. lanc.—Invol. hemisph., hairy ; scales all lax—Styles yellow.
- iv. St. silky, one-headed—Rad. ls. blunt, hairy on both sides—Invol. turbinate, silky ; inner scales adpr.—Styles yellow.
- v. St. floccose, one or two-headed—Rad. ls. lanc., acute, sharply dent.—Invol. truncate, shaggy ; scales attenuate—Styles livid.
- v. 2. St. short—Rad. ls. lin.-lanc.—Styles yellow.
- vi. Like No. 5, but heads larger and generally solitary—Rad. ls. ovate, spath., obt.

B. St. one or few-headed—Invol. villous or hairy ; bracts adpressed, or a few outer lax—Florets nearly glab., with a hairy tip.

- vii. St. one or few-headed—Rad. ls. obl.-lanc. ; st. ls. large—Invol. ventricose, densely setose ; inner bracts broad—Styles livid.
- viii. St. one or few-headed—Rad. ls. ovate, shortly stalked ; st. ls. few, small—Invol. glob. hairy ; bracts lin., attenu.—Styles yellow.

ORD. XLVI.—COMPOSITÆ—*cont.*

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
ix. <i>H. nigrescens</i> , <i>Willd.</i>	. . . .	Sc. mount.	Y.	6-12 in.	P.	7-8
x. <i>H. lingulatum</i> , <i>Back.</i>	. . . .	Sc. mount.	Y.	1-2 ft.	P.	7-8
xi. <i>H. senescens</i> , <i>Back.</i>	. . . .	Sc. mount.	Y.	15-18 in.	P.	7-8
xii. <i>H. chrysanthum</i> , <i>Back.</i>	. . . .	Sc. mount.	Y.	9-15 in.	P.	7-8
v. 2. <i>microcephalum</i> .	. . . .	Cumb. .	Y.	. . .	. . .	. . .
xiii. <i>H. cerinthoides</i> , <i>L.</i>	. . . .	Sc. mount.	Y.	12-18 in.	P.	7-8
v. 2. <i>anglicum</i> , <i>Fries.</i>	. . . .	Teesdale, &c.	. .	. .	. .	. .
v. 3. <i>acutifolium</i> , <i>Back.</i>	. . . .	Sco. . . .	. .	. .	. .	. .
xiv. <i>H. iricum</i> , <i>Fries.</i> .	. . . .	Mount. .	Y.	1-3 ft.	P.	7-8
<i>C.</i>						
*						
xv. <i>H. pallidum</i> , <i>Fr.</i> .	. . . .	Mount. .	Y.	6-18 in.	P.	6-9
xvi. <i>H. lasiophyllum</i> , <i>Koch.</i>	. . . .	Sc. mount.	Y.	. .	. .	. .
xvii. <i>H. Gibsoni</i> , <i>Back.</i> .	. . . .	Settle. .	Y.	6-18 in.	. .	. .
xviii. <i>H. argenteum</i> , <i>Fr.</i>	. . . .	Mount. .	Y.	1-2 ft.	P.	7-8
xix. <i>H. nitidum</i> , <i>Back.</i> .	. . . .	Sc. mount.	Y.	1-2 ft.	P.	6-7
xx. <i>H. aggregatum</i> , <i>Back.</i>	. . . .	Mount. .	Y.	1-2 ft.	P.	6-7
* *						
xxi. <i>H. murorum</i> , <i>L.</i> .	. . . .	Rocky pl.	Y.	12-18 in.	P.	6-7
xxii. <i>H. cæsium</i> , <i>Fr.</i> . .	. . . .	Mount. .	Y.	12-18 in.	P.	6-7
v. 2. <i>obtusifolium</i> . . .	. . . .	. . .	. .	. .	. .	. .
xxiii. <i>H. stelligerum</i> , <i>Froel.</i>	. . . .	Sc. mount.	Y.	1-2 ft.	P.	6-7
xxiv. <i>H. vulgatum</i> , <i>Fr.</i> .	. . . .	Woods .	Y.	1-3 ft.	P.	7-9
v. 2. <i>cinereum</i>	. . . .	. . .	. .	. .	. .	. .
v. 3. <i>rubescens</i>	. . . .	. . .	. .	. .	. .	. .

CHARACTERISTICS OF SPECIES AND VARIETIES.

- ix.** St. few headed—Rad. ls. ov.-lanc., irregularly and coarsely dent. with slender stalks; st. ls. few, shortly stalked—Invol. ov.; bracts broad, outer obt., erect in bud—Styles brown.
- x.** St. few headed—Rad. ls. obl. dentic., apic., coarsely hairy, with short stalks; st. ls. few, sess.—Invol. trunc. at base; bracts atten. acute, erect in bud—Sty. livid.
- xi.** St. few headed—Rad. ls. ellip.-lanc., remotely and evenly dent.; st. ls. lin.-lanc., with short slender stalks—Invol. turbinate; bracts attenuate, inflexed in bud.
- xii.** St. few headed—Rad. ls. ov., acu., coarsely dent. on long slend. stalks; st. ls. subu., stalked—Fls. large, drooping—Invol. rounded at base; bracts lin., attenu.; outer ones lax., inflexed in bud.
- v. 2. St. simple—Rad. ls. more evenly dent.—Fls. smaller, erect—Invol. narrower.*
- xiii.** Glauous green—St. few-headed—Rad. ls. ov.-lanc., with long shaggy stalks; st. ls. few, ov., acu., amplex.—Florets pilose at tip—Invol. ventr.; bracts acum., erect in buds—Styles livid.
- v. 2. St. with one large sol. leaf, and long asc., lat., fl.-stalks—Bracts all adpr.*
- v. 3. Very glauc.—St. branched from base—Ls. very ac., sharply toothed, glab.—Invol. bracts adpr.*
- xiv.** Glauous gr.—St. leafy, corymb.—Rad. ls. ov. with short stalks; st. ls. broadly ovate, amplex.—Florets with glab. tips—Invol. trunc.; bracts blunt—Styles livid.
- C. St. few headed—Invol. hairy, with adpressed bracts—Florets externally nearly hairless, with glab. tips.*
- * *Sty. yellow.*
- xv.** Very glauc.—St. glab.—Rad. ls. ovate lanc., coarsely ciliate, stalked; st. ls. (o-1) lanc., sess.—Invol. ov.; bracts acute.
- xvi.** Ash-coloured—St. brittle, pilose—Rad. ls. oval, coarsely pilose on both sides; st. ls. (rare) acute at each end—Invol. vent.; bracts cuspl.
- xvii.** Glau.—St. rigid, leafless—Rad. ls. broadly ov., spotted—Pedunc. bifid, straight—Invol. trunc.; bracts broad, blunt.
- xviii.** Very glauc.—St. hollow, glab.—Rad. ls. lanc., almost ent.; st. ls. (1-3) lin.-lanc., sess.—Pedunc. long, rigid—Invol. broadly ventr., bracts blunt.
- xix.** Dark green—St. glab.—Rad. ls. lanc., acum., coarsely toothed; st. ls. (o-1)—Pedunc. spreading, rigid, scaly—Invol. scales acum.
- xx.** Dark green—St. sub-umbell., floccose—Rad. ls. broad, ov., glab.; st. ls. (1) lanc., sess.—Pedunc. erect, aggregate—Invol. narrow, conical, hairless: bracts blunt.
- * * *Styles livid.*
- xxi.** Light green—St. corymb.—Rad. ls. thin, ov., dent.: st. ls. (o-1) stalked—Pedunc. arc. short—Invol. gr.; bracts ac., inner erect in bud.
- xxii.** Dull green—St. few headed, leafless—Rad. ls. coriaceous, ov.—Pedunc. straight—Invol. vent.; outer bracts blunt, inner ac. inflexed in bud.
- v. 2. Yellowish gr.—Rad. ls. soft, rounded at each end; st. l. (1)—Pedunc. spreading—Fls. pale.*
- xxiii.** Ashy green—St. leafy, few headed, corymb.—Rad. ls. ov., floccose on both sides, long stalked; st. ls. large, semi-amplex.—Pedunc. floccose—Invol. ov.; bracts acum., inflexed in bud.
- xxiv.** Purplish or glau.—St. leafy, branched, corymb.—Rad. ls. obl. or lanc. dent. at middle, narrowed to base; lower st. ls. stalked, upper sess.—Pedunc. straight, floccose—Invol. cylind. in bud; bracts equally attenu.
- v. 2. Ashy-green—St. short, much branched—Rad. ls. few, ob.-ov. lanc.*
- v. 3. St. stout, purplish—Rad. ls. many; st. ls. (1-2) semi-amplex.*
- (*Several other varieties closely resembling each other occur of this p'ant.*)

ORD. XLVI.—COMPOSITÆ—cont.

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
<i>D.</i>						
xxv. <i>H. gothicum</i> , <i>Fr.</i> .	. . . .	Heaths .	Y.	1-4 ft.	P.	7-8
v. 2. <i>latifolium</i> . . . . * * *	. . . .	. . . .	. .	. .	. .	. .
xxvi. <i>H. tridentatum</i> , <i>Fr.</i>	. . . .	Banks. .	Y.	2-5 ft.	P.	8-9
xxvii. <i>H. prenanthoides</i> , <i>Vill.</i>	. . . .	Riv. banks N. of E.	Y.	1-3 ft.	P.	8-9
xxviii. <i>H. strictum</i> , <i>Fr.</i> .	. . . .	Mount. .	Y.	1-3 ft.	P.	8-9
xxix. <i>H. umbellatum</i> , <i>L.</i> .	. . . .	San. pl. .	Y.	1-4 ft.	P.	7-9
v. 2. <i>filifolium</i>	. . . .	. . . .	. .	. .	. .	. .
xxx. <i>H. crocatum</i> , <i>Fr.</i> .	. . . .	Mount. .	Y.	1-3 ft.	P.	7-9
xxxi. <i>H. rigidum</i> , <i>Fr.</i> . .	. . . .	Mount. .	Y.	3-4 ft.	P.	7-9
xxxii. <i>H. corymbosum</i> , <i>Fr.</i>	. . . .	Sc. mount.	Y.	3-4 ft.	P.	7-9
xxxiii. <i>H. boreale</i> , <i>Fr.</i> . .	. . . .	Banks .	Y.	2-4 ft.	P.	7-9
Gen. XI. <i>Barkhausia</i> , <i>Manch.</i>	<i>Barkhausia.</i>					
i. <i>B. foetida</i> , <i>DC.</i> . . .	<i>stinking B.</i> . . .	Ch. downs	Y.	1-2 ft.	B.	6-7
ii. <i>B. taraxifolia</i> , <i>DC.</i> . .	<i>small rough B.</i> .	Ch. pas. .	Y.	1-2 ft.	P.	6-7
[iii. <i>B. setosa</i> , <i>DC.</i>	. . . .	Clo. fie. .	Y.	1-3 ft.	B.	7-8
Gen. XII. <i>Taraxacum</i> , <i>Fuss.</i>	<i>Dandelion.</i>					
i. <i>T. officinale</i> , <i>Wigg.</i> .	<i>common D.</i> . . .	Mead. . .	Y.	6-12 in.	P.	3-10
v. 1. <i>Dens-leonis</i> , <i>Desf.</i> .	. . . .	. . . .	. .	. .	. .	. .
v. 2. <i>erythrospermum</i> , <i>Andrz.</i>	. . . .	Dry pl. .	. .	. .	. .	. .
v. 3. <i>lævigatum</i> , <i>DC.</i> . . .	. . . .	San. pl. .	. .	. .	. .	. .
v. 4. <i>palustre</i>	. . . .	. . . .	. .	. .	. .	. .
Gen. XIII. <i>Arnoseris</i> , <i>Gaertn.</i>	<i>Nipplewort.</i>					
i. <i>A. pusilla</i> , <i>Gaertn.</i> . .	<i>Dwarf N.</i> . . .	Corn fie.!	Y.	4-12 in.	A.	6-7
Gen. XIV. <i>Lapsana</i> , <i>L.</i>	<i>Nipplewort.</i>					
i. <i>L. communis</i> , <i>L.</i> . . .	<i>common N.</i> . . .	Was. pl. .	Y.	1-2 ft.	A.	7-9

CHARACTERISTICS OF SPECIES AND VARIETIES.

D. Invol. almost glab.—Root ls. generally withering before the plant flowers.

- xxv.** Dark green—St. rigid, leafy, few headed—Rad. ls. ov., lanc.; st. ls. sess., lower ones aggreg.—Invol. ventri.; bracts broad, imbric.—Sty. yellow, with livid hairs.
- v. 2. St. branched above—Ls. broadly lanc, rad. ones generally persistent.*
* * * No radical ls.—Plants forming in autumn buds which the next season become leafy stems—Invol. bracts in many rows.
- xxvi.** St. sometimes forming a small rosette—Ls. lanc., dent. in middle; st. ls. (10-20) gradually diminishing to bracts, three-nerved—Invol. trunc., constricted in fr.—Styles livid.
- xxvii.** Ls. all amplex., loosely veined, hairy on both sides—Heads small, in a spreading panic.; pedunc. short, densely flocc.—Invol. very setose; outer bracts few, obt., short; inner equal, blunt.
- xxviii.** Ls. obl.-lanc., sess., glauc. beneath; upper sess.—Heads irreg. corymb.; pedunc. scarcely setose—Invol. ellip., setose; bracts irreg. imbric.—Sty. with dark hairs.
- xxix.** Ls. lin.-lanc., sess., narrowed below—Heads sub-umbell.—Invol. broadly ovate; bracts reflexed at point.
- v. 2. Ls. very narrow, revolute at margin.*
- xxx.** St. densely leafy—Ls. lin.-lanc., sess., truncate at base—Pedunc. not scaly, with few heads—Invol. sub-cylind.; bracts blunt, adpr.
- xxxi.** St. rigid—Ls. obl.-lanc., sess., rounded at base, glab.—Heads in a branched, leafy, corymb.; pedunc. not scaly—Invol. ovate; bracts obt., adpr.
- xxxii.** St. densely leafy—Ls. ov-lanc., irreg. toothed, sess., loosely veined, glauc. beneath—Heads in a branched spreading panic.; pedunc. scaly.
- xxxiii.** Ls. ovate; upper broad, scarcely clasping; lower sub-peti.—Heads in a corymb.; pedunc. scaly—Invol. blackish gr.—Sty. dark livid.

Gen. XI.

- i. St. round—Buds nodding—Central fruits long beaked.
 ii. St. ang., furrowed—Buds erect—Fruits all equally beaked.
 iii. St. ang., furrowed—Outer scales of invol. lin., spreading.]

Gen. XII.

- i. Ls. toothed, sinu. or pinnfid.
v. 1. Ls. bright green—Outer invol. bracts lin., recurved.
v. 2. Ls. dull green—Outer invol. bracts patent or recurved.
v. 3. Ls. dull green—Outer invol. bracts ovate, erect.
v. 4. Ls. almost ent.—Outer invol. bracts ovate, acum. spreading or adpr.

Gen. XIII.

- i. St. leafless, fistulous upwards—Ls. ob.-ov., obt., toothed.

Gen. XIV.

- i. St. branched, paniced—Ls. lyr.-pinnfid.; upper ls. ent.

ORD. XLVI.—COMPOSITÆ—cont.

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
Gen. XV. Cichorium, L.	<i>Chicory or Succory.</i>					
i. <i>C. Intybus, L.</i>	<i>common S.</i>	Waysides	B.	1-3 ft.	P.	7-10
Gen. XVI. Arctium, L.	<i>Burdock.</i>					
i. <i>A. lappa, L.</i>	<i>common B.</i>	Was. pl. .	Pu.	2-6 ft.	B.	7-8
a. <i>majus, Schk.</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
b. <i>minus, Schk.</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
c. <i>intermedium, Lan.</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
d. <i>nemorosum, Lan.</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
Gen. XVII. Saussurea, DC.	<i>Saussurea.</i>					
i. <i>S. alpina, DC.</i>	<i>Alpine S.</i>	Alp. rocs.!	Pu.	8-12 in.	P.	8-9
Gen. XVIII. Serratula, L.	<i>Saw-wort</i>					
i. <i>S. tinctoria, L.</i>	<i>common S.</i>	Thick. .	Pu.	2-3 ft.	P.	8-9
v. 2. <i>monticola, Bor.</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
Gen. XIX. Carduus, L.	<i>Thistle.</i>					
*						
i. <i>C. nutans, L.</i>	<i>Musk T.</i>	Downs .	Pu.	1-2 ft.	B.	5-10
ii. <i>C. crispus, L.</i>	<i>welted T.</i>	Was. pl. .	Pu.	1-3 ft.	B.	6-8
a. <i>genuinus, Gr.</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
b. <i>polyanthemus, Godr.</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
c. <i>litigiosus, Gr.</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
iii. <i>C. tenuiflorus, Curt.</i>	<i>slender flowered T.</i>	San. pl. .	♂ Pu.	1-3 ft.	B.	6-8
iv. <i>C. Marianus, L.</i>	<i>Milk T.</i>	Was. pl. .	Pu.	1-5 ft.	B.	7-8
* *						
v. <i>C. lanceolatus, L.</i>	<i>Spear T.</i>	Was. pl. .	Pu.	1-5 ft.	B.	7-8
vi. <i>C. eriophorus, L.</i>	<i>woolly-headed T.</i>	Ch. fie. .	♂ Pu.	3-5 ft.	B.	7-8
vii. <i>C. palustris, L.</i>	<i>Marsh T.</i>	Mar. . .	Pu.	2-4 ft.	B.	7-8
viii. <i>C. arvensis, Curt.</i>	<i>creeping T.</i>	Waysides	Pu.	2-4 ft.	P.	7-8
v. 2. <i>setosus.</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
ix. <i>C. pratensis, Huds.</i>	<i>Meadow T.</i>	Wet pas.	Pu.	12-18 in.	P.	6-8
x. <i>C. tuberosus, L.</i>	<i>tuberosus T.</i>	Mead.!	Pu.	12-18 in.	P.	6-8
xi. <i>C. acaulis, L.</i>	<i>dwarf T.</i>	Downs .	Pu.	1-12 in.	P.	7-9
xii. <i>C. heterophyllus, L.</i> . . .	<i>melancholy T.</i> . . .	Moi. pas.	P.	2-3 ft.	P.	7-8
Gen. XX. Onopordium, L.	<i>Cotton Thistle.</i>					
i. <i>O. acanthium, L.</i>	<i>common C.</i>	Was. pl. .	♂ Pu.	2-5 ft.	B.	7-9

CHARACTERISTICS OF SPECIES AND VARIETIES.

Gen. XV.

- i. St. strong, erect, viscid—Lower ls. runc.; upper lanc., amplex.—Heads axill., in pairs.

Gen. XVI.

- i. Ls. large, cord., stalked—Inner invol. bracts with hooked points.
a. Heads long stalked, glob., glab. .
b. Heads shortly stalked, glob., greenish, slightly webbed.
c. Heads sub-sess., ovate, purplish, cottony.
d. Heads sess., glob., very cottony.

Gen. XVII.

- i. Ls. obl.-lanc., toothed, cottony below—Heads in a dense corymb.

Gen. XVIII.

- i. Ls. lyr.-pinnfid., finely serr.—Heads. obl., corymb.
v. 2. *Heads nearly sess.*

Gen. XIX.

* *Pappus rough.*

- i. Heads large, sol., drooping—Invol. bracts lanc., cottony, outer spreading, reflexed.
 ii. Heads small, round, clustered—Invol. bracts lin., erect or asc
a. Ls. hoary beneath—Invol. sub-glob.
b. Ls. pubescent on nerves only—Invol. ovoid.
c. Heads sub-sol.—Ls. as Var. 2—Invol. glob.
 iii. Heads cylind., clustered—Invol. bracts, ov.-lanc. attenuate.
 iv. Heads very large glob., sol.—Invol. bracts leafy, recurved—Ls. veined with white.
 * * *Pappus feathery.*
 v. St. winged—Ls. pinnfid.—Invol. ovate, cottony; invol. bracts lanc., spreading.
 vi. St. not winged—Ls. pinnfid.; lobes bifid.—Heads very large, cottony; invol. bracts lanc., with a long spinous point.
 vii. St. winged—Ls. pinnfid.—Heads close clustered; invol. bracts, ov.-lanc., adpressed.
 viii. St. not winged; rootstock creeping—Ls. pinnfid. invol. ov., glab.; scales broad, lanc., adpr., with a spreading spine.
v. 2. *Ls. lanc. ent. or slightly pinnfid.*
 ix. St. nearly leafless, one-flowered; rootstock fibrous—Ls. lanc., wavy, ciliated—Invol. glob., cottony; bracts adpressed.
 x. St. as last; but rootstock of tuberos fibres—Ls. deeply pinnfid., ciliate—Invol. ovate; bracts adpr., glab.
 xi. St. o. or short; one-flowered—Ls. pinnfid., glab.—Invol. ovoid, glab.; bracts acute, adpr.
 xii. St. branched, cottony—Ls. large, lanc., serr. white beneath—Heads large, sol.; invol. bracts acuminate.

(*Various hybrids occur between several species of this genus.*)

Gen. XX.

- i. St. erect, branched, with spinous wings—Ls. ov.-obl., pinnfid., woolly on both sides.

ORD. XLVI.—COMPOSITÆ—cont.

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
Gen. XXI. Carlina, L.	<i>Carlina Thistle.</i>					
i. <i>C. vulgaris, L.</i> . . .	<i>common C.</i> . . .	Downs .	pu	6-18 in.	B.	6-10
Gen. XXII. Centaurea, L.	<i>Knap-weed, Blue-bottle, Star-thistle.</i>		Br.			
* i. <i>C. nigra, L.</i>	<i>black K.</i>	Mead. .	Pu.	½-3 ft.	P.	6-9
a. <i>vulgaris</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
b. <i>decipiens, Thuil.</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
ii. <i>C. Cyanus, L.</i>	<i>Corn B.</i>	Fie. . .	B.	1-2 ft.	A.	6-8
iii. <i>C. Scabiosa, L.</i>	<i>great K.</i>	Fie. . .	Pu.	2-3 ft.	P.	7-9
* *						
iv. <i>C. aspera, L.</i>	<i>Jersey S.</i>	Jersey ! .	y Br.	1-2 ft.	P.	7-9
v. <i>C. Calcitrapa, L.</i>	<i>common S.</i>	Was. pl.	ro Pu.	1-2 ft.	A.	7-8
vi. <i>C. solstitialis, L.</i>	. . . . .	Fields ! S. of E.	Y.	1-2 ft.	A.	7-9
Gen. XXIII. Bidens, L.	<i>Bur-marigold.</i>					
i. <i>B. cernua, L.</i>	<i>nodding B.</i>	Wet pl. .	g Y.	1-2 ft.	A.	7-10
v. 2. <i>radiata</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
ii. <i>B. tripartita, L.</i>	<i>trifid B.</i>	Wet pl. .	g Y.	1-2 ft.	A.	7-10
Gen. XXIV. Eupatorium, L.	<i>Hemp Agrimony.</i>					
i. <i>E. cannabinum, L.</i>	<i>common H.</i>	Riv. bks.	p Pu.	2-4 ft.	P.	7-9
Gen. XXV. Chrysocoma, L.	<i>Goldilocks.</i>					
i. <i>C. Linosyris, L.</i>	<i>flax-leaved G.</i>	Li. rocks !	Y.	6-18 in.	P.	8-9
Gen. XXVI. Dioitis, Desf.	<i>Cotton weed.</i>					
i. <i>D. maritima, Cass.</i>	<i>Seaside C.</i>	San. sh. ! .	Y.	6-12 in.	P.	8-9
Gen. XXVII. Tanacetum, L.	<i>Tansy.</i>					
i. <i>T. vulgare, L.</i>	<i>common T.</i>	Was. pl. .	Y.	2-3 ft.	P.	8-9
Gen. XXVIII. Artemisia, L.	<i>Worm-wood, Mugwort.</i>					
i. <i>A. campestris, L.</i>	<i>Field W.</i>	San. hths. ! E. of E.	Y.	2-3 ft.	P.	8-9
ii. <i>A. maritima, L.</i>	<i>Sea W.</i>	Sea sh. .	r W.	10-18 in.	P.	8-9
v. 2. <i>gallica, Willd.</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
iii. <i>A. Absinthium, L.</i>	<i>common W.</i>	Was. pl. .	Y.	1-2 ft.	P.	8-9
iv. <i>A. vulgaris, L.</i>	<i>Mugwort</i>	Was. pl. .	br Y.	3-4 ft.	P.	7-9
Gen. XXIX. Antennaria, Gaert.	<i>Everlasting.</i>					
i. <i>A. dioicum, L.</i>	<i>Mountain E., or Cat's-foot.</i>	Mount. hths.	pk W.	3-8 in.	P.	6-7

CHARACTERISTICS OF SPECIES AND VARIETIES.

Gen. XXI.

- i. St. erect, branched—Outer invol. bracts pinnfid. ; inner lin., white.

Gen. XXII.

* *Invol. not spinous.*

- i. Ls. lanc., ent. ; lower lyr.—Invol. bracts ovate, pectinate.
a. Pedunc. stout—Bracts with dark brown appendages.
b. Pedunc. slender—Bracts with pale appendages.
 ii. Ls. lin.-lanc., ent. ; lower ones toothed—Invol. bracts with pectinate margins.
 iii. Ls. pinnfid. ; seg. lanc.—Invol. bracts with ciliate, downy margins.
 * * *Invol. spinous.*
 iv. Heads term., sol.—Ls. lin.—Invol. spines equal.
 v. Heads lat., sess.—Ls. deeply pinnfid.—Central invol. spine long, channelled.
 vi. Heads term., sol.—Ls. lin.-lanc., hoary ; rad. ls. lyr.—Central invol. spine long, slender.]

Gen. XXIII.

- i. Ls. lanc., ent., sess.—Heads drooping.
 v. 2. *Heads rayed.*
 ii. Ls. three-cleft, stalked—Heads sub-erect.

Gen. XXIV.

- i. Ls. opp. 3-5 part ; lfts. lanc., serr.

Gen. XXV.

- i. St. simp., leafy—Ls. lin., ent., glab.—Heads corymb.

Gen. XXVI.

- i. Densely toment.—Ls. sess., obt.—Heads corymb.

Gen. XXVII.

- i. Ls. bipinnfid. ; seg. inciso.-serr.—Fls. in a term. corymb.

Gen. XXVIII.

- i. St. twiggy—Ls. bipinn. ; seg. lin., silky beneath—Heads glab., drooping, outer—flowers fertile.
 ii. Ls. woolly, pinnfid. ; seg. lin., obt.—Heads drooping—Fls. few, all perfect.
 v. 2. *Heads dense, erect.*
 iii. Ls. silky, pinnfid. ; seg. lanc., obt.—Heads drooping, silky ; outer flowers fert.
 iv. Ls. woollv, pinnfid. ; seg. lan., acute, serr.—Heads erect, woolly ; fls. all perfect.

Gen. XXIX.

- i. St. herb., procumb.—Ls. spath., glab. above, cottony below.

ORD. XLVI.—COMPOSITÆ—*cont.*

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
v. 2. <i>hyperboreum</i>						
[ii. <i>A. margaritaceum</i> , <i>L.</i>	<i>pearly E.</i>	Moi. mead. W. of E.!	br W.	2-3 ft.	P.	8-9
Gen. XXX. Gnaphalium, <i>L.</i> <i>Cudweed.</i>						
i. <i>G. luteo-album</i> , <i>L.</i> . . .	<i>Jersey C.</i>	San. fie.!	♂ Y.	6-12 in.	A.	7-8
ii. <i>G. sylvaticum</i> , <i>L.</i> . . .	<i>Highland C.</i> . . .	Copses .	Br.	2-12 in.	P.	7-9
v. 2. <i>norvegicum</i> , <i>Gunn.</i> . .		Sc. mount.				
iii. <i>G. supinum</i> , <i>L.</i>	<i>dwarf C.</i>	Alp. rocks	Br.	1-3 in.	P.	7-8
iv. <i>G. uliginosum</i> , <i>L.</i> . . .	<i>marsh C.</i>	Wet pl. .	♂ Br.	2-6 in.	A.	7-9
v. 2. <i>pilulare</i> , <i>Wahl.</i> . . .	. . . . .	. . . . .	. . .	. . .	. . .	. . .
Gen. XXXI. Filago, <i>L.</i> <i>Filago.</i>						
[i. <i>F. gallica</i> , <i>L.</i>	<i>narrow-leaved F.</i>	San. fie.!	Y.	3-6 in.	A.	7-9
ii. <i>F. minima</i> , <i>Fries.</i> . . .	<i>least F.</i>	Dry pl. .	Y.	4-6 in.	A.	6-9
iii. <i>F. germanica</i> , <i>L.</i> . . .	<i>common F.</i>	San. pl. .	♂ Br.	6-18 in.	A.	7-9
a. <i>canescens</i> , <i>Ford.</i>	. . . . .	. . . . .	. . .	. . .	. . .	. . .
b. <i>apiculata</i> , <i>C. E. Sm.</i> . .	. . . . .	. . . . .	. . .	. . .	. . .	. . .
c. <i>spathulata</i> , <i>Presl.</i> . . .	. . . . .	. . . . .	. . .	. . .	. . .	. . .
Gen. XXXII. Petasites, <i>Gaert.</i> <i>Butter-bur.</i>						
i. <i>P. vulgaris</i> , <i>Desf.</i> . . .	<i>common B.</i>	Wet fie. .	Pk.	4-12 in.	P.	3-5
Gen. XXXIII. Tussilago, <i>L.</i> <i>Colt's-foot.</i>						
i. <i>T. Farfara</i> , <i>L.</i>	<i>common C.</i>	Moi. pl. .	Y.	4-10 in.	P.	3-5
Gen. XXXIV. Erigeron, <i>L.</i> <i>Flea-bane.</i>						
i. <i>E. alpinus</i> , <i>L.</i>	<i>Alpine F.</i>	Alp. rocks	Pu.	6-8 in.	P.	7-8
ii. <i>E. acris</i> , <i>L.</i>	<i>blue F.</i>	Dry past.	Pu.	1-1½ in.	B.	7-8
[iii. <i>E. canadensis</i> , <i>L.</i> . . .	<i>Canada F.</i>	Was. pl..	♂ Y.	1-2 ft.	A.	8-9
Gen. XXXV. Aster, <i>L.</i> <i>Star-wort, Mi- chaelmas Daisy.</i>						
i. <i>A. Tripolium</i> , <i>L.</i>	<i>Sea S. or M.</i> . . .	Salt mar.	Pu.	2-3 ft.	P.	8-9
v. 2. <i>discoideus</i>	. . . . .	. . . . .	. . .	. . .	. . .	. . .
Gen. XXXVI. Solidago, <i>L.</i> <i>Golden-rod.</i>						
i. <i>S. Virgaurea</i> , <i>L.</i>	<i>common G.</i>	Thick. .	Y.	6-24 in.	P.	7-9
v. 2. <i>angustifolia</i> , <i>Gaud.</i> . .	. . . . .	. . . . .	. . .	. . .	. . .	. . .
v. 3. <i>canadica</i> , <i>Huds.</i> . . .	. . . . .	Mount. .	. . .	. . .	. . .	. . .

CHARACTERISTICS OF SPECIES AND VARIETIES.

- v. 2. *Ls. cottony on both sides (I. of Skye).*
 ii. St. erect, shrubby—*Ls. lin.-lanc.*]

Gen. XXX.

- i. St. simp., herb.—Heads in dense, leafless corymbs.
 ii. St. simp., woolly—Heads axill. and term., forming a leafy rac.
 v. 2. *Ls. broader—Spike dense, term.*
 iii. St. tufted, one or two-headed—Heads sess., erect.
 iv. St. spreading, much branched—Heads in term., crowded tufts.
 v. 2. *Fr. papillose (Toft. Camb.).*

Gen. XXXI.

- i. St. erect—*Ls. lin., revolute*—Heads axill., shorter than the ls.]
 ii. St. branched—*Ls. lin., flat*—Heads term. and axill., longer than the ls.
 iii. St. erect, proliferous at summit—*Ls. lin., obt., waved*—Heads five-angled in term., glab. heads.
 a. Heads obscurely five-angled—Tips of invol. bracts yell.
 b. Taller—*Ls. broader*—Heads acutely five-angled, overtopped by blunt ls.—Tips of invol. bracts red.
 c. Short—*Ls. spath.*—Heads overtopped by acute ls.—Tips of invol. bracts pale.

Gen. XXXII.

- i. *Ls. very large, orb., cord., toothed, glab. above*—Fls. in a dense spike, appearing before the ls.

Gen. XXXIII.

- i. *Ls. cord., ang., downy beneath*—St. single-flowered.

Gen. XXXIV.

- i. *Ls. mostly rad.*—St. one or two-headed—Rays (purple) much longer than disk.
 ii. St. leafy, branched—Florets of the ray purplish, scarcely longer than disk.
 iii. St. leafy, branched—Florets of the ray yellowish.]

Gen. XXXV.

- i. *Ls. spath., fleshy*—Heads corymb., ray pu. ; disk yell.
 v. 2. *Ray wanting.*

Gen. XXXVI.

- i. *Ls. ellip.-lanc., upper narrower*—Fls. in paniced racemes.
 v. 2. *Tall—Ls. all lanc.*
 v. 3. *Small—Rad. ls. broader—Fls. larger.*

ORD. XLVI.—COMPOSITÆ—cont.

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
Gen. XXXVII. Senecio, <i>L.</i>	<i>Groundsel, Rag- wort.</i>					
*						
i. <i>S. vulgaris, L.</i>	<i>common G.</i>	Was. pl..	Y.	6-12 in.	A.	1-12
ii. <i>S. sylvaticus, L.</i>	<i>Mountain G.</i>	Dry pas.	Y.	½-3 ft.	A.	7-9
iii. <i>S. viscosus, L.</i>	<i>stinking G.</i>	Was. pl..	Y.	1-2 ft.	A.	7-9
* *						
<i>A.</i>						
iv. <i>S. squalidus, L.</i>	<i>inelegant R.</i>	Old walls!	Y.	8-12 in.	A.	6-10
v. <i>S. erucifolius, L.</i>	<i>hoary R.</i>	Waysides	Y.	1-4 ft.	P.	7-8
vi. <i>S. Jacobœa, L.</i>	<i>common R.</i>	Waysides	Y.	1-4 ft.	P.	7-8
vii. <i>S. aquaticus, Huds.</i>	<i>Marsh R.</i>	Wet pl. .	Y.	1-4 ft.	P.	7-8
v. 2. <i>pinnatifidus</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
<i>B.</i>						
viii. <i>S. paludosus, L.</i>	<i>great Fen R.</i>	Fens! E. of E.	Y.	3-6 ft.	P.	6-7
[ix. <i>S. saracenicus, L.</i>	<i>broad-leaved R.</i>	Moist mead.!	Y.	3-5 ft.	P.	6-7
Gen. XXXVIII. Cinera- ria, L.	<i>Flea-wort.</i>					
i. <i>C. palustris, L.</i>	<i>Marsh F.</i>	Fens! E. of E.	Y.	2-3 ft.	P.	6-7
ii. <i>C. campestris, Retz.</i>	<i>Field F.</i>	Ch. dwns.!	Y.	4-12 in.	P.	5-6
v. 2. <i>maritima, Sy.</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
Gen. XXXIX. Doronicum, <i>L.</i>	<i>Leopard's-bane.</i>					
[i. <i>D. Pardalianches, L.</i>	<i>great L.</i>	Dp. wds.!	Y.	2-3 ft.	P.	5-7
ii. <i>D. plantagineum, L.</i>	<i>plaintain-leaved L.</i>	Dp. wds.!	Y.	2-3 ft.	P.	5-7
Gen. XL. Inula, L.	<i>Inula, &c.</i>					
i. <i>I. Helenium, L.</i>	<i>Elecampane</i>	Moi. mead.!	Y.	2-5 ft.	P.	7-8
ii. <i>I. Conyza, DC.</i>	<i>Ploughman's Spikenard.</i>	Ch. pas..	Y.	2-5 ft.	B.	7-9
iii. <i>I. salicina, L.</i>	. . . . .	I.!	y W.	12-18 in.	P.	7-9
iv. <i>I. crithmoides, L.</i>	<i>golden Samphire</i>	Sea sh.!	Y.	6-18 in.	P.	7-8
v. <i>I. dysenterica, L.</i>	<i>Flea-bane</i>	SW. of E.	Y.	6-12 in.	P.	7-9
vi. <i>I. Pulicaria, L.</i>	<i>small F.</i>	Moi. pl. .	Y.	6-12 in.	A.	8-9
Gen. XLI. Bellis, L.	<i>Daisy.</i>					
i. <i>B. perennis, L.</i>	<i>common D.</i>	Past. . .	W.	2-5 in.	P.	1-12
Gen. XLII. Chrysanthem- um, L.	<i>Ox-eye.</i>					
i. <i>C. segetum, L.</i>	<i>yellow O., or Corn Marigold.</i>	Corn fie.	Y.	12-18 in.	A.	6-9
ii. <i>C. Leucanthemum, L.</i>	<i>great white O.</i>	Dry pas.	W.	1-2 ft.	P.	6-8

CHARACTERISTICS OF SPECIES AND VARIETIES.

Gen. XXXVII.

* *Florets of the ray revolute or wanting.*

- i. Glab.—Ray florets wanting.
- ii. Slightly viscid—Ray floret revolute—Outer invol. bracts very short.
- iii. Viscid and hairy—Ray florets revolute—Outer invol. bracts half as long as inner.
* * *Florets of the ray spreading.*
A. Ls. pinnfid. or lyrate.
- iv. Ls. pinnfid., glab. ; seg. lin.—Outer invol. bracts few, small, very short.]
- v. St. cottony—Ls. pinnfid., downy beneath—Outer invol. bracts half as long as inner
—Fr. all hairy.
- vi. St. smooth—Ls. lyr.-pinnfid., glab.—Fr. of the ray glab.
- vii. Root ls. ent. ; upper lyr., serr.—Fr. all glab.
- v. 2. *Ls. all pinnfid.*
B. Leaves entire.
- viii. St. hollow—Ls. lanc., sharply serr., woolly beneath.
- ix. St. solid—Ls. lanc., irregularly serr., glab.]

Gen. XXXVIII.

- i. St. branched—Ls. lanc., semi-amplex. ; lower sinu. dent.
- ii. St. simple—Rad. ls. oval, ent. ; st. ls. lanc.—Heads corymb.
- v. 2. *Larger—Heads many, umbellate—Lower ls. toothed (Holyhead).*

Gen. XXXIX.

- i. Rad. ls. cord. ; st. ls. with clasping aur.—Heads 3-5.]
- ii. Rad. ls. ovate ; st. ls. sess.—Heads sol.]

Gen. XL.

- i. Ls. large, cord.-ov., unequally toothed, wrinkled—Heads very large ; invol. bracts leafy, ovate, reflexed.
- ii. Ls. oval-lanc., pubesc., serr. ; upper ent.—Invol. bracts lin., reflexed.
- iii. Ls. lin., obl., toothed ; upper sess.—Heads sol.—(Shores of Lough Derg, Galway).
- iv. Ls. lin., fleshy, pointed—Invol. bracts lin., adpressed.
- v. St. woolly—Ls. obl. cord.—Heads corymb. ; ray much longer than disk.
- vi. Ls. lanc., wavy, base narrow—Heads term. and axill. ; ray scarcely longer than disk.

Gen. XLI.

- i. Ls. ob.-ov., glab.—St. one-flowered.

Gen. XLII.

- i. Ray yellow—Invol. bracts with broad membr. margins.
- ii. Ray white—Invol. bracts with narrow membr. margins.

ORD. XLVI.—COMPOSITÆ—*cont.*

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
Gen. XLIII. <i>Matricaria</i>, L.	<i>Wild Chamomile, Feverfew.</i>					
i. <i>M. Parthenium</i> , L. . .	<i>common F.</i> . . .	Was. pl..	W.	1-2 ft.	P.	7-9
ii. <i>M. inodorum</i> , L. . .	<i>Corn F., or scent- less Mayweed.</i>	Was. pl..	W.	1-2 ft.	A.	6-10
v. 2. <i>maritimum</i> , Pers. . .		Sea sh. .	W.	12-18 in.	A.	6-8
iii. <i>M. Chamomilla</i> , L. . .	<i>wild C.</i>	Was. pl..	W.	12-18 in.	A.	6-8
Gen. XLIV. <i>Anthemis</i>, L.	<i>Chamomile.</i>					
i. <i>A. nobilis</i> , L.	<i>common C.</i>	Downs .	W.	6-12 in.	P.	7-9
ii. <i>A. arvensis</i> , L. . . .	<i>Corn C.</i>	Corn fie.	W.	1-2 ft.	A.	6-8
v. 2. <i>anglica</i> , Spr.			W.	8-16 in.	A.	6-9
iii. <i>A. Cotula</i> , L.	<i>stinking C.</i>	Corn fie.	W.	8-16 in.	A.	6-9
Gen. XLV. <i>Achillea</i>, L.	<i>Yarrow, Milfoil.</i>					
i. <i>A. Ptarmica</i> , L.	<i>Sneeze-wort.</i> . . .	Mead. . .	W.	1-2 ft.	P.	7-8
ii. <i>A. Millefolium</i> , L. . .	<i>common Y., or M.</i>	Waysides	W. or Pk.	6-18 in.	P.	6-9
Gen. XLVI. <i>Galinsoga</i>, Ruiz.						
[i. <i>G. parviflora</i> , Cav. . . .	. . . . .	Waysides, Kew.	Y.	1-2 ft.	A.	8-10

ORD. XLVII.—CAMPANULACEÆ.

Gen. I. <i>Campanula</i>, L.	<i>Bell-flower.</i>					
i. <i>C. rotundifolia</i> , L. . . .	<i>round-leaved C., or Hair-bell.</i>	Heaths .	B.	1-2 ft.	P.	7-9
v. 2. <i>montana</i> , Sy.			W.	1-3 ft.	A.	7-9
ii. <i>C. patula</i> , L.	<i>spreading B.</i> . . .	Copses !.	B.	1-3 ft.	A.	7-9
iii. <i>C. Rapunculus</i> , L. . . .	<i>Rampion B.</i>	Grav. waysides.	B.	2-3 ft.	P.	7-8
iv. <i>C. latifolia</i> , L.	<i>Giant B.</i>	Moi. wds.	B. or W.	3-5 ft.	P.	7-8
v. <i>C. rapunculoides</i> , L. . .	<i>creeping B.</i>	Woods .	B.	1-2 ft.	P.	7-8
vi. <i>C. Trachelium</i> , L. . . .	<i>nettle-leaved B.</i> . .	Woods .	B.	1-3 ft.	P.	7-9
vii. <i>C. glomerata</i> , L.	<i>clustered B.</i>	Downs .	B.	6-18 in.	B.	7-8
[viii. <i>C. persicifolia</i> , L. . . .	<i>Peach-leaved B.</i> . .	Woods ! Cullen.	B.	1-2 ft.	P.	7
ix. <i>C. hederacea</i> , L.	<i>Ivy-leaved B.</i>	Moi. wds.	β B.	Cr.	P.	7-8
x. <i>C. hybrida</i> , L.	<i>Corn B.</i>	Cha. fie..	B.	6-10 in.	A.	7-9

ORD. XLVI.—COMPOSITÆ—*cont.*

CHARACTERISTICS OF SPECIES AND VARIETIES.

Gen. XLIII.

- i. Ls. pinn., stalked ; seg. ovate, lobed—Heads corymb.
  - ii. Ls. pinnfid., sess. ; seg. capill.—Heads sol.
- v. 2. *Seg. of ls. fleshy.*
- iii. Ls. bipinn. ; seg. lin.—Heads sol., long-stalked—Recep. con., hollow.

Gen. XLIV.

- i. St. procumb.—Ls. bipinn. ; seg. lin., slightly downy—Invol. bracts blunt—Aromatic.
  - ii. St. erect—Ls. bi-pinnfid., hoary—Invol. bracts acute—Recep. con.
- v. 2. *Ls. fleshy—Recep. flat—(Sea coast, Northumb.).*
- iii. St. erect, much branched—Ls. bipinnfid., glab. ; seg. subul.—Invol. bracts lin., setaceous—Fœtid.

Gen. XLV.

- i. Ls. lin.-lanc., serr., glab.
- ii. Ls. bipinnfid., woolly.

Gen. XLVI.

- i. St. slender—Ls. ov., bluntly serr.—Fls. in dichot cymes.]

ORD. XLVII.—CAMPANULACEÆ.

Gen. I.

- i. St. slender, glab.—Rad. ls. cord. ; st. ls., lin. or lanc.—Fls. rac., drooping.
- v. 2. *Lower st. ls. obl.—Fls. sol.*
- ii. St. ang., scabr.—Rad. ls. ob.-ov. ; st. ls. lin.—Fls. pani., erect, long stalked—Cal seg. toothed at base.
  - iii. Hispid—St. angled—St. ls. obl.-lanc.—Fls. pani., erect—Cal. seg. narrow ent.
  - iv. St. simple, furrowed, leafy—Ls. ov.-lanc., doubly serr., lower stalked, upper sess.—Pedunc. one-flowered—Cal. seg. lin.
  - v. St. slightly branched—Ls. scabr., unequally serr., lower long-stalked, upper lanc., sess.—Fls. unilat.—Cal. seg. lin.
  - vi. St. angled, rough—Ls. coarsely serr. ; lower cord., long stalked ; upper lanc., acum., nearly sess.—Pedunc. 2-3 flowered—Cal. segm. lanc., rough.
  - vii. St. simp., ang.—Ls. scab., obl.-lanc. ; st. ls. semi-amplex.—Fls. erect, sess., in term. and axill. clusters.
  - viii. Glab.—St. round, few-flowered—Rad. ls. ob.-ov. stalked ; st. ls. lin., sess.—Cal. seg. lanc., ent.]
  - ix. St. fili.—Ls. alt., stalked, cord., lobed, glab.—Pedunc. sol.
  - x. St. erect, hispid.—Ls. obl., waved—Fls. sess. axill.—Caps fusi., angled.

ORD. XLVII.—CAMPANULACEÆ—*cont.*

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
Gen. II. <i>Phyteuma</i>, L.	<i>Rampion.</i>					
i. <i>P. orbiculare</i> , L. . . .	<i>round-leaved R.</i> . .	Ch. downs!	B.	6-18 in.	P.	7-8
ii. <i>P. spicatum</i> , L. . . .	<i>spiked R.</i>	Thick!	Y.	2-3 ft.	P.	6-7
Gen. III. <i>Jasione</i>, L.	<i>Sheep's-bit.</i>					
i. <i>J. montana</i> , L. . . .	<i>annual S.</i>	Heaths .	B.	6-10 in.	A.	6-9
Gen. IV. <i>Lobelia</i>, L.	<i>Lobelia.</i>					
i. <i>L. urens</i> , L.	<i>acrid L.</i>	Heaths!	Pu.	12-18 in.	P.	8-9
ii. <i>L. Dortmannia</i> , L. . .	<i>Water L.</i>	Lakes .	♂ B.	1-2 ft.	P.	7-8

ORD. XLVIII.—ERICACEÆ.

Gen. I. <i>Erica</i>, L.	<i>Heath.</i>					
*						
i. <i>E. Tetralix</i> , L. . . .	<i>cross-leaved H.</i> . .	Heaths .	Ro.	12-18 in.	Shb.	7-8
v. 2. <i>Mackiana</i> , Bab. . .	. . . .	. . . .	. .	. .	. .	. .
v. 3. <i>Watsoni</i> , Benth. . .	. . . .	. . . .	. .	. .	. .	. .
ii. <i>E. ciliaris</i> , L.	<i>ciliate H.</i>	Heaths! S.W. of E.	Cr.	12-18 in.	Shb.	6-9
iii. <i>E. cinerea</i> , L.	<i>fine-leaved H.</i> . .	Heaths .	Cr.	1-2 ft.	Shb.	7-9
* *						
iv. <i>E. hybernica</i> , Sy. . .	. . . .	Boggy heaths! I.	Pk.	2-5 ft.	Shb.	4-5
v. <i>E. vagans</i> , L.	<i>Cornish H.</i>	Heaths! Corn.	Cr.	1-3 ft.	Shb.	7-8
Gen. II. <i>Calluna</i>, Salisb.	<i>Ling.</i>					
i. <i>C. vulgaris</i> , Salisb. . .	<i>common L.</i>	Heaths .	Ro. or W.	1-2 ft.	Shb.	7-9
a. <i>glabrata</i>	. . . .	. . . .	. .	. .	. .	. .
b. <i>incana</i>	. . . .	. . . .	. .	. .	. .	. .
Gen. III. <i>Menziesia</i>, Sm.	<i>Menziesia.</i>					
i. <i>M. cærulea</i> , Swtz. . .	<i>Scottish M.</i>	Sow of Athol.	Li.	5-10 in.	Shb.	6-7
ii. <i>M. polifolia</i> , Fuss. . .	<i>Irish M., or St. Dabeocs H.</i>	Moist heaths, I.	Pu.	1-2 ft.	Shb.	7-8
Gen. IV. <i>Loiseleuria</i>, Desv.	<i>Azalea.</i>					
i. <i>L. procumbens</i> , Desv.	<i>trailing A.</i>	Sc. mount.!	Pk.	2-3 in.	Shb.	5-6
Gen. V. <i>Andromeda</i>, L.	<i>Andromeda.</i>					
i. <i>A. polifolia</i> , L. . . .	<i>Marsh A.</i>	Peat bogs!	Pk.	6-12 in.	Shb.	6-8

ORD. XLVII.—CAMPANULACEÆ—*cont.*

CHARACTERISTICS OF SPECIES AND VARIETIES.

Gen. II.

- i. Heads of fls. glob., blue ; of fr. oval.
- ii. Heads of fls. obl., yellowish ; of fr. elongate (Sussex).

Gen. III.

- i. Ls. obl., wavy, blunt—Fls. small in term., hemisph., heads.

Gen. IV.

- i. St. erect, leafy—Rad. ls. ob.-ov. ; upper lan., sess. (Axminster).
- ii. St. simple—Ls. all rad., submerged, cylind., of two parallel tubes.

ORD. XLVIII.—ERICACEÆ.

Gen. I.

* *Anthers included.*

- i. Ls. four in a whorl, lanc., downy above—Fls. sub-umbell. ; sep. lin., downy—Ovary downy.
- v. 2. *Ls. glab. above—Sep. ov.-lanc., glab.—Ovary glab.—(Galway.)*
- v. 3. *Like E. Tetralix, but coroll. ventri. and fls. racemed.—(Truro.)*
- ii. Ls. four in a whorl, ov., ciliate—Fls. in a term., unilat. rac.

- iii. Ls. three in a whorl, lin., keeled, glab.—Fls. in a dense, whorled rac.

* * *Anthers exserted.*

- iv. Ls. four in a whorl, lin., glab.—Fls. in a dense rac.—Cor. urc. ; anth. half exserted.
- v. Ls. 4-5 in a whorl, lin., glab.—Fls. axill., crowded—Cor. short, campan.

Gen. II.

- i. Ls. small, opp., imbri. in four rows—Fls. small, drooping, shortly stalked.

a. Ls. glabrous.

b. Ls. hoary.

(*The two forms are connected by intermediate varieties.*)

Gen. III.

- i. Ls. lin., toothed—Fls. few, term., aggreg. ; cor. five-cleft.
- ii. Ls. oval, cili., downy beneath—Fls. in term., leafy rac. ; cor. four-cleft.

Gen. IV.

- i. St. procumb., woody—Ls. opp., lin.-obl., revol.—Fls. term., aggreg.

Gen. V.

- i. Ls. alt., lan., glab., glauc. beneath, margins revol.—Fls. in short, term. rac.

ORD. XLVIII.—ERICACEÆ—*cont.*

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
Gen. VI. Arbutus, L.	<i>Strawberry Tree.</i>					
i. <i>A. Unedo, L.</i>	<i>austere S.</i>	Woods, I.!	Cr.	8-10 ft.	Shb.	9-10
Gen. VII. Arctostaphylos, Adanson.	<i>Bear-berry.</i>					
i. <i>A. alpina, Spreng.</i> . .	<i>black B.</i>	Sc. mount.	W.	3-4 in.	Shb.	6-8
ii. <i>A. Uva-Ursi, Spreng.</i>	<i>red B.</i>	Sc. mount.	Pk.	4-6 in.	Shb.	5-6
Gen. VIII. Vaccinium, L.	<i>Bilberry, Whortleberry, &c.</i>					
i. <i>V. Myrtillus, L.</i> . . .	<i>B. or W.</i>	Heaths .	Ro.	6-18 in.	Shb.	4-6
ii. <i>V. uliginosum, L.</i> . .	<i>great B. or bog W.</i>	Mount. bogs!	♂Pk.	6-10 in.	Shb.	5-6
iii. <i>V. Vitis-Idæa, L.</i> . .	<i>red W. or Cowberry.</i>	Mount. heaths.	♂Pk.	6-18 in.	Shb.	6-8
iv. <i>V. Oxycoccus, L.</i> . .	<i>Cranberry.</i> . . .	Peat bogs	R.	Cr	Shb.	7-8
Gen. IX. Pyrola, L.	<i>Winter-Green.</i>					
i. <i>P. rotundifolia, L.</i> . .	<i>round-leaved W.</i>	Moi. wds.!	W.	8-12 in.	P.	7-9
v. 2. <i>arenaria, Koch.</i> . .	. . . .	San. sh. Lancas.	. .	. .	. .	. .
ii. <i>P. media, Swartz.</i> . .	<i>intermediate W.</i>	Woods! .	W.	8-12 in.	P.	6-7
iii. <i>P. minor, L.</i>	<i>lesser W.</i>	Woods .	W.	8-12 in.	P.	6-7
iv. <i>P. secunda, L.</i>	<i>serrate W.</i>	Mount. woods!	g W.	2-5 in.	P.	7-8
v. <i>P. uniflora, L.</i>	<i>large-flowered W., or Moneses.</i>	Sco. fir. woods!	W.	1-3 in.	P.	7-8
Gen. X. Monotropa, L.	<i>Bird's-nest.</i>					
i. <i>M. Hypopitys, L.</i> . .	<i>yellow B.</i>	Beech woods!	Y.	6-12 in.	P.	7-8
a. <i>glabra, Roth.</i>	. . . .	. . . .	. .	. .	. .	. .
b. <i>hirsuta, Roth.</i>	. . . .	. . . .	. .	. .	. .	. .

ORD. XLIX.—AQUIFOLIACEÆ.

Gen. I. Ilex, L.	<i>Holly.</i>					
i. <i>I. Aquifolium, L.</i> . .	<i>common H.</i>	Woods .	W.	10-30ft.	T.	5-8

ORD. L.—OLEACEÆ.

Gen. I. Ligustrum, L.	<i>Privet.</i>					
i. <i>L. vulgare, L.</i>	<i>common P.</i>	Thick. .	W.	4-10 ft.	Shb.	6-7

ORD. XLVIII.—ERICACEÆ—*cont.*

CHARACTERISTICS OF SPECIES AND VARIETIES.

Gen. VI.

- i. St. rough—Ls. ellip.-lanc., sess.—Fls. pend.—Fr. tubercled.

Gen. VII.

- i. Ls. wrinkled, serr.—Fr. black.
 ii. Ls. ob.-ov., everg., ent.—Fr. red.

Gen. VIII.

* *Ls. deciduous.*

- i. St. erect, ang.—Ls. ov., serr.
 ii. St. procumb., round—Ls. ob.-ov., ent., glaucous beneath.
 * * *Ls. evergreen.*
 iii. St. woody, procumb.—Ls. ob.-ov., dotted beneath; margins slightly revol.—Cor. campan.
 iv. St. fili. cr.—Ls. ov.-obl., ent., glauc. beneath; marg. very revol.—Cor. rotate, seg. reflexed.

Gen. IX.

* *Fls. racemose.*

- i. Sta. erect, shorter than the decurved sty. which is bent at the extremity.
 v. 2. *Ls. small—Scape with bracts.*
 ii. Sta. incurved, shorter than the decurved, straight style.
 iii. Sta. incurved as long as the short, straight style.
 iv. Sta. shorter than the long, ascending style—Ls. serr.—Fls. unilat.
 * * *Fls. sol.*
 v. Ls. few, round, serr.—Fls. large, drooping.

Gen. X.

- i. St. scaly, succ.—Fls. rac.
 a. Fls. glab.
 b. Inside of pet., sep., filaments and ovary hairy.

ORD. XLIX.—AQUIFOLIACEÆ.

Gen. I.

- i. Ls. spiny, ovate, evergreen.

ORD. L.—OLEACEÆ.

Gen. I.

- i. Ls. ellip.-lanc., ent., glab., opp.—Fls. panicled.

ORD. L.—OLEACEÆ—*cont.*

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering
Gen. II. Fraxinus, L.	<i>Ash.</i>					
i. <i>F. excelsior, L.</i>	<i>common A.</i>	Woods .	..	..	T.	4-5
v. 2. <i>heterophyllus, Vahl.</i>	. . . .	. . . .	..	..	..	..

ORD. LI.—APOCYNACEÆ.

Gen. I. Vinca, L.	<i>Periwinkle.</i>					
i. <i>V. minor, L.</i>	<i>lesser P.</i>	Thick. .	B.	1-2 ft.	P.	4-5
ii. <i>V. major, L.</i>	<i>great P.</i>	Thick. .	B.	6-12 in.	P.	4-5

ORD. LII.—GENTIANACEÆ.

Gen. I. Gentiana, L.	<i>Gentian.</i>					
*						
i. <i>G. verna, L.</i>	<i>Spring G.</i>	Alp. pas.!	B.	1-2 in.	P.	4-6
ii. <i>G. Pneumonanthe, L.</i>	<i>Marsh G.</i>	Moi. hths.!	B.	4-10 in.	P.	8-9
iii. <i>G. nivalis, L.</i>	<i>small alpine G.</i> .	Sc.mount.!	B.	1-6 in.	A.	7-8
* *						
iv. <i>G. Amarella, L.</i>	<i>small-flowered G.</i>	Dry hths.	♂ Pu.	3-12 in.	A.	8-9
v. 2. <i>germanica, Willd.</i> . .					..	..
v. <i>G. campestris, L.</i> . . .	<i>Field G.</i>	Cha. hills	♂ Pu.	3-10 in.	A.	8-9
Gen. II. Cicendia, Adams.	<i>Cicendia.</i>					
i. <i>C. filiformis, Reich.</i> .	<i>least C.</i>	San. bogs! S.W. of E.	Y.	4-8 in.	A.	7-10
ii. <i>C. pusilla, Griseb.</i> . .	. . . .	San. com.! Guernsey.	Pk.	1-2 in.	A.	7-9
Gen. III. Erythrœa, Ren.	<i>Centaury.</i>					
*						
i. <i>E. Centaurium, Pers.</i>	<i>common C.</i>	Dry pas.	Ro.	6-18 in.	A.	6-9
v. 2. <i>latifolia, E. B. S.</i> . .	. . . .	. . . .	..	..	..	..
ii. <i>E. pulchella, Fries.</i> . .	<i>dwarf C.</i>	San. sh. .	Ro.	1-4 in.	A.	6-9
* *						
iii. <i>E. littoralis, Fries.</i> . .	<i>tufted C.</i>	San. sh.!	Ro.	2-6 in.	A.	6-9
iv. <i>E. latifolia, Sm.</i>	<i>broad-leaved C.</i> .	San. sh.!	Ro.	2-6 in.	A.	6-9
Gen. IV. Chlora, L.	<i>Yellow-wort.</i>					
i. <i>C. perfoliata, L.</i>	<i>perfoliate Y.</i> . .	Ch. pas..	Y.	6-18 in.	A.	6-9
Gen. V. Villarsia, Vent.	<i>Villarsia.</i>					
i. <i>V. nymphæoides, Vent.</i>	<i>Nymphæa-like V.</i>	Ponds! .	Y.	Fl.	P.	7-8

ORD. L.—OLEACEÆ—*cont.*

CHARACTERISTICS OF SPECIES AND VARIETIES.

Gen. II.

- i. Ls. pinn. ; lfts. obl.-lanc., serr., sess.—Sep. and pet. wanting.
v. 2. *Ls. simp.*

ORD. LI.—APOCYNACEÆ.

Gen. I.

- i. Ls. ellip.-lanc. ; margins glab.
 ii. Ls. ov., cord. ; margins ciliate.]

ORD. LII.—GENTIANACEÆ.

Gen. I.

* *Throat of calyx naked.*

- i. St tufted, one-flowered—Ls. ov.—Cor. salver-shaped, with five large and five small, bifid. segm.
 ii. St. erect—Ls. lin., obt.—Fls. term. and axill. ; cor. large, bell-shaped, five-cleft.
 iii. St. branched—Ls. ellip.—Fls. sol. ; cor. funnel-shaped, with five large and five small, bifid. seg.
 * * *Throat of calyx fringed.*
 iv. Corolla 4-5 cleft—Cal. five-cleft, seg. equal.
v. 2. *Larger—Fls. ob.-conical—Cal. lobes unequal, shorter than cor. tube.*
 v. Corolla four-cleft—Cal. four-cleft ; two outer seg. large, ovate.

Gen. II.

- i. St. slend., forked—Fls. yell.—Cal. camp.
 ii. St. much branched—Fls. pink—Cal. four-part ; seg. lin., adpr.

Gen. III.

* *Cal. about half as long as tube of opening corolla.*

- i. St. erect, simple below—Ls. ellip.-obl.—Fls. corym., paniced.
v. 2. *A stunted broad-leaved variety.*
 ii. St. much branched—Ls. ov.-obl.—Fls. term. and axill.
 * * *Cal. as long as tube of opening corolla.*
 iii. St. simp.—Ls. obl.-lin., lower ones spath.—Fls. corymb.
 iv. St. branched—Ls. broadly ellip.—Fls. in dense, forked tufts.

Gen. IV.

- i. Glau.—St. simp.—Ls. ov., conn.

Gen. V.

- i. Ls. flo., heart-shaped—Pedunc. one-flowered—Cor. cili.

ORD. LII.—GENTIANACEÆ—*cont.*

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
Gen. VI. Menyanthes, L.	<i>Buckbean.</i>					
i. <i>M. trifoliata, L. . . .</i>	<i>B. or Marsh Trefoil.</i>	Bogs . .	W.	3-8 in.	P.	5-7

ORD. LIII.—POLEMONIACEÆ.

Gen. I. Polemonium, L.	<i>Jacob's Ladder.</i>					
i. <i>P. cœruleum, L. . . .</i>	<i>blue J. . . . .</i>	Copses !.	B.	1-3 ft.	P.	6-7

ORD. LIV.—CONVOLVULACEÆ.

Gen. I. Convolvulus, L.	<i>Bindweed.</i>					
i. <i>C. arvensis, L. . . .</i>	<i>small B. . . . .</i>	Fie. . . .	Pk.	6-24 in.	P.	6-8
ii. <i>C. sepium, L. . . .</i>	<i>great B. . . . .</i>	Hed. . .	W.	3-5 ft.	P.	6-8
iii. <i>C. Soldanella, L. . .</i>	<i>Sea-side B. . . .</i>	San. sh. .	∂ Pu.	6-12 in.	P.	6-8
Gen. II. Cuscuta, L.	<i>Dodder.</i>					
* i. <i>C. Europœa, L. . . .</i>	<i>great D. . . . .</i>	Waysides !	∂ Pk.	Cl.	A.	7-9
ii. <i>C. Epilinium, Weihe.</i>	<i>flax D. . . . .</i>	Flax fie. !	∂ Pk.	Cl.	A.	8
* * iii. <i>C. Epithymum, L. . .</i>	<i>lesser D. . . . .</i>	Heaths .	∂ Pk.	Cl.	A.	8-9
iv. <i>C. Trifolii, Bab. . . .</i>	<i>Clover D. . . . .</i>	Clo. fie. !	∂ Pk.	Cl.	A.	8-9

ORD. LV.—SOLANACEÆ.

Gen. I. Hyoscyamus, L.	<i>Henbane.</i>					
i. <i>H. niger, L. . . . .</i>	<i>common H. . . . .</i>	Was. pl..	Y.	1-2 ft.	B.	6-9
Gen. II. Solanum, L.	<i>Nightshade.</i>					
i. <i>S. nigrum, L. . . . .</i>	<i>common N. . . . .</i>	Was. pl..	W.	6-24 in.	B.	6-11
v. 2. <i>miniatum, Bernh. . .</i>	. . . . .	Jersey .	. .	. .	. .	. .
ii. <i>S. Dulcamara, L. . .</i>	<i>Woody N., or Bittersweet.</i>	Hedges .	Pu.	4-6 ft.	P.	6-8
v. 2. <i>marinum . . . . .</i>	. . . . .	Sea co. .	. .	. .	. .	. .
Gen. III. Atropa, L.	<i>Deadly Night- shade, or Dwale.</i>					
i. <i>A. Belladonna, L. . .</i>	. . . . .	Was. pl. !	Pu.	2-3 ft.	P.	6-8

ORD. LII.—GENTIANACEÆ—*cont.*

CHARACTERISTICS OF SPECIES AND VARIETIES.

Gen. VI.

- i. Ls. tern. ; lfts. ob.-ov., wavy—Fls. clustered, densely bearded.

ORD. LIII.—POLEMONIACEÆ.

Gen. I.

- i. Ls. pinn. ; lfts. obl.-lanc.—Pan. downy.

ORD. LIV.—CONVOLVULACEÆ.

Gen. I.

- i. St. cli.—Ls. sagitt. ; lobes acu.—Bracts small, distant from fls.
 ii. St. cli.—Ls. sagitt. ; lobes trunc.—Bracts large, close to fls.
 iii. St. prost.—Ls. reni., fleshy—Bracts ov., close to fls.

Gen. II.

* *Cor. inflated—Scales adpr.—Sta. included.*

- i. Cal. much shorter than cor. ; seg. obt.
 ii. Cal. nearly as long as tube of cor. ; seg. acu.]
 * * *Cor. cylind.—Scales spreading—Sta. exserted.*
 iii. Cal. reddish, shorter than cor. tube.
 iv. Cal. whitish, nearly as long as cor. tube.]

ORD. LV.—SOLANACEÆ.

Gen. I.

- i. Ls. obl., sess., sinu. —Fls. sess., axill., drooping, unilat.

Gen. II.

- i. St. herbaceous with tubercled angles—Ls. ov., wavy—Berries glo., blk.
 v. 2. *Ls. toothed—Hairs spreading—Berry scarlet.*
 ii. St. shrubby, zigzag, cli.—Ls. cord., upper ones aur.—Berries ov., red.
 v. 2. *St. prost., branched—Ls. fleshy, all cord.*

Gen. III.

- i. St. herb—Ls. ent., ov.—Fls. sol., axill., lurid.

ORD. LVI.—SCHROPHULARIACEÆ.

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
Gen. I. Verbascum, L.	<i>Mullein.</i>					
* i. V. Thapsus, L. . . .	great M.	Ch. banks	Y.	2-4 ft.	B.	6-8
ii. V. Lychnitis, L. . . .	white M.	Ch. banks	W.	2-3 ft.	B.	7-8
iii. V. pulverulentum, Vill. * *	yellow hoary M.	Ch. banks	Y.	2-3 ft.	B.	7-8
iv. V. nigrum, L.	dark M.	Ch. banks	Y.	2-3 ft.	P.	6-10
v. V. Blattaria, L.	Moth M.	Was. pl.!	Y.	1-4 ft.	B.	6-10
vi. V. virgatum, With. . .	large-flowered M.	Was. pl.!	p Y.	1-4 ft.	B.	7-8
Gen. II. Veronica, L.	<i>Speedwell.</i>					
A. * i. V. spicata, L.	spiked S.	Ch. pas.!	B.	6-12 in.	P.	7-8
v. 2. <i>hybrida</i> , L. . . . . * *	. . . . .	. . . . .	. . .	. . .	. . .	. . .
ii. V. serpyllifolia, L. . . .	Thyme-leaved S. . .	Waysides	Li.	3-10 in.	P.	5-6
v. 2. <i>humifusa</i> , Dicks. . . .	. . . . .	Sc.mount.!	. . .	. . .	. . .	. . .
iii. V. alpina, L.	Alpine S.	Sc.mount.!	d B.	3-10 in.	P.	7-8
iv. V. saxatilis, L. . . . . B.	blue rock S.	Sc.mount.!	B.	2-4 in.	P.	7-8
v. V. scutellata, L.	Marsh S.	Ditches .	Pk.	6-12 in.	P.	7-8
vi. V. Anagallis, L.	Water S.	Marshes.	Li.	1-2 ft.	P.	7-8
vii. V. Beccabunga, L. . . .	Brooklime	Ditches .	B.	1-2 ft.	P.	5-9
viii. V. officinalis, L.	common S.	Dry pl. .	p B.	2-10 in.	P.	5-7
v. 2. <i>hirsuta</i> , Hopk.	. . . . .	!	. . .	. . .	. . .	. . .
ix. V. montana, L.	Mountain S. . . .	Moi. wds.	B.	1-2 ft.	P.	5-7
x. V. Chamaedrys, L. . . . . C.	Germander S. . . .	Banks . .	B.	1-2 ft.	P.	5-6
xi. V. arvensis, L.	Wall S.	Dry pl. .	p B.	4-12 in.	A.	4-7
xii. V. verna, L.	vernal S.	San. hths. E. of E.!	p B.	2-4 in.	A.	4-5
xiii. V. triphyllous, L. . . . . D.	blunt-fingered S.	San. hths. E. of E.!	B.	3-4 in.	A.	4
xiv. V. hederifolia, L. . . .	Ivy-leaved S. . . .	Banks . .	p L.	6-18 in.	A.	3-8
xv. V. agrestis, L.	field S.	Was. pl. .	B.	4-8 in.	A.	4-9
xvi. V. polita, Fries.	gray field S. . . .	Was. pl. .	B.	4-8 in.	A.	4-9
v. 2. <i>grandiflora</i>	. . . . .	. . . . .	. . .	. . .	. . .	. . .
xvii. V. Buxbaumii, Pen. . .	. . . . .	Fie. . . .	B.	6-12 in.	A.	4-9
Gen. III. Bartsia, L.	<i>Bartsia.</i>					
i. B. alpina, L.	Alpine B.	Alp. pas.!	b Pu.	4-8 in.	P.	6-8
ii. B. viscosa, L.	yellow B.	Mead.!	Y.	6-18 in.	A.	6-10
iii. B. Odontites, Huds. . . .	red B.	Was. pl. .	R.	6-18 in.	A.	6-8
a. verna, Reich.	. . . . .	. . . . .	. . .	. . .	. . .	. . .
b. scrotina, Reich.	. . . . .	. . . . .	. . .	. . .	. . .	. . .
c. rotundata, Ball	. . . . .	. . . . .	. . .	. . .	. . .	. . .
d. divergens, Ford.	. . . . .	. . . . .	. . .	. . .	. . .	. . .

ORD. LVI.—SCHROPHULARIACEÆ.

CHARACTERISTICS OF SPECIES AND VARIETIES.

Gen. I.

* *Hairs of the filaments white.*

- i. Ls. decurr., woolly on both sides—Fls. in dense spikes.
- ii. St. ang.—Ls. not decurr., glab. above—Fls. in a narrow pan.
- iii. St. terete—Ls. not decurr., mealy above and below—Fls. in a pyram. pan.
* * *Hairs of the filaments purple.*
- iv. Ls. not decurr., glab. above ; lower ones long-stalked—Fls. in a spicate pan.
- v. Ls. decurr., glab.—Fls. racemed—Pedunc. sol., longer than the cal.
- vi. Ls. decurr., gland.-hairy—Pedunc. 1-5 flowered, shorter than cal.

Gen. II.

A. Root perennial—Rac. term.* *Tube of coroll. long.*

- i. Ls. ov.-lanc. ; lower ls. with a cun. base—Spike dense.
- v. 2. *Larger—Lower ls. with a cord. base.*
* * *Tube of cor. short.*
- ii. St. asc.—Ls. broadly ov.—Spike many-flowered—Caps. ob.-cord.—Sty. long.
- v. 2. *St. prost.—Rac. short.*
- iii. St. erect—Ls. ellip.-ov.—Spike few-flowered—Caps. ob.-ov.—Sty. short.
- iv. St. woody, decumb.—Ls. ellip.—Spike few-flowered—Caps. ov.—Sty. long.
B. Root perennial—Rac. axill.
- v. St. glab.—Ls. lin.-lanc.—Rac. alt.—Caps. of two orb. lobes.
- vi. St. erect, succ.—Ls. lan., serr., sess. acute—Rac. opp.—Caps. ellip., slightly notched.
- vii. St. procumb. at base, succ.—Ls. ellip., obt.—Rac. opp.—Caps. roundish, tumid, slightly notched
- viii. St. hairy—Ls. ellip.—Rac. alt., spic.—Caps. ob.-cord., deeply notched.
- v. 2. *Smaller—Ls. ov.-lanc.—Caps. ent.—(Scot.)*
- ix. St. hairy, diff.—Ls. ob.-cord., stalked serr.—Rac. alt., lax—Caps. orb., flat, notched, with crenulate margins.
- x. St. hairy on alt. sides—Ls. ov., sess., deeply serr.—Rac. opp.—Caps. ob.-cord.
C. Root annual—Fls. spicate.
- xi. Ls. cord., ov.—Rac. lax, many-flowered.
- xii. Ls. pinnfid.—Pedi. shorter than sep.—Seeds flat.

- xiii. Ls. digit.—Pedi. longer than sep.—Seeds concave.

D. Root annual—Fls. axill., sol.

- xiv. Ls. cord., 5-7 lobed—Fls. pale lilac.
- xv. Ls. cord., serr.—Sep. obl., obt.—Lower pet. white—Cap. of two turgid lobes.
- xvi. Ls. cord., serr.—Sep. ov., ac.—Pet. all blue—Cap. of two turgid lobes.
- v. 2. *Larger—Cor. large.*
- xvii. Ls. cord., serr.—Sep. lanc., ac.—Caps. of two divar. lobes compressed upwards and keeled.

Gen. III.

- i. Ls. opp., cord., bluntly serr.—Fls. in a short, leafy spike.
- ii. Ls. lanc., serr.—Fls. axill., upper crowded.
- iii. Ls. lanc., remotely serr.—Fls. in unilat. rac.
  - a. Ls. rounded at base—Bracts longer than fls.—Cal. teeth as long as tube.
  - b. Ls. narrow at base—Bracts shorter than fls.—Cal. teeth as long as tube.
  - c. Ls. broader at base—Bracts shorter than fls.—Cal. teeth half as long as tube.
  - d. St. much branched—Branches widely spreading.

ORD. LVI.—SCHROPHULARIACEÆ—*cont.*

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
Gen. IV. Euphrasia, L.	<i>Eyebright.</i>					
i. <i>E. officinalis, L.</i> . . .	<i>common E.</i> . . .	Heaths .	W.	1-8 in.	A.	5-9
v. 2. <i>gracilis, Fries.</i> . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
v. 3. <i>maritima</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
Gen. V Rhinanthus, L.	<i>Yellow-rattle.</i>					
i. <i>R. Crista-Galli, L.</i> . . .	<i>common Y.</i> . . .	Dp. pas..	Y.	6-18 in.	A.	5-7
ii. <i>R. major, Ehrh.</i> . . .	<i>bushy Y.</i>	Corn fie. N. of E.	Y.	1-2 ft.	A.	5-7
Gen. VI. Melampyrum, L.	<i>Cow-wheat.</i>					
i. <i>M. cristatum, L.</i> . . .	<i>crested C.</i>	Copses! E. of E.	Y.	6-24 in.	A.	7
ii. <i>M. arvense, L.</i>	<i>purple C.</i>	Corn fie.!	Ro.	1-2 ft.	A.	6-8
iii. <i>M. pratense, L.</i> . . .	<i>yellow C.</i>	Copses .	Y.	6-18 in.	A.	5-8
v. 1. <i>vulgaris.</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
v. 2. <i>latifolium</i>	. . . . .	Ch. pl. !.	. . . . .	. . . . .	. . . . .	. . . . .
v. 3. <i>montanum</i>	. . . . .	Mount. .	. . . . .	. . . . .	. . . . .	. . . . .
iv. <i>M. sylvaticum, L.</i> . . .	<i>Wood C.</i>	Alp. wds.!	Y.	6-24 in.	A.	7-8
Gen. VII. Pedicularis, L.	<i>Louse-wort.</i>					
i. <i>P. palustris, L.</i> . . .	<i>Marsh L.</i>	Marshes.	Cr.	6-18 in.	A.	5-9
ii. <i>P. sylvatica, L.</i> . . .	<i>Field L.</i>	Moi. pas.	Ro.	3-10 in.	P.	4-7
Gen. VIII. Scrophularia, L.	<i>Fig-wort.</i>					
i. <i>S. nodosa, L.</i>	<i>knotted F.</i>	Moi. pl. .	g Br.	1-3 ft.	P.	7-9
v. 2. <i>Ehrharti, Stev.</i> . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
ii. <i>S. aquatica, L.</i> . . .	<i>Water F.</i>	Marshes.	g Br.	2-4 ft.	P.	7-9
iii. <i>S. Scorodoina, L.</i> . . .	<i>balm-leaved F.</i> . .	Moi. pl. ! SW. of E.	Pu.	2-3 ft.	P.	7-8
[iv. <i>S. vernalis, L.</i>	<i>yellow F.</i>	Was. pl. !	Y.	1-3 ft.	P.	4-6
Gen. IX. Digitalis, L.	<i>Fox-glove.</i>					
i. <i>D. purpurea, L.</i> . . .	<i>common F.</i> . . .	Woods .	Pu.	2-4 ft.	P.	6-8
Gen. X. Antirrhinum, L.	<i>Snap-Dragon.</i>					
i. <i>A. majus, L.</i>	<i>great S.</i>	Old walls	var.	1-2 ft.	P.	7-9
ii. <i>A. Orontium, L.</i>	<i>lesser S.</i>	Corn fie..	roPu.	6-18 in.	B.	7-10
Gen. XI. Linaria, Tourn.	<i>Toad-flax, Fluellen.</i>					
*						
i. <i>L. cymbalaria, L.</i> . . .	<i>Ivy-leaved T.</i> . . .	Old walls	p B.	6-24 in.	P.	5-9
ii. <i>L. spuria, Mill.</i> . . .	<i>round-leaved F.</i> . .	Corn fie..	Y.	4-18 in.	A.	7-10
iii. <i>L. Elatine, Mill.</i> . . .	<i>sharp-leaved F.</i> . .	Corn fie..	Y.	6-24 in.	A.	7-10
* * . . .						
iv. <i>L. repens, Ait.</i>	<i>creeping T.</i>	Ch. banks !	Li.	1-2 ft.	P.	7-9
v. <i>L. vulgaris, Mill.</i> . . .	<i>yellow T.</i>	Corn fie..	Y.	1-2 ft.	P.	7-10

ORD. LVI.—SCHROPHULARIACEÆ—*cont.*

CHARACTERISTICS OF SPECIES AND VARIETIES.

Gen. IV.

- i. Ls. ov., deeply serr., sess.—Fls. in leafy spikes—Bracts broad at base.
- v. 2. *Slenderer*—Bracts narrowed at base.
- v. 3. *Caps. longer than cal.*—(*Shetland*).

Gen. V.

- i. Ls. lanc., serr.—Fls. in lax spikes—Bracts green.
- ii. Ls. lin.-lanc., serr.—Fls. in crowded spikes—Bracts yellow.

Gen. VI.

- i. Fls. in dense, imbric., four-sided spikes—Bracts cord., pect., coloured.
- ii. Fls. in lax, coni., spikes—Bracts broadly lanc., pinnfid., coloured.
- iii. Fls. axill., secund.—Cor. four times as long as cal.—Upper bracts ent., or toothed at base, green.
- v. 1. *Ls. lin.-lanc., acum.*—Bracts all toothed; teeth long, slender.
- v. 2. *Ls. ov.-lanc., acum.*—Bracts all toothed; teeth short.
- v. 3. *Ls. lin.-lanc.; lower obt.*—Bracts ent.
- iv. Fls. axill., secund.—Cor. twice as long as cal.—Bracts all ent.

Gen. VII.

- i. St. sol., erect—Cal. hairy.
- ii. St. branched, spreading—Cal. glab.

Gen. VIII.

- i. St. four-angled—Ls. glab., ac.
- v. 2. *St. winged above*—Bracts leafy.
- ii. St. winged—Ls. glab., obt.
- iii. St. bluntly four-angled—Ls. downy, cord.-trian.—Cor. purplish.
- iv. St. winged—Ls. downy, cord.—Cor. yell.]

Gen. IX.

- i. Ls. ov., lanc., downy, veiny—Fls. spic., drooping.

Gen. X.

- i. Ls. lanc.—Rac. dense—Sep. ov., shorter than cor.]
- ii. Ls. lin.-lanc.—Rac. lax—Sep. lin., longer than cor.

Gen. XI.

* *Stem trailing.*

- i. Ls. five-lobed, glab.—Fls. long-stalked, sol.
- ii. Ls. roundish-ov., downy—Spur curved.
- iii. Ls. ovate-hast., downy—Spur straight.
- * * *St. erect.*
- iv. Ls. lin., whorled—Fls. in a long, lax pan.—Cor. violet.
- v. Ls. lin., crowded—Fls. in a dense, imbric. rac.—Cor. yell.

ORD. LVI.—SCHROPHULARIACEÆ—*cont.*

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
v. 2. <i>speciosa</i> , Bromf. . .						
vi. <i>L. Pelisseriana</i> , Mill.	purple T.	St. Ouen's, Jersey.	Pu.	6-10 in.	A.	6-7
vii. <i>L. minor</i> , Desf. . . .	least T.	San. fie. .	puY	6-18 in.	A.	5-10
Gen. XII. <i>Limosella</i> , L.	Mud-wort.					
i. <i>L. aquatica</i> , L. . . .	common M. . . .	Mud pl.!	pkW.	1-2 in.	A.	7-9
Gen. XIII. <i>Sipthorpia</i> , L.	Money-wort.					
i. <i>S. europæa</i> , L. . . .	Cornish M. . . .	Sh. banks S. of E.!	Pk.	6-12 in.	P.	7-8
Gen. XIV. <i>Mimulus</i> , L.						
[i. <i>M. luteus</i> , L.	yellow M. . . .	Riv. banks	Y.	6-24 in.	P.	7-9

ORD. LVII.—OROBANCHACEÆ.

Gen. I. <i>Orobanche</i> , L.	Broom-rape.					
*						
i. <i>O. Rapum</i> , Thuil. . .	great B.	Broom furze.	W.	1-1½ ft.	P.	5-7
ii. <i>O. caryophyllacea</i> , Sm.	Clove B.	Galium Mollugo.	W.	6-12 in.	P.	5-7
iii. <i>O. elatior</i> , Sutton . .	tall B.	Centaurea scabiosa.	W.	6-12 in.	P.	6-8
* *						
iv. <i>O. minor</i> , L.	least B.	. . . .	W.	½-2 ft.	A.	6-10
a. <i>minor</i> , L.	. . . . .	Clover .	. .	. .	. .	. .
b. <i>amethystea</i> , Thuil. .	. . . . .	Daucus maritimus, S. of E.!	. .	. .	. .	. .
c. <i>Picridis</i> , F. Schultz .	. . . . .	Picris hie- racioides!	. .	. .	. .	. .
d. <i>Hederæ</i> , Duby	. . . . .	Ivy! . .	. .	. .	. .	. .
v. <i>O. rubra</i> , Sm.	red B.	Thymus serpyllum!	dull R.	4-8 in.	P.	7-8
* * *						
vi. <i>O. cærulea</i> , Vill. . .	purple B.	Achillea mille- folium! S. of E.	b Pu.	6-12 in.	P.	6-8
vii. <i>O. arenaria</i> , Bork. . .	Sand B.	Jersey! .	b Pu.	6-12 in.	P.	6-8
[viii. <i>O. ramosa</i> , L.	branched B. . . .	Hemp! .	. .	4-8 in.	A.	8-9
Gen. II. <i>Lathræa</i> , L.	Tooth-wort.					
i. <i>L. squamaria</i> , L. . .	great T.	Woods .	bPu.	4-10 in.	P.	4-5

ORD. LVI.—SCHROPHULARIACEÆ—*cont.*

CHARACTERISTICS OF SPECIES AND VARIETIES

v. 2. *Ls. broader*—*Fls. much larger.*

vi. Ann.—*Ls. lin.*, lower ones vertic.—*Fls. in short, few-flowered rac.*—*Cor. pur.*

vii. *Ls. lin.-lanc.*, obt.—*Fls. sol.*, axill.

Gen. XII.

i. Rhizome cr., rooting—*Ls. lan.*, spath., long-stalked.

Gen. XIII.

i. St. fili., hairy, prost.—*Ls. reni.* 5-7 lobed—*Fls. minute.*

Gen. XIV.

i. St. erect, holl.—*Ls. obl.*, coarsely-toothed, upper sess.]

ORD. LVII.—OROBANCHACEÆ.

Gen. I.

* *Bracts one*—*Sepals bifid.*

i. Cor. campan.; upper lip nearly ent.; lower three-lobed, middle lobe much the longest.

ii. Cor. campan.; upper lip two-lobed; lower of three nearly equal, obtuse lobes.

iii. Cor. tubular, curved; upper lip of two inflexed lobes; lower of three nearly equal, acute lobes.

* * *Bracts one*—*Sep. entire.*

iv. Yellowish brown—*Sep. one or many-veined*, ovate, with long slender points.

a. *Sep. many-veined*—*Cor. tubular*, slightly curved—*Lobes of stig. (purple)* just touching.

b. *Sep. many-veined*—*Cor. much curved at base*—*Lobes of stig. (purple)* divari.

c. Very pale—*Sep. 1-3 veined*—*Cor. slightly curved at each end*—*Lobes of stig. (purple)* nearly close.

d. Rac. lax—*Sep. one-veined*—*Lobes of stig. (yellow)* nearly entire.

v. Dull red—*Sep. one-veined, lanc., subu.*, longer than the cor.—*Lobes of lower lip* nearly equal.

* * * *Bracts three.*

vi. St. simple—*Lobes of cor. acute*—*Anth. glab.*

vii. St. simple—*Lobes of cor. obt.*—*Anth. pilose.*

viii. St. branched—*Sep. 4.*]

Gen. II.

i. *Fls. drooping, unilat.*—*Lower lip of cor. three-cleft.*

ORD. LVIII.—VERBENACEÆ.

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
Gen. I. Verbena, L.	<i>Vervain.</i>					
i. <i>V. officinalis, L.</i> . . .	<i>common V.</i> . . .	Dry pl. .	Li.	1-2 ft.	P.	7-9

ORD. LIX.—LABIATÆ.

Gen. I. Salvia, L.	<i>Sage, Clary.</i>					
i. <i>S. Verbenaca, L.</i> . . .	<i>wild S.</i>	Ch. downs	b Pu.	1-2 ft.	P.	5-8
ii. <i>S. pratensis, L.</i> . . .	<i>Meadow C.</i>	Dry fie.!	B.	2-5 ft.	P.	6-8
Gen. II. Lycopus, L.	<i>Gipsy-wort.</i>					
i. <i>L. Europæus, L.</i> . . .	<i>common G.</i>	Ditches .	W.	1-3 ft.	P.	6-9
Gen. III. Mentha, L.	<i>Mint.</i>					
<i>A.</i>						
* i. <i>M. rotundifolia, L.</i> . .	<i>round-leaved M.</i> .	Wet pl. .	Li.	2-3 ft.	P.	8-9
v. 2. <i>alopecuroides, Hull.</i>						
ii. <i>M. sylvestris, L.</i> . . .	<i>Horse M.</i>	Wet pl. .	Li.	2-3 ft.	P.	8-9
v. 2. <i>nemorosa, Willd.</i> . .						
v. 3. <i>mollissima, Borkh.</i>						
[iii. <i>M. viridis, L.</i>	<i>Spear M.</i>	Marshes !	Li.	2-3 ft.	P.	8-9
* *						
iv. <i>M. piperata, L.</i>	<i>Pepper M.</i>	Dp. pl. .	Li.	6-18 in.	P.	8-9
a. <i>officinalis, Hull</i>						
b. <i>vulgaris, Sole</i>						
v. <i>M. aquatica, L.</i>	<i>Water M.</i>	Marshes.	Li.	1-4 ft.	P.	8-9
a. <i>pubescens, Willd.</i>		S. of E.!				
b. <i>hircina, Hull</i>						
c. <i>hirsuta, L.</i>						
d. <i>subglabra, Baker</i>						
e. <i>citrate, Ehrh.</i>	<i>Bergamot M.</i> . . .					
<i>B.</i>						
vi. <i>M. sativa, L.</i>	<i>Marsh whorled M.</i>	Wet. pl. .	Li.	2-5 ft.	P.	7-8
a. <i>genuina, Sy.</i>						
v. 2. <i>paludosa, Sol.</i>						
v. 3. <i>subglabra, Baker</i> . . .						
b. <i>rubra, Sm.</i>						
c. <i>gracilis, Sm.</i>		Wilts.!				
v. 2. <i>cardiaca, Baker</i>						
d. <i>pratensis, Sole</i>						
e. <i>gentilis, L.</i>						
v. 2. <i>Wirtgeniana,</i> <i>F. Schultz.</i>						

ORD. LVIII.—VERBENACEÆ.

CHARACTERISTICS OF SPECIES AND VARIETIES.

Gen. I.

- i. St. erect, four-angled—Ls. opp., trifold and ent.—Fls. in long spikes.

ORD. LIX.—LABIATÆ.

Gen. I.

- i. Upper ls. broad, cord.—Cor. narrow, about as long as cal.
 ii. Upper ls. ov.-lanc., ac.—Cor. thrice as long as cal.

Gen. II.

- i. Ls. deeply serr. or pinnfid., wrinkled—Fls. in dense whorls.

Gen. III.

A. Throat of calyx naked—Fls. in term. heads.

* *Ls. sess. or only lower ls. stalked.*

- i. Ls. ellip., ov., serr., wrinkled, shaggy beneath—Spikes interrupted—Bracts ov.-acum.
v. 2. Spikes short, stout—Bracts narrower.
 ii. Ls. ov.-lanc., serr., ac., silky beneath—Spikes slender, continuous—Bracts subu.
v. 2. Spikes stouter—Ls. broadly obl., tomentose beneath.
v. 3. Ls. broadly ov.-cord. at base, felted beneath.
 iii. Ls. glab., lanc., ac.—Spikes cylind., interrupted—Bracts subul.—Smell very pungent.]
 * * *Ls. stalked.*
 iv. Glab.—Ls. ov.-lanc., upper smaller—Spikes lax, interrupted below.
a. Ls. rounded at base—Spikes slender.
b. Ls. sub-cord. at base—Spikes shorter.
 v. Hairy (mostly)—Ls. ov., serr.; upper ls., bracteiform, shorter than fls.—Spikes axill. and term., dense
a. Ls. rounded at base, pubesc. above, woolly beneath.
b. Ls. rounded at base, glab. above, hairy beneath.
c. Ls. sub-cord., hairy on both sides.
d. Ls. sub-cord., glab., except on the nerves beneath.
e. Whole plant glab.!
B. Throat of calyx naked—Fls. in axill. whorls.
 vi. Ls. ellip., serr.; upper ls. smaller—Cal. teeth lanc., ac.
a. Cor. and cal. hairy—Whorls separate—Bracts leafy.
v. 2. Upper whorls continuous—Bracts smaller.
v. 3. Sub-glab.
b. Cor. glab.; cal. hairy—Nerves of ls. purple—Ls. glab.
c. Cor. glab.; cal. ciliate—Ls. glab.—Whorls all separate—Bracts 5-6 times as long as whorls.
v. 2. Bracts 2-4 times as long as whorls.
d. Ls. nearly sess., drooping, hairy above, and hairy on the nerves only beneath—Whorls distant, sub-glob.—(New Forest !)
e. Ls. all nearly of one size, hairy above, and on the few nerves beneath—Cor. glab.—cal. teeth densely hairy—St. hairy.
v. 2. St. almost glab.—Cal. teeth sparingly hairy.

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
<i>v. 3. Pauliana, F. Schultz</i>						
vii. <i>M. arvensis, L.</i> . . .	<i>Corn M.</i>	Corn fie..	Li.	1-2 ft.	P.	8-9
<i>a. genuina</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
<i>b. nummularia, Shreb.</i> . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
<i>c. agrestis, Sole</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
<i>d. præcox, Sole</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
<i>e. Allionii, Bor.</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
<i>f. parietariifolia, Beck.</i> . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
<i>C.</i>						
viii. <i>M. Pulegium, L.</i> . . .	<i>Penny-royal</i>	Dp. hths.	Li.	2-10 in.	P.	8-9
Gen. IV. Thymus, L.	<i>Thyme.</i>					
<i>i. T. serpyllum, L.</i> . . .	<i>wild T.</i>	Dry hths.	Pu.	2-8 in.	P.	6-8
<i>ii. T. Chamædrys, Fr.</i> . .	. . . . .	Heaths .	Pu.	2-8 in.	P.	6-8
Gen. V. Origanum, L.	<i>Marjoram.</i>					
<i>i. O. vulgare, L.</i>	<i>common M.</i>	Dry hills	Pu.	1-2 ft.	P.	7-9
Gen. VI. Calamintha, <i>Mænoch.</i>	<i>Calamint, Basil</i> <i>Thyme.</i>					
<i>*</i>						
<i>i. C. Acinos, Clairv.</i> . . .	<i>common B.</i>	Corn fie.	<i>b</i> Pu.	6-8 in.	A.	7
<i>ii. C. Clinopodium, Sp.</i> . .	<i>wild B.</i>	Dry hills	Pu.	12-18 in.	P.	7-9
<i>* *</i>						
<i>iii. C. Nepeta, Clairv.</i> . . .	<i>lesser C.</i>	Was. pl..	. .	1-2 ft.	P.	7-8
<i>iv. C. officinalis, Auct.</i> . .	<i>common C.</i>	Was. pl..	. .	1-2 ft.	P.	7-8
<i>a. menthifolia, Host.</i> . . .	. . . . .	. . . . .	. .	. .	. .	. .
<i>v. 2. Briggsii, Sy.</i>	. . . . .	. . . . .	. .	. .	. .	. .
<i>b. sylvatica, Bromf.</i> . . .	. . . . .	I. of W..	. .	. .	. .	. .
Gen. VII. Melissa, L.	<i>Balm.</i>					
<i>[i. M. officinalis, L.]</i>	<i>common B.</i>	S. of E.!	W.	1-3 ft.	P.	7-8
Gen. VIII. Melittis, L.	<i>Bastard-Balm.</i>					
<i>i. M. Mellissophyllum, L.</i>	<i>Bastard-Balm.</i> . .	Copses !	<i>pu</i> W.	1-2 ft.	P.	5-6
Gen. IX. Teucrium, L.	<i>Germander.</i>					
<i>i. T. Scorodonia, L.</i> . . .	<i>Wood G.</i>	Heaths .	<i>y</i> W.	1-2 ft.	P.	7-8
<i>ii. T. Scordium, L.</i>	<i>Water G.</i>	Wet mead !	Pu.	4-10 in.	P.	7-8
<i>v. 2. Scordioides, Bab.</i> . .	. . . . .	. . . . .	. .	. .	. .	. .
iii. <i>T. Botrys, L.</i>	<i>Cut-leaved annual</i> <i>G.</i>	Box Hill !	Ro.	4-8 in.	B.	8
iv. <i>T. Chamædrys, L.</i> . . .	<i>Wall G.</i>	Old walls !	Ro.	6-18 in.	P.	7-9
Gen. X. Ajuga, L.	<i>Bugle.</i>					
<i>1. A. reptans, L.</i>	<i>common B.</i>	Moi. wds.	<i>pu</i> B.	6-12 in.	P.	5-6
<i>b. pseudo-alpina</i>	. . . . .	. . . . .	. .	. .	. .	. .

CHARACTERISTICS OF SPECIES AND VARIETIES.

v. 3. *St. almost glab.*—*Cal. teeth densely hairy.*

vii. Ls. obl.-lanc., upper same size—*Cal. teeth triang., as broad as long.*

a. *St. short, very hairy.*

b. *St. long, not so hairy*—Ls. and cal. sparingly hairy.

c. *St. long, hairy*—Ls. cord., wrinkled—*Cal. very hairy.*

d. *St. thick, hairy*—Ls. slightly hairy—*Cal. teeth longer.*

e. *St. tall, hairy above, glab. below*—*Cal. teeth very short.*

f. *St. long, nearly glab.*—Ls. glab.

(*Many other intermediate varieties occur.*)

C. Throat of cal. hairy.

viii. *St. prost.*—Ls. ellip., obt.—Fls. in distant whorls.

Gen. IV.

i. *St. prost.* ; flowering sts. erect—Ls. obl.—Upper lip of cor. obl.

ii. Sts. all asc.—Ls. ov.—Upper lip of cor. broad, short.

Gen. V.

i. Ls. ov., obt.—Fls. panicled ; bracts ov.

Gen. VI.

* *Whorls simp.*

i. Whorls 5-6 flowered—*Cal. gibbous at base.*

ii. Whorls dense, many-flowered—*Cal. equal at base.*

* * *Whorls compound in forked cymes.*

iii. Hairs on mouth of cal. prominent—Ls. shortly stalked.

iv. Hairs on mouth of cal. included.

a. Teeth of upper lip of cal. straight—Seg. of lower lip of cor. distant, centre one longest.

v. 2. *Larger—Pedunc. of lower whorls longer than the stalk of central fl.*

b. Teeth of upper lip of cal. reflexed—Seg. of lower lip of cor. overlapping, equal.

Gen. VII.

i. Ls. ov., ac., serr.—Fls. in secund., axill., whorls.]

Gen. VIII.

i. Ls. obl., ov.—Cor. large (1-1½ in. long).

Gen. IX.

i. Ls. stalked, ov.-cord., wrinkled—Fls. unilat. in lat. and term. rac.

ii. Ls. sess., obl., all similar—Fls. in axill. whorls.

v. 2. *Ls. cord., amplex.*

iii. Ls. pinnfid. ; seg. obl. or cut—Fls. in axill. whorls.

iv. Ls. stalked, ov., wedge-shaped at base—Fls. in axill. whorls.

Gen. X.

i. Glab.—St. with creeping scions—Fls. whorled, in a loose spike.

b. A form without creeping scions.

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
ii. <i>A. pyramidalis</i> , <i>L.</i> . . .	<i>pyramidal B.</i> . . .	Alp. pas.!	<i>bu</i> B.	6-12 in.	P.	5-7
iii. <i>A. Chamæpitys</i> , <i>Shered</i>	<i>yellow B., or</i> <i>Ground-pine.</i>	Ch. fie.!	Y.	3-6 in.	P.	5-8
Gen. XI. Ballota , <i>L.</i>	<i>Horehound.</i>					
i. <i>B. nigra</i> , <i>L.</i>	<i>black H.</i>	Was. pl..	Pu.	2-3 ft.	P.	6-10
v. 2. <i>ruderalis</i> , <i>Fries.</i> . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
Gen. XII. Leonurus , <i>L.</i>	<i>Mother-wort.</i>					
[i. <i>L. cardiaca</i> , <i>L.</i>	<i>common M.</i>	Was. pl.!	<i>r</i> W.	2-3 ft.	P.	7-9
Gen. XIII. Lamium , <i>L.</i>	<i>Dead-nettle.</i>					
*						
i. <i>L. Galeobdolon</i> , <i>Crantz.</i>	<i>yellow Weasel-</i> <i>snout, or Arch-</i> <i>angel.</i>	Thick. .	Y.	6-18 in.	P.	5-6
*						
*						
<i>A.</i>						
ii. <i>L. album</i> , <i>L.</i>	<i>white D.</i>	Was. pl..	W.	6-18 in.	P.	5-9
iii. <i>L. maculatum</i> , <i>L.</i> . . .	. . . . .	Was. pl.!	Pu.	6-18 in.	P.	5-9
		N. of E.				
<i>B.</i>						
iv. <i>L. amplexicaule</i> , <i>L.</i> . .	<i>Henbit D.</i>	San. fie. .	Ro.	4-10 in.	A.	5-8
v. 2. <i>intermedium</i> , <i>Fr.</i> . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
v. <i>L. purpureum</i> , <i>L.</i> . . .	<i>red D.</i>	Was. pl..	<i>ro</i> Pu.	6-18 in.	A.	4-10
<i>a. genuinum</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
<i>b. decipiens</i> , <i>Son.</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
vi. <i>L. incisum</i> , <i>Willd.</i> . . .	<i>cut-leaved D.</i> . . .	Was. pl..	<i>ro</i> Pu.	6-8 in.	A.	4-10
Gen. XIV. Galeopsis , <i>L.</i>	<i>Hemp-nettle.</i>					
*						
i. <i>G. Ladanum</i> , <i>L.</i>	<i>red H.</i>	Ch. fie. .	Ro.	10-12 in.	A.	7-10
<i>b. canescens</i>	. . . . .	Shingly shores.	. . . . .	. . . . .	. . . . .	. . . . .
<i>c. intermedia</i> , <i>Vill.</i>	. . . . .	Moray. .	. . . . .	. . . . .	. . . . .	. . . . .
ii. <i>G. ochroleuca</i> , <i>Lam.</i> . .	<i>downy H.</i>	San. fie.!	Y.	10-12 in.	A.	7-10
*						
iii. <i>G. Tetralcit</i> , <i>L.</i>	<i>common H.</i>	Corn fie..	Pu. or W.	1-2 ft.	A.	7-9
v. 2. <i>bifida</i> , <i>Boeun.</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
iv. <i>G. versicolor</i> , <i>Curt.</i> . .	<i>large-flowered H.</i>	Corn fie.!	Y.	2-3 ft.	A.	7-8
Gen. XV. Stachys , <i>L.</i>	<i>Woundwort,</i> <i>Betony.</i>					
i. <i>S. Betonica</i> , <i>Benth.</i> . .	<i>Wood B.</i>	Thick. .	Pu.	1-2 ft.	P.	6-8

CHARACTERISTICS OF SPECIES AND VARIETIES.

- ii. Hairy—Scions none—Fls. whorled in a compact, pyram. spike.
- iii. Ls. trifid ; segm. lin.—Fls. sol., axill.

Gen. XI.

- i. Ls. ov., serr.—Fls. whorled ; bracts lin.—Teeth of cal. ov., suddenly acum.
- v. 2. *Teeth of cal. lanc., gradually acum.—(St. Vincent's Rocks.)*

Gen. XII.

- i. Lower ls. palm, five cleft ; upper ent.—Fls. in crowded whorls ; upper lip of cor. hairy.]

Gen. XIII.

* *Lobes of lower lip of cor. equal.*

- i. Ls. ov., ac., coarsely serr.—Fls. whorled, yell.

* * *Lower lip of cor. with one ob.-cord. lobe, with one or two teeth at base.*

A. Perenn.—Tube of cor. curved.

- ii. Fls. white—Hairs in tube of cor. oblique.
- iii. Fls. purple—Ls. spotted with white—Hairs in tube of cor. transverse.

B. Annual—Tube of cor. straight.

- iv. Upper ls. sess., orb., wrinkled—Cor. long, slender—Cal. teeth connivent in fr., as long as cal. tube.
- v. 2. *Habit of amplexicaule, but cal. teeth always spreading, and much longer than tube.*
- v. Ls. all stalked, cord., cren.—Tube of cor. longer than cal. ; cal. teeth spreading.
- a. Ls. and bracts slightly crenate.
- b. Ls. and bracts deeply crenate.
- vi. Ls. stalked, cord., deeply inciso.-serr.—Tube of cor. shorter than cal.

Gen. XIV.

* *Softly hairy—St. not swollen at joints.*

- i. Ls. lanc., downy—Cor. (Rose), its upper lip slightly notched.
- b. Very pubesc.—Ls. lin.-lanc., hairs on cal. spreading.
- c. Ls. ov.-lanc., serr.—Whorls all sep.—Tube of cor. shorter.
- ii. Cor. (yell.) much longer than cal. ; its upper lip deeply notched.
- * * *Hispid—St. thickened at joints.*
- iii. Cor. (Pu. or W.), its tube as long as cal.—Teeth of cal. twice as long as its tube.
- v. 2. *Cor. half as long as cal. ; its lower lip notched.*
- iv. Cor. (Yell.) twice as long as cal.—Teeth of cal. shorter than its tube.

Gen. XV.

- i. Lower ls. cord.-obl., long-stalked ; upper ones few, lin., sess.—Fls. in dense, term. spikes.

ORD. LIX.—LABIATÆ—*cont.*

Botanical Name.	Common Name.	Soil or Situation.	Colour	Growth.	Dura- tion.	Time of flowering.
ii. <i>S. palustris</i> , <i>L.</i> . . .	<i>Marsh W.</i> . . .	Wet pl. .	Pu.	1-3 ft.	P.	7-8
v. 2. <i>ambigua</i> , <i>Sm.</i> . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
iii. <i>S. sylvatica</i> , <i>L.</i> . . .	<i>Hedge W.</i> . . .	Sh. pl. .	Pu.	1-3 ft.	P.	7-8
iv. <i>S. germanica</i> , <i>L.</i> . . .	<i>downy W.</i> . . .	Ch. fie.! .	roPu.	1-3 ft.	B.	7-8
v. <i>S. arvensis</i> , <i>L.</i>	<i>Corn W.</i>	Corn fie..	p Pu.	6-18 in.	A.	4-11
Gen. XVI. <i>Nepeta</i>, <i>L.</i>						
i. <i>N. Glechoma</i> , <i>Benth.</i>	<i>Ground-ivy</i> . . .	Was. pl..	B.	Cr.	P.	3-5
a. <i>hederacea</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
b. <i>parviflora</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
c. <i>hirsuta</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
ii. <i>N. Cataria</i> , <i>L.</i>	<i>Cat-mint.</i>	Was. pl.!	W.	2-3 ft.	P.	7-9
Gen. XVII. <i>Marrubium</i>, <i>L.</i>						
	<i>White Horehound.</i>					
i. <i>M. vulgare</i> , <i>L.</i>	<i>common W.</i>	Was. pl.!	W.	12-18 in.	P.	8-9
Gen. XVIII. <i>Prunella</i>, <i>L.</i>						
	<i>Self-heal.</i>					
i. <i>P. vulgaris</i> , <i>L.</i>	<i>common S.</i>	Moi. pas.	Vi.	4-10 in.	P.	7-8
Gen. XIX. <i>Scutellaria</i>, <i>L.</i>						
	<i>Skull Cap.</i>					
i. <i>S. galericulata</i> , <i>L.</i> . . .	<i>common S.</i>	Riv. banks	B.	8-12 in.	P.	7-8
ii. <i>S. minor</i> , <i>L.</i>	<i>lesser S.</i>	Moi. hths.	Pk.	4-6 in.	P.	7-10

ORD. LX.—BORAGINACEÆ.

Gen. I. <i>Myosotis</i>, <i>L.</i>						
	<i>Scorpion Grass.</i>					
i. <i>M. palustris</i> , <i>With.</i> .	<i>Water S., or</i> <i>Forget-me-not.</i>	Wet pl. .	B.	12-18 in.	P.	6-8
v. 2. <i>strigulosa</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
ii. <i>M. repens</i> , <i>Don.</i> . . .	<i>creeping Water S.</i>	Wet pl. .	B.	4-15 in.	P.	6-8
iii. <i>M. cæspitosa</i> , <i>Schultz.</i>	<i>tufted Water S.</i> .	Wet pl. .	B.	6-18 in.	P.	6-8
	* *					
	<i>A.</i>					
iv. <i>M. alpestris</i> , <i>Schmidt.</i>	<i>Rock S.</i>	Sc. mount.	B.	5-10 in.	P.	7-8
v. <i>M. sylvatica</i> , <i>Ehrh.</i> .	<i>Wood S.</i>	Dry woods	B.	1-2 ft.	P.	6-8
vi. <i>M. arvensis</i> , <i>Hoffm.</i> .	<i>Field S.</i>	Was. pl..	B.	6-18 in.	A.	6-8
v. 2. <i>umbrosa</i> , <i>Bab.</i> . . .	. . . . .	Sh. pl. .	. . .	. . .	. . .	. . .
	<i>B.</i>					
vii. <i>M. collina</i> , <i>Hoffm.</i> . .	<i>early Field S.</i> . . .	Dry banks	B.	3-10 in.	A.	4-5
viii. <i>M. versicolor</i> , <i>Lehm.</i> .	<i>yellow and blue S.</i>	Dp. mead.	y B.	3-10 in.	A.	4-6
Gen. II. <i>Lithospermum</i>, <i>L.</i>						
	<i>Gromwell.</i>					
i. <i>L. officinale</i> , <i>L.</i> . . .	<i>common G.</i>	Was. pl. .	p Y.	12-18 in.	P.	6-7

ORD. LIX.—LABIATÆ—*cont.*

CHARACTERISTICS OF SPECIES AND VARIETIES.

- ii. St. hollow—Ls. lin.-lanc., cord., sess. ; lower very shortly stalked.
- v. 2. *Ls. shortly stalked (a hybrid).*
- iii. St. solid—Ls. all stalked, ov.-cord., acum. ; upper ones lin., ent.
- iv. Densely silky—Lower ls. obl.-ov., cord., wrinkled, stalked ; upper lanc., sess.
- v. St. weak, decumb.—Ls. cord.-ov., obt.—Cor. scarcely longer than cal.

Gen. XVI.

- i. St. procumb.—Ls. reni., ent., deeply cren.—Whorls 3-4 flowered.
- a. Glab.—Cal. half as long as cor. tube.
- b. Glab.—Cal. as long as cor. tube.
- c. Hairy—Cal. as long as cor. tube.
- ii. St. erect—Ls. ov.-cord., serr., pubes. beneath—Whorls dense.

Gen. XVII.

- i. St. erect, hoary—Ls. ov., cren., woolly, wrinkled—Whorls dense.

Gen. XVIII.

- i. St. erect—Ls. ov.-obl.—Whorls forming a dense spike, with two ls. at base.

Gen. XIX.

- i. Fls. blue—Ls. all obl.-lanc., cord., cren.
- ii. Fls. pink—Lower ls. broadly ov. ; upper lanc. ent.

ORD. LX.—BORAGINACEÆ.

Gen. I.

* *Hairs of cal. straight, adpr.*

- i. Cor. large—Cal. divided to one-third ; its teeth short, triang.—Hairs of st. spreading.
- v. 2. *Fls. smaller—St. more erect, with adpr. hairs.*
- ii. Cor. large—Cal. divided to half ; its teeth lanc.—Hairs of st. spreading.
- iii. Cor. scarcely longer than cal.—Cal. teeth lanc.—Hairs of st. adpr.
- * * *Hairs of cal. hooked, spreading.*
- A. *Pedi. longer than cal.*
- iv. Fls. large—Rad. ls. pointed, long-stalked—Cal. deeply five cleft, atten. at base—Sty. half as long as cal.
- v. Fls. large—Rad. ls. blunt, long-stalked—Cal. deeply five cleft, obt. at base—Sty. nearly as long as cal.
- vi. Fls. small—Ls. ob.-ov., obt.—Cal. five cleft to middle—Sty. very short.
- v. 2. *Fls. larger.*
- B. *Pedi. shorter than cal.*
- vii. Cor. bright blue ; tube short—Cal. of fr. open, ventri.
- viii. Cor. yell., changing to dull blue, tube long—Cal. of fr. closed, obl.

Gen. II.

- i. St. erect—Cor. as long as cal., with small scales in throat—Ach. polished.

ORD. LX.—BORAGINACEÆ—cont.

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
ii. <i>L. arvense</i> , <i>L.</i>	<i>Corn G., or Bas- tard Alkanet.</i>	Was. pl. .	g W.	10-16in.	A.	5-6
iii. <i>L. purpureo-cœruleum</i> , <i>L.</i>	<i>purple G.</i>	Thick.! .	B.	1-2 ft.	P.	6-7
Gen. III. <i>Mertensia</i> , <i>Roth.</i>	<i>Smooth Gromwell.</i>					
i. <i>M. maritima</i> , <i>Don.</i> . . .	<i>Sea-side S.</i>	Sea sh.! .	b Pu.	1-2 ft.	P.	5-6
Gen. IV. <i>Symphytum</i> , <i>L.</i>	<i>Comfrey.</i>					
i. <i>S. officinale</i> , <i>L.</i>	<i>common C.</i>	Riv. banks	y W.	2-3 ft.	P.	5-6
v. 2. <i>patens</i> , <i>Siphth.</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
ii. <i>S. tuberosum</i> , <i>L.</i>	<i>tuberous C.</i>	Sha. wds. N. of E.	y W.	1-2 ft.	P	5-6
Gen. V. <i>Borago</i> , <i>L.</i>	<i>Borage.</i>					
i. <i>B. officinalis</i> , <i>L.</i>	<i>common B.</i>	Was. pl. .	B.	1-2 ft.	B.	6-7
Gen. VI. <i>Lycopsis</i> , <i>L.</i>	<i>Bugloss.</i>					
i. <i>L. arvensis</i> , <i>L.</i>	<i>small B.</i>	Corn fie.	B.	6-18 in.	A.	6-7
Gen. VII. <i>Anchusa</i> , <i>L.</i>	<i>Alkanet.</i>					
[i. <i>A. officinalis</i> , <i>L.</i>	<i>common A.</i>	Was. pl.! .	Pu.	1-2 ft.	P.	6-7
[ii. <i>A. sempervirens</i> , <i>L.</i> . .	<i>evergreen A.</i> . . .	Was. pl.! .	B.	1-2 ft.	P.	5-6
Gen. VIII. <i>Asperugo</i> , <i>L.</i>	<i>Mad-wort.</i>					
[i. <i>A. procumbens</i> , <i>L.</i> . . .	<i>German M.</i>	Was. pl.! .	b Pu.	1-2 ft.	A.	6-7
Gen. IX. <i>Cynoglossum</i> , <i>L.</i>	<i>Hound's-tongue.</i>					
i. <i>C. officinale</i> , <i>L.</i>	<i>common H.</i>	Was. pl. .	r Pu.	1-2 ft.	B.	6-7
ii. <i>C. montanum</i> , <i>Lam.</i> . .	<i>green-leaved H.</i> . .	Sha. pl. S. of E.!	r Pu.	1-2 ft.	B.	6-7
Gen. X. <i>Pulmonaria</i> , <i>L.</i>	<i>Lung-wort.</i>					
i. <i>P. angustifolia</i> , <i>L.</i> . . .	<i>narrow-leaved L.</i>	Thick.! S. of E.	Pk.	12-18in.	P.	5-6
[ii. <i>P. officinalis</i> , <i>L.</i>	<i>common L.</i>	Thick.! .	p Pu.	12-18in.	P.	5-6
Gen. XI. <i>Echium</i> , <i>L.</i>	<i>Viper's Bugloss.</i>					
i. <i>E. vulgare</i> , <i>L.</i>	<i>common V.</i>	Ch. fie. .	r Pu.	2-3 ft.	B.	6-7
ii. <i>E. plantagineum</i> , <i>L.</i> . .	<i>purple-flowered V.</i>	Jersey .	b Pu.	1-3 ft.	B.	6-7

ORD. LXI.—LENTIBULARIACEÆ.

Gen. I. <i>Pinguicula</i> , <i>L.</i>	<i>Butter-wort.</i>					
i. <i>P. grandiflora</i> , <i>Lam.</i> . .	<i>large-flowered B.</i>	Bogs ! I..	Vi.	4-6 in.	P.	5-6
ii. <i>P. vulgaris</i> , <i>L.</i>	<i>common B.</i>	Bogs . .	Vi.	2-4 in.	P.	5-7

ORD. LX.—BORAGINACEÆ—*cont.*

CHARACTERISTICS OF SPECIES AND VARIETIES.

- ii. St. erect—Cor. rather longer than cal., throat naked—Ach. wrinkled.
- iii. Barren st. prost.—Cor. (bluish Pu.) much longer than cal.

Gen. III.

- i. St. procumb.—Ls. ov., ac., rough, glauc., fleshy.

Gen. IV.

- i. St. winged above—Upper ls. lanc., very decurrent.
- v. 2. *Cor. purple—Cal. seg. spreading.*
- ii. St. scarcely winged—Upper ls. ov.-lanc., slightly decurr.

Gen. V.

- i. Very hisp.—Lower ls. ob.-ov., atten. at base—Seg. of cor. acu., spread.

Gen. VI.

- i. Very hisp.—Ls. lanc., wavy ; bristles with tuberc. bases.

Gen. VII.

- i. Ls. lanc.—Spikes unilat., crowded.]
- ii. Ls. ov. ; lower long-stalked—Spikes in pairs, axill.]

Gen. VIII.

- i. St. procumb., prickly—Ls. obl., hisp., lower-stalked ; upper sess.—Fls. sol., axill.]

Gen. IX.

- i. Ls. downy ; upper narrower at base.
- ii. Ls. glab. above, scab. beneath ; upper ls. less narrowed at base.

Gen. X.

- i. Ls. lanc., rarely spotted.
- ii. Ls. ov.-lanc., cord., spotted.]

Gen. XI.

- i. St. simp.—Ls. lanc. ; st. ls. narrowed below—Spikes short, lat.
- ii. St. branched—Lower ls. obl., stalked ; upper cord., semi-amplex. long, panicked.

ORD. LXI.—LENTIBULARIACEÆ.

Gen. I.

- i. Spur subu., shorter than the veined cor.—Segm. of cor. contiguous.
- ii. Spur subu., as long as the veinless cor.—Segm. of cor. separated.

ORD. LXI.—LENTIBULARIACEÆ—*cont.*

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
iii. <i>P. alpina</i> , <i>L.</i>	<i>Alpine B.</i>	Scs. bogs!	<i>p</i> Y.	2-4 in.	P.	6-7
iv. <i>P. lusitanica</i> , <i>L.</i>	<i>pale B.</i>	Bogs W. of E.	<i>p</i> Y.	2-4 in.	P.	6-10
Gen. II. Utricularia, <i>L.</i>	<i>Bladder-wort.</i>					
i. <i>U. vulgaris</i> , <i>L.</i>	<i>greater B.</i>	Pools . .	Y.	Fl.	P.	6-7
v. 2. <i>neglecta</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
ii. <i>U. intermedia</i> , <i>Hayne</i>	. . . . .	Pools! . .	Y.	Fl.	P.	6-7
iii. <i>U. minor</i> , <i>L.</i>	<i>lesser B.</i>	Pools . .	<i>p</i> Y.	Fl.	P.	6-7

ORD. LXII.—PRIMULACEÆ.

Gen. I. Primula, <i>L.</i>	<i>Primrose, Cowslip, &c.</i>					
i. <i>P. vulgaris</i> , <i>L.</i>	<i>common P.</i>	Woods .	<i>p</i> Y.	4-8 in.	P.	4-5
ii. <i>P. elatior</i> , <i>Jacq.</i>	<i>Oxlip.</i>	Copses E. of E.	<i>p</i> Y.	4-12 in.	P.	4-5
iii. <i>P. veris</i> , <i>L.</i>	<i>Cowslip.</i>	Past. . .	Y.	6-12 in.	P.	5-6
iv. <i>P. farinosa</i> , <i>L.</i>	<i>Bird's Eye P.</i>	Mount. past!	Li.	2-8 in.	P.	5-7
v. <i>P. scotica</i> , <i>Hook.</i>	<i>Scottish P.</i>	Sc. sea sh.	Li.	2-4 in.	P.	6-7
Gen. II. Cyclamen, <i>L.</i>	<i>Sow-bread.</i>					
[i. <i>C. hederifolium</i> , <i>Willd.</i>	<i>Sow-bread</i>	Woods! Suffolk.	Pk.	4-8 in.	P.	9
Gen. III. Trientalis, <i>L.</i>	<i>Chickweed Winter-green.</i>					
i. <i>T. europœa</i> , <i>L.</i>	<i>European C.</i>	Sc. woods	W.	4-6 in.	P.	6
Gen. IV. Hottonia, <i>L.</i>	<i>Water Violet.</i>					
i. <i>H. palustris</i> , <i>L.</i>	<i>common W., or Featherfoil.</i>	Pools . .	<i>p</i> Pu.	Fl.	P.	5-6
Gen. V. Lysimachia, <i>L.</i>	<i>Loosestrife.</i>					
i. <i>L. vulgaris</i> , <i>L.</i>	<i>great yell. L.</i>	Riv. banks	Y.	2-3 ft.	P.	7-8
ii. <i>L. thyrsiflora</i> , <i>L.</i>	<i>tufted L.</i>	Mar.! . .	Y.	1-3 ft.	P.	7-8
iii. <i>L. nummularia</i> , <i>L.</i>	<i>creeping L., or Money-wort.</i>	Damp pl.	Y.	Cr.	P.	6-7
iv. <i>L. nemorum</i> , <i>L.</i>	<i>Wood L., or yellow Pimpernel.</i>	Sh. pl. .	Y.	3-12 in.	P.	5-8
Gen. VI. Anagallis, <i>L.</i>	<i>Pimpernel.</i>					
i. <i>A. arvensis</i> , <i>L.</i>	<i>scarlet P., or poor man's Weather- glass.</i>	Corn fie..	Sc.	6-18 in.	A.	5-11

ORD. LXI.—LENTIBULARIACEÆ—*cont.*

CHARACTERISTICS OF SPECIES AND VARIETIES.

- iii. Spur conical, shorter than the cor.—Scape glab.
- iv. Spur cylind., obt., decurved, shorter than cor.—Seg. of cor. equal—Scape hairy.

Gen. II.

- i. Spur conic.—Bladders sess. on the ls.—Upper lip of cor. as long as palate.
- v. 2. *Slenderer*—*Pedi. longer*—*Upper lip of cor. three times as long as palate (Essex)*.
- ii. Spur conic.—Bladders on the leafless branches—Upper lip of cor. twice as long as palate.
- iii. Spur short, obt.—Bladders on the ls.—Upper lip of cor. as long as palate.

ORD. LXII.—PRIMULACEÆ.

Gen. I.

- i. Ls. wrinkled—*Pedi. single flowered*; limb of cor. flat.
- ii. Ls. wrinkled—Fls. umbell., erect—Cal. teeth lanc., acu.
- iii. Ls. wrinkled—Fls. umbell., drooping—Cal. teeth short, ov.
- iv. Ls. mealy—Seg. of cor. distant, as long as tube—Cal. teeth lin.
- v. Ls. mealy—Seg. of cor. contiguous, half as long as tube—Cal. teeth short, ov obt.

Gen. II.

- i. Ls. cord., ang., finely toothed, blotched.]

Gen. III.

- i. St. with ls. mostly at top—Ls. obl., ob.-ov.—Fls. on slender stalks.

Gen. IV.

- i. Ls. submerged, pecti.—Fl. st. rising from water; fls. whorled.

Gen. V.

- i. St. erect—Ls. ov.-lanc., opp., or whorled—Fls. in term. and axill. pan.
- ii. St. erect—Ls. lanc., opp.—Fls. in axill., dense rac.
- iii. St. prost.—Ls. orb., obt.—Fls. axill., sol.—Sep. ova., acu.
- iv. St. prost.—Ls. ova., acu.—Fls. axill., sol.—Sep. lin.

Gen. VI.

- i. Ls. ov., sess.—Pet. scarcely longer than cal., fringed.

ORD. LXII.—PRIMULACEÆ.—*cont.*

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
<i>a. phoenicea</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
<i>b. pallida</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
<i>c. cærulea</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
11. <i>A. tenella</i> , <i>L.</i>	<i>Bag P.</i>	Bogs . .	Pk.	3-4 in.	P.	7-8
Gen. VII. Centunculus, <i>L.</i>	<i>Bastard Pimper- nel, or Chaff- weed.</i>					
i. <i>C. minimus</i> , <i>L.</i>	<i>small C., or B.</i> . .	San. hea.	<i>p</i> Ro.	1-2 in.	A.	6-7
Gen. VIII. Samolus, L.	<i>Brookweed.</i>					
i. <i>S. Valerandi</i> , <i>L.</i>	<i>B., or Water-pim- pernel.</i>	Wat. pl..	W.	8-10 in.	P.	6-9
Gen. IX. Glaux, L.	<i>Sea Milk-wort.</i>					
i. <i>G. maritima</i> , <i>L.</i>	<i>Sea M., or black Saltwort.</i>	Mud sh. .	Ro.	3-6 in.	B.	6-7

ORD. LXIII.—PLUMBAGINACEÆ.

Gen. I. Armeria, Willd. . .	<i>Thrift, Sea Pink.</i>					
i. <i>A. maritima</i> , <i>Auct.</i> . .	<i>common T., or S.</i>	Mud sh. & mount.	Ro.	3-12 in.	P.	4-9
v. 2. <i>planifolia</i> , <i>Sy.</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
11. <i>A. plantaginea</i> , <i>Willd.</i>	<i>Plaintain-leaved I.</i>	Jersey .	<i>p</i> Pu.	6-18 in.	P.	6-7
Gen. II. Statice, L.	<i>Sea Lavender.</i>					
i. <i>S. Limonium</i> , <i>L.</i>	<i>common S.</i>	Mud sh. .	<i>b</i> Pu.	4-12 in.	P.	7-8
<i>a. Behen</i> , <i>Fries.</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
<i>b. pyramidalis</i> , <i>Sy.</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
ii. <i>S. bahusiensis</i> , <i>Fries.</i>	<i>remote-flowered S.</i>	Mud sh. .	<i>b</i> Pu.	4-12 in.	P.	7-8
iii. <i>S. binervosa</i> , <i>G. E. Sm.</i>	<i>upright-stalked S.</i>	Rocky sh.	<i>p</i> Pu.	4-10 in.	P.	7-8
<i>a. occidentalis</i> , <i>Lloyd.</i> . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
<i>b. intermedia</i> , <i>Syme.</i> . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
<i>c. Dodartii</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
iv. <i>S. caspia</i> , <i>Willd.</i>	<i>matted S.</i>	Salt mar. E. of E.	<i>p</i> Li.	4-8 in.	P.	7-8

ORD. LXII.—PRIMULACEÆ—*cont.*

CHARACTERISTICS OF SPECIES AND VARIETIES.

- a.* Cor. scarlet.
- b.* Cor. white, with a purple eye.
- c.* Cor. blue.
- ii.** Ls. round, stalked—Pet. much longer than cal., ent., not fringed.

Gen. VII.

- i.** Ls. ov., alt., acu.—Fls. sol., axill., sess.

Gen. VIII.

- i.** Ls. ob.-ov., obt.; fleshy—Rac. many-flowered; erect.

Gen. IX.

- i.** St. procumb.—Ls. ov., opp., glab., fleshy—Fls. sess., axill.

ORD. LXIII.—PLUMBAGINACEÆ.

Gen. I.

- i.** Ls. lin., one-nerved—Cal. teeth short.
- v.* **2.** *Ls. broader, flat.*
- ii.** Ls. lan. 3-5 nerved—Cal. teeth long, setaceous.

Gen. II.

- i.** Ls. ellip.-lanc., one-ribbed—Scape branched from above the middle—Spikes short, dense flowered.
  - a.* Spikes lax on a compact, level topped corymb., with short stout branches.
  - b.* Spikes lax on a spread., pyram. corymb., with flexuous branches.
- ii.** Ls. obl.-lanc., one-ribbed—Scape branched from near the base—Spikes long; fls. distant.
- iii.** Ls. spath., narrowed into the stalk—Scape branched from near the middle—Branches nearly all flowering.
  - a.* Ls. ob. ov.-lanc.—Lower branches of pan. sometimes flowerless; spikes slender.
  - b.* Ls. as Var. *a*—Branches all flowering; spikes dense.
  - c.* Ls. spath.—Branches all flowering; spikes dense, imbric.
- iv.** Ls. spath.—Scape branched from base; upper branches flowering, lower barren, tangled—Fls. small, crowded.

ORD. LXIV.—PLANTAGINACEÆ.

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
Gen. I. Plantago, L.	<i>Plantain.</i>					
i. <i>P. major, L.</i>	<i>greater P.</i>	Past. . .	G.	2-5 in.	P.	6-8
v. 2. <i>microstachya</i>	. . . . .	. . . . .	. . .	. . .	. . .	. . .
ii. <i>P. media, L.</i>	<i>hoary P.</i>	Ch. pas..	G.	6-12 in.	P.	6-10
iii. <i>P. lanceolata, L.</i>	<i>Ribwort P.</i>	Past. . .	G.	2-12 in.	P.	6-7
v. 2. <i>Timbali</i>	. . . . .	. . . . .	. . .	. . .	. . .	. . .
iv. <i>P. maritima, L.</i>	<i>Sea-side P.</i>	Sea sh. & Sc. mount.	G.	1-12 in.	P.	6-9
v. <i>P. Coronopus, L.</i>	<i>Bucks-horn P.</i> . . .	Dry pl. .	G.	1-8 in.	A.	6-7
Gen. II. Littorella, L. . . .	<i>Shore-weed.</i>					
i. <i>L. lacustris, L.</i>	<i>Plantain S.</i>	San. sh. of lakes.	W.	2-3 in.	P.	6-7

ORD. LXV.—AMARANTHACEÆ.

Gen. I. Amaranthus, L.	<i>Amaranth.</i>					
[i. <i>A. Blitum, L.</i>	<i>wild A.</i>	Was. pl..	. .	1-2 ft.	A.	8

ORD. LXVI.—CHENOPODIACEÆ.

Gen. I. Chenopodium, L.	<i>Goose-foot.</i>					
<i>A.</i> *						
i. <i>C. olidum, Curt.</i>	<i>stinking G.</i>	Was. pl..	. .	6-18 in.	A.	8-9
ii. <i>C. polyspermum, L.</i> . . .	<i>many-seeded G.</i> . .	Was. pl..	. .	6-18 in.	A.	8-9
a. <i>acutifolium, Sm.</i>	. . . . .	Cornwall!	. .	. .	. .	. .
b. <i>obtusifolium</i>	. . . . .	(common)	. .	. .	. .	. .
* *						
†						
iii. <i>C. urbicum, L.</i>	<i>upright G.</i>	Was. pl..	. .	1-3 ft.	A.	8-9
v. 2. <i>intermedium</i>	. . . . .	. . . . .	. .	. .	. .	. .
iv. <i>C. murale, L.</i>	<i>Nettle-leaved G.</i> . .	Was. pl..	. .	6-18 in.	A.	8-9
v. <i>C. hybridum, L.</i>	<i>Maple-leaved G.</i> . .	Was. pl..	. .	1-3 ft.	A.	8-9
† †						
vi. <i>C. album, L.</i>	<i>white G.</i>	Was. pl..	. .	1-3 ft.	A.	7-9
a. <i>candicans</i>	. . . . .	. . . . .	. .	. .	. .	. .
b. <i>viride, L.</i>	. . . . .	. . . . .	. .	. .	. .	. .
c. <i>virens</i>	. . . . .	. . . . .	. .	. .	. .	. .
vii. <i>C. filicifolium, Sm.</i> . . .	<i>Fig-leaved G.</i> . . .	. . . . .	. .	. .	. .	. .

ORD. LXIV.—PLANTAGINACEÆ.

CHARACTERISTICS OF SPECIES AND VARIETIES.

Gen. I.

- i. Ls. broadly ov., on long grooved stalks—Scape round—Spike long; sep. with a prominent rib.
- v. 2. *Scape short—Spike 3-6 flowered (Connemara).*
- ii. Ls. ellip., sess., or shortly stalked—Scape round—Spike cylind.; sep. not keeled.
- iii. Ls. lanc., five-ribbed, narrowed at each end—Scape furrowed—Spike glob.
- v. 2. *Sep. and bracts with silvery margins.*
- iv. Ls. lin., fleshy—Scape round—Spike cylind.; sep. not keeled.
- v. Ls. lin., pinnfid.—Scape round, hairy—Spike cylind., slend.

Gen. II.

- i. Ls. all rad., lin., fleshy—Fert. fls. sess.; barren fls. sol., stalked, axill.

ORD. LXV.—AMARANTHACEÆ.

Gen. I.

- i. St. decumb.—Ls. cb.-ov., long-stalked—Fls. minute, in small axill. clusters.]

ORD. LXVI.—CHENOPODIACEÆ.

Gen. I.

A. Seeds horizontal.

* *Ls. ent.*

- i. Ls. stalked, ov., mealy, fetid—Fls. in dense, leafless spikes.
- ii. Ls. sess., ov.-ellip.—Fls. in long, axill., leafless rac.
- a. St. ascend.—Ls. acute.
- b. St. decumb.—Ls. obtuse.
- * * *Ls. angled, lobed or toothed.*
- † *Ls. green beneath.*
- iii. Ls. triang., toothed, their base narrowed into the stalk—Spikes long, erect—Fls. in small, remote clusters.
- v. 2. *Ls. with large acute teeth.*
- iv. Ls. ov.-rhomb. sharply and unequally toothed—Branches of spike spreading; fls. cymose; seeds keeled.
- v. Ls. cord., angled, teeth large, distant—Spikes in axill., lax corymbs—Seeds not keeled.
- † † *Ls. glauc. or mealy beneath.*
- vi. Ls. ov.-rhomb., ent at base, mealy; upper lanc. ent.—Spikes leafy, branched; with distant clusters.
- a. Ls. mealy, toothed—Spikes dense, simp.
- b. Ls. sparingly mealy, nearly ent.—Spikes lax, branched.
- c. Lower ls. bluntly serr., mealy below—Spikes lax, erect.
- vii. Ls. ov.-obl. unequally three-lobed; base wedge-shaped—Upper lin.-lanc.—Spike leafy at base, with erect branches.

ORD. LXVI.—CHENOPODIACEÆ—cont.

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
<i>B.</i>						
viii. <i>C. rubrum</i> , <i>L.</i>	<i>red G.</i>	Was. pl..	..	1-3 ft.	A.	8-9
v. 2. <i>pseudo-botryoides</i>	. . . . .	Sea co. .	..	..	..	..
ix. <i>C. botryoides</i> , <i>Sm.</i>	. . . . .	San. sh. .	..	4-5 in.	A.	8-9
x. <i>C. glaucum</i> , <i>L.</i>	<i>Oak-leaved G.</i>	Was. pl.!	..	6-18 in.	A.	8-9
xi. <i>C. Bonus-Henricus</i> , <i>L.</i>	<i>Mercury G., or Good King Henry.</i>	Was. pl..	..	1-3 ft.	P.	6-8
Gen. II. Obione , <i>Gaert.</i> <i>Sea-Purslane.</i>						
i. <i>O. portulacoides</i> , <i>Moq.</i>	<i>Shrubby S.</i>	Mud. sh.	..	1-2 ft.	P.	8-10
ii. <i>O. pedunculata</i> , <i>Moq.</i>	<i>Stalked S.</i>	Mud. sh.	..	3-8 in.	A.	8-10
Gen. III. Atriplex , <i>L.</i> <i>Orache.</i>						
<i>A.</i>						
i. <i>A. arenaria</i> , <i>Woods</i> .	<i>Frosted Sea O.</i> .	San. sh. .	..	4-10 in.	A.	7-10
<i>B.</i>						
*						
ii. <i>A. Babingtonii</i> , <i>Woods</i>	<i>spreading-fruited O.</i>	Sea sh. .	..	½-3 ft.	A.	7-9
iii. <i>A. hastata</i> , <i>L.</i>	<i>Halberd-leaved O.</i>	Was. pl..	..	½-3 ft.	A.	7-9
a. <i>hastata</i>	. . . . .	. . . . .	..	..	..	..
b. <i>deltoidea</i> , <i>Bab.</i>	. . . . .	. . . . .	..	..	..	..
iv. <i>A. paluta</i> , <i>L.</i>	. . . . .	Was. pl..	..	1-3 ft.	A.	7-9
a. <i>angustifolia</i>	. . . . .	. . . . .	..	..	..	..
b. <i>erecta</i>	. . . . .	. . . . .	..	..	..	..
* *						
v. <i>A. littoralis</i> , <i>L.</i>	<i>Grass-leaved O.</i> .	Salt mar.	..	1-2 ft.	A.	7-9
v. 2. <i>marina</i>	. . . . .	. . . . .	..	..	..	..
Gen. IV. Beta , <i>L.</i> <i>Beet.</i>						
i. <i>B. maritima</i> , <i>L.</i>	<i>common B.</i>	Mud. sh.	..	1-2 ft.	P.	6-10
Gen. V. Salsola , <i>L.</i> <i>Saltwort.</i>						
i. <i>S. Kali</i> , <i>L.</i>	<i>prickly S.</i>	San. sh. .	..	6-18 in.	A.	7
Gen. VI. Suaeda , <i>Forsk.</i> <i>Sea-blite.</i>						
i. <i>S. maritima</i> , <i>Dum.</i>	<i>annual S.</i>	Salt mar.	..	3-18 in.	A.	7-9
ii. <i>S. fruticosa</i> , <i>Forsk.</i>	<i>Shrubby S.</i>	San. sh.!	..	1-3 ft.	P.	7-9
Gen. VII. Salicornia , <i>L.</i> <i>Glass-wort.</i>						
i. <i>S. herbacea</i> , <i>L.</i>	<i>jointed-G.</i>	Salt mar.	..	6-18 in.	A.	8-9
v. 2. <i>procumbens</i> , <i>Sm.</i>	. . . . .	. . . . .	..	..	..	..
ii. <i>S. radicans</i> , <i>Sm.</i>	<i>creeping G.</i>	Salt mar.!	..	6-18 in.	P.	8-9

ORD. LXVI.—CHENOPODIACEÆ—*cont.*

CHARACTERISTICS OF SPECIES AND VARIETIES.

B. Seeds vertical.

- viii. Ls. all triang., irregularly toothed, glab.—Spikes erect, leafy—Fls. crowded.
 v. 2. *Small, reddish—Ls. rhomb., spikes shorter.*
 ix. St. prost.—Ls. triang., scarcely toothed, fleshy—Fls. in dense, leafy cymes.
 x. St. prost.—Ls. all obl., toothed, mealy beneath—Spikes erect, simp., leafless.
 xi. Perenn.—Ls. triang.-hast., acu.—Fls. in term, compound spikes.

Gen. II.

- i. Shrubby—Ls. ov.-lanc., ent., silvery white—Fr. sess.
 ii. Herb—Ls. ob.-ov., ent., mealy—Fr. long-stalked.

Gen. III.

A. St. buff.

- i. Ls. triang.-rhomb., lacin., silvery white—Fls. in axill., sess. clusters.

*B. St. green, with reddish stripes.*** Lower ls. with lateral lobes.*

- ii. St. procumb.—Ls. ov.-triang., pale green, mealy ; upper lanc., three-lobed at base—Fls. in axill. and term clusters.
 iii. Ls. hast.-triang. ; lobes spreading—Fls. densely panicled, in simp. spikes.
 a. Upper ls. not hast.—Spikes lax.
 b. Upper ls. hast.—Spikes dense.
 iv. Ls. with two ascending lobes from a cuneate base, upper ent.—Fls. in simple, interrupted spikes.
 a. St. procumb.—Lower ls. ent.—Spikes long, lax.
 b. St. erect—Lower ls. sinu.—Spikes short, dense.
* * Ls. all ent.
 v. Ls. lin.-lanc.—Fls. in axill., sess. clusters.
 v. 2. *Ls. ov.-lanc., serr.*

Gen. IV.

- i. Sts. many, procumb. at base—Ls. ov.-cun., fleshy, narrowed into the stalk—Spikes long, simp., leafy.

Gen. V.

- i. St. prost.—Ls. subu., fleshy, spiny—Fls. axill., sol.

Gen. VI.

- i. St. herb., ann.—Ls. acute.
 ii. St. shrubby, perenn.—Ls. obt.

Gen. VII.

- i. St. erect, herb, leafy ; its joints thickened upwards—Spikes cylind., tapering.
 v. 2. *St. procumb.*
 ii. St. procumb., woody, leafless ; its joints scarcely thickened—Spikes obt.

ORD. LXVII.—POLYGONACEÆ.

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
Gen. I. Polygonum, L.	<i>Persicaria, Bistorta, Knot-grass, Buckwheat.</i>					
<i>A.</i>						
i. <i>P. Bistorta, L.</i>	<i>common B., or Snake-weed.</i>	Moi. mead.	Fl.	12-18 in.	P.	6-9
ii. <i>P. viviparum, L.</i>	<i>alpine B.</i>	Mount. past.!	♂ Fl.	4-8 in.	P.	6-7
<i>B.</i>						
iii. <i>P. amphibium, L.</i>	<i>amphibious B.</i>	Dit. & mud. pl.	Ro.	2-3 ft.	P.	7-8
<i>a. aquaticum</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
<i>b. terrestre</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
iv. <i>P. lapathifolium, L.</i>	<i>pale-flowered P.</i>	Was. pl..	gr Pk.	1-4 ft.	A.	7-8
<i>v. 2. laxum</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
v. <i>P. Persicaria, L.</i>	<i>spotted P.</i>	Was. pl..	gr Ro.	1-2 ft.	A.	7-10
<i>* *</i>						
vi. <i>P. mite, Schrank.</i>	<i>lax-flowered P.</i>	Wet pl.!	Pk.	1-2 ft.	A.	7-9
vii. <i>P. Hydropiper, L.</i>	<i>biting P.</i>	Wet pl. .	Pk.	1-3 ft.	A.	8-9
viii. <i>P. minus, Huds.</i>	<i>creeping P.</i>	Mar. ! . .	Pk.	6-18 in.	A.	8-9
<i>C.</i>						
ix. <i>P. aviculare, L.</i>	<i>common K.</i>	Was. pl..	Var.	1-3 ft.	A.	7-9
x. <i>P. Raii, Bab.</i>	<i>Ray's K.</i>	San. sh. .	Var.	1-3 ft.	A.	7-9
xi. <i>P. maritimum, L.</i>	<i>Sea-side K.</i>	Sea sh. .	Var.	1-3 ft.	P.	8-9
<i>D.</i>						
xii. <i>P. Convolvulus, L.</i>	<i>climbing B.</i>	Corn fie..	Gr.	1-4 ft.	A.	7-9
<i>v. 2. pseudo-dumetorum.</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
xiii. <i>P. dumetorum, L.</i>	<i>Copse B.</i>	Thick. .	Gr.	1-4 ft.	A.	8-9
Gen. II. Rumex, L.	<i>Dock, Sorrel.</i>					
<i>A.</i>						
<i>*</i>						
i. <i>R. obtusifolius, L.</i>	<i>broad-leaved D.</i>	Was. pl..	r G.	2-3 ft.	P.	7-9
ii. <i>R. pratensis, M. & K.</i>	<i>Meadow D.</i>	Mar. . .	r G.	2-3 ft.	P.	7-9
iii. <i>R. pulcher, L.</i>	<i>Fiddle D.</i>	Waysides	r G.	6-18 in.	P.	6-10
iv. <i>R. maritimus, L.</i>	<i>golden D.</i>	Mar. ! . .	y G.	1-2 ft.	P.	7-8
v. <i>R. palustris, Sm.</i>	<i>yellow Marsh D.</i>	Mar. ! . .	y G.	1-2 ft.	P.	7-8
<i>* *</i>						
vi. <i>R. crispus, L.</i>	<i>curled D.</i>	Was. pl..	r G.	2-3 ft.	P.	6-8
vii. <i>R. nemorosus, Schrad.</i>	. . . . .	Sh. pl. .	r G.	1-4 ft.	P.	7-8
<i>a. viridis, Sibth.</i>	<i>green-veined D.</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
<i>b. sanguineus, L.</i>	<i>bloody-veined D.</i>	Woods ! .	. . . . .	. . . . .	. . . . .	. . . . .

ORD. LXVII.—POLYGONACEÆ.

CHARACTERISTICS OF SPECIES AND VARIETIES.

Gen. I.

A. St. simple—*Spikes term.*

i. Ls. ov., sub-cord., waved; rad. ls. with a winged stalk—Spikes dense.

ii. Ls. lin.-lanc., margins revolute—Spikes lax.

B. St. branched—*Fls. racemed or paniced.*** Spike short, dense, not leafy at base.*

iii. Ls. lanc., stalked—Spike dense, sol. term.

a. Floating—Ls. broadly lanc., glab.

b. Creeping—Ls. narrow lanc., with adpr. hairs.

iv. Ls. ov.-lanc., not spotted—Spikes term. and lat.—Pedunc. and cal. gland., scab; sty. free—Lower stip. not fringed.

v. 2. *Spikes drooping in pairs*—*Joints of st. tumid.*—*Stip. slightly fringed.*

v. Ls. lanc., usually spotted—Spikes term. and axill.—Pedunc. and cal. glab.—Sty. half united—Stip. fringed.

* * Spike lax, interrupted, leafy at base.

vi. Spikes erect—Ls. lanc., wavy—Stip. hairy.

vii. Spikes drooping—Ls. lanc., wavy—Stip. glab.—Stig. half united.

viii. Spikes erect—St. procumb.—Ls. lin.-lanc., flat—Sty. united.

C. St. procumb.—*Fls. axill.*

ix. St. herb—Ls. flat—Stip. shorter than the internodes, with few distant veins—Fr. shorter than cal.

x. Habit of *P. aviculare*, but fr. longer than cal.

xi. St. woody—Ls. fleshy, glauc., margins revolute—Stip. as long as internodes, with many branched veins—Fr. longer than cal.

D. St. climbing—*Fls. racemose.*

xii. St. ang.—Segm. of cal. obtusely keeled.

v. 2. *Seg. of cal. winged.*

xiii. St. round—Segm. of cal. winged.

Gen. II.

A. Fls. perfect—*Ls. not acid.*** Inner sep. strongly toothed.*N.B. *The inner enlarged sep. are alluded to in all cases in the Genus Rumex.*

i. Ls. cord.-ov., blunt; upper lanc.—Whorls dist., leafless above—Sep. ov. with a long point; upper tubercled.

ii. Ls. obl.-lanc., acu.; lower rather cord.—Whorls distinct, leafless—Sep. cord., unequal, with a small triang. point; upper one tubercled.

iii. St. procumb.—Lower ls. fiddle-shaped; upper lanc. ac.—Whorls distant, leafy—Sep. triang.-ov., veined; one tubercled.

iv. Ls. lin.-lanc.—Whorls crowded, leafy—Sep. delt., with 2-4 setaceous teeth on each side, and a narrow tubercle.

v. Ls. lin.-lanc.—Whorls dist., leafy—Sep. ov.-obl., with 2-3 short setaceous teeth on each side, and a narrow tubercle.

* * Inner sep. ent., or slightly toothed.

vi. Ls. lanc., acu., wavy—Whorls crowded, leafless—Sep. cord., upper one tubercled.

vii. Ls. ov.-lanc.; lower cord.—Whorls distant, leafless—Sep. obl., obt., upper tubercled.

a. Ls. veined with green.

b. Ls. veined with red.

ORD. LXVII.—POLYGONACEÆ—*cont.*

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
viii. <i>R. conglomeratus</i> , <i>Mur.</i>	<i>sharp D.</i>	Wet pl. .	r G.	1-3 ft.	P.	7-8
ix. <i>R. Hydrolapathum</i> , <i>Huds.</i>	<i>great Water D.</i>	Riv. sides	r G.	3-6 ft.	P.	7-8
x. <i>R. aquaticus</i> , <i>L.</i> . . .	<i>grainless Water D.</i>	Mar. . .	r G.	1-3 ft.	P.	7-8
[xi. <i>R. alpinus</i> , <i>L.</i> . . . . <i>B.</i>	<i>Alpine D., or Monk's-Rhubarb.</i>	Sco ! . .	r G.	2-4 ft.	P.	7-8
xii. <i>R. Acetosa</i> , <i>L.</i> . . .	<i>common S.</i>	Mead. . .	puR.	1-2 ft.	P.	5-7
xiii. <i>R. Acetosella</i> , <i>L.</i> . .	<i>Sheep's S.</i>	Dry pas.	puR.	3-12 in.	P.	5-7
Gen. III. Oxyria, Hill.						
i. <i>O. reniformis</i> , <i>Hook.</i> .	<i>Mountain Sorrel.</i> <i>Mountain S.</i> . . .	Mount. rocks.	r G.	6-10 in.	P.	7-8

ORD. LXVIII.—ELEAGNACEÆ.

Gen. I. Hippophaë, L.						
i. <i>H. rhamnoides</i> , <i>L.</i> . .	<i>Sea Buck-thorn.</i> <i>Sea B.</i>	San. cliffs!	G.	1-8 ft.	Shb.	5-6

ORD. LXIX.—THYMELACEÆ.

Gen. I. Daphne, L.						
	<i>Mezereon, Spurge Laurel.</i>					
i. <i>D. mezereum</i> , <i>L.</i> . . .	<i>common M.</i>	Woods ! .	p R.	1-3 ft.	Shb.	3-4
ii. <i>D. laureola</i> , <i>L.</i>	<i>common S.</i>	Woods .	y G.	2-4 ft.	Shb.	2-4

ORD. LXX.—SANTALACEÆ.

Gen. I. Thesium, L.						
	<i>Bastard Toad- flax.</i>					
i. <i>T. humifusum</i> , <i>L.</i> . . .	<i>Lint-leaved B.</i> . .	Ch. downs	W.	6-12 in.	P.	5-7

ORD. LXXI.—ARISTOLOCHIACEÆ.

Gen. I. Asarum, L.						
	<i>Asarabacca.</i>					
i. <i>A. europæum</i> , <i>L.</i> . . .	<i>A.</i>	Woods! N. of E.	g B.	Cr.	P.	5

ORD. LXVII.—POLYGONACEÆ—*cont.*

CHARACTERISTICS OF SPECIES AND VARIETIES.

- viii. Ls. ov.-lanc., not contracted above the base—Whorls distant, leafy—Sep. lin.-obl., blunt, each tubercled.
- ix. Ls. lanc., ac., ent., tapering into the stalk—Whorls crowded, leafless—Sep. ov.-triang., all tubercled.
- x. Ls. lan., lower cord., wavy—Whorls crowded, leafless—Sep. cord., membranous, not tubercled.
- xi. Ls. roundish-cord., blunt; stalk furrowed; upper ov.—Sep. cord., ov., not tubercled.]
- B. Flowers diacious—Leaves acid.*
- xii. Ls. obl.-sagitt.—Outer sep. reflexed—Inner enlarged, slightly tubercled.
- xiii. Lower ls. hast.—Outer sep. ascend.—Inner hardly enlarged, not tubercled.

Gen. III.

- i. Ls. rad., renif., long-stalked—Pan. leafless, slender—Pedi. thickened upwards.

ORD. LXVIII.—ELEAGNACEÆ.

Gen. I.

- i. Thorny—Ls. lin.-lanc. silvery—Fls. small, appearing with the young ls.

ORD. LXIX.—THYMELACEÆ.

Gen. I.

- i. Ls. decid.—Fls. lat., sess. in threes.
- ii. Ls. evergreen—Fls. in axill., few-flowered racemes.

ORD. LXX.—SANTALACEÆ.

Gen. I.

- i. Ls. lin.-lanc., one-nerved—Fls. in leafy, paniced racemes.

ORD. LXXI.—ARISTOLOCHIACEÆ.

Gen. I.

- i. Ls. reni., obt., in pairs, in the axills of which is a sol. drooping fl.

ORD. LXXI.—ARISTOLOCHIACEÆ—cont.

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
Gen. II. Aristolochia, L.	<i>Birthwort.</i>					
i. <i>A. Clematitis, L.</i> . . .	<i>common B.</i> . . .	Ruins . .	g Y.	3-6 in.	P.	7-8

ORD. LXXII.—EMPETRACEÆ.

Gen. I. Empetrum, L.	<i>Crowberry.</i>					
i. <i>E. nigrum, L.</i>	<i>Black C., or Crake- berry.</i>	Mount. hths.	Pu.	6-18 in.	Shb.	5-6

ORD. LXXIII.—EUPHORBIACEÆ.

Gen. I. Euphorbia, L.	<i>Spurge.</i>					
A.						
i. <i>E. Peplis, L.</i>	<i>purple S.</i>	San. sh.!	Y.	6-12 in.	A.	7-9
B.		S. of E.				
*						
ii. <i>E. Helioscopia, L.</i> . . .	<i>Sun S.</i>	Was. pl.	g Y.	6-18 in.	A.	6-10
iii. <i>E. platyphylla, L.</i> . . .	<i>broad-leaved Warted S.</i>	Corn fie.!	g Y.	½-3 ft.	A.	7-10
		S. of E.				
v. 2. <i>stricta</i>		Li. woods,	. .	. .	. .	. .
		Glos. & Mon.				
iv. <i>E. hiberna, L.</i>	<i>Irish S.</i>	Thick.!	g Y.	1-2 ft.	P.	5-6
v. <i>E. pilosa, L.</i>	<i>Marsh S.</i>	Sh. pl.!	g Y.	2-4 ft.	P.	5-6
[vi. <i>E. coralloides, L.</i> . .	<i>Coral-like hairy S.</i>	Hedges, Slinfold.!	g Y.	1-2 ft.	P.	5-6
* *						
†						
vii. <i>E. amygdaloides, L.</i> . .	<i>Wood S.</i>	Thick. .	Y.	6-12 in.	P.	3-5
† †						
[viii. <i>E. Esula, L.</i>	<i>leafy-branched S.</i>	Woods !	Y.	1-2 ft.	P.	7-8
		Sco.				
[ix. <i>E. Cyparissias, L.</i> . . .	<i>Cypress S.</i>	Woods !.	Y.	1-2 ft.	P.	6-7
x. <i>E. Paralias, L.</i>	<i>Sea S.</i>	San. sh. .	Y.	1-2 ft.	P.	7-10
xi. <i>E. portlandica, L.</i> . . .	<i>Portland S.</i>	San. sh. .	Y.	6-18 in.	P.	5-9
xii. <i>E. exigua, L.</i>	<i>dwarf S.</i>	Corn fie. .	Y.	6-12 in.	A.	7-10
xiii. <i>E. Peplus, L.</i>	<i>petty S.</i>	Was. pl. .	Y.	6-12 in.	A.	7-11
* * *						
[xiv. <i>E. Lathyris, L.</i>	<i>Caper S.</i>	Thick.!	Pu.	3-4 ft.	P.	6-7
Gen. II. Buxus, L.	<i>Box.</i>					
i. <i>B. sempervirens, L.</i> . .	<i>common B.</i>	Ch. hills !	g W.	6-12 ft.	Shb.	4-6

ORD. LXXI.—ARISTOLOCHIACEÆ.—*cont.*

CHARACTERISTICS OF SPECIES AND VARIETIES.

Gen. II.

- i. St. erect, simp.—Ls. cord., stalked, glab.—Fls. crowded, erect.

ORD. LXXII.—EMPETRACEÆ.

Gen. I.

- i. St. procumb.—Ls. lin.-obl. ; margins meeting behind—Fls. axill., small.

ORD. LXXIII.—EUPHORBIACEÆ.

Gen. I.

A. Ls. stipuled.

- i. St. procumb., glauc.—Ls. opp., oblong ; heart-shaped, ent.—Fls. axill., sol.

B. Ls. not stipuled.

* *Ls. alt.—Invol. glands transversely oval.*

- ii. Ls. ob.-ov., serr. upwards—Bracts ob.-cord.—Caps. glab.—Seeds netted.

- iii. Ls. ob.-ov.-lanc., finely serr.—Bracts cord.—Caps. with small warts—Seeds smooth, olive-brown.

- v. 2. *Invol., caps., and seeds smaller—Caps. with prominent warts—Seeds red brown.*

- iv. Hairy—Ls. ellip.-lanc., ent.—Caps. warted, glab.—Seeds smooth.

- v. Pubesc.—Ls. ellip.-lanc., obt., finely serr.—Bracts all ellip., glab.—Caps. warted, hairy.

- vi. Hairy—Ls. lanc., minutely serr.—Bracts ov.-obl., hairy—Caps. not warted, woolly.]

* * *Ls. alt.—Invol. bracts lunate, triang. or two-horned.*

† *Bracts united.*

- vii. Ls. ob.-ov.-lanc., ent.—Caps. glab.—Seeds smooth.

† † *Bracts separate.*

- viii. Ls. obl.-lanc., dentic.—Bracts cord., mucro.—Caps. scab.—Seeds smooth.]

- ix. Ls. lin., ent., almost glauc.—Bracts obt.—Caps. granulate—Seeds smooth.]

- x. Ls. ellip.-obl., lanc., coriac., imbric.—Bracts broadly cord.—Caps. wrinkled—Seeds smooth.

- xi. Ls. ob.-ov.-lanc., blunt, glauc.—Bracts triang.-cord.—Caps. slightly rough—Seeds rough.

- xii. Ls. lin., blunt, or ac.—Bracts lanc., rough—Seeds rough.

- xiii. Ls. broadly ov., stalked—Bracts ov.—Caps. smooth, with rough keels—Seeds rough.

* * * *Ls. opp.*

- xiv. Ls. lin.-obl., decussate ; upper cord.—Caps. smooth.]

Gen. II.

- i. Ls. obl., shining ; stalks rather hairy—Fls. sess.

ORD. LXXIII.—EUPHORBIACEÆ—*cont.*

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
Gen. III. Mercurialis, L.	<i>Mercury.</i>					
i. <i>M. perennis, L.</i> . . .	<i>perennial, or Dog's M.</i>	Sh. pl. .	Gr.	6-18 in.	P.	3-5
v. 2. <i>ovata, Steud.</i> . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
ii. <i>M. annua, L.</i>	<i>annual M.</i> . . .	Was. pl..	Gr.	6-18 in.	A.	7-11
v. 2. <i>ambigua, L. fil.</i> . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .

ORD. LXXIV.—URTICACEÆ.

Gen. I. Urtica, L.	<i>Nettle.</i>					
i. <i>U. urens, L.</i>	<i>small N.</i>	Was. pl..	. .	1-2 ft.	A.	6-9
ii. <i>U. dioica, L.</i>	<i>great N.</i>	Was. pl..	. .	1-4 ft.	P.	6-9
iii. <i>U. pilulifera, L.</i>	<i>Roman N.</i> . . .	Was. pl..	. .	1-2 ft.	A.	6-9
v. 2. <i>Dodartii, L.</i>	. . . . .	. . . . .	. .	. .	. .	. .
Gen. II. Parietaria, L.	<i>Wall-Pellitory.</i>					
i. <i>P. officinalis, L.</i>	<i>common W.</i> . . .	Walls . .	Gr.	1-2 ft.	P.	6-9
a. <i>diffusa</i>	. . . . .	. . . . .	. .	. .	. .	. .
b. <i>erecta</i>	. . . . .	. . . . .	. .	. .	. .	. .
Gen. III. Humulus, L.	<i>Hop.</i>					
i. <i>H. Lupulus, L.</i>	<i>common H.</i> . . .	Hedges .	g Y.	Cl.	P.	7-9

ORD. LXXV.—ULMACEÆ.

Gen. I. Ulmus, L.	<i>Elm.</i>					
i. <i>U. montana, Sm.</i> . . .	<i>Wych E.</i>	Woods .	. .	. .	T.	4-5
ii. <i>U. campestris, Sm.</i> . .	<i>common E.</i> . . .	Woods .	. .	. .	T.	4-5

ORD. LXXVI.—CUPULIFERÆ.

Gen. I. Quercus, L.	<i>Oak.</i>					
i. <i>Q. Robur, L.</i>	<i>British O.</i>	Woods .	. .	. .	T.	4-5
a. <i>pedunculata, Ehr.</i> . . .	. . . . .	. . . . .	. .	. .	. .	. .
b. <i>sessiflora, Sm.</i>	. . . . .	. . . . .	. .	. .	. .	. .
Gen. II. Fagus, L.	<i>Beech.</i>					
i. <i>F. sylvatica, L.</i>	<i>common B.</i> . . .	Woods .	. .	. .	T.	4-5
Gen. III. Carpinus, L.	<i>Hornbeam.</i>					
i. <i>C. Betulus, L.</i>	<i>common H.</i> . . .	Woods .	. .	. .	T.	5-6

ORD. LXXIII.—EUPHORBIACEÆ—*cont.*

CHARACTERISTICS OF SPECIES AND VARIETIES.

Gen. III.

- i. St. simp.—Ls. obl.-lanc., hairy—Fert. fls. stalked.

v. 2. *Ls. broader, sess.*

- ii. St. branched—Ls. ov., glab.—Fertile fls. sess.

v. 2. *Ls. lanc.—Fertile and sterile fls. mixed, whorled.—(Jersey!)*

ORD. LXXIV.—URTICACEÆ.

Gen. I.

- i. Ls. ellip.—Spikes simp., shorter than the peti.

- ii. Ls. ov.-cord.—Spikes branched, longer than the peti.

- iii. Ls. ov., coarsely toothed—Fert. fls. in glob., heads.

v. 2. *Ls. ent.!*

Gen. II.

- i. Ls. ov.-lanc., three-nerved—Fls. in axill. clusters.

- a. St. prost. or ascend.—Invol. three-flowered.

- b. St. erect—Invol. seven-flowered.

Gen. III.

- i. St. long, climb., scab.—Ls. opp., 3-5 lobed, rough.

ORD. LXXV.—ULMACEÆ.

Gen. I.

- i. Seed in the centre of the sub-orb., or obl. samara.

- ii. Seeds above the centre of ob.-ov. samara.

(Many varieties of each occur barely distinguishable.)

ORD. LXXVI.—CUPULIFERÆ.

Gen. I.

- i. Ls. ob.-ov., sinu.-lobed—Catk. pend.

- a. Ls. sess.—Pedunc. long.

- b. Ls. stalked—Pedunc. short.

Gen. II.

- i. Ls. ov., glab., ciliate on margins.

Gen. III.

- i. Ls. ellip.-ov., doubly serr., pil. beneath.

ORD. LXXVI.—CUPULIFERÆ—*cont.*

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
Gen. IV. <i>Corylus</i>, L.	<i>Hazel.</i>					
i. <i>C. Avellana</i> , L. . . .	<i>common H.</i> . . .	Woods .	..	..	T.	3-4

ORD. LXXVII.—BETULACEÆ.

Gen. I. <i>Alnus</i>, Tourn.	<i>Alder.</i>					
i. <i>A. glutinosa</i> , Gærtn. .	<i>common A.</i> . . .	Wet pl. .	..	..	T.	3-4
Gen. II. <i>Betula</i>, L.	<i>Birch.</i>					
i. <i>B. alba</i> , L.	<i>common B.</i> . . .	Woods .	..	..	T.	4-5
a. <i>verrucosa</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
b. <i>glutinosa</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
v. 2. <i>pubescens</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
ii. <i>B. nana</i> , L.	<i>dwarf B.</i>	Sc. mount.!	..	1-2 ft.	Shb.	5

ORD. LXXVIII.—SALICACEÆ.

Gen. I. <i>Populus</i>, L.	<i>Poplar.</i>					
i. <i>P. alba</i> , L.	<i>great white P.</i> .	Moi. woods	..	..	T.	3-4
ii. <i>P. canescens</i> , Sm. . .	<i>gray P.</i>	Moi. woods S. of E.	..	..	T.	3-4
iii. <i>P. tremula</i> , L. . . .	<i>Aspen</i>	Moi. woods	..	..	T.	3-4
iv. <i>P. nigra</i> , L.	<i>black P.</i>	Riv. banks	..	..	T.	3-4
Gen. II. <i>Salix</i>, L.	<i>Willow, Sallow, Osier, &c.</i>					
i. <i>S. triandra</i> , L. . . .	<i>Almond-leaved W.</i>	Riv. banks	..	20 ft.	T.	4-5
ii. <i>S. pentandra</i> , L. . .	<i>Bay-leaved W.</i> .	Riv. banks	..	8-20 ft.	T.	5-6
iii. <i>S. fralilis</i> , L. . . .	<i>crack W., or Withy.</i>	Marshes.	..	80 ft.	T.	4-5
iv. <i>S. alba</i> , L.	<i>white W.</i>	Marshes.	..	80 ft.	T.	4-5
v. <i>S. Caprea</i> , L.	<i>S., or Goat-Willow</i>	Woods .	..	30 ft.	T.	4-5
vi. <i>S. aurita</i> , L.	<i>round-eared S.</i> .	Moi. thick.	..	2-4 ft.	Shb.	4-5

ORD. LXXVI.—CUPULIFERÆ—*cont.*

CHARACTERISTICS OF SPECIES AND VARIETIES.

Gen. IV.

- i. Ls. orb., doubly serr.

ORD. LXXVII.—BETULACEÆ.

Gen. I.

- i. Ls. orb.-cun., green on both sides.

Gen. II.

- i. Tree—Ls. rhomb.-triang., doubly serr.—Fr. broadly winged.

a. Lat. lobes of fruiting scales decurved.

b. Ls. cord. ov.—Lat. lobes of fruiting scales ascending

v. 2. *Ls. and twigs downy.*

- ii. A bush—Ls. orb.-cren. shortly stalked—Fr. narrowly winged.

ORD. LXXVIII.—SALICACEÆ.

Gen. I.

- i. Ls. roundish-cord., cottony beneath; those of the young shoots five-lobed—Stig. (yellow) bipart., with lin. seg.
- ii. Ls. roundish, toothed, waved, cottony beneath; those of the young shoots ent.—Stig. (purple) 3-4 lobed.
- iii. Ls. nearly round, acu., glab. beneath—Catk. dense—Sty. bifid.
- iv. Ls. triang., acum., glab. beneath—Catk. lax—Sty. two-lobed at apex.]

Gen. II.

- i. Ls. obl.-lanc., glab.—Catk. obl.-ov., stalked, appearing with the ls.; scales persist., glab.—Stam. three—Stig. sess.

Vars. Hoffmanniana, Sm.—amygdalina, Sm.—contorta, Crowe.

- ii. Ls. ellip.-lanc., gland., shining, pale beneath—Catk. ov.-atten., stalked; with the ls., fragrant; scales obl., decid.—Stam: five—Stig. bifid.

- iii. Ls. lan., pointed, serr., glab.—Caps. ob.-ov. lanc., stalked, glab.; with the ls.; scales decid.—Stam. two—Stig. bifid.

Vars. decipiens, Hoffm.—Russelliana, Sm.

- iv. Ls. lanc., acum., silky above and below—Catk. slend., lax, erect, sess.; with the ls.; scales lin., decid.—Sty. short.

Vars. cærulea, Sm.—vitellina, Sm.

- v. Ls. ov.-ellip., acu., deep green, cottony beneath—Catk. thick, blunt, sess.; before the ls.; scales persist.—Sty. short.

Vars. sphacelata, Sm.—cinerea, L. Sp.—aquatica, Sm.—oleifolia, Sm.

- vi. Ls. ob.-ov., cren., wrinkled, pubesc. beneath—Catk. lanc., short, sess.; before the ls.; scales persis.—Sty. short.

ORD. LXXVIII.—SALICACEÆ—*cont.*

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
vii. <i>S. repens</i> , <i>L.</i> . . .	<i>dwarf silky W.</i> .	Heaths .	. .	1-3 ft.	Shb.	4-5
viii. <i>S. nigricans</i> , <i>Sm.</i> . .	<i>dark-leaved S.</i> . .	Riv. banks	. .	6-10 ft.	Shb.	4-5
ix. <i>S. phylicifolia</i> , <i>L.</i> . .	<i>Tea-leaved W.</i> .	Mount.str.	. .	6-10 ft.	Shb.	4-5
x. <i>S. viminalis</i> , <i>L.</i> . . .	<i>Osier</i>	Wet pl. .	. .	30 ft.	Shb.	4-6
xi. <i>S. lanata</i> , <i>L.</i>	<i>woolly broad- leaved W.</i>	Sc. mount.	. .	1-2 ft.	Shb.	5-6
xii. <i>S. Lapponum</i> , <i>L.</i> . .	<i>downy mount W.</i>	Sc. mount.	. .	2-3 ft.	Shb.	6-7
xiii. <i>S. Myrsinites</i> , <i>L.</i> . .	<i>Whortle-leaved W.</i>	Sc. mount.	. .	1-2 ft.	Shb.	6-7
xiv. <i>S. arbuscula</i> , <i>L.</i> . . .	<i>small Tree W.</i> .	Sc. mount.	. .	1-2 ft.	Shb.	6-7
xv. <i>S. herbacea</i> , <i>L.</i>	<i>least W.</i>	Mount. .	. .	2-6 in.	Shb.	6-7
xvi. <i>S. reticulata</i> , <i>L.</i> . . .	<i>reticulate W.</i> . .	Sc. mount.	. .	1-2 ft.	Shb.	6-7
xvii. <i>S. purpurea</i> , <i>L.</i> . . .	<i>purple O.</i>	Riv. banks	. .	5-10 ft.	Shb.	4-5

CHARACTERISTICS OF SPECIES AND VARIETIES.

- vii.** St. procumb. or ascend.—Ls. small, obl.-lanc., margins recurved, shining above, silky beneath—Catk. cylind.-obl., erect, short, sess.; before the ls.; scales spath., silky—Sty. short.
Vars. *fusca*, L.—*prostrata*, Sm.—*ascendens*, Sm.—*parviflora*, Sm.—*argentea*, Sm.—*rosmarinifolia*, L.—*incubacea*, Sy.—*S. herbacea*, Ehrh., is probably a hybrid between *S. repens* and *S. aurita*, and *S. spathulata*, Willd., is a hairy form of *S. herbacea*.
- viii.** Ls. ov.-ellip., thin, glauc. beneath, turning black in drying—Catk. ov.-cylind., erect, sess.; scales lin.-obl., pilose—Sty. long.
Vars. *cotinifolia*, Sm.—*Forsteriana*, Sm.—*rupestris*, Don.—*Andersoniana*, Sm.—*damascena*, Forbes—*petraea*, And.—*hirta*, Sm.
- ix.** Ls. ellip.-lanc., glab., shining, glauc. beneath—Catk. sess.; scales lin.-obl., acu.—Sty. distinct.
Vars. *Davalliana*, Sm.—*Weigliana*, Willd. (*Wulfeniana*, Sm.)—*nitens*, G. Anders.—*Croweana*, Sm.—*Dicksoniana*, Sm.—*tenuior*, Borr.—*laxiflora*, G. Anders.—*propinqua*, Borr.—*tetrapla*, Walk.—*Borreriana*, Sm.—*phillyreifolia*, Borr.—*tenuifolia*, Borr.—*radicans*, Sm.—*S. laurina*, Sm. (*bicolor*, Sm.); *laxiflora*, Borr., is probably a hybrid between *S. phyllicifolia* and *S. caprea*.
- x.** Ls. lin.-lanc., silky beneath—Catk. lanc., long, nearly sess.; before the ls.; scales obt., brown—Sty. lin., ent.
Vars. *Intricata*, Leefe—*stipularis*, Leefe.
S. stipularis, Sm., is probably a hybrid between *S. viminalis* and *S. caprea*.
S. Smithiana, Willd. (*S. mollissima*, Sm.) also probably a hybrid between the same, of which *acuminata*, Sm.—*ferruginea* (G. Anders.) and *rugosa* (Sm.) are varieties.
- xi.** Ls. broadly ov., coriaceous, ent., shaggy—Catk. term., sess., stout, dense; with the ls.; scales obt., with long yellow hairs—Sty. long.
- xii.** Ls. ellip.-lanc., silky above, cottony beneath—Catk. sess.; with the ls.; scales acu., with long white hairs—Sty. very long.
Vars. *arenaria*, Sm.—*Stuartiana*, Sm.—*glaucia*, Sm.
- xiii.** Ls. small, rigid, lanc., with prominent veins—Catk. term. on leafy pedunc.; with the ls.; scales spath.—Sty. long.
Vars. *procumbens*, Forbes (*levis*, Hook)—*arbutifolia*, Sy.—*serrata*, Sy.
- xiv.** Ls. ov.-lanc., glab., veiny above, finely serr., glauc. beneath—Catk. lat., sess., on leafy pedunc.; before the ls.; scales obt., hairy—Sty. long.
Vars. *carinata*, Sm.—*prunifolia*, Sm.—*verulosa*, Sm.—*vaccinifolia*, Walk. (*livida*, Sm.)—*angustifolia*, Wulf.
- xv.** Very small; st. underground—Ls. obl. or orb., obt., veined above—Catk., term. on two-leaved pedunc.; with the ls.; scales glab.
- xvi.** Small; st. underground—Ls. orb.-obl., strongly veined, glab. above, glauc. beneath—Catk., on long leafy pedunc.; scales ob-ov.
- xvii.** Ls. lin.-lanc., thin, gland., serr.—Catk. ov., sess.; before the ls.; very downy; scales small, purple—Stig. ov.
Vars. *Woollgariana*, Borr.—*ramulosa*, Borr.—*Lambertiana*, Sm.
S. rubra is a hybrid between *S. purpurea* and *S. viminalis*, of which *S. Forbyana*, Sm., and *S. Helix*, L., are varieties.
S. Doniana is a hybrid between *S. purpurea* and *S. repens*.
S. cuspidata, Schultz, is a hybrid between *S. pentandra* and *S. fragilis*.
S. acutifolia, Willd., (*S. Daphnoides*, Vill.) is not indigenous.
S. undulata, Ehrh., (*S. lanceolata*, Sm.) is a hybrid between *S. triandra* and *S. viminalis*.

ORD. LXXIX.—MYRICACEÆ.

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
Gen. I. Myrica, L.	<i>Sweet-gale, Bog-myrtle.</i>					
i. <i>M. Gale, L.</i>	<i>S. or B.</i>	Mount. & bogs.	..	2-3 ft.	Shb.	6-7

ORD. LXXX.—CONIFERÆ.

Gen. I. Pinus, L.	<i>Pine, Fir.</i>					
i. <i>P. sylvestris, L.</i>	<i>Scotch Fir</i>	Scot. . .	..	100 ft.	T.	5-6
Gen. II. Juniperus, L.	<i>Juniper.</i>					
i. <i>J. communis, L.</i>	<i>common J.</i>	Downs .	..	1-4 ft.	Shb.	5-6
v. 2. <i>nana, Willd.</i>	<i>dwarf J.</i>	Mount. .	..	..	..	..
Gen. III. Taxus, L.	<i>Yew.</i>					
i. <i>T. baccata, L.</i>	<i>common Y.</i>	Mnt. wds.	..	50 ft.	T.	3-4
v. 2. <i>fastigiata, Lindl.</i> .	. . . . .	..	..	..	..	..

ORD. LXXXI.—ORCHIDACEÆ.

Gen. I. Goodyera, Br.	<i>Goodyera.</i>					
i. <i>G. repens, Br.</i>	<i>creeping G.</i>	Sco. fir woods.	crW.	4-8 in.	P.	8
Gen. II. Spiranthes, Rich.	<i>Lady's-tresses.</i>					
i. <i>S. autumnalis, Rich.</i> .	<i>Autumnal L.</i>	Downs .	W.	4-8 in.	P.	8-9
ii. <i>S. æstivalis, Rich.</i> . .	<i>Summer L.</i>	Bogs ! S. of E.	W.	6-18 in.	P.	7-8
iii. <i>S. Gemmipara, Lindl.</i>	<i>drooping L.</i>	Mead. Ire.!	W.	6-10 in.	P.	8-9
Gen. III. Neottia, L.	<i>Bird's-nest.</i>					
i. <i>N. Nidus-avis, Rich.</i> .	<i>common B.</i>	Sh. woods	Pk.	4-10 in.	P.	6-7
Gen. IV. Listera, Br.	<i>Sway-blade.</i>					
i. <i>L. cordata, Br.</i>	<i>Heart-leaved T.</i>	Turfy moors !	gr Br.	4-8 in.	P.	6-8
ii. <i>L. ovata, Br.</i>	<i>common T.</i>	Moi. pas.	y Gr.	1-2 ft.	P.	5-7
Gen. V. Epipactis, Hall.	<i>Helleborine.</i>					
i. <i>E. latifolia, Sw.</i>	<i>broad-leaved H.</i>	Ch. woods	gPu.	1-3 ft.	P.	7-8
b. <i>media, Fries.</i>	. . . . .	..	..	..	..	..

ORD. LXXIX.—MYRICACEÆ.

CHARACTERISTICS OF SPECIES AND VARIETIES.

Gen. I.

- i. Ls. lanc., serr., broader upwards, fragrant catk., sess., erect.

ORD. LXXX.—CONIFERÆ.

Gen. I.

- i. Ls. rigid, in pairs—Cones ov.-acum., stalked.

Gen. II.

- i. Ls. lin., mucro., three in a whorl, longer than the fr.
v. 2. *St. procumb.*—Ls. broader and shorter—Fr. longer.

Gen. III.

- i. Ls. lin., acum., crowded—Fls. axill., sess.—Fr. roundish.
v. 2. *Branches fastigate*—Ls. scattered—Fr. obl.

ORD. LXXXI.—ORCHIDACEÆ.

Gen. I.

- i. Root creeping—Ls. stalked, ov., netted—Sep. and lip ov.-lanc.

Gen. II.

- i. Rad. ls. in a lat. rosette, ov., ac.—Spike dense; fls. in one rank—Tubers ovoid.
ii. Rad. ls. at base of st., obl.-lanc.—Spike lax; fls. in one rank—Tubers cylind.

- iii. Rad. ls. lin.-lanc.—Spike dense; fls. in three ranks.

Gen. III.

- i. St. leafless, with brown scales—Lip lin., obt., with two spread. lobes.

Gen. IV.

- i. St. short, glab.—Ls. cord.—Lip bifid. with two lat. lobes.
ii. St. tall, hairy above—Ls. ov.—Lip bifid.

Gen. V.

- i. Ls. ovate—Bracts mostly longer than fls.—Lower lobe of lip acute, shorter than ovate sep.
b. Ls. longer and narrower—Lower lobe of lip triang.-cord., as long as the lanc. sep.

ORD. LXXXI.—ORCHIDACEÆ—cont.

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
<i>c. purpurata</i> , Sm. . . .	. . . .	. . . .	. .	. .	. .	. .
<i>d. ovalis</i> , Bab.	. . . .	. . . .	. .	. .	. .	. .
ii. <i>E. palustris</i> , Sw. . .	<i>Marsh H.</i> . . .	Marshes.	W.	6-18 in.	P.	7-8
Gen. VI. <i>Cephalanthera</i> , <i>Rich.</i>	<i>Helleborine.</i>					
i. <i>C. grandiflora</i> , Bab. .	<i>large white H.</i> .	Ch. woods	C.	6-18 in.	P.	5-6
ii. <i>C. ensifolia</i> , Rich. .	<i>narrow-leaved H.</i>	Mount. woods!	W.	6-18 in.	P.	5-6
iii. <i>C. rubra</i> , Rich. . . .	<i>purple H.</i> . . .	Mount. woods!	puR.	6-12 in.	P.	6-7
Gen. VII. <i>Epipogium</i> , <i>Gm.</i>						
i. <i>E. Gmelini</i> , Rich. . .	. . . .	. . . .	p Y.	4-8 in.	P.	8
Gen. VIII. <i>Corallorhiza</i> , <i>Hall.</i>	<i>Coral-root.</i>					
i. <i>C. innata</i> , Br.	<i>spurless C.</i> . . .	Bog. wds! Sco.	y G.	6-12 in.	P.	7
Gen. IX. <i>Orchis</i> , L. SEC. I. *	<i>Orchis</i>					
A.						
i. <i>O. Morio</i> , L.	<i>green-winged Meadow O.</i>	Past. . .	gPu.	6-12 in.	P.	5-6
ii. <i>O. mascula</i> , L. . . .	<i>early purple O.</i> .	Past. . .	Pu.	6-12 in.	P.	4-6
iii. <i>O. ustulata</i> , L. . . .	<i>dwarf O.</i>	Ch. past.	puW.	4-5 in.	P.	5-6
iv. <i>O. fusca</i> , Jacq. . . .	<i>great brown- winged O.</i>	Ch. past.! Kent.	gPu.	1-2 ft.	P.	5-6
v. <i>O. militaris</i> , L. . . .	<i>military O.</i> . . .	Ch. hills!	Pu.	12-18 in.	P.	5-6
v. 2. <i>Simia</i> , Lam. . . .	<i>Monkey O.</i> . . .	Ch. hills!	. .	. .	. .	. .
B.						
vi. <i>O. laxiflora</i> , Lam. . .	<i>lax-flowered O.</i> .	Jersey & Guernsey.	r Pu.	1-2 ft.	P.	5-6
**						
vii. <i>O. latifolia</i> , L. . . .	<i>Marsh O.</i>	Moi.mead.	roPu.	1-3 ft.	P.	6-7
v. 2. <i>incarnata</i> , L. . . .	. . . .	. . . .	. .	. .	. .	. .
viii. <i>O. maculata</i> , L. . .	<i>spotted palmate O.</i>	Heaths .	pPu.	6-18 in.	P.	6-8
SEC. II.						
ix. <i>O. pyramidalis</i> , L. . .	<i>pyramidal O.</i> . .	Ch. past.	Ro.	6-18 in.	P.	7-8
x. <i>O. hircina</i> , Scop. . . .	<i>Lizard O.</i>	Ch. hills!	puW.	1-2 ft.	P.	7-8
Gen. X. <i>Gymnadenia</i> , Br.						
i. <i>G. conopsea</i> , Br. . . .	<i>fragrant G.</i> . . .	Ch. heaths	Ro.	12-18 in.	P.	6-8

CHARACTERISTICS OF SPECIES AND VARIETIES.

- c.* As Var. *b.*, but st. and ls. tinged with purple, and lower lobe of lip shorter than sep.
d. Ls. ov.-lanc.—Fls. small, dingy purple—Lower lobe of lip broader than long, as long as the ov., acute sep.
ii. Ls. lanc.—Bracts shorter than fls.—Lower lobe of lip obt.

Gen. VI.

- i.* Fls. creamy white—Ls. ov.—Ovary glab., shorter than bracts.
ii. Fls. pure white—Ls. lanc.—Ovary glab., longer than the minute bracts.
iii. Fls. rosy purple—Ls. lanc.—Ovary downy.

Gen. VII.

- i.* Leafless—St. swollen at base—Fls. pale yellow, streaked with white.
(Once found at Tedstone, Delamere Forest, Hereford.)

Gen. VIII.

- i.* Root of thick fleshy fibres—St. with 2-3 lanc., sheathing leaves—Fls. in a lax spike.

Gen. IX.

SEC. I.—*Pollen-glands divided.*

* *Tubers of root two, undivided.*

A. Bracts one-veined.

- i.* Spike lax—Lat. sep. obt., connivent—Spur shorter than germen.
ii. Ls. spotted—Spike lax—Lat. sep. acute, reflexed upwards—Spur obt., as long as germen.
iii. Spike dense—Seg. of lip deep, lin. ; lower one without a point in notch.
iv. Spike dense—Lower seg. of lip with a point in notch ; lat. sep. obt., connivent, dent., greenish purple.
v. Spike dense—Lower seg. of lip with a point in notch ; lat. seg. acute, connivent, pale purple.
v. 2. *Lat. sep. and lip white—Lobes of lip long, narrow, crimson.*
B. Bracts three-veined.
vi. Ls. lanc.—Spike lax—Lat. sep., reflexed.
 * * *Tubers of root palmate.*
vii. Lip indistinctly three-lobed ; its sides reflexed—St. hollow—Bracts longer than ovary.
v. 2. *Ls. concave at tip, unspotted.*
viii. Lip deeply three-lobed, flat—Ls. spotted—St. solid—Bracts shorter than ovary.
 SEC. II.—*Pollen-glands united.*
ix. Spike dense—Sep. spread.—Lip with three equal lobes.
x. Spike lax—Sep. connivent—Middle lobe of lip long, lin., twisted.

Gen. X.

- i.* Spike dense ; fls. very fragrant—Lat. sep. spread.—Spur long, slender, flexuous.

ORD. LXXXI.—ORCHIDACEÆ—*cont.*

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
Gen. XI. Habenaria, Br.	<i>Butterfly-Orchis.</i>					
i. <i>H. bifolia, Br.</i>	<i>lesser B.</i>	Moi. mead.	W.	6-18 in.	P.	6-8
v. 2. <i>chlorantha, Bab.</i>	<i>great B.</i>	• • • •	• W.	• 6-8 in.	• P.	• 6-8
ii. <i>H. viridis, Br.</i>	<i>Frog-O., or green H.</i>	Dry past.	g W.	6-8 in.	P.	6-8
iii. <i>H. albida, Br.</i>	<i>small white H.</i> . .	Mnt. past.	W.	6-12 in.	P.	6-8
Gen. XII. Aceras, Br.	<i>Man Orchis.</i>					
i. <i>A. anthropophora, Br.</i>	<i>green M.</i>	Ch. past.	g Y.	6-12 in.	P.	6-7
Gen. XIII. Neotinea, Reich., fil.						
i. <i>N. intacta</i>	. . . . .	Ch. past. ! Galway.	Pk.	4-10 in.	P.	6-7
Gen. XIV. Herminium, Br.	<i>Musk Orchis.</i>					
i. <i>H. monorchis, Br.</i> . . .	<i>green M.</i>	Ch. past.	Gr.	4-6 in.	P.	6-7
Gen. XV. Ophrys, L.	<i>Insect Orchis.</i>					
i. <i>O. apifera, Huds.</i> . . .	<i>Bee O.</i>	Ch. past.	Pu.	6-12 in.	P.	6-7
v. 2. <i>arachnites, Willd.</i> . .	<i>late Spider O.</i> . . .	Ch. downs !	• •	• • •	• •	• •
ii. <i>O. aranifera, Huds.</i> . .	<i>Spider O.</i>	Ch. past. !	pu Br.	6-12 in.	P.	4-5
v. 2. <i>fucifera</i>						
iii. <i>O. muscifera, Huds.</i> . .	<i>Fly O.</i>	Ch. past.	b Pu.	6-12 in.	P.	5-7
Gen. XVI. Malaxis, Sw.	<i>Bog Orchis.</i>					
i. <i>M. paludosa, Sw.</i> . . .	<i>Bog O.</i>	Bogs ! Scot.	Gr.	2-4 in.	P.	7-9
Gen. XVII. Liparis, Rich.						
i. <i>L. Loeselii, Rich.</i> . . .	<i>two-leaved L.</i> . . .	San. bogs E. of E. !	y G.	6-8 in.	P.	7-8
Gen. XVIII. Cypripedium, L.	<i>Lady's Slipper.</i>					
i. <i>C. calceolus, L.</i>	<i>common L.</i>	Woods ! N. of E.	y Br.	12-18 in.	P.	5-6

ORD. LXXXII.—IRIDACEÆ.

Gen. I. Iris, L.	<i>Iris, Flower-de-luce.</i>					
i. <i>I. fœtidissima, L.</i> . . .	<i>fœtid I., or Glad-don.</i>	Thick. ! .	Pu.	2-3 ft.	P.	5-7
ii. <i>I. Pseudacorus, L.</i> . . .	<i>yellow I., or Flag.</i>	Wet pl. .	Y.	2-3 ft.	P.	5-8

ORD. LXXXI.—ORCHIDACEÆ—*cont.*

CHARACTERISTICS OF SPECIES AND VARIETIES.

Gen. XI.

- i. Ls. two—Sep. spread.—Spur long—Lip entire—Anther cells parallel.
- v. 2. *Larger—Fls. greener—Anther cells diverging at base.*
- ii. Ls. several—Fls. green—Sep. connivent—Spur short—Lip lin., bifid, greenish brown.
- iii. Ls. several—Fls. white—Sep. connivent—Lip acutely three-lobed.

Gen. XII.

- i. Spike lax—Lat. sep. ovate, connivent, green—Lip three-part ; lobes lin., lower one bifid.

Gen. XIII.

- i. Spike dense—Ls. ellip.-obl., spotted—Lat. sep. and pet. connivent—Lip three-lobed.

Gen. XIV.

- i. Spike lax, slender—Fls. small—Sep. broad ; Pet. narrower—Lip three-lobed.

Gen. XV.

- i. Pet. downy, lin.—Sep. pink inside—Lip tumid., four-lobed, not longer than sep. with an intermediate, acute, reflexed appendage in notch.
- v. 2. *Pet. deltoid—Lip longer than sep.—Appendage of notch cord., flat.—(Kent.)*
- ii. Pet. glab., lin.—Sep. yellowish green inside—Lip. tumid., four-lobed, without an intermediate appendage in notch.
- v. 2. *Pet. rough inside—Lip undivided, convex.*
- iii. Pet. filiform—Sep. yellowish green—Lip oblong, three-lobed ; middle lobe bifid.

Gen. XVI.

- i. Spike dense—Ls. 3-5, ob.-ov., concave, papillose at tip—Lip ent., acute, concave.

Gen. XVII.

- i. Spike lax—St. triang.—Ls. two, obl.-lanc., keeled—Lip ent., ob.-ov.

Gen. XVIII.

- i. St. leafy—Ls. obl., ribbed—Fls. 1-2, large—Sep. and pet. brown ; lip yellow, inflated.

ORD. LXXXII.—IRIDACEÆ.

Gen. I.

- i. St. angled—Fls. livid, purple.
- ii. St. round—Fls. yell.

ORD. LXXXII.—IRIDACEÆ—*cont.*

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
Gen. II. Gladiolus, L.	<i>Gladiolus, or Corn- flag.</i>					
i. <i>G. communis, L.</i> . . .	<i>common G., or C.</i>	New Forest!	Cr.	2-3 ft.	P.	6-7
Gen. III. Sisyrinchium, L.						
[i. <i>S. anceps, Lam.</i> . . .	. . .	Galway!	B.	6-18 in.	P.	7-8
Gen. IV. Trichonema, Ker.						
i. <i>T. Columnæ, Reich.</i>	. . . .	Channel Islands!	b Pu.	2-4 in.	P.	3-4
Gen. V. Crocus, L. . . .	<i>Crocus.</i>					
[i. <i>C. vernus, Willd.</i> . . .	<i>purple Spring C.</i>	Mead.!	Vi.	2-6 in.	P.	3-4
ii. <i>C. nudiflorus, Sm.</i> . . .	<i>naked-flowered C.</i>	Mead.!	Pu.	2-6 in.	P.	9-10

ORD. LXXXIII.—AMARYLLIDACEÆ.

Gen. I. Narcissus, L.	<i>Narcissus, Daffo- dil.</i>					
[i. <i>N. biflorus, Curt.</i> . . .	<i>pale N.</i>	San. fie.!	p Y.	12-18 in.	P.	4-5
ii. <i>N. pseudo-narcissus, L.</i>	<i>common D.</i>	Moi. wds.	Y.	12-18 in.	P.	3-4
Gen. II. Galanthus, L.	<i>Snow-drop.</i>					
i. <i>G. nivalis, L.</i>	<i>common S.</i>	Woods .	W.	6-10 in.	P.	2-3
Gen. III. Leucojum, L.	<i>Snow-flake.</i>					
[i. <i>L. vernal, L.</i>	<i>Spring S.</i>	Copses! .	g W.	12-18 in.	P.	4-5
ii. <i>L. æstivum, L.</i>	<i>Summer S.</i>	Wet mead.!	g W.	1-2 ft.	P.	5-7

ORD. LXXXIV.—LILIACEÆ.

Gen. I. Lilium, L.	<i>Martagon Lily.</i>					
[i. <i>L. martagon, L.</i>	<i>purple M. L.</i> . . .	Mickle- ham!	Pu.	2-3 ft.	P.	7-8
Gen. II. Tulipa, L.	<i>Tulip.</i>					
[i. <i>T. sylvestris, L.</i>	<i>wild Tulip</i>	Ch. pits!	Y.	1-2 ft.	P.	4-5

ORD. LXXXII.—IRIDACEÆ—*cont.*

CHARACTERISTICS OF SPECIES AND VARIETIES.

Gen. II.

- i. Ls. glauc., ensi.—Fls. unilat, crim.-purp.

Gen. III.

- i. Ls. rad., ensi.—Scape two-edged—Fls. umbell.]

Gen. IV.

- i. Ls. rad., fili., furrowed—Scape one-flowered, drooping.

Gen. V.

- i. Fls. vernal—Stig. three fid. ; lobes cuneate.]
 ii. Fls. autumn—Stig. deeply laciniate.
(Several other species of Crocus are naturalized in various parts of England, as C. minimus, C. aureus, C. sativus, &c. ; but C. nudiflorus is the only one which has any claim to be truly indigenous.)

ORD. LXXXIII.—AMARYLLIDACEÆ.

Gen. I.

- i. Ls. keeled—St. 1-3 flowered—Fls. salver-shaped.]
 ii. Ls. flat—St. one-flowered—Fls. campan.
(Several other species have been naturalized in various parts of England, but only N. pseudo-narcissus has any claim to be indigenous.)

Gen. II.

- i. Ls. glauc., keeled—Fls. sol., drooping.

Gen. III.

- i. Fls. in spring—St. 1-2 flowered.]
 ii. Fls. in summer—St. many-flowered.

ORD. LXXXIV.—LILIACEÆ.

Gen. I.

- i. Ls. whorled, ellip.-lanc.—Fls. drooping ; perianth reflexed.]

Gen. II.

- i. Ls. lin.-lanc.—Fls. sub-erect—Seg. of perianth ellip.-lanc., bearded.]

ORD. LXXXIV.—LILIACEÆ—cont.

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
Gen. III. Fritillaria, L.	<i>Fritillary.</i>					
i. <i>F. meleagris, L.</i> . . .	<i>common F., or Snake's-head.</i>	Past.! . .	brPu.	6-12 in.	P.	5-6
Gen. IV. Allium, L.	<i>Onion, Leek, Garlic, &c.</i>					
* A.						
[i. <i>A. ampeloprasum, L.</i>	<i>wild Leek</i>	Rocky pl.	gW.	3-4 ft.	P.	7-8
v. 2. <i>Babingtonii, Borr.</i>						
ii. <i>A. Scorodoprasum, L.</i>	<i>Sand G.</i>	Sand. wds. N. of E.	rPu.	2-3 ft.	P.	5-7
B.						
iii. <i>A. oleraceum, L.</i> . . .	<i>Field G.</i>	Waysides!	brW.	10-18 in.	P.	7-8
v. 2. <i>carinatum, Sm.</i> . . .	. . . .	. . . .	. . .	. . .	. . .	. . .
C.						
iv. <i>A. vineale, L.</i>	<i>Crow Garlic.</i> . . .	Was. pl.	Pk.	1½-2 ft.	P.	6-7
v. 2. <i>compactum, Thuill.</i>	. . . .					
v. <i>A. sphaerocephalum,</i>	<i>small round-</i>	Bristol,	rPu.	1½-2 ft.	P.	6-7
L.	<i>headed G.</i>	Jersey!				
vi. <i>A. Schoenoprasum, L.</i>	<i>Chives</i>	Rock.pas.!	pPu.	6-12 in.	P.	6-7
v. 2. <i>sibiricum, L.</i>	. . . .	. . . .	. . .	. . .	. . .	. . .
* *						
vii. <i>A. ursinum, L.</i>	<i>broad-leaved G., or Ransoms.</i>	Woods .	W.	6-12 in.	P.	4-6
viii. <i>A. triquetrum, L.</i> . .	<i>triangular stalked G.</i>	Guernsey!	W.	10-18 in.	P.	4-6
Gen. V. Gagea, Salisb.	<i>Gagea.</i>					
i. <i>G. lutea, Ker.</i>	<i>yellow G.</i>	Woods! .	Y.	6-8 in.	P.	3-4
Gen. VI. Ornithogallum,	<i>Star of Bethlehem.</i>					
L.						
i. <i>O. pyrenaicum, L.</i> . . .	<i>spiked S.</i>	Copses! .	gW.	1½-2 ft.	P.	6-7
[ii. <i>O. nutans, L.</i>	<i>drooping S.</i>	Copses! .	W.	8-12 in.	P.	4-5
[iii. <i>O. umbellatum, L.</i> . . .	<i>common S.</i>	Copses .	W.	8-12 in.	P.	4-5
Gen. VII Scilla, L.	<i>Squill.</i>					
i. <i>S. verna, Huds.</i>	<i>vernal S.</i>	Rocky past.!	B.	4-6 in.	P.	4-5
ii. <i>S. autumnalis, L.</i> . . .	<i>autumnal S.</i> . . .	Dry past.	puB.	4-6 in.	P.	8-9
Gen. VIII. Hyacinthus,	<i>Blue-bell, Wild Hyacinth.</i>					
L.						
i. <i>H. nonscriptus, L.</i> . . .	<i>Blue-bell, or W.</i> . .	Woods .	B.	6-18 in.	P.	4-6
Gen. IX. Muscari, Tourn.	<i>Grape-Hyacinth.</i>					
i. <i>M. racemosum, Mill.</i>	<i>Starch G.</i>	Sand. fie.! E. of E.	B.	6-12 in.	P.	5-6
Gen. X. Lloydia, Salisb.						
i. <i>L. serotina, Reich.</i> . . .	<i>Mountain L.</i> . . .	Welsh mount.!	W.	2-6 in.	P.	6-7

CHARACTERISTICS OF SPECIES AND VARIETIES.

Gen. III.

- i. Ls. lin.-lanc., ent.—Fls. sol., drooping—Points of perianth inflexed.

Gen. IV.

* *Stem leafy.*

A. Ls. flat.

- i. Ls. lin., keeled; sheaths cylind.—Umbels compact; fls. many; bulbs o.; sta exserted.

v. 2. *Umbels irregular, with few flowers and many bulbs.]*

- ii. Ls. lin., flat; sheaths two-edged—Sta. included.

B. Ls. fleshy.

- iii. Ls. grooved above, ribbed beneath—Umbels lax, bulbiferous.

v. 2. *Ls. thick, nearly flat.—(N. of E.!)*

C. Ls. fistulous.

- iv. Umbels bulbiferous.

v. 2. *Umbels compact, bearing bulbs only.*

- v. Umbels without bulbs—Stam. exserted; the alternate filaments three-cleft.

- vi. Umbels without bulbs—Stam. included, all simp.

v. 2. *Larger—Ls. spreading, arching.—(Cornwall!)*

* * *Ls. all radical.*

- vii. Ls. ov.-lanc.—Stem bluntly triang.

- viii. Ls. lin., keeled—Stem acutely triang

Gen. V.

- i. St. ang., shorter than the lin.-lanc. ls.—Fls. umbell.

Gen. VI.

- i. Ls. vernal—Fls. small, erect, in a long raceme.

ii. Fls. drooping in a pend. raceme—Pedunc. shorter than the bracts.]

- iii. Fls. erect, corymbose—Pedunc. longer than the bracts.]

Gen. VII.

- i. Ls. channelled—Rac. few-flowered, corymb.—Bracts lanc.

- ii. Ls. lin.—Rac. lax—Bracts o.

Gen. VIII.

- i. Ls. long, acum., channelled—Fls. drooping.

Gen. IX.

- i. Ls. lin., flaccid, keeled, longer than st.—Fls. crowded, nodding.

Gen. X.

- i. Ls. semi-cylind.; st. ls. sheathing—Fls. sol.

ORD. LXXXIV.—LILIACEÆ—*cont.*

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
Gen. XI. Simethis, Kunth.						
i. <i>S. bicolor, Kunth.</i> . . .	<i>variegated S.</i> . . .	Fir woods! Bourne- mouth.	Pu.	6-18 in.	P.	5-6
Gen. XII. Asparagus, L.	<i>Asparagus.</i>					
i. <i>A. officinalis, L.</i> . . .	<i>common A.</i> . . .	Sea coast!	y W.	1-2 ft.	P.	6-8
Gen. XIII. Ruscus, L.	<i>Butcher's Broom.</i>					
i. <i>R. aculeatus, L.</i> . . .	<i>common B.</i> . . .	Heaths!.	W.	1-2 ft.	Shb.	3-4
Gen. XIV. Maianthe- mum, Mæench.	<i>May Lily.</i>					
i. <i>M. bifolium, DC.</i> . . .	<i>two-leaved M.</i> . . .	Woods!.	. .	4-8 in.	P.	5-6
Gen. XV. Convallaria, L.	<i>Lily of the Valley.</i>					
i. <i>C. majalis, L.</i>	<i>common L.</i>	Woods!.	W.	6-8 in.	P.	5-6
Gen. XVI. Polygonatum, Tourn.	<i>Solomon's Seal.</i>					
i. <i>P. verticillatum, All.</i> .	<i>narrow-leaved S.</i>	Woods!.	G.	2-3 ft.	P.	6-7
ii. <i>P. multiflorum, All.</i> .	<i>common S.</i>	Woods!.	g W.	2-3 ft.	P.	6-7
iii. <i>P. officinale, All.</i> . .	<i>angular S.</i>	Woods!.	g W.	6-12 in.	P.	5-6

ORD. LXXXV.—TRILLIACEÆ.

Gen. I. Paris, L.	<i>Herb-Paris.</i>					
i. <i>P. quadrifolia, L.</i> . . .	<i>common H.</i>	Woods .	g W.	6-12 in.	P.	5-6

ORD. LXXXVI.—DIOSCOREACEÆ.

Gen. I. Tamus, L.	<i>Black Bryony.</i>					
i. <i>T. communis, L.</i> . . .	<i>common B.</i>	Hedges .	y G.	Cl.	P.	5-6

ORD. LXXXVII.—MELANTHIACEÆ.

Gen. I. Colchicum, L.	<i>Meadow Saffron.</i>					
i. <i>C. autumnale, L.</i> . . .	<i>common M.</i>	Mead.! .	p Pu.	2-6 in.	P.	8-10

ORD. LXXXIV.—LILIACEÆ—*cont.*

CHARACTERISTICS OF SPECIES AND VARIETIES.

Gen. XI.

- i. Ls. lin., recurved—Scape dichotomously branched—Fls. corymb.

Gen. XII.

- i. St. erect, much branched—Ls. setaceous, fasciculate.

Gen. XIII.

- i. St. branched, rigid—Ls. ov.-acum., rigid, evergreen, with fl. in centre.

Gen. XIV.

- i. Ls. two, triang.-cord., alt., stalked—Rac. spicate.

Gen. XV.

- i. Ls. two, rad., ov.-lanc.—Fls. nodding, fragrant.

Gen. XVI.

- i. St. ang.—Ls. lin.-lanc., whorled.
 ii. St. round—Ls. ellip.-lanc.—Fila. hairy.
 iii. St. ang.—Ls. ellip.-lanc.—Fila. glab.

ORD. LXXXV.—TRILLIACEÆ.

Gen. I.

- i. Ls. ov., four or five in a whorl, at summit of st.—Fls. sol., stalked.

ORD. LXXXVI.—DIOSCOREACEÆ.

Gen. I.

- i. St. long, twining—Ls. cord., acute.

ORD. LXXXVII.—MELANTHIACEÆ.

Gen. I.

- i. Ls. lanc., erect, vernal—Fls. autumnal.

ORD. LXXXVII.—MELANTHACEÆ—*cont.*

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
Gen. II. Tofieldia, Huds.	<i>Scottish-Asphodel.</i>					
i. <i>T. palustris, Huds.</i> . .	<i>Mount S.-A.</i> . .	Mount. bogs.	p G.	4-8 in.	P.	7-8

ORD. LXXXVIII.—HYDROCHARIDACEÆ.

Gen. I. Hydrocharis, L.	<i>Frog-bit.</i>					
i. <i>H. Morsus-ranæ, L.</i> . .	<i>common F.</i> . . .	Ponds .	W.	Fl.	P.	7-8
Gen. II. Elodea, Rich.	<i>Water-thyme.</i>					
[i. <i>E. Canadensis, Rich.</i> ..	<i>long-leaved W.</i> .	Streams, &c.	g Pu.	Fl.	P.	7-10
Gen. III. Stratiotes, L.	<i>Water-Soldier.</i>					
i. <i>S. aloides, L.</i>	<i>Water-S.</i>	Ponds .	W.	4-6 in.	P.	7-8

ORD. LXXXIX.—ALISMACEÆ.

Gen. I. Alisma, L.	<i>Water-Plantain.</i>					
i. <i>A. Plantago, L.</i>	<i>great W.</i>	Ditches, &c.	p Ro.	1-3 ft.	P.	6-8
v. 2. <i>lanceolata, With.</i> .						
ii. <i>A. ranunculoides, L.</i> .	<i>lesser W.</i>	Ditches .	p Ro.	6-18 in.	P.	5-9
v. 2. <i>repens, Dav.</i>						
iii. <i>A. natans, L.</i>	<i>floating W.</i> . . .	Lakes, N. of E.!	W.	Fl.	P.	7-8
Gen. II. Actinocarpus, Br.	<i>Star-fruit.</i>					
i. <i>A. Damosonium, Br.</i> .	<i>common S.</i>	Ditches ! S. of E.	W.	4-6 in.	P.	5-7
Gen. III. Sagittaria, L.	<i>Arrow-head.</i>					
i. <i>S. sagittifolia, L.</i> . .	<i>common A.</i>	Ditches .	W.	6-18 in.	P.	7-9
Gen. IV. Butomus, L.	<i>Flowering Rush.</i>					
i. <i>B. umbellatus, L.</i> . .	<i>common F.</i>	Ditches .	Ro.	3-4 ft.	P.	6-7
Gen. V. Triglochin, L.	<i>Arrow-grass.</i>					
i. <i>maritimum, L.</i>	<i>Sea-side A.</i>	Salt mar.	Gr.	10-12 in.	P.	5-9
ii. <i>palustre, L.</i>	<i>Marsh A.</i>	Mar. . .	Gr.	6-18 in.	P.	6-8

ORD. LXXXVII.—MELANTHACEÆ—*cont.*

CHARACTERISTICS OF SPECIES AND VARIETIES.

Gen. II.

- i. Ls. rad., lin.—St. slender—Fls. in an ovate spike.

ORD. LXXXVIII.—HYDROCHARIDACEÆ

Gen. I.

- i. Ls. reni., stalked, ent.—Fls. delicate white, umbellate.

Gen. II.

- i. Ls. lin., 3-4 in a whorl—St. brittle, rooting at joints.]

Gen. III.

- i. Ls. sword-shaped, spinous, serrate.

ORD. LXXXIX.—ALISMACEÆ.

Gen. I.

- i. Ls. all rad., ov.-lanc., sub-cord.—Scape panicled with compound branches—Fr. compressed, obt.
 v. 2. *Ls. lanc., narrowed at base.*
 ii. Ls. all rad., lin.-lanc.—Scape with simple branches—Fr. ribbed, acute.
 v. 2. *Fls. larger—Scape procumb., rooting.*
 iii. Rad. ls. lin., sess. ; flo. ls. ellip., obt.—Pedunc. simp., from the st. joints.

Gen. II.

- i. Ls. rad., ellip., five-ribbed, long-stalked—Scape with 2-3 whorls of fls.

Gen. III.

- i. Ls. arrow-shaped—Scape with 3-5 distant whorls.

Gen. IV.

- i. Ls. rad., lin., trigonous—Scape terete, longer than ls.—Fls. umbell., with a three-leaved invol.

Gen. V.

- i. Ls. rad., lin., fleshy—Fls. in a lax, simp. spike—Fr. ovate, six-celled.
 ii. Similar but smaller—Fr. lin., ang., three-celled.

ORD. LXXXIX.—ALISMACEÆ—*cont.*

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
Gen. VI. <i>Scheuchzeria</i> , <i>L.</i>						
i. <i>S. palustris</i> , <i>L.</i> . . .	<i>Marsh S.</i>	Mar.! . .	Gr.	6-8 in.	P.	7-8

ORD. XC.—NAIADACEÆ.

Gen. I. <i>Potamogeton</i> , <i>L.</i>	<i>Pond-weed.</i>					
SEC. I.						
A.						
i. <i>P. densus</i> , <i>L.</i>	<i>opposite-leaved P.</i>	Ditches .	. .	Fl.	P.	6-7
B.						
ii. <i>P. pectinatus</i> , <i>L.</i> . . .	<i>Fennel-leaved P.</i>	Streams .	. .	Fl.	P.	6-7
v. 2. <i>flabellatus</i> , <i>Bab.</i> . . .						
iii. <i>P. filiformis</i> , <i>Pers.</i> . .	<i>slender-leaved P.</i>	Lakes! .	. .	Fl.	P.	6-7
C.						
iv. <i>P. trichioides</i> , <i>Cham.</i> .	<i>hair-like P.</i>	Ponds, Bexley!	. .	Fl.	P.	7-8
v. <i>P. pusillus</i> , <i>L.</i>	<i>small P.</i>	Ditches .	. .	Fl.	P.	6-7
vi. <i>P. compressus</i> , <i>L.</i> . . .	. . . . .	Ditches .	. .	Fl.	P.	6-7
vii. <i>P. gramineus</i> , <i>Auct.</i> . .	<i>grassy P.</i>	Ditches .	. .	Fl.	P.	6-7
a. <i>obtusifolius</i> , <i>M. & K.</i> . .	. . . . .	. . . . .	. .	. .	. .	. .
b. <i>acutifolius</i> , <i>Link.</i> . . .	. . . . .	. . . . .	. .	. .	. .	. .
c. <i>zosterifolius</i> , <i>Schum.</i> . .	. . . . .	. . . . .	. .	. .	. .	. .
D.						
viii. <i>P. crispus</i> , <i>L.</i>	<i>curly P.</i>	Ditches .	. .	Fl.	P.	6-7
ix. <i>P. perfoliatus</i> , <i>L.</i> . . .	<i>perfoliate P.</i> . . .	Ditches .	. .	Fl.	P.	6-7
x. <i>P. lucens</i> , <i>L.</i>	<i>shining P.</i>	Lakes . .	. .	Fl.	P.	6-7
b. <i>acuminatus</i> , <i>Schum.</i> . .	. . . . .	. . . . .	. .	. .	. .	. .
c. <i>decipiens</i> , <i>Nolte</i>	. . . . .	. . . . .	. .	. .	. .	. .
xi. <i>P. prælongus</i> , <i>Wulf.</i> .	<i>long-stalked P.</i> . .	Lakes! .	. .	Fl.	P.	7-8
SEC. II.						
A.						
xii. <i>P. heterophyllus</i> , <i>Schreb.</i>	<i>various-leaved P.</i>	Ditches .	. .	Fl.	P.	6-7
a. <i>heterophyllus</i>	. . . . .	. . . . .	. .	. .	. .	. .
b. <i>nitens</i> , <i>Weber</i>	. . . . .	. . . . .	. .	. .	. .	. .
c. <i>lanceolatus</i> , <i>Sm.</i>	. . . . .	Anglesea!	. .	. .	. .	. .
xiii. <i>P. rufescens</i> , <i>Schrad.</i>	<i>reddish P.</i>	Ditches!	. .	Fl.	P.	7
B.						
xiv. <i>P. natans</i> , <i>L.</i>	<i>broad-leaved P.</i> . .	Ponds . .	. .	Fl.	P.	6-7
xv. <i>P. polygonifolius</i> , <i>Pourr.</i>	. . . . .	Ponds . .	. .	Fl.	P.	6-7
xvi. <i>P. plantagineus</i> , <i>Ducr.</i>	<i>Plantain-leaved P</i>	Peaty ditches	. .	Fl.	P.	6-7

ORD. LXXXIX.—ALISMACEÆ—*cont.*

CHARACTERISTICS OF SPECIES AND VARIETIES.

Gen. VI.

- i. Ls. few, alt., semi-cylind.—Scape with a term. rac. of about five fis.

ORD. XC.—NAIADACEÆ.

Gen. I.

SEC. I.—*Ls. all submerged.*

A. Ls. opp.—Stip. o.

- i. Ls. ov.-lanc., amplex, pellucid—Spikes short, few-flowered.

B. Ls. alt., lin.—Stip. adnate.

- ii. St. fili., channelled—Ls. setaceous, one-nerved, sheathing—Spikes interrupted—Fr. with conspicuous keels.

v. 2. *Ls. flat, three-nerved—Keel of fr. obscure.*

- iii. St. fili.—Ls. channelled, one-nerved—Spike very interrupted—Fr. small, not keeled.

C. Ls. alt., lin.—Stipules free.

- iv. St. fili., much branched—Ls. setaceous, one-veined.

- v. St. fili., cylind.—Ls. three-veined, acute.

- vi. St. compressed, winged—Ls. five-veined—Spike short, lax, interrupted.

- vii. St. slightly compressed—Ls. three-veined—Spike ovate, dense, continuous.

a. Ls. obt.—Spike oval—Pedunc. short.

b. Ls. acu.—Spike oval—Pedunc. short.

c. Ls. acu.—Spike cylind.—Pedunc. long.

D. Ls. alt., ov. or obl.—Stip. free.

- viii. St. flattened—Ls. obl., obt., wavy—Fr. with a long beak.

- ix. St. round, stout—Ls. cord.-ov., amplex, pellucid.

- x. St. stout—Ls. stalked, ellip.-lanc., pellucid—Stip. winged—Spike cylind., many flowered.

(*The largest British species.*)

b. Ls. longer and narrower, the mid-rib running out into a point.

c. Ls. sess., cusp.—Stip. not winged—Pedunc. shorter.

- xi. Ls. elong.-obl., semi-amplex, concave at tip—Stip. not winged—Spike cylind., dense—Pedunc. long.

SEC. II.—*Ls. alternate, upper ones floating—Stip. free.*

A. Submerged Ls. sess.

- xii. St. slender—Submerged Ls. lanc.—Flo. Ls. ellip.—Stip. small, ribbed—Pedunc. thickened upwards.

a. Ls. flat, not amplex; upper coriac., opp., fl.

b. Ls. recurved, all submerged, with many rooting branches from the upper axils.

c. Ls. dark gr.; obt.—Pedunc. not thickened.

- xiii. St. round, simp.—Ls. reddish, lanc., many-nerved—Pedunc. equal.

B. Submerged Ls. stalked.

- xiv. Ls. all coriaceous, ellip.; lower Ls. lin., often absent—Fr. large, keeled when dry.

- xv. Upper Ls. coriaceous, ellip.-obl., rounded at base; lower Ls. lin.-lanc., membr.—Fr. small, rounded when dry.

- xvi. Ls. all membr., pellucid, beautifully netted—Fr. small, keeled when dry.

ORD. XC.—NAIADACEÆ—*cont.*

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
Gen. II. <i>Ruppia</i>, L.	<i>Tassel Pond-weed.</i>					
i. <i>R. maritima</i> , L.	<i>Sea T.</i>	Ditches in salt mar.	..	Fl.	P.	7-8
a. <i>maritima</i>	.. . . .	.. . . .	..	..	..	..
b. <i>rostellata</i> , Koch.	.. . . .	.. . . .	..	..	..	..
Gen. III. <i>Zannichellia</i>, L.	<i>Horned Pond- weed.</i>					
i. <i>Z. palustris</i> , L.	<i>common H.</i>	Ditches .	..	Fl.	A.	5-8
a. <i>palustris</i>	.. . . .	.. . . .	..	..	..	..
b. <i>pedicellata</i> , Fries.	.. . . .	.. . . .	..	..	..	..
Gen. IV. <i>Zostera</i>, L.	<i>Grass-wrack.</i>					
i. <i>Z. marina</i> , L.	<i>broad-leaved G.</i> .	Salt waters.	..	Fl.	P.	7-8
v. 2. <i>angustifolia</i>	.. . . .	.. . . .	..	..	..	..
ii. <i>Z. nana</i> , Roth.	<i>dwarf G.</i>	Salt waters.	..	Fl.	P.	4-8
Gen. V. <i>Naias</i>, L.	<i>Naias.</i>					
i. <i>N. flexilis</i> , Rostk.	<i>flexible N.</i>	Lakes! I.	..	Fl.	P.	8

ORD. XCI.—LEMNACEÆ.

Gen. I. <i>Lemna</i>, L.	<i>Duckweed.</i>					
i. <i>L. minor</i> , L.	<i>lesser D.</i>	Ponds . .	..	Fl.	A.	7-8
ii. <i>L. gibba</i> , L.	<i>gibbous D.</i>	Ponds . .	..	Fl.	A.	6-9
iii. <i>L. polyrhiza</i> , L.	<i>greater D.</i>	Ponds . .	..	Fl.	A.	7-8
iv. <i>L. trisulca</i> , L.	<i>Ivy-leaved D.</i> . . .	Ponds . .	..	Fl.	A.	6-7
Gen. II. <i>Wolffia</i>, Horkel.						
i. <i>W. arrhiza</i> , Wimm.	.. . . .	Ponds! .	..	Fl.	A.	6-7

ORD. XCII.—ARACEÆ.

Gen. I. <i>Arum</i>, L.	<i>Cuckoo-pint.</i>					
i. <i>A. maculatum</i> , L.	<i>Cuckoo-pint, Wake Robin, or Lords & Ladies.</i>	Waysides	..	6-10 in.	P.	4-5
[ii. <i>A. italicum</i> , Mill.	.. . . .	I. of W. .	..	8-12 in.	P.	7-8
Gen. II. <i>Acorus</i>, L.	<i>Sweet-flag.</i>					
i. <i>A. calamus</i> , L.	<i>common S.</i>	Marshes.	..	3-6 ft.	P.	6-7

ORD. XC.—NAIADACEÆ—*cont.*

CHARACTERISTICS OF SPECIES AND VARIETIES.

Gen. II.

- i. St. fili., much branched—Ls. lin.—Fls. on a spadix arising from the leaf-sheath.
- a.* Sheaths inflated—Pedunc. long, spirally twisted.
- b.* Sheaths small, close—Ls. narrower—Pedunc. short, flexuous.

Gen. III.

- i. Ls. opp., fili.—Fls. axill., sess.
- a.* Clusters of fr. sess.—Sty. half as long as fr.
- b.* Clusters of fr. stalked—Sty. as long as fr.

Gen. IV.

- i. Ls. (1-3 ft.) lin., 3-7 veined—Spadix many-flowered, without bands.
- v.* 2. *Ls. narrower (1-3 veined).*
- ii. Ls. (6-8 in.) lin., 1-3 veined—Spadix few-flowered, with clasping bands.

Gen. V.

- i. Ls. lin., minutely dentic., opp. or three in a whorl—Fls. axill., sol.

ORD. XCI.—LEMNACEÆ.

Gen. I.

- i. Fronds ob.-ov., compressed—Root sol.
- ii. Fronds ob.-ov., tumid below—Root sol.
- iii. Fronds broadly ob.-ov.—Roots many, clustered.
- iv. Fronds thin, ellip.-lanc., tip serr.—Roots sol.

Gen. II.

- i. Fronds obl., very minute, like a grain of sand.
(*The smallest known flowering plant.*)

ORD. XCII.—ARACEÆ.

Gen. I.

- i. Ls. vernal, sagitt., spotted, with deflexed lobes.
- ii. Ls. appearing before winter, triang., with diverging lobes.

Gen. II.

- i. Ls. ensif.—Scape leaf-like, with spadix in middle—Sweet scented

ORD. XCIII.—TYPHACEÆ.

Botanical Name.	Common Name.	Soil or Situation.	Colour	Growth.	Dura- tion.	Time of flowering.
Gen. I. Sparganium, L.	<i>Bur-weed.</i>					
i. <i>S. natans, L.</i>	<i>floating B.</i>	Lakes! .	..	Fl.	P.	8-9
a. <i>natans, L.</i>	. . . . .	. . . . .	..	. . .	. . .	. . .
b. <i>minimum, Fries.</i>	. . . . .	. . . . .	..	. . .	. . .	. . .
ii. <i>S. simplex, Huds.</i>	<i>unbranched B.</i>	Ditches .	..	1-2 ft.	P.	7-8
iii. <i>S. ramosum, Huds.</i>	<i>branched B.</i>	Ditches .	..	2-4 ft.	P.	6-7
Gen. II. Typha, L.	<i>Reed-mace, Cat's- tail, Bulrush.</i>					
i. <i>T. latifolia, L.</i>	<i>great B., &c.</i>	Ponds . .	..	3-7 ft.	P.	7-8
ii. <i>T. angustifolia, L.</i>	<i>lesser B., &c.</i>	Ponds! .	..	1-3 ft.	P.	7-8

ORD. XCIV.—RESTIACEÆ.

Gen. I. Eriocaulon, L.	<i>Pipe-wort.</i>					
i. <i>E. septangulare, With.</i>	<i>jointed P.</i>	Lakes! .	..	Fl.	P.	8-9

ORD. XCV.—JUNCACEÆ.

Gen. I. Narthecium, Huds.	<i>Bog-Asphodel.</i>					
i. <i>N. ossifragum, Huds.</i>	<i>Lancashire B.</i> . .	Mount. bogs.	Y.	6-8 in.	P.	6-8
Gen. II. Juncus, L.	<i>Rush.</i>					
* i. <i>J. communis, Mey.</i>	<i>common R.</i>	Wet pl. .	..	1-3 ft.	P.	7-8
a. <i>conglomeratus, L.</i>	. . . . .	. . . . .	..	. . .	. . .	. . .
b. <i>effusus, L.</i>	. . . . .	. . . . .	..	. . .	. . .	. . .
c. <i>diffusus, Hoppe.</i>	. . . . .	. . . . .	..	. . .	. . .	. . .
ii. <i>J. glaucus, Ehrh.</i>	<i>hard R.</i>	Wet pl. .	..	1-3 ft.	P.	7-8
iii. <i>J. balticus, Willd.</i>	<i>Baltic R.</i>	San. sh.!	..	1-2 ft.	P.	7-8
iv. <i>J. filiformis, L.</i>	<i>Thread R.</i>	Lakes sh.!	..	4-8 in.	P.	7-8
* * *						
v. <i>J. maritimus, Sm.</i>	<i>lesser Sea R.</i>	Salt mar.	..	1-3 ft.	P.	7-8
vi. <i>J. acutus, L.</i>	<i>great Sea R.</i>	San. sh.!	..	2-4 ft.	P.	7-8

ORD. XCIII.—TYPHACEÆ.

CHARACTERISTICS OF SPECIES AND VARIETIES.

Gen. I.

- i. St. simple—Ls. floating, flat, concave at base.
- a. Barren heads several, sess.—Fr. stalked, beak long.
- b. Barren head sol.—Fr. sess., shortly beaked.
- ii. St. simple—Ls. erect, triang. at base, with flat sides.
- iii. St. branched—Ls. erect, triang. at base, with concave sides.

Gen. II.

- i. Ls. lin., flat—Spikes contiguous.
- ii. Ls. lin., channelled below—Spikes separate..

ORD. XCIV.—RESTIACEÆ.

Gen. I.

- i. Ls. subu., glab.—Scape long ; fls. in a compact scaly head.

ORD. XCV.—JUNCACEÆ

Gen. I.

- i. Ls. rigid, ribbed, acum.—Pedunc. with one bract at base and one above the middle.

Gen. II.

* *Ls. none—Pan. lat.—Fls. not clustered—Seeds with a close testa.*

- i. St. soft, faintly striate ; pith continuous—Pan. very much branched, variable—Seg. of perian. lanc. longer than the retuse caps.—Sta. three.
- a. Perian. olive—Pan. dense, glob.—Fr. apicu.
- b. Perian. olive-gr.—Pan. lax—Fr. not apicu.
- c. St. rigid, striate—Caps. ob.-ov., mucro.—(*Probably a hybrid between J. communis and J. glaucus.*)
- ii. St. rigid, glauc.—Pith interr.—Pan. lax, much branched—Seg. of perian. narrow, lin. ; as long as the ellip.-obl. caps.—Sta. six.
- iii. Root creeping, not tufted—St. rigid—Pith continuous—Pan. slightly branched—Seg. of perian. ov.-lanc., acu. as long as the ellip., abruptly mucro. caps. —Sta. three.
- iv. St. fili., slend.—Pan. small, few-flowered, about middle of st.—Caps. obt., mucro.—Sta. six.
- * * *Ls. none—Pan. lat.—Fls. clustered—Seeds with a loose testa.*
- v. St. rigid—Segm. of peri. lanc., acu., without scarious margins, as long as the ellip. caps.
- vi. St. rigid—Segm. of peri. ov.-lanc., with a scarious margin, half as long as the oval caps.

ORD. XCV.—JUNCACEÆ—*cont.*

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
* * *						
vii. <i>J. acutiflorus</i> , Ehrh. .	<i>sharp-flowered jointed R.</i>	Bogs . .	. .	1-2 ft.	P.	6-8
viii. <i>J. lamprocarpus</i> , Ehr.	<i>shining-fruited jointed R.</i>	Bogs . .	. .	1-2 ft.	P.	7-8
v. 2. <i>nigritellus</i> , Don. . .	. . . . .	. . . . .	. .	. .	. .	. .
ix. <i>J. uliginosus</i> , Siph. .	<i>lesser jointed R.</i>	Bogs . .	. .	1-2 ft.	P.	7-8
x. <i>J. obtusifolius</i> , Ehrh.	<i>blunt-flowered jointed R.</i>	Bogs . .	. .	2-3 ft.	P.	7-8
* * * *						
†						
xi. <i>J. castaneus</i> , Sm. . .	<i>clustered Alp. R.</i>	Sc.mount.!	. .	6-12 in.	P.	7-8
xii. <i>J. trifidus</i> , L. . . .	<i>three-leaved R.</i>	Sc. bogs!	. .	. .	. .	. .
††						
xiii. <i>J. compressus</i> , Jacq. .	<i>round-fruited R.</i>	Bogs . .	. .	1-2 ft.	P.	6-8
v. 2. <i>Gerardi</i> , Lois. . . .	. . . . .	. . . . .	. .	. .	. .	. .
xiv. <i>J. bufonius</i> , L. . . .	<i>Toad R.</i>	Wat. pl.	. .	1-8 in.	A.	7-8
* * * * *						
A.						
xv. <i>J. squarrosus</i> , L. . . .	<i>Heath R.</i>	Heaths .	. .	4-10 in.	P.	6-7
xvi. <i>J. capitatus</i> , Wei. . .	<i>capitate R.</i>	Jersey .	. .	1-4 in.	P.	5-7
B.						
xvii. <i>J. biglumis</i> , L. . . .	<i>two-flowered R.</i>	Sc.mount.!	. .	2-6 in.	P.	7-8
xviii. <i>J. triglumis</i> , L. . . .	<i>three-flowered R.</i>	Sc.mount.!	. .	6-12 in.	P.	7-8
Gen. III. <i>Luzula</i> , DC.						
	<i>Woodrush.</i>					
i. <i>L. sylvatica</i> , Bich. . .	<i>great hairy W.</i>	Woods .	. .	12-18 in.	P.	5-6
ii. <i>L. pilosa</i> , Willd. . . .	<i>broad-leaved hairy W.</i>	Woods .	. .	6-12 in.	P.	5-6
v. 2. <i>Borreri</i> , Bromf. . . .	. . . . .	. . . . .	. .	. .	. .	. .
iii. <i>L. Forsteri</i> , DC. . . .	<i>narrow-leaved hairy W.</i>	Woods .	. .	6-12 in.	P.	3-6
iv. <i>L. campestris</i> , Willd.	<i>Field W.</i>	Heaths .	. .	4-10 in.	P.	4-6
a. <i>campestris</i>	. . . . .	. . . . .	. .	. .	. .	. .
b. <i>multiflora</i> , Lej.	. . . . .	. . . . .	. .	. .	. .	. .
v. <i>L. arcuata</i> , Hook. . . .	<i>curved mount W.</i>	Sc.mount.!	. .	1-4 in.	P.	7
vi. <i>L. spicata</i> , DC.	<i>spiked mount W.</i>	Sc. mount.	. .	6-12 in.	P.	7

ORD. XCVI.—CYPERACEÆ.

Gen. I. <i>Cyperus</i> , L.						
	<i>Cyperus</i> , <i>Galin- gale.</i>					
i. <i>C. longus</i> , L.	<i>sweet C.</i>	Marshes! S. of E.	. .	2-3 ft.	P.	8-9

CHARACTERISTICS OF SPECIES AND VARIETIES.

* * * *Sts. leafy—Ls. terete, or compressed, jointed internally—Pan. term.—Fls. fasciculate; seeds not appendaged.*

- vii. St. and ls. sub-compressed—Pan. pyram., compound; segm. of peri. acum., as long as the acum., pale brown caps.
- viii. St. and ls. compressed—Pan. repeatedly forked—Outer segm. of peri. acute; inner obt.; shorter than the ob.-lanc., dark brown caps.
- v. 2. *Outer sep. acute, shorter than the lin.-obl., rostr., black caps.*
- ix. St. slender—Ls. setaceous, slightly grooved—Pan. nearly simp., irreg.—Seg. of peri. obl.; outer three acute, inner obt., nearly as long as the ellip., obt., pale brown caps.
- x. St. and ls. terete—Pan. spreading—Seg. of peri., obt., as long as the ov., acum., caps.

* * * * *St. leafy—Ls. flat or grooved, not jointed.*

† *Seeds appendaged.*

- xi. St. with two channelled ls.—Pan. term.—Seg. of peri. ellip.-lanc., half as long as the ov.-obl. caps.
- xii. St. one-leaved—Pan. term. 1-3 flowered, with two long leafy bracts.
†† *Seeds not appendaged.*
- xiii. St. tufted, with one leaf in middle—Pan. compact—Seg. of peri. ov.-obl., obt., shorter than the ov., mucro. caps.
- v. 2. *Sts. more dist.—Seg. of peri. as long as caps.*
- xiv. St. slend., dichotomously branched—Fls. sol., sess.—Seg. of peri. longer than the obl., obt. caps.

* * * * * *Ls. all rad.—Fls. term.*

A. *Seeds not appendaged.*

- xv. Ls. rigid—Pan. term., compound—Seg. of peri. ov.-lanc., as long as the obt. caps.
- xvi. Ls. soft, fili.—Seg. of peri. unequal, arist.; twice as long as the acum. caps.
B. *Seeds appendaged.*
- xvii. Sts. scattered, channelled on one side—Ls. not channelled—Caps. turb., retuse.
- xviii. Sts. tufted, round—Ls. channelled—Caps. ellip., acute.

Gen. III.

- i. Pan. cymose, doubly compound—Pedunc. long, three-flowered—Sep. arist., as long as the ov., acute, beaked caps.
- ii. Pan. little branched—Pedunc. one-flowered, reflexed in fr.—Sep. acum., shorter than the broadly ov., obt. caps.
- v. 2. *Caps. shorter, acute, not ripening.*
- iii. As last, but more slender—Ls. lin., pedunc. erect and caps. acute.
- iv. Pan. of three or four round, term., sess. or stalked spikes—Caps. blunt.
  - a. Filaments much shorter than the anth.—Seeds glob.
  - b. Filaments nearly as long as anth.—Seeds twice as long as broad.
- v. Pan. sub-umbell., of few 3-5 flowered clusters on a long, drooping pedunc.
- vi. Pan. spike-like, dense-flowered, drooping.

ORD. XCVI.—CYPERACEÆ.

Gen. I.

- i. Spikelets in erect, doubly compound, umbels—Root creeping.

ORD. XCVI.—CYPERACEÆ—cont.

Botanical Name.	Common Name.	Soil or Situation.	Colour	Growth.	Dura- tion.	Time of flowering.
ii. <i>C. fuscus</i> , <i>L.</i>	<i>brown C.</i>	Wetmead! Surrey.	..	2-6 in.	A.	8-9
Gen. II. <i>Cladium</i>, <i>Schrad.</i> <i>Twig-rush.</i>						
i. <i>C. mariscus</i> , <i>Br.</i>	<i>prickly T.</i>	Bogs! . .	..	3-5 ft.	P.	7-8
Gen. III. <i>Schoenus</i>, <i>L.</i> <i>Bog-rush.</i>						
i. <i>S. nigricans</i> , <i>L.</i>	<i>black B.</i>	Bogs! . .	..	6-18 in.	P.	6-7
Gen. IV. <i>Rhynchospora</i>, <i>Vahl.</i> <i>Beak-rush.</i>						
i. <i>R. alba</i> , <i>Vahl.</i>	<i>white B.</i>	Bogs . .	..	6-18 in.	P.	6-7
ii. <i>R. fusca</i> , <i>Sm.</i>	<i>brown B.</i>	Bogs! S.W. of E.	..	6-12 in.	P.	7-8
Gen. V. <i>Blysmus</i>, <i>Panz.</i> <i>Blysmus.</i>						
i. <i>B. compressus</i> , <i>Panz.</i>	<i>broad-leaved B.</i> .	Bogs . .	..	4-10 in.	P.	6-7
ii. <i>B. rufus</i> , <i>Link.</i>	<i>narrow-leaved B.</i>	Bogs . .	..	3-12 in.	P.	6-7
Gen. VI. <i>Scirpus</i>, <i>L.</i> <i>Club-rush, Bul- rush, &c.</i>						
i. <i>S. lacustris</i> , <i>L.</i>	<i>lake C., or B.</i> . .	Marshes	..	1-8 ft.	P.	7-8
b. <i>Tabernæmontani</i> , <i>Gmel.</i>	. . . .	Sea sh.!	..	..	..	..
c. <i>carinatus</i> , <i>Sm.</i>	. . . .	Riv. bnks.!	..	..	..	..
ii. <i>S. triqueter</i> , <i>L.</i>	<i>triangular</i> . .	Riv. bnks.!	..	1-3 ft.	P.	8-9
iii. <i>S. pungens</i> , <i>Vahl.</i> . .	<i>sharp C.</i>	Jersey! .	..	6-18 in.	P.	6-7
iv. <i>S. maritimus</i> , <i>L.</i>	<i>Salt marsh C.</i> . .	Salt mar.	..	1-3 ft.	P.	7-8
v. <i>S. sylvaticus</i> , <i>L.</i>	<i>Wood C.</i>	Woods .	..	1-3 ft.	P.	7-8
Gen. VII. <i>Isolepis</i>, <i>Br.</i> <i>Mud-rush.</i>						
i. <i>I. setacea</i> , <i>Br.</i>	<i>bristle-stalked M.</i>	Moi. pl. .	..	3-6 in.	P.	7-8
ii. <i>I. Savii</i> , <i>Schultz.</i> . .	<i>Savi's M.</i>	Bogs! . .	..	3-10 in.	P.	7-8
iii. <i>I. fluitans</i> , <i>Br.</i>	<i>floating M.</i>	Ditches .	..	Flo.	P.	6-7
iv. <i>I. Holoschoenus</i> , <i>L.</i> . .	<i>cluster-headed M.</i>	San. sh.!	..	2-3 ft.	P.	9-10
Gen. VIII. <i>Eleocharis</i>, <i>Br.</i> <i>Spike-rush.</i>						
i. <i>E. palustris</i> , <i>Br.</i>	<i>creeping S.</i>	Bogs . .	..	6-12 in.	P.	6-7
a. <i>palustris.</i>	. . . . .	. . . .	..	..	..	..
b. <i>uniglumis</i> , <i>Link.</i> . . .	. . . . .	. . . .	..	..	..	..
c. <i>Watsoni.</i>	. . . . .	. . . .	..	..	..	..

CHARACTERISTICS OF SPECIES AND VARIETIES.

- ii. Spikelets in small, round heads—Root tufted.

Gen. II.

- i. St. leafy—Edges of ls. rough—Cymes, term., and lat., much branched ; spikelets capit.

Gen. III.

- i. St. naked, round—Spikelets 5-10 in a term., round head, shorter than the lower bract.

Gen. IV.

- i. Spikelets in a compact head, as long as the outer bracts—Sta. two.
ii. Spikelets in an oval head, much shorter than the outer bracts—Sta. three.

Gen. V.

- i. St. triang.—Ls. lin., flat, rough—Spikelets 6-8 flowered, longer than bract.
ii. St. round—Ls. channelled, smooth—Spikelets 2-4 flowered, not longer than bract.

Gen. VI.

* *Stem leafless.*

- i. St. terete—Ls. often flo.—Cymes term., doubly compound—Glu. notched, mucro., fringed, glab.—Fr. trigonous.
b. St. glauc.—Flo. ls. absent—Glu. rough—Fr. compressed.
c. St. obscurely triang. above—Glu. glab.—Fr. compressed.
ii. St. acutely triang.—Spikes ov., clustered—Glu. glab., notched, mucro.—Fr. obtusely lobed.
iii. St. acutely triang.—Spikes few, sess.—Glu. bifid, with acute lobes.
* * *Stem leafy.*
iv. Spike in a dense, term., clust.—Glu. bifid, with acute lobes.
v. Spike in a very compact pan.—Glu. ent., obt..

Gen. VII.

* *Spikes 1-3 sess.*

- i. Lower bract very long—Fr. transversely striate.
ii. Lower bracts not longer than spikelets—Fr. rough, with raised points.
* * *Spikelets sol., term.—St. floating.*
iii. St. much branched, leafy, flo.—Spikes ovate.
* * * *Spikelets many, in glob. heads.*
iv. St. round—Ls. channelled.

Gen. VIII.

- i. Root creeping, with many tufts—St. round, with membr. transversely trunc. sheaths—Fr. plano-convex—Term. glu. acute—Stig. two.
a. Lower glu. half surrounding the spikelet.
b. Lower glu. almost surrounding the spikelet.
c. Smaller and more rigid—Lower glu. surrounding the spikelet.

ORD. XCVI.—CYPERACEÆ—*cont.*

Botanical Name.	Common Name.	Soil . or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
ii. <i>E. multicaulis</i> , <i>Sm.</i> .	<i>many-stalked S.</i> .	Bogs . .	. .	6-12 in.	P.	6-7
iii. <i>E. pauciflora</i> , <i>Link.</i> .	<i>Chocolate-headed S.</i>	Bogs Sco.!	. .	6-12 in.	P.	7-8
iv. <i>E. cæspitosa</i> , <i>Link.</i> .	<i>scaly-stalked S., or Deer's-hair.</i>	Moors. .	. .	3-12 in.	P.	6-7
v. <i>E. acicularis</i> , <i>L.</i> . . .	<i>least S.</i>	San. sh. of lakes.	. .	2-6 in.	P.	7-8
vi. <i>E. parvula</i> , <i>L.</i>	. . . . .	Mud. sh. I.!	. .	1-2 in.	A.	7-8
Gen. IX. Eriophorum, L. <i>Cotton-grass.</i>						
*						
i. <i>E. alpinum</i> , <i>L.</i>	<i>Alpine C.</i>	Alp. bogs!	. .	6-10 in.	P.	6
ii. <i>E. vaginatum</i> , <i>L.</i> . . .	<i>Hare's-tail C.</i> . . .	Boggy moors.	. .	6-10 in.	P.	4-5
* *						
iii. <i>E. polystachyon</i> , <i>L.</i> .	<i>common C.</i>	Bogs . .	. .	6-15 in.	P.	5-6
a. <i>angustifolium</i> , <i>Roth.</i> .	. . . . .	. . . . .	. .	. . . . .	. .	. .
b. <i>latifolium</i> , <i>Hoppe.</i> . .	. . . . .	. . . . .	. .	. . . . .	. .	. .
iv. <i>E. gracile</i> , <i>Koch.</i> . . .	<i>slender C.</i>	Bogs! . .	. .	1-2 ft.	P.	6-7
Gen. X. Kobresia, Willd.						
i. <i>K. caricina</i> , <i>Willd.</i> . .	. . . . .	Moors! N. of E.	. .	4-8 in.	P.	8-9
Gen. XI. Carex, L. <i>Sedge.</i>						
SEC. I.						
A.						
i. <i>C. dioica</i> , <i>L.</i>	<i>separate-headed S.</i>	Bogs . .	. .	6-12 in.	P.	5-6
ii. <i>C. pulicaris</i> , <i>L.</i>	<i>Flea S.</i>	Bogs . .	. .	3-8 in.	P.	5-6
B.						
iii. <i>C. rupestris</i> , <i>All.</i> . . .	<i>Rock S.</i>	Alp. rocs.!	. .	4-6 in.	P.	7-8
iv. <i>C. pauciflora</i> , <i>Lightf.</i> .	<i>few-flowered S.</i> . .	Sc.mount.!	. .	3-10 in.	P.	6-7
SEC. II.						
A.						
v. <i>C. incurva</i> , <i>Lightf.</i> . . .	<i>curved S.</i>	San. sh.!	. .	2-4 in.	P.	6-7
B.						
vi. <i>C. ovalis</i> , <i>Good.</i>	<i>oval-spiked S.</i> . . .	Bogs . .	. .	8-12 in.	P.	6-7
vii. <i>C. stellulata</i> , <i>Good.</i> . .	<i>little prickly S.</i> . .	Marshes.	. .	6-12 in.	P.	5-6
viii. <i>C. curta</i> , <i>Good.</i> . . .	<i>white S.</i>	Bogs . .	. .	6-12 in.	P.	6-7
v. 2. <i>alpicola</i> , <i>Wahl.</i> . . .						
ix. <i>C. lagopina</i> , <i>Wahl.</i> . .	<i>Hare's-foot S.</i> . . .	Sc.mount.!	. .	. . . . .	. .	. .
x. <i>elongata</i> , <i>L.</i>	<i>elongated S.</i>	Sc.mount.!	. .	4-8 in.	P.	7-8
		Marshes!	. .	1-2 ft.	P.	7-8

CHARACTERISTICS OF SPECIES AND VARIETIES.

- ii. Root hardly creeping, with one tuft—St. round, with obliquely trunc., acu., leafless, sheaths—Fr. ob.-ov., three-sided—Term. glu., obt.—Stig. three—Bristles 4-6, longer than fr.
- iii. Stems tufted, with long stolons, round, striate, with narrow, leafless sheaths—Fr. striate, obl., mucro.—Lower glu. obt., shorter than spikelets—Bristles 3-6, shorter than fr.
- iv. Root densely tufted, wiry—St. round, with leafless sheaths, and short, subu. ls. below—Stig. three—Fr. ob.-ov., acum.; two lower glu. awned, longer than the spikelet.
- v. Root with fili., stolons—St. quadrangular, with leafless sheaths—Spikelets minute—Stig. three—Glu. obt.—Fr. ob.-ov., ribbed and transversely striate.
- vi. Very small—Root fili., with dense tufts—St. round, with one leafless sheath—Ls. fili., rad.

Gen. IX.

* *Spikelets sol.*

- i. St. triang.—Spikelet obl., few-flowered—Bristles 4-6, crisped.
- ii. St. triang. above, round below—Spikelet ov., many-flowered—Bristles many, straight.

* * *Spikelets several, lateral.*

- iii. St. rigid—Ls. flat—Fer. spikelets 4-12—Fr. ob.-ov., mucro.
  - a. Ls. channelled near the base—Pedunc. glab.—Glu. ov.
  - b. Ls. quite flat—Pedunc. scab.—Glu. lanc.
- iv. St. slender—Ls. trique.—Pedunc. downy—Fr. ob.-ov., obt.

Gen. X.

- i. St. densely tufted—Ls. wiry, grooved, shorter than st.—Spikelets aggregate.

Gen. XI.

SEC. I.—*Spikelets solitary, simple.*A. *Stig. two.*

- i. Diæcious—Spikelet ov., dense—Fr. ov., ascend., serrulated at margin.
- ii. Androgynous—Spikelet with upper part barren—Fr. lanc.-triang., deflexed.
- B. *Stig. three.*
- iii. Ls. flat, ending in a twisted point—Upper half of spikelet barren—Fr. ob.-ov., adpr.—Glu. persist.
- iv. One or two term. fls. barren—Fr. few, lan.-subu., reflexed.

SEC. II.—*Spikelets compound, continuous.*A. *Spikelets capitate, barren at top—Root creeping—Bractæas membranous.*

- v. Spikelets in a glob. head—St. curved—Fr. broadly ov.—Ls. channelled.

B. *Spikelets barren at base—Root tufted—Bractæas membranous.*

- vi. Spikelets fuscous, about six, alt., ov., approx.—Fr. ov.-acum., erect, with a membr. marg.; beak bifid.
- vii. Spikelets pale, three or four, glob., remote—Fr. broadly ov., spreading, much beaked; beak bifid.
- viii. Spikelets pale, about six, ellip., rather remote—Fr. broadly ov., compressed, acute, faintly striate; beak emarg.
- v. 2. *Smaller—Beak split to base.*
- ix. Spikelets fuscous, ellip., contig.—Fr. erect, ellip.; beak entire, scarious at apex.
- x. Spikelets pale, many, obl., approx.—Fr. spread., obl.-acum., many ribbed; beak almost entire, with rough edges.

ORD. XCVI.—CYPERACEÆ—*cont.*

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering
<i>C.</i>						
xi. <i>C. remota</i> , <i>L.</i> . .	<i>distant-spiked S.</i>	Sh. pl. . .	. .	1-2 ft.	P.	6-7
xii. <i>C. axillaris</i> , <i>Good.</i>	<i>axillary clustered S.</i>	Marshes !	. .	1-2 ft.	P.	6-7
xiii. <i>C. Boenninghausiana</i> , <i>Weihe.</i>	. . . .	Marshes !	. .	1-2 ft.	P.	6-7
<i>D.</i>						
xiv. <i>C. paniculata</i> , <i>L.</i>	<i>great panicled S.</i>	Marshes.	. .	1-4 ft.	P.	6-7
xv. <i>C. paradoxa</i> , <i>Willd.</i>	. . . .	Marshes !	. .	1-3 ft.	P.	6-7
xvi. <i>C. teretiuscula</i> , <i>Good.</i>	<i>lesser panicled S.</i>	Marshes.	. .	1-2 ft.	P.	6-7
v. 2. <i>Ehrartiana</i> , <i>Hop.</i>	. . . .	. . . .	. .	. .	. .	. .
xvii. <i>C. vulpina</i> , <i>L.</i> . .	<i>great S.</i>	Marshes.	. .	2-4 ft.	P.	6-7
<i>E.</i>						
xviii. <i>C. divulsa</i> , <i>Good.</i>	<i>gray S.</i>	Marshes.	. .	1-2 ft.	P.	5-6
xix. <i>C. muricata</i> , <i>L.</i> .	<i>great prickly S.</i> .	Marshes.	. .	1-2 ft.	P.	5-6
* *						
xx. <i>C. arenaria</i> , <i>L.</i> .	<i>Sea S.</i>	San. sh. .	. .	8-10 in.	P.	6-7
xxi. <i>C. intermedia</i> , <i>Good.</i>	<i>soft brown S.</i> . .	Marshes.	. .	1-3 ft.	P.	6-7
xxii. <i>C. divisa</i> , <i>Huds.</i> .	<i>bracteate Marsh S.</i>	Marshes.	. .	1-3 ft.	P.	6-7
<i>F.</i>						
xxiii. <i>C. VahlII</i> , <i>Schk.</i> .	<i>close-headed Alpine S.</i>	Sc.mount.!	. .	6-18 in.	P.	6-7
xxiv. <i>C. canescens</i> , <i>L.</i>	<i>hoary S.</i>	Lough Neagh, I.!	. .	1-2 ft.	P.	7-8
xxv. <i>C. atrata</i> , <i>L.</i> . .	<i>black S.</i>	Mount.!	. .	1-2 ft.	P.	7-8
SEC. III.						
*						
<i>A.</i>						
xxvi. <i>C. rigida</i> , <i>Good.</i> .	<i>rigid S.</i>	Mount.!	. .	4-6 in.	P.	6-8
xxvii. <i>C. acuta</i> , <i>L.</i> . . .	<i>slender-spiked S.</i>	Wet pl. .	. .	2-3 ft.	P.	6-8
xxviii. <i>C. stricta</i> , <i>Good.</i> .	<i>tufted Bag S.</i> . .	Bogs . .	. .	1-3 ft.	P.	5-6
<i>B.</i>						
xxix. <i>C. aquatilis</i> , <i>Wahl.</i>	<i>Water S.</i>	Sco. bogs.	. .	1-2 ft.	P.	6-8

CHARACTERISTICS OF SPECIES AND VARIETIES.

C. Spikelets barren at their base, or at both ends ; lower distant—Root tufted—Bracteas leafy.

- xi.** Spikelets all simp.; upper crowded—Fr. ov.-lanc., longer than the glu.; beak notched—Ls. channelled.
- xii.** Spikelets ; upper simp., crowded ; lower remote, compound—Fr. ov.-lanc., serr. above, as long as the glu.; beak deeply bifid—Ls. flat.
- xiii.** Spikelets ; upper simp.; lower compound—Fr. lanc., as long as the glu., serr. from below the middle ; beak almost ent.

(Probably a hybrid between eleven and twelve.)

D. Spikelets barren at top, compound, approx., paniced—Root tufted—Bracteas membranous.

- xiv.** Spikelet thrice compound, paniced, branches spreading—Root densely tufted—St. trique, with flat sides—Fr. ov., many veined, gibb.; beak winged.
  - xv.** Spikelet elongate, lower branched, dist.—St. slender, trique, with convex sides—Fr. ov., gibb., with many short veins near the base ; beak not winged.
  - xvi.** Spikelet obl., compact, acute—Root forming scattered tufts—St. trique, with convex sides—Fr. ov., gibb., with 3-4 central nerves ; beak winged.
- v. 2.** *Root more tufted—Spikelet longer, interr.*

- xvii.** Spikelet crowded, cylind., obt.—St. triquet, with concave sides—Fr. ov.-acum., spreading, not gibb.; beak bifid, finely serr.

E. Spikelets barren at top, alt., simp.

* Root tufted.

- xviii.** Spikelets (5-6) dist., lax, upper approx.—Fr. ov., acute, erect, smooth, bifid.
- xix.** Spikelets (4-6), compact, approx., in an oval spike—Fr. ov.-acum., spreading, finely serr. above, bifid.

* * Root creeping.

- xx.** Spikelets in an obl., interr., spike : upper barren ; interm. barren at end—Fr. ov., veined, winged, finely serr. above ; shorter than the glu.
- xxi.** Spikelets in an obl., interr., spike ; upper and lower fert.; intermediate barren—Fr. ov.-lanc. narrowly winged, longer than the glu.
- xxii.** Spikelets in a crowded head, with a leafy bract at base—Fr. plano-convex, veined, not winged ; beak finely serr., acutely bifid.

F. Tern spikelet with barren and fertile fls.; the rest fertile.

- xxiii.** Spikelets (1-4), roundish, contig., sess.—Fr. ob.-ov., three-angled, shortly beaked, scabr. above, longer than the ov.-obt. glu.
- xxiv.** Spikelets (3-4) obl., contig., lower rather dist. and shortly stalked—Fr. ov., compressed, at length three-angled, bifid, shorter than the ov., cusp. glu.—Leaf sheaths with a filamentous edge.
- xxv.** Spikelets (3-4) ov.-obl., contig., shortly stalked, at length drooping—Fr. ellip., three-angled when ripe, rather shorter than the glu.; beak terete, short, slightly notched.

SEC. III.—Terminal spikelets barren—The lower ones fertile.

* Stig. two (rarely three in thirty).

A. Margins of ls. revolute when dry.

- xxvi.** Barren spikelet one ; fert. short, ov., approx.—Bracts sheathing at base—Fr. ellip., obt., not ribbed.
- xxvii.** Barren spikelets 1-3 ; fert. slender, cylind., remote, erect in fr.—Lower bract long, with long aur.—Fr. obl., ribbed.
- xxviii.** Barren spikelet one ; fert. (2-3), approx., erect, thick, lower bract short—Sheaths of ls. filamentous—Fr. ellip.-obl., acute, compr., veined.

B. Margins of ls. involute when dry.

- xxix.** Fert., spikelets elong., narrowed below—Fr. ellip., doubly convex, not veined.

ORD. XCVI.—CYPERACEÆ—*cont.*

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
xxx. <i>C. vulgaris</i> , <i>Fries.</i> * * A. † †	<i>common S.</i> . . .	Marshes.	. .	½-2 ft.	P.	6-8
xxxxi. <i>C. limosa</i> , <i>L.</i> . . v. 2. <i>irrigua</i> , <i>Sm.</i> . .	<i>Mud S.</i>	Bogs!	. . . .	6-12 in. . .	P. . .	6-8 . .
xxxii. <i>C. rariflora</i> , <i>Sm.</i> .	<i>loose-flowered Alpine S.</i>	Alp. bogs!	. .	6-12 in.	P.	6-8
xxxiii. <i>C. glauca</i> , <i>Scop.</i> .	<i>glaucous heath S.</i>	Moors. .	. .	½-2 ft.	P.	6-7
xxxiv. <i>C. capillaris</i> , <i>L.</i> .	<i>dwarf capillary S.</i>	Sc.mount.!	. .	4-8 in.	P.	6-7
xxxv. <i>C. pendula</i> , <i>Huds.</i>	<i>great pendulous S.</i>	Wet wds.	. .	3-6 ft.	P.	5-6
xxxvi. <i>C. vaginata</i> , <i>Tausch.</i> † †	<i>short brown spiked S.</i>	Sc.mount.!	. .	1-2 ft.	P.	6-7
xxxvii. <i>C. pallescens</i> , <i>L.</i> .	<i>pale S.</i>	Marshes.	. .	1-2 ft.	P.	6-7
xxxviii. <i>C. panicea</i> , <i>L.</i> . .	<i>pink-leaved S.</i> .	Marshes.	. .	1-2 ft.	P.	6-7
† † †						
xxxix. <i>C. præcox</i> , <i>Jacq.</i>	<i>vernal S.</i>	Heaths .	. .	6-12 in.	P.	4-5
xl. <i>C. tomentosa</i> , <i>L.</i>	<i>downy-fruited S.</i>	Wet fie., Wilts!	. .	12-18 in.	P.	6-7
xli. <i>C. pilulifera</i> , <i>L.</i> .	<i>round-headed S.</i> .	Heaths .	. .	6-18 in.	P.	6-7
xlii. <i>C. montana</i> , <i>L.</i> .	<i>Mountain S.</i> . .	Woods! W. of E.	. .	6-18 in.	P.	4-5
xliii. <i>C. ericetorum</i> , <i>Poll.</i>	. . . . .	Li. banks! Gogmagog hills.	. .	2-6 in.	P.	5-6
† † †						
xliv. <i>C. digitata</i> , <i>L.</i> . .	<i>fingered S.</i>	Li. woods! W. of E.	. .	6-10 in.	P.	5-6
xlv. <i>C. clandestina</i> , <i>L.</i>	<i>dwarf silvery S.</i> .	Li. rocks! W. of E.	. .	1-3 in.	P.	5-6
B. †						
xlvi. <i>C. hirta</i> , <i>L.</i> . . .	<i>hairy S.</i>	Damp woods.	. .	1-2 ft.	P.	5-6
xlvii. <i>C. filiformis</i> , <i>L.</i> .	<i>slender-leaved S.</i>	Bogs!	. .	2-3 ft.	P.	5-6
xlviii. <i>C. extensa</i> , <i>Good.</i>	<i>long bracteate S.</i> .	Salt marshes!	. .	10-18 in.	P.	6-7

CHARACTERISTICS OF SPECIES AND VARIETIES.

- xxx. Fert. spikelets, cylind., sub-sess.—Fr. ellip., plano-convex, many veined.
 * * *Stigmas three (two in saxatilis).*
A. Beak of fruit entire, or obliquely notched.
 † *Fr. glabrous.*
 † *Fertile spikelets drooping.*
- xxxxi. Fert. spikelets (1-2) long-stalked, ov., dense-flowered—Rt. cr.—Ls. narrow, rough at edge—Bracts auricled—Fr. round, ov., compr., ribbed.
 v. 2. *Ls. broader, rough only at point—Lower glu. narrow, attenuate; upper glu. broader and shorter.*
- xxxii. Fert. spikelets (2-3) obl., lax, few-flowered—Rt. cr.—Bracts with short sheaths—Fr. obl., acum., shorter than the broad, blunt glu.
- xxxiii. Fert. spikelets (2-3) dense-flowered, at length drooping—Bracts not sheathing, —Fr. ellip., obt., not veined, scab., as long as the acu. glu.
- xxxiv. Fert. spikelets, lax., few-flowered—Rt. tufted—Lower bracts sheathing, and including several pedunc.—Fr. obl.-acum. as long as the ov., obt. glu.
- xxxv. Fert. spikelets (4-5) cylind., very long stalked, dense-flowered—Bracts sheathing—Fr. imbric., ellip., tumid., with a short emarg. beak.
 † † *Fert. spikes erect.*
- xxxvi. Fert. spikelets lax, distant—Fr. ellip.-lanc., three-edged, beak trunc., decurved, obliquely notched.
- xxxvii. Fert. spikelets ov., approx.—Fr. ov.-obl., obt., doubly convex, veined; beak o.
- xxxviii. Fert. spikelets remote, lax-flowered—Fr. ov., tumid., not veined; beak short, terete.
 † † *Fruit hairy.*
 † *Bracts not sheathing.*
- xxxix. Fert. spikelets (1-3) ov.-obl., sess., approx.—Rt. cr.—Bracts clasping—Fr. ov., three-sided, as long as the ov.-acum., glu.
- xl. Fert. spikelets (1-2) short, cylind., shortly stalked—Rt. cr.—Lower bracts leaf-like, slightly sheathing—Fr. ob.-ov., densely downy, longer than the small acu. glu.; beak notched.
- xli. Fert. spikelets (2-3) round, sess., approx.—Rt. fibrous—Bracts small, subu.—Fr. sub-glob., stalked, as long as the broadly ov., mucro., glu.; beak short, notched.
- xl. Fert. spikelets (1-3) ov., sess., approx.—Rt. fibrous—Bracts small, membran.—Fr. obl.-ov., longer than the ob.-ov., minute glu.; beak short, notched.
- xl. Fert. spikelets few, capit., small, sess.—St. curved, short—Fr. ob.-ov., three-sided, as long as the broadly ov., ciliate, obt., glu.; beak ent.
 † † *Bracts sheathing.*
- xl. Ls. revolute—Fert. spikelets, dist., erect, lin., lax—Lower bract with a setac. point.
- xl. Ls. involute—Fert., spikelets very remote, of 3-4 fls. enclosed in the memb. bracts.
B. Beak of fruit long, bifid.
 † *Fr. hairy—Barren spikelet sol.*
- xl. Fert. spikelets (2-3) cylind., stalked—Lower bracts with long sheaths—Ls. flat, hairy—Fr. ov.-obl., with a long beak, deeply bifid at point.
- xl. Fert. spikelets (3-4) obl., sess.—Lower bracts slightly sheathed—Ls. channelled—Fr. ov.-obl. narrowing into a short deeply bifid beak.
 † † *Fr. glabrous—Barren spikelets sol.; fertile erect.*
- xl. Fert. spikelets, obl., approx., sub-sess.—Bracts long, leafy, with short sheaths—Ls. convolute—Fr.-ov., three-sided, ribbed, inflated; beak short, smooth, straight.

ORD. XCVI.—CYPERACEÆ—*cont.*

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
xlix. <i>C. flava</i> , <i>L.</i>	<i>yellow S.</i>	Boggy heaths.	..	3-18 in.	P.	5-6
v. 2. <i>Æderi</i> , <i>Auct.</i>	.. . . .	.. . . .	..	..	..	..
v. 3. <i>lepidocarpa</i> , <i>Tausch.</i>	.. . . .	.. . . .	..	..	..	..
1. <i>C. distans</i> , <i>L.</i>	<i>loose S.</i>	Marshes!	..	1-2 ft.	P.	6-7
li. <i>C. fulva</i> , <i>Good.</i>	<i>tawny S.</i>	Marshes.	..	12-18 in.	P.	6-7
v. 2. <i>speirostachya</i>	.. . . .	.. . . .	..	..	..	..
lii. <i>C. binervis</i> , <i>Sm.</i>	<i>green-ribbed S.</i> . .	Heaths .	..	1-3 ft.	P.	6-7
liii. <i>C. lævigata</i> , <i>Sm.</i> . . .	<i>smooth-stalked beaked S.</i>	Marshes.	..	1-3 ft.	P.	6-7
liv. <i>C. punctata</i> , <i>Gaud.</i> . .	<i>dotted-fruited S.</i>	Marshes! S. of E. & I.	..	12-18 in.	P.	6-7
lv. <i>C. depauperata</i> , <i>Good.</i>	<i>starved wood S.</i> . .	Dry wds.!	..	1-2 ft.	P.	5-6
† † †						
lvi. <i>C. sylvatica</i> , <i>Huds.</i> . .	<i>pendulous Wood S.</i>	Damp woods.	..	1-3 ft.	P.	5-6
lvii. <i>C. strigosa</i> , <i>Huds.</i> . .	<i>loose pendulous S.</i>	Woods .	..	1-2 ft.	P.	5-6
† † † †						
*						
lviii. <i>C. saxatilis</i> , <i>L.</i> . . .	<i>russet S.</i>	Sc. mount.	..	4-10 in.	P.	5-6
v. 2. <i>Grahami</i> , <i>Boott.</i> . .	.. . . .	.. . . .	..	..	..	..
* *						
†						
lix. <i>C. vesicaria</i> , <i>L.</i>	<i>short beaked Bladder S.</i>	Bogs . .	..	1-2 ft.	P.	5-6
lx. <i>C. ampullacea</i> , <i>Good.</i> .	<i>slender beaked Bladder S.</i>	Bogs . .	..	1-2 ft.	P.	5-6
† †						
lxi. <i>C. Pseudo-cyperus</i> , <i>L.</i>	<i>Cyperus-like S.</i> . .	Riv. banks	..	1-3 ft.	P.	6-7
lxii. <i>C. paludosa</i> , <i>Good.</i> . .	<i>lesser common S.</i>	Ditches .	..	2-3 ft.	P.	5-6
lxiii. <i>C. riparia</i> , <i>Curt.</i> . . .	<i>great common S.</i>	Ditches .	..	3-5 ft.	P.	5-6

ORD. XCVII.—GRAMINEÆ.

Gen. I. <i>Leersia</i> , <i>Soland.</i>	<i>Cut-grass.</i>					
i. <i>L. oryzoides</i> , <i>Sw.</i> . .	<i>European C.</i> . .	Ditches!	..	2-3 ft.	P.	8-10

CHARACTERISTICS OF SPECIES AND VARIETIES.

- xliv.** Fertile spikelets, roundish-ov., approx.—Bracts long, leafy, with short sheaths—Fr. ob.-ov., inflated, spread., ribbed, much longer than the glu.; beak deflexed, scabrid.
- v. 2. Smaller—Spikelets all approx.—Beak straight.*
- v. 3. Barren spikelets long stalked—Fr. small, suddenly narrowed in a short, smooth beak.*
- 1.** Fert. spikelets obl., erect, dist., upper stalks included in sheath of bracts—Fr. ov., equally ribbed, smooth; beak narrow, rough edged.
- li.** Fert. spikelets ov.-obl., dist.—Bracts with long sheaths; stalks exserted—Fr. broadly ov., ribbed; beak straight, rough edged.
- v. 2. Fertile spikelets longer stalked—Beak smoother.*
- lii.** Fert. spikelets, cylind., dist.—Stalks included in sheaths—Fr. ov., with two green ribs on outer surface; beak broad, smooth.
- liii.** Fertile spikelets cylind., dist.; stalks exserted—Fr. ov.-atten., striate; beak long, with scabrous edges.
- liv.** Fertile spikelets, cylind.; stalks slightly exserted—Fr. ov., tumid., obsoletely veined, pellucidly punctate; beak lin., smooth.
- lv.** Fert. spikelets erect, 3-4 flowered; stalks exserted—Fr. large, glob.; beak long.
† † † *Fruit glabrous—Barren spikelets sol.—Fertile drooping.*
- lvi.** Fertile spikelets fili.; stalks half exserted—Fr. ellip., obscurely veined; beak long, smooth.
- lvii.** Fertile spikelets slender; stalks exserted—Fr. obl.-lanc., veined, narrowed at each end; beak obscure, obliquely trunc.
† † † † *Fruit glabrous—Barren spikelets several—Fertile usually drooping, stout.*
* *Stigmas two.*
- lviii.** Fertile spikelets (1-3) roundish-ov., lower stalked—Fr. ov., inflated, obscurely veined; beak short, notched.
- v. 2. Taller—Fr. prominently veined.*
(Probably an Alpine form of *C. vesicaria*, to which the *Var. 2, Grahami*, forms the passage.)
* * *Stigmas three.*
‡ *Fr. inflated.*
- lix.** Fertile spikelets, cylin., distant—Fr. broadly ov., gradually narrowed into the beak—St. sharply angled—Ls. flat.
- lx.** Fertile spikelets cylind., dist.—Fr. sub-glob., suddenly narrowed into a long slender beak—St. with obt. angles—Ls. channelled.
‡ ‡ *Fruit not inflated.*
- lxi.** Fertile spikelets (3-6) dense, drooping, on long stalks—Fr. obl., very attenu., ribbed; beak long.
- lxii.** Fertile spikelets about three, cylind., obt., erect—Fr. obl.-ov., compressed—Glu. of barren spikelets obt.
- lxiii.** Fertile spikelets cylind., acute, erect—Fr. ob.-ov., convex on both sides—Glu. of the barren spikelets acute.

ORD. XCVII.—GRAMINEÆ.

Gen. I.

- i.** Pan. diff., few-flowered, with wavy branches, mostly enclosed in sheath—Ls. broad, glauc., rough edged.

ORD. XCVII.—GRAMINEÆ—*cont.*

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
Gen. II. <i>Spartina</i>, Schred.	<i>Cord-grass.</i>					
i. <i>S. stricta</i> , Roth. . . .	<i>Cord-grass</i> . . .	Salt mar.	. .	1-2 ft.	P.	8-9
ii. <i>S. alternifolia</i> , Loisel.	<i>many-spiked C.</i> .	Salt mar.!	. .	1-2 ft.	P.	8-9
Gen. III. <i>Cynodon</i>, Rich.	<i>Dog's-tooth-grass.</i>					
i. <i>C. Dactylon</i> , Pers. . .	. . . .	San. sh.!	. .	4-8 in.	P.	7-8
Gen. IV. <i>Digitaria</i>, Scop.	<i>Finger-grass.</i>					
i. <i>D. humifusa</i> , Pers. . .	. . . .	San. fie.!	. .	6-8 in.	A.	7-8
Gen. V. <i>Panicum</i>, L.	<i>Panic-grass.</i>					
[i. <i>P. Crus-Galli</i> , L. . .	<i>loose P.</i>	Fields! .	. .	2-3 ft.	A.	7-8
Gen. VI. <i>Setaria</i>, Beauv.	<i>Bristle-grass.</i>					
i. <i>S. viridis</i> , Beauv. . .	<i>green B.</i>	Fields! .	. .	1-2 ft.	A.	7-8
[ii. <i>S. verticillata</i> , Beauv.	<i>rough B.</i>	Fields! .	. .	1-2 ft.	A.	7-8
Gen. VII. <i>Phalaris</i>, L.	<i>Canary-grass.</i>					
i. <i>P. arundinacea</i> , L. . .	<i>reed C.</i>	San. sh. .	. .	2-3 ft.	P.	7-8
[ii. <i>P. canariensis</i> , L. . .	<i>Canary-grass</i> . .	Was. pl. .	. .	1-2 ft.	P.	6-8
Gen. VIII. <i>Anthoxan- thum</i>, L.	<i>Vernal-grass.</i>					
i. <i>A. odoratum</i> , L. . . .	<i>sweet V.</i>	Past. . .	. .	1-2 ft.	P.	6-7
Gen. IX. <i>Hierochloe</i>, Gmel.	<i>Holy-grass.</i>					
i. <i>H. borealis</i> , R. & S.	<i>northern H.</i> . . .	Thurso .	. .	10-12 in	P.	5-6
Gen. X. <i>Phleum</i>, L.	<i>Cat's-tail-grass.</i>					
* i. <i>P. alpinum</i> , L.	<i>Mountain C.</i> . .	Mount. .	. .	1-2 ft.	P.	7-8
ii. <i>P. pratense</i> , L. . . .	<i>common C.</i> . . .	Past. . .	. .	2-3 ft.	P.	6-7
* * [iii. <i>P. arenarium</i> , L. . . .	<i>Sea C.</i>	San. sh. .	. .	6-12 in.	A.	6-7
iv. <i>P. Boehmeri</i> , Wibel. .	<i>purple-stalked C.</i>	Past. . .	. .	1-2 ft.	P.	7-8
Gen. XI. <i>Alopecurus</i>, L.	<i>Fox-tail-grass.</i>					
i. <i>A. alpinus</i> , Sm. . . .	<i>Alpine F.</i>	Mount.!. .	. .	1-2 ft.	P.	7-8
ii. <i>A. pratensis</i> , L. . . .	<i>common F.</i> . . .	Past. . .	. .	1-2 ft.	P.	4-6
iii. <i>A. geniculatus</i> , L. . .	<i>floating F.</i> . . .	Ditches .	. .	6-12 in.	P.	4-8
iv. <i>A. fulvus</i> , Sm.	<i>Orange-spiked F.</i>	Ditches!	. .	6-12 in.	P.	7-8

CHARACTERISTICS OF SPECIES AND VARIETIES.

Gen. II.

- i. Spikes 2-3, longer than the ls.—Outer glu. one-nerved, hairy—Tip of rachis as long as last spikelet.
- ii. Spikes many, shorter than the ls.—Outer glu. five-nerved, glab. except on keel—Tip of rachis longer than last spikelet.

Gen. III.

- i. Spikes 3-5, digitate, purplish—Ls. downy beneath—Root creeping.

Gen. IV.

- i. Spikes 3-5, digitate—Spikelets purplish, ovate, in pairs.

Gen. V.

- i. Spike with alt. or opp. branches—Spikelets crowded—Fls. unilat.]

Gen. VI.

- i. Pan. cylind.—Bristles of the invol. ascending.
- ii. Pan. narrow, lobed—Bristles of the invol. deflexed.]

Gen. VII.

- i. Pan. erect, with spread. branches—Glu. not keeled.
- ii. Pan. ovate, spike-like—Glu. with broad keels.]

Gen. VIII.

- i. Spike ov.-obl.—Fls. longer than their awns.—Sta. two.

Gen. IX.

- i. Pan. erect—Spike ov., brown—Paleæ awnless.

Gen. X.

- * *Glu. truncate, with a bristly point.*
- i. Pan. ov.-cylind.—Awn as long as glu.
- ii. Pan. cylind., long—Awn not half as long as glu.
- * * *Glu. acute at each end.*
- iii. Pan. ov.-lanc., blunt, narrowed at base—Glu. lanc., fringed.
- iv. Pan. cylind.—Glu. lin.-lanc., obliquely trunc.

Gen. XI.

- i. St. erect—Pan. obl.—Glu. ov., suddenly acute, conn. at base—Awn scarcely longer than the paleæ.
- ii. St. erect—Pan. cylind., obt.—Glu. lanc., acu., hairy, conn. at base—Awn twice as long as paleæ.
- iii. St. asc., bent at joints—Pan. cylind., obt., glu. obt., conn. at base—Awn longer than paleæ, inserted at base.
- iv. Very like A. geniculatus, but the awn is not longer than palea, and is inserted at middle—Spike pale, slender.

ORD. XCVII.—GRAMINEÆ—*cont.*

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
v. <i>A. bulbosus</i> , <i>L.</i> . . .	<i>tuberous F.</i> . . .	Salt mar.!	. .	12-18in.	P.	5-7
vi. <i>A. agrestis</i> , <i>L.</i> . . .	<i>slender F.</i> . . .	Past. . .	. .	1-2 ft.	P.	4-6
Gen. XII. Knappia , <i>Smith.</i>	<i>Knappia.</i>					
i. <i>K. agrostidea</i> , <i>Sm.</i> . .	. . . .	San. sh.!	. .	2-4 in.	P.	3-4
Gen. XIII. Gastridium , <i>Beauv.</i>	<i>Nit-grass.</i>					
i. <i>G. lendigerum</i> , <i>Gaud.</i>	<i>awned N.</i>	Fie.!	. .	12-18in.	P.	6-10
Gen. XIV. Polypogon , <i>Desf.</i>	<i>Beard-grass.</i>					
i. <i>P. littoralis</i> , <i>Sm.</i> . . .	<i>perennial B.</i> . .	Salt mar.!	. .	1-2 ft.	P.	7-8
ii. <i>P. monspeliensis</i> , <i>Desf.</i>	<i>annual B.</i>	Fie.!	. .	12-18in.	A.	7-8
Gen. XV. Miliun , <i>L.</i>	<i>Millet-grass.</i>					
i. <i>M. effusum</i> , <i>L.</i> . . .	. . . .	Sha. woods	. .	1-2 ft.	P.	6-7
Gen. XVI. Apera , <i>Beauv.</i>	<i>Bent-grass.</i>					
i. <i>A. Spica-venti</i> , <i>Beauv.</i>	<i>silky B.</i>	Sand fie.	. .	2-3 ft.	A.	7-8
v. 2. <i>interrupta</i>	. . . .	. . .	. .	. .	. .	. .
Gen. XVII. Agrostis , <i>L.</i>	<i>Bent-grass.</i>					
*						
i. <i>A. setacea</i> , <i>Curt.</i> . .	<i>bristle-leaved B.</i> .	Heaths .	. .	1-2 ft.	P.	7-8
ii. <i>A. canina</i> , <i>L.</i>	<i>brown B.</i>	Fields .	. .	1-2 ft.	P.	6-7
***						
iii. <i>A. vulgaris</i> , <i>With.</i> . .	<i>fine B.</i>	Past. . .	. .	12-18in.	P.	7-8
v. 2. <i>pumila</i> , <i>Light.</i> . .	. . . .	. . .	. .	. .	. .	. .
iv. <i>A. alba</i> , <i>L.</i>	<i>Marsh B.</i>	Past. . .	. .	1-2 ft.	P.	7-8
v. 2. <i>stolonifera</i> , <i>L.</i> . . .	. . . .	. . .	. .	. .	. .	. .
Gen. XVIII. Ammophila , <i>Host.</i>	<i>Marram</i>					
i. <i>A. arundinacea</i> , <i>Host.</i>	<i>M., Sea-reed, or Mat-grass.</i>	San. sh. .	. .	2-4 ft.	P.	7-8
Gen. XIX. Arundo , <i>L.</i>	<i>Reed.</i>					
i. <i>A. Phragmites</i> , <i>L.</i> . .	<i>common R.</i> . . .	Ditches, &c.	. .	6-10 ft.	P.	7-8
Gen. XX. Calamagrostis , <i>Adans.</i>	<i>Small-reed.</i>					
*						
i. <i>C. Epigejos</i> , <i>L.</i> . . .	<i>Wood S.</i>	Moi. pl. .	. .	3-5 ft.	P.	6-7
ii. <i>C. lanceolata</i> , <i>Roth.</i> .	<i>purple flowered S.</i>	Moi. pl. .	. .	3-4 ft.	P.	6-7

CHARACTERISTICS OF SPECIES AND VARIETIES.

- v. St. erect—Pan. cylin., slender—Glu. abruptly acum., free at base—Awn half as long again as palea.
 vi. St. erect, rough—Pan. cylind., acum.—Glu. acu., united to middle, glab.—Awn more than half as long again as palea.

Gen. XII.

- i. Sts. many—Ls. short, rough—Glu. longer than the paleæ.

Gen. XIII.

- i. Pan. large, dense—Glu. lanc., acum.—Awns twice as long as glu.

Gen. XIV.

- i. Root cr.—Glu. acu., as long as awns.
 ii. Root fibr.—Glu. obt., half as long as awns.

Gen. XV.

- i. Pan. diff.; branches spread., sub-verticillate, remote at base.

Gen. XVI.

- i. Pan. spread., with very slender branches—Awn three or four times longer than paleæ.
 v. 2. *Pan. close, interrupted—Anthers shorter.*

Gen. XVII.

* *Paleæ awned.*

- i. Ls. setac.; sheath rough—Pan. close, obt.
 ii. Ls. narrow; sheaths smooth—Pan. spread. in fr.; otherwise close.

* * *Paleæ awnless.*

- iii. Pan. always spread.—Ligule short, trunc.—Sheaths smooth.
 v. 2. *Short, stout, tufted (2-3 in. high).*
 iv. Pan. compact after flowering—Ligule long, acute—Sheaths rough.
 v. 2. *Stems with long prostrate stolons.*

Gen. XVIII.

- i. Ls. long, rigid, glauc., invol.—Pan. white, cylind., acum.—Pairs one-third as long as acute glu.

Gen. XIX.

- i. Pan. large and spread., purple, nodding—Spikelets five-flowered, coloured, ribbed with a tuft of silky hairs at base.

Gen. XX.

* *Hairs longer than the paleæ.*

- i. Pan. erect, close—Spikelets crowded, unilat.—Awn projecting beyond the paleæ.
 ii. Pan. loose—Spikelets spreading—Ls. narrower and smoother than the last—Awn scarcely projecting.

ORD. XCVII.—GRAMINEÆ—*cont.*

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
* *						
iii. <i>C. stricta</i> , <i>Nutt.</i> . . .	<i>narrow S.</i> . . .	Bogs! . .	. .	1-3 ft.	P.	6-7
Gen. XXI. Sesleria , <i>Scop.</i> <i>Moor-grass.</i>						
i. <i>S. cœrulea</i> , <i>Scop.</i> . . .	<i>blue M.</i>	Mount. .	. .	1 ft.	P.	5-6
Gen. XXII. Aira , <i>L.</i> <i>Hair-grass.</i>						
*						
i. <i>A. cœspitosa</i> , <i>L.</i> . . .	<i>tufted H.</i>	Past. . .	. .	1-3 ft.	P.	7-8
ii. <i>A. alpina</i> , <i>L.</i>	<i>Alpine H.</i>	Mount.! .	. .	6-12 in.	P.	6-7
iii. <i>A. flexuosa</i> , <i>L.</i> . . .	<i>waved H.</i>	Heaths .	. .	1-2 ft.	P.	6-7
* *						
iv. <i>A. caryophyllæa</i> , <i>L.</i> .	<i>Silver H.</i>	Hilly past.	. .	6-9 in.	P.	6-7
v. <i>A. præcox</i> , <i>L.</i>	<i>early H.</i>	Dry pl. .	. .	1-4 in.	A.	5-6
* * *						
vi. <i>A. canescens</i> , <i>L.</i> . . .	<i>grey H.</i>	San. fie. .	. .	1-2 ft.	P.	7-8
en. XXIII. Lagurus , <i>L.</i> <i>Hare's-tail-grass.</i>						
i. <i>L. ovatus</i> , <i>L.</i>	. . . . .	San. sh.! Guernsey.	. .	6-12 in.	A.	6-7
Gen. XXIV. Avena , <i>L.</i> <i>Oat-grass.</i>						
*						
i. <i>A. fatua</i> , <i>L.</i>	<i>wild O., or Havers.</i>	Corn fie..	. .	2-3 ft.	A.	6-7
ii. <i>A. strigosa</i> , <i>Shreb.</i> . .	<i>Bristle O.</i>	Corn fie.!	. .	2-3 ft.	A.	7-8
* *						
iii. <i>A. pratensis</i> , <i>L.</i> . . .	<i>narrow-leaved O.</i>	Dry past.	. .	12-18 in.	P.	7-8
v. 2. <i>alpina</i> , <i>Sm.</i>	. . . . .	. . . . .	. .	. . . . .	. .	. .
iv. <i>A. pubescens</i> , <i>L.</i> . . .	<i>downy O.</i>	Cha. pas.	. .	1-2 ft.	P.	6-7
v. <i>A. flavescens</i> , <i>L.</i> . . .	<i>yellow O.</i>	Cha. pas.	. .	1-2 ft.	P.	6-7
Gen. XXV. Arrhenathe- <i>False Oat-grass.</i>						
rum , <i>Beauv.</i>						
i. <i>A. avenaceum</i> , <i>Beauv.</i>	. . . . .	Waysides	. .	2-3 ft.	P.	6-7
Gen. XXVI. Holcus , <i>L.</i> <i>Soft-grass.</i>						
i. <i>H. lanatus</i> , <i>L.</i>	<i>Meadow S.</i>	Mead. .	. .	12-18 in.	P.	7-8
ii. <i>H. mollis</i> , <i>L.</i>	<i>creeping S.</i>	Mead. .	. .	6-12 in.	P.	7-8
Gen. XXVII. Triodia , <i>Heath-grass.</i>						
<i>Brown.</i>						
i. <i>P. decumbens</i> , <i>Beauv.</i>	<i>decumbent H.</i> . .	Mount. past.	. .	6-12 in.	P.	7-8
Gen. XXVIII. Koeleria , <i>Pers.</i>						
i. <i>K. cristata</i> , <i>Pers.</i> . .	. . . . .	Dry past.	. .	6-8 in.	P.	6-7

CHARACTERISTICS OF SPECIES AND VARIETIES.

* * *Hairs shorter than the paleæ.*

- iii. Pan. narrow, erect, close—Glu. lanc., acum.

Gen. XXI.

- i. Much tufted—Ls. lin., obt., with a small rough point—Pan., ov.-obl.; fls. imbricated.

Gen. XXII.

* *Lower paleæ trunc., toothed.*

- i. Pan. spread.—Ls. flat—Awn straight, from below middle of palea, scarcely exceeding the glu.
 ii. Pan. close—Ls. (mostly) invol.—Awn bent, from above the middle of palea, scarcely exceeding the glu.
 iii. Pan. spread., with wavy branches—Ls. setaceous—Awn bent, from near the base of palea, and much exceeding it.

* * *Lower paleæ bifid.*

- iv. Pan. very spread.—Awn from below the middle of palea—Sheaths scabrous.

- v. Pan. obt., spike-like—Awn from near base of palea—Sheaths glab.

* * * *Lower paleæ entire—Awn jointed with tuft of hairs at base.*

- vi. Pan. long, rather dense—Ls. setaceous.

Gen. XXIII.

- i. Spikelets soft, ov.—Ls. broadly lanc., downy.

Gen. XXIV.

* *Inner glu. 5-7 ribbed—Spikelets ultimately drooping.*

- i. Spikelets three-flowered—Lower paleæ bifid at end, with fulvous hairs at base.

- ii. Spikelets two-flowered—Lower paleæ glab., ending in two long straight bristles.]

* * *Inner glu. with three ribs—Spikelets erect.*

- iii. Rad. ls., glauc., glab.—Pan. lax; spikelets 3-6 flowered.

- v. 2. *Lower ls short, suddenly acum.—Spikelets 5-6 flowered.*

- iv. Rad. ls. hairy—Pan. nearly simp.—Outer paleæ jagged—Ligule acute.

- v. Rad. ls. hairy—Pan. much branched—Outer paleæ with two term. bristles—Ligule obt.

Gen. XXV.

- i. Pan. simple, lax—Ls. rough, flat, scabrous.

Gen. XXVI.

- i. Inner glu. obt.—Awn included, smooth except at tip—Rt. fibr.

- ii. Inner glu. acute—Awn exserted, rough throughout—Root cr.

Gen. XXVII.

- i. Pan. racemose—Spikelets few—Ligule only a tuft of hairs.

Gen. XXVIII.

- i. Pan. interrupted below—Lower palea acute, with a short awn.

ORD. XCVII.—GRAMINEÆ—*cont.*

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
Gen. XXIX. Melica, L.	<i>Melic-grass.</i>					
i. <i>M. uniflora, Retz.</i> . . .	<i>Wood M.</i>	Sh. woods	..	1-2 ft.	P.	5-7
ii. <i>M. nutans, L.</i>	<i>Mountain M.</i> . . .	Mount. woods.	..	1-2 ft.	P.	5-6
Gen. XXX. Molinia, Schränk.						
i. <i>M. cærulea, Mænnch.</i> . .	<i>purple M.</i>	Moors. .	..	1-2 ft.	P.	7-8
Gen. XXXI. Catabrosa, Beauv.	<i>Whorl-grass.</i>					
i. <i>C. aquatica, Beauv.</i> . . .	<i>Water W.</i>	Ditches .	..	½-2 ft.	P.	5-6
Gen. XXXII. Glyceria, Br.	<i>Meadow-grass.</i>					
i. <i>G. aquatica, Sm.</i>	<i>Reed M.</i>	Ditches .	..	4-6 ft.	P.	4-8
ii. <i>G. fluitans, Br.</i>	<i>floating M., or Manna-croup G.</i>	Ditches .	..	1-2 ft.	P.	7-8
a. <i>fluitans</i>	.....	.....	..	..	..	..
b. <i>plicata</i>	.....	.....	..	..	..	..
Gen. XXXIII. Schleroch- loa, Lindl.	<i>Meadow-grass.</i>					
* A.						
i. <i>S. maritima, Lindl.</i>	<i>Sea M.</i>	Salt mar.	..	6-12 in.	P.	7-9
B.						
ii. <i>S. distans, Bab.</i>	<i>reflexed M.</i>	San. pl. .	..	1-2 ft.	P.	7-10
v. 2. <i>Borreri, Bab.</i>	.....	Salt mar.	..	..	..	..
iii. <i>S. procumbens, Beauv.</i>	<i>procumbent M.</i> . .	Salt mar.	..	6-8 in.	A.	7-8
* *						
iv. <i>S. rigida, Link.</i>	<i>hard M.</i>	Dry pl. .	..	4-6 in.	A.	6-7
v. <i>S. loliacea, Woods</i> . . .	<i>Wheat M.</i>	San. sh. .	..	6-12 in.	A.	6-7
Gen. XXXIV. Poa, L.	<i>Meadow-grass.</i>					
A.						
i. <i>P. annua, L.</i>	<i>annual M.</i>	Was. pl. .	..	2-10 in.	A.	3-10
B.						
* ii. <i>P. pratensis, L.</i>	<i>smooth M.</i>	Mead. . .	..	1-2 ft.	P.	6-7
iii. <i>P. compressa, L.</i>	<i>flat-stalked M.</i> . .	Dry pl. .	..	6-8 in.	P.	6-7
* *						
iv. <i>P. trivialis, L.</i>	<i>rough M.</i>	Mead. . .	..	1-2 ft.	P.	6-7

CHARACTERISTICS OF SPECIES AND VARIETIES.

Gen. XXIX.

- i. Pan. lax, slightly drooping—Spikelets erect, of one perfect fl.
- ii. Pan. a unilat, nearly simp. raceme—Spikelets drooping, with two perfect fls.

Gen. XXX.

- i. Pan. long, narrow, erect, purplish—Spikelets 1-3 flowered.

Gen. XXXI.

- i. Pan. long, pyram. ; branches whorled, patent—Ls. lanc., obt.

Gen. XXXII.

- i. Pan. erect, much branched—Spikelets obt., 5-10 flowered.
- ii. Pan. secund., slightly branched—Spikelets 7-12 flowered.

- a. Branches erect in fr.—Sheaths striate.
- b. Branches spread. in fr.—Sheaths furrowed.

Gen. XXXIII.

* *Glumes three-veined.*

A. Root creeping.

- i. Pan. erect, rigid; lower branches in pairs—Ls. invol.—Lower paleæ obt., apic., mid-rib reaching to tip.

B. Root tufted.

- ii. Perenn.—Pan. long, branched; branches at length deflexed—Ls. flat—Lower paleæ obt., mid-rib not reaching to tip.
- v. 2. *Branches ascending in fr.—Spikelets smaller, crowded (midway between S. maritima and S. distans).*
- iii. Ann.—Pan. ob.-lanc., compact, secund.; branches short, rigid—Lower paleæ five-ribbed.

** *Glumes one-veined.*

- iv. Pan. compound, rigid, distichous—Spikelets lin.-acum.—Upper glu. reaching to base of third floret.
- v. Pan. racemose, simp., with alt., sol., lin.-obl. spikelets—Upper glu. reaching to base of fourth floret.

Gen. XXXIV.

A. Root annual.

- i. Pan. spread.—Spikelets five-flo., ov.—Ls. invol.

B. Root perennial.

* *Root creeping; with stolons.*

- ii. St. terete—Pan. spread.—Upper leaf shorter than sheath.
- iii. St. compressed—Pan. slightly secund., condensed; spikelets ov.—Upper leaf as long as sheath.

** *Root fibrous; or creeping with short stolons.*

- iv. St. and ls. rough—Pan. spread.; spikelets ov. of three fls.—Paleæ webbed at base.

ORD. XCVII.—GRAMINEÆ—*cont.*

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
v. <i>P. nemoralis</i> , <i>L.</i> . . .	<i>Wood M.</i>	Woods .	..	1-3 ft.	P.	7-8
<i>a.</i> <i>nemoralis</i>	. . . . .	. . . .	..	..	..	..
<i>b.</i> <i>Pamelii</i> , <i>Bab.</i>	. . . . .	. . . .	..	..	..	..
<i>c.</i> <i>glauca</i> , <i>Sm.</i>	<i>glaucous M.</i> . .	Mount. .	..	..	..	..
vi. <i>P. laxa</i> , <i>Hacuke</i> . . .	<i>wavy M.</i>	Mount.!. .	..	6-8 in.	P.	7-8
<i>a.</i> <i>laxa</i>	. . . . .	. . . .	..	..	..	..
<i>b.</i> <i>minor</i> , <i>Gaud.</i>	. . . . .	. . . .	..	..	..	..
vii. <i>P. alpina</i> , <i>L.</i>	<i>Alpine M.</i> . . .	Mount.!. .	..	6-8 in.	P.	7-8
viii. <i>P. bulbosa</i> , <i>L.</i>	<i>bulbous M.</i> . . .	San. pl.!. .	..	6-8 in.	P.	4-5
Gen. XXXV. Briza , <i>L.</i>	<i>Quaking-grass.</i>					
<i>i.</i> <i>B. media</i> , <i>L.</i>	<i>common Q.</i> . . .	Downs .	..	6-12 in.	P.	6-7
<i>ii.</i> <i>B. minor</i> , <i>L.</i>	<i>least Q.</i>	Fields! S. of E.	..	6-12 in.	A.	7-8
Gen. XXXVI. Cynosurus , <i>L.</i>	<i>Dogs-tail-grass.</i>					
<i>i.</i> <i>C. cristatus</i> , <i>L.</i>	<i>common D.</i> . . .	Past. . .	..	1-2 ft.	P.	7-8
<i>ii.</i> <i>C. echinatus</i> , <i>L.</i>	<i>rough D.</i>	Jersey! .	..	6-10 in.	A.	7-8
Gen. XXXVII. Dactylis , <i>L.</i>	<i>Cock's-foot-grass.</i>					
<i>i.</i> <i>D. glomerata</i> , <i>L.</i> . .	<i>rough C.</i>	Fields .	..	1-2 ft.	P.	6-8
Gen. XXXVIII. Festuca , <i>L.</i>	<i>Fescue-grass.</i>					
* <i>A.</i>						
<i>i.</i> <i>F. uniglumis</i> , <i>Sol.</i> . .	<i>single husked F.</i> .	San. sh. .	..	8-12 in.	A. or B.	6-7
<i>ii.</i> <i>F. Myurus</i> , <i>L.</i>	<i>Wall F.</i>	Dry pl. .	..	6-8 in.	A.	6-7
<i>a.</i> <i>sciuroides</i> , <i>Roth.</i> . . .	. . . . .	. . . .	..	..	..	..
<i>b.</i> <i>pseudo-myurus</i> , <i>Soy.</i> .	. . . . .	. . . .	..	..	..	..
<i>B.</i>						
<i>iii.</i> <i>F. ovina</i> , <i>L.</i>	<i>Sheep's F.</i>	Dry past.	..	6-8 in.	P.	6-7
<i>a.</i> <i>ovina</i>	. . . . .	. . . .	..	..	..	..
<i>v.</i> <i>2. tenuifolia</i>	. . . . .	. . . .	..	..	..	..
<i>v.</i> <i>3. vivipara</i>	. . . . .	. . . .	..	..	..	..
<i>b.</i> <i>duriuscula</i> , <i>L.</i>	<i>hard F.</i>	. . . .	..	..	..	..
<i>c.</i> <i>rubra</i> , <i>L.</i>	<i>creeping F.</i> . . .	. . . .	..	..	..	..
* * <i>A.</i>						
iv. <i>F. sylvatica</i> , <i>Vill.</i> . .	<i>Reed F.</i>	Moi. woods	..	2-3 ft.	P.	7-8

CHARACTERISTICS OF SPECIES AND VARIETIES.

- v. St. and ls. smooth—Pan. spread.; branches slender, wavy—Spikelets lanc. of three fls.—Paleæ three-ribbed, webbed at base.
 a. Upper ls. as long as sheath—Ligule short—Paleæ webbed.
 b. Upper ls. shorter than sheath—Ligule very short—Paleæ slightly webbed.
 c. Ls. glauc.—Upper leaf as long as sheath—Ligule very short—Paleæ slightly webbed.

(*P. casia*, *Sm.* is a doubtful plant.)

- vi. St. smooth, slightly compressed—Pan. contracted, a little drooping—Spikelets ov., 3-4 fld.
 a. Ls. flat—Paleæ with hairs at base.
 b. Ls. keeled—Glu. nerved—Paleæ webbed.
 vii. St. glab., terete—Ls. broad, firm, obt.—Pan. loose; spikelets erect, 4-5 fld.—Ligules long, pointed.
 viii. St. terete, swollen at base—Ls. finely serr.—Pan. close; spikelets ov., 3-4 fld. paleæ webbed at base.

Gen. XXXV.

- i. Spikelets ov., about five-fld.—Ligule short, trunc.
 ii. Spikelets triang., about seven-fld.—Ligule long, acute.

Gen. XXXVI.

- i. Spike simp., elong., unilat.—Glu. with a short awn.
 ii. Spike ov., dense—Glu. with awn as long as paleæ.

Gen. XXXVII.

- i. Pan. branched; branches distinct, distant; with crowded, unilat., clusters of spikelets.

Gen. XXXVIII.

* *Radical ls. narrow.*

A. Root annual—Awns longer than the fls.

- i. Pan. upright, simp., unilat.—Fls. compressed, awned; lower glu. obsolete.
 ii. Pan. upright, unilat., branched at base only—Fls. terete, awned; lower glu. distinct.

a. St. leafless above—Pan. erecto-patent.

b. St. leafy to near the pan.—Pan. drooping at end.

B. Root perennial—Awns shorter than fls.

iii. Glau.—Pan. sub-unilat.—Spikelets obl., purplish.

a. St. tufted, four angled, rough below the pan.—Ls. all setac.

v. 2. *Ls. very slend.—Glu. not awned.*

v. 3. *Fls. viviparous:*

b. Less tufted—St. many angled, glab. below the pan.—St.-ls. flat—Glu. narrow.

c. Creeping—St. many angled, glab. below the pan.—St.-ls. flat, afterwards invol.—Glu. broad.

* * *Radical ls. broader than the st.-ls.*

A. Ligules of upper sheath long.

- iv. Pan. spread., erect, repeatedly branched—Fls. 2-5 ac., awnless, scab.

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
<i>B.</i>						
v. <i>F. elatior</i> , <i>Auct.</i> . . .	<i>tall F.</i>	Moi. past.	..	3-5 ft.	P.	7-8
v. 2. <i>arundinacea</i> . . .	..	..	..	..	..	..
vi. <i>F. pratensis</i> , <i>Huds.</i> .	<i>Meadow F.</i> . . .	Mead..	..	1-2 ft.	P.	6-7
v. 2. <i>lohiacea</i> , <i>Huds.</i> . .	<i>spiked F.</i>	..	..	..	..	..
vii. <i>F. gigantea</i> , <i>L.</i> . . .	<i>giant F.</i>	Sh. pl.	..	3-4 ft.	P.	7-8
Gen. XXXIX. <i>Bromus</i>, <i>L.</i>	<i>Brome-grass.</i>					
<i>*</i>						
<i>A.</i>						
i. <i>B. erectus</i> , <i>Huds.</i> . .	<i>upright B.</i> . . .	San. fie. .	..	2-3 ft.	P.	7-8
ii. <i>B. asper</i> , <i>L.</i>	<i>hairy B.</i>	Woods .	..	4-6 ft.	A.	7-8
<i>B.</i>						
iii. <i>B. sterilis</i> , <i>L.</i> . . .	<i>barren B.</i>	Fields .	..	1-2 ft.	A.	7-8
iv. <i>B. madritensis</i> , <i>L.</i> . .	<i>upright B.</i> . . .	Dry pl.!	..	1-2 ft.	A.	6-7
v. <i>B. maximus</i> , <i>Desf.</i> . .	<i>great B.</i>	Jersey .	..	1-2 ft.	A.	6-7
<i>* *</i>						
vi. <i>B. secalinus</i> , <i>L.</i> . . .	<i>Rye B.</i>	Fields .	..	2-3 ft.	A.	7-9
v. 2. <i>velutinus</i> , <i>Sm.</i> . . .	..	..	..	..	..	..
vii. <i>B. racemosus</i>	<i>smooth B.</i>	Past. . .	..	2-3 ft.	A.	7-9
v. 2. <i>commutatus</i> , <i>Schrad.</i>	..	..	..	..	..	..
viii. <i>B. mollis</i> , <i>L.</i>	<i>soft B.</i>	Waysides	..	1-2 ft.	A.	5-6
[ix. <i>B. arvensis</i> , <i>L.</i> . . .	<i>Field B.</i>	Sea co. .	..	2-3 ft.	A.	7-8
<i>* * *</i>						
[x. <i>B. squarrosus</i> , <i>L.</i> . .	<i>Corn B.</i>	Corn fie.!	..	1-2 ft.	A.	6-7
Gen. XL. <i>Brachypodium</i>, <i>Beauv.</i>	<i>False Brome- grass.</i>					
i. <i>B. sylvaticum</i> , <i>Beauv.</i>	<i>slender F.</i>	Woods .	..	1-2 ft.	P.	6-7
ii. <i>B. pinnatum</i> , <i>Beauv.</i> ..	<i>Heath F.</i>	Heaths .	..	1-2 ft.	P.	7-8
Gen. XLI. <i>Triticum</i>, <i>L.</i>	<i>Wheat-grass, &c.</i>					
i. <i>T. caninum</i> , <i>Huds.</i> . .	<i>bearded W.</i> . . .	Woods .	..	1-2 ft.	P.	6-7
v. 2. <i>biflorum</i> , <i>Mitten</i> . .	..	..	..	..	..	..
ii. <i>T. repens</i> , <i>L.</i>	<i>Couch-grass</i> . . .	Fields .	..	1-2 ft.	P.	6-7
v. 2. <i>littoralis</i>	..	..	..	..	..	..
iii. <i>T. junceum</i> , <i>L.</i> . . .	<i>rusky W.</i>	San. sh..	..	2-3 ft.	P.	7-8
<i>b. acutum</i> , <i>DC.</i>	..	..	..	..	..	..
<i>c. pungens</i> , <i>Pers.</i>	..	..	..	..	..	..
Gen. XLII. <i>Lolium</i>, <i>L.</i>	<i>Rye-grass.</i>					
i. <i>L. perenne</i> , <i>L.</i>	<i>Darnel, or R.</i> . .	Past. . .	..	1-2 ft.	P.	6-7

CHARACTERISTICS OF SPECIES AND VARIETIES.

B. Ligules of upper sheath very short.

- v. Pan. spread., drooping, much branched—Spikelets ov.-lanc. of 5-10 fls.—Awn very short—Ls. lin.-lanc.
- v. 2. *Branches of pan. divaricate after flowering.*
- vi. Pan. close, simp., sub-secund.—Spikelets lin., of 5-10 fls.—Awn very short.
- v. 2. *Spikelets sol., sess.—Awn. obsolete.*
- vii. Pan. drooping, spread., branched—Spikelets of 3-6 fls.—Awn long.

Gen. XXXIX.

* *Lower glu. one-nerved—Upper 3-5 nerved.**A. Awn short.*

- i. Pan. erect, slightly branched—Spikelets lin.-lanc.—Fls. many, distant, compressed.
- ii. Pan. drooping—Spikelets lin.-lanc.—Fls. cylind., remote—Lower ls. hairy.

B. Awn long.

- iii. Pan. drooping; pedunc. long, simp., ov., slightly branched—Spikelets lin.-lanc.—Ls. hairy.
- iv. Pan. erect; pedunc. short, simp., or slightly branched—Spikelets lin.-lanc.—Stam. usually two—Awn not longer than fls.
- v. Pan. erect, lax., ultimately drooping, slightly branched—Spikelets lanc., downy, long-stalked after flowering.

* * *Lower glu. 3-5 nerved—Upper 7-9 nerved.*

- vi. Pan. slightly branched, loose, spreading—Spikelets ov., compressed, about ten-flowered, glab.—Awns shorter than fls.
- v. 2. *Spikelets downy.*
- vii. Pan. simp., nearly erect—Spikelets ov.-obl., 6-10 flowered, smooth—Awns as long as fls.
- v. 2. *Stouter—Lower pedunc. often branched—Spikelets shorter.*
- viii. Pan. erect, close, slightly branched—Spikelets ov.; fls. closely imbric., hairy—Awn as long as fls.
- ix. Pan. spreading, drooping; lower pedunc. branched—Spikelets lanc., compr., pointed—Ls. and sheaths hairy—Fls. imbric. in fr., as long as awns.]
- * * * *Lower glu. nine-nerved.*
- x. Pan. simp., drooping—Spike ov.-obl., about twelve-flowered—Ls. downy—Awn spreading.]

Gen. XL.

- i. Root fibr.—Ls. broad—Pan. drooping—Awn as long as glu.
- ii. Root creeping—Ls. narrow—Pan. erect—Awn shorter than glu.

Gen. XLI.

- i. Root fibr.—Spikelets of about five-fls.—Glu. 3-5 ribbed, acu.—Awns long, rough.
- v. 2. *or T. alpinum (Don.); this is only T. repens.*
- ii. Root cr.—Spikelets of about five-fls.—Glu. acum., 5-7 ribbed—Awns short.
- v. 2. *Glauc.—Ls. invol.—Rachis smooth.*
- iii. Glauc.—Root cr.—Spikelets of 4-6 fls.—Glu. obt., 6-10 ribbed—Awns o.
- b. Ls. scabrous—Fls. 5-8; glu., mucro. ov. slightly awned.
- c. Ls. very scabr., pungent—Spikelets smaller, fls. 5-12; glu. acute.

Gen. XLII.

- i. Perenn.—Glu. shorter than the spikelet—Awns o.
(*L. italicum* is a cultivated form of this.)

ORD. XCVII.—GRAMINEÆ—*con*

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
ii. <i>L. temulentum</i> , <i>L.</i> . . .	<i>bearded D.</i> . . .	Past. . .	. .	1-2 ft.	P.	6-7
v. 2. <i>arvense</i> , <i>With.</i> . . .	. . . . .	. . . . .	. .	. .	. .	. .
Gen. XLIII. <i>Elymus</i> , <i>L.</i>	<i>Lyme-grass.</i>					
i. <i>E. arenarius</i> , <i>L.</i>	<i>Sea L.</i>	San. sh. .	. .	3-4 ft.	P.	7-8
Gen. XLIV. <i>Hordeum</i> , <i>L.</i>	<i>Barley.</i>					
* i. <i>H. sylvaticum</i> , <i>Huds.</i>	<i>Wood B.</i>	Woods .	. .	1-2 ft.	P.	7-8
* *						
ii. <i>H. pratense</i> , <i>Huds.</i> .	<i>Meadow B.</i>	Moi. past.	. .	1-2 ft.	P.	6-7
iii. <i>H. murinum</i> , <i>L.</i>	<i>Wall B.</i>	Dry pl. .	. .	6-12 in.	A.	6-7
iv. <i>H. maritimum</i> , <i>With.</i>	<i>Sea B.</i>	Sea co. .	. .	6-12 in.	A.	6-7
Gen. XLV. <i>Nardus</i> , <i>L.</i>	<i>Mat-grass.</i>					
j. <i>N. stricta</i> , <i>L.</i>	<i>common M.</i>	Moors. .	. .	8-10 in.	P.	6-7
Gen. XLVI. <i>Lepturus</i> ,	<i>Hard-grass.</i>					
<i>R. Brown.</i>						
i. <i>L. filiformis</i> , <i>Trin.</i> . .	<i>Sea H.</i>	Sea co. .	. .	4-8 in.	P.	7-9

ORD. XCVIII.—FILICES.

Gen. I. <i>Ceterach</i> , <i>Willd.</i>	<i>Ceterach.</i>					
i. <i>C. officinarum</i> , <i>Willd.</i>	<i>common C., or Scale-fern.</i>	Walls . .	. .	4-6 in.	P.	4-10
Gen. II. <i>Gymnogramma</i> ,	<i>Gymnogram.</i>					
<i>Desv.</i>						
i. <i>G. leptophylla</i> , <i>Desv.</i>	<i>slender G.</i>	Jersey! .	. .	1-3 in.	A.	3-5
Gen. III. <i>Woodsia</i> , <i>Br.</i>	<i>Woodsia.</i>					
i. <i>W. ilvensis</i> , <i>Br.</i>	<i>oblong W.</i>	Mount.! .	. .	3-6 in.	P.	7-8
ii. <i>W. hyperborea</i> , <i>Br.</i> .	<i>round-leaved W.</i>	Mount.! .	. .	3-6 in.	P.	7-8
Gen. IV. <i>Polypodium</i> , <i>L.</i>	<i>Polypody, &c.</i>					
i. <i>P. vulgare</i> , <i>L.</i>	<i>common P.</i>	Walls & banks.	. .	6-12 in.	P.	7-8
ii. <i>P. Phegopteris</i> , <i>L.</i> . . .	<i>Beech-fern</i>	Dmp. wds.	. .	6-9 in.	P.	6-8
iii. <i>P. Dryopteris</i> , <i>L.</i> . . .	<i>Oak-fern</i>	Dry woods	. .	6-12 in.	P.	6-8
iv. <i>P. calcareum</i> , <i>Sm.</i> . . .	<i>limestone P.</i>	Ch. woods	. .	6-12 in.	P.	6-8
Gen. V. <i>Allosorus</i> , <i>Bernh.</i>	<i>Rock-brake.</i>					
i. <i>A. crispus</i> , <i>Bernh.</i> . . .	<i>curled R., or Parsley-fern.</i>	Mount.! .	. .	3-8 in.	P.	7-8

ORD. XCVII.—GRAMINEÆ—*cont.*

CHARACTERISTICS OF SPECIES AND VARIETIES.

- ii. Ann.—Glu. longer than the spikelet—Awns long.
v. 2. Awn o., or short.

Gen. XLIII.

- i. Glauc.—Root cr.—Pan. erect, stout—Ls. rigid.

Gen. XLIV.

* *Lat. fls. perfect—Middle barren.*

- i. Glu. all awl-shaped ; awn of outer glu. twice as long as glu.
 * * *Lateral fls. imperfect—Middle perfect.*
 ii. Perenn.—Glu. all setaceous, rough.
 iii. Ann.—Glu. of the lat. fls. setaceous, rough ; of the middle one lin.-lanc., cil.
 iv. Ann.—Glu. scab. ; inner ones of the lat. fls. half-ov. ; the rest setaceous.

Gen. XLV.

- i. Spike erect, slender ; spikelets sess., unilat.—Ls. setaceous.

Gen. XLVI.

- i. Spike cylind., slender, erect—Glu. united at base.

ORD. XCVIII.—FILICES.

Gen. I.

- i. Fro. pinnfid.—pinnæ ov.-obl., covered beneath with chaffy scales.

Gen. II.

- i. Fro. bi-pinnfid. ; pinnæ obov., cun., bifid, or three-lobed ; lobes obt.—Root ann.

Gen. III.

- i. Fro. lanc., pinn. ; pinnæ obl., deeply pinnfid., with many obl., cren. lobes.
 ii. Fro. lanc., pinn. ; pinnæ ov., with few broad, ent. lobes ; less scaly.

Gen. IV.

- i. Fro. deeply pinnfid. ; pinnæ lanc., cren.
 ii. Fro. pinn. ; pinnæ pinnfid., the lowest deflexed ; seg. lin.-lanc., obt.
 iii. Fro. deltoid, tern., pinn., flaccid ; pinnæ pinnfid.
 iv. Fro. sub-tern, pinn., coriac., gland-pubesc. ; pinnæ pinnfid.

Gen. V.

- i. Barren fro. bi-tripinn. ; pinnules cun., often bifid—Fertile pinnules obl.

ORD. XCVIII.—FILICES—*cont.*

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
Gen. VI. Cystopteris, <i>Bern.</i>	<i>Bladder-fern.</i>					
i. <i>C. fragilis, Bernh.</i> . . .	<i>brittle B.</i>	Rocks & walls.	..	4-8 in.	P.	7-8
<i>a. fragilis</i>	.....	..	..	..	..	..
<i>b. dentata, Hook.</i>	.....	..	..	..	..	..
<i>c. Dichieana, Hook.</i> . . .	.....	..	..	..	..	..
ii. <i>C. montana, Link.</i> . . .	<i>Mountain B.</i> . . .	Sc.mount.!	..	4-6 in.	P.	8-9
Gen. VII. Polystichum, <i>Roth.</i>	<i>Shield-fern.</i>					
i. <i>P. Lonchitis, Roth.</i> . .	<i>Holly-fern</i>	Alp.rocks!	..	6-18 in.	P.	6-8
ii. <i>P. aculeatum, Roth.</i> . .	<i>prickly S.</i>	Woods .	..	6-18 in.	P.	6-8
<i>a. aculeatum</i>	.....	..	..	..	..	..
<i>b. lobatum, Sw.</i>	.....	..	..	..	..	..
<i>c. angulare, Willd.</i>	.....	..	..	..	..	..
Gen. VIII. Lastræa, Presl.	<i>Buckler-fern, &c.</i>					
i. <i>L. Thelypteris, Presl.</i>	<i>Marsh-fern</i>	Marshes!	..	1-2 ft.	P.	7-8
ii. <i>L. Oreopteris, Presl.</i> .	<i>Mountain F.</i> . . .	Mount. .	..	1-2 ft.	P.	7-8
iii. <i>L. Filix-mas, Presl.</i> .	<i>Male-fern</i>	Sh. pl. .	..	1-3 ft.	P.	7-8
<i>b. incisum, Newm.</i>	.....	..	..	..	..	..
<i>c. Borreri, Newm.</i>	.....	..	..	..	..	..
<i>d. abbreviata, Newm.</i> . . .	.....	..	..	..	..	..
iv. <i>L. rigida, Presl.</i>	<i>rigid B.</i>	Mount.!	..	1-2 ft.	P.	7-8
v. <i>L. cristata, Presl.</i> . . .	<i>crested B.</i>	Boggy heaths.!	..	1-1½ ft.	P.	7-8
<i>b. uliginosa, Newm.</i>	.....	..	..	..	..	..
vi. <i>L. spinulosa, Presl.</i> . .	<i>spinulose B.</i>	Dmp. wds.	..	12-18 in.	P.	7-8
vii. <i>L. dilatata, Presl.</i> . .	.....	..	..	..	..	..
<i>b. glandulosa, Newm.</i> . . .	.....	..	..	..	..	..
<i>c. collina, Newm.</i>	.....	..	..	..	..	..
<i>d. nana, Newm.</i>	.....	..	..	..	..	..
viii. <i>L. æmula, Brach.</i> . . .	.....	Mount.!	..	12-18 in.	P.	7-9
Gen. IX. Pseudathyrium, <i>Newm.</i>						
i. <i>P. alpestre, Newm.</i> . . .	.....	Sc.mount.!	..	6-12 in.	P.	7-8
v. 2. <i>flexile, Newm.</i>	.....	..	..	..	..	..
Gen. X. Athyrium, Roth.	<i>Lady-fern.</i>					
i. <i>A. filix-fæmina, Bernh.</i>	<i>Lady-fern</i>	Moi. pl. .	..	1-5 ft.	P.	7-8

ORD. XCVIII.—FILICES—*cont.*

CHARACTERISTICS OF SPECIES AND VARIETIES.

Gen. VI.

- i. Fro. ov.-lanc., bipinn.; pinnules ov., ac. lobed or toothed.
- a. Sori. central.
- b. Pinnules obt., unequally toothed—Sori. marginal.
- c. Pinnules broad, overlapping—Sori. marginal.
- ii. Fro. triang., tripinn., on long stalks—Lower pinnæ the largest.

Gen. VII.

- i. Fro. lin.-lanc., pinn., rigid; pinnæ lanc., falc., serr.; lower auricled.
- ii. Fro. lanc. bi-tripinn.; pinnules ov., sub-falc., auricled.
- a. Fro. lanc., rigid—Pinnules obliquely decurr.
- b. Fro. narrower, rigid—Pinnules decurr., sess., less divided.
(*Louchitioides* is a starved form of this.)
- c. Fro. lax—Pinnules thin, membran., distinctly stalked, teeth awned.

Gen. VIII.

- i. Fro. pinn.; pinnæ pinnfid. not glandular; lower pinnæ not shorter than the upper—Rt. creeping.
- ii. Fro. pinn.; pinnæ pinnfid., glandular; sweet-scented; gradually decreasing from the middle of fro.—Rt. tufted.
- iii. Fro. ov.-lanc., pinn.—Pinnæ pinnfid., lower ones rather smaller—Pinnules obl., blunt, serr.
- b. Pinnules less crowded, obl.-lanc., deeply serr.
- c. St. covered with golden yellow scales—Pinnules almost trunc., less serr.
- d. Small—Fro. pinn.; pinnules very blunt, crowded.
- iv. Fro. obl.-lanc., bi-pinn., gland.; pinnules obl., blunt serr., not spinulose—Stalk covered with chaffy scales.
- v. Fro. lin.-obl., erect, pinn.; pinnæ, short, distant, deeply pinnfid.; pinnules obl.-ov., acutely serr., but not spinulose.
- b. Pinnules more divided, teeth slightly spinulose.
- vi. Fro. pale green, lanc., pinn.; pinnæ pinnfid.; pinnules acute, serr., spinulose—Scales on st. ov., few, uniformly coloured.
- vii. Fro. dark green, ov.-triang., bipinn.; pinnules, serr., spinulose—Scales on st. lanc., acum., centre dark brown.
- b. Fro. broader, more gland. beneath.
- c. Fro. obl.-lanc.—Pinnules ov., blunt, sub-entire, less spinulose.
- d. Small—Fro. ov.-triang.—Scales narrow, pale.
- viii. Fro. ov.-triang., bi-pinn.; pinnules obl.-ov., lobed, their margins recurved—Scales of st. many, small, narrow, jagged, pale.

Gen. IX.

- i. Fro. obl.-lanc., bipinn.; pinnules crowded, ov.-obl., acute, lobed or pinnfid.: sharply toothed.
- v. 2. Stem short—Pinnæ short, ov.-lanc.; pinnules distant.

Gen. X.

- i. Fro. obl.-lanc., flaccid, bi-pinn.—Pinnules lin. obl., deeply serr., lobed or pinnfid.; sharply toothed.

ORD. XCVIII.—FILICES—*cont.*

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
<i>a. rhæticum</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
<i>b. latifolium</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
<i>c. incisum</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
Gen. XI. Asplenium, L.	<i>Spleenwort.</i>					
i. <i>A. viride, Huds.</i>	<i>green S.</i>	Rocks .	. . . . .	6-12 in.	P.	6-9
ii. <i>A. Trichomanes, L.</i> . .	<i>common S., or Maidenhair S.</i>	Walls & rocks.	. . . . .	6-12 in.	P.	6-9
v. 2. <i>anceps</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
iii. <i>A. marinum, L.</i>	<i>Sea S.</i>	Sea cliffs	. . . . .	3-10 in.	P.	6-10
iv. <i>A. lanceolatum, Huds.</i>	<i>lanceolate S.</i>	Wet rocks!	. . . . .	6-9 in.	P.	6-9
v. <i>A. Adiantum-nigrum, L.</i>	<i>black S.</i>	Banks & walls.	. . . . .	6-12 in.	P.	6-10
v. 2. <i>acutum, Bory.</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
vi. <i>A. ruta-muraria, L.</i> . .	<i>Wall-rue</i>	Walls, &c.	. . . . .	1-2 in.	P.	6-10
vii. <i>A. septentrionale, Hull.</i>	<i>forked S.</i>	Mount. rocks!	. . . . .	1-2 in.	P.	6-10
viii. <i>A. germanicum, Weiss.</i>	<i>alternate-leaved S.</i>	Rocks! .	. . . . .	2-3 in.	P.	6-10
Gen. XII. Scolopendrium, Sw.	<i>Hart's-tongue.</i>					
i. <i>S. vulgare, Sym.</i>	<i>common H.</i>	Sh. pl. .	. . . . .	1-2 ft.	P.	7-8
Gen. XIII. Blechnum, L.	<i>Hard-fern.</i>					
i. <i>B. boreale, Sw.</i>	<i>northern H.</i>	Woods .	. . . . .	6-9 in.	P.	7-8
Gen. XIV. Pteris, L.	<i>Brake.</i>					
i. <i>P. aquilina, L.</i>	<i>common B.</i>	San. pl. .	. . . . .	2-6 ft.	P.	7-8
Gen. XV. Adiantum, L.	<i>Maiden-hair.</i>					
i. <i>A. Capillus-Veneris, L.</i>	<i>true M.</i>	Rocks! .	. . . . .	4-12 in.	P.	7-8
Gen. XVI. Trichomanes, L.	<i>Bristle-fern.</i>					
i. <i>T. radicans, Sw.</i>	<i>rooting B.</i>	Wet rocks! I.	. . . . .	4-10 in.	P.	6-9
v. 2. <i>Andrewsii.</i>	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .	. . . . .
Gen. XVII. Hymenophyllum, Sm.	<i>Filmy-fern.</i>					
i. <i>H. Tunbridgense, Sm.</i>	<i>Tunbridge F.</i>	Wet rocks!	. . . . .	1-3 in.	P.	6-7
ii. <i>H. Wilsonii, Hook.</i> . .	<i>Scottish F.</i>	Wet rocks!	. . . . .	1-3 in.	P.	6-7
Gen. XVIII. Osmunda, L.	<i>Osmund.</i>					
i. <i>O. regalis, L.</i>	<i>royal-fern</i>	Bogs . .	. . . . .	2-5 ft.	P.	6-8
Gen. XIX. Botrychium, Sw.	<i>Moon-wort.</i>					
i. <i>B. lunaria, Sw.</i>	<i>common M.</i>	Dry past.! .	. . . . .	3-6 in.	P.	6-8

ORD. XCVIII.—FILICES—*cont.*

CHARACTERISTICS OF SPECIES AND VARIETIES.

- a.* Fro. narrow, lanc.—Pinnules narrow, distinct, convex.
- b.* Fro. narrow, lanc.—St. short—Pinnules ov., crowded, irregularly lobed.
- c.* Fro. very large, dark green—Pinnules obl.-lanc., flat, deeply pinnfid.

Gen. XI.

- i.* Fro. lin., pinn., their stalk green—Pinnæ roundish-ov., obtusely serr.
- ii.* Fro. lin., pinn., their stalk black—Pinnæ roundish-obl., cren.
- v.* 2. *Larger.*
- iii.* Fro. obl.-lanc., pinn., coriac.; pinnæ obl., serr.—Rachis winged.
- iv.* Fro. broadly lanc., bipinn.—Pinnules obov., acu., serr.—Rachis scaly, not winged.
- v.* Fro. triang., bi-tri-pinn.; pinnules cun., deeply serr. or toothed.
- v.* 2. *Fro. tripinn., very acum.—Pinnules lin., acu.*
- vi.* Fro. ov., bi-pinn. below; pinnæ cuneate, lobed at apex.
- vii.* Fro. lin.-lanc., 2-3 cleft—Seg. alt., lanc., toothed.
- viii.* Fro. lin.-pinn.—Pinn. alt., narrow cuneate, toothed above.

Gen. XII.

- i.* Fro. ent., strap-shaped; base cord.

Gen. XIII.

- i.* Fert. fro. pinn.; pinnæ lin., acu.—Barren fro. pinnfid.; pinnæ lanc., obt.

Gen. XIV.

- i.* Fro. tripart.—Branches bi-pinn.; pinnules lanc., blunt, the lower pinnfid.

Gen. XV.

- i.* Fro. bi-pinn.—Pinnules round, wedge-shaped, lobed, stalked.

Gen. XVI.

- i.* Fro. triang., glab., 3-4 pinnfid.—Seg. lin. entire, membran.
- v.* 2. *Fro. lanc.—Invol. winged.*

Gen. XVII.

- i.* Fro. pinn.—Pinnæ spreading—Invol. serr.
- ii.* Fro. pinn.—Pinnæ deflexed—Invol. entire.

Gen. XVIII.

- i.* Fro. bi-pinn.; pinnæ obl.-ov., serr., lower auricled.

Gen. XIX.

- i.* Fro. sol., pinn.—Pinnæ fan-shaped, cren.

ORD. XCVIII.—FILICES—*cont.*

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
Gen. XX. Ophioglossum, <i>L.</i>	<i>Adder's-tongue.</i>					
i. <i>O. vulgatum, L.</i> . . .	<i>common A.</i> . . .	Moi. past.	..	2-8 in.	P.	6-8
v. 2. <i>lusitanicum, L.</i> . . .	. . . . .	. . . . .	..	..	..	..

ORD. XCIX.—LYCOPODIACEÆ.

Gen. I. Lycopodium, L.	<i>Club-moss.</i>					
i. <i>L. clavatum, L.</i> . . .	<i>common C.</i> . . .	Heaths .	..	1-3 ft.	P.	7-8
ii. <i>L. annotinum, L.</i> . . .	<i>interrupted C.</i> . . .	Sc.mount.!	..	6-18 in.	P.	7-8
iii. <i>L. inundatum, L.</i> . . .	<i>Marsh C.</i>	Moi. hths.	..	2-6 in.	P.	6-8
iv. <i>L. alpinum, L.</i>	<i>Alpine C.</i>	Mount.!	..	6-18 in.	P.	6-8
v. <i>L. Selago, L.</i>	<i>Fir C.</i>	Heaths .	..	2-6 in.	P.	6-8
Gen. II. Selaginella, <i>Beauv.</i>						
i. <i>S. selaginoides, L.</i> . . .	. . . . .	Sc. bogs.	..	2-6 in.	P.	6-8
Gen. III. Isoetes, L.	<i>Quill-wort.</i>					
i. <i>I. lacustris, L.</i>	<i>European Q.</i> . . .	Bottoms of lakes!	..	2-6 in.	P.	5-7
<i>a. lacustris, L.</i>	. . . . .	. . . . .	..	..	..	..
<i>b. echinospora, Dur.</i> . . .	. . . . .	. . . . .	..	..	..	..
ii. <i>I. Hystrix, Dur.</i>	. . . . .	Guernsey!	..	1-2 in.	P.	5-6

ORD. C.—MARSILEACEÆ.

Gen. I. Pilularia, L.	<i>Pill-wort.</i>					
i. <i>P. globulifera, L.</i> . . .	<i>creeping P.</i>	Lake shores!	..	2-4 in.	P.	5-7

ORD. CI.—EQUISETACEÆ.

Gen. I. Equisetum, L. *	<i>Horse-tail.</i>					
i. <i>E. maximum, Lam.</i> . .	<i>great Water H.</i> . .	Bogs . .	..	3-6 ft.	P.	4-5
ii. <i>E. umbrosum, Willd.</i>	<i>blunt-topped H.</i> . .	Bogs! . .	..	1-2 ft.	P.	4-5
iii. <i>E. arvense, L.</i>	<i>Corn H.</i>	Fie. . . .	..	3-8 in.	P.	4-5

ORD. XCVIII.—FILICES—*cont.*

CHARACTERISTICS OF SPECIES AND VARIETIES.

Gen. XX.

- i. Fro. ov., ent., obt.—Spike club-shaped.
 v. 2. *Fro. lin.-lanc., fleshy (Guernsey).*

ORD. XCIX.—LYCOPODIACEÆ.

Gen. I.

- i. St. long, cr.—Ls. scattered, incurved, hair pointed—Spikes (2-3) stalked.
 ii. St. long, cr.—Ls. scattered, lanc., mucro., serr.—Spike sol., sess.
 iii. St. short, cr.—Ls. scattered, lin., acu., curved upwards—Spike sol., sess.
 iv. St. long, cr.—Ls. in four rows, obl., keeled, acu., adpr.—Spikes sol., sess.
 v. St. erect, forked, fastigiate—Ls. in eight rows, lin.-lanc., acu., imbric.—Caps not spiked, but in the axils of the common ls.

Gen. II.

- i. St. cr., slender—Ls. lanc., cili.—Spikes on short erect branches.

Gen. III.

- i. Aquatic—Ls. subu., bluntly quadrang.
 a. Ls. erect, green—Caps. half covered by the leaf-sheath.
 b. Ls. spreading, pale—Caps. almost covered by the leaf-sheath.
 ii. Terrestrial—Ls. filiform.

ORD. C.—MARSILEACEÆ.

Gen. I.

- i. Rhizome slend., cr.—Ls. fili., erect.

ORD. CI.—EQUISETACEÆ.

Gen. I.

* *Fert. sts. simp., succulent, appearing before the whorled, barren stem.*

- i. Barren st. with about thirty striæ, smooth—Branches erect; teeth of sheaths two-ribbed.
 ii. Barren st. with about twenty striæ, rough—Branches spreading; teeth of sheaths blunt, with one rib, not reaching to tip—Sheaths of fert. sts. adpr.
 iii. Barren st. with about 10-15 striæ, slightly rough—Branches spreading; teeth of sheaths acu., with one rib reaching to tip—Sheaths of fert. sts. lax.

ORD. CL.—EQUISETACEÆ—*cont.*

Botanical Name.	Common Name.	Soil or Situation.	Colour.	Growth.	Dura- tion.	Time of flowering.
* *						
<i>A.</i>						
iv. <i>E. sylvaticum</i> , <i>L.</i> . . .	<i>Wood H.</i>	Damp woods	. . .	1-2 ft.	P.	4-5
<i>B.</i>						
v. <i>E. palustre</i> , <i>L.</i>	<i>Marsh H.</i>	Bogs . .	. . .	6-18 in.	P.	6-7
v. 2. <i>alpinum</i>	. . . . .	. . . . .	. . .	. . .	. . .	. . .
v. 3. <i>polystachion</i>	. . . . .	. . . . .	. . .	. . .	. . .	. . .
vi. <i>E. limosum</i> , <i>L.</i>	<i>smooth naked H.</i>	Bogs . .	. . .	1-3 ft.	P.	6-7
<i>a. limosum</i>	. . . . .	. . . . .	. . .	. . .	. . .	. . .
<i>b. fluviatile</i>	. . . . .	. . . . .	. . .	. . .	. . .	. . .
vii. <i>E. hyemale</i> , <i>L.</i>	<i>rough H.</i>	Boggy woods.	. . .	1-3 ft.	P.	7-8
viii. <i>E. trachyodon</i> , <i>Braun.</i>	<i>long-stemmed H.</i>	. . . . .	. . .	. . .	. . .	. . .
v. 2. <i>Moorei</i> , <i>Newm.</i>	. . . . .	. . . . .	. . .	. . .	A.	. . .
ix. <i>E. variegatum</i> , <i>Sch.</i> . .	<i>variegated H.</i> . .	Wet pl. .	. . .	1-2 ft.	P.	7-8
<i>b. Wilsoni</i> , <i>Newm.</i>	. . . . .	. . . . .	. . .	. . .	. . .	. . .
<i>c. arenarium</i>	. . . . .	San. sh. .	. . .	. . .	. . .	. . .

ORD. CI.—EQUISETACEÆ—*cont.*

CHARACTERISTICS OF SPECIES AND VARIETIES.

* * *Fertile and barren stems similar.*

A. Branches compound.

iv. Sts. with many whorls of compound deflexed branches.

B. Branches simple or none.

v. Sts. with 4-10 deep furrows—Sheaths loose, their teeth wedge-shaped, acu., with membran. tips.

v. 2. *A stunted form—St. almost simple.*

v. 3. *Upper branches all fertile.*

vi. St. smooth, with 15-20 faint striæ—Branches often none, or irregular—Sheaths short, adpr.; teeth short, rigid.

a. Smooth—Branches short or o., rigid, as long as the internodes.

b. Slightly scab. above—Branches longer than the internodes.

vii. St. rough, with 15-20 distinct striæ, glauc.—Branches o., or all rad.; sheaths close, white, with black base and tip; teeth black, deciduous.

viii. St. rough, with 8-12 distinct striæ—Branches rad., long, flexuous; sheaths close, black; teeth slender, persist., white.

v. 2. *St. ann.—Sheaths loose.*

ix. St. rough, with 4-10 grooves, fili., simp., or with rad. branches—Sheaths green below, black above, with obt., bristle-pointed teeth.

δ. Tall (2-3 ft.) erect—Teeth of sheaths short, obt.

c. St. decumb., slend.—Teeth of sheaths cuneate.

APPENDIX.

EXCLUDED SPECIES.

RANUNCULACEÆ.

- Anemone ranunculoides**, *L.* Sep. 5, yellow—Ls. ternate or quinate.—*King's Langley, Herts, and Wrotham, Kent.*
Pæonia corallina, *Retz.* Ls. biternate—Fls. large, crimson.—*Steep Holmes Island, in the Severn.—Introduced.*

BERBERIDACEÆ.

- Epimedium alpinum**, *L.* Leaves all cauline, biternate—Fls. red.—*Sub-alpine woods, &c.—An escape.*

CARYOPHYLLACEÆ.

- Silene italica**, *Pers.* Resembles *S. nutans*, but has a long, clavate calyx, and the seg. of the corolla broader and not crowned.—*Between Darenth and Dartford, Kent.*
Cucubalus bacciferus, *L.* Ls. ovate, acute—Fruit fleshy.—*Isle of Dogs.—Introduced.*

HYPERICACEÆ.

- Hypericum hircinum**, *L.*, and **H. elatum**, *L.* Both introduced in shrubberies, with no claim to be indigenous.

MALVACEÆ.

- Malva verticillata**, *L.* St. erect—Outer sep. lin.—Pet. not longer than sep.—*Near Llanelly, South Wales.*

GERANIACEÆ.

- Geranium nodosum**, *L.*, and **G. striatum**, *L.* Ls. 3-5 lobed; the stem of the former glabrous, of the latter hairy.—*Escapes from gardens in various places.*

BALSAMINACEÆ.

- Impatiens parviflora**, *DC.* A weed resembling the other species of *Impatiens*, but with small flowers.—*Naturalized in many places.*

CELASTRACEÆ.

- Staphyllea pinnata**, *L.* Ls. pinnate—Capsule bladdery—Fls. yellowish white.—*Shrubberies.—An escape.*

LEGUMINOSÆ.

- Trifolium resupinatum**, *L.* Flowers resupinate.—*A casual*.—Near Liverpool, &c.
Lathyrus latifolius, *L.* The common everlasting pea of gardens; an escape in several parts.

ROSACEÆ.

- Several varieties of *Rosa* are found wild in various parts, which have escaped from gardens.

UMBELLIFERÆ.

- Angelica archangelica**, *L.* Ls. very large; lfts. ov.-lanc.—Fls. greenish white.—Watery places.—An escape.

RUBIACEÆ, *L.*

- Asperula taurina**, *L.* Ls. four in a whorl, elliptical, three-veined.—Cadeby, Leicestershire; Casterton Woods, Westmoreland.

VALERIANACEÆ.

- Centranthus Calcitrapa**, *Dufr.* Rad. ls. ov., ent.; st. ls. pinnfid—Spur short.—Eltham.

DIPSACEÆ.

- Dipsacus fullonum**, *Mill.* Scales of the involucre hooked at the end.—*W. of E.*—An escape.

COMPOSITÆ.

- Petasites fragrans**, *Presl.* Outer florets ligulate.—Shrubberies in many places.—Naturalized.

- Anthemis tinctoria**, *L.* Receptacle hemispherical—Rays and disk yellow.—Ballast heaps.—An escape.

- Anacyclus radiatus**, *Lois.* Ls. woolly, bi-pinnfid.—Ray yellow.—Dunboy Quay, Bearhaven, Cork.

- Achillea tanacetifolia**, *All.* Ls. bi-pinnfid., with a lanc. outline—Rachis winged, serrate.—Ringing Low, Sheffield; Cromford Moor, Derbyshire.

- Achillea decolorans**, *Sch.* Ls. lin.-lanc., downy, opaque—Ray buff coloured.—Near Matlock.

- Achillea tomentosa**, *L.* Ls. woolly, pinnfid.—Corymbs much branched—Rays golden yellow.—Many places in Scotland (possibly native).

- Xanthium strumarium**, *L.* Lower ls. heart-shaped, three lobed at base—Involucre prickly.—Waste places.—*A casual*.

CONVOLVULACEÆ

- Cuscuta hassiaca**, *Pfeiff.* Flowers stalked, scented like Heliotrope.—*A casual* on Lucerne, in Essex.

SOLANACEÆ.

- Datura stramonium**, *L.* Ls. ovate, unequally serr.—Flowers large, white.—*A casual* on dung-hills, &c.

- Lycium barbatum**, *L.* An escape in many places from cottage gardens.

SCHROPHULARIACEÆ.

Linaria supina, *Desf.* Ls. lin., blunt, whorled—Fls. capitate, yellow.—*Devonshire, Cornwall, &c.*

Linaria purpurea, *L.* Ls. lin.-lanc., scattered—Spur long—Fls. racemose, yellow or purple.—*Old walls.—An escape.*

LABIATÆ.

Salvia clandestina, *L.* Upper ls. obl.-acute—Tube of corolla longer than calyx.—*Cornwall and Jersey.*

Stachys annua, *L.* Lower ls. ovate, oblong—Floral ls. lanc.—Fls. yellowish—*Cornfields near Gadskill, Kent.*

PRIMULACEÆ.

Lysimachia ciliata, *L.* St. erect—Ls. ov.-lanc.—Fls. in axill. rac.—Sta. ten; five of them sterile.—*Naturalized in several places.*

Lysimachia punctata, *L.* Resembles *L. vulgaris*, but has the flowers solitary and the segments of the corolla ciliate.—*Dulverton, Devon.*

POLYGONACEÆ.

Polygonum Fagopyrum, *L.* St. erect—Ls. sagittate—Fls. pale reddish, in spreading panicles.—*Dung-hills, &c.*

CHENOPODIACEÆ.

Atriplex nitens, *L.* St. erect—Ls. triang.; glauc. beneath.—*Sea-shore near Ryde, I. of W.—Introduced.*

CUPULIFERÆ.

Castanea vulgaris, *Lamk.* *The Spanish chestnut.—Shrubberies, parks, &c.—Introduced.*

Many other species have been described by various botanists; they are either now extinct, errors, or of such a casual character as not to warrant their insertion in the list of British plants.

INDEX OF THE ORDERS AND GENERA.

ACER, 28
 Aceraceæ, 28
 Aceras, 138
 Achillea, 88, 189
 Aconitum, 6
 Acorus, 150
 Actæa, 6
 Actinocarpus, 146
 Adiantum, 182
 Adonis, 2
 Adoxa, 58
 Ægopodium, 62
 Æthusa, 64
 Agrimonia, 46
 Agrostis, 168
 Aira, 170
 Ajuga, 106
 Alchemilla, 48
 Alisma, 146
 Alismaceæ, 146
 Allium, 142
 Allosorus, 178
 Allysum, 12
 Alnus, 130
 Alopecurus, 166
 Althæa, 26
 Alyssum, 12
 Amaranthaceæ, 118
 Amaranthus, 118
 Amaryllidaceæ, 140
 Ammophila, 168
 Anacyclus, 189
 Anagallis, 114
 Anchusa, 112
 Andromeda, 90
 Anemone, 2, 188
 Angelica, 64, 189
 Antennaria, 82
 Anthemis, 88, 189
 Anthoxanthum, 166
 Anthriscus, 66
 Anthyllis, 32
 Antirrhinum, 100
 Apera, 168
 Apium, 60
 Apocynaceæ, 94

Aquifoliaceæ, 92
 Aquilegia, 4
 Arabis, 12
 Araceæ, 150
 Araliaceæ, 58
 Arbutus, 92
 Arctium, 80
 Arctostaphylos, 92
 Arenaria, 22
 Aristolochia, 126
 Aristolochiaceæ, 124
 Armeria, 116
 Arnoseris, 78
 Arrhenatherum, 170
 Artemisia, 82
 Arthrolobium, 36
 Arum, 150
 Arundo, 168
 Asarum, 124
 Asparagus, 144
 Asperugo, 112
 Asperula, 70, 189
 Asplenium, 182
 Aster, 84
 Astragalus, 36
 Astrantia, 60
 Athyrium, 180
 Atriplex, 120, 190
 Atropa, 96
 Avena, 170

Ballota, 108
 Balsaminaceæ, 30
 Barbarea, 12
 Barkhausia, 78
 Bartsia, 98
 Bellis, 86
 Berberaceæ, 6
 Berberis, 6
 Beta, 120
 Betula, 130
 Betulaceæ, 130
 Bidens, 82
 Blechnum, 182
 Blysmus, 156
 Boraginaceæ, 110

Borago, 112
 Botrychium, 182
 Brachypodium, 176
 Brassica, 14
 Briza, 174
 Bromus, 176
 Bryonia, 52
 Bunium, 62
 Bupleurum, 62
 Butomus, 146
 Buxus, 126

Cakile, 8
 Calamagrostis, 168
 Calamintha, 106
 Callitriche, 50
 Calluna, 90
 Caltha, 4
 Camelina, 12
 Campanula, 88
 Campanulaceæ, 88
 Caprifoliaceæ, 68
 Capsella, 10
 Cardamine, 12
 Carduus, 80
 Carex, 158
 Carlina, 82
 Carpinus, 128
 Carum, 62
 Caryophyllaceæ, 20
 Castanea, 190
 Catabrosa, 172
 Caulis, 66
 Celastraceæ, 30
 Centaurea, 82
 Centranthus, 70, 189
 Centunculus, 116
 Cephalanthera, 136
 Cerastium, 24
 Ceratophyllum, 52
 Ceterach, 178
 Chærophyllosum, 66
 Cheiranthus, 14
 Chelidonium, 8
 Chenopodiaceæ, 118
 Chenopodium, 118

Cherleria, 26
 Chlora, 94
 Chrysanthemum, 86
 Chrysocoma, 82
 Chrysosplenium, 58
 Cicendia, 94
 Cichorium, 80
 Cicuta, 60
 Cineraria, 86
 Circoea, 50
 Cistaceæ, 16
 Cladium, 156
 Clematis, 2
 Cochlearia, 10
 Colchicum, 144
 Comarum, 40
 Compositæ, 72
 Coniferæ, 134
 Conium, 60
 Convallaria, 144
 Convolvulaceæ, 96
 Convolvulus, 96
 Corallorhiza, 136
 Coriandrum, 66
 Cornaceæ, 60
 Cornus, 60
 Coronopus, 8
 Corrigiola, 54
 Corydalis, 8
 Corylus, 130
 Cotoneaster, 48
 Cotyledon, 56
 Crambe, 8
 Crassulaceæ, 54
 Cratægus, 48
 Crepis, 74
 Crithmum, 64
 Crocus, 140
 Cruciferæ, 8
 Cucubalus, 188
 Cucurbitaceæ, 52
 Cupuliferæ, 128
 Cuscuta, 96, 189
 Cyclamen, 114
 Cynodon, 166
 Cynoglossum, 112
 Cynosorus, 174
 Cyperaceæ, 154
 Cyperus, 154
 Cyripedium, 138
 Cystopteris, 180

 Dactylis, 174
 Daphne, 124
 Datura, 189
 Daucus, 66
 Delphinium, 4
 Dentaria, 12
 Dianthus, 20
 Digitalis, 100

Digitaria, 166
 Dioscoreaceæ, 144
 Dioitis, 82
 Diplotaxis, 14
 Dipsaceæ, 72
 Dipsacus, 72, 189
 Doronicum, 86
 Draba, 10
 Drosera, 18
 Droseraceæ, 18
 Dryas, 40

 Echium, 112
 Elatinaceæ, 20
 Elatine, 20
 Eleagnaceæ, 124
 Eleocharis, 156
 Elodea, 146
 Elymus, 178
 Empetraceæ, 126
 Empetrum, 126
 Epilobium, 50
 Epimedium, 188
 Epipactis, 134
 Epipogium, 136
 Equisetaceæ, 184
 Equisetum, 184
 Eranthis, 4
 Erica, 90
 Ericaceæ, 90
 Erigeron, 84
 Eriocaulon, 152
 Eriophorum, 158
 Erodium, 30
 Eryngium, 60
 Erysimum, 14
 Erythraea, 94
 Euonymus, 30
 Eupatorium, 82
 Euphorbia, 126
 Euphorbiaceæ, 126
 Euphrasia, 100

 Fagus, 128
 Festuca, 174
 Filago, 84
 Filices, 178
 Fœniculum, 64
 Fragaria, 42
 Frankenia, 18
 Frankeniaceæ, 18
 Fraxinus, 94
 Fritillaria, 142
 Fumaria, 8
 Fumariaceæ, 8

 Gagea, 142
 Galanthus, 140
 Galeopsis, 108
 Galinsoga, 88

Galium, 68
 Gastridium, 168
 Genista, 32
 Gentiana, 94
 Gentianaceæ, 94
 Geraniaceæ, 28
 Geranium, 28, 188
 Geum, 40
 Gladiolus, 140
 Glaucium, 8
 Glaux, 116
 Glyceria, 172
 Gnaphalium, 84
 Goodyera, 134
 Gramineæ, 164
 Grossulariaceæ, 54
 Gymnadenia, 136
 Gymnogramma, 178

 Habenaria, 138
 Haloragiaceæ, 50
 Hedera, 58
 Helianthemum, 16
 Helleborus, 4
 Helminthia, 72
 Helosciadium, 62
 Heracleum, 64
 Herminium, 138
 Herniaria, 54
 Hesperis, 14
 Hieracium, 74
 Hierochloë, 166
 Hippocrepis, 36
 Hippophae, 124
 Hippuris, 50
 Holcus, 170
 Holosteum, 24
 Honckenya, 22
 Hordeum, 178
 Hottonia, 114
 Humulus, 128
 Hutchinsia, 10
 Hyacinthus, 142
 Hydrocharidaceæ, 146
 Hydrocharis, 146
 Hydrocotyle, 60
 Hymenophyllum, 182
 Hyoscyamus, 96
 Hypericaceæ, 28
 Hypericum, 28, 188
 Hypochæris, 72

 Iberis, 10
 Ilex, 92
 Illicebrum, 54
 Impatiens, 30, 188
 Inula, 86
 Iridaceæ, 138
 Iris, 138
 Isatis, 8

Isnardia, 50
Isoetes, 184
Isoplepis, 156

Jasione, 90
Juncaceæ, 152
Juncus, 152
Juniperus, 134

Knappia, 168
Knautia, 72
Kobresia, 158
Koeleria, 170
Koniga, 12

Labiatae, 104
Lactuca, 72
Lagurus, 170
Lamium, 108
Lapsana, 78
Lastræa, 180
Lathræa, 102
Lathyrus, 38, 189
Lavatera, 26
Leersia, 164
Leguminosæ, 32
Lemna, 150
Lemnaceæ, 150
Lentibulariaceæ, 112
Leontodon, 72
Leonurus, 108
Lepidium, 10
Lepturus, 178
Leucojum, 140
Ligusticum, 64
Ligustrum, 92
Liliaceæ, 140
Lilium, 140
Limosella, 102
Linaceæ, 26
Linaria, 103, 190
Linnæa, 68
Linum, 26
Liparis, 138
Listera, 134
Lithospermum, 110
Littorella, 118
Lloydia, 142
Lobelia, 90
Loiseleuria, 90
Lolium, 176
Lonicera, 68
Loranthaceæ, 66
Lotus, 36
Luzula, 154
Lychnis, 20
Lycium, 189
Lycopodiaceæ, 184
Lycopodium, 184
Lycopsis, 112

Lycopus, 104
Lysimachia, 114, 190
Lythraceæ, 52
Lythrum, 52

Maianthemum, 144
Malaxis, 138
Malva, 26, 188
Malvaceæ, 26
Marrubium, 110
Marsileaceæ, 184
Matricaria, 88
Matthiola, 14
Meconopsis, 6
Medicago, 32
Melampyrum, 100
Melanthiaceæ, 144
Melica, 172
Melissa, 106
Melitotus, 34
Melittis, 106
Mentha, 104
Menyanthes, 96
Menziesia, 90
Mercurialis, 126
Mertensia, 112
Mespilus, 48
Meum, 64
Miliun, 168
Mimulus, 102
Mœchia, 22
Molinia, 172
Monotropa, 92
Montia, 52
Mulgedium, 74
Muscari, 142
Myosotis, 110
Myosurus, 2
Myrica, 134
Myricaceæ, 134
Myriophyllum, 50
Myrrhis, 66

Naiadaceæ, 148
Naias, 150
Narcissus, 140
Nardus, 178
Narthecium, 152
Nasturtium, 12
Neotinea, 138
Neottia, 134
Nepeta, 110
Nuphar, 6
Nymphæa, 6
Nymphæaceæ, 6

Obione, 120
Oenanthe, 62
Oenothera, 50
Oleaceæ, 92

Onagraceæ, 50
Onobrychis, 36
Ononis, 32
Onopordium, 80
Ophioglossum, 184
Ophrys, 138
Orchidaceæ, 134
Orchis, 136
Origanum, 106
Ornithogallum, 142
Ornithopus, 36
Orobanchaceæ, 102
Orobanche, 102
Osmunda, 182
Oxalidaceæ, 30
Oxalis, 30
Oxyria, 124
Oxytropis, 36

Pæonia, 188
Panicum, 166
Papaver, 6
Papaveraceæ, 6
Parietaria, 128
Paris, 144
Parnassia, 58
Pastinaca, 64
Pedicularis, 100
Peplis, 52
Petasites, 84, 189
Petroselinum, 60
Peucedanum, 64
Phalaris, 166
Phleum, 166
Physospermum, 60
Phyteuma, 90
Picris, 72
Pilularia, 184
Pimpinella, 62
Pinguicula, 112
Pinus, 134
Plantaginaceæ, 118
Plantago, 118
Plumbaginaceæ, 116
Poa, 172
Poleinomaceæ, 96
Polemonium, 96
Polycarpon, 54
Polygala, 18
Polygalaceæ, 18
Polygonaceæ, 122
Polygonatum, 144
Polygonum, 122, 190
Polypodium, 178
Polypogon, 168
Polystichum, 180
Populus, 130
Portulacaceæ, 52
Potamogeton, 148
Potentilla, 40

Poterium, 48
 Primula, 114
 Primulaceæ, 114
 Prunella, 110
 Prunus, 38
 Pseudathyrium, 180
 Pteris, 182
 Pulmonaria, 112
 Pyrola, 92
 Pyrus, 48

Quercus, 128

Radiola, 26
 Ranunculaceæ,
 Ranunculus, 2
 Raphanus, 16
 Reseda, 16
 Resedaceæ, 16
 Restiaceæ, 152
 Rhamnaceæ, 32
 Rhamnus, 32
 Rhinanthus, 100
 Rhyncospora, 156
 Ribes, 54
 Rosa, 46
 Rosaceæ, 38
 Rubia, 68
 Rubiaceæ, 68
 Rubus, 42
 Rumex, 122
 Ruppia, 150
 Ruscus, 144

Sagina, 22
 Sagittaria, 146
 Salicaceæ, 130
 Salicornia, 120
 Salix, 130
 Salsola, 120
 Salvia, 140, 190
 Sambucus, 68
 Samolus, 116
 Sanguisorba, 48
 Sanicula, 60
 Santalaceæ, 124
 Saponaria, 20
 Saromanthus, 32
 Saussurea, 80
 Saxifraga, 56
 Saxifragaceæ, 56
 Scabiosa, 72
 Scandix, 66
 Scheuchzeria, 148
 Schoenus, 156
 Scilla, 142
 Scirpus, 156

Scleranthaceæ, 54
 Scleranthus, 54
 Sclerochloa, 172
 Schrophularia, 100
 Schrophulariaceæ, 98
 Scolopendrium, 182
 Scutellaria, 110
 Sedum, 56
 Selaginella, 184
 Sempervivum, 56
 Senecio, 86
 Serratula, 80
 Seseli, 64
 Sesleria, 170
 Setaria, 166
 Sherardia, 70
 Sibbaldia, 40
 Sibthorpia, 102
 Silaus, 64
 Silene, 20, 188
 Simethis, 144
 Sinapis, 14
 Sison, 62
 Sisymbrium, 14
 Sisyrinchium, 140
 Sium, 62
 Smyrnium, 60
 Solanaceæ, 96
 Solanum, 96
 Solidago, 84
 Sonchus, 74
 Sparganium, 152
 Spartina, 166
 Spergula, 22
 Spergularia, 22
 Spiræa, 40
 Spiranthes, 134
 Stachys, 108, 190
 Staphyllea, 188
 Statice, 116
 Stellaria, 24
 Stratiotes, 146
 Subularia, 10
 Succeda, 120
 Symphytum, 112

Tamariscaceæ, 52
 Tamarix, 52
 Tamus, 144
 Tanacetum, 82
 Taraxacum, 78
 Taxus, 134
 Teesdalia, 10
 Teucrium, 106
 Thalictum, 2
 Thesium, 124
 Thlaspi, 10

Thymelaceæ, 124
 Thymus, 106
 Tilia, 26
 Tiliaceæ, 26
 Tillæa, 54
 Tofieldia, 146
 Tordylium, 66
 Torilis, 66
 Tragopogon, 72
 Trichomanes, 182
 Trichonema, 140
 Trientalis, 114
 Trifolium, 34, 189
 Triglochin, 146
 Trigonella, 34
 Trilliaceæ, 144
 Trinia, 60
 Triodia, 170
 Triticum, 176
 Trollius, 4
 Tulipa, 140
 Tussilago, 84
 Typha, 152
 Typhaceæ, 152

Ulex, 32
 Ulmaceæ, 128
 Ulmus, 128
 Umbelliferæ, 60
 Urtica, 128
 Urticaceæ, 128
 Utricularia, 114

Vaccinium, 92
 Valeriana, 70
 Valerianaceæ, 70
 Valerianella, 70
 Verbascum, 98
 Verbena, 104
 Verbenaceæ, 104
 Veronica, 98
 Viburnum, 68
 Vicia, 36
 Villarsia, 94
 Vinca, 94
 Viola, 16
 Violaceæ, 16
 Viscum, 66

Woodsia, 178
 Woolfia, 150

Xanthium, 189

Zannichellia, 150
 Zostera, 150

INDEX OF THE COMMON NAMES.

-
ACONITE, 4	Birch, 130	Buttercup, 2
Adder's-tongue, 184	Bird-cherry, 40	Butterfly Orchis, 138
Agrimony, 46	Bird's-foot, 36	Butter-wort, 112
Alder, 130	Bird's-nest, 92, 134	
Alexanders, 60	Birth-wort, 126	Cabbage, 14
Alkanet, 112	Bishop-weed, 62	Calamint, 106
Allseed, 54	Bistort, 122	Campion, 20
Allysum, 12	Bitter-cress, 12	Canary-grass, 166
Amaranth, 118	Bitter-sweet, 95	Candy-tuft, 10
Andromeda, 90	Bitter-vetch, 38	Caraway, 62
Anemone, 2	Blackberry, 42	Carline Thistle, 82
Angelica, 64	Black Salt-wort, 116	Carnation, 20
Apple, 48	Blackthorn, 38	Carrot, 66
Archangel, 108	Bladder-fern, 180	Catch-fly, 20
Arrow-grass, 146	Bladder-seed, 60	Cat-mint, 110
Arrow-head, 146	Bladder-wort, 114	Cat's-ear, 72
Asarabacca, 124	Blinks, 52	Cat's-foot, 82
Ash, 94	Blue-bell, 142	Cat's-tail, 152
Asparagus, 144	Blue-bottle, 82	Cat's-tail-grass, 166
Aspen, 130	Bog-Asphodel, 152	Celandine, 8
Avens, 40	Bog-Myrtle, 134	Celery, 60
Awl-wort, 10	Bog Orchis, 138	Centaury, 94
Azalea, 90	Bog-rush, 156	Ceterach, 178
	Borage, 112	Chaff-weed, 116
Bald-money, 64	Box, 126	Chamomile, 88
Balm, 106	Brake, 182	Charlock, 14
Balsam, 30	Bristle-fern, 182	Cherry, 38
Baneberry, 6	Bristle-grass, 166	Chervil, 66
Barberry, 6	Brome-grass, 176	Chickweed, 24
Barley, 178	Brook-lime, 98	Chickweed winter-green,
Bartsia, 98	Brook-weed, 116	114
Basil, 106	Broom, 32	Chicory, 80
Bastard Alkanet, 112	Broom-rape, 102	Chives, 142
Bastard Balm, 106	Bryony, 52, 144	Cicely, 66
Bastard Pimpernel, 116	Buck-bean, 96	Cinque-foil, 40
Bastard Toad-flax, 124	Buckler-fern, 180	Clary, 104
Beaked Parsley, 66	Buck-thorn, 32	Cleavers, 70
Beak-rush, 156	Buck-wheat, 122	Cloudberry, 42
Bear-berry, 92	Bugle, 106	Clover, 34
Beard-grass, 168	Bugloss, 112	Club-moss, 184
Bedstraw, 68	Bullace, 38	Club-rush, 156
Beech, 128	Bulrush, 152	Cock's-foot-grass, 174
Beech-fern, 178	Burdock, 80	Colt's-foot, 84
Bee Orchis, 138	Bur-marigold, 82	Columbine, 4
Beet, 120	Burnet, 48	Comfrey, 112
Bell-flower, 88	Burnet Saxifrage, 62	Coral-root, 136
Bent-grass, 168	Bur-parsley, 66	Coral-wort, 12
Betony, 108	Bur-weed, 34, 152	Cord-grass, 166
Bilberry, 92	Butcher's-broom, 144	Coriander, 66
Bindweed, 96	Butter-bur, 84	Corn-cockle, 22

- Cornel, 60
 Corn-flag, 140
 Corn Marigold, 86
 Corn-salad, 70
 Corydalis, 8
 Cotoneaster, 48
 Cotton-grass, 158
 Cotton-thistle, 80
 Couch-grass, 176
 Cowbane, 60
 Cowberry, 92
 Cow-parsnep, 64
 Cowslip, 114
 Cow-wheat, 100
 Crab-tree, 48
 Crake-berry, 126
 Cran-berry, 92
 Crane's-bill, 28
 Crocus, 140
 Cross-wort, 68
 Crowberry, 126
 Crowfoot, 2
 Cuckoo-flower, 12
 Cuckoo-pint, 150
 Cudweed, 84
 Currant, 54
 Cut-grass, 164
 Cyphel, 26

 Daffodil, 140
 Daisy, 86
 Dame's Violet, 14
 Dandelion, 78
 Dane-wort, 68
 Darnel, 176
 Deadly Nightshade, 96
 Dead-nettle, 108
 Devil's-bit Scabious, 72
 Dock, 122
 Dodder, 96
 Dog-rose, 46
 Dog's-tail-grass, 174
 Dog's-tooth-grass, 166
 Dog-violet, 16
 Dog-wood, 60
 Drop-wort, 62
 Duck-weed, 150
 Dwale, 96
 Dyer's-weed, 16, 32
 Dyer's-wood, 8

 Earth-nut, 62
 Earth-nut-pea, 38
 Elder, 68
 Elecampane, 86
 Elm, 128
 Enchanter's Nightshade, 50
 Eryngo, 60
 Evening Primrose, 50
 Everlasting Pea, 38
 Eye-bright, 100

 Feather-foil, 114
 Fennel, 64
 Fennugreek, 34
 Fescue-grass, 174
 Feverfew, 88
 Fig-wort, 100
 Filago, 84
 Filmy-fern, 182
 Finger-grass, 166
 Fir, 134
 Flag, 138
 Flax, 26
 Flax-seed, 26
 Flea-bane, 84, 86
 Flea-wort, 86
 Flix-weed, 14
 Flowering-rush, 146
 Fluellen, 100
 Fly-Orchis, 138
 Fool's-Parsley, 64
 Forget-me-not, 110
 Fox-glove, 100
 Fox-tail-grass, 166
 Fritillary, 142
 Frog-bit, 146
 Frog-Orchis, 138
 Fumitory, 8
 Furze, 32

 Gagea, 142
 Gale, 134
 Galingale, 154
 Garlic, 142
 Garlic-Mustard, 14
 Gentian, 94
 Germander, 106
 Gipsy-wort, 104
 Gladdon, 138
 Gladiolus, 140
 Glass-wort, 120
 Globe-flower, 4
 Goat's-beard, 72
 Golden-rod, 84
 Golden-Samphire, 86
 Golden-Saxifrage, 58
 Goldilocks, 4, 82
 Gold-of-pleasure, 12
 Good King Henry, 120
 Gooseberry, 54
 Goose-foot, 118
 Goose-grass, 70
 Gorse, 32
 Gout-weed, 62
 Grape-hyacinth, 142
 Grass-of-Parnassus, 58
 Grass-vetch, 38
 Grass-wrack, 150
 Green-weed, 32
 Gromwell, 110, 112
 Ground-ivy, 110
 Ground-pine, 108

 Groundsel, 86
 Guelder-rose, 68
 Gymnogramm, 178

 Habenaria, 138
 Hairbell, 88
 Hair-grass, 170
 Hard-fern, 182
 Hard-grass, 178
 Hare's-ear, 14, 62
 Hare's-tail-grass, 170
 Hart's-tongue, 182
 Hart-wort, 66
 Havers, 170
 Hawk-bit, 72
 Hawk's-beard, 74
 Hawk-weed, 74
 Hawthorn, 48
 Hazel, 130
 Heartsease, 16
 Heath, 90
 Heath-grass, 170
 Hedge-mustard, 14
 Hedge-parsley, 66
 Hellebore, 4
 Helleborine, 134, 136
 Hemlock, 60
 Hemp-Agrimony, 82
 Hemp-nettle, 108
 Henbane, 96
 Henbit, 108
 Herb-Christopher, 6
 Herb-Paris, 144
 Hog's-fennel, 64
 Hog-weed, 64
 Holly, 92
 Holly-fern, 180
 Holy-grass, 166
 Honeysuckle, 68
 Hone-wort, 60
 Hop, 128
 Horehound, 108, 110
 Hornbeam, 128
 Horned Poudweed, 150
 Horned Poppy, 8
 Horn-wort, 52
 Horse-radish, 10
 Horse-shoe Vetch, 36
 Horse-tail, 184
 Hound's-tongue, 112
 House-leek, 56
 Hutchinsia, 10

 Inula, 86
 Iris, 138
 Isle-of-Man Cabbage, 14
 Ivy, 58

 Jacob's Ladder, 96
 Joint-vetch, 36
 Juniper, 134

 Kale, 8

Kidney-vetch, 32
 Knap-weed, 82
 Knautia, 72
 Knawel, 54
 Knot-grass, 54, 122
 Koniga, 12
 Lady-fern, 180
 Lady's Fingers, 32
 Lady's Mantle, 48
 Lady's Slipper, 138
 Lady's Tresses, 134
 Lamb's Lettuce, 70
 Larkspur, 4
 Leek, 142
 Leopard's-bane, 86
 Lettuce, 72
 Lily, 140
 Lily-of-the-Valley, 144
 Lime, 26
 Ling, 90
 Livelong, 56
 Lobelia, 90
 London Pride, 58
 London Rocket, 14
 Loosestrife, 52, 114
 Lords and Ladies, 150
 Louse-wort, 100
 Lovage, 64
 Lucerne, 32
 Lung-wort, 112
 Lyme-grass, 178
 Madder, 68, 70
 Mad-wort, 112
 Maiden-hair, 182
 Male-fern, 180
 Mallow, 26
 Manna-Croup-grass, 172
 Man-Orchis, 138
 Maple, 28
 Mare's-tail, 50
 Marjoram, 106
 Marram, 168
 Marsh-fern, 180
 Marsh-mallow, 26
 Marsh-marigold, 4
 Marsh-trefoil, 96
 Marsh-wort, 62
 Martagon-Lily, 140
 Master-wort, 64
 Mat-grass, 168, 178
 May, 48
 May Lily, 144
 May-weed, 88
 Meadow-grass, 172
 Meadow-Rue, 2
 Meadow-Saffron, 144
 Meadow-Saxifrage, 64
 Meadow-Sweet, 40
 Medick, 32
 Medlar, 48

Melic-grass, 172
 Melilot, 34
 Menziesia, 90
 Mercury, 128
 Meu, 64
 Mezereon, 124
 Michaelmas Daisy, 84
 Mignonette, 16
 Milfoil, 50, 88
 Milk-parsley, 64
 Milk-thistle, 80
 Milk-vetch, 36
 Milk-wort, 18
 Millet-grass, 168
 Mint, 104
 Mistletoe, 66
 Moenchia, 22
 Money-wort, 102, 114
 Monk's-hood, 6
 Monk's-Rhubarb, 124
 Moon-fern, 182
 Moor-grass, 170
 Moschatel, 58
 Mother-wort, 108
 Mountain-Ash, 48
 Mountain-fern, 180
 Mountain-Sorrel, 124
 Mouse-ear Chickweed, 24
 Mouse-tail, 2
 Mud-rush, 156
 Mud-wort, 102
 Mug-wort, 68, 82
 Mullen, 98
 Musk-Orchis, 138
 Mustard, 14
 Narcissus, 140
 Navew, 14
 Needle-whin, 32
 Nettle, 128
 Nightshade, 96
 Nipple-wort, 78
 Nit-grass, 168
 None-so-pretty, 58
 Nonsuch, 32
 Oak, 128
 Oak-fern, 178
 Oat-grass, 170
 Onion, 142
 Orache, 120
 Orchis, 136
 Orpine, 56
 Osier, 132
 Ox-eye, 86
 Ox-lip, 114
 Ox-tongue, 72
 Oxytropis, 36
 Panic-grass, 166
 Pansy, 16
 Parsley, 60

Parsley-fern, 178
 Parsley-Piert, 48
 Parsnep, 64
 Pasque-flower, 2
 Pea, 38
 Pearl-wort, 22
 Pear-tree, 48
 Pellitory, 128
 Penny-cress, 10
 Penny-royal, 106
 Penny-wort, 56, 60
 Pepper-Saxifrage, 64
 Pepper-wort, 10
 Periwinkle, 94
 Persicaria, 122
 Petty-Whin, 32
 Pheasant's-eye,
 Picris, 72
 Pile-wort, 4
 Pill-wort, 184
 Pimpernel, 114
 Pine, 134
 Pink, 20
 Pipe-wort, 152
 Plaintain, 118
 Plowman's Spikenard, 86
 Plum, 38
 Polypody, 173
 Pond-weed, 148
 Poplar, 130
 Poppy, 6
 Primrose, 114
 Privet, 92
 Purple Loosestrife, 52
 Purslane, 120
 Quaking-grass, 174
 Quill-wort, 184
 Radish, 16
 Ragged-Robin, 20
 Rag-wort, 86
 Rampion, 90
 Ransoms, 142
 Rape, 14
 Raspberry, 42
 Reed, 168
 Reed-mace, 152
 Rest-harrow, 32
 Rib-wort, 118
 Rock-brake, 178
 Rock-cress, 12
 Rocket, 14
 Rock-rose, 16
 Rose, 46
 Rose-bay, 50
 Rose-root, 56
 Royal-fern, 182
 Rupture-wort, 54
 Rush, 152
 Rye-grass, 176

- Saffron, 144
 Sage, 104
 Sainfoin, 36
 Saint Dabeoc's Heath, 90
 Sallow, 130
 Salsify, 72
 Salt-wort, 116, 120
 Samphire, 64
 Sand-wort, 22
 Sanicle, 60
 Saw-wort, 80
 Saxifrage, 56
 Scabious, 72
 Scorpion-grass, 110
 Scotch Fir, 134
 Scottish Asphodel, 146
 Scurvy-grass, 10
 Sea-blite, 120
 Sea Buckthorn, 124
 Sea-heath, 18
 Sea-holly, 60
 Sea-kale, 8
 Sea-lavender, 116
 Sea Milk-wort, 116
 Sea-pink, 116
 Sea-purslane, 22, 120
 Sea-radish, 16
 Sea-reed, 168
 Sea-rocket, 8
 Sedge, 158
 Self-heal, 110
 Service-tree, 48
 Sheep's-bit, 90
 Shepherd's-needle, 66
 Shepherd's-purse, 10
 Shield-fern, 180
 Shore-weed, 118
 Silver-weed, 40
 Skull-cap, 110
 Sloe, 38
 Small-reed, 168
 Snake's-head, 142
 Snake-weed, 122
 Snap-dragon, 100
 Sneeze-wort, 88
 Snow-drop, 140
 Snow-flake, 140
 Soap-wort, 20
 Soft-grass, 170
 Solomon's-seal, 144
 Sorrel, 122
 Sow-bread, 114
 Sow-thistle, 74
 Spear-wort, 2
 Speedwell, 98
 Spider-Orchis, 138
 Spignel, 64
 Spike-rush, 156
 Spindle-tree, 30
 Spleen-wort, 182
 Spurge, 126
 Spurge Laurel, 124
 Spurrey, 22
 Squill, 142
 Squinancy-wort, 70
 Star-fruit, 146
 Star-of-Bethlehem, 142
 Star-thistle, 82
 Star-wort, 84
 Stitch-wort, 24
 St. John's-wort, 28
 Stock, 14
 Stone-crop, 56
 Stone-parsley, 62
 Stork's-bill, 28
 Strap-wort, 54
 Strawberry, 42
 Strawberry-tree, 92
 Succory, 80
 Sulphur-weed, 64
 Sulphur-wort, 62
 Sundew, 18
 Sweet-briar, 46
 Sweet-flag, 150
 Sycamore, 28
 Tamarisk, 52
 Tansy, 82
 Tare, 36
 Tassel Pond-weed, 150
 Teasel, 72
 Teesdalia, 10
 Thale-cress, 14
 Thistle, 80
 Thrift, 116
 Thyme, 106
 Tillæa, 54
 Toad-flax, 100
 Tooth-wort, 102
 Tower-mustard, 12
 Tower-wort, 12
 Traveller's-joy, 2
 Treacle-mustard, 14
 Tree-mallow, 26
 Trefoil, 34
 Tulip, 140
 Turk's-cap Lily, 140
 Turnip, 14
 Tutsan, 28
 Tway-blade, 134
 Twig-rush, 156
 Valerian, 70
 Venus' Comb, 66
 Vernal-grass, 166
 Vervain, 104
 Vetch, 36
 Violet, 16
 Viper's Bugloss, 112
 Wake-Robin, 150
 Wallflower, 14
 Wall-Pellitory, 128
 Wall-pepper, 56
 Wall-rue, 182
 Wart-cress, 8
 Water-cress, 12
 Water-dropwort, 62
 Water-lily, 6
 Water-Milfoil, 50
 Water-parsnep, 62
 Water-pimpernel, 116
 Water-plaintain, 146
 Water-purslane, 52
 Water-Soldier, 146
 Water Star-wort, 50
 Water-thyme, 146
 Water-violet, 114
 Water-wort, 20
 Wayfaring-tree, 68
 Weasel-snout, 108
 Welch-Poppy, 6
 Wheat-grass, 176
 Whin, 32
 White Beam-tree, 48
 White-rot, 60
 Whitlow-grass, 10
 Whorl-grass, 172
 Whortle-berry, 92
 Wild Hyacinth, 142
 Willow, 130
 Willow-herb, 50
 Winter-aconite, 4
 Winter-cress, 12
 Winter-green, 92
 Woad, 8
 Wood-bine, 68
 Wood-ruff, 70
 Wood-rush, 154
 Woodsia, 178
 Wood-sorrel, 30
 Worm-seed, 14
 Worm-wood, 82
 Wound-wort, 108
 Yarrow, 88
 Yellow-cress, 12
 Yellow-rattle, 100
 Yellow-wort, 94
 Yew, 134