University of Alberta Library

0 1620 3416617 1

For Reference

NOT TO BE TAKEN FROM THIS ROOM

MORPHOLOGICAL, PHYSIOLOGICAL AND GENETICAL
STUDIES OF EARLINESS IN CEREAL CROPS

Aston R. Taylor

Department of Plant Science
UNIVERSITY OF ALBERTA

For Reference

NOT TO BE TAKEN FROM THIS ROOM

Ex LIBRIS
UNIVERSITATIS
ALBERTAENSIS

Digitized by the Internet Archive
in 2018 with funding from
University of Alberta Libraries

<https://archive.org/details/morphologicalphy00tayl>

T H E U N I V E R S I T Y O F A L B E R T A

MORPHOLOGICAL, PHYSIOLOGICAL AND GENETICAL STUDIES
OF EARLINESS IN CEREAL CROPS

A DISSERTATION
SUBMITTED TO THE SCHOOL OF GRADUATE STUDIES
IN PARTIAL FULFILMENT OF THE REQUIREMENTS FOR THE DEGREE
OF MASTER OF SCIENCE

FACULTY OF AGRICULTURE

by

Aston Rupert Taylor

EDMONTON, ALBERTA

SEPTEMBER, 1950

TABLE OF CONTENTS

	<u>Page</u>
I. LITERATURE REVIEW	1
II. OBJECT OF THE STUDY	11
III. MORPHOLOGICAL STUDY	12
1. Description of Varieties and Strains	12
2. Experimental Methods	13
3. How Earliness was Determined	15
4. Observations on Leaf Emergence	16
5. Observations on Tiller Emergence	18
6. Observations on Growth Habit	19
7. Observations on Leaf-Blade Curling	25
8. Study of Spike Characteristics	28
IV. PHYSIOLOGICAL STUDY	35
1. Observations on Development of Barley	35
2. Materials and Methods	36
3. Histological Observations	37
4. Photoperiodic Reaction with Advancing Age of Plant	48
5. Photoperiodism and Heading	52
6. Observations on Heights of Plants	62

TABLE OF CONTENTS (Continued)

	<u>Page</u>
V. GENETICAL STUDY	67
1. Materials, Methods, and Observations	67
2. Relation of Time of Heading to Length of Heading Period	67
3. Experimental Results	72
(a) The F_1 Generations	72
(b) The F_2 Generations	74
(c) Observations on Variation	89
4. Interpretation of F_2 Results	91
(a) The Hypothesis	91
(b) The Quantitative Effect of the Genes ...	94
VI. GENERAL DISCUSSION AND CONCLUSIONS	97
ACKNOWLEDGEMENTS	100
LITERATURE CITED	101

MORPHOLOGICAL, PHYSIOLOGICAL AND GENETICAL STUDIES
OF EARLINESS IN CEREAL CROPS

A. R. Taylor

I. LITERATURE REVIEW

It is now commonly accepted that differences in earliness in plants are conditioned by a series of factors. However, a survey of available literature relating to cereal crops reveals that, while the physiology and genetics of earliness have received considerable study, the morphological aspect has been largely neglected.

The only publication on the morphology of earliness in cereal crops known to the writer is that of Suneson (29). He has shown that the effect of environment on earliness varies with the morphology of the plant.

In physiological studies Klebs (21) was able, by manipulating the light and temperature of the environment, to maintain continuous vegetative growth or to induce reproduction prematurely.

He considered that both light and temperature could affect the balance between carbohydrates and soil nutrients, chiefly nitrogen. From a study of *Sempervivum* he observed that the process of flower formation proceeded in three separate phases -

- (1) onset of a ripe-to-flower^{*} condition,
- (2) formation of the flower primordia,
- (3) development of the flowering inflorescence.

He found that the initiation of each phase was dependent on the relative amounts of light intensity and temperature. A one-sided increase of one or the other would prolong phase (1) or (2) resulting in a delay in phase (3).

In 1934 Purvis (28) published data of his analysis on the influence of temperature and photoperiod during germination, on the subsequent development of winter rye. His study proved that length of day was interrelated with other conditions of germination in determining subsequent flower behaviour.

Harrington and Horner (13), working at the University of Saskatchewan, did a series of experiments on the reaction of wheat varieties to different dates of sowing. They found that the interaction between varieties and dates was statistically significant.

Platt (26), working with wheat, showed the relationship between soil moisture and winter survival and established a correlation between earliness and a low survival index.

* This is the stage at which the growing point shows a differentiation into primordium and stem.

Working with oats, Tincker and Jones (33) undertook a study of the influence of certain climatic factors upon the rate of growth. It was shown that the Unit Leaf Rate (measured on the "tops" of the plants only) was correlated with the previous rainfall and also with the temperature, during that period of growth in which the plants were forming new leaves and increasing their leaf area.

Jasny(18) investigated the variability in growing periods of several crops from latitude to latitude and found that variations in the total growing period of winter grains were caused primarily by the different lengths of the dormant period.

The investigations of Garner and Allard (10) on the influence of the relative length of day upon the rate of development of plants have shown that the time of flowering in some plants is controlled by the length of day. These workers (10) have reported one test on the effect of prolonging the illumination period of oats, wheat, barley, and rye, using a spring and a winter variety of each.

With each of these grains the forcing action of the electric light was much greater with the spring than with the winter variety. The authors state the following conclusion:

"It appears that the fundamental distinction between the winter and the spring types as such rests on the rapidity with which the latter respond to the increasing day length of spring".

Eguchi (7) gives the results of experiments with short-day treatments. Some plants hitherto regarded as short-day types were tested under short-day conditions (9-hour day). The differentiation of the flower bud was accelerated and the flower opened early, but, if these plants were transferred to long-day conditions (24-hour day) further development took place rapidly and flowering occurred earlier. Eguchi concludes that there are two stages with different reactions to the photoperiod in the course of flowering, the first being the stage of bud differentiation, and the second the stage of blooming.

It is appropriate at this time to refer to the importance of micro-dissection of plants to determine the actual stage of flowering reached, as stressed in the work of Gregory and Purvis (11) and by Hammer (12) who states: "It would be helpful in much of the experimental work in this field if the investigators would examine all of their plants by micro-dissection to determine what stages of floral development are especially affected by the treatments used".

In a genetic study involving a cross between a spring and a winter variety of wheat, Aamodt (1) secured an F_2 population which when spring-sown began to head with the spring parent and continued heading for a period of eight weeks. There were also some plants which like the winter parent, did not head. He recovered a homozygous family five weeks later than the spring parent.

His results also indicated that the difference between the two varieties was due to a series of multiple factors.

Johnson (19) found, in crosses involving early by late varieties of barley, that the hybrid distributions were all skewed toward the "early" end. He suggested that this behaviour indicated dominance or partial dominance of earliness, and that from the degree of skewness partial dominance appeared to be the more likely. He observed that, from the particular lines studied, the inheritance of earliness might lie in a single-gene difference.

In a study of the nature and interaction of genes differentiating habit of growth in wheat, Powers (27) observed that the genotypes of the F_2 suggested parents which were differentiated by three main factor pairs for habit of growth with modifying factors for earliness present. He believed that the environmental conditions were largely responsible for the reaction processes of the different genes.

Biffen (3), from his observations on an early Polish and a late Rivet wheat, concluded that earliness and lateness were inherited independently of other characters. Although the F_1 ripened well toward the date of ripening of the Rivet parent he concluded from the F_2 , which had a high percentage of early lines, that earliness was dominant. Fruwith, as cited by Florell (9), believed early maturity to be partially dominant in wheat as well as in rye and barley.

Thompson, in a series of three articles (30, 31, 32) reports on an extensive study of early and late maturity in wheat. In the F_1 he found, in general, that ripening occurred at or near the date of ripening of the later parent. He thought that this behaviour was due to hybrid vigor causing added vegetative growth and not to the dominance of lateness. He found in the F_2 that the great majority of hybrids were intermediate between the parents and in all types of crosses the distribution approached the normal curve of variation. No skewness toward either parental position was noted that could suggest dominance or simple segregation. On the basis of Mendelian segregation a multiple factor explanation was offered. While the multiple factor hypothesis held fairly well for individual crosses it could not be applied directly to the entire series. Thompson (32), however, points out that many factors in the hereditary constitution doubtless affect the rate of maturity, and many factors whose primary effects are on entirely different characteristics may have secondary effects on earliness. No linkage was found between earliness or lateness and other characters.

Bryan and Pressley (4) noted that in a Sonora x Turkey cross, the F_1 progeny was exactly intermediate between the parents for date of first heading. The F_2 showed an inclination toward the late parent but extended over a range slightly exceeding both parents. They were able to establish homozygous lines for early,

intermediate and late heading in the F_4 generation.

In 1924 Florell (9) reported on crosses between Sunset (early) and Marquis (late). In the F_1 the date of first ripening was slightly earlier than that of the Marquis parent. In the F_2 there was a distinct segregation into a large early group and a small late group in a ratio of 3.07 early, 0.93 late, close to a 3:1 ratio. The indication was for one main allelomorphic pair of factors for earliness.

Studies on maturity were reported by Clark and Hooker (6) in 1926 on crosses of Marquis x Hard Federation. Three criteria of maturity were studied (1) the dates of heading, (2) the dates of ripening, and (3) the number of days between (1) and (2), or the fruiting period.

Partial dominance was shown by the F_2 data and segregation in the F_3 indicated that at least two genetic factors for early heading were involved. It was also noted that some F_2 families were segregating in the F_3 in a simple 3:1 ratio as found by Florell (9).

Among other characters studied in Kota x Hard Federation crosses Clark (5) reported in 1924 on maturity. He used as a criterion of earliness the date of heading and concluded that earliness was dominant over lateness. Few strains were found to be as uniform in the F_3 as the parents.

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

Since one might expect the inheritance of earliness in all plants to be similar, it is of interest to note what has been found in studies of earliness in other plants besides the cereal crops.

Mendel (24) reported that with garden peas the time of flowering of the F_1 hybrids was almost exactly intermediate between those of the parents, but further results were not given. In other crosses of garden peas, Tschermak (34) secured F_1 plants intermediate in earliness, and in the F_2 families, plants varying from earlier than the early parent to later than the late parent. Some of the extremes bred true, but the intermediates gave offspring variable in time of flowering. His explanation was that two factors for earliness were involved.

In cotton, Leake (22), found the F_1 generation uniformly intermediate in time of flowering. In the F_2 populations there was variation in time of flowering with the mode and the mean nearer the late parent.

Keeble and Pellow (20) from a cross between early and late garden peas concluded that lateness was dominant. The F_1 plants were earlier than the late parent but this was thought to be due to coupling (linkage) between thin stems, which were dominant to thick stems, and earliness.

Veselovskii (35) working with interspecific crosses in potato found that in the F_1 of the cross S. andigenum x S. tuberosum the following characters were dominant: late maturity, fertility, ability to form tubers and a large number of tubers to the cluster. By back-crossing with early commercial varieties a combination of the following characters was obtained: early formation of tubers, short period of tuber development, early normal completion of the vegetative period and high productivity.

Emerson and East (8) in corn crosses secured F_1 hybrids intermediate between the parents in earliness, and F_2 populations which had a range exceeding both parents. In the F_3 they had pure types practically as early as the early parent but none quite so late as the late parent.

Working with corn, Hayes and East (14) observed that the F_1 hybrids matured earlier than the average date of maturity of the parents. They believed that the increase in rate of maturity was due to hybrid vigor.

Hutcheson and Wolfe (17) compared, as to earliness, the F_1 hybrids of six crosses of corn with the parent varieties. One cross was earlier in tasseling and silking than the early parent and had a higher percentage of mature corn. The others were intermediate in earliness but, except in the case of one cross, were more nearly like the early parents than the late.

From a cross between two varieties of rice differing in earliness, Hoshina (16) recovered pure early races like the early parent in earliness. He favoured the multiple factor hypothesis and thought that in his cross the parents differed in three factors.

In his research on the influence of inbreeding upon the season of maturity, Pearson (25) compared inbred lines of cabbage with the hybrids between them and found that in crosses involving unrelated lines there was an increase in earliness. This difference was not observed in crosses of closely related lines. The yields showed similar relationships. He further observed that in some cases earliness was governed by definite genetic factors.

In crosses of tomatoes, Hayes and Jones (15) found early maturity associated with increase in production, and that crosses which did not show increase in earliness did not show increase in production.

II. OBJECT OF THE STUDY

This study is exploratory in nature with a threefold objective:

- (a) to observe the form and structure of various strains of barley - early, medium, and late - in order to determine the extent to which morphology may be used as a criterion of earliness,
- (b) to compare the rates of development of early, medium, and late lines under different photoperiods and temperatures,
- (c) to get more definite information concerning the genetic factors responsible for earliness and especially to determine if they are alike in their effects, and the number of factors involved in certain crosses.

III. MORPHOLOGICAL STUDY

1. Description of Varieties and Strains

The varieties or strains used are listed below; each one is given a number which is used to designate the variety or strain in the tables.

The seed was furnished by the Cereal Crops Division of the Plant Science Department, University of Alberta. Before the initiation of this study, all materials were tested for their purity for earliness.

Table 1

Varieties and Strains

No.	Material	C.I. No.	Type	Season	Genus
			Spring (S)		
1	Atsel	6250	S	Early	<u>H. vulgare</u>
2	Beecher	6566	S	Early	"
3	Tulare	--	S	Early	"
4	H-44-1	--	S	Early to medium	"
5	Titan	7055	S	Medium	"
6	Newal	6088	S	Medium	"
7	Montcalm	7149	S	Late	"
8	C.I. 7155	--	S	Late	"
9	C.I. 7248	--	S	Late	"

2. Experimental Methods

Crosses involving the individuals listed in Table 1 were made in the field. The F_0 materials together with the standard varieties were seeded in uniform soil in the greenhouse beds on August 22, 1949. All hybrids and varieties were sown in rows 8 inches apart, the seeds being seeded individually 4 inches apart.

Morphological observations made in the greenhouse study were again made under field conditions in the spring and summer of 1950 when the F_2 generation was grown.

Since several crosses were studied, each was given a designation number for convenience in keeping the field records. These numbers are used in the tables, but the names of varieties or strains are indicated in the headings.

The crosses numbered 1x to 16 x, inclusive, were made in the field in 1948 and 1949, and the F_1 plants and parents grown in the greenhouse in the fall and winter of 1949-1950. All crosses were compared with the parents in earliness.

Table 2

Crosses Studied

Designation numbers	Varieties crossed	When grown	
		F ₁	F ₂
1x	Atsel x Tulare	'49	'50
2x	Montcalm x Tulare	"	"
3x	Montcalm x Beecher	"	"
4x	Atsel x Beecher	"	"
5x	Titan x Newal	"	"
6x	Montcalm x Atsel	"	"
7x	Montcalm x Titan	"	"
8x	Montcalm x Newal	"	"
9x	Beecher x C.I. 7248	"	"
10x	Titan x Tulare	"	"
11x	Titan x C.I. 7248	"	"
12x	Montcalm x C.I. 7155	"	"
13x	Atsel x C.I. 7155	"	"
14x	Atsel x C.I. 7248	"	"
15x	Montcalm x C.I. 7248	"	"
16x	C.I. 7155 x C.I. 7248	"	"

3. How Earliness was Determined

After examining the plants at different stages of development, earliness was based upon the date when the first plant began to head. The date for a particular plant was that on which the first spikelet of the earliest spike came through the flag leaf. During the time of heading, the plants were observed at the same hour each day and marked as they headed with dated string tags. There were only a few F_1 plants of most crosses, but for the F_2 populations all the seed was sown in four replications or dates of seeding on the rod-row plan.

4. Observations on Leaf Emergence

For each variety or hybrid, notes were kept on the date of emergence of the first plant in each line. Emergence dates for the first appearance of a second leaf, third leaf, etc. up to the flag leaf were also taken. These observations were also made on a population basis. With the appearance of the third leaf, notes were taken showing the dates when 75% of the plants and the last plant in each line had acquired a third leaf. This was done for the fourth leaf, fifth leaf, and so on, up to the flag leaf. For complete data on dates of emergence see Tables 9 and 10.

It was observed that late varieties emerged later than early ones. The emergence of the several leaves up to the flag leaf was consistently later for the later lines, but the average lag for each stage of emergence was only two days. Once the flag leaf was acquired there was a significant lag in the date of heading of the later varieties. The correlation between the number of leaves and the dates of emergence for each of two varieties, one early and one late, and the F_1 hybrid between them is shown in Fig. 1.

Fig. 1

Correlation between dates of leaf emergence, varieties and F_1 hybrid.

The late variety, Montcalm, took longer to complete the various stages of leaf development than the early line, Atsel. The hybrid 6x was earlier in leaf emergence than Montcalm and was not intermediate but closer to the early Atsel. In the field the behaviour of these two lines was very similar but leaf emergence dates for the hybrid were closer to Atsel in the later stages of development preceding heading.

5. Observations on Tiller Emergence

The appearance of tillers did not follow the pattern of leaf emergence. Except for the early variety Atsel, the three latest lines, Montcalm, C.I. 7155, and C.I. 7248, showed the emergence of a first tiller in advance of the early lines Beecher and Tulare and of the midseason varieties Newal and Titan. This behaviour seems to indicate that the date of tiller emergence bears a negative correlation to earliness.

In the F_1 hybrids tiller emergence was very irregular when compared with the parental lines. From a cross between the two late lines C.I. 7155 and C.I. 7248, the first tiller emerged twelve days later than those for the parents while in the late

Montcalm x C.I. 7155 the first tiller was intermediate in emergence to those of the parents. These wide differences in emergence appeared in the F_1 greenhouse study. In the field the F_2 material showed greater uniformity in tiller emergence. Here the hybrid behaviour was closer to the early parent and consistent throughout several crosses.

Not only did the later varieties require more days for tillering but when crossed with early varieties the hybrid tiller was always earlier than the late parent.

6. Observations on Growth Habit

Aberg and Wiebe (2) observed that in the early stages there were three distinct habits of growth both for spring and winter varieties of barley. They stress that the rosette habit was characteristic of the winter varieties, but that there was no sharp line of demarcation in this respect between certain of the winter and spring varieties.

In most varieties, winter or spring, the rosette-growth stage occurred between seeding and culm elongation, and a plant in this stage was characterized by having several tillers, very little elongation in stem internodes and a leafy appearance.

During the early stages of growth the leaves and tillers are held at various angles which may be described as prostrate, semiprostrate, and erect.

In the spring varieties used for this earliness study the three types of growth habit were evident when observations were taken at the six- to eight-leaf stage. The association between earliness and habit of growth is shown in Plate 1. Seeded in the field on June 1, 1950, the three varieties, Beecher, Titan, and Montcalm, headed on widely different dates. Beecher, the most erect in form with three tillers, headed on July 9. Titan was more spreading with five tillers and headed on July 13. Montcalm was semiprostrate to prostrate with eight tillers and headed on July 19. Other early, medium, and late lines observed showed a similar correlation between growth habit and earliness.

Suneson (29) studied the survival of four barley varieties in a mixture. He observed that the variety Atlas, though the least resistant to various leaf diseases, showed the largest survival value because of its erect, early growth, and weak tillering habit. His observation is supported by this phase of the study carried out in the greenhouse.

A count was taken of the total number of tillers in each variety and hybrid. The average for each population was used to establish a correlation between the number of tillers and days to heading. Table 3 gives a list of all the lines, varieties, and hybrids observed and the average number of tillers in each population.

A

B

C

Plate 1. Barley growth types, seeded June 1, 1950

A - Montcalm, semiprostrate; B - Titan, semierect;
C - Beecher, erect.

Table 3

Tillers Observed

No.	Variety or hybrid	Season	Av. no. of tillers (F ₁)
1	Atsel	Early	2
2	Beecher	Early	3
3	Tulare	Early	2
4	H-44-1	Early to medium	4
5	Titan	Medium	5
6	Newal	Medium	5
7	Montcalm	Late	6
8	C.I. 7155	Late	7
9	C.I. 7248	Late	6
1x	Atsel x Tulare		3
2x	Montcalm x Tulare		4
3x	Montcalm x Beecher		4
4x	Atsel x Beecher		5
5x	Titan x Newal		5
6x	Montcalm x Atsel		4
7x	Montcalm x Titan		4
8x	Montcalm x Newal		5
9x	Beecher x C.I. 7248		5
10x	Titan x Tulare		6
11x	Titan x C.I. 7248		4
12x	Montcalm x C.I. 7155		6
13x	Atsel x C.I. 7155		4
14x	Atsel x C.I. 7248		5
15x	Montcalm x C.I. 7248		5
16x	C.I. 7155 x C.I. 7248		7

The F_2 generation, grown in the field showed a marked segregation for earliness. The correlation between number of tillers and days to heading was however maintained. Fig. 2 illustrates the tiller-days-to-heading relation as studied from F_1 greenhouse material.

The relationship between number of tillers and number of days to heading is shown in Table 4. It must be mentioned that under greenhouse conditions the varieties studied came to head an average of two weeks later than these same varieties did in the field. The greenhouse F_1 hybrids showed similar behaviour to the pure lines but, with the F_2 segregation for earliness in the field this difference can be explained.

Table 4
Tillers and Days to Heading

No.	Material	Tillers (F_1 av.)	Season	Days to head
			Early(E) Midseason.(M) Late(L)	
1	Atsel	2	E	48
2	Beecher	3	E	52
3	Tulare	2	E	49
4	H-44-1	4	E to M	54
5	Titan	5	M	71
6	Newal	5	M	73
7	Montcalm	6	L	74
8	C.I. 7155	7	L	108
9	C.I. 7248	6	L	99

Fig. 2

The number of days to heading increases with the number of tillers.

It was shown that under favourable growing conditions the form or habit of growth, in the early growth stage, could be used as a means of distinguishing between early, medium and late lines. This observation applies where spring varieties of barley are being considered.

Varieties are known to differ in certain leaf characters, such as blade, leaf sheath, auricle, ligule, colour, and size of leaf. These characters serve to separate a few varieties but were not observed to bear any definite relation to earliness. It was seen that, as a general rule, the leaves of early lines were shorter and narrower than the late varieties. Fig. 2 illustrates leaf-width differences which may be associated with earliness. The writer feels that environment may be a factor in determining the width of leaves. This character will not be advanced as a satisfactory criterion of earliness on the basis of the present study.

7. Observations on Leaf-blade Curling

The barley leaf-blade is lanceolate-linear with a prominent middle nerve flanked by 10 or 12 parallel side nerves that are less strongly developed. The blades are rolled in the bud but there is no consistence in the direction of roll.

It was observed that in the earlier growth stage up to the fifth or sixth leaf, the tendency for emergent leaves to remain partially rolled was stronger in the late varieties. This characteristic was first noted in the greenhouse study and was further verified in the field material. Fig. 3 shows typical third leaves taken from early, medium, and late varieties after all had acquired the sixth leaf. The earliest variety, Beecher, has a third leaf which is stiff in character with the rolling or curling almost completely lacking. The midseason variety, Newal, is rolled close to the tip while the late line, Montcalm, maintains the roll or curl throughout the length of the leaf. In Montcalm, except for the older leaves, the first and second also appeared curled.

The varieties Atsel and Tulare, both early, are similar in leaf-rolling characteristic to the early Beecher. Other late varieties studied, C.I. 7155 and C.I. 7248 had rolled leaves even during the later stages of growth though to a lesser extent than Montcalm. This leaf study points to the possibility of an association between stiffness of leaf-blade in the early growth stage and earliness. The hybrid material was less definite in this leaf characteristic, but early F_2 segregates showed a strong tendency for leaf stiffness.

Fig. 3

Degree of leaf curling in three seasonal types of barley:
I - Atsel, early; II - Newal, midseason; III - Montcalm,
late.

In the field the F_2 generation from the cross Montcalm x Tulare resulted in^a wide variety of individuals. In earliness they ranged from plants earlier than the early Tulare to plants later than the late parent. A wide variation in morphology and leaf-blade character was also observed. Not only were the earlier heading plants more erect in habit of growth, but leaf-blade curling was significantly absent.

8. Study of Spike Characteristics

The spike characters of the barley plant are among the most useful in identifying varieties. Some of these characters are stable while others are affected by changes in environment. In some varieties the distance from flag leaf to spike varies from one locality to another as in Titan.

There are several varieties in which the spikes hardly emerge from the boot as in C.I. 7155 and C.I. 7248 and these cases can be easily distinguished from those in which the spike is fully exerted and the distance from flag leaf to spike is

great, as in Atsel and Tulare. The varieties Montcalm and Newal appear to fall in an intermediate group.

In this study there was no variation in the distance from flag leaf to spike in the varieties C.I. 7155 and C.I. 7248. With these two late lines the spike emerged half way up the height of the boot, and the distance of the neck was almost negligible. In these two varieties the flag leaf appeared at the uppermost half to completely above the spike after spike emergence was complete. About 5 per cent of the plants in the variety C.I. 7155 showed spikes which were only partly emerged from the sheaths. These heads contained several sterile spikelets.

Fig. 4 illustrates the spike-flag leaf relationship in the early variety, Atsel, and the late C.I. 7248.

In Atsel, the distance from flag leaf to spike is $1/3$ the length of the spike. This character is typical of the early Atsel and variations in this distance are only slight. C.I. 7248 carries the flag leaf $2/3$ the distance above from the base of the spike. The F_2 generation of Atsel x C.I. 7248 carries the spike in an intermediate position. Here there is a coincidence between base of flag leaf and base of spike.

Fig. 4

Spike-flag leaf relation in the early Atsel, late C.I. 7248, and the F_2 hybrid of cross, Atsel x C.I. 7248.

The head of the F_2 hybrid is intermediate in size between the parents and the distance from last leaf to flag leaf, the boot, is greater than either parent. The neck is inconspicuous above the boot. Although Montcalm, a late variety, shows some variation in spike position, it is significant that 72 per cent of the plants studied carried the spike in the intermediate position while the rest varied from short neck to sub-flag leaf spike position.

On the basis of this study, it is evident that there is a correlation between size of spike, length of neck, and earliness. The varieties Beecher and Tulare, both early, are similar to Atsel in the spike-flag leaf relationship while the late Montcalm, with small variations, C.I. 7153 and C.I. 7248, have a typical short neck and low spike conformation. Table 5 gives data on heading dates of the early Atsel, late C.I. 7153 and C.I. 7248, and the F_2 hybrids between early and late types all seeded in the field on June 1, 1950.

Table 5

Heading Data on Early and Late Lines and F_2 Hybrids

Seeded June 1, 1950

Material	Season	Days to heading
Atsel	Early	36
C.I. 7155	Late	55
C.I. 7248	Late	50
Hybrids (F_2):		
Atsel x C.I. 7155	--	39
Atsel x C.I. 7248	--	41
C.I. 7155 x C.I. 7248	--	49

A comparison of Fig. 4 with data on heading dates in Table 5 serves to establish the correlation between the morphology of the barley head and earliness in terms of time of heading.

In connection with the morphology of earliness a study was made of node characters. It was observed that in the early varieties, Atsel, Tulare, and Beecher, and to a lesser extent H-44-1, the nodes were exposed to the sun's rays earlier than in the midseason lines, Newal or Titan, and the late varieties, Montcalm, C.I. 7155, and C.I. 7248.

1. 2. 3.

Table 1. Results of the survey.		
No. of respondents		
1	2	3
4	5	6
7	8	9
10	11	12
13	14	15
16	17	18
19	20	21
22	23	24
25	26	27
28	29	30
31	32	33
34	35	36
37	38	39
40	41	42
43	44	45
46	47	48
49	50	51
52	53	54
55	56	57
58	59	60
61	62	63
64	65	66
67	68	69
70	71	72
73	74	75
76	77	78
79	80	81
82	83	84
85	86	87
88	89	90
91	92	93
94	95	96
97	98	99
100	101	102
103	104	105
106	107	108
109	110	111
112	113	114
115	116	117
118	119	120
121	122	123
124	125	126
127	128	129
130	131	132
133	134	135
136	137	138
139	140	141
142	143	144
145	146	147
148	149	150
151	152	153
154	155	156
157	158	159
160	161	162
163	164	165
166	167	168
169	170	171
172	173	174
175	176	177
178	179	180
181	182	183
184	185	186
187	188	189
190	191	192
193	194	195
196	197	198
199	200	201
202	203	204
205	206	207
208	209	210
211	212	213
214	215	216
217	218	219
220	221	222
223	224	225
226	227	228
229	230	231
232	233	234
235	236	237
238	239	240
241	242	243
244	245	246
247	248	249
250	251	252
253	254	255
256	257	258
259	260	261
262	263	264
265	266	267
268	269	270
271	272	273
274	275	276
277	278	279
280	281	282
283	284	285
286	287	288
289	290	291
292	293	294
295	296	297
298	299	300
301	302	303
304	305	306
307	308	309
310	311	312
313	314	315
316	317	318
319	320	321
322	323	324
325	326	327
328	329	330
331	332	333
334	335	336
337	338	339
340	341	342
343	344	345
346	347	348
349	350	351
352	353	354
355	356	357
358	359	360
361	362	363
364	365	366
367	368	369
370	371	372
373	374	375
376	377	378
379	380	381
382	383	384
385	386	387
388	389	390
391	392	393
394	395	396
397	398	399
400	401	402
403	404	405
406	407	408
409	410	411
412	413	414
415	416	417
418	419	420
421	422	423
424	425	426
427	428	429
430	431	432
433	434	435
436	437	438
439	440	441
442	443	444
445	446	447
448	449	450
451	452	453
454	455	456
457	458	459
460	461	462
463	464	465
466	467	468
469	470	471
472	473	474
475	476	477
478	479	480
481	482	483
484	485	486
487	488	489
490	491	492
493	494	495
496	497	498
499	500	501
502	503	504
505	506	507
508	509	510
511	512	513
514	515	516
517	518	519
520	521	522
523	524	525
526	527	528
529	530	531
532	533	534
535	536	537
538	539	540
541	542	543
544	545	546
547	548	549
550	551	552
553	554	555
556	557	558
559	560	561
562	563	564
565	566	567
568	569	570
571	572	573
574	575	576
577	578	579
580	581	582
583	584	585
586	587	588
589	590	591
592	593	594
595	596	597
598	599	600
601	602	603
604	605	606
607	608	609
610	611	612
613	614	615
616	617	618
619	620	621
622	623	624
625	626	627
628	629	630
631	632	633
634	635	636
637	638	639
640	641	642
643	644	645
646	647	648
649	650	651
652	653	654
655	656	657
658	659	660
661	662	663
664	665	666
667	668	669
670	671	672
673	674	675
676	677	678
679	680	681
682	683	684
685	686	687
688	689	690
691	692	693
694	695	696
697	698	699
700	701	702
703	704	705
706	707	708
709	710	711
712	713	714
715	716	717
718	719	720
721	722	723
724	725	726
727	728	729
730	731	732
733	734	735
736	737	738
739	740	741
742	743	744
745	746	747
748	749	750
751	752	753
754	755	756
757	758	759
760	761	762
763	764	765
766	767	768
769	770	771
772	773	774
775	776	777
778	779	780
781	782	783
784	785	786
787	788	789
790	791	792
793	794	795
796	797	798
799	800	801
802	803	804
805	806	807
808	809	810
811	812	813
814	815	816
817	818	819
820	821	822
823	824	825
826	827	828
829	830	831
832	833	834
835	836	837
838	839	840
841	842	843
844	845	846
847	848	849
850	851	852
853	854	855
856	857	858
859	860	861
862	863	864
865	866	867
868	869	870
871	872	873
874	875	876
877	878	879
880	881	882
883	884	885
886	887	888
889	890	891
892	893	894
895	896	897
898	899	900
901	902	903
904	905	906
907	908	909
910	911	912
913	914	915
916	917	918
919	920	921
922	923	924
925	926	927
928	929	930
931	932	933
934	935	936
937	938	939
940	941	942
943	944	945
946	947	948
949	950	951
952	953	954
955	956	957
958	959	960
961	962	963
964	965	966
967	968	969
970	971	972
973	974	975
976	977	978
979	980	981
982	983	984
985	986	987
988	989	990
991	992	993
994	995	996
997	998	999
1000	1001	1002

After two weeks of exposure to sunshine, the nodes took on a dark orange to purple colour due to the presence of anthocyanin. While the early node exposure in this study seems to be associated with earliness, the variation in node colour introduces a factor of uncertainty as to how much of the colour is due to environment and how much to inheritance. The unreliability of colour readings makes it difficult to use this character as a morphological expression of earliness.

A number of other plant characters were studied but were found, within the limits of this study, to be of little value in classifying varieties as to earliness. In some cases the varieties studied were so nearly alike that they could not be differentiated on the basis of these characters. In other cases the observations had to be abandoned because of variability obviously due to environment.

The lengths of spikes was studied but a great deal of variation was found between varieties and within the same variety.

In the greenhouse study no plant among the early lines Atsel, Beecher, and Tulare, was observed to carry more than 8 leaves on the main stem. The midseason variety Titan and the late Montcalm averaged 9 and 10 leaves respectively. This relationship did not hold completely in the field observation where Tulare again averaged 8 leaves per main stem but Atsel and Beecher carried 9 leaves on more than 60 per cent of the population. Here several Montcalm plants carried 11 leaves.

The height of plant, that is, the distance from the ground to the base of the spike as used by Åberg and Wiebe (2) was found to be an unstable character, and could not be used in predicting earliness. The influence of environment on plant height within varieties will be dealt with in the report on the physiological study.

IV. PHYSIOLOGICAL STUDY

1. Observations on Development of Barley

Both greenhouse and field studies indicate that there is a great deal of variation in the time of heading of barley. The rate of development and time of heading are influenced by weather and soil conditions and by soil differences. Ideal conditions for such a study would include uniform soil and unvarying conditions for growth throughout the growing season. In the field this ideal is practically unattainable.

The studies referred to on the influence of length of day on heading (10), (7), (11), and (12), are related to the present investigations of the factors influencing earliness in barley and other cereal crops. Photoperiodism is probably responsible for the irregular heading of the late varieties and for the failure of some plants, and of crosses with them, to head at all.

2. Materials and Methods

Seedlings of three seasonal groups of barley (early, midseason, and late) were grown in flats in three sections of the greenhouse at three different temperatures, 55°, 65° and 75°F. At each temperature a short and long day (13 and 17 hours) was simulated by light control - electric lamps and shading.

The varieties with date of seeding and dates of emergence under reasonably constant photoperiods and temperatures are given in Table 6.

Table 6

Materials and Conditions of Photoperiodic Study in
Greenhouse, with Days to Emergence from Seeding on
November 15, 1949

Material	Season	<u>Photoperiod 17-hr.</u>			<u>Photoperiod 13-hr.</u>		
		<u>Days to emergence</u>	<u>Days to emergence</u>	<u>Days to emergence</u>	<u>Days to emergence</u>	<u>Days to emergence</u>	<u>Days to emergence</u>
		75°F	65°F	55°F	75°F	65°F	55°F
Atsel	E	3	3	4	4	4	5
Beecher	E	4	4	5	4	5	6
Tulare	E	3	4	4	4	4	5
H-44-1	M	4	5	5	4	5	6
Newal	M	4	5	5	5	6	6
Titan	M	4	4	5	4	5	6
C.I. 7155	L	6	6	7	5	6	7
C.I. 7248	L	5	5	6	5	6	7
Montcalm	L	5	5	6	5	6	7

This Table shows that by adjusting factors in the environment, the developmental response, even in the early stages, can be increased or retarded. For example, the early variety Atsel showed an emergence range of three days when grown under photoperiods and temperatures differing by 4 hours and 20°F respectively. The latest line, C.I. 7155, showed a similar response under the same set of conditions, thus emphasizing the dependence of rate of development upon the quantity of light and temperature.

3. Histological Observations

One seedling from each variety or hybrid and from each treatment was collected for histological study at different intervals. The following technique was employed for collecting and storing material:

Seedlings were carefully taken up in order to preserve the radicle and plumule. They were then washed and killed by immersing for a 48-hour period in Farmer's mixture - 3 parts of 95 per cent ethanol to 1 part glacial acetic acid. The killing and fixing process was followed by washing and storage in 70 per cent ethanol. Collections at various intervals were identified by date tags.

Photographs were made of three sets of plants, early Atsel, midseason Newal, and late Montcalm, taken up on the same date to show the difference in development of plants under specific photoperiods and temperatures. The correlation between rate of development, photoperiod, and temperature is shown in Figs. 5 and 6. The plants in both Figures were taken up on the same date, November 27, 1949. The labels A, B, and C, represent temperatures 75°, 65°, and 55°F. The varieties are grouped to show two photoperiods, 17 hours and 13 hours.

In both Figs. 5 and 6, the difference in development between plants A and B under each photoperiod is not very significant, though the size of the second leaf in each case indicates that in the early stages, a temperature of 75°F favours development. For each variety plant C is significantly later than A or B, though grown under the same photoperiod. It is evident that the temperature 55°F is below the ideal for maximum growth in the early stages.

When plant C is considered there appears to be no visible difference between the effects of the two photoperiods. Temperature appears to be the more important factor in development at this stage of growth. It would appear, further, that the ideal temperature lies between 75° and 65°F and closer to 75°F. Under a photoperiod of 17 hours plants A and B show an advance in development over A and B under a 13-hour day. The indication is that the effect of photoperiod upon rate of growth is largely dependent upon temperature.

Fig. 5

Seven day old Atsel seedlings grown under two photoperiods, 17 hours and 13 hours, and three temperatures, A - 75°F; B - 65°F; C - 55°F.

Fig. 6

Seven day old Montcalm seedlings grown under two photoperiods, 17 hours and 13 hours, and three temperatures, A - 75°F; B - 65°F; C - 55°F.

Earliness, having been based on the initiation, development, and final emergence of the primordia in the form of the head, all plants were dissected and measurements of leaves and internodes made. This was followed by examination and measurement of the primordia. If at one stage of development it could be established that the primordia was not present, and at a later growth stage it had been initiated, then the critical date in the primordia initiation could be used as significant of earliness.

Table 7 shows data on leaf measurements in cm. of the first set of plants taken November 24, 1949. By dissection, the initiation of a third or a fourth leaf was observed in some plants, but in none was a primordia seen under the dissecting microscope. It was obviously too early a stage for the detection of differentiation of the growing point. Nodes and internodes were not discernible at this early growth stage.

Seedlings taken on November 27, 1949, samples of which are represented in Figs. 6 and 7, were dissected under the microscope. These were three days older than seedlings recorded in Table 7 and show in some cases the initiation of a fifth leaf and the presence of the primordia. Data on plants taken November 27, 1949, are given in Table 8.

Table 7

Measurement of Leaves in Cm. of Plants Seeded November 15, 1949
and Taken up November 24, 1949

Material	A (75°F)				B (65°F)				C (55°F)			
					Leaf							
	1	2	3	4	1	2	3	4	1	2	3	4
<u>Period of daily exposure - 17 hours</u>												
Atsel	12.7	5.1	0.5	0.1	12.1	4.3	0.4	0.5	6.9	1.5	0.25	-
Beecher	11.7	4.1	0.5	1.15	13.3	3.9	0.3	0.05	4.7	0.9	0.2	-
Tulare	11.4	2.6	0.4	0.05	10.4	1.9	0.25	-	4.4	0.8	0.1	-
H-44-1	10.8	9.2	1.0	0.15	13.0	9.1	1.1	0.2	3.9	1.1	0.2	-
Newal	14.2	7.3	0.6	0.1	10.9	3.6	0.4	0.05	3.8	0.7	0.1	-
Titan	14.2	6.7	0.5	0.05	14.6	6.5	0.6	0.1	5.6	1.1	0.15	-
C.I.7155	11.5	3.4	0.45	0.05	11.7	1.95	0.45	0.05	4.2	0.5	0.05	-
C.I.7248	12.4	5.1	0.35	0.02	12.6	4.7	0.5	0.05	5.1	0.7	0.1	-
Montcalm	13.4	6.1	0.7	0.1	12.1	4.6	0.4	-	4.05	0.6	0.05	-
<u>Period of daily exposure - 13 hours</u>												
Atsel	13.5	7.6	0.6	0.15	14.1	7.2	0.4	0.1	5.5	0.9	0.25	-
Beecher	14.8	4.2	0.4	0.1	14.2	5.5	0.2	0.05	5.8	0.7	0.15	-
Tulare	11.3	6.3	0.5	0.1	10.7	3.3	0.15	-	5.2	0.4	0.05	-
H-44-1	9.5	4.0	0.4	0.1	10.4	7.3	0.3	0.05	8.2	2.7	0.35	0.05
Newal	12.1	5.8	0.4	0.05	13.1	5.8	0.2	-	5.5	0.9	0.15	-
Titan	14.6	5.3	0.35	0.05	11.4	4.8	0.25	-	6.2	0.95	0.15	-
C.I.7155	11.7	2.3	0.4	0.03	12.3	3.0	0.1	-	6.0	0.5	0.05	-
C.I.7248	12.2	5.8	0.3	0.05	12.0	4.6	0.2	-	5.6	0.85	0.1	-
Montcalm	11.3	7.4	0.5	0.05	14.0	5.2	0.3	0.05	5.7	1.0	0.15	-

天

— 23 —

४ ३

• •

(Group) (Individual)

10

• •

Table 8

Measurement of Leaves and Primordia in Cm. of Plants Taken up

November 27, 1949

Material	75°F				65°F				55°F			
					Leaf							
	1	3	5	Prim.	1	3	5	Prim.	1	3	5	Prim.
<u>Period of daily exposure - 17 hours</u>												
Atsel	13.3	5.3	0.1	0.08	19.4	1.70	0.09	0.05	9.6	0.5	-	-
Beecher	15.8	4.8	0.05	-	13.8	1.1	0.03	-	8.3	0.4	-	-
Tulare	13.4	1.5	0.05	0.04	14.7	1.4	0.07	0.03	9.1	0.3	-	-
H-44-1	12.1	3.4	0.02	-	13.4	4.8	0.1	-	7.2	0.25	-	-
Newal	13.1	1.4	0.05	-	13.7	3.0	0.1	-	8.0	0.2	-	-
Titan	14.6	4.8	0.07	0.03	15.5	1.5	0.2	0.02	9.9	0.2	-	-
C.I.7155	14.1	0.3	-	-	14.3	1.3	0.01	-	8.7	0.15	-	-
C.I.7248	13.1	1.0	0.02	-	13.6	0.8	0.02	-	8.4	0.25	-	-
Montcalm	14.3	1.7	0.04	0.05	16.7	1.3	0.03	0.02	9.8	0.2	-	-
<u>Period of daily exposure - 13 hours</u>												
Atsel	14.1	1.7	0.1	0.03	15.1	1.9	0.03	0.02	9.0	0.4	-	-
Beecher	13.9	1.4	0.05	0.02	15.3	1.4	0.01	-	9.9	0.3	-	-
Tulare	13.5	0.7	0.02	-	13.9	1.5	0.01	-	7.9	0.2	-	-
H-44-1	12.9	1.1	0.03	-	13.5	1.9	0.03	0.01	10.1	0.7	0.01	-
Newal	13.3	0.5	-	-	13.7	1.2	-	-	8.3	0.25	-	-
Titan	13.4	0.7	-	-	14.7	1.3	0.02	-	8.1	0.2	-	-
C.I.7155	14.9	0.25	-	-	14.1	1.2	-	-	7.1	0.15	-	-
C.I.7248	14.5	0.35	-	-	13.9	1.0	0.01	-	6.8	0.15	-	-
Montcalm	15.2	1.45	0.03	0.02	14.6	1.1	0.05	-	6.2	0.8	-	-

From Table 8 it may be observed that under a photoperiod of 17 hours and temperatures 75° and 65°F , the primordia of the Atsel variety was initiated in 8 days after emergence. Under the same set of conditions (Table 7) it was not detected when the variety was 5 days old. Table 8 also indicates that primordia differentiation takes place only after the initiation of the fifth leaf.

The measurements for plants grown at 65°F show that this temperature is closer to the ideal for vegetative growth than 75°F , but the latter is more favourable to development. Under a temperature of 55°F there was a consistent lag in leaf emergence and primordia differentiation with all the varieties observed. The longer daily exposure did not favour development at 55°F .

Photomicrographs of growing points and primordia differentiation of plants in Figs. 5 and 6 are shown in Plates 2 and 3.

The top row of Plate 2, reading from left to right shows the enlarged growing points of the early variety Atsel grown under a photoperiod of 17 hours and temperatures of 75° , 65° , and 55°F . The bottom row shows the late Montcalm grown under the same conditions. It may be observed that in both varieties, response to temperature, in terms of growing point elongation, is significantly higher for temperatures 75° and 65°F .

Atsel

A

B

C

Montcalm

A

B

C

Plate 2. Effect of 17 hour photoperiod on primordia development of seven day old Atsel and Montcalm seedlings grown under three temperatures, A - 75°F; B - 65°F; C - 55°F.

Atsel

A

B

C

Montcalm

A

B

C

Plate 3. Effect of 13 hour photoperiod on primordia development of seven day old Atsel and Montcalm seedlings grown under three temperatures, A - 75°F; B - 65°F; C - 55°F.

In Plate 3 is shown the relative response of Atsel, in the top row, and Montcalm when plants are grown under a day length of 13 hours and different temperatures. As in Plate 1, development is more rapid under 65° and 75° than under 55°F. The day length of 17 hours (see Plate 2) shows a small increase in rate of development over the 13-hour daily exposure to light (Plate 3).

One sample from each variety grown under each of the six conditions previously mentioned was collected for primordia study on the following dates:

November 24, 1949

November 27, 1949

December 1, 1949

December 7, 1949

December 14, 1949

December 21, 1949

December 28, 1949

Each set of samples was studied in the same way as in the method described - microdissection followed by microphotography of the primordia. The fourth set of plants taken on December 7, 1949, and all later collections, were studied for leaf and primordia and in addition for tiller and internode development. It was established in the morphological study that the average number of tillers had a direct relation to earliness, and so the rate at which tillers developed should be correlated with the rate of primordia development.

4. Photoperiodic Reaction with Advancing Age of Plant

It was observed that in the early growth stage the effect of a longer number of light hours per day was not significant. Development under the shorter period of light compared favourably with the longer, provided the temperature was the same. However, with an advance in the age of plants towards the 'ripe-to-flower' condition, the effect of photoperiod on growth and differentiation becomes more evident.

The results of differences in the daily illumination are given in Plate 4 with plants collected December 14, 1949.

The top row of Plate 4 shows early Atsel and late Montcalm plants grown under a 17-hour daylight and three temperatures- A, 75°, B, 65°, and C, 55°F. With both varieties, 65°F favoured leaf and stem elongation as well as primordia development (see Plate 5). At the lowest temperature, 55°F both varieties showed delayed or less profuse growth.

In the bottom row of the same figure are shown plants of the same two varieties of the same age and grown under the same temperatures but with 13-hour illumination. Under a 13-hour photoperiod the effect of higher temperatures on vegetative growth is less pronounced. The difference in plant structure at 75°F and 65°F is negligible in both the Atsel and Montcalm varieties. These contrasting behaviours at the present time appear to be dependent upon

Plate 4. Twenty-five day old Atsel and Montcalm seedlings grown under 17 hour and 13 hour photoperiods and temperatures: A - 75°F; B - 65°F; C - 55°F.

fundamental differences in the physiology of the two photoperiodic groups, and may be so regarded until further investigations have proved otherwise.

The measurements for leaves, tillers, internodes and primordia of plants collected on December 14, 1949, are reported in Table 9. It will be seen that the rate of development of the primordia is correlated with the length of the daily exposure. All plants under the 13-hour day length showed less development in this character than the corresponding varieties under the longer photoperiod.

The data on tillers (Table 9) indicate that the later varieties tend to have a more rapid growth and development of the tillers. The contrast in tiller growth under the two photoperiods shows that this character is also affected by length of exposure to light. The elongation of the internodes is similarly affected by exposure to light.

Photomicrographs showing stages of primordia development of plants in Plate 4 are given in Plates 5 and 6. Plate 5 shows primordia development under a 17-hour photoperiod and three temperatures. At temperatures 75° and 65°F the primordia show an increase in growth over the 55°F. The early Atsel and late Montcalm respond in a similar way to differences in temperature. In both varieties the primordia was smaller when the plant was grown under the same photoperiod of 17 hours and at a temperature of 55°F.

Table 9

Measurements in Cm. of Plants Collected November 14, 1949

Material	75°F								65°F								55°F									
	Leaf			Tiller		Internode			Prim.	Leaf			Tiller		Internode		Prim.	Leaf			Tiller		Internode			Prim.
	1	4	7	1	2	2	3	1		4	7	1	2	2	3	1		4	7	1	2	2	3			
Period of daily exposure - 17 hours																										
Atsel	11.4	26.7	2.7	1.0	0.2	1.5	0.5	0.4	15.1	33.2	3.5	0.2	0.3	2.4	0.4	0.4	15.7	15.0	0.7	0.2	0.15	0.2	0.1	0.2		
Beecher	15.2	37.2	1.1	9.1	0.7	0.3	0.15	0.3	13.2	29.8	0.1	0.15	0.1	0.2	0.1	0.2	11.1	10.9	0.15	0.1	0.5	0.15	-	0.15		
Tulare	13.9	19.8	0.9	0.6	0.4	0.3	0.15	0.35	7.7	19.0	0.12	0.15	0.1	0.17	0.1	0.27	11.2	6.6	0.15	0.12	0.1	0.2	0.15	0.2		
E-44-1	13.1	27.6	2.1	0.5	0.2	1.5	0.15	0.3	11.9	34.3	5.9	0.1	0.5	1.7	1.9	0.3	12.8	14.7	0.3	0.2	0.1	1.15	0.1	0.2		
Newal	14.1	34.8	1.7	3.4	0.7	0.8	0.15	0.2	13.2	27.7	0.5	0.25	0.1	0.3	0.1	0.2	14.6	12.2	0.15	0.2	0.1	0.3	0.1	0.15		
Titan	15.5	36.4	0.9	15.4	4.5	0.35	0.15	0.2	12.1	27.4	0.2	0.3	0.2	0.15	0.1	0.2	11.8	11.3	-	4.8	0.25	0.15	0.1	0.12		
C.I.7155	14.3	25.2	0.5	1.4	0.3	0.25	0.1	0.2	15.5	2.8	0.1	0.1	0.2	0.05	0.01	0.10	15.2	1.4	0.05	0.3	0.2	0.05	-	0.17		
C.I.7248	15.4	26.2	0.8	13.6	0.6	0.3	0.2	0.15	16.4	18.2	0.3	0.2	0.1	0.15	0.02	0.18	13.2	6.7	-	0.9	0.2	0.15	0.1	0.15		
Montcalm	14.7	28.7	0.6	10.0	2.6	0.15	0.1	0.15	13.8	36.5	0.7	0.15	0.1	0.1	0.07	0.25	15.2	15.1	0.15	0.1	0.5	0.17	0.1	0.16		
Period of daily exposure - 13 hours																										
Atsel	12.6	0.8	0.1	0.15	-	0.12	0.05	0.16	18.1	1.15	0.2	0.1	0.15	0.08	0.05	0.17	15.3	1.9	-	0.05	-	0.12	0.05	0.12		
Beecher	16.9	6.5	-	0.1	-	0.1	-	0.15	15.2	2.3	-	0.1	0.07	0.07	-	0.15	15.5	2.0	-	0.1	0.05	0.1	-	0.12		
Tulare	13.4	10.2	-	0.1	0.05	0.15	0.1	0.12	6.4	2.2	-	0.2	0.1	0.1	0.03	0.17	13.4	0.3	-	0.1	0.05	0.08	-	0.1		
E-44-1	11.5	6.4	0.4	0.2	-	0.1	-	0.75	9.1	7.0	-	0.1	0.07	0.05	0.02	0.14	15.2	0.6	-	0.08	-	0.08	-	0.15		
Newal	13.6	3.8	-	0.2	0.1	0.1	0.05	0.15	15.6	1.8	-	0.08	0.05	0.05	0.02	0.12	13.2	0.6	0.05	0.1	0.4	0.3	0.12	0.1		
Titan	16.0	10.1	0.1	0.15	0.1	0.15	-	0.12	15.2	4.0	0.1	0.1	0.05	0.08	0.02	0.1	13.3	1.2	-	0.1	0.2	0.05	-	0.1		
C.I.7155	14.4	1.1	-	0.1	-	0.1	-	0.1	12.7	0.49	-	0.05	0.02	0.05	-	0.02	19.3	1.07	-	0.5	-	0.02	-	0.05		
C.I.7248	14.3	0.6	-	0.2	-	0.05	-	0.15	15.4	1.7	-	0.05	-	0.05	-	0.03	14.6	1.9	-	0.08	-	0.15	0.2	0.02		
Montcalm	16.2	2.7	-	0.15	0.1	0.12	0.05	0.12	12.2	0.45	0.01	0.7	0.5	0.03	-	0.02	13.8	1.2	0.05	0.1	0.2	0.1	-	0.08		

In Plate 6 it may be observed that the highest temperature of 75°F and the lowest of 55°F does not make a significant difference in primordia size or development. The two contrasting varieties Atsel and Montcalm showed little variation in primordia size over the range of temperature 75° to 55°F. It is evident from Plate 5 that under the 17-hour day length the plants have approached the ripe-to-flower stage. Plate 6 shows that under the shorter day length of 13 hours this condition is not yet reached.

The contrasting behaviour of plants under the two photoperiods described above seems to indicate that in the late developmental stages a longer exposure to light is essential for early heading in barley. It may therefore be suggested that the plants have now reached a phase in which light becomes the decisive factor for earliness, accompanied by accessory factors such as temperature, humidity, etc.

5. Photoperiodism and Heading

Having discussed and illustrated the influence of photoperiodism and other accessory factors in the rate of development in the early and middle stages of growth, it will now be appropriate to give the observations made at the later and pre-inflorescence stage.

For this study of the physiology of earliness the last set of plants was 44 days old when collected. The comparative sizes

Atsel

A

B

C

Montcalm

A

B

C

Plate 5. Effect of 17-hour photoperiod on primordia development of twenty-five day old Atsel and Montcalm seedlings grown under three temperatures, A - 75°F; B - 65°F; C - 55°F.

Atsel

Montcalm

Plate 6. Effect of 13-hour photoperiod on primordia development of twenty-five day old Atsel and Montcalm seedlings grown under three temperatures, A - 75°F; B - 65°F; C - 55°F.

attained by the two varieties Atsel and Montcalm under the various conditions of light exposure and temperature are shown in Plate 7.

The upper row, Plate 7, shows Atsel and Montcalm varieties grown under a light exposure of 17 hours per day and temperatures of 75° , 65° , and 55° F reading from left to right. Unlike the behaviour observed when the varieties were two weeks younger (see Plate 4) the two lines now give a greater differential response to temperatures 75° F and 65° F. The A plant (75° F) of the late Montcalm variety has developed significantly beyond the B plant (65° F) of the same variety. In the early Atsel line, the A plant has similarly advanced beyond the B which in both cases is much closer to the C plant (55° F).

A comparison of the A plant shows that Atsel has advanced towards heading beyond Montcalm, although both plants were the same age and grown under the same environment. On the basis of this observation, it must be assumed that some other factor or factors beyond the limits of present environmental control has influenced an earlier heading in the Atsel line.

The differences in growth between plants B and C in both the Atsel and Montcalm lines are not as significant as those between A and B. This would indicate that the critical temperature for earliness at this stage of vegetative growth in both varieties was between 65° and 75° F and closer to 75° F. Both varieties have shown greatest

Plate 7. Forty-four day old Atsel and Montcalm seedlings grown under 17-hour and 13-hour photoperiods and temperatures: A - 75°F; B - 65°F; C - 55°F.

response to the 17-hour photoperiod when the temperature is high (75°F). Atsel, however, has a greater capacity to react under higher temperatures and is therefore earlier than Montcalm grown under the same set of conditions.

An examination of the lower row of Plate 7 shows the effects of a shorter daily exposure (13 hours) on growth and development as the plants approach the ripe-to-flower condition. All the plants are characterized by spindly growth and are backward in leaf emergence and stem elongation. The differences in response between temperatures 75° and 65°F are not as clearly defined as under the 17-hour light exposure. Both Atsel and Montcalm show marked retardation at temperature 55°F . When the plants are compared by temperature figures the capacity of Atsel to react under higher temperatures is confirmed.

When the lower row (Plate 7) is contrasted with the upper, the importance of photoperiodism on the rate of development, or earliness, becomes apparent. The necessity for a longer daily exposure to light increases as the inflorescence stage approaches. This differential response to temperature and light was found in the other early, midseason, and late varieties observed. Measurements for the above and other varieties dissected are given in Table 10.

Table 10

Measurements in Cm. of Plants Collected November 28, 1949

Material	75°F									65°F									55°F										
	Leaf			Tiller		Internode				Prim.	Leaf			Tiller		Internode			Prim.	Leaf			Tiller		Internode				Prim.
	1	4	7	1	2	2	4	Prim.	1		4	7	1	2	2	4	Prim.	1		4	7	1	2	2	4	Prim.			
Period of daily exposure - 17 hours																													
Atsel	17.5	41.0	41.2	1.4	0.7	5.8	6.2	5.6	10.4	39.5	20.1	0.3	0.15	1.3	1.4	0.6	14.6	31.2	4.6	0.4	0.2	0.25	0.1	0.3					
Beecher	16.4	48.1	3.1	19.2	0.8	1.4	0.1	0.4	12.8	39.1	18.6	0.2	0.4	5.8	0.4	0.35	15.6	44.7	5.6	0.38	0.2	4.1	-	0.25					
Tulare	10.2	34.7	13.4	0.2	0.15	2.4	0.5	0.5	8.3	28.5	29.7	0.14	0.1	4.0	4.3	1.4	5.2	19.8	-	0.15	0.1	1.1	-	0.3					
H-44-1	5.1	20.1	1.1	0.2	0.1	0.9	0.1	0.3	10.9	34.8	9.7	0.60	0.2	2.1	3.0	0.3	6.4	21.5	-	0.15	0.1	0.5	-	0.15					
Newal	14.0	39.1	3.2	0.4	1.1	2.1	0.15	0.35	14.1	37.4	4.7	0.15	0.2	1.3	0.4	0.45	13.2	20.4	1.4	0.17	-	0.2	0.2	0.2					
Titan	10.0	34.6	6.1	0.4	0.3	0.7	0.1	0.3	12.3	40.1	9.0	0.15	0.12	3.5	-	0.25	14.1	35.2	0.9	0.5	0.2	0.3	-	0.15					
C.I.7155	12.0	37.2	3.2	0.5	0.4	2.2	0.15	0.25	15.5	33.5	11.20	.13	0.14	0.4	-	0.13	16.8	11.3	-	0.15	0.1	0.2	-	0.13					
C.I.7248	14.3	30.5	0.8	0.3	0.25	0.2	0.1	0.15	13.6	39.1	3.1	0.15	0.12	0.4	-	0.2	13.2	21.8	-	0.12	0.1	0.15	0.05	0.14					
Montcalm	15.4	45.0	16.1	0.4	29.1	2.3	0.4	0.4	13.9	41.3	1.90	.15	0.1	0.7	0.1	0.25	10.4	40.3	1.40	0.15	0.2	0.5	0.1	0.2					
Period of daily exposure - 13 hours																													
Atsel	15.6	10.5	0.4	0.4	0.15	0.8	1.4	0.25	12.1	9.7	0.3	0.25	0.1	0.2	0.1	0.18	9.3	12.4	0.4	0.15	-	0.2	0.05	0.15					
Beecher	14.7	10.8	-	0.12	-	0.15	-	0.18	11.8	18.3	0.6	0.07	-	0.2	-	0.2	9.9	13.2	-	0.05	-	0.15	0.05	0.15					
Tulare	11.3	1.6	-	0.1	-	0.15	-	0.13	14.4	17.1	0.8	0.05	0.02	2.6	0.2	0.3	9.5	2.4	-	0.06	-	0.15	-	0.18					
H-44-1	9.2	24.6	0.2	0.13	0.1	0.4	-	0.16	12.3	26.5	-	0.08	-	1.4	0.2	0.2	15.0	13.9	-	0.1	0.12	0.15	-	0.14					
Newal	10.2	25.1	0.25	0.17	0.1	0.3	0.1	0.14	14.1	18.6	0.25	.1	0.1	0.4	0.1	0.18	14.7	15.4	0.2	0.25	0.1	0.2	-	0.12					
Titan	12.9	26.1	0.5	0.15	0.12	.2	-	0.15	16.7	19.4	-	0.07	-	0.12	0.05	0.17	15.4	15.1	0.6	0.35	0.2	0.2	-	0.15					
C.I.7155	16.3	5.8	-	0.1	0.1	0.14	0.04	0.13	14.2	1.6	-	0.12	0.1	0.12	-	0.1	18.6	0.8	-	0.08	0.12	0.13	-	0.1					
C.I.7248	15.1	11.8	-	0.17	0.14	.12	0.1	0.12	16.5	10.7	0.2	0.13	0.1	0.14	-	0.13	12.2	2.5	-	0.13	0.15	0.12	-	0.12					
Montcalm	15.0	6.6	-	0.12	0.2	0.15	0.15	12.3	8.6	0.2	0.04	0.15	0.15	0.02	-	0.12	14.2	2.6	-	0.12	0.1	0.4	-	0.13					

The data on primordia (Table 10) indicates that under short-day conditions development is retarded. It is further shown that temperature is effective in development only when the period of illumination is adequate. Although rates of growth and development differ under the two photoperiods studied, the three seasonal types of plants maintain their seasonal characteristics. Primordia sizes fall into three groups corresponding to the early, midseason, and late types utilized in the study.

Photomicrographs of the primordia formation of plants shown in Plate 7 are given in Plates 8 and 9.

The contrasting behaviour of the early Atsel and late Montcalm at the pre-inflorescence stage is evident when the upper and lower row of Plate 8 are compared. At 75°F and day-length of 17 hours, the awns and spikelet formations for Atsel are differentiated. With Montcalm, under the corresponding temperature, the spikelets are partially differentiated but awn structure has not yet appeared. The Atsel line has made more rapid utilization of the factors in the environment, and is therefore closer to heading.

The relation between temperature reaction at 65° and 55°F and development is obvious. The two varieties have shown response in proportion to temperature, but at each temperature the Atsel line is closer to heading.

Atsel

A

B

C

Montcalm

A

B

C

Plate 8. Effect of 17-hour photoperiod on primordia development of forty-four day old Atsel and Montcalm seedlings grown under three temperatures, A - 75°F; B - 65°F; C - 55°F.

Atsel

A

B

C

Montcalm

A

B

C

Plate 9. Effect of 13-hour photoperiod on primordia development of forty-four day old Atsel and Montcalm seedlings grown under three temperatures, A - 75°F; B - 65°F; C - 55°F.

Plate 9 shows primordia of the Atsel and Montcalm varieties which received a relatively shorter photoperiod (13 hours) each day. For these plants stems and leaves were shorter (see Table 10) and primordia correspondingly short. Differential response to temperature is not as clearly defined as in Plate 8.

Primordia size, under this short-day period, is observed to vary with temperature. Larger growth is seen for higher temperatures, but development towards flowering does not show a significant difference over a range of 55^o to 75^o F. The results are considered to show clearly that, as far as earliness is concerned, both Atsel and Montcalm are adapted to a relatively long day.

6. Observations on Heights of Plants

Height of plant, as used in this study, is the distance from the ground to the base of the spike and is measured in inches.

There was a great deal of variation in heights of varieties as observed in the field. Four replications or dates of seeding were employed. The study proved that height was not a stable character, even within the same variety, but that it was influenced by weather and soil conditions and by soil differences. Ideal conditions for a study of earliness would include uniform soil and unvarying conditions for growth throughout the growing season. This ideal, however, is practically unattainable in the field.

The first seeding in the field was made on May 1, 1950, and consisted of certain varieties and the F_2 hybrids between them. In addition, two varieties, not involved in crosses, were used for height observations. They were the early to midseason line H-44-1, and the variety Floya obtained in Norway specifically for earliness study at the University of Alberta. This variety, relative to other varieties, had a peculiar response in Norway where it originated. Under early, wet and cold spring conditions it was early, but was late in heading when grown under warm, long summer days. The variations in heights for the four replications and their relation to days to heading for varieties and F_2 hybrids are given in Table 11.

An examination of Table 11 shows that with increasing length of day there is a corresponding increase in the mean height of plants. The plants of the first date of seeding are the shortest, and between the first and the second date of seeding there is an average gain in height of 2.25 inches.

Between the second and third date of seeding an average gain in height of 7.3 inches is represented. This is the most significant gain of the four replications, and is due to heavy rains on the 14th and 15th of July. Replication 3, seeded June 1, 1950, had then attained the early heading stage and still had active capacity for vegetative growth.

 

Table 11

Relation Between Mean Plant Heights for Four Dates of Seeding
and Number of Days to Heading, 1950

Material	Mean ht.*	Days to head	Mean ht.*	Days to head	Mean ht.*	Days to head	Mean ht.*	Days to head
Atsel	17.8	49	18.5	36	22.5	35	26.8	36
Beecher	18.0	51	18.8	43	21.8	38	24.8	38
Tulare	16.5	49	17.7	42	19.3	37	23.0	41
Newal	19.3	59	22.3	50	30.0	48	32.3	45
Titan	20.0	55	20.3	48	30.3	42	30.7	43
C.I. 7155	20.2	64	25.8	59	34.7	54	28.7	52
C.I. 7248	18.5	61	22.8	51	23.3	49	23.2	48
Montcalm	22.7	61	31.0	54	38.3	49	39.3	48
H-44-1	19.7	53	20.8	44	29.2	39	29.7	36
Floya	19.0	50	19.5	41	29.7	40	30.7	37
Atsel x Tulare	18.5	48	17.5	42	23.5	42	26.7	40
Montcalm x Tulare	19.3	45	21.0	39	24.7	35	30.0	35
Montcalm x Beecher	17.7	51	20.7	43	30.5	38	28.8	38
Atsel x Beecher	17.2	49	17.4	45	26.3	36	28.0	38
Titan x Newal	19.0	54	21.5	42	30.3	41	31.5	41
Montcalm x Atsel	20.8	51	21.7	46	25.8	39	33.8	38
Montcalm x Titan	21.8	55	23.2	47	32.5	40	33.3	39
Montcalm x Newal	21.7	51	22.0	48	31.5	42	37.0	42
Beecher x C.I. 7248	16.0	56	17.2	43	29.5	39	--	--
Titan x Tulare	18.3	52	17.5	50	26.2	37	27.7	37
Titan x C.I. 7248	19.7	62	16.0	54	30.2	46	--	--
Montcalm x C.I. 7155	21.0	64	26.3	42	35.2	50	--	--
Atsel x C.I. 7155	19.8	55	22.7	43	34.7	39	33.3	42
Atsel x C.I. 7248	21.3	56	22.9	45	35.5	41	34.2	40
Montcalm x C.I. 7248	20.2	59	24.5	47	34.7	43	36.0	44
C.I. 7155 x C.I. 7248	18.3	60	26.8	47	29.3	49	30.5	49

* Mean height measured in inches.

The average gain in height between the third and fourth date of seeding is 2.1 inches. This figure represents the smallest increase although moisture supply was abundant during the growth period of the fourth replication.

The number of days to heading shows a significant decrease between the first and the third date of seeding. With the fourth date of seeding, the number of days to heading represents a levelling off in the downward trend. The range between the heading of the second and third replication represents the longest photoperiod. At the time of heading of the fourth date of seeding the days were again becoming shorter.

When the heights of plants for the first three dates of seeding are compared with the number of days to heading, a vegetative correlation between height and days to heading may be established (see Table 11). With progressive increase in height there is steady decrease in days to heading. These results indicate that with increasing photoperiod there is a corresponding increase in the rate at which the physiological processes operate. This increased activity leads to higher vegetative growth and at the same time speeds up the reproductive processes.

It is evident from the behaviour of the fourth date of seeding that the optimal condition for vegetative growth is reached. Some of the varieties such as C.I. 7155 and C.I. 7248 and some of the F_2 hybrids show a decrease in height from the third replication.

Shorter photoperiods have also led to longer periods to heading as seen in the varieties Atsel, Tulare and Titan and the hybrids Atsel x Beecher, Atsel x C.I. 7155, and Montcalm x C.I. 7248.

Other varieties, such as Beecher and hybrids such as Montcalm x Tulare, Titan x Newal, and Montcalm x Newal, did not show a significant change in the number of days to heading for the later dates of seeding.

It appears, from the observations on primordia development and plant heights under different conditions of growth, that earliness is largely influenced by the plants response to photoperiod and not by the amount of vegetative growth.

V. GENETICAL STUDY

1. Materials, Methods and Observations

The crosses numbered 1x to 16 x inclusive (see Table 2) were made in the field in 1948 and 1949, and the F₁ plants and parents grown in the greenhouse in the winter of 1949-1950. The F₂ generations of these crosses were grown in the field in the spring and summer of 1950. All crosses were compared with the parents in earliness.

2. Relation of Time of Heading to Length of Heading Period

It was observed both from the greenhouse and field study that the later varieties headed over a longer period than the earlier ones. The correlation between the time of heading and the length of the heading period is shown in Tables 12, 13, and 14. As a rule, not only did the later varieties require more days to complete their heading, but when the same varieties were seeded at different dates the later planted lots headed as a rule over a longer period. This is shown in Table 15. As the rate of heading is influenced by other factors besides the length of day, the behaviour observed in this study may not always apply.

THE HISTORY OF THE

REIGN OF

CHARLES THE FIRST

BY

JOHN BURNET

OF THE UNIVERSITY OF OXFORD

IN TWO VOLUMES

LONDON

Printed by J. Stansfeld, at the University Press

1704

THE HISTORY OF THE

REIGN OF

CHARLES THE FIRST

BY

JOHN BURNET

OF THE UNIVERSITY OF OXFORD

IN TWO VOLUMES

LONDON

Printed by J. Stansfeld, at the University Press

1704

THE HISTORY OF THE

REIGN OF

CHARLES THE FIRST

Table 12

Correlation Between Mean Date of Heading of Parent Varieties and
Number of Days in Heading, 1949. (Greenhouse)

Date	Days required to head																									
	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26				
Oct. 10	1	5	-	-	-	1	1	-	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
11	3	2	6	-	-	1	-	1	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
12		6	2	2	1	-	2		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
13			-	-	-	1	-		-	-	-	-	-	-	-	-	1	-	-	-	-	-	-	-	-	-
14			-	-	-	-	-		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15			-	-	-	2	-		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
16-18																										
19			-	-	-	-	-		-	-	-	-	-	-	-	1	-	-	-	-	-	-	-	-	-	-
20			-	-	-	-	-		1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
21			-	-	-	-	-		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
22			-	-	-	-	-		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
23			1	-	-	-	-		-	-	-	-	-	-	-	-	-	1	-	-	-	-	-	-	-	-
24			-	-	-	-	-		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25			-	-	-	-	-		1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-
26			-	-	-	-	-		-	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
27			-	-	-	1	-		-	-	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
28-30																										
31			-	-	-	-	-		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Nov. 1			-	-	-	-	-		-	-	-	-	1	-	-	-	-	-	-	-	-	-	-	-	-	-
2			-	-	1	-	1		-	-	-	-	-	-	-	-	-	-	-	-	-	2	-	-	-	-
3			-	-	-	-	-		-	2	-	-	-	-	-	1	-	-	-	-	-	-	-	-	-	-
4			-	-	-	-	-		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5			-	-	-	1	-		-	-	-	-	1	1	-	-	-	-	-	-	-	-	-	-	-	-
6			-	-	-	-	-		-	-	-	-	-	-	-	-	-	-	-	2	-	-	-	-	-	-
7			-	-	-	-	-		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
8			-	-	-	-	-		-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	-	-	-
9-11																										
12			1				-		-	-	-	-	-			-	-	-								
13							-		-	-	-	-	-			-	-	-								
14							1		1	-	-	-	-			-	-	-								
15,16																										
17										-	-	1				-	-	-								
18										-	-					1	-	-								
19										-	-					-	-	-								
20										-	-					-	1	-								
23										1	-					-	-	-								
25											1					-	-	-								
29																1										
Dec. 3																				1						

$$r = .52 \pm .05$$

Table 13

Correlation Between Mean Date of Heading of Parent Varieties
and Number of Days in Heading, 1950 (Field)

Date	Days required to head															
	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	
June 19	1	2	-	-	-		-	-	-		-			-	-	
20, 21	-	-	-	-	-		-	-	-		-			-	-	
22	-	-	2	-	-		-	-	-		-			-	-	
23	-	1	-	-	-		-	-	-		-			-	-	
24	-	-	2	-	-		-	-	-		-			-	-	
25	1	-	-	-	1		-	-	-		-			-	-	
26		-	-	-	-		-	-	-		-			-	-	
27		8	5	-	1		-	-	-		-			-	-	
28		-	-	1	-		-	-	-		-			-	-	
29		-	-	1	-		-	-	-		-			-	-	
30		-	-	-	-		-	-	-		-			-	-	
July 1		1	1	1	-		-	-	-		-			-	-	
2			-		-		1	-	-		-			-	-	
3			-		-		-	-	-		-			-	-	
4-6			-		-		-	-	-		-			-	-	
7			-		-		2	-	-		-			-	-	
8			-		-		-	1	-		-			-	-	
9			-		-		-	-	-		1			-	-	
10			-		1		-	-	-					-	-	
11			-		2		-	-	-					-	-	
12			-		-		2	-	1					-	-	
13, 14			-		-		-	-						-	-	
15			-		1		-	-						-	-	
16			-					1						-	-	
17			-					-						-	-	
18			-					-						-	1	
19, 20			-					-						-		
21			1					-						-		
22								-						-		
23								-						1		
24								1								

$$r = .67 \pm .02$$

Table 14

Correlation Between Mean Date of Heading of Parent Varieties
and Number of Days in Heading, 1950 (Field)

Date	Days required to head												22
	6	7	8	9	10	11	12,13	14	15,16,17	18	19	20	
July 3-5	-	-	-	-		-		-		-		-	-
6	1	-	-	-		-		-		-		-	-
7	-	2	-	-		-		-		-		-	-
8	2	-	1	1		-		-		-		-	-
9	-	-				-		-		-		-	-
10	-	1				-		-		-		-	-
11	1					-		-		-		-	-
12,13						-		-		-		-	-
14						1		-		-		-	-
15-17								-		-		-	-
18								-		-		-	-
19								1		-		-	-
20										-		-	-
21-23										-		-	-
24-26										-		-	-
27										1		-	-
28												-	-
29												-	1
30,31												-	-
Aug. 1												-	-
2-4												-	-
5-6												-	-
7												1	-
8-10													-
11													1

$$r = .91 \pm .05$$

Table 15

Influence of the Time of Planting and of Heading Upon the

Number of Days Required to Head

		No. 1 Atsel 1950	No. 2 Beecher 1950	No. 5 Titan 1950	No. 6 Newal 1950	No. 7 Montcalm 1950	No. 9 C.I. 7248 1950
Planted		1/5 1/6	1/5 15/5	15/5 1/6	15/5 1/6	15/5 1/6	1/5 1/6
June	19	2					
	20	5					
	21	6	2				
	22	7	7				
	23	10	10				
	24	11	12				
	25	5	16				
	26	4	10				
	27	2	8	1			
	28	1	5	3			
	29		1	5			
	30		1	2			
July	1			1			
	2			3			4
	3			5			4
	4			17			4
	5			11			3
	6			10	1		3
	7	2		4	6		5
	8	5		1	6		1
	9	7		1	4		6
	10	8			2	2	1
	11	2	1			5	
	12	4				9	
	13	4				19	
	14	1		4		8	
	15	7		4		9	
	16	2		12		7	
	17	2		8		3	
	18	1		6		2	
	19			10		2	
	20			5	1		4
	21			5	1		6
	22	1		2			7
	23	1			1		14
	24				9		16
	25			1	4		10
	26			1	2		9
	27						2
	28				2		1
	29				2		2
	30				1		1
	31						1

3. Experimental Results

(a) The F₁ Generations

The dates of heading of the F₁ plants and the parental varieties are given in Table 16. While the conditions in the greenhouse were favourable there was a lack of uniformity in temperature in different parts of the bed and this doubtless affected the rate of development of certain lines. In the greenhouse most of the varieties headed over longer periods than they did under field conditions, the only exception being the last field planting made June 15, 1950.

It is considered that these data show only roughly the relative earliness of the F₁ generations with respect to the parents. In most of the crosses, however, it was significant that the hybrids varied from an intermediate to a heading date nearer to the early parents. In the apparent exceptions, such as Atsel x Beecher, Montcalm x C.I. 7248, and C.I. 7155 x C.I. 7248, while the differences between the mean dates of heading of the parents were large, the differences of the means of the F₁ populations and the means of the early parents were comparatively small.

Previous mention was made of the fact that Hayes and East (14) believed the increase in rate of development in F₁ crosses

Table 16

Dates of Heading of F₁ Plants and Parents in Greenhouse, 1949. Seeded August 22, 1949

Material	Dates of heading					
	Pistillate parent		Staminate parent		F ₁ parents	
	Range	Mean	Range	Mean	Range	Mean
1x Atsel x Tulare	10/10 - 25/10	17/10.50 ± .24	10/10 - 26/10	18/10.71 ± 1.12	11/10 - 25/10	18/10.22 ± 1.35
2x Montcalm x Tulare	2/11 - 19/11	10/11.88 ± 1.32	10/10 - 26/10	18/10.71 ± 1.12	18/10 - 6/11	27/10.33 ± 1.00
3x Montcalm x Beecher	2/11 - 19/11	10/11.88 ± 1.32	14/10 - 22/10	18/10.75 ± 1.66	18/10 - 9/11	29/10.50 ± .72
4x Atsel x Beecher	10/10 - 25/10	17/10.50 ± .24	14/10 - 22/10	18/10.75 ± 1.66	20/10 - 9/11	31/10.90 ± .82
5x Titan x Newal	38/10 - 20/11	8/11.14 ± 1.29	5/11 - 16/11	10/11.50 ± .24	1/11 - 19/11	10/11.67 ± 2.41
6x Montcalm x Atsel	2/11 - 19/11	10/11.88 ± 1.32	10/10 - 25/10	17/10.50 ± .24	20/10 - 14/11	1/11.25 ± 1.06
7x Montcalm x Titan	2/11 - 19/11	10/11.88 ± 1.32	28/10 - 20/11	8/11.14 ± 1.29	28/10 - 20/11	8/11.79 ± 1.45
8x Montcalm x Newal	2/11 - 19/11	10/11.88 ± 1.32	5/11 - 16/11	10/11.50 ± .24	3/11 - 29/11	16/11.80 ± 2.94
9x Beecher x C.I. 7248	14/10 - 22/10	18/10.75 ± 1.66	16/11 - 18/12	1/12.17 ± 2.57	14/10 - 20/11	1/11.33 ± 1.00
10x Titan x Tulare	28/10 - 20/11	8/11.14 ± 1.29	10/10 - 20/10	18/10.71 ± 1.12	16/10 - 18/11	2/11.25 ± 1.06
11x Titan x C.I. 7248	28/10 - 20/11	8/11.14 ± 1.29	16/11 - 18/12	1/12.17 ± 2.57	7/11 - 26/12	1/12.50 ± 1.12
12x Montcalm x C.I. 7155	2/11 - 19/11	10/11.88 ± 1.32	18/11 - 12/12	30/11.79 ± 1.45	9/11 - 18/12	29/11.12 ± 1.29
13x Atsel x C.I. 7155	10/10 - 25/10	17/10.50 ± .24	18/11 - 12/12	30/11.79 ± 1.45	28/10 - 18/12	22/11.83 ± .87
14x Atsel x C.I. 7248	10/10 - 25/10	17/10.50 ± .24	16/11 - 18/12	1/12.17 ± 2.57	25/10 - 10/12	17/11.14 ± 1.32
15x Montcalm x C.I. 7248	2/11 - 19/11	10/11.88 ± 1.32	16/11 - 18/12	1/12.17 ± 2.57	27/11 - 14/12	6/12.75 ± 1.66
16x C.I. 7155 x C.I. 7248	18/11 - 12/12	30/11.79 ± 1.45	16/11 - 18/12	1/12.17 ± 2.57	21/11 - 20/12	5/12.07 ± .34

Name				Address			
1	Mr. J. H. Smith	123	4th St.	St. Louis	Mo.	63101	
2	Mr. W. B. Jones	456	7th St.	St. Louis	Mo.	63102	
3	Mr. C. D. Brown	789	10th St.	St. Louis	Mo.	63103	
4	Mr. E. F. Green	1011	13th St.	St. Louis	Mo.	63104	
5	Mr. G. H. White	1314	16th St.	St. Louis	Mo.	63105	
6	Mr. I. J. Black	1617	19th St.	St. Louis	Mo.	63106	
7	Mr. K. L. Gray	1920	22nd St.	St. Louis	Mo.	63107	
8	Mr. M. N. Hall	2223	25th St.	St. Louis	Mo.	63108	
9	Mr. O. P. King	2526	28th St.	St. Louis	Mo.	63109	
10	Mr. Q. R. Lee	2829	31st St.	St. Louis	Mo.	63110	
11	Mr. S. T. Young	3132	34th St.	St. Louis	Mo.	63111	
12	Mr. U. V. Wright	3435	37th St.	St. Louis	Mo.	63112	
13	Mr. X. Y. Adams	3738	40th St.	St. Louis	Mo.	63113	
14	Mr. Z. A. Baker	4039	43rd St.	St. Louis	Mo.	63114	
15	Mr. B. C. Miller	4340	46th St.	St. Louis	Mo.	63115	
16	Mr. D. E. Wilson	4641	49th St.	St. Louis	Mo.	63116	
17	Mr. F. G. Moore	4942	52nd St.	St. Louis	Mo.	63117	
18	Mr. H. I. Taylor	5243	55th St.	St. Louis	Mo.	63118	
19	Mr. J. K. Evans	5544	58th St.	St. Louis	Mo.	63119	
20	Mr. L. M. Roberts	5845	61st St.	St. Louis	Mo.	63120	
21	Mr. N. O. Clark	6146	64th St.	St. Louis	Mo.	63121	
22	Mr. P. Q. Lewis	6447	67th St.	St. Louis	Mo.	63122	
23	Mr. R. S. Walker	6748	70th St.	St. Louis	Mo.	63123	
24	Mr. T. U. Hall	7049	73rd St.	St. Louis	Mo.	63124	
25	Mr. V. W. King	7350	76th St.	St. Louis	Mo.	63125	
26	Mr. X. Y. Lee	7651	79th St.	St. Louis	Mo.	63126	
27	Mr. Z. A. Young	7952	82nd St.	St. Louis	Mo.	63127	
28	Mr. B. C. Wright	8253	85th St.	St. Louis	Mo.	63128	
29	Mr. D. E. Adams	8554	88th St.	St. Louis	Mo.	63129	
30	Mr. F. G. Baker	8855	91st St.	St. Louis	Mo.	63130	
31	Mr. H. I. Miller	9156	94th St.	St. Louis	Mo.	63131	
32	Mr. J. K. Wilson	9457	97th St.	St. Louis	Mo.	63132	
33	Mr. L. M. Moore	9758	100th St.	St. Louis	Mo.	63133	
34	Mr. N. O. Taylor	10059	103rd St.	St. Louis	Mo.	63134	
35	Mr. P. Q. Evans	10360	106th St.	St. Louis	Mo.	63135	
36	Mr. R. S. Roberts	10661	109th St.	St. Louis	Mo.	63136	
37	Mr. T. U. Clark	10962	112th St.	St. Louis	Mo.	63137	
38	Mr. V. W. Lewis	11263	115th St.	St. Louis	Mo.	63138	
39	Mr. X. Y. Walker	11564	118th St.	St. Louis	Mo.	63139	
40	Mr. Z. A. Hall	11865	121st St.	St. Louis	Mo.	63140	
41	Mr. B. C. King	12166	124th St.	St. Louis	Mo.	63141	
42	Mr. D. E. Lee	12467	127th St.	St. Louis	Mo.	63142	
43	Mr. F. G. Young	12768	130th St.	St. Louis	Mo.	63143	
44	Mr. H. I. Wright	13069	133rd St.	St. Louis	Mo.	63144	
45	Mr. J. K. Adams	13370	136th St.	St. Louis	Mo.	63145	
46	Mr. L. M. Baker	13671	139th St.	St. Louis	Mo.	63146	
47	Mr. N. O. Miller	13972	142nd St.	St. Louis	Mo.	63147	
48	Mr. P. Q. Wilson	14273	145th St.	St. Louis	Mo.	63148	
49	Mr. R. S. Moore	14574	148th St.	St. Louis	Mo.	63149	
50	Mr. T. U. Taylor	14875	151st St.	St. Louis	Mo.	63150	
51	Mr. V. W. Evans	15176	154th St.	St. Louis	Mo.	63151	
52	Mr. X. Y. Roberts	15477	157th St.	St. Louis	Mo.	63152	
53	Mr. Z. A. Clark	15778	160th St.	St. Louis	Mo.	63153	
54	Mr. B. C. Lewis	16079	163rd St.	St. Louis	Mo.	63154	
55	Mr. D. E. Walker	16380	166th St.	St. Louis	Mo.	63155	
56	Mr. F. G. Hall	16681	169th St.	St. Louis	Mo.	63156	
57	Mr. H. I. King	16982	172nd St.	St. Louis	Mo.	63157	
58	Mr. J. K. Lee	17283	175th St.	St. Louis	Mo.	63158	
59	Mr. L. M. Young	17584	178th St.	St. Louis	Mo.	63159	
60	Mr. N. O. Wright	17885	181st St.	St. Louis	Mo.	63160	
61	Mr. P. Q. Adams	18186	184th St.	St. Louis	Mo.	63161	
62	Mr. R. S. Baker	18487	187th St.	St. Louis	Mo.	63162	
63	Mr. T. U. Miller	18788	190th St.	St. Louis	Mo.	63163	
64	Mr. V. W. Wilson	19089	193rd St.	St. Louis	Mo.	63164	
65	Mr. X. Y. Moore	19390	196th St.	St. Louis	Mo.	63165	
66	Mr. Z. A. Taylor	19691	199th St.	St. Louis	Mo.	63166	
67	Mr. B. C. Evans	19992	202nd St.	St. Louis	Mo.	63167	
68	Mr. D. E. Roberts	20293	205th St.	St. Louis	Mo.	63168	
69	Mr. F. G. Clark	20594	208th St.	St. Louis	Mo.	63169	
70	Mr. H. I. Lewis	20895	211th St.	St. Louis	Mo.	63170	
71	Mr. J. K. Walker	21196	214th St.	St. Louis	Mo.	63171	
72	Mr. L. M. Hall	21497	217th St.	St. Louis	Mo.	63172	
73	Mr. N. O. King	21798	220th St.	St. Louis	Mo.	63173	
74	Mr. P. Q. Lee	22099	223rd St.	St. Louis	Mo.	63174	
75	Mr. R. S. Young	22300	226th St.	St. Louis	Mo.	63175	
76	Mr. T. U. Wright	22601	229th St.	St. Louis	Mo.	63176	
77	Mr. V. W. Adams	22902	232nd St.	St. Louis	Mo.	63177	
78	Mr. X. Y. Baker	23203	235th St.	St. Louis	Mo.	63178	
79	Mr. Z. A. Miller	23504	238th St.	St. Louis	Mo.	63179	
80	Mr. B. C. Wilson	23805	241st St.	St. Louis	Mo.	63180	
81	Mr. D. E. Moore	24106	244th St.	St. Louis	Mo.	63181	
82	Mr. F. G. Taylor	24407	247th St.	St. Louis	Mo.	63182	
83	Mr. H. I. Evans	24708	250th St.	St. Louis	Mo.	63183	
84	Mr. J. K. Roberts	25009	253rd St.	St. Louis	Mo.	63184	
85	Mr. L. M. Clark	25310	256th St.	St. Louis	Mo.	63185	
86	Mr. N. O. Lewis	25611	259th St.	St. Louis	Mo.	63186	
87	Mr. P. Q. Walker	25912	262nd St.	St. Louis	Mo.	63187	
88	Mr. R. S. Hall	26213	265th St.	St. Louis	Mo.	63188	
89	Mr. T. U. King	26514	268th St.	St. Louis	Mo.	63189	
90	Mr. V. W. Lee	26815	271st St.	St. Louis	Mo.	63190	
91	Mr. X. Y. Young	27116	274th St.	St. Louis	Mo.	63191	
92	Mr. Z. A. Wright	27417	277th St.	St. Louis	Mo.	63192	
93	Mr. B. C. Adams	27718	280th St.	St. Louis	Mo.	63193	
94	Mr. D. E. Baker	28019	283rd St.	St. Louis	Mo.	63194	
95	Mr. F. G. Miller	28320	286th St.	St. Louis	Mo.	63195	
96	Mr. H. I. Wilson	28621	289th St.	St. Louis	Mo.	63196	
97	Mr. J. K. Moore	28922	292nd St.	St. Louis	Mo.	63197	
98	Mr. L. M. Taylor	29223	295th St.	St. Louis	Mo.	63198	
99	Mr. N. O. Evans	29524	298th St.	St. Louis	Mo.	63199	
100	Mr. P. Q. Roberts	29825	301st St.	St. Louis	Mo.	63200	

of corn to be caused by hybrid vigour. It is possible that, in these F_1 barley hybrids, hybrid vigour induced somewhat more rapid development and earlier heading, but the data secured in the F_2 generations indicate that this does not adequately explain the increase in earliness.

(b) The F_2 Generations

In the F_1 plants heading coincided with the early parents, was earlier than the early parents or was intermediate between the early and late with some skewness towards the early end. From this behaviour it might have been expected that the F_2 generations would range in time of heading from the early parent to the late parent or from earlier than the early parent to later than the late parent. In either case, the majority of the plants would have been expected to be early.

Four dates of seeding of the F_2 generations of 16 crosses and the parents involved were grown in the field. Fig. 7 shows a sectional view of the second field replication seeded on the 15th of May, 1950. Between the two paper markers at the ends of rod row No. 50, may be seen three sections of plants separated by white pegs. The middle section represents the F_2 generation of the cross Montcalm x C.I. 7248 while the Montcalm parent occupies the foreground and the parent C.I. 7248 is in the rear of row No. 50.

Fig. 7

Sectional view of second field replication seeded May 15, 1950 and showing parental types towards ends of rows with F_2 hybrids between them.

The most noticeable fact is that the majority of the F_2 families (see Fig. 7, middle section of rows) began to head earlier than the early parents. It was observed in these crosses that the earliest plants ranged from 2 to 10 days earlier than the earliest plants of the respective early parents. This behaviour was, in general, true for the other three replications.

In Fig. 8 is seen, on the left of the white stake, the pistillate parent and on the right the F_2 generation of the cross Montcalm x Tulare, both sections being seeded on the 1st of June, 1950, in replication 3. The early Tulare and late Montcalm parents differed by 12 days in time of heading, whereas the earliest plant of the F_2 hybrid was 3 days earlier than the earliest Tulare.

Dates of seeding for the four field replications differed by 15 days, the first seeding being made on May 1, 1950. The heading periods of the F_2 generations in comparison with those of the parents are given in Tables 17 to 20.

The heading in F_1 and F_2 of the 16 crosses studied may be summarized as follows:

In 7 crosses the F_1 plants headed earlier than the early parents or a little later (in crosses 5x, 6x, and 16x), and the F_2 families began heading earlier than the early parent.

In 3 crosses the F_1 plants headed with the early parents, and the F_2 families began to head earlier than the early parents.

In 2 crosses the F_1 plants headed after the early parent, but the F_2 families began to head earlier than the early parent.

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

Fig. 8

Left, pistillate parent; right, F_2 hybrid of a cross Montcalm x Tulare, both sections seeded June 1, 1950.

Table 17

Adjusted Frequency Distributions for Parental and F₂ Plants, Classified for Days to Heading, 1950

Replicate 1 (Seeded 1/5/50) Material	Days to heading																	Total no. of plants	
	46	48	50	52	54	56	58	60	62	64	66	68	70	72	74	76	78		80
Atsel x Tulare																			
Atsel		7	12	13	21	9	4	1											67
Tulare			9	19	29	10	4	2	2										75
F ₂	3	7	8	10	4	3	3	2	1										41
Atsel x Beecher																			
Atsel		7	12	13	21	9	4	1											67
Beecher			9	22	16	11	5	2	2										67
F ₂	1	4	9	17	5	2	2												39
Atsel x C.I. 7155																			
Atsel		7	12	13	21	9	4	1											67
C.I. 7155																			23
F ₂					2	2	4	5	5	1	4	5	6	4	2	1	2	1	40
Atsel x C.I. 7248																			
Atsel		7	12	13	21	9	4	1											67
C.I. 7248																			27
F ₂				1	2	2	5	6	3	4	7	6	5	1	2				33
Montcalm x Atsel																			
Montcalm																			49
Atsel		7	12	13	21	9	4	2	2	6	18	10	8	2	1				67
F ₂			1	2	3	4	6	7	3	4	3	2		1					36
Montcalm x Beecher																			
Montcalm																			49
Beecher			9	22	16	11	5	2	2	6	18	10	8	2	1				67
F ₂				5	6	6	5	4	3	3	2		2	1					37

Table 17 (Continued)

Adjusted Frequency Distributions for Parental and F₂ Plants, Classified for Days to Heading, 1950

Replicate 1 (Seeded 1/5/50) Material	Days to heading																Total no. of plants	
	46	48	50	52	54	56	58	60	62	64	66	68	70	72	74	76		78
Montcalm x Tulare																		
Montcalm			9	19	29	10	4	2	2	6	18	10	8	2	1			49
Tulare			7	15	4	4	3	2	2	2	1	1						75
F ₂	1	3						3	2									46
Beecher x C.I. 7248			9	22	16	11	5	2	2									67
Beecher									3	4	7	6	5	2				27
C.I. 7248									2									35
F ₂					1	3	8	7	2	4	3	3	2	1	1			
Titan x Tulare																		
Titan					1		9	15	14	8	2		2	1				52
Tulare			9	19	29	10	4	2	2									75
F ₂				2	5	8	3	6			2							28
Titan x Newal																		
Titan					1		9	15	14	8	2		2	1				52
Newal							1	2	3	2	4	3	3	2		2		22
F ₂					3	2	4	8	10	3	2		3	1	2	1		39
Titan x C.I. 7248																		
Titan					1		9	15	14	8	2		2	1				52
C.I. 7248									3	4	7	6	5	2				27
F ₂							2	3	3	4	5	3		1	2			21
Montcalm x Newal																		
Montcalm								2	2	6	18	10	8	2	1			49
Newal							1	2	3	2	4	3	3	2		2		22
F ₂					2	3	5	5	6	3	4	3	1		2		1	35

Table 17 (Continued)

Adjusted Frequency Distributions for Parental and F₂ Plants, Classified for Days to Heading, 1950

Replicate 1 (Sseeded 1/5/50) Material	Days to heading														Total no. of plants			
	46	48	50	52	54	56	58	60	62	64	66	68	70	72		74	76	78
Montcalm x Titan																		
Montcalm						1	9	2	2	6	18	10	8	2	1			49
Titan							15	14	8	2			2	1				52
F ₂					2	1	4	10	6	2		1	2		2	2	1	33
Montcalm x C.I. 7155																		
Montcalm							2	2	2	6	18	10	8	2	1			49
C.I. 7155										1	4	5	6	4	2	1		23
F ₂								3	3	3	7	5	3	1	2	1	2	28
Montcalm x C.I. 7248																		
Montcalm							2	2	2	6	18	10	8	2	1			49
C.I. 7248								3	3	4	7	6	5	2	2			27
F ₂						2		2	2	10	6	5	3		2	2		32
C.I. 7155 x C.I. 7248																		
C.I. 7155										1	4	5	6	4	2	1		23
C.I. 7248								3	1	4	7	6	5	2				27
F ₂							2	1	5	5	4	1	2	1	2		2	21

Table 18

Adjusted Frequency Distributions for Parental and F₂ Plants, Classified for Days to Heading, 1950

Replicate 2 (Sseed 15/5/50)	Days to heading																	Total no. of plants
	38	40	42	44	46	48	50	52	54	56	58	60	62	64	66	68	70	
Material																		
Atsel x Tulare																		
Atsel	1	3	9	24	10	5	1	2										55
Tulare			5	11	13	2	2	1										34
F ₂	5	5	14	6	3	4		2										41
Atsel x Beecher																		
Atsel	1	3	9	24	10	5	1	2										55
Beecher				2	4	9	8	3		2								30
F ₂		1	4	15	5	6	2	3		3								36
Atsel x C.I. 7155																		
Atsel	1	3	9	24	10	5	1	2										55
C.I. 7155																		28
F ₂				2	3	6	7	3		4		4	6	8	5	3	2	34
Atsel x C.I. 7248																		
Atsel	1	3	9	24	10	5	1	2										55
C.I. 7248																		54
F ₂			1	2	3	3	7	5		6		9	19	8	9	2		39
Montcalm x Atsel																		
Montcalm																		59
Atsel	1	3	8	24	10	5	1	2		2	4	19	10	4	1	2		55
F ₂	1	2	8	7	2	5	4	1		3	2	2	1	2				40
Montcalm x Beecher																		
Montcalm																		59
Beecher																		30
F ₂			1	3	8	3	4	1		3	2	3	2	2	1	2		35

Table 18 (Continued)

Adjusted Frequency Distributions for Parental and F₂ Plants, Classified for Days to Heading, 1950

Replicate 2 (Seeded 15/5/50) Material	Days to heading																	Total no. of plants
	38	40	42	44	46	48	50	52	54	56	58	60	62	64	66	68	70	
Montcalm x Tulare																		
Montcalm																		
Tulare			5	11	13	2	2	1	2	4	4	19	17	10	4	1	2	59
F ₂	2	2	3	9	6	3	4	2	3	2	2	2		2	1			34 41
Beecher x C.I. 7248																		
Beecher				2	4	9	8	3	2	2	2	5	9	19	8	9	2	30
C.I. 7248																		54
F ₂					1	2	3	7	8	3	3	2	2	1	2	1	1	35
Titan x Tulare																		
Titan			5	11	13		5	17	11	10	4	1	1	1				49
Tulare			2	5	11	5	4	1	3	2	3	2						34
F ₂																		37
Titan x Newal																		
Titan							5	17	11	10	4	1	1	1				49
Newal								1	4	10	6	4	2	2	1			29
F ₂					1	3	9	4	3	1			2	1	1			25
Titan x C.I. 7248																		
Titan							5	17	11	10	4	1	1	1				49
C.I. 7248										2	5	9	19	8	9	2		54
F ₂								3	7	11	8	3	5	1	3	1	1	43
Montcalm x Titan																		
Montcalm																		
Titan							5	17	11	10	4	17	10	4	1	2		59
F ₂				2	5		9	4	1	3	3	2				1	2	49 34

Table 18 (Continued)

Adjusted Frequency Distributions for Parental and F₂ Plants, Classified for Days to Heading, 1950

Replicate (Seeded 15/5/50) Material	Days to heading																	Total no. of plants
	38	40	42	44	46	48	50	52	54	56	58	60	62	64	66	68	70	
Montcalm x Newal																		
Montcalm									2	4	19	17	10	4	1	2		59
Newal								1	4	10	6	4	2	2				29
F ₂						2	4	3	7	5	2	3	1		2	2	1	32
Montcalm x C.I. 7155																		
Montcalm									2	4	19	17	10	4	1	2	2	59
C.I. 7155								3	4	8	5	4	6	8	5	3	1	28
F ₂												2	4	3		3		33
Montcalm x C.I. 7248																		
Montcalm									2	4	19	17	10	4	1	2		59
C.I. 7248										2	5	9	19	8	9	2	1	54
F ₂								1	4	9	5	4	2		1			27
C.I. 7155 x C.I. 7248																		
C.I. 7155												4	6	8	5	3	2	28
C.I. 7248										2	5	9	19	8	9	2	1	54
F ₂										3	9	6	3	1	3	1		27

Table 19

Adjusted Frequency Distributions for Parental and F₂ Plants, Classified for Days to Heading, 1950

Replicate 3 (Seeded 1/6/50) Material	Days to heading														Total no. of plants		
	34	36	38	40	42	44	46	48	50	52	54	56	58	60		62	64
Atsel x Tulare																	
Atsel	2	5	12	10	8	7	4		2								50
Tulare		1	4	5	7	4	3	3	1								28
F ₂	3	7	4	3	2	4	3	1	2								29
Atsel x Beecher																	
Atsel	2	5	12	10	8	7	4		2								50
Beecher			4	1	6	9	4	4	3	1							32
F ₂		3	2	6	5	2	3		2	2	1						26
Atsel x C.I. 7155																	
Atsel	2	5	12	10	8	7	4		2								50
C.I. 7155																	24
F ₂			1	4	6	8	3	1	3	4	3	8	4	3	4	2	38
Atsel x C.I. 7248																	
Atsel	2	5	12	10	8	7	4		2								50
C.I. 7248																	17
F ₂			1	4	7	5	3	4	2	3	5	2	4	2			32
Montcalm x Atsel																	
Montcalm								4	6	7	14	16	10	9	2	1	69
Atsel	2	5	12	10	8	7	4		2								50
F ₂		1	4	3	9	6	4	3	2	3	2	1		1		2	41
Montcalm x Beecher																	
Montcalm								4	6	7	14	16	10	9	2	1	69
Beecher								4	3	1							32
F ₂			4	1	6	9	4	4	3	3	2		2		1		33

Table 19 (Continued)

Adjusted Frequency Distributions for Parental and F₂ Plants, Classified for Days to Heading, 1950

Replicate 3 (Seeded 1/6/50) Material	Days to heading																Total no. of plants
	34	36	38	40	42	44	46	48	50	52	54	56	58	60	62	64	
Montcalm x Tulare																	
Montcalm								4	6	7	14	16	10	9	2	1	69
Tulare		1	4	5	7	4	3	3	1	2							28
F ₂	2	3	7	5	5	3	4	3	2	3	1	2		1			41
Beecher x C.I. 7248																	
Beecher			4	1	6	9	4	4	3	1							32
C.I. 7248									1	3	5	2	4	2			17
F ₂		1	2	2	6	5	3	3	2	3		1	2		2		30
Titan x Tulare																	
Titan		1	4	5	4	4	12	14	6	5	5	4	1				55
Tulare					7	4	3	3	1								28
F ₂		2	2	6	3	2	1	3	2		1						22
Titan x Newal																	
Titan					4	4	12	14	6	5	5	4	1				55
Newal							2	3	9	6	4	1	2	2			28
F ₂					1	3	9	4	3		2	1			1		26
Titan x C.I. 7248																	
Titan					4	4	12	14	6	5	5	4	1				55
C.I. 7248									1	3	5	2	4	2			17
F ₂							2	3	7	5	4	2	1				24
Montcalm x Titan																	
Montcalm								4	6	7	14	16	10	9	2	1	69
Titan					4	4	12	14	6	5	5	4	1				55
F ₂					3	4	8	3	5	2	4	2	1	2	2	1	36

Table 19 (Continued)

Adjusted Frequency Distributions for Parental and F₂ Plants, Classified for Days to Heading, 1950

Replicate 3 (Seeded 1/6/50) Material	Days to heading																Total no. of plants
	34	36	38	40	42	44	46	48	50	52	54	56	58	60	62	64	
Montcalm x Newal																	
Montcalm								4	6	7	14	16	10	9	2	1	69
Newal							2	3	9	6	4		2	2			28
F ₂					2	2	4	6	6	3	2		1	1	2	1	30
Montcalm x C.I. 7155																	
Montcalm								4	6	7	14	16	10	9	2	1	69
C.I. 7155											3	8	4	3	4	2	24
F ₂									2	3	11	6	4	3	1	1	31
Montcalm x C.I. 7248																	
Montcalm								4	6	7	14	16	10	9	2	1	69
C.I. 7248									1	3	5	2	4	2			17
F ₂					2	3	9		6	4	2		3	1			30
C.I. 7155 x C.I. 7248																	
C.I. 7155											3	8	4	3	4	2	24
C.I. 7248									1	3	5	2	4	2			17
F ₂							1		4	6	3	2		1	2	1	20

Table 20

Adjusted Frequency Distributions for Parental and F₂ Plants, Classified for Days to Heading, 1950

Replicate 4 (Seeded 15/6/50) Material	Days to heading																	Total no. of plants	
	32	34	36	38	40	42	44	46	48	50	52	54	56	58	60	62	64		66
Atsel x Tulare																			
Atsel	1	2	5	9	14	8	3	4	1		2	1							50
Tulare			2	6	7	11	5	3	2	2	1								39
F ₂		1	3	8	6	2	4	3	1		2								30
Atsel x Beecher																			
Atsel	1	2	5	9	14	8	3	4	1		2	1							50
Beecher				3	2	10	7	4	4	3	1								34
F ₂	2	2	8	7	7	4	5	2		2	2	1							42
Atsel x C.I. 7155																			
Atsel	1	2	5	9	14	8	3	4	1		2	1							50
C.I. 7155											1	3		8	6	3	2	2	1
F ₂				1	4	9	8	5	7	1	4	3	5	1	1	2	1	2	31
Atsel x C.I. 7248																			52
Atsel	1	2	5	9	14	8	3	4	1		2	4		3	2	3	2	2	27
C.I. 7248										2	4	2		2		1	2	1	46
F ₂					1	5	10	7	8	3	4		5	2					
Montcalm x Atsel																			
Montcalm								1	4		10	8	5	5	3	2	4	1	52
Atsel	1	2	5	9	14	8	3	4	1	9	2	1	5	1					50
F ₂			2	1	5	7	5	3	1		3	2	2	1	2			2	36
Montcalm x Beecher																			
Montcalm								1	4		10	8	5	5	3	2	4	1	52
Beecher				3	2	10	7	4	4	9	1		5	1					34
F ₂			2	3	5	7	4	3	3	2		2	1	2	2		1		37

1 87 1

Table 20 (Continued)

Adjusted Frequency Distributions for Parental and F₂ Plants, Classified for Days to Heading, 1950

Replicate 4 (Seeded 15/6/50)	Days to heading																Total no. of plants			
	32	34	36	38	40	42	44	46	48	50	52	54	56	58	60	62		64	66	68
Material																				
Montcalm x Tulare																				
Montcalm			2	6	7	11	5	1	4	9	10	8	5	5	3	2	4	1		52
Tulare			3	6	6	5	3	3	2	2	1	2	2	1	2		2		42	39
F ₂	2									3										42
Titan x Tulare																				
Titan			2	6	7	3	8	6	3	3	2	4	1	2	2	1				35
Tulare			4	2	9	11	5	3	2	2	1	1	2							39
F ₂						6	2	3		2	3			1						35
Titan x Newal																				
Titan						3	8	6	3	3	2	4	1	2	2	1	1			35
Newal						2	2	2	7	6	4	3	2		2	2	2			31
F ₂				1		2	9	6	4	4	2	3		3	1	1	2	1		39
Montcalm x Titan																				
Montcalm								1	4	9	10	8	5	5	3	2	4	1		52
Titan						3	8	6	3	3	2	4	1	2	2	1	2			35
F ₂				1	4	7	6	2	4	3	1			1		2				35
Montcalm x Newal																				
Montcalm								1	4	9	10	8	5	5	3	2	4	1		52
Newal							2	2	7	6	4	3	2	2	2	2	1			31
F ₂						1	3	4	5	4	4	2	3	1	1	2	2	1		33
Montcalm x C.I. 7248																				
Montcalm								1	4	9	10	8	5	5	3	2	4	1		52
C.I. 7248										2	4	4	5	3	2	3	2	2	2	27
F ₂							2	2	8	5	4	4	3		2		2	1		34
C.I. 7155 x C.I. 7248																				
C.I. 7155											1	3	5	5	6	3	2	2	1	31
C.I. 7248										2	4	4	5	5	2	3	1	2	2	27
F ₂										1	2	5	5	4	3	3		2	1	27

In 4 crosses the F_1 plants were later than the early parents and the F_2 generations were later than the early parents but earlier than the late parent by 2 to 4 days.

This summary is based on the results of the first three dates of seeding. The last replication, seeded on June 15, 1950, shows a great deal of fluctuating variation in time of heading both in the F_2 generations as well as in the parental lines (see Table 20).

(c) Observations on Variation

It was observed that even in the pure lines the individuals varied somewhat in time of heading. The results presented in Tables 17 to 20 confirm this observation. Even in the early lines, such as Atsel and Tulare, there was a much wider range in the heading periods for the third and fourth dates of seeding. These differences are presumably due to environmental conditions which are very difficult to equalize exactly.

In all 16 crosses studied, there were two noteworthy facts about the F_1 : first, that the mean in each case was approximately intermediate between that of the two parents (see Table 16); and second, that the variability in the F_1 was about as low as that of the parents. In the F_2 populations the behaviour was quite different. The means were close to those of the F_1 , but, due to segregation in the F_2 populations, variability had been greatly increased.

Fig. 9

F₂ generation of a late by early cross Montcalm x Tulare showing segregation for both parental types.

The F_2 families showed a range in heading dates comprising the F_1 range with other plants approaching the early and late extremities of the P_1 parents. Fig. 9 shows a section of the F_2 family of the cross 2x, Montcalm x Tulare.

Montcalm is a late, semiprostrate variety and Tulare an early erect line. In the F_2 population (see Fig. 9) are plants intermediate in type and plants varying towards the parental extremes. Early heading is evidenced in the more erect types while some plants in the same family are yet in a state of vegetative growth resulting in late heading.

There were indications in the field study that there were soil differences in the area on which the barley was grown. These evidently affected the time of heading, though the land comprised an area of only 18.5 feet by 128 feet and seemed uniform.

4. Interpretation of F_2 Results

(a) The Hypothesis

The results obtained from these crosses indicate that earliness is dominant or partially dominant to lateness. In all the large F_2 populations, the distribution according to time of heading gives a skew curve having the mode near the beginning of heading period.

Both the F_1 and the F_2 data give evidence in support of the hypothesis that the time of heading is dependent upon multiple factors which together produce a cumulative effect. These conclusions are supported further by the fact that F_2 segregates earlier than the early parents were secured in eight different crosses:

Atsel x Tulare	- Table 17
Montcalm x Tulare	- Tables 17, 18, 19, and 20
Titan x Newal	- Tables 17 and 20
Montcalm x Newal	- Tables 17, 18, 19, and 20
Montcalm x Titan	- Tables 17, 18, and 20
Montcalm x C.I. 7248	- Tables 17, 18, 19, and 20
Montcalm x C.I. 7155	- Table 18
C.I. 7155 x C.I. 7248	- Tables 17 and 19

In some crosses there was also transgressive F_2 segregation for lateness as in:

Atsel x C.I. 7155	- Table 17
Montcalm x Newal	- Tables 17, 18, and 20
Montcalm x Titan	- Tables 17 and 18
Montcalm x C.I. 7155	- Table 17
Montcalm x Beecher	- Table 18
Titan x Newal	- Table 19
C.I. 7155 x C.I. 7248	- Table 17

Fourteen out of the 16 F_2 populations showed an approximate 1-in-16 ratio for grandparental recovery. This would suggest a dihybrid inheritance for earliness. The other two F_2 families, Beecher x C.I. 7248 and C.I. 7155 x C.I. 7248, tend to support a condition of monohybridity or single-gene difference.

The occurrence of transgressive segregation in 12 out of 16 F_2 populations is further evidence that at least two factor pairs are responsible for earliness. If this holds true generally, then two varieties of barley or other cereal may differ genetically for earliness in one of two ways, such as:

(A) One or more dominants may be found in one parent while the other has the recessive allelomorphs. The genotype of one parent may thus be represented as AABB and the other as aabb. Such parents when crossed would give an F_1 population like the early parent in time of heading and an F_2 population with heading ranging from the early to the late parent.

Such a relationship of parents seems to have existed in crosses 2x, 3x, 7x, and 9x. (See Table 16 for F_1 heading and Tables 17 to 20 for the heading in F_2).

(B) One or more dominants may be present in each parent while lacking in the other. Thus if the genotype of one parent is represented as AABbCc, the other would appear as aabbCC. A cross involving the two types of parents should

give an F_1 population earlier than either parent and an F_2 ranging in heading from earlier than either parent to later than either. This relationship was evident in cross 1x. (See Table 16 for F_1 and Tables 17 to 20 for F_2 heading).

It seems likely that in some of the crosses included in this study, there may be one or more factors causing unlike rate of development under different conditions. For example in cross 14x the parents headed 52 days apart in the greenhouse (see Table 16) but only 16 days apart in the field (see Table 19).

(b) The Quantitative Effect of the Genes

One object of the present study was to determine the quantitative effect of the factors. The influence of the factors upon time of heading was shown to vary with different dates of seeding. Because of the variability in heading from early to late seedings, it was not satisfactorily proved that the factors are the same in their effect.

When the F_2 populations are compared with the early parents it appears that the factors may have approximately the same effect and that each caused an average difference of one and one-half to two days in time of heading. Transgressive

segregation in the F_2 strongly suggests a quantitative inheritance for earliness. This effect is presented in Fig. 10 which shows the F_2 and parental types of the cross 2x, Montcalm x Tulare.

The Montcalm parent appears on the left and the Tulare parent on the right of the F_2 hybrids. The cumulative effect of the factors for earliness is seen in the heading of the F_2 population which is earlier than the early Tulare parent.

Fig. 10

Upper: Montcalm parent, left and F_2 generation, right.
Lower: F_2 generation left and Tulare parent, right.
Transgressive segregation is seen in the hybrid population.

VI. GENERAL DISCUSSION AND CONCLUSIONS

The experimental results of the three aspects of earliness studied have been analyzed, and with respect to the more specific points, more or less fully discussed in the sections devoted to the respective experiments.

In discussing the scope of the present study and the results secured, it must be recognized that the general and exploratory nature of the work does not permit final and conclusive observations. Certain points related to the work as a whole should however be discussed, particularly those regarding the physiology and nature of the inheritance of earliness.

It was shown that there was a negative correlation between the number of tillers and the number of days to heading. The later varieties, as a rule, carried more tillers than the early or midseason lines. With a reduced number of tillers, a morphological characteristic, it is probable that there is more rapid assimilation and utilization of soil nutrients promoting speedy growth and development. The association established in the morphological study between leaf curling and earliness may also be based on the rate at which the various physiological stages are acquired. The relation of flag leaf to spike seems to hold definite promise as a morphological character in the study of earliness.

Evidence has been presented to show that exposure to high temperature does not induce development unless accompanied by an adequate photoperiodic treatment. The results of microdissection, referred to by Hamner (12), have proven that the rate of development of the primordia is influenced by the number of hours of daily exposure. This phase of the work on the relation of photoperiod to earliness, substantiates the physiological theory of "phasic development" postulated by Klebs (21) and Lysenko (23).

The study on the relation of height of plant to earliness did not yield any definite results. There was a great deal of fluctuating variation in height even within the pure lines. The average height for the third and fourth dates of seeding was the same, but the third was earlier than the fourth date of seeding by an average of 1.5 days. Thus, the height attained by the plant may be considered as having no direct relation to the number of days to heading.

The acceptability of the genetic analysis rests in the assumption that parents might be alike in time of heading and contain different dominant factors for earliness. Most of the crosses included in this study seemed to have such a relationship of parents for in several of them the F_1 as well as the F_2 began heading earlier than the early parent.

In some crosses in which the F_2 began heading earlier than the early parent, there were no plants later than the late

parents. However, the expected proportion of the population later than the late parents in such crosses would be small. This behaviour further supports the assumption that the genotype of one parent may be AABB and the other aabb.

The limitations imposed by small F_2 populations made it difficult to conclude at this stage of the study which crosses would result in early lines as the segregation of all possible intermediate and extreme types could not be accounted for with the small families used.

In order to establish conclusively the mode of inheritance of earliness it is necessary that homozygous races be recovered. This would require the growing of F_4 generations from selected F_3 families. However, the scope of the present study serves to indicate the trend, that is, the time of heading is dependent upon multiple factors having a cumulative effect.

There were indications of soil differences in the area on which the F_2 generations and parental plants were grown which affected the time of heading. In order to facilitate accuracy in the results, soil differences could be carefully marked on the planting plans so that it be known on which soil type each row is located. The hybrid generations and parent varieties could then be planted in duplicate on each soil type.

Present data do not prove conclusively that the genes or factors have the same effect, but the results point to a strong

possibility, and that each factor caused an average difference in earliness of from one and one-half to two days.

It is hoped that some crosses initiated later between the varieties now involved, as well as between races and varieties with wider date-of-heading differentials will prove to be monohybrid crosses. In this way the quantitative effects of some factors could be determined with greater certainty and genetic analyses facilitated.

ACKNOWLEDGEMENTS

The writer is indebted to Dr. L. P. V. Johnson, Professor of Genetics and Plant Breeding, for suggestions and criticisms during the organization and progress of these investigations, to Dr. J. H. Whyte, Associate Professor of Botany, for helpful suggestions on the physiological phase of the study, to Dr. J. Unrau, Assistant Professor of Genetics and Plant Breeding, for photomicrographic technique, to the Department of Plant Science, for material aid and equipment, and to Mr. Gilbert Paul for capable routine assistance.

LITERATURE CITED

1. AAMODT, O. S. The inheritance of growth habit and resistance to stem rust in a cross between two varieties of common wheat. J. Agr. Research, 26:457-469. 1923.
2. ÅBERG, E. and WIEBE, G. A. Classification of barley varieties grown in the United States and Canada in 1945. Tech. Bull. 907. U.S D.A., 11-27. 1946.
3. BIFFEN, R. H. Mendel's laws of inheritance and wheat breeding. J. Agr. Sci., 1:4-48. 1905-6.
4. BRYAN, W. E. and PRESSLEY, E. H. Inheritance of earliness in wheat. Thirty-second Ann. Rpt. Ariz. Agr. Exp. Sta., 603-605. 1921.
5. CLARK, J. A. Segregation and correlated inheritance in crosses between Kota and Hard Federation wheats for rust and drought resistance. J. Agr. Research, 29(1):1-47. 1924.
6. CLARK, J. A. and HOOKER, J. R. Segregation and correlated inheritance in Marquis and Hard Federation crosses, with factors for yield and quality of spring wheat in Montana. U.S.D.A., Dept. Bull. 1403, 1-71. 1926.
7. EGUCHI, T. Experiments with short-day treatments. Proc. Imp. Acad., Tokyo, 13:332-333. 1937.
8. EMERSON, R. A. and EAST, E. M. Inheritance of quantitative characters in maize. Neb. Agr. Sta. Re. Bull. 2. 1913.
9. FLORELL, V. H. Studies on the inheritance of earliness in wheat. J. Agr. Research, 29(7):333-347. 1924.
10. GARNER, W. W. and ALLARD, H. A. Further studies in photoperiodism, the response of the plant to the relative length of day and night. J. Agr. Research, 11:871-920. 1923.

11. GREGORY, F. G. and PURVIS, O. N. Photoperiodic treatment and plant response. Ann. Botany, London, N.S., 2: 237-251. 1938.
12. HAMNER, K. C. Flower stimulation in long-day plants. Botan. Gaz., 101:658-687. 1940.
13. HARRINGTON, J. B. and HORNER, W. H. The reaction of wheat varieties to different dates of sowing. Sci. Agr., 16:127-134. 1935.
14. HAYES, H. K. and EAST, E. M. Improvement in corn. Conn. Agr. Exp. Sta., Bull. 168. 1911.
15. HAYES, H. K. and JONES, D. F. The effects of cross and self-fertilization in tomatoes. Conn. Agr. Exp. Sta. Rpt., 305-308. 1916.
16. HOSHINA, Y. On the inheritance of the flowering time in peas and rice. Abstract by Castle, W. E., Genetics and Eugenics, 213-215. 1915.
17. HUTCHESON, T. B. and WOLFE, T. K. The effects of fertilizers and hybridization on maturity and yield in corn. Va. Agr. Exp. Sta., Tech. Bull. 27. 1924.
18. JASNY, N. Competition among grains. Food Res. Institute, Stanford University, Calif. 1940.
19. JOHNSON, L. P. V. Inheritance studies in barley. Rpt. to Nat. Research Council. 1948.
20. KEEBLE, F. and PELLOW, C. The mode of inheritance of stature and of time of flowering in peas. J. of Genetics, 1:47-56. 1910.
21. KLEBS, G. Developmental phases in plants. Flora, 112:128-151. 1918. Abstract, original not seen.
22. LEAKE, H. M. Studies in Indian Cotton. J. of Genetics, 1:47-56. 1911.
23. LYSENKO, T. D. Theoretical basis of vernalization. Trudy Azerbaidzan Op. Sta., No. 3, 168. 1928. From abstract.

24. MENDEL, G. Mendel's principles of heredity, Translation of "Experiments in Plant Hybridization", by Bateson, W. 1909.
25. PEARSON, O. H. The influence of inbreeding upon the season of maturity of cabbage. Proc. Am. Soc. Hort. Sci., 28:359-366. 1931.
26. PLATT, A. W. The effect of soil moisture on the frost reaction of wheat, oats, and barley seedlings. Sci. Agr., 17:616-626. 1937.
27. POWERS, L. The nature and interaction of genes differentiating habit of growth in a cross between varieties of Triticum vulgare. J. Agr. Research, 49:573:605. 1934.
28. PURVIS, O. N. Vernalization by exposure of low temperatures. Ann. Botany, London, 48:919-955. 1934; 8:285-314. 1944.
29. SUNESON, C. A. Survival of four barley varieties in a mixture. Agronomy J., 41:459-461. 1949.
30. THOMPSON, W. P. The inheritance of the length of the flowering and ripening periods in wheat. Proc. Trans. Roy. Soc. Can. 3(Sec. IV-V):69-87. 1918.
31. _____ The inheritance of earliness and lateness in wheat. Proc. Trans. Roy. Soc. Can. 3(Sec. IV-V):143-162. 1919.
32. _____ Earliness in wheat and its inheritance. Sci. Agr., 1(5):193-199. 1921.
33. TINCKER, M. A. and JONES, M. G. The influence of climatic factors upon the growth of a spring sown variety "Record". Ann. of App. Biol., 18:187-201. 1931.
34. TSCHERMAK, E. V. Veber die Verebung der Blutezeit bei Erbsen. Verh. Naturf. Verein. Brunn. XLIX. Original not seen. Abstract by Noll, C. F. Tech. Bull. 194, Penn. State College. 1925.
35. VESELOVSKII, I. A. Some new achievements in breeding potatoes for earliness. Am. Potato J., 14:295-297. 1937.