FIRST BOOK

OF

ZOÖLOGY

PL 62431

Library
of the
University of Toronto

First ed. is NY 1875

Digitized by the Internet Archive
in 2018 with funding from
University of Toronto

ZOÖLOGY.

STANDARD SCIENTIFIC BOOKS.

THE OBSERVING POWERS OF CHILDREN, an Essay on the Culture of, especially in connection with the Study of Botany. By ELIZA A. YOUMANS. Edited, with Notes and a Supplement, by Joseph Payne, F.C.P., Author of 'Lectures on the Science and Art of Education,' &c. Crown 8vo. 2s. 6d.

FIRST BOOK OF BOTANY. Designed to cultivate the Observing Powers of Children. By ELIZA A. YOUMANS. With 300 Engravings. New and Enlarged Edition. Crown 8vo. 5s.

'It has been everywhere welcomed as a timely and invaluable contribution to the improvement of primary education.'—PALL MALL GAZETTE.

HISTORY OF THE ENGLISH REVOLUTION OF 1688. By C. D. YONGE, Regius Professor, Queen's College, Belfast. Crown 8vo. 6s.

'The volume before us is, on the whole, a fair, succinct, useful, and masterly summary of the main causes, circumstances, and history of the Revolution.'

STANDARD.

THE EXPANSE OF HEAVEN. By R. A. PROCTOR, B.A. A Series of Essays on the Wonders of the Firmament. With a Frontispiece. Second Edition. Crown 8vo. 6s.

'The worth of the book consists in its easy adaptation of modern scientific knowledge to the needs of a large class of readers. It may be confidently recommended.'

DAILY NEWS.

OUR PLACE AMONG INFINITIES. By R. A. PROCTOR, B.A., Author of 'Saturn and its Systems,' 'The Universe,' 'The Expanse of Heaven,' &c. To which are added Essays on 'ASTROLOGY,' and 'THE JEWISH SABBATH.' Crown 8vo. price 6s.

SCIENTIFIC CULTURE. By JOSIAH P. COOKE, Jun., Professor of Chemistry and Mineralogy in Harvard College. Cr. 8vo. cloth, 1s.

A CLASS BOOK OF CHEMISTRY. On the Basis of the New System. By EDWARD L. YOUMANS, M.D. With 200 Illustrations. [Immediately.]

*** A List of Elegant Gift Books, suitable for Presents and Prizes in Schools, may be had on application to

HENRY S. KING & CO., London.

FIRST BOOK OF ZÖÖLOGY.

BY

EDWARD S. MORSE, PH.D.

LATE PROFESSOR OF COMPARATIVE ANATOMY AND ZÖÖLOGY IN BOWDOIN COLLEGE.

‘As for your pretty little seed-cups, or vases, they are a sweet confirmation of the pleasure Nature seems to take in superadding an elegance of form to most of her works, wherever you find them. How poor and bungling are all the imitations of art! When I have the pleasure of seeing you next we will sit down—nay, kneel down if you will—and admire these things.’

HOGARTH in a *Letter to Ellis*.

HENRY S. KING & CO., LONDON.

1876.

(The rights of translation and of reproduction are reserved.)

P R E F A C E .

THE "First Book of Zoölogy" is expressly prepared for the use of pupils who wish to gain a general knowledge of the structure, habits, modes of growth, and other leading features concerning the common animals of the country.

Particular attention has been given to the lower animals, as these are more often neglected in text-books. Directions for collecting, the preparation of specimens for the cabinet, and the haunts of the animals to be studied are given, and the pupil is expected to study, with the book in one hand, and the specimens in the other. The figures illustrating this work, with a few exceptions, have been drawn from Nature by the author, and have been prepared with especial reference to their being copied by the pupil. To facilitate this the figures are made in outline, with the shaded side of the figure indicated by darker lines.

The necessity of the pupils copying (however poorly) the figures, either upon the slate, or upon paper, cannot be too strongly urged.

From his own experience, the author has learned that a specimen or figure may oftentimes be carefully studied, and yet only an imperfect idea be formed of it; but, when it had been once copied, the new points gained repaid all the trouble spent in the task.

It makes but little difference whether the pupil is proficient in drawing or not; it should be strenuously insisted upon by the teacher that the pupils copy, as far as possible, the figures contained in each lesson.

To collect in the field, to make a cabinet, and then to examine and study the specimens collected, are the three stages that naturalists, with few exceptions, have passed through in their boyhood.

If one recalls the way in which boys first manifest their taste for such studies, he will remember that first a few examples were brought together; a collection was made. It may have been birds'-eggs, insects, or shells; then little boxes, a case of drawers, or shelves, were secured to hold their treasures. In thus collecting and arranging and rearranging the cabinet, the eye becomes familiar with the outline and general character of the objects, and in this way the mind is finally prepared to comprehend the relations existing

between animals, and to appreciate the leading points upon which classification is founded.

Agassiz invariably placed before his students a single specimen, or a box full of specimens, and told them to look and see what they could find out.

It has seemed, therefore, that the way to commence the study of zoölogy is to follow the course one naturally pursues when he is led to the study by predisposition. Nor is it essential, at the outset, to present the entire range of the animal kingdom. Teach the characters of one or two great divisions first, and then the pupil is better prepared to grasp in turn the other divisions. The persistent attempt, in all text-books of this kind, to give some attention to every large group in the animal kingdom, has often resulted in wearying and confusing the minds of those who take up the study for the first time.

A very serious difficulty is encountered in those books which give a more or less complete view of systematic zoology for beginners. In some, the authors commence with the lowest forms, and end with the highest. In others, the highest animals are dealt with first, and the lessons end with the lowest. The first mistake made is the attempt to teach systematic zoölogy, where the pupil is quite ignorant of the material to be classified; and proper familiarity with the objects, the author contends, can only be acquired

by collecting the specimens and forming a little cabinet of them.

The difficulty, however, arises in commencing the lessons with either the lowest or the highest animals. If the author commences with the lowest animals, he deals at the outset with creatures which the pupil in certain cases can never see, as many of the animals to be considered are microscopic, and most of them of such a nature that their soft parts cannot be preserved. On the other hand, if the author commence with the vertebrates, he presents, point-blank, some of the forms of structure most difficult to understand.

The main thing at the outset is to teach the pupil how to collect the objects for study; this leads him to observe them in Nature, and here the best part of the lesson is learned: methods of protection for the young, curious habits, modes of fabricating nests, and many little features are here observed, which can never be studied from an ordinary collection. Hence, collecting in the field is of paramount importance. Next, the forming of a little collection at home prompts the pupil to seek out certain resemblances among his objects, in order to bring those of a kind together. In this way he is prepared to understand and appreciate methods of classification. Finally, having grasped the leading features of a few groups, he is enabled to comprehend the character of cognate groups with less difficulty. Thus,

an inland student, having got the typical idea of an insect from the study of a common grasshopper, for example, is much better prepared to understand the general structure of the Crustacea, though he may never have seen the few forms peculiar to fresh water. In the same way after having studied the common earthworm, he can form a better idea of the complicated structure of many marine worms, though these he may never see. After long deliberation, and some hesitancy, the author is forced to depart from common usage, and present, in this first book, only a few of the leading groups in the animal kingdom.

From the abundance of material, and the comparative ease with which the specimens may be preserved for cabinet use, shells and insects have always formed the favorite collections of children. They are the most common objects in nearly all collections, and it has seemed to the author that here the pupil ought to commence his studies.

Having learned to collect and prepare specimens for the cabinet, and to observe the relations and differences existing among them, the pupil is then prepared to go on to forms less familiar, or to study in detail the material already gone over.

Great pains have been taken to present, in every case, drawings made from the animal, expressly for this book. With but very few exceptions, these illustrations are en-

tirely new. It is believed that teachers will appreciate the absence of those hackneyed illustrations which have too long done service in text-books on the subject.

To those especially interested in the study, many figures are given which have never before been published, even in scientific works, as, for example, the rare *Lymnæa megasoma*, *Lymnæa ampla*, *Ptyelus lineatus*, and many others.

The propriety of giving the technical names has been carefully considered, and the decision to leave them out has been only arrived at after long deliberation. All who desire to know these names can find access to some of the books given in a list for reference on the last page of this volume, wherein full descriptions of the species oftentimes accompany the names. The main reason, however, in not giving the technical names was, that at present the names are changing so rapidly that only the specialist in each department pretends to keep up with modern names. An eminent naturalist has stated that in the present condition of nomenclature the common name of an animal was oftentimes more reliable than the scientific one. The names learned to-day would with few exceptions be of no use ten years hence. More than this, they would be an absolute hindrance.

An extreme instance might be given regarding one of our most common animals of the coast—the common sea-urchin.

This creature is still known by many as *Echinus granulatus*, this name being given within twelve years by Agassiz in his "Methods of Study in Natural History." The following list shows the successive changes in its name since that time :

Echinus granulatus.

Euryechinus granulatus.

Toxopneustes dröbachiensis.

Euryechinus dröbachiensis.

And at this date of going to press this unfortunate creature bears the name of

STRONGYLOCENTROTUS Dröbachiensis.

And yet this little animal will be known by the name of Sea-urchin long after the time when the systematists and their distracting nomenclature will have been forgotten.

E. S. M.

November 1875.

CONTENTS.

CHAPTER I.

	PAGE
FRESH-WATER SHELLS	1
Where to collect, 1; Empty or Dead Shells, 2; Parts of a Shell named, 3; Forms of Spire, Dextral, and Sinistral Shells, 4; Lines of Growth, Growth of a Shell, 5, 6.	

CHAPTER II.

FRESH-WATER SNAILS	7
Where and how to collect, 7, 8; Live Shells, 8; Head, Tentacles, Foot, how they breathe, 9, 10; Operculum, 12; Gills, 13.	

CHAPTER III.

LAND SNAILS	14
Where to collect, 14; Parts of the Animal, 15, 16; Tentacles, Jaw, 17; Explanation about an Object being enlarged, 18; Slug, 19; Eggs and Young, 20; Mantle, 21.	

CHAPTER IV.

SEA SNAILS	21
Where to collect, 22; Entire Aperture, Aperture Notched or Canaliculated, 23; Siphon, and its Uses, 24; Flesh and Animal Feeders, 25; Egg-cases or Capsules, 26, 27.	

CHAPTER V.

FRESH-WATER MUSSELS	27
Where found, 27; Ligament, Lines of Growth, 28; Regions of Mussel, 29; How they crawl, 30; Excurrent and Incurrent Orifices, 31; Siphons, Mantle 32; How Pearls are formed, 33; Internal Markings of Shell, Hinge, 34; How the Shells open and Ligament acts, 36.	

CHAPTER VI.

	PAGE
CLAMS, MUSSELS, AND OYSTERS	37
Salt-water Bivalves, 37; Parts of a Clam, Siphons, Currents of Water, and how they eat, 38, 39; Ligament, and Internal Markings of Shell, 40; Byssus, 42; Siphonal Tubes separate, 44; Oysters, 45; Heart, Palpi, Liver, Gills, Mouth of Oyster, 46, 47; Relations between Clams, Mussels, and Snails, 48.	

CHAPTER VII.

COLLECTING INSECTS	49
How to collect and arrange, 49; To make Insects' Boxes, 50; To kill and pin Insects, 51, 52; Where to find Insects, 52, 53.	

CHAPTER VIII.

PARTS OF AN INSECT	53
True Insects or Hexapods, Mouth-parts, 55; Antennæ, Eyes, 56, 57; Regions of Body, 58; Wings, 59, 60; Position of Wings, 61; How to Spread the Wings, 62; Beetle spread, 63; Abdomen, 63, 64.	

CHAPTER IX.

PARTS OF AN INSECT (continued)	65
Parts of the Thorax, 65; Beetle dissected, Mouth-parts, 66-69; Appendages of the Segments, 70.	

CHAPTER X.

GROWTH OF INSECTS	71
Collecting Eggs, Caterpillars, Chrysalides, 71, 72; Where to collect, 73; Eggs, 74; Raising Caterpillars, 75, 77; Chrysalis, 78, 79; Development of Chrysalis, 80; Incomplete Changes, 81; Complete and Incomplete Metamorphosis, 82; Difference between Larva and True Worms, 83-85.	

CHAPTER XI.

HABITS AND STRUCTURE OF INSECTS	85
Mud-wasp, 85-87; Mosquito, 88.	

CHAPTER XII.

HABITS AND STRUCTURE OF INSECTS (continued)	89
Young Grasshopper, 89; Cast-off Skin of Grasshopper, 90; Curious Accidents, 91; Air-tubes, 92; How Insects breathe, and how some rest, 93; Sounds produced by Insects, 94; Stridulation of Grasshopper, 95, 96.	

CHAPTER XIII.

	PAGE
HABITS AND STRUCTURE OF INSECTS (continued) . . .	97
Spittle Insect, 97, 98; Cicada or Seventeen-year Locust, 99, 100.	

CHAPTER XIV.

HABITS AND STRUCTURE OF INSECTS (concluded) . . .	101
May Fly or Ephemera, 100-103; Caddis Worm, 104; Gall Fly, 104, 105; Other Galls, 106.	

CHAPTER XV.

SPIDERS	109
Parts of a Spider, 110; Mouth Parts and Eyes, 111; Spinnerets, 112; Legs, 113; Habits of some, 114; Net Building, 115; Ballooning Spiders, 116, 117; Egg Cases, Young Spiders, 117, 118, Care for the Young, 119, 120.	

CHAPTER XVI.

DADDY-LONG-LEGS, CENTIPEDES, AND MILLIPEDES . . .	121
Daddy-long-legs, 121-123; Centipedes, 123, 124; Mouth. Parts of, 125; Millepedes, 126; How they lay their Eggs, and Young, 127; Comparison between Myriopods and Insects, 127, 128; Some Characters of Insects, Spiders, and Myriopods, 129.	

CHAPTER XVII.

CRAW-FISH AND LOBSTER	130
Craw-Fish, where found, 130; Regions of, 131; Mouth Parts and Appendages, 132, 133; Gills, 134; How to arrange Parts, 135, 136; Moulting, 137; Young Lobster, 138.	

CHAPTER XVIII.

CRABS, HERMIT-CRABS, AND OTHER CRUSTACEANS . . .	138
Common Crab, 139; Eggs, and how they carry them, 140; Fin-like Legs, Oyster Crab, 141; Hermit Crab, 142-144; Fresh-water Crustaceans, 144; Sowbug and Eggs, 145, 146; Regions of Crustacea, 146, 147.	

CHAPTER XIX.

BIVALVE CRUSTACEANS AND BARNACLES	147
Characters of, 148, 149; Where found, 150; Appendages, how they secure their Food, 151; Young Barnacles, 152, 153.	

CHAPTER XX.

	PAGE
WORMS	153
Angle Worms, 154 ; How they move, 154, 155 ; Leech, 155 ; Sea-Worms, 156, 157 ; Tube-building Worms, 159.	

CHAPTER XXI.

CONCERNING NATURAL GROUPS	160
Characters of Articulates, 160 ; Of Worms and Mollusks, 161 ; Type Characters, 161 ; Typical Animals, 162 ; Systematic Tables and Classification, 163.	

CHAPTER XXII.

CHARACTERS OF VERTEBRATES	163
External Characters of Salamanders, 163, 164 ; Comparison with Cat, 165 ; Differences between the two, 166 ; Young Salamander, 166 ; Skeleton of Cat, 167 ; Vertebral Column and Spinal Cord, 169, 170 ; Bones of the Ankle and Wrist, 171, 172 ; Groups of Vertebrates, 173, 174.	

CHAPTER XXIII.

BONES OF THE LEG AND WING OF BIRDS	174
Importance of studying Young Forms, 174 ; Wing Bones of an Adult Bird, 175 ; Embryo Birds, and Wing Bones of same, 176 ; Leg Bones of an Adult Bird, 177 ; Leg Bones of Embryo Bird, 178 ; Leg Bones of Young Toad, 180.	

CHAPTER XXIV.

CLASSES AND SUB-KINGDOMS	181
Sub-Kingdom Mollusca, Class Gasteropoda, 181 ; Class Acephala, 182 ; Sub-Kingdom Arthropoda, Class Insecta, 183 ; Class Myriopoda, 184 ; Class Arachnida, 185 ; Sub-Kingdom Vermes, Class Annelida, 185 ; Class Crustacea, 186 ; Sub-Kingdom Vertebrata, 187.	
NOTICE TO TEACHERS	189

FIRST BOOK OF ZOÖLOGY.

CHAPTER I.

FRESH-WATER SHELLS.

For these lessons, it has been deemed best to commence with the shells of mollusks, such as snail-shells and mussels. They are better objects to examine than insects, being more simple in structure, and less liable to be broken in handling. When found alive, their habits can be readily studied, as they can easily be kept alive in jars filled with water.

1. Let the pupils first make a collection along the shores of some lake or river, picking up all the different kinds of shells they meet with. The waves will have thrown them up on the shores, or in times of drought the waters will have left them exposed. Certain kinds are very small, though they will be found by sharp looking. Most of the shells collected will be empty, and these shells are called *dead shells*, because the soft-bodied creatures once contained in them have died and decayed, leaving the hard, limy shells. Some of the shells collected may contain the animal, and at one time each of them possessed a little creature within, which was the fabricator of the shell.

2. Remember that the shell is not a house built by the snail, as a wasp builds its nest, but the shell is a part of the animal, and is connected to it by certain muscles, so that it cannot leave the shell, as many suppose.

The empty, or dead, shells, are to be studied first.

Looking over the shells collected, we shall find some of the following kinds :

FIG. 1.—FRESH-WATER SHELLS.

A number of fresh-water mussel-shells, also, will probably be collected.

FIG. 2.—FRESH-WATER MUSSEL-SHELL.

These are to be reserved for future examination. Those

having a spiral turn or twist are called snail shells, and are to be studied first.

3. Let the pupils pick out from their collections the shells like these :

FIG. 3.—FRESH-WATER SNAIL-SHELLS.

The different spiral turns, or twists, are called *whorls*, and the whorls together form the *spire*. The opening into the shell is called the *aperture*, and the line separating the whorls is called the *suture*. The pointed end of the spire is called the *apex*.

FIG. 4.—SHELL, WITH PARTS NAMED.

In some shells the spire is *elongated*; in others the spire is *short*; in others still the spire is *depressed* or *flattened*.

FIG. 5.

4. If the shell is held in the hand, with the aperture toward the holder, and the spire pointing upward, as in the figures drawn, the aperture will be either toward the right hand, or toward the left hand. In the figures already given, the aperture is on the right hand, and these shells are called *dextral*, or right-handed, shells.

Shells having the aperture on the left hand when held in the way above described, are called *sinistral*, or left-handed, shells. Let the pupils here examine all the shells they have collected, holding each one with the spire pointing upward, and the aperture toward them, and separate the *dextral* shells from the *sinistral* shells. As sinistral shells are not so common as the other kind, it may be that none will be found in the first collections made by the pupils. The following is a figure of a sinistral shell:

FIG. 6.—SINISTRAL SHELL.

5. If the surface of the shell be examined closely, delicate lines running from one suture to another will be seen, as in the figures already given ; and, if the shell be looked at from the side of the aperture, these lines will be found running parallel to the edge of the aperture, or *lip*, as it is called.

These delicate lines are called *lines of growth*.

FIG. 7.—SHOWING LINES OF GROWTH RUNNING PARALLEL TO THE EDGE OF THE APERTURE.

The shell is increased in size by successive layers of shelly matter added to the borders of the aperture. In this way the shell grows.

A clearer idea of the growth of a shell may be obtained by studying the next figure (Fig. 8): *A* representing in outline a young shell ; *B* representing the full-grown shell in outline ; and *C* representing the same outline as *B*, with a number of lines of growth represented upon it.

If the shell were now to continue its growth a single half-turn, or *whorl*, the dotted lines would indicate the increased stages it would assume : *a* representing the first increase in size, *b* the next stage, and *c* the appearance of the shell when the additional half-whorl has been completed.

FIG. 8.—ILLUSTRATING THE INCREASE IN SIZE OF A SHELL.

Among the lot of shells collected by the pupil, different-sized ones will be found. Now, if a number of these, of different sizes, can be picked out, provided they belong to the same kind, or species, it will be noticed that the apex of all of them will be the same; but that the shells have increased in size at the aperture, and the aperture will be larger, and the larger shells will have more whorls than the smaller shells. The following figures illustrate four different ages of the same species of shell:

FIG. 9.—ILLUSTRATING DIFFERENT AGES OF THE SAME SHELL—THE LOWER FIGURES REPRESENTING A VIEW OF THE SHELL FROM THE APEX.

6. The axis around which the whorls revolve is called the *columella*. This axis is generally solid, though in many shells it is hollow, as if the whorls had turned around a shaft which had afterward been withdrawn. This hollow axis looks like an opening in the base of the shell, as in the following figure :

FIG. 10.—A SNAIL-SHELL SEEN FROM BELOW.

This opening is called the *umbilicus*. The apex of the shell is sometimes called the *nucleus*, because the shell commences to grow from this point.

CHAPTER II.

FRESH-WATER SNAILS.

7. THE pupils will now be required to bring in some live snails. Let them examine bits of bark, chips, or branches, found in ditches, or muddy brooks. Under lily-pads and on the stems and leaves of other aquatic plants, and on stones in rivers, snails of various kinds will be found. A dipper with the bottom perforated, or made into a sieve, and attached to a wooden handle four or five feet in length, will be found useful in scooping up the sand or mud from the

bottom of rivers and ditches. The dirt having been sifted out, the shells and other objects will be left behind. The dipper may be made as in the figure.

FIG. 11.—DIPPER ATTACHED TO A WOODEN HANDLE FOR COLLECTING SNAILS.

Shells collected with the snails inside, and cleaned for the cabinet, are called *live shells*. They are always more fresh and perfect than dead shells.

Having made the collection, the snails should be kept alive in a wide-mouthed jar, or bottle, care being taken not to have more than fifteen or twenty in a jar holding a quart of water.

8. The pupils will have secured some of the following forms:

FIG. 12.—FRESH-WATER SNAILS.

The broad, creeping disk upon which the snail rests, and by which it retains its hold to the glass, is called the *foot*. The snail moves about, and crawls or glides slowly along, by means of the foot.

The two little horns or feelers, in front, are called *tentacles*, and, as the snail moves, the tentacles are seen stretched out in front, and occasionally bending, as if the creature were feeling its way along. The eyes are seen at the base of the tentacles, as two minute black dots. The mouth is between the tentacles, and below. The part from which the tentacles spring is called the *head*, and the opposite end of the body is called the *tail*. The surface upon which the snail rests is called the *ventral* or lower surface, and consequently that portion of the body which is above is called the *dorsal* surface, or back.

9. The pupil, in watching the habits of the snails he has collected, will notice some of them crawling to the surface of the water to breathe air. The snail accomplishes this by raising the outer edge of the aperture to the water's edge, and then opening a little orifice in the side, through which the air enters to the simple lung within.

This orifice is on the right side in those snails having dextral shells, and on the left side in those snails having sinistral shells.

Many kinds of snails which live in fresh water are called air-breathers, because they are forced to come to the surface of the water to breathe air. In doing so they first expel a

bubble of air, which may be seen escaping from the breathing-orifice, as in Fig. 14, *B*.

A view from the back,
or *dorsal view*.

side-view.

ventral view.

t t, Tentacles.
e e, Eyes.
b, Breathing-orifice.
m, Mouth.

FIG. 13.—A FRESH-WATER SNAIL SEEN FROM ABOVE, FROM THE SIDE, AND FROM BELOW.

These fresh-water air-breathing snails may be kept under water for many hours before life is extinct.

FIG. 14. —Jar of water, in which is contained a number of species of mollusks, some of which have already been studied. Some of them are near the surface, breathing air: *A* and *C* are taking in air; *B* is just expelling a bubble of air from the lung; *D* is crawling on the surface of the water; *E*, *G*, and *I*, are in the act of crawling up, to get a fresh supply of air; and *J* is a water-breather, having gills, but no lung.

10. Among the snails collected, there will probably be found some which have a peculiar scale on the hinder part of

the body. When the snail crawls, this scale will be seen just behind the shell, as in Fig. 15, *o*.

This scale is called the *operculum*, and when the snail has contracted, or drawn within the shell, the operculum is seen to fit the aperture of the shell, closing the shell as a stopper closes the mouth of a bottle.

FIG. 15.—SNAIL WITH OPERCULUM.—*o*, Operculum; *e*, Eye; *r*, Rostrum; *g*, Entrance to Gill-Cavity.

Nearly all sea-snails, that is, snails which live in salt-water, and many species of fresh-water snails, and also many snails which live in damp places on the land, and which are called land snails, have an operculum.

When the snail has retired within the shell, the operculum will look like this in the aperture of the shell (Fig. 16):

A series of concentric lines will be seen marking the operculum, and these are the lines of growth, the operculum growing around the outer edge by successive additions, just as the shell grows by successive additions to its outer margin.

The Western rivers teem with species of snails having opercula.

11. If the pupil has any of these operculated snails alive,

FIG. 16.—APERTURE OF SHELL CLOSED BY OPERCULUM, *o*.

he will observe that they do not come to the surface to breathe air.

Instead of a lung, the snail has a cavity containing an organ, or part, called the gill, by means of which it is capable of getting from the water what the air-breathing snail gets from the air, namely, *oxygen*.¹

It will be seen that the head of the snail is shaped differently in the snails having an operculum, the mouth being at the end of a sort of proboscis or *rostrum*. (See Fig. 15.) The shells, too, are, as a general thing, more solid.

12. Thus far the pupils have examined those snails which live in fresh water. Some of these were air-breathers, and came to the surface of the water at intervals to breathe air. He has studied other fresh-water snails which did not breathe air directly, but performed this function by means of

¹ If the class is sufficiently advanced, the teacher may here explain about oxygen and what the blood requires, and gets by respiration.

an organ called the gill. And these snails were operculated, that is, they all possessed a little scale called the operculum, which closed the aperture tightly when the snail contracted within the shell.

He has also learned that the shells grow in size by successive additions of limy matter deposited around the free border of the aperture, and that the delicate lines which mark the surface of the shell, and which run parallel to the outer edge of the aperture, are lines of accretion, or lines of growth.

CHAPTER III.

LAND SNAILS.

13. THERE are many other species of snails which live out of the water altogether, though they are generally found in damp places; and these are called land snails, because they live on the land.

Let the pupils now endeavor to collect some land snails. By going to some hard-wood grove of maple, beech, or oak, and turning over the layers of dead leaves, old rotten logs, or pieces of bark, they will be sure to find some specimens of land snails. Some of them do not grow larger than a pin's-head, others have shells as large as a walnut.

They are generally light brown in color, and the smaller species often have highly-polished or shiny shells.

The spire is generally depressed or flattened. In many,

the border of the aperture has a thickened white rim, or lip, as it is called. Such a collection having been made, the pupil will find among them some of the following kinds :

FIG. 17.—THE SHELLS OF LAND SNAILS.

14. The snail, as it crawls along, leaves a slimy trail after it. This trail consists of a fluid, which flows not only from the creeping disk, but also from the surface of the body. If the back of the snail is irritated by a sharp-pointed stick, a little whitish mass of this fluid, or mucus, will adhere to the end of the stick.

By placing the snail on a piece of glass, and allowing it to adhere and crawl on it, a good view may be obtained of the peculiar movements of this creeping disk, by looking through the glass from the other side.

The breathing-orifice may be found just within the aperture of the shell, on the right side of the snail.

This orifice will be seen opening and closing at intervals. (See Fig. 18, *b*.)

15. In the fresh-water snails there are but two tentacles upon the head. In the land snails with few exceptions, the

tentacles are four in number, a larger and a smaller pair. The larger tentacles are called the *superior tentacles*; the

FIG. 18.—LAND-SNAILS CRAWLING.—*b*, Breathing Orifice; *s t*, Superior Tentacles; *i t*, Inferior Tentacles; *m*, Mouth.

smaller ones, often appearing as mere tubercles, are called the *inferior tentacles*. (See Fig. 18, *s t* and *i t*.) As the snail crawls, the superior tentacles are seen in constant motion, as if the creature were feeling its way about with them.

The eyes, instead of being at the base of the tentacles, as in the fresh-water snails, are found at the tip of the superior tentacles.

In the land snails, with few exceptions, the tentacles can be drawn within the head, and for this reason they are also called *retractile* tentacles.

While the snail is crawling, if the pupil touch the end of the tentacle with his finger, or, even if he alarm the snail by a sudden jar, he will see the tentacles quickly withdraw within the head. The pupil will observe that the bulbous

end containing the eye disappears first, as the end of a glove-finger disappears, when the hand is withdrawn from the glove, the glove turning wrong-side out.

FIG. 19.—SHOWING SNAIL WITH TENTACLES RETRACTED, *A*; AND TENTACLES PROTRUDED, *B*.

16. Something may now be learned as to the way in which land snails eat.

By placing before the snail the tender leaves of lettuce or cabbage, the head will be seen to move, as little mouthfuls of the leaf are bitten off. The upper lip of the mouth is furnished with a hardened piece called the *buccal plate*. It is crescent-shaped, and, in some species, the cutting edge is notched, so that it acts like an upper set of teeth, by which it bites off little bits of the leaf. The floor of the mouth is lined with a membrane having upon it rows of little points which enable the snail to rasp and grind its food. These parts

FIG. 20.—JAW, OR BUCCAL PLATE OF A LAND SNAIL, HIGHLY MAGNIFIED.—(It can just be discerned without a microscope.)

are so minute that they can be studied only by the aid of a microscope. If the pupil will watch his fresh-water snail

as it crawls around on the sides of the jar, he will see at intervals the mouth open, and a glistening tongue appear, as the snail laps up the scum which forms upon the surface of the glass.

NOTE FOR TEACHERS AND PUPILS.—Let the teacher here explain to the pupils what is meant by an object's being magnified.

If the teacher has a common magnifying-glass, let each pupil in turn observe its magnifying effect, by looking at a common house-fly, or the printed page of a book. If a microscope can be shown to the class, it will be better still.

Let it be explained, also, what is meant by an object being enlarged two, or three, or more times. To be enlarged two or three times, is to make the object two or three times as long as it was before, and of proportional size.

Oftentimes the object has to be reduced in size in the figure, as in pictures of large animals, in the picture of an elephant, for example.

In representations of very small animals, however, the figure has to be enlarged in order to show parts plainly that could not otherwise be seen.

17. In searching for snails, the pupil will come across snail-like animals, which have no coiled shells on their backs. Let the pupil examine the under side of damp boards or plank walks in gardens, and he will be sure to find them.

They are very common in old gardens in cities. These

FIG. 21.—A SLUG CONTRACTED.

creatures will be found clinging to the board or upon the ground, and will present this appearance (Fig. 21). Soon, however, they will stretch out their tentacles, and commence

crawling, and then their resemblance to the shell-bearing snails will be seen at once.

Instead of having a coiled shell into which they can retreat when alarmed, they have a little limy scale imbedded in a portion of the back, called the *mantle*. The breathing-orifice is on the right side of the body, and the tentacles, mouth-parts, creeping disk, and other features, are quite similar to the land snails already studied.

FIG. 22.--A NAKED LAND SNAIL, OR SLUG, FULLY EXPANDED.—*a*, Mantle; *b*, Breathing-Orifice.

18. On the approach of winter, land snails bury themselves in the ground, and those that have shells retire within the shell as far as possible, and close the aperture of the shell with a film of the mucus which the body secretes so abundantly. In this condition they remain dormant until the warm weather of spring revives them again.

If the pupil will collect some snails in the early spring, and keep them confined in a box, with earth, damp leaves, or bits of rotten wood or bark, the snails in the course of a few weeks will lay a number of little eggs. These eggs will be white and round, about the size of a pin's-head. By careful tending, that is, by keeping the leaves slightly moist, the

eggs will hatch out tiny snails, and these will attain half their mature size the first season.

FIG. 23.—*a*, Eggs of Land Snail; *b*, Young Snail just hatched; *c*, Young Snail somewhat advanced: *b* and *c* are magnified.

19. If the pupil will also collect a lot of fresh-water air-breathing snails, and keep them alive, they will deposit their eggs upon the sides of the glass jar in which they are confined. These eggs will be oval in shape, and transparent, and will be inclosed in a transparent, jelly-like substance. Fig. 14, *II*, represents the appearance of a cluster of these eggs.

Fig. 24 shows a cluster of eggs with the appearance of two eggs highly magnified, showing the young snails as they appear within the egg.

FIG. 24.—*a*, Cluster of Eggs of a Fresh-Water Snail; *b*, *c*, Eggs enlarged, showing the young Snails within the Eggs.

With the aid of a magnifying-glass, the eggs may be watched from day to day, and the young snail can be seen in its various stages of growth.

20. If a land-snail is taken out of its shell (and this can be done if boiling water be first poured upon it, and then with a pin the animal can be readily picked out), it will present this appearance (Fig. 25):

FIG. 25.—LAND SNAIL REMOVED FROM ITS SHELL.—*m*, Mantle.

The portion contained within the shell presents the same general appearance as the shell itself. A free border, or collar, is seen which corresponds to the aperture of the shell. This border is called the *mantle*, and is a characteristic feature of all the snails thus far studied. It is the edge of the mantle which deposits the successive layers of the shell, and increases its size. In the slug, there is only the limy scale; this is buried in the mantle, which is plainly seen covering a portion of the back, like a shield (Fig. 22, *a*).

CHAPTER IV.

SEA SNAILS.

21. CLASSES that live near the sea-coast may now study the marine, or salt-water snails. These may be collected alive at low tide, upon rocks, or under the sea-weed. By

examining pools left at low tide, many little sea snails may be seen creeping about.

A good place to collect dead shells may be found along an exposed beach. After a violent storm, when the waves have been running high, a great many animals are thrown up from the sea, and among them many shells may be secured.

The following will be some of the shells collected :

FIG. 26.—COMMON SEA SNAILS.

22. With very few exceptions, all sea snails are water-breathers; that is, they are furnished with gills, instead of a simple lung. Most of them are operculated.

In the shells collected the pupil will find two well-marked groups.

In the two lower right-hand figures of Fig. 26, the shells have an aperture with a continuous border; that is,

there is no notch, or fold, in it. In the three upper figures there is a notch, or fold, in the base of the aperture.

FIG. 27.—SHELL SHOWING LONG CANAL.

This notch is called the *canal*, and in some species it is very short, while in other species it is very long, as in Fig. 27.

The mantle of the animal is prolonged in a fold which occupies this canal, and is called the *siphon*. Through this fold or *siphon* the water finds access to the gills. (Fig. 28 shows another species. The siphon is seen as a fold of the mantle running into the canal of the shell.)

23. A very common species, found in the greatest abundance from Maine to Florida, on mud-flats, will give a good

illustration of the uses of the siphon. In this particular species, the siphon is much longer than the canal, and, when the snail is crawling, the siphon is bent upward. As the

FIG. 28.—A SEA SNAIL SEEN FROM BELOW.—*e*, Eye; *m*, Mouth; *s*, Siphon.

habit of this species is to crawl along partly buried in the mud, the siphon, projecting above the level of the mud, conducts the pure sea-water to the gills of the snail below. Fig. 29 illustrates the appearance of this snail :

FIG. 29.—*A*, the Snail crawling upon the Surface of the Mud; *B*, the same slightly buried; *C*, the same nearly buried; the Siphon, *s*, is seen curved upward.

Such shells are called canaliculated shells.

The aperture of the shell is said to be *entire*, when it does not possess this notch, or canal. Let the pupil separate the shells having the aperture entire, from those shells having a canal.

These differences in the shell, as slight as they appear, are accompanied by corresponding differences in the character and habits of the animal.

Those snails having the aperture of the shell entire are with few exceptions vegetable-feeders, while those having a canal to the shell are flesh-feeders. The mouth-parts, and opercula, too, are different in the two groups.

24. Other shells will be found differing greatly in appearance from those thus far studied. One of these is represented in the lower left-hand corner of Fig. 26.

Another species, called the limpet, looks like this (Fig. 30) :

FIG. 30.—SHELL OF LIMPET.

These shells will be found sticking with great tenacity to the rocks, and some skill and force will be required to remove them. This can be done by using the large blade of a pocket-knife, and suddenly scraping them from the rock. If they are then placed in a saucer of water, with the shell downward, the animal within will be found to have the broad, creeping disk, head, tentacles, and other parts, peculiar to the snails already studied.

25. In the land snails, it was learned that the eggs were deposited separately, while in the air-breathing, fresh-water snails the eggs were inclosed in a gelatinous substance. Among the sea snails there are many species which inclose their eggs in pods, or capsules, as is also the case in the operculated fresh-water snails. Sometimes the capsules are clustered together in large masses, as in those of the whelk (Fig. 31):

FIG. 31.— A VERY SMALL CLUSTER OF EGGS OF THE WHELK DEPOSITED ON THE STEM OF A LARGE SEA-WEED.

In others they are united in a long string (Fig. 32).

FIG. 32.—*A*, a Portion of a String of Egg-Capsules, from a Florida Species of Whelk similar to Fig. 27; *B*, a single Capsule separated, showing Outlet, *o*, through which the Young escape; *s*, Stem.

The common cockle sticks its separate egg-capsules side by side upon the rock (Fig. 33).

The beach-cockle deposits its eggs in a broad ribbon of sand cemented together, looking very much like a deep saucer, with the bottom broken out, and the side separated (Fig. 34).

FIG. 33.—EGG-CAPSULES OF COMMON COCKLE.—(A shell of the animal which produces these capsules is shown in the upper central figure of Fig. 26.)

This ribbon is elastic when wet, and, if it is held up to the light, the little transparent spaces for the eggs will be plainly seen.

FIG. 34.—EGG-RIBBON OF BEACH-COCKLE (the shell of which is shown in the lower right-hand figure of Fig. 26).

CHAPTER V.

FRESH-WATER MUSSELS.

26. LOOKING over our fresh-water shells again, we find many that are known as mussels, or clams. These shells are common everywhere along the margins of brooks, rivers, and lakes. The musk-rats feed upon the soft parts of the mus-

sels, and the remains of their feasts may be found in piles of mussel-shells, all along the shores of certain lakes.

The shell is composed of two pieces, or *valves*, as they are called. The two valves are often found united, and the margin along which they are connected is called the *hinge-margin*, because the shells hinge at this part, and will open and shut as a door swings upon its hinges.

Let the pupil now examine a perfect fresh-water mussel, that is, a mussel in which the valves are united in this way, and he will observe that they are connected by a brownish substance, which is quite elastic when the shell is alive, but becomes brittle when dried. The shells are held together as the covers of a book are held together by the back.

This substance is called the *ligament*, and the position of this ligament will indicate the back, or *dorsal region* of the animal.

27. On the outside of the shell will be seen fine lines, which run nearly parallel to the outside margin of the shell. These lines are the *lines of growth*, and indicate the successive stages of growth, or increase of the shell, as in the lines of growth in the snail-shell already studied, and, as in the snails, the growth takes place at the margin of the shells.

The pupil may trace these concentric lines back, as they grow smaller and smaller, till they are found to start from one point at the back of the shell, and this point is called the *beak* or *umbone*. It represents the starting-point in the growth of the shell. In fresh-water mussels, the umbones are eaten away by some corrosive action of the water, and the

early stages in the growth of the shell are usually destroyed. In very young shells, however, the early stages can be plainly seen.

FIG. 35.—A FRESH-WATER MUSSEL.—*l*, Ligament; *u*, Umbone; *f*, Foot; *ex.*, Excurrent Orifice; *in.*, Incurrent Orifice.

28. The ligament is always behind the beak, or umbone, in fresh-water mussels, and in nearly all *bivalve* shells (so called, because they have two valves or pieces, while the snail-shells are sometimes called *univalve* shells, because they have but one valve or piece).

Let the pupil now hold a perfect mussel-shell in his hand (that is, a mussel in which both valves are together, and united across the back), with the ligament uppermost, and the umbone away from him, or beyond the ligament, and the valve on his left hand is the one which covers the left side of the animal, while the valve on his right hand covers the

right side of the animal. The forward end will be the end away from him, and the hinder end will, of course, be the end toward him. (*See Fig. 35.*)

29. Let the pupils now endeavor to collect some fresh-water mussels alive. These may be found partly buried in the sand or mud of rivers and lakes. As they crawl along partly buried in this way, they plough up the sand, leaving a well-marked furrow or groove behind them. Every boy that goes in bathing is familiar with the peculiar furrow left by the fresh-water mussel. By following such a furrow, the mussel that made it will soon be found.

Fig. 36 represents the appearance of a common fresh-water mussel in the act of crawling.

FIG. 36.—SHOWING POSITION OF MUSSEL WHEN CRAWLING.—*f*, foot buried below the surface of the sand *s*. Above the line *s* is supposed to be water, the line representing the bottom of a lake or river.

Having collected a few in this way, they may be placed in a large, shallow pan of water, and allowed to remain quiet for a while. Gradually the shells will open a little, and from the hinder end a curious fringed border appears; on examin-

ing this border, it will soon be found that the border forms two openings which lead into the shell.

Great care must be taken not to jar the dish, or the table upon which it rests. The slightest jar will cause the shells to instantly close. If some indigo, or small particles of dirt, be dropped near these openings, currents of water will be revealed; one current pouring out of the opening nearest the back, and another current as steadily pouring in at the other opening. The opening into which the current of water is passing is called the *incurrent orifice*, while the orifice from which a current of water is passing is called the *excurrent orifice*. The incurrent orifice is sometimes called the respiratory orifice, because the water is taken in to supply the gills which are the breathing or respiratory organs of the mussel, and this orifice corresponds to the siphon in the sea snails already studied. This current of water, besides bathing the gills, also carries in minute particles which are floating in the water, and these particles are conducted to the mouth of the creature, and swallowed as food. At the opposite end of the shell from these openings, or the forward end, a whitish, fleshy mass will be seen protruding. This is called the *foot*, and corresponds to the foot or creeping disk in the snails. By means of this foot the mussel crawls through the sand.

The mouth is above the foot, and always concealed within the shell. In Fig. 35 the foot is shown, and also the excurrent and incurrent orifices, with arrows drawn to indicate the direction of the currents of water.

In some small species of fresh-water bivalves, the excur-

rent and incurrent orifices are prolonged into tubes, and then they are called *siphons*. Fig. 37 represents a common species which the pupils may find in muddy brooks and ditches. By using the long-handled dipper already described, some specimens will probably be found. They are quite small, from the size of a pea to that of a nickel cent. The siphonal tubes are prominent, and the foot is long and tongue-shaped, and the animal is very active in crawling about; also in Fig. 14 *K* and *L* represent two of these small animals with bivalve shells.

FIG. 37.

30. The foot of these creatures resembles in appearance and action the foot of a fresh-water snail, only there is no mouth nor tentacles in sight. These parts are present, but are never protruded beyond the edges of the shell.

When the fresh-water mussels are partly open, a fleshy border will be seen just within the edges of the shell, and this is the border of the *mantle*, and corresponds to the same parts described in the snails; the fringed membrane which formed the openings at the hinder part of the mussel is simply a continuation of the mantle.

When the shells are removed from the animal, the mantle will be found lining the shells, just as the blank pages line the inside of a book-cover. While the edge of the mantle deposits the successive layers, which increase the size of

the shell, the entire surface of the mantle deposits the pearly substance which lines the inner surface of the shells, and which is so characteristic of the fresh-water mussels.

31. Grains of sand, or other particles, getting in between the mantle and the shell, are soon covered by layers of pearly substance poured out, or secreted by the mantle. In this way pearls are formed.

If pearls are broken open, a centre, or nucleus, will be found, consisting of some particle of dirt or sand, or some substance which had found its way by accident between the mantle and the shell, and around which the pearly matter has been formed in successive layers.

FIG. 38.—A, Pearly Concretions from a Fresh-water Mussel; B, Pearly Concretions from the Common Oyster.

In shells having a brilliant, pearly lining, or *nacre*, the pearls obtained are oftentimes very beautiful, and from certain Oriental species living in the sea, called *Avicula*, the most brilliant pearls of commerce are obtained. If, on the other hand, the nacre lining the shell is dull white, as in the common oyster, the pearls are dull-colored. This kind of pearls is often found in oysters.

The Chinese have long been familiar with the art of

making artificial pearls. By partly opening the shells of certain fresh-water mussels, and inserting little lead images, or other objects, between the mantle and the shell, the objects soon become covered with a natural layer of pearl.

32. Let the pupils now study the markings on the inner surface of the shells of river-mussels. The shells of these creatures are called *valves*, and are spoken of as right or left valves, according to whether they are on the right or left side of the animal.

Certain ridges and prominences will be seen at the hinge, and, when the valves are carefully joined, the ridges in one valve will correspond to grooves in the other valve. These ridges are called *teeth*. The short ones, near the beak, are called *cardinal teeth*, and the long ones *lateral teeth*. The margin upon which they occur is called the *hinge-margin*, for it is upon this margin that the valves turn. (See Fig. 39.)

33. Certain scars, or impressions, will be found marking the inside of the valves, and these indicate the point of the attachment of certain muscles to move the valves, and to enable the animal to protrude its foot, and crawl along.

These marks are hence called muscular marks, or *muscular impressions*, and will be found to correspond in the right and left valves.

An irregular, round impression will be found at each end of the valve, near the hinge-margin. These show where the muscles are attached to move and close the valves, and hold them firmly together. The muscles run directly across from one valve to the other; and, to open a live

mussel, it is necessary to pass a sharp blade between the valves, and cut through the muscles, before the valves will open. These muscles are called the *adductor muscles*, and the scars or impressions on the valves are called the *adductor muscular impressions*. Very close to the adductor muscular impressions are seen smaller impressions, and these indicate where the muscles are attached which move the foot. These muscles are called the *pedal muscles*, and the impressions are called the *pedal muscular impressions*. One occurs just behind the anterior adductor impression; the other will be found just above, and in front of the posterior adductor impression.

FIG. 39.—THE RIGHT VALVE OF A FRESH-WATER MUSSEL.—*c*, Cardinal Teeth; *l*, Lateral Tooth; *li*, Ligament; *aa*, Anterior Adductor Impression; *pa*, Posterior Adductor Impression; *ap*, Anterior Pedal Muscular Impression; *pp*, Posterior Pedal Muscular Impression; *p*, Pallial Line.

34. Besides these marks, the pupil will see a delicate and slightly irregular line running from the anterior to the

posterior muscular impression, just inside, and nearly parallel with the lower margin of the shell. This line is called the *pallial line*, and indicates where the mantle is attached to the shell. It will be observed that, when the soft parts are removed from the shell, the mantle adheres along this line.

The pupil may mark with a pen the names of all the parts upon the inside of a fresh-water mussel.

35. When the mussel is opened by separating the adductor muscles with a knife, the valves slowly open, and after the animal is removed the valves still remain partly open, and, to preserve them closed, a string has to be tied around them, and in this condition, if the ligament is allowed to dry, the valves will then remain closed. From this it is evident that the ligament acts upon the valves to draw them apart. To keep them closed, then, the animal must continually exert itself by contracting the adductor muscles; and it will be found that, when these creatures are left in the water, undisturbed for a while, the muscles relax, and the valves partly open. The ligament is elastic, and is stretched as it were from one valve to the other, over the back. A possible imitation of the action might be represented by partly opening the lids of a book, and then gluing across the back, from one lid to the other, a sheet of elastic rubber. If, now, the lids are tightly closed, the rubber is drawn out, or stretched across the back, and, if allowed to regain its elasticity, the lids are pulled apart. This experiment illustrates the way in which the ligament acts in those shells which have the ligament external.

CHAPTER VI.

CLAMS, MUSSELS, AND OYSTERS.

36. CLASSES having access to salt-water may now collect some bivalves, as the clam, mussel, razor-shell, oyster, scallop, and whatever species they can find belonging to this group. A much greater variety of forms will be found in salt-water than in fresh-water.

Among some of the common species met with will be the following :

FIG. 40.—SALT-WATER BIVALVES.

In these the pupil may trace out the muscular impressions within the shell, and make out their relations to the impressions already described in the fresh-water mussels.

Many differences will be observed in the muscular impressions, as well as in the teeth and the position of the ligament.

37. As the common soft-shelled clam can be readily procured in the fish-markets, it will be well to study this first. A live specimen must be selected, and, as the clam lives a long time after it has been removed from the water, there will be no difficulty in getting the proper specimen. Upon pressing the valves together, or touching the soft parts which partly protrude from between the valves, the creature will show signs of life, by drawing the shells closer together, and this will assure the pupil that the specimen is alive.

A large shallow dish may now be filled with pure seawater, and in this the clam may be placed. After it has remained there for some time, the black end of the animal, which is incorrectly called the head, will slowly stretch out from between the shells, and the end, unfolding, will display two openings fringed with little feelers (*see* Fig. 42). Into one of these openings the water will be seen flowing, while from the other a current of water will be seen issuing. And these openings are called the *incurrent* and *excurrent orifices*, and correspond to similar parts previously described in the fresh-water mussels. In the latter creature, the openings just protruded beyond the edge of the shell. In some very small species of fresh-water bivalves, one of which was shown in Fig. 37, these openings were at the end of separate tubes. In the clam the tubes are inclosed in one sheath.

The clam can protrude this apparatus to a length equaling that of the shell two or three times. As the clam lives

buried at some depth below the level of the sand or mud in which it occurs, it requires this extension of the openings to reach the sea-water above.

38. It may be stated here, that the current of water passing into the general cavity of the shell not only carries the particle of food upon which the animal subsists, but conveys the pure sea-water to the gills by which it breathes, the gills performing the same function for animals living immersed in water as the lungs perform for creatures which breathe air. All bivalves depend upon currents of water to convey their food to them.

While, in the snails, the creatures could go in quest of food, having the power of protruding the head from the shell, and mouths furnished with means to bite or rasp their food, in the bivalves there is really no head, they having only a little opening directly under the anterior adductor muscle, which is the mouth, and into which the particles of food are swept.

39. If, now, the clam is opened, the edges of the mantle will be found much thickened and united, except a small slit near the front edge, through which can be protruded a small, tongue-shaped foot. Powerful muscles will be found at the base of the united siphons or tubes, which move the siphons in and out, and an examination of the inside of the shell will show where these muscles are attached. The pallial line, instead of running directly from the anterior adductor impression to the posterior one, is abruptly curved back, and forms a sharp bend, as it turns again to the posterior ad-

ductor impression. This mark is called the *sinus*, and, when present, indicates the siphons to be of considerable size, and having large muscles to contract them.

Let these parts now be marked with a pen upon the shell, with their names, as in Fig. 41 :

FIG. 41.—RIGHT VALVE OF A COMMON CLAM.—*z*, Ligament; *aa*, Anterior Adductor Impression; *pa*, Posterior Adductor Impression.

40. On opening the shell, a prominent tooth is seen on one of the valves near the hinge, while on the other valve there is a corresponding depression. When the valves are forcibly separated, there is left attached either to the tooth, or in the depression, a substance resembling dark glue, very elastic, and firmly attached to its place. This is the ligament, and is said to be *internal*, because it is within the shell, and not upon the outside, as in the fresh-water mussel already studied. When the animal closes the shell by contracting the adductors, the ligament is compressed, by being jammed between the prominent tooth, and the recess into which it fits. When the muscles relax, the ligament expands,

FIG. 42.—COMMON CLAM.—*A*, showing Siphons partly extended; *in.*, Incurrent Orifice; *ex.*, Excurrent Orifice; *f.* Foot; *m.* thickened border of the Mantle projecting beyond the Edge of the Shell; *B*, Siphons, greatly extended. (The shell is not drawn, as there was no room for it on the page.) The excurrent and incurrent orifices are more open than in *A*.

forcing the valves apart. The way in which it works might be illustrated by placing a piece of rubber inside the hinge of a door: when the door is closed, the rubber is squeezed, and the tendency would, of course, be for the rubber, in expanding, to again push the door open.

That it requires a continual effort for the clam to keep the valves closed, is seen in the fact that when these creatures are allowed to remain out of water for a while, as they are when in the market, the muscles get tired, and, relaxing, the shells partly open. If, now, the basket or barrel which they are in be suddenly shaken, the clams will as suddenly close, and a rustling sound is made, as the water is forced out from the gill-cavity, the water often squirting out in a stream from the siphonal openings.

41. On the rocks between high and low water mark, and adhering to the piles of the wharves, may be found clusters of mussels which are attached to these places, and to each other, by little brown threads which issue from between the valves below. These threads are made at will by the creature, one by one, and are fastened to the substances upon which they rest. The threads are called *byssal-threads*, and, combined, form the *byssus*.

If the pupils will collect a number of these salt-water mussels, and place them in a large glass dish or bowl filled with salt-water, they may watch the mussels as the creatures attach themselves to the sides of the vessel. In the figure, the byssus is seen like threads coming from between the valves below, with their ends adhering to the stone.

42. Fig. 44 represents an animal which is often thrown up on beaches along the coasts, after a storm, and whose shells are very common in the *débris* thrown up by the waves.

FIG. 43.—A MUSSEL ATTACHED TO A STONE BY ITS BYSSUS.—*f*, Foot.

These shells are very thin and delicate, and the valves are strengthened by a thickened rib which runs from the umbone toward the lower margin of the valve within.

In this species the siphons are united, the mantle projects beyond the edge of the valves, and the foot is flattened in front.

FIG. 44.—*l*, Ligament; *f*, Foot; *m*, Mantle; *s*, Siphons; *ex.*, Excurrent Orifice; *in.*, Incurrent Orifice.

Fig. 45 represents another animal which is often abundant on the sea-beach. In this species the siphonal tubes are

separate, instead of being united. This figure represents the tubes only partly extended. They can be thrown out to twice the length represented in the figure.

FIG. 45.

Another species quite similar to the above occurs on mud flats, in company with the common clam. If this be collected alive and placed in sea-water, the creature will extend its siphons, which are long and separate, and bend them in coils.

43. The pupils have now learned, among other things, a few features regarding the position which certain bivalves occupy in their native haunts: the fresh-water mussel creeping by means of its foot through the mud or sand in which it lives partly buried; the salt-water mussel, fastened to some place by means of its byssus; the soft-shelled clam, lying buried at some depth in the mud, and extending its siphons to conduct the pure sea-water to its gills, and food to its mouth.

Oysters differ considerably from the animals to which they are related, and which have just been studied. Instead of being free, they grow attached by one of their valves to the rock, or to one another; clusters of a dozen or more individuals of different sizes are found growing, attached to each other, and forming large masses. At any oyster-market

the pupil may get these clusters. Before studying a specimen, it is best to clean the shell thoroughly in water by means of a coarse brush.

44. Instead of having two adductor muscles, it has but one (though this muscle, it seems, is composed of two elements). A single dark-purple mark on the inside of each valve shows the point of attachment of the adductor muscle. When the oyster is opened, the mantle contracts somewhat, so that the edge of the mantle is some way from the margin of the shell, as shown in Fig. 47.

The left valve is the larger, and is the one that becomes attached; the right valve is flattened, and somewhat smaller. The mantle has its margins free; that is, the edges are not united as in the common clam, where they are not only united, but greatly thickened. Neither is the mantle prolonged into siphons as in other species; consequently, the water flows in at one portion of the shell, and pours out of another portion, not being definitely conducted by special channels, as in those forms heretofore given. The oyster can be readily studied, as specimens may be got in almost every village in the country.

In looking over canned specimens, be sure and pick out a large one, and one that does not appear to be mutilated, as they frequently are when taken out of the shell by the oysterman, or jammed, as they often are in packing.

To those who can get them alive, it is well to have the oysterman open the specimen, being sure that he removes the larger valve, leaving the oyster attached, and resting in

the smaller and flat valve, which is the *right* one. To examine it properly, the specimen must be placed in a deep saucer filled with water, so as to cover it. A number of rinsings will remove the mucus with which the oyster is covered, and this will render the specimen in better condition to examine. In placing it under water in this way, the membranes float apart, and can be more readily studied.

45. The adductor muscle is near the middle of the animal. It is composed of two elements, one half being a glistening white, and the other half being grayish. Immediately adjoining the grayish portion of the muscle, a translucent space is seen, and this space contains the heart, composed of a body constricted in the centre, as if a tube had been tied in the middle by a string. This is the heart proper, and in specimens freshly opened the heart may be seen to slowly pulsate, or beat.

By raising the mantle, the gills will be seen as delicate, leaf-like membranes.

At the smaller end of the oyster, and that portion which comes next to the beak or hinge, the mouth will be found having on each side two delicate lappets, which are called the *palpi*. It will be difficult to find the mouth, and some patience will be demanded in lifting the mantle and following up between the palpi to where the mouth is.

The dark region just back of the mouth contains the stomach and liver; the dark or blackish portion, showing so conspicuously in cooked specimens, being the liver.

By referring to the accompanying figures, these parts may be readily made out :

FIG. 46.—OYSTER ATTACHED BY LEFT VALVE TO A STONE.

FIG. 47.—OYSTER WITH THE LEFT VALVE REMOVED.—*h*, Heart; *l*, Ligament; *m*, Position of Mouth.

46. The pupil will now recall some characters in common between the snails, and the mussels, clams, and oysters, thus far studied in these lessons, namely : they all have the body protected by a limy shell (except the slug), this shell either composed of one piece, as in the snails, or of two pieces or

FIG. 48.—RIGHT VALVE OF AN OYSTER, the Dark Semicircular Mark near the Middle of the Shell being the Adductor Muscular Impression, the Pallial Line showing faintly.

valves, as in the clams, mussels, and oysters. All of them increase the size of their shells, or grow, by the addition of layers of shell-material to the edge of the aperture, or the margins of their valves, and these layers are indicated by delicate lines on the outside of the shell, and called lines of growth. They all, excepting the oyster and a few other forms, have the creeping disk or foot. In the snails, this is broad and flat; in the mussel and clam the foot is flattened sideways, and variously shaped. In the snails, the creature projects, with the foot, a head furnished with feelers, or tentacles, and the mouth is possessed of certain hard parts by which food can be eaten. In the mussels and clams there is no definite head, the mouth being hidden away within the mantle, and the creature projecting, from the forward end, only the foot. In all of these animals thus far studied there is a cavity within, containing the gills to which water has access, or else there is a simple

lung, as in the air-breathing snails. These, with the cuttle-fishes, which we will not consider here, belong to a branch of the Animal Kingdom called Mollusca.

CHAPTER VII.

COLLECTING INSECTS.

47. THESE lessons, as well as the preceding ones, are prepared with the understanding that the pupils shall, so far as possible, make a collection of the species of animals studied. In fact, it is a part of the lesson to know how and where to collect, and above all to know how to preserve the specimens collected. To enable the pupils to do this, the briefest directions are given for the making of boxes, nets, etc., accompanied with the simplest methods of preserving the collections made.

In many cases the directions given are by no means the professional methods; thus the pupils are directed to use common pins for insects, while the professional collector uses only the true insect-pins made expressly for the purpose, but these are oftentimes difficult to procure, and are more expensive than the common ones.

In commencing these lessons, each pupil must first be provided with a number of common pins, and a box properly arranged in which to pin the insects collected.

Some holiday afternoon, or an hour before school-time

in the morning, may be spent in making the insect-boxes. These may be of any convenient size, having a depth of not over two and a half or three inches, and furnished with a lid. A shallow cigar-box will answer the purpose. The bottom of the box may be lined with strips of corn-pith, or slices of cork, into which the pins can be easily stuck. Large cork-stoppers will do, and these may be cut into lozenge-shaped pieces like this:

FIG. 49.—SLICED CORK FOR INSECT-BOX.

These pieces are to be fastened to the bottom of the box by gluing. If strips of corn-pith are used, they may be tacked or glued to the bottom of the box. The box, when finished, will look something like this:

FIG. 50.—INSECT-BOX.

48. The insects, when collected, are to be pinned to the

cork in the way figured, leaving the head of the pin sufficiently above the insect to grasp with the fingers.

Care must be taken not to have the insect too far down on the pin, as its legs in that case would touch the bottom of the box, and break off. Insects may be killed by immersing them in alcohol for a few minutes.

FIG. 51.—INSECT PINNED

Butterflies may be killed by compressing the body between the thumb and forefinger, as shown in the figure, using just force enough to kill, without crushing them. The fumes of benzine, or ether, and of certain poisons, will also kill insects, but these substances should not be suggested to young pupils, as their use is dangerous. (Teachers will here use their judgment according to the character of their classes.) The rude box and common pins are offered simply for experimental collections. The ingenuity of a pupil, where neater collections are desired, will readily secure better ways of making them.

49. The pupils may go out in a class in quest of material

for study, and this is the best way, as the activity and success of one will act as a stimulus to the others.

FIG. 52.

In the country, the best places to collect are by the roadsides, or borders of woods and groves; in the gardens, and by the fences, or along the shores of lakes and brooks, under stones and stumps, the bark of fallen trees, or beneath the layers of dead leaves. Insects are scarce in deep woods, and in large, open tracts of pasture-land.

In the cities, the parks and gardens will afford good collecting-grounds, as under plank-walks and boards many insects find shelter. Alongside of railroad-tracks the discarded sleepers often hide many a curious beetle. In short,

let the pupil peer under any object large enough to afford shelter to these creatures. By following the furrows made by a plow, certain kinds will surely be met with. The pupil must be urged to pick up every thing that he thinks is an insect, such as grasshoppers, beetles, flies, ants, spiders, etc.

In a single holiday afternoon the pupil will have gathered some of the following animals:

FIG. 53.—SOME OF THE ANIMALS COLLECTED.

CHAPTER VIII.

PARTS OF AN INSECT.

50. THE animals are now to be carefully examined. Let the pupils pick out, and arrange together in one portion of the box, all of those which have three pairs of legs. In

some, the legs will be closely drawn to the body, but by sharp looking they will be found.

After studying these carefully, the pupil will observe that those insects which have three pairs of legs have the body divided into three regions, or parts, called respectively the *head*, *thorax*, and *abdomen*, and that, with few exceptions, they all have wings. Insects having these characters are called *Insects proper*. They are also called *Hexapods*, a word meaning six legs.

These are to be studied first. The other animals collected may be saved for future study.

51. Some insects have the three parts of the body distinctly separated, as in the ants, flies, and wasps. In other insects the parts of the body are very close together, so that it is difficult to distinguish the dividing line, as in certain beetles. Let the pupils examine each insect, and make out the head, thorax, and abdomen.

FIG. 54.

In the head, we find the *mouth*, the *eyes*, and the *feelers*, or *antennæ*.

The mouth is on the under side of the head, and is surrounded by certain parts called *mouth-parts*. These parts differ greatly in different insects.

52. In those insects that chew their food, such as the beetles and grasshoppers, certain of the mouth-parts act as teeth, or jaws, and, being joined to the right and left sides of the mouth, move sideways, and not up and down, as in other animals. In other insects some of the mouth-parts are very long and slender, so as to form a long, sharp sting, as in the bugs proper, so that they use them to suck the juices of plants upon which they feed. Or, the parts are again modified in shape to form a long, slender tube, by which the nectar of flowers may be sucked, as in the butterflies.

FIG. 55.—SHOWING MOUTH-PARTS OF A FEW INSECTS.—THE HEADS ARE SEPARATED FROM THE BODIES, AND ARE FACING THE LEFT, AND DRAWN IN PROFILE AS SEEN FROM THEIR LEFT SIDES.

In the common house-fly, the mouth-parts appear as a proboscis, a kind of fleshy appendage which is bent up when not in use. When the fly feeds, the proboscis unbends, and

the food is lapped up by it. Let the pupils carefully watch a fly as it feeds upon a bit of sugar, or as it laps the hand.

In the butterfly and moth the pupil may uncoil the long tongue with a pin. It resembles in appearance a watch-spring.

53. On the front of the head are two horns, or feelers, called *antennæ*.

These are variously jointed, and vary greatly in different insects.

In butterflies, they are generally long and slender, and swollen at the tips, like drum-sticks. Sometimes they are thread-like, and in others the antennæ are barbed on the sides, and look like a feather, as in certain moths. In some beetles they are strongly jointed. In the common house-fly, they hang down in front of the head.

Below are given figures of the left antenna of several different insects showing how different they are in different kinds of insects. The pupils might save the antennæ of different insects and glue them to a card, writing opposite each one the name of the insect, whether fly, beetle, or locust.

54. On the sides of the head are round, smooth places, and these are the eyes. They are entirely different from the eyes of most animals, for, when examined under the microscope, they are seen to be divided into little spaces, looking very much like the surface of honey-comb. Each of these little spaces represents a separate eye. Some insects have hundreds and even thousands of these little spaces, or eyes. For this reason, such kinds of eyes are called *compound eyes*.

Under the microscope three minute black dots may be

seen on top of the head between the compound eyes, and these are called *simple eyes*.

FIG. 56.—ANTENNÆ OF VARIOUS KINDS OF INSECTS.

A, Fly.

B, Bee.

C, Green Grasshopper.

D, Dragon-fly.

E, Moth.

F, Butterfly.

G, Bug.

H, Beetle.

I, Beetle.

In many insects, as in the dragon-flies for example, the compound eyes are very prominent and cover the sides of the head, enabling the insect to look backward as well as forward.

FIG. 57.—SHOWING COMPOUND AND SIMPLE EYES.

A, Head of Fly, natural size.

B, Head of Fly enlarged eight times; c, compound eye; s, simple eyes.

C, Portion of the surface of a compound eye highly magnified.

55. Thus far we have learned that an *insect proper*, or *true insect*, has the body divided into three parts or regions, called the head, the thorax, and the abdomen.

Let the pupil take a dead fly, and first pull off carefully the legs and wings, and afterward separate the head from the thorax, and the thorax from the abdomen. Having separated the parts in this way, they may be pasted to a card in this manner, writing the correct name beside each part, or region, as shown in Fig. 58.

FIG. 58.—CARD, WITH REGIONS OF AN INSECT GLUED TO IT, AND MARKED.

The principal parts of the head are the mouth-parts, compound eyes, simple eyes, and antennæ.

56. In studying the thorax, the pupil may select some common insect (a large fly, or a bee, will answer the purpose), and pull off the head and abdomen. A common house-fly separated in this way may be stuck upon a card. By experimenting with a number of insects in this manner, the pupil will soon learn that insects not only have the body divided into three sections, but that the thorax invariably has attached to it the legs and wings—the legs being at-

tached to the under side of the thorax, while the wings are attached to the upper side thereof.

FIG. 59.—Head, having mouth-parts, antennæ, compound eyes, *c* ; and simple eyes, *s*.

Thorax, having legs and wings.

Abdomen, never having legs or wings, but having certain appendages at the extremity.

57. The wings of insects are never more than four in number, and these are arranged in two pairs.

The group of insects to which the house-fly belongs has but two wings, or a single pair, and in this group (and other groups of insects as well) there are some which have no wings.

The wings are very different in shape and structure in distinct kinds of insects. In the common fly they are quite small, and transparent. In the butterflies they are large and broad, and are covered with minute scales which rub off on the fingers like dust. In the dragon-fly the wings are long and narrow.

In all these wings the pupil will observe a net-work of lines, which stiffen the wing and support the delicate membrane constituting the wing, just as the frame of a kite stiffens and supports the paper that is stretched upon it. These lines are called *veins*, or *nervures*. To study the *vena-*

FIG. 60.—AN IN-
SECT WITHOUT
WINGS.

FIG. 61.—AN INSECT WITH
TWO WINGS.

FIG. 62.—AN INSECT WITH FOUR WINGS.

tion of the wings, is to study the way in which these veins are arranged. It would be well for the pupils to stick upon a card a number of different kinds of wings, such as those of the grasshoppers, beetles, flies, wasps, and label them accordingly.

58. In many insects the forward and hinder pair of wings are of the same nature, as in the butterflies, moths, bees, wasps, and dragon-flies. In other insects, however, the forward-wings differ in character from the hind-wings. Thus, in the grasshopper the forward pair of wings are more dense in structure than the hind-wings, though the little veins may be seen closely crowded together. They differ as well in form. (*See Fig. 64.*)

In other insects, as in the squash-bug, the front-wings

have the half nearer the body dense and stiff, while the remaining portion of the wing is very thin, or membranous.

In the beetles the front wings are hard throughout, and in most of them are bent and moulded to the shape of the body, and, when closed, form a tight cover over the hind-wings.

The forward-wings of a beetle are so unlike ordinary wings, that they are not called wings, but are known as *elytra*, a single one being called an *elytron*.

59. When insects are at rest, they generally bring their wings into a position different from that taken by them in flight. In certain dragon-flies, however, the wings when

Butterfly at rest with wings
meeting over the back.

Moth at rest with the wings
sloping on the sides of the
body.

FIG. 63.—INSECTS AT REST.

at rest assume the same position as they do when flying. In the butterfly the wings are brought together over the back when at rest, while the moths with few exceptions rest them sloping over the abdomen, the front-wings covering the hinder-wings.

In the grasshopper, the front-wings are long and narrow, while the hind-wings are large and broad. When the grasshopper is at rest, the hind-wings are folded together precisely like a fan, and, when closed, rest against the sides of the abdomen, the long, narrow front-wings closing down upon them, and covering them.

60. Let the pupils prepare a grasshopper, with the wings spread as in the act of flying. A specimen which is dry may be moistened by wrapping it up in a piece of wet cloth, and letting it remain a day or two.

FIG. 64.—GRASSHOPPER WITH THE WINGS OF ONE SIDE EXPANDED.—*f*, Forward-wing; *h*, Hinder-wing.

Having softened the joints of the insect in this way, it may then be pinned to a piece of cork, or a pin-cushion, and, the wings having been stretched, they may be pinned in this position, using triangular bits of card through which the pins are passed to hold the wings in place, as represented in Fig. 64, which shows a grasshopper with the wings on

one side of the body pinned in the way described. When the insect becomes perfectly dry the wings will remain in the position in which they were pinned.

A common beetle should be prepared in the same way.

In the beetle the front-wings are very hard and are closed tightly over the hind-wings. With a pin, or the blade of a knife, the upper or front wings may be opened, and beneath these will be seen the hind-wings, not folded like a fan as in the grasshopper, but folded or bent in the middle, as the arm is bent at the elbow.

FIG. 65.—A BEETLE WITH THE ELYTRON AND HIND-WING OF THE RIGHT SIDE OPEN, AND THE ELYTRON OF THE LEFT SIDE OPEN WITH THE LEFT HIND-WING FOLDED IN ITS NATURAL POSITION WHEN CLOSED.

61. The abdomen has no wings or legs, but is plainly marked with lines running across the abdomen transversely.

FIG. 63.—ABDOMEN OF A DRAGON-FLY, SHOWING RINGS OR SEGMENTS.

These lines show the separation of the abdomen into rings, or segments. In insects with lengthened and slender abdomens the segments are long, and the abdomen, when bent or curved, bends at these joints, as shown in Fig. 67.

FIG. 67.—INSECT WITH A LONG, SLENDER ABDOMEN.

If the pupil can handle these parts delicately enough, he may be able to separate the abdomen at these joints, into a series of rings, or segments, and glue them on a card, marked "*Rings or segments of the abdomen.*" In the grasshoppers the segments show very plainly. On the hinder part of the abdomen there are various appendages, sometimes so short as to be scarcely perceptible, sometimes long, and thread-like, as in the May-fly (Fig. 98); again, in the shape of a sharp sting, as in the hornet. In the cricket, they are quite long and conspicuous. These appendages vary greatly in different insects.

CHAPTER IX.

PARTS OF AN INSECT (CONTINUED).

62. THE pupils have learned that the abdomen is divided into rings or segments, and the division between these segments is plainly seen in most insects.

The thorax is divided in a similar manner, only the lines which divide the thorax are not so plainly seen.

The number of segments in the thorax is three. To the first segment, the head and first pair of legs are attached; to the second segment, the second pair of legs and the first pair of wings are attached; and to the third segment, the hind pair of legs, the hind pair of wings, and the abdomen, are attached. The three segments of the thorax have special names: the *prothorax*, this being the forward segment, next to the head; *mesothorax*, being the middle segment; and *metathorax*, being the last segment. Arranging these segments with the appendages attached to them in a table, they would appear as follows:

THE THORAX IS COM- POSED OF THREE SEG- MENTS.	{	1st Segment, <i>Prothorax</i> ,	{ Has attached to it the first pair of legs.
		2d Segment, <i>Mesothorax</i> ,	{ Has attached to it the second pair of legs and the first pair of wings.
		3d Segment, <i>Metathorax</i> ,	{ Has attached to it the third pair of legs and the second pair of wings.

63. Let the pupils now endeavor to dissect a beetle, carefully separating the segments of the thorax, and, if possible, the minute jaws and other mouth-parts, and stick them on

a large card, with the names of the different parts neatly marked upon the card, as in the accompanying figure (Fig. 68):

FIG. 68.—A COMMON BROWN BEETLE, WITH THE PARTS SEPARATED.

Having now learned something about the parts of an insect, and having seen how much these parts vary in size and appearance in different insects, the pupils are better prepared to understand the surprising modification which the *mouth-parts* undergo in the different groups.

64. The parts of an insect's mouth, generally speaking, consist of an upper lip, called the *labrum*; a pair of jaws, called *mandibles*; a pair of smaller jaws, called *maxillæ*, to which are attached little jointed feelers, called *maxillary*

palpi; and a lower lip, called the *labium*, which represents still another pair of jaws joined together; to this joined piece, or *labium*, are attached a pair of jointed feelers, called *labial palpi*. In Fig. 68 these parts are shown separated from the head.

The numberless varieties of mouth-parts, peculiar to different insects, are in reality made up by modifications of the parts above described. Thus, in one group of insects, the mandibles are lengthened out into a piercing-like sting, while some of the other parts are reduced in size, or become almost obsolete. In another group the maxillæ are greatly elongated, with their edges joined to form a tube, while the other parts of the mouth are scarcely to be discerned. In another group the labium is greatly lengthened to form a tongue-like organ for lapping up food, while the mandibles—so big and hard in some insects—are barely perceptible, and of no use to the insect.

Not only, then, do these parts assume different proportions and different shapes in the different groups of insects, but they also vary greatly in being very hard or very soft.

If the pupils will examine the different kinds of insects' wings, taking the front-pair of wings for example, they will find a marked difference between them, some being very large and transparent, as in the dragon-fly, others being hard and opaque, as in the front wings or elytra of a beetle. Compare the broad and brilliant-colored wing of the butterfly with the straight and narrow fore-wing of a common

grasshopper. And yet these are all wings. In a similar way do the mouth-parts of an insect vary.

65. In the head of a mosquito, what appears to be a single sting, by which the animal pierces the flesh and sucks the blood, is in reality composed of long, delicate, thread-like parts, which represent the mandibles, maxillæ, and the tongue, or *ligula*, which represents a prolongation of the labium. In the bugs the mouth-parts are compacted into a hard beak—the *piercer*, so called, consisting of mandibles, maxillæ, and labium, the labrum being represented by an acutely triangular piece.

The mouth-parts of a beetle have already been described in general terms. They are represented as separated from the head in Fig. 68, while in Fig. 69 a side-view of another beetle is given in which the mandible shows very prominently, while the labrum, labium, and maxillæ, do not show at all, as they are concealed by the other parts. The maxillary and labial palpi of one side are seen, however.

66. In the butterfly the labial palpus is seen very large and prominent, while the coiled, thread-like tongue represents the pair of maxillæ lengthened and joined, forming a long elastic tube which can be coiled or uncoiled by the insect, and through which it sips the nectar of flowers. In the mouth-parts of a house-fly the parts are soft and fleshy, and united together to form a sort of proboscis; the maxillæ are minute; the maxillary palpi are present as simple jointed appendages; the mandibles are minute, and useless; while the labium is greatly developed, having a broad end which is

divided into two lobes at the extremity, by means of which the fly laps up its liquid food. The insides of these lobes are rough, and the irritation which flies produce when they alight upon the hand is caused by the scratching of these rough surfaces.

FIG. 69.—HEADS OF A FEW INSECTS SEEN FROM THE SIDE, SHOWING MOUTH-PARTS, NAMED.

67. Much more may be learned about the mouth-parts of insects, and the pupils might attempt the separation of the mouth-parts of such insects as the grasshopper, beetle, wasp, and butterfly, sticking these parts, when separated, upon a piece of white card, as shown in Fig. 68.

It will be advisable, also, for the pupils to utilize the broken specimens of insects by selecting the wings of different insects and gluing them upon a card, labeling each one. Different kinds of antennæ might be fastened upon another card, and the legs of some widely-different insects may be arranged in the same way. For example, let them take the hind-pair of legs of a grasshopper and of a water-beetle. It will be instructive to observe how different these two kinds are, and how admirably one is adapted for jumping, while the other is so perfectly fitted for a paddle. By making comparative collections of this kind the pupils will learn a great deal regarding the structure of insects.

68. The pupils have thus far learned that a true insect has the body divided into regions called the head, thorax, and abdomen; that the head bears the mouth-parts, antennæ, and eyes. The thorax has the legs and wings, while the abdomen has only the caudal or tail appendages, and these are not often apparent. They have also learned that the thorax is made up of three segments, not often plainly marked, while the abdomen is composed of a greater number of segments, in most cases very plainly apparent. As each segment of the thorax is characterized by having attached to it a pair of appendages, and as the head contains a number of appendages, it is believed by many naturalists that an insect's head is composed of a number of segments, so closely merged together, however, as not to be distinguished, except theoretically. As naturalists, however, differ in their estimate of the number, we will leave this difficult problem for more advanced students to study.

CHAPTER X.

GROWTH OF INSECTS.

69. As the study of the growth of an insect, from the egg to maturity, requires some time and considerable care, the different stages of such growth may be described and taught with what examples the pupil may be able to collect.

An afternoon may be spent exclusively in collecting the following objects :

FIG. 70.—EGGS: *A*, Eggs attached to Outside of a Cocoon (*B* and *C* copied from figures by C. V. Riley).

FIG. 71.—CATERPILLAR AND GRUB-WORMS.

FIG. 72.—CHRYSIDES.

FIG. 73.—COCOONS: *A*, showing Inside of Cocoon, containing the Remains of a Chrysalis-Skin.

In the spring and fall the eggs of the canker-worm moth may be found in abundance on fences and trees in cities. They are very minute, and are found in clusters arranged like stones in a pavement, but with greater regularity. With a sharp knife a shaving of wood may be cut off, taking the eggs with it. (*See* eggs in Fig. 77, *b*.)

If they are collected in the spring-time, little worms will hatch from them in the course of a few weeks, and these may be fed on the young leaves of the elm-tree.

Eggs of other insects may be found on fences, leaves, and twigs of plants; also on the leaves of the squash-vine, and other plants in the garden.

Certain eggs may be found upon the twigs of apple-trees, covered with a shiny coating, like varnish. For all these objects the pupils will have to hunt carefully, as only the keenest eyes will find them out.

70. *Caterpillars* and *grub-worms* are found everywhere, so common indeed that the pupil has only to examine the fences as he goes to school to secure some. For certain kinds of grub-worms, he may dig in the garden, follow the furrow made by a plough, or tear the bark from some dead tree, and discover the specimens he is in search of. For *chrysalides* and *cocoons* the pupils may be directed to search on old garden-fences, under stones and dry boards.

Some chrysalides will be found hanging with the large end downward, as in Fig. 72; others will be found suspended by the small end, with a delicate thread around the middle,

holding the chrysalis horizontally, or vertically, against the fence, as in Fig. 80, *C*.

Some of them are encased in a mesh of threads, which may be built against the fence, or under the edges of clapboards on the sides of houses. And within the cocoons collected, the chrysalides, or their empty cases, will be found, as in Fig. 73.

FIG. 74.—PORTION OF A FENCE, HAVING UPON IT, AMONG OTHER THINGS, EGGS, CATERPILLARS AND CHRYSALIDES OF INSECTS.

Rich collecting-places may always be found on old garden-fences in cities and towns. Fences surrounded by trees and bushes will oftentimes have a great many insects lurking under projecting edges—caterpillars climbing up the posts,

and chrysalides attached to the rails of the fence. Fig. 74 shows a portion of a fence of this kind.

With the eggs, caterpillars, and chrysalides on hand, the pupils are ready to study the life-history of an insect.

71. Many animals, as fishes, snakes, and birds, lay eggs, and from these eggs little creatures are hatched that resemble the animals which laid the eggs.

Insects also lay eggs, but from these eggs come little creatures which do not at all resemble the insect that produced them.

The different kinds of eggs collected by the pupils were laid by insects of different kinds. For example, if they have collected eggs like the following (Fig. 75), the insect that laid them was a moth, and looked like Fig. 76.

FIG. 75.—EGGS LAID ON THE
TWIG OF AN APPLE-TREE.

FIG. 76.—INSECT WHICH LAID THE EGGS IN
FIG. 75.

Now, if the pupil will keep these eggs in a box, there will hatch from them little animals resembling worms, very tiny at first, but growing rapidly if supplied with appropriate

food. Insects generally lay their eggs in such places that the worm, or caterpillar, coming from them can easily find access to its proper food, and this food in the case of most caterpillars consists of leaves, or the wood, bark, or juices, of plants and trees.

72. Commencing with the egg, the pupil should get, if possible, the insect while in the act of depositing her eggs, and this will not be difficult to do in the case of the canker-worm moth, whose caterpillars commit such ravages upon the elm-trees.

FIG. 77.—CANKER-WORM MOTH, EGGS, AND WORMS: *a*, Female Canker-worm Moth laying her Eggs, *b*; *c*, Top-View, and *d*, Side-View of an Egg magnified; *e*, Canker-worm eating its way out of the Egg, magnified; *f*, Magnified View of Canker-worm; *g*, Natural Size of Canker-worm after leaving the Egg; *h*, Male Canker-worm Moth.

The female of the canker-worm moth has no wings. They are very common in early spring and in the fall, laying their eggs on fences. Fig. 77, *a*, represents the female moth depositing the eggs; *b*, the eggs being deposited upon a chip which was cut from a fence while the female was at work; *c*, represents a top view of one egg magnified; *d*,

represents a side-view of the same egg; *e*, represents another egg with the canker-worm eating its way out; *f*, represents the canker-worm highly magnified, after it has crawled out from the egg; *g*, shows the natural size of the worm; *h*, shows the appearance of the male canker-worm moth. The female moth which is laying the eggs differs from the male in having no wings.

Now, if fresh elm-leaves are placed in the box with the worms, they will commence to feed on them. The eggs hatch out just as the leaves commence to grow, and consequently the young worms have tender leaves to feed on at the outset.

The worm grows rapidly, and after a few weeks ceases feeding, and, dropping to the ground, or lowering itself down by a thread spun from the head, buries itself just below the surface of the ground, and there changes into a chrysalis, forming a rude cocoon of earth about it. At the proper time there comes from the chrysalis a male canker-worm moth with wings, or a female canker-worm moth without wings.

73. From the eggs, then, come worms or caterpillars. The worms or caterpillars change into chrysalides, and sometimes these are inclosed in cases or cocoons. From the chrysalides come the perfect insects similar to the insects which first laid the eggs.

If it is desired to keep the caterpillars alive, the pupil should try to find them when they are feeding, and to observe the kind of leaf they are eating, and then, by giving them fresh leaves of the same kind as they need them, the

caterpillars will grow rapidly, and ultimately change into chrysalides. It is a common thing to see boys collect caterpillars and place them in a box, with grass to feed upon; the grass dries, and of course becomes unfit for food, and even if it were fresh the caterpillar would not eat it, unless it were its natural food. If the pupils wish to raise caterpillars, they must be sure and furnish them with the kind of leaf they are accustomed to feed upon. It may be an elm-leaf, or a cabbage-leaf. Thus, there is a common worm which they will find feeding on the leaves of the Tartarian honeysuckle, as in Fig. 78. To raise this worm, therefore, it is necessary to furnish it, from time to time, with the leaves of this honeysuckle.

FIG. 78.—WORM FEEDING ON THE LEAF OF THE TARTARIAN HONEYSUCKLE.
(Copied from figure by J. H. Emerton, in Packard's "Guide to the Study of Insects.")

In its growth, the caterpillar usually sheds its skin three or four times.

After the caterpillar has become full-grown, it stops eating, and crawls about in a restless manner.

In the summer and fall, various kinds of caterpillars are seen crawling over the sidewalks and along fences. They are not now searching for food, but are seeking an appropriate place where they can change into the chrysalis state.

From the long, active, and often devastating caterpillar, having three pairs of small legs in front, and other pairs of blunter legs behind, the creature gradually changes into a body, blunt in front, tapering behind, with no indications of legs, head, or any of the appearances seen in the caterpillar, except that the hinder part still shows the division of that portion into rings or segments as in the caterpillar, and signs of life are still manifested by this portion moving from side to side, when touched. Many caterpillars spin a case or cocoon, as it is called, in which it incloses itself previous to changing to a chrysalis. The thread with which they make this case issues from a little tube in the lower part of the mouth or labium. Silk is made from the thread composing the silk-worm's cocoon.

FIG. 79.—*A*, Hinder Portion of Chrysalis-skin hanging; *B*, *C*, Enlarged View of Hinder End, to show the little Hooks by which it hangs.

74. Fig. 79, *A*, shows the hinder portion of a chrysalis-skin, the insect having escaped from it, and the forward part

having fallen to the ground. The chrysalis is seen suspended by the tail, and is held there by little hooks on the end of the tail, which become entangled in a sort of web previously made by the caterpillar; *B* and *C* represent different views of the chrysalis-tail enlarged so as to show the hooks. Pupils will be sure to find these empty chrysalis-skins attached under the projecting portions of fences.

FIG. 80.—*A*, Caterpillar getting ready to change into a Chrysalis; *B*, Just ready to shed its Skin previous to changing; *b*, little Band to hold it up; *C*, Chrysalis; *D*, Butterfly just escaping from Chrysalis, the Wings just being unfolded.

Fig. 80 shows different stages of a cabbage-worm, from the worm stage to the chrysalis stage: *A*, representing the worm as it assumes a position under the projecting edge of a fence; *B*, after it has supported itself round the body by a delicate thread, *b*, and attached itself by the tail at the same time; and, *C*, representing its complete chrysalis condition; *D*, shows the butterfly just escaping from the chrysalis, the wings still being ruffled. After having escaped from the chrysalis, the butterfly generally clings to the empty case till the wings have expanded and dried, when it flies away. Fig. 80 represents a cabbage-butterfly introduced from Eu-

rope, and now common in certain parts of New England. Its wings are yellow, with two blackish spots on the forward wing, and one on the hinder wing. The chrysalides of this species are very common on fences, and, when collected in the fall, may be kept through the winter. During February and March the butterflies will come out, and these may be fed on honey or sugar mixed with water, and in this way may be kept alive for some time.

75. The caterpillar, then, having changed into the chrysalis, remains in this condition a few weeks, or even many months, and then the skin slowly cracks open, and out crawls a creature no longer like a caterpillar, but having three pairs of long, jointed legs, the body divided into three very distinct regions—the head, thorax, and abdomen—the thorax having wings, and the head furnished with long antennæ, and provided with mouth-parts suitable for sipping nectar, and no longer like the heavy jaws of the caterpillar, suited only to chewing coarse leaves; in short, a creature resembling the insect which first laid the eggs from which the caterpillar came.

76. Other names are given to these three stages of an insect. The worm, or caterpillar, is called the *larva*; the chrysalis is called the *pupa*; while the perfect insect is called the *imago*. These terms are necessary, for without them the proper condition of an insect could not be easily described.

Take, for example, the caterpillar stage of a butterfly: the same stage in a common fly is known by the name of

maggot, and in other insects the same conditions are known by the name of borer, grub-worm, and many other terms. If the pupil learns that all these various names describe a similar stage in the lives of these insects, it is much more convenient to have some general term describing all these stages, such as *larva*, or *larval stage*.

77. While most insects pass through changes similar to those above described, there are others, such as the grasshoppers, crickets, roaches, and bugs proper (a group of insects which includes the squash-bug, chinch-bug, and bed-bug, all of which have a disagreeable taste and odor, and to which naturalists restrict the name of bug), which do not pass through a caterpillar and chrysalis state. The young hatch from the egg, and closely resemble the adult insect,

FIG. 81.—DIFFERENT STAGES OF THE CHINCH-BUG: *a*, Egg; *b*, Newly-hatched Larva; *c*, Larva after First Moulting; *d*, Larva after Second Moulting; *e*, Pupa; *f*, Perfect Insect.

[These figures are copied from the Seventh Annual Report of C. V. RILEY, State Entomologist of Missouri.]

except that it has no wings, and is of course much smaller than the parent. In its growth it moults or sheds its skin, and each moult reveals its wings more advanced in growth,

till finally, on the last moult, it attains the size and features of the mature insect. And even in this growth, so unlike the moth and butterfly, the terms larva and pupa are applied to certain stages of its history.

The foregoing figure represents the egg and successive stages of the chinch-bug, an insect which has been so destructive to various crops in the West. The figures are all enlarged; the little line at the lower left-hand side of each figure represents the natural size.

78. Many insects, as the beetles, flies, moths, butterflies, bees, and wasps, pass through complete and distinct changes from their early condition to maturity, as above described. Other insects, as the grasshoppers, crickets, roaches, and bugs, hatch out from the egg, as little six-footed insects, and not as worms, and in their growth do not pass through an *inactive pupa* or chrysalis stage, but slowly acquire wings, and ultimately attain full growth as above stated. Hence these changes are not so completely defined as the changes in the insects first mentioned. For this reason the term *complete metamorphosis* is used to define the mode of growth of the beetles, flies, and other insects having a similar mode of growth; while the term *incomplete metamorphosis* defines the mode of growth of the grasshoppers, crickets, and others.

79. Many of the larvæ of insects look like worms—so much so, indeed, that they are commonly called worms, such as cut-worms, canker-worms, currant-worms, and the like. The pupils have learned that these are not true worms, but only the larval condition of certain insects.

True worms, however, never change into any thing else. Such, for example, is the earthworm, hair-worm, and leech, and worms which live in the sea.

FIG. 82.—EARTHWORM.

FIG. 83.—MARINE WORM.

These worms, and other true worms, generally speaking, have the body divided into a great many segments or rings, as in the earthworm. In the larvæ of insects, on the contrary, the segments are limited in number. With few exceptions the larvæ of insects have legs, and these legs in the fore-part of the body are jointed. In worms, jointed legs do not occur. The jointed legs of the larvæ number three pairs, and are on the three successive rings back of the head, and consequently correspond to the three pairs of legs in the adult insect.

In certain larvæ other legs occur, but these are not jointed, though often having special structures at their extremities.

by which they are enabled to cling. These are called *pro-legs*, or *false legs*.

It would be well for the pupils to collect some leeches and earthworms, and, if they live near the sea-shore, a few worms may be collected under stones at low tide.

Having collected these, let the pupils compare them with the larvæ of insects.

As the larva of an insect comes from the egg, it has its full number of segments at the outset. In the larva of a butterfly, for example, there may be counted, besides the head, twelve segments or rings, and this number does not increase as the creature grows, but remains constant; and, as we have already learned, the creature does not long remain in its worm-like stage, but assumes other conditions, ultimately becoming a creature unlike, in form and habits, the larval condition in which it spent a portion of its life.

80. The true worm, on the contrary, comes from the egg with a very limited number of segments, and as it grows new segments are formed, till in some worms as many as four or five hundred segments are developed before the animal has attained mature proportions, and in this condition it remains; that is, it is complete, never changing or passing through larval or pupal stages to develop into something quite unlike the worm.

Briefly, then, a larva may be distinguished from a true worm generally by its limited number of segments, and, when supplied with legs, having three pairs of jointed ones on the anterior rings of the body.

With few exceptions worms live in the water, and even those which live on the land are limited to damp earth or moist places. While the larvæ of insects are in many cases aquatic in their habits, and breathe or respire by means of gills, the larger number live on the land, feeding on leaves, wood, and substances of a similar nature, and are air-breathers.

CHAPTER XI.

HABITS AND STRUCTURE OF INSECTS.

81. LET the pupils now endeavor to study the habits of certain insects by direct observation. The following sketches are given as aids to the pupils in making independent observations on special insects.

Most insects make no provision for the larvæ, but leave them to take care of themselves, though usually the egg is deposited where the larva coming from it may find proper food at hand.

Other insects prepare cells or cavities in which they deposit their eggs, just as a bird builds a nest to hold its eggs. Certain insects, in preparing these cells, also lay up a store of food ready for the larva when it shall have hatched from the egg. Such is the case with the common mud-wasp. This insect makes a number of little chambers of mud, generally sticking them to the sides of a wall, or to the ceilings of sheds, barns, and attics. These pellets of mud are seen firmly plas-

tered to the wall, rough and irregular in appearance, and at first sight might be mistaken for the work of some mischievous boy.

FIG. 84.—A MUD-WASP'S NEST, HAVING TWO CELLS.

Fig. 84 represents their general appearance, for, rough as they appear, on close examination they are seen to be constructed very systematically—the soft clay of which they are built being laid on in alternate layers, looking something like a braid; while the upper portion, being built of looser and coarser material, is put on in irregular lumps. With a thin-bladed knife these mud-cells may be scraped off, and sometimes can be pulled away with the fingers. Upon opening them they will be found either filled with little spiders, or containing yellow-colored larvæ, pupæ in brown skins, or wasps. Their history is as follows:

82. When the mother-wasp gets ready to lay her eggs, she first builds these curious nests of mud, which she collects from the streets, or by the sides of brooks in clayey soil. Having finished one cell, she deposits therein an egg, and then collects a number of small spiders with which she completely fills the cell. After this is done she closes up the top of the

cell with looser mud. Thus she proceeds, constructing cell after cell, going through the same manœuvres with each one. It has been observed that the wasp stings the spiders so as to paralyze, but not to kill them. Hence they remain alive but cannot struggle, and when the egg hatches, the little larva coming from it finds in these spiders a store of food on which to feed. These are gradually eaten, and thus room is made for the rapidly-growing larva which, having eaten all the spiders, passes into its pupa state surrounded by its brown chrysalis case, and finally emerges a perfect wasp, when it softens the mud-walls of its nest, by a fluid poured from its mouth, and gnaws its way out.

FIG. 85.—SHOWING A NEST OF FOUR CELLS CUT OPEN: *a*, representing a Cell with the Egg at the Bottom, and the remaining Space filled with Spiders; *b*, the Larva full-grown, after having consumed all the Spiders; *c*, the Pupa; and *d*, the Imago, or Perfect Wasp, ready to come out.

Fig. 51 shows one of these mud-wasps pinned.

The pupils may collect these nests or cells in April or May, and by June the wasps will be ready to come out. If collected soon after they are made, the eggs may be found; if

a little later, the larva will be found feeding on the spiders; and, still later, the full-grown larvæ and pupæ appear.

In collecting for the cabinet, one nest should be cut open to show the cells and their contents.

83. The mosquito deposits her eggs on the surface of the water, sticking them together in such a way as to form a raft. From these eggs little black creatures hatch, which swim about with a quick, jerking motion. In this condition they represent the larvæ. If the pupils will examine pools and ditches, or even the tubs and barrels of water which often stand about farm-houses, they will be very sure to find some of these animals. They are small and black, and by their

FIG. 86.—*A*, Larva; and *B*, Pupa of a Mosquito; *c*, Line showing Natural Size of Larva; *d*, representing Water-Line.—The larva is seen with the hinder end of the body just projecting above the surface of the water, so that the air may enter through the little tube. The pupa is seen with the back just level with the surface of the water, and through the two tubes, on the back of the thorax, the air is being admitted to the body.

rapid, jerking motion cannot be mistaken. A number may be collected and placed in a jar of water, where all their changes, from the larval to the perfect state, may be watched. They will be seen coming to the surface of the water for air,

which they breathe in through openings in the hinder part of the body. Changes soon take place by which they assume the pupa state, and at this time they no longer breathe through the hinder portion of the body, but through two tubes on the back of the thorax. Finally, the pupal skin cracks open, and out crawls the perfect mosquito, for a while resting on the empty pupal case which floats in the water like a raft, and the insect remains supported in this way till the wings become fully expanded and dry, when it flies away.

CHAPTER XII.

HABITS AND STRUCTURE OF INSECTS (CONTINUED).

84. In certain groups of insects the young hatches from the egg, not as a caterpillar, but as a little insect having the body divided into three regions, possessing three pairs of jointed legs, and looking very much like the mature insect, except that it is very much smaller and has no wings.

FIG. 87.—YOUNG GRASSHOPPER.—*w*, Wing just appearing.

In the grasshopper, for example, the animal does not pass through a series of abrupt changes, but the creature comes from the egg with the general proportion of the adult insect,

save that it has no wings, these growing gradually as the creature increases in age. Fig. 87 represents a young grasshopper with the wings just appearing. At intervals, the insect sheds its skin, or *moults*, the wings continually increasing in length until mature size is reached.

Let the pupils endeavor to collect some young grasshoppers representing various stages of growth, and place these in their collections beside the full-grown one.

By searching in the grass, the cast-off skins of grasshoppers may be occasionally found still clinging to the spears of grass, where they were left when the grasshoppers shed them.

FIG. 88.—CAST-OFF SKIN OF A GRASSHOPPER.—The grasshopper has shed its skin while clinging to a blade of grass. The skin is imperfect, the antennæ and parts of its legs are broken; the abdomen is shriveled, and does not show.

Fig. 88 shows the appearance of one of these cast-off skins.

85. Grasshoppers are often infested with parasites. Frequently the grasshoppers, in a sickly condition, are met with clinging to the grass, or bushes. A careful examination of them will show a number of little bright-red mites crawling on them, or attached near the base of the wings, and evidently the cause of their weakness.

Curious cases have been found wherein these creatures had met with fatal accidents. In their headlong fall to the

FIG. 89.—GRASSHOPPER PIERCED WITH SPEAR OF GRASS.

ground, after one of their reckless jumps, they are liable to have their armor pierced with the dried spears of grass. Fig. 89 represents a grasshopper which had been pierced in this

way, the dried point of the grass probably striking the head, and then glancing off, and entering between the head and the thorax.

86. In studying the early stages of the mosquito, it was found that at the outset the animal breathed air through an opening in the hinder part of the body; that soon after this the opening closed, and air was taken in by two openings on the back, but in no case did the insect breathe through its mouth. In the perfect insect, as well as in most larvæ, there are little openings along the sides of the body. These little openings communicate with tubes which branch, and subdivide again and again, sending their little air-twigs into every part of the body, even into the legs and the veins of the wings. These little tubes represent the lungs of an insect. They necessarily render the body very light besides.

FIG. 90.—INSECT SHOWING THE SPIRACLES.—Grasshopper with the wings and two hinder pairs of legs removed to show spiracles, or openings in the sides of the body which communicate with the air-tubes within the body: *w*, showing where the wings were attached; *h* and *m*, where hind and middle legs were attached; *s*, spiracle on thorax; *t*, tympanum.

In large insects like the grasshopper the minute openings in the sides of the body can be plainly seen without the aid of a glass. The segments of the abdomen have each a little

opening, which is represented in the figure ; and where the abdomen joins the thorax, a cavity lined with a delicate skin will be found, which is called the *tympanum*, and is supposed to be an organ of hearing. If the softer parts within the body of an insect be removed and slightly compressed between two pieces of thin glass, the air-tubes, looking like fine white threads, may be seen with an ordinary pocket-lens.

The air-tubes are called *tracheæ*, and the openings on the outside of the body which communicate with them are called *spiracles*.

87. Insects breathe by dilating and contracting the abdominal segments. The act of breathing can be plainly seen in the grasshopper or the honey-bee, and it will be noticed that after violent exercise, as in a long flight, the insect breathes more rapidly than when it has been at rest for some time, just as a boy after running finds himself compelled to breathe rapidly for a while.

After violent exercise the insect gets tired and rests. Bees may often be seen, after a long flight, to alight in the grass near a flower, and for a while appear so fatigued that they cannot reach the flower, but remain breathing very rapidly. Insects have curious ways of resting and sleeping. A species of wasp has been observed soundly sleeping while holding on to a blade of grass by its jaws alone, the fore-legs just touching the grass, while the body and the middle and hinder pair of legs were hanging downward, and not bearing against the grass at all, as shown in Fig. 91.

88. In this connection it may be well to allude briefly to

the manner in which the various sounds emitted by insects are made. It is obvious that the vibration of the wings produces the loud buzzing sound made by certain insects. But there are other sounds which are traced directly to the effect of the air rushing in and out of the spiracles, and impinging on certain plates whose sharp edges border the spiracle. The experiment has been made of closing the

FIG. 91.--WASP SLEEPING WHILE HOLDING ON TO A BLADE OF GRASS WITH ITS JAWS.

spiracles with varnish, when all noise ceased. It is believed that the mosquito produces its remarkable tones in this way. Such noises have always been associated with the vibration of the wings, because the noise seems to be made when the insect is flying, but the cause of this is explained by supposing that the violent muscular action of moving the wings also causes the air to be violently thrown out of the spiracles, and as a proof of this it has been found that cutting off the

wings of such an insect, while modifying the sound, by no means prevented it being made ; and it is a fact also that no sound is produced by other insects whose wings vibrate with great energy.

89. The peculiar chirp of the cricket is made by the fore-wings being rubbed rapidly against the hind-wings upon which they rest ; one of the large veins in the fore-wing being thickened and notched like a file, and the wing itself acting as a resonant body in augmenting the sound. The males, only, make this sound ; the females are silent ; and if the fore-wing of the female be examined, the vein in question is not thickened, nor is it rough, like a file.

Mr. Samuel H. Scudder has stated that the grasshoppers produce their sound, or *stridulate*, in four different ways : “ 1. By rubbing the base of one wing-cover upon the other, using for that purpose the veins running through the middle portion of the wing ; 2. By a similar method, but using the veins of the inner part of the wing ; 3. By rubbing the inner surface of the hind-legs against the outer surface of the wing-covers ; and 4. By rubbing together the upper surface of the front edge of the wings and the under surface of the wing-covers. The insects which employ the fourth method stridulate during flight—the others while at rest. To the first group belong the crickets ; to the second, the green or long-horned grasshoppers ; to the third and fourth, certain kinds of short-horned or jumping grasshoppers.”

90. Harris, in describing the third method of stridulation,

says that “their instruments may rather be likened to violins, their hind-legs being the bows, and the projecting edge of the wing-covers the strings,” and adds that when a grasshopper begins to play “he bends the shank of one hind-leg beneath the thigh, where it is lodged in a furrow designed to receive it, and then draws the leg briskly up and down several times against the projecting lateral edge and veins of the wing-cover. He does not play both fiddles together, but alternately, for a little time, first upon one, and then on the other, standing meanwhile upon the four anterior legs and the hind-leg which is not employed.”

FIG. 92.—Leg of a Grasshopper magnified, showing Ridge of Fine Teeth on the Inside of the Leg, marked *a*, by which the Insect rasps the Wing; *b*, *c*, Different Views of Ridge of Fine Teeth, highly magnified.

A figure is here given of the hind-leg of a common grasshopper, showing the row of minute teeth which occur on the inside of the leg, and which are drawn across the edge of the wing. The pupils may imitate the sound thus produced by drawing a coarse file, or the teeth of a comb, rapidly across the edge of a stiff sheet of paper.

CHAPTER XIII.

HABITS AND STRUCTURE OF INSECTS (CONTINUED).

91. A LITTLE bug, called the tree-hopper, has a peculiar history in its young state. The eggs of certain species are laid in the ground, and, as soon as these hatch, the young ones crawl up the stems of grass, and, piercing the grass with their mouth-parts, commence to suck the juices contained therein. While this action is going on, a clear, watery fluid escapes from certain pores in the body, and in a short time the young insect is completely immersed in it. As it is obliged to breathe air, it secures this by turning up the hinder part of the body, and by means of little appendages, clasping a bubble of air, which then flows along the under side of the abdomen; here it is taken in through the spiracles. The air having been so used, is allowed to escape in the fluid in which the insect is immersed. This operation is repeated over and over again, fresh bubbles of air being thus secured, and then escaping in the fluid. After a while the fluid becomes filled with these little bubbles, which soon convert it into a frothy substance, and this is the origin of the white flecks which occur so thickly on grass, and which is here commonly called *frog-spit*, and, in England, *cuckoo-spittle*.

92. There are certain insects belonging to the same group which are aquatic, and whose young come to the surface of

the water, and in the same manner secure air. So this little tree-hopper, while in the young state sucking the juices of grass, and completely immersed in a watery fluid, may be looked upon as an aquatic larva during this stage.

FIG. 93.—GRASS, WITH THE FROTH UPON IT, *a, a*, and a Young Insect exposed at *b*.

FIG. 94.—A PORTION OF A GRASS-STEM, WITH THE YOUNG INSECTS MAGNIFIED: *a*, the Insect reaching out the Hinder Part of the Body to secure a Bubble of Air; *b*, an Insect allowing a Bubble of Air to escape in the Fluid, the dotted line *b* indicates the bubble; *c*, the Mouth-parts, like a Sting, piercing the Grass.

Let the pupils collect and examine this froth, and, by carefully wiping it away, they may expose the young insect fastened to the grass.

93. The insect called the seventeen-year cicada, or seven-

teen year locust as it is improperly called, has an interesting life-history.

The perfect insect is shown in Fig. 95. They may be known by the peculiar loud, buzzing sound emitted by the male. This sound can oftentimes be heard at a great distance.

FIG. 95.—SEVENTEEN-YEAR CICADA.

The seventeen-year cicada is found rarely in Southern New England, but is common in the Southern and Western States. This species exists in great numbers, and does immense damage to the trees which it infests. The female deposits her eggs in the twigs and smaller branches of oaks. Little furrows are made in the twig, side by side, and into these furrows the eggs are laid. The leaves wither on the trees from the injuries inflicted in this way. Little insects hatch from the eggs, entirely different in appearance from the parent; and these, running to the end of the twig, fling themselves off, and falling to the ground dig their way down, till they come to some root upon which they fasten, and with a piercing sting suck the juices of the root. Here they remain for nearly seventeen years, slowly growing. At the end of that time they assume the

appearance represented in Fig. 96. At the proper time they crawl out of the ground, and their skins splitting open along the back, out come the perfect insects, with wings and all complete, to fly away, lay their eggs, and devastate the forests.

There are certain species which do not occur in such numbers, and which pass through all their changes in a single year. The cast-off skins of the pupæ of such species may be often found clinging to apple-trees and fences in New England.

FIG 96.—PUPA-CASE OF A SPECIES OF NEW ENGLAND CICADA OR HARVEST-FLY, CLINGING TO A TWIG.

The pupils should, if possible, collect a twig in which the eggs have been deposited, a pupa-skin, and the perfect insect.

CHAPTER XIV.

HABITS AND STRUCTURE OF INSECTS (CONCLUDED).

94. THE May-fly, or *Ephemera*, is one of the most common insects in the Western States. They live only a few weeks in their perfect state, oftentimes but a few days. Their eggs are laid in the water, and the larvæ live in the water two or three years. At the end of this time they come to the surface in immense numbers, and, shedding their skins, come out as winged insects. At this time they resemble their perfect state so closely, that the name *sub-*

FIG. 97.—LARVA OF EPHEMERA.

(Reduced from Figure by J. H. Emerton, in Dr. A. S. Packard's Directions for collecting Insects, Smithsonian Institution.)

imago is given to them in this condition. They often fly a considerable distance from the water, alighting on the ground and trees. Here they again shed their skin, and then have attained their perfect state.

These insects occur in prodigious numbers in certain parts of the world. In some regions of Europe they are so

FIG. 98.--EPIHEMERA.

abundant that the inhabitants collect them in heaps, and use them as dressing for the land. In the cities bordering the

great lakes it is a common sight to see the gas-posts and adjoining buildings blackened by the myriads of *Ephemera* which have been blown in from the lakes and have been attracted by the lights. The following figure represents a gas-post, in Cleveland, Ohio, as it appeared with *Ephemera* clinging to it:

FIG. 99.—GAS-POST, WITH EPHEMERA CLINGING TO IT.

95. Another group of insects somewhat resembling the *Ephemera* pass their larval state in the water.

Some of their larvæ are called *caddis-worms*, or case-

worms, and are inclosed in cases of cylindrical and other shapes. These are variously made of grains of sand, bits of bark and sticks, and other fragments of convenient size cemented together. Some of these cases, built of small

FIG. 100.—CADDIS-WORM, WITH ITS CASE.

grains of sand, look like coiled snail-shells. Other larvæ shelter themselves in bits of straw, or the fragments of hollow stems of plants.

Fig. 100 represents the larva of one of these insects in its case, which is made of bits of sticks arranged in a spiral course. The larva drags about this case, and as it grows collects material for the enlargement of its tube.

In almost any quiet pool or running stream these curious cases may be found, containing the larvæ within.

FIG. 101.—GALL-FLY.

96. The curious round balls called gall-nuts, which are found on the leaves of the oak and of other trees, are produced by an insect called the gall-fly.

The eggs are deposited in the substance of the leaf, and

it may be that the larva, by its presence there, causes the unnatural growth of the leaf, resulting in a wart or tumor, and sometimes in a large round nut. It is believed, however, that the adult insect, in depositing the egg, also stings the leaf, and, poisoning it at the same time, induces the abnormal growth of the leaf. Within this the larva feeds,

FIG. 102.—GALL-NUT ON OAK-LEAF.

(Copied from Harris's "Insects injurious to Vegetation," third edition.)

and changes into the pupa state, and finally into the perfect insect, when it gnaws its way out.

In the autumn the pupils will find the gall-nuts abundantly in the woods. Let them collect a number of these, and, on carefully cutting them open, they will find within a tiny oval case, and upon opening this they will discover snugly stowed away a little, polished black fly having four wings. The creature when liberated is ready to fly away.

Some of the nuts will be empty, because the gall-insects have already escaped.

Galls are also produced by other kinds of insects. The following figure, which represents a gall common on the golden-rod, is produced by a two-winged fly. The figure represents the stem or stalk unnaturally swollen, the swollen portion being the gall, within which the larva, pupa, or perfect insect, may be found if the creature has not already escaped.

FIG. 103.—GALL ON GOLDEN-ROD STALK.—*p*, Pupa removed from the Gall.

Let the pupils arrange in their collecting-box a leaf with the nut attached, a nut cut open showing the pupa-case, and the insect pinned.

97. Only a few brief lessons have been given representing the life-history of a butterfly, mud-wasp, mosquito, spit-

tle-insect, seventeen-year cicada, May-fly, and gall-fly. Let the pupils endeavor from their own observations to make additional life-histories, or record facts, concerning other insects, such as the honey-bee, paper-wasp, and a great many other common insects, of which no mention has been made here. The turning over of stones and logs in the woods will oftentimes expose the burrows of ants, and the ants will probably be found busily engaged in carrying off long, white, oval cases, which look like eggs; let the pupils collect some of these, and see if they can find out what stage in the history of the insect they represent.

98. An instinct which appears wonderful to us, prompts the insect to seek appropriate places for the deposition of her eggs. The butterfly, for example, seeks for food the nectar of flowers; its larvæ, however, must have leaves upon which to feed, and the instinct of the butterfly impels it to deposit its eggs in a place where the young shall find their appropriate food. It has been learned also that other insects store up animal food for their young, as in the case of the mud-wasp, where spiders are imprisoned in cells in which the eggs have been previously laid.

The gall-flies deposit their eggs directly in the substance of the leaf.

99. Another group of insects, much resembling the gall-flies, deposit their eggs directly in the bodies of the larvæ and pupæ of other insects. They are called *ichneumon-flies*. These insects have on the hinder part of the body a

sharp, piercing sting, and with this organ the necessary hole is made through which the egg is deposited.

A caterpillar soon hatches from the egg thus deposited by the ichneumon-fly, and feeds upon the fatty portions of the body of the larva in which it has been so placed. But this larva containing the ichneumon-caterpillar, meanwhile, completes its growth and changes into a chrysalis, when the inclosed ichneumon-larva devours the entire contents of the chrysalis, and then changing into the pupa state soon emerges as an ichneumon-fly, to go in quest of caterpillars, in which to deposit its eggs. Thus it will often happen that a number of cocoons have been collected, from which ought to appear a certain kind of moth, for example, but from many of them a brown ichneumon-fly will emerge, a sight quite as startling, to one not familiar with insects, as if a robin should be seen to hatch from a hen's egg.

If the pupils will collect from the fences a large number of the chrysalides of the common yellow cabbage-butterfly, and keep them in a box, with a piece of glass for a cover, they will observe that while butterflies come from many, from others, which have already changed to a lighter color, little black flies will appear, crawling out of holes in the side of the chrysalis which have been made by some of the imprisoned ichneumons. (*See Fig. 104.*)

100. Nearly every species of insect is infested with one or more species of ichneumons, which deposit their eggs either within the pupæ, larvæ, or the eggs themselves.

There are some species of ichneumons which deposit their

eggs within the eggs of the canker-worm moth, and, as tiny as these eggs are, they are still large enough to furnish nourishment and room for the complete development of the insect feeding within.

In Fig. 74 an ichneumon-fly is shown on the wing, in search of caterpillars wherein to deposit her eggs.

Fig. 67 also represents an ichneumon-fly of large size.

Fig. 104 represents ichneumon-flies escaping from the chrysalis of the cabbage-worm butterfly.

FIG. 104. — CHRYSALIS OF THE CABBAGE-WORM, FROM WHICH ARE SEEN ESCAPING ICHNEUMON-FLIES.

CHAPTER XV.

SPIDERS.

101. For this lesson the pupils are to collect a number of spiders, securing, if possible, the largest specimens. A wide-mouthed bottle, with a little alcohol, will answer to collect them in. Let each pupil select the largest specimen to study, and pin it to a piece of cork, or to a soft pine strip. The legs are to be arranged with two pairs pointing forward and two pairs pointing backward, as shown in Fig. 105.

Let them study the following characters with the specimen before them :

The spider is divided into two regions. That region or part to which the legs are attached is called the cephalo-

thorax. The hinder region is called the *abdomen*. Instead of having a separate head, as in true insects, the spider has its head and thorax combined, and hence this part is called the cephalo-thorax, a compound word meaning head-thorax.

FIG. 105.—COMMON GARDEN SPIDER.—*A*, as seen from above; *B*, as seen from below; *p*, Palpi; *m*, Mandibles; *s*, Spinnerets from which the Spider's Thread issues.

102. The spider has four pairs of legs, instead of three pairs of legs as in the true insects. Projecting in front are a pair of jointed feelers called *palpi* (see Fig. 105, *p*). These look very much like legs, and in very young spiders can scarcely be distinguished from them.

FIG. 106.—SIDE-VIEW OF COMMON GARDEN SPIDER.—*C*, Cephalo-thorax; *A*, Abdomen; 1, 2, 3, 4, First, Second, Third, and Fourth Pairs of Legs; *s*, Spinnerets; *m*, Mandibles; *p*, Palpi.

The mouth is armed with a pair of jaws which are attached above the mouth and hang down in front, at the end of which are the poison-fangs. With these they are enabled to secure and kill the flies and other insects upon which they feed. The following figure represents the jaws or mandibles.

FIG. 107.—FRONT PORTION OF COMMON GARDEN SPIDER GREATLY ENLARGED, showing, *pp*, Palpi; *m*, Mandibles; *ll*, a Portion of the First Pair of Legs, and above, the Front of the Cephalo-thorax, with the Eight Eyes upon it.

Directly behind the mandibles, are two smaller jaws, called maxillæ (*see* Fig. 108), which aid in crushing the food and arranging it for the mouth.

FIG. 108.—INNER JAWS, OR MAXILLÆ, OF A COMMON GARDEN SPIDER.—The first Joints of the Palpi are seen also.

The spider has eight eyes, situated on the front part of the cephalo-thorax. They look like little black beads, and in

large spiders can be easily seen without the aid of a magnifying-glass.

103. The *abdomen* has little appendages at its hinder end called *spinnerets*, and from these the spider produces the thread with which it builds its nests and nets, the nets being commonly called spiders' webs.

Highly magnified the spinnerets appear as blunt protuberances arranged together in pairs, and capable of being contracted or expanded. These spinnerets are covered with hundreds of jointed hairs which are perforated and through which the web-forming material escapes. This material is fluid and something like the white of an egg. Escaping from the body, through hundreds of these minute openings, the strands of this fluid dry almost instantly, and, uniting, form the delicate, yet comparatively strong, thread of the spider. Thus it will be seen that the thread of the spider is composed of hundreds of strands, which may be often separated just as the fibres of a rope may be pulled apart. Under the microscope the posterior end of the abdomen with the spinnerets looks like this.

FIG. 109.—SPINNERETS OF A SPIDER.—*t*, one of the Tubular Hairs from the Spinnerets, highly magnified.

104. As the thread issues from the spinnerets, the spider guides it with its hind pair of feet, and these are curiously adapted for the purpose not only of holding and guiding the thread, but also of enabling the spider to run rapidly across its nets without getting entangled, while other animals become helplessly ensnared in attempting the same thing.

The ends of the legs terminate in three claws, a pair of larger ones generally notched like a comb, and a third one like a spine which acts as a thumb. Other notched spines or hairs also aid in securing a hold upon the web, and even if these fail to secure a footing, the leg itself is covered with long stiff bristles pointing downward which are sure to catch in the web. The two large notched claws, as well as the other claw and spines, are highly polished, and consequently present no roughened surface to which the thread will adhere.

The following figure (Fig. 110) represents the end of a spider's leg magnified, showing the arrangement of hooks and claws.

FIG. 110.—END OF A COMMON GARDEN SPIDER'S LEG MAGNIFIED.—*o*, Outer Claws; *m*, Middle Claw; *t*, Toothed Hairs.

105. By observing the spiders which build their nets across the openings of windows and in other convenient

places, and while at work, they may be seen to use their hind-feet in apparently drawing out the thread as it were from the spinnerets. It will be observed that the thread issues in a broad band, and, when these spiders are sluggish, their thread may be caught on the end of a pencil by gently rubbing the spinnerets with it, and then by withdrawing the pencil the thread may be reeled off.

The various kinds of nets are adapted to entrap the spider's food, which consists of flies and other insects.

Certain kinds of spiders do not build nets, but go in search of their prey by stealthily creeping up and pouncing upon it unawares.

It is a very interesting sight to watch the little black-and-white spider (so common on the sides of houses) slyly approach a fly which has alighted near it. If the spider is on the side of a window-sill and a fly has alighted near it, the spider instantly turns round, facing the fly, cautiously and very slowly moves backward, till it gets on the upper side of the window-sill and out of sight, when it rapidly approaches, now and then peering over the edge of the sill, to see where the fly is, and, finally getting directly above the fly, it gathers its legs for a jump, securing its thread to the window-sill at the same time, and then with a sudden spring seizes the fly in its jaws. Sometimes the insect is much larger than the spider, and flies away, with the spider tightly clinging to it; the thread, however, holds fast, though sometimes run out to the length of a foot or more. Soon the poison of the spider takes effect, and the fly gradually weakens, and ceases its strug-

gles, when the spider carries it off to some nook, there to devour it.

106. One of the most common spider-nets is like the one shown in Fig. 73. If the place selected is in the opening of a window or similar place, the spider first runs a few threads as a sort of framework, to which are to be afterward attached the radiating threads, that is, those which run from the centre of the net to the sides. Having arranged these so near together that the spider can easily reach from one radiating thread to the other, the creature commences at the centre of the net, and runs a thread from one radiating thread to the other in a rapidly-unwinding spiral till it reaches the outer edge of the net. This is to form a staging, and also the better to hold the radiating threads in place. It then commences at the outside, and going back over its last course carefully constructs the permanent mesh; and, as it comes to each radiating thread, it will be seen to attach to it the thread it is now making, by simply pressing the spinnerets against it. As it goes around again and again, continually lessening the circle, it gathers up the thread which was first laid as a staging, and, rolling it up in little balls, drops it to the ground. This habit has led to the impression that the spider eats its web. The circular threads are glutinous, while the radiating threads are smooth, and this can be proved by throwing dust through the net, when the cross-threads will catch and hold the dust, while the radiating threads will remain clean. The actual centre of the net is not the geometrical or true centre, but a little above it.

It may be observed, too, that the net does not stand vertical, but leans a little, and the spider having completed the net takes a position in the actual centre of the net, head downward and on the inclining side of the net. With its legs outstretched, and resting on the radiating lines, it can feel the slightest jar or agitation made by a struggling insect. The spider being above the true centre of the net and on the inclining side, if the fly has become entangled below the centre, it can instantly drop to the desired point suspended by the ever-ready thread which it makes, and, swinging to the net, it almost instantly catches the fly.

The pupils would do well to watch the spiders while they are constructing their nets, and to observe and describe, or sketch in outline, the different kinds of nets they find and the kinds of spiders which construct them.

107. Besides the nets made by spiders to ensnare insects, some species have the power of running out a long thread which answers the purpose of a balloon in raising them from the ground and carrying them floating a long distance in the air. In constructing this buoyant means of transportation, the spider does it at peculiar times of the day, and in peculiar positions. Selecting some place where the heated air is rising from the ground or from the side of a fence, it turns up its abdomen and allows the rising current of air to carry upward the thread which is being made, and, when this thread is of sufficient length for its buoyancy to overcome the weight of the spider, it floats away with the spider hanging below.

The following represents the young spider in the atti-

tude of throwing out its thread for the purpose of sailing in the air.

Voyagers often meet with these spiders in myriads as the wind sweeps them from the land.

FIG. 111.—YOUNG SPIDER GREATLY ENLARGED, SHOWING ITS ATTITUDE IN THROWING OUT THE THREAD, PREVIOUS TO RISING FROM THE GROUND. (Copied from a Figure by J. H. Emerton.)

103. The spider also constructs cases to hold her eggs, and lines them warmly with the finest web. These nests vary greatly in appearance. A very common variety, somewhat oval in shape, may be found suspended in barns and sheds. The pupils should collect and open these cases or nests, and they will be found to contain little eggs, sometimes rolling out like beads into the hand, or, the eggs having hatched, hundreds of little spiders will appear moving within the nest.

Nests, or, more properly speaking, egg-cases of different kinds, may be collected under stones and logs, and wherever spiders' nests occur. The little spiders hatching from the egg

will grow to twice their size in the nest, without apparent food, and it becomes evident that, in some cases, they must eat each other, as Prof. Wilder has observed within some of the egg-cases a far less number of spiders than there were eggs in the nest at the outset. These nests may be kept in boxes, and the eggs will hatch in due time.

FIG. 112.—SPIDERS' NESTS OF DIFFERENT KINDS CONTAINING EGGS.—*A* and *C* are common nests in sheds and barns; *B* was found under a board in the field, the part containing the eggs stands upon a stalk.

109. The young spider comes from the egg resembling in form the parent spider, except that the legs are much shorter in proportion to his relative size, and the palpi appear so large that they look like another pair of legs, as they then are in fact, but they afterward become modified to feelers.

FIG. 113.—ENLARGED FIGURE OF A YOUNG SPIDER JUST FROM THE EGG, WITH THE FIRST MOULT, *m*, adhering to the Hinder Part of the Body; *y*, the Natural Size of the Spider; *l*, extremity of a Leg highly magnified, showing an Outer Skin which has not been shed.

As the young spider grows, it sheds its skin at short intervals of time. If the pupils will examine the young spider soon after it is hatched from the egg, they will find attached to the hinder part of the body the skin which has just been shed. This curious process of shedding the skin, or *moulting*, occurs at intervals, till the spider has reached adult size.

FIG. 114.—THE CAST-OFF SKIN OF AN ADULT SPIDER.

110. The cast-off skins of spiders are very common in their webs, and, if the pupils examine any barn-window which is covered with spider's webs, they will be sure to find some of these cast-off skins, like the one represented in Fig. 114.

The mother-spider, generally so timid, overcomes her fear during the time she has the care of her eggs, and with many spiders the egg-cases are directly cared for by the mother, she oftentimes carrying them about with her or holding on to them and showing the greatest solicitude for their safety. Let the pupils try to separate the egg-case from the mother-spider, and they will then learn how courageous the spider is at this time, and how persistently she remains by her eggs. Some species of spiders carry their young on their backs, and move about with them.

A small black spider was picked up in the woods, which had her body entirely covered with young spiders, which were evidently newly hatched. When the mother-spider was picked up, all the little spiders becoming frightened jumped off, but just before jumping each one attached a tiny thread to its mother's back, and as the spider was held up in the air there hung below, suspended by invisible threads, the whole progeny looking like little black beads. The mother-spider was then thrown down among the dead leaves, sticks, and pine-cones. She did not run away, however, but waited till all of the young ones had found their way through this tangled wilderness, safely back to their mother, and this they accomplished by means of their threads, one end of which they had previously attached to her back. Having waited till all had been gathered in this way, she continued her journey.

111. The spider has no power of throwing or ejecting its thread to distant objects, as many suppose. When threads are seen stretching from one tree to another, the spider has caused the thread to issue from the spinnerets, and the wind has then caught it and borne it along, till finally it gets entangled with some object, and in this way the spider is enabled to cross from one point to another.

These creatures are not so dangerous as many suppose, and but very few authenticated cases are known of man having been bitten by these animals; though the larger spiders at the South, and in California, as the tarantula, for example, can inflict a dangerous wound.

CHAPTER XVI.

DADDY-LONG-LEGS, CENTIPEDES, AND MILLEPEDES.

112. IN the insects proper, or true insects, the pupil has learned that the head, thorax, and abdomen, are separated into three regions or parts. In the spiders, it has been seen that the head and thorax are combined, forming a single region or part, and called the cephalo-thorax, while the abdomen appears as a distinct part. There is another group of animals allied to the spiders, the individuals thereof having four pairs of legs, and the head, thorax, and abdomen, more or less merged together. The animals belonging to this group are called in various parts of the country, *daddy-long-legs*, *granddaddy-long-legs*, *grandfather-graybeards*, and *harvest-men*, and in northern New York are known by the name of “grab for gray bears.”

Certain species are common around houses and sheds, others are found in the woods. They are easily recognized by their small bodies and extremely long and slender legs. It is difficult to hold them in the fingers, as some of the legs are liable to drop off on the slightest effort made to retain the animals.

In the middle and on the back of the cephalo-thorax, there is a slight eminence, upon which are situated the eyes, two in number. The abdomen appears distinctly segmented.

113. Their food consists of small insects, such as flies and

mosquitoes; and these they go in quest of, slyly approaching and pouncing on their victim and seizing it with their mandibles, which are furnished at their ends with a pair of nippers, which enable them to retain their prey. (*See Fig. 115, m.*)

They build no net to entrap their prey, and are weak and helpless compared with their higher relatives, the true spiders. They are dependent then for food upon such insects as they can overcome, and these they devour, differing in this respect from the rapacious spiders which suck the fluid contents of their prey, rejecting the rest. Certain species are known to

FIG. 115. —YOUNG DADDY-LONG-LEGS, ENLARGED: *n*, showing Natural Size; *B*, Under Side of Body still more enlarged; *m*, Mandible of Left Side: *p*, Palpus of Right Side: *P*, Palpus, greatly enlarged: *P C*, End of Palpus, showing Notched Claw; *L*, End of Leg, showing Claw; *E*, Eye-Prominence, with the Two Eyes. (The three last-mentioned Figures are greatly magnified.)

be cannibals, as some have been seen to pounce upon a brother daddy-long-legs and devour it, leaving only the legs.

It is believed that in the Northern States they do not survive the winter, as in the spring only young ones are seen, and these attain full size by autumn. At this season, the eggs are laid under stones and in the cracks of boards and other protected places, where they remain to hatch out in the following spring.

114. Under old boards in gardens and hidden beneath stones and dead leaves in the fields and woods, the pupils will find the other creatures to be studied in this lesson. They are commonly known as centipedes, and in the Eastern States, at least, are also known as earwigs; though the earwig in England is an entirely different animal, being a true six-legged insect.

The centipede belongs to a group of animals called *Myriapods*, and the animals belonging to this group are composed of a great many similar segments, some species having as few as ten segments, others having over two hundred segments. In this latter respect, these creatures resemble the worms, but differ from the worms in having jointed legs and antennæ, in these last-named characters resembling the insects, besides having other affinities with them in breathing air through spiracles and tracheæ which run through the body.

There are two very distinct groups of Myriapods; one group comprising the true centipedes, in which the body is flattened, and segments loosely joined, and the legs gener-

ally equaling, and sometimes exceeding, the width of the body.

The segments in many cases are unequal in length, some of them being very short and alternating with long ones, though all bear a pair of legs below. The antennæ are much longer than the legs, and are often composed of a great many joints. A pair of modified legs reach out behind and look like a hinder pair of antennæ.

In a few forms the eyes are compound as in the insects, while in others the eyes are separate as in the spider, and are called *ocelli*. These are grouped on each side of the head, at the base of the antennæ.

FIG. 116.—COMMON CENTIPEDE,
NATURAL SIZE.

A magnified view of the head, showing group of eyes at the base of antennæ. A few joints only of the antennæ are shown.

115. The jaws or mandibles are large and jointed, with the terminal joint long and sharp as in the spiders. The other pairs of jointed appendages act also as mouth-parts. The under lip is notched with fine teeth, as shown in the following figure, which represents the under surface of the head of the species of centipede shown in Fig. 116.

These creatures are active in their motions, and rapacious in their habits. Some of them feed on small insects, others attack earthworms. Their bite is venomous to insects, and one species having very long legs will produce by its bite a severe pain lasting several hours. A large species found in the Southern States, and in the tropics, and commonly known

FIG. 117.—SHOWING MAGNIFIED VIEW OF THE UNDER SURFACE OF THE HEAD OF A CENTIPEDE: *a*, Antenna; *mp*, Maxillary Palpus; *fj*, Foot-jaw; *f*, Poison-Fang of Foot-jaw; *l*, Labium; *m*, Maxilla. The Mandibles are hidden behind the other parts, and do not show.

as the centipede, is considered a dangerous animal from its bite. The feet of this species are supposed to poison by their touch, since, when they run over the flesh, small ulcers appear where the feet have come in contact with the skin. The pupils may collect these animals, and either dry them and stick them to cards, or preserve the specimens in vials filled with alcohol.

116. The other group of myriapods, commonly known as

millepedes, have a long, cylindrical, and oftentimes shiny body, composed of a great many segments so smoothly joined together that it is difficult to see the separation between them.

The antennæ are short, there are no long caudal appendages, and the legs are short and feeble. At first sight it would appear that these creatures were exceptional among insects and spiders, in having two pairs of legs to one segment; but it has been learned, by studying the very young millepede, that there is really but one pair of legs to a segment, but that the segments grow together in pairs, so that each apparent segment is really two segments united.

FIG. 118.—A COMMON MILLEPEDE. The line underneath the figure represents the length of the specimen from which the drawing was made. *A*, a Magnified View of the Head of the Millepede represented above; *B*, a Magnified View of the Left Jaw.

These creatures live on decaying matter, and are slow and weak in all their movements. When touched, or alarmed, they coil up in a closely-wound roll. The body is hard, and the animal can be stuck on a card for the cabinet. The eggs, to the number of sixty or more, are laid in little burrows

previously prepared by the creature in earth that is neither too moist nor yet too dry. In preparing the burrow the female makes use of the fluid which comes from her mouth, and which enables her to stick the earth together in little balls, and these she passes up from her burrow by means of the little legs which grasp the pellet and convey it from one pair of legs to the next pair, and so on till it is thrown out of the burrow. After the burrow is completed, and the eggs laid, the entrance to the nest is carefully filled up with clay, or dirt, moistened with fluid from the mouth.

117. It has been learned, in studying the development of the insect proper, that the worm-like larva comes from the egg with its full number of rings or segments, and that, as the creature matures, some of these segments are so merged into other parts, particularly with some of the caudal ones, that it seems as if the perfect insect has a less number of rings than the larva. In the myriapods, however, the young creature as it hatches from the egg possesses only a few seg-

FIG. 119.—HIGHLY-MAGNIFIED FIGURE OF A VERY YOUNG MILLEPEDE, SHORTLY AFTER HATCHING FROM THE EGG.

(Reduced from a figure by Elias Metschnikoff.)

ments, but as it grows, new segments are from time to time formed near the hinder part of the body, until the creature

attains adult size, when it may possess over a hundred segments.

Like true insects, however, the young myriapod makes its appearance from the egg with three pairs of legs. The body, however, is never divided into a thoracic portion, and an abdominal portion, as in the true insects, or into two regions as in the spiders, but after the head there succeeds a continuous row of similar segments to the tail.

118. In studying the insects, spiders, and centipedes, or myriopods, the pupils have learned something about three groups of animals which have in common a body composed of segments, and possessing jointed legs. They all breathe air through holes in the side of the body, called spiracles, the air, thus breathed, finding its way through various parts of the body by means of little tubes called tracheæ, except in the spiders, where little sacs, called pulmonary sacs, take the place of tracheæ.

In the true insects the segments of the body are gathered into three regions, called respectively the head, thorax, and abdomen. In the spiders the segments of the body are gathered into two regions, called respectively the cephalo-thorax, and abdomen, the head being merged in the thorax. In the myriapods the head is again distinct as in the true insects, but the remaining segments of the body are distinct and are not grouped into regions.

The true insects have three pairs of legs. The spiders have four pairs of legs, while the myriapods have no definite number of legs. In some species there are nearly two hundred

pairs of legs, and in no species are there less than ten pairs of legs. The true insects alone have wings.

119. In the growth or development of the true insects and spiders, the young animal comes from the egg with its full number of segments complete, while in the myriapods the young animal comes from the egg with a few segments, and new ones are added as the animal grows.

Some of the characters of the insects, spiders, and myriapods, may be represented as follows:

Three pairs of legs, and having wings.
TRUE INSECT.—Body divided into Three Regions.

Four pairs of legs.
SPIDER.—Body divided into Two Regions, Head not separate.

No definite number of legs
MYRIOPOD.—Body not divided into Regions, but Head separate.

FIG. 120.—ANIMALS WHOSE BODIES ARE COMPOSED OF SEGMENTS POSSESSING JOINTED LEGS, AND BREATHING AIR THROUGH OPENINGS IN THE SIDES OF THE BODY.

On account of some of these characteristics above mentioned, with others not mentioned, being held in common by the true insects, spiders, and myriapods, these creatures form a natural group in the animal kingdom, just as the snails,

mussels, oysters, and clams, possessing certain characters in common, together form a natural group of animals.

There are, however, many other animals which are not insects, spiders, or myriapods, and still possess a body composed of segments, and also have jointed legs, and these animals are to furnish the subject for the next lesson.

CHAPTER XVII.

CRAWFISH AND LOBSTER.

120. THE fresh-water Crawfish, or fresh-water Lobster as it is sometimes called, is very common in many of the Western rivers. It may be collected in little pools by the river-side, and kept alive for a long while in a jar of water. It may be fed on fresh-water snails and the larvæ of insects. It would be well to keep the animal alive for a while, so that its motions in swimming and crawling may be observed. For the cabinet, it can be dried with the legs outstretched, or specimens may be preserved in alcohol.

The general form of the body is much like that of the salt-water lobster, differing, however, greatly in size; the crawfish varying from three to five inches in length, and the lobster attaining a much larger size.

The animal is divided into two regions, the body proper, to which the legs and big claws are attached, and the abdomen, consisting of the jointed portion behind. The head

does not appear separated from the body as in the insects, but is combined with the thorax, and hence this part is called the cephalo-thorax, as in the spiders. The cephalo-thorax is covered by a continuous shield, or shell, called the *carapace*, while the abdomen is divided into a series of segments. This part can be bent snugly beneath the body (*see* Fig. 121).

FIG. 121.—FRESH-WATER CRAWFISH FROM THE MISSISSIPPI RIVER.

At the hinder end of the abdomen are five flattened appendages, which serve as fins, by means of which the animal can swim vigorously backward. (*See* Fig. 122.)

FIG. 122.—TAIL OF CRAWFISH SHOWING FLATTENED APPENDAGES FOR SWIMMING.

There are two compound eyes in front, which rest upon little jointed stalks, so that the creature can turn them in various directions.

121. Just below the eyes are two long, and two short and double antennæ, or feelers, and directly below these are six pairs of variously shaped and jointed appendages closely packed together. They surround the mouth and assist in securing and preparing the food for the stomach. The first pair are called jaws, or *mandibles*, and are furnished with sharp cutting edges for biting the food, and a flattened surface for grinding or crushing it. The next two pairs are called *maxillæ*, and are accessory jaws. The pair of mandibles and the two pairs of *maxillæ*, with another pair just behind, making four pairs in all, belong to the head, the other two pairs of mouth-parts belong to the thorax, and are so evidently modified claws or feet that they, with the pair just in front of them, are called foot-jaws or *maxillipedes*. (See Fig. 126, in which these parts are all named.)

122. From the under side of the body project five pairs of jointed legs, and these differ in shape and size. The first pair are much larger than the rest, and in the lobster are

FIG. 123.—A BIG CLAW OF THE LOBSTER, SHOWING THE WOODEN WEDGE, *a*.

called the *big claws*. They carry at their extremities big pincer-like jaws capable of giving a sharp nip, and these are used as weapons of defence, and also to hold on to their prey. The lobster can bite very severely with these big

claws, and for this reason the fisherman drives in a little wedge of wood to prevent the animal from opening the movable part, so that he can handle it without being bitten.

The other legs are long and slender. The two forward pairs end in slender nippers, while the two hinder pairs end in a single projecting claw. With these four smaller pairs of legs the crawfish and lobster crawl or walk.

On the under side of the abdomen are little flattened appendages arranged in pairs, a pair to each ring or segment. The animal not only swims backward by means of the broad

FIG. 124.—ONE OF THE FLATTENED OR ABDOMINAL APPENDAGES OF A LOBSTER.

fins on the end of the tail, or abdomen, but has the power besides to swim in a forward direction by extending the abdomen, and using the little fins below as swimming organs.

123. The crawfish, like the lobster, breathes in the water by means of gills. These are attached to the base of the legs and are concealed on the sides of the thorax by the carapace, which covers them. By forcibly tearing up the side of the carapace, there will be exposed the gills which

look like plumes. This space may be called the gill-chamber, and the water flows into it by passing under the edge of the carapace back of the big claws, and passes out of an opening near the mouth-parts. The currents of water flowing in to the gill-chamber are induced by a stiff appendage attached to the base of the second pair of maxillæ called the *flabellum* (see Fig. 125), and which swings back and forth and scoops the water into this chamber. These gills are shown as they appear in the 'crawfish. In tearing off the claws of the lobster, the gills are often drawn out too, and remain attached to the base of the legs. In the lobster the carapace can be easily bent up, so as to show the gills.

In the following figure a crawfish is shown with a portion

FIG. 125.—CRAWFISH SEEN FROM THE SIDE, WITH THAT PORTION OF THE CARAPACE REMOVED WHICH COVERS THE BRANCHIÆ OR GILLS. THE APPENDAGES OF THE LEFT SIDE ONLY SHOWN.—s, Region of Stomach; A, Abdominal Appendages; B, Bases of the Four Small Legs; C, Base of Large Claw; f, Flabellum attached to the Second Maxillipede; e, Eye.

of the carapace on the left side removed to show the gills as they appear in the gill-space. The big claw and the four smaller claws or legs are cut off, so that the other parts can be plainly shown.

124. If the pupils are skillful enough, it will be a good exercise for them to separate the various appendages of the crawfish, and arrange them upon a card, just as the beetle was arranged, as shown on page 66, fastening the carapace in the middle of the card with the back uppermost, then gluing the abdomen to the card, or, if possible, separating each ring of the abdomen and gluing each one separately to the card, one behind the other, and then arranging the appendages on each side of the thorax and abdomen, and with a pen marking the names of the various parts on the card.

Figure 126 represents the way in which the mouth-parts of a crawfish or lobster may be arranged.

As the lobster is a much larger animal than the crawfish, it will be easier to separate its appendages, and these may as well be taken from a specimen which has already been boiled, or as it may be obtained in the market. They may then be dried and fastened to a card with glue. The red color of the lobster appears only when the animal is boiled. When alive the color of the creature is a reddish-yellow mottled with bluish or greenish-black.

125. The eggs are carried by the crawfish and lobster glued in masses to the swimming appendages which are attached to the lower surface of the abdomen, and the creatures retain them in this way till the young hatch out.

This feature is characteristic of the class to which these animals belong. How different in this respect from the creatures already studied, in which the eggs are deposited and left by the animal ! It has been learned that certain spiders too carry their eggs round with them, and protect them.

FIG. 126.—MOUTH-PARTS OF A CRAWFISH FROM THE LEFT SIDE.

Fig. 128 represents a crab carrying its eggs glued to the appendages on the under surface of the abdomen. Pupils having access to lobsters in the markets, will, by looking over them, find some specimens in which the eggs are being carried in this way.

126. The young animal in growing sheds its entire shell,

in this respect again resembling the spider. This process is called *moulting*, a term used in describing a similar process in the spider. The lobster and crawfish continue to shed their shells at different periods, till they attain full growth. It is stated that the crawfish sheds its shell annually. If the pupils will keep some of these creatures alive, they will probably have an opportunity to observe this curious process of moulting, in which the entire outer skin, or shell, is discarded, so that there is left, complete in all its parts, the empty crust, like a discarded garment. The carapace separates from the abdomen above and cracks along the back, and by a series of efforts the animal pulls its way out. Great trouble is experienced in withdrawing the legs, and oftentimes a leg is left behind, and cases are recorded wherein the animal has perished in the struggles to liberate itself from the old skin. For some time the animal shows great timidity, and the lobster, when it has freshly shed its skin, retires to some secluded place, and there remains till the soft and tender skin has become thickened and hardened, so as to enable it to withstand the attack of its enemies. Lobsters often lose their legs in fighting, and on a sudden alarm are capable of dropping them off. The loss of the leg in this way is made good by the curious property the stump has of reproducing another leg, which grows out again, jointed and shaped like the one lost, only much smaller than the original one. At each succeeding moult, however; the leg becomes larger and larger. If the pupils will now carefully examine a lot of lobsters, they will notice among them some specimens in

which some one of the legs will be much smaller than its mate on the other side. This shows where a new leg has grown, to replace one previously lost.

127. The young passes through a remarkable series of moults, or shedding of the shell, and each moult brings it nearer in appearance to the general form of a lobster. The following figure presents the appearance of a young lobster which has undergone several such moults.

FIG. 127.—A YOUNG LOBSTER MAGNIFIED.—*a* shows the Natural Size of the Creature.
(Reduced from a Figure drawn by Sidney I. Smith.)

CHAPTER XVIII.

CRABS, HERMIT-CRABS, AND OTHER CRUSTACEANS.

128. THE class of animals to which the crawfish and lobster belong is called *Crustacea*, a name derived from a Latin word, *crusta*, meaning a crust, or the shell with which the animals of this class are covered.

To this class belong the crabs, hermit-crabs, shrimps, and an infinite variety of forms found in salt-water, certain little creatures found in the great lakes and other fresh

waters, as well as a little creature known as the sowbug, which is common under stones and boards in damp places. By far the larger proportion of these animals are found in salt-water.

Pupils having access to the sea-coast will find along the shore, and in pools of water left at low tide, a number of species unlike any thing found in fresh water.

The common crab may be studied and compared with the lobster. It will be found that the crab has the large claws, little legs, mouth-parts, antennæ, and other details similar to the lobster and crawfish. The body, however, is entirely unlike in shape; instead of being long and cylindrical, it is wide and flattened, and the long, jointed abdomen so characteristic of the lobster and crawfish is quite concealed in the crab. At first sight, the creature would appear to have no portion corresponding to this part in the lobster, but beneath the body there will be found a close-fitting piece composed of segments or joints, and, if this be raised or opened, the relation between this small piece and the large jointed abdomen of the lobster becomes at once apparent.

The crab carries her eggs attached in masses to the abdominal appendages which are arranged in pairs on the segments of the abdomen, as in the lobster.

In the following figure, which represents a crab carrying her eggs, a comparison of parts may be made between it and the crawfish or lobster.

The eggs as they are laid are covered with a sticky fluid, which thickens into threads and holds the eggs together and also holds them in masses to the abdominal appendages.

These appendages, having long hairs, retain the eggs all the more securely.

FIG. 128.—SIDE-VIEW OF COMMON CRAB, WITH THE ABDOMEN EXTENDED AND CARRYING A MASS OF EGGS BENEATH.—*e*, Eggs.

Under the microscope the eggs appear like bunches of berries or currants. The following figure represents a few eggs from a common crab:

FIG. 129.—*A*, a few Eggs from a Common Crab, enlarged; *B*, Single Egg greatly enlarged, showing more plainly the hardened Thread *t*, by which they are attached to each other. This Egg shows the young crab just beginning to form.

129. The small legs of a crab terminate in a single claw. There are no nipper or pincer-like ends as in the two forward

pairs of claws of the lobster, and in studying the crustaceans generally an infinite variety of modification will be found in these parts. In the crab which is so much sought after for food, and which is known as the soft-shell crab (a condition which indicates that the crab has just moulted, or shed its hard shell), the hinder pair of legs have the last or terminal joints flattened, and these flattened joints are used as fins by means of which the creature swims through the water.

FIG. 130.—RIGHT HIND-LEG OF THE EDIBLE CRAB.

The above figure shows the appearance of the right hind-leg of one of these crabs. Compare this with the common crab shown in Fig. 128.

130. A curious little crab, called the oyster-crab, makes its home within the shell of the oyster, living in the gill-cavity

FIG. 131.—OYSTER-CRAB. The Tail is hidden beneath the Body, one Segment only showing.

of the animal. Specimens may sometimes be found in canned oysters, and, to those who do not have access to the sea-shore, these creatures will furnish objects from which an idea of the crabs or short-tailed crustaceans may be

obtained. The tail will be found flattened against the under side of the body. Another species occurs in the salt-water mussel.

In the female crab, a figure of which is given (128), this part is very large and will be oftentimes found holding a mass of eggs. With care the creatures may be dried, and their various parts separated and stuck upon cards for the cabinet.

FIG. 132.—HERMIT-CRAB IN THE SHELL OF A SEA-SNAIL.

131. The hermit-crab possesses the general features of the common crab and lobster. The abdominal portion is long and cylindrical, and, instead of being encased in a hardened shell as in the lobster, it is soft and pliant, with scarcely a trace of hardened parts to indicate the segments. The creature, having this defenseless part, protects itself by securing the hard shell of some sea-snail as a house in which it constantly lives. The caudal appendages are curiously modified

to enable it to retain its hold on the shell, and the other abdominal appendages are rudimentary or wanting on the right side, or that side which comes most against the inside of the shell, as if they had been worn off. Wherever the creature goes, it drags the shell after it as a house.

FIG. 133.—HERMIT-CRAB REMOVED FROM ITS SHELL: *r*, Hardened Ridge which bears against the Inner Edge of the Aperture of the Shell; *a*, *a*, Appendages to which the Eggs are attached.

As the hermit-crab grows, it passes through the same features of moulting which characterize the crustaceans generally. The shell which protects its soft defenseless abdomen has, of course, no power of growth, and is abandoned when

the hermit-crab gets too big for it. The creature has, therefore, to go in search of another house slightly bigger than the one ready to be discarded. It is said that it does not always content itself with the dead shells that strew the beach, but has been seen attacking a live snail and eating it for the purpose of occupying its vacant shell, and this is rendered probable by the fact that they so often occupy fresh and perfect shells.

Not unfrequently they are found living in old and beach-worn shells which they have dragged about so long as to have worn the shell nearly through at the place where it rests and rubs against the sand. The pupils may collect hermit-crabs of all sizes on the shores.

The figure on the preceding page represents a hermit-crab after its removal from the shell. The creature is drawn as it appears lying on its back.

132. There are comparatively few species of crustaceans found in fresh water; and, with the exception of the species of crawfish and a few others, the fresh-water crustaceans are of small size.

FIG. 134.—FRESH-WATER CRUSTACEAN. The Line below represents the Natural Length of the Animal.

(Reduced from a Figure, in S. I. Smith's Report. of Fresh-Water Crustacea, published by U. S. Fish Commission.)

The figure 124 represents a small species which is common in stagnant pools in nearly all the Northern States.

From this species the pupils may study a form in which the segments of the thorax are not covered by a continuous shield.

The sowbug is a crustacean which lives out of water, though always requiring damp surroundings. It may be collected under logs and stones. In this creature the seven segments of the thorax are easily counted.

FIG. 135.—COMMON SOWBUG.—The line shows the length of the specimen from which this figure was made.

The eggs of the sowbug, as well as those of other species of crustaceans of the same group, are carried on the under side of the thorax and between the legs, in a little *brooding cavity* made by leaf-like parts which lap over each other and hold the eggs in place.

The eggs of these crustaceans may be found by examining the under side of the body, and observing a lightish-colored space between the legs. With a pin or the point of a knife-blade they may be scraped away without injuring the animal. They are very minute, and only under the

microscope can the development of the young creature be watched. The following figure represents a single egg of the sowbug highly magnified:

FIG. 136.—EGG OF SOWBUG, HIGHLY MAGNIFIED.—The little dot, at one side, represents the natural size of the egg. The head faces the left.

Around the upper edge of the *embryo* (as a young animal in the egg is called), from eighteen to twenty little blunt appendages may be seen; these represent the legs and other appendages of the body—the one longer than the rest is an antenna. As the creature grows, these appendages become jointed and variously modified to form the legs, mouth-parts, antennæ, and the appendages on the tail, which differ greatly from each other, though at the outset they are all alike.

133. Having studied a few of the many different kinds of crustaceans, let the pupils examine them together to find some points characteristic of them all.

Their bodies, in common with the insects, are composed of segments to which are attached jointed appendages of various kinds. This body is divided into two regions, the cephalo-thorax and the abdomen. In some the cephalo-thorax is covered by a continuous shield, called the carapace, as in the crawfish, crab, lobster, and shrimp. In others the segments of the cephalo-thorax are distinctly separate, and movable upon each other, as in the sowbug and certain other

crustaceans, one of which is figured on page 145. The cephalo-thorax is composed of fourteen segments, seven of these belonging to the head, judging from the number of appendages which arise from that part. The abdomen is supposed to possess seven segments, though the last one is so rudimentary that its existence as a true segment has been denied by some.

The deep line running across the back of the carapace, in the crawfish and lobster, is called the *cervical suture*.

The following outline represents a sowbug, with the regions of the body marked. Compare this with Fig. 125.

FIG. 137.—SOWBUG, WITH REGIONS OF BODY MARKED.

CHAPTER XIX.

BIVALVE CRUSTACEANS AND BARNACLES.

134. THE crustaceans during their growth shed their hard outer covering at intervals, and, in this as in many other respects, resemble the spiders. In the possession of gills, in-

stead of air-cavities, the crustaceans show a marked difference between themselves and spiders.

Now, there are hundreds of minute crustaceans in which it would be hard to recognize any close relations between them and the crustaceans already studied.

Among these odd forms may be mentioned certain little animals abundant in ditches and pools throughout the country. Some of these creatures are smaller than a pin's-head; others are as large as an ordinary white bean.

At first sight they might readily be mistaken for bivalve mollusks, as the body is covered with a bivalve shell, which partly opens and shuts, and is firmly attached to the body within. If the pupils are fortunate enough to collect some of these creatures in a net and watch them as they actively dart about in a jar of water, they will at once see the difference between them and the clam or mussel.

Instead of the animal's projecting a soft and fleshy foot with which to creep slowly along, as in the mussels, they will see numbers of little jointed, swimming legs partly protruded, and jointed antennæ thrust out in front; and, if their eyes are keen enough, may detect a little black speck just above the antennæ, which represents the eye.

The following figures represent a species collected in Dubuque, Iowa, and another form from Lynn, Massachusetts.

135. The concentric lines on the shell appear like lines of growth, and such they really are; but they are not made like the lines of growth on the mussel. When the creature moults, the delicate skin covering the antennæ and swim-

ming legs is discarded. The moulting process also takes place with the bivalve shell, but, instead of its being discarded, the moult is held or cemented to the new shell which forms underneath. Moult after moult of the shell is thus retained, the increasing size of each moult showing as sep-

FIG. 133.—FRESH-WATER CRUSTACEANS; *D*, Species from Dubuque, Iowa; *e*, Fye. The line below indicates the natural length of the specimen; *L*, Species from Lynn, Mass.: this is figured the natural size; *S* presents a highly-magnified section of one of the shells, to show the successive moults, these being numbered in their order of moulting; *B* shows the appearance of a portion of the edge of the shell along the back, representing the successive moults lapping one over the other.

arate concentric lines of growth. If the shell is cut into and the cut edge is examined with a microscope, the successive moults will be seen resting one upon the other, like the leaves of a book. By reading carefully the description accompanying Fig. 138, the pupils will learn more about it.

The very young stages of these creatures have some resemblance to the young of the barnacle shown in Fig. 141.

136. Another group of animals classed with the crustacea is still more unlike the forms already studied. These are

the barnacles. They are found in immense numbers covering the rocks and piers in places between high and low water mark. In nearly all places along the coast the rocks are whitened by their numbers. Pupils living inland can get specimens of the barnacle by visiting places where oysters are received in the shell; and, by examining the shells as they are thrown away, may now and then come across good specimens. With a stiff brush and some water the mud may be washed off the shell, and then the creature will present the following appearance.

FIG. 139.—BARNACLE ON OYSTER-SHELL.—The circular scar on the shell indicates the place from which another barnacle had been taken.

The barnacle-shell is round and conical, broad at the base by which it is firmly attached to the rock or whatever object it grows upon. The walls of this shell are composed of six pieces whose edges overlap each other. The opening in the top of the shell is closed by four plates which tightly fit together.

In collecting these creatures for the purpose of studying them alive, care must be taken in breaking them from the

rock. It will be much better to take them from the wooden piers, or, if possible, specimens should be collected attached to some pebble. These may now be placed in a jar or bowl of salt-water; and, if they are watched closely, there will be seen a set of fringed arms, like hairs, thrust out at the top of the shell, which, stretching out with graceful curves, close and partly retract within the opening. This motion will be constantly and rapidly repeated. There are six pairs of these appendages, and they are flung out in this way to grasp the minute particles in the water, which serve them as food. The arms being jointed and fringed with delicate hairs, the whole combined forms a sort of net. In Fig. 140, *C*, the tip of one of these arms is shown, highly magnified.

FIG. 140.—*A*, Side-View of Barnacle, Natural Size, showing Appendages protruded for Food; *B*, Top-View of same closed; *C*, highly-magnified View of the Tip of one of the Appendages.

137. Inland pupils may break open the barnacles collected on the oyster-shell, and, taking out the soft parts, may examine them by placing the parts in water, when the fringed arms become apparent; and under the microscope the delicate hairs

which fringe the arms may be seen. The jointed structure of these appendages and their arrangement in pairs show that the creature does not belong to the shell-fish or mollusks, as its shell might seem to indicate.

In past times many able naturalists classed these creatures with the mollusks, because they judged from the external appearances of the shell, which was limy. A careful study of their anatomy and development proved their relations to the crustaceans, and that they had no affinities whatever with the mollusks. In their growth they moult, in this act shedding all the skin, and at certain times in the summer the water will contain myriads of their cast-off skins. The shell, however, is not shed.

FIG. 141.—EARLY STAGES OF A BARNACLE: *A*, shortly after leaving the Egg; *e*, Eyes; *B*, having acquired a Bivalve Shell, and just before becoming attached, represented upside down; *C*, Appearance after becoming attached—Side-View; *D*, Top-View of still later Stage, with the Shell forming around it; *E*, Side-View of Later Stage, showing Appendages protruded. (The little marks at the sides of the figures indicate the natural size of the object. *A*, *B*, highly magnified; all of these Views are magnified, and, with the exception of *D*, are reduced from figures of C. Spence Bate.)

138. The young come from eggs as free-swimming animals, furnished with eyes and jointed appendages provided with hairs. In this condition they swim about for a while, and then acquire a bivalve shell, and in this state remotely resemble the little creatures shown in Fig. 138.

In Fig. 141, the letter *A* represents its first appearance from the egg. Its next appearance with the bivalve shell is shown at *B*. After remaining a free-swimming animal for a while it becomes attached to the rock, adhering by means of appendages on the head, and, then moulting, it loses its former appearance; the appendages change their proportions, new parts are added, the eyes disappear, a limy shell gradually forms around it, and it assumes characters entirely unlike those of its earlier stages.

CHAPTER XX.

WORMS.

139. AMONG the animals thus far studied, having a body composed of segments, the pupils have seen that in all cases the appendages were jointed, that is, the legs and antennæ were composed of distinct segments or joints; and, with the exception of the myriapods, or centipedes, the animals possessed a limited number of segments to the body.

In the group of animals now to be studied—the worms—the body has, generally speaking, an indefinite number of segments, and there are no jointed appendages attached to it.

The most accessible worm is the common earthworm. Specimens can be collected by digging for them in damp earth; and they may be found under almost any board or rock which has lain for some time.

The worms may be washed by placing them in a bowl of water, where their movements will soon remove the dirt.

The body is composed of a series of rings or segments, which are alike in form, except those at the extremities, which differ. The body tapers at both ends. The forward or anterior end tapers to a blunt point, while the hinder end becomes broad and flattened.

By carefully watching the worm when it shortens up after a long stretch, there will be seen, projecting from the sides and lower portion of each ring, minute points, which are the ends of little bristles protruding from openings in the sides. These bristles as they move project backward.

FIG. 142.—COMMON EARTHWORM.

The worm moves along the ground by the aid of these bristles, or *setæ*, as they are called. The body alternately lengthens and shortens. When the body lengthens, the *setæ*

on the hinder segments of the body prevent its stretching backward, because the setæ pointing backward stick into the ground; the body therefore can lengthen in only one direction, and that is in a forward one. Then, when the body shortens, the setæ in the forward segments stick into the ground, and the hinder part of the body is drawn up, and by this method the creature moves along.

FIG. 143.—EGG-CAPSULES OF LEECH, *A*; AND YOUNG LEECH, *B*.—The egg-capsules are drawn natural size. The young leech, which was taken from the egg-capsule, is drawn greatly enlarged, the little line above showing its natural size.

140. Another very common worm is the leech. Specimens may be collected in almost any pond or lake, and kept alive in jars of water. The creature is flat and broad, and in some species is furnished, at the hinder end of the body, with a sucker, while others have a sucker at each end of the body. It crawls by means of these suckers, and swims through the water by an undulating movement of the body. The eggs of the leech are laid on the leaves and stems of plants which grow in the water. The eggs are contained in little oval and flattened capsules, and these capsules are laid side by side. In Figure 143 *A* represents two capsules in which the little leeches can be seen; *B* represents the figure of a young leech greatly enlarged, showing

the eyes and mouth, at one end, and the sucker at the hinder end. The body being nearly transparent, the internal organs show through.

The species of worms in fresh water are few in number and quite small. The ocean seems to be their true home, and all along the sea-coast occur a great variety of worms — many of them of large size.

141. Pupils who have access to the sea-coast may collect them between high and low water mark. Certain species may be got by turning over stones and others by digging either in muddy or sandy places. After a violent storm from the ocean, many kinds of worms are thrown up, and may be found in pools left by the receding tide. The roots of large sea weeds also afford shelter to certain kinds. All of these creatures may be kept alive for a few days, though considerable care is required, and those not experienced in keeping salt-water aquaria are warned to exclude these animals.

They may be best studied by being placed in shallow bowls or plates, and there will be much to admire in their graceful motions and curious ways.

142. A very common form is found under stones at low tide. The body is composed of a great many segments, from the sides of which project little appendages of various shapes, and also bunches of bristles which can be plainly seen as the creature moves. The head, instead of being simple as in the earthworm, is surmounted by various feelers.

Another species very common on the sea-shore, under stones, is much shorter than the one just described. It has two rows of oval scales along the back, and the mouth is fur-

FIG. 144.—A COMMON SEA-WORM.

nished with powerful jaws, which work up and down. When they are placed in alcohol these jaws generally protrude.

143. Some species have a curious way of protruding their œsophagus when they seize their food, at the end of

FIG. 145.—A SEA-WORM WITH SCALES.—The eyes may be seen, four in number, between the forward scales.

which appear the jaws, like sharp claw-shaped teeth. The following figure shows the anterior end of one of these worms, with the œsophagus protruded, in the act of securing its prey.

Certain other sea-worms build tubes of mud or sand in which they live, and many of these have bunches of

thread-like feelers on the head. If these worms are taken from their tubes and placed in a plate of sea-water in which

FIG. 146.—ANTERIOR PORTION OF A SEA-WORM, WITH THE ŒSOPHAGUS PROTRUDED.—œ, Œsophagus; *j*, Jaws; *h*, Head,

are contained also particles of dirt or sand, they will commence to build a new tube, and for this purpose the threads on the head will stretch out like delicate rubber cords, and, becoming entangled in the particles of dirt, will draw them toward the head, when the appendages on the body will mould it around them in the shape of a tube.

FIG. 147.—SEA-WORMS WHICH MAKE LIMY TUBES.—*A*, Worm with Irregular Tube, the line *l* indicates natural length; *B*, Worm with Spiral Tube enlarged; *C*, Piece of Sea-Weed, showing the Appearance and Natural Size of these Spiral Tubes attached to it.

Other worms deposit a hard, shelly tube of lime. Some of these are irregular in shape, as in Fig. 147, *A*. Other

species build the tube in a coil, looking very much like a snail-shell, as in Fig. 147, *B*. This form is very common on the ordinary rock-weed, and may be collected in the *débris* thrown up by the waves. Pupils should collect these, and, if possible, watch the little creatures as they protrude the feathery appendages which surround the head. In the forms figured, one of the appendages is modified into a sort of plug, and, when the worm retires within the tube, the plug stops up the aperture securely, as certain snails close the aperture of their shells with an operculum.

144. In studying the affinities existing between animals in order to group them together naturally, the pupils should learn how unsafe it is to judge by the external appearances of the animals to be classified. For example, the little worm *B* just described has a coiled, limy shell, which might easily be mistaken for the shell of a snail. Yet the slightest examination of the soft parts within shows that the animal is made up of segments, and that, minute as it is, there are bunches of bristles, or setæ, projecting from the sides of the segments, and from these and other characteristics the creature is proved to be a true worm, having no affinities with the snails. The barnacles too have a limy shell; yet, when the creatures within are examined, their affinities with the crustaceans are seen at once; and, although distinguished naturalists in past times grouped them with the shell-fish, or mollusks, they properly belong to that class which includes the lobster and crab.

CHAPTER XXI.

CONCERNING NATURAL GROUPS.

146. THE pupils have seen, thus far, not only that the various creatures studied differ greatly in their structure, but that some are complex or elaborate in their characters, while others are quite simple. But, while these animals differ so much among themselves, there are certain characteristics which many of them have in common, as in the crustaceans and insects, for example, where all of them have the body divided into transverse segments, and the appendages are all jointed. These features, which are common in large assemblages of animals, are the essential characters by which they are brought together in great groups or divisions. Thus, all those animals which have the body jointed, that is to say, divided into a series of segments, as in the worms, crustaceans, and insects, form the great branch of *Articulates* of Cuvier, because Cuvier, the celebrated French naturalist, first applied the name *Articulata* to an assemblage of animals which included the worms, crustaceans, and insects. Since then naturalists have separated the worms from the crustaceans and insects, and have made a great branch of them called *Vermes*.

The worms differ from the other two classes with which

they were associated by Cuvier in not having jointed legs, and generally speaking in not having the segments grouped together into regions.

In worms, too, the segments are far more numerous, and there is no fixed or definite number of them as in the crustaceans and insects. The name *Articulates* is therefore abandoned, and the crustaceans and insects are united in one branch or sub-kingdom, and called *Arthropods*, a word derived from two Greek words, meaning *jointed foot*, while the worms are embraced under another sub-kingdom, *Vermes*. The clams, oysters, mussels, snails, and some other creatures which have not been mentioned in this book, have certain essential features in common, and so they are included in another great branch called *Mollusks*, from a Latin word *mollis*, meaning soft, because the bodies of these animals are soft. A very inexact name, because there are many other animals which are soft bodied having no relation with the shell-fish or Mollusca.

146. Now, these divisions or branches not only include animals which are simple in their structure, but animals which are very elaborate. All the animals in each great division, however, must embrace creatures that possess the same essential characters. With a knowledge of these essential features, it has been customary to make a diagram of a theoretical animal out of these characters only. This theoretical figure is called an *archetype*, meaning an ancient type, or first type, and the characters composing it are hence called *type-characters*, or *typical characters*, and that animal which

possesses most of these characters, in the plainest manner, is called a *typical animal*. This mode of presentation applies as well to smaller groups as to larger ones. Thus in the crustaceans, the lobster and crawfish might be called typical crustaceans, as being the types or representatives of the class, while a barnacle would certainly not be looked upon as a typical animal of this class, though belonging to it. In the same way an insect without wings would not be looked upon as a type of the insects, because one of the leading characters of the class of insects is the possession of wings.

In making systematic tables to show the relative grade an animal occupies, the simplest groups may be placed lowest in the list to indicate their inferior position. For instance, if the arthropods were to be arranged in a systematic table, those which have no lungs, but gills instead, would be placed lowest, because it has been found in other classes of animals that oftentimes the young or immature animal has gills which are afterward replaced by cavities for the purpose of breathing air direct, and the immature animal is regarded as less perfect, or lower in its organization than the mature or adult form. Consequently the crustaceans would be placed lowest in the scale. Then would come the air-breathing arthropods, and lowest among these would come the spiders, as the head is not specialized from the thorax. Next would come the myriapods, as in these the head is specialized as in the insects. And, finally, the true insects would come highest, as here the legs are reduced to three pairs, the head as well as the thorax is definitely sepa-

rated, and now the creature has added to it wings by means of which it has new means of locomotion. This, however, refers only to the most prominent types.

If the table is to be arranged to show the highest animals in the highest part of the column, it would be arranged as follows :

ARTHROPODS.	{	AIR-BREATHING ARTHROPODS.	{	<i>Insects.</i>
			{	<i>Myriapods.</i>
				<i>Spiders.</i>
	{	WATER-BREATHING ARTHROPODS.	{	<i>Crustaceans.</i>

This is *classification*: to classify animals is to bring those creatures together which have certain leading features in common. And these classes may be divided again and again into smaller groups.

As, for example, the insects proper: if the pupils were to divide them into smaller groups, the beetles would come together as one group, no matter what their shape or size; moths would form another group; the bugs another group, and so on.

CHAPTER XXII.

CHARACTERS OF VERTEBRATES.

147. THE lessons thus far presented have been upon creatures belonging to three great divisions of the animal kingdom, the Mollusks, Arthropods, and Worms. There are many other groups which represent other great divisions of the animal kingdom, of which no mention has yet been made,

and upon some of which a few brief lessons will be given. The material to be collected for this lesson should consist either of salamanders (or lizards, as they are incorrectly termed) or water-newts. The salamanders may be found in groves and forests, under rotten logs or bark. They are absolutely harmless, though many people regard them as poisonous. Water-newts are similar to the salamanders, except that they live in the water, and the tail is often provided with a fin.

The following figure (Fig. 148) represents a species of

FIG. 148.—COMMON YELLOW SPOTTED SALAMANDER.

salamander common to the Northern States, and also found at the South. Its color is bluish black, with a row of irregular shaped yellow spots on each side of the body.

In studying the external characters of the salamander, the pupils will observe the following features:

The animal has a head with a slight constriction between it and the body. The head has two eyes, capable of being closed by movable lids. The mouth opens transversely; that is, the lower jaw is on the under part of the head, and moves up and down. In the insects and crustacea, the jaws are on the sides of the head and open sideways. The jaws have minute teeth, and in large salamanders the teeth can be felt by rubbing the finger along the edge of the mouth. On the front of the head there are two holes representing the nostrils. The creature has four short legs, a forward pair and a hinder pair; it has short feet also, with four toes on each forward foot, and five toes on each hinder foot. The tail, which is quite as long as the body at its commencement, is nearly as wide as the body, but tapers gradually, and becomes flattened at its end.

148. How different in every respect is this creature from the animals thus far studied in this book! and yet, if the salamander is compared with a dog or cat, the characters mentioned above will be found in each. The cat has also a head containing two eyes with movable lids; the lower jaw is on the under part of the head and moves up and down and the mouth opens transversely; it is furnished with teeth, there are four legs with feet and toes, and the creature has a tail. But there are also important differences between the two: the cat has external ears, while the salamander has none, though it has parts which enable it to hear. The cat's skin is covered with hair and is dry, while the salamander has no hair upon it and the skin is always moist. The cat has sharp claws,

while the salamander now being described has none. On touching the cat it feels warm, while the salamander feels cold to the touch; with the cat the young are born alive, and the little kittens have the same general features as the parent-cat. The salamander lays a lot of eggs, and most of the species lay their eggs in the water; when these hatch, the creatures coming from them have no lungs and cannot breathe air directly, but have gills instead. More curious still is the fact that, when the young creature hatches from the egg, it has no legs, these appearing afterward as the animal grows.

The following figure represents the appearance of a young salamander ten days after hatching from the egg. The gills, appearing like feathers, are seen on the sides of the neck.

FIG. 149.—YOUNG SALAMANDER.—*g, g*, Gills.—(After a figure by Dr. P. R. Hoy.)

149. Thus, while there are important differences between the cat and the salamander, there are also many points of resemblance; and, if the arrangement of bones constituting the skeleton be examined, a still closer resemblance may be seen. Within the body the salamander has a series of bones which together form the *skeleton*. The most important part of the skeleton consists of a row of bones which runs along the central line of the back of the body from the head to the tip of

the tail. This row, or column of bones, is called the *vertebral column*, and the bones composing it are called the *vertebræ*. The bones of the head combined form the skull or *cranium*. The ribs, which in the salamander are rudimentary, are attached to the sides of the body *vertebræ*. The bones of the fore and hind legs are similar, though they are called by different names.

A series of bones just back of the head, the longest of which is the shoulder-blade, forms the *pelvic girdle*; to this girdle the first bone of the fore-leg is joined. Other bones at the hinder part of the body form the *pelvic arch*, and to this the first bone of the hind-leg is joined.

150. It will be a difficult task for the pupils to remove the flesh from a salamander so as to show the bones united, and it will also be difficult to prepare the skeleton of a cat; but the pupils may learn something about the bones and their attachments by gently handling the creature. Strauss-Durkheim, a celebrated naturalist, when he was writing his famous work on the cat, used to hold one of these animals for hours in his lap, while he felt of the muscles and other portions of the body.

The following figure of a cat shows the position of the bones in outline.

Along the back is a series of prominences which indicates the vertebral column, or, as it is usually called, the backbone or spine. On the sides of the body the ribs may be felt like bars or ridges. The shoulder-blades, or *scapulae*, are prominent bones, forming the shoulders, and from these the bones of the fore-legs start.

The bones of the pelvic arch, or *pelvis*, may be readily detected, and from these the hind-legs start.

FIG. 150.—CAT, WITH BONES OF RIGHT SIDE DRAWN.—*Cra*, Cranium; *sc*, Scapula or Shoulder-Blade; 1, Humerus; 2, Radius and Ulna; 3, Carpus; 4, Phalanges; 5, Femur; 6, Tibia and Fibula; 7, Tarsus; 8, Metatarsus; 9, Phalanges; *i*, Innominate Bone—a number of Bones combined, forming the Pelvic Arch; *v*, Vertebral Column.

151. This mode of examining the bones is suggested, since it would hardly be possible for pupils to prepare a proper skeleton, and because few of the museums of the country to which they would have access possess skeletons of this kind. Now and then there may be found upon the beach a nicely-cleaned skeleton of a dog or cat, made so by little creatures which have fed upon the flesh, and this may be used for study.

As unlike as the salamander and cat are in some respects, in many characters, both external and internal, they are remarkably alike.

If one of the body vertebræ be examined, there will be found a central bony mass, in the upper side of which will be found a hole which is made by the bone growing up from

each side of the bony mass, arching over, and uniting above. All of the vertebræ, except those in the tail, possess this channel or tube.

Within the skull is a mass called the brain, and running from this through a hole in the skull is a long white cord called the spinal cord or *cerebro-spinal cord*, and this always runs along on the back of the spine or vertebral column, passing through the arch or hole of each vertebra just described. In fact, the spinal cord is protected from injury by passing through this bony tube.

These features with various modifications will be found in all animals having a vertebral column, that is, a central longitudinal axis either of bone or cartilage above which runs the spinal cord.

The impulses of the animal to move originate in the brain and, passing along the spinal cord, run off by means of nerves, to animate the movements of the muscles. As a proof of this, if the spinal cord be injured, the parts behind and below it are rendered helpless.

The ribs passing from the vertebræ arch below, and form another and much larger cavity, in which are contained the organs which contribute to the body's growth, such as the lungs for breathing, the heart for propelling the blood, the stomach for digesting the food, and so on. The following figure represents a body vertebra to which are attached a pair of ribs with the cavity above, in which is contained the spinal cord or cerebro-spinal cord, and the cavity below, in which are contained the lungs, heart, stomach, etc.

152. If the legs of the salamander are now examined, the following characteristics will be noticed :

FIG. 151.—A VERTEBRA AND A PAIR OF RIBS, FROM A CAT.—*C S*, Cavity for Cerebro-spinal Cord.

If we compare the hind-leg of the salamander with our own leg, we shall find essentially the same arrangement of parts, namely: the leg bending at a joint in its middle, or the knee-joint, and below this joint another one called the ankle-joint. In that portion of the leg above the knee-joint there is one long bone called the *femur* which joins or articulates with the pelvic arch. In the leg below the knee-joint there are two long bones side by side, called the *tibia* and *fibula*. In the ankle-joint there are a number of small bones closely packed together; these are called the *tarsal bones*, and together form the *tarsus*. In the body of the foot there are several bones upon which the toes rest, and

these are called the *metatarsal* bones, and together form the *metatarsus*; and finally come the bones of the toes called *phalanges*—a long series of names to remember, yet they apply to every animal possessing a vertebral column and having legs.

153. The fore-legs of the salamander have similar joints, the elbow-joint corresponding with the knee-joint, the wrist-joint corresponding with the ankle-joint. Above the elbow there is one long bone called the *humerus*, and this articulates with the shoulder-blade, or *scapula*, as it is called. Below the elbow are two long bones side by side, called the *radius* and *ulna*. In the wrist are a number of small bones called *carpal* bones, which together form the *carpus*; and then follow longer bones corresponding with the metatarsal bones, and these are called the *metacarpal* bones, and together form the *metacarpus*; and finally the bones of the fingers which are called phalanges, the same name which applies to the bones of the toes. The following figures represent the bones of the right fore and hind leg of a species of salamander common to the Northeastern and Middle States.

With the aid of a good hand-lens the pupils may see these bones in the leg of any small salamander by observing the following directions: having secured a live salamander, the animal may be killed with ether; now, if the leg be cut off and gently pressed between two thin pieces of glass, the flesh is sufficiently translucent to show all the principal bones quite distinctly:

FIG. 152.—HIND AND FORE LEG OF A COMMON WOOD SALAMANDER. (These figures are greatly enlarged.)

154. All animals possessing a vertebral column have essentially the same external features as have been given in the preceding lessons on the salamander and the cat; that is, they have a head containing two eyes with movable lids, a mouth opening transversely and generally furnished with teeth; and, with the exception of the snakes and a few other creatures, possessing two pairs of legs—a fore-pair and a

hind-pair—and internally having a skeleton embracing the features already noticed.

In the fishes the head is continuous with the body, the fore and hind legs are represented by fins, one pair directly back of the head, corresponding with the front-legs, and another pair of fins, which represent the hind-legs, variously situated behind, below, or even in front of the first pair.

155. The animals which possess a vertebral column, and hence called *Vertebrates* (from the Latin word *verto*, “I turn”), are the fishes, frogs, toads, salamanders, snakes, lizards, turtles, crocodiles, birds, and the warm-blooded, four-legged beasts, such as the cat, dog, sheep, horse, and elephant. These last-mentioned creatures belong to a large class of animals called *Mammalia*, the leading features of which are, that the young are born alive, and the mother nurses the young.

The fishes, frogs, toads, salamanders, snakes, lizards, turtles, and crocodiles, and others like them, are cold-blooded, while the birds and mammals are warm-blooded, and all of them, except the mammals, lay eggs from which their young hatch. The fishes, snakes, lizards, and other reptiles are, generally speaking, covered with scales. The frogs, toads, and salamanders, are smooth-skinned; the birds are covered with feathers, while the mammals, with few exceptions, are clothed with fur. In the general grouping of the vertebrates the fishes and amphibians—namely, the toads, frogs, and salamanders—form one group, the reptiles and birds form another group, and the mammals a third group.

With the exception of the fishes, the similarity in the character and arrangement of the bones of the skeleton of every vertebrate is remarkable. Even the birds, which are apparently so different from the mammalia on the one hand and the turtle and salamander on the other, are yet quite similar to each in the general character and arrangement of their bones; and if the young bird, while yet in the egg, is examined, the presence and affinities of certain bones are very clearly seen.

CHAPTER XXIII.

BONES OF THE LEG AND WING OF BIRDS.

156. OUR pupils have learned by this time how important it is to study the very young animal in order to determine its relationships. Thus in studying the young barnacle the affinities of the creature were more readily recognized, and in the affinities of the parts of the animal, by studying the young spider the palpi were more easily seen to be modified legs. Now, as an interesting example of the necessity of studying the young or early condition of an animal, a bird is cited.

The fore-leg in the bird is represented by its wing, and in studying the bones of the wing of an adult bird but little resemblance can be seen between them and similar parts in other vertebrates. The humerus, radius, and ulna are plain enough, it is true, and, when the pupil has a chance to pick

the flesh from a chicken's or a turkey's wing, he may observe these bones easily enough. The bones of the wrist and hand, however, seem to be few in number and curiously grown together.

FIG. 153.—RIGHT WING OF AN ADULT BIRD SEEN FROM THE INSIDE.—*H*, Humerus; *R*, Radius; *U*, Ulna.

157. If, however, a young bird is taken from the egg before the parts are fully formed, the bones of the wing will be found separate and distinct, and the relation between the wing of the bird and the fore-leg of other vertebrates becomes more fully apparent.

So constant are the characters of these parts in all birds, that a robin, a sparrow, a pigeon, or a chicken, will reveal the parts quite as distinctly as the larger birds.

In the embryo bird, that is, a bird while still in the egg, the wing and leg appear far more alike than in the adult, as may be seen by looking at the following figures of embryo birds :

FIG. 154.—EMBRYO BIRDS IN VARIOUS STAGES OF DEVELOPMENT.—A, Chipping Sparrow; B, Petrel; C, Tern seen from above.

If, now, the bones of the wing of an embryo bird be examined, the bones of the extremity of the wing, instead of

FIG. 155.—RIGHT WING OF AN EMBRYO BIRD GREATLY ENLARGED.—Only a portion of the humerus is seen.

being closely welded together as in Fig. 153, are found to be separate, and the carpal bones, of which only two were apparent in the full-grown wing, are now separate and four in number. The wing at this stage looks more like a three-toed foot. Fig. 155 shows the appearance of the wing of an embryo bird.

As the bird develops, the bones of the fingers gradually approach and some of the bones grow together till they present the appearance shown in Fig. 153.

158. The bones of the leg grow together in the same way.

FIG. 156.—BONES OF THE RIGHT LEG OF AN ADULT BIRD.—Only the lower portion of the tibia and fibula is drawn.

The foregoing figure (*see* page 177) represents the lower portion of the leg-bones of a bird.

It will be seen that three of the metatarsal bones, corresponding to the three long toes, are combined; their ends, where the toes join on, appearing separate, while the metatarsal bone of the short toe on the side remains separate from the others.

At the ankle-joint, or tarsus, there are no separate tarsal bones to be seen, nor would their existence be known, except theoretically, without an examination of the embryo.

The following figure represents the appearance of the leg

FIG. 157.—RIGHT LEG OF AN EMBRYO BIRD GREATLY ENLARGED.—Only the lower portion of the tibia and fibula is drawn.

of an embryo bird greatly enlarged, showing the bones in place.

At this early stage, not only the metatarsal bones are seen separate, but the bones of the tarsus, four in number, are well marked and distinct.¹

The many strange modifications in the shape and proportions of the legs of vertebrate animals are accompanied by similar changes in the arrangements, number, and proportion of the bones of these parts.

Thus, in the short leg of the salamander (*see* Fig. 152) the bones of the tarsus occupy a space about as long as that of the metatarsus. In birds, on the contrary, the tarsus is very short, while the metatarsus is very long.

159. If the pupils will observe the hind-leg of the toad or frog, they will see that not only is it much longer than the fore-leg, but that an extra joint seems to be added in the foot. An examination of the bones shows that the first two tarsal bones are very long, while the other tarsal bones are very short; so that in this case the tarsus combined is much longer than the metatarsus. The following figure represents the right hind-leg of a young toad while still in the tadpole state, though the adult shows the same features.

The above brief lessons on the vertebrates do not even embrace an outline of the structure and habits of any of the classes, and they are given only as suggestions toward a method of study, and to point out the essential paths to fol-

¹ The author has seen the tarsal bone, marked *i*, in the heron, tern, and sea-pigeon.

low, in order to gain an insight into the affinities existing between the various groups composing this great branch of animals.

FIG. 158.—BONES OF THE RIGHT LEG OF A YOUNG TOAD GREATLY ENLARGED.—The femur is not shown in this drawing; the tibia and fibula are combined together.

There are still other large classes of animals forming branches or sub-kingdoms quite distinct from those already studied, and represented by animals which live in the sea, such as the star-fish, sea-urchin, jelly-fish, and sea-anemone, of which no mention will be made in this book. There are also many classes, belonging to branches already touched upon, which have not been alluded to. These will be fully dealt with in the second book, now in preparation.

CHAPTER XXIV.

CLASSES AND SUB-KINGDOMS.

THE following figures illustrate the classes and sub-kingsdoms which have been dealt with in this book, with their technical names :

SUB-KINGDOM MOLLUSCA (*Clams, Snails, Squids, etc.*).

Class Gasteropoda (snails, periwinkles, limpets, etc.).—Animals whose bodies are generally inclosed in a mantle or sac, which usually secretes a shell composed of one piece, and this often assuming a spiral shape. The body rests upon a broad, creeping disk called the *foot*, and this part represents the ventral portion or belly of the animal. The name *Gasteropoda* is derived from two Greek words, *gaster*, the belly, and *pous*, foot. The following figures represent a few animals of this class :

GASTEROPODS.

Class Acephala (clams, oysters, mussels, etc.).—Animals whose bodies are protected by a mantle which secretes a bi-valve shell, or a shell composed of two pieces. They have no well-defined head, and hence the name *Acephala*, derived from two Greek words, *a*, without, and *cephale*, head. These animals are also called *Lamellibranchiates*, because the gills form leaf-like membranes or plates on the sides of the body ; the word being derived from a Latin and a Greek word, *lamella*, a plate, and *branchia*, gill. The following figures represent a few animals of this class :

ACEPHALA.

SUB-KINGDOM ARTHROPODA (*Insects, Spiders, Centipedes, Crabs, etc.*).

Class Insecta (beetles, bugs, butterflies, etc.).—Animals whose bodies are made up of segments grouped together in three regions, the head, thorax, and abdomen ; having three pairs of jointed legs, and one or two pairs of wings, and breathing air through openings in the sides of the body. The word *insecta* comes from a Latin word, *inseco*, I cut into, referring to the distinct separation of the body into regions.

INSECTS.

Class Myriapoda (centipedes, millepedes). Animals composed of many segments. These not apparently combined into regions, except the head, which is distinct. The number of pairs of legs coinciding with the number of segments. Breathing air through openings in the sides of the body.

The word *myriapoda* is derived from two Greek words, *myrioi*, ten thousand, and *pous*, foot.

A MYRIAPOD.

Class Arachnida (spiders). Animals whose bodies are segmented. The segments grouped together into two regions, the cephalo-thorax and the abdomen. Having four pairs of legs, and breathing air through openings in the body.

The word *arachnida* is derived from a Greek word, *arachne*, spider.

ARACHNIDE.

Class Crustacea (lobsters, crabs, barnacles, etc.). Difficult to define, but including animals which pass through a series

of moults in their growth, though in this respect resembling the spiders, and breathing by means of gills, and in this respect differing from other *arthropods*.

CRUSTACEANS.

SUB-KINGDOM VERMES (worms). *Class Annelida* (angle-worms, leeches, certain sea-worms, etc.). Animals whose bodies are made up of an indefinite number of segments, bearing appendages which are not jointed, and in the larger number of groups having bunches of bristles or *setæ* upon the sides of the body which act as supplementary organs of locomotion. The name *annelida* is derived from the Latin word *annulus*, a ring.

WORMS.

SUB-KINGDOM VERTEBRATA, with the following classes, which have been only briefly alluded to : *Fishes*, *Amphibians*, *Reptiles*, *Birds*, *Mammals*. According to a late classification of Professor Huxley's, these classes would stand *Ichthyopsida*, which includes the *Fishes* and *Amphibians*; *Sau-*

ropsida, which includes the Reptiles, and Birds; and lastly *Mammalia*.

Ichthyopsida is derived from two Greek words, *ichthus*, a fish, and *opsis*, appearance. *Sauropsida* comes from two Greek words, *sauros*, a lizard, and *opsis*, appearance.

VERTEBRATES.

NOTICE TO TEACHERS.

To those who care to pursue the subject more in detail with their classes, or to present the history of those groups of animals of which no mention has been made in this book, the author would suggest the following publications, among many others of value, as works of reference :

Woodward's Manual of Mollusca.

Marine Mammalia and American Whale-Fishery. By Captain C. M. Scammon,
U. S. R. M.

Sea-side Studies in Natural History. By Mrs. Agassiz and Alexander Agassiz.

Corals and Coral Islands. By Prof. J. D. Dana.

Packard's Guide to the Study of Insects.

Insects injurious to Vegetation. Harris.

The Annual Reports of the State of Missouri on the Noxious, Beneficial, and
other Insects. By Prof. C. V. Riley.

United States Fish Commissioners' Report for 1871-'72. By Prof. Spencer F.
Baird. Containing valuable illustrated chapters on the Mollusca, Crustacea,
Worms, etc., by Prof. A. E. Verrill and S. I. Smith.

Key to North American Birds. By Dr. Elliot Coues, U. S. A.

Osteology of Mammalia. Flower.

A History of North American Birds. By Baird, Brewer, and Ridgeway.

Canadian Entomologist.

Forms of Animal Life. Rolleston.

Methods of Study in Natural History. By Prof. L. Agassiz.

And for general information, the works of Darwin, Huxley, Owen, Wallace, Mivart, Lubbock, and the Duke of Argyle, and also

Nature, a weekly magazine published in London, and containing an infinite variety of contributions from English and Continental naturalists.

The American Naturalist, a popular illustrated magazine of natural history. Edited by Dr. A. S. Packard and F. W. Putnam, and published in Salem, Massachusetts; containing a vast amount of information by Lockwood, Allen, Coues, Verrill, Smith, Stearns, Scudder, Emerton, Gill, Putnam, Packard, Hyatt, Mann, Marsh, Dall, Cooper, Gill, Cope, Ridgeway, Wood, Abbott, Trippe, Le Conte, Wyman, Dawson, Grote, Mayer, Gentry, Shaler, Wilder, Aikin, Treat, Perkins, Riley, Agassiz, Dana, Hill, Uhler, Edwards, Tuttle, Tenney, Ward, Hagen, Hartt, Shimmer, Hartshorne, Ritchie, Tilsdale, Hoy, Orton, Lewis, Leidy, Brigham, Scammon, Binney, Stimpson, Collins, Fowler, Walker, Jordan, Wright, Norton, Maynard, Canfield, Fellowes, Endicott, and others.

The Popular Science Monthly, edited by Prof. E. L. Youmans, and published by D. Appleton & Co., New York; containing valuable illustrated articles by American and European naturalists.

Every school library should, if possible, contain a complete set of *Nature*, the *Naturalist*, and *The Popular Science Monthly*. For special descriptions of species, the miscellaneous collection of the Smithsonian Institution, with contributions by Stimpson, Gill, Bland, Binney, Prime, Tryon, and others. Also the *American Journal of Science and Art*, and the publications of the Boston Society of Natural History, Philadelphia Academy of Sciences, New York Lyceum of Natural History, Buffalo Academy of Natural Sciences, California Academy of Natural Sciences, Portland Society of Natural History, Museum of Comparative Zoölogy, Peabody Academy of Science, etc.

All these last-named publications should be found in the larger libraries of the country.

THE INTERNATIONAL SCIENTIFIC SERIES.

The INTERNATIONAL SCIENTIFIC SERIES is a series of Monographs on Selected Scientific Topics, contributed by some of the leading thinkers of different countries. It gives distinctive prominence to those branches of Biological, Psychological, and Social Science which are undergoing rapid development in the present age, and which are ever more and more involved with the highest human interests.

The books, though thoroughly scientific, are also designed for the use of the non-scientific reader. They are explanatory in character, and as free from technicality as is compatible with thorough accuracy.

NOW READY.

I. The FORMS of WATER in CLOUDS and RIVERS, ICE and GLACIERS. By J. TYNDALL, LL.D., F.R.S. Fifth Edition. With Twenty-six Illustrations. 5s.

'In works of this kind almost everything depends upon the writer's powers of exposition, and those of Professor Tyndall are almost unsurpassed. Felicitous illustration, pertinent anecdote, and a crystal style make his expositions read like a romance, while, we scarcely need add, he speaks with a scientific knowledge that few are competent to question. We simply report, therefore, a fascinating book dealing largely with the phenomena of glaciers and snow, with which, more than any living man, Professor Tyndall has made us familiar.'

BRITISH QUARTERLY REVIEW.

II. PHYSICS and POLITICS; or, Thoughts on the Application of the Principles of 'Natural Selection' and 'Inheritance' to Political Society. By WALTER BAGEHOT. Third Edition. 4s.

'Mr. Bagehot writes in a graceful style, and has much to say upon political topics that is well worth attention. We can recommend the book as well deserving to be read by thoughtful students of politics.'—SATURDAY REVIEW.

'A work of really original and interesting speculation.'—GUARDIAN.

III. FOODS. By EDWARD SMITH, M.D., LL.B., F.R.S. Third Edition, Profusely Illustrated. 5s.

'Dr. Smith discusses, with much minuteness, the economical and sanitary value of the various kinds of animal and vegetable food which are commonly in use, and of drinks, both alcoholic and other.'—SPECTATOR.

IV. MIND AND BODY: the Theories of their Relation. By ALEXANDER BAIN, LL.D. Fourth Edition. With Four Illustrations: 4s.

'The work before us seeks to complete the doctrine of the relation between consciousness and bodily organism by noting and classifying the normal instances. It proposes to show not how curiously disease or any sudden change in bodily function affects the brain, and through the brain the mind, but how completely all the familiar processes of sensation, thought, and emotion flow, so to speak, upon the surface of nervous currents, which sustain and fashion their ever-varying shapes.'—SATURDAY REVIEW.

V. The STUDY of SOCIOLOGY. By HERBERT SPENCER. Fifth Edition. 5s.

'There is no lack of interest even to those who are not inclined to concern themselves with the projected social science.'—ACADEMY.

'It contains, as any writing of Mr. Spencer's needs must, a great amount of interesting and suggestive matter.'—SATURDAY REVIEW.

VI. On the CONSERVATION of ENERGY. By BALFOUR STEWART, M.D., LL.D., F.R.S. Third Edition. With Fourteen Engravings. 5s.

'Thorough and simple. . . A boon to science and the world at large. . . Even the uneducated can obtain an accurate knowledge, provided they are taught by as sympathetic and suggestive a teacher as Professor Stewart.'—SATURDAY REVIEW.

VII. ANIMAL LOCOMOTION; or, Walking, Swimming, and Flying. By J. B. PETTIGREW, M.D., F.R.S. Second Edition. With 119 Illustrations. 5s.

'It is a clear and comprehensive *résumé* of the present advanced state of our knowledge of animal locomotion, as shown by the most recent successful experiments and discoveries.'

STANDARD.

VIII. RESPONSIBILITY in MENTAL DISEASE. By HENRY MAUDSLEY, M.D. Second Edition. 5s.

'This volume contains in a clear and popular form the results of all the most recent investigations into the nature of insanity. The volume is altogether one of the best of the International Scientific Series which has yet appeared, and will add to its author's high literary reputation.'—ACADEMY.

IX. The NEW CHEMISTRY. By Professor JOSIAH P. COOKE, of the Harvard University. Second Edition. With Thirty-one Illustrations. 5s.

'Many portions of the book are eminently suggestive to the student.'—ACADEMY.

'The science it contains is popular science in the best sense of the term. The great ideas of modern chemistry are presented with singular clearness and with very varied illustration.'—LANCET.

X. The SCIENCE of LAW. By Professor SHELDON AMOS. Second Edition. 5s.

'Mr. Sheldon Amos has written a work on the Science of Law in a style adapted to the ordinary reader who wishes to become familiar with the landmarks of jurisprudence. . . . Much would remain unsaid though we discussed the author's very interesting volume at far greater length; but this we can say, that it does "open out to novices an unsuspected region of interest," and that we hope with Mr. Amos it will "whet their curiosity and stimulate them to further research."'—LAW TIMES.

XI. ANIMAL MECHANISM: a Treatise on Terrestrial and Aerial Locomotion. By Professor E. J. MAREY. Second Edition. With 117 Illustrations. 5s.

'It is not only to a passage here and a passage there that we have had to refer, but to the thorough exposition of intricate problems of mechanical physiology, which have been worked out with a degree of ability rarely to be found in a single author.'—NATURE.

XII. The DOCTRINE of DESCENT and DARWINISM. By Professor OSCAR SCHMIDT (Strasburg University). Second Edition. With Twenty-six Illustrations. 5s.

'The book is one of very conspicuous ability, and will be read with profit even by those who may take occasion to dissent from its special conclusions.'—SCOTSMAN.

XIII. The HISTORY of the CONFLICT between RELIGION and SCIENCE. By Professor J. W. DRAPER. Fifth Edition. 5s.

'Certainly it is long since a book with such an important bearing on society, and so thoroughly scientific, and therefore impartial, was placed in the hands of the British public as that of Professor Draper's "Conflict between Religion and Science." And assuredly very few men could have been found so thoroughly able to criticise the scientific side, and yet to enter freely upon the literary, as the learned Professor of Physiology in the University of New York.'—POPULAR SCIENCE REVIEW.

XIV. FUNGI: their Nature, Influences, Uses, &c. By M. C. COOKE, M.A. LL.D. Edited by the Rev. M. J. BERKELEY, M.A., F.L.S. Second Edition. Crown 8vo. With Illustrations. 5s.

XV. The CHEMICAL EFFECTS of LIGHT and PHOTOGRAPHY, in their APPLICATION to ART, SCIENCE, and INDUSTRY. By Dr. HERMANN VOGEL (Polytechnic Academy of Berlin). Third Edition. The Translation thoroughly revised. With 100 Illustrations, including some beautiful Specimens of Photography. 5s.

XVI. The LIFE and GROWTH of LANGUAGE. By W. D. WHITNEY, Professor of Sanskrit and Comparative Philology in Yale College, New Haven. Second Edition. 5s.

XVII. MONEY and the MECHANISM of EXCHANGE. By W. STANLEY JEVONS, Professor of Logic and Political Economy in the Owen's College, Manchester. Second Edition. 5s.

XVIII. The NATURE of LIGHT; with a General Account of PHYSICAL OPTICS. By Dr. EUGENE LOMMEL, Professor of Physics in the University of Erlangen. Second Edition. With 188 Illustrations and a Plate of Spectra in Chromolithography. 5s.

October, 1875.

AN ALPHABETICAL LIST
OF
HENRY S. KING & CO.'S
PUBLICATIONS.

*65 Cornhill, and 12 Paternoster Row, London,
October, 1875.*

A LIST OF
HENRY S. KING & CO.'S
PUBLICATIONS.

ABBEY (Henry).

BALLADS OF GOOD DEEDS. Fcap. 8vo. Cloth gilt. 5s.

ADAMS (A. L.), M.A.

**FIELD AND FOREST RAMBLES OF A NATURALIST IN
NEW BRUNSWICK.** With Notes and Observations on the
Natural History of Eastern Canada. 8vo., cloth. Illustrated. 14s.

ADAMS (F. O.), H.B.M.'s Secretary of Embassy at
Paris, formerly H.B.M.'s Chargé d'Affaires, and Secretary of
Legation at Yedo.

THE HISTORY OF JAPAN. From the Earliest Period to the
Present Time. New Edition, revised. Demy 8vo. In 2 vols.
With Maps and Plans. 21s. each.

ADAMS (W. Davenport, Jun.)

LYRICS OF LOVE, from Shakespeare to Tennyson. Selected
and arranged by. Fcap. 8vo., cloth extra, gilt edges, 3s. 6d.

ADON.

THROUGH STORM AND SUNSHINE. Illustrated by M. E.
Edwards, A. T. H. Paterson, and the Author. 1 vol. Crown 8vo.
Price 7s. 6d.

A. K. H. B.

A SCOTCH COMMUNION SUNDAY, to which are added Cer-
tain Discourses from a University City. By the Author of
"The Recreations of a Country Parson." Second Edition.
Crown 8vo. 5s.

ALLEN (Rev. R.), M.A.

ABRAHAM: HIS LIFE, TIMES, AND TRAVELS, as told by a Contemporary 3800 years ago. Post 8vo., with Map. Cloth. Price 10s. 6d.

AMOS (Sheldon).

THE SCIENCE OF LAW. Crown 8vo. 5s.
Being Vol. X. of the International Scientific Series.

ANDERSON (Rev. Charles), M.A.

CHURCH THOUGHT AND CHURCH WORK. Edited by. Containing articles by the Revs. J. M. Capes, Professor Cheetham, J. Ll. Davis, Harry Jones, Brooke, Lambert, A. J. Ross, the Editor, and others. Second Edition. Demy 8vo. 7s. 6d.

WORDS AND WORKS IN A LONDON PARISH. Edited by. Second Edition. Demy 8vo. 6s.

THE CURATE OF SHYRE. Second Edition. Demy 8vo. Cloth. 7s. 6d.

ANDERSON (Colonel R. P.)

VICTORIES AND DEFEATS. An Attempt to explain the Causes which have led to them. An Officer's Manual. Demy 8vo. 14s.

ANSON (Lieut.-Col. The Hon. A.), V.C., M.P.

THE ABOLITION OF PURCHASE AND THE ARMY REGULATION BILL OF 1871. Crown 8vo. 1s.

ARMY RESERVES AND MILITIA REFORMS. Crown 8vo. 1s.

THE STORY OF THE SUPERSESSIONS. Crown 8vo. 6d.

ARMY OF THE NORTH GERMAN CONFEDERATION. A Brief Description of its Organization, of the Different Branches of the Service and their rôle in War, of its Mode of Fighting, etc., etc. Translated from the Corrected Edition, by permission of the author, by Colonel Edward Newdegate. Demy 8vo. 5s.

ARBUTHNOT (Sir Alexander).

SIR THOMAS MUNRO, BART., K.C.B., sometime Governor of Madras. A Selection from his Minutes and other Official Writings.

ARGYLE (Duke of).

SPEECHES ON THE SECOND READING OF THE CHURCH PATRONAGE (SCOTLAND) BILL IN THE HOUSE OF LORDS, June 2, 1874; and Earl of Camperdown's Amendment, June 9, 1874, placing the Election of Ministers in the hands of Rate-payers. Crown 8vo. Sewed. 1s.

ASHANTEE WAR (The).

A Popular Narrative. By the Special Correspondent of the "Daily News." Crown 8vo. 6s.

ASHE (T.) Author of "The Sorrows of Hypsipyle."

EDITH; OR, LOVE AND LIFE IN CHESHIRE. Sewed. 6d.

ASHTON (John).

ROUGH NOTES OF A VISIT TO BELGIUM, SEDAN, AND PARIS, in September, 1870-71. Crown 8vo. 3s. 6d.

AUNT MARY'S BRAN PIE.

By the author of "St. Olave's," "When I was a Little Girl," etc. Illustrated. 3s. 6d.

SUNNYLAND STORIES. Fcap. 8vo. Illustrated. 3s. 6d.

Being two of Henry S. King and Co.'s Three and Sixpenny Series of Children's Books.

AURORA: A Volume of Verse. 5s.

AYRTON (J. C.)

A SCOTCH WOOLING. 2 vols. Crown 8vo.

BAGEHOT (Walter).

PHYSICS AND POLITICS; or, Thoughts on the Application of the Principles of "Natural Selection" and "Inheritance" to Political Society. Third Edition. Crown 8vo. 4s.

Volume II. of the International Scientific Series.

THE ENGLISH CONSTITUTION. A New Edition, Revised and Corrected, with an Introductory Dissertation on Recent Changes and Events. Crown 8vo. 7s. 6d.

LOMBARD STREET. A Description of the Money Market. Crown 8vo. Sixth Edition. 7s. 6d.

BAIN (Alexander), LL.D.

MIND AND BODY. The Theories of their Relation. Third Edition. Crown 8vo. 4s.

Volume IV. of the International Scientific Series.

BANKS (Mrs. G. Linnæus).

GOD'S PROVIDENCE HOUSE. Crown 8vo. 3s. 6d.

One of the volumes of the Cornhill Library of Fiction.

BARING (T. C., Esq.) M.P., late Fellow of Brasenose College, Oxford.

THE ODES OF PINDAR. A Translation.

BARLEE (Ellen).

LOCKED OUT; A Tale of the Strike. With a Frontispiece.
1s. 6d.

BAYNES (Rev. Canon R. H.), Editor of "Lyra Anglicana," etc.

HOME SONGS FOR QUIET HOURS. Second Edition. Fcap. 8vo. Cloth extra, 3s. 6d.

* * *This may also be had handsomely bound in Morocco with gilt edges.*

BECKER (Bernard H.)

THE SCIENTIFIC SOCIETIES OF LONDON. 1 vol. Crown 8vo. 5s.

BENNETT (Dr. W. C.)

SONGS FOR SAILORS. Dedicated by Special Request to H.R.H. the Duke of Edinburgh. Crown 8vo. 3s. 6d. With Steel Portrait and Illustrations.

An Edition in Illustrated Paper Covers, 1s.

BABY MAY. Home Poems and Ballads. With Frontispiece. Cloth elegant. Crown 8vo. 6s.

BABY MAY AND HOME POEMS. Fcap. 8vo. Sewed in Coloured Wrapper. 1s.

NARRATIVE POEMS AND BALLADS. Fcap. 8vo. Sewed in Coloured Wrapper. 1s.

BENNIE (Rev. Jas. Noble), M.A.

THE ETERNAL LIFE. Sermons preached during the last twelve years. Crown 8vo. 6s.

BERNARD (Bayle).

SAMUEL LOVER, THE LIFE AND UNPUBLISHED WORKS OF. In 2 vols. Post 8vo. With a Steel Portrait. 21s.

BETHAM-EDWARDS (Miss M.)

KITTY. Crown 8vo. With a Frontispiece. 3s. 6d.
One of the volumes of the Cornhill Library of Fiction.

MADEMOISELLE JOSEPHINE'S FRIDAYS, AND OTHER STORIES. Crown 8vo. 7s. 6d.

BISCOE (A. C.)

THE EARLS OF MIDDLETON, Lords of Clairmont and of Fettercairne, and the Middleton Family. 1 vol. Crown 8vo. 10s. 6d.

BLANC (Henry), M.D.

CHOLERA: HOW TO AVOID AND TREAT IT. Popular and Practical Notes. Crown 8vo. 4s. 6d.

BLUME (Major William).

THE OPERATIONS OF THE GERMAN ARMIES IN FRANCE, from Sedan to the end of the war of 1870-71. With Map. From the Journals of the Head-quarters Staff. Translated by the late E. M. Jones, Maj. 20th Foot, Prof. of Mil. Hist., Sandhurst. Demy 8vo. 9s.

BOGUSLAWSKI (Captain A. von).

TACTICAL DEDUCTIONS FROM THE WAR OF 1870-71. Translated by Colonel Lumley Graham, late 18th (Royal Irish) Regiment. Third Edition, Revised and Corrected. Demy 8vo. 7s.
A volume of Henry S. King and Co.'s Series of Military Works.

BONWICK (James).

THE TASMANIAN LILY. Cr. 8vo. With Frontispiece. 5s.

MIKE HOWE, THE BUSHRANGER OF VAN DIEMEN'S LAND. Crown 8vo. With Frontispiece. 5s.

BOSWELL (R. B.), M.A., Oxon.

METRICAL TRANSLATIONS FROM THE GREEK AND LATIN POETS, and other Poems. Crown 8vo. 5s.

BOTHMER (Countess Von).

CRUEL AS THE GRAVE. A Novel. 3 vols.

BOWRING (L.), C.S.I., Lord Canning's Private Secretary, and for many years Chief Commissioner of Mysore and Coorg.

EASTERN EXPERIENCES. Illustrated with Maps and Diagrams. Demy 8vo. 16s.

BRAVE MEN'S FOOTSTEPS. By the Editor of "Men who have Risen." A Book of Example and Anecdote for Young People. With Four Illustrations by C. Doyle. Third Edition. Crown 8vo. 3s. 6d.

BRIALMONT (Colonel A.)

HASTY INTRENCHMENTS. Translated by Lieut. Charles A. Empsom, R.A. With nine Plates. Demy 8vo. 6s.

BRIEFS AND PAPERS. Being Sketches of the Bar and the Press. By Two Idle Apprentices. Crown 8vo. 7s. 6d.

BROOKE (Rev. Stopford A.), M.A., Chaplain in Ordinary to Her Majesty the Queen.

THE LATE REV. F. W. ROBERTSON, M.A., LIFE AND LETTERS OF. Edited by Stopford Brooke, M.A.

I. In 2 vols., uniform with the Sermons. Steel Portrait. 7s. 6d.

II. Library Edition. 8vo. Two Steel Portraits. 12s.

III. A Popular Edition, in 1 vol. 8vo. 6s.

BROOKE (Rev. Stopford A.), M.A.

THEOLOGY IN THE ENGLISH POETS.—COWPER, COLERIDGE, WORDSWORTH, and BURNS. Second Edition. Post 8vo. 9s.

CHRIST IN MODERN LIFE. Sermons Preached in St. James's Chapel, York Street, London. Eighth Edition. Crown 8vo. 7s. 6d.

FREEDOM IN THE CHURCH OF ENGLAND. Six Sermons suggested by the Voysey Judgment. Second Edition. Crown 8vo. 3s. 6d.

SERMONS Preached in St. James's Chapel, York Street, London. Eighth Edition. Crown 8vo. 6s.

SERMONS Preached in St. James's Chapel, York Street, London. Second Series. By the Rev. Stopford A. Brooke, M.A., Chaplain in Ordinary to the Queen. Second Edition. Crown 8vo. 7s.

FREDERICK DENISON MAURICE: The Life and Work of. A Memorial Sermon. Crown 8vo. Sewed. 1s.

BROOKE (W. G.), M.A., Barrister-at-Law, Author of "Six Privy Council Judgments," &c.

THE PUBLIC WORSHIP REGULATION ACT. With a Classified Statement of its Provisions, Notes, and Index. Third Edition, revised and corrected. Crown 8vo. 3s. 6d.

SIX PRIVY COUNCIL JUDGMENTS—1850-1872. Annotated by. Third Edition. Crown 8vo. 9s.

BROWN (Rev. J. Baldwin), B.A., Author of "First Principles of Ecclesiastical Truth," etc.

THE HIGHER LIFE. Its Reality, Experience, and Destiny. Fourth Edition. Crown 8vo. 7s. 6d.

THE DOCTRINE OF ANNIHILATION IN THE LIGHT OF THE GOSPEL OF LOVE. Five Discourses by the Rev. J. Baldwin Brown, B.A., Author of "The Higher Life," etc. Second Edition. Crown 8vo. 2s. 6d.

BROWN (John Croumbie), LL.D., etc.

REBOISEMENT IN FRANCE; or, Records of the Replanting of the Alps, the Cevennes, and the Pyrenees with Trees, Herbage, and Bush, with a view to arresting and preventing the destructive consequences and effects of Torrents. 1 vol. Demy 8vo.

THE HYDROLOGY OF SOUTHERN AFRICA. 10s. 6d.

BROWNE (Rev. Marmaduke E.)

UNTIL THE DAY DAWN. Four Advent Lectures delivered in the Episcopal Chapel, Milverton, Warwickshire, on the Sunday evenings during Advent, 1870. Crown 8vo. 2s. 6d.

BRYANT (William Cullen).

POEMS. Red-line Edition. Handsomely bound. With 24 Illustrations and Portrait of the Author. 7s. 6d.

A Cheaper Edition, with Frontispiece. 3s. 6d.

BUCHANAN (Robert).

POETICAL WORKS. Collected Edition, in 3 Vols., price 6s. each.

Vol. I.—“Ballads and Romances;” “Ballads and Poems of Life,” and a Portrait of the Author.

Vol. II.—“Ballads and Poems of Life;” “Allegories and Sonnets.”

Vol. III.—“Cruiskeen Sonnets;” “Book of Orm;” “Political Mystics.”

MASTER-SPIRITS. Post 8vo. 10s. 6d.

BULKELEY (Rev. Henry J.)

WALLED IN, and other Poems. Crown 8vo. 5s.

BUNNETT (F. E.)

LEONORA CHRISTINA, MEMOIRS OF, Daughter of Christian IV. of Denmark; Written during her Imprisonment in the Blue Tower of the Royal Palace at Copenhagen, 1663-1685. Translated by F. E. Bunnètt. With an Autotype Portrait of the Princess. Medium 8vo. A New and Cheaper Edition. 5s.

LINKED AT LAST. 1 vol. Crown 8vo.

UNDER A CLOUD; OR, JOHANNES OLAF. By E. D. Wille. Translated by F. E. Bunnètt. 3 vols.

BURTON (Mrs. Richard).

THE INNER LIFE OF SYRIA, PALESTINE, AND THE HOLY LAND. 2 vols. Demy 8vo. 24s.

BUTLER (Josephine E.)

JOHN GREY (of Dilston): MEMOIRS. By his Daughter. New and Cheaper Edition. Crown 8vo. 3s. 6d.

CADELL (Mrs. H. M.)

IDA CRAVEN. A Novel. 3 vols. Crown 8vo.

CAMDEN (Charles).

HOITY TOITY, THE GOOD LITTLE FELLOW. With Eleven Illustrations. Crown 8vo. 3s. 6d.

THE TRAVELLING MENAGERIE. By Charles Camden, Author of “Hoity Toity.” With Ten Illustrations by J. Mahoney. Crown 8vo. 3s. 6d.

The above form part of Henry S. King & Co.'s Three and Sixpenny Series of Children's Books.

CARLISLE (A. D.), B.A., Trin. Coll., Camb.

ROUND THE WORLD IN 1870. A Volume of Travels, with Maps. New and Cheaper Edition. Demy 8vo. 6s.

CARNE (Miss E. T.)

THE REALM OF TRUTH. Crown 8vo. 5s. 6d.

CARPENTER (E.)

NARCISSUS AND OTHER POEMS. Fcap. 8vo. 5s.

CARPENTER (W. B.), LL.D., M.D., F.R.S., etc.

THE PRINCIPLES OF MENTAL PHYSIOLOGY. With their Applications to the Training and Discipline of the Mind, and the Study of its Morbid Conditions. 8vo. Illustrated. 12s.

CARR (Lisle).

JUDITH GWYNNE. 3 vols. Crown 8vo. Second Edition.

CHRISTOPHERSON (The late Rev. Henry), M.A.,
Assistant Minister at Trinity Church, Brighton.

SERMONS. Crown 8vo. Cloth. 7s. 6d. With an Introduction by John Rae, LL.D., F.S.A.

CLAYTON (Cecil).

EFFIE'S GAME; HOW SHE LOST AND HOW SHE WON.
A Novel. 2 vols.

CLERK (Mrs. Godfrey), Author of "The Antipodes and Round the World."

'ILAM EN NAS. Historical Tales and Anecdotes of the Times of the Early Khalifahs. Translated from the Arabic Originals. Illustrated with Historical and Explanatory Notes. Crown 8vo. 7s.

CLERY (C.), Captain 32nd Light Infantry, Professor of Tactics Royal Military College, Sandhurst.

MINOR TACTICS. Demy 8vo. New Edition. With 26 Plans. 16s.

CLODD (Edward), F.R.A.S.

THE CHILDHOOD OF THE WORLD: a Simple Account of Man in Early Times. New Edition. Crown 8vo. 3s.
A Special Edition for Schools. 1s.

THE CHILDHOOD OF RELIGIONS. Including a Simple Account of the Birth and Growth of Myths and Legends. Crown 8vo. 5s.

COLERIDGE (Sara).

PRETTY LESSONS IN VERSE FOR GOOD CHILDREN, with some Lessons in Latin, in Easy Rhyme. A New Edition. Illustrated. 3s. 6d.

COLERIDGE (Sara).

PHANTASMION. A Fairy Romance. With an Introductory Preface by the Right Hon. Lord Coleridge of Ottery St. Mary. A New Edition. Illustrated. 7s. 6d.

MEMOIR AND LETTERS OF SARA COLERIDGE. Edited by her Daughter. Third Edition, Revised and Corrected. With Index. 2 vols. Crown 8vo. With Two Portraits. 24s.
Cheap Edition. With one Portrait. 7s. 6d.

COLLINS (Mortimer).

THE PRINCESS CLARICE. A Story of 1871. 2 vols.

SQUIRE SILCHESTER'S WHIM. By Mortimer Collins, Author of "Marquis and Merchant," etc. 3 vols.

MIRANDA. A Midsummer Madness. 3 vols.

THE INN OF STRANGE MEETINGS, AND OTHER POEMS. Crown 8vo. 5s.

THE SECRET OF LONG LIFE. Dedicated by special permission to Lord St. Leonard's. Fourth Edition. Large crown 8vo. 5s.

COLLINS (Rev. Richard), M.A.

MISSIONARY ENTERPRISE IN THE EAST. With special reference to the Syrian Christians of Malabar, and the results of modern Missions. With Four Illustrations. Crown 8vo. 6s.

CONWAY (Moncure D.)

REPUBLICAN SUPERSTITIONS. Illustrated by the Political History of the United States. Including a Correspondence with M. Louis Blanc. Crown 8vo. 5s.

CONYERS (Ansley).

CHESTERLEIGH. 3 vols. Crown 8vo.

COOKE (M. C.), M.A., LL.D.

FUNGI; their Nature, Influences, Uses, etc. Edited by the Rev. M. J. Berkeley, M.A., F.L.S. Second Edition. Crown 8vo. With Illustrations. 5s.

Being Vol. XIV. of the International Scientific Series.

COOKE (Professor Josiah P.), of the Harvard University.

THE NEW CHEMISTRY. Second Edition. With Thirty-one Illustrations. 5s.

Vol. IX. of the International Scientific Series.

SCIENTIFIC CULTURE. Crown 8vo. Cloth. 1s.

COOPER (T. T.)

THE MISHMEE HILLS: an Account of a Journey made in an Attempt to Penetrate Thibet from Assam, to open New Routes for Commerce. Second Edition. With Four Illustrations and Map. Demy 8vo. 10s. 6d.

CALDERON.

CALDERON'S DRAMAS: The Wonder-Working Magician—Life is a Dream—The Purgatory of St. Patrick. Translated by Denis Florence MacCarthy. Post 8vo. 10s.

CORNHILL LIBRARY OF FICTION, The. 3s. 6d. per Volume.

HALF-A-DOZEN DAUGHTERS. By J. Masterman.

THE HOUSE OF RABY. By Mrs. G. Hooper.

A FIGHT FOR LIFE. By Moy Thomas.

ROBIN GRAY. By Charles Gibbon.

KITTY. By Miss M. Betham-Edwards.

HIRELL. By John Saunders.

ONE OF TWO; or, The Left-Handed Bride. By J. Hain Friswell.

READY-MONEY MORTIBOY. A Matter-of-Fact Story.

GOD'S PROVIDENCE HOUSE. By Mrs. G. L. Banks.

FOR LACK OF GOLD. By Charles Gibbon.

ABEL DRAKE'S WIFE. By John Saunders.

COSMOS. A Poem. Fcap. 8vo. 3s. 6d.

SUBJECT.—Nature in the Past and in the Present—Man in the Past and in the Present—The Future.

COTTON (Robert Turner).

MR. CARINGTON. A Tale of Love and Conspiracy. 3 vols. Crown 8vo.

CROFT (Herbert), Barrister-at-Law.

SIR THOMAS ELYOT'S "GOVERNOUR." A Reprint of the Original Edition, Revised and Corrected, with Copious Notes and a Life of the Author.

*** No Complete Reprint of the above Work is known to exist.

CUMMINS (Henry Irwin), M.A.

PAROCHIAL CHARITIES OF THE CITY OF LONDON. Sewed. 1s.

CURWEN (Henry).

SORROW AND SONG: Studies of Literary Struggle. Henry Mürger—Novalis—Alexander Petöfi—Honoré de Balzac—Edgar Allan Poe—André Chénier. 2 vols. Crown 8vo. 15s.

DAVIDSON (Samuel), D.D., LL.D.

THE NEW TESTAMENT, TRANSLATED FROM THE LATEST GREEK TEXT OF TISCHENDORF. Post 8vo. 10s. 6d.

DAVIES (G. Christopher).

MOUNTAIN, MEADOW, AND MERE: a Series of Outdoor Sketches of Sport, Scenery, Adventures, and Natural History. With Sixteen Illustrations by Bosworth W. Harcourt. Crown 8vo. 6s.

RAMBLES AND ADVENTURES OF OUR SCHOOL FIELD CLUB. Crown 8vo. With 4 Illustrations. 5s.

DAVIES (Rev. J. Llewelyn), M.A.

THEOLOGY AND MORALITY. Essays on Questions of Belief and Practice. Crown 8vo. 7s. 6d.

DE KERKADEC (Vicomtesse Solange).

A CHEQUERED LIFE, being Memoirs of the Vicomtesse de Leoville Meilhan. Edited by. Crown 8vo. 7s. 6d.

Containing many recollections of the First Emperor Napoleon and his Court.

DE L'HOSTE (Colonel E. P).

THE DESERT PASTOR, JEAN JAROUSSEAU. Translated from the French of Eugène Pelletan. In fcap. 8vo., with an Engraved Frontispiece. New Edition. 3s. 6d.

One of Henry S. King and Co.'s Three and Sixpenny Books for the Young.

DE LIEFDE (Jacob).

THE GREAT DUTCH ADMIRALS. Crown 8vo. With Eleven Illustrations by Townley Green and others. 5s.

One of Henry S. King and Co.'s Five Shilling Books for the Young.

DE REDCLIFFE (Viscount Stratford), P.C., K.G., G.C.B.

WHY AM I A CHRISTIAN? Fifth Edition. Crown 8vo. 3s.

DE TOCQUEVILLE (Alexis).

CORRESPONDENCE AND CONVERSATIONS OF, WITH NASSAU WILLIAM SENIOR. 2 vols. Post 8vo. 21s.

DE VERE (Aubrey).

ALEXANDER THE GREAT. A Dramatic Poem. Small crown 8vo. 5s.

THE INFANT BRIDAL, AND OTHER POEMS. A New Edition. Fcap. 8vo.

THE LEGENDS OF ST. PATRICK, and other Poems. Small crown 8vo. 5s.

DE WILLE (E.)

UNDER A CLOUD; OR, JOHANNES OLAF. A Novel. Translated by F. E. Bunnètt. 3 vols. Crown 8vo.

DENNIS (John).

ENGLISH SONNETS. Collected and Arranged. Fcap. 8vo. Elegantly bound. 3s. 6d.

DOBSON (Austin).

VIGNETTES IN RHYME AND VERS DE SOCIÉTÉ. Second Edition. Fcap. 8vo. 5s.

DONNÉ (Alphonse), M.D.

CHANGE OF AIR AND SCENE. A Physician's Hints about Doctors, Patients, Hygiene, and Society; with Notes of Excursions for Health in the Pyrenees, and amongst the Watering-places of France (Inland and Seaward), Switzerland, Corsica, and the Mediterranean. A New Edition. Large post 8vo. 9s.

DOWDEN (Edward), LL.D.

SHAKSPERE: a Critical Study of his Mind and Art. Post 8vo. 12s.

DOWNTON (Rev. Henry), M.A.

HYMNS AND VERSES. Original and Translated. Small crown 8vo. 3s. 6d.

DRAPER (John William), M.D., LL.D. Professor in the University of New York; Author of "A Treatise on Human Physiology."

HISTORY OF THE CONFLICT BETWEEN RELIGION AND SCIENCE. Fourth Edition. 5s.

Vol. XIII. of the International Scientific Series.

DREW (Rev. G. S.), M.A., Vicar of Trinity, Lambeth.

SCRIPTURE LANDS IN CONNECTION WITH THEIR HISTORY. Second Edition. 8vo. 10s. 6d.

NAZARETH: ITS LIFE AND LESSONS. Second Edition. Crown 8vo. 5s.

DREW (Rev. G. S.), M.A.

THE DIVINE KINGDOM ON EARTH AS IT IS IN HEAVEN.
8vo. 10s. 6d.

THE SON OF MAN: His Life and Ministry. Crown 8vo.
7s. 6d.

DREWRY (G. Overend), M.D.

THE COMMON-SENSE MANAGEMENT OF THE STOMACH.
Fcap. 8vo. 2s. 6d.

DURAND (Lady).

**IMITATIONS FROM THE GERMAN OF SPITTA AND
TERSTEGEN.** Fcap. 8vo. 4s.

DU VERNON (Colonel von Verdy).

STUDIES IN LEADING TROOPS. An authorized and accurate
Translation by Lieutenant H. J. T. Hildyard, 71st Foot. Parts I.
and II. Demy 8vo. 7s.

This is one of Henry S. King and Co.'s Series of Military Works.

E. A. V.

JOSEPH MAZZINI: A Memoir. With Two Essays by
Mazzini—"Thoughts on Democracy," and "The Duties of
Man." Dedicated to the Working Classes by P. H. Taylor, M.P.
Crown 8vo. With Two Portraits. 3s. 6d.

EDEN (Frederic).

THE NILE WITHOUT A DRAGOMAN. Second Edition.
Crown 8vo. 7s. 6d.

EDWARDS (Rev. Basil).

MINOR CHORDS; OR, SONGS FOR THE SUFFERING: a
Volume of Verse. Fcap. 8vo. Cloth, 3s. 6d.; paper, 2s. 6d.

EILOART (Mrs.)

LADY MORETOUN'S DAUGHTER. 3 vols. Crown 8vo.

ENGLISH CLERGYMAN.

**AN ESSAY ON THE RULE OF FAITH AND CREED OF
ATHANASIUS.** Shall the Rubric preceding the Creed be
removed from the Prayer-book? 8vo. Sewed. 1s.

EROS AGONISTES. Poems. By E. B. D. Fcap. 8vo.
3s. 6d.

EVANS (Mark).

THE STORY OF OUR FATHER'S LOVE, told to Children;
being a New and Enlarged Edition of Theology for Children.
3s. 6d.

EYRE (Maj.-Gen. Sir Vincent), C.B., K.C.S.I., etc.

LAYS OF A KNIGHT-ERRANT IN MANY LANDS. Square crown 8vo. With Six Illustrations. 7s. 6d.

Pharaoh Land. | Home Land. | Wonder Land. | Rhine Land.

FAITHFULL (Mrs. Francis G.)

LOVE ME, OR LOVE ME NOT. 3 vols. Crown 8vo.

FARQUHARSON (Martha).

I. ELSIE DINSMORE. Crown 8vo. 3s. 6d.

II. ELSIE'S GIRLHOOD. Crown 8vo. 3s. 6d.

III. ELSIE'S HOLIDAYS AT ROSELANDS. Crown 8vo. 3s. 6d.

These are volumes of Henry S. King and Co.'s Series of Three and Sixpenny Books for the Young.

FAVRE (Mons. Jules).

THE GOVERNMENT OF THE NATIONAL DEFENCE. From the 30th June to the 31st October, 1870. The Plain Statement of a Member. 1 vol. Demy 8vo. 10s. 6d.

FISHER (Alice).

HIS QUEEN. 3 vols. Crown 8vo.

FORBES (Archibald).

SOLDIERING AND SCRIBBLING. A Series of Sketches. Crown 8vo. 7s. 6d.

FOTHERGILL (JESSIE).

HEALEY. A Romance. 3 vols. Crown 8vo.

ALDYTH. A Novel. 3 vols. Crown 8vo.

FOWLE (Rev. T. W.), M.A.

THE RECONCILIATION OF RELIGION AND SCIENCE. Being Essays on Immortality, Inspiration, Miracles, and the Being of Christ. Demy 8vo. 10s. 6d.

FRASER (Donald), Accountant to the British-Indian Steam Navigation Company, Limited.

EXCHANGE TABLES OF STERLING AND INDIAN RUPEE CURRENCY, upon a new and extended system, embracing Values from One Farthing to One Hundred Thousand Pounds, and at Rates progressing, in Sixteenths of a Penny, from 1s. 9d. to 2s. 3d. per Rupee. Royal 8vo. 10s. 6d.

FRERE (Sir H. Bartle E.), G.C.B., G.C.S.I., etc.

THE THREATENED FAMINE IN BENGAL; How it may be Met, and the Recurrence of Famines in India Prevented. Being No. 1 of "Occasional Notes on Indian Affairs." Crown 8vo. With 3 Maps. 5s.

FRISWELL (J. Hain).

THE BETTER SELF. Essays for Home Life. Crown 8vo. 6s.
Contents :—Beginning at Home—The Girls at Home—The Wife's Mother—Pride in the Family—Discontent and Grumbling—Domestic Economy—On Keeping People Down—Likes and Dislikes—On Falling Out—Peace.

ONE OF TWO; or, The Left-Handed Bride. Crown 8vo. With a Frontispiece. 3s. 6d.

Being a Volume of the Cornhill Library of Fiction.

GARDNER (John), M.D.

LONGEVITY; THE MEANS OF PROLONGING LIFE AFTER MIDDLE AGE. Third Edition, revised and enlarged. Small crown 8vo. 4s.

GARRETT (Edward).

BY STILL WATERS. A Story for Quiet Hours. Crown 8vo. With Seven Illustrations. 6s.

GIBBON (Charles).

FOR LACK OF GOLD. Crown 8vo. With a Frontispiece. 3s. 6d.

ROBIN GRAY. Crown 8vo. With a Frontispiece. 3s. 6d.

The above Volumes form part of the Cornhill Library of Fiction.

GILBERT (Mrs.)

MRS. GILBERT, FORMERLY ANN TAYLOR, AUTOBIOGRAPHY AND OTHER MEMORIALS OF. Edited by Josiah Gilbert. In 2 vols. Post 8vo. With 2 Steel Portraits and several Wood Engravings. 24s.

GILL (Rev. W. W.)

MYTHS AND SONGS OF THE SOUTH PACIFIC. With a Preface by F. Max Müller, M.A., Professor of Comparative Philology at Oxford. 1 vol. Post 8vo.

GODKIN (James).

THE RELIGIOUS HISTORY OF IRELAND: Primitive, Papal, and Protestant. Including the Evangelical Missions, Catholic Agitations, and Church Progress of the last half Century. 1 vol. 8vo. 12s.

GODWIN (William).

WILLIAM GODWIN: his Friend's Contemporaries. By C. Kegan Paul. 2 vols. Demy 8vo. With Portraits.

THE GENIUS OF CHRISTIANITY UNVEILED. Being Essays never before published. Edited, with a Preface, by C. Kegan Paul. 1 vol. Crown 8vo. 7s. 6d.

GOETZE (Capt. A. von), Captain of the Prussian Corps of Engineers attached to the Engineer Committee, and Instructor at the Military Academy.

OPERATIONS OF THE GERMAN ENGINEERS DURING THE WAR OF 1870-1871. Published by Authority, and in accordance with Official Documents. Translated from the German by Colonel G. Graham, V.C., C.B., R.E. Demy 8vo. Cloth. With 6 large Maps. 21s.

GOODMAN (Walter).

CUBA, THE PEARL OF THE ANTILLES. Crown 8vo. 7s. 6d.

GOSSE (Edmund W.)

ON VIOL AND FLUTE. With Title-page specially designed by William B. Scott. Crown 8vo. 5s.

GRANVILLE (A. B.), M.D., F.R.S., etc.

AUTOBIOGRAPHY OF A. B. GRANVILLE, F.R.S., etc. Edited, with a brief account of the concluding years of his life, by his youngest Daughter, Paulina B. Granville. 2 vols. Demy 8vo. With a Portrait. 32s.

GRAY (Mrs. Russell).

LISETTE'S VENTURE. A Novel. 2 vols. Crown 8vo. 21s.

GREEN (T. Bowden).

FRAGMENTS OF THOUGHT. Dedicated by permission to the Poet Laureate. Crown 8vo. 7s. 6d.

GREENWOOD (James), "The Amateur Casual."

IN STRANGE COMPANY; or, The Note Book of a Roving Correspondent. Second Edition. Crown 8vo. 6s.

GREY (John), of Dilston.

JOHN GREY (of Dilston): MEMOIRS. By Josephine E. Butler. New and Cheaper Edition. Crown 8vo. 3s. 6d.

GRIFFITH (Rev. T.), A.M., Prebendary of St. Paul's.

STUDIES OF THE DIVINE MASTER. Demy 8vo. 12s.

GRIFFITHS (Captain Arthur).

THE QUEEN'S SHILLING. A Novel. 2 vols. 21s.

MEMORIALS OF MILLBANK, AND CHAPTERS IN PRISON HISTORY. 2 vols. Post 8vo. 21s. With Illustrations.

GRUNER (M. L.)

STUDIES OF BLAST FURNACE PHENOMENA. Translated by L. D. B. Gordon, F.R.S.E., F.G.S. Demy 8vo. 7s. 6d.

GURNEY (Rev. Archer Thompson).

WORDS OF FAITH AND CHEER. A Mission of Instruction and Suggestion. 1 vol. Crown 8vo. 6s.

FIRST PRINCIPLES IN CHURCH AND STATE. Demy 8vo. Sewed. 1s. 6d.

HAECKEL (Professor Ernst), of the University of Jena.

THE HISTORY OF CREATION. A Popular Account of the Development of the Earth and its Inhabitants, according to the Theories of Kant, Laplace, Lamarck, and Darwin. The Translation revised by Professor E. Ray Lankester, M.A., F.R.S. With Coloured Plates and Genealogical Trees of the various groups of both plants and animals. 2 vols. Post 8vo. 32s.

THE HISTORY OF THE EVOLUTION OF MAN. Translated by E. A. Van Rhyn and L. Elsberg, M.D. (University of New York), with Notes and Additions sanctioned by the Author. Post 8vo.

HARCOURT (Capt. A. F. P.)

THE SHAKESPEARE ARGOSY: Containing much of the wealth of Shakespeare's Wisdom and Wit, alphabetically arranged and classified. Crown 8vo. 6s.

Haweis (Rev. H. R.), M.A.

SPEECH IN SEASON. Third Edition. Crown 8vo. 9s.

THOUGHTS FOR THE TIMES. Eighth Edition. Crown 8vo. 7s. 6d.

UNSECTARIAN FAMILY PRAYERS, for Morning and Evening for a Week, with short selected passages from the Bible. Square crown 8vo. 3s. 6d.

HAWTHORNE (Julian).

BRESSANT. A Romance. 2 vols. Crown 8vo. 21s.

IDOLATRY. A Romance. 2 vols. Crown 8vo. 21s.

HAWTHORNE (Nathaniel).

NATHANIEL HAWTHORNE. A Memoir, with Stories now first published in this country. By H. A. Page. Post 8vo. 7s. 6d.

SEPTIMUS. A Romance. Second Edition. Crown 8vo. 9s.

HAYMAN (Henry), D.D., late Head Master of Rugby School.

RUGBY SCHOOL SERMONS. With an Introductory Essay on the Indwelling of the Holy Spirit. Crown 8vo. 7s. 6d.

HEATHERGATE. A Story of Scottish Life and Character. By a New Author. 2 vols. Crown 8vo. 21s.

HELLWALD (Baron F. Von).

THE RUSSIANS IN CENTRAL ASIA. A Critical Examination, down to the present time, of the Geography and History of Central Asia. Translated by Lieut.-Col. Theodore Wirgman, LL.B. In 1 vol. Large post 8vo. With Map. 12s.

HELVIG (Captain Hugo).

THE OPERATIONS OF THE BAVARIAN ARMY CORPS. Translated by Captain G. S. Schwabe. With Five large Maps. In 2 vols. Demy 8vo. 24s.

This is one of Henry S. King and Co.'s Series of Military Books.

HINTON (James), late Aural Surgeon to Guy's Hospital.

THE PLACE OF THE PHYSICIAN. Being the Introductory Lecture at Guy's Hospital, 1873-74; to which is added **ESSAYS ON THE LAW OF HUMAN LIFE, AND ON THE RELATION BETWEEN ORGANIC AND INORGANIC WORLDS.** Crown 8vo. 3s. 6d.

PHYSIOLOGY FOR PRACTICAL USE. By various writers. Second Edition. With 50 Illustrations. 2 vols. Crown 8vo. 12s. 6d.

AN ATLAS OF DISEASES OF THE MEMBRANA TYMPANI. With Descriptive Text. Post 8vo. £6 6s.

THE QUESTIONS OF AURAL SURGERY. Post 8vo. With Illustrations. 2 vols. 12s. 6d.

HOCKLEY (W. B.)

TALES OF THE ZENANA; or, A Nuwab's Leisure Hours. By the Author of "Pandurang Hari." With a Preface by Lord Stanley of Alderley. 2 vols. Crown 8vo. 21s.

PANDURANG HARI; or, Memoirs of a Hindoo. A Tale of Mahratta Life sixty years ago. With a Preface by Sir H. Bartle E. Frere, G.C.S.I., etc. 2 vols. Crown 8vo. 21s.

HOFFBAUER (Captain).

THE GERMAN ARTILLERY IN THE BATTLES NEAR METZ. Based on the official reports of the German Artillery. Translated by Capt. E. O. Hollist. Demy 8vo. With Map and Plans. 21s.

This is one of the volumes in Henry S. King and Co.'s Military Series.

HOLROYD (Captain W. R. M.), Bengal Staff Corps, Director of Public Instruction, Punjab.

TAS-HIL UL KALAM; or, Hindustani made Easy. Crown 8vo. 5s.

HOPE (Lieut. James).

IN QUEST OF COOLIES. With Illustrations. Crown 8vo. 6s.

HOOPER (Mrs. G.)

THE HOUSE OF RABY. With a Frontispiece. Crown 8vo. 3s. 6d.

One of the volumes of the Cornhill Library of Fiction.

HOOPER (Mary).

LITTLE DINNERS: HOW TO SERVE THEM WITH ELEGANCE AND ECONOMY. Seventh Edition. 1 vol. Crown 8vo. 5s.

COOKERY FOR INVALIDS. Crown 8vo.

HOPKINS (Manley).

THE PORT OF REFUGE; or, Counsel and Aid to Shipmasters in Difficulty, Doubt, or Distress. Crown 8vo. 6s.

HOWARD (Mary M.), Author of "Brampton Rectory."

BEATRICE AYLMER, AND OTHER TALES. Crown 8vo. 6s.

HOWARD (Rev. G. B.)

AN OLD LEGEND OF ST. PAUL'S. Fcap. 8vo. 4s. 6d.

HOWE (Cupples), Master Mariner.

THE DESERTED SHIP. A real story of the Atlantic. Illustrated by Townley Green. Crown 8vo. 3s. 6d.

One of Henry S. King and Co.'s Three and Sixpenny Books for the Young.

HOWELL (James).

A TALE OF THE SEA, SONNETS, AND OTHER POEMS. Fcap. 8vo. 5s.

HUGHES (Allison).

PENELOPE, AND OTHER POEMS. Fcap. 8vo. 4s. 6d.

HULL (Edmund C. P.)

THE EUROPEAN IN INDIA. A Handbook of Practical Information for those proceeding to, or residing in, the East Indies, relating to Outfits, Routes, Time for Departure, Indian Climate, etc. With a **MEDICAL GUIDE FOR ANGLO-INDIANS.** By R. R. S. Mair, M.D., F.R.C.S.E., late Deputy Coroner of Madras. Second Edition, Revised and Corrected. In 1 vol. Post 8vo. 6s.

HUMPHREY (Rev. W.), of the Congregation of the Oblates of St. Charles.

MR. FITZJAMES STEPHEN AND CARDINAL BELLARMINE. Demy 8vo. Sewed. 1s.

HUTTON (James).

MISSIONARY LIFE IN THE SOUTHERN SEAS. With Illustrations. Crown 8vo. 7s. 6d.

INTERNATIONAL SCIENTIFIC SERIES (The).

- I. THE FORMS OF WATER IN CLOUDS AND RIVERS, ICE AND GLACIERS.** By J. Tyndall, LL.D., F.R.S. With 14 Illustrations. Fifth Edition. 5s.
- II. PHYSICS AND POLITICS;** or, Thoughts on the Application of the Principles of "Natural Selection" and "Inheritance" to Political Society. By Walter Bagehot. Third Edition. 4s.
- III. FOODS.** By Edward Smith, M.D., LL.B., F.R.S. Profusely Illustrated. Third Edition. 5s.
- IV. MIND AND BODY:** The Theories of their Relation. By Alexander Bain, LL.D. Third Edition. With Four Illustrations. 4s.
- V. THE STUDY OF SOCIOLOGY.** By Herbert Spencer. Fourth Edition. 5s.
- VI. ON THE CONSERVATION OF ENERGY.** By Balfour Stewart, M.D., LL.D., F.R.S. With 14 Engravings. Third Edition. 5s.
- VII. ANIMAL LOCOMOTION;** or, Walking, Swimming, and Flying. By J. B. Pettigrew, M.D., F.R.S. Second Edition. With 119 Illustrations. 5s.
- VIII. RESPONSIBILITY IN MENTAL DISEASE.** By Henry Maudsley, M.D. Second Edition. 5s.
- IX. THE NEW CHEMISTRY.** By Professor J. P. Cooke, of the Harvard University. Second Edition. With 31 Illustrations. 5s.
- X. THE SCIENCE OF LAW.** By Professor Sheldon Amos. Second Edition. 5s.
- XI. ANIMAL MECHANISM.** A Treatise on Terrestrial and Aerial Locomotion. By Professor E. J. Marey. With 117 Illustrations. Second Edition. 5s.
- XII. THE DOCTRINE OF DESCENT AND DARWINISM.** By Professor Oscar Schmidt (Strasburg University). Second Edition. With 26 Illustrations. 5s.
- XIII. THE HISTORY OF THE CONFLICT BETWEEN RELIGION AND SCIENCE.** By Professor J. W. Draper. Fifth Edition. 5s.

INTERNATIONAL SCIENTIFIC SERIES (The).

XIV. FUNGI; their Nature, Influences, Uses, etc. By M. C. Cooke, M.A., LL.D. Edited by the Rev. M. J. Berkeley, M.A., F.L.S. Second Edition. Crown 8vo. With numerous Illustrations. 5s.

XV. THE CHEMICAL EFFECTS OF LIGHT AND PHOTOGRAPHY. By Dr. Hermann Vogel (Polytechnic Academy of Berlin). Second Edition. With 100 Illustrations. 5s.

XVI. THE LIFE AND GROWTH OF LANGUAGE. By William Dwight Whitney, Professor of Sanskrit and Comparative Philology in Yale College, New Haven. Second Edition. Crown 8vo. 5s.

XVII. MONEY AND THE MECHANISM OF EXCHANGE. By Prof. W. Stanley Jevons. Crown 8vo. 5s.

XVIII. THE NATURE OF LIGHT: With a General Account of Physical Optics. By Dr. Eugene Lommel, Professor of Physics in the University of Erlangen. With 148 Illustrations and a table of Spectra in Chromolithography. 5s.

XIX. ANIMAL PARASITES AND MESSMATES. By Monsieur Van Beneden, Professor of the University of Louvain, Correspondent of the Institute of France. Crown 8vo. With 83 Illustrations. 5s.

XX. THE FIVE SENSES OF MAN. By Professor Bernstein, of the University of Halle. Crown 8vo.

XXI. FERMENTATIONS. By Professor Schutzenberger, Director of the Chemical Laboratory at the Sorbonne. Crown 8vo.

INTERNATIONAL SCIENTIFIC SERIES (The).*Forthcoming Volumes.*

Prof. W. KINGDON CLIFFORD, M.A. The First Principles of the Exact Sciences explained to the Non-mathematical.

Prof. T. H. HUXLEY, LL.D., F.R.S. Bodily Motion and Consciousness.

Dr. W. B. CARPENTER, LL.D., F.R.S. The Physical Geography of the Sea.

Prof. WILLIAM ODLING, F.R.S. The Old Chemistry viewed from the New Standpoint.

W. LAUDER LINDSAY, M.D., F.R.S.E. Mind in the Lower Animals.

Sir JOHN LUBBOCK, Bart., F.R.S. On Ants and Bees.

Prof. W. T. THISELTON DYER, B.A., B.Sc. Form and Habit in Flowering Plants.

Mr. J. N. LOCKYER, F.R.S. Spectrum Analysis.

Prof. MICHAEL FOSTER, M.D. Protoplasm and the Cell Theory.

H. CHARLTON BASTIAN, M.D., F.R.S. The Brain as an Organ of Mind.

Prof. A. C. RAMSAY, LL.D., F.R.S. Earth Sculpture: Hills, Valleys, Mountains, Plains, Rivers, Lakes; how they were Produced, and how they have been Destroyed.

Prof. RUDOLPH VIRCHOW (Berlin Univ.) Morbid Physiological Action.

Prof. CLAUDE BERNARD. History of the Theories of Life.

Prof. H. SAINTE-CLAIRE DEVILLE. An Introduction to General Chemistry.

Prof. WURTZ. Atoms and the Atomic Theory.

Prof. De QUATREFAGES. The Human Race.

Prof. LACAZE-DUTHIERS. Zoology since Cuvier.

Prof. BERTHELOT. Chemical Synthesis.

INTERNATIONAL SCIENTIFIC SERIES (The).

(Forthcoming Volumes.)

Prof. J. ROSENTHAL. General Physiology of Muscles and Nerves.

Prof. JAMES D. DANA, M.A., LL.D. On Cephalization; or, Head-Characters in the Gradation and Progress of Life.

Prof. S. W. JOHNSON, M.A. On the Nutrition of Plants.

Prof. AUSTIN FLINT, Jr. M.D. The Nervous System, and its Relation to the Bodily Functions.

Prof. FERDINAND COHN (Breslau Univ.) Thallophytes (Algæ, Lichens, Fungi).

Prof. HERMANN (University of Zurich). Respiration.

Prof. LEUCKART (University of Leipsic). Outlines of Animal Organization.

Prof. LIEBREICH (University of Berlin). Outlines of Toxicology.

Prof. KUNDT (University of Strasburg). On Sound.

Prof. REES (University of Erlangen). On Parasitic Plants.

Prof. STEINTHAL (University of Berlin). Outlines of the Science of Language.

P. BERT (Professor of Physiology, Paris). Forms of Life and other Cosmical Conditions.

E. ALGLAVE (Professor of Constitutional and Administrative Law at Douai, and of Political Economy at Lille). The Primitive Elements of Political Constitutions.

P. LORAIN (Professor of Medicine, Paris). Modern Epidemics.

Mons. FREIDEL. The Functions of Organic Chemistry.

Mons. DEBRAY. Precious Metals.

Prof. CORFIELD, M.A., M.D. (Oxon.) Air in its relation to Health.

Prof. A. GIARD. General Embryology.

IGNOTUS.

CULMSHIRE FOLK. A Novel. New and Cheap Edition.
In 1 vol. Crown 8vo. 6s.

INGELOW (Jean).

THE LITTLE WONDER-HORN. A Second Series of "Stories Told to a Child." With Fifteen Illustrations. Square 24mo. 3s. 6d.

OFF THE SKELLIGS. (Her First Romance.) 4 vols. Crown 8vo. 42s.

JACKSON (T. G.)

MODERN GOTHIC ARCHITECTURE. Crown 8vo. 5s.

JACOB (Maj.-Gen. Sir G. Le Grand), K.C.S.I., C.B.

WESTERN INDIA BEFORE AND DURING THE MUTINIES.
Pictures drawn from life. Second Edition. Crown 8vo. 7s. 6d.

JENKINS (E.) and RAYMOND (J.), Esqs.

A LEGAL HANDBOOK FOR ARCHITECTS, BUILDERS, AND BUILDING OWNERS. Second Edition Revised. Crown 8vo. 6s.

JENKINS (Rev. R. C.), M.A., Rector of Lyminge, and
Honorary Canon of Canterbury.

THE PRIVILEGE OF PETER, Legally and Historically Examined, and the Claims of the Roman Church compared with the Scriptures, the Councils, and the Testimony of the Popes themselves. Fcap. 8vo. 3s. 6d.

JENKINS (Edward), M.P.

GLANCES AT INNER ENGLAND. A Lecture delivered in the United States and Canada. Crown 8vo. 5s.

GINX'S BABY: His Birth and other Misfortunes. Thirty-fourth Edition. Crown 8vo. 2s.

LUCHMEE AND DILLO. A Story of West Indian Life. 2 vols. Demy 8vo. Illustrated. [Preparing.]

LITTLE HODGE. A Christmas Country Carol. Fourteenth Thousand. With Five Illustrations. Crown 8vo. 5s.
A Cheap Edition in paper covers, price 1s.

LORD BANTAM. Seventh Edition. Crown 8vo. 2s. 6d.

JEVONS (Prof. W. Stanley).

MONEY AND THE MECHANISM OF EXCHANGE. Crown 8vo. 5s.

Being Vol. XVII. of the International Scientific Series.

KAUFMANN (Rev. M.), B.A.

SOCIALISM: Its Nature, its Dangers, and its Remedies considered. Crown 8vo. 7s. 6d.

KEATING (Mrs.)

HONOR BLAKE: The Story of a Plain Woman. 2 vols. Crown 8vo. 21s.

KER (David).

ON THE ROAD TO KHIVA. Illustrated with Photographs of the Country and its Inhabitants, and a copy of the Official Map in use during the Campaign, from the Survey of Captain Leusilin. 1 vol. Post 8vo. 12s.

THE BOY SLAVE IN BOKHARA. A Tale of Central Asia. Crown 8vo. With Illustrations. 5s.

THE WILD HORSEMAN OF THE PAMPAS. Crown 8vo. Illustrated. 5s.

One of Henry S. King and Co.'s Five Shilling Books for the Young.

KING (Alice).

A CLUSTER OF LIVES. Crown 8vo. 7s. 6d.

KING (Mrs. Hamilton).

THE DISCIPLES. A New Poem. Second Edition, with some Notes. Crown 8vo. 7s. 6d.

ASPRONTE, AND OTHER POEMS. Second Edition. Cloth. 4s. 6d.

KINGSFORD (Rev. F. W.), M.A., Vicar of St. Thomas's, Stamford Hill; late Chaplain H. E. I. C. (Bengal Presidency).

HARTHAM CONFERENCES; or, Discussions upon some of the Religious Topics of the Day. "Audi alteram partem." Crown 8vo. 3s. 6d.

KNIGHT (Annette F. C.)

POEMS. Fcap. 8vo. Cloth. 5s.

LACORDAIRE (Rev. Père).

LIFE: Conferences delivered at Toulouse. Crown 8vo. A New and Cheaper Edition. 3s. 6d.

LADY OF LIPARI (The).

A Poem in Three Cantos. Fcap. 8vo. 5s.

LAURIE (J. S.), of the Inner Temple, Barrister-at-Law ; formerly H.M. Inspector of Schools, England ; Assistant Royal Commissioner, Ireland ; Special Commissioner, African Settlement ; Director of Public Instruction, Ceylon.

EDUCATIONAL COURSE OF SECULAR SCHOOL BOOKS FOR INDIA.

The following Works are now ready :—

THE FIRST HINDUSTANI READER. Stiff linen wrapper, 6*d.*

THE SECOND HINDUSTANI READER. Stiff linen wrapper, 6*d.*

GEOGRAPHY OF INDIA ; with Maps and Historical Appendix, tracing the growth of the British Empire in Hindustan. 128 pp. fcap. 8vo. Cloth. 1*s.* 6*d.*

In the Press :—

ELEMENTARY GEOGRAPHY OF INDIA.

FACTS AND FEATURES OF INDIAN HISTORY, in a series of alternating Reading Lessons and Memory Exercises.

LAYMANN (Captain), Instructor of Tactics at the Military College, Neisse.

THE FRONTAL ATTACK OF INFANTRY. Translated by Colonel Edward Newdigate. Crown 8vo. 2*s.* 6*d.*

L. D. S.

LETTERS FROM CHINA AND JAPAN. 1 vol. Crown 8vo., with Illustrated Title-page. 7*s.* 6*d.*

LEANDER (Richard).

FANTASTIC STORIES. Translated from the German by Paulina B. Granville. With Eight full-page Illustrations by M. E. Fraser-Tytler. Crown 8vo. 5*s.*

One of Henry S. King and Co.'s Five Shilling Books for the Young.

LEATHES (Rev. Stanley), M.A.

THE GOSPEL ITS OWN WITNESS. Being the Hulsean Lectures for 1873. 1 vol. Crown 8vo. 5*s.*

LEE (Rev. Frederick George), D.C.L.

THE OTHER WORLD ; or, Glimpses of the Supernatural. Being Facts, Records, and Traditions, relating to Dreams, Omens, Miraculous Occurrences, Apparitions, Wraiths, Warnings, Second-sight, Necromancy, Witchcraft, etc. 2 vols. A New Edition. Crown 8vo. 15*s.*

LEE (Holme).

HER TITLE OF HONOUR. A Book for Girls. New Edition. Crown 8vo. With a Frontispiece. 5s.

LENOIR (J).

FAYOUM; or, Artists in Egypt. A Tour with M. Gérôme and others. Crown 8vo. A New and Cheaper Edition. With 13 Illustrations. 3s. 6d.

LISTADO (J. T.)

CIVIL SERVICE. A Novel. 2 vols. Crown 8vo.

LORIMER (Peter), D.D.

JOHN KNOX AND THE CHURCH OF ENGLAND: His work in her Pulpit and his influence upon her Liturgy, Articles, and Parties. Demy 8vo. 12s.

LOVER (Samuel), R.H.A.

THE LIFE OF SAMUEL LOVER, R.H.A.; Artistic, Literary, and Musical. With Selections from his Unpublished Papers and Correspondence. By Bayle Bernard. 2 vols. Post 8vo. With a Portrait. 21s.

LOWER (Mark Antony), M.A., F.S.A.

WAYSIDE NOTES IN SCANDINAVIA. Being Notes of Travel in the North of Europe. Crown 8vo. 9s.

LYONS (R. T.), Assistant-Surgeon, Bengal Army.

A TREATISE ON RELAPSING FEVER. Post 8vo. 7s. 6d.

MACAULAY (James), M.A., M.D., Edin.

IRELAND. A Tour of Observation, with Remarks on Irish Public Questions. Crown 8vo. A New and Cheaper Edition. 3s. 6d.

MAC CARTHY (Denis Florence).

CALDERON'S DRAMAS. Translated from the Spanish. Post 8vo. Cloth, gilt edges. 10s.

MAC DONALD (George).

GUTTA-PERCHA WILLIE, THE WORKING GENIUS. With Nine Illustrations by Arthur Hughes. Second Edition. Crown 8vo. 3s. 6d.

One of Henry S. King and Co.'s Three and Sixpenny Books for the Young.

MALCOLM. A Novel. Second Edition. 3 vols. Crown 8vo.

ST. GEORGE AND ST. MICHAEL. 3 vols. Crown 8vo.

MY SISTER ROSALIND. By the author of "Christina North." A Novel. 2 vols.

MAC KENNA (Stephen J.)

PLUCKY FELLOWS. A Book for Boys. With Six Illustrations. Second Edition. Crown 8vo. 3s. 6d.

One of Henry S. King and Co.'s Three and Sixpenny Books for the Young.

AT SCHOOL WITH AN OLD DRAGOON. Crown 8vo. With Six Illustrations. 5s.

One of Henry S. King and Co.'s Five Shilling Books for the Young.

MACLACHLAN (Archibald Neil Campbell), M.A.

WILLIAM AUGUSTUS, DUKE OF CUMBERLAND; being a Sketch of his Military Life and Character, chiefly as exhibited in the General Orders of his Royal Highness, 1745—1747. Post 8vo. With Illustrations. 15s.

MAIR (R. S.), M.D., F.R.C.S.E., late Deputy Coroner of Madras.

THE MEDICAL GUIDE FOR ANGLO-INDIANS. Being a Compendium of Advice to Europeans in India, relating to the Preservation and Regulation of Health. With a Supplement on the Management of Children in India. Crown 8vo. Limp cloth. 3s. 6d.

MANNING (The Most Rev. Archbishop).

ESSAYS ON RELIGION AND LITERATURE. By various Writers. Demy 8vo. 10s. 6d.

CONTENTS:—The Philosophy of Christianity—Mystic Elements of Religion—Controversy with the Agnostics—A Reasoning Thought—Darwinism brought to Book—Mr. Mill on Liberty of the Press—Christianity in relation to Society—The Religious Condition of Germany—The Philosophy of Bacon—Catholic Laymen and Scholastic Philosophy.

MAREY (E. J.)

ANIMAL MECHANICS. A Treatise on Terrestrial and Aerial Locomotion. Second Edition. With 117 Illustrations. 5s.
Volume XI. of the International Scientific Series.

MARKEWITCH (B.)

THE NEGLECTED QUESTION. Translated from the Russian, by the Princess Ourousoff, and dedicated by Express Permission to Her Imperial and Royal Highness Marie Alexandrovna, the Duchess of Edinburgh. 2 vols. Crown 8vo. 14s.

MARRIOTT (Maj.-Gen. W. F.), C.S.I.

A GRAMMAR OF POLITICAL ECONOMY. Crown 8vo. 6s.

MARSHALL (Hamilton).

THE STORY OF SIR EDWARD'S WIFE. A Novel. 1 vol.
Crown 8vo. 10s. 6d.

MARZIALS (Theophile).

THE GALLERY OF PIGEONS, and other Poems. Crown 8vo.
4s. 6d.

MASTERMAN (J.)

HALF-A-DOZEN DAUGHTERS. Crown 8vo. With a Frontis-
piece. 3s. 6d.

One of the Cornhill Library of Fiction.

MAUDSLEY (Dr. Henry).

RESPONSIBILITY IN MENTAL DISEASE. Second Edition. 5s.
Vol. VIII. of the International Scientific Series.

MAUGHAN (William Charles).

THE ALPS OF ARABIA; or, Travels through Egypt, Sinai,
Arabia, and the Holy Land. Demy 8vo. With Map. A New
and Cheaper Edition. 5s.

MAURICE (C. Edmund).

LIVES OF ENGLISH POPULAR LEADERS. No. 1.—STEPHEN
LANGTON. Crown 8vo. 7s. 6d.

No. 2.—TYLER, BALL, and OLDCASTLE. Crown 8vo. 7s. 6d.

MEDLEY (Lieut.-Col. J. G.), Royal Engineers.

AN AUTUMN TOUR IN THE UNITED STATES AND CANADA.
Crown 8vo. 5s.

MENZIES (Sutherland).

MEMOIRS OF DISTINGUISHED WOMEN. 2 vols. Post 8vo.

MEUGENS (Miss N. R.)

LITTLE MINNIE'S TROUBLES. An Every-day Chronicle,
Illustrated by Mr. W. H. Hughes. Fcap. 3s. 6d.
A Simple Chronicle of a Child's Life.

MICKLETHWAITE (J. T.), F.S.A.

MODERN PARISH CHURCHES: Their Plan, Design, and
Furniture. Crown 8vo. 7s. 6d.

MIRUS (Major-General von).

CAVALRY FIELD DUTY. Translated by Captain Frank S.
Russell, 14th (King's) Hussars. Crown 8vo. Cloth limp. 7s. 6d.
This work is one of Henry S. King and Co.'s Military Series.

MOORE (Rev. Daniel), M.A.

CHRIST AND HIS CHURCH. A Course of Lent Lectures, delivered in the Parish Church of Holy Trinity, Paddington. By the author of "The Age and the Gospel: Hulsean Lectures," etc. Crown 8vo. 3s. 6d.

MOORE (Rev. Thomas), Vicar of Christ Church, Chesham.

SERMONETTES: on Synonymous Texts, taken from the Bible and Book of Common Prayer, for the Study, Family Reading, and Private Devotion. Small Crown 8vo. 4s. 6d.

MORELL (J. R.)

EUCLID SIMPLIFIED IN METHOD AND LANGUAGE. Being a Manual of Geometry on the French System. Fcap. 8vo. 2s. 6d.

MORICE (Rev. F. D.), M.A., Fellow of Queen's College, Oxford.

THE OLYMPIAN AND PYTHIAN ODES OF PINDAR. A New Translation in English Verse.

MORLEY (Susan).

AILEEN FERRERS. A Novel. 2 vols. Crown 8vo.

THROSTLETHWAITE. 3 vols. Crown 8vo.

MORSE (Edward S.), Ph. D., late Professor of Comparative Anatomy and Zoology in Bowdoin College.

FIRST BOOK OF ZOOLOGY. With numerous Illustrations.

MOSTYN (Sydney).

PERPLEXITY. A Novel. 3 vols. Crown 8vo.

MUSGRAVE.

STUDIES IN POLITICAL ECONOMY. 6s.

NAAKÈ (John T.), of the British Museum.

SLAVONIC FAIRY TALES. From Russian, Servian, Polish, and Bohemian Sources. With Four Illustrations. Crown 8vo. 5s.

NEWMAN (John Henry), D.D.

CHARACTERISTICS FROM THE WRITINGS OF DR. J. H. NEWMAN. Being Selections, Personal, Historical, Philosophical, and Religious, from his various Works. Arranged with the Author's personal approval. Second Edition. Crown 8vo. With Portrait. 6s.

NEWMAN (Mrs.)

TOO LATE. A Novel. 2 vols. Crown 8vo.

NOBLE (James Ashcroft).

THE PELICAN PAPERS. Reminiscences and Remains of a Dweller in the Wilderness. Crown 8vo. 6s.

NORMAN PEOPLE (The).

THE NORMAN PEOPLE, and their Existing Descendants in the British Dominions and the United States of America. One handsome volume. 8vo. 21s.

NORRIS (Rev. A.)

THE INNER AND OUTER LIFE POEMS. Fcap. 8vo. 6s.

NOTREGE (John), A.M.

THE SPIRITUAL FUNCTION OF A PRESBYTER IN THE CHURCH OF ENGLAND. Crown 8vo. Red edges. 3s. 6d.

ORIENTAL SPORTING MAGAZINE (The).

THE ORIENTAL SPORTING MAGAZINE. A Reprint of the first 5 Volumes, in 2 Volumes. Demy 8vo. 28s.

PAGE (H. A.)

NATHANIEL HAWTHORNE, A MEMOIR OF, with Stories now first published in this country. Large post 8vo. 7s. 6d.

PAGE (Capt. S. Flood).

DISCIPLINE AND DRILL. Four Lectures delivered to the London Scottish Rifle Volunteers. Cheaper Edition. Crown 8vo. 1s.

PALGRAVE (W. Gifford).

HERMANN AGHA. An Eastern Narrative. 2 vols. Crown 8vo. Cloth, extra gilt. 18s.

PARKER (Joseph), D.D.

THE PARACLETE: An Essay on the Personality and Ministry of the Holy Ghost, with some reference to current discussions. Demy 8vo. 12s.

PARR (Harriett).

ECHOES OF A FAMOUS YEAR. Crown 8vo. 8s. 6d.

PAUL (C. Kegan).

GOETHE'S FAUST. A New Translation in Rime. Crown 8vo. 6s.

WILLIAM GODWIN: HIS FRIENDS AND CONTEMPORARIES. 2 vols. With Portraits. Demy 8vo.

PAYNE (John).

SONGS OF LIFE AND DEATH. Crown 8vo. 5s.

PAYNE (Professor).

LECTURES ON EDUCATION. 6*d.* each.

- I. Pestalozzi: the Influence of His Principles and Practice.
- II. Fröbel and the Kindergarten System. Second Edition.
- III. The Science and Art of Education.
- IV. The True Foundation of Science Teaching.

PELLETAN (Eugène).

THE DESERT PASTOR, JEAN JAROUSSEAU. Translated from the French. By Colonel E. P. De L'Hoste. With an Engraved Frontispiece. Fcap. 8vo. New Edition. 3*s.* 6*d.*

PENRICE (Major J.), B.A.

A DICTIONARY AND GLOSSARY OF THE KO-RAN. With copious Grammatical References and Explanations of the Text. 4to. 2*l*s.

PERCEVAL (Rev. P.)

TAMIL PROVERBS, WITH THEIR ENGLISH TRANSLATION Containing upwards of Six Thousand Proverbs. Third Edition. 8vo. Sewed. 9*s.*

PERRIER (Amelia).

A WINTER IN MOROCCO. With Four Illustrations. Crown 8vo. A New and Cheaper Edition. 3*s.* 6*d.*

A GOOD MATCH. A Novel. 2 vols. Crown 8vo.

PESCHEL (Dr.)

MANKIND: A Scientific Study of the Races and Distribution of Man, considered in their Bodily Variations, Languages, Occupations, and Religions.

PETTIGREW (J. B.), M.D., F.R.S.

ANIMAL LOCOMOTION; or, Walking, Swimming, and Flying. Second Edition. With 119 Illustrations. 5*s.*

Volume VII. of the International Scientific Series.

PIGGOT (John), F.S.A, F.R.G.S.

PERSIA—ANCIENT AND MODERN. Post 8vo. 10*s.* 6*d.*

POUSHKIN (Alexander Serguevitch).

RUSSIAN ROMANCE. Translated from the Tales of Belkin, etc. By Mrs. J. Buchan Telfer (*née* Mouravieff). Cr. 8vo. 7*s.* 6*d.*

POWER (Harriet).

OUR INVALIDS: HOW SHALL WE EMPLOY AND AMUSE THEM? Fcap 8vo. 2s. 6d.

POWLETT (Lieut. Norton), Royal Artillery.

EASTERN LEGENDS AND STORIES IN ENGLISH VERSE.
Crown 8vo. 5s.

PRESBYTER.

UNFOLDINGS OF CHRISTIAN HOPE. An Essay showing that the Doctrine contained in the Damnatory Clauses of the Creed commonly called Athanasian is unscriptural. Small crown 8vo. Cloth. 4s. 6d.

PRICE (Prof. Bonamy).

CURRENCY AND BANKING. One Vol. Crown 8vo. 6s.

PROCTOR (Richard A.)

OUR PLACE AMONG INFINITIES. A Series of Essays contrasting our little abode in space and time with the Infinities around us. To which are added "Essays on Astrology," and "The Jewish Sabbath." 6s.

THE EXPANSE OF HEAVEN. A Series of Essays on the Wonders of the Firmament. With a Frontispiece. Second Edition. Crown 8vo. 6s.

RANKING (B. Montgomerie).

STREAMS FROM HIDDEN SOURCES. Crown 8vo. 6s.

READY-MONEY MORTIBOY.

READY-MONEY MORTIBOY. A Matter-of-Fact Story. Crown 8vo. With frontispiece. 3s. 6d.

This is one of the volumes of the Cornhill Library of Fiction.

REANEY (Mrs. G. S.)

WAKING AND WORKING; OR, FROM GIRLHOOD TO WOMANHOOD. With a Frontispiece. Crown 8vo. 5s.

One of Henry S. King and Co.'s Five Shilling Books for the Young.

SUNBEAM WILLIE, AND OTHER STORIES, for Home Reading and Cottage Meetings. Small square, uniform with "Lost Gip," etc. 3 Illustrations. 1s. 6d.

REGINALD BRAMBLE.

REGINALD BRAMBLE. A Cynic of the Nineteenth Century. An Autobiography. 1 vol. Crown 8vo. 10s. 6d.

REID (T. Wemyss).

CABINET PORTRAITS. Biographical Sketches of Statesmen of the Day. 1 vol. Crown 8vo. 7s. 6d.

RHOADES (James).

TIMOLEON. A Dramatic Poem. Fcap. 8vo. 5s.

RIBOT (Professor Th.)

CONTEMPORARY ENGLISH PSYCHOLOGY. La. post 8vo. 9s.
An analysis of the views and opinions of the following metaphysicians, as expressed in their writings:—James Mill, Alexander Bain, John Stuart Mill, George H. Lewes, Herbert Spencer, Samuel Bailey.

HEREDITY: A Psychological Study on its Phenomena, its Laws, its Causes, and its Consequences. 1 vol. Large crown 8vo. 9s.

ROBERTSON (The Late Rev. F. W.), M.A.

THE LATE REV. F. W. ROBERTSON, M.A., LIFE AND LETTERS OF. Edited by the Rev. Stopford Brooke, M.A., Chaplain in Ordinary to the Queen.

- I. 2 vols., uniform with the Sermons. With Steel Portrait. 7s. 6d.
- II. Library Edition, in Demy 8vo. with Two Steel Portraits. 12s.
- III. A Popular Edition, in 1 vol. 6s.

New and Cheaper Editions :—

SERMONS.

Vol. I. Small crown 8vo. 3s. 6d.

Vol. II. Small crown 8vo. 3s. 6d.

Vol. III. Small crown 8vo. 3s. 6d.

Vol. IV. Small crown 8vo. 3s. 6d.

EXPOSITORY LECTURES ON ST. PAUL'S EPISTLE TO THE CORINTHIANS. Small crown 8vo. 5s.

AN ANALYSIS OF MR. TENNYSON'S "IN MEMORIAM." (Dedicated by Permission to the Poet-Laureate.) Fcap. 8vo. 2s.

THE EDUCATION OF THE HUMAN RACE. Translated from the German of Gotthold Ephraim Lessing. Fcap. 8vo. 2s. 6d.

The above Works can also be had bound in half morocco.

* * * A Portrait of the late Rev. F. W. Robertson, mounted for framing, can be had, price 2s. 6d.

NOTES ON GENESIS. Uniform with the Sermons.

LECTURES AND ADDRESSES, with other literary remains. A New Edition. 5s.

ROSS (Mrs. Ellen), ("Nelsie Brook.")

DADDY'S PET. A Sketch from Humble Life. Square crown 8vo. Uniform with "Lost Gip." With Six Illustrations. 1s.

ROXBURGHE LOTHIAN.

DANTE AND BEATRICE FROM 1282 TO 1290. A Romance. 1 vol. Post 8vo. Cloth.

RUSSELL (William Clark).

MEMOIRS OF MRS. LÆTITIA BOOTHBY. Crown 8vo. 7s. 6d.

RUSSELL (E. R.)

IRVING AS HAMLET. Demy 8vo. Second Edition. Sewed. 1s.

SADLER (S. W.), R.N., Author of "Marshall Vavasour."

THE AFRICAN CRUISER. A Midshipman's Adventures on the West Coast. A Book for Boys. With Three Illustrations. Second Edition. Crown 8vo. 3s. 6d.

One of Henry S. King and Co.'s Three and Sixpenny Books for the Young.

SAMAROW (Gregor).

FOR SCEPTRE AND CROWN. A Romance of the Present Time. Translated by Fanny Wormald. 2 vols. Crown 8vo. 15s.

SAUNDERS (Katherine).

THE HIGH MILLS. A Novel. 3 vols. Crown 8vo.

GIDEON'S ROCK, and other Stories. 1 vol. Crown 8vo. 6s.

JOAN MERRYWEATHER, and other Stories. 1 vol. Crown 8vo. 6s.

MARGARET AND ELIZABETH. A Story of the Sea. 1 vol. Crown 8vo. 6s.

TOO LONG UNTOLD, and other Stories. 2 vols. Crown 8vo.

CONTENTS:—Too Long Untold—The Harpers of Men-y-don—Ida's Story—Little Missy—The Shaken Nest.

SAUNDERS (John).

HIRELL. Crown 8vo. With Frontispiece. 3s. 6d.

ABEL DRAKE'S WIFE. Crown 8vo. With Frontispiece. 3s. 6d.

These works form separate volumes of the Cornhill Library of Fiction.

ISRAEL MORT, OVERMAN. The Story of the Mine. 3 vols. Crown 8vo.

SCHELL (Major von).

THE OPERATIONS OF THE FIRST ARMY UNDER GEN. VON GOEBEN. Translated by Col. C. H. von Wright. Four Maps. Demy 8vo. 9s.

THE OPERATIONS OF THE FIRST ARMY UNDER GEN. VON STEINMETZ. Translated by Captain E. O. Hollist. Demy 8vo. 10s. 6d.

These works form separate volumes of Henry S. King and Co.'s Military Series.

SCHERFF (Major W. von).

STUDIES IN THE NEW INFANTRY TACTICS. Parts I. and II. Translated from the German by Colonel Lumley Graham. Demy 8vo. 7s. 6d.

This work is one of Henry S. King and Co.'s Military Series.

SCHMIDT (Prof. Oscar), Strasburg University.

THE DOCTRINE OF DESCENT AND DARWINISM. Second Edition. 26 Illustrations. 5s.

Being Vol. XII. of the International Scientific Series.

HANDBOOK OF COMPARATIVE ANATOMY. Crown 8vo.

SCOTT (Patrick).

THE DREAM AND THE DEED, and other Poems. Fcap. 8vo. 5s.

SEEKING HIS FORTUNE, and other Stories.

SEEKING HIS FORTUNE, and other Stories. Crown 8vo. With Four Illustrations. 3s. 6d.

One of Henry S. King and Co.'s Three and Sixpenny Books for the Young.

SENIOR (Nassau William).

ALEXIS DE TOCQUEVILLE. Correspondence and Conversations with Nassau W. Senior, from 1833 to 1859. Edited by M. C. M. Simpson. 2 vols. Large post 8vo. 21s.

JOURNALS KEPT IN FRANCE AND ITALY. From 1848 to 1852. With a Sketch of the Revolution of 1848. Edited by his Daughter, M. C. M. Simpson. 2 vols. Post 8vo. 24s.

SEVEN AUTUMN LEAVES.

SEVEN AUTUMN LEAVES FROM FAIRYLAND. Illustrated with 9 Etchings. Square crown 8vo. 3s. 6d.

SHADWELL (Major-General), C.B.

MOUNTAIN WARFARE. Illustrated by the Campaign of 1799 in Switzerland. Being a Translation of the Swiss Narrative compiled from the Works of the Archduke Charles, Jomini, and others. Also of Notes by General H. Dufour on the Campaign of the Valtelline in 1635. With Appendix, Maps, and Introductory Remarks. Demy 8vo. 16s.

SHELDON (Philip).

WOMAN'S A RIDDLE; or, Baby Warmstrey. A Novel. 3 vols. Crown 8vo.

SHERMAN (Gen. W. T.)

MEMOIRS OF GEN. W. T. SHERMAN, Commander of the Federal Forces in the American Civil War. By Himself. 2 vols. Demy 8vo. With Map. 24s. *Copyright English Edition.*

SHELLEY (Lady).

SHELLEY MEMORIALS FROM AUTHENTIC SOURCES. With (now first printed) an Essay on Christianity by Percy Bysshe Shelley. Third Edition. Crown 8vo. With Portrait. 5s.

SHIPLEY (Rev. Orby), M.A.

STUDIES IN MODERN PROBLEMS. By various Writers. Crown 8vo. 2 vols. 5s. each.

CONTENTS.—VOL. I.

Sacramental Confession.
Abolition of the Thirty-nine Articles. Part I.
The Sanctity of Marriage.
Creation and Modern Science.

Retreats for Persons Living in the World.
Catholic and Protestant.
The Bishops on Confession in the Church of England.

CONTENTS.—VOL. II.

Some Principles of Christian Ceremonial.
A Layman's View of Confession of Sin to a Priest. Parts I. and II.
Reservation of the Blessed Sacrament.

Missions and Preaching Orders.
Abolition of the Thirty-nine Articles. Part II.
The First Liturgy of Edward VI. and our own office contrasted and compared.

SMEDLEY (M. B.)

BOARDING-OUT AND PAUPER SCHOOLS FOR GIRLS. Crown 8vo. 3s. 6d.

SMITH (Edward) M.D., LL.B., F.R.S.

HEALTH AND DISEASE, as influenced by the Daily, Seasonal, and other Cyclical Changes in the Human System. A New Edition. 7s. 6d.

FOODS. Third Edition. Profusely Illustrated. 5s.
Volume III. of the International Scientific Series.

PRACTICAL DIETARY FOR FAMILIES, SCHOOLS, AND THE LABOURING CLASSES. A New Edition. 3s. 6d.

CONSUMPTION IN ITS EARLY AND REMEDIABLE STAGES. A New Edition. 7s. 6d.

SMITH (Hubert).

TENT LIFE WITH ENGLISH GIPSIES IN NORWAY. With Five full-page Engravings and Thirty-one smaller Illustrations by Whymper and others, and Map of the Country showing Routes. Second Edition. Revised and Corrected. 8vo. 21s.

SONGS FOR MUSIC.

SONGS FOR MUSIC. By Four Friends. Square crown 8vo. 5s.
Containing Songs by Reginald A. Gatty, Stephen H. Gatty, Greville J. Chester, and Juliana H. Ewing.

SOME TIME IN IRELAND.

SOME TIME IN IRELAND. A Recollection. Crown 8vo. 7s. 6d.

SONGS OF TWO WORLDS.

SONGS OF TWO WORLDS. By a New Writer. First Series. Second Edition. Fcap. 8vo. 5s.

SONGS OF TWO WORLDS. By a New Writer. Second Series. Second Edition. Fcap. 8vo. 5s.

SONGS OF TWO WORLDS. By a New Writer. Third Series. Fcap. 8vo. 5s.

SPENCER (HERBERT).

THE STUDY OF SOCIOLOGY. Fourth Edition. Crown 8vo. 5s.
Volume V. of the International Scientific Series.

STEVENSON (Rev. W. Fleming).

HYMNS FOR THE CHURCH AND HOME. Selected and Edited by the Rev. W. Fleming Stevenson.

The most complete Hymn Book published.

The Hymn Book consists of Three Parts:—I. For Public Worship.—II. For Family and Private Worship.—III. For Children.

* * * Published in various forms and prices, the latter ranging from 8d. to 6s. Lists and full particulars will be furnished on application to the Publishers.

STEWART (Professor Balfour).

ON THE CONSERVATION OF ENERGY. Third Edition.
With Fourteen Engravings. 5s.
Volume VI. of the International Scientific Series.

STRETTON (Hesba). Author of "Jessica's First Prayer."

CASSY. Twenty-first Thousand. With Six Illustrations.
Square crown 8vo. 1s. 6d.

THE KING'S SERVANTS. With Eight Illustrations. Twenty-sixth Thousand. Square crown 8vo. 1s. 6d.

LOST GIP. Thirty-eighth Thousand. With Six Illustrations.
Square crown 8vo. 1s. 6d.

* * * *Also a handsomely-bound Edition, with Twelve Illustrations, price 2s. 6d.*

THE WONDERFUL LIFE. Seventh Thousand. Fcap. 8vo.
2s. 6d.

HESTER MORLEY'S PROMISE. 3 vols. Crown 8vo.

THE DOCTOR'S DILEMMA. 3 vols. Crown 8vo.

FRIENDS TILL DEATH. With Illustration. 6d.

STUBBS (Major Francis W.), Royal (late Bengal)
Artillery.

THE REGIMENT OF BENGAL ARTILLERY: The History of
its Organization, Equipment, and War Services. With Maps
and Plans. 2 vols. 8vo. [Preparing.

SULLY (James).

SENSATION AND INTUITION. Demy 8vo. 10s. 6d.

TAYLOR (Rev. J. W. Augustus), M.A.

POEMS. Fcap. 8vo. 5s.

TAYLOR (Sir Henry).

EDWIN THE FAIR AND ISAAC COMNENUS. 3s. 6d.

A SICILIAN SUMMER AND OTHER POEMS. 3s. 6d.

TAYLOR (Colonel Meadows), C.S.I., M.R.I.A.

SEETA. A Novel. 3 vols.

THE CONFESSIONS OF A THUG.

TARA: a Mahratta Tale.

RALPH DARNELL.

TIPPOO SULTAN.

A NOBLE QUEEN. 3 vols. Crown 8vo.

New and Cheaper Edition in one vol. crown 8vo. with Frontis-
piece. Each 6s.

TENNYSON (Alfred).**QUEEN MARY.** A Drama. Crown 8vo. 6s.**TENNYSON'S (Alfred) Works.** Cabinet Edition. Ten Volumes. Each with Portrait. 2s. 6d.

CABINET EDITION. 10 vols. Complete in handsome Ornamental Case. 28s.

TENNYSON'S (Alfred) Works. Author's Edition. Complete in Five Volumes. Cloth gilt, 6s. each; half-morocco, Roxburgh style, 7s. 6d. each.**EARLY POEMS, and ENGLISH IDYLLS.—VOL. I.****LOCKSLEY HALL, LUCRETIVS, and other Poems.—VOL. II.****THE IDYLLS OF THE KING (Complete).—VOL. III.****THE PRINCESS, and MAUD.—VOL. IV.****ENOCH ARDEN, and IN MEMORIAM.—VOL. V.****TENNYSON'S IDYLLS OF THE KING, and other Poems.** Illustrated by Julia Margaret Cameron. 1 vol. Folio. Half-bound morocco, cloth sides. Six Guineas.**TENNYSON'S (Alfred) Works.** Original Editions.**POEMS.** Small 8vo. 6s.**MAUD, and other Poems.** Small 8vo. 3s. 6d.**THE PRINCESS.** Small 8vo. 3s. 6d.**IDYLLS OF THE KING.** Small 8vo. 5s.**IDYLLS OF THE KING.** Collected. Small 8vo. 7s.**THE HOLY GRAIL, and other Poems.** Small 8vo. 4s. 6d.**GARETH AND LYNETTE.** Small 8vo. 3s.**ENOCH ARDEN, etc.** Small 8vo. 3s. 6d.**SELECTIONS FROM THE ABOVE WORKS.** Square 8vo. Cloth, 3s. 6d. Cloth gilt, extra, 4s.**SONGS FROM THE ABOVE WORKS.** Square 8vo. Cloth extra, 3s. 6d.**IN MEMORIAM.** Small 8vo. 4s.**LIBRARY EDITION.** In 6 vols. 8vo. 10s. 6d. each.**POCKET VOLUME EDITION.** 11 vols. In neat case, 31s. 6d.

Ditto, ditto. Extra cloth gilt, in case, 35s.

POEMS. Illustrated Edition. 4to. 25s.

THOMAS (Moy).

A FIGHT FOR LIFE. Crown 8vo. With Frontispiece. 3s. 6d.
This is one of the volumes of the Cornhill Library of Fiction.

THOMSON (J. T.), F.R.G.S.

HAKAYIT ABDULLA. The Autobiography of a Malay Munshi, between the years 1808 and 1843. Demy 8vo. 12s.

THOMPSON (A. C.)

PRELUDES. A Volume of Poems. Illustrated by Elizabeth Thompson (Painter of "The Roll Call"). 8vo. 7s. 6d.

THOMPSON (Rev. A. S.), British Chaplain at St. Petersburg.

HOME WORDS FOR WANDERERS. A Volume of Sermons. Crown 8vo. 6s.

THOUGHTS IN VERSE. Small crown 8vo. 1s. 6d.

THRING (Rev. Godfrey), B.A.

HYMNS AND SACRED LYRICS. 1 vol. Fcap. 8vo. 5s.

TODD (Herbert), M.A.

ARVAN; or, The Story of the Sword. A Poem. Crown 8vo. 7s. 6d.

TRAHERNE (Mrs. Arthur).

THE ROMANTIC ANNALS OF A NAVAL FAMILY. Crown 8vo. A New and Cheaper Edition. 5s.

TRAVERS (Mar.)

THE SPINSTERS OF BLATCHINGTON. A Novel. 2 vols. Crown 8vo.

TREVANDRUM OBSERVATIONS.

OBSERVATIONS OF MAGNETIC DECLINATION MADE AT TREVANDRUM AND AGUSTIA MALLEY in the Observatories of his Highness the Maharajah of Travancore, G.C.S.I., in the Years 1852 to 1860. Being Trevandrum Magnetical Observations, Volume I. Discussed and Edited by John Allan Brown, F.R.S., late Director of the Observatories. With an Appendix. Imp. 4to. Cloth. £3 3s.

* * *The Appendix, containing Reports on the Observatories and on the Public Museum, Public Park, and Gardens at Trevandrum, pp. xii.-116, may be had separately.* 21s.

TURNER (Rev. Charles).

SONNETS, LYRICS, AND TRANSLATIONS. Crown 8vo. 4s. 6d.

TYNDALL (J.), LL.D., F.R.S.

THE FORMS OF WATER IN CLOUDS AND RIVERS, ICE AND GLACIERS. With Twenty-six Illustrations. Fourth Edition. Crown 8vo. 5s.

Volume I. of the International Scientific Series.

UMBRA OXONIENSIS.

RESULTS OF THE EXPOSTULATION OF THE RIGHT HONOURABLE W. E. GLADSTONE, in their Relation to the Unity of Roman Catholicism. Large fcap. 8vo. 5s.

UPTON (Roger D.), Captain late 9th Royal Lancers.

NEWMARKET AND ARABIA. An Examination of the Descent of Racers and Coursers. With Pedigrees and Frontispiece. Post 8vo. 9s.

VAMBERY (Prof. Arminius), of the University of Pesth.

BOKHARA : Its History and Conquest. Demy 8vo. 18s.

VANESSA. By the Author of "Thomasina," etc. A Novel. Second Edition. 2 vols. Crown 8vo.

VAUGHAN (Rev. C. J.), D.D.

WORDS OF HOPE FROM THE PULPIT OF THE TEMPLE CHURCH. Third Edition. Crown 8vo. 5s.

THE SOLIDITY OF TRUE RELIGION, and other Sermons Preached in London during the Election and Mission Week, February, 1874. Crown 8vo. 3s. 6d.

FORGET THINE OWN PEOPLE. An Appeal for Missions. Crown 8vo. 3s. 6d.

THE YOUNG LIFE EQUIPPING ITSELF FOR GOD'S SERVICE. Being Four Sermons Preached before the University of Cambridge, in November, 1872. Fourth Edition. Crown 8vo. 3s. 6d.

VINCENT (Capt. C. E. H.), late Royal Welsh Fusiliers.

ELEMENTARY MILITARY GEOGRAPHY, RECONNOITRING, AND SKETCHING. Compiled for Non-Commissioned Officers and Soldiers of all Arms. Square crown 8vo. 2s. 6d.

RUSSIA'S ADVANCE EASTWARD. Based on the Official Reports of Lieutenant Hugo Stumm, German Military Attaché to the Khivan Expedition. With Map. Crown 8vo. 6s.

VIZCAYA ; or, Life in the Land of the Carlists.

VIZCAYA ; or, Life in the Land of the Carlists at the Outbreak of the Insurrection, with some Account of the Iron Mines and other Characteristics of the Country. With a Map and Eight Illustrations. Crown 8vo. 9s.

VOGEL (Prof.), Polytechnic Academy of Berlin.

THE CHEMICAL EFFECTS OF LIGHT AND PHOTOGRAPHY, in their application to Art, Science, and Industry. Third Edition. The translation thoroughly revised. With 100 Illustrations, including some beautiful Specimens of Photography. 5s.

Volume XV. of the International Scientific Series.

VYNER (Lady Mary).

EVERY DAY A PORTION. Adapted from the Bible and the Prayer Book, for the Private Devotions of those living in Widowhood. Collected and Edited by Lady Mary Vyner. Square crown 8vo. Elegantly bound. 5s.

WAITING FOR TIDINGS.

WAITING FOR TIDINGS. By the Author of "White and Black." 3 vols. Crown 8vo.

WARTENSLEBEN (Count Hermann von), Colonel in the Prussian General Staff.

THE OPERATIONS OF THE SOUTH ARMY IN JANUARY AND FEBRUARY, 1871. Compiled from the Official War Documents of the Head-quarters of the Southern Army. Translated by Colonel C. H. von Wright. With Maps. Demy 8vo. 6s.

THE OPERATIONS OF THE FIRST ARMY UNDER GEN. VON MANTEUFFEL. Translated by Colonel C. H. von Wright. Uniform with the above. Demy 8vo. 9s.

These works form separate volumes of Henry S. King and Co.'s Military Series.

WEDMORE (Frederick).

TWO GIRLS. 2 vols. Crown 8vo.

WELLS (Captain John C.), R.N.

SPITZBERGEN—THE GATEWAY TO THE POLYNIA ; or, A Voyage to Spitzbergen. With numerous Illustrations by Whympers and others, and Map. 8vo. 21s.

WHAT 'TIS TO LOVE. By the Author of "Flora Adair," "The Value of Fostertown." 3 vols. Crown 8vo.

WHITNEY (William Dwight). Professor of Sanskrit and Comparative Philology in Yale College, New Haven.

THE LIFE AND GROWTH OF LANGUAGE. Second Edition. Crown 8vo. 5s. *Copyright Edition.*
Volume XVI. of the International Scientific Series.

WHITTLE (J. Lowry), A.M., Trin. Coll., Dublin.

CATHOLICISM AND THE VATICAN. With a Narrative of the Old Catholic Congress at Munich. Second Edition. Crown 8vo. 4s. 6d.

WILBERFORCE (Henry W.)

THE CHURCH AND THE EMPIRES. Historical Periods. Preceded by a Memoir of the Author by John Henry Newman, D.D., of the Oratory. With Portrait. Post 8vo. 10s. 6d.

WILKINSON (T. Lean).

SHORT LECTURES ON THE LAND LAWS. Delivered before the Working Men's College. Crown 8vo. 2s.

WILLIAMS (Rev. Rowland), D.D.

LIFE AND LETTERS OF ROWLAND WILLIAMS, D.D., with Selections from his Note-books. Edited by Mrs. Rowland Williams. With a Photographic Portrait. 2 vols. Large post 8vo. 24s.

WILLOUGHBY (The Hon. Mrs.)

ON THE NORTH WIND—THISTLEDOWN. A Volume of Poems. Elegantly bound. Small crown 8vo. 7s. 6d.

WILSON (H. Schütz).

STUDIES AND ROMANCES. Crown 8vo. 7s. 6d.

WINTERBOTHAM (Rev. R.), M.A., B.Sc.

SERMONS AND EXPOSITIONS. Crown 8vo. 7s. 6d.

WOOD (C. F.)

A YACHTING CRUISE IN THE SOUTH SEAS. Demy 8vo. With Six Photographic Illustrations. 7s. 6d.

WRIGHT (Rev. W.), of Stoke Bishop, Bristol.

MAN AND ANIMALS: A Sermon. Crown 8vo. Stitched in wrapper. 1s.

WAITING FOR THE LIGHT, AND OTHER SERMONS. Crown 8vo. 6s.

WYLD (R. S.), F.R.S.E.

THE PHYSICS AND PHILOSOPHY OF THE SENSES; or,
The Mental and the Physical in their Mutual Relation. Illustrated by several Plates. Demy 8vo. 16s.

YONGE (C. D.), Regius Professor, Queen's College, Belfast.

HISTORY OF THE ENGLISH REVOLUTION OF 1688. Crown 8vo. 6s.

YORKE (Stephen), Author of "Tales of the North Riding."

CLEVEDEN. A Novel. 2 vols. Crown 8vo.

YOUMANS (Eliza A.)

AN ESSAY ON THE CULTURE OF THE OBSERVING POWERS OF CHILDREN, especially in connection with the Study of Botany. Edited, with Notes and a Supplement, by Joseph Payne, F.C.P., Author of "Lectures on the Science and Art of Education," etc. Crown 8vo. 2s. 6d.

FIRST BOOK OF BOTANY. Designed to cultivate the Observing Powers of Children. With 300 Engravings. New and Enlarged Edition. Crown 8vo. 5s.

YOUMANS (Edward L.), M.D.

A CLASS BOOK OF CHEMISTRY, on the Basis of the new System. With 200 Illustrations.

ZIMMERN (Helen).

STORIES IN PRECIOUS STONES. With Six Illustrations. Third Edition. Crown 8vo. 5s.