A-P [Phillips
J.C.]

Quail

no date

Quail breeding.

Phillips, J.C.

HARVARD UNIVERSITY.

LIBRARY

OF THE

MUSEUM OF COMPARATIVE ZOÖLOGY

54,162

BEQUEST OF

WILLIAM BREWSTER

May 10, 1920.

Facts About Quail Breeding

BY J. C. PHILLIPS

Photographs by the Author

Hen vs. Incubator—Some Practical Hints

WITH regard to game and its ownership, we are passing through a transition period. We have got far enough to know that, with our democratic and frontier methods, the game cannot last, but we might have known this fifty years ago. Thus far the attempt has been made to balance the increase of guns by a decrease of the open season and the bag limits, with the result that legitimate sport is almost a joke. Fancy keeping a dog and gun in Massachusetts for the purpose of shooting six pheasants a year (two only in one day) and fifteen grouse. I do not include woodcock, for they are too few to depend upon in most parts of this state.

We have legislated the old New England grouse dog out of existence because the man that bred and trained him, the market shooting type, has had to give way before the fury of the law makers. But now a change is slowly taking place. The unprotected farmer is worse off than he ever was because the cheap motor car has brought the city to his very doors, and if he happens to be the unfortunate owner of a good grouse or pheasant cover, he is safe only if he digs himself a trench and stays in it for the first of the open season; yes, and takes his family and his domestic animals in after him.

So far, I say, everything has gone against the farmer. His infrequent posting has been more or less ignored by types of men who, ignorant of farmers and farm life, think that once outside the radius of the police officer, they may help themselves to everything in sight. I am speaking of New England conditions only.

As far as I can see, these conditions will continue to get worse until one of these two things happens: either the plague of irresponsible shooters will

become so unbearable for the land owner that the rabid sentimentalists will have their say at last, and all shooting will be stopped by law; or else the farmer will learn the value of his birds, will be helped out by stricter trespass laws, and have something to say about his own property.

In the latter case the average man will have to plan for his shooting. He will have to organize into clubs or groups of sportsmen and make arrangements with farmers for his fall campaign. This does not mean that we are in any danger of Anglicizing our shooting, the farms are too small for that, but it does mean that the owner of the land will own the non-migratory game and will be encouraged to breed and protect game, and do with it as he sees fit. If the land owner has no rights with game reared on his own land, then, I should like to ask, who has?

At this point we are met by the old query: Then, where are the great army of poor men to get their shooting? The answer is, they won't get it, not all of them, for it is ridiculous to expect something for nothing. Which is better, to have no game at all, and no sport except for those who can afford long trips; or a moderate amount of nearby shooting for a large number of men who are willing to sacrifice a little for it? Another way out would be to issue licenses only to responsible people, and to raise the price. Such a measure would have small chance of going through at present.

Now some of us, sick and tired of the game law situation and the antics of the professional game law jugglers, are trying to find a way out through the propagation of game. By attempting to make this popular, we hope not only to increase the game and to

Mating pens should get plenty of sun and be placed where the grass does not grow too long and rank. A couple of pine boughs are put in for shelter, sand for dusting, charcoal and shell grit to aid digestion, and a crockery fountain for water

A twelve foot pen may be built in portable sections which will stow compactly when not in use, though it has few other advantages over the smaller size

The openings in the sliding door of the wooden coop allow the young quail to run in and out, while their foster mother is confined

stimulate the farmer's interest, but to eventually give the non-shooting public game to eat, and plenty of it. Already laws have been modified in a good many states to permit the rearing of game under license, and the sale of such raised game under proper supervision, but still a bitter war is being fought out between the two opposed camps of game protectors. On the one hand are the restrictive cranks who want to legislate all game and sport out of existence, and on the other hand the extreme game breeders who would abolish all laws except those covered by the Federal migratory statutes, and let game flow into the markets again. Obviously it is not quite time to adopt the latter course, but perhaps we are coming to it.

In the meantime, then, let us see what may be done in a small way with our native bob-white. These (and the California quail) are the only members of our American gallinae, (barring turkeys), which have proved easy to rear, and the bob-white is so remarkably easy that the wonder is he has not been raised before on a large scale.

Bob White and the Hen

In OUTING for October, 1915, there was an article devoted to quail raising which lauded the incubator and brooder methods, and dismissed the humble hen with scant attention. The writer is an advocate of the bantam method, and knows of many other quail farms that are using bantams with marked success. We see no disadvantage in using hens if they are the proper type, and we feel sure that natural methods are best for game birds whenever it is possible to use them.

Suppose then, that you have your stock of breeding quail bought and paid for and wintered in a fairly large pen, vermin-proof. Along in April or early May is the time to mate up your birds, and this is a rather delicate operation, because quail are very soft feathered and require to be carefully handled.

You have already provided your-

self with mating pens, one for each pair. I have tried various sizes. My first were 3 x 6 feet on the bottom and 3½ feet high, covered in with one-inch mesh, and fitted with a board a foot high around the bottom, and with a door in one end. It is a good thing to board up the end opposite the door, and also, if you want still snugger quarters, to attach a wooden box 22 inches square to the back end of the pen outside, with a small door leading into it. Then when you are gathering eggs or cleaning the pen, the birds will run into the box and hide until you are through.

We have made other pens in portable sections. They are 12 feet by 5 on the bottom, and 4½ feet high, but they cost about ten dollars each and have few, if any, advantages. We have gotten just as many eggs from the smaller pens, and the birds have kept fully as healthy, if properly fed. The advantage of the sectional pens is that they can be taken down and stored in piles.

Quail placed in large runs, even with only one pair to a run are not apt to do well, in my experience.

Placing the Pens

The placing of your mating pens is very important. They should get lots of sun and very little, or no, shade, and you should avoid ground where the grass will grow up long and rank in the pens. I made a great mistake formerly in placing pens in the woods; now I put them out in the most exposed situations and on old worn-out pasture ground. We put in a couple of thick pine boughs for shelter, a little heap of sand for dusting, a little charcoal and shell grit to promote digestion and a crockery fountain for water. A narrow strip of wire netting should be pegged down to the ground all around the bottom of the pen to keep vermin from burrowing under. The mated birds should be disturbed as little as possible, and fed once a day only enough for their actual needs. Never put more than one pair in a pen.

Facts About Quail Breeding

Some little time before laying begins, the hen birds shape nesting hollows in the ground, often several. Let them get a good start laying before you gather the eggs. At the first nest robbing, leave one egg in the nests that are full. Don't disturb the birds that have just begun to lay. We have gathered eggs at various intervals, but I think about every eight or ten days is a good rule, depending of course on how you are supplied with brooding bantams at the time. You want to hatch as many as you can early in the season, for then you will have strong and wild birds by the time the hard weather begins.

In exceptional cases, quail lay a great many eggs, but there are always some pairs that do not lay at all. My best pen this year produced 45 eggs, the next best 34. The average for the southern or southwestern stock was 15 eggs, and for the northern or New England stock, 11 eggs. I may add that a number of these pens were placed in bad situations where rank grass grew, that made too much shade and dampness. Besides this, my birds were broken up July 13, while still laying well.

My average, therefore, is much too low, and should easily be improved upon. These estimates were based on 24 pens. Two birds escaped during the laying season, and two died, so that there were only about 20 pens in full working order. The total produced was 350 eggs (a few eggs were gathered from some odd quail in a large run).

From these eggs we were able to rear to maturity about 175 quail. Exact estimate is impossible because the birds when grown are too wild and scattered to count accurately. We also sold about 100 eggs at a good price, ranging from 75 cents to over a dollar each.

I have now supposed that the quail breeder has on hand some good fresh quail eggs. The bantams are important. We have used mostly small mongrel cochins, and various crosses with silky bantams. The nest should be carefully shaped out in a sod, and about 16 eggs placed under each hen.

It would take too much space to follow all the details of this stage and of the hatching and rearing, so I wish to refer anyone who is especially interested to the very good routine account to be found in Job's "Propagation of Wild Birds." I will from now on dwell on my personal experience, and touch on some points that seem to me important.

When American breeders first began to rear quail they tried to confine them and handle them like pheasants in rearing yards. The little birds soon exhausted the insect food and developed the dread quail disease, which when once started will often sweep off a whole brood in one day. No progress was made until the free range system was developed. This consists briefly of the following procedure:

The hen is moved to the brooder coop as soon as the chicks are dry and strong, or as soon as the weather suits. The brooder coop is the same as that used for pheasants. Ours are 22 inches square, with sides 14 inches high. The photo shows the front of the coop with sliding door admitting chicks, but not bantams. At first the young quail are only let out into the guard, which is three sided and fits on the front of the coop. It extends out three feet in front of the coop and is one foot high. The presence of this guard enables the chicks to become used to the hen and prevents them from getting lost before they know their quarters.

Keeping them Healthy

If the weather is good, the guard ought to come off in four or five days. At this time we take a roll of 18 inch cellar wire and set it up on end with stakes, forming a circular pen with the coop inside. The larger this is, the better, and the grass or clover of the rearing field, if too long, should have broad paths cut through it, so that the young birds may have plenty of sun when they want it. As soon as the guard is removed, let the hen slip out into the new wire yard and range as she pleases. She will go back into

the coop at night and then can be closed in with the young brood. They are let out again about half-past six or seven o'clock in the morning.

The only objection to the cellar wire yard is its expense, but you only have to leave it on for about a week or ten days, and can then use it for other broods. It keeps the different sizes of quail apart until they get large enough to look out for themselves, and also keeps out vermin to a certain extent. The cellar wire ought to be removed at the latest when the young birds begin to fly over it, for they often get out, and are unable to get back.

After the wire is taken up, the young quail tend to pack together in large bunches, followed by several bantams, and then you run great danger from hawks, cats, etc. The free range system demands very careful attention to vermin trapping, although quail are less liable to attack from vermin than pheasants are, in my opinion. If the weather be very bad, with long storms, the quail will have to be kept longer under the guard, and even at times protected by old glass windows and shelters, covering over the guard or the ground in front of the coop. The ground must not be allowed to get wet under the coops, and the coops should be moved every other day to fresh ground.

Under the best conditions, some hens are sure to kill young quail, usually by scratching about on the floor of the coops, but sometimes by direct attacks. If the trouble is of the former kind, place a burlap on the ground under the coop; and if the latter, remove the hen as soon as possible, and distribute the chicks among other hens.

I have found it a good plan to sow a strip of oats along one side of the rearing field. This provides dense shade in July and good food in August and September, and even much later.

The young quail will, under the conditions of hand-rearing, or rather hen-rearing, tend to pack together so that as a sporting proposition they cannot be said to amount to much the first season, at least not till late in the autumn.

In my experience the bantams should be removed as soon as possible, for they only attract hawks, and other robbers, and tend to keep the quail too tame. Most of the bantams can be taken away at the end of about eight weeks. Bantams will usually bring the quail back to the brooder coops each evening and about dark they can all be closed in. When the birds are about two-thirds grown, they begin to sit out at night and are by then well able to look out for themselves.

Quail should not be caught up for confinement until they are full grown. It would be best to trap them in November, December, or January, having previously fed them at certain spots so that they come regularly to these places morning and afternoon. By catching your birds late, you run much less risk of disease and have larger and stronger stock for breeders another year.

The great trouble for the beginner has been in getting stock. The laws make the shipment of quail extremely difficult, and the prices have risen accordingly. At one time they cost three or four dollars a dozen; now they bring from \$22 to \$35 a dozen.

I have been asked whether quail breeding could be made profitable. It is too early to answer that question because nobody has handled the birds yet in very large numbers, and we do not know how much we could crowd them, or how long the land would last before diseases made a change to fresh ground necessary. Undoubtedly the price of adult birds would have to be high, at least \$4 or \$5 a pair, and there would have to be a good market for eggs. As far as breeding for preserves goes, the thing is perfectly practicable, and the sale of spare eggs and birds would greatly curtail the expenses of such an undertaking.

Hand-reared birds have a great advantage over wild birds in winter, because you can teach them where to get their food, and so keep them from starvation in the severest weather. You will begin to coax them to a few good feeding-places before the coops are all taken up, and the coop-feeding

stopped. But while the insect life lasts, and your patches of oats or buckwheat hold out, you cannot expect to centralize them very much. A little later you can build your traps and make ready to catch what you need.

A small flock of quail that I reared a year ago never went more than a few hundred yards from the feeding station and not one bird was missed between October and May. The winter yard for your breeding stock should, of course, be as large as possible, for it is occupied from the end of July until April.

In regard to feeding the young birds, this subject has been well covered in Mr. Job's book, but I will state briefly the method which I have used and found successful. For the old stock, almost any mixture of grain, cracked corn, wheat, kaffir corn, split peas, oats, etc., with green feed twice a week when possible, and a slightly more stimulating ration, including ant's eggs and Spratt's crissel during the laying season.

The Chick's Diet

For the young birds, about as follows: *First week*, custard and Spratt's pheasant meal,—say one part custard to three parts meal. Boil the custard until it is thoroughly set, and steam the pheasant meal for half an hour. Chop the custard very fine, mix it with boiled meal, and feed four times a day a very little at a time, on a perfectly clean board.

Second week. Hard boiled egg and steamed pheasant meal, about one part to four. By this time the young birds will be following the hen around the cellar wire yard and getting an abundance of insect food. They will even tackle huge grasshoppers, which they can hardly hold in their bills.

At the *third week*, feed hard-boiled eggs, pheasant meal, and Spratt's chick grain, mixing the pheasant meal and chick grain together before steaming. Use about one egg for five litters of quail, the feed should be mealy and not soft and sloppy, and the food should never be allowed to stand

around unused. This mixture can be fed until the birds are about ten weeks old, and then they can be started on a variety of small grains twice a day.

For starting quail on tracts of land where no wild breeding stock exists, it is far better to go to a little trouble and hand-rear your birds. A dozen young birds raised on your land are worth a hundred adult wild birds simply turned loose. Thousands of dollars have been wasted in buying and turning out adult quail, and only very rarely do we hear good results reported. My own experiments in this line have always been total failures.

This sketch was not intended to be an exhaustive account of quail breeding; it was meant only to show the possibilities that now confront the energetic sportsman. Thousands of acres of farm lands in southern New England are adapted to quail rearing, and the business man with comparatively little trouble and expense could have abundant sport near home.

So far as I can see, pheasant and quail rearing can be carried on side by side, but when a good stock of quail has been reared and is properly looked after, it might not be necessary to hand-rear large numbers of them every year. Quail are splendid mothers, and can be depended upon to raise large families. All that is necessary is to provide patches of oats and buckwheat, and keep down stray cats and other enemies.

Let us then face this subject of diminishing game supply by direct attack rather than by multiplication of laws. It is up to the sportsmen themselves to see that the irresponsible shooters do not put an end to all sport by so annoying the land-owners that they are obliged to go to extremes. Only the other day a farmer told me that he had given up working in his field during the open season because he was not safe there. In some cases town societies have united to post a whole town, and not without reason. A state of affairs like this is unfair, unnatural, and cannot last.

Gaylord Bros.
Makers
Syracuse, N Y.
PAT. JAN. 21, 1908

Date Due

~~JUN 23 1986~~