Projected Changes in the Energy Release Component Under Climate Change in Northwest Predictive Services Areas

January 2015

A Report to the Bureau of Land Management Oregon-Washington State Office

Prepared by

The Oregon Climate Change Research Institute (OCCRI)

Authors:

Meghan M. Dalton, OCCRI, Oregon State University, Corvallis, OR

John T. Abatzoglou, Department of Geography, University of Idaho, Moscow, ID

Louisa Evers, Bureau of Land Management, Oregon-Washington State Office, Portland, OR

Katherine Hegewisch, Department of Geography, University of Idaho, Moscow, ID

Table of Contents

ABSTRACT	3
1. INTRODUCTION	4
2. DATA	5
2.1 STATION DATA	6
2.2 GLOBAL CLIMATE MODEL DATA	7
3. METHODS	9
3.1 HISTORICAL ERC	9
3.2 FUTURE ERC	10
4. RESULTS	11
4.1 CLIMATOLOGY	11
4.2 OBSERVED TRENDS	11
4.3 FUTURE PROJECTIONS	12
5. DISCUSSION	14
6. CONCLUSION	16
REFERENCES	17
TABLES	23
FIGURES	38
APPENDIX A: SEASONAL ERC CLIMATOLOGY	45
APPENDIX B: PERCENT OF DAYS EXCEEDING 90TH PERCENTILE ERC	52
APPENDIX C: TRENDS IN MEAN FIRE SEASON ERC	55
APPENDIX D: TRENDS IN % OF DAYS EXCEEDING 90TH PERCENTILE ERC	62
APPENDIX E: SEASONAL CLIMATE PROJECTIONS	75
APPENDIX F: CHANGE IN ERC VS. CHANGE IN CONTRIBUTING VARIABLES	80

Abstract

Fire management agencies in the United States (US) use fire danger indices from National Fire Danger Rating Systems as decision-support tools. A widely used index tracking seasonal dryness in the western US is the Energy Release Component (ERC). Historically, ERC percentile thresholds provide information on the potential for large and uncontrollable fires, but climate change may alter the utility of specific percentiles in identifying large fire potential in the future. We investigated two management-focused questions: 1) how has seasonal ERC and percentile values changed, and 2) how will ERC and the number of high fire danger days change under projected future climatic conditions. We calculated historical and future ERC for Predictive Service Areas in Oregon and Washington using station observations and downscaled climate models, respectively. ERC and frequency of high fire danger days during the fire season increased for most PSAs over the 1981-2010 period. These increases were consistent with projected increases in ERC and high fire danger days using climate projections, with the most acute increases projected east of the Cascades and during the peak of the fire season. Irrespective of changes in vegetation and fire management, these changes in fire danger have and will continue to increase wildfire exposure across the region.

1. Introduction

Wildland fire managers have long recognized the interannual variability in regional fire activity and seasonal dryness. A number of different fire-danger indices have been developed to monitor seasonal fire potential due to variability in weather and climate (e.g., McArthur, 1967; Van Wagner, 1987; Stocks et al., 1989). The National Fire Danger Rating System (NFDRS) was developed in the United States in the 1970s as a statistical-based system used to model fire behavior, initiation, and spread (Bradshaw et al., 1984; Cohen and Deeming, 1985; Burgan, 1988). These fire danger indices and systems all utilize daily surface meteorological data from weather stations, but differ in their sensitivities to forcing variables and serial correlation thus resulting in different application potential to vegetation types (e.g., Wotton, 2009; Mölders, 2010).

Federal, state and local fire management agencies in the United States use NFDRS indices as measures of fire danger, including Energy Release Component (ERC), a weather-climate hybrid index of daily fire energy intensity that considers the cumulative drying effect of weather in the previous days to weeks to estimate live and dead fuel moisture values (Bradshaw et al. 1984; Cohen and Deeming 1985). ERC is the most widely used index in the western United States to track seasonal dryness (Brown et al., 2002; Andrews et al., 2003). These numerical values are used as fire business decision support tools to guide suppression and fuel treatments activities. The historical distribution of ERC helps define periods of high fire danger; above the historical 90th percentile the potential for large fires increases and above the 97th percentile uncontrollable fires are expected. ERC is among the key indicators used in developing pocket cards used by firefighters to compare the current season with notable fire seasons and large fires of the past in a local area (Andrews et al. 1998, see also <http://fam.nwcg.gov/fam-web/pocketcards/default.htm>).

Annual mean temperature in the Pacific Northwest increased by approximately 0.6°-0.8°C from 1901 to 2012, with nearly all of the increase in summer temperature occurring in the past three decades (Abatzoglou et al., 2014). In the absence of long-term coherent changes in precipitation, warming temperatures and an earlier commencement of the growing season can promote declines in soil and fuel moisture, increased ERC, and fuels that are more receptive to fire.

Studies have noted an increase in the length of the fire season in the western United States due to earlier snowmelt and increased temperatures (Westerling et al. 2006), although increased vapor pressure deficit and ERC during the fire season have also contributed (Williams et al., 2014; Abatzoglou and Kolden, 2013).

Fire managers in the western United States have begun to wonder how both observed and projected changes in climate may be manifest in terms of fire danger indices such as ERC. Large fires have become more frequent and total area burned has increased across the West during the last 30 years (Dennison et al., 2014) and over the past century (Littell et al., 2009). Significant increasing trends in ERC during 1979-2012 occurred throughout the Southwest US and in some areas in southeast Oregon and eastern Washington, though the trend in most areas in Oregon and Washington were not distinguishable from zero (Abatzoglou and Kolden, 2013). Applying statistical relationships between very large fires and weekly variability in temperature, ERC, and other weather variables, to future climate conditions, the probability of very large wildfires is projected to increase by at least 30% by the end of the century in the West (Stavros et al., 2014). ERC percentiles are a good predictor of frequency of large fires and area burned (Riley et al., 2013), but some managers suspect that the 90th and 97th percentile values will increase as new data is included such that what is considered high fire danger today may occur at lower percentile values in the future. We describe the historical seasonal ERC and 90th and 97th percentile values, examine recent ERC trends, and present projections of ERC and frequency of high fire danger days under future climate conditions for Oregon and Washington.

2. Data

Our region of interest includes the US states of Washington and Oregon. Strong longitudinal gradients in temperature and moisture exist across the region and are further reinforced by the north-south oriented Cascade Mountains. The region receives the majority of its annual precipitation during the cool season (Oct-May) with over an order of magnitude difference in precipitation between the windward slopes of the Cascades, coast range and Olympics and the rain-shadowed deserts in central to eastern Washington and Oregon. Summer temperature and humidity are moderated in western Oregon and Washington due to the maritime influence,

whereas east of the Cascade crest summer temperature and humidity are more continental. The Northwest Coordination Center (NWCC) developed 12 Predictive Service Areas (PSAs) covering the region (Figure 1, Table 1) to track subregional changes in fire danger and fire activity. Each PSA represents an area of similar seasonal trends in fire danger and wildfire activity. Climate provides a template for the ecoregions that cover the region. Western PSAs on the windward side of the Cascade Mountain Range (NW01-NW04) and in northeast Washington (NW08, NW09) are generally characterized by dense, closed canopy, and multi-layered forests, dominated by coastal Douglas-fir, western hemlock, and several species of true firs. Central and eastern forest PSAs on the lee side of the Cascade Mountains (NW05-NW07) and the Blue Mountains (NW11) are generally characterized by more open forests with grassy or shrubby understories; the most common tree species are ponderosa pine, interior Douglas-fir and either grand fir or white fir. The semi-arid PSAs (NW10 and NW12) consist of grassland, sagebrush-steppe and juniper woodlands.

2.1 Station Data

To support fire danger rating, federal and state firefighting agencies initially established a system of manual weather stations (Fisher and Bradley 1976) that have subsequently been replaced with remote automated weather stations (RAWS). NWCC identified from 3 to 9 representative RAWS within each PSA that are used to assess changes in fire danger across the geographic area to support decisions concerning movement of firefighting resources and prioritization of large fires. We obtained station catalogs and daily datasets from station start year to 2010 for the 73 key stations used operationally by NWCC.

Typically, trend analyses use all available years, but as daily records were missing a considerable amount of data prior to 1980, we constrained our analyses to data from 1981-2010 to reduce the impact of missing data on the results. For NW10, data was available starting only in 1987. Many RAWS records (and thus PSA records) are missing considerable data before and after the declared fire season (June 1 to October 31) as not all RAWS are operated year-round and of those that are most lack heated rain gauges so generally do not accurately measure winter precipitation, so we restricted our analysis to the fire season for the same reason. Some missing

data remained, but preliminary procedures to fill in the gaps suggest that filling gaps would have little effect on the results so is not pursued further here.

We conducted an extensive quality control procedure on the station data following Serreze et al (1999) and including manual removal of erroneous data. Daily temperatures outside of state records of -48°C (-54°F) and 48°C (119°F) were set to missing along with any internal inconsistencies (e.g., the daily maximum temperature being less than the daily minimum temperature). Daily temperatures exceeding 3 standard deviations of monthly means were also removed. This procedure was performed a second time with recomputed means and standard deviations. For precipitation, negative values and values larger than 38.1 cm (15 in) were set to missing values as well as values outside of 5 standard deviations of monthly means for square root of precipitation following Serreze et al (1999). Any spurious values that remained were checked manually and removed if found to be inconsistent with nearby stations. For relative humidity, negative and zero values and values larger than 100% were set to missing values along with any internal inconsistencies.

The maximum amount of data removed from a station record by this procedure was ~3% for temperature data and ~2% for precipitation data. In half of the stations, less than 1% of temperature data and less than 0.1% of precipitation data was removed. For relative humidity variables, up to a quarter of the data was removed for a few stations, but for half of the stations, less than 2% was removed. Historical trends were computed prior to quality control, but the quality-controlled version of the station data was used to bias-correct the downscaled GCM data where erroneous values would have a larger impact. We also did not correct RAWS data for station biases (e.g., station moves, vegetation canopy changes, instrumental changes, etc.) that may affect trends in the observational record.

2.2 Global Climate Model Data

Historical (1950-2005) and future (2006-2100) simulations were acquired from 18 global climate models (GCMs) that participated in the Coupled Model Intercomparison Project phase 5 (CMIP5, Taylor et al., 2012; Table 2). We considered two future scenarios: Representative Concentration Pathways 4.5 and 8.5 (RCP4.5 and RCP8.5), corresponding to future pathways of

radiative forcing (additional energy trapped by the Earth-Atmosphere system by 2100, measured in W m^{-2} , quantified relative to preindustrial conditions circa 1765). RCP 4.5 represents a reduction in the current rate of greenhouse gas emissions while our current trajectory aligns most closely with RCP8.5, which represents continued emissions.

Daily output of maximum and minimum temperature (TMAX, TMIN), maximum and minimum relative humidity (RMAX, RMIN), precipitation (PPT), and downward shortwave solar radiation (RSDS) from the first ensemble member of each model was statistically downscaled using the Multivariate Adaptive Constructed Analogs (MACA) method (Abatzoglou and Brown, 2012). Downscaling was performed first to the gridded surface meteorological dataset of Abatzoglou (2013) at 1/24 deg (~4-km) spatial resolution that contains the requisite variables for computing ERC. The incongruence between gridded data and station data can be rather pronounced in complex terrain, necessitating that we adjust the downscaled projections for the station location. Secondly, we extracted data from the pixel closest to each of the RAWS and performed a bias correction of each of the downscaled fields, except RSDS, to ensure that the historical modeled data have the same statistics as the observed RAWS.

We derived daily values for ‘precipitation duration’ (PDUR) and ‘state of the weather’ (SOW) using the downscaled data at the stations following Abatzoglou and Kolden (2013). We estimated PDUR using formula (1),

$$(1) PDUR = 24 * (1 - e^{-\beta * PPT})$$

Where PPT is the daily precipitation total and β is a function of latitude and longitude and can be thought of as the inverse of average precipitation intensity (M. Jolly, personal communications). The SOW was determined by daily precipitation and percent of maximum daily solar insolation (Table 3). For example, the state of the weather was assigned SOW 6 (rain) when PPT was greater than 0.13 cm (0.05 in) and for days with PPT less than 0.02 cm (0.01 in), the state of the weather was assigned SOW 3 (overcast) when RSDS was less than 50% of maximum, and 0 (clear) when RSDS was greater than 91% of maximum.

3. Methods

3.1 Historical ERC

We imported data from the 73 key RAWS into FireFamilyPlus 4.1 beta (FFP) (RMRS, 2002; Bradshaw and Tirmenstein, 2009) and created Special Interest Groups (SIGs) for each PSA and assigned equal weight to all reporting stations in each PSA, in keeping with the procedures used by NWCC. Thus, if only one key station had data for a specific day, the PSA dataset had data for that day. Missing data in the PSA dataset arose when none of the key stations used within that PSA contained a data point. Less than 10% of fire season days during 1981-2010 were missing for each PSA record, except NW12 in which 13% of days were missing (Table 1). Key stations within a PSA began recording data in different years resulting in an increasing number of reporting key stations over time, which could influence PSA weather records and trends. For example, four out of the nine key stations in NW01 didn't start recording data until sometime after 1981, with one station starting as late as 2001. Key stations within a PSA were all reporting data by the mid-1990s for most PSAs (Table 1).

We followed the standards used by NWCC, by (i) using fuel model G since it best reflects the seasonal dynamics of the fire season regardless of vegetation differences between PSAs, (ii) assuming herbaceous plants are mainly perennial in all PSAs, which allows for more direct comparisons between PSAs because different cure rates for annual and perennial grasses affect calculated ERC (Bradshaw et al. 1984), and (iii) using constant green-up dates. This procedure resulted in daily ERC for the 30-year period from 1981-2010 during the fire season (Jun-Oct).

We calculated the 90th and 97th percentiles of fire season ERC for each PSA and rounded this number to the nearest integer. We also calculated the fire season mean ERC and the number of fire season days greater than the 1981-2010 90th and 97th ERC percentiles. The number of fire season days per year exceeding thresholds was converted to a percentage by dividing by the number of non-missing data points per year. We also computed the mean ERC and percent of days greater than the 1981-2010 fire season 90th and 97th percentile thresholds for each month of the fire season (Jun-Oct).

We analyzed trends in the mean fire season ERC and the percent of fire season days exceeding the 1981-2010 90th and 97th percentile thresholds using the statistical software R (R Core Team, 2013). We also evaluated trends for each month from Jun-Oct. We computed the least-square linear regression and evaluated the significance of the trend using the non-parametric Mann-Kendall trend test. In the case of a time series with significant autocorrelation, we bootstrapped the Mann-Kendall statistic (n=500) to construct the 95% confidence interval. Several time series of percent of days greater than 90th and 97th percentile thresholds displayed heteroskedasticity (non-constant variance of residuals) due to the presence of zeros in the early part of the record. This implies that the linear model of the trend is inefficient and the significance may be overestimated, calling for caution in the interpretation of trends in these time series.

3.2 Future ERC

Using the downscaled variables of daily TMIN, TMAX, RMIN, RMAX, PPT, PDUR, and SOW at each station, we estimated ERC with the method utilized in Abatzoglou and Kolden (2013). In this method, fuel model G is used with dynamic green-up dates defined by the first day of each year when the normalized growing season index for each pixel >0.5 (Jolly et al., 2005; M. Jolly, personal communications). We replaced missing values in the bias corrected data with default values which should put the fuels to ‘sleep’, namely temperatures of 0 °C, relative humidity of 100% and zero precipitation. Since these missing values all occur in winter before green-up, these should have very small influences on any results.

We analyzed future ERC for early- (2010-2039) and mid- (2040-2069) century and compared against the modeled historical baseline period (1950-2005). The downscaled data were adjusted to match the statistics of the RAWs observations from 1981-2010, so it is proper to use the entire modeled historical baseline in comparison with observations. As in the RAWs analysis, we computed the fire season mean, 90th, and 97th percentile thresholds for each time period for each model and each scenario and each PSA. We computed the percent of days during the fire season and during each month of the fire season above the modeled historical (1950-2005) fire season thresholds, and daily mean ERC climatology over each time period. Results below are presented as the multi-model mean change (that is, computing the future change for each model compared to each model's historical period, then averaging the amount of change).

4. Results

4.1 Climatology

Climatological ERC increases throughout spring and into summer before reaching its peak in August and subsiding thereafter (Appendix A). ERC values are lower west of the Cascades and higher in the semi-arid central and eastern parts of the region and at more southerly latitudes (Table 4). Similarly, the fire season 90th (97th) ERC percentiles for the 1981-2010 period vary geographically across the region (Figure 2, Tables 5, 6). By definition, approximately 10% and 3% of fire season days during 1981-2010 exceed the 90th and 97th percentile thresholds, respectively. High fire danger days are primarily concentrated in July-September, with some regional differences (e.g., greater percent of days in Sep-Oct in the southwestern PSAs) (Table 7, Appendix B).

4.2 Observed Trends

Linear trends in mean fire-season ERC exhibited increases of 3 to 7 units per decade over the 1981-2010 period (Table 8, Appendix C), with the largest increases in northeastern WA (NW08 and NW09) and northeastern OR (NW11). Trends were significant at the 95% level for all but three PSAs (NW01, NW10, NW12). Likewise, significant positive trends in ERC were observed for the months of July, August, and September in most PSAs, as well as for October in six PSAs (NW04, NW06, NW07, NW08, NW09, NW11). Conversely, no significant trends were found for June ERC. The nearly significant negative June ERC trend in NW12 is associated with a positive trend in June precipitation (not shown). No significant trends in ERC were observed in the wettest PSA (NW01 – northwestern WA) as well as two of the driest PSAs in the eastern part of the region (NW10, NW12 – Columbia Basin, southeast OR) for the fire season as a whole, although an increase in August ERC was noted for the latter. Abatzoglou and Kolden (2013) also found significant trends in ERC in some Northwest locations, but the magnitude was smaller on the order of 3-4 units per decade. This difference in trend magnitude may be due to several factors including the increasing number of reporting stations within a PSA over time and station moves during the switch from manual to automated reporting.

Increases in percent of the fire season and individual months of July-September exceeding historical 90th percentile ERC thresholds were found for all but three PSAs, the same three without a significant trend in the mean (Table 9, Appendix D). Likewise, an increase in the percent of days exceeding the 97th was noted for several PSAs east of the Cascades (Table 10, Appendix D). While linear trends oversimplify the time evolution of such time series, much of the increase appears to arise due to differences since the year 2000. While western PSAs did not have significant trends in percent of fire season days exceeding the 97th percentile, there were significant trends in August or September for NW02, NW03, and NW04.

4.3 Future Projections

The multi-model mean downscaled historical (1950-2005) ERC climatology (Figure 3) was broadly consistent with the observed (1981-2010) ERC climatology with a few exceptions. Downscaled historical mean fire season ERC and 90th and 97th percentile values were within 6 units of observed station based ERC (Tables 4-6). The downscaled historical ERC exhibited a slight shift in the timing of high fire danger days relative to observations with more days in June and July exceeding ERC thresholds in downscaled data versus observations and vice-versa for September and October. While simulated historical ERCs may not have the exact same characteristics of observed ERCs due to differences in how GCMs simulate sequences in variables and joint variability across variables, overall, the differences are small.

Projected changes in ERC climatology by the 2020s and 2050s were pronounced during the peak of the fire season (Figure 3) when climate change projections (e.g., increased temperature, decreased relative humidity and precipitation; Appendix E) were most conducive to drying fuels. Projections for the 2050s under RCP8.5 showed the largest magnitude of change with the most models agreeing on the sign of the change. We describe results for the 2050s under RCP8.5. Projections for RCP4.5 and the 2020s under both scenarios were similar albeit with a smaller magnitude and less model agreement and are presented in the tables. By the 2050s, multi-model mean fire season ERC increased by around 3 units for all PSAs under RCP8.5 compared to the historical baseline (1950-2005) (Figure 4, Table 4). The 90th and 97th percentile ERC value increased by a larger margin of between 4-7 units (Tables 5, 6) with larger increases generally in central and eastern PSAs (Figure 5). The larger increase in extreme ERC values compared with

the mean resulted in an increased standard deviation of fire season ERC values by up to 2 units on average (not shown).

The multi-model mean projected increase in fire season days exceeding the 90th percentile historical baseline for the 2050s under RCP8.5 was 6-11% (9-17 days) depending on the PSA (Table 11, Figure 6; 2020s in Table 12) with at least three-quarters of the models agreeing on the sign of change. These increases in high fire danger days were most pronounced from July through September with actual increases of 8-13% (2-4 days), 12-23% (4-7 days), and 7-14% (2-4 days), respectively, with at least three-quarters model agreement. Much smaller increases were noted during the fire season shoulders of June and October (Table 11). Depending on the PSA, approximately 20-30% of August days (6-9 days) exceeded the 90th percentile value during the historical baseline and by the 2050s under the high emissions scenario (RCP8.5) some models projected that 50-75% of August days (15-23 days) would exceed the historic baseline 90th percentile threshold (Figure 7). Results are similar for actual change in percent of fire season days exceeding the 97th percentile except changes are of a smaller magnitude with projected fire season increases of 3-8% (4-12 days) by the 2050s for RCP8.5 (Table 13; 2020s in Table 14) and the largest changes also occur in July, August, and September.

We examined the contributions of each variable used in the ERC calculation to the variance of future changes in mean fire season ERC. Aggregating the future fire season changes for the 2020s and 2050s and both RCPs, we computed the square of Pearson's correlation coefficient between ERC changes and changes in TMIN, TMAX, RMIN, RMAX, PPT, PDUR, RSDS, and SOW during the fire season (Table 15). ERC increases were correlated with increases in TMIN, TMAX, and RSDS, and decreases in RMIN, RMAX, PPT, PDUR, and SOW (Appendix F). ERC was most sensitive to relative humidity, found in earlier studies (e.g., Brown et al., 2004): changes in RMAX explained from 29% to 56% of the variance in ERC changes and RMIN explained from 40% to 64% of the variance depending on the PSA. Changes in ERC in northern and western PSAs tended to depend more heavily on changes in temperature, relative humidity, and precipitation than southern and eastern PSAs.

5. Discussion

Large and rapidly spreading fires are expected when the ERC exceeds 90th and 97th percentile values. Thus, the firefighting community uses these values to support requests for additional funds to hire or activate additional firefighting resources, extend the hours of operation for existing resources, and to shift resources from areas with lower fire danger into areas with higher danger, especially when a triggering event, such as thunderstorms or high atmospheric instability are forecast. Incident management teams and agency administrators use the current ERC percentile to support decisions on how to manage individual wildfires and complexes of wildfires, such as determining what values may be at risk in the near future, how long the fire may burn, and how aggressively to manage the fire as well as to support the development of contingency plans. The projected changes in ERC by mid-century indicates that how managers use ERC may need to be re-evaluated: reliance on the percentile value may result in an underestimation of risks in the future as the current value associated with a given percentile increases in the future.

The greatest amount of change in mean ERC and the percent of days above the long-term 90th and 97th percentiles occurred historically and is expected to occur in the future primarily in August and September. Small increases during the beginning and end of the fire season (June and October) are consistent with other research findings that the fire season is beginning earlier and lasting longer (e.g., Westerling et al., 2006, Abatzoglou and Kolden, 2011). The larger increases in ERC during the peak fire season are associated with larger changes during the peak season in relative humidity and to a lesser degree, temperature consistent with regional climate studies indicating larger changes in these variables in summer than in spring or fall (Mote et al., 2013; Collins et al., 2013).

Our results of increasing fire danger days in the future are consistent with previous fire danger studies in the Western US and other parts of the world using different fire danger indices. Under doubled-CO₂ simulations, the number of high fire danger days increased across Australia (Williams et al., 2001; Beer and Williams, 1995) and fire danger increased in areal extent across Russian and Canadian boreal forests, particularly in June and July with an earlier start to the fire

season (Stocks et al., 1998). In the western US, projected increases in high fire danger days (based on ERC) associated with drying over most of the West occurred throughout the 21st century and were most pronounced in the northern Rockies, Great Basin, and the Southwest (Brown et al., 2004). By mid-century, the largest increases in extreme fire danger days (defined as $ERC \geq 60$), of about 2 weeks, occurred in eastern Oregon and Washington (Brown et al., 2004). Brown et al. (2004) also found that increases in high fire danger days tended to occur within the fire season rather than substantially lengthen it, consistent with our results.

However, the observed trend in mean October ERC and projected increases (with model agreement) in percent of October days exceeding the 90th and 97th percentile for some southern, central, and eastern PSAs suggests that further attention to fire behavior and management toward the end of the fire season may be warranted under future climate change. Although the reduction in solar zenith angle and shortening of the day beginning around mid-September decreases fire potential, large growth events have occurred in the past in October. These remain very rare events associated with extreme October weather conditions (e.g., very high temperatures, very low relative humidity, and very strong winds) and tend to last only a few hours as opposed to days.

The magnitude of projected increase in ERC values is greater in the cooler, moister PSAs with a greater marine influence year-round, than in the warmer, drier PSAs on the east side of the Cascades Mountains. Central and eastern PSAs tend to exhibit larger projected future increases in percent of days greater than the 90th or 97th percentiles than western PSAs, consistent with Abatzoglou and Kolden (2011) and the larger projected increases in area burned for the eastern Cascade Range and non-forested systems compared to the western Cascade Range (Littell et al., 2013).

Changes in ERC in forest-dominated PSAs are more sensitive to changes in relative humidity, temperature, and precipitation than rangeland-dominated PSAs, suggesting a bias in the equations. NFRDS was developed in a time when little weather and fuels data were available for rangelands, range fires were not perceived to be significant problems, crown fires in forests were rare, and computing ability was limited. Fire research at the time was focused on surface fires in

dead continuous fuels, conducted primarily in laboratories, and with measures easily collected in the field (e.g., Rothermel 1972; Deeming et al. 1974; Albini 1976). Relative humidity and even low amounts of precipitation affect the moisture status of dead fuels, especially fine dead fuels (Rothermel 1972), but have little effect on the moisture status of live fuels. Instead, live fuel moisture responds to the interaction between evapotranspiration demand and soil water availability (Dennison et al. 2003; Dimitrakopoulos and Bemmerzouk 2003), of which temperature and precipitation may serve as proxies. Given the importance of live fuels in crown fires in forests and conifer woodlands and most fires in shrublands and shrub-steppe (Jolly 2007; Dennison and Moritz 2009), a re-examination of the NFDRS equations may be warranted to better reflect the role of live fuels in fire danger. Recent research has focused on the use of remote sensing to detect changes in the relative water content or effective water thickness in leaves to determine live fuel moisture (e.g., Chuvieco et al. 2003; Dennison et al. 2003; Dennison et al. 2005; Yebra et al. 2008).

6. Conclusion

The energy release component (ERC) is a widely used fire danger index assessing the potential for large, rapidly spreading fires. Fire managers in the Northwest are concerned about using ERC percentile thresholds to determine high fire danger days instead of a set value as the climate continues to warm in the future. The 90th and 97th percentile values are projected to increase by a greater margin than the mean by mid-century indicating that the thresholds currently used to assess fire danger are a “moving target” as the climate changes, motivating a reexamination of the understanding of fire behavior at critical ERC values. Future changes in ERC and high fire danger days are expected to occur during the existing peak fire season and be largest in areas already experiencing the highest number of high fire danger days. Future projections show an exacerbation of existing fire danger conditions. Our results suggest that managers should consider using a threshold value based on the historical percentiles (1981-2010). Comparing large fire growth events against both the threshold value and the current percentile value may not reveal much difference in the short-term, particularly given how the percentile value is calculated. However, over time as the actual number associated with a given percentile increases, managers may well see a difference emerge.

References

- Abatzoglou JT, Rupp DE, Mote PW (2014) Seasonal climate variability and change in the Pacific Northwest of the United States. *Journal of Climate* **27**, 2125-2142. DOI: 10.1175/JCLI-D-13-00218.1.
- Abatzoglou JT (2013) Development of gridded surface meteorological data for ecological applications and modeling. *International Journal of Climatology* **33**, 121-131. DOI: 10.1002/joc.3413
- Abatzoglou JT, Brown TJ (2012) A comparison of statistical downscaling methods suited for wildfire applications. *International Journal of Climatology* **32**, 772-780. DOI: 10.1002/joc.2312.
- Abatzoglou JT, Kolden CA (2013) Relationships between climate and macro scale area burned in the western United States. *International Journal of Wildland Fire* **22**, 1003-1020. DOI:10.1071/WF13019
- Abatzoglou JT, Kolden CA (2011) Climate change in western US deserts: Potential for increased wildfire and invasive annual grasses. *Rangeland Ecology & Management* **64**, 471-478. DOI: 10.2111/REM-D-09-00151.1.
- Albini FA (1976) Estimating wildfire behavior and effects. USDA Forest Service, Intermountain Forest and Range Experiment Station General Technical Report INT-30. (Ogden, UT)
- Andrews PL, Loftsgaarden DO, Bradshaw LS (2003) Evaluation of fire danger rating indexes using logistic regression and percentile analysis. *International Journal of Wildland Fire* **12**, 213-226. DOI: 10.1071/WF02059.
- Andrews PL, Bradshaw LS, Bunnell, DL, Curcio GM (1998) Fire danger rating pocket card for firefighter safety. (Extended Abstract) 2nd Symposium on Fire and Forest Meteorology: Jan 11-16, 1998. (Phoenix, AZ) Available at http://fam.nwcg.gov/fam-web/pocketcards/firedanger_pc_ams_1998.pdf
- Beer T, Williams A (1995) Estimating Australian forest fire danger under conditions of doubled carbon dioxide concentrations. *Climatic Change* **29**, 169-188. DOI: 10.1007/BF1094015.
- Bradshaw LS, Deeming JE, Burgan RE, Cohen JD (Comps) (1984) The 1978 National Fire-Danger Rating System: Technical documentation. USDA Forest Service, Intermountain Forest and Range Experiment Station, General Technical Report INT-169. (Ogden, UT)

- Bradshaw LS, Tirmenstein D (2009) FireFamilyPlus version 4 user guide (Draft). USDA Forest Service, National Fire and Aviation Management Information Systems Team (Boise, ID). Available at http://www.firemodels.org/downloads/firefamilyplus/publications/FFP-4_1_Draft_Users_Guide.pdf
- Brown TJ, Hall BL, Westerling AL (2004) The impact of twenty-first century climate change on wildland fire danger in the western United States: An applications perspective. *Climatic Change* **62**, 365-388. DOI: 10.1023/B:CLIM.0000013680.07783.de.
- Burgan RE (1988) 1988 revisions to the 1978 National Fire-Danger Rating System. USDA Forest Service, Southeastern Forest Experiment Station, Research Paper SE-273 (Asheville, NC)
- Cohen JD, Deeming JE (1985) The National Fire-Danger Rating System: basic equations. USDA Forest Service, Southwest Forest and Range Experiment Station, General Technical Report PSW-82 (Berkeley, CA)
- Collins M, Knutti R, Arblaster J, Dufresne J-L, Fichetef T, Friedlingstein P, Gao X, Gutowski WJ, Johns T, Krinner G, Shongwe M, Tebaldi C, Weaver AJ, Wehner M (2013) Long-term climate change: Projections, commitments and irreversibility. In 'Climate change 2013: The physical science basis. Contribution of working group I to the fifth assessment report of the Intergovernmental Panel on Climate Change' (Eds TF Stocker, D Qin, G-K Plattner, M Tignor, SK Allen, J Boschung, A Nauels, Y Xia, V Bex, PM Midgley) pp. 1029–1136. (Cambridge University Press: Cambridge, United Kingdom and New York, NY, USA) DOI:10.1017/CBO9781107415324.024.
- Chuvieco E, Aguado I, Cocero D, Riaño D (2003) Design of an empirical index to estimate fuel moisture content from NOAA-AVHRR images in forest fire danger studies. *International Journal of Remote Sensing* **24**, 1621-1637. DOI: 10.1080/01431160210144660.
- Deeming JE, Lancaster JW, Fosberg MA, Furman RW, Schroeder MJ (1974) The National Fire Danger Rating System. USDA Forest Service, Rocky Mountain Forest and Range Experiment Station Research Paper RM-84 (Rev.) (Ft. Collins, CO)
- Dennison PE, Roberts DA, Peterson SH, Rechel J (2005) Use of normalized difference water index for monitoring live fuel moisture. *International Journal of Remote Sensing* **26**, 1035-1042. DOI:10.1080/0143116042000273998.

- Dennison PE, Roberts DA, Thorgusen SR, Regelbrugge JC, Weise D, Lee C (2003) Modeling seasonal changes in live fuel moisture and equivalent water thickness using a cumulative water balance index. *Remote Sensing of the Environment* **88**, 442-452. DOI: 10.1016/j.rse.2003.08.015
- Dennison PE, Moritz MA (2009) Critical live fuel moisture in chaparral ecosystems: A threshold for fire activity and its relationship to antecedent precipitation. *International Journal of Wildland Fire* **18**, 1021-1027.
- Dennison PE, Brewer SC, Arnold JD, Moritz MA (2014) Large wildfire trends in the western United States, 1984–2011. *Geophysical Research Letters* **41**, 2928–2933. DOI: 10.1002/2014GL059576.
- Dimitrakopoulos AP, Bemmerzouk AM (2003) Predicting live herbaceous moisture content from a seasonal drought index. *International Journal of Biometeorology* **47**, 73-79. DOI: 10.1007/s00484-002-0151-1.
- Jolly WM, Nemani R, Running SW (2005) A generalized, bioclimatic index to predict foliar phenology in response to climate. *Global Change Biology* **11**: 619-632.
- Jolly WM (2007) Sensitivity of a surface fire spread model and associated fire behaviour fuel models to changes in live fuel moisture. *International Journal of Wildland Fire* **16**, 503-509.
- Littell JS, McKenzie D, Peterson DL, Westerling AL (2009) Climate and wildfire area burned in western U.S. ecoprovinces, 1916–2003. *Ecological Applications* **19**, 1003–1021. DOI: 10.1890/07-1183.1.
- Littell JS, Hicke JA, Shafer SL, Capalbo SM, Houston LL, Glick P (2013) Forest ecosystems: Vegetation, disturbance, and economics. In ‘Climate change in the Northwest: Implications for our landscapes, waters, and communities.’ (Eds MM Dalton, PW Mote, AK Snover) pp. 110-148. (Island Press: Washington, DC)
- McArthur AG (1967) Fire behaviour in eucalypt forests. Commonw. Aust. For. And Timber Bur. Leaflet Number 107. 36pp.
- Mölders N (2010) Comparison of Canadian Forest Fire Danger Rating System and National Fire Danger Rating System fire indices derived from Weather Research and Forecasting (WRF) model data for the June 2005 Interior Alaska wildfires. *Atmospheric Research* **95**, 290-306.
- Mote PW, Abatzoglou JT, Kunkel KE (2013) Climate: Variability and change in the past and the future. In ‘Climate change in the Northwest: Implications for our landscapes, waters, and

- communities.’ (Eds MM Dalton, PW Mote, AK Snover) pp. 25-40. (Island Press: Washington, DC)
- R Core Team (2013). R: A language and environment for statistical computing. R Foundation for Statistical Computing (Vienna, Austria) ISBN 3-900051-07-0, URL <http://www.R-project.org/>.
- Riley KL, Abatzoglou JT, Grenfell IC, Klene AE, Heinsch FA (2013) The relationship of large fire occurrence with drought and fire danger indices in the western USA, 1984-2008: The role of temporal scale. *International Journal of Wildland Fire* **22**, 894-909. DOI: 10.1071/WF12149.
- Rocky Mountain Research Station and Systems for Environmental Management (RMRS) (2002) Fire Family Plus user’s guide version 3.0. USDA Forest Service, National Fire and Aviation Management Information Systems Team (Boise, ID). Available at http://www.firemodels.org/downloads/firefamilyplus/publications/ffp_v3_user_guide.pdf
- Rothermel RC (1972) A mathematical model for predicting fire spread in wildland fuels. USDA Forest Service, Intermountain Forest and Range Experiment Station Research Paper INT-115. (Ogden, UT)
- Serreze MC, Clark MP, Armstrong RL, McGinnis DA, Pulwarty, RS (1999) Characteristics of the western United States snowpack from snowpack telemetry (SNOTEL) data. *Water Resources Research* **35**. 2145-2160. DOI: 10.1029/1999WR900090.
- Stavros EN, Abatzoglou JT, McKenzie D, Larkin NK (2014) Regional projections of the likelihood of very large wildland fires under a changing climate in the contiguous Western United States. *Climatic Change* **126**, 455–468. DOI: 10.1007/s10584-014-1229-6
- Stocks BJ, Lawson BD, Alexander ME, Van Wagner CE, McAlpine RS, Lynham TJ, Dube DE (1989) The Canadian Forest Fire Danger Rating System: An overview. *The Forestry Chronicle* **65**, 450-457.
- Stocks BJ, Fosberg MA, Lynham TJ, Mearns L, Wotton BM, Yang Q, Jin J-Z, Lawrence K, Hartley GR, Mason JA, McKenney DW (1998) Climate change and forest fire potential in Russian and Canadian boreal forests. *Climatic Change* **38**, 1-13, DOI: 10.1023/A:1005306001055.

- Taylor KE, Stouffer RJ, Meehl GA (2012) An overview of CMIP5 and the experiment design. *Bulletin of the American Meteorological Society* **93**, 485-498, DOI: 10.1175/BAMS-D-11-00094.1.
- Williams AJ, Karoly DJ, Tapper N (2001) The sensitivity of Australian fire danger to climate change. *Climatic Change* **49**, 171-191, DOI: 10.1023/A:1010706116176.
- Williams AP, Seager R, Berkelhammer M, Macalady AK, Crimmins MA, Swetnam TW, Trugman AT, Buenning N, Hryniw N, McDowell NG, Noone D, Mora CI, Rahn T (2014) Causes and implications of extreme atmospheric moisture demand during the record-breaking 2011 wildfire season in the southwestern United States. *Journal of Applied Meteorology and Climatology* **53**, 2671-2684. DOI:10.1175/JAMC-D-14-0053.1.
- Wotton BM (2009) Interpreting and using outputs from the Canadian Forest Fire Danger Rating System in research applications. *Environmental and Ecological Statistics* **16**, 107-131.
- van Wagner CE (1987) Development and structure of the Canadian Forest Fire Weather Index System. Canadian Forest Service Technical Report No. 35.
- Yebra M, Chuvieco E, Riaño D (2008) Estimation of live fuel moisture content from MODIS images for fire risk assessment. *Agriculture and Forest Meteorology* **148**, 523-536. DOI: 10.1016/j.agrformet.2007.12.00

Tables

Table 1. Description of Predictive Services Areas (PSAs) in Oregon and Washington. The third column lists the number of key stations within each PSA, the first common year in which all key stations report data, and the percent of fire season days during 1981-2010 in which all stations are missing data.

Designation	Subregional Area	# of Key RAWS	Elevation Range (m)	Slope Range	Climate Classes	Green-up Date Range
		(year; %msg)				
NW01	Northwestern WA	9 (2001; 7.6%)	610-1036	26-75%	Humid-Wet	Apr 15–Jun 23
NW02	Northwest OR/ Southwest WA	8 (1997; 2.2%)	610-762	26-75%	Humid	Apr 13–May 1
NW03	Central Western OR	8 (1995; 2.1%)	154-1326	41-55%	Humid	Apr 15–May 18
NW04	Southwestern OR	9 (1996; 3.3%)	358-1252	26-55%	Subhumid-Humid	Mar 1–Jun 9
NW05	Central WA	4 (1987; 9.2%)	1116-1382	41-55%	Subhumid	May 1–Jun 15
NW06	North Central OR	6 (1991; 2.0%)	1012-1431	26-44%	Subhumid	Apr 22–Jun 2
NW07	South Central OR	5 (1994; 1.8%)	1385-1798	26-44%	Subhumid	May 13–Jun 1
NW08	North Central WA	4 (1993; 1.5%)	517-1979	26-44%	Subhumid	Apr 7–Jun 28
NW09	Northeast WA	4 (1995; 6.1%)	399-1413	26-44%	Subhumid	Apr 25–May 26
NW10	Southeast WA	3 (2001; 4.1%)	276-771	0-25%	Semi-arid	Apr 6
NW11	Northeast OR	8 (1998; 2.8%)	1158-1850	26-55%	Subhumid	May 15–Jun 16
NW12	Southeast OR	5 (1994; 13.1%)	1341-1984	0-44%	Semi-arid	Apr 22–Jun 1

Table 2. Eighteen CMIP5 global climate models downscaled using the MACA approach.

Model	Center
BCC-CSM1-1	Beijing Climate Center, China Meteorological Administration, China
BCC-CSM1-1-M	Beijing Climate Center, China Meteorological Administration, China
BNU-ESM	College of Global Change and Earth System Science, Beijing Normal University, China
CanESM2	Canadian Centre for Climate Modeling and Analysis
CNRM-CM5	National Centre of Meteorological Research, France
CSIRO-Mk3.6.0	Commonwealth Scientific and Industrial Research Organization/Queensland Climate Change Centre of Excellence, Australia
GFDL-ESM2G	NOAA Geophysical Fluid Dynamics Laboratory, USA
GFDL-ESM2M	NOAA Geophysical Fluid Dynamics Laboratory, USA
HadGEM2-CC*	Met Office Hadley Center, UK
HadGEM2-ES*	Met Office Hadley Center, UK
INM-CM4	Institute for Numerical Mathematics, Russia
IPSL-CM5A-LR	Institute Pierre-Simon Laplace, France
IPSL-CM5A-MR	Institute Pierre-Simon Laplace, France
IPSL-CM5B-LR	Institute Pierre-Simon Laplace, France
MIROC5	Atmosphere and Ocean Research Institute (The University of Tokyo), National Institute for Environmental Studies, and Japan Agency for Marine-Earth Science and Technology
MIROC-ESM	Japan Agency for Marine-Earth Science and Technology, Atmosphere and Ocean Research Institute (The University of Tokyo), and National Institute for Environmental Studies
MIROC-ESM-CHEM	Japan Agency for Marine-Earth Science and Technology, Atmosphere and Ocean Research Institute (The University of Tokyo), and National Institute for Environmental Studies
MRI-CGCM3	Meteorological Research Institute, Japan

*The HadGEM2 models 360-day outputs were both interpolated to 365-day calendars before the MACA downscaling process.

Table 3. Definition of state of weather (SOW) codes based on daily precipitation (PPT) and percent of maximum daily solar insolation ($\%RSDS_{max}$).

SOW	Description	PPT	$\%RSDS_{max}$
0	Clear	$<0.01''$	≥ 91
1	Scattered clouds	$<0.01''$	$73 \leq \%RSDS_{max} < 91$
2	Broken clouds	$<0.01''$	$50 < \%RSDS_{max} < 73$
3	Overcast	$<0.01''$	≤ 50
4	Fog	---	---
5	Drizzle	$0.01'' < PPT < 0.05''$	---
6	Rain	$\geq 0.05''$	---

Table 4. Mean fire season ERC value for each Predictive Services Area (PSA) based on a June 1-October 31 fire season. The first column shows the 1981-2010 observed mean fire season ERC with the multi-model average fire season ERC value for the modeled historical period (1950-2005) in parentheses. Also shown are the multi-model mean projected changes in mean fire season for RCP4.5 and RCP8.5 for the 2020s and 2050s. Highlighted cells indicate that at least three-quarters (14 out of 18) of the models agreed on the sign of change.

PSA	Historical Baseline	2020s		2050s	
		RCP4.5	RCP8.5	RCP 4.5	RCP8.5
NW01	15 (17)	<1	1	2	3
NW02	16 (20)	<1	1	2	3
NW03	22 (24)	<1	1	2	3
NW04	36 (37)	<1	1	2	3
NW05	47 (48)	<1	1	2	3
NW06	50 (50)	<1	1	2	3
NW07	49 (49)	1	1	2	3
NW08	43 (45)	1	1	2	4
NW09	41 (41)	<1	1	2	3
NW10	62 (60)	1	1	2	3
NW11	47 (48)	1	1	2	3
NW12	65 (62)	1	1	2	3

Table 5. Fire season ERC 90th percentile value for each Predictive Services Area (PSA) based on a June 1 through October 31 fire season. The first column shows the 1981-2010 observed value with the multi-model average fire season ERC 90th percentile value for the modeled historical period (1950-2005) in parentheses. Also shown are the multi-model mean projected increases in fire season ERC 90th percentile for RCP4.5 and RCP8.5 for the 2020s and 2050s. Highlighted cells indicate that at least three-quarters (14 out of 18) of the models agreed on the sign of change.

PSA	Historical Baseline	2020s		2050s	
		RCP4.5	RCP8.5	RCP 4.5	RCP8.5
NW01	31 (35)	1	1	3	5
NW02	34 (40)	<1	1	3	4
NW03	42 (43)	<1	1	2	4
NW04	57 (58)	<1	1	2	4
NW05	67 (69)	1	1	4	5
NW06	71 (72)	1	2	3	5
NW07	70 (71)	1	2	3	5
NW08	65 (66)	1	2	4	6
NW09	67 (65)	1	1	4	6
NW10	79 (80)	2	3	4	5
NW11	72 (73)	1	2	4	5
NW12	86 (87)	2	2	3	5

Table 6. Fire season ERC 97th percentile value for each Predictive Services Area (PSA) based on a June 1 through October 31 fire season. The first column shows the 1981-2010 observed value with the multi-model average fire season ERC 97th percentile value for the modeled historical period (1950-2005) in parentheses. Also shown are the multi-model mean projected increases in fire season ERC 97th percentile for RCP4.5 and RCP8.5 for the 2020s and 2050s. Highlighted cells indicate that at least three-quarters (14 out of 18) of the models agreed on the sign of change.

PSA	Historical Baseline	2020s		2050s	
		RCP4.5	RCP8.5	RCP 4.5	RCP8.5
NW01	38 (43)	<1	1	3	6
NW02	42 (48)	<1	1	2	5
NW03	49 (50)	<1	1	3	4
NW04	63 (64)	<1	1	2	4
NW05	73 (76)	1	2	4	6
NW06	76 (78)	1	2	4	6
NW07	75 (78)	1	2	4	6
NW08	73 (74)	1	2	4	7
NW09	75 (73)	1	1	4	6
NW10	83 (85)	2	3	4	6
NW11	79 (81)	1	2	4	5
NW12	91 (94)	2	2	4	6

Table 7. Percent of days exceeding the 90th and 97th percentile ERC value each month during the declared fire season for each Predictive Services Area based on data from 1981 through 2010.

PSA	June		July		August		September		October	
	90 th	97 th	90 th	97 th	90 th	97 th	90 th	97 th	90 th	97 th
NW01	2.0	0.0	5.6	1.2	20.0	5.4	14.4	5.3	3.7	1.0
NW02	0.0	0.0	4.9	0.4	20.2	5.7	14.3	5.3	7.6	1.6
NW03	0.0	0.0	2.6	0.0	17.6	2.6	18.5	8.1	8.7	2.4
NW04	0.0	0.0	2.9	0.0	19.3	4.2	18.2	6.6	6.6	1.6
NW05	0.5	0.0	5.0	0.5	20.5	5.9	14.7	6.2	4.2	0.8
NW06	0.0	0.0	5.8	0.4	24.4	7.0	13.4	4.8	3.0	0.2
NW07	0.0	0.0	2.0	0.1	16.6	4.2	18.2	7.1	8.9	2.4
NW08	1.1	0.0	8.3	1.5	23.4	8.5	13.9	5.0	1.4	0.0
NW09	0.0	0.0	4.1	0.7	24.9	6.6	17.5	5.8	2.2	0.1
NW10	0.0	0.0	9.1	2.5	22.7	6.9	10.4	3.3	1.9	0.0
NW11	0.0	0.0	1.9	0.0	22.4	6.7	17.3	6.0	4.3	0.4
NW12	0.0	0.0	4.8	1.0	19.7	7.8	18.7	5.0	4.9	0.9

Table 8. Annual trend (units/year) in mean ERC value for the fire season and for each month in the fire season for each Predictive Services Area over the analysis period (1981-2010). Trends that are statistically significant at the 95% level are bolded with the Mann-Kendall p-value or the 95% confidence interval in parentheses. Confidence intervals from a bootstrapping procedure are displayed instead of p-values for those time series with significant autocorrelation.

PSA	Fire Season	June	July	August	September	October
NW01	0.00 (1)	-0.10 (0.2319)	0.07 (0.3178)	0.13 (0.3678)	0.21 (0.3110)	0.06 (0.4530)
NW02	0.36 (0.0007)	0.04 (1)	0.40 (0.0060)	0.57 (0.0039)	0.57* (0.0020)	0.15 (0.2319)
NW03	0.33 (0.0102)	-0.04 (0.8028)	0.37 (0.0257)	0.59 (0.0015)	0.49 (0.1250)	0.16 (0.2390)
NW04	0.42 (0.0102)	-0.08 (0.7784)	0.46* (0.0420)	0.60* (0.0022)	0.65* (0.0075)	0.61 (0.0409)
NW05	0.51* (0.0002)	0.38 (0.0811)	0.79 (0.0000)	0.71 (0.0001)	0.86 (0.0003)	0.45 (0.1095)
NW06	0.56 (0.0001)	0.14 (0.3724)	0.65 (0.0015)	0.73 (0.56, 1.28)	0.72* (0.0017)	0.61 (0.0138)
NW07	0.53 (0.43, 1.22)	0.08 (0.7079)	0.49 (0.0168)	0.69 (0.0008)	0.78 (0.0007)	0.67 (0.0270)
NW08	0.74 (0.52, 1.43)	0.41 (0.1338)	0.88 (0.44, 1.25)	0.95 (0.45, 1.36)	0.86* (0.0012)	0.62 (0.0043)
NW09	0.62 (0.0007)	0.21 (0.4325)	0.54 (0.0176)	0.99 (0.0002)	0.84 (0.0052)	0.65 (0.0153)
NW10	0.21 (0.1886)	-0.17 (0.7850)	0.42 (0.0782)	0.60 (0.0131)	0.16 (0.9013)	-0.01 (0.8741)
NW11	0.63 (0.0002)	0.29 (0.1868)	0.68 (0.0012)	0.84 (0.0005)	0.77 (0.0113)	0.67 (0.0048)
NW12	0.02 (0.8025)	-0.56 (0.0578)	0.15 (0.6315)	0.37 (0.0454)	0.35 (0.1131)	-0.22 (0.4536)

*Residuals of linear regression model display non-constant variance (i.e., Breusch-Pagan p-value <0.05); linear model inefficient for this time series

Table 9. Annual trend (units/year) in percent of days greater than the long-term fire season 90th percentile ERC value for the fire season and for each month in the fire season for each Predictive Services Area over the analysis period (1981-2010). Trends that are statistically significant at the 95% level are bolded with the Mann-Kendall p-value or the 95% confidence interval in parentheses. Confidence intervals from a bootstrapping procedure are displayed instead of p-values for those time series with significant autocorrelation.

PSA	Fire Season	June	July	August	September	October
NW01	0.06 (0.92872)	-0.01 (0.6073)	-0.29 (0.7324)	0.76 (-0.11, 0.49)	0.35 (0.5980)	-0.15 (0.8211)
NW02	0.68* (0.0039)	0 ---	0.23 (0.0254)	1.9* (0.30, 1.03)	1.03* (0.0317)	0.14 (0.2411)
NW03	0.46 (0.0355)	0 ---	0.22 (0.0768)	1.57* (0.16, 0.94)	0.64 (0.0418)	-0.20 (0.5954)
NW04	0.45 (0.0110)	0 ---	0.14 (0.0757)	0.98 (0.0239)	0.98 (0.0645)	0.31 (0.0789)
NW05	0.75* (0.0001)	-0.03 (0.6757)	0.52* (0.0062)	1.81* (0.0006)	1.21* (0.0003)	0.27 (0.0761)
NW06	0.72* (0.0001)	0 ---	0.65* (0.0119)	1.82* (0.0006)	0.86 (0.0058)	0.23 (0.0628)
NW07	0.67* (0.0005)	0 ---	0.11 (0.1716)	1.36* (0.0006)	1.61* (0.0002)	0.26 (0.1007)
NW08	1.01* (0.61, 1.59)	0.19* (0.0378)	0.95* (0.42, 1.17)	2.49* (0.67, 1.64)	1.25* (0.0063)	0.13 (0.2802)
NW09	0.61* (0.34, 1.16)	0 ---	0.08 (0.3291)	1.70* (0.0324)	1.25* (0.38, 1.07)	0.01 (0.5610)
NW10	0.46 (0.1189)	0 ---	0.24 (0.2321)	1.78* (0.0106)	0.40 (0.2538)	-0.21 (0.5133)
NW11	0.62* (0.0012)	0 ---	0.23 (0.2017)	1.69* (0.0027)	1.05* (0.0131)	0.16 (0.1818)
NW12	0.33 (0.1546)	0 ---	0.17 (0.8967)	0.85 (0.3342)	0.70 (0.1865)	-0.08 (1)

*Residuals of linear regression model display non-constant variance (i.e., Breusch-Pagan p-value <0.1); linear model inefficient for this time series

Table 10. Annual trend (units/year) in percent of days greater than the long-term fire season 97th percentile ERC value for the fire season and for each month in the fire season for each Predictive Services Area over the analysis period (1981-2010). Trends that are statistically significant at the 95% level are bolded with the Mann-Kendall p-value or the 95% confidence interval in parentheses. Confidence intervals from a bootstrapping procedure are displayed instead of p-values for those time series with significant autocorrelation.

PSA	Fire Season	June	July	August	September	October
NW01	0.03 (0.7770)	0 ---	-0.17 (0.2480)	0.19 (0.5026)	0.32 (0.7373)	-0.10 (0.3700)
NW02	0.23 (0.1355)	0 ---	-0.06 (0.2480)	0.70* (0.0225)	0.56* (0.0189)	-0.08 (0.6929)
NW03	0.08 (0.2591)	0 ---	0 ---	0.29* (0.0150)	0.21 (0.4549)	-0.13 (0.5123)
NW04	0.12 (0.0929)	0 ---	0 ---	0.12 (0.6950)	0.48 (0.0229)	0.07 (0.2313)
NW05	0.28* (0.0019)	0 ---	0.04 (0.6182)	0.68* (0.0025)	0.63* (0.0050)	0.06 (0.4148)
NW06	0.26* (0.0018)	0 ---	0.05 (0.1145)	0.79* (0.0008)	0.46* (0.0303)	0.02 (0.3390)
NW07	0.25* (0.0014)	0 ---	-0.002 (0.9080)	0.40 (0.0768)	0.63* (0.0048)	0.22* (0.0284)
NW08	0.32 (0.0003)	0 ---	0.20 (0.0139)	0.87 (0.0037)	0.55* (0.0194)	0 ---
NW09	0.22* (0.0431)	0 ---	0.04 (0.8681)	0.61* (0.0426)	0.46 (0.0554)	-0.01 (0.6440)
NW10	0.23 (0.1192)	0 --	0.002 (0.6955)	0.85* (0.0099)	0.26 (0.4110)	0 ---
NW11	0.25* (0.0035)	0 ---	0 ---	0.71* (0.0049)	0.54 (0.0761)	0.02 (0.4075)
NW12	0.03 (0.6994)	0 ---	0.02 (1)	0.13 (0.6286)	0.09 (0.7451)	-0.09 (0.7794)

*Residuals of linear regression model display non-constant variance (i.e., Breusch-Pagan p-value <0.1); linear model inefficient for this time series

Table 11. Multi-model mean change in percent (%) of days exceeding the **90th percentile** ERC value each month during the declared fire season for each Predictive Services Area for the **2050s** under both scenarios (RCP4.5 and RCP8.5). Highlighted cells indicate that at least three-quarters (14 out of 18) of the models agreed on the sign of change.

PSA	Fire Season		June		July		August		September		October	
	4.5	8.5	4.5	8.5	4.5	8.5	4.5	8.5	4.5	8.5	4.5	8.5
NW01	5.1	7.6	1.6	3.2	8.5	12.3	10.7	14.4	3.6	7.3	0.8	0.7
NW02	4.2	6.3	1.1	1.7	6.6	10.1	9.2	12.2	3.5	6.5	0.7	0.8
NW03	4.3	6.4	1	1.2	6.6	10	9.7	12.9	3.3	6.5	0.6	1
NW04	4.2	6.3	1.2	1.1	5.5	9	9.5	11.5	4.1	7.5	0.6	2.3
NW05	5.7	8.8	1.8	2.5	8.6	12.7	11.2	16.8	5.6	10.1	1.1	1.7
NW06	5.7	8.4	1.8	2.3	8	11.6	11.9	15.7	5.9	10.6	1	1.8
NW07	5.4	7.5	1.9	2.1	6.4	9.6	12.4	13.5	5.2	9.2	1	3.1
NW08	6.5	9.7	2.9	4.3	9.4	13.4	13	19	6.4	10.7	0.9	1.2
NW09	5.4	7.6	1.6	1.7	6.8	9.5	11.6	15.4	5.6	9.9	1	1.4
NW10	7.4	11.1	2.1	2.5	9.9	15.9	16.2	22.5	7.8	13.5	0.6	1.1
NW11	5.3	7.1	0.9	0.9	6.3	8.3	12.5	13.3	5.7	10.7	1.2	2.3
NW12	5.7	7.7	0.9	0.9	7.1	10	13.3	14.1	5.8	10.7	1.2	2.6

Table 12. Multi-model mean change in percent (%) of days exceeding the **90th percentile** ERC value each month during the declared fire season for each Predictive Services Area for the **2020s** under both scenarios (RCP4.5 and RCP8.5). Highlighted cells indicate that at least three-quarters (14 out of 18) of the models agreed on the sign of change.

PSA	Fire Season		June		July		August		September		October	
	4.5	8.5	4.5	8.5	4.5	8.5	4.5	8.5	4.5	8.5	4.5	8.5
NW01	0.9	1.9	0.4	0.6	1.5	2.7	1.1	3.6	1.8	2	-0.5	0.5
NW02	0.5	1.5	0.2	0.1	0.4	1.5	1.7	3.1	0.4	2.4	-0.4	0.3
NW03	0.5	1.7	0	0.2	0	1.4	3	4.5	-0.1	2.5	-0.6	-0.3
NW04	0.6	2	0	0.2	-0.2	1.9	3.4	5.2	0.3	2.8	-0.6	-0.1
NW05	1.3	2.1	0.8	0.4	1.5	2.3	2.6	3.8	1.3	3.1	0.1	0.7
NW06	1.6	3	0.5	0.7	0.8	2.8	4.9	6.4	1.7	5	0.2	0.2
NW07	2	3.3	0.2	0.7	0.9	2.5	6.3	8.2	2.5	5	0.2	0
NW08	1.8	2.9	1.2	1	1.6	3.2	3.8	6.3	2.4	3.6	0.2	0.5
NW09	1.3	1.7	0.8	0.3	1.1	1.2	2.9	3.9	1.6	2.7	0.2	0.4
NW10	3	5.2	0.9	0.8	3.1	6.6	7.4	12	3.1	6.1	0.2	0.3
NW11	1.6	2.6	0.1	0.3	1	2.1	4.3	6.1	1.9	4.2	0.4	0.1
NW12	2.4	3.5	0.2	0.4	1.6	3.9	6.1	7.9	2.9	5.1	0.9	0.3

Table 13. Multi-model mean change in percent (%) of days exceeding the **97th percentile** ERC value each month during the declared fire season for each Predictive Services Area for the **2050s** under both scenarios (RCP4.5 and RCP8.5). Highlighted cells indicate that at least three-quarters (14 out of 18) of the models agreed on the sign of change.

PSA	Fire Season		June		July		August		September		October	
	4.5	8.5	4.5	8.5	4.5	8.5	4.5	8.5	4.5	8.5	4.5	8.5
NW01	2.8	4.6	0.7	1.4	3.9	7.1	6.4	9	1.9	4.8	0.9	0.7
NW02	1.9	3.2	0.5	0.7	2.3	4.9	4.7	5.9	1.5	3.8	0.6	0.5
NW03	2.1	3.3	0.5	0.5	2.8	5.2	5.5	6.4	1.4	4.1	0.3	0.4
NW04	2.2	3.5	0.4	0.4	2.9	5	5.4	6	2.2	4.7	0.3	1.3
NW05	3.6	5.8	0.9	1.1	5	8.2	7.6	11.8	3.8	6.5	0.8	1.1
NW06	3.6	5.4	1	1	4.6	7.6	8.3	11.1	3.3	6.3	0.4	0.8
NW07	3.2	4.5	1	0.7	3.7	6.1	7.5	8.9	3.1	5.2	0.6	1.6
NW08	4.2	6.6	1.5	2.1	6.2	9.7	9.5	14.5	3.2	5.8	0.4	0.6
NW09	3.5	5.4	0.4	0.8	3.7	6.4	9.1	12.4	4	6.4	0.3	0.6
NW10	5	7.9	1	0.9	7.3	11.7	12.3	19	3.9	7.5	0	0.3
NW11	3.3	4.6	0.3	0.3	3.3	5.1	8.9	11.1	3.3	5.9	0.4	0.8
NW12	4	5.8	0.4	0.4	4.1	7	11	13.3	3.7	6.8	0.5	1

Table 14. Multi-model mean change in percent (%) of days exceeding the **97th percentile** ERC value each month during the declared fire season for each Predictive Services Area for the **2020s** under both scenarios (RCP4.5 and RCP8.5). Highlighted cells indicate that at least three-quarters (14 out of 18) of the models agreed on the sign of change.

PSA	Fire Season		June		July		August		September		October	
	4.5	8.5	4.5	8.5	4.5	8.5	4.5	8.5	4.5	8.5	4.5	8.5
NW01	0.3	0.7	0.3	0.2	0.6	0.8	0.4	1.6	0.6	0.9	-0.3	0.2
NW02	-0.1	0.5	0.2	0.1	-0.2	0.3	0	1.2	0.1	0.8	-0.4	-0.1
NW03	0	0.7	0.1	0.1	-0.2	0.2	0.5	2.2	-0.1	1.2	-0.3	-0.2
NW04	0.3	1	0.1	-0.1	0.2	0.9	1	2.6	0.4	1.9	-0.2	-0.1
NW05	0.6	1.1	0.5	0.2	0.8	1.5	1.3	2.2	0.5	1.7	0	0.2
NW06	0.9	1.7	0.3	0.2	0.4	1.4	2.7	4.5	1.2	2.5	0.1	0
NW07	1.1	1.7	0.1	0.2	0.6	1.4	2.9	4.5	1.5	2.3	0.2	0
NW08	1.1	1.6	0.8	0.4	1.7	1.9	2.2	4	0.6	1.5	0.1	0
NW09	0.9	1.1	0.3	0.1	1.3	0.8	2.5	3.1	0.2	1.2	0.1	0.1
NW10	1.7	3.2	0.5	0.3	2	4	4.8	8.6	1.3	2.8	0.1	0
NW11	0.9	1.5	0.1	0	0.9	1.3	2.8	4.6	0.8	1.8	0	0
NW12	1.5	2.3	0.2	0.1	1.4	2.6	4	6.1	1.8	2.6	0.3	0.1

Table 15. Fraction of variance in future ERC changes explained by future changes in tmax, tmin, rmax, rmin, ppt, pdur, rsds, and sow during the fire season for each PSA. Values are the square of Pearson's correlation coefficient.

PSA	TMAX	TMIN	RMAX	RMIN	PPT	PDUR	RSDS	SOW
NW01	0.36	0.39	0.56	0.51	0.18	0.27	0.18	0.30
NW02	0.36	0.38	0.50	0.48	0.28	0.34	0.15	0.31
NW03	0.36	0.36	0.43	0.47	0.29	0.35	0.22	0.26
NW04	0.34	0.29	0.38	0.40	0.19	0.31	0.26	0.33
NW05	0.32	0.34	0.47	0.59	0.02	0.01	0.20	0.30
NW06	0.28	0.31	0.44	0.53	0.16	0.17	0.33	0.43
NW07	0.22	0.20	0.39	0.41	0.08	0.11	0.20	0.40
NW08	0.25	0.30	0.54	0.64	0.05	0.04	0.19	0.40
NW09	0.21	0.25	0.50	0.62	0.04	0.07	0.23	0.46
NW10	0.19	0.21	0.45	0.57	<0.01	0.01	0.17	0.32
NW11	0.15	0.14	0.31	0.49	0.03	0.06	0.08	0.37
NW12	0.09	0.11	0.29	0.44	<0.01	<0.01	0.06	0.30

Figures

NWCC PSA Stations

Figure 1. Predictive Service Areas and the key Remote Automated Weather Stations in Washington and Oregon.

Figure 2. 90th percentile ERC by PSA calculated over fire season months during 1981-2010.

Mean ERC Climatology

Figure 3. Seasonal ERC Climatology for downscaled data

Figure 4. Mean ERC by PSA averaged over fire season months (Jun-Oct) during the modeled historic baseline (1950-2005), 2020s, and 2050s, for RCP4.5 and RCP8.5. Boxes display the mean and plus/minus one standard deviation and whiskers display the extremes of the 17 GCMs.

Figure 5. Future changes in fire season 90th percentile ERC for RCP8.5 for the 2020s (top) and 2050s (bottom) compared to the historical baseline (1950-2005) for each PSA. Shaded regions indicate that more than 75% (at least 14 out of 18) of the models agreed on the sign of change.

Figure 6. Future changes in % of fire season days exceeding the baseline 90th percentile ERC for RCP8.5 for the 2020s (top) and 2050s (bottom) compared to the historical baseline (1950-2005) for each PSA. Shaded regions indicate that more than 75% (at least 14 out of 18) of the models agreed on the sign of change.

% August Days Greater Than 90th Percentile ERC

Figure 7. Percent of August days exceeding the historic fire season ERC 90th percentile value by PSA during the modeled historic baseline (1950-2005), 2020s, and 2050s, for RCP4.5 and RCP8.5. Boxes display the mean and plus/minus one standard deviation and whiskers display the extremes of the 18 GCMs.

Appendix A: Seasonal ERC Climatology

Figures A1-A12 display the daily climatology of ERC calculated over 1981-2010 for each PSA. Plotted are the mean (solid black), median (dashed black), lower and upper quartiles (blue), and maximum (green). Vertical lines demarcate the start (June 1) and end (October 31) of the fire season. Horizontal red lines display the long-term fire season mean, 90th percentile and 97th percentile values.

Figure A1. NW01 ERC Climatology

Figure A2. NW02 ERC Climatology

NW02 ERC Climatology (1981-2010)

Figure A3. NW03 ERC Climatology

NW03 ERC Climatology (1981-2010)

Figure A4. NW04 ERC Climatology

Figure A5. NW05 ERC Climatology

Figure A6. NW06 ERC Climatology

Figure A7. NW07 ERC Climatology

Figure A8. NW08 ERC Climatology

Figure A9. NW09 ERC Climatology

Figure A10. NW10 ERC Climatology

Figure A11. NW11 ERC Climatology

Figure A12. NW12 ERC Climatology

Appendix B: Percent of Days Exceeding 90th Percentile ERC

Figures B1-B5 display the percent of days in each fire season month (June-October) during 1981-2010 that exceed the 1981-2010 fire season 90th percentile ERC threshold.

Figure B1. Percent of June days greater than 90th percentile ERC threshold.

Figure B2. Percent of July days greater than 90th percentile ERC threshold.

Figure B3. Percent of August days greater than 90th percentile ERC threshold.

Figure B4. Percent of September days greater than 90th percentile ERC threshold.

Figure B5. Percent of October days greater than 90th percentile ERC threshold.

Appendix C: Trends in Mean Fire Season ERC

Figures C1-C12 display the annual time series and linear least square trends (red) in fire season mean ERC from 1981-2010 for each PSA. The slope of the trend in ERC units per year along with the Mann-Kendall p-value is noted in the top left corner.

Figure C1. NW01 Fire Season Mean ERC. There are no significant trends for the fire season or for any month of the fire season.

Figure C2. NW02 Fire Season Mean ERC. There is a significant increasing trend in the fire season ERC mean of about 0.4/year. Similar increasing trends are seen in July, August, and September.

Figure C3. NW03 Fire Season Mean ERC. There is a significant increasing trend in the fire season ERC mean of about 0.3/year. Similar increasing trends are seen in July and August.

Figure C4. NW04 Fire Season Mean ERC. There is a significant increasing trend in the fire season mean of about 0.4/year. Similar significant increasing trends are seen in July, August, September, and October.

Figure C5. NW05 Fire Season Mean ERC. There is a significant increasing trend in the fire season ERC mean of about 0.5/year. Similar increasing trends are seen in July, August, and September.

Figure C6. NW06 Fire Season Mean ERC. There is a significant increasing trend in the fire season ERC mean of about 0.6/year. Similar increasing trends are seen in all fire season months except June.

Figure C7. NW07 Fire Season Mean ERC. There is a significant increasing trend in the fire season ERC mean of about 0.5/year. Similar increasing trends are seen in all fire season months except June.

Figure C8. NW08 Fire Season Mean ERC. There is a significant increasing trend in the fire season ERC mean of about 0.7/year. Similar increasing trends are seen in all fire season months except June.

Figure C9. NW09 Fire Season Mean ERC. There is a significant increasing trend in the fire season ERC mean of about 0.6/year. Similar increasing trends are seen in all fire season months except June.

Figure C10. NW10 Fire Season Mean ERC. There is no significant trend in the mean fire season ERC, but there is a positive and significant trend in August of about 0.6/year.

Figure C11. NW11 Fire Season Mean ERC. There is a significant increasing trend in the fire season ERC mean of about 0.6/year. Similar significant increasing trends are seen in all fire season months except June.

Figure C12. NW12 Fire Season Mean ERC. There are no significant increasing trends in mean fire season ERC, but there is a positive and significant trend in August of about 0.4/year.

Appendix D: Trends in % of Days Exceeding 90th Percentile ERC

Figures D1-D12 display the annual time series from 1981-2010 of the percent of fire season days greater than the 90th and 97th percentile thresholds for 1981-2010 for each PSA.

Figure D1. Percent of fire season days greater than the 90th (top) and 97th (bottom) percentile thresholds for NW01. There are no significant trends for the fire season or for any month of the fire season. The two years with the largest percent of fire season days above the 90th and 97th percentiles are 2003 and 1985 with the majority of those days occurring in July and August.

Figure D2. Percent of fire season days greater than the 90th (top) and 97th (bottom) percentile thresholds for NW02. There is a significant positive trend in percent of fire season days greater than the 90th, but it appears to be dominated by generally larger values in the early- to mid-2000s that occur mostly in August and September. July, August, and September also have significant positive trends. There are also significant positive trends in percent of days above the 97th percentile in August and September, but not for the entire fire season.

Figure D3. Percent of fire season days greater than the 90th (top) and 97th (bottom) percentile thresholds for NW03. The significant positive trend in percent of fire season days greater than the 90th appears to be dominated by generally larger percentages in the early- to mid- 2000s that occur mostly in August. August and September also have significant positive trends. Only in August is the positive trend in percent of days above the 97th percentile significant.

Figure D4. Percent of fire season days greater than the 90th (top) and 97th (bottom) percentile thresholds for NW04. The positive trend in percent of fire season days greater than the 90th suggests the percent of days is increasing with contributions mainly from August and September. The August trend is also significant. Only the September trend is significantly increasing for percent of days above the 97th percentile.

Figure D5. Percent of fire season days greater than the 90th (top) and 97th (bottom) percentile thresholds for NW05. There is a positive trend in percent of fire season days greater than the 90th with significant positive trends also in July, August, and September. Percent of days above the 97th percentile has an increasing trend during the fire season and August and September, though virtually no days exceed the 97th percentile prior to 1993.

Figure D6. Percent of fire season days greater than the 90th (top) and 97th (bottom) percentile thresholds for NW06. There is a positive trend in percent of fire season days greater than the 90th and 97th, similarly in August and September, and in July only for the 90th percentile.

Figure D7. Percent of fire season days greater than the 90th (top) and 97th (bottom) percentile thresholds for NW07. There is a positive trend in percent of fire season days greater than the 90th, similarly in August and September. There is also a significant positive trend in percent of fire season days exceeding the 97th percentile, similarly in September and October.

Figure D8. Percent of fire season days greater than the 90th (top) and 97th (bottom) percentile thresholds for NW08. The positive trend in percent of fire season days greater than the 90th is dominated by markedly larger percentages during the 2000s compared to the rest of the period. The percent of August days greater than the 90th increases in a similar fashion. Significant positive trends occur in all fire season months except October. Significant positive trends in percent of days greater than the 97th percentile are seen for the fire season and the months of July, August, and September.

Figure D9. Percent of fire season days greater than the 90th (top) and 97th (bottom) percentile thresholds for NW09. There are significant positive trends in percent of fire season days greater than the 90th and 97th percentiles with similar trends in August and for the 90th percentile, also in September.

Figure D10. Percent of fire season days greater than the 90th (top) and 97th (bottom) percentile thresholds for NW10. There are no significant trends in the percent of fire season days greater than the 90th or 97th percentiles, but the positive trends in August are significant.

Figure D11. Percent of fire season days greater than the 90th (top) and 97th (bottom) percentile thresholds for NW11. There are increasing trends in percent of fire season days greater than the 90th and 97th percentiles, similarly in August and for the 90th also in September.

Figure D12. Percent of fire season days greater than the 90th (top) and 97th (bottom) percentile thresholds for NW12. There are no significant trends in percent of days greater than the 90th or 97th percentiles for the fire season or for any month.

Appendix E: Seasonal Climate Projections

Figure E1. Daily climatology of TMAX for modeled historical baseline (black), 2020s (blue), and 2050s (orange) for RCP4.5 (dashed) and RCP8.5 (solid) for each PSA.

Mean TMAX Climatology

Figure E2. As in Figure E1, but for TMIN.

Mean TMIN Climatology

Figure E3. As in Figure E1, but for RMAX.

Mean RMAX Climatology

Figure E4. As in Figure E1, but for RMIN.

Mean RMIN Climatology

Figure E5. As in Figure E1, but for PPT.

Mean PPT Climatology

Appendix F: Change in ERC vs. Change in Contributing Variables

Figure F1. Change in fire season mean ERC as a function of change in fire season mean RMAX for RCP8.5 (plus), RCP4.5 (circle), 2020s (blue), and 2050s (orange) for each PSA.

Figure F2. As in Figure F1, but for RMIN.

Δ RMIN vs Δ ERC

Figure F3. As in Figure F1, but for TMAX.

Δ TMAX vs Δ ERC

Figure F4. As in Figure F1, but for TMIN.

Δ TMIN vs Δ ERC

Figure F5. As in Figure F1, but for PPT.

Δ PPT vs Δ ERC

Figure F6. As in Figure F1, but for PDUR.

Δ PDUR vs Δ ERC

Figure F7. As in Figure F1, but for RSDS.

Δ RSDS vs Δ ERC

Figure F8. As in Figure F1, but for SOW.

Δ SOW vs Δ ERC