The City Library Association, Springfield, Mass.
LITERATURE: ART: SCIENCE.

No. 64. 10-02-2M.

V.6

FIELD COLUMBIAN MUSEUM

PUBLICATION 105

ZOOLOGICAL SERIES

VOL. VI

A

CHECK LIST

OF

MAMMALS

OF THE

NORTH AMERICAN CONTINENT

THE

WEST INDIES

AND THE

NEIGHBORING SEAS

BY

DANIEL GIRAUD ELLIOT, F. R. S. E., ETC.

Curator of Department.

CHICAGO, U. S. A.

1905

PUBLICATION
OF THE
FIELD COLUMBIAN MUSEUM

ZOÖLOGICAL SERIES
VOL. VI.

CHICAGO, U. S. A.

1905

All rights reserved

STACK.

570.6

F45

v.6

159900¹²/₀₅

Y8A88U Y710

22AM 010700842

DISCARDED

by

THE CITY LIBRARY ASSOCIATION
SPRINGFIELD, MASSACHUSETTS

279
23

OVIBOS MOSCHATUS NIPHÆCUS ELLIOT.
THE BLACK MUSK-OX.

FIELD COLUMBIAN MUSEUM

PUBLICATION 105

ZOOLOGICAL SERIES

VOL. VI

A
CHECK LIST
OF
MAMMALS
OF THE
NORTH AMERICAN CONTINENT
THE
WEST INDIES
AND THE
NEIGHBORING SEAS

BY

DANIEL GIRAUD ELLIOT, F. R. S. E., ETC.

Curator of Department.

CHICAGO, U. S. A.

1905

PREFACE.

The present volume is supposed to include all the species and races of Mammals (that have been described up to the date of publication) which inhabit the North American Continent from the Arctic Ocean to the Province of Cauca in Colombia, South America, together with those of the West India Islands, and the adjoining seas. These are arranged systematically according to the author's views, and descriptions have been given of those forms that have received names since the two previous works (the Synopsis of North American Mammals, and the Land and Sea Mammals of Middle America and the West Indies) were published.

In the works above named, in order that their bulk should not be unduly increased, generic names were omitted from the synonymy of the species and races. These have been supplied in the present volume, thus making all references complete, and increasing their usefulness for those who may have occasion to consult them. Mammalian nomenclature being still in a transitory condition, names that have perhaps become familiar to many have been obliged, in certain cases, to give place to others less known on account of either the same name having been previously employed in other branches of Zoölogy, or having been antedated by a different one. Changes, therefore, are continually occurring, as this volume will testify, and new names in various instances have taken the place of those employed even so recently as in the lately published volumes on the Mammals of Middle America and the West Indies. This Check List consequently should always be consulted for the latest accepted name, from that of the Family down to the subspecies.

The present volume does not profess to be more than its title implies, a CHECK LIST of all known North American Mammals, without descriptions, save of those named forms not included in previous volumes, but with a moderate amount of synonymy added, and the type locality and geographical distribution in every instance given, if known. For all other information regarding the Orders, Families, Genera, species, and races, the author's other published works on the same subject should be consulted. The Check List is merely the natural sequence of the three previous volumes, and concludes the whole work as originally conceived.

Great care has been taken to correct all errors, typographical, as well as those of other kinds, which have been observed, and although the author is well aware that a complete elimination of these is practically impossible in works of this description, yet it is hoped that in this volume errors and omissions have been reduced to a minimum.

The number of species enumerated in the present list is 1,308, + No. 936a = 1,309, and 676 races of species found within the geographical limits embraced in the work; added to which are 22 races of exotic species, originating in South America. These all make a grand total of 2,007 species and races with a recognized standing at the present time, inhabiting the North American Continent, with its islands and seas. Many groups of these have never been critically examined, or the specific or racial value of their members clearly determined, and when this is done, doubtless the total number will be considerably reduced and the synonymy correspondingly increased. The *Furor specificus* has probably reached its zenith, and the Sober Second Thought, it is hoped, may soon institute its benign sway.

The Index is intended to answer in a certain degree for the two previous works on the Mammals of the North American Continent, with its islands and seas, as well as for the present volume, and the number of the page given under its respective head, when indicated by heavy type, is that on which the genus, species, or race is described. As probably new names will be bestowed on specimens from time to time, it is proposed to issue at stated intervals, as may be deemed expedient, a list of these forms with descriptions, and such information concerning them as is given of those contained in the volumes already published, and any additional knowledge of the older and longer known forms that may have been gained will be also included, thus keeping the entire work regularly up to date.

My thanks are due to the Assistant Curators of the Department of Zoölogy in this institution, Dr. S. E. Meek and Mr. W. J. Gerhard, for much valuable aid in reading the proofs and constructing the Index; and also to my Assistant, Mr. E. B. Chope.

D. G. E.

1st July, 1905.

ERRATA.

- Page 25, 19th line from bottom, for *Delphinus phocæna*, read *Delphinus phocæna*.
- Page 32, 14th line from bottom, for *Delphinapteris borealis*, read *Delphinapterus borealis*.
- Page 33, 14th line from top, for 1904, read 1901.
- Page 64, 3d line from top, for *Sciurus collæi*, read *Sciurus colliæi*.
- Page 76, top line, for *Sciurus adolphei dorsalis*, read *Sciurus adolphii dorsalis*.
- Page 123, 4th line from bottom, for Suppl., read Append.
- Page 125, 25th line from top, for Suppl., read Append.
- Page 319, 5th line from bottom, for *Heteromys phæurus*, read *Heteromys phæura*.
- Page 420, 22d line from top, for *Mustela americana brumalis*, read *Mustela atrata brumalis*; and 12th line from bottom, insert *Mustela atrata actiosa* before Elliot, Syn. N. Am. Mamm. etc., and begin separate paragraph.
- Page 473, 12th line from top, for Italy, read Ital.
- Page 495, 8th line from bottom, for Mag. Zoöl. Bot., read Mag. Zoöl. & Bot.

CLASS MAMMALIA.

Order I. **Marsupialia. Marsupials.**

Family I. **Didelphyidae. Opossums.**

Genus 1. **Chironectes.**

Chironectes Illig., Prodr. Syst. Mar. et Av., 1811, p. 76. Type

Lutra minima Zimmermann.

Memina G. Fisch., Zoogn., III, 1814, p. 611.

1. **Chironectes minimus** (Zimmermann).

Lutra minima Zimm., Geog. Gesch., II, 1780, p. 317.

Mustela paraguensis Kerr, Linn. Anim. Kingd., 1792, p. 172.

Mustela guianensis Kerr, Linn. Anim. Kingd., 1792, p. 174.

Didelphis memina Cuv., Tabl. Élément., 1798, p. 125.

Lutra sarcovienna Shaw, Gen. Zoöl., I, Pt. II, 1809, p. 447.

Chironectes variegatus Illig., Abh. Ak. Wiss. Berl., 1811, p. 107.

Chironectes palmata Cuv., Règn. Anim., I, 1817, p. 174.

Chironectes yapock Desm., Mamm., I, 1820, p. 261.

Chironectes minimus Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., IV, Pt. I, 1904, p. 3, figs. 1, 1. Zoöl. Ser.

WATER OPOSSUM, YAPOCK. Zarro de Agua in Costa Rica; Tlacuazin de Agua in Guatemala.

Type locality. Guiana.

Geogr. Distr. Guatemala, south through Central America to southern Brazil.

Genus 2. **Marmosa.**

Marmosa Gray, Lond. Med. Repos., xv, 1821, p. 308. Type *Didelphis murina* Linnæus.

Micoureus Less., Tabl. Règn. Anim., Mamm., 1842, p. 186.

Asagis and *Notagopus* Glog., Hand- u. Hilfsb. Naturg., I, 1841, p. 82.

Thylamys (sic) Gray, List. Mamm. Brit. Mus., 1843, p. 101.
Grymæomys Burm., Syst. Uebers. Thiere Bras., 1, 1854, p. 135.
Cuica Liais, Climats, Géol., Faun., Géog., et Botan. Brésil, 1872,
 p. 427.

2. *Marmosa murina* (Linnæus).

Didelphis! murina Linn., Syst. Nat., 1, 1758, p. 55.
Didelphis! dorsigera Linn., Syst. Nat., 1, 1758, p. 55.
Didelphys cayopollin Schreb., Säugth., III, 1778, p. 544, pl. CXLVIII.
Didelphys macrotarsus Wagn., Archiv f. Naturg., VIII, 1842, p. 359.
 (nec Schreber.)
Didelphys microtarsus Wagn., Archiv f. Naturg., VIII, 1842, p. 359.
 (nec Schreber.)
Didelphys imparida Tschudi, Faun. Peruana, 1844, p. 149, pl. IX.
Didelphys ruscuius Cap., Schomb. Brit. Guiana, III, 1848, p. 778.
Grymæomys murinus Burm., Erläut. Faun. Bras., 1856, pp. 80-81.
Grymæomys dorsiger Burm., Erläut. Faun. Bras., 1856, pp. 80-81.
Grymæomys imparida Burm., Erläut. Faun. Bras., 1856, pp. 80-81.
Didelphys murinus True, Proc. U. S. Nat. Mus., VII, 1885, p. 587.
Marmosa murina Thomas, Ann. Mag. Nat. Hist., 6th Ser., 1895,
 p. 58. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M.
 Pub., IV, Pt. 1, 1904, p. 5. Zoöl. Ser.

MURINE OPOSSUM. *Tlacuazin Raton* in Guatemala.

Type locality. Unknown.

Geogr. Distr. Central America to Brazil.

a. *Marmosa murina mexicana* Merriam.

Marmosa murina mexicana Merr., Proc. Biol. Soc. Wash., XI,
 1897, p. 44. Elliot, Mamm. Middle Amer. & W. Indies,
 F. C. M. Pub., IV, Pt. 1, 1904, p. 6. Zoöl. Ser.

MEXICAN MURINE OPOSSUM.

Type locality. Juquila, State of Oaxaca, Mexico.

Geogr. Distr. States of Oaxaca and Chiapas, Mexico.

3. *Marmosa canescens* (Allen).

Didelphis! (Micoureus) canescens Allen, Bull. Am. Mus. Nat. Hist.
 N. Y., 1893, p. 235.

Marmosa canescens Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1897,
 p. 58. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M.
 Pub., IV, Pt. 1, 1904, p. 6. Zoöl. Ser.

ASHY OPOSSUM.

Type locality. Santo Domingo de Guzman, Isthmus of Tehuantepec, Mexico.

Geogr. Distr. Hacienda Magdalena, State of Colima, through States of Michoacan, Guerrero, and Oaxaca, to the Isthmus of Tehuantepec.

4. *Marmosa sinaloæ* Allen.

Marmosa sinaloæ Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1898, p. 143. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 6. Zoöl. Ser.

SINALOA OPOSSUM.

Type locality. Tatemales, State of Sinaloa, Mexico.

Geogr. Distr. State of Sinaloa, Mexico.

5. *Marmosa cinerea* (Temminck).

Didelphis! cinerea Temm., Monog. Mamm., I, 1827, p. 46.

Didelphys cinerea True, Proc. U. S. Nat. Mus., VII, 1885, p. 587.

Marmosa cinerea Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1897, p. 43. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 7, fig. 11. Zoöl. Ser.

GRAY OPOSSUM.

Type locality. Brazil.

Geogr. Distr. Costa Rica to Brazil.

6. *Marmosa insularis* Merriam.

Marmosa insularis Merr., Proc. Biol. Soc. Wash., XII, 1898, p. 14.

Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 7. Zoöl. Ser.

MARIA MADRE ISLAND OPOSSUM.

Type locality. Maria Madre Island, State of Jalisco, Mexico.

Geogr. Distr. Tres Marias Islands, State of Jalisco, Mexico.

7. *Marmosa oaxacæ* Merriam.

Marmosa oaxacæ Merr., Proc. Biol. Soc. Wash., XI, 1897, p. 43.

Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 8. Zoöl. Ser.

OAXACA OPOSSUM.

Type locality. City of Oaxaca, State of Oaxaca, Mexico.

Geogr. Distr. Range unknown. "Sonoran fauna of highlands of the State of Oaxaca, Mexico."

8. *Marmosa fulviventer* Bangs.

Marmosa fulviventer Bangs, Amer. Nat., XXXV, 1901, p. 632.

Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 8. Zoöl. Ser.

FULVOUS-BELLIED OPOSSUM.

Type locality. San Miguel Island, Bay of Panama.

Geogr. Distr. Known only from San Miguel Island.

Genus 3. **Caluromys.**

Caluromys Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1900, p. 189.
Type *Didelphis! philander* Linnæus.

9. **Caluromys alstoni** Allen.

Caluromys alstoni Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1900,
p. 189. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M.
Pub., IV, Pt. 1, 1904, p. 9, pl. 1. Zool. Ser.

Didelphys cinereus Alston, Biol. Centr. Amer., Mamm., 1, 1880,
p. 199, pl. XXI. (nec Temm.)

ALSTON'S OPOSSUM.

Type locality. Tres Rios, Costa Rica.

Geogr. Distr. Costa Rica, Central America.

10. **Caluromys derbianus** (Waterhouse).

Didelphys derbianus Waterh., Jard. Nat. Libr., Mamm., XI, 1841,
p. 97.

Caluromys derbianus Elliot, Mamm. Middle Amer. & W. Indies,
F. C. M. Pub., IV, Pt. 1, 1904, p. 9, fig. III. Zool. Ser.

EARL OF DERBY'S OPOSSUM.

Type locality. Unknown.

Geogr. Distr. Nicaragua in Central America to Peru in South
America.

Caluromys laniger pallidus (Thomas).

Philander laniger pallidus Thomas, Ann. Mag. Nat. Hist., 7th Ser.,
IV, 1899, p. 286.

Caluromys laniger pallidus Allen, Bull. Am. Mus. Nat. Hist. N. Y.,
1900, p. 189. Elliot, Mamm. Middle Amer. & W. Indies,
F. C. M. Pub., IV, Pt. 1, 1904, p. 10. Zool. Ser.

PALE WOOLLY OPOSSUM.

Type locality. Bogava, Chiriqui, Panama. Altitude, about 750
feet.

Geogr. Distr. Southeastern Central America.

Genus 4. **Metachirus.**

Metachirus Burm., Syst. Uebers. Thier. Bras., 1, 1854, p. 135.
Type *Didelphys nudicaudata* E. Geoffroy.

11. **Metachirus nudicaudatus** (E. Geoffroy).

Didelphys nudicaudata E. Geoff., Cat. Mus. Paris, 1803, p. 42.

Didelphis! myosurus Temm., Mon. Mamm., 1, 1827, p. 38.

Metachirus nudicaudatus Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1900, p. 197. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 12. Zoöl. Ser.

RAT-TAILED OPOSSUM.

Type locality. Cayenne, French Guiana.

Geogr. Distr. Costa Rica to Brazil.

12. *Metachirus fuscogriseus* Allen.

Metachirus fuscogriseus Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1900, p. 194. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 12, figs. 3, IV. Zoöl. Ser.

Didelphys quica True, Proc. U. S. Nat. Mus., VII, 1885, p. 587. (nec Temm.)

ALLEN'S OPOSSUM. *Tlacuazin* in Guatemala.

Type locality. Unknown.

Geogr. Distr. Southern Mexico to Costa Rica.

a. *Metachirus fuscogriseus pallidus* Allen.

Metachirus fuscogriseus pallidus Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1901, p. 215. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 13. Zoöl. Ser.

ORIZABA OPOSSUM.

Type locality. Orizaba, State of Vera Cruz, Mexico.

Geogr. Distr. States of Vera Cruz, Puebla, and Tabasco, south-eastern Mexico.

Genus 5. **Didelphys.**

Didelphis (*sic*) Linnæus, Syst. Nat., 1758, p. 54. Type *Didelphis! marsupialis* Linnæus.

13. *Didelphys yucatanensis* Allen.

Didelphis! yucatanensis Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1901, p. 178.

Didelphys yucatanensis Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 14. Zoöl. Ser.

YUCATAN OPOSSUM.

Type locality. Chichen Itza, Yucatan, Mexico.

Geogr. Distr. States of Campeche and Yucatan, Mexico.

a. *Didelphys yucatanensis cozumelæ* Merriam.

Didelphis! yucatanensis cozumelæ Merr., Proc. Biol. Soc. Wash., XIV, 1901, p. 101.

Didelphys yucatanensis cozumelæ Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 15. Zoöl. Ser.

ISLAND OF COZUMEL OPOSSUM.

Type locality. Cozumel Island, Yucatan.

Geogr. Distr. Known only from the type locality.

14. *Didelphys mesamericana* Oken.

Didelphis! mesamericana Oken, Lehrb. der Zool., II, 1816, p. 1152.

Didelphis! californica Bennett, Proc. Zool. Soc., 1833, p. 40.

Didelphys californica Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 3. Zool. Ser.

Didelphys mesamericana Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, p. 15, fig. 5, pl. II. Zool. Ser.

LINNEAN OPOSSUM.

Type locality. "Northwestern Mexico, adjacent to California."

Geogr. Distr. Oklahoma Territory, through Mexico generally into Guatemala.

a. *Didelphys mesamericana texensis* (Allen).

Didelphis! marsupialis texensis Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1901, p. 172.

Didelphys marsupialis texensis Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, Suppl., p. 519. Zool. Ser.

Didelphis! mesamericana texensis Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1902, p. 256.

Didelphys mesamericana texensis Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 16. Zool. Ser.

TEXAS OPOSSUM.

Type locality. Brownsville, Cameron County, Texas.

Geogr. Distr. Coast region of Texas from Neuces Bay southward, and the lower Rio Grande Valley as far at least as Del Rio, Val Verde County, sporadically northward to San Antonio; Matamoros, State of Tamaulipas, Mexico.

b. *Didelphys mesamericana tabascensis* (Allen).

Didelphis! marsupialis tabascensis Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1901, p. 173.

Didelphis! mesamericana tabascensis Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1902, p. 257.

Didelphys mesamericana tabascensis Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 16. Zool. Ser.

TABASCO OPOSSUM.

Type locality. Teapa, State of Tabasco, Mexico.

Geogr. Distr. State of Vera Cruz to that of Tabasco, and across State of Chiapas, Mexico, to northern Guatemala.

15. *Didelphys virginiana* Kerr.

Didelphis! virginiana Kerr, Anim. Kingd., 1792, p. 193.

Didelphys virginiana Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 1, fig. 1. Zoöl. Ser.

**Didelphis! virginiana pigra* Bangs, Proc. Bost. Soc. Nat. Hist., 1898, p. 172.

Didelphys virginiana pigra Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 1. Zoöl. Ser. *Id.* Pub. Field Colum. Mus., III, 1901, p. 32. Zoöl. Ser.

VIRGINIAN OPOSSUM.

Type locality. Virginia.

Geogr. Distr. Atlantic coast of United States from New York to southern Florida, and west to Mississippi, and Texas.

***Didelphys marsupialis insularis* Allen.**

Didelphis! marsupialis insularis Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1902, p. 259.

Didelphys marsupialis insularis Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 17. Zoöl. Ser.

ISLAND OPOSSUM.

Type locality. Caparo, Island of Trinidad.

Geogr. Distr. Islands of Dominica, Grenada & St. Vincent, West Indies. Trinidad.

***Didelphys marsupialis battyi* Thomas.**

Didelphis! marsupialis battyi Thomas, Novitat. Zool., IX, 1902, p. 137.

Didelphys marsupialis battyi Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 17. Zoöl. Ser.

*I see no reason whatever to alter the opinion expressed in Pub. F. C. M., III, 1901, p. 32, Zoöl. Ser., that *D. v. pigra* Bangs is not entitled to even a subspecific rank, but should be considered a synonym of *D. virginiana*. That the latter species would assume a black pelage in certain parts of its dispersion, just as *D. marsupialis* Linn. does, is to be expected, and color in this animal and its near allies cannot be deemed alone to possess specific value. Dr. Allen in his review of *Didelphys* (Bull. Am. Mus. Nat. Hist. N. Y., 1901, p. 165), can only say in favor of *pigra* that "it is not sharply or very strongly differentiated from *D. virginiana* (typica), but south Florida and Louisiana specimens are quite unlike the northern form through their much darker general coloration," and also slight differences in extent of facial markings, black on ears and tail, and white on toes and ears. It may be remarked that in a series of specimens the last differences are very slight indeed. No CHARACTERS are produced by which *pigra* can be distinguished from the species *D. virginiana*, worthy of consideration, save a "much darker general coloration" in some examples, which, if allowed, at once necessitates raising to a subspecific rank the black individuals of *D. marsupialis* Linn., found so frequently in certain parts of its dispersion, and thus far no one has had the temerity to do this. *D. v. pigra* apparently has not one qualification to entitle it to a subspecific rank, and if it has, no one has, as yet, been able to discover it.

BATTY'S OPOSSUM.

Type locality. Coiba Island, west coast of Panama.

Geogr. Distr. Known only from type locality.

Didelphys marsupialis etensis Allen.

Didelphis! karkinophaga! caucæ Bangs, Amer. Nat., xxxv, 1901, p. 633.

Didelphis! marsupialis etensis Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1902, p. 262.

Didelphys marsupialis etensis Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 18. Zool. Ser.

ETEN OPOSSUM.

Type locality. Eten, Piura, Peru. Altitude, 50 feet.

Geogr. Distr. Low coast belt of Ecuador and Peru, bordering the Gulf of Guayaquil and probably northward near the coast of Chiriqui, Panama; San Miguel Island, Bay of Panama.

16. *Didelphys richmondi* Allen.

Didelphis! richmondi Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1901, p. 175.

Didelphys aurita Alston, Biol. Centr. Amer., Mamm., 1, 1881, p. 197. (nec Max.)

Didelphys richmondi Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 16. Zool. Ser.

RICHMOND'S OPOSSUM:

Type locality. Greytown, Nicaragua.

Geogr. Distr. Nicaragua and Costa Rica, Central America.

Order II. **Edentata. Edentates.**

Family I. **Bradypodidae. Sloths.**

Genus 6. **Choloepus.**

Choloepus Illig., Prodr. Syst. Mamm. et Av., 1811, p. 108. Type
Bradypus didactylus Linnaeus.

17. **Choloepus hoffmanni** Peters.

Choloepus hoffmanni Peters, Monatsb. K. Preuss. Ak. Wiss. Berlin, 1858, p. 128. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 20, figs. 4, VI. Zoöl. Ser.

HOFFMANN'S SLOTH.

Type locality. Costa Rica.

Geogr. Distr. Costa Rica, Central America.

Genus 7. **Bradypus.**

Bradypus Linn., Syst. Nat., 1, 1758, p. 34. Type *Bradypus tri-*
dactylus Linnaeus.

Ignavus Frisch, Nat. Syst. vierfüß. Thiere, in Tabellen, Tab. Gen., 1775.

Arctopithecus Virey, Nouv. Dist. Hist. Nat., xxxi, 1819, p. 279.
Gray, Proc. Zoöl. Soc., 1871, p. 436.

18. **Bradypus castaneiceps** (Gray).

Arctopithecus castaneiceps Gray, Proc. Zoöl. Soc., 1871, p. 444.

Bradypus castaneiceps Alston, Biol. Centr. Amer., Mamm., 1, 1881, p. 184. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 22, fig. VII. Zoöl. Ser.

CHESTNUT-HEADED SLOTH. *Camaleon* in Nicaragua.

Type locality. Chontales, Nicaragua.

Geogr. Distr. Nicaragua, Central America.

19. **Bradypus infuscatus** Wagler.

Bradypus infuscatus Wagl., Isis, 1831, p. 611. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 23, fig. 5. Zoöl. Ser.

Arctopithecus griseus Gray, Proc. Zoöl. Soc., 1871, p. 446, pl. xxxvi.

DUSKY SLOTH.

Type locality. Western Brazil.

Geogr. Distr. Costa Rica, Central America, to Brazil and Bolivia.

Family II. **Myrmecophagidæ. Anteaters.**Genus 8. **Cyclopes. Two-toed Anteaters.**

Cyclopes Gray, Lond. Med. Repos., xv, 1821, p. 305. Type

Myrmecophaga didactyla Linnæus.

Didactyles F. Cuv., Dict. Scien. Nat., LIX, 1829, p. 501.

Myrmydon Wagl., Nat. Syst. Amph., 1830, p. 36.

Dionyx I. Geoff., Rés. Leçons Mamm. Mus. Paris, 1835, p. 54.

Myrmecolichnus Reich., K. Säch. Naturh. Mus. Dresden, Ein Leitfaden, 1836, p. 51.

Eurypterna Glog., Hand- u. Hilfsb. Naturg., I, 1841, pp. xxxi, 112.

Cyclothurus (sic) Less., Nouv. Tabl. Règn. Anim., Mamm., 1842, p. 152.

Didactyla Liais, Climats, Géol., Faun., Géog., Botanique Brésil, 1872, p. 356.

20. **Cyclopes dorsalis** (Gray).

Cyclothurus dorsalis Gray, Proc. Zoöl. Soc., 1865, p. 385, pl. 19.

Cyclopes dorsalis Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 25, figs. 6, VIII. Zoöl. Ser.

LITTLE OR TWO-TOED ANTEATER.

Type locality. Costa Rica.

Geogr. Distr. Guatemala through Central America to South America.

Genus 9. **Tamandua. Three-toed Anteaters.**

Tamandua Frisch, Nat. Syst. vierfüß. Thiere, in Tabellen, 5, Tab.

Gen., 1775. Less., Nouv. Tabl. Règn. Anim., Mamm., 1842, p. 152. Gray, Proc. Zoöl. Soc., 1865, p. 383. Type *Myrmecophaga tetradactyla* Linnæus.

Tamanduas F. Cuv., Dict. Scien. Nat., LIX, 1829, p. 501.

Uroleptes Wagl., Nat. Syst. Amphib., 1830, p. 36.

21. **Tamandua tetradactyla** (Linnæus).

Myrmecophaga tetradactyla Linn., Syst. Nat., I, 1758, p. 35.

Myrmecophaga myosura Pall., Miscell., 1766, p. 64.

Ursine Anteater Griff., Anim. Kingd., III, 1827, p. 304, pl.

Myrmecophaga crispus Rupp., Mus. Senck., III, 1845, p. 179.

Tamandua bivittata Gray, Proc. Zoöl. Soc., 1865, p. 384.

Myrmecophaga quadridactyla True, Proc. U. S. Nat. Mus., 1885, p. 588.

Tamandua tetradactyla Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 27, figs. 7, IX, X. Zool. Ser.

THREE-TOED ANTEATER. *Tejon, Oso Colmenero* in Costa Rica.

Type locality. "America Meridionali," Brazil?

Geogr. Distr. Mexico, through Central America to Peru and Paraguay, South America.

a. * *Tamandua tetradactyla tenuirostris* Allen.

Tamandua tetradactyla tenuirostris Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1904, p. 394.

SLENDER NOSE ANTEATER.

Type locality. Passa Nueva, State of Vera Cruz, Mexico.

Geogr. Distr. Range unknown.

Genl. Char. Size large; tail long as head and body.

Color. "Light areas white faintly tinged with yellowish, the yellow tint a little stronger on the nuchal-interscapular space; median light dorsal line extending to hips; dark area purplish black sharply defined against the white, and reaching the base of the tail above and below."

Measurements. Total length, 1185; tail, 600; hind foot, 95. Skull: occipito-nasal length, 133; width of braincase, 42; length of nasals, 55.

b. * *Tamandua tetradactyla chiriquensis* Allen,

Tamandua tetradactyla chiriquensis Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1904, p. 295.

CHIRIQUI ANTEATER.

Type locality. Boqueron, Chiriqui, Panama.

Geogr. Distr. Range unknown.

Genl. Char. Size large; tail as long as head and body.

Color. Dark areas deep brownish black; light areas dull brownish buffy white; deeper on nuchal-interscapular region. The black extending for nearly four inches on the basal portion of the tail. Light median dorsal stripe terminates considerably in front of hips.

Measurements. Total length, 1180; tail, 555; hind foot, 100; ear, 50. Skull: occipito-nasal length, 132.5; width of braincase, 42; length of nasals, 55.

22. *Tamandua sellata* (Cope).

Myrmecophaga sellata Cope, Amer. Nat., XXIII, 1889, p. 183.

*The specific standing of these two named forms depends upon the results of a comparison between them and specimens from Honduras, the type locality of *M. sellata* Cope, for it may be that one, perhaps both, will prove to be synonymous with that species.

Tamandua sellata Elliot, Mamm. Middle Amer. & W. Indies,
F. C. M. Pub., IV, Pt. I, 1904, p. 28. Zoöl. Ser.

SADDLE-BACKED ANTEATER.

Type locality. Honduras.

Geogr. Distr. Honduras to French Guiana? South America;
range unknown.

Genus 10. *Myrmecophaga*. Great Anteaters.

Myrmecophaga Linn., Syst. Nat., I, 1758, p. 35. Type *Myrmecophaga tridactyla* Linnæus.

Falcifer Rehn, Amer. Nat., XXXIV, 1900, p. 576.

23. *Myrmecophaga tridactyla* Linnæus.

Myrmecophaga tridactyla Linn., Syst. Nat., I, 1758, p. 35. Elliot,
Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt.
I, 1904, p. 29, figs. 8, XI. Zoöl. Ser.

Myrmecophaga jubata Linn., Syst. Nat., I, 1766, p. 52.

Falcifer jubata Rehn, Amer. Nat., XXXIV, 1900, p. 576.

GREAT ANTEATER.

Type locality. Brazil.

Geogr. Distr. Guatemala to Brazil.

Family III. *Dasypodidæ*. Armadillos.

Subfamily I. *Dasypodinaæ*.

Genus 11. *Cabassous*.

Cabassous McMurtrie, Cuv. Anim. Kingd., I, 1831, p. 164. Type
Dasypus uncinatus Linnæus.

Xenurus Wagl., Nat. Syst. Amph., 1830, p. 36. (nec Boie, Aves,
1826.)

Arizostus Glog., Hand- u. Hilfsb. Naturg., I, 1841, pp. XXXII, 114.

Tatona Gray, Proc. Zoöl. Soc., 1865, p. 378.

Lysiurus Ameghino, Rev. Argent. Hist. Nat., I, 1891, entr. 4a,
p. 254.

24. *Cabassous centralis* (Miller).

Tatona (Ziphila) centralis Miller, Proc. Biol. Soc. Wash., XIII,
1899, pp. 4, 7.

Xenurus cinereus hispidus True, Proc. U. S. Nat. Mus., 1896, p. 345.

Cabassous centralis Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 32, fig. XII, pl. III. Zoöl. Ser.

MILLER'S ARMADILLO.

Type locality. Chamelicon, Honduras.

Geogr. Distr. Honduras, Central America; range unknown.

Subfamily II. **Tatuinæ.**

Genus 12. **Tatu.**

Tatu Frisch, Nat. Syst. vierfüss. Thiere, in Tabellen, 5, Tab. Gen., 1775. Blumenb. Handb. Naturg., 1799, p. 73. Type *Dasy-
pus novemcinctus* Linnæus.

Tatusia F. Cuvier, in Less. Man. de Mamm., 1827, p. 309.

Cachicamus McMurtrie, G. Cuv. Anim. Kingd., 1, 1831, p. 163.

25. Tatu novemcinctum (Linnæus).

*Dasy-
pus novemcinctus* Linn., Syst. Nat., I, 1758, p. 51; I, 1766, p. 51.

*Dasy-
pus octo-cinctus* (Linn.), Schreb. Säugth., II, 1775, p. 222, tab. LXXIII, LXXVI.

*Dasy-
pus peba* Desm., Mamm., 1820, p. 368.

*Dasy-
pus longicaudus* Max., Breit. Naturg. Bras., II, 1825, p. 531.

*Dasy-
pus mexicanus fenestratus* Peters, Monatsb. K. Preuss. Akad. Wiss. Berl., 1864, p. 180.

Tatusia leptorhyncha Gray, Handl. Edent., 1873, p. 14, pl. 2, figs. 3, 4.

Cabassous novemcinctus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 4, pl. 1. Zoöl. Ser.

Tatu novemcinctum Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 33, figs. 9, XIII. Zoöl. Ser.

NINE-BANDED ARMADILLO.

Type locality. "America meridionali," Brazil?

Geogr. Distr. Texas, through Mexico and Central America to Paraguay, South America.

Order III. **Sirenia. Sirenians.**

Family I. **Hydrodamalidae.**

Genus 13. **Hydrodamalis. Sea Cows.**

Hydrodamalis Retzius, Kongl. Vetensk. Acad. Nya. Handl. Stockholm, xv, 1794, p. 292. Type *Hydrodamalis stelleri* Retzius = *Manatus gigas* Zimmermann.

Rytina Illig., Prodr. Syst. Mamm. et Av., 1811, p. 141.

Rhytina Glog., Hand- u. Hilfsb. Naturg., 1841, pp. xxxiv, 165.

26. **Hydrodamalis gigas** (Zimmermann).

Manatus gigas Zimm., Geogr. Gesch., II, 1780, p. 426, fig. 5.

Hydrodamalis gigas Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 5, pl. II. Zoöl. Ser.

STELLER'S SEA COW.

Type locality. Shores of Bering Island, Bering Sea.

Geogr. Distr. Bering and Copper Islands, Bering Sea.

Family II. **Trichechidae. Manatees.**

Genus 14. **Trichechus.**

Trichechus Linn., Syst. Nat., I, 1758, p. 34. Type *Trichechus manatus* Linnæus.

Manatus Brünn., Zool. Fund., 1772, pp. 34, 38, 39. Scopoli, Intr. Hist. Nat., 1777, p. 490. Storr, Prodr. Meth. Mamm., 1780, p. 41.

27. **Trichechus manatus** Linnæus.

Trichechus manatus Linn., Syst. Nat., I, 1758, p. 34; I, 1766, p. 49. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 37, fig. xiv, pls. iv, v. Zoöl. Ser.

Manatus latirostris Harl., Journ. Acad. Nat. Scien. Phil., 1824, p. 111. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 6, pl. III. Zoöl. Ser.

Manatus australis Tilesius, Jahrb. Naturg., I, 1802, p. 23.

Manatus americanus Desm., Dict. Hist. Nat., 1817, p. 262, pl. 96. (Part.)

Manatus fluviatilis Wagn., Schreb. Säugth., Suppl., 1846, pl. 379.

MANATEE. *Vacca de Agua* in Guatemala.

Type locality. East coast of Florida.

Geogr. Distr. Florida coast to Yucatan, Mexico.

Order IV. **Cetacea. Cetaceans.**

Family I. **Balænidæ. Baleen Whales.**

Subfamily I. **Balæninæ.**

Genus 15. **Balæna.**

Balæna Linn., Syst. Nat., 1, 1758, p. 75; 1, 1766, p. 105. Type
Balæna mysticetus Linnæus.

Leiobalæna Eschr., K. Danske Vidensk. Selsk. Skrifter, Naturv.
Math. Afd. Kjöbenh., 5te Række, 1, 1849, p. 108; Unters. Nord.
Wallth., 1849, p. 7.

Eubalæna Gray, Proc. Zoöl. Soc., 1864, p. 201.

Hunterus Gray, Ann. Mag. Nat. Hist., 3d Ser., xiv, 1864, p. 349.

Hunterius Gray, Cat. Seals and Whales, 1866, p. 98.

Macleayius Gray, Proc. Zoöl. Soc., 1864, p. 589, figs. 1, 2. *Id.* Cat.
Seals and Whales, 1866, p. 78.

28. **Balæna glacialis** Bonnaterre.

Balæna glacialis Bonn., Tab. Encycl. Méthod. des trois Règnes de
la Nature, Cétol., 1789, p. 3. Elliot, Syn. N. Am. Mamm.,
F. C. M. Pub., 11, 1901, p. 8, fig. 2. *Id.* Mamm. Middle
Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 39,
fig. 10. Zoöl. Ser.

Balæna biscayensis Eschr., Rev. Mag. Zool., 2me Sér., 1860, p. 229.

Eubalæna australis Gray, Proc. Zoöl. Soc., 1864, p. 202.

Balæna cisarctica Cope, Proc. Acad. Nat. Scien. Phil., 1865, pp.
168-169.

Balæna britannicus Gray, Ann. Mag. Nat. Hist., 4th Ser., 1870,
p. 200.

Balæna australis var. *biscayensis* Lydekk., Brit. Mamm., 1895,
p. 260.

ICE BALEEN WHALE.

Type locality. Coast of Maryland.

Geogr. Distr. North Atlantic Ocean, south to Maryland on
American coast.

29. **Balæna sieboldii** Gray.

Balæna sieboldii Gray, Ann. Mag. Nat. Hist., 3d Ser., 1864, p. 349.

Balæna antarctica Schleg., Abh. Geb., 1841, p. 37. (Part.)

Balæna australis Temm., Faun. Japon., Mamm., 1842, pls. 28, 29.
(nec Desmoul.)

Balæna aleoutiensis Van Ben., Bull. Acad. Belg., xxv, 1868, p. 16.

Balæna sieboldii Elliot, Syn. N. Amer. Mamm., F. C. M. Pub., 11, 1901, p. 8. Zoöl. Ser.

SIEBOLD'S BALEEN WHALE.

Type locality. Coast of Japan.

Geogr. Distr. North Pacific Ocean, Bering Sea, coast of North America from California to the Aleutian Islands, and westward to longitude 150°.

30. *Balæna mysticetus* Linnæus.

Balæna mysticetus Linn., Faun. Seuc., 11, 1761, p. 16. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 8. Zoöl. Ser.

Balæna groenlandica Linn., Mus. Ad. Frid., 1, p. 51.

Balæna mysticetus pittekajensis Malmg., Bikang. Svensk. Ak. Handl., VIII, 1883, fig.

BOWHEAD. ARCTIC RIGHT WHALE.

Type locality. Greenland Seas.

Geogr. Distr. Arctic Ocean, Bering Sea, south to north latitude 64° on east coast of Greenland, and also to coast of Labrador; in Bering Sea to north latitude 55°, and in Sea of Okhotsk to latitude 54°.

31. *Balæna kuzira* (Gray).

Megaptera kuzira Gray, Cat. Seals and Whales, 1866, p. 130.

Balæna allamak Pall., Zoog. Rosso-Asiat., 1, 1831, p. 288.

Balæna antarctica Temm., Faun. Japon., 1842, p. 21, pl. 30. (nec F. Cuv.)

Balæna australis Schrenck, Reis. Amur-Land, 1, 1858, p. 193. (nec Gray.)

Balæna kuzira Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 9. Zoöl. Ser.

GRAY'S JAPANESE BALEEN WHALE.

Type locality. Coast of Japan.

Geogr. Distr. Japan Seas, North Pacific, Bering Sea.

Genus 16. **Rhachianectes.**

Rhachianectes Cope, Proc. Acad. Nat. Scien. Phil., 1869, p. 15.

Type Agaphelus glaucus Cope.

32. **Rhachianectes glaucus** (Cope).

Agaphelus glaucus Cope, Proc. Acad. Nat. Scien. Phil., 1868, p. 159; 1869, p. 15.

Rhachianectes glaucus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 9, and Suppl., p. 478. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 40, pls. VI, VII, VIII. Zoöl. Ser.

GRAY'S BALEEN WHALE.

Type locality. Coast of California.

Geogr. Distr. Coast of Lower California, Mexico, to the North Pacific and Arctic Oceans; Okhotsk and Bering Seas.

Genus 17. **Megaptera.**

Megaptera Gray, Erebus & Terror, Zoöl., 1846, p. 16. Type *Balæna longimana* Rudolphi = *Balæna nodosa* Bonnaterre.

Megapteron Wagn., Archiv f. Naturg., 1847, Bd. II, p. 38.

Kyphobalæna Eschr., K. Danske Vidensk. Selsk. Skrifter, Naturv. & Math. Afd. Kjöbenh., 5te Række, I, 1849, p. 108. *Id.* Unter. Nord. Wallth., 1849, p. 56.

Megapteropsis Van Ben., Bull. Acad. Roy. Sci. de Belgique, 2e Sér, xxxiv, 1872, p. 15.

Pæscopia Gray, Proc. Zoöl. Soc., 1864, p. 207, fig. 3. *Id.* Cat. Seals and Whales Brit. Mus., 1866, pp. 113, 125, fig. 19.

33. **Megaptera nodosa** (Bonnat).

Balæna nodosa Bonn., Tabl. Encyclop. and Méth. des trois Règnes de la Nature, Cétol., 1789, p. 5.

Balæna longimana Rudolphi, Abh. Ak. Berl., 1829, pls. 1-5.

Balæna boops (nec Linn.) (*keporkak*) Eschr., K. Dansk. Vid. Selskabs. Ath., XL, 1845, p. 239, t. 1, 3, 4.

Megaptera americana Gray, Erebus & Terror, Zoöl. 1846, p. 17.

Megaptera nodosa Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 10. Zoöl. Ser.

ARCTIC BALEEN WHALE.

Type locality. Coast of New England States.

Geogr. Distr. Arctic and North Atlantic Oceans.

a. *Megaptera nodosa osphyia* Cope.

Megaptera nodosa osphyia Cope, Proc. Acad. Nat. Scien. Phil., 1865, pp. 178-180. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 10.

COPE'S BALEEN WHALE.

Type locality. Coast of Maine forty miles from Petit Menan lighthouse.

Geogr. Distr. Atlantic Ocean.

b. *Megaptera nodosa bellicosa* Cope.

Megaptera nodosa bellicosa Cope, Proc. Am. Phil. Soc., XII, 1870,
p. 103. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M.
Pub., IV, Pt. 1, 1904, p. 41, pls. IX, X, XI. Zoöl. Ser.

FIGHTING BALEEN WHALE.

Type locality. Vicinity of St. Bartholomew's Island, West Indies.
Geogr. Distr. Caribbean Sea, Gulf of Mexico, South Atlantic.

34. *Megaptera versabilis* Cope.

Megaptera versabilis Cope, Proc. Acad. Nat. Scien. Phil., 1869,
p. 15. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901,
p. 10. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub.,
IV, Pt. 1, 1904, p. 41. Zoöl. Ser.

MOBILE BALEEN WHALE.

Type locality. Northwestern coast of North America.
Geogr. Distr. North Pacific Ocean to Alaska.

Subfamily II. *Balænopterinae*. Finback Whales.

Genus 18. *Balænoptera*.

Balænoptera Lacép., Hist. Nat. Cét., 1804, pp. xxxvi-xxxvii,
114-141, pls. IV figs. 1, 2, V fig. 1, VI-VIII. Type *Balænoptera*
gibbar Desmoulins.

Physalus Lacép., Hist. Nat. Cét., Ordres, Genres et Espèces, 1804,
pp. XL, 219-226.

Cetoptera Rafin., Analyse Nat., Adden., 1815, p. 219.

Benedenia Gray, Proc. Zoöl. Soc., 1864, p. 211, figs. 16, 17.

Sibbaldus Gray, Proc. Zoöl. Soc., 1864, p. 222.

Sibbaldius Flower, Proc. Zoöl. Soc., 1864, p. 391.

Cuvierius Gray, Cat. Seals and Whales, 1866, pp. 114, 164; 1871,
p. 54.

**Agaphelus* Cope, Proc. Acad. Nat. Scien. Phil., 1868, p. 223.

35. *Balænoptera acuto-rostrata* Lacépède.

Balænoptera acuto-rostrata Lacép., Hist. Nat. Cét., XII, 1, 1803-4,
p. 197, pl. 8. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II,
1901, p. 11, fig. 3. Zoöl. Ser.

Rorqualus minor Knox, Jour. Inst., 1834, p. 336.

Agaphelus gibbosus Cope, Proc. Acad. Nat. Scien. Phil., 1869,
p. 15. (nec Erxl.) Elliot, Syn. N. Am. Mamm., F. C. M.
Pub., II, 1901, p. 11. Zoöl. Ser.

*See True, Smith. Contr. Knowledge, xxxiii, 1904, p. 105.

Balænoptera Mondini Capell., Mem. Acad. Bologn., 1877, VII, pl. 1-4.
SHARP-NOSED FINBACK WHALE.

Type locality. —?

Geogr. Distr. North Atlantic, Davis Straits.

36. *Balænoptera davidsoni* Scammon.

Balænoptera davidsoni Scamm., Proc. Calif. Acad. Scien., IV, 1872, p. 269. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 12. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 42, pls. XII, XIII, XIV. Zoöl. Ser.

DAVIDSON'S FINBACK WHALE.

Type locality. Admiralty Inlet, coast of Washington, North Pacific Ocean.

Geogr. Distr. West coast of North America, Mexico to Bering Sea.

37. *Balænoptera velifera* Cope.

Balænoptera velifera Cope, Proc. Acad. Nat. Scien. Phil., 1869, p. 16. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 12. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 43. Zoöl. Ser.

PACIFIC FINBACK WHALE.

Type locality. Shores of Oregon and California.

Geogr. Distr. North Pacific Ocean; coasts of California and Oregon; Commander Islands.

a. Balænoptera velifera copei Elliot.

Balænoptera borealis Dall, in Scammon Mar. Mamm. N. W. Amer., 1874, p. 303, fig. p. 37. (nec Lesson.)

Balænoptera velifera copei Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 13. Zoöl. Ser.

COPE'S FINBACK WHALE.

Type locality. Shumagin Islands, Alaska.

Geogr. Distr. Northern Seas and about the Aleutian Islands.

38. *Balænoptera physalus* (Linnæus).

Balæna physalus Linn., Syst. Nat., I, 1766, p. 106.

Balæna boops Linn., Syst. Nat., I, 1766, p. 106.

Balænoptera roqual Lacép., Hist. Nat. Cét., 1803-4, pp. 37, 126, t. I, fig. 3.

Physalus duguidii Heddle, Proc. Zoöl. Soc., 1856, p. 187. Gray, Proc. Zoöl. Soc., 1864, p. 221.

Pterobalæna communis Van Ben., Bull. Acad. Brux., 1re Ser., I, 1857, p. 403.

Benedenia knoxii Gray, Proc. Zoöl. Soc., 1864, p. 212.

Balænoptera physalus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 13. Zoöl. Ser.

NORTHERN FINBACK WHALE.

Type locality. Unknown. Atlantic coast of Europe.

Geogr. Distr. Arctic and Atlantic Oceans; Mediterranean Sea. On North American coast from Greenland to New Jersey.

39. *Balænoptera borealis* (Lesson).

Balæna borealis Less., Hist. Nat. Cét., 1828, p. 342.

Balænoptera borealis Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 13. Zoöl. Ser.

LESSON'S FINBACK WHALE.

Type locality. — ?

Geogr. Distr. Arctic and North Atlantic Oceans; Mediterranean Sea.

40. *Balænoptera musculus* (Linnæus).

Balæna musculus Linn., Syst. Nat., I, 1766, p. 106.

Balæna physalus Pall., Zoogr. Rosso-Asiat., 1830, p. 289.

Physalus (Rorqualus) sibbaldii Gray, Proc. Zoöl. Soc., 1847, p. 92.

Sibbaldius tectirostris Cope, Proc. Acad. Nat. Scien. Phil., 1869, p. 17.

Balænoptera musculus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 13. Zoöl. Ser.

BLUE FINBACK WHALE.

Type locality. Coast of Yorkshire, England.

Geogr. Distr. Arctic and North Atlantic Oceans.

41. *Balænoptera sulfurea* (Cope).

Sibbaldius sulfureus Cope, Proc. Acad. Nat. Scien. Phil., 1869, p. 20.

Balænoptera sulfurea Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 14. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W.

Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 43. Zoöl. Ser.

YELLOW-BELLIED FINBACK WHALE.

Type locality. Northwest coast of North America.

Geogr. Distr. North Pacific Ocean to Bering Sea.

Family II. **Physeteridæ. Sperm Whales.**

Subfamily I. **Physeterinæ.**

Genus 19. **Physeter.**

Physeter Linn., Syst. Nat., I, 1758, p. 76; I, 1766, p. 107. *Type* *Physeter macrocephalus* Linnæus.

Catodon Linn., Fauna Suecica, 2d ed., 1761, p. 18.

Tursio Fleming, Phil. Zoöl., 11, 1822, p. 211. (nec Wagler, nec Gray.)

42. *Physeter macrocephalus* Linnæus.

Physeter macrocephalus Linn., Syst. Nat., 1, 1758, p. 76; 1, 1766, p. 107, fig. 4. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 15. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., 14, Pt. 1, 1904, p. 45, figs. 11, xvi. Zoöl. Ser.

Catodon macrocephalus Gray, Cat. Seals and Whales, 1866, p. 202.

CACHALOT SPERM WHALE.

Type locality. North Atlantic.

Geogr. Distr. All seas.

Genus 20. *Hyperoodon*.

Hyperoodon Lacép., Hist. Nat. Cét., 1804, p. XLIV. Type *Delphinus butskopf* Bonn. = *Balæna rostrata* Müller.

Ancylodon Illig., Prodr. Syst. Mamm. et Av., 1811, p. 142.

Uranodon Illig., Prodr. Syst. Mamm. et Av., 1811, p. 143.

Cetodiodon Jacobs, Dublin Phil. Jour. and Scient. Rev., 1825, t.

Chænodelphinus Eschr., Forhandl. Skandinav. Naturforsk., 3die möte, Stockholm, 1843, pp. 651-655.

Chænocetus Eschr., Oversigt K. Danske Vidensk. Selsk. Forhandl. Kjöbenh., 1846. *Id.* K. Danske Vidensk. Selsk. Skrifter, Naturv. and Math. Afd. Kjöbenh. 5te Række, 1, 1849, p. 97.

Lagenocetus Gray, Proc. Zoöl. Soc., 1863, p. 200.

43. *Hyperoodon rostratus* (Müller).

Balæna rostrata Müll., Zool. Dan. Prodr., 1776, p. 7.

Delphinus bidentatus Bonn., Tab. Encycl. and Méth. des trois Règnes de la Nature, Cétol., 1789, p. 25.

Delphinus butskopf Bonn., Tab. Encycl. and Méth. des trois Règnes de la Nature, Cétol., 1789, p. 25.

Delphinus anarnacus Desm., Mamm., 1820, p. 550, No. 780.

Delphinus hunteri Desm., Mamm., 1820, p. 520, No. 782.

Hyperoodon latifrons Gray, Erebus & Terror, Zoöl., 1846, p. 27, pl. 4.

Lagenocetus latifrons Gray, Proc. Zoöl. Soc., 1864, p. 241.

Delphinus hyperoodon Schleg., Zoogd. Nederl., 1870, p. 94, pl. 18.

Hyperoodon rostratus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 16, pl. iv. Zoöl. Ser.

BEAKED SPERM WHALE.

Type locality. — ?

Geogr. Distr. North Atlantic Ocean, Davis Straits, Hudson Straits.

Genus 21. **Cogia.**

Kogia ! Gray, Voy. Erebus and Terror, Zoöl., Mamm., I, 1846, p. 22.

Type *Physeter breviceps* Blainville.

44. **Cogia breviceps** (Blainville).

Physeter breviceps Blainv., Ann. Anat. Phys., II, 1838, p. 337.

Kogia floweri Gill, Amer. Nat., IV, 1871, p. 738, fig. 172.

Kogia breviceps Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, Suppl., p. 479, pls. L, LI, LII. Zoöl. Ser.

Cogia breviceps Elliot, Mamm. Middle Amer. & W. Indies, F.

C. M. Pub., IV, Pt. I, 1904, p. 46, pls. XV, XVI, XVII. Zoöl. Ser.

PIGMY SPERM WHALE.

Type locality. Cape of Good Hope.

Geogr. Distr. Indian and Pacific Oceans; coast of southern California, possibly of Mexico and Atlantic coasts of North America.

Genus 22. **Ziphius. Two-toothed Whales.**

Ziphius G. Cuv., Rech. Oss. Foss., 2d ed., v, Pt. I, 1823, p. 352, pl. xxvii, figs. 3, 4, 7, 9. Type *Ziphius cavirostris* G. Cuvier.

Aliama Gray, Proc. Zoöl. Soc., 1864, p. 242.

Petrorhynchus Gray, Proc. Zoöl. Soc., 1865, p. 524.

Ziphiorrhynchus Burm., Revista Farmac., Bull. Acad. Belg., 1865.

Id. Ann. Mag. Nat. Hist., 3d Ser., xvii, 1866, p. 94.

Mesoodon Brandt, Mém. Acad. Imp. Scien. St. Pétersb., 7th Ser., xx, 1873, p. 220.

45. **Ziphius cavirostris** G. Cuvier.

Ziphius cavirostris G. Cuv., Rech. Oss. Foss., 2d ed., v, 1823, p. 353. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 17. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 47, pls. xviii, xix, xx. Zoöl. Ser.

TWO-TOOTHED SPERM WHALE.

Type locality. Coast of France, Mediterranean Sea.

Geogr. Distr. All seas.

46. Ziphius grebnitzkii Stejneger.

Ziphius grebnitzkii Stejneger, Proc. U. S. Nat. Mus., 1883, p. 77.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, Suppl.,
p. 480. Zoöl. Ser.

GREBNITZKY'S TWO-TOOTHED SPERM WHALE.

Type locality. Bering Island, Bering Sea.

Geogr. Distr. Bering and adjacent northern seas.

Genus 23. **Mesoplodon.**

Mesoplodon Gerv., Ann. Sciën. Nat. Paris, 1850, 3d Ser., xiv, p. 16.

Zoöl. *Type Delphinus sowerbiensis* Blainville = *Physeter bidens*
Sowerby.

47. Mesoplodon bidens (Sowerby).

Physeter bidens Sowerby, Brit. Miscell., 1, 1804, p. 1.

Delphinus sowerbyi Blainv., Nouv. Dict. Hist. Nat., ix, 1817,
p. 197.

Delphinus sowerbiensis Blainv., Nouv. Dict. Hist. Nat., ix, 1817,
p. 197.

Delphinus epidon Desm., Mamm., 1821, p. 521, No. 786.

Dauphindo dale Blainv., Bull. Soc. Philom., 1825, p. 139, desc.

Delphinus micropterus Cuv., Règn. Anim., 1829, p. 288.

Mesoplodon bidens Elliot, Syn. N. Am. Mamm., F. C. M. Pub.,
11, 1901, p. 18. Zoöl. Ser.

SOWERBY'S SPERM WHALE.

Type locality. Coast of Elginshire, England.

Geogr. Distr. North Atlantic Ocean.

48. Mesoplodon stejnegeri True.

Mesoplodon stejnegeri True, Proc. U. S. Nat. Mus., 1885, p. 384,
pl. 25.

Balænoptera stejnegeri Elliot, Syn. N. Am. Mamm., F. C. M.
Pub., 11, 1901, p. 14. Zoöl. Ser.

STEJNEGER'S SPERM WHALE.

Type locality. Coast of the Commander Islands.

Geogr. Distr. Bering Sea.

Genus 24. **Berardius.**

Berardius Duvernoy, Ann. Scien. Nat. Paris, 3me Sér., Zool., xv,
1851, p. 41. *Type Berardius arnuxii* Duvernoy.

49. Berardius bairdi Stejneger.

Berardius bairdi Stejneger, Proc. U. S. Nat. Mus., 1883, p. 75.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, Suppl.,
p. 481, pls. LIII, LIV, LV. Zoöl. Ser.

BAIRD'S SPERM WHALE.

Type locality. Stare Gavan, Bering Island.

Geogr. Distr. Northern Pacific Ocean; Bering Sea.

Family III. **Delphinidae. Dolphins, Porpoises, etc.**Subfamily I. **Delphinapterinæ.**Genus 25. **Monodon.**

Monodon Linn., Syst. Nat., 1, 1766, p. 105. Type *Monodon monoceros* (*sic*) Linnæus.

Ceratodon Briss., Règn. Anim., 1, 2d ed., 1762, p. 231.

Diodon Storr., Prodr. Meth. Mamm., 1780, p. 42, Tab. C.

Narwalus Lacép., Hist. Nat. Cét., 1804, pp. xxxvii-xxxviii, 142-163, pl. ix, fig. 1.

Oryx Oken, Lehrb. Naturg., 3ter Theil, 2te Abth., 1816, p. 672.

Tachynices Brookes, Cat. Anat. and Zoöl. Mus. Jos. Brookes, Lond., 1828, p. 40.

50. Monodon monoceras Linnæus.

Monodon monoceros (*sic*) Linn., Syst. Nat., 1, 1766, p. 105.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 19,
fig. 6. Zoöl. Ser.

Narwalus vulgaris Lacép., Hist. Nat. Cét., 1804, p. 142.

Narwalus andersoni Lacép., Hist. Nat. Cét., 1804, p. 163, pl. 9,
fig. 1.

Monodon microcephalus Flem., Mem. Wern. Nat. Hist. Soc., 1,
1811, t.

Tachynices megacephalus Brookes, Cat. Anat. Zoöl. Mus., 1828,
p. 40.

NARWHAL.

Type locality. Greenland Sea.

Geogr. Distr. Arctic Sea.

Genus 26. **Delphinapterus.**

Delphinapterus Lacép., Hist. Nat. Cét., 1804, pp. xli, 243-249.

Type *Delphinapterus beluga* Lacépède = *Delphinus leucas* Pallas.

51. Delphinapterus leucas (Pallas).

Delphinus leucas Pallas, Reise in Russisch. Reich., III, 1776, p. 92, pl. 79.

Delphinapterus beluga Lacép., Hist. Nat. Cét., 1804, p. 243.

Cetus albicans Briss., Règn. Anim., 17, p. 359.

Delphinus canadensis Desm., Mamm., 2d Pt., 1822, p. 516.

Beluga rhinodon Cope, Proc. Acad. Nat. Scien. Phil., 1865, p. 278.

Beluga concreta Cope, Proc. Acad. Nat. Scien. Phil., 1865, p. 278.

Beluga declivis Cope, Proc. Acad. Nat. Scien. Phil., 1865, p. 278.

Beluga angustata Cope, Proc. Acad. Nat. Scien. Phil., 1869, p. 25.

Delphinapterus leucas Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 19, pl. v. Zoöl. Ser.

WHITE WHALE.

Type locality. Coast of Siberia.

Geogr. Distr. Arctic and sub-Arctic seas; south to Cape Cod, Massachusetts.

Genus 27. Phocæna. Porpoises.

Phocæna G. Cuv., Nouv. Dict. Hist. Nat., 2d ed., IX, 1817, p. 163.

Id. Règn. Anim., 1829, p. 289. *Type Delphinus phocæna* Linnæus.

52. Phocæna phocæna (Linnæus).

Delphinus phocæna Linn., Syst. Nat., I, 1758, p. 77.

Phocæna communis G. Cuv., Règn. Anim., 1817, p. 279.

Phocæna vomerina Gill, Proc. Acad. Nat. Scien. Phil., 1865, p. 178.

Phocæna brachycium Cope, Proc. Acad. Nat. Scien. Phil., 1865, p. 279; 1876, p. 134.

Phocæna lineata? Cope, Proc. Acad. Nat. Scien. Phil., 1876, pp. 134, 135.

Phocæna communis Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 20, pl. v.

Phocæna phocæna Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 48, fig. xvii, pl. xxi. Zoöl. Ser.

COMMON PORPOISE.

Type locality. Coast of Europe.

Geogr. Distr. North Atlantic and North Pacific Oceans.

53. Phocæna dalli True.

Phocæna dalli True, Proc. U. S. Nat. Mus., 1885, p. 95, pls. 2-6.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 21, fig. 7. Zoöl. Ser.

DALL'S PORPOISE. . .

Type locality. Strait west of Adakh Island, Aleutian Group.

Geogr. Distr. North Pacific Ocean, Alaska, Aleutian Group.

Genus 28. **Orcinus.** Killer Whales.

Orcinus Fitzin., Wiss-Populäre Naturg. Säugeth., VI, 1860, pp. 204, 217. Type *Delphinus orca* Linnæus.

54. **Orcinus orca** (Linnæus).

Delphinus orca Linn., Syst. Nat., I, 1758, p. 77.

Delphinus gladiator Bonnat., Tab. Encycl. & Méth. des trois Règnes de la Nature, Cétol., 1789, p. 23.

Orca schlegelii Lilljeb., Ray Soc., 1866, p. 235.

Orca arcticus Gerv. & Van Ben., Ost. Cét., 1868, p. 314.

Orca europæus Gerv. & Van Ben., Ost. Cét., 1868, p. 314.

Orca latirostris Gerv., Ost. Cét., 1868, p. 543.

Orca stenorhyncha Gray, Proc. Zool. Soc., 1870, p. 71.

Orcinus gladiator Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 22, pl. VI. Zool. Ser.

Orcinus orca Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 51, fig. XVIII, pl. XXII. Zool. Ser.

COMMON KILLER WHALE..

Type locality. Coast of Europe.

Geogr. Distr. All seas.

55. **Orcinus rectipinna** (Cope).

Orca rectipinna Cope Proc. Acad. Nat. Scien. Phil., 1869, p. 22, figs. 15, 16.

Orca ater! Cope, Proc. Acad. Nat. Scien. Phil., 1869, pp. 22, 57, fig. 17.

Orcinus rectipinna Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 22. Zool. Ser.

COPE'S KILLER WHALE.

Type locality. Pacific coast of the United States.

Geogr. Distr. Pacific Ocean. California to the Aleutian Islands.

Genus 29. **Pseudorca.**

Pseudorca Reinh., Overs. K. Dan. Vidensk. Selsk. Forh. Kjöbenh., 1862, p. 151. Type *Phocæna crassidens* Owen.

56. Pseudorca crassidens (Owen).

Phocæna crassidens Owen, Brit. Foss. Mamm., 1846, p. 516.

Orca (*Pseudorca*?) *meridionalis* Flower, Proc. Zool. Soc., 1864, p. 420.

Orca destructor Cope, Proc. Acad. Nat. Scien. Phil., 1866, p. 293.

Globiocephalus grayi Burm., Ann. Mus. Pub. Buen. Aires, I, 1864-69, p. 367, pl. XXI.

Pseudorca crassidens Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 23, pl. VI. Zool. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 51, pl. XXII. Zool. Ser.

LARGE-TOOTHED KILLER WHALE.

Type locality. Coast of Lincolnshire, England.

Geogr. Distr. All seas.

Genus 30. Globicephalus. Blackfish.

Globicephala Less., Hist. Nat. Mamm. Ois., I, 1828, 441. *Id.* Nouv. Tabl. Règn. Anim., Mamm., 1842, p. 200. *Type Delphinus deductor* Scoresby = *Delphinus melas* Traill.

Sphærocephalus Gray, Cat. Seals & Whales Brit. Mus., 1866, p. 323.

57. Globicephalus melas (Traill).

Delphinus melas Traill, Nichols Journ., 1809, XXIII, pl. 3.

Delphinus globiceps Cuv., Ann. Mus. Paris, 1812, p. 14. pl. I, figs. 1, 2.

Delphinus deductor Scoresby, Arct. Reg., I, 1820, t. 13, fig. 1.

Delphinus intermedius Harl., Jour. Acad. Nat. Scien. Phil., 1829, p. 51, pl. I, fig. 13.

Globiocephalus! incrassatus Gray, Proc. Zool. Soc., 1861, p. 309, fig. 1.

Sphærocephalus incrassatus Gray, Cat. Seals & Whales Brit. Mus., 1866, p. 323.

Globiocephalus! macrorhynchus Hector, Trans. N. Zeal. Inst., VII, 1874, pl. 16, figs. 3, 3a. (nec Gray.)

Globiocephalus! melas Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 23, pl. VII. Zool. Ser.

Globicephalus melas Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 52, fig. XIX, pl. XXIII. Zool. Ser.

PILOT OR CA'ING WHALE.

Type locality. Coast of England.

Geogr. Distr. South and North Atlantic Oceans; Gulf of Mexico.

58. *Globicephalus brachypterus* Cope.

Globiocephalus! brachypterus Cope, Proc. Acad. Nat. Scien. Phil., 1876, p. 129, fig. 131. Elliot; Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 24. Zool. Ser.

Globicephalus brachypterus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., 14, Pt. 1, 1904, p. 53. Zool. Ser.

SHORT-FINNED BLACKFISH.

Type locality. East coast of Delaware Bay, at the mouth of Meurice River.

Geogr. Distr. Gulf of Mexico, north to New Jersey, Atlantic Ocean.

59. *Globicephalus scammoni* Cope.

Globiocephalus! scammoni Cope, Proc. Acad. Nat. Scien. Phil., 1869, p. 21. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 24. Zool. Ser.

Globicephalus scammoni Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., 14, Pt. 1, 1904, p. 54. Zool. Ser.

SCAMMON'S BLACKFISH.

Type locality. Coast of Lower California.

Geogr. Distr. Coast of California southward; coast of South America.

Genus 31. *Grampus*.

***Grampus* Gray, Spicil. Zool., 1, 1828, p. 2. Type *Delphinus griseus* Cuvier.**

60. *Grampus griseus* (Cuvier).

Delphinus griseus Cuv., Ann. Mus. Paris, 1812, p. 14, pl. 1, fig. 1.

Delphinus rissoanus Desm., Mamm., 1822, p. 519.

Delphinus intermedius Harl., Journ. Acad. Nat. Scien. Phil., 1829, p. 51, t. 1.

Grampus cuvieri Gray, Ann. Mag. Nat. Hist., 1st Ser., 1846, p. 85.

Grampus stearnsii Dall, Proc. Calif. Acad. Scien., v, 1873, p. 13.

Grampus griseus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 25, pl. VII. Zool. Ser.

GRAY BLACKFISH.

Type locality. Brest, coast of France.

Geogr. Distr. North Atlantic and North Pacific Oceans. Coasts of United States, Cape Cod to Atlantic City, New Jersey; California.

Genus 32. **Lagenorhynchus. Bottle-nosed Whales.**

Lagenorhynchus Gray, Voy. Erebus & Terror, Zool., 1846, p. 34, pl. 6, fig. 3-5. Type *Delphinus leucopleurus* Rasch = *Delphinus acutus* Gray.

Leucopleurus Gray, Proc. Zool. Soc., 1866, p. 216.

61. Lagenorhynchus acutus (Gray).

Delphinus acutus Gray, Spicil. Zool., 1828, p. 2.

Delphinus eschrichti Schleg., Abh. Geb. Zool., 1841, p. 23.

Delphinus leucopleurus Rasch, Nyt. Mag. für Naturv., IV, 1843, p. 97, pls. 2, 3.

Leucopleurus arcticus Gray, Syn. Cét., 1868, p. 7.

Lagenorhynchus perspicillatus Cope, Proc. Acad. Nat. Scien. Phil., 1876, p. 136, pl. 4.

Lagenorhynchus gubernator Cope, Proc. Acad. Nat. Scien. Phil., 1876, p. 138.

Lagenorhynchus acutus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 25, pl. VIII. Zool. Ser.

SHARP-BEAKED BOTTLE-NOSED WHALE.

Type locality. Farøe Islands, North Sea.

Geogr. Distr. North Atlantic Ocean; northern coasts of United States; Cape Cod, Greenland, Farøe Islands, North Sea.

62. Lagenorhynchus thicola Gray.

Lagenorhynchus thicola Gray, Proc. Zool. Soc., 1849, p. 2. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 26. Zool. Ser.

GRAY'S BOTTLE-NOSED WHALE.

Type locality. West coast of North America.

Geogr. Distr. West coast of North America.

63. Lagenorhynchus albirostris (Gray).

Delphinus albirostris Gray, Ann. Mag. Nat. Hist., 1st Ser., 1846, p. 84.

?*Delphinus ibsenii* Eschricht, Undersøg. over Hvaldy., 5te Afh., p. 73.

Lagenorhynchus albirostris Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 26, fig. 8. Zool. Ser.

WHITE-BEAKED BOTTLE-NOSED WHALE.

Type locality. Coast of Norfolk, England.

Geogr. Distr. North Atlantic Ocean, Greenland, North and Baltic Seas, Farøe Islands, Davis Straits.

64. Lagenorhynchus obliquidens Gill.

Lagenorhynchus obliquidens Gill, Proc. Acad. Nat. Scien. Phil., 1865, p. 177. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 27. Zoöl. Ser.

Delphinus longidens Cope, Proc. Acad. Nat. Scien. Phil., 1866, p. 295.

Clymene similis Gray, Proc. Zoöl. Soc., 1868, p. 147.

GILL'S BOTTLE-NOSED WHALE.

Type locality. Coast of California.

Geogr. Distr. North Pacific Ocean, Monterey to Puget Sound, United States coast.

Genus 33. **Delphinus. Dolphins.**

Delphinus Linn., Syst. Nat., I, 1758, p. 77; I, 1766, p. 108. Type *Delphinus delphis* Linnaeus.

Rhinodelphis Wagn., Schreb. Säugth., VII, 1846, pp. 281, 316-349.

Eudelphinus Van Ben. & Gerv., Ostéog. Cét. Viv. et Foss., 1880, p. 600.

65. Delphinus delphis Linnaeus.

Delphinus delphis Linn., Syst. Nat., I, 1758, p. 77; I, 1766, p. 108. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 28, pl. ix. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 54, pl. xxiv. Zoöl. Ser.

Delphinus fulvi-fasciatus Wagn., Schreb. Säugth., 1846, pl. 361, fig. 1.

Delphinus novæ-zelandiæ Quoy & Gaim., Voy. Astrolabe, Mamm., 1830, p. 149.

Delphinus janira Gray, Voy. Erebus & Terror, Zoöl., 1846, p. 474, pl. 22, fig. 1.

Delphinus forsteri Gray, Cat. Seals & Whales Brit. Mus., 1866, p. 248.

Delphinus major Gray, Cat. Seals & Whales Brit. Mus., 1866, p. 396.

Delphinus moorii Gray, Cat. Seals & Whales Brit. Mus., 1866, p. 396.

Delphinus walkeri Gray, Cat. Seals & Whales Brit. Mus., 1866, p. 397.

Delphinus pomeegra Owen, Trans. Zoöl. Soc., VI, 1866, p. 23, pls. 6, 8.

Delphinus bairdi Dall, Proc. Calif. Acad. Scien., v, 1873, p. 12.

Delphinus microps Burm., Desc. Phys. Argent., III, 1879, p. 534.
(nec Gray.)

Delphinus fuscus Fisch., Act. Soc. Linn. Bord., v, 1881, p. 127,
pl. IV.

Delphinus sowerbianus Fisch., Act. Soc. Linn. Bord., v, 1881,
p. 127, pl. IV.

Delphinus variegatus Fisch., Act. Soc. Linn. Bord., v, 1881, p.
127, pl. VI.

Delphinus balteatus Fisch., Act. Soc. Linn. Bord., v, 1881, p. 127,
pl. —.

Delphinus moschatus Fisch., Act. Soc. Linn. Bord., v, 1881, p.
127, pl. V.

Clymene marginatus Fisch., Act. Soc. Linn. Bord., v, 1881, p. 150.
(nec Pucheran.)

Delphinus delphis var. *curvirostris* Riggio, Nat. Sicil., II, 1883, p.
157, pl. 3.

COMMON DOLPHIN.

Type locality. Coast of Europe.

Geogr. Distr. Atlantic Ocean; Gulf of Mexico.

Genus 34. **Tursiops. Dolphins.**

Tursiops Gerv., Hist. Nat. Mamm., II, 1855, p. 323. Type *Delphinus tursio* Fabricius.

Tursio Gray, List Spec. Mamm. Brit. Mus., 1843, pp. XXIII, 105.

Id. Cat. Seals & Whales, 1866, p. 254.

66. **Tursiops truncatus** (Montagu).

Delphinus truncatus Montagu, Mem. Wern. Soc., III, 1821, p. 73.

True, Proc. Acad. Nat. Scien. Phil., 1903, p. 313.

Phocæna compressicauda Less., Nat. Hist. Cét., 1828, p. 199.

Delphinus metis Gray, Voy. Erebus & Terror, Zoöl., 1846, p. 38,
pl. 17.

Delphinus cymodoce Gray, Voy. Erebus & Terror, Zoöl., 1846,
p. 38, pl. 19.

Delphinus eurynome Gray, Voy. Erebus & Terror, Zoöl., 1846,
p. 38, pl. 18.

Tursiops tursio Elliot, Syn. N. Am. Mamm., F. C. M. Pub.,
II, 1901, p. 29, pl. IX. Zoöl. Ser.

Tursiops truncatus Elliot, Mamm. Middle Amer. and W. Indies,
F. C. M. Pub., IV, Pt. 1, 1904, p. 56, fig. XX, pl. XXIV (mis-
printed on plate *Tursiops tursio*). Zoöl. Ser.

BOTTLE-NOSED PORPOISE OR DOLPHIN.

Type locality. Coast of Greenland.

Geogr. Distr. Atlantic Ocean, Mediterranean Sea, widely distributed; coast of Atlantic States, Maine to Florida; Gulf of Mexico.

67. *Tursiops gilli* Dall.

Tursiops gilli Dall, Proc. Calif. Acad. Scien., v, 1873, p. 13. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 29. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 56. Zoöl. Ser.

GILL'S DOLPHIN. COW FISH.

Type locality. North Pacific Ocean.

Geogr. Distr. Shores of southern and Lower California to northern part of Pacific Ocean.

Genus 35. *Lissodelphis*.

Lissodelphis Glog., Hand- u. Hilfsb. Naturg., 1841, pp. xxxiv, 169.

Type Delphinus peronii Lacépède.

Delphinapterus Less. & Garnot, Voy. Coquille, Zool., 1, 1re Pt., 1826, p. 179, pl. 9, fig. 1. Gray, Voy. Erebus & Terror, Zoöl., 1846, p. 35. (nec Lacépède.)

Tursio Wagl., Nouv. Syst. Amph., 1830, p. 34. (nec Fleming.)

68. *Lissodelphis borealis* (Peale).

Delphinapteris borealis Peale, U. S. Expl. Exped., Mamm., 1848, p. 35, pl. viii, fig. 2.

Lissodelphis borealis Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1904, p. 30, pl. x. Zoöl. Ser.

PACIFIC OCEAN DOLPHIN.

Type locality. Off coast of Oregon.

Geogr. Distr. North Pacific Ocean; San Diego, coast of California to Bering Sea.

Genus 36. *Prodelphinus*. Dolphins.

Prodelphinus Van Ben. & Gerv., Ostéog. Cét., Viv. et Foss., 1880, p. 604, pl. xxxviii. *Type Delphinus dubius?* Cuvier.

Clymene Gray, Proc. Zoöl. Soc., 1864, p. 237. (nec Munster, 1839.)

Clymenia Gray, Syn. Whales & Dolphins, 1868, p. 6. (nec Munster, 1839.)

69. Prodelphinus crotaphiscus (Cope).

Delphinus crotaphiscus Cope, Proc. Acad. Nat. Scien. Phil., 1865,
p. 203.

Prodelphinus crotaphiscus Elliot, Syn. N. Am. Mamm., F. C. M.
Pub., II, 1901, p. 31.

TEMPORAL DOLPHIN.

Type locality. Unknown. Skull in Museum at Salem, Massachusetts.

Geogr. Distr. Unknown.

70. Prodelphinus plagiodon (Cope).

Delphinus plagiodon Cope, Proc. Acad. Nat. Scien. Phil., 1866,
p. 296.

Prodelphinus plagiodon Elliot, Syn. N. Am. Mamm., F. C. M.
Pub., II, 1904, p. 31, fig. 9. Zool. Ser. *Id.* Mamm. Middle
Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 58,
fig. XXI. Zool. Ser.

SHARP-TOOTHED DOLPHIN.

Type locality. Eastern coast of the United States.

Geogr. Distr. Atlantic coast of the United States. Cape Hatteras
to the Gulf of Mexico.

71. Prodelphinus longirostris (Gray).

Delphinus longirostris Gray, Spicil. Zool., 1828, p. 1.

Delphinus microps Gray, Spicil. Zool., 1828, p. 1.

Delphinus alope Gray, Voy. Erebus & Terror, Zool., 1846, p. 42,
pl. 25.

Delphinus stenorhynchus Gray, Cat. Seals & Whales Brit. Mus.,
1866, p. 396.

Prodelphinus longirostris Elliot, Mamm. Middle Amer. & W.
Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 58. Zool. Ser.

LONG-NOSED DOLPHIN.

Type locality. Cape of Good Hope.

Geogr. Distr. Southern Oceans. Taken off the Tres Marias
Islands, Mexico.

72. Prodelphinus euphrosyne (Gray).

Delphinus euphrosyne Gray, Voy. Erebus & Terror, Zool., 1846,
p. 40, pl. 22. *Id.* Cat. Seals & Whales Brit. Mus., 1866, p.
251.

Delphinus styx Gray, Voy. Erebus & Terror, Zool., 1846, p. 39,
pl. 2.

Delphinus tethyos Gerv., Bull. Soc. Agr. Herault, 1853, XL, p. 150,
pl. 1.

Delphinus marginatus (Duvern.) Pucher., Rev. Zool., 1854, pl. 547.

Delphinus eurynome Blyth, Cat. Mus. Asiat. Soc. Beng., 1863, p. 90. (nec Gray.)

Tursio doreides Gray, Cat. Seals & Whales Brit. Mus., 1866, p. 400.

Clymenia euphrosinoides Gray, Syn. Whales & Dolphins, 1868, p. 6.

Clymenia novæ-zelandiæ Hector, Trans. N. Zeal. Inst., 1873, v, p. 159.

Prodelphinus euphrosyne Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 30, pl. x. Zool. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 58, pl. XXIII. Zool. Ser.

GRAY'S DOLPHIN.

Type locality. Unknown.

Geogr. Distr. Atlantic Ocean; Mediterranean Sea; coast of Jamaica.

73. *Prodelphinus frenatus* (Cuvier).

Delphinus frenatus Cuv., Mamm., 1825, pl. 426.

Delphinus frontalis (Duss.) Cuv., Règn. Anim., I, 1829, p. 288.

Delphinus doris Gray, Voy. Erebus & Terror, Zool., 1846, p. 39, pl. 20.

Delphinus clymene Gray, Cat. Seals & Whales Brit. Mus., 1866, p. 249.

Clymene normalis Gray, Proc. Zool. Soc., 1866, p. 214.

Prodelphinus frenatus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 32. Zool. Ser.

BRIDLED DOLPHIN.

Type locality. ———?

Geogr. Distr. Atlantic and Indian Oceans.

Order V. **Ungulata. Hoofed Quadrupeds.**

Family I. **Tagassuidæ. Peccaries.**

Genus 37. **Tagassu.**

Tagassu Frisch, Nat. Syst. vierfüß. Thiere, in Tabellen, 3 Tab.

Gen., 1775. Type *Sus tajacu* Linnæus.

Tayassu Fisch., Zoogn., III, 1814, p. 284.

Dicotyles G. Cuv., Règn. Anim., I, 1817, p. 327.

Notophorus Fisch., Mém. Soc. Imp. Nat. Moscou, v, 1817, pp. 373, 418.

Adenonotus Brookes, Prodr. Syst. Anim., II, 1828.

Subgenus A. **Tagassu.**

74. Tagassu nanus (Merriam).

Tayassu nanus Merr., Proc. Biol. Soc. Wash., XIV, 1901, p. 102.

Tagassu nanus Elliot, Mamm. Middle Amer. & W. Indies, F. C.

M. Pub., IV, Pt. I, 1904, p. 62. Zoöl. Ser.

DWARF PECCARY.

Type locality. Island of Cozumel, Yucatan.

Geogr. Distr. Only known from the Island of Cozumel, Yucatan.

75. Tagassu angulatum (Cope).

Dicotyles angulatus Cope, Amer. Nat., XXIII, 1889, p. 147.

Dicotyles tajacu Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 33, pl. XI. Zoöl. Ser. (nec Linn.)

Tagassu angulatum Elliot, Mamm. Middle Amer. & W. Indies, F.

C. M. Pub., IV, Pt. I, 1904, p. 63, pls. XXV, XXVI. Zoöl. Ser.

TEXAN PECCARY. *Moran, Jabali* in Mexico.

Type locality. Guadalupe River, Texas.

Geogr. Distr. Northeastern Mexico into Texas.

a. Tagassu angulatum humerale (Merriam).

Tayassu angulatus humeralis Merr., Proc. Biol. Soc. Wash., XIV, 1901, p. 122.

Tagassu angulatum humerale Elliot, Mamm. Middle Amer. & W.

Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 63. Zoöl. Ser.

ARMERIA PECCARY.

Type locality. Armeria, State of Colima, Mexico.

Geogr. Distr. State of Colima to Tehuantepec, Mexico.

b. *Tagassu angulatum yucatanense* (Merriam).

Tayassu angulatus yucatanensis Merr., Proc. Biol. Soc. Wash., XIV, 1901, p. 123.

Tagassu angulatum yucatanense Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 63. Zoöl. Ser.

YUCATAN PECCARY.

Type locality. Tunkas, Yucatan.

Geogr. Distr. Arid peninsula of Yucatan.

c. *Tagassu angulatum crassum* (Merriam).

Tayassu angulatus crassus Merr., Proc. Biol. Soc. Wash., XIV, 1901, p. 124.

Tagassu angulatum crassum Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 64. Zoöl. Ser.

HEAVY PECCARY.

Type locality. Metlatoyuca, State of Puebla, Mexico.

Geogr. Distr. States of Puebla and Chiapas, Mexico.

d. *Tagassu angulatum sonoriense* (Mearns).

Dicotyles angulatus sonoriensis Mearns, Proc. U. S. Nat. Mus., XX, 1897, p. 469.

Tagassu angulatum sonoriense Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 64. Zoöl. Ser.

SONORA PECCARY.

Type locality. San Bernardino River, State of Sonora, Mexico, near Monument No. 77, Mexican boundary line.

Geogr. Distr. State of Sonora, Mexico; Arizona.

76. *Tagassu crusnigrum* (Bangs).

Tayassu crusnigrum Bangs, Bull. Mus. Comp. Zoöl., XXXIX, 1902, p. 20.

Tagassu crusnigrum Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 65. Zoöl. Ser.

BOQUETE PECCARY.

Type locality. Boquete, Chiriqui, Panama, Central America. Altitude, 4,000 feet.

Geogr. Distr. Panama, Central America.

Subgenus B. *Olidosus*.

Dicotyles Gray (nec Cuv.), Proc. Zoöl. Soc., 1868, p. 45.

Olidosus Merr., Proc. Biol. Soc. Wash., XIV, 1901, p. 120. Type

Tayassu pecari Fischer.

77. *Tagassu pecari* (Fischer).

Tagassu pecari Fisch., Zoogn., III, 1814, p. 285.

Sus albirostris Illig., Abhandl. K. Preuss. Akad. Wiss. Berlin, 1815, p. 115.

Dicotyles labiatus Cuv., Règn. Anim., I, 1817, p. 238.

Tagassu pecari Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 66, fig. XXII, pls. XXVII, XXVIII. Zool. Ser.

WHITE-LIPPED PECCARY. Warree, Caribbanco, in Costa Rica.

Type locality. Paraguay.

Geogr. Distr. Guatemala, Central America, to Paraguay in South America.

a. *Tagassu pecari ringens* (Merriam).

Tagassu albirostris ringens Merr., Proc. Biol. Soc. Wash., XIV, 1901, p. 121.

Tagassu pecari ringens Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 66. Zool. Ser.

SAVAGE PECCARY.

Type locality. Apazote, near Yohaltem, State of Campeche, Mexico.

Geogr. Distr. State of Campeche. Range unknown.

Family II. *Cervidae*.**Subfamily I. *Cervinae*.****Genus 38. *Rangifer*. Caribou.**

Rangifer Frisch, Das Nat. Syst. vierfüß. Thiere, in Tabellen, 3 Tab.

Gen., 1775. Type *Rangifer lapponicus* Frisch = *Cervus tarandus* Linnæus.

Tarandus Billberg, Syn. Faun. Scand., Mamm., I, 1828, Consp. A, D, p. 22.

78. *Rangifer caribou* (Gmelin).

Cervus tarandus caribou Gmel., Syst. Nat., I, 1788, p. 177.

Cervus tarandus sylvestris Rich., Faun. Bor. Amer., Mamm., 1829, p. 250.

Cervus hastalis Agass., Sillim. Journ., 1847, p. 436.

Rangifer caribou Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 35, pl. XIII. Zool. Ser.

CARIBOU.

Type locality. Eastern Canada.

Geogr. Distr. Labrador, Lower Canada south to Nova Scotia, New Brunswick and Maine in the east; west through Quebec and Ontario to Montana, in forest lands.

79. *Rangifer terrænovæ* Bangs.

Rangifer terrænovæ Bangs, Private Pub., Boston, 1896. Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1896, p. 233. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 36. Zool. Ser.

NEWFOUNDLAND CARIBOU.

Type locality. Grand Lake, Newfoundland (Allen), Codroy, Newfoundland (Bangs).

Geogr. Distr. Newfoundland. Nova Scotia?

80. *Rangifer montanus* Seton-Thompson.

Rangifer montanus Seton-Thompson, Ottawa Nat., XIII, 1899, p. 129. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 36. Zool. Ser.

MOUNTAIN CARIBOU.

Type locality. Illecillewaet watershed, near Revelstoke, Selkirk Range, British Columbia.

Geogr. Distr. Mountains of British Columbia, north into south-eastern Alaska, east into the Rocky Mountains of Alberta, and south into Idaho.

81. **Rangifer osborni* Allen.

Rangifer osborni Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1902, p. 149.

OSBORN'S CARIBOU.

Type locality. Gold Range Mountains, near Sicamous, British Columbia.

Geogr. Distr. Gold Range and Cassiar Mountains, northern British Columbia.

Genl. Char. Size large; antlers large, heavy.

Color. Above brown, darkest on breast and limbs; neck brownish gray, nearly white in front; rump and ventral surface white; tail white, with a central dark brown stripe; white band above hoofs; ears dark brown.

Measurements. Total length, 2057-2210; tail vertebræ, 152.4; hind foot, 597-635; height at shoulder, 1270-1397; ear from crown,

*Specific distinctness not satisfactorily established. Increased knowledge and more specimens required.

140. Skull: basal length, 360-430; length of nasals, 122-142; zygomatic breadth, 136-157; mastoid breadth, 146-149; upper tooth row, 99-101; length of mandible, 310-330. Antlers: length of beam, 745-1235; spread at tips, 393-790.

82. **Rangifer stonii* Allen.

Rangifer stonei Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1901, p. 143. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, Suppl., p. 483. Zool. Ser. *Id.* Pub. Field Columb. Mus., 1901, p. 61, pls. XI, XII, XIII. Zool. Ser.

STONE'S CARIBOU.

Type locality. Kenai Peninsula, Alaska.

Geogr. Distr. Unknown.

83. **Rangifer granti* Allen.

Rangifer granti Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1902, p. 122.

GRANT'S CARIBOU.

Type locality. Alaskan Peninsula opposite Popoff Island.

Geogr. Distr. Barren Lands, Alaskan Peninsula; Unimak Island.

Extent of range unknown.

Genl. Char. Size small; antlers light and slender; color similar to that of other caribou.

Color. Adult male at end of October. Above dark brown; top of nose blackish brown; sides of head lighter brown; shoulders and sides of neck light gray (probably white in winter); rump patch, ventral surface, and inner side of thighs white; breast dark brown; tail white, with a dark band above continued from the back through the white rump patch; legs and feet as in other caribou. Female lighter, broccoli brown.

Measurements. Total length, 1651-1905; tail vertebræ, 140-203; tarsus and foot, 470-584; height at shoulders, 903-1245. Skull: basal length, 350-375; length of nasals, 118-126; zygomatic breadth, 133-141; mastoid breadth, 111-135; upper tooth row, 96-104; length of mandible, 266-300. Antlers: beam along curve to tip, 780-1010; greatest spread, 640-890.

84. *Rangifer groenlandicus* (Gmelin).

Cervus groenlandicus Gmel., Syst. Nat., 1, 1788, p. 177.

Rangifer groenlandicus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 37. Zool. Ser.

GREENLAND OR ARCTIC CARIBOU.

*Specific distinctness not satisfactorily established. Increased knowledge and more specimens required.

Type locality. Greenland.

Geogr. Distr. Restricted to Greenland, if distinct from the animal obtained on the mainland, which cannot as yet be said to be completely established.

85. Rangifer arcticus (Richardson).

Cervus tarandus var. *arctica* Rich., Faun. Bor. Amer., Mamm., 1829, p. 241.

Rangifer dawsoni Seton-Thompson, Ottawa Nat., XIII, 1900, p. 260. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 36. Zool. Ser. Osgood, N. Am. Faun., No. 21, 1901, p. 26.

Rangifer arcticus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 37.

BARREN GROUND CARIBOU.

Type locality. Eastern Canada; shores of Hudson Bay?

Geogr. Distr. Barren grounds of Arctic America north of the tree limit, to the shores and islands of the Arctic Ocean, Greenland excepted, if the specific distinctness of the mainland animal can be proven.

86. *Rangifer pearyi Allen.

Rangifer pearyi Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1902, p. 409.

PEARY'S CARIBOU.

Type locality. Ellesmere Land, N. Lat. 79°.

Geogr. Distr. Known only from the type locality.

Color. Pure white, except an oval grayish brown patch over posterior half of dorsal surface; legs and feet white; ears tinged with gray. Flat skin, in winter pelage.

Measurements. Total length of flat skin, 1660.

Genus 39. **Alce. Moose.**

Alce Frisch, Das Natur-Syst. vierfüss. Thiere, in Tabellen, 3 Tab.

Gen., 1775. H. Smith, Griff. Anim. Kingd., v, 1827, p. 303.

Type *Cervus alces* Linnæus.

Alces Jard., Nat. Libr., Mamm., III, 1835, p. 125. Type *Alces americanus* Jardine.

Alcelaphus Glog., Handb. u. Hilfsb. Naturg., 1841, p. 143. (nec de Blainville, 1816.)

Paralces Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1902, p. 160.

*Specific distinctness of this animal by no means established.

87. Alce alces (Linnæus).

Cervus alces Linn., Syst. Nat., 1, 1758, p. 66.

Alces americanus Jard., Nat. Libr., Mamm., 1835, p. 125. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 38, pl. xiv. Zoöl. Ser.

Alces machlis Ogilby, Proc. Zoöl. Soc., 1836, p. 135.

Alces palmatus Gray, List. Mamm. Brit. Mus., 1843, p. 182.

Cervus lobatus Agass., Proc. Bost. Soc. Nat. Hist., 1846, p. 188.

Alces muswa Rich., Zoöl. of Herald, Mamm., 1852, p. 66.

MOOSE.

Type locality. Eastern Canada.

Geogr. Distr. Labrador, Nova Scotia, New Brunswick, Maine, Quebec, Ontario, Minnesota, Montana, and north to latitude 69° east of the Rocky Mountains.

88. Alce gigas (Miller).

Alces gigas Miller, Proc. Biol. Soc. Wash., XIII, 1899, p. 57. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 38. Zoöl. Ser.

ALASKAN MOOSE.

Type locality. Tustumena Lake, Kenai Peninsula, Alaska.

Geogr. Distr. Territory of Alaska.

Genus 40. Cervus. Wapiti. American Elk.

Cervus Linn., Syst. Nat., 1, 1758, p. 66. Type *Cervus elaphus* Linnæus.

Elaphus H. Smith, Griff. Anim. Kingd., v, 1827, p. 307.

Pseudocervus Hodgson, Journ. Asiat. Soc. Bengal, x, 1841, p. 914.

Strongyloceros! Owen, Brit. Foss. Mamm. & Birds, 1846, p. 470.

89. Cervus canadensis (Erxleben).

Cervus elaphus canadensis Erxl., Syst. Regn. Anim., 1, 1777, p. 305.

Cervus strongyloceros Schreb., Säugth., pl. CCXLVII, 1792?

Cervus wapiti Barton, Am. Phil. Trans., 1869, p. 70.

Cervus canadensis Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 34, pl. XII. Zoöl. Ser.

WAPITI. AMERICAN ELK.

Type locality. Eastern Canada.

Geogr. Distr. Rocky Mountain region from northern boundary of New Mexico to British Columbia, and from the eastern base of the range westward to about the 120th meridian. East of the westward borders of the great plains it is practically extinct, although it may still linger in northern Minnesota and possibly in North Dakota.

a. *Cervus canadensis occidentalis* (Smith).

Cervus occidentalis H. Smith, Griff. Cuv. Anim. Kingd., IV, 1827, p. 101.

Cervus roosevelti Merr., Proc. Biol. Soc. Wash., 1897, p. 271.

Cervus canadensis occidentalis Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 34. Zool. Ser.

PACIFIC COAST WAPITI.

Type locality. Northwestern America. No definite locality given.

Geogr. Distr. Vancouver Island, British Columbia; coast ranges of Washington, and Oregon to northwestern California.

90. *Cervus merriami* Nelson.

Cervus merriami Nels., Bull. Am. Mus. Nat. Hist. N. Y., 1902, pp. 1-12.

MERRIAM'S WAPITI.

Type locality. Black River, White Mountains, Arizona.

Geogr. Distr. Nearly extinct. Formerly throughout the high portion of the White and Mogollon Mountains of Arizona and New Mexico. Now limited to a small area of these.

Genl. Char. Nose darker, head and legs more reddish than *C. canadensis*, but paler than *C. c. occidentalis*. Nasals broader and flatter; upper molar series heavier and more curved. Antlers similar to those of *C. canadensis* but tines straighter.

Color. Summer Pelage. Head above reddish chestnut brown, paler on crown and sides; top of neck, back, and sides grizzled yellowish brown shading into pale dingy yellowish on flanks; rump patch dingy yellowish white bordered on lower edge with seal brown; under side of neck and body dark brown; legs dark reddish brown, paler on sides.

Winter Pelage. Body above, beneath, and on sides pale yellowish brown; head and neck seal brown, pale around eyes; ears whitish at base, liver brown on outer side; inner side of legs dark seal brown, outer side cinnamon rufous.

Measurements. Skull: occiput to front of premaxillæ, 498; palatal length, 288; length of nasals, 183; breadth of nasals, 83; orbital breadth, 194; breadth of premaxillæ, 99; zygomatic breadth, 203.

91. *Cervus nannodes* Merriam.

Cervus nannodes Merr., Proc. Biol. Soc. Wash., XVIII, 1905, p. 23. SAN JOAQUIN VALLEY ELK.

Type locality. Buttonwillow, Kern County, California.

Genl. Char. Size small; legs short; color pale; rump patch small; skull comparatively different from that of *C. canadensis*.

Color. Head and shoulders grizzled grayish brown; back grizzled whitish buffy; sides pale fulvous; rump patch and tail soiled whitish; anterior portion of legs bright golden fulvous, posterior portion buffy fulvous; mane grizzled grayish brown; thigh stripe faint; ears buffy ochraceous to ochraceous fulvous; the inside, borders, and outer base buffy white.

Measurements. "Type (two-year-old male). Total length, 2030; tail vertebræ, 140; hind foot, 620. Skull: basilar length, 358; zygomatic breadth, 155; occipito-sphenoid length, 79; palatal length, 230; palatal floor of premaxillaries, 82."

Genus 41. **Odontocælus. Deer.**

Odocoileus (*sic*) Rafin., Atlantic Journ., 1, 1832, No. 3, p. 109, fig. Type *Odocoileus! speleus* Rafin. = *Cervus dama americanus?* Erxl.

Mazama H. Smith, Griff. Anim. Kingd., v, 1827, p. 314. (nec Rafin.)

Dorcelaphus Glog., Hand- u. Hilfsb. Naturg., 1841, pp. xxxiii, 140.

Cariacus Less., Nouv. Tab. Règn. Anim., 1842, p. 173.

Oplacernus Haldeman, Proc. Acad. Nat. Scien. Phil., 1, 1842, p. 188.

Reduncina Wagn., Schreb. Säugth., Suppl., iv, 1844, p. 373, Taf. ccxlvii H.

Macrotis Wagn., Schreb. Säugth., Suppl., v, 1855, p. 368-372.

Eucervus Gray, Ann. Mag. Nat. Hist., 3d Ser., xviii, 1866, p. 338.

Otelaphus Fitzin., Sitzungsab. Math.-Nat. Cl. K. Akad. Wiss. Wien, lxxviii, Abth. 1, 1874, p. 356.

Gymnotis Fitzin., Sitzungsab. Math.-Nat. Cl. K. Akad. Wiss. Wien, lxxviii, Heft 11, Abth. 1, 1879, p. 343.

92. **Odontocælus americanus** (Erxleben).

Cervus dama americanus Erxl., Syst. Regn. Anim., 1777, p. 312.

Cervus virginianus Bodd., Elench. Anim., 1, 1785, p. 136.

Odocoileus! americanus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 39, pl. xv. Zool. Ser.

VIRGINIAN WHITE-TAILED DEER.

Type locality. Virginia.

Geogr. Distr. Eastern North America from New York to Florida; west to the Missouri River, excepting possibly Louisiana.

a. *Odontocælus americanus borealis* Miller.

Odocoileus! americanus borealis Miller, Bull. N. Y. State Museum, 1900, p. 83.

NORTHERN WHITE-TAILED DEER.

Type locality. Bucksport, Maine.

Geogr. Distr. Canadian zone, northern New York, New England, and southeastern Canada.

Genl. Char. Size large; lower tooth row 85 mm. in length. Winter pelage different from that of Summer.

Color. Like that of *O. americanus*.

b. *Odontocælus americanus macrourus* (Rafinesque).

Cervus macrourus Rafin., Amer. Month. Mag., 1817, p. 436.

Cervus leucurus Dougl., Zool. Journ., IV, 1829, p. 330.

Odocoileus! americanus macrourus Elliot, Syn. N. Am. Mamm.,

F. C. M. Pub., II, 1901, p. 39. Zool. Ser.

RAFINESQUE'S WHITE-TAILED DEER.

Type locality. Plains of the Kansas River, Upper Mississippi Valley.

Geogr. Distr. Kansas, Nebraska, and the Dakotas westward to California, Oregon, and Washington.

c. *Odontocælus americanus osceola* (Bangs).

Cariacus osceola Bangs, Proc. Biol. Soc. Wash., IX, 1896, p. 26.

Odocoileus! americanus osceola Elliot, Syn. N. Am. Mamm.,

F. C. M. Pub., II, 1901, p. 40. Zool. Ser.

OSCEOLA'S WHITE-TAILED DEER.

Type locality. Citronelle, Citrus County, Florida.

Geogr. Distr. Florida.

d. *Odontocælus americanus louisianæ* (Allen).

Odocæus! virginianus louisianæ G. M. Allen, Amer. Nat., xxxv, 1901, p. 449.

LOUISIANA WHITE-TAILED DEER.

Type locality. Mer Rouge, Morehouse County, Louisiana.

Geogr. Distr. Louisiana. Range undetermined.

Genl. Char. "Size large; skull long, slender; lower tooth row long; antlers heavy, high; color in winter pale."

Color. Top and sides of head grizzled brownish black and white; usual black patch about nostrils and lower jaw; chin, upper part of throat, and orbital ring white; upper parts grayish buff and black; sides of belly, throat, and chest ochraceous buff; inner sides of limbs, axillæ, and thighs white; tail above like the back, beneath white; metatarsal gland small, surrounded by white hairs, and a brownish black border; ears outside grizzled black and white, inside white.

Measurements. Nose to end of tail, 1695-2072; tail, 220-285; hind foot, 375-490; ear from crown, 122-134.5.

e. Odontocælus americanus texensis (Mearns).

Dorcelaphus texanus Mearns, Proc. Biol. Soc. Wash., 1898, p. 23.

Odocoileus! americanus texensis Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 40. Zoöl. Ser.

Odontocælus americanus texensis Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 70, pls. xxix, xxx. Zoöl. Ser.

TEXAN WHITE-TAILED DEER. *Venado* in Mexico for all deer.

Type locality. Fort Clark, Kinney County, Texas.

Geogr. Distr. Texas and northern Mexico.

f. Odontocælus americanus couesi (Rothrock Ms.)

Cariacus americanus var. *couesi* Rothrock, Ms. in Coues and Yar-row., Wheeler's Geog. and Geol. Surv. West 100th Merid., v, 1875, p. 72.

Odocoileus! americanus couesi Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 40. Zoöl. Ser.

Odontocælus americanus couesi Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 70, fig. xxiii. Zoöl. Ser.

COUES' WHITE-TAILED DEER.

Type locality. Camp Crittenden, Pima County, Arizona.

Geogr. Distr. Arizona, southwestern end of New Mexico, and State of Sonora to City of Mexico.

93. *Odontocælus battyi* Allen.

Odocoileus! battyi Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1903, p. 591.

Odontocælus battyi Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 71. Zoöl. Ser.

BATTY'S WHITE-TAILED DEER.

Type locality. Rancho Santuario, State of Durango, Mexico.

Geogr. Distr. State of Durango; range unknown.

94. *Odontocælus lichtensteini* (Allen).

Cervus mexicanus Lichtenstein, Darst. Thiere, 1827, p. 34, pl. xviii. (nec Gmelin.)

Dama lichtensteini Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1902, p. 20.

Odontocælus lichtensteini Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 72. Zoöl. Ser.

LICHTENSTEIN'S OR MEXICAN WHITE-TAILED DEER.

Type locality. Unknown.

Geogr. Distr. Southern Mexico; range unknown.

95. *Odontocælus rothschildi* (Thomas).

Dama rothschildi Thomas, Novitat. Zool., IX, 1902, p. 136.

Odontocælus rothschildi Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., IV, Pt. 1, 1904, p. 72. Zoöl. Ser.

ROTHSCHILD'S WHITE-TAILED DEER.

Type locality. Coiba Island, off west coast of Panama.

Geogr. Distr. Apparently restricted to Coiba Island.

96. *Odontocælus truii* Merriam.

Odocoileus! clavatus True, Proc. U. S. Nat. Mus., 1888, p. 417.

(nec H. Smith.)

Odocoileus! truei Merr., Proc. Biol. Soc. Wash., XII, 1898, p. 103.

(Footnote.)

Odontocælus truii Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., IV, Pt. 1, 1904, p. 73. Zoöl. Ser.

TRUE'S WHITE-TAILED DEER.

Type locality. Segovia River, Honduras.

Geogr. Distr. Honduras, Central America; range unknown.

97. *Odontocælus costaricensis* Miller.

Odocoileus! costaricensis Miller, Proc. Biol. Soc. Wash., XIV, 1901,

p. 35.

Odontocælus costaricensis Elliot, Mamm. Middle Amer. & W.

Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 73. Zoöl. Ser.

COSTA RICA WHITE-TAILED DEER.

Type locality. Talamanca, eastern side of Costa Rica, and the foot of the Cordilleras.

Geogr. Distr. Known only from Costa Rica.

98. *Odontocælus nemoralis* (H. Smith).

Cervus nemoralis H. Smith, Griff. Anim. Kingd., IV, 1827, p. 137.

Odontocælus nemoralis Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., IV, Pt. 1, 1904, p. 74. Zoöl. Ser.

HAMILTON SMITH'S WHITE-TAILED DEER.

Type locality. "Virginia"?

Geogr. Distr. Honduras to Panama, Central America.

99. *Odontocælus toltecus* (Saussure).

Cervus toltecus Sauss., Rev. Mag. Zool., 2me Sér., XII, 1860, p. 247.

Cervus yucatanensis Hays, Ann. Lyc. N. Y., X, 1874, p. 218.

Cervus acapulcensis Caton, Antel. & Deer Amer., 1877, p. 113.

Odontocælus toltecus Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., IV, Pt. 1, 1904, p. 74. Zoöl. Ser.

YUCATAN WHITE-TAILED DEER.

Type locality. Near Orizaba, State of Vera Cruz, Mexico.

Geogr. Distr. State of Vera Cruz to Yucatan, southeastern Mexico.

100. *Odontocælus thomasi* Merriam.

Odocoileus! thomasi Merr., Proc. Biol. Soc. Wash., XII, 1898, p. 102.

Odontocælus thomasi Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., IV, Pt. I, 1904, p. 75. Zoöl. Ser.

THOMAS' WHITE-TAILED DEER.

Type locality. Huehuetan, State of Chiapas, Mexico.

Geogr. Distr. States of Oaxaca and Chiapas, Mexico. Range unknown.

101. *Odontocælus nelsoni* Merriam.

Odocoileus! nelsoni Merr., Proc. Biol. Soc. Wash., XII, 1898, p. 103.

Odontocælus nelsoni Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., IV, Pt. I, 1904, p. 75. Zoöl. Ser.

NELSON'S WHITE-TAILED DEER.

Type locality. San Christobal, highlands of State of Chiapas, Mexico.

Geogr. Distr. State of Chiapas, Mexico.

102. *Odontocælus cerrosensis* Merriam.

Odocoileus! cerrosensis Merr., Proc. Biol. Soc. Wash., XII, 1898, p. 101.

Odontocælus cerrosensis Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., IV, Pt. I, 1904, p. 76. Zoöl. Ser.

CERROS ISLAND WHITE-TAILED DEER.

Type locality. Cerros or Cedros Island, off coast of Lower California, Mexico.

Geogr. Distr. Type locality only.

103. *Odontocælus sinaloæ* Allen.

Odocoileus! sinaloæ Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1903, p. 613.

Odontocælus sinaloæ Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., IV, Pt. I, 1904, p. 78. Zoöl. Ser.

SINALOA WHITE-TAILED DEER.

Type locality. Escuinapa, State of Sinaloa, Mexico.

Geogr. Distr. State of Sinaloa, Mexico; limits of range unknown.

104. *Odontocælus crooki* (Mearns.)

Dorcelaphus crooki Mearns, Proc. U. S. Nat. Mus., 1897, p. 468.

Odocoileus! crooki Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 41. Zoöl. Ser.

CROOK'S BLACK-TAILED DEER.

Type locality. Dog Mountains, Grant County, New Mexico.

Geogr. Distr. New Mexico, extent of range unknown.

105. *Odontocælus columbianus* (Richardson).

Cervus macrotis var. *columbiana* Rich., Faun. Bor. Amer., 1829, p. 257.

Cervus lewisii Peale, U. S. Expl. Exped., Mamm., VIII, 1848, p. 39, fig. p. 43.

Cervus richardsoni Aud. & Bach., Quad. N. Amer., III, 1853, p. 27, pl. CVI.

Cariacus punctulatus Gray, Proc. Zool. Soc., 1850, p. 239, pl. 28.

Odocoileus! columbianus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 41. Zool. Ser.

COLUMBIAN BLACK-TAILED DEER.

Type locality. Mouth of the Columbia River.

Geogr. Distr. British Columbia, Vancouver Island, and west of the Cascade Mountains in Washington, Oregon, and California.

a. *Odontocælus columbianus sitkensis* Merriam.

Odocoileus! columbianus sitkensis Merr., Proc. Biol. Soc. Wash., XII, 1898, p. 100. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 42. Zool. Ser.

SITKA BLACK-TAILED DEER.

Type locality. Sitka, Alaska.

Geogr. Distr. Alaska, limits of range unknown.

b. *Odontocælus columbianus scaphiotus* Merriam.

Odocoileus! columbianus scaphiotus Merr., Proc. Biol. Soc. Wash., XII, 1898, p. 101. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 42. Zool. Ser.

BLACK-TAILED DEER.

Type locality. Laguna Ranch, Gabilan Range, Riverside County, California.

Geogr. Distr. Gabilan Range, California; limits of dispersion unknown.

106. *Odontocælus hemionus* (Rafinesque).

Cervus hemionus Rafin., Amer. Month. Mag., I, 1817, p. 436.

Cervus auritus Ward., Desc. États Unis, v, 1820, p. 540.

Cervus macrotis Say, Narr. Long's Exped. Rocky Mts., II, 1823, p. 88.

Odocoileus! hemionus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 42. Zool. Ser.

Odontocælus hemionus Elliot, Mamm. Middle Amer. & W. Indies,
F. C. M. Pub., IV, Pt. I, 1904, p. 76. Zoöl. Ser.

MULE DEER.

Type locality. "Sioux River," probably on eastern border of South Dakota.

Geogr. Distr. West of Missouri River from Fort George south to Texas, westward through Nevada to latitude of San Francisco, including North and South Dakota, Nebraska, Kansas, Texas, Colorado, Wyoming, Montana, Idaho, California, Oregon, and Washington; Lower California, Mexico.

a. Odontocælus hemionus virgultus (Hallock).

Cariacus virgultus Hallock, Forest and Stream, 27th May, 1899,
p. 404.

MINNESOTA MULE DEER.

Type locality. Minnesota.

Geogr. Distr. Northeastern Minnesota and northwestern Manitoba, Rainy River, Lake of the Woods, Red Lake, and Roseau Region.

Genl. Char. Tail short, thick, uniform throughout its length. Antlers small, allied to *O. hemionus*.

Color. Body dark gray sprinkled with long black and white hairs; rump light gray; forehead black; face and throat light gray; breast and belly jet black; tail white on basal half, black on remainder; hocks tawny.

Measurements. Male. Height of shoulders, three feet five inches; nose to end of tail, five feet; girth of body, four feet; around neck, two feet. Antlers smaller than those of the Mule Deer.

b. Odontocælus hemionus californicus (Caton).

Cervus macrotis var. *californicus* Caton, Amer. Nat., x, 1876, p. 464.

Odocoileus! hemionus californicus Elliot, Syn. N. Am. Mamm.,
F. C. M. Pub., II, 1901, p. 43. Zoöl. Ser.

Odontocælus hemionus californicus Elliot, Mamm. Middle Amer. &
W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 77. Zoöl. Ser.

CALIFORNIA MULE DEER.

Type locality. St. Julian Ranch, summit of Gaviota Pass, Coast Range, forty miles from Santa Barbara, California.

Geogr. Distr. In the Coast Range south of San Francisco into Lower California, Mexico.

c. Odontocælus hemionus eremicus (Mearns).

Dorcelaphus hemionus eremicus Mearns, Proc. U. S. Nat. Mus.,
XX, 1897, p. 470.

Odontocælus hemionus eremicus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 77. Zoöl. Ser.
DESERT MULE DEER.

Type locality. Sierra Seri, near the Gulf of California, State of Sonora, Mexico.

Geogr. Distr. Lower California and State of Sonora, Mexico. Limits of range unknown.

d. *Odontocælus hemionus canus* Merriam.

Odocoileus! hemionus canus Merr., Proc. Wash. Acad. Scien., III, 1901, p. 560.

Odontocælus hemionus canus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 78. Zoöl. Ser.

CHIHUAHUA MULE DEER.

Type locality. Sierra en Media, State of Chihuahua, Mexico.

Geogr. Distr. State of Chihuahua, limits of range unknown.

e. *Odontocælus hemionus peninsulæ* (Lydekker).

Mazama hemionus peninsulæ Lydekk., Proc. Zoöl. Soc., 1897, pp. 899-900.

Odontocælus hemionus peninsulæ Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 78. Zoöl. Ser.

LOWER CALIFORNIA MULE DEER.

Type locality. La Paz, Lower California, Mexico.

Geogr. Distr. Cape Region of Lower California.

Genus 42. **Hippocamelus.** The Brockets.

Hippocamelus Leuckart, Dissert. Inaug. de *Equo bisulco* Molinæ, 1816, p. 24. Type *Equus bisulcus* Molina = *Hippocamelus dubius* Leuckart.

Mazama Rafin., Amer. Month. Mag., I, 1817, p. 363.

107. **Hippocamelus sartori** (Saussure).

Cervus sartorii Sauss., Rev. Mag. Zool., 2me Sér., XII, 1860, p. 252.

Mazama sartori Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 80, fig. xxiv, pl. xxxi. Zoöl. Ser.

SARTORI'S BROCKET. *Cabra del Monte* in Costa Rica.

Type locality. Mirador, State of Vera Cruz, Mexico.

Geogr. Distr. Southern Mexico and Central America.

108. Hippocamelus pandora (Merriam).

Mazama pandora Mérr., Proc. Biol. Soc. Wash., XIV, 1901, p. 105.

Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV,

Pt. I, 1904, p. 80. Zoöl. Ser.

TUNKAS BROCKET.

Type locality. Tunkas, Yucatan, Mexico.

Geogr. Distr. State of Campeche, and Yucatan, Mexico.

Family III. **Antilocapridæ. Prong-horn Antelopes.**Genus 43. **Antilocapra.**

Antilocapra Ord, Journ. de Phys., 1818, p. 149. Type *Antilope americana* Ord.

**Dicranocerus!* H. Smith, Griff. Anim. Kingd., IV, 1827, p. 169.

109. Antilocapra americana (Ord).

Antilope americana Ord, Guth. Geog., 2d Am. ed., II, 1815, p. 292, desc. p. 308.

Antilocapra americana Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 43, pl. XVI. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 82. Zoöl. Ser.

PRONG-HORN ANTELOPE.

Type locality. Plains east of the Missouri? Black Mountains?

Geogr. Distr. Valley of the Saskatchewan, latitude 53°, south to Mexico, and from the Missouri River on the plains, westward to the Rocky Mountains and the Cascade Range in Oregon and Washington.

a. Antilocapra americana mexicana Merriam.

Antilocapra americana mexicana Merr., Proc. Biol. Soc. Wash., XIV, 1901, p. 31. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 82, fig. XXV, pls. XXXII, XXXIII. Zoöl. Ser.

MEXICAN PRONG-HORN. *Berendo* in Mexico.

Type locality. Sierra en Media, State of Chihuahua, Mexico.

Geogr. Distr. Northern Mexico, in States of Sonora, Chihuahua, and Tamaulipas; Lower California.

*διεφραγος, two-headed, + κέρας, horn, = Dicranoceras.

Family IV. **Bovidae. Oxen, Sheep, Goats.**Subfamily I. **Rupicaprinæ.**Genus 44. **Oreamnus. Goats.**

Oreamnos! Rafin., Amer. Month. Mag., 11, 1817, p. 44. Type

Ovis montanus Ord.

Aplocerus (sic) H. Smith, Griff. Anim. Kingd., v, 1827, p. 354.

110. Oreamnus montanus (Ord).

Ovis montanus Ord, Guth. Geogr., 2d Amer. ed., 11, 1815, p. 292.

Oreamnus montanus Elliot, Syn. N. Am. Mamm., F. C. M. Pub.,

11, 1901, p. 44, pls. XVII, XVIII, XIX, XX. Zoöl. Ser.

MOUNTAIN GOAT.

Type locality. Cascade Range near Brant Island, Columbia River, Oregon.

Geogr. Distr. High elevations in Rocky Mountains and other ranges between them and the Pacific, to southern British Columbia. Not found on the Olympic Mountains, Washington.

a. Oreamnus montanus columbianus Allen.

Oreamnos! montanus columbianus Allen, Bull. Am. Mus. Nat. Hist.

N. Y., 1904, p. 20.

COLUMBIAN MOUNTAIN GOAT.

Type locality. Shesley Mountains, northern British Columbia.

Genl. Char. Size large, skull narrow.

Color. White.

Measurements. Femur to humerus, 914; height at shoulder, 1088. Skull: occipito-nasal length, 336; Hensel, 293; zygomatic breadth, 119; interorbital width, 98; width of maxillary region above second upper molar, 87; nasals, 122 x 38; palatal length, 184; length of upper tooth row, 77; ratio of interorbital breadth to basal length, 35; hoof, outer digit on fore foot, 74 x 35; outer digit on hind foot, 66 x 33.

b. Oreamnus montanus missoulæ Allen.

Oreamnos! montanus missoulæ Allen, Bull. Am. Mus. Nat. Hist.

N. Y., 1904, p. 20.

MISSOULA MOUNTAIN GOAT.

Type locality. Missoula, Montana.

Genl. Char. Size small; skull narrow.

Color. White.

Measurements. Skull: occipito-nasal length, 299; zygomatic breadth, 108; interorbital width, 87; width of intermaxillary region

above second upper molar, 81; nasals, 104 x 33; length of upper tooth row, 69; hoof, at outer digit of fore foot, 56 x 28; at outer digit of hind foot, 52 x 26.

111. *Oreamnus kennedyi* Elliot.

Oreamnus kennedyi Elliot, Pub. Field Columb. Mus., III, 1900, p. 3. Zoöl. Ser. *Id. Syn. N. Am. Mamm.*, F. C. M. Pub., II, 1901, p. 44, pls. XVII, XVIII, XIX, XX.

KENNEDY'S MOUNTAIN GOAT.

Type locality. Mountains at the mouth of Copper River, opposite Kyak Island, Alaska.

Geogr. Distr. Coast mountains of Alaska from Copper River to vicinity of Knik near Cook's Inlet.

Genus 45. *Ovis*. Sheep.

Ovis Linn., *Syst. Nat.*, I, 1758, p. 70; I, 1766, p. 97. Type *Ovis aries* Linnæus.

112. *Ovis cervina* Desmarest.

Ovis cervina Desm., *Nouv. Dict. Hist. Nat.*, 1804, p. 5; nouv. ed., 1818, XXI, p. 553. Elliot, *Syn. N. Am. Mamm.*, F. C. M.

Pub., II, 1901, p. 46, pl. XXI. Zoöl. Ser.

Ovis canadensis Shaw, *Nat. Misc.*, 1804? pl. 610.

Ovis ammon Mitch., *Med. Rep.*, 1807, p. 35. (nec Linn.)

Ovis pygargus H. Smith, *Griff. Anim. Kingd.*, v, 1827, p. 318.

Ovis californianus Dougl., *Zoöl. Journ.*, 1828, p. 332.

MOUNTAIN SHEEP.

Type locality. Rocky Mountains between the Kootenai Pass and Mount Mead, N. lat. 50°, W. long. 115, Alberta, Canada.

Geogr. Distr. High mountain ranges from the Colorado River and Arizona north into British Columbia.

a. **Ovis cervina auduboni* (Merriam).

Ovis canadensis auduboni Merr., *Proc. Biol. Soc. Wash.*, XIV, 1901, p. 31.

Ovis cervina auduboni Elliot, *Syn. N. Am. Mamm.*, F. C. M. Pub., II, 1901, Suppl., p. 484. Zoöl. Ser.

AUDUBON'S MOUNTAIN SHEEP.

Type locality. "Upper Missouri."

Geogr. Distr. Unknown.

*Very doubtfully separable from *O. cervina*.

b. *Ovis cervina nelsoni* (Merriam).

Ovis nelsoni Merr., Proc. Biol. Soc. Wash., XI, 1897, p. 217.

Ovis cervina nelsoni Elliot, Syn. N. Am. Mamm., F. C. M. Pub.,
II, 1901, p. 46. Zoöl. Ser.

NELSON'S MOUNTAIN SHEEP.

Type locality. Grapevine Mountains, boundary line of Nevada and California.

Geogr. Distr. Mountains of southeastern California.

c. *Ovis cervina cremnobates* Elliot.

Ovis cervina cremnobates Elliot, Pub. Field Columb. Mus., III, 1903,
p. 239. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies,
F. C. M. Pub., IV, Pt. I, 1904, p. 84, figs. xxvi, xxvii. Zoöl.
Ser.

LOWER CALIFORNIA MOUNTAIN SHEEP.

Type locality. Mattomi, San Pedro Martir Mountains, Lower California, Mexico.

Geogr. Distr. San Pedro Martir Range, and probably also the Laguna Mountains, Lower California, Mexico.

d. *Ovis cervina mexicanus* (Merriam).

Ovis mexicanus Merr., Proc. Biol. Soc. Wash., XIV, 1901, p. 30.

Ovis cervina mexicanus Elliot, Mamm. Middle Amer. & W. Indies,
F. C. M. Pub., IV, Pt. I, 1904, p. 86, pls. xxxiv, xxxv. Zoöl.
Ser.

MEXICAN MOUNTAIN SHEEP.

Type locality. Mountains about Lake Santa Maria, State of Chihuahua.

Geogr. Distr. Sierra Madre and Guadalupe Mountains of northern Mexico. Limits of range unknown.

113. *Ovis stonii* Allen.

Ovis stonei Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1897, p. 111,
pls. II, III. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II,
1901, p. 47. Zoöl. Ser.

Ovis canadensis liardensis Lydekk., Wild Oxen, Sheep & Goats,
1898, p. 215.

STONE'S SHEEP.

Type locality. Che-on-nee Mountains, Coast Range, British Northwest Territory.

Geogr. Distr. From the Cassiar Mountains 61° north latitude and south to the head waters of the Nelson and Peace Rivers, Rocky Mountains; west to longitude 134°. Also found in the Stickeen, Cheonnees, and Etsazas Mountains, Alaska.

114. *Ovis dalli (Nelson).

Ovis montana dalli Nelson, Proc. U. S. Nat. Mus., 1884, p. 12.

Ovis dalli Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 47. Zoöl. Ser.

DALL'S MOUNTAIN SHEEP.

Type locality. Fort Reliance, Upper Yukon River, Alaska.

Geogr. Distr. Alaskan Mountains, to near the Arctic coast west of the Mackenzie River, and south possibly to north latitude 60° and westward to Kotzebue Sound.

a. *Ovis dalli kenaiensis Allen.

Ovis dalli kenaiensis Allen, Bull. Am. Mus. Nat. Hist., 1902, p. 145.

KENAI MOUNTAIN SHEEP.

Type locality. Kenai Peninsula, Alaska.

Geogr. Distr. Limits of range unknown. Probably not restricted to the Kenai Peninsula.

Genl. Char. Color white. Skull: posterior nares short; teeth broad and short; tooth rows convex outwardly; palate broad; upper side of tail white.

Color. White.

Measurements. Skull: base of postoccipital process to third upper molar, 84; posterior nares, 58-60.

115. *Ovis fannini Hornaday.

Ovis fannini Hornaday, Rep. N. Y. Zoöl. Soc., 1901, Append. 1, p. 2. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, Append., p. 425. Zoöl. Ser.

FANNIN'S MOUNTAIN SHEEP.

Type locality. Rocky Mountains east side of the Yukon about seventy-five miles east of Dawson, Yukon, N. W. Territory.

Geogr. Distr. In the mountains above named, but the extent of range unknown.

Genus 46. Ovibos. Musk Oxen.

Ovibos De Blainv., Bull. Soc. Philom. Paris, 1816, p. 76. Type *Bos moschatus* Zimmermann.

116. Ovibos moschatus (Zimmermann).

Bos moschatus Zimm., Geog. Geschichte, 11, 1780, p. 86.

Ovibos pallantis H. Smith, Griff. Anim. Kingd., IV, 1827, p. 375.

*The relative specific standing of these sheep is not yet satisfactorily established.

Bos pallasi De Kay, Ann. N. Y. Lyc. Nat. Hist., II, 1828, p. 29.
(nec Baer.)

Bos canaliculatis Fisch., Mém. Acad. Mosc., III, 1834, p. 287.

Ovibos moschatus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II,
1901, p. 48, pl. XXII. Zoöl. Ser.

MUSK OX.

Type locality. Churchill River, Kewatin, Canada.

Geogr. Distr. Arctic America north of 60th parallel from Mackenzie River to region of Hudson Bay.

a. *Ovibos moschatus wardi* Lydekker.

Ovibos moschatus wardi Lydekk., Nature, LXIII, 1900, p. 157.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, Suppl.
p. 485. Zoöl. Ser.

WARD'S MUSK OX.

Type locality. East Greenland.

Geogr. Distr. "Southern border of Ellesmere Land northward through Grinnell Land to the Polar Sea, and on the Greenland coast from about latitude 78°, on the western side northward to and around the northern end of Greenland to about latitude 75° N. on the east coast."

b. *Ovibos moschatus niphæcus* Elliot.

Ovibos moschatus niphæcus Elliot, Proc. Biol. Soc. Wash., XVIII,
1905, p. 35.

BLACK MUSK OX.

Type locality. About 600 miles north of Hudson Bay

Genl. Char. In color intermediate between *O. moschatus* and *O. m. wardi*; cranial characters different.

Color. Type, bull in prime of life. A narrow whitish band between ears, behind the bases of horns, and a small brown saddle on middle of back; rest of head, neck, and body jet black. Fore legs grayish in front, black behind, changing into grayish white above the hoofs; hind legs black in front, becoming gray above hoof; grayish on sides and hinder part. Long black hairs on body covering the legs to the knees. Nose and lips and chin grayish white; ears black.

The young bull, female, and calves resemble the type in their jet black hue and in the varying color of the legs. The old bull is of a dark brown hue, becoming black upon the flanks, but no white whatever showing anywhere. The old cow is of the same general color as the aged bull, but has a little white upon the sides of the nose.

Measurements. Skull: total length, 430 mm.; occipito-nasal length, 355; greatest breadth across orbits, 245; length of nasals, median, 121; greatest width posteriorly, 64; anterior width (at tips), 13; zygomatic width, 168; palatal arch to end of premaxillæ, 249; greatest width of palatal floor between fourth premolars, 76; length of horn core from edge on top of head to tip, 230; length of upper tooth row, alveolar border outer side, 140; posterior width of basi-occipital, 62; anterior width in front of bullæ, 32; length of mandible, 345; length of lower tooth row, alveolar border outer side, 145.

Genus 47. **Bison. Bison, American Buffalo.**

Bison H. Smith, Griff. Anim. Kingd., v, 1827, p. 373. Type *Bos bison* Linnæus.

117. Bison bison (Linnæus).

Bos bison Linn., Syst. Nat., I, 1758, p. 72.

Bison americanus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 49, pl. XXIII. Zoöl. Ser.

Bison bison Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, Suppl., p. 486. Zoöl. Ser.

BISON. AMERICAN BUFFALO.

Type locality. Texas.

Geogr. Distr. Practically extinct in the wild state; a few survive in the National Park, and some are said to be in the Pan Handle of Texas.

a. Bison bison athabascæ Rhoads.

Bison bison athabascæ Rhoads, Proc. Acad. Nat. Scien. Phil., 1897, p. 498. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, Suppl., p. 486. Zoöl. Ser.

Bison americanus athabascæ Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 49. Zoöl. Ser.

WOOD BISON.

Type locality. Fifty miles southwest of Fort Resolution, Great Slave Lake, Mackenzie, Canada.

Geogr. Distr. "Wooded uplands of Northwest Territories. Formerly from the east slope of the Rocky Mountains to the 95th meridian, and from latitude 63° to latitude 55°; probably south along the Rocky Mountains to the United States."

Family V. **Tapiridae. Tapirs.**Genus 48. **Tapirella.**

Tapirella Palmer, Science, N. Ser., xvii, 1903, p. 873.

Elasmognathus Gill, Proc. Acad. Nat. Scien. Phil., 1865, p. 183.
(nec Fieber, Hemiptera.) Type *Elasmognathus bairdii* Gill.

118. **Tapirella bairdi** (Gill).

Elasmognathus bairdii Gill, Proc. Acad. Scien. Phil., 1865, p. 183.

Tapirella bairdi Elliot, Mamm. Middle Amer. & W. Indies, F. C. M.
Pub., iv, Pt. 1, 1904, p. 87, pls. xxxvi, xxxvii. (Name on
plate misprinted *Tapirella dowi*.) Zool. Ser.

BAIRD'S TAPIR.

Type locality. Isthmus of Panama.

Geogr. Distr. Southern Mexico to Panama, Central America.

119. **Tapirella dowi** (Gill).

Elasmognathus dowii Gill, Amer. Journ. Scien. Arts, L, 1870, p. 142.

Tapirella dowi Elliot, Mamm. Middle Amer. & W. Indies, F. C.
M. Pub., iv, Pt. 1, 1904, p. 88, fig. xxviii. Zool. Ser.

DOW'S TAPIR.

Type locality. Guatemala.

Geogr. Distr. Guatemala and Nicaragua, Central America.

Order VI. **Rodentia. Rodents.**

Family I. **Sciuridæ. Squirrels, Marmots, etc.**

Subfamily I. **Sciurinae.**

Genus 49. **Synthetosciurus.**

Syntheosciurus (*sic*) Bangs, Bull. Mus. Comp. Zool., xxxix, 1902, p. 25.

Type *Syntheosciurus! brochus* Bangs.

120. **Synthetosciurus brochus** Bangs.

Syntheosciurus! brochus Bangs, Bull. Mus. Comp. Zool., xxxix, 1902, p. 25.

Synthetosciurus brochus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 91, fig. 12. Zool. Ser.

PROJECTING-TEETH SQUIRREL.

Type Locality. Boquete, Chiriqui, Panama. Altitude, 7,000 feet.

Geogr. Distr. Panama, Central America.

Genus 50. **Sciurus. Tree Squirrels.**

Sciurus Linn., Syst. Nat., I, 1758, p. 63. Type *Sciurus vulgaris* Linnæus.

Guerlinguetus Gray, Lond. Med. Rep., xv, 1821, p. 304.

Macroxus F. Cuv., Dent's des Mamm., 1823, p. 162, pl. 56.

Rhinosciurus Gray, List Spec. Mamm. Brit. Mus., 1843, pp. xxv, 195. *Id.* Ann. Mag. Nat. Hist., 3d Ser., xx, 1867, p. 286.

Rheithrosciurus (*sic*) Gray, Ann. Mag. Nat. Hist., 3d Ser., xx, 1867, p. 272.

Neosciurus Treuss., Le Nat., II, 1880, p. 292.

Parasciurus Treuss., Le Nat., II, 1880, p. 292.

Echinosciurus Treuss., Le Nat., II, 1880, p. 292.

Tamiasciurus Treuss., Le Nat., II, 1880, p. 292.

Microsciurus Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1895, p. 332.

Hesperosciurus Nelson, Proc. Wash. Acad. Scien., I, 1899, p. 27.

Otosciurus Nelson, Proc. Wash. Acad. Scien., I, 1899, p. 28.

Aræosciurus Nelson, Proc. Wash. Acad. Scien., I, 1899, p. 29.

Baiosciurus Nelson, Proc. Wash. Acad. Scien., I, 1899, p. 31.

Subgenus A. *Microsciurus*.**121. *Sciurus alfari* Allen.**

Sciurus alfari Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1895, p. 333. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 99. Zoöl. Ser.

ALFARO'S PIGMY SQUIRREL.

Type locality. Jimenez, Costa Rica, Central America.

Geogr. Distr. Northern, eastern, and southwestern Costa Rica.

122. *Sciurus browni* Bangs.

Sciurus browni Bangs, Bull. Mus. Comp. Zoöl., xxxix, 1902, p. 24. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 100, fig. 13. Zoöl. Ser.

BROWN'S PIGMY SQUIRREL.

Type Locality. Bogava, Chiriqui, Panama.

Geogr. Distr. Panama, Central America.

123. *Sciurus boquetensis* Nelson.

Sciurus boquetensis Nelson, Proc. Biol. Soc. Wash., xvi, 1903, p. 121. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 100. Zoöl. Ser.

CHIRIQUI PIGMY SQUIRREL.

Type locality. Boquete, Chiriqui, Panama. Altitude, 6,000 feet.

Geogr. Distr. Panama, Central America.

Subgenus B. *Baiosciurus*.**124. *Sciurus deppii* Peters.**

Sciurus deppei Peters, Monatsb. K. Preuss. Akad. Wiss. Berl., 1863, p. 654.

Sciurus tephrogaster Gray, Ann. Mag. Nat. Hist., 3d Ser., xx, 1867, p. 431.

Sciurus griseogena Gray, Ann. Mag. Nat. Hist., 3d Ser., xx, 1867, p. 429.

Sciurus tæniurus Gray, Ann. Mag. Nat. Hist., 3d Ser., xx, 1867, p. 431.

Sciurus deppii Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 101, fig. 14. Zoöl. Ser.

DEPPE'S SQUIRREL.

Type locality. Papantla, State of Vera Cruz, Mexico.

Geogr. Distr. State of Vera Cruz from Papantla on east coast to Isthmus of Tehuantepec, Mexico, and into Guatemala, Central America. Altitude, 6,000-9,000 feet.

a. *Sciurus deppii vivax* Nelson.

Sciurus deppii vivax Nelson, Proc. Biol. Soc. Wash., xiv, 1901, p. 131.

Sciurus deppii vivax Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 102. Zoöl. Ser.

APAZOTE SQUIRREL.

Type locality. Apazote, State of Campeche, Mexico.

Geogr. Distr. Lowland forest in the States of eastern Tabasco, southern Campeche, and southern and eastern Yucatan, Mexico.

125. *Sciurus negligens* Nelson.

Sciurus negligens Nelson, Proc. Biol. Soc. Wash., xii, 1898, p. 147. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 102. Zoöl. Ser.

Sciurus arizonensis Alston, Biol. Centr. Amer., Mamm., 1, 1880, p. 125. (Part.)

Sciurus deppii Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1891, p. 222. (Part.)

LITTLE GRAY SQUIRREL.

Type locality. Alta Mira, State of Tamaulipas, Mexico.

Geogr. Distr. Southern Tamaulipas, through eastern San Luis Potosi into northern Vera Cruz, Mexico.

Subgenus C. *Guerlinguetus*.

Sciurus æstuans hoffmanni Peters.

Sciurus æstuans var. *hoffmanni* Peters, Monatsb. K. Preuss. Akad. Wiss. Berlin, 1863, p. 654. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 104, fig. 15. Zoöl. Ser.

Sciurus xanthotus Gray, Ann. Mag. Nat. Hist., 3d Ser., xx, 1867, p. 429.

Sciurus griseogena Gray, Ann. Mag. Nat. Hist., 3d Ser., xx, 1867, p. 430. (Part. Costa Rica.)

Sciurus rufoniger Allen, Mon. N. Am. Rod., 1877, pp. 757-763. (Part. Costa Rica.)

Sciurus griseogenys Alston, Proc. Zoöl. Soc., 1878, p. 667. (Part. Costa Rica, Veragua, Panama.)

HOFFMANN'S SQUIRREL.

Type locality. Costa Rica, Central America.

Geogr. Distr. Costa Rica south to upper Cauca River, Columbia.

Sciurus æstuans chiriquensis Bangs.

Sciurus æstuans chiriquensis Bangs, Bull. Mus. Comp. Zool.,
xxxix, 1902, p. 22. Elliot, Mamm. Middle Amer. & W.
Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 104. Zool. Ser.

CHIRIQUI SQUIRREL.

Type locality. Divala, Chiriqui, Panama.

Geogr. Distr. Panama, Central America.

126. *Sciurus richmondi* Nelson.

Sciurus richmondi Nelson, Proc. Biol. Soc. Wash., xii, 1898, p.
146. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M.
Pub., iv, Pt. 1, 1904, p. 105. Zool. Ser.

RICHMOND'S SQUIRREL.

Type locality. Escondido River, fifty miles above Bluefields,
Nicaragua.

Geogr. Distr. Tropical lowland forests along the Escondido River,
Nicaragua.

Sciurus variabilis morulus Bangs.

Sciurus variabilis morulus Bangs, Proc. N. Eng. Zool. Club, ii,
1900, p. 43. Elliot, Mamm. Middle Amer. & W. Indies,
F. C. M. Pub., iv, Pt. 1, 1904, p. 105. Zool. Ser.

Sciurus variabilis True, Proc. U. S. Nat. Mus., vii, 1884, p. 596.

LION HILL SQUIRREL.

Type locality. Loma del Leon, Panama.

Geogr. Distr. Panama, Central America.

Subgenus D. *Aræosciurus*.127. *Sciurus oculus* Peters.

Sciurus oculus Peters, Monatsb. K. Preuss. Akad. Wiss. Berlin,
1863, p. 653. Elliot, Mamm. Middle Amer. & W. Indies,
F. C. M. Pub., iv, Pt. 1, 1904, p. 107. Zool. Ser.

Sciurus capistratus Licht., Abh. K. Akad. Wiss., 1827, p. 116.
(nec Bosc.)

Sciurus carolinensis Sauss., Rev. Mag. Zool. 2me Sér., xiii, 1861,
p. 4. (nec Gmel.)

Sciurus hypopyrrhus Allen, Bull. U. S. Geol. Surv. Terr., iv,
1878, p. 881. (Part.)

Sciurus niger melanonotus Thomas, Proc. Zool. Soc., 1890, p. 73,
pl. vi.

BLACK-BACKED SQUIRREL.

Type locality. Las Vigas? State of Vera Cruz, Mexico.

Geogr. Distr. Northwestward through States of Vera Cruz, Hidalgo, Queretaro to southeastern San Luis Potosi, Mexico. Altitude, 6,000 to 12,000 feet.

a. *Sciurus oculatus tolucae* Nelson.

Sciurus oculatus tolucae Nelson, Proc. Biol. Soc. Wash., xii, 1898, p. 148. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 107. Zool. Ser.

TOLUCA SQUIRREL.

Type locality. North slope of the Volcano of Toluca, State of Mexico, Mexico.

Geogr. Distr. Sierra Madre from Toluca Volcano, State of Mexico, to border of the State of Michoacan; southern and western parts of State of Queretaro; central and eastern parts of State of Guanajuato, and southern part of State of San Luis Potosi, Mexico.

128. *Sciurus alleni* Nelson.

Sciurus alleni Nelson, Proc. Biol. Soc. Wash., xii, 1898, p. 147. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 108. Zool. Ser.

Sciurus carolinensis? Baird, N. Am. Mamm., 1857, p. 263.

Sciurus carolinensis var. *carolinensis* Allen, Mon. N. Am. Rod., 1877, p. 706.

Sciurus carolinensis Alston, Proc. Zool. Soc., 1878, p. 658.

Sciurus arizonensis Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1891, p. 222. (Part.)

ALLEN'S SQUIRREL.

Type locality. Monterey, State of Nuevo Leon, Mexico.

Geogr. Distr. From Monterey, State of Nuevo Leon, into State of Tamaulipas, Mexico. Altitude, 2,000-8,500 feet.

129. *Sciurus nayaritensis* Allen.

Sciurus nayaritensis Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1890, p. vii, footnote, and p. 185. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 108, fig. 16. Zool. Ser.

Sciurus alstoni Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1889, p. 167. (nec Anderson.)

NAYARIT SQUIRREL.

Type locality. Sierra de Valparaiso, State of Zacatecas, Mexico.

Geogr. Distr. States of Jalisco and Zacatecas, Mexico.

130. *Sciurus arizonensis* Coues.

Sciurus arizonensis Coues, Amer. Nat., i, 1867, p. 357. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., ii, 1901, p. 59, fig. 14.

Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies., F. C. M. Pub., IV, Pt. 1, 1904, p. 109. Zoöl. Ser.

Sciurus collaei Allen, Mon. N. Am. Rod., 1877, p. 738. (nec Rich.)
ARIZONA GRAY SQUIRREL.

Type locality. Fort Whipple, Yavapai County, Arizona.

Geogr. Distr. State of Nuevo Leon, Mexico (San Pedro Mines, Allen), to Arizona. Texas?

a. *Sciurus arizonensis huachuca* Allen.

Sciurus arizonensis huachuca Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1894, p. 349. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 60. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 109. Zoöl. Ser.

HUACHUCA SQUIRREL.

Type locality. Huachuca Mountains, southern Arizona.

Geogr. Distr. Huachuca Mountains, Arizona, into State of Sonora, Mexico.

131. *Sciurus apache* Allen.

Sciurus apache Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1893, p. 29. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 58. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 110. Zoöl. Ser.

Sciurus griseoflavus Thomas, Proc. Zoöl. Soc., 1882, p. 372. (nec Gray.)

Sciurus niger ludovicianus Thomas, Proc. Zoöl. Soc., 1890, p. 73 (footnote).

APACHE SQUIRREL.

Type locality. Mountains of northwestern Chihuahua, Mexico.

Geogr. Distr. Sierra Madre in States of western Durango, northwestern Chihuahua, eastern Sonora, and northeastern Sinaloa, Mexico; also in Chiricahua Mountains, southern Arizona.

Subgenus E. *Parasciurus*.

132. *Sciurus rufiventer* E. Geoffroy.

Sciurus rufiventer E. Geoff., Cat. Mamm. Mus. Hist. Nat. Paris, 1803, p. 176. Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1902, p. 167.

Sciurus ludovicianus Custis, Bart. Med. & Phys. Journ., II, Pt. II, 1806, p. 47. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 52. Zoöl. Ser.

Sciurus macroura Say, Long's Exped. Rocky Mountains, I, 1823, p. 115. (Kansas.)

Sciurus magnicaudatus Harlan, Faun. Amer., 1825, p. 178.

Sciurus subauratus Bachm., Proc. Zool. Soc., 1838, p. 87.

Sciurus auduboni Bachm., Proc. Zool. Soc., 1838, p. 97.

Sciurus occidentalis Aud. & Bachm., Journ. Acad. Nat. Scien. Phila., VIII, 1842, p. 317.

Sciurus sayi Aud. & Bachm., Quad. N. Amer., II, 1851, p. 274, pl. LXXXIX.

FOX SQUIRREL.

Type locality. Red River of Louisiana?

Geogr. Distr. Mississippi Valley from Louisiana to South Dakota and Michigan; and from the western Alleghanies to eastern border of the plains.

a. *Sciurus rufiventer texensis* (Bachman).

Sciurus texianus! Bach., Proc. Zool. Soc., 1838, p. 86.

Sciurus ludovicianus limitis Baird, Proc. Acad. Nat. Scien. Phil., 1855, p. 331. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 53. Zool. Ser.

Sciurus rufiventer texianus! Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1902, p. 167.

Sciurus rufiventer texensis Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 110, fig. 17. Zool. Ser.

TEXAS FOX SQUIRREL.

Type locality. Devil's River, Valverde County, Texas.

Geogr. Distr. Texas and adjacent parts of the States of Nuevo Leon and Coahuila, Mexico.

b. *Sciurus rufiventer neglectus* (Gray).

Macroxus neglectus Gray, Ann. Mag. Nat. Hist., 3d Ser., xx, 1867, p. 425.

Sciurus vulpinus Schreb., Säugth., IV., 1792, p. 772. (nec Gmel.)

Sciurus cinereus Le Conte, App. McMurt., Cuv. Règn. Anim., 1831, p. 433.

Sciurus ludovicianus vicinus Bangs, Proc. Biol. Soc. Wash., x, 1896, p. 150. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 52. Zool. Ser.

Sciurus ludovicianus neglectus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, Append., p. 426. Zool. Ser.

Sciurus rufiventer neglectus Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1902, p. 22.

EASTERN FOX SQUIRREL.

Type locality. Wilmington, Delaware.

Geogr. Distr. Formerly from northern Virginia to southern New England; now from mountains of West Virginia northward into western Pennsylvania.

133. *Sciurus niger* Linnæus.

Sciurus niger Linn., Syst. Nat., 1, 1758, p. 64. Elliot, Syn. N. Am.

Mamm., F. C. M. Pub., 11, 1901, p. 51, fig. 10. Zool. Ser.

Sciurus cinereus Linn., Syst. Nat., 1, 1758, p. 64. (Part.)

Sciurus vulpinus Gmel., Syst. Nat., 1, 1788, p. 147.

Sciurus capistratus Bosc., Ann. Mus. Paris, 1, 1808, p. 281.

BLACK FOX SQUIRREL.

Type locality. South Carolina?

Geogr. Distr. Virginia to Florida east of the Alleghanies; and on the coast of the Gulf of Mexico to Louisiana.

Subgenus F. *Otosciurus*.

134. *Sciurus aberti* Woodhouse.

Sciurus aberti Woodh., Proc. Acad. Nat. Scien. Phil., 1852, p.

220. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901,

p. 57, fig. 13. Zool. Ser.

Sciurus dorsalis Woodh., Proc. Acad. Nat. Scien. Phil., 1852, p.

110. (nec Gray.)

ABERT'S SQUIRREL.

Type locality. San Francisco Mountains, Coconino County, Arizona.

Geogr. Distr. New Mexico, Arizona, and mountains of Colorado.

a. *Sciurus aberti mimus* Merriam.

Sciurus aberti mimus Merr., Proc. Biol. Soc. Wash., xvii, 1904,

p. 130.

MIMIC SQUIRREL.

Type locality. Hall Peak, Cimarron Mountains, New Mexico.

Genl. Char. Similar to *S. aberti*, but paler on upper parts, and red dorsal area nearly obsolete; tail paler; ear tufts pale fulvous.

Measurements. Total length, 485; tail vertebræ, 215; hind foot, 70.

b. *Sciurus aberti ferreus* True.

Sciurus aberti concolor True, Proc. U. S. Nat. Mus., 1894, p. 241.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 57.

Zool. Ser.

Sciurus aberti ferreus True, Proc. Biol. Soc. Wash., xiii, 1900,

p. 183.

PLAIN-BACKED SQUIRREL.

Type locality. Loveland, Larimer County, Colorado.

Geogr. Distr. Northeastern Colorado.

135. *Sciurus kaibabensis* Merriam.

Sciurus kaibabensis Merr., Proc. Biol. Soc. Wash., xvii, 1904, p. 129.

KAIBAB SQUIRREL.

Type locality. Bright Angel Creek, Kaibab Plateau, Grand Cañon of the Colorado, Arizona.

Genl. Char. Similar to *S. aberti*, but black beneath, and tail all white.

Color. Entire upper parts grizzled gray; dorsal region ferruginous; lower sides, upper part of arms, and thighs black; under parts on middle of body black; tail with base gray; rest white, with a median buffy gray streak and black subterminal bands above; ears in summer blackish; gray with black tips in winter; hands mixed gray and black; hind feet gray in summer, black in winter.

136. *Sciurus durangi* (Thomas).

Sciurus aberti durangi Thomas, Ann. Mag. Nat. Hist., 6th Ser., xi, 1893, p. 49. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 112, fig. 18. Zoöl. Ser.

Sciurus aberti Thomas, Proc. Zoöl. Soc., 1882, p. 372. (nec Woodhouse.)

DURANGO SQUIRREL.

Type locality. Cuidad Ranch (100 miles west of Durango City), State of Durango, Mexico.

Geogr. Distr. Sierra Madre, west parts of States of Durango and Chihuahua, Mexico.

a. *Sciurus durangi phæurus* Allen.

Sciurus durangi phæurus Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1904, p. 205. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 11, 1904, Appendix, p. 741. Zoöl. Ser.

LA CIENAGA SQUIRREL.

Type locality. La Cienaga, State of Durango, Mexico.

Geogr. Distr. State of Durango, Mexico. Limits of range unknown.

137. *Sciurus barberi* (Allen).

Sciurus aberti barberi Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1904, p. 207.

Sciurus barberi Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 11, 1904, Appendix, p. 741. Zoöl. Ser.

BARBER'S SQUIRREL.

Type locality. Colonia Garcia, State of Chihuahua, Mexico.

Geogr. Distr. State of Chihuahua, Mexico. Limits of range unknown.

Subgenus G. *Echinosciurus*.

138. *Sciurus aureigaster* F. Cuvier.

Sciurus aureigaster F. Cuv., Hist. Nat. Mamm., VI, 1829, Livr.

LIX, pl. Text. Gray, Ann. Mag. Nat. Hist., 3d Ser., XX, 1867, p. 423.

Sciurus niger Erxl., Syst. Regn. Anim., 1777, p. 417. (nec Linn.)

Sciurus variegatus Desm., Nouv. Dict. Hist. Nat., X, 1817, pp. 103-104. (nec Erxleben, 1777.)

Sciurus rufiventer Licht., Abh. K. Ak. Wiss. Berl., (1827), 1830, p. 116.

Sciurus leucogaster F. Cuv., Suppl. Hist. Nat. Buffon, I, 1831, pp. 300-301.

Sciurus mustelinus Aud. & Bach., Proc. Acad. Nat. Scien. Phil., 1841, p. 100.

Sciurus ferruginiventris! Aud. & Bach., Proc. Acad. Nat. Scien. Phil., 1841, p. 101.

Sciurus aurogaster! Aud. & Bach., Quad. N. Am., III, 1851, p. 344.

Sciurus hypoxanthus I. Geoff., Voy. de la Vénus, Zool., 1855, p. 158.

Sciurus hypopyrrhus Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1891, p. 222. (nec Wagl.)

Sciurus leucops Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1897, p. 198.

Sciurus aureigaster Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 115, figs. 19, XXIX. Zool. Ser.

GOLDEN-BELLIED SQUIRREL. *Ardilla* in Mexico.

Type locality. Eastern Mexico.

Geogr. Distr. Eastern Mexico from southern part of State of Tamaulipas to northern side of Isthmus of Tehuantepec, through State of Vera Cruz, eastern San Luis Potosi, Queretaro, Puebla, and northeastern Hidalgo and northern Oaxaca, up to 4,000 and 8,000 feet.

a. *Sciurus aureigaster hypopyrrhus* (Wagler).

Sciurus hypopyrrhus Wagl., Isis, 1831, p. 510.

Macroxus morio Gray, Ann. Mag. Nat. Hist., 3d Ser., XX, 1867, p. 424.

Macroxus maurus Gray, Ann. Mag. Nat. Hist., 3d Ser., xx, 1867, p. 425.

Sciurus variegatus Sumichrast, La Naturaleza, v, 1882, p. 324.

Sciurus rufiventris Rovirosa, La Naturaleza, vii, 1887, p. 360.

Sciurus aureogaster Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1890, p. 181. (nec Cuvier.)

Sciurus aureigaster hypopyrrhus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 116. Zoöl. Ser.

FIRE-BELLIED SQUIRREL.

Type locality. State of Vera Cruz, Mexico.

Geogr. Distr. Southern part of the State of Vera Cruz and States of Oaxaca, Tabasco, and Chiapas, Mexico.

b. *Sciurus aureigaster frumentor* Nelson.

Sciurus aureogaster frumentor Nelson, Proc. Biol. Soc. Wash., xii, 1898, p. 154.

Sciurus aureigaster frumentor Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 117. Zoöl. Ser.

PEROTE SQUIRREL.

Type locality. Las Vigas, State of Vera Cruz, Mexico.

Geogr. Distr. "East and north base of the Cape de Perote, and eastern slope of the Cordillera near Las Vigas, State of Vera Cruz, Mexico, in pine and oak forests, at 6,000-8,000 feet."

139. *Sciurus poliopus* (Fitzinger).

Sciurus albipes Wagn., Abhandl. Math.-Phys. Cl. K. Bayer Akad. Wiss. München, ii, 1837, pp. 501-506. (nec Kerr.)

Sciurus varius Wagn., Schreb. Säugth., Suppl., iii, 1843, p. 168.

Sciurus variegatus poliopus Fitz., Sitzung. K. Akad. Wiss. Wien, i, 1867, p. 478.

Sciurus variegatus rufipes Fitz., Sitzung. K. Akad. Wiss. Wien, i, 1867, p. 478.

Macroxus leucops Gray, Ann. Mag. Nat. Hist., 3d Ser., xx, 1867, p. 427.

Sciurus variegatus Alston, Proc. Zoöl. Soc., 1878, p. 660.

Sciurus wagneri Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1898, p. 453.

Sciurus poliopus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 117. Zoöl. Ser.

OAXACA SQUIRREL.

Type locality. Cerro San Felipe, State of Oaxaca, Mexico.

Geogr. Distr. Mountains about Valley of Oaxaca, except those on the west, at 7,500-11,000 feet altitude.

a. *Sciurus poliopus hernandezi* (Nelson).

Sciurus albipes hernandezi Nelson, Science, VIII, 1898, p. 783.

Sciurus albipes quercinus Nelson, Proc. Biol. Soc. Wash., XII, 1898, p. 150. (nec Erxl.)

Sciurus wagneri quercinus Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1898, p. 453.

Sciurus poliopus hernandezi Nelson, Proc. Wash. Acad. Scien., I, 1899, p. 48. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 118. Zool. Ser.

OAK WOODS SQUIRREL.

Type locality. Mountains 15 miles west of the city of Oaxaca, State of Oaxaca, Mexico.

Geogr. Distr. Mountains west of Oaxaca Valley into southern part of the State of Puebla and southeastern portion of State of Guerrero, Mexico. Altitude, 8,000-9,000 feet.

b. *Sciurus poliopus peregrinator* Nelson.

Sciurus poliopus peregrinator Nelson, Proc. Biol. Soc. Wash., XVII, 1904, p. 149.

PUEBLA SQUIRREL.

Type locality. Piaxtla, State of Puebla.

Geogr. Distr. Southern part of State of Puebla, northwestern portion of State of Oaxaca and adjacent parts of State of Guerrero.

Genl. Char. Like *S. p. hernandezi*, but brighter.

Color. Top of nose and fore part of crown iron gray; back part of crown and nape dark ochraceous buffy and black; rest of upper parts light iron gray mixed with ochraceous; sides of body paler; rump patch mixed with black; top of feet white; under parts of body cream buff; tail above black washed with white, bottom of hairs rusty ochraceous, beneath ochraceous bordered with black and edged with white.

Measurements. Total length, 535; tail vertebrae, 273; hind foot, 69.

c. *Sciurus poliopus nemoralis* (Nelson).

Sciurus albipes nemoralis Nelson, Proc. Biol. Soc. Wash., XII, 1898, p. 151.

Sciurus wagneri nemoralis Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1898, p. 454.

Sciurus poliopus nemoralis Nelson, Proc. Wash. Acad. Scien., I, 1899, p. 50. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 118. Zool. Ser.

MICHOACAN SQUIRREL.

Type locality. Patzcuaro, State of Michoacan, Mexico.

Geogr. Distr. Volcano of Toluca, State of Mexico, to Nahuatzin, State of Michoacan, Mexico. Altitude 7,000-12,000 feet.

d. Sciurus poliopus senex Nelson.

Sciurus poliopus senex Nelson, Proc. Biol. Soc. Wash., xvii, 1904, p. 148.

LA SALADA SQUIRREL.

Type locality. La Salada, 40 miles south of Uruapan, southern part of the State of Michoacan, Mexico.

Geogr. Distr. Valley of the Balsas River, below 4,000 feet; in central and southern part of the State of Michoacan and adjacent parts of the State of Guerrero, Mexico.

Genl. Char. Like *S. p. nemoralis*, but paler; nuchal patch more yellowish; rump patch less distinct; rostrum heavier; braincase narrower, more constricted posteriorly.

Color. Top of head iron gray; nape ochraceous and black mixed; rest of back pale grizzled gray; sides of body paler, under parts white; tail above black washed with white, beneath dull gray bordered with blackish, and edged with white; top of feet white.

Measurements. Total length, 543; tail vertebræ, 275; hind foot, 69.

e. Sciurus poliopus cervicalis (Allen).

Sciurus leucops Allen, Mon. N. Am. Rod., 1877, p. 753. (nec Gray.)

Sciurus variegatus Alston, Biol. Centr. Amer., Mamm., 1, 1880, p. 127. (nec Erxl.)

Sciurus aureogaster leucops Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1889, p. 166.

Sciurus cervicalis Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1890, p. 183.

Sciurus wagneri cervicalis Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1898, p. 454.

Sciurus poliopus cervicalis Nelson, Proc. Wash. Acad. Scien., 1, 1899, p. 51. Elliot, Mamm. Middle Amer. & W. Indies. F. C. M. Pub., iv Pt. 1, 1904, p. 119. Zool. Ser.

COLIMA MOUNTAIN SQUIRREL.

Type locality. Hacienda San Marcos, Tonila, State of Jalisco, at east base of Sierra Nevada de Colima, Mexico.

Geogr. Distr. From the Sierra Nevada de Colima, along the high mountains north to Ameca, State of Jalisco, and east into the western part of the State of Michoacan, Mexico. Altitude, 6,000 to 12,000 feet, and down occasionally to 4,000 feet.

f. Sciurus poliopus colimensis (Nelson).

Sciurus albipes colimensis Nelson, Proc. Biol. Soc. Wash., xii, 1898, p. 152.

Sciurus poliopus colimensis Nelson, Proc. Wash. Acad. Scien.,
1, 1899, p. 52. Elliot, Mamm. Middle Amer. & W. Indies,
F. C. M. Pub., IV, Pt. 1, 1904, p. 119. Zoöl. Ser.

Sciurus leucops Allen, N. Am. Rod., 1877, p. 753. (Part.)

Sciurus albipes colimensis Nelson, Proc. Biol. Soc. Wash., XII,
1898, p. 152.

Sciurus wagneri colimensis Allen, Bull. Am. Mus. Nat. Hist.,
1898, p. 454.

COLIMA SQUIRREL.

Type locality. Hacienda Magdalena, State of Colima, Mexico.

Geogr. Distr. Coast region of the State of Colima, Mexico.

g. *Sciurus poliopus effugius* (Nelson).

Sciurus albipes effugius Nelson, Proc. Biol. Soc. Wash., XII, 1898,
p. 152.

Sciurus poliopus effugius Nelson, Proc. Wash. Acad. Scien., 1,
1899, p. 54. Elliot, Mamm. Middle Amer. & W. Indies,
F. C. M. Pub., IV, Pt. 1, 1904, p. 120. Zoöl. Ser.

Sciurus wagneri effugius Allen, Bull. Am. Mus. Nat. Hist. N. Y.,
1898, p. 454.

GUERRERO SQUIRREL.

Type locality. Mountains west of Chilpancingo, State of Guerrero, Mexico. Altitude, 7,500-9,500 feet.

Geogr. Distr. Forests of Cordillera, State of Guerrero, Mexico.

140. *Sciurus nelsoni* Merriam.

Sciurus nelsoni Merr., Proc. Biol. Soc. Wash., VIII, 1893, p. 144.

Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub.,
IV, Pt. 1, 1904, p. 120. Zoöl. Ser.

NELSON'S SQUIRREL.

Type locality. Huitzilac, State of Morelos, Mexico.

Geogr. Distr. Mountains south and west of the Valley of Mexico
and south of Valley of Toluca, in the States of Mexico and
Morelos.

a. *Sciurus nelsoni hirtus* Nelson.

Sciurus nelsoni hirtus Nelson, Proc. Biol. Soc. Wash., XII, 1898,

p. 153. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M.
Pub., IV, Pt. 1, 1904, p. 121. Zoöl. Ser.

MOUNTAIN SQUIRREL.

Type locality. Tochmilco, State of Puebla, Mexico.

Geogr. Distr. Mounts Popocatepetl and Iztaccihuatl, Mexico.
Altitude, 8,000-12,000 feet, in States of Mexico, Puebla and Morelos.

141. *Sciurus colliæi* Richardson.

Sciurus colliæi Rich., Voy. Blossom, Zoöl., 1839, p. 8. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 121. Zoöl. Ser.

Sciurus hypopyrrhus Alston, Proc. Zoöl. Soc., 1878, p. 662. (Part.)
COLLIE'S SQUIRREL.

Type locality. San Blas, Territorio de Tepic, Mexico.

Geogr. Distr. From northern border of Territorio de Tepic to Bay of Banderas, State of Jalisco, Mexico, below 2,500 feet.

a. *Sciurus colliæi nuchalis* Nelson.

Sciurus colliæi nuchalis Nelson, Proc. Wash. Acad. Scien., 1, 1899, p. 59. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 122. Zoöl. Ser.

MANZANILLO SQUIRREL.

Type locality. Manzanillo, State of Colima, Mexico.

Geogr. Distr. From State of Michoacan to Bay of Banderas, State of Jalisco, and inland to mountains near San Sebastian in the same state, below 3,000 feet.

142. *Sciurus sinaloensis* Nelson.

Sciurus sinaloensis Nelson, Proc. Wash. Acad. Scien., 1, 1899, p. 60. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 122. Zoöl. Ser.

SINALOA SQUIRREL.

Type locality. Mazatlan, State of Sinaloa, Mexico.

Geogr. Distr. Southern and central parts of the State of Sinaloa, Mexico, below 2,500 feet.

143. *Sciurus truei* Nelson.

Sciurus truei Nelson, Proc. Wash. Acad. Scien., 1, 1899, p. 61.

Sciurus truei Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 123. Zoöl. Ser.

TRUE'S SQUIRREL.

Type locality. Camoa, Rio Mayo, State of Sonora, Mexico.

Geogr. Distr. Southwestern part of the State of Sonora, and northern portion of the State of Sinaloa, Mexico.

144. *Sciurus socialis* Wagner.

Sciurus socialis Wagn., Abh. Math.-Phys. Cl. K. Bayer Akad. Wiss. München, 11, 1837, pp. 501-7, pl. v. Elliot, Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 123. Zoöl. Ser.

Sciurus aureigaster Allen, Mon. N. Am. Rod., 1877, p. 750. (Part.)

Sciurus leucops Allen, Mon. N. Am. Rod., 1877, p. 753. (Part.)

Sciurus variegatus Alston, Proc. Zool. Soc., 1878, p. 660. (Part.)

TEHUANTEPEC SQUIRREL.

Type locality. Near Tehuantepec City, State of Oaxaca, Mexico.

Geogr. Distr. Puerto Angel, State of Oaxaca, South to Tonalá and Tuxtla, State of Chiapas, Mexico, and up the river valley to Newton, Guatemala, below 3,000 feet.

a. *Sciurus socialis cocos* Nelson.

Sciurus socialis cocos Nelson, Proc. Biol. Soc. Wash., XII, 1898, p. 155. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 124.

Sciurus aureogaster I. Geoff., Voy. Vénus, Zool., Atlas, 1846, pl. 10. (nec Cuvier.)

ACAPULCO SQUIRREL.

Type locality. Acapulco, State of Guerrero, Mexico.

Geogr. Distr. From Acapulco, State of Guerrero, to Jamiltepec, State of Oaxaca, Mexico, below 1,500 feet.

145. *Sciurus griseiflavus* (Gray).

Macroxus griseoflavus Gray, Ann. Mag. Nat. Hist., 3d Ser., XX, 1867, p. 427.

Sciurus ludovicianus Tomes, Proc. Zool. Soc., 1861, p. 281. (nec Custis.)

Sciurus aureigaster Allen, Mon. N. Am. Rod., 1877, p. 750. (Part.)

Sciurus affinis Alston, Proc. Zool. Soc., 1878, p. 660.

Sciurus griseiflavus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 124. Zool. Ser.

GUATEMALA SQUIRREL.

Type locality. Guatemala.

Geogr. Distr. Southeastern part of the State of Chiapas, Mexico, and northwestern Guatemala. Altitude, 7,000-10,000 feet.

a. *Sciurus griseiflavus chiapensis* Nelson.

Sciurus griseoflavus chiapensis Nelson, Proc. Wash. Acad. Scien., I, 1899, p. 69.

Sciurus griseiflavus chiapensis Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 125. Zool. Ser.

CHIAPAS SQUIRREL.

Type locality. San Cristobal, State of Chiapas, Mexico.

Geogr. Distr. Mountains of central portion of State of Chiapas, Mexico. Altitude, 7,500-9,500 feet.

146. *Sciurus yucatanensis* (Allen).

Sciurus carolinensis yucatanensis Allen, Mon. N. Am. Rod., 1877, p. 705.

Sciurus yucatanensis Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1897, p. 5. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 125. Zoöl. Ser.

Sciurus carolinensis Alston, Proc. Zoöl. Soc., 1878, p. 658.

YUCATAN SQUIRREL.

Type locality. Merida, Yucatan, Mexico.

Geogr. Distr. Peninsula of Yucatan, Mexico.

a. Sciurus yucatanensis baliolus Nelson.

Sciurus yucatanensis baliolus Nelson, Proc. Biol. Soc. Wash., XIV, 1901, p. 131. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 126. Zoöl. Ser.

SWARTHY SQUIRREL.

Type locality. Apazote, State of Campeche, Mexico.

Geogr. Distr. Southern part of the State of Campeche, and eastern part of the State of Tabasco, Mexico.

147. *Sciurus thomasi* Nelson.

Sciurus thomasi Nelson, Proc. Wash. Acad. Scien., 1, 1899, p. 71. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 126. Zoöl. Ser.

Sciurus boothiae Allen, Mon. N. Am. Rod., 1877, pp. 741-746. (Part. Costa Rica.) (nec Gray.)

Sciurus hypopyrrhus Alston, Proc. Zoöl. Soc., 1878, pp. 662-664. (Part. Costa Rica.)

THOMAS' SQUIRREL.

Type locality. Talamanca, Costa Rica.

Geogr. Distr. "Humid tropical forests of eastern Costa Rica."

148. *Sciurus adolphii* (Lesson).

Macroxus adolpheii Less., Nouv. Tabl. Règn. Anim., Mamm., 1842, p. 112.

Sciurus boothiae Allen, Mon. N. Am. Rod., 1879, p. 741. (Part. Nicaragua.)

Sciurus hypopyrrhus Alston, Biol. Centr. Amer., 1, Mamm., 1880, p. 128.

Sciurus adolpheii Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 126. Zoöl. Ser.

NICARAGUA SQUIRREL.

Type locality. Realejo, Nicaragua.

Geogr. Distr. Lowlands on the west coast of Nicaragua.

a. Sciurus adolphei dorsalis (Gray).

Sciurus dorsalis Gray, Proc. Zool. Soc., 1848, p. 138.

Sciurus rigidus Peters, Monatsb. K. Preuss. Akad. Wiss. Berlin, 1863, p. 652.

Sciurus intermedius Gray, Ann. Mag. Nat. Hist., 3d Ser., xx, 1867, p. 421.

Sciurus nicoyana Gray, Ann. Mag. Nat. Hist., 3d Ser., xx, 1867, p. 423.

Sciurus boothiæ Allen, Mon. N. Am. Rod., 1877, p. 741. (Part. West Costa Rica specimens.)

Sciurus hypopyrrhus Allen, Mon. N. Am. Rod., 1877, p. 746. (Part. West Costa Rica specimens.)

Sciurus adolphei dorsalis Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 127. Zool. Ser.

BANDED-BACK SQUIRREL.

Type locality. Liberia, Costa Rica, nec Caracas, Venezuela.

Geogr. Distr. Western Costa Rica, Alajuela to Liberia, Peninsula of Nicoya, and possibly adjacent parts of Nicaragua.

149. *Sciurus melania* (Gray).

Macroxus melania Gray, Ann. Mag. Nat. Hist., 3d Ser., xx, 1867, p. 425.

Sciurus melania Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 128. Zool. Ser.

GRAY'S BLACK SQUIRREL.

Type locality. Point Burica, Costa Rica.

Geogr. Distr. Costa Rica, Chiriqui, Panama. Altitude, 2,000 feet.

150. *Sciurus boothiæ* Gray.

Sciurus boothiæ Gray, List Spec. Mamm. Brit. Mus., 1843, p. 139.

Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 128. Zool. Ser.

Sciurus richardsoni Gray, Ann. Mag. Nat. Hist., 1st Ser., x, 1842, p. 264. (nec Bachman.)

Sciurus fuscovariegatus Schinz., Syst. Mamm., II, 1845, p. 15.

Sciurus hypopyrrhus Alston, Proc. Zool. Soc., 1878, p. 662. (Part. Honduras specimens.)

HONDURAS SQUIRREL.

Type locality. Honduras, San Pedro Sula?

Geogr. Distr. Humid coast forests of northern Honduras and border of Guatemala.

a. Sciurus boothiae belti Nelson.

Sciurus boothiae belti Nelson, Proc. Wash. Acad. Scien., 1, 1899, p. 78. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 128. Zoöl. Ser.

ESCONDIDO RIVER SQUIRREL.

Type locality. Escondido River, fifty miles above Bluefields, Nicaragua.

Geogr. Distr. East coast region of Nicaragua, and north to Segovia River, Honduras.

151. *Sciurus variegatoides* Ogilby.

Sciurus variegatoides Ogilby, Proc. Zoöl. Soc., 1839, p. 117. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 129. Zoöl. Ser.

Sciurus pyladei Less., Nouv. Tabl. Règn. Anim., Mamm., 1842, p. 112.

Sciurus griseocaudatus Gray, Voy. 'Sulphur,' Mamm., II, 1843, p. 34, pl. 13, fig. 2, pl. 18, figs. 7-12.

Sciurus colliæi Gray, Ann. Mag. Nat. Hist., 3d Ser., XX, 1867, p. 746. (Part. var. 1.)

Sciurus hypopyrrhus Allen, Mon. N. Am. Rod., 1877, pp. 746-750. (Part. Salvador specimens.)

VARIEGATED SQUIRREL.

Type locality. Salvador, west coast of Central America.

Geogr. Distr. Tropical forests of Salvador, west coast of Central America.

152. *Sciurus managuensis* (Nelson).

Sciurus boothiae managuensis Nelson, Proc. Biol. Soc. Wash., XII, 1898, p. 150.

Sciurus managuensis Nelson, Proc. Wash. Acad. Scien., 1, 1899, p. 81. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 129. Zoöl. Ser.

RIO MANAGUA SQUIRREL.

Type locality. Managua River, Guatemala.

Geogr. Distr. Northern Guatemala, along the Managua River.

153. *Sciurus goldmani* Nelson.

Sciurus goldmani Nelson, Proc. Biol. Soc. Wash., XII, 1898, p. 149. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 130. Zoöl. Ser.

GOLDMAN'S SQUIRREL.

Type locality. Huehuetan, State of Chiapas, Mexico.

Geogr. Distr. Southeastern part of the State of Chiapas, Mexico, into Guatemala.

Subgenus H. *Hesperosciurus*.**154. *Sciurus griseus* Ord.**

Sciurus griseus Ord, Guth. Geogr., 2d Am. ed., II, 1815, p. 292, mss. annot. author's copy. *Id.* Journ. Phys., LXXXVII, 1818, p. 150. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 55, fig. 12. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 130, fig. 20. Zoöl. Ser.

Sciurus fessor Peale, U. S. Expl. Exped., Mamm. and Birds, 1848, p. 55.

Sciurus heermanni Le Conte, Proc. Acad. Nat. Scien. Phil., 1852, p. 149.

Sciurus leporinus Hensh., Ann. Rep. Engin., 1876, p. 25.

Sciurus fessor anthonyi Mearns, Proc. U. S. Nat. Mus., XX, 1898, p. 501.

Sciurus griseus anthonyi Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 57.

CALIFORNIA GRAY SQUIRREL.

Type locality. The Dalles, Columbia River, Wasco County, Oregon.

Geogr. Distr. Northern California, Oregon, and Washington.

a. Sciurus griseus nigripes (Bryant.)

Sciurus fessor nigripes Bryant, Proc. Calif. Acad. Scien., 1889, p. 25.

Sciurus griseus nigripes Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 56. Zoöl. Ser.

BLACKFOOTED GRAY SQUIRREL.

Type locality. San Mateo County, California, south of San Francisco.

Geogr. Distr. Red Wood Belt, coast region of California, south of San Francisco.

Subgenus I. *Neosciurus*.**155. *Sciurus carolinensis* Gmelin.**

Sciurus carolinensis Gmel., Syst. Nat., I, 1788, p. 148. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 53, fig. 11. Zoöl. Ser.

CAROLINA GRAY SQUIRREL.

Type locality. Carolina.

Geogr. Distr. Eastern United States from Southern New York to northern Florida, and westward south of Pennsylvania to Missouri, Indian Territory, and to edge of the plains.

a. *Sciurus carolinensis leucotis* (Gapper).

Sciurus leucotis Gapper, Zoöl. Journ., v, 1830, p. 206.

Sciurus cinereus Schreb., Säugth, iv, 1792, p. 766, pl. ccxcii.
(nec Linn.)

?**Sciurus pennsylvanicus* Ord, Guth. Geogr., 2d Am. ed., II, 1815,
p. 292.

Sciurus hiemalis Ord, Guth. Geogr., 2d Am. ed., II, 1815, p. 292.

Sciurus niger Godman, Am. Nat. Hist., II, 1826, p. 133. (nec
Linn.)

Sciurus vulpinus DeKay, Zoöl. N. Y., I, 1842, p. 59. (nec Gmel.,
nec Schreb.)

Sciurus migratorius Aud. & Bachm., Quad. N. Am., I, 1849, p. 265,
pl. xxxv.

Sciurus carolinensis leucotis Elliot, Syn. N. Am. Mamm., F. C. M.
Pub., II, 1901, p. 53. Zoöl. Ser.

WHITE-EARED GRAY SQUIRREL.

Type locality. Region between York and Lake Simcoe, Ontario,
Canada.

Geogr. Distr. Alleghany Mountains in Pennsylvania north through
New York and New England to southern New Brunswick and southern
Canada, west to Minnesota.

b. *Sciurus carolinensis hypophæus* Merriam.

Sciurus carolinensis hypophæus Merr., Scien., VIII, 1886, p. 351.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 54.
Zoöl. Ser.

MINNESOTA GRAY SQUIRREL.

Type locality. Elk River, Sherburne County, Minnesota.

Geogr. Distr. Minnesota forest belt. Range unknown.

c. *Sciurus carolinensis fuliginosus* (Bachman).

Sciurus fuliginosus Bachm., Proc. Zoöl. Soc., 1838, p. 97.

Sciurus carolinensis fuliginosus Elliot, Syn. N. Am. Mamm.,
F. C. M. Pub., II, 1901, p. 55. Zoöl. Ser.

DUSKY GRAY SQUIRREL.

Type locality. Near New Orleans, Louisiana.

Geogr. Distr. Coast region of Louisiana.

**Sciurus pennsylvanicus* Ord, if it could be proved to have been applied
to the northern form of the gray squirrel would have fifteen years precedence
over *leucotis* Gapper, and that name would necessarily become a synonym.
But as Ord's name is without description, and therefore a *nomen nudum*, it
cannot be determined whether he intended to apply it to the southern or
northern form of gray squirrel.

d. Sciurus carolinensis extimus Bangs. .

Sciurus carolinensis extimus Bangs, Proc. Biol. Soc. Wash., x,
1896, p. 158. Elliot, Syn. N. Am. Mamm., F. C. M. Pub.,
II, 1901, p. 55. Zoöl. Ser.

FLORIDA GRAY SQUIRREL.

Type locality. Miami, Dade County, Florida.

Geogr. Distr. South Florida.

Subgenus J. *Tamiasciurus*.**156. *Sciurus hudsonius* (Erxleben).**

Sciurus vulgaris hudsonicus Exrl., Syst. Reg. Anim., Mamm.,
1777, p. 416.

Sciurus hudsonius Elliot, Syn. N. Am. Mamm., F. C. M. Pub.,
II, 1901, p. 60, fig. 15. Zoöl. Ser.

HUDSON BAY CHICKAREE.

Type locality. Hudson Strait.

Geogr. Distr. Boreal North America, Labrador, west to Rocky Mountains and Alaska.

a. Sciurus hudsonius gymnicus (Bangs).

Sciurus hudsonicus gymnicus Bangs, Proc. N. Eng. Zoöl. Club,
I, 1899, p. 28.

Sciurus hudsonius gymnicus Elliot, Syn. N. Am. Mamm., F. C. M.
Pub., II, 1901, p. 61. Zoöl. Ser.

EASTERN CHICKAREE.

Type locality. Greenville, Maine.

Geogr. Distr. Eastern North America, south of Labrador to northern New York, through Canada to northern Michigan and northern Minnesota.

b. Sciurus hudsonius loquax (Bangs).

Sciurus hudsonicus loquax Bangs, Proc. Biol. Soc. Wash., x,
1896, p. 161.

Sciurus carolinensis Ord, Guth. Geogr., 2d Am. ed., II, 1815,
p. 292. (nec Gmel.)

Sciurus hudsonius loquax Elliot, Syn. N. Am. Mamm., F. C. M.
Pub., II, 1901, p. 62. Zoöl. Ser.

NOISY CHICKAREE.

Type locality. Liberty Hill, New London County, Connecticut.

Geogr. Distr. Ontario to North Carolina in the east, and west to Minnesota.

c. *Sciurus hudsonius minnesota* (Allen).

Sciurus hudsonicus minnesota Allen, Amer. Nat., xxxiii, 1899, p. 640.

Sciurus hudsonius minnesota Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 62. Zool. Ser.

MINNESOTA CHICKAREE.

Type locality. Fort Snelling, Hennepin County, Minnesota.

Geogr. Distr. Southern Minnesota and Wisconsin, Iowa? eastward to northern Indiana.

d. *Sciurus hudsonius dakotensis* (Allen).

Sciurus hudsonicus dakotensis Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1894, p. 325.

Sciurus hudsonius dakotensis Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 62. Zool. Ser.

SOUTH DAKOTA CHICKAREE.

Type locality. Squaw Creek, Black Hills, South Dakota.

Geogr. Distr. Black Hills of South Dakota and adjoining parts of Wyoming.

e. *Sciurus hudsonius baileyi* (Allen).

Sciurus hudsonicus baileyi Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1898, p. 261.

Sciurus hudsonius baileyi Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 63. Zool. Ser.

BAILEY'S CHICKAREE.

Type locality. Big Horn Mountains, Wyoming. Altitude, 8,400 feet.

Geogr. Distr. Mountain ranges of Central Wyoming and eastern Montana, north into Alberta, in eastern foothills of the Rocky Mountains.

f. *Sciurus hudsonius ventorum* (Allen).

Sciurus hudsonicus ventorum Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1898, p. 263.

Sciurus hudsonius ventorum Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 63. Zool. Ser.

WIND RIVER MOUNTAINS CHICKAREE.

Type locality. South Pass City, Fremont County, Wyoming.

Geogr. Distr. Wind River Mountains north to Mystic Lake, west to head of Snake River in Idaho, south to Wahsatch Range, north-eastern Utah.

g. *Sciurus hudsonius richardsoni* (Bachman).

Sciurus richardsonii Bach., Proc. Zool. Soc., 1838, p. 100.

Sciurus hudsonius richardsoni Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 64. Zool. Ser.

RICHARDSON'S CHICKAREE.

Type locality. Head of Big Lost River, Idaho.

Geogr. Distr. Western border of northern Montana, central and northern Idaho, northeastern Washington and Oregon, northward into British Columbia.

h. *Sciurus hudsonius streatori* (Allen).

Sciurus hudsonicus streatori Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1898, p. 267.

Sciurus hudsonius streatori Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 64. Zool. Ser.

STREATOR'S CHICKAREE.

Type locality. Ducks, British Columbia.

Geogr. Distr. Northern Washington from the Columbia River to central British Columbia.

i. *Sciurus hudsonius petulans* (Osgood).

Sciurus hudsonicus petulans Osgood, N. Am. Faun., No. 19, 1900, p. 27.

Sciurus hudsonius petulans Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, Append., p. 426. Zool. Ser.

SCOLDING CHICKAREE.

Type locality. Glacier, White Pass, Alaska. Altitude, 1,870 feet.

Geogr. Distr. Alaska, range unknown.

j. *Sciurus hudsonius vancouverensis* Allen.

Sciurus hudsonius vancouverensis Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1890, p. 165.

Sciurus hudsonius vancouverensis Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 64. Zool. Ser.

VANCOUVER ISLAND CHICKAREE.

Type locality. Duncan Station, Vancouver Island, British Columbia, Canada.

Geogr. Distr. Vancouver Island, British Columbia, north to Sitka, Alaska.

157. *Sciurus douglasii* Bachman.

Sciurus douglasii Bachm., Proc. Zool. Soc., 1838, p. 99. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 65. Zool. Ser.

Sciurus belcheri Gray, Ann. Mag. Nat. Hist., 1st Ser., 1842, p. 263.

Sciurus suckleyi Baird, Proc. Acad. Nat. Scien. Phil., 1855, p. 333.

DOUGLAS' CHICKAREE.

Type locality. Columbia River, Oregon.

Geogr. Distr. Coast region of Oregon and Washington from Cape Blanco to Puget Sound.

a. *Sciurus douglasi mollipilosus* (Audubon and Bachman).

Sciurus mollipilosus Aud. & Bachm., Proc. Acad. Nat. Scien. Phil., 1841, p. 42.

Sciurus douglasi mollipilosus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 65. Zoöl. Ser.

SOFT-FURRED CHICKAREE.

Type locality. Coast of northern California.

Geogr. Distr. Coast of northern California from Sonoma County into Curry County, Oregon.

b. *Sciurus douglasi cascadiensis* Allen.

Sciurus douglasii cascadiensis Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1898, p. 277. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 66. Zoöl. Ser.

CASCADE MOUNTAINS CHICKAREE.

Type locality. Mt. Hood, Oregon.

Geogr. Distr. Cascade Mountains of Oregon and Washington, north into British Columbia. Also coast region from Fraser River to River's Inlet north of Vancouver Island.

c. *Sciurus douglasi albolimbatus* Allen.

Sciurus hudsonius californicus Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1890, p. 165.

Sciurus douglasi albolimbatus Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1898, p. 453. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 66. Zoöl. Ser.

CALIFORNIA CHICKAREE.

Type locality. Blue Cañon, Placer County, California.

Geogr. Distr. Sierra Nevada region of northern California and Oregon, east of the Cascades to the Maury Mountains and Strawberry Butte.

d. *Sciurus douglasi mearnsi* (Townsend).

Sciurus hudsonius mearnsi Town., Proc. Biol. Soc. Wash., XI, 1897, p. 146.

Sciurus douglasi mearnsi Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 133, fig. 21. Zoöl. Ser.

MEARNS' CHICKAREE.

Type locality. San Pedro Martir Mountains, Lower California. Altitude, 7,000 feet.

Geogr. Distr. Forests of San Pedro Martir Mountains, Lower California. Altitude, 7,000 feet.

158. *Sciurus fremonti* Audubon and Bachman.

Sciurus fremonti Aud. & Bachm., Quad. N. Am., III, 1853, p. 237.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 66.

Zoöl. Ser.

FREMONT'S CHICKAREE.

Type locality. South Pass, Rocky Mountains, Park Region, Central Colorado.

Geogr. Distr. Mountains of Colorado and the Uintah Mountains, Utah, to the southern border of Wyoming.

a. Sciurus fremonti mogollonensis (Mearns).

Sciurus hudsonius mogollonensis Mearns, Bull. Am. Mus. Nat. Hist.

N. Y., 1887-90, p. 277.

Sciurus fremonti mogollonensis Elliot, Syn. N. Am. Mamm., F. C.

M. Pub., II, 1901, p. 67. Zoöl. Ser.

ARIZONA CHICKAREE.

Type locality. Quaking Asp Settlement, summit of Mogollon Mountains, Arizona.

Geogr. Distr. High Mountains and plateaus of central Arizona.

b. Sciurus fremonti neomexicanus Allen.

Sciurus fremonti neomexicanus Allen, Bull. Am. Mus. Nat. Hist.

N. Y., 1898, p. 291. Elliot, Syn. N. Am. Mamm., F. C. M.

Pub., II, 1901, p. 67. Zoöl. Ser.

TAOS CHICKAREE.

Type locality. Rayado Cañon, Colfax County, New Mexico.

Geogr. Distr. Taos Range, New Mexico.

c. Sciurus fremonti lychnuchus Stone and Rehn.

Sciurus fremonti lychnuchus Stone and Rehn, Proc. Acad. Nat.

Scien. Phil., 1903, p. 18.

RUIDOSO CHICKAREE.

Type locality. Forks of the Ruidoso (White Mountains), Lincoln County, New Mexico.

Geogr. Distr. New Mexico, range unknown.

Genl. Char. Similar to *S. f. neomexicanus*, but larger and more rufous; nasals longer.

Color. Above dull ferrugineous, deepest on rump; sides pale rufescent with ochraceous tinge; lateral line narrow, indistinct; head washed with blackish; orbital ring white; ear ferrugineous, tips blackish; arms rufescent, lateral fringe of hands white; legs dull fer-

rugineous, rufescent on upper surface of feet; tail above ferrugineous, fringed with ochraceous, end of tail black with the tips of hairs ochraceous, beneath pale grayish ochraceous fringed with ochraceous; under parts with inner surface of limbs white.

Measurements. Total length, 350; tail vertebræ, 140; hind foot, 55. Skull: total length, 50; basilar length, 42; zygomatic width, 29; length of nasals, 16; width of nasals, 8.2.

d. Sciurus fremonti grahamensis (Allen).

Sciurus hudsonicus grahamensis Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1894, p. 350.

Sciurus fremonti grahamensis Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 67. Zoöl. Ser.

GRAHAM MOUNTAINS CHICKAREE.

Type locality. Graham Mountains, Graham County, Arizona.

Geogr. Distr. Graham Mountains, southeastern Arizona.

Genus 51. **Tamias.** Chipmunks.

Tamias Illig., Prodr. Syst. Mamm. et Aves, 1811, p. 83. Type *Sciurus striatus* Linnæus.

Tenotis Rafin., Am. Month. Mag., I, 1817, p. 362.

Tamia Less., Man. Mamm., 1827, p. 230.

Eutamias Trouess., Cat. Mamm. Viv. et Foss., Bull. Soc. Études Angers, x, 1880, pp. 86-87.

• Subgenus A. **Eutamias.**

159. **Tamias dorsalis** Baird.

Tamias dorsalis Baird, Proc. Acad. Nat. Scien. Phil., 1855, p. 332.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 68, fig. 16. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 135. Zoöl. Ser.

Tamias quadrivittatus pallidus Coues and Yarr., in Wheeler's Rep. Geogr. Expl. & Surv. West of 100th Merid., v, 1876, p. 118.

GILA CHIPMUNK.

Type locality. Fort Webster, Grant County, New Mexico.

Geogr. Distr. Northern Utah to northern Mexico; west to the Sierra Nevada, east to the Mimbres in New Mexico.

a. Tamias dorsalis utahensis (Merriam).

Eutamias dorsalis utahensis Merr., Proc. Biol. Soc. Wash., 1897, p. 210.

Tamias dorsalis utahensis Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 69. Zoöl. Ser.

UTAH CHIPMUNK.

Type locality. Ogden, Weber County, Utah.

Geogr. Distr. Northern Utah and northwestern Nevada through Arizona to New Mexico.

160. *Tamias canescens* (Allen).

Eutamias canescens Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1904, p. 208.

*Tamias canescens*¹ Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, Append., p. 741. Zoöl. Ser.

GUANACEVI CHIPMUNK.

Type locality. Guanacevi, State of Durango, Mexico. Altitude, 8,000 feet.

Geogr. Distr. State of Durango, Mexico, range unknown.

161. *Tamias palmeri* (Merriam).

Eutamias palmeri Merr., Proc. Biol. Soc. Wash., 1897, p. 208.

Tamias palmeri Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 69. Zoöl. Ser.

PALMER'S CHIPMUNK.

Type locality. Charleston Peak, Lincoln County, Nevada. Altitude, 8,000 feet.

Geogr. Distr. Restricted to type locality.

162. *Tamias townsendi* Bachman.

Tamias townsendii Bachm., Jour. Acad. Nat. Scien. Phil., 1839, p. 68. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 69. Zoöl. Ser.

Tamias cooperi Baird, Proc. Acad. Nat. Scien. Phil., 1855, p. 334.

TOWNSEND'S CHIPMUNK.

Type locality. Mouth of the Columbia River.

Geogr. Distr. Coast region of Oregon, Washington, and British Columbia, from the mouth of the Columbia River northward; east in northern Cascades to the head of Lake Chelan.

a. Tamias townsendi hindsi (Gray).

Tamias hindsi Gray, Ann. Mag. Nat. Hist. 1st Ser., X, 1842, p. 264.

Tamias townsendi hindsi Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 70. Zoöl. Ser.

HIND'S CHIPMUNK.

Type locality. Nicasio? California.

Geogr. Distr. Coast of California from San Francisco north.

b. Tamias townsendi pricei (Allen).

Tamias pricei Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1895, p. 333.

Tamias townsendi pricei Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 70. Zoöl. Ser.

PRICE'S CHIPMUNK.

Type locality. Portola, San Mateo County, California.

Geogr. Distr. Coast region of California, from mountains of San Diego and San Bernardino County north to San Francisco.

c. *Tamias townsendi ochrogenys* (Merriam).

Eutamias townsendii ochrogenys Merriam, Proc. Biol. Soc. Wash., XI, 1897, p. 206.

Tamias townsendi ochrogenys Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 71. Zoöl. Ser.

BUFF-CHEEKED CHIPMUNK.

Type locality. Mendocino, Mendocino County, California.

Geogr. Distr. Cazadero, north of San Francisco Bay, California, to the mouth of Rogue River, Oregon.

d. *Tamias townsendi littoralis* Elliot.

Tamias townsendi littoralis Elliot, Pub. Field Columb. Mus., III, 1903, p. 153. Zoöl. Ser.

COAST CHIPMUNK.

Type locality. Marshfield, Coos County, Oregon.

Geogr. Distr. Coast of Oregon from Rogue River to the mouth of the Columbia.

Genl. Char. Size large; no white or grayish dorsal streaks; five distinct black stripes on back, intermediate ones slightly paler than the upper parts; antorbital stripe distinct; face ochraceous; under parts white.

Color. Upper parts and sides uniform dark ochre; top of head and nose black and ochraceous; five black dorsal stripes, central one from occiput to root of tail; space between stripes slightly paler than body color; face stripes and nose black; cheeks ochraceous; postauricular spot grayish white, under parts white; tail above dark ochraceous at base with black hairs intermingled, rest black tipped with white, tip black edged with white, beneath dark ochre at base, rest reddish ochre with black lateral band fringed with white; hands and feet pale grayish buff; ear black, with broad external band, including tip, white, external basal edge ochraceous.

Measurements. Total length, 262; tail vertebræ, 115; hind foot, 37; ear, 23. Skull: occipito-nasal length, 38; Hensel, 30; zygomatic width, 21; interorbital constriction, 8.5; length of nasals, 11; palatal length, 16; length of upper tooth row, alveolar border, 6; length of mandible, 16.

163. *Tamias merriami* (Allen).

Tamias asiaticus merriami Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1899, p. 176.

Tamias merriami Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 71. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 137. Zoöl. Ser.

MERRIAM'S CHIPMUNK.

Type locality. San Bernardino Mountains, California.

Geogr. Distr. Mountains of southern California from Tulare and Monterey Counties south into Lower California, Mexico.

164. *Tamias bulleri* (Allen).

Tamias asiaticus bulleri Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1889, p. 173.

Tamias bulleri Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 136. Zoöl. Ser.

MEXICAN CHIPMUNK.

Type locality. Sierra de Valparaiso, State of Zacatecas, Mexico.

Geogr. Distr. State of Zacatecas, Mexico.

165. *Tamias durangæ* (Allen).

Eutamias durangæ Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1903, p. 595.

Tamias durangæ Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 137. Zoöl. Ser.

DURANGO CHIPMUNK.

Type locality. Arroyo de Bucy, State of Durango, Mexico.

Geogr. Distr. State of Durango, Mexico, range unknown.

166. *Tamias obscurus* Allen.

Tamias obscurus Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1890, p. 70. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 136, figs. 22, xxx. Zoöl. Ser.

LOWER CALIFORNIA CHIPMUNK.

Type locality. San Pedro Martir Mountains, Lower California, Mexico.

Geogr. Distr. San Pedro Martir and Hanson Laguna Mountains, Lower California.

167. *Tamias quadrimaculatus* Gray.

Tamias quadrimaculatus Gray, Ann. Mag. Nat. Hist., 3d Ser., xx, 1867, p. 435. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 71. Zoöl. Ser.

Tamias macrorhabdotes Merr., Proc. Biol. Soc. Wash., 111, 1886, p. 25, fig. p. 26.

FOUR-BANDED CHIPMUNK.

Type locality. Michigan Bluff, Sacramento Valley, California.

Geogr. Distr. Central California in the Sierra Nevada, Placer County, to Fort Klamath, Oregon, through the valley of the Sacramento River.

168. *Tamias senex* Allen.

Tamias senex Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1890, p. 83.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 72.
Zoöl. Ser.

GRAY CHIPMUNK.

Type locality. Summit of Donner Pass, Placer County, California

Geogr. Distr. Cascade Mountains from Farewell Bend on the Des Chutes River in Oregon, south to the head waters of the Merced River in the Yosemite National Park.

169. *Tamias speciosus* Allen.

Tamias speciosus Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1890,

p. 86. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901,
p. 72. Zoöl. Ser.

SHOWY CHIPMUNK.

Type locality. San Bernardino Mountains, San Bernardino County, California.

Geogr. Distr. San Bernardino and San Jacinto Mountains, California.

a. *Tamias speciosus inyoensis* (Merriam).

Eutamias speciosus inyoensis Merr., Proc. Biol. Soc. Wash., XI,
1897, p. 208.

Tamias callipeplus inyoensis Elliot, Syn. N. Am. Mamm., F. C. M.

Pub., 11, 1901, p. 73. Zoöl. Ser. (Misplaced on page.)

INYO MOUNTAINS CHIPMUNK.

Type locality. White Mountains, Inyo County, California.

Geogr. Distr. Summits of the White and Inyo Mountains, California.

170. *Tamias callipeplus* Merriam.

Tamias callipeplus Merr., Proc. Biol. Soc. Wash., VIII, 1893, p. 136.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 73.
Zoöl. Ser.

BEAUTIFULLY GARBED CHIPMUNK.

Type locality. Mount Pinos, Ventura County, California.

Geogr. Distr. Mount Pinos and head waters of Yule River, north nearly to the Yosemite Valley.

171. *Tamias frater* Allen.

Tamias frater Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1890, p. 88.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 73.

Zoöl. Ser.

RELATED CHIPMUNK.

Type locality. Sierra Nevada, Placer County, California.

Geogr. Distr. Western Nevada to Central California and region of Lake Tahoe.

172. *Tamias panamintinus* Merriam.

Tamias panamintinus Merr., Proc. Biol. Soc. Wash., VIII, 1893,

p. 134. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901,

p. 74. Zoöl. Ser.

PANAMINT MOUNTAINS CHIPMUNK.

Type locality. Johnson Cañon, Panamint Mountains, Inyo County, California.

Geogr. Distr. West side of Great Basin in California and Nevada.

173. *Tamias amœnus* Allen.

Tamias amœnus Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1890,

p. 90. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901,

p. 74. Zoöl. Ser.

CHARMING CHIPMUNK.

Type locality. Fort Klamath, Klamath County, Oregon.

Geogr. Distr. Fort Klamath, Oregon, to Placer County, California, and western Nevada about Lake Tahoe.

174. *Tamias caurinus* (Merriam).

Eutamias caurinus Merr., Proc. Acad. Nat. Scien. Phil., 1898,

p. 352.

Tamias caurinus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II,

1901, p. 74. Zoöl. Ser.

NORTHWEST CHIPMUNK.

Type locality. Solduck Trail, Clallam County, Olympic Mountains, Washington.

Geogr. Distr. Olympic Mountains, Washington; timber line.

175. *Tamias cinereicollis* Allen.

Tamias cinereicollis Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1890,

p. 94. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901,

p. 75. Zoöl. Ser.

ASH-COLORED CHIPMUNK.

Type locality. San Francisco Mountains, Coconino County, Arizona.

Geogr. Distr. San Francisco and Mogollon Mountains, Arizona.

a. *Tamias cinereicollis canipes* (Bailey).

Eutamias cinereicollis canipes Bailey, Proc. Biol. Soc. Wash., xv, 1902, p. 117.

GRAY-FOOTED CHIPMUNK.

Type locality. Guadalupe Mountains, El Paso County, Texas. Altitude, 7,000 feet.

Genl. Char. Similar to *T. cinereicollis* in fresh post-breeding pelage but grayer; paler ochraceous on sides; tips of long hairs on rump and upper surface of tail white instead of yellowish; feet clear gray; black dorsal stripe reaching to between ears.

Measurements. Total length, 230; tail vertebrae, 104; hind foot, 35. Skull: basal length, 30; length of nasals, 11; zygomatic width, 19.3; mastoid breadth, 15.8; length of upper molar series, alveolar border, 5.8.

176. *Tamias umbrinus* Allen.

Tamias umbrinus Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1890, p. 96. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 75. Zool. Ser.

DARK CHIPMUNK.

Type locality. Uintah Mountains, Utah, south of Fort Bridger, Sweetwater County, Wyoming.

Geogr. Distr. Wahsatch and Uintah Ranges of northern and central Utah, north into Wyoming.

177. *Tamias quadrivittatus* (Say).

Sciurus quadrivittatus Say, Long's Exped. Rocky Mts., 11, 1823, p. 45.

Tamias quadrivittatus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 75. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 138. Zool. Ser.

COLORADO CHIPMUNK.

Type locality. Arkansas River, Colorado, "near where it breaks through the foothills." Park County?

Geogr. Distr. State of Durango, Mexico, north to southern boundary of Colorado, northward through Wyoming to and including the Yellowstone National Park.

a. *Tamias quadrivittatus gracilis* Allen.

Tamias quadrivittatus gracilis Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1890, p. 99. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 76. Zool. Ser.

GRACEFUL CHIPMUNK.

Type locality. San Pedro, Socorro County, New Mexico.

Geogr. Distr. Socorro County, New Mexico, to Apache County, Arizona.

b. Tamias quadrivittatus luteiventris Allen.

Tamias quadrivittatus luteiventris Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1890, p. 101. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 76. Zoöl. Ser.

BUFF-BELLIED CHIPMUNK.

Type locality. Chief Mountain Lake, northwestern Montana.

Geogr. Distr. Rocky Mountains in Montana from Helena northward; probably into British America.

c. Tamias quadrivittatus felix Rhoads.

Tamias quadrivittatus felix Rhoads, Amer. Nat., xxix, 1895, p. 941. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 76. Zoöl. Ser.

HAPPY CHIPMUNK.

Type locality. Mt. Baker Range, Westminster District, British Columbia, Canada. Altitude, 7,000 feet.

Geogr. Distr. Mt. Baker Range, British Columbia; range unknown.

d. Tamias quadrivittatus affinis Allen.

Tamias quadrivittatus affinis Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1890, p. 103. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 77. Zoöl. Ser.

ALLIED CHIPMUNK.

Type locality. Ashcroft, British Columbia, Canada.

Geogr. Distr. Interior of British Columbia, east of the Cascade Mountains.

e. Tamias quadrivittatus borealis (Allen).

Tamias asiaticus borealis Allen, Mon. N. Am. Rod., 1877, p. 793.

Tamias quadrivittatus borealis Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 77. Zoöl. Ser.

NORTHERN CHIPMUNK.

Type locality. Fort Liard, Mackenzie, Canada.

Geogr. Distr. Northern North America, Hudson Bay to eastern base of Rocky Mountains; also in North and South Dakota and Montana.

f. Tamias quadrivittatus neglectus Allen.

Tamias quadrivittatus neglectus Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1890, p. 106. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 78. Zoöl. Ser.

NEGLECTED CHIPMUNK.

**Type locality.* Eastern end of Lake Superior, Ontario, Canada.

Geogr. Distr. Northeastern Minnesota, northern Wisconsin, northern peninsula of Michigan, and northern shore of Lake Superior.

178. *Tamias caniceps* (Osgood).

Eutamias caniceps Osgood, N. Am. Faun. No. 19, 1900, p. 28.

Tamias caniceps Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, Append., p. 426. Zoöl. Ser.

GRAY-HEADED CHIPMUNK.

Type locality. Lake Lebarge, Yukon, Canada.

Geogr. Distr. Head waters of the Yukon to Fort Selkirk, Long. 137° N., Lat. 62° W.

179. *Tamias canicaudus* (Merriam).

Eutamias canicaudus Merr., Proc. Biol. Soc. Wash., xvi, 1903, p. 77.

GRAY-TAILED CHIPMUNK.

Type locality. Spokane, Spokane County, Washington.

Geogr. Distr. Washington. Limits of range unknown.

Genl. Char. Size large; tail long; outer pair of stripes white.

Color. Spring Pelage. Above with middle pair of light stripes vinaceous gray; other stripes black, outer pair washed with fulvous; tail above grizzled gray edged with whitish; beneath with a median buffy band bordered with black and edged with whitish.

Post-breeding Pelage. Neck, sides, and edges of stripes washed with ochraceous or light fulvous.

Measurements. Total length, 228; tail vertebræ, 98; hind foot, 32.

180. *Tamias minimus* Bachman.

Tamias minimus Bachm., Journ. Acad. Nat. Scien. Phil., 1839, p. 71. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901,

p. 78. Zoöl. Ser.

LEAST CHIPMUNK.

Type locality. Near Green River City, southwestern Wyoming.

Geogr. Distr. Bad Lands and plains of the Dakotas, Montana, and Wyoming.

a. Tamias minimus consobrinus (Allen).

Tamias asiaticus var. *pallidus* Allen, Mon. N. A. Roden., 1877, p. 793. (Part.)

Tamias minimus consobrinus Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1890, p. 112. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 78. Zoöl. Ser.

*Definite type locality of this species is not given.

RELATED CHIPMUNK.

Type locality. Wahsatch foothills near Salt Lake City, Utah.

Geogr. Distr. Eastern border of the Great Basin, in eastern Utah, western and southern Colorado, and northwestern New Mexico.

b. *Tamias minimus melanurus* Merriam.

Tamias minimus melanurus Merr., N. Am. Faun., No. 4, 1890, p. 22. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 79. Zoöl. Ser.

BLACK-TAILED CHIPMUNK.

Type locality. Near Blackfoot, Bingham County, Idaho.

Geogr. Distr. Known only from type locality.

c. *Tamias minimus pictus* Allen.

Tamias minimus pictus Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1890, p. 115. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 79. Zoöl. Ser.

PAINTED CHIPMUNK.

Type locality. Kelton, Boxelder County, Utah.

Geogr. Distr. Western border of Great Salt Lake westward, and from southern Utah and southern Nevada to the Snake Plains of eastern Washington.

d. *Tamias minimus alpinus* (Merriam).

Tamias alpinus Merr., Proc. Biol. Soc. Wash., VIII, 1893, p. 137.

Tamias minimus alpinus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 79. Zoöl. Ser.

ALPINE CHIPMUNK.

Type locality. Big Cottonwood Meadows, high Sierras, Tulare County, California, south of Mt. Whitney.

Geogr. Distr. Alpine summit of high Sierras, California.

181. *Tamias lectus* (Allen).

Eutamias lectus Allen, Bull. Brookl. Inst. Arts & Scien., I, No. 5, 1905, p. 117.

BEAVER VALLEY CHIPMUNK.

Type locality. Beaver Valley, Beaver County, Utah.

Geogr. Distr. Unknown.

Color. Post-breeding Pelage. Similar in size to *T. m. pictus*, but colors stronger; flanks, shoulders, and sides of neck, deep rufous; face stripes broader and darker, and light stripes narrower; post-auricular patch gray; central area of tail beneath bright rufous.

Breeding Pelage. Closely resembling the coloration of *T. cinerei-*

collis, but distinguished from that species by the small size of the present form.

Measurements. None given.

182. *Tamias adsitus* (Allen).

Eutamias adsitus Allen, Bull. Brookl. Inst. Arts & Scien., 1, No. 5, 1905, p. 118.

BEAVER MOUNTAINS CHIPMUNK.

Type locality. Briggs Meadows, Beaver Range Mountains, Utah. Altitude, 10,000 feet.

Geogr. Distr. Unknown.

Color. Similar to *T. lectus* in coloration, but one-third larger.

Measurements. None given.

183. *Tamias oreocetes* (Merriam).

Eutamias oreocetes Merr., Proc. Biol. Soc. Wash., XI, 1897, p. 207.

Tamias oreocetes Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 80. Zoöl. Ser.

MOUNTAIN CHIPMUNK.

Type locality. Timber line near Summit, Teton Mountains, Teton County, Montana.

Geogr. Distr. Known only from type locality.

Subgenus B. *Tamias*.

184. *Tamias striatus* (Linnæus).

Sciurus striatus Linn., Syst. Nat., 1, 1758, p. 64.

Sciurus americanus Gmel., Syst. Nat., 1, 1788, p. 150.

Tamias striatus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 80. Zoöl. Ser.

STRIPED CHIPMUNK.

Type locality. Virginia?

Geogr. Distr. Southern New York through Atlantic States to Georgia.

a. Tamias striatus lysteri (Richardson).

Sciurus (Tamias) lysteri Rich., Faun. Bor. Amer., Mamm., 1, 1829, p. 181, pl. 15.

Tamias striatus lysteri Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 80. Zoöl. Ser.

LYSTER'S STRIPED CHIPMUNK.

Type locality. "Penetanguishene," Georgian Bay, Ontario, Canada.

Geogr. Distr. North latitude 50° in eastern North America south to northern New York, west to Michigan, and Ontario, Canada.

b. *Tamias striatus griseus* Mearns.

Tamias striatus griseus Mearns, Bull. Am. Mus. Nat. Hist. N. Y., 1891, p. 231. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 81. Zoöl. Ser.

GRAY STRIPED CHIPMUNK.

Type locality. Fort Snelling, Hennepin County, Minnesota.

Geogr. Distr. Upper Mississippi Valley, west of Great Lakes.

c. *Tamias striatus venustus* Bangs.

Tamias striatus venustus Bangs, Proc. Biol. Soc. Wash., x, 1896, p. 137. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 81. Zoöl. Ser.

CHARMING STRIPED CHIPMUNK.

Type locality. Stillwell, Boston Mountains, Indian Territory.

Geogr. Distr. Known only from the type locality.

Genus 52. **Citellus. Spermophiles.**

Citellus Oken, Lehrb. der Naturg., Zool., 11, 1816, p. 842. Type *Mus citellus* Linnæus.

Spermophilus F. Cuv., Dents Mamm., 1825, pp. 160-161, pls. LV, 255.

Spermatophilus Wagl., Nat. Syst. Amphib., 1830, p. 22.

Citillus Lichten., Darst. neuer oder wenig bekannt. Säugeth., Heft 5, 1827-34, Taf. xxxi, fig. 2.

Colobotis Brandt, Bull. Classe Phys.-math. Acad. Imp. Scien. St. Pétersb., 11, 1844, p. 365.

Otospermophilus Brandt, Bull. Classe Phys.-math. Acad. Imp. Scien. St. Pétersb., 11, 1844, p. 379.

Otocolobus Brandt, Bull. Classe Phys.-math. Acad. Imp. Scien. St. Pétersb., 11, 1844, p. 382.

Ictidomys Allen, Mon. N. Am. Rod., 1877, p. 821.

Spermophilopsis Blasius, Tag. 57ten Vers. Deut. Nat. Aertz., 1884, p. 324.

Ammospermophilus Merr., Proc. Biol. Soc. Wash., VII, 1892, p. 27.

Xerospermophilus Merr., Proc. Biol. Soc. Wash., VII, 1892, p. 27.

Callospermophilus Merr., Proc. Biol. Soc. Wash., XI, 1897, p. 189 (note).

Subgenus A. *Ammospermophilus*.

185. **Citellus harrisi** (Audubon and Bachman).

Spermophilus harrisii Aud. & Bachm., Quad. N. Am., III, 1854, p. 267, pl. 144, fig. 1. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 85. Zoöl. Ser.

Citellus harrisi Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 141. Zoöl. Ser.

HARRIS' SPERMOPHILE.

Type locality. Unknown.

Geogr. Distr. Southern Utah and Nevada into California and south into Arizona and northwestern New Mexico.

a. *Citellus harrisi saxicola* (Mearns).

Spermophilus harrisii saxicolus! Mearns, Proc. U. S. Nat. Mus., 1895, p. 444. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 86.

Citellus harrisi saxicola Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 142, fig. 23. Zoöl. Ser.

ROCK SPERMOPHILE.

Type locality. Tinajas Atlas, Gila Mountains, Yuma County, Arizona.

Geogr. Distr. Gila Mountains in southwestern Arizona, into Sonora.

186. *Citellus leucurus* (Merriam).

Tamias leucurus Merr., N. Am. Faun., No. 2, 1889, p. 20.

Spermophilus leucurus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 86, fig. 18. Zoöl. Ser.

Citellus leucurus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 142. Zoöl. Ser.

WHITE-TAILED SPERMOPHILE.

Type locality. San Gorgonio Pass, Riverside County, California.

Geogr. Distr. Lower California, Mexico into California, Utah, Arizona, and New Mexico.

a. *Citellus leucurus cinnamomeus* (Merriam).

Tamias leucurus cinnamomeus Merr., N. Am. Faun., No. 3, 1890, pp. 51, 52.

Spermophilus leucurus cinnamomeus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 86. Zoöl. Ser.

CINNAMON-HUED SPERMOPHILE.

Type locality. Echo Cliffs, Painted Desert, Coconino County, Arizona.

Geogr. Distr. Desert region of Colorado and Arizona.

b. *Citellus leucurus peninsulæ* (Allen).

Tamias leucurus peninsulæ Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1893, p. 197.

Citellus leucurus peninsulæ Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 143, fig. 24. Zoöl. Ser.

LOWER CALIFORNIA SPERMOPHILE.

Type locality. San Telmo, Lower California, Mexico.

Geogr. Distr. Lower California, Mexico; range unknown.

187. *Citellus vinnulus* Elliot.

Citellus vinnulus Elliot, Pub. Field Columb. Mus., III, 1903, p. 241.

Zoöl. Ser.

CHARMING SPERMOPHILE.

Type locality. Keeler, Owens Lake, Inyo County, California.

Geogr. Distr. Panamint, Coso, and Inyo Mountains to Keeler, Owens Lake, Inyo County, California.

Genl. Char. Similar to *C. l. peninsulae*, but body and tail beneath white instead of pale yellow. Generally darker than *C. leucurus*, with smaller hind feet.

Color. Top of head and upper parts mixed black and vinaceous, the latter hue predominating; nape and between shoulders grayish; two white stripes from shoulders to rump; shoulders, top of fore legs and feet to toes, thighs and upper surface of hind legs and feet vinaceous; toes white on fore feet, buffy vinaceous at tips of hind feet; sides and face and neck and entire under parts of body, legs and feet silvery white, base of fur plumbeous; tail above on basal third like back, remainder black with white hairs intermingled, and edged with white, beneath white with a subapical black bar.

Measurements. Total length, 215; tail vertebræ, 66; hind foot, 38.5; ear, 12. Skull: total length, 37.3; Hensel, 30; zygomatic width, 19; interorbital constriction, 10; length of nasals, 10.5; palatal length, 12; length of upper tooth row, 7; length of mandible, 22; length of lower tooth row, 7.

188. *Citellus interpres* (Merriam).

Tamias interpres Merr., N. Am. Faun., No. 4, 1890, p. 21.

Spermophilus interpres Elliot, Syn. N. Am. Mamm., F. C. M. Pub.,

II, 1901, p. 86. Zoöl. Ser.

Citellus interpres Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., IV, Pt. 1, 1904, p. 143. Zoöl. Ser.

TRADER SPERMOPHILE.

Type locality. El Paso, El Paso County, Texas.

Geogr. Distr. State of Chihuahua, Mexico, north to Utah.

189. *Citellus nelsoni* (Merriam).

Spermophilus nelsoni Merr., Proc. Biol. Soc. Wash., VIII, 1893,

p. 129. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901,

p. 87. Zoöl. Ser.

NELSON'S SPERMOPHILE.

Type locality. Tipton, San Joaquin Valley, Tulare County, California.

Geogr. Distr. South central California.

Subgenus B. *Xerospermophilus*.190. *Citellus obsoletus* (Kennicott).

Spermophilus obsoletus Kennicott, Proc. Acad. Nat. Scien. Phil., 1863, p. 157. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 95. Zoöl. Ser.

KENNICOTT'S SPERMOPHILE.

Type locality. Western Nebraska.

Geogr. Distr. Black Hills, South Dakota into Nebraska and westward to Utah.

191. *Citellus spilosoma* (Bennett).

Spermophilus spilosoma Bennett, Proc. Zoöl. Soc., 1833, p. 40. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 96. Zoöl. Ser.

Spermophilus mexicanus Aud. & Bachm., Quad. N. Am., III, 1853, p. 42, pl. CIX.

Citellus spilosoma Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 145. Zoöl. Ser.

SPOTTED SPERMOPHILE.

Type locality. California near Mexican boundary.

Geogr. Distr. New Mexico and Texas south into Sonora, Mexico.

a. *Citellus spilosoma macrospilotus* (Merriam).

Spermophilus spilosoma macrospilotus Merr., N. Am. Faun., No. 4, 1890, p. 38. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 96, fig. 21. Zoöl. Ser. (Misprinted *microspilotus*.)

Citellus spilosoma macrospilotus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 145, fig. 25. Zoöl. Ser. (Misprinted *microspilotus*.)

SPOTTED SPERMOPHILE.

Type locality. Oracle, Pinal County, Arizona.

Geogr. Distr. Southern Arizona into State of Sonora, Mexico.

b. *Citellus spilosoma major* (Merriam).

Spermophilus spilosoma major Merr., N. Am. Faun., No. 4, 1890, p. 39. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 96. Zoöl. Ser.

LARGE SPOTTED SPERMOPHILE.

Type locality. Albuquerque, Bernalillo County, New Mexico.

Geogr. Distr. Eastern New Mexico.

c. *Citellus spilosoma pratensis* (Merriam).

Spermophilus spilosoma pratensis Merr., N. Am. Faun., No. 3, 1890, p. 55. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 96. Zool. Ser.

PARK SPERMOPHILE.

Type locality. Pine plateau at north foot of San Francisco Mountain, Coconino County, Arizona.

Geogr. Distr. San Francisco Mountain, northwestern Arizona.

d. *Citellus spilosoma marginatus* (Bailey).

Spermophilus spilosoma marginatus Bailey, Proc. Biol. Soc. Wash., XV, 1902, p. 118.

BAILEY'S SPERMOPHILE.

Type locality. Alpina, Brewster County, Texas.

Geogr. Distr. Limits of range unknown.

Genl. Char. Similar to typical *C. spilosoma* from New Mexico, but smaller and colors brighter.

Color. Above bright cinnamon brown; back from the ears spotted with whitish edged with black.

Measurements. Total length, 225; tail vertebræ, 67; hind foot, 32. Skull: basal length, 33.7; length of nasals, 13.3; zygomatic breadth, 23.5; mastoid breadth, 19; alveolar length of upper molar series, 7.2.

e. *Citellus spilosoma obsidianus* (Merriam).

Spermophilus spilosoma obsidianus Merr., N. Am. Faun., No. 3, 1890, p. 56. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 97. Zool. Ser.

DARK SPOTTED SPERMOPHILE.

Type locality. Cedar Belt, northeast of San Francisco Mountain, Coconino County, Arizona.

Geogr. Distr. Limits of range unknown.

f. *Citellus spilosoma annectens* (Merriam).

Spermophilus spilosoma annectens Merr., Proc. Biol. Soc. Wash., VIII, 1893, p. 132. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 97. Zool. Ser.

PADRE ISLAND SPERMOPHILE.

Type locality. Padre Island, Cameron County, Texas.

Geogr. Distr. Mustang and Padre Islands, Cameron County, Texas; the former island probably the northern limit of the race.

g. *Citellus spilosoma arens* (Bailey).

Spermophilus spilosoma arens Bailey, Proc. Biol. Soc. Wash., xv, 1902, p. 118.

LANGUISHING SPERMOPHILE.

Type locality. El Paso, El Paso County, Texas.

Geogr. Distr. Limits of range unknown.

Genl. Char. Similar to *C. cryptospilotus* but larger, spots coarser.

Skull: longer, nasals wider, dentition heavier.

Color. Above light clay or dark buffy, spotted over back and rump with white without black edgings.

Measurements. Total length, 237; tail vertebræ, 85; hind foot, 35. Skull: basal length, 33; length of nasals, 12; zygomatic breadth, 18; length of upper molar series, 7.

192. *Citellus cryptospilotus* (Merriam).

Spermophilus cryptospilotus Merr., N. Am. Faun., No. 3, 1890, p. 57. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 97. Zoöl. Ser.

DESERT SPERMOPHILE.

Type locality. Tenebitto Wash, Painted Desert, 25 miles east of the Little Colorado, Coconino County, Arizona.

Geogr. Distr. Painted Desert, Arizona.

193. *Citellus canescens* (Merriam).

Spermophilus canescens Merr., N. Am. Faun., No. 4, 1890, p. 38. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 97. Zoöl. Ser.

DRAB GRAY SPERMOPHILE.

Type locality. Wilcox, Cochise County, Arizona.

Geogr. Distr. Southeastern Arizona.

194. *Citellus eremonomus* Elliot.

Citellus eremonomus Elliot, Pub. Field Columb. Mus., III, 1903, p. 243. Zoöl. Ser.

DEATH VALLEY SPERMOPHILE.

Type locality. Furnace Creek, Death Valley, Inyo County, California.

Genl. Char. Size small; color dark; tail with only basal half like the back. Skull: similar to that of *C. t. mohavensis*, but processes of the pterygoids do not touch the bullæ.

Color. Upper parts grizzled vinaceous cinnamon; sides of face, nose, and body, inner sides of limbs and entire under parts silvery white; fore feet pale brown, hind feet whitish; tail above, basal half grizzled vinaceous cinnamon like the back, terminal half blackish

mixed with white hairs and narrowly edged with white, beneath silvery white at base, remainder buff mixed with black, bordered and tipped with black and narrowly fringed with white. Ear very small, similar in color to the back.

Measurements. Total length, 252; tail vertebræ, 89; hind foot, 35; ear, 8.5. Skull: total length, 36; occipito-nasal length, 35; Hensel, 30; zygomatic width, 23; interorbital constriction, 9.5; palatal length, 17; length of nasals, 12; length of upper tooth row, 7; length of mandible, 2.5; of lower tooth row, 7.

195. *Citellus tereticaudus* (Baird).

Spermophilus tereticaudus Baird, N. Am. Mamm., 1857, p. 315.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 98.
Zoöl. Ser.

Citellus tereticaudus Elliot, Mamm. Middle Amer. & W. Indies,
F. C. M. Pub., IV, Pt. I, 1904, p. 144. Zoöl. Ser.

ROUND-TAILED SPERMOPHILE.

Type locality. Old Fort Yuma, San Diego County, California.

Geogr. Distr. Central California to Tucson Plain, Arizona; Lower California and State of Sonora, Mexico.

a. *Citellus tereticaudus mohavensis* (Merriam).

Spermophilus mohavensis Merr., N. Am. Faun., No. 2, 1889, p. 15.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 98.
Zoöl. Ser.

Citellus tereticaudus mohavensis Elliot, Pub. Field Columb. Mus.,
III, 1903, p. 291. Zoöl. Ser.

MOHAVE DESERT SPERMOPHILE.

Type locality. Mohave River, San Bernardino County, California.

Geogr. Distr. Mohave Desert, California.

b. *Citellus tereticaudus sonoriensis* (Ward).

Spermophilus sonoriensis Ward, Amer. Nat., xxv, 1891, p. 158.

Citellus tereticaudus sonoriensis Elliot, Mamm. Middle Amer. &
W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 144. Zoöl. Ser.

SONORAN SPERMOPHILE.

Type locality. Hermosillo, State of Sonora, Mexico.

Geogr. Distr. State of Sonora. Range unknown.

196. *Citellus perotensis* (Merriam).

Spermophilus perotensis Merr., Proc. Biol. Soc. Wash., VIII, 1893,
p. 131.

Citellus perotensis Elliot, Mamm. Middle Amer. & W. Indies,
F. C. M. Pub., IV, Pt. I, 1904, p. 145. Zoöl. Ser.

PEROTE SPERMOPHILE.

Type locality. Perote, State of Vera Cruz, Mexico.

Geogr. Distr. State of Vera Cruz, Mexico, extreme eastern border of tableland. Altitude, 8,000 to 10,000 feet.

197. *Citellus neglectus* (Merriam).

Spermophilus neglectus Merr., N. Am. Faun., No. 2, 1889, p. 17.

Elliot; Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 98.

Zoöl. Ser.

NEGLECTED SPERMOPHILE.

Type locality. Dolan's Springs, Mohave County, Arizona.

Geogr. Distr. Arizona.

193. *Citellus chlorus* Elliot.

Citellus chlorus Elliot, Pub. Field Columb. Mus., III, 1903, p. 242.

Zoöl. Ser.

PALE SPERMOPHILE.

Type locality. Palm Springs, Riverside County, California.

Geogr. Distr. Riverside and San Diego Counties, California.

Genl. Char. Size equal to *C. tereticaudus*, color entirely different.

Skull heavy, tail long.

Color. Entire upper parts and sides olive gray with a brownish sheen in certain lights; upper parts of arms and thighs olive gray; entire under parts grayish white; hands brownish, feet whitish; tail above, basal half like back, slightly more brownish, apical half blackish mixed with brown and edged with white, beneath pale brown margined narrowly with black and fringed with white; ears very small, blackish.

Measurements. Total length, 255; tail vertebræ, 100; hind foot, 37; ear, 8. Skull: total length, 32; Hensel, 30; interorbital width, 8; zygomatic width, 22; width of braincase, 18; length of nasals, 8; palatal length, 17; length of upper tooth row, 7; length of mandible, 24; length of lower tooth row, 6.5.

199. *Citellus mexicanus* (Erxleben).

Sciurus mexicanus Erxl., Syst. Regn. Anim., I, 1777, p. 428.

Citellus mexicanus Licht., Darst. Neu. oder wenig bekannt.

Säugeth., 1830, pl. 31, fig. 2. (ex Toluca.) Elliot, Mamm.

Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 146. Zoöl. Ser.

Spermophilus mexicanus Elliot, Syn. N. Am. Mamm., F. C. M.

Pub., II, 1901, p. 98. Zoöl. Ser.

MEXICAN SPERMOPHILE.

Type locality. Toluca, Mexico.

Geogr. Distr. South Central Mexico.

a. *Citellus mexicanus parvidens* (Mearns).

Spermophilus mexicanus parvidens Mearns, Proc. U. S. Nat. Mus., XVIII, 1896, p. 443. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 99. Zoöl. Ser.

Citellus mexicanus parvidens Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 146. Zoöl. Ser.

SMALL-TOOTHED SPERMOPHILE.

Type locality. Fort Clark, Kinney County, Texas.

Geogr. Distr. Texas into Mexico.

Subgenus C. *Ictidomys*.

200. *Citellus tredecimlineatus* (Mitchill).

Sciurus tredecimlineatus (sic) Mitch., Lond. Med. Repos., N. S., 1821, p. 248.

Arctomys hoodi Sabine, Trans. Linn. Soc., 1822, p. 590, pl. 29.

Spermophilus tredecimlineatus! Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 100, fig. 22. Zoöl. Ser.

13-STRIPED SPERMOPHILE.

Type locality. Upper Mississippi. Central Minnesota?

Geogr. Distr. Central North America; eastern Michigan to Montana and Colorado, and central Texas north to Saskatchewan Plains in Canada, and in prairie region of the Mississippi from Ohio to Minnesota.

a. *Citellus tredecimlineatus pallidus* (Allen).

Spermophilus tredecimlineatus! *pallidus* Allen, Mon. N. Am. Rod., 1877, p. 872. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 100. Zoöl. Ser.

PALLID SPERMOPHILE.

Type locality. Plains of the lower Yellowstone River.

Geogr. Distr. Wyoming south to Texas and east to Missouri, on the plains and desert regions.

b. *Citellus tredecimlineatus olivaceus* (Allen).

Spermophilus tredecimlineatus! *olivaceus* Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1895, p. 337. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 100. Zoöl. Ser.

OLIVACEOUS SPERMOPHILE.

Type locality. Custer, Black Hills, South Dakota.

Geogr. Distr. South Dakota.

c. *Citellus tredecimlineatus parvus* (Allen).

Spermophilus tredecimlineatus! *parvus* Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1895, p. 337. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 100. Zoöl. Ser.

LITTLE SPERMOPHILE.

Type locality. Uncompahgre Indian Reservation, northeastern Utah.

Geogr. Distr. Utah and Wyoming.

d. *Citellus tredecimlineatus alleni* (Merriam).

Spermophilus tridecemlineatus! alleni Merr., Proc. Biol. Soc. Wash., XII, 1898, p. 71. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 101. Zoöl. Ser.

ALLEN'S SPERMOPHILE.

Type locality. Bighorn Mountains, Wyoming.

Geogr. Distr. Wyoming. Range unknown.

e. *Citellus tredecimlineatus texensis* (Merriam).

Spermophilus tridecemlineatus! texensis Merr., Proc. Biol. Soc. Wash., XII, 1898, p. 71. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 101. Zoöl. Ser.

TEXAS SPERMOPHILE.

Type locality. Gainesville, Cooke County, Texas.

Geogr. Distr. Northern Texas and Oklahoma Territory. Range unknown.

f. *Citellus tredecimlineatus badius* (Bangs).

Spermophilus tridecemlineatus! badius Bangs, Proc. N. Eng. Zoöl. Club, 1899, p. 1. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 101. Zoöl. Ser.

BAY SPERMOPHILE.

Type locality. Stotesburg, Vernon County, Missouri.

Geogr. Distr. Missouri. Range unknown.

201. *Citellus franklini* (Sabine).

Arctomys franklinii Sabine, Trans. Linn. Soc., 1822, p. 587, pl. 27.

Spermophilus franklini Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 101. Zoöl. Ser.

FRANKLIN'S SPERMOPHILE.

Type locality. Near Cumberland House, Saskatchewan, Canada.

Geogr. Distr. Valley of the Saskatchewan, and through those of the Red River and Mississippi to the Dakotas and central Kansas. Eastern limit western Indiana.

Subgenus D. *Callospermophilus*.202. *Citellus lateralis* (Say).

Sciurus lateralis Say, Long's Exped. Rocky Mts., II, 1823, p. 46.

Spermophilus lateralis Elliot, Syn. N. Am. Mamm., F. C. M. Pub., I, 1901, p. 83, fig. 17. Zoöl. Ser.

STRIPED SPERMOPHILE.

Type locality. Arkansas River, near Cañon City, Colorado.

Geogr. Distr. Colorado, Utah (Uintah Mts.) and Arizona (White and San Francisco Mts.).

a. *Citellus lateralis saturatus* (Rhoads).

Tamias lateralis saturatus Rhoads, Proc. Acad. Nat. Scien. Phil., 1895, p. 43.

Spermophilus lateralis saturatus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 83. Zoöl. Ser.

DARK SPERMOPHILE.

Type locality. Lake Kichelos, Kittitas County, Washington.

Geogr. Distr. Central Washington. Range unknown.

203. *Citellus madrensis* (Merriam).

Callospermophilus madrensis Merr., Proc. Wash. Acad. Scien., III, 1901, p. 563.

Citellus madrensis Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 147, fig. 26. Zoöl. Ser.

SIERRA MADRE SPERMOPHILE.

Type locality. Sierra Madre near Guadalupe y Calvo, State of Chihuahua, Mexico. Altitude, 7,000 feet.

Geogr. Distr. State of Chihuahua, Mexico. Range unknown.

204. *Citellus cinerascens* (Merriam).

Tamias cinerascens Merr., N. Am. Faun., No. 4, 1890, p. 20.

Spermophilus cinerascens Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 83. Zoöl. Ser.

ASHY SPERMOPHILE.

Type locality. Helena, Lewis and Clarke County, Montana. Altitude, 4,500 feet.

Geogr. Distr. Montana, Idaho, and northward into Alberta, Canada.

205. *Citellus castanurus* (Merriam).

Tamias castanurus Merr., N. Am. Faun., No. 4, 1890, p. 19.

Spermophilus castanurus Elliot, N. Am. Mamm., F. C. M. Pub., II, 1901, p. 84. Zoöl. Ser.

CHESTNUT-TAILED SPERMOPHILE.

Type locality. Park City, Wahsatch Mountains, Summit County, Utah. Altitude, 7,000 feet.

Geogr. Distr. Wahsatch Mountains, Utah.

206. *Citellus chrysodeirus* (Merriam).

Tamias chrysodeirus Merr., N. Am. Faun., No. 4, 1890, p. 19.

Spermophilus chrysodeirus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 84. Zoöl. Ser.

GOLDEN-NECK SPERMOPHILE.

Type locality. Fort Klamath, Klamath County, Oregon.

Geogr. Distr. Oregon, northern California, and western Nevada.

a. *Citellus chrysodeirus trinitatis* (Merriam).

Callospermophilus chrysodeirus trinitatis Merr., Proc. Biol. Soc. Wash., XIV, 1901, p. 126.

TRINITY MOUNTAINS SPERMOPHILE.

Type locality. Trinity Mountains, east of Hoopa Valley, Humboldt County, California. Altitude, 5,700 feet.

Geogr. Distr. Trinity Mountains, California. Range unknown.

Genl. Char. Large. Similar to *C. chrysodeirus*, but darker.

Color. Like *C. chrysodeirus* in winter pelage, no golden hue on head or shoulders; tail beneath dark chestnut.

Measurements. Total length, 290; tail vertebræ, 105; hind foot, 44.

b. *Citellus chrysodeirus bernardinus* (Merriam).

Spermophilus chrysodeirus brevicaudus Merr., Proc. Biol. Soc. Wash., VIII, 1893, p. 134. (Preoccupied.)

Spermophilus bernardinus Merriam, Science, N. S., VIII, 1898, p. 782.

Spermophilus chrysodeirus bernardinus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 84. Zool. Ser.

SAN BERNARDINO SPERMOPHILE.

Type locality. San Bernardino Peak, San Bernardino County, California.

Geogr. Distr. Southern California.

207. *Citellus wortmani* (Allen).

Tamias wortmani Allen, Bull. Amer. Mus. Nat. Hist. N. Y., 1895, p. 335.

Spermophilus wortmani Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 84. Zool. Ser.

WORTMAN'S SPERMOPHILE.

Type locality. Kinney Ranch, Bitter Creek, Sweetwater County, Wyoming.

Geogr. Distr. Wyoming. Limits of range unknown.

Subgenus E. *Colobotis*.208. *Citellus parryi* (Richardson).

Arctomys parryi Richard., Append. Parry's Sec. Voy., 1825, p. 316.

Arctomys parryi var. *phæognatha* Rich., Faun. Bor. Am., Mamm., I, 1829, p. 161.

Arctomys kennicottii Ross, Canad. Nat. and Geol., 1861, p. 434.

Spermophilus empetra Elliot, Syn. N. Am. Mamm., F. C. M. Pub.,
11, 1901, p. 89. Zoöl. Ser. (nec Pallas.)

PARRY'S SPERMOPHILE.

Type locality. Five Hawser Bay, Lyon Inlet, Melville Peninsula, Keewatin, Canada.

Geogr. Distr. Arctic America from Melville Peninsula to Icy Cape, Bering Sea, southwest along the Yukon River for a considerable distance, and in the east to the verge of the Barren Grounds, Latitude 65°.

a. *Citellus parryi kadiacensis* (Allen).

Spermophilus parryi var. *kadiacensis* Allen, Proc. Bost. Soc. Nat. Hist., XVI, 1874, p. 292.

Spermophilus empetra var. *kadiacensis* Allen, Mon. N. Am. Rod., 1877, p. 839.

Spermophilus empetra kadiacensis Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 90, fig. 20. Zoöl. Ser.

KADIAK ISLAND SPERMOPHILE.

Type locality. Island of Kadiak, Alaska.

Geogr. Distr. Kadiak Island, Alaska, only.

209. *Citellus erythroglutæus* (Richardson).

Arctomys parryi var. *B. erythrogluteia*! Richard., Faun. Bor. Amer., Mamm., 1, 1829, p. 161.

Citellus erythrogluteius! Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1903, p. 534.

RED-THIGHED SPERMOPHILE.

Type locality. Head of Elk River, Rocky Mountains. Latitude 57° N.

Geogr. Distr. From Elk River, British Columbia, possibly northward to Lake Bennett.

Genl. Char. Resembles *C. columbianus* in size and general coloration. Lower surface of tail rufous. Skull narrow, slender.

Color. Male Post-breeding Pelage. Dorsal area iron gray blotched with small white spots, the hairs tipped with whitish; top of nose to middle of eyes bright rusty chestnut; sides of neck, shoulders, forearms and thighs and the entire ventral surface deep rusty ochraceous; central area of the tail beneath rufous.

Measurements. Total length, 318-349; tail vertebræ, 75-92; hind foot, 47-56; ear, 11-12.

210. **Citellus plesius* (Osgood).

Spermophilus empetra plesius Osgood, N. Am. Faun., No. 19,

*This may prove to be the same as *C. erythroglutæus* Richardson.

1900, p. 29. Elliot, Syn. N. Am. Mamm., F. C. M. Pub.,
II, 1901, Append., p. 427. Zool. Ser.

OSGOOD'S SPERMOPHILE.

Type locality. Bennett City, Lake Bennett, British Columbia, Canada.

Geogr. Distr. South Side of White Pass to Fort Selkirk, Alaska.

a. Citellus plesius ablusus Osgood.

Citellus plesius ablusus Osgood, Proc. Biol. Soc. Wash., xvi,
1903, p. 25.

NUSHAGAK SPERMOPHILE.

Type locality. Nushagak, Alaska.

Genl. Char. Similar to *C. plesius*, but larger. Skull larger, nasals longer, narrower, and more restricted posteriorly; molars about the size of those of *C. plesius*.

Color. Forehead and crown burnt umber; sides of head, nape, shoulders and neck pale gray, blackish slate under fur showing through in places; orbital ring and subauricular spot white; middle of back and rump Vandyke brown spotted with grayish white; under parts dull grayish white; tail beneath tawny in center, lateral hairs with three or four dusky rings and grayish white tip; pencil with black subterminal section, then one or two narrow dusky rings; tip white; hands and feet creamy white.

Measurements. Total length, 394; tail vertebræ, 103; hind foot, 60. Skull: Hensel, 49; occipito-nasal length, 57.5; zygomatic breadth, 38; length of nasals, alveolar border, 13.3.

211. *Citellus nebulicola* Osgood.

Citellus nebulicola Osgood, Proc. Biol. Soc. Wash., xvi, 1903, p. 26.

NAGAI ISLAND SPERMOPHILE.

Type locality. Nagai Island, Shumagin Islands, Alaska.

Genl. Char. Similar to *C. p. kadiacensis*, but smaller; tail shorter, paler. Skull: small; basi-occipital narrower; audital bullæ higher; braincase narrower.

Color. Similar to *C. p. kadiacensis*, but paler; tail with less black; under parts ochraceous, as are also the arms, neck, and face; tail beneath, tawny margined with buff and submargined with black for distal half; forehead and crown mars brown; upper parts mixed black and russet spotted with creamy white.

Measurements. Total length, 340; tail vertebræ, 82; hind foot, 53. Skull: Hensel, 42; occipito-nasal length, 49; zygomatic width, 32; length of nasals, 18; upper molar series, alveolar border, 12.

212. *Citellus beringensis* (Merriam).

Spermophilus beringensis Merr., Proc. Wash. Acad. Scien., 11, 1900, p. 20. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 91. Zoöl. Ser.

BERING'S SPERMOPHILE.

Type locality. Cape Lisbourne, Alaska.

Geogr. Distr. Northern Alaska.

213. *Citellus osgoodi* (Merriam).

Spermophilus osgoodi Merr., Proc. Wash. Acad. Scien., 11, 1900, p. 18. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 91. Zoöl. Ser.

OSGOOD'S SPERMOPHILE.

Type locality. Fort Yukon, Alaska.

Geogr. Distr. Northern Alaska.

214. *Citellus barrowensis* (Merriam).

Spermophilus barrowensis Merr., Proc. Wash. Acad. Scien., 11, 1900, p. 19. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 91. Zoöl. Ser.

BARROW'S SPERMOPHILE.

Type locality. Point Barrow, Alaska.

Geogr. Distr. Northwestern Alaska.

215. *Citellus stonii* Allen.

Citellus stonei Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1903, p. 537.

STONE'S SPERMOPHILE.

Type locality. Wrangel Island, Alaska.

Geogr. Distr. Known only from the type locality.

Genl. Char. Size large; colors gray and rufous.

Color. Above gray suffused with rufous, increasing in intensity on lower back and rump, the hairs being black basally, then a bright buff band, then a dark brown band suffused with rusty, and tip white; among these are dispersed long black hairs; front of hairs rusty cinnamon; posterior half rusty and blackish; sides of face and neck gray; arms paler gray and buffy; sides of shoulders suffused with pale rufous; thighs pale rufous; feet above deep rusty, beneath gray; front and sides of nose, underside of head, throat, breast, and inside of arms whitish; ventral surface washed with pale rufous; tail above at base tinged with rusty, rest gray, bordered with black and fringed broadly with pale yellowish white, and a broad subapical band of black; beneath pale brownish rusty, with a black subapical band and a yellowish white fringe.

Measurements. Total length, 350; tail vertebræ, 100; hind foot, 59. Skull: total length, 59; zygomatic breadth, 37; length of nasals, 22; width of nasals at base, 5.

216. *Citellus columbianus* (Ord).

Arctomys columbianus Ord, Guth. Geogr., 2d Am. ed., II, 1815, p. 292.

Spermophilus columbianus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 91. Zoöl. Ser.

COLUMBIAN SPERMOPHILE.

Type locality. Camas prairie, between the forks of the Clearwater and Kooskooskie Rivers, about forty miles from Moscow, Lincoln County, Washington.

Geogr. Distr. Western Montana to Washington and north through British Columbia and Alaska to Plover Bay in Siberia.

a. Citellus columbianus albertæ Allen.

Citellus columbianus albertæ Allen, Am. Mus. Nat. Hist. N. Y., 1903, p. 537. (Footnote.)

ALBERTA SPERMOPHILE.

Type locality. Canadian National Park, Alberta, Canada.

Genl. Char. Similar to *C. columbianus*. Skull more massive; interorbital foramen triangular instead of circular and postpalatal border more produced.

Color. Like *C. columbianus*, but paler.

Measurements. Total length, 321; tail vertebræ, 108; hind foot, 57. Skull: total length, 52; zygomatic breadth, 33.

217. *Citellus richardsoni* (Sabine).

Arctomys richardsonii Sabine, Trans. Linn. Soc., 1822, p. 589, pl. 28.

Arctomys (Spermophilus) guttatus Rich., Faun. Bor. Amer., Mamm., I, 1829, p. 162. (nec Temm.)

Spermophilus richardsoni Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 92. Zoöl. Ser.

RICHARDSON'S SPERMOPHILE.

Type locality. Carlton House, Saskatchewan, Canada.

Geogr. Distr. From Aberdeen, South Dakota, to North Dakota and Montana, west to Rocky Mountains and north to the Saskatchewan River to latitude 55°.

218. *Citellus townsendi* (Bachman).

Spermophilus townsendii Bachm., Jour. Acad. Nat. Scien. Phil., 1839, p. 61. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 92. Zoöl. Ser.

TOWNSEND'S SPERMOPHILE.

Type locality. Plains of the Columbia near the mouth of the Walla Walla River, Walla Walla County, Washington.

Geogr. Distr. Nebraska westward to plains of the Columbia River, and from Wyoming and Utah to Montana, Idaho, and Oregon.

219. Citellus mollis (Kennicott).

Spermophilus mollis Kennicott, Proc. Acad. Nat. Scien. Phil., 1863, p. 158. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 92. Zoöl. Ser.

KENNICOTT'S SPERMOPHILE.

Type locality. Camp Floyd near Fairfield, Wasatch County, Utah.

Geogr. Distr. Utah and Nevada, limits of range unknown.

a. Citellus mollis stephensi (Merriam).

Spermophilus mollis stephensi Merr., Proc. Biol. Soc. Wash., 11, 1898, p. 69. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 93. Zoöl. Ser.

STEPHEN'S SPERMOPHILE.

Type locality. Queen's Station, Owens Valley, Esmeralda County, Nevada. Altitude, 6,000 feet.

Geogr. Distr. Nevada.

b. Citellus mollis yakimensis (Merriam).

Spermophilus mollis yakimensis Merr., Proc. Biol. Soc. Wash., 11, 1898, p. 70. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 93. Zoöl. Ser.

YAKIMA SPERMOPHILE.

Type locality. Mabton, Yakima County, Washington.

Geogr. Distr. Eastern Washington.

c. Citellus mollis canus (Merriam).

Spermophilus mollis canus Merr., Proc. Biol. Soc. Wash., 11, 1898, p. 70. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 93. Zoöl. Ser.

HOARY SPERMOPHILE.

Type locality. Antelope, Wasco County, Oregon.

Geogr. Distr. Northern Oregon, range unknown.

220. Citellus armatus (Kennicott).

Spermophilus armatus Kennicott, Proc. Acad. Nat. Scien. Phil., 1863, p. 158. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 93. Zoöl. Ser.

ARMED SPERMOPHILE.

Type locality. Fort Bridger, foothills of the Uinta Mountains, Sweetwater County, Utah.

Geogr. Distr. Utah, Wyoming, Idaho, and Montana. Precise range unknown.

221. *Citellus elegans* (Kennicott).

Spermophilus elegans Kennicott, Proc. Acad. Nat. Scien. Phil., 1863, p. 158. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 94. Zoöl. Ser.

Spermophilus richardsoni var. *townsendi* Allen, Mon. Rod., 1877, p. 848. (nec. Bachm.)

ELEGANT SPERMOPHILE.

Type locality. Fort Bridger, Sweetwater County, Wyoming.

Geogr. Distr. Wyoming and Utah.

222. *Citellus beldingi* (Merriam).

Spermophilus beldingi Merr., Ann. Acad. Nat. Scien. N. Y., 14, 1888, p. 317. Elliot, N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 94. Zoöl. Ser.

BELDING'S SPERMOPHILE.

Type locality. Donner, Placer County, California.

Geogr. Distr. California.

223. *Citellus oregonus* (Merriam).

Spermophilus oregonus Merr., Proc. Biol. Soc. Wash., XII, 1898, p. 69. Elliott, N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 94. Zoöl. Ser.

OREGON SPERMOPHILE.

Type locality. Swan Lake Valley, Klamath Basin, Oregon.

Geogr. Distr. Known only from type locality.

Subgenus F. *Otospermophilus*.

224. *Citellus variegatus* (Erxleben).

Sciurus variegatus Erxl., Syst. Regn. Anim., 1, 1777, p. 421.

Citellus variegatus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 148, figs. 27, xxxi. Zoöl. Ser.

LONG-TAILED SPERMOPHILE.

Type locality. State of Jalisco?

Geogr. Distr. States of Sonora and Jalisco, Mexico.

a. *Citellus variegatus grammurus* (Say).

Sciurus grammurus Say, Long's Exped. Rocky Mts., 11, 1823, p. 72.

Spermophilus macrourus Bennett, Proc. Zoöl. Soc., 1833, p. 41.

Spermophilus couchii Baird, N. Am. Mamm., 1857, p. 311, pl. 81.

Spermophilus grammurus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 88, fig. 19. (Part.) Zoöl. Ser.

Citellus variegatus grammurus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 149. Zoöl. Ser.

BUSHY-TAILED SPERMOPHILE.

Type locality. Purgatory Creek, Arkansas River near Bent's Fort, Otero County, Colorado.

Geogr. Distr. Parks of Central Colorado, south into State of Sonora, Mexico.

b. *Citellus variegatus beecheyi* (Richardson).

Arctomys beecheyi Rich., Faun. Bor. Amer., Mamm., I, 1829, p. 170.

Sciurus (Macroxus) californicus Less., Mamm. & Ois. Nouv., 1847, p. 143.

Spermophilus grammurus beecheyi Elliot, Syn. N. Am. Mamm., F.

C. M. Pub., II, 1901, p. 88. Zoöl. Ser.

BEECHY'S SPERMOPHILE.

Type locality. Vicinity of San Francisco, or Monterey.

Geogr. Distr. West of the Sierra Nevada, northern California.

c. *Citellus variegatus fisheri* (Merriam).

Spermophilus beecheyi fisheri Merr., Proc. Biol. Soc. Wash., VIII, 1893, p. 133.

Spermophilus grammurus fisheri Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 88. Zoöl. Ser.

Citellus variegatus fisheri Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 150. Zoöl. Ser.

FISHER'S SPERMOPHILE.

Type locality. Kern Valley, twenty-five miles above Kernville, Tulare County, California.

Geogr. Distr. Central and southern California to western border of Nevada. Lower California, Mexico.

d. *Citellus variegatus atricapillus* (Bryant).

Spermophilus grammurus atricapillus Bryant, Proc. Calif. Acad. Scien., 2d Ser., II, 1889, p. 26.

Citellus variegatus atricapillus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 150. Zoöl. Ser.

BLACK-HEADED SPERMOPHILE.

Type locality. Comondu, Lower California, Mexico.

Geogr. Distr. Lower California, Mexico.

e. *Citellus variegatus rupestris* Allen.

Citellus grammurus rupestris Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1903, p. 595.

Citellus variegatus rupestris Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 150. Zoöl. Ser.

ROCK SPERMOPHILE.

Type locality. Rio Sestin, State of Durango, Mexico.

Geogr. Distr. State of Durango, Mexico.

f. *Citellus variegatus utah* Merriam.

Citellus variegatus utah Merr., Proc. Biol. Soc. Wash., xvi, 1903,
p. 77.

UTAH SPERMOPHILE.

Type locality. Wasatch Mountains near Ogden, Utah.

Geogr. Distr. Utah.

Genl. Char. "Similar to *C. variegatus*, but smaller, ears longer, back redder, tail darker, nasals longer, and extending more behind premaxillæ, point of incisors pale yellow."

g. *Citellus variegatus douglasi* (Richardson).

Arctomys (Spermophilus) douglasii Rich., Faun. Bor. Amer.,
Mamm., i, 1829, p. 172.

Spermophilus grammurus douglasi Elliot, Syn. N. Am. Mamm.,
F. C. M. Pub., ii, 1901, p. 89. Zoöl. Ser.

DOUGLAS' SPERMOPHILE.

Type locality. Plains of the Columbia River, Oregon.

Geogr. Distr. Northern California, Oregon.

h. *Citellus variegatus buckleyi* (Slack).

Spermophilus buckleyi Slack, Proc. Acad. Nat. Scien. Phil., 1861,
p. 314.

Spermophilus grammurus buckleyi Elliot, Syn. N. Am. Mamm.,
F. C. M. Pub., ii, 1901, p. 89. Zoöl. Ser.

BUCKLEY'S SPERMOPHILE.

Type locality. Pack-saddle Mountain, Llano County, Texas.

Geogr. Distr. Central Texas.

225. *Citellus nesioticus* Elliot.

Citellus nesioticus Elliot, Field Columb. Mus. Pub., iii, 1904, p.
263. Zoöl. Ser.

ISLAND SPERMOPHILE.

Type locality. Santa Catalina Island, California.

Geogr. Distr. Santa Catalina Island, California.

Genl. Char. Color dark; little or none of the yellowish brown hue so conspicuous in *C. v. beecheyi*. Skull shorter and broader; molar broad and heavy; nasals short; braincase short, contracted anteriorly; antorbital processes very long; pterygoid fossa very short, and processes of pterygoid very long and turning sharply outward at tips; basi-occipital short and broad; bullæ large; palate very broad;

zygomatic arch widely expanded; occipital region low and broad; azygos process on palatal arch.

Color. Top of head from forehead to between ears jet black mixed with a few tawny hairs; nose tawny ochraceous; sides of head mixed tawny ochraceous and black; back and sides black and tawny ochraceous, the black predominating; under parts buffy ochraceous; hands and feet buff, ankles tawny; limbs mixed buff and black; tail above mixed buff and black appearing gray with a black border and pale buff fringe, beneath mixed buff and black; ears black.

Measurements. Total length, 476.2; tail vertebræ, 184; hind foot, 60. Skull: occipito-nasal length, 60.5; zygomatic width, 39; inter-orbital constriction, 15.5; width of postorbital processes, 27; braincase above zygomata, 25.3; length of braincase, 31; median length of nasals, 19; lateral length of nasals, 20; length of pterygoids, 11; palatal length, 30; length of upper tooth row, 12; length of mandible, 36; length of lower tooth row, 11.

226. *Citellus annulatus* (Audubon and Bachman).

Spermophilus annulatus Aud. & Bachm., Jour. Acad. Nat. Scien. Phil., 1842, p. 319.

Citellus annulatus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 151. Zoöl. Ser.

RING-TAILED SPERMOPHILE.

Type locality. Western Mexico?

Geogr. Distr. Plains of State of Colima and Territory of Tepic, western Mexico; extent of range unknown.

a. *Citellus annulatus goldmani* (Merriam).

Spermophilus annulatus goldmani Merr., Proc. Biol. Soc. Wash., xv, 1902, p. 69.

Citellus annulatus goldmani Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 151. Zoöl. Ser.

GOLDMAN'S SPERMOPHILE.

Type locality. Santiago, Territory of Tepic, Mexico.

Geogr. Distr. Limits of range unknown.

227. *Citellus adocetus* Merriam.

Citellus adocetus Merr., Proc. Biol. Soc. Wash., xvi, 1903, p. 79.

Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 152. Zoöl. Ser.

PLAIN-TAILED SPERMOPHILE.

Type locality. La Salada, forty miles south of Uruapan, State of Michoacan, Mexico.

Geogr. Distr. State of Michoacan, Mexico.

Genus 53. **Cynomys.**

Cynomys Rafin., Am. Month. Mag., II, 1817, p. 45.. Type *Arctomys ludovicianus* Ord.

Anisonyx Rafin., Am. Month. Mag., II, 1817, p. 45. (nec Latreille, Coleopt., 1807.)

Monax Warden, Statist. Polit. Hist. Acc. U. S., I, 1819, pp. 225-228.

228. Cynomys ludovicianus (Ord).

Arctomys ludovicianus Ord, Guth. Geogr., 2d Am. ed., II, 1815, p. 292.

Cynomys socialis Rafin., Am. Month. Mag., II, 1817, p. 45.

Cynomys grisea Rafin., Am. Month. Mag., II, 1817, p. 45.

Arctomys missouriensis Warden, Desc. États Unis, v, 1820, p. 627.

Arctomys latrans Harlan, Faun. Am., 1825, p. 306.

Cynomys ludovicianus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 102, fig. 23. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 155, figs. 28, XXXII. Zoöl. Ser.

PRAIRIE-DOG. *Perrito*, *Perrito del Campo* in Mexico.

Type locality. Plains of the Upper Missouri.

Geogr. Distr. From 49th parallel southwest of the Mississippi through western Kansas and Texas into the State of Chihuahua, Mexico.

229. Cynomys pyrrotrichus Elliot.

Cynomys pyrrotrichus Elliot, Proc. Biol. Soc. Wash., XVIII, 1905, p. 139.

RED PRAIRIE-DOG.

Type locality. White Horse Spring, Oklahoma Territory.

Genl. Char. General color cinnamon rufous, altogether different from that of *C. ludovicianus*. Skull compared with that of above-named species has a narrower palate and larger bullæ; and with *arizonensis* has a broader braincase and wider nasals.

Color. Head, upper parts of body and sides cinnamon rufous, darkest on head and dorsal region, each hair tipped with whitish, most conspicuous on sides of rump and flanks; under parts and inner side of legs pale yellowish washed or tinged with rufous on chest and abdomen grading on sides into the color of the flanks; upper parts of fore legs and feet pale vinaceous cinnamon; outer side of thighs pale cinnamon rufous; feet pale vinaceous buff; claws black; tail cinnamon rufous, grading to Van Dyke brown at tips.

Measurements. Total length, 385; tail vertebræ, 70; hind foot, 58. Skull, total length, 63; Hensel, 52; zygomatic width, 45; greatest width of braincase, 25; length of nasals, 21; palatal length, 31.5; length of upper molar series, 13; length of mandible, angle to top of incisors, 18; of lower molar series, 16.

230. *Cynomys arizonensis* Mearns.

Cynomys arizonensis Mearns, Bull. Am. Mus. Nat. Hist. N. Y., 1890, p. 305. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 104. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 154. Zoöl. Ser.

ARIZONA PRAIRIE-DOG.

Type locality. Point of Mountain near Wilcox, Cochise County, Arizona.

Geogr. Distr. Arizona into State of Chihuahua.

231. *Cynomys gunnisoni* (Baird).

Spermophilus gunnisoni Baird, Proc. Acad. Nat. Scien. Phil., 1855, p. 334.

Cynomys gunnisoni Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 103. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 156. Zoöl. Ser.

GUNNISON'S PRAIRIE-DOG.

Type locality. Cochetopa Pass, Rocky Mountains, Saguache County, Colorado.

Geogr. Distr. Colorado, Arizona, and New Mexico, into State of Chihuahua, Mexico.

232. *Cynomys lewisi* (Audubon and Bachman).

Arctomys lewisii Aud. & Bachm., Quad. N. Am., III, 1853, p. 32, pl. cvii.

Cynomys leucurus Merr., N. Am. Faun., No. 4, 1890, p. 33.

Cynomys lewisi Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 104. Zoöl. Ser.

LEWIS' PRAIRIE-DOG.

Type locality. Fort Bridger, Sweetwater County, Wyoming.

Geogr. Distr. Wyoming, Utah in the Uncompahgre Indian Reservation.

233. *Cynomys parvidens* Allen.

Cynomys parvidens Allen, Bull. Brookl. Inst. Arts & Scien., 1, No. 5, 1905, p. 119.

UTAH PRAIRIE-DOG.

Type locality. Buckskin Valley, Iron County, Utah.

Geogr. Distr. Unknown.

Genl. Char. Two-thirds of tail white; dentition weak.

Color. Above dull rufous, hairs posteriorly tipped with dusky; top of head and cheeks mixed with black; black spot above eye; thighs and fore legs yellowish rufous; beneath yellowish buff; tail, basal third like body, remainder white.

Measurements. Hind foot (skin), 53; none other given.

234. *Cynomys mexicanus* Merriam.

Cynomys mexicanus Merr., Proc. Biol. Soc. Wash., VII, 1892, p.

157. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M.

Pub., IV, Pt. 1, 1904, p. 156. Zoöl. Ser.

MEXICAN PRAIRIE-DOG.

Type locality. La Ventura, State of Coahuila, Mexico.

Geogr. Distr. Northern and eastern Mexico.

Genus 54. **Marmota. Woodchucks.**

* **Marmota** Frisch, Natur-Syst. vierfüß. Thiere, in Tabellen, 9, 1775.

Type ———— ?

Mus Linn., Syst. Nat., I, 1758, p. 60. (Part.)

Glis Erxleb., Syst. Regn. Anim., 1777, p. 358. (Part. nec Brisson, 1762.)

Arctomys Schreb., Säugth., IV, 1780, pp. 721-743; pls. CCVII-CCXI.

Lipura Illig., Prodr. Syst. Mamm. et Av., 1811, p. 95.

235. *Marmota monax* (Linnæus).

Mus monax Linn., Syst. Nat., I, 1758, p. 60.

Arctomys monax Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 105, fig. 24. Zoöl. Ser.

WOODCHUCK.

Type locality. Maryland.

Geogr. Distr. From New York to Georgia, west to the Dakotas, intergrading in the Alleghanian and Canadian faunæ with next three species.

a. Marmota monax ignavus (Bangs).

Arctomys ignavus Bangs, Proc. N. Eng. Zoöl. Club, 1899, p. 13.

Arctomys monax ignavus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 105. Zoöl. Ser.

**Marmota* of Frisch includes the following species: *Mus alpinus*, *Marmota polonica*, *Cricetus*, Europe; and *Gerboa*, Africa.

LABRADOR WOODCHUCK.

Type locality. Black Bay, Straits of Belle Isle, Labrador.

Geogr. Distr. Labrador. Range unknown.

b. *Marmota monax canadensis* (Erxleben).

Glis canadensis Erxl., Syst. Regn. Anim., 1777, p. 363.

Mus empetra Pallas, Nov. Spec. Quad. Glir. Ord., 1778, p. 75.

Arctomys melanopus Kuhl, Beitr., 1820, p. 64.

Arctomys monax canadensis Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 106. Zoöl. Ser.

CANADIAN WOODCHUCK.

Type locality. Hudson Bay.

Geogr. Distr. Newfoundland to Rocky Mountains in Arctic America, and from 57° N. latitude to central New England, west to northeastern Minnesota.

236. *Marmota pruinosus* (Gmelin).

Arctomys pruinosus Gmel., Syst. Nat., 1, 1788, p. 144. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 106. Zoöl. Ser.

Arctomys caligatus Eschsch., Zool. Atlas, 11, 1829, pl. 6.

HOARY WOODCHUCK.

Type locality. Hudson Bay region?

Geogr. Distr. Columbia River northward to the Barren Grounds, Arctic America, east to Hudson Bay.

237. *Marmota flaviventer* (Audubon and Bachman).

Arctomys flaviventer Aud. & Bach., Proc. Acad. Nat. Scien. Phil., 1841, p. 99. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 106. Zoöl. Ser.

YELLOW-BELLIED WOODCHUCK.

Type locality. "Mountains between Texas and California."

Geogr. Distr. Western Texas, New Mexico, and Arizona north to 49th parallel.

a. *Marmota flaviventer avarus* (Bangs).

Arctomys flaviventer avarus Bangs, Proc. N. Eng. Zoöl. Club, 1899, p. 68. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, Suppl., p. 427. Zoöl. Ser.

GREEDY WOODCHUCK.

Type locality. Okanagan, British Columbia, Canada.

Geogr. Distr. British Columbia. Range unknown.

238. *Marmota dakota* (Merriam).

Arctomys dacota Merr., N. Am. Faun., No. 2, 1889, p. 8. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 106. Zoöl. Ser.

DAKOTA WOODCHUCK.

Type locality. Custer, Black Hills, South Dakota.

Geogr. Distr. South Dakota. Range unknown.

239. *Marmota engelhardti* Allen.

Marmota engelhardti Allen, Bull. Brookl. Inst. Arts & Scien., 1, No. 5, 1905, p. 120.

ENGELHARDT'S WOODCHUCK.

Type locality. Briggs Meadows, Beaver Range Mountains, Utah. Altitude, 10,000 feet.

Geogr. Distr. Unknown.

Genl. Char. Similar to *M. flaviventer* and *M. dakota*, but darker in color; palate more produced posteriorly; dentition weaker.

Color. Side of nose, chin, and anterior part of throat white; top of head black; upper parts, shoulders and thighs, grizzled gray and dusky brown; under fur blackish slate basally, tipped broadly with cinnamon; long hairs black tipped with white; orbital ring, and patch on side of neck, pale rusty yellow; ventral surface deep chestnut, basal third of hairs posteriorly blackish; upper surface of hind feet deep chestnut; tail long, proximal half varied with gray and rusty-tipped hairs; apical half black.

Measurements. (Dried skin.) Total length, 640; tail vertebrae, 160; hind foot, 70. Skull: total length, 74; zygomatic breadth, 48.3; length of nasals, 29; breadth of rostrum in front, 15.

240. *Marmota olympus* (Merriam).

Arctomys olympus Merr., Proc. Acad. Nat. Scien. Phil., 1898, p. 352. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 107. Zool. Ser.

OLYMPIC MOUNTAINS WOODCHUCK.

Type locality. Solduc Trail, timber line, Olympic Mountains, Clallam County, Washington.

Geogr. Distr. High Mountains, northwestern Washington.

Subfamily II. **Pteromyinæ.**Genus 55. **Sciuropterus. Flying Squirrels.**

Sciuropterus F. Cuv., Dents Mammifères, 1825, pp. 161-162, 255, pl. 56. Type *Sciurus volans* Linnæus.

241. *Sciuropterus alpinus* (Richardson).

Pteromys alpinus Rich., Zool. Journ., III, 1828, p. 519.

Pteromys sabrinus var. *β. alpinus* Richard., Faun. Bor. Amer., 1, Mamm., 1829, p. 195, pl. 16.

Sciuropterus volans alpinus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 109. Zoöl. Ser.

RICHARDSON'S FLYING SQUIRREL.

Type locality. Head waters of the Athabasca and Peace Rivers, Alberta, Canada.

Geogr. Distr. From the Mackenzie River along east side of the Rocky Mountains to the United States border.

a. *Sciuropterus alpinus zaphæus* Osgood.

Sciuropterus alpinus zaphæus Osgood, Proc. Biol. Soc. Wash., XVIII, 1905, p. 133.

DUSKY ALASKAN FLYING SQUIRREL.

Type locality. Helm Bay, Cleveland Peninsula, Alaska.

Geogr. Distr. Only known from the type locality.

Genl. Char. Similar to *S. alpinus*, but darker.

Color. Above pale russet; dusky line from base of whiskers to and around eye, thence gray to posterior base of ear; lateral line black; upper side of forearm blackish brown; sides of throat grayish; axillary region creamy buff; throat whitish; breast and middle of belly creamy buff; sides of belly and under side of hind legs pale creamy mixed with dusky; anal region pale russet mixed with dusky; hairs of tail above subterminally brocoli brown, overlaid with black; beneath more rufescent, with an indistinct grayish median line; upper side of fore and hind feet blackish brown.

Measurements. "Total length, 311; tail vertebræ, 133; hind foot, 42. Skull: basilar length of Hensel, 32.3; occipito-nasal length, 41.4; zygomatic breadth, 26.2; constriction in front of postorbital processes, 8.3; median length of nasals, 12.8; upper tooth row, 8.2."

b. *Sciuropterus alpinus californicus* Rhoads.

Sciuropterus alpinus californicus Rhoads, Proc. Acad. Nat. Scien. Phil., 1897, p. 323.

Sciuropterus volans californicus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 110. Zoöl. Ser.

CALIFORNIAN FLYING SQUIRREL.

Type locality. San Bernardino Mountains, southern California.

Geogr. Distr. Mountain ranges, southeastern California.

c. *Sciuropterus alpinus oregonensis* (Bachman).

Pteromys oregonensis Bachm., Jour. Acad. Nat. Scien. Phil., 1839, p. 101.

Sciuropterus volans oregonensis Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 110. Zoöl. Ser.

OREGON FLYING SQUIRREL.

Type locality. Probably near St. Helen, Columbia County, Oregon.

Geogr. Distr. Southern Alaska to northern California, at lower elevations on the Pacific slopes.

d. *Sciuropterus alpinus olympicus* Elliot.

Sciuropterus alpinus olympicus Elliot, Pub. Field Columb. Mus.

I, 1899, p. 225. Zoöl. Ser.

Sciuropterus volans olympicus Elliot, Syn. N. Am. Mamm., F. C. M.

Pub., II, 1901, p. 110. Zoöl. Ser.

OLYMPIC MOUNTAINS FLYING SQUIRREL.

Type locality. Happy Lake, Clallam County, Olympic Mountains, Washington.

Geogr. Distr. Higher parts of the Olympic Mountains, northwestern Washington.

e. *Sciuropterus alpinus fuliginosus* Rhoads.

Sciuropterus alpinus fuliginosus Rhoads, Proc. Acad. Nat. Scien.

Phil., 1897, p. 32.

Sciuropterus volans fuliginosus Elliot, Syn. N. Am. Mamm.,

F. C. M. Pub., II, 1901, p. 111. Zoöl. Ser.

DUSKY FLYING SQUIRREL.

Type locality. Martin Station, Kittitass County, Cascade Mountains, Washington.

Geogr. Distr. Cascade, Coast, and Sierra Nevada Ranges, at high elevations.

f. *Sciuropterus alpinus klamathensis* Merriam.

Sciuropterus alpinus klamathensis Merr., Proc. Biol. Soc. Wash.,

XI, 1897, p. 225.

Sciuropterus volans klamathensis Elliot, Syn. N. Am. Mamm.,

F. C. M. Pub., II, 1901, p. 111. Zoöl. Ser.

KLAMATH FLYING SQUIRREL.

Type locality. Fort Klamath, Klamath County, Oregon. Altitude, 4,200 feet.

Geogr. Distr. Oregon. Range unknown.

g. *Sciuropterus alpinus lascivus* Bangs.

Sciuropterus alpinus lascivus Bangs, Proc. N. Engl. Zoöl. Club, I,

1899, p. 69. Elliot, Syn. N. Am. Mamm., F. C. M. Pub.,

II, 1901, Suppl., p. 428. Zoöl. Ser.

FRISKY FLYING SQUIRREL.

Type locality. Tallac, El Dorado County, California.

Geogr. Distr. California. Range unknown.

h. Sciuropterus alpinus stephensi (Merriam).

Sciuropterus oregonensis stephensi Merr., Proc. Biol. Soc. Wash., XIII, 1900, p. 151.

Sciuropterus stephensi Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 111. Zoöl. Ser.

STEPHENS' FLYING SQUIRREL.

Type locality. Sherwood, Mendocino County, California. Altitude, 2,500 feet.

Geogr. Distr. Western California.

i. Sciuropterus alpinus bangsi Rhoads.

Sciuropterus alpinus bangsi Rhoads, Proc. Acad. Nat. Scien. Phil., 1897, p. 321. (Footnote.)

BANGS' FLYING SQUIRREL.

Type locality. Idaho County, Idaho.

Geogr. Distr. Idaho.

Genl. Char. Smaller and darker than *S. alpinus*.

Color. Sides of face, lips, eyelids, and ears smoky gray shaded with brown; upper parts of body wood brown; underparts tinged with wood brown, darkest on sides of abdomen; tail heavily shaded with black above and below; hands and feet grayish plumbaceous.

Measurements. Total length, 304; tail vertebræ, 146; hind foot, 40; ear from crown, 16.

242. Sciuropterus sabrinus (Shaw).

Sciurus sabrinus Shaw, Gen. Zoöl., I, 1801, p. 157.

Sciurus hudsonius Gmel., Syst. Nat., I, 1788, p. 153. (nec Pallas.)

Sciurus acrobates Wagn., Schreb. Säugth., Suppl., IV, 1843, p. 231, pl. CCXXII, B.

Sciuropterus sabrinus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 107. Zoöl. Ser.

BOREAL FLYING SQUIRREL.

Type locality. Severn River, Keewatin, Canada.

Geogr. Distr. Eastern boreal North America, south to northern New York and southern New Hampshire.

a. Sciuropterus sabrinus macrotis Mearns.

Sciuropterus sabrinus macrotis Mearns, Proc. U. S. Nat. Mus., XXI, 1899, (1898), p. 353, figs. 1, 2, 3. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 108. Zoöl. Ser.

LARGE-EARED FLYING SQUIRREL.

Type locality. Hunter Mountain, Catskill Mountains, Greene County, New York.

Geogr. Distr. Northern border of the United States, east of the Great Lakes.

b. Sciuropterus sabrinus makkovikensis Sornborger.

Sciuropterus sabrinus makkovikensis Sornborger, Ottawa Nat., XIV, 1900, p. 48.

MAKKOVIK FLYING SQUIRREL.

Type locality. Makkovik, Labrador.

Geogr. Distr. Labrador. Range unknown.

Genl. Char. Size large, color dark.

Color. Similar to *S. sabrinus* but darker above, beneath whitish; tail sooty, basal half darkest.

Measurements. Total length, —; tail vertebræ, —; hind foot, 40.5-42. Skull: occipito-nasal length, 35-41.4; Hensel, 27.9-33; zygomatic width, 21-26.2; interorbital constriction, 7.1-8.9; length of mandible, 21-25.9; length of lower molar series, 6.7-8.2.

c. Sciuropterus sabrinus silus (Bangs).

Sciuropterus silus Bangs, Proc. Biol. Soc. Wash., x, 1896, p. 163.

Sciuropterus sabrinus silus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 108. Zoöl. Ser.

PUG-NOSED FLYING SQUIRREL.

Type locality. Katis Mountain, White Sulphur Springs, Greenbrier County, West Virginia.

Geogr. Distr. Probably the higher southern Alleghanies.

243. *Sciuropterus yukonensis* Osgood.

Sciuropterus yukonensis Osgood, N. Am. Faun., No. 19, 1900, p. 25. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, Suppl., p. 428. Zoöl. Ser.

YUKON FLYING SQUIRREL.

Type locality. Camp Davidson, Yukon River near Alaska, Canada boundary, Yukon, Canada.

Geogr. Distr. Possibly from Chilkoot Inlet, Alaska, north to Canada boundary.

244. *Sciuropterus volans* (Linnæus).

Mus volans Linn., Syst Nat., I, 1758, p. 63; I, 1766, p. 85. (*Mus nec Sciurus*, p. 88.)

Sciurus volucella Pallas, Nov. Spec. Quad. Glir., 1778, p. 351.

Sciuropterus volans Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 109, fig. 25. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 158, figs. 29, xxxiii. Zoöl. Ser.

COMMON FLYING SQUIRREL.

Type locality. Virginia.

Geogr. Distr. Northern New York and southern New Hampshire south to Florida, west to the plains.

a. *Sciuropterus volans querceti* Bangs.

Sciuropterus volans querceti Bangs, Proc. Biol. Soc. Wash., x, 1896, p. 166. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 109. Zoöl. Ser.

FLORIDA FLYING SQUIRREL.

Type locality. Citronelle, Citrus County, Florida.

Geogr. Distr. Southern Georgia and Florida, west to Louisiana?

b. *Sciuropterus volans goldmani* Nelson.

Sciuropterus volans goldmani Nelson, Proc. Biol. Soc. Wash., xvii, 1904, p. 148.

MEXICAN FLYING SQUIRREL.

Type locality. Twenty miles southeast of Teopisca, State of Chiapas, Mexico.

Geogr. Distr. Highlands of the State of Chiapas, Mexico, and Guatemala.

Genl. Char. Like *S. v. querceti*; top of nose white; membrane beneath ochraceous buff.

Color. Upper parts reddish brown; top and side of nose, lower part of cheeks and sides of neck to back of ears whitish; between eye and ear dusky, lower part of cheeks and sides of head below ears dingy grayish brown; supraloral spot whitish; underside of neck and body white tinted with buff; upper surface of membrane blackish slate color. under surface deep ochraceous buff; tail above cinnamon brown, beneath dingy buff; top of fore feet dingy whitish; top of hind feet dusky, toes dingy white.

Measurements. Total length, 237; tail vertebræ, 112; hind foot, 30. Skull: Hensel, 28; interorbital width, 7.5; zygomatic width, 22; greatest width of braincase, 17.5; length of nasals, 10.2; width of rostrum, 6; depth of rostrum, 7.

Family II. **Haplodontidæ. Sewellels.**

Genus 56. **Haplodontia.**

Aplodontia (*sic*) Rich., Zoöl. Journ., iv, 1829, p. 334. Type *Aplodontia! leporina* Richardson = *Anisonyx rufa* Rafinesque.

Anisonyx Rafin., Am. Month. Mag., II, 1817, p. 45. (Part.)

245. Haplodontia rufa (Rafinesque).

Anisonyx rufa Rafin., Am. Month. Mag., II, 1817, p. 45.

Aplodontia! leporina Rich., Faun. Bor. Amer., Mamm., I, 1829, p. 211, pl. 18 C.

Haplodontia rufa Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 112. Zoöl. Ser.

RUFIOUS SEWELLEL.

Type locality. Cascades of the Columbia River, Oregon.

Geogr. Distr. Eastern Washington and Oregon.

a. *Haplodontia rufa californica* Peters.

Haplodon californica Peters, Monatsb. K. Preus. Akad. Wiss. Berl., 1864, p. 179.

Aploadontia! major Merr., Ann. Acad. Scien N. Y., III, 1886, p. 316.

Haplodontia rufa californica Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 112, fig. 26. Zoöl. Ser.

CALIFORNIA SEWELLEL.

Type locality. Sierra Nevada Range, Placer County, California.

Geogr. Distr. California. Range unknown.

b. *Haplodontia rufa rainieri* (Merriam).

Aploadontia! major rainieri Merr., Proc. Biol. Soc. Wash., XIII, 1899, p. 21.

Haplodontia rufa rainieri Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 112. Zoöl. Ser.

MOUNT RAINIER SEWELLEL.

Type locality. Paradise Creek, south of Mount Rainier, Lewis County, Washington. Altitude, 5,200 feet.

Geogr. Distr. Washington. Range unknown.

246. *Haplodontia pacifica* (Merriam).

Aploadontia! pacifica Merr., Proc. Biol. Soc. Wash., XIII, 1899, p. 19.

Haplodontia pacifica Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 114. Zoöl. Ser.

PACIFIC SEWELLEL.

Type locality. Newport, Yaquina Bay, Lincoln County, Oregon.

Geogr. Distr. Oregon. Range unknown.

247. *Haplodontia phæa* (Merriam).

Aploadontia! phæa Merr., Proc. Biol. Soc. Wash., XIII, 1899, p. 20.

Haplodontia phæa Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 114. Zoöl. Ser.

DARK SEWELLEL.

Type locality. Point Reyes, Marin County, California.

Geogr. Distr. Western California. Range unknown.

248. *Haplodontia olympica* (Merriam).

Aploadontia! olympica Merr., Proc. Biol. Soc. Wash., XIII, 1899, p. 20.

Haplodontia olympica Elliot, Syn. N. Am. Mamm., F. C. M. Pub.,
II, 1901, p. 114. Zoöl. Ser.

OLYMPIC MOUNTAINS SEWELLEL. *Farmer*.

Type locality. Queniult Lake, Chehalis County, Olympic Mountains, Washington.

Geogr. Distr. Olympic Mountains, Washington. Range undetermined.

Family III. **Castoridae. Beavers.**

Genus 57. **Castor.**

Castor Linn., Syst. Nat., I, 1758, p. 58; I, 1766, p. 78. Type *Castor fiber* Linnæus.

249. **Castor canadensis** Kuhl.

Castor canadensis Kuhl, Beit. Zool., 1820, p. 64. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 116, fig. 27. Zoöl. Ser.

CANADA BEAVER.

Type locality. Hudson Bay.

Geogr. Distr. Northeastern North America from northern limit of trees to United States, and west to Cascade Mountains.

a. *Castor canadensis carolinensis* Rhoads.

Castor canadensis carolinensis Rhoads, Trans. Am. Philos. Soc., 1898, p. 420. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 116. Zoöl. Ser.

CAROLINA BEAVER.

Type locality. Dan River, near Danbury, Stokes County, North Carolina.

Geogr. Distr. North Carolina west to Louisiana.

b. *Castor canadensis frondator* Mearns.

Castor canadensis frondator Mearns, Proc. U. S. Nat. Mus., 1897, p. 502. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 116. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 159, figs. 30, xxxiv. Zoöl. Ser.

SONORAN BEAVER.

Type locality. San Pedro River, State of Sonora, Mexico, near monument No. 98 of the Mexican boundary line.

Geogr. Distr. From State of Sonora, Mexico, to Wyoming and Montana, intergrading into the other forms wherever the limits of their dispersion meet.

c. Castor canadensis pacificus Rhoads.

Castor canadensis pacificus Rhoads, Trans. Am. Philos., 1898, p. 422. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 117. Zoöl. Ser.

PACIFIC COAST BEAVER.

Type locality. Lake Kichelos, Cascade Mountains, Kittitass County, Washington.

Geogr. Distr. Pacific Slope; California to Alaska.

Family IV. **Muridæ. Rats, Mice, Voles.**

Subfamily I. **Murinæ.**

Genus 58. **Mus.**

Mus Linn., Syst. Nat., I, 1758, p. 59; I, 1766, p. 79. Type *Mus rattus* Linnaeus.

250. Mus rattus Linnaeus.

Mus rattus Linn., Syst. Nat., I, 1758, p. 61; I, 1766, p. 83. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 117, fig. 28. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 163, figs. 31, xxxv. Zoöl. Ser.

BLACK RAT.

Type locality. Sweden.

Geogr. Distr. Cosmopolitan.

251. Mus norvegicus Erxleben.

Mus norvegicus Erxl., Syst. Regn. Anim., 1777, p. 381. Elliot, Syn. N. Am. Mamm., II, 1901, Append., p. 428. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 164. Zoöl. Ser.

Mus decumanus Pall., Nov. Sp. Quad. Glir., 1778, p. 91. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 117. Zoöl. Ser.

NORWAY RAT.

Type locality. Norway.

Geogr. Distr. Cosmopolitan.

252. Mus alexandrinus I. Geoffroy.

Mus alexandrinus I. Geoff., Descr. Égypte, II, 1813, p. 733; Atl. pl. v, fig. 1. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 118. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 164. Zoöl. Ser.

Mus tectorum Savi, Nov. Giorn. di Lett., 1825.

BROWN RAT.

Type locality. Alexandria, Egypt.

Geogr. Distr. Cosmopolitan.

253. *Mus musculus* Linnæus.

Mus musculus Linn., Syst. Nat., 1, 1758, p. 62; 1, 1766, p. 83.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 118.

Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M.

Pub., 1V, Pt. 1, 1904, p. 162. Zoöl. Ser.

HOUSE MOUSE.

Type locality. Sweden.

Geogr. Distr. Cosmopolitan.

a. *Mus musculus jalapæ* Allen and Chapman.

Mus musculus jalapæ Allen and Chapman, Bull. Am. Mus. Nat.

Hist. N. Y., 1897, p. 198. Elliot, Mamm. Middle Amer. &

W. Indies, F. C. M. Pub., 1V, Pt. 1, 1904, p. 163. Zoöl. Ser.

JALAPA HOUSE MOUSE.

Type locality. Jalapa, State of Vera Cruz, Mexico.

Geogr. Distr. State of Vera Cruz, Mexico. Range unknown.

Subfamily II. *Cricetinae*.Genus 59. *Onychomys*.

Onychomys Baird, N. Am. Mamm., 1857, p. 458. Type *Hypudæus leucogaster* Maximilian.

Hypudæus Max., Reise N. Am., 11, 1841, p. 99. (nec Auct.)

254. *Onychomys leucogaster* (Maximilian).

Hypudæus leucogaster Max., Reise N. Am., 11, 1841, p. 99.

Mus missouriensis Aud. & Bach., 11, 1851, p. 327, pl. 100.

Onychomys leucogaster Elliot, Syn. N. Am. Mamm., F. C. M.

Pub., 11, 1901, p. 119, fig. 29. Zoöl. Ser.

WHITE-BELLIED MOLE MOUSE.

Type locality. Clark, Clark County, South Dakota.

Geogr. Distr. South Dakota, Missouri, Nebraska, Kansas, Oklahoma and Indian Territories.

a. *Onychomys leucogaster breviceaudus* Merriam.

Onychomys leucogaster breviceaudus Merr., N. Am. Faun., No. 5, Montan., 1891, p. 52. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., their dispers., 1901, p. 120. Zoöl. Ser.

SHORT-TAILED MOLE MOUSE.

Type locality. Blackfoot, Bingham County, Idaho.

Geogr. Distr. Idaho. Range unknown.

b. *Onychomys leucogaster longipes* (Merriam).

Onychomys longipes Merr., N. Am. Faun., No. 2, 1889, p. 1.

Onychomys leucogaster longipes Elliot, Syn. N. Am. Mamm.,

F. C. M. Pub., II, 1901, p. 120. Zool. Ser.

LONG-FOOTED MOLE MOUSE.

Type locality. Concho County, Texas.

Geogr. Distr. West central Texas.

c. *Onychomys leucogaster albescens* Merriam.

Onychomys leucogaster albescens Merr., Proc. Biol. Soc. Wash.,

XVII, 1904, p. 124. Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., IV, Pt. II, 1904, Append., p. 742. Zool. Ser.

SAMALAYUCA MOLE MOUSE.

Type locality. Samalayuca, State of Chihuahua, Mexico.

Geogr. Distr. State of Chihuahua, Mexico. Range unknown.

255. *Onychomys melanophrys* (Merriam).

Onychomys leucogaster melanophrys Merr., N. Am. Faun., No. 2,

1889, p. 2. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II,

1901, p. 120. Zool. Ser.

Onychomys melanophrys Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., IV, Pt. I, 1904, p. 166. Zool. Ser.

BLACK-BROWED MOLE MOUSE.

Type locality. Kanab, Kane County, Utah.

Geogr. Distr. Utah. Range unknown.

a. *Onychomys melanophrys pallescens* Merriam.

Onychomys melanophrys pallescens Merr., N. Am. Faun., No. 3,

1890, p. 61. Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., IV, Pt. I, 1904, p. 166. Zool. Ser.

Onychomys leucogaster pallescens Elliot, Syn. N. Am. Mamm.,

F. C. M. Pub., II, 1901, p. 121. Zool. Ser.

PALE MOLE MOUSE.

Type locality. Moki Pueblos, Apache County, Arizona.

Geogr. Distr. Northeastern Arizona.

256. *Onychomys ruidosæ* Stone & Rehn.

Onychomys ruidosæ Stone & Rehn, Proc. Acad. Nat. Scien. Phil.,

1903, p. 22.

RUIDOSO MOLE MOUSE.

Type locality. Ruidoso, Lincoln County, New Mexico.

Geogr. Distr. New Mexico. Range unknown.

Genl. Char. Nearest to *O. melanophrys*, but redder; head and ears darker; pterygoid fossa narrower.

Color. Above dull ferrugineous, rump clear rufous; head and nape blackish gray washed with ferrugineous; ears blackish gray mixed with silvery hairs, most numerous towards tip; chin pale grayish pink; under parts and limbs dull white; tail mixed brown and white above.

Measurements. Total length, 156; tail vertebræ, 47; hind foot, 22.

257. *Onychomys arcticeps* Rhoads.

Onychomys arcticeps Rhoads, Proc. Acad. Nat. Scien. Phil., 1898, p. 194. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 121. Zoöl. Ser.

RHOADS' MOLE MOUSE.

Type locality. Clapham, Union County, New Mexico.

Geogr. Distr. Northeastern New Mexico.

258. *Onychomys longicaudus* Merriam.

Onychomys longicaudus Merr., N. Am. Faun., No. 2, 1889, p. 2.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 121.

Zoöl. Ser.

LONG-TAILED MOLE MOUSE.

Type locality. St. George, Washington County, Utah.

Geogr. Distr. Utah. Range unknown.

a. *Onychomys longicaudus tularensis* (Merriam).

Onychomys torridus tularensis Merr., Proc. Biol. Soc. Wash., XVII, 1904, p. 123.

TULARE MOLE MOUSE.

Type locality. Bakersfield, Kern County, California.

Geogr. Distr. Tulare Basin from Huron and Alcade south, the Carrizo Plains on the west, and Kern Valley on the east.

Genl. Char. Similar to *O. longicaudus*, but paler and without ochraceous suffusion; size small.

Color. Pale drab gray tinged with buff.

Measurements. Average of 5 specimens: Total length, 143; tail vertebræ, 50; hind foot, 21.

259. *Onychomys ramona* Rhoads.

Onychomys ramona Rhoads, Amer. Nat., 1893, p. 833. Elliot,

Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 121. Zoöl.

Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub.,

IV, Pt. I, 1904, p. 167. Zoöl. Ser.

RAMONA MOLE MOUSE.

Type locality. San Bernardino Valley, California.

Geogr. Distr. Southeastern California.

260. *Onychomys torridus* (Coues).

Hesperomys (Onychomys) torridus Coues, Proc. Acad. Nat. Scien. Phil., 1874, p. 183.

Onychomys torridus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 122. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 167, figs. 32, xxxvi. Zoöl. Ser.

TROPICAL MOLE MOUSE.

Type locality. Camp Grant, Graham County, Arizona.

Geogr. Distr. Between the Mississippi River and the Rocky Mountains. Upper Missouri, Red River of the North, Wyoming, Kansas, and Arizona.

a. *Onychomys torridus yakiensis* Merriam.

Onychomys torridus yakiensis Merr., Proc. Biol. Soc. Wash., xvii, 1904, p. 124. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, Append., p. 742. Zoöl. Ser.

YAKI MOLE MOUSE.

Type locality. Camoa Rio Mayo, State of Sonora, Mexico.

Geogr. Distr. Western part of the State of Sonora, and northern part of the State of Sinaloa, Mexico.

b. *Onychomys torridus canus* Merriam.

Onychomys torridus canus Merr., Proc. Biol. Soc. Wash., xvii, 1904, p. 124. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, Append., p. 742. Zoöl. Ser.

HOARY MOLE MOUSE.

Type locality. San Juan Capistrano, State of Zacatecas, Mexico.

Geogr. Distr. States of Zacatecas and San Luis Potosi, Mexico.

c. *Onychomys torridus arenicola* Mearns.

Onychomys torridus arenicola Mearns, Proc. U. S. Nat. Mus., 1897, p. 139. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 122. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 168. Zoöl. Ser.

SAND-LOVING MOLE MOUSE.

Type locality. Rio Grande, near El Paso, El Paso County, Texas.

Geogr. Distr. Texas into State of Chihuahua, Mexico.

d. *Onychomys torridus perpallidus* Mearns.

Onychomys torridus perpallidus Mearns, Proc. U. S. Nat. Mus., 1897, p. 140. Elliot, Syn. N. Am. Mamm., F. C. M. Pub.,

II, 1901, p. 122. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 168. Zoöl. Ser.

DRAB-GRAY MOLE MOUSE.

Type locality. Colorado River at Monument No. 204, Mexican boundary line, Yuma County, Arizona.

Geogr. Distr. Gila City, across Yuma and Colorado Deserts to the Coast Range.

261. *Onychomys fuliginosus* Merriam.

Onychomys fuliginosus Merr., N. Am. Faun., No. 3, 1890, p. 60.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 123. Zoöl. Ser.

DUSKY MOLE MOUSE.

Type locality. Black Tank lava beds, northeast of San Francisco Mountain, Coconino County, Arizona.

Geogr. Distr. Arizona. Range unknown.

262. *Onychomys macrotis* Elliot.

Onychomys macrotis Elliot, Pub. Field Columb. Mus., III, 1903, p. 155. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 169. Zoöl. Ser.

LARGE-EARED MOLE MOUSE.

Type locality. Head of San Antonio River, Lower California, Mexico.

Geogr. Distr. San Quentin to San Antonio River, Trinidad Valley and plain of El Alamo to about 5,000 feet elevation in the San Pedro Martir Mountains, Lower California, Mexico.

263. *Onychomys pulcher* Elliot.

Onychomys pulcher Elliot, Pub. Field Columb. Mus., III, 1903, p. 243. Zoöl. Ser.

BEAUTIFUL MOLE MOUSE.

Type locality. Morongo Pass, San Bernardino Mountains, Riverside County, California.

Genl. Char. Size medium; color pale.

Color. Upper parts buff inclining to pinkish, darker on rump, where the tint becomes almost a salmon buff; nose, sides of face, lips, entire under parts, legs and feet pure white; tail above soiled white; sides and under parts white; ears whitish at base, probably flesh color in life, apical half brownish black. A tuft of whitish hairs covers the base of the ear. Orbital ring black.

Measurements. Total length, 150; tail vertebrae, 55; hind foot, 21; ear, 18.5. Skull: total length, 25.5; Hensel, 20; zygomatic width, 13.5; interorbital constriction, 5; length of nasals, 9; palatal length,

10; length of upper tooth row, 4; length of mandible, 15; length of lower tooth row, 4.

Genus 60. **Peromyscus.**

Peromyscus Gloger, Hand- u. Hilfsb. Naturg., 1, 1841, pp. xxx, 95.

Type *Peromyscus arboreus* Gloger = *Mus sylvaticus noveboracensis* Fischer.

Calomys Aud. & Bach., Quad. N. Am., 11, 1851, p. 303. (nec Waterhouse, 1837.)

Sitomys Fitzin., Sitzungsab. Math.-Nat. Cl. K. Akad. Wiss. Wien, LVI, 1867, p. 97.

**Vesperimus* Coues, Proc. Acad. Nat. Scien. Phil., 1874, p. 178.

Baiomys True, Proc. U. S. Nat. Mus., xvi, 1894, p. 758.

Trinodontomys Rhoads, Proc. Acad. Nat. Scien. Phil., 1894, p. 257.

Haplomyiomys Osgood, Proc. Biol. Soc. Wash., xvii, 1904, p. 53.

Subgenus A. **Baiomys.**

"Size very small. Ascending ramus of mandible short and erect. Condyle terminal coronoid process well developed, uncinate, and near condyle. Plantar tubercles six; soles hairy. Tail short."

264. Peromyscus taylori (Thomas).

Hesperomys taylori Thomas, Ann. Mag. Nat. Hist., 5th Ser., xix, 1887, p. 66.

Peromyscus taylori Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 141. Zool. Ser.

TAYLOR'S MOUSE.

Type locality. San Diego, Duval County, Texas.

Geogr. Distr. Southern Texas.

265. Peromyscus allex Osgood.

Peromyscus allex Osgood, Proc. Biol. Soc. Wash., xvii, 1904, p. 76. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 175. Zool. Ser.

DWARF MOUSE.

Type locality. Colima, State of Colima, Mexico.

Geogr. Distr. Unknown.

266. Peromyscus musculus (Merriam).

Sitomys musculus Merr., Proc. Biol. Soc. Wash., viii, 1892, p. 170.

*A subgenus of *Hesperomys* Coues. (nec Waterhouse.)

Peromyscus musculus Elliot, Mamm. Middle Amer. & W. Indies,
F. C. M. Pub., IV, Pt. 1, 1904, p. 175. Zoöl. Ser.

COLIMA CITY MOUSE.

Type locality. Near Colima City, State of Colima, Mexico.

Geogr. Distr. States of Colima and Jalisco, Mexico.

a. *Peromyscus musculus brunneus* Allen & Chapman.

Peromyscus musculus brunneus Allen & Chapman, Bull. Am. Mus.

Nat. Hist. N. Y., 1897, p. 203. Elliot, Mamm. Middle Amer.

& W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 176. Zoöl. Ser.

JALAPA BROWN MOUSE.

Type locality. Jalapa, State of Vera Cruz, Mexico.

Geogr. Distr. State of Vera Cruz. Range unknown.

b. *Peromyscus musculus nigrescens* Osgood.

Peromyscus musculus nigrescens Osgood, Proc. Biol. Soc. Wash.,

XVII, 1904, p. 76. Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., IV, Pt. 1, 1904, p. 176. Zoöl. Ser.

SOOTY MOUSE.

Type locality. Valley of Comitán, State of Chiapas, Mexico.

Geogr. Distr. State of Chiapas. Range unknown.

267. *Peromyscus paulus* Allen.

Peromyscus paulus Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1903,

p. 598. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M.

Pub., IV, Pt. 1, 1904, p. 176. Zoöl. Ser.

TINY MOUSE.

Type locality. Rio Sestín, State of Durango, Mexico.

Geogr. Distr. Known only from the type locality.

Subgenus B. *Haplomylomys*.

"Size medium or small; pelage soft and silky; tail longer than head and body, subterete, rather thinly haired; soles of hind feet naked (in median line) to calcaneum, six tuberculate and paved with minute imbricate scales; skull with cranium rather large and rostral region relatively weak; first and second upper molars with three salient and two reëntrant outer angles at all stages of wear; small secondary tubercles never present between outer primary tubercles; lower molars correspondingly simple."

268. *Peromyscus eremicus* (Baird).

Hesperomys eremicus Baird, N. Am. Mamm., 1857, p. 479.

Peromyscus eremicus Elliot, Syn. N. Am. Mamm., F. C. M. Pub.,

II, 1901, p. 136. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W.

Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 191. Zoöl. Ser.

DESERT MOUSE.

Type locality. Old Fort Yuma, San Diego County, California.

Geogr. Distr. New Mexico, Arizona, and California, into State of Sonora and Lower California, Mexico.

a. *Peromyscus eremicus arenarius* Mearns.

Peromyscus eremicus arenarius Mearns, Proc. U. S. Nat. Mus., 1896, p. 138. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 136. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 191.

SAND-LOVING MOUSE.

Type locality. Rio Grande near El Paso, El Paso County, Texas.

Geogr. Distr. Southern Texas and State of Chihuahua, Mexico.

b. *Peromyscus eremicus phæurus* Osgood.

Peromyscus eremicus phæurus Osgood, Proc. Biol. Soc. Wash., XVII, 1904, p. 75. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 191. Zoöl. Ser.

BLACK-TAILED MOUSE.

Type locality. Hacienda La Parada, State of San Luis Potosi, Mexico.

Geogr. Distr. Middle portion of tableland in the States of San Luis Potosi, Zacatecas, and Nuevo Leon, Mexico.

269. *Peromyscus propinquus* (Allen).

Peromyscus eremicus propinquus Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1898, p. 154.

Peromyscus propinquus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 185. Zoöl. Ser.

ALLIED DESERT MOUSE.

Type locality. San Pablo Point, San Pablo Bay, Lower California, Mexico.

Geogr. Distr. Lower California, Mexico.

270. *Peromyscus anthonyi* (Merriam).

Hesperomys anthonyi Merr., Proc. Biol. Soc. Wash., IV, 1887, p. 5.

Peromyscus anthonyi Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 141. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 185. Zoöl. Ser.

ANTHONY'S MOUSE.

Type locality. Camp Apache, Grant County, New Mexico.

Geogr. Distr. New Mexico and Arizona.

271. *Peromyscus californicus* (Gambel).

Mus californicus Gambel, Proc. Acad. Nat. Scien. Phil., 1848, p. 78.

Peromyscus californicus Elliot, Syn. N. Am. Mamm., F. C. M. Pub.,
 II, 1901, p. 137. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W.
 Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 203. Zoöl. Ser.

CALIFORNIA MOUSE.

Type locality. Monterey, Monterey County, California.

Geogr. Distr. Coast region of California.

272. *Peromyscus goldmani* Osgood.

Peromyscus goldmani Osgood, Proc. Biol. Soc. Wash., xvii, 1904,
 p. 75. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M.
 Pub., IV, Pt. I, 1904, p. 186. Zoöl. Ser.

GOLDMAN'S MOUSE.

Type locality. Alamos, State of Sonora, Mexico.

Geogr. Distr. Known only from the type locality.

Subgenus C. *Peromyscus*.

273. *Peromyscus metallicola* Elliot.

Peromyscus metallicola Elliot, Pub. Field Columb. Mus., III, 1903,
 p. 245. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies,
 F. C. M. Pub., IV, Pt. I, 1904, p. 192. Zoöl. Ser.

MINE-HAUNTING MOUSE.

Type locality. Providentia Mines, northwestern part of State of
 Sonora, Mexico.

Geogr. Distr. Known only from the type locality.

274. *Peromyscus niveiventris* (Chapman).

Hesperomys niveiventris Chapman, Bull. Am. Mus. Nat. Hist.
 N. Y., 1889, p. 117.

Peromyscus niveiventris Elliot, Syn. N. Am. Mamm., F. C. M. Pub.,
 II, 1901, p. 141. Zoöl. Ser.

WHITE-BELLIED MOUSE.

Type locality. Opposite Micco, Brevard County, Florida.

Geogr. Distr. Eastern part of the peninsula of Florida.

a. Peromyscus niveiventris phasma (Bangs).

Peromyscus phasma Bangs, Proc. Bost. Soc. Nat. Hist., 1898,
 p. 199. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901,
 p. 142. Zoöl. Ser.

Peromyscus niveiventris phasma Elliot, Pub. Field Columb. Mus.,
 III, 1901, p. 41. Zoöl. Ser. (June.)

ANASTASIA ISLAND MOUSE.

Type locality. Point Romo, Anastasia Island, St. John County;
 Florida.

Geogr. Distr. Known only from Anastasia Island.

275. *Peromyscus subgriseus* (Chapman).

Sitomys subgriseus Chapman, Bull. Am. Mus. Nat. Hist. N. Y., 1893, p. 340.

Peromyscus subgriseus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 142. Zoöl. Ser.

CHAPMAN'S MOUSE.

Type locality. Gainesville, Alachua County, Florida.

Geogr. Distr. Central Florida.

a. Peromyscus subgriseus rhoadsi Bangs.

Peromyscus subgriseus rhoadsi Bangs, Proc. Bost. Soc. Nat. Hist., 1898, p. 201. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 142. Zoöl. Ser.

RHOADS' MOUSE.

Type locality. Anclote River, Hillsboro County, Florida.

Geogr. Distr. Southwestern portion of the peninsula of Florida.

b. Peromyscus subgriseus baliolus Bangs.

Peromyscus subgriseus arenarius Bangs, Proc. Bost. Soc. Nat. Hist., 1898, p. 202. (nec Mearns.)

Peromyscus subgriseus baliolus Bangs, Science, N. Ser., VIII, 1898, p. 214. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 142. Zoöl. Ser.

SWARTHY MOUSE.

Type locality. Hursman's Lake, Savannah River, near Bascom, Scriven County, Georgia.

Geogr. Distr. Sand hill region of northern Georgia and southern South Carolina.

276. *Peromyscus hemionotis* Elliot.

Peromyscus hemionotis Elliot, Pub. Field Columb. Mus., III, 1903, p. 157. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 199. Zoöl. Ser.

MULE-EARED MOUSE.

Type locality. Rosarito Divide, San Pedro Martir Mountains, Lower California, Mexico.

Geogr. Distr. Known only from the type locality.

277. *Peromyscus oresterus* Elliot.

Peromyscus oresterus Elliot, Pub. Field Columb. Mus., III, 1903, p. 159. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 200. Zoöl. Ser.

MOUNTAIN-LOVING MOUSE.

Type locality. Vallecitos, San Pedro Martir Mountains, Lower California, Mexico. Altitude, 9,000 feet.

Geogr. Distr. San Pedro Martir and Hanson Laguna Ranges, Lower California, Mexico.

278. *Peromyscus leucopus* (Rafinesque).

Musculus leucopus Rafin., Am. Month. Mag., III, 1818, p. 446.

Mus agrarius americanus Kerr, Anim. Kingd., I, 1792, p. 231.

Peromyscus americanus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 124, fig. 30. Zoöl. Ser.

Peromyscus leucopus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, Suppl., p. 492. Zoöl. Ser.

WHITE-FOOTED DEER MOUSE.

Type locality. Pine Barrens of Kentucky.

Geogr. Distr. New Jersey to South Carolina, west to Mississippi, north to Minnesota.

a. *Peromyscus leucopus noveboracensis* (Fischer).

Mus sylvaticus noveboracensis Fisch., Syn. Mamm., 1829, p. 318.

Cricetus myoides Gapper, Zoöl. Journ., v, 1830, p. 204, pl. x.

Peromyscus leucopus myodes! Rhoads, Proc. Acad. Nat. Scien. Phil., 1897, p. 27.

Peromyscus americanus noveboracensis Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 125. Zoöl. Ser.

NORTHERN WHITE-FOOTED DEER MOUSE.

Type locality. New York State.

Geogr. Distr. From Lake Simcoe, Ontario and Digby, Nova Scotia, south through New York except in the boreal area of that state.

b. *Peromyscus leucopus deserticola* (Mearns).

Hesperomys leucopus deserticolus Mearns, Bull. Am. Mus. Nat. Hist. N. Y., 1890, p. 285.

Peromyscus americanus deserticolus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 125. Zoöl. Ser.

Peromyscus texensis deserticola Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 188. Zoöl. Ser.

DESERT WHITE-FOOTED MOUSE.

Type locality. Mohave Desert, San Bernardino County, California.

Geogr. Distr. Desert region of California and Arizona.

c. *Peromyscus leucopus arizonæ* (Allen).

Sitomys americanus arizonæ Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1894, p. 321.

Peromyscus americanus arizonæ Elliot, Syn. N. Am. Mamm. F. C. M. Pub., II, 1901, p. 125. Zoöl. Ser.

Peromyscus texensis arizonæ Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 186. Zoöl. Ser.

ARIZONA WHITE-FOOTED DEER MOUSE.

Type locality. Fairbank, Cochise County, Arizona.

Geogr. Distr. Mountains of Arizona; White, Chiricahua, and Graham Ranges.

d. *Peromyscus leucopus artemisiæ* (Rhoads).

Sitomys americanus artemisiæ Rhoads, Proc. Acad. Nat. Scien. Phil., 1894, p. 260.

Peromyscus americanus artemisiæ Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 126. Zoöl. Ser.

PACIFIC COAST WHITE-FOOTED DEER MOUSE.

Type locality. Ashcroft, British Columbia, Canada.

Geogr. Distr. Southern Idaho, western Washington, to British Columbia, Canada.

e. *Peromyscus leucopus minnesotæ* Mearns.

Peromyscus leucopus minnesotæ Mearns, Proc. Biol. Soc. Wash., XIV, 1901, p. 154.

MINNESOTA WHITE-FOOTED DEER MOUSE.

Type locality. Fort Snelling, Hennepin County, Minnesota.

Geogr. Distr. Minnesota. Range unknown.

Genl. Char. Like *P. leucopus*.

Color. Summer Pelage. Above light bistre lined with black, under parts gray; tail and feet nearly naked; tail beneath gray; ear sepia.

Winter Pelage. Above cinnamon lined with black, beneath white; ear light brown, with whitish tuft at base not always present; tail above blackish brown, beneath white; feet white.

Measurements. Total length, 175; tail vertebræ, 75; hind foot, 21.5; ear, 11.5. Skull: total length, 25; Hensel, 18; occipito-nasal length, 25; interorbital constriction, 4; zygomatic width, 13; palatal length, 10; length of nasals, 9; length of upper tooth row, 4; length of mandible, 15; length of lower tooth row, 4.

279. *Peromyscus sonoriensis* (Le Conte).

Hesperomys sonoriensis Le Conte, Proc. Acad. Nat. Scien. Phil., 1853, p. 413.

Peromyscus americanus sonoriensis Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 125. Zoöl. Ser.

Peromyscus leucopus sonoriensis Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 181, fig. xxxvii. Zoöl. Ser.

SONORA WHITE-FOOTED FIELD MOUSE.

Type locality. Santa Cruz, State of Sonora, Mexico.

Geogr. Distr. Arctic regions into Mexico, west of Rocky Mountains.

a. *Peromyscus sonoriensis blandus* Osgood.

Peromyscus sonoriensis blandus Osgood, Proc. Biol. Soc. Wash., xvii, 1904, p. 56.

Peromyscus leucopus blandus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 182. Zoöl. Ser.

GENTLE FIELD MOUSE.

Type locality. Escalon, State of Chihuahua, Mexico.

Geogr. Distr. State of Chihuahua. Range unknown.

b. *Peromyscus sonoriensis fulvus* Osgood.

Peromyscus sonoriensis fulvus Osgood, Proc. Biol. Soc. Wash., xvii, 1904, p. 57.

Peromyscus leucopus fulvus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 183. Zoöl. Ser.

FULVOUS FIELD MOUSE.

Type locality. Near Oaxaca City, State of Oaxaca, Mexico.

Geogr. Distr. State of Oaxaca. Range unknown.

280. *Peromyscus labecula* Elliot.

Peromyscus labecula Elliot, Pub. Field Columb. Mus., III, 1903, p. 143. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 178. Zoöl. Ser.

WHITE-SPOT FIELD MOUSE.

Type locality. Ocotlan, State of Jalisco, Mexico.

Geogr. Distr. State of Jalisco. Range unknown.

281. *Peromyscus tornillo* Mearns.

Peromyscus tornillo Mearns, Proc. U. S. Nat. Mus., 1896, p. 445.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 126.

Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 188. Zoöl. Ser.

RIO GRANDE WHITE-FOOTED MOUSE.

Type locality. Rio Grande, six miles above El Paso, El Paso County, Texas.

Geogr. Distr. State of Chihuahua, Mexico; north into Texas and New Mexico.

282. *Peromyscus rufinus* (Merriam).

Hesperomys leucopus rufinus Merr., N. Am. Faun., No. 3, 1890, p. 65, pl. 3.

Peromyscus rufinus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 126. Zoöl. Ser.

TAWNY FIELD MOUSE.

Type locality. San Francisco Mountain, Coconino County, Arizona. Altitude, 9,000 feet.

Geogr. Distr. Northwestern Arizona to New Mexico, Utah, and Colorado.

283. *Peromyscus* **gossypinus* (Le Conte).

Hesperomys gossypinus! Le Conte, Proc. Acad. Nat. Scien. Phil., 1853, p. 411.

Peromyscus gossypinus! Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 126. Zool. Ser.

COTTON TREE MOUSE.

Type locality. Near Riceboro, Liberty County, Georgia.

Geogr. Distr. North Florida to Bertie County, North Carolina, west to Alabama and Mississippi.

a. *Peromyscus gossypinus mississippiensis* Rhoads.

Peromyscus gossypinus mississippiensis Rhoads, Proc. Acad. Nat. Scien. Phil., 1896, p. 189. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 127. Zool. Ser.

TENNESSEE COTTON MOUSE.

Type locality. Samburg, Reelfoot Lake, Tennessee.

Geogr. Distr. Mississippi bottoms in Tennessee. Range unknown.

b. *Peromyscus gossypinus palmarius* Bangs.

Peromyscus gossypinus palmarius Bangs, Proc. Biol. Soc. Wash., x, 1896, p. 124. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 127. Zool. Ser.

FLORIDA FIELD MOUSE.

Type locality. Oak Lodge, opposite Micco, Brevard County, Florida.

Geogr. Distr. Peninsula of Florida.

c. *Peromyscus gossypinus nigriculus* Bangs.

Peromyscus gossypinus nigriculus Bangs, Proc. Biol. Soc. Wash., x, 1896, p. 124. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 127. Zool. Ser.

LOUISIANA FIELD MOUSE.

Type locality. Burbridge, Plaquemines Parish, Louisiana.

Geogr. Distr. Bayou Region, Louisiana coast.

284. *Peromyscus anastasie* Bangs.

Peromyscus anastasæ! Bangs, Proc. Bost. Soc. Nat. Hist., xxviii, 1898, p. 195.

**Gossypinus*, the cotton tree; also *gossypion*, nec *gossypinus!*

Peromyscus anastasiæ Elliot, Syn. N. Am. Mamm., F. C. M. Pub.,
 II, 1901, p. 128. Zoöl. Ser.

ANASTASIA ISLAND MOUSE.

Type locality. Point Romo, Anastasia Island, St. John's County, Florida.

Geogr. Distr. Anastasia Island, Florida.

285. *Peromyscus insulanus* Bangs.

Peromyscus insulanus Bangs, Proc. Bost. Soc. Nat. Hist., xxviii,
 1898, p. 196. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II,
 1901, p. 128. Zoöl. Ser.

CUMBERLAND ISLAND FIELD MOUSE.

Type locality. Cumberland Island, Camden County, Georgia.

Geogr. Distr. Known only from the type locality.

286. *Peromyscus canadensis* (Miller).

? *Hesperomys gracilis* Le Conte, Proc. Acad. Nat. Scien. Phil.,
 1855, p. 442.

Hesperomys myoides Baird, N. Am. Mamm., 1857, p. 472. (nec
 Gapper.)

Sitomys americanus canadensis Miller, Proc. Biol. Soc. Wash.,
 VIII, 1893, p. 55.

Peromyscus canadensis Elliot, Syn. N. Am. Mamm., F. C. M. Pub.,
 II, 1901, p. 128. Zoöl. Ser.

CANADIAN FIELD MOUSE.

Type locality. Peterboro, Madison County, New York.

Geogr. Distr. Spruce forests of New Brunswick, south to central
 New York and western Massachusetts.

a. *Peromyscus canadensis abietorum* Bangs.

Peromyscus canadensis abietorum Bangs, Proc. Biol. Soc. Wash., x,
 1896, p. 49. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II,
 1901, p. 128. Zoöl. Ser.

BANGS' FIELD MOUSE.

Type locality. Banks of James River, Nova Scotia.

Geogr. Distr. Nova Scotia. Range unknown.

b. *Peromyscus canadensis umbrinus* Miller.

Peromyscus canadensis umbrinus Miller, Proc. Bost. Soc. Nat.
 Hist., xxviii, 1897, p. 23. Elliot, Syn. N. Am. Mamm.,
 F. C. M. Pub., II, 1901, p. 129. Zoöl. Ser.

SHADED FIELD MOUSE.

Type locality. Peninsula Harbor, north shore of Lake Superior,
 Ontario, Canada.

Geogr. Distr. Unknown.

c. *Peromyscus canadensis nubiterræ* Rhoads.

Peromyscus canadensis nubiterræ Rhoads, Proc. Acad. Nat. Scien. Phil., 1896, p. 187. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 129. Zoöl. Ser.

ROAN MOUNTAIN MOUSE.

Type locality. Roan Mountain, Mitchell County, North Carolina.
Geogr. Distr. Summit of southern Alleghanies. Altitude, 6,370 feet.

287. *Peromyscus texensis* (Woodhouse).

Hesperomys texana! Woodh., Proc. Acad. Nat. Scien. Phil., 1853, p. 242.

Peromyscus texensis Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 130. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 186. Zoöl. Ser.

TEXAS FIELD MOUSE.

Type locality. Rio Grande near El Paso, El Paso County, Texas.
Geogr. Distr. Southern Texas into State of Chihuahua, Mexico.

a. *Peromyscus texensis clementis* Mearns.

Peromyscus texanus clementis Mearns, Proc. U. S. Nat. Mus., 1896, p. 446.

Peromyscus texensis clementis Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 130. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 187. Zoöl. Ser.

SAN CLEMENTE ISLAND MOUSE.

Type locality. San Clemente Island, Los Angeles County, California.

Geogr. Distr. Coronados Islands, Mexico, and San Clemente Island, California.

b. *Peromyscus texensis gambeli* (Baird).

Hesperomys gambelii Baird, N. Am. Mamm., 1857, p. 464.

Peromyscus texensis gambeli Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 130. Zoöl. Ser.

GAMBEL'S MOUSE.

Type locality. Monterey, Monterey County, California.

Geogr. Distr. California, Nevada, Oregon, and Washington.

c. *Peromyscus texensis saturatus* Bangs.

Peromyscus texanus saturatus Bangs, Amer. Nat., xxxi, 1897, p. 75.

Peromyscus texensis saturatus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 131. Zoöl. Ser.

RICH-COLORED MOUSE.

Type locality. Saturna Island, Gulf of Georgia, British Columbia, Canada.

Geogr. Distr. Unknown.

d. *Peromyscus texensis arcticus* (Mearns).

Hesperomys leucopus arcticus Mearns, Bull. Am. Mus. Nat. Hist. N. Y., 1890, p. 285.

Peromyscus texensis arcticus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 131. Zoöl. Ser.

ARCTIC FIELD MOUSE.

Type locality. Fort Simpson, Mackenzie, Canada.

Geogr. Distr. Hudson Bay to Rocky Mountains.

e. *Peromyscus texensis subarcticus* Allen.

Peromyscus texanus subarcticus Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1899, p. 15.

Peromyscus texensis subarcticus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 131. Zoöl. Ser.

SUBARCTIC FIELD MOUSE.

Type locality. Deerlodge County, Montana.

Geogr. Distr. From Utah and Colorado, through Wyoming, the Dakotas, and Montana to the Saskatchewan Valley, Alberta, Canada.

f. *Peromyscus texensis nebrascensis* (Mearns).

Hesperomys leucopus nebrascensis Mearns, Bull. Am. Mus. Nat. Hist. N. Y., 1890, p. 287.

Peromyscus texensis nebrascensis Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 130. Zoöl. Ser.

NEBRASKA FIELD MOUSE.

Type locality. Calf Creek, central Montana.

Geogr. Distr. From Kansas and Nebraska, through the Dakotas and Wyoming into Montana.

g. *Peromyscus texensis thurberi* (Allen).

Sitomys americanus thurberi Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1893, p. 185.

Peromyscus texanus medius Mearns, Proc. U. S. Nat. Mus., 1896, p. 446.

Peromyscus texensis medius Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 130. Zoöl. Ser.

Peromyscus thurberi Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 176. Zoöl. Ser.

THURBER'S FIELD MOUSE.

Type locality. San Pedro Martir Mountains, Lower California.

Geogr. Distr. Region of San Pedro Martir Mountains, Lower California, Mexico, north into southern California to Colorado desert and possibly further.

h. *Peromyscus texensis flaccidus* Allen.

Peromyscus texanus flaccidus Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1903, p. 599.

Peromyscus texensis flaccidus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 187. Zoöl. Ser.

FEEBLE MOUSE.

Type locality. Rio Sestin, State of Durango, Mexico.

Geogr. Distr. State of Durango, Mexico. Range unknown.

i. *Peromyscus texensis coolidgei* (Thomas).

Peromyscus leucopus coolidgei Thomas, Ann. Mag. Nat. Hist., 7th Ser., 1, 1898, p. 45.

Peromyscus texensis coolidgei Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 187. Zoöl. Ser.

COOLIDGE'S FIELD MOUSE.

Type locality. Santa Anita, Cape Region, Lower California.

Geogr. Distr. Cape Region, Lower California.

j. *Peromyscus texensis mesomelas* Osgood.

Peromyscus texanus mesomelas Osgood, Proc. Biol. Soc. Wash., XVII, 1904, p. 57.

Peromyscus thurberi mesomelas Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 177. Zoöl. Ser.

OSGOOD'S FIELD MOUSE.

Type locality. Orizaba, State of Vera Cruz, Mexico.

Geogr. Distr. Known only from the type locality.

k. *Peromyscus texensis castaneus* Osgood.

Peromyscus texanus castaneus Osgood, Proc. Biol. Soc. Wash., XVII, 1904, p. 58.

Peromyscus thurberi castaneus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 177. Zoöl. Ser.

YOHALTUN FIELD MOUSE.

Type locality. Yohaltun, State of Campeche, Mexico.

Geogr. Distr. Known only from the type locality.

288. *Peromyscus luteus* Osgood.

Peromyscus luteus Osgood, Proc. Biol. Soc. Wash., XVIII, 1905, p. 77.

BUFF MOUSE.

Type locality. Kennedy, Nebraska.

Geogr. Distr. Western Nebraska and adjacent regions.

Color. "Upper parts bright ochraceous buff very lightly mixed with dusky; subauricular spots pure white, large, and conspicuous; under parts pure white."

Measurements. "Total length, 152; tail vertebræ, 62; hind foot, 20."

289. *Peromyscus exiguus* Allen.

Peromyscus exiguus Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1898, p. 157. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 183. Zoöl. Ser.

SAN MARTIN ISLAND MOUSE.

Type locality. San Martin Island, Lower California, Mexico.

Geogr. Distr. Known only from the type locality.

290. *Peromyscus akeleyi* Elliot.

Peromyscus akeleyi Elliot, Pub. Field Columb. Mus., I, 1899, p. 226. Zoöl. Ser. *Id.* Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 129. Zoöl. Ser.

AKELEY'S FIELD MOUSE.

Type locality. Johnson's Ranch, Elwah River, Clallam County, Washington.

Geogr. Distr. Region of Olympic Mountains, Washington.

291. *Peromyscus homochroia* Elliot.

Peromyscus homochroia Elliot, Pub. Field Columb. Mus., III, 1903, p. 158. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 200. Zoöl. Ser.

SAN QUENTIN MOUSE.

Type locality. San Quentin, Lower California, Mexico.

Geogr. Distr. Known only from the type locality.

292. *Peromyscus gaurus* Elliot.

Peromyscus gaurus Elliot, Pub. Field Columb. Mus., III, 1903, p. 157. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 199. Zoöl. Ser.

FRISKY MOUSE.

Type locality. San Antonio, San Pedro Martir Mountains, Lower California, Mexico.

Geogr. Distr. Known only from the type locality.

293. *Peromyscus maniculatus* (Wagner).

Hesperomys maniculatus Wagn., Archiv für Naturg., XI, Band I, 1845, p. 148.

Peromyscus maniculatus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 131. Zoöl. Ser.

WAGNER'S FIELD MOUSE.

Type locality. Labrador.

Geogr. Distr. Range unknown.

294. *Peromyscus catalinæ* Elliot.

Peromyscus catalinæ Elliot, Pub. Field Columb. Mus., III, 1903,
p. 160. Zoöl. Ser.

CATALINA ISLAND MOUSE.

Type locality. Santa Catalina Island, Santa Barbara Islands, California.

Geogr. Distr. Known only from the type locality.

Genl. Char. Size of *P. t. gambeli*, but with longer tail, ears, and feet.

Color. Upper parts grayish buff lined with black, lower back darkest; rump grayish buff; cheeks, nose, and sides buff; upper lip, lower sides of face, entire under parts, hands, and feet pure white; tail above dusky, beneath white; ears brownish black.

Measurements. Total length, 176; tail vertebræ, 92; hind foot, 23; ear, 18. Skull: occipito-nasal length, 25; Hensel, 19; zygomatic width, 13; interorbital constriction, 4; mastoid width, 10.2; length of nasals, 8; width of braincase, 10; length of upper tooth row, 3.

295. *Peromyscus austerus* (Baird).

Hesperomys austerus Baird, Proc. Acad. Nat. Scien. Phil., 1855,
p. 336.

Peromyscus austerus Elliot, Syn. N. Am. Mamm., F. C. M. Pub.,
II, 1901, p. 132. Zoöl. Ser.

TROUBLESOME FIELD MOUSE.

Type locality. Spokane Plain, Spokane County, Washington.

Geogr. Distr. Pacific coast, California to Washington.

296. *Peromyscus perimecurus* Elliot.

Peromyscus perimecurus Elliot, Pub. Field Columb. Mus., III,
1903, p. 156. Zoöl. Ser.

LONG-TAILED MOUSE.

Type locality. Goldbeach, Curry County, Oregon.

Geogr. Distr. Coast region of Oregon.

Genl. Char. Similar to *P. austerus* but larger and lighter in color; tail very long. Skull with larger bullæ, wider pterygoid fossa, and narrower rostrum.

Color. Back of head and dorsal region buff lined with black; fore part of head, nose, sides of head and body grayish buff; lateral line bright buff; lips and under parts plumbeous washed with white; hands and feet white; tail brown above, flesh color beneath, end

black; ear inside orange at base, rest black, outside pale brown on basal half, rest blackish with white edge.

Measurements. Total length, 220; tail vertebræ, 116; hind foot, 23; ear, 19. Skull: occipito-nasal length, 25.5; Hensel, 19; zygomatic width, 12; interorbital constriction, 4; palatal length, 9.5; width of braincase, 11.5; length of nasals, 9.5; width of rostrum, 3; length of upper tooth row, 3; length of mandible, 11.5; height at condyle, 4.5.

297. *Peromyscus mearnsi* (Allen).

Vesperimus mearnsii Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1891, p. 300.

Peromyscus mearnsi Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 132. Zoöl. Ser.

MEARNS' FIELD MOUSE.

Type locality. Brownsville, Cameron County, Texas.

Geogr. Distr. Texas to Arizona.

298. *Peromyscus canus* Mearns.

Peromyscus canus Mearns, Proc. U. S. Nat. Mus., 1896, p. 445.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 132. Zoöl. Ser.

HOARY FIELD MOUSE.

Type locality. Fort Clark, Kinney County, Texas.

Geogr. Distr. Texas to Oklahoma Territory.

299. *Peromyscus boylii* (Baird).

Hesperomys boylii Baird, Proc. Acad. Nat. Scien. Phil., 1855, p. 335.

Sitomys gilberti Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1893, p. 188.

Peromyscus boylii Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 132. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W.

Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 195. Zoöl. Ser.

BOYLE'S FIELD MOUSE.

Type locality. American River, Eldorado County, California.

Geogr. Distr. Central and eastern California.

a. *Peromyscus boylii penicillatus* Mearns.

Peromyscus boylii penicillatus Mearns, Proc. U. S. Nat. Mus., 1896, p. 139. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11,

1901, p. 133. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 196. Zoöl. Ser.

PENCILED FIELD MOUSE.

Type locality. Foothills of Franklin Mountains, near El Paso, El Paso County, Texas.

Geogr. Distr. Northwestern Texas.

300. *Peromyscus rowleyi* (Allen).

Sitomys rowleyi Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1893, p. 76.

Peromyscus rowleyi Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 134. Zoöl. Ser.

ROWLEY'S FIELD MOUSE.

Type locality. Nolan's Ranch, San Juan County, Utah.

Geogr. Distr. Utah to Arizona.

a. *Peromyscus rowleyi pinalis* (Miller).

Sitomys rowleyi pinalis Miller, Bull. Am. Mus. Nat. Hist. N. Y., 1893, p. 331.

Peromyscus rowleyi pinalis Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 135. Zoöl. Ser. *Id.* Mamm. Middle Amer.

& W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 189. Zoöl. Ser.

ROWLEY'S PINE MOUSE.

Type locality. Granite Gap, Grant County, New Mexico.

Geogr. Distr. State of Sonora, Mexico, north to New Mexico and Arizona.

301. *Peromyscus parasiticus* Elliot.

Peromyscus parasiticus Elliot, Pub. Field Columb. Mus., III, 1903, p. 244. Zoöl. Ser.

PARASITE MOUSE.

Type locality. Lone Pine, Inyo County, California.

Geogr. Distr. Known only from the type locality.

Genl. Char. Similar to *P. r. pinalis*, but larger in all of its dimensions. Skull with larger rostrum, longer nasals, broader between orbits, and larger, differently shaped braincase.

Color. Top of head and dorsal region dusky cinnamon, becoming pale cinnamon on sides of face beneath eyes; shoulders, flanks, and sides of rump, lips, sides of nose, lower part of flanks, thighs, hands, and feet and entire under parts white; base of fur plumbeous; tail above dusky, beneath yellowish white.

Measurements. Total length, 214; tail vertebræ, 119; hind foot, 23.5; ear, 20.5. Skull: total length, 28; Hensel, 19; zygomatic width, 13; interorbital constriction, 4.5; width of braincase, 12.5; length of braincase, 14; palatal length, 11; length of nasals, 11; length of upper tooth row, 4; length of mandible, 16; length of lower tooth row, 4.

302. *Peromyscus keeni* (Rhoads).

Sitomys keeni Rhoads, Proc. Acad. Nat. Scien. Phil., 1894, p. 258.

Peromyscus keeni Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 133. Zoöl. Ser.

KEEN'S MOUSE.

Type locality. Masset, Graham Island, Queen Charlotte Islands, British Columbia, Canada.

Geogr. Distr. Known only from the type locality.

303. *Peromyscus macrorhinus* (Rhoads).

Sitomys macrorhinus Rhoads, Proc. Acad. Nat. Scien. Phil., 1894, p. 259.

Peromyscus macrorhinus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 133. Zoöl. Ser.

LONG-NOSED MOUSE.

Type locality. Skeena River, British Columbia, Canada.

Geogr. Distr. British Columbia, Canada, possibly south into Washington.

304. *Peromyscus prevostensis* Osgood.

Peromyscus prevostensis Osgood, N. Am. Faun., No. 21, 1901, p. 29.

PREVOST ISLAND MOUSE.

Type locality. Prevost Island, Queen Charlotte Islands, British Columbia, Canada.

Geogr. Distr. Known only from the type locality.

Genl. Char. Similar to *P. sitkensis*, tail shorter; nasals shorter; posterior palatine foramina much longer.

Color. In front of and around eyes black; faint dusky dorsal stripe; beneath grayish white; occasionally a buff stripe in center of the breast; hind feet dusky; tail bicolor.

Measurements. "Average of 47 adults, total length, 217; tail vertebræ, 104; hind foot, 26." (Osgood.)

305. *Peromyscus sitkensis* Merriam.

Peromyscus sitkensis Merr., Proc. Biol. Soc. Wash., XI, 1897, p. 223. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901,

p. 134. Zoöl. Ser.

SITKA MOUSE.

Type locality. Sitka, Alaska.

Geogr. Distr. Baranof Island, Alaska.

306. *Peromyscus oreas* Bangs.

Peromyscus oreas Bangs, Proc. Biol. Soc. Wash., XII, 1898, p. 84.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 134.

Zoöl. Ser.

MOUNTAIN MOUSE.

Type locality. Mount Baker Range, British Columbia, Canada, near Whatcom County, Washington.

Geogr. Distr. Higher mountains of northern Washington and southern British Columbia.

a. Peromyscus oreas rubidus Osgood.

Peromyscus oreas rubidus Osgood, Proc. Biol. Soc. Wash., xiv, 1901, p. 193.

REDDISH MOUSE.

Type locality. Mendocino City, Mendocino County, California.

Geogr. Distr. Coast region of California and southern Oregon, south to Cazadero, California, in the red wood strip.

Genl. Char. Similar to *P. oreas*, tail shorter; hind foot smaller.

Color. Summer Pelage. Above and on sides brownish fawn; dorsal line dark; under parts white; ears edged with white; tail bicolor.

Measurements. Total length, 180-203; tail vertebræ, 90-100; hind foot, 21-22.

307. *Peromyscus auripectus* (Allen).

Sitomys auripectus Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1893, p. 75.

Peromyscus auripectus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 134. Zoöl. Ser.

GOLDEN BREASTED MOUSE.

Type locality. Bluff City, San Juan County, Utah.

Geogr. Distr. Utah, Arizona, and New Mexico.

308. *Peromyscus stephensi* Mearns.

Peromyscus stephensi Mearns, Proc. U. S. Nat. Mus., 1897, p. 721. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 136.

Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 190. Zoöl. Ser.

**Peromyscus petræus* Elliot, Pub. Field Columb. Mus., III, 1903, p. 244. Zoöl. Ser.

STEPHENS' MOUSE.

Type locality. Wagon-road in cañon at east base of the Coast Range, San Diego County, southern California.

Geogr. Distr. From San Diego County and Palm Springs, Riverside County, to Lone Pine, Inyo County.

309. *Peromyscus polius* Osgood.

Peromyscus polius Osgood, Proc. Biol. Soc. Wash., xvii, 1904, p. 61. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M.

Pub., IV, Pt. 1, 1904, p. 196. Zoöl. Ser.

*Mr. W. H. Osgood kindly compared my specimens with the type of *P. stephensi* in Coll. Biol. Survey, Wash., and advised me they were of the same species.

GRIZZLED MOUSE.

Type locality. Colonia Garcia, State of Chihuahua, Mexico.

Geogr. Distr. Known only from the type locality.

310. *Peromyscus attwateri* Allen.

Peromyscus attwateri Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1895, p. 330. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 135. Zoöl. Ser.

ATTWATER'S FIELD MOUSE.

Type locality. Turtle Creek, Kerr County, Texas.

Geogr. Distr. Texas to Indian Territory.

a. *Peromyscus attwateri pectoralis* Osgood.

Peromyscus attwateri pectoralis Osgood, Proc. Biol. Soc. Wash., XVII, 1904, p. 59. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 192. Zoöl. Ser.

JALAPAN MOUSE.

Type locality. Jalapan, State of Queretaro, Mexico.

Geogr. Distr. State of Queretaro, Mexico. Range unknown.

b. *Peromyscus attwateri eremicoides* Osgood.

Peromyscus attwateri eremicoides Osgood, Proc. Biol. Soc. Wash., XVII, 1904, p. 60. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 192. Zoöl. Ser.

ALLIED FIELD MOUSE.

Type locality. Mapimi, State of Durango, Mexico.

Geogr. Distr. Unknown.

311. *Peromyscus bellus* Bangs.

Peromyscus bellus Bangs, Proc. Biol. Soc. Wash., x, 1896, p. 137. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 135. Zoöl. Ser.

BEAUTIFUL MOUSE.

Type locality. Stillwell, Boston Mountains, Indian Territory.

Geogr. Distr. Known only from the type locality.

312. *Peromyscus insolatus* (Rhoads).

Sitomys insolatus Rhoads, Proc. Acad. Nat. Scien. Phil., 1894, p. 256.

Peromyscus insolatus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 135. Zoöl. Ser.

SUNNY MOUSE.

Type locality. Oro Grande, Mohave Desert, Kern County, California.

Geogr. Distr. Mohave Desert, California. Range unknown.

313. *Peromyscus fraterculus* (Miller).

Vesperimus fraterculus Miller, Amer. Nat., xxvi, 1892, p. 261.

Peromyscus fraterculus Elliot, Syn. N. Am. Mamm., F. C. M. Pub.,
 11, 1901, p. 136. Zoöl. Ser. *Id.* Mamm. Middle Amer. &
 W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 184. Zoöl. Ser.

RELATED WHITE-FOOTED MOUSE.

Type locality. Dulzura, San Diego County, California.

Geogr. Distr. Southern California into Lower California, Mexico.

314. *Peromyscus cedrosensis* Allen.

Peromyscus cedrosensis Allen, Bull. Am. Mus. Nat. Hist. N. Y.,
 1898, p. 154. Elliot, Mamm. Middle Amer. & W. Indies,
 F. C. M. Pub., iv, Pt. 1, 1904, p. 185. Zoöl. Ser.

CERROS ISLAND MOUSE.

Type locality. Cerros or Cedros Island, Lower California, Mexico.

Geogr. Distr. Known only from the type locality.

315. *Peromyscus major* (Rhoads).

Sitomys major Rhoads, Amer. Nat., xxvii, 1893, p. 831.

Peromyscus major Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11,
 1901, p. 137. Zoöl. Ser.

SQUIRREL INN MOUSE.

Type locality. Squirrel Inn, San Bernardino County, California.

Geogr. Distr. Southern California. Range unknown.

316. *Peromyscus herroni* (Rhoads).

Sitomys herronii Rhoads, Amer. Nat., xxvii, 1893, p. 832.

Peromyscus herroni Elliot, Syn. N. Am. Mamm., F. C. M. Pub.,
 11, 1901, p. 137. Zoöl. Ser.

HERRON'S FIELD MOUSE.

Type locality. San Bernardino Valley, California.

Geogr. Distr. Southern California.

a. Peromyscus herroni nigellus (Rhoads).

Sitomys herroni nigellus Rhoads, Proc. Acad. Nat. Scien. Phil.,
 1894, p. 257.

Peromyscus herroni nigellus Elliot, Syn. N. Am. Mamm., F. C. M.
 Pub., 11, 1901, p. 137. Zoöl. Ser.

DARK FIELD MOUSE.

Type locality. West Cajon Pass, San Bernardino Mountains,
 San Bernardino County, California.

Geogr. Distr. High mountains, southern California.

317. *Peromyscus insignis* Rhoads.

Peromyscus insignis Rhoads, Proc. Acad. Nat. Scien. Phil., 1895,
 p. 33. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901,

p. 138. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies,
F. C. M. Pub., IV, Pt. 1, 1904, p. 203. Zoöl. Ser.

SACHEM OR CHIEF MOUSE.

Type locality. Dulzura, San Diego County, California.

Geogr. Distr. Southern California into Lower California, Mexico.

318. *Peromyscus dyselius* Elliot.

Peromyscus dyselius Elliot, Pub. Field Columb. Mus., 1, 1898,
p. 207. Zoöl. Ser. *Id.* Syn. N. Am. Mamm., F. C. M. Pub.,

II, 1901, p. 138. Zoöl. Ser.

BLACKISH FIELD MOUSE.

Type locality. Portola, San Mateo County, California.

Geogr. Distr. Coast region of California south of San Francisco.

319. *Peromyscus robustus* (Allen).

Sitomys robustus Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1893,
p. 335.

Peromyscus robustus Elliot, Syn. N. Am. Mamm., F. C. M. Pub.,
II, 1901, p. 138. Zoöl. Ser.

ROBUST FIELD MOUSE.

Type locality. Lakeport, Lake County, California.

Geogr. Distr. Unknown.

320. *Peromyscus floridanus* (Chapman).

Hesperomys floridanus Chapman, Bull. Am. Mus. Nat. Hist.
N. Y., 1889, p. 117.

Peromyscus floridanus Elliot, Syn. N. Am. Mamm., F. C. M. Pub.,
II, 1901, p. 138. Zoöl. Ser.

FLORIDA FIELD MOUSE.

Type locality. Gainesville, Alachua County, Florida.

Geogr. Distr. Florida Peninsula.

321. *Peromyscus spicilegus* Allen.

Peromyscus spicilegus Allen, Bull. Am. Mus. Nat. Hist. N. Y.,
1897, p. 50. Elliot, Mamm. Middle Amer. & W. Indies,
F. C. M. Pub., IV, Pt. 1, 1904, p. 179. Zoöl. Ser.

JALISCO SMALL-EARED MOUSE.

Type locality. Mineral San Sebastian, Mascota, State of Jalisco,
Mexico.

Geogr. Distr. States of Jalisco and Sinaloa. Range unknown.

a. Peromyscus spicilegus evides Osgood.

Peromyscus spicilegus evides Osgood, Proc. Biol. Soc. Wash.,
xvii, 1904, p. 64. Elliot, Mamm. Middle Amer. & W. Indies,
F. C. M. Pub., IV, Pt. 1, 1904, p. 179. Zoöl. Ser.

JUQUILA MOUSE.

Type locality. Juquila, State of Oaxaca, Mexico.

Geogr. Distr. Known only from the type locality.

b. *Peromyscus spicilegus simulus* Osgood.

Peromyscus spicilegus simulus Osgood, Proc. Biol. Soc. Wash., xvii, 1904, p. 64. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 180. Zoöl. Ser.

MIMIC MOUSE.

Type locality. San Blas, Territorio de Tepic, Mexico.

Geogr. Distr. Lowlands of Territorio de Tepic, Mexico. Range unknown.

322. *Peromyscus megalotis* (Merriam).

Hesperomys megalotis Merr., N. Am. Faun., No. 3, 1890, p. 64.

Peromyscus megalotis Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 139. Zoöl. Ser.

BIG-EARED MOUSE.

Type locality. Black Tank, Desert of the Little Colorado, Cocino County, Arizona.

Geogr. Distr. Desert of the Little Colorado, Arizona.

323. *Peromyscus montipinoris* Elliot.

Peromyscus montipinoris Elliot, Pub. Field Columb. Mus., 111, 1904, p. 264. Zoöl. Ser.

MOUNT PINOS MOUSE.

Type locality. Mount Pinos, Lockwood Valley, Ventura County, California.

Geogr. Distr. Known only from the type locality.

Genl. Char. Size medium; tail very long, hairy, penicillate; ears large; hind foot long; color pale; skull similar to that of *P. truei*, but with longer nasals, broader and longer braincase, and narrower inter-orbital region.

Color. Top of head and back mixed cinereous and buff, darker than rest of upper parts; upper sides buffy; entire under parts, hands, and feet white; ankles dusky; tail above black, beneath yellowish white; ears dark brown.

Measurements. Total length, 214; tail vertebræ, 118; hind foot, 24.5; ear, 24.5. Skull: occipito-nasal length, 28; Hensel, 22; inter-orbital constriction, 4.5; zygomatic width, 13; width of braincase, 13; length of nasals, 10; palatal length, 11; length of upper tooth row, 4; length of mandible, 12; length of lower tooth row, 4.

324. *Peromyscus lasius* Elliot.

Peromyscus lasius Elliot, Pub. Field Columb. Mus., 111, 1904, p. 265. Zoöl. Ser.

WOOLLY MOUSE.

Type locality. Hannopee Cañon, Panamint Mountains, Inyo County, California.

Geogr. Distr. Panamint and Inyo Mountains, California.

Genl. Char. Allied to *P. megalotis* and *P. truei*; colors pale; tail long; skull heavy; rostrum long.

Color. Nose and sides of face gray tinged with buff; orbital ring black; top of head and upper part of body pale ochraceous buff, mixed slightly with black on dorsal region; sides, and lateral line from beneath eyes to root of tail, ochraceous buff; entire under parts, hands, and feet pure white; under fur plumbeous; tail hairy, penicillate, with a pale dusky line above, remainder white; ears large, naked, dusky brown.

Measurements. Total length, 208; tail vertebræ, 105; hind foot, 24; ear from notch, 27.5. Skull: total length, 29; Hensel, 21.5; zygomatic width, 14; interorbital constriction, 4; length of nasals, 9.5; length of braincase, 18.5; width of braincase above root of zygomatic, 13.5; palatal length, 11; length of upper tooth row, 4; length of mandible, 12.5; length of lower tooth row, 4.

325. *Peromyscus difficilis* (Allen).

Vesperimus difficilis Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1891, p. 298.

Peromyscus difficilis Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 193. Zoöl. Ser.

TROUBLESOME MOUSE.

Type locality. Sierra de Valparaiso, State of Zacatecas, Mexico.

Geogr. Distr. State of Zacatecas, Mexico. Range unknown.

326. *Peromyscus bullatus* Osgood.

Peromyscus bullatus Osgood, Proc. Biol. Soc. Wash., xvii, 1904, p. 63. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 193. Zoöl. Ser.

PEROTE MOUSE.

Type locality. Perote, State of Vera Cruz, Mexico.

Geogr. Distr. Known only from the type locality.

327. *Peromyscus sagax* Elliot.

Peromyscus sagax Elliot, Pub. Field Columb. Mus., III, 1903, p. 142. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 194. Zoöl. Ser.

LA PALMA FIELD MOUSE.

Type locality. La Palma, State of Michoacan, Mexico.

Geogr. Distr. Unknown.

328. *Peromyscus truei* (Shufeldt).

Hesperomys truei Shufeldt, Proc. U. S. Nat. Mus., 1886, p. 407,
pl. XXI, figs. 1, 5, 8.

Peromyscus truei Elliot, Syn. N. Am. Mamm., F. C. M. Pub.,
11, 1901, p. 139. Zoöl. Ser.

TRUE'S MOUSE.

Type locality. Fort Wingate, Valencia County, New Mexico.

Geogr. Distr. Nevada, Utah, and New Mexico.

a. Peromyscus truei nasutus (Allen).

Vesperimus nasutus Allen, Bull. Am. Mus. Nat. Hist. N. Y.,
1891, p. 299.

Peromyscus truei nasutus Elliot, Syn. N. Am. Mamm., F. C. M.
Pub., 11, 1901, p. 139. Zoöl. Ser.

LARGE-NOSED MOUSE.

Type locality. Estes Park, Larimer County, Colorado.

Geogr. Distr. Colorado.

b. Peromyscus truei crinitus (Merriam).

Hesperomys crinitus Merr., N. Am. Faun., No. 5, 1891, p. 53,
fig. 3.

Peromyscus truei crinitus Elliot, Syn. N. Am. Mamm., F. C. M.
Pub., 11, 1901, p. 140. Zoöl. Ser.

LONG-HAIRED MOUSE.

Type locality. Shoshone Falls, Snake River, Lincoln County,
Idaho.

Geogr. Distr. Idaho. Range undetermined.

c. Peromyscus truei scitulus (Bangs).

Peromyscus crinitus scitulus Bangs, Proc. N. Engl. Zoöl. Club, 1,
1899, p. 67.

Peromyscus truei scitulus Elliot, Syn. N. Am. Mamm., F. C. M.
Pub., 11, 1901, Append., p. 429. Zoöl. Ser.

NEAT FIELD MOUSE.

Type locality. Gardnerville, Douglas County, Nevada.

Geogr. Distr. Nevada. Range unknown.

329. *Peromyscus martirensis* (Allen).

Sitomys martirensis Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1893,
p. 187.

Peromyscus martirensis Elliot, Mamm. Middle Amer. & W. Indies,
F. C. M. Pub., IV, Pt. 1, 1904, p. 189. Zoöl. Ser.

SAN PEDRO MARTIR MOUNTAINS MOUSE.

Type locality. San Pedro Martir Mountains, Lower California,
Mexico. Altitude, above 5,000 feet.

Geogr. Distr. San Pedro Martir and Hanson Laguna Mountains, Lower California, Mexico.

330. *Peromyscus nuttalli* (Harlan).

Arvicola nuttalli Harlan, Am. Month. Journ. Geol. & Nat. Scien. Phil., 1832, p. 446.

Peromyscus nuttalli Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 140. Zoöl. Ser.

NUTTALL'S GOLDEN MOUSE.

Type locality. Norfolk, Norfolk County, Virginia.

Geogr. Distr. Southern Virginia, North and South Carolina, and Georgia.

331. *Peromyscus cherrii* (Allen).

Hesperomys cherrii Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1891, p. 211.

Peromyscus cherrii, Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 183. Zoöl. Ser.

CHERRIE'S COSTA RICA MOUSE.

Type locality. La Carpintera, Costa Rica.

Geogr. Distr. Costa Rica, Central America.

332. *Peromyscus michiganensis* (Audubon & Bachman).

Mus michiganensis Aud. & Bachm., Jour. Acad. Nat. Scien. Phil., 1842, p. 304.

Peromyscus michiganensis Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 140. Zoöl. Ser.

MICHIGAN FIELD MOUSE.

Type locality. Erie County, Ohio.

Geogr. Distr. Indiana north to Minnesota and Dakota, and west and south to Kansas.

a. Peromyscus michiganensis pallescens Allen.

Peromyscus michiganensis pallescens Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1896, p. 238. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 140. Zoöl. Ser.

PALE MICHIGAN MOUSE.

Type locality. San Antonio, Bexar County, Texas.

Geogr. Distr. Texas.

333. *Peromyscus tiburonensis* Mearns.

Peromyscus tiburonensis Mearns, Proc. U. S. Nat. Mus., 1897, p. 720. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 175. Zoöl. Ser.

ISLAND OF TIBURON MOUSE.

Type locality. Tiburon Island, Gulf of California. Mexico.

Geogr. Distr. Known only from the type locality.

334. *Peromyscus ceciliae* Thomas.

Peromyscus ceciliae Thomas, Ann. Mag. Nat. Hist., 7th Ser., xi, 1903, p. 486. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 178. Zool. Ser.

CECIL'S MOUSE.

Type locality. Santa Barbara Camp, southern slope of Mt. Orizaba, State of Vera Cruz, Mexico. Altitude, 12,500 feet.

Geogr. Distr. State of Vera Cruz, Mexico. Range unknown.

335. *Peromyscus melanotis* Allen & Chapman.

Peromyscus melanotis Allen & Chapman, Bull. Am. Mus. Nat. Hist. N. Y., 1897, p. 203. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 178. Zool. Ser.

BLACK-EARED FIELD MOUSE.

Type locality. Las Vigas, State of Vera Cruz, Mexico.

Geogr. Distr. Range unknown.

a. *Peromyscus melanotis zamelas* Osgood.

Peromyscus melanotis zamelas Osgood, Proc. Biol. Soc. Wash., xvii, 1904, p. 59. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 179. Zool. Ser.

JET MOUSE.

Type locality. Colonia Garcia, State of Chihuahua, Mexico. Altitude, 6,700 feet.

Geogr. Distr. Range undetermined.

336. *Peromyscus affinis* (Allen).

Hesperomys affinis Allen, Proc. U. S. Nat. Mus., 1891, p. 195.

Peromyscus affinis Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 180. Zool. Ser.

ALLIED FIELD MOUSE.

Type locality. Barrio, State of Oaxaca, Mexico.

Geogr. Distr. Southeastern Mexico.

a. *Peromyscus affinis musculoides* (Merriam).

Peromyscus musculoides Merr., Proc. Biol. Soc. Wash., xii, 1898, p. 124. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 180. Zool. Ser.

ALLIED HOUSE MOUSE.

Type locality. Cuicatlan, State of Oaxaca, Mexico.

Geogr. Distr. State of Oaxaca, Mexico. Range unknown.

337. *Peromyscus cozumelæ* Merriam.

Peromyscus cozumelæ Merr., Proc. Biol. Soc. Wash., xiv, 1901, p. 103. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 181. Zoöl. Ser.

ISLAND OF COZUMEL MOUSE.

Type locality. Island of Cozumel, Yucatan, Mexico.

Geogr. Distr. Known only from the type locality.

338. *Peromyscus geronimensis* Allen.

Peromyscus geronimensis Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1898, p. 156. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 181. Zoöl. Ser.

SAN GERONIMO ISLAND FIELD MOUSE.

Type locality. San Geronimo Island, Lower California, Mexico.

Geogr. Distr. Lower California, Mexico.

339. *Peromyscus dubius* Allen.

Peromyscus dubius Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1898, p. 157. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 181. Zoöl. Ser.

TODOS SANTOS ISLAND MOUSE.

Type locality. Todos Santos Island, Lower California, Mexico.

Geogr. Distr. Known only from the type locality.

340. *Peromyscus aztecus* (Saussure).

Hesperomys aztecus Sauss., Rev. Mag. Zool., 2me Sér., xii, 1860, p. 105.

Peromyscus aztecus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 184. Zoöl. Ser.

AZTEC MOUSE.

Type locality. "Mexico."

Geogr. Distr. Cape St. Lucas, Lower California, and Sierra de Juanacutlan, State of Jalisco, Mexico. Limits of range unknown.

341. *Peromyscus beatæ* Thomas.

Peromyscus beatæ Thomas, Ann. Mag. Nat. Hist., 7th Ser., xi, 1903, p. 485. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 184. Zoöl. Ser.

XOMETLA MOUSE.

Type locality. Xometla Camp, Mt. Orizaba, State of Vera Cruz, Mexico.

Geogr. Distr. Known only from the type locality.

342. *Peromyscus gymnotis* Thomas.

Peromyscus gymnotis Thomas, Ann. Mag. Nat. Hist., 6th Ser., xiv, 1894, p. 365. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 188. Zoöl. Ser.

NAKED-EARED MOUSE.

Type locality. Guatemala.

Geogr. Distr. Guatemala, Central America.

343. *Peromyscus banderanus* Allen.

Peromyscus banderanus Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1897, p. 51. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 189. Zoöl. Ser.

BANDERAS FIELD MOUSE.

Type locality. Terro Tepic, Valle de Banderas, State of Jalisco, Mexico.

Geogr. Distr. State of Jalisco. Range unknown.

a. Peromyscus banderanus viciniior Osgood.

Peromyscus banderanus viciniior Osgood, Proc. Biol. Soc. Wash., XVII, 1904, p. 68. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 190. Zoöl. Ser.

LA SALADA DEER MOUSE.

Type locality. La Salada, State of Michoacan, Mexico.

Geogr. Distr. State of Michoacan. Range unknown.

b. Peromyscus banderanus angelensis Osgood.

Peromyscus banderanus angelensis Osgood, Proc. Biol. Soc. Wash., XVII, 1904, p. 69. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 190. Zoöl. Ser.

PUERTO ANGEL MOUSE.

Type locality. Puerto Angel, State of Oaxaca, Mexico.

Geogr. Distr. State of Oaxaca, Mexico. Range unknown.

344. *Peromyscus yucatanicus* Allen & Chapman.

Peromyscus yucatanicus Allen & Chapman, Bull. Am. Mus. Nat. Hist. N. Y., 1897, p. 8. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 194. Zoöl. Ser.

YUCATAN MOUSE.

Type locality. Chichen Itza, Yucatan, Mexico.

Geogr. Distr. Yucatan, Mexico.

a. Peromyscus yucatanicus badius Osgood.

Peromyscus yucatanicus badius Osgood, Proc. Biol. Soc. Wash., XVII, 1904, p. 70. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 194. Zoöl. Ser.

APAZOTE MOUSE.

Type locality. Apazote, State of Campeche, Mexico.

Geogr. Distr. State of Campeche. Range unknown.

345. *Peromyscus cineritius* Allen.

Peromyscus cineritius Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1898, p. 155. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 195. Zoöl. Ser.

ASHY GRAY FIELD MOUSE.

Type locality. San Roque Island, Lower California, Mexico.

Geogr. Distr. Known only from the type locality.

346. *Peromyscus levipes* Merriam.

Peromyscus levipes Merr., Proc. Biol. Soc. Wash., XII, 1898, p. 123. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 195. Zoöl. Ser.

MOUNT MALINCHE MOUSE.

Type locality. Mount Malinche, State of Tlaxcala, Mexico. Altitude, 8,400 feet.

Geogr. Distr. Limits of range unknown.

347. *Peromyscus gratus* Merriam.

Peromyscus gratus Merr., Proc. Biol. Soc. Wash., X, 1898, p. 123. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 196. Zoöl. Ser.

TLALPAM FIELD MOUSE.

Type locality. Tlalpam, Valley of Mexico.

Geogr. Distr. State of Mexico. Range unknown.

a. *Peromyscus gratus gentilis* Osgood.

Peromyscus gratus gentilis Osgood, Proc. Biol. Soc. Wash., XVII, 1904, p. 61. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 197. Zoöl. Ser.

FOREIGN MOUSE.

Type locality. Lagos, State of Jalisco, Mexico.

Geogr. Distr. Limits of range unknown.

348. *Peromyscus pavidus* Elliot.

Peromyscus pavidus Elliot, Pub. Field Columb. Mus., III, 1903, p. 142. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 197. Zoöl. Ser.

SHY FIELD MOUSE.

Type locality. Patzcuaro, State of Michoacan, Mexico.

Geogr. Distr. State of Michoacan, Mexico. Range unknown.

349. *Peromyscus madrensis* Merriam.

Peromyscus madrensis Merr., Proc. Biol. Soc. Wash., XII, 1898, p. 16. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 197. Zoöl. Ser.

MARIA MADRE ISLAND MOUSE.

Type locality. Maria Madre Island, Tres Marias Islands, State of Jalisco, Mexico.

Geogr. Distr. Known only from the type locality.

350. *Peromyscus eva* Thomas.

Peromyscus eva Thomas, Ann. Mag. Nat. Hist., 7th Ser., 1, 1898, p. 44. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 198. Zool. Ser.

EVE'S MOUSE.

Type locality. San José del Cabo, Lower California, Mexico.

Geogr. Distr. Cape Region, Lower California, Mexico.

351. *Peromyscus nudipes* (Allen).

Hesperomys nudipes Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1891, p. 213.

Peromyscus nudipes Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 198. Zool. Ser.

LA CARPINTERA FIELD MOUSE.

Type locality. La Carpintera, Costa Rica.

Geogr. Distr. Costa Rica, Central America.

352. *Peromyscus merriami* Mearns.

Peromyscus merriami Mearns, Proc. U. S. Nat. Mus., 1896, p. 138. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 198. Zool. Ser.

SONOYTA DESERT MOUSE.

Type locality. Sonoyta, on Sonoyta River, State of Sonora, Mexico.

Geogr. Distr. State of Sonora, Mexico. Range unknown.

353. *Peromyscus leucurus* Thomas.

Peromyscus leucurus Thomas, Ann. Mag. Nat. Hist., 6th Ser., XIV, 1894, p. 364. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 200. Zool. Ser.

WHITE-TAILED MOUSE.

Type locality. Tehuantepec, Mexico.

Geogr. Distr. Southern Mexico.

a. *Peromyscus leucurus gadovi* Thomas.

Peromyscus leucurus gadovi Thomas, Ann. Mag. Nat. Hist. N. Y., 7th Ser., XI, 1903, p. 484. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 201. Zool. Ser.

GADOW'S MOUSE.

Type locality. San Carlos, Yautepec, State of Oaxaca, Mexico. Altitude, 2,250 feet.

Geogr. Distr. Limits of range unknown.

354. *Peromyscus melanophrys* (Coues).

Hesperomys melanophrys Coues, Proc. Acad. Nat. Scien. Phil., 1874, p. 181.

Peromyscus melanophrys Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 201. Zoöl. Ser.

BLACK-BROWED MOUSE.

Type locality. Santa Efigenia, Tehuantepec, State of Oaxaca, Mexico.

Geogr. Distr. Southern Mexico, Tehuacan and Tehuantepec, States of Puebla and Oaxaca.

a. Peromyscus melanophrys zamoræ Osgood.

Peromyscus melanophrys zamoræ Osgood, Proc. Biol. Soc. Wash., XVII, 1904, p. 65. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 202. Zoöl. Ser.

ZAMORA MOUSE.

Type locality. Zamora, State of Michoacan, Mexico.

Geogr. Distr. Limits of range unknown.

b. Peromyscus melanophrys consobrinus Osgood.

Peromyscus melanophrys consobrinus Osgood, Proc. Biol. Soc. Wash., XVII, 1904, p. 66. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 202. Zoöl. Ser.

KINDRED MOUSE.

Type locality. Berriozabal, State of Zacátecas, Mexico.

Geogr. Distr. Unknown.

355. *Peromyscus xenurus* Osgood.

Peromyscus xenurus Osgood, Proc. Biol. Soc. Wash., XVII, 1904, p. 67. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 202. Zoöl. Ser.

ODD-TAILED MOUSE.

Type locality. Durango, State of Durango, Mexico.

Geogr. Distr. State of Durango, Mexico. Limits unknown.

356. *Peromyscus zelotes* Osgood.

Peromyscus zelotes Osgood, Proc. Biol. Soc. Wash., XVII, 1904, p. 67. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 203. Zoöl. Ser.

ENERGETIC MOUSE.

Type locality. Querendaro, State of Michoacan, Mexico.

Geogr. Distr. Known only from the type locality.

357. *Peromyscus hylocetes* Merriam.

Peromyscus hylocetes Merr., Proc. Biol. Soc. Wash., XII, 1898, p. 124. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 204. Zool. Ser.

DRYAD MOUSE.

Type locality. Patzcuaro, State of Michoacan, Mexico. Altitude, 8,000 feet.

Geogr. Distr. State of Michoacan, Mexico. Range unknown.

358. *Peromyscus lepturus* Merriam.

Peromyscus lepturus Merr., Proc. Biol. Soc. Wash., XII, 1898, p. 118. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 204. Zool. Ser.

MOUNT ZEMPOALTEPEC FIELD MOUSE.

Type locality. Mount Zempoaltepec, State of Oaxaca, Mexico. Altitude, 8,200 feet.

Geogr. Distr. Known only from the type locality.

359. *Peromyscus lophurus* Osgood.

Peromyscus lophurus Osgood, Proc. Biol. Soc. Wash., XVII, 1904, p. 72. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 204. Zool. Ser.

CRESTED-TAILED MOUSE.

Type locality. Todos Santos, Guatemala.

Geogr. Distr. Guatemala, Central America.

360. *Peromyscus simulatus* Osgood.

Peromyscus simulatus Osgood, Proc. Biol. Soc. Wash., XVII, 1904, p. 72. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 205. Zool. Ser.

IMITATOR MOUSE.

Type locality. Jico, State of Vera Cruz, Mexico.

Geogr. Distr. Known only from the type locality.

361. *Peromyscus felipensis* Merriam.

Peromyscus felipensis Merr., Proc. Biol. Soc. Wash., XII, 1898, p. 122. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 205. Zool. Ser.

SAN FELIPE FIELD MOUSE.

Type locality. Cerro San Felipe, State of Oaxaca, Mexico. Altitude, 10,200 feet.

Geogr. Distr. State of Oaxaca, Mexico. Range unknown.

362. *Peromyscus amplus* Osgood.

Peromyscus amplus Osgood, Proc. Biol. Soc. Wash., XVII, 1904,

p. 62. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 205. Zoöl. Ser.

CHUBBY MOUSE.

Type locality. Coixtlahuaca, State of Oaxaca, Mexico.

Geogr. Distr. Unknown.

363. *Peromyscus tehuantepecus* Merriam.

Peromyscus tehuantepecus Merr., Proc. Biol. Soc. Wash., XII, 1898, p. 122. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 206. Zoöl. Ser.

TEHUANTEPEC FIELD MOUSE.

Type locality. Tehuantepec, State of Oaxaca, Mexico.

Geogr. Distr. State of Oaxaca, Mexico. Range unknown.

364. *Peromyscus oaxacensis* Merriam.

Peromyscus oaxacensis Merr., Proc. Biol. Soc. Wash., XII, 1898, p. 122. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 206. Zoöl. Ser.

OAXACA FIELD MOUSE.

Type locality. Cerro San Felipe, State of Oaxaca, Mexico.

Geogr. Distr. Known only from the type locality.

365. *Peromyscus comptus* Merriam.

Peromyscus comptus Merr., Proc. Biol. Soc. Wash., XII, 1898, p. 120. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 206. Zoöl. Ser.

CHILPANCINGO MOUSE.

Type locality. Mountains near Chilpancingo, State of Guerrero, Mexico.

Geogr. Distr. State of Guerrero, Mexico. Range unknown.

366. *Peromyscus mexicanus* (Saussure).

Hesperomys mexicanus Sauss., Rev. Mag. Zool., 2me Sér., XII, 1860, p. 103, pl. IX, figs. 1, 1a.

Peromyscus mexicanus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 207. Zoöl. Ser.

MEXICAN MOUSE.

Type locality. State of Vera Cruz, Mexico.

Geogr. Distr. Not determined.

a. Peromyscus mexicanus teapensis Osgood.

Peromyscus mexicanus teapensis Osgood, Proc. Biol. Soc. Wash., XVII, 1904, p. 69. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 207. Zoöl. Ser.

TEAPA MOUSE.

Type locality. Teapa, State of Tabasco, Mexico.

Geogr. Distr. Known only from the type locality.

b. *Peromyscus mexicanus orizabæ* Merriam.

Peromyscus mexicanus orizabæ Merr., Proc. Biol. Soc. Wash., XII, 1898, p. 121. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 207. Zoöl. Ser.

ORIZABA FIELD MOUSE.

Type locality. Orizaba, State of Vera Cruz, Mexico. Altitude, 4,200 feet.

Geogr. Distr. State of Vera Cruz. Range unknown.

c. *Peromyscus mexicanus saxatilis* Merriam.

Peromyscus mexicanus saxatilis Merr., Proc. Biol. Soc. Wash., XII, 1898, p. 121. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 207. Zoöl. Ser.

ROCK MOUSE.

Type locality. Jacaltenango, Huehuetenango, Guatemala.

Geogr. Distr. Guatemala, Central America.

d. *Peromyscus mexicanus totontepecus* Merriam.

Peromyscus mexicanus totontepecus Merr., Proc. Biol. Soc. Wash., XII, 1898, p. 120. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 208. Zoöl. Ser.

TOTONTEPEC FIELD MOUSE.

Type locality. Totontepec, State of Oaxaca.

Geogr. Distr. State of Oaxaca, Mexico. Range unknown.

367. *Peromyscus allophylus* Osgood.

Peromyscus allophylus Osgood, Proc. Biol. Soc. Wash., XVII, 1904, p. 71. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 208. Zoöl. Ser.

ALIEN MOUSE.

Type locality. Huehuetan, State of Chiapas, Mexico.

Geogr. Distr. State of Chiapas, Mexico.

368. *Peromyscus furvus* Allen & Chapman.

Peromyscus furvus Allen & Chapman, Bull. Am. Mus. Nat. Hist. N. Y., 1897, p. 201. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 208, fig. 33. Zoöl. Ser.

JALAPA FIELD MOUSE.

Type locality. Jalapa, State of Vera Cruz, Mexico.

Geogr. Distr. State of Vera Cruz, Mexico.

369. *Peromyscus zarhynchus* Merriam.

Peromyscus zarhynchus Merr., Proc. Biol. Soc. Wash., xii, 1898, p. 117. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 209. Zoöl. Ser.

LONG-NOSED MOUSE.

Type locality. Tumbala, State of Chiapas, Mexico.

Geogr. Distr. State of Chiapas, Mexico.

a. Peromyscus zarhynchus cristobalensis Merriam.

Peromyscus zarhynchus cristobalensis Merr., Proc. Biol. Soc. Wash., xii, 1898, p. 117. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 209. Zoöl. Ser.

SAN CRISTOBAL FIELD MOUSE.

Type locality. San Cristobal, State of Chiapas, Mexico.

Geogr. Distr. State of Chiapas, Mexico. Range unknown.

370 *Peromyscus auritus* Merriam.

Peromyscus auritus Merr., Proc. Biol. Soc. Wash., xii, 1898, p. 119. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 209. Zoöl. Ser.

LARGE-EARED FIELD MOUSE.

Type locality. Mountains west of Oaxaca City, State of Oaxaca, Mexico.

Geogr. Distr. Known only from the type locality.

371. *Peromyscus megalops* Merriam.

Peromyscus megalops Merr., Proc. Biol. Soc. Wash., xii, 1898, p. 119. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 210. Zoöl. Ser.

OZOLOTEPEC FIELD MOUSE.

Type locality. Mountains near Ozolotepec, State of Oaxaca, Mexico.

Geogr. Distr. Known only from the type locality.

372. *Peromyscus melanocarpus* Osgood.

Peromyscus melanocarpus Osgood, Proc. Biol. Soc. Wash., xvii, 1904, p. 73. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 210. Zoöl. Ser.

BLACK-FOOTED MOUSE.

Type locality. Mount Zempoaltepec, State of Oaxaca, Mexico.

Geogr. Distr. Known only from the type locality.

373. *Peromyscus altilaneus* Osgood.

Peromyscus altilaneus Osgood, Proc. Biol. Soc. Wash., xvii, 1904, p. 74. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 210. Zoöl. Ser.

FLUFFY MOUSE.

Type locality. Todos Santos, Guatemala. Altitude, 10,000 feet.
Geogr. Distr. Guatemala, Central America.

374. *Peromyscus guatemalensis* Merriam.

Peromyscus guatemalensis Merr., Proc. Biol. Soc. Wash., XII, 1898,
p. 118. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M.
Pub., IV, Pt. I, 1904, p. 211. Zool. Ser.

TODOS SANTOS MOUSE.

Type locality. Todos Santos, Guatemala. Altitude, 10,000 feet.
Geogr. Distr. Guatemala, Central America.

375. *Peromyscus cacabatus* Bangs.

Peromyscus cacabatus Bangs, Bull. Mus. Comp. Zool., XXXIX, 1902,
p. 29. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M.
Pub., IV, Pt. I, 1904, p. 211. Zool. Ser.

SOOTY MOUSE.

Type locality. Boquete, Chiriqui, Panama.
Geogr. Distr. Panama.

376. *Peromyscus mecisturus* Merriam.

Peromyscus mecisturus Merr., Proc. Biol. Soc. Wash., XII, 1898,
p. 124.

Peromyscus mecisturus Elliot, Mamm. Middle Amer. & W. Indies,
F. C. M. Pub., IV, Pt. I, 1904, p. 212. Zool. Ser.

GREAT-TAILED MOUSE.

Type locality. Chalchicomula, State of Puebla, Mexico. Altitude,
8,400 feet.

Geogr. Distr. State of Puebla, Mexico.

Genus 61. **Megadontomys.**

Megadontomys Merr., Proc. Biol. Soc. Wash., XII, 1898, p. 115.

Type Peromyscus thomasi Merriam.

377. *Megadontomys thomasi* (Merriam).

Peromyscus thomasi Merr., Proc. Biol. Soc. Wash., XII, 1898, p.
116.

Megadontomys thomasi Elliot, Mamm. Middle Amer. & W. Indies,
F. C. M. Pub., IV, Pt. I, 1904, p. 213, fig. 34. Zool. Ser.

THOMAS' GUERRERO FIELD MOUSE.

Type locality. Mountains near Chilpancingo, State of Guerrero,
Mexico. Altitude, 9,700 feet.

Geogr. Distr. State of Guerrero, Mexico. Range unknown.

378. *Megadontomys nelsoni* (Merriam).

Peromyscus nelsoni Merr., Proc. Biol. Soc. Wash., xii, 1898, p. 116.

Megadontomys nelsoni Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 214. Zoöl. Ser.

NELSON'S FIELD MOUSE.

Type locality. Jico, State of Vera Cruz, Mexico. Altitude, 6,000 feet.

Geogr. Distr. State of Vera Cruz, Mexico. Range unknown.

379. *Megadontomys flavidus* Bangs.

Megadontomys flavidus Bangs, Bull. Mus. Comp. Zoöl., xxxix, 1902, p. 27. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 214. Zoöl. Ser.

BOQUETE FIELD MOUSE.

Type locality. Boquete, Chiriqui, Panama. Altitude, 4,000 feet.

Geogr. Distr. Panama, Central America.

Genus 62. ***Nyctomys*.**

***Nyctomys* Sauss., Rev. Mag. Zool., 2me Sér., xii, 1860, p. 106. Type *Hesperomys sumichrasti* Saussure.**

Myxomys Tomes, Proc. Zoöl. Soc., 1861, p. 284, pl. xxxi.

380. *Nyctomys sumichrasti* (Saussure).

Hesperomys sumichrasti Sauss., Rev. Mag. Zool., 2me Sér., 1860, p. 107, pl. 9, fig. 2.

Nyctomys sumichrasti Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 216, fig. 35. Zoöl. Ser.

SUMICHRAST'S VESPER RAT.

Type locality. "Habite le versant oriental de la Cordillère," State of Vera Cruz? Mexico.

Geogr. Distr. Eastern Mexico, State of Vera Cruz to Isthmus of Tehuantepec, Mexico.

381. *Nyctomys decolorus* (True).

Sitomys (Rhipidomys) decolorus True, Proc. U. S. Nat. Mus., xvi, 1894, p. 689.

Nyctomys decolorus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 216. Zoöl. Ser.

FADED VESPER RAT.

Type locality. Rio de las Piedras, Honduras.

Geogr. Distr. Tehuantepec, Mexico, into Honduras, Central America.

382. *Nyctomys nitellinus* Bangs.

Nyctomys nitellinus Bangs, Bull. Mus. Comp. Zool., xxxix, 1902, p. 30. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. I, 1904, p. 216. Zool. Ser.

BOQUETE VESPER RAT.

Type locality. Boquete, Chiriqui, Panama. Altitude, 4,000 feet.
Geogr. Distr. Panama, Central America.

Genus 63. *Tylomys*.

***Tylomys* Peters**, Monatsb. K. Preuss. Akad. Wiss. Berl., 1866, p. 404. Type *Hesperomys nudicaudus* Peters.

383. *Tylomys nudicaudus* (Peters).

Hesperomys nudicaudus Peters, Monatsb. K. Preuss. Akad. Wiss. Berl., 1866, p. 404, pl. 1, figs. 1-4.

Tylomys nudicaudus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. I, 1904, p. 217, fig. 36. Zool. Ser.

NAKED-TAILED RAT.

Type locality. Guatemala.

Geogr. Distr. Southeastern Mexico (State of Chiapas), into Guatemala.

384. *Tylomys tumbalensis* Merriam.

Tylomys tumbalensis Merr., Proc. Wash. Acad. Scien., iii, 1901, p. 560. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. I, 1904, p. 218. Zool. Ser.

TUMBALA RAT.

Type locality. Tumbala, State of Chiapas, Mexico.

Geogr. Distr. State of Chiapas, Mexico.

385. *Tylomys bullaris* Merriam.

Tylomys bullaris Merr., Proc. Wash. Acad. Scien., iii, 1901, p. 561. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. I, 1904, p. 219. Zool. Ser.

TUXTLA RAT.

Type locality. Tuxtla, State of Chiapas, Mexico.

Geogr. Distr. State of Chiapas, Mexico.

386. *Tylomys watsoni* Thomas.

Tylomys watsoni Thomas, Ann. Mag. Nat. Hist., 7th Ser., iv, 1897, p. 278. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. I, 1904, p. 219. Zool. Ser.

WATSON'S RAT.

Type locality. Bogava, Chiriqui, Panama. Altitude, 8,000 feet.
Geogr. Distr. Panama, Central America.

387. *Tylomys panamensis* (Gray).

Neomys panamensis Gray, Ann. Mag. Nat. Hist., 4th Ser., XII, 1873, p. 417.

Tylomys panamensis Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 219. Zoöl. Ser.

PANAMA RAT.

Type locality. Panama.
Geogr. Distr. Panama, Central America.

Genus 64. *Otodylomys*.

Otodylomys Merr., Proc. Wash. Acad. Scien., III, 1901, p. 561.
Type Otodylomys phyllotis Merriam.

388. *Otodylomys phyllotis* Merriam.

Otodylomys phyllotis Merr., Proc. Wash. Acad. Scien., III, 1901, p. 562. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 221, fig. 37. Zoöl. Ser.

TUNKAS RAT.

Type locality. Tunkas, Yucatan, Mexico.
Geogr. Distr. Yucatan.

a. *Otodylomys phyllotis phæus* Merriam.

Otodylomys phyllotis phæus Merr., Proc. Wash. Acad. Scien., III, 1901, p. 563. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 221. Zoöl. Ser.

APAZOTE RAT.

Type locality. Apazote, near Yohaltun, State of Campeche, Mexico.
Geogr. Distr. State of Campeche, Mexico.

Genus 65. *Sigmodon*. Cotton Rats.

Sigmodon Say and Ord, Journ. Acad. Nat. Scien. Phil., IV, 1825, p. 352, pl. xxii, figs. 5-8. Type *Sigmodon hispidus* Say and Ord.

389. *Sigmodon hispidus* Say and Ord.

Sigmodon hispidus Say and Ord, Journ. Acad. Nat. Scien. Phil., 1825, p. 354. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 144, fig. 31. Zoöl. Ser.

Arvicola ferrugineus Harl., Am. Journ. Scien., 1828, p. 285.

Arvicola hortensis Harl., Med. & Phys. Res., 1835, p. 49, pl., fig. 58.

HISPID COTTON RAT.

Type locality. St. John's River, east Florida.

Geogr. Distr. North Carolina to northern Florida and west to southern Louisiana.

a. **Sigmodon hispidus littoralis* Chapman.

Sigmodon hispidus littoralis Chapm., Bull. Am. Mus. Nat. Hist.

N. Y., 1889, p. 118. Elliot, Syn. N. Am. Mamm., F. C. M.

Pub., 11, 1901, p. 144. Zool. Ser.

COAST COTTON RAT.

Type locality. Micco, Brevard County, Florida.

Geogr. Distr. Southeastern Florida.

b. *Sigmodon hispidus spadicipygus* Bangs.

Sigmodon hispidus spadicipygus Bangs, Proc. Bost. Soc. Nat. Hist., 1898, p. 192.

Sigmodon spadicipygus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 145. Zool. Ser.

BROWN-RUMPED COTTON RAT.

Type locality. Cape Sable, Monroe County, Florida.

Geogr. Distr. Extreme southern part of the peninsula of Florida.

c. *Sigmodon hispidus texensis* (Audubon & Bachman).

Arvicola texiana! Aud. & Bach., Quad. N. Am., 111, 1853, p. 229.

Sigmodon hispidus texensis Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 144. Zool. Ser.

TEXAN COTTON RAT.

Type locality. Brazos River, Texas.

Geogr. Distr. Eastern Texas as far west as Vernon and San Antonio, in Wilbarger and Bexar Counties, Texas, north to Cairo, Pratt County, Kansas.

d. *Sigmodon hispidus berlandieri* (Baird).

Sigmodon berlandieri Baird, Proc. Acad. Nat. Scien. Phil., 1855, p. 333.

Sigmodon hispidus pallidus Mearns, Proc. U. S. Nat. Mus., 1897, p. 504. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 145. Zool. Ser. *Id.* Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., 14, Pt. 1, 1904, p. 228. Zool. Ser.

Sigmodon hispidus berlandieri Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., 14, Pt. 1, 1904, p. 227. Zool. Ser.

*Very doubtfully separable from *S. hispidus*. See F. C. M. Pub., 11, 1901, p. 42.

BERLANDIER'S COTTON RAT.

Type locality. Rio Nazas, State of Coahuila, Mexico.

Geogr. Distr. Rio Grande and Pecos Valleys from Brownsville and El Paso, Texas, and Carlsbad, New Mexico, south to southern part of State of Jalisco, Mexico.

e. *Sigmodon hispidus eremicus* Mearns.

Sigmodon hispidus eremicus Mearns, Proc. U. S. Nat. Mus., 1897, p. 504. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 145. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 227. Zoöl. Ser.

WESTERN DESERT COTTON RAT.

Type locality. Cienega Well, State of Sonora, Mexico, on east bank of the Colorado River, 30 miles south of monument No. 204 on the United States and Mexico boundary line.

Geogr. Distr. Both sides of the Lower Colorado River.

f. *Sigmodon hispidus arizonæ* Mearns.

Sigmodon hispidus arizonæ Mearns, Bull. Am. Mus. Nat. Hist. N. Y., 1890, p. 287. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 144. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 228. Zoöl. Ser.

ARIZONA COTTON RAT.

Type locality. Fort Verde, Yavapai County, Arizona.

Geogr. Distr. Arizona, limits of range unknown.

g. *Sigmodon hispidus mascotensis* (Allen).

Sigmodon mascotensis Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1897, p. 54.

Sigmodon colimæ Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1897, p. 55.

Sigmodon hispidus mascotensis Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 227. Zoöl. Ser.

MASCOTA COTTON RAT.

Type locality. San Sebastian, near Mascota, State of Jalisco, Mexico.

Geogr. Distr. West coast of Mexico from western part of the State of Jalisco to southern portion of the State of Oaxaca.

h. *Sigmodon hispidus tonalensis* Bailey.

Sigmodon hispidus tonalensis Bailey, Proc. Biol. Soc. Wash., xv, 1902, p. 109. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 229. Zoöl. Ser.

TONALA COTTON RAT.

Type locality. Tonalá, State of Chiapas, Mexico.

Geogr. Distr. Western part of the State of Chiapas, and eastern part of the State of Oaxaca, Mexico.

i. *Sigmodon hispidus inexoratus* Elliot.

Sigmodon hispidus inexoratus Elliot, Pub. Field. Columb. Mus., III, 1903, p. 144. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 229, fig. xxxviii. Zoöl. Ser.

OCOTLAN COTTON RAT.

Type locality. Ocotlán, State of Jalisco, Mexico.

Geogr. Distr. State of Jalisco, Mexico. Range unknown.

j. *Sigmodon hispidus major* Bailey.

Sigmodon hispidus major Bailey, Proc. Biol. Soc. Wash., xv, 1902, p. 109. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 228. Zoöl. Ser.

LARGE COTTON RAT.

Type locality. Sierra de Choix, 50 miles northeast of Choix, State of Sinaloa, Mexico.

Geogr. Distr. Province of Tepic to southern part of the State of Sonora, Mexico.

k. *Sigmodon hispidus baileyi* (Allen).

Sigmodon baileyi Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1903, p. 601.

Sigmodon hispidus baileyi Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 226. Zoöl. Ser.

BAILEY'S COTTON RAT.

Type locality. La Ciénega de las Vacas, State of Durango, Mexico. Altitude, 8,500 feet.

Geogr. Distr. State of Durango, Mexico. Range unknown.

l. *Sigmodon hispidus toltecus* (Saussure).

Hesperomys toltecus Sauss., Rev. Mag. Zool., 2me Sér., 1860, p. 98.

Sigmodon hispidus toltecus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 226. Zoöl. Ser.

TOLTEC COTTON RAT.

Type locality. "Cordilleras of Vera Cruz," Mexico.

Geogr. Distr. Eastern Mexico from Alta Mira, southern part of the State of Tamaulipas, to Orizaba, State of Vera Cruz, Mexico.

m. *Sigmodon hispidus saturatus* Bailey.

Sigmodon hispidus saturatus Bailey, Proc. Biol. Soc. Wash., xv, 1902, p. 111. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 225. Zoöl. Ser.

TEAPA COTTON RAT.

Type locality. Teapa, State of Tabasco, Mexico.

Geogr. Distr. States of Tabasco, Chiapas, and Vera Cruz.

n. *Sigmodon hispidus furrus* Bangs.

Sigmodon hispidus furrus Bangs, Bull. Mus. Comp. Zool., xxxiv, 1903, p. 158. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 225. Zool. Ser.

CEIBA COTTON RAT.

Type locality. Ceiba, Honduras.

Geogr. Distr. Honduras. Range unknown.

o. *Sigmodon hispidus microdon* Bailey.

Sigmodon hispidus microdon Bailey, Proc. Biol. Soc. Wash., xv, 1902, p. 111. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 226. Zool. Ser.

SMALL-TOOTHED COTTON RAT.

Type locality. Puerto Morelos, Yucatan.

Geogr. Distr. Northern Yucatan and State of Campeche, Mexico.

p. *Sigmodon hispidus borucæ* (Allen).

Sigmodon borucæ Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1897, p. 40.

Sigmodon hispidus borucæ Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 224, fig. 38. Zool. Ser.

BORUCA COTTON RAT.

Type locality. Boruca, Costa Rica, Central America.

Geogr. Distr. Costa Rica, limits of range unknown.

q. *Sigmodon hispidus chiriquensis* (Allen).

Sigmodon borucæ chiriquensis Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1904, p. 68.

Sigmodon hispidus chiriquensis Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 225. Zool. Ser.

BOQUERON COTTON RAT.

Type locality. Boqueron, Chiriqui, Panama.

Geogr. Distr. Panama, Central America.

390. *Sigmodon minimus* Mearns.

Sigmodon minimus Mearns, Proc. U. S. Nat. Mus., xvii, 1894, p. 130. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., ii, 1901, p. 146. Zool. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 230. Zool. Ser.

LEAST COTTON RAT.

Type locality. Northern boundary line between New Mexico and Mexico, 100 miles west of the Initial Monument in Grant County, on the west bank of the Rio Grande.

Geogr. Distr. Mountains of southern New Mexico, Arizona, and northern Mexico.

391. *Sigmodon alticola* Bailey.

Sigmodon alticola Bailey, Proc. Biol. Soc. Wash., xv, 1902, p. 116.

Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 231. Zoöl. Ser.

ALPINE COTTON RAT.

Type locality. Cerro San Felipe, State of Oaxaca, Mexico. Altitude, 10,000 feet.

Geogr. Distr. Mountains in State of Oaxaca, Mexico.

a. Sigmodon alticola amoles Bailey.

Sigmodon alticola amoles Bailey, Proc. Biol. Soc. Wash., xv, 1902,

p. 116. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 231. Zoöl. Ser.

AMOLES COTTON RAT.

Type locality. Pinal de Amoles, State of Queretaro, Mexico.

Geogr. Distr. Known only from the type locality.

392. *Sigmodon ochrognathus* Bailey.

Sigmodon ochrognathus Bailey, Proc. Biol. Soc. Wash., xv, 1902,

p. 115. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 230. Zoöl. Ser.

OCHRACEOUS-FACED COTTON RAT.

Type locality. Chisos Mountains, Brewster County, Texas. Altitude, 8,000 feet.

Geogr. Distr. Chisos Mountains, Texas, south of Parral, State of Chihuahua, Mexico.

393. *Sigmodon leucotis* Bailey.

Sigmodon leucotis Bailey, Proc. Biol. Soc. Wash., xv, 1902, p. 115.

Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 230. Zoöl. Ser.

WHITE-EARED COTTON RAT.

Type locality. Valparaiso Mountains, State of Zacatecas, Mexico. Altitude, 8,700 feet.

Geogr. Distr. Known only from the type locality.

394. *Sigmodon alleni* Bailey.

Sigmodon alleni Bailey, Proc. Biol. Soc. Wash., xv, 1902, p. 112.

Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 224. Zoöl. Ser.

ALLEN'S COTTON RAT.

Type locality. San Sebastian, Mascota, State of Jalisco, Mexico.

Geogr. Distr. Western portion of State of Jalisco, and southern part of Province of Tepic, Mexico.

395. *Sigmodon fulviventer* Allen.

Sigmodon fulviventer Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1889, p. 180. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 232. Zoöl. Ser.

FULVOUS-BELLIED COTTON RAT.

Type locality. Zacatecas, State of Zacatecas, Mexico.

Geogr. Distr. States of Durango and Zacatecas, Mexico.

396. *Sigmodon melanotis* Bailey.

Sigmodon melanotis Bailey, Proc. Biol. Soc. Wash., xv, 1902, p. 114. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 232. Zoöl. Ser.

BLACK-EARED COTTON RAT.

Type locality. Patzcuaro, State of Michoacan, Mexico.

Geogr. Distr. State of Michoacan, range unknown.

397. *Sigmodon austerulus* Bangs.

Sigmodon austerulus Bangs, Bull. Mus. Comp. Zoöl., xxxix, 1902, p. 32. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 231. Zoöl. Ser.

VOLCAN DE CHIRIQUI COTTON RAT.

Type locality. Volcan de Chiriqui, Chiriqui, Panama. Altitude, 10,000 feet.

Geogr. Distr. Panama, Central America.

Genus 66. **Oryzomys. Rice Rats.**

Oryzomys Baird, N. Am. Mamm., 1857, p. 458. Type *Mus palustris* Harlan.

Oligoryzomys Bangs, Proc. N. Eng. Zoöl. Club, 1, 1900, p. 94, pl. 1, fig. 2.

Subgenus A. **Oryzomys.**

398. *Oryzomys palustris* (Harlan).

Mus palustris Harl., Am. Journ. Scien. & Arts, xxxi, 1837, pp. 385-386.

Hesperomys palustris Wagn., Schreb. Säugth., Suppl., III, 1843, p. 543.

Arvicola oryzivora Aud. & Bachm., N. Am. Quad., III, 1853, p. 214, pl. CXLIV, fig. 3.

Oryzomys palustris texensis Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1894, p. 177. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 147. Zoöl. Ser.

Oryzomys palustris Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 147. Zoöl. Ser.

SWAMP RICE RAT.

Type locality. "Fast Land," near Salem, Salem County, Delaware Bay, southern New Jersey.

Geogr. Distr. New Jersey to Georgia.

a. *Oryzomys palustris coloratus* Bangs.

Oryzomys palustris coloratus Bangs, Proc. Bost. Soc. Nat. Hist., 1898, p. 189.

Oryzomys aquaticus coloratus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 148. Zoöl. Ser.

CAPE SABLE RICE RAT.

Type locality. Cape Sable, Florida.

Geogr. Distr. Southern portion of the Florida Peninsula.

399. **Oryzomys natator* (Chapman).

Oryzomys palustris natator Chapman, Bull. Am. Mus. Nat. Hist. N. Y., 1893, p. 44. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 147. Zoöl. Ser.

SWIMMING RICE RAT.

Type locality. Gainesville, Alachua County, Florida.

Geogr. Distr. Florida west to Texas.

a. *Oryzomys natator floridanus* Merriam.

Oryzomys natator floridanus Merr., Proc. Wash. Acad. Scien., III, 1901, p. 277.

FLORIDA RICE RAT.

Type locality. Everglade, Monroe County, Florida.

Geogr. Distr. Florida.

Genl. Char. Similar to *O. natator* but smaller. Skull shorter, zygomata broadly spreading, outer sides nearly parallel.

Color. Like *O. natator*, but the cheeks and sides more inclining to fulvous.

Measurements. Average of 6 males: total length, 267; tail vertebrae, 137; hind foot, 33.

400. *Oryzomys aquaticus* Allen.

Oryzomys aquaticus Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1891, p. 289. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 147, fig. 32. Zoöl. Ser.

*Raised to specific rank by Dr. Merriam, Proc. Wash. Acad. Scien., III, 1901, p. 277.

AQUATIC RICE RAT.

Type locality. Brownsville, Cameron County, Texas.

Geogr. Distr. Southeastern Texas; limits of range unknown.

401. *Oryzomys peninsulæ* Thomas.

Oryzomys peninsulæ Thomas, Ann. Mag. Nat. Hist., 6th Ser., xx, 1897, p. 548. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 236. Zoöl. Ser.

LOWER CALIFORNIA RICE RAT.

Type locality. Santa Anita, Lower California, Mexico.

Geogr. Distr. Cape Region, Lower California, Mexico.

402. *Oryzomys nelsoni* Merriam.

Oryzomys nelsoni Merr., Proc. Biol. Soc. Wash., XII, 1898, p. 15. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 235. Zoöl. Ser.

MARIA MADRE ISLAND RICE RAT.

Type locality. Maria Madre Island, Tres Marias Islands, State of Jalisco, Mexico.

Geogr. Distr. Tres Marias Islands, State of Jalisco, Mexico.

403. *Oryzomys albiventer* Merriam.

Oryzomys albiventer Merr., Proc. Wash. Acad. Scien., III, 1901, p. 279. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 236. Zoöl. Ser.

WHITE-BELLIED RICE RAT.

Type locality. Ameca, State of Jalisco, Mexico.

Geogr. Distr. State of Jalisco, Mexico. Extent of range unknown.

404. *Oryzomys cozumelæ* Merriam.

Oryzomys cozumelæ Merr., Proc. Biol. Soc. Wash., XIV, 1901, p. 103. *Id.* Proc. Wash. Acad. Scien., III, 1901, p. 280. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 241. Zoöl. Ser.

COZUMEL ISLAND RICE RAT.

Type locality. Cozumel Island, Yucatan, Mexico.

Geogr. Distr. Known only from the type locality.

405. *Oryzomys panamensis* Thomas.

Oryzomys panamensis Thomas, Ann. Mag. Nat. Hist., 7th Ser., VIII, 1901, p. 252. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 241. Zoöl. Ser.

PANAMA RICE RAT.

Type locality. Near City of Panama, Panama.

Geogr. Distr. Panama, Central America.

406. *Oryzomys talamancæ* Allen.

Oryzomys talamancæ Allen, Proc. U. S. Nat. Mus., xiv, 1891, p. 193. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 241. Zoöl. Ser.

TALAMANCA RICE RAT.

Type locality. Talamanca, Costa Rica.

Geogr. Distr. Costa Rica, Central America.

407. *Oryzomys alfari* (Allen).

Hesperomys alfari Allen, Bull. Am. Mus. Nat. Hist., N. Y., 1891, p. 214.

Oryzomys alfari Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 242. Zoöl. Ser.

ALFARO'S RICE RAT.

Type locality. San Carlos, Costa Rica, Central America.

Geogr. Distr. Costa Rica, Central America.

408. *Oryzomys fulgens* Thomas.

Oryzomys fulgens Thomas, Ann. Mag. Nat. Hist., 6th Ser., xi, 1893, p. 403. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 240. Zoöl. Ser.

SHINING RICE RAT.

Type locality. Unknown, "Mexico."

Geogr. Distr. Unknown.

409. *Oryzomys molestus* Elliot.

Oryzomys molestus Elliot, Pub. Field Columb. Mus., iii, 1903, p. 145. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 240, fig. xxxix. Zoöl. Ser.

OCOTLAN RICE RAT.

Type locality. Ocotlan, State of Jalisco, Mexico.

Geogr. Distr. State of Jalisco, Mexico; range unknown.

410. *Oryzomys crinitus* Merriam.

Oryzomys crinitus Merr., Proc. Wash. Acad. Scien., iii, 1901, p. 281. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 245. Zoöl. Ser.

LONG-HAIRED RICE RAT.

Type locality. Talpam, Federal District, Mexico.

Geogr. Distr. Unknown.

*a. *Oryzomys crinitus aztecus* Merriam.*

Oryzomys crinitus aztecus Merr., Proc. Wash. Acad. Scien., iii, 1901, p. 282. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 245. Zoöl. Ser.

AZTEC RICE RAT.

Type locality. Yautepec, State of Morelos, Mexico.

Geogr. Distr. Southeastern Mexico, from States of Morelos and Puebla to Isthmus of Tehuantepec.

411. *Oryzomys mexicanus* Allen.

Oryzomys mexicanus Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1897, p. 52. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 242. Zoöl. Ser.

TONILA RICE RAT.

Type locality. Hacienda San Marcos, Tonila, State of Jalisco, Mexico.

Geogr. Distr. Southern part of the State of Sinaloa to Isthmus of Tehuantepec, Mexico.

a. *Oryzomys mexicanus peragrus* Merriam.

Oryzomys mexicanus peragrus Merr., Proc. Wash. Acad. Scien., III, 1901, p. 283. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 243. Zoöl. Ser.

WANDERING RICE RAT.

Type locality. Rio Verde, State of San Luis Potosi, Mexico.

Geogr. Distr. State of San Luis Potosi, Mexico. Limits of range unknown.

412. *Oryzomys richmondi* Merriam.

Oryzomys richmondi Merr., Proc. Wash. Acad. Scien., III, 1901, p. 284. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 248. Zoöl. Ser.

ESCONDIDO RIVER RICE RAT.

Type locality. Escondido River, 50 miles above Bluefields, Nicaragua.

Geogr. Distr. Nicaragua. Range unknown.

413. *Oryzomys jalapæ* Allen & Chapman.

Oryzomys jalapæ Allen & Chapman, Bull. Am. Mus. Nat. Hist. N. Y., 1897, p. 206. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 246. Zoöl. Ser.

JALAPA RICE RAT.

Type locality. Jalapa, State of Vera Cruz, Mexico.

Geogr. Distr. State of Vera Cruz, and northwestern part of State of Puebla, Mexico.

a. *Oryzomys jalapæ apatelius* Elliot.

Oryzomys jalapæ apatelius Elliot, Pub. Field Columb. Mus., III, 1904, p. 266. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 246. Zoöl. Ser.

DECEITFUL RICE RAT.

Type locality. San Carlos, State of Vera Cruz, Mexico.

Geogr. Distr. State of Vera Cruz, Mexico. Range unknown.

b. *Oryzomys jalapæ rufinus* Merriam.

Oryzomys jalapæ rufinus Merr., Proc. Wash. Acad. Scien., III, 1901, p. 285. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 247. Zoöl. Ser.

CATEMACO RICE RAT.

Type locality. Catemaco, State of Vera Cruz, Mexico.

Geogr. Distr. State of Vera Cruz, Mexico. Range unknown.

414. *Oryzomys zygomatus* Merriam.

Oryzomys zygomatus Merr., Proc. Wash. Acad. Scien., III, 1901, p. 285. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 241. Zoöl. Ser.

GUATEMALAN RICE RAT.

Type locality. Nentón, Guatemala.

Geogr. Distr. Guatemala.

415. *Oryzomys teapensis* Merriam.

Oryzomys teapensis Merr., Proc. Acad. Scien. Wash., III, 1901, p. 286. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 247. Zoöl. Ser.

TEAPA RICE RAT.

Type locality. Teapa, State of Tabasco, Mexico.

Geogr. Distr. State of Tabasco, possibly also the State of Chiapas, Mexico.

416. *Oryzomys bulleri* Allen.

Oryzomys bulleri Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1897, p. 53. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 239. Zoöl. Ser.

BULLER'S RICE RAT.

Type locality. Valle de Banderas, Territorio de Tepic, State of Jalisco, Mexico.

Geogr. Distr. States of Jalisco and Colima, Mexico.

417. *Oryzomys rufus* Merriam.

Oryzomys rufus Merr., Proc. Wash. Acad. Scien., III, 1901, p. 287. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 239. Zoöl. Ser.

RUFIOUS RICE RAT.

Type locality. Santiago, Territorio de Tepic, State of Jalisco, Mexico.

Geogr. Distr. Territorio de Tepic, State of Jalisco, Mexico.

418. *Oryzomys couesi* (Alston).

Hesperomys couesi Alston, Proc. Zool. Soc., 1876, p. 756.

Oryzomys couesi Elliot, Mamm. Middle Amer. & W. Indies, F. C. M.

Pub., IV, Pt. 1, 1904, p. 236. Zool. Ser.

COUES' RICE RAT.

Type locality. Coban, Guatemala.

Geogr. Distr. State of Chiapas, Mexico, into Guatemala, Central America.

419. *Oryzomys goldmani* Merriam.

Oryzomys goldmani Merr., Proc. Wash. Acad. Scien., III, 1901,

p. 288. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M.

Pub., IV, Pt. 1, 1904, p. 246. Zool. Ser.

GOLDMAN'S RICE RAT.

Type locality. Coatzacoalcos, State of Vera Cruz, Mexico.

Geogr. Distr. State of Vera Cruz. Range unknown.

420. *Oryzomys chapmani* Thomas.

Oryzomys chapmani Thomas, Ann. Mag. Nat. Hist., 7th Ser., 1,

1898, p. 179. Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., IV, Pt. 1, 1904, p. 238. Zool. Ser.

CHAPMAN'S RICE RAT.

Type locality. Jalapa, State of Vera Cruz, Mexico.

Geogr. Distr. State of Vera Cruz, Mexico.

a. *Oryzomys chapmani caudatus* Merriam.

Oryzomys chapmani caudatus Merr., Proc. Wash. Acad. Scien., III,

1901, p. 289. Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., IV, Pt. 1, 1904, p. 238. Zool. Ser.

LONG-TAILED RICE RAT.

Type locality. Comaltepec, State of Oaxaca, Mexico.

Geogr. Distr. State of Oaxaca. Range unknown.

b. *Oryzomys chapmani dilutior* Merriam.

Oryzomys chapmani dilutior Merr., Proc. Wash. Acad. Scien., III,

1901, p. 290. Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., IV, Pt. 1, 1904, p. 238. Zool. Ser.

PALE RICE RAT.

Type locality. Huauchinango, State of Puebla, Mexico.

Geogr. Distr. State of Puebla, Mexico. Range unknown.

c. *Oryzomys chapmani saturator* Merriam.

Oryzomys chapmani saturator Merr., Proc. Wash. Acad. Scien.,

III, 1901, p. 290. Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., IV, Pt. 1, 1904, p. 239. Zool. Ser.

TUMBALA RICE RAT.

Type locality. Tumbala, State of Chiapas, Mexico.

Geogr. Distr. State of Chiapas, Mexico. Range unknown.

421. *Oryzomys palatinus* Merriam.

Oryzomys palatinus Merr., Proc. Wash. Acad. Scien., III, 1901, p. 290. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 237. Zool. Ser.

TABASCAN RICE RAT.

Type locality. Teapa, State of Tabasco, Mexico.

Geogr. Distr. State of Tabasco. Range unknown.

422. *Oryzomys hylocetes* Merriam.

Oryzomys hylocetes Merr., Proc. Wash. Acad. Scien., III, 1901, p. 291. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 237. Zool. Ser.

MARSH RICE RAT.

Type locality. Chicharras, State of Chiapas, Mexico.

Geogr. Distr. State of Chiapas, Mexico. Range unknown.

423. *Oryzomys rhabdops* Merriam.

Oryzomys rhabdops Merr., Proc. Wash. Acad. Scien., III, 1901, p. 291. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 237. Zool. Ser.

STRIPED-FACE RICE RAT.

Type locality. Cael, Guatemala. Altitude, 10,000 feet?

Geogr. Distr. State of Chiapas, Mexico, into Guatemala, Central America.

424. *Oryzomys angusticeps* Merriam.

Oryzomys angusticeps Merr., Proc. Wash. Acad. Scien., III, 1901, p. 292. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 245. Zool. Ser.

SANTA MARIA VOLCAN RICE RAT.

Type locality. Volcan Santa Maria, Guatemala. Altitude, 9,000 feet.

Geogr. Distr. Guatemala, Central America.

425. *Oryzomys melanotis* Thomas.

Oryzomys melanotis Thomas, Ann. Mag. Nat. Hist., 6th Ser., XI, 1893, p. 404. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 243. Zool. Ser.

BLACK-EARED RICE RAT.

Type locality. Mineral San Sebastian, State of Jalisco, Mexico.

Geogr. Distr. State of Jalisco, Mexico.

426. *Oryzomys rostratus* Merriam.

Oryzomys rostratus Merr., Proc. Wash. Acad. Scien., III, 1901, p. 293. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 243. Zoöl. Ser.

BROAD-NOSED RICE RAT.

Type locality. Metlaltoyuca, State of Puebla, Mexico.

Geogr. Distr. Eastern Mexico from State of Puebla to Isthmus of Tehuantepec.

a. *Oryzomys rostratus megadon* Merriam.

Oryzomys rostratus megadon Merr., Proc. Wash. Acad. Scien., III, 1901, p. 294. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 243. Zoöl. Ser.

LARGE-TOOTHED RICE RAT.

Type locality. Teapa, State of Tabasco, Mexico.

Geogr. Distr. State of Tabasco, westward probably to State of Puebla, Mexico.

427. *Oryzomys victus* Thomas.

Oryzomys victus Thomas, Ann. Mag. Nat. Hist., 7th Ser., I, 1898, p. 177. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 244. Zoöl. Ser.

ST. VINCENT RICE RAT.

Type locality. Island of St. Vincent, Lesser Antilles.

Geogr. Distr. Island of St. Vincent.

428. *Oryzomys devius* Bangs.

Oryzomys devius Bangs, Bull. Mus. Comp. Zoöl., XXXIX, 1902, p. 34. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 244. Zoöl. Ser.

LONELY RICE RAT.

Type locality. Boquete, Chiriqui, Panama. Altitude, 5,000 feet.

Geogr. Distr. Panama, Central America.

429. *Oryzomys antillarum* Thomas.

Oryzomys antillarum Thomas, Ann. Mag. Nat. Hist., 7th Ser., I, 1898, p. 177. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 247. Zoöl. Ser.

JAMAICA RICE RAT.

Type locality. Island of Jamaica.

Geogr. Distr. Known only from the type locality.

430. *Oryzomys tectus* Thomas.

Oryzomys tectus Thomas, Ann. Mag. Nat. Hist., 7th Ser., VIII, 1901, p. 251. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 245. Zoöl. Ser.

BOGAVA RICE RAT.

Type locality. Bogava, Chiriqui, Panama. Altitude, 800 feet.

Geogr. Distr. Panama, Central America.

431. *Oryzomys yucatanensis* Merriam.

Oryzomys yucatanensis Merr., Proc. Wash. Acad. Scien., III, 1901,

p. 294. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M.

Pub., IV, Pt. 1, 1904, p. 236. Zoöl. Ser.

CHICHEN ITZA RICE RAT.

Type locality. Chichen Itza, Yucatan, Mexico.

Geogr. Distr. Yucatan, Mexico.

432. *Oryzomys fulvescens* (Saussure).

Hesperomys fulvescens Sauss., Rev. Mag. Zool., 2me Sér., XII, 1860,

p. 102.

Oryzomys fulvescens Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., IV, Pt. 1, 1904, p. 248. Zoöl. Ser.

FULVOUS RICE RAT.

Type locality. "Mexique," Orizaba, State of Vera Cruz? Mexico.

Geogr. Distr. Jalapa and vicinity, State of Vera Cruz, south to Yucatan, Mexico, and Guatemala, Central America.

Subgenus B. *Oligoryzomys*.**433. *Oryzomys costaricensis* Allen.**

Oryzomys costaricensis Allen, Bull. Am. Mus. Nat. Hist. N. Y.,

1893, p. 239. Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., IV, Pt. 1, 1904, p. 249, fig. 39. Zoöl. Ser.

Oryzomys vegetus Bangs, Bull. Mus. Comp. Zoöl., XXXIX, 1902,

p. 35. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M.

Pub., IV, Pt. 1, 1904, p. 249, fig. 40. Zoöl. Ser.

EL GENERAL RICE RAT.

Type locality. El General, Costa Rica.

Geogr. Distr. Costa Rica, Central America. Altitude, from 2,150 to 10,342 feet.

Genus 67. **Moschophoromys. Musk-Bearing Rats.**

Moschophoromys Elliot, Pub. Field Columb. Mus., III, 1904, p. 270.

Zoöl. Ser. *Type Mus desmaresti* Fischer.

Megalomys Trouess., Le Naturaliste, III, No. 45, 1881, p. 5. *Id.*

Ann. Scien. Nat. Paris, Zool., XIX, No. 5, 1885, p. 13, pl. 1.

(nec D'Orbigny & Laurillard, 1842, Paleontology.)

Moschomys Trouess., Ann. Mag. Nat. Hist., 7th Ser., XI, 1903,

p. 387. (nec Billberg, 1828, Microtinæ.)

434. Moschophoromys desmaresti (Fischer).

Mus desmaresti Fischer, Syn. Mamm., 1829, p. 316.

Mus pilorides Desm., Dict. Scien. Nat., XLIV, 1826, p. 483. (nec Pallas ex Ceylon.)

Moschophoromys desmaresti Elliot, Mamm. Middle Amer. & W.

Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 250, fig. 41. Zoöl. Ser.

BLACK RICE RAT.

Type locality. Island of Martinique, West Indies.

Geogr. Distr. Known only from the Island of Martinique.

435. Moschophoromys luciae (Forsyth-Major).

Oryzomys luciae Forsyth-Major, Ann. Mag. Nat. Hist., 7th Ser., VII, 1901, p. 206.

Moschophoromys luciae Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., IV, Pt. I, 1904, p. 251. Zoöl. Ser.

ST. LUCIA RICE RAT.

Type locality. Island of St. Lucia, West Indies.

Geogr. Distr. Known only from the Island of St. Lucia.

Genus 68. Zygodontomys.

Zygodontomys Allen, Bull. Am. Mus. Nat. Hist. N. Y., IX, 1897, p. 38, pl. 1, figs. 1-7. Type "*Oryzomys cherriei*" Allen.

436. Zygodontomys cherrii (Allen).

Oryzomys cherriei Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1895, p. 329.

Zygodontomys cherrii Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., IV, Pt. I, 1904, p. 253, figs. 42, XL. Zoöl. Ser.

CHERRIE'S RICE RAT.

Type locality. Boruca, Costa Rica.

Geogr. Distr. Costa Rica, Central America. Range unknown.

437. Zygodontomys chrysomelas (Allen).

Oryzomys chrysomelas Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1897, p. 37.

Hesperomys (Habrothrix) caliginosus Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1891, p. 210. (nec Tomes.)

Zygodontomys chrysomelas Elliot, Mamm. Middle Amer. & W.

Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 253. Zoöl. Ser.

SUERRE RICE RAT.

Type locality. Suerre, Costa Rica.

Geogr. Distr. Costa Rica, Central America.

438. *Zygodontomys seorsus* Bangs.

Zygodontomys seorsus Bangs, Amer. Nat., xxxv, 1901, p. 642.
 Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv,
 Pt. 1, 1904, p. 254. Zoöl. Ser.

ISOLATED RICE RAT.

Type locality. San Miguel Island, Bay of Panama.
Geogr. Distr. Known only from San Miguel Island.

Genus 69. *Sigmodontomys*.

***Sigmodontomys* Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1897, p. 38.**
 pl. 1, figs. 8-14. Type *Sigmodontomys alfari* Allen.

439. *Sigmodontomys alfari* Allen.

Sigmodontomys alfari Allen, Bull. Am. Mus. Nat. Hist. N. Y.,
 1897, p. 39. Elliot, Mamm. Middle Amer. & W. Indies,
 F. C. M. Pub., iv, Pt. 1, 1904, p. 255. Zoöl. Ser.

ALFARO'S RICE RAT.

Type locality. Jimenez, Costa Rica. Altitude, 7,000 feet.
Geogr. Distr. Costa Rica, Central America. Range unknown.

Genus 70. *Rhithrodontomys*. Harvest Mice.

***Reithrodontomys* (sic) Giglioli, Ricer. intern. alla Distrib. Geogr. Gen.**
 Roma, 1873, p. 160. Type *Mus lecontei* Audubon & Bachman.

440. *Rhithrodontomys lecontei* (Audubon & Bachman).

Mus lecontei Aud. & Bachm., Jour. Acad. Nat. Scien. Phil., 1842,
 p. 307.

Reithrodontomys lecontei Elliot, Syn. N. Am. Mamm., F. C. M.
 Pub., II, 1901, p. 148, fig. 33. Zoöl. Ser.

LE CONTE'S HARVEST MOUSE.

Type locality. Probably near Riceboro, Liberty County, Georgia.
Geogr. Distr. Coast district of South Carolina and Georgia,
 southward into Florida.

a. *Rhithrodontomys lecontei impiger* Bangs.

Reithrodontomys lecontei impiger Bangs, Proc. Biol. Soc. Wash.,
 XII, 1898, p. 167. Elliot, Syn. N. Am. Mamm., F. C. M. Pub.,
 II, 1901, p. 149. Zoöl. Ser.

DILIGENT HARVEST MOUSE.

Type locality. White Sulphur Springs, Greenbrier County, West
 Virginia.

Geogr. Distr. West Virginia.

b. Rhithrodontomys lecontei dickinsoni (Rhoads).

Reithrodontomys humilis dickinsoni Rhoads, Amer. Nat., xxix, 1895, p. 590.

Reithrodontomys! lecontei dickinsoni Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 150. Zoöl. Ser.

DICKINSON'S HARVEST MOUSE.

Type locality. Willow Oak, Pasco County, Florida.

Geogr. Distr. Southern portion of the Peninsula of Florida.

441. *Rhithrodontomys merriami* Allen.

Reithrodontomys! merriami Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1895, p. 119. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 150. Zoöl. Ser.

MERRIAM'S HARVEST MOUSE.

Type locality. Austin Bayou, near Alvin, Brazoria County, Texas.

Geogr. Distr. Coast district of southwestern Louisiana to Brazoria County, Texas.

442. *Rhithrodontomys dychei* Allen.

Ochetodon humilis Coues, Proc. Acad. Nat. Scien. Phil., 1874, p. 185. (nec Aud. & Bachm.)

Reithrodontomys! dychei Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1895, p. 120. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 150. Zoöl. Ser.

DYCHE'S HARVEST MOUSE.

Type locality. Lawrence, Douglas County, Kansas.

Geogr. Distr. Kansas east to St. Louis, Missouri; south to western part of Oklahoma Territory; north to Nebraska and southwestern Iowa.

a. Rhithrodontomys dychei nebrascensis Allen.

Reithrodontomys! dychei nebrascensis Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1895, p. 122. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 150. Zoöl. Ser.

NEBRASKA HARVEST MOUSE.

Type locality. Kennedy, Cherry County, Nebraska.

Geogr. Distr. Fremont County, Colorado, north to Custer, Montana, east to central and northeastern Nebraska.

443. **Rhithrodontomys albescens* Cary.

Reithrodontomys! albescens Cary, Proc. Biol. Soc. Wash., xvi, 1903, p. 53.

*As the type locality of this form is practically that of *R. d. nebrascensis*, the two probably are not distinct.

WHITISH HARVEST MOUSE.

Type locality. Eighteen miles northwest of Kennedy, Cherry County, Nebraska.

Geogr. Distr. Northern Nebraska. Range unknown.

Genl. Char. Size small; tail short; color pale; ears with two black spots.

Color. Above plain gray, or tinged with buff; sides pale buffy gray; lateral line pinkish or cream buff; sides of nose, lower part of cheeks, throat, forelegs, and under parts pure white; tail bicolor; feet above white.

Measurements. Total length, 125; tail vertebræ, 53.5; hind foot, 16.6. Skull: basal length, 15.7; occipito-nasal length, 19.7; length of nasals, 17.4; zygomatic breadth, 10.5; mastoid breadth, 9.

444. *Rhithrodontomys montanus* (Baird).

Reithrodon! montanus Baird, Proc. Acad. Nat. Scien. Phil., 1855, p. 335.

Reithrodontomys! montanus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 151. Zool. Ser.

MOUNTAIN HARVEST MOUSE.

Type locality. Upper end of the San Luis Valley, Saguache County, Colorado.

Geogr. Distr. Known only from the type locality.

445. *Rhithrodontomys megalotis* (Baird).

Reithrodon! megalotis Baird, N. Am. Mamm., 1857, p. 451.

Reithrodontomys! aztecus Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1893, p. 79.

Reithrodontomys! megalotis deserti Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1895, p. 127.

Reithrodontomys! megalotis Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 151. Zool. Ser.

Rhithrodontomys megalotis Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 259, figs. 43, XLI. Zool. Ser.

BIG-EARED HARVEST MOUSE.

Type locality. Between Janos and San Luis Spring, State of Sonora, Mexico, near boundary of Grant County, New Mexico.

Geogr. Distr. States of Sonora and Chihuahua, Mexico, north to Nevada.

a. *Rhithrodontomys megalotis sestiniensis* Allen.

Reithrodontomys! megalotis sestiniensis Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1903, p. 602.

Rhithrodontomys megalotis sestinensis Elliot, Mamm. Middle Amer.

& W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 260. Zoöl. Ser.
RIO SESTIN MEADOW MOUSE.

Type locality. Rio Sestin, State of Durango, Mexico. Altitude,
7,500 feet.

Geogr. Distr. State of Durango, Mexico. Range unknown.

b. *Rhithrodontomys megalotis zacatecae* Merriam.

Reithrodontomys! megalotis zacatecae Merr., Proc. Wash. Acad.
Scien., III, 1901, p. 557.

Rhithrodontomys megalotis zacatecae Elliot, Mamm. Middle Amer.

& W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 260. Zoöl. Ser.
VALPARAISO HARVEST MOUSE.

Type locality. Valparaiso Mountains, State of Zacatecas, Mexico.

Geogr. Distr. State of Zacatecas, Mexico. Range unknown.

c. *Rhithrodontomys megalotis obscurus* Merriam.

Reithrodontomys! megalotis obscurus Merr., Proc. Wash. Acad.
Scien., III, 1901, p. 558.

Rhithrodontomys megalotis obscurus Elliot, Mamm. Middle Amer.

& W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 261. Zoöl. Ser.
DUSKY HARVEST MOUSE.

Type locality. Sierra Madre, near Guadalupe y Calvo, State of
Chihuahua, Mexico.

Geogr. Distr. State of Chihuahua, Mexico. Range unknown.

446. *Rhithrodontomys australis* Allen.

Reithrodontomys! australis Allen, Bull. Am. Mus. Nat. Hist. N. Y.,
1895, p. 328.

Rhithrodontomys australis Elliot, Mamm. Middle Amer. & W.

Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 259. Zoöl. Ser.
IRAZÚ VOLCANO HARVEST MOUSE.

Type locality. Volcano of Irazú, Costa Rica.

Geogr. Distr. Costa Rica, Central America. Range unknown.

a. *Rhithrodontomys australis vulcanius* Bangs.

Reithrodontomys! australis vulcanius Bangs, Bull. Mus. Comp.
Zoöl., XXXIX, 1902, p. 38.

Rhithrodontomys australis vulcanius Elliot, Mamm. Middle Amer.

& W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 259. Zoöl.
Ser.

CHIRIQUI HARVEST MOUSE.

Type locality. Volcan de Chiriqui, Chiriqui, Panama. Altitude,
10,300 feet.

Geogr. Distr. Panama. Range unknown

447. *Rhithrodontomys longicaudus* (Baird).

Reithrodon! longicauda Baird, N. Am. Mamm., 1857, p. 451.

Reithrodontomys! longicaudus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 151. Zoöl. Ser.

Rhithrodontomys longicaudus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 261. Zoöl. Ser.

LONG-TAILED HARVEST MOUSE.

Type locality. Petaluma, Sonora County, California.

Geogr. Distr. Lower California, Mexico, north to California; west to the Sierra Nevada from coast region of Monterey County; north to Mendocino and Humboldt Counties; and in the interior from San Joaquin County, north to Tehama County. Probably farther south irregularly in the Coast and San Bernardino ranges of mountains. (Allen.)

a. *Rhithrodontomys longicaudus pallidus* (Rhoads).

Reithrodontomys! pallidus Rhoads, Amer. Nat., 1893, p. 835.

Reithrodontomys! longicaudus pallidus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 152. Zoöl. Ser.

Rhithrodontomys longicaudus pallidus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 262. Zoöl. Ser.

LOWER CALIFORNIA OR PALLID HARVEST MOUSE.

Type locality. Santa Ysabel, San Jacinto Mountains, San Diego County, California.

Geogr. Distr. Southern California and northern Lower California.

448. *Rhithrodontomys chrysotis* Elliot.

Reithrodontomys! chrysotis Elliot, Pub. Field Columb. Mus., I, 1899, p. 281. Zoöl. Ser. *Id.* Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 152. Zoöl. Ser.

GOLDEN-EARED HARVEST MOUSE.

Type locality. Dougherty, Indian Territory.

Geogr. Distr. Known only from the type locality; and Alva, Oklahoma Territory.

449. *Rhithrodontomys catalinæ* Elliot.

Rhithrodontomys catalinæ Elliot, Pub. Field Columb. Mus., III, 1903, p. 246. Zoöl. Ser.

SANTA CATALINA ISLAND HARVEST MOUSE.

Type locality. Santa Catalina Island, Santa Barbara Islands, California.

Geogr. Distr. Known only from Santa Catalina Island.

Genl. Char. Similar to *R. longicaudus*, but larger; hind foot very much larger.

Color. Top of head and dorsal region brownish black or buff mixed, the black predominating; sides cream buff; lateral line indistinct cream buff; under parts, hands, and feet white, plumbeous of under fur showing; large cream buff spot on breast; tail above blackish, beneath soiled white; ears brown.

Measurements. Total length, 155.7; tail vertebræ, 83.8; hind foot, 19.5.

450. *Rhithrodontomys klamathensis* Merriam.

Reithrodontomys! klamathensis Merr., N. Am. Faun., No. 16, 1899, p. 93. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 152. Zoöl. Ser.

KLAMATH HARVEST MOUSE.

Type locality. Big Spring, Shasta Valley, Siskiyou County, California.

Geogr. Distr. Shasta Valley, California, and Klamath County, Oregon.

451. *Rhithrodontomys arizonensis* Allen.

Reithrodontomys! longicaudus Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1894, p. 320. Ex Arizona. (nec Baird.)

Reithrodontomys! arizonensis Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1895, p. 134. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 153. Zoöl. Ser.

ARIZONA HARVEST MOUSE.

Type locality. Chiricahua Mountains, Cochise County, Arizona. Altitude, 8,000 feet.

Geogr. Distr. Southeastern Arizona.

452. *Rhithrodontomys saturatus* Allen and Chapman.

Reithrodontomys! saturatus Allen & Chapman, Bull. Am. Mus. Nat. Hist. N. Y., 1897, p. 201.

Rhithrodontomys saturatus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 262. Zoöl. Ser.

LAS VIGAS HARVEST MOUSE.

Type locality. Las Vigas, State of Vera Cruz, Mexico.

Geogr. Distr. State of Vera Cruz, Mexico. Range unknown.

a. Rhithrodontomys saturatus cinereus Merriam.

Reithrodontomys! saturatus cinereus Merr., Proc. Wash. Acad. Scien., 111, 1901, p. 556.

Rhithrodontomys saturatus cinereus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 262. Zoöl. Ser.

CINEREOUS HARVEST MOUSE.

Type locality. Chalchicomula, State of Puebla, Mexico.

Geogr. Distr. State of Puebla, Mexico. Range unknown.

b. *Rhithrodontomys saturatus alticola* Merriam.

Reithrodontomys! saturatus alticola Merr., Proc. Wash. Acad. Scien., III, 1901, p. 556.

Rhithrodontomys saturatus alticola Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 263. Zoöl. Ser.

MOUNTAIN HARVEST MOUSE.

Type locality. Cerro San Felipe, near Oaxaca City, State of Oaxaca, Mexico. Altitude, 10,000 feet.

Geogr. Distr. State of Oaxaca. Range unknown.

453. *Rhithrodontomys peninsulæ* Elliot.

Rhithrodontomys peninsulæ Elliot, Pub. Field Columb. Mus., III, 1903, p. 164. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 263. Zoöl. Ser.

PENINSULAR HARVEST MOUSE.

Type locality. San Quentin, Lower California, Mexico.

Geogr. Distr. Known only from the type locality.

454. *Rhithrodontomys difficilis* Merriam.

Reithrodontomys! difficilis Merr., Proc. Wash. Acad. Scien., III, 1901, p. 556.

Rhithrodontomys difficilis Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 263. Zoöl. Ser.

CAPTIOUS HARVEST MOUSE.

Type locality. Orizaba, State of Vera Cruz, Mexico. Altitude, 4,500 feet.

Geogr. Distr. State of Vera Cruz. Range unknown.

455. *Rhithrodontomys mexicanus* (Saussure).

Reithrodon! mexicanus Sauss., Rev. Mag. Zool., 2me Sér., XII, 1860, p. 109, pl. IX, fig. 1.

Reithrodon! sumichrasti Sauss., Rev. Mag. Zool., 2me Sér., XIII, 1861, p. 3.

Ochetodon mexicanus Coues, Proc. Acad. Nat. Scien. Phil., 1874, p. 186.

Rhithrodontomys mexicanus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 264. Zoöl. Ser.

MEXICAN HARVEST MOUSE.

Type locality. Mountains in State of Vera Cruz.

Geogr. Distr. State of Vera Cruz, Mexico.

a. *Rhithrodontomys mexicanus aurantius* Allen.

Ochetodon mexicanus Coues, Mon. N. Am. Rod., 1877, p. 128.
(Louisiana specimens.)

Reithrodontomys! mexicanus aurantius Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1895, p. 137.

Reithrodontomys! arizonensis aurantius Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 153. Zoöl. Ser.

GOLDEN HARVEST MOUSE.

Type locality. Lafayette, Lafayette Parish, Louisiana.

Geogr. Distr. Coast region of Texas from Matagorda County northward, and thence eastward to Houma, Louisiana, and north to Beebe, Arkansas.

b. *Rhithrodontomys mexicanus intermedius* Allen.

Reithrodontomys! mexicanus intermedius Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1895, p. 136.

Reithrodontomys! arizonensis intermedius Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 153. Zoöl. Ser.

Rhithrodontomys mexicanus intermedius Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 264. Zoöl. Ser.

BROWNSVILLE HARVEST MOUSE.

Type locality. Brownsville, Cameron County, Texas.

Geogr. Distr. Northern Mexico, into Texas.

c. *Rhithrodontomys mexicanus fulvescens* Allen.

Reithrodontomys! mexicanus fulvescens Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1894, p. 319.

Rhithrodontomys mexicanus fulvescens Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 264. Zoöl. Ser.

OPOSURA HARVEST MOUSE.

Type locality. Oposura, State of Sonora, Mexico.

Geogr. Distr. State of Sonora, Mexico. Range unknown.

d. *Rhithrodontomys mexicanus gracilis* Allen & Chapman.

Reithrodontomys! mexicanus gracilis Allen & Chapman, Bull. Am. Mus. Nat. Hist. N. Y., 1897, p. 9.

Rhithrodontomys mexicanus gracilis Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 265. Zoöl. Ser.

CHICHEN ITZA HARVEST MOUSE.

Type locality. Chichen Itza, Yucatan, Mexico.

Geogr. Distr. Yucatan, Mexico.

456. *Rhithrodontomys laceyi* Allen.

Reithrodontomys! laceyi Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1896, p. 235. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 153. Zoöl. Ser.

LACEY'S HARVEST MOUSE.

Type locality. Watson's Ranch, fifteen miles south of San Antonio, Bexar County, Texas.

Geogr. Distr. Southeastern Texas.

457. *Rhithrodontomys tenuis* Allen.

Reithrodontomys! tenuis Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1899, p. 15.

Rhithrodontomys tenuis Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 265. Zoöl. Ser.

SLENDER HARVEST MOUSE.

Type locality. Rosario, State of Sinaloa, Mexico.

Geogr. Distr. State of Sinaloa, Mexico. Range unknown.

458. *Rhithrodontomys chrysopsis* Merriam.

Reithrodontomys! chrysopsis Merr., Proc. Biol. Soc. Wash., XIII, 1900, p. 152.

Rhithrodontomys chrysopsis Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 265. Zoöl. Ser.

MOUNT POPOCATEPETL HARVEST MOUSE.

Type locality. Mt. Popocatepetl, Mexico.

Geogr. Distr. State of Mexico. Range unknown.

a. *Rhithrodontomys chrysopsis tolucae* Merriam.

Reithrodontomys! chrysopsis tolucae Merr., Proc. Wash. Acad. Scien., III, 1901, p. 549.

Rhithrodontomys chrysopsis tolucae Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 266. Zoöl. Ser.

VOLCAN TOLUCA HARVEST MOUSE.

Type locality. North slope of the Volcan Toluca, State of Michoacan, Mexico. Altitude, 11,500 feet.

Geogr. Distr. Known only from the type locality.

459. *Rhithrodontomys perotensis* Merriam.

Reithrodontomys! perotensis Merr., Proc. Wash. Acad. Scien., III, 1901, p. 550.

Rhithrodontomys perotensis Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 266. Zoöl. Ser.

COFRE DE PEROTE HARVEST MOUSE.

Type locality. Cofre de Perote, State of Vera Cruz, Mexico. Altitude, 9,500 feet.

Geogr. Distr. Known only from the type locality.

460. *Rhithrodontomys orizabæ* Merriam.

Reithrodontomys! orizabæ Merr., Proc. Wash. Acad. Scien., III, 1901, p. 550.

Rhithrodontomys orizabæ Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 266. Zoöl. Ser.

ORIZABA HARVEST MOUSE.

Type locality. Mount Orizaba, State of Puebla, Mexico.

Geogr. Distr. Known only from the type locality.

461. *Rhithrodontomys colimæ* Merriam.

Reithrodontomys! colimæ Merr., Proc. Wash. Acad. Scien., III, 1901, p. 551.

Rhithrodontomys colimæ Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 266. Zoöl. Ser.

ALPINE HARVEST MOUSE.

Type locality. Sierra Nevada de Colima, State of Jalisco, Mexico. Altitude, 12,000 feet.

Geogr. Distr. Known only from the type locality.

a. *Rhithrodontomys colimæ nerterus* Merriam.

Reithrodontomys! colimæ nerterus Merr., Proc. Wash. Acad. Scien., III, 1901, p. 551.

Rhithrodontomys colimæ nerterus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 267. Zoöl. Ser.

COLIMA HARVEST MOUSE.

Type locality. Sierra Nevada de Colima, State of Jalisco, Mexico. Altitude, 6,500 feet.

Geogr. Distr. Known only from the type locality.

462. *Rhithrodontomys costaricensis* Allen.

Reithrodontomys! costaricensis Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1895, p. 139.

Rhithrodontomys costaricensis Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 267. Zoöl. Ser.

COSTA RICA HARVEST MOUSE.

Type locality. La Carpintera, Costa Rica. Altitude, 6,000 feet.

Geogr. Distr. Costa Rica, Central America. Range unknown.

a. *Rhithrodontomys costaricensis jalapæ* Merriam.

Reithrodontomys! costaricensis jalapæ Merr., Proc. Wash. Acad. Scien., III, 1901, p. 552.

Rhithrodontomys costaricensis jalapæ Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 267. Zoöl. Ser.

JALAPA HARVEST MOUSE.

Type locality. Jalapa, State of Vera Cruz, Mexico. Altitude, 4,000 feet.

Geogr. Distr. State of Vera Cruz. Range unknown.

463. *Rhithrodontomys goldmani* Merriam.

Reithrodontomys! goldmani Merr., Proc. Wash. Acad. Scien., III, 1901, p. 552.

Rhithrodontomys goldmani Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 268. Zool. Ser.

GOLDMAN'S HARVEST MOUSE.

Type locality. Metlatoyuca, State of Puebla, Mexico. Altitude, 800 feet.

Geogr. Distr. State of Puebla. Range unknown.

464. *Rhithrodontomys rufescens* Allen & Chapman.

Reithrodontomys! rufescens Allen & Chapman, Bull. Am. Mus. Nat. Hist. N. Y., 1897, p. 199.

Rhithrodontomys rufescens Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 268. Zool. Ser.

RUFIOUS HARVEST MOUSE.

Type locality. Jalapa, State of Vera Cruz, Mexico.

Geogr. Distr. State of Vera Cruz, Mexico. Range unknown.

465. *Rhithrodontomys tenuirostris* Merriam.

Reithrodontomys! tenuirostris Merr., Proc. Wash. Acad. Scien., III, 1901, p. 547.

Rhithrodontomys tenuirostris Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 268. Zool. Ser.

SLENDER-NOSED HARVEST MOUSE.

Type locality. Todos Santos, Guatemala. Altitude, 10,000 feet.

Geogr. Distr. Guatemala, Central America. Range unknown.

a. *Rhithrodontomys tenuirostris aureus* Merriam.

Reithrodontomys! tenuirostris aureus Merr., Proc. Wash. Acad. Scien., III, 1901, p. 548.

Rhithrodontomys tenuirostris aureus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 269. Zool. Ser.

CALEL HARVEST MOUSE.

Type locality. Calal, Guatemala. Altitude, 10,200 feet.

Geogr. Distr. Known only from the type locality.

466. *Rhithrodontomys microdon* Merriam.

Reithrodontomys! microdon Merr., Proc. Wash. Acad. Scien., III, 1901, p. 548.

Rhithrodontomys microdon Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 269. Zoöl. Ser.

TODOS SANTOS HARVEST MOUSE.

Type locality. Todos Santos, Guatemala. Altitude, 10,000 feet.
Geogr. Distr. Guatemala. Range unknown.

a. *Rhithrodontomys microdon albilabris* Merriam.

Reithrodontomys! microdon albilabris Merr., Proc. Wash. Acad. Scien., III, 1901, p. 549.

Rhithrodontomys microdon albilabris Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 269. Zoöl. Ser.

WHITE-LIPPED HARVEST MOUSE.

Type locality. Cerro San Felipe, State of Oaxaca, Mexico. Altitude, 10,000 feet.

Geogr. Distr. State of Oaxaca. Range unknown.

467. *Rhithrodontomys hirsutus* Merriam.

Reithrodontomys! hirsutus Merr., Proc. Wash. Acad. Scien., III, 1901, p. 553.

Rhithrodontomys hirsutus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 269. Zoöl. Ser.

HISPID HARVEST MOUSE.

Type locality. Ameca, State of Jalisco, Mexico. Altitude, 5,500 feet.

Geogr. Distr. State of Jalisco, Mexico. Range unknown.

468. *Rhithrodontomys griseiflavus* Merriam.

Reithrodontomys! griseoflavus Merr., Proc. Wash. Acad. Scien., III, 1901, p. 553.

Rhithrodontomys griseiflavus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 270. Zoöl. Ser.

AMECA HARVEST MOUSE.

Type locality. Ameca, State of Jalisco, Mexico. Altitude, 4,000 feet.

Geogr. Distr. State of Jalisco. Range unknown.

a. *Rhithrodontomys griseiflavus helvolus* Merriam.

Reithrodontomys! griseoflavus helvolus Merr., Proc. Wash. Acad. Scien., III, 1901, p. 554.

Rhithrodontomys griseiflavus helvolus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 270. Zoöl. Ser.

OAXACA HARVEST MOUSE.

Type locality. Near Oaxaca City, State of Oaxaca, Mexico.

Geogr. Distr. State of Oaxaca. Range unknown.

469. *Rhithrodontomys levipes* Merriam.

Reithrodontomys! levipes Merr., Proc. Wash. Acad. Scien., III, 1901, p. 554.

Rhithrodontomys levipes Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 270. Zoöl. Ser.

SAN SEBASTIAN HARVEST MOUSE.

Type locality. San Sebastian, State of Jalisco, Mexico. Altitude, 3,000 feet.

Geogr. Distr. State of Jalisco, Mexico. Range unknown.

a. *Rhithrodontomys levipes otus* Merriam.

Reithrodontomys! levipes otus Merr., Proc. Wash. Acad. Scien., III, 1901, p. 555.

Rhithrodontomys levipes otus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 271. Zoöl. Ser.

LARGE-EARED HARVEST MOUSE.

Type locality. Sierra Nevada de Colima, State of Jalisco, Mexico. Altitude, 6,500 feet.

Geogr. Distr. State of Jalisco, Mexico. Range unknown.

b. *Rhithrodontomys levipes toltecus* Merriam.

Reithrodontomys! levipes toltecus Merr., Proc. Wash. Acad. Scien., III, 1901, p. 555.

Rhithrodontomys levipes toltecus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 271. Zoöl. Ser.

TLALPAM HARVEST MOUSE.

Type locality. Tlalpam, Federal District, Mexico.

Geogr. Distr. Known only from the type locality.

470. *Rhithrodontomys inexpectatus* Elliot.

Rhithrodontomys inexpectatus Elliot, Pub. Field Columb. Mus., III, 1903, p. 145. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 271. Zoöl. Ser.

PATZCUARO HARVEST MOUSE.

Type locality. Patzcuaro, State of Michoacan, Mexico.

Geogr. Distr. Known only from the type locality.

471. *Rhithrodontomys dorsalis* Merriam.

Reithrodontomys! dorsalis Merr., Proc. Wash. Acad. Scien., III, 1901, p. 557.

Rhithrodontomys dorsalis Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 272. Zoöl. Ser.

GRAY-BACKED HARVEST MOUSE.

Type locality. Caelé, Guatemala.

Geogr. Distr. Guatemala. Range unknown.

472. Rhithrodontomys creper Bangs.

Reithrodontomys! creper Bangs, Bull. Mus. Comp. Zool., xxxix, 1902, p. 39.

Rhithrodontomys creper Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 272. Zool. Ser.

DUSKY HARVEST MOUSE.

Type locality. Volcan de Chiriqui, Chiriqui, Panama. Altitude, 11,000 feet.

Geogr. Distr. Panama, Central America. Range unknown.

Genus 71. Acodon.

Akodon! Meyen, Nova Acta Phys. Med. Acad. Cæs. Leop.-Carol., xvi, 1833, p. 599, tab. XLIII, fig. 1. Type *Akodon boliviense* Meyen.

Abrothrix (sic) Waterh., Proc. Zool. Soc., 1837, p. 21.

473. Acodon teguina (Alston).

Hesperomys teguina Alston, Proc. Zool. Soc., 1876, p. 755.

Acodon teguina Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 273, figs. 44, XLII. Zool. Ser.

ALSTON'S MOUSE.

Type locality. Coban, Guatemala.

Geogr. Distr. Guatemala. Range unknown.

Acodon teguina apricus Bangs.

Akodon! teguina apricus Bangs, Bull. Mus. Comp. Zool., xxxix, 1902, p. 40.

Acodon teguina apricus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 274. Zool. Ser.

BOQUETE MOUSE.

Type locality. Boquete, Chiriqui, Panama. Altitude, 4,000-5,000 feet.

Geogr. Distr. Panama, Central America. Range unknown.

474. Acodon irazu Allen.

Akodon! irazu Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1904, p. 46.

Acodon irazu Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 274. Zool. Ser.

VOLCANO OF IRAZU MOUSE.

Type locality. Volcan de Irazu, Costa Rica.

Geogr. Distr. Costa Rica, Central America. Range unknown.

475. *Acodon xerampelinus* Bangs.

Akodon! xerampelinus Bangs, Bull. Mus. Comp. Zool., xxxix, 1902, p. 41.

Acodon xerampelinus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 275. Zool. Ser.

BANGS' RED MOUSE.

Type locality. Volcan de Chiriqui, Chiriqui, Panama. Altitude, 10,300 feet.

Geogr. Distr. Panama, Central America. Range unknown.

Subfamily III. Neotominæ. Wood Rats.**Genus 72. *Neotoma*.**

Neotoma Say and Ord, Journ. Acad. Nat. Scien. Phil., iv, 1825, p. 345, pls. xxi, xxii. *Type Mus floridana* Ord.

476. *Neotoma cumulator* Mearns.

Neotoma cumulator Mearns, Proc. U. S. Nat. Mus., 1898, p. 503.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., ii, 1901, p. 154.

Zool. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M.

Pub., iv, Pt. 1, 1904, p. 280. Zool. Ser.

COLLECTOR WOOD RAT.

Type locality. Old Fort Yuma, San Diego County, California.

Geogr. Distr. Lower California and State of Sonora, Mexico; southern California.

477. *Neotoma palatina* Goldman.

Neotoma palatina Goldm., Proc. Biol. Soc. Wash., xviii, 1905, p. 27.

BOLAÑOS WOOD RAT.

Type locality. Bolaños, State of Jalisco, Mexico. Altitude, 2,800 feet.

Genl. Char. Size large; ears small; speno-palatine vacuities closed; vomer prolonged as a thin plate along median line of presphenoid, partly dividing posterior nares, and ending in a point at suture between presphenoid and basisphenoid; bullæ small and heavy.

Color. Above pale cinnamon, darker on dorsal region; cheeks and sides suffused with buffy; under parts dull white; hairs on the throat, breast, and inguinal region white to the roots; ears brownish; feet white; tail scantily haired, above blackish, beneath soiled whitish.

Measurements. Length, head and body, 187; tail vertebræ, incomplete, 60; hind foot, 37.

478. *Neotoma fuscipes* Baird.

Neotoma fuscipes Baird, N. Am. Mamm., 1857, p. 495. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 158. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 278. Zoöl. Ser.

Neotoma monochloura! Rhoads, Amer. Nat., XXVIII, 1894, p. 67.

Neotoma splendens True, Proc. U. S. Nat. Mus., 1894, p. 353.

DUSKY-FOOTED WOOD RAT.

Type locality. Petaluma, Sonoma County, California.

Geogr. Distr. Coast region of California and Oregon from Monterey Bay to the Columbia River.

a. *Neotoma fuscipes macrotis* (Thomas).

Neotoma macrotis Thomas, Ann. Mag. Nat. Hist., 6th Ser., XII, 1893, p. 234.

Neotoma fuscipes macrotis Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 159. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 279. Zoöl. Ser.

Neotoma macrotis simplex True, Proc. U. S. Nat. Mus., 1894, p. 354.

LARGE-EARED WOOD RAT.

Type locality. San Diego, San Diego County, California.

Geogr. Distr. Lower California, Mexico, and southern California.

b. *Neotoma fuscipes streatori* Merriam.

Neotoma fuscipes streatori Merr., Proc. Biol. Soc. Wash., IX, 1894, p. 124. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 159. Zoöl. Ser.

STREATOR'S WOOD RAT.

Type locality. Carbondale, Amador County, California.

Geogr. Distr. Coast region of California, south of Monterey Bay.

c. *Neotoma fuscipes dispar* Merriam.

Neotoma fuscipes dispar Merr., Proc. Biol. Soc. Wash., IX, 1894, p. 124. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 160. Zoöl. Ser.

DISSIMILAR WOOD RAT.

Type locality. Lone Pine, Owens Valley, Inyo County, California.

Geogr. Distr. East base of Sierra Nevada in Owens Valley, and western edge of Mohave Desert, California.

d. *Neotoma fuscipes annectens* Elliot.

Neotoma fuscipes annectens Elliot, Pub. Field. Columb. Mus., I, 1898, p. 201. Zoöl. Ser. *Id.* Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 160. Zoöl. Ser.

ALLIED WOOD RAT.

Type locality. Portola, San Mateo County, California.

Geogr. Distr. Coast line of California south of San Francisco Bay to Monterey County.

e. Neotoma fuscipes affinis Elliot.

Neotoma fuscipes affinis Elliot; Pub. Field Columb. Mus., I, 1898, p. 202. Zoöl. Ser. *Id.* Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 160. Zoöl. Ser.

RELATED WOOD RAT.

Type locality. Alum Rock Park, Santa Clara County, California.

Geogr. Distr. Known only from the type locality.

479. *Neotoma distincta* Bangs.

Neotoma distincta Bangs, Proc. Biol. Soc. Wash., xvi, 1903, p. 89. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 278. Zoöl. Ser.

TEXOLO WOOD RAT.

Type locality. Texolo, State of Vera Cruz, Mexico.

Geogr. Distr. State of Vera Cruz, Mexico. Range unknown.

480. *Neotoma mohavensis* (Elliot).

Neotoma fuscipes mohavensis Elliot, Pub. Field Columb. Mus., III, 1903, p. 246. Zoöl. Ser.

MOHAVE DESERT WOOD RAT.

Type locality. Oro Grande, Mohave Desert, San Bernardino County, California.

Geogr. Distr. Mohave Desert, southern California.

Genl. Char. Smaller than *N. f. macrotis*, more grayish in color, and with smaller feet.

Color. Upper parts dark drab gray, darkest on top of head and on the dorsal line; sides paler, inclined to buffy; arms buffy gray; thighs light gray or dark plumbeous; chin, throat, inner sides of fore legs, thighs, and ventral region, with the lower parts of hands and feet white; hairs on sides plumbeous at base; all the others on under parts white to the roots; tail above blackish brown, beneath whitish brown, line of demarcation very distinct; ears naked, dark brown.

Measurements. Total length, 384; tail vertebræ, 173; hind foot, 40; ear from notch, 31. Skull: occipito-nasal length, 47; Hensel, 39; zygomatic width, 24; interorbital constriction, 5; length of nasals, 16; palatal length, 21; length of upper tooth row, 9; length of mandible, 31; length of lower tooth row, 8.

481. *Neotoma cnemophila* (Elliot).

Neotoma fuscipes cnemophila Elliot, Pub. Field Columb. Mus., III, 1904, p. 267. Zoöl. Ser.

MOUNT PINOS WOOD RAT.

Type locality. Mt. Pinos, Lockwood Valley, Ventura County, California.

Geogr. Distr. Southern California. Range unknown.

Genl. Char. Size large, tail long, black above; general line of upper parts buff.

Color. Upper parts and sides buff, with base of hairs dark plumbeous; limbs mixed plumbeous and buff; chin, throat, and between thighs white; rest of under parts white to root of hairs, save a line along the sides; hands and feet white; wrists and ankles dusky; tail above and on sides dusky, beneath grayish white; ears dark brown.

Measurements. Total length, 380; tail vertebræ, 176; hind foot, 39.5; ear, 33. Skull: occipito-nasal length, 46; Hensel, 38; zygomatic width, 23; interorbital constriction, 5.6; width of braincase, 18; length of nasals, 18.5; palatal length, 19; length of incisive foramen, 9; length of upper tooth row, 9; length of mandible, angle to incisors, 23; length of lower tooth row, 8.

482. *Neotoma torquata* Ward.

Neotoma torquata Ward, Amer. Nat., xxv, 1891, p. 160. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 279. Zool. Ser.

COLLARED WOOD RAT.

Type locality. Between Tetela del Volcan and Zacualpan Amilpas, State of Morelos, Mexico.

Geogr. Distr. Known only from the type locality.

483. *Neotoma bryanti* Merriam.

Neotoma bryanti Merr., Amer. Nat., xxi, 1887, p. 191. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 280. Zool. Ser.

CERROS ISLAND WOOD RAT.

Type locality. Cerros or Cedros Island, Lower California, Mexico.

Geogr. Distr. Known only from the type locality.

484. *Neotoma micropus* Baird.

Neotoma micropus Baird, Proc. Acad. Nat. Scien. Phil., 1855, p. 333. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 155. Zool. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 281, figs. 45, XLIII. Zool. Ser. *Neotoma micropus canescens* Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1891, p. 285.

SMALL-FOOTED WOOD RAT.

Type locality. Charco Escondido, State of Tamaulipas, Mexico.

Geogr. Distr. States of Coahuila and Tamaulipas, Mexico, northward to Oklahoma and New Mexico.

a. *Neotoma micropus surberi* Elliot.

Neotoma micropus surberi Elliot, Pub. Field Columb. Mus., I, 1899, p. 279. Zoöl. Ser. *Id. Syn. N. Am. Mamm.*, F. C. M. Pub., II, 1901, p. 156. Zoöl. Ser.

SURBER'S WOOD RAT.

Type locality. Cañon three miles west of Alva, Oklahoma Territory.

Geogr. Distr. Northwestern Oklahoma.

b. *Neotoma micropus littoralis* Goldman.

Neotoma micropus littoralis Goldm., Proc. Biol. Soc. Wash., XVIII, 1905, p. 31.

COAST WOOD RAT.

Type locality. Alta Mira, State of Tamaulipas, Mexico. Altitude, 100 feet.

Genl. Char. Similar to *N. micropus*; but somewhat smaller.

Color. Above grayish brown, darkest on median line; under parts white; fur plumbeous at base except on throat, breast, and inguinal regions; foot white; tail blackish above, whitish below; ears brownish.

Measurements. Total length, 366; tail vertebræ, 166; hind foot, 39.

c. *Neotoma micropus planiceps* Goldman.

Neotoma micropus planiceps Goldm., Proc. Biol. Soc. Wash., XVIII, 1905, p. 32.

FLAT-HEADED WOOD RAT.

Type locality. Rio Verde, State of San Luis Potosi, Mexico. Altitude, 3,000 feet.

Genl. Char. Slightly smaller than *N. micropus*; color more buffy; skull flatter; "nasals pointed, deeply emarginate posteriorly."

Color. Above pale buffy gray, darkest on median line; under parts white; base of hairs plumbeous along sides of belly and inner surface of hind legs; nose, eyelids, and ankles dusky; feet white; tail brownish black above, grayish below.

Measurements. Total length, 351; tail vertebræ, 167; hind foot, 38.

485. *Neotoma floridana* (Ord).

Mus floridana Ord, Bull. Soc. Philom. Paris, 1818, p. 181.

Arvicola floridana Harl., Faun. Amer., 1825, p. 141.

Neotoma floridana Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 157, fig. 34. Zoöl. Ser.

FLORIDA WOOD RAT.

Type locality. St. John's River, near Jacksonville, Duval County, Florida.

Geogr. Distr. South Atlantic and Gulf Coasts, and lower Mississippi Valley.

a. *Neotoma floridana attwateri* (Mearns).

Neotoma attwateri Mearns, Proc. U. S. Nat. Mus., 1897, p. 721.

Neotoma floridana attwateri Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 157. Zoöl. Ser.

ATTWATER'S WOOD RAT.

Type locality. Lacey's Ranch, Turtle Creek, Kerr County, Texas.

Geogr. Distr. Southern Texas. Range unknown.

b. *Neotoma floridana rubida* Bangs.

Neotoma floridana rubida Bangs, Proc. Bost. Soc. Nat. Hist., 1898, p. 185.

Neotoma campestris rubida Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 156. Zoöl. Ser.

REDDISH WOOD RAT.

Type locality. Gibson, Terrebonne Parish, Louisiana.

Geogr. Distr. Lower Mississippi Valley; east to western Florida.

486. *Neotoma baileyi* Merriam.

Neotoma baileyi Merr., Proc. Biol. Soc. Wash., 19, 1894, p. 123.

Neotoma campestris baileyi Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 156. Zoöl. Ser.

BAILEY'S WOOD RAT.

Type locality. Valentine, Cherry County, Nebraska.

Geogr. Distr. Nebraska, Kansas, and Oklahoma Territory.

487. *Neotoma pennsylvanica* Stone.

Neotoma pennsylvanica Stone, Proc. Acad. Nat. Scien. Phil., 1893, p. 16. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 157. Zoöl. Ser.

PENNSYLVANIA WOOD RAT.

Type locality. Lewis Cave, South Mountain, Cumberland County, Pennsylvania.

Geogr. Distr. New York to Virginia in the Alleghany Mountains.

488. *Neotoma ferruginea* Tömes.

Neotoma ferruginea Tömes, Proc. Zoöl. Soc., 1861, p. 282. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., 14, Pt. 1, 1904, p. 280. Zoöl. Ser.

RUSTY WOOD RAT.

Type locality. Dueñas, Guatemala.

Geogr. Distr. Tehuantepec, State of Oaxaca, Mexico, south into Guatemala.

a. *Neotoma ferruginea ochracea* Goldman.

Neotoma ferruginea ochracea Goldm., Proc. Biol. Soc. Wash.,
XVIII, 1905, p. 30.

ATEMAJAC WOOD RAT.

Type locality. Atemajac, near Guadalajara, State of Jalisco, Mexico. Altitude, 4,000 feet.

Genl. Char. Smaller than *N. ferruginea*, tail sharply bicolor; skull with comparative differences.

Color. Above light ochraceous buff, darkest on dorsal region; under parts dingy white washed with buff; pectoral region ochraceous buff; nose and upper lip grayish white; tail brownish above, whitish below.

Measurements. Total length, 348; tail vertebræ, 161; hind foot, 38.

b. *Neotoma ferruginea solitaria* Goldman.

Neotoma ferruginea solitaria Goldm., Proc. Biol. Soc. Wash.,
XVIII, 1905, p. 31.

NENTON WOOD RAT.

Type locality. Nenton, Guatemala.

Genl. Char. Similar to *N. ferruginea* but smaller; outer side of forearms and hind legs not dusky. Differences of skull comparative.

Color. Above tawny ochraceous; top of head ochraceous buff; outer sides of forearms and hind legs blackish; under parts dull white; chin and throat pure white; fore feet white; hind feet clouded with dusky to toes which are white; tail brownish black above, dull gray beneath; tip brownish.

Measurements. Total length, 338; tail vertebræ, 156; hind foot, 35.

489. *Neotoma pinetorum* Merriam.

Neotoma pinetorum Merr., Proc. Biol. Soc. Wash., VIII, 1893,
p. 111. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901,
p. 158. Zoöl. Ser.

PINE WOOD RAT.

Type locality. San Francisco Mountain, Coconino County, Arizona.

Geogr. Distr. Pine covered plateau of Arizona.

490. *Neotoma nelsoni* Goldman.

Neotoma nelsoni Goldm., Proc. Biol. Soc. Wash., XVIII, 1905, p. 29.
NELSON'S WOOD RAT.

Type locality. Perote, State of Vera Cruz, Mexico.

Genl. Char. Size large; similar to *N. leucodon*, but darker; palate shorter than incisor foramina and with short median projection; premaxillæ reaching beyond nasals.

Color. Above pale cinnamon mixed with Prout's brown; cheeks and middle of face grayish brown; under parts dull white; base of fur plumbeous; pectoral spot white to roots of hairs; tail smoky brown above, paler beneath.

Measurements. Total length, 335; tail vertebræ, 145; hind foot, 37.

491. *Neotoma leucodon* Merriam.

Neotoma leucodon Merr., Proc. Biol. Soc. Wash., IX, 1894, p. 120.

Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 281. Zoöl. Ser.

WHITE-TOOTHED WOOD RAT.

Type locality. San Luis Potosi, State of San Luis Potosi, Mexico.

Geogr. Distr. States of Zacatecas, San Luis Potosi, and Vera Cruz, Mexico, from Berriozabal to Perote.

a. *Neotoma leucodon zacatecæ* Goldman.

Neotoma leucodon zacatecæ Goldm., Proc. Biol. Soc. Wash., XVIII, 1905, p. 30.

PLATEADO WOOD RAT.

Type locality. Plateado, State of Zacatecas, Mexico. Altitude, 7,600 feet.

Genl. Char. Similar to *N. leucodon* but darker; skull more arched; rostrum more decurved; frontals longer; upper incisors smaller.

Color. Above pale cinnamon, darkest on dorsal line, redder on rump and lightest on cheeks and sides; chest and inguinal region pure white; belly dull creamy white; feet white; upper lip dusky; tail brownish black above to white beneath.

Measurements. Total length, 358; tail vertebræ, 162; hind foot, 37.

492. *Neotoma latifrons* Merriam.

Neotoma latifrons Merr., Proc. Biol. Soc. Wash., IX, 1894, p. 121.

Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 282. Zoöl. Ser.

QUERENDARO WOOD RAT.

Type locality. Querendaro, State of Michoacan, Mexico.

Geogr. Distr. Valley of Querendaro, south side of Lake Cuitzeo, State of Michoacan, Mexico. Range unknown.

493. *Neotoma mexicana* Baird.

Neotoma mexicana Baird, Proc. Acad. Nat. Scien. Phil., 1855, p. 333. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 158. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 282. Zoöl. Ser.

MEXICAN WOOD RAT.

Type locality. Mountains near city of Chihuahua, State of Chihuahua, Mexico.

Geogr. Distr. State of Chihuahua, Mexico north into New Mexico and Texas, in latter state the southwestern part from Davis Mountains to Paisano.

a. *Neotoma mexicana bullata* Merriam.

Neotoma mexicana bullata Merr., Proc. Biol. Soc. Wash., IX, 1894, p. 122. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 158. Zoöl. Ser.

SANTA CATALINA WOOD RAT.

Type locality. Santa Catalina Mountains, southeastern Arizona.

Geogr. Distr. Santa Catalina Mountains, Arizona.

b. *Neotoma mexicana madrensis* Goldman.

Neotoma mexicana madrensis Goldm., Proc. Biol. Soc. Wash., XVIII, 1905, p. 31.

SIERRA MADRE WOOD RAT.

Type locality. Sierra Madre, near Guadalupe y Calvo, State of Chihuahua, Mexico. Altitude, 7,000 feet.

Genl. Char. Smaller than *N. mexicana*; color different.

Color. Above pale cinnamon, darkest on dorsal region, beneath dull white; fur plumbeous at base; axillæ ochraceous buff; fore and hind feet white; tail brownish above, grayish beneath.

Measurements. Total length, 320; tail vertebræ, 150; hind foot, 33.

494. *Neotoma navus* Merriam.

Neotoma navus Merr., Proc. Biol. Soc. Wash., XVI, 1903, p. 47.

Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 282. Zoöl. Ser.

ACTIVE WOOD RAT.

Type locality. Sierra Guadalupe, State of Coahuila, Mexico.

Geogr. Distr. State of Coahuila, Mexico. Range unknown.

495. *Neotoma sinaloæ* Allen.

Neotoma sinaloæ Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1898,

p. 149. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 283. Zoöl. Ser.

SINALOA WOOD RAT.

Type locality. Tatameles, State of Sinaloa, Mexico.

Geogr. Distr. State of Sinaloa. Range unknown.

496. *Neotoma arenacea* Allen.

Neotoma arenacea Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1898, p. 150. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 283. Zool. Ser.

LOWER CALIFORNIA WOOD RAT.

Type locality. San José del Cabo, Lower California, Mexico.

Geogr. Distr. Cape Region, Lower California, Mexico.

497. *Neotoma nudicauda* Goldman.

Neotoma nudicauda Goldm., Proc. Biol. Soc. Wash., XVIII, 1905, p. 28.

NAKED-TAILED WOOD RAT.

Type locality. Carmen Island, Lower California, Mexico.

Genl. Char. Similar to *N. arenacea* but paler, with comparative differences in cranial characters.

Color. Above pale grayish buff, slightly darker on back, and the sides tinged with brownish; under parts white, base of hairs plumbeous on belly; feet white, tail bicolor, brownish on upper third, whitish on lower two-thirds; ears grayish brown.

Measurements. Total length, 330; tail vertebræ, 91; hind foot, 39.

498. *Neotoma anthonyi* Allen.

Neotoma anthonyi Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1898, p. 151. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 283. Zool. Ser.

TODOS SANTOS WOOD RAT.

Type locality. Todos Santos Island, Lower California, Mexico.

Geogr. Distr. Known only from the type locality.

499. *Neotoma martinensis* Goldman.

Neotoma martinensis Goldm., Proc. Biol. Soc. Wash., XVIII, 1905, p. 28.

SAN MARTIN ISLAND WOOD RAT.

Type locality. San Martin Island, Lower California, Mexico.

Genl. Char. Size medium; color resembling that of *N. anthonyi*; Skull: temporal ridges faint, widely separated; nasal long, narrowing abruptly posteriorly, extending beyond lachrymals; frontal region constricted anteriorly, without well-developed ridges; bullæ small; meatus large, widely open.

Color. Above grayish brown, mixed with dusky on dorsal region; sides suffused with pale buff; under parts creamy white, base of fur

plumbeous; upper sides of forearms dusky brown; outer sides of hind legs and inner sides of ankles blackish; feet white; tail above brownish black, beneath grayish.

Measurements. Total length, 342; tail vertebræ, 167; hind foot, 35.7.

500. *Neotoma grandis* (Elliot).

Neotoma desertorum grandis Elliot, Pub. Field Columb. Mus., III, 1903, p. 247. Zoöl. Ser.

CAMERON LAKE WOOD RAT.

Type locality. Cameron Lake, Sierra Nevada, Kern County, California.

Geogr. Distr. Known only from the type locality.

Genl. Char. Similar to *N. desertorum*, but larger; tail more hairy and blacker above.

Color. Upper parts mixed buff and black; sides and thighs bright buff with less black; nose and sides of face buff; under parts and feet white; under fur along sides and thighs plumbeous; hairs on throat, chest, and middle of ventral surface white to the roots; tail very hairy, black above, white beneath; ears pale brown, the tuft of hair at base succeeded by a band of black hairs near middle of ear.

Measurements. Total length, 385; tail vertebræ, 185; hind foot, 38; ear, 30. Skull: total length, 47; Hensel, 40; zygomatic width, 23; interorbital constriction, 6; length of nasals, 17; palatal length, 21; length of upper molar series, 8; length of mandible, 30; length of lower molar series, 9.

501. *Neotoma desertorum* Merriam.

Neotoma desertorum Merr., Proc. Biol. Soc. Wash., IX, 1894, p. 125.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 161. Zoöl. Ser.

DEATH VALLEY WOOD RAT.

Type locality. Furnace Creek, Death Valley, Inyo County, California.

Geogr. Distr. Mohave, Colorado, and Sonoran Deserts of south-eastern California, western Nevada, and western Utah, north to east Humboldt Valley, and Kelton, Utah.

a. Neotoma desertorum sola Merriam.

Neotoma desertorum sola Merr., Proc. Biol. Soc. Wash., IX, 1894, p. 126. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 161. Zoöl. Ser.

SOLITARY WOOD RAT.

Type locality. San Emigdio, Kern County, California.

Geogr. Distr. Head of San Joaquin Valley, California.

502. *Neotoma lepida* Thomas.

Neotoma lepida Thomas, Ann. Mag. Nat. Hist., 6th Ser., xii, 1893, p. 235.

Neotoma cinerea lepida Elliot, Syn. N. Am. Mamm., F. C. M. Pub., ii, 1901, p. 164. Zoöl. Ser.

NEAT BUSHY-TAILED WOOD RAT.

Type locality. Williams Spring, about lat. 40°, long. 113°, just south of Champlin Mts., Juab County, Utah. Altitude, 4,558 feet.

Geogr. Distr. Utah. Range unknown.

503. *Neotoma stephensi* Goldman.

Neotoma stephensi Goldm., Proc. Biol. Soc. Wash., xviii, 1905, p. 32.

STEPHENS' WOOD RAT.

Type locality. Hualpai Mountains, Arizona. Altitude, 6,300 feet.

Genl. Char. Size small. Similar to *N. lepida*, hind foot larger; front upper molar with antero-internal sulcus obsolete. Compared with *N. lepida* the tooth row is longer; interparietal larger and bullæ smaller.

Color. Above grayish buff, palest on head; cheeks and sides pinkish buff; dorsal region mixed with brownish; under parts washed with pinkish buff; pectoral and inguinal regions, sometimes also the throat, pure white; feet white, ankles dusky; tail grayish brown above, paler beneath; ears grayish brown.

Measurements. Total length, 310; tail vertebræ, 139; hind foot, 31.

504. *Neotoma fallax* Merriam.

Neotoma fallax Merr., Proc. Biol. Soc. Wash., ix, 1894, p. 123.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., ii, 1901, p. 161. Zoöl. Ser.

DECEITFUL WOOD RAT.

Type locality. Gold Hill, Boulder County, Colorado.

Geogr. Distr. Eastern base of Rocky Mountains in Colorado, to an altitude of 7,000 feet.

505. *Neotoma intermedia* Rhoads.

Neotoma intermedia Rhoads, Amer. Nat., xxviii, 1894, p. 69.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., ii, 1901, p. 161.

Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 284. Zoöl. Ser.

Neotoma californica Price, Proc. Calif. Acad. Scien., 1894, p. 154, pl. xi.

RHOADS' WOOD RAT.

Type locality. Dulzura, San Diego County, California.

Geogr. Distr. Lower California and southern California south of the Bay of Monterey.

a. Neotoma intermedia melanura Merriam.

Neotoma intermedia melanura Merr., Proc. Biol. Soc. Wash., ix, 1894, p. 126. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 284. Zoöl. Ser.

BLACK-TAILED WOOD RAT.

Type locality. Ortiz, State of Sonora, Mexico.

Geogr. Distr. State of Sonora, Mexico, near west base of the Sierra Madre.

b. Neotoma intermedia angusticeps Merriam.

Neotoma intermedia angusticeps Merr., Proc. Biol. Soc. Wash., ix, 1894, p. 127. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., ii, 1901, p. 162. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 285. Zoöl. Ser.

NARROW-HEADED WOOD RAT.

Type locality. Southwest corner of Grant County, New Mexico.

Geogr. Distr. Southwestern New Mexico into the State of Chihuahua, Mexico, northwestern part.

c. Neotoma intermedia gilva Rhoads.

Neotoma intermedia gilva Rhoads, Amer. Nat., xxviii, 1894, p. 70. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., ii, 1901, p. 162. Zoöl. Ser.

Neotoma venusta True, Proc. U. S. Nat. Mus., 1894, p. 354.

YELLOW WOOD RAT.

Type locality. Banning, San Bernardino County, California.

Geogr. Distr. San Gorgonio Pass and Mohave Desert, San Bernardino County, California.

d. Neotoma intermedia albigula (Hartley).

Neotoma albigula Hartley, Proc. Calif. Acad. Scien., 2d Ser., iv, 1894, p. 157, pl. xii.

Neotoma intermedia albigula Elliot, Syn. N. Am. Mamm., F. C. M. Pub., ii, 1901, p. 162. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 285. Zoöl. Ser.

WHITE-THROATED WOOD RAT.

Type locality. Vicinity of Fort Lowell, near Tucson, Pima County, Arizona.

Geogr. Distr. Southern and western Arizona south into the States of Sonora and Coahuila, Mexico.

e. Neotoma intermedia durangæ Allen.

Neotoma intermedia durangæ Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1903, p. 602. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 285. Zoöl. Ser.

DURANGO WOOD RAT.

Type locality. San Gabriel, State of Durango, Mexico. Altitude, 7,000 feet.

Geogr. Distr. State of Durango, Mexico. Range unknown.

506. *Neotoma campestris* Allen.

Neotoma campestris Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1894, p. 322. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 150. Zoöl. Ser.

FIELD WOOD RAT.

Type locality. Pendennis, Lane County, Kansas.

Geogr. Distr. Kansas and Colorado.

507. *Neotoma bella* Bangs.

Neotoma bella Bangs, Proc. Biol. Soc. Wash., XII, 1899, p. 66. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, Append., p. 429. Zoöl. Ser.

BEAUTIFUL WOOD RAT.

Type locality. Palm Springs, Riverside County, California.

Geogr. Distr. Mohave and Colorado Deserts and San Bernardino Mountains, southeastern California.

a. Neotoma bella felipensis Elliot.

Neotoma bella felipensis Elliot, Pub. Field Columb. Mus., III, 1903, p. 217. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 286. Zoöl. Ser.

SAN FELIPE DESERT RAT.

Type locality. San Felipe, Gulf of California, Lower California, Mexico.

Geogr. Distr. Known only from the type locality, and Palomar in the Hanson Laguna Mountains.

508. *Neotoma montezumæ* Goldman.

Neotoma montezumæ Goldm., Proc. Biol. Soc. Wash., XVIII, 1905, p. 29.

MONTEZUMA WOOD RAT.

Type locality. Zimapan, State of Hidalgo, Mexico. Altitude, 7,500 feet.

Genl. Char. Resembling *N. i. melanura* but darker, with comparative differences in crania.

Color. Above dull ochraceous buff; palest on head and sides, mixed with blackish hairs; under parts white to roots on throat, chest, and inguinal region, rest plumbeous at base of fur; nose, upper side of forearms and outer side of legs, grayish brown; feet white, tail above blackish, beneath whitish; ears brownish.

Measurements. Length head and body, 181; tail vertebræ, 91, broken; hind foot, 39.

509. *Neotoma orizabæ* Merriam.

Neotoma orizabæ Merr., Proc. Biol. Soc. Wash., ix, 1894, p. 122.

Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 286. Zoöl. Ser.

ORIZABA WOOD RAT.

Type locality. Mt. Orizaba, State of Puebla, Mexico.

Geogr. Distr. States of Puebla, Tlaxcala, and Vera Cruz, Mexico.

510. *Neotoma tenuicauda* Merriam.

Neotoma tenuicauda Merr., Proc. Biol. Soc. Wash., vii, 1892,

p. 169. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 286. Zoöl. Ser.

SLENDER-TAILED WOOD RAT.

Type locality. North slope of the Sierra Nevada de Colima, State of Jalisco, Mexico. Altitude, 12,000 feet.

Geogr. Distr. Sierra Nevada de Colima, State of Jalisco, Mexico.

511. *Neotoma picta* Goldman.

Neotoma picta Goldm., Proc. Biol. Soc. Wash., xvii, 1904, p. 79.

Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 287. Zoöl. Ser.

PAINTED WOOD RAT.

Type locality. Chilpancingo, State of Guerrero, Mexico.

Geogr. Distr. State of Guerrero, Mexico. Range unknown.

512. *Neotoma isthmica* Goldman.

Neotoma isthmica Goldm., Proc. Biol. Soc. Wash., xvii, 1904,

p. 80. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 287. Zoöl. Ser.

ISTHMIAN WOOD RAT.

Type locality. Huilotepec, eight miles south of Tehuantepec, State of Oaxaca, Mexico.

Geogr. Distr. State of Oaxaca. Range unknown.

513. *Neotoma parvidens* Goldman.

Neotoma parvidens Goldm., Proc. Biol. Soc. Wash., xvii, 1904,

p. 81. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 288. Zoöl. Ser.

JUQUILA WOOD RAT.

Type locality. Juquila, State of Oaxaca, Mexico.

Geogr. Distr. State of Oaxaca, Mexico. Range unknown.

514. *Neotoma tropicalis* Goldman.

Neotoma tropicalis Goldm., Proc. Biol. Soc. Wash., xvii, 1904, p. 81. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 288. Zool. Ser.

TROPICAL WOOD RAT.

Type locality. Totontepec, State of Oaxaca, Mexico.

Geogr. Distr. State of Oaxaca, Mexico.

515. *Neotoma goldmani* Merriam.

Neotoma goldmani Merr., Proc. Biol. Soc. Wash., viii, 1903, p. 48. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 288. Zool. Ser.

GOLDMAN'S WOOD RAT.

Type locality. Saltillo, State of Coahuila, Mexico.

Geogr. Distr. Saltillo and Jaral, State of Coahuila, Mexico. Range unknown.

516. *Neotoma fulviventer* Merriam.

Neotoma fulviventer Merr., Proc. Biol. Soc. Wash., ix, 1904, p. 121. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 288. Zool. Ser.

FULVOUS-BELLIED WOOD RAT.

Type locality. Toluca Valley, State of Mexico, Mexico.

Geogr. Distr. State of Mexico, Mexico. Range unknown.

Genus 73. *Teonoma*. Bushy-tailed Wood Rats.

Teonoma Gray, List of Spec. Mamm. Brit. Mus., 1843, p. 117. Type *Myoxus drummondi* Richardson.

517. *Teonoma cinerea* (Ord).

Mus cinerea Ord, Guth. Geog., 2d Am. ed., ii, 1815, p. 292.

Neotoma cinerea Elliot, Syn. N. Am. Mamm., F. C. M. Pub., ii, 1901, p. 163, fig. 35. Zool. Ser.

GRAY BUSHY-TAILED WOOD RAT.

Type locality. Near Great Falls, Cascade County, Montana.

Geogr. Distr. Northern Rocky Mountain region from Utah to Wyoming; east to the Black Hills and plains of North Dakota, west of the Missouri River; west in southern British Columbia to Cascade Range, and south through the Sierra Nevada to Mt. Whitney in southern California.

a. *Teonoma cinerea drummondi* (Richardson).

Myoxus drummondi Rich., Zool. Journ., III, 1828, p. 517.

Neotoma cinerea drummondi Elliot, Syn. N. Am. Mamm., F. C. M.

Pub., II, 1901, p. 164. Zool. Ser.

DRUMMOND'S BUSHY-TAILED WOOD RAT.

Type locality. Rocky Mountains, British Columbia, Canada, N. Lat. 57°.

Geogr. Distr. Western Canada. Range unknown.

b. *Teonoma cinerea occidentalis* (Baird).

Neotoma occidentalis Baird, Proc. Acad. Nat. Scien. Phil., 1855, p. 335.

Neotoma cinerea occidentalis Elliot, Syn. N. Am. Mamm., F. C. M.

Pub., II, 1901, p. 164. Zool. Ser.

WESTERN BUSHY-TAILED WOOD RAT.

Type locality. Shoalwater Bay, Pacific County, Washington.

Geogr. Distr. Pacific coast region of Oregon and Washington and thence easterly to the Snake Plains of central Idaho.

c. *Teonoma cinerea columbiana* (Elliot).

Neotoma cinerea columbiana Elliot, Pub. Field Columb. Mus., I, 1899, p. 255. (Footnote.) Zool. Ser. *Id.* Syn. N. Am.

Mamm., F. C. M. Pub., II, 1901, p. 165. Zool. Ser.

BRITISH COLUMBIA BUSHY-TAILED WOOD RAT.

Type locality. Ducks, British Columbia, Canada.

Geogr. Distr. Western British Columbia, Canada. Range unknown.

518. *Teonoma saxamans* (Osgood).

Neotoma saxamans Osgood, N. Am. Faun., No. 19, 1900, p. 33.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, Append., p. 429. Zool. Ser.

ROCK-LOVING WOOD RAT.

Type locality. Bennett City, Lake Bennett, British Columbia, Canada.

Geogr. Distr. White Pass to the Semenow Hills, Alaska.

519. *Teonoma acraea* (Elliot).

Teonoma cinerea acraea Elliot, Pub. Field Columb. Mus., III, 1903, p. 247. Zool. Ser.

MOUNTAIN BUSHY-TAILED WOOD RAT.

Type locality. Hot Springs, Long Cañon, Mount Whitney, Inyo County, California. Altitude, 8,000 feet.

Geogr. Distr. Mount Whitney, Inyo County, California. Range unknown.

Genl. Char. Similar to *T. cinerea* but much paler; tail paler; feet smaller.

Color. Upper parts pinkish buff lined with black on top of the head and dorsal region, lightest on rump; sides of face and flanks pinkish buff, with very little black showing, this color extending over shoulders and thighs; orbital ring black; entire under parts, hands, and feet white; base of fur on sides only, plumbeous; tail above like back, slightly darker towards tip, beneath yellowish white; ears naked, dark brown; soles of feet naked.

Measurements. Total length, 360; tail vertebræ, 150; hind foot, 40; ear, 33.5. Skull: total length, 45.5; Hensel, 40; zygomatic width, 25; interorbital constriction, 6; braincase above roots of zygomata, 19; palatal length, 22; length of upper tooth row, 9; lower tooth row, alveolar border, 9.

520. *Teonoma fusca* (True).

Neotoma occidentalis fusca True, Proc. U. S. Nat. Mus., 1894, p. 354.

Neotoma cinerea fusca Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 165. Zoöl. Ser.

Teonoma fusca Elliot, Pub. Field Columb. Mus., III, 1903, p. 187. Zoöl. Ser.

FUSCOUS BUSHY-TAILED WOOD RAT.

Type locality. Fort Umpqua, Douglas County, Oregon.

Geogr. Distr. Coast region of Oregon. Range unknown.

a. *Teonoma fusca apicalis* (Elliot).

Neotoma fusca apicalis Elliot, Pub. Field Columb. Mus., III, 1903, p. 160. Zoöl. Ser.

WHITE-TIPPED BUSHY-TAILED WOOD RAT.

Type locality. Gardiner, Coos County, Oregon.

Geogr. Distr. Known only from the type locality.

Genl. Char. Size large, tail bushy, tip white. Skull long, narrow, orbital constriction considerable, rostrum long.

Color. Top of head and upper parts mixed black and yellowish; sides of head dark gray and yellowish; sides of body yellowish buff mixed with black; chin and upper part of the throat pure white, rest of throat plumbeous washed with white; inside of arms and a narrow line across breast deep ochraceous; white patch on breast and anal region; center of abdomen buff; rest of under parts plumbeous washed with white; outer side of hind legs black to ankles; hands and feet pure white; tail above blackish with white tip, beneath ochraceous at base and in a thin line in center of basal half, remainder white.

Measurements. Total length, 425; tail vertebrae, 202; hind foot, 47; ear, 32. Skull: occipito-nasal length, 51; Hensel, 43; zygomatic width, 27; interorbital constriction, 5; length of nasals, 20; palatal length, 24; length of upper tooth row, 9.5; length of mandible, 25; height at condyle, 10.5; at coronoid process, 15.

521. *Teonoma arizonæ* (Merriam).

Neotoma arizonæ Merr., Proc. Biol. Soc. Wash., VIII, 1903, p. 110.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 165.
Zoöl. Ser.

ARIZONA BUSHY-TAILED WOOD RAT.

Type locality. Keam's Cañon, Apache County, Arizona.

Geogr. Distr. Moki region, northeastern Arizona, northwestern New Mexico, southeastern Utah, and southwestern Colorado.

522. *Teonoma grangeri* (Allen).

Neotoma grangeri Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1894,
p. 324. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901,
p. 165. Zoöl. Ser.

GRANGER'S BUSHY-TAILED WOOD RAT.

Type locality. Custer, Black Hills, Custer County, South Dakota.

Geogr. Distr. South Dakota. Range unknown.

523. *Teonoma oreolestes* (Merriam).

Neotoma oreolestes (sic) Merr., Proc. Biol. Soc. Wash., IX, 1894,
p. 128. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901,
p. 166. Zoöl. Ser.

COLORADO BUSHY-TAILED WOOD RAT.

Type locality. Sagauche Valley, twenty miles west of Sagauché, Sagauche County, Colorado.

Geogr. Distr. Rocky Mountains, Colorado, Wyoming, and New Mexico.

524. *Teonoma rupicola* (Allen).

Neotoma rupicola Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1894,
p. 323. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901,
p. 166. Zoöl. Ser.

ROCK-DWELLING BUSHY-TAILED WOOD RAT.

Type locality. Corral Draw, Pine River Indian Reservation, southeastern base of the Black Hills, Custer County, South Dakota.

Geogr. Distr. South Dakota. Range unknown.

525. *Teonoma cinnamomea* (Allen).

Neotoma cinnamomea Allen, Bull. Am. Mus. Nat. Hist. N. Y.,
1895, p. 331. Elliot, Syn. N. Am. Mamm., F. C. M. Pub.,
II, 1901, p. 166. Zoöl. Ser.

CINNAMON BUSHY-TAILED WOOD RAT.

Type locality. Kinney Ranch, Bitter Creek, Sweetwater County, Wyoming.

Geogr. Distr. Wyoming, Uncompahgre Indian Reservation, Utah; New Mexico.

Genus 74. **Nelsonia.**

Nelsonia Merr., Proc. Biol. Soc. Wash., XI, 1897, p. 277, figs. 14-15.

Type *Nelsonia neotomodon* Merriam.

526. **Nelsonia neotomodon** Merriam.

Nelsonia neotomodon Merr., Proc. Biol. Soc. Wash., XI, 1897, p.

278. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M.

Pub., IV, Pt. I, 1904, p. 290, fig. 46. Zoöl. Ser.

ZACATECAS WOOD RAT.

Type locality. Mountains near Plateado, State of Zacatecas, Mexico. Altitude, 8,200 feet.

Geogr. Distr. State of Zacatecas, Mexico. Range unknown.

527. **Nelsonia goldmani** Merriam.

Nelsonia goldmani Merr., Proc. Biol. Soc. Wash., XVI, 1903, p.

80. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M.

Pub., IV, Pt. I, 1904, p. 290. Zoöl. Ser.

MT. TANCITARO WOOD RAT.

Type locality. Mt. Tancitaro, State of Michoacan, Mexico.

Geogr. Distr. State of Michoacan, Mexico. Range unknown.

Genus 75. **Xenomys.**

Xenomys Merr., Proc. Biol. Soc. Wash., VII, 1892, p. 160. Type

Xenomys nelsoni Merriam.

528. **Xenomys nelsoni** Merriam.

Xenomys nelsoni Merr., Proc. Biol. Soc. Wash., VII, 1892, p. 161.

Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV,

Pt. I, 1904, p. 292, fig. 47. Zoöl. Ser.

NELSON'S WOOD RAT.

Type locality. Hacienda Magdalena, between City of Colima and Manzanillo, State of Colima, Mexico.

Geogr. Distr. State of Colima, Mexico. Range unknown.

Genus 76. **Neotomodon.**

Neotomodon Merr., Proc. Biol. Soc. Wash., XII, 1898, p. 127. Type
Neotomodon alstoni Merriam.

529. Neotomodon perotensis Merriam.

Neotomodon perotensis Merr., Proc. Biol. Soc. Wash., XII, 1898,
p. 129. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M.
Pub., IV, Pt. 1, 1904, p. 293. Zoöl. Ser.

PEROTE WOOD RAT.

Type locality. Cofre de Perote, State of Vera Cruz, Mexico. Altitude, 9,500 feet.

Geogr. Distr. State of Vera Cruz, Mexico. Range unknown.

530. Neotomodon orizabæ Merriam.

Neotomodon orizabæ Merr., Proc. Biol. Soc. Wash., XII, 1898, p.
129. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M.
Pub., IV, Pt. 1, 1904, p. 294. Zoöl. Ser.

MOUNTAIN WOOD RAT.

Type locality. Mt. Orizaba, State of Puebla, Mexico. Altitude, 9,500 feet.

Geogr. Distr. State of Puebla. Range unknown.

531. Neotomodon alstoni Merriam.

Neotomodon alstoni Merr., Proc. Biol. Soc. Wash., XII, 1898, p.
128. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M.
Pub., IV, Pt. 1, 1904, p. 294, fig. 48. Zoöl. Ser.

ALSTON'S WOOD RAT.

Type locality. Nahuatzin, State of Michoacan, Mexico. Altitude, 8,500 feet.

Geogr. Distr. States of Michoacan, Morelos, and Mexico, Mexico.

Genus 77. **Teanopus.**

Teanopus Merr., Proc. Biol. Soc. Wash., XVI, 1903, p. 81. Type
Teanopus phenax Merriam.

532. Teanopus phenax Merriam.

Teanopus phenax Merr., Proc. Biol. Soc. Wash., XVI, 1903, p. 81.
Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV,
Pt. 1, 1904, p. 295, fig. 49. Zoöl. Ser.

CHEATING WOOD RAT.

Type locality. Camoa, Rio Mayo, State of Sonora, Mexico.

Geogr. Distr. State of Sonora, Mexico. Range unknown.

Genus 78. **Hodomys.**

Hodomys Merr., Proc. Acad. Nat. Scien. Phil., 1894, p. 232. Type
Neotoma alleni Merriam.

533. Hodomys alleni (Merriam).

Neotoma alleni Merr., Proc. Biol. Soc. Wash., VII, 1892, p. 168.

Hodomys alleni Elliot, Mamm. Middle Amer. & W. Indies, F. C. M.

Pub., IV, Pt. 1, 1904, p. 297, figs. 50, XLIV. Zoöl. Ser.

ALLEN'S WOOD RAT.

Type locality. Manzanillo, State of Colima, Mexico.

Geogr. Distr. State of Colima, Mexico.

534. Hodomys vetulus Merriam.

Hodomys vetulus Merr., Proc. Acad. Nat. Scien. Phil., 1894, p.

236. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M.

Pub., IV, Pt. 1, 1904, p. 297. Zoöl. Ser.

GRAY-FACED WOOD RAT.

Type locality. Tehuacan, State of Puebla, Mexico.

Geogr. Distr. State of Puebla. Range unknown.

Subfamily IV. **Microtinæ.**Genus 79. **Phenacomys. Voles.**

Phenacomys Merr., N. Am. Faun., No. 2, 1889, p. 28. Type *Phenacomys intermedius* Merriam.

535. Phenacomys intermedius Merriam.

Phenacomys intermedius Merr., N. Am. Faun., No. 2, 1889, p. 32.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 167.

Zoöl. Ser.

INTERMEDIATE VOLE.

Type locality. Twenty miles N. N. W. of Kamloops, British Columbia, Canada. Altitude, 5,500 feet.

Geogr. Distr. Known only from the type locality.

536. Phenacomys oreophilus Merriam.

Phenacomys orophilus (*sic*) Merr., N. Am. Faun., No. 5, 1891,

p. 65. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901,

p. 168, fig. 36. Zoöl. Ser.

Phenacomys truei Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1894,

p. 331.

Phenacomys oramontis Rhoads, Amer. Nat., XXIX, 1895, p. 941.

MOUNTAIN VOLE.

Type locality. Salmon River Mountains, near head of Timber Creek, Idaho. Altitude, 10,500 feet.

Geogr. Distr. Alberta, British Columbia, Canada, and western United States to southern central Oregon, central Idaho, and southern Wyoming.

537. *Phenacomys olympicus* Elliot.

Phenacomys olympicus Elliot, Pub. Field Columb. Mus., 1, 1899, p. 225. Zoöl. Ser. *Id. Syn. N. Am. Mamm., F. C. M. Pub.*, 11, 1901, p. 168. Zoöl. Ser.

Microtus pumilus Elliot, Pub. Field Columb. Mus., 1899, p. 226, juv. Zoöl. Ser.

OLYMPIC MOUNTAIN VOLE.

Type locality. Happy Lake, Olympic Mountains, Clallam County, Washington. Altitude, 6,500 feet.

Geogr. Distr. Known only from the type locality and Boulder Lake, Olympic Mountains, Washington.

538. *Phenacomys prebliei* Merriam.

Phenacomys preblei Merr., Proc. Biol. Soc. Wash., XI, 1897, p. 45. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 168. Zoöl. Ser.

PREBLE'S VOLE.

Type locality. Twin Peak near Long's Peak, Boulder County, Colorado. Altitude, 9,000 feet.

Geogr. Distr. Known only from the type locality.

539. *Phenacomys celatus* Merriam.

Phenacomys celatus Merr., N. Am. Faun., No. 2, 1889, p. 33. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 169. Zoöl. Ser.

Phenacomys latimanus Merr., N. Am. Faun., No. 2, 1889, p. 34.

Phenacomys ungava Merr., N. Am. Faun., No. 2, 1889, p. 35.

SECRETIVE VOLE.

Type locality. Fort Chimo, Ungava, Labrador.

Geogr. Distr. Northwestern Labrador to Peninsula Harbor, southeastern shore of Lake Superior.

a. Phenacomys celatus crassus Bangs.

Phenacomys celatus crassus Bangs, Proc. New Eng. Zoöl. Club, 1900, p. 39. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, Append., p. 430. Zoöl. Ser.

STUPID VOLE.

Type locality. Rigoulette, Hamilton Inlet, Labrador.

Geogr. Distr. Labrador. Range unknown.

540. *Phenacomys mackenzii* Preble.

Phenacomys mackenzii Preble, Proc. Biol. Soc. Wash., xv, 1902, p. 182.

MACKENZIE'S VOLE.

Type locality. Fort Smith, Slave River (near Athabasca, Mackenzie boundary line), Canada.

Genl. Char. Similar to *P. celatus* and *P. c. crassus*; cranial characters different; interorbital beads prominent, nearly meeting in old adults.

Color. Above mixed black and gray, tips of hairs yellowish brown, beneath grayish white; tail above brownish, beneath grayish white.

Measurements. Total length, 142; tail vertebræ, 34; hind foot, 17.

541. *Phenacomys constablii* Allen.

Phenacomys constablei Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1899, p. 4. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 169. Zool. Ser.

CONSTABLE'S VOLE.

Type locality. Telegraph Creek, Yukon, Canada.

Geogr. Distr. Known only from the type locality.

542. *Phenacomys longicaudus* True.

Phenacomys longicaudus True, Proc. U. S. Nat. Mus., 1890, p. 303. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 169. Zool. Ser.

LONG-TAILED VOLE.

Type locality. Marshfield, Coos County, Oregon.

Geogr. Distr. Coast district of Oregon. Range unknown.

543. *Phenacomys albipes* Merriam.

Phenacomys albipes Merr., Proc. Biol. Soc. Wash., xiv, 1901, p. 125.

WHITE-FOOTED VOLE.

Type locality. Redwoods near Arcata, Humboldt Bay, Humboldt County, California.

Genl. Char. Large; tail long, scantily haired; anterior root of zygoma sloping backward; jugal nearly parallel; nasals truncate posteriorly; bullæ large; interpterygoid fossa long; palate with median azygos process.

Color. Above grizzled bister; head, shoulders, and sides washed with brown, beneath grayish plumbeous washed with buffy; feet white; ankles dusky; tail above dusky, beneath whitish.

Measurements. Total length, 168; tail vertebræ, 62; hind foot, 19.

Genus 80. **Evotomys. Red-backed Voles.**

Evotomys Coues, Proc. Acad. Nat. Scien. Phil., 1874, p. 186. Type
Mus rutilus Pallas.

544. Evotomys rutilus (Pallas).

Mus rutilus Pall., Nov. Sp. Quad. Glir., 1778, p. 246, pl. xiv, B.

Evotomys rutilus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11,
1901, p. 170. Zoöl. Ser.

RUFIOUS RED-BACKED VOLE.

Type locality. Siberia, east of the Obi.

Geogr. Distr. Arctic regions of America, Europe, and Siberia.

545. Evotomys wrangeli Bailey.

Evotomys wrangeli Bailey, Proc. Biol. Soc. Wash., xi, 1897, p.

120. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901,
p. 171. Zoöl. Ser.

WRANGEL RED-BACKED VOLE.

Type locality. Wrangel, Wrangel Island, Alaska.

Geogr. Distr. Wrangel and Revillagigedo Islands, Alaska.

546. Evotomys caurinus Bailey.

Evotomys caurinus Bailey, Proc. Biol. Soc. Wash., xii, 1898, p.

21. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p.
171. Zoöl. Ser.

NORTHWESTERN RED-BACKED VOLE.

Type locality. Lund, east shore of Malaspina Inlet, British Columbia, Canada.

Geogr. Distr. Coast region of British Columbia east of the Strait of Georgia, and south of the Frazer River, Canada.

547. Evotomys dawsoni Merriam.

Evotomys dawsoni Merr., Amer. Nat., xxii, 1888, p. 650. Elliot,

Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 172. Zoöl.
Ser.

DAWSON'S RED-BACKED VOLE.

Type locality. Finlayson River, Northwest Territory, lat. 61° 30' N., long. 129° 30' West. Altitude, 3,000 feet.

Geogr. Distr. Finlayson River and Fort Liard, west to Yakutak and Juneau, and north along the coast to Prince William Sound.

a. Evotomys dawsoni alascensis (Miller).

Evotomys alascensis Miller, Proc. Acad. Nat. Scien. Phil., 1898,

p. 364, fig. 2. Elliot, Syn. N. Am. Mamm., F. C. M. Pub.,
11, 1901, p. 177. Zoöl. Ser.

Evotomys rutilus True, Proc. U. S. Nat. Mus., 1885, p. 596. (nec *Mus rutilus* Pallas.)

ALASKA RED-BACKED VOLE.

Type locality. St. Michaels, Norton Sound, Alaska.

Geogr. Distr. Alaska. Fort Yukon to St. Michaels. Limits of range unknown.

548. *Evotomys orca* Merriam.

Evotomys orca Merr., Proc. Wash. Acad. Scien., 11, 1900, p. 24.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 172.

Zoöl. Ser.

ORCA RED-BACKED VOLE.

Type locality. Orca, Prince William Sound, Alaska.

Geogr. Distr. Known only from the type locality.

549. *Evotomys gapperi* (Vigors).

Arvicola gapperi Vigors, Zoöl. Journ., v, 1830, p. 204, pl. ix.

Evotomys fuscodorsalis Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1894, p. 103.

Evotomys gapperi Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 172, fig. 37. Zoöl. Ser.

GAPPER'S RED-BACKED VOLE.

Type locality. Vicinity of Lake Simcoe, Ontario, Canada.

Geogr. Distr. Pennsylvania, New Jersey, and Massachusetts northward, and in Canada west to the Rocky Mountains.

a. Evotomys gapperi ochraceus Miller.

Evotomys gapperi ochraceus Miller, Proc. Bost. Soc. Nat. Hist., xxvi, 1894, p. 193. Elliot, Syn. N. Am. Mamm., F. C. M.

Pub., 11, 1901, p. 172. Zoöl. Ser.

OCHRACEOUS RED-BACKED VOLE.

Type locality. Mount Washington, Coos County, New Hampshire. Altitude, 5,500 feet.

Geogr. Distr. White Mountains of New Hampshire, and probably east to Nova Scotia.

b. Evotomys gapperi rhoadsi Stone.

Evotomys gapperi rhoadsi Stone, Amer. Nat., xxvii, 1893, p. 55.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 173. Zoöl. Ser.

RHOADS' RED-BACKED VOLE.

Type locality. May's Landing, Atlantic County, New Jersey.

Geogr. Distr. Known only from the type locality.

c. *Evotomys gapperi loringi* Bailey.

Evotomys gapperi loringi Bailey, Proc. Biol. Soc. Wash., XI, 1897, p. 125. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 173. Zoöl. Ser.

LORING'S RED-BACKED VOLE.

Type locality. Portland, Traill County, North Dakota.

Geogr. Distr. Timbered valleys along edge of plains in Minnesota, and eastern North and South Dakota.

d. *Evotomys gapperi galei* (Merriam).

Evotomys galei Merr., N. Am. Faun., No. 4, 1890, p. 23, pl. II, fig. 3.

Evotomys gapperi galei Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 173. Zoöl. Ser.

GALE'S RED-BACKED VOLE.

Type locality. Ward, Boulder County, Colorado. Altitude, 9,000 feet.

Geogr. Distr. Mountains of Colorado, north on eastern ranges of Rocky Mountains to northern Montana.

e. *Evotomys gapperi saturatus* Rhoads.

Evotomys gapperi saturatus Rhoads, Proc. Acad. Nat. Scien. Phil., 1894, p. 284. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 174. Zoöl. Ser.

RICH-COLOURED RED-BACKED VOLE.

Type locality. Nelson, British Columbia, Canada, on the Kootenay River, thirty miles north of the Washington boundary line.

Geogr. Distr. Blue Mountains of Oregon, mountains of northern Idaho, and northward into British Columbia to Cariboo Lake, Canada.

550. *Evotomys breviceaudus* (Merriam).

Evotomys gapperi breviceaudus Merr., N. Am. Faun., No. 5, 1891, p. 119, pl. III, figs. 7, 8.

Evotomys breviceaudus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 174. Zoöl. Ser.

SHORT-TAILED RED-BACKED VOLE.

Type locality. Near Custer, Black Hills, Custer County, South Dakota. Altitude, 6,000 feet.

Geogr. Distr. Black Hills, Custer County, South Dakota.

551. *Evotomys carolinensis* Merriam.

Evotomys carolinensis Merr., Am. Jour. Scien., 3d Ser., xxxvi, 1888, p. 460. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 174. Zoöl. Ser.

CAROLINA RED-BACKED VOLE.

Type locality. Roan Mountain, Mitchell County, North Carolina. Altitude, 6,000 feet.

Geogr. Distr. High portions of the Alleghany Mountains of North Carolina, Tennessee, and West Virginia.

552. *Evotomys ungava* Bailey.

Evotomys ungava Bailey, Proc. Biol. Soc. Wash., XI, 1897, p. 130.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 175.
Zool. Ser.

UNGAVA RED-BACKED VOLE.

Type locality. Fort Chimo, Ungava, Labrador.

Geogr. Distr. Ungava, Labrador. Range unknown.

553. *Evotomys idahoensis* Merriam.

Evotomys idahoensis Merr., N. Am. Faun., No. 5, 1891, p. 67.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 175.
Zool. Ser.

IDAHO RED-BACKED VOLE.

Type locality. Sawtooth or Alturas Lake, east base of the Sawtooth Mountains, Idaho. Altitude, 7,200 feet.

Geogr. Distr. Mountains of south central Idaho, between Snake and Salmon Rivers.

554. *Evotomys mazama* Merriam.

Evotomys mazama Merr., Proc. Biol. Soc. Wash., XI, 1897, p. 71.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 175.
Zool. Ser.

MT. MAZAMA RED-BACKED VOLE.

Type locality. Crater Lake, Mt. Mazama, Klamath County, Oregon. Altitude, 7,000 feet.

Geogr. Distr. Crest of the Cascade Mountains in Oregon.

555. *Evotomys obscurus* Merriam.

Evotomys obscurus Merr., Proc. Biol. Soc. Wash., XI, 1897, p. 72.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 176.
Zool. Ser.

DUSKY RED-BACKED VOLE.

Type locality. Prospect, Upper Rogue River Valley, Jackson County, Oregon.

Geogr. Distr. West slope of the southern Cascade Range and northern Sierra Nevada in southern Oregon and northern California.

556. *Evotomys californicus* Merriam.

Evotomys californicus Merr., N. Am. Faun., No. 4, 1890, p. 26,

pl. II, fig. 2. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 176. Zoöl. Ser.

CALIFORNIA RED-BACKED VOLE.

Type locality. Eureka, Humboldt County, California.

Geogr. Distr. Coast of northern California and Oregon.

557. *Evotomys occidentalis* Merriam.

Evotomys occidentalis Merr., N. Am. Faun., No. 4, 1890, p. 25,

pl. II, fig. 1. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 176. Zoöl. Ser.

WESTERN RED-BACKED VOLE.

Type locality. Aberdeen, Chehalis County, Washington.

Geogr. Distr. Coast and Puget Sound region of Washington and southern British Columbia.

558. *Evotomys nivarius* Bailey.

Evotomys nivarius Bailey, Proc. Biol. Soc. Wash., XI, 1897, p.

136. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 177. Zoöl. Ser.

SNOW-LOVING RED-BACKED VOLE.

Type locality. On the northwest slope of Mt. Ellinor, Olympic Mountains, Washington. Altitude, 4,000 feet.

Geogr. Distr. Mt. Ellinor and high altitudes in the Olympic Mountains, Washington.

559. *Evotomys proteus* Bailey.

Evotomys proteus Bailey, Proc. Biol. Soc. Wash., XI, 1897, p. 137.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 177. Zoöl. Ser.

CUNNING RED-BACKED VOLE.

Type locality. Hamilton Inlet, Labrador.

Geogr. Distr. Labrador. Range unknown.

Genus 81. *Microtus*. Meadow Voles.

Microtus Schrank, Fauna Boica, I, 1ste Abth., 1798, pp. 66, 72. Type

Mus arvalis Pallas.

Arvicola Lacép., Tab. Div. Sous-div. Ord. et Gen. Mamm., 1799, p. 10.

Mynomes Rafin., Am. Month. Mag., II, 1817, p. 45.

Psammomys Le Conte, Ann. N. Y. Lyc. Nat. Hist., 1830, p. 132. (nec Cretzschmar.)

Pitymys McMurt., Am. Ed. Cuv. Anim. King., I, App., 1831, p. 434. (Footnote.)

Ammomys Bon., Sagg. Dist. Met. degli Anim. Vert., 1831, p. 20. (Footnote.)

- Pinemys* Less., Hist. Nat. Mamm. et Ois. decouv. depuis 1788, in Ouvre de Buff., v, 1836, p. 436.
- Hemiotomys* Selys Longchamps, Essai Mon. Campagn. Envir. Liege, 1836, p. 7, pl. 1.
- Lagurus* Glog., Hand- u. Hilfsb. Naturgesch., I, 1841, p. 97.
- Neodon* Hodg., Ann. Mag. Nat. Hist., 2d Ser., III, 1849, p. 203.
- Agricola* Blas., Naturgesch. Säugeth. Deutsch., 1857, pp. 334-335, 368-374, figs. 202-206.
- Paludicola* Blas., Naturgesch. Säugeth. Deutsch., 1857, pp. 333-334, 343-368, figs. 183-201.
- Chilotus* Baird, N. Am. Mamm., 1857, p. 516.
- Pedomys* Baird, N. Am. Mamm., 1857, p. 517.
- Sylvicola* Fatio, Les Campagn. Bass. Léman, Ass. Zool. Léman, 1867, pp. 63-72, 75, pl. 1, figs. 18-25, pl. VI.
- Praticola* Fatio, Les Campagn. Bass. Léman, Ass. Zool. Léman, 1867, pp. 36-63, 75, pl. 1, figs. 5-17, pls. III-V. (Part.)
- Terricola* Fatio, Les Campagn. Bass. Léman, Ass. Zool. Léman, 1867, pp. 73, 75. (Part.)
- Ochetomys* Fitzin., Sitzungb., K. Akad. Wiss. Wien., LVI, 1867, pp. 103-105.
- Micrurus* Forsyth-Major, Att. dell. Soc. Tosc. Scien. Nat., Pisa, III, 1877, p. 126.
- Eremiomys* Polyakoff, Mém. Acad. Imp. Scien. St. Pétersb., xxxix, Suppl., 1881, pp. 35, 38, figs. 1-2.
- Neofiber* True, Science, IV, 1884, p. 34.
- Campicola* Schulze, Schrift. Natur. Ver. Harz. Wernig., v, 1890, pp. 24-25.
- Tetramerodon* Rhoads, Proc. Acad. Nat. Scien. Phil., 1894, pp. 282-283.
- Aulacomys* Rhoads, Amer. Nat., xxviii, 1894, pp. 182-185, figs. 1-5.
- Orthriomys* Merr., Proc. Biol. Soc. Wash., xii, 1898, pp. 106-107.
- Herpetomys* Merr., Proc. Biol. Soc. Wash., xii, 1898, pp. 107-108.

Subgenus A. *Microtus*.

560. *Microtus pennsylvanicus* (Ord).

- Mus pennsylvanica* Ord, Guth. Geog., 2d Am. ed., II, 1815, p. 292.
- Mynomes pratensis* Rafin., Am. Month. Mag., II, 1817, p. 45.
- Lemmus noveboracensis* Rafin., Ann. Nat., 1820, p. 3.
- Arvicola palustris* Harl., Faun. Amer., 1825, pp. 136-138.
- Arvicola riparius* Ord, Jour. Acad. Nat. Scien. Phil., IV, Pt. II, 1825, p. 305-306.

Arvicola hirsutus Emmons, Rept. Quad. Mass., 1840, p. 60.

Arvicola alborufescens! Emmons, Rept. Quad. Mass., 1840, pp. 60-61.

Arvicola fulva Aud. & Bachm., Proc. Acad. Nat. Scien. Phil., 1841, p. 96.

Arvicola nasuta Aud. & Bachm., Proc. Acad. Nat. Scien. Phil., 1841, pp. 96-97.

Arvicola rufescens DeKay, Zool. N. Y., Mamm., 1, 1842, p. 85.

Arvicola oneida DeKay, Zool. N. Y., Mamm., 1, 1842, pp. 88-89.

Arvicola dekayi Aud. & Bachm., Quad. N. Am., III, 1854, pp. 287-288.

Arvicola riparia var. *longipilis* Baird, N. Am. Mamm., 1857, p. 524.

Arvicola rufidorsum Baird, N. Am. Mamm., 1857, p. 526.

Microtus pennsylvanicus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 179, fig. 38. Zool. Ser.

PENNSYLVANIA MEADOW VOLE.

Type locality. Pennsylvania (meadows below Philadelphia).

Geogr. Distr. Eastern United States westward to the Dakotas and Nebraska.

a. *Microtus pennsylvanicus nigrans* Rhoads.

Microtus pennsylvanicus nigrans Rhoads, Proc. Acad. Nat. Scien. Phil., 1897, p. 307. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 180. Zool. Ser.

ALBEMARLE MEADOW VOLE.

Type locality. Currituck, Currituck County, North Carolina.

Geogr. Distr. Coast region of southern Virginia and northern North Carolina.

b. *Microtus pennsylvanicus acadicus* Bangs.

Microtus pennsylvanicus acadicus Bangs, Amer. Nat., xxxi, 1897, p. 239. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 180. Zool. Ser.

ACADIAN VOLE.

Type locality. Digby, Nova Scotia.

Geogr. Distr. Nova Scotia and Prince Edward Island.

c. *Microtus pennsylvanicus shattucki* Howe.

Microtus pennsylvanicus shattucki Howe, Proc. Portl. Soc. Nat. Hist., II, Pt. 6, 1901, p. 201, pl. v.

PENOBSCOT FIELD VOLE.

Type locality. Tumble Down Dick Island, near Long Island, Penobscot Bay, Maine.

Geogr. Distr. Known only from the type locality.

Genl. Char. "Size large; tail very long; ears prominent; bullæ very large and globular; palatine foramina markedly broad and bottle-shaped."

Color. Similar to that of *M. pennsylvanicus*, but slightly darker.

Measurements. "Total length, 176-198; tail vertebræ, 54-62; fore (hind?) foot, 12-13; ear, 12-18."

d. Microtus pennsylvanicus modestus (Baird).

Arvicola modesta Baird, N. Am. Mamm., 1857, p. 535.

Arvicola insperatus Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1894, p. 347.

Microtus pennsylvanicus modestus Elliot, Syn. N. Am. Mamm.,

F. C. M. Pub., 11, 1901, p. 180. Zoöl. Ser.

SAWATCH MEADOW VOLE.

Type locality. Cochetopa ("Sawatch") Pass, Saguache County, Colorado.

Geogr. Distr. Rocky Mountains and western plains from New Mexico to British Columbia, Canada, and from the Black Hills of South Dakota to central Idaho and the plains of the Columbia.

e. Microtus pennsylvanicus fontigenus (Bangs).

Microtus fontigenus Bangs, Proc. Biol. Soc. Wash., x, 1896, p. 48.

Microtus pennsylvanicus fontigenus Elliot, Syn. N. Am. Mamm.,

F. C. M. Pub., 11, 1901, p. 181. Zoöl. Ser.

FOREST VOLE.

Type locality. Lake Edward, Quebec, Canada.

Geogr. Distr. Quebec Province, Canada, west to Peninsula Harbor and Nepigon, north shore of Lake Superior.

f. Microtus pennsylvanicus labradorius Bailey.

Microtus pennsylvanicus labradorius Bailey, Proc. Biol. Soc. Wash.,

xii, 1898, p. 88. Elliot, Syn. N. Am. Mamm., F. C. M. Pub.,

11, 1901, p. 181. Zoöl. Ser.

LITTLE LABRADOR MEADOW VOLE.

Type locality. Fort Chimo, Ungava, Labrador.

Geogr. Distr. Known only from the type locality.

561. *Microtus drummondi* (Audubon & Bachman).

Arvicola drummondi Aud. & Bachm., Quad. N. Am., iii, 1854, p. 166.

Arvicola (Mynomes) microcephalus Rhoads, Proc. Acad. Nat. Scien. Phil., 1894, p. 286.

Microtus stonei Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1899, p. 5.

Microtus drummondi Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 181. Zoöl. Ser.

DRUMMOND'S MEADOW VOLE.

Type locality. Rocky Mountains, near Jasper House, Alberta, Canada.

Geogr. Distr. Hudson Bay to Alaska, and from the United States northern boundary to Fort Anderson, Northwest Territory.

562. *Microtus aztecus* (Allen).

Arvicola (Mynomes) aztecus Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1893, p. 73.

Microtus aztecus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 182. Zoöl. Ser.

AZTEC MEADOW VOLE.

Type locality. Aztec, San Juan County, New Mexico. Altitude, 5,900 feet.

Geogr. Distr. Estes Park (Allen), Colorado, to valley of the San Juan River, New Mexico.

563. *Microtus enixus* Bangs.

Microtus enixus Bangs, Amer. Nat., xxx, 1896, p. 1051. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 182. Zoöl. Ser.

LARGE LABRADOR MEADOW VOLE.

Type locality. Hamilton Inlet, Labrador.

Geogr. Distr. Eastern coast of Labrador from Hamilton Inlet to Ungava Bay, Labrador.

564. *Microtus aphorodemus* Preble.

Microtus aphorodemus Preble, N. Am. Faun., No. 22, 1901, p. 52.

BARREN-GROUND MEADOW VOLE.

Type locality. Barren Grounds near the mouth of the Thlewiaza River, Keewatin, Canada.

Geogr. Distr. Barren Grounds, Arctic America. Range unknown.

Genl. Char. Similar to *M. drummondi*, but larger and skull heavier.

Color. Above dark yellowish bister mixed with yellow tipped hairs; beneath white or light plumbeous, sometimes tinged with light brown.

Measurements. Total length, 190; tail vertebræ, 60; hind foot, 20. Skull: basal length, 28; length of nasals, 7.6; zygomatic breadth, 16.5; mastoid breadth, 12.6; alveolar length of upper molar series, 7.

565. *Microtus terrænovæ* (Bangs).

Arvicola terrænovæ Bangs, Proc. Biol. Soc. Wash., ix, 1894, p. 129.

Microtus terrænovæ Elliot, Syn. N. Am. Mamm., F. C. M. Pub.,
 II, 1901, p. 182. Zoöl. Ser.

NEWFOUNDLAND MEADOW VOLE.

Type locality. Codroy, Newfoundland.

Geogr. Distr. Newfoundland and Penguin Island.

566. *Microtus breweri* (Baird).

Arvicola breweri Baird, N. Am. Mamm., 1857, p. 525.

Microtus breweri Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II,
 1901, p. 183. Zoöl. Ser.

BEACH MEADOW VOLE.

Type locality. Muskeget Island, off Nantucket, Massachusetts.

Geogr. Distr. Known only from the type locality.

567. *Microtus nesophilus* Bailey.

Microtus insularis Bailey, Proc. Biol. Soc. Wash., XII, 1898, p. 86.
 (nec Nilson.)

Microtus nesophilus Bailey, Science, N. S., VIII, 1898, p. 782.
 Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 183
 Zoöl. Ser.

GULL ISLAND MEADOW VOLE.

Type locality. Great Gull Island, off eastern end of Long Island,
 Suffolk County, New York.

Geogr. Distr. Known only from the type locality.

568. *Microtus montanus* (Peale).

Arvicola montanus Peale, U. S. Expl. Exped., Mamm., VIII, 1848,
 p. 44.

Arvicola longirostris Baird, N. Am. Mamm., 1857, p. 530.

Microtus montanus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II,
 1901, p. 183. Zoöl. Ser.

PEALE'S MEADOW VOLE.

Type locality. Headwaters of the Sacramento River, near Mount
 Shasta, Siskiyou County, California.

Geogr. Distr. Northern Utah, Nevada, northeastern California,
 and eastern Oregon.

a. *Microtus montanus arizonensis* Bailey.

Microtus montanus arizonensis Bailey, Proc. Biol. Soc. Wash.,
 XII, 1898, p. 88. Elliot, Syn. N. Am. Mamm., F. C. M. Pub.,
 II, 1901, p. 184. Zoöl. Ser.

ARIZONA MEADOW VOLE.

Type locality. Springerville, Apache County, Arizona.

Geogr. Distr. Plateau County of eastern Arizona at head of the Little Colorado River.

b. Microtus montanus rivularis (Bailey).

Microtus nevadensis rivularis Bailey, Biol. Soc. Wash., xii, 1898, p. 87.

Microtus montanus rivularis Elliot, Syn. N. Am. Mamm., F. C. M. Pub., ii, 1901, p. 184. Zoöl. Ser.

UTAH MEADOW VOLE.

Type locality. St. George, Washington County, Utah.

Geogr. Distr. Known only from the type locality.

569. *Microtus nanus* (Merriam).

Arvicola nanus Merr., N. Am. Faun., No. 5, 1891, p. 62, pl. ii, figs. 5, 6.

Microtus nanus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., ii, 1901, p. 184. Zoöl. Ser.

DWARF MEADOW VOLE.

Type locality. Pahsimeroi Mountains, Idaho. Altitude, 9,350 feet.

Geogr. Distr. Rocky Mountains and other ranges from central Idaho, southward to Central Nevada and southern Colorado.

a. Microtus nanus canescens Bailey.

Microtus nanus canescens Bailey, Proc. Biol. Soc. Wash., xii, 1898, p. 87. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., ii, 1901, p. 185. Zoöl. Ser.

GRAY MEADOW VOLE.

Type locality. Conconully, Okanogan County, Washington.

Geogr. Distr. Northern Washington and southern British Columbia, east of the Cascades.

570. *Microtus canicaudus* Miller.

Microtus canicaudus Miller, Proc. Biol. Soc. Wash., xi, 1897, p. 67. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., ii, 1901, p. 185. Zoöl. Ser.

GRAY-TAILED MEADOW VOLE.

Type locality. McCoy, Willamette Valley, Polk County, Oregon.

Geogr. Distr. Willamette Valley, Oregon, and southern Washington, on eastern slopes of the Cascade Mountains.

571. *Microtus dutcheri* Bailey.

Microtus dutcheri Bailey, Proc. Biol. Soc. Wash., xii, 1898, p. 85. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., ii, 1901, p. 185. Zoöl. Ser.

DUTCHER'S MEADOW VOLE.

Type locality. Big Cottonwood Meadows, near Mount Whitney, Tulare County, California. Altitude, 10,000 feet.

Geogr. Distr. Inyo and Tulare Counties, southeastern California. "Hudsonian zone of the southern Sierra Nevada."

572. *Microtus nevadensis* Bailey.

Microtus nevadensis Bailey, Proc. Biol. Soc. Wash., XII, 1898, p. 86. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 185. Zool. Ser.

NEVADA MEADOW VOLE.

Type locality. Ash Meadows, Nye County, Nevada.

Geogr. Distr. Type locality, and Pahrangat Valley, southeastern Nevada.

573. *Microtus californicus* (Peale).

Arvicola californica Peale, U. S. Expl. Exped., VIII, Mamm. & Ornith., 1848, p. 46.

Arvicola trowbridgi Baird, N. Am. Mamm., 1857, p. 529.

Microtus californicus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 186. Zool. Ser.

CALIFORNIA MEADOW VOLE.

Type locality. Vicinity of San Francisco Bay, California.

Geogr. Distr. California west of the Colorado Desert and the Sierra Nevada, and from Santa Isabel, San Diego County, California, to the Rogue River and Umpqua Valleys, Oregon.

a. *Microtus californicus hyperythrus* Elliot.

Microtus californicus hyperythrus Elliot, Pub. Field Columb. Mus., III, 1903, p. 161. Zool. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 302, figs. 51, XLV. Zool. Ser.

REDDISH MEADOW VOLE.

Type locality. San Quentin, Lower California, Mexico.

Geogr. Distr. Sea coast in the vicinity of San Quentin, up to 8,000 feet elevation in the San Pedro Martir Mountains, Lower California, Mexico.

b. *Microtus californicus vallicola* Bailey.

Microtus californicus vallicola Bailey, Proc. Biol. Soc. Wash., XII, 1898, p. 89. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 186. Zool. Ser.

VALLEY MEADOW VOLE.

Type locality. Lone Pine, Inyo County, California.

Geogr. Distr. Valleys east of the Sierra Nevada, California.

c. Microtus californicus constrictus Bailey.

Microtus californicus constrictus Bailey, N. Am. Faun., No. 17, 1900, p. 36. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 186. Zool. Ser.

COAST MEADOW VOLE.

Type locality. Cape Mendocino, Humboldt County, California.

Geogr. Distr. Coast region near Cape Mendocino, California.

574. *Microtus edax* (Le Conte).

Arvicola edax Le Conte, Proc. Acad. Nat. Scien. Phil., 1853, p. 405.

Microtus edax Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 187. Zool. Ser.

TULE MEADOW VOLE.

Type locality. California south of San Francisco.

Geogr. Distr. San Joaquin and Sacramento Valleys, California.

575. *Microtus scirpensis* Bailey.

Microtus scirpensis Bailey, N. Am. Faun., No. 17, 1900, p. 38.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 187. Zool. Ser.

DESERT MEADOW VOLE.

Type locality. Amargosa River, near Nevada line, Inyo County, California.

Geogr. Distr. Known only from the type locality.

576. *Microtus operarius* (Nelson).

Arvicola operarius Nelson, Proc. Biol. Soc. Wash., VIII, 1893, p. 139.

Microtus operarius Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 187. Zool. Ser.

TUNDRA MEADOW VOLE.

Type locality. St. Michaels, Norton Sound, Alaska.

Geogr. Distr. Coast tundras of Bering Sea from Cape Vancouver to Bering Straits, including Nelson, St. Michaels, and Stewart Islands; eastward in Alaska to Anderson River.

577. *Microtus macfarlanii* Merriam.

Microtus macfarlani Merr., Proc. Wash. Acad. Scien., 11, 1900, p. 24. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 187. Zool. Ser.

MACFARLANE'S MEADOW VOLE.

Type locality. Fort Anderson, Anderson River, Mackenzie, Canada.

Geogr. Distr. Tundra region of Arctic America, east to the Mackenzie River.

578. *Microtus yakutatensis* Merriam.

Microtus yakutatensis Merr., Proc. Wash. Acad. Scien., 11, 1900,
p. 22. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901,
p. 188. Zoöl. Ser.

YAKUTAT MEADOW VOLE.

Type locality. Yakutat Bay (north shore), Alaska.

Geogr. Distr. Mainland of Alaska from Glacier Bay to Prince William Sound.

579. *Microtus kadiacensis* Merriam.

Microtus kadiacensis Merr., Proc. Biol. Soc. Wash., 11, 1897,
p. 222. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901,
p. 188. Zoöl. Ser.

KADIAK MEADOW VOLE.

Type locality. Kadiak Island, Alaska.

Geogr. Distr. Known only from the type locality.

580. *Microtus unalascensis* Merriam.

Microtus unalascensis Merr., Proc. Biol. Soc. Wash., 11, 1897,
p. 222. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901,
p. 188. Zoöl. Ser.

UNALASKA ISLAND MEADOW VOLE.

Type locality. Ilulink village, Unalaska Island, Alaska.

Geogr. Distr. Unalaska Island, Alaska.

a. *Microtus unalascensis popofensis* Merriam.

Microtus unalascensis popofensis Merr., Proc. Wash. Acad. Scien.,
11, 1900, p. 22. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11,
1901, p. 188. Zoöl. Ser.

POPOF ISLAND MEADOW VOLE.

Type locality. Popof Island, Shumagin Group, Alaska.

Geogr. Distr. Known only from Popof Island.

581. *Microtus sitkensis* Merriam.

Microtus sitkensis Merr., Proc. Biol. Soc. Wash., 11, 1897, p. 221.
Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 189.
Zoöl. Ser.

SITKA MEADOW VOLE.

Type locality. Sitka, Alaska.

Geogr. Distr. Baranof Island, Alaska.

582. *Microtus innuitus* Merriam.

Microtus innuitus Merr., Proc. Wash. Acad. Scien., 11, 1900, p. 21.
Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 189.
Zoöl. Ser.

INNUIT MEADOW VOLE.

Type locality. St. Lawrence Island, Bering Sea, Alaska.

Geogr. Distr. Known only from the type locality.

583. *Microtus abbreviatus* Miller.

Microtus abbreviatus Miller, Proc. Biol. Soc. Wash., XIII, 1899, p. 13. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 189. Zoöl. Ser.

HALL ISLAND MEADOW VOLE.

Type locality. Hall Island, Bering Sea, Alaska.

Geogr. Distr. Known only from Hall Island.

a. Microtus abbreviatus fisheri Merriam.

Microtus abbreviatus fisheri Merr., Proc. Wash. Acad. Scien., II, 1900, p. 23. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 190. Zoöl. Ser.

ST. MATTHEW ISLAND MEADOW VOLE.

Type locality. St. Matthew Island, Bering Sea, Alaska.

Geogr. Distr. Known only from St. Matthew Island.

584. *Microtus townsendi* (Bachman).

Arvicola townsendi Bachm., Journ. Acad. Nat. Scien. Phil., 1839, p. 60, pl. 1.

Arvicola occidentalis Peale, U. S. Expl. Exped., Mamm. & Ornith., VIII, 1848, p. 45.

Microtus townsendi Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 190. Zoöl. Ser.

TOWNSEND'S MEADOW VOLE.

Type locality. Lower Columbia River, near mouth of the Willamette, or near Wappatoo or Sauvie Island.

Geogr. Distr. Port Moody, British Columbia, Canada, south to the Willamette Valley and to Yaquina Bay, Oregon.

585. *Microtus tetramerus* (Rhoads).

Arvicola (Tetramerodon) tetramerus Rhoads, Proc. Acad. Nat. Scien. Phil., 1894, p. 283.

Microtus tetramerus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 190. Zoöl. Ser.

VANCOUVER MEADOW VOLE.

Type locality. Beacon Hill Park, Victoria, British Columbia.

Geogr. Distr. Southern end of Vancouver Island, British Columbia, Canada.

586. *Microtus longicaudus* (Merriam).

Arvicola (Mynomes) longicaudus Merriam, Amer. Nat., XXII, 1888, p. 934.

Microtus longicaudus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 191. Zoöl. Ser.

LONG-TAILED MEADOW VOLE.

Type locality. Custer, Black Hills, Custer County, South Dakota. Altitude, 5,500 feet.

Geogr. Distr. Western South Dakota and Wyoming.

587. *Microtus mordax* (Merriam).

Arvicola (Mynomes) mordax Merr., N. Am. Faun., No. 5, 1891, p. 61.

Microtus vellerosus Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1899 p. 7.

Microtus cantus Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1899, p. 7.

Microtus mordax Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 191. Zoöl. Ser.

CANTANKEROUS MEADOW VOLE.

Type locality. Sawtooth (or Alturas) Lake, east base of Sawtooth Mountains, Idaho. Altitude, 7,200 feet.

Geogr. Distr. Rocky Mountains and contiguous ranges from latitude 60° to northern New Mexico, and in the Cascades and Sierra Nevada to Kaweah and Kern rivers, California.

588. *Microtus macrurus* Merriam.

Microtus macrurus Merr., Proc. Acad. Nat. Scien. Phil., 1898, p. 353. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 191. Zoöl. Ser.

OLYMPIC MEADOW VOLE.

Type locality. Lake Cushman, Olympic Mountains, Washington.

Geogr. Distr. Olympic Mountains, Washington, and along the coast of British Columbia north to Yakutat in Alaska.

589. *Microtus angusticeps* Bailey.

Microtus angusticeps Bailey, Proc. Biol. Soc. Wash., XII, 1898, p. 86. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 192. Zoöl. Ser.

COAST MEADOW VOLE.

Type locality. Crescent City, Del Norte County, California.

Geogr. Distr. Coast region of northwestern California and southwestern Oregon.

590. *Microtus alticola* (Merriam).

Arvicola (Mynomes) alticolus! Merr., N. Am. Faun., No. 3, 1890, p. 67, pl. v, figs. 1, 2, and pl. vi, figs. 1, 2, 3, 4.

Microtus alticola Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 192. Zoöl. Ser.

MOUNTAIN MEADOW VOLE.

Type locality. Little Spring, San Francisco Mountain, Coconino County, Arizona. Altitude, 8,200 feet.

Geogr. Distr. San Francisco Mountain, at an altitude of 8,200 to 11,000 feet.

a. Microtus alticola leucophæus (Allen).

Arvicola leucophæus! Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1894, p. 320.

Microtus alticola leucophæus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 192. Zoöl. Ser.

GRAHAM MOUNTAIN MEADOW VOLE.

Type locality. Graham Mountains, Graham County, Arizona.

Geogr. Distr. Known only from the type locality.

591. *Microtus mexicanus* (Saussure).

Arvicola (Hemiotomys) mexicanus Sauss., Rev. Mag. Zool., 2me Sér., XIII, 1861, p. 3.

Microtus mexicanus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 301. Zoöl. Ser.

MEXICAN MEADOW VOLE.

Type locality. Mount Orizaba, State of Puebla, Mexico.

Geogr. Distr. Eastern part of the State of Puebla, Mexico. Range unknown.

a. Microtus mexicanus phæus (Merriam).

Arvicola phæus! Merr., Proc. Biol. Soc. Wash., VII, 1892, p. 171.

Microtus mexicanus phæus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 302. Zoöl. Ser.

COLIMA OR DARK MEADOW VOLE.

Type locality. North slope of the Sierra Nevada de Colima, State of Jalisco, Mexico. Altitude, 10,000 feet.

Geogr. Distr. Known only from type locality.

b. Microtus mexicanus guadalupensis Bailey.

Microtus mexicanus guadalupensis Bailey, Proc. Biol. Soc. Wash., XV, 1902, p. 118.

GUADALUPE MOUNTAINS MEADOW VOLE.

Type locality. Guadalupe Mountains, El Paso County, Texas. Altitude, 7,800 feet.

Genl. Char. Larger and lighter colored than *M. mexicanus*. Skull long; nasals short, wide, posterior ends rounded; incisors dark orange; upper and middle molars like that of *M. mogollonensis*, with projecting inner point at base of posterior triangle.

Color. Above dull umber brown; belly buffy gray; feet and tail brownish gray.

Measurements. Total length, 152; tail vertebræ, 34; hind foot, 20. Skull: basal length, 24.5; length of nasals, 7.5; zygomatic breadth, 16; mastoid breadth, 12.4; alveolar length of upper molar series, 7.2.

592. *Microtus fulviventer* Merriam.

Microtus fulviventer Merr., Proc. Biol. Soc. Wash., XII, 1898, p. 106. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 301. Zoöl. Ser.

FULVOUS-BELLIED OR OAXACA MEADOW VOLE.

Type locality. Cerro San Felipe, State of Oaxaca, Mexico. Altitude, 10,200 feet.

Geogr. Distr. State of Oaxaca, Mexico.

593. *Microtus mogollonensis* (Mearns).

Arvicola mogollonensis Mearns, Bull. Am. Mus. Nat. Hist. N. Y. 1890, p. 283.

Microtus mogollonensis Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 192. Zoöl. Ser.

MOGOLLON MOUNTAINS MEADOW VOLE.

Type locality. Baker's Butte, Mogollon Mountains, Yavapai County, Arizona.

Geogr. Distr. Plateau of central Arizona.

594. *Microtus xanthognathus* (Leach).

Arvicola xanthognatha Leach, Zoöl. Misc., I, 1814, p. 60.

Microtus xanthognathus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 193. Zoöl. Ser.

YELLOW-CHEEKED MEADOW VOLE.

Type locality. Hudson Bay.

Geogr. Distr. Northwestern Canada and Alaska, from central Alberta north to the Arctic Coast, and west to Central Alaska.

595. *Microtus chrotorrhinus* (Miller).

Arvicola chrotorrhinus Miller, Proc. Bost. Soc. Nat. Hist., xxvi, 1894, p. 190.

Microtus chrotorrhinus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 193. Zoöl. Ser.

ROCK MEADOW VOLE.

Type locality. Head of Tuckerman's Ravine, Mount Washington, Coos County, New Hampshire. Altitude, 5,300 feet.

Geogr. Distr. The Catskills, New York, to New Hampshire northward into New Brunswick to Trowser's Lake and also central Quebec.

a. Microtus chrotorrhinus rarus Bangs.

Microtus chrotorrhinus rarus Bangs, Proc. Biol. Soc. Wash., XII, 1898, p. 188. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 193. Zoöl. Ser.

GRAY ROCK MEADOW VOLE.

Type locality. Black Bay, north shore of the Strait of Belle Isle, Labrador.

Geogr. Distr. Known only from the type locality.

596. *Microtus borealis* (Richardson).

Arvicola borealis Rich., Zoöl. Journ., III, 1828, p. 517.

Microtus borealis Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 194. Zoöl. Ser.

NORTHERN MEADOW VOLE.

Type locality. Great Bear Lake, Mackenzie, Canada.

Geogr. Distr. Arctic America. Range unknown.

Subgenus B. *Lagurus*.

597. *Microtus pauperrimus* (Cooper).

Arvicola pauperrima Cooper, Amer. Nat., II, 1868, p. 535.

Microtus pauperrimus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 195. Zoöl. Ser.

PIGMY MEADOW VOLE.

Type locality. Plains of the Columbia near the Snake River, southwestern Washington.

Geogr. Distr. Washington and Oregon, central Idaho, and the north slope of the Uinta Mountains, Utah.

598. *Microtus curtatus* (Cope).

Arvicola curtata Cope, Proc. Acad. Nat. Scien. Phil., 1868, p. 2.

Arvicola decurtata Coues, Mon. N. Am. Rod., 1877, p. 215.

Microtus curtatus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 195, fig. 39. Zoöl. Ser.

SHORT-TAILED MEADOW VOLE.

Type locality. Pigeon Spring, Mount Magruder, Nevada.

Geogr. Distr. Eastern California, north of Death Valley and western Nevada.

599. *Microtus pallidus* (Merriam).

Arvicola pallidus! Merr., Amer. Nat., XXII, 1888, p. 702.

Microtus pallidus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 195. Zoöl. Ser.

PALLID MEADOW VOLE.

Type locality. Fort Buford, Howard County, North Dakota.

Geogr. Distr. North Dakota, Montana, north to Calgary, Alberta.

Subgenus C. *Pedomys*.**600. *Microtus austerus* (Le Conte).**

Arvicola austerus! Le Conte, Proc. Acad. Nat. Scien. Phil., 1853,
p. 405.

Arvicola (Pedomys) cinnamomea Baird, N. Am. Mamm., 1857,
p. 541.

?*Hypudæus ochrogaster* Wagn., Schreb. Säugth., Suppl., III, 1843,
p. 592.

Microtus austerus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II,
1901, p. 196, fig. 40. Zoöl. Ser.

PRAIRIE MEADOW VOLE.

Type locality. Racine, Racine County, Wisconsin.

Geogr. Distr. Mississippi Valley, from southern Wisconsin to southern Missouri and Fort Reno, Oklahoma; west to eastern Kansas and Nebraska.

601. *Microtus ludovicianus* Bailey.

Microtus ludovicianus Bailey, N. Am. Faun., No. 17, 1900, p. 74.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 197.
Zoöl. Ser.

LOUISIANA MEADOW VOLE.

Type locality. Iowa, Calcasieu Parish, Louisiana.

Geogr. Distr. Coast prairie of southwestern Louisiana.

602. *Microtus haydeni* (Baird).

Arvicola haydenii Baird, N. Am. Mamm., 1857, p. 543.

Microtus haydeni Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II,
1901, p. 197. Zoöl. Ser.

HAYDEN'S MEADOW VOLE.

Type locality. Fort Pierre, Stanley County, South Dakota.

Geogr. Distr. Fort Pierre and the Black Hills, South Dakota, to the North Platte River, western Nebraska; eastern Colorado, Wyoming, and southern Montana.

603. *Microtus minor* (Merriam).

Arvicola austerus! minor Merr., Amer. Nat., xxii, 1888, p. 508.

Microtus minor Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II,
1901, p. 197. Zoöl. Ser.

LEAST UPLAND MEADOW VOLE.

Type locality. Bottineau at base of Turtle Mountains, Bottineau County, North Dakota.

Geogr. Distr. From Edmonton, Alberta, Canada, to northwestern North Dakota, and southeastwardly to Minneapolis, Minnesota.

Subgenus D. *Pitymys*.**604. *Microtus pinetorum* (Le Conte).**

Psammomys pinetorum Le Conte, Ann. N. Y. Lyc. Nat. Hist., III, 1830, p. 133, pl. II.

Microtus pinetorum Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 198, fig. 41. Zool. Ser.

PINE MEADOW VOLE.

Type locality. Probably near Riceboro, Liberty County, Georgia.
Geogr. Distr. North and South Carolina, Georgia, and Florida.

a. Microtus pinetorum scalopsoides (Audubon & Bachman).

Arvicola scalopsoides Aud. & Bachm., Proc. Acad. Nat. Scien. Phil., 1841, p. 97.

Arvicola apella Le Conte, Proc. Acad. Nat. Scien. Phil., 1853, p. 405.

Arvicola kennicotti Baird, N. Am. Mamm., 1857, p. 547.

Microtus pinetorum scalopsoides Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 198. Zool. Ser.

MOLE-LIKE MEADOW VOLE.

Type locality. Long Island, New York.

Geogr. Distr. Southern New York, grading along the coast into *M. pinetorum*; westward to Illinois.

b. Microtus pinetorum auricularis Bailey.

Microtus pinetorum auricularis Bailey, Proc. Biol. Soc. Wash., XII, 1898, p. 90. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 198. Zool. Ser.

BLUEGRASS MEADOW VOLE.

Type locality. Washington, Adams County, Mississippi.

Geogr. Distr. Alleghany Mountains to the Mississippi River, in the States of Tennessee, Kentucky, and northern Mississippi.

605. *Microtus nemoralis* (Bailey).

Microtus pinetorum nemoralis Bailey, Proc. Biol. Soc. Wash., XII, 1898, p. 89.

Microtus nemoralis Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 199. Zool. Ser.

WOODLAND MEADOW VOLE.

Type locality. Stilwell, Boston Mountains, Indian Territory.

Geogr. Distr. Indian Territory and Arkansas north to Council Bluffs, Iowa.

606. *Microtus quasiater* (Coues).

Arvicola (Pitymys) pinetorum quasiater Coues, Proc. Acad. Nat. Scien. Phil., 1874, p. 191.

Microtus quasiater Elliot, Mamm. Middle Amer. & W. Indies,
F. C. M. Pub., IV, Pt. 1, 1904, p. 303, fig. 52. Zoöl. Ser.

JALAPA MEADOW VOLE.

Type locality. Jalapa, State of Vera Cruz, Mexico.

Geogr. Distr. Central portions of the State of Vera Cruz, and eastern part of State of Puebla, Mexico. Altitude, 4,000 to 5,000 feet.

Subgenus E. *Chilotus*.

607. *Microtus oregoni* (Bachman).

Arvicola oregoni Bachm., Journ. Acad. Nat. Scien. Phil., 1839,
p. 60, pl. 1.

Microtus morosus Elliot, Pub. Field Columb. Mus., 1, 1899, p. 227.
Zoöl. Ser.

Microtus oregoni Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11,
1901, p. 200, fig. 42. Zoöl. Ser.

OREGON MEADOW VOLE.

Type locality. Astoria, Clatsop County, Oregon.

Geogr. Distr. Pacific coast region from northern California to Puget Sound.

608. *Microtus serpens* Merriam.

Microtus serpens Merr., Proc. Biol. Soc. Wash., XI, 1897, p. 75.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 200.
Zoöl. Ser.

CREEPING MEADOW VOLE.

Type locality. Agassiz, British Columbia, Canada.

Geogr. Distr. Southern British Columbia and northern Washington, between the Cascade Mountains and Puget Sound.

609. *Microtus bairdi* Merriam.

Microtus bairdi Merr., Proc. Biol. Soc. Wash., XI, 1897, p. 74.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 200.
Zoöl. Ser.

BAIRD'S MEADOW VOLE.

Type locality. Glacier Peak, Crater Lake, Klamath County, Oregon. Altitude, 7,800 feet.

Geogr. Distr. Known only from the type locality.

Subgenus F. *Arvicola*.

610. *Microtus richardsoni* (De Kay).

Arvicola richardsoni De Kay, Zoöl. N. York, Mamm., 1842, p. 91.

Microtus richardsoni Elliot, Syn. N. Am. Mamm., F. C. M. Pub.,
11, 1901, p. 200. Zoöl. Ser.

RICHARDSON'S MEADOW VOLE.

Type locality. "Near the foot of the Rocky Mountains"; vicinity of Jasper House, Alberta, Canada.

Geogr. Distr. Vicinity of Jasper House and of Henry House, Alberta, Canada.

a. *Microtus richardsoni macropus* (Merriam).

Arvicola macropus Merr., N. Am. Faun., No. 5, 1891, p. 59.

Microtus richardsoni macropus Elliot, Syn. N. Am. Mamm.,

F. C. M. Pub., II, 1901, p. 201, fig. 43 (*M. macropus*). Zoöl. Ser.

BIG-FOOTED MEADOW VOLE.

Type locality. Pahsimeroi Mountains, Idaho. Altitude, 9,700 feet.

Geogr. Distr. Rocky Mountains from the Wasatch north to Canada, the Wind River Mountains of Wyoming, the Blue Mountains of Oregon, and intermediate ranges. High altitudes.

b. *Microtus richardsoni arvicoloides* (Rhoads).

Aulacomys arvicoloides Rhoads, Amer. Nat., xxviii, 1894, p. 182.

Microtus principalis Rhoads, Amer. Nat., xxix, 1895, p. 940.

Microtus richardsoni arvicoloides Elliot, Syn. N. Am. Mamm.,

F. C. M. Pub., II, 1901, p. 202. Zoöl. Ser.

CASCADE MOUNTAINS MEADOW VOLE.

Type locality. Lake Kichelos, Kittitas County, Washington. Altitude, 8,000 feet.

Geogr. Distr. Cascade range in Oregon and Washington. High elevations.

Subgenus G. *Orthriomys*.611. *Microtus umbrosus* Merriam.

Microtus umbrosus Merr., Proc. Biol. Soc. Wash., xii, 1898, p. 107.

Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv,

Pt. I, 1904, p. 305, fig. 53. Zoöl. Ser.

MOUNT ZEMPOALTEPEC MEADOW VOLE.

Type locality. Mount Zempoaltepec, State of Oaxaca, Mexico. Altitude, 8,200 feet.

Geogr. Distr. Known only from the type locality.

Subgenus H. *Herpetomys*.612. *Microtus guatemalensis* Merriam.

Microtus guatemalensis Merr., Proc. Biol. Soc. Wash., xii, 1898,

p. 108. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M.

Pub., iv, Pt. I, 1904, p. 306, fig. 54. Zoöl. Ser.

GUATEMALAN MEADOW VOLE.

Type locality. Todos Santos, Huehuetenango, Guatemala. Altitude, 10,000 feet.

Geogr. Distr. Known only from the type locality.

Subgenus I. *Neofiber*.**613. *Microtus alleni* (True).**

Neofiber alleni True, Science, IV, 1884, p. 34.

Microtus alleni Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 203, fig. 44. Zoöl. Ser.

FLORIDA WATER VOLE.

Type locality. Georgiana, Merritt's Island, Brevard County, Florida.

Geogr. Distr. Eastern Florida from Indian River on the north to Eden on the south, and to Gainesville in the interior. The limits of this species are not exactly known; and it may go beyond the boundaries given.

Genus 82. **Fiber. Musk Rats.**

Fiber G. Cuv., Leçons d'Anat. Comp., I, 1800, tab. I. Type *Castor zibethicus* Linnæus.

Ondatra Lacép., Tab. Div. Sous-div. Ord. et Gen. Mamm., 1799, p. 9. (nec Link.)

Moschomys Billb., Syn. Faun. Scandinav., I, 1828, Mamm., Consp. A.

614. *Fiber zibethicus* (Linnæus).

Castor zibethicus Linn., Syst. Nat., I, 1766, p. 79.

Fiber osoyoosensis Lord, Proc. Zoöl. Soc., 1863, p. 97.

Fiber zibethicus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 212, fig. 49. Zoöl. Ser.

MUSK RAT. *Ondatra*, Indian name.

Type locality. Eastern Canada.

Geogr. Distr. Labrador to the Gulf States, excepting possibly the Dismal Swamp, Virginia, and from the Atlantic Coast to the Rocky Mountains north of the Gulf States and Arizona, and south of Keewatin, Canada.

a. *Fiber zibethicus ripensis* Bailey.

Fiber zibethicus ripensis Bailey, Proc. Biol. Soc. Wash., xv, 1902, p. 119.

PECOS RIVER MUSK RAT.

Type locality. Pecos River, at Carlsbad, Eddy County, New Mexico.

Geogr. Distr. Unknown.

Genl. Char. Size small; color paler than *F. zibethicus*. Skull narrow; nasals short, rounded at posterior end; incisors heavy, molars light.

Color. Above light brown; belly buff brown; throat gray with a dusky stripe on the chin; white spot over each nostril.

**Measurements.* Total length, 470; tail vertebræ, 202; hind foot, 67. Skull: basal length, 55; length of nasals, 18; zygomatic width, 35; mastoid breadth, 25; alveolar length of upper molar series, 15.

b. Fiber zibethicus pallidus Mearns.

Fiber zibethicus pallidus Mearns, Bull. Am. Mus. Nat. Hist. N. Y., 1890, p. 280. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 213. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 308, fig. 55. Zoöl. Ser.

PALE MUSK RAT.

Type locality. Fort Verde, Yavapai County, Arizona.

Geogr. Distr. Lower California and State of Sonora, Mexico, north to Montana.

c. Fiber zibethicus rivalicius Bangs.

Fiber zibethicus rivalicius Bangs, Proc. Bost. Soc. Nat. Hist., 1895, p. 541. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 213. Zoöl. Ser.

RELATED MUSK RAT.

Type locality. Burbridge, Plaquemines Parish, Louisiana.

Geogr. Distr. Lower Mississippi region and swampy coasts of Louisiana and Alabama.

d. Fiber zibethicus aquilonius Bangs.

Fiber zibethicus aquilonius Bangs, Proc. N. Engl. Zoöl. Club, 1, 1899, p. 11. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 213. Zoöl. Ser.

NORTHERN MUSK RAT.

Type locality. Rigoulette, Hamilton Inlet, Labrador.

Geogr. Distr. Labrador. Range unknown.

e. Fiber zibethicus hudsonius Preble.

Fiber zibethicus hudsonius Preble, N. Am. Faun., No. 22, 1902, p. 53.

HUDSON BAY MUSK RAT.

Type locality. Fort Churchill, Keewatin, Canada.

Geogr. Distr. Keewatin, Canada. Range unknown.

Genl. Char. Size small. Skull with high interorbital crest; molars small.

Color. Above dull yellowish brown, dorsal region darker; throat yellowish white; cheeks yellowish fawn; middle of chin dusky; under parts yellowish brown.

Measurements. Total length, 539; tail vertebræ, 225; hind foot, 74. Skull: occipito-nasal, 56; zygomatic breadth, 38; interorbital breadth, 6; length of nasals, 20; widest part of nasals, 9.5; length of upper molar series, 13.5.

615. *Fiber spatulatus* Osgood.

Fiber spatulatus Osgood, N. Am. Faun., No. 19, 1900, p. 36.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 214.

Zoöl. Ser.

NORTHWEST TERRITORY MUSK RAT.

Type locality. Lake Marsh, Yukon, Canada.

Geogr. Distr. Northwest Canada. Range unknown.

616. *Fiber occipitalis* Elliot.

Fiber occipitalis Elliot, Pub. Field Columb. Mus., 111, 1903, p. 162,

fig. Zoöl. Ser.

OREGON MUSK RAT.

Type locality. Florence, Oregon.

Geogr. Distr. Known only from the type locality.

Genl. Char. Largest species of the genus. Superior outline of occipital bowed inward at parietal, nearly forming a half circle; interorbital constriction greatly pronounced; frontal crest marked; postorbital process pointing downward, acute; brain case narrow posteriorly; interparietal nearly as long as broad; parietals pointed anteriorly, very narrow posteriorly; face of occipital broader in proportion to its length than that of *F. zibethicus*; coronoid process of mandible narrow, acute; condyle broad and heavy; molars smaller than those of *F. zibethicus*; incisors short, heavy.

Color. Above reddish brown with black hairs mixed; top of head, nose, and around eyes blackish; lips white; beneath rufous; inner side of thighs and anal region white; hands and feet blackish brown; tail black.

Measurements. Total length, 597; tail vertebræ, 295; hind foot, 85; ear, 21. Skull: occipito-nasal length, 67; Hensel, 60; zygomatic width, 43; interorbital constriction, 6.5; across postorbital processes, 23; length of nasals, 27; width of nasals posteriorly, 2.5; anteriorly, 8; greatest width of rostrum, 11; palatal length, 22.

617. *Fiber obscurus* Bangs.

Fiber obscurus Bangs, Proc. Biol. Soc. Wash., 19, 1894, p. 133.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 214.

Zoöl. Ser.

DUSKY MUSK RAT.

Type locality. Codroy, Newfoundland.

Geogr. Distr. Newfoundland. Range unknown.

618. *Fiber macrodon* Merriam.

Fiber macrodon Merr., Proc. Biol. Soc. Wash., XI, 1897, p. 143.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 214.

Zoöl. Ser.

LARGE-TOOTHED MUSK RAT.

Type locality. Lake Drummond, Dismal Swamp, Norfolk County, Virginia.

Geogr. Distr. Known only from the Dismal Swamp, Virginia.

Genus 83. *Synaptomys*. Lemming Voles.

Synaptomys Baird, N. Am. Mamm., 1857, p. 558. Type *Synaptomys cooperi* Baird.

Subgenus A. *Synaptomys*.619. *Synaptomys cooperi* Baird.

Synaptomys cooperi Baird, N. Am. Mamm., 1857, p. 558. Elliot,

Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 203, fig. 45.

Zoöl. Ser.

COOPER'S LEMMING VOLE.

Type locality. Unknown. Southern New York, or northern New Jersey.

Geogr. Distr. Eastern Massachusetts to Minnesota, south to North Carolina, Tennessee, Indiana, and Iowa.

a. *Synaptomys cooperi stonii* (Rhoads).

Synaptomys stonei Rhoads, Amer. Nat., XXVII, 1893, p. 53.

Synaptomys helaletes Merr., Proc. Biol. Soc. Wash., X, 1896, p. 59.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 204.

Zoöl. Ser.

STONE'S LEMMING VOLE.

Type locality. Mays Landing, Atlantic County, New Jersey.

Geogr. Distr. Sphagnum bogs of the Carolina fauna.

b. *Synaptomys cooperi gossi* (Merriam).

Synaptomys helaletes gossii Merr., Proc. Biol. Soc. Wash., X, 1896,

p. 60. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901,

p. 204. Zoöl. Ser.

Synaptomys cooperi gossi Rhoads, Proc. Acad. Nat. Scien. Phil.,

1897, p. 307.

GOSS' LEMMING VOLE.

Type locality. Neosho Falls, Woodson County, Kansas.

Geogr. Distr. Kansas. Range unknown.

620. *Synaptomys fatuus* Bangs.

Synaptomys fatuus Bangs, Proc. Biol. Soc. Wash., x, 1896, p. 47.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 204.

Zoöl. Ser.

SILLY LEMMING VOLE.

Type locality. Lake Edward, Quebec, Canada.

Geogr. Distr. Lake Edward to Victoria County, New Brunswick, and Godbout, Quebec, and west to Peninsula Harbor, northeastern Lake Superior, Ontario.

Subgenus B. *Mictomys*.621. *Synaptomys innuitus* (True).

Mictomys innuitus True, Proc. U. S. Nat. Mus., 1894, p. 243.

Synaptomys innuitus Elliot, Syn. N. Am. Mamm., F. C. M. Pub.,

II, 1901, p. 205, fig. 46. Zoöl. Ser.

INNUIT LEMMING VOLE.

Type locality. Fort Chimo, Ungava, Labrador.

Geogr. Distr. Labrador. Range unknown.

a. *Synaptomys innuitus medioximus* Bangs.

Synaptomys innuitus medioximus Bangs, Proc. N. Engl. Zoöl.

Club, II, 1900, p. 40. Elliot, Syn. N. Am. Mamm., F. C. M.

Pub., II, 1901, Append., p. 430. Zoöl. Ser.

LABRADOR LEMMING VOLE.

Type locality. Lance au Loup, Strait of Belle Isle, Labrador.

Geogr. Distr. Labrador. Range unknown.

622. *Synaptomys sphagnicola* Preble.

Synaptomys sphagnicola Preble, Proc. Biol. Soc. Wash., XIII, 1899,

p. 43. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901,

p. 206. Zoöl. Ser.

WHITE MOUNTAINS LEMMING VOLE.

Type locality. Fabyans, Coos County, New Hampshire.

Geogr. Distr. White Mountains, New Hampshire. Range unknown.

623. *Synaptomys wrangeli* Merriam.

Synaptomys wrangeli Merr., Proc. Biol. Soc. Wash., x, 1896, p. 63.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 206.

Zoöl. Ser.

WRANGEL'S LEMMING VOLE.

Type locality. Wrangel, Alaska.

Geogr. Distr. Southwestern Alaska. Range unknown.

624. *Synaptomys andersoni* Allen.

Synaptomys andersoni Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1903, p. 554.

ANDERSON'S LEMMING VOLE.

Type locality. Level Mountain, northern British Columbia, Canada.

Geogr. Distr. British Columbia. Range unknown.

Genl. Char. Skull similar to that of *S. wrangeli*, but bullæ more-inflated, especially anteriorly; posterior loop of last upper molar about two-thirds as large as the middle loop; reëtrant angle on outer side of last molar nearly as deep as that in similar tooth of *S. truii*.

Color. Above dark brown varied with blackish and suffused with clay color; head and cheeks like back; flanks yellowish brown; under surface ashy gray; tail with dorsal third blackish; sides and lower surface dull grayish; upper surface of hands and feet blackish brown.

Measurements. Total length, 120; tail vertebræ, 25; hind foot, 18; ear, 11.5. Skull: occipito-nasal length, 24.6; length of nasals, 23; zygomatic width, 14.5; mastoid breadth, 11.5; interorbital constriction, 3; length of braincase, 14; length of rostrum, 11; length of nasals, 6; length of incisive foramina, 4.2; length of upper tooth row, 6.6.

625. *Synaptomys chapmani* Allen.

Synaptomys chapmani Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1903, p. 555.

CHAPMAN'S LEMMING VOLE.

Type locality. Glacier, Selkirk Range, British Columbia, Canada.

Geogr. Distr. British Columbia. Range unknown.

Genl. Char. Larger than *S. andersoni*, and skull narrower; rostrum longer; incisive foramina longer and narrower, and bullæ less inflated.

Color. Above grayish brown tinged with buff, strongest on the head; beneath dark gray, base of fur plumbeous; feet grayish brown; tail dark; side glands in front of hips covered with whitish fur.

Measurements. Total length, 134; tail vertebræ, 21; hind foot, 20; ear (from notch in dry skin), 11. Skull: occipito-nasal length, 26; basal length, 24.5; zygomatic breadth, 15.2; mastoid breadth, 11.5; interorbital constriction, 3; length of braincase, 13.5; length of rostrum, 12.2; length of nasals, 6.6; length of palatal foramina, 5; length of upper tooth row, 7.

626. *Synaptomys dalli* Merriam.

Synaptomys dalli Merr., Proc. Biol. Soc. Wash., x, 1896, p. 62.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 206.

Zoöl. Ser.

DALL'S LEMMING VOLE.

Type locality. Nulato, Alaska.

Geogr. Distr. Valley of the Yukon, western Alaska. Range unknown.

627. *Synaptomys bullatus* Preble.

Synaptomys bullatus Preble, Proc. Biol. Soc. Wash., xv, 1902,

p. 181.

PREBLE'S LEMMING VOLE.

Type locality. Trout Rock near Fort Rae, Great Slave Lake, Mackenzie, Canada.

Geogr. Distr. Unknown.

Genl. Char. Skull similar to that of *S. dalli*, but pelage less reddish; hind foot smaller.

Color. Above black and reddish brown, black predominating; sides lighter; beneath ashy plumbeous; feet dusky brown; tail above dusky, beneath ashy.

Measurements. Total length 130; tail vertebræ, 24; hind foot, 18. Skull: occipito-nasal length, 26; zygomatic width, 16; inter-orbital constriction, 3; length of upper molar series, 7.

628. *Synaptomys truei* Merriam.

Synaptomys truei Merr., Proc. Biol. Soc. Wash., x, 1896, p. 62.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 207.

Zoöl. Ser.

TRUE'S LEMMING VOLE.

Type locality. Skagit Valley, northwestern Washington.

Geogr. Distr. Northwestern Washington. Range unknown.

Genus 84. **Lemmus. Lemmings.**

Lemmus Link, Zool. Beyt. Natur., I, Pt. 2, 1795, p. 75. Type *Mus lemmus* Linnæus.

Myodes Pallas, Zoog. Rosso-Asiat., I, 1811, p. 172.

Lemmus Rocherb., Actes Soc. Linn. Bordeaux, XII, 1841, No. 42, p. 216.

629. *Lemmus trimucronatus* (Richardson).

Arvicola trimucronatus Rich., Journ. Parry's 2d Voy., 1825, Append., p. 309.

- Myodes albigularis* Wagn., Schreb. Säugth., Suppl., III, 1843, p. 602.
Lemmus alascensis Merr., Proc. Wash. Acad. Scien., II, 1900, p. 26.
Lemmus trimucronatus Elliot, Syn. N. Am. Mamm., F. C. M. Pub.,
 II, 1901, p. 207, fig. 47. Zoöl. Ser.

RICHARDSON'S LEMMING.

Type locality. Point Lake, N. Lat. 65°, Mackenzie, Canada.
Geogr. Distr. Arctic regions of North America.

a. *Lemmus trimucronatus helvolus* (Richardson).

Arvicola (*Georychus*?) *helvolus* Rich., Faun. Bor. Amer., Mamm.,
 1829, p. 128.
Lemmus trimucronatus helvolus Elliot, Syn. N. Am. Mamm.,
 F. C. M. Pub., II, 1901, p. 208. Zoöl. Ser.

YELLOWISH LEMMING.

Type locality. Alpine swamps, Arctic America, 56° north latitude.
Geogr. Distr. Arctic regions of North America.

b. *Lemmus trimucronatus yukonensis* (Merriam).

Lemmus yukonensis Merr., Proc. Wash. Acad. Scien., II, 1900, p. 27.
Lemmus trimucronatus yukonensis Elliot, Syn. N. Am. Mamm.,
 F. C. M. Pub., II, 1901, p. 208. Zoöl. Ser.

YUKON LEMMING.

Type locality. Charlie Creek, Yukon River, Alaska.
Geogr. Distr. Yukon River District, Alaska. Range unknown.

630. *Lemmus minusculus* Osgood.

Lemmus minusculus Osgood, N. Am. Faun., No. 24, 1904, p. 36.

SMALL LEMMING.

Type locality. Kakhtul River near junction with the Malchatna River, Alaska.

Geogr. Distr. Alaska. Range unknown.

Genl. Char. "Similar in general to *L. alascensis* (*L. trimucronatus*) but much smaller." Skull with naso-frontal region elevated; rostrum depressed.

Color. Upper parts mixed ochraceous and black, becoming light chestnut on the rump; blackish median line from nose to shoulders; lower sides and under parts ochraceous or tawny ochraceous; ears dusky; hands and feet seal brown; tail sometimes uniform pale buff above and below, again in other specimens blackish above, light buff below.

Measurements. Average of 10 males. Total length, 131; tail vertebræ, 12; hind foot, 19. Skull: greatest length, 28.5; Hensel, 25.4; zygomatic width, 19; mastoid width, 15; length of nasals, 8.9; postpalatal length, 12.2; length of upper molars, 8.

631. Lemmus nigripes (True).

Myodes nigripes True, Proc. U. S. Nat. Mus., xvii, 1894, p. 242.

Lemmus nigripes Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 209. Zool. Ser.

BLACK-FOOTED LEMMING.

Type locality. St. George Island, Pribilof Islands, Bering Sea.

Geogr. Distr. Known only from the type locality.

Genus 85. Dicrostonyx. Forked-claw Lemmings.

Dicrostonyx Glog., Gem. Hand- u. Hilfsb. Naturg., 1841, pp. xxxi, 97. Type *Mus hudsonius* Pallas.

Cuniculus Wagl., Nat. Syst. Amph., 1830, p. 121.

Myolemmus Pomel, Cat. Méth. Vert. Foss. Bass. Loire, 1854, p. 27.

Misothermus Hensel, Zeitsch. Deutsch. Geol. Gesellsch., VII, 1855, p. 492. (Part.)

Borioikon Poliakoff, Ann. au tome xxxix, Mém. Acad. St. Pétersb., No. 2, 1881, p. 35.

Tylonyx Schulze, Mamm. Europ. Hel. Abhandl. u. Vorträge Gesammtg. Naturw., Berlin, xiv, 1897, p. 83.

632. Dicrostonyx hudsonius (Pallas).

Mus hudsonius Pall., Nov. Spec. Quad. Glir. Ord., 1778, p. 208.

Dicrostonyx hudsonius Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 209. Zool. Ser.

HUDSONIAN LEMMING.

Type locality. Labrador.

Geogr. Distr. Labrador. Range unknown.

a. Dicrostonyx hudsonius unalascensis (Merriam).

Dicrostonyx unalascensis Merr., Proc. Wash. Acad. Scien., II, 1900, p. 25.

Dicrostonyx hudsonius unalascensis Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 210. Zool. Ser.

UNALASKA LEMMING.

Type locality. Unalaska, Alaska.

Geogr. Distr. Range unknown.

b. Dicrostonyx hudsonius nelsoni (Merriam).

Dicrostonyx nelsoni Merr., Proc. Wash. Acad. Scien., II, 1900, p. 25.

Dicrostonyx hudsonius alascensis Stone, Proc. Acad. Nat. Scien. Phil., 1900, p. 37.

Dicrostonyx hudsonius nelsoni Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 210, fig. 48. Zool. Ser.

NELSON'S LEMMING.

Type locality. St. Michaels, Alaska.

Geogr. Distr. Range unknown.

c. *Dicrostonyx hudsonius richardsoni* (Merriam).

Dicrostonyx richardsoni Merr., Proc. Wash. Acad. Scien., II, 1900, p. 26.

Dicrostonyx hudsonius richardsoni Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 211. Zool. Ser.

FORT CHURCHILL LEMMING.

Type locality. Fort Churchill, west shore of Hudson Bay, Keewatin, Canada.

Geogr. Distr. Hudson Bay region. Range unknown.

Family V. **Geomyidae. Pouched Rats.**

Genus 86. **Geomys. Pocket Gophers.**

Geomys Rafin., Am. Month. Mag., II, No. 1, 1817, p. 45. Type *Geomys pinetis* Rafin. = *Mus tuza* Ord.

Diplostoma Rafin., Am. Month. Mag., II, No. 1, 1817, p. 44.

Saccophorus Kuhl, Beitr. Zool. und vergl. Anat., 1820, p. 65.

Pseudostoma Say, Long's Exped. Rocky Mts., I, 1823, p. 406.

Ascomys Licht., Abhand. K. Akad. Wiss. Berl., 1825, p. 20, fig. 2.

633. **Geomys tuza** (Ord).

Mus tuza Ord, Guthr. Geogr., 2d Amer. ed., II, 1815, p. 292.

Geomys pinetis Rafin., Am. Month. Mag., II, No. 1, 1817, p. 45.

Geomys tuza Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 215, fig. 50. Zool. Ser.

PINE BARRENS POCKET GOPHER.

Type locality. Pine barrens near Augusta, Richmond County, Georgia.

Geogr. Distr. Pine barrens of Georgia, and possibly northern Florida.

a. *Geomys tuza floridanus* (Audubon & Bachman).

Pseudostoma floridana Aud. & Bachm., N. Am. Quad., III, 1854, p. 242.

Geomys tuza floridanus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 215. Zool. Ser.

FLORIDA POCKET GOPHER.

Type locality. St. Augustine, St. John County, Florida.

Geogr. Distr. Peninsula of Florida.

b. *Geomys tuza mobilensis* Merriam.

Geomys tuza mobilensis Merr., N. Am. Faun., No. 8, 1895, p. 119, pls. and figs. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 216. Zoöl. Ser.

Geomys floridanus austrinus Bangs, Proc. Bost. Soc. Nat. Hist., 1898, p. 177.

Geomys tuza austrinus Elliot, Pub. Field Columb. Mus., 111, 1901, p. 49. Zoöl. Ser. *Id.* Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 217. Zoöl. Ser.

MOBILE POCKET GOPHER.

Type locality. Mobile Bay, Baldwin County, Alabama.

Geogr. Distr. Southern Alabama, and adjacent northwest and western Florida.

634. *Geomys colonus* Bangs.

Geomys colonus Bangs, Proc. Bost. Soc. Nat. Hist., 1898, p. 178.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 217. Zoöl. Ser.

FARMER POCKET GOPHER.

Type locality. St. Mary's, Camden County, Georgia.

Geogr. Distr. Georgia. Range unknown.

635. *Geomys cumberlandius* Bangs.

Geomys cumberlandius Bangs, Proc. Bost. Soc. Nat. Hist., 1898.

p. 180. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 217. Zoöl. Ser.

CUMBERLAND POCKET GOPHER.

Type locality. "Stafford Place," Cumberland Island, Camden County, Georgia.

Geogr. Distr. Cumberland Island, Camden County, Georgia.

636. *Geomys bursarius* (Shaw).

Mus bursarius Shaw, Trans. Linn. Soc., v, 1800, p. 227, pl. 8.

Diplostoma fusca Rafin., Am. Month. Mag., 11, No. 1, 1817, p. 45.

Geomys cinerea Rafin., Am. Month. Mag., 11, No. 1, 1817, p. 45.

Saccophorus bursarius Kuhl, Beitr. Zool., 1820, p. 65.

Mus saccatus Mitch., N. Y. Med. Repos., vi, 1821, p. 249.

Ascomys canadensis Licht., Abh. K. Akad. Wiss. Berlin, 1825, p. 20, fig. 2.

Geomys oregonensis Le Conte, Proc. Acad. Nat. Scien. Phil., vi, 1852, p. 160.

Geomys bursarius Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 217. Zoöl. Ser.

SHAW'S POCKET GOPHER.

Type locality. Unknown. Mississippi Valley?

Geogr. Distr. Canadian boundary, longitude 97°, south to eastern Kansas, southeastern Missouri and southern Illinois, east nearly to Lake Michigan and west to the Dakotas and Nebraska.

637. *Geomys lutescens* (Merriam).

Geomys bursarius lutescens Merr., N. Am. Faun., No. 4, 1890, p. 51.

Geomys lutescens Merr., Mon. Geomyidæ, N. Am. Faun., No. 8, 1895, p. 127. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 218. Zoöl. Ser.

PALE POCKET GOPHER.

Type locality. Sandhills on Birdwood Creek, Lincoln County, western Nebraska.

Geogr. Distr. Great Plains, and southwestern South Dakota, southward to Colorado, western Nebraska, western Kansas, eastern Colorado, western Oklahoma, and western Texas.

638. *Geomys breviceps* Baird.

Geomys breviceps Baird, Proc. Acad. Nat. Scien. Phil., 1855, p. 335. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 218. Zoöl. Ser.

SHORT-HEADED POCKET GOPHER.

Type locality. Prairie Mer Rouge, Morehouse Parish, Louisiana.

Geogr. Distr. Mississippi Valley and Gulf Coast, in southern Arkansas, Louisiana, and Texas, and valley of the Arkansas River, north nearly to southern Kansas, and west to near 98th meridian.

*a. *Geomys breviceps sagittalis* Merriam.*

Geomys breviceps sagittalis Merr., Mon. Geomyidæ, N. Am. Faun., No. 8, 1895, p. 134, pl. 9, fig. 4. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 218. Zoöl. Ser.

CRESTED POCKET GOPHER.

Type locality. Clear Creek, Galveston Bay, Texas.

Geogr. Distr. Gulf Coast of Texas about Galveston Bay.

*b. *Geomys breviceps attwateri* Merriam.*

Geomys breviceps attwateri Merr., Mon. Geomyidæ, N. Am. Faun., No. 8, 1895, p. 135, pl. 9, fig. 3. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 219. Zoöl. Ser.

ATTWATER'S POCKET GOPHER.

Type locality. Rockport, Aransas County, Texas.

Geogr. Distr. Coast and islands of Texas, between Matagorda and Nueces Bay; also in interior and vicinity of San Antonio.

639. *Geomys texensis* Merriam.

Geomys texensis Merr., Mon. Geomyidæ, N. Am. Faun., No. 8, 1895, p. 137, pls. and figs. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 219. Zoöl. Ser.

TEXAS POCKET GOPHER.

Type locality. Mason, Mason County, Texas.

Geogr. Distr. Mason County, central Texas, and probably south to the Rio Grande.

640. *Geomys arenarius* Merriam.

Geomys arenarius Merr., Mon. Geomyidæ, N. Am. Faun., No. 8, 1895, p. 139, pls. and figs. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 219. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 310, fig. 56. Zoöl. Ser.

SAND-FREQUENTING POCKET GOPHER.

Type locality. El Paso, El Paso County, Texas.

Geogr. Distr. Valley of the Upper Rio Grande from El Paso and Juarez, State of Chihuahua, Mexico, north to Las Cruces and west to Deming, New Mexico.

641. *Geomys personatus* True.

Geomys personatus True, Proc. U. S. Nat. Mus., XI, 1889, p. 159. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 220. Zoöl. Ser.

MASKED POCKET GOPHER.

Type locality. Padre Island, Cameron County, Texas.

Geogr. Distr. Padre Island and mainland to Carrizo on the Rio Grande.

a. Geomys personatus fallax Merriam.

Geomys personatus fallax Merr., Mon. Geomyidæ, N. Am. Faun., No. 8, 1895, p. 144, pl. 12, fig. 3. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 220. Zoöl. Ser.

DECEITFUL POCKET GOPHER.

Type locality. South side of Nueces Bay, Cameron County, Texas.

Geogr. Distr. South shore of Nueces Bay and lower Nueces River, Texas.

Genus 87. **Cratogeomys. Powerful Pocket Gophers.**

Cratogeomys Merr., Mon. Geomyidæ, N. Am. Faun., No. 8, 1895, p. 150. Type *Geomys merriami* Thomas.

642. *Cratogeomys merriami* (Thomas).

Geomys merriami Thomas, Ann. Mag. Nat. Hist., 6th Ser., xii, 1893, p. 271. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 313. Zoöl. Ser.

MERRIAM'S POCKET GOPHER.

Type locality. "Southern Mexico." Valley of Mexico?

Geogr. Distr. Valley of Mexico and Toluca Valley, State of Mexico, and States of Puebla and Hidalgo. "South end of Valley of Mexico and adjacent mountain slopes from just below the lower edge of the lower pine belt to an altitude of 10,000 to 11,000 feet; east to Atlitxco, State of Puebla, north to Irolo, State of Hidalgo, and west to Lerma in Toluca Valley." (Merr.)

643. *Cratogeomys perotensis* Merriam.

Cratogeomys perotensis Merr., Mon. Geomyidæ, N. Am. Faun., No. 8, 1895, p. 154, pl. 8, fig. 6. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 313. Zoöl. Ser.

PEROTE POCKET GOPHER.

Type locality. Cofre de Perote, State of Vera Cruz, Mexico. Altitude, 9,500 feet.

Geogr. Distr. State of Vera Cruz, Mexico.

644. *Cratogeomys estor* Merriam.

Cratogeomys estor Merr., Mon. Geomyidæ, N. Am. Faun., No. 8, 1895, p. 155, pl. 8, figs. 4, 5. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 314. Zoöl. Ser.

LAS VIGAS POCKET GOPHER.

Type locality. Las Vigas, State of Vera Cruz, Mexico. Altitude, 8,000 feet.

Geogr. Distr. Northeastern foothills of the Cofre de Perote, and the wooded hills to the north, State of Vera Cruz, Mexico.

645. *Cratogeomys oreocetes* Merriam.

Cratogeomys oreocetes Merr., Mon. Geomyidæ, N. Am. Faun., No. 8, 1895, p. 156, pl. 8, figs. 1, 2. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 314. Zoöl. Ser.

MOUNTAIN POCKET GOPHER.

Type locality. Mount Popocatepetl, State of Mexico. Altitude, 11,500 feet.

Geogr. Distr. Higher slopes of Mount Popocatepetl, Mexico.

646. *Cratogeomys peregrinus* Merriam.

Cratogeomys peregrinus Merr., Mon. Geomyidæ, N. Am. Faun., No. 8, 1895, p. 158, pl. 8, fig. 3. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 314. Zoöl. Ser.

MOUNT IZTACCIHUATL POCKET GOPHER.

Type locality. Mount Iztaccihuatl, State of Mexico, Mexico.
Altitude, 11,500 feet.

Geogr. Distr. Higher slopes of Mount Iztaccihuatl, Mexico.

647. *Cratogeomys castanops* (Baird).

Pseudostoma castanops Baird, Rep. Stansb. Exped. to Great Salt Lake, 1852, p. 313.

Geomys clarkii Baird, Proc. Acad. Nat. Scien. Phil., 1855, p. 332.

Cratogeomys castanops Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 220, fig. 51. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 315, figs. 57, XLVI. Zoöl. Ser.

CHESTNUT-FACED POCKET GOPHER.

Type locality. Prairie road to Bent's Fort, near the present town of Las Animas, Bent County, Colorado, on the Arkansas River.

Geogr. Distr. States of Chihuahua and Coahuila, Mexico, north to Colorado.

a. *Cratogeomys castanops goldmani* Merriam.

Cratogeomys castanops goldmani Merr., Mon. Geomyidæ, N. Am. Faun., No. 8, 1895, p. 160. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 316. Zoöl. Ser.

GOLDMAN'S POCKET GOPHER.

Type locality. Cañitas, State of Zacatecas, Mexico.

Geogr. Distr. Known only from the type locality.

648. *Cratogeomys fulvescens* Merriam.

Cratogeomys fulvescens Merr., Mon. Geomyidæ, N. Am. Faun., No. 8, 1895, p. 161, pl. 12, fig. 2. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 316. Zoöl. Ser.

FULVOUS POCKET GOPHER.

Type locality. Chalchicomula, State of Puebla, Mexico.

Geogr. Distr. States of Puebla and Tlaxcala, from "Esperanza north to Perote, and west to the northeast base of Mount Malinche in Tlaxcala," Mexico.

Genus 88. *Platygeomys*. Broad-headed Pocket Gophers.

Platygeomys Merr., Mon. Geomyidæ, N. Am. Faun., No. 8, 1895, p. 162, pls. and figs. Type *Geomys gymnurus* Merriam.

649. *Platygeomys gymnurus* (Merriam).

Geomys gymnurus Merr., Proc. Biol. Soc. Wash., VII, 1892, p. 166.

Platygeomys gymnurus Merr., Mon. Geomyidæ, N. Am. Faun., No. 8, 1895, p. 164, pls. and figs. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 318, fig. 58. Zoöl. Ser.

NAKED-TAILED POCKET GOPHER.

Type locality. Zapotlan, State of Jalisco, Mexico.

Geogr. Distr. Valley of Zapotlan and slopes of the Sierra Nevada de Colima, State of Jalisco, Mexico.

650. *Platygeomys fumosus* (Merriam).

Geomys fumosus Merr., Proc. Biol. Soc. Wash., VII, 1892, p. 165.

Platygeomys fumosus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 318. Zoöl. Ser.

SOOTY POCKET GOPHER.

Type locality. Colima City, State of Colima, Mexico.

Geogr. Distr. State of Colima, Mexico.

651. *Platygeomys tylorhinus* Merriam.

Platygeomys tylorhinus Merr., Mon. Geomyidæ, N. Am. Faun., No.

8, 1895, p. 167, pl. 13, fig. 1. Elliot, Mamm. Middle Amer. &

W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 319. Zoöl. Ser.

TOUGH-SKINNED POCKET GOPHER.

Type locality. Tula, State of Hidalgo, Mexico.

Geogr. Distr. States of Hidalgo and Michoacan, on north slope of the Sierra Madre, Mexico.

a. *Platygeomys tylorhinus angustirostris* Merriam.

Platygeomys tylorhinus angustirostris Merr., Proc. Biol. Soc. Wash.,

XVI, 1903, p. 81. Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., IV, Pt. 1, 1904, p. 319. Zoöl. Ser.

SLENDER-NOSED POCKET GOPHER.

Type locality. Patamban, State of Michoacan, Mexico.

Geogr. Distr. State of Michoacan. Range unknown.

652. *Platygeomys planiceps* Merriam.

Platygeomys planiceps Merr., Mon. Geomyidæ, N. Am. Faun., No. 8,

1895, p. 168, pls. and figs. Elliot, Mamm. Middle Amer. &

W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 319. Zoöl. Ser.

FLAT-HEADED POCKET GOPHER.

Type locality. North slope of the Volcano Toluca, State of Mexico, Mexico. Altitude, 8,600 feet.

Geogr. Distr. Volcano Toluca to City of Toluca, State of Mexico, Mexico.

653. *Platygeomys neglectus* Merriam.

Platygeomys neglectus Merr., Proc. Biol. Soc. Wash., xv, 1902, p. 68. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 319. Zoöl. Ser.

NEGLECTED POCKET GOPHER.

Type locality. Mount Cerro de la Calentura, eight miles northwest of Pinal de Amoles, State of Queretaro, Mexico. Altitude, 9,000 feet.

Genus 89. *Pappogeomys*. Ancient Pocket Gophers.

Pappogeomys Merr., Mon. Geomyidæ, N. Am. Faun., No. 8, 1895, p. 145, pls. and figs. Type *Geomys bulleri* Thomas.

654. *Pappogeomys bulleri* (Thomas).

Geomys bulleri Thomas, Ann. Mag. Nat. Hist., 6th Ser., 1892, p. 196.

Geomys nelsoni Merr., Proc. Biol. Soc. Wash., vii, 1892, p. 164.

Pappogeomys bulleri Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 320, fig. 59. Zoöl. Ser.

BULLER'S POCKET GOPHER.

Type locality. Near Talpa, west slope of Sierra de Mascota, State of Jalisco, Mexico. Altitude, 8,500 feet.

Geogr. Distr. State of Jalisco, Mexico.

655. *Pappogeomys albinasus* Merriam.

Pappogeomys albinasus Merr., Mon. Geomyidæ, N. Am. Faun., No. 8, 1895, p. 149. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 321. Zoöl. Ser.

WHITE-NOSED POCKET GOPHER.

Type locality. Atemajac, a suburb of Guadalajara, State of Jalisco, Mexico. Altitude, 8,500 feet.

Geogr. Distr. State of Jalisco, Mexico.

Genus 90. *Orthogeomys*. Straight-headed Pocket Gophers.

Orthogeomys Merr., Mon. Geomyidæ, N. Am. Faun., No. 8, 1895, p. 172, pls. and figs. Type *Geomys scalops* Thomas.

656. *Orthogeomys grandis* (Thomas).

Geomys grandis Thomas, Ann. Mag. Nat. Hist., 6th Ser., xii, 1893, p. 270.

Orthogeomys grandis Elliot, Mamm. Middle Amer. & W. Indies,
F. C. M. Pub., IV, Pt. 1, 1904, p. 322. Zoöl. Ser.

GIANT POCKET GOPHER.

Type locality. Dueñas, Guatemala.

Geogr. Distr. Guatemala. Range unknown.

657. *Orthogeomys nelsoni* Merriam.

Orthogeomys nelsoni Merr., Mon. Geomyidæ, N. Am. Faun., No. 8,
1895, p. 176, fig. 63. Elliot, Mamm. Middle Amer. & W. In-
dies, F. C. M. Pub., IV, Pt. 1, 1904, p. 324. Zoöl. Ser.

NELSON'S POCKET GOPHER.

Type locality. Mount Zempoaltepec, State of Oaxaca, Mexico.
Altitude, 8,000 feet.

Geogr. Distr. Mount Zempoaltepec and adjacent region, State of
Oaxaca, Mexico.

658. *Orthogeomys scalops* (Thomas).

Geomys scalops Thomas, Ann. Mag. Nat. Hist., 6th Ser., XIII,
1894, p. 437.

Orthogeomys scalops Elliot, Mamm. Middle Amer. & W. Indies,
F. C. M. Pub., IV, Pt. 1, 1904, p. 324, fig. 60. Zoöl. Ser.

DIGGER POCKET GOPHER.

Type locality. Tehuantepec, State of Oaxaca, Mexico.

Geogr. Distr. State of Oaxaca, and possibly in State of Chiapas,
Mexico.

659. *Orthogeomys latifrons* Merriam.

Orthogeomys latifrons Merr., Mon. Geomyidæ, N. Am. Faun., No. 8,
1895, p. 178, pls. and figs. Elliot, Mamm. Middle Amer. &
W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 324. Zoöl. Ser.

BRISTLED POCKET GOPHER.

Type locality. Guatemala. Exact locality unknown.

Geogr. Distr. Guatemala. Range unknown.

Genus 91. **Heterogeomys. Distinct Pocket Gophers.**

Heterogeomys Merr., Mon. Geomyidæ, N. Am. Faun., No. 8, 1895,
p. 179, pls. and figs. Type *Geomys hispidus* Le Conte.

660. *Heterogeomys hispidus* (Le Conte).

Geomys hispidus Le Conte, Proc. Acad. Nat. Scien. Phil., 1852,
p. 158.

Heterogeomys hispidus Elliot, Mamm. Middle Amer. & W. Indies,
F. C. M. Pub., IV, Pt. 1, 1904, p. 326. Zoöl. Ser.

HARSH-COATED POCKET GOPHER.

Type locality. Near Jalapa, State of Vera Cruz, México.

Geogr. Distr. State of Vera Cruz, Mexico. Altitude, 4,000 feet.

661. *Heterogeomys torridus* Merriam.

Heterogeomys torridus Merr., Mon. Geomyidæ, N. Am. Faun., No. 8, 1895, p. 183, pls. and figs. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 326, fig. 61. Zoöl. Ser.

TROPICAL POCKET GOPHER.

Type locality. Chichicaxtle, State of Vera Cruz, Mexico.

Geogr. Distr. Lowlands, State of Vera Cruz, through States of Oaxaca and Chiapas, Mexico, into Guatemala.

Genus 92. **Macrogeomys. Large Size Pocket Gophers.**

Macrogeomys Merr., Mon. Geomyidæ, N. Am. Faun., No. 8, 1895, p. 185. Type *Geomys heterodus* Peters.

662. *Macrogeomys heterodus* (Peters).

Geomys heterodus Peters, Monatsb. K. Preuss. Akad. Wiss. Berl., 1865, p. 177, pls. and figs.

Macrogeomys heterodus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 328. Zoöl. Ser.

IRAZU POCKET GOPHER.

Type locality. Costa Rica. Locality unknown.

Geogr. Distr. Irazu Range and possibly other parts of Costa Rica.

663. *Macrogeomys dolichocephalus* Merriam.

Macrogeomys dolichocephalus Merr., Mon. Geomyidæ, N. Am. Faun., No. 8, 1895, p. 189, pls. and figs. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 328. Zoöl. Ser.

NARROW-HEADED POCKET GOPHER.

Type locality. San José, Costa Rica.

Geogr. Distr. Costa Rica. Range unknown.

664. *Macrogeomys cavator* Bangs.

Macrogeomys cavator Bangs, Bull. Mus. Comp. Zoöl., xxxix, 1902, p. 42. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 329. Zoöl. Ser.

BOQUETE POCKET GOPHER.

Type locality. Boquete, Chiriqui, Panama. Altitude, 4,800 feet.

Geogr. Distr. Panama. Range unknown.

665. *Macrogeomys pansa* Bangs.

Macrogeomys pansa Bangs, Bull. Mus. Comp. Zool. xxxix, 1902, p. 44. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 329. Zool. Ser.

BROAD-FOOTED POCKET GOPHER.

Type locality. Bogaba, Chiriqui, Panama. Altitude, 600 feet.

Geogr. Distr. Panama. Range unknown.

666. *Macrogeomys costaricensis* Merriam.

Macrogeomys costaricensis Merr., Mon. Geomyidæ, N. Am. Faun., No. 8, 1895, p. 192, pls. and figs. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 330. Zool. Ser.

PACUARE POCKET GOPHER.

Type locality. Pacuare, Costa Rica.

Geogr. Distr. Costa Rica. Range unknown.

667. *Macrogeomys cherrii* (Allen).

Geomys cherriei Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1893, p. 337.

Macrogeomys cherrii Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 330, fig. 62. Zool. Ser.

CHERRIE'S POCKET GOPHER.

Type locality. Santa Clara, Costa Rica.

Geogr. Distr. Costa Rica. Range unknown.

Genus 93. *Zygogeomys*. *Zygomata* Pocket Gophers.

Zygogeomys Merr., Mon. Geomyidæ, N. Am. Faun., No. 8, 1895, p. 195, pls. and figs. Type *Zygogeomys trichopus* Merriam.

668. *Zygogeomys trichopus* Merriam.

Zygogeomys trichopus Merr., Mon. Geomyidæ, N. Am. Faun., No. 8, 1895, p. 195, pls. and figs. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 332, fig. 63. Zool. Ser.

PINE ZONE POCKET GOPHER.

Type locality. Nahuatzin, State of Michoacan, Mexico.

Geogr. Distr. Sierra Madre, State of Michoacan, Mexico. Altitude, 6,800 to 9,500 feet.

Genus 94. *Thomomys*. Pocket Gophers.

Thomomys Max., Nov. Act. Phys. Med. Akad. Cæs. Leop.-Carol., xix, 1839, p. 377. Type *Thomomys rufescens* Maximilian.

Diplostoma Rich., Faun. Bor. Amer., Mamm., 1, 1829, p. 206. (nec Rafin.)

Oryctomys Eyd. & Gerv., Mag. Zool., vi, 1836, p. 23. (Part.)
Megascapheus Elliot, Pub. Field Columb. Mus., iii, 1903, p. 190.
 Zoöl. Ser.

Subgenus A. *Megascapheus*.

669. *Thomomys bulbivorus* (Richardson).

Diplostoma bulbivorum Rich., Faun. Bor. Amer., i, 1829, p. 206,
 pl. 18, B.

Thomomys bulbivorus Elliot, Syn. N. Am. Mamm., F. C. M. Pub.,
 ii, 1901, p. 222. Zoöl. Ser.

GREAT POCKET GOPHER.

Type locality. Banks of the Columbia River, Oregon.

Geogr. Distr. Shores of the Lower Columbia River, Oregon, to
 coast of California north of San Francisco.

Subgenus B. *Thomomys*.

670. *Thomomys fulvus* (Woodhouse).

Geomys fulvus Woodh., Proc. Acad. Nat. Scien. Phil., 1852, p. 201.

Thomomys fulvus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., ii,
 1901, p. 227. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. In-
 dies, F. C. M. Pub., iv, Pt. 1, 1904, p. 334, fig. XLVII. Zoöl. Ser.

Ascomys umbrinus Wagn., Schreb. Säugth., Suppl., iv, 1843,
 p. 339.

Pseudostoma umbrinus Aud. & Bach., Quad. N. Am., iii, 1854,
 p. 307.

Thomomys umbrinus Baird, N. Am. Mamm., 1857, p. 399.

REDDISH BROWN POCKET GOPHER.

Type locality. San Francisco Mountains, Coconino County, Ari-
 zona.

Geogr. Distr. State of Sonora, Mexico, north into New Mexico,
 and Arizona to central California. Lower California?

671. *Thomomys nevadensis* Merriam.

Thomomys nevadensis Merr., Proc. Biol. Soc. Wash., xi, 1897,
 p. 213. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., ii, 1901,
 p. 223. Zoöl. Ser.

NEVADA POCKET GOPHER.

Type locality. Austin, Lander County, Nevada.

Geogr. Distr. Nevada. Range unknown.

672. *Thomomys uinta* Merriam.

Thomomys uinta Merr., Proc. Biol. Soc. Wash., xiv, 1901, p. 112.

UINTA MOUNTAINS POCKET GOPHER.

Type locality. Uinta Mountains, Summit County, Utah. Altitude, 10,000 feet.

Geogr. Distr. Uinta Mountains. Range unknown.

Genl. Char. Nasals short, faintly notched behind, not so long as premaxillæ; interparietal large, pentagonal; no temporal ridges; auditory tubes conspicuous; bullæ large.

Color. Above grayish brown suffused with fulvous and mixed with black tipped hairs; cheeks and sides of neck grizzled bister; sides grayish brown; nose, ear patch, and throat dusky; fore legs and feet grayish dusky; hind feet whitish; tail above dark.

Measurements. Total length, 220; tail vertebræ, 68; hind foot, 30.

673. *Thomomys leucodon* Merriam.

Thomomys leucodon Merr., Proc. Biol. Soc. Wash., XI, 1897, p. 215.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 223.

Zoöl. Ser.

WHITE-TOOTHED POCKET GOPHER.

Type locality. Grant Pass, Rogue River Valley, Josephine County, Oregon.

Geogr. Distr. Oregon. Range unknown.

a. Thomomys leucodon navus Merriam.

Thomomys leucodon navus Merr., Proc. Biol. Soc. Wash., XIV, 1901, p. 112.

ACTIVE POCKET GOPHER.

Type locality. Red Bluff, Tehama County, California.

Geogr. Distr. Range unknown.

Genl. Char. Similar to *T. leucodon*; size smaller; incisors yellow; nasals cuneate, notched behind.

Color. Above pale fulvous; under parts buffy ochraceous.

Measurements. Total length, 200; tail vertebræ, 67; hind foot, 27.

674. *Thomomys operarius* Merriam.

Thomomys operarius Merr., Proc. Biol. Soc. Wash., XI, 1897, p. 215.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 223.

Zoöl. Ser.

WORKING POCKET GOPHER.

Type locality. Keeler, east side of Owens Lake, Inyo County, California.

Geogr. Distr. Eastern California, apparently restricted to the vicinity of Owens Lake.

675. *Thomomys scapterus* Elliot.

Thomomys scapterus Elliot, Pub. Field Columb. Mus., III, 1903,
p. 248. Zoöl. Ser.

DIGGER POCKET GOPHER.

Type locality. Hannopee Cañon, Panamint Mountains, Inyo County, California.

Geogr. Distr. Panamint, Coso and Inyo Mountains, Inyo County, California.

Genl. Char. Similar to *T. operarius*, but much darker in color, and with much shorter nasals.

Color. Upper parts and sides wood brown, heavily lined with black on top of head and dorsal region, in some specimens nearly forming a dorsal band; lower sides and entire under parts white, the plumbeous under fur showing; hands and feet whitish, tail unicolor, white; ears, and small spot behind ear, black.

Measurements. Total length, 229; tail vertebræ, 74; hind foot, 29; ear, 6. Skull: total length, 37; Hensel, 33; zygomatic width, 23; interorbital constriction, 6; greatest width of braincase, 9; palatal length, 33; length of nasals, 10; anterior width of nasals, 5; length of mandible, 24.

676 *Thomomys bottæ* (Eydoux and Gervais).

Oryctomys bottæ Eyd. and Gerv., Mag. Zool., VI, 1836, p. 23, pl. XXI,
fig. 4. (teeth.)

Thomomys bottæ Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II,
1901, p. 223, fig. 52. Zoöl. Ser.

BOTTA'S POCKET GOPHER.

Type locality. Coast of California, Monterey.

Geogr. Distr. Central and southern California.

a. *Thomomys bottæ pallescens* Rhoads.

Thomomys bottæ pallescens Rhoads, Proc. Acad. Nat. Scien. Phil.,
1895, p. 36. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II,
1901, p. 224. Zoöl. Ser.

PALLID POCKET GOPHER.

Type locality. Grapelands, San Bernardino Valley, California.

Geogr. Distr. Southern California.

677. *Thomomys laticeps* Baird.

Thomomys laticeps Baird, Proc. Acad. Nat. Scien. Phil., 1855,
p. 335. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901,
p. 224. Zoöl. Ser.

BROAD-HEADED POCKET GOPHER.

Type locality. Humboldt Bay, Humboldt County, California.

Geogr. Distr. Northern California.

678. *Thomomys townsendi* (Bachman).

Geomys townsendii Bachm., Journ. Acad. Nat. Scien. Phil., 1839,
p. 105.

Geomys borealis Bachm., Journ. Acad. Nat. Scien. Phil., 1839,
p. 103. (nec Rich.)

Thomomys townsendi Elliot, Syn. N. Am. Mamm., F. C. M. Pub.,
II, 1901, p. 224. Zoöl. Ser.

TOWNSEND'S POCKET GOPHER.

Type locality. "Rocky Mountains"; probably southwestern Wyoming.

Geogr. Distr. Southwestern Wyoming. Range unknown.

679. *Thomomys monticola* Allen.

Thomomys monticolus! Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1893,
p. 48.

Thomomys monticola Elliot, Syn. N. Am. Mamm., F. C. M. Pub.,
II, 1901, p. 225. Zoöl. Ser.

MOUNTAIN POCKET GOPHER.

Type locality. Mt. Tallac, El Dorado County, California. Altitude, 7,500 feet.

Geogr. Distr. Northern California from San Francisco Bay to northwestern Nevada.

a. *Thomomys monticola pinetorum* Merriam.

Thomomys monticolus! pinetorum Merr., N. Am. Faun., No. 16,
1899, p. 97.

Thomomys monticola pinetorum Elliot, Syn. N. Am. Mamm.,
F. C. M. Pub., II, 1901, p. 225. Zoöl. Ser.

PINE POCKET GOPHER.

Type locality. Sisson, Siskiyou County, California.

Geogr. Distr. Northern California. Range unknown.

680. *Thomomys mazama* Merriam.

Thomomys mazama Merr., Proc. Biol. Soc. Wash., XI, 1897, p. 214.
Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 225.
Zoöl. Ser.

MT. MAZAMA POCKET GOPHER.

Type locality. Crater Lake, Mount Mazama, Klamath County, Oregon. Altitude, 6,000 feet.

Geogr. Distr. Mt. Mazama. Range unknown.

681. *Thomomys quadratus* Merriam.

Thomomys quadratus Merr., Proc. Biol. Soc. Wash., XI, 1897, p. 214. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 225. Zool. Ser.

THE DALLES POCKET GOPHER.

Type locality. The Dalles, Wasco County, Oregon.

Geogr. Distr. Oregon. Plains of the Columbia.

682. *Thomomys myops* Merriam.

Thomomys myops Merr., Proc. Biol. Soc. Wash., XIV, 1901, p. 112. NEAR-SIGHTED POCKET GOPHER.

Type locality. Conconully, east base of Cascade Range, Okanogan County, Washington.

Geogr. Distr. Washington. Range unknown.

Genl. Char. Size small. Skull similar to that of *T. quadratus*, but with comparative differences, such as a less quadrate zygomata with shorter posterior root; broader nasals, equal in length to pre-maxillæ and truncated posteriorly.

Color. Upper parts dull rufous brown; chin, throat, hands, and feet white; under parts buffy ochraceous, base of fur dark slate; sides of mouth and orbital ring plumbeous.

Measurements. Total length, 197; tail vertebræ, 63; hind foot, 26.

683. *Thomomys nasicus* Merriam.

Thomomys nasicus Merr., Proc. Biol. Soc. Wash., XI, 1897, p. 216.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 226.

Zool. Ser.

LARGE-NOSED POCKET GOPHER.

Type locality. Farewell Bend, Des Chutes River, Crook County, Oregon.

Geogr. Distr. Oregon. Range unknown.

684. *Thomomys niger* Merriam.

Thomomys niger Merr., Proc. Biol. Soc. Wash., XIV, 1901, p. 117. BLACK POCKET GOPHER.

Type locality. Seaton, near mouth of Umpqua River, Douglas County, Oregon.

Geogr. Distr. Known only from the type locality.

Genl. Char. Size medium; feet large. Skull with temporal ridges; interparietal oval; nasals emarginate, narrowed posteriorly.

Color. Slaty black with greenish reflections; feet and tail white, sometimes blotched with dusky. Probably melanistic individuals of *T. douglasi*.

Measurements. Total length, 225; tail vertebræ, 81; hind foot, 30.

685. *Thomomys douglasi* (Richardson).

Geomys douglasii Rich., Faun. Bor. Amer., 1, 1829, p. 200.

Thomomys douglasi Elliot, Syn. N. Am. Mamm., F. C. M. Pub.,
11, 1901, p. 226. Zoöl. Ser.

DOUGLAS' POCKET GOPHER.

Type locality. Near mouth of the Columbia River, Oregon.

Geogr. Distr. Lower Columbia River, northward into British Columbia.

a. *Thomomys douglasi yelmensis* Merriam.

Thomomys douglasi yelmensis Merr., Proc. Biol. Soc. Wash., XIII,
1899, p. 21. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11,
1901, p. 226. Zoöl. Ser.

YELM PRAIRIE POCKET GOPHER.

Type locality. Tenino, Yelm Prairie, Thurston County, Washington.

Geogr. Distr. Washington. Range unknown.

b. *Thomomys douglasi fuscus* (Merriam).

Thomomys clusius fuscus Merr., N. Am. Faun., No. 5, 1891, p. 70.

Thomomys douglasi fuscus Elliot, Syn. N. Am. Mamm., F. C. M.
Pub., 11, 1901, p. 226. Zoöl. Ser.

SWARTHY POCKET GOPHER.

Type locality. Mountains at head of Big Lost River, Custer County, Idaho.

Geogr. Distr. Mountains of central Idaho.

c. *Thomomys douglasi fisheri* (Merriam).

Thomomys fuscus fisheri Merr., Proc. Biol. Soc. Wash., XIV, 1901,
p. 111.

FISHER'S POCKET GOPHER.

Type locality. Beckwith, Sierra Valley, Plumas County, California.

Geogr. Distr. California. Range unknown.

Genl. Char. Similar to *T. d. fuscus*, but paler; comparative differences in skull.

Color. Upper parts and beneath grayish brown.

Measurements. Total length, 191; tail vertebræ, 62; hind foot, 25.

d. *Thomomys douglasi oregonus* Merriam.

Thomomys douglasi oregonus Merr., Proc. Biol. Soc. Wash., XIV,
1901, p. 115.

OREGON POCKET GOPHER.

Type locality. Oregon City, Willamette Valley, Clackamas County, Oregon.

Geogr. Distr. Oregon. Range unknown.

Genl. Char. Like *T. douglasi*. Skull with broader zygomata, more bowed outwardly, the outer sides parallel; interparietal larger, subtriangular, with straight portions emerging; pterygoid notch V-shaped.

Color. Exactly like that of *T. douglasi*.

Measurements. Total length, 220; tail vertebrae, 70; hind foot, 30.

686. *Thomomys helleri* Elliot.

Thomomys helleri Elliot, Pub. Field Columb. Mus., III, 1903, p. 165. Zool. Ser.

HELLER'S GOPHER.

Type locality. Goldbeach, Oregon.

Geogr. Distr. Oregon. Range undetermined.

Genl. Char. Nearest to *T. douglasi*; color different; tail and hind foot longer. Skull shorter; molars small; rostrum broad.

Color. Head and upper parts reddish brown; spot behind ears, line beneath eyes, nose, lips, and margin of pouches, black; cheeks, sides, throat, and under parts buff; base of fur plumbeous; feet yellowish white; tail above dark gray, beneath white.

Measurements. Total length, 209; tail vertebrae, 67; hind foot, 28; ear, 8. Skull: occipito-nasal length, 36; Hensel, 28; zygomatic width, 19; interorbital constriction, 6; greatest width of braincase, 14; mastoid breadth, 13.5; length of nasals, 11; width of rostrum, 6; palatal length, 5; length of mandible, 19; height at condyle, 7; at coronoid process, 11; length of upper tooth row, 7.

687. *Thomomys limosus* Merriam.

Thomomys limosus Merr., Proc. Biol. Soc. Wash., XIV, 1901, p. 116. MUD-LOVING POCKET GOPHER.

Type locality. White Salmon, Gorge of the Columbia, Klickitat County, Washington.

Geogr. Distr. Washington. Range unknown.

Genl. Char. Similar to *T. douglasi*, but darker. Skull with zygomata bowed outward, nasals restricted behind anterior third, expanded and emarginate posteriorly; interpterygoid space V-shaped; angular process of mandible spreading.

Measurements. Total length, 224; tail vertebrae, 68; hind foot, 29.

688. *Thomomys talpoides* (Richardson).

Cricetus talpoides Rich., Zool. Journ., III, 1828, p. 518.

Geomys borealis Rich., Rep. Brit. Ass. for 1836, VI, 1837, p. 156.

Thomomys talpoides Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 227. Zool. Ser.

RICHARDSON'S POCKET GOPHER.

Type locality. Probably Saskatchewan, Canada.

Geogr. Distr. Saskatchewan region south to Missouri region.

a. *Thomomys talpoides rufescens* (Maximilian).

Thomomys rufescens Max., Nov. Act. Phys. Med. Acad. Cæs. Leop.-Carol., XIX, Pt. I, 1839, p. 378.

Thomomys talpoides rufescens Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 227. Zoöl. Ser.

RUFESCENT POCKET GOPHER.

Type locality. Missouri River; exact locality unknown.

Geogr. Distr. Upper Missouri and Valley of the Saskatchewan, Canada.

689. *Thomomys bridgeri* Merriam.

Thomomys bridgeri Merr., Proc. Biol. Soc. Wash., XIV, 1901, p. 113.

BRIDGER'S POCKET GOPHER.

Type locality. Fort Bridger, Uinta County, Wyoming.

Geogr. Distr. Wyoming. Range unknown.

Genl. Char. Size large. Skull with parallel temporal ridges; rostrum and nasals long, notched posteriorly, otherwise similar to those of *T. talpoides*.

Color. Above chestnut to buffy brown mixed with black; beneath washed with buffy fulvous; chin, ear patch, and entrance to pouches dusky; feet whitish.

Measurements. Total length, 237; tail vertebræ, 71; hind foot, 34.

690. *Thomomys melanops* Merriam.

Thomomys melanops Merr., Proc. Biol. Soc. Wash., XIII, 1899, p. 21. Elliot, Syn. N. Am. Mamm., II, 1901, p. 227. Zoöl. Ser.

BLACK-FACED POCKET GOPHER.

Type locality. Solduck Trail, Olympic Mountains, Clallam County, Washington.

Geogr. Distr. Olympic Mountains, Washington. High elevation.

691. *Thomomys hesperus* Merriam.

Thomomys hesperus Merr., Proc. Biol. Soc. Wash., XIV, 1901, p. 116.

WESTERN POCKET GOPHER.

Type locality. Tillamook, Tillamook County, Oregon.

Geogr. Distr. Oregon. Range unknown.

Genl. Char. Size small. Interparietal large, triangular; bullæ small.

Color. Above deep rufous; beneath buffy to salmon fulvous; nose, ear patch, and orbital ring dusky; tail above dark, beneath whitish; fore and hind feet whitish.

Measurements. Total length, 175; tail vertebræ, 54; hind foot, 24.

692. *Thomomys fulvus* (Woodhouse).

Geomys fulvus Woodh., Proc. Acad. Nat. Scien. Phil., 1852, p. 201.

Geomys umbrinus Rich., Faun. Bor. Amer., I, 1829, p. 202. (Undeterminable.)

Thomomys umbrinus Baird, N. Am. Mamm., 1857, p. 399.

Thomomys fulvus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 227. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 334, fig. XLVII. Zoöl. Ser.

FULVOUS POCKET GOPHER.

Type locality. San Francisco Mountain, Coconino County, Arizona.

Geogr. Distr. Central California south into Arizona, New Mexico, and State of Chihuahua, Mexico. Lower California, Mexico?

a. *Thomomys fulvus anitæ* Allen.

Thomomys fulvus anitæ Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1898, p. 146. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 335. Zoöl. Ser.

SANTA ANITA POCKET GOPHER.

Type locality. Santa Anita, Lower California, Mexico.

Geogr. Distr. Cape region of Lower California, Mexico.

b. *Thomomys fulvus alticola* Allen.

Thomomys fulvus alticola Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1899, p. 13. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 335. Zoöl. Ser.

SIERRA LAGUNA POCKET GOPHER.

Type locality. Sierra Laguna, Lower California, Mexico. Altitude, 7,000 feet.

Geogr. Distr. High altitudes in the Sierra Laguna range, Lower California, Mexico.

c. *Thomomys fulvus nigricans* Rhoads.

Thomomys fulvus nigricans Rhoads, Proc. Acad. Nat. Scien. Phil., 1895, p. 36. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 228. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 335. Zoöl. Ser.

BLACKISH POCKET GOPHER.

Type locality. Witch Creek, San Diego County, California.

Geogr. Distr. Southern California, south into Lower California, Mexico.

d. *Thomomys fulvus intermedius* Mearns.

Thomomys fulvus intermedius Mearns, Proc. U. S. Nat. Mus., XIX, 1897, p. 719. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II,

1901, p. 228. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 336. Zoöl. Ser.

INTERMEDIATE POCKET GOPHER.

Type locality. Summit of Huachuca Mountains, southern Arizona. Altitude, 9,000 feet.

Geogr. Distr. Mountains connecting Colorado Plateau with that of Mexico.

e. Thomomys fulvus texensis Bailey.

Thomomys fulvus texensis Bailey, Proc. Biol. Soc. Wash., xv, 1902, p. 119.

TEXAS POCKET GOPHER.

Type locality. Limpia Creek, Davis Mountain, Jeff Davis County, Texas. Altitude, 5,500 feet.

Geogr. Distr. Texas. Range unknown.

Genl. Char. Like *T. fulvus*, but smaller, less dusky. Skull narrower, pterygoids shorter; lateral palatal pits shallower.

Color. Above dusky fulvous; ear patch and nose blackish; beneath pale bright fulvous; feet and lips gray; pouch linings white.

Measurements. Total length, 204; tail vertebrae, 63; hind foot, 26. Skull: basal length, 34.5; length of nasals, 13; zygomatic breadth 22; mastoid breadth, 18; alveolar length of upper molar series, 7.

693. *Thomomys desertorum* Merriam.

Thomomys desertorum Merr., Proc. Biol. Soc. Wash., xiv, 1901, p. 114.

DESERT POCKET GOPHER.

Type locality. Mud Spring, Detrital Valley, Mohave County, Arizona.

Geogr. Distr. Arizona. Range unknown.

Genl. Char. Size small; zygomata much bowed; interparietal subquadrate; nasals not as long as premaxillæ, notched posteriorly; bullæ large.

Color. Above buffy to orange fulvous; dark nose patch reaching to eyes or ears; beneath buffy to salmon fulvous; ear patch dusky; feet whitish; tail like back.

Measurements. None given.

694. *Thomomys angularis* Merriam.

Thomomys angularis Merr., Proc. Biol. Soc. Wash., xi, 1897, p. 214. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 228. Zoöl. Ser.

ANGULAR POCKET GOPHER.

Type locality. Los Baños, Merced County, California.

Geogr. Distr. Western California.

a. Thomomys angularis pascalis Merriam.

Thomomys angularis pascalis Merr., Proc. Biol. Soc. Wash., xiv
1901, p. 111.

GRAZING POCKET GOPHER.

Type locality. Fresno, San Joaquin Valley, Fresno County, California.

Geogr. Distr. Joaquin Valley, California. Range unknown.

Genl. Char. Similar to *T. angularis*, but smaller. Skull: temporal ridges not forming a sagittal crest; bullæ large.

Color. Upper parts buffy yellowish; under parts paler, sometimes with patches of white; wrists and ankles white.

Measurements. Total length, 236; tail vertebræ, 80; hind foot, 32.

695. *Thomomys martirensis* (Allen).

Thomomys fulvus martirensis Allen, Bull. Am. Mus. Nat. Hist.
N. Y., 1898, p. 147.

Thomomys martirensis Elliot, Mamm. Middle Amer. & W. Indies,
F. C. M. Pub., iv, Pt. 1, 1904, p. 336. Zoöl. Ser.

SAN PEDRO MARTIR POCKET GOPHER.

Type locality. San Pedro Martir Mountains, Lower California, Mexico. Altitude, 8,200 feet.

Geogr. Distr. San Pedro Martir Mountains, Lower California, Mexico. Altitude, 7,000 to 8,200 feet.

696. *Thomomys aphrastus* Elliot.

Thomomys aphrastus Elliot, Pub. Field Columb. Mus., III, 1903,
p. 219. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies,
F. C. M. Pub., iv, Pt. 1, 1904, p. 336. Zoöl. Ser.

TROUBLESOME POCKET GOPHER.

Type locality. San Tomas, Lower California, Mexico.

Geogr. Distr. Lower California, Mexico. Range unknown.

697. *Thomomys peregrinus* Merriam.

Thomomys peregrinus Merr., Proc. Biol. Soc. Wash., VIII, 1893,
p. 146. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M.
Pub., iv, Pt. 1, 1904, p. 337. Zoöl. Ser.

WANDERING POCKET GOPHER.

Type locality. Salazar, State of Mexico, Mexico.

Geogr. Distr. State of Mexico. Range unknown.

698. *Thomomys orizabæ* Merriam.

Thomomys orizabæ Merr., Proc. Biol. Soc. Wash., VIII, 1893, p. 145.
Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv,
Pt. 1, 1904, p. 337. Zoöl. Ser.

ORIZABA POCKET GOPHER.

Type locality. Mt. Orizaba, State of Puebla, Mexico. Altitude, 9,500 feet.

Geogr. Distr. State of Puebla, Mexico. Range unknown.

699. *Thomomys atrovarius* Allen.

Thomomys atrovarius Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1898, p. 148. Elliot, Mamm. Middle Amer. & W. Indies F. C. M. Pub., IV, Pt. I, 1904, p. 338, fig. 64 Zoöl. Ser.

TATEMALES POCKET GOPHER.

Type locality. Tatemaes, State of Sinaloa, Mexico.

Geogr. Distr. State of Sinaloa, Mexico. Range unknown.

700. *Thomomys toltecus* Allen.

Thomomys toltecus Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1893, p. 52, pl. I, fig. 13. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 338. Zoöl. Ser.

JUAREZ POCKET GOPHER.

Type locality. Juarez, northern part of State of Chihuahua, Mexico.

Geogr. Distr. State of Chihuahua, Mexico. Range unknown.

701. *Thomomys perditus* Merriam.

Thomomys perditus Merr., Proc. Biol. Soc. Wash., XIV, 1901, p. 108. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 338. Zoöl. Ser.

LOST POCKET GOPHER.

Type locality. Lampazos, State of Nuevo León, Mexico.

Geogr. Distr. Range unknown.

702. *Thomomys goldmani* Merriam.

Thomomys goldmani Merr., Proc. Biol. Soc. Wash., XIV, 1901, p. 108. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 339. Zoöl. Ser.

GOLDMAN'S POCKET GOPHER.

Type locality. Mapimi, State of Durango, Mexico.

Geogr. Distr. Range unknown.

703. *Thomomys alpinus* Merriam.

Thomomys alpinus Merr., Proc. Biol. Soc. Wash., XI, 1897, p. 216. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 229. Zoöl. Ser.

ALPINE POCKET GOPHER.

Type locality. Big Cottonwood Meadows, Mount Whitney, eight miles southeast of the Peak, Tulare County, California. Altitude, 10,000 feet.

Geogr. Distr. High Sierra, eastern California.

704. *Thomomys clusius* Coues.

Thomomys clusius Coues, Proc. Acad. Nat. Scien. Phil., 1875,
p. 138. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901,
p. 229. Zoöl. Ser.

COUES' POCKET GOPHER.

Type locality. Bridger Pass, Sweetwater County, Wyoming.

Geogr. Distr. Uncompahgre Reservation, Utah, through southwestern Wyoming into southern and central Idaho.

a. *Thomomys clusius ocus* Merriam.

Thomomys clusius ocus Merr., Proc. Biol. Soc. Wash., XIV, 1901,
p. 114.

SWIFT POCKET GOPHER.

Type locality. Fort Bridger, Uinta County, Wyoming.

Geogr. Distr. Wyoming. Range unknown.

Genl. Char. Similar to *T. clusius*; interparietal and bullæ larger.

Color. Above pale buffy; cheeks pale buffy gray; sides buffy white; under parts and feet white.

Measurements. Total length, 204; tail vertebræ, 197; hind foot, 25.

705. *Thomomys idahoensis* Merriam.

Thomomys idahoensis Merr., Proc. Biol. Soc. Wash., XIV, 1901, p. 114.

IDAHO POCKET GOPHER.

Type locality. Birch Creek, Fremont County, Idaho.

Geogr. Distr. Idaho. Range unknown.

Genl. Char. Size small; similar to *T. clusius*. Skull: bullæ large; zygomata narrow; nasals long, slender.

Color. Above grayish buff washed with buffy fulvous; under parts, tail, and feet buffy white.

Measurements. Total length, 179; tail vertebræ, 47; hind foot, 23.

706. *Thomomys pygmæus* Merriam.

Thomomys pygmæus Merr., Proc. Biol. Soc. Wash., XIV, 1901,
p. 115.

PYGMY POCKET GOPHER.

Type locality. Montpelier Creek, Bear County, Idaho. Altitude, 6,700 feet.

Geogr. Distr. Idaho. Range unknown.

Genl. Char. Size small. Skull similar to *T. idahoensis*, smaller; interparietal transversely oval; nasals short, emarginate posteriorly; bullæ small.

Color. Above dark rufous brown; beneath buffy fulvous; feet whitish.

Measurements. Total length, 177; tail vertebræ, 46; hind foot, 22.

707. *Thomomys aureus* Allen.

Thomomys aureus Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1893,
p. 49. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901,
p. 229. Zoöl. Ser.

GOLDEN POCKET GOPHER.

Type locality. Bluff City, San Juan County, Utah.
Geogr. Distr. Southeastern Utah.

a. *Thomomys aureus lachuguilla* Bailey.

Thomomys aureus lachuguilla Bailey, Proc. Biol. Soc. Wash., xv,
1902, p. 120.

BAILEY'S POCKET GOPHER.

Type locality. Foothills near El Paso, Texas.
Geogr. Distr. Texas. Range unknown.
Genl. Char. Size small; bullæ large; basi-occipital narrow; skull slender.

Color. Summer Pelage. Above dull ochraceous; beneath buffy or pale cinnamon.

Winter Pelage. Brighter ochraceous above; light buffy beneath.

Measurements. Total length, 215; tail vertebræ, 65; hind foot, 29.

b. *Thomomys aureus pervagus* Merriam.

Thomomys aureus pervagus Merr., Proc. Biol. Soc. Wash., xiv,
1901, p. 110.

ROAMING POCKET GOPHER.

Type locality. Espanola, Santa Fé County, New Mexico.

Geogr. Distr. Range unknown.

Genl. Char. Nasals broader than those of *T. aureus*.

Color. Above chestnut fulvous lined with black on dorsal region; nose and mouth dusky; chin white; under parts salmon.

Measurements. Total length, 244; tail vertebræ, 73; hind foot, 32.

c. *Thomomys aureus perpes* Merriam.

Thomomys aureus perpes Merr., Proc. Biol. Soc. Wash., xiv, 1901,
p. 111.

LONE PINE POCKET GOPHER.

Type locality. Lone Pine, Owens Valley, Inyo County, California.

Geogr. Distr. East slope of the Sierras in Owens Valley, Inyo County, California.

Genl. Char. Size small; skull generally like that of *T. aureus*, but smaller.

Color. Winter Pelage. Above buffy gray, darkest on head; sides buffy; under parts, feet, and tail whitish.

Summer Pelage. Like that of *T. aureus*.

Measurements. Total length, 215; tail vertebræ, 65; hind foot, 28.

708. *Thomomys baileyi* Merriam.

Thomomys baileyi Merr., Proc. Biol. Soc. Wash., XIV, 1901, p. 109.
SIERRA BLANCA POCKET GOPHER.

Type locality. Sierra Blanca, El Paso County, Texas.

Genl. Char. Size small; upper incisors projecting forwards. Skull resembles that of *T. toltecus* but differs in the projecting incisors, and longer nasals, wider interorbital region, and more open post-coronoid notch of mandible.

Color. Above buffy fulvous lined with black; beneath buff; mouth dusky; pouches and feet white.

Measurements. Total length, 212; tail vertebræ, 168; hind foot, 30.

709. *Thomomys nelsoni* Merriam.

Thomomys nelsoni Merr., Proc. Biol. Soc. Wash., XIV, 1901, p. 109.

Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 339. Zoöl. Ser.

NELSON'S POCKET GOPHER.

Type locality. Parral, State of Chihuahua, Mexico.

Geogr. Distr. State of Chihuahua, Mexico. Range unknown.

710. *Thomomys latirostris* Merriam.

Thomomys latirostris Merr., Proc. Biol. Soc. Wash., XIV, 1901, p. 107.
BROAD-NOSED POCKET GOPHER.

Type locality. Little Colorado River, Painted Desert, Coconino County, Arizona.

Geogr. Distr. Arizona. Range unknown.

Genl. Char. Size medium; rostrum broad; nasals shorter than the premaxillæ; interparietal pentagonal.

Color. Above pale ochraceous buff; under parts, feet, and tail whitish.

Measurements. Total length, 232; tail vertebræ, 79; hind foot, 33.

711. *Thomomys perpallidus* (Merriam).

Thomomys talpoides perpallidus Merr., Science, VIII, 1886, p. 588.

Thomomys perpallidus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 229. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 340. Zoöl. Ser.

PALE POCKET GOPHER.

Type locality. Colorado Desert. San Diego County, California.

Geogr. Distr. Lower California, Mexico, Colorado Desert, southern California, and northeastward to the Painted Desert, Arizona.

712. *Thomomys *cabezona* Merriam.

Thomomys cabezonæ Merr., Proc. Biol. Soc. Wash., XIV, 1901, p. 110.

*Mr. Heller in his letters spells the name of the place from which this species came, Cabazon.

CABEZON POCKET GOPHER.

Type locality. Cabezon, San Gorgonio Pass, Riverside County, California.

Geogr. Distr. Known only from the type locality.

Genl. Char. Smaller than *T. perpallidus*; tail long, ears large; interparietal quadrangular; bullæ small.

Color. Above and on sides buffy ochraceous or drab brown; nose, lips, chin, and pouches dusky; under parts white or pale salmon.

Measurements. Total length, 235; tail vertebræ, 79; hind foot, 30.

713. *Thomomys cervinus* Allen.

Thomomys cervinus Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1895, p. 203, fig. 1. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 230. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 340. Zoöl. Ser.

CERVINE POCKET GOPHER.

Type locality. Phoenix, Maricopa County, Arizona.

Geogr. Distr. Southern Arizona into State of Sonora, Mexico.

714. *Thomomys sinaloæ* Merriam.

Thomomys sinaloæ Merr., Proc. Biol. Soc. Wash., XIV, 1901, p. 108. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 339. Zoöl. Ser.

SINALOA POCKET GOPHER.

Type locality. Altata, State of Sinaloa, Mexico.

Geogr. Distr. Known only from the type locality.

715. *Thomomys altivallis* Rhoads.

Thomomys altivallis Rhoads, Proc. Acad. Nat. Scien. Phil., 1895, p. 34. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 230. Zoöl. Ser.

RHOADS' POCKET GOPHER.

Type locality. San Bernardino Mountains, California. Altitude, 5,000 feet.

Geogr. Distr. Mountains of southeastern California.

716. *Thomomys fossor* Allen.

Thomomys fossor Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1893, p. 51. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 230. Zoöl. Ser.

DELVING POCKET GOPHER.

Type locality. Florida, La Plata County, Colorado. Altitude, 7,200 feet.

Geogr. Distr. Upper San Juan region. Colorado.

Family VI. **Heteromyidae. Kangaroo Rats. Pocket Mice.**Subfamily I. **Dipodomysinae.**Genus 95. **Dipodomys. Four-toed Kangaroo Rats.****Dipodomys** Gray, Ann. Mag. Nat. Hist., 1st Ser., VII, 1841, p. 521.Type *Dipodomys phillipsi* Gray.*Macrocolus* Wagn., Schreb. Säugth., Suppl., IV, 1844 (no text), pl. CCXXXIX. (Fig. of teeth.)**717. *Dipodomys merriami* Mearns.***Dipodomys merriami* Mearns, Bull. Am. Mus. Nat. Hist. N. Y., 1890, p. 290. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 232, fig. 53. Zool. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 345, fig. XLVIII. Zool. Ser.**MERRIAM'S KANGAROO RAT.***Type locality.* New River, between Phoenix and Prescott, Arizona.*Geogr. Distr.* Arizona south into States of Sonora and Chihuahua, Mexico.*a. *Dipodomys merriami nevadensis* Merriam.**Dipodomys merriami nevadensis* Merr., Proc. Biol. Soc. Wash., IX, 1894, p. 111. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 232. Zool. Ser.**NEVADA KANGAROO RAT.***Type locality.* Pyramid Lake, Washoe County, Nevada.*Geogr. Distr.* Northwestern Nevada. Range unknown.*b. *Dipodomys merriami nitratus* Merriam.**Dipodomys merriami nitratus* Merr., Proc. Biol. Soc. Wash., IX, 1894, p. 112. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 233. Zool. Ser.**KEELER KANGAROO RAT.***Type locality.* Keeler, Owens Lake, Inyo County, California.*Geogr. Distr.* Type locality only.*c. *Dipodomys merriami nitratoides* Merriam.**Dipodomys merriami nitratoides* Merr., Proc. Biol. Soc. Wash., IX, 1894, p. 112. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 233. Zool. Ser.**SIMILAR KANGAROO RAT.***Type locality.* Tipton, San Joaquin Valley, Tulare County, California.*Geogr. Distr.* South central California.

d. *Dipodomys merriami exilis* Merriam.

Dipodomys merriami exilis Merr., Proc. Biol. Soc. Wash., ix, 1894, p. 113. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 233. Zoöl. Ser.

SLENDER KANGAROO RAT.

Type locality. Fresno, San Joaquin Valley, Fresno County, California.

Geogr. Distr. South central California.

e. *Dipodomys merriami ambiguus* (Merriam).

Dipodomys ambiguus Merr., N. Am. Faun., No. 4, 1890, p. 42.

Dipodomys merriami ambiguus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 234. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 346. Zoöl. Ser.

DOUBTFUL KANGAROO RAT.

Type locality. El Paso, El Paso County, Texas.

Geogr. Distr. Northwestern Texas, south to State of Chihuahua, Mexico.

f. *Dipodomys merriami parvus* (Rhoads).

Dipodomys parvus Rhoads, Amer. Nat., xxviii, 1894, p. 70.

Dipodomys merriami parvus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 234. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 346. Zoöl. Ser.

SMALL KANGAROO RAT.

Type locality. San Bernardino, San Bernardino County, California.

Geogr. Distr. Southern California south into Lower California, Mexico.

g. *Dipodomys merriami simiolus* (Rhoads).

Dipodomys simiolus Rhoads, Proc. Acad. Nat. Scien. Phil., 1894, p. 410.

Dipodomys similis Rhoads, Proc. Acad. Nat. Scien. Phil., 1894, p. 411.

Dipodomys merriami simiolus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 234. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 347. Zoöl. Ser.

Dipodomys merriami similis Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 234. Zoöl. Ser.

ALLIED KANGAROO RAT.

Type locality. Agua Caliente, Mohave Desert, California.

Geogr. Distr. Southern California into Lower California and State of Sonora, Mexico.

h. Dipodomys merriami mortivallis Elliot.

Dipodomys merriami mortivallis Elliot, Pub. Field Columb. Mus.,
III, 1903, p. 250. Zoöl. Ser.

DEATH VALLEY KANGAROO RAT.

Type locality. Furnace Creek, Death Valley, Inyo County, California.

Geogr. Distr. Death Valley and near-by region on the west.

Genl. Char. Similar to *D. m. simiolus*, but the dorsal and ventral stripe and pencil of the tail vary from a purplish tint to pale russet, quite different from the blackish tail of *D. m. simiolus*. The general color of the upper part of the body is also darker. The skulls of the two races are much alike save that of the present one has much longer and broader nasals, widening anteriorly; extreme width of parietal is greater and the mastoids are broader.

Color. Above russet, darkest on the rump, plumbeous under fur showing; sides dark russet; spots behind ear, superciliary stripe, face in front of eye, nose, entire under parts, stripes across thighs and feet, pure white; inner side of thighs deep russet like rump; black bar across rump; tail bushy on apical third, with dorsal and ventral stripes and also bushy portion or pencil pale russet.

Measurements. Total length, 240; tail vertebrae, 142; hind foot, 37.5; ear, 14. Skull: total length, 36; Hensel, 22; greatest width of mastoids, 22; interorbital constriction, 13; width of interparietal, 17; length of nasals, 13; width of nasals posteriorly, 2; anteriorly, 3; length of upper tooth row, 3.5; height of coronoid process from angle, 6.

i. Dipodomys merriami arenivagus Elliot.

Dipodomys merriami arenivagus Elliot, Pub. Field Columb. Mus.,
III, 1903, p. 249. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W.

Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 347. Zoöl. Ser.

SAN FELIPE KANGAROO RAT.

Type locality. San Felipe, Lower California, Mexico.

Geogr. Distr. Northeastern coast of Lower California, Mexico.

j. Dipodomys merriami atronasus Merriam.

Dipodomys merriami atronasus Merr., Proc. Biol. Soc. Wash., IX,
1894, p. 113. Elliot, Mamm. Middle Amer. & W. Indies,
F. C. M. Pub., IV, Pt. 1, 1904, p. 348. Zoöl. Ser.

BLACK-NOSED KANGAROO RAT.

Type locality. Hacienda La Parada, twenty-five miles northwest of the City of San Luis Potosi, State of San Luis Potosi, Mexico.

Geogr. Distr. Known only from the type locality.

k. Dipodomys merriami melanurus Merriam.

Dipodomys merriami melanurus Merr., Proc. Calif. Acad. Scien., 2d Ser., 1893, p. 345. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 348. Zoöl. Ser.

BLACK-TAILED KANGAROO RAT.

Type locality. San José del Cabo, Lower California, Mexico.

Geogr. Distr. Cape region of Lower California, Mexico.

718. *Dipodomys mitchelli* Mearns.

Dipodomys mitchelli Mearns, Proc. U. S. Nat. Mus., XIX, 1897, p. 719. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 344. Zoöl. Ser.

TIBURON ISLAND KANGAROO RAT.

Type locality. Tiburon Island, Gulf of California, Mexico.

Geogr. Distr. Known only from the type locality.

719. *Dipodomys phillipsi* Gray.

Dipodomys phillipsi Gray, Ann. Mag. Nat. Hist., VII, 1st Ser., 1841, p. 522. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 343, fig. 65. Zoöl. Ser.

PHILLIPS' KANGAROO RAT.

Type locality. Real del Monte, north end of the Valley of Mexico.

Geogr. Distr. Valley of Mexico, State of Mexico, and States of Puebla, Tlaxcala, and Vera Cruz, Mexico.

720. *Dipodomys ornatus* Merriam.

Dipodomys ornatus Merr., Proc. Biol. Soc. Wash., IX, 1894, p. 110. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 344. Zoöl. Ser.

ORNAMENTED KANGAROO RAT.

Type locality. Berriozabal, State of Zacatecas, Mexico.

Geogr. Distr. State of Zacatecas, Mexico.

721. *Dipodomys perotensis* Merriam.

Dipodomys perotensis Merr., Proc. Biol. Soc. Wash., IX, 1894, p. 111. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 344. Zoöl. Ser.

PEROTE KANGAROO RAT.

Type locality. Perote, State of Vera Cruz, Mexico.

Geogr. Distr. State of Vera Cruz.

722. *Dipodomys californicus* Merriam.

Dipodomys californicus Merr., N. Am. Faun., No. 4, 1890, p. 49. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 236. Zoöl. Ser.

CALIFORNIA KANGAROO RAT.

Type locality. Ukiah, Mendocino County, California.

Geogr. Distr. Northwestern California.

a. *Dipodomys californicus pallidulus* Bangs.

Dipodomys californicus pallidulus Bangs, Proc. New Engl. Zool. Club, I, 1899, p. 65. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, Append., p. 431. Zool. Ser.

PALER KANGAROO RAT.

Type locality. Sites, Colusa County, California.

Geogr. Distr. Known only from the type locality.

723. *Dipodomys elator* Merriam.

Dipodomys elator Merr., Proc. Biol. Soc. Wash., IX, 1894, p. 109.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 235. Zool. Ser.

PROUD KANGAROO RAT.

Type locality. Henrietta, Clay County, Texas.

Geogr. Distr. Northern Texas. Range unknown.

724. *Dipodomys deserti* Stephens.

Dipodomys deserti Stephens, Amer. Nat., XXI, 1887, p. 42. Elliot,

Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 235. Zool.

Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 345. Zool. Ser.

DESERT KANGAROO RAT.

Type locality. Mohave River, San Bernardino County, California.

Geogr. Distr. Mohave and Colorado deserts, southeastern California, south to the State of Sonora and Lower California, Mexico.

a. *Dipodomys deserti helleri* Elliot.

Dipodomys deserti helleri Elliot, Pub. Field Columb. Mus., III, 1903, p. 249. Zool. Ser.

HELLER'S KANGAROO RAT.

Type locality. Keeler, Owens Lake, Inyo County, California.

Geogr. Distr. Known only from the type locality.

Genl. Char. Size similar to that of *D. deserti*; color pale ochraceous instead of pale yellowish brown; upper part of tail like the back, with no black markings; no black on face.

Color. Upper parts pale ochraceous or dark pinkish buff, a similar tint to that of *D. m. nitratus*; line over eyes, sides of nose, fore part of face, entire under parts, limbs and feet, pure white; tail with pale ochraceous line above, becoming ochraceous near the tip; sides and under part of lip pure white; ear same color as the back.

Measurements. Total length, 333; tail vertebrae, 195; hind foot, 53.5; ear, 16. Skull: total length, 43; Hensel, 37; zygomatic width, 21; width of mastoids, 29; greatest width of parietal, 20; length of nasals, 14; palatal length, 14; length of upper tooth row, 5; length of mandible, 21; length of lower tooth row, 5.

725. *Dipodomys spectabilis* Merriam.

Dipodomys spectabilis Merr., N. Am. Faun., No. 4, 1890, p. 46.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 235.

Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M.

Pub., IV, Pt. 1, 1904, p. 345. Zoöl. Ser.

NOTABLE KANGAROO RAT.

Type locality. Dos Cabezos, Cochise County, Arizona.

Geogr. Distr. Portions of Arizona and Texas, eastward to Sierra Blanca, Texas, and south into the States of Chihuahua and Sonora, Mexico.

Genus 96. *Perodipus*. Five-toed Kangaroo Rats.

***Perodipus* Fitzin., Sitz. Math.-Natur. Classe K. Akad. Wiss. Wien,**

LVI, 1867, p. 126. *Type* *Dipodomys agilis* Gambel.

Dipodops Merr., N. Am. Faun., No. 3, 1890, p. 72.

726. *Perodipus agilis* (Gambel).

Dipodomys agilis Gambel, Proc. Acad. Nat. Scien. Phil., 1848,

p. 77.

Dipodomys hermanni Le Conte, Proc. Acad. Nat. Scien. Phil.,

1853, p. 224.

? *Dipodomys wagneri* Le Conte, Proc. Acad. Nat. Scien. Phil.,

1853, p. 224.

Perodipus agilis Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11,

1901, p. 236, fig. 54. Zoöl. Ser. *Id.* Mamm. Middle Amer.

& W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 351. Zoöl.

Ser.

NIMBLE KANGAROO RAT.

Type locality. Los Angeles, Los Angeles County, California.

Geogr. Distr. Middle and southern California, south to Lower California, Mexico.

a. Perodipus agilis tularensis Merriam.

Perodipus agilis tularensis Merr., Proc. Biol. Soc. Wash., XVII,

1904, p. 143.

TULARE KANGAROO RAT.

Type locality. Alila, Tulare County, California.

Geogr. Distr. Known only from the type locality and Tejon Cañon, California.

Genl. Char. Like *P. agilis*. Maxillary arches more developed and with a prominent recurved angle on outer side.

Color. Externally like *P. agilis*.

Measurements. Type. Total length, 308; tail vertebræ, 182; hind foot, 41.

727. *Perodipus venustus* Merriam.

Perodipus venustus Merr., Proc. Biol. Soc. Wash., XVII, 1904, p. 142.
GRACEFUL KANGAROO RAT.

Type locality. Santa Cruz, Santa Cruz County, California.

Geogr. Distr. Santa Cruz and Santa Lucia Mountains, California.

Genl. Char. Similar to *P. agilis*, but darker in color; skull slightly longer; maxillary arch of zygoma broader on outer side, with an outer angle; nasals longer; premaxillæ broader; incisors heavier.

Color. Nose black; base of whiskers black; top of head, back, and thigh patches dusky, finely grizzled with ochraceous; ears large and nearly black, with pale spots at base and top of fold; ankles, side of heels, soles, and tail stripes nearly black; black patch in front of basal white ring of tail.

Measurements. Type. Total length, 339; tail vertebræ, 211; hind foot, 46.

728. *Perodipus goldmani* Merriam.

Perodipus goldmani Merr., Proc. Biol. Soc. Wash., XVII, 1904, p. 143.
GOLDMAN'S KANGAROO RAT.

Type locality. Salinas, mouth of Salinas Valley, Monterey County, California.

Geogr. Distr. Known only from the type locality.

Genl. Char. Size large; tail short; ears small.

Color. Upper parts finely mixed dusky and buffy ochraceous; flanks grizzled ochraceous; conspicuous dusky marks at base of whiskers and on ankles.

Measurements. Type. Total length, 312; tail vertebræ, 185; hind foot, 46.

729. *Perodipus streatori* Merriam.

Perodipus streatori Merr., Proc. Biol. Soc. Wash., IX, 1894, p. 113.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 237.

Zoöl. Ser.

STREATOR'S KANGAROO RAT.

Type locality. Carbondale, Mariposa County, California.

Geogr. Distr. West foot of Sierra Nevada, California.

a. Perodipus streatori simulans Merriam.

Perodipus streatori simulans Merr., Proc. Biol. Soc. Wash., xvii, 1904, p. 144.

COUNTERFEIT KANGAROO RAT.

Type locality. Dulzura, San Diego County, California.

Geogr. Distr. San Diego County to Morro in San Luis Obispo County, California.

Genl. Char. End of tail not white; maxillaries separated.

Color. Like *P. streatori*.

Measurements. Total length, 280; tail vertebræ, 165; hind foot, 40.

730. *Perodipus panamintinus* Merriam.

Perodipus panamintinus Merr., Proc. Biol. Soc. Wash., ix, 1894, p. 114. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 239. Zoöl. Ser.

PANAMINT KANGAROO RAT.

Type locality. Head of Willow Creek, Panamint Mountains, Inyo County, California.

Geogr. Distr. Panamint Mountains, eastern California.

731. *Perodipus cabezonæ* Merriam.

Perodipus cabezonæ Merr., Proc. Biol. Soc. Wash., xvii, 1904, p. 144.

CABEZON KANGAROO RAT.

Type locality. Cabezon, Colorado Desert, California.

Geogr. Distr. Known only from the type locality.

Genl. Char. Size small; ears large. Skull rather long and narrow; sides of fronto-parietal shield approximating anteriorly; maxillary arches compressed.

Color. "Buffy ochraceous. Externally most like *P. panamintinus*."

Measurements. Type. Total length, 275; tail vertebræ, 162; hind foot, 42.3.

732. *Perodipus montanus* (Baird).

Dipodomys montanus Baird, Proc. Acad. Nat. Scien. Phil., 1855, p. 334. Mearns, Proc. Biol. Soc. Wash., xvii, 1904, p. 140. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, Appendix, p. 431. Zoöl. Ser.

Perodipus montanus Merr., Proc. Biol. Soc. Wash., xvii, 1904, p. 140.

BAIRD'S KANGAROO RAT.

Type locality. Near Fort Garland (Old Fort Massachusetts), San Luis Valley, south central Colorado.

Geogr. Distr. Known only from the type locality. Range undetermined.

Genl. Char. Size medium; tail rather short; ears small. Skull: rostrum, nasals, and premaxillæ rather narrow; bullæ rather large; maxillary arch weak, narrow outer angle well developed; supraoccipital and interparietal broad.

Color. Upper parts dull buffy ochraceous, lined on head with dark-tipped hairs; end of nose dusky; patch at base of whiskers dusky; usual pale spots at base and fold of ear; dorsal tail stripe pale brownish drab; under stripe narrowing towards tip, often not passing end of vertebræ.

Measurements. Average of 40 specimens from type locality. Total length, 250; tail vertebræ, 140; hind foot, 40.8.

a. Perodipus montanus utahensis Merriam.

Perodipus montanus utahensis Merr., Proc. Biol. Soc. Wash., xvii, 1904, p. 143.

UTAH KANGAROO RAT.

Type locality. Ogden, Utah.

Geogr. Distr. Range unknown.

Genl. Char. Similar to *P. montanus*; hind foot smaller; upper parts less fulvous; ears darker, the anterior fold dusky except at extreme tip; ventral tail stripe continuous to tip of pencil.

Measurements. Total length, 260; tail vertebræ, 150; hind foot, 41.

733. *Perodipus ordi* (Woodhouse).

Dipodomys ordii Woodh., Proc. Acad. Nat. Scien. Phil., 1853, p. 235.

Perodipus ordi Elliot, Syn. N. Am. Mamm., F. C. M. Pub., ii, 1901, p. 238. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 350. Zoöl. Ser.

ORD'S KANGAROO RAT.

Type locality. El Paso, El Paso County, Texas.

Geogr. Distr. From the Snake Plains and their northward extensions south into the State of Chihuahua, Mexico.

a. Perodipus ordi palmeri (Allen).

Dipodops ordii palmeri Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1891, p. 276.

Perodipus ordi palmeri Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 350. Zoöl. Ser.

PALMER'S KANGAROO RAT.

Type locality. San Luis Potosi, State of San Luis Potosi, Mexico.

Geogr. Distr. State of San Luis Potosi, Mexico.

b. Perodipus ordi columbianus Merriam.

Perodipus ordi columbianus Merr., Proc. Biol. Soc. Wash., ix, 1894, p. 115. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 238. Zoöl. Ser.

COLUMBIAN KANGAROO RAT.

Type locality. Umatilla, at mouth of Umatilla River, Plains of the Columbia, Umatilla County, Oregon.

Geogr. Distr. Plains of the Columbia, Oregon.

734. *Perodipus richardsoni* (Allen).

Dipodops richardsoni Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1891, p. 277.

Perodipus richardsoni Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 238. Zoöl. Ser.

RICHARDSON'S KANGAROO RAT.

Type locality. Source of the Beaver River, northwestern corner of Indian Territory.

Geogr. Distr. Oklahoma and Indian Territories.

735. *Perodipus chapmani* (Mearns).

Dipodomys chapmani Mearns, Bull. Am. Mus. Nat. Hist. N. Y., 1890, p. 291.

Perodipus chapmani Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 237. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 350, fig. 66. Zoöl. Ser.

CHAPMAN'S KANGAROO RAT.

Type locality. Fort Verde, Yavapai County, Arizona.

Geogr. Distr. Elevated central tract, Arizona.

736. *Perodipus sennetti* (Allen).

Dipodops sennetti Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1891, p. 226.

Perodipus sennetti Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 239. Zoöl. Ser.

SENNETT'S KANGAROO RAT.

Type locality. Near Brownsville, Cameron County, Texas.

Geogr. Distr. Southeastern Texas.

737. *Perodipus microps* Merriam.

Perodipus microps Merr., Proc. Biol. Soc. Wash., xvii, 1904, p. 145.

SMALL-FACE KANGAROO RAT.

Type locality. Lone Pine, Owens Valley, Inyo County, California.

Geogr. Distr. Known only from the type locality.

Genl. Char. Size small; ears small; color pale. Skull small and rather narrow; breadth of parietal less than the length; maxillary arch weak and slender, without external angle.

Color. General color, "pale buffy ochraceous, much paler than *ordi* or *columbianus*."

Measurements. Type. Total length, 282; tail vertebræ, 165; hind foot, 41.

a. Perodipus microps levipes Merriam.

Perodipus microps levipes Merr., Proc. Biol. Soc. Wash., xvii, 1904, p. 145.

LIGHT-FOOTED KANGAROO RAT.

Type locality. Perognathus Flat, Emigrant Gap, Panamint Mountains, California.

Geogr. Distr. Panamint Mountains, California.

Genl. Char. Size small; color as in *P. microps*. Skull small; bullæ bulging posteriorly; maxillary arches weak.

Color. Like *P. microps*.

Measurements. Type. Total length, 288; tail vertebræ, 164; hind foot, 42.4.

738. *Perodipus ingens* Merriam.

Perodipus ingens Merr., Proc. Biol. Soc. Wash., xvii, 1904, p. 141.

GREAT KANGAROO RAT.

Type locality. Painted Rock, twenty miles southeast of Simmler, Carrizo Plain, San Luis Obispo County, California.

Geogr. Distr. Carrizo Plain, and southwestern border of San Joaquin Valley, California.

Genl. Char. Size large; ears small.

Color. Buffy ochraceous; upper and lower tail bands black, uniting at end of vertebræ; pencil white washed with dusky.

Measurements. Total length, 360; tail vertebræ, 191; hind foot, 54. Skull: occipito-nasal length, 45; basal length, 34; zygomatic breadth, 23.5; breadth across bullæ, 30.5; breadth of frontals posteriorly, 17; behind lachrymals, 15; length of nasals, 18.

739. *Perodipus longipes* (Merriam).

Dipodops longipes Merr., N. Am. Faun., No. 3, 1890, p. 72.

Perodipus longipes Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 239. Zool. Ser.

LONG-FOOTED KANGAROO RAT.

Type locality. Foot of Echo Cliffs, Painted Desert, Coconino County, Arizona.

Geogr. Distr. Utah, Wyoming, Nebraska, New Mexico, and Arizona.

740. *Perodipus obscurus* Allen.

Perodipus obscurus Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1903, p. 603. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 351. Zool. Ser.

DUSKY KANGAROO RAT.

Type locality. Rio Sestin, State of Durango, Mexico.

Geogr. Distr. State of Durango.

741. *Perodipus compactus* (True).

Dipodomys compactus True, Proc. U. S. Nat. Mus., 1888, p. 160.

Perodipus compactus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 240. Zool. Ser.

THICK-SET KANGAROO RAT.

Type locality. Padre Island, Cameron County, Texas.

Geogr. Distr. Known only from the type locality.

Genus 97. *Microdipodops*. Small Kangaroo Rats.

***Microdipodops* Merr., N. Am. Faun., No. 5, 1891, p. 115. Type**

Microdipodops megacephalus Merriam.

742. *Microdipodops megacephalus* Merriam.

Microdipodops megacephalus Merr., N. Am. Faun., No. 5, 1891, p. 116. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 241, fig. 55. Zool. Ser.

LARGE-HEADED KANGAROO RAT.

Type locality. Halleck, East Humboldt Valley, Elko County, Nevada.

Geogr. Distr. Central Nevada.

a. *Microdipodops megacephalus oregonus* Merriam.

Microdipodops megacephalus oregonus Merr., Proc. Biol. Soc. Wash., XIV, 1901, p. 127.

OREGON KANGAROO RAT.

Type locality. Lake Alvord, Alvord Desert, Harney County, Oregon.

Geogr. Distr. Known only from the type locality.

Genl. Char. Similar to *M. megacephalus*, but tail longer.

Color. Above olivaceous, slightly lined with black; beneath white; under side of tail with white streak.

Measurements. Total length, 153; tail vertebræ, 88; hind foot, 21.

743. *Microdipodops pallidus* Merriam.

Microdipodops pallidus Merr., Proc. Biol. Soc. Wash., XIV, 1901, p. 127.

PALLID KANGAROO RAT.

Type locality. Ten miles east of Stillwater, near sink of the Humboldt and Carson, Churchill County, Nevada.

Geogr. Distr. Known only from the type locality.

Genl. Char. Tail long and without dark tip.

Color. Above pale buffy fulvous, lined with black; beneath white; side of nose, face and legs white; tail above buffy, beneath white.

Measurements. Total length, 171; tail vertebræ, 102; hind foot, 25.5.

744. *Microdipodops californicus* Merriam.

Microdipodops californicus Merr., Proc. Biol. Soc. Wash., xiv, 1901, p. 128.

CALIFORNIA KANGAROO RAT.

Type locality. Sierra Valley near Vinton, Plumas County, California.

Geogr. Distr. Known only from the type locality.

Genl. Char. Size of *M. megacephalus*; tail and hind foot longer.

Color. Above olivaceous, lined with black; beneath white; sides of nose, spot over the eye, behind and on the fold of the ear and feet white; tail above pale fulvous, blackish at tip, beneath white; sides of face and inner side of fore leg washed with pale buffy fulvous.

Measurements. Total length, 158; tail-vertebræ, 91; hind foot, 25.

Subfamily II. *Heteromyinæ*.**Genus 98. *Perognathus*. Pocket Mice.**

Perognathus Max., Reise Innere N. Amer., 1, 1839, pp. 449-450; *Id.* Nov. Act. Phys. Med. Acad. Cæs. Leop.-Carol., xix, Pt. 1, 1839, pp. 368-373, pl. xxxiv. Type *Perognathus fasciatus* Maximilian.

Cricetodipus Peale, Rep. Mamm. and Ornith. U. S. Expl. Exped., 1848, p. 53, pl. 13, fig. 2.

Abromys (sic) Gray, Proc. Zool. Soc., 1868, p. 202.

Otognosis Coues, Proc. Acad. Nat. Scien. Phil., 1875, p. 305.

Chætodipus Merr., N. Am. Faun., No. 1, 1889, p. 5, pl. III, fig. 15.

Subgenus A. *Perognathus*.**745. *Perognathus fasciatus* Maximilian.**

Perognathus fasciatus Max., Nov. Act. Acad. Cæs. Leop.-Carol., xix, Pt. 1, 1839, p. 369, pl. xxxiv. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 242. Zool. Ser.

MAXIMILIAN'S POCKET MOUSE.

Type locality. Upper Missouri River near junction with the Yellowstone River.

Geogr. Distr. Eastern Montana and Wyoming, east into North and South Dakota.

a. Perognathus fasciatus infraluteus (Thomas).

Perognathus infraluteus Thomas, Ann. Mag. Nat. Hist., 6th Ser., XI, 1893, p. 406.

Perognathus fasciatus infraluteus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 242. Zoöl. Ser.

BUFF-BELLIED POCKET MOUSE.

Type locality. Loveland, Larimer County, Colorado.

Geogr. Distr. Known only from the type locality.

746. *Perognathus flavescens* Merriam.

Perognathus flavescens Merr., N. Am. Faun., No. 1, 1889, p. 11.
Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 243.
Zoöl. Ser.

Perognathus copei Rhoads, Proc. Acad. Nat. Scien. Phil., 1893, p. 404.

PLAINS POCKET MOUSE.

Type locality. Kennedy, Cherry County, Nebraska.

Geogr. Distr. Plains of South Dakota, Nebraska, and Kansas, south possibly to northern Texas, and west to base of the Rocky Mountains.

a. Perognathus flavescens perniger Osgood.

Perognathus flavescens perniger Osgood, Proc. Biol. Soc. Wash., XVII, 1904, p. 127.

BLACK POCKET MOUSE.

Type locality. Vermillion, Clay County, South Dakota.

Geogr. Distr. Known only from the type locality.

Genl. Char. Like *P. flavescens* in size; upper parts black.

Color. Dorsal region black; sides and head black and ochraceous buff mixed; lateral line ochraceous buff; chin and narrow stripe on throat and breast white; rest of under parts ochraceous buff; subauricular spot buff; tail above dusky, beneath white; sides buffy; feet buffy; ear brownish black, edged with buff.

Measurements. Total length, 140; tail vertebræ, 68; hind foot, 17.

747. *Perognathus merriami* Allen.

Perognathus merriami Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1892, p. 45, pl. III, figs. 1-6. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 243. Zoöl. Ser. *Id.* Mamm. Middle

Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 353, figs. 67 and XLIX. Zoöl. Ser.

Cricetodipus flavus Thomas, Proc. Zoöl. Soc., 1888, p. 449.

Perognathus mearnsi Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1896, p. 237.

MERRIAM'S POCKET MOUSE.

Type locality. Brownsville, Cameron County, Texas.

Geogr. Distr. Washburn, Texas, to Roswell, New Mexico, east to San Antonio, west to Comstock on the Rio Grande, and south to Alta Mira, State of Tamaulipas, Mexico.

a. *Perognathus merriami gilvus* Osgood.

Perognathus merriami gilvus Osgood, N. Am. Faun., No. 18, 1900, p. 22. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 243. Zoöl. Ser.

DUTCHER'S POCKET MOUSE.

Type locality. Eddy, Eddy County, New Mexico.

Geogr. Distr. Western Texas and southeastern New Mexico.

748. *Perognathus flavus* Baird.

Perognathus flavus Baird, Proc. Acad. Nat. Scien. Phil., 1855, p. 332. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 244, fig. 56. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 354. Zoöl. Ser.

BAIRD'S POCKET MOUSE.

Type locality. El Paso, El Paso County, Texas.

Geogr. Distr. Northeastern Colorado and western Nebraska to northern Mexico, westward into central Arizona, eastward to western Texas, and south into the State of Chihuahua, Mexico.

a. *Perognathus flavus bimaculatus* (Merriam).

Perognathus bimaculatus Merr., N. Am. Faun., No. 1, 1889, p. 12.

Perognathus apache Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1893, p. 71. (Part.)

Perognathus flavus bimaculatus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 244. Zoöl. Ser.

YAVAPAI POCKET MOUSE.

Type locality. Fort Whipple, Yavapai County, Arizona.

Geogr. Distr. Central and northeastern Arizona, and southeastern Utah.

b. *Perognathus flavus fuliginosus* (Merriam).

Perognathus fuliginosus Merr., N. Am. Faun., No. 3, 1890, p. 74.

Perognathus flavus fuliginosus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 244. Zoöl. Ser.

DUSKY POCKET MOUSE.

Type locality. Cedar belt northeast of San Francisco Mountain, Coconino County, Arizona. Altitude, 7,000 feet.

Geogr. Distr. Lava beds near San Francisco Mountain, Coconino County, northern Arizona.

c. *Perognathus flavus mexicanus* Merriam.

Perognathus flavus mexicanus Merr., Proc. Acad. Nat. Scien. Phil., 1894, p. 265. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 355. Zoöl. Ser.

MEXICAN POCKET MOUSE.

Type locality. Tlalpam, Federal District, Mexico.

Geogr. Distr. States of Zacatecas and San Luis Potosi to the State of Jalisco on the west, and State of Hidalgo on the east, and south to the State of Mexico.

749. *Perognathus apache* Merriam.

Perognathus apache Merr., N. Am. Faun., No. 1, 1889, p. 14. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 244. Zoöl. Ser.

APACHE POCKET MOUSE.

Type locality. Keam Cañon, Apache County, Arizona.

Geogr. Distr. Eastern Arizona, western New Mexico, and southern Utah.

a. *Perognathus apache melanotis* Osgood.

Perognathus apache melanotis Osgood, N. Am. Faun., No. 18, 1900, p. 27. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 356. Zoöl. Ser.

BLACK-EARED POCKET MOUSE.

Type locality. Casas Grandes, State of Chihuahua, Mexico.

Geogr. Distr. Known only from the type locality.

750. *Perognathus pericalles* Elliot.

Perognathus pericalles Elliot, Pub. Field Columb. Mus., III, 1903, p. 252. Zoöl. Ser.

BEAUTIFUL POCKET MOUSE.

Type locality. Keeler, Owens Lake, Inyo County, California.

Geogr. Distr. Range unknown.

Genl. Char. Size small; color pale; ear rather large.

Color. Entire upper parts deep cream buff, tinged with reddish, darkest on head and rump; sides pale cream buff; upper lip and entire under parts white; feet buffy white; tail above pale brown, beneath yellowish white; ear pale brown, with a buffy tuft of hair at base.

Measurements. Total length, 130; tail vertebrae, 73; hind foot, 19; ear, 6.5. Skull: total length, 21.5; Hensel, 14.5; zygomatic width, 11; interorbital width, 5; mastoid width, 12; greatest width of parietal, 10; length of interparietal, 2.5; length of nasals, 8.4; palatal length, 7.5; length of upper tooth row, 4; length of mandible, angle to tips of incisors, 10.5; length of lower tooth row, 3.

751. *Perognathus callistus* Osgood.

Perognathus callistus Osgood, N. Am. Faun., No. 18, 1900, p. 28.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 245. Zoöl. Ser.

OSGOOD'S POCKET MOUSE.

Type locality. Kinney Ranch, Green River Basin, near Bitter Creek, Sweetwater County, Wyoming.

Geogr. Distr. Known only from the type locality.

752. *Perognathus elibatus* Elliot.

Perognathus elibatus Elliot, Pub. Field Columb. Mus., III, 1903, p. 252. Zoöl. Ser.

MOUNT PINOS POCKET MOUSE.

Type locality. Mount Pinos, Ventura County, California. Altitude, 5,000 feet.

Geogr. Distr. Known only from the type locality.

Genl. Char. Size small; color dark; tail long.

Color. Upper parts black and buff, the former color predominating; nose, sides of face, line over eye, and lateral line cream buff; under parts, hands, and feet white; tail, basal half above buff, tinged with dusky, remainder dusky, beneath yellowish white; ears brown, a white spot over each side of margin near notch; whiskers black; line on sides of nose black.

Measurements. Total length, 146; tail, 77; hind foot, 20.5; ear, 7. Skull: total length, 22; Hensel, 15; zygomatic width, 11; interorbital constriction, 5; mastoid width, 12; greatest width of parietal, 9; length of nasals, 7; palatal length, 9; length of upper tooth row, 3; length of mandible, 11; length of lower tooth row, 3.

753. *Perognathus panamintinus* (Merriam).

Perognathus longimembris panamintinus Merr., Proc. Acad. Nat. Scien. Phil., 1894, p. 265.

Perognathus panamintinus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 245. Zoöl. Ser.

PANAMINT POCKET MOUSE.

Type locality. Perognathus Flat, Panamint Mountains, Inyo County, California. Altitude, 5,200 feet.

Geogr. Distr. Panamint Mountains, California, and eastward through southern Nevada to St. George, Utah.

a. *Perognathus panamintinus bangsi* (Mearns).

Perognathus longimembris bangsi Mearns, Bull. Am. Mus. Nat. Hist. N. Y., 1898, p. 300.

Perognathus panamintinus bangsi Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 245. Zool. Ser.

BANGS' POCKET MOUSE.

Type locality. Palm Springs, Colorado Desert, Riverside County, California.

Geogr. Distr. Desert valleys in southern and southeastern California.

b. *Perognathus panamintinus arenicola* Stephens.

Perognathus panamintinus arenicola Steph., Proc. Biol. Soc. Wash., XIII, 1900, p. 153. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 246. Zool. Ser.

SAND-DWELLING POCKET MOUSE.

Type locality. San Felipe Narrows, San Diego County, California.

Geogr. Distr. Known only from the type locality.

c. *Perognathus panamintinus brevinasus* Osgood.

Perognathus panamintinus brevinasus Osgood, N. Am. Faun., No. 18, 1900, p. 30. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 246. Zool. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 355. Zool. Ser.

SHORT-NOSED POCKET MOUSE.

Type locality. San Bernardino, San Bernardino County, California.

Geogr. Distr. Southeastern California from Coso Valley south to Oro Grande in the Mohave Desert, and westward to Fort Tejon; Daggett, and the Calico Mountains appearing to be the eastern limit of the range. Lower California.

754. *Perognathus nevadensis* Merriam.

Perognathus nevadensis Merr., Proc. Acad. Nat. Scien. Phil., 1894, p. 264. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 246. Zool. Ser.

NEVADA POCKET MOUSE.

Type locality. Halleck, East Humboldt Valley, Elko County, Nevada.

Geogr. Distr. Central Nevada north to southern Oregon, and northern Utah.

755. *Perognathus pacificus* Mearns.

Perognathus pacificus Mearns, Bull. Am. Mus. Nat. Hist. N. Y., 1898, p. 299. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 246. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., 14, Pt. 1, 1904, p. 355. Zoöl. Ser.

PACIFIC POCKET MOUSE.

Type locality. Mouth of Tijuana River, Mexican boundary Monument, No. 258, San Diego County, California.

Geogr. Distr. Edge of the Pacific Ocean on a flat at the mouth of the Tijuana River. Range unknown.

756. *Perognathus amplus* Osgood.

Perognathus amplus Osgood, N. Am. Faun., No. 18, 1900, p. 32. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 247. Zoöl. Ser.

LORING POCKET MOUSE.

Type locality. Fort Verde, Yavapai County, Arizona.

Geogr. Distr. Known only from the type locality.

757. *Perognathus longimembris* (Coues).

Otognosis longimembris Coues, Proc. Acad. Nat. Scien. Phil., 1875, p. 305.

Cricetodipus parvus True, Proc. U. S. Nat. Mus., 1882, p. 474.

Perognathus inornatus Merr., N. Am. Faun., No. 1, 1889, p. 15.

Perognathus longimembris Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 247. Zoöl. Ser.

SAN JOAQUIN POCKET MOUSE.

Type locality. Old Fort Tejon, Kern County, California.

Geogr. Distr. San Joaquin Valley, southern California.

758. *Perognathus parvus* (Peale).

Cricetodipus parvus Peale, U. S. Expl. Exped., VIII, 1848, p. 53, pl. 13, fig. 2, juv.

Perognathus parvus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 247. Zoöl. Ser.

OREGON POCKET MOUSE.

Type locality. The Dalles? Wasco County, Oregon.

Geogr. Distr. Range unknown.

759. *Perognathus monticola* Baird.

Perognathus monticola Baird, N. Am. Mamm., 1857, p. 422. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 248. Zoöl. Ser.

Perognathus parvus Osgood, N. Am. Faun., No. 18, 1900, p. 34. (nec Peale.)

BAIRD'S POCKET MOUSE.

Type locality. St. Mary's Mission? Montana, west of Rocky Mountains.

Geogr. Distr. Valley of the Yakima River, Washington, south to central and southeastern Oregon.

a. *Perognathus monticola mollipilosus* (Coues).

Perognathus mollipilosus Coues, Proc. Acad. Nat. Scien. Phil., 1875, p. 296.

Perognathus monticola Towns., Proc. U. S. Nat. Mus., x, 1888, p. 177. (nec Baird.)

Perognathus monticola mollipilosus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 249. Zool. Ser.

COUES' POCKET MOUSE.

Type locality. Fort Crook, Shasta County, California.

Geogr. Distr. Northeastern California to Klamath Basin, Oregon.

b. *Perognathus monticola olivaceus* (Merriam).

Perognathus olivaceus Merr., N. Am. Faun., No. 1, 1889, p. 15.

Perognathus olivaceus amœnus Merr., N. Am. Faun., No. 1, 1889, p. 16.

Perognathus monticola olivaceus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 249. Zool. Ser.

GREAT BASIN POCKET MOUSE.

Type locality. Kelton, near north end of Great Salt Lake, Boxelder County, Utah.

Geogr. Distr. Northern Utah and southern Idaho to Owens Valley, southwestern California, west to southern Oregon and northeastern California.

c. *Perognathus monticola magruderensis* (Osgood).

Perognathus parvus magruderensis Osgood, N. Am. Faun., No. 18, 1900, p. 38.

Perognathus monticola magruderensis Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 249. Zool. Ser.

MOUNT MAGRUDER POCKET MOUSE.

Type locality. Mount Magruder, Nevada, near boundary between Inyo County, California, and Esmeralda County, Nevada. Altitude, 8,000 feet.

Geogr. Distr. Southern Nevada and adjacent parts of eastern California.

760. *Perognathus alticola* Rhoads.

Perognathus alticola Rhoads, Proc. Acad. Nat. Scien. Phil., 1893, p. 412. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 249. Zool. Ser.

WHITE-EARED POCKET MOUSE.

Type locality. Squirrel Inn, San Bernardino Mountains, San Bernardino County, California.

Geogr. Distr. Known only from the type locality.

761. *Perognathus lordi* (Gray).

Abromys (sic) lordi Gray, Proc. Zool. Soc., 1868, p. 202.

Perognathus lordi Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 250. Zool. Ser.

NORTHWEST POCKET MOUSE.

Type locality. Southern British Columbia, Canada.

Geogr. Distr. Southern British Columbia to the plains of the Columbia River, Washington.

a. *Perognathus lordi columbianus* (Merriam).

Perognathus columbianus Merr., Proc. Acad. Nat. Scien. Phil., 1894, p. 263.

Perognathus lordi columbianus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 250. Zool. Ser.

COLUMBIAN POCKET MOUSE.

Type locality. Pasco, Franklin County, Washington.

Geogr. Distr. Known only from the type locality.

762. *Perognathus formosus* Merriam.

Perognathus formosus Merr., N. Am. Faun., No. 1, 1889, p. 17, Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 250. Zool. Ser.

LONG-TAILED POCKET MOUSE.

Type locality. St. George, Washington County, Utah.

Geogr. Distr. Southwestern Utah, southern Nevada, and adjacent portions of California as far south as Copper City.

763. *Perognathus mesembrinus* Elliot.

Perognathus mesembrinus Elliot, Pub. Field Columb. Mus., 111, 1903, p. 251. Zool. Ser.

SOUTHERN POCKET MOUSE.

Type locality. Palm Springs, Colorado Desert, Riverside County, California.

Geogr. Distr. Southern California, probably into Lower California, Mexico.

Genl. Char. Color pale; tail hairy, pencil large, bushy; size medium.

Color. Upper parts mixed drab gray and buff; no lateral line; thighs like the back; lips, entire under parts, fore legs, fore and hind feet white; tail above with pencil brownish drab, beneath whitish; ear dark brown; bases covered with tufts of drab gray.

Measurements. Total length, 195; tail vertebrae, 114; hind foot, 23; ear, 11. Skull: total length, 21; Hensel, 18; zygomatic width, 13; interorbital constriction, 7; mastoid width, 14; greatest parietal width, 10.5; length of nasals, 9.4; palatal length, 10; length of upper tooth row, 4; length of mandible, 12.5; length of lower tooth row, 3.5.

Subgenus B. *Chatodipus*.

764. *Perognathus baileyi* Merriam.

Perognathus baileyi Merr., Proc. Acad. Nat. Scien. Phil., 1894, p. 262, fig. 1. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 251. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 361. Zoöl. Ser.

BAILEY'S POCKET MOUSE.

Type locality. Magdalena, State of Sonora, Mexico.

Geogr. Distr. South central Arizona, south into the State of Sonora and Lower California, Mexico.

a. Perognathus baileyi rhydinorhis Elliot.

Perognathus baileyi rhydinorhis Elliot, Pub. Field Columb. Mus., III, 1903, p. 167. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 362. Zoöl. Ser.

SAN QUENTIN POCKET MOUSE.

Type locality. San Quentin, Lower California, Mexico.

Geogr. Distr. West coast of Lower California, Mexico, in the vicinity of San Quentin.

765. *Perognathus cneus* Elliot.

Perognathus cneus Elliot, Pub. Field Columb. Mus., III, 1903, p. 169. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 362. Zoöl. Ser.

BUFF-COLORED POCKET MOUSE.

Type locality. Rosarito, San Pedro Martir Mountains, Lower California, Mexico.

Geogr. Distr. Known only from the type locality.

766. *Perognathus hispidus* Baird.

Perognathus fasciatus Baird, N. Am. Mamm., 1857, p. 420. (nec Max.)

Perognathus hispidus Baird, N. Am. Mamm., 1857, p. 421.

Perognathus paradoxus spilotos Merr., N. Am. Faun., No. 1, 1889, p. 25.

Perognathus hispidus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 251. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 360. Zoöl. Ser.

HISPID POCKET MOUSE.

Type locality. Charco Escondido, State of Tamaulipas, Mexico.

Geogr. Distr. Southern and western Texas, north to Oklahoma and south to the border states of Mexico.

a. Perognathus hispidus paradoxus (Merriam).

Perognathus paradoxus Merr., N. Am. Faun., No. 1, 1889, p. 24.

Perognathus latirostris Rhoads, Amer. Nat., xxviii, 1894, p. 185.

Perognathus conditi Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1894, p. 318.

Perognathus hispidus paradoxus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 252. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 361. Zoöl. Ser.

KANSAS POCKET MOUSE.

Type locality. Banner, Trego County, Kansas.

Geogr. Distr. The Dakotas to Texas, westward to the base of the Rocky Mountains.

b. Perognathus hispidus maximus Elliot.

Perognathus hispidus maximus Elliot, Pub. Field Columb. Mus., III, 1903, p. 253. Zoöl. Ser.

GREAT POCKET MOUSE.

Type locality. Noble, Oklahoma Territory.

Geogr. Distr. Oklahoma Territory. Range unknown.

Genl. Char. Similar to *P. h. paradoxus*, but brighter and richer in color; hind foot and other dimensions larger.

Color. Upper parts mixed ochraceous and black, the latter color predominating; lateral line from nose to rump, including shoulders and upper parts of arms and hind legs very bright ochraceous buff; face and orbital region bright ochraceous buff lightly lined with black; under parts, hands, and feet white; ear buff on outside, dusky inside; tail above blackish brown, sides buff, beneath white.

Measurements. Total length, 243; tail vertebræ, 110; hind foot, 29. Skull: total length, 34; Hensel, 25; zygomatic width, 16.5; mastoid width, 16; length of parietal, 5; greatest width of parietal, 14; length of nasals, 10.5; palatal length, 14; length of upper tooth row, 5; length of mandible, 20; length of lower tooth row, 4.

c. Perognathus hispidus zacatecæ Osgood.

Perognathus hispidus zacatecæ Osgood, N. Am. Faun., No. 18, 1900, p. 45. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 361. Zoöl. Ser.

ZACATECAS POCKET MOUSE.

Type locality. Valparaiso, State of Zacatecas, Mexico.

Geogr. Distr. States of Zacatecas to Celaya in State of Guana-
juato, Mexico.

767. *Perognathus penicillatus* Woodhouse.

Perognathus penicillatus Woodh., Proc. Acad. Nat. Scien. Phil.,
1852, p. 200. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II,
1901, p. 252. Zoöl. Ser.

DESERT POCKET MOUSE.

Type locality. San Francisco Mountain, Coconino County, Ari-
zona.

Geogr. Distr. Vicinity of the Colorado River from Bunkerville,
Nevada, to Yuma, Arizona.

a. Perognathus penicillatus angustirostris Osgood.

Perognathus penicillatus angustirostris Osgood, N. Am. Faun., No.
18, 1900, p. 47. Elliot, Syn. N. Am. Mamm., F. C. M. Pub.,
II, 1901, p. 252. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W.
Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 358. Zoöl. Ser.

SLENDER-NOSED POCKET MOUSE.

Type locality. Carriso Creek, Colorado Desert, San Diego County,
California.

Geogr. Distr. Colorado Desert, California, to southwestern Ari-
zona, and northern Lower California, Mexico.

b. Perognathus penicillatus pricei (Allen).

Perognathus pricei Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1894,
p. 318.

Perognathus obscurus Allen, Bull. Am. Mus. Nat. Hist. N. Y.,
1895, p. 216. (nec Merr.)

Perognathus penicillatus pricei Elliot, Syn. N. Am. Mamm., F. C. M.
Pub., II, 1901, p. 253. Zoöl. Ser. *Id.* Mamm. Middle Amer.
& W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 358. Zoöl. Ser.

PRICE'S POCKET MOUSE.

Type locality. Oposura, State of Sonora, Mexico.

Geogr. Distr. South central Arizona, into northern Mexico, west
of the Sierra Madre.

c. Perognathus penicillatus eremicus (Mearns).

Perognathus eremicus Mearns, Bull. Am. Mus. Nat. Hist. N. Y.,
1898, p. 300. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II,
1901, p. 253. Zoöl. Ser.

Perognathus penicillatus eremicus Elliot, Mamm. Middle Amer.
& W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 359. Zoöl. Ser.

EASTERN DESERT POCKET MOUSE.

Type locality. Fort Hancock, El Paso County, Texas.

Geogr. Distr. Western Texas south to north central Mexico east of the Sierra Madre, to La Ventura, State of Coahuila.

d. Perognathus penicillatus arenarius (Merriam).

Perognathus arenarius Merr., Proc. Cal. Acad. Scien., 2d Ser., IV, 1894, p. 461.

Perognathus penicillatus arenarius Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 358. Zoöl. Ser.

LITTLE DESERT POCKET MOUSE.

Type locality. San Jorge, near Comondu, Lower California, Mexico.

Geogr. Distr. East coast of Lower California.

768. *Perognathus helleri* Elliot.

Perognathus helleri Elliot, Pub. Field Columb. Mus., III, 1903, p. 166. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 360. Zoöl. Ser.

HELLER'S POCKET MOUSE.

Type locality. San Quentin, Lower California, Mexico.

Geogr. Distr. Known only from the type locality.

769. *Perognathus stephensi* Merriam.

Perognathus stephensi Merr., Proc. Acad. Nat. Scien. Phil., 1894, p. 267. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 253. Zoöl. Ser.

STEPHENS' POCKET MOUSE.

Type locality. Mesquite Valley, northwest arm of Death Valley, Inyo County, California.

Geogr. Distr. Known only from the type locality.

770. *Perognathus pernix* Allen.

Perognathus pernix Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1898, p. 149. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 359. Zoöl. Ser.

SINALOA POCKET MOUSE.

Type locality. Rosario, State of Sinaloa, Mexico.

Geogr. Distr. States of Sinaloa and Jalisco, Mexico.

a. Perognathus pernix rostratus Osgood.

Perognathus pernix rostratus Osgood, N. Am. Faun., No. 18, 1900, p. 51. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 359. Zoöl. Ser.

BROAD-NOSED POCKET MOUSE.

Type locality. Camoa, Rio Mayo, State of Sonora, Mexico.

Geogr. Distr. Southern part of the State of Sonora, and northern portion of the State of Sinaloa, plains of the coast, Mexico.

771. *Perognathus intermedius* Merriam.

Perognathus intermedius Merr., N. Am. Faun., No. 1, 1889, p. 18, pl. 11, fig. 13. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 253. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., 14, Pt. 1, 1904, p. 364. Zoöl. Ser.

Perognathus obscurus Merr., N. Am. Faun., No. 1, 1889, p. 20.

INTERMEDIATE POCKET MOUSE.

Type locality. Mud Spring, Mohave County, Arizona.

Geogr. Distr. Arizona, New Mexico, and northern Mexico.

772. *Perognathus nelsoni* Merriam.

Perognathus nelsoni Merr., Proc. Acad. Nat. Scien. Phil., 1894, p. 266. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., 14, Pt. 1, 1904, p. 364. Zoöl. Ser.

NELSON'S POCKET MOUSE.

Type locality. Hacienda La Parada, 25 miles northwest of the city of San Luis Potosi, State of San Luis Potosi, Mexico.

Geogr. Distr. States of San Luis Potosi and Durango, south into the State of Jalisco and east into the State of Tamaulipas, Mexico.

a. *Perognathus nelsoni canescens* (Merriam).

Perognathus intermedius canescens Merr., Proc. Acad. Nat. Scien. Phil., 1894, p. 267.

Perognathus nelsoni canescens Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., 14, Pt. 1, 1904, p. 365. Zoöl. Ser.

JARAL POCKET MOUSE.

Type locality. Jaral, State of Coahuila, Mexico.

Geogr. Distr. Known only from the type locality

773. *Perognathus goldmani* Osgood.

Perognathus goldmani Osgood, N. Am. Faun., No. 18, 1900, p. 54.

Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., 14, Pt. 1, 1904, p. 365. Zoöl. Ser.

GOLDMAN'S POCKET MOUSE.

Type locality. Sinaloa, State of Sinaloa, Mexico.

Geogr. Distr. States of Sonora and Sinaloa, Mexico; coast plains.

a. *Perognathus goldmani artus* (Osgood).

Perognathus artus Osgood, N. Am. Faun., No. 18, 1900, p. 55.

Perognathus goldmani artus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., 14, Pt. 1, 1904, p. 365. Zoöl. Ser.

BATOPILAS POCKET MOUSE.

Type locality. Batopilas, State of Chihuahua, Mexico.

Geogr. Distr. States of Chihuahua, Durango, and Sinaloa, Mexico.

774. *Perognathus fallax* Merriam.

Perognathus fallax Merr., N. Am. Faun., No. 1, 1889, p. 19.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 254.

Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M.

Pub., IV, Pt. 1, 1904, p. 366. Zoöl. Ser.

SHORT-EARED CALIFORNIA POCKET MOUSE.

Type locality. Reche Cañon, three miles southeast of Colton, San Bernardino County, California.

Geogr. Distr. Southern California, and northern Lower California, Mexico.

a. *Perognathus fallax pallidus* Mearns.

Perognathus fallax pallidus Mearns, Proc. Biol. Soc. Wash., xiv,

1901, p. 135. Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., IV, Pt. 1, 1904, p. 366. Zoöl. Ser.

PALLID POCKET MOUSE.

Type locality. Mountain Spring, east slope of the Coast Range Mountains on the Mexican boundary line, San Diego County, California.

Geogr. Distr. Riverside County, southern California, into Lower California.

775. *Perognathus anthonyi* Osgood.

Perognathus anthonyi Osgood, N. Am. Faun., No. 18, 1900, p. 56.

Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV,

Pt. 1, 1904, p. 366. Zoöl. Ser.

ANTHONY'S POCKET MOUSE.

Type locality. South Bay, Cerros or Cedros Island, Lower California, Mexico.

Geogr. Distr. Known only from the type locality.

776. *Perognathus femoralis* Allen.

Perognathus femoralis Allen, Bull. Am. Mus. Nat. Hist. N. Y.,

1891, p. 281. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11,

1901, p. 254. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. In-

dies F. C. M. Pub., IV, Pt. 1, 1904, p. 367. Zoöl. Ser.

GREAT CALIFORNIA POCKET MOUSE.

Type locality. Dulzura, San Diego County, California.

Geogr. Distr. Northern Lower California, Mexico, and adjoining portions of southern California.

a. Perognathus femoralis mesopolius Elliot.

Perognathus femoralis mesopolius Elliot, Pub. Field Columb. Mus., III, 1903, p. 168. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 367, fig. 68. Zoöl. Ser.

GRAY POCKET MOUSE.

Type locality. Piñon, San Pedro Martir Mountains, Lower California, Mexico. Altitude, 5,000 feet.

Geogr. Distr. San Pedro Martir Range at high elevations.

777. *Perognathus californicus* Merriam.

Perognathus californicus Merr., N. Am. Faun., No. 1, 1889, p. 26.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 254.

Zoöl. Ser.

CALIFORNIA POCKET MOUSE.

Type locality. Berkeley, Alameda County, California.

Geogr. Distr. Coast region of California from San Francisco Bay to Bear Valley, San Benito Valley, San Benito County, California.

a. Perognathus californicus dispar Osgood.

Perognathus californicus dispar Osgood, N. Am. Faun., No. 18,

1900, p. 58. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II,

1901, p. 255. Zoöl. Ser.

ALLEN'S POCKET MOUSE.

Type locality. Carpenteria, Santa Barbara County, California.

Geogr. Distr. Coast valleys of California from San Bernardino to San Benito County, and north along the foothills of the west slope of the Sierras to Placer County.

b. Perognathus californicus ochrus Osgood.

Perognathus californicus ochrus Osgood, Proc. Biol. Soc. Wash.,

XVII, 1904, p. 128.

YELLOW POCKET MOUSE.

Type locality. Santiago Springs, Kern County, California.

Genl. Char. Like *P. c. dispar*, but paler.

Color. Upper parts mixed with pinkish buff and dusky; lateral line pinkish buff; under parts creamy white; tail above hair brown, beneath white; hands and feet white.

Measurements. Total length, 200; tail vertebræ, 108.7; hind foot, 25.

778. *Perognathus spinatus* Merriam.

Perognathus spinatus Merr., N. Am. Faun., No. 1, 1889, p. 21.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 255.

Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M.

Pub., IV, Pt. I, 1904, p. 363. Zoöl. Ser.

SPINY POCKET MOUSE.

Type locality. Twenty-five miles below the Needles, Colorado River, San Bernardino County, California.

Geogr. Distr. Southern California, and northern Lower California Mexico.

a. Perognathus spinatus peninsulæ Merriam.

Perognathus spinatus peninsulæ Merr., Proc. Cal. Acad. Scien., 2d Ser., IV, 1894, p. 460. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 363. Zoöl. Ser.

CAPE ST. LUCAS POCKET MOUSE.

Type locality. San José del Cabo, Lower California, Mexico.

Geogr. Distr. Cape region of Lower California.

b. Perognathus spinatus bryanti (Merriam).

Perognathus bryanti Merr., Proc. Cal. Acad. Scien., 2d Ser., IV, 1894, p. 458.

Perognathus spinatus bryanti Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 364. Zoöl. Ser.

BRYANT'S POCKET MOUSE.

Type locality. San José Island, Lower California, Mexico.

Geogr. Distr. Known only from the type locality.

779. *Perognathus margaritæ* Merriam.

Perognathus margaritæ Merr., Proc. Cal. Acad. Scien., 2d Ser., IV, 1894, p. 459. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 363. Zoöl. Ser.

MARGARITA POCKET MOUSE.

Type locality. Santa Margarita Island, Lower California, Mexico.

Geogr. Distr. Known only from the type locality.

Genus 99. **Heteromys. Spiny Pouched Rats.**

Heteromys Desm., Nouv. Dict. Hist. Nat. Mamm., 1817, p. 313.

Type *Mus anomalus* Thompson.

Dasynotus Wagl., Nat. Syst. Amphib., 1830, p. 21.

Sacomys F. Cuv., Mém. Mus. Hist. Nat. Paris, x, 1823, p. 419.

Xylomys Merr., Proc. Biol. Soc. Wash., xv, 1902, p. 43.

Liomys Merr., Proc. Biol. Soc. Wash., xv, 1902, p. 44.

Subgenus A. **Heteromys.**780. **Heteromys irroratus** Gray.

Heteromys irroratus Gray, Proc. Zoöl. Soc., 1868, p. 205. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 369. Zoöl. Ser.

LA PARADA SPINY RAT.

Type locality. La Parada, State of Oaxaca, Mexico.

Geogr. Distr. State of Oaxaca, Mexico.

781. *Heteromys bulleri* Thomas.

Heteromys bulleri Thomas, Ann. Mag. Nat. Hist., 6th Ser., xi, 1893, p. 330. Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., iv, Pt. 1, 1904, p. 369. Zoöl. Ser.

BULLER'S SPINY RAT.

Type locality. La Laguna, Sierra de Juanacatlan, State of Jalisco, Mexico. Altitude, 7,000 feet.

Geogr. Distr. High elevations, State of Jalisco, Mexico.

782. *Heteromys salvini* Thomas.

Heteromys longicaudatus Alston, Biol. Centr. Amer., Mamm., i, 1880, p. 167, pl. xvii, fig. 2. (nec Gray.)

Heteromys salvini Thomas, Ann. Mag. Nat. Hist., 6th Ser., xi, 1893, p. 331. Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., iv, Pt. 1, 1904, p. 370. Zoöl. Ser.

SALVIN'S SPINY RAT.

Type locality. Dueñas, Guatemala.

Geogr. Distr. Guatemala, Central America.

a. *Heteromys salvini nigrescens* Thomas.

Heteromys salvini nigrescens Thomas, Ann. Mag. Nat. Hist., 6th Ser., xii, 1893, p. 234. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 370. Zoöl. Ser.

BLACK SPINY RAT.

Type locality. Costa Rica.

Geogr. Distr. Costa Rica, Central America.

783. *Heteromys gaumeri* Allen and Chapman.

Heteromys gaumeri Allen and Chapman, Bull. Am. Mus. Nat. Hist. N. Y., 1897, p. 9. Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., iv, Pt. 1, 1904, p. 371, fig. 69. Zoöl. Ser.

GAUMER'S SPINY RAT.

Type locality. Chichen Itza, Yucatan, Mexico.

Geogr. Distr. Yucatan, Mexico.

784. *Heteromys annectens* Merriam.

Heteromys annectens Merr., Proc. Biol. Soc. Wash., xv, 1902, p. 43.

Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 371. Zoöl. Ser.

ALLIED SPINY RAT.

Type locality. Pluma, State of Oaxaca, Mexico.

Geogr. Distr. State of Oaxaca, Mexico.

785. *Heteromys hispidus* Allen.

Heteromys hispidus Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1897, p. 56. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 371. Zoöl. Ser.

HISPID SPINY RAT.

Type locality. Compostella, Territorio de Tepic, Mexico.

Geogr. Distr. Territorio de Tepic, Mexico.

786. *Heteromys desmarestianus* Gray.

Heteromys desmarestianus Gray, Proc. Zoöl. Soc., 1868, p. 204. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 372. Zoöl. Ser.

COBAN SPINY RAT.

Type locality. Coban, Guatemala.

Geogr. Distr. Guatemala, Central America.

787. *Heteromys longicaudatus* Gray.

Heteromys longicaudatus Gray, Proc. Zoöl. Soc., 1868, p. 204. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 372. Zoöl. Ser.

LONG-TAILED SPINY RAT.

Type locality. Mexico.

Geogr. Distr. State of Oaxaca, Mexico. Range unknown.

788. *Heteromys repens* Bangs.

Heteromys repens Bangs, Bull. Mus. Comp. Zoöl., xxxix, 1902, p. 45. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 372. Zoöl. Ser.

BOQUETE SPINY RAT.

Type locality. Boquete, Chiriqui, Panama. Altitude, 4,000 feet.

Geogr. Distr. Panama. Range unknown.

789. *Heteromys goldmani* Merriam.

Heteromys goldmani Merr., Proc. Biol. Soc. Wash., xv, 1902, p. 41. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 373. Zoöl. Ser.

GOLDMAN'S SPINY RAT.

Type locality. Chicharras, State of Chiapas, Mexico.

Geogr. Distr. State of Chiapas, Mexico.

a. *Heteromys goldmani lepturus* Merriam.

Heteromys goldmani lepturus Merr., Proc. Biol. Soc. Wash., xv, 1902, p. 42. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 373. Zoöl. Ser.

SHORT-TAILED SPINY RAT.

Type locality. Mountains near Santo Domingo, State of Oaxaca, Mexico.

Geogr. Distr. State of Oaxaca, Mexico.

790. *Heteromys griseus* Merriam.

Heteromys griseus Merr., Proc. Biol. Soc. Wash., xv, 1902, p. 42.

Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv,

Pt. I, 1904, p. 373. Zoöl. Ser.

GRAY SPINY RAT.

Type locality. Mountains near Tonalá, State of Chiapas, Mexico.

Geogr. Distr. State of Chiapas, Mexico.

791. *Heteromys adpersus* Peters.

Heteromys adpersus Peters, Monatsb. K. Preuss. Akad. Wiss.

Berl., 1874, p. 357. Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., iv, Pt. I, 1904, p. 373. Zoöl. Ser.

SPOTTED SPINY RAT.

Type locality. Panama.

Geogr. Distr. Panama.

Subgenus B. *Liomys*.**792. *Heteromys veræcrucis* (Merriam).**

Liomys veræcrucis Merr., Proc. Biol. Soc. Wash., xv, 1902, p. 47.

Heteromys veræcrucis Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., iv, Pt. I, 1904, p. 379. Zoöl. Ser.

VERA CRUZ SPINY MOUSE.

Type locality. San Andreas Tuxtla, State of Vera Cruz, Mexico.

Geogr. Distr. State of Vera Cruz, Mexico.

793. *Heteromys obscurus* (Merriam).

Liomys obscurus Merr., Proc. Biol. Soc. Wash., xv, 1902, p. 48.

Heteromys obscurus Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., iv, Pt. I, 1904, p. 379. Zoöl. Ser.

DUSKY SPINY MOUSE.

Type locality. Carrizal, State of Vera Cruz, Mexico.

Geogr. Distr. State of Vera Cruz, Mexico.

794. *Heteromys phæurus* (Merriam).

Liomys phæura! Merr., Proc. Biol. Soc. Wash., xv, 1902, p. 48.

Heteromys phæurus Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., iv, Pt. I, 1904, p. 379. Zoöl. Ser.

DARK-TAILED SPINY MOUSE.

Type locality. Pinotepa, State of Oaxaca, Mexico.

Geogr. Distr. State of Oaxaca, Mexico.

795. *Heteromys orbitalis* (Merriam).

Liomys orbitalis Merr., Proc. Biol. Soc. Wash., xv, 1902, p. 48.

Heteromys orbitalis Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., iv, Pt. 1, 1904, p. 380. Zoöl. Ser.

CATEMACO SPINY MOUSE.

Type locality. Catemaco, State of Vera Cruz, Mexico.

Geogr. Distr. State of Vera Cruz, Mexico.

796. *Heteromys crispus* (Merriam).

Liomys crispus Merr., Proc. Biol. Soc. Wash., xv, 1902, p. 49.

Heteromys crispus Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., iv, Pt. 1, 1904, p. 380. Zoöl. Ser.

CURLY SPINY MOUSE.

Type locality. Tonalá, State of Chiapas, Mexico.

Geogr. Distr. State of Chiapas, Mexico.

a. *Heteromys crispus setosus* (Merriam).

Liomys crispus setosus Merr., Proc. Biol. Soc. Wash., xv, 1902,
p. 49.

Heteromys crispus setosus Elliot, Mamm. Middle Amer. & W. In-
dies, F. C. M. Pub., iv, Pt. 1, 1904, p. 380. Zoöl. Ser.

HUEHUETAN SPINY MOUSE.

Type locality. Huehuetan, State of Chiapas, Mexico.

Geogr. Distr. State of Chiapas, Mexico.

797. *Heteromys heterothrix* (Merriam).

Liomys heterothrix Merr., Proc. Biol. Soc. Wash., xv, 1902, p. 50.

Heteromys heterothrix Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., iv, Pt. 1, 1904, p. 380. Zoöl. Ser.

HONDURAS SPINY MOUSE.

Type locality. San Pedro Sula, Honduras.

Geogr. Distr. Honduras, Central America.

798. *Heteromys paralius* Elliot.

Heteromys paralius Elliot, Pub. Field Columb. Mus., iii, 1903,
p. 233. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., iv, Pt. 1, 1904, p. 381. Zoöl. Ser.

LITTORAL SPINY MOUSE.

Type locality. San Carlos, State of Vera Cruz, Mexico.

Geogr. Distr. Coast district of the State of Vera Cruz, Mexico.

799. *Heteromys albolimbatus* Gray.

Heteromys albolimbatus Gray, Proc. Zoöl. Soc., 1868, p. 205.

Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv,
Pt. 1, 1904, p. 375, fig. 70. Zoöl. Ser.

GRAY'S SPINY MOUSE.

Type locality. Undetermined. "La Parda, Mexico?"

Geogr. Distr. Undetermined. Specimens from the States of Michoacan and Morelos.

800. *Heteromys canus* (Merriam).

Liomys canus Merr., Proc. Biol. Soc. Wash., xv, 1902, p. 44.

Heteromys canus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M.

Pub., iv, Pt. 1, 1904, p. 375. Zoöl. Ser.

HOARY SPINY MOUSE.

Type locality. Parral, State of Chihuahua, Mexico.

Geogr. Distr. State of Chihuahua, Mexico.

801. *Heteromys alleni* Coues.

Heteromys alleni Coues, Bull. Mus. Comp. Zoöl., viii, 1881, p. 187.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., ii, 1901, p. 256,

fig. 58. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., iv, Pt. 1, 1904, p. 376, fig. 100. Zoöl. Ser.

ALLEN'S SPINY MOUSE.

Type locality. Hacienda Angostura, Rio Verde, State of San Luis Potosi, Mexico.

Geogr. Distr. State of San Luis Potosi, Mexico.

802. *Heteromys torridus* (Merriam).

Liomys torridus Merr., Proc. Biol. Soc. Wash., xv, 1902, p. 45.

Heteromys torridus Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., iv, Pt. 1, 1904, p. 376. Zoöl. Ser.

TORRID SPINY MOUSE.

Type locality. Cuicatlan, State of Oaxaca, Mexico.

Geogr. Distr. Known only from the type locality.

a. *Heteromys torridus minor* (Merriam).

Liomys torridus minor Merr., Proc. Biol. Soc. Wash., xv, 1902, p. 45.

Heteromys torridus minor Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 377. Zoöl. Ser.

SMALLEST SPINY MOUSE.

Type locality. Huajuapam, State of Oaxaca.

Geogr. Distr. State of Oaxaca, Mexico.

803. *Heteromys exiguus* Elliot.

Heteromys exiguus Elliot, Pub. Field Columb. Mus., iii, 1903, p. 146. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., iv, Pt. 1, 1904, p. 377. Zoöl. Ser.

LITTLE SPINY MOUSE.

Type locality. Puente de Ixtla, State of Morelos, Mexico.

Geogr. Distr. State of Morelos, Mexico.

804. *Heteromys plantinarensis* Merriam.

Liomys plantinarensis Merr., Proc. Biol. Soc. Wash., xv, 1902, p. 46.

Heteromys plantinarensis Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 377. Zoöl. Ser.

PLANTINAR SPINY MOUSE.

Type locality. Plantinar, State of Jalisco, Mexico.

Geogr. Distr. State of Jalisco, Mexico.

805. *Heteromys pictus* Thomas.

Heteromys pictus Thomas, Ann. Mag. Nat. Hist., 6th Ser., xii, 1893, p. 233. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 377. Zoöl. Ser.

PAINTED SPINY MOUSE.

Type locality. Mineral San Sebastian, State of Jalisco, Mexico. Altitude, 4,300 feet.

Geogr. Distr. State of Jalisco, Mexico.

a. *Heteromys pictus rostratus* (Merriam).

Liomys pictus rostratus Merr., Proc. Biol. Soc. Wash., xv, 1902, p. 46.

Heteromys pictus rostratus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 378. Zoöl. Ser.

LONG-NOSED SPINY MOUSE.

Type locality. Ometepepec, State of Guerrero, Mexico.

Geogr. Distr. State of Guerrero, Mexico.

b. *Heteromys pictus isthmius* (Merriam).

Liomys pictus isthmius Merr., Proc. Biol. Soc. Wash., xv, 1902, p. 46.

Heteromys pictus isthmius Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 378. Zoöl. Ser.

ISTHMIAN SPINY MOUSE.

Type locality. Tehuantepec, State of Oaxaca, Mexico.

Geogr. Distr. State of Oaxaca, Mexico.

806. *Heteromys parviceps* (Goldman).

Liomys parviceps Goldm., Proc. Biol. Soc. Wash., xvii, 1904, p. 82.

Heteromys parviceps Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 378. Zoöl. Ser.

URUAPAN SPINY MOUSE.

Type locality. Uruapan, State of Michoacan, Mexico.

Geogr. Distr. State of Michoacan, Mexico.

807. *Heteromys sonorana* (Merriam).

Liomys sonorana Merr., Proc. Biol. Soc. Wash., xv, 1902, p. 47.

Heteromys sonorana Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., iv, Pt. I, 1904, p. 379. Zoöl. Ser.

SONORA SPINY MOUSE.

Type locality. Alamos, State of Sonora, Mexico.

Geogr. Distr. State of Sonora, Mexico.

808. *Heteromys texensis* (Merriam).

Liomys texensis Merr., Proc. Biol. Soc. Wash., xv, 1902, p. 44.

TEXAN SPINY MOUSE.

Type locality. Brownsville, Cameron County, Texas.

Geogr. Distr. Southern Texas. Range unknown.

Genl. Char. Size small; hind foot with five tubercles on soles; rostrum narrow; interparietal small, oval.

Color. Above grayish brown or drab, grizzled with red and blackish hairs, red predominating; under parts yellowish white; ears edged with whitish; hind feet and under side of tail white.

Measurements. Total length, 231; tail vertebræ, 114; hind foot, 30.

Subgenus C. *Xylomys*.**809. *Heteromys nelsoni* (Merriam).**

Xylomys nelsoni Merr., Proc. Biol. Soc. Wash., xv, 1902, p. 43.

Heteromys nelsoni Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., iv, Pt. I, 1904, p. 381. Zoöl. Ser.

NELSON'S SPINY MOUSE.

Type locality. Pinabete, State of Chiapas, Mexico.

Geogr. Distr. State of Chiapas, Mexico. Range unknown.

Family VII. ***Zapodidae. Jumping Mice.**Subfamily I. **Zapodinae.**Genus 100. **Zapus.**

Zapus Coues, Bull. U. S. Geol. Surv. Terr., 2d Ser., 1, 1875, p. 253.

Type *Dipus hudsonius* Zimmermann.

*See Lyon, Proc. U. S. Nat. Mus., 1901, pp. 659-668, pls. 25-27, in which paper the author decides that *Zapus* and *Sminthus* should be associated in one group as the family *Zapodidae*.

- Dipus* Zimm., Geog. Gesch. Mensch. u. vierfüß. Thiere, II, 1780, p. 358. (Old World Species.)
Gerbillus Rafin., Précis des Découv. Somologiques, 1814, p. 14. (nec Demarest, 1804.)
Meriones F. Cuv., Dents de Mamm., 1823, pp. 187, 256. (nec Illiger, 1811.)
Jaculus Wagl., Nat. Syst. Amphib., 1830, p. 23. (nec Erxleben, 1777.)
Napæozapus Preble, N. Am. Faun., No. 15, 1899, p. 33.
Eozapus Preble, N. Am. Faun., No. 15, 1899, p. 37.

Subgenus A. *Zapus*.

810. *Zapus hudsonius* (Zimmermann).

- Dipus hudsonius* Zimm., Geog. Gesch. Mensch. u. vierfüß. Thiere, II, 1780, p. 358.
Dipus labradorius Kerr, Anim. Kingd., 1792, p. 276.
Dipus canadensis Davies, Trans. Linn. Soc. Lond., IV, 1798, p. 157.
Jaculus americanus Wagl., Nat. Syst. Amphib., 1830, p. 23.
Meriones microcephalus Harlan, Proc. Zool. Soc., 1839, p. 1.
Meriones acadicus Dawson, Edinb. New Philos. Journ., N. S., III, 1856, p. 2.
Zapus hudsonius hardyi Batchelder, Proc. N. Eng. Zool. Club, I, 1899, p. 5.
Zapus hudsonius Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 257, fig. 59. Zool. Ser.

HUDSON BAY JUMPING MOUSE.

Type locality. Hudson Bay.

Geogr. Distr. Hudson Bay south to New Jersey; mountains of North Carolina west to Iowa and Missouri, and northwest to Alaska.

a. *Zapus hudsonius ladas* Bangs.

- Zapus hudsonius ladas* Bangs, Proc. N. Eng. Zool. Club, I, 1899, p. 10. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 258. Zool. Ser.

LABRADOR JUMPING MOUSE.

Type locality. Rigoulette, Hamilton Inlet, Labrador.

Geogr. Distr. Eastern Quebec, north to Hamilton Inlet, Labrador.

b. *Zapus hudsonius alascensis* Merriam.

- Zapus hudsonius alascensis* Merr., Proc. Biol. Soc. Wash., XI, 1897, p. 223. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 259. Zool. Ser.

ALASKA JUMPING MOUSE.

Type locality. Yakutat Bay, Alaska.

Geogr. Distr. Yakutat Bay, north to Yukon River. Limits unknown.

c. *Zapus hudsonius americanus* (Barton).

Dipus americanus Barton, Trans. Am. Philos. Soc., iv, 1799, p. 115.

Zapus hudsonius americanus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 259. Zoöl. Ser.

CAROLINIAN JUMPING MOUSE.

Type locality. Near Philadelphia, Pennsylvania.

Geogr. Distr. Raleigh, North Carolina, north to southern Connecticut and valley of the lower Hudson.

d. *Zapus hudsonius campestris* Preble.

Zapus hudsonius campestris Preble, N. Am. Faun., No. 15, 1899, p. 20. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 259. Zoöl. Ser.

PRAIRIE JUMPING MOUSE.

Type locality. Bear Lodge Mountains, Crook County, Wyoming.

Geogr. Distr. Great Plains from Manitoba southward to Nebraska and westward to Colorado and Wyoming.

811. *Zapus tenellus* Merriam.

Zapus tenellus Merr., Proc. Biol. Soc. Wash., xi, 1897, p. 103. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 259. Zoöl. Ser.

KAMLOOPS JUMPING MOUSE.

Type locality. Kamloops, British Columbia, Canada.

Geogr. Distr. Known only from the type locality.

812. *Zapus princeps* Allen.

Zapus princeps Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1893, p. 71. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 260. Zoöl. Ser.

ROCKY MOUNTAIN JUMPING MOUSE.

Type locality. Florida, La Plata County, Colorado.

Geogr. Distr. Rocky Mountain region from northern New Mexico northward to Henry House, Alberta, Canada.

a. *Zapus princeps minor* Preble.

Zapus princeps minor Preble, N. Am. Faun., No. 15, 1899, p. 23. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 260. Zoöl. Ser.

SASKATCHEWAN JUMPING MOUSE.

Type locality. Wingard, near Carlton House, Saskatchewan, Canada.

Geogr. Distr. Plains of Saskatchewan.

b. *Zapus princeps oregonus* Preble.

Zapus princeps oregonus Preble, N. Am. Faun., No. 15, 1899, p.

24. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p.

260. Zoöl. Ser.

BLUE MOUNTAINS JUMPING MOUSE.

Type locality. Elgin, Blue Mountains, Union County, Oregon.

Geogr. Distr. Blue Mountains, Oregon.

813. *Zapus major* Preble.

Zapus major Preble, N. Am. Faun., No. 15, 1899, p. 24. Elliot,

Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 260. Zoöl.

Ser.

WARNER MOUNTAIN JUMPING MOUSE.

Type locality. Warner Mountains, Lake County, Oregon.

Geogr. Distr. Known only from the type locality.

814. *Zapus nevadensis* Preble.

Zapus nevadensis Preble, N. Am. Faun., No. 15, 1899, p. 25.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 261.

Zoöl. Ser.

NEVADA JUMPING MOUSE.

Type locality. Ruby Mountains, Elko County, Nevada.

Geogr. Distr. Known only from the type locality.

815. *Zapus trinotatus* Rhoads.

Zapus trinotatus Rhoads, Proc. Acad. Nat. Scien. Phil., 1894,

p. 421. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901,

p. 261. Zoöl. Ser.

Zapus imperator Elliot, Pub. Field Columb. Mus., 1, 1899, p. 228.

Zoöl. Ser.

NORTHWEST JUMPING MOUSE.

Type locality. Lulu Island, mouth of the Fraser River, British Columbia.

Geogr. Distr. Coast region of southern British Columbia, Canada; Washington, Oregon west of Cascade Mountains, northern California to Humboldt Bay.

a. *Zapus trinotatus alleni* (Elliot).

Zapus alleni Elliot, Pub. Field Columb. Mus., 1, 1898, p. 212.

Zoöl. Ser.

Zapus trinotatus alleni Elliot, Syn. N. Am. Mamm., F. C. M. Pub.,
11, 1901, p. 261. Zoöl. Ser.

ALLEN'S JUMPING MOUSE.

Type locality. Pyramid Peak, Lake Tahoe, Eldorado County, California.

Geogr. Distr. Mount Shasta, southward in the Sierra Nevada to Mammoth, and north fork of Kern River, California.

816. *Zapus montanus* (Merriam).

Zapus trinotatus montanus Merr., Proc. Biol. Soc. Wash., XI, 1897,
p. 104. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901,
p. 262. Zoöl. Ser.

MOUNTAIN JUMPING MOUSE.

Type locality. Crater Lake, Mount Mazama, Klamath County, Oregon.

Geogr. Distr. Cascade Range in Oregon.

817. *Zapus orarius* Preble.

Zapus orarius Preble, N. Am. Faun., No. 15, 1899, p. 29. Elliot,
Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 262. Zoöl. Ser.

COAST JUMPING MOUSE.

Type locality. Point Reyes, Marin County, California.

Geogr. Distr. Coast of California from Point Reyes, Marin County, north to Mad River, Humboldt County. Limits of range unknown.

818. *Zapus pacificus* Merriam.

Zapus pacificus Merr., Proc. Biol. Soc. Wash., XI, 1897, p. 104.
Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 262.
Zoöl. Ser.

PACIFIC JUMPING MOUSE.

Type locality. Prospect, Rogue River Valley, Jackson County, Oregon.

Geogr. Distr. Interior valleys of southwestern Oregon and northwestern California.

819. *Zapus saltator* Allen.

Zapus saltator Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1899, p. 3.
Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 262.
Zoöl. Ser.

STICKEEN JUMPING MOUSE.

Type locality. Telegraph Creek, Stikine River, British Columbia, Canada.

Geogr. Distr. Telegraph Creek, south to mouth of Skeena River and Tschimshian Peninsula. Limits of range unknown.

Subgenus B. *Napæozapus*.

820. *Zapus insignis* Miller.

Meriones labradorius Dawson, Edinb. New Philos. Journ., N. S., III, 1856, p. 2. (nec Kerr.)

Jaculus hudsonius Baird, N. Am. Mamm., 1857, p. 467. (Part.)

Zapus insignis Miller, Amer. Nat., xxv, 1891, p. 742. Elliot, Syn.

N. Am. Mamm., F. C. M. Pub., II, 1901, p. 263, fig. 60. Zoöl. Ser.

WOODLAND JUMPING MOUSE.

Type locality. Restigouche River, New Brunswick, Canada.

Geogr. Distr. Eastern Canada, south to western Maryland.

a. Zapus insignis roanensis Preble.

Zapus insignis roanensis Preble, N. Am. Faun., No. 15, 1899, p. 35.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 263.

Zoöl. Ser.

ROAN MOUNTAIN JUMPING MOUSE.

Type locality. Roan Mountain, Mitchell County, North Carolina.

Geogr. Distr. Known only from the type locality.

b. Zapus insignis abietorum Preble.

Zapus insignis abietorum Preble, N. Am. Faun., No. 15, 1899,

p. 36. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901,

p. 264. Zoöl. Ser.

NORTHERN WOODLAND JUMPING MOUSE.

Type locality. Peninsula Harbor, north shore of Lake Superior, Ontario, Canada.

Geogr. Distr. Eastern Canada. Range unknown.

Family VIII. Octodontidæ. The Octodonts.

Subfamily I. Loncherinæ.

Genus 101. Loncheres.

Loncheres Illig., Prodr. Syst. Mamm. et Av., 1811, p. 90. Type

Myoxus chrysurus Zimmermann = *Echinomys cristatus* Desmarest.

Lonchetes Billberg, Syn. Faun. Scandinav., I, 1828, Mamm., Conspectus A.

821. *Loncheres labilis* Bangs.

Loncheres labilis Bangs, Amer. Nat., xxxv, 1901, p. 638. Elliot,

Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1,

1904, p. 384, figs. 71, LI. Zoöl. Ser.

GLIDING SPINY RAT.

Type locality. San Miguel Island, Bay of Panama.

Geogr. Distr. Known only from the type locality.

Genus 102. **Proechinomys.**

Proechimys (*sic*) Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1899, p. 264.

Type Echimys! trinitatis Allen & Chapman.

Echimys (sic) Geoff. St. Hilare, Ann. Scien. Nat., 2me Sér., x, 1838, p. 125. (nec G. Cuvier, 1809.)

822. Proechinomys semispinosus (Tomes).

Echimys! semispinosus Tomes, Proc. Zool. Soc., 1860, p. 265.

Proechinomys semispinosus Elliot, Mamm. Middle Amer. & W.

Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 386. Zool. Ser.

SHORT-SPINED RAT.

Type locality. Ecuador.

Geogr. Distr. Nicaragua, south to Ecuador.

823. Proechinomys centralis (Thomas).

Echinomys centralis Thomas, Ann. Mag. Nat. Hist., 6th Ser., XVIII, 1896, p. 312.

Proechinomys centralis Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., IV, Pt. 1, 1904, p. 386, fig. 72. Zool. Ser.

NICARAGUAN SPINY RAT.

Type locality. San Emilio, north end of Lake Nicaragua, Nicaragua.

Geogr. Distr. Costa Rica, Nicaragua, Central America.

a. Proechinomys centralis chiriquinus Thomas.

Proechimys! centralis chiriquinus Thomas, Ann. Mag. Nat. Hist., 7th Ser., v, 1900, p. 220.

Proechinomys centralis chiriquinus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 387. Zool. Ser.

BOGAVA SPINY RAT.

Type locality. Bogava, Chiriqui, Panama. Altitude, 800 feet.

Geogr. Distr. Panama, Central America.

b. Proechinomys centralis panamensis Thomas.

Proechimys! centralis panamensis Thomas, Ann. Mag. Nat. Hist., 7th Ser., v, 1900, p. 220.

Proechinomys centralis panamensis Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 387. Zool. Ser.

PANAMA SPINY RAT.

Type locality. Savanna near City of Panama, Panama.

Geogr. Distr. Panama, Central America.

824. *Proechinomys burrus* Bangs.

Proechimys! burrus Bangs, Amer. Nat., xxxv, 1901, p. 640.

Proechinomys burrus Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., IV, Pt. I, 1904, p. 388. Zoöl. Ser.

SAN MIGUEL SPINY RAT.

Type locality. San Miguel Island, Bay of Panama.

Geogr. Distr. Known only from the type locality.

Subfamily II. **Capromyinae.**Genus 103. **Capromys.**

Capromys Desmarest, Mém. Soc. Hist. Nat. Paris, I, 1822, p. 43.

Type *Capromys fournieri* Desmarest = *Isodon pilorides* Say.

Geocapromys Chapman, Bull. Am. Mus. Nat. Hist. N. Y., 1901,
p. 314.

Procapromys Chapman, Bull. Am. Mus. Nat. Hist. N. Y., 1901,
p. 322.

Subgenus A. *Capromys*.825. **Capromys pilorides** (Say).

Isodon pilorides Say, Journ. Acad. Nat. Scien. Phil., II, 1822, p. 333,
fig.

Capromys fournieri Desm., Mém. Soc. Hist. Nat. Paris, 1822, p. 43,
pl. I.

Capromys quemi Fisch., Syn. Mamm., 1829, p. 312.

Capromys pilorides Fisch., Syn. Mamm., 1830, p. 313. Elliot,
Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I,
1904, p. 390, figs. 73, LII. Zoöl. Ser.

HAIRY HUTIA.

Type locality. "South America, or one of the West Indian
Islands." Cuba?

Geogr. Distr. Cuba.

826. **Capromys melanurus** Poey.

Capromys melanurus Poey, Monatsb. K. Preuss. Akad. Wiss.
Berl., 1864, p. 384. Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., IV, Pt. I, 1904, p. 391. Zoöl. Ser.

BLACK-TAILED HUTIA.

Type locality. Manzanillo, Cuba.

Geogr. Distr. Island of Cuba.

827. *Capromys prehensilis* Poeppig.

Capromys prehensilis Poeppig, Journ. Acad. Nat. Scien. Phil., 1824, p. 11. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 391. Zool. Ser.

PREHENSILE-TAILED HUTIA.

Type locality. Southern Cuba.

Geogr. Distr. Island of Cuba.

a. *Capromys prehensilis gundlachi* Chapman.

Capromys prehensilis gundlachi Chapman, Bull. Am. Mus. Nat. Hist. N. Y., 1901, p. 317. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 392. Zool. Ser.

GUNDLACH'S HUTIA.

Type locality. Nueva Gerona, Isle of Pines, near Cuba.

Geogr. Distr. Known only from the type locality.

828. *Capromys elegans* Latorre.

Capromys elegans Latorre, Bol. Soc. Españ. Hist. Nat. Madrid, 1, 1901, p. 372. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 392. Zool. Ser.

LANCEOLATE-SPOT HUTIA.

Type locality. Cuba?

Geogr. Distr. Range unknown.

Subgenus B. *Geocapromys*.829. *Capromys browni* Fischer.

Capromys brownii Fisch., Syn. Mamm., Addend., 1830, p. 389. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 393, fig. 74. Zool. Ser.

SHORT-TAILED HUTIA.

Type locality. Jamaica.

Geogr. Distr. Known only from the type locality.

830. *Capromys thoracatus* (True).

Capromys brachyurus thoracatus True, Proc. U. S. Nat. Mus., 1888, p. 469. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 394. Zool. Ser.

WHITE-BANDED HUTIA.

Type locality. Little Swan Island, Gulf of Honduras.

Geogr. Distr. Known from the type locality only.

831. Capromys ingrahami Allen.

Capromys ingrahami Allen, Bull. Am. Mus. Nat. Hist., 1891, p. 329. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV., Pt. 1, 1904, p. 394. Zoöl. Ser.

INGRAHAM'S HUTIA.

Type locality. Plana Key, Bahama Islands.

Geogr. Distr. Bahama Islands.

Genus 104. **Plagiodontia.**

Plagiodontia F. Cuv., Ann. Scien. Nat., 2me Sér., VI, 1836, p. 347. Zoöl. *Type Plagiodontia aedium* F. Cuvier.

832. Plagiodontia aedium F. Cuvier.

Plagiodontia aedium F. Cuvier, Ann. Scien. Nat., 2me Sér., VI, 1836, p. 347. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 396, figs. 75, LIII. Zoöl. Ser.

HAITIAN HUTIA.

Type locality. Island of Haiti.

Geogr. Distr. Island of Haiti.

Family IX. **Erethizontidæ.**Subfamily I. **Erethizontinæ.**Genus 105. **Erethizon. Long-spined Porcupines.**

Erethizon F. Cuv., Mém. Mus. Hist. Nat. Paris, IX, 1822, p. 436.

Type Hystrix dorsatus Linnæus.

Echinothrix Brookes, Cat. Anat. & Zoöl. Mus. J. Brookes, Lond., 1828, p. 54.

Echinoprocta Gray, Proc. Zoöl. Soc., 1865, pp. 321-322, desc.

833. Erethizon epixanthum Brandt.

Erethizon epixanthum Brandt, Mém. Acad. Imp. Scien. St. Pétersb., 6th Ser., 1835, p. 390, pls. 1, 9. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 265. Zoöl. Ser.

Erethizon epixanthum Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 397, figs. 76, LIV. Zoöl. Ser.

WESTERN PORCUPINE.

Type locality. California ? Unalaska ?

Geogr. Distr. State of Sonora, Mexico, into New Mexico, eastward to Missouri, west to the Pacific and north to Alaska and the limit of trees.

a. Erethizon epixanthum couesi Mearns.

Erethizon epixanthus couesi Mearns, Proc. U. S. Nat. Mus., 1897, p. 723. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 265. Zoöl. Ser.

COUES' TREE PORCUPINE.

Type locality. Fort Whipple, Yavapai County, Arizona.

Geogr. Distr. Arizona and New Mexico.

b. Erethizon epixanthum myops Merriam.

Erethizon epixanthus myops Merr., Proc. Wash. Acad. Scien., 11, 1900, p. 27. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 266. Zoöl. Ser.

ALASKA PORCUPINE.

Type locality. Portage Bay, Alaska Peninsula, Alaska.

Geogr. Distr. Range unknown.

c. Erethizon epixanthum nigrescens Allen.

Erethizon epixanthus! nigrescens Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1903, p. 558.

DUSKY PORCUPINE.

Type locality. Shesley River, British Columbia, Canada.

Genl. Char. Skull markedly depressed at fronto-parietal region; lateral border of interorbital region forming a high obtuse knob.

Color. Front and sides of head, black, lightest on the nape; sides of lower back and thighs with tips of hairs yellowish; underparts sooty black; long hairs on top of head and nape broadly tipped with pale yellow; on shoulders and lower back the long hairs are mostly black; long hairs on thighs and sides of tail broadly tipped with yellowish white; spines white tipped with black except on the shoulders where they are plumbeous at tips.

Measurements. Total length, 740; tail vertebræ, 210; hind foot, 90; ear, 27. Skull: total length, 105; basal length, 99; zygomatic breadth, 67; mastoid breadth, 42; upper molar series, 26.

834. *Erethizon dorsatum* (Linnaeus).

Hystrix dorsata Linn., Syst. Nat., 1, 1758, p. 57; 1, 1766, p. 76.

Hystrix pilosus Richards., Faun. Bor. Amer., Mamm., 1, 1829, p. 214. (nec Peale.)

Hystrix hudsonius De Kay, N. Y. Zoöl., 1, 1842, p. 27, pl. xxv, fig. 1, pl. viii, 2, a. b. c.

Erethizon dorsatus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 264, pl. XXIV. Zoöl. Ser.

CANADIAN TREE PORCUPINE.

Type locality. Eastern Canada.

Geogr. Distr. Canada, west to the Saskatchewan Plains, south to central Maine and northern New Hampshire and west of the Connecticut River nearly to the border of Massachusetts. Formerly it extended along the Alleghanies through Pennsylvania into Virginia and possibly west to Kentucky. North in Canada it goes to the Barren Grounds, and the limit of trees.

a. *Erethizon dorsatum picinum* Bangs.

Erethizon dorsatus picinus Bangs, Proc. N. E. Zoöl. Club, II, 1900, p. 37. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, Append., p. 431. Zoöl. Ser.

LABRADOR PORCUPINE.

Type locality. Lance au Loop, Strait of Belle Isle, Labrador.

Geogr. Distr. Labrador. Range unknown.

Genus 106. **Coendu.**

Coendou Lacép., Disc. d'ouvert. et de cloture du Cours Hist. Nat., Suppl., 1799, p. 11. Type *Hystrix prehensilis* Linnæus.

Eucrius Fisch., Mém. Soc. Imp. Mosc., v, 1817, pp. 372, 411.

Sinetheres (sic) F. Cuv., Mém. Mus. Hist. Nat. Paris, IX, 1822, p. 433.

Laboura Billb., Syn. Faun. Scandinav., Mamm., I, 1828, Consp. A.

Cercolabes Brandt, Mamm. Exot. Nov., in Mém. Acad. Imp. St.

Pétersb., Sér. 3, III, 1835, p. 55.

835. Coendu pallidum (Waterhouse).

Cercolabes pallidus Waterh., Nat. Hist. Mamm., II, 1848, p. 434.

Coendu pallidum Elliot, Mamm. Middle Amer. & W. Indies, F.

C. M. Pub., IV, Pt. 1, 1904, p. 401. Zoöl. Ser.

LIGHT-COLORED PORCUPINE.

Type locality. "West Indies."

Geogr. Distr. Undetermined.

836. Coendu rothschildi Thomas.

Coendou rothschildi Thomas, Ann. Mag. Nat. Hist., 7th Ser., x, 1902, p. 169.

Coendu rothschildi Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 401. Zoöl. Ser.

ROTHSCHILD'S TREE PORCUPINE.

Type locality. Sevilla Island, off Chiriqui, Panama.

Geogr. Distr. Known only from the type locality.

837. *Coendu mexicanum* (Kerr).¹

Hystrix mexicana Kerr, Anim. Kingd., 1792, p. 214.

Hystrix novæ-hispaniæ Briss., Règn. Anim., 1756, p. 127.

Hystrix prehensilis Schreb., Säugth., IV, 1792, p. 603.

Coendu mexicanum Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., IV, Pt. 1, 1904, p. 402, figs. 77, LV. Zoöl. Ser.

PREHENSILE-TAILED PORCUPINE.

Type locality. Mountains of Mexico.

Geogr. Distr. Mirador, State of Vera Cruz, Mexico, south to Costa Rica, Central America.

a. *Coendu mexicanum yucatanicæ* Thomas.

Coendou mexicanum yucatanicæ Thomas, Ann. Mag. Nat. Hist., 7th Ser., x, 1902, p. 249.

Coendu mexicanum yucatanicæ Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 402. Zoöl. Ser.

YUCATAN TREE PORCUPINE.

Type locality. Izamal, Yucatan.

Geogr. Distr. Yucatan. Range unknown.

838. *Coendu lænatum* Thomas.

Coendou lænatus Thomas, Ann. Mag. Nat. Hist., 7th Ser., XI, 1903, p. 381.

Coendu lænatum Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 402. Zoöl. Ser.

CHIRIQUI PORCUPINE.

Type locality. Boquete, Chiriqui, Panama. Altitude, 5,000 feet.

Geogr. Distr. Panama. Range unknown.

Family X. **Agoutiidae. Agoutis, Paca.**Genus 107. **Dasyprocta. Agoutis.**

Dasyprocta Illig., Prodr. Syst. Mamm. et Av., 1811, p. 93. Type undetermined.

Cutia Liais, Climats, Géol., Faun., Geog. et Bot. Brésil, 1872, p. 534.

Myoprocta Thomas, Ann. Mag. Nat. Hist., 7th Ser., XII, 1903, p. 464.

839. *Dasyprocta punctata* Gray.

Dasyprocta punctata Gray, Ann. Mag. Nat. Hist., 1st Ser., x, 1842, p. 264. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 405. Zoöl. Ser.

SPOTTED AGOUTI. *Cotusa* in Guatemala.

Type locality. "South America."

Geogr. Distr. Guatemala and Costa Rica, Central America.

840. *Dasyprocta ruatanica* Thomas.

Dasyprocta ruatanica Thomas, Ann. Mag. Nat. Hist., 7th Ser., VIII, 1901, p. 272. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 405. Zoöl. Ser.

RUATAN ISLAND AGOUTI.

Type locality. Island of Ruatan, Bay of Honduras.

Geogr. Distr. Known only from the type locality.

841. *Dasyprocta mexicana* Saussure.

Dasyprocta mexicana Sauss., Rev. Mag. Zoöl., 2me Sér., XII, 1860, p. 53. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 405, fig. 78. Zoöl. Ser.

MEXICAN AGOUTI.

Type locality. "Hot zone of Mexico." State of Vera Cruz?

Geogr. Distr. Range unknown.

842. *Dasyprocta callida* Bangs.

Dasyprocta callida Bangs, Amer. Nat., xxxv, 1901, p. 635. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 406. Zoöl. Ser.

CUNNING AGOUTI.

Type locality. San Miguel Island, Bay of Panama.

Geogr. Distr. Known only from the type locality.

843. *Dasyprocta isthmica* Alston.

Dasyprocta isthmica Alston, Proc. Zoöl. Soc., 1876, p. 347. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 407, fig. LVI. Zoöl. Ser.

ISTHMIAN AGOUTI.

Type locality. Colon, Colombia.

Geogr. Distr. Unknown; probably Costa Rica to Colombia.

844. *Dasyprocta coibæ* Thomas.

Dasyprocta coibæ Thomas, Novitat. Zool., IX, 1902, p. 136. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 407. Zoöl. Ser.

COIBA AGOUTI.

Type locality. Coiba Island, West Coast of Panama.

Geogr. Distr. Known only from the type locality.

845. *Dasyprocta cristata* Desmarest.

Dasyprocta cristata Desm., Nouv. Dict. Hist. Nat., I, 1816, p. 213.

Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV,

Pt. I, 1904, p. 407. Zoöl. Ser.

Dasyprocta antillensis Sclat., Proc. Zoöl. Soc., 1874, p. 666.

CRESTED AGOUTI.

Type locality. "Surinam." Erroneous.

Geogr. Distr. Islands of St. Vincent, St. Lucia, and St. Thomas, West Indies.

Genus 108. *Agouti*. *Paca*.

Agouti Lacép., Tableau des Divis. Sous-divis. Ordres et Genres

Mamm., Suppl. Disc. d'ouvert et de cloture du Cours d'Hist.

Nat., etc., 1799, p. 11. Type *Mus paca* Linnæus.

Coelogenus (sic) F. Cuv., Ann. Mus. Hist. Nat. Paris, x, 1807, p. 203.

Paca Fisch., Zoogn., III, 1814, p. 85.

Osteopera Harlan, Faun. Amer., 1825, p. 126.

Genyscælus Liais, Climats, Géol., Faun., Geogr. et Bot. Brésil, 1872,

p. 537.

Agouti paca virgata Bangs.

Agouti paca virgatus Bangs, Bull. Mus. Comp. Zoöl., xxxix, 1902,

p. 47.

Agouti paca virgata Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., IV, Pt. I, 1904, p. 408, pls. xxxviii, xxxix,

and fig. LVII. Zoöl. Ser.

CENTRAL AMERICAN PACA.

Type locality. Divala, Chiriqui, Panama.

Geogr. Distr. Panama, Central America.

Family XI. *Ochotonidae*.Genus 109. *Ochotona*. *Pikas*, *Chief Hares*.

Ochotona Link, Zool. Beytr. Nat., I, Pt. II, 1795, p. 74. Type undetermined.

Lagomys Storr, Prodr. Meth. Mamm., 1780, pp. 39-40, tab. B.

Ogotoma Gray, Ann. Mag. Nat. Hist., 3d Ser., xx, 1867, p. 220.

846. Ochotona princeps (Richardson).

Lepus princeps Richards., Zoöl. Journ., III, 1828, p. 520.

Ochotona princeps Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 267, fig. 61. Zoöl. Ser.

CHIEF PIKA.

Type locality. "Stony places in the Rocky Mountains"; probably in northern British Columbia, Canada.

Geogr. Distr. Rocky Mountains in British Columbia, to the south branch of the Mackenzie River.

847. Ochotona schisticeps (Merriam).

Lagomys schisticeps Merr., N. Am. Faun., No. 2, 1889, p. 11.

Ochotona schisticeps Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 267. Zoöl. Ser.

GRAY-HEADED PIKA.

Type locality. Donner, Placer County, California.

Geogr. Distr. Sierra Nevada Mountains, California.

848. Ochotona cinnamomea Allen.

Ochotona cinnamomea Allen, Bull. Brookl. Inst. Arts & Scien., I, No. 5, 1905, p. 121.

CINNAMON CHIEF HARE.

Type locality. Briggs Meadows, Beaver Range, Beaver County, Utah. Altitude, 11,000 feet.

Geogr. Distr. Unknown.

Genl. Char. Size small. Skull similar to that of *O. schisticeps*, less convex, more depressed; audital bullæ smaller.

Color. Above pale cinnamon rufous, varied with black on dorsal region; sides dull cinnamon; top of head and nape gray tinged with buff; hairs tipped with black; under parts grayish washed with buff; ears blackish externally, narrowly edged with white; inside grayish dusky; fore feet above buffy gray, soles silvery gray; hind feet soles black.

Measurements. (Dried skin.) Hind foot, 27; ear, 23; none other given.

849. Ochotona collaris (Nelson).

Lagomys collaris Nels., Proc. Biol. Soc. Wash., VIII, 1893, p. 117.

Ochotona collaris Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 268. Zoöl. Ser.

COLLARED PIKA.

Type locality. Near head of Tanana River, 200 miles south of Fort Yukon, Alaska.

Geogr. Distr. Mountains south of Fort Yukon from the head of Tanana River to the Chigmit Mountains, near the head of Bristol Bay, Alaska.

850. *Ochotona cuppes* Bangs.

Ochotona cuppes Bangs, Proc. N. Eng. Zoöl. Club, 1, 1899, p. 40.
 Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 268.
 Zoöl. Ser.

DAINTY PIKA.

Type locality. Monishee Divide, Gold Range, British Columbia, Canada. Altitude, 4,000 feet.

Geogr. Distr. Gold Range, British Columbia, Canada.

851. *Ochotona saxatilis* Bangs.

Ochotona saxatilis Bangs, Proc. N. Eng. Zoöl. Club, 1, 1899, p. 41.
 Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 268.
 Zoöl. Ser.

ROCK PIKA.

Type locality. Snowy Range, Montgomery, Park County, Colorado.

Geogr. Distr. Snowy Range, Colorado.

852. *Ochotona minimus* (Lord).

Lagomys minimus Lord, Proc. Zoöl. Soc., 1863, p. 98.
Ochotona minimus Elliot, Syn. N. Am. Mamm., F. C. M. Pub.,
 11, 1901, p. 269. Zoöl. Ser.

LEAST PIKA.

Type locality. Ptarmigan Hill, Cascade Range, British Columbia, Canada. Altitude, 7,000 feet.

Geogr. Distr. Cascade Range, British Columbia, Canada. Limits unknown.

Family XII. **Leporidae. Hares. Rabbits.**

Genus 110. **Romerolagus.**

Romerolagus Merr., Proc. Biol. Soc. Wash., x, 1896, p. 173. Type
Romerolagus nelsoni Merriam.

853. *Romerolagus nelsoni* Merriam.

Romerolagus nelsoni Merr., Proc. Biol. Soc. Wash., x, 1896, p.
 173. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M.
 Pub., IV, Pt. 1, 1904, p. 412, fig. 79. Zoöl. Ser.

NELSON'S HARE.

Type locality. West slope of Mt. Popocatepetl, State of Mexico, Mexico. Altitude, 10,000 to 12,000 feet.

Geogr. Distr. Mt. Popocatepetl and Mt. Iztaccihuatl, State of Mexico, Mexico.

Genus III. **Lepus.**

Lepus Linn., Syst. Nat., I, 1758, p. 57; I, 1766, p. 77. Type *Lepus timidus* Linnæus.

Hydrolagus Gray, Ann. Mag. Nat. Hist., 3d Ser., xx, 1867, p. 221.

Sylvilagus Gray, Ann. Mag. Nat. Hist., 3d Ser., xx, 1867, p. 222.

Tapeti Gray, Ann. Mag. Nat. Hist., 3d Ser., xx, 1867, p. 224.

Macrotolagus Mearns, Science, I, 1895, p. 698.

Microlagus Trouess., Cat. Mamm. vivent. quam fossil., 1897, fasc. III, p. 660.

Limnolagus Mearns, Science, N. S., v, 1897, p. 393.

Brachylagus Miller, Proc. Biol. Soc. Wash., XIII, 1900, p. 157.

Subgenus A. *Limnolagus*. Swamp Hares.**854. Lepus aquaticus** Bachman.

Lepus aquaticus Bachm., Journ. Acad. Nat. Scien. Phil., 1837, p. 319. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 277. Zool. Ser.

GREAT SWAMP HARE.

Type locality. Alabama.

Geogr. Distr. Gulf states, from Alabama to Texas, and northward to southern Illinois in the interior about Mound City and Cairo; up the Ohio to the Wabash and "on the Mississippi to Cape Girardeau, and possibly higher where swamps extend."

a. *Lepus aquaticus attwateri* Allen.

Lepus aquaticus attwateri Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1895, p. 327. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 278. Zool. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 414, pl. XL. Zool. Ser.

ATTWATER'S SWAMP HARE.

Type locality. Medina River, eighteen miles south of San Antonio, Bexar County, Texas.

Geogr. Distr. Yucatan, Mexico, to southeastern Texas.

855. *Lepus palustris* Bachman.

Lepus palustris Bachm., Journ. Acad. Nat. Scien. Phil., 1837, p. 194. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 279. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., 14, Pt. 1, 1904, p. 415. Zoöl. Ser.

EASTERN SWAMP HARE. *Conejo* in Mexico.

Type locality. Eastern South Carolina.

Geogr. Distr. North Carolina to Florida and Texas and south to Yucatan, Mexico.

a. *Lepus palustris paludicola* (Miller & Bangs).

Lepus paludicola Miller & Bangs, Proc. Biol. Soc. Wash., 19, 1894, p. 105.

Lepus palustris paludicola Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 279. Zoöl. Ser.

FLORIDA SWAMP HARE.

Type locality. Fort Island, near Crystal River, Citrus County, Florida.

Geogr. Distr. Southern Florida.

856. *Lepus telmalemonus* Elliot.

Lepus telmalemonus Elliot, Pub. Field Columb. Mus., 1, 1899, p. 285. Zoöl. Ser. *Id.* Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 278, pl. xxv. Zoöl. Ser.

WANDERING SWAMP HARE.

Type locality. Washita River, near Dougherty, Chickasaw Nation, Indian Territory.

Geogr. Distr. Indian Territory. Range unknown.

857. *Lepus truei* Allen.

Lepus truei Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1890, p. 192.

Lepus truei Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., 14, Pt. 1, 1904, p. 415. Zoöl. Ser.

TRUE'S SWAMP HARE.

Type locality. Mirador, State of Vera Cruz, Mexico.

Geogr. Distr. State of Vera Cruz, Mexico. Range unknown.

Subgenus B. **Sylvilagus*. Forest Hares.

858. *Lepus insonus* Nelson.

Lepus insonus Nelson, Proc. Biol. Soc. Wash., xvii, 1904, p. 103. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., 14, Pt. 11, 1904, Append., p. 743. Zoöl. Ser.

**Sylva*—more properly written *silva*.

OMILTEME COTTON-TAIL.

Type locality. Omilteme, State of Guerrero, Mexico.

Geogr. Distr. Known only from the type locality.

859. *Lepus floridanus* (Allen).

Lepus sylvaticus floridanus Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1890, p. 160.

Lepus floridanus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 280, pl. xxvi. Zoöl. Ser.

FLORIDA COTTON-TAIL.

Type locality. Sebastian River, Brevard County, Florida.

Geogr. Distr. North Carolina to Florida, west to Louisiana.

a. *Lepus floridanus mallurus* (Thomas).

Lepus sylvaticus Bachm., Journ. Acad. Nat. Scien. Phil., 1837, p. 403. (nec Nillson.)

Lepus nuttalli mallurus Thomas, Ann. Mag. Nat. Hist., 7th Ser., 11, 1898, p. 320.

Lepus floridanus mallurus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 280. Zoöl. Ser.

COMMON COTTON-TAIL.

Type locality. Raleigh, Wake County, North Carolina.

Geogr. Distr. Southern Maine, along the coast to Florida and the Gulf States, west to eastern Kansas, Nebraska, and the Indian Territory. In the states bordering on the Mississippi it grades into *L. nuttalli* and in Arizona into *L. arizonæ*. In the interior its range is limited apparently by the northern boundary of the United States.

b. *Lepus floridanus mearnsi* Allen.

Lepus sylvaticus mearnsi Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1894, p. 171.

Lepus floridanus mearnsi Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1899, p. 13. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 280. Zoöl. Ser.

MEARNS' COTTON-TAIL.

Type locality. Fort Snelling, Hennepin County, Minnesota.

Geogr. Distr. Iowa, Wisconsin, and Minnesota.

c. *Lepus floridanus transitionalis* (Bangs).

Lepus sylvaticus transitionalis Bangs, Proc. Bost. Soc. Nat. Hist., 1895, p. 405.

Lepus floridanus transitionalis Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 281. Zoöl. Ser.

BANGS' COTTON-TAIL.

Type locality. Liberty Hill, New London County, Connecticut.

Geogr. Distr. East side of lower Hudson River, New York, along the Alleghanies to West Virginia, north to southern New Hampshire, and west to Hudson River.

d. *Lepus floridanus alacer* (Bangs).

Lepus sylvaticus alacer Bangs, Proc. Biol. Soc. Wash., x, 1896, p. 136.

Lepus floridanus alacer Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 282. Zoöl. Ser.

LIVELY COTTON-TAIL.

Type locality. Stilwell, Boston Mountains, Indian Territory.

Geogr. Distr. Missouri, south to Indian and Oklahoma Territories.

e. *Lepus floridanus chapmani* (Allen).

Lepus sylvaticus bachmani Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1894, p. 170. (nec Waterhouse.)

Lepus floridanus chapmani Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1899, p. 12. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 282. Zoöl. Ser.

CHAPMAN'S COTTON-TAIL.

Type locality. Corpus Christi, Nueces County, Texas.

Geogr. Distr. Texas; common in chapparal regions south of Bexar County.

f. *Lepus floridanus connectens* Nelson.

Lepus floridanus connectens Nelson, Proc. Biol. Soc. Wash., xvii, 1904, p. 105. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, Append., p. 743. Zoöl. Ser.

ALTA MIRA COTTON-TAIL.

Type locality. Chichicaxtle, State of Vera Cruz, Mexico.

Geogr. Distr. Tropical parts of eastern Mexico from southern Tamaulipas, throughout the coast lowlands of the Papaloapam River in central Vera Cruz and along the east slope of the Cordillera of eastern San Luis Potosi, eastern Puebla, and eastern Oaxaca, and south to Mt. Zempoaltepec.

g. *Lepus floridanus simplicianus* (Miller).

Lepus simplicianus Miller, Proc. Biol. Soc. Wash., xv, 1902, p. 81.

ALLIED COTTON-TAIL.

Type locality. Brownsville, Cameron County, Texas.

Geogr. Distr. Texas. Range undetermined.

Genl. Char. Like *L. f. chapmani*; "fur shorter, grizzle of head and anterior portion of body much less coarse, owing to the reduced length of the pale ring on each hair."

Color. Above finely grizzled with black and cream buff; sides and rump paler; head and ears clear yellowish gray; nape patch rufous; under parts, tail and feet like *L. f. chapmani*, but with rufous tints duller.

Measurements. Total length, 420; tail vertebræ, 68; hind foot, 91; ear, 90. (Skin.) From fresh specimen. Total length, 416; tail vertebræ, 68; hind foot, 87; ear, 76.

h. Lepus floridanus caniclunis Miller.

Lepus floridanus caniclunis Miller, Proc. Acad. Nat. Scien. Phil., 1899, p. 388. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 283. Zool. Ser.

HOARY-RUMPED COTTON-TAIL.

Type locality. Fort Clark, Kinney County, Texas.

Geogr. Distr. Southwestern Texas. Range unknown.

i. Lepus floridanus aztecus (Allen).

Lepus sylvaticus aztecus Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1890, p. 188.

Lepus floridanus aztecus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 418. Zool. Ser.

TEHUANTEPEC COTTON-TAIL.

Type locality. City of Tehuantepec, State of Oaxaca, Mexico.

Geogr. Distr. State of Oaxaca, Mexico.

j. Lepus floridanus chiapensis Nelson.

Lepus floridanus chiapensis Nelson, Proc. Biol. Soc. Wash., XVII, 1904, p. 106. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, Append., p. 744. Zool. Ser.

CHIAPAS COTTON-TAIL.

Type locality. San Cristobal, State of Chiapas, Mexico.

Geogr. Distr. Interior of the State of Chiapas and Guatemala, from 2,500 to 10,000 feet altitude.

k. Lepus floridanus subcinctus Miller.

Lepus floridanus subcinctus Miller, Proc. Acad. Nat. Scien. Phil., 1899, p. 386. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 417, fig. 80. Zool. Ser.

MICHOACAN COTTON-TAIL.

Type locality. Hacienda El Molino, near Negrete, State of Michoacan, Mexico.

Geogr. Distr. State of Michoacan, Mexico. Range unknown.

l. *Lepus floridanus persultator* Elliot.

Lepus floridanus persultator Elliot, Pub. Field Columb. Mus., III, 1903, p. 147. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 418. Zoöl. Ser.

PUEBLA COTTON-TAIL.

Type locality. Puebla, State of Puebla, Mexico.

Geogr. Distr. State of Puebla, Mexico. Range unknown.

m. *Lepus floridanus yucatanicus* Miller.

Lepus floridanus yucatanicus Miller, Proc. Acad. Nat. Scien. Phil., 1899, p. 384. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 419. Zoöl. Ser.

YUCATAN COTTON-TAIL.

Type locality. Merida, Yucatan, Mexico.

Geogr. Distr. Yucatan, Mexico. Range unknown.

n. *Lepus floridanus holzneri* (Mearns).

Lepus sylvaticus holzneri Mearns, Proc. U. S. Nat. Mus., 1896, p. 554.

Lepus floridanus rigidus Mearns, Proc. U. S. Nat. Mus., 1896, p. 555. (Footnote.)

Lepus holzneri Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 284. Zoöl. Ser.

Lepus durangæ Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1903, p. 609. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 424, Index, p. XI. Zoöl. Ser.

Lepus floridanus holzneri Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 419. Zoöl. Ser.

HOLZNER'S COTTON-TAIL.

Type locality. Near summit of Huachuca Mountains, southern Arizona. Douglas Spruce zone.

Geogr. Distr. Southern Arizona and northern New Mexico to the States of Sonora, Chihuahua, and Durango in northern Mexico.

860. *Lepus auduboni* Baird.

Lepus auduboni Baird, N. Am. Mamm., 1857, p. 608, pl. 58, fig.

2. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 283. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 422. Zoöl. Ser.

AUDUBON'S COTTON-TAIL.

Type locality. San Francisco, California.

Geogr. Distr. Coast of California from San Francisco to San Diego.

861. *Lepus sanctidiegi* (Miller).

Lepus floridanus sanctidiegi Miller, Proc. Acad. Nat. Scien. Phil., 1899, p. 389.

Lepus auduboni sanctidiegi Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 283. Zoöl. Ser.

Lepus sanctidiegi Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 422. Zoöl. Ser.

SAN DIEGO COTTON-TAIL.

Type locality. Mexican boundary line, Monument No. 258, shore of Pacific Ocean, San Diego County, California.

Geogr. Distr. Northern Mexico to southwestern California.

862. *Lepus laticinctus* Elliot.

Lepus laticinctus Elliot, Pub. Field Columb. Mus., III, 1903, p. 254. Zoöl. Ser.

BROAD-BANDED COTTON-TAIL.

Type locality. Oro Grande, Mohave Desert, San Bernardino County, California.

Geogr. Distr. Range undetermined.

Genl. Char. Pelage pale; dark band in front of thigh; soles of feet bister; nasals short.

Color. Upper part of head and dorsal region pinkish buff, the black bases of the hairs showing, giving these parts a streaked appearance of black and pinkish buff; nape tawny ochraceous; rump French gray, darkest in the middle; sides cream buff; broad band in front of thighs like dorsal region; sides of head mixed buff and black; orbital ring pale buff; pectoral band buff; lips, throat, and rest of under parts, under part of fore legs, and upper part of hind legs and feet white; some white on fore feet about toes; soles of all feet bister; tail above blackish, the hairs tipped with buff, beneath white; ears externally mixed buff and black, with the edges white; internally lead color, nearly naked.

Measurements. Total length, 395; tail vertebræ, 62; hind foot, 88; ear, 79. Skull: total length, 70; Hensel, 52; interorbital width, 19; length of nasals, 20; lateral length of nasals, 29; width of nasals posteriorly, 14; anteriorly, 9; palatal length, 25; length of upper tooth row, 11; length of mandible, 54; length of lower tooth row, alveolar border, 13.

a. Lepus laticinctus rufipes Elliot.

Lepus laticinctus rufipes Elliot, Pub. Field Columb. Mus., III, 1903, p. 254. Zoöl. Ser.

RUSSET-FOOTED COTTON-TAIL.

Type locality. Furnace Creek, Death Valley, Inyo County, California.

Geogr. Distr. Death Valley, Inyo County, California. Range undetermined.

Genl. Char. Similar to *L. laticinctus*, but paler and smaller; soles of feet russet; ear shorter.

Color. Upper parts buffy white, base of fur lead color, then pale brown and hairs tipped with white; nape buff, darkest on lower part; top of head like back; sides grayish white; faint brown stripes in front of thighs; rump pale gray; upper part of fore legs, lower part of shoulders and thighs and soles of feet russet; pectoral band pale buff; entire under parts and upper surface of hind feet white; base of fur plumbeous; ears mixed buff and black, edges white; tail above similar to rump, beneath white.

Measurements. Total length, 355; tail vertebræ, 59; hind foot, 85; ear, 74. Skull: total length, 66; Hensel, 50; zygomatic width, 32; interorbital width, 16; median length of nasals, 15; lateral length of nasals, 25; anterior width of nasals, 7; posterior width, 12; palatal length, 24; length of upper tooth row, 11.5; length of lower tooth row, 10.

b. Lepus laticinctus perplicatus Elliot.

Lepus laticinctus perplicatus Elliot, Pub. Field Columb. Mus., III, 1903, p. 255. Zoöl. Ser.

MIXED COTTON-TAIL.

Type locality. Hannopee Cañon, Panamint Mountains, Inyo County, California. Altitude, 7,500 feet.

Geogr. Distr. Panamint, Coso and Inyo Mountains, Inyo County, California.

Genl. Char. Similar to *L. laticinctus* and *L. l. rufipes*, but smaller than the former and larger than the latter, with the soles of the feet Prout's brown. Ears shorter in proportion to the other dimensions.

Color. Upper parts mixed black and buff; rump dark gray, hairs tipped with white; upper part of fore legs vinaceous cinnamon; soles of feet Prout's brown; pectoral band dark buff; throat whitish plumbeous; rest of under parts white.

Measurements. Total length, 380; tail vertebræ, 69; hind foot, 98; ear, 73. Skull: total length, 64.5; Hensel, 49; zygomatic width, 33; interorbital width, 16; median length of nasals, 21; anterior width, 11; palatal length, 23; length of upper tooth row, 8; of lower tooth row, 9.

863. *Lepus veræcrucis* Thomas.

Lepus veræcrucis Thomas, Proc. Zoöl. Soc., 1890, p. 74, pl. VII.

Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV,

Pt. I, 1904, p. 420. Zoöl. Ser.

VERA CRUZ COTTON-TAIL.

Type locality. Las Vigas, Jalapa, State of Vera Cruz, Mexico.

Geogr. Distr. State of Vera Cruz, Mexico. Range unknown.

a. *Lepus veracrucis pacificus* Nelson.

Lepus veracrucis pacificus Nelson, Proc. Biol. Soc. Wash., xvii, 1904, p. 104.

Lepus pacificus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. II, 1904, Append., p. 744. Zoöl. Ser.

ACAPULCO COTTON-TAIL.

Type locality. Acapulco, State of Guerrero, Mexico.

Geogr. Distr. Pacific coast region of State of Guerrero and adjacent part of the State of Oaxaca, Mexico.

864. *Lepus russatus* Allen.

Lepus russatus Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1904, p. 31. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. I, 1904, p. 420. Zoöl. Ser.

RUSSET COTTON-TAIL.

Type locality. Pasa Nueva, State of Vera Cruz, Mexico.

Geogr. Distr. State of Vera Cruz, Mexico. Range unknown.

865. *Lepus parvulus* Allen.

Lepus parvulus Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1904, p. 134. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. I, 1904, p. 421. Zoöl. Ser.

LITTLE COTTON-TAIL.

Type locality. Apam, southern part of the State of Hidalgo, Mexico.

Geogr. Distr. State of Hidalgo, Mexico. Range unknown.

866. *Lepus insolitus* Allen.

Lepus insolitus Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1890, p. 189. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. I, 1904, p. 421. Zoöl. Ser.

PLAINS COTTON-TAIL.

Type locality. Plains of Colima, State of Jalisco, Mexico.

Geogr. Distr. State of Jalisco, Mexico.

867. *Lepus nuttalli* Bachman.

Lepus nuttalli Bachm., Journ. Acad. Nat. Scien. Phil., 1837, p. 345, pl. 22. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 284. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. I, 1904, p. 425. Zoöl. Ser.

NUTTALL'S COTTON-TAIL.

Type locality. Plains near the junction of the Snake and Columbia Rivers, Washington.

Geogr. Distr. From the 49th parallel south to Texas and the State of Sonora, Mexico, and west through the Sierra Nevada to Oregon and Washington.

868. *Lepus baileyi* Merriam.

Lepus baileyi Merr., Proc. Biol. Soc. Wash., XI, 1897, p. 148.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 284.

Zoöl. Ser.

BAILEY'S COTTON-TAIL.

Type locality. Spring Creek, east side of Bighorn Basin, Bighorn County, Wyoming.

Geogr. Distr. Bighorn Basin in Montana, across the Owl Creek Mountains into Wind River Basin, and east into northeastern Wyoming. Exact range not determined.

869. *Lepus grangeri* (Allen).

Lepus sylvaticus grangeri Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1895, p. 264.

Lepus grangeri Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1896, p.

246. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 285. Zoöl. Ser.

GRANGER'S COTTON-TAIL.

Type locality. Hill City, Black Hills, Custer County, South Dakota.

Geogr. Distr. Mountains of South Dakota. Range unknown.

870. *Lepus arizonæ* (Allen).

Lepus sylvaticus var. *arizonæ* Allen, N. Am. Rod., 1877, p. 332.

Lepus arizonæ Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II,

1901, p. 285. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 423. Zoöl. Ser.

ARIZONA COTTON-TAIL.

Type locality. Beal's Spring, fifty miles west of Fort Whipple, Yavapai County, Arizona.

Geogr. Distr. Deserts of Arizona and the Chiricahua and Huachuca Mountains (but not to the White Mountains), up to 8,500 feet, south into Lower California and the State of Sonora, Mexico.

***a. Lepus arizonæ goldmani* Nelson.**

Lepus arizonæ goldmani Nelson, Proc. Biol. Soc. Wash., XVII, 1904, p. 107.

Lepus goldmani Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, Append., p. 745. Zoöl. Ser.

SINALOA COTTON-TAIL.

Type locality. Culiacan, State of Sinaloa, Mexico.

Geogr. Distr. Southern part of the State of Sonora to central part of the State of Sinaloa.

b. **Lepus arizonæ pinetis* (Allen).

Lepus sylvaticus pinetis Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1894, p. 348.

Lepus grangeri pinetis Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 285. Zoöl. Ser.

PINE WOOD COTTON-TAIL.

Type locality. White Mountains, Apache County, Arizona.

Geogr. Distr. White Mountains, southeastern Arizona north to Colorado.

c. *Lepus arizonæ major* Mearns.

Lepus arizonæ major Mearns, Proc. U. S. Nat. Mus., 1896, p. 557. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 286. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., 14, Pt. 1, 1904, p. 423. Zoöl. Ser.

GREATER DESERT COTTON-TAIL.

Type locality. Calabasas, Pima County, Arizona.

Geogr. Distr. Colorado, through plateau of Arizona south to the Mimbres, State of Chihuahua, and west to Poso de Luis, State of Sonora, Mexico.

d. *Lepus arizonæ minor* Mearns.

Lepus arizonæ minor Mearns, Proc. U. S. Nat. Mus., 1896, p. 557. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 286. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., 14, Pt. 1, 1904, p. 423. Zoöl. Ser.

LESSER DESERT COTTON-TAIL.

Type locality. El Paso, El Paso County, Texas.

Geogr. Distr. Plains of Colorado southward to the Rio Grande, and westward to the elevated central tract, and intergrades with *L. a. major* in the pass between the southern end of the Rocky Mountains and the northern extremity of the Sierra Nevada; south into the State of Chihuahua, Mexico.

e. *Lepus arizonæ confinis* Allen.

Lepus arizonæ confinis Allen, Bull. Am. Mus. Nat. Hist. N. Y., x, 1898, p. 146. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., 14, Pt. 1, 1904, p. 424. Zoöl. Ser.

ALLIED COTTON-TAIL.

Type locality. Playa Maria, Lower California, Mexico.

Geogr. Distr. Lower California, Mexico. Range unknown.

*More appropriately a sub-species of *L. arizonæ* than of *L. grangeri*, as given in the Synopsis.

871. *Lepus orizabæ* Merriam.

Lepus orizabæ Merr., Proc. Biol. Soc. Wash., VIII, 1893, p. 143.

Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV,

Pt. I, 1904, p. 425. Zoöl. Ser.

ORIZABA COTTON-TAIL.

Type locality. Mt. Orizaba, State of Puebla, Mexico. Altitude, about 9,500 feet.

Geogr. Distr. State of Puebla, Mexico.

872. *Lepus graysoni* Allen.

Lepus graysoni Allen, N. Am. Rod., 1877, p. 347. Elliot, Mamm.

Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p.

426. Zoöl. Ser.

GRAYSON'S COTTON-TAIL.

Type locality. Maria Madre Island, Tres Marias Islands, State of Jalisco, Mexico.

Geogr. Distr. Tres Marias Islands, State of Jalisco, Mexico.

Subgenus C. *Brachylagus*. Short-tailed Cotton-tails.

873. *Lepus idahoensis* Merriam.

Lepus idahoensis Merr., N. Am. Faun., No. 5, 1891, p. 76, fig. 4,

p. 77. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p.

287. Zoöl. Ser.

IDAHO COTTON-TAIL.

Type locality. Pahsimeroi Valley, central Idaho.

Geogr. Distr. Sage Plains on the Snake River, Idaho, and the northward extension of these plains; south into northern Nevada, and west into eastern Oregon and Washington.

Subgenus D. *Tapeti*. Tailless Hares.

874. *Lepus gabbi* (Allen).

Lepus brasiliensis var. *gabbi* Allen, N. Am. Rod., 1877, p. 349.

Lepus gabbi Elliot, Mamm. Middle Amer. & W. Indies, F. C. M.

Pub., IV, Pt. I, 1904, p. 426, fig. 81. Zoöl. Ser.

GABB'S HARE.

Type locality. Talamanca, Costa Rica.

Geogr. Distr. Costa Rica, Central America.

875. *Lepus incitatus* Bangs.

Lepus incitatus Bangs, Amer. Nat., xxxv, 1901, p. 633. Elliot,

Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I,

1904, p. 428. Zoöl. Ser.

FLEET HARE.

Type locality. San Miguel Island, Bay of Panama.

Geogr. Distr. Known only from the type locality.

Subgenus E. *Microlagus*. Short-tailed Hares.

876. *Lepus bachmani* Waterhouse.

Lepus bachmani Waterh., Proc. Zool. Soc., 1838, p. 103. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 429. Zool. Ser.

Lepus floridanus bachmani Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 281. Zool. Ser.

BACHMAN'S COTTON-TAIL. *Conejo* in Mexico.

Type locality. Southwest coast of North America, probably California.

Geogr. Distr. Fort Crook, California, south to Cape St. Lucas, Lower California, Mexico.

a. *Lepus bachmani ubericolor* Miller.

Lepus bachmani ubericolor Miller, Proc. Acad. Nat. Scien. Phil., 1899, p. 383.

Lepus floridanus ubericolor Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 282. Zool. Ser.

RICH-COLORED COTTON-TAIL.

Type locality. Beaverton, Washington County, Oregon.

Geogr. Distr. Northwestern Oregon. Range unknown.

877. *Lepus cerrosensis* Allen.

Lepus cerrosensis Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1898, p. 145. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 428. Zool. Ser.

CERROS ISLAND COTTON-TAIL.

Type locality. Cerros or Cedros Island, off west coast of Lower California, Mexico.

Geogr. Distr. Known only from the type locality.

878. *Lepus cinerascens* Allen.

Lepus cinerascens Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1890, p. 159. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 287, Pl. XXVII. Zool. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 429, fig. 82. Zool. Ser.

ASH-COLORED COTTON-TAIL.

Type locality. San Fernando, Los Angeles County, California.

Geogr. Distr. Lower California, Mexico, and desert of southern California.

879. *Lepus peninsularis* Allen.

Lepus peninsularis Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1898, p. 144. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 430. Zoöl. Ser.

LOWER CALIFORNIA COTTON-TAIL.

Type locality. Santa Anita, Lower California, Mexico.

Geogr. Distr. Cape Region of Lower California, Mexico.

Subgenus F. *Macrotolagus*. Jack Rabbits.

880. *Lepus alleni* Mearns.

Lepus alleni Mearns, Bull. Am. Mus. Nat. Hist. N. Y., 1890, p. 294. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 288. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 433, fig. LVIII. Zoöl. Ser.

ALLEN'S JACK RABBIT.

Type locality. Rillito Station, Southern Pacific Railroad, Pima County, Arizona.

Geogr. Distr. Desert region between Phoenix and Benson, Arizona. Mexico?

a. Lepus alleni palitans Bangs.

Lepus alleni palitans Bangs, Proc. N. Eng. Zoöl. Club, I, 1900, p. 85. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 434. Zoöl. Ser.

WANDERING JACK RABBIT.

Type locality. Agua Caliente, forty miles southeast of Mazatlan, State of Sinaloa, Mexico.

Geogr. Distr. State of Sinaloa, Mexico. Range unknown.

881. *Lepus callotis* Wagler.

Lepus callotis Wagl., Nat. Syst. Amph., 1830, p. 23. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 288. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 431. Zoöl. Ser.

Lepus nigricaudatus Bennett, Proc. Zoöl. Soc., 1833, p. 41.

Lepus mexicanus Rich., 6th Rep. Brit. Assoc., 1837, p. 150.

**Lepus flavigularis* Wagn., Schreb. Säugth., Suppl., IV, 1844, p. 107

BEAUTIFUL-EARED JACK RABBIT.

Type locality. "Mexico."

Geogr. Distr. Through Mexico, from State of Chihuahua to Tehuantepec.

* See Appendix, p. 543.

882. *Lepus festinus* Nelson.

Lepus festinus Nelson, Proc. Biol. Soc. Wash., xvii, 1904, p. 108.

Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. II, 1904, Append., p. 746. Zoöl. Ser.

HIDALGO JACK RABBIT.

Type locality. Irolo, State of Hidalgo, Mexico.

Geogr. Distr. Southeastern part of Mexican tableland in southern and eastern parts of State of Queretaro, throughout most of the State of Hidalgo, extreme northern part of State of Mexico and Valley of Mexico, State of Tlaxcala, and adjacent northern part of the State of Puebla.

883. *Lepus merriami* Mearns.

Lepus texianus! Aud. & Bachm., N. Am. Quad., III, 1853, p. 156, pl. cxxxiii. (nec Waterhouse.)

Lepus merriami Mearns, Proc. U. S. Nat. Mus., 1896, p. 444.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 289, pl. xxvii. Zoöl. Ser. *Id.*, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. I, 1904, p. 432. Zoöl. Ser.

MERRIAM'S JACK RABBIT.

Type locality. Fort Clark, Kinney County, Texas.

Geogr. Distr. Northern Mexico near boundary line and "Lower Gulf Coast to the mouth of the Rio Grande, and up that stream to the mouth of the Devil's River."

a. *Lepus merriami altimiræ* Nelson.

Lepus merriami altimiræ Nelson, Proc. Biol. Soc. Wash., xvii, 1904, p. 109.

Lepus altimiræ Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. II, 1904, Append., p. 745. Zoöl. Ser.

ALTA MIRA JACK RABBIT.

Type locality. Alta Mira, State of Tamaulipas, Mexico.

Geogr. Distr. Coast plains in southern part of the State of Tamaulipas, extreme northern part of the State of Vera Cruz, and eastern part of the State of San Luis Potosi.

884. *Lepus gaillardi* Mearns.

Lepus gaillardi Mearns, Proc. U. S. Nat. Mus., 1896, p. 560. Elliot,

Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 289. Zoöl.

Ser. *Id.*, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. I, 1904, p. 433. Zoöl. Ser.

GAILLARD'S JACK RABBIT.

Type locality. Playas Valley, near its west fork, near Monument No. 63, Mexican boundary line, Grant County, New Mexico.

Geogr. Distr. State of Sonora, Mexico, into plain east of the San Luis Mountains, at the head of the Rio Yaqui, and east of that river's watershed. Extent of range unknown.

a. Lepus gaillardi battyi Allen.

Lepus gaillardi battyi Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1903, p. 607. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 433. Zoöl. Ser.

BATTY'S JACK RABBIT.

Type locality. Rancho Santuario, State of Durango, Mexico.

Geogr. Distr. State of Durango, Mexico. Range unknown.

885. *Lepus asellus* Miller.

Lepus asellus Miller, Proc. Acad. Nat. Scien. Phil., 1899, p. 380.

Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 435. Zoöl. Ser.

DONKEY JACK RABBIT.

Type locality. Near city of San Luis Potosi, State of San Luis Potosi, Mexico.

Geogr. Distr. State of San Luis Potosi, Mexico. Range unknown.

886. *Lepus melanotis* Mearns.

Lepus melanotis Mearns, Bull. Am. Mus. Nat. Hist. N. Y., 1890,

p. 297. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 290. Zoöl. Ser.

BLACK-EARED JACK RABBIT.

Type locality. Independence, Montgomery County, Kansas.

Geogr. Distr. Kansas, western Texas, Oklahoma, and Indian Territories.

887. *Lepus texensis* (Waterhouse).

Lepus texianus! Waterh., Nat. Hist. Mamm., II, 1848, p. 136.

Lepus texensis Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 290. Zoöl. Ser.

TEXAS JACK RABBIT.

Type locality. Unknown.

Geogr. Distr. Southeastern Oregon to Mexico, and from the Sierra Nevada to eastern border of plains east of Rocky Mountains?

a. Lepus texensis eremicus (Allen).

Lepus texianus! eremicus Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1894, p. 347.

Lepus texensis eremicus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 291. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. I, 1904, p. 435, pl. xli. Zoöl. Ser.

DESERT JACK RABBIT.

Type locality. Fairbank, Cochise County, Arizona.

Geogr. Distr. White Mountains region, southeastern Arizona, south into State of Sonora, Mexico.

b. *Lepus texensis griseus* (Mearns).

Lepus texianus! griseus Mearns, Proc. U. S. Nat. Mus., 1896, p. 562.

Lepus texensis griseus Elliot, Syn. N. Am. Mamm., F. C. M. Pub.,

II, 1901, p. 291. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W.

Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 436. Zoöl. Ser.

GRAY DESERT JACK RABBIT.

Type locality. Fort Hancock, El Paso County, Texas.

Geogr. Distr. Upper Rio Grande from Maverick to Kinney County, Texas, west to Grant County, New Mexico, and south to the States of Coahuila and Chihuahua, Mexico.

c. *Lepus texensis micropus* (Allen).

Lepus texianus! micropus Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1903, p. 605.

Lepus texensis micropus Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., IV, Pt. 1, 1904, p. 436. Zoöl. Ser.

SMALL-FOOTED HARE.

Type locality. Rio del Bocas, State of Durango, Mexico. Altitude, 6,800 feet.

Geogr. Distr. State of Durango.

d. *Lepus texensis deserticola* (Mearns).

Lepus texianus! deserticola Mearns, Proc. U. S. Nat. Mus., 1896, p. 564.

Lepus texensis deserticola Elliot, Syn. N. Am. Mamm., F. C. M.

Pub., II, 1901, p. 291. Zoöl. Ser. *Id.* Mamm. Middle Amer.

& W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 437. Zoöl. Ser.

WESTERN DESERT JACK RABBIT.

Type locality. Western edge of the Colorado Desert, at the base of the Coast Range Mountains, San Diego County, California.

Geogr. Distr. Desert region east of the high Sierra, including the Mohave and Colorado deserts and the region between the Sonoyta Valley of Arizona and the State of Sonora, Lower California, Mexico.

e. *Lepus texensis wallawalla* (Merriam).

Lepus texianus! wallawalla Merr., Proc. Biol. Soc. Wash., XVII, 1904, p. 137.

WALLAWALLA JACK RABBIT.

Type locality. Touchet, Plains of the Columbia, Washington.

Geogr. Distr. Plains of the Columbia in Oregon and Washington

Genl. Char. Similar to *L. texensis deserticola*, but darker above in summer pelage; in winter pelage like *L. t. eremicus* or *L. tularensis*. Ears shorter; hind foot slightly larger.

Color. Fall Pelage. Upper parts mixed black and buffy fulvous, deeper on the sides; pectoral band dark fulvous; top of head and sides of face grizzled gray; orbital ring and sides of face pale fulvous; ears finely grizzled fulvous brown. In summer the orbital ring is nearly white; cheeks are pale buffy gray, and the pectoral collar pale yellowish buffy.

Measurements. Type. Total length, 555; tail vertebræ, 95; hind foot, 126.

888. *Lepus tularensis* Merriam.

Lepus tularensis Merr., Proc. Biol. Soc. Wash., xvii, 1904, p. 136.
TULARE JACK RABBIT.

Type locality. Alila (San Joaquin Valley), Tulare County, California.

Geogr. Distr. Baker's field, Tulare Basin, south end of San Joaquin Valley, California.

Genl. Char. Similar to *L. t. deserticola*, but paler and more yellowish; size rather small.

Color. General color pale buffy grayish toned with yellowish; back slightly grizzled with black; nape patch whitish; face and neck pale buffy yellowish or yellowish buff slightly grizzled with dark hairs; pectoral collar pale yellowish; thighs grayish clay color; under parts white tinged with pale yellow; buffy on sides.

Measurements. Total length, 558; tail vertebræ, 84; hind foot, 117.

889. *Lepus californicus* Gray.

Lepus californica Gray, Charlesw. Mag. Nat. Hist., i, 1837, p. 586.

Lepus californicus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., ii, 1901, p. 291. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 1, 1904, p. 437. Zoöl. Ser.

Lepus bennetti Gray, Voy. Sulphur, Zoöl., 1844, p. 36, pl. 14.

CALIFORNIA JACK RABBIT.

Type locality. "St. Antoine," Santa Barbara County? California.

Geogr. Distr. Northern California south to Cape St. Lucas, Lower California, Mexico.

a. *Lepus californicus xanti* Thomas.

Lepus californicus xanti Thomas, Ann. Mag. Nat. Hist., 7th Ser.,

i, 1898, p. 45. Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., iv, Pt. 1, 1904, p. 438. Zoöl. Ser.

LOWER CALIFORNIA JACK RABBIT.

Type locality. Santa Anita, Lower California, Mexico.

Geogr. Distr. Cape Region of Lower California, Mexico.

890. *Lepus insularis* Bryant.

Lepus insularis Bryant, Proc. Calif. Acad. Scien., 2d Ser., III, 1891, p. 92. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 438. Zool. Ser.

Lepus edwardsi St. Loup, Bull. Mus. Hist. Nat. Paris, 1895, p. 5.

ESPERITO SANTO ISLAND JACK RABBIT.

Type locality. Esperito Santo Island, coast of Lower California, Mexico.

Geogr. Distr. Known only from the type locality.

891. *Lepus martirensis* Stowell.

Lepus martirensis Stowell, Proc. Calif. Acad. Scien., 2d Ser., v, 1895, p. 51. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 1, 1904, p. 439. Zool. Ser.

SAN PEDRO MARTIR JACK RABBIT.

Type locality. La Grulla, San Pedro Martir Mountains, Lower California, Mexico. Altitude, 8,000 feet.

Geogr. Distr. San Pedro Martir Range, Lower California, Mexico.

Subgenus G. *Lepus*. Changeable Hares.

892. *Lepus arcticus* "Leach" Ross.

Lepus arcticus "Leach" Ross, Voy. Arct. Reg., II, 1819, Append., IV, p. 151. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 270. Zool. Ser.

Lepus glacialis Leach, in Ross' Voy. Arct. Reg., II, 1819, Append., IV, p. 170.

Lepus timidus var. *arcticus* Allen, Mon. N. Am. Rod., 1877, p. 288.

POLAR HARE.

Type locality. Baffin Land, southeast of Cape Bowen.

Geogr. Distr. Labrador, west shore of Hudson Bay, north shore of Great Slave Lake, and Barren Grounds to the Arctic Ocean; west of the mountains in the Upper Yukon Valley.

a. *Lepus arcticus bangsi* Rhoads.

Lepus arcticus bangsi Rhoads, Amer. Nat., xxx, 1896, p. 236. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 270. Zool. Ser.

BANGS' POLAR HARE.

Type locality. Codroy, Newfoundland.

Geogr. Distr. Newfoundland, northern Labrador. Range unknown.

b. Lepus arcticus canus Preble.

Lepus arcticus canus Preble, N. Am. Faun., No. 22, 1902, p. 59.

HOARY POLAR HARE.

Type locality. Hubbard Point, 75 miles north of Fort Churchill, Keewatin, Canada.

Geogr. Distr. Barren Grounds from Fort Churchill northwards in Arctic America.

Color. Upper parts with the hairs light plumbeous at base, then yellowish fawn, then dusky and tips white; throat, rump, and thighs light plumbeous; chin grayish white; head grizzled fawn; nape grayish; occiput dusky; soles of feet clay color; ears anteriorly dusky grayish, edged at the center with white; interior dusky with the tips white. Winter pelage white, ears tipped with black.

Measurements. Total length, 570; tail vertebræ, 80; hind foot, 154. Skull: greatest length, 98; zygomatic breadth, 50; breadth across postorbital processes, 39; length of the upper molar series, 19.

893. *Lepus labradorius* Miller.

Lepus labradorius Miller, Proc. Biol. Soc. Wash., XIII, 1899, p. 39.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 271. Zoöl. Ser.

LABRADOR HARE.

Type locality. Fort Chimo, Ungava, Labrador.

Geogr. Distr. Ungava, Labrador. Range unknown.

894. *Lepus grænlændicus* Rhoads.

Lepus grænlændicus Rhoads, Amer. Nat., xxx, 1896, p. 236. Elliot,

Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 271. Zoöl. Ser.

GREENLAND HARE.

Type locality. Robertson's Bay (N. lat. 78°), Greenland.

Geogr. Distr. Greenland.

895. *Lepus tschuktschorum* Nordquist.

Lepus tschuktschorum Nordquist, Vega Exped., II, 1883, p. 84,

figs. 8, 9, 10. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 271. Zoöl. Ser.

SIBERIAN HARE.

Type locality. Pitlekaj, lat. 67° N., long. 173° W., northeast Siberia.

Geogr. Distr. Northwest Alaska from the mouth of the Kuskokwim River to the Arctic Ocean. Siberia.

896. *Lepus othus* Merriam.

Lepus othus Merr., Proc. Wash. Acad. Scien., II, 1900, p. 28. Elliot,

Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 272. Zoöl. Ser.

ST. MICHAELS' HARE.

Type locality. St. Michaels, Norton Sound, Alaska.

Geogr. Distr. Northwestern Alaska. Range unknown.

897. *Lepus poadromus* Merriam.

Lepus poadromus Merr., Proc. Wash. Acad. Scien., 11, 1900, p. 29.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 272.

Zoöl. Ser.

SWIFT HARE.

Type locality. Stepovak Bay, Alaska Peninsula, Alaska.

Geogr. Distr. Alaska Peninsula. Range unknown.

898. *Lepus americanus* Erxleben.

Lepus americanus Erxl., Syst. Reg. Anim., 1, 1777, p. 330.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 273.

Zoöl. Ser.

Lepus hudsonius Pallas, Nov. Spec. Quad. Glires, 1778, p. 30.

Lepus nanus Schreb., Säugth., 11, 1792, pl. 234 B.

Lepus borealis Schinz, Synop., 11, 1845, p. 286.

Lepus campestris Hayden, Amer. Nat., 111, 1869, p. 115. (Part.)

AMERICAN HARE.

Type locality. Vicinity of Hudson Bay, Canada.

Geogr. Distr. New Brunswick and Nova Scotia northward to the limit of trees, and westward to Alaska.

a. *Lepus americanus virginianus* (Harlan).

Lepus virginianus Harlan, Faun. Amer., 1825, p. 196.

Lepus americanus virginianus Elliot, Syn. N. Am. Mamm., F. C. M.

Pub., 11, 1901, p. 273. Zoöl. Ser.

VIRGINIAN HARE.

Type locality. Blue Mountains, northeast of Harrisburg, Pennsylvania.

Geogr. Distr. Atlantic coast from Nova Scotia to Connecticut, and in the Alleghanies south to Virginia; through the northern part of the border tier of States and the southern half of Canada west to the north shore of Lake Superior and northern Minnesota to the Missouri River.

b. *Lepus americanus phæonotus* Allen.

Lepus americanus phæonotus Allen, Bull. Am. Mus. Nat. Hist.

N. Y., 1899, p. 11. Elliot, Syn. N. Am. Mamm., F. C. M. Pub.,

11, 1901, p. 273. Zoöl. Ser.

WESTERN HARE.

Type locality. Hallock, Kittson County, Minnesota.

Geogr. Distr. Central and northern Minnesota. Range unknown.

c. *Lepus americanus struthopus* Bangs.

Lepus americanus struthopus Bangs, Proc. Biol. Soc. Wash., xii, 1898, p. 81. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., ii, 1901, p. 274. Zoöl. Ser.

GREAT-FOOTED HARE.

Type locality. Digby, Nova Scotia, Canada.

Geogr. Distr. Province of Nova Scotia. Range unknown.

d. *Lepus americanus bairdi* (Hayden).

Lepus bairdi Hayden, Amer. Nat., iii, 1869, p. 115.

Lepus americanus bairdi Elliot, Syn. N. Am. Mamm., F. C. M. Pub., ii, 1901, p. 274. Zoöl. Ser.

BAIRD'S HARE.

Type locality. Columbia Valley, Wind River Mountains, Fremont County, Wyoming.

Geogr. Distr. Higher portions of the Rocky Mountains from Great Slave Lake, Arctic America, south to New Mexico.

e. *Lepus americanus washingtoni* (Baird).

Lepus washingtonii Baird, Proc. Acad. Nat. Scien. Phil., 1855, p. 333, pl. 15.

Lepus americanus washingtoni Elliot, Syn. N. Am. Mamm., F. C. M. Pub., ii, 1901, p. 274. Zoöl. Ser.

WASHINGTON HARE.

Type locality. Vicinity of Puget Sound, Washington.

Geogr. Distr. North latitude 55°, south to the mouth of the Willamette River, Oregon.

f. *Lepus americanus columbiensis* Rhoads.

Lepus americanus columbiensis Rhoads, Proc. Acad. Nat. Scien. Phil., 1895, p. 242. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., ii, 1901, p. 275. Zoöl. Ser.

BRITISH COLUMBIAN HARE.

Type locality. Vernon, British Columbia, Canada.

Geogr. Distr. Arid interior basin of southern British Columbia, Canada.

g. *Lepus americanus dalli* Merriam.

Lepus americanus dalli Merr., Proc. Wash. Acad. Scien., ii, 1900, p. 29. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., ii, 1901, p. 275. Zoöl. Ser.

DALL'S HARE.

Type locality. Nulato River, Alaska.

Geogr. Distr. Unknown.

h. Lepus americanus macfarlanii Merriam.

Lepus americanus macfarlani Merr., Proc. Wash. Acad. Scien., 11, 1900, p. 30. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 275. Zoöl. Ser.

MACFARLANE'S HARE.

Type locality. Fort Anderson, near mouth of Anderson River, Mackenzie, Arctic America, Canada.

Geogr. Distr. Arctic America. Range unknown.

899. *Lepus saliens* Osgood.

Lepus saliens Osgood, N. Am. Faun., No. 19, 1900, p. 39. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 276. Zoöl. Ser.

LEAPING HARE.

Type locality. Caribou Crossing, between Lake Bennett and Lake Tagish, Northwest Territory, Canada.

Geogr. Distr. Limits of range unknown.

900. *Lepus bishopi* Allen.

Lepus bishopi Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1899, p. 11. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 276. Zoöl. Ser.

BISHOP'S HARE.

Type locality. Mill Lake, Turtle Mountains, near northern border of North Dakota.

Geogr. Distr. North Dakota. Range unknown.

901. *Lepus klamathensis* Merriam.

Lepus klamathensis Merr., N. Am. Faun., No. 16, 1899, p. 100. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 276. Zoöl. Ser.

KLAMATH HARE.

Type locality. Fort Klamath, Klamath County, Oregon.

Geogr. Distr. Oregon. Range unknown.

902. *Lepus campestris* Bachman.

Lepus campestris Bachm., Journ. Acad. Nat. Scien. Phil., 1837, p. 349. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 277. Zoöl. Ser.

PLAINS HARE.

Type locality. Plains of the Saskatchewan, Canada.

Geogr. Distr. Twin Plains of Saskatchewan southward to Kansas, and from Minnesota westward to the Rocky Mountains; and from Green River in southwestern Wyoming over eastern Idaho, northern Utah, and northeastern Nevada.

a. *Lepus campestris richardsoni* (Bachman).

Lepus richardsoni Bachm., Journ. Acad. Nat. Scien. Phil., 1839, p. 88. Merr., Proc. Biol. Soc. Wash., xvii, 1904, p. 136.

RICHARDSON'S HARE.

Type locality. California.

Geogr. Distr. Salinas Valley, and Ranges on either side, and mountains around south end of Joaquin Valley, passing north in the foothills of the Sierra to about latitude of San Francisco.

Color. Head, cheeks, and upper parts mixed yellowish white, black, and pale buff, giving a mottled gray effect; back of ears, neck, breast, and upper part of fore and hind legs brownish gray; chin and throat white; entire under surface of the body white, tinged with yellowish on the sides; soles of feet pale dark brown; tail above black, beneath pale brownish yellow; outside of ears grizzled black and pale yellow, edged with white, but not reaching the tip; interior of ears white, becoming black near the edge.

Measurements. Total length, 481.6; tail vertebræ, 33; hind foot to end of nails, 119.3; ear, 175.8.

b. *Lepus campestris townsendi* (Bachman).

Lepus townsendi Bachm., Journ. Acad. Nat. Scien. Phil., 1839, p. 90.

Lepus campestris townsendi Merr., Proc. Biol. Soc. Wash., xvii, 1904, p. 132.

TOWNSEND'S HARE.

Type locality. Walla Walla, Washington.

Geogr. Distr. Plains of the Columbia and Oregon in Washington.

Color. Upper parts clear gray; thighs and hind legs deep gray; top of fore leg and foot buffy gray grizzled with black; top of head and front of ears gray tinged slightly with pale buffy fulvous; pectoral collar buffy gray; edge of ear black.

Measurements. "Average of 5. Total length, 576; tail vertebræ, 81; hind foot, 147."

c. *Lepus campestris sierræ* Merriam.

Lepus campestris sierræ Merr., Proc. Biol. Soc. Wash., xvii, 1904, p. 132.

SIERRA HARE.

Type locality. Hope Valley, Alpine County, California. Altitude, 7,800 feet.

Geogr. Distr. Sierra Nevada from Lake Tahoe to south of Mono Lake, in summer. Winter range, east side of the Sierra in Nevada and California.

Genl. Char. Size large; similar to *L. c. townsendi* generally.

Color. Back, thighs, and pectoral collar gray; gray median band on tail above; orbital ring broad, white; upper lip and sides of nose buffy fulvous; flanks gray, encroaching on the belly; top of fore legs grizzled buffy fulvous; wrists and fore feet dirty yellowish white; hind feet white.

Measurements. Total length, 635; tail vertebræ, 112; hind foot, 167.

Order VII. **Carnivora. Carnivores.**

Family I. **Felidae. Cats.**

Genus 112. **Felis.**

Felis Linn., Syst. Nat., 1, 1758, p. 41; 1, 1766, p. 60. Type *Felis catus* Linnæus.

Linx (sic) Frisch., Das Natur-syst. vierfüss. Thiere, in Tabellen, 12, Tab. Gen., 1775.

Lynx Kerr, Anim. Kingd. Syst. Catal., pp. 32-33, 1792, desc. p. 155.

Lynceus Gray, Lond. Med. Repos., xv, 1821, 302.

Puma Jard., Nat. Libr., Mamm., 11, 1834, p. 266.

Lynchus Jard., Nat. Libr., Mamm., 11, 1834, pp. 274, 275.

Leopardus Gray, Ann. Mag. Nat. Hist., x, 1842, 260. (nec 1867).

Herpailurus Severtz., Rev. Mag. Zool., 2me Sér., x, 1858, p. 385.

Jaguaris Severtz., Rev. Mag. Zool., 2me Sér., x, 1858, pp. 386, 390.

Lynchailurus Severtz., Rev. Mag. Zool., 2me Sér., x, 1858, pp. 386, 390.

Noctifelis Severtz., Rev. Mag. Zool., 2me Sér., x, 1858, pp. 386, 390.

Oncifelis Severtz., Rev. Mag. Zool., 2me Sér., x, 1858, pp. 386, 390.

Leopardus Gray, Proc. Zoöl. Soc., 1867, p. 263.

Margay Gray, Proc. Zoöl. Soc., 1867, p. 271.

Pardalis Gray, Proc. Zoöl. Soc., 1867, p. 272.

Cervaria Gray, Proc. Zoöl. Soc., 1867, p. 276.

Subgenus A. **Leopardus. Jaguars, Ocelots, etc.**

Felis onca centralis (Mearns).

Felis centralis Mearns, Proc. Biol. Soc. Wash., xiv, 1901, p. 139.

Felis onca centralis Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. 11, 1904, p. 446, pls. XLII and XLIII. Zoöl. Ser.

COSTA RICA JAGUAR.

Type locality. Talamanca, Costa Rica.

Geogr. Distr. Costa Rica, north to Honduras, Central America.

Felis onca hernandezii (Gray).

Leopardus hernandezii Gray, Proc. Zoöl. Soc., 1857, p. 278.

Felis onca Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 294. Zoöl. Ser. (Part.)

Felis onca hernandezi Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, 1904, p. 446. Zoöl. Ser. *Id.* Mon. Felidæ, pl. v (*Felis onca*), fig. LIX, No. 2.

HERNANDEZ'S JAGUAR.

Type locality. Mazatlan, State of Sinaloa, Mexico.

Geogr. Distr. Apparently western Mexico from State of Colima north to San Blas.

Felis onca goldmani (Mearns).

Felis hernandezii goldmani Mearns, Proc. Biol. Soc. Wash., XIV, 1901, p. 142.

Felis onca goldmani Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 447. Zoöl. Ser.

GOLDMAN'S JAGUAR.

Type locality. Yohatlan, State of Campeche, Mexico.

Geogr. Distr. Unknown.

903. *Felis pardalis* Linnæus.

Felis pardalis Linn., Syst. Nat., I, 1758, p. 42; I, 1766, p. 62. Elliot, Mon. Felidæ, 1881, pl. XVIII. *Id.* Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 294. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 447. Zoöl. Ser.

OCELOT. *Manigordo* in Costa Rica.

Type locality. Mexico.

Geogr. Distr. Southern Texas to Mexico and southward to Buenos Ayres, east of the Andes. Boquete, Chiriqui, Panama. Central America. (Bangs.)

a. *Felis pardalis mearnsi* Allen.

Felis costaricensis Mearns, Proc. U. S. Nat. Mus., 1902, p. 245. (nec Merr.)

Felis pardalis mearnsi Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1904, p. 71. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 448. Zoöl. Ser.

COSTA RICA OCELOT.

Type locality. Talamanca, Costa Rica.

Geogr. Distr. Costa Rica. Range unknown.

904. *Felis carrikeri* Allen.

Felis carrikeri Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1904, p. 47. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 448. Zoöl. Ser. Melanistic specimen.

CARRIKER'S OCELOT.

Type locality. Pozo Azul, Pirris Province, Costa Rica.

Geogr. Distr. Costa Rica. Range unknown.

905. *Felis tigrina* Erxleben.

Felis tigrina Erxl., Syst. Regn. Anim., I, 1777, p. 517. Elliot, Mon. Felidæ, 1881, pl. XIX. *Id.* Mamm. Middle Amer. & W.

Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 449. Zoöl. Ser.

Felis mitis F. Cuv., Hist. Nat. Mamm., II, 1820, pl. 137.

Felis brasiliensis F. Cuv., Hist. Nat. Mamm., 1820, pl. 139.

Felis macroura Max., Beitr. Naturg. Bras., II, 1826, p. 371.

Felis chati Griff., Anim. King., II, 1827, p. 479.

Felis elegans Less., Cent. Zool., 1830, p. 69, pl. 21.

THE MARGAY. *Cauzel* in Costa Rica.

Type locality. Unknown. "South America."

Geogr. Distr. Honduras to Paraguay.

906. *Felis glaucula* Thomas.

Felis glaucula Thomas, Ann. Mag. Nat. Hist., 7th Ser., XII, 1903, p. 235. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M.

Pub., IV, Pt. II, 1904, p. 450. Zoöl. Ser.

SMALL SPOTTED CAT.

Type locality. Beltran, State of Jalisco, Mexico.

Geogr. Distr. Unknown.

Felis pardinoides oncella Thomas.

Felis pardinoides oncella Thomas, Ann. Mag. Nat. Hist., 7th Ser., XII, 1903, p. 237. Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., IV, Pt. II, 1904, p. 450. Zoöl. Ser.

OUNCE-LIKE CAT.

Type locality. Volcan de Irazú, Costa Rica.

Geogr. Distr. Unknown.

907. *Felis jaguarondi* Fischer.

Felis jaguarondi Fisch., Zoogn., 1814, p. 228. Elliot, Mon.

Felidæ, 1881, pl. XIII. *Id.* Syn. N. Am. Mamm., F. C. M. Pub.,

II, 1901, p. 295. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W.

Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 451, fig. 83. Zoöl.

Ser.

JAGUARONDI. *Leon monero*, *Leon miquero* in Costa Rica.

Type locality. Paraguay.

Geogr. Distr. Rio Grande, Texas, southward to Paraguay, east of the Andes.

a. *Felis jaguarondi cacomitli* (Berlandier).

Felis cacomitli Berl., in Baird, Rep. U. S. & Mex. Bound. Surv., Mamm., II, 1857, p. 12.

Felis jaguarondi cacomitli Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 451. Zoöl. Ser.

GRAY JAGUARONDI.

Type locality. Matamoros, State of Tamaulipas, Mexico.

Geogr. Distr. Unknown.

b. *Felis jaguarondi tolteca* Thomas.

Felis jaguarondi tolteca Thomas, Ann. Mag. Nat. Hist., 7th Ser.,
I, 1898, p. 41. Elliot, Mamm. Middle Amer. & W. Indies,
F. C. M. Pub., IV, Pt. II, 1904, p. 452. Zoöl. Ser.

SINALOA JAGUARONDI.

Type locality. Tatemelas, State of Sinaloa, Mexico.

Geogr. Distr. State of Sinaloa, Mexico. Range unknown.

908. *Felis panamensis* Allen.

Felis panamensis Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1904,
p. 71. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M.
Pub., IV, Pt. II, 1904, p. 452. Zoöl. Ser.

BOQUERON JAGUARONDI.

Type locality. Boqueron, Chiriqui, Panama.

Geogr. Distr. Panama, Central America.

909. *Felis eyra* Fischer.

Felis eyra Fisch., Zoogn., 1814, p. 228. Elliot, Mon. Felidæ, 1881,
pl. xiv. *Id.* Syn. N. Am. Mamm., F. C. M. Pub., II, 1901,
p. 295. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies,
F. C. M. Pub., IV, Pt. II, 1904, p. 453. Zoöl. Ser.

THE EYRA. *Apache*, *Onza* in Mexico.

Type locality. Paraguay.

Geogr. Distr. Brownsville, Texas, south to Paraguay; east of
the Andes.

a. *Felis eyra apache* (Mearns).

Felis eyra Baird, Rep. U. S. & Mex. Bound. Surv., Mamm., 1857,
p. 88, pl. LXII, fig. 1. (nec Fischer.)

Felis apache Mearns, Proc. Biol. Soc. Wash., XIV, 1901, p. 150.

Felis eyra apache Elliot, Mamm. Middle Amer. & W. Indies,
F. C. M. Pub., IV, Pt. II, 1904, p. 453. Zoöl. Ser.

TAMAULIPAS EYRA.

Type locality. Rio Grande del Norte, State of Tamaulipas,
Mexico.

Geogr. Distr. State of Tamaulipas, Mexico. Range unknown.

910. *Felis fossata* Mearns.

Felis fossata Mearns, Proc. Biol. Soc. Wash., XIV, 1901, p. 150.

Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV,
Pt. II, 1904, p. 453. Zoöl. Ser.

YUCATAN EYRA.

Type locality. Merida, Yucatan.

Geogr. Distr. Yucatan. Range unknown.

Subgenus B. Felis. Pumas.

Felis concolor oregonensis (Rafinesque).

Felix! oregonensis Rafin., Atlantic Journ., I, 1832, p. 62.

Felis hipolestes Merr., Proc. Biol. Soc. Wash., XI, 1897, p. 219.

Felis hipolestes olympus Merr., Proc. Biol. Soc. Wash., XI, 1897, p. 220.

Felis hipolestes aztecus Merr., Proc. Wash. Acad. Scien., III, 1901, p. 592.

?**Felis aztecus browni* Merr., Proc. Biol. Soc. Wash., XVI, 1903, p. 73.

Felis concolor oregonensis Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 294. Zoöl. Ser. *Id.* Mamm. Middle Amer.

& W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 454, pls. XLIV, XLV. Zoöl. Ser.

NORTHWESTERN PUMA.

Type locality. Northwest coast.

Geogr. Distr. West coast of North America, west of Rocky Mountains south into northern Mexico.

911. *Felis cougar* Kerr.

Felis cougar Kerr, Anim. Kingd., 1792, p. 151.

Felis pennsylvanica Link, Beyt. Naturgesch., II, 1795, p. 90.

Felis concolor floridana Cory, Hunt. & Fish. Florida, 1896, p. 109.

Felis coryi Bangs, Proc. Biol. Soc. Wash., XIII, 1899, p. 15.

Felis concolor Auct., Eastern North America.

EASTERN PUMA.

Type locality. Pennsylvania.

Geogr. Distr. Adirondack Mountains, New York; Green Mountains, Vermont; higher Alleghanies of Pennsylvania, Maryland, and West Virginia; and Florida.

Genl. Char. Size various, medium to large; color various.

Color. General hue fulvous or tawny, changing with the seasons; belly rufous or whitish; ears behind blackish, upper lip with the usual black patch.

Measurements. Total length, 1828.8; tail, 609.6. Killed in the Blue Mountains, Berks County, Pennsylvania. See Forest and Stream, III, 1874, p. 67. No skull is available for measurements.

*I have not seen a specimen of this form, but I find nothing in Dr. Merriam's description to warrant a distinct rank. I have placed the name with a query. *Felis concolor* by some has been restricted to South America.

* *Felis bangsi costaricensis* Merriam.

Felis bangsi costaricensis Merr., Proc. Wash. Acad. Scien., III, 1901, p. 596. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 456. Zoöl. Ser.

CENTRAL AMERICAN PUMA.

Type locality. Boquete, Chiriqui, Panama.

Geogr. Distr. Unknown.

Subgenus C. Lynx. Lynx.

912. *Felis canadensis* (Kerr).

Lynx canadensis Kerr, Anim. Kingd., 1792, p. 32, desc. p. 157.

Felis canadensis Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 295. Zoöl. Ser.

CANADA LYNX.

Type locality. Eastern Canada.

Geogr. Distr. Boreal America, from Maine to 60° north latitude, and from the Atlantic to the Pacific Ocean, possibly not including Alaska.

a. *Felis canadensis mollipilosa* (Stone).

Lynx canadensis mollipilosus Stone, Proc. Acad. Nat. Scien. Phil., 1900, p. 48.

Felis canadensis mollipilosus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 295. Zoöl. Ser.

SOFT-FURRED LYNX.

Type locality. Wainwright Inlet, Point Barrow, Alaska.

Geogr. Distr. Alaska. Range undetermined.

b. *Felis canadensis subsolana* (Bangs).

Lynx subsolanus Bangs, Proc. Biol. Soc. Wash., XI, 1897, p. 49.

Felis canadensis subsolana Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 296. Zoöl. Ser.

EASTERN LYNX.

Type locality. Codroy, Newfoundland.

Geogr. Distr. Newfoundland. Range unknown.

Subgenus D. Cervaria.

913. *Felis rufa* Gldenstaedt.

Felis rufa (sic) Gldenst., Nov. Comm. Acad. Imp. Scien. St. Ptersb., XX, 1776, p. 484.

Felis rufa Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 296. Zoöl. Ser.

*Of a very doubtful subspecific standing.

RED LYNX. WILD CAT.

Type locality. New York.

Geogr. Distr. Maine to southern Georgia; west to Texas.

a. *Felis rufa texensis* (Allen).

Felis maculata Vig. & Horsf., Zoöl. Journ., IV, 1829, p. 381, pl. XIII.

Lynx texensis Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1895, p. 188.

Felis rufa maculata Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 296. Zoöl. Ser.

Felis rufa texensis Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 298, and Suppl., p. 506 (note). Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 458. Zoöl. Ser.

SPOTTED LYNX. WILD CAT. *Gato montes* in Mexico for all Lynxes.

Type locality. Mexico.

Geogr. Distr. Mexico to California.

b. *Felis rufa fasciata* (Rafinesque).

Lynx fasciatus Rafin., Am. Month. Mag., II, 1817, p. 46.

Felis rufa fasciata Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 297. Zoöl. Ser.

BARRED LYNX.

Type locality. "Northwest Coast," probably near the mouth of the Columbia River.

Geogr. Distr. Oregon and Washington.

c. *Felis rufa pallescens* (Merriam).

Lynx fasciatus pallescens Merr., N. Am. Faun., No. 16, 1899, p. 104.

Felis rufa pallescens Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 297. Zoöl. Ser.

PALLID LYNX.

Type locality. South side of Mount Adams, near Trout Lake, Skamania County, Washington.

Geogr. Distr. Region about Shasta Mountain, California; Oregon, and Washington.

d. *Felis rufa oculatea* (Bangs).

Lynx fasciatus oculus Bangs, Proc. N. Eng. Zoöl. Club, I, 1899, p. 23.

Felis rufa oculatea Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 297. Zoöl. Ser.

SHARP-SIGHTED LYNX.

Type locality. Nicasio, Marin County, California.*

Geogr. Distr. Coast region of California, north of San Francisco Bay.

e. * *Felis rufa baileyi* (Merriam).

Lynx baileyi Merr., N. Am. Faun., No. 3, 1890, p. 79.

Felis rufa baileyi Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 297. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 459, figs. 84, LX. Zoöl. Ser.

BAILEY'S LYNX.

Type locality. Moccasin Spring, Coconino County, Arizona.

Geogr. Distr. Arizona and northern Mexico. (State of Chihuahua.)

f. *Felis rufa eremica* (Mearns).

Lynx rufus eremicus Mearns, Proc. U. S. Nat. Mus., xx, 1897, p. 457.

Felis rufa eremica Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 298. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 458. Zoöl. Ser.

DESERT LYNX.

Type locality. New River, near Laguna Station, Colorado Desert, San Diego County, California.

Geogr. Distr. Eastern and Western Desert Tracts on the Mexican Line; probably in the States of Sonora and Chihuahua, Mexico.

g. *Felis rufa californica* (Mearns).

Lynx rufus californicus Mearns, Proc. U. S. Nat. Mus., xx, 1897, p. 458.

Felis rufa californica Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 298. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 458. Zoöl. Ser.

CALIFORNIA LYNX.

Type locality. San Diego, San Diego County, California.

Geogr. Distr. Pacific coast tract of California, and Lower California.

h. *Felis rufa floridana* (Rafinesque).

Lynx floridanus Rafin., Am. Month. Mag., II, 1817, p. 46.

Felis rufa floridana Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 299. Zoöl. Ser.

FLORIDA LYNX.

Type locality. Florida.

Geogr. Distr. Florida to Louisiana, Gulf Coast.

i. *Felis rufa escuinapæ* (Allen).

Lynx rufus (sic) *escuinapæ* Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1903, p. 614.

*Does this not = *F. maculata* Vigors and Horsfield ex Mexico?

Felis rufa escuinapæ Elliot, Mamm. Middle Amer. & W. Indies,
F. C. M. Pub., IV, Pt. II, 1904, p. 456. Zoöl. Ser.

ESCUINAPA LYNX.

Type locality. Escuinapa, State of Sinaloa, Mexico.

Geogr. Distr. State of Sinaloa, Mexico. Range unknown.

914. *Felis uinta* (Merriam).

Lynx uinta Merr., Proc. Biol. Soc. Wash., xv, 1902, p. 71.

UINTA MOUNTAINS LYNX.

Type locality. Bridger Pass, south slope of the Uinta Mountains, Uinta County, Wyoming.

Geogr. Distr. Utah. Range unknown.

Genl. Char. Size large; tail very long with two black subterminal bands, black tips. Skull long, narrow; sagittal crest present; ascending arm of maxillary broad, under jaw heavy, teeth small.

Color. Above buffy grizzled with gray and black; beneath white with black spots, these becoming bands on inner sides of arms and thighs; throat fulvous, crossed by a fulvous brown band; V-shaped black mark on anterior part of throat; tail with two or three blackish bands above and black tip.

Measurements. Hind foot, 200; tail, 195.

915. *Felis gigas* (Bangs).

Lynx gigas Bangs, Proc. Biol. Soc. Wash., xi, 1897, p. 50.

Felis gigas Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 299. Zoöl. Ser.

GIANT LYNX.

Type locality. Fifteen miles back of Bear River, Nova Scotia.

Geogr. Distr. Nova Scotia: Range unknown.

916. *Felis peninsularis* (Thomas).

Lynx rufus peninsularis Thomas, Ann. Mag. Nat. Hist., 7th Ser.,
I, 1898, p. 42.

Felis peninsularis Elliot, Mamm. Middle Amer. & W. Indies,
F. C. M. Pub., IV, Pt. II, 1904, p. 460. Zoöl. Ser.

LOWER CALIFORNIA LYNX.

Type locality. Santa Anita, Lower California, Mexico.

Geogr. Distr. Cape Region of Lower California, Mexico.

Family II. **Viverridæ. Mongoose, Civets, etc.**

Genus 113. **Herpestes. Ichneumon.**

Herpestes Illig., Prodr. Syst. Mamm. et Av., 1811, p. 135. Type
Viverra ichneumon Gmelin.

917. *Herpestes mungo* (Gmelin).

Viverra mungo Gmel., Syst. Nat., I, 1788, p. 84.

Herpestes mungo Elliot, Mamm. Middle Amer. & W. Indies,
F. C. M. Pub., IV, Pt. II, 1904, p. 462, figs. 85, LXI. Zool. Ser.

COMMON INDIAN MUNGOOSE.

Type locality. India.

Geogr. Distr. India. Introduced into the islands of Jamaica,
Porto Rico, St. Kitts, etc.

Family III. **Canidae. Dogs, Wolves, Foxes.**

Subfamily I. **Caninae.**

Genus 114. **Canis. Dogs, Wolves.**

Canis Linn., Syst. Nat., I, 1758, p. 38; and I, 1766, p. 56. Type
Canis familiaris Linnæus.

Lupus Frisch, Nat.-Syst. vierfüss. Thiere, in Tabellen, 14, Tab.
Gen., 1775.

Lyciscus H. Smith, Jard. Nat. Libr., Mamm., 2d ed., IX, 1839,
pp. 160-166.

918. *Canis pambasileus* Elliot.

Canis pambasileus Elliot, Proc. Biol. Soc. Wash., XVIII, 1905, p. 79.
AUTOCRAT TIMBER WOLF.

Type locality. Sushitna River, region of Mount McKinley, Alaska.
No. 13,481, Field Columbian Mus., Chicago.

Genl. Char. Color from nearly uniform black to white and black
in various mixtures. Skull: size larger than those of the timber
wolves of Canada or United States; ridge of sagittal and occipital
crest nearly on a level with frontal and with only a very slight descent
at occiput and very deep at that point; maxillæ very broad and
rounded posteriorly at junction with the frontals, much broader than
in *C. occidentalis*; nasals pointed posteriorly, but considerably broader
than those of southern timber wolves; premaxillæ extending con-
siderably over one-half the length of the nasals, while in *C. occiden-
talis* in some cases this bone does not reach over one third the length
of the nasals and occasionally not even that length; intertemporal
width considerable, with a deep median depression between the
frontals; the basisphenoid is very broad, and the postglenoid processes
very wide and flattened; the mandible is massive, heavy, very deep,
and of nearly uniform height on horizontal portion, with the inferior
outline nearly straight, very different from the curving outline of the

mandible of *C. occidentalis*: the coronoid process is very broad, high, of nearly equal width throughout, and with the posterior outline nearly straight, not curving like that of the southern species; teeth in both jaws large and heavy, exceeding in size those of *C. occidentalis* in the same proportion as do the skulls.

Color. Type: General hue of body, tail, legs, and feet black; the bases of the long hairs a smoky white, showing rather conspicuously on the sides; a pure white spot on the breast and lower part of the abdomen, and large spots on the fore part of chest behind fore legs grayish white; head and ears shining black; tail black, with a few white hairs at tip; claws black.

Other specimens are all black with very few grayish white hairs showing. From this sable hue, which in certain sections seems to be the prevailing one, the color varies in different mixtures of black and white to that in which the white is the prominent color with jet black lines on dorsal surface and neck. The tail of this style is mostly white with black mixed in places and with the tip black.

Measurements. Skull: total length, 263; Hensel, 233; zygomatic width, 143; intertemporal width, 64; lateral length of nasals, 97; median length, 85; width of rostrum at canines, 55.5; palatal length, 126; width of palate between sectorials at base, 64; length of upper tooth row, anterior edge of canine to posterior edge of last molar, 114; length of canine, 36; alveolar length of carnassial, 27; width of row of incisors at tips, 42; total length of mandible, 206; depth beneath carnassial, 32; height at coronoid process from lower edge of angle, 78; width of coronoid process at tip, 29.

919. **Canis griseus* (Sabine).

Canis lupus griseus Sabine, Frankl. Narr. Journ. Polar Sea, 1823, p. 654.

Canis lupus albus Sabine, Frankl. Narr. Journ. Polar Sea, 1823, p. 655.

Lupus griseus var. A. Richard., Faun. Bor. Amer., Mamm., 1, 1829, p. 66.

Lupus albus var. B. Richard., Faun. Bor. Amer., Mamm., 1, 1829, p. 68.

Lupus sticticus var. C. Richard., Faun. Bor. Amer., Mamm., 1, 1829, p. 68.

*The status of the North American Timber Wolves has never been satisfactorily determined, and it is not yet decided whether they are all of one species, or represent various species and races. *Canis griseus* of Sabine appears to be the earliest name for the Wolves north of Mexico, preceding *Canis albus* of the same author by one page of the publication in which they both appear. If the Timber Wolves of the North American continent prove eventually to be of the same species, *Canis mexicanus* Linnæus is apparently the name by which they will be known.

Lupus ater var. E. Richard., Faun. Bor. Amer., Mamm., 1, 1829, p. 70.

Canis occidentalis Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 300. (Part.) Zoöl. Ser.

ARCTIC WOLF.

Type locality. Cumberland House, N. L. 54°? Saskatchewan, Canada.

Geogr. Distr. Arctic America.

920. *Canis occidentalis* (Richardson).

Canis lupus occidentalis Richard., Faun. Bor. Amer., Mamm., 1, 1829, p. 60.

Canis occidentalis Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 300. (Part.) Zoöl. Ser.

WESTERN WOLF. TIMBER WOLF.

Type locality. — ?

Geogr. Distr. Western North America, Pacific Coast to Nebraska, and from Canada to Mexico. Rare east of the Mississippi.

921. *Canis nubilus* Say.

Canis nubilus Say, Long's Exped. Rocky Mts., 1, 1823, p. 169.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 300. Zoöl. Ser.

DUSKY WOLF.

Type locality. Vicinity of Council Bluffs, Pottawattamie County, Iowa.

Geogr. Distr. Nebraska northwestward through Idaho to Puget Sound. California?

922. *Canis mexicanus* Linnaeus.

Canis mexicanus Linn., Syst. Nat., 1, 1766, p. 60. Elliot, Mamm.

Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 11, 1904, p. 471, fig. LXII, pl. XLVI. Zoöl. Ser.

MEXICAN TIMBER WOLF. *Lobo* in Spanish America.

Type locality. Mexico.

Geogr. Distr. States of Sonora and Chihuahua, Mexico. Range unknown.

923. *Canis latrans* Say.

Canis latrans Say, Long's Exped. Rocky Mts., 1, 1823, p. 168.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 301, pl. XXIX. Zoöl. Ser.

BARKING COYOTE.

Type locality. Vicinity of Council Bluffs, Pottawattamie County, Iowa.

Geogr. Distr. Northern Valley of the Mississippi, westward on northern edge of plains to Rocky Mountains in Alberta Province, Canada.

924. *Canis lestes* Merriam.

Canis lestes Merr., Proc. Biol. Soc. Wash., xi, 1897, p. 25. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., ii, 1901, p. 301. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. II, 1904, p. 465. Zoöl. Ser.

ROBBER COYOTE.

Type locality. Toyabe Mountains, near Cloverdale, Nye County, Nevada.

Geogr. Distr. Northern Mexico, probably through Arizona and New Mexico to Washington and southern British Columbia.

925. *Canis nebrascensis* Merriam.

Canis pallidus Merr., Proc. Biol. Soc. Wash., xi, 1897, p. 24. (pre-occupied.)

Canis nebracensis! Merr., Science, N. Ser., viii, 1897, p. 782. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., ii, 1901, p. 301. Zoöl. Ser.

NEBRASKA COYOTE.

Type locality. Johnstown, Brown County, Nebraska.

Geogr. Distr. Great Plains, eastern Colorado to Canada.

926. *Canis frustror* Woodhouse.

Canis frustror Woodh., Proc. Acad. Nat. Scien. Phil., 1851, p. 147. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., ii, 1901, p. 302. Zoöl. Ser.

TRICKY COYOTE.

Type locality. Fort Gibson, at junction of the Neosho and Arkansas Rivers, Indian Territory.

Geogr. Distr. Gulf region of Texas from Nueces Bay northward, and possibly through the intervening States into Oklahoma and Indian Territory.

927. *Canis mearnsi* Merriam.

Canis mearnsi Merr., Proc. Biol. Soc. Wash., xi, 1897, p. 30. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., ii, 1901, p. 302. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. II, 1904, p. 468. Zoöl. Ser.

MEARNS' COYOTE.

Type locality. Quitobaquita, Pima County, Arizona.

Geogr. Distr. Southern Arizona south into State of Sonora, Mexico.

928. *Canis estor* Merriam.

Canis estor Merr., Proc. Biol. Soc. Wash., XI, 1897, p. 31. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 302. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 469. Zoöl. Ser.

NOLAND'S RANCH COYOTE.

Type locality. Noland's Ranch, San Juan River, San Juan County, Utah.

Geogr. Distr. Deserts of eastern California, Nevada, and Utah.

929. *Canis cagottis* (H. Smith).

Lyciscus cagottis H. Smith, Jard. Nat. Libr., Mamm., XVIII, 1839, p. 164.

Canis cagottis Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 466. Zoöl. Ser.

SMITH'S COYOTE.

Type locality. Rio Frio, between City of Mexico and Puebla, Mexico.

Geogr. Distr. States of Mexico and Oaxaca. Range unknown.

930. *Canis peninsulæ* Merriam.

Canis peninsulæ Merr., Proc. Biol. Soc. Wash., XI, 1897, p. 28. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 466. Zoöl. Ser.

LOWER CALIFORNIA COYOTE.

Type locality. Santa Anita, Cape St. Lucas, Lower California, Mexico.

Geogr. Distr. Cape Region of Lower California, Mexico.

931. *Canis vigilis* Merriam.

Canis vigilis Merr., Proc. Biol. Soc. Wash., XI, 1897, p. 33. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 467. Zoöl. Ser.

COLIMA COYOTE.

Type locality. Manzanillo, State of Colima, Mexico.

Geogr. Distr. State of Colima, Mexico. Range unknown.

932. *Canis goldmani* Merriam.

Canis goldmani Merr., Proc. Biol. Soc. Wash., XVII, 1904, p. 157.

GOLDMAN'S COYOTE.

Type locality. San Vincente, State of Chiapas, Mexico.

Geogr. Distr. State of Chiapas, Mexico. Range unknown.

Genl. Char. Size large; muzzle broad; frontals deeply sulcate; bullæ very large, flattened on outer side; teeth large, except lower carnassial, which is smaller than that of *C. cagottis*.

Color. Muzzle, top of head, ears, and legs fulvous; face grizzled grayish fulvous; back grizzled buffy gray and fulvous; under fur pale fulvous.

Measurements. Total length, 1,220; tail vertebræ, 355; hind foot, 216.

933. *Canis clepticus* Elliot.

Canis clepticus Elliot, Pub. Field Columb. Mus., III, 1903, p. 225.

Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 467. Zoöl. Ser.

THIEVISH COYOTE.

Type locality. Vallecitos, San Pedro Martir Mountains, Lower California, Mexico. Altitude, 9,000 feet.

Geogr. Distr. San Pedro Martir Mountains, Lower California, Mexico.

934. *Canis impavidus* Allen.

Canis impavidus Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1903, p. 609. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M.

Pub., IV, Pt. II, 1904, p. 468. Zoöl. Ser.

BOLD COYOTE.

Type locality. Rio del Bocas, State of Durango, Mexico. Altitude, 7,000 feet.

Geogr. Distr. State of Durango, Mexico. Range unknown.

935. *Canis microdon* Merriam.

Canis microdon Merr., Proc. Biol. Soc. Wash., XI, 1897, p. 29.

Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 469. Zoöl. Ser.

TAMAULIPAS COYOTE.

Type locality. Mier, on the Rio Grande, State of Tamaulipas, Mexico.

Geogr. Distr. State of Tamaulipas, Mexico. Range unknown.

936. *Canis ochropus* Eschscholtz.

Canis ochropus Eschsch., Zool. Atlas, III, 1829, pp. 1-2, pl. II.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 303.

Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 469. Zoöl. Ser.

OCHRACEOUS-FOOTED COYOTE.

Type locality. California.

Geogr. Distr. San Joaquin Valley, California, south into Lower California, Mexico.

Genus 115. **Vulpes. Red Foxes.**

- Vulpes** Frisch, Natur. Syst. vierfüß. Thiere, in Tabellen, Tab. Gen., 1775. Type *Canis vulpes* Linnæus.
Leucocyon Gray, Proc. Zool. Soc., 1868, p. 521.

936a. Vulpes harrimani Merriam.

- Vulpes harrimani* Merr., Proc. Wash. Acad. Scien., 1900, p. 14.
 Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 303.
 Zool. Ser.

HARRIMAN'S FOX.

- Type locality.* Kadiak Island, Alaska.
Geogr. Distr. Only known from Kadiak Island.

937. Vulpes fulva (Desmarest).

- Canis fulvus* Desm., Mamm., 1, 1820, p. 203.
Canis decussatus Desm., Mamm., 1, 1820, p. 203. (Cross Fox.)
Vulpes pennsylvanica Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 304. Zool. Ser.

RED FOX.

- Type locality.* Virginia.
Geogr. Distr. Canadian boundary to Georgia, westward to the Great Plains.

a. Vulpes fulva argentata (Shaw).

- Canis argentatus* Shaw, Gen. Zool., 1, 1800, p. 325.
Vulpes pennsylvanica argentata Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 304. Zool. Ser.

SILVER FOX. BLACK FOX.

- Type locality.* Louisiana!
Geogr. Distr. Boreal regions of North America.

b. Vulpes fulva macroura (Baird).

- Vulpes macrourus* Baird, Stansb. Expl. Great Salt Lake, Utah, 1852, p. 309.
Vulpes pennsylvanica macroura Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 305. Zool. Ser.

LONG-TAILED FOX.

- Type locality.* Wasatch Mountains, near Great Salt Lake, Utah.
Geogr. Distr. Nebraska to California south to Arizona and Oklahoma Territory.

c. Vulpes fulva rubricosa (Bangs).

- Vulpes pennsylvanica vafra* Bangs, Proc. Biol. Soc. Wash., 11, 1897, p. 53. (nec *V. vafra* Leidy.)
Vulpes pennsylvanica rubricosa Bangs, Scien., N. S., VII, 1898, p. 272. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 305. Zool. Ser.

REDDISH FOX.

Type locality. Digby, Nova Scotia, Canada.

Geogr. Distr. Nova Scotia. Range unknown.

d. *Vulpes fulva bangsi* (Merriam).

Vulpes rubricosa bangsi Merr., Proc. Wash. Acad. Scien., II, 1900, p. 667.

Vulpes pennsylvanica bangsi Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, Append., p. 434. Zoöl. Ser.

BANGS' RED FOX.

Type locality. Lance au Loup, Strait of Belle Isle, Labrador.

Geogr. Distr. Labrador.

938. *Vulpes alascensis* Merriam.

Vulpes alascensis Merr., Proc. Wash. Acad. Scien., II, 1900, p. 668.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, Append., p. 434. Zoöl. Ser.

ALASKA RED FOX.

Type locality. Andreafski, near mouth of the Yukon River, Alaska.

Geogr. Distr. Northern Alaska.

a. *Vulpes alascensis abietorum* Merriam.

Vulpes alascensis abietorum Merr., Proc. Wash. Acad. Scien., II, 1900, p. 669. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, Append., p. 434. Zoöl. Ser.

FIR-WOOD FOX.

Type locality. Stuart Lake, British Columbia, Canada.

Geogr. Distr. Interior of British Columbia, Canada, and probably northeastern Alaska.

939. *Vulpes deletrix* Bangs.

Vulpes deletrix Bangs, Proc. Biol. Soc. Wash., XII, 1898, p. 36.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 305. Zoöl. Ser.

DESTRUCTIVE FOX.

Type locality. Bay St. George, Newfoundland.

Geogr. Distr. Newfoundland. Labrador?

940. *Vulpes necator* Merriam.

Vulpes necator Merr., Proc. Wash. Acad. Scien., II, 1900, p. 664.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, Append., p. 433. Zoöl. Ser.

MURDERING FOX.

Type locality. Whitney Meadows, near Mount Whitney, Tulare County, California. Altitude, 9,500 feet.

Geogr. Distr. High Sierra, southern California.

941. *Vulpes cascadenis* Merriam.

Vulpes cascadenis Merr., Proc. Wash. Acad. Scien., 11, 1900, p. 665. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, Append., p. 433. Zoöl. Ser.

CASCADE MOUNTAINS FOX.

Type locality. Trout Lake, south base of Mt. Adams, Cascade Mountains, Skamania County, Washington.

Geogr. Distr. Cascade Range in Washington and Oregon, and northern Sierra in California.

942. *Vulpes regalis* Merriam.

Vulpes regalis Merr., Proc. Wash. Acad. Scien., 11, 1900, p. 672. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, Append., p. 432. Zoöl. Ser.

ROYAL FOX.

Type locality. Elk River, Sherburn County, Minnesota.

Geogr. Distr. Northern Plains from Dakota to Alberta; east of Manitoba and Minnesota. Range limits unknown.

943. *Vulpes kenaiensis* Merriam.

Vulpes kenaiensis Merr., Proc. Wash. Acad. Scien., 11, 1900, p. 670. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, Append., p. 432. Zoöl. Ser.

KENAI FOX.

Type locality. Kenai Peninsula, Alaska.

Geogr. Distr. Kenai Peninsula. Range unknown.

944. *Vulpes hallensis* Merriam.

Vulpes hallensis Merr., Proc. Wash. Acad. Scien., 11, 1900, p. 15. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 307. Zoöl. Ser.

HALL ISLAND FOX.

Type locality. Hall Island, Bering Sea, Alaska.

Geogr. Distr. Known only from Hall Island.

Vulpes lagopus innuitus Merriam.

Vulpes lagopus innuitus Merr., Proc. Biol. Soc. Wash., xv, 1902, p. 170.

SUMMIT ARCTIC FOX.

Type locality. Karogar River, Point Barrow, Alaska.

Geogr. Distr. Alaska to Baffin Bay, Arctic America.

Genl. Char. Braincase broader and more pyriform than that of *V. lagopus*, and tapering more abruptly behind broadest part.

Color. Like that of *V. lagopus*.

Measurements. Skull: basal length, 117; zygomatic width, 69; palatal length, 56; rostrum width at second premolar, 23.5; length of upper carnassial, 12.5; transverse diameter in outer side, 10.

Vulpes lagopus ungava Merriam.

Vulpes lagopus ungava Merr., Proc. Biol. Soc. Wash., xv, 1902, p. 170.

UNGAVA ARCTIC FOX.

Type locality. Fort Chimo, Ungava, Labrador.

Geogr. Distr. Labrador.

Genl. Char. Similar to *V. l. innuitus*, but slightly larger; "rostrum longer; frontal shield less flat; postorbital processes more pointed and less deeply excavated posteriorly; under jaw broader vertically and deeply bellied under sectorial."

Color. Like "*V. l. innuitus* and *V. lagopus*."

Measurements. Skull: basal length, 118; zygomatic breadth, 70; palatal length, 63; post palatal length, 55; width of rostrum at second premolar, 26; length of upper carnassial, 12.5; length of upper molar, 9.5.

945. *Vulpes pribilofensis* Merriam.

Vulpes pribilofensis Merr., Proc. Biol. Soc. Wash., xv, 1902, p. 171.

PRIBILOF ISLAND ARCTIC FOX.

Type locality. St. George Island, Pribilof Islands, Bering Sea.

Geogr. Distr. St. George and St. Paul Islands, Pribilof Island, Bering Sea.

Genl. Char. Size large. Skull elongate; frontals narrow; postorbital processes slightly developed; braincase long and flat; sagittal crest well developed; teeth as in *V. l. innuitus*.

Measurements. Skull: basal length, 127; zygomatic breadth, 72; palatal length, 66; postpalatal length, 61; width of rostrum at second premolar, 25; length of upper carnassial, 12.5; transverse diameter of first upper molar, 9.5.

946. *Vulpes beringensis* Merriam.

Vulpes beringensis Merr., Proc. Biol. Soc. Wash., xv, 1902, p. 171.

BERING ISLAND ARCTIC FOX.

Type locality. Bering Island, Bering Sea.

Genl. Char. Size large. Rostrum broad, massive; frontals high anteriorly and swollen on each side of median sulcus; premolars large and swollen.

Color. Not given.

Measurements. Skull: basal length, 123; zygomatic breadth, 71; palatal length, 66; postpalatal length, 56; width of rostrum at second premolar, 27.

947. *Vulpes velox* (Say).

Canis velox Say, in Long's Exped. Rocky Mts., I, 1823, p. 487.

Canis cinereo-argentatus Sabine, Frankl. Narr. Journ. Polar Sea, 1823, p. 658.

Canis microtus Reichenb., Wagn. in Weigm. Archiv, III, 1837; II, p. 162.

Vulpes velox Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 306. Zoöl. Ser.

SWIFT FOX.

Type locality. Vicinity of Cherry Creek, Laramie County, Wyoming.

Geogr. Distr. Colorado and Nebraska north to the plains of the Saskatchewan, Assiniboia, Canada.

a. Vulpes velox hebes Merriam.

Vulpes velox hebes Merr., Proc. Biol. Soc. Wash., xv, 1902, p. 73.

SLUGGISH FOX.

Type locality. Calgary, Alberta, Canada.

Geogr. Distr. Range unknown.

Genl. Char. Like *V. velox*, but larger and grayer. Skull larger, palate longer; mandible longer, heavier, more bellied under sectorial; coronoid process higher.

Measurements. Total length, 844; tail vertebræ, 312; hind foot, 130.

948. *Vulpes macrotis* Merriam.

Vulpes macrotis Merr., Proc. Biol. Soc. Wash., iv, 1888, p. 135.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 306.

Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 473, figs. 86, LXIII. Zoöl. Ser.

LARGE-EARED KIT FOX.

Type locality. Riverside, San Bernardino County, California.

Geogr. Distr. Southern California, south to States of Sonora and Chihuahua and Lower California, Mexico.

a. Vulpes macrotis neomexicana Merriam.

Vulpes macrotis neomexicanus! Merr., Proc. Biol. Soc. Wash., xv, 1902, p. 74.

NEW MEXICAN KIT FOX.

Type locality. San Andreas Range (about fifty miles north of El Paso, Texas), Donna Ana County, New Mexico.

Geogr. Distr. New Mexico and Arizona. Range unknown.

Genl. Char. Like *V. macrotis* but larger. Skull heavier, broader; rostrum thicker; frontals broader; zygomata more spreading; bullæ larger, with anteriorly projecting point on inner side; lower jaw decidedly heavier and more bellied; coronoid process higher.

Measurements. Total length, 820; tail vertebræ, 300; hind foot, 135.

949. *Vulpes mutica* Merriam.

Vulpes muticus! Merr., Proc. Biol. Soc. Wash., xv, 1902, p. 74.

CURTAILED FOX.

Type locality. Tracy, San Joaquin Valley, San Joaquin County, California.

Genl. Char. Similar to *V. macrotis*, but larger; hind foot and tail longer; chin and under lip blacker; top of head and middle of back with more reddish brown showing through the grayish tips; outer side of hind feet more fulvous; top of hind foot less white; pectoral collar more pronounced; black tip of tail less extensive on under side. Skull larger, broader, and more massive; under jaw much heavier and more deeply bellied under sectorial and last premolar; carnassial teeth and molars larger and much heavier; premolars thicker, but not otherwise larger.

Measurements. Total length, 950; tail vertebræ, 350; hind foot, 123. (Merr. l. c.)

950. *Vulpes arsipus* Elliot.

Vulpes arsipus Elliot, Pub. Field Columb. Mus., III, 1903, p. 256.

Zoöl. Ser.

SWIFT-FOOTED KIT FOX.

Type locality. Daggett, Mohave Desert, San Bernardino County, California.

Geogr. Distr. Southern California. Range unknown.

Genl. Char. Similar to *V. macrotis* but paler and smaller; post-orbital processes longer; pterygoid fossa narrower.

Color. Top of head mixed gray and brownish fulvous, darker than the back; upper parts of body pale grizzled gray; sides paler grading into buff; outer sides of fore legs and thighs, and down outside of hind legs to the toes, pale fulvous; narrow pectoral collar pale fulvous; black patch on side of nose from eye, and one on either side of chin; postocular stripe of brown; under parts, inner side of thighs and front of hind legs whitish; tail above pale gray tinged with buff, beneath buffy, tip brownish black; ears externally pale cinnamon and narrowly edged with white.

Measurements. Total length, 810; tail vertebræ, 310; hind foot, 128; ear from notch, 86. Skull: occipito-nasal length, 103; Hensel, 104; zygomatic width, 61; interorbital constriction, 20.5; across postorbital processes, 28; palatal length, 56; length of nasals, 39; length of upper molar series, 44; length of mandible, 82.5; length of lower molar series, anterior edge of first premolar to posterior edge of last molar, 47.5.

Genus 116. **Urocyon.**

Urocyon Baird, N. Am. Mamm., 1857, p. 121. Type *Canis virginianus* Erxleben = *Canis cinereo-argenteus* Müller.

951. Urocyon cinereo-argenteus (Müller).

Canis cinereoargenteus Müll., Natur. Suppl., 1776, p. 29. (Desc.)

Canis cinereoargenteus Schreb., Säugth., III, 1776, p. 361, pl. XCII, 1775.

Canis virginianus Erxl., Syst. Regn. Anim., 1777, p. 567.

Canis griseus Bodd., Elench. Anim., I, 1784, p. 97.

Urocyon cinereo-argenteus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 307. Zoöl. Ser.

GRAY FOX.

Type locality. Eastern North America. Exact place unknown.

Geogr. Distr. Eastern North America from Georgia north to New England States, west to Mississippi Valley.

a. Urocyon cinereo-argenteus floridanus Rhoads.

Urocyon cinereo-argenteus floridanus Rhoads, Proc. Acad. Nat. Scien. Phil., 1895, p. 42. Elliot, Syn. N. Am. Mamm., F. C. M.

Pub., II, 1901, p. 307. Zoöl. Ser.

FLORIDA GRAY FOX.

Type locality. Tarpon Springs, Hillsboro County, Florida.

Geogr. Distr. Florida generally, north into Georgia where it interbreeds with *U. cinereo-argenteus*.

b. Urocyon cinereo-argenteus scotti Mearns.

Urocyon cinereo-argenteus scotti Mearns, Bull. Am. Mus. Nat. Hist. N. Y., 1891, p. 236. Elliot, Syn. N. Am. Mamm.,

F. C. M. Pub., II, 1901, p. 308, pl. xxxi. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II,

1904, p. 477. Zoöl. Ser.

SCOTT'S GRAY FOX.

Type locality. Pinal County, Arizona.

Geogr. Distr. Southern California, Arizona, New Mexico, and northern Mexico. State of Chihuahua.

c. Urocyon cinereo-argenteus texensis Mearns.

Urocyon cinereo-argenteus texensis Mearns, Proc. U. S. Nat. Mus., xx, 1897, p. 459. Elliot, Syn. N. Am. Mamm., F. C. M. Pub.,

II, 1901, p. 308. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 478. Zoöl. Ser.

TEXAS GRAY FOX.

Type locality. San Pedro, near Eagle Pass, Maverick County, Texas.

Geogr. Distr. Texas.

d. *Urocyon cinereo-argenteus californicus* Mearns.

Urocyon cinereo-argenteus californicus Mearns, Proc. U. S. Nat. Mus., xx, 1897, p. 459. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 308. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 477. Zoöl. Ser.

CALIFORNIA GRAY FOX.

Type locality. San Jacinto Mountains, Riverside County, California. Altitude, 8,000 feet.

Geogr. Distr. Washington south into Lower California.

e. *Urocyon cinereo-argenteus townsendi* (Merriam).

Urocyon californicus townsendi Merr., N. Am. Faun., No. 16, 1899, p. 103.

Urocyon cinereo-argenteus townsendi Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 309. Zoöl. Ser.

TOWNSEND'S GRAY FOX.

Type locality. Baird, Shasta County, California.

Geogr. Distr. Known only from the type locality.

f. *Urocyon cinereo-argenteus inyoensis* Elliot.

Urocyon cinereo-argenteus inyoensis Elliot, Pub. Field Columb. Mus., III, 1904, p. 268. Zoöl. Ser.

INYO MOUNTAINS GRAY FOX.

Type locality. Beveridge Cañon, Inyo Mountains, Inyo County, California.

Geogr. Distr. Southern California. Range unknown.

Genl. Char. Paler than *U. c. californicus*; no ochraceous lines; black on lower back only.

Color. Upper parts, sides, and breast silvery gray, with black streaks on lower part of back; chin and nose above black; sides of neck and bar across lower part of neck buff; fore legs silvery gray above, wood brown on outer side and whitish on inner side; hind legs silvery gray above, drab gray on outer side, and whitish on inner side; under parts white; tail silvery gray slightly tinged in places with buff, and with a black line curving around it; ears brownish mixed with silver white hairs on the outside; inside white; head back of ears buffy.

Measurements. Total length, 950; tail vertebræ, 385; hind foot, 140; ear, 82. Skull: total length, 124; occipito-nasal length, 118;

Hensel, 118; zygomatic width, 66; interorbital width, 23.5; palatal length, 56; palatal arch to tip of pterygoid process, 20; anterior width of pterygoid fossa, 9; posterior width of basioccipital, 13; length of nasals, 40; length of upper tooth row, anterior edge of canines to posterior edge of last molar, 51; length of lower tooth row, 57; length of mandible, 88.

g. *Urocyon cinereo-argenteus ocythous* Bangs.

Urocyon cinereo-argenteus ocythous Bangs, N. Engl. Zool. Club, 1, 1899, p. 43. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 309. Zool. Ser.

SWIFT GRAY FOX.

Type locality. Platteville, Grant County, Wisconsin.

Geogr. Distr. Upper Mississippi Valley.

h. *Urocyon cinereo-argenteus borealis* Merriam.

Urocyon cinereo-argenteus borealis Merr., Proc. Biol. Soc. Wash., XVI, 1903, p. 74.

NEW ENGLAND GRAY FOX.

Type locality. Marlboro, seven miles from Monadnock, Cheshire County, New Hampshire.

Genl. Char. Larger than *U. cinereo-argenteus*. Skull with longer nasals, more slender posteriorly; palate and pterygoid fossa broader; last upper molar and fourth lower premolar, large.

Color. Like *U. cinereo-argenteus*.

Measurements. Skull: basal length, 126; Hensel 118.5; zygomatic width, 73; palatal length, 63; postpalatal length, 55; front of canine to back of last upper molar, 56.

i. *Urocyon cinereo-argenteus littoralis* (Baird).

Vulpes (Urocyon) littoralis Baird, N. Am. Mamm., 1857, p. 143.

Urocyon cinereo-argenteus littoralis Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, Suppl., p. 507. Zool. Ser.

SAN MIGUEL ISLAND GRAY FOX.

Type locality. San Miguel Island, Santa Barbara Islands, Santa Barbara County, California.

Geogr. Distr. Known only from San Miguel Island.

j. *Urocyon cinereo-argenteus santacruzæ* (Merriam).

Urocyon littoralis santacruzæ Merr., Proc. Biol. Soc. Wash., XVI, 1903, p. 75.

SANTA CRUZ ISLAND GRAY FOX.

Type locality. Santa Cruz Island, Santa Barbara Islands, Santa Barbara County, California.

Geogr. Distr. Known from type locality only.

Genl. Char. Like *U. c. littoralis*. Skull smaller, nasals narrower tapering posteriorly; frontal spines long; rostrum narrower; bullæ smaller, inflated; basioccipital and upper carnassial narrower.

Measurements. Average of 3. Total length, 708; tail vertebrae, 266; hind foot, 109. (Merr. l. c.)

k. Urocyon cinereo-argenteus guatemalæ (Miller).

Urocyon guatemalæ Miller, Proc. Acad. Nat. Scien. Phil., 1899, p. 278.

Urocyon cinereo-argenteus guatemalæ Elliot, Mamm. Middle Amer.

& W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 475. Zoöl. Ser.

GUATEMALAN GRAY FOX.

Type locality. Nentón, Guatemala.

Geogr. Distr. Tropical region of Guatemala and State of Chiapas, Mexico.

l. Urocyon cinereo-argenteus fraterculus Elliot.

Urocyon cinereo-argenteus fraterculus Elliot, Pub. Field Columb.

Mus., I, 1896, p. 80. Zoöl. Ser. *Id.* Mamm. Middle Amer. &

W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 475, figs. 87,

LXIV. Zoöl. Ser.

LITTLE GRAY FOX.

Type locality. San Felipe, Yucatan.

Geogr. Distr. Yucatan, Tehuantepec, Mexico. Range unknown.

m. Urocyon cinereo-argenteus parvidens (Miller).

Urocyon parvidens Miller, Proc. Acad. Nat. Scien. Phil., 1899,

p. 276.

Urocyon cinereo-argenteus parvidens Elliot, Mamm. Middle Amer.

& W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 476. Zoöl. Ser.

SMALL-TOOTHED FOX.

Type locality. Merida, Yucatan, Mexico.

Geogr. Distr. Yucatan, Mexico. Range unknown.

952. *Urocyon catalinæ* Merriam.

Urocyon catalinæ Merr., Proc. Biol. Soc. Wash., XVI, 1903, p. 74.

SANTA CATALINA ISLAND GRAY FOX.

Type locality. Santa Catalina Island, Santa Barbara Islands, Santa Barbara County, California.

Geogr. Distr. Santa Catalina Island only.

Genl. Char. Similar to *U. c. littoralis*, but nasals narrower, and not constricted in the middle, but broad posteriorly and obliquely truncated; frontal spines short.

Color. Like *U. c. littoralis*.

Measurements. Total length, 795; tail vertebrae, 298; hind foot, 112.

953. *Urocyon clementæ* Merriam.

Urocyon clementæ Merr., Proc. Biol. Soc. Wash., xvi, 1903, p. 75.
SAN CLEMENTE ISLAND GRAY FOX.

Type locality. San Clemente Island, Santa Barbara Islands, Santa Barbara County, California.

Geogr. Distr. San Clemente Island only.

Genl. Char. Similar to *U. c. littoralis*, but smaller; nasals broadest in the middle, tapering posteriorly; bullæ more inflated; carnassial and molars small.

Color. Like *U. c. littoralis*.

Measurements. Average of 6. Total length, 688; tail vertebrae, 250; hind foot, 100. (Dry skin.) (Merr. l. c.)

Family IV. Ursidae. Bears.

Subfamily I. Ursinae.

Genus 117. Ursus.

Ursus Linn., Syst. Nat., i, 1758, p. 47; i, 1766, p. 69. Type *Ursus arctos*! Linnæus.

Danis Gray, in Thoms. Ann. of Phil., xxvi, 1825, pp. 60, 339.

Thalarchtos! Gray, in Thoms. Ann. of Phil., xxvi, 1825, p. 62.

Thalassarctos! Gray, in Thoms. Ann. of Phil., xxvi, 1825, p. 339.

Euarctos! Gray, Proc. Zool. Soc., 1864, p. 692.

Subgenus A. Thalassarctus. Polar Bear.

954. *Ursus maritimus* Phipps.

Ursus maritimus Phipps, Voy. N. Pole, 1774, p. 185. Elliot, Syn.

N. Am. Mamm., F. C. M. Pub., ii, 1901, p. 310, pl. xxxii.

Zool. Ser.

POLAR BEAR.

Type locality. Spitzbergen.

Geogr. Distr. Arctic regions of both hemispheres.

Subgenus B. Ursus. Brown Bears.

955. *Ursus middendorffi* Merriam.

Ursus middendorffi Merr., Proc. Biol. Soc. Wash., x, 1896, p. 69.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., ii, 1901, p. 310.

Zool. Ser.

MIDDENDORFF'S BROWN BEAR.

Type locality. Kadiak Island, Alaska.

Geogr. Distr. Kadiak Island, Alaskan Peninsula, Kenai Peninsula, Prince William Sound, Sushitna, and Kuik River regions and the Alaska Mountains, Alaska.

956. *Ursus kidderi* Merriam.

Ursus kidderi Merr., Proc. Biol. Soc. Wash., xv, 1902, p. 78.

KIDDER'S BROWN BEAR.

Type locality. Chinitna Bay, Cook Inlet, Alaska.

Geogr. Distr. Unknown.

Genl. Char. Size large; claws rather short. Skull large, massive; lacrymal large, high, reaching to maxillary protuberance on inner side of orbit; teeth comparatively small; lower sectorial rectangular, two cusps only on inner side of heel of last upper molar.

Color. Yellowish brown, darkest on belly and legs. June.

Measurements. Skull: basal length, 301; occipito-sphenoid length, 82; length of upper molar series, 74; length of lower molar series, 70; width across upper incisors, 44.

957. *Ursus dalli* Merriam.

Ursus dalli Merr., Proc. Biol. Soc. Wash., x, 1896, p. 71, figs. 8, 9, pl. v, fig. 1; pl. vi, fig. 5. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 311. Zool. Ser.

Ursus sitkensis Merr., Proc. Biol. Soc. Wash., x, 1896, p. 73, pl. vi, fig. 1; pl. v, fig. 3.

DALL'S BROWN BEAR.

Type locality. Yakutat Bay, Alaska.

Geogr. Distr. Coast region between Baranof Island and Copper River, Alaska.

a. *Ursus dalli gyas* Merriam.

Ursus dalli gyas Merr., Proc. Biol. Soc. Wash., xv, 1902, p. 78.

Ursus merriami Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1902, p. 141.

GIANT BROWN BEAR.

Type locality. Pavlof Bay, Alaska Peninsula, Alaska.

Geogr. Distr. Unknown.

Genl. Char. Similar to *U. dalli* but larger; postpalatal region longer; frontals and vault of cranium higher, less flat; rostrum longer, more elevated; anterior nares larger, higher; teeth larger; heel of last upper molar cut off obliquely on outer side.

Measurements. Skull: basal length, 390; zygomatic breadth, 285; palatal length, 212; postpalatal length, 177; occipito-sphenoid length, 112.

958. *Ursus eulophus* Merriam.

Ursus eulophus Merr., Proc. Biol. Soc. Wash., xvii, 1904, p. 153.

ADMIRALTY ISLAND BROWN BEAR.

Type locality. Admiralty Island, southeastern Alaska.

Geogr. Distr. Unknown.

Genl. Char. Size large. Skull: sagittal crest high anteriorly; frontals much elevated posteriorly, rather narrow interorbitally; frontal shield long and high, sloping strongly upward from anterior third of nasals almost to fronto-parietal suture; braincase narrowed anteriorly; rostrum rather narrow; maxillæ long, reaching to beyond plane of nasals; interpterygoid fossa long and narrow; molars large; lower carnassial slender.

Measurements. None given!

959. **Ursus kenaiensis* Merriam.

Ursus kenaiensis Merr., Proc. Biol. Soc. Wash., xvii, 1902, p. 154.

KENAI PENINSULA BROWN BEAR.

Type locality. Cape Elizabeth, west end of Kenai Peninsula.

Geogr. Distr. Unknown.

Genl. Char. Size large. Skull broad, flat, and massive, with broad rostrum; zygomata wide; jugal broad anteriorly; frontals depressed, low posteriorly; postorbital large, projecting outwardly; palate very broad; nasals broad; anterior nares rather small; canines small; incisors and molars large.

Measurements. None given!

Subgenus C. Danis. Grizzly Bears.

960. *Ursus horribilis* Ord.

Ursus horribilis Ord, Guthr. Geog., 2d Amer. ed., II, 1815, p. 291, desc. p. 299. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 312, pl. XXXIII. Zool. Ser.

Ursus horribilis californicus Merr., Proc. Biol. Soc. Wash., x, 1896, p. 76.

GRIZZLY BEAR.

Type locality. Montana.

Geogr. Distr. California, and Rocky Mountains north of Wyoming and Utah. British Columbia to Alaska.

*The specific values of the Brown Bears of Alaska cannot be considered as satisfactorily established. The roaming habits of these animals frequently cause them to travel very great distances, even swimming wide rivers and estuaries of the sea. It is not probable, therefore, that closely allied animals, as are all those heretofore described, would be able to maintain their distinctive characters, as they must at certain seasons mingle promiscuously together, and bears, like pigs, are not particular in their choice of a mate. Great variations in skulls and in size of individuals occur among all bears. It is most probable that when these forms are treated monographically with ample materials, the number of species and races will be conspicuously reduced.

a. *Ursus horribilis alascensis* Merriam.

Ursus horribilis alascensis Merr., Proc. Biol. Soc. Wash., x, 1896,
p. 74. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901,
p. 312. Zoöl. Ser.

ALASKAN GRIZZLY BEAR.

Type locality. Norton Sound, Alaska.

Geogr. Distr. Not determined.

b. *Ursus horribilis horriæus* Baird.

Ursus horribilis var. *horriæus* Baird, Rep. Mex. Bound. Surv., II,
Pt. II, 1859, p. 24.

Ursus horribilis horriæus Elliot, Syn. N. Am. Mamm., F. C. M.
Pub., II, 1901, p. 312. Zoöl. Ser.

Ursus horriæus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M.
Pub., IV, Pt. II, 1904, p. 480, fig. LXV, pl. XLVII. Zoöl. Ser.

MEXICAN GRIZZLY BEAR.

Type locality. Los Nogales, State of Sonora, Mexico, near border
of Pima County, Arizona.

Geogr. Distr. Northern Mexico and southern New Mexico.
Range undetermined.

c. *Ursus horribilis phæonyx* Merriam.

Ursus horribilis phæonyx Merr., Proc. Biol. Soc. Wash., XVII, 1904,
p. 154.

DARK-CLAWED GRIZZLY BEAR.

Type locality. Comet Creek, a tributary of Forty Mile Creek,
near Eagle, Alaska.

Geogr. Distr. Unknown.

Genl. Char. "Similar to *U. horribilis*, but claws shorter, more
strongly curved (!) and dark blue black. Ears rather short and
densely haired on both sides. Skull: zygomata more spreading, muz-
zle broader, shorter, canines heavier, incisors larger."

Color. Back and legs very dark brown, almost blackish brown;
unworn tips of hairs on back grizzled; under parts and muzzle pale
brown.

Measurements. "Middle claw of fore foot, over curve, 93; top of
base to tip, 75; bottom of base to tip, 55.

961. *Ursus richardsoni* Swainson.

Ursus richardsoni Swainson, Anim. Menag., 1838, p. 54. Elliot,

Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 313. Zoöl. Ser.

BARREN GROUND GRIZZLY BEAR.

Type locality. Shore of the Arctic Ocean, Mackenzie, Canada.

Geogr. Distr. Barren Ground between Hudson Bay and Mackenzie

River, hilly regions east and north of Great Bear Lake; lower Rocky Mountains west of the Mackenzie; possibly to the upper tributaries of the Yukon.

Subgenus D. *Euarctus*. Black Bears.

962. *Ursus americanus* Pallas.

Ursus americanus Pallas, Spicileg. Zool., fasc. xiv, 1780, p. 5.
 Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 313,
 pl. xxxiv. Zool. Ser.

AMERICAN BLACK BEAR.

Type locality. Eastern North America.

Geogr. Distr. North America, eastern part, where forests are found, except in Louisiana, Florida, and Labrador.

a. *Ursus americanus sornborgeri* Bangs.

Ursus americanus sornborgeri Bangs, Amer. Nat., xxxii, 1898,
 p. 500. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901,
 p. 314. Zool. Ser.

SORNBORGER'S BLACK BEAR.

Type locality. Okak, Labrador.

Geogr. Distr. Labrador north to the limit of trees.

b. *Ursus americanus carlottæ* (Osgood).

Ursus carlottæ Osgood, N. Am. Faun., No. 21, 1901, p. 30, pl. iv,
 fig. 1.

QUEEN CHARLOTTE ISLANDS BLACK BEAR.

Type locality. Massett, Graham Island, Queen Charlotte Islands, British Columbia, Canada.

Geogr. Distr. Queen Charlotte Islands, British Columbia, Canada.

Genl. Char. Skull longer than that of *U. americanus*; rostrum more slender; cranium less arched; teeth larger and heavier; heel of last molar elongate.

Color. Black? Skull only known.

Measurements. Skull: basilar length, 282; palatal length, 155; postpalatal length, 127; zygomatic width, 193; width across post-orbital processes, 116; length of nasals, 82.

c. **Ursus americanus eremicus* Merriam.

Ursus americanus eremicus Merr., Proc. Biol. Soc. Wash., xvii,
 1904, p. 154.

*Dr. Merriam compares this type specimen with *Ursus amblyceps* Baird. As I have already shown, Pub. Field Columb. Mus., III, 1903, p. 235, Zool. Ser., this name of Baird's, as it represents no characters whatever, is a nomen nudum, and cannot be used. Baird had only a flat skin of a cinnamon bear, and no skull, and was unable to give any recognizable description.

DESERT BLACK BEAR.

Type locality. Sierra Guadalupe, State of Coahuila, Mexico.

Geogr. Distr. State of Coahuila, Mexico, possibly north to Davis Mountain, Texas.

Genl. Char. Frontals depressed, flat; nasals nearly straight; rostrum slender anteriorly; naked nose pad very long.

Color. Black; the under fur dark brown; muzzle dark brown.

Measurements. Total length, 1.290; tail vertebræ, 66; hind foot, 210.

963. *Ursus floridanus* Merriam.

Ursus floridanus Merr., Proc. Biol. Soc. Wash., x, 1896, p. 81.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 314.

Zoöl. Ser.

FLORIDA BLACK BEAR.

Type locality. Key Biscayne, Dade County, Florida.

Geogr. Distr. Everglades of Florida.

964. *Ursus luteolus* Griffith.

Ursus luteolus Griffith, Carniv. Anim., 1821, p. 236. Elliot, Syn.

N. Am. Mamm., F. C. M. Pub., II, 1901, p. 314. Zoöl. Ser.

LOUISIANA BLACK BEAR.

Type locality. Louisiana.

Geogr. Distr. Louisiana to Texas. Range undetermined.

965. *Ursus altifrontalis* Elliot.

Ursus altifrontalis Elliot, Pub. Field Columb. Mus., III, 1903, p. 234.

NORTHWESTERN BLACK BEAR.

Type locality. Shore of Lake Crescent, Clallam County, Washington.

Geogr. Distr. Washington. Range unknown.

Genl. Char. Skull: forehead very broad and high, bulging on the sides anteriorly, rounded above; nasals long, broad, elevated posteriorly and extending considerably beyond the ends of the maxillæ; outline of skull above convex, highest at anterior portion of frontals; occipital crest prominent; zygomata widely spread; basioccipital and basisphenoid very broad, the latter rounded anteriorly; pterygoid fossa short; muzzle broad, heavy, greatly swollen at base of canines.

Color. Black; nose tan color

Measurements. Skull: total length, 300; occipito-nasal length, 287; zygomatic width, 190; interorbital constriction, 75; width across postorbital processes, 106; length of nasals, 80; width of nasals anteriorly, 32; palatal length, 138; width of palate at posterior end

of last molar, 47; at anterior end of first molar, 45; at posterior edge of canines, 40; breadth of muzzle at outer sides of canines, 68; greatest breadth of basioccipital, 49; of basisphenoid, 30; palatal arch to end of pterygoid, 50; length of crown of last molar, 27; width, 15; from anterior edge of canine to posterior edge of last molar, alveolar border, 100; length of mandible from angle to symphysis on top, 202; height at condyle, 32; at coronoid process, 86; length of lower molar series, alveolar border, 66.

966. *Ursus hylodromus* Elliot.

Ursus hylodromus Elliot, Pub. Field Columb. Mus., III, 1903, p. 257.

Zoöl. Ser.

ALBERTA BLACK BEAR.

Type locality. Alberta, Canada.

Genl. Char. Skull: forehead prominent, elevated above the face; highest part of braincase anterior to line from the roots of the zygomata; width across postorbital processes very considerable; nasals concave in outline, posterior ends elevated; braincase greatly swollen on sides outward; zygomatic arches moderately expanded; narial opening very large; occipital crest prominent; bullæ wider than long; tubular meatus elongate and narrow; pterygoid fossa broad, narrowest anteriorly at palatal arch; the pterygoid processes broad and turned inward at their tips; palate of equal width to last molar where it narrows gradually to the pterygoid fossa; mandible very heavy, upper end of coronoid process curving downward forming a hook.

Measurements. Skull: total length, 312; occipito-nasal length, 257; Hensel, 270; zygomatic width, 173; width across postorbital processes, 96; between orbits, 69; greatest width of braincase, 102; length of nasals, 76; width of basioccipital, 47; of basisphenoid, 36; length of pterygoid fossa, 46; posterior width at processes, 24; palatal length, 153; width between last molars, 45; length of upper molars, alveolar border, 67; length of mandible, 217; depth at center of second molar, 37.5; height at coronoid process, 88; breadth of coronoid process at hook, 40; beneath hook, 41; length of three lower molars, alveolar border, 65.

967. *Ursus machetes* Elliot.

Ursus machetes Elliot, Pub. Field Columb. Mus., III, 1903, p. 235.

Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M.

Pub., IV, Pt. II, 1904, p. 481, pls. XLVIII, XLIX. Zoöl. Ser.

FIGHTING BEAR.

Type locality. Casas Grandes, Sierra Madre, State of Chihuahua, Mexico.

Geogr. Distr. Northern Mexico.

968. *Ursus emmonsii* (Dall).

Ursus americanus var. *emmonsii* Dall, Science, N. S., II, 1895, p. 87.

Ursus emmonsii Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 315. Zoöl. Ser.

EMMONS' GLACIER BEAR.

Type locality. Saint Elias Alps, near Yakutat Bay, Alaska.

Geogr. Distr. Saint Elias range of mountains southeasterly to Juneau. Exact range unknown.

969. *Ursus kermodei* Hornaday.

Ursus kermodei Hornaday, 9th Ann. Rept. N. York Zoöl. Soc., 1905, p. 6.

INLAND WHITE BEAR.

Type locality. Gribble Island, Lat. $53^{\circ} 25'$, Long. 129° W., British Columbia, Canada.

Geogr. Distr. District south of Skeena River to Rivers Inlet and Kitimat Arm, Gribble Island, British Columbia, Canada. The distribution only imperfectly known.

Genl. Char. Size small; color white.

Color. Clear creamy white down to the roots of the hair; the cream tint deepest on head, upper neck, and fore legs.

Measurements. Skin. Total length, 1384.3; ear, 76.2; Skull: length of "molar tooth row, including large premolar," 45.4.

Family V. *Procyonidae*. Raccoons, Coatis, etc.**Subfamily I. *Procyoninae*.****Genus 118. *Bassariscus*.**

Bassariscus Coues, Science, IX, 1887, p. 516. *Type Bassaris astuta* Lichtenstein.

Bassariscus Licht., Isis, 1831. (nec Hübner, 1816.)

Wagneria Jentink, Notes Leyden Mus., VIII, 1886, p. 129.

970. *Bassariscus astutus* (Lichtenstein).

Bassariscus astutus Licht., Isis, XXIV, 1831, p. 513.

Bassariscus astutus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 316. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, p. 484, figs. 88 and LXVI. Zoöl. Ser.

COMMON RACCOON-FOX. *Cacomistl* in Mexico. *Cat Squirrel* in Texas.

Type locality. Southern Mexico.

Geogr. Distr. California through Arizona, New Mexico, and Texas; southern Mexico.

a. *Bassariscus astutus raptor* (Baird).

Bassariscus raptor Baird, Rep. U. S. Mex. Bound. Surv., Mamm., 11, 1850, p. 19.

Bassariscus flavus oregonus Rhoads, Proc. Acad. Nat. Scien. Phil., 1893, p. 416.

Bassariscus astutus raptor Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 316, pl. xxxv. Zool. Ser.

Bassariscus astutus oregonus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 317. Zool. Ser.

ROBBIE RACCOON-FOX.

Type locality. California?

Geogr. Distr. California and Oregon.

b. *Bassariscus astutus flavus* Rhoads.

Bassariscus astutus flavus Rhoads, Proc. Acad. Nat. Scien. Phil., 1893, p. 417. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 316. Zool. Ser.

Id. Mamm. Middle Amer. & W. Indies, F. C. M. Pub., 14, Pt. 11, 1904, p. 485. Zool. Ser.

TAWNY RACCOON-FOX.

Type locality. Texas.

Geogr. Distr. Texas south into northern Mexico.

971. *Bassariscus saxicola* Merriam.

Bassariscus saxicola Merr., Proc. Biol. Soc. Wash., 11, 1897, p. 185. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., 14, Pt. 11, 1904, p. 485. Zool. Ser.

ESPIRITU SANTO RACCOON-FOX.

Type locality. Espiritu Santo Island, Lower California, Mexico.

Geogr. Distr. Known only from the type locality.

972. *Bassariscus* * *sumichrasti* (Saussure).

Bassariscus sumichrasti Sauss., Rev. Mag. Zool., 2me Sér., 11, 1860, p. 7, pl. 1.

Bassariscus variabilis Peters, Monatsb. K. Preuss. Akad. Wiss. Berl., 1874, p. 704, pls. 1, 11.

Bassariscus monticola Cordero, La Natural., 11, 1875, p. 270, pl. —.

*In the Mammals of Middle America and the West Indies, Pt. 11, p. 487, *Bassariscus sumichrasti* was given as a synonym of *annulatus* Wagner. On further investigation I am inclined to adopt Mr. Thomas' view, Am. Mag. Nat. Hist., 7th Ser., 11, 1903, p. 380, that Wagner's type was too young to definitely determine whether or not it was the same as Saussure's species, and therefore *B. sumichrasti* would assume its specific rank and the synonyms placed under *annulatus* must be restored to it. A description of *B. sumichrasti* is, therefore, here given.

SAUSSURE'S RACCOON-FOX. *Tepichiclu del Cofre de Perote, Cacomistle de Monte* in Mexico.

Type locality. Tehuantepec, Mexico.

Geogr. Distr. Unknown.

Genl. Char. Skull longer than that of *B. astutus*; zygomatic arches more bowed and heavier; sagittal crest present; the carnassial is shorter than that of the species named, the heel less oblique and pointed, and the second premolar is longer.

Color. General hue fawn mixed with black; flanks yellowish spotted with black; under parts yellowish. Head on top gray varied with black; nose brownish black; a buff spot above and below eye; cheeks gray spotted with black; throat and sides of neck whitish buff; chin blackish; a blackish buff band on inner side of fore legs, outer side of fore legs grizzled; outer side of hind legs like back, a mixed buff and black; feet blackish, becoming grayish buff at the toes; tail black, with four or five grayish buff rings on basal half, covered by the long hairs of the black portion; ears grayish buff at tips, brown at base.

Measurements. Total length, 782; tail vertebræ, 431.8.

a. *Bassariscus sumichrasti notinus* Thomas.

Bassariscus sumichrasti notinus Thomas, Ann. Mag. Nat. Hist., 7th Ser., XI, 1903, p. 379. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 485. Zoöl. Ser.

BOQUETE RACCOON-FOX.

Type locality. Boquete, Chiriqui, Panama.

Geogr. Distr. Panama, Central America. Range unknown.

973. *Bassariscus albipes* Elliot.

Bassariscus albipes Elliot, Pub. Field Columb. Mus., III, 1903, p. 258. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 486. Zoöl. Ser.

WHITE-FOOTED RACCOON-FOX.

Type locality. Near Vera Cruz, State of Vera Cruz, Mexico.

Geogr. Distr. State of Vera Cruz, Mexico.

974. *Bassariscus annulatus* (Wagner).

Paradoxurus annulatus Wagn., Schreb. Säugth., Suppl., II, 1841, p. 353.

Bassariscus annulatus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 487. Zoöl. Ser.

SOUTHERN RACCOON-FOX. *Tepachiche del Cofre de Perote, Cacomistl de Monte* in Mexico, *Muyus* in Guatemala.

Type locality. Unknown.

Geogr. Distr. Mexico to Costa Rica.

Genus 119. **Bassaricyon.**

Bassaricyon Allen, Proc. Acad. Nat. Scien. Phil., 1876, p. 20. Type
Bassaricyon gabbii Allen.

975. Bassaricyon gabbii Allen.

Bassaricyon gabbii Allen, Proc. Acad. Nat. Scien. Phil., 1876, p.
23. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub.,
IV, Pt. II, 1904, p. 489, fig. 89. Zool. Ser.

GABB'S COON.

Type locality. Costa Rica.

Geogr. Distr. Unknown.

Genus 120. **Procyon.**

Procyon Storr, Prodr. Meth. Mamm., 1780, p. 35, Tab. A. Type
Ursus lotor Linnæus.

Euprocyon Gray, Proc. Zool. Soc., 1864, pp. 705-706.

976. Procyon lotor (Linnæus).

Ursus lotor Linn., Syst. Nat., I, 1758, p. 48.

Procyon lotor Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901,
p. 317, fig. 62. Zool. Ser.

COMMON RACCOON.

Type locality. Eastern United States.

Geogr. Distr. Eastern North America from Canada to western
Georgia, and west to the Rocky Mountains north of Texas.

*a. *Procyon lotor elucus* Bangs.

Procyon lotor elucus Bangs, Proc. Bost. Soc. Nat. Hist., 1898, p.
219. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p.
319. Zool. Ser.

SLEEPY RACCOON.

Type locality. Oak Lodge, opposite Micco, Brevard County,
Florida.

Geogr. Distr. Eastern Georgia and Florida.

b. Procyon lotor hernandezii (Wagler).

Procyon hernandezii Wagl., Isis, XXIV, 1831, p. 514.

Procyon nivea Gray, Charls. Mag. Nat. Hist., I, 1837, p. 580.

Procyon psora Gray, Ann. Mag. Nat. Hist., 1st Ser., 1842, p. 261.

*Doubtfully separable from *P. lotor*. See Elliot, Pub. Field Columb.
Mus., III, 1901, p. 51, under *P. lotor*.

Procyon lotor hernandezi Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 317. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 491, fig. LXVII, pl. L. Zoöl. Ser.

HERNANDEZ'S RACCOON.

Type locality. Southern Mexico.

Geogr. Distr. Puget Sound, Washington, west of the mountains, south and west to Texas and south to southern Mexico.

c. *Procyon lotor pallidus* (Merriam).

Procyon pallidus Merr., Proc. Biol. Soc. Wash., XIII, 1900, p. 151.

Procyon lotor pallidus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 318. Zoöl. Ser.

PALE RACCOON.

Type locality. New River, Colorado Desert, San Diego County, California.

Geogr. Distr. Southern California. Range unknown.

d. *Procyon lotor pacificus* (Merriam).

Procyon psora pacifica Merr., N. Am. Faun., No. 16, 1899, p. 107.

Procyon lotor pacificus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 319. Zoöl. Ser.

PACIFIC COAST RACCOON.

Type locality. Lake Kichelos, Kittitas County, Washington. Altitude, 8,000 feet.

Geogr. Distr. Puget Sound and basal slopes of northern Cascades, Washington.

e. *Procyon lotor insularis* Merriam.

Procyon lotor insularis Merr., Proc. Biol. Soc. Wash., XII, 1898, p. 17. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M.

Pub., IV, Pt. II, 1904, p. 492. Zoöl. Ser.

ISLAND RACCOON.

Type locality. Maria Madre Island, Tres Marias Islands, State of Jalisco, Mexico.

Geogr. Distr. Tres Marias Islands, State of Jalisco, Mexico.

977. *Procyon pygmaeus* Merriam.

Procyon pygmaeus Merr., Proc. Biol. Soc. Wash., XIV, 1901, p. 101. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M.

Pub., IV, Pt. II, 1904, p. 490. Zoöl. Ser.

LITTLE RACCOON.

Type locality. Island of Cozumel, Yucatan, Mexico.

Geogr. Distr. Known only from the type locality.

978. *Procyon maynardi* Bangs.

Procyon maynardi Bangs, Proc. Biol. Soc. Wash., XII, 1898, p. 92. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 490. Zoöl. Ser.

MAYNARD'S RACCOON.

Type locality. New Providence Island, Bahama Islands.

Geogr. Distr. Known only from the type locality.

Subgenus A. *Euprocyon*.**979. *Procyon cancrivorus* (Cuvier).**

Ursus cancrivorus Cuv., Tab. Élém. Hist. Nat., 1798, p. 113.

Procyon cancrivorus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 492, fig. 90. Zoöl. Ser.

CRAB-EATING RACCOON.

Type locality. Cayenne, French Guiana.

Geogr. Distr. Tres Marias Islands, State of Jalisco, Mexico. (Alston.) Panama; northern South America.

Genus 121. *Nasua*. *Coatis*.

***Nasua* Storr, Prodr. Meth. Mamm., 1780, p. 35. Type *Viverra nasua* Linnæus.**

980. *Nasua nelsoni* Merriam.

Nasua nelsoni Merr., Proc. Biol. Soc. Wash., XIV, 1901, p. 100.

Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV., Pt. II, 1904, p. 495. Zoöl. Ser.

NELSON'S COATI.

Type locality. Cozumel Island, Yucatan, Mexico.

Geogr. Distr. Known only from the type locality.

981. *Nasua nasica* (Linnæus).

Viverra narica (sic) Linn., Syst. Nat., I, 1766, p. 64.

Nasua nasica Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 497, figs. 91 and LXVIII. Zoöl. Ser.

COATI. *Pisoti*, *Tejon*, in Spanish America.

Type locality. "In America."

Geogr. Distr. Northern boundary of Mexico generally into Central America. Exact range not yet well defined.

a. *Nasua nasica molaris* (Merriam).

Nasua narica! molaris Merr., Proc. Biol. Soc. Wash., XV, 1902, p. 68.

Nasua nasica molaris Elliot, Mamm. Middle Amer. & W. Indies,
F. C. M. Pub., IV, Pt. II, 1904, p. 497. Zoöl. Ser.

MANZANILLO COATI.

Type locality. Manzanillo, State of Colima, Mexico.

Geogr. Distr. State of Colima, Mexico. Range unknown.

b. *Nasua nasica bullata* (Allen).

Nasua narica! bullata Allen, Bull. Am. Mus. Nat. Hist. N. Y.,
1904, p. 48.

Nasua nasica bullata Elliot, Mamm. Middle Amer. & W. Indies,
F. C. M. Pub., IV, Pt. II, 1904, p. 497. Zoöl. Ser.

DARK COATI.

Type locality. Pozo Azul, Pirris Province, Costa Rica.

Geogr. Distr. Costa Rica, Central America. Range unknown.

c. *Nasua nasica panamensis* (Allen).

Nasua narica! panamensis Allen, Bull. Am. Mus. Nat. Hist. N. Y.,
1904, p. 51.

Nasua nasica panamensis Elliot, Mamm. Middle Amer. & W.
Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 498. Zoöl. Ser.

BOQUERON COATI.

Type locality. Boqueron, Chiriqui, Panama.

Geogr. Distr. Panama, Central America.

d. *Nasua nasica yucatanica* (Allen).

Nasua narica! yucatanica Allen, Bull. Am. Mus. Nat. Hist. N. Y.,
1904, p. 52.

Nasua nasica yucatanica Elliot, Mamm. Middle Amer. & W.
Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 498. Zoöl. Ser.

YUCATAN COATI.

Type locality. Chichen Itza, Yucatan, Mexico.

Geogr. Distr. Yucatan, Mexico. Range unknown.

e. *Nasua nasica pallida* (Allen).

Nasua narica! pallida Allen, Bull. Am. Mus. Nat. Hist. N. Y.,
1904, p. 53.

Nasua nasica pallida Elliot, Mamm. Middle Amer. & W. Indies,
F. C. M. Pub., IV, Pt. II, 1904, p. 498. Zoöl. Ser.

PALLID COATI.

Type locality. Near Guadalupe y Calva, Sierra Nevada, State of
Chihuahua, Mexico.

Geogr. Distr. State of Chihuahua, Mexico. Range unknown.

Genus 122. ***Potos. Kinkajous.**

Potos Cuv. & Geoff., Méth. Mamm., in Mag. Encyl., II, 1795, p. 187.

Type *Viverra caudivoluta* Schreber.

Cercoleptes Illig., Prodr. Syst. Mamm. et Av., 1811, p. 127.

Potos flavus aztecus Thomas.

Potos flavus aztecus Thomas, Ann. Mag. Nat. Hist., 7th Ser., IX, 1902, pp. 266, 268, desc. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 500, figs. (*Potos flavus*) 92 and LXIX. Zool. Ser.

AZTEC KINKAJOU. *Martica* in Mexico; *Micoleon* in Guatemala.

Type locality. Atoyca, State of Vera Cruz, Mexico.

Geogr. Distr. Central America, south to Guatemala.

Potos flavus chiriquensis Allen.

Potos flavus chiriquensis Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1904, p. 72. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 501. Zool. Ser.

CHIRIQUE KINKAJOU.

Type locality. Boqueron, Chiriqui, Panama.

Geogr. Distr. Panama. Range unknown.

Family VI. **Mustelidæ. Badgers, Weasels, Otters, etc.**Subfamily I. **Melinæ.**Genus 123. **Taxidea. Badgers.**

Taxidea Storr, Prodr. Meth. Mamm., 1780, p. 34. Type *Ursus taxus* Schreber.

982. Taxidea taxus (Schreber).

Ursus taxus Schreb., Säugth., III, 1778, p. 520.

Meles taxus var. *B. americanus* Bodd., Elench. Anim., I, 1785, p. 136.

Ursus labradoria Gmel., Syst. Nat., I, 1788, p. 102.

Meles jeffersoni Harl., Faun. Am., 1825, p. 309.

Taxidea americana Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 320, fig. 63. Zool. Ser.

**Potos flavus megalotus* (Martin), which Thomas has restricted to Colombia and Costa Rica, has been reconsidered by Allen, who assigns the name *megalotus* to the form found in eastern Colombia.

AMERICAN BADGER.

Type locality. Hudson Bay.

Geogr. Distr. Hudson Bay, British North America, in north latitude 58°, and in the United States from Wisconsin south to Oklahoma Territory, west to Rocky Mountains. Formerly it was found as far east as Ohio.

a. Taxidea taxus neglecta (Mearns).

Taxidea americana neglecta Mearns, Bull. Am. Mus. Nat. Hist. N. Y., 1891, p. 250. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 321. Zoöl. Ser.

CALIFORNIA BADGER.

Type locality. Fort Crook, Shasta County, California.

Geogr. Distr. California.

b. Taxidea taxus berlandieri (Baird).

Taxidea berlandieri Baird, N. Am. Mamm., 1857, p. 205.

Taxidea americana berlandieri Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 321. Zoöl. Ser.

Taxidea taxus berlandieri Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 503, figs. 93 and LXX. Zoöl. Ser.

BERLANDIER'S BADGER.

Type locality. Llano Estacado, Texas, near border of New Mexico.

Geogr. Distr. Texas, Oklahoma Territory, New Mexico, and Arizona, and State of Tamaulipas, Mexico.

c. Taxidea taxus infusca Thomas.

Taxidea taxus infusca Thomas, Proc. Zoöl. Soc., 1898, p. 899. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 505. Zoöl. Ser.

LOWER CALIFORNIA BADGER.

Type locality. Santa Anita, Lower California, Mexico.

Geogr. Distr. Known only from the type locality.

Genus 124. **Mephitis. Skunks.**

Mephitis Cuv., Leçons d'Anat. Comp., Class Mamm., I, 1800, Tabl.

1. Type? "Les Moufettes."

Chincha Less., Nouv. Tabl. Règn. Anim., 1842, p. 67.

Leucomitra Howell, N. Am. Faun., No. 20, 1901, p. 39, pls. IV,

Subgenus A. *Chincha*. Large-striped Skunks.

983. *Mephitis mephitis* (Schreber).

Viverra mephitis Schreb., Säugth., III, 1776, p. 444, tab. 121.

Mephitis mephitica Elliot, Syn. N. Am. Mamm., F. C. M. Pub.,

II, 1901, p. 322, fig. 64. (Part.) Zoöl. Ser.

CANADA SKUNK.

Type locality. North America.

Geogr. Distr. Eastern Canada—Nova Scotia, Quebec, and northern Ontario, west and north to Oxford House, Keewatin, Canada.

984. *Mephitis hudsonia* (Richardson).

Mephitis americana var. *hudsonica* Rich., Faun. Bor. Amer., I Mamm., 1829, p. 55.

Mephitis mephitica hudsonia Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 322. Zoöl. Ser.

NORTHERN PLAINS SKUNK.

Type locality. Plains of the Saskatchewan.

Geogr. Distr. Western Canada from Manitoba to British Columbia (east of the Cascades); south in the United States to Colorado, Nebraska, and Minnesota.

985. *Mephitis *olida* Boitard.

Mephitis putida G. Cuv., Tab. Élem. Hist. Nat. Anim., 1798, pp. 116, 117. (Indeterminable.) Boitard, Jard. Plantes, Paris, Mamm., 1842, p. 147. (Untenable.)

Mephitis olida Boitard, Jard. Plantes, Paris, Mamm., 1842, p. 147.

Mephitis fetidissima Boitard, Jard. Plantes, Paris, Mamm., 1842, p. 147.

Mephitis mephitica Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 322. (Part.) Zoöl. Ser.

EASTERN SKUNK.

Type locality. New Jersey.

Geogr. Distr. New England and Middle Atlantic States, south to Virginia; west to Indiana.

986. *Mephitis elongata* (Bangs).

Mephitis mephitica elongata Bangs, Proc. Bost. Soc. Nat. Hist., 1895, p. 531.

Mephitis elongata Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 324. Zoöl. Ser.

*As both *putida* Cuvier, and *putida* Boitard are inadmissible, the first because it is indeterminable and the latter because it was previously used by Cuvier, *olida* Boitard (l. c.), becomes the name that should properly be employed. (On this subject, see Allen, Proc. Biol. Soc. Wash., xv, 1902, p. 66.)

FLORIDA SKUNK.

Type locality. Micco, Brevard County, Florida.

Geogr. Distr. Florida, vicinity of Lake Worth, to North Carolina; mountains of West Virginia; west on Gulf coast to the Mississippi River.

987. *Mephitis mesomelas* Lichtenstein.

Mephitis mesomelas Licht., Darst. Säugeth., 1832, Text, pl. 45, fig. 2.

Mephitis mephitica scrutator Bangs, Proc. Biol. Soc. Wash., x, 1896, p. 141. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 324. Zoöl. Ser.

LOUISIANA SKUNK.

Type locality. Louisiana.

Geogr. Distr. West side of Mississippi Valley from southern Louisiana to Missouri; westward along coast of Texas to Matagorda Island; and up the Red River Valley as far at least as Wichita Falls.

a. *Mephitis mesomelas avia* (Bangs).

Mephitis avia Bangs, Proc. Biol. Soc. Wash., XII, 1898, p. 32.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 327. Zoöl. Ser.

ILLINOIS SKUNK.

Type locality San José, Mason County, Illinois.

Geogr. Distr. Prairie region of Illinois, western Indiana, and eastern Iowa. Range not entirely determined.

b. *Mephitis mesomelas varians* (Gray).

Mephitis varians Gray, Charlesw. Mag. Nat. Hist., I, 1837, p. 581.

Mephitis mesomelas Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 325. Zoöl. Ser. (nec Lichten.)

LONG-TAILED TEXAN SKUNK.

Type locality. Texas. Lower Grande Valley specimens typical.

Geogr. Distr. Southern and western Texas, eastern New Mexico, and adjacent parts of Mexico; north into Oklahoma, Colorado, Kansas, and Nebraska.

988. *Mephitis estor* Merriam.

Mephitis estor Merr., N. Am. Faun., No. 3, 1890, p. 81. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 326. Zoöl. Ser.

Mephitis mesomelas estor Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 509. Zoöl. Ser.

ARIZONA SKUNK.

Type locality. San Francisco Mountain, Coconino County, Arizona.

Geogr. Distr. Arizona, western New Mexico, States of Sonora, and Chihuahua south in the Sierra Madre in latter state; Lower California, Mexico.

989. *Mephitis occidentalis* Baird.

Mephitis occidentalis Baird, N. Am. Mamm., 1857, p. 194. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1904, p. 324. Zool. Ser.

CALIFORNIA SKUNK.

Type locality. Petaluma, Sonoma County, California.

Geogr. Distr. Northern and central California; vicinity of Monterey Bay northward, west of the Sierra and Cascades to the Willamette Valley, Oregon.

a. Mephitis occidentalis spissigrada (Bangs).

Mephitis spissigrada Bangs, Proc. Biol. Soc. Wash., 11, 1898, p. 31. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 326. Zool. Ser.

Mephitis fatulenta Elliot, Pub. Field Columb. Mus., 1, 1899, p. 269. Zool. Ser. *Id.* Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 326. Zool. Ser.

PUGET SOUND SKUNK.

Type locality. Sumas, British Columbia, Canada.

Geogr. Distr. Shores of Puget Sound and coast region of Washington and northern Oregon.

b. Mephitis occidentalis notata (Howell).

Chincha occidentalis notata Howell, N. Am. Faun., No. 20, 1901, p. 36.

CASCADE SKUNK.

Type locality. Trout Lake, Mount Adams, Skamania County, Washington.

Geogr. Distr. Southern Washington and northern Oregon, east of the Cascades.

Genl. Char. Similar to *M. occidentalis*, but tail shorter; body stripes narrow, and separated for their entire length. Skull: nasals short, lower carnassial broad; palate without notch.

Color. Black; white stripes narrow, not confluent, but usually joined to the nape; tail black exteriorly.

Measurements. Total length, average of 3 ad., 633; tail vertebræ, 249; hind foot, 76.

c. Mephitis occidentalis major (Howell).

Chincha occidentalis major Howell, N. Am. Faun., No. 20, 1901, p. 37.

GREAT BASIN SKUNK.

Type locality. Fort Klamath, Klamath County, Oregon. Altitude, 4,200 feet.

Geogr. Distr. Eastern Oregon, northern California, and Nevada east to the Wahsatch Mountains.

Genl. Char. White stripes broad and extending only short way on the tail; tail short, black exteriorly. Skull large; palate sometimes without notch; rostrum broad, flattened; premaxillæ short, broad.

Color. Like *M. occidentalis*; white stripes broad, bifurcating near middle of the back, and extending only a short distance on the tail.

Measurements. Average of 5 males. Total length, 705; tail vertebræ, 306; hind foot, 84.

d. Mephitis occidentalis holzneri Mearns.

Mephitis occidentalis holzneri Mearns, Proc. U. S. Nat. Mus., xx, 1897, p. 461. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 507, fig. 94. Zool. Ser.

LOWER CALIFORNIA SKUNK.

Type locality. San Isidro Ranch, Lower California, Mexico; near border of San Diego County, California.

Geogr. Distr. Monterey Bay south into Lower California, Mexico; east to the Sierra Nevada and San Bernardino Range, and the Mohave Desert.

990. *Mephitis platyrhina* (Howell).

Chincha platyrhina Howell, N. Am. Faun., No. 20, 1901, p. 39.

BROAD-NOSED SKUNK.

Type locality. South Fork of Kern River, twenty-five miles east of Kernville, Kern County, California.

Geogr. Distr. Known only from the type locality and from Owens Valley, Inyo County, California.

Genl. Char. Like *M. occidentalis*; rostrum very broad; zygomata spreading evenly, nearly parallel to skull; nasals short, broad.

Color. Black; white stripe of median width passing but short distance on the tail, which is black, an indistinct white band on the upper surface of apical half.

Measurements. Total length, 750; tail vertebræ, 320; hind foot, 90.

Subgenus B. *Leucomitra*. Hooded Skunks.991. *Mephitis macrura* Lichtenstein.*Mephitis macroura* Licht., Darst. Säugeth., 1832, pl. XLVI.*Mephitis macrura* Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 510, figs. 95 and LXXI. Zoöl. Ser.

HOODED SKUNK.

Type locality. Mountains northwest of the City of Mexico, Mexico.*Geogr. Distr.* Highlands of central and southern Mexico, south to Guatemala.a. *Mephitis macrura milleri* (Mearns).*Mephitis milleri* Mearns, Proc. U. S. Nat. Mus., xx, 1897, p. 467.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 325. Zoöl. Ser.

Mephitis macrura milleri Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 511. Zoöl. Ser.

MILLER'S SKUNK.

Type locality. Fort Lowell, near Tucson, Pima County, Arizona.*Geogr. Distr.* Known only from the type locality.b. *Mephitis macrura vittata* (Lichtenstein).*Mephitis vittata* Licht., Darst. Säugeth., 1832, pl. 47.*Mephitis concolor* Gray, Proc. Zoöl. Soc., 1865, p. 149.*Mephitis vittata* var. *b. intermedia* Gray, Cat. Carniv. Brit. Mus., 1869, p. 138.*Mephitis macrura vittata* Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 512. Zoöl. Ser.

BRIDLED SKUNK.

Type locality. San Mateo del Mar, State of Oaxaca, Mexico.*Geogr. Distr.* Known only from the type locality.Genus 125. *Conepatus*.*Conepatus* Gray, Charlesw. Mag. Nat. Hist., 1837, p. 581. Type*Conepatus humboldtii* Gray.*Ictonyx (sic)* Kaup, Thierr., I, 1835, p. 352.*Marputius* Gray, Charlesw. Mag. Nat. Hist., 1837, p. 581.*Thiosmus* Licht., Abh. K. Akad. Wiss. Berl., 1838, p. 270.*Ozolicis* Gloger, Hand- u. Hilfsb. Naturg., 1841, p. 57.*Rhinozolis* Gloger, Hand- u. Hilfsb. Naturg., 1841, p. 58.*Oryctogale* Merr., Proc. Biol. Soc. Wash., xv, 1902, p. 161.

Subgenus A. *Conepatus*.

992. *Conepatus sonoriensis* Merriam.

Conepatus sonoriensis Merr., Proc. Biol. Soc. Wash., xv, 1902, p. 162. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. II, 1904, p. 514. Zoöl. Ser.

SONORAN SKUNK.

Type locality. Camoa, Rio Mayo, State of Sonora, Mexico.

Geogr. Distr. Valparaiso Mountains, State of Zacatecas, north-westerly into the State of Sonora.

993. *Conepatus leuconotus* (Lichtenstein).

Mephitis leuconota Licht., Darst. neuer oder wenig bekannter Säugeth., 1832, text, pl. XLIV, fig. 1.

Conepatus leuconotus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. II, 1904, p. 515. Zoöl. Ser.

WHITE-SPOTTED SKUNK.

Type locality. Upper waters of Rio Alvarado, State of Vera Cruz, Mexico.

Geogr. Distr. States of Vera Cruz, Mexico. Range unknown.

a. Conepatus leuconotus texensis Merriam.

Conepatus leuconotus texensis Merr., Proc. Biol. Soc. Wash., xv, 1902, p. 162. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. II, 1904, p. 515. Zoöl. Ser.

TEXAN SKUNK.

Type locality. Brownsville, Lower Rio Grande, Cameron County, Texas.

Geogr. Distr. Coast of Texas from Rockport, Aransas County, to mouth of the Rio Grande. State of Tamaulipas, Mexico?

994. *Conepatus mesoleucus* (Lichtenstein).

Mephitis mesoleuca Licht., Darst. neuer oder wenig bekannter Säugeth., 1832, pl. XLIV, fig. 2.

Conepatus mesoleucus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. II, 1904, p. 515. Zoöl. Ser.

WHITE-BACKED SKUNK.

Type locality. Chico, State of Vera Cruz, Mexico.

Geogr. Distr. State of Vera Cruz, Mexico. Range unknown.

a. Conepatus mesoleucus mearnsi Merriam.

Conepatus mesoleucus mearnsi Merr., Proc. Biol. Soc. Wash., xv, 1902, p. 163. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. II, 1904, p. 516, figs. 96 and LXXII. Zoöl. Ser.

Conepatus mapurito Elliot, Syn. N. Am. Mamm., F. C. M. Pub.,
II, 1901, Suppl., p. 510, pl. LVI. Zoöl. Ser. (nec Gmel.)

MEARNS' SKUNK.

Type locality. Mason, Mason County, Texas.

Geogr. Distr. Tableland of Mexico from the States of Jalisco and southwestern San Luis Potosi, northward to central Texas and central Arizona.

995. *Conepatus felipensis* Merriam.

Conepatus felipensis Merr., Proc. Biol. Soc. Wash., xv, 1902, p.
163. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M.
Pub., IV, Pt. II, 1904, p. 517. Zoöl. Ser.

SAN FELIPE SKUNK.

Type locality. Cerro San Felipe, State of Oaxaca, Mexico.

Geogr. Distr. Cerro San Felipe, State of Oaxaca, Mexico. Altitude, 7,000 to 10,500 feet.

996. *Conepatus pediculus* Merriam.

Conepatus pediculus Merr., Proc. Biol. Soc. Wash., xv, 1902, p.
164. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M.
Pub., IV, Pt. II, 1904, p. 517. Zoöl. Ser.

GUADALUPE SKUNK.

Type locality. Sierra Guadalupe, State of Coahuila, Mexico.

Geogr. Distr. State of Coahuila, Mexico. Range unknown.

Subgenus B. *Marputius*.

997. *Conepatus tropicalis* Merriam.

Conepatus tropicalis Merr., Proc. Biol. Soc. Wash., xv, 1902, p.
164. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M.
Pub., IV, Pt. II, 1904, p. 517, fig. 97. Zoöl. Ser.

MOTZORONGO SKUNK.

Type locality. Motzorongo, State of Vera Cruz, Mexico.

Geogr. Distr. East coast of the State of Vera Cruz, southeasterly to Yucatan, Mexico.

Genus 126. **Spilogale. Little Striped or Spotted Skunks.**

Spilogale Gray, Proc. Zoöl. Soc., 1865, p. 150. Type *Mephitis interrupta* Rafinesque.

998. *Spilogale interrupta* (Rafinesque).

Mephitis interrupta Rafin., Ann. Nat., I, 1820, p. 3.

Mephitis bicolor Gray, Charles. Mag. Nat. Hist., I, 1837, p. 581.

Mephitis quaterlinearis Winans, Coues Fur-bear. Anim., 1877, p. 239.

Spilogale interrupta Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 328. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 523, figs. 98, LXXIII. Zoöl. Ser.

WHITE-SPOTTED SKUNK.

Type locality. Upper Missouri?

Geogr. Distr. Kansas, Oklahoma Territory to Texas, south through Mexico to Guatemala.

999. *Spilogale ringens* Merriam.

Spilogale ringens Merr., N. Am. Faun., No. 4, 1890, p. 9. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 329. Zoöl. Ser.

GROWLING SPOTTED SKUNK.

Type locality. Greensborough, Hale County, Alabama.

Geogr. Distr. Alabama, western Georgia, north to West Virginia and Virginia.

1000. *Spilogale indianola* Merriam.

Spilogale indianola Merr., N. Am. Faun., No. 4, 1890, p. 10. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 329. Zoöl. Ser.

INDIANOLA SPOTTED SKUNK.

Type locality. Indianola, Matagorda Bay, Matagorda County, Texas.

Geogr. Distr. Texas.

1001. *Spilogale leucoparia* Merriam.

Spilogale leucoparia Merr., N. Am. Faun., No. 4, 1890, p. 11. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 329. Zoöl. Ser.

MERRIAM'S SPOTTED SKUNK.

Type locality. Mason, Mason County, Texas.

Geogr. Distr. Texas.

1002. *Spilogale tenuis* Howell.

Spilogale tenuis Howell, Proc. Biol. Soc. Wash., xv, 1902, p. 241.

SLENDER SPOTTED SKUNK.

Type locality. Arkins, Larimer County, Colorado.

Geogr. Distr. Colorado. Range unknown.

Genl. Char. Similar to *S. leucoparia*; lateral stripes narrower; frontal patch long and narrow; terminal third of tail white. Skull:

braincase broad and very flat; rostrum, postorbital region, and anterior part of braincase very narrow.

Measurements. Total length, 450; tail vertebræ, 165; hind foot, 51. Skull: Hensel, 52; zygomatic width, 34.7; mastoid breadth, 32.5; interorbital breadth, 14.3.

1003. *Spilogale saxatilis* Merriam.

Spilogale saxatilis Merr., N. Am. Faun., No. 4, 1890, p. 13. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 330. Zool. Ser.

ROCK SPOTTED SKUNK.

Type locality. Provo, Utah County, Utah.

Geogr. Distr. Utah.

1004. *Spilogale phenax* Merriam.

Spilogale phenax Merr., N. Am. Faun., No. 4, 1890, p. 13. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 331. Zool. Ser.

CHEATING SPOTTED SKUNK.

Type locality. Nicasio, Marin County, California.

Geogr. Distr. California.

a. Spilogale phenax latifrons Merriam.

Spilogale phenax latifrons Merr., N. Am. Faun., No. 4, 1890, p. 15. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 331. Zool. Ser.

BROAD-HEADED SPOTTED SKUNK.

Type locality. Roseburg, Douglas County, Oregon.

Geogr. Distr. Oregon.

1005. *Spilogale arizonæ* (Mearns).

Spilogale phenax arizonæ Mearns, Bull. Am. Mus. Nat. Hist. N. Y., 1891, p. 256. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 331. Zool. Ser.

ARIZONA SPOTTED SKUNK.

Type locality. Fort Verde, Arizona.

Geogr. Distr. Central Arizona.

a. Spilogale arizonæ martirensis Elliot.

Spilogale arizonæ martirensis Elliot, Pub. Field Columb. Mus., 111, 1903, p. 170. Zool. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 11, 1904, p. 521. Zool. Ser.

SAN PEDRO MARTIR SPOTTED SKUNK.

Type locality. Vallecitos, San Pedro Martir Mountains, Lower California, Mexico. Altitude, 9,000 feet.

Geogr. Distr. San Pedro Martir Range, Lower California, Mexico.

1006. *Spilogale olympica* Elliot.

Spilogale olympica Elliot, Pub. Field Columb. Mus., 1, 1899, p. 270. Zoöl. Ser. *Id.* Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 332. Zoöl. Ser.

OLYMPIC MOUNTAINS SPOTTED SKUNK.

Type locality. Lake Southerland, Olympic Mountains, Clallam County, Washington.

Geogr. Distr. Northwest Washington.

1007. *Spilogale ambarvalis* Bangs.

Spilogale ambarvalis Bangs, Proc. Bost. Soc. Nat. Hist., 1898, p. 222. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 332. Zoöl. Ser.

FIELD SPOTTED SKUNK.

Type locality. Oak Lodge, opposite Micco, Brevard County, Florida.

Geogr. Distr. East Peninsula, Indian River, Florida.

1008. *Spilogale gracilis* Merriam.

Spilogale gracilis Merr., N. Am. Faun., No. 4, 1890, p. 83. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 330. Zoöl. Ser.

GRACEFUL SKUNK.

Type locality. Grand Cañon of the Colorado, north of San Francisco Mountain, Coconino County, Arizona.

Geogr. Distr. Arizona. Range undetermined.

1009. *Spilogale pygmæa* Thomas.

Spilogale pygmæa Thomas, Proc. Zoöl. Soc., 1898, p. 898. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., 1v, Pt. 11, 1904, p. 519. Zoöl. Ser.

PIGMY SPOTTED SKUNK.

Type locality. Rosario, State of Sinaloa, Mexico.

Geogr. Distr. State of Sinaloa, Mexico.

1010. *Spilogale ambigua* Mearns.

Spilogale ambigua Mearns, Proc. U. S. Nat. Mus., xx, 1897, p. 460. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., 1v, Pt. 11, 1904, p. 521. Zoöl. Ser.

CHIHUAHUA LITTLE SPOTTED SKUNK.

Type locality. Eagle Mountain, State of Chihuahua, Mexico.

Geogr. Distr. State of Chihuahua, Mexico; Mimbres Valley and Lake Palomas, boundary line.

1011. *Spilogale angustifrons* Howell.

Spilogale angustifrons Howell, Proc. Biol. Soc. Wash., xv, 1902, p. 242. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. II, 1904, p. 521. Zoöl. Ser.

NARROW-HEADED SPOTTED SKUNK.

Type locality. Tlalpam, Federal District, Mexico.

Geogr. Distr. Tableland in the State of Guanajuato to State of Chiapas, Mexico.

a. *Spilogale angustifrons tropicalis* Howell.

Spilogale angustifrons tropicalis Howell, Proc. Biol. Soc. Wash., xv, 1902, p. 242. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. II, 1904, p. 521. Zoöl. Ser.

OAXACA SPOTTED SKUNK.

Type locality. San Mateo del Mar, State of Oaxaca, Mexico.

Geogr. Distr. State of Oaxaca, Mexico. Range unknown.

1012. *Spilogale lucasana* Merriam.

Spilogale lucasana Merr., N. Am. Faun., No. 4, 1890, p. 11. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. II, 1904, p. 523. Zoöl. Ser.

CAPE ST. LUCAS SPOTTED SKUNK.

Type locality. Cape St. Lucas, Lower California, Mexico.

Geogr. Distr. Cape Region of Lower California, Mexico.

Subfamily II. *Mustelinæ*.**Genus 127. *Gulo*. *Wolverine*.**

Gulo Frisch, Das Natur-syst. vierfüss. Thiere, in Tabellen, 17, Tab. Gen., 1775. Type *Mustela gulo* Linnæus.

1013. *Gulo luscus* (Linnæus).

Mustela gulo Linn., Syst. Nat., 1, 1758, p. 45.

Ursus luscus Linn., Syst. Nat., 1, 1758, p. 47.

Gulo luscus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 333. Zoöl. Ser.

WOLVERINE.

Type locality. Hudson Bay.

Geogr. Distr. Boreal North America from Atlantic Ocean to Alaska, south to the Basin of the Great Salt Lake, Utah.

1014. *Gulo luteus* Elliot.

Gulo luteus Elliot, Pub. Field Columb. Mus., III, 1903, p. 200.
Zool. Ser.

BUFF-COLORED WOLVERINE.

Type locality. Mount Whitney, Inyo County, California.

Geogr. Distr. Mount Whitney to Yakutat Bay? Alaska.

Genl. Char. General hue of hind part of head, sides of body, and base of tail buff color.

Color. Nose, lips, cheeks to and including eyes jet black; top of head and back of eyes pale gray; nape and space between shoulders chestnut; lower part of back and rump seal brown in the center, grading to chestnut on the edges; band across middle of back and encircling the dark patch and sides buff; under parts blackish chestnut with a small white spot on throat; legs and feet black; tail, basal half buff, remainder black; ears chestnut with broad buff edging.

Measurements. Total length, 850; tail vertebrae, 205; hind foot, 165; ear, 53.

1015. *Gulo hylaëus* Elliot.

Gulo hylaëus Elliot, Proc. Biol. Soc. Wash., XVIII, 1905, p. 81.

MOUNT MCKINLEY WOLVERINE.

Type locality. Upper waters of Sushitna River, region of Mount McKinley, Alaska. No. 9883, Field Columbian Mus., Chicago.

Genl. Char. Type: General color very dark; no buff hue on body; or gray on head anywhere visible. Skull compared with that of specimens from New Brunswick, Canada, is narrower, especially at intertemporal region, also at postorbital constriction and posterior region of braincase; and the zygomatic width is less; but the audital bullae are nearly twice as large, and swollen greatly on their inner margin, whereas those of the Canadian animal are much flattened and small in size.

Color. Head, throat, sides of neck, and body, and base of tail chestnut; hind part of neck, back, under parts, legs, and feet black; chest spotted or blotched with white or orange, and orange spot on anal region; nose darker chestnut than head; tail, except at base, black.

Measurements. Skull: Total length, 157; Hensel, 133; zygomatic width, 104; intertemporal width, 47; postorbital constriction, 32; palatal length, 75; greatest width of braincase, 67.5; mastoid width, 89; post-palatal length, 57; length of nasals, 21; length of upper pre-molar and molar series, 41; length of mandible, 101; height of coronoid process above angle, 35; length of lower premolar and molar series 54; length of lower carnassial, alveolar border, 21.

Genus 128. **Galera. The Grisons.**

Galera Browne, Civil & Naval Hist. Jamaica, 2d Ser., 1789, p. 485, Tab. 49, fig. 1. Type *Mustela barbara* Linnæus.

**Grison* Oken, Lehrb. der Naturgesch., Zool., II, 1816, pp. XI, 1000.

Tayra Oken, Lehrb. der Naturgesch., Zool., II, 1816, p. 1001.

Grisonia Gray, Ann. Phil., XXVI, 1825, p. 339.

Laira F. Cuv., Hist. Nat. Mamm., v, 1826, livr. XLV.

Galictis Bell, Zoöl. Journ., II, 1826, p. 551.

Galidictis I. Geoff., Mag. Zool., Mamm., 1839, p. 32.

Subgenus A. *Tayra*.

Galera barbara biologiae (Thomas).

Galictis barbara biologiae Thomas, Ann. Mag. Nat. Hist., 7th Ser., v, 1900, p. 146.

Grison barbara biologiae Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 524, fig. 99 (*G. barbara*). Zoöl. Ser.

CALOVEVORA GRISON. *Chulomuco, Tulomuco* in Costa Rica.

Type locality. Calovevora, Veragua, Panama.

Geogr. Distr. Central America generally.

Galera barbara senex (Thomas).

Galictis barbara senex Thomas, Ann. Mag. Nat. Hist., 7th Ser., v, 1900, p. 146.

Grison barbara senex Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 525. Zoöl. Ser.

GRAY-HEADED GRISON.

Type locality. Hacienda Tortugas, Jalapa, State of Vera Cruz, Mexico. Altitude, about 600 feet.

Geogr. Distr. Known only from the type locality.

Subgenus B. *Galictis*.

1016. **Galera allamandi** (Bell).

Galictis allamandi Bell, Proc. Zoöl. Soc., 1837, p. 47.

Grison allamandi Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 526. Zoöl. Ser.

*In the Mamm. Middle Amer. & W. Indies, *Grison* was employed as the generic term for the allied forms of *Mustela barbara* Linn., but this appears to be an error, as *Galera* (l. c.) Brown, clearly antedates *Grison* Oken, and should be the term employed. The fact that *Galera* has been used in various ways by authors, and some confusion has been thereby created, in no way invalidates its claim to take precedence over all other names proposed for this genus of mammals.

ALLAMAND'S GRISON.

Type locality. Unknown. Collection of the Zoölogical Society of London.

Geogr. Distr. Unknown.

1017. *Galera canaster* (Nelson).

Galictis canaster Nelson, Proc. Biol. Soc. Wash., XIV, 1901, p. 129.

Grison canaster Elliot, Mamm. Middle Amer. & W. Indies, F. C. M.

Pub., IV, Pt. II, 1904, p. 526, figs. 100, LXXIV. Zoöl. Ser.

GRIZZLED, OR NELSON'S GRISON.

Type locality. Tunkas, State of Yucatan, Mexico.

Geogr. Distr. From Orizaba, State of Puebla, to Yucatan, Mexico.

Genus 129. *Mustela*. Martens, or Sables. Fishers.

Mustela Linn., Syst. Nat., I, 1758, p. 45; I, 1766, p. 66. *Type Mustela martes* Linnæus.

1018. *Mustela americana* Turton.

Mustela martes Forst., Philos. Trans., 1772, p. 372. (nec Linn.)

Mustela zibellina Pallas, Spicil. Zool., XIV, 1780?, p. 57. *Id.* Zoög.

Rosso-Asiat., I, 1831, p. 84. (nec Linn.)

Mustela americanus Turton, Linn. Syst. Nat., I, 1806, p. 60.

Mustela vulpina Rafin., Am. Journ. Scien., I, 1819, p. 82.

Mustela leucopus Kuhl, Beitr., 1820, p. 74.

Mustela huro F. Cuv., Dict. Scien. Nat., XXIX, 1823, p. 256.

Mustela americana Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 334, fig. 66. Zoöl. Ser.

AMERICAN MARTEN. HUDSON BAY SABLE.

Type locality. Eastern North America.

Geogr. Distr. Boreal North America, west to the Rocky Mountains, south to New York.

a. *Mustela americana abieticola* Preble.

Mustela americana abieticola Preble, N. Am. Faun., No. 22, 1902, p. 68.

SASKATCHEWAN MARTEN.

Type locality. Cumberland House, Saskatchewan, Canada.

Genl. Char. Larger than *M. americana*; dentition heavier, except last molar.

Color. General color dark yellowish brown, darkest on dorsal region; legs dark brown; tail dark brown, tip usually black; spot on chest ochraceous; face and cheeks grayish brown, ears edged with whitish.

Measurements. Total length, 640; tail vertebræ, 210; hind foot, 95. Skull: average of six adults; occipito-nasal length, 78.9; zygomatic breadth, 47.8; interorbital constriction, 16.7; width across postorbital processes, 22.5; mastoid breadth, 37.8.

b. Mustela americana abietinoides (Gray).

Martes americana var. 1. *abietinoides* Gray, Proc. Zool. Soc., 1865, p. 106.

SELKIRK MARTEN.

Type locality. "Rocky Mountains."

Geogr. Distr. Interior region of British Columbia, Selkirk, and Gold Mountain Ranges, Canada.

Color. "Black brown; ears pale; head grizzled with white hairs, more or less gray; throat yellow or yellow spotted; throat spots large or broken up into small spots; the head sometimes with only a few gray hairs, and the throat with only a few distinct small spots." (Gray.)

Measurements. Total length, 584; tail vertebræ, 178; hind foot, 87. Skull: basal length, 70; zygomatic breadth, 43; postorbital constriction, 17.

c. Mustela americana brumalis (Bangs).

Mustela brumalis Bangs, Amer. Nat., xxxii, 1898, p. 502, fig.

Mustela americana brumalis Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 336. Zool. Ser.

LABRADOR MARTEN.

Type locality. Okak, Labrador.

Geogr. Distr. Northern Labrador.

d. Mustela americana actiosa Osgood.

Mustela americana actiosa Osgood, N. Am. Faun., No. 19, 1900, p. 43. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 336. Zool. Ser.

ACTIVE MARTEN.

Type locality. Fort Yukon, Alaska.

Geogr. Distr. Alaska. Range unknown.

1019. *Mustela nesophila* Osgood.

Mustela nesophila Osgood, N. Am. Faun., No. 21, 1901, p. 33.

QUEEN CHARLOTTE ISLAND MARTEN.

Type locality. Graham Island, Queen Charlotte Islands, British Columbia, Canada.

Genl. Char. Rostrum thick, heavy.

Measurements. Skull: basilar length, 73; palatal length, 39; post-

palatal length, 34; zygomatic breadth, 45.5; postorbital width across processes, 23; interorbital constriction, 19; skull only known.

1020. *Mustela boria* Elliot.

Mustela boria Elliot, Proc. Biol. Soc. Wash., XVIII, 1905, p. 139.

FARTHEST NORTH MARTEN.

Type locality. Lower Mackenzie River District, towards Arctic Ocean.

Genl. Char. Darkest in color of American Martens. Tail long, black.

Color. Sides of head grayish white; nose and top of head light brown and gray mixed; entire body above and below burnt umber, blackish on dorsal region; entire throat beneath blotched with orange; fore and hind legs black; tail black, base of hairs on proximal half burnt umber; ears like body externally, white internally.

Measurements. Total length, nose to end of hairs of tail, 785; tail, end of hairs, 267; hind foot, 85. (Skin.) Skull (broken): zygomatic width, 50; intertemporal width, 16; length of nasals, 15; width of rostrum at canines, 18; palatal length, 41; length of upper tooth row, outer edge of canine to posterior edge of last molar, 30; length of mandible, 51; length of lower tooth row, 36.

1021. *Mustela kenaiensis* (Elliot).

Mustela americana kenaiensis Elliot, Pub. Field. Columb. Mus., III, 1903, p. 151, pl. XXXII. Zool. Ser.

KENAI PENINSULA MARTEN.

Type locality. Kenai Peninsula, Alaska.

Geogr. Distr. Range unknown.

Genl. Char. Smaller than *M. a. actiosa*; tail longer; feet shorter; color darker; no gular or abdominal patches; audital bullæ large, ridged, rectangular, longitudinally produced; molar heavy; sagittal crest prominent.

Color. Above tawny buff and black; top of head dark gray; forehead, top of nose, and sides of head brownish white; sides of nose dark brown; chin and throat uniform dark ochre; under parts dark chocolate brown; inguinal region pale buff; outer fore legs above chocolate brown; feet black with grayish patch at toes beneath; hind legs and feet blackish; chocolate patch at base of toes above; tail black, chocolate at base; ears brownish black.

Measurements. Total length, 790; tail vertebræ, 293; hind foot, 90. (Skin.) Skull: occipito-nasal length, 79.5; Hensel, 75; zygomatic width, 50; interorbital breadth, 18; mastoid breadth, 35.5; length of nasals, 8; palatal length, 39.5; post-palatal length, 34.5; length of pterygoid fossa, 12; length of bullæ, 17; length of upper

tooth row, 25; length of mandible, 55; height at condyle, 26; length of lower tooth row, 30.

1022. *Mustela atrata* Bangs.

Mustela atrata Bangs, Amer. Nat., xxxi, 1897, p. 162. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 335. Zoöl. Ser.

NEWFOUNDLAND MARTEN.

Type locality. Bay St. George, Newfoundland.

Geogr. Distr. Newfoundland.

1023. *Mustela caurina* Merriam.

Mustela caurina Merr., N. Am. Faun., No. 4, 1890, p. 27.

Mustela atrata caurina Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 335. Zoöl. Ser.

PACIFIC MARTEN.

Type locality. Near Gray's Harbor, Chehalis County, Washington.

Geogr. Distr. Northwest coast of California to Puget Sound.

a. Mustela caurina origenes Rhoads.

Mustela caurina origenes Rhoads, Proc. Acad. Nat. Scien. Phil., 1902, p. 458.

ROCKY MOUNTAIN MARTEN.

Type locality. Marvine Mountain, Garfield County, Colorado.

Geogr. Distr. Southern Rocky Mountain region; new Mexico?

Color. Upper parts and sides wood brown; long hairs on dorsal region and rump broccoli brown; gular patch extensive; buffy orange; rest of under parts lighter than back; tail long, dark; ears edged with ochraceous buff.

Measurements. Total length, 676; tail vertebræ, 230; hind foot, 85. Skull: basal length, 65; zygomatic breadth, 41.

1024. *Mustela canadensis* Schreber.

Mustela canadensis Schreber, Säugth., III, Text, p. 492, 1777, pl. CXXIV, 1776.

Mustela pennanti Erxl., Syst. Anim., 1777, p. 470. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 337. Zoöl. Ser.

Mustela melanorhyncha Bodd., Elench. Anim., 1784, p. 88.

Viverra piscator Shaw, Gen. Zoöl., I, 1800, p. 414.

Mustela nigra Turton, Linn. Syst. Nat., I, 1806, p. 60.

Mustela godmani Fischer, Syn. Mamm., 1829, p. 217.

PENNANT'S FISHER.

Type locality. "New York and Pennsylvania." (Pennant.)

Geogr. Distr. North America north of 35°, in forest-covered country, east of the Cascade Mountains. Intergrading with the next

species on the Pacific slope. Typical form restricted to Alleghanian specimens.

a. *Mustela canadensis pacifica* Rhoads.

Mustela canadensis pacifica Rhoads, Trans. Am. Philos. Soc., 1898, p. 435.

Mustela pennanti pacifica Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 337. Zoöl. Ser.

PACIFIC FISHER.

Type locality. Lake Kichelos, Kittitas County, Washington. Altitude, 8,000 feet.

Geogr. Distr. Pacific slope, California to Alaska.

Genus 130. **Putorius. Mink, Weasels.**

Putorius Frisch., Nat. Syst. vierfüß. Thiere, in Tab., 11, Tab. Gen., 1775. Type *Mustela putorius* Linnæus.

Arctogale Kaup (nec Peters), Entw.-Gesch. Nat. Syst. der Europ. Thierw., 1, 1829, p. 30.

Ictis Kaup, Entw.-Gesch. Nat. Syst. der Europ. Thierw., 1, 1829, p. 40.

Gale Wagn., Schreb. Säugth., Suppl., 11, 1841, p. 234.

Lutreola Wagn., Schreb. Säugth., Suppl., 11, 1841, p. 239.

Vison Gray, List Spec. Mamm. Brit. Mus., 1843, p. 64.

Neogale Gray, Proc. Zoöl. Soc., 1865, p. 114.

Cynomyonax (sic) Coues, Fur-bear. Anim., 1877, p. 147.

Subgenus A. *Lutreola*. Mink.

1025. **Putorius vison** (Schreber).

Mustela lutreola Forst., Phil. Trans., LXII, 1772, p. 371. (nec Linn.)

Mustela canadensis Erxl., Syst. Reg. Anim., 1, 1777, p. 455. (nec *M. l. canadensis* of Schreber.)

Mustela vison Schreber, Säugth., III, 1777, p. 463.

Mustela winingus Barton, Am. Philos. Trans., VI, 1809, p. 70.

Mustela minx Ord, Guthrie's Geogr., 2d Am. ed., 11, 1815, p. 291.

Mustela lutreocephala Harlan, Faun. Amer., 1825, p. 63.

Putorius nigrescens Aud. & Bach., Quad. N. Am., III, 1853, p. 104, pl. cxxiv.

Putorius vison Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 338, fig. 67. Zoöl. Ser.

Putorius vison lutreocephalus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 339. Zoöl. Ser.

COMMON MINK.

Type locality. Eastern Canada.

Geogr. Distr. Eastern North America through Canada north to the Arctic Sea east of Hudson Bay. In United States north of Gulf States to eastern boundary of the Pacific coast.

a. *Putorius vison lacustris* (Preble).

Lutreola vison lacustris Preble, N. Am. Faun., No. 22, 1902, p. 66.
KEEWATIN MINK.

Type locality. Echimamish River, Keewatin, Canada.

Genl. Char. Similar to *P. vison*, but larger; skull more angular; sagittal crest highly developed; zygomatics greatly bowed outward; bullæ large, flat, inner anterior corner produced toward pterygoids, terminating in acute angle.

Color. Above dark chocolate brown; beneath lighter; chin and narrow stripe to breast white; also some white between hind legs.

Measurements. Total length, 560; tail vertebrae, 190; hind foot, 67. Skull: average 10 adults. Occipito-nasal length, 62.5; zygomatic breadth, 41.25; interorbital constriction, 11.9; width across postorbital processes, 18.25; mastoid breadth, 36.

b. *Putorius vison vulgivagus* (Bangs).

Putorius (Lutreola) vulgivagus Bangs, Proc. Bost. Soc. Nat. Hist., 1895, p. 536.

Putorius vison vulgivagus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 340. Zoöl. Ser.

RAMBLING MINK.

Type locality. Burbridge, Plaquemines Parish, Louisiana.

Geogr. Distr. Shores of the Gulf of Mexico from Texas to Florida and probably on the Atlantic coast of Georgia and South Carolina.

c. *Putorius vison energumenus* Bangs.

Putorius vison energumenus Bangs, Proc. Bost. Soc. Nat. Hist., 1896, p. 5.

Putorius vison energumenus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 340. Zoöl. Ser.

DEVILISH MINK.

Type locality. Sumas, British Columbia, Canada.

Geogr. Distr. Pacific slope from northern California to Alaska.

d. *Putorius vison ingens* (Osgood).

Lutreola vison ingens Osgood, N. Am. Faun., No. 19, 1900, p. 42.

Putorius vison ingens Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 340. Zoöl. Ser.

LARGE MINK.

Type locality. Fort Yukon, Alaska.

Geogr. Distr. Region of the River Yukon, Alaska.

1026. *Putorius melampeplus* (Elliot).

Putorius vison melampeplus Elliot, Pub. Field Columb. Mus., III, 1903, p. 170. Zoöl. Ser.

DARK-HUED MINK.

Type locality. Kenai Peninsula, Alaska.

Genl. Char. Size large, but smaller than *P. v. ingens*; color dark without reddish tint. Skull: compared with that of *P. v. energumenus* has flatter audital bullæ; palate broader posteriorly; broader pterygoid fossa and comparatively shorter; rostrum wider and heavier; interorbital constriction greater as is also the width across post-orbital processes; palate wider at incisors, and last upper molar smaller and narrower. Well developed sagittal and occipital crests.

Color. Entire upper parts, legs, and feet dark chocolate without any red tints; under parts slightly paler; white spot on chin, and sometimes a smaller one on chest; tail blackish.

Measurements. Total length, 710; tail vertebræ, 180; hind foot, 65. (Skin.) Skull: occipito-nasal length, 59; Hensel, 60; zygomatic width, 40; interorbital constriction, 10; mastoid width, 32; width of braincase, 30; palatal length, 28; length of audital bullæ, 15; breadth of rostrum, 15; length of nasals, 9; width across post-orbital processes, 18; length of mandible, 35; height at angle, 6; at coronoid process, 19.

1027. *Putorius lutensis* Bangs.

Putorius (Lutreola) lutensis Bangs, Proc. Bost. Soc. Nat. Hist., 1898, p. 229.

Putorius lutensis Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 340. Zoöl. Ser.

MUD-LOVING MINK.

Type locality. Salt marsh off Matanzas Inlet, St. John County, Florida.

Geogr. Distr. South Atlantic States, South Carolina to Florida.

Subgenus B. *Oynomyanax*.

1028. *Putorius nigripes* Audubon & Bachman.

Putorius nigripes Aud. & Bach., N. Am. Quadr., II, 1851, p. 297.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 341, fig. 68. Zoöl. Ser.

BLACK-FOOTED MINK.

Type locality. Plains of the Platte River, Nebraska.

Geogr. Distr. Kansas to Montana and North Dakota, east of the Rocky Mountains.

Subgenus C. *Arctogale*. Weasels.

1029. *Putorius cicognani* (Bonaparte).

Mustela (Putorius) vulgaris Rich., Faun. Bor. Amer., Mamm., 1, 1829, p. 45. (nec Brisson.)

Mustela cicognanii Bonap., Iconog. Faun. Ital., 1, fasc. xxii, 1838, p. 4.

Putorius cicognani Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 344, fig. 69. Zoöl. Ser.

COMMON WEASEL.

Type locality. Northeastern North America. Carlton House?

Geogr. Distr. Forest country from New England and Labrador to coast of southeastern Alaska and south in the Rocky Mountains to Colorado. In the United States, it is common in New England and New York and in parts of Minnesota. Probably also in northern Michigan and Wisconsin.

a. Putorius cicognani richardsoni (Bonaparte).

Mustela richardsonii Bonap., Charlesw. Mag. Nat. Hist., 11, 1838, p. 38.

Putorius cicognani richardsoni Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 344. Zoöl. Ser.

RICHARDSON'S WEASEL.

Type locality. Fort Franklin, Great Bear Lake, Mackenzie, Canada.

Geogr. Distr. Hudson Bay to interior of Alaska and British Columbia, Canada.

b. Putorius cicognani alascensis (Merriam).

Putorius richardsoni alascensis Merr., N. Am. Faun., No. 11, 1896, p. 12.

Putorius cicognani alascensis Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 344. Zoöl. Ser.

ALASKAN WEASEL.

Type locality. Juneau, Alaska.

Geogr. Distr. Alaska. Range unknown.

1030. *Putorius microtis* Allen.

Putorius microtis Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1903, p. 563.

SMALL-EARED WEASEL.

Type locality. Shesley, headwaters of the Shesley River, British Columbia, Canada.

Geogr. Distr. Range unknown.

Genl. Char. Similar to *P. c. richardsoni*, but smaller and with smaller ears. Skull: with very broad rostral and interorbital portions and broad nasals; postpalatal fossa small, U-shaped; audital bullæ small, flat.

Color. Above dark brown tinged with golden; beneath white tinged with sulphur yellow; toes white on all feet.

Measurements. Total length, 295; tail vertebræ, 82; hind foot, 37; ear from crown, 19. Skull: total length, 40; zygomatic breadth, 20.5; length of nasals, 7; width of nasals at middle, 3.5; palatal length, 14; interorbital breadth, 9; postorbital breadth, 9.5; across postorbital processes, 11; mastoid breadth, 14; length of bullæ, 12; width of bullæ, 7; length of mandible, 20; length of molar series, 9; height of mandible at coronoid process, 9.

1031. *Putorius streator* Merriam.

Putorius streator Merr., N. Am. Faun., No. 11, 1896, p. 13. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 345. Zool. Ser.

STREATOR'S WEASEL.

Type locality. Mount Vernon, Skagit Valley, Skagit County, Washington.

Geogr. Distr. Coast of Oregon and Washington to Puget Sound.

a. Putorius streator leptus Merriam.

Putorius streator leptus Merr., Proc. Biol. Soc. Wash., XVI, 1903, p. 76.

FEEBLE WEASEL.

Type locality. Silverton, Colorado.

Geogr. Distr. Range unknown.

Genl. Char. Similar to *P. streator*, but smaller; black tip to tail, 15 mm.

Color. Summer Pelage. Above uniform drab brown; end of tail black; under parts and inner side of limbs to wrists and ankles, white; rest of limbs and feet brown like the back; toes on fore feet white; those of hind feet mixed brown and white.

Measurements. Total length, 243; tail vertebræ, 64; hind foot, 31. Skull: basal length, 31.5; Hensel, 30.5; zygomatic breadth, 17.5; interorbital breadth, 7.5; palatal length, 13; postpalatal length, 18; length of tooth row with canine, 9.

1032. *Putorius muricus* Bangs.

Putorius (Arctogale) muricus Bangs, Proc. New Engl. Zool. Club, 1, 1899, p. 71. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, Append., p. 434. Zool. Ser.

MOUSING WEASEL.

Type locality. Echo, El Dorado County, Colorado.

Geogr. Distr. Colorado. Range unknown.

1033. *Putorius alleghanensis* Rhoads.

Putorius alleghanensis Rhoads, Proc. Acad. Nat. Scien. Phil., 1901, p. 751. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, Suppl., p. 511. Zoöl. Ser.

ALLEGHANY MOUNTAINS WEASEL.

Type locality. Beallsville, Washington County, Pennsylvania.

Geogr. Distr. Known only from Washington and Allegheny Counties, Pennsylvania.

1034. *Putorius rixosus* Bangs.

Putorius rixosus Bangs, Proc. Biol. Soc. Wash., x, 1896, p. 21.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 345.

Zoöl. Ser.

PUGNACIOUS WEASEL.

Type locality. Osler, Saskatchewan, Canada.

Geogr. Distr. Hudson Bay to coast of Alaska; northern Minnesota to Montana.

a. *Putorius rixosus eskimo* Stone.

Putorius rixosus eskimo Stone, Proc. Acad. Nat. Scien. Phil.,

1900, p. 44. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11,

1901, p. 345. Zoöl. Ser.

ESKIMO WEASEL.

Type locality. Point Barrow, Alaska.

Geogr. Distr. Northwestern Alaska. Range unknown.

1035. *Putorius arcticus* Merriam.

Putorius arcticus Merr., N. Am. Faun., No. 11, 1896, p. 15. Elliot,

Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 345. Zoöl.

Ser.

ARCTIC WEASEL.

Type locality. Point Barrow, Alaska.

Geogr. Distr. Arctic coast and Tundras.

a. *Putorius arcticus kadiacensis* Merriam.

Putorius arcticus kadiacensis Merr., N. Am. Faun., No. 11, 1896,

p. 16. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901,

p. 346. Zoöl. Ser.

KADIAK ISLAND WEASEL.

Type locality. Kadiak Island, Alaska.

Geogr. Distr. Known only from the type locality.

1036. *Putorius haidarum* Preble.

Putorius haidarum Preble, Proc. Biol. Soc. Wash., xii, 1898,
p. 169. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., ii, 1901,
p. 351. Zoöl. Ser.

HAIDA WEASEL.

Type locality. Massett, Graham Island, Queen Charlotte Islands,
British Columbia, Canada.

Geogr. Distr. Queen Charlotte Islands, British Columbia, Canada.

1037. *Putorius noveboracensis* Emmons.

Putorius noveboracensis Emmons, Rep. Quadr. Mass., 1840, p. 45.
Elliot, Syn. N. Am. Mamm., F. C. M. Pub., ii, 1901, p. 346.
Zoöl. Ser.

NORTHERN WEASEL.

Type locality. Southern New York.

Geogr. Distr. Maine, south to North Carolina, west to Illinois.

a. *Putorius noveboracensis notius* Bangs.

Putorius noveboracensis notius Bangs, Proc. N. Engl. Zoöl. Club,
i, 1899, p. 53. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., ii,
1901, p. 346. Zoöl. Ser.

SOUTHERN WEASEL.

Type locality. Weaverville, Buncombe County, North Carolina.

Geogr. Distr. Eastern Middle States.

1038. *Putorius occisor* Bangs.

Putorius occisor Bangs, Proc. N. Engl. Zoöl. Club, i, 1899, p. 54.

Putorius noveboracensis occisor Elliot, Syn. N. Am. Mamm., F. C.
M. Pub., ii, 1901, p. 347. Zoöl. Ser.

MURDEROUS WEASEL.

Type locality. Bucksport, near mouth of Penobscot River, Han-
cock County, Maine.

Geogr. Distr. New England States; Canada? Range unknown.

1039. *Putorius washingtoni* Merriam.

Putorius washingtoni Merr., N. Am. Faun., No. 11, 1896, p. 18. Elliot,
Syn. N. Am. Mamm., F. C. M. Pub., ii, 1901, p. 347. Zoöl. Ser.

WASHINGTON WEASEL.

Type locality. Trout Lake, south base of Mt. Adams, Skamania
County, Washington.

Geogr. Distr. Washington. Range unknown.

1040. *Putorius peninsulæ* Rhoads.

Putorius peninsulæ Rhoads, Proc. Acad. Nat. Scien. Phil., 1894,
p. 152. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., ii, 1901,
p. 347. Zoöl. Ser.

FLORIDA WEASEL.

Type locality. Hudson, Pasco County, fourteen miles north of Tarpon Springs, Hillsboro County, Florida.

Geogr. Distr. Florida Peninsula.

1041. *Putorius longicaudus* (Bonaparte).

Mustela (Putorius) erminea Rich., Faun. Bor. Amer., Mamm., 1, 1829, p. 46. (Part.)

Mustela longicauda Bonap., Charlesw. Mag. Nat. Hist., 11, 1838, p. 38.

Putorius longicaudus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 348. Zoöl. Ser.

LONG-TAILED WEASEL.

Type locality. Carlton House, on North Saskatchewan River, Saskatchewan, Canada.

Geogr. Distr. Great Plains from Kansas northward.

a. *Putorius longicaudus oribasus* Bangs.

Putorius (Arctogale) longicauda oribasus Bangs, Proc. N. Engl. Zoöl. Club, 1, 1899, p. 81.

Putorius longicaudus oribasus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 348. Zoöl. Ser.

MOUNTAIN WEASEL.

Type locality. Source of Kettle River, British Columbia, Canada. Altitude, 7,500 feet.

Geogr. Distr. British Columbia, Canada. Range unknown.

b. *Putorius longicaudus spadix* Bangs.

Putorius longicauda spadix Bangs, Proc. Biol. Soc. Wash., x, 1896, p. 8.

Putorius spadix Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 348. Zoöl. Ser.

CHESTNUT-BROWN WEASEL.

Type locality. Fort Snelling, Hennepin County, Minnesota.

Geogr. Distr. Minnesota. Range unknown.

1042. *Putorius saturatus* Merriam.

Putorius saturatus Merr., N. Am. Faun., No. 11, 1896, p. 21.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 348. Zoöl. Ser.

RICHLY-COLORED WEASEL.

Type locality. Siskiyou, Jackson County, Oregon. Altitude, 4,000 feet.

Geogr. Distr. Oregon, Washington, and part of British Columbia, in the Cascade and Siskiyou Mountains.

1043. *Putorius arizonensis* Mearns.

Putorius arizonensis Mearns, Bull. Am. Mus. Nat. Hist. N. Y., 1891, p. 234. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 349. Zoöl. Ser.

ARIZONA WEASEL.

Type locality. San Francisco forest near Flagstaff, Coconino County, Arizona.

Geogr. Distr. Sierra Nevada and Rocky Mountains into British Columbia, not north of Siskiyou Mountains in the Cascade Range.

1044. *Putorius alleni* Merriam.

Putorius alleni Merr., N. Am. Faun., No. 11, 1896, p. 24. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 349. Zoöl. Ser.

ALLEN'S WEASEL.

Type locality. Custer, Black Hills, Custer County, South Dakota.

Geogr. Distr. Black Hills, South Dakota. Range unknown.

1045. *Putorius xanthogenys* (Gray).

Mustela xanthogenys Gray, Ann. Mag. Nat. Hist., XI, 1st Ser., 1843, p. 118.

Putorius xanthogenys Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 349. Zoöl. Ser.

YELLOW-CHEEKED WEASEL.

Type locality. Southern California; San Diego? San Diego County.

Geogr. Distr. Both sides of the Sierra Nevada, California.

a. Putorius xanthogenys mundus Bangs.

Putorius xanthogenys mundus Bangs, Proc. N. Engl. Zoöl. Club, 1, 1899, p. 56. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 350. Zoöl. Ser.

DECORATED WEASEL.

Type locality. Point Reyes, Marin County, California.

Geogr. Distr. Northern California.

b. Putorius xanthogenys oregonensis Merriam.

Putorius xanthogenys oregonensis Merr., N. Am. Faun., No. 11, 1896, p. 25. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 350. Zoöl. Ser.

OREGON WEASEL.

Type locality. Grant's Pass, Rogue River Valley, Josephine County, Oregon.

Geogr. Distr. Oregon. Range unknown.

1046. *Putorius frenatus* (Lichtenstein).

Mustela frenata Licht., Darst. neuer oder wenig bekannt. Säugeth., 1832, pl. XLII.

Putorius frenatus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 350. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 532, figs. 101, LXXV. Zoöl. Ser.

BRIDLED WEASEL.

Type locality. Valley of Mexico, near City of Mexico, Mexico.

Geogr. Distr. From State of Mexico north into Texas.

a. *Putorius frenatus neomexicanus* Barber & Cockerell.

Putorius frenatus neomexicanus Barb. & Cockerell, Proc. Acad. Nat. Scien. Phil., 1898, p. 188. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 351. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 533. Zoöl. Ser.

RIO GRANDE BRIDLED WEASEL.

Type locality. Armstrong's Lake, Mesilla Valley, Donna Ana County, New Mexico.

Geogr. Distr. Northeastern Mexico to 100 miles north in the Rio Grande Valley, New Mexico.

b. *Putorius frenatus goldmani* Merriam.

Putorius frenatus goldmani Merr., N. Am. Faun., No. 11, 1896, p. 28. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 533. Zoöl. Ser.

GOLDMAN'S BRIDLED WEASEL.

Type locality. Pinabete, State of Chiapas, Mexico. Altitude, 8,200 feet.

Geogr. Distr. State of Chiapas, Mexico. Range unknown.

c. *Putorius frenatus leucoparia* Merriam.

Putorius frenatus leucoparia Merr., N. Am. Faun., No. 11, 1896, p. 29. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 534. Zoöl. Ser.

MICHOACAN BRIDLED WEASEL.

Type locality. Patzcuaro, State of Michoacan, Mexico.

Geogr. Distr. State of Michoacan, Mexico. Range unknown.

1047. *Putorius tropicalis* Merriam.

Putorius tropicalis Merr., N. Am. Faun., No. 11, 1896, p. 30. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 531. Zoöl. Ser.

TROPICAL WEASEL.

Type locality. Jico, State of Vera Cruz, Mexico.

Geogr. Distr. State of Vera Cruz, Mexico, and along the coast to Guatemala.

a. *Putorius tropicalis perdis* Merriam.

Putorius tropicalis perdis Merr., Proc. Biol. Soc. Wash., xv, 1902, p. 67. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 531. Zool. Ser.

TEAPA WEASEL.

Type locality. Teapa, State of Tabasco, Mexico.

Geogr. Distr. Known only from the type locality.

1048. *Putorius affinis* (Gray).

Mustela affinis Gray, Ann. Mag. Nat. Hist., 4th Ser., xiv, 1874, p. 375.

Putorius affinis Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 534. Zool. Ser.

ALLIED WEASEL. *Collaraja* in Costa Rica.

Type locality. Colombia.

Geogr. Distr. Costa Rica south into South America.

Subfamily III. **Lutrinæ.**

Genus 131. **Lutra. Otters.**

Lutra Briss., Regn. Anim., 2d ed., 1762, p. 201. Type *Mustela lutra* Linnæus.

Lataxina Gray, List. Spec. Mamm. Brit. Mus., 1843, pp. xxi, 70.

Lataxia Gerv., Hist. Nat. Mamm., II, 1855, p. 118.

1049. *Lutra canadensis* (Schreber).

Mustela lutra canadensis Schreb., Säugth., 1776, pl. cxxvi B; Text, 1778.

Mustela (Lutra) canadensis Kerr, Anim. Kingd., 1792, No. 337.

Lutra hudsonica F. Cuv., Suppl. Buffon, I, 1831, p. 194.

Lutra canadensis Aud. & Bachm., N. Am. Quad., III, 1853, p. 97, pl. cxxii (*Lataxina mollis* Gray). Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 352. Zool. Ser.

CANADA OTTER.

Type locality. Eastern Canada. Exact locality unknown.

Geogr. Distr. Eastern North America, north of North Carolina on the Atlantic coast.

a. *Lutra canadensis lataxina* (F. Cuvier).

Lutra lataxina F. Cuv., Dict. Scien. Nat., xxvii, 1823, p. 242.

Lutra canadensis lataxina Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 352. Zool. Ser.

SOUTHERN OTTER.

Type locality. South Carolina.

Geogr. Distr. Eastern North America from North Carolina to Alabama and Mississippi.

b. Lutra canadensis vaga (Bangs).

Lutra hudsonica vaga Bangs, Proc. Bost. Soc. Nat. Hist., 1898, p. 224.

Lutra canadensis vaga Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 352. Zoöl. Ser.

FLORIDA OTTER.

Type locality. Micco, Brevard County, Florida.

Geogr. Distr. Eastern Georgia and Florida.

c. Lutra canadensis pacifica (Rhoads).

Lutra hudsonica pacifica Rhoads, Trans. Am. Philos. Soc., N. S., XIX, 1898, p. 429.

Lutra canadensis pacifica Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 353. Zoöl. Ser.

PACIFIC OTTER.

Type locality. Lake Kichelos, Kittitas County, Washington. Altitude, 8,000 feet.

Geogr. Distr. Pacific slopes. California to Alaska.

d. Lutra canadensis sonora (Rhoads).

Lutra hudsonica sonora Rhoads, Trans. Am. Philos. Soc., N. S., XIX, 1898, p. 431.

Lutra canadensis sonora Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 353, pl. xxxvii. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 11, 1904, p. 535, fig. LXXVI. Zoöl. Ser.

MEXICAN OTTER.

Type locality. Montezuma Well, Beaver Creek, Yavapai County, Arizona.

Geogr. Distr. State of Sonora, Mexico, to Wyoming.

1050. *Lutra periclyzomæ* Elliot.

Lutra periclyzomæ Elliot, Proc. Biol. Soc. Wash., XVIII, 1905, p. 80.

SEA-GIRT OR ISLAND OTTER.

Type locality. From Gawi, west coast of Moresby Isl., Queen Charlotte Islands, British Columbia, Canada. No. 491, Field Columbian Mus., Chicago.

Genl. Char. Skull: size large; intertemporal region long and narrow, greatly constricted for its entire length; in young animals this constriction is not so apparent. Rostrum short and very broad;

postorbital processes greatly elongated; braincase without crests, rounded, and widest posteriorly; bullæ small and flat, barely rising above the level of the basioccipital; pterygoid fossa broad for its entire length, the sides nearly straight, but widening gradually to tips of pterygoid processes where the width is greatest; upper molars very large, the cusps on exterior side very high and acute; last molar square-shaped, slightly widest on interior edge.

Measurements. Skull: total length, 122; Hensel, 107; zygomatic width, 81; intertemporal constriction, 20; width across postorbital processes, 40; width of rostrum at canines, 30.5; palatal length, 55; length of upper tooth row, anterior edge of canine to posterior edge of last molar, 40; length of nasals, 17.5; total length of mandible, angle to tips of incisors, 76; length of lower tooth row, anterior edge of canine to posterior edge of last molar, alveolar border, 46; length of canine, 13; length of lower carnassial, 15.

1051. *Lutra annectens* Forsyth-Major.

Lutra annectens Forsyth-Major, Zool. Anzeiger, xx, 1897, p. 142.

Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv,

Pt. II, 1904, p. 536, pl. LI. Zoöl. Ser.

CENTRAL AMERICAN OTTER. *Nutria* in Costa Rica.

Type locality. Rio de Tepic, Territorio de Tepic, State of Jalisco, Mexico.

Geogr. Distr. Mexico to Costa Rica. Limits of range unknown.

1052. *Lutra degener* Bangs.

Lutra degener Bangs, Proc. Biol. Soc. Wash., xii, 1898, p. 35.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 353.

Zoöl. Ser.

DEGENERATE OTTER.

Type locality. Bay St. George, Newfoundland.

Geogr. Distr. Newfoundland.

Genus 132. **Latax. Sea Otters.**

Latax Gloger, Nov. Acta Phys. Med. Acad. Cæs. Leopold.-Carol., xiii, Pt. II, 1827, p. 511. Type *Lutra marina* Erxleben = *Mustela lutris* Linnæus.

Pusa Oken, Lehrb. Naturg., Theil III, 2te Abth., 1816, p. 986. (nec Scopoli.)

Enhydra Flem., Philos. Zoöl., II, 1822, p. 187. (nec *Enhydris* Merrem, Rept.)

1053. *Latax lutris* (Linnæus).

Mustela lutris Linn., Syst. Nat., I, 1758, p. 45; I, 1766, p. 66.

Lutra marina Erxl., Syst. Regn. Anim., 1777, p. 445.

Pusa orientalis Oken, Lehrb. Naturg., Theil III, 2te Abth., 1816, p. 986.

Lutra stelleri Less., Man. Mamm., 1827, p. 156.

Enydriis (sic) gracilis Fisch., Syn. Mamm., 1829, p. 229.

Latax lutris Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 354, pl. XXXVIII. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 537, pl. LII. Zoöl. Ser.

NORTHERN SEA OTTER.

Type locality. "America Septentrionali." Coasts of Bering Sea.

Geogr. Distr. Coasts of Bering Sea and of the north Pacific Ocean, south on American coast possibly to California.

a. *Latax lutris nereis* Merriam.

Latax lutris nereis Merr., Proc. Biol. Soc. Wash., XVII, 1904, p. 159. SAN MIGUEL ISLAND SEA OTTER.

Type locality. San Miguel Island, Santa Barbara Islands, Santa Barbara County, California.

Geogr. Distr. Santa Barbara Islands; also probably south to the coast of Lower California, and it may be the species found rarely among the Kelp beds of San Tomas and San Quentin Bays.

Cranial Characters. Skull alone known. This is "broad and high, with long and high sagittal crest and swollen braincase. Compared with *lutris* the following differences appear: Skull as a whole less flattened, braincase more swollen and rounded, the sides (viewed from above) more convex and swollen, especially behind the constriction; anterior part of zygomata more broadly and squarely expanded; basioccipital forming an angle with basisphenoid; coronoid process sloping strongly backward; sagittal crest much higher and more decurved posteriorly; inner cusp of large upper premolar (*pm.* 3) elongated along anterior part of inner lobe (instead of conical) and showing a tendency to subdivide into two parts; first lower molar broader and more broadly truncate posteriorly."

Measurements. "The specimen in the flesh measures six feet in length." (Merr. l. c.)

Order VIII. **Pinnipedia. Sea-lions, Seals, etc.**

Family I. **Otariidae. Sea-lions.**

Genus 133. **Eumetopias.**

Eumetopias Gill, Proc. Essex Institute, v, 1866, pp. 7, 11. Type
Otaria stelleri Lesson.

1054. **Eumetopias jubata** (Schreber).

Leo marinus Steller, Nov. Comm. Petrop., xi, 1751, p. 360.

Phoca jubata Schreber, Säugth., iii, 1778, p. 300, pl. LXXXIII B.

Otaria jubata Péron, Voy. Terr. Austr., ii, 1816, p. 40.

Otaria stelleri Lesson, Dict. Class. Hist. Nat., xiii, 1828, p. 420.

Phoca leonina Pall., Zoog. Rosso-Asiat., i, 1831, p. 104.

?*Arctocephalus monteriensis* Gray, Proc. Zool. Soc., 1859, p. 358,
pl. LXXII.

Eumetopias stelleri Elliot, Syn. N. Am. Mamm., F. C. M. Pub., ii,
1901, p. 355, pl. xxxix. Zool. Ser.

STELLER'S SEA-LION.

Type locality. Bay of San Francisco.

Geogr. Distr. Shores of California and of North Pacific Ocean to
Berings Straits, Japan.

Genus 134. **Zalophus.**

Zalophus Gill, Proc. Essex Institute, v, 1866, pp. 7, 11. Type
Otaria gillespii McBain = *Otaria californiana* Lesson.

Neophoca Gray, Ann. Mag. Nat. Hist., 3d Ser., xviii, 1866, p.
231.

1055. **Zalophus californianus** (Lesson).

Otaria californiana Less., Dict. Class. Hist. Nat., xiii, 1828, p. 420.

Otaria gillespii McBain, Proc. Roy. Soc. Edinb., i, 1858, p. 422.

Zalophus californianus Elliot, Syn. N. Am. Mamm., F. C. M. Pub.,
ii, 1901, p. 356, pl. xxxix. Zool. Ser. *Id.* Mamm. Middle
Amer. & W. Indies, F. C. M. Pub., iv, Pt. ii, 1904, p. 539, fig.
LXXVII, pl. LIII. Zool. Ser.

CALIFORNIA SEA-LION.

Type locality. California.

Geogr. Distr. West coast of the United States from Bay of San
Francisco to Coronado and Tres Marias Islands, and coast of Lower
California, Mexico.

Genus 135. **Callotaria. Fur Seal.**

Callotaria Palmer, Proc. Biol. Soc. Wash., VII, 1892, p. 156. Type
Phoca ursina Linnæus.

Callorhinus Gray, Proc. Zool. Soc., 1859, p. 359. (nec Blanchard,
Entomol.; nec Girard, Herpetology.)

Arctocephalus Gill, Proc. Essex Inst., v, 1866, pp. 7, 11. (nec
F. Cuv.)

Otoes Palmer, Proc. Biol. Soc. Wash., XIV, 1901, p. 133. (nec
Fischer.)

1056. Callotaria alascana (Jordan & Clark).

Callorhinus alascanus Jord. & Clark, The Fur Seals and Fur Seal
Isl., Pt. 3, 1899, p. 2.

Callotaria ursina Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11,
1901, p. 357, pl. XL. (nec Linnæus.)

PRIBILOF ISLANDS FUR SEAL.

Type locality. Pribilof Islands, Bering Sea.

Geogr. Distr. Pacific shores of North America; Pribilof Islands,
Bering Sea.

Genus 136. **Arctocephalus. Sea Bears.**

Arctocephalus F. Cuv., Dict. Scien. Nat., xxxix, 1826, p. 554. Type

**Phoca antarctica* Thunberg.

Halarctus Gill, Proc. Essex Inst., v, 1866, pp. 7, 11.

Arctophoca Peters, Monatsb. K. Preuss. Akad. Wiss. Berl., 1866,
p. 276, Tab. 11, A. B. C.

Euotaria Gray, Ann. Mag. Nat. Hist., 3d Ser., xviii, 1866,
p. 236.

Gypsophoca Gray, Ann. Mag. Nat. Hist., 3d Ser., xviii, 1866, p. 236.

1057. Arctocephalus townsendi Merriam.

Arctocephalus townsendi Merr., Proc. Biol. Soc. Wash., xi, 1897,
p. 178. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M.
Pub., IV, Pt. 11, 1904, p. 544, fig. LXXIX, pls. LX, LXI, LXII.
Zool. Ser.

GUADALUPE FUR SEAL.

Type locality. Guadalupe Island, off coast Lower California,
Mexico.

Geogr. Distr. Guadalupe and San Benito Island, off coast Lower
California, Mexico.

*See Allen, Princt. Univ. Patagon. Exped., Zool., III, 1905, p. 120.

Family II. **Odontobænidae.**Genus 137. **Odontobænus. Walrus.**

Odobenus (*sic*) Briss., Reg. Anim., 2d ed., 1762, pp. 12, 30. Type
Phoca rosmarus Linnæus.

1058. Odontobænus rosmarus (Linnæus).

Phoca rosmarus Linn., Syst. Nat., 1, 1758, p. 38.

Trichechus rosmarus Linn., Syst. Nat., 1, 1766, p. 49. Elliot, Syn.
N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 358, pl. xli. Zoöl.
Ser.

ATLANTIC WALRUS.

Type locality. Eastern part of Arctic Sea. North Atlantic.

Geogr. Distr. Coast of Labrador north to Arctic Ocean; shores
of Greenland, boreal islands and polar areas.

1059. Odontobænus obesus (Illiger).

Trichechus rosmarus Shaw, Gen. Zoöl., 1, 1800, p. 234, fig. 68.
(Part.)

Trichechus obesus Illig., Abhandl. d. Akad. Berl., 1815, pp. 64, 70,
75. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901,
p. 358. Zoöl. Ser.

Trichechus divergens Illig., Abhandl. d. Akad. Berl., 1815, p. 68.

Rosmarus cookii Fremery, Bydrag. tot de naturk. Wetensch., vi,
1831, p. 385.

Rosmarus arcticus Pall., Zoog. Rosso-Asiat., 1831, p. 269, pls. xxviii,
xxix.

PACIFIC WALRUS.

Type locality. Western Arctic Seas.

Geogr. Distr. Northwest coast of North America in Arctic Ocean
and Bering Sea and certain of the latter's islands; Kotzbue and Norton
sounds and Bristol Bay; also on northeastern coast of Asia.

Family III. **Phocidae.**Subfamily I. **Cystophorinæ.**Genus 138. **Mirounga. Sea Elephants.**

Mirounga Gray, in Griff., Cuvier's Anim. Kingd., v, 1827, p. 179.
(Part.) Type *Phoca proboscidea* Péron = *Phoca leonina* Lin-
næus.

Macrorhinus F. Cuv., Dict. Scien. Nat., xxxix, 1826, p. 552. (nec
Latreille, Coleopt., 1825.)

Rhinophoca Wagl., Nat. Syst. Amph., 1830, p. 27.

Morunga Gray, List Ost. Spec. Brit. Mus., 1847, p. 33.

1060. *Mirounga angustirostris* (Gill).

Macrorhinus angustirostris Gill, Proc. Chicago Acad. Scien., 1866, p. 33. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 359, pls. XLII, XLIII, XLIV. Zool. Ser.

Mirounga angustirostris Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 545, fig. LXXX, pls. LXIII, LXIV, LXV, LXVI. Zool. Ser.

ELEPHANT SEAL.

Type locality. St. Bartholomew's Bay, Lower California, Mexico.

Geogr. Distr. Formerly from Cape San Lazaro, Lower California, Mexico, to Point Reyes, Marine County, California.

Genus 139. **Cystophora. Hooded Seals.**

Cystophora Nilss., Skand. Faun., I, 1820, p. 382. Type *Cystophora borealis* Nilsson = *Phoca cristata* Erxleben.

Stemmatope F. Cuv., Mém. Mus. Hist. Nat. Paris, XI, 1824, p. 196.

Stemmatopus F. Cuv., Dict. Scien. Nat., XXXIX, 1826, p. 550.

1061. *Cystophora cristata* (Erxleben).

Phoca cristata Erxl., Syst. Reg. Anim., 1777, p. 590.

Phoca cucullata Bodd., Elench. Anim., 1785, p. 107.

Cystophora borealis Nilss., Skand. Faun., 1820, p. 383.

Phoca mitrata G. Cuv., Oss. Foss., 3me ed., v, 1825, p. 210, pl. XVIII, fig. 3.

Phoca isidorei Less., Rev. Zool., 1843, p. 256.

Cystophora cristata Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 360, pl. XLV. Zool. Ser.

HOODED SEAL.

Type locality. Southern Greenland.

Geogr. Distr. Colder portions of the North Atlantic Ocean; Arctic Ocean. Rarely found south of Newfoundland.

Subfamily II. **Phocinæ.**

Genus 140. **Halichærus. Gray Seals.**

Halichærus Nilss., Skand. Faun., I, 1820, p. 376. Type *Phoca grypus* Fabricius.

Pusa Gill, Johns. New Univ. Cyclop., 181, 1877, p. 1226. (nec Scopoli.)

1062. Halichærus grypus (Fabricius).

Phoca grypus Fabr., Skriv. af Natur.-Selsk., I, 2, 1791, p. 167, pl. XIII, fig. 4. (Skull.)

Halichærus grypus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 361, pl. XLVI. Zoöl. Ser.

GRAY SEAL.

Type locality. Iceland?

Geogr. Distr. North Atlantic from Nova Scotia to Greenland, east to British Islands.

Genus 141. **Phoca. Hair Seals.**

Phoca Linn., Syst. Nat., I, 1758, p. 37. Type *Phoca vitulina* Linnæus.

Pusa Scop., Intr. Hist. Nat., 1777, p. 490.

Calocephalus F. Cuv., Dict. Scien. Nat., xxxix, 1826, p. 544.

Pagophilus Gray, Voy. Erebus & Terror, Zoöl., 1844, p. 3.

Pagomys Gray, Proc. Zoöl. Soc., 1864, p. 31.

Erignathus Gill, Proc. Essex Inst., v, 1866, pp. 5, 9.

Haliphilus Gray, Ann. Mag. Nat. Hist., xvii, 4th Ser., 1866, p. 446.

Histriophoca Gill, Amer. Nat., vii, 1873, p. 179.

Subgenus A. **Erignathus.****1063. Phoca barbata** Fabricius.

Phoca barbata Fabr., Müller's Zool. Dan. Prod., viii, 1776. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 362, pl. XLVI. Zoöl. Ser.

Phoca leporina Lepech., Act. Acad. Petrop., I, 1777, p. 264, pls. 8, 9. (juv.)

Phoca parsonsi Less., Dict. Class. Hist. Nat., xiii, 1828, p. 414.

Phoca lepechini Less., Dict. Class. Hist. Nat., xiii, 1828, p. 415.

Phoca nautica Pallas, Zoog. Rosso-Asiat., I, 1831, p. 108.

Phoca albigena Pallas, Zoog. Rosso-Asiat., 1831, p. 109.

Phoca naurica Gray, Cat. Seals & Whales, Suppl., 1871, p. 3.

BEARDED SEAL.

Type locality. Coast of Cornwall, England.

Geogr. Distr. Circumpolar, boreal; south on eastern coast of North America to Newfoundland.

Subgenus B. **Histriophoca.****1064. Phoca fasciata** Zimmermann.

Phoca fasciata Zimm., Geog. Gesch., III, 1783, p. 277. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 362, pls. XLVII, XLVIII. Zoöl. Ser.

Phoca equestris Pallas, Zoog. Rosso-Asiat., 1, 1831, p. 111.

RIBBON SEAL.

Type locality. Kurile Islands, Japan.

Geogr. Distr. Bering Sea south to the Aleutian Islands and to the island of Saghalien on the Asiatic coast, and Kurile Islands, Japan.

Subgenus C. *Phoca*.

1065. *Phoca vitulina* Linnæus.

Phoca vitulina Linn., Syst. Nat., 1, 1758, p. 38; 1, 1766, p. 56.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 363, pl. XLIX. Zoöl. Ser.

Phoca variegata Nilss., Skand. Faun., 1, 1820, p. 359.

Phoca scopulicola Thienem., Reis. Nord. Eur., 1, 1824, p. 59, pl. v. (ad male.)

Phoca littorea Thienem., Reis. Nord. Eur., 1, 1824, p. 61, pl. vi (male), pls. VII, VIII (skulls).

Phoca pealei Gill, Proc. Essex Inst., v, 1866, pp. 4, 13.

Phoca fætida Bartl., Proc. Zoöl. Soc., 1868, p. 402.

HARBOR SEAL.

Type locality. Coast of Europe.

Geogr. Distr. Coast of North Atlantic Ocean from New York to the Arctic Ocean; coast of Greenland and other eastern boreal islands. Northern coast of Europe.

1066. *Phoca richardi* (Gray).

Halicyon richardii Gray, Proc. Zoöl. Soc., 1864, p. 28, fig. 1.

Phoca richardii Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1902, p. 491.

Phoca largha Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 363. Zoöl. Ser. (nec Pallas.)

RICHARD'S HARBOR SEAL.

Type locality. Vancouver Island.

Geogr. Distr. Northern Pacific Ocean, Bering Sea.

a. Phoca richardi pribilofensis Allen.

Phoca richardii pribilofensis Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1902, p. 495.

PRIBILOF ISLANDS SEAL.

Type locality. St. Paul Island, Pribilof Island, Bering Sea.

Geogr. Distr. Bering Sea. Range unknown.

Genl. Char. Dentition weaker than that of *P. richardi*.

Color. Above silvery yellowish white, spotted with dark brown and blackish, confluent from the back of the head to middle of back

where the general color is blackish; beneath pale ochraceous varied with paler streaks and blotched with dusky, the spots blacker and more distinct on the throat and sides of neck; sides paler and heavily blotched with blackish.

Measurements. None given.

b. Phoca richardii geronimensis Allen.

Phoca richardii geronimensis Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1902, p. 495. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 542, fig. LXXVIII (*P. richardii*), pls. LIV, LV (*P. r. geronimensis*). Zoöl. Ser.

SAN GERONIMO HARBOR SEAL.

Type locality. San Geronimo Island, Lower California, Mexico.

Geogr. Distr. San Geronimo Island to Santa Barbara Island, Santa Barbara County, California. Range unknown in North Pacific Ocean.

1067. *Phoca stejnegeri* Allen.

Phoca stejnegeri Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1902, p. 485.

STEJNEGER'S HARBOR SEAL.

Type locality. Shores of Bering Island.

Geogr. Distr. Bering Sea and shores of Kamschatka.

Genl. Char. Similar to *P. vitulina*, but larger. A deep groove for the maxillary nerve runs from the infraorbital foramen to a point opposite the middle of upper premolar third.

Color. Light phase. Above deep straw yellow spotted with black, beneath deeper yellow and more sparsely spotted. Whiskers yellowish brown, nails brownish black.

Dark phase. Ground color deep yellow, but the black spots occupy four-fifths of dorsal surface, and over one-half of the ventral surface, beneath the spots on the median half of the dorsal region are irregular spots or bands of same color. Dark patches on limbs.

Measurements. Nose to end of tail, 1,545; to end of hind flipper, 1,760; tail, 140; hind foot, 335.

Subgenus D. *Pusa*.

1068. *Phoca hispida* Schreber.

Phoca hispida Schreb., Säugth., III, 1776, p. 312, pl. LXXXVI. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 364. Zoöl. Ser.

Phoca fetida Fabr., Müller's Zool. Dan. Prod., VIII, 1776.

Phoca annellata Nilss., Skand. Faun., I, 1820, p. 362.

Calocephalus discolor F. Cuv., Dict. Sci. Nat., xxxix, 1826, p. 554.

Phoca schreberi Less., Dict. Class. Hist. Nat., xiii, 1828, p. 414.

Phoca frederici Less., Dict. Class. Hist. Nat., xiii, 1828, p. 416.

Callocephalus dimidiatus Gray, Cat. Seals & Whales, 1866, p. 22.

RINGED SEAL.

Type locality. Greenland.

Geogr. Distr. Circumpolar; south to Labrador on the Atlantic coast, and northern shores of Bering Sea on the Pacific coast.

Subgenus E. *Pagophilus*.

1069. *Phoca groenlandica* Fabricius.

Phoca groenlandica Fabr., Müller's Zool. Dan. Prod., viii, 1776.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., ii, 1901, p. 364, pl. XLIX. Zoöl. Ser.

Phoca oceanica Lepech., Act. Acad. Petrop., i, 1777, p. 295, pls. vii, viii.

Phoca groenlandica var. *nigra* Kerr., Anim. Kingd., 1792, p. 125.

Phoca semilunaris Bodd., Elench. Anim., 1785, p. 170.

Phoca albicauda Desm., Mamm., Suppl., 1822, p. 839.

Phoca lagura G. Cuv., Oss. Foss., v, 1825, p. 206. (juv.)

Phoca mulleri Less., Dict. Class. Hist. Nat., xiii, 1828, p. 413.

Phoca desmaresti Less., Dict. Class. Hist. Nat., xiii, 1828, p. 416.

Phoca pilayi Less., Dict. Class. Hist. Nat., xiii, 1828, p. 416.

Phoca dorsata Pallas, Zoog. Rosso-Asiat., i, 1811, p. 112.

Phoca albinii Alex., Mem. Acad. Torino, ii, 1850, p. 141, pls. 1-4. (Skeleton.)

Phoca annellata Gaim., Voy. en Islands, Atlas, 1851, pl. xi, fig. 9. (nec Nilsson.)

HARP SEAL.

Type locality. Greenland.

Geogr. Distr. Circumpolar; south to Newfoundland in Atlantic Ocean, and in Bering Sea and north Pacific Ocean possibly to Sitka.

Genus 142. *Monachus*.

Monachus Flem., Phil. Zoöl., ii, 1822, p. 187. Type *Phoca monachus* Hermann.

Pelagios F. Cuv., Mém. Mus. Hist. Nat. Paris, xi, 1824, p. 193, pl. 13, fig. 2.

Pelagocyon Glog., Hand-u. Hilfsb. Naturg., 1841, pp. xxxiv, 163.

Rigoon Gistel, Naturg. Thiere, 1848, p. x.

Heliophoca Gray, Ann. Mag. Nat. Hist., 2d Ser., xiii, 1854, p. 201.

1070. *Monachus tropicalis* (Gray).

Phoca tropicalis Gray, Cat. Seals Brit. Mus., 1850, p. 28.

Monachus tropicalis Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., IV, Pt. II, 1904, p. 543, pls. LVI, LVII, LVIII, LIX.

Zoöl. Ser.

WEST INDIAN SEAL.

Type locality. Island of Jamaica.

Geogr. Distr. Formerly from islands off the coast of Yucatan, Mexico, to Bay of Honduras, and eastward to Jamaica, Cuba, Florida Keys, and the Bahamas. At present found only on some Keys north of Cuba and on some scattered islands between Cuba and Yucatan.

Order IX. **Insectivora. The Insectivores.**

Family I. **Soricidae. Shrews.**

Subfamily I. **Soricinae.**

Genus 143. **Sorex. Common Shrews.**

Sorex Linn., Syst. Nat., I, 1758, p. 53; I, 1766, p. 73. Type *Sorex araneus* Linnæus.

1071. **Sorex personatus** I. Geoff. St. Hilaire.

Sorex personatus I. Geoff. St. Hil., Mém. du Mus. Hist. Nat. Paris, xv, 1827, pp. 122-125. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 366. Zoöl. Ser.

Sorex forsteri Rich., Zoöl. Journ., III, 1828, p. 516.

Sorex cooperi Bachm., Journ. Acad. Nat. Scien. Phil., 1837, p. 388, pl. xxiv, fig. 7.

Sorex fimbripes Bachm., Journ. Acad. Nat. Scien. Phil., 1837, p. 391, pl. xxiv, fig. 8.

Amphisorex lesneuri Duvern., Mag. Zool., Mamm., 1842, p. 33, pl. L.

Sorex platyrhynchus Linsley, Sillim. Am. Journ. Scien., XLIII, 1842, p. 346.

Otisorex platyrhinus DeKay, Zoöl. N. Y., I, 1842, p. 22, pl. v, fig. 1.

Sorex haydeni Baird, N. Am. Mamm., 1857, p. 29.

Sorex richardsoni Dobson, Mon. Insectiv., I, Pt. III, 1890, pl. xxiii, fig. 9. (nec Bachm.)

Sorex idahoensis Merr., N. Am. Faun., No. 5, 1891, p. 32.

MASKED SHREW.

Type locality. Eastern United States.

Geogr. Distr. New England to Alaska and south in high mountains to Tennessee and North Carolina. Not found in the southern Rocky Mountains or in the Cascade or Sierra Ranges.

a. Sorex personatus streatori Merriam.

Sorex personatus streatori Merr., N. Am. Faun., No. 10, 1895, p. 62.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 367. Zoöl. Ser.

STREATOR'S SHREW.

Type locality. Yakutat, Alaska.

Geogr. Distr. Range unknown.

b. Sorex personatus arcticus Merriam.

Sorex personatus arcticus Merr., Proc. Wash. Acad. Sci., 11, 1900, p. 17. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 368. Zoöl. Ser.

ARCTIC SHREW.

Type locality. St. Michaels, Norton Sound, Alaska.

Geogr. Distr. Northwestern Alaska.

1072. *Sorex richardsoni* Bachman.

Sorex parvus Richard., Faun. Bor. Amer., 1, Mamm., 1829, p. 8. (nec Say.)

Sorex richardsonii Bachm., Jour. Acad. Nat. Scien. Phil., VII, 1837, p. 383, pl. xxiv, fig. 5. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 368. Zoöl. Ser.

RICHARDSON'S SHREW.

Type locality. Unknown. Plains of the Saskatchewan? Canada.

Geogr. Distr. Northern Minnesota, and Saskatchewan Plains.

1073. *Sorex tundrensis* Merriam.

Sorex tundrensis Merr., Proc. Wash. Acad. Scien., 11, 1900, p. 16. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 368. Zoöl. Ser.

TUNDRA SHREW.

Type locality. St. Michaels, Norton Sound, Alaska.

Geogr. Distr. Northwest coast of Alaska. Range unknown.

1074. *Sorex sphagnicola* Coues.

Sorex sphagnicola Coues, Bull. U. S. Geol. & Geogr. Surv., III, 1877, p. 650. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 368. Zoöl. Ser.

SPHAGNUM MOSS SHREW.

Type locality. Lat. 60°, vicinity of Fort Liard, northwestern British Columbia, Canada.

Geogr. Distr. British Columbia to Hudson Bay, Arctic America, Canada.

1075. *Sorex fumeus* Miller.

Sorex platyrhinus Dobson, Mon. Insectiv., 1, Pt. III, 1890, pl. xxiii, fig. 5. (nec DeKay.)

Sorex fumeus Miller, N. Am. Faun., No. 10, 1895, p. 50. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 369. Zoöl. Ser.

SMOKY SHREW.

Type locality. Peterboro, Madison County, New York.

Geogr. Distr. Eastern United States, Nova Scotia, New Brunswick west to Ontario and the Great Lakes.

1076. *Sorex macrurus* Batchelder.

Sorex macrurus Batchelder., Proc. Biol. Soc. Wash., x, 1896,
p. 133. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901,
p. 369. Zool. Ser.

LONG-TAILED SHREW.

Type locality. Beedes (Keene Heights), Essex County, New York.

Geogr. Distr. Northern New York. Range unknown.

1077. *Sorex setosus* Elliot.

Sorex setosus Elliot, Pub. Field Columb. Mus., I, 1899, p. 274.
Zool. Ser. *Id.* Syn. N. Am. Mamm., F. C. M. Pub., II, 1901,
p. 369. Zool. Ser.

SPINY SHREW.

Type locality. Happy Lake, Olympic Mountains, Clallam County, Washington.

Geogr. Distr. Known only from the type locality.

1078. *Sorex vagrans* Baird.

Sorex vagrans Baird, N. Am. Mamm., 1857, p. 15, pl. xxvi, fig.
1675. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901,
p. 370. Zool. Ser.

Sorex suckleyi Baird, N. Am. Mamm., 1857, p. 18, pl. xxvii,
fig. 1677.

WANDERING SHREW.

Type locality. Shoalwater Bay, Washington.

Geogr. Distr. Southern British Columbia, western Washington and Oregon, and northern California to Monterey on the coast and to Fort Crook in the mountains.

a. *Sorex vagrans dobsoni* (Merriam).

Sorex dobsoni Merr., N. Am. Faun., No. 5, 1891, p. 33, pl. iv, fig. 2.

Sorex vagrans dobsoni Elliot, Syn. N. Am. Mamm., F. C. M. Pub.,
II, 1901, p. 370. Zool. Ser.

DOBSON'S SHREW.

Type locality. Saw Tooth or Alturas Lake, east base of the Saw Tooth Mountains, central Idaho. Altitude, 7,200 feet.

Geogr. Distr. Rocky Mountains in northern Idaho, and western Montana, also in Big Snowy and Prior Mountains, Montana, Big Horn Mountains in Wyoming, and Wahsatch Mountains, Utah.

b. *Sorex vagrans monticola* (Merriam).

Sorex monticolus Merr., N. Am. Faun., No. 3, 1890, p. 43.

Sorex vagrans monticola Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 371. Zool. Ser.

MOUNTAIN SHREW.

Type locality. San Francisco Mountains, Coconino County, Arizona. Altitude, 11,500 feet.

Geogr. Distr. Arizona. Range unknown.

1079. *Sorex amœnus* Merriam.

Sorex amœnus Merr., N. Am. Faun., No. 10, 1895, p. 69. Elliot,

Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 371. Zool. Ser.

CHARMING SHREW.

Type locality. Mammoth Pass, head of Owens River, east slope of the Sierra Nevada, California. Altitude, 10,000 feet.

Geogr. Distr. Northern California.

1080. *Sorex vancouverensis* Merriam.

Sorex vancouverensis Merr., N. Am. Faun., No. 10, 1895, p. 70.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 371.

Zool. Ser.

VANCOUVER ISLAND SHREW.

Type locality. Goldstream, Vancouver Island, British Columbia, Canada.

Geogr. Distr. Vancouver Island, British Columbia, Canada.

1081. *Sorex nevadensis* Merriam.

Sorex nevadensis Merr., N. Am. Faun., No. 10, 1895, p. 71. Elliot,

Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 371. Zool. Ser.

NEVADA SHREW.

Type locality. Reese River, Lander County, Nevada.

Geogr. Distr. Great Basin, Nevada.

1082. *Sorex orizabæ* Merriam.

Sorex orizabæ Merr., N. Am. Faun., No. 10, 1895, p. 71. Elliot,

Mamm. Middle Amer. & W. Indies, F. C. M. Pub., 11, Pt. 11,

1904, p. 550. Zool. Ser.

MINUTE SHREW.

Type locality. Mount Orizaba, State of Puebla, Mexico. Altitude, 9,500 feet.

Geogr. Distr. Southern Mexico, States of Puebla, Tlaxcala, Vera Cruz, Mexico, and Michoacán.

1083. *Sorex obscurus* Merriam.

Sorex vagrans similis Merr., N. Am. Faun., No. 5, 1891, p. 34, pl. 11, fig. 3. (nec Hensel.)

Sorex obscurus Merr., N. Am. Faun., No. 10, 1895, p. 72, pl. 11, figs. 1, 1a. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11,

1901, p. 371. Zool. Ser.

DUSKY SHREW.

Type locality. Timber Creek, Salmon River Mountains, Idaho. Altitude, 8,200 feet.

Geogr. Distr. British Columbia, and mountains of western Washington, Montana, Wyoming, Utah, and Colorado, south in California, to Mt. Whitney.

a. *Sorex obscurus longicaudus* Merriam.

Sorex obscurus longicauda Merr., N. Am. Faun., No. 10, 1895, p. 74.

Sorex obscurus longicaudus Elliot, Syn. N. Am. Mamm., F. C. M.

Pub., 11, 1901, p. 372. Zoöl. Ser.

LONG-TAILED SHREW.

Type locality. Wrangel, Alaska.

Geogr. Distr. Coast of southeast Alaska from Wrangel southward; also the coast of Washington, including Puget Sound and Skagit Valley.

b. *Sorex obscurus ventralis* Merriam.

Sorex obscurus ventralis Merr., N. Am. Faun., No. 10, 1895, p. 75.

Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., 14,

Pt. 11, 1904, p. 550, fig. LXXXI. Zoöl. Ser.

CHESTNUT-BELLIED SHREW.

Type locality. Cerro San Felipe, State of Oaxaca, Mexico. Altitude, 10,000 feet.

Geogr. Distr. State of Oaxaca, Mexico.

c. *Sorex obscurus elassodon* (Osgood).

Sorex longicauda elassodon Osgood, N. Am. Faun., No. 21, 1901,

p. 35.

QUEEN CHARLOTTE SHREW.

Type locality. Cumshewa Inlet, Moresby Island, Queen Charlotte Islands.

Geogr. Distr. Queen Charlotte Islands, British Columbia, Canada.

Genl. Char. Similar to *S. o. longicaudus*, but smaller and tail shorter.

Color. Like *S. o. longicaudus*, underparts paler.

Measurements. Total length, 123; tail vertebræ, 52; hind foot, 14.

d. *Sorex obscurus prevostensis* (Osgood).

Sorex longicauda prevostensis Osgood, N. Am. Faun., No. 21,

1901, p. 35.

PREVOST ISLAND SHREW.

Type locality. Prevost Island, Queen Charlotte Islands, British Columbia, Canada.

Geogr. Distr. Queen Charlotte Islands, British Columbia, Canada.

Genl. Char. Similar to *S. o. longicaudus*, tail shorter; teeth distinctive.

Color. Slightly darker than *S. o. longicaudus*.

Measurements. Total length, 133; tail vertebræ, 58; hind foot, 15.

1084. *Sorex oreopolus* Merriam.

Sorex oreopolus Merr., Proc. Biol. Soc. Wash., VII, 1892, p. 173.

Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV,

Pt. II, 1904, p. 550. Zoöl. Ser.

LOFTY MOUNTAIN SHREW.

Type locality. North slope of the Sierra Nevada de Colima, State of Jalisco, Mexico. Altitude, 10,000 feet.

Geogr. Distr. State of Jalisco, Mexico.

1085. *Sorex glacialis* Merriam.

Sorex glacialis Merr., Proc. Wash. Acad. Scien., II, 1900, p. 16.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 372.

Zoöl. Ser.

GLACIER BAY SHREW.

Type locality. Point Gustavus, east side of Glacier Bay, Alaska.

Geogr. Distr. Alaska. Range unknown.

a. Sorex glacialis alascensis (Merriam).

Sorex obscurus alascensis Merr., N. Am. Faun., No. 10, 1895, p. 76.

Sorex glacialis alascensis Elliot, Syn. N. Am. Mamm., F. C. M.

Pub., II, 1901, p. 372. Zoöl. Ser.

YAKUTAT BAY SHREW.

Type locality. Yakutat Bay, Alaska.

Geogr. Distr. Juneau to Yakutat, Alaska.

b. Sorex glacialis shumaginensis (Merriam).

Sorex alascensis shumaginensis Merr., Proc. Wash. Acad. Scien., II, 1900, p. 18.

Sorex glacialis shumaginensis Elliot, Syn. N. Am. Mamm., F. C. M.

Pub., II, 1901, p. 373. Zoöl. Ser.

POPOF ISLAND SHREW.

Type locality. Popof Island, Shumagin Group, Alaska.

Geogr. Distr. Shumagin Islands, Alaska.

1086. *Sorex bairdi* Merriam.

Sorex bairdi Merr., N. Am. Faun., No. 10, 1895, p. 77, pl. VII, figs. 3, 3a. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II,

1901, p. 373. Zoöl. Ser.

BAIRD'S SHREW.

Type locality. Astoria, mouth of the Columbia River, Clatsop County, Oregon.

Geogr. Distr. Coast of Oregon. Range unknown.

1087. *Sorex trowbridgii* Baird.

Sorex trowbridgii Baird, N. Am. Mamm., 1857, p. 13. Elliot, Syn.

N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 373. Zoöl. Ser.

TROWBRIDGE'S SHREW.

Type locality. Astoria, mouth of the Columbia River, Clatsop County, Oregon.

Geogr. Distr. Western Washington and Oregon, west of Cascade Range.

1088. *Sorex montereyensis* Merriam.

Sorex montereyensis Merr., N. Am. Faun., No. 10, 1895, p. 79.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 373, fig. 70. Zoöl. Ser.

MONTEREY SHREW.

Type locality. Monterey, Monterey County, California.

Geogr. Distr. Coast, and Sierra Nevada of California; south on coast to Morro and San Luis Obispo; south in the Sierra to Sequoia National Park and East Fork, Kaweah River.

1089. *Sorex californicus* Merriam.

Sorex californicus Merr., N. Am. Faun., No. 10, 1895, p. 80, pl. XII,

figs. 6, 7. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 374. Zoöl. Ser.

CALIFORNIA SHREW.

Type locality. Walnut Creek, Contra Costa County, California.

Geogr. Distr. California.

1090. *Sorex ornatus* Merriam.

Sorex ornatus Merr., N. Am. Faun., No. 10, 1895, p. 79, pl. VIII,

figs. 3, 3a. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 374. Zoöl. Ser.

ORNAMENTED SHREW.

Type locality. San Emigdio Cañon, Mount Pinos, Monterey County, California.

Geogr. Distr. Southern California from head of Ventura River and Mount Pinos, easterly to San Bernardino Peak, and south through the San Jacinto Range to Santa Isabel.

1091. *Sorex shastensis* Merriam.

Sorex shastensis Merr., N. Am. Faun., No. 16, 1899, p. 87. Elliot,

Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 374. Zoöl. Ser.

MOUNT SHASTA SHREW.

Type locality. Mount Shasta, Siskiyou County, California. Altitude, 5,700 feet.

Geogr. Distr. Known only from the type locality.

1092. *Sorex tenellus* Merriam.

Sorex tenellus Merr., N. Am. Faun., No. 10, 1895, p. 81, pl. XII, figs. 8, 9. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 374. Zool. Ser.

DELICATE SHREW.

Type locality. Alabama Hills, near Lone Pine, Owens Valley, Inyo County, California.

Geogr. Distr. Southeastern California.

a. *Sorex tenellus lyelli* Merriam.

Sorex tenellus lyelli Merr., Proc. Biol. Soc. Wash., xv, 1902, p. 75.

MOUNT LYELL SHREW.

Type locality. Mount Lyell, Tuolumne County, California.

Geogr. Distr. Known only from the type locality.

Genl. Char. Similar to *S. tenellus*, but browner above; 3d unicuspid larger than 4th; unicuspid series heavier; 1st and 2d broader.

Measurements. Total length, 103; tail vertebræ, 41; hind foot, 12.

b. *Sorex tenellus myops* Merriam.

Sorex tenellus myops Merr., Proc. Biol. Soc. Wash., xv, 1902, p. 76.

WHITE MOUNTAINS SHREW.

Type locality. White Mountains, Mono County, California.

Geogr. Distr. Southeastern California.

Genl. Char. Like *S. tenellus*, but smaller; ears larger; color paler. Skull shorter and broader; rostrum broadened behind unicuspid; molars deeply excavated posteriorly.

Measurements. Total length, 98; tail vertebræ, 41; hind foot, 12.

c. *Sorex tenellus nanus* Merriam.

Sorex tenellus nanus Merr., N. Am. Faun., No. 10, 1895, p. 81, pl. VIII, figs. 5, 5a. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 375. Zool. Ser.

DWARF SHREW.

Type locality. Estes Park, Larimer County, Colorado.

Geogr. Distr. Colorado; Montana.

1093. *Sorex longirostris* Bachman.

Sorex longirostris Bachm., Jour. Acad. Nat. Scien. Phil., 1837, p. 370, pl. XXIII, fig. 2. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 375. Zool. Ser.

LONG-NOSED SHREW.

Type locality. Swamps of the Santee River, South Carolina.

Geogr. Distr. North and South Carolina; probably also Georgia and Florida.

1094. *Sorex fisheri* Merriam.

Sorex fisheri Merr., N. Am. Faun., No. 10, 1895, p. 86, pl. iv, fig. 4. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 375. Zoöl. Ser.

FISHER'S SHREW.

Type locality. Lake Drummond, Dismal Swamp, Norfolk County, Virginia.

Geogr. Distr. Virginia and North Carolina.

1095. *Sorex pacificus* Coues.

Sorex pacificus Coues, Bull. U. S. Geol. & Geog. Surv., III, 1877, p. 650. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 375. Zoöl. Ser.

PACIFIC SHREW.

Type locality. Fort Umpqua, mouth of Umpqua River, Douglas County, Oregon.

Geogr. Distr. Pacific coast from Point Reyes, Marin County, California, to Yaquina Bay, Oregon.

1096. *Sorex pribilofensis* Merriam.

Sorex pribilofensis Merr., N. Am. Faun., No. 10, 1895, p. 87, pl. ix, figs. 3, 3a. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 376. Zoöl. Ser.

ST. PAUL ISLAND SHREW.

Type locality. St. Paul Island, Pribilof Islands, Bering Sea, Alaska.

Geogr. Distr. Pribilof Islands, Bering Sea, Alaska.

1097. *Sorex merriami* Dobson.

Sorex merriami Dobson, Mon. Insect., fasc. 1, Pt. III, 1890, pl. xxiii, fig. 6. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 376. Zoöl. Ser.

MERRIAM'S SHREW.

Type locality. Little Big Horn River, near Fort Custer, Crow Indian Reservation, Montana.

Geogr. Distr. Known only from the type locality.

a. *Sorex merriami miscix* (Bangs).

Sorex personatus miscix Bangs, N. Engl. Zoöl. Club, 1, 1899, p. 15.

Sorex merriami miscix Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 376. Zoöl. Ser.

LABRADOR SHREW.

Type locality. Black Bay, Strait of Belle Isle, Labrador.

Geogr. Distr. Labrador. Range unknown.

1098. *Sorex salvini* Merriam.

Sorex salvini Merr., Proc. Biol. Soc. Wash., XI, 1897, p. 229.

Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 551. Zoöl. Ser.

SALVIN'S SHREW.

Type locality. Calel, Totonicapan, Guatemala. Altitude, 10,200 feet.

Geogr. Distr. Guatemala. Range unknown.

1099. *Sorex stizodon* Merriam.

Sorex stizodon Merr., N. Am. Faun., No. 10, 1895, p. 98. Elliot,

Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 551. Zoöl. Ser.

SAN CRISTOBAL SHREW.

Type locality. San Cristobal, State of Chiapas, Mexico.

Geogr. Distr. State of Chiapas, Mexico. Range unknown.

1100. *Sorex veræpaci* Alston.

Sorex veræpaci Alston, Proc. Zoöl. Soc., 1877, p. 445. Elliot,

Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 551. Zoöl. Ser.

GUATEMALAN SHREW.

Type locality. Coban, Guatemala.

Geogr. Distr. Guatemala, Central America.

1101. *Sorex macrodon* Merriam.

Sorex macrodon Merr., N. Am. Faun., No. 10, 1895, p. 82. Elliot,

Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 552. Zoöl. Ser.

LARGE-TOOTHED SHREW.

Type locality. Orizaba, State of Vera Cruz, Mexico. Altitude, 4,200 feet.

Geogr. Distr. State of Vera Cruz, Mexico.

1102. *Sorex godmani* Merriam.

Sorex godmani Merr., Proc. Biol. Soc., Wash., XI, 1897, p. 229.

Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 552. Zoöl. Ser.

GODMAN'S SHREW.

Type locality. Volcano Santa Maria, Quezaltenango, Guatemala. Altitude, 9,000 feet.

Geogr. Distr. Guatemala, Central America. Range unknown.

1103. *Sorex sclateri* Merriam.

Sorex sclateri Merr., Proc. Biol. Soc. Wash., XI, 1897, p. 228.

Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 552. Zoöl. Ser.

SCLATER'S SHREW.

Type locality. Tumbala, State of Chiapas, Mexico. Altitude, 5,000 feet.

Geogr. Distr. State of Chiapas, Mexico.

1104. *Sorex saussurii* Merriam.

Sorex saussurei Merr., Proc. Biol. Soc. Wash., VII, 1892, p. 173.

Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 553. Zoöl. Ser.

SAUSSURE'S SHREW.

Type locality. North slope of the Sierra Nevada de Colima, State of Jalisco, Mexico. Altitude, 8,000 feet.

Geogr. Distr. State of Jalisco, Mexico.

a. *Sorex saussurii mutabilis* Merriam.

Sorex caudatus Merr., N. Am. Faun., No. 10, 1895, p. 84. (nec Hodgson, 1851.)

Sorex saussurei mutabilis Merr., Science, N. S., VIII, 1898, p. 782.

Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 553. Zoöl. Ser.

CHANGEABLE SHREW.

Type locality. Reyes, State of Oaxaca, Mexico. Altitude, 10,200 feet.

Geogr. Distr. From State of Vera Cruz to State of Oaxaca, Mexico.

1105. *Sorex orinus* Elliot.

Sorex orinus Elliot, Pub. Field Columb. Mus., III, 1903, p. 172.

Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 553, fig. 102. Zoöl. Ser.

MOUNTAIN SHREW.

Type locality. Aguaje de las Fresas, San Pedro Martir Mountains, Lower California, Mexico. Altitude, 6,000 feet.

Geogr. Distr. Known only from the type locality.

Genus 144. *Microsorex*. Small Shrews.

Microsorex Coues, Notes on Am. Insectiv. Mamm., Bull. U. S. Geol. and Geogr. Surv., III, 1877, No. 3, p. 646. Type *Sorex hoyi* Baird.

1106. *Microsorex hoyi* (Baird).

Sorex hoyi Baird, N. Am. Mamm., 1857, p. 32, pl. xxviii.

Sorex thompsoni Baird, N. Am. Mamm., 1857, p. 34.

Microsorex hoyi Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 377, fig. 71. Zoöl. Ser.

HOY'S SHREW.

Type locality. Racine, Racine County, Wisconsin.

Geogr. Distr. Labrador to British Columbia; northern Minnesota and North Dakota.

1107. *Microsorex eximius* (Preble).

Sorex eximius Preble, N. Am. Faun., No. 21, 1901, p. 71.

EXTRAORDINARY SHREW.

Type locality. Tyonek, Cook Inlet, Alaska.

Geogr. Distr. Known only from the type locality.

Genl. Char. Similar to *M. hoyi*, but larger and paler. Rostrum narrow, elongate; braincase high, compressed anteriorly, elevated above plane of rostrum; palate long, narrow, excavated; fourth unicuspid small; first upper incisor, long, slightly decurved.

Color. Head, back, and sides uniform pearly sepia; under parts pale drab; tail bicolor.

Measurements. Total length, 98; tail vertebræ, 31; hind foot, 11. Skull: basal length, 15; palatal length, 6.5; mastoid width, 7.1; antorbital breadth, 4.3.

1108. *Microsorex alnorum* (Preble).

Sorex alnorum Preble, N. Am. Faun., No. 22, 1902, p. 72.

KEEWATIN SHREW.

Type locality. Robinson's Portage, Keewatin, Canada.

Geogr. Distr. Keewatin, Canada.

Genl. Char. Larger than *M. hoyi*. Skull larger.

Color. Upper parts sepia brown; beneath ashy gray; hind foot dusky above, whitish below, and dusky toward tip.

Measurements. Total length, 98; tail vertebræ, 35; hind foot, 12. Skull: greatest length, 16.5; width of braincase, 7; length of mandible, 10.

Genus 145. **Neosorex. Swimming Shrews.**

Neosorex Baird, N. Am. Mamm., 1857, p. 11. Type *Neosorex navigator* Baird.

1109. *Neosorex palustris* (Richardson).

Sorex palustris Rich., Zoöl. Journ., 111, 1828, p. 517.

Neosorex palustris Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 378, fig. 72. Zoöl. Ser.

MARSH SHREW.

Type locality. Marshy places from Hudson Bay to the Rocky Mountains.

Geogr. Distr. Central Minnesota to east base of the Rocky Mountains.

a. *Neosorex palustris navigator* (Baird).

Neosorex navigator Baird, N. Am. Mamm., 1857, p. 11.

Sorex palustris Merr., N. Am. Faun., No. 5, 1891, p. 35. (nec Richardson.)

Neosorex palustris navigator Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 379. Zool. Ser.

NAVIGATOR SHREW.

Type locality. Northern Idaho?

Geogr. Distr. Rocky Mountains from British Columbia to southern Colorado; Sierra Nevada of California to Sequoia National Park.

b. *Neosorex palustris alaskanus* (Merriam).

Sorex navigator alaskanus Merr., Proc. Wash. Acad. Scien., 11, 1900, p. 18.

Neosorex palustris alaskanus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 379. Zool. Ser.

ALASKA SWIMMING SHREW.

Type locality. Point Gustavus, east side of Glacier Bay, Alaska.

Geogr. Distr. Alaska. Range unknown.

1110. *Neosorex albibarbis* Cope.

Neosorex albibarbis Cope, Proc. Acad. Nat. Scien. Phil., 1862, p. 188.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 379. Zool. Ser.

WHITE-LIPPED SWIMMING SHREW.

Type locality. Profile Lake, Franconia Mountains, Grafton County, New Hampshire.

Geogr. Distr. Pennsylvania northward to Labrador.

1111. *Neosorex hydrodromus* (Dobson).

Sorex hydrodromus Dobson, Ann. Mag. Nat. Hist., 6th Ser., iv, 1889, p. 373.

Neosorex hydrodromus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 379. Zool. Ser.

DOBSON'S SWIMMING SHREW.

Type locality. Unalaska Island, Alaska.

Geogr. Distr. Aleutian Islands, Alaska. Range unknown.

Genus 146. **Atophrax. Anomalous Shrews.**

Atophrax Merr., Trans. Linn. Soc. N. Y., 11, 1884, p. 217. Type
Atophrax bendirii Merriam.

1112. Atophrax bendirii Merriam.

Atophrax bendirii Merr., Trans. Linn. Soc. N. Y., 11, 1884, p. 217.
Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 381,
fig. 73. Zoöl. Ser. *Id.* Pub. Field Columb. Mus., 111, 1903,
p. 195. Zoöl. Ser.

Sorex bendirii palmeri Merr., N. Am. Faun., No. 10, 1895, p. 97.

Atophrax bendirii palmeri Elliot, Syn. N. Am. Mamm., F. C. M.
Pub., 11, 1901, p. 381. Zoöl. Ser.

BENDIRE'S SHREW.

Type locality. Near Williamson River, eighteen miles southwest
of Fort Klamath, Klamath County, Oregon.

Geogr. Distr. From Sonoma County, California, and from east
side of Cascade Range to the Pacific Ocean; north to Puget Sound,
Washington.

a. Atophrax bendirii albiventer (Merriam).

Sorex bendirii albiventer Merr., N. Am. Faun., No. 10, 1895, p. 97.

Atophrax bendirii albiventer Elliot, Syn. N. Am. Mamm., F. C. M.
Pub., 11, 1901, p. 381. Zoöl. Ser.

WHITE-BELLIED SHREW.

Type locality. Lake Cushman, Olympic Mountains, Mason
County, Washington.

Geogr. Distr. Olympic Mountains, Washington.

Genus 147. **Notiosorex. Southern Shrews.**

Notiosorex Coues, Notes Insectiv. Mamm., Bull. U. S. Geol. & Geogr.
Surv. Terr., 111, 1877, p. 646. Type *Sorex crawfordi* Coues.

1113. Notiosorex crawfordi (Coues).

Sorex crawfordi Coues, Bull. U. S. Geol. & Geogr. Surv. Terr., 111,
1877, p. 651.

Notiosorex crawfordi Elliot, Syn. N. Am. Mamm., F. C. M. Pub.,
11, 1901, p. 385, fig. 76. Zoöl. Ser. *Id.* Mamm. Middle Amer.
& W. Indies, F. C. M. Pub., IV, Pt. 11, 1904, p. 554. Zoöl. Ser.

CRAWFORD'S SHREW.

Type locality. Near Old Fort Bliss, two miles above El Paso,
El Paso County, Texas.

Geogr. Distr. Eastern Texas to southern California, and on northern boundary line of Mexico at Monument 66.

a. Notiosorex crawfordi evotis (Coues).

Sorex evotis Coues, Bull. U. S. Geol. & Geogr. Surv. Terr., III, 1877, p. 652.

Notiosorex crawfordi evotis Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 555. Zoöl. Ser.

MAZATLAN SHREW.

Type locality. Mazatlan, State of Sinaloa, Mexico.

Geogr. Distr. State of Sinaloa, Mexico. Range unknown.

1114. Notiosorex gigas Merriam.

Notiosorex gigas Merr., Proc. Biol. Soc. Wash., XI, 1897, p. 227.

Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 555, fig. 103. Zoöl. Ser.

GIANT SHREW.

Type locality. Mountains of Milpillas near San Sebastian, State of Jalisco, Mexico.

Geogr. Distr. Mountains of Milpillas, State of Jalisco, Mexico. Range unknown.

Genus 148. **Blarina. Short-tailed Shrews.**

Blarina Gray, Proc. Zoöl. Soc., 1837, p. 124. Type *Sorex talpoides* Gapper = *Sorex brevicaudus* Say.

Brachysorex Duvern., Mag. Zool., 2me Sér., IV, 1842, p. 37, pl. 52.

Blaria Gray, List Spec. Mamm. Brit. Mus., 1843, p. XXI.

Talposorex! Pomel, Archiv. Scien. Phys. & Nat. Bibl. Univ. Genève, IX, 1848, p. 248. (nec Schinz, 1821.)

Cryptotis Pomel, Archiv. Scien. Phys. & Nat. Bibl. Univ. Genève, IX, 1848, p. 249.

Galemys Pomel, Archiv. Scien. Phys. & Nat. Bibl. Univ. Genève, IX, 1848, p. 249. (nec Kaup.)

Anotus Wagn., Schreb. Säugth., Suppl., 1855, p. 550. (nec Rafin., 1815.)

Soriciscus Coues, Bull. U. S. Geol. & Geogr. Surv. Terr., 1877, p. 649.

Subgenus A. **Blarina.**

1115. Blarina brevicauda (Say).

Sorex brevicaudus Say, Long's Exped. Rocky Mts., I, 1823, p. 164.

Sorex talpoides Gapper, Zoöl. Journ., V, 1830, p. 202, pl. VIII.

Sorex dekayi Bachm., Journ. Acad. Nat. Scien. Phil., 1837, p. 377.

Galemys (Brachysorex) micrurus Pomel, Archiv. Scien. Phys. & Nat. Bibl. Univ. Genève, ix, 1848, p. 249.

Sorex angusticeps Baird, N. Am. Mamm., 1857, p. 47.

Blarina costaricensis Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1891, p. 205. (Not from Costa Rica.)

Blarina brevicauda Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 382, fig. 74. Zoöl. Ser.

Blarina brevicauda talpoides Bangs, Proc. N. Engl. Zoöl. Club, III, 1902, p. 75.

SHORT-TAILED SHREW.

Type locality. Near Blair, Missouri River, Washington County, Nebraska.

Geogr. Distr. Atlantic coast west to Manitoba, Canada, and Nebraska.

a. *Blarina brevicauda carolinensis* (Bachman).

Sorex carolinensis Bachm., Journ. Acad. Nat. Scien. Phil., 1837, p. 366.

Blarina brevicauda carolinensis Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 383. Zoöl. Ser.

CAROLINA SHREW.

Type locality. Eastern South Carolina.

Geogr. Distr. Chesapeake Bay, Maryland, to Arkansas.

b. *Blarina brevicauda *hylophaga* Elliot.

Blarina brevicauda hulo-phaga Elliot, Pub. Field Columb. Mus., I, 1899, p. 287. Zoöl. Ser. *Id.* Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 383. Zoöl. Ser.

WOOD SHREW.

Type locality. Dougherty, Washita River, Chickasaw Nation, Indian Territory.

Geogr. Distr. Known only from the type locality.

c. *Blarina brevicauda peninsulæ* (Merriam).

Blarina carolinensis peninsulæ Merr., N. Am. Faun., No. 10, 1895, p. 14.

Blarina brevicauda peninsulæ Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 383. Zoöl. Ser.

PENINSULA SHREW.

Type locality. Miami River, Dade County, Florida.

Geogr. Distr. Peninsula of Florida, south of latitude 28°.

*Misprinted originally *hulophaga*.

d. *Blarina brevicauda aloga* Bangs.

Blarina brevicauda aloga Bangs, Proc. N. Engl. Zool. Club, III, 1902, p. 76.

IRRATIONAL SHREW.

Type locality. West Tisbury, Martha's Vineyard, Massachusetts.

Geogr. Distr. Known only from the type locality.

Genl. Char. Rostrum very short, massive, and wide; nasal aperture wide.

Color. Above dark brownish drab; beneath silvery gray; hands and feet white; tail above dusky, beneath gray.

Measurements. Total length, adult males, 115-128; tail vertebræ, 24-29; hind foot, 14-15. Skull: greatest length, 22; basal length, 19.6; interorbital breadth, 6; length of tooth row, 10; length of single half of mandible, 12.2.

e. *Blarina brevicauda compacta* Bangs.

Blarina brevicauda compacta Bangs, Proc. N. Engl. Zool. Club, III, 1902, p. 77.

THICK-SET SHREW.

Type locality. Nantucket, Nantucket Island, Massachusetts.

Geogr. Distr. Nantucket Island, Massachusetts.

Genl. Char. Like *B. b. aloga* in size; color different.

Color. Above and beneath slate gray shaded with drab; hands and feet grayish; tail dusky, unicolor.

Measurements. Total length, adult males, 119-125; tail vertebræ, 23-25; hind foot, 14.5-15. Skull: greatest length, 22.4; basal length, 19.8; interorbital breadth, 6; length of tooth row, 10; length of single half of mandible, 12.2.

f. *Blarina brevicauda berlandieri* (Baird).

Blarina berlandieri Baird, N. Am. Mamm., 1857, p. 20.

Blarina brevicauda berlandieri Elliot, Mamm. Middle Amer. & W.

Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 558. Zool. Ser.

BERLANDIER'S SHREW.

Type locality. Matamoros, State of Tamaulipas, Mexico.

Geogr. Distr. Range unknown. State of Tamaulipas, Mexico, possibly north into Texas.

1116. *Blarina telmalestes* Merriam.

Blarina telmalestes Merr., N. Am. Faun., No. 10, 1895, p. 15, pl. 11, fig. 5. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 384. Zool. Ser.

DISMAL SWAMP SHREW.

Type locality. Lake Drummond, Dismal Swamp, Norfolk County, Virginia.

Geogr. Distr. Dismal Swamp, Virginia.

Subgenus B. *Cryptotis*.1117. *Blarina parva* (Say).

Sorex parvus Say, Long's Exped. Rocky Mts., I, 1823, p. 163.

Sorex cinereus Bachm., Journ. Acad. Nat. Scien. Phil., 1837, p. 373, pl. XXIII, fig. 3.

Blarina exilipes Baird, N. Am. Mamm., 1857, p. 51.

Blarina eximius Baird, N. Am. Mamm., 1857, p. 52.

Blarina parva Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 384, fig. 75. Zoöl. Ser.

SMALL SHREW.

Type locality. Near Blair, Missouri River, Washington County, Nebraska.

Geogr. Distr. Atlantic coast west to eastern Nebraska and Texas.

1118. *Blarina floridana* Merriam.

Blarina floridana Merr., N. Am. Faun., No. 10, 1895, p. 19, pl. 1, fig. 7. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 385. Zoöl. Ser.

FLORIDA SHREW.

Type locality. Chester Shoal, eleven miles north of Cape Canaveral, Brevard County, Florida.

Geogr. Distr. Peninsula of Florida, south of latitude 29°.

1119. *Blarina pergracilis* Elliot.

Blarina pergracilis Elliot, Pub. Field Columb. Mus., III, 1903, p. 149. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 557, figs. 104, LXXXII. Zoöl. Ser.

OCOTLAN SLENDER SHREW.

Type locality. Ocotlan, State of Jalisco, Mexico.

Geogr. Distr. Known only from the type locality.

1120. *Blarina nigrescens* Allen.

Blarina nigrescens Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1895, p. 339. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 558. Zoöl. Ser.

BLACKISH SHREW.

Type locality. San Isidro (San José), Costa Rica.

Geogr. Distr. Known only from the type locality.

1121. *Blarina tropicalis* Merriam.

Blarina tropicalis Merr., N. Am. Faun., No. 10, 1895, p. 21.

Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV,

Pt. II, 1904, p. 359. Zoöl. Ser.

TROPICAL SHREW.

Type locality. Coban, Guatemala. Altitude, 4,400 feet.

Geogr. Distr. States of Vera Cruz and Oaxaca, Mexico, into Guatemala.

1122. *Blarina soricina* Merriam.

Blarina soricina Merr., N. Am. Faun., No. 10, 1895, p. 22. Elliot,

Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II,

1904, p. 560. Zoöl. Ser.

TALPAM SHREW.

Type locality. Talpam, Valley of Mexico, ten miles south of Mexico City. Altitude, 7,600 feet.

Geogr. Distr. State of Mexico. Range unknown.

1123. *Blarina oreophila* Allen.

Blarina oreophila Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1895, p. 340.

Blarina oreophila Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., IV, Pt. II, 1904, p. 559. Zoöl. Ser.

VOLCANO OF IRAZÚ SHREW.

Type locality. Volcan de Irazú, Costa Rica, Central America.

Geogr. Distr. Known only from the type locality.

1124. *Blarina obscura* Merriam.

Blarina obscura Merr., N. Am. Faun., No. 10, 1895, p. 23. Elliot,

Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II,

1904, p. 559. Zoöl. Ser.

DUSKY SHREW.

Type locality. Tulancingo, State of Hidalgo, Mexico. Altitude, 8,500 feet.

Geogr. Distr. State of Hidalgo, Mexico. Range unknown.

1125. *Blarina mexicana* Coues.

Blarina mexicana Coues, Bull. U. S. Geol. & Geogr. Surv. Terr., III, 1877, p. 652. Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., IV, Pt. II, 1904, p. 560. Zoöl. Ser.

COUES' MEXICAN SHREW.

Type locality. Jalapa, State of Vera Cruz, Mexico.

Geogr. Distr. States of Vera Cruz and Oaxaca.

a. *Blarina mexicana goldmani* Merriam.

Blarina mexicana goldmani Merr., N. Am. Faun., No. 10, 1895,
p. 25. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M.
Pub., IV, Pt. II, 1904, p. 560. Zool. Ser.

GOLDMAN'S SHREW.

Type locality. Mountains near Chilpancingo, State of Guerrero,
Mexico. Altitude, 10,000 feet.

Geogr. Distr. State of Guerrero, Mexico. Range unknown.

b. *Blarina mexicana peregrina* Merriam.

Blarina mexicana peregrina Merr., N. Am. Faun., No. 10, 1895,
p. 24. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M.
Pub., IV, Pt. II, 1904, p. 560. Zool. Ser.

WANDERING SHREW.

Type locality. Mountains fifteen miles west of Oaxaca, State of
Oaxaca, Mexico. Altitude, 9,500 feet.

Geogr. Distr. State of Oaxaca, Mexico. Range unknown.

c. *Blarina mexicana machetes* Merriam.

Blarina mexicana machetes Merr., N. Am. Faun., No. 10, 1895,
p. 26. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M.
Pub., IV, Pt. II, 1904, p. 561. Zool. Ser.

WARRING SHREW.

Type locality. Mountains near Ozolotepec, State of Oaxaca,
Mexico. Altitude, 10,000 feet.

Geogr. Distr. State of Oaxaca, Mexico. Range unknown.

1126. *Blarina mayensis* Merriam.

Blarina mayensis Merr., Proc. Wash. Acad. Scien., III, 1901, p. 559.
Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV,
Pt. II, 1904, p. 561. Zool. Ser.

MAYA SHREW.

Type locality. Maya ruin, Chichen Itza, Yucatan, Mexico.

Geogr. Distr. Yucatan, Mexico. Range unknown.

1127. *Blarina nelsoni* Merriam.

Blarina nelsoni Merr., N. Am. Faun., No. 10, 1895, p. 26. Elliot,
Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II,
1904, p. 561. Zool. Ser.

NELSON'S SHREW.

Type locality. Volcano of Tuxtla, State of Vera Cruz, Mexico.
Altitude, 4,800 feet.

Geogr. Distr. State of Vera Cruz, Mexico.

1128. *Blarina alticola* Merriam.

Blarina alticola Merr., N. Am. Faun., No. 10, 1895, p. 27.
 Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV,
 Pt. II, 1904, p. 561. Zoöl. Ser.

MT. POPOCATEPETL SHREW.

Type locality. Mt. Popocatepetl, State of Mexico. Altitude,
 11,500 feet.

Geogr. Distr. State of Mexico, Mexico.

1129. *Blarina fossor* Merriam.

Blarina fossor Merr., N. Am. Faun., No. 10, 1895, p. 28. Elliot,
 Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II,
 1904, p. 562. Zoöl. Ser.

MT. ZEMPOALTEPEC SHREW.

Type locality. Mt. Zempoaltepec, State of Oaxaca, Mexico.
 Altitude, 10,500 feet.

Geogr. Distr. Mt. Zempoaltepec, 8,200-10,500 feet altitude, State
 of Oaxaca, Mexico.

1130. *Blarina magna* Merriam.

Blarina magna Merr., N. Am. Faun., No. 10, 1895, p. 28. Elliot,
 Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II,
 1904, p. 562. Zoöl. Ser.

GREAT SHREW.

Type locality. Totontepec, State of Oaxaca, Mexico. Altitude,
 6,800 feet.

Geogr. Distr. Mountains around Totontepec and Zempoaltepec,
 State of Oaxaca, Mexico. Altitude, 6,800-8,000 feet.

Family II. Talpidæ. Moles.**Subfamily I. Myogalinæ.****Genus 149. *Neurotrichus*. Shrew-like Moles.**

Neurotrichus Günther, Proc. Zoöl. Soc., 1880, p. 441. Type *Urotrichus gibbsii* Baird.

1131. *Neurotrichus gibbsii* (Baird).

Urotrichus gibbsii Baird, N. Am. Mamm., 1857, p. 76.

Neurotrichus gibbsii Elliot, Syn. N. Am. Mamm., F. C. M. Pub.,
 II, 1901, p. 387, fig. 77. Zoöl. Ser.

GIBBS' SHREW MOLE.

Type locality. White River Pass, Cascade Mountains, Pierce
 County, Washington.

Geogr. Distr. Fraser River Region, British Columbia, west of Cascade and Sierra Nevada Mountains to Shasta County, and thence along the coast of San Francisco Bay, California.

a. Neurotrichus gibbsi major Merriam.

Neurotrichus gibbsi major Merr., N. Am. Faun., No. 16, 1899, p. 88. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 387. Zoöl. Ser.

LARGE SHREW MOLE.

Type locality. Carberry Ranch, Shasta County, California.

Geogr. Distr. Canadian zone, Shasta County, California.

b. Neurotrichus gibbsi hyacinthinus Bangs.

Neurotrichus gibbsi hyacinthinus Bangs, Amer. Nat., xxxi, 1897, p. 240. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 388. Zoöl. Ser.

HYACINTHINE SHREW MOLE.

Type locality. Nicasio, Marin County, California.

Geogr. Distr. California. Range unknown.

Subfamily II. **Talpinæ.**

Genus 150. **Condylura. Star-nosed Moles.**

Condylura Illig., Prodr. Syst. Mamm. et Av., 1811, p. 125. Type *Sorex cristatus* Linnæus.

Rhinaster Wagl., Nat. Syst. Amphib., 1830, p. 14.

Astromyctes Gray, List Mamm. Brit. Mus., 1843, pp. xxi, 76. (*nomen nudum.*)

Astromydes Blyth, Cat. Mamm. Asiat. Soc. Mus., 1863, p. 87.

1132. Condylura cristata (Linnæus).

Sorex cristatus Linn., Syst. Nat., I, 1758, p. 53.

Talpa longicaudata Erxl., Syst. Reg. Anim., I, 1777, p. 118.

Talpa radiata Shaw, Gen. Zoöl., Mamm., I, 1800, p. 523.

Condylura macroura Harl., Faun. Amer., 1825, p. 36.

Condylura prasinata Harris, Bost. Journ. Phil. & Arts, 1825, p. 582.

Condylura cristata Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 395, fig. 81. Zoöl. Ser.

STAR-NOSED MOLE.

Type locality. Pennsylvania.

Geogr. Distr. Hudson Bay to Manitoba, south to Minnesota on the west, and on the Atlantic coast to South Carolina in the Alleghanies.

Genus 151. **Scapanus. Western Moles.**

Scapanus Pomel, Archiv. Scien. Phys. Nat. Bibl. Univ. Genève, 4th Ser., ix, 1848, p. 247. Type by elimination, *Scalops townsendii* Bachman.

1133. Scapanus townsendi (Bachman).

Scalops canadensis Rich., Faun. Bor. Am., i, 1829, p. 9. (nec Desmarest, nec Harlan.)

Scalops townsendii Bach., Journ. Acad. Nat. Scien. Phil., 1839, p. 58.

Scalops latimanus Bach., Bost. Journ. Nat. Hist., 1843, p. 41.

Scalops metallescens Cass., Proc. Acad. Nat. Scien. Phil., 1853, p. 242.

Scalops æneus Cass., Proc. Acad. Nat. Scien. Phil., 1853, p. 299.

Talpa tæniata Le Conte, Proc. Acad. Nat. Scien. Phil., 1853, p. 327.

Scapanus townsendi Elliot, Syn. N. Am. Mamm., F. C. M. Pub., ii, 1901, p. 391, fig. 79. Zoöl. Ser.

TOWNSEND'S MOLE.

Type locality. Vicinity of Vancouver, Clarke County, Washington.

Geogr. Distr. Coast line of Oregon and Washington, and north-western California to Crescent City.

1134. Scapanus orarius True.

Scapanus orarius True, Proc. U. S. Nat. Mus., xix, 1896, p. 52.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., ii, 1901, p. 392. Zoöl. Ser.

COAST MOLE.

Type locality. Shoalwater Bay, Pacific County, Washington.

Geogr. Distr. Sea coast of Washington and Oregon from Coquilla River northward, and shores of Puget Sound to Sumas, British Columbia; Fort Walla Walla, Walla Walla County, Washington.

1135. Scapanus californicus (Ayres).

Scalops californicus Ayres, Proc. Calif. Acad. Nat. Scien., i, 1855, p. 54.

Scapanus dilatatus True, Proc. U. S. Nat. Mus., 1894, p. 242.

Scapanus californicus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., ii, 1901, p. 392. Zoöl. Ser.

CALIFORNIA MOLE.

Type locality. San Francisco, California.

Geogr. Distr. California west of the Coast Range and Sierra Nevada; Fort Klamath, Klamath County, Oregon.

a. Scapanus californicus minusculus Bangs.

Scapanus californicus minusculus Bangs, Proc. N. Engl. Zool. Club, 1, 1899, p. 70. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, Append., p. 435. Zool. Ser.

SMALLER CALIFORNIA MOLE.

Type locality. Fyffe, El Dorado County, California.

Geogr. Distr. California. Range unknown.

1136. *Scapanus alpinus* Merriam.

Scapanus alpinus Merr., Proc. Biol. Soc. Wash., 11, 1897, p. 102.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 393. Zool. Ser.

ALPINE MOLE.

Type locality. Crater Lake, Mount Mazama, Klamath County, Oregon. Altitude, 7,000 feet.

Geogr. Distr. Oregon. Range unknown.

1137. *Scapanus truei* Merriam.

Scapanus truei Merr., Proc. Biol. Soc. Wash., 11, 1897, p. 102.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 393. Zool. Ser.

TRUE'S MOLE.

Type locality. Lake City, Modoc County, California.

Geogr. Distr. California. Range unknown.

1138. *Scapanus anthonyi* Allen.

Scapanus anthonyi Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1893, p. 200. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901,

p. 393. Zool. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 11, 1904, p. 564, figs. 105, LXXXIII. Zool. Ser.

LOWER CALIFORNIA MOLE.

Type locality. San Pedro Martir Mountains, Lower California, Mexico. Altitude, 7,000 feet.

Geogr. Distr. Known only from the type locality.

Genus 152. *Scalopus*. Common Moles.

Scalopus Geoffroy, Cat. Mamm. Mus. Hist. Nat. Paris, 1803, pp. 77-78.

Type *Scalopus virginianus* Geoffroy = *Sorex aquaticus* Linnæus.

Scalops Illig., Prodr. Syst. Mamm. et Avium, 1811, p. 126.

1139. *Scalopus aquaticus* (Linnæus).

Sorex aquaticus Linn., Syst. Nat., 1, 1758, p. 53.

Talpa europæa flavescens Erxl., Syst. Reg. Anim., 1777, p. 118.

- Talpa europæa flava* Gmel., Syst. Nat., 1, 1788, p. 112.
Talpa flava Kerr, Anim. Kingd., 1792, p. 201.
Talpa fusca Kerr, Anim. Kingd., 1792, p. 202.
Talpa cupreata Rafin., Préc. découv. trav. somiol. Palerme, 1814
 p. 14.
Scalops canadensis Desm., Mamm., 1820, p. 155.
Scalops pennsylvanica Harl., Faun. Amer., 1825, p. 33.
Talpa pennantii Le Conte, Proc. Acad. Nat. Scien. Phil., 1853, p. 327.
Scalops aquaticus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11,
 1901, p. 389, fig. 78. Zoöl. Ser.

COMMON MOLE.

- Type locality.* Eastern United States.
Geogr. Distr. Massachusetts to Florida, west to the Mississippi.

a. *Scalopus aquaticus machrinus* (Rafinesque).

- Talpa machrina* Rafin., Atlantic Journ., 1, 1832, p. 61.
Talpa sericea Rafin., Atl. Jour., 1832, p. 61.
Scalops argentatus Aud. & Bachm., Journ. Acad. Nat. Scien. Phil.,
 1842, p. 292.
Scalops aquaticus machrinus Elliot, Syn. N. Am. Mamm., F. C. M.
 Pub., 11, 1901, p. 389. Zoöl. Ser.

LARGE-NOSED MOLE.

- Type locality.* Near Lexington, Fayette County, Kentucky.
Geogr. Distr. Wisconsin and Minnesota to Tennessee and Missouri; west to eastern Kansas, Nebraska, and southwestern South Dakota.

b. *Scalopus aquaticus intermedius* (Elliot).

- Scalops machrinus intermedius* Elliot, Pub. Field Columb. Mus.,
 1, 1899, p. 280. Zoöl. Ser.
Scalops aquaticus intermedius Elliot, Syn. N. Am. Mamm., F. C. M.
 Pub., 11, 1901, p. 390. Zoöl. Ser.

INTERMEDIATE MOLE.

- Type locality.* Alva, Woods County, Oklahoma Territory.
Geogr. Distr. Oklahoma and Indian Territories.

c. *Scalopus aquaticus australis* (Chapman).

- Scalops aquaticus australis* Chapman, Bull. Am. Mus. Nat. Hist.
 N. Y., 1893, p. 339. Elliot, Pub. Field Columb. Mus., 111,
 1901, p. 56. Zoöl. Ser.
Scalops parvus Rhoads, Proc. Acad. Nat. Scien. Phil., 1894, p. 157.
Scalops anastasiæ Bangs, Proc. Bost. Soc. Nat. Hist., 1898, p. 212.
 Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 391.
 Zoöl. Ser.

Scalops aquaticus parvus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 390. Zoöl. Ser.

FLORIDA MOLE.

Type locality. Gainesville, Alachua County, Florida.

Geogr. Distr. East Florida, south to Lake Worth.

d. *Scalopus aquaticus texensis* (Allen).

Scalops argentatus texanus! Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1891, p. 221.

Scalops aquaticus texensis Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 390. Zoöl. Ser.

TEXAS MOLE.

Type locality. Texas (Aransas County?).

Geogr. Distr. Coast of Texas.

1140. **Scalopus aereus* (Bangs).

Scalops texanus! aereus Bangs, Proc. Biol. Soc. Wash., x, 1896, p. 138.

Scalops aereus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 390. Zoöl. Ser.

COPPER-COLORED MOLE.

Type locality. Stilwell, Boston Mountains, Indian Territory.

Geogr. Distr. Known only from the type locality.

Genus 153. *Parascalops*. Hairy-tailed Moles.

Parascalops True, Proc. U. S. Nat. Mus., xvii, 1894, p. 242. Type *Scalops breweri* Bachman.

1141. *Parascalops breweri* (Bachman).

Scalops breweri Bachm., Bost. Journ. Nat. Hist., 1841, p. 32.

Parascalops breweri Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 394, fig. 80. Zoöl. Ser.

BREWER'S HAIRY-TAILED MOLE.

Type locality. Martha's Vineyard, Massachusetts?

Geogr. Distr. Atlantic coast from New Brunswick to North Carolina, West Virginia, and southeastern Ohio. Shores of Lakes Erie and Ontario, and banks of the St. Lawrence to Quebec, possibly to the Atlantic.

* Doubtfully distinct from *S. a. texensis*.

Family III. **Solenodontidae.**Genus 154. **Solenodon. Almiqui and Agouta.****Solenodon** Brandt, Mém. Acad. Imp. Scien. St. Pétersb., 6th Ser., 11, 1833, p. 459. Type *Solenodon paradoxus* Brandt.**1142. Solenodon cubanus** Peters.*Solenodon cubanus* Peters, Abh. K. Preuss. Akad. Wiss. Berl., 1863, p. 2, pl. III. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 567, figs. 106, LXXXIV. Zool. Ser.**ALMIQUI.***Type locality.* Island of Cuba.*Geogr. Distr.* Island of Cuba. West Indies.**1143. Solenodon paradoxus** Brandt.*Solenodon paradoxus* Brandt, Mém. Acad. Imp. Scien. St. Pétersb., 6th Ser., 11, 1833, p. 459, pls. 1, 2. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 568. Zool. Ser.**AGOUTA.***Type locality.* Island of Haiti.*Geogr. Distr.* Island of Haiti.

Order X. **Chiroptera. Bats.**

Family I. **Vespertilionidæ. Common Bats.**

Subfamily I. **Vespertilioninæ.**

Genus 155. **Myotis.**

Myotis Kaup, Skizzirte Entw.-Gesch. u. Natürl. Syst. Europ. Thierw., 1, 1829, p. 106. Type *Vespertilio murinus* Schreber = *Vespertilio myotis* Bechstein.

Nystactes Kaup, Skizzirte Entw.-Gesch. u. Natürl. Syst. Europ. Thierw., 1, 1829, pp. 106, 108-109.

Vespertilio Keyserl. & Blas., Wieg. Archiv f. Naturg., 1839, p. 306. (nec Linn.)

Selysius Bonap., Iconog. Faun. Italy, 1, 1841, Intr. p. 3.

Brachyotus Kolen., Allgem. Deutsch. Naturh. Zeit. Dresden, neue Folge, 11, 1856, pp. 131, 174-177.

Isotus Kolen., Allgem. Deutsch. Naturh. Zeit. Dresden, neue Folge, 11, 1856, pp. 131, 177-179.

Aeorestes Fitzing., Sitzungs. Math.-Nat. Cl. K. Akad. Wiss. Wien, LXII, 1870, abth. 1, p. 427.

Comastes Fitzing., Sitzungs. Math.-Nat. Cl. K. Akad. Wiss. Wien, LXII, 1870, abth. 1, p. 565.

1144. **Myotis velifer** (J. A. Allen).

Vespertilio velifer J. A. Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1890, p. 177.

Myotis velifer Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 401. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 11, 1904, p. 573. Zoöl. Ser.

LARGE-WINGED BAT.

Type locality. Santa Cruz del Valle, near Guadalajara, State of Jalisco, Mexico.

Geogr. Distr. From the State of Mexico north throughout Mexico to Missouri.

1145. **Myotis incautus** (J. A. Allen).

Vespertilio incautus J. A. Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1896, p. 239.

HEEDLESS BAT.

Type locality. San Antonio, Bexar County, Texas.

Geogr. Distr. Texas. Range unknown.

Genl. Char. Size large; color without reddish brown tint.

Color. Above dull hair brown shaded with olive; beneath grayish tinged with buff, base of fur darker, ears and membranes blackish brown, posterior edge of wing and membrane lighter.

Measurements. Forearm, 42; first finger to end of claw, 7; second metacarpal, 40; fourth metacarpal, 39; fifth metacarpal, 39; tibia, 21; foot, 15; ear, 13; tragus, 8. Skull: length, 17; greatest width of braincase, 9.

1146. *Myotis evotis* (H. Allen).

Vespertilio evotis H. Allen, Mon. N. Am. Bats, 1864, p. 48.

Myotis evotis Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 406. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., 14, Pt. 11, 1904, p. 574. Zoöl. Ser.

PROMINENT-EARED BAT.

Type locality. Unknown; possibly Monterey, Monterey County, California.

Geogr. Distr. State of Vera Cruz, Mexico, north and west to the Pacific coast of the United States.

1147. *Myotis peninsularis* Miller.

Myotis peninsularis Miller, Ann. Mag. Nat. Hist., 7th Ser., 11, 1898, p. 124. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., 14, Pt. 11, 1904, p. 573. Zoöl. Ser.

LOWER CALIFORNIA BAT.

Type locality. San José del Cabo, Lower California, Mexico.

Geogr. Distr. Cape Region of Lower California, Mexico.

1148. *Myotis vivesi* Menegaux.

Myotis vivesi Meneg., Bull. Mus. Hist. Nat. Paris, 1901, p. 323. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., 14, Pt. 11, 1904, p. 574. Zoöl. Ser.

CARDONAL ISLAND BAT.

Type locality. Cardonal Island, Archipelago of Salsi Pudes, off east coast of Lower California, Mexico.

Geogr. Distr. Archipelago of Salsi Pudes, Lower California, Mexico.

1149. *Myotis milleri* Elliot.

Myotis milleri Elliot, Pub. Field Columb. Mus., 111, 1903, p. 172. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., 14, Pt. 11, 1904, p. 575. Zoöl. Ser.

MILLER'S BAT.

Type locality. La Grulla, San Pedro Martir Mountains, Lower California, Mexico. Altitude, 8,000 feet.

Geogr. Distr. San Pedro Martir Mountains, Lower California, Mexico.

1150. *Myotis nigricans* (Maximilian).

Vespertilio nigricans Max., Beitr. Natur. Bras., II, 1826, p. 266.

Vespertilio concinnus H. Allen, Proc. Acad. Nat. Scien. Phil., 1866, p. 280.

Myotis nigricans Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 575, figs. 107, LXXXV. Zoöl. Ser.

GRIZZLED BROWN BAT.

Type locality. Fazenda de Aga, near the Iritiba River, south-eastern Brazil.

Geogr. Distr. Tres Marias Islands, State of Jalisco, Mexico, south to Colombia and Brazil; West Indies.

1151. *Myotis chiriquensis* J. A. Allen.

Myotis chiriquensis J. A. Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1904, p. 77. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 576. Zoöl. Ser.

CHIRIQUEI BAT.

Type locality. Boqueron, Chiriqui, Panama.

Geogr. Distr. Panama, Central America.

1152. *Myotis dominicensis* Miller.

Myotis dominicensis Miller, Proc. Biol. Soc. Wash., xv, 1902, p. 243. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 576. Zoöl. Ser.

DOMINICAN COMMON BAT.

Type locality. Island of Dominica, West Indies.

Geogr. Distr. Known only from the type locality.

1153. *Myotis yumanensis* (H. Allen).

Vespertilio yumanensis H. Allen, Mon. N. Am. Bats, 1864, p. 58.

Myotis yumanensis Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 403. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 576. Zoöl. Ser.

FORT YUMA BAT.

Type locality. Old Fort Yuma, San Diego County, California.

Geogr. Distr. State of San Luis Potosi, Mexico, into southwestern United States.

*a. **Myotis yumanensis saturatus** Miller.*

Myotis yumanensis saturatus Miller, N. Am. Faun., No. 13, 1897, p. 68. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 403. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 577. Zoöl. Ser.

DARK YUMA BAT.

Type locality. Hamilton, Skagit County, Washington.

Geogr. Distr. State of Michoacan, Mexico, north to Oregon, Washington, and British Columbia.

1154. *Myotis orinomus* Elliot.

Myotis orinomus Elliot, Pub. Field Columb. Mus., III, 1903, p. 228.

Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M.

Pub., IV, Pt. II, 1904, p. 577. Zoöl. Ser.

LA GRULLA BROWN BAT.

Type locality. La Grulla, San Pedro Martir Mountains, Lower California, Mexico. Altitude, 8,000 feet.

Geogr. Distr. Lower California, Mexico. Range unknown.

1155. *Myotis californicus* (Audubon & Bachman).

Vespertilio californicus Aud. & Bachm., Journ. Acad. Nat. Scien. Phil., 1842, p. 285.

Vespertilio nitidus H. Allen, Proc. Acad. Nat. Scien. Phil., 1862, p. 247.

Vespertilio oregonensis H. Allen, Mon. N. Am. Bats, 1864, p. 61.

Vespertilio obscurus H. Allen, Proc. Acad. Nat. Scien. Phil., 1866, p. 281.

Vespertilio volans H. Allen, Proc. Acad. Nat. Scien. Phil., 1866, p. 282.

Vespertilio exilis H. Allen, Proc. Acad. Nat. Scien. Phil., 1866, p. 283.

Vespertilio tenuidorsalis H. Allen, Proc. Acad. Nat. Scien. Phil., 1866, p. 283.

Vespertilio yumanensis H. Allen, Proc. Acad. Nat. Scien. Phil., 1866, p. 283. (nec H. Allen, 1864.)

Vespertilio melanorhinus Merr., N. Am. Faun., No. 3, 1890, p. 46.

Vespertilio nitidus henshawi H. Allen, Mon. N. Am. Bats, 1894, p. 103.

Vespertilio nigricans H. Allen, Mon. N. Am. Bats, 1894, p. 97. (nec Max., 1826.)

Myotis californicus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 403. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W.

Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 578. Zoöl. Ser.

LITTLE CALIFORNIA BAT.

Type locality. California.

Geogr. Distr. From Wyoming south into Lower California and northern Mexico, Mexico.

a. *Myotis californicus caurinus* Miller.

Myotis californicus caurinus Miller, N. Am. Faun., No. 13, 1897,
p. 72. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901,
p. 404. Zoöl. Ser.

GRAHAM ISLAND BAT.

Type locality. Massett, Graham Island, Queen Charlotte Islands,
British Columbia, Canada.

Geogr. Distr. Humid coast districts of British Columbia, Wash-
ington, Oregon, and possibly northern California.

b. *Myotis californicus ciliolabrum* (Merriam).

Vespertilio ciliolabrum Merr., Proc. Biol. Soc. Wash., IV, 1886, p. 2.

Myotis californicus ciliolabrum Elliot, Syn. N. Am. Mamm.,
F. C. M. Pub., II, 1901, p. 404. Zoöl. Ser.

HAIRY-LIPPED BAT.

Type locality. Near Banner, Trego County, Kansas.

Geogr. Distr. Kansas and central South Dakota; limits of
range unknown.

c. *Myotis californicus jaliscensis* Menegaux.

Myotis californicus jaliscensis Meneg., Bull. Mus. Hist. Nat. Paris,
1901, p. 321. Elliot, Mamm. Middle Amer. & W. Indies,
F. C. M. Pub., IV, Pt. II, 1904, p. 579. Zoöl. Ser.

JALISCO BROWN BAT.

Type locality. Vicinity of Lake Zacoalco, State of Jalisco, Mexico.

Geogr. Distr. State of Jalisco, Mexico. Range unknown.

d. *Myotis californicus mexicanus* (Saussure).

Vespertilio mexicanus Sauss., Rev. Mag. Zool., 2me Sér., XII, 1860,
p. 282.

Myotis californicus mexicanus Elliot, Mamm. Middle Amer. & W.
Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 579. Zoöl. Ser.

LITTLE MEXICAN BAT.

Type locality. Unknown.

Geogr. Distr. States of San Luis Potosi, Michoacan, and Oaxaca,
Mexico. Limits of range unknown.

e. *Myotis californicus pallidus* Stephens.

Myotis californicus pallidus Steph., Proc. Biol. Soc. Wash., XIII,
1900, p. 153. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II,
1901, p. 405. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. In-
dies, F. C. M. Pub., IV, Pt. II, 1904, p. 579. Zoöl. Ser.

PALE BAT.

Type locality. Vallecito, San Diego County, California.

Geogr. Distr. Southern California, into Lower California, Mexico.

f. Myotis californicus durangæ J. A. Allen.

Myotis californicus durangæ J. A. Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1903, p. 612. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 579. Zoöl. Ser.

DURANGO BROWN BAT.

Type locality. Rio Sestin, State of Durango, Mexico.

Geogr. Distr. State of Durango, Mexico. Range unknown.

1156. *Myotis subulatus* (Say).

Vespertilio subulatus Say, in Long's Exped. Rocky Mts., II, 1823, p. 65.

Vespertilio gryphus var. (b) H. Allen, Mon. N. Am. Bats, 1894, p. 80.

Vespertilio gryphus var. *septentrionalis* Trouess., Cat. Mamm. t. Viv. q. Foss., 1897, p. 131.

Myotis subulatus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 405. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 580. Zoöl. Ser.

SMALL-WINGED BAT.

Type locality. Near La Junta, Arkansas River, Otero County, Colorado.

Geogr. Distr. United States and Canada, east of Rocky Mountains south to State of Sonora, Mexico.

a. Myotis subulatus keenii (Merriam).

Vespertilio subulatus keenii Merr., Amer. Nat., XXIX, 1895, p. 860.

Myotis subulatus keenii Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 405. Zoöl. Ser.

KEENE'S BROWN BAT.

Type locality. Massett, Graham Island, Queen Charlotte Islands, British Columbia, Canada.

Geogr. Distr. Known only from the type locality.

1157. *Myotis lucifugus* (Le Conte).

Vespertilio lucifugus Le Conte, McMurtrie, Cuv. Anim. Kingd., I, App., 1831, p. 431.

Vespertilio affinis H. Allen, Mon. N. Am. Bats, 1864, p. 53.

Vespertilio carolinæ Dobson, Cat. Chirop. Brit. Mus., 1878, p. 325.

Vespertilio lucifugus austroriparius Rhoads, Proc. Acad. Nat. Scien. Phil., 1897, p. 227.

Myotis lucifugus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 402, fig. 84. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 580. Zoöl. Ser.

LITTLE BROWN BAT.

Type locality. Near Riceboro, southern part of Liberty County, Georgia.

Geogr. Distr. Hudson Bay to Brazil, South America.

a. *Myotis lucifugus longicrus* (True).

Vespertilio longicrus True, Science, VIII, 1886, p. 588.

Vespertilio albescens H. Allen, Mon. N. Am. Bats, 1893, p. 92.
(nec Geoffroy.)

Myotis lucifugus longicrus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 402. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 581. Zoöl. Ser.

TRUE'S BAT.

Type locality. Puget Sound, Washington.

Geogr. Distr. Puget Sound, east to Wyoming, south to southern California and Arizona; also Lower California and northern Mexico.

b. *Myotis lucifugus alascensis* Miller.

Myotis lucifugus alascensis Miller, N. Am. Faun., No. 13, 1897, p. 63. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 402. Zoöl. Ser.

ALASKA BAT.

Type locality. Sitka, Alaska.

Geogr. Distr. Coast district of southern Alaska, and also of northern British Columbia, Canada.

1158. *Myotis thysanodes* Miller.

Vespertilio albescens evotis H. Allen, Mon. N. Am. Bats, 1894, p. 90.

Vespertilio albescens velifer H. Allen, Mon. N. Am. Bats, 1894, p. 93.

Myotis thysanodes Miller, N. Am. Faun., No. 13, 1897, p. 80.
Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 572. Zoöl. Ser.

Myotis evotis thysanodes Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 406. Zoöl. Ser.

FRINGED BAT.

Type locality. Old Fort Tejon, in mountains south of Kern Lake, Kern County, California.

Geogr. Distr. Southern California to State of Michoacan, Mexico.

Genus 156. **Lasionycteris.**

Lasionycteris Peters, Monatsb. K. Preuss. Akad. Wiss. Berlin, 1865, p. 648. Type *Vespertilio noctivagans* Le Conte.

Scotophilus H. Allen, Mon. N. Am. Bats, 1864, p. 27. (nec Leach, Part.)

- Cnephaiophilus* Fitzin., Sitzung. K. Akad. Wiss. Wien, LXII, 1870, abth. 1, p. 8.
Vesperides Coues, in Coues & Yarrow, Zoöl. Wheeler's Exped., 1875, p. 83.
Vesperugo Dobson, Cat. Chirop. Brit. Mus., 1878, p. 183. (Part.)

1159. *Lasionycteris noctivagans* (Le Conte).

- Vespertilio noctivagans* Le Conte, McMurtrie, Cuv. Anim. Kingd., Am. ed., 1, 1831, p. 31.
Vespertilio auduboni Harl., Am. Month. Journ. Geol. & Nat. Hist., 1831, p. 220, pl. II.
Vespertilio pulverulentus Temm., Mon. Mamm., II, 1835, p. 325.
Lasionycteris noctivagans Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 407, fig. 85. Zoöl. Ser.

NIGHT-WANDERING BAT.

- Type locality.* Eastern United States; exact locality unknown.
Geogr. Distr. North America, north of Mexico, from the Atlantic to the Pacific Ocean.

Genus 157. *Pipistrellus*.

- Pipistrellus*** Kaup, Skizzirte Entwick.-Gesch. u. Natürl. Syst. d. Europ. Thierw., Th. 1, 1829, pp. 97, 98. Type *Vespertilio pipistrellus* Schreber.
Vesperugo Keyserl. & Blas., Wieg. Archiv f. Naturg., 5ter Jahrg., Bd. 1, 1839, p. 312. (Part.)
Nannugo Kolenati, Allgem. Deutsch. Naturh. Zeitg. Dresden, neue Folge II, 1856, pp. 131, 169-172.
Hypsugo Kolenati, Allgem. Deutsch. Naturh. Zeitg. Dresden, neue Folge II, 1856, pp. 131, 167-169.
Scotophilus H. Allen, Mon. N. Am. Bats, 1864, p. 27. (Part, nec Leach.)

1160. *Pipistrellus hesperus* (H. Allen).

- Scotophilus hesperus* H. Allen, Mon. N. Am. Bats, 1864, p. 43.
Vesperugo merriami Dobson, Cat. Chirop. Brit. Mus., 1878, p. 228.
Pipistrellus hesperus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 409, fig. 86. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 582, figs. 108, LXXXVI. Zoöl. Ser.

WESTERN BAT.

- Type locality.* Old Fort Yuma, San Diego County, California.

Geogr. Distr. Western Texas to Pacific coast and southern California, south into Lower California, Mexico.

a. Pipistrellus hesperus apus Elliot.

Pipistrellus hesperus apus Elliot, Pub. Field Columb. Mus., III, 1904, p. 269. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 583. Zoöl. Ser.

SWIFT BAT.

Type locality. Providentia Mines, State of Sonora, Mexico.

Geogr. Distr. State of Sonora, Mexico. Range unknown.

b. Pipistrellus hesperus australis Miller.

Pipistrellus hesperus australis Miller, N. Am. Faun., No. 13, 1897, p. 90. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 584. Zoöl. Ser.

ALLIED BAT.

Type locality. Barranca Ibarra, State of Jalisco, Mexico.

Geogr. Distr. Known only from the type locality.

1161. *Pipistrellus subflavus* (F. Cuvier).

Vespertilio subflavus F. Cuv., Nouv. Ann. Mus. Hist. Nat. Paris, 1832, p. 17.

Vespertilio erythrodactylus Temm., Mon. Mamm., II, 1835-41, p. 238.

Vesperugo carolinensis H. Allen, Mon. N. Am. Bats, 1894, p. 121. (nec Geoffroy.)

Pipistrellus subflavus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 409. Zoöl. Ser.

YELLOWISH-BROWN BAT.

Type locality. Eastern United States. Georgia?

Geogr. Distr. Atlantic coast west to Iowa, south to eastern and southern Texas.

a. Pipistrellus subflavus obscurus Miller.

Pipistrellus subflavus obscurus Miller, N. Am. Faun., No. 13, 1897, p. 93. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 409. Zoöl. Ser.

DUSKY BAT.

Type locality. Lake George, Warren County, New York.

Geogr. Distr. Known only from the type locality.

1162. *Pipistrellus veræcrucis* (Ward).

Vesperugo veræcrucis Ward, Amer. Nat., xxv, 1891, p. 745.

Pipistrellus veræcrucis Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 584. Zoöl. Ser.

LAS VIGAS BAT.

Type locality. Las Vigas, Jalapa, State of Vera Cruz, Mexico.

Geogr. Distr. Known only from the type locality.

1163. *Pipistrellus vagans* (Dobson).

Vesperugo vagans Dobson, Ann. Mag. Nat. Hist., 5th Ser., IV, 1879, p. 135.

Pipistrellus vagans Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 584. Zool. Ser.

WANDERING BERMUDA BAT.

Type locality. Bermuda.

Geogr. Distr. Bermuda.

1164. *Pipistrellus cinnamomeus* Miller.

Pipistrellus cinnamomeus Miller, Proc. Acad. Nat. Scien. Phil., 1902, p. 390. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 585. Zool. Ser.

CINNAMON BAT.

Type locality. Montechristo, State of Tabasco, Mexico.

Geogr. Distr. Known only from the type locality.

Genus 158. *Vespertilio*.

Vespertilio Linn., Syst. Nat., I, 1758, p. 31; I, 1766, p. 46. Type *Vespertilio murinus* Linnæus. (nec Schreber.)

Eptesicus Rafin., Ann. Nat., I, 1820, p. 2.

Cnephæus Kaup, Skizzirte Entw.-Gesch. u. Natürl. Syst. Europ. Thierw., I, 1829, p. 103.

Noctula Bonap., Iconog. Faun. Ital., I, 1837, fasc. XXI.

Vesperugo Keyserl. & Blas., Wieg. Archiv f. Naturg., Bd. I, 1839, p. 312. (Part.)

Vesperus Keyserl. & Blas., Wieg. Archiv f. Naturg., Bd. I, 1839, p. 313.

Cateorus Kolenati, Allgem. Deutsch. Naturh. Zeit. Dresden, neue Folge, II, 1856, pp. 131, 162-163.

Meteoros Kolenati, Allgem. Deutsch. Naturh. Zeit. Dresden, neue Folge, II, 1856, pp. 131, 163-167.

Marsipolæmus Peters, Monatsb. K. Preuss. Akad. Wiss. Berl., 1872, p. 260.

Adelonycteris H. Allen, Proc. Acad. Nat. Scien. Phil., 1892, p. 466.

Subgenus A. *Vespertilio*.1165. *Vespertilio fuscus* Beauvois.

Vespertilio fuscus Beauv., Cat. Peale's Mus. Phil., 1796, p. 14.

Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 410.

fig. 87. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 587, figs. 109, LXXXVII. Zoöl. Ser. *Vespertilio carolinensis* Geoff., Ann. Mus. Hist. Nat. Paris, VIII, 1806, p. 193.

Vespertilio phaiops Rafin., Am. Month. Mag., III, 1818, p. 445.

Eptesicus melanops Rafin., Ann. Nat., 1820, p. 2.

Vespertilio arquatus Say, Long's Exped. Rocky Mts., I, 1823, p. 167. (note.)

Vespertilio ursinus Temm., Mon. Mamm., II, 1835, p. 235.

Scotophilus greenii Gray, List. Spec. Mamm. Brit. Mus., I, 1843, p. 30.

BROWN BAT.

Type locality. Philadelphia, Pennsylvania.

Geogr. Distr. Eastern United States north of Florida, and in the adjoining British Provinces.

a. *Vespertilio fuscus osceola* (Rhoads).

Eptesicus fuscus osceola Rhoads, Proc. Acad. Nat. Scien. Phil., 1901, p. 618.

FLORIDA BROWN BAT.

Type locality. Tarpon Springs, Hillsboro County, Florida.

Geogr. Distr. Florida.

Genl. Char. Similar to *V. fuscus*, but of a cinnamon brown color instead of sepia.

Color. Cinnamon brown, uniform.

Measurements. Total length, 110; tail, 38; hind foot, 9.5.

b. *Vespertilio fuscus melanopterus* (Rehn).

Eptesicus fuscus melanopterus Rehn, Proc. Acad. Nat. Scien. Phil., 1904, p. 590.

BLACK-WINGED BAT.

Type locality. Mount Tallac, California.

Geogr. Distr. Known only from the type locality.

Genl. Char. Similar to *V. f. osceola* Rhoads, color different.

Color. Above dark cinnamon; lightest on top of head and on shoulders; beneath reddish wood brown; membranes and face blackish.

Measurements. Total length, 104; tail vertebræ, 40; ear, 12; tibia, 17; forearm (average of five), 47.

c. *Vespertilio fuscus bernardinus* (Rhoads).

Eptesicus fuscus bernardinus Rhoads, Proc. Acad. Nat. Scien. Phil., 1901, p. 619.

Vespertilio fuscus bernardinus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 589. Zoöl. Ser.

SAN BERNARDINO BROWN BAT.

Type locality. San Bernardino Valley, California.

Geogr. Distr. Southern California and Lower California, Mexico.

d. *Vespertilio fuscus cubensis* (Gray).

Scotophilus cubensis Gray, Ann. Nat. Hist., IV, 1839, p. 7.

Vespertilio dutertreus Gerv., in Ram. de la Sagra's Hist. Isle de Cuba, Mamm., 1840, p. 6. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 588. Zoöl. Ser.

CUBAN BROWN BAT.

Type locality. Cuba.

Geogr. Distr. Island of Cuba.

e. *Vespertilio fuscus miradorensis* (H. Allen).

Scotophilus miradorensis H. Allen, Proc. Acad. Nat. Scien. Phil., 1866, p. 287.

Vespertilio fuscus miradorensis Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 588. Zoöl. Ser.

MIRADOR BROWN BAT.

Type locality. Mirador, State of Vera Cruz, Mexico.

Geogr. Distr. Southern Mexico into Guatemala and Costa Rica.

f. *Vespertilio fuscus bahamensis* Miller.

Vespertilio fuscus bahamensis Miller, N. Am. Faun., No. 13, 1897, p. 101. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 588. Zoöl. Ser.

BAHAMA BROWN BAT.

Type locality. Nassau, New Providence Island, Bahamas.

Geogr. Distr. Known only from the type locality.

g. *Vespertilio fuscus peninsulæ* Thomas.

Vespertilio fuscus peninsulæ Thomas, Ann. Mag. Nat. Hist., 7th Ser., 1898, p. 43. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 589. Zoöl. Ser.

LOWER CALIFORNIA BROWN BAT.

Type locality. Sierra Laguna, Lower California, Mexico.

Geogr. Distr. Lower California, Mexico.

h. *Vespertilio fuscus propinquus* (Peters).

Vesperus propinquus Peters, Monatsb. K. Preuss. Akad. Wiss. Berl., 1872, p. 262.

Vespertilio fuscus propinquus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 589. Zoöl. Ser.

ALLIED BROWN BAT.

Type locality. Santa Isabel, Guatemala.

Geogr. Distr. Guatemala and Nicaragua, Central America.

1166. Vespertilio gaumeri (J. A. Allen).

Adelonycteris gaumeri J. A. Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1897, p. 231.

Vespertilio gaumeri Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 590. Zoöl. Ser.

YUCATAN BROWN BAT.

Type locality. Izamal, Yucatan, Mexico.

Geogr. Distr. Yucatan, Mexico.

Subgenus B. **Marsipolæmus.**

1167. Vespertilio albigularis (Peters).

Vesperus albigularis Peters, Monatsb. K. Preuss. Akad. Wiss. Berl., 1872, p. 260.

Vespertilio albigularis Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 590. Zoöl. Ser.

WHITE-THROATED BROWN BAT.

Type locality. Mexico?

Geogr. Distr. Unknown.

Genus 159. **Lasiurus.**

Lasiurus Gray, Zoöl. Misc., 1831, No. 1, p. 38. Type *Vespertilio borealis* Müller.

Atalapha Peters, Monatsb. K. Preuss. Akad. Wiss. Berl., 1870, p. 907. (nec Rafin.)

1168. Lasiurus borealis (Müller).

Vespertilio borealis Müll., Natursyst., Suppl., 1776, p. 21.

Vespertilio noveboracensis Erxl., Syst. Reg. Anim., I, 1777, p. 155.

Vespertilio lasiurus Schreb., Säugth., Abth. I, 1781, Taf LXII B.

Vespertilio noveboracensis Bodd., Elench. Anim., I, 1785, p. 71.

Vespertilio rubellus Beauv., Cat. Peale's Mus., 1796, p. 204.

Vespertilio rubra Ord, Guthrie's Geogr., 2d Amer. ed., II, 1815, p. 291.

Vespertilio tessellatus Rafin., Am. Month. Mag., III, 1818, p. 445.

Vespertilio monachus Rafin., Am. Month. Mag., III, 1818, p. 445.

Vespertilio rufus Warden, Desc. États-Unis Amér., v, 1820, p. 606.

Lasiurus funebris Fitzin., Sitzungsab. K. Akad. Wiss. Wien., LXII, 1870, p. 46.

Atalapha borealis Rhoads, Amer. Nat., xxviii, 1894, p. 523.

Lasiurus borealis Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 411, fig. 88. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 592. Zoöl. Ser.

RED BAT.

Type locality. State of New York.

Geogr. Distr. North America east of Rocky Mountains, from Canada southward through Mexico and Central America into South America to Brazil, and Chili.

a. *Lasiurus borealis pfeifferi* (Gundlach).

Atalapha pfeifferi Gundl., Monatsb. K. Preuss. Akad. Wiss. Berl., 1861, p. 152.

Lasiurus borealis pfeifferi Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 593. Zoöl. Ser.

PFEIFFER'S BAT.

Type locality. Cuba.

Geogr. Distr. Island of Cuba.

b. *Lasiurus borealis seminolus* (Rhoads).

Atalapha borealis seminola Rhoads, Proc. Acad. Nat. Scien. Phil., 1895, p. 32.

Lasiurus borealis seminolus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 413. Zoöl. Ser.

FLORIDA RED BAT.

Type locality. Tarpon Springs, Hillsboro County, Florida.

Geogr. Distr. South Carolina to southern Texas; Nassau, New Providence Island, Bahamas.

c. *Lasiurus borealis teliotis* (H. Allen).

Atalapha teliotis H. Allen, Proc. Am. Phil. Soc., XXIX, 1891, p. 5.

Lasiurus borealis teliotis Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 413. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 593. Zoöl. Ser.

CALIFORNIA RED BAT.

Type locality. Unknown. California?

Geogr. Distr. Sacramento Valley, California, south to Comondu, Lower California, Mexico.

d. *Lasiurus borealis mexicanus* (Saussure).

Atalapha mexicana Sauss., Rev. Mag. Zool., 2me Sér., XIII, 1861, p. 97.

Atalapha frantzii Peters, Monatsb. K. Preuss. Akad. Wiss. Berl., 1871, p. 908.

Lasiurus borealis mexicanus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 594. Zoöl. Ser.

MEXICAN RED BAT.

Type locality. Southern Mexico; exact locality unknown.

Geogr. Distr. Tres Marias Islands, State of Jalisco to State of Vera Cruz, south into Central America.

1169. Lasiurus cinereus (Beauvois).

Vespertilio cinereus Beauv., Cat. Peale's Mus. Phil., 1796, p. 15.

Vespertilio pruinosus Say, Long's Exped. Rocky Mts., 1, 1823, p. 167. (Footnote.)

Lasiurus cinereus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 413. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 11, 1904, p. 595, figs. 110, LXXXVIII. Zoöl. Ser.

HOARY BAT.

Type locality. Philadelphia, Pennsylvania.

Geogr. Distr. North America from the Atlantic to the Pacific, at different seasons, south through Mexico, Central and South America to Chili.

Genus 160. **Dasypterus.**

Dasypterus Peters, Monatsb. K. Preuss. Akad. Wiss. Berl., 1871, p. 912. Type *Lasiurus intermedius* H. Allen.

1170. Dasypterus intermedius (H. Allen).

Lasiurus intermedius H. Allen, Proc. Acad. Nat. Scien. Phil., 1862, p. 246.

Dasypterus intermedius Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 414, fig. 89. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 11, 1904, p. 595, fig. LXXXIX. Zoöl. Ser.

INTERMEDIATE BAT.

Type locality. Matamoros, State of Tamaulipas, Mexico.

Geogr. Distr. State of Tamaulipas, Mexico, north to the Gulf States of the United States, except possibly Florida.

Dasypterus ega xanthinus Thomas.

Dasypterus ega xanthinus Thomas, Ann. Mag. Nat. Hist., 6th Ser., 1897, p. 544. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 11, 1904, p. 597, fig. 111. Zoöl. Ser.

SIERRA LAGUNA BAT.

Type locality. Sierra Laguna, Lower California, Mexico.

Geogr. Distr. Lower California, Mexico.

Dasypterus ega panamensis Thomas.

Dasypterus ega panamensis Thomas, Ann. Mag. Nat. Hist., 7th Ser., VIII, 1901, p. 246. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. 11, 1904, p. 597. Zoöl. Ser.

PANAMA BAT.

Type locality. Bogava, Chiriqui, Panama. Altitude, 1,000 feet.
Geogr. Distr. South Central America. Range unknown.

1171. *Dasypterus floridanus* Miller.

Dasypterus floridanus Miller, Proc. Acad. Nat. Scien. Phil., 1902,
 p. 392.

MILLER'S FLORIDA BAT.

Type locality. Lake Kissimmee, Osceola County, Florida.

Geogr. Distr. Florida.

Genl. Char. Like *D. intermedius*, but smaller.

Color. Same as *D. intermedius*.

Measurements. Total length, 129; tail, 52; forearm, 49; thumb, 10; second finger, 61; third finger, 98; fourth finger, 80; fifth finger, 65; tibia, 20; foot, 9; ear from crown, 11; from meatus, 17; width of ear, 11.

Genus 161. *Nycticeius*.

Nycticeius Rafin., Journ. Phys., LXXXVIII, 1819, p. 417. Type (by elimination) *Vespertilio humeralis* Rafinesque.

1172. *Nycticeius humeralis* (Rafinesque).

Vespertilio humeralis Rafin., Amer. Month. Mag., III, 1818, p. 445.

Nycticeia crepuscularis Le Conte, McMurtrie, Cuv. Anim. King., I,
 1831, p. 432.

Nycticejus humeralis Elliot, Syn. N. Am. Mamm., F. C. M. Pub.,
 II, 1901, p. 415, fig. 90. Zoöl. Ser.

Nycticeius humeralis Elliot, Mamm. Middle Amer. & W. Indies,
 F. C. M. Pub., IV, Pt. II, 1904, p. 598, figs. 112, xc. Zoöl. Ser.

RAFINESQUE'S BAT.

Type locality. Kentucky.

Geogr. Distr. Eastern United States west to Arkansas and southern Texas.

a. *Nycticeius humeralis cubanus* (Gundlach).

Vesperus cubanus Gund., Monatsb. K. Preuss. Akad. Wiss. Berl.,
 1861, p. 150.

Nycticeius humeralis cubanus Elliot, Mamm. Middle Amer. & W.
 Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 599. Zoöl. Ser.

GUNDLACH'S BAT.

Type locality. Near Cardenas, Cuba.

Geogr. Distr. Island of Cuba.

Genus 162. ***Rhogoessa**.

Rhogeëssa! H. Allen, Proc. Acad. Nat. Scien. Phil., 1866, p. 285.
Type *Rhogeëssa! tumida* H. Allen.

1173. Rhogoessa tumida H. Allen.

Rhogeëssa! tumida H. Allen, Proc. Acad. Nat. Scien. Phil., 1866,
p. 286.

Rhogōessa tumida Elliot, Mamm. Middle Amer. & W. Indies,
F. C. M. Pub., IV, Pt. II, 1904, p. 601, figs. 113, xci. Zoöl. Ser.

RESTLESS BAT.

Type locality. Mirador, State of Vera Cruz, Mexico.

Geogr. Distr. Southern Mexico into Central America.

1174. Rhogoessa parvula H. Allen.

Rhogeëssa! parvula H. Allen, Proc. Acad. Nat. Scien. Phil., 1866,
p. 285.

Rhogōessa parvula Elliot, Mamm. Middle Amer. & W. Indies,
F. C. M. Pub., IV, Pt. II, 1904, p. 601. Zoöl. Ser.

TRES MARIAS ISLANDS BAT.

Type locality. Tres Marias Islands, State of Jalisco, Mexico.

Geogr. Distr. Tres Marias Islands, State of Jalisco, through
Mexico to Costa Rica, Central America.

1175. Rhogoessa gracilis Miller.

Rhogeëssa! gracilis Miller, N. Am. Faun., No. 13, 1897, p. 126.

Rhogōessa gracilis Elliot, Mamm. Middle Amer. & W. Indies,
F. C. M. Pub., IV, Pt. II, 1904, p. 602. Zoöl. Ser.

GRACEFUL BAT.

Type locality. Piaxtla, State of Puebla, Mexico.

Geogr. Distr. State of Puebla, Isthmus of Tehuantepec, Mexico.

1176. Rhogoessa alleni Thomas.

Rhogeëssa! alleni Thomas, Ann. Mag. Nat. Hist., 6th Ser., x, 1892,
p. 477.

Rhogōessa alleni Elliot, Mamm. Middle Amer. & W. Indies, F. C. M.
Pub., IV, Pt. II, 1904, p. 602. Zoöl. Ser.

SANTA ROSALIA BAT.

Type locality. Santa Rosalia, near Autlin, State of Jalisco,
Mexico.

Geogr. Distr. Known only from the type locality.

* ρῶξ, ῥωγος-έσσα = *rhogoessa*, having a cleft or rent.

Subfamily II. **Plecotinæ.**Genus 163. **Euderma.**

Euderma H. Allen, Proc. Acad. Nat. Scien. Phil., 1892, p. 467. Type

Histiotus maculatus J. A. Allen.

Histiotus J. A. Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1891,
p. 195. (nec Gervais.)

1177. **Euderma maculatum** (J. A. Allen).

Histiotus maculatus J. A. Allen, Bull. Am. Mus. Nat. Hist. N. Y.,
1891, p. 195.

Euderma maculata Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11,
1901, p. 398. Zool. Ser.

SPOTTED GREAT-EARED BAT.

Type locality. Near Piru, Ventura County, California.

Geogr. Distr. Known only from the type locality.

Genus 164. **Corynorhinus. Club-nosed or Big-eared Bats.**

Corynorhinus H. Allen, Proc. Acad. Nat. Scien. Phil., 1865, p. 173.

Type *Plecotus macrotis* Le Conte.

Plecotus Le Conte, McMurtrie, Cuv. Anim. King., 1, App., 1831,
p. 431. (nec Geoffroy.)

Synotus H. Allen, Mon. N. Am. Bats, 1864, p. 62. (nec Keyserl.
& Blasius.)

1178. **Corynorhinus macrotis** (Le Conte).

Plecotus macrotis Le Conte, McMurtrie, Cuv. Anim. Kingd., 1, 1831,
p. 431.

Plecotus lecontii Cooper, Ann. N. Y. Lyc. Nat. Hist., 14, 1837,
p. 72.

Corynorhinus macrotis Elliot, Syn. N. Am. Mamm., F. C. M. Pub.,
11, 1901, p. 399. Zool. Ser.

BIG-EARED BAT.

Type locality. Probably Le Conte's plantation near Riceboro,
Georgia.

Geogr. Distr. South Atlantic States to Louisiana.

a. **Corynorhinus macrotis pallescens** Miller.

Synotus townsendi H. Allen, Mon. N. Am. Bats, 1864, p. 65. (nec
P. townsendi Cooper.)

Corynorhinus macrotis pallescens Miller, N. Am. Faun., No. 13, 1897, p. 52. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 399. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., 14, Pt. 11, 1904, p. 604, figs. 114, xcii (*C. macrotis*). Zoöl. Ser.

PALLID BIG-EARED BAT.

Type locality. Keam Cañon, Navajo County, Arizona.

Geogr. Distr. From California and Colorado south and east to western Texas, and south to State of Oaxaca, Mexico.

b. *Corynorhinus macrotis townsendi* (Cooper).

Plecotus townsendii Cooper, Ann. N. Y. Lyc. Nat. Hist., 14, 1837, p. 73.

Corynorhinus macrotis townsendi Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 400, fig. 83. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., 14, Pt. 11, 1904, p. 604. Zoöl. Ser.

TOWNSEND'S BIG-EARED BAT.

Type locality. Region of the Columbia River, Oregon.

Geogr. Distr. Humid coast region of Oregon, Washington, and southern British Columbia, Canada.

Subfamily III. **Antrozoinæ.**

Genus 165. **Antrozous.**

Antrozous H. Allen, Proc. Acad. Nat. Scien. Phil., 1862, p. 248.
Type *Vespertilio pallidus* Le Conte.

1179. **Antrozous pallidus** (Le Conte).

Vespertilio pallidus Le Conte, Proc. Acad. Nat. Scien. Phil., 1856, p. 437.

Antrozous pallidus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 396, fig. 82. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., 14, Pt. 11, 1904, p. 605, figs. 115, xciii. Zoöl. Ser.

BIG-EARED PALE BAT.

Type locality. El Paso, El Paso County, Texas.

Geogr. Distr. Nevada, eastern Texas, south to western Mexico and to Cape St. Lucas, Lower California, Mexico.

a. *Antrozous pallidus pacificus* Merriam.

Antrozous pallidus pacificus Merr., Proc. Biol. Soc. Wash., 11, 1897, p. 180. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., 11, 1901, p. 397. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., 14, Pt. 11, 1904, p. 607. Zoöl. Ser.

PACIFIC BIG-EARED BAT.

Type locality. Old Fort Tejon, in mountains south of Kern Lake, Kern County, California.

Geogr. Distr. Southern California, into Lower California, Mexico.

1180. *Antrozous minor* Miller.

Antrozous minor Miller, Proc. Acad. Nat. Scien. Phil., 1902, p. 389. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 607. Zoöl. Ser.

LITTLE COMONDU BAT.

Type locality. Comondu, Lower California, Mexico.

Geogr. Distr. Southern Lower California, Mexico.

Family II. **Noctilionidæ.**Subfamily I. **Emballonurinae.**Genus 166. **Rhynchonycteris.**

Rhynchonycteris Peters, Monatsb. K. Preuss. Akad. Wiss. Berl., 1867, p. 477. Type *Vespertilio naso* Maximilian.

1181. **Rhynchonycteris naso** (Maximilian).

Vespertilio naso Max., Schinz's Thierr., I, 1821, p. 179.

Proboscidea saxatilis Spix, Simiar. et Vespert. Bras., 1823, p. 62, pl. xxxv, fig. VIII.

Proboscidea rivalis Spix, Simiar. et Vespert. Bras., 1823, p. 62, pl. xxxv.

Emballonura lineata Temm., Mon. Mamm., II, 1835-41, p. 297.

Rhynchonycteris naso Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 609, figs. 116, xciv. Zoöl. Ser.

LONG-NOSED BAT.

Type locality. East coast of Brazil.

Geogr. Distr. Guatemala through Central America, to Brazil, South America.

Genus 167. **Saccopteryx.**

Saccopteryx Illig., Prodr. Syst. Mamm. et Av., 1811, p. 121. Type

Vespertilio lepturus Schreber.

Urocryptus Temm., V. d. Hoev., Tijds. Nat. Gesch., v, 1838-39, pp. 31-34, pl. II, figs. 3, 4.

Centronycteris Gray, Mag. Zoöl. Bot., II, 1838, p. 499.

1182. Saccopteryx bilineata (Temminck).

Urocryptus bilineatus Temm., V. d. Hoev., Tijds. Nat. Gesch. en Physiol., v, 1838, p. 33, figs. 3, 4.

Saccopteryx bilineata Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 610, fig. 117. Zoöl. Ser.

WHITE-STRIPED BAT.

Type locality. Surinam.

Geogr. Distr. Guatemala, Central America to Dutch Guiana, South America.

Genus 168. Balantiopteryx.

Balantiopteryx Peters, Monatsb. K. Preuss. Akad. Wiss. Berl., 1867, p. 476. Type *Balantiopteryx plicata* Peters.

1183. Balantiopteryx plicata Peters.

Balantiopteryx plicata Peters, Monatsb. K. Preuss. Akad. Wiss. Berl., 1867, p. 476. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 611, figs. 118, xcv. Zoöl. Ser.

POUCHED BAT.

Type locality. Puente Arenas, Costa Rica, Central America.

Geogr. Distr. State of Jalisco, opposite Tres Marias Islands Mexico, into Central America.

1184. Balantiopteryx infusca (Thomas).

Saccopteryx infusca Thomas, Ann. Mag. Nat. Hist., 6th Ser., xx, 1897, p. 546.

Balantiopteryx infusca Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 612. Zoöl. Ser.

THOMAS' BAT.

Type locality. Cachavi, northern Ecuador.

Geogr. Distr. Coban, Guatemala to Ecuador, South America.

Genus 169. Peropteryx.

Peropteryx Peters, Monatsb. K. Preuss. Akad. Wiss. Berl., 1867, p. 472. Type *Vespertilio caninus* Maximilian.

1185. Peropteryx canina (Maximilian).

Vespertilio caninus Max., Schinz's Thierr., I, 1821, p. 179.

Peropteryx canina Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 614, figs. 119, xcvi. Zoöl. Ser.

DOG-LIKE BAT.

Type locality. East coast of Brazil.

Geogr. Distr. Guatemala, Central America to Brazil.

Genus 170. * **Diclidurus.**

Diclidurus Max., Isis, 1820, p. 1629. Type *Diclidurus albus* Maximilian.

1186. **Diclidurus albus** Maximilian.

Diclidurus albus Max., Isis, 1820, p. 1630. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 615, figs. 120, xcvi. Zoöl. Ser.

Diclidurus frëyreissii Max., Abbild. zur Naturg. Bras., 1838, pl. xvi.

WHITE BAT.

Type locality. Mouth of Rio Pardo, east coast of Brazil.

Geogr. Distr. Costa Rica, Central America, south to Brazil, South America.

1187. **Diclidurus virgo** Thomas.

Diclidurus virgo Thomas, Ann. Mag. Nat. Hist., 7th Ser., 1903, p. 377. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 615. Zoöl. Ser.

ESCAZU WHITE BAT.

Type locality. Escazu, Costa Rica.

Geogr. Distr. Central America. Range unknown.

Subfamily II. **Noctilioninæ.**Genus 171. **Noctilio.**

Noctilio Linn., Syst. Nat., I, 1766, p. 88. Type *Vespertilio leporinus* Linnæus.

Noctilio leporinus mastivus (Dahl).

Vespertilio mastivus Dahl, Skrift. Naturh. Selsk. Kjöbenh., IV, 1797, p. 132, pl. 7.

Noctilio dorsatus Max., Beit. Natur. Bras., 1826, p. 223.

Noctilio leporinus Burm., Thier. Bras., 1854, p. 60. (nec Linn.)

Noctilio leporinus mastivus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 617, figs. 121, xcvi. Zoöl. Ser.

*Misprinted in Mamm. Middle Amer., etc., p. 614, *Declidurus*.

MUSKY BAT.

Type locality. Island of St. Croix, West Indies.

Geogr. Distr. Islands of Jamaica, Mona and St. Croix, West Indies, and in South America, to Brazil.

Family III. **Molossidae.**Genus 172. **Molossus.**

Molossus E. Geoff., Ann. Mus. Hist. Nat. Paris, vi, 1805, p. 151.

Type Molossus rufus E. Geoffroy.

Dysopes Illig., Prodr. Syst. Mamm. et Av., 1811, p. 122.

1188. **Molossus rufus** E. Geoffroy.

Molossus rufus E. Geoff., Ann. Mus. Hist. Nat. Paris, vi, 1805, p. 155. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M.

Pub., iv, Pt. II, 1904, p. 619, figs. 122, xcix. Zoöl. Ser.

Molossus ursinus Spix, Simiar. et Vespert. Bras., 1823, p. 58, pl. xxv, fig. 4.

Dysopes alecto Temm., Mon. Mamm., i, 1827, p. 231.

Dysopes holosericeus Wagn., Archiv f. Naturg., 1843, p. 368.

Dysopes albus Wagn., Archiv f. Naturg., 1843, p. 368.

RUFIOUS MASTIFF BAT.

Type locality. Paraguay, South America.

Geogr. Distr. Southern Mexico, through Central America to Paraguay and Bolivia.

a. *Molossus rufus obscurus* (E. Geoffroy).

Molossus obscurus E. Geoff., Ann. Mus. Hist. Nat. Paris, vi, 1805, p. 155.

Molossus longicaudatus E. Geoff., Ann. Mus. Hist. Nat. Paris, vi, 1805, p. 155.

Molossus acaticaudatus Desm., Mamm., 1820, p. 116.

Dysopes velox Temm., Mon. Mamm., i, 1827, p. 234.

Molossus fuliginosus Gray, Mag. Zoöl. Bot., II, 1838, p. 501.

Dysopes olivaceo-fuscus Wagn., Abh. Akad. Münch., v, 1844, p. 202.

Dysopes fumarius Burm., Thier. Bras., i, 1854, p. 71.

Molossus aztecus Sauss., Rev. Mag. Zool., 2me Sér., XII, 1860, p. 285.

Molossus rufus obscurus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. II, 1904, p. 620. Zoöl. Ser.

DUSKY MASTIFF BAT.

Type locality. Paraguay.

Geogr. Distr. Guatemala, through Central America to Paraguay and Bolivia, South America. West Indies.

1189. *Molossus coibensis* Allen.

Molossus obscurus Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1904, p. 78. (nec E. Geoffroy.)

Molossus coibensis Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1904, p. 227.

COIBA ISLAND BAT.

Type locality. Coiba Island, Panama.

Geogr. Distr. Restricted to Coiba Island.

Genl. Char. Similar to *M. r. obscurus*, but smaller and blacker, with heavier skull and dentition.

Color. Above deep blackish brown, beneath lighter.

Measurements. Forearm, 34.5-36; first metacarpal, 37.

1190. *Molossus tropidorhynchus* Gray.

Molossus tropidorhynchus Gray, Ann. Nat. Hist., iv, 1839, p. 6.

Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv,

Pt. II, 1904, p. 620. Zool. Ser.

PUG-NOSED MASTIFF BAT.

Type locality. Cuba.

Geogr. Distr. West Indian Islands.

1191. *Molossus nigricans* Miller.

Molossus nigricans Miller, Proc. Acad. Nat. Scien. Phil., 1902,

p. 395. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M.

Pub., iv, Pt. II, 1904, p. 620. Zool. Ser.

BLACKISH MASTIFF BAT.

Type locality. Acaponeta, Territorio de Tepic, Mexico.

Geogr. Distr. Known only from the type locality.

Genus 173. *Promops*.

Promops Gerv., Expéd. du Comte de Castelnau, Zool., Mamm., II, 1855, p. 58, pl. XII, figs. 3, 3a. Type *Promops ursinus* Gervais = *Molossus nasutus* Spix.

1192. *Promops nasutus* (Spix).

Molossus nasutus Spix, Simiar. et Vespert. Bras., 1823, p. 60, pl. xxxv, fig. VII.

Molossus fumarius Spix, Simiar. et Vespert. Bras., 1823, p. 60, pl. xxxv, figs. v, VI.

Promops ursinus Gerv., Expéd. du Com. de Casteln., Zool., Mamm., 1855, p. 59, pl. XII, figs. 3, 3a.

Promops nasutus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 621, fig. c. Zoöl. Ser.

LARGE-NOSED MASTIFF BAT.

Type locality. Near the Rio San Francisco, Brazil.

Geogr. Distr. Guatemala, through Central America to Brazil, South America.

1193. *Promops abrasus* (Temminck).

Dysopes abrasus Temm., Mon. Mamm., I, 1827, p. 232.

Dysopes longimanus Wagn., Archiv f. Naturg., 1843, p. 367.

Dysopes leucopleura Wagn., Archiv f. Naturg., 1843, p. 367.

Promops abrasus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 623. Zoöl. Ser.

SHAVED MASTIFF BAT.

Type locality. Brazil.

Geogr. Distr. Guatemala, through Central America to Brazil, South America.

1194. *Promops orthotis* (H. Allen).

Nyctinomus orthotis H. Allen, Proc. Amer. Philos. Soc., xxvi, 1889, p. 561.

Promops orthotis Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 623. Zoöl. Ser.

STRAIGHT-EARED MASTIFF BAT.

Type locality. Spanishtown, Island of Jamaica.

Geogr. Distr. Known only from the type locality.

1195. *Promops glaucinus* (Wagner).

Dysopes glaucinus Wagn., Archiv f. Naturg., 1843, Pt. I, p. 368.

Molossus ferox Gundl., Monatsb. K. Preuss. Akad. Wiss. Berl., 1861, p. 149.

Promops glaucinus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 624, figs. 123, ci. Zoöl. Ser.

CHESTNUT MASTIFF BAT.

Type locality. Cuyaba, Matto Grosso, Brazil.

Geogr. Distr. Island of Jamaica; in South America to Brazil and Bolivia.

1196. *Promops nanus* Miller.

Promops nanus Miller, Ann. Mag. Nat. Hist., 7th Ser., vi, 1900, p. 470. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. II, 1904, p. 624. Zoöl. Ser.

DWARF MASTIFF BAT.

Type locality. Bogava, Chiriqui, Panama.

Geogr. Distr. Panama, Central America. Range unknown.

1197. *Promops californicus* (Merriam).

Molossus californicus Merr., N. Am. Faun., No. 4, 1890, p. 31.

Promops californicus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 421, fig. 94. Zoöl. Ser.

CALIFORNIA MASTIFF BAT.

Type locality. Alhambra, Los Angeles County, California.

Geogr. Distr. Southern California. Range unknown.

Genus 174. *Nyctinomops*.***Nyctinomops* Miller, Proc. Acad. Nat. Scien. Phil., 1902, p. 393.**

Type Nyctinomus femorosaccus Merriam.

1198. *Nyctinomops femorosaccus* (Merriam).

Nyctinomus femorosaccus Merr., N. Am. Faun., No. 2, 1889, p. 23. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 417. Zoöl. Ser.

Nyctinomops femorosaccus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. II, 1904, p. 626. Zoöl. Ser.

PALM SPRINGS FREE-TAILED BAT.

Type locality. Agua Caliente, Palm Springs, Riverside County, California.

Geogr. Distr. Southern California to Ensinada, Lower California, Mexico.

1199. *Nyctinomops yucatanicus* Miller.

Nyctinomops yucatanicus Miller, Proc. Acad. Nat. Scien. Phil., 1902, p. 393. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. II, 1904, p. 626, fig. 124. Zoöl. Ser.

YUCATAN FREE-TAILED BAT.

Type locality. Chichen Itza, Yucatan.

Geogr. Distr. Yucatan. Range unknown.

1200. *Nyctinomops gracilis* (Wagner).

Dysopes gracilis Wagn., Archiv f. Naturg., 1843, p. 368.

Nyctinomops gracilis Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. II, 1904, p. 626. Zoöl. Ser.

GRACEFUL FREE-TAILED BAT.

Type locality. Cuyaba, Matto Grosso, Brazil.

Geogr. Distr. Guatemala, through Central America to Brazil, South America.

1201. *Nyctinomops depressus* (Ward).

Nyctinomus depressus Ward, Amer. Nat., xxv, 1891, p. 747, fig.

Nyctinomus macrotis nevadensis H. Allen, Mon. N. Am. Bats, 1894, p. 171. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 417. Zoöl. Ser.

Nyctinomops depressus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 627. Zoöl. Ser.

TUCUBAYA FREE-TAILED BAT.

Type locality. Tucubaya, Federal District, Mexico.

Geogr. Distr. Known only from the type locality.

1202. *Nyctinomops macrotis* (Gray).

Nyctinomus macrotis Gray, Ann. Nat. Hist., IV, 1840, p. 5.

Dysopes auritus Wagn., Archiv f. Naturg., 1843, p. 368.

Dysopes laticaudatus Rengg., Säugeth. Parag., 1830, p. 88.

Dysopes aurispinosus Peale, U. S. Expl. Exp., VIII, 1848, p. 21.

Nyctinomops macrotis Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 627. Zoöl. Ser.

CUBAN FREE-TAILED BAT.

Type locality. Cuba.

Geogr. Distr. Islands of Cuba and Jamaica, West Indies to Brazil and Paraguay.

Genus 175. *Nyctinomus*.

Nyctinomus E. Geoff., Desc. Égypte, Mamm., II, 1813, pp. 114, 128-130, pl. 2, No. 2. Type *Nyctinomus ægyptiacus* E. Geoffroy.

Dinops Savi, Nouv. Giorn. de Lett., 1825, p. 230.

Dysopes Cretzsch., Rüpp. Atl. Reis. Nordl. Afr., Zool., 1826, p. 69. (nec Illig.)

1203. *Nyctinomus antillularum* Miller.

Nyctinomus antillularum Miller, Proc. Acad. Nat. Scien. Phil., 1902, p. 398. Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., IV, Pt. II, 1904, p. 629. Zoöl. Ser.

DOMINICAN FREE-TAILED BAT.

Type locality. Roseau, Island of Dominica.

Geogr. Distr. Islands of Dominica, Montserrat, St. Kitts and St. Lucia, West Indies.

1204. *Nyctinomus mexicanus* Saussure.

Nyctinomus mexicanus Sauss., Rev. Mag. Zool., 2me Sér., XII, 1860, p. 283. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 629, fig. 125. Zoöl. Ser.

Nyctinomus mohavensis Merr., N. Am. Faun., No. 2, 1889, p. 25. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 418. Zoöl. Ser.

?*Nyctinomus brasiliensis californicus* H. Allen, Mon. N. Am. Bats, 1894, p. 166.

MEXICAN FREE-TAILED BAT.

Type locality. Cofre de Perote, State of Vera Cruz, Mexico. Altitude, 13,000 feet.

Geogr. Distr. Volcano of Popocatepetl, through northern Mexico and the United States. Lower California.

1205. **Nyctinomus cynocephalus* (Le Conte).

Nycticea cynocephala Le Conte, McMurtrie, Cuv. Anim. Kingd., I, 1831, p. 432.

Nyctinomus cynocephalus Bangs, Proc. Bost. Soc. Nat. Hist., 1898, p. 218.

DOG-HEADED FREE-TAILED BAT.

Type locality. Probably the Le Conte plantation near Riceboro, Liberty County, Georgia.

Geogr. Distr. Southeastern United States. Range unknown.

Genl. Char. Size moderately large. Color without any red tint.

Color. Above dull brown; of a grayish hue in some lights; beneath paler.

Measurements. Total length, 97-102; forearm, 38-41; longest finger, 80-84.

1206. *Nyctinomus bahamensis* Rehn.

Nyctinomus bahamensis Rehn, Proc. Acad. Nat. Scien. Phil., 1902, p. 641. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 630. Zoöl. Ser.

BAHAMAN FREE-TAILED BAT.

Type locality. Governor's Harbor, Eleuthera Island, Bahamas.

Geogr. Distr. Eleuthera, Little and Great Abaco, Bahama Islands.

1207. *Nyctinomus musculus* Gundlach.

Nyctinomus musculus Gundl., Monatsb., K. Preuss. Akad. Wiss. Berl., 1861, p. 149. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 630. Zoöl. Ser.

*While this may be distinct from *N. brasiliensis* Geoff., why is it not the same as *N. mexicanus* Saussure?

FREE-TAILED MOUSE BAT.

Type locality. Cuba.

Geogr. Distr. Islands of Cuba, Porto Rico, St. Kitts, and Trinidad.

1208. *Nyctinomus minutus* Miller.

**Nyctinomus brasiliensis* Chapman, Bull. Am. Mus. Nat. Hist. N. Y., 1892, p. 316. (Part.)

Nyctinomus minutus Miller, Bull. Am. Mus. Nat. Hist. N. Y., 1899, p. 173. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 630. Zool. Ser.

LITTLE FREE-TAILED BAT.

Type locality. Trinidad, Island of Cuba.

Geogr. Distr. Island of Cuba.

Family IV. **Thyropteridæ. Funnell-eared Bats.**Genus 176. **Spectrellum.**

Spectrellum Gerv., Expéd. du Comte de Casteln. l'Amér. Sud, Mamm., Zool., 1855, p. 51. Type *Spectrellum macrurum* Gervais.

†*Natalus* (misprint for *Natalis*) Gray, Mag. Zool. Bot., II, 1838, p. 496. Type *Natalus! stramineus* Gray. (nec Lap., Coleopt., 1836, *Natalis*.)

1209. *Spectrellum stramineum* (Gray).

Natalus! stramineus Gray, Mag. Zool. Bot., II, 1838, p. 496.

Spectrellum macrurum Gerv., Compt. Rend. Acad. Scien. Paris, 1856, p. 347.

Natalis stramineus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 632, figs. 126, cII. Zool. Ser.

STRAW-COLORED BAT.

Type locality. Unknown.

Geogr. Distr. Island of Dominica, West Indies; Guatemala, Central America, to Brazil, South America.

a. *Spectrellum stramineum major* (Miller).

Natalus! major Miller, Proc. Acad. Nat. Scien. Phil., 1902, p. 398.

Natalis stramineus major Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 633. Zool. Ser.

**Nyctinomus brasiliensis* Is. Geoffroy does not seem to be found beyond the boundaries of South America.

† The genus *Natalus* Gray, 1838, being preoccupied by *Natalis* Lap., 1836, necessitates a new name for the Family, which, by adopting the oldest proposed genus **THYROPTERA**, according to the rule governing such cases, would be **THYROPTERIDÆ**.

GREAT BAT.

Type locality. Savaneta, San Domingo, West Indies.

Geogr. Distr. Island of Haiti.

1210. Spectrellum mexicanum (Miller).

Natalus! mexicanus Miller, Proc. Acad. Nat. Scien. Phil., 1902, p. 399.

Natalis mexicanus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 633. Zoöl. Ser.

SANTA ANITA BAT.

Type locality. Santa Anita, Lower California, Mexico.

Geogr. Distr. Lower California, Mexico.

Genus 177. **Nyctiellus.**

Nyctiellus Gerv., Expéd. du Comte de Castelnau Amér. Sud., Mamm., 1855, p. 84, pl. xv, fig. 6. Type *Vespertilio lepidus* Gervais.

1211. Nyctiellus lepidus (Gervais).

Vespertilio lepidus Gerv., in Ramon de la Sagra's Hist. de l'Isle de Cuba, Mamm., 1838, p. 22, pl. I, figs. 1-4.

Vespertilio barbatus Gundl., Wiegmann Archiv f. Naturg., 1840, p. 356.

Nyctiellus lepidus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 634, fig. 127. Zoöl. Ser.

GRACEFUL BAT.

Type locality. Cuba.

Geogr. Distr. Island of Cuba; Isle of Pines.

Genus 178. **Chilonatalis.**

Chilonatalus (sic) Miller, Proc. Acad. Nat. Scien. Phil., 1898, p. 326. Type *Natalus! micropus* Dobson.

1212. Chilonatalis micropus (Dobson).

Natalus! micropus Dobson, Proc. Zoöl. Soc., 1880, p. 443.

Chilonatalis micropus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 635. Zoöl. Ser.

SMALL-FOOTED BAT.

Type locality. Environs of Kingston, Island of Jamaica.

Geogr. Distr. Known only from the type locality.

1213. Chilonatalis brevimanus (Miller).

Natalus! brevimanus Miller, Proc. Acad. Nat. Scien. Phil., 1898, p. 328.

Chilonatalis brevimanus Elliot, Mamm. Middle Amer. & W. Indies,
F. C. M. Pub., IV, Pt. II, 1904, p. 636, figs. 128, CIII. Zoöl. Ser.

SHORT-HANDED BAT.

Type locality. Old Providence Island, Caribbean Sea.

Geogr. Distr. Known only from the type locality.

1214. *Chilonatalis tumidifrons* Miller.

Chilonatalus! tumidifrons Miller, Proc. Biol. Soc. Wash., XVI, 1903,
p. 119.

Chilonatalis tumidifrons Elliot, Mamm. Middle Amer. & W. Indies,
F. C. M. Pub., IV, Pt. II, 1904, p. 637. Zoöl. Ser.

WATLING'S ISLAND BAT.

Type locality. Cave near Sandy Point, at the southwest corner
of Watling's Island, Bahamas.

Geogr. Distr. Watling's and Great Abaco Islands, Bahamas.

Genus 179. **Thyroptera.**

Thyroptera Spix, Simiar. et Vespert. Brasil., 1823, p. 61, tab. xxxvi,
fig. 9. *Type Thyroptera tricolor* Spix.

Hyonycteris Licht. & Peters, Monatsb. K. Preuss. Akad. Wiss.
Berl., 1854, p. 335.

1215. **Thyroptera discifera** (Lichtenstein & Peters).

Hyonycteris discifera Licht. & Peters, Monatsb. K. Preuss. Akad.
Wiss. Berl., 1854, p. 335.

Thyroptera discifera Elliot, Mamm. Middle Amer. & W. Indies,
F. C. M. Pub., IV, Pt. II, 1904, p. 637, figs. 129, CIV. Zoöl. Ser.

DISK-BEARING BAT.

Type locality. Puerto Caballos, District of Cortez, Honduras,
Central America.

Geogr. Distr. Honduras, Central America, south into South
America.

Family V. **Phyllostomatidae. Vampire Bats.**

Subfamily I. **Mormopinæ.**

Genus 180. **Chilonycteris.**

Chilonycteris Gray, Ann. Nat. Hist., IV, 1839, p. 4. *Type Chilonyc-*
teris macleayii Gray.

Lobostoma Gundl., Wieg. Archiv f. Naturg., VI, 1840, p. 356.

Phyllodia Gray, Proc. Zoöl. Soc., 1843, p. 50.

1216. *Chilonycteris macleayi* Gray.

Chilonycteris macleayi Gray, Ann. Mag. Nat. Hist., iv, 1839, p. 5, pl. 1, fig. 2. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. II, 1904, p. 641, fig. cv. Zoöl. Ser.

Lobostoma quadridens Gundl., Wieg. Archiv f. Naturg., vi, 1840, p. 357.

MACLEAY'S BAT.

Type locality. Cuba.

Geogr. Distr. Islands of Cuba, Haiti, and Jamaica, West Indies.

a. *Chilonycteris macleayi inflata* Rehn.

Chilonycteris macleayi Gundl., Ann. Soc. Espan. Hist. Nat. vii, 1878, Cuad. 1, p. 140. (nec Gray.)

Chilonycteris macleayi inflata Rehn, Proc. Acad. Nat. Scien. Phil., 1904, p. 190. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. II, 1904, Append., p. 747. Zoöl. Ser.

INFLATED-NOSE BAT.

Type locality. Cueva di Fari, near Pueblo Viejo, Porto Rico.

Geogr. Distr. Island of Porto Rico.

b. *Chilonycteris macleayi grisea* (Gosse).

Chilonycteris grisea Gosse, Nat. Sojourn in Jamaica, 1851, p. 326, pl. vi, fig. 1.

Chilonycteris quadridens Tomes, Proc. Zoöl. Soc., 1861, p. 65. (nec Gundlach.)

Chilonycteris macleayi grisea Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. II, 1904, Append., p. 747. Zoöl. Ser.

GRAY BAT.

Type locality. Phoenix Park, St. Ann's Parish, Jamaica.

Geogr. Distr. Island of Jamaica.

c. *Chilonycteris macleayi fuliginosa* Gray.

Chilonycteris fuliginosa Gray, Proc. Zoöl. Soc., 1843, p. 20.

Chilonycteris macleayi Peters, Monatsb. K. Preuss. Akad. Wiss. Berl., 1872, p. 360. (Part.)

Chilonycteris macleayi fuliginosa Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. II, 1904, Append., p. 748. Zoöl. Ser.

DUSKY BAT.

Type locality. Port au Prince, Haiti.

Geogr. Distr. Island of Haiti.

1217. *Chilonycteris personata* Wagner.

Chilonycteris personata Wagn., Archiv f. Naturg., ix, 1843, p. 367. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., iv, Pt. II, 1904, p. 641. Zoöl. Ser.

MASKED BAT.

Type locality. Matto Grosso, Brazil.

Geogr. Distr. Guatemala, Central America, to Brazil, South America.

1218. *Chilonycteris psilotis* Dobson.

Chilonycteris psilotis Dobson, Cat. Chirop. Brit. Mus., 1878, p. 451, pl. xxiii, fig. 2. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 642. Zoöl. Ser.

SHAGGY-EARED BAT.

Type locality. Unknown.

Geogr. Distr. Isthmus of Tehuantepec, Mexico. Range unknown.

1219. *Chilonycteris parnelli* (Gray).

Phyllodia parnellii Gray, Proc. Zoöl. Soc., 1843, p. 50.

Chilonycteris osburnii Tomes, Proc. Zoöl. Soc., 1861, p. 66, pl. xiii.

Chilonycteris parnelli Elliot, Mamm. Middle Amer. & W. Indies. F. C. M. Pub., IV, Pt. II, 1904, p. 642. Zoöl. Ser.

PARNELL'S BAT.

Type locality. Island of Jamaica

Geogr. Distr. Island of Jamaica.

1220. *Chilonycteris boothi* Gundlach.

Chilonycteris boothi Gundl., Monatsb. K. Preuss. Akad. Wiss. Berl., 1861, p. 154. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 642, fig. 130. Zoöl. Ser.

BOOTH'S BAT.

Type locality. Fundador, Island of Cuba.

Geogr. Distr. Island of Cuba.

1221. *Chilonycteris portoricensis* Miller.

Chilonycteris portoricensis Miller, Proc. Acad. Nat. Scien. Phil., 1902, p. 400. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 643. Zoöl. Ser.

PORTO RICAN BAT.

Type locality. Cave near Pueblo Viejo, Island of Porto Rico.

Geogr. Distr. Island of Porto Rico.

1222. *Chilonycteris rubiginosa* Wagner.

Chilonycteris rubiginosa Wagn., Archiv f. Naturg., ix, bd. I, 1843, p. 367. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 643. Zoöl. Ser.

DARK BROWN BAT.

Type locality. Caicara, Brazil.

Geogr. Distr. States of Vera Cruz and Michoacan, Mexico, south to Brazil.

1223. *Chilonycteris mexicana* Miller.

Chilonycteris mexicana Miller, Proc. Acad. Nat. Scien. Phil., 1902, p. 401. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 644. Zoöl. Ser.

MEXICAN BAT.

Type locality. San Blas, Territorio de Tepic, Mexico.

Geogr. Distr. Central and southern Mexico from State of Durango through Province of Tepic to State of Oaxaca in the south.

Genus 181. *Dermonotus*.

Dermonotus Gill, Proc. Biol. Soc. Wash., XIV, 1901, p. 177. Type *Pteronotus davyi* Gray.

Pteronotus Gray, Mag. Zoöl. Bot., II, 1838, p. 500. (nec Rafinesque.)

1224. *Dermonotus davyi* (Gray).

Pteronotus davyi Gray, Mag. Zoöl. Bot., II, 1838, p. 500.

Dermonotus davyi Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 644, figs. 131, CVI. Zoöl. Ser.

DAVY'S BAT.

Type locality. Island of Trinidad.

Geogr. Distr. Islands of Dominica and Trinidad, south to Brazil.

a. *Dermonotus davyi fulvus* (Thomas).

Chilonycteris davyi fulvus Thomas, Ann. Mag. Nat. Hist., 6th Ser., X, 1892, p. 410.

Dermonotus davyi fulvus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 646. Zoöl. Ser.

LAS PEÑAS BAT.

Type locality. Las Peñas, State of Jalisco, Mexico.

Geogr. Distr. State of Jalisco, Mexico. Range unknown.

Genus 182. *Mormops*.

Mormoops (*sic*) Leach, Trans. Linn. Soc., XIII, 1820, p. 76. Type *Mormoops* (*sic*) *blainvillii* Leach.

1225. *Mormops megalophylla* Peters.

Mormops blainvillii Peters, Abhandl. Akad. Berl., 1856, p. 289, pl. I, figs. 1-5. (nec Leach.)

Mormops megalophylla Peters, Monatsb. K. Preuss. Akad. Wiss. Berl., 1864, p. 381. Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 418, fig. 92. Zoöl. Ser. *Id.* Mamm. Middle

Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 648, figs. 132, CVII. Zoöl. Ser.

BIG-EARED BAT.

Type locality. Mexico.

Geogr. Distr. Islands of Cuba and Jamaica; eastern and southern Mexico to Brazil.

a. Mormops megalophylla senicula (Rehn).

Mormoops! megalophylla senicula Rehn, Proc. Acad. Nat. Scien. Phil., 1902, p. 169.

Mormops megalophylla senicula Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 648. Zoöl. Ser.

ANTIQUE BAT.

Type locality. Fort Clark, Kinney County, Texas.

Geogr. Distr. Southern Texas, into northern and central Mexico, to State of Morelos.

1226. Mormops blainvillii (Leach).

Mormoops! blainvillii Leach, Trans. Linn. Soc., XIII, 1820, p. 77, pl. VII, figs. 1-4.

Mormops blainvillii Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 649. Zoöl. Ser.

DE BLAINVILLE'S BAT.

Type locality. Island of Jamaica.

Geogr. Distr. Islands of Cuba and Jamaica.

a. Mormops blainvillii cinnamomea (Gundlach).

Lobostoma cinnamomeum Gundl., Wieg. Archiv f. Naturg., VI, 1840, p. 357.

Mormops blainvillii cinnamomea Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 649. Zoöl. Ser.

CINNAMON BAT.

Type locality. Cafetal San Antonio el Fundador, Cuba.

Geogr. Distr. Islands of Cuba, Haiti, and Mona.

Subfamily II. **Phyllostomatinae.**

Genus 183. **Lonchorina.**

Lonchorina Tomes, Proc. Zoöl. Soc., 1863, p. 81, pl. 22. Type *Lonchorina aurita* Tomes.

1227. *Lonchorina aurita* Tomes.

Lonchorina aurita Tomes, Proc. Zool. Soc., 1863, p. 83. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 650, figs. 133, CVIII. Zool. Ser.

TOMES' LONG-EARED BAT.

Type locality. West Indies. Islands unknown.

Geogr. Distr. Unknown.

Genus 184. **Otopterus.**

Otopterus Lydekker, in Flow. & Lydek. Mamm. Living and Extinct, 1891, p. 673. Type *Macrotus waterhousii* Gray.

Macrotus Gray, Proc. Zool. Soc., 1843, p. 21. (nec Leach, Vespertil., 1816; nec Dejean, Coleopt., 1833; nec Reid, Marsupialia, 1836.)

1228. *Otopterus waterhousii* (Gray).

Macrotus waterhousii Gray, Proc. Zool. Soc., 1843, p. 21.

Otopterus waterhousii Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 652, figs. 134, CIX. Zool. Ser.

WATERHOUSE'S LARGE-EARED BAT.

Type locality. Island of Haiti.

Geogr. Distr. Island of Haiti, West Indies.

a. Otopterus waterhousii jamaicensis (Rehn).

Macrotus waterhousii jamaicensis Rehn, Proc. Acad. Nat. Scien. Phil., 1904, p. 432.

JAMAICA LARGE-EARED BAT.

Type locality. Spanishtown, Jamaica.

Geogr. Distr. Island of Jamaica.

Genl. Char. Similar to *O. waterhousii*, but color duller, skull smaller and narrower, and weaker upper tooth row.

Color. Above Prout's brown, basal half of hairs white; beneath drab and wood brown, basal half of hairs white; ears and uropatagium Prout's brown; endopatagium, mesopatagium, and ectopatagium clove brown.

Measurements. Total length, 90.2-103; forearm, 52-54.5; thumb, 12-15; third finger, 88-94; tibia, 21.5-24; calcaneum, 9.2-12; foot, 12.2-16; tail, 27-36; ear, 23-29; tragus, 8-9.5. Skull; total length, 24.2-24.5; greatest zygomatic width, 12-12.2; interorbital width, 4.1-4.3; height of braincase, 9-9.1; width of braincase above roots of zygomata, 9.4-9.5; length of palate, 9.3-11; width of palate, 7.8-8; length of mandible, 16-16.5.

b. Otopterus waterhousii compressus (Rehn).

Macrotus waterhousii compressus Rehn, Proc. Acad. Nat. Scien. Phil., 1904, p. 434.

BAHAMA LONG-EARED BAT.

Type locality. Eleuthera, Bahamas.

Geogr. Distr. Eleuthera, Providence, Long Island, and Conch Sound, Andros, Bahamas.

Genl. Char. Rostrum narrower anteriorly than the interorbital region.

Color. Above Prout's brown, the hairs with more than basal half white; base of ears and along the internal margins whitish; beneath hairs white, basally ringed with pale drab, and tipped with silvery white; membranes sepia, ears and uropatagium paler.

Measurements. Total length, 94; forearm, 52; thumb, 13.5; third finger, 85; tibia, 21; calcaneum, 11; foot, 13; tail, 25; ear, 24; tragus, 8.5. Skull: total length, 25-25.1; greatest zygomatic width, 10.6-12; interorbital width, 4.2-4.5; height of braincase, 8-8.6; width of braincase, above roots of zygomata, 8.5-9.5; length of palate, 10.5-11; width of palate, 7.5-7.6; greatest length of mandible, 16.8-17.

c. Otopterus waterhousii minor (Gundlach).

Macrotus minor Gundl., Monatsb. K. Preuss. Akad. Wiss. Berl., 1864, p. 382.

CUBAN LARGE-EARED BAT.

Type locality. Cuba.

Geogr. Distr. Cuba and Isle of Pines.

Genl. Char. Size smaller than *O. waterhousii*, color darker.

Color. Two phases. *Brown phase.* Above seal brown, with over half of the basal portion white; between ears and posterior to the connecting membrane blackish; beneath, hairs white basally, then annulated with drab, and tipped with silvery; membranes blackish brown.

Red phase. Above and beneath dull ferruginous unicolor; patch between ears blackish; membranes black.

Measurements. Total length, 84.5-96.5; forearm, 49-54; thumb, 12.5-15; third finger, 81-88; tibia, 19-22.5; calcaneum, 8.5-12.3; foot, 10.5-15.5; tail, 20-37; ear, 21.5-30; tragus, 7.8-10. Skull: total length, 23.5-24.3; greatest zygomatic width, 11-12; interorbital width, 3.5-4.1; height of braincase, 8-8.8; width of braincase above roots of zygomata, 9-9.2; length of palate, 9.2-10; width of palate, 7-8; length of mandible, 15-16.

1229. *Otopterus californicus* (Baird).

Macrotus californicus Baird, Proc. Acad. Nat. Scien. Phil., 1858, p. 116.

Otopterus californicus Elliot, Syn. N. Am. Mamm., F. C. M. Pub., II, 1901, p. 420. Zoöl. Ser. *Id.* Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 653. Zoöl. Ser.

CALIFORNIA LARGE-EARED BAT.

Type locality. Old Fort Yuma, San Diego County, California.

Geogr. Distr. Southern California into Lower California, Mexico.

1230. *Otopterus mexicanus* (Saussure).

Macrotus mexicanus Sauss., Rev. Mag. Zool. 2me Sér., XII, 1860, p. 486.

Otopterus mexicanus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 653. Zoöl. Ser.

SAUSSURE'S LARGE-EARED BAT.

Type locality. District of Yautepec, near Cuautla, State of Morelos, Mexico.

Geogr. Distr. Tres Marias Islands, State of Jalisco, south to States of Mexico and Oaxaca, Mexico.

1231. **Otopterus bocourtianus* (Dobson).

Macrotus bocourtianus Dobson, Ann. Mag. Nat. Hist., 4th Ser., 1876, p. 436.

Macrotus mexicanus J. A. Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1891, p. 179.

Otopterus bocourtianus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 654. Zoöl. Ser.

BOCOURT'S LARGE-EARED BAT.

Type locality. Vera Paz, Guatemala.

Geogr. Distr. Southern Mexico and Guatemala.

1232. *Otopterus bulleri* H. Allen.

Macrotus bulleri H. Allen, Proc. Am. Phil. Soc., xxviii, 1890, p. 73. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 654. Zoöl. Ser.

*Mr. J. A. G. Rehn in his Monograph of the genus *Otopterus*, Proc. Acad. Nat. Scien. Phil., 1904, p. 437 (erroneously called by him *Macrotus*), makes Dobson's species a synonym of *O. mexicanus*, and retains *O. bulleri* as a race of the last named. Dr. J. A. Allen, in a "Note on Mexican Species of *Otopterus*," Bull. Am. Mus. Nat. Hist. N. Y., 1904, p. 235, after a careful examination, decides that *O. bocourtianus* is a good species, and that *O. bulleri* should become a synonym of *O. mexicanus*. Here we have two diametrically opposed opinions entertained by two naturalists working independently of each other, and in view of the *impasse* thus created, it seems best to consider the two forms distinct species, as they have heretofore been by most investigators, especially as the specimens in this Museum appear to sustain that view of the case. Although not a matter of supreme importance, it will, however, probably require, after the decisions mentioned above, a consensus of opinions among mammalogists before the question at issue shall be satisfactorily settled.

BULLER'S LARGE-EARED BAT.

Type locality. Bolaños, State of Jalisco, Mexico.

Geogr. Distr. State of Guadalajara to States of Durango and Chihuahua, west to the Tres Marias Islands, State of Jalisco.

1233. *Otopterus pygmæus* (Rehn).

Macrotus pygmæus Rehn, Proc. Acad. Nat. Scien. Phil., 1904, p. 444.

LITTLE LARGE-EARED BAT.

Type locality. Izamal, Yucatan.

Geogr. Distr. Known only from the type locality.

Genl. Char. Size very small; skull with broad and flat interorbital region; teeth large; ear rounded, internal margin arcuate; tragus lanceolate, tip blunt, slight basal accessory longitudinal lobule on external margin; nose leaf high, lanceolate; width of nose pad twice that of its height; nostrils slightly crescentic; shallow median depression on the triangular pad on chin.

Color. Above mars brown, basally grayish white; beneath écru drab, unicolor; wing membrane clove brown.

Measurements. Total length, 79.3; forearm, 35.5; thumb, 10; third finger, 65.5; tibia, 14.9; calcaneum, 9; foot, 10.5; tail, 28; ear, 17.2; tragus, 7. Skull: greatest zygomatic width, 9.2; interorbital width, 4.2; height of braincase, 7.6; width of braincase above the roots of zygomata, 8; length of palate, 8; width of palate, 6.5; length of mandible, 12.8.

Genus 185. **Vampyrus.**

Vampyrus Leach, Trans. Linn. Soc., XIII, 1821, p. 79. Type *Vespertilio spectrum* Linnæus.

1234. *Vampyrus spectrum* (Linnæus).

Vespertilio spectrum Linn., Syst. Nat. I, 1758, p. 31; I, 1766, p. 46.

Vampyrus spectrum Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., IV, Pt. II, 1904, p. 656, fig. 135. Zool. Ser.

SPECTER BAT.

Type locality. South America.

Geogr. Distr. Guatemala, Central America to Brazil, South America; Island of Jamaica.

Genus 186. **Chropterus.**

Chropterus Peters, Monatsb. K. Preuss. Akad. Wiss. Berl., 1865, p. 505. Type *Vampyrus auritus* Peters.

1235. *Chrotopterus auritus* (Peters).

Vampyrus auritus Peters, Abhandl. K. Preuss. Akad. Wiss. Berl., 1856, p. 415, pl. II, figs. 1-5.

Chrotopterus auritus Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., IV, Pt. II, 1904, p. 656, figs. 136, cx. Zoöl. Ser.

PETERS' VAMPIRE BAT.

Type locality. Mexico.

Geogr. Distr. Eastern Mexico into South America to southern Brazil.

Genus 187. **Tonatia.**

Tonatia Gray, in Griff., Cuv. Anim. Kingd., v, 1827, p. 71 (footnote).

Type Vampyrus bidens Spix.

Lophostoma D'Orbigny, Voy. Amér. Mérid., 1847, p. II.

1236. *Tonatia amblyotis* Wagner.

Phyllostoma amblyotis Wagn., Archiv f. Naturg., 1843, p. 365.

Tonatia amblyotis Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., IV, Pt. II, 1904, p. 658, figs. 137, cxI. Zoöl. Ser.

ROUND-EARED BAT.

Type locality. Matto Grosso, Brazil.

Geogr. Distr. Bogava, Chiriqui, Panama to Brazil, South America.

Genus 188. **Micronycteris.**

Micronycteris Gray, Proc. Zoöl. Soc., 1866, p. 113. *Type Phyllorhina megalotis* Gray.

Schizostoma Gerv., Expéd. du Comt. de Casteln. Amér. Sud., Zool., 1855, p. 49. (nec Bronn, Mollusca, 1835.)

Vampyrella Reinh., Vidensk. Meddels. Naturh. Foren. Kjöbenh., 3 Aartis, IV, 1872, p. 111.

1237. *Micronycteris hirsutus* (Peters).

Schizostoma hirsutum Peters, Monatsb. K. Preuss. Akad. Wiss. Berl., 1869, p. 396.

Micronycteris hirsutus Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., IV, Pt. II, 1904, p. 661. Zoöl. Ser.

HAIRY BAT.

Type locality. Unknown.

Geogr. Distr. Pozo Azul, Costa Rica. Altitude, 200 meters. Range unknown.

1238. *Micronycteris megalotis* (Gray).

Phyllophora megalotis Gray, Ann. Mag. Nat. Hist., 1st Ser., x, 1842, p. 257.

Phyllostoma elongata Gray, Ann. Mag. Nat. Hist., 1st Ser., x, 1842, p. 257. (nec Geoffroy.)

Phyllostoma scrobiculatum Wagn., Schreb. Säugth., Suppl., v, 1855, p. 627.

Micronycteris megalotis Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 662, figs. 138, CXII. Zool. Ser.

LARGE-EARED VAMPIRE BAT.

Type locality. "Brazil."

Geogr. Distr. Mexico through Central America to Brazil, South America.

***a. Micronycteris megalotis mexicanus* Miller.**

Micronycteris megalotis mexicanus Miller, Proc. Acad. Nat. Scien. Phil., 1898, p. 329. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 662. Zool. Ser.

MEXICAN VAMPIRE BAT.

Type locality. Plantinar, State of Jalisco, Mexico.

Geogr. Distr. State of Jalisco, western Mexico. Range unknown.

1239. *Micronycteris microtis* Miller.

Micronycteris microtis Miller, Proc. Acad. Nat. Scien. Phil., 1898, p. 328. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 663. Zool. Ser.

SMALL-EARED NICARAGUAN BAT.

Type locality. Greytown, Nicaragua.

Geogr. Distr. Nicaragua, Central America. Range unknown.

Genus 189. *Glyphonycteris*.

Glyphonycteris Thomas, Ann. Mag. Nat. Hist., 6th Ser., XVIII, 1896, p. 302. Type *Glyphonycteris sylvestris* Thomas.

1240. *Glyphonycteris sylvestris* Thomas.

Glyphonycteris sylvestris Thomas, Ann. Mag. Nat. Hist., 6th Ser., XVIII, 1896, p. 303. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 664. Zool. Ser.

FOREST BAT.

Type locality. Imravallas, Costa Rica.

Geogr. Distr. Costa Rica, Central America. Range unknown.

Genus 190. **Trachyops.**

Trachops (*sic*, misprint) Gray, Proc. Zool. Soc., 1847, p. 14. Type
Trachops! fuliginosus Gray = *Vampyrus cirrhosus* Spix.

1241. Trachyops cirrhosus (Spix).

Vampyrus cirrhosus Spix, Simiar. et Vespert. Bras., 1823, p. 64,
pl. xxxvi, fig. iii.

Trachops (sic) fuliginosus Gray, Proc. Zool. Soc., 1847, p. 14.

Tylostoma mexicana Sauss., Rev. Mag. Zool., xii, 1860, p. 484.

Trachyops cirrhosus Elliot, Mamm. Middle Amer. & W. Indies,
F. C. M. Pub., iv, Pt. II, 1904, p. 665, figs. 139, cxiii. Zool.
Ser.

FRINGED-FACED BAT.

Type locality. Brazil.

Geogr. Distr. Mexico to Brazil, South America.

Genus 191. **Phyllostoma.**

Phyllostomus Lacép., Tabl. Div. Sous-div. Ordres et Genres des
Mamm., 1799, p. 16. Type *Vespertilio hastatus* Pallas.

Phyllostoma Geoff., Ann. Mus. Hist. Nat. Paris, xv, 1810, p. 174.

1242. Phyllostoma hastatum (Pallas).

Vespertilio hastatus Pall., Spicil. Zool., fasc. iii, 1767, p. 7.

Phyllostomus maximus Max., Reise Bras., II, 1820-21, p. 242.

Phyllostoma hastatum Elliot, Mamm. Middle Amer. & W. Indies,
F. C. M. Pub., iv, Pt. II, 1904, p. 666, figs. 140, cxiv. Zool.
Ser.

SPEAR-NOSED BAT.

Type locality. Unknown.

Geogr. Distr. Panama, Central America to Brazil.

a. Phyllostoma hastatum panamensis (Allen).

Phyllostomus hastatus panamensis Allen, Bull. Am. Mus. Nat.
Hist. N. Y., 1904, p. 233.

PANAMA SPEAR-NOSED BAT.

Type locality. Boqueron, Chiriqui, Panama.

Geogr. Distr. Panama. Range unknown.

Genl. Char. Similar to *P. hastatum*, but larger.

Color. Above dark seal brown, slightly varied with gray on
shoulders, sides of neck, and front of head; beneath light brown
washed with grayish; ears, nose-leaf, and membranes blackish brown.

Measurements. Forearm, 90; third finger metacarpal, 83; tibia, 34; foot, 24; calcaneum, 24. Skull: total length, 39; basal length, 30; zygomatic breadth, 21.2; mastoid breadth, 19; postorbital constriction, 7.5; width of rostrum at base of canines, 9.5; length of upper tooth row, 14.3.

Genus 192. **Hemiderma.**

Hemiderma Gerv., Expéd. du Comt. de Castelnau Amér. Sud., Mamm., Zool., 1855, p. 43. Type *Phyllostoma brevicaudum* Maximilian. *Carollia* Gray, Mag. Zoöl. & Bot., II, 1838, p. 488. (nec Cantraine, Mollusca, 1837.)

1243. Hemiderma perspicillatum (Linnæus).

Vespertilio perspicillatus Linn., Syst. Nat., I, 1758, p. 31; I, 1766, p. 47.

Phyllostoma brevicaudum Max., Schinz's Thierr., I, 1821, p. 164.

Vampyrus soricinus Spix, Simiar. et Vespert. Bras., 1823, p. 66, pl. xxxvi.

Phyllostoma grayi Waterh., Voy. Beagle, Mamm., *1838, p. 20, pl. VIII.

Phyllostoma lanceolatum Temm., Gray's List. Spec. Mamm. Brit. Mus., 1843, p. 20.

Carollia verrucata Gray, Voy. "Sulphur," Mamm., 1844, p. 20, pl. VIII.

Phyllostoma bicolor Wagn., Schreb. Säugth., Suppl., I, 1844, p. 400; v, 1855, p. 626.

Phyllostoma calcaratum Wagn., Abhandl. Akad. Münch., v, p. 168.

Phyllostoma brachyotum Burm., Thiere Brasil, 1854, p. 46.

Hemiderma brevicaudum Gerv., Expéd. du Comt. de Castelnau Amér. Sud., Zool., 1855, p. 43, pl. VII.

Carollia azteca Sauss., Rev. Mag. Zool., 2me Sér., XII, 1860, p. 480, pl. xx, figs. 1, 1a.

Rhinops minor Gray, Proc. Zoöl. Soc., 1866, p. 115.

Hemiderma perspicillatum Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 668, figs. 141, cxv. Zoöl. Ser.

SHORT-TAILED BAT.

Type locality. Unknown.

Geogr. Distr. Mexico to southern Brazil; West Indies.

*Sherborn, Ann. Mag. Nat. Hist., xx, 1897, p. 483.

1244. Hemiderma castaneum (H. Allen).

Carollia castanea H. Allen, Proc. Am. Phil. Soc., 1890, p. 19.

Hemiderma castaneum Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., IV, Pt. II, 1904, p. 670. Zoöl. Ser.

CHESTNUT-COLORED BAT.

Type locality. Costa Rica.

Geogr. Distr. Costa Rica, Central America.

Subfamily III. **Glossophaginæ.**Genus 193. **Glossophaga.**

Glossophaga E. Geoff., Mém. Mus. Hist. Nat. Paris, IV, 1818, pp. 413, 418, pls. 17, 18. Type *Vespertilio soricinus* Pallas.

Phyllophora Gray, Mag. Zoöl. & Bot., II, 1838, p. 489.

Nicon Gray, Proc. Zoöl. Soc., 1847, p. 15.

1245. Glossophaga mutica Merriam.

Glossophaga mutica Merr., Proc. Biol. Soc. Wash., XII, 1898, p. 18.

Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV,

Pt. II, 1904, p. 671. Zoöl. Ser.

MARIA MADRE ISLAND BAT.

Type locality. Maria Madre Island, Tres Marias Islands, State of Jalisco, Mexico.

Geogr. Distr. State of Jalisco, Mexico. (Tres Marias Islands.)

1246. Glossophaga soricina (Pallas):

Vespertilio soricinus Pall., Misc. Zool., 1766, p. 48, pl. IV, figs. 16-18, pl. V.

Glossophaga amplexicaudata E. Geoff., Mém. Mus. Hist. Nat. Paris, IV, 1818, p. 418, pl. XVIII.

Phyllophora nigra Gray, Voy. "Sulphur," Mamm., 1844, p. 18, pl. V, fig. 1.

Monophyllus leachii Gray, Voy. "Sulphur," Mamm., 1844, p. 18.

Nicon caudifer Gray, Ann. Mag. Nat. Hist., 1st Ser., XIX, 1847, p. 407.

Glossophaga soricina Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 672, figs. 142, CXVI. Zoöl. Ser.

SHREW-LIKE BAT.

Type locality. Unknown.

Geogr. Distr. Southern Mexico to Brazil and Bolivia.

a. Glossophaga soricina antillarum Rehn.

Glossophaga soricina antillarum Rehn, Proc. Acad. Nat. Scien. Phil., 1902, p. 37. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 672. Zoöl. Ser.

ANTILLES BAT.

Type locality. Port Antonio, Island of Jamaica.

Geogr. Distr. Island of Jamaica and the Bahamas.

Genus 194. **Chæronycteris.**

Chæronycteris Licht., Tschudi's Faun. Peruana, 1844, p. 70, pl. III.

Type Chæronycteris mexicana Tschudi.

1247. Chæronycteris mexicana Tschudi.

Chæronycteris mexicana Tschudi, Faun. Peruana, 1844, p. 72, pl. III, fig. 3. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 673, fig. CXVII. Zoöl. Ser.

TRES MARIAS ISLANDS BAT.

Type locality. Mexico.

Geogr. Distr. Tres Marias Islands, State of Jalisco, Mexico, to Guatemala.

1248. Chæronycteris minor Peters.

Chæronycteris minor Peters, Monatsb. K. Preuss. Akad. Wiss. Berl., 1868, p. 366. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 674. Zoöl. Ser.

SMALL BAT.

Type locality. Surinam.

Geogr. Distr. Guatemala, Central America, to Brazil, South America.

1249. Chæronycteris godmani Thomas.

Chæronycteris godmani Thomas, Ann. Mag. Nat. Hist., 7th Ser., XI, 1903, p. 288. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 674. Zoöl. Ser.

GODMAN'S BAT.

Type locality. Guatemala.

Geogr. Distr. Unknown.

Genus 195. **Hylonycteris.**

Hylonycteris Thomas, Ann. Mag. Nat. Hist., 7th Ser., XI, 1903, p. 286.

Type Hylonycteris underwoodi Thomas.

1250. Hylonycteris underwoodi Thomas.

Hylonycteris underwoodi Thomas, Ann. Mag. Nat. Hist., 7th Ser., XI, 1903, p. 287. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 675. Zoöl. Ser.

UNDERWOOD'S BAT.

Type locality. Rancho Redondo, Costa Rica.

Geogr. Distr. Rancho Redondo and Tarbaca, Costa Rica Central America. Range unknown.

Genus 196. **Monophyllus.**

Monophyllus Leach, Trans. Linn. Soc., XIII, 1821, p. 75. Type *Monophyllus redmani* Leach.

1251. Monophyllus portoricensis Miller.

Monophyllus portoricensis Miller, Proc. Wash. Acad. Scien., II, 1900, p. 34. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 677, figs. 143, CXVIII. Zoöl. Ser.

PORTO RICO NOSE-LEAF BAT.

Type locality. Cave near Bayamon, Porto Rico.

Geogr. Distr. Island of Porto Rico.

1252. Monophyllus plethodon Miller.

Monophyllus plethodon Miller, Proc. Wash. Acad. Scien., II, 1900, p. 35. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 677. Zoöl. Ser.

BARBADOES NOSE-LEAF BAT.

Type locality. St. Michael's Parish, Barbadoes.

Geogr. Distr. Island of Barbadoes.

1253. Monophyllus luciae Miller.

Monophyllus luciae Miller, Proc. Acad. Nat. Scien. Phil., 1902, p. 411. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 678. Zoöl. Ser.

SANTA LUCIA NOSE-LEAF BAT.

Type locality. Island of Santa Lucia.

Geogr. Distr. Island of Santa Lucia.

1254. Monophyllus clinedaphus Miller.

Monophyllus clinedaphus Miller, Proc. Wash. Acad. Scien., II, 1900, p. 36. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 678. Zoöl. Ser.

MILLER'S NOSE-LEAF BAT.

Type locality. Unknown.

Geogr. Distr. Unknown.

1255. *Monophyllus cubanus* Miller.

Monophyllus cubanus Miller, Proc. Acad. Nat. Scien. Phil., 1902, p. 410. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 678. Zool. Ser.

CUBAN NOSE-LEAF BAT.

Type locality. Baracoa, Island of Cuba.

Geogr. Distr. Restricted to the Island of Cuba.

1256. *Monophyllus redmani* Leach.

Monophyllus redmani Leach, Trans. Linn. Soc., XIII, 1821, p. 76. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 679. Zool. Ser.

REDMAN'S NOSE-LEAF BAT.

Type locality. Island of Jamaica.

Geogr. Distr. Island of Jamaica and possibly Island of Cuba.

Genus 197. *Leptonycteris*.

Leptonycteris Lydekker, in Flow. & Lydek. Mamm. Living and Extinct, 1891, p. 674. Type *Ischnoglossa nivalis* Saussure. *Ischnoglossa* Sauss., Rev. Mag. Zool., 2me Sér., XII, 1860, p. 492. (nec Kraatz, Coleopt., 1856.)

1257. *Leptonycteris nivalis* (Saussure).

Ischnoglossa nivalis Sauss., Rev. Mag. Zool., 2me Sér., XII, 1860, p. 492, pl. XX, figs. 2, 2c.

Leptonycteris nivalis Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 679, figs. 144, CXIX. Zool. Ser.

SNOWY BAT.

Type locality. Snow line on the peak of Mount Orizaba, State of Puebla, Mexico.

Geogr. Distr. Southern Mexico into Guatemala, Central America.

Genus 198. *Anura*.

Anoura (sic) Gray, Mag. Zool. & Bot., II, 1838, p. 490. Type *Anoura geoffroyi* Gray = *Glossophaga caudifera* E. Geoffroy. *Glossonycteris* Peters, Monatsb. K. Preuss. Akad. Wiss. Berl., 1868, p. 365.

1258. *Anura geoffroyi* Gray.

Glossophaga caudifera E. Geoff., Mém. Mus. Hist. Nat. Paris, IV, 1818, p. 418, pls. XVII, XVIII, fig. B.

Anoura! geoffroyi Gray, Mag. Zool. & Bot., II, 1838, p. 490.

Charonycteris peruana Tschudi, Faun. Peruana, I, 1844, p. 71.

Anoura! ecaudata Sauss., Rev. Mag. Zool., 2me Sér., XII, 1860, p. 493. (nec Geoffroy.)

Glossonycteris lasiopyga Peters, Monatsb. K. Preuss. Akad. Wiss. Berl., 1868, p. 368.

Anura geoffroyi Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 682, figs. 145, CXX. Zool. Ser.

GEOFFROY'S BAT.

Type locality. Brazil.

Geogr. Distr. Mexico through Central America into Brazil and Peru.

Genus 199. *Lichonycteris*.

***Lichonycteris* Thomas**, Ann. Mag. Nat. Hist., 6th Ser., XVI, 1895, p. 55. Type *Lichonycteris obscurus* Thomas.

1259. *Lichonycteris obscurus* Thomas.

Lichonycteris obscurus Thomas, Ann. Mag. Nat. Hist., 6th Ser., XVI, 1895, p. 56. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 683, fig. 146. Zool. Ser.

DUSKY BAT.

Type locality. Managua, Nicaragua.

Geogr. Distr. Nicaragua, Central America.

Genus 200. *Phyllonycteris*.

***Phyllonycteris* Gundl.**, Monatsb. K. Preuss. Akad. Wiss. Berl., 1860, p. 817. Type *Phyllonycteris poeyi* Gundlach.

1260. *Phyllonycteris bombifrons* Miller.

Phyllonycteris bombifrons Miller, Proc. Biol. Soc. Wash., XIII, 1899, p. 36. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 685. Zool. Ser.

BAYAMON BAT.

Type locality. Cave near Bayamon, Porto Rico.

Geogr. Distr. Island of Porto Rico.

1261. *Phyllonycteris planifrons* Miller.

Phyllonycteris planifrons Miller, Proc. Biol. Soc. Wash., XIII, 1899, p. 34. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 685. Zool. Ser.

FLAT FOREHEAD BAHAMA BAT.

Type locality. Nassau, New Providence; Great Abaco, Bahamas.
Geogr. Distr. Bahama Islands.

1262. *Phyllonycteris sezekorni* Gundlach.

Phyllonycteris sezekorni Gundl., Monatsb. K. Preuss. Akad. Wiss. Berl., 1860, p. 818. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 686. Zoöl. Ser.

SEZEKORN'S BAT.

Type locality. Cuba.
Geogr. Distr. Islands of Cuba and Jamaica.

1263. *Phyllonycteris poeyi* Gundlach.

Phyllonycteris poeyi Gundl., Monatsb. K. Preuss. Akad. Wiss. Berl., 1860, p. 817. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 686, figs. 147, CXXI. Zoöl. Ser.

POEY'S BAT.

Type locality. Cuba.
Geogr. Distr. Island of Cuba.

Genus 201. ***Rhithronycteris*.**

***Reithronycteris* (sic) Miller**, Proc. Acad. Nat. Scien. Phil., 1898, p. 333. Type *Reithronycteris aphylla* Miller.

1264. *Rhithronycteris aphylla* Miller.

Reithronycteris aphylla Miller, Proc. Acad. Nat. Scien. Phil., 1898, p. 334.
Rhithronycteris aphylla Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 688, figs. 148, CXXII. Zoöl. Ser.

LEAFLESS OR BLUNT-NOSED BAT.

Type locality. Island of Jamaica.
Geogr. Distr. Restricted to the Island of Jamaica.

Genus 202. ***Brachyphylla*.**

***Brachyphylla* Gray**, Proc. Zoöl. Soc., 1834, p. 122. Type *Brachyphylla cavernarum* Gray.

1265. *Brachyphylla cavernarum* Gray.

Brachyphylla cavernarum Gray, Proc. Zoöl. Soc., 1834, p. 123. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 689, fig. 149. Zoöl. Ser.

CAVERN NOSE-LEAF BAT.

Type locality. St. Vincent, Lesser Antilles.

Geogr. Distr. Islands of Cuba, Porto Rico, and St. Vincent.

1266. *Brachyphylla nana* Miller.

Brachyphylla nana Miller, Proc. Acad. Nat. Scien. Phil., 1902, p. 409. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 690, figs. 150, CXXIII. Zool. Ser.

DWARF NOSE-LEAF BAT.

Type locality. El Guama, Island of Cuba.

Geogr. Distr. Island of Cuba.

Subfamily IV. *Stenodermatinæ*.Genus 203. *Artibeus*.

Artibeus Leach, Trans. Linn. Soc., XIII, 1821, p. 75. Type *Artibeus jamaicensis* Leach.

Madatæus Leach, Trans. Linn. Soc., XIII, 1821, p. 81.

Pteroderma Gerv., Expéd. du Comte de Castelnau Amér. Sud, Zool., 1855, p. 34.

1267. *Artibeus coryi* J. A. Allen.

Artibeus coryi J. A. Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1890, p. 173. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 693. Zool. Ser.

CORY'S BAT.

Type locality. St. Andrew's Island, Caribbean Sea.

Geogr. Distr. Known only from the type locality.

1268. *Artibeus jamaicensis* Leach.

Artibeus jamaicensis Leach, Trans. Linn. Soc., XIII, 1821, p. 75. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 693, figs. 151, CXXIV. Zool. Ser.

Artibeus carpolegus Gosse, Nat. Sojourn in Jamaica, 1851, p. 271, pl. VI, fig. 5.

JAMAICA BAT.

Type locality. Island of Jamaica.

Geogr. Distr. Mexico to Brazil and Bolivia; West Indies.

1269. *Artibeus insularis* J. A. Allen.

Artibeus insularis J. A. Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1904, p. 231.

ST. KITTS BAT.

Type locality. Island of St. Kitts, West Indies.

Geogr. Distr. Restricted to the Island of St. Kitts, W. I.

Genl. Char. Size large; coloration pale, no head stripes; nose-leaf large.

Color. Above pale reddish brown, beneath paler; membranes pale brown.

Measurements. Length of head and body, 82; nose-leaf, 13.5 x 8; forearm, 61; thumb, 15; third metacarpal, 58; third finger, 78; tibia, 25.5; foot, 16.5; ear from crown, 15. Skull: width across upper second molar, 13; width of rostrum at base of canines, 8.6; length of upper tooth row, 11; length of mandible, 19.6; height at coronoid, 8; length of lower tooth row; 11.

1270. *Artibeus intermedius* J. A. Allen.

Artibeus carpolegus Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1891, p. 205. (nec Gosse.)

Artibeus intermedius J. A. Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1897, p. 33. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 694. Zoöl. Ser.

INTERMEDIATE BAT.

Type locality. San José, Costa Rica.

Geogr. Distr. Costa Rica, Central America. Range unknown.

1271. *Artibeus parvipes* Rehn.

Artibeus parvipes Rehn, Proc. Acad. Nat. Scien. Phil., 1902, p. 639.

Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 694. Zoöl. Ser.

SMALL-FOOTED BAT.

Type locality. Santiago de Cuba, Cuba.

Geogr. Distr. Island of Cuba.

1272. *Artibeus yucatanicus* Allen.

Artibeus yucatanicus Allen, Bull. Am. Mus. Nat. Hist. N. Y., 1904, p. 232.

YUCATAN BAT.

Type locality. Chichen Itza, Yucatan.

Geogr. Distr. Yucatan. Range unknown.

Genl. Char. Similar to *A. parvipes*.

Color. Similar to *A. parvipes*.

Measurements. "Forearm, 55; third metacarpal, 57; tibia, 24; foot, 14. Skull: total length, 27.5; zygomatic breadth, 17.5; mastoid breadth, 15; interorbital constriction, 7."

1273. Artibeus planirostris (Spix).

Phyllostoma planirostre Spix, Simiar. et Vespertil. Bras., 1823, p. 66, pl. xxxvi, fig. 1.

Phyllostoma obscurum Max., Beitr. Naturg. Bras., II, 1826, p. 203.

Phyllostoma perspicillatum Burm., Thiere Bras., 1854, p. 45.

Artibeus fallax Peters, Monatsb. K. Preuss. Akad. Wiss. Berl., 1865, p. 355.

Artibeus planirostris Elliot, Mamm. Middle Amer. & W. Indies. F. C. M. Pub., IV, Pt. II, 1904, p. 695, figs. 152, cxxv. Zool. Ser.

FLAT-NOSED BAT.

Type locality. Near Bahia, Brazil.

Geogr. Distr. State of Guerrero, Mexico, to Brazil; Island of Grenada.

1274. Artibeus watsoni Thomas.

Artibeus watsoni Thomas, Ann. Mag. Nat. Hist., 7th Ser., VII, 1901, p. 542. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M.

Pub., IV, Pt. II, 1904, p. 696. Zool. Ser.

WATSON'S BAT.

Type locality. Bogava, Chiriqui, Panama.

Geogr. Distr. Panama, Central America.

1275. Artibeus eva (Cope).

Dermanura eva Cope, Amer. Nat., xxIII, 1889, p. 130.

Artibeus eva Elliot, Mamm. Middle Amer. & W. Indies, F. C. M.

Pub., IV, Pt. II, 1904, p. 697. Zool. Ser.

ST. MARTIN'S BAT.

Type locality. Island of St. Martin, West Indies.

Geogr. Distr. Restricted to St. Martin's Island, W. I.

Genus 204. **Uroderma.**

Uroderma Peters, Monatsb. K. Preuss. Akad. Wiss. Berl., 1865, p. 588. (Footnote.) Type *Uroderma bilobatum* Peters.

1276. Uroderma convexum Lyon.

Uroderma convexum Lyon, Proc. Biol. Soc. Wash., xv, 1902, p. 83.

Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV,

Pt. II, 1904, p. 697, figs. 153, cxxvi. Zool. Ser.

COLON BAT.

Type locality. Colon, Panama.

Geogr. Distr. Known only from the type locality.

Genus 205. **Dermanura.**

Dermanura Gerv., Expéd. du Comte de Castelnau Amér. Sud, Zool., Mamm., 1855, p. 36. Type *Dermanura cinereum* Gervais.

1277. **Dermanura phæotis** Miller.

Dermanura phæotis Miller, Proc. Acad. Nat. Scien. Phil., 1902, p. 405. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 699. Zoöl. Ser.

DARK-EARED BAT.

Type locality. Chichen Itza, Yucatan.

Geogr. Distr. Yucatan. Range unknown.

1278. **Dermanura cinereum** Gervais.

Dermanura cinereum Gerv., Expéd. du Comte de Castelnau Amér. Sud, Zool., Mamm., 1855, p. 36, pl. VIII, fig. 4; pl. IX, figs. 4, 4a; pl. XI, fig. 3. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 699, figs. 154, CXXVII (D. 4—vittata). Zoöl. Ser.

Stenoderma tolteca Sauss., Rev. Mag. Zool., 2me Sér., XII, 1860, p. 427, pl. xv, fig. 4.

CINEREOUS BAT.

Type locality. Brazil.

Geogr. Distr. Mexico to Brazil.

Genus 206. **Vampyrops.**

Vampyrops Peters, Monatsb. K. Preuss. Akad. Wiss. Berl., 1865, p. 356. Type *Phyllostoma lineatum* E. Geoffroy.

Vampyressa Thomas, Ann. Mag. Nat. Hist., 7th Ser., v, 1900, p. 270.

Vampyriscus Thomas, Ann. Mag. Nat. Hist., 7th Ser., v, 1900, p. 270.

Vampyrodes Thomas, Ann. Mag. Nat. Hist., 7th Ser., v, 1900, p. 270.

1279. **Vampyrops lineatus** (E. Geoffroy).

Phyllostoma lineatum E. Geoff., Ann. Mus. Hist. Nat. Paris, xv, 1810, p. 180.

Vampyrops lineatus Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 702. Zoöl. Ser.

WHITE-STRIPED BAT.

Type locality. Paraguay.

Geogr. Distr. Mexico to Paraguay.

1280. *Vampyrops vittatus* (Peters).

Artibeus vittatus Peters, Monatsb. K. Preuss. Akad. Wiss. Berl.
1859, p. 225.

Vampyrops vittatus Elliot, Mamm. Middle Amer. & W. Indies
F. C. M. Pub., IV, Pt. II, 1904, p. 704, fig. CXXVIII. Zoöl. Ser.

PETERS' WHITE-STRIPED BAT.

Type locality. Puerto Cabello, Venezuela.

Geogr. Distr. Costa Rica, Central America, to northern South America.

1281. *Vampyrops helleri* Peters.

Vampyrops helleri Peters, Monatsb. K. Preuss. Akad. Wiss. Berl.,
1866, p. 392. Elliot, Mamm. Middle Amer. & W. Indies,
F. C. M. Pub., IV, Pt. II, 1904, p. 703, fig. 155. Zoöl. Ser.

HELLER'S WHITE-STRIPED BAT.

Type locality. Mexico.

Geogr. Distr. Mexico to South America; Venezuela; Chiriqui, Panama, and San Miguel Island, Bay of Panama.

Genus 207. ****Stenoderma*.**

Stenoderma E. Geoff., Descr. l'Égypte, Mamm., II, 1818, p. 114.

Type Stenoderma rufum E. Geoffroy.

Artibeus Gray, Mag. Zoöl. & Bot., II, 1838, p. 491.

Histiops Peters, Monatsb. K. Preuss. Akad. Wiss. Berl., 1869,
p. 339.

Artibeus Gerv., Expéd. du Comte de Castelnau Amér. Sud, Zool.,
Mamm., 1855, p. 34. (nec Leach.)

Peltorhinus Peters, Monatsb. K. Preuss. Akad. Wiss. Berl., 1876,
p. 433.

Subgenus A. ***Stenoderma*.**

1282. *Stenoderma montserratense* Thomas.

Stenoderma montserratense Thomas, Proc. Zoöl. Soc., 1894, p. 133.
Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV,
Pt. II, 1904, p. 705. Zoöl. Ser.

MONTSERRAT BAT.

Type locality. Island of Montserrat, West Indies.

Geogr. Distr. Known only from the type locality.

1283. *Stenoderma nicholli* Thomas.

Stenoderma nicholli Thomas, Ann. Mag. Nat. Hist., 6th Ser., VII,
1891, p. 529. Elliot, Mamm. Middle Amer. & W. Indies,
F. C. M. Pub., IV, Pt. II, 1904, p. 706. Zoöl. Ser.

*Misprinted *Sternoderma* in the Mamm. Middle Amer. & W. Indies.

NICHOLLS' BAT.

Type locality. Island of Dominica, West Indies.

Geogr. Distr. Known only from the type locality.

1284. *Stenoderma luciae* Miller.

Stenoderma luciae Miller, Proc. Acad. Nat. Scien. Phil., 1902, p. 407.

Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV,

Pt. II, 1904, p. 706, fig. CXXIX. Zool. Ser.

SANTA LUCIA BAT.

Type locality. Island of Santa Lucia, West Indies.

Geogr. Distr. Restricted to the Island of Santa Lucia, W. I.

Subgenus B. *Peltorhinus*.1285. *Stenoderma achradophilum* (Gosse).

Artibeus achradophilus Gosse, Nat. Sojourn in Jamaica, 1851,
p. 271, pl. VI, fig. 4.

Artibeus sulphureus Gosse, Nat. Sojourn in Jamaica, 1851, p. 271,
pl. VI, fig. 5.

Artibeus jamaicensis Gosse, Nat. Sojourn in Jamaica, 1851, p. 271,
pl. VI, fig. 3.

Artibeus flavescens Gray, Proc. Zool. Soc., 1866, p. 117.

Stenoderma achradophilum Elliot, Mamm. Middle Amer. & W. In-
dies, F. C. M. Pub., IV, Pt. II, 1904, p. 707, fig. 156. Zool. Ser.

FRUIT-LOVING BAT.

Type locality. Content, Island of Jamaica.

Geogr. Distr. Islands of Cuba and Jamaica.

Genus 208. *Phyllops*.

Phyllops Peters, Monatsb. K. Preuss. Akad. Wiss. Berl., 1865, p. 356.

Type *Arctibeus (sic) falcatus* Gray.

1286. *Phyllops falcatus* (Gray).

Arctibeus! falcatus Gray, Ann. Mag. Nat. Hist., IV, 1839, p. 1.

Phyllostoma albomaculatum Gundl., Monatsb. K. Preuss. Akad.
Wiss. Berl., 1861, p. 155.

Phyllops falcatum Elliot, Mamm. Middle Amer. & W. Indies,
F. C. M. Pub., IV, Pt. II, 1904, p. 708, fig. 157. Zool. Ser.

FALCATE BAT.

Type locality. Cuba.

Geogr. Distr. Island of Cuba.

Genus 209. **Ectophylla.**

Ectophylla H. Allen, Proc. U. S. Nat. Mus., xv, 1892, p. 441, 2 figs. in text. Type *Ectophylla alba* H. Allen.

1287. **Ectophylla alba** H. Allen.

Ectophylla alba H. Allen, Proc. U. S. Nat. Mus., xv, 1892, p. 442.

Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV,

Pt. II, 1904, p. 710, fig. CXXX. Zool. Ser.

WHITE HONDURAS BAT.

Type locality. Segovia River, Honduras.

Geogr. Distr. Honduras, Central America. Range unknown.

Genus 210. **Chiroderma.**

Chiroderma Peters, Monatsb. K. Preuss. Akad. Wiss. Berl., 1860, p. 747. Type *Chiroderma villosum* Peters.

1288. **Chiroderma salvini** Dobson.

Chiroderma salvini Dobson, Cat. Chiropt. Brit. Mus., 1878, p. 532,

pl. 29, figs. 3, 3a. Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., IV, Pt. II, 1904, p. 710, figs. 158, CXXXI. Zool.

Ser.

SALVIN'S NOSE-LEAFED BAT.

Type locality. Costa Rica.

Geogr. Distr. Costa Rica, Central America. Range unknown.

Genus 211. **Pygoderma.**

Pygoderma Peters, Monatsb. K. Preuss. Akad. Wiss. Berl., 1863, p. 83; 1865, p. 357. Type *Phyllostoma bilabiatum* Wagner.

1289. **Pygoderma bilabiatum** (Wagner).

Phyllostoma bilabiatum Wagn., Archiv f. Naturg., IX, 1843, bd.

I, p. 366.

Artibeus (sic) *leucomus* Gray, Proc. Zool. Soc., 1848, p. 57.

Stenoderma (*Pygoderma*) *microdon* Peters, Monatsb. K. Preuss.

Akad. Wiss. Berl., 1863, p. 83.

Pygoderma bilabiatum Elliot, Mamm. Middle Amer. & W. Indies

F. C. M. Pub., IV, Pt. II, 1904, p. 712, figs. 159, CXXXII. Zool. Ser.

YPANEMA NOSE-LEAF BAT.

Type locality. Ypanema, São Paulo, Brazil.

Geogr. Distr. Mexico to Brazil.

Genus 212. **Sturnira.**

Sturnira Gray, Ann. Mag. Nat. Hist., 1st Ser., x, 1842, p. 257. Type
Sturnira spectrum Gray = *Phyllostoma lilium* E. Geoffroy.
Nyctiplanus Gray, Proc. Zool. Soc., 1849, p. 57.

1290. **Sturnira lilium** (E. Geoffroy).

Phyllostoma lilium E. Geoff., Ann. Mus. Hist. Nat. Paris, xv, 1810,
 p. 181.

Phyllostoma spiculatus Illig., Licht. Verz. der Doubl., 1823, p. 3.

Sturnira spectrum Gray, Ann. Mag. Nat. Hist., 1st Ser., 1842, p. 257.

Phyllostoma excisum Wagn., Archiv f. Naturg., 1842, p. 358.

Phyllostoma albescens Wagn., Abhandl. Münch. Akad., v, p. 178.

Phyllostoma fumarium Wagn., Abhandl. Münch. Akad., v, p. 178.

Phyllostoma erythromos Tschudi, Faun. Peruana, 1844-46, p. 64
 pl. 1.

Phyllostoma oporaphilum Tschudi, Faun. Peruana, 1844-46, p. 64,
 pl. 11.

Stenoderma chilense Gerv., Gay's Hist. Chile, Mamm., 1847, p.
 30, pl. 1, fig. 1.

Nyctiplanus rotundatus Gray, Proc. Zool. Soc., 1848, p. 57.

Phyllostoma chrysocomus Wagn., Schreb. Säugth., Suppl., v, 1855,
 p. 635.

Sturnira lilium Elliot, Mamm. Middle Amer. & W. Indies, F. C. M.
 Pub., IV, Pt. II, 1904, p. 715, figs. 160, cxxxiii. Zool. Ser.

GEOFFROY'S BAT.

Type locality. Paraguay.

Geogr. Distr. States of Jalisco and Vera Cruz, Mexico, to Para-
 guay and Chili. West Indies.

Subfamily V. **Centurioninæ.**Genus 213. **Centurio.**

Centurio Gray, Ann. Mag. Nat. Hist., 1st Ser., x, 1842, p. 259. Type
Centurio senex Gray.

Trichocoryes! H. Allen, Proc. Acad. Nat. Scien. Phil., 1861, p. 359.

1291. **Centurio senex** Gray.

Centurio senex Gray, Ann. Mag. Nat. Hist., 1st Ser., x, 1842, p.
 259. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M.
 Pub., IV, Pt. II, 1904, p. 717, figs. 161, cxxxiv, cxxxv. Zool.
 Ser.

Centurio flavogularis Licht. & Peters, Monatsb. K. Preuss. Akad. Wiss. Berl., 1854, p. 335.

Centurio mexicanus Sauss., Rev. Mag. Zool., 2me Sér., XII, 1860, p. 381.

Centurio mcmurtrii H. Allen, Proc. Acad. Nat. Scien. Phil., 1861, p. 360. (Adult male.)

Centurio minor Ward, Amer. Nat., xxv, 1891, p. 750. (Female.)

WRINKLED-FACE BAT.

Type locality. Unknown.

Geogr. Distr. Eastern Mexico, State of Vera Cruz to Costa Rica, Central America. Exact range unknown.

Subfamily VI. **Desmodontinæ.**

Genus 214. * **Desmodon.**

Desmodus (sic) Max., Abbild. Naturg. Bras., 5te Lief., 1824, pl. and text. Type *Desmodus rufus* Maximilian = *Phyllostoma rotundum* E. Geoffroy.

1292. **Desmodon rotundum** (E. Geoffroy).

Phyllostoma rotundum E. Geoffroy, Ann. Mus. Hist. Nat. Paris, 1810, p. 181.

Rhinolophus ecaudatus Schinz, Cuv. Thiere., I, 1821, p. 168.

Desmodus rufus Max., Beitr. Naturg. Bras., 1826, p. 233.

Edostoma cinerea D'Orbigny, Voy. l'Amér. Mérid., IV, 1847, p. 11, pl. VIII.

Desmodus d'orbignyi Waterh., Voy. Beagle, Mamm., 1838, p. 1, pls. 5, xxxv, fig. 1.

Desmodus murinus Wagn., Schreb. Säugth., Suppl., I, 1847, p. 377.

Desmodus fuscus Lund, Burm. Thiere Bras., 1854, p. 57.

Desmodus rotundus Elliot, Mamm. Middle Amer. & W. Indies, F.C.M. Pub., IV, Pt. II, 1904, p. 718, figs. 162, cxxxvi. Zool. Ser.

RUFIOUS VAMPIRE BAT.

Type locality. Paraguay.

Geogr. Distr. Southern Mexico south to Chili and Paraguay.

Genus 215. **Diphylla.**

Diphylla Spix, Simiar. et Vespert. Bras., 1823, p. 68. Type *Diphylla ecaudata* Spix.

Hæmatonycteris H. Allen, Proc. U. S. Nat. Mus., xviii, 1896, p. 777. Type *Diphylla ecaudata* Dobson. (nec Spix?)

* δεσμότης, οδοῦς = DESMODON Lat. nec *Desmodus*.

1293. *Diphylla ecaudata* Spix.

Diphylla ecaudata Spix, Simiar. et Vespert. Bras., 1823, p. 68, pl. xxxvi. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 720, fig. 163. Zool. Ser.

TAILLESS BAT.

Type locality. Brazil.

Geogr. Distr. Southern Mexico? Brazil.

1294. *Diphylla centralis* Thomas.

Diphylla centralis Thomas, Ann. Mag. Nat. Hist., 7th Ser., XI, 1903, p. 378. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 721. Zool. Ser.

BOQUETE TAILLESS BAT.

Type locality. Boquete, Chiriqui, Panama.

Geogr. Distr. Panama into South America, possibly to Brazil.

Order XI. **Primates. Primates.**

Sub-order Anthropoidea.

Family I. **Callitrichidæ. Marmosets.**

Genus 216. **Ædipomidas.**

- Ædipomidas** Reichenb. Vollständ. Naturgesch. Affen, 1862, pp. 5-6, pl. II, figs. 18-20. Type *Simia ædipus* Linnæus.
Midas E. Geoff., Ann. Mus. Hist. Nat. Paris, XIX, 1812, p. 120. (nec Latreille, 1796.)
Marikina Reich., Vollständ. Naturgesch. Affen, 1862, pp. 7-9, pl. II, figs. 25-31.
Ædipus Gray, Cat. Monkeys, Lemurs and Fruit-eating Bats Brit. Mus., 1870, p. 65. (nec Tschudi, 1838.)
Seniocebus Gray, Cat. Monkeys, Lemurs and Fruit-eating Bats Brit. Mus., 1870, p. 68.

1295. **Ædipomidas geoffroyi** (Pucheran).

- Hapale geoffroyi* Pucheran, Rev. Zoöl., 1845, p. 336.
Hapale ædipus Sclat., Nat. Hist. Rev., 1861, p. 509. (nec Linn.)
Midas geoffroyi Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 724, figs. 164, CXXXVII. Zoöl. Ser.
GEOFFROY'S TITI MONKEY.
Type locality. Panama.
Geogr. Distr. Panama to Colombia.

Family II. **Cebidæ. Prehensile-tailed Monkeys.**

Subfamily I. **Alouattinæ.**

Genus 217. **Alouatta. Howling Monkeys.**

- Alouatta** Lacépède, Tabl. Div. Sous-div. Ordres et Genres Mamm., 1799, p. 4. Type *Simia belzebul* Linnæus.
Mycetes Illig., Prodr. Syst. Mamm. et Av., 1811, p. 70.

1296. **Alouatta villosa** (Gray).
Mycetes villosus Gray, Ann. Mag. Nat. Hist., 1st Ser., XVI, 1845, p. 220.
Alouatta villosa Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 726, pl. LXVII. Zoöl. Ser.

VILLOUS HOWLER. *Mono* in Guatemala.

Type locality. Brazil.

Geogr. Distr. Guatemala and Honduras, Central America, into South America.

1297. *Alouatta palliata* (Gray).

Mycetes palliatus Gray, Proc. Zool. Soc., 1848, p. 138, pl. vi.

Alouatta palliata Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., IV, Pt. II, 1904, p. 726, fig. cxxxviii. Zool. Ser.

MANTLED HOWLER. *Congo* in Costa Rica; *Congo, o'Mono Chillon* in Nicaragua.

Type locality. Caracas, Venezuela.

Geogr. Distr. Nicaragua to South America.

a. Alouatta palliata mexicana Merriam.

Alouatta palliata mexicana Merr., Proc. Biol. Soc. Wash., xv,

1902, p. 67. Elliot, Mamm. Middle Amer. & W. Indies,

F. C. M. Pub., IV, Pt. II, 1904, p. 727. Zool. Ser.

MEXICAN HOWLER.

Type locality. Minatitlan, State of Vera Cruz, Mexico.

Geogr. Distr. Eastern Mexico.

b. Alouatta palliata coibensis Thomas.

Alouatta palliata coibensis Thomas, Novit. Zool., ix, 1902, p. 135.

Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV,

Pt. II, 1904, p. 727. Zool. Ser.

ISLAND OF COIBA HOWLER.

Type locality. Coiba Island, off west coast of Panama.

Geogr. Distr. Restricted to Coiba Island.

Subfamily II. **Aotinae. Squirrel Monkeys.**

Genus 218. **Aotus.**

Aotes Humboldt, Rec. d'obs. Zool. et Anat. Comp., I, 1811, pp.

306-311, 358, (for *Aotus*) pl. xxviii. Type *Simia trivirgata*

Humboldt.

Nyctipithecus Spix, Simiar. et Vespert. Bras., 1823, p. 25.

1298. *Aotus rufipes* (Sclater).

Nyctipithecus rufipes Slater, Proc. Zool. Soc., 1872, p. 3.

Aotus rufipes Elliot, Mamm. Middle Amer. & W. Indies, F. C. M.

Pub., IV, Pt. II, 1904, p. 729, figs. 165 (*Aotus azarae*), cxxxix.

Zool. Ser.

RUFIOUS-FOOT SQUIRREL MONKEY.

Type locality. San Juan del Norte, Nicaragua.

Geogr. Distr. Nicaragua. Range unknown.

1299. *Aotus vociferans* (Spix).

Nyctipithecus vociferans Spix, Simiar. et Vespert. Bras., 1823, p. 25, pl. 19.

Aotus vociferans Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 729. Zoöl. Ser.

NOISY SQUIRREL MONKEY. *Mico-dormilon* in Colombia.

Type locality. Tabatinga, Upper Amazon, Brazil.

Geogr. Distr. Costa Rica, Central America to Brazil.

Genus 219. *Saimiri*.

Saimiri Voigt, Cuvier's Thierreich, I, 1831, p. 95. Type *Simia sciurea* Linnæus.

1300. *Saimiri ærstedii* (Reinhardt).

Chrysothrix ærstedii Reinh., Vidensk. Medd. Nat. For. Kjöbenh., 1872, p. 157, pl. III.

Saimiris entomophaga Sclat., Proc. Zoöl. Soc., 1872, p. 3. (nec D'Orbigny.)

Chrysothrix sciurea Sclat., Nat. Hist. Rev., 1861, p. 510. (nec Linn.)

Saimiri ærstedii Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 731, figs. 166, cXL. Zoöl. Ser.

ÆRSTED'S TITI MONKEY. *Titi*, *Cuistiti* in Costa Rica.

Type locality. Cartago, Costa Rica.

Geogr. Distr. Guatemala to Panama, Central America.

Subfamily III. *Cebinæ*.Genus 220. *Ateles*. Spider Monkeys.

Ateles E. Geoff., Ann. Mus. Hist. Nat. Paris, VII, 1806, p. 262. Type *Simia paniscus* Linnæus.

1301. *Ateles vellerosus* Gray.

Ateles vellerosus Gray, Proc. Zoöl. Soc., 1865, p. 733. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 733, fig. cXLI, pl. LXVIII. Zoöl. Ser.

Ateles fuliginosus Schleg., Mus. Pays-Bas, III, 1876, p. 179. (nec Kuhl.)

Ateles pan. Schleg., Mus. Pays-Bas, III, 1876, p. 180.

MEXICAN SPIDER MONKEY.

Type locality. Unknown.

Geogr. Distr. Volcano of Orizaba, State of Vera Cruz, south to Guatemala, Central America.

1302. *Ateles geoffroyi* Kuhl.

Ateles geoffroyi Kuhl, Beit. Zool., 1820, p. 26. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 733. Zoöl. Ser.

Ateles melanochir Desm., Mamm., 1820, p. 76.

Eriodes frontatus Gray, Ann. Mag. Nat. Hist., 1st Ser., x, 1842, p. 256.

Ateles variegatus Frantz., Archiv f. Naturg., xxxv, I, 1869, p. 257. (nec Wagn.)

Ateles hybridus Gray, Cat. Monkeys, Lemurs and Fruit-eating Bats Brit. Mus., 1870, p. 43.

Ateles ornatus Gray, Cat. Monkeys, Lemurs and Fruit-eating Bats Brit. Mus., 1870, p. 44.

Ateles albifrons Gray, Cat. Monkeys, Lemurs and Fruit-eating Bats Brit. Mus., 1870, p. 44.

GEOFFROY'S SPIDER MONKEY. *Mono Colorado* in Costa Rica.

Type locality. Unknown.

Geogr. Distr. Nicaragua, Central America, to Colombia, South America.

1303. *Ateles rufiventris* Sclater.

Ateles rufiventris Sclat., Proc. Zoöl. Soc., 1872, p. 688, pl. LVII, juv. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 734. Zoöl. Ser.

FULVOUS-BELLIED SPIDER MONKEY.

Type locality. Colon, Atrato River, northern Colombia.

Geogr. Distr. Panama? into Colombia, South America.

1304. *Ateles ater* F. Cuvier.

Ateles ater F. Cuv., Hist. Nat. Mamm., 2d ed., III, 1823, Livr. xxxix, p. 107, pl. 56. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 734. Zoöl. Ser.

BLACK SPIDER MONKEY.

Type locality. Cayenne, French Guiana.

Geogr. Distr. Panama to eastern Peru.

1305. *Ateles grisescens* Gray.

Ateles grisescens Gray, Proc. Zoöl. Soc., 1865, p. 732. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 734. Zoöl. Ser.

GRIZZLED SPIDER MONKEY.

Type locality. Unknown.*Geogr. Distr.* Central America.Genus 221. **Cebus.****Cebus** Erxl., Syst. Règn. Anim., 1777, p. 44. Type —?**1306. Cebus hypoleucus** (Humboldt).*Simia hypoleuca* Humb., Recueil, Obs. Zool. Anat. Comp., 1, 1811, p. 337.*Cebus hypoleucus* Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 736, figs. 167, CLXII. Zool. Ser.WHITE-THROATED CAPUCHIN. *Mono carablanca.**Type locality.* Rio Sinu, Bolívar, Colombia.*Geogr. Distr.* Nicaragua to Colombia.**1307. Cebus imitator** Thomas.*Cebus imitator* Thomas, Ann. Mag. Nat. Hist., 7th Ser., XI, 1903, p. 376. Elliot, Mamm. Middle Amer. & W. Indies, F. C. M. Pub., IV, Pt. II, 1904, p. 737. Zool. Ser.

ALLIED SAPAJOU.

Type locality. Boquete, Chiriqui, Panama. Altitude, 4,000 feet.*Geogr. Distr.* Panama, Central America.Family III. **Cercopithecidae.**Subfamily I. **Cercopithecinae.**Genus 222. **Cercopithecus.****Cercopithecus** Gronovius, Zooph. Gron., 1, 1763, p. 5. Type —?**1308. *Cercopithecus callitrichus** Is. Geoffroy.*Le Callitriche* F. Cuv., Mamm., 1, 1824, pl. 21.*Cercopithecus callitrichus* Is. Geoffroy, Cat. Mamm., 1851, p. 23.

Schleg., Mus. Pays-Bas, Simiæ, 1876, p. 73.

*Not having been able to obtain any reliable information regarding this species in the West Indies, it was omitted from the work on the mammals of Middle America and the West Indies. I have lately been informed by Mr. A. H. Clark of Boston that it is common in certain districts of Barbadoes, and also in St. Kitts, and is, therefore, included in this list. I have heard that another species of monkey from the Old World has been introduced into the Island of Grenada, but although I have tried to obtain some information regarding it, and written to persons on the Island, yet I am unable to give its name, or learn anything about it.

GREEN MONKEY.

Type locality. Senegambia, West Africa.

Geogr. Distr. West Africa, Senegambia to River Niger. Introduced into the Islands of St. Kitts and Barbadoes, West Indies.

Color. Upper parts of body and sides yellowish green, the hairs ringed with black and yellow; under parts, inner side of arms and legs, throat and neck yellowish white; outer side of arms and legs, and also hands and feet gray; tail like the body, with a long yellow pencil; under side paler; whiskers, eyebrows, and anal region yellow; face, ears, and palms of hands and soles of feet black.

Measurements. Total length, 1066.8; tail, 673.

APPENDIX.

THE FOLLOWING DESCRIPTIONS WERE PUBLISHED TOO LATE
TO BE INCLUDED IN THEIR PROPER PLACES IN
THE BODY OF THE WORK.

Order VI. **Rodentia.**

Family I. **Sciuridæ.**

Subfamily I. **Sciurinaë.**

Genus 51. **Tamias.**

173a. *Tamias amœnus operarius* (Merriam).

Eutamias amœnus operarius Merr., Proc. Biol. Soc. Wash., xviii, 1905, p. 164.

BUSY CHIPMUNK.

Type locality. Gold Hill, Colorado. Altitude, 7,400 feet.

Genl. Char. "Similar to *amœnus*, but lacking the black on posterior two-thirds of ear; lower dark cheek stripe and stripe between eye and ear usually narrower; tail longer, the upper side more strongly fulvous; black beneath less marked; submarginal black band beneath narrower; median dorsal stripe blacker and more pronounced on back of neck and occiput; outer lateral stripe more pronounced." (Merriam, l. c.)

Fall and Post-breeding Pelage. Middle pair of light dorsal stripes and rump gray; in post-breeding pelage, the dorsal stripes are less gray, the rump more olivaceous, and the shoulders and anterior part of back more fulvous. "Some specimens have the under parts suffused with yellowish and fulvous, as in the larger *luteiventris*."

Measurements. Total length, 200; tail vertebræ, 93; hind foot, 31.

177aa. *Tamias hopiensis* (Merriam).

Eutamias hopiensis Merr., Proc. Biol. Soc. Wash., xviii, 1905, p. 165.

HOPi CHIPMUNK.

Type locality. Keam Cañon, Painted Desert, Arizona.

Genl. Char. Size medium; general color pale golden fulvous; head and rump pale.

Color. Head above pale buffy gray; rump pale grizzled ochraceous gray; outer dorsal stripes white; inner light stripes grizzled white and gray, sometimes tinged with buffy; dark dorsal stripes in winter bright fulvous, in summer with black centers; tail deep rufous, narrow black submarginal band beneath.

Measurements. Total length, 224; tail vertebræ, 101; hind foot, 33.5.

Family IV. **Muridæ.**Subfamily II. **Cricetinaë.**Genus 60. **Peromyscus.****278aa.** *Peromyscus leucopus fusus* Bangs.

Peromyscus leucopus fusus Bangs, N. Eng. Zoöl. Club, IV, 1905,
p. 13.

MARTHA'S VINEYARD ISLAND DEER MOUSE.

Type locality. West Tisbury, Martha's Vineyard Island, Massachusetts.

Geogr. Distr. Martha's Vineyard Island, Massachusetts.

Genl. Char. Similar to *P. l. noveboracensis*, but larger.

Color. *Summer Pelage.* Above rich brown, dorsal region darkest; beneath white; hands and feet white; ears dusky; tail blackish above, white below.

Measurements. Total length, 187-203; tail vertebræ, 85-96; hind foot, 21.5-23.5. Skull: basal length, 25.8; occipito-nasal length, 28.4; length of nasals, 10.6; palatal length, 11.4; zygomatic width, 14.6; interorbital width, 4.4; length of mandible, 15.8; of upper molar series, 3.8; of lower molar series, 3.6.

278aaa. *Peromyscus leucopus ammodytes* Bangs.

Peromyscus leucopus ammodytes Bangs, Proc. N. Eng. Zoöl. Club,
IV, 1905, p. 14.

MONOMOY ISLAND DEER MOUSE.

Type locality. Monomoy Island, Massachusetts.

Geogr. Distr. Known only from the type locality.

Genl. Char. Smaller and paler than *P. l. noveboracensis*.

Color. Above pale gray, beneath white; ill-defined dorsal stripe sometimes present; hands and feet white; tail pale gray above, white below, ears pale drab.

Measurements. Total length, 151.5-188; tail vertebræ, 67-88; hind foot, 18.8-21. Skull: basal length, 23.8; occipito-nasal length, 26; length of nasals, 10; palatal length, 10.4; zygomatic width, 13.6; interorbital width, 4; length of mandible, 14.2; length of upper molar series, 3.4; of lower molar series, 3.

Family XII. **Leporidae.**Genus III. **Lepus.**Subgenus F. **Macrotolagus.**

881a. **Lepus callotis flavigularis* Wagner.

Lepus callotis var. *flavigularis* Wagn., Schreb. Säugth., Suppl.,
IV, 1844, p. 107.

YELLOW-THROATED JACK RABBIT.

Type locality. San Mateo del Mar, Tehuantepec.

Geogr. Distr. Southern Mexico.

Genl. Char. Similar to *L. callotis*, gray of the rump pale, and very sharply defined from the black and yellow of the dorsal tract; throat patch greatly elongated, much more extensive than that of *L. callotis*, and a pale straw yellow; fore and hind legs iron gray.

Measurements. Total length, 525; tail vertebræ, 70; hind foot, 135; ear, 122.

*A fairly marked subspecies.

INDEX OF LATIN NAMES .

FOR THE

CHECK-LIST OF MAMMALS &c.

THE MAMMALS OF MIDDLE AMERICA, WEST INDIES &c.

AND THE

SYNOPSIS OF NORTH AMERICAN MAMMALS &c.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
abbreviatus. (<i>Microtus</i>).....	243		189
aberti. (<i>Sciurus</i>).....	66, 67	112	57
abieticola. (<i>Mustela a.</i>).....	419		
abietinoides. (<i>Mustela a.</i>).....	420		
abietorum. (<i>Peromyscus c.</i>).....	144		128
abietorum. (<i>Vulpes a.</i>).....	381		
abietorum. (<i>Zapus i.</i>).....	328		264
ablusus. (<i>Citellus p.</i>).....	109		
abrasus. (<i>Dysopes</i>).....	497	623	
abrasus. (<i>Promops</i>).....	497	623	417
Abromys.....	300	352	241
Abromys lordi.....	308		250
Abrothrix.....	204	272	
acadicus. (<i>Meriones</i>).....	324		257
acadicus. (<i>Microtus p.</i>).....	235		180
acapulcensis. (<i>Cervus</i>).....	46	74	
achradophilum. (<i>Stenoderma</i>).....	527	707	
achradophilus. (<i>Artibeus</i>).....	527	707	
Acodon.....	204	272	
Acodon irazu.....	204	274	
Acodon teguina.....	204	273	
Acodon t. apricus.....	204	274	
Acodon xerampelinus.....	205	275	
acræa. (<i>Teonoma</i>).....	221		
acræa. (<i>Teonoma c.</i>).....	221		
acrobates. (<i>Sciurus</i>).....	124		107
actuosa. (<i>Mustela a.</i>).....	420, 421		336
actuosa. (<i>Mustela c.</i>).....			*510
acuticaudatus. (<i>Molossus</i>).....	495	620	
acuto-rostrata. (<i>Balenoptera</i>).....	18		11
acutus. (<i>Delphinus</i>).....	29		25
acutus. (<i>Lagenorhynchus</i>).....	29		25
Adelonycteris.....	482	586	410
Adelonycteris gaumeri.....	485	590	
Adenonotus.....	35	62	
adocetus. (<i>Citellus</i>).....	116	152	

* Supplement.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
adolphei. (Macroxus)	75	126	
adolphii. (Sciurus)	75	126	
adsitus. (Eutamias)	95		
adsitus. (Tamias)	95		
adpersus. (Heteromys)	319	373	
aedium. (Plagiodontia)	332		
ægyptiacus. (Nyctinomus)	499	628	
æneus. (Scalops)	468		391
Aeorestes	473	571	400
aereus. (Scalops)	471		390
aereus. (Scalops t.)	471		
aereus. (Scalopus)	471		390
affinis. (Hesperomys)	161	180	
affinis. (Mustela)	433		
affinis. (Neotoma f.)	207		160
affinis. (Peromyscus)	161	180	
affinis. (Putorius)	433	534	
affinis. (Sciurus)	74	124	
affinis. (Tamias q.)	92		77
affinis. (Vespertilio)	478		402
Agaphelus	18		11
Agaphelus gibbosus	18		11
Agaphelus glaucus	16	40	9
agilis. (Dipodomys)	293	351	236
agilis. (Perodipus)	293, 294	351	236
Agouti.	337	408	
Agouti p. virgata	337	408	
Agoutidæ	335	403	
Agricola	234	299	178
akeleyi. (Peromyscus)	148		129
Akodon	204	272	
Akodon boliviense	204	272	
Akodon irazu	204	274	
Akodon t. apricus	204	274	
Akodon xerampelinus	205	275	
alacer. (Lepus f.)	343		282
alascana. (Callotaria)	438		
alascanus. (Callorhinus)	438		
alascensis. (Dicrostonyx h.)	260, 261		210
alascensis. (Evotomys)	229		177
alascensis. (Evotomys d.)	229		
alascensis. (Lemmus)	259		207
alascensis. (Myotis l.)	479		402
alascensis. (Putorius c.)	426		344
alascensis. (Putorius r.)	426		
alascensis. (Sorex g.)	451		372
alascensis. (Sorex o.)	451		
alascensis. (Ursus h.)	393		312
alascensis. (Vulpes)	381		
alascensis. (Zapus h.)	324		259
alaskanus. (Neosorex p.)	458		

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
alaskanus. (Sorex n.)	458		
alba. (Ectophylla)	528	710	
albertæ. (Citellus c.)	111		
albescens. (Onychomys l.)	131	†742	
albescens. (Phyllostoma)	529	715	
albescens. (Reithrodontomys)	192		
albescens. (Rhithrodontomys)	192		
albescens. (Vespertilio)	479		
albibarbis. (Neosorex)	458		379
albicans. (Cetus)	25		19
albicauda. (Phoca)	444		364
albifrons. (Ateles)	535	733	
albigena. (Phoca)	441		362
albigula. (Neotoma)	217		
albigula. (Neotoma i.)	217	285	162
albigularis. (Myodes)	259		207
albigularis. (Vespertilio)	485	590	
albigularis. (Vesperus)	485	590	
albilabris. (Reithrodontomys m.)	202	269	
albilabris. (Rhithrodontomys m.)	202	269	
albimanus. (Delphinus)		55	28
albinasus. (Pappo)	268	321	
alбини. (Phoca)	444		364
albipes. (Bassariscus)	399	486	
albipes. (Phenacomys)	228		
albipes. (Sciurus)	69	117	
albirostris. (Delphinus)	29		26
albirostris. (Lagenorhynchus)	29		26
albirostris. (Sus)	37	66	
albiventer. (Atophrax b.)	459		381
albiventer. (Oryzomys)	182	236	
albiventer. (Sorex b.)	459		381
albolimbatus. (Heteromys)	320	375	
albolimbatus. (Sciurus d.)	83		66
albomaculatum. (Phyllostoma)	527	708	
alborufescens. (Arvicola)	235		179
albus. (Canis)	375		
albus. (Canis l.)	375		
albus. (Dielidurus)	494	615	
albus. (Dysopes)	495	619	
albus. (Lupus)	375		
Alce	40		
Alce alces	41		
Alce gigas	41		
Alcelaphus	40		
Alces	40		37
Alces americanus	41		38
Alces gigas	41		38
Alces machlis	41		37

†Appendix.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Alces muswa</i>	41		38
<i>Alces palmatus</i>	41		38
<i>alces.</i> (<i>Alce</i>)	41		
<i>alces.</i> (<i>Cervus</i>)	41		
<i>alecto.</i> (<i>Dysopes</i>)	495	619	
<i>aleoutiensis.</i> (<i>Balæna</i>)	16		8
<i>alexandrinus.</i> (<i>Mus</i>)	129	164	118
<i>alfari.</i> (<i>Oryzomys</i>)	183	242	
<i>alfari.</i> (<i>Sciurus</i>)	60	99	
<i>alfari.</i> (<i>Sigmodontomys</i>)	191	254, 255	
<i>alfaroi.</i> (<i>Hesperomys</i>)	183, 184	242	
<i>algeriensis.</i> (<i>Delphinus</i>)		55	28
<i>Aliama</i>	22	47	16
<i>allamak.</i> (<i>Balæna</i>)	16		9
<i>allamandi.</i> (<i>Galera</i>)	418		
<i>allamandi.</i> (<i>Galictis</i>)	418	526	
<i>allamandi.</i> (<i>Grison</i>)	418	526	
<i>allegghanensis.</i> (<i>Putorius</i>)	428		
<i>allegghenensis.</i> (<i>Putorius</i>)	428		*511
<i>alleni.</i> (<i>Citellus t.</i>)	105		
<i>alleni.</i> (<i>Heteromys</i>)	321	376	256
<i>alleni.</i> (<i>Hodomys</i>)	226	297	
<i>alleni.</i> (<i>Lepus</i>)	353	433	288
<i>alleni.</i> (<i>Microtus</i>)	252		203
<i>alleni.</i> (<i>Neofiber</i>)	252		203
<i>alleni.</i> (<i>Neotoma</i>)	226	297	
<i>alleni.</i> (<i>Putorius</i>)	431		349
<i>alleni.</i> (<i>Rhogeëssa</i>)	489		
<i>alleni.</i> (<i>Rhogoëssa</i>)	489	602	
<i>alleni.</i> (<i>Rhogoëssa</i>)	489		
<i>alleni.</i> (<i>Sciurus</i>)	63	108	
<i>alleni.</i> (<i>Sigmodon</i>)	179	224	
<i>alleni.</i> (<i>Spermophilus</i>)	105		101
<i>alleni.</i> (<i>Zapus</i>)	326		261
<i>alleni.</i> (<i>Zapus t.</i>)	326, 327		261
<i>alex.</i> (<i>Peromyscus</i>)	135	175	
<i>allophylus.</i> (<i>Peromyscus</i>)	169	208	
<i>alnorum.</i> (<i>Microsorex</i>)	457		
<i>alnorum.</i> (<i>Sorex</i>)	457		
<i>aloga.</i> (<i>Blarina b.</i>)	462		
<i>alope.</i> (<i>Delphinus</i>)	33	58	
<i>Alopex</i>		465	
<i>Alouatta</i>	532	726	
<i>Alouatta palliata.</i>	533	726	
<i>Alouatta p. coibensis</i>	533	727	
<i>Alouatta p. mexicana.</i>	533	727	
<i>Alouatta villosa.</i>	532	726	
<i>Alouattinæ</i>	532	725	
<i>alpinus.</i> (<i>Mus</i>)	119		

*Supplement.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
alpinus. (Pteromys s.)	121		109
alpinus. (Scapanus)	469		393
alpinus. (Sciuropterus)	121, 124		
alpinus. (Sciuropterus v.)	121		109
alpinus. (Tamias)	94		
alpinus. (Tamias m.)	94		79
alpinus. (Thomomys)	283		229
alstoni. (Caluromys)	4	9	
alstoni. (Neotomodon)	225	294	
alstoni. (Sciurus)	63	108	
alticola. (Blarina)	466	561	
alticola. (Microtus)	244		192
alticola. (Perognathus)	307		249
alticola. (Reithrodontomys s.)	197	263	
alticola. (Rhithrodontomys s.)	197	263	
alticola. (Sigmodon)	179	231	
alticola. (Thomomys f.)	280	335	
alticolus. (Arvicola)	244		192
altifrontalis. (Ursus)	395		
altilaneus. (Peromyscus)	170	210	
altimiræ. (Lepus)	354	†745	
altimiræ. (Lepus m.)	354		
altivallis. (Thomomys)	287		230
ambarvalis. (Spilogale)	415		332
ambigua. (Spilogale)	415	521	
ambiguus. (Dipodomys)	289		234
ambiguus. (Dipodomys m.)	289		234
amblyceps. (Ursus)	394, 396		
amblyotis. (Phyllostoma)	572	659	
amblyotis. (Tonatia)	512	658	
americana. (Antilocapra)	51	82, 83	43
americana. (Antilope)	51	81, 82	43
americana. (Megaptera)	17		10
americana. (Mustela)	419	334	
americana. (Taxidea)	404		320
americanus. (Alces)	41		38
americanus. (Bison)	57		49
americanus. (Cervus)		68	
americanus. (Cervus d.)	43		
americanus. (Dipus)	325		259
americanus. (Jaculus)	324		257
americanus. (Lepus)	360		273
americanus. (Manatus)	14	37	6
americanus. (Meles t.)	404		
americanus. (Mus a.)	140		124
americanus. (Mustela)	419		
americanus. (Odocoileus)	43		39
americanus. (Odontocelus)	43	70, 75	
americanus. (Peromyscus)	140		124

†Appendix.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
americanus. (Sciurus).....	95		80
americanus. (Ursus).....	394		313
americanus. (Zapus h.).....	325		259
ammodytes. (Peromyscus l.).....	†542		
Ammomys.....	233	299	178
ammon. (Ovis).....	53		46
Ammospermophilus.....	96	139, 141	82, 85
amœnus. (Perognathus o.).....	307		
amœnus. (Sorex).....	449		371
amœnus. (Tamias).....	90		74
amoles. (Sigmodon a.).....	179	231	
Amphisorex lesueuri.....	446		366
amplexicaudata. (Glossophaga).....	576	672	
amplus. (Perognathus).....	306		247
amplus. (Peromyscus).....	167	205	
anarnacus. (Delphinus).....	21		16
anastasæ. (Peromyscus).....	143		
anastasiæ. (Peromyscus).....	143, 144		128
anastasiæ. (Scalops).....	470		390
Ancylodon.....	21		
andersoni. (Narwalus).....	24		19
andersoni. (Synaptomys).....	257		
angelensis. (Peromyscus b.).....	163	190	
angularis. (Thomomys).....	281		228
angulatum. (Tagassu).....	35	63	
angulatus. (Dicotyles).....	35	63	
angustata. (Beluga).....	25		
angusticeps. (Microtus).....	244		192
angusticeps. (Neotoma i.).....	217	285	162
angusticeps. (Oryzomys).....	187	245	
angusticeps. (Sorex).....	461		382
angustifrons. (Spilogale).....	416	521	
angustirostris. (Macrorhinus).....	440		359
angustirostris. (Mirounga).....	440	545	
angustirostris. (Perognathus p.).....	311	358	252
angustirostris. (Platygeomys t.).....	267	319	
Anisonyx.....	117, 126		102, 112
Anisonyx rufa.....	126		112
anitæ. (Thomomys f.).....	280	335	
annectens. (Citellus s.).....	100		
annectens. (Heteromys).....	317	371	
annectens. (Lutra).....	435	536	
annectens. (Neotoma f.).....	206		160
annectens. (Spermophilus s.).....	100		97
annellata. (Phoca).....	443, 444		364
annulatus. (Bassariscus).....	399	487	
annulatus. (Citellus).....	116	151, 152	
annulatus. (Paradoxurus).....	399	487	
annulatus. (Spermophilus).....	116	151	

†Appendix.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
anomalus. (Mus)	316		
Anotus	460	549, 556	382
Anoura	579	681	
Anoura ecaudata	520	682	
Anoura geoffroyi	519		
antarctica. (Balæna)	15, 16		9
antarctica. (Phoca)	438		
anthonyi. (Hesperomys)	137	185	141
anthonyi. (Neotoma)	214	283	
anthonyi. (Perognathus)	314	366	
anthonyi. (Peromyscus)	137	185	141
anthonyi. (Scapanus)	469	564	393
anthonyi. (Sciurus f.)	78	130	
anthonyi. (Sciurus g.)	78		57
Anthropoidea	532	723	
antillarum. (Glossophaga s.)	517	672	
antillarum. (Oryzomys)	188	247	
antillensis. (Dasypsecta)	337	407	
antillularum. (Nyctinomus)	499	629	
Antilocapra	51	81	43
Antilocapra americana	51	82, 83	43
Antilocapra a. mexicana	51	82	
Antilocapridæ	51	81	43
Antilope americana	51	81, 82	43
Antrozoinæ	491	605	396
Antrozous	491	605	396
Antrozous minor	492	607	
Antrozous pallidus	491	605	396
Antrozous p. pacificus	491	607	397
Anura	519	681	
Anura geoffroyi	519, 520	682	
Aotes	533	728	
Aotinæ	533	728	
Aotus	533	728	
Aotus azaræ	533	728	
Aotus rufipes	533	729	
Aotus vociferans	534	729	
apache. (Felis)	368		
apache. (Felis e.)	368	453	
apache. (Perognathus)	302, 303		244
apache. (Sicurus)	64	110	58
apatelius. (Oryzomys j.)	184	246	
apella. (Arvicola)	249		198
aphorodemus. (Microtus)	237		
aphrastus. (Thomomys)	282	336	
aphylla. (Reithronycteris)	521	688	
aphylla. (Rhithronycteris)	521	688	
apicalis. (Neotoma f.)	222		
apicalis. (Teonoma f.)	222		
Aplocerus	52		

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
Aplodontia	126		112
Aplodontia leporina	126		112
Aplodontia major	127		112
Aplodontia m. rainieri	127		112
Aplodontia olympica	127		114
Aplodontia pacifica	127		114
Aplodontia phæa	127		114
apricus. (Acodon t.)	204	274	
apricus. (Akodon t.)	204	274	
apus. (Pipistrellus h.)	481	583	
aquaticus. (Lepus)	340		277
aquaticus. (Oryzomys)	181	241	147
aquaticus. (Scalops)	470		388
aquaticus. (Scalopus)	469		
aquaticus. (Sorex)	469		388
aquilonius. (Fiber z.)	253		213
Aræosciurus	59, 62	94, 106	50, 58
araneus. (Sorex)	446	548	366
arboreus. (Peromyscus)	135	170	
Arctibeus falcatus	527	707	
Arctibeus leucomus	528	712	
arctica. (Cervus t.)	40		
arcticeps. (Onychomys)	132		121
arcticus. (Hesperomys l.)	146		131
arcticus. (Lepus)	358		270
arcticus. (Lepus t.)	358		270
arcticus. (Leucopleurus)	29		25
arcticus. (Orca)	26	51	21
arcticus. (Peromyscus)	146		131
arcticus. (Putorius)	428		345
arcticus. (Rangifer)	40		37
arcticus. (Rosmarus)	439		358
arcticus. (Sorex p.)	447		368
Arctocephalus	438	543	357
Arctocephalus monteriensis	437		357
Arctocephalus townsendi	438	544	
Arctogale	423, 426	529	338, 343
Arctomys	119		104
Arctomys avarus	120		
Arctomys beecheyi	114		88
Arctomys caligatus	120		106
Arctomys columbianus	111		91
Arctomys dacota	120		106
Arctomys douglasii	115		89
Arctomys flaviventer	120		106
Arctomys franklinii	105		101
Arctomys guttatus	111		92
Arctomys hoodi	104		
Arctomys ignavus	119		
Arctomys kennicottii	107		89

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Arctomys latrans</i>	117		102
<i>Arctomys lewisii</i>	118		104
<i>Arctomys ludovicianus</i>	117		102
<i>Arctomys melanopus</i>	120		106
<i>Arctomys missouriensis</i>	117		102
<i>Arctomys monax</i>	119		105
<i>Arctomys m. canadensis</i>	120		106
<i>Arctomys m. ignavus</i>	119		105
<i>Arctomys olympus</i>	121		107
<i>Arctomys parryi</i>	107		89
<i>Arctomys p. erythroglutea</i>	108		91
<i>Arctomys p. phœognatha</i>	107		89
<i>Arctomys pruinosus</i>	120		106
<i>Arctomys richardsonii</i>	111		92
<i>Arctophoca</i>	438		
<i>Arctopithecus</i>	9	21	
<i>Arctopithecus castaneiceps</i>	9	22	
<i>Arctopithecus griseus</i>	9	23	
<i>arctos.</i> (<i>Ursus</i>)	390	479	309
<i>arenacea.</i> (<i>Neotoma</i>)	214	283	
<i>arenarius.</i> (<i>Geomys</i>)	264	310	219
<i>arenarius.</i> (<i>Perognathus</i>)	312		
<i>arenarius.</i> (<i>Perognathus p.</i>)	312	358	
<i>arenarius.</i> (<i>Peromyscus e.</i>)	137	191	136
<i>arenicola.</i> (<i>Onychomys t.</i>)	133	168	122
<i>arenicola.</i> (<i>Perognathus p.</i>) ..	305		246
<i>arenivagus.</i> (<i>Dipodomys m.</i>) ..	290	347	
<i>arens.</i> (<i>Citellus s.</i>)	101		
<i>arens.</i> (<i>Spermophilus s.</i>)	101		
<i>argentata.</i> (<i>Vulpes f.</i>)	380		304
<i>argentata.</i> (<i>Vulpes p.</i>)	380		
<i>argentatus.</i> (<i>Canis</i>)	380		304
<i>argentatus.</i> (<i>Scalops</i>)	470		389
<i>aries.</i> (<i>Ovis</i>)	53	83	
<i>Ariteus</i>	526	704	
<i>Ariteus flavescens</i>	527	707	
<i>arizonæ.</i> (<i>Lepus</i>)	342, 349, 350	423	285
<i>arizonæ.</i> (<i>Lepus s.</i>)	349		
<i>arizonæ.</i> (<i>Neotoma</i>)	223		165
<i>arizonæ.</i> (<i>Peromyscus a.</i>)	140	186	125
<i>arizonæ.</i> (<i>Peromyscus l.</i>)	140		
<i>arizonæ.</i> (<i>Peromyscus t.</i>)	141	186	
<i>arizonæ.</i> (<i>Sigmodon h.</i>)	176	228	144
<i>arizonæ.</i> (<i>Sitomys a.</i>)	140		125
<i>arizonæ.</i> (<i>Spilogale</i>)	414		331
<i>arizonæ.</i> (<i>Spilogale p.</i>)	414		331
<i>arizonæ.</i> (<i>Teonoma</i>)	223		
<i>arizonensis.</i> (<i>Cynomys</i>)	118		103
<i>arizonensis.</i> (<i>Microtus m.</i>) ..	238		183
<i>arizonensis.</i> (<i>Putorius</i>)	431		349

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
arizonensis. (Reithrodontomys).....	196		
arizonensis. (Rhithrodontomys).....	196		
arizonensis. (Sciurus).....	61, 63	102, 109	59
Arizostus	12	31	
armatus. (Citellus).....	112		
armatus. (Spermophilus).....	112		93
arnuxii. (Berardius).....	23		
arquatus. (Vespertilio).....	483	587	410
arsipus. (Vulpes).....	385		
artemisia. (Lepus)		425	
artemisiae. (Peromyscus a.).....	141		126
artemisiae. (Peromyscus l.).....	141		
artemisiae. (Sitomys a.).....	141		126
Artibeus.....	522, 526	691, 704	
Artibeus achradoophilus	527	707	
Artibeus carpolegus	522, 523	694	
Artibeus coryi	522	693	
Artibeus eva	524	697	
Artibeus fallax	524	695	
Artibeus insularis.....	522		
Artibeus intermedius	523	694	
Artibeus jamaicensis	522	693	
Artibeus parvipes.....	523	694	
Artibeus planirostris.....	524	695	
Artibeus sulphureus.....	527	707	
Artibeus vittatus	526	704	
Artibeus watsoni	524	696	
Artibeus yucatanicus.....	523		
artus. (Perognathus).....	313		
artus. (Perognathus g.)	313	365	
arvalis. (Mus).....	233	298	178
Arvicola	233, 250	298	178, 200
Arvicola alborufescens	235		179
Arvicola alticolus	244		192
Arvicola apella	249		198
Arvicola austerus	248		196
Arvicola a. minor	248		197
Arvicola aztecus	237		182
Arvicola borealis	247		194
Arvicola breweri.....	238		
Arvicola californica	240		186
Arvicola chrotorrhinus	246		
Arvicola cinnamomea.....	248		196
Arvicola curtata	247		195
Arvicola decurtata.....	247		
Arvicola dekayi	235		180
Arvicola drummondi	236		181
Arvicola edax	241		187
Arvicola ferrugineus	175		144
Arvicola floridana	209, 210		

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Arvicola fulva</i>	235		180
<i>Arvicola gapperi</i>	230		172
<i>Arvicola haydenii</i>	248		197
<i>Arvicola helvolus</i>	259		208
<i>Arvicola hirsutus</i>	235		175
<i>Arvicola hortensis</i>	175		144
<i>Arvicola insperatus</i>	236		180
<i>Arvicola kennicotti</i>	249		198
<i>Arvicola leucophæus</i>	245		192
<i>Arvicola longicaudus</i>	243		191
<i>Arvicola longirostris</i>	238		182
<i>Arvicola mexicanus</i>	245	301	
<i>Arvicola microcephalus</i>	236		181
<i>Arvicola modesta</i>	236		180
<i>Arvicola mogollonensis</i>	246		192
<i>Arvicola montanus</i>	238		182
<i>Arvicola mordax</i>	244		191
<i>Arvicola nanus</i>	239		184
<i>Arvicola nasuta</i>	235		180
<i>Arvicola nuttalli</i>	160		140
<i>Arvicola occidentalis</i>	243		190
<i>Arvicola oneida</i>	235		180
<i>Arvicola operarius</i>	241		187
<i>Arvicola oregoni</i>	250		200
<i>Arvicola oryzivora</i>	180		147
<i>Arvicola pallidus</i>	247		195
<i>Arvicola palustris</i>	234		179
<i>Arvicola pauperrima</i>	247		195
<i>Arvicola phæus</i>	245	302	
<i>Arvicola p. quasiater</i>	249		
<i>Arvicola richardsoni</i>	250		200
<i>Arvicola riparius</i>	234		179
<i>Arvicola r. longipilis</i>	235		180
<i>Arvicola rufescens</i>	235		180
<i>Arvicola rufidorsum</i>	235		180
<i>Arvicola scalopsoides</i>	249		198
<i>Arvicola terrænovæ</i>	238		
<i>Arvicola tetramerus</i>	243		190
<i>Arvicola texiana</i>	175		144
<i>Arvicola townsendi</i>	243		190
<i>Arvicola trimucronatus</i>	258		207
<i>Arvicola trowbridgi</i>	240		186
<i>Arvicola xanthognatha</i>	246		193
<i>arvicoloides</i> . (<i>Microtus r.</i>)	251		202
<i>Asagis</i>	1	5	
<i>Ascomys</i>	261	310	215
<i>Ascomys canadensis</i>	262		217
<i>Ascomys umbrinus</i>	272		
<i>asellus</i> . (<i>Lepus</i>)	355	435	
<i>Astromyctes</i>	467		394
<i>Astromydes</i>	467		394

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
astuta. (Bassaris).	397	484	
astutus. (Bassariscus).	397, 398	484	316
Atalapha	485	591	411
Atalapha borealis	485	593	
Atalapha b. seminola	486		
Atalapha frantzii	486	594	
Atalapha mexicana.	486	594	
Atalapha pfeifferi	486	593	
Atalapha teliotis.	486	593	413
Ateles	534	732	
Ateles albifrons.	535	733	
Ateles ater.	535	734	
Ateles fuliginosus	534	733	
Ateles geoffroyi.	535	733	
Ateles griseus.	535	734	
Ateles hybridus	535	733	
Ateles melanochir.	535	733	
Ateles ornatus.	535	733	
Ateles pan	534	733	
Ateles rufiventris	535	734	
Ateles variegatus	535	733	
Ateles vellerosus.	534	733	
ater. (Ateles).	535	734	
ater. (Lupus).	376		
ater. (Orca).	26		
athabascæ. (Bison a.)	57		49
athabascæ. (Bison b.)	57		
Atophrax.	459		380
Atophrax benderii.	459		381
Atophrax b. albiventer	459		381
Atophrax b. palmeri	459		381
atra. (Orca).			22
atrata. (Mustela)	422		335
atricapillus. (Citellus v.)	114	150	
atricapillus. (Spermophilus g)	114	150	
atronasus. (Dipodomys m.)	290	348	
atrovarius. (Thomomys)	283	338	
attwateri. (Geomys b.)	263		219
attwateri. (Lepus a.)	340	414	278
attwateri. (Neotoma)	210		
attwateri. (Neotoma f.)	210		157
attwateri. (Peromyscus)	154		135
auduboni. (Lepus)	345	422	283
auduboni. (Ovis c.)	53		*484
auduboni. (Sciurus)	65		52
auduboni. (Vespertilio)	480		407
Aulacomys.	234		179
Aulacomys arvicoloides.	251		202
aurantius. (Reithrodontomys a.)	198		153

* Supplement.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
aurantius. (Reithrodontomys m.)	198		153
aurantius. (Rhithrodontomys m.)	198		
aureigaster. (Sciurus)	68, 74	115, 124	
aureogaster. (Sciurus)	69, 74	116, 124	
aureus. (Reithrodontomys t.)	201	269	
aureus. (Rhithrodontomys t.)	201	269	
aureus. (Thomomys)	285		229
auricularis. (Microtus p.)	249		198
auripectus. (Peromyscus)	153		134
auripectus. (Sitomys)	153		134
aurispinosus. (Dysopes)	499	627	
aurita. (Didelphys)	8	17	
aurita. (Lonchorina)	507, 508	650	
auritus. (Cervus)	48	76	42
auritus. (Chrotopterus)	512	656	
auritus. (Dysopes)	499	627	
auritus. (Peromyscus)	170	209	
auritus. (Vampyrus)	511, 512	656	
aurogaster. (Sciurus)	68	115	
austerulus. (Sigmodon)	180	231	
austerus. (Arvicola)	248		196
austerus. (Hesperomys)	149		132
austerus. (Microtus)	248		196
austerus. (Peromyscus)	149		132
australis. (Balæna)	15, 16		8, 9
australis. (Eubalæna)	15		8
australis. (Hydrodamalis)		36	
australis. (Manatus)	14	37	6
australis. (Pipistrellus h.)	481	584	
australis. (Reithrodontomys)	194		
australis. (Rhithrodontomys)	194	259	
australis. (Scalops a.)	470		
australis. (Scalopus a.)	470		390
austrinus. (Geomys f.)	262		
austrinus. (Geomys t.)	262		217
austroriparius. (Vespertilio l.)	478		402
avarus. (Arctomys f.)	120		†427
avarus. (Marmota f.)	120		
avia. (Mephitis)	407		327
avia. (Mephitis m.)	407		
azaræ. (Aotus)	533	728	
azteca. (Carollia)	515	669	
aztecus. (Arvicola)	237		182
aztecus. (Felis h.)	369	454	
aztecus. (Hesperomys)	162	184	
aztecus. (Lepus f.)	344	418	
aztecus. (Lepus s.)	344		
aztecus. (Microtus)	237		182
aztecus. (Molossus)	495	620	

† Appendix.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
aztecus. (<i>Oryzomys</i> c.)	183	245	
aztecus. (<i>Peromyscus</i>)	162	184	
aztecus. (<i>Potos</i> f.)	404	500	
aztecus. (<i>Reithrodontomys</i>)	193	259	151
bachmani. (<i>Lepus</i>)	352	429	
bachmani. (<i>Lepus</i> f.)	352		281
bachmani. (<i>Lepus</i> s.)	343		
badius. (<i>Citellus</i> t.)	105		
badius. (<i>Peromyscus</i> y.)	163	194	
badius. (<i>Spermophilus</i> t.)	105		101
bahamensis. (<i>Nyctinomus</i>)	500	630	
bahamensis. (<i>Vespertilio</i> f.)	484	588	
baileyi. (<i>Felis</i> r.)	372	459	297
baileyi. (<i>Lepus</i>)	349		284
baileyi. (<i>Lynx</i>)	372		
baileyi. (<i>Neotoma</i>)	210		156
baileyi. (<i>Nebtoma</i> c.)	210		
baileyi. (<i>Perognathus</i>)	309	361	251
baileyi. (<i>Sciurus</i> h.)	81		63
baileyi. (<i>Sigmodon</i>)	177	226	
baileyi. (<i>Sigmodon</i> h.)	177	226	
baileyi. (<i>Thomomys</i>)	286		
Baiomys	135	170	123
Baiosciurus	59, 60	94, 100	50
bairdi. (<i>Berardius</i>)	24		*481
bairdi. (<i>Delphinus</i>)	30	55	28
bairdi. (<i>Lepus</i> a.)	361		274
bairdi. (<i>Microtus</i>)	250		200
bairdi. (<i>Sorex</i>)	451		373
bairdi. (<i>Tapirella</i>)	58	87	
bairdii. (<i>Elasmognathus</i>)	58	87	
bairdii. (<i>Lepus</i>)	361		274
Balæna	15		7
Balæna aleoutiensis	16		8
Balæna allamak	16		9
Balæna antarctica	15, 16		9
Balæna australis	15, 16		8, 9
Balæna a. biscayensis	15		8
Balæna biscayensis	15		8
Balæna boops	17, 19	40	13
Balæna borealis	20		13
Balæna britannicus	15		8
Balæna cisarctica	15		8
Balæna glacialis	15		8
Balæna groenlandica	16		8
Balæna keporkak	17		10
Balæna kuzira	16		9
Balæna longimana	17		10
Balæna musculus	20		13

*Supplement.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Balæna mysticetus</i>	16	39	8
<i>Balæna m. pittekajensis</i>	16		8
<i>Balæna nodosa</i>	17		10
<i>Balæna physalus</i>	19, 20		13
<i>Balæna rostrata</i>	21		11, 16
<i>Balæna sieboldii</i>	15		8
<i>Balænidæ</i>	15	38, 39	7
<i>Balæninæ</i>	15		
<i>Balænoptera</i>	18	42	11
<i>Balænoptera acuto-rostrata</i>	18		11
<i>Balænoptera borealis</i>	19, 20		13
<i>Balænoptera davidsoni</i>	19	42	12
<i>Balænoptera gibbar</i>	18	42	
<i>Balænoptera mondini</i>	19		
<i>Balænoptera musculus</i>	20		13
<i>Balænoptera physalus</i>	19, 20		13
<i>Balænoptera roqual</i>	19		13
<i>Balænoptera stejneri</i>	23		
<i>Balænoptera sulfurea</i>	20	38, 43	14
<i>Balænoptera velifera</i>	19	43	12
<i>Balænoptera v. copei</i>	19		13
<i>Balænopterinæ</i>	18	42	11
<i>Balantiopteryx</i>	493	611	
<i>Balantiopteryx infusca</i>	493	611	
<i>Balantiopteryx plicata</i>	493	611	
<i>baliolus</i> . (<i>Peromyscus s.</i>).....	139		141
<i>baliolus</i> . (<i>Sciurus y.</i>).....	75	126	
<i>balteatus</i> . (<i>Delphinus</i>).....	31	55	28
<i>banderanus</i> . (<i>Peromyscus</i>).....	163	189	
<i>bangsi</i> . (<i>Lepus a.</i>).....	358		270
<i>bangsi</i> . (<i>Perognathus l.</i>).....	305		
<i>bangsi</i> . (<i>Perognathus p.</i>).....	305		245
<i>bangsi</i> . (<i>Sciuropterus a.</i>).....	124		
<i>bangsi</i> . (<i>Vulpes f.</i>).....	381		
<i>bangsi</i> . (<i>Vulpes p.</i>).....	381		†434
<i>bangsi</i> . (<i>Vulpes r.</i>).....	381		
<i>barbara</i> . (<i>Mustela</i>).....	418		
<i>barbata</i> . (<i>Phoca</i>).....	441		362
<i>barbatus</i> . (<i>Vespertilio</i>).....	502	634	
<i>barberi</i> . (<i>Sciurus</i>).....	67	†741	
<i>barberi</i> . (<i>Sciurus a.</i>).....	67		
<i>barrowensis</i> . (<i>Citellus</i>).....	110		91
<i>barrowensis</i> . (<i>Spermophilus</i>).....	110		91
<i>Bassaricyon</i>	400	487	
<i>Bassaricyon gabbi</i>	400	487, 489	
<i>Bassaricyon gabbi</i>	400		
<i>Bassaris</i>	397	483	315
<i>Bassaris astuta</i>	397	484	
<i>Bassaris monticola</i>	398	487	

†Appendix.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Bassaris raptor</i>	398		
<i>Bassaris variabilis</i>	398	487	
<i>Bassariscus</i>	397	482	315
<i>Bassariscus albipes</i>	399	486	
<i>Bassariscus annulatus</i>	399	487	
<i>Bassariscus astutus</i>	397, 398	484	316
<i>Bassariscus a. flavus</i>	398	485	316
<i>Bassariscus a. raptor</i>	398		316
<i>Bassariscus f. oregonus</i>	398		317
<i>Bassariscus saxicola</i>	398	485	
<i>Bassariscus sumichrasti</i>	398		
<i>Bassariscus s. notinus</i>	399	485	
<i>battyi</i> . (<i>Didelphys m.</i>)	7	17	
<i>battyi</i> . (<i>Lepus g.</i>)	355	433	
<i>battyi</i> . (<i>Odocoileus</i>)	45	71	
<i>battyi</i> . (<i>Odontocœlus</i>)	45	71	
<i>beatae</i> . (<i>Peromyscus</i>)	162	184	
<i>beecheyi</i> . (<i>Arctomys</i>)	114		88
<i>beecheyi</i> . (<i>Citellus v.</i>)	114	150	
<i>beecheyi</i> . (<i>Spermophilus g.</i>)	114		88
<i>belcheri</i> . (<i>Sciurus</i>)	82		65
<i>beldingi</i> . (<i>Citellus</i>)	113		
<i>beldingi</i> . (<i>Spermophilus</i>)	113		94
<i>bella</i> . (<i>Neotoma</i>)	218		†429
<i>belli</i> . (<i>Sorex</i>)			368
<i>bellicosa</i> . (<i>Megaptera n.</i>)	18	41	
<i>bellus</i> . (<i>Peromyscus</i>)	154		135
<i>belti</i> . (<i>Sciurus b.</i>)	77	128	
<i>Beluga angustata</i>	25		
<i>Beluga concreta</i>	25		20
<i>Beluga declivis</i>	25		20
<i>Beluga rhinodon</i>	25		19
<i>beluga</i> . (<i>Delphinapterus</i>)	25		19
<i>belzebul</i> . (<i>Simia</i>)	532	726	
<i>bendirii</i> . (<i>Atophyrax</i>)	459		381
<i>Benedenia</i>	18	42	11
<i>Benedenia knoxii</i>	20		13
<i>bennetti</i> . (<i>Lepus</i>)	357	437	
<i>Berardius</i>	23		*481
<i>Berardius arnuxii</i>	23		
<i>Berardius bairdi</i>	24		*481
<i>beringensis</i> . (<i>Citellus</i>)	110		
<i>beringensis</i> . (<i>Spermophilus</i>)	110		91
<i>beringensis</i> . (<i>Vulpes</i>)	383		
<i>berlandieri</i> . (<i>Blarina</i>)	462	558	
<i>berlandieri</i> . (<i>Blarina b.</i>)	462	558	
<i>berlandieri</i> . (<i>Sigmodon</i>)	175	227	
<i>berlandieri</i> . (<i>Sigmodon h.</i>)	175	225, 227	

*Appendix.

*Supplement.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
berlandieri. (Taxidea)	405		
berlandieri. (Taxidea a.)	405		321
berlandieri. (Taxidea t.)	405	503	
bernardinus. (Citellus c.)	107		84
bernardinus. (Eptesicus f.)	483		
bernardinus. (Spermophilus)	107		
bernardinus. (Spermophilus c.)	107		
bernardinus. (Vespertilio f.)	483	589	
bicolor. (Mephitis)	412		328
bicolor. (Phyllostoma)	515	669	
bidens. (Mesopiodon)	23		*480
bidens. (Physeter)	23		19
bidens. (Vampyrus)	512	658	
bidentatus. (Delphinus)	21		16
bilabiatum. (Phyllostoma)	528	712	
bilabiatum. (Pygoderma)	528	712	
bilineata. (Saccopteryx)	493	610	
bilineatus. (Urocryptus)	493	*610	
bilobatum. (Uroderma)	524	697	
bimaculatus. (Perognathus)	302		244
bimaculatus. (Perognathus f.)	302		244
biologiæ. (Galera b.)	418		
biologiæ. (Galictis b.)	418		
biologiæ. (Grison b.)	418	524	
biscayensis. (Balæna)	15		8
biscayensis. (Balæna a.)	15		8
bishopi. (Lepus)	362		276
Bison	57		48
Bison americanus	57		49
Bison a. athabascæ	57		149
Bison bison	57		
Bison b. athabascæ	57		
bison. (Bison)	57		
bison. (Bos)	57		
bisulcus. (Equus)	50		
bivittata. (Tamandua)	10	27	
blainvillii. (Mormoops)	506	646, 649	
blainvillii. (Mormops)	507	649	419
blandus. (Peromyscus l.)		182	
blandus. (Peromyscus s.)	142	182	
Blaria	460		381
Blarina	460	555	381
Blarina alticola	466	561	
Blarina berlandieri	462	558	
Blarina breviceauda	460, 461		382
Blarina b. aloga	462		
Blarina b. berlandieri	462	558	
Blarina b. carolinensis	461		383
Blarina b. compacta	462		

*Supplement

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
Blarina b. hulophaga	461		383
Blarina b. hylophaga	461		
Blarina b. peninsulæ	461		383
Blarina b. talpoides	461		
Blarina costaricensis	461		382
Blarina c. peninsulæ	461		
Blarina exilipes	463		384
Blarina eximius	463		384
Blarina floridana	463		385
Blarina fossor	466	562	
Blarina magna	466	562	
Blarina mayensis	465	561	
Blarina mexicana	464	560	
Blarina m. goldmani	465	560	
Blarina m. machetes	465	561	
Blarina m. peregrina	465	560	
Blarina micrura		559	
Blarina nelsoni	465	561	
Blarina nigrescens	463	558	
Blarina obscura	464	559	
Blarina oreophila	464	559	
Blarina orophila	464	559	
Blarina parva	463		384
Blarina pergracilis	463	559	
Blarina soricina	464	560	
Blarina telmalestes	462		384
Blarina tropicalis	464	559	
bocourtianus. (Macrotus)	510	654	
bocourtianus. (Otopterus)	510	654	
boliviense. (Akodon)	204	272	
bombifrons. (Phyllonycteris)	520	685	
bonasus. (Bos)			48
boops. (Balæna)	17, 19	40	13
boothi. (Chilonycteris)	505	642	
boothia. (Sciurus)	75, 76	127, 128	
boquetensis. (Sciurus)	60	100	
borealis. (Arvicola)	247		194
borealis. (Atalapha)	485	593	
borealis. (Balæna)	20		13
borealis. (Balanoptera)	19, 20		13
borealis. (Cystophora)	440		360
borealis. (Delphinapteris)	32		30
borealis. (Geomys)	275, 278		224, 227
borealis. (Lasiurus)	485	592	411
borealis. (Lepus)	360		273
borealis. (Lissodelphis)	32		30
borealis. (Microtus)	247		194
borealis. (Odocoileus a.)	43		
borealis. (Odontocœlus a.)	43		
borealis. (Tamias a.)	92		77

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
borealis. (Tamias q.)	92		77
borealis. (Urocyon c.)	388		
borealis. (Vespertilio)	485	592	411
boria. (Mustela)	421		
Boriotkon	260		
borucæ. (Sigmodon)	178	224	
borucæ. (Sigmodon h.)	178	224, 232	
Bos bison	57		
Bos bonasus			48
Bos canaliculatis	56		48
Bos moschatus	55		
Bos pallasi	56		48
bottæ. (Oryctomys)	274		223
bottæ. (Thomomys)	274		223
Bovidæ	52	83	44
boylii. (Hesperomys)	150	195	132
boylii. (Peromyscus)	150	195	132
brachycium. (Phocæna)	25	48	20
Brachylagus	340, 351		287
brachyotum. (Phyllostoma)	515	669	
Brachyotus	473	571	400
Brachyphylla	521	688	
Brachyphylla cavernarum	521	689	
Brachyphylla nana	522	690	
brachypterus. (Globicephalus)	28	53	
brachypterus. (Globiocephalus)	28	53	24
Brachysorex	460	555	381
Bradypodidæ	9	19	4
Bradypus	9	21	
Bradypus castaneiceps	9	22	
Bradypus didactylus	9	19	
Bradypus griseus		24	
Bradypus infuscatus	9	23	
Bradypus tridactylus	9	21	
brasiliensis. (Felis)	367	449	
brasiliensis. (Nyctinomus)	501	629	417
brevicauda. (Blarina)	460, 461		382
brevicauda. (Zygodontomys)		254	
brevicaudum. (Hemiderma)	515	669	
brevicaudum. (Phyllostoma)	515	668	
brevicaudus. (Evotomys)	231		174
brevicaudus. (Evotomys g.)	231		
brevicaudus. (Onychomys l.)	130		120
brevicaudus. (Sorex)	460		382
brevicaudus. (Spermophilus c.)	107		84
breviceps. (Cogia)	22	46	
breviceps. (Geomys)	263		218
breviceps. (Kogia)	22		*479
breviceps. (Physeter)	22	46	*479

* Supplement.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
brevimanus. (Chilonatalis)	502, 503	636	
brevimanus. (Natalus)	502		
brevinasus. (Perognathus p.)	305	355	246
breweri. (Arvicola)	238		
breweri. (Microtus)	238		182
breweri. (Parascalops)	471		394
breweri. (Scalops)	471		394
bridgeri. (Thomomys)	279		
britannicus. (Balæna)	15		8
brochus. (Syntheosciurus)	59	91	
brochus. (Synthetosciurus)	59	91	
browni. (Capromys)	331	393	
browni. (Felis a.)	369		
browni. (Sciurus)	60	100	
brumalis. (Mustela)	420		336
brumalis. (Mustela a.)	420		336
brunneus. (Peromyscus m.)	136	176	
bryanti. (Neotoma)	208	280	
bryanti. (Perognathus)	316		
bryanti. (Perognathus s.)	316	364	
buccatus. (Citillus)		149	
buckleyi. (Citellus v.)	115		
buckleyi. (Spermophilus)	115		89
buckleyi. (Spermophilus g.)	115		89
bulbivorum. (Diplostoma)	272		
bulbivorus. (Thomomys)	272		222
bullaris. (Tylomys)	173	219	
bullata. (Nasua n.)	403	497	
bullata. (Neotoma m.)	213	282	158
bullatus. (Peromyscus)	158	193	
bullatus. (Synaptomys)	258		
bulleri. (Geomys)	268	320	
bulleri. (Heteromys)	317	369	
bulleri. (Macrotus)	510	654	
bulleri. (Oryzomys)	185	239	
bulleri. (Otopterus)	510	654	
bulleri. (Pappogeomys)	268	320	
bulleri. (Tamias)	88	136, 137	
bulleri. (Tamias a.)	88		
burrus. (Proechimys)	330	388, 396	
burrus. (Proechinomys)	330	388	
bursarius. (Geomys)	262, 263		217
bursarius. (Mus)	262		217
bursarius. (Saccophorus)	262		
butskopf. (Delphinus)	21		16
Cabassous.	12	31	4
Cabassous centralis.	12	32	
Cabassous novemcinctus.	13	34	4
cabezonæ. (Perodipus)	295		

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
cabezonæ. (Thomomys)	286		
cacabatus. (Peromyscus)	171	211	
Cachicamus.....	13	32	
cacomitli. (Felis)	367		
cacomitli. (Felis j.)	367	451	
cæcus. (Dysopes)		627	
cagottis. (Canis)	378	465	
cagottis. (Lyciscus)	378		
calcaratum. (Phyllostoma)	515	669	
californiana. (Otaria)	437		356
californianus. (Ovis)	53		46
californianus. (Zalophus)	437		356
californica. (Arvicola)	240		186
californica. (Didelphis)	6	15	
californica. (Didelphys)	6		3
californica. (Felis r.)	372	458	298
californica. (Haplodon)	127		
californica. (Haplodontia r.)	127		112
californica. (Lepus)	357	437	
californica. (Neotoma)	216		
californicus. (Cariacus h.)	49	77	43
californicus. (Cervus m.)	49	77	
californicus. (Dipodomys)	291		236
californicus. (Evotomys)	232		176
californicus. (Lepus)	357	437	291
californicus. (Lynx r.)	372	458	298
californicus. (Macrotus)	509	653	420
californicus. (Microdipodops)	300		
californicus. (Microtus)	240		186
californicus. (Molossus)	498		
californicus. (Mus)	137	203	137
californicus. (Myotis)	476	578	
californicus. (Nyctinomus b.)	500		
californicus. (Odocoileus h.)	49	77	43
californicus. (Odontocælus h.)	49	76, 77, 78	
californicus. (Otopterus)	509, 510	653	420
californicus. (Perognathus)	315		254
californicus. (Peromyscus)	137, 138	203	137
californicus. (Promops)	498		421
californicus. (Scalops)	468		392
californicus. (Scapanus)	468		392
californicus. (Sciuropterus a.)	122		
californicus. (Sciuropterus v.)	122		110
californicus. (Sciurus)	114		
californicus. (Sciurus h.)	83		66
californicus. (Sorex)	452		374
californicus. (Urocyon c.)	387	477	308
californicus. (Ursus h.)	392		
californicus. (Vespertilio)	476	578	
caligatus. (Arctomys)	120		106

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
caliginosus. (<i>Hesperomys</i>)	190	253	
callida. (<i>Dasyprocta</i>)	336	406	
callipeplus. (<i>Tamias</i>)	89		73
callistus. (<i>Perognathus</i>)	304		245
Callitrichidæ	532		
callitrichus. (<i>Cercopithecus</i>)	536		
Callocephalus dimidiatus.....	444		364
Callocephalus discolor.....	444		364
Callorhinus.....	438		357
Callorhinus alascanus.....	438		
Callospermophilus.....	96, 105	139, 147	82
Callospermophilus c. trinitatis.....	107		
Callospermophilus madrensis.....	106	147	
Callotaria.....	438		357
Callotaria alascana.....	438		
Callotaria ursina.....	438		357
callotis. (<i>Lepus</i>)	353	431	288
Calocephalus.....	441	541	361
Calomys.....	135	170	123
Caluromys.....	4	9	
Caluromys alstoni.....	4	9	
Caluromys derbianus.....	4	9	
Caluromys l. pallidus.....	4	10	
campestri. (<i>Lepus</i>)	360		273
campestris. (<i>Lepus</i>)	362		277
campestris. (<i>Neotoma</i>)	218		156
campestris. (<i>Zapus</i> h.)	325		259
Campicola	234		
canadensis. (<i>Arctomys</i> m.)	120		106
canadensis. (<i>Ascomys</i>)	262		217
canadensis. (<i>Castor</i>)	128		116
canadensis. (<i>Cervus</i>)	41, 42		34
canadensis. (<i>Cervus</i> e.)	41		
canadensis. (<i>Delphinus</i>)	25		19
canadensis. (<i>Dipus</i>)	324		257
canadensis. (<i>Felis</i>)	370		295
canadensis. (<i>Glis</i>)	120		106
canadensis. (<i>Lutra</i>)	433		352
canadensis. (<i>Lynx</i>)	370		
canadensis. (<i>Marmota</i> m.)	120		
canadensis. (<i>Mustela</i>)	422, 423		335, 338
canadensis. (<i>Mustela</i> l.)	433		
canadensis. (<i>Ovis</i>)	53		46
canadensis. (<i>Peromyscus</i>)	144		128
canadensis. (<i>Scalops</i>)	468, 470		388, 391
canadensis. (<i>Sitomys</i> a.)	144		128
canaliculatis. (<i>Bos</i>)	56		48
canaster. (<i>Galera</i>)	419		
canaster. (<i>Galictis</i>)	419	526	
canaster. (<i>Grison</i>)	419	526	

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
carnivorus. (Procyon)	402	492	
carnivorus. (Ursus)	402	492	
canescens. (Citellus)	101		
canescens. (Didelphis)	2	6	
canescens. (Eutamias)	86		
canescens. (Marmosa)	2	6	
canescens. (Microtus n.)	239		185
canescens. (Neotoma m.)	208		
canescens. (Perognathus i.)	313		
canescens. (Perognathus n.)	313	365	
canescens. (Spermophilus)	101		97
canescens. (Tamias)	86		
canicaudus. (Eutamias)	93		
canicaudus. (Microtus)	239		185
canicaudus. (Tamias)	93		
caniceps. (Eutamias)	93		
caniceps. (Tamias)	93		426
caniculus. (Lepus f.)	344		283
Canidae	374	464	299
canina. (Peroptryx)	493	614	
Caninae	374	464	299
caninus. (Vespertilio)	493	614	
canipes. (Eutamias c.)	91		
canipes. (Tamias c.)	91		
Canis	374	464	299
Canis albus	375		
Canis argentatus	380		304
Canis cagottis	378	465	
Canis cinereo-argentatus	384		
Canis cinereoargenteus	386		
Canis clepticus	379	467	
Canis decussatus	380		304
Canis estor	378	469	302
Canis familiaris	374	464	
Canis frustror	377		302
Canis fulvus	380		304
Canis goldmani	378		
Canis griseus	375, 386		307
Canis impavidus	379	468	
Canis latrans	376		300
Canis lestes	377	465	301
Canis lupus			299
Canis l. albus	375		
Canis l. griseus	375		300
Canis l. occidentalis	376		
Canis mearnsi	377	468	302
Canis mexicanus	375, 376	471	
Canis microdon	379	469	
Canis microtus	384		
Canis nebrascensis	377		301

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Canis nubilus</i>	376		300
<i>Canis occidentalis</i>	375, 376		300
<i>Canis ochropus</i>	379	469	303
<i>Canis pallidus</i>	377		301
<i>Canis pambasileus</i>	374		
<i>Canis peninsulæ</i>	378	466	
<i>Canis velox</i>	384		
<i>Canis vigilis</i>	378	467	
<i>Canis virginianus</i>	386		307
<i>Canis vulpes</i>	380	471	
<i>canus</i> . (<i>Citellus m.</i>).....	112		
<i>canus</i> . (<i>Heteromys</i>).....	321	375	
<i>canus</i> . (<i>Lepus a.</i>).....	359		
<i>canus</i> . (<i>Liomys</i>).....	321	375	
<i>canus</i> . (<i>Odocoileus h.</i>).....	50	78	
<i>canus</i> . (<i>Odontocœlus h.</i>).....	50	78	
<i>canus</i> . (<i>Onychomys t.</i>).....	133		
<i>canus</i> . (<i>Peromyscus</i>).....	150		132
<i>canus</i> . (<i>Spermophilus m.</i>).....	112		93
<i>capistratus</i> . (<i>Sciurus</i>).....	62	107	
<i>Capromyinae</i>	330	389	
<i>Capromys</i>	330	389	
<i>Capromys b. thoracatus</i>	331		
<i>Capromys browni</i>	331	393	
<i>Capromys brownii</i>	331	393	
<i>Capromys elegans</i>	331	392	
<i>Capromys fourrieri</i>	330	390	
<i>Capromys ingrahami</i>	332	394	
<i>Capromys melanurus</i>	330	391	
<i>Capromys pilorides</i>	330	390	
<i>Capromys prehensilis</i>	331	391	
<i>Capromys p. gundlachi</i>	331	392	
<i>Capromys quemi</i>	330	390	
<i>Capromys thoracatus</i>	331	394	
<i>Cariacus</i>	43	68	
<i>Cariacus a. couesi</i>	45	70	40
<i>Cariacus h. californicus</i>	49	77	43
<i>Cariacus osceola</i>	44		
<i>Cariacus punctulatus</i>	48		41
<i>Cariacus virginianus</i>		70	
<i>Cariacus virgultus</i>	49		
<i>caribou</i> . (<i>Cervus t.</i>).....	37		
<i>caribou</i> . (<i>Rangifer</i>).....	37		35
<i>carlottæ</i> . (<i>Ursus</i>).....	394		
<i>carlottæ</i> . (<i>Ursus a.</i>).....	394		
<i>Carnivora</i>	365	441	293
<i>carolii</i> . (<i>Vespertilio</i>).....	478		402
<i>carolinensis</i> . (<i>Blarina b.</i>).....	461		383
<i>carolinensis</i> . (<i>Castor c.</i>).....	128		116
<i>carolinensis</i> . (<i>Evotomys</i>).....	231		174

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals etc.
carolinensis. (Rhinopoma)			417
carolinensis. (Sciurus)	75, 78, 80	108, 125	53, 62
carolinensis. (Sciurus c.)	63		
carolinensis. (Sorex)	461		
carolinensis. (Vespertilio)	483	587	410
carolinensis. (Vesperugo)	481		409
Carollia	515	667	
Carollia azteca	515	669	
Carollia castanea	516	670	
Carollia verrucata	515	669	
carpolegus. (Artibeus)	522, 523	694	
carrikeri. (Felis)	366	448	
cascadensis. (Sciurus d.)	83		66
cascadensis. (Vulpes)	382		433
castanea. (Carollia)	516	670	
castaneiceps. (Arctopithecus)	9	22	
castaneiceps. (Bradypus)	9	22	
castaneum. (Hemiderma)	516	670	
castaneus. (Peromyscus t.)	147	177	
castanonotus. (Sciurus)			57
castanops. (Cratogeomys)	266	315	220
castanops. (Pseudostoma)	266	315	
castanurus. (Citellus)	106		84
castanurus. (Spermophilus)	106		84
castanurus. (Tamias)	106		85
Castor	128	159	115
Castor canadensis	128		116
Castor c. carolinensis	128		116
Castor c. frondator	128	159	116
Castor c. pacificus	129	159	117
Castor fiber	128		115
Castor zibethicus	252		212
Castoridae	128	159	114
catalinae. (Peromyscus)	149		
catalinae. (Rhithrodontomys)	195		
catalinae. (Urocyon)	389		
Cateorus	482	586	410
Catodon	21		
Catodon macrocephalus	21		
catus. (Felis)	365	443	
caucæ. (Didelphis c.)		18	
caucæ. (Didelphis k.)	8	18	
caucæ. (Didelphis m.)		14, 18	
caudatus. (Oryzomys c.)	186	238	
caudatus. (Sorex)	456	552	
caudifer. (Nicon)	516	672	
caudifera. (Glossophaga)	519, 520		
caudivolvula. (Viverra)	404	499	
caurina. (Mustela)	422		335
caurinus. (Eutamias)	90		

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
caurinus. (Evotomys)	229		171
caurinus. (Myotis c.)	477		404
caurinus. (Tamias)	90		74
cautus. (Microtus)	244		191
cavator. (Macrogeomys)	270	329	
cavernarum. (Brachyphylla)	521	689	
cavirostris. (Ziphius)	22	47	17
cayopollin. (Didelphys)	2		
Cebidæ.	532	725	
Cebinae.	534	732	
Cebus.	536	735	
Cebus hypoleucus.	536	736	
Cebus imitator.	536	737	
cecilii. (Peromyscus)	161	178	
cedrosensis. (Peromyscus)	155	185	
celatus. (Phenacomys)	227, 228		169
centralis. (Cabassous)	12	32	
centralis. (Diphylla)	531	721	
centralis. (Echinomys)	329	386	
centralis. (Felis)	365		
centralis. (Felis o.)	365	446	
centralis. (Proechinomys)	329	386	
centralis. (Tatoua)	12	32	
Centronycteris.	492	610	
Centurio.	529	716	
Centurio flavogularis.	530	717	
Centurio mcMurtrii.	530	717	
Centurio mexicanus.	530	717	
Centurio minor.	530	717	
Centurio senex.	529	717	
Centurioninæ.	529	716	
Ceratodon.	24		18
Cercolabes.	334	400	
Cercolabes pallidus.	334	401	
Cercoleptes.	404	499	
Cercopithecidæ.	536		
Cercopithecinae.	536		
Cercopithecus	536		
Cercopithecus callitrichus.	536		
cerrosensis. (Lepus)	352	428	2
cerrosensis. (Odocoileus)	47	76	
cerrosensis. (Odontocælus)	47	76	
Cervaria	365, 370	443	
cervicalis. (Sciurus)	71		
cervicalis. (Sciurus p.)	71	119	
cervicalis. (Sciurus w.)	71	119	
Cervidæ	37	68	33
cervina. (Ovis)	53	86	46
Cervinæ.	37	68	34
cervinus. (Thomomys)	287	340	230

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
Cervus.....	41		34
Cervus acapulcensis.....	46	74	
Cervus alces.....	40, 41		
Cervus americanus.....		68	
Cervus auritus.....	48	76	42
Cervus canadensis.....	41, 42		34
Cervus c. occidentalis.....	42		34
Cervus d. americanus.....	43		
Cervus elaphus.....	41		
Cervus e. canadensis.....	41		
Cervus groenlandicus.....	39		37
Cervus hastalis.....	37		35
Cervus hemionus.....	48	76	42
Cervus leucurus.....	44		39
Cervus lewisi.....	48		41
Cervus lobatus.....	41		38
Cervus macrotis.....	48	76	42
Cervus m. californicus.....	49	77	
Cervus m. columbiana.....	48		41
Cervus macrourus.....	44		39
Cervus merriami.....	42		
Cervus mexicanus.....	45	70, 72	40
Cervus nannodes.....	42		
Cervus nemoralis.....	46	74	
Cervus occidentalis.....	42		
Cervus richardsoni.....	48		41
Cervus roosevelti.....	42		34
Cervus rufinus.....		79	
Cervus sartorii.....	50	80	
Cervus stronglyloceros.....	41		34
Cervus tarandus.....	37		35
Cervus t. arctica.....	40		
Cervus t. caribou.....	37		
Cervus t. sylvestris.....	37		35
Cervus toltecus.....	46		
Cervus virginianus.....	43		39
Cervus wapiti.....	41		34
Cervus yucatanensis.....	46	74	
Cetacea.....	15	38	7
Cetodiodon.....	21		
Cetoptera.....	18	42	
Cetus albicans.....	25		19
Chænocetus.....	21		
Chænodelphinus.....	21		
Chæronycteris peruana.....	520	682	
Chætodipus.....	300, 309	352, 356	241, 251
chapmani. (Dipodomys).....	297	350	239
chapmani. (Lepus f.).....	343, 344		282
chapmani. (Oryzomys).....	186	237, 238	
chapmani. (Perodipus).....	297	350	237

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
chapmani. (Synaptomys)	257		
chati. (Felis)	367	449	
cherriei. (Geomys)	271		
cherriei. (Oryzomys)	190	252, 253	
cherrii. (Hesperomys)	160	183	
cherrii. (Macrogeomys)	271	330	
cherrii. (Peromyscus)	160	183, 267	
cherrii. (Zygodontomys)	190	253	
chiapensis. (Lepus f.)	344	†744	
chiapensis. (Sciurus g.)	74	f. f.	
childi. (Oryzomys)		244	
chilense. (Stenoderma)	529	715	
Chilonatalis	502	635	
Chilonatalis brevimanus	502, 503	636	
Chilonatalis micropus	502	635	
Chilonatalis tumidifrons	503	637	
Chilonatalus	502	635	
Chilonatalus tumidifrons	503	637	
Chilonycteris	503	639	
Chilonycteris boothi.	505	642	
Chilonycteris d. fulvus	506		
Chilonycteris fuliginosa	504	641	
Chilonycteris grisea	504		
Chilonycteris macleayi	503, 504	641	
Chilonycteris m. fuliginosa	504	641, †748	
Chilonycteris m. grisea.	504	†747	
Chilonycteris m. inflata.	504	†747	
Chilonycteris mexicana	506	644	
Chilonycteris osburnii	505	642	
Chilonycteris parnelli.	505	642	
Chilonycteris personata	504	641	
Chilonycteris portoricensis	505	643	
Chilonycteris psilotis	505	642	
Chilonycteris quadridens.	504	641	
Chilonycteris rubiginosa	505	643	
Chilotus	234, 250	299	178, 199
Chincha.	405, 406	507	
Chincha o. major	409		
Chincha o. notata.	408		
Chincha platyrhina	409		
chiriquensis. (Myotis)	475	576	
chiriquensis. (Potos f.)	404	501	
chiriquensis. (Sciurus æ.)	62	104	
chiriquensis. (Sigmodon b.)	178	225	
chiriquensis. (Sigmodon h.)	178	225	
chiriquensis. (Tamandua t.)	11		
chiriquinus. (Proechimys c.)	329	387	
chiriquinus. (Proechinomys c.)	329	387	
Chiroderma	528	710	

* Appendix.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
Chiroderma salvini	528	710	
Chiroderma villosum	528	710	
Chironectes	1	2	
Chironectes minimus	1	3	
Chironectes palmata	1	3	
Chironectes variegatus	1	3	
Chironectes yapock	1	3	
Chiroptera	473	569	396
chlorus. (Citellus)	103		
Chæronycteris	517	673	
Chæronycteris godmani	517	674	
Chæronycteris mexicana	517	673	
Chæronycteris minor	517	674	
Choloepus	9	19	
Choloepus hoffmanni	9	20	
Chrotopterus	511	656	
Chrotopterus auritus	512	656	
chrotorrhinus. (Arvicola)	246		
chrotorrhinus. (Microtus)	246		193
chrysocomus. (Phyllostoma)	529	715	
chrysodeirus. (Citellus)	106		84
chrysodeirus. (Spermophilus)	106		84
chrysodeirus. (Tamias)	106		84
chrysomelas. (Oryzomys)	190	253	
chrysomelas. (Zygodontomys)	190	253	
chrysopsis. (Reithrodontomys)	199	265	
chrysopsis. (Rhithrodontomys)	199	265, 266	
Chrysothrix ærstedii	534	731	
Chrysothrix sciurea	534	731	
chrysotis. (Reithrodontomys)	195		152
chrysotis. (Rhithrodontomys)	195		
chrysurus. (Myoxus)	328	383	
cicognani. (Putorius)	426		344
cicognanii. (Mustela)	426		344
ciliolabrum. (Myotis c.)	477		404
ciliolabrum. (Vespertilio)	477		
cinerascens. (Citellus)	106		83
cinerascens. (Lepus)	352	429	287
cinerascens. (Spermophilus)	106		83
cinerascens. (Tamias)	106		83
cinerea. (Didelphis)	3	7	
cinerea. (Didelphys)	3		
cinerea. (Edostoma)	530	718	
cinerea. (Geomys)	262		217
cinerea. (Marmosa)	3	7	
cinerea. (Mus)	220		
cinerea. (Neotoma)	220		163
cinerea. (Teonoma)	220, 221		
cinereicollis. (Tamias)	90		75
cinereoargenteus. (Canis)	384, 386		

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
cinereo-argenteus. (Urocyon)	386		307
cinereum. (Dermanura)	525	699	
cinereus. (Didelphys)	4	9	
cinereus. (Lasiurus)	487	595	413
cinereus. (Rhithrodontomys s.).....	196	262	
cinereus. (Sciurus)	65, 66, 79		51, 52, 53
cinereus. (Sorex)	463		384
cinereus. (Vespertilio)	487	595	413
cineritius. (Peromyscus)	164	195	
cinnamomea. (Arvicola).....	248		196
cinnamomea. (Mormops b.)	507	649	
cinnamomea. (Neotoma)	223		166
cinnamomea. (Ochotona)	338		
cinnamomea. (Teonoma)	223		
cinnamomeum. (Lobostoma).....	507	649	
cinnamomeus. (Citellus l.)	97		
cinnamomeus. (Pipistrellus)	482	585	
cinnamomeus. (Spermophilus l.)	97		86
cinnamomeus. (Tamias l.)	97		86
cirrhosus. (Trachyops)	514	665	
cirrhosus. (Vampyrus)	514	665	
cisarctica. (Balæna)	15		8
Citellus.....	96	138, 139	
Citellus adocetus	116	152	
Citellus annulatus	116	151, 152	
Citellus a. goldmani	116	151	
Citellus armatus	112		
Citellus barrowensis	110		
Citellus beldingi	113		
Citellus beringensis	110		
Citellus canescens	101		
Citellus castanurus	106		
Citellus chlorus	103		
Citellus chrysodeirus	106		
Citellus c. bernardinus	107		
Citellus c. trinitatis	107		
Citellus cinerascens	106		
Citellus columbianus	111		
Citellus c. albertæ	111		
Citellus cryptospilotus	101	144	
Citellus elegans	113	145	
Citellus eremonomus	101		
Citellus erythroglutæus	108		
Citellus erythrogluteius	108		
Citellus franklini	105		
Citellus g. rupestris	114		
Citellus harrisi	96	141, 142	
Citellus h. saxicola	97	142	
Citellus interpres	98	143	
Citellus lateralis	105	147	

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Citellus l. saturatus</i>	106		
<i>Citellus leucurus</i>	97, 98	142, 143	
<i>Citellus l. cinnamomeus</i>	97		
<i>Citellus l. peninsulæ</i>	97	143	
<i>Citellus madrensis</i>	106	147	
<i>Citellus mexicanus</i>	103	146	
<i>Citellus m. parvidens</i>	104	146	
<i>Citellus mollis</i>	112		
<i>Citellus m. canus</i>	112		
<i>Citellus m. stephensi</i>	112		
<i>Citellus m. yakimensis</i>	112		
<i>Citellus nebulicola</i>	109		
<i>Citellus neglectus</i>	103		
<i>Citellus nelsoni</i>	98		
<i>Citellus nesioticus</i>	115		
<i>Citellus obsoletus</i>	99		
<i>Citellus oregonus</i>	113		
<i>Citellus osgoodi</i>	110		
<i>Citellus parryi</i>	107		
<i>Citellus p. kadiacensis</i>	108, 109		
<i>Citellus perotensis</i>	102	145	
<i>Citellus plesius</i>	108, 109		
<i>Citellus p. ablusus</i>	109		
<i>Citellus richardsoni</i>	111		
<i>Citellus spilosoma</i>	99, 100	145	
<i>Citellus s. annectens</i>	100		
<i>Citellus s. arens</i>	101		
<i>Citellus s. macrospilotus</i>	99	145	
<i>Citellus s. major</i>	99		
<i>Citellus s. marginatus</i>	100		
<i>Citellus s. obsidianus</i>	100		
<i>Citellus s. pratensis</i>	100		
<i>Citellus stonei</i>	110		
<i>Citellus stonii</i>	110		
<i>Citellus tereticaudus</i>	102	144	
<i>Citellus t. mohavensis</i>	101, 102		
<i>Citellus t. sonoriensis</i>	102	144	
<i>Citellus townsendi</i>	111		
<i>Citellus tredecimlineatus</i>	104		
<i>Citellus t. alleni</i>	105		
<i>Citellus t. badius</i>	105		
<i>Citellus t. olivaceus</i>	104		
<i>Citellus t. pallidus</i>	104		
<i>Citellus t. parvus</i>	104		
<i>Citellus t. texensis</i>	105		
<i>Citellus variegatus</i>	113, 115	148	
<i>Citellus v. atricapillus</i>	114	150	
<i>Citellus v. beecheyi</i>	114, 115	150	
<i>Citellus v. buckleyi</i>	115		
<i>Citellus v. douglasi</i>	115		

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Citellus v. fisheri</i>	114	150	
<i>Citellus v. grammurus</i>	113, 114	149, 150	
<i>Citellus v. rupestris</i>	114	150	
<i>Citellus v. utah</i>	115		
<i>Citellus vinnulus</i>	98		
<i>Citellus wortmani</i>	107		
<i>citellus. (Mus)</i>	96	139	81
<i>Citillus</i>	96	139	81
<i>Citillus buccatus</i>		149	
<i>Citillus mexicanus</i>	103	146	
<i>clarkii. (Geomys)</i>	266	315	220
<i>clavatus. (Odocoileus)</i>	46	73	
<i>clementæ. (Urocyon)</i>	390		
<i>clementis. (Peromyscus t.)</i>	145	187	130
<i>clepticus. (Canis)</i>	379	467	
<i>clinedaphus. (Monophyllus)</i>	518	678	
<i>clusius. (Thomomys)</i>	284		229
<i>Clymene</i>	32	57	30
<i>Clymene marginatus</i>	31	55	28
<i>Clymene normalis</i>	34		32
<i>Clymene similis</i>	30		27
<i>clymene. (Delphinus)</i>	34		32
<i>Clymenia</i>	32	57	30
<i>Clymenia euphrosinoides</i>	34	59	30
<i>Clymenia novæ-zelandiæ</i>	34	59	30
<i>cnecus. (Perognathus)</i>	309	362	
<i>cnemophila. (Neotoma)</i>	207		
<i>cnemophila. (Neotoma f.)</i>	207		
<i>Cnephæus</i>	482	585	410
<i>Cnephaiophilus</i>	480		407
<i>cocos. (Sciurus s.)</i>	74	124	
<i>Coelogenus</i>	337	408	
<i>Coendou</i>	334		
<i>Coendou lænatus</i>	335	402	
<i>Coendou m. yucatanicæ</i>	335	402	
<i>Coendou rothschildi</i>	334		
<i>Coendu</i>	334	399	
<i>Coendu lænatum</i>	335	402	
<i>Coendu mexicanum</i>	335	402	
<i>Coendu m. yucatanicæ</i>	335	402	
<i>Coendu pallidum</i>	334	401	
<i>Coendu rothschildi</i>	334	401	
<i>Cogia</i>	22	46	
<i>Cogia breviceps</i>	22	46	
<i>cognatus. (Hesperomys)</i>			127
<i>coibæ. (Dasyprocta)</i>	336	407	
<i>coibensis. (Alouatta p.)</i>	533	727	
<i>coibensis. (Molossus)</i>	496		
<i>colimæ. (Reithrodontomys)</i>	200	266	
<i>colimæ. (Rhithrodontomys)</i>	200	266, 267	

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
colimæ. (Sigmodon)	176	227	
colimensis. (Sciurus a.)	71, 72	120	
colimensis. (Sciurus p.)	71	119	
colimensis. (Sciurus w.)	72	120	
collaris. (Lagomys)	338		268
collaris. (Ochotona)	338		268
colliæi. (Sciurus)	73, 77	121, 129	60
Colobotis	96, 107	139	82, 89
colonus. (Geomys)	262		217
coloratus. (Oryzomys a.)	181		148
coloratus. (Oryzomys p.)	181		
columbiana. (Cervus m.)	48		41
columbiana. (Neotoma c.)	221		165
columbiana. (Teonoma c.)	221		
columbianus. (Arctomys)	111		91
columbianus. (Citellus)	111		
columbianus. (Odocoileus)	48		41
columbianus. (Odontocelus)	48		
columbianus. (Oreamnos m.)	52		
columbianus. (Oreamnos m.)	52		
columbianus. (Perodipus o.)	297		238
columbianus. (Perognathus)	308		256
columbianus. (Perognathus l.)	308		250
columbianus. (Spermophilus)	111		91
columbiensis. (Lepus a.)	361		275
Comastes	473	571	400
communis. (Phocæna)	25	48	20
communis. (Pterobalæna)	19		13
compacta. (Blarina b.)	462		
compactus. (Dipodomys)	299		240
compactus. (Perodipus)	299		240
compressicauda. (Phocæna)	31	56	29
compressus. (Macrotus w.)	509		
compressus. (Otopterus w.)	509		
comptus. (Peromyscus)	168	206	
concinus. (Vespertilio)	475	575	
concolor. (Felis)	369		293
concolor. (Mephitis)	410	512	
concolor. (Sciurus a.)	66		57
concreta. (Beluga)	25		20
conditi. (Perognathus)	310		252
Condylura	467		394
Condylura cristata	467		395
Condylura macroura	467		395
Condylura prasinata	467		395
Conepatus	410, 411	512	†509
Conepatus felipensis	412	517	
Conepatus humboldtii	410	512	
Conepatus leuconotus	411	515	

†Appendix.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Conepatus l. texensis</i>	411	515	
<i>Conepatus mapurito</i>	412	516	510
<i>Conepatus mesoleucus</i>	411	515	
<i>Conepatus m. mearnsi</i>	411	516	
<i>Conepatus pediculus</i>	412	517	
<i>Conepatus sonoriensis</i>	411	514	
<i>Conepatus tropicalis</i>	412	517	
<i>confinis.</i> (<i>Lepus a.</i>)	350	424	
<i>connectens.</i> (<i>Lepus f.</i>)	343		
<i>consobrinus.</i> (<i>Peromyscus m.</i>)	166	202	
<i>consobrinus.</i> (<i>Tamias m.</i>)	93		78
<i>constabiei.</i> (<i>Phenacomys</i>)	228		169
<i>constablii.</i> (<i>Phenacomys</i>)	228		
<i>constrictus.</i> (<i>Microtus c.</i>)	241		186
<i>convexum.</i> (<i>Uroderma</i>)	524	697	
<i>cookii.</i> (<i>Rosmarus</i>)	439		358
<i>coolidgei.</i> (<i>Peromyscus l.</i>)	147		
<i>coolidgei.</i> (<i>Peromyscus t.</i>)	147	187	
<i>cooperi.</i> (<i>Sorex</i>)	446		366
<i>cooperi.</i> (<i>Synaptomys</i>)	255		203
<i>cooperi.</i> (<i>Tamias</i>)	86		69
<i>copei.</i> (<i>Balænoptera v.</i>)	19		13
<i>copei.</i> (<i>Perognathus</i>)	301		243
<i>Corsira tropicalis.</i>		559	
<i>coryi.</i> (<i>Artibeus</i>)	522	693	
<i>coryi.</i> (<i>Felis</i>)	369		293
<i>Corynorhinus</i>	490	603	398
<i>Corynorhinus macrotis</i>	490		399
<i>Corynorhinus m. pallescens</i>	490	604	399
<i>Corynorhinus m. townsendi</i>	491	604	400
<i>costaricensis.</i> (<i>Blarina</i>)	461		382
<i>costaricensis.</i> (<i>Felis</i>)	366	448	
<i>costaricensis.</i> (<i>Felis b.</i>)	370	456	
<i>costaricensis.</i> (<i>Macrogeomys</i>)	271	330	
<i>costaricensis.</i> (<i>Odocoileus</i>)	46	73	
<i>costaricensis.</i> (<i>Odontocelus</i>)	46	73	
<i>costaricensis.</i> (<i>Oryzomys</i>)	189	249, 250	
<i>costaricensis.</i> (<i>Reithrodontomys</i>)	200	267	
<i>costaricensis.</i> (<i>Rhithrodontomys</i>)	200	266, 267	
<i>couchii.</i> (<i>Spermophilus</i>)	113	149	188
<i>couesi.</i> (<i>Cariacus a.</i>)	45	70	40
<i>couesi.</i> (<i>Erethizon c.</i>)	332, 333		265
<i>couesi.</i> (<i>Hesperomys</i>)	186	236	
<i>couesi.</i> (<i>Odocoileus a.</i>)	45	70	40
<i>couesi.</i> (<i>Odontocelus a.</i>)	45	70, 71	
<i>couesi.</i> (<i>Oryzomys</i>)	186	236, 239	
<i>cougar.</i> (<i>Felis</i>)	369		
<i>cozumelæ.</i> (<i>Didelphis y.</i>)	5	15	
<i>cozumelæ.</i> (<i>Didelphys y.</i>)	5	15	
<i>cozumelæ.</i> (<i>Oryzomys</i>)	182	241	

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
cozumelæ. (<i>Peromyscus</i>)	162	181	
crassidens. (<i>Phocæna</i>)	27	51	23
crassidens. (<i>Pseudorca</i>)	27	51	23
crassum. (<i>Tagassu a.</i>)	36	64	
crassus. (<i>Phenacomys c.</i>)	227, 228		†430
crassus. (<i>Tayassu a.</i>)	36	64	
<i>Cratogeomys</i>	264	311	220
<i>Cratogeomys castanops</i>	266	315	220
<i>Cratogeomys c. goldmani</i>	266	316	
<i>Cratogeomys estor</i>	265	314	
<i>Cratogeomys fulvescens</i>	266	316	
<i>Cratogeomys merriami</i>	265	313	
<i>Cratogeomys oreocetes</i>	265	314	
<i>Cratogeomys peregrinus</i>	265	314	
<i>Cratogeomys perotensis</i>	265	313	
crawfordi. (<i>Notiosorex</i>)	459	554	385
crawfordi. (<i>Sorex</i>)	459	554	385
cremnobates. (<i>Ovis c.</i>)	54	84	
creper. (<i>Reithrodontomys</i>)	204	272	
creper. (<i>Rhithrodontomys</i>)	204	272	
crepuscularis. (<i>Nycticea</i>)	488	598	415
<i>Cricetinae</i>	130		119
<i>Cricetodipus</i>	300	352	241
<i>Cricetodipus flavus</i>	302		243
<i>Cricetodipus parvus</i>	306		247
<i>Cricetus</i>	119		
<i>Cricetus myoides</i>	140		125
<i>Cricetus talpoides</i>	278		227
crinitus. (<i>Hesperomys</i>)	159		140
crinitus. (<i>Oryzomys</i>)	183	245	
crinitus. (<i>Peromyscus t.</i>)	159		140
crispus. (<i>Heteromys</i>)	320	380	
crispus. (<i>Liomys</i>)	320	380	
crispus. (<i>Myrmecophaga</i>)	10	27	
cristata. (<i>Condylura</i>)	467		395
cristata. (<i>Cystophora</i>)	440		360
cristata. (<i>Dasyprocta</i>)	337	407	
cristata. (<i>Phoca</i>)	440		360
cristatus. (<i>Echinomys</i>)	328	383	
cristatus. (<i>Sorex</i>)	467		395
cristobalensis. (<i>Peromyscus z.</i>)	170	209	
crooki. (<i>Dorcelaphus</i>)	47		41
crooki. (<i>Odocoileus</i>)	47		41
crooki. (<i>Odontocælus</i>)	47		
crotaphiscus. (<i>Delphinus</i>)	33		31
crotaphiscus. (<i>Prodelphinus</i>)	33		31
crusnigrum. (<i>Tagassu</i>)	36	65	
crusnigrum. (<i>Tayassu</i>)	36	65	
cryptospilotus. (<i>Citellus</i>)	101	144	

†Appendix.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
cryptospilotus. (Spermophilus)	101		97
Cryptotis	460, 463	556, 557	382, 384
cubanus. (Monophyllus)	519	678	
cubanus. (Nycticeius h.)	488	599	
cubanus. (Solenodon)	472	567	
cubanus. (Vesperus)	488	599	
cubensis. (Scotophilus)	484	588	
cubensis. (Vespertilio f.)	484	588	
cucullata. (Phoca)	440		360
Cuica	2	5	
cumberlandius. (Geomys)	262		217
cumulator. (Neotoma)	205	280	154
Cuniculus	260		209
cupes. (Ochotona)	339		268
cupreata. (Talpa)	470		388
curtata. (Arvicola)	247		195
curtatus. (Microtus)	247		195
curvirostris. (Delphinus d.)	31	55	28
Cutia	335	403	
cuvieri. (Grampus)	28		25
Cuvierius	18	42	11
Cyclopes	10	24	
Cyclopes dorsalis	10	25	
Cyclothurus	10	25	
Cyclothurus dorsalis	10	25	
cymodoce. (Delphinus)	31	56	29
cynocephala. (Nycticea)	500		417
cynocephalus. (Nyctinomus)	500		417
Cynomyanax	425		
Cynomyonax	423	530	338, 341
Cynomys	117		102
Cynomys arizonensis	118		103
Cynomys grisea	117		102
Cynomys gunnisoni	118		103
Cynomys leucurus	118		104
Cynomys lewisi	118		104
Cynomys ludovicianus	117		102
Cynomys mexicanus	119	156	
Cynomys parvidens	118		
Cynomys pyrrotrichus	117		
Cynomys socialis	117		102
Cystophora	440		360
Cystophora borealis	440		360
Cystophora cristata	440		360
Cystophorinæ	439		359
dacota. (Arctomys)	120		106
dakota. (Marmota)	120		
dakotensis. (Sciurus h.)	81		62
dale. (Dauphindo)	23		18

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
dalli. (<i>Lepus a.</i>)	361		275
dalli. (<i>Ovis</i>)	55		47
dalli. (<i>Ovis m.</i>)	55		
dalli. (<i>Phocæna</i>)	25		21
dalli. (<i>Synaptomys</i>)	258		206
dalli. (<i>Ursus</i>)	391		311
Dama		68	
Dama lichtensteini.	45	70	
Dama rothschildi	46	72	
Danis	390, 392	479	309
Dasynotus	316	368	
Dasypodidæ	12	31	4
Dasypodinae	12	31	
Dasyprocta	335	403	
Dasyprocta antillensis	337	407	
Dasyprocta callida	336	406	
Dasyprocta coibæ	336	407	
Dasyprocta cristata	337	407	
Dasyprocta isthmica	336	407	
Dasyprocta mexicana	336	405	
Dasyprocta punctata	336	405	
Dasyprocta ruatanica	336	405	
Dasypterus	487	595	414
Dasypterus e. panamensis	487	597	
Dasypterus e. xanthinus	487	597	
Dasypterus floridanus	488		
Dasypterus intermedius	487, 488	595	414
Dasypus longicaudus	13	33	4
Dasypus m. fenestratus	13	33	4
Dasypus novemcinctus	13	32, 33	4
Dasypus octo-cinctus	13	33	4
Dasypus peba	13	33	4
Dasypus unicinctus	12	31	
Dauphindo dale	23		18
davidsoni. (<i>Balænoptera</i>)	19	42	12
davyi. (<i>Dermonotus</i>)	506	644	
davyi. (<i>Pteronotus</i>)	506	644	
dawsoni. (<i>Evotomys</i>)	229		172
dawsoni. (<i>Rangifer</i>)	40		36
Declidurus	494		
declivis. (<i>Beluga</i>)	25		20
decolorus. (<i>Nyctomys</i>)	172	216	
decolorus. (<i>Sitomys</i>)	172	216	
decumanus. (<i>Mus</i>)	129	164	117
decurtata. (<i>Arvicola</i>)	247		
decussatus. (<i>Canis</i>)	380		304
deductor. (<i>Delphinus</i>)	27	52	23
degener. (<i>Lutra</i>)	435		353
dekayi. (<i>Arvicola</i>)	235		180
dekayi. (<i>Sorex</i>)	461		382

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
deletrix. (Vulpes)	381		305
Delphinapterinæ	24	48	
Delphinapterus	24, 32		19, 29
Delphinapterus beluga	24, 25		19
Delphinapterus borealis	32		30
Delphinapterus leucas	25		19
Delphinidæ	24	47, 48	18
Delphinus	30	54	27
Delphinus acutus	29		25
Delphinus albimanus		55	28
Delphinus albirostris	29		26
Delphinus algeriensis		55	28
Delphinus alope	33	58	
Delphinus anarnacus	21		16
Delphinus bairdi	30	55	28
Delphinus balteatus	31	55	28
Delphinus bidentatus	21		16
Delphinus butskopf	21		16
Delphinus canadensis	25		19
Delphinus clymene	34		32
Delphinus crotaphiscus	33		31
Delphinus cymodoce	31	56	29
Delphinus deductor	25	52	23
Delphinus delphis	30	54	28
Delphinus d. curvirostris	31	55	28
Delphinus doris	34		32
Delphinus dubius	32		30
Delphinus epidon	23		18
Delphinus eschrichti	29		25
Delphinus euphrosyne	33	58	30
Delphinus eurynome	31, 34	56	29
Delphinus forsteri	30	55	28
Delphinus frenatus	34		32
Delphinus frontalis	34		32
Delphinus fulvi-fasciatus	30	54	28
Delphinus fulvo-fasciatus	30	54	28
Delphinus fuscus	31	55	28
Delphinus gladiator	26	51	22
Delphinus globiceps	27	52	23
Delphinus griseus	28		25
Delphinus hunteri	21		16
Delphinus hyperoodon	21		16
Delphinus ibsenii	29		26
Delphinus intermedius	27, 28	52	23, 25
Delphinus janira	30	55	28
Delphinus leucas	25		19
Delphinus leucopleurus	29		25
Delphinus longidens	30		27
Delphinus longirostris	33	58	
Delphinus major	30	55	28

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Delphinus marginatus</i>	34	58	30
<i>Delphinus melas</i>	27	52	
<i>Delphinus metis</i>	31	56	29
<i>Delphinus microps</i>	31, 33	55, 58	28
<i>Delphinus micropterus</i>	23		18
<i>Delphinus moorii</i>	30	55	28
<i>Delphinus moschatus</i>	31	55	28
<i>Delphinus novæ-zelandiæ</i>	30	55	28
<i>Delphinus orca</i>	26	50, 51	
<i>Delphinus peronii</i>	32		29
<i>Delphinus phocæna</i>	25	48	20
<i>Delphinus plagiodon</i>	33	58	31
<i>Delphinus pomeegra</i>	30	55	28
<i>Delphinus rissoanus</i>	28		25
<i>Delphinus sowerbianus</i>	31	55	28
<i>Delphinus sowerbiensis</i>	23		18
<i>Delphinus sowerbyi</i>	23		18
<i>Delphinus stenorhynchus</i>	33	58	
<i>Delphinus styx</i>	33	58	30
<i>Delphinus tethyos</i>	34	58	30
<i>Delphinus truncatus</i>	31	56	29
<i>Delphinus tursio</i>	31	55	29
<i>Delphinus variegatus</i>	31	55	28
<i>Delphinus walkeri</i>	30	55	28
<i>delphis</i> . (<i>Delphinus</i>).....	30	54	28
<i>deppei</i> . (<i>Sciurus</i>).....	60, 61	101, 102	
<i>deppii</i> . (<i>Sciurus</i>).....	60	101	
<i>depressus</i> . (<i>Nyctinomops</i>).....	499	627	
<i>depressus</i> . (<i>Nyctinomus</i>).....	499	627	
<i>derbianus</i> . (<i>Caluromys</i>).....	4	9	
<i>derbianus</i> . (<i>Didelphys</i>).....	4	9	
<i>Dermanura</i>	525	699	
<i>Dermanura cinereum</i>	525	699	
<i>Dermanura eva</i>	524	697	
<i>Dermanura phæotis</i>	525	699	
<i>Dermonotus</i>	506	644	
<i>Dermonotus davyi</i>	506	644	
<i>Dermonotus d. fulvus</i>	506	646	
<i>deserti</i> . (<i>Dipodomys</i>).....	292	345	235
<i>deserti</i> . (<i>Reithrodontomys m.</i>).....	193	259	151
<i>deserticola</i> . (<i>Lepus t.</i>).....	356, 357	437	291
<i>deserticola</i> . (<i>Peromyscus l.</i>).....	140	188	
<i>deserticola</i> . (<i>Peromyscus t.</i>).....	140	188	
<i>deserticolus</i> . (<i>Hesperomys l.</i>).....	140	188	
<i>deserticolus</i> . (<i>Peromyscus a.</i>).....	140	188	125
<i>desertorum</i> . (<i>Neotoma</i>).....	215		161
<i>desertorum</i> . (<i>Thomomys</i>).....	281		
<i>desmaresti</i> . (<i>Moschophoromys</i>).....	190	250, 251	
<i>desmaresti</i> . (<i>Mus</i>).....	189, 190	250	
<i>desmaresti</i> . (<i>Phoca</i>).....	444		364

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
desmarestianus. (Heteromys)	318	372	
Desmodon.....	530		
Desmodon rotundum.....	530		
Desmodontinae.....	530	718	
Desmodus.....	530	718	
Desmodus d'orbignyi.....	530	718	
Desmodus fuscus.....	530	719	
Desmodus murinus.....	530	719	
Desmodus rotundus.....	530	718	
Desmodus rufus.....	530	718	
destructor. (Lutra)			352
destructor. (Orca)	27	51	23
devius. (Oryzomys)	188	244	
diazi. (Lepus)		412	
dickinsoni. (Reithrodontomys h.)	192		
dickinsoni. (Reithrodontomys l.)	192		150
dickinsoni. (Rhithrodontomys l.)	192		
Diclidurus.....	494	614	
Diclidurus albus.....	494	615	
Diclidurus freyreissii.....	494	615	
Diclidurus virgo.....	494	615	
Dicotyles.....	36	65	33
Dicotyles.....	35	62	33
Dicotyles angulatus.....	35	63	
Dicotyles a. sonoriensis.....	36	64	
Dicotyles labiatus.....	37	66	
Dicotyles tajacu.....	35		33
Dicotylidæ.....			33
Dicranocerus.....	51	81	43
Dicrostonyx.....	260		209
Dicrostonyx hudsonius.....	260		209
Dicrostonyx h. alascensis.....	260, 261		210
Dicrostonyx h. nelsoni.....	261		210
Dicrostonyx h. richardsoni.....	261		211
Dicrostonyx h. unalascensis.....	260		210
Dicrostonyx nelsoni.....	260		210
Dicrostonyx richardsoni.....	261		211
Dicrostonyx unalascensis.....	260		210
Didactyla.....	10	25	
didactyla. (Myrmecophaga)	10	24	
Didactyles.....	10	25	
didactylus. (Bradypus)	9	19	
Didelphis.....	5	14	
Didelphis californica.....	6	15	
Didelphis canescens.....	2	6	
Didelphis c. caucæ.....		18	
Didelphis cinerea.....	3	7	
Didelphis dorsigera.....	2		
Didelphis k. caucæ.....	8		
Didelphis marsupialis.....	5	14	

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Didelphis m. battyi</i>	7	17	
<i>Didelphis m. caucæ</i>		14, 18	
<i>Didelphis m. etensis</i>	8	18	
<i>Didelphis m. insularis</i>	7	17	
<i>Didelphis m. tabascensis</i>	6		
<i>Didelphis m. texensis</i>	6	16	
<i>Didelphis memina</i>	1	3	
<i>Didelphis mesamericana</i>	6	15	
<i>Didelphis m. tabascensis</i>	6	16	
<i>Didelphis m. texensis</i>	6	16	
<i>Didelphis murina</i>	2	4, 5	
<i>Didelphis myosurus</i>	4		
<i>Didelphis philander</i>	4	9	
<i>Didelphis richmondi</i>	8	16	
<i>Didelphis virginiana</i>	7		
<i>Didelphis v. pigra</i>	7		I
<i>Didelphis yucatanensis</i>	5	14	
<i>Didelphis y. cozumelæ</i>	5	15	
<i>Didelphyidæ</i>	1	I	I
<i>Didelphys</i>	5, 7	13	I
<i>Didelphys aurita</i>	8	17	
<i>Didelphys californica</i>	6		3
<i>Didelphys cayopollin</i>	2		
<i>Didelphys cinerea</i>	3		
<i>Didelphys cinereus</i>	4	9	
<i>Didelphys derbianus</i>	4	9	
<i>Didelphys impavida</i>	2		
<i>Didelphys macrotarsus</i>	2		
<i>Didelphys marsupialis</i>	7		
<i>Didelphys m. battyi</i>	7	17	
<i>Didelphys m. etensis</i>	8	18	
<i>Didelphys m. insularis</i>	7	17	
<i>Didelphys m. texensis</i>	6		
<i>Didelphys mesamericana</i>	6	15, 16, 17	
<i>Didelphys m. tabascensis</i>	6	16	
<i>Didelphys m. texensis</i>	6	16	
<i>Didelphys microtarsus</i>	2		
<i>Didelphys murinus</i>	2		
<i>Didelphys musculus</i>	2		
<i>Didelphys nudicaudata</i>	4	11	
<i>Didelphys quica</i>	5	12	
<i>Didelphys richmondi</i>	8	16	
<i>Didelphys virginiana</i>	7		I
<i>Didelphys v. pigra</i>	7		I
<i>Didelphys yucatanensis</i>	5	14, 15	
<i>Didelphys y. cozumelæ</i>	5	15	
<i>difficilis</i> . (<i>Peromyscus</i>)	158	193	
<i>difficilis</i> . (<i>Reithrodontomys</i>)	197	263	
<i>difficilis</i> . (<i>Rhithrodontomys</i>)	197	263	
<i>difficilis</i> . (<i>Vesperimus</i>)	158	193	

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
dilatus. (Scapanus)	468		392
dilutior. (Oryzomys c.)	186	238	
dimidiatus. (Callocephalus)	444		364
Dinops.....	499	628	416
Diodon.....	24		18
Dionyx.....	10	25	
Diphylla.....	530	720	
Diphylla centralis.....	531	741	
Diphylla ecaudata	530, 531	720	
Diplostoma	261, 271	332	221
Diplostoma bulbivorum.....	272		
Diplostoma fusca.....	262		217
Dipodomysinae.....	288	341	231
Dipodomys.....	288	341	231
Dipodomys agilis	293	351	236
Dipodomys ambiguus.....	289		234
Dipodomys californicus.....	291		236
Dipodomys c. pallidulus.....	292		†431
Dipodomys chapmani	297	350	237
Dipodomys compactus.....	299		240
Dipodomys deserti.....	292	345	235
Dipodomys d. helleri.....	292		
Dipodomys elator.....	292		235
Dipodomys hermanni.....	293		236
Dipodomys merriami.....	288	345	232
Dipodomys m. ambiguus.....	289		234
Dipodomys m. arenivagus.....	290	347	
Dipodomys m. atronasus.....	290	348	
Dipodomys m. exilis.....	289		233
Dipodomys m. melanurus.....	291	348	
Dipodomys m. mortivallis.....	290		
Dipodomys m. nevadensis.....	288		232
Dipodomys m. nitratoides.....	288		233
Dipodomys m. nitratus.....	288		233
Dipodomys m. parvus.....	289		234
Dipodomys m. similis.....	289		234
Dipodomys m. simiolus.....	289, 290		234
Dipodomys mitchelli.....	291	344	
Dipodomys montanus.....	295		
Dipodomys ordii.....	296		
Dipodomys ornatus.....	291	344	
Dipodomys parvus.....	289		234
Dipodomys perotensis.....	291	344	
Dipodomys phillipsi.....	288, 291	343	231
Dipodomys similis	289		234
Dipodomys simiolus.....	289		234
Dipodomys spectabilis.....	293	345	235
Dipodomys wagneri.....	293		236
Dipodops.....	295		

†Appendix.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
Dipodops longipes.....*	298		239
Dipodops o. palmeri.....	296		
Dipodops richardsoni.....	297		238
Dipodops sennetti.....	297		239
Dipus.....	324		
Dipus americanus.....	325		259
Dipus canadensis.....	324		257
Dipus hudsonius.....	323, 324		257
Dipus labradorius.....	324		257
discifera. (Hyonycteris).....	503	637	
discifera. (Thyroptera).....	503	637	
discolor. (Callocephalus).....	444		364
discolor. (Felis).....			293
dispar. (Neotoma f.).....	206		160
dispar. (Perognathus c.).....	315		255
distincta. (Neotoma).....	207	278	
divergens. (Trichechus).....	439		358
dobsoni. (Sorex).....	448		370
dobsoni. (Sorex v.).....	448		370
dolichocephalus. (Macrogeomys).....	270	328	
dominicensis. (Myotis).....	475	576	
d'orbigny. (Desmodus).....	530	718	
Dorcelaphus.....	43	68	
Dorcelaphus crooki.....	47		41
Dorcelaphus h. eremicus.....	49	77	
Dorcelaphus texanus.....	45	70	40
doreides. (Tursio).....	34	58	30
doris. (Delphinus).....	34		32
dorsalis. (Cyclopes).....	10	25	
dorsalis. (Cyclothurus).....	10	25	
dorsalis. (Reithrodontomys).....	203	272	
dorsalis. (Rhithrodontomys).....	203	272	
dorsalis. (Sciurus).....	66, 76	127	57
dorsalis. (Sciurus a.).....	76	127	
dorsalis. (Tamias).....	85	135	68
dorsata. (Hystrix).....	333		
dorsata. (Phoca).....	444		364
dorsatum. (Erethizon).....	333		
dorsatus. (Erethizon).....	334		264
dorsatus. (Hystrix).....	332	397	264
dorsatus. (Noctilio).....	494	617	
dorsiger. (Grymæomys).....	2		
dorsigera. (Didelphis).....	2		
douglasi. (Citellus v.).....	115		
douglasi. (Lepus).....		415	
douglasi. (Sciurus).....	82		65
douglasi. (Spermophilus g.).....	115		89
douglasi. (Thomomys).....	277, 278		226
douglasii. (Arctomys).....	115		89
douglasii. (Geomys).....	277		226

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals etc.
douglasii. (Sciurus)	82		
dowi. (Tapirella)	56	88	
dowii. (Elasmognathus)	58	88	
drummondi. (Arvicola)	236		181
drummondi. (Microtus)	236, 237		181
drummondi. (Myoxus)	220, 221		
drummondi. (Neotoma c.)	221		164
drummondi. (Teonoma c.)	221		
dubius. (Delphinus)	32		30
dubius. (Hippocamelus)	50		
dubius. (Peromyscus)	162	181	
dugong. (Hydrodamalis)		36	
duguidii. (Physalus)	19		13
durangæ. (Eutamias)	88		
durangæ. (Lepus)	345	424	
durangæ. (Myotis c.)	478	579	
durangæ. (Neotoma i.)	217	285	
durangæ. (Tamias)	88	137	
durangi. (Sciurus)	67	112	
durangi. (Sciurus a.)	67		
dutcheri. (Microtus)	239		185
dutertreus. (Vespertilio)	484	588	
dychei. (Reithrodontomys)	194		
dychii. (Rhithrodontomys)	192		150
dyselius. (Peromyscus)	156		138
Dysopes.....	495, 499	619, 628	416
Dysopes abrasus.....	497	623	
Dysopes albus.....	495	619	
Dysopes alecto.....	495	619	
Dysopes aurispinosus.....	499	627	
Dysopes auritus.....	499	627	
Dysopes cæcus.....		627	
Dysopes fumarius.....	495	620	
Dysopes glaucinus.....	497	623	
Dysopes gracilis.....	498	626	
Dysopes holosericeus.....	495	619	
Dysopes laticaudatus.....	499	627	
Dysopes leucopleura.....	497	623	
Dysopes longimanus.....	497	623	
Dysopes naso.....			417
Dysopes olivaceo-fuscus.....	495	620	
Dysopes velox.....	495	620	
ecaudata. (Anoura)	520	682	
ecaudata. (Diphylla)	530, 531	720	
ecaudatus. (Rhinolophus)	530	718	
Echimys.....	329	385	
Echimys semispinosus.....	329	386	
Echimys trinitatis.....	329	385	
Echinomys centralis.....	320	386	

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
Echinomys cristatus.....	328	383	
Echinoprocta.....	332	397	264
Echinosciurus.....	59, 68	94, 113	50
Echinothrix.....	332	397	
Ectophylla.....	528	709	
Ectophylla alba.....	528	710	
edax. (Arvicola).....	241		187
edax. (Microtus).....	241		187
Edentata.....	9	19	4
Edostoma cinerea.....	530	718	
edwardsi. (Lepus).....	358		
effugius. (Sciurus a.).....	72	120	
effugius. (Sciurus p.).....	72	120	
effugius. (Sciurus w.).....	72	120	
Elaphus.....	41		
elaphus. (Cervus).....	41		
Elasmognathus.....	58	87	
Elasmognathus bairdii.....	58	87	
Elasmognathus dowii.....	58	88	
elassodon. (Sorex l.).....	450		
elassodon. (Sorex o.).....	450		
elator. (Dipodomys).....	292		235
elegans. (Capromys).....	331	392	
elegans. (Citellus).....	113	145	
elegans. (Felis).....	367	449	
elegans. (Spermophilus).....	113		94
elibatus. (Perognathus).....	304		
elongata. (Mephitis).....	406		324
elongata. (Mephitis m.).....	406		
elongata. (Phyllostoma).....	513	662	
elucus. (Procyon l.).....	400		319
Emballonura lineata.....	492	609	
Emballonurinae.....	492	608	
emmonsii. (Ursus).....	397		315
emmonsii. (Ursus a.).....	397		
empetra. (Mus).....	120		
empetra. (Spermophilus).....	108		89
energumenos. (Putorius v.).....	424, 425		
energumenus. (Putorius v.).....	424		340
engelhardtii. (Marmota).....	121		
Enhydra.....	435	537	354
enixus. (Microtus).....	237		182
entomophaga. (Saimiris).....	534	731	
Enydris gracilis.....	436	537	354
Eozapus.....	324		257
epidon. (Delphinus).....	23		18
epixanthum. (Erethizon).....	332		397
epixanthus. (Erethizon).....	332		265
Eptesicus.....	482	585	410
Eptesicus f. bernardinus.....	483		

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Eptesicus f. melanopterus</i>	483		
<i>Eptesicus f. osceola</i>	483		
<i>Eptesicus melanops</i>	483	587	410
<i>equestris</i> . (<i>Phoca</i>).....	442		362
<i>Equus bisulcus</i>	50		
<i>eremica</i> . (<i>Felis r.</i>).....	372	458	298
<i>eremicoides</i> . (<i>Peromyscus a.</i>).....	154	192	
<i>eremicus</i> . (<i>Dorcelaphus h.</i>).....	49	77	
<i>eremicus</i> . (<i>Hesperomys</i>).....	136	191	136
<i>eremicus</i> . (<i>Lepus t.</i>).....	355, 357	435	291
<i>eremicus</i> . (<i>Lynx r.</i>).....	372	458	298
<i>eremicus</i> . (<i>Odontocelus h.</i>).....	49, 50	77	
<i>eremicus</i> . (<i>Perognathus</i>).....	311		253
<i>eremicus</i> . (<i>Perognathus p.</i>).....	311	359	253
<i>eremicus</i> . (<i>Peromyscus</i>).....	136	191	136
<i>eremicus</i> . (<i>Sigmodon h.</i>).....	176	227	145
<i>eremicus</i> . (<i>Ursus a.</i>).....	394		
<i>Eremiomys</i>	234	299	178
<i>eremonomus</i> . (<i>Citellus</i>).....	101		
<i>Erethizon</i>	332	397	264
<i>Erethizon dorsatum</i>	333		
<i>Erethizon d. picinum</i>	334		
<i>Erethizon dorsatus</i>	334		264
<i>Erethizon d. picinus</i>	334		†431
<i>Erethizon epixanthum</i>	332	397	
<i>Erethizon e. couesi</i>	333		
<i>Erethizon e. myops</i>	333		
<i>Erethizon e. nigrescens</i>	333		
<i>Erethizon epixanthus</i>	332		265
<i>Erethizon e. couesi</i>	332		265
<i>Erethizon e. myops</i>	333		266
<i>Erethizon e. nigrescens</i>	333		
<i>Erethizontidæ</i>	332	397	264
<i>Erethizontinæ</i>	332	397	264
<i>Erignathus</i>	441	541	361
<i>Eriodes frontatus</i>	535	733	
<i>erminea</i> . (<i>Mustela</i>).....	430		
<i>erythrodactylus</i> . (<i>Vespertilio</i>).....	281		409
<i>erythroglutæus</i> . (<i>Citellus</i>).....	108		
<i>erythrogluteia</i> . (<i>Arctomys p.</i>).....	108		91
<i>erythrogluteius</i> . (<i>Citellus</i>).....	108		
<i>erythromos</i> . (<i>Phyllostoma</i>).....	529	715	
<i>eschrichti</i> . (<i>Delphinus</i>).....	29		25
<i>escuinapæ</i> . (<i>Felis r.</i>).....	372, 373	456	
<i>escuinapæ</i> . (<i>Lynx r.</i>).....	372		
<i>eskimo</i> . (<i>Putorius r.</i>).....	428		345
<i>estor</i> . (<i>Canis</i>).....	378	469	302
<i>estor</i> . (<i>Cratogeomys</i>).....	265	314	
<i>estor</i> . (<i>Mephitis</i>).....	407		326

†Appendix.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
estor. (Mephitis m.)	407	509	
etensis. (Didelphis m.)	8	18	
etensis. (Didelphys m.)	8	18	
Euarctos.	390	479	309
Euarctus.	394		313
Eubalæna.	15		7
Eubalæna australis.	15		8
Eucervus	43	68	
Eucritus.	334		
Eudelphinus.	30	54	27
Euderma	490		398
Euderma maculata.	490		398
Euderma maculatum.	490		
eulophus. (Ursus)	392		
Eumetopias.	437		555
Eumetopias jubata.	437		
Eumetopias stelleri.	437		355
Euotaria.	438		
euphrosinoides. (Clymenia)	34	59	30
euphrosyne. (Delphinus)	33	58	30
euphrosyne. (Prodelphinus)	34	58	30
Euprocyon.	400, 402	492	
europæus. (Orca)	26	51	22
eurynome. (Delphinus)	31, 34	56	29
Eurypterna	10	25	
Eutamias.	85	134	68
Eutamias adsitus	95		
Eutamias a. operarius	†541		
Eutamias canescens.	86		
Eutamias canicaudus.	93		
Eutamias caniceps.	93		
Eutamias caurinus.	90		
Eutamias c. canipes.	91		
Eutamias d. utahensis.	85		
Eutamias durangæ.	88		
Eutamias hopiensis	†541		
Eutamias lectus	94		
Eutamias orocetes.	95		
Eutamias palmeri.	86		
Eutamias s. inyoensis.	89		
Eutamias t. ochrogenys.	87		
eva. (Artibeus)	524	697	
eva. (Dermanura)	524	697	
eva. (Peromyscus)	165	198	
evides. (Peromyscus s.)	156	179	
evotis. (Myotis)	474	574	
evotis. (Notiosorex c.)	460	555	
evotis. (Sorex)	460	555	
evotis. (Vespertilio)	474	574	

†Appendix.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
evotis. (<i>Vespertilio a.</i>)	479		
Evotomys.....	229		170
Evotomys alascensis.....	229		177
Evotomys breviceaudus	231		174
Evotomys californicus.....	232		176
Evotomys carolinensis.....	231		174
Evotomys caurinus.....	229		171
Evotomys dawsoni.....	229		172
Evotomys d. alascensis.....	229		
Evotomys fuscordorsalis.....	230		172
Evotomys galei.....	231		173
Evotomys gapperi	230		172
Evotomys g. breviceaudus.....	231		
Evotomys g. galei.....	231		173
Evotomys g. galii.....	231		
Evotomys g. loringi	231		173
Evotomys g. ochraceus.....	230		172
Evotomys g. rhoadsi	230		173
Evotomys g. saturatus.....	231		174
Evotomys idahoensis.....	232		175
Evotomys mazama.....	232		175
Evotomys nivarius.....	233		177
Evotomys obscurus	232		176
Evotomys occidentalis.. ..	233		176
Evotomys orca.....	230		172
Evotomys proteus	233		177
Evotomys rutilus.....	229, 230		170
Evotomys ungava	232		175
Evotomys wrangeli.....	229		171
excisum. (<i>Phyllostoma</i>)	529	715	
exiguus. (<i>Heteromys</i>)	321	377	
exiguus. (<i>Peromyscus</i>)	148	183	
exilipes. (<i>Blarina</i>)	463		384
exilis. (<i>Dipodomys m.</i>)	289		233
exilis. (<i>Vespertilio</i>)	476	578	403
eximius. (<i>Blarina</i>)	463		384
eximius. (<i>Microsorex</i>)	457		
eximius. (<i>Sorex</i>)	457		
extimus. (<i>Sciurus c.</i>)	80		55
eyra. (<i>Felis</i>)	368	453	295
falcatum. (<i>Phyllops</i>)	527	708	
falcatus. (<i>Arctibeus</i>)	527	707	
falcatus. (<i>Phyllops</i>)	527	707	
Falcifer	12	29	
Falcifer jubata	12	29	
fallax. (<i>Artibeus</i>)	524	695	
fallax. (<i>Geomys p.</i>)	264		220
fallax. (<i>Neotoma</i>)	216		161
fallax. (<i>Perognathus</i>)	314	366	254

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
familiaris. (Canis)	374	464	
fannini. (Ovis)	55		†425
fasciata. (Felis r.)	371		297
fasciata. (Phoca)	441		362
fasciatus. (Lynx)	371		297
fasciatus. (Perognathus)	300, 309		242, 251
fatuus. (Synaptomys)	256		204
Felidæ	365	442	293
felipensis. (Conepatus)	412	517	
felipensis. (Neotoma b.)	218	286	
felipensis. (Peromyscus)	167	205	
Felis	365, 369	443	293
Felis apache	368		
Felis a. browni	369		
Felis b. costaricensis	370	456	
Felis brasiliensis	367	449	
Felis cacomitli	367		
Felis canadensis	370		295
Felis c. mollipilosa	370		295
Felis c. subsolana	370		296
Felis carrikeri	366	448	
Felis catus	365	443	
Felis centralis	365		
Felis chati	367	449	
Felis concolor	369	293	293
Felis c. floridana	369		293
Felis c. oregonensis	369	454	294
Felis coryi	369		293
Felis costaricensis	366		
Felis cougar	369		
Felis discolor			293
Felis elegans	367	449	
Felis eyra	368	453	295
Felis c. apache	368	453	
Felis floridana			293
Felis fossata	368	453	
Felis gigas	373		299
Felis glaucula	367	450	
Felis hernandezi			294
Felis h. goldmani	366		
Felis hippolestes	369	454	294
Felis h. aztecus	369	454	
Felis h. olympus	369	454	294
Felis jaguarondi	367	451	295
Felis j. cacomitli	367	451	
Felis j. tolteca	368	452	
Felis macroura	367	449	
Felis maculata	371, 372		296
Felis mexicana			294

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Felis mitis</i>	367	449	
<i>Felis onca</i>	365		294
<i>Felis o. centralis</i>	365	446	
<i>Felis o. goldmani</i>	366	447	
<i>Felis o. hernandezi</i>	365, 366	446	
<i>Felis panamensis</i>	368	452	
<i>Felis pardalis</i>	366	447	294
<i>Felis p. mearnsi</i>	366	448	
<i>Felis p. oncilla</i>	367	450	
<i>Felis peninsularis</i>	373	460	
<i>Felis pennsylvanica</i>	369		
<i>Felis puma</i>			293
<i>Felis rufa</i>	370		296
<i>Felis r. baileyi</i>	372	459	297
<i>Felis r. californica</i>	372	458	298
<i>Felis r. eremica</i>	372	458	298
<i>Felis r. escuinapæ</i>	372, 373	456	
<i>Felis r. fasciata</i>	371		297
<i>Felis r. floridana</i>	372		299
<i>Felis r. maculata</i>	371		298
<i>Felis r. oculatea</i>	371		297
<i>Felis r. pallescens</i>	371		297
<i>Felis r. texensis</i>	371	458	298
<i>Felis ruffa</i>	370		
<i>Felis tigrina</i>	367	449	
<i>Felis uinta</i>	373		
<i>Felix oregonensis</i>	369		
<i>felix. (Tamias q.)</i>	92		76
<i>femoralis. (Perognathus)</i>	314	367	254
<i>femorosaccus. (Nyctinomops)</i>	498	626	
<i>femorosaccus. (Nyctinomus)</i>	498	626, 628	417
<i>fenestratus. (Dasypus m.)</i>	13	33	4
<i>ferox. (Molossus)</i>	497	623	
<i>ferreus. (Sciurus a.)</i>	66		
<i>ferruginea. (Neotoma)</i>	210, 211	280	
<i>ferrugineus. (Arvicola)</i>	175		144
<i>ferruginiventris. (Sciurus)</i>	68	115	
<i>festinus. (Lepus)</i>	354	†746	
<i>fetidissima. (Mephitis)</i>	406		
<i>Fiber</i>	252	307	211
<i>Fiber macrodon</i>	255		214
<i>Fiber obscurus</i>	254		214
<i>Fiber occipitalis</i>	254		
<i>Fiber osoyoosensis</i>	252		212
<i>Fiber spatulatus</i>	254		214
<i>Fiber zibethicus</i>	252, 254	307	212
<i>Fiber z. aquilonius</i>	253		213
<i>Fiber z. hudsonius</i>	253		
<i>Fiber z. pallidus</i>	253	308	213

†Appendix.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
Fiber z. ripensis	252		
Fiber z. rivalicius	253		213
fiber. (Castor)	128		115
fimbripes. (Sorex)	446		366
fisheri. (Citellus v.)	114	150	
fisheri. (Microtus a.)	243		190
fisheri. (Sorex)	454		375
fisheri. (Spermophilus b.)	114		
fisheri. (Spermophilus g.)	114	150	88
fisheri. (Thomomys d.)	277		
fisheri. (Thomomys f.)	277		
flaccidus. (Peromyscus t.)	147	189	
flava. (Talpa)	470		388
flava. (Talpa e.)	470		388
flavescens. (Ariteus)	527	707	
flavescens. (Perognathus)	301		243
flavescens. (Talpa e.)	469		388
flavescens. (Talpa f.)	469		388
flavidus. (Megadontomys)	172	214	
flavigularis. (Lepus)	353	431	288
flavigularis. (Lepus c.)	†543		
flaviventer. (Arctomys)	120		106
flaviventer. (Marmota)	120		
flavogularis. (Centurio)	530	717	
flavus. (Bassariscus a.)	398	485	316
flavus. (Cricetodipus)	302		243
flavus. (Perognathus)	302	354	244
floridana. (Arvicola)	209, 210		
floridana. (Blarina)	463		385
floridana. (Felis)			293
floridana. (Felis c.)	369		293
floridana. (Felis r.)	372		299
floridana. (Mus)	205, 209	276	154, 157
floridana. (Neotoma)	209		157
floridana. (Pseudostoma)	261		215
floridanus. (Dasypterus)	488		
floridanus. (Geomys t.)	261		215
floridanus. (Hesperomys)	156		138
floridanus. (Lepus)	342		280
floridanus. (Lepus s.)	340		
floridanus. (Lynx)	372		
floridanus. (Oryzomys n.)	181		
floridanus. (Peromyscus)	156		138
floridanus. (Urocyon c.)	386		307
floridanus. (Ursus)	395		314
floweri. (Kogia)	22	46	*479
fluviatilis. (Manatus)	14	37	6
foetida. (Phoca)	442, 443		363, 364
foetulenta. (Mephitis)	408		326

†Appendix.

*Supplement.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
fontigenus. (Microtus)	236		181
fontigenus. (Microtus p.)	236		181
formosus. (Perognathus)	308		250
forsteri. (Delphinus)	30	55	28
forsteri. (Sorex)	446		366
fossata. (Felis)	368	453	
fossor. (Blarina)	466	562	
fossor. (Sciurus)	78	130	55
fossor. (Thomomys)	287		230
fournieri. (Capromys)	330	390	
franklini. (Citellus)	105		
franklini. (Spermophilus)	105		101
franklinii. (Arctomys)	105		101
frantzii. (Atalapha)	486	594	
frater. (Tamias)	90		73
fraterculus. (Peromyscus)	155	184	136
fraterculus. (Urocyon c.)	389	475	
fraterculus. (Vesperimus)	155		136
frederici. (Phoca)	444		364
fremonti. (Sciurus)	84		66
frenata. (Mustela)	431		350
frenatus. (Delphinus)	34		32
frenatus. (Prodelphinus)	34		32
frenatus. (Putorius)	431, 432	532	350
freyreissii. (Didelurus)	494	615	
frondator. (Castor c.)	128	163	117
frontalis. (Delphinus)	34		32
frontatus. (Eriodes)	535	733	
frumentor. (Sciurus a.)	69	117	
frustror. (Canis)	377		302
fulgens. (Oryzomys)	183	240	
fuliginosa. (Chilonycteris)	504	641	
fuliginosa. (Chilonycteris m.)	504	†748	
fuliginosus. (Ateles)	534	733	
fuliginosus. (Molossus)	495	620	417
fuliginosus. (Onychomys)	134		123
fuliginosus. (Perognathus)	302		244
fuliginosus. (Perognathus f.)	302		244
fuliginosus. (Sciuropterus a.)	123		
fuliginosus. (Sciuropterus v.)	123		111
fuliginosus. (Sciurus)	79		
fuliginosus. (Sciurus c.)	79		55
fuliginosus. (Trachops)	514	664	
fulva. (Arvicola)	235		180
fulva. (Vulpes)	380		
fulvescens. (Cratogeomys)	266	316	
fulvescens. (Hesperomys)	189	248	
fulvescens. (Oryzomys)	189	248	
fulvescens. (Reithrodontomys m.)	198	264	

†Appendix.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
fulvescens. (Rhithrodontomys m.).....	198	264, 265	
fulvi-fasciatus. (Delphinus).....	30	54	28
fulviventer. (Marmosa).....	3	8	
fulviventer. (Microtus).....	246	301	
fulviventer. (Neotoma).....	220	288	
fulviventer. (Sigmodon).....	180	232	
fulvo-fasciatus. (Delphinus).....	30	54	28
fulvus. (Canis).....	380		304
fulvus. (Chilonycteris d.).....	506		
fulvus. (Dermonotus d.).....	506	646	
fulvus. (Geomys).....	272, 280		227
fulvus. (Peromyscus l.).....	142	183	
fulvus. (Peromyscus s.).....	142	183	
fulvus. (Thomomys).....	272, 280	334	227
fumarium. (Phyllostoma).....	529		
fumarius. (Dysopes).....	495	620	
fumarius. (Molossus).....	496	621	
fumeus. (Sorex).....	447		369
fumosus. (Geomys).....	267	318	
fumosus. (Platygeomys).....	267	318	
funebri. (Lasiurus).....	485	593	412
furvus. (Peromyscus).....	169	208	
furvus. (Sigmodon h.).....	178	225	
fusca. (Diplostoma).....	262		217
fusca. (Neotoma c.).....	222		165
fusca. (Neotoma o.).....	222		
fusca. (Talpa).....	470		388
fusca. (Teonoma).....	222		
fuscipes. (Neotoma).....	206	278, 279	158
fuscodorsalis. (Evotomys).....	230		172
fuscogriseus. (Metachirus).....	5	12	
fuscovariegatus. (Sciurus).....	76	128	
fuscus. (Delphinus).....	31	55	28
fuscus. (Desmodus).....	530	719	
fuscus. (Thomomys c.).....	277		
fuscus. (Thomomys d.).....	277		226
fuscus. (Vespertilio).....	482, 483	587	410
fuscus. (Peromyscus l.).....	†542		
gabbi. (Bassaricyon).....	400	487, 489	
gabbi. (Lepus).....	351	426	
gabbi. (Lepus b.).....	351		
gabbii. (Bassaricyon).....	400		
gadovi. (Peromyscus l.).....	165	201	
gaillardi. (Lepus).....	354	433	
Gale.....	423	530	338
galei. (Evotomys).....	231		173
galei. (Evotomys g.).....	231		173
Galemys.....	460	556	382

†Appendix.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Galemys micrusus</i>	461		382
<i>Galera</i>	418	524	
<i>Galera allamandi</i>	418		
<i>Galera b. biologiae</i>	418		
<i>Galera b. senex</i>	418		
<i>Galera canaster</i>	419		
<i>Galictis</i>	418	524	
<i>Galictis allamandi</i>	418	526	
<i>Galictis b. biologiae</i>	418		
<i>Galictis b. senex</i>	418	525	
<i>Galictis canaster</i>	419	526	
<i>Galidictis</i>	418	524	
<i>galii</i> . (<i>Evotomys g.</i>).....	231		
<i>gambeli</i> . (<i>Peromyscus t.</i>).....	145, 149		130
<i>gambelii</i> . (<i>Hesperomys</i>).....	145		130
<i>gapperi</i> . (<i>Arvicola</i>).....	230		172
<i>gapperi</i> . (<i>Evotomys</i>).....	230		172
<i>gaumeri</i> . (<i>Adelonycteris</i>).....	485	590	
<i>gaumeri</i> . (<i>Heteromys</i>).....	317	371	
<i>gaumeri</i> . (<i>Vespertilio</i>).....	485	590	
<i>gaurus</i> . (<i>Peromyscus</i>).....	148	199	
<i>gentilis</i> . (<i>Peromyscus g.</i>).....	164	197	
<i>Genyscelus</i>	337	408	
<i>Geocapromys</i>	330, 331	389, 393	
<i>geoffroyi</i> . (<i>Anoura</i>).....	519		
<i>geoffroyi</i> . (<i>Anura</i>).....	519, 520	682	
<i>geoffroyi</i> . (<i>Ateles</i>).....	535	733	
<i>geoffroyi</i> . (<i>Hapale</i>).....	532	724	
<i>geoffroyi</i> . (<i>Midas</i>).....	532	724	
<i>geoffroyi</i> . (<i>Edipomidas</i>).....	532		
<i>Geomyidae</i>	261	309	215
<i>Geomys</i>	261	310	215
<i>Geomys arenarius</i>	264	310	219
<i>Geomys borealis</i>	275, 278		224, 227
<i>Geomys breviceps</i>	263		218
<i>Geomys b. atwateri</i>	263		219
<i>Geomys b. sagittalis</i>	263		218
<i>Geomys bulleri</i>	268	320	
<i>Geomys bursarius</i>	262, 263		217
<i>Geomys b. lutescens</i>	263		218
<i>Geomys cherriei</i>	271	330	
<i>Geomys cinerea</i>	262		217
<i>Geomys clarkii</i>	266	315	
<i>Geomys colonus</i>	262		217
<i>Geomys cumberlandius</i>	262		217
<i>Geomys douglasii</i>	277		226
<i>Geomys f. austrinus</i>	262		
<i>Geomys fulvus</i>	272, 280	334	227
<i>Geomys fumosus</i>	267	318	
<i>Geomys grandis</i>	268	322	

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
Geomys gymnurus.....	266	318	
Geomys heterodus.....	270	327	
Geomys hispidus.....	269	326	
Geomys lutescens.....	263		218
Geomys merriami.....	264		
Geomys nelsoni.....	268	320	
Geomys oregonensis.....	262		217
Geomys personatus.....	264		220
Geomys p. fallax.....	264		220
Geomys pinetis.....	261		215
Geomys scalops.....	269	324	
Geomys texensis.....	264		219
Geomys townsendii.....	275		224
Geomys tuza.....	261		215
Geomys t. austrinus.....	262		217
Geomys t. floridanus.....	261		215
Geomys t. mobilensis.....	262		216
Geomys umbrinus.....	280		
Gerbillus.....	324		
Gerboa.....	119		
geronimensis. (Peromyscus).....	162	181	
geronimensis. (Phoca r.).....	443	542	
gibbar. (Balænoptera).....	18	42	
gibbosus. (Agaphelus).....	18		11
gibbsi. (Neurotrichus).....	466		387
gibbsii. (Urotrichus).....	466		387
gigas. (Alce).....	41		38
gigas. (Alces).....	41		
gigas. (Felis).....	373		299
gigas. (Hydrodamalis).....	14	35	5
gigas. (Lynx).....	373		299
gigas. (Manatus).....	14		5
gigas. (Notiosorex).....	460	555	
gilberti. (Sitomys).....	150	195	132
gillespii. (Otaria).....	437	539	356
gilli. (Tursiops).....	32	56	29
gilva. (Neotoma i.).....	217		162
gilvus. (Perognathus m.).....	302		243
glacialis. (Balæna).....	15		8
glacialis. (Lepus).....	358		270
glacialis. (Sorex).....	451		372
gladiator. (Delphinus).....	26	51	22
gladiator. (Orcinus).....	26	51	22
glaucinus. (Dysopes).....	497	623	
glaucinus. (Promops).....	497	623	
glaucula. (Felis).....	367	450	
glaucus. (Agaphelus).....	16	40	9
glaucus. (Rhachianectes).....	16	40	9
Glis.....	119		104
Glis canadensis.....	120		106

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
Globicephala.....	27	52	
Globicephalus.....	27	52	
Globicephalus brachypterus.....	28	53, 54	
Globicephalus melas.....	27	47, 52	
Globicephalus scammoni.....	28	54	
globiceps. (Delphinus).....	27	52	23
Globiocephalus.....			23
Globiocephalus brachypterus.....	28	53	24
Globiocephalus grayi.....	27	51	23
Globiocephalus incrassatus.....	27	53	23
Globiocephalus macrorhynchus.....	27	53	23
Globiocephalus melas.....	27	53	23
Globiocephalus scammoni.....	28	54	24
Glossonycteris.....	519	681	
Glossonycteris lasiopyga.....	520	682	
Glossophaga.....	516	670	
Glossophaga amplexicaudata.....	516	672	
Glossophaga caudifera.....	519, 520		
Glossophaga mutica.....	516	671	
Glossophaga soricina.....	516	672	
Glossophaga s. antillarum.....	517	672	
Glossophaginae.....	516	670	
Glyphonycteris.....	513	663	
Glyphonycteris sylvestris.....	513	664	
godmani. (Chceronycteris).....	517	674	
godmani. (Mustela).....	422		335
godmani. (Sorex).....	455	552	
goldmani. (Blarina m.).....	465	560	
goldmani. (Canis).....	378		
goldmani. (Citellus a.).....	116	151	
goldmani. (Cratogeomys c.).....	266	316	
goldmani. (Felis h.).....	366		
goldmani. (Felis o.).....	366	447	
goldmani. (Heteromys).....	318	373	
goldmani. (Lepus).....	349		†745
goldmani. (Lepus a.).....	349		
goldmani. (Nelsonia).....	224	290	
goldmani. (Neotoma).....	220	288	
goldmani. (Oryzomys).....	186	246	
goldmani. (Perodipus).....	294		
goldmani. (Perognathus).....	313	365	
goldmani. (Peromyscus).....	138	186	
goldmani. (Putorius f.).....	432	534	
goldmani. (Reithrodontomys).....	201	268	
goldmani. (Rhithrodontomys).....	201	268	
goldmani. (Sciuropterus v.).....	126		
goldmani. (Sciurus).....	77	130	
goldmani. (Spermophilus a.).....	116	151	
goldmani. (Thomomys).....	283	339	

†Appendix.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
gossi. (<i>Synaptomys c.</i>)	255		204
gossii. (<i>Synaptomys h.</i>)	255		204
Gossypinus.	143		
gossypinus. (<i>Peromyscus</i>)	143		
Gossypinus.	143		
gossypinus. (<i>Hesperomys</i>)	143		126
gossypinus. (<i>Peromyscus</i>)	143		126
Gossypion.	143		
gracilis. (<i>Dysopes</i>)	498	626	
gracilis. (<i>Enydris</i>)	436	537	354
gracilis. (<i>Hesperomys</i>)	144		128
gracilis. (<i>Nyctinomops</i>)	498	626	
gracilis. (<i>Reithrodontomys m.</i>)	198	265	
gracilis. (<i>Rhithrodontomys m.</i>)	198	265	
gracilis. (<i>Rhogeessa</i>)	498		
gracilis. (<i>Rhogeessa</i>)	489	602	
gracilis. (<i>Spilogale</i>)	415		330
gracilis. (<i>Tamias a.</i>)	91		76
grahamensis. (<i>Sciurus f.</i>)	85		67
grahamensis. (<i>Sciurus h.</i>)	85		
grammurus. (<i>Citellus v.</i>)	113, 114	149, 150	
grammurus. (<i>Sciurus</i>)	113	149	88
grammurus. (<i>Spermophilus</i>)	113	149	
Grampus.	28		24
Grampus cuvieri.	28		25
Grampus griseus.	28		25
Grampus stearnsii	28		25
grandis. (<i>Geomys</i>)	268	322	
grandis. (<i>Neotoma</i>)	215		
grandis. (<i>Neotoma d.</i>)	215		
grandis. (<i>Orthogeomys</i>)	268, 269	322	
grangeri. (<i>Lepus</i>)	349, 350		285
grangeri. (<i>Lepus s.</i>)	349		
grangeri. (<i>Neotoma</i>)	223		165
grangeri. (<i>Teonoma</i>)	223		
granti. (<i>Rangifer</i>)	39		
gratus. (<i>Peromyscus</i>)	164	196	
grayi. (<i>Globiocephalus</i>)	27	51	23
grayi. (<i>Phyllostoma</i>)	515	669	
graysoni. (<i>Lepus</i>)	351	426	
grebnitzkii. (<i>Ziphius</i>)	23		*480
greenii. (<i>Scotophilus</i>)	483	587	410
grisea. (<i>Chilonycteris</i>)	504		
grisea. (<i>Chilonycteris m.</i>)	504	†747	
grisea. (<i>Cynomys</i>)	117		102
griseiflavus. (<i>Rhithrodontomys</i>)	202	270	
griseiflavus. (<i>Sciurus</i>)	74	124	
griseocaudatus. (<i>Sciurus</i>)	77		

*Supplement.

†Appendix.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
griseoflavus. (Macroxus)	74		
griseoflavus. (Reithrodontomys)	202	270	
griseoflavus. (Sciurus)	64	110	58
griseogena. (Sciurus)	60, 61	101, 104	
griseogenys. (Sciurus)	61	104	
grisescens. (Ateles)	535	734	
griseus. (Arctopithecus)	9	23	
griseus. (Bradypus)		24	
griseus. (Canis)	375, 386		307
griseus. (Canis l.)	375		300
griseus. (Delphinus)	28		25
griseus. (Grampus)	28		25
griseus. (Heteromys)	319	373	
griseus. (Lepus t.)	356	436	
griseus. (Lupus)	375		
griseus. (Sciurus)	78	130	55
griseus. (Tamias s.)	96		81
Grison	418	524	
Grison allamandi	418	526	
Grison b. biologiæ	418	524	
Grison b. senex	418	525	
Grison canaster	419	526	
Grisonia	418	524	
groenlandicus. (Lepus)	359		271
groenlandicus. (Mus)			209
groenlandica. (Phoca)	444		364
groenlandicus. (Cervus)	39		37
groenlandicus. (Rangifer)	39		37
Grymæomys	2	5	
Grymæomys dorsiger	2		
Grymæomys impavida	2		
Grymæomys murinus	2		
gryphus. (Vespertilio)	478		405
grypus. (Halichoerus)	441		361
grypus. (Phoca)	441		363
guadalupensis. (Microtus m.)	245		
guatemalæ. (Urocyon)	389		
guatemalæ. (Urocyon c.)	389	475	
guatemalensis. (Microtus)	251	306	
guatemalensis. (Peromyscus)	171	211	
gubernator. (Lagenorhynchus)	29		25
Guerlinguetus	59, 61	93, 103	50
guianensis. (Mustela)	1	3	
Gulo	416		333
Gulo hylæus	417		
Gulo luscus	416		333
Gulo luteus	417		
gulo. (Mustela)	416		
gundlachi. (Capromys p.)	331	392	
gunnisoni. (Cynomys)	118		103

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
gunnisoni. (Spermophilus)	118		
guttatus. (Arctomys).....	111		92
gyas. (Ursus d.)	391		
gymnicus. (Sciurus h.)	80		61
Gymnotis.....	43	68	
gymnotis. (Peromyscus).....	162	188	
gymnurus. (Geomys)	266	318	
gymnurus. (Platygeomys)	266, 267	318	
Gypsophoca	438		
Haematomycteris	530	720	
haidarum. (Putorius).....	429		351
Halarctus	438		
Halichoerus	440		361
Halichoerus grypus	441		361
Halicyon richardii.....	442		
Halophilus	441	541	361
hallensis. (Vulpes)	382		307
Hapale geoffroyi.....	532	724	
Hapale œdipus	532	724	
Haplodon californica	127		
Haplodontia	126		112
Haplodontia olympica.....	127, 128		114
Haplodontia pacifica	127		114
Haplodontia phæa	127		114
Haplodontia rufa	126, 127		112
Haplodontia r. californica.....	127		112
Haplodontia r. rainieri	127		112
Haplodontidæ	126		111
Haplomylomys	135, 136	170	
hardyi. (Zapus h.)	324		257
harrimani. (Vulpes).....	380		303
harrisi. (Citellus)	96	141, 143	
harrisii. (Spermophilus)	96	141	85
hastalis. (Cervus)	37		35
hastatum. (Phyllostoma)	514	666	
hastatus. (Vespertilio)	514	666	
haydeni. (Microtus)	248		197
haydeni. (Sorex).....	446		367
haydenii. (Arvicola).....	248		197
hebes. (Vulpes v.)	384		
heermanni. (Sciurus)	78	130	55
helaletes. (Synaptomys)	255		204
Heliophoca	444	542	
helleri. (Dipodomys d.)	292		
helleri. (Perognathus.).....	312	360	
helleri. (Thomomys)	278		
helleri. (Vampyrops).....	526	703	
helvolus. (Arvicola)	259		208
helvolus. (Lemmus t.)	259		208

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
helvolus. (Reithrodontomys g.)	202	270	
helvolus. (Rhithrodontomys g.)	202	270	
Hemiderma	515	667	
Hemiderma brevicaudum	515	669	
Hemiderma castaneum	516	670	
Hemiderma perspicillatum	515	668	
hemionotis. (Peromyscus)	139	199	
hemionus. (Cervus)	48	76	42
hemionus. (Odocoileus)	48	76	42
hemionus. (Odontocœlus)	48	76, 77, 78	
Hemiotomys	234	299	178
henshawi. (Vespertilio n.)	476	578	404
hermanni. (Dipodomys)	293		236
hernandezi. (Felis)			294
hernandezi. (Felis o.)	365, 366	446	
hernandezi. (Procyon l.)	401	491	317
hernandezi. (Sciurus a.)	70	118	
hernandezi. (Sciurus p.)	70	118	
hernandezii. (Leopardus)	365	446	
hernandezii. (Procyon)	400		
Herpailurus	365	443	
Herpestes	373	461	
Herpestes mungo	374	462	
Herpetomys	234, 251	306	179
herroni. (Peromyscus)	155		137
herronii. (Sitomys)	155		137
Hesperomys	135		
Hesperomys affinis	161	180	
Hesperomys alfaroi	183, 184	242	
Hesperomys anthonyi	137	185	141
Hesperomys austerus	149		132
Hesperomys aztecus	162	184	
Hesperomys boylii	150	195	132
Hesperomys caliginosus	190	253	
Hesperomys cherrii	160	183	
Hesperomys cognatus			127
Hesperomys couesi	186	236	
Hesperomys crinitus	159		140
Hesperomys eremicus	136	191	136
Hesperomys floridanus	156		138
Hesperomys fulvescens	189	248	
Hesperomys gambelii	145		130
Hesperomys gossypinus	143		126
Hesperomys gracilis	144		128
Hesperomys l. arcticus	146		131
Hesperomys l. deserticolus	140	188	
Hesperomys l. nebrascensis	146		
Hesperomys l. rufinus	142		126
Hesperomys maniculatus	148		131
Hesperomys megalotis	157		139

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Hesperomys melanophrys</i>	166	201	
<i>Hesperomys mexicanus</i>	168	207	
<i>Hesperomys myoides</i>	144		128
<i>Hesperomys niveiventris</i>	138		141
<i>Hesperomys nudicaudus</i>	173	217	
<i>Hesperomys nudipes</i>	165	198	
<i>Hesperomys palustris</i>	180		
<i>Hesperomys sonoriensis</i>	141	181	125
<i>Hesperomys sumichrasti</i>	172	216	
<i>Hesperomys taylori</i>	135		141
<i>Hesperomys teguina</i>	204	273	
<i>Hesperomys texana</i>	145	186	
<i>Hesperomys toltecus</i>	177	226	
<i>Hesperomys torridus</i>	133		
<i>Hesperomys truei</i>	159		139
<i>Hesperosciurus</i>	59, 78	94, 130	50, 55
<i>hesperus</i> . (<i>Pipistrellus</i>)	480	582	409
<i>hesperus</i> . (<i>Scotophilus</i>)	480	582	409
<i>hesperus</i> . (<i>Thomomys</i>)	279		
<i>heterodus</i> . (<i>Geomys</i>)	270	327	
<i>heterodus</i> . (<i>Macrogeomys</i>)	270	328	
<i>Heterogeomys</i>	269	325	
<i>Heterogeomys hispidus</i>	269	326	
<i>Heterogeomys torridus</i>	270	326	
<i>Heteromyidæ</i>	288	341	231
<i>Heteromyinæ</i>	300	352	241
<i>Heteromys</i>	316	368, 369	255
<i>Heteromys adspersus</i>	319	373	
<i>Heteromys albolimbatus</i>	320	375	
<i>Heteromys alleni</i>	321	376	256
<i>Heteromys annectens</i>	317	371	
<i>Heteromys bulleri</i>	317	369	
<i>Heteromys canus</i>	321	375	
<i>Heteromys crispus</i>	320	380	
<i>Heteromys c. setosus</i>	320	380	
<i>Heteromys desmarestianus</i>	318	372	
<i>Heteromys exiguus</i>	321	377	
<i>Heteromys gaumeri</i>	317	371	
<i>Heteromys goldmani</i>	318	373	
<i>Heteromys g. lepturus</i>	318	373	
<i>Heteromys griseus</i>	319	373	
<i>Heteromys heterothrix</i>	320	380	
<i>Heteromys hispidus</i>	318	371	
<i>Heteromys irroratus</i>	316	369	
<i>Heteromys longicaudatus</i>	317, 318	370, 372	
<i>Heteromys nelsoni</i>	323	381	
<i>Heteromys obscurus</i>	319	379	
<i>Heteromys orbitalis</i>	320	380	
<i>Heteromys paralius</i>	320	381	
<i>Heteromys parviceps</i>	322	378	

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Heteromys phæura</i>	319	379	
<i>Heteromys phæurus</i>	319		
<i>Heteromys pictus</i>	322	377	
<i>Heteromys p. isthmius</i>	322	378	
<i>Heteromys p. rostratus</i>	322	378	
<i>Heteromys plantinarenensis</i>	322	377	
<i>Heteromys repens</i>	318	372	
<i>Heteromys salvini</i>	317	370	
<i>Heteromys s. nigrescens</i>	317	370	
<i>Heteromys sonorana</i>	323	379	
<i>Heteromys texensis</i>	323		
<i>Heteromys torridus</i>	321	376	
<i>Heteromys t. minor</i>	321	377	
<i>Heteromys veræcrucis</i>	319	379	
<i>heterothrix</i> . (<i>Heteromys</i>)	320	380	
<i>heterothrix</i> . (<i>Liomys</i>)	320	380	
<i>hiemalis</i> . (<i>Sciurus</i>)	79		53
<i>hindsii</i> . (<i>Tamias</i>)	86		
<i>hindsii</i> . (<i>Tamias t.</i>)	86		70
<i>hindsii</i> . (<i>Tamias t.</i>)	86		
<i>Hippocamelus</i>	50		
<i>Hippocamelus dubius</i>	50		
<i>Hippocamelus pandora</i>	51		
<i>Hippocamelus sartori</i>	50		
<i>hippolestes</i> . (<i>Felis</i>)	369	454	294
<i>hirsutum</i> . (<i>Schizostoma</i>)	512	661	
<i>hirsutus</i> . (<i>Arvicola</i>)	235		179
<i>hirsutus</i> . (<i>Micronycteris</i>)	512	661	
<i>hirsutus</i> . (<i>Reithrodontomys</i>)	202	269	
<i>hirsutus</i> . (<i>Rhithrodontomys</i>)	202	269, 270	
<i>hirtus</i> . (<i>Sciurus n.</i>)	72	121	
<i>hispida</i> . (<i>Phoca</i>)	443		364
<i>hispidus</i> . (<i>Geomys</i>)	269	326	
<i>hispidus</i> . (<i>Heterogeomys</i>)	269	326	
<i>hispidus</i> . (<i>Heteromys</i>)	318	371	
<i>hispidus</i> . (<i>Perognathus</i>)	309	360	251
<i>hispidus</i> . (<i>Sigmodon</i>)	174, 175	222, 232	144
<i>hispidus</i> . (<i>Xenurus c.</i>)	12	32	
<i>Histiops</i>	526	704	
<i>Histiotus</i>	490		398
<i>Histiotus maculatus</i>	490		398
<i>Histriophoca</i>	441		361
<i>Hodomys</i>	226	296	
<i>Hodomys alleni</i>	226	297	
<i>Hodomys vetulus</i>	226	297	
<i>hoffmanni</i> . (<i>Choloepus</i>)	9	20	
<i>hoffmanni</i> . (<i>Sciurus æ.</i>)	61	104	
<i>Holochilus</i>		221	
<i>Holochilus pilorides</i>		221	

	Check-List of Mammals.	Mammals. of Middle America, etc.	Synopsis of Mammals, etc.
holosericeus. (Dysopes).....	495	619	
holzneri. (Lepus).....	345		284
holzneri. (Lepus f.).....	345	419	
holzneri. (Lepus s.).....	345		
holzneri. (Mephitis o.).....	409	507	
homochroia. (Peromyscus).....	148	200	
hoodi. (Arctomys).....	104		
hopiensis. (Eutamias).....	†541		
hopiensis. (Tamias).....	†541		
horriæus. (Ursus).....	393	480	
horriæus. (Ursus h.).....	393		312
horribilis. (Ursus).....	392, 393		312
hortensis. (Arvicola).....	175		144
hoyi. (Microsorex).....	457		377
hoyi. (Sorex).....	456, 457		377
huachuca. (Sciurus a.).....	64	109	60
hudsonia. (Mephitis).....	406		322
hudsonia. (Mephitis m.).....	406		
hudsonica. (Lutra).....	433		352
hudsonica. (Mephitis a.).....	406		
hudsonicus. (Sciurus v.).....	80		
hudsonius. (Dicrostonyx).....	260		209
hudsonius. (Dipus).....	323, 324		257
hudsonius. (Fiber z.).....	253		
hudsonius. (Hystrix).....	333		265
hudsonius. (Jaculus).....	328		
hudsonius. (Lepus).....	360		273
hudsonius. (Mus).....	260		209
hudsonius. (Sciurus).....	80, 124		60, 107
hudsonius. (Zapus).....	324		257
hulophaga. (Blarina b.).....	461		
humboldtii. (Conepatus).....	410	512	
humerales. (Tagassu a.).....	35	63	
humeralis. (Nycticeius).....	488	598	
humeralis. (Nycticejus).....	488	598	415
humeralis. (Tayassu a.).....	35	63	
humeralis. (Vespertilio).....	488	597	415
humilis. (Ochetodon).....	192		151
hunteri. (Delphinus).....	21		16
Hunterius.....	15		7
Hunterus.....	15		
huro. (Mustela).....	419	335	
hyacinthinus. (Neurotrichus g.).....	467		388
hybridus. (Ateles).....	535	733	
Hydrodamalidæ.....	14		5
Hydrodamalis.....	14		5
Hydrodamalis australis.....		36	
Hydrodamalis dugong.....		36	
Hydrodamalis gigas.....	14	35	5

†Appendix.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
Hydrodamalis stelleri.....	14		
Hydrodamalis tabernaculi.....		36	
hydrodromus. (Neosorex).....	458		379
hydrodromus. (Sorex).....	458		379
Hydrolagus.....	340	413	269
hylæus. (Gulo).....	417		
hylocetes. (Oryzomys).....	187	237	
hylocetes. (Peromyscus).....	167	204	
hylodromus. (Ursus).....	396		
Hylonycteris.....	517	675	
Hylonycteris underwoodi.....	517, 518	675	
hylophaga. (Blarina b.).....	461		383
Hyonycteris.....	503	637	
Hyonycteris discifera.....	503	637	
hyperythrus. (Microtus c.).....	240	302	
hypoleuca. (Simia).....	536	736	
hypoleucus. (Cebus).....	536	736	
hypophæus. (Sciurus c.).....	79		54
hypopyrrhus. (Sciurus).....	62, 68, 73	121, 126	
.....	75, 76, 77	127, 128	
hypopyrrhus. (Sciurus a.).....	68	116	
hypoxanthus. (Sciurus).....	68	115	
Hypsugo.....	480	582	408
Hypudæus.....	130		119
Hypudæus leucogaster.....	130		119
Hypudæus ochrogaster.....	248		196
Hystrix dorsata.....	333		
Hystrix dorsatus.....	332	397	264
Hystrix hudsonius.....	333		265
Hystrix mexicana.....	335	402	
Hystrix novæ-hispaniæ.....	335	402	
Hystrix pilosus.....	333	397	264
Hystrix prehensilis.....	334, 335	399, 402	
Hyperoodon.....	21		15
Hyperoodon latifrons.....	21		16
Hyperoodon rostratus.....	21		16
hyperoodon. (Delphinus).....	21		16
ibsenii. (Delphinus).....	29		26
ichneumon. (Viverra).....	373	461	
Ictidomys.....	96, 104	139	82, 99
Ictis.....	423	529	338
Ictonyx.....	410	512	
idahoensis. (Evotomys).....	232		175
idahoensis. (Lepus).....	351		287
idahoensis. (Sorex).....	446		367
idahoensis. (Thomomys).....	284		
Ignavus.....	9	21	
ignavus. (Arctomys).....	119		

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
ignavus. (Arctomys m.).....	119		105
ignavus. (Marmota m.).....	119		
imitator. (Cebus).....	536	737	
impavida. (Didelphys).....	2		
impavida. (Grymæomys).....	2		
impavidus. (Canis).....	379	468	
imperator. (Zapus).....	326		261
impiger. (Reithrodontomys l.).....	191		149
impiger. (Rhithrodontomys l.).....	191		
incautus. (Myotis).....	473		
incautus. (Vespertilio).....	473		
incitatus. (Lepus).....	351	428	
incrassatus. (Globiocephalus).....	27	53	23
incrassatus. (Sphærocephalus).....	27	53	23
indianola. (Spilogale).....	413		329
inexoratus. (Sigmodon h.).....	177	229	
inexpectatus. (Rhithrodontomys).....	203	271	
inflata. (Chilonycteris m.).....	504	†747	
infraluteus. (Perognathus).....	301		242
infraluteus. (Perognathus f.).....	301		242
infusca. (Balantiopteryx).....	493	611	
infusca. (Saccopteryx).....	493	611	
infusca. (Taxidea t.).....	405	505	
infuscatus. (Bradypus).....	9	23	
ingens. (Lutreola v.).....	424		
ingens. (Perodipus).....	298		
ingens. (Putorius v.).....	424, 425		340
ingrahami. (Capromys).....	332	394	
innuitus. (Microtus).....	242		189
innuitus. (Mictomys).....	256		205
innuitus. (Synaptomys).....	256		205
innuitus. (Vulpes l.).....	382, 383		
inornatus. (Perognathus).....	306		247
Insectivora.....	446	547	366
insignis. (Peromyscus).....	155	203	138
insignis. (Zapus).....	328		263
insolatus. (Peromyscus).....	154		135
insolatus. (Sitomys).....	154		135
insolitus. (Lepus).....	348	421	
insonus. (Lepus).....	341	†743	
insperatus. (Arvicola).....	236		180
insulanus. (Peromyscus).....	144		128
insularis. (Artibeus).....	522		
insularis. (Didelphis m.).....	7	17	
insularis. (Didelphys m.).....	7	17	
insularis. (Lepus).....	358	438	
insularis. (Marmosa).....	3	7	
insularis. (Microtus).....	238		182
insularis. (Procyon l.).....	401	492	

†Appendix.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
intermedia. (Mephitis v.)	410		
intermedia. (Neotoma)	216	284	161
intermedius. (Artibeus)	523	694	
intermedius. (Dasypterus)	487, 488	595	414
intermedius. (Delphinus)	27, 28	52	23, 25
intermedius. (Lasiurus)	487	595	414
intermedius. (Perognathus)	313	364	253
intermedius. (Phenacomys)	226		167
intermedius. (Reithrodontomys a.)	198		
intermedius. (Reithrodontomys m.)	198	264	153
intermedius. (Rhithrodontomys)	198	264	
intermedius. (Scalops a.)	470		390
intermedius. (Scalops m.)	470		
intermedius. (Scalopus a.)	470		
intermedius. (Sciurus)	76	127	
intermedius. (Thomomys f.)	280	336	228
interpres. (Citellus)	98	143	
interpres. (Spermophilus)	98	143	86
interpres. (Tamias)	98		86
interrupta. (Mephitis)	412		
interrupta. (Spilogale)	412	523	328
inunguis. (Trichechus)		35	
inoensis. (Eutamias s.)	89		
inoensis. (Tamias c.)	89		73
inoensis. (Tamias s.)	89		
inoensis. (Urocyon c.)	387		
irazu. (Acodon)	204	274	
irazu. (Akodon)	204	274	
irroratus. (Heteromys)	316	369	
Ischnoglossa	519	679	
Ischnoglossa nivalis	519	679	
isidorei. (Phoca)	440		360
Isodon pilorides	330	390	
Isotus	473	571	400
isthmica. (Dasypsecta)	336	407	
isthmica. (Neotoma)	219	287	
isthmus. (Heteromys p.)	322	378	
Jaculus	324		
Jaculus americanus	324		257
Jaculus hudsonius	328		
Jaguaris	365	443	
jaguarondi. (Felis)	367	451	295
jalapæ. (Mus m.)	130	163	
jalapæ. (Oryzomys)	184	246, 247	
jalapæ. (Reithrodontomys c.)	200	267	
jalapæ. (Rhithrodontomys c.)	200	267	
jaliscensis. (Myotis c.)	477	579	
jamaicensis. (Artibeus)	522, 527	693, 707	
jamaicensis. (Macrotus w.)	508		

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
jamaicensis. (Otopterus w.).....	508		
janira. (Delphinus).....	30	55	28
jeffersoni. (Meles).....	404		320
jubata. (Eumetopias).....	437		
jubata. (Myrmecophaga).....	12	29	
jubata. (Otaria).....	437		
jubata. (Phoca).....	437		355
kadiacensis. (Citellus p.).....	108		
kadiacensis. (Microtus).....	242		188
kadiacensis. (Putorius a.).....	428		346
kadiacensis. (Spermophilus e.).....	108		90
kaibabensis. (Sciurus).....	67		
keeni. (Peromyscus).....	151		133
keeni. (Sitomys).....	151		133
keenii. (Myotis s.).....	478		405
keenii. (Vespertilio s.).....	478		405
kenaiensis. (Mustela).....	421		
kenaiensis. (Mustela a.).....	421		
kenaiensis. (Ovis d.).....	55		
kenaiensis. (Ursus).....	392		
kenaiensis. (Vulpes).....	382		†432
kennedyi. (Oreamnus).....	53		44
kennicotti. (Arvicola).....	249		198
kennicottii. (Arctomys).....	107		89
keporkak. (Balæna).....	17		10
kermodei. (Ursus).....	397		
kermodii. (Ursus).....	397		
kidderi. (Ursus).....	391		
klamathensis. (Lepus).....	362		276
klamathensis. (Reithrodontomys).....	196		152
klamathensis. (Rhithrodontomys).....	196		
klamathensis. (Sciuropterus a.).....	123		
klamathensis. (Sciuropterus v.).....	123		111
knoxii. (Benedenia).....	20		13
kodiacensis. (Spermophilus e.).....	108		
kodiacensis. (Spermophilus p.).....	108		
Kogia.....	22	46	*479
Kogia breviceps.....	22		*479
Kogia floweri.....	22	46	*479
kracheninikowi. (Otaria).....			357
kuzira. (Balæna).....	16		9
kuzira. (Megaptera).....	16		9
Kyphobalæna.....	17	40	9
labecula. (Peromyscus).....	142	178	
labiatus. (Dicotyles).....	37	66	
labilis. (Loncheres).....	328	384	

†Appendix.

*Supplement.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
Laboura.....	334	400	
labradoria. (Ursus).....	404		320
labradorius. (Dipus).....	324		257
labradorius. (Lepus).....	359		271
labradorius. (Meriones).....	328		263
labradorius. (Microtus p.).....	236		181
laceyi. (Reithrodontomys).....	199		153
laceyi. (Rhithrodontomys).....	199		
lachuguilla. (Thomomys a.).....	285		
lacustris. (Lutreola v.).....	424		
lacustris. (Putorius v.).....	424		
ladas. (Zapus h.).....	324		258
lænatum. (Coendu).....	335	402	
lænatus. (Coendou).....	335	402	
Lagenocetus.....	21		
Lagenocetus latifrons.....	21		16
Lagenorhynchus.....	29		25
Lagenorhynchus acutus.....	29		25
Lagenorhynchus albirostris.....	29		26
Lagenorhynchus gubernator.....	29		25
Lagenorhynchus obliquidens.....	30		27
Lagenorhynchus perspicillatus.....	29		25
Lagenorhynchus thicola.....	29		26
Lagomys.....	337		266
Lagomys collaris.....	338		268
Lagomys minimus.....	339		269
Lagomys schisticeps.....	338		267
lagopus. (Canis).....			306
lagopus. (Vulpes).....	383		306
lagura. (Phoca).....	444		364
Lagurus.....	234, 247	299	178, 194
Laira.....	418	524	
lanceolatum. (Phyllostoma).....	515	669	
lapponicus. (Rangifer).....	37		
largha. (Phoca).....	442		363
lascivus. (Sciuropterus a.).....	123		†428
Lasionycteris.....	479		406
Lasionycteris noctivagans.....	480		407
lasiopyga. (Glossonycteris).....	520	682	
Lasiurus.....	485	591	411
Lasiurus borealis.....	485	592	411
Lasiurus b. mexicanus.....	486	594	
Lasiurus b. pfeifferi.....	486	593	
Lasiurus b. seminolus.....	486		413
Lasiurus b. teliotis.....	486	593	413
Lasiurus cinereus.....	487	595	413
Lasiurus funebris.....	485	593	412
Lasiurus intermedius.....	487	595	414
lasiurus. (Vespertilio).....	485	592	411

†Appendix.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
lasius. (Peromyscus).....	157		
Latax.....	435	537	354
Latax lutris.....	435	537	354
Latax l. nereis.....	436		
Lataxia.....	433	535	
Lataxina.....	433	535	
Lataxina mollis.....	433		352
lataxina. (Lutra c.).....	433		352
lateralis. (Citellus).....	105	147	83
lateralis. (Sciurus).....	105		83
lateralis. (Spermophilus).....	105		83
laticaudatus. (Dysopes).....	499	627	
laticeps. (Thomomys).....	274		224
laticinctus. (Lepus).....	346, 347		
latifrons. (Hyperoodon).....	21		16
latifrons. (Lagenocetus).....	21		16
latifrons. (Neotoma).....	212	282	
latifrons. (Orthogeomys).....	269	324	
latifrons. (Spilogale p.).....	414		331
latimanus. (Phenacomys).....	227		169
latimanus. (Scalops).....	468		391
latirostris. (Manatus).....	14	37	6
latirostris. (Orca).....	26	51	22
latirostris. (Perognathus).....	310		252
latirostris. (Thomomys).....	286		
Latra minima.....	1	2, 3	
latrans. (Arctomys).....	117		102
latrans. (Canis).....	376		301
leachii. (Monophyllus).....	516	672	
lecontei. (Reithrodontomys).....	191		148
lecontei. (Mus).....	191	256	
lecontei. (Plecotus).....	490		399
lecontei. (Rhithrodontomys).....	191		
lectus. (Eutamias).....	94		
lectus. (Tamias).....	94		
Leiobalæna.....	15		7
Lemmus.....	258		206
Lemmus alascensis.....	259		207
Lemmus minusculus.....	259		
Lemmus nigripes.....	260		209
Lemmus noveboracensis.....	234		179
Lemmus trimucronatus.....	258, 259		207
Lemmus t. helvolus.....	259		208
Lemmus t. yukonensis.....	259		208
Lemmus yukonensis.....	259		
lemmus. (Mus).....	258		206
Leo marinus.....	437		
leonina. (Phoca).....	437, 439	545	355, 359
Leopardus.....	365	443, 444	294
Leopardus hernandezii.....	365	446	

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
lepechini. (Phoca).....	441		362
lepida. (Neotoma).....	216		
lepida. (Neotoma c.).....	216		164
lepidus. (Nyctiellus).....	502	634	
lepidus. (Vespertilio).....	502	634	
Leporidae.....	339	411	269
leporina. (Aplodontia).....	126		112
leporina. (Phoca).....	441		362
leporinus. (Noctilio).....	494	617	
leporinus. (Sciurus).....	78	130	55
leporinus. (Vespertilio).....	494	617	
Leptonycteris.....	519	679	
Leptonycteris nivalis.....	519	679	
leptorhyncha. (Tatusia).....	13	34	4
lepturus. (Heteromys g.).....	318	373	
lepturus. (Peromyscus).....	167	204	
lepturus. (Vespertilio).....	492	610	
leptus. (Putorius s.).....	427		
Lepus.....	340, 358	413	269
Lepus alleni.....	353	433	288
Lepus a. palitans.....	353	434	
Lepus altimiræ.....	354	†745	
Lepus americanus.....	360		273
Lepus a. bairdi.....	361		274
Lepus a. columbiensis.....	361		275
Lepus a. dalli.....	361		275
Lepus a. macfarlani.....	362		275
Lepus a. macfarlanii.....	362		
Lepus a. phæonotus.....	360		273
Lepus a. struthopus.....	361		274
Lepus a. virginianus.....	360		273
Lepus a. washingtoni.....	361		274
Lepus aquaticus.....	340		277
Lepus a. attwateri.....	340	414	278
Lepus arcticus.....	358		270
Lepus a. bangsi.....	358		270
Lepus a. canus.....	359		
Lepus arizonæ.....	349, 350	423	285
Lepus a. confinis.....	350	424	
Lepus a. goldmani.....	349		
Lepus a. major.....	350	423	286
Lepus a. minor.....	350	423	286
Lepus a. pinetis.....	350		
Lepus artemesia.....		425	
Lepus asellus.....	355	435	
Lepus auduboni.....	345	422	283
Lepus a. sanctidiegi.....	346		
Lepus bachmani.....	352	429	
Lepus b. ubericolor.....	352		

†Appendix.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Lepus baileyi</i>	349		284
<i>Lepus bairdii</i>	361		274
<i>Lepus bennetti</i>	357	437	
<i>Lepus bishopi</i>	362		276
<i>Lepus borealis</i>	360		273
<i>Lepus b. gabbi</i>	351		
<i>Lepus californica</i>	357	437	
<i>Lepus californicus</i>	357	437	291
<i>Lepus c. xanti</i>	357	438	
<i>Lepus callotis</i>	353	431	288
<i>Lepus c. flavigularis</i>	†543		
<i>Lepus campestris</i>	360, 362		273, 277
<i>Lepus c. richardsoni</i>	363		
<i>Lepus c. sierræ</i>	363		
<i>Lepus c. townsendi</i>	363, 364		
<i>Lepus cerrosensis</i>	352	428	
<i>Lepus cinerascens</i>	352	429	287
<i>Lepus diazi</i>		412	
<i>Lepus douglasi</i>		415	
<i>Lepus durangæ</i>	345	424	
<i>Lepus edwardsi</i>	358		
<i>Lepus festinus</i>	354	†746	
<i>Lepus flavigularis</i>	353	431	288
<i>Lepus floridanus</i>	342		280
<i>Lepus f. alacer</i>	343		282
<i>Lepus f. aztecus</i>	344	418	
<i>Lepus f. bachmani</i>	352		281
<i>Lepus f. caniculus</i>	344		283
<i>Lepus f. chapmani</i>	343, 344		282
<i>Lepus f. chiapensis</i>	344	†744	
<i>Lepus f. connectens</i>	343	†743	
<i>Lepus f. holzneri</i>	345	419	
<i>Lepus f. mallurus</i>	342		280
<i>Lepus f. mearnsi</i>	342		280
<i>Lepus f. persultator</i>	345	418	
<i>Lepus f. rigidus</i>	345	419	
<i>Lepus f. sanctidiegi</i>	345		
<i>Lepus f. simplicianus</i>	343		
<i>Lepus f. subcinctus</i>	344	417	
<i>Lepus f. transitionalis</i>	342		281
<i>Lepus f. ubericolor</i>	352		282
<i>Lepus f. yucatanicus</i>	345	419	
<i>Lepus gabbi</i>	351	426	
<i>Lepus gaillardi</i>	354	433	
<i>Lepus g. battyi</i>	355	433	
<i>Lepus glacialis</i>	358		270
<i>Lepus goldmani</i>	349	†745	
<i>Lepus grangeri</i>	349, 350		285
<i>Lepus g. pinetis</i>	350		285

†Appendix.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Lepus graysoni</i>	351	426	
<i>Lepus grœnlandicus</i>	359		271.
<i>Lepus holzneri</i>	345		284
<i>Lepus hudsonius</i>	360		273
<i>Lepus idahoensis</i>	351		287
<i>Lepus incitatus</i>	351	428	
<i>Lepus insolitus</i>	348	421	
<i>Lepus insonus</i>	341	†743	
<i>Lepus insularis</i>	358	438	
<i>Lepus klamathensis</i>	362		276
<i>Lepus labradorius</i>	359		271
<i>Lepus laticinctus</i>	346, 347		
<i>Lepus l. perplicatus</i>	347		
<i>Lepus l. rufipes</i>	346		
<i>Lepus martirensis</i>	358	439	
<i>Lepus melanotis</i>	355		290
<i>Lepus merriami</i>	354	432	289
<i>Lepus m. altimiræ</i>	354		
<i>Lepus mexicanus</i>	353	431	288
<i>Lepus nanus</i>	360		273
<i>Lepus nigricaudatus</i>	353	431	288
<i>Lepus nuttalli</i>	348	425	284
<i>Lepus n. mallurus</i>	342		
<i>Lepus orizabæ</i>	351	425	
<i>Lepus othus</i>	359		272
<i>Lepus pacificus</i>	348	†744	
<i>Lepus paludicola</i>	341		279
<i>Lepus palustris</i>	341	415	279
<i>Lepus p. paludicola</i>	341		279
<i>Lepus parvulus</i>	348	421	
<i>Lepus peninsularis</i>	353	430	
<i>Lepus podromus</i>	360		272
<i>Lepus princeps</i>	338		267
<i>Lepus richardsoni</i>	363	437	
<i>Lepus russatus</i>	348	420	
<i>Lepus saliens</i>	362		276
<i>Lepus sanctidiegi</i>	345, 346	422	283
<i>Lepus simplicianus</i>	343		
<i>Lepus sylvaticus</i>	342		280
<i>Lepus s. alacer</i>	343		
<i>Lepus s. arizonæ</i>	349		
<i>Lepus s. aztecus</i>	344		
<i>Lepus s. bachmani</i>	343		
<i>Lepus s. floridanus</i>	342		
<i>Lepus s. grangeri</i>	349		
<i>Lepus s. holzneri</i>	345		
<i>Lepus s. mearnsi</i>	342		
<i>Lepus s. pinetis</i>	350		
<i>Lepus s. transitionalis</i>	342		

†Appendix.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Lepus telmalemonus</i>	341		278
<i>Lepus texensis</i>	355		290
<i>Lepus t. deserticola</i>	356, 357	437	291
<i>Lepus t. eremicus</i>	355, 357	435	291
<i>Lepus t. griseus</i>	356	436	
<i>Lepus t. micropus</i>	356	436	
<i>Lepus t. wallawalla</i>	356		
<i>Lepus texianus</i>	354, 355		
<i>Lepus t. deserticola</i>	356		
<i>Lepus t. eremicus</i>	355		
<i>Lepus t. griseus</i>	356		
<i>Lepus t. micropus</i>	356		
<i>Lepus t. wallawalla</i>	356		
<i>Lepus timidus</i>	340	413	269
<i>Lepus t. arcticus</i>	358		270
<i>Lepus townsendi</i>	363		
<i>Lepus trowbridgei</i>			281
<i>Lepus truei</i>	341		
<i>Lepus truii</i>	341	415	
<i>Lepus tschuktschorum</i>	359		271
<i>Lepus tularensis</i>	357		
<i>Lepus veræcrucis</i>	347	420	
<i>Lepus v. pacificus</i>	348		
<i>Lepus virginianus</i>	360		273
<i>Lepus washingtonii</i>	361		274
<i>lestes</i> . (<i>Canis</i>).....	377	465	301
<i>lesueuri</i> . (<i>Amphisorex</i>).....	446		366
<i>leucas</i> . (<i>Delphinapterus</i>).....	25		19
<i>leucas</i> . (<i>Delphinus</i>).....	25		19
<i>Leucocyon</i>	380	465	
<i>leucodon</i> . (<i>Neotoma</i>).....	212	281	
<i>leucodon</i> . (<i>Thomomys</i>).....	273		223
<i>leucogaster</i> . (<i>Hypudæus</i>).....	130		119
<i>leucogaster</i> . (<i>Onychomys</i>).....	130		119
<i>leucogaster</i> . (<i>Sciurus</i>).....	68	115	
<i>Leucomitra</i>	405, 410	507, 509	
<i>leucomus</i> . (<i>Arctibeus</i>).....	528	712	
<i>leuconotus</i> . (<i>Conepatus</i>).....	411	515	
<i>leucoparia</i> . (<i>Putorius f.</i>).....	432	535	
<i>leucoparia</i> . (<i>Spilogale</i>).....	413		329
<i>leucophæus</i> . (<i>Arvicola</i>).....	245		192
<i>leucophæus</i> . (<i>Microtus a.</i>).....	245		192
<i>leucopla</i> . (<i>Phoca</i>).....			360
<i>leucopleura</i> . (<i>Dysopes</i>).....	497	623	
<i>Leucopleurus</i>	29		
<i>Leucopleurus arcticus</i>	29		25
<i>leucopleurus</i> . (<i>Delphinus</i>).....	29		25
<i>leucops</i> . (<i>Macroxus</i>).....	69	117	
<i>leucops</i> . (<i>Sciurus</i>).....	68, 72, 74	119, 123	
<i>leucops</i> . (<i>Sciurus a.</i>).....	71	119	

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
leucopus. (Musculus)	140		
leucopus. (Mustela)	419	335	
leucopus. (Peromyscus)	140, 141		124, *492
leucotis. (Sciurus)	79		
leucotis. (Sciurus c.)	79		53
leucotis. (Sigmodon)	179	230, 231	
leucurus. (Cervus)	44		39
leucurus. (Citellus)	97	142, 143	
leucurus. (Cynomys)	118		104
leucurus. (Peromyscus)	165	200	
leucurus. (Spermophilus)	97	142	86.
leucurus. (Tamias)	97	142	86
levipes. (Perodipus m.)	298		
levipes. (Peromyscus)	164	195	
levipes. (Reithrodontomys)	203	270	
levipes. (Rhithrodontomys)	203	270, 271	
lewisi. (Cervus)	48		41
lewisi. (Cynomys)	118		104
lewisi. (Arctomys)	118		104
liardensis. (Ovis c.)	54		47
Lichonycteris	520	683	
Lichonycteris obscurus	520	683	
lichtensteini. (Dama)	45	70	
lichtensteini. (Odontocælus)	45	72	
lilium. (Phyllostoma)	529	715	
lilium. (Sturnira)	529	715	
limitis. (Sciurus l.)	65	110	53
Limnolagus	340	413	277
limosus. (Thomomys)	278		
lineata. (Emballonura)	492	609	
lineata. (Phocæna)	25	48	20
lineatum. (Phyllostoma)	525	702	
lineatus. (Vampyrops)	525	702	
Linx	365	443	293
Liomys	316, 319	368, 374	
Liomys canus	321	375	
Liomys crispus	320	380	
Liomys c. setosus	320		
Liomys heterothrix	320	380	
Liomys obscurus	319	379	
Liomys orbitalis	320	380	
Liomys parviceps	322	378	
Liomys phæura	319	379	
Liomys pictus	323		
Liomys p. isthmius	322	378	
Liomys p. rostratus	322	378	
Liomys plantinarenis	322	377	
Liomys sonorana	323	379	
Liomys texensis	323		

*Supplement.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Liomys torridus</i>	321	376	
<i>Liomys t. minor</i>	321	377	
<i>Liomys verærucis</i>	319	379	
<i>Lipura</i>	119		
<i>Lissodelphis</i>	32		29
<i>Lissodelphis borealis</i>	32		30
<i>littoralis</i> . (<i>Neotoma m.</i>)	209		
<i>littoralis</i> . (<i>Sigmodon h.</i>)	175		144
<i>littoralis</i> . (<i>Tamias t.</i>)	87		
<i>littoralis</i> . (<i>Urocyon c.</i>)	388, 389		†507
<i>littoralis</i> . (<i>Vulpes</i>)	388		
<i>littorea</i> . (<i>Phoca</i>)	442		363
<i>lobatus</i> . (<i>Cervus</i>)	41		38
<i>Lobostoma</i>	503	639	
<i>Lobostoma cinnamomeum</i>	507	649	
<i>Lobostoma quadridens</i>	504	641	
<i>Loncheres</i>	328	383	
<i>Loncheres labilis</i>	328	384	
<i>Loncherinæ</i>	328	382	
<i>Lonchetes</i>	328	383	
<i>Lonchorina</i>	507	649	
<i>Lonchorina aurita</i>	507, 508	650	
<i>longicauda</i> . (<i>Mustela</i>)	430		
<i>longicauda</i> . (<i>Reithrodon</i>)	195	261	151
<i>longicauda</i> . (<i>Sorex o.</i>)	450		
<i>longicaudata</i> . (<i>Talpa</i>)	467		
<i>longicaudatus</i> . (<i>Heteromys</i>)	317, 318	370, 372	
<i>longicaudatus</i> . (<i>Molossus</i>)	495	620	
<i>longicaudus</i> . (<i>Arvicola</i>)	243		191
<i>longicaudus</i> . (<i>Dasytus</i>)	13	33	4
<i>longicaudus</i> . (<i>Microtus</i>)	243, 244		191
<i>longicaudus</i> . (<i>Onychomys</i>)	132		121
<i>longicaudus</i> . (<i>Phenacomys</i>)	228		169
<i>longicaudus</i> . (<i>Putorius</i>)	430		348
<i>longicaudus</i> . (<i>Reithrodontomys</i>)	195, 196	261	151, 153
<i>longicaudus</i> . (<i>Rhithrodontomys</i>)	195, 196	261, 262	
<i>longicaudus</i> . (<i>Sorex o.</i>)	450, 451		372
<i>longicrus</i> . (<i>Myotis l.</i>)	479	581	402
<i>longicrus</i> . (<i>Vespertilio</i>)	479	581	402
<i>longidens</i> . (<i>Delphinus</i>)	30		27
<i>longimana</i> . (<i>Balæna</i>)	17		10
<i>longimana</i> . (<i>Megaptera</i>)		41	10
<i>longimanus</i> . (<i>Dysopes</i>)	497	623	
<i>longimembris</i> . (<i>Otognosis</i>)	306		247
<i>longimembris</i> . (<i>Perognathus</i>)	306		247
<i>longipes</i> . (<i>Dipodops</i>)	298		239
<i>longipes</i> . (<i>Onychomys</i>)	131	167	120
<i>longipes</i> . (<i>Onychomys l.</i>)	131		120
<i>longipes</i> . (<i>Perodipus</i>)	298		239

†Appendix.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
longipilis. (Arvicola r.)	235		180
longirostris. (Arvicola)	238		182
longirostris. (Delphinus)	33	58	
longirostris. (Prodelphinus)	33	58	
longirostris. (Sorex)	453		375
Lophostoma	572	658	
lophurus. (Peromyscus)	167	204	
loquax. (Sciurus h.)	80		62
lordi. (Abromys)	308		250
lordi. (Perognathus)	308		250
loringi. (Eutamias g.)	231		173
lotor. (Procyon)	400		317
lotor. (Ursus)	400		
louisianæ. (Odocoileus v.)	44		
louisianæ. (Odontocercus a.)	44		
lucasana. (Spilogale)	416	523	
luciæ. (Monophyllus)	518	678	
luciæ. (Moschophoromys)	190	251	
luciæ. (Oryzomys)	190	251	
luciæ. (Stenoderma)	527	706	
lucifugus. (Myotis)	478	580	402
lucifugus. (Vespertilio)	478		402
ludovicianus. (Arctomys)	117		102
ludovicianus. (Cynomys)	117		102
ludovicianus. (Microtus)	248		197
ludovicianus. (Sciurus)	64, 74	124	52
ludovicianus. (Sciurus n.)	64	110	
Lupus	374	464	
Lupus albus	375		
Lupus ater	376		
Lupus griseus	375		
Lupus sticticus	375		
lupus. (Canis)			299
luscus. (Gulo)	416		333
luscus. (Ursus)	416		
luteiventris. (Tamias q.)	92		76
lutensis. (Putorius)	425		340
luteolus. (Ursus)	395		314
lutescens. (Geomys)	263		218
lutescens. (Geomys b.)	263		298
luteus. (Gulo)	417		
luteus. (Peromyscus)	147		
Lutra	433	535	352
Lutra annectens	435	536	
Lutra canadensis	433		352
Lutra c. lataxina	433		352
Lutra c. pacifica	434		353
Lutra c. sonora	434	535	353
Lutra c. vaga	434		352

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc
<i>Lutra degener</i>	435		353
<i>Lutra destructor</i>			35 ²
<i>Lutra hudsonica</i>	433		35 ²
<i>Lutra h. pacifica</i>	434		
<i>Lutra h. sonora</i>	434		
<i>Lutra h. vaga</i>	434		
<i>Lutra marina</i>	435	537	354
<i>Lutra periclyzomæ</i>	434		
<i>Lutra sarcovienna</i>	1	3	
<i>Lutra stelleri</i>	436	537	354
<i>lutra</i> . (<i>Mustela</i>)	433	535	35 ²
<i>lutrecephala</i> . (<i>Mustela</i>)	423		
<i>lutrecephalus</i> . (<i>Putorius v.</i>)	423		339
<i>Lutreola</i>	423	530	338
<i>Lutreola v. ingens</i>	424		
<i>Lutreola v. lacustris</i>	424		
<i>lutreola</i> . (<i>Mustela</i>)	423		
<i>Lutrinæ</i>	433	535	351
<i>lutris</i> . (<i>Mustela</i>)	435	537	354
<i>Lyciscus</i>	374	465	
<i>Lyciscus cagottis</i>	378		
<i>lyelli</i> . (<i>Sorex t.</i>)	453		
<i>Lynceus</i>	365	443	
<i>Lynchailurus</i>	365	443	
<i>Lynchus</i>	365	443	293
<i>Lynx</i>	365, 370	443, 456	295
<i>Lynx baileyi</i>	372		
<i>Lynx canadensis</i>	370		
<i>Lynx c. mollipilosus</i>	370		295
<i>Lynx fasciatus</i>	371		297
<i>Lynx f. oculus</i>	371		297
<i>Lynx f. pallescens</i>	371		297
<i>Lynx floridanus</i>	372		
<i>Lynx gigas</i>	373		299
<i>Lynx r. californicus</i>	372	458	298
<i>Lynx r. eremicus</i>	372	458	298
<i>Lynx r. escuinapæ</i>	372		
<i>Lynx r. peninsularis</i>	373		
<i>Lynx subsolanus</i>	370		296
<i>Lynx texensis</i>	371		
<i>Lynx uinta</i>	373		
<i>Lysiurus</i>	12		
<i>lysteri</i> . (<i>Sciurus</i>)	95		
<i>lysteri</i> . (<i>Tamias s.</i>)	95		80
<i>macfarlani</i> . (<i>Lepus a.</i>)	362		275
<i>macfarlani</i> . (<i>Microtus</i>)	241		187
<i>macfarlanii</i> . (<i>Lepus a.</i>)	362		
<i>macfarlanii</i> . (<i>Microtus</i>)	241		

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
machetes. (Blarina m.)	465	561	
machetes. (Ursus)	396	481	
machlis. (Alces)	41		37
machrina. (Talpa)	470		389
machrinus. (Scalops a.)	470		389
machrinus. (Scalopus a.)	470		389
mackenzii. (Phenacomys)	228		
macleayi. (Chilonycteris)	504	641	
macleayii. (Chilonycteris)	503, 504		
Macleayius	15		8
macrocephalus. (Catodon)	21		
macrocephalus. (Physeter)	21	45	
Macrocolus	288		231
macrodon. (Fiber)	255		214
macrodon. (Sorex)	455	552	
Macrogeomys	270	326	
Macrogeomys cavator	270	329	
Macrogeomys cherrii	271	330	
Macrogeomys costaricensis	271	330	
Macrogeomys dolichocephalus	270	328	
Macrogeomys heterodus	270	328	
Macrogeomys pansa	271	329	
macropus. (Arvicola)	251		201
macropus. (Microtus r.)	251		201
macrorhabdotes. (Tamias)	88		71
Macrorhinus	439	545	359
Macrorhinus angustirostris	440		359
macrorhinus. (Peromyscus)	152		133
macrorhinus. (Sitomys)	152		133
macrorhynchus. (Globiocephalus)	27	53	23
macrospilotus. (Citellus s.)	99	145	
macrospilotus. (Spermophilus s.)	99	145	96
macrotarsus. (Didelphys)	2		
Macrotis	43	68	
macrotis. (Cervus)	48	76	42
macrotis. (Corynorhinus)	490		399
macrotis. (Neotoma)	206		
macrotis. (Neotoma f.)	206, 207	279	159
macrotis. (Nyctinomops)	499	627	
macrotis. (Nyctinomus)	499	627	
macrotis. (Onychomys)	134	169	
macrotis. (Plecotus)	490		399
macrotis. (Sciuropterus s.)	124		108
macrotis. (Vulpes)	384, 385	472	306
Macrotolagus	340, 353	413	269, 288
Macrotus	508	652	
Macrotus bocourtianus	510	654	
Macrotus bulleri	510	654	
Macrotus californicus	509	653	420
Macrotus mexicanus	510	653	

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
Macrotus minor.....	509		
Macrotus pygmaeus.....	511		
Macrotus waterhousii.....	508	652	
Macrotus w. compressus.....	509		
Macrotus w. jamaicensis.....	508		
macroura. (Condylura).....	467		395
macroura. (Felis).....	367	449	
macroura. (Mephitis).....	410		
macroura. (Sciurus).....	64		52
macroura. (Vulpes f.).....	380		305
macroura. (Vulpes p.).....	380		
macrourus. (Cervus).....	44		39
macrourus. (Odocoileus a.).....	44		39
macrourus. (Odontocælus a.).....	44		
macrourus. (Spermophilus).....	113	149	88
macrourus. (Vulpes).....	380		305
Macroxus.....	59	93	50
Macroxus adolphei.....	75	126	
Macroxus griseoflavus.....	74		
Macroxus leucops.....	69	117	
Macroxus maurus.....	69	116	
Macroxus melania.....	76		
Macroxus morio.....	68	116	
Macroxus neglectus.....	65		
macrura. (Mephitis).....	410	510	
macrurum. (Spectrellum).....	501	634	
macrurus. (Microtus).....	244		191
macrurus. (Sorex).....	448		369
maculata. (Euderma).....	490		398
maculata. (Felis).....	371, 372		296
maculata. (Felis r.).....	371		296
maculatum. (Euderma).....	490		
maculatus. (Felis r.).....			298
maculatus. (Histiotus).....	490		398
Madataeus.....	522	691	
madrensis. (Callospermophilus).....	106	147	
madrensis. (Citellus).....	106	147	
madrensis. (Neotoma m.).....	213		
madrensis. (Peromyscus).....	164	197	
magna. (Blarina).....	466	562	
magnicaudatus. (Sciurus).....	65		52
magruderensis. (Perognathus m.).....	307		249
magruderensis. (Perognathus p.).....	307		
major. (Aplodontia).....	127		112
major. (Chincha o.).....	409		
major. (Citellus s.).....	99		
major. (Delphinus).....	30	55	28
major. (Lepus a.).....	350	423	286
major. (Mephitis o.).....	409		
major. (Natalus).....	501	633	

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
major. (<i>Natalis</i> s.).....	501	633	
major. (<i>Neurotrichus</i> g.).....	467		387
major. (<i>Peromyscus</i>).....	155		137
major. (<i>Sigmodon</i> h.).....	177	226, 228	
major. (<i>Sitomys</i>).....	155		137
major. (<i>Spectrellum</i> s.).....	501		
major. (<i>Spermophilus</i> s.).....	99		96
major. (<i>Zapus</i>).....	326		260
makkovikensis. (<i>Sciuropterus</i> s.).....	125		
mallurus. (<i>Lepus</i> f.).....	342		280
mallurus. (<i>Lepus</i> n.).....	342		
Mammalia.....	I	I	I
managuensis. (<i>Sciurus</i>).....	77	129	
managuensis. (<i>Sciurus</i> b.).....	77	129	
Manatidæ.....			5
Manatus.....	14	36	5
Manatus americanus.....	14	37	6
Manatus australis.....	14	37	6
Manatus fluviatilis.....	14	37	6
Manatus gigas.....	14		5
Manatus latirostris.....	14	37	6
manatus. (<i>Trichechus</i>).....	14	35, 37	6
maniculatus. (<i>Hesperomys</i>).....	148		131
maniculatus. (<i>Peromyscus</i>).....	148		131
mapurito. (<i>Conepatus</i>).....	412	516	*510
margaritæ. (<i>Perognathus</i>).....	316	363	
Margay.....	365	443	
marginatus. (<i>Citellus</i> s.).....	100		
marginatus. (<i>Clymene</i>).....	31	55	28
marginatus. (<i>Delphinus</i>).....	34	58	30
marginatus. (<i>Spermophilus</i> s.).....	100		
Marikina.....	532	724	
marina. (<i>Lutra</i>).....	435	537	354
marinus. (<i>Leo</i>).....	437		
maritimus. (<i>Ursus</i>).....	390		310
Marmosa.....	I	4	
Marmosa canescens.....	2	6, 8	
Marmosa cinerea.....	3	7	
Marmosa fulviventer.....	3	8	
Marmosa insularis.....	3	7	
Marmosa mitis.....		8	
Marmosa murina.....	2	5, 6	
Marmosa m. mexicana.....	2	6	
Marmosa oaxacæ.....	3	8	
Marmosa sinaloæ.....	3	6	
Marmota.....	119		
Marmota dakota.....	120, 121		
Marmota engelhardti.....	121		
Marmota flaviventer.....	120, 121		

*Supplement.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
Marmota f. avarus.....	120		
Marmota monax.....	119		
Marmota m. canadensis.....	120		
Marmota m. ignavus.....	119		
Marmota olympus.....	121		
Marmota polonica.....	119		
Marmota pruinosa.....	120		
Marputius.....	410, 412	512, 517	
Marsipolæmus.....	482, 485	586, 590	410
Marsupialia.....	1	1	1
marsupialis. (Didelphis).....	5	14	
marsupialis. (Didelphys).....	7		
Martes a. abietinoides.....	420		
martes. (Mustela).....	419	334	
martinensis. (Neotoma).....	214		
martirensis. (Lepus).....	358	439	
martirensis. (Sitomys).....	159	189	
martirensis. (Spilogale a.).....	414	521	
martirensis. (Thomomys).....	282	336	
martirensis. (Thomomys f.).....	282		
mascotensis. (Sigmodon).....	176	227, 229	
mascotensis. (Sigmodon h.).....	176	227	
mastivus. (Noctilio l.).....	494	617	
mastivus. (Vespertilio).....	494	617	
maurus. (Macroxus).....	69	116	
maximus. (Perognathus h.).....	310		
maximus. (Phyllostomus).....	514	666	
mayensis. (Blarina).....	465	561	
maynardi. (Procyon).....	402	490	
Mazama.....	43, 50	68, 79	
Mazama h. peninsulæ.....	50	78	
Mazama pandora.....	51	80	
Mazama sartori.....	50	80	
Mazama tema.....		80	
mazama. (Evotomys).....	232		175
mazama. (Thomomys).....	275		225
mcmurtrii. (Centurio).....	530	717	
mearnsi. (Canis).....	377	468	302
mearnsi. (Conepatus m.).....	411	516	
mearnsi. (Felis p.).....	366	448	
mearnsi. (Lepus f.).....	342		280
mearnsi. (Lepus s.).....	342		
mearnsi. (Perognathus).....	302	353	243
mearnsi. (Peromyscus).....	150		132
mearnsi. (Sciurus d.).....	83	133	
mearnsi. (Sciurus h.).....	83		
mearnsii. (Vesperimus).....	150		132
mecisturus. (Peromyscus).....	171	212	
medioximus. (Synaptomys i.).....	256		†430

†Appendix.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
medius. (<i>Peromyscus t.</i>).....	146		130
megacephalus. (<i>Microdipodops</i>).....	299, 300		241
megacephalus. (<i>Sitomys</i>).....			127
megacephalus. (<i>Tachynices</i>).....	24		
megadon. (<i>Oryzomys r.</i>).....	188	243	
Megadontomys.....	171	212	
Megadontomys flavidus.....	172	214	
Megadontomys nelsoni.....	172	214	
Megadontomys thomasi.....	171	213	
Megalomys.....	189	250	
megalophylla. (<i>Mormops</i>).....	506	648	418
megalops. (<i>Peromyscus</i>).....	170	210	
megalotis. (<i>Hesperomys</i>).....	157		139
megalotis. (<i>Micronycteris</i>).....	513	662	
megalotis. (<i>Peromyscus</i>).....	157		139
megalotis. (<i>Phyllophora</i>).....	513	662	
megalotis. (<i>Potos f.</i>).....	404		
megalotis. (<i>Reithrodon</i>).....	193	259	151
megalotis. (<i>Reithrodontomys</i>).....	193		151
megalotis. (<i>Rhithrodontomys</i>).....	193	259	
Megaptera.....	17	40	9
Megaptera americana.....	17		10
Megaptera kuzira.....	16		9
Megaptera longimana.....		41	10
Megaptera nodosa.....	17		10
Megaptera n. bellicosa.....	18	41	
Megaptera n. osphyia.....	17		10
Megaptera versabilis.....	18	41	10
Megapteron.....	17	40	9
Megapteropsis.....	17	41	9
Megascapheus.....	272		
mekisturus. (<i>Peromyscus</i>).....	171	212	
melampeplus. (<i>Putorius</i>).....	425		
melampeplus. (<i>Putorius v.</i>).....	425		
melania. (<i>Macroxus</i>).....	76		
melania. (<i>Sciurus</i>).....	76	128	
melanocarpus. (<i>Peromyscus</i>).....	170	210	
melanochir. (<i>Ateles</i>).....	535	733	
melanonotus. (<i>Sciurus n.</i>).....	62	107	
melanophrys. (<i>Hesperomys</i>).....	166	201	
melanophrys. (<i>Onychomys</i>).....	131, 132	166	
melanophrys. (<i>Onychomys l.</i>).....	131		120
melanophrys. (<i>Peromyscus</i>).....	166	201	
melanops. (<i>Eptesicus</i>).....	483	587	410
melanops. (<i>Thomomys</i>).....	279		227
melanopterus. (<i>Eptesicus f.</i>).....	483		
melanopterus. (<i>Vespertilio f.</i>).....	483		
melanopus. (<i>Arctomys</i>).....	120		106
melanorhinus. (<i>Vespertilio</i>).....	476	578	
melanorhyncha. (<i>Mustela</i>).....	422		335

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
melanotis. (Lepus).....	355		290
melanotis. (Oryzomys).....	187	244, 243	
melanotis. (Perognathus a.).....	303	356	
melanotis. (Peromyscus).....	161	178	
melanotis. (Sigmodon).....	180	232	
melanura. (Neotoma i.).....	217, 218	284	
melanurus. (Capromys).....	330	391	
melanurus. (Dipodomys m.).....	291	348	
melanurus. (Tamias m.).....	94		79
melas. (Delphinus).....	27	52	
melas. (Globicephalus).....	27	52	
melas. (Globiocephalus).....	27	53	23
Meles jeffersoni.....	404		320
Meles t. americanus.....	404		
Melinæ.....	404	503	319
Memina.....	1		
memina. (Didelphis).....	1	3	
mephitica. (Mephitis).....	406		322
Mephitis.....	405	507	322
Mephitis a. hudsonica.....	406		
Mephitis avia.....	407		327
Mephitis bicolor.....	412		328
Mephitis concolor.....	410	512	
Mephitis elongata.....	406		324
Mephitis estor.....	407		326
Mephitis fetidissima.....	406		
Mephitis fœtulenta.....	408		326
Mephitis hudsonia.....	406		322
Mephitis interrupta.....	412		
Mephitis leuconota.....	411		
Mephitis macroura.....	410		
Mephitis macrura.....	410	510	
Mephitis m. milleri.....	410	511	
Mephitis m. vittata.....	410	512	
Mephitis mephitica.....	406		322
Mephitis m. elongata.....	406		
Mephitis m. hudsonia.....	406		
Mephitis m. scrutator.....	407		324
Mephitis mephitis.....	406		
Mephitis mesoleuca.....	411		
Mephitis mesomelas.....	407		325
Mephitis m. avia.....	407		
Mephitis m. estor.....	407	509	
Mephitis m. varians.....	407		
Mephitis milleri.....	410		325
Mephitis occidentalis.....	408, 409		324
Mephitis o. holzneri.....	409	507	
Mephitis o. major.....	409		
Mephitis o. notata.....	408		
Mephitis o. spissigrada.....	408		

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
Mephitis olida.....	406		
Mephitis platyrhina.....	409		
Mephitis putida.....	406		
Mephitis quaterlinearis.....	413		328
Mephitis spissigrada.....	408		326
Mephitis varians.....	407		322
Mephitis vittata.....	410		
Mephitis v. intermedia.....	410		
mephitis. (Viverra).....	406		
meridionalis. (Orca).....	27	51	23
Meriones.....	324		
Meriones acadicus.....	324		257
Meriones labradorius.....	328		263
Meriones microcephalus.....	324		257
merriami. (Cervus).....	42		
merriami. (Cratogeomys).....	265	313	
merriami. (Dipodomys).....	288	345	232
merriami. (Geomys).....	264		
merriami. (Lepus).....	354	432	289
merriami. (Perognathus).....	301	353	243
merriami. (Peromyscus).....	165	198	
merriami. (Reithrodontomys).....	192		150
merriami. (Rhithrodontomys).....	192		
merriami. (Sorex).....	454		376
merriami. (Tamias).....	88	136, 137	71
merriami. (Tamias a.).....	88		
merriami. (Ursus).....	391		
merriami. (Vesperugo).....	480	582	409
mesamericana. (Didelphis).....	6	15	
mesamericana. (Didelphys).....	6	15	
mesembrinus. (Perognathus).....	308		
mesoleuca. (Mephitis).....	411		
mesoleucus. (Conepatus).....	411	515	
mesomelas. (Mephitis).....	407		325
mesomelas. (Peromyscus t.).....	147	177	
Mesoodon.....	22		
Mesoplodon.....	23		17
Mesoplodon bidens.....	23		18, *480
Mesoplodon stejnegeri.....	23		14, *481
mesopolius. (Perognathus f.).....	315	367	
Metachirus.....	4	11	
Metachirus fuscogriseus.....	5	12, 13	
Metachirus f. pallidus.....	5	13	
Metachirus nudicaudatus.....	4, 5	12	
Metachirus opossum.....		12	
metallescens. (Scalops).....	468		391
metallicola. (Peromyscus).....	138	192	
Meteorus.....	482	586	410
metis. (Delphinus).....	31	56	29

*Supplement.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
mexicana. (Alouatta p.).....	533	727	
mexicana. (Antilocapra a.).....	51	82	
mexicana. (Atalapha).....	486	594	
mexicana. (Blarina).....	464	560	
mexicana. (Chilonycteris).....	506	644	
mexicana. (Chæronycteris).....	517	673	
mexicana. (Dasypsecta).....	336	405	
mexicana. (Felis).....			294
mexicana. (Hystrix).....	335	402	
mexicana. (Marmosa m.).....	2	6	
mexicana. (Neotoma).....	213	282	158
mexicana. (Tylostoma).....	514	665	
mexicanum. (Coendu).....	335	402	
mexicanum. (Spectrellum).....	502		
mexicanus. (Arvicola).....	245	301	
mexicanus. (Canis).....	375, 376	471	
mexicanus. (Centurio).....	530	717	
mexicanus. (Cervus).....	45	70, 72	40
mexicanus. (Citellus).....	103	146	
mexicanus. (Citillus).....	103	146	
mexicanus. (Cynomys).....	119	156	
mexicanus. (Hesperomys).....	168	207	
mexicanus. (Lasiurus b.).....	486	594	
mexicanus. (Lepus).....	353	431	288
mexicanus. (Macrotus).....	510	653	
mexicanus. (Micronycteris m.).....	513	662	
mexicanus. (Microtus).....	245	301	
mexicanus. (Myotis c.).....	477	579	
mexicanus. (Natalis).....	502	633	
mexicanus. (Natalus).....	502	633	
mexicanus. (Nyctinomus).....	500	629	417
mexicanus. (Ochetodon).....	198		
mexicanus. (Oryzomys).....	184	242, 246	
mexicanus. (Otopterus).....	510	653	
mexicanus. (Ovis).....	54		
mexicanus. (Ovis c.).....	54	86	
mexicanus. (Perognathus f.).....	303	355	
mexicanus. (Peromyscus).....	168	207	
mexicanus. (Reithrodon).....	197	264	
mexicanus. (Rhithrodontomys).....	197	264, 265	
mexicanus. (Sciurus).....	103	146	
mexicanus. (Spermophilus).....	99, 103	145, 146	96, 98
mexicanus. (Vespertilio).....	477	579	
michiganensis. (Mus).....	160		140
michiganensis. (Peromyscus).....	160		140
Micoureus.....	1	5	
microcephalus. (Arvicola).....	236		181
microcephalus. (Meriones).....	324		257
microcephalus. (Monodon).....	24		19
Microdipodops.....	299		240

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Microdipodops californicus</i>	300		
<i>Microdipodops megacephalus</i>	299, 300		241
<i>Microdipodops m. oregonus</i>	299		
<i>Microdipodops pallidus</i>	299		
<i>microdon</i> . (<i>Canis</i>).....	379	469	
<i>microdon</i> . (<i>Reithrodontomys</i>).....	201	269	
<i>microdon</i> . (<i>Rhithrodontomys</i>).....	201, 202	269	
<i>microdon</i> . (<i>Sigmodon h.</i>).....	178	226	
<i>microdon</i> . (<i>Stenoderma</i>).....	528		
<i>Microlagus</i>	340, 352	413, 428	269, 287
<i>Micronycteris</i>	512	660	
<i>Micronycteris hirsutus</i>	512	661	
<i>Micronycteris megalotis</i>	513	662	
<i>Micronycteris m. mexicanus</i>	513	662	
<i>Micronycteris microtis</i>	513	663	
<i>microps</i> . (<i>Delphinus</i>).....	31, 33	55, 58	28
<i>microps</i> . (<i>Perodipus</i>).....	297, 298		
<i>micropterus</i> . (<i>Delphinus</i>).....	23		18
<i>micropus</i> . (<i>Chilonatalis</i>).....	502	635	
<i>micropus</i> . (<i>Lepus t.</i>).....	356	436	
<i>micropus</i> . (<i>Natalus</i>).....	502	635	
<i>micropus</i> . (<i>Neotoma</i>).....	208	281	155
<i>Microsciurus</i>	59, 60	94, 98	
<i>Microsorex</i>	456		377
<i>Microsorex alnorum</i>	457		
<i>Microsorex eximius</i>	457		
<i>Microsorex hoyi</i>	457		377
<i>Microspilotus</i>	99		
<i>microtarsus</i> . (<i>Didelphys</i>).....	2		
<i>Microtinæ</i>	226	298	167
<i>microtis</i> . (<i>Micronycteris</i>).....	573	663	
<i>microtis</i> . (<i>Putorius</i>).....	426		
<i>Microtus</i>	233, 234	298	178, 179
<i>Microtus abbreviatus</i>	243		189
<i>Microtus a. fisheri</i>	243		190
<i>Microtus alleni</i>	252		203
<i>Microtus alticola</i>	244		192
<i>Microtus a. leucophæus</i>	245		192
<i>Microtus angusticeps</i>	244		192
<i>Microtus aphorodemus</i>	237		
<i>Microtus austerus</i>	248		196
<i>Microtus aztecus</i>	237		182
<i>Microtus bairdi</i>	250		200
<i>Microtus borealis</i>	247		194
<i>Microtus breweri</i>	238		182
<i>Microtus californicus</i>	240		186
<i>Microtus c. constrictus</i>	241		186
<i>Microtus c. hyperythrus</i>	240	302	
<i>Microtus c. vallicola</i>	240		186
<i>Microtus canicaudus</i>	239		185

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Microtus cautus</i>	244		191
<i>Microtus chrotorrhinus</i>	246		193
<i>Microtus c. ravus</i>	247		193
<i>Microtus curtatus</i>	247		195
<i>Microtus drummondi</i>	236, 237		181
<i>Microtus dutcheri</i>	239		185
<i>Microtus edax</i>	241		187
<i>Microtus enixus</i>	237		182
<i>Microtus fontigenus</i>	236		181
<i>Microtus fulviventris</i>	246	301	
<i>Microtus guatemalensis</i>	251	306	
<i>Microtus haydeni</i>	248		197
<i>Microtus innuitus</i>	242		189
<i>Microtus insularis</i>	238		182
<i>Microtus kadiacensis</i>	242		188
<i>Microtus longicaudus</i>	243, 244		191
<i>Microtus ludovicianus</i>	248		197
<i>Microtus macfarlandi</i>	241		187
<i>Microtus macfarlandii</i>	241		
<i>Microtus macrurus</i>	244		191
<i>Microtus mexicanus</i>	245	301	
<i>Microtus m. guadalupensis</i>	245		
<i>Microtus m. phæus</i>	245	302	
<i>Microtus minor</i>	248		197
<i>Microtus mogollonensis</i>	245, 246		192
<i>Microtus montanus</i>	238		182
<i>Microtus m. arizonensis</i>	238		183
<i>Microtus m. rivularis</i>	239		183
<i>Microtus mordax</i>	244		191
<i>Microtus morosus</i>	250		200
<i>Microtus nanus</i>	239		184
<i>Microtus n. canescens</i>	239		185
<i>Microtus nemoralis</i>	249		199
<i>Microtus nesophilus</i>	238		182
<i>Microtus nevadensis</i>	240		185
<i>Microtus n. rivularis</i>	239		
<i>Microtus operarius</i>	241		187
<i>Microtus oregoni</i>	250		200
<i>Microtus pallidus</i>	247		195
<i>Microtus pauperrimus</i>	247		195
<i>Microtus pennsylvanicus</i>	234, 235		179
<i>Microtus p. acadicus</i>	235		180
<i>Microtus p. fontigenus</i>	236		181
<i>Microtus p. labradorius</i>	236		181
<i>Microtus p. modestus</i>	236		180
<i>Microtus p. nigrans</i>	235		180
<i>Microtus p. shattucki</i>	235		
<i>Microtus pinetorum</i>	249		198
<i>Microtus p. auricularis</i>	249		198
<i>Microtus p. nemoralis</i>	249		199

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Microtus p. scalopoides</i>	249		198
<i>Microtus principalis</i>	251		
<i>Microtus pumilus</i>	227		168
<i>Microtus quasiater</i>	249, 250	303	
<i>Microtus richardsoni</i>	250		200
<i>Microtus r. arvicoloides</i>	251		202
<i>Microtus r. macropus</i>	251		201
<i>Microtus scirpensis</i>	241		187
<i>Microtus serpens</i>	250		200
<i>Microtus sitkensis</i>	242		189
<i>Microtus stonei</i>	236		181
<i>Microtus terrænovæ</i>	238		182
<i>Microtus tetramerus</i>	243		190
<i>Microtus townsendi</i>	243		190
<i>Microtus umbrosus</i>	251	305	
<i>Microtus unalascensis</i>	242		188
<i>Microtus u. popofensis</i>	242		188
<i>Microtus vellerosus</i>	244		191
<i>Microtus xanthognathus</i>	246		193
<i>Microtus yakutatensis</i>	242		188
<i>microtus. (Canis)</i>	384		
<i>micrura. (Blarina)</i>		559	
<i>Micurus</i>	234	299	178
<i>micurus. (Galemys)</i>	461		382
<i>Mictomys</i>	256		205
<i>Mictomys innuitus</i>	256		205
<i>Midas</i>	532	723	
<i>Midas geoffroyi</i>	532	724	
<i>middendorffi. (Ursus)</i>	390		310
<i>migratorius. (Sciurus)</i>	79		54
<i>milleri. (Mephitis)</i>	410		325
<i>milleri. (Mephitis m.)</i>	410	511	
<i>milleri. (Myotis)</i>	474	575	
<i>mimus. (Sciurus a.)</i>	66		
<i>minima. (Lutra)</i>	1	3	
<i>minimus. (Chironectes)</i>	1	3	
<i>minimus. (Lagomys)</i>	339		269
<i>minimus. (Ochotona)</i>	339		269
<i>minimus. (Sigmodon)</i>	178	230	146
<i>minimus. (Tamias)</i>	93		78
<i>minnesota. (Sciurus h.)</i>	81		62
<i>minnesota. (Peromyscus l.)</i>	141		
<i>minor. (Antrozous)</i>	492	607	
<i>minor. (Arvicola a.)</i>	248		197
<i>minor. (Centurio)</i>	530	717	
<i>minor. (Chæronycteris)</i>	517	674	
<i>minor. (Heteromys t.)</i>	321	377	
<i>minor. (Lepus a.)</i>	350	423	286
<i>minor. (Liomys t.)</i>	321	377	
<i>minor. (Macrotus)</i>	509		

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
minor. (<i>Microtus</i>).....	248		197
minor. (<i>Otopterus</i> w.).....	509		
minor. (<i>Rhinops</i>).....	515	669	
minor. (<i>Rorqualus</i>).....	18		11
minor. (<i>Zapus</i> p.).....	325		260
minusculus. (<i>Lemmus</i>).....	259		
minusculus. (<i>Scapanus</i> c.).....	469		†435
minutus. (<i>Nyctinomus</i>).....	501	630	
minx. (<i>Mustela</i>).....	423		
miradorensis. (<i>Scotophilus</i>).....	484	588	
miradorensis. (<i>Vespertilio</i> f.).....	484	588	
Mirounga.....	439	545	
Mirounga angustirostris.....	440	545	
miscix. (<i>Sorex</i> m.).....	454		376
miscix. (<i>Sorex</i> p.).....	454		
Misothermus.....	260		209
mississippiensis. (<i>Peromyscus</i> g.).....	143		127
missoulæ. (<i>Oreamnos</i> m.).....	52		
missoulæ. (<i>Oreamnus</i> m.).....	52		
missouriensis. (<i>Arctomys</i>).....	117		102
missouriensis. (<i>Mus</i>).....	130		119
mittelli. (<i>Dipodomys</i>).....	291	344	
mitis. (<i>Felis</i>).....	367	449	
mitis. (<i>Marmosa</i>).....		8	
mitrata. (<i>Phoca</i>).....	440		360
mobilensis. (<i>Geomys</i> t.).....	262		216
modesta. (<i>Arvicola</i>).....	236		180
modestus. (<i>Microtus</i> p.).....	236		180
mogollónensis. (<i>Arvicola</i>).....	246		192
mogollónensis. (<i>Microtus</i>).....	245, 246		192
mogollónensis. (<i>Sciurus</i> f.).....	84		67
mogollónensis. (<i>Sciurus</i> h.).....	84		67
mohavensis. (<i>Citellus</i> t.).....	102		
mohavensis. (<i>Neotoma</i>).....	207		
mohavensis. (<i>Neotoma</i> f.).....	207		
mohavensis. (<i>Nyctinomus</i>).....	500	629	418
mohavensis. (<i>Spermophilus</i>).....	102		98
molaris. (<i>Nasua</i> n.).....	402, 403	497	
molestus. (<i>Oryzomys</i>).....	183	240	
mollipilosa. (<i>Felis</i> c.).....	370		
mollipilosus. (<i>Felis</i> c.).....	370		295
mollipilosus. (<i>Lynx</i> c.).....	370		295
mollipilosus. (<i>Perognathus</i>).....	307		249
mollipilosus. (<i>Perognathus</i> m.).....	307		249
mollipilosus. (<i>Sciurus</i>).....	83		
mollipilosus. (<i>Sciurus</i> d.).....	83		65
mollis. (<i>Citellus</i>).....	112		
mollis. (<i>Lataxina</i>).....	433		352
mollis. (<i>Spermophilus</i>).....	112		92

†Appendix.

	Check-List of Mammals.	Mammals. of Middle America, etc.	Synopsis of Mammals, etc.
Molossidæ.....	495	618	
Molossinæ.....			421
Molossus.....	495	618	
Molossus acuticaudatus.....	495	620	
Molossus aztecus.....	495	620	
Molossus californicus.....	498		
Molossus coibensis.....	496		
Molossus ferox.....	497	623	
Molossus fuliginosus.....	495	620	417
Molossus fumarius.....	496	621	
Molossus longicaudatus.....	495	620	
Molossus nasutus.....	496		
Molossus nigricans.....	496	620	
Molossus obscurus.....	495, 496	620	
Molossus rufus.....	495	619	
Molossus r. obscurus.....	495	620	
Molossus tropidorhynchus.....	496	620	
Molossus ursinus.....	495	619	
Monachus.....	444	542	
Monachus tropicalis.....	445	543	
monachus. (Phoca).....	444		
monachus. (Vespertilio).....	485	593	412
Monax.....	117		102
monax. (Arctomys).....	119		105
monax. (Marmota).....	119		
monax. (Mus.).....	119		
mondini. (Balænoptera).....	19		
monoceras. (Monodon).....	24		19
monoceros. (Monodon).....	24		
monochroua. (Neotoma).....	206	278	
Monodon.....	24		18
Monodon microcephalus.....	24		19
Monodon monoceras.....	24		19
Monodon monoceros.....	24		
Monophyllus.....	518	675	
Monophyllus clinedaphus.....	518	678	
Monophyllus cubanus.....	519	678	
Monophyllus leachii.....	516	672	
Monophyllus luciae.....	518	678	
Monophyllus plethodon.....	518	677	
Monophyllus portoricensis.....	518	677	
Monophyllus redmani.....	518, 519	679	
montanus. (Arvicola).....	238		182
montanus. (Dipodomys).....	295		†431
montanus. (Microtus).....	238		183
montanus. (Oreamnus).....	52		44
montanus. (Ovis).....	52		44
montanus. (Perodipus).....	295		
montanus. (Rangifer).....	38		36

†Appendix.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
montanus. (Reithrodon).....	193		
montanus. (Reithrodontomys)	193		151
montanus. (Rhithrodontomys)	193		
montanus. (Zapus)	327		262
montanus. (Zapus t.)	327		262
montereyensis. (Sorex)	452		373
monteriensis. (Arctocephalus)	437		357
montezumæ. (Neotoma).....	218		
monticola. (Bassaridæ).....	398	487	
monticola. (Perognathus).....	306, 307		248
monticola. (Sorex v.)	448		371
monticola. (Thomomys)	275		225
monticolus. (Sorex).....	448		371
monticolus. (Thomomys)	275		
montipinoris. (Peromyscus).....	157		
montserratense. (Stenoderma).....	526	705	
moorii. (Delphinus).....	30	55	28
mordax. (Arvicola)	244		191
mordax. (Microtus)	244		191
morio. (Macroxus).....	68	116	
Mormoops	506	646	
Mormoops blainvillii	506	646, 649	
Mormoops m. senicula	507	648	
Mormopinæ	503	639	
Mormops	506	646	418
Mormops blainvillii	507	649	419
Mormops b. cinnamomea	507	649	
Mormops megalophylla	506	648	418
Mormops m. senicula	507	648	
morosus. (Microtus).....	250		200
mortivallis. (Dipodomys m.)	290		
morulus. (Sciurus v.)	62	105	
Morunga	440	545	359
moschatus. (Bos).....	55		
moschatus. (Delphinus)	31	55	28
moschatus. (Ovibos)	55		48
Moschomys	189, 252	250	
Moschophoromys	189	250	
Moschophoromys desmaresti	190	250, 251	
Moschophoromys luciae	190	251	
mulleri. (Phoca).....	444		364
mundus. (Putorius x.).....	431		350
mungo. (Herpestes).....	374	462	
mungo. (Viverra)	374		
muricus. (Putorius).....	427		*434
Muridæ.....	129	162	117
murina. (Didelphis).....	2	5	
murina. (Marmosa)	2	5	
Murinae.....	129	162	117

* Supplement.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
murinus. (Desmodus)	530	719	
murinus. (Didelphys)	2		
murinus. (Grymæomys)	2		
murinus. (Vespertilio)	473	570	400
Mus.	119, 129	162	104, 117
Mus a. americanus	140		124
Mus alexandrinus	129	164	118
Mus alpinus.	119		
Mus anomalus	316	368	
Mus arvalis	233	298	178
Mus bursarius	262		217
Mus californicus	137	203	137
Mus cinerea	220		
Mus citellus	96	139	81
Mus decumanus	129	164	117
Mus desmaresti	189, 190	250	
Mus empetra	120		
Mus floridana	205, 209	276	154, 157
Mus groenlandicus			209
Mus hudsonius	260		209
Mus leontii	191	256	
Mus lemmus	258		206
Mus michiganensis	160		140
Mus missouriensis	130		119
Mus monax	119		
Mus musculus	130	162	118
Mus m. jalapæ	130	163	
Mus norvegicus	129	164	†428
Mus paca	337	408	
Mus palustris	180	233	146
Mus pennsylvanica	234		179
Mus pilorides	190	250	
Mus rattus	129	163	117
Mus rutilus	229		170
Mus saccatus	262		217
Mus s. noveboracensis	135, 141	70	125
Mus tectorum	129	164	118
Mus tuza	261		215
Mus volans	125	158	109
musculoides. (Peromyscus)	161		
musculoides. (Peromyscus a.)	161	180	
musculus. (Balæna)	20		13
musculus. (Balænoptera)	20		13
musculus. (Didelphys)	2		
Musculus leucopus	140		
musculus. (Mus)	130	162	118
musculus. (Nyctinomus)	500	639	
musculus. (Peromyscus)	135, 136	175	
musculus. (Sitomys)	135	175	

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Mustela</i>	419		333
<i>Mustela affinis</i>	433		
<i>Mustela americana</i>	419		334
<i>Mustela a. abieticola</i>	419		
<i>Mustela a. abietinoides</i>	420		
<i>Mustela a. actiosa</i>	420, 421		
<i>Mustela a. brumalis</i>	420		
<i>Mustela a. kenaiensis</i>	421		
<i>Mustela americanus</i>	419		
<i>Mustela atrata</i>	422		335
<i>Mustela a. actiosa</i>	420		336
<i>Mustela a. brumalis</i>	420		336
<i>Mustela a. caurina</i>	422		335
<i>Mustela barbara</i>	418		
<i>Mustela boria</i>	421		
<i>Mustela brumalis</i>	420		336
<i>Mustela canadensis</i>	422, 423		338
<i>Mustela c. pacifica</i>	423		
<i>Mustela caurina</i>	422		335
<i>Mustela c. actiosa</i>			*510
<i>Mustela c. atrata</i>			*510
<i>Mustela c. brumalis</i>			*510
<i>Mustela c. origenes</i>	422		335
<i>Mustela cicognanii</i>	426		344
<i>Mustela erminea</i>	430		
<i>Mustela frenata</i>	431		350
<i>Mustela godmani</i>	422		335
<i>Mustela guianensis</i>	1	3	
<i>Mustela gulo</i>	416		
<i>Mustela huro</i>	419	335	
<i>Mustela kenaiensis</i>	421		
<i>Mustela leucopus</i>	419	335	
<i>Mustela longicauda</i>	430		348
<i>Mustela lutra</i>	433	535	352
<i>Mustela l. canadensis</i>	433		
<i>Mustela lutrecephala</i>	423		
<i>Mustela lutreola</i>	423		
<i>Mustela lutris</i>	435	537	354
<i>Mustela martes</i>	419		334
<i>Mustela melanorhyncha</i>	422		335
<i>Mustela minx</i>	423		
<i>Mustela nesophila</i>	420		
<i>Mustela nigra</i>	422		335
<i>Mustela paraguensis</i>	1	3	
<i>Mustela pennanti</i>	422		337
<i>Mustela p. pacifica</i>	423		337
<i>Mustela putorius</i>	423	529	
<i>Mustela richardsonii</i>	426		344
<i>Mustela vison</i>	423		

* Supplement.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Mustela vulgaris</i>	426		344
<i>Mustela vulpina</i>	419	334	
<i>Mustela winingus</i>	423		
<i>Mustela xanthogenys</i>	431		349
<i>Mustela zibellina</i>	419	335	
<i>Mustelidæ</i>	404	502	319
<i>Mustelinæ</i>	416		
<i>mustelinus</i> . (<i>Sciurus</i>).....	68	115	
<i>muswa</i> . (<i>Alces</i>).....	41		38
<i>mutabilis</i> . (<i>Sorex s.</i>).....	456	553	
<i>mutica</i> . (<i>Glossophaga</i>).....	516	671	
<i>mutica</i> . (<i>Vulpes</i>).....	385		
<i>muticus</i> . (<i>Vulpes</i>).....	385		
<i>Mycetes</i>	532	*726	
<i>Mycetes palliatus</i>	533	726	
<i>Mycetes villosus</i>	532	726	
<i>Mynomes</i>	233	298	178
<i>Mynomes pratensis</i>	234		179
<i>Myodes</i>	258		206
<i>Myodes albigularis</i>	259		207
<i>Myodes nigripes</i>	260		209
<i>myodes</i> . (<i>Peromyscus l.</i>).....	140		125
<i>Myogalinæ</i>	466		386
<i>myoides</i> . (<i>Cricetus</i>).....	140		125
<i>myoides</i> . (<i>Hesperomys</i>).....	144		128
<i>Myolemmus</i>	260		209
<i>Myoprocta</i>	335	403	
<i>myops</i> . (<i>Erethizon e.</i>).....	333		266
<i>myops</i> . (<i>Sorex t.</i>).....	453		
<i>myops</i> . (<i>Thomomys</i>).....	276		
<i>myosura</i> . (<i>Myrmecophaga</i>).....	10	27	
<i>myosurus</i> . (<i>Didelphis</i>).....	4		
<i>Myotis</i>	473	570	400
<i>Myotis californicus</i>	476	578	403
<i>Myotis c. caurinus</i>	477		404
<i>Myotis c. ciliolabrum</i>	477		404
<i>Myotis c. durangæ</i>	478	579	
<i>Myotis c. jaliscensis</i>	477	579	
<i>Myotis c. mexicanus</i>	477	579	
<i>Myotis c. pallidus</i>	477	579	405
<i>Myotis chiriquensis</i>	475	576	
<i>Myotis dominicensis</i>	475	576	
<i>Myotis evotis</i>	474	574	
<i>Myotis e. thysanode</i>	479		
<i>Myotis incautus</i>	473		
<i>Myotis lucifugus</i>	478	580	402
<i>Myotis l. alascensis</i>	479		402
<i>Myotis l. longicus</i>	479	581	402
<i>Myotis milleri</i>	474	575	
<i>Myotis nigricans</i>	475	575	

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
Myotis orinomus.....	476	577	
Myotis peninsularis.....	474	573	
Myotis s. obscurus.....			409
Myotis subulatus.....	478	580	405
Myotis s. keenii.....	478		405
Myotis thysanodes.....	479	572	
Myotis velifer.....	473	573	401
Myotis vivesi.....	474	574	
Myotis yumanensis.....	475	576	403
Myotis y. saturatus.....	475	577	403
myotis. (Vespertilio).....	473	570	
Myoxomys.....		214	
Myoxus chrysurus.....	328	383	
Myoxus drummondi.....	220, 221		
Myrmecolichnus.....	10	25	
Myrmecophaga.....	12	29	
Myrmecophaga crispus.....	10	27	
Myrmecophaga didactyla.....	10	24	
Myrmecophaga jubata.....	12	29	
Myrmecophaga myosura.....	10	27	
Myrmecophaga quadridactyla.....	10		
Myrmecophaga sellata.....	11	28	
Myrmecophaga tetradactyla.....	10	26, 27	
Myrmecophaga tridactyla.....	12	24, 29	
Myrmecophagidæ.....	10	24	
Myrmydon.....	10	25	
mysticetus. (Balæna).....	16	39	8
nana. (Brachyphylla).....	522	690	
nannodes. (Cervus).....	42		
Nannugo.....	480	581	408
nanus. (Arvicola).....	239		184
nanus. (Lepus).....	360		273
nanus. (Microtus).....	239		184
nanus. (Promops).....	498	624	
nanus. (Sorex t.).....	453		375
nanus. (Tagassu).....	35	62	
nanus. (Tayassu).....	35	62	
Napæozapus.....	324, 328		257, 263
narica. (Viverra).....	402		
Narwalus.....	24		18
Narwalus andersoni.....	24		19
Narwalus vulgaris.....	24		19
nasica. (Nasua).....	402	497	
nasicus. (Thomomys).....	276		226
naso. (Dysopes).....			417
naso. (Rhynchonycteris).....	492	609	
naso. (Vespertilio).....	492		
Nasua.....	402	494	
Nasua n. bullata.....	403		

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Nasua n. molaris</i>	402		
<i>Nasua n. pallida</i>	403		
<i>Nasua n. panamensis</i>	403		
<i>Nasua n. yucatanica</i>	403		
<i>Nasua nasica</i>	402	497	
<i>Nasua n. bullata</i>	403	497	
<i>Nasua n. molaris</i>	402, 403	497	
<i>Nasua n. pallida</i>	403	498	
<i>Nasua n. panamensis</i>	403	498	
<i>Nasua n. yucatanica</i>	403	498	
<i>Nasua nelsoni</i>	402	495	
<i>Nasua thersites</i>		486	
<i>nasua. (Viverra)</i>	402	494	
<i>nasuta. (Arvicola)</i>	235		180
<i>nasutus. (Molossus)</i>	496		
<i>nasutus. (Peromyscus t.)</i>	159		139
<i>nasutus. (Promops)</i>	496, 497	621	
<i>nasutus. (Vesperimus)</i>	159		139
<i>Natalis</i>	501	631	
<i>Natalis mexicanus</i>	502		
<i>Natalis stramineus</i>	501	632	
<i>Natalis s. major</i>	501	633	
<i>Natalus</i>	501	631	
<i>Natalus brevipennis</i>	502		
<i>Natalus major</i>	501	633	
<i>Natalus mexicanus</i>	502		
<i>Natalus micropus</i>	502	635	
<i>Natalus stramineus</i>	501	632	
<i>natator. (Oryzomys)</i>	181		
<i>natator. (Oryzomys p.)</i>	181		147
<i>naurica. (Phoca)</i>	441		362
<i>nautica. (Phoca)</i>	441		362
<i>navigator. (Neosorex)</i>	457, 458		378, 379
<i>navigator. (Neosorex p.)</i>	458		379
<i>navus. (Neotoma)</i>	213	282	
<i>navus. (Thomomys l.)</i>	273		
<i>nayaritensis. (Sciurus)</i>	63	108	
<i>nebrascensis. (Canis)</i>	377		301
<i>nebrascensis. (Hesperomys l.)</i>	146		
<i>nebrascensis. (Peromyscus t.)</i>	146		130
<i>nebrascensis. (Reithrodontomys d.)</i>	192		150
<i>nebrascensis. (Rhithrodontomys d.)</i>	192		
<i>nebulicola. (Citellus)</i>	109		
<i>necator. (Vulpes)</i>	381		†433
<i>neglecta. (Taxidea a.)</i>	405		321
<i>neglecta. (Taxidea t.)</i>	405		
<i>neglectus. (Citellus)</i>	103		
<i>neglectus. (Macroxus)</i>	65		
<i>neglectus. (Platygeomys)</i>	268	319	
<i>neglectus. (Sciurus l.)</i>	65		†426

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
neglectus. (Sciurus r.)	65		
neglectus. (Spermophilus)	103		98
neglectus. (Tamias q.)	92		78
negligens. (Sciurus)	61	102	
nelsoni. (Blarina)	465	561	
nelsoni. (Citellus)	98		
nelsoni. (Dicrostonyx)	260		210
nelsoni. (Dicrostonyx h.)	261		210
nelsoni. (Geomys)	268	320	
nelsoni. (Heteromys)	323	381	
nelsoni. (Megadontomys)	172	214	
nelsoni. (Nasua)	402	495	
nelsoni. (Neotoma)	212		
nelsoni. (Odocoileus)	47	75	
nelsoni. (Odontocelus)	47	75	
nelsoni. (Orthogeomys)	269	324	
nelsoni. (Oryzomys)	182	235	
nelsoni. (Ovis)	54		
nelsoni. (Ovis c.)	54	84, 85	46
nelsoni. (Perognathus)	313	364	
nelsoni. (Peromyscus)	172	214	
nelsoni. (Romerolagus)	339	412	
nelsoni. (Sciurus)	72	120	
nelsoni. (Spermophilus)	98		87
nelsoni. (Thomomys)	286	339	
nelsoni. (Xenomys)	224	292	
nelsoni. (Xylomys)	323		
Nelsonia	224	289	
Nelsonia goldmani	224	290	
Nelsonia neotomodon	224	290	
memoralis. (Cervus)	46	74	
memoralis. (Microtus)	249		199
memoralis. (Microtus p.)	249		199
memoralis. (Odontocelus)	46	73, 74	
memoralis. (Sciurus a.)	70	118	
memoralis. (Sciurus p.)	70	118	
memoralis. (Sciurus w.)	70	118	
Neocyon		465	
Neodon	234	299	178
Neofiber	234, 252		202
Neofiber alleni	252		203
Neogale	423	530	338
neomexicana. (Vulpes m.)	384		
neomexicanus. (Putorius f.)	432	534	351
neomexicanus. (Sciurus f.)	84		
neomexicanus. (Vulpes m.)	384		
Neomys panamensis	174	219	
Neophoca	437		356
Neosciurus	59, 78	93	50, 53
Neosorex	457		378

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Neosorex albibarbis</i>	458		379
<i>Neosorex hydrodromus</i>	458		379
<i>Neosorex navigator</i>	457, 458		378, 379
<i>Neosorex palustris</i>	457		378
<i>Neosorex p. alaskanus</i>	458		379
<i>Neosorex p. navigator</i>	458		379
<i>Neotoma</i>	205	276	154
<i>Neotoma albigula</i>	217		
<i>Neotoma alleni</i>	226	297	
<i>Neotoma anthonyi</i>	214	283	
<i>Neotoma arenacea</i>	214	283	
<i>Neotoma arizonæ</i>	223		165
<i>Neotoma attwateri</i>	210		
<i>Neotoma baileyi</i>	210		156
<i>Neotoma bella</i>	218		†429
<i>Neotoma b. felipensis</i>	218	286	
<i>Neotoma bryanti</i>	208	280	
<i>Neotoma californica</i>	216		
<i>Neotoma campestris</i>	218		156
<i>Neotoma c. baileyi</i>	210		
<i>Neotoma c. rubida</i>	210		156
<i>Neotoma cinerea</i>	220		163
<i>Neotoma c. columbiana</i>	221		165
<i>Neotoma c. drummondi</i>	221		164
<i>Neotoma c. fusca</i>	222		165
<i>Neotoma c. lepida</i>	216		164
<i>Neotoma c. occidentalis</i>	221		164
<i>Neotoma cinnamomea</i>	223		166
<i>Neotoma cnemophila</i>	207		
<i>Neotoma cumulator</i>	205	280	154
<i>Neotoma desertorum</i>	215		161
<i>Neotoma d. grandis</i>	215		
<i>Neotoma d. sola</i>	215		161
<i>Neotoma distincta</i>	207	278	
<i>Neotoma fallax</i>	216		161
<i>Neotoma ferruginea</i>	210, 211	280	
<i>Neotoma f. ochracea</i>	211		
<i>Neotoma f. solitaria</i>	211		
<i>Neotoma floridana</i>	209		157
<i>Neotoma f. attwateri</i>	210		157
<i>Neotoma f. rubida</i>	210		
<i>Neotoma fulviventer</i>	220	288	
<i>Neotoma f. apicalis</i>	222		
<i>Neotoma fuscipes</i>	206	279, 278	158
<i>Neotoma f. affinis</i>	207		160
<i>Neotoma f. annectens</i>	206		160
<i>Neotoma f. cnemophila</i>	207		
<i>Neotoma f. dispar</i>	206		160
<i>Neotoma f. macrotis</i>	206, 207	279	159

†Appendix.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Neotoma f. mohavensis</i>	207		
<i>Neotoma f. streatori</i>	206		159
<i>Neotoma goldmani</i>	220	288	
<i>Neotoma grandis</i>	215		
<i>Neotoma grangeri</i>	223		165
<i>Neotoma intermedia</i>	216	284	161
<i>Neotoma i. albigula</i>	217	285	162
<i>Neotoma i. angusticeps</i>	217	285	162
<i>Neotoma i. durangæ</i>	217	285	
<i>Neotoma i. gilva</i>	217		162
<i>Neotoma i. melanura</i>	217, 218	284	
<i>Neotoma isthmica</i>	219	287	
<i>Neotoma latifrons</i>	212	282	
<i>Neotoma lepida</i>	216		164
<i>Neotoma leucodon</i>	212	281	
<i>Neotoma l. zacatecæ</i>	212		
<i>Neotoma macrotis</i>	206		
<i>Neotoma m. simplex</i>	206		
<i>Neotoma martinensis</i>	214		
<i>Neotoma mexicana</i>	213	282	158
<i>Neotoma m. bullata</i>	213		158
<i>Neotoma m. madrensis</i>	213		
<i>Neotoma micropus</i>	208	281	155
<i>Neotoma m. canescens</i>	208		
<i>Neotoma m. littoralis</i>	209		
<i>Neotoma m. planiceps</i>	209		
<i>Neotoma m. surberi</i>	209		156
<i>Neotoma mohavensis</i>	207		
<i>Neotoma monochroura</i>	206	278	
<i>Neotoma montezumæ</i>	218		
<i>Neotoma navus</i>	213	282	
<i>Neotoma nelsoni</i>	212		
<i>Neotoma nudicauda</i>	214		
<i>Neotoma occidentalis</i>	221		
<i>Neotoma o. fusca</i>	222		
<i>Neotoma orizabæ</i>	219	286	
<i>Neotoma orolestes</i>	223		166
<i>Neotoma palatina</i>	205		
<i>Neotoma parvidens</i>	219	288	
<i>Neotoma pennsylvanica</i>	210		157
<i>Neotoma picta</i>	219	287	
<i>Neotoma pinetorum</i>	211		158
<i>Neotoma rupicola</i>	223		166
<i>Neotoma saxamans</i>	221		†429
<i>Neotoma sinalcæ</i>	213	283	
<i>Neotoma splendens</i>	206	278	
<i>Neotoma stephensi</i>	216		
<i>Neotoma tenuicauda</i>	219	286	
<i>Neotoma torquata</i>	208	279	
<i>Neotoma tropicalis</i>	220	288	

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
Neotominae		276	
Neotomodon	225	292	
Neotomodon alstoni.....	225	294	
Neotomodon orizabæ.....	225	294	
Neotomodon perotensis.....	225	293	
neotomodon. (Nelsonia).....	224	290	
nereis. (Latax l.).....	436		
nerterus. (Reithrodontomys c.).....	200	267	
nerterus. (Rhithrodontomys c.).....	200	267	
nesioticus. (Citellus).....	115		
nesophila. (Mustela).....	420		
nesophilus. (Microtus).....	238		182
Neurotrichus	466		386
Neurotrichus gibbsi.....	466		387
Neurotrichus g. hyacinthinus.....	467		388
Neurotrichus g. major	467		387
nevadensis. (Dipodomys m.).....	288		232
nevadensis. (Microtus).....	240		185
nevadensis. (Nyctinomus m.).....	499	627	417
nevadensis. (Perognathus).....	305		246
nevadensis. (Sorex).....	449		371
nevadensis. (Thomomys).....	272		223
nevadensis. (Zapus).....	326		261
nicholli. (Stenoderma).....	526	706	
Nicon	516	670	
Nicon caudifer.....	516	672	
nicoyana. (Sciurus).....	76	127	
nigellus. (Peromyscus h.).....	155		137
nigellus. (Sitomys h.).....	155		137
niger. (Sciurus).....	66, 68, 79	115	51, 53
niger. (Thomomys).....	276		
nigra. (Mustela).....	422		335
nigra. (Phoca).....			357
nigra. (Phoca g.).....	444		364
nigra. (Phyllophora).....	516	672	
nigrans. (Microtus p.).....	235		180
nigrescens. (Blarina).....	463	558	
nigrescens. (Erethizon e.).....	333		
nigrescens. (Heteromys s.).....	317	370	
nigrescens. (Peromyscus m.).....	136	176	
nigrescens. (Putorius).....	423		338
nigricans. (Molossus).....	496	620	
nigricans. (Myotis).....	475	575	
nigricans. (Thomomys f.).....	280	335	228
nigricans. (Vespertilio).....	475, 476	575	404
nigricaudatus. (Lepus).....	353	431	288
nigriculus. (Peromyscus g.).....	143		127
nigripes. (Lemmus).....	260		209
nigripes. (Myodes).....	260		209

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
nigripes. (Putorius)	425		341
nigripes. (Sciurus f.)	78		
nigripes. (Sciurus g.)	78		56
niphæcus. (Ovibos m.)	56		
nitellinus. (Nyctomys)	173	216	
nitidus. (Vespertilio)	476	578	403
nitratoides. (Dipodomys m.)	288		233
nitratius. (Dipodomys m.)	288		233
nivalis. (Ischnoglossa)	519	679	
nivalis. (Leptonycteris)	519	679	
nivarius. (Eutamias)	233		177
nivea. (Procyon)	400		
niveiventris. (Hesperomys)	138		141
niveiventris. (Peromyscus)	138		141
Noctifelis	365	443	
Noctilio	494	617	
Noctilio dorsatus	494	617	
Noctilio leporinus	494	617	
Noctilio l. mastivus	494	617	
Noctilionidæ	492	608	421
Noctilioninæ	494	617	
noctivagans. (Lasionycteris)	480		407
noctivagans. (Vespertilio)	479, 480		407
Noctula	482	585	410
nodosa. (Balæna)	17		10
nodosa. (Megaptera)	17		10
normalis. (Clymene)	34		32
norvegicus. (Mus)	129	164	†428
Notagogus	1	5	
notata. (Chincha o.)	408		
notata. (Mephitis o.)	408		
notinus. (Bassariscus s.)	399	485	
Notiosorex	459	554	385
Notiosorex crawfordi	459	554	385
Notiosorex c. evotis	460	555	
Notiosorex gigas	460	555	
notius. (Putorius n.)	429		346
Notophorus	35	62	
novæ-hispaniæ. (Hystrix)	335	402	
novæ-zelandiæ. (Clymenia)	34	59	30
novæ-zelandiæ. (Delphinus)	30	55	28
noveboracensis. (Lemmus)	234		179
noveboracensis. (Mus s.)	135, 141	170	125
noveboracensis. (Peromyscus a.)	140		125
noveboracensis. (Peromyscus l.)	140		
noveboracensis. (Putorius)	429		346
noveboracensis. (Vespertilio)	485	592	411
noveboracensis. (Vespertilio)	485	593	411
novemcinctum. (Tatu)	13	33	

†Appendix.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
novemcinctus. (Cabassous).....	13	34	4
novemcinctus. (Dasypus).....	13	32, 33	4
nubilus. (Canis).....	376		300
nubiterræ. (Peromyscus c.).....	145		129
nuchalis. (Sciurus c.).....	73	122	
nudicauda. (Neotoma).....	214		
nudicaudata. (Didelphys).....	4	11	
nudicaudatus. (Metachirus).....	4, 5	12	
nudicaudus. (Hesperomys).....	173	217	
nudicaudus. (Tylomys).....	173	217	
nudipes. (Hesperomys).....	165	198	
nudipes. (Peromyscus).....	165	198	
nuttalli. (Arvicola).....	160		140
nuttalli. (Lepus).....	348	425	284
nuttalli. (Peromyscus).....	160		140
Nycticea crepuscularis.....	488	598	415
Nycticea cynocephala.....	500		417
Nycticeius.....	488	597	
Nycticeius humeralis.....	488	598	
Nycticeius h. cubanus.....	488	599	
Nycticejus.....			415
Nycticejus humeralis.....	488	598	415
Nycticeyx.....			415
Nyctiellus.....	502	634	
Nyctiellus lepidus.....	502	634	
Nyctinomops.....	498	625	
Nyctinomops depressus.....	499	627	
Nyctinomops femorosaccus.....	498	626	
Nyctinomops gracilis.....	498	626	
Nyctinomops macrotis.....	499	627	
Nyctinomops yucatanicus.....	498	626	
Nyctinomus.....	499	628	416
Nyctinomus ægyptiacus.....	499	628	
Nyctinomus antillarum.....	499	629	
Nyctinomus bahamensis.....	500	630	
Nyctinomus brasiliensis.....	500, 501	649	417
Nyctinomus b. californicus.....	500		
Nyctinomus cynocephalus.....	500		
Nyctinomus depressus.....	499	627	
Nyctinomus femorosaccus.....	498	626	417
Nyctinomus macrotis.....	499	627	
Nyctinomus m. nevadensis.....	499	627	417
Nyctinomus mexicanus.....	500	629	417
Nyctinomus minutus.....	500	630	
Nyctinomus mohavensis.....	500	629	418
Nyctinomus musculus.....	500	630	
Nyctinomus orthotis.....	497	623	
Nyctipithecus.....	533	728	
Nyctipithecus rufipes.....	533	729	
Nyctipithecus vociferans.....	534	729	

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
Nyctiplanus.....	529	714	
Nyctiplanus rotundatus.....	529	715	
Nyctomys.....	172	214	
Nyctomys decolorus.....	172	216	
Nyctomys nitellinus.....	173	216	
Nyctomys sumichrasti.....	172	216	
Nystactes.....	473	570	400
oaxacæ. (Marmosa).....	3	8	
oaxacensis. (Peromyscus).....	168	206	
obesus. (Odontobænus).....	439		
obesus. (Trichechus).....	439		358
obliquidens. (Lagenorhynchus).....	30		27
obscura. (Blarina).....	464	559	
obscurum. (Phyllostoma).....	524	695	
obscurus. (Evotomys).....	232		176
obscurus. (Fiber).....	254		214
obscurus. (Heteromys).....	319	379	
obscurus. (Lichonycteris).....	520	683	
obscurus. (Liomys).....	319	379	
obscurus. (Molossus).....	495, 496	620	
obscurus. (Molossus r.).....	495	620	
obscurus. (Myotis s.).....			409
obscurus. (Perodipus).....	299	351	
obscurus. (Perognathus).....	311, 313		253
obscurus. (Pipistrellus s.).....	481		409
obscurus. (Reithrodontomys m.).....	194	261	
obscurus. (Rhithrodontomys m.).....	194	261	
obscurus. (Sorex).....	449		371
obscurus. (Tamias).....	88	136	
obscurus. (Vespertilio).....	476	578	403
obsidianus. (Citellus s.).....	100		
obsidianus. (Spermophilus s.).....	100		97
obsoletus. (Citellus).....	99		
obsoletus. (Spermophilus).....	99		95
occidentalis. (Arvicola).....	243		190
occidentalis. (Canis).....	374, 376		300
occidentalis. (Canis l.).....	376		
occidentalis. (Cervus).....	42		
occidentalis. (Cervus c.).....	42		34
occidentalis. (Evotomys).....	233		176
occidentalis. (Mephitis).....	408, 409		324
occidentalis. (Neotoma).....	221		
occidentalis. (Neotoma c.).....	221		164
occidentalis. (Sciurus).....	65		52
occidentalis. (Teonoma c.).....	221		
occipitalis. (Fiber).....	254		
occisor. (Putorius).....	429		347
occisor. (Putorius n.).....	429		347
oceanica. (Phoca).....	444		364

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
Ochetodon	191	256	148
Ochetodon humilis	192		151
Ochetodon mexicanus	198		
Ochetomys	234	299	178
Ochotona	337		266
Ochotona cinnamomea	338		
Ochotona collaris	338		268
Ochotona cuppes	339		268
Ochotona minimus	339		269
Ochotona princeps	338		267
Ochotona saxatilis	339		268
Ochotona schisticeps	338		267
Ochotonidæ	337		266
ochracea. (<i>Neotoma</i> f.)	211		
ochraceus. (<i>Evotomys</i> g.)	230		172
ochrogaster. (<i>Hypudæus</i>)	248		196
ochrogenys. (<i>Eutamias</i> t.)	87		
ochrogenys. (<i>Tamias</i> t.)	87		71
ochrognathus. (<i>Sigmodon</i>)	179	230	
ochropus. (<i>Canis</i>)	379	469	303
ochrus. (<i>Perognathus</i> c.)	315		
ocius. (<i>Thomomys</i> c.)	284		
octo-cinctus. (<i>Dasypus</i>)	13	33	4
Octodontidæ	328	382	
oculatus. (<i>Sciurus</i>)	62	107	
oculea. (<i>Felis</i> r.)	371		297
oculeus. (<i>Lynx</i> f.)	371		297
ocythous. (<i>Urocyon</i> c.)	388		309
Odobenus	439		
Odocoileus v. louisianæ	44		
Odocoileus	43	68	38
Odocoileus americanus	43, 44		39
Odocoileus a. borealis	43		
Odocoileus a. couesi	45	70	40
Odocoileus a. macroirus	44		39
Odocoileus a. osceola	44		40
Odocoileus a. texensis	45	70	40
Odocoileus battyi	45	71	
Odocoileus cerrosensis	47	76	
Odocoileus clavatus	46	73	
Odocoileus columbianus	48		41
Odocoileus c. scaphiotus	48		42
Odocoileus c. sitkensis	48		42
Odocoileus costaricensis	46	73	
Odocoileus crooki	47		41
Odocoileus hemionus	48	76	42
Odocoileus h. californicus	49	77	43
Odocoileus h. canus	50	78	
Odocoileus nelsoni	47	75	
Odocoileus sinaloæ	47	78	

	Check-List of Mammals.	Mammals. of Middle America, etc.	Synopsis of Mammals, etc.
<i>Odocoileus speleus</i>	43	68	38
<i>Odocoileus thomasi</i>	47	75	
<i>Odocoileus truei</i>	46	73	
<i>Odontobænida</i>	439		
<i>Odontobænus</i>	439		
<i>Odontobænus obesus</i>	439		
<i>Odontobænus rosmarus</i>	439		
<i>Odontocælus</i>	43	68	
<i>Odontocælus americanus</i>	43	70, 75	
<i>Odontocælus a. borealis</i>	43		
<i>Odontocælus a. couesi</i>	45	70, 71	
<i>Odontocælus a. louisianæ</i>	44		
<i>Odontocælus a. macrourus</i>	44		
<i>Odontocælus a. osceola</i>	44		
<i>Odontocælus a. texensis</i>	45	70	
<i>Odontocælus battyi</i>	45	71	
<i>Odontocælus cerrosensis</i>	47	76	
<i>Odontocælus columbianus</i>	48		
<i>Odontocælus c. scaphiotus</i>	48		
<i>Odontocælus c. sitkensis</i>	48		
<i>Odontocælus costaricensis</i>	46	73	
<i>Odontocælus crooki</i>	47		
<i>Odontocælus hemionus</i>	48, 49	76, 77, 78	
<i>Odontocælus h. californicus</i>	49	76, 77, 78	
<i>Odontocælus h. canus</i>	50	78	
<i>Odontocælus h. eremicus</i>	49, 50	77	
<i>Odontocælus h. peninsulæ</i>	50	78	
<i>Odontocælus h. virgultus</i>	49		
<i>Odontocælus lichtensteini</i>	45	72	
<i>Odontocælus nelsoni</i>	47	75	
<i>Odontocælus nemoralis</i>	46	73, 74	
<i>Odontocælus rothschildi</i>	46	72	
<i>Odontocælus sinaloæ</i>	47	78	
<i>Odontocælus thomasi</i>	47	75	
<i>Odontocælus toltecus</i>	46	74	
<i>Odontocælus truii</i>	46	73, 74	
<i>Ædipomidas</i>	532		
<i>Ædipomidas geoffroyi</i>	532		
<i>Ædipus</i>	532		
<i>ædipus</i> . (Hapale)	532	724	
<i>ædipus</i> . (Simia)	532		
<i>ærstedii</i> . (Saimiri).....	534	731	
<i>Ogotoma</i>	337		266
<i>olida</i> . (Mephitis).....	406		
<i>Olidosus</i>	36	65	
<i>Oligoryzomys</i>	180, 189	233, 248	
<i>olivaceo-fuscus</i> . (Dysopes)	495	620	
<i>olivaceus</i> . (Citellus t.).....	104		
<i>olivaceus</i> . (Perognathus).....	307		249
<i>olivaceus</i> . (Perognathus m.).....	307		249

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
olivaceus. (Spermophilus t.).....	104		100
olympica. (Haplodontia).....	127		114
olympica. (Spilogale).....	415		332
olympicus. (Phenacomys).....	227		168
olympicus. (Sciuropterus a.).....	123		
olympicus. (Sciuropterus v.).....	123		110
olympus. (Arctomys).....	121		107
olympus. (Felis h.).....	369	454	294
olympus. (Marmota).....	121		
onca. (Felis).....	365		
Oncifelis.....	365	443	
onchilla. (Felis p.).....	367	450	
Ondatra.....	252		211
oneida. (Arvicola).....	235		180
Onychomys.....	130	165	119
Onychomys arcticeps.....	132		121
Onychomys fuliginosus.....	134		123
Onychomys leucogaster.....	130		119
Onychomys l. albescens.....	131	†742	
Onychomys l. brevicaudus.....	130		120
Onychomys l. longipes.....	131		120
Onychomys l. melanophrys.....	131		120
Onychomys l. pallescens.....	131		121
Onychomys longicaudus.....	132		121
Onychomys l. tularensis.....	132		
Onychomys longipes.....	131		120
Onychomys macrotis.....	134	169	
Onychomys melanophrys.....	131, 132	166	
Onychomys m. pallescens.....	131	166	
Onychomys pulcher.....	134		
Onychomys ramona.....	132	167	121
Onychomys ruidosæ.....	131		
Onychomys torridus.....	133	167	122
Onychomys t. arenicola.....	133	168	122
Onychomys t. canus.....	133	†742	
Onychomys t. perpallidus.....	133	168	122
Onychomys t. tularensis.....	132		
Onychomys t. yakiensis.....	133	†742	
operarius. (Arvicola).....	241		187
operarius. (Eutamias a.).....	†541		
operarius. (Microtus).....	241		187
operarius. (Tamias a.).....	†541		
operarius. (Thomomys).....	273		223
Oplacerus.....	43	68	
oporaphilum. (Phyllostoma).....	529	715	
opossum. (Metachirus).....		12	
oramontis. (Phenacomys).....	226		168
orarius. (Scapanus).....	468		392
orarius. (Sciurus).....			65

†Appendix.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
orarius. (Zapus)	327		262
orbitalis. (Heteromys).....	320	380	
orbitalis. (Liomys).....	320	380	
Orca		50	22
Orca arcticus.....	26	51	22
Orca ater	26		
Orca atra			22
Orca destructor.....	27	51	23
Orca europæus	26	51	22
Orca latirostris	26	51	22
Orca meridionalis	27	51	23
Orca rectipinna.....	26		22
Orca schlegelii.....	26	51	22
Orca stenorhyncha.....	26	51	22
orca. (Delphinus).....	26	50, 51	
orca. (Evotomys).....	230		172
orca. (Orcinus).....	26	51	
Orcinus.....	26	50	21
Orcinus gladiator	26	51	22
Orcinus orca.....	26	51	
Orcinus rectipinna	26		22
ordi. (Perodipus).....	296	350	238
ordii. (Dipodomys)	296		
Oreamnos.....	52		44
Oreamnos m. columbianus.....	52		
Oreamnos m. missoulæ	52		
Oreamnus	52		44
Oreamnus kennedyi.....	53		44
Oreamnus montanus	52		44
Oreamnus m. columbianus	52		
Oreamnus m. missoulæ	52		
oreas. (Peromyscus)	152		134
oregonensis. (Felis c.).....	369	454	294
oregonensis. (Felix).....	369		
oregonensis. (Geomys)	262		217
oregonensis. (Pteromys)	122		110
oregonensis. (Putorius x.).....	431		350
oregonensis. (Sciuropterus a.)	122		
oregonensis. (Sciuropterus v.)	122		110
oregonensis. (Vespertilio)	476	578	403
oregoni. (Arvicola).....	250		200
oregoni. (Microtus)	250		200
oregonus. (Bassariscus a.)	398		317
oregonus. (Bassariscus f.)	398		
oregonus. (Citellus)	113		
oregonus. (Microdipodops m.)	299		
oregonus. (Spermophilus).....	113		94
oregonus. (Thomomys d.).....	277		
oregonus. (Zapus p.).....	326		260
oreocetes. (Cratogeomys)	265	314	

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
oreocetes. (Eutamias)	95		
oreocetes. (Tamias)	95		80
oreolestes. (Teonoma)	223		
oreophila. (Blarina)	464	559	
oreophilus. (Phenacomys)	226		168
oreopolus. (Sorex)	451	550	
oresterus. (Peromyscus)	139	200	
oribasus. (Putorius l.)	430		348
orientalis. (Pusa)	436	537	354
origenes. (Mustela c.)	422		
orinomus. (Myotis)	476	577	
orinus. (Sorex)	456	553	
orizabæ. (Lepus)	351	425	
orizabæ. (Neotoma)	219	286	
orizabæ. (Neotomodon)	225	294	
orizabæ. (Peromyscus m.)	169	207	
orizabæ. (Reithrodontomys)	200	266	
orizabæ. (Rhithrodontomys)	200	266	
orizabæ. (Sorex)	449	550	
orizabæ. (Thomomys)	282	337	
ornatus. (Ateles)	535	733	
ornatus. (Dipodomys)	291	344	
ornatus. (Sorex)	452		374
orolestes. (Neotoma)	223		166
orophila. (Blarina)	464	559	
Orthogeomys	268	322	
Orthogeomys grandis	268, 269	322	
Orthogeomys latifrons	269	324	
Orthogeomys nelsoni	269	324	
Orthogeomys scalops	269	324	
orthotis. (Nyctinomys)	497	623	
orthotis. (Promops)	497	623	
Orthriomys	234, 251	304	179
Oryctogale.	410	512	
Oryctomys	272	332	221
Oryctomys bottæ	274		223
Oryx	24		18
oryzivora. (Arvicola)	180		147
Oryzomys	180	232, 233	146
Oryzomys albiventer	182	236	
Oryzomys alfari	183	242	
Oryzomys angusticeps	187	245	
Oryzomys antillarum	188	247	
Oryzomys aquaticus	181	241	147
Oryzomys a. coloratus	181		148
Oryzomys bulleri	185	239	
Oryzomys chapmani	186	237, 238	
Oryzomys c. caudatus	186	238	
Oryzomys c. dilutior	186	238	
Oryzomys c. saturatior	186	239	

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Oryzomys cherriei</i>	190	252, 253	
<i>Oryzomys childi</i>		244	
<i>Oryzomys chrysomelas</i>	190	253	
<i>Oryzomys costaricensis</i>	189	249, 250	
<i>Oryzomys couesi</i>	186	236, 239	
<i>Oryzomys cozumelæ</i>	182	241	
<i>Oryzomys crinitus</i>	183	245	
<i>Oryzomys c. aztecus</i>	183	245	
<i>Oryzomys devius</i>	188	244	
<i>Oryzomys fulgens</i>	183	240	
<i>Oryzomys fulvescens</i>	189	248	
<i>Oryzomys goldmani</i>	186	246	
<i>Oryzomys hylocetes</i>	187	237	
<i>Oryzomys jalapæ</i>	184	246, 247	
<i>Oryzomys j. apatelius</i>	184	246	
<i>Oryzomys j. rufinus</i>	185	247	
<i>Oryzomys lucie</i>	190	251	
<i>Oryzomys melanotis</i>	187	243, 244	
<i>Oryzomys mexicanus</i>	184	235, 242	
<i>Oryzomys m. peragrus</i>	184	243	
<i>Oryzomys molestus</i>	183	240	
<i>Oryzomys natator</i>	181		
<i>Oryzomys n. floridanus</i>	181		
<i>Oryzomys nelsoni</i>	182	235	
<i>Oryzomys palatinus</i>	187	237	
<i>Oryzomys palustris</i>	180	242	147
<i>Oryzomys p. coloratus</i>	181		
<i>Oryzomys p. natator</i>	181		147
<i>Oryzomys p. texensis</i>	181		147
<i>Oryzomys panamensis</i>	182	241	
<i>Oryzomys peninsulæ</i>	182	236	
<i>Oryzomys rhabdops</i>	187	237, 246	
<i>Oryzomys richmondi</i>	184	248	
<i>Oryzomys rostratus</i>	188	243, 244	
<i>Oryzomys r. megadon</i>	188	243	
<i>Oryzomys rufus</i>	185	239	
<i>Oryzomys talamancæ</i>	183	241	
<i>Oryzomys teapensis</i>	185	247	
<i>Oryzomys tectus</i>	188	244	
<i>Oryzomys vegetus</i>	189	249	
<i>Oryzomys victus</i>	188	244	
<i>Oryzomys yucatanensis</i>	189	236	
<i>Oryzomys zygomatiscus</i>	185	241	
<i>osborni</i> . (Rangifer).....	38		
<i>osburnii</i> . (Chilonycteris).....	505	642	
<i>osceola</i> . (Cariacus).....	44		
<i>osceola</i> . (Eptesicus f.).....	483		
<i>osceola</i> . (Odocoileus a.).....	44		40
<i>osceola</i> . (Odontocelus a.).....	44		
<i>osceola</i> . (Vespertilio f.).....	483		

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
osgoodi. (Citellus)	110		
osgoodi. (Spermophilus)	110		91
osoyoosensis. (Fiber)	252		212
osphyia. (Megaptera n.)	17		10
Osteopera.	337	408	
Otaria californiana.	437		356
Otaria gillespii.	437	539	356
Otaria jubata.	437		
Otaria kracheninikowi.			357
Otaria stelleri.	437		
Otariidæ.	437	538	1355
Otelaphus.	43	68	
othus. (Lepus)	359		272
Otisorex platyrhinus	446		367
Otocolobus.	96	139	82
Otognosis.	300	352	241
Otognosis longimembris.	306		247
Otopterus.	508, 510	651	419
Otopterus bocourtianus.	510	654	
Otopterus bulleri	510	654	
Otopterus californicus	509, 510	653	420
Otopterus mexicanus.	510	653	
Otopterus pygmæus.	511		
Otopterus waterhousii	508	652	
Otopterus w. compressus.	509		
Otopterus w. jamaicensis	508		
Otopterus w. minor	509		
Otosciurus	59, 66	94, 111	50, 57
Otospermophilus.	96, 113	139, 148	82, 87
Ototylomys	174	220	
Ototylomys phyllotis.	179	220, 221	
Ototylomys p. phæus.	174	221	
otus. (Reithrodontomys l.)	203	271	
otus. (Rhithrodontomys l.)	203	271	
Ovibos	55		48
Ovibos moschatus.	55, 56		48
Ovibos m. niphæcus	56		
Ovibos m. wardi.	56		*485
Ovibos pallantis	55		48
Ovis	53	83	46
Ovis ammon	53		46
Ovis aries.	53	83	
Ovis californianus.	53		46
Ovis canadensis	53		46
Ovis c. auduboni.	53		
Ovis c. liardensis.	54		47
Ovis cervina	53	86	46
Ovis c. auduboni.	53		
Ovis c. cremnobates.	54	84	

* Supplement.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Ovis c. mexicanus</i>	54	86	
<i>Ovis c. nelsoni</i>	54	84, 85	46
<i>Ovis dalli</i>	55		47
<i>Ovis d. kenaiensis</i>	55		
<i>Ovis fannini</i>	55		†425
<i>Ovis mexicanus</i>	54		
<i>Ovis montanus</i>	52		44
<i>Ovis m. dalli</i>	55		
<i>Ovis nelsoni</i>	54		
<i>Ovis pygargus</i>	53		46
<i>Ovis stonei</i>	54		47
<i>Ovis stonii</i>	54	85	
<i>Ozolictis</i>	410	512	
<i>Paca</i>	337	408	
<i>paca</i> . (<i>Mus</i>).....	337	408	
<i>pacifica</i> . (<i>Aplodontia</i>).....	127		114
<i>pacifica</i> . (<i>Haplodontia</i>).....	127		114
<i>pacifica</i> . (<i>Lutra c.</i>).....	434		353
<i>pacifica</i> . (<i>Lutra l.</i>).....	434		
<i>pacifica</i> . (<i>Mustela c.</i>).....	423		
<i>pacifica</i> . (<i>Mustela p.</i>).....	423		337
<i>pacifica</i> . (<i>Procyon p.</i>).....	401		
<i>pacificus</i> . (<i>Antrozous p.</i>).....	491	607	397
<i>pacificus</i> . (<i>Lepus</i>).....	348	†744	
<i>pacificus</i> . (<i>Lepus v.</i>).....	348		
<i>pacificus</i> . (<i>Perognathus</i>).....	306	355	246
<i>pacificus</i> . (<i>Procyon l.</i>).....	401		319
<i>pacificus</i> . (<i>Sorex</i>).....	454		375
<i>pacificus</i> . (<i>Zapus</i>).....	327		262
<i>Pagomys</i>	441	541	361
<i>Pagophilus</i>	441, 444	541	361, 364
<i>palatina</i> . (<i>Neotoma</i>).....	205		
<i>palatinus</i> . (<i>Oryzomys</i>).....	187	237	
<i>palitans</i> . (<i>Lepus a.</i>).....	353	434	
<i>pallantis</i> . (<i>Ovibos</i>).....	55		48
<i>pallasi</i> . (<i>Bos</i>).....	56		48
<i>pallascens</i> . (<i>Corynorhinus m.</i>).....	490	604	399
<i>pallascens</i> . (<i>Felis r.</i>).....	371		297
<i>pallascens</i> . (<i>Lynx f.</i>).....	371		297
<i>pallascens</i> . (<i>Onychomys l.</i>).....	131		121
<i>pallascens</i> . (<i>Onychomys m.</i>).....	131	166	
<i>pallascens</i> . (<i>Peromyscus m.</i>).....	160		140
<i>pallascens</i> . (<i>Thomomys b.</i>).....	274		224
<i>palliata</i> . (<i>Alouatta</i>).....	533	726	
<i>palliatas</i> . (<i>Mycetes</i>).....	533	726	
<i>pallida</i> . (<i>Nasua n.</i>).....	403	498	
<i>pallidulus</i> . (<i>Dipodomys c.</i>).....	292		†431
<i>pallidum</i> . (<i>Coendu</i>).....	334	401	

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
pallidus. (Antrozous).....	491	605	396
pallidus. (Arvicola)	247		195
pallidus. (Caluromys l.)	4	10	
pallidus. (Canis).....	377		301
pallidus. (Cercolabes)	334	401	
pallidus. (Citellus t.)	104		
pallidus. (Fiber z.).....	253	308	213
pallidus. (Metachirus f.)	5	13	
pallidus. (Microdipodops).....	299		
pallidus. (Microtus).....	247		195
pallidus. (Myotis c.)	477	579	405
pallidus. (Perognathus f.).....	314	366	
pallidus. (Philander l.)	4	10	
pallidus. (Procyon)	401		
pallidus. (Procyon l.)	401		318
pallidus. (Reithrodontomys)	195	262	
pallidus. (Reithrodontomys l.)	195	262	152
pallidus. (Rhithrodontomys l.)	195	262	
pallidus. (Sigmodon h.).....	175	228	145
pallidus. (Spermophilus t.).....	104		100
pallidus. (Tamias a.).....	93		
pallidus. (Tamias q.).....	85	135	68
pallidus. (Vespertilio)	491		396
palmarius. (Peromyscus g.)	143		127
palmata. (Chironectes)	1	3	
palmatus. (Alces).....	41		38
palmeri. (Atophyrax b.).....	459		381
palmeri. (Eutamias)	86		
palmeri. (Perodipus o.).....	296	350	
palmeri. (Sorex b.)	459		381
palmeri. (Tamias)	86		69
Paludicola	234	299	178
paludicola. (Lepus)	341		279
paludicola. (Lepus p.).....	341		279
palustris. (Arvicola)	234		179
palustris. (Hesperomys)	180		
palustris. (Lepus).....	341	415	279
palustris. (Mus)	180	233	146
palustris. (Neosorex).....	457		378
palustris. (Oryzomys)	180	242	147
palustris. (Sorex).....	457, 458		379
pambasileus. (Canis).....	374		
pan. (Ateles).....	534	733	
panamensis. (Dasyptherus e.).....	487	597	
panamensis. (Felis).....	368	452	
panamensis. (Nasua n.).....	403	498	
panamensis. (Neomys).....	174	219	
panamensis. (Oryzomys).....	182	241	
panamensis. (Phyllostoma h.).....	514		
panamensis. (Proechimys c.)	329	387	

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
panamensis. (<i>Proechinomys</i> c.).....	329	387	
panamensis. (<i>Tylomys</i>).....	174	219	
panamintinus. (<i>Perodipus</i>).....	295		239
panamintinus. (<i>Perognathus</i>).....	304		245
panamintinus. (<i>Perognathus</i> l.).....	304		
panamintinus. (<i>Tamias</i>).....	90		74
pandora. (<i>Hippocamelus</i>).....	51		
pandora. (<i>Mazama</i>).....	51	80	
paniscus. (<i>Simia</i>).....	534	732	
pansa. (<i>Macrogeomys</i>).....	271	329	
Pappogeomys.....	268	320	
Pappogeomys albinasus.....	268	321	
Pappogeomys bulleri.....	268	320	
Paradoxurus annulatus.....	399	489	
paradoxus. (<i>Perognathus</i>).....	310		252
paradoxus. (<i>Perognathus</i> h.).....	310		252
paradoxus. (<i>Solenodon</i>).....	472	568	
paraguënsis. (<i>Mustela</i>).....	1	3	
Paralces.....	40		
paralius. (<i>Heteromys</i>).....	320	381	
Parascalops.....	471		393
Parascalops breweri.....	471		394
Parasciurus.....	59, 64	94, 110	50, 51
parasiticus. (<i>Peromyscus</i>).....	151		
Pardalis.....	365	443	
pardalis. (<i>Felis</i>).....	366	447	294
parnelli. (<i>Chilonycteris</i>).....	505	642	
parnellii. (<i>Phyllodia</i>).....	505	642	
parryi. (<i>Citellus</i>).....	107		
parryii. (<i>Arctomys</i>).....	107		89
parsonsi. (<i>Phoca</i>).....	441		362
parva. (<i>Blarina</i>).....	463		384
parviceps. (<i>Heteromys</i>).....	322	378	
parviceps. (<i>Liomys</i>).....	322	398	
parvidens. (<i>Citellus</i> m.).....	104	146	
parvidens. (<i>Cynomys</i>).....	118		
parvidens. (<i>Neotoma</i>).....	219	288	
parvidens. (<i>Spermophilus</i> m.).....	104	146	99
parvidens. (<i>Urocyon</i>).....	389		
parvidens. (<i>Urocyon</i> c.).....	389	476	
parvipes. (<i>Artibeus</i>).....	523	694	
parvula. (<i>Rhogeassa</i>).....	489		
parvula. (<i>Rhogoessa</i>).....	489	601	
parvulus. (<i>Lepus</i>).....	348	421	
parvus. (<i>Citellus</i> t.).....	104		
parvus. (<i>Cricetodipus</i>).....	306		247
parvus. (<i>Dipodomys</i>).....	289		234
parvus. (<i>Dipodomys</i> m.).....	289		234
parvus. (<i>Perognathus</i>).....	306		247
parvus. (<i>Scalops</i>).....	470		390

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
parvus. (Scalops a.).....	471		390
parvus. (Sorex).....	447, 463		368, 384
parvus. (Spermophilus t.).....	104		100
pascalis. (Thomomys a.).....	282		
paulus. (Peromyscus).....	136	176	
pauperrima. (Arvicola).....	247		195
pauperrimus. (Microtus).....	247		195
pavidus. (Peromyscus).....	164	197	
pealei. (Phoca).....	442		363
pearyi. (Rangifer).....	40		
peba. (Dasypus).....	13	33	4
pecari. (Tagassu).....	37	66, 67	
pecari. (Tayassu).....	37	65, 66	
pectoralis. (Peromyscus a.).....	154	192	
pediculus. (Conepatus).....	412	517	
Pedomys.....	234	299	178
Pelagios.....	444	542	
Pelagocyon.....	444	542	
Peltorhinus.....	526, 527	704, 707	
penicillatus. (Perognathus).....	311		252
penicillatus. (Peromyscus b.).....	150	196	133
peninsulæ. (Blarina b.).....	461		383
peninsulæ. (Blarina c.).....	461		
peninsulæ. (Canis).....	378	466	
peninsulæ. (Citellus l.).....	97	143	
peninsulæ. (Mazama h.).....	50	78	
peninsulæ. (Odontocælus h.).....	50	78	
peninsulæ. (Oryzomys).....	182	236	
peninsulæ. (Perognathus s.).....	316	363	
peninsulæ. (Putorius).....	429		347
peninsulæ. (Rhithrodontomys).....	197	263	
peninsulæ. (Tamias l.).....	97	143	
peninsulæ. (Vespertilio f.).....	484	589	
peninsularis. (Feliş).....	373	460	
peninsularis. (Lepus).....	353	430	
peninsularis. (Lynx r.).....	373		
peninsularis. (Myotis).....	474	573	
pennanti. (Mustela).....	422		335
pennantii. (Talpa).....	470		388
pennsylvanica. (Felis).....	369		
pennsylvanica. (Mus).....	234		179
pennsylvanica. (Neotoma).....	210		157
pennsylvanica. (Scalops).....	470		388
pennsylvanica. (Vulpes).....	380		304
pennsylvanicus. (Microtus).....	234, 236		179
pennsylvanicus. (Sciurus).....	79		53
peragus. (Oryzomys m.).....	184	243	
perditus. (Thomomys).....	283	338	
perdus. (Putorius t.).....	433	531	
peregrina. (Blarina m.).....	465	560	

	Check-List of Mammals.	Mammals of Middle America, etc	Synopsis of Mammals, etc.
peregrinator. (<i>Sciurus p.</i>).....	70		
peregrinus. (<i>Cratogeomys</i>).....	265	314	
peregrinus. (<i>Thomomys</i>).....	282	337	
pergracilis. (<i>Blarina</i>).....	463	557	
pericalles. (<i>Perognathus</i>).....	303		
periclyzomæ. (<i>Lutra</i>).....	434		
perimecurus. (<i>Peromyscus</i>).....	149		
perniger. (<i>Perognathus f.</i>).....	301		
pernix. (<i>Perognathus</i>).....	312	359	
Perodipus.....	293	349	236
Perodipus agilis.....	293, 294	351	236
Perodipus a. tularensis.....	293		
Perodipus cabezonæ.....	295		
Perodipus chapmani.....	297	350	237
Perodipus conipactus.....	299		240
Perodipus goldmani.....	294		
Perodipus ingens.....	298		
Perodipus longipes.....	298		239
Perodipus microps.....	297, 298		
Perodipus m. levipes.....	298		
Perodipus montanus.....	295		
Perodipus m. utahensis.....	296		
Perodipus obscurus.....	299	351	
Perodipus ordi.....	296	350	238
Perodipus o. columbianus.....	297		238
Perodipus o. palmeri.....	296	350	
Perodipus panamintinus.....	295		239
Perodipus richardsoni.....	297		238
Perodipus sennetti.....	297		239
Perodipus streatori.....	294		237
Perodipus s. simulans.....	295		
Perodipus venustus.....	294		
Perognathus.....	300	352	241
Perognathus alticola.....	307		249
Perognathus amplus.....	306		247
Perognathus anthonyi.....	314	366	
Perognathus apache.....	302, 303		244
Perognathus a. melanotis.....	303	356	
Perognathus arenarius.....	312		
Perognathus artus.....	313		
Perognathus baileyi.....	309	361	251
Perognathus b. rhydinorhis.....	309	362	
Perognathus bimaculatus.....	302		
Perognathus bryanti.....	316		
Perognathus californicus.....	315		254
Perognathus c. dispar.....	315		255
Perognathus c. ochrus.....	315		
Perognathus callistus.....	304		245
Perognathus cnecus.....	309	362	
Perognathus columbianus.....	308		250

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Perognathus conditi</i>	310		252
<i>Perognathus copei</i>	301		243
<i>Perognathus elibatus</i>	304		
<i>Perognathus eremicus</i>	311		253
<i>Perognathus fallax</i>	314	366.	254
<i>Perognathus f. pallidus</i>	314	366	
<i>Perognathus fasciatus</i>	300, 309		242, 251
<i>Perognathus f. infraluteus</i>	301		242
<i>Perognathus femoralis</i>	314	367	254
<i>Perognathus f. mesopolius</i>	315	367	
<i>Perognathus flavescens</i>	301		243
<i>Perognathus f. perniger</i>	301		
<i>Perognathus flavus</i>	302	354	244
<i>Perognathus f. bimaculatus</i>	302		244
<i>Perognathus f. fuliginosus</i>	302		244
<i>Perognathus f. mexicanus</i>	303	355	
<i>Perognathus formosus</i>	308		250
<i>Perognathus fuliginosus</i>	302		244
<i>Perognathus goldmani</i>	313	365	
<i>Perognathus g. artus</i>	313	365	
<i>Perognathus helleri</i>	312	360	
<i>Perognathus hispidus</i>	309	360	251
<i>Perognathus h. maximus</i>	310		
<i>Perognathus h. paradoxus</i>	310		252
<i>Perognathus h. zacatecæ</i>	310	361	
<i>Perognathus infraluteus</i>	301		242
<i>Perognathus inornatus</i>	306		247
<i>Perognathus intermedius</i>	313	364	253
<i>Perognathus i. canescens</i>	313		
<i>Perognathus latirostris</i>	310		252
<i>Perognathus longimembris</i>	306		247
<i>Perognathus l. bangsi</i>	305		
<i>Perognathus l. panamintinus</i>	304		
<i>Perognathus lordi</i>	308		250
<i>Perognathus l. columbianus</i>	308		250
<i>Perognathus margaritæ</i>	316	363	
<i>Perognathus mearnsi</i>	302	353	243
<i>Perognathus merriami</i>	301	353	243
<i>Perognathus m. gilvus</i>	302		243
<i>Perognathus mesembrinus</i>	308		
<i>Perognathus mollipilosus</i>	307		249
<i>Perognathus monticola</i>	306, 307		248
<i>Perognathus m. magruderensis</i>	307		249
<i>Perognathus m. mollipilosus</i>	307		249
<i>Perognathus m. olivaceus</i>	307		249
<i>Perognathus nelsoni</i>	313	364	
<i>Perognathus n. canescens</i>	313	365	
<i>Perognathus nevadensis</i>	305		246
<i>Perognathus obscurus</i>	311, 313		253
<i>Perognathus olivaceus</i>	307		249

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Perognathus o. amoenus</i>	307		
<i>Perognathus pacificus</i>	306	355	246
<i>Perognathus panamintinus</i>	304		245
<i>Perognathus p. arenicola</i>	305		246
<i>Perognathus p. bangsi</i>	305		245
<i>Perognathus p. brevinasus</i>	305	355	246
<i>Perognathus paradoxus</i>	310		252
<i>Perognathus p. spilotus</i>	309		251
<i>Perognathus parvus</i>	306		247
<i>Perognathus p. magruderensis</i>	307		
<i>Perognathus penicillatus</i>	311		252
<i>Perognathus p. angustirostris</i>	311	358	252
<i>Perognathus p. arenarius</i>	312	358	
<i>Perognathus p. eremicus</i>	311	359	253
<i>Perognathus p. pricii</i>	311	358	253
<i>Perognathus pericalles</i>	303		
<i>Perognathus pernix</i>	312	359	
<i>Perognathus p. rostratus</i>	312	359	
<i>Perognathus pricei</i>	311		253
<i>Perognathus spinatus</i>	315	363	255
<i>Perognathus s. bryanti</i>	316	364	
<i>Perognathus s. peninsulae</i>	316	363	
<i>Perognathus stephensi</i>	312		253
<i>Peromyscus</i>	135, 138	170	123
<i>Peromyscus affinis</i>	161	180	
<i>Peromyscus a. musculoides</i>	161	180	
<i>Peromyscus akeleyi</i>	148		129
<i>Peromyscus allex</i>	135	175	
<i>Peromyscus allophylus</i>	169	208	
<i>Peromyscus altilaneus</i>	170	210	
<i>Peromyscus americanus</i>	140		124
<i>Peromyscus a. arizonæ</i>	140	186	125
<i>Peromyscus a. artemisiæ</i>	141		126
<i>Peromyscus a. deserticolus</i>	140	188	125
<i>Peromyscus a. noveboracensis</i>	140		125
<i>Peromyscus a. sonoriensis</i>	141	182	125
<i>Peromyscus amplus</i>	167	205	
<i>Peromyscus anastase</i>	143		
<i>Peromyscus anastasiæ</i>	143, 144		128
<i>Peromyscus anthonyi</i>	137	185	141
<i>Peromyscus arboreus</i>	135	170	
<i>Peromyscus attwateri</i>	154		135
<i>Peromyscus a. eremicoides</i>	154	192	
<i>Peromyscus a. pectoralis</i>	154	192	
<i>Peromyscus auripectus</i>	153		134
<i>Peromyscus auritus</i>	170	209	
<i>Peromyscus austerus</i>	149		132
<i>Peromyscus aztecus</i>	162	184	
<i>Peromyscus banderanus</i>	163	189	
<i>Peromyscus b. angelensis</i>	163	190	

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Peromyscus b. vicinior</i>	163	190	
<i>Peromyscus beatæ</i>	162	184	
<i>Peromyscus bellus</i>	154		135
<i>Peromyscus boylii</i>	150	195	132
<i>Peromyscus b. penicillatus</i>	150	196	133
<i>Peromyscus bullatus</i>	158	193	
<i>Peromyscus cacabatus</i>	171	211	
<i>Peromyscus californicus</i>	137, 138	203	137
<i>Peromyscus canadensis</i>	144		128
<i>Peromyscus c. abietorum</i>	144		128
<i>Peromyscus c. nubiterræ</i>	145		129
<i>Peromyscus c. umbrinus</i>	144		129
<i>Peromyscus canus</i>	150		132
<i>Peromyscus catalinæ</i>	149		
<i>Peromyscus ceciliæ</i>	161	178	
<i>Peromyscus cedrosensis</i>	155	185	
<i>Peromyscus cherrii</i>	160	183, 267	
<i>Peromyscus cineritius</i>	164	195	
<i>Peromyscus comptus</i>	168	206	
<i>Peromyscus cozumelæ</i>	162	181	
<i>Peromyscus c. scitulus</i>	159		
<i>Peromyscus difficilis</i>	158	193	
<i>Peromyscus dubius</i>	162	181	
<i>Peromyscus dyselius</i>	156		138
<i>Peromyscus eremicus</i>	136	191	136
<i>Peromyscus e. arenarius</i>	137	191	136
<i>Peromyscus e. phæurus</i>	137	191	
<i>Peromyscus e. propinquus</i>	137	185	
<i>Peromyscus eva</i>	165	198	
<i>Peromyscus exiguus</i>	148	183	
<i>Peromyscus felipensis</i>	167	205	
<i>Peromyscus floridanus</i>	156		138
<i>Peromyscus fraterculus</i>	155	184	136
<i>Peromyscus furvus</i>	169	208	
<i>Peromyscus gaurus</i>	148	199	
<i>Peromyscus geronimensis</i>	162	181	
<i>Peromyscus goldmani</i>	138	186	
<i>Peromyscus gossypinus</i>	143		
<i>Peromyscus g. nigriculus</i>	143		
<i>Peromyscus g. palmarius</i>	143		
<i>Peromyscus gossypinus</i>	143		126
<i>Peromyscus g. mississippiensis</i>	143		127
<i>Peromyscus g. nigriculus</i>	143		127
<i>Peromyscus g. palmarius</i>	143		127
<i>Peromyscus gratus</i>	164	196	
<i>Peromyscus g. gentilis</i>	164	197	
<i>Peromyscus guatemalensis</i>	171	211	
<i>Peromyscus gymnotis</i>	162	188	
<i>Peromyscus hemionotis</i>	139	199	
<i>Peromyscus herroni</i>	155		137

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Peromyscus h. nigellus</i>	155		137
<i>Peromyscus homochroia</i>	148	200	
<i>Peromyscus hylocetes</i>	167	204	
<i>Peromyscus insignis</i>	155	203	138
<i>Peromyscus insolatus</i>	154		135
<i>Peromyscus insulanus</i>	144		128
<i>Peromyscus keeni</i>	151		133
<i>Peromyscus labecula</i>	142	178	
<i>Peromyscus lasius</i>	157		
<i>Peromyscus lepturus</i>	167	204	
<i>Peromyscus leucopus</i>	140, 141		124, *492
<i>Peromyscus l. ammodytes</i>	†542		
<i>Peromyscus l. arizonæ</i>	140		
<i>Peromyscus l. artemisiæ</i>	141		
<i>Peromyscus l. blandus</i>	142	182	
<i>Peromyscus l. coolidgei</i>	147		
<i>Peromyscus l. deserticola</i>	140	188	
<i>Peromyscus l. fulvus</i>	142	183	
<i>Peromyscus l. fusus</i>	†542		
<i>Peromyscus l. minnesotæ</i>	141		
<i>Peromyscus l. myodes</i>	140		125
<i>Peromyscus l. noveboracensis</i>	141		125
<i>Peromyscus l. sonoriensis</i>	141	181	
<i>Peromyscus leucurus</i>	165	200	
<i>Peromyscus l. gadovi</i>	165	201	
<i>Peromyscus levipes</i>	164	195	
<i>Peromyscus lophurus</i>	167	204	
<i>Peromyscus luteus</i>	147		
<i>Peromyscus macrorhinus</i>	152		133
<i>Peromyscus madrensis</i>	164	197	
<i>Peromyscus major</i>	155		137
<i>Peromyscus maniculatus</i>	148		131
<i>Peromyscus martirensis</i>	159	189	
<i>Peromyscus mearnsi</i>	150		132
<i>Peromyscus mecisturus</i>	171	212	
<i>Peromyscus megalops</i>	170	210	
<i>Peromyscus megalotis</i>	157		139
<i>Peromyscus mekisturus</i>	171	212	
<i>Peromyscus melanocarpus</i>	170	210	
<i>Peromyscus melanophrys</i>	166	201	
<i>Peromyscus m. consobrinus</i>	166	202	
<i>Peromyscus m. zamoræ</i>	166	202	
<i>Peromyscus melanotis</i>	161	178	
<i>Peromyscus m. zamelas</i>	161	179	
<i>Peromyscus merriami</i>	165	198	
<i>Peromyscus metallicola</i>	138	192	
<i>Peromyscus mexicanus</i>	168	207	
<i>Peromyscus m. orizabæ</i>	169	207	

* Supplement.

† Appendix.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Peromyscus m. saxatilis</i>	169	207	
<i>Peromyscus m. teapensis</i>	168	207	
<i>Peromyscus m. totontepecus</i>	169	208	
<i>Peromyscus michiganensis</i>	160		140
<i>Peromyscus m. pallescens</i>	160		140
<i>Peromyscus montipinoris</i>	157		
<i>Peromyscus musculoides</i>	161		
<i>Peromyscus musculus</i>	135, 136	175	
<i>Peromyscus m. brunneus</i>	136	176	
<i>Peromyscus m. nigrescens</i>	136	176	
<i>Peromyscus niveiventris</i>	138		141
<i>Peromyscus n. phasma</i>	138		
<i>Peromyscus nudipes</i>	165	198	
<i>Peromyscus nuttalli</i>	160		140
<i>Peromyscus oaxacensis</i>	168	206	
<i>Peromyscus oreas</i>	152		134
<i>Peromyscus o. rubidus</i>	153		
<i>Peromyscus oresterus</i>	139	200	
<i>Peromyscus parasiticus</i>	151		
<i>Peromyscus paulus</i>	136	176	
<i>Peromyscus pavidus</i>	164	197	
<i>Peromyscus perimecurus</i>	149		
<i>Peromyscus petraeus</i>	153		
<i>Peromyscus phasma</i>	138		141
<i>Peromyscus polius</i>	153	196	
<i>Peromyscus prevostensis</i>	152		
<i>Peromyscus propinquus</i>	137	185	
<i>Peromyscus robustus</i>	156		138
<i>Peromyscus rowleyi</i>	151		134
<i>Peromyscus r. pinalis</i>	151	189	135
<i>Peromyscus rufinus</i>	142		126
<i>Peromyscus sagax</i>	158	194	
<i>Peromyscus simulatus</i>	167	205	
<i>Peromyscus sitkensis</i>	152		134
<i>Peromyscus sonoriensis</i>	141		
<i>Peromyscus s. blandus</i>	142	182	
<i>Peromyscus s. fulvus</i>	142	183	
<i>Peromyscus spicilegus</i>	156	179	
<i>Peromyscus s. evides</i>	156	179	
<i>Peromyscus s. simulus</i>	157	180	
<i>Peromyscus stephensi</i>	153	190	136
<i>Peromyscus subgriseus</i>	139		142
<i>Peromyscus s. baliolus</i>	139		142
<i>Peromyscus s. rhoadsi</i>	139		142
<i>Peromyscus taylori</i>	135		141
<i>Peromyscus tehuantepecus</i>	168	206	
<i>Peromyscus texanus castaneus</i>	147		
<i>Peromyscus t. clementis</i>	145		
<i>Peromyscus t. flaccidus</i>	147		

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Peromyscus t. medius</i>	146		
<i>Peromyscus t. mesomelas</i>	147		
<i>Peromyscus t. saturatus</i>	145		
<i>Peromyscus t. subarcticus</i>	146		
<i>Peromyscus texensis</i>	145	186	130
<i>Peromyscus t. arcticus</i>	146		131
<i>Peromyscus t. arizonæ</i>	141	186	
<i>Peromyscus t. castaneus</i>	147		
<i>Peromyscus t. clementis</i>	145	187	130
<i>Peromyscus t. coolidgei</i>	147	187	
<i>Peromyscus t. deserticola</i>	140	188	
<i>Peromyscus t. flaccidus</i>	147	187	
<i>Peromyscus t. gambeli</i>	145		130
<i>Peromyscus t. medius</i>	146		130
<i>Peromyscus t. mesomelas</i>	147		
<i>Peromyscus t. nebrascensis</i>	146		130
<i>Peromyscus t. saturatus</i>	145		131
<i>Peromyscus t. subarcticus</i>	146		131
<i>Peromyscus t. thurberi</i>	146		
<i>Peromyscus thomasi</i>	171	213	
<i>Peromyscus thurberi</i>	146	176	
<i>Peromyscus t. castaneus</i>	147	177	
<i>Peromyscus t. mesomelas</i>	147	177	
<i>Peromyscus tiburonensis</i>	160	175	
<i>Peromyscus tornillo</i>	142	188	126
<i>Peromyscus truei</i>	159		
<i>Peromyscus t. crinitus</i>	159		140
<i>Peromyscus t. nasutus</i>	159		
<i>Peromyscus t. scitulus</i>	159		†429
<i>Peromyscus truii</i>	159		139
<i>Peromyscus t. crinitus</i>	159		
<i>Peromyscus t. nasutus</i>	159		
<i>Peromyscus t. scitulus</i>	159		
<i>Peromyscus xenurus</i>	166	202	
<i>Peromyscus yucatanicus</i>	163	194	
<i>Peromyscus y. badius</i>	163	194	
<i>Peromyscus zarhynchus</i>	170	209	
<i>Peromyscus z. cristobalensis</i>	170	209	
<i>Peromyscus zelotes</i>	166	203	
<i>peronii</i> . (<i>Delphinus</i>).....	32		
<i>Peropteryx</i>	493	613	
<i>Peropteryx canina</i>	493	614	
<i>perotensis</i> . (<i>Citellus</i>)	102	145	
<i>perotensis</i> . (<i>Cratogeomys</i>)	265	313	
<i>perotensis</i> . (<i>Dipodomys</i>).....	291	344	
<i>perotensis</i> . (<i>Neotomodon</i>).....	225	293	
<i>perotensis</i> . (<i>Reithrodontomys</i>)	199	266	
<i>perotensis</i> . (<i>Rhithrodontomys</i>)	199	266	
<i>perotensis</i> . (<i>Spermophilus</i>).....	102	145	

†Appendix.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
perpallidus. (<i>Onychomys t.</i>).....	133	168	122
perpallidus. (<i>Thomomys</i>).....	286, 287	340	229
perpallidus. (<i>Thomomys t.</i>).....	286		
perpes. (<i>Thomomys a.</i>).....	285		
perplicatus. (<i>Lepus l.</i>).....	347		
personata. (<i>Chilonycteris</i>).....	504	641	
personatus. (<i>Geomys</i>).....	264		220
personatus. (<i>Sorex</i>).....	446		366
perspicillatum. (<i>Hemiderma</i>).....	515	668	
perspicillatum. (<i>Phyllostoma</i>).....	524	695	
perspicillatus. (<i>Lagenorhynchus</i>).....	29		25
perspicillatus. (<i>Vespertilio</i>).....	515	667	
persultator. (<i>Lepus f.</i>).....	345	418	
peruana. (<i>Chæronycteris</i>).....	520	682	
pervagus. (<i>Thomomys a.</i>).....	285		
petræus. (<i>Peromyscus</i>).....	153		
Petrorhynchus.....	22	47	16
petulans. (<i>Sciurus h.</i>).....	82		†426
Peudostoma floridana.....	261		
pfeifferi. (<i>Atalapha</i>).....	486	593	
pfeifferi. (<i>Lasiurus b.</i>).....	486	593	
phæa. (<i>Aplodontia</i>).....	127		114
phæa. (<i>Haplodontia</i>).....	127		114
phæognatha. (<i>Arctomys p.</i>).....	107		89
phæonotus. (<i>Lepus a.</i>).....	360		273
phæonyx. (<i>Ursus h.</i>).....	393		
phæotis. (<i>Dermanura</i>).....	525	699	
phæura. (<i>Heteromys</i>).....	319	379	
phæura. (<i>Liomys</i>).....	319	379	
phæurus. (<i>Heteromys</i>).....	319		
phæurus. (<i>Peromyscus e.</i>).....	137	191	
phæurus. (<i>Sciurus d.</i>).....	67	†741	
phæus. (<i>Arvicola</i>).....	245	302	
phæus. (<i>Microtus m.</i>).....	245	302	
phæus. (<i>Ototylomys p.</i>).....	174	221	
phaiops. (<i>Vespertilio</i>).....	483	587	410
phasma. (<i>Peromyscus</i>).....	138		141
phasma. (<i>Peromyscus n.</i>).....	138		
Phenacomys.....	226		167
Phenacomys albipes.....	228		
Phenacomys celatus.....	227, 228		169
Phenacomys c. crassus.....	227, 228		†430
Phenacomys constablii.....	228		169
Phenacomys intermedius.....	226		167
Phenacomys latimanus.....	227		169
Phenacomys longicaudus.....	228		169
Phenacomys mackenzii.....	228		
Phenacomys olympicus.....	227		168
Phenacomys oramontis.....	226		168

†Appendix.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
Phenacomys oreophilus.....	226		
Phenacomys preblei.....	227		168
Phenacomys preblui.....	227		168
Phenacomys truei.....	226		168
Phenacomys ungava.....	227		169
phenax. (Spilogale).....	414		381
phenax. (Teanopus).....	225	295	
philander. (Didelphis).....	4	9	
Philander l. pallidus.....	4	10	
phillipsi. (Dipodomys).....	288, 291	343	231
Phoca.....	441, 442	541	361
Phoca albicauda.....	444		364
Phoca albigena.....	441		362
Phoca albiini.....	444		364
Phoca annellata.....	443, 444		364
Phoca antarctica.....	438		
Phoca barbata.....	441		362
Phoca cristata.....	440		360
Phoca cucullata.....	440		360
Phoca desmaresti.....	444		364
Phoca dorsata.....	444		364
Phoca equestris.....	442		362
Phoca fasciata.....	441		362
Phoca foetida.....	442, 443		363, 364
Phoca frederici.....	444		364
Phoca groenlandica.....	444		364
Phoca g. nigra.....	444		364
Phoca grypus.....	441		361
Phoca hispida.....	443		364
Phoca isidorei.....	440		360
Phoca jubata.....	437		
Phoca lagura.....	444		364
Phoca largha.....	442		363
Phoca leonina.....	437, 439	545	355, 359
Phoca lepechini.....	441		362
Phoca leporina.....	441		362
Phoca leucopha.....			360
Phoca littorea.....	442		363
Phoca mitrata.....	440		360
Phoca monachus.....	444		
Phoca mulleri.....	444		364
Phoca naurica.....	441		362
Phoca nautica.....	441		362
Phoca nigra.....			357
Phoca oceanica.....	444		364
Phoca parsonsi.....	441		362
Phoca pealei.....	442		363
Phoca pilayi.....	444		364
Phoca proboscidea.....	439	545	
Phoca richardi.....	442		

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Phoca r. geronimensis</i>	443	542	
<i>Phoca r. pribilofensis</i>	442		
<i>Phoca rosmarus</i>	439		
<i>Phoca schreberi</i>	444		364
<i>Phoca scopulicola</i>	442		363
<i>Phoca semilunaris</i>	444		364
<i>Phoca stejnegeri</i>	443		
<i>Phoca tropicalis</i>	445		
<i>Phoca ursina</i>	438		357
<i>Phoca variegata</i>	442		363
<i>Phoca vitulina</i>	441, 442	541	363
<i>Phocæna</i>	25	48	20
<i>Phocæna brachycium</i>	25	48	20
<i>Phocæna communis</i>	25	48	20
<i>Phocæna compressicauda</i>	31	56	29
<i>Phocæna crassidens</i>	27	51	23
<i>Phocæna dalli</i>	25		21
<i>Phocæna lineata</i>	25	48	20
<i>Phocæna phocæna</i>	25	48	
<i>Phocæna vomerina</i>	25	48	20
<i>phocæna. (Delphinus)</i>	25	48	20
<i>phocæna. (Phocæna)</i>	25	48	
<i>Phocidæ</i>	439	541	359
<i>Phocinæ</i>	440		361
<i>Phyllodia</i>	503	639	
<i>Phyllodia parnellii</i>	505	642	
<i>Phyllonycteris</i>	520	684	
<i>Phyllonycteris bombifrons</i>	520	685	
<i>Phyllonycteris planifrons</i>	520	685	
<i>Phyllonycteris poeyi</i>	520, 521	686	
<i>Phyllonycteris sezekorni</i>	521	686	
<i>Phyllophora</i>	516	670	
<i>Phyllophora megalotis</i>	512, 513	660, 662	
<i>Phyllophora nigra</i>	516	672	
<i>Phyllops</i>	527	707	
<i>Phyllops falcatum</i>	527	708	
<i>Phyllops falcatus</i>	527	707	
<i>Phyllostoma</i>	514	666	
<i>Phyllostoma albescens</i>	529	715	
<i>Phyllostoma albomaculatum</i>	527	708	
<i>Phyllostoma amblyotis</i>	512	659	
<i>Phyllostoma bicolor</i>	515	669	
<i>Phyllostoma bilabiatum</i>	528	712	
<i>Phyllostoma brachyotum</i>	515	669	
<i>Phyllostoma brevicaudum</i>	515	668	
<i>Phyllostoma calcaratum</i>	515	669	
<i>Phyllostoma chrysocomus</i>	529	715	
<i>Phyllostoma elongata</i>	513	662	
<i>Phyllostoma erythromos</i>	529	715	
<i>Phyllostoma excisum</i>	529	715	

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
Phyllostoma fumarium	529		
Phyllostoma grayi	515	669	
Phyllostoma hastatum	514	666	
Phyllostoma h. panamensis	514		
Phyllostoma lanceolatum	515	669	
Phyllostoma lilium	529	715	
Phyllostoma lineatum	525	702	
Phyllostoma obscurum	524	695	
Phyllostoma oporaphilum	529	715	
Phyllostoma perspicillatum	524	695	
Phyllostoma planirostre	524		
Phyllostoma rotundum	530		
Phyllostoma scrobiculatum	513	662	
Phyllostoma spiculatus	529	715	
Phyllostomatidæ	503	639	418
Phyllostomatinae	507	649	419
Phyllostomus	514	666	
Phyllostomus h. panamensis	514		
Phyllostomus maximus	514	666	
phylotis. (Ototylomys)	174	220, 221	
Physalus	18	42	11
Physalus duguidii	19		13
Physalus sibbaldii	20		13
physalus. (Balæna)	19, 20		13
physalus. (Balænoptera)	19		13
Physeter	20	45	
Physeter bidens	23		18
Physeter breviceps	22	46	*479
Physeter macrocephalus	20, 21	45	
Physeteridæ	20	44	14
Physeterinae	20		14
picinum. (Erethizon d.)	334		
picinus. (Erethizon d.)	334		†431
picta. (Neotoma)	219	287	
pictus. (Heteromys)	322	377	
pictus. (Liomys)	323		
pictus. (Tamias m.)	94		79
pigra. (Didelphis v.)	7		1
pigra. (Didelphys v.)	7		1
pilayi. (Phoca)	444		364
pilorides. (Capromys)	330	390	
pilorides. (Holochilus)		221	
pilorides. (Isodon)	330	390	
pilorides. (Mus)	190	250	
pilosus. (Hystrix)	333	397	264
pinalis. (Peromyscus r.)	151	189	135
pinalis. (Sitomys r.)	151	189	135
Pinemys	234	299	178

*Supplement.

†Appendix.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
pinetis. (Geomys).....	261		215
pinetis. (Lepus a.).....	350		
pinetis. (Lepus g.).....	350		285
pinetis. (Lepus s.).....	350		
pinetorum. (Microtus).....	249		198
pinetorum. (Neotoma).....	211		158
pinetorum. (Psammomys).....	249		198
pinetorum. (Thomomys m.).....	275		225
Pinnipedia.....	437	538	355
Pipistrellus.....	480	581	408
Pipistrellus cinnamomeus.....	482	585	
Pipistrellus hesperus.....	480	582	409
Pipistrellus h. apus.....	481	583	
Pipistrellus h. australis.....	481	584	
Pipistrellus subflavus.....	481		409
Pipistrellus s. obscurus.....	481		409
Pipistrellus vagans.....	482	584	
Pipistrellus verærucis.....	481	584	
pipistrellus. (Vespertilio).....	480	581	408
piscator. (Viverra).....	422		335
pittekajensis. (Balæna m.).....	16		8
Pitymys.....	233, 249	299	178, 197
plagiodon. (Delphinus).....	33	58	31
plagiodon. (Prodelphinus).....	33	58	31
Plagiodontia.....	332	395	
Plagiodontia ædium.....	332	396	
planiceps. (Neotoma m.).....	209		
planiceps. (Platygeomys).....	267	319	
planifrons. (Phyllonycteris).....	520	685	
planirostre. (Phyllostoma).....	524		
planirostris. (Artibeus).....	524	695	
plantinarenensis. (Heteromys).....	322	377	
plantinarenensis. (Liomys).....	322	377	
Platygeomys.....	266	316	
Platygeomys fumosus.....	267	318	
Platygeomys gymnurus.....	266, 267	318	
Platygeomys neglectus.....	268	319	
Platygeomys planiceps.....	267	319	
Platygeomys tylorhinus.....	267	319	
Platygeomys t. angustirostris.....	267	319	
platyrhina. (Chincha).....	409		
platyrhina. (Mephitis).....	409		
platyrhinus. (Otisorex).....	446		367
platyrhinus. (Sorex).....	447		369
platyrhynchus. (Sorex).....	446		366
Plecotinæ.....	490		
Plecotus.....	490	603	398
Plecotus lecontii.....	490		399
Plecotus macrotis.....	490		399
Plecotus townsendii.....	490, 491	604	400

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
plesius. (Citellus)	108		
plesius. (Spermophilus e.)	108		†427
plethodon. (Monophyllus)	518	677	
plicata. (Balantiopteryx)	493	611	
poadromus. (Lepus)	360		272
Pœscopia	17	41	10
poeyi. (Phyllonycteris)	520, 521	686	
poliopus. (Sciurus)	69	117	
poliopus. (Sciurus v.)	69		
polius. (Peromyscus)	153	196	
polonica. (Marmota)	119		
pomeegra. (Delphinus)	30	55	28
popofensis. (Microtus u.)	242		188
portoricensis. (Chilonycteris)	505	643	
portoricensis. (Monophyllus)	518	677	
Potos	404	499	
Potos f. aztecus	404	500	
Potos f. chiriquensis	404	501	
Potos f. megalotus	404		
prasinata. (Condylura)	467		395
pratensis. (Citellus s.)	100		
pratensis. (Mynomes)	234		179
pratensis. (Spermophilus s.)	100		96
Praticola	234	299	178
preblei. (Phenacobius)	227		168
prebii. (Phenacomys)	227		
prehensilis. (Capromys)	331	391	
prehensilis. (Hystrix)	334, 335	399, 402	
prevostensis. (Peromyscus)	152		
prevostensis. (Sorex l.)	450		
prevostensis. (Sorex o.)	450		
pribilofensis. (Phoca r.)	442		
pribilofensis. (Sorex)	454		376
pribilofensis. (Vulpes)	383		
pricei. (Perognathus)	311		253
pricei. (Tamias)	86		
pricei. (Tamias t.)	87		70
pricii. (Perognathus p.)	311	358	253
pricii. (Tamias t.)	86		70
Primates	532	723	
princeps. (Lepus)	338		267
princeps. (Ochotona)	338		267
princeps. (Zapus)	325		260
principalis. (Microtus)	251		
Proboscidea rivalis	492	609	
Proboscidea saxatilis	492	609	
proboscidea. (Phoca)	439	545	
Procapromys	330	389	
Procyon	400	490	317

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Procyon cancrivorus</i>	402	492	
<i>Procyon hernandezii</i>	400		
<i>Procyon lotor</i>	400		317
<i>Procyon l. elucus</i>	400		319
<i>Procyon l. hernandezi</i>	401	491	317
<i>Procyon l. insularis</i>	401	492	
<i>Procyon l. pacificus</i>	401		319
<i>Procyon l. pallidus</i>	401		318
<i>Procyon maynardi</i>	402	490	
<i>Procyon nivea</i>	400		
<i>Procyon pallidus</i>	401		
<i>Procyon psora</i>	400		
<i>Procyon p. pacifica</i>	401		
<i>Procyon pygmaeus</i>	401	490	
<i>Procyonidae</i>	397	482	315
<i>Procyoninae</i>	397	482	315
<i>Prodelphinus</i>	32	57	30
<i>Prodelphinus crotaphiscus</i>	33		31
<i>Prodelphinus euphrosyne</i>	34	58	30
<i>Prodelphinus frenatus</i>	34		32
<i>Prodelphinus longirostris</i>	33	58	
<i>Prodelphinus plagiodon</i>	33	58	31
<i>Proechimys</i>	329	385	
<i>Proechimys burrus</i>	330	388	
<i>Proechimys c. chiriquinus</i>	329	387	
<i>Proechimys c. panamensis</i>	329	387	
<i>Proechinomys</i>	329	385	
<i>Proechinomys burrus</i>	330	388	
<i>Proechinomys centralis</i>	329	386	
<i>Proechinomys c. chiriquinus</i>	329	387	
<i>Proechinomys c. panamensis</i>	329	387	
<i>Proechinomys semispinosus</i>	329	386	
<i>Promops</i>	496	621	421
<i>Promops abrasus</i>	497	623	
<i>Promops californicus</i>	498		421
<i>Promops glaucinus</i>	497	623	
<i>Promops nanus</i>	498	624	
<i>Promops nasutus</i>	496, 497	621	
<i>Promops orthotis</i>	497	623	
<i>Promops ursinus</i>	497	621	
<i>propinquus. (Peromyscus)</i>	137	185	
<i>propinquus. (Peromyscus e.)</i>	137	185	
<i>propinquus. (Vespertilio f.)</i>	484	589	
<i>propinquus. (Vesperus)</i>	484		
<i>proteus. (Evotomys)</i>	233		177
<i>pruinosa. (Marmota)</i>	120		
<i>pruinosis. (Arctomys)</i>	120		106
<i>pruinosis. (Vespertilio)</i>	487	595	413
<i>Psammomys</i>	233	298	178
<i>Psammomys pinetorum</i>	249		198

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Pseudocervus</i>	41		
<i>Pseudorca</i>	26	51	22
<i>Pseudorca crassidens</i>	27	51	23
<i>Pseudostoma</i>	261	310	215
<i>Pseudostoma castanops</i>	266	315	
<i>Pseudostoma floridana</i>	261		215
<i>Pseudostoma umbrinus</i>	272	334	
<i>psilotis</i> . (<i>Chilonycteris</i>).....	505	642	
<i>psora</i> . (<i>Procyon</i>).....	400		
<i>Pterobakena communis</i>	19		13
<i>Pteroderma</i>	522	691	
<i>Pteromyinæ</i>	121	157	107
<i>Pteromys alpinus</i>	121		109
<i>Pteromys oregonensis</i>	122		110
<i>Pteromys s. alpinus</i>	121		
<i>Pteronotus</i>	506	644	
<i>Pteronotus davayi</i>	506	644	
<i>pulcher</i> . (<i>Onychomys</i>).....	134		
<i>pulverulentus</i> . (<i>Vespertilio</i>).....	480		407
<i>Puma</i>	365	443	
<i>puma</i> . (<i>Felis</i>).....			293
<i>pumilus</i> . (<i>Microtus</i>).....	227		168
<i>punctata</i> . (<i>Dasyprocta</i>).....	336	405	
<i>punctulatus</i> . (<i>Cariacus</i>).....	48		41
<i>Pusa</i>	435, 441	537	354, 363
<i>Pusa orientalis</i>	436	357	354
<i>putida</i> . (<i>Mephitis</i>).....	406		
<i>Putorius</i>	423	529	339
<i>Putorius affinis</i>	433	534	
<i>Putorius alleghanensis</i>	428		*511
<i>Putorius alleghenensis</i>	428		*511
<i>Putorius alleni</i>	431		349
<i>Putorius arcticus</i>	428		345
<i>Putorius a. kadiacensis</i>	428		346
<i>Putorius arizonensis</i>	431		349
<i>Putorius cicognani</i>	426		344
<i>Putorius c. alascensis</i>	426		344
<i>Putorius c. richardsoni</i>	426, 427		344
<i>Putorius frenatus</i>	431, 432	532	350
<i>Putorius f. goldmani</i>	432	534	
<i>Putorius f. leucoparia</i>	432	535	
<i>Putorius f. neomexicanus</i>	432	534	351
<i>Putorius haidarum</i>	429		351
<i>Putorius longicaudus</i>	430		348
<i>Putorius l. oribasus</i>	430		348
<i>Putorius l. spadix</i>	430		
<i>Putorius lütensis</i>	425		340
<i>Putorius melampeplus</i>	425		
<i>Putorius microtis</i>	426		

*Supplement.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Putorius muricus</i>	427		†434
<i>Putorius nigrescens</i>	423		338
<i>Putorius nigripes</i>	425		341
<i>Putorius noveboracensis</i>	429		346
<i>Putorius n. notius</i>	429		346
<i>Putorius n. occisor</i>	429		347
<i>Putorius occisor</i>	429		347
<i>Putorius peninsulæ</i>	429		347
<i>Putorius r. alascensis</i>	426		
<i>Putorius rixosus</i>	428		345
<i>Putorius r. eskimo</i>	428		345
<i>Putorius saturatus</i>	430		348
<i>Putorius spadix</i>	430		348
<i>Putorius streatori</i>	427		345
<i>Putorius s. leptus</i>	427		
<i>Putorius tropicalis</i>	432	531	
<i>Putorius t. perdis</i>	433	531	
<i>Putorius vison</i>	423		338
<i>Putorius v. energumenos</i>	424		
<i>Putorius v. energumenus</i>	424, 425		340
<i>Putorius v. ingens</i>	424, 425		340
<i>Putorius v. lacustris</i>	424		
<i>Putorius v. lutrecephalus</i>	423		339
<i>Putorius v. melampeplus</i>	425		
<i>Putorius v. vulgivagus</i>	424		340
<i>Putorius vulgivagus</i>	424		
<i>Putorius washingtoni</i>	429		347
<i>Putorius xanthogenys</i>	431		349
<i>Putorius x. mundus</i>	341		350
<i>Putorius x. oregonensis</i>	431		350
<i>putorius. (Mustela)</i>	423	529	
<i>putorius. (Spilogale)</i>			327
<i>pygargus. (Ovis)</i>	53		46
<i>pygmæa. (Spilogale)</i>	415	519	
<i>pygmæus. (Macrotus)</i>	511		
<i>pygmæus. (Otopterus)</i>	511		
<i>pygmæus. (Procyon)</i>	401	490	
<i>pygmæus. (Thomomys)</i>	284		
<i>Pygoderma</i>	528	712	
<i>Pygoderma bilabiatum</i>	528	712	
<i>pyladei. (Sciurus)</i>	77	129	
<i>pyrrotrichus. (Cynomys)</i>	117		
<i>quadratus. (Thomomys)</i>	276		225
<i>quadridactyla. (Myrmecophaga)</i>	10		
<i>quadridens. (Chilonycteris)</i>	504	641	
<i>quadridens. (Lobostoma)</i>	504	641	
<i>quadrinaculatus. (Tamias)</i>	88		71
<i>quadrivittatus. (Sciurus)</i>	91	138	

†Appendix.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
quadrivittatus. (Tamias)	91	138	75
quasiater. (Arvicola p.)	249		
quasiater. (Microtus)	249, 250	303	
quaterlinearis. (Mephitis)	413		328
quem. (Capromys)	330	390	
querceti. (Sciuropterus v.)	126		109
quercinus. (Sciurus a.)	70	118	
quercinus. (Sciurus w.)	70	118	
quica. (Didelphys)	5	12	
radiata. (Talpa)	467		395
rainieri. (Aplodontia m.)	127		112
rainieri. (Haplodontia r.)	127		112
ramona. (Onychomys)	132	167	121
Rangifer	37		35
Rangifer arcticus	40		37
Rangifer caribou	37		35
Rangifer dawsoni	40		36
Rangifer granti	39		
Rangifer groenlandicus	39		37
Rangifer lapponicus	37		
Rangifer montanus	38		36
Rangifer osborni	38		
Rangifer pearyi	40		
Rangifer stonei	39		
Rangifer stonii	39		
Rangifer terrænovæ	38		36
raptor. (Bassaris)	398		
raptor. (Bassariscus a.)	398		316
rattus. (Mus)	120	163	117
ravus. (Microtus c.)	247		193
rectipinna. (Orca)	26		22
rectipinna. (Orcinus)	26		22
redmani. (Monophyllus)	581	679	
Reduncina	43	68	
regalis. (Vulpes)	382		†432
Reithrodon	191	256	148
Reithrodon longicauda	195	261	
Reithrodon megalotis	193	259	151
Reithrodon mexicanus	197	264	
Reithrodon montanus	193		
Reithrodon sumichrasti	197	264	
Reithrodontomys	191	256	148
Reithrodontomys albescens	192		
Reithrodontomys arizonensis	196		153
Reithrodontomys a. aurantius	198		153
Reithrodontomys a. intermedius	198		153
Reithrodontomys australis	194		
Reithrodontomys a. vulcanius	194	259	

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Reithrodontomys aztecus</i>	193	259	151
<i>Reithrodontomys chrysopsis</i>	199	265	
<i>Reithrodontomys c. tolucae</i>	199	266	
<i>Reithrodontomys chrysotis</i>	195		152
<i>Reithrodontomys colimæ</i>	200	266	
<i>Reithrodontomys c. nerterus</i>	200	267	
<i>Reithrodontomys costaricensis</i>	200	267	
<i>Reithrodontomys c. jalapæ</i>	200	267	
<i>Reithrodontomys creper</i>	204	272	
<i>Reithrodontomys difficilis</i>	197	263	
<i>Reithrodontomys dorsalis</i>	203	272	
<i>Reithrodontomys dychei</i>	192		150
<i>Reithrodontomys d. nebrascensis</i>	192		150
<i>Reithrodontomys goldmani</i>	201	268	
<i>Reithrodontomys griseoflavus</i>	202	270	
<i>Reithrodontomys g. helvolus</i>	202	270	
<i>Reithrodontomys hirsutus</i>	202	269	
<i>Reithrodontomys h. dickinsoni</i>	192		
<i>Reithrodontomys klamathensis</i>	196		152
<i>Reithrodontomys laceyi</i>	199		153
<i>Reithrodontomys lecontei</i>	191		148
<i>Reithrodontomys l. dickinsoni</i>	192		150
<i>Reithrodontomys l. impiger</i>	191		149
<i>Reithrodontomys levipes</i>	203	270	
<i>Reithrodontomys l. otus</i>	203	271	
<i>Reithrodontomys l. toltecus</i>	203	271	
<i>Reithrodontomys longicaudus</i>	195, 196	261	151, 153
<i>Reithrodontomys l. pallidus</i>	195	262	152
<i>Reithrodontomys megalotis</i>	193		151
<i>Reithrodontomys m. deserti</i>	193	259	151
<i>Reithrodontomys m. obscurus</i>	194	261	
<i>Reithrodontomys m. sestinensis</i>	193	260	
<i>Reithrodontomys m. zacatecæ</i>	194	260	
<i>Reithrodontomys merriami</i>	192		150
<i>Reithrodontomys m. aurantius</i>	198		
<i>Reithrodontomys m. fulvescens</i>	198	264	
<i>Reithrodontomys m. gracilis</i>	198	265	
<i>Reithrodontomys m. intermedius</i>	198	264	153
<i>Reithrodontomys microdon</i>	201	269	
<i>Reithrodontomys m. albilabris</i>	202	269	
<i>Reithrodontomys montanus</i>	193		151
<i>Reithrodontomys orizabæ</i>	200	266	
<i>Reithrodontomys pallidus</i>	195	262	
<i>Reithrodontomys perotensis</i>	199	266	
<i>Reithrodontomys rufescens</i>	201	268	
<i>Reithrodontomys saturatus</i>	196		
<i>Reithrodontomys s. alticola</i>	197	263	
<i>Reithrodontomys s. cinereus</i>	196	262	
<i>Reithrodontomys tenuirostris</i>	201	268	
<i>Reithrodontomys t. aureus</i>	201	269	

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Reithrodontomys tenuis</i>	199	265	
<i>Reithronycteris</i>	521	687	
<i>Reithronycteris aphylla</i>	521	688	
<i>repens</i> . (<i>Heteromys</i>)	318	372	
<i>rhabdops</i> . (<i>Oryzomys</i>)	187	237, 246	
<i>Rhachianectes</i>	16	40	9
<i>Rhachianectes glaucus</i>	16, 17	40	9
<i>Rheithrosciurus</i>	59	93	50
<i>Rhinaster</i>	467		394
<i>Rhinodelphis</i>	30	54	
<i>rhinodon</i> . (<i>Beluga</i>)	25		19
<i>Rhinolophus ecaudatus</i>	530	718	
<i>Rhinophoca</i>	440	545	359
<i>Rhinopoma carolinensis</i>			417
<i>Rhinops minor</i>	515	669	
<i>Rhinosciurus</i>	59	93	50
<i>Rhinozolis</i>	410	512	
<i>Rhithrodontomys</i>	191	255	
<i>Rhithrodontomys albescent</i>	192		
<i>Rhithrodontomys arizonensis</i>	196		
<i>Rhithrodontomys australis</i>	194	259	
<i>Rhithrodontomys a. vulcanius</i>	194	259	
<i>Rhithrodontomys catalinae</i>	195		
<i>Rhithrodontomys chrysopsis</i>	199	265, 266	
<i>Rhithrodontomys c. tolucae</i>	199	266	
<i>Rhithrodontomys chrysotis</i>	195		
<i>Rhithrodontomys colimae</i>	200	266, 267	
<i>Rhithrodontomys c. nerterus</i>	200	267	
<i>Rhithrodontomys costaricensis</i>	200	266, 267	
<i>Rhithrodontomys c. jalapae</i>	200	267	
<i>Rhithrodontomys creper</i>	204	272	
<i>Rhithrodontomys difficilis</i>	197	263	
<i>Rhithrodontomys dorsalis</i>	203	272	
<i>Rhithrodontomys dychii</i>	192		
<i>Rhithrodontomys d. nebrascensis</i>	192		
<i>Rhithrodontomys goldmani</i>	201	268	
<i>Rhithrodontomys griseiflavus</i>	202	270	
<i>Rhithrodontomys g. helvolus</i>	202	270	
<i>Rhithrodontomys hirsutus</i>	202	269, 270	
<i>Rhithrodontomys inexpectatus</i>	203	271	
<i>Rhithrodontomys klamathensis</i>	196		
<i>Rhithrodontomys laceyi</i>	199		
<i>Rhithrodontomys lecontii</i>	191		
<i>Rhithrodontomys l. dickinsoni</i>	192		
<i>Rhithrodontomys l. impiger</i>	191		
<i>Rhithrodontomys levipes</i>	203	270, 271	
<i>Rhithrodontomys l. otus</i>	203	271	
<i>Rhithrodontomys l. toltecus</i>	203	271	
<i>Rhithrodontomys longicaudus</i>	195, 196	261, 262	
<i>Rhithrodontomys l. pallidus</i>	195	262	

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Rhithrodontomys megalotis</i>	193	259, 262	
<i>Rhithrodontomys m. obscurus</i>	194	261	
<i>Rhithrodontomys m. sestinensis</i>	193, 194	260	
<i>Rhithrodontomys m. zacatecæ</i>	194	260	
<i>Rhithrodontomys merriami</i>	192		
<i>Rhithrodontomys mexicanus</i>	197	264, 265	
<i>Rhithrodontomys m. aurantius</i>	198		
<i>Rhithrodontomys m. fulvescens</i>	198	264, 265	
<i>Rhithrodontomys m. gracilis</i>	198	265	
<i>Rhithrodontomys m. intermedius</i>	198	264	
<i>Rhithrodontomys microdon</i>	201, 202	269	
<i>Rhithrodontomys m. albilabris</i>	202	269	
<i>Rhithrodontomys montanus</i>	193		
<i>Rhithrodontomys orizabæ</i>	200	266	
<i>Rhithrodontomys peninsulæ</i>	197	263	
<i>Rhithrodontomys perotensis</i>	199	266	
<i>Rhithrodontomys rufescens</i>	201	268	
<i>Rhithrodontomys saturatus</i>	196	262, 263	
<i>Rhithrodontomys s. alticola</i>	197	263	
<i>Rhithrodontomys s. cinereus</i>	196	262	
<i>Rhithrodontomys tenuirostris</i>	201	268, 269	
<i>Rhithrodontomys t. aureus</i>	201	269	
<i>Rhithrodontomys tenuis</i>	199	265	
<i>Rhithronycteris</i>	521	687	
<i>Rhithronycteris aphylla</i>	521	688	
<i>rhoadsi</i> . (<i>Evotomys g.</i>).....	230		173
<i>rhoadsi</i> . (<i>Peromyscus s.</i>).....	139		141
<i>Rhogeëssa</i>	489	600	
<i>Rhogeëssa alleni</i>	489		
<i>Rhogeëssa gracilis</i>	489		
<i>Rhogeëssa parvula</i>	489		
<i>Rhogeëssa tumida</i>	489		
<i>Rhogoessa</i>	489	600	
<i>Rhogoessa alleni</i>	489	602	
<i>Rhogoessa gracilis</i>	489	602	
<i>Rhogoessa parvula</i>	489	601	
<i>Rhogoessa tumida</i>	489	601	
<i>rhydinorhis</i> . (<i>Perognathus b.</i>).....	309	362	
<i>Rhynchonycteris</i>	492	608	
<i>Rhynchonycteris naso</i>	492	609	
<i>Rhytina</i>	14		5
<i>richardi</i> . (<i>Phoca</i>).....	442		
<i>richardii</i> . (<i>Halicyon</i>).....	442		
<i>richardii</i> . (<i>Phoca</i>).....	442		
<i>richardsoni</i> . (<i>Arvicola</i>).....	250		200
<i>richardsoni</i> . (<i>Cervus</i>).....	48		41
<i>richardsoni</i> . (<i>Citellus</i>).....	111		
<i>richardsoni</i> . (<i>Dicrostonyx</i>).....	261		211
<i>richardsoni</i> . (<i>Dicrostonyx h.</i>).....	261		211
<i>richardsoni</i> . (<i>Dipodops</i>).....	297		238

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
richardsoni. (Lepus)	363	437	
richardsoni. (Lepus c.)	363		
richardsoni. (Microtus)	250		200
richardsoni. (Mustela)	426		344
richardsoni. (Perodipus)	297		238
richardsoni. (Putorius c.)	426, 427		344
richardsoni. (Sciurus)	76	128	
richardsoni. (Sciurus h.)	82		64
richardsoni. (Sorex)	446, 447		367, 368
richardsoni. (Spermophilus)	111		92
richardsoni. (Ursus)	393		313
richardsonii. (Arctomys)	111		92
richardsonii. (Mustela)	426		344
richardsonii. (Sciurus)	82		
richardsonii. (Sorex)	447		
richmondi. (Didelphis)	8	16	
richmondi. (Didelphys)	8	16	
richmondi. (Orzomys)	184	248	
richmondi. (Sciurus)	62	105	
rigidus. (Lepus f.)	345	419	
rigidus. (Sciurus)	76	127	
Rigoon	444	542	
ringens. (Spilogale)	413		329
ringens. (Tagassu a.)	37	66	
ringens. (Tagassu p.)	37	66	
riparius. (Arvicola)	234		179
ripensis. (Fiber z.)	252		
rissoanus. (Delphinus)	28		25
rivalicius. (Fiber z.)	253		213
rivalis. (Proboscidea)	492	609	
rivularis. (Microtus m.)	239		183
rivularis. (Microtus n.)	239		
rixosus. (Putorius)	428		345
roanensis. (Zapus i.)	328		263
robustus. (Peromyscus)	156		138
robustus. (Sitomys)	156		138
Rodentia	59	89	50
Romerolagus	339	411	
Romerolagus nelsoni	339	412	
roosevelti. (Cervus)	42		34
roqual. (Balænoptera)	19		13
Rorqualus minor	18		11
Rosmarus arcticus	439		358
Rosmarus cookii	439		358
rosmarus. (Odontobænus)	439		
rosmarus. (Phoca)	439		
rosmarus. (Trichechus)	439		358
rostrata. (Balæna)	21		11, 16
rostratus. (Heteromys p.)	322	378	
rostratus. (Hyperoodon)	21		16

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
rostratus. (Liomys p.).....	322	378	
rostratus. (Oryzomys).....	188	243, 244	
rostratus. (Perognathus p.).....	312	359	
rothschildi. (Coendou).....	334	401	
rothschildi. (Coendu).....	334	401	
rothschildi. (Dama).....	46	72	
rothschildi. (Odontocœlus).....	46	72	
rotundatus. (Nyctiplanus).....	529	715	
rotundum. (Desmodon).....	530		
rotundum. (Phyllostoma).....	530		
rotundus. (Desmodus).....	530	718	
rowleyi. (Peromyscus).....	151		134
rowleyi. (Sitomys).....	151		134
ruatanica. (Dasyprocta).....	336	405	
rubellus. (Vespertilio).....	485	593	411
rubida. (Neotoma c.).....	210		156
rubida. (Neotoma f.).....	210		
rubidus. (Peromyscus o.).....	153		
rubiginosa. (Chilonycteris).....	505	643	
rubra. (Vespertilio).....	485	593	417
rubricaudatus. (Sciurus).....			52
rubricosa. (Vulpes f.).....	380		
rubricosa. (Vulpes p.).....	380		305
rufa. (Anisonyx).....	126		112
rufa. (Felis).....	370		296
rufa. (Haplodontia).....	126, 127		112
rufescens. (Arvicola).....	235		180
rufescens. (Reithrodontomys).....	201	268	
rufescens. (Rhithrodontomys).....	201	268	
rufescens. (Thomomys).....	279	332	221, 227
rufescens. (Thomomys t.).....	279		227
ruffa. (Felis).....	370		
rufidorsum. (Arvicola).....	235		180
rufinus. (Cervus).....		79	
rufinus. (Hesperomys l.).....	142		126
rufinus. (Oryzomys j.).....	185	247	
rufinus. (Peromyscus).....	142		126
rufipes. (Aotus).....	533	729	
rufipes. (Lepus l.).....	346		
rufipes. (Nyctipithecus).....	533	729	
rufipes. (Sciurus v.).....	69	117	
rufiventer. (Sciurus).....	64, 68	115	52
rufiventris. (Ateles).....	535	734	
rufiventris. (Sciurus).....	69		
rufoniger. (Sciurus).....	61	104	
rufum. (Stenoderma).....	526	704	
rufus. (Desmodus).....	530	718	
rufus. (Molossus).....	495	619	
rufus. (Oryzomys).....	185	239	
rufus. (Vespertilio).....	485	593	412

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
ruidosæ. (Onychomys)	131		
rupestris. (Citellus g.)	114		
rupestris. (Citellus v.)	114	150	
Rupicaprinæ	52		44
rupicola. (Neotoma)	223		166
rupicola. (Teonoma)	223		
russatus. (Lepus)	348	420	
rutilus. (Evotomys)	229		170
rutilus. (Mus)	229		170
Rytina	14		
sabrinus. (Sciuropterus)	124		107
sabrinus. (Sciurus)	124		107
saccatus. (Mus)	262		217
Sacomys	316	368	
Saccophorus	261	310	215
Saccophorus bursarius	262		
Saccopteryx	492	610	
Saccopteryx bilineata	493	610	
Saccopteryx infusca	493	611	
sagax. (Peromyscus)	158	194	
sagittalis. (Geomys b.)	263		218
Saimiri	534	730	
Saimiri ærstedii	534	731	
Saimiris entomophaga	534	731	
saliens. (Lepus)	362		276
saltator. (Zapus)	327		262
salvini. (Chiroderma)	528	710	
salvini. (Heteromys)	317	370	
salvini. (Sorex)	455	551	
sanctidiegi. (Lepus)	345, 346	422	283
sanctidiegi. (Lepus a.)	346		
sanctidiegi. (Lepus f.)	345		
santacruzæ. (Urocyon c.)	388		
santacruzæ. (Urocyon l.)	388		
sarcovienna. (Lutra)	1	3	
sartori. (Hippocamelus)	50		
sartori. (Mazama)	50	80	
sartorii. (Cervus)	50	80	
saturatior. (Oryzomys c.)	186	239	
saturatus. (Citellus l.)	106		
saturatus. (Evotomys g.)	231		174
saturatus. (Myotis y.)	475	577	403
saturatus. (Peromyscus t.)	145		131
saturatus. (Putorius)	430		348
saturatus. (Reithrodontomys)	196		
saturatus. (Rhithrodontomys)	196	262, 263	
saturatus. (Sigmodon h.)	177	225	
saturatus. (Spermophilus l.)	106		83
saturatus. (Tamias l.)	106		83

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
saussurei. (Sorex).....	456		
saussurii. (Sorex).....	456	553	
saxamans. (Neotoma).....	221		†429
saxamans. (Teonoma).....	221		
saxatilis. (Ochotona).....	339		268
saxatilis. (Peromyscus m.).....	169	207	
saxatilis. (Proboscidea).....	492	609	
saxatilis. (Spilogale).....	414		330
saxicola. (Bassariscus).....	398	485	
saxicola. (Citellus h.).....	97	142	
saxicolus. (Spermophilus h.).....	97	142	86
sayi. (Sciurus).....	65		52
Scalops.....	469		388
Scalops æneus.....	468		391
Scalops aereus.....	471		390
Scalops anastasie.....	490		391
Scalops aquaticus.....	470		388
Scalops a. australis.....	470		390
Scalops a. machrinus.....	470		389
Scalops a. parvus.....	471		390
Scalops a. texensis.....	471		390
Scalops argentatus.....	470		389
Scalops a. texanus.....	471		
Scalops breweri.....	471		394
Scalops californicus.....	468		392
Scalops canadensis.....	468, 470		388, 391
Scalops latimanus.....	468		391
Scalops m. intermedius.....	470		
Scalops metallescens.....	468		391
Scalops parvus.....	470		390
Scalops pennsylvanica.....	470		388
Scalops t. aereus.....	471		
Scalops townsendii.....	468		391
scalops. (Geomys).....	269	324	
scalops. (Orthogeomys).....	269	324	
scalopsoides. (Arvicola).....	249		198
scalopsoides. (Microtus p.).....	249		198
Scalopus.....	469		
Scalopus aereus.....	471		
Scalopus aquaticus.....	469		
Scalopus a. australis.....	470		
Scalopus a. intermedius.....	470		
Scalopus a. machrinus.....	470		
Scalopus a. texensis.....	471		
Scalopus virginianus.....	469		
scammoni. (Globicephalus).....	28	54	
scammoni. (Globiocephalus).....	28	54	24
Scapanus.....	468	564	391
Scapanus alpinus.....	469		393
Scapanus anthonyi.....	469	564	393

	Check-List of Mammals.	Mammals. of Middle America, etc.	Synopsis of Mammals, etc.
<i>Scapanus californicus</i>	468		392
<i>Scapanus c. minusculus</i>	469		†435
<i>Scapanus dilatus</i>	468		392
<i>Scapanus orarius</i>	468		392
<i>Scapanus townsendi</i>	468		391
<i>Scapanus truei</i>	469		393
<i>Scapanus truii</i>	469		
<i>scaphiotus</i> . (<i>Odocoileus c.</i>).....	48		42
<i>scaphiotus</i> . (<i>Odontocælus c.</i>).....	48		
<i>scapterus</i> . (<i>Thomomys</i>).....	274		
<i>schisticeps</i> . (<i>Lagomys</i>).....	338		267
<i>schisticeps</i> . (<i>Ochotona</i>).....	338		267
<i>Schizostoma</i>	512	660	
<i>Schizostoma hirsutum</i>	512	661	
<i>schlegelii</i> . (<i>Orca</i>).....	26	51	22
<i>schreberi</i> . (<i>Phoca</i>).....	444		364
<i>scirpensis</i> . (<i>Microtus</i>).....	241		187
<i>scitulus</i> . (<i>Peromyscus c.</i>).....	159		
<i>scitulus</i> . (<i>Peromyscus t.</i>).....	159		†429
<i>sciurea</i> . (<i>Chrysothrix</i>).....	534	731	
<i>sciurea</i> . (<i>Simia</i>).....	534	730	
<i>Sciuridæ</i>	59	90	50
<i>Sciurinæ</i>	59	90	50
<i>Sciuropterus</i>	121	157	107
<i>Sciuropterus alpinus</i>	121, 122		
<i>Sciuropterus a. bangsi</i>	124		
<i>Sciuropterus a. californicus</i>	122		
<i>Sciuropterus a. fuliginosus</i>	123		
<i>Sciuropterus a. klamathensis</i>	123		
<i>Sciuropterus a. lascivus</i>	123		†428
<i>Sciuropterus a. olympicus</i>	123		
<i>Sciuropterus a. oregonensis</i>	122		
<i>Sciuropterus a. stephensi</i>	124		
<i>Sciuropterus a. zaphæus</i>	122		
<i>Sciuropterus o. stephensi</i>	124		
<i>Sciuropterus sabrinus</i>	124, 125		107
<i>Sciuropterus s. macrotis</i>	124		108
<i>Sciuropterus s. makkovikensis</i>	125		
<i>Sciuropterus s. silus</i>	125		108
<i>Sciuropterus silus</i>	125		
<i>Sciuropterus stephensi</i>	124		111
<i>Sciuropterus volans</i>	125	158	109
<i>Sciuropterus v. alpinus</i>	122		109
<i>Sciuropterus v. californicus</i>	122		110
<i>Sciuropterus v. fuliginosus</i>	123		111
<i>Sciuropterus v. goldmani</i>	126		
<i>Sciuropterus v. klamathensis</i>	123		111
<i>Sciuropterus v. olympicus</i>	123		110
<i>Sciuropterus v. oregonensis</i>	122		110

†Appendix.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Sciuropterus v. querceti</i>	126		109
<i>Sciuropterus yukonensis</i>	125		†428
<i>Sciurus</i>	59	93	50
<i>Sciurus aberti</i>	66, 67	112	57
<i>Sciurus a. barberi</i>	67		
<i>Sciurus a. concolor</i>	66		57
<i>Sciurus a. durangi</i>	67		
<i>Sciurus a. ferreus</i>	66		
<i>Sciurus a. mimus</i>	66		
<i>Sciurus acrobates</i>	124		107
<i>Sciurus adolpheï</i>	75	126	
<i>Sciurus a. dorsalis</i>	76	127	
<i>Sciurus adolphii</i>	75		
<i>Sciurus a. dorsalis</i>	76		
<i>Sciurus æstuans</i>		104	
<i>Sciurus æ. chiriquensis</i>	62	104	
<i>Sciurus æ. hoffmanni</i>	61	104	
<i>Sciurus affinis</i>	74	124	
<i>Sciurus albipes</i>	69	117	
<i>Sciurus a. colimensis</i>	71, 72	129	
<i>Sciurus a. effugius</i>	72	120	
<i>Sciurus a. hernandezi</i>	70	118	
<i>Sciurus a. nemoralis</i>	70	118	
<i>Sciurus a. quercinus</i>	70	118	
<i>Sciurus alfari</i>	60	99	
<i>Sciurus alleni</i>	63	108	
<i>Sciurus alstoni</i>	63	108	
<i>Sciurus americanus</i>	95		80
<i>Sciurus apache</i>	64	110	58
<i>Sciurus arizonensis</i>	61, 63	102, 109	59
<i>Sciurus a. huachuca</i>	64	109	60
<i>Sciurus auduboni</i>	65		52
<i>Sciurus aureigaster</i>	68, 74	115, 124	
<i>Sciurus a. frumentor</i>	69	117	
<i>Sciurus a. hypopyrrhus</i>	68, 69	116	
<i>Sciurus aureogaster</i>	69, 74	116	
<i>Sciurus a. frumentor</i>	69		
<i>Sciurus a. leucops</i>	71	119	
<i>Sciurus aurogaster</i>	68	115	
<i>Sciurus barberi</i>	67	†741	
<i>Sciurus belcheri</i>	82		65
<i>Sciurus boothiæ</i>	75	126, 128	
<i>Sciurus b. belti</i>	77	128	
<i>Sciurus b. managuensis</i>	77	129	
<i>Sciurus boquetensis</i>	60	100	
<i>Sciurus browni</i>	60	100	
<i>Sciurus californicus</i>	114		
<i>Sciurus capistratus</i>	62, 66	107	51
<i>Sciurus carolinensis</i>	63, 75	108, 125	53, 61

†Appendix.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Sciurus c. carolinensis</i>	63		
<i>Sciurus c. extimus</i>	80		55
<i>Sciurus c. fuliginosus</i>	79		55
<i>Sciurus c. hypophæus</i>	79		54
<i>Sciurus c. leucotis</i>	79		53
<i>Sciurus c. yucatanensis</i>	75	125	
<i>Sciurus castanonotus</i>			57
<i>Sciurus cervicalis</i>	71		
<i>Sciurus cinereus</i>	65, 66		51, 52
<i>Sciurus colliæi</i>	73, 77	121, 129	60
<i>Sciurus c. nuchalis</i>	73	122	
<i>Sciurus deppei</i>	61	102	
<i>Sciurus d. vivax</i>	61		
<i>Sciurus deppii</i>	60	101	
<i>Sciurus d. vivax</i>	61	102	
<i>Sciurus dorsalis</i>	66, 76	127	57
<i>Sciurus douglasi</i>	82		65
<i>Sciurus d. albolimbatus</i>	83		66
<i>Sciurus d. cascadenis</i>	83		66
<i>Sciurus d. mearnsi</i>	83	133	
<i>Sciurus d. mollipilosus</i>	83		65
<i>Sciurus douglasii</i>	82		
<i>Sciurus durangi</i>	67	112	
<i>Sciurus d. phæurus</i>	67	†741	
<i>Sciurus ferruginiventris</i>	68	115	
<i>Sciurus fossor</i>	78	130	55
<i>Sciurus f. anthonyi</i>	78	130	
<i>Sciurus f. nigripes</i>	78		
<i>Sciurus fremonti</i>	84		66
<i>Sciurus f. grahamensis</i>	85		67
<i>Sciurus f. lychnuchus</i>	84		
<i>Sciurus f. mogollonensis</i>	84		67
<i>Sciurus f. neomexicanus</i>	84		67
<i>Sciurus fuliginosus</i>	79		
<i>Sciurus fuscovariegatus</i>	76	128	
<i>Sciurus goldmani</i>	77	130	
<i>Sciurus grammurus</i>	113	149	88
<i>Sciurus griseiflavus</i>	74	124	
<i>Sciurus g. chiapensis</i>	74	125	
<i>Sciurus griseocaudatus</i>	77		
<i>Sciurus griseoflavus</i>	64	110	58
<i>Sciurus g. chiapensis</i>	74		
<i>Sciurus griseogena</i>	60, 61	101, 104	
<i>Sciurus griseogenys</i>	61	104	
<i>Sciurus griseus</i>	78	130	55
<i>Sciurus g. anthonyi</i>	78		57
<i>Sciurus g. nigripes</i>	78		56
<i>Sciurus heermanni</i>	78	130	55
<i>Sciurus hiemalis</i>	79		53

†Appendix.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Sciurus hudsonicus baileyi</i>	81		
<i>Sciurus h. dakotensis</i>	81		
<i>Sciurus h. grahamensis</i>	85		
<i>Sciurus h. gymnicus</i>	80		
<i>Sciurus h. loquax</i>	80		
<i>Sciurus h. minnesota</i>	81		
<i>Sciurus h. petulans</i>	82		
<i>Sciurus h. streatori</i>	82		
<i>Sciurus h. ventorum</i>	81		
<i>Sciurus hudsonius</i>	80, 124		60, 107
<i>Sciurus h. baileyi</i>	81		63
<i>Sciurus h. californicus</i>	83		66
<i>Sciurus h. dakotensis</i>	81		62
<i>Sciurus h. grahamensis</i>	85		
<i>Sciurus h. gymnicus</i>	80		61
<i>Sciurus h. loquax</i>	80		62
<i>Sciurus h. mearnsi</i>	83		
<i>Sciurus h. minnesota</i>	81		62
<i>Sciurus h. petulans</i>	82		†426
<i>Sciurus h. richardsoni</i>	82		64
<i>Sciurus h. streatori</i>	82		64
<i>Sciurus h. vancouverensis</i>	82		64
<i>Sciurus h. ventorum</i>	81		63
<i>Sciurus hypopyrrhus</i>	62, 68, 73	121, 126	
.....	75, 76, 77	127, 128	
<i>Sciurus hypoxanthus</i>	68	115	
<i>Sciurus intermedius</i>	76	127	
<i>Sciurus kaibabensis</i>	67		
<i>Sciurus lateralis</i>	105		83
<i>Sciurus leporinus</i>	78	130	55
<i>Sciurus leucogaster</i>	68	115	
<i>Sciurus leucops</i>	71, 74	119, 123	
<i>Sciurus leucotis</i>	79		
<i>Sciurus ludovicianus</i>	64, 74	124	52
<i>Sciurus l. limitis</i>	65	110	53
<i>Sciurus l. neglectus</i>	65		
<i>Sciurus l. vicinus</i>	65		52
<i>Sciurus lysteri</i>	95		
<i>Sciurus macroura</i>	64		52
<i>Sciurus magnicaudatus</i>	65		52
<i>Sciurus managuensis</i>	77	129	
<i>Sciurus melania</i>	76	128	
<i>Sciurus mexicanus</i>	103	146	
<i>Sciurus migratorius</i>	79		54
<i>Sciurus mollipilosus</i>	83		
<i>Sciurus mustelinus</i>	68	115	
<i>Sciurus nayaritensis</i>	63	108	
<i>Sciurus negligens</i>	61	102	
<i>Sciurus nelsoni</i>	72	120	

*Appendix.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Sciurus n. hirtus</i>	72	121	
<i>Sciurus nicoyana</i>	76	127	
<i>Sciurus niger</i>	66, 79	115	51, 53
<i>Sciurus n. ludovicianus</i>	64	110	
<i>Sciurus n. melanonotus</i>	62	107	
<i>Sciurus occidentalis</i>	65		52
<i>Sciurus oculatus</i>	62	107	
<i>Sciurus o. toluæ</i>	63	107	
<i>Sciurus orarius</i>			65
<i>Sciurus pennsylvanicus</i>	79		53
<i>Sciurus poliopus</i>	69	117	
<i>Sciurus p. cervicalis</i>	71	119	
<i>Sciurus p. colimensis</i>	71, 72	119	
<i>Sciurus p. effugius</i>	72	120	
<i>Sciurus p. hernandezi</i>	70	118	
<i>Sciurus p. nemoralis</i>	70	118	
<i>Sciurus p. peregrinator</i>	70		
<i>Sciurus p. senex</i>	71		
<i>Sciurus pyladei</i>	77	129	
<i>Sciurus quadrivittatus</i>	91	138	
<i>Sciurus richardsoni</i>	76	128	
<i>Sciurus richardsonii</i>	82		
<i>Sciurus richmondi</i>	62	105	
<i>Sciurus rigidus</i>	76	127	
<i>Sciurus rubricaudatus</i>			52
<i>Sciurus rufiventer</i>	64, 68	115	52
<i>Sciurus r. neglectus</i>	65		
<i>Sciurus r. texensis</i>	65	110	
<i>Sciurus r. texianus</i>	65		
<i>Sciurus rufiventris</i>	69		
<i>Sciurus rufoniger</i>	61	104	
<i>Sciurus sabrinus</i>	124		107
<i>Sciurus sayi</i>	65		52
<i>Sciurus sinaloensis</i>	73	122	
<i>Sciurus socialis</i>	73	123	
<i>Sciurus s. cocos</i>	74	124	
<i>Sciurus striatus</i>	85, 95	134	
<i>Sciurus subauratus</i>	65		52
<i>Sciurus suckleyi</i>	82		65
<i>Sciurus tæniurus</i>	60	101	
<i>Sciurus tephrogaster</i>	60	101	
<i>Sciurus texianus</i>	65		
<i>Sciurus thomasi</i>	75	126	
<i>Sciurus tridecemlineatus</i>	104		
<i>Sciurus truei</i>	73		
<i>Sciurus truei</i>	73	123	
<i>Sciurus variabilis</i>	62	105	
<i>Sciurus v. morulus</i>	62	105	
<i>Sciurus variegatoides</i>	77	129	
<i>Sciurus variegatus</i>	69, 113	116, 148	

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Sciurus v. poliopus</i>	69		
<i>Sciurus v. rufipes</i>	69	117	
<i>Sciurus varius</i>	69	117	
<i>Sciurus volans</i>	121	157	107
<i>Sciurus volucella</i>	125	158	109
<i>Sciurus vulgaris</i>	59		
<i>Sciurus v. hudsonicus</i>	80		
<i>Sciurus vulpinus</i>	66, 79		51, 53
<i>Sciurus wagneri</i>	69	117	
<i>Sciurus w. cervicalis</i>	71	119	
<i>Sciurus w. colimensis</i>	72	120	
<i>Sciurus w. effugius</i>	72	120	
<i>Sciurus w. nemoralis</i>	70	118	
<i>Sciurus w. quercinus</i>	70	118	
<i>Sciurus xanthotus</i>	61	104	
<i>Sciurus yucatanensis</i>	75	125	
<i>Sciurus y. baliolus</i>	75	126	
<i>sclateri</i> . (<i>Sorex</i>).....	456	552	
<i>scopulicola</i> . (<i>Phoca</i>).....	442		363
<i>Scotophilus</i>	479		408
<i>Scotophilus cubensis</i>	484	588	
<i>Scotophilus greenii</i>	483	587	410
<i>Scotophilus hesperus</i>	480	582	409
<i>Scotophilus miradorensis</i>	484	588	
<i>scotti</i> . (<i>Urocyon c.</i>).....	386	477	308
<i>scrobiculatum</i> . (<i>Phyllostoma</i>).....	513	662	
<i>scrutator</i> . (<i>Mephitis m.</i>).....	407		324
<i>sellata</i> . (<i>Myrmecophaga</i>).....	11	28	
<i>sellata</i> . (<i>Tamandua</i>).....	11	28	
<i>Selysius</i>	473	570	400
<i>semilunaris</i> . (<i>Phoca</i>).....	444		364
<i>seminola</i> . (<i>Atalapha b.</i>).....	486		
<i>seminolus</i> . (<i>Lasiurus b.</i>).....	486		413
<i>semispinosus</i> . (<i>Echimys</i>).....	329	386	
<i>semispinosus</i> . (<i>Proechinomys</i>).....	329	386	
<i>senegalensis</i> . (<i>Trichechus</i>).....		35	
<i>senex</i> . (<i>Centurio</i>).....	529	717	
<i>senex</i> . (<i>Galera b.</i>).....	418		
<i>senex</i> . (<i>Galictis b.</i>).....	418	525	
<i>senex</i> . (<i>Grison b.</i>).....	418	525	
<i>senex</i> . (<i>Sciurus p.</i>).....	71		
<i>senex</i> . (<i>Tamias</i>).....	89		72
<i>senicula</i> . (<i>Mormoops m.</i>).....	507	648	
<i>senicula</i> . (<i>Mormops m.</i>).....	507	648	
<i>Seniocebus</i>	532	724	
<i>sennetti</i> . (<i>Dipodops</i>).....	297		239
<i>sennetti</i> . (<i>Perodipus</i>).....	297		239
<i>seorsus</i> . (<i>Zygodontomys</i>).....	191	254	
<i>septentrionalis</i> . (<i>Vespertilio g.</i>).....	478		405

*Appendix.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
sericea. (Talpa).....	470		389
serpens. (Microtus).....	250		200
sestinensis. (Reithrodontomys m.).....	193	260	
sestinensis. (Rhithrodontomys m.).....	193, 194	260	
setosus. (Heteromys c.).....	320	380	
setosus. (Liomys c.).....	320	380	
setosus. (Sorex).....	448		369
sezekorni. (Phyllonycteris).....	521	686	
shastensis. (Sorex).....	452		374
shattucki. (Microtus p.).....	235		
shumaginensis. (Sorex a.).....	451		
shumaginensis. (Sorex g.).....	451		373
sibbaldii. (Physalus).....	20		13
Sibbaldius.....	18	42	11
Sibbaldius sulfureus.....	20	43	
Sibbaldius tectirostris.....	20		13
Sibbaldus.....	18	42	11
sieboldii. (Bakena).....	15		8
sierræ. (Lepus c.).....	363		
Sigmodon.....	174	222	143
Sigmodon alleni.....	179	224	
Sigmodon alticola.....	179	231	
Sigmodon a. amoles.....	179	231	
Sigmodon austerulus.....	180	231	
Sigmodon baileyi.....	177	226	
Sigmodon berlandieri.....	175	227	
Sigmodon borucæ.....	178	224	
Sigmodon b. chiriquensis.....	178	225	
Sigmodon colimæ.....	176	227	
Sigmodon fulviverter.....	180	232	
Sigmodon hispidus.....	174	228	144
Sigmodon h. arizonæ.....	176	228	144
Sigmodon h. baileyi.....	177	226	
Sigmodon h. berlandieri.....	175	225, 227	
Sigmodon h. borucæ.....	178	224, 232	
Sigmodon h. chiriquensis.....	178	225	
Sigmodon h. eremicus.....	176	227	145
Sigmodon h. furvus.....	178	225	
Sigmodon h. inexoratus.....	177	229	
Sigmodon h. littoralis.....	175		144
Sigmodon h. major.....	177	226, 228	
Sigmodon h. mascotensis.....	176	227	
Sigmodon h. microdon.....	178	226	
Sigmodon h. pallidus.....	175	228	145
Sigmodon h. saturatus.....	177	225	
Sigmodon h. spadicipygus.....	175		145
Sigmodon h. texensis.....	175	228	144
Sigmodon h. toltecus.....	177	226	
Sigmodon h. tonalensis.....	176	229	
Sigmodon leucotis.....	179	230, 231	

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Sigmodon mascotensis</i>	176	227, 229	
<i>Sigmodon melanotis</i>	180	232	
<i>Sigmodon minimus</i>	178	230	146
<i>Sigmodon ochrognathus</i>	179	230	
<i>Sigmodon spadicipygus</i>	175		145
<i>Sigmodontomys</i>	191	254	
<i>Sigmodontomys alfari</i>	191	254, 255	
<i>silus</i> . (<i>Sciuropterus</i>).....	121		
<i>silus</i> . (<i>Sciuropterus s.</i>).....	125		108
<i>Simia belzebul</i>	532	726	
<i>Simia hypoleuca</i>	536	736	
<i>Simia œdipus</i>	532		
<i>Simia paniscus</i>	534	732	
<i>Simia sciurea</i>	534	730	
<i>Simia trivirgata</i>	533	728	
<i>similis</i> . (<i>Clymene</i>).....	30		27
<i>similis</i> . (<i>Dipodomys</i>).....	289		234
<i>similis</i> . (<i>Dipodomys m.</i>).....	289		234
<i>similis</i> . (<i>Sorex v.</i>).....	449		372
<i>simiolus</i> . (<i>Dipodomys</i>).....	289		234
<i>simiolus</i> . (<i>Dipodomys m.</i>).....	289, 290		234
<i>simplex</i> . (<i>Neotoma m.</i>).....	206		159
<i>simplicicanus</i> . (<i>Lepus</i>).....	343		
<i>simplicicanus</i> . (<i>Lepus f.</i>).....	343		
<i>simulans</i> . (<i>Perodipus s.</i>).....	295		
<i>simulatus</i> . (<i>Peromyscus</i>).....	167	205	
<i>simulus</i> . (<i>Peromyscus s.</i>).....	157	180	
<i>sinaloæ</i> . (<i>Marmosa</i>).....	3	6	
<i>sinaloæ</i> . (<i>Neotoma</i>).....	213	283	
<i>sinaloæ</i> . (<i>Odocoileus</i>).....	47	78	
<i>sinaloæ</i> . (<i>Odontocœlus</i>).....	47	78	
<i>sinaloæ</i> . (<i>Thomomys</i>).....	287	339	
<i>sinaloensis</i> . (<i>Sciurus</i>).....	73	122	
<i>Sinetheres</i>	334	399	
<i>Sirenia</i>	14	35	5
<i>sitkensis</i> . (<i>Microtus</i>).....	242		189
<i>sitkensis</i> . (<i>Odocoileus c.</i>).....	48		42
<i>sitkensis</i> . (<i>Odontocœlus c.</i>).....	48		
<i>sitkensis</i> . (<i>Peromyscus</i>).....	152		134
<i>sitkensis</i> . (<i>Ursus</i>).....	391		
<i>Sitomys</i>	135		
<i>Sitomys a. arizonæ</i>	140		125
<i>Sitomys a. artemisæ</i>	141		126
<i>Sitomys a. canadensis</i>	144		128
<i>Sitomys a. thurberi</i>	146		
<i>Sitomys auripectus</i>	153		134
<i>Sitomys decolorus</i>	172	216	
<i>Sitomys gilberti</i>	150	195	132
<i>Sitomys herronii</i>	155		137
<i>Sitomys h. nigellus</i>	155		137

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Sitomys insolatus</i>	154		135
<i>Sitomys keeni</i>	151		133
<i>Sitomys macroshinus</i>	152		133
<i>Sitomys major</i>	155		137
<i>Sitomys martirensis</i>	159	189	
<i>Sitomys megacephalus</i>			127
<i>Sitomys musculus</i>	135	175	
<i>Sitomys robustus</i>	156		138
<i>Sitomys rowleyi</i>	151		134
<i>Sitomys r. pinalis</i>	151	189	135
<i>Sitomys subgriseus</i>	139		141
<i>Sminthus</i>	323		
<i>socialis</i> . (<i>Cynomys</i>)	117		102
<i>socialis</i> . (<i>Sciurus</i>)	73	123	
<i>sola</i> . (<i>Neotoma d.</i>)	215		161
<i>Solenodon</i>	472	567	
<i>Solenodon cubanus</i>	472	567	
<i>Solenodon paradoxus</i>	472	568	
<i>Solenodontidæ</i>	472	566	
<i>Solitaria</i> . (<i>Neotoma f.</i>)	211		
<i>sonora</i> . (<i>Lutra c.</i>)	434	535	353
<i>sonora</i> . (<i>Lutra h.</i>)	434		
<i>sonorana</i> . (<i>Heteromys</i>)	323	379	
<i>sonorana</i> . (<i>Liomys</i>)	323	379	
<i>sonoriense</i> . (<i>Tagassu a.</i>)	36	64	
<i>sonoriensis</i> . (<i>Citellus t.</i>)	102	144	
<i>sonoriensis</i> . (<i>Conepatus</i>)	411	514	
<i>sonoriensis</i> . (<i>Dicotyles a.</i>)	36	64	
<i>sonoriensis</i> . (<i>Hesperomys</i>)	141	181	125
<i>sonoriensis</i> . (<i>Peromyscus</i>)	141		
<i>sonoriensis</i> . (<i>Peromyscus a.</i>)	141	182	125
<i>sonoriensis</i> . (<i>Peromyscus l.</i>)	141	181	
<i>sonoriensis</i> . (<i>Spermophilus</i>)	102		
<i>sonoriensis</i> . (<i>Spermophilus t.</i>)	102	144	
<i>Sorex</i>	446	548	366
<i>Sorex a. shumaginensis</i>	451		
<i>Sorex alnorum</i>	457		
<i>Sorex amœnus</i>	449		371
<i>Sorex angusticeps</i>	461		382
<i>Sorex aquaticus</i>	469		388
<i>Sorex araneus</i>	446	549	366
<i>Sorex bairdi</i>	451		373
<i>Sorex belli</i>			368
<i>Sorex bendirii albiventer</i>	459		381
<i>Sorex b. palmeri</i>	459		381
<i>Sorex brevicandus</i>	460		382
<i>Sorex californicus</i>	452		374
<i>Sorex carolinensis</i>	461		
<i>Sorex caudatus</i>	456	552	
<i>Sorex cinereus</i>	463		384

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Sorex cooperi</i>	446		366
<i>Sorex crawfordi</i>	459	554	385
<i>Sorex cristatus</i>	467		395
<i>Sorex dekayi</i>	461		382
<i>Sorex dobsoni</i>	448		370
<i>Sorex evotis</i>	460	555	
<i>Sorex eximius</i>	457		
<i>Sorex fimbripes</i>	446		366
<i>Sorex fisheri</i>	454		375
<i>Sorex forsteri</i>	446		366
<i>Sorex fumeus</i>	447		369
<i>Sorex glacialis</i>	451		372
<i>Sorex g. alascensis</i>	451		372
<i>Sorex g. shumaginensis</i>	451		373
<i>Sorex godmani</i>	455	552	
<i>Sorex haydeni</i>	446		367
<i>Sorex hoyi</i>	456, 457		377
<i>Sorex hydrodromus</i>	458		379
<i>Sorex idahoensis</i>	446		367
<i>Sorex l. elassodon</i>	450		
<i>Sorex l. prevostensis</i>	450		
<i>Sorex longirostris</i>	453		375
<i>Sorex macrodon</i>	455	552	
<i>Sorex macrurus</i>	448		369
<i>Sorex merriami</i>	454		376
<i>Sorex m. miscix</i>	454		376
<i>Sorex montereyensis</i>	452		373
<i>Sorex monticolus</i>	448		371
<i>Sorex n. alaskanus</i>	458		
<i>Sorex nevadensis</i>	449		371
<i>Sorex obscurus</i>	449		371
<i>Sorex o. alascensis</i>	451		
<i>Sorex o. elassodon</i>	450		
<i>Sorex o. longicauda</i>	450		
<i>Sorex o. longicaudus</i>	450, 451		372
<i>Sorex o. prevostensis</i>	450		
<i>Sorex o. ventralis</i>	450	550	
<i>Sorex oreopolus</i>	451	550	
<i>Sorex orinus</i>	456	553	
<i>Sorex orizabæ</i>	449	550	
<i>Sorex ornatus</i>	452		374
<i>Sorex pacificus</i>	454		375
<i>Sorex palustris</i>	457, 458		379
<i>Sorex parvus</i>	449, 463		368, 384
<i>Sorex personatus</i>	446		366
<i>Sorex p. arcticus</i>	447		368
<i>Sorex p. miscix</i>	454		
<i>Sorex p. streator</i>	446		367
<i>Sorex platyrhinus</i>	447		369
<i>Sorex platyrhynchus</i>	446		366

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Sorex pribilofensis</i>	454		376
<i>Sorex richardsoni</i>	446, 447		367, 368
<i>Sorex salyini</i>	455	551	
<i>Sorex saussurei</i>	456		
<i>Sorex s. mutabilis</i>	456		
<i>Sorex saussurii</i>	456	553	
<i>Sorex s. mutabilis</i>	456	553	
<i>Sorex sclateri</i>	456	552	
<i>Sorex setosus</i>	448		369
<i>Sorex shastensis</i>	452		374
<i>Sorex sphagnicola</i>	447		368
<i>Sorex stizodon</i>	455	551	
<i>Sorex suckleyi</i>	448		370
<i>Sorex talpoides</i>	460	555	382
<i>Sorex tenellus</i>	453		374
<i>Sorex t. lyelli</i>	453		
<i>Sorex t. myops</i>	453		
<i>Sorex t. nanus</i>	453		375
<i>Sorex thompsoni</i>	457		377
<i>Sorex trowbridgii</i>	452		373
<i>Sorex tundrensis</i>	447		368
<i>Sorex vagrans</i>	448		370
<i>Sorex v. dobsoni</i>	448		370
<i>Sorex v. monticola</i>	448		371
<i>Sorex v. similis</i>	449		372
<i>Sorex vancouverensis</i>	449		371
<i>Sorex verapacis</i>	455	551	
<i>Soricidæ</i>	446	548	366
<i>soricina</i> . (<i>Blarina</i>)	464	560	
<i>soricina</i> . (<i>Glossophaga</i>)	516	672	
<i>Soricinæ</i>	446	548	366
<i>soricinus</i> . (<i>Vampyrus</i>)	515	669	
<i>soricinus</i> . (<i>Vespertilio</i>)	516	672	
<i>Soriciscus</i>	460	556	382
<i>sornborgeri</i> . (<i>Ursus a.</i>)	394		314
<i>sowerbianus</i> . (<i>Delphinus</i>)	31	55	28
<i>sowerbiensis</i> . (<i>Delphinus</i>)	23		18
<i>sowerbyi</i> . (<i>Delphinus</i>)	23		18
<i>spadicipygus</i> . (<i>Sigmodon</i>)	175		145
<i>spadicipygus</i> . (<i>Sigmodon h.</i>)	175		145
<i>spadix</i> . (<i>Putorius</i>)	430		348
<i>spadix</i> . (<i>Putorius l.</i>)	430		
<i>spatulatus</i> . (<i>Fiber</i>)	254		214
<i>speciosus</i> . (<i>Tamias</i>)	89		72
<i>spectabilis</i> . (<i>Dipodomys</i>)	293	345	235
<i>Spectrellum</i>	501	631	
<i>Spectrellum macrurum</i>	501	634	
<i>Spectrellum mexicanum</i>	502		
<i>Spectrellum stramineum</i>	501		
<i>Spectrellum s. major</i>	501		

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
spectrum. (<i>Sturnira</i>)	529	715	
spectrum. (<i>Vampyrus</i>)	511	656	
spectrum. (<i>Vespertilio</i>)	511	655	
speleus. (<i>Odocoileus</i>)	43	68	38
Spermatophilus	96	139	81
Spermophilopsis	96	139	
Spermophilus	96	139	81
Spermophilus annulatus	116	151	
Spermophilus a. goldmani	116	151	
Spermophilus armatus	112		93
Spermophilus barrowensis	110		91
Spermophilus beecheyi fisheri	114		
Spermophilus beldingi	113		94
Spermophilus beringensis	110		91
Spermophilus bernardinus	107		
Spermophilus buckleyi	115		89
Spermophilus canescens	101		97
Spermophilus castaneus	106		84
Spermophilus chrysodeirus	106		84
Spermophilus c. bernardinus	107		84
Spermophilus c. brevicaudus	107		84
Spermophilus cinerascens	106		83
Spermophilus columbianus	111		91
Spermophilus couchii	113	149	88
Spermophilus cryptospilotus	101		97
Spermophilus elegans	113		94
Spermophilus empetra	108		89
Spermophilus e. kadiacensis	108		90
Spermophilus e. plesius	108		
Spermophilus franklini	105		101
Spermophilus grammurus	113	149	88
Spermophilus g. atricapillus	114	150	
Spermophilus g. beecheyi	114		88
Spermophilus g. buckleyi	115		89
Spermophilus g. douglasi	115		89
Spermophilus g. fisheri	114	150	88
Spermophilus gunnisoni	118		
Spermophilus harrisii	96	141	85
Spermophilus h. saxicolus	97	142	86
Spermophilus interpres	98	143	86
Spermophilus lateralis	105		83
Spermophilus l. saturatus	106		83
Spermophilus leucurus	97	142	86
Spermophilus l. cinnameus	97		86
Spermophilus macrourus	113	149	88
Spermophilus mexicanus	99, 103	145, 146	96, 98
Spermophilus m. parvidens	104	146	99
Spermophilus mohavensis	102		98
Spermophilus mollis	112		92
Spermophilus m. canus	112		93

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Spermophilus m. stephensi</i>	112		93
<i>Spermophilus m. yakimensis</i>	112		93
<i>Spermophilus neglectus</i>	103		98
<i>Spermophilus nelsoni</i>	98		87
<i>Spermophilus obsoletus</i>	99		95
<i>Spermophilus oregonus</i>	113		94
<i>Spermophilus osgoodi</i>	110		91
<i>Spermophilus p. kadiacensis</i>	108		
<i>Spermophilus perotensis</i>	102	145	
<i>Spermophilus richardsoni</i>	111		92
<i>Spermophilus r. townsendi</i>	113		94
<i>Spermophilus sonoriensis</i>	102		
<i>Spermophilus spilosoma</i>	99	145	96
<i>Spermophilus s. annectens</i>	100		97
<i>Spermophilus s. arens</i>	101		
<i>Spermophilus s. macrospilotus</i>	99	145	96
<i>Spermophilus s. major</i>	99		96
<i>Spermophilus s. marginatus</i>	100		
<i>Spermophilus s. obsidianus</i>	100		97
<i>Spermophilus s. pratensis</i>	100		96
<i>Spermophilus tereticaudus</i>	102, 103	144	98
<i>Spermophilus t. sonoriensis</i>	102	144	
<i>Spermophilus townsendii</i>	111		92
<i>Spermophilus tridecemlineatus</i>	104		100
<i>Spermophilus t. alleni</i>	105		101
<i>Spermophilus t. badius</i>	105		101
<i>Spermophilus t. olivaceus</i>	104		100
<i>Spermophilus t. pallidus</i>	104		100
<i>Spermophilus t. parvus</i>	104		100
<i>Spermophilus t. texensis</i>	105		101
<i>Spermophilus wortmani</i>	107		
<i>Sphærocephalus</i>	27		
<i>Sphærocephalus incrassatus</i>	27	53	23
<i>sphagnicola</i> . (<i>Sorex</i>)	447		368
<i>sphagnicola</i> . (<i>Synaptomys</i>)	256		206
<i>spicilegus</i> . (<i>Peromyscus</i>)	156	179	
<i>spiculatus</i> . (<i>Phyllostoma</i>)	529	715	
<i>Spilogale</i>	412	519	327
<i>Spilogale ambarvalis</i>	415		332
<i>Spilogale ambigua</i>	415	521	
<i>Spilogale angustifrons</i>	416	521	
<i>Spilogale a. tropicalis</i>	416	521	
<i>Spilogale arizonæ</i>	414		331
<i>Spilogale a. martirensis</i>	414	521	
<i>Spilogale gracilis</i>	415		330
<i>Spilogale indianola</i>	413		329
<i>Spilogale interrupta</i>	412, 413	523	237
<i>Spilogale leucoparia</i>	413		329
<i>Spilogale lucasana</i>	416	523	
<i>Spilogale olympica</i>	415		332

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
Spilogale phenax.....	414		331
Spilogale p. arizonæ.....	414		331
Spilogale p. latifrons.....	414		331
Spilogale putorius.....			327
Spilogale pygmæa.....	415	519	
Spilogale ringens.....	413		329
Spilogale saxatilis.....	414		330
Spilogale tenuis.....	413		
spilosoma. (Citellus).....	99	145	
spilosoma. (Spermophilus).....	99	145	96
spilotus. (Perognathus p.).....	309		251
spinatus. (Perognathus).....	315	363	255
spissigrada. (Mephitis).....	408		326
spissigrada. (Mephitis o.).....	408		
splendens. (Neotoma).....	206	278	
stearnsii. (Grampus).....	28		25
stejnegeri. (Balænoptera).....	23		
stejnegeri. (Mesoplodon).....	23		*481
stejnegeri. (Phoca).....	443		
stelleri. (Eumetopias).....	437		355
stelleri. (Hydrodamalis).....	14		
stelleri. (Lutra).....	436	537	354
stelleri. (Otaria).....	437		
Stemmatope.....	440		
Stemmatopus.....	440		360
Stenoderma.....	526	704	
Stenoderma achradoophilum.....	527	707	
Stenoderma chilense.....	529	715	
Stenoderma lucie.....	527	706	
Stenoderma microdon.....	528		
Stenoderma montserratense.....	526	705	
Stenoderma nicholli.....	526	706	
Stenoderma rufum.....	526	704	
Stenoderma tolteca.....	525	699	
Stenodermatinæ.....	522	691	
stenorhyncha. (Orca).....	26	51	22
stenorhynchus. (Delphinus).....	33	58	
stephensi. (Citellus m.).....	112		
stephensi. (Neotoma).....	216		
stephensi. (Perognathus).....	312		253
stephensi. (Peromyscus).....	153	190	136
stephensi. (Sciuropterus).....	124		111
stephensi. (Sciuropterus a.).....	124		
stephensi. (Sciuropterus o.).....	124		
stephensi. (Spermophilus m.).....	112		93
sticte. (Lupus).....	375		
stizodon. (Sorex).....	455	551	
stonei. (Citellus).....	110		
stonei. (Microtus).....	236		181

*Supplement.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
stonei. (Ovis)	54		47
stonei. (Rangifer)	39		*483
stonei. (Synaptomys)	255		203
stonii. (Citellus)	110		
stonii. (Ovis)	54	85	
stonii. (Rangifer)	39		
stonii. (Synaptomys c.)	255		
stramineum. (Spectrellum)	501		
stramineus. (Natalis)	501	632	
stramineus. (Natalus)	501	632	
streatori. (Neotoma f.)	206		159
streatori. (Perodipus)	294		237
streatori. (Putorius)	427		345
streatori. (Sciurus h.)	82		64
streatori. (Sorex p.)	446		367
striatus. (Sciurus)	85, 95	134	
striatus. (Tamias)	95		80
Strongyloceros	41		
strongyloceros. (Cervus)	41		34
struthopus. (Lepus a.)	361		274
Sturnira	529	714	
Sturnira lilium	529	715	
Sturnira spectrum	529	715	
styx. (Delphinus)	33	58	30
subarcticus. (Peromyscus t.)	146		131
subauratus. (Sciurus)	65		52
subcinctus. (Lepus f.)	344	417	
subflavus. (Pipistrellus)	481		409
subflavus. (Vespertilio)	481		400
subgriseus. (Peromyscus)	139		142
subgriseus. (Sitomys)	139		145
subsolana. (Felis c.)	370		296
subsolanus. (Lynx)	370		296
subulatus. (Myotis)	478	580	405
subulatus. (Vespertilio)	478	580	405
suckleyi. (Sciurus)	82		65
suckleyi. (Sorex)	448		370
Suidæ		61	
sulfurea. (Bakenoptera)	20	38, 43	14
sulfureus. (Sibbaldius)	20	43	
sulphureus. (Artibeus)	527	707	
sumichrasti. (Bassariscus)	398		
sumichrasti. (Hesperomys)	172	216	
sumichrasti. (Nyctomys)	172	216	
sumichrasti. (Reithrodon)	197	264	
surberi. (Neotoma m.)	209		156
Sus albirostris	37	66	
Sus tajacu	35	61	
sylvaticus. (Lepus)	342		280

*Supplement.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
sylvestris. (<i>Cervus t.</i>).....	37		35
sylvestris. (<i>Glyphonycteris</i>).....	513	664	
Sylvicola.....	234	299	178
Sylvilagus.....	340, 341	413, 416	269
Synaptomys.....	255		203
Synaptomys andersoni.....	257		
Synaptomys bullatus.....	258		
Synaptomys chapmani.....	257		
Synaptomys cooperi.....	255		203
Synaptomys c. gossi.....	255		
Synaptomys c. stonii.....	255		203
Synaptomys dalli.....	258		206
Synaptomys fatuus.....	256		204
Synaptomys helaletes.....	255		204
Synaptomys h. gossii.....	255		204
Synaptomys innuitus.....	256		205
Synaptomys i. medioximus.....	256		†430
Synaptomys sphagnicola.....	256		206
Synaptomys stonei.....	255		203
Synaptomys truei.....	258		207
Synaptomys truii.....	257, 258		
Synaptomys wrangeli.....	256, 257		206
Synotus.....	490	603	398
Synotus townsendi.....	490	604	399
Syntheosciurus.....	59	91	
Syntheosciurus brochus.....	59	91	
Synthetosciurus.....	59	91	
Synthetosciurus brochus.....	59	91	
tabascensis. (<i>Didelphis m.</i>).....	6	16	
tabascensis. (<i>Didelphys m.</i>).....	6	16	
tabernaculi. (<i>Hydrodamalis</i>).....		36	
Tachynices.....	24		
Tachynices megacephalus.....	24		
tæniata. (<i>Talpa</i>).....	468		391
tæniurus. (<i>Sciurus</i>).....	60	101	
Tagassu.....	35	61	
Tagassu angulatum.....	35	62, 63	
Tagassu a. crassum.....	36	64	
Tagassu a. humerale.....	35	63	
Tagassu a. sonoriense.....	36	64	
Tagassu a. yucatanense.....	36	63	
Tagassu crusnigrum.....	36	65	
Tagassu nanus.....	35	62	
Tagassu pecari.....	37	66, 67	
Tagassu p. ringens.....	37	66	
Tagassuidæ.....	35	61	
tajacu. (<i>Dicotyles</i>).....	35		33
tajacu. (<i>Sus</i>).....	35	61	

†Appendix.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
talamancæ. (<i>Oryzomys</i>)	183	241	
<i>Talpa cupreata</i>	470		388
<i>Talpa e. flava</i>	470		388
<i>Talpa e. flavescens</i>	469		388
<i>Talpa flava</i>	470		388
<i>Talpa fusca</i>	470		388
<i>Talpa longicaudata</i>	467		
<i>Talpa machrina</i>	470		389
<i>Talpa pennantii</i>	470		388
<i>Talpa radiata</i>	467		395
<i>Talpa sericea</i>	470		389
<i>Talpa tæniata</i>	468		391
<i>Talpidae</i>	466	563	386
<i>Talpinæ</i>	467	564	388
<i>talpoides</i> . (<i>Blarina</i> b.)	461		
<i>talpoides</i> . (<i>Cricetus</i>)	278		227
<i>talpoides</i> . (<i>Sorex</i>)	460	555	382
<i>talpoides</i> . (<i>Thomomys</i>)	278		227
<i>Talposorex</i>	460		381
<i>Tamandua</i>	10	26	
<i>Tamandua bivittata</i>	10	27	
<i>Tamandua sellata</i>	11, 12	28	
<i>Tamandua tetradactyla</i>	10, 11	27	
<i>Tamandua t. chiriquensis</i>	11		
<i>Tamandua t. tenuirostris</i>	11		
<i>Tamanduas</i>	10	26	
<i>Tamia</i>	85		
<i>Tamias</i>	85	134	67
<i>Tamias adsitus</i>	95		
<i>Tamias alpinus</i>	94		
<i>Tamias amœnus</i>	90		74
<i>Tamias a. borealis</i>	92		
<i>Tamias a. bulleri</i>	88		
<i>Tamias a. merriami</i>	88		
<i>Tamias a. operarius</i>	†541		
<i>Tamias a. pallidus</i>	93		
<i>Tamias bulleri</i>	88	136, 137	
<i>Tamias callipeplus</i>	89		73
<i>Tamias c. inyoensis</i>	89		73
<i>Tamias canescens</i>	86	†741	
<i>Tamias canicaudus</i>	93		
<i>Tamias caniceps</i>	93		†426
<i>Tamias castanurus</i>	106		85
<i>Tamias caurinus</i>	90		74
<i>Tamias chrysodeirus</i>	106		84
<i>Tamias cinerascens</i>	106		83
<i>Tamias cinereicollis</i>	90		75
<i>Tamias c. canipes</i>	91		
<i>Tamias cooperi</i>	86		69

†Appendix.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Tamias dorsalis</i>	85	135	68
<i>Tamias d. utahensis</i>	85		69
<i>Tamias durangæ</i>	88	137	
<i>Tamias frater</i>	90		73
<i>Tamias hindsi</i>	86		
<i>Tamias hopiensis</i>	†541		
<i>Tamias interpretes</i>	98		86
<i>Tamias l. saturatus</i>	106		83
<i>Tamias lectus</i>	94		
<i>Tamias leucurus</i>	97	142	86
<i>Tamias l. cinnamomeus</i>	97		86
<i>Tamias l. peninsulae</i>	97	143	
<i>Tamias macrorhabdotes</i>	88		71
<i>Tamias m. melanurus</i>	94		79
<i>Tamias merriami</i>	88	136, 137	71
<i>Tamias minimus</i>	93		78
<i>Tamias m. alpinus</i>	94		79
<i>Tamias m. consobrinus</i>	93		78
<i>Tamias m. pictus</i>	94		79
<i>Tamias obscurus</i>	88	136	
<i>Tamias oreocetes</i>	95		80
<i>Tamias palmeri</i>	86		69
<i>Tamias panamintinus</i>	90		74
<i>Tamias pricei</i>	86		
<i>Tamias quadrimaculatus</i>	88		†71
<i>Tamias quadrivittatus</i>	91	138	75
<i>Tamias q. affinis</i>	92		77
<i>Tamias q. borealis</i>	92		77
<i>Tamias q. felix</i>	92		76
<i>Tamias q. gracilis</i>	91		76
<i>Tamias q. luteiventris</i>	92		76
<i>Tamias q. neglectus</i>	92		78
<i>Tamias q. pallidus</i>	85	135	68
<i>Tamias senex</i>	89		72
<i>Tamias speciosus</i>	89		72
<i>Tamias s. inyoensis</i>	89		73
<i>Tamias striatus</i>	95		80
<i>Tamias s. griseus</i>	96		81
<i>Tamias s. lysteri</i>	95		80
<i>Tamias s. venustus</i>	96		81
<i>Tamias townsendi</i>	86		69
<i>Tamias t. hindsi</i>	86		70
<i>Tamias t. hindsii</i>	86		
<i>Tamias t. littoralis</i>	87		
<i>Tamias t. ochrogenys</i>	87		71
<i>Tamias t. pricei</i>	87		70
<i>Tamias t. pricii</i>	86		
<i>Tamias townsendii</i>	86		
<i>Tamias umbrinus</i>	91		75

†Appendix.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Tamias wortmani</i>	107		84
<i>Tamiasciurus</i>	59, 80	94	60
<i>Tapeti</i>	340, 357	413, 426	
<i>Tapirella</i>	58	87	
<i>Tapirella bairdi</i>	58	87	
<i>Tapirella dowi</i>	58	88	
<i>Tapiridæ</i>	58	87	
<i>Tarandus</i>	37		
<i>tarandus</i> . (<i>Cervus</i>)	37		35
<i>Tatoua</i>	12	31	
<i>Tatoua centralis</i>	12	32	
<i>Tatu</i>	13	32	
<i>Tatu novemcinctum</i>	13	33	
<i>Tatuinae</i>	13	32	
<i>Tatusia</i>	13	32	4
<i>Tatusia leptorhyncha</i>	13	34	4
<i>Taxidea</i>	404	503	319
<i>Taxidea americana</i>	404		320
<i>Taxidea a. berlandieri</i>	405		321
<i>Taxidea a. neglecta</i>	405		321
<i>Taxidea berlandieri</i>	405		
<i>Taxidea taxus</i>	404		
<i>Taxidea t. berlandieri</i>	405	503	
<i>Taxidea t. infusca</i>	405	505	
<i>Taxidea t. neglecta</i>	405		
<i>taxus</i> . (<i>Taxidea</i>)	404		
<i>taxus</i> . (<i>Ursus</i>)	404		
<i>Tayassu</i>	35	61	*483
<i>Tayassu a. ringens</i>	37	66	
<i>Tayassu angulatus crassus</i>	36	64	
<i>Tayassu a. humeralis</i>	35	63	
<i>Tayassu a. yucatanensis</i>	36	63	
<i>Tayassu crasnigrum</i>	36	65	
<i>Tayassu nanus</i>	35	62	
<i>Tayassu pecari</i>	36, 37	65, 66	
<i>taylori</i> . (<i>Hesperomys</i>)	135		141
<i>taylori</i> . (<i>Peromyscus</i>)	135		141
<i>Tayra</i>	418	24	
<i>Teanopus</i>	225	294	
<i>Teanopus phenax</i>	225	295	
<i>teapensis</i> . (<i>Oryzomys</i>)	185	247	
<i>teapensis</i> . (<i>Peromyscus m.</i>)	168	207	
<i>tectirostris</i> . (<i>Sibbaldius</i>)	20		
<i>tectorum</i> . (<i>Mus</i>)	129	164	118
<i>tectus</i> . (<i>Oryzomys</i>)	188	244	
<i>teguina</i> . (<i>Acodon</i>)	204	273, 274	
<i>teguina</i> . (<i>Hesperomys</i>)	204	273	
<i>tehuantepecus</i> . (<i>Peromyscus</i>)	168	206	
<i>teliotis</i> . (<i>Atalapha</i>)	486	593	413

*Supplement.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
teliotis. (<i>Lasiurus b.</i>).....	486	593	413
telmalemonus. (<i>Lepus</i>).....	341		278
telmalestes. (<i>Blarina</i>).....	462		384
tema. (<i>Mazama</i>).....		80	
tenellus. (<i>Sorex</i>).....	453		374
tenellus. (<i>Zapus</i>).....	325		259
Tenotis.....	85	134	68
tenuicauda. (<i>Neotoma</i>).....	219	286	
tenuidorsalis. (<i>Vespertilio</i>).....	476	578	404
tenuirostris. (<i>Reithrodontomys</i>).....	201	268	
tenuirostris. (<i>Rhithrodontomys</i>).....	201	268, 269	
tenuirostris. (<i>Tamandua t.</i>).....	11		
tenuis. (<i>Reithrodontomys</i>).....	199	265	
tenuis. (<i>Rhithrodontomys</i>).....	199	265	
tenuis. (<i>Spilogale</i>).....	413		
Teonoma.....	220		163
Teonoma acraea.....	221		
Teonoma arizonae.....	223		
Teonoma cinerea.....	220, 221		
Teonoma c. acraea.....	221		
Teonoma c. columbiana.....	221		
Teonoma c. drummondi.....	221		
Teonoma c. occidentalis.....	221		
Teonoma cinnamomea.....	223		
Teonoma fusca.....	222		
Teonoma f. apicalis.....	222		
Teonoma grangeri.....	223		
Teonoma oreolestes.....	223		
Teonoma rupicola.....	223		
Teonoma saxamans.....	221		
tephrogaster. (<i>Sciurus</i>).....	60	101	
tereticaudus. (<i>Citellus</i>).....	102	144	
tereticaudus. (<i>Spermophilus</i>).....	102	144	98
terrænovæ. (<i>Arvicola</i>).....	238		
terrænovæ. (<i>Microtus</i>).....	238		182
terrænovæ. (<i>Rangifer</i>).....	38		36
Terricola.....	234	299	178
tesselatus. (<i>Vespertilio</i>).....	485	593	412
tethyos. (<i>Delphinus</i>).....	34	58	30
tetradactyla. (<i>Myrmecophaga</i>).....	10	26, 27	
tetradactyla. (<i>Tamandua</i>).....	10	27	
Tetramerodon.....	234		179
tetramerus. (<i>Arvicola</i>).....	243		190
tetramerus. (<i>Microtus</i>).....	243		190
texana. (<i>Hesperomys</i>).....	145	186	
texanus. (<i>Dorcelaphus</i>).....	45	70	40
texanus. (<i>Scalops a.</i>).....	471		
texensis. (<i>Citellus t.</i>).....	105		
texensis. (<i>Conepatus l.</i>).....	411	515	
texensis. (<i>Didelphis m.</i>).....	6	16	

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
texensis. (Didelphys m.).....	6	16	
texensis. (Felis r.).....	371	458	298
texensis. (Geomys).....	264		219
texensis. (Heteromys).....	323		
texensis. (Lepus).....	355		290
texensis. (Liomys).....	323		
texensis. (Lynx).....	371		
texensis. (Odocoileus a.).....	45	70	
texensis. (Odontocælus a.).....	45	70	40
texensis. (Oryzomys p.).....	181		147
texensis. (Peromyscus).....	145	186	130
texensis. (Scalops a.).....	471		390
texensis. (Scalopus a.).....	471		
texensis. (Sciurus r.).....	65	110	
texensis. (Sigmodon h.).....	175	228	144
texensis. (Spermophilus t.).....	105		101
texensis. (Thomomys f.).....	288		
texensis. (Urocyon c.).....	386	478	308
texiana. (Arvicola).....	175		144
texianus. (Lepus).....	354, 355		
texianus. (Sciurus).....	65		
texianus. (Sciurus r.).....	65		
Thalarctos.....	390	479	
Thalassarctos.....	390	479	309
Thalassarcus.....	390		310
thersites. (Nasua).....		486	
thicola. (Lagenorhynchus).....	29		26
Thiosmus.....	410	512	
thomasi. (Megadontomys).....	171	213	
thomasi. (Odocoileus).....	47	75	
thomasi. (Odontocælus).....	47	75	
thomasi. (Peromyscus).....	171	213	
thomasi. (Sciurus).....	75	126	
Thomomys.....	271, 272	332	221
Thomomys alpinus.....	283		229
Thomomys altivallis.....	287		230
Thomomys angularis.....	281		228
Thomomys a. pascalis.....	282		
Thomomys aphrastus.....	282	336	
Thomomys atrovarius.....	283	338	
Thomomys aureus.....	285		229
Thomomys a. lachuguilla.....	285		
Thomomys a. perpes.....	285		
Thomomys a. pervagus.....	285		
Thomomys baileyi.....	286		
Thomomys bottæ.....	274		223
Thomomys b. pallescens.....	274		224
Thomomys bridgeri.....	279		
Thomomys bulbivorus.....	272		222
Thomomys cabezonæ.....	286		

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Thomomys cervinus</i>	287	340	230
<i>Thomomys clusius</i>	284		229
<i>Thomomys c. fuscus</i>	277		
<i>Thomomys c. ocius</i>	284		
<i>Thomomys desertorum</i>	281		
<i>Thomomys douglasi</i>	277, 278		226
<i>Thomomys d. fisheri</i>	277		
<i>Thomomys d. fuscus</i>	277		226
<i>Thomomys d. oregonus</i>	277		
<i>Thomomys d. yelmensis</i>	277		226
<i>Thomomys fossor</i>	287		230
<i>Thomomys fulvus</i>	272, 280	334	227
<i>Thomomys f. alticola</i>	280	335	
<i>Thomomys f. anitæ</i>	280	335	
<i>Thomomys f. fisheri</i>	277		
<i>Thomomys f. intermedius</i>	280	336	228
<i>Thomomys f. martirensis</i>	282		
<i>Thomomys f. nigricans</i>	280	335	228
<i>Thomomys f. texensis</i>	281		
<i>Thomomys goldmani</i>	283	339	
<i>Thomomys helleri</i>	278		
<i>Thomomys hesperus</i>	279		
<i>Thomomys idahoensis</i>	284		
<i>Thomomys laticeps</i>	274		224
<i>Thomomys latirostris</i>	286		
<i>Thomomys leucodon</i>	273		223
<i>Thomomys l. navus</i>	273		
<i>Thomomys limosus</i>	278		
<i>Thomomys martirensis</i>	282	336	
<i>Thomomys mazama</i>	275		225
<i>Thomomys melanops</i>	279		227
<i>Thomomys monticola</i>	275		225
<i>Thomomys m. pinetorum</i>	275		225
<i>Thomomys monticolus</i>	275		
<i>Thomomys m. pinetorum</i>	275		
<i>Thomomys myops</i>	276		
<i>Thomomys nasicus</i>	276		226
<i>Thomomys nelsoni</i>	286	339	
<i>Thomomys nevadensis</i>	272		223
<i>Thomomys niger</i>	276		
<i>Thomomys operarius</i>	273		223
<i>Thomomys orizabæ</i>	282	337	
<i>Thomomys perditus</i>	283	338	
<i>Thomomys peregrinus</i>	282	337	
<i>Thomomys perpallidus</i>	286, 287	340	229
<i>Thomomys pygmæus</i>	284		
<i>Thomomys quadratus</i>	276		225
<i>Thomomys rufescens</i>	279	332	221, 227
<i>Thomomys scapterus</i>	274		
<i>Thomomys sinaloæ</i>	287	339	

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
Thomomys talpoides	278		227
Thomomys t. perpallidus.....	286		
Thomomys t. rufescens	279		227
Thomomys toltecus	283	338	
Thomomys townsendi	275		224
Thomomys uinta.....	272		
Thomomys umbrinus.....	272, 280		
thompsoni. (Sorex)	457		377
thoracatus. (Capromys)	331	394	
thoracatus. (Capromys b.)	331	394	
thurberi. (Peromyscus).....	146	176	
thurberi. (Peromyscus t.).....	146		
thurberi. (Sitomys a.).....	146		
Thylamys.....	2	5	
Thyroptera.....	503	637	
Thyroptera discifera.....	503	637	
Thyroptera tricolor.....	503	637	
Thyropteridæ.....	501		
thysanodes. (Myotis).....	479	572	
thysanodes. (Myotis e.).....	479		
tiburonensis. (Peromyscus)	160	175	
tigrina. (Felis)	367	449	
timidus. (Lepus).....	340	413	269
tolteca. (Felis j.)	368	452	
tolteca. (Stenoderma)	525	699	
toltecus. (Cervus).....	46		
toltecus. (Hesperomys).....	177	226	
toltecus. (Odontocœlus)	46	74	
toltecus. (Reithrodontomys l.)	203	271	
toltecus. (Rhithrodontomys l.)	203	271	
toltecus. (Sigmodon h.).....	177	226	
toltecus. (Thomomys).....	283	338	
toluçæ. (Reithrodontomys c.)	199	266	
toluçæ. (Rhithrodontomys c.)	199	266	
toluçæ. (Sciurus o.)	63	107	
tonalensis. (Sigmodon h.).....	176	229	
Tonatia.....	512	658	
Tonatia amblyotis.....	512	658	
tornillo. (Peromyscus).....	142	188	126
torquata. (Neotoma).....	208	279	
torridus. (Hesperomys).....	133		
torridus. (Heterogeomys).....	270	326	
torridus. (Heteromys).....	321	376	
torridus. (Liomys).....	321	376	
torridus. (Onychomys)	133	167	122
totontepecus. (Peromyscus m.).....	160	608	
townsendi. (Arctocephalus)	438	544	
townsendi. (Arvicola)	243		190
townsendi. (Citellus)	111		
townsendi. (Corynorhinus m.).....	491	604	400

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
townsendi. (<i>Lepus</i>)	363		
townsendi. (<i>Lepus c.</i>)	363		
townsendi. (<i>Microtus</i>)	243		190
townsendi. (<i>Scapanus</i>)	468		391
townsendi. (<i>Spermophilus r.</i>)	113		94
townsendi. (<i>Synotus</i>)	490	604	399
townsendi. (<i>Tamias</i>)	86		69
townsendi. (<i>Thomomys</i>)	275		224
townsendi. (<i>Urocyon c.</i>)	387		309
townsendii. (<i>Geomys</i>)	275		224
townsendii. (<i>Plecotus</i>)	490, 491	604	400
townsendii. (<i>Scalops</i>)	468		391
townsendii. (<i>Spermophilus</i>)	111		92
townsendii. (<i>Tamias</i>)	86		
Trachops'	514	664	
Trachops fuliginosus	514	664	
Trachyops	514	664	
Trachyops cirrhosus	514	665	
transitionalis. (<i>Lepus f.</i>)	342		281
transitionalis. (<i>Lepus s.</i>)	342		
tredecimlineatus. (<i>Citellus</i>)	104		
Trichechidæ	14	36	357
Trichechus	14	36	358
Trichechus divergens	439		358
Trichechus inunguis		35	
Trichechus manatus	14	35, 37	6
Trichechus obesus	437		358
Trichechus rosmarus	439		358
Trichechus senegalensis		35	
Trichocorys'	529	716	
trichopus. (<i>Zygoeomys</i>)	271	332	
tricolor. (<i>Thyroptera</i>)	503	637	
tridactyla. (<i>Myrmecophaga</i>)	12	24, 29	
tridactylus. (<i>Bradypus</i>)	9	21	
tridecemlineatus. (<i>Sciurus</i>)	104		
tridecemlineatus. (<i>Spermophilus</i>)	104		100
trimucronatus. (<i>Arvicola</i>)	258		207
trimucronatus. (<i>Lemmus</i>)	258, 259		207
trinitatis. (<i>Callospermophilus c.</i>)	107		
trinitatis. (<i>Citellus c.</i>)	107		
trinitatis. (<i>Echimys</i>)	329	385	
Trinodontomys	135	170	123
trinitatus. (<i>Zapus</i>)	326		261
trivirgata. (<i>Simia</i>)	533	728	
tropicalis. (<i>Blarina</i>)	464	559	
tropicalis. (<i>Conepatus</i>)	412	517	
tropicalis. (<i>Corsira</i>)		559	
tropicalis. (<i>Monachus</i>)	445	543	
tropicalis. (<i>Neotoma</i>)	220	288	
tropicalis. (<i>Phoca</i>)	445		

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
tropicalis. (Putorius).....	432	531	
tropicalis. (Spilogale a.).....	416	521	
tropidorhynchus. (Molossus).....	496	620	
trowbridgei. (Lepus).....			281
trowbridgi. (Arvicola).....	240		186
trowbridgii. (Sorex).....	452		373
truei. (Hesperomys).....	159		139
truei. (Lepus).....	341		
truei. (Odocoileus).....	46	73	
truei. (Peromyscus).....	159		139
truei. (Phenacomys).....	226		168
truei. (Scapanus).....	469		393
truei. (Sciurus).....	73		
truei. (Synaptomys).....	258		207
truii. (Lepus).....	341	415	
truii. (Odontocælus).....	46	73, 74	
truii. (Peromyscus).....	159		
truii. (Scapanus).....	469		
truii. (Sciurus).....	73	123	
truii. (Synaptomys).....	257, 258		
truncatus. (Delphinus).....	31	56	29
truncatus. (Tursiops).....	31	56	29
tschuktschorum. (Lepus).....	359		271
tularensis. (Lepus).....	357		
tularensis. (Onychomys l.).....	132		
tularensis. (Onychomys t.).....	132		
tularensis. (Perodipus a.).....	293		
tumbalensis. (Tylomys).....	173	218	
tumida. (Rhogeessa).....	489		
tumida. (Rhogoessa).....	489	601	
tumidifrons. (Chilonatalis).....	503	637	
tumidifrons. (Chilonatalus).....	503	637	
tundrensis. (Sorex).....	447		368
Tursio.....	21, 31	45, 55	28
Tursio doreides.....	34	58	30
tursio. (Delphinus).....	31	55	29
tursio. (Tursiops).....	31	56	29
Tursiops.....	31	55, 57	28
Tursiops gilli.....	32	47, 56	29
Tursiops truncatus.....	31	56	29
Tursiops tursio.....	31	56	29
tuza. (Geomys).....	261		215
tuza. (Mus).....	261		215
Tylomys.....	173	217, 221	
Tylomys bullaris.....	173	219	
Tylomys nudicaudus.....	173	217	
Tylomys panamensis.....	174	219	
Tylomys tumbalensis.....	173	218	
Tylomys watsoni.....	173	219	
Tylonyx.....	260		

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
tylorhinus. (<i>Platygeomys</i>)	267	319	
Tylostoma mexicana	514	665	
ubericolor. (<i>Lepus</i> b.)	352		
ubericolor. (<i>Lepus</i> f.)	352		282
uinta. (<i>Felis</i>)	373		
uinta. (<i>Lynx</i>)	373		
uinta. (<i>Thomomys</i>)	272		
umbrinus. (<i>Ascomys</i>)	272		
umbrinus. (<i>Geomys</i>)	280		
umbrinus. (<i>Peromyscus</i> c.)	144		129
umbrinus. (<i>Pseudostoma</i>)	272	334	
umbrinus. (<i>Tamias</i>)	91		75
umbrinus. (<i>Thomomys</i>)	272, 280		
umbrosus. (<i>Microtus</i>)	251	305	
unalascensis. (<i>Dicrostonyx</i>)	260		210
unalascensis. (<i>Dicrostonyx</i> h.)	260		210
unalascensis. (<i>Microtus</i>)	242		188
underwoodi. (<i>Hylonycteris</i>)	517, 518	675	
ungava. (<i>Evotomys</i>)	232		175
ungava. (<i>Phenacomys</i>)	227		169
ungava. (<i>Vulpes</i> l.)	383		
Ungulata	35	60	33
unicinctus. (<i>Dasypus</i>)	12	31	
Uranodon	21		
Urocryptus	492	610	
Urocryptus bilineatus	493	610	
Urocyon	386	474	307
Urocyon c. townsendi	387		
Urocyon catalinae	389		
Urocyon cinereo-argenteus	386		307
Urocyon c. californicus	387	477	308
Urocyon c. borealis	388		
Urocyon c. floridanus	386		307
Urocyon c. fraterculus	389	475	
Urocyon c. guatemalæ	389	475	
Urocyon c. inyoensis	387		
Urocyon c. littoralis	388, 389		*507
Urocyon c. ocythous	388		309
Urocyon c. parvidens	389	476	
Urocyon c. santacruzæ	388		
Urocyon c. scotti	386	477	308
Urocyon c. texensis	386	478	308
Urocyon c. townsendi	387		309
Urocyon clementæ	390		
Urocyon guatemalæ	389		
Urocyon l. santacruzæ	388		
Urocyon parvidens	389		
Uroderma	524	697	

* Supplement.

	Check-List of Mammals.	Mammals of Middle America, etc	Synopsis of Mammals, etc.
Uroderma bilobatum.....	524	697	
Uroderma convexum.....	524	697	
Uroleptes.....	10	26	
Urotrichus gibbsii.....	466		387
Ursidæ.....	390	479	309
ursina. (Callotaria).....	438		357
ursina. (Phoca).....	438		357
Ursinæ.....	390	479	309
ursinus. (Molossus).....	495	619	
ursinus. (Promops).....	496, 497	621	
ursinus. (Vespertilio).....	483	587	410
Ursus.....	390	479	309
Ursus altifrontalis.....	395		
Ursus amblyceps.....	394, 396		
Ursus americanus.....	394		313
Ursus a. carlottæ.....	394		
Ursus a. emmonsii.....	397		
Ursus a. eremicus.....	394		
Ursus a. sornborgeri.....	394		314
Ursus arctos.....	390	479	309
Ursus cancrivorus.....	402	492	
Ursus carlottæ.....	394		
Ursus dalli.....	391		311
Ursus d. gyas.....	391		
Ursus emmonsii.....	397		315
Ursus eulophus.....	392		
Ursus floridanus.....	395		314
Ursus horriæus.....	393	480	
Ursus horribilis.....	392, 393		312
Ursus h. alascensis.....	393		312
Ursus h. californicus.....	392		
Ursus h. horriæus.....	393		312
Ursus h. phæonyx.....	393		
Ursus hylodromus.....	396		
Ursus kenaicensis.....	392		
Ursus kermodei.....	397		
Ursus kermodii.....	397		
Ursus kidderi.....	391		
Ursus labradoria.....	404		320
Ursus lotor.....	400		
Ursus iuscus.....	416		
Ursus luteolus.....	395		314
Ursus machetes.....	396	481	
Ursus maritimus.....	390		310
Ursus merriami.....	391		
Ursus middendorffi.....	390		310
Ursus richardsoni.....	393		313
Ursus sitkensis.....	391		
Ursus taxus.....	404		
utah. (Citellus v.).....	115		

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
utahensis. (<i>Eutamias d.</i>).....	85		
utahensis. (<i>Perodipus m.</i>).....	296		
utahensis. (<i>Tamias d.</i>).....	85		69
 vafra. (<i>Vulpes p.</i>).....	380		305
vaga. (<i>Lutra c.</i>).....	434		352
vaga. (<i>Lutra h.</i>).....	434		
vagans. (<i>Pipistrellus</i>).....	487	584	
vagans. (<i>Vesperugo</i>).....	482		
vagrans. (<i>Sorex</i>).....	448		370
vallicola. (<i>Microtus c.</i>).....	240		186
Vampyrella.....	512	660	
Vampyressa.....	525	701	
Vampyriscus.....	525	701	
Vampyrodes.....	525	701	
Vampyrops.....	525	701	
Vampyrops helleri.....	526	703	
Vampyrops lineatus.....	525	702	
Vampyrops vittatus.....	526	704	
Vampyrus.....	511	655	
Vampyrus auritus.....	511, 512	656	
Vampyrus bidens.....	512	658	
Vampyrus cirrhosus.....	514	665	
Vampyrus soricinus.....	515	669	
Vampyrus spectrum.....	511	656	
vancouverensis. (<i>Sciurus h.</i>).....	82		64
vancouverensis. (<i>Sorex</i>).....	449		371
variabilis. (<i>Bassaris</i>).....	398	487	
variabilis. (<i>Sciurus</i>).....	62	105	
varians. (<i>Mephitis</i>).....	407		322
varians. (<i>Mephitis m.</i>).....	407		
variegata. (<i>Phoca</i>).....	442		363
variegatoides. (<i>Sciurus</i>).....	77	129	
variegatus. (<i>Ateles</i>).....	535	733	
variegatus. (<i>Chironectes</i>).....	1	3	
variegatus. (<i>Citellus</i>).....	113	148	
variegatus. (<i>Delphinus</i>).....	31	55	28
variegatus. (<i>Sciurus</i>).....	69, 113	116, 148	
varius. (<i>Sciurus</i>).....	69	117	
velifer. (<i>Myotis</i>).....	473	573	401
velifer. (<i>Vespertilio</i>).....	473	573	401
velifer. (<i>Vespertilio a.</i>).....	479		
velifera. (<i>Balanoptera</i>).....	19	43	12
vellerosus. (<i>Ateles</i>).....	534	733	
vellerosus. (<i>Microtus</i>).....	244		191
velox. (<i>Canis</i>).....	384		
velox. (<i>Dysopes</i>).....	495	620	
velox. (<i>Vulpes</i>).....	384		306
ventorum. (<i>Sciurus h.</i>).....	81		63
ventralis. (<i>Sorex o.</i>).....	450	550	

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
venustus. (Perodipus)	294		
venustus. (Tamias s.)	96		81
veræcrucis. (Heteromys)	319	379	
veræcrucis. (Lepus)	347	420	
veræcrucis. (Liomys)	319	379	
veræcrucis. (Pipistrellus)	481	584	
veræcrucis. (Vesperugo)	481	584	
veræpaci. (Sorex)	455	551	
verrucata. (Carollia)	515	669	
versabilis. (Megaptera)	18	41	10
Vesperides	480		407
Vesperimus	135	170	123
Vesperimus difficilis	158	193	
Vesperimus fraterculus	155		136
Vesperimus mearnsii	150		132
Vesperimus nasutus	159		139
Vespertilio	473, 482	570, 585	400, 410
Vespertilio affinis	478		402
Vespertilio albescens	479		
Vespertilio a. evotis	479		
Vespertilio a. velifer	479		
Vespertilio albicularis	485	590	
Vespertilio arquatus	483	587	410
Vespertilio auduboni	480		407
Vespertilio barbatus	502	634	
Vespertilio borealis	485	592	411
Vespertilio californicus	476	578	
Vespertilio caninus	493	614	
Vespertilio carolii	478		402
Vespertilio carolinensis	483	587	410
Vespertilio ciliolabrum	477		
Vespertilio cinereus	487	595	413
Vespertilio concinnus	475	575	
Vespertilio dutertreus	484	588	
Vespertilio erythrodactylus	481		409
Vespertilio evotis	474	574	
Vespertilio exilis	476	578	403
Vespertilio fuscus	482, 483	589	410
Vespertilio f. bahamensis	484	588	
Vespertilio f. bernardinus	483	589	
Vespertilio f. cubensis	484	588	
Vespertilio f. melanopterus	483		
Vespertilio f. miradorensis	484	588	
Vespertilio f. osceola	483		
Vespertilio f. peninsule	484	589	
Vespertilio f. propinquus	484	589	
Vespertilio gaumeri	485	590	
Vespertilio gryphus	478		405
Vespertilio g. septentrionalis	478		405
Vespertilio hastatus	514	666	

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Vespertilio humeralis</i>	488	597	415
<i>Vespertilio incautus</i>	473		
<i>Vespertilio lasiurus</i>	485	592	411
<i>Vespertilio lepidus</i>	502	634	
<i>Vespertilio leporinus</i>	494	617	
<i>Vespertilio lepturus</i>	492	610	
<i>Vespertilio longicrus</i>	479	581	402
<i>Vespertilio lucifugus</i>	478		402
<i>Vespertilio l. austroriparius</i>	478		402
<i>Vespertilio mastivus</i>	494	617	
<i>Vespertilio melanorhinus</i>	476	578	
<i>Vespertilio mexicanus</i>	477	579	
<i>Vespertilio monachus</i>	485	593	412
<i>Vespertilio murinus</i>	473, 482	570, 585	400, 410
<i>Vespertilio myotis</i>	473	570	
<i>Vespertilio naso</i>	492		
<i>Vespertilio nigricans</i>	475, 476	575	404
<i>Vespertilio nitidus</i>	476	578	403
<i>Vespertilio n. henshawi</i>	476	578	404
<i>Vespertilio noctivagans</i>	479, 480		407
<i>Vespertilio noveboracensis</i>	485	592	411
<i>Vespertilio noveboracis</i>	485	593	411
<i>Vespertilio obscurus</i>	476	578	403
<i>Vespertilio oregonensis</i>	476	578	403
<i>Vespertilio pallidus</i>	491		396
<i>Vespertilio perspicillatus</i>	515	667	
<i>Vespertilio phaiops</i>	483	587	410
<i>Vespertilio pipistrellus</i>	480	581	408
<i>Vespertilio pruinosus</i>	487	595	413
<i>Vespertilio pulverulentus</i>	480		407
<i>Vespertilio rubellus</i>	485	593	411
<i>Vespertilio rubra</i>	485	593	412
<i>Vespertilio rufus</i>	485	593	412
<i>Vespertilio soricinus</i>	516	672	
<i>Vespertilio spectrum</i>	511	655	
<i>Vespertilio subflavus</i>	481		409
<i>Vespertilio subulatus</i>	478	580	405
<i>Vespertilio s. keenii</i>	478		405
<i>Vespertilio tenuidorsalis</i>	476	578	404
<i>Vespertilio tessellatus</i>	485	593	412
<i>Vespertilio ursinus</i>	483	587	410
<i>Vespertilio velifer</i>	473	573	401
<i>Vespertilio volans</i>	476	578	403
<i>Vespertilio yumanensis</i>	475, 476	576, 578	403
<i>Vespertilionidæ</i>	473	570	396
<i>Vespertilioninæ</i>	473	570	
<i>Vesperugo</i>	483	585	410
<i>Vesperugo carolinensis</i>	481		409
<i>Vesperugo merriami</i>	480	582	409
<i>Vesperugo vagans</i>	482		

	Check-List of Mammals	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
Vesperugo veraecrucis.....	481	584	
Vesperus.....	482	580	410
Vesperus albigularis.....	485	590	
Vesperus cubanus.....	488	599	
Vesperus propinquus.....	484		
vetulus. (Hodomys).....	226	297	
vicinior. (Peromyscus b.).....	163	190	
vicinus. (Sciurus l.).....	65		52
victus. (Oryzomys).....	188	244	
vigilis. (Canis).....	378	467	
villosa. (Alouatta).....	532	726	
villosum. (Chiroderma).....	528	710	
villosus. (Mycetes).....	532	726	
vinnulus. (Citellus).....	98		
virgata. (Agouti p.).....	337	408	
virginiana. (Didelphis).....	7		
virginiana. (Didelphys).....	7		1
virginianus. (Canis).....	386		307
virginianus. (Cariacus).....		70	
virginianus. (Cervus).....	43		39
virginianus. (Lepus).....	360		273
virginianus. (Lepus a.).....	360		273
virginianus. (Scalopus).....	469		
virgo. (Dididurus).....	494	615	
virgultus. (Cariacus).....	49		
virgultus. (Odontocælus h.).....	49		
Vison.....	423	530	338
vison. (Mustela).....	423		
vison. (Putorius).....	423		338
vittata. (Mephitis).....	410		
vittata. (Mephitis m.).....	410	512	
vittata. (Viverra).....	524		
vittatus. (Artibeus).....	526	704	
vittatus. (Vampyrops).....	526	704	
vitulina. (Phoca).....	411, 442	541	363
vivax. (Sciurus d.).....	61	102	
Viverra caudivolvula.....	404	499	
Viverra ichneumon.....	373	461	
Viverra mephitis.....	406		
Viverra mungo.....	374		
Viverra narica.....	402		
Viverra nasua.....	402	494	
Viverra piscator.....	422		535
Viverra vittata.....	524		
Viverridæ.....	373	460	
vivesi. (Myotis).....	474	574	
vociferans. (Aotus).....	534	729	
vociferans. (Nyctipithceus).....	534	729	
volans. (Mus).....	125	158	109
volans. (Sciuropterus).....	125	158	109

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
volans. (Sciurus).....	121	157	
volans. (Vespertilio).....	476	578	403
volucella. (Sciurus).....	125	158	109
vomerina. (Phocæna).....	25	48	20
vulcanius. (Reithrodontomys a.) [†]	194	259	
vulcanius. (Rhithrodontomys a.).....	194	259	
vulgaris. (Mustela).....	426		344
vulgaris. (Narwalus).....	24		19
vulgaris. (Sciurus).....	59		
vulgivagus. (Putorius).....	424		
Vulpes.....	380	471	303
Vulpes alascensis.....	381		†434
Vulpes a. abietorum.....	381		†434
Vulpes arsipus.....	385		
Vulpes beringensis.....	383		
Vulpes cascadiensis.....	382		433
Vulpes deletrix.....	381		305
Vulpes fulva.....	380		
Vulpes f. argentata.....	380		
Vulpes f. bangsi.....	381		
Vulpes f. macroura.....	380		
Vulpes f. rubricosa.....	380		
Vulpes hallensis.....	382		307
Vulpes harrimani.....	380		303
Vulpes kenaicensis.....	382		†432
Vulpes lagopus.....	383		306
Vulpes l. innuitus.....	382 , 383		
Vulpes l. ungava.....	383		
Vulpes littoralis.....	388		
Vulpes macrotis.....	384 , 385	472	306
Vulpes m. neomexicana.....	384		
Vulpes macrourus.....	380		305
Vulpes mutica.....	385		
Vulpes muticus.....	385		
Vulpes necator.....	381		†433
Vulpes pennsylvanica.....	380		304
Vulpes p. argentata.....	380		
Vulpes p. bangsi.....	381		
Vulpes p. macroura.....	380		
Vulpes p. rubricosa.....	380		
Vulpes p. vafra.....	380		305
Vulpes pribilofensis.....	383		
Vulpes regalis.....	382		†433
Vulpes rubricosa bangsi.....	381		
Vulpes velox.....	384		306
Vulpes v. hebes.....	384		
vulpes. (Canis).....	380	471	
vulpina. (Mustela).....	419	334	
vulpinus. (Sciurus).....	65, 79		51, 53

†Appendix.

	Check-List of Mammals.	Mammals of Middle America, etc	Synopsis of Mammals, etc.
wagneri. (Dipodomys).....	293		236
wagneri. (Sciurus).....	69	117	
Wagneria.....	397	483	
walkeri. (Delphinus).....	30	55	28
wallawalla. (Lepus t.).....	356		
wapiti. (Cervus).....	41		34
wardi. (Ovibos m.).....	56		*485
washingtoni. (Lepus a.).....	361		274
washingtoni. (Putorius).....	429		347
washingtonii. (Lepus).....	361		274
waterhousii. (Macrotus).....	508	652	
waterhousii. (Otopterus).....	508	652	
watsoni. (Artibeus).....	524	696	
watsoni. (Tylomys).....	173	219	
winingus. (Mustela).....	423		
wortmani. (Citellus).....	107		
wortmani. (Spermophilus).....	107		84
wortmani. (Tamias).....	107		84
wrangeli. (Evotomys).....	229		171
wrangeli. (Synaptomys).....	256, 257		206
xanthinus. (Dasypterus e.).....	487	597	
xanthogenys. (Mustela).....	431		349
xanthogenys. (Putorius).....	431		349
xanthognatha. (Arvicola).....	246		193
xanthognathus. (Microtus).....	246		193
xanthotus. (Sciurus).....	61	104	
xanti. (Lepus c.).....	357	438	
Xenomys.....	224	291	
Xenomys nelsoni.....	224	292	
Xenurus.....	12	31	
Xenurus hispidus.....	12	32	
xenurus. (Peromyscus).....	166	202	
xerampelinus. (Acodon).....	205	275	
xerampelinus. (Akodon).....	205	275	
Xerospermophilus.....	96, 99	139, 143	82
Xylomys.....	316, 323	368, 381	
Xylomys nelsoni.....	323		
yakiensis. (Onychomys t.).....	133	†742	
yakimensis. (Citellus m.).....	112		
yakimensis. (Spermophilus m.).....	112		93
yakutatensis. (Microtus).....	242		188
yapock. (Chironectes).....	1	3	
yelmensis. (Thomomys d.).....	277		226
yucatanense. (Tagassu a.).....	36	63	
yucatanensis. (Cervus).....	46	74	
yucatanensis. (Didelphis).....	5	14	

*Supplement.

†Appendix.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
yucatanensis. (Didelphys).....	5	14	
yucatanensis. (Oryzomys).....	189	236	
yucatanensis. (Sciurus).....	75	125	
yucatanensis. (Sciurus c.).....	75	125	
yucatanensis. (Tayassu a.).....	36	63	
yucatania. (Coendou m.).....	335	402	
yucatania. (Coendu m.).....	335	402	
yucatanica. (Nasua n.).....	403	498	
yucatanicus. (Artibeus).....	523		
yucatanicus. (Lepus f.).....	345	419	
yucatanicus. (Nyctinomops).....	498	626	
yucatanicus. (Peromyscus).....	163	194	
yukonensis. (Lemmus).....	259		208
yukonensis. (Lemmus t.).....	259		208
yukonensis. (Sciuropterus).....	125		†428
yumanensis. (Myotis).....	475	576	403
yumanensis. (Vespertilio).....	475	576	403
zacatecæ. (Neotoma l.).....	212		
zacatecæ. (Perognathus h.).....	310	361	
zacatecæ. (Reithrodontomys m.).....	194	260	
zacatecæ. (Rhithrodontomys m.).....	194	260	
Zalophus.....	437		356
Zalophus californianus.....	437		356
zamelas. (Peromyscus m.).....	161	179	
zamora. (Peromyscus m.).....	166	202	
zaphæus. (Sciuropterus a.).....	122		
Zapodidæ.....	323		256
Zapodina.	323		257
Zapus.....	323, 324		257
Zapus alleni.....	326		261
Zapus hudsonius.....	324		257
Zapus h. alascensis.....	324		250
Zapus h. americanus.....	325		259
Zapus h. campestris.....	325		259
Zapus h. hardyi.....	324		257
Zapus h. ladæ.....	324		258
Zapus imperator.....	326		261
Zapus insignis.....	328		263
Zapus i. abietorum.....	328		264
Zapus i. roanensis.....	328		263
Zapus major.....	326		260
Zapus montanus.....	327		262
Zapus nevadensis.....	326		261
Zapus orarius.....	327		262
Zapus pacificus.....	327		262
Zapus princeps.....	325		260
Zapus p. minor.....	325		260
Zapus p. oregonus.....	326		260

†Appendix.

	Check-List of Mammals.	Mammals of Middle America, etc.	Synopsis of Mammals, etc.
<i>Zapus saltator</i>	327		262
<i>Zapus tenellus</i>	325		259
<i>Zapus trinotatus</i>	326		261
<i>Zapus t. alleni</i>	326, 327		261
<i>Zapus t. montanus</i>	327		262
<i>zarhynchus</i> . (<i>Peromyscus</i>).....	170	209	
<i>zelotes</i> . (<i>Peromyscus</i>).....	166	203	
<i>zibellina</i> . (<i>Mustela</i>).....	419	335	
<i>zibethicus</i> . (<i>Castor</i>).....	252		212
<i>zibethicus</i> . (<i>Fiber</i>).....	252, 254	307	212
<i>Ziphiorrhynchus</i>	22	47	16
<i>Ziphius</i>	22	47	16
<i>Ziphius cavirostris</i>	21, 22	47	17
<i>Ziphius grebnitzkii</i>	23		*480
<i>Zygodontomys</i>	190	252	
<i>Zygodontomys brevicauda</i>		254	
<i>Zygodontomys cherrii</i>	190	253	
<i>Zygodontomys chrysomelas</i>	190	253	
<i>Zygodontomys seorsus</i>	191	254	
<i>Zygogeomys</i>	271	330	
<i>Zygogeomys trichopus</i>	271	332	
<i>zygomatichus</i> . ^a (<i>Oryzomys</i>).....	185	241	

*Supplement.

INDEX OF COMMON NAMES.

	PAGE.		PAGE.
Abert's Squirrel.....	66	Allen's Wood Rat.....	226
Acadian Vole.....	235	Allied Bat.....	481
Acapulco Cotton-tail.....	345	Allied Brown Bat.....	484
Acapulco Squirrel.....	74	Allied Chipmunk.....	92
Active Marten.....	420	Allied Cotton-tail.....	343, 350
Active Pocket Gopher.....	273	Allied Desert Mouse.....	137
Active Wood Rat.....	213	Allied Field Mouse.....	154, 161
Admiralty Island Brown Bear....	392	Allied House Mouse.....	161
Agouta.....	472	Allied Kangaroo Rat.....	289
Agouti, Coiba.....	337	Allied Sapajou.....	536
Agouti, Crested.....	337	Allied Spiny Rat.....	317
Agouti, Cunning.....	336	Allied Weasel.....	433
Agouti, Isthmian.....	336	Allied Wood Rat.....	207
Agouti, Mexican.....	336	Almiqui.....	472
Agouti, Ruatan Island.....	330	Alpine Chipmunk.....	94
Agouti, Spotted.....	336	Alpine Cotton Rat.....	179
Agoutis.....	335	Alpine Harvest Mouse.....	200
Akeley's Field Mouse.....	148	Alpine Mole.....	469
Alaska Bat.....	479	Alpine Pocket Gopher.....	283
Alaska Jumping Mouse.....	325	Alston's Mouse.....	204
Alaska Porcupine.....	333	Alston's Opossum.....	4
Alaska Red Fox.....	381	Alston's Wood Rat.....	225
Alaska Red-backed Vole.....	230	Alta Mira Cotton-tail.....	343
Alaska Swimming Shrew.....	458	Alta Mira Jack Rabbit.....	354
Alaskan Grizzly Bear.....	393	Ameca Harvest Mouse.....	202
Alaskan Moose.....	41	American Badger.....	405
Alaskan Weasel.....	426	American Black Bear.....	394
Albemarle Meadow Vole.....	235	American Buffalo.....	57
Alberta Black Bear.....	396	American Elk.....	41, 42
Alberta Spermophile.....	111	American Hare.....	360
Alfaro's Pigmy Squirrel.....	60	American Marten.....	419
Alfaro's Rice Rat.....	183, 191	Amoles Cotton Rat.....	179
Alien Mouse.....	169	Anastasia Island Mouse.....	138, 144
Allamand's Grison.....	419	Ancient Pocket Gophers.....	268
Alleghany Mountains Weasel.....	428	Anderson's Lemming Vole.....	257
Allen's Cotton Rat.....	179	Angular Pocket Gopher.....	281
Allen's Jack Rabbit.....	353	Anomalous Shrews.....	459
Allen's Jumping Mouse.....	327	Anteater, Chiriqui.....	11
Allen's Opossum.....	5	Anteater, Great.....	12
Allen's Pocket Mouse.....	315	Anteater, Little or Two-toed.....	10
Allen's Spermophile.....	105	Anteater, Saddle-backed.....	12
Allen's Spiny Mouse.....	321	Anteater, Slender Nose.....	11
Allen's Squirrel.....	63	Anteater, Three-toed.....	11
Allen's Weasel.....	431	Anteater, Ursine.....	10

	PAGE.		PAGE.
Anteaters	10	Autocrat Timber Wolf.....	374
Antelope, Prong-horn.....	51	Aztec Kinkajou.....	404
Anthony's Mouse.....	137	Aztec Meadow Vole.....	237
Anthony's Pocket Mouse.....	314	Aztec Mouse.....	162
Antilles Bat.....	517	Aztec Rice Rat.....	184
Antique Bat.....	507		
<i>Apache</i>	368	Bachman's Cotton-tail	352
Apache Pocket Mouse.....	303	Badger, American	405
Apache Squirrel.....	64	Badger, Berlandier's.....	405
Apazote Mouse.....	163	Badger, California.....	405
Apazote Rat.....	174	Badger, Lower California.....	405
Apazote Squirrel.....	61	Badgers.....	404
Aquatic Rice Rat.....	182	Bahama Brown Bat	484
Arctic Baleen Whale.....	17	Bahama Long-eared Bat	509
Arctic Caribou.....	39	Bahaman Free-tailed Bat.....	500
Arctic Field Mouse.....	146	Bailey's Chickaree.....	81
Arctic Right Whale.....	16	Bailey's Cotton Rat.....	177
Arctic Shrew	447	Bailey's Cotton-tail.....	349
Arctic Weasel.....	428	Bailey's Lynx.....	372
Arctic Wolf.....	376	Bailey's Pocket Gopher	285
<i>Ardilla</i>	68	Bailey's Pocket Mouse	309
Arizona Bushy-tailed Wood Rat..	223	Bailey's Spermophile.....	100
Arizona Chickaree.....	84	Bailey's Wood Rat	210
Arizona Cotton Rat.....	176	Baird's Hare.....	361
Arizona Cotton-tail.....	349	Baird's Kangaroo Rat.....	295
Arizona Gray Squirrel.....	64	Baird's Meadow Vole	250
Arizona Harvest Mouse.....	196	Baird's Pocket Mouse	302, 307
Arizona Meadow Vole.....	238	Baird's Shrew.....	452
Arizona Prairie-dog	118	Baird's Sperm Whale	24
Arizona Skunk.....	408	Baird's 'Tapir'	58
Arizona Spotted Skunk.....	414	Baleen Whales.....	15
Arizona Weasel.....	431	Banded-back Squirrel.....	76
Arizona White-footed Deer Mouse.	141	Banderas Field Mouse.....	163
Armadillo, Miller's.....	13	Bangs' Cotton-tail.....	343
Armadillo, Nine-banded.....	13	Bangs' Field Mouse.....	144
Armadillos	12	Bangs' Flying Squirrel.....	124
Armed Spermophile.....	112	Bangs' Pocket Mouse	305
Armeria Peccary.....	35	Bangs' Polar Hare.....	358
Ash-colored Chipmunk.....	90	Bangs' Red Fox.....	381
Ash-colored Cotton-tail.....	352	Bangs' Red Mouse.....	205
Ashy Gray Field Mouse.....	164	Barbadoes Nose-leaf Bat	518
Ashy Opossum.....	2	Barber's Squirrel.....	68
Ashy Spermophile.....	106	Barking Coyote	376
Atemajac Wood Rat.....	211	Barred Lynx.....	371
Atlantic Walrus.....	439	Barren Ground Caribou.....	40
Attwater's Field Mouse.....	154	Barren Ground Grizzly Bear.....	393
Attwater's Pocket Gopher.....	263	Barren Ground Meadow Vole	237
Attwater's Swamp Hare.....	340	Barrow's Spermophile.....	110
Attwater's Wood Rat.....	210	Bat, Alaska.....	479
Audubon's Cotton-tail.....	345	Bat, Allied.....	481
Audubon's Mountain Sheep.....	53	Bat, Allied Brown.....	484

	PAGE.		PAGE.
Bat, Antilles.....	517	Bat, Flat-nosed.....	524
Bat, Antique.....	507	Bat, Florida Brown.....	483
Bat, Bahama Brown.....	484	Bat, Florida Red.....	486
Bat, Bahama Long-eared.....	509	Bat, Forest.....	513
Bat, Bahaman Free-tailed.....	500	Bat, Fort Yuma.....	475
Bat, Barbadoes Nose-leaf.....	518	Bat, Free-tailed Mouse.....	501
Bat, Bayamon.....	520	Bat, Fringed.....	479
Bat, Big-eared.....	490, 507	Bat, Fringed-faced.....	514
Bat, Big-eared Pale.....	491	Bat, Fruit-loving.....	527
Bat, Blackish Mastiff.....	496	Bat, Geoffroy's.....	520, 529
Bat, Black-winged.....	483	Bat, Godman's.....	517
Bat, Bocourt's Large-eared.....	510	Bat, Graceful.....	502
Bat, Booth's.....	505	Bat, Graceful Free-tailed.....	499
Bat, Boquete Tailless.....	531	Bat, Graham Island.....	477
Bat, Brown.....	483	Bat, Gray.....	504
Bat, Buller's Large-eared.....	511	Bat, Great.....	501
Bat, California Large-eared.....	510	Bat, Grizzled Brown.....	475
Bat, California Mastiff.....	498	Bat, Gundlach's.....	488
Bat, California Red.....	486	Bat, Hairy.....	512
Bat, Cardonal Islands.....	474	Bat, Hairy-lipped.....	477
Bat, Cavern Nose-leaf.....	522	Bat, Heedless.....	473
Bat, Chestnut-colored.....	516	Bat, Heller's White-striped.....	526
Bat, Chestnut Mastiff.....	497	Bat, Hoary.....	487
Bat, Chiriqui.....	475	Bat, Inflated-nose.....	504
Bat, Cinerous.....	525	Bat, Intermediate.....	487, 523
Bat, Cinnamon.....	482, 507	Bat, Jalisco Brown.....	477
Bat, Coiba Island.....	496	Bat, Jamaica.....	522
Bat, Colon.....	524	Bat, Jamaica Large-eared.....	508
Bat, Cory's.....	522	Bat, Keene's Brown.....	478
Bat, Cuban Brown.....	484	Bat, La Grulla Brown.....	476
Bat, Cuban Free-tailed.....	499	Bat, Large-eared Vampire.....	513
Bat, Cuban Large-eared.....	509	Bat, Large-nosed Mastiff.....	497
Bat, Cuban Nose-leaf.....	519	Bat, Large-winged.....	473
Bat, Dark Brown.....	505	Bat, Las Penas.....	506
Bat, Dark-eared.....	525	Bat, Las Vigas.....	482
Bat, Dark Yuma.....	476	Bat, Leafless or Blunt-nosed.....	521
Bat, Davy's.....	506	Bat, Little Brown.....	479
Bat, De Blainville's.....	507	Bat, Little California.....	476
Bat, Disk-bearing.....	503	Bat, Little Comodu.....	492
Bat, Dog-headed Free-tailed.....	500	Bat, Little Free-tailed.....	501
Bat, Dog-like.....	494	Bat, Little Large-eared.....	511
Bat, Dominican Common.....	475	Bat, Little Mexican.....	477
Bat, Dominican Free-tailed.....	499	Bat, Long-nosed.....	492
Bat, Durango Brown.....	478	Bat, Lower California.....	474
Bat, Dusky.....	481, 504, 520	Bat, Lower California Brown.....	484
Bat, Dusky Mastiff.....	496	Bat, Macleay's.....	504
Bat, Dwarf Mastiff.....	498	Bat, Maria Madre Island.....	516
Bat, Dwarf Nose-leaf.....	522	Bat, Masked.....	505
Bat, Escazu White.....	494	Bat, Mexican.....	506
Bat, Falcate.....	527	Bat, Mexican Free-tailed.....	500
Bat, Flat Forehead Bahama.....	521	Bat, Mexican Red.....	486

	PAGE.		PAGE.
Bat, Mexican Vampire.....	513	Bat, Small-winged.....	478
Bat, Miller's.....	474	Bat, Snowy.....	519
Bat, Miller's Florida.....	488	Bat, Spear-nosed.....	514
Bat, Miller's Nose-leaf.....	519	Bat, Specter.....	511
Bat, Mirador Brown.....	484	Bat, Spotted Great-eared.....	490
Bat, Montserrat.....	526	Bat, Straight-eared Mastiff.....	497
Bat, Musky.....	495	Bat, Straw-colored.....	501
Bat, Nicholls'.....	527	Bat, Swift.....	481
Bat, Night-wandering.....	480	Bat, Tailless.....	531
Bat, Pacific Big-eared.....	492	Bat, Thomas'.....	493
Bat, Pale.....	477	Bat, Tomes' Long-eared.....	508
Bat, Pallid Big-eared.....	491	Bat, Townsend's Big-eared.....	491
Bat, Palm Springs Free-tailed.....	498	Bat, Tres Marias Islands.....	489, 517
Bat, Panama.....	488	Bat, True's.....	479
Bat, Panama Spear-nosed.....	514	Bat, Tucubaya Free-tailed.....	499
Bat, Parnell's.....	505	Bat, Underwood's.....	518
Bat, Peters' Vampire.....	512	Bat, Wandering Bermuda.....	482
Bat, Peters' White-striped.....	526	Bat, Waterhouse's Large-eared.....	508
Bat, Pfeiffer's.....	486	Bat, Watling's Island.....	503
Bat, Poey's.....	421	Bat, Watson's.....	524
Bat, Porto Rican.....	505	Bat, Western.....	480
Bat, Porto Rico Nose-leaf.....	518	Bat, White.....	494
Bat, Pouched.....	493	Bat, White Honduras.....	528
Bat, Prominent-eared.....	474	Bat, White-striped.....	493, 525
Bat, Pug-nosed Mastiff.....	496	Bat, White-throated Brown.....	485
Bat, Rafinesque's.....	488	Bat, Wrinkled-face.....	530
Bat, Red.....	485	Bat, Yellowish-brown.....	481
Bat, Redman's Nose-leaf.....	519	Bat, Ypanema Nose-leaf.....	528
Bat, Restless.....	489	Bat, Yucatan.....	523
Bat, Round-eared.....	512	Bat, Yucatan Brown.....	485
Bat, Rufous Mastiff.....	495	Bat, Yucatan Free-tailed.....	498
Bat, Rufous Vampire.....	530	Batopilas Pocket Mouse.....	314
Bat, St. Kitts.....	523	Bats.....	473
Bat, St. Martin's.....	524	Bats, Big-eared.....	490
Bat, Salvin's Nose-leafed.....	528	Bats, Club-nosed.....	490
Bat, San Bernardino Brown.....	484	Bats, Common.....	473
Bat, Santa Anita.....	502	Bats, Funnell-eared.....	501
Bat, Santa Lucia.....	527	Bats, Vampire.....	503
Bat, Santa Lucia Nose-leaf.....	518	Batty's Jack Rabbit.....	355
Bat, Santa Rosalia.....	489	Batty's Opossum.....	8
Bat, Saussure's Large-eared.....	510	Batty's White-tailed Deer.....	45
Bat, Sezekorn's.....	521	Bay Spermophile.....	105
Bat, Shaggy-eared.....	505	Bayamon Bat.....	520
Bat, Shaved Mastiff.....	497	Beach Meadow Vole.....	238
Bat, Short-handed.....	503	Beaked Sperm Whale.....	22
Bat, Short-tailed.....	515	Bear, Admiralty Island Brown.....	392
Bat, Shrew-like.....	516	Bear, Alaskan Grizzly.....	393
Bat, Sierra Laguna.....	487	Bear, Alberta Black.....	396
Bat, Small.....	517	Bear, American Black.....	394
Bat, Small-eared Nicaraguan.....	513	Bear, Barren Ground Grizzly.....	393
Bat, Small-footed.....	502, 523	Bear, Dall's Brown.....	391

	PAGE.		PAGE.
Bear, Dark-clawed Grizzly.....	393	Bishop's Hare.....	362
Bear, Desert Black.....	395	Bison.....	57
Bear, Emmons' Glacier.....	397	Bison, Wood.....	57
Bear, Fighting.....	396	Black Bears.....	394
Bear, Florida Black.....	395	Black Fox.....	380
Bear, Giant Brown.....	391	Black Fox Squirrel.....	66
Bear, Inland White.....	397	Black Musk Ox.....	56
Bear, Kenai Peninsula Brown.....	392	Black Pocket Gopher.....	276
Bear, Kidder's Brown.....	391	Black Pocket Mouse.....	301
Bear, Louisiana Black.....	395	Black Rat.....	129
Bear, Mexican Grizzly.....	393	Black Rice Rat.....	190
Bear, Middendorff's Brown.....	391	Black Spider Monkey.....	535
Bear, Northwestern Black.....	395	Black Spiny Rat.....	317
Bear, Polar.....	390	Black-backed Squirrel.....	62
Bear, Queen Charlotte Islands Black.....	394	Black-browed Mole Mouse.....	131
Bear, Sornborger's Black.....	394	Black-browed Mouse.....	166
Bearded Mole Mouse.....	134	Black-eared Cotton Rat.....	180
Bearded Seal.....	441	Black-eared Field Mouse.....	161
Bearing Island Artic Fox.....	383	Black-eared Jack Rabbit.....	355
Bears.....	390	Black-eared Pocket Mouse.....	303
Bears, Black.....	394	Black-eared Rice Rat.....	187
Bears, Brown.....	390	Black-faced Pocket Gopher.....	279
Bears, Grizzly.....	392, 393	Blackfish.....	27
Bears, Sea.....	438	Blackfish, Gray.....	28
Beautiful Mouse.....	154	Blackfish, Scammon's.....	28
Beautiful Pocket Mouse.....	303	Blackfish, Short-finned.....	28
Beautiful Wood Rat.....	218	Black-footed Gray Squirrel.....	78
Beautiful-eared Jack Rabbit.....	353	Black-footed Lemming.....	260
Beautifully Garbed Chipmunk.....	89	Black-footed Mink.....	425
Beaver, Canada.....	127	Black-footed Mouse.....	170
Beaver, Carolina.....	128	Black-headed Sperrophile.....	114
Beaver Mountains Chipmunk.....	95	Blackish Field Mouse.....	156
Beaver, Pacific Coast.....	129	Blackish Mastiff Bat.....	496
Beaver, Sonoran.....	128	Blackish Pocket Gopher.....	280
Beaver Valley Chipmunk.....	95	Blackish Shrew.....	463
Beavers.....	127	Black-nosed Kangaroo Rat.....	290
Beechey's Sperrophile.....	114	Black-tailed Chipmunk.....	94
Belding's Sperrophile.....	113	Black-tailed Deer.....	48
Bendire's Shrew.....	459	Black-tailed Hutia.....	331
<i>Berendo</i>	51	Black-tailed Kangaroo Rat.....	291
Bering's Sperrophile.....	110	Black-tailed Mouse.....	137
Berlandier's Badger.....	405	Black-tailed Wood Rat.....	217
Berlandier's Cotton Rat.....	176	Black-winged Bat.....	483
Berlandier's Shrew.....	462	Blue Finback Whale.....	20
Big-eared Bat.....	490, 507	Blue Mountains Jumping Mouse.....	326
Big-eared Bats.....	490	Bluegrass Meadow Vole.....	249
Big-eared Harvest Mouse.....	193	Bocourt's Large-eared Bat.....	510
Big-eared Mouse.....	157	Bogava Rice Rat.....	189
Big-eared Pale Bat.....	491	Bogava Spiny Rat.....	329
Big-footed Meadow Vole.....	251	Bolaños Wood Rat.....	205
		Bold Coyote.....	379

	PAGE.		PAGE.
Booth's Bat	505	Buff-cheeked Chipmunk	87
Boqueron Coati	403	Buff-colored Pocket Mouse	309
Boqueron Cotton Rat	178	Buff-colored Wolverine	417
Boqueron Jaguarondi	368	Buller's Large-eared Bat	511
Boquete Field Mouse	172	Buller's Pocket Gopher	268
Boquete Mouse	204	Buller's Rice Rats	185
Boquete Peccary	36	Buller's Spiny Rat	317
Boquete Pocket Gopher	270	Bushy-tailed Spermophile	114, 149
Boquete Raccoon-fox	399	Bushy-tailed Wood Rats	220
Boquete Spiny Rat	318		
Boquete Tailless Bat	531	Cabezon Kangaroo Rat	295
Boquete Vesper Rat	173	Cabezon Pocket Gopher	287
Boreal Flying Squirrel	124	<i>Cabra del Monte</i>	50
Boruca Cotton Rat	178	Cachalot Sperm Whale	21
Botta's Pocket Gopher	274	<i>Cacomistl</i>	397
Bottle-nosed Porpoise	32	<i>Cacomistl de Monte</i>	399
Bottle-nosed Whales	29	Calé Harvest Mouse	201
Bowhead	16	California Badger	405
Boyle's Field Mouse	150	California Chickaree	83
Brewer's Hairy-tailed Mole	471	California Gray Fox	387
Bridger's Pocket Gopher	279	California Gray Squirrel	78
Bridled Dolphin	34	California Jack Rabbit	357
Bridled Skunk	410	California Kangaroo Rat	292, 300
Bridled Weasel	432	California Large-eared Bat	510
Bristled Pocket Gopher	269	California Lynx	372
British Columbia Bushy-tailed Wood Rat	221	California Mastiff Bat	498
British Columbian Hare	361	California Meadow Vole	240
Broad-banded Cotton-tail	346	California Mole	468
Broad-footed Pocket Gopher	271	California Mouse	138
Broad-headed Pocket Gopher	275	California Mule Deer	49
Broad-headed Pocket Gophers	266	California Pocket Mouse	315
Broad-headed Spotted Skunk	414	California Red Bat	486
Broad-nosed Pocket Gopher	286	California Red-backed Vole	233
Broad-nosed Pocket Mouse	313	California Sea-lion	437
Broad-nosed Rice Rat	188	California Sewellel	127
Broad-nosed Skunk	409	California Shrew	452
Brocket, Sartori's	50	California Skunk	408
Brocket, Tunkas	51	Californian Flying Squirrel	122
Brockets	50	<i>Callitriche</i>	536
Brown Bat	483	Calovevora Grison	418
Brown Rat	130	<i>Camaleon</i>	9
Brown-rumped Cotton Rat	175	Cameron Lake Wood Rat	215
Brown's Pigmy Squirrel	60	Canada Beaver	127
Brownsville Harvest Mouse	198	Canada Lynx	370
Bryant's Pocket Mouse	316	Canada Otter	433
Buckley's Spermophile	115	Canada Skunk	406
Buff Mouse	147	Canadian Field Mouse	144
Buffalo, American	57	Canadian Tree Porcupine	334
Buff-bellied Chipmunk	92	Canadian Woodchuck	120
Buff-bellied Pocket Mouse	301	Cantankerous Meadow Vole	244
		Cape Sable Rat	181

	PAGE.		PAGE.
Cape St. Lucas Pocket Mouse.....	316	Chapman's Cotton-tail.....	343
Cape St. Lucas Spotted Skunk....	416	Chapman's Kangaroo Rat.....	297
Captious Harvest Mouse.....	197	Chapman's Lemming Vole.....	257
Capuchin, White-throated.....	536	Chapman's Mouse.....	139
<i>Carablanca, Mono</i>	536	Chapman's Rice Rat.....	186
Cardonal Island Bat.....	474	Charming Chipmunk.....	90
<i>Caribbanco</i>	37	Charming Shrew.....	449
Caribou.....	37, 38	Charming Spermophile.....	98
Caribou, Arctic.....	39	Charming Striped Chipmunk.....	96
Caribou, Barren Ground.....	39	Cheating Spotted Skunk.....	414
Caribou, Grant's.....	39	Cheating Wood Rat.....	225
Caribou, Greenland or Artic.....	39	Cherrie's Costa Rica Mouse.....	160
Caribou, Mountain.....	38	Cherrie's Pocket Gopher.....	271
Caribou, Newfoundland.....	38	Cherrie's Rice Rat.....	190
Caribou, Osborn's.....	38	Chestnut Mastiff Bat.....	497
Caribou, Peary's.....	40	Chestnut-bellied Shrew.....	450
Caribou, Stone's.....	39	Chestnut-brown Weasel.....	430
Carnivores.....	365	Chestnut-colored Bat.....	516
Carolina Beaver.....	128	Chestnut-faced Pocket Gopher....	266
Carolina Gray Squirrel.....	78	Chestnut-headed Sloth.....	9
Carolina Red-backed Vole.....	232	Chestnut-tailed Spermophile.....	106
Carolina Shrew.....	461	Chiapas Cotton-tail.....	344
Carolinian Jumping Mouse.....	325	Chiapas Squirrel.....	74
Carriker's Ocelot.....	366	Chichen Itza Harvest Mouse.....	198
Cascade Mountain Meadow Vole...	251	Chichen Itza Rice Rat.....	189
Cascade Mountains Chickaree....	83	Chickaree, Arizona.....	84
Cascade Mountains Fox.....	382	Chickaree, Bailey's.....	81
Cascade Skunk.....	408	Chickaree, California.....	83
Cat, Duncce-like.....	367	Chickaree, Cascade Mountains....	83
Cat, Small Spotted.....	367	Chickaree, Douglas.....	82
<i>Cat Squirrel</i>	397	Chickaree, Eastern.....	80
Cat, Wild.....	371	Chickaree, Fremont's.....	84
Catalina Island Mouse.....	149	Chickaree, Graham Mountains....	85
Catemaco Rice Rat.....	185	Chickaree, Hudson Bay.....	80
Catemaco Spiny Mouse.....	320	Chickaree, Mearns'.....	83
Cats.....	365	Chickaree, Minnesota.....	81
<i>Cauzel</i>	367	Chickaree, Noisy.....	80
Cavern Nose-leaf Bat.....	522	Chickaree, Richardson's.....	82
Cecil's Mouse.....	161	Chickaree, Ruidoso.....	84
Ceiba Cotton Rat.....	178	Chickaree, Scolding.....	82
Central American Otter.....	435	Chickaree, Soft-furred.....	83
Central American Paca.....	337	Chickaree, South Dakota.....	81
Central American Puma.....	370	Chickaree, Streater's.....	82
Cerros Island Cotton-tail.....	352	Chickaree, Taos.....	84
Cerros Island Mouse.....	155	Chickaree, Vancouver Island....	82
Cerros Island White-tailed Deer...	47	Chickaree, Wind River Mountains.	81
Cerros Island Wood Rat.....	208	Chief Hares.....	337
Cervine Pocket Gopher.....	287	Chief Pika.....	338
Cetaceans.....	15	Chihuahua Mule Deer.....	50
Changeable Hares.....	358	Chihuahuan Little Spotted Skunk.	415
Changeable Shrew.....	456	<i>Chillon, o' Mono</i>	533

	PAGE.		PAGE.
Chilpancingo Mouse.....	168	Chiriqui Pigmy Squirrel.....	60
Chipmunk, Allied.....	92	Chiriqui Porcupine.....	335
Chipmunk, Alpine.....	94	Chiriqui Squirrel.....	62
Chipmunk, Ash-colored.....	90	Chubby Mouse.....	168
Chipmunk, Beautifully Garbed.....	89	<i>Chulomuco</i>	418, 419
Chipmunk, Beaver Mountains.....	95	Cinereous Bat.....	525
Chipmunk, Beaver Valley.....	95	Cinereous Harvest Mouse.....	197
Chipmunk, Black-tailed.....	94	Cinnamon Bat.....	482, 507
Chipmunk, Buff-bellied.....	92	Cinnamon Bushy-tailed Wood Rat.....	224
Chipmunk, Buff-cheeked.....	87	Cinnamon Chief Hare.....	338
Chipmunk, Charming.....	90	Cinnamon-hued Spermophile.....	97
Chipmunk, Charming Striped.....	96	Civets.....	373
Chipmunk, Coast.....	87	Club-nosed Bats.....	490
Chipmunk, Colorado.....	91	Coast Chipmunk.....	87
Chipmunk, Dark.....	91	Coast Cotton Rat.....	175
Chipmunk, Durango.....	88	Coast Jumping Mouse.....	327
Chipmunk, Four-banded.....	89	Coast Meadow Vole.....	241, 244
Chipmunk, Gila.....	85	Coast Mole.....	468
Chipmunk, Graceful.....	91	Coast Wood Rat.....	209
Chipmunk, Gray.....	89	Coati.....	402
Chipmunk, Gray Striped.....	96	Coati, Boqueron.....	403
Chipmunk, Gray-footed.....	91	Coati, Dark.....	403
Chipmunk, Gray-headed.....	93	Coati, Manzanillo.....	403
Chipmunk, Gray-tailed.....	93	Coati, Nelson's.....	402
Chipmunk, Guanacevi.....	86	Coati, Pallid.....	403
Chipmunk, Happy.....	92	Coati, Yucatan.....	403
Chipmunk, Hind's.....	86	Coatis.....	397, 402
Chipmunk, Inyo Mountains.....	89	Coban Spiny Rat.....	318
Chipmunk, Least.....	93	Cofre de Perote Harvest Mouse.....	199
Chipmunk, Lower California.....	88	Coiba Agouti.....	337
Chipmunk, Lyster's Striped.....	95	Coiba Island Bat.....	496
Chipmunk, Merriam's.....	88	Colima City Mouse.....	136
Chipmunk, Mexican.....	88	Colima Coyote.....	378
Chipmunk, Mountain.....	95	Colima Harvest Mouse.....	200
Chipmunk, Neglected.....	93	Colima or Dark Meadow Vole.....	245
Chipmunk, Northern.....	92	Colima Mountain Squirrel.....	71
Chipmunk, Northwest.....	90	Colima Squirrel.....	72
Chipmunk, Painted.....	94	<i>Collaraja</i>	432, 433
Chipmunk, Palmier's.....	86	Collared Pika.....	338
Chipmunk, Panamint Mountains.....	90	Collared Wood Rat.....	208
Chipmunk, Price's.....	87	Collector Wood Rat.....	205
Chipmunk, Related.....	90, 94	Collie's Squirrel.....	73
Chipmunk, Showy.....	89	Colon Bat.....	524
Chipmunk, Striped.....	95	Colorado Bushy-tailed Wood Rat.....	223
Chipmunk, Townsend's.....	86	Colorado Chipmunk.....	91
Chipmunk, Utah.....	86	<i>Colorado, Mono</i>	535
Chipmunks.....	85	Columbian Black-tailed Deer.....	48
Chiriqui Anteater.....	11	Columbian Kangaroo Rat.....	297
Chiriqui Bat.....	475	Columbian Mountain Goat.....	52
Chiriqui Harvest Mouse.....	194	Columbian Pocket Mouse.....	308
Chiriqui Kinkajou.....	404	Columbian Spermophile.....	111

	PAGE.		PAGE.
Common Bats.....	473	Cotton-tail, Lesser Desert.....	350
Common Cotton-tail.....	342	Cotton-tail, Little.....	348
Common Dolphin.....	31	Cotton-tail, Lively.....	343
Common Flying Squirrel.....	125	Cotton-tail, Lower California.....	352
Common Indian Mongoose.....	374	Cotton-tail, Mearns'.....	342
Common Killer Whale.....	26	Cotton-tail, Michoacan.....	344
Common Mink.....	423	Cotton-tail, Mixed.....	347
Common Mole.....	470	Cotton-tail, Nuttall's.....	348
Common Moles.....	469	Cotton-tail, Omilteme.....	342
Common Porpoise.....	25	Cotton-tail, Orizaba.....	351
Common Raccoon.....	400	Cotton-tail, Pine Wood.....	350
Common Raccoon-fox.....	397	Cotton-tail, Plains.....	348
Common Shrews.....	446	Cotton-tail, Puebla.....	345
Common Weasel.....	426	Cotton-tail, Rich-colored.....	352
<i>Conejo</i>	341, 352	Cotton-tail, Russet.....	348
<i>Congo</i>	533	Cotton-tail, Russet-footed.....	346
Constable's Vole.....	228	Cotton-tail, San Diego.....	346
Coolidge's Field Mouse.....	147	Cotton-tail, Sinaloa.....	349
Coon, Gabb's.....	400	Cotton-tail, Tehuantepec.....	344
Cooper's Lemming Vole.....	255	Cotton-tail, Vera Cruz.....	347
Cope's Baleen Whale.....	17	Cotton-tail, Yucatan.....	345
Cope's Finback Whale.....	19	Cotton-tails, Short-tailed.....	351
Cope's Killer Whale.....	26	Cotton Tree Mouse.....	143
Copper-colored Mole.....	471	<i>Cotusa</i>	336
Cory's Bat.....	522	Coues' Mexican Shrew.....	464
Costa Rica Harvest Mouse.....	200	Coues' Pocket Gopher.....	284
Costa Rica Jaguar.....	365	Coues' Pocket Mouse.....	307
Costa Rica Ocelot.....	366	Coues' Rice Rat.....	186
Costa Rica White-tailed Deer.....	46	Coues' Tree Porcupine.....	333
Cotton Rats.....	174	Coues' White-tailed Deer.....	45
Cotton-tail, Acapulco.....	348	Counterfeit Kangaroo Rat.....	295
Cotton-tail, Allied.....	343, 350	Cow Fish.....	32
Cotton-tail, Alta Mira.....	343	Cow, Steller's Sea.....	14
Cotton-tail, Arizona.....	349	Coyote, Barking.....	377
Cotton-tail, Ash-colored.....	352	Coyote, Bold.....	379
Cotton-tail, Audubon's.....	345	Coyote, Colima.....	378
Cotton-tail, Bachman's.....	352	Coyote, Goldman's.....	378
Cotton-tail, Bailey's.....	349	Coyote, Lower California.....	378
Cotton-tail, Bangs'.....	343	Coyote, Mearns'.....	377
Cotton-tail, Broad-banded.....	346	Coyote, Nebraska.....	377
Cotton-tail, Cerros Island.....	352	Coyote, Noland's Ranch.....	378
Cotton-tail, Chapman's.....	343	Coyote, Ochraceous-footed.....	379
Cotton-tail, Chiapas.....	344	Coyote, Robber.....	377
Cotton-tail, Common.....	342	Coyote, Smith's.....	378
Cotton-tail, Florida.....	342	Coyote, Tamaulipas.....	379
Cotton-tail, Granger's.....	349	Coyote, Thievish.....	378
Cotton-tail, Grayson's.....	351	Coyote, Tricky.....	377
Cotton-tail, Greater Desert.....	350	Cozumel Island Rice Rat.....	182
Cotton-tail, Hoary-rumped.....	344	Crab-eating Raccoon.....	402
Cotton-tail, Holzner's.....	345	Crawford's Shrew.....	459
Cotton-tail, Idaho.....	351	Creeping Meadow Vole.....	250

	PAGE.		PAGE.
Crested Agouti.....	337	Deer, Chihuahua Mule.....	50
Crested Pocket Gopher.....	263	Deer, Columbian Back-tailed.....	48
Crested-tailed Mouse.....	167	Deer, Costa Rica White-tailed.....	46
Crook's Black-tailed Deer.....	48	Deer, Coues' White-tailed.....	45
Cuban Brown Bat.....	484	Deer, Crook's Black-tailed.....	48
Cuban Free-tailed Bat.....	499	Deer, Desert Mule.....	50
Cuban Large-eared Bat.....	509	Deer, Hamilton Smith's White-	
Cuban Nose-leaf Bat.....	519	tailed.....	46
<i>Cuistiti</i>	534	Deer, Louisiana White-tailed.....	44
Cumberland Island Field Mouse...	144	Deer, Lower California Mule.....	50
Cumberland Pocket Gopher.....	262	Deer, Mexican White-tailed.....	45
Cunning Agouti.....	336	Deer, Minnesota Mule.....	49
Cunning Red-backed Vole.....	233	Deer, Mule.....	49
Curly Spiny Mouse.....	320	Deer, Nelson's White-tailed.....	47
Curtailed Fox.....	385	Deer, Northern White-tailed.....	44
		Deer, Osceola's White-tailed.....	44
Dainty Pika.....	399	Deer, Rafinesque's White-tailed...	44
Dakota Woodchuck.....	121	Deer, Rothschild's White-tailed...	46
Dall's Brown Bear.....	391	Deer, Sinaloa White-tailed.....	47
Dall's Hare.....	361	Deer, Stika Black-tailed.....	48
Dall's Lemming Vole.....	258	Deer, Texan White-tailed.....	45
Dall's Mountain Sheep.....	55	Deer, Thomas' White-tailed.....	47
Dall's Porpoise.....	26	Deer, True's White-tailed.....	46
Dark Brown Bat.....	505	Deer, Virginian White-tailed.....	43
Dark Chipmunk.....	91	Deer, Yucatan White-tailed.....	47
Dark Coati.....	403	Degenerate Otter.....	435
Dark Field Mouse.....	155	Delicate Shrew.....	453
Dark Sewellel.....	127	Delving Pocket Gopher.....	287
Dark Sperophile.....	106	Deppe's Squirrel.....	60
Dark Spotted Sperophile.....	100	Desert Black Bear.....	395
Dark Yuma Bat.....	476	Desert Jack Rabbit.....	356
Dark-clawed Grizzly Bear.....	393	Desert Kangaroo Rat.....	292
Dark-eared Bat.....	525	Desert Lynx.....	372
Dark-hued Mink.....	425	Desert Meadow Vole.....	241
Dark-tailed Spiny Mouse.....	319	Desert Mouse.....	137
Davidson's Finback Whale.....	19	Desert Mule Deer.....	50
Davy's Bat.....	506	Desert Pocket Gopher.....	281
Dawson's Red-backed Vole.....	229	Desert Pocket Mouse.....	311
Death Valley Kangaroo Rat.....	290	Desert Sperophile.....	101
Death Valley Sperophile.....	101	Desert White-footed Mouse.....	140
Death Valley Wood Rat.....	215	Destructive Fox.....	381
De Blainville's Bat.....	507	Devilish Mink.....	424
Deceitful Pocket Gopher.....	264	Dickinson's Harvest Mouse.....	192
Deceitful Rice Rat.....	185	Digger Pocket Gopher.....	269, 274
Deceitful Wood Rat.....	216	Diligent Harvest Mouse.....	191
Decorated Weasel.....	431	Disk-bearing Bat.....	503
Deer.....	43	Dismal Swamp Shrew.....	463
Deer, Batty's White-tailed.....	45	Dissimilar Wood Rat.....	206
Deer, Black-tailed.....	48	Distinct Pocket Gophers.....	269
Deer, California Mule.....	49	Dobson's Shrew.....	448
Deer, Cerros Island White-tailed..	47	Dobson's Swimming Shrew.....	458

	PAGE.		PAGE.
Dog-headed Free-tailed Bat.....	500	Dwarf Meadow Vole.....	239
Dog-like Bat.....	494	Dwarf Mouse.....	135
Dogs.....	374	Dwarf Nose-leaf Bat.....	522
Dolphin.....	32	Dwarf Peccary.....	35
Dolphin, Bridled.....	34	Dwarf Shrew.....	453
Dolphin, Common.....	31	Dyche's Harvest Mouse.....	192
Dolphin, Gill's.....	32		
Dolphin, Gray's.....	34	Earl of Derby's Opossum.....	4
Dolphin, Long-nosed.....	33	Eastern Chickaree.....	80
Dolphin, Pacific Ocean.....	32	Eastern Desert Pocket Mouse.....	312
Dolphin, Sharp-toothed.....	33	Eastern Fox Squirrel.....	65
Dolphin, Temporal.....	33	Eastern Lynx.....	370
Dolphins.....	24, 30	Eastern Puma.....	369
Dominican Common Bat.....	475	Eastern Skunk.....	406
Dominican Free-tailed Bat.....	499	Eastern Swamp Hare.....	341
Donkey Jack Rabbit.....	355	Edentates.....	9
Doubtful Kangaroo Rat.....	289	El General Rice Rat.....	189
Douglas' Chickaree.....	87	Elegant Spermophile.....	113
Douglas' Pocket Gopher.....	277	Elephant Seal.....	440
Douglas' Spermophile.....	115	Elephants, Sea.....	439
Dow's Tapir.....	58	Elk, American.....	41, 42
Drab Gray Mole Mouse.....	134	Elk, San Joaquin Valley.....	42
Drab Gray Spermophile.....	101	Emmons' Glacier Bear.....	397
Drummond's Bushy-tailed Wood Rat.....	221	Energetic Mouse.....	166
Drummond's Meadow Vole.....	237	Engelhardt's Woodchuck.....	121
Dryad Mouse.....	167	Escazu White Bat.....	494
Durango Brown Bat.....	478	Escondido River Rice Rat.....	184
Durango Chipmunk.....	88	Escondido River Squirrel.....	77
Durango Squirrel.....	67	Escuinapa Lynx.....	373
Durango Wood Rat.....	218	Eskimo Weasel.....	428
Dusky Alaskan Flying Squirrel.....	122	Esperitu Santo Island Jack Rabbit.....	358
Dusky Bat.....	504, 520, 481	Esperitu Santo Raccoon Fox.....	398
Dusky Flying Squirrel.....	123	Eten Opossum.....	8
Dusky Gray Squirrel.....	79	Eve's Mouse.....	165
Dusky Harvest Mouse.....	194, 204	Extraordinary Shrew.....	457
Dusky Kangaroo Rat.....	299	Eyra.....	368
Dusky Mastiff Bat.....	496	Eyra, Tamaulipas.....	368
Dusky Mole Mouse.....	134	Eyra, Yucatan.....	369
Dusky Musk Rat.....	255		
Dusky Pocket Mouse.....	303	Faded Vesper Rat.....	172
Dusky Porcupine.....	333	Falcate Bat.....	527
Dusky Red-backed Vole.....	232	Fannin's Mountain Sheep.....	55
Dusky Shrew.....	450, 464	<i>Farmer</i>	127
Dusky Sloth.....	9	Farmer Pocket Gopher.....	262
Dusky Spiny Mouse.....	319	Farthest North Marten.....	421
Dusky Wolf.....	376	Feeble Mouse.....	147
Dusky-footed Wood Rat.....	206	Feeble Weasel.....	427
Dutcher's Meadow Vole.....	240	Field Spotted Skunk.....	415
Dutcher's Pocket Mouse.....	302	Field Wood Rat.....	218
Dwarf Mastiff Bat.....	498	Fighting Baleen Whale.....	18
		Fighting Bear.....	396

	PAGE.		PAGE.
Finback Whales.....	18	Fox, Cascade Mountains.....	382
Fire-bellied Squirrel.....	69	Fox, Curtailed.....	385
Fir-wood Fox.....	381	Fox, Destructive.....	382
Fish, Cow.....	32	Fox, Fir-wood.....	381
Fisher, Pacific.....	423	Fox, Florida Gray.....	386
Fisher, Pennant's.....	422	Fox, Gray.....	386
Fisher's Pocket Gopher.....	277	Fox, Guatemalan Gray.....	369
Fisher's Shrew.....	454	Fox, Hall Island.....	382
Fisher's Sperophile.....	114	Fox, Inyo Mountains Gray.....	387
Fishers.....	419	Fox, Kenai.....	382
Five-toed Kangaroo Rats.....	296	Fox, Large-eared Kit.....	384
Flat Forehead Bahama Bat.....	521	Fox, Little Gray.....	389
Flat-headed Pocket Gopher.....	267	Fox, Long-tailed.....	381
Flat-headed Wood Rat.....	209	Fox, Murdering.....	381
Flat-nosed Bat.....	524	Fox, New England Gray.....	388
Fleet Hare.....	352	Fox, New Mexican Kit.....	384
Florida Black Bear.....	395	Fox, Pribilof Island Arctic.....	383
Florida Brown Bat.....	483	Fox, Red.....	380
Florida Cotton-tail.....	342	Fox, Reddish.....	381
Florida Field Mouse.....	143, 156	Fox, Royal.....	382
Florida Flying Squirrel.....	126	Fox, San Clemente Island Gray ...	390
Florida Gray Fox.....	386	Fox, San Miguel Island Gray.....	388
Florida Gray Squirrel.....	80	Fox, Santa Catalina Island Gray..	389
Florida Lynx.....	372	Fox, Santa Cruz Island Gray.....	388
Florida Mole.....	471	Fox, Scott's Gray.....	386
Florida Otter.....	434	Fox, Silver.....	380
Florida Pocket Gopher.....	262	Fox, Sluggish.....	384
Florida Red Bat.....	486	Fox, Small-toothed.....	389
Florida Rice Rat.....	181	Fox Squirrel.....	65
Florida Shrew.....	463	Fox, Summit Arctic.....	382
Florida Skunk.....	407	Fox, Swift.....	384
Florida Swamp Hare.....	341	Fox, Swift Gray.....	388
Florida Water Vole.....	252	Fox, Swift-footed Kit.....	385
Florida Weasel.....	430	Fox, Texas Gray.....	387
Florida Wood Rat.....	210	Fox, Townsend's Gray.....	387
Fluffy Mouse.....	171	Fox, Ungava Arctic.....	383
Flying Squirrels.....	121	Foxes.....	374
Foreign Mouse.....	164	Foxes, Red.....	380
Forest Bat.....	513	Franklin's Sperophile.....	105
Forest Hares.....	341	Free-tailed Mouse Bat.....	501
Forest Vole.....	236	Fremont's Chickaree.....	84
Forked-claw Lemmings.....	260	Fringed Bat.....	479
Fort Churchill Lemming.....	261	Fringed-face Bat.....	514
Fort Yuma Bat.....	475	Frisky Flying Squirrel.....	123
Four-banded Chipmunk.....	89	Frisky Mouse.....	148
Four-toed Kangaroo Rats.....	288	Fruit-loving Bat.....	527
Fox, Alaska Red.....	381	Fulvous Field Mouse.....	142
Fox, Bangs' Red.....	381	Fulvous Pocket Gopher.....	266, 280
Fox, Bering Island Arctic.....	383	Fulvous Rice Rat.....	189
Fox, Black.....	380	Fulvous-bellied Cotton Rat.....	180
Fox, California Gray.....	387	Fulvous-bellied Opossum.....	3

	PAGE.		PAGE.
Fulvous-bellied Spider Monkey . . .	535	Goldman's Rice Rat	186
Fulvous-bellied Vole	246	Goldman's Shrew	465
Fulvous-bellied Wood Rat	220	Goldman's Spermophile	116
Funnell-eared Bats	501	Goldman's Spiny Rat	318
Fur Seal	438	Goldman's Squirrel	77, 130
Fuscous Bushy-tailed Wood Rat . .	222	Goldman's Wood Rat	220
Gabb's Coon	400	Gopher, Active Pocket	273
Gabb's Hare	351	Gopher, Alpine Pocket	283
Gadow's Mouse	165	Gopher, Angular Pocket	281
Gaillard's Jack Rabbit	354	Gopher, Attwater's Pocket	263
Gale's Red-backed Vole	231	Gopher, Bailey's Pocket	285
Gambel's Mouse	145	Gopher, Black Pocket	276
Gapper's Red-backed Vole	230	Gopher, Black-faced Pocket	279
<i>Gato Montes</i>	371	Gopher, Blackish Pocket	280
Gaumer's Spiny Rat	317	Gopher, Boquete Pocket	270
Gentle Field Mouse	142	Gopher, Botta's Pocket	274
Geoffroy's Bat	520, 529	Gopher, Bridger's Pocket	279
Geoffroy's Spider Monkey	535	Gopher, Bristled Pocket	269
Geoffroy's Titi Monkey	532	Gopher, Broad-footed Pocket	271
Giant Brown Bear	391	Gopher, Broad-headed Pocket	275
Giant Lynx	373	Gopher, Broad-nosed Pocket	286
Giant Pocket Gopher	269	Gopher, Buller's Pocket	268
Giant Shrew	460	Gopher, Cabezon Pocket	287
Gibbs' Shrew Mole	466	Gopher, Cervine Pocket	287
Gila Chipmunk	85	Gopher, Cherrie's Pocket	271
Gill's Bottle-nosed Whale	30	Gopher, Chestnut-faced Pocket	266
Gill's Dolphin	32	Gopher, Coues' Pocket	284
Glacier Bay Shrew	451	Gopher, Crested Pocket	263
Gliding Spiny Rat	329	Gopher, Cumberland Pocket	262
Goat, Columbian Mountain	52	Gopher, Deceitful Pocket	264
Goat, Kennedy's Mountain	53	Gopher, Delving Pocket	287
Goat, Missoula Mountain	52	Gopher, Dessert Pocket	281
Goat, Mountain	52	Gopher, Digger Pocket	269, 274
Goats	52	Gopher, Douglas' Pocket	277
Godman's Bat	517	Gopher, Farmer Pocket	262
Godman's Shrew	455	Gopher, Fisher's Pocket	277
Golden Breasted Mouse	153	Gopher, Flat-headed Pocket	267
Golden Harvest Mouse	198	Gopher, Florida Pocket	262
Golden Pocket Gopher	285	Gopher, Fulvous Pocket	266, 280
Golden-bellied Squirrel	68	Gopher, Giant Pocket	269
Golden-eared Harvest Mouse	195	Gopher, Golden Pocket	285
Golden-neck Spermophile	107	Gopher, Goldman's Pocket	266, 283
Goldman's Bridled Weasel	432	Gopher, Grazing Pocket	282
Goldman's Coyote	378	Gopher, Great Pocket	272
Goldman's Harvest Mouse	201	Gopher, Harsh-coated Pocket	270
Goldman's Jaguar	366	Gopher, Heller's	278
Goldman's Kangaroo Rat	294	Gopher, Idaho Pocket	284
Goldman's Mouse	138	Gopher, Intermediate Pocket	281
Goldman's Pocket Gopher	266, 283	Gopher, Irazu Pocket	270
Goldman's Pocket Mouse	313	Gopher, Juarez Pocket	283
		Gopher, Large-nosed Pocket	276

	PAGE.		PAGE.
Gopher, Las Vigas Pocket	265	Gopher, Wandering Pocket	282
Gopher, Lone Pine Pocket	285	Gopher, Western Pocket	279
Gopher, Lost Pocket	283	Gopher, White-nosed Pocket	268
Gopher, Masked Pocket	264	Gopher, White-toothed Pocket	273
Gopher, Merriam's Pocket	265	Gopher, Working Pocket	273
Gopher, Mobile Pocket	262	Gopher, Yelm Prairie Pocket	277
Gopher, Mountain Pocket	265, 275	Gophers, Ancient Pocket	268
Gopher, Mount Iztaccihuatl Pocket	266	Gophers, Broad-headed Pocket	266
Gopher, Mt. Mazama Pocket	275	Gophers, Distinct Pocket	269
Gopher, Mud-loving Pocket	278	Gophers, Large Size Pocket	270
Gopher, Naked-tailed Pocket	267	Gophers, Pocket	261, 271
Gopher, Narrow-headed Pocket	270	Gophers, Powerful Pocket	264
Gopher, Near-sighted Pocket	276	Gophers, Straight-headed Pocket	268
Gopher, Neglected Pocket	268	Gophers, Zygomata Pocket	271
Gopher, Nelson's Pocket	269, 286	Goss' Lemming Vole	256
Gopher, Nevada Pocket	272	Graceful Bat	502
Gopher, Oregon Pocket	277	Graceful Chipmunk	91
Gopher, Orizaba	283	Graceful Free-tailed Bat	499
Gopher, Pacuare Pocket	271	Graceful Kangaroo Rat	294
Gopher, Pale Pocket	263, 286	Graceful Skunk	415
Gopher, Pallid Pocket	274	Graham Island Bat	477
Gopher, Peroté Pocket	265	Graham Mountain Meadow Vole	245
Gopher, Pine Barrens Pocket	261	Graham Mountains Chickaree	85
Gopher, Pine Pocket	275	Granger's Bushy-tailed Wood Rat	223
Gopher, Pine Zone Pocket	271	Granger's Cotton-tail	349
Gopher, Pygmy Pocket	284	Grant's Caribou	39
Gopher, Reddish Brown Pocket	272	Gray Bat	504
Gopher, Rhoads' Pocket	287	Gray Blackfish	28
Gopher, Richardson's Pocket	279	Gray Bushy-tailed Wood Rat	220
Gopher, Roaming Pocket	285	Gray Chipmunk	89
Gopher, Rufescent Pocket	279	Gray Desert Jack Rabbit	356
Gopher, San Pedro Martir Pocket	282	Gray Fox	386
Gopher, Sand-frequenting Pocket	264	Gray Jaguarondi	368
Gopher, Santa Anita Pocket	280	Gray Meadow Vole	239
Gopher, Shaw's Pocket	263	Gray Opossum	3
Gopher, Short-headed Pocket	263	Gray Pocket Mouse	315
Gopher, Sierra Blanca Pocket	286	Gray Rock Meadow Vole	247
Gopher, Sierra Laguna Pocket	280	Gray Seal	440
Gopher, Sinaloa Pocket	287	Gray Seals	440
Gopher, Slender-nosed Pocket	267	Gray Spiny Rat	319
Gopher, Sooty Pocket	267	Gray Striped Chipmunk	96
Gopher, Swarthy Pocket	277	Gray-backed Harvest Mouse	203
Gopher, Swift Pocket	284	Gray-faced Wood Rat	226
Gopher, Tatemales Pocket	283	Gray-footed Chipmunk	91
Gopher, Texas Pocket	264, 281	Gray-headed Chipmunk	93
Gopher, The Dalles Pocket	276	Gray-headed Grison	418
Gopher, Tough-skinned Pocket	269	Gray-headed Pika	338
Gopher, Townsend's Pocket	275	Gray's Baleen Whale	17
Gopher, Tropical Pocket	270	Gray's Black Squirrel	76
Gopher, Troublesome Pocket	282	Gray's Bottle-nosed Whale	29
Gopher, Uinta Mountains Pocket	273	Gray's Dolphin	34

	PAGE.		PAGE.
Gray's Japanese Baleen Whale....	16	Haida Weasel.....	429
Gray's Spiny Mouse.....	321	Hair Seals.....	441
Grayson's Cotton-tail.....	351	Hairy Bat.....	512
Gray-tailed Chipmunk.....	93	Hairy Hutia.....	330
Gray-tailed Meadow Vole.....	239	Hairy-lipped Bat.....	477
Grazing Pocket Gopher.....	282	Hairy-tailed Moles.....	471
Great Anteater.....	12	Haitian Huita.....	332
Great Basin Pocket Mouse.....	307	Hall Island Fox.....	382
Great Basin Skunk.....	409	Hall Island Meadow Vole.....	243
Great Bat.....	501	Hamilton Smith's White-tailed Deer.....	46
Great California Pocket Mouse...	314	Happy Chipmunk.....	92
Great Kangaroo Rat.....	298	Harbor Seal.....	442
Great Pocket Gopher.....	272	Hare, American.....	360
Great Pocket Mouse.....	310	Hare, Attwater's Swamp.....	340
Great Shrew.....	466	Hare, Baird's.....	361
Great Swamp Hare.....	340	Hare, Bangs' Polar.....	358
Greater Desert Cotton-tail.....	350	Hare, Bishop's.....	362
Great-footed Hare.....	361	Hare, British Columbian.....	361
Great-tailed Mouse.....	171	Hare, Cinnamon Chief.....	338
Grebnitzky's Two-toothed Sperm Whale.....	23	Hare, Dall's.....	361
Greedy Woodchuck.....	120	Hare, Eastern Swamp.....	341
Green Monkey.....	536	Hare, Fleet.....	352
Greenland or Arctic Caribou.....	39	Hare, Florida Swamp.....	341
Greenland Hare.....	359	Hare, Gabb's.....	351
Grison, Allamand's.....	419	Hare, Great-footed.....	361
Grison, Calovevora.....	418	Hare, Great Swamp.....	340
Grison, Gray-headed.....	418	Hare, Greenland.....	359
Grison, Grizzled.....	419	Hare, Hoary Polar.....	358
Grison, Nelson's.....	419	Hare, Klamath.....	362
Grisons.....	418	Hare, Labrador.....	359
Grizzled Brown Bat.....	475	Hare, Leaping.....	362
Grizzled Grison.....	419	Hare, MacFarlane's.....	362
Grizzled Mouse.....	154	Hare, Nelson's.....	340
Grizzled Spider Monkey.....	536	Hare, Plains.....	362
Grizzly Bear.....	392	Hare, Polar.....	358
Growling Spotted Skunk.....	413	Hare, Richardson's.....	363
Guadalupe Fur Seal.....	438	Hare, St. Michaels'.....	360
Guadalupe Mountains Meadow Vo'e	245	Hare, Siberian.....	359
Guadalupe Skunk.....	412	Hare, Sierra.....	363
Guanacevi Chipmunk.....	86	Hare, Small-footed.....	356
Guatemala Squirrel.....	74	Hare, Swift.....	360
Guatemalan Gray Fox.....	389	Hare, Townsend's.....	363
Guatemalan Meadow Vole.....	252	Hare, True's Swamp.....	341
Guatemalan Rice Rat.....	185	Hare, Virginian.....	360
Guatemalan Shrew.....	455	Hare, Wandering Swamp.....	341
Guerrero Squirrel.....	72	Hare, Washington.....	361
Gull Island Meadow Vole.....	238	Hare, Western.....	360
Gundlach's Bat.....	488	Hares.....	339
Gundlach's Hutia.....	331	Hares, Changeable.....	358
Gunnison's Prairie-dog.....	118	Hares, Chief.....	337

	PAGE.		PAGE.
Hares, Forest.....	341	Hudson Bay Musk Rat	253
Hares, Short-tailed.....	352	Hudson Bay Sable	419
Hares, Swamp.....	340	Hudsonian Lemming	260
Hares, Tailless.....	351	Huehuetan Spiny Mouse	320
Harp Seal.....	444	Hutia, Black-tailed.....	331
Harris' <i>Spermophile</i>	97	Hutia, Gundlach's.....	331
Harsh-coated Pocket Gopher	270	Hutia, Hairy	330
Harvest Mice.....	191	Hutia, Haitian.....	332
Hayden's Meadow Vole.....	248	Hutia, Ingraham's.....	332
Heavy Peccary	36	Hutia, Lanceolate-spot.....	331
Heedless Bat.....	473	Hutia, Prehensile-tailed.....	331
Heller's Gopher.....	278	Hutia, Short-tailed.....	331
Heller's Kangaroo Rat.....	292	Hutia, White-banded.....	331
Heller's Pocket Mouse.....	312	Hyacinthine Shrew Mole.....	467
Heller's White-striped Bat.....	526		
Hernandez's Jaguar.....	366	Ice Baleen Whale	15
Hernandez's Raccoon.....	401	Ichneumon	173
Herron's Field Mouse.....	155	Idaho Cotton-tail.....	351
Hidalgo Jack Rabbit	354	Idaho Pocket Gopher.....	284
Hind's Chipmunk.....	86	Idaho Red-backed Vole.....	232
Hispid Cotton Rat	175	Illinois Skunk.....	407
Hispid Harvest Mouse	202	Imitator Mouse.....	167
Hispid Pocket Mouse.....	310	Indianola Spotted Skunk.....	413
Hispid Spiny Rat.....	318	Inflated-nosed Bat	514
Hoary Bat.....	487	Ingraham's Hutia	332
Hoary Field Mouse.....	150	Inland White Bear.....	397
Hoary Mole Mouse.....	133	Innuity Lemming Vole.....	256
Hoary Polar Hare.....	359	Innuity Meadow Vole.....	243
Hoary <i>Spermophile</i>	112	Intermediate Bat	487, 523
Hoary Spiny Mouse	321	Intermediate Mole.....	470
Hoary Woodchuck.....	120	Intermediate Pocket Gopher.....	281
Hoary-rumped Cotton-tail.....	344	Intermediate Pocket Mouse.....	313
Hoffmann's Sloth	9	Intermediate Vole.....	226
Hoffmann's Squirrel.....	61	Inyo Mountains Chipmunk.....	89
Holzner's Cotton-tail	345	Inyo Mountains Gray Fox.....	387
Honduras Spiny Mouse	320	Irazu Pocket Gopher	270
Honduras Squirrel.....	76	Irazu Volcano Harvest Mouse.....	194
Hooded Seal.....	440	Irrational Shrew	462
Hooded Seals.....	440	Island of Coiba Howler	533
Hooded Skunk.....	410	Island of Cozumel Mouse.....	162
Hoofed Quadrupeds.....	35	Island of Cozumel Opossum	6
House Mouse.....	130	Island of Tiburon Mouse.....	161
Howler, Island of Coiba.....	533	Island Opossum.....	7
Howler, Mantled.....	533	Island Raccoon	401
Howler, Mexican.....	533	Island <i>Spermophile</i>	115
Howler, Villous.....	533	Isolated Rice Rat	191
Howling Monkeys.....	532	Isthmian Agouti	336
Hoy's Shrew	457	Isthmian Spiny Mouse.....	322
Huachuca Squirrel.....	64	Isthmian Wood Rat.....	219
Hudson Bay Chickaree.....	80		
Hudson Bay Jumping Mouse.....	324	<i>Jabali</i>	35

	PAGE.		PAGE.
Jack Rabbit, Donkey.....	355	Kinkajou, Chiriqui.....	404
Jack Rabbits.....	353	Kinkajous.....	397, 404
Jaguar, Costa Rica.....	365	Klamath Flying Squirrel.....	123
Jaguar, Goldman's.....	366	Klamath Hare.....	362
Jaguar, Hernandez's.....	366	Klamath Harvest Mouse.....	196
Jaguarondi.....	367		
Jaguarondi, Boqueron.....	368	La Carpintera Field Mouse.....	165
Jaguarondi, Gray.....	368	La Cienaga Squirrel.....	67
Jaguarondi, Sinaloa.....	368	La Grulla Brown Bat.....	476
Jaguars.....	365	La Palma Field Mouse.....	158
Jalapa Brown Mouse.....	136	La Parada Spiny Rat.....	317
Jalapa Field Mouse.....	169	La Salada Deer Mouse.....	163
Jalapa Harvest Mouse.....	201	La Salada Squirrel.....	71
Jalapa House Mouse.....	130	Labrador Hare.....	359
Jalapa Meadow Vole.....	250	Labrador Jumping Mouse.....	324
Jalapa Rice Rat.....	184	Labrador Lemming Vole.....	256
Jalapan Mouse.....	154	Labrador Marten.....	420
Jalisco Brown Bat.....	477	Labrador Porcupine.....	334
Jalisco Small-eared Mouse.....	156	Labrador Shrew.....	455
Jamaica Bat.....	522	Labrador Woodchuck.....	120
Jamaica Large-eared Bat.....	508	Lacey's Harvest Mouse.....	199
Jamaica Rice Rat.....	188	Lanceolate-spot Hutia.....	331
Jaral Pocket Mouse.....	313	Languishing Spermophile.....	101
Jet Mouse.....	161	Large Cotton Rat.....	177
Juarez Pocket Gopher.....	283	Large Labrador Meadow Vole.....	237
Jumping Mice.....	323	Large Mink.....	424
Juquila Mouse.....	157	Large Shrew Mole.....	467
Juquila Wood Rat.....	220	Large Size Pocket Gophers.....	270
		Large Spotted Spermophile.....	100
Kadiak Island Spermophile.....	108	Large-eared Field Mouse.....	170
Kadiak Island Weasel.....	428	Large-eared Flying Squirrel.....	124
Kadiak Meadow Vole.....	242	Large-eared Harvest Mouse.....	203
Kaibab Squirrel.....	67	Large-eared Kit Fox.....	384
Kamloops Jumping Mouse.....	325	Large-eared Mole Mouse.....	134
Kangaroo Rats.....	288	Large-eared Vampire Bat.....	513
Kansas Pocket Mouse.....	310	Large-eared Wood Rat.....	206
Keeler Kangaroo Rat.....	288	Large-headed Kangaroo Rat.....	299
Keene's Brown Bat.....	478	Large-nosed Mastiff Bat.....	497
Keen's Mouse.....	152	Large-nosed Mole.....	470
Keewatin Mink.....	424	Large-nosed Mouse.....	159
Keewatin Shrew.....	457	Large-nosed Pocket Gopher.....	276
Kenai Fox.....	382	Large-striped Skunks.....	406
Kenai Mountain Sheep.....	55	Large-toothed Killer Whale.....	27
Kenai Peninsula Brown Bear.....	392	Large-toothed Musk Rat.....	255
Kenai Peninsula Marten.....	421	Large-toothed Rice Rat.....	188
Kennedy's Mountain Goat.....	53	Large-toothed Shrew.....	455
Kennicott's Spermophile.....	99, 112	Large-winged Bat.....	473
Kidder's Brown Bear.....	391	Las Peñas Bat.....	506
Killer Whales.....	26	Las Vigas Bat.....	482
Kindred Mouse.....	166	Las Vigas Harvest Mouse.....	196
Kinkajou, Aztec.....	404	Las Vigas Pocket Gopher.....	265

	PAGE.		PAGE.
Leafless or Blunt-nosed Bat	521	Long-footed Mole Mouse	131
Leaping Hare	362	Long-haired Mouse	159
Least Chipmunk	93	Long-haired Rice Rat	183
Least Cotton Rat	178	Long-nosed Bat	492
Least Pika	339	Long-nosed Dolphin	33
Least Upland Meadow Vole	248	Long-nosed Mouse	170, 152
Le Conte's Harvest Mouse	191	Long-nosed Shrew	454
Lemming, Black-footed	260	Long-nosed Spiny Mouse	322
Lemming, Fort Churchill	261	Long-spined Porcupines	332
Lemming, Hudsonian	260	Long-tailed Fox	380
Lemming, Nelson's	260	Long-tailed Harvest Mouse	194
Lemming, Richardson's	259	Long-tailed Meadow Vole	244
Lemming, Small	259	Long-tailed Mole Mouse	132
Lemming, Unalaska	260	Long-tailed Mouse	149
Lemming Voles	255	Long-tailed Pocket Mouse	308
Lemming, Yellowish	259	Long-tailed Rice Rat	186
Lemming, Yukon	259	Long-tailed Shrew	448, 450
Lemmings	258	Long-tailed Spemophile	113
<i>Leon Miquero</i>	367	Long-tailed Spiny Rat	318
<i>Leon Monero</i>	367	Long-tailed Texan Skunk	407
Lesser Desert Cotton-tail	350	Long-tailed Vole	228
Lesson's Finback Whale	20	Long-tailed Weasel	430
Lewis' Prairie-dog	118	Loring Pocket Mouse	306
Lichtenstein's White-tailed Deer	45	Loring's Red-backed Vole	231
Light-colored Porcupine	334	Lost Pocket Gopher	283
Light-footed Kangaroo Rat	298	Louisiana Black Bear	395
Linnean Opossum	6	Louisiana Field Mouse	143
Lion Hill Squirrel	62	Louisiana Meadow Vole	248
Little Brown Bat	479	Louisiana Skunk	407
Little California Bat	476	Louisiana White-tailed Deer	44
Little Comodu Bat	492	Lower California Badger	405
Little Cotton-tail	348	Lower California Bat	474
Little Desert Pocket Mouse	312	Lower California Brown Bat	484
Little Free-tailed Bat	501	Lower California Chipmunk	88
Little Gray Fox	389	Lower California Cotton-tail	353
Little Gray Squirrel	61	Lower California Coyote	378
Little Labrador Meadow Vole	236	Lower California Jack Rabbit	358
Little Large-eared Bat	511	Lower California Lynx	373
Little Mexican Bat	477	Lower California Mole	469
Little or Two-toed Anteater	10	Lower California Mountain Sheep	54
Little Raccoon	401	Lower California Mule Deer	50
Little Spemophile	105	Lower California or Pallid Harvest	
Little Spiny Mouse	322	Mouse	195
Little Striped Skunk	412	Lower California Rice Rat	182
Littoral Spiny Mouse	320	Lower California Skunk	409
Lively Cotton-tail	343	Lower California Spemophile	98
<i>Lobo</i>	376	Lower California Wood Rat	214
Lofty Mountain Shrew	451	Lynx	370
Lone Pine Pocket Gopher	285	Lynx, Bailey's	372
Lonely Rice Rat	188	Lynx, Barred	371
Long-footed Kangaroo Rat	298	Lynx, California	372

	PAGE.		PAGE.
Lynx, Canada	370	Masked Pocket Gopher.....	264
Lynx, Desert.....	372	Masked Shrew.....	446
Lynx, Eastern.....	370	Maximilian's Pocket Mouse.....	301
Lynx, Escuinapa.....	373	Maya Shrew.....	465
Lynx, Florida.....	372	Maynard's Raccoon.....	402
Lynx, Giant.....	373	Mazatlan Shrew.....	460
Lynx, Lower California	373	Meadow Voles	233
Lynx, Pallid.....	371	Mearns' Chickaree.....	83
Lynx, Red	317	Mearns' Cotton-tail.....	342
Lynx, Sharp-sighted.....	371	Mearns' Coyote	377
Lynx, Soft-furred	370	Mearns' Field Mouse.....	150
Lynx, Spotted.....	371	Mearns' Skunk.....	412
Lynx, Uinta Mountains.....	373	Merriam's Chipmunk	88
Lyster's Striped Chipmunk.....	95	Merriam's Harvest Mouse	192
MacFarlane's Hare.	362	Merriam's Jack Rabbit.....	354
MacFarlane's Meadow Vole.....	241	Merriam's Kangaroo Rat.....	288
Mackenzie's Vole.....	228	Merriam's Pocket Gopher	265
Macleay's Bat	504	Merriam's Pocket Mouse	302
Makkovik Flying Squirrel.....	125	Merriam's Shrew.....	454
Manatee	14	Merriam's Spotted Skunk	413
Manatees.....	14	Merriam's Wapiti	42
<i>Mamgordo</i>	366	Mexican Agouti.....	336
Mantled Howler.....	533	Mexican Bat.....	506
Manzanillo Coati.....	403	Mexican Chipmunk.....	88
Manzanillo Squirrel	73	Mexican Flying Squirrel.....	126
Margarita Pocket Mouse	316	Mexican Free-tailed Bat.....	500
Margay	367	Mexican Grizzly Bear.....	393
Maria Madre Island Bat.....	516	Mexican Harvest Mouse.....	197
Maria Madre Island Mouse	165	Mexican Howler.....	533
Maria Madre Island Opossum.....	3	Mexican Meadow Vole.....	245
Maria Madre Island Rice Rat.....	182	Mexican Mountain Sheep.....	54
Marmosets	531	Mexican Mouse	168
Marmots.....	59	Mexican Murine Opossum.....	2
Marsh Rice Rat.....	187	Mexican Otter.....	434
Marsh Shrew	458	Mexican Pocket Mouse.....	303
Marten, Active	420	Mexican Prairie-dog.....	119
Marten, American	419	Mexican Prong-horn.....	51
Marten, Farthest North.....	421	Mexican Red Bat	486
Marten, Kenai Peninsula.....	421	Mexican Spermophile.....	103
Marten, Labrador.....	420	Mexican Spider Monkey.....	535
Marten, Newfoundland.....	422	Mexican Timber Wolf.....	376
Marten, Pacific.....	422	Mexican Vampire Bat.....	513
Marten, Queen Charlotte Island.....	420	Mexican White-tailed Deer.....	45
Marten, Rocky Mountain.....	422	Mexican Wood Rat.....	213
Marten, Saskatchewan.....	419	Mice.....	129
Marten, Selkirk.....	420	Mice, Harvest.....	191
Martens.....	419	Mice, Jumping.....	323
<i>Martica</i>	404	Mice, Pocket.....	288, 300
Mascota Cotton Rat.....	176	Michigan Field Mouse.....	160
Masked Bat	505	Michoacan Bridled Weasel.....	432
		Michoacan Cotton-tail	344

	PAGE.		PAGE.
Michoacan Squirrel.....	70	Monkeys, Spider.....	534
<i>Mico-dormilon</i>	534	Monkeys, Squirrel.....	533
<i>Micoleon</i>	404	Mole, Alpine.....	469
Middendorff's Brown Bear.....	391	Mole, Brewer's Hairy-tailed.....	471
Miller's Armadillo.....	13	Mole, California.....	468
Miller's Bat.....	474	Mole, Coast.....	468
Miller's Florida Bat.....	488	Mole, Common.....	470
Miller's Nose-leaf Bat.....	519	Mole, Copper-colored.....	471
Miller's Skunk.....	410	Mole, Florida.....	471
Mimic Mouse.....	157	Mole, Gibbs' Shrew.....	466
Mimic Squirrel.....	66	Mole, Hyacinthine Shrew.....	467
Mine-haunting Mouse.....	138	Mole, Intermediate.....	470
Mink.....	423	Mole, Large-nosed.....	470
Mink, Black-footed.....	425	Mole, Large Shrew.....	467
Mink, Common.....	423	Mole, Lower California.....	469
Mink, Dark-hued.....	425	Mole, Smaller California.....	469
Mink, Devilish.....	424	Mole, Star-nosed.....	467
Mink, Keewatin.....	424	Mole, Texas.....	471
Mink, Large.....	424	Mole, Townsend's.....	468
Mink, Mud-loving.....	425	Mole, True's.....	469
Mink, Rambling.....	424	Moles.....	466
Minnesota Chickaree.....	81	Moles, Common.....	469
Minnesota Gray Squirrel.....	79	Moles, Hairy-tailed.....	471
Minnesota Mule Deer.....	49	Moles, Shrew-like.....	466
Minnesota White-footed Deer		Moles, Star-nosed.....	467
Mouse.....	141	Moles, Western.....	468
Minute Shrew.....	449	<i>Mono</i>	533
<i>Minero, Leon</i>	367	<i>Mono Carablanca</i>	536
Mirador Brown Bat.....	484	<i>Mono Colorada</i>	535
Missoula Mountain Goat.....	52	<i>Monte, Cacomistl de</i>	399
Mixed Cotton-tail.....	347	<i>Monte, Cacomistle de</i>	399
Mobile Baleen Whale.....	18	Monterey Shrew.....	452
Mobile Pocket Gopher.....	262	<i>Montes Gato</i>	371
Mogollon Mountains Meadow Vole.....	246	Montezuma Wood Rat.....	218
Mohave Desert Spermophile.....	103	Montserrat Bat.....	526
Mohave Desert Wood Rat.....	207	Moose.....	40
Mole-like Meadow Vole.....	249	Moose, Alaskan.....	41
Moles.....	466	<i>Moran</i>	35
<i>Monero, Leon</i>	367	Motzorongo Skunk.....	412
Monkey, Black Spider.....	535	Mount Iztaccihuatl Pocket Gopher.....	266
Monkey, Fulvous-bellied Spider... ..	535	Mount Lyell Shrew.....	453
Monkey, Geoffroy's Spider.....	535	Mount Magruder Pocket Mouse... ..	307
Monkey, Geoffroy's Titi.....	532	Mount Malinche Mouse.....	164
Monkey, Green.....	536	Mt. Mazama Pocket Gopher.....	275
Monkey, Grizzled Spider.....	536	Mt. Mazama Red-backed Vole... ..	232
Monkey, Mexican Spider.....	535	Mount McKinley Wolverine.....	417
Monkey, Noisy Squirrel.....	534	Mount Pinos Mouse.....	157
Monkey, CErsted's Titi.....	534	Mount Pinos Pocket Mouse.....	304
Monkey, Rufous-foot Squirrel.....	534	Mount Pinos Wood Rat.....	208
Monkeys, Howling.....	532	Mount Popocatepetl Harvest Mouse.....	199
Monkeys, Prehensile-tailed.....	532	Mount Popocatepetl Shrew.....	466

	PAGE.
Mount Rainier Sewellel	127
Mount Shasta Shrew	453
Mt. Tancitaro Wood Rat	224
Mount Zempoaltepec Field Mouse	167
Mount Zempoaltepec Meadow Vole	251
Mount Zempoaltepec Shrew	466
Mountain Bushy-tailed Wood Rat	221
Mountain Caribou	38
Mountain Chipmunk	95
Mountain Goat	52
Mountain Harvest Mouse	193
Mountain Harvest Mouse	197
Mountain Jumping Mouse	327
Mountain Meadow Vole	245
Mountain Mouse	152
Mountain Pocket Gopher	205, 275
Mountain Sheep	53
Mountain Shrew	449, 456
Mountain Squirrel	72
Mountain Vole	227
Mountain Weasel	430
Mountain Wood Rat	225
Mountain-loving Mouse	139
Mouse, Akeley's Field	148
Mouse, Alaska Jumping	325
Mouse, Alien	169
Mouse, Allen's Jumping	327
Mouse, Allen's Pocket	315
Mouse, Allen's Spiny	321
Mouse, Allied Desert	137
Mouse, Allied Field	154, 161
Mouse, Allied House	161
Mouse, Alpine Harvest	200
Mouse, Alston's	204
Mouse, Ameca Harvest	202
Mouse, Anastasia Island	138, 144
Mouse, Anthony's	137
Mouse, Anthony's Pocket	314
Mouse, Apache Pocket	303
Mouse, Apazote	163
Mouse, Arizona Harvest	196
Mouse, Arizona White-footed Deer	141
Mouse, Ashy Gray Field	164
Mouse, Attwater's Field	154
Mouse, Aztec	162
Mouse, Bailey's Pocket	309
Mouse, Baird's Pocket	302, 307
Mouse, Banderas Field	163
Mouse, Bangs' Field	144
Mouse, Bangs' Pocket	305
Mouse, Bangs' Red	205

	PAGE.
Mouse, Batopilas Pocket	314
Mouse, Beautiful	154
Mouse, Beautiful Mole	134
Mouse, Beautiful Pocket	303
Mouse, Big-eared	157
Mouse, Big-eared Harvest	193
Mouse, Black Pocket	301
Mouse, Black-browed	166
Mouse, Black-browed Mole	131
Mouse, Black-eared Field	161
Mouse, Black-eared Pocket	303
Mouse, Black-footed	170
Mouse, Blackish Field	156
Mouse, Black-tailed	137
Mouse, Blue Mountains Jumping	326
Mouse, Boquete	204
Mouse, Boquete Field	172
Mouse, Boyle's Field	150
Mouse, Broad-nosed Pocket	313
Mouse, Brownsville Harvest	198
Mouse, Bryant's Pocket	316
Mouse, Buff	147
Mouse, Buff-bellied Pocket	301
Mouse, Buff-colored Pocket	309
Mouse, Cael Harvest	201
Mouse, California	138
Mouse, California Pocket	315
Mouse, Canadian Field	144
Mouse, Cape St. Lucas Pocket	316
Mouse, Captious Harvest	197
Mouse, Carolinian Jumping	325
Mouse, Catalina Island	149
Mouse, Catemaco Spiny	320
Mouse, Cecil's	161
Mouse, Cerros Island	155
Mouse, Chapman's	139
Mouse, Cherrie's Costa Rica	160
Mouse, Chichen Itza Harvest	198
Mouse, Chilpancingo	168
Mouse, Chiriqui Harvest	194
Mouse, Chubby	168
Mouse, Cinereous Harvest	197
Mouse, Coast Jumping	327
Mouse, Cofre de Perote Harvest	199
Mouse, Colima City	136
Mouse, Colima Harvest	200
Mouse, Columbian Pocket	308
Mouse, Coolidge's Field	147
Mouse, Costa Rica Harvest	200
Mouse, Cotton Tree	143
Mouse, Coues' Pocket	307

	PAGE.		PAGE.
Mouse, Crested-tailed.....	167	Mouse, Hoary Spiny.....	321
Mouse, Cumberland Island Field..	144	Mouse, Honduras Spiny.....	320
Mouse, Curly Spiny.....	320	Mouse, House.....	130
Mouse, Dark Field.....	155	Mouse, Hudson Bay Jumping....	324
Mouse, Dark-tailed Spiny.....	319	Mouse, Huehuetan Spiny.....	320
Mouse, Desert.....	137	Mouse, Imitator.....	167
Mouse, Desert Pocket.....	311	Mouse, Intermediate Pocket.....	313
Mouse, Desert White-footed.....	140	Mouse, Irazú Volcano Harvest....	194
Mouse, Dickinson's Harvest.....	192	Mouse, Island of Cozumel.....	162
Mouse, Diligent Harvest.....	191	Mouse, Island of Tiburon.....	161
Mouse, Drab-gray Mole.....	134	Mouse, Isthmian Spiny.....	322
Mouse, Dryad.....	167	Mouse, Jalapa Brown.....	136
Mouse, Dusky Harvest.....	194, 204	Mouse, Jalapa Field.....	169
Mouse, Dusky Mole.....	134	Mouse, Jalapa Harvest.....	201
Mouse, Dusky Pocket.....	303	Mouse, Jalapa House.....	130
Mouse, Dusky Spiny.....	319	Mouse, Jalapan.....	154
Mouse, Dutcher's Pocket.....	302	Mouse, Jalisco Small-eared.....	156
Mouse, Dwarf.....	135	Mouse, Jaral Pocket.....	313
Mouse, Dyche's Harvest.....	192	Mouse, Jet.....	161
Mouse, Eastern Desert Pocket....	312	Mouse, Juquila.....	157
Mouse, Energetic.....	166	Mouse, Kamloops Jumping.....	325
Mouse, Eve's.....	165	Mouse, Kansas Pocket.....	310
Mouse, Feeble.....	147	Mouse, Keen's.....	152
Mouse, Florida Field.....	143, 156	Mouse, Kindred.....	166
Mouse, Fluffy.....	171	Mouse, Klamath Harvest.....	196
Mouse, Foreign.....	164	Mouse, La Carpintera Field.....	165
Mouse, Frisky.....	118	Mouse, La Palma Field.....	158
Mouse, Fulvous Field.....	142	Mouse, La Salada Deer.....	163
Mouse, Gadow's.....	165	Mouse, Labrador Jumping ⁴	324
Mouse, Gambel's.....	145	Mouse, Lacey's Harvest.....	199
Mouse, Gentle Field.....	142	Mouse, Large-eared Field.....	170
Mouse, Golden Breasted.....	153	Mouse, Large-eared Harvest.....	203
Mouse, Golden Harvest.....	198	Mouse, Large-eared Mole.....	134
Mouse, Golden-eared Harvest.....	195	Mouse, Large-nosed.....	159
Mouse, Goldman's.....	138	Mouse, Las Vigas Harvest.....	196
Mouse, Goldman's Harvest.....	201	Mouse, Le Conte's Harvest.....	191
Mouse, Goldman's Pocket.....	313	Mouse, Little Desert Pocket.....	312
Mouse, Gray Pocket.....	315	Mouse, Little Spiny.....	322
Mouse, Gray-backed Harvest.....	203	Mouse, Littoral Spiny.....	320
Mouse, Gray's Spiny.....	321	Mouse, Long-footed Mole.....	131
Mouse, Great Basin Pocket.....	307	Mouse, Long-haired.....	159
Mouse, Great California Pocket ...	314	Mouse; Long-nosed.....	152, 170
Mouse, Great Pocket.....	310	Mouse, Long-nosed Spiny.....	322
Mouse, Great Tailed.....	171	Mouse, Long-tailed.....	149
Mouse, Grizzled.....	154	Mouse, Long-tailed Harvest.....	194
Mouse, Heller's Pocket.....	312	Mouse, Long-tailed Mole.....	132
Mouse, Herron's Field.....	155	Mouse, Long-tailed Pocket.....	308
Mouse, Hispid Harvest.....	202	Mouse, Loring Pocket.....	306
Mouse, Hispid Pocket.....	310	Mouse, Louisiana Field.....	143
Mouse, Hoary Field.....	150	Mouse, Lower California or Pallid	
Mouse, Hoary Mole.....	133	Harvest.....	195

	PAGE.		PAGE.
Mouse, Margarita Pocket.....	316	Mouse, Pacific Coast White-footed	
Mouse, Maria Madre Island.....	165	Deer.....	141
Mouse, Maximilian's Pocket.....	301	Mouse, Pacific Jumping.....	327
Mouse, Mearns' Field.....	150	Mouse, Pacific Pocket.....	305
Mouse, Merriam's Harvest.....	192	Mouse, Painted Spiny.....	322
Mouse, Merriam's Pocket.....	302	Mouse, Pale Michigan.....	160
Mouse, Mexican.....	168	Mouse, Pale Mole.....	131
Mouse, Mexican Harvest.....	197	Mouse, Pallid Pocket.....	314
Mouse, Mexican Pocket.....	303	Mouse, Panamint Pocket.....	304
Mouse, Mimic.....	157	Mouse, Parasite.....	151
Mouse, Mine-haunting.....	138	Mouse, Patzcuaro Harvest.....	203
Mouse, Minnesota White-footed		Mouse, Penciled Field.....	150
Deer.....	141	Mouse, Peninsular Harvest.....	197
Mouse, Mount Magruder Pocket....	307	Mouse, Perote.....	158
Mouse, Mount Malinche.....	164	Mouse, Plains Pocket.....	301
Mouse, Mount Pinos.....	157	Mouse, Plantinar Spiny.....	322
Mouse, Mount Pinos Pocket.....	304	Mouse, Prairie Jumping.....	325
Mouse, Mount Popocatepetl Har-		Mouse, Prevost Island.....	152
vest.....	199	Mouse, Price's Pocket.....	311
Mouse, Mount Zempoaltepec Field.	167	Mouse, Puerto Angel.....	163
Mouse, Mountain.....	152	Mouse, Ramona Mole.....	133
Mouse, Mountain Harvest.....	193, 197	Mouse, Reddish.....	153
Mouse, Mountain Jumping.....	327	Mouse, Related White-footed....	155
Mouse, Mountain-loving.....	139	Mouse, Rhoads'.....	139
Mouse, Mule-eared.....	139	Mouse, Rhoads' Mole.....	132
Mouse, Naked-eared.....	163	Mouse, Rich-colored.....	146
Mouse, Neat Field.....	159	Mouse, Rio Grande White-footed..	142
Mouse, Nebraska Field.....	146	Mouse, Rio Sestin Meadow.....	194
Mouse, Nebraska Harvest.....	192	Mouse, Roan Mountain.....	145
Mouse, Nelson's Field.....	172	Mouse, Roan Mountain Jumping ..	328
Mouse, Nelson's Pocket.....	313	Mouse, Robust Field.....	156
Mouse, Nelson's Spiny.....	323	Mouse, Rock.....	169
Mouse, Nevada Jumping.....	326	Mouse, Rocky Mountain Jumping.	325
Mouse, Nevada Pocket.....	315	Mouse, Rowley's Field.....	151
Mouse, Northern White-footed		Mouse, Rowley's Pine.....	151
Deer.....	140	Mouse, Rufous Harvest.....	201
Mouse, Northern Woodland Jump-		Mouse, Ruidoso Mole.....	131
ing.....	328	Mouse, Sachem or Chief.....	156
Mouse, Northwest Jumping.....	326	Mouse, Samalayuca Mole.....	131
Mouse, Northwest Pocket.....	308	Mouse, San Clemente Island.....	145
Mouse, Nuttall's Golden.....	160	Mouse, San Cristobal Field.....	170
Mouse, Oaxaca Field.....	168	Mouse, San Felipe Field.....	167
Mouse, Oaxaca Harvest.....	202	Mouse, San Geronimo Island Field.	162
Mouse, Odd-tailed.....	166	Mouse, San Joaquin Pocket.....	306
Mouse, Oposura Harvest.....	198	Mouse, San Martin Island.....	148
Mouse, Oregon Pocket.....	306	Mouse, San Pedro Martir Mountains	159
Mouse, Orizaba Field.....	169	Mouse, San Quentin.....	148
Mouse, Orizaba Harvest.....	200	Mouse, San Quentin Pocket.....	309
Mouse, Osgood's Field.....	147	Mouse, San Sebastian Harvest....	203
Mouse, Osgood's Pocket.....	304	Mouse, Sand-dwelling Pocket.....	305
Mouse, Ozolotepec Field.....	170	Mouse, Sand-loving.....	137

	PAGE.		PAGE.
Mouse, Sand-loving Mole.....	133	Mouse, Uruapan Spiny.....	322
Mouse, Santa Catalina Island Har-		Mouse, Valparaiso Harvest.....	194
vest.....	195	Mouse, Vera Cruz Spiny.....	319
Mouse, Saskatchewan Jumping....	326	Mouse, Volcan Toluca Harvest....	199
Mouse, Shaded Field.....	144	Mouse, Volcano of Irazu.....	204
Mouse, Short-eared California		Mouse, Wagner's Field.....	149
Pocket.....	314	Mouse, Warner Mountain Jump-	
Mouse, Short-nosed Pocket.....	305	ing.....	326
Mouse, Short-tailed Mole.....	131	Mouse, White-bellied.....	138
Mouse, Shy Field.....	164	Mouse, White-bellied Mole.....	130
Mouse, Sinaloa Pocket.....	312	Mouse, White-eared Pocket.....	308
Mouse, Sitka.....	152	Mouse, White-footed Deer.....	140
Mouse, Slender Harvest.....	199	Mouse, White-lipped Harvest....	202
Mouse, Slender-nosed Harvest....	201	Mouse, White-spot Field.....	142
Mouse, Slender-nosed Pocket.....	311	Mouse, White-tailed.....	165
Mouse, Smallest Spiny.....	321	Mouse, Whitish Harvest.....	193
Mouse, Sonora Spiny.....	323	Mouse, Woodland Jumping.....	328
Mouse, Sonora White-footed Field.	142	Mouse, Woolly.....	158
Mouse, Sonoyta Desert.....	165	Mouse, Xometla.....	162
Mouse, Sooty.....	136, 171	Mouse, Yaki Mole.....	133
Mouse, Southern Pocket.....	308	Mouse, Yavapai Pocket.....	302
Mouse, Spiny Pocket.....	316	Mouse, Yellow Pocket.....	315
Mouse, Squirrel Inn.....	155	Mouse, Yohaltun Field.....	147
Mouse, Stephens'.....	153	Mouse, Yucatan.....	163
Mouse, Stephens' Pocket.....	312	Mouse, Zacatecas Pocket.....	311
Mouse, Stickeen Jumping.....	327	Mouse, Zamora.....	166
Mouse, Subarctic Field.....	146	Mousing Weasel.....	428
Mouse, Sunny.....	154	Mud-loving Mink.....	425
Mouse, Swarthy.....	139	Mud-loving Pocket Gopher.....	278
Mouse, Tawny Field.....	143	Mule Deer.....	49
Mouse, Taylor's.....	135	Mule-eared Mouse.....	139
Mouse, Teapa.....	169	Mongoose.....	373
Mouse, Tehuantepec Field.....	168	Mongoose, Common Indian.....	374
Mouse, Tennessee Cotton.....	143	Murdering Fox.....	381
Mouse, Texan Spiny.....	323	Murderous Weasel.....	429
Mouse, Texas Field.....	145	Murine Opossum.....	2
Mouse, Thomas' Guerrero Field....	171	Musk Ox.....	56
Mouse, Thurber's Field.....	146	Musk Oxen.....	55
Mouse, Tiny.....	136, 176	Musk Rat.....	254
Mouse, Tlalpam Field.....	164	Musk Rats.....	252
Mouse, Tlalpam Harvest.....	203	Musk-bearing Rats.....	189
Mouse, Todos Santos.....	171	Musky Bat.....	495
Mouse, Todos Santos Harvest....	202	<i>Muyus</i>	399
Mouse, Todos Santos Island.....	162		
Mouse, Torrid Spiny.....	321	Nagai Island Sperophile.....	109
Mouse, Totontepec Field.....	169	Naked-eared Mouse.....	163
Mouse, Tropical Mole.....	133	Naked-tailed Pocked Gopher.....	267
Mouse, Troublesome.....	158	Naked-tailed Rat.....	173
Mouse, Troublesome Field.....	149	Naked-tailed Wood Rat.....	214
Mouse, True's.....	159	Narrow-headed Pocket Gopher....	270
Mouse, Tulare Mole.....	132	Narrow-headed Spotted Skunk....	416

	PAGE.		PAGE.
Narrow-headed Wood Rat.....	217	Northern Musk Rat.....	253
Narwhal.....	24	Northern Plains Skunk.....	406
Navigator Shrew.....	458	Northern Sea Otter.....	436
Nayarit Squirrel.....	63	Northern Weasel.....	429
Near sighted Pocket Gopher.....	276	Northern White-footed Deer Mouse	140
Neat Bushy-tailed Wood Rat.....	216	Northern White-tailed Deer.....	44
Neat Field Mouse.....	159	Northern Woodland Jumping Mouse	328
Nebraska Coyote.....	377	Northwest Chipmunk.....	90
Nebraska Field Mouse.....	146	Northwest Jumping Mouse.....	326
Nebraska Harvest Mouse.....	192	Northwest Pocket Mouse.....	308
Neglected Chipmunk.....	93	Northwest Territory Musk Rat...	254
Neglected Pocket Gopher.....	268	Northwestern Black Bear.....	395
Neglected Spermophile.....	103	Northwestern Puma.....	369
Nelson's Coati.....	402	Northwestern Red-backed Vole...	229
Nelson's Field Mouse.....	172	Notable Kangaroo Rat.....	293
Nelson's Grison.....	419	Norway Rat.....	129
Nelson's Hare.....	340	Nushagak Spermophile.....	109
Nelson's Lemming.....	260	<i>Nutria</i>	435
Nelson's Mountain Sheep.....	54	Nuttall's Cotton-tail.....	348
Nelson's Pocket Gopher.....	269, 286	Nuttall's Golden Mouse.....	160
Nelson's Pocket Mouse.....	313		
Nelson's Shrew.....	465	Oak Woods Squirrel.....	70
Nelson's Spermophile.....	99	Oaxaca Field Mouse.....	168
Nelson's Spiny Mouse.....	323	Oaxaca Harvest Mouse.....	202
Nelson's Squirrel.....	72	Oaxaca Meadow Vole.....	246
Nelson's White-tailed Deer.....	47	Oaxaca Opossum.....	3
Nelson's Wood Rat.....	212, 224	Oaxaca Spotted Skunk.....	416
Nenton Wood Rat.....	211	Oaxaca Squirrel.....	69
Nevada Jumping Mouse.....	326	Ocelot.....	366
Nevada Kangaroo Rat.....	288	Ocelot, Carriker's.....	366
Nevada Meadow Vole.....	240	Ocelot, Costa Rica.....	366
Nevada Pocket Gopher.....	272	Ocelots, etc.....	365
Nevada Pocket Mouse.....	305	Ochraceous Red-backed Vole.....	230
Nevada Shrew.....	449	Ochraceous-faced Cotton Rat.....	179
New England Gray Fox.....	388	Ochraceous-footed Coyote.....	379
New Mexican Kit Fox.....	384	Ocotlan Cotton Rat.....	177
Newfoundland Caribou.....	38	Ocotlan Rice Rat.....	183
Newfoundland Marten.....	422	Ocotlan Slender Shrew.....	463
Newfoundland Meadow Vole.....	238	Octodonts.....	328
Nicaragua Squirrel.....	75	Odd-tailed Mouse.....	166
Nicaraguan Spiny Rat.....	329	Orsted's Titi Monkey.....	534
Nicholls' Bat.....	527	Olivaceous Spermophile.....	104
Night-wandering Bat.....	480	Olympic Meadow Vole.....	244
Nimble Kangaroo Rat.....	293	Olympic Mountain Vole.....	227
Nine-banded Armadillo.....	13	Olympic Mountains Flying Squirrel	123
Noisy Chickaree.....	80	Olympic Mountains Sewellel.....	127
Noisy Squirrel Monkey.....	534	Olympic Mountains Spotted Skunk	415
Noland's Ranch Coyote.....	378	Olympic Mountains Woodchuck...	121
Northern Chipmunk.....	92	Omlilteme Cotton-tail.....	342
Northern Finback Whale.....	20	<i>o' Mono Chillon</i>	533
Northern Meadow Vole.....	247	<i>Ondatra</i>	254

	PAGE.		PAGE.
<i>Onza</i>	368	<i>Oso Colmenero</i>	11
Opossum, Allen's	5	Otter, Canada	434
Opossum, Alston's	4	Otter, Central American	435
Opossum, Ashy	2	Otter, Degenerate	435
Opossum, Batty's	8	Otter, Florida	434
Opossum, Earl of Derby's	4	Otter, Mexican	434
Opossum, Eten	8	Otter, Pacific	434
Opossum, Fulvous-bellied	3	Otter, San Miguel Island Sea	436
Opossum, Gray	3	Otter, Sea-girt or Island	437
Opossum, Island	7	Otter, Southern	434
Opossum, Island of Cozumel	6	Otters	404, 433
Opossum, Linnean	6	Otters, Sea	435
Opossum, Maria Madre Island	3	Ounce-like Cat	367
Opossum, Mexican Murine	2	Ox, Black Musk	56
Opossum, Murine	2	Ox, Musk	56
Opossum, Oaxaca	3	Ox, Ward's Musk	56
Opossum, Orizaba	5	Oxen	52
Opossum, Pale Woolly	4	Oxen, Musk	55
Opossum, Rat-tailed	5	Ozolotepec Field Mouse	170
Opossum, Richmond's	8		
Opossum, Sinaloa	3	Paca	335, 337
Opossum, Tabasco	6	Paca, Central American	337
Opossum, Texas	6	Pacific Big-eared Bat	492
Opossum, Virginian	7	Pacific Coast Beaver	129
Opossum, Water	1	Pacific Coast Raccoon	401
Opossum, Yucatan	5	Pacific Coast Wapiti	42
Opossums	1	Pacific Coast White-footed Deer	
Opusura Harvest Mouse	198	Mouse	141
Orca Red-backed Vole	230	Pacific Finback Whale	19
Ord's Kangaroo Rat	296	Pacific Fisher	423
Oregon Flying Squirrel	123	Pacific Jumping Mouse	327
Oregon Kangaroo Rat	299	Pacific Marten	422
Oregon Meadow Vole	250	Pacific Ocean Dolphin	32
Oregon Musk Rat	254	Pacific Otter	434
Oregon Pocket Gopher	277	Pacific Pocket Mouse	306
Oregon Pocket Mouse	306	Pacific Sewellel	127
Oregon Sperophile	113	Pacific Shrew	454
Oregon Weasel	431	Pacific Walrus	439
Orizaba Cotton-tail	351	Pacuare Pocket Gopher	271
Orizaba Field Mouse	169	Padre Island Sperophile	100
Orizaba Harvest Mouse	200	Painted Chipmunk	94
Orizaba Opossum	5	Painted Spiny Mouse	322
Orizaba Pocket Gopher	283	Painted Wood Rat	219
Orizaba Wood Rat	219	Pale Bat	477
Ornamented Kangaroo Rat	291	Pale Michigan Mouse	160
Ornamented Shrew	452	Pale Mole Mouse	131
Osborn's Caribou	38	Pale Musk Rat	253
Osceola's White-tailed Deer	44	Pale Pocket Gopher	263, 286
Osgood's Field Mouse	147	Pale Raccoon	401
Osgood's Pocket Mouse	304	Pale Rice Rat	186
Osgood's Sperophile	109, 110	Pale Sperophile	103

	PAGE.
Pale Woolly Opossum.....	4
Paler Kangaroo Rat.....	292
Pallid Big-eared Bat	491
Pallid Coati	403
Pallid Kangaroo Rat	300
Pallid Lynx	371
Pallid Meadow Vole	247
Pallid Pocket Gopher.....	274
Pallid Pocket Mouse.....	314
Pallid Spermophile.....	104
Palm Springs Free-tailed Bat.....	498
Palmer's Chipmunk.....	86
Palmer's Kangaroo Rat.....	296
Panama Bat.....	488
Panama Rat.....	174
Panama Rice Rat.....	182
Panama Spear-nosed Bat.....	514
Panama Spiny Rat.....	330
Panamint Kangaroo Rat.....	295
Panamint Mountains Chipmunk...	90
Panamint Pocket Mouse	304
Parasite Mouse.....	151
Park Spermophile.....	100
Parnell's Bat.....	505
Parry's Spermophile.....	108
Patzcuaro Harvest Mouse.....	203
Peale's Meadow Vole.....	238
Peary's Caribou.....	40
Peccary, Armeria.....	35
Peccary, Boquete.....	36
Peccary, Dwarf.....	35
Peccary, Heavy.....	36
Peccary, Savage.....	37
Peccary, Sonora.....	36
Peccary, Texan.....	35
Peccary, White-lipped.....	37
Peccary, Yucatan.....	36
Peccaries.....	35
Pecos River Musk Rat.....	252
Penciled Field Mouse	150
Peninsula Shrew.....	461
Peninsular Harvest Mouse	197
Pennant's Fisher.....	422
Pennsylvania Meadow Vole.....	235
Pennsylvania Wood Rat.....	210
Penobscot Field Vole	235
Perote Kangaroo Rat.....	291
Perote Mouse.....	158
Perote Pocket Gopher.....	265
Perote Spermophile.....	103
Perote Squirrel.....	69

	PAGE.
<i>Perote, Tepachiche del Cofre de</i>	399
<i>Perote, Tepichichi del Cofre de</i>	399
Perote Wood Rat.....	225
<i>Perrito</i>	117
<i>Perrito del Campo</i>	117
Peters' Vampire Bat.....	512
Peters' White-striped Bat	526
Pfeiffer's Bat	486
Phillips' Kangaroo Rat.....	291
Pigmy Meadow Vole	247
Pigmy Sperm Whale.....	22
Pigmy Spotted Skunk.....	415
Pika, Chief.....	338
Pika, Collared.....	338
Pika, Dainty.....	339
Pika, Gray-headed.....	338
Pika, Least.....	339
Pika, Rock.....	339
Pikas.....	337
Pilot or Ca'ing Whale.....	27
Pine Barrens Pocket Gopher.....	261
Pine Meadow Vole.....	249
Pine Pocket Gopher	275
Pine Wood Cotton-tail	350
Pine Wood Rat.....	211
Pine Zone Pocket Gopher.....	271
<i>Pisoti</i>	402
Plain-backed Squirrel.....	67
Plains Cotton-tail.....	348
Plains Hare.....	362
Plains Pocket Mouse.....	301
Plain-tailed Spermophile	116
Plantain Spiny Mouse.....	322
Plateado Wood Rat.....	212
Pocket Gophers.....	261, 271
Pocket Mice.....	288, 300
Poey's Bat	521
Polar Bear.....	390
Polar Hare.....	358
Popof Island Meadow Vole	242
Popof Island Shrew.....	451
Porcupine, Alaska.....	333
Porcupine, Canadian Tree.....	334
Porcupine, Chiriqui.....	335
Porcupine, Coues' Tree.....	333
Porcupine, Dusky.....	333
Porcupine, Labrador.....	334
Porcupine, Light-colored.....	334
Porcupine, Prehensile-tailed.....	335
Porcupine, Rothschild's Tree.....	335
Porcupine, Western.....	332

	PAGE.		PAGE.
Porcupine, Yucatan Tree.....	335	Quadrupeds, Hoofed.....	35
Porcupines, Long-spined.....	332	Queen Charlotte Island Marten....	420
Porpoise, Bottle-nosed.....	32	Queen Charlotte Islands Black	
Porpoise, Common.....	25	Bear.....	394
Porpoise, Dall's.....	26	Queen Charlotte Shrew.....	450
Porpoises.....	24, 25	Querendaro Wood Rat.....	212
Porto Rican Bat.....	505		
Porto Rico Nose-leaf Bat.....	518	Rabbit, Allen's Jack.....	353
Pouched Bat.....	493	Rabbit, Alta Mira Jack.....	354
Pouched Rats.....	261	Rabbit, Batty's Jack.....	355
Powerful Pocket Gophers.....	264	Rabbit, Beautiful-eared Jack.....	353
Prairie Jumping Mouse.....	325	Rabbit, Black-eared Jack.....	355
Prairie Meadow Vole.....	248	Rabbit, California Jack.....	357
Prairie-dog.....	117	Rabbit, Desert Jack.....	356
Prairie-dog, Arizona.....	118	Rabbit, Esperito Santo Island Jack	358
Prairie-dog, Gunnison's.....	118	Rabbit, Gaillard's Jack.....	354
Prairie-dog, Lewis'.....	118	Rabbit, Gray Desert Jack.....	356
Prairie-dog, Mexican.....	119	Rabbit, Hidalgo Jack.....	354
Prairie-dog, Red.....	117	Rabbit, Lower California Jack.....	358
Prairie-dog, Utah.....	118	Rabbit, Merriam's Jack.....	354
Preble's Lemming Vole.....	258	Rabbit, San Pedro Martir Jack....	358
Preble's Vole.....	227	Rabbit, Texas Jack.....	355
Prehensile-tailed Hutia.....	331	Rabbit, Tulare Jack.....	357
Prehensile-tailed Monkeys.....	532	Rabbit, Wallawalla Jack.....	356
Prehensile-tailed Porcupine.....	335	Rabbit, Wandering Jack.....	353
Prevost Island Mouse.....	152	Rabbit, Western Desert Jack.....	356
Prevost Island Shrew.....	450	Rabbits.....	339
Pribilof Island Arctic Fox.....	383	Rabbits, Jack.....	353
Pribilof Islands Fur Seal.....	438	Raccoon, Common.....	400
Pribilof Islands Seal.....	442	Raccoon, Crab-eating.....	402
Price's Chipmunk.....	87	Raccoon, Hernandez's.....	401
Price's Pocket Mouse.....	311	Raccoon, Island.....	401
Primates.....	531	Raccoon, Little.....	401
Projecting-teeth Squirrel.....	59	Raccoon, Maynard's.....	402
Prominent-eared Bat.....	474	Raccoon, Pacific Coast.....	401
Prong-horn Antelope.....	51	Raccoon, Pale.....	401
Prong-horn Antelopes.....	51	Raccoon, Sleepy.....	400
Prong-horn, Mexican.....	51	Raccoon-fox, Boquete.....	399
Proud Kangaroo Rat.....	292	Raccoon-fox, Common.....	397
Puebla Cotton-tail.....	345	Raccoon-fox, Esperito Santo.....	398
Puebla Squirrel.....	70	Raccoon-fox, Robber.....	398
Puerto Angel Mouse.....	163	Raccoon-fox, Saussure's.....	399
Puget Sound Skunk.....	408	Raccoon-fox, Southern.....	399
Pugnacious Weasel.....	428	Raccoon-fox, Tawny.....	398
Pug-nosed Flying Squirrel.....	125	Raccoon-fox, White-footed.....	399
Pug-nosed Mastiff Bat.....	496	Raccoons.....	397
Puma, Central American.....	370	Rafinesque's Bat.....	488
Puma, Eastern.....	370	Rafinesque's White-tailed Deer....	44
Puma, Northwestern.....	369	Rambling Mink.....	424
Pumas.....	369	Ramona Mole Mouse.....	133
Pygmy Pocket Gopher.....	284	Rat, Active Wood.....	213

	PAGE.		PAGE.
Rat, Alfaro's Rice.....	183, 191	Rat, Cheating Wood.....	225
Rat, Allen's Cotton.....	179	Rat, Cherrie's Rice.....	190
Rat, Allen's Wood.....	225	Rat, Chichen Itza Rice.....	189
Rat, Allied Kangaroo.....	289	Rat, Cinnamon Bushy-tailed Wood	224
Rat, Allied Spiny.....	317	Rat, Coast Cotton.....	175
Rat, Allied Wood.....	207	Rat, Coast Wood.....	209
Rat, Alpine Cotton.....	179	Rat, Coban Spiny.....	318
Rat, Alston's Wood.....	225	Rat, Collared Wood.....	208
Rat, Amoles Cotton.....	179	Rat, Collector Wood.....	205
Rat, Apazote.....	174	Rat, Colorado Bushy-tailed Wood..	223
Rat, Aquatic Rice.....	182	Rat, Colombian Kangaroo.....	297
Rat, Arizona Busy-tailed Wood...	223	Rat, Coues' Rice.....	186
Rat, Arizona Cotton.....	176	Rat, Counterfeit Kangaroo	295
Rat, Atemajac Wood.....	211	Rat, Cozumel Island Rice	182
Rat, Attwater's Wood	210	Rat, Death Valley Kangaroo.....	290
Rat, Aztec Rice.....	184	Rat, Death Valley Wood	215
Rat, Bailey's Cotton.....	177	Rat, Deceitful Rice.....	185
Rat, Bailey's Wood.....	210	Rat, Deceitful Wood.....	216
Rat, Baird's Kangaroo.....	295	Rat, Desert Kangaroo	292
Rat, Beautiful Wood.....	218	Rat, Dissimilar Wood.....	206
Rat, Berlandier's Cotton	176	Rat, Doubtful Kangaroo.....	289
Rat, Black	129	Rat, Drummond's Bushy-tailed	
Rat, Black Rice.....	190	Wood.....	221
Rat, Black Spiny.....	317	Rat, Durango Wood.....	218
Rat, Black-eared Cotton.....	180	Rat, Dusky Kangaroo.....	299
Rat, Black-eared Rice.....	187	Rat, Dusky Musk	255
Rat, Black-nosed Kangaroo.....	290	Rat, Dusky-footed Wood.....	206
Rat, Black-tailed Kangaroo.....	291	Rat, El General Rice.....	189
Rat, Black-tailed Wood.....	217	Rat, Escondido River Rice.....	184
Rat, Bogava Rice.....	189	Rat, Faded Vesper	172
Rat, Bogava Spiny.....	329	Rat, Field Wood.....	218
Rat, Bolaños Wood.....	205	Rat, Flat-headed Wood	209
Rat, Boqueron Cotton	178	Rat, Florida Rice.....	181
Rat, Boquete Spiny.....	318	Rat, Florida Wood.....	210
Rat, Boquete Vesper.....	171	Rat, Fulvous-bellied.....	220
Rat, British Columbia Bushy-tailed		Rat, Fulvous-bellied Cotton.....	180
Wood.....	221	Rat, Fulvous Rice.....	189
Rat, Broad-nosed Rice.....	188	Rat, Fuscous Bushy-tailed Wood..	221
Rat, Brown.....	130	Rat, Gaumer's Spiny.....	317
Rat, Brown-rumped Cotton.....	175	Rat, Gliding Spiny	329
Rat, Buller's Rice.....	185	Rat, Goldman's Kangaroo.....	294
Rat, Buller's Spring.....	317	Rat, Goldman's Rice.....	186
Rat, Cabezon Kangaroo.....	295	Rat, Goldman's Spiny.....	318
Rat, California Kangaroo	292, 300	Rat, Goldman's Wood	220
Rat, Cameron Lake Wood.....	215	Rat, Graceful Kangaroo.....	294
Rat, Cape Sable	181	Rat, Granger's Bushy-tailed Wood	223
Rat, Catemaco Rice	185	Rat, Gray Bushy-tailed Wood.....	220
Rat, Ceiba Cotton.....	178	Rat, Gray Spiny	319
Rat, Cerros Island Wood.....	208	Rat, Gray-faced Wood.....	226
Rat, Chapman's Kangaroo.....	297	Rat, Great Kangaroo.....	298
Rat, Chapman's Rice.....	186	Rat, Guatemalan Rice.....	185

	PAGE.		PAGE.
Rat, Heller's Kangaroo.....	292	Rat, Ocotlan Cotton.....	177
Rat, Hispid Cotton.....	175	Rat, Ocotlan Rice.....	183
Rat, Hispid Spiny.....	318	Rat, Ord's Kangaroo.....	296
Rat, Hudson Bay Musk.....	253	Rat, Oregon Kangaroo.....	299
Rat, Isolated Rice.....	191	Rat, Oregon Musk.....	254
Rat, Isthmian Wood.....	219	Rat, Orizaba Wood.....	219
Rat, Jalapa Rice.....	184	Rat, Ornamented Kangaroo.....	291
Rat, Jamaica Rice.....	188	Rat, Painted Wood.....	219
Rat, Juquila Wood.....	220	Rat, Pale Musk.....	253
Rat, Keeler Kangaroo.....	288	Rat, Pale Rice.....	186
Rat, La Parada Spiny.....	317	Rat, Paler Kangaroo.....	292
Rat, Large Cotton.....	177	Rat, Pallid Kangaroo.....	300
Rat, Large-eared Wood.....	206	Rat, Palmer's Kangaroo.....	296
Rat, Large-headed Kangaroo.....	299	Rat, Panama.....	174
Rat, Large-toothed Musk.....	255	Rat, Panama Rice.....	182
Rat, Large-toothed Rice.....	188	Rat, Panama Spiny.....	330
Rat, Least Cotton.....	178	Rat, Panamint Kangaroo.....	295
Rat, Light-footed Kangaroo.....	298	Rat, Pecos River Musk.....	252
Rat, Lonely Rice.....	188	Rat, Pennsylvania Wood.....	210
Rat, Long-footed Kangaroo.....	298	Rat, Perote Kangaroo.....	291
Rat, Long-haired Rice.....	183	Rat, Perote Wood.....	224
Rat, Long-tailed Rice.....	186	Rat, Phillips' Kangaroo.....	291
Rat, Long-tailed Spiny.....	318	Rat, Pine Wood.....	211
Rat, Lower California Rice.....	182	Rat, Plateado Wood.....	212
Rat, Lower California Wood.....	214	Rat, Proud Kangaroo.....	292
Rat, Maria Madre Island Rice.....	182	Rat, Querendaro Wood.....	212
Rat, Marsh Rice.....	187	Rat, Reddish Wood.....	210
Rat, Mascota Cotton.....	176	Rat, Related Musk.....	253
Rat, Merriam's Kangaroo.....	288	Rat, Related Wood.....	207
Rat, Mexican Wood.....	213	Rat, Rhoads' Wood.....	216
Rat, Mohave Desert Wood.....	207	Rat, Richardson's Kangaroo.....	297
Rat, Montezuma Wood.....	218	Rat, Rock-dwelling Bushy-tailed Wood.....	223
Rat, Mt. Tancitaro Wood.....	224	Rat, Rock-loving Wood.....	221
Rat, Mt. Pinos Wood.....	208	Rat, Rufous Rice.....	185
Rat, Mountain Bushy-tailed Wood.....	221	Rat, Rusty Wood.....	211
Rat, Mountain Wood.....	225	Rat, St. Lucia Rice.....	190
Rat, Musk.....	254	Rat, St. Vincent Rice.....	188
Rat, Naked-tailed.....	173	Rat, Salvin's Spiny.....	317
Rat, Naked-tailed Wood.....	214	Rat, San Felipe Desert.....	218
Rat, Narrow-headed Wood.....	217	Rat, San Felipe Kangaroo.....	290
Rat, Neat Bushy-tailed Wood.....	216	Rat, San Martin Island Wood.....	214
Rat, Nelson's Wood.....	212, 224	Rat, San Miguel Spiny.....	330
Rat, Nenton Wood.....	211	Rat, Santa Catalina Wood.....	213
Rat, Nevada Kangaroo.....	288	Rat, Santa Maria Volcan Rice.....	187
Rat, Nicaraguan Spiny.....	329	Rat, Sennett's Kangaroo.....	297
Rat, Nimble Kangaroo.....	293	Rat, Shining Rice.....	183
Rat, Northern Musk.....	253	Rat, Sierra Madre Wood.....	213
Rat, Northwest Territory Musk.....	254	Rat, Similar Kangaroo.....	288
Rat, Norway.....	129	Rat, Sinaloa Wood.....	214
Rat, Notable Kangaroo.....	293	Rat, Short-spined.....	329
Rat, Ochraceous-faced Cotton.....	179		

	PAGE.		PAGE.
Rat, Short-tailed Spiny	319	Rats, Bushy-tailed Wood.....	220
Rat, Slender Kangaroo.....	289	Rats, Cotton.....	174
Rat, Slender-tailed Wood.....	219	Rats, Five-toed Kangaroo.....	296
Rat, Small Kangaroo.....	289	Rats, Four-toed Kangaroo.....	288
Rat, Small-face Kangaroo.....	297	Rats, Kangaroo.....	288
Rat, Small-footed Wood.....	209	Rats, Musk.....	252
Rat, Small-toothed Cotton.....	178	Rats, Musk-bearing.....	189
Rat, Solitary Wood.....	215	Rats, Pouched.....	261
Rat, Spotted Spiny.....	319	Rats, Rice.....	180
Rat, Stephens' Wood.....	216	Rats, Small Kangaroo.....	299
Rat, Streater's Kangaroo.....	294	Rats, Spiny Pouched.....	316
Rat, Streater's Wood.....	206	Rats, Wood.....	205
Rat, Striped-face Rice	187	Rat-tailed Opossum.....	5
Rat, Surre Rice.....	190	Red Bat.....	486
Rat, Sumichrast's Vesper.....	172	Red Fox.....	380
Rat, Surber's Wood.....	209	Red Lynx.....	371
Rat, Swamp Rice.....	181	Red Prairie-dog.....	117
Rat, Swimming Rice.....	181	Red-backed Voles.....	229
Rat, Tabascan Rice.....	187	Reddish Brown Pocket Gopher....	272
Rat, Talamanca Rice.....	183	Reddish Fox.....	380
Rat, Teapa Cotton.....	178	Reddish Meadow Vole.....	240
Rat, Teapa Rice.....	185	Reddish Mouse.....	153
Rat, Texan Cotton	175	Reddish Wood Rat.....	210
Rat, Texolo Wood.....	207	Redman's Nose-leaf Bat.....	519
Rat, Thick-set Kangaroo.....	299	Red-thighed Spermophile.....	108
Rat, Tiburon Island Kangaroo....	291	Related Chipmunk.....	90, 94
Rat, Todos Santos Wood.....	214	Related Musk Rat.....	253
Rat, Toltec Cotton	177	Related White-footed Mouse.....	155
Rat, Tonalá Cotton.....	177	Related Wood Rat	207
Rat, Tonalá Rice.....	184	Restless Bat.....	489
Rat, Tropical Wood	220	Rhoads' Mole Mouse.....	132
Rat, Tulare Kangaroo	293	Rhoads' Mouse.....	139
Rat, Tumbala.....	173	Rhoads' Pocket Gopher.....	287
Rat, Tumbala Rice.....	187	Rhoads' Red-backed Vole.....	230
Rat, Tunkas.....	174	Rhoads' Wood Rat.....	216
Rat, Tuxtla.....	173	Ribbon Seal.....	442
Rat, Utah Kangaroo.....	296	Rice Rats.....	180
Rat, Volcan de Chiriqui Cotton ..	180	Richard's Harbor Seal.....	442
Rat, Wandering Rice.....	184	Richardson's Chickaree.....	82
Rat, Watson's	174	Richardson's Flying Squirrel.....	122
Rat, Western Bushy-tailed Wood .	221	Richardson's Hare.....	363
Rat, Western Desert Cotton.....	176	Richardson's Kangaroo Rat.....	297
Rat, White-bellied Rice.....	182	Richardson's Lemming.....	259
Rat, White-eared Cotton.....	179	Richardson's Meadow Vole	251
Rat, White-throated Wood.....	217	Richardson's Pocket Gopher.....	279
Rat, White-tipped Bushy-tailed		Richardson's Shrew.....	447
Wood.....	221	Richardson's Spermophile.....	111
Rat, White-toothed Wood.....	212	Richardson's Weasel.....	426
Rat, Yellow Wood.....	217	Rich-colored Cotton-tail.....	352
Rat, Zacatecas Wood.....	226	Rich-colored Mouse.....	146
Rats.....	129	Rich-colored Red-backed Vole	231

	PAGE.		PAGE.
Richly-colored Weasel.....	430	Sachem or Chief Mouse.....	156
Richmond's Opossum.....	8	Saddle-backed Anteater.....	12
Richmond's Squirrel.....	62	St. Kitts Bat.....	523
Ringed Seal.....	444	St. Lucia Rice Rat.....	190
Ring-tailed Sperophile.....	116	St. Martin's Bat.....	524
Rio Grande Bridled Weasel.....	432	St. Matthew Island Meadow Vole..	243
Rio Grande White-footed Mouse ..	142	St. Michaels' Hare.....	360
Rio Managua Squirrel.....	77	St. Paul Island Shrew.....	454
Rio Sestin Meadow Mouse.....	194	St. Vincent Rice Rat.....	188
Roaming Pocket Gopher.....	285	Salvin's Nose-leafed Bat.....	528
Roan Mountain Jumping Mouse...	328	Salvin's Shrew.....	455
Roan Mountain Mouse.....	145	Salvin's Spiny Rat.....	317
Robber Coyote.....	377	Samalayuca Mole Mouse.....	131
Robber Raccoon-fox.....	398	San Bernardino Brown Bat.....	484
Robust Field Mouse.....	160	San Bernardino Sperophile.....	107
Rock Meadow Vole.....	246	San Clemente Island Gray Fox....	390
Rock Mouse.....	169	San Clemente Island Mouse.....	145
Rock Pika.....	339	San Cristobal Field Mouse.....	170
Rock Sperophile.....	97, 115	San Cristobal Shrew.....	455
Rock Spotted Skunk.....	414	San Diego Cotton-tail.....	346
Rock-dwelling Bushy-tailed Wood Rat.....	223	San Felipe Desert Rat.....	218
Rock-loving Wood Rat.....	221	San Felipe Field Mouse.....	167
Rocky Mountain Jumping Mouse..	325	San Felipe Kangaroo Rat.....	290
Rocky Mountain Marten.....	422	San Felipe Skunk.....	412
Rodents.....	59	San Geronimo Harbor Seal.....	443
Rothschild's Tree Porcupine.....	335	San Geronimo Island Field Mouse..	162
Rothschild's White-tailed Deer....	46	San Joaquin Pocket Mouse.....	306
Round-eared Bat.....	512	San Joaquin Valley Elk.....	42
Round-tailed Sperophile.....	102	San Martin Island Mouse.....	148
Rowley's Field Mouse.....	151	San Martin Island Wood Rat.....	214
Rowley's Pine Mouse.....	151	San Miguel Island Gray Fox.....	388
Royal Fox.....	382	San Miguel Island Sea Otter.....	436
Ruatan Island Agouti.....	336	San Miguel Spiny Rat.....	330
Rufescent Pocket Gopher.....	279	San Pedro Martir Jack Rabbit....	358
Rufous Harvest Mouse.....	201	San Pedro Martir Mountains Mouse	159
Rufous Mastiff Bat.....	495	San Pedro Martir Pocket Gopher..	282
Rufous Red-backed Vole.....	229	San Pedro Martir Spotted Skunk..	414
Rufous Rice Rat.....	185	San Quentin Mouse.....	148
Rufous Sewellel.....	127	San Quentin Pocket Mouse.....	309
Rufous Vampire Bat.....	530	San Sebastian Harvest Mouse	203
Rufous-foot Squirrel Monkey.....	534	Sand-dwelling Pocket Mouse.....	305
Ruidoso Chickaree.....	84	Sand-frequenting Pocket Gopher..	264
Ruidoso Mole Mouse.....	131	Sand-loving Mole Mouse.....	133
Russet Cotton-tail.....	348	Sand-loving Mouse.....	137
Russet-footed Cotton-tail.....	346	Santa Anita Bat.....	502
Rusty Wood Rat.....	211	Santa Anita Pocket Gopher.....	280
		Santa Catalina Island Gray Fox...	389
		Santa Catalina Island Harvest Mouse.....	195
Sable, Hudson Bay.....	419	Santa Catalina Wood Rat.....	213
Sables.....	419	Santa Cruz Island Gray Fox.....	388

	PAGE.		PAGE.
Santa Lucia Bat.....	527	Sewellel, Olympic Mountains.....	127
Santa Lucia Nose-leaf Bat.....	518	Sewellel, Pacific.....	127
Santa Maria Volcan Rice Rat.....	187	Sewellel, Rufous.....	127
Santa Rosalia Bat.....	489	Sewellels.....	126
Sapajou, Allied.....	536	Sezekorn's Bat.....	521
Sartori's Brocket.....	50	Shaded Field Mouse.....	144
Saskatchewan Jumping Mouse.....	326	Shaggy-eared Bat.....	505
Saskatchewan Marten.....	419	Sharp-beaked Bottle-nosed Whale.....	29
Saussure's Large-eared Bat.....	510	Sharp-nosed Finback Whale.....	19
Saussure's Raccoon Fox.....	399	Sharp-sighted Lynx.....	371
Saussure's Shrew.....	456	Sharp-toothed Dolphin.....	33
Savage Peccary.....	37	Shaved Mastiff Bat.....	497
Sawatch Meadow Vole.....	236	Shaw's Pocket Gopher.....	263
Scammon's Blackfish.....	28	Sheep.....	52, 53
Sclater's Shrew.....	456	Sheep, Audubon's Mountain.....	53
Scolding Chickaree.....	82	Sheep, Dall's Mountain.....	55
Scott's Gray Fox.....	386	Sheep, Fannin's Mountain.....	55
Sea Bears.....	438	Sheep, Kenai Mountain.....	55
Sea Cows.....	14	Sheep, Lower California Mountain.....	54
Sea Elephants.....	439	Sheep, Mexican Mountain.....	54
Sea Otters.....	435	Sheep, Mountain.....	53
Sea-girt or Island Otter.....	433	Sheep, Nelson's Mountain.....	54
Seal, Bearded.....	441	Sheep, Stone's.....	54
Seal, Elephant.....	440	Shining Rice Rat.....	183
Seal, Fur.....	438	Short - eared California Pocket	
Seal, Gray.....	440	Mouse.....	314
Seal, Guadalupe Fur.....	438	Short-finned Blackfish.....	28
Seal, Harbor.....	442	Short-handed Bat.....	503
Seal, Harp.....	444	Short-headed Pocket Gopher.....	263
Seal, Hooded.....	440	Short-nosed Pocket Mouse.....	305
Seal, Pribilof Islands.....	442	Short-spined Rat.....	329
Seal, Pribilof Islands Fur.....	438	Short-tailed Bat.....	515
Seal, Ribbon.....	442	Short-tailed Cotton-tails.....	351
Seal, Richard's Harbor.....	442	Short-tailed Hares.....	352
Seal, Ringed.....	443	Short-tailed Hutia.....	331
Seal, San Geronimo Harbor.....	443	Short-tailed Meadow Vole.....	247
Seal, Stejneger's Harbor.....	443	Short-tailed Mole Mouse.....	131
Seal, West Indian.....	445	Short-tailed Red-backed Vole.....	231
Sea-lion, California.....	437	Short-tailed Shrew.....	461
Sea-lion, Steller's.....	437	Short-tailed Shrews.....	460
Sea-lions.....	437	Short-tailed Spiny Rat.....	319
Seals.....	437	Showy Chipmunk.....	89
Seals, Gray.....	440	Shrew, Alaska Swimming.....	458
Seals, Hair.....	440	Shrew, Arctic.....	447
Seals, Hooded.....	440	Shrew, Baird's.....	452
Secretive Vole.....	227	Shrew, Bendire's.....	459
Selkirk Marten.....	420	Shrew, Berlandier's.....	462
Sennett's Kangaroo Rat.....	297	Shrew, Blackish.....	463
Sewellel, California.....	127	Shrew, California.....	452
Sewellel, Dark.....	127	Shrew, Carolina.....	461
Sewellel, Mt. Rainier.....	127	Shrew, Changeable.....	456

	PAGE.		PAGE.
Shrew, Charming.....	449	Shrew, Salvin's.....	455
Shrew, Chestnut-bellied.....	450	Shrew, San Cristobal.....	455
Shrew, Coues' Mexican.....	464	Shrew, Saussure's.....	456
Shrew, Crawford's.....	459	Shrew, Sclater's.....	455
Shrew, Delicate.....	453	Shrew, Short-tailed.....	461
Shrew, Dismal Swamp.....	463	Shrew, Small.....	463
Shrew, Dobson's.....	448	Shrew, Smoky.....	447
Shrew, Dobson's Swimming.....	458	Shrew, Sphagnum Moss.....	447
Shrew, Dusky.....	450, 464	Shrew, Spiny.....	448
Shrew, Dwarf.....	453	Shrew, Streator's.....	447
Shrew, Extraordinary.....	457	Shrew, Thick-set.....	462
Shrew, Fisher's.....	454	Shrew, Tlalpam.....	464
Shrew, Florida.....	463	Shrew, Tropical.....	464
Shrew, Giant.....	460	Shrew, Trowbridge's.....	452
Shrew, Glacier Bay.....	451	Shrew, Tundra.....	447
Shrew, Godman's.....	455	Shrew, Vancouver Island.....	449
Shrew, Goldman's.....	465	Shrew, Volcano of Irazú.....	464
Shrew, Great.....	466	Shrew, Wandering.....	465, 448
Shrew, Guatemalan.....	455	Shrew, Warring.....	465
Shrew, Hoy's.....	457	Shrew, White Mountains.....	453
Shrew, Irrational.....	462	Shrew, White-bellied.....	459
Shrew, Keewatin.....	457	Shrew, White-tipped Swimming...	458
Shrew, Labrador.....	455	Shrew, Wood.....	461
Shrew, Large-toothed.....	455	Shrew, Yakutat Bay.....	451
Shrew, Lofty Mountain.....	451	Shrew-like Bat.....	516
Shrew, Long-nosed.....	454	Shrew-like Moles.....	466
Shrew, Long-tailed.....	448, 450	Shrews.....	446
Shrew, Marsh.....	458	Shrews Anomalous.....	459
Shrew, Masked.....	446	Shrews, Common.....	446
Shrew, Maya.....	465	Shrews, Short-tailed.....	460
Shrew, Mazatlan.....	460	Shrews, Small.....	456
Shrew, Merriam's.....	454	Shrews, Southern.....	459
Shrew, Minute.....	449	Shrews, Swimming.....	457
Shrew, Monterey.....	452	Shy Field Mouse.....	164
Shrew, Mt. Lyell.....	453	Siberian Hare.....	359
Shrew, Mt. Popocatepetl.....	466	Siebold's Baleen Whale.....	16
Shrew, Mt. Shasta.....	453	Sierra Blanca Pocket Gopher.....	286
Shrew, Mt. Zempoaltepec.....	466	Sierra Hare.....	363
Shrew, Mountain.....	449, 456	Sierra Laguna Bat.....	487
Shrew, Navigator.....	458	Sierra Laguna Pocket Gopher.....	280
Shrew, Nelson's.....	465	Sierra Madre Spermathophile.....	106
Shrew, Nevada.....	449	Sierra Madre Wood Rat.....	213
Shrew, Ocotlan Slender.....	463	Silly Lemming Vole.....	256
Shrew, Ornamented.....	452	Silver Fox.....	380
Shrew, Pacific.....	454	Similar Kangaroo Rat.....	288
Shrew, Peninsula.....	461	Sinaloa Cotton-tail.....	349
Shrew, Popof Island.....	451	Sinaloa Jaguarondi.....	368
Shrew, Prevost Island.....	450	Sinaloa Opossum.....	3
Shrew, Queen Charlotte.....	450	Sinaloa Pocket Gopher.....	287
Shrew, Richardson's.....	447	Sinaloa Pocket Mouse.....	312
Shrew, St. Paul Island.....	454	Sinaloa Squirrel.....	73

	PAGE.		PAGE.
Sinaloa White-tailed Deer.....	47	Skunks, Little-striped.....	412
Sinaloa Wood Rat.....	214	Skunks, Spotted.....	412
Sirenians.....	14	Sleepy Raccoon.....	400
Sitka Black-tailed Deer.....	48	Slender Harvest Mouse.....	199
Sitka Meadow Vole.....	242	Slender Kangaroo Rat.....	289
Sitka Mouse.....	152	Slender Spotted Skunk.....	413
Skunk, Arizona.....	406	Slender-nose Anteater.....	11
Skunk, Arizona Spotted.....	414	Slender-nosed Harvest Mouse.....	201
Skunk, Bridled.....	410	Slender-nosed Pocket Gopher.....	267
Skunk, Broad-headed Spotted.....	414	Slender-nosed Pocket Mouse.....	311
Skunk, Broad-nosed.....	409	Slender-tailed Wood Rat.....	219
Skunk, California.....	406	Sloth, Chestnut-headed.....	9
Skunk, Canada.....	406	Sloth, Dusky.....	9
Skunk, Cape St. Lucas Spotted.....	416	Sloth, Hoffmann's.....	9, 11
Skunk, Cascade.....	406	Sloths.....	9
Skunk, Cheating Spotted.....	414	Sluggish Fox.....	384
Skunk, Chihuahuan Little Spotted.....	415	Small Bat.....	517
Skunk, Eastern.....	406	Small Kangaroo Rat.....	289
Skunk, Field Spotted.....	415	Small Kangaroo Rats.....	299
Skunk, Florida.....	406	Small Lemming.....	259
Skunk, Graceful.....	415	Small Shrew.....	456, 463
Skunk, Great Basin.....	406	Small Spotted Cat.....	367
Skunk, Growling Spotted.....	413	Small-eared Nicaraguan Bat.....	513
Skunk, Guadalupe.....	412	Small-eared Weasel.....	426
Skunk, Hooded.....	410	Smaller California Mole.....	469
Skunk, Illinois.....	406	Smallest Spiny Mouse.....	321
Skunk, Indianola Spotted.....	413	Small-faced Kangaroo Rat.....	297
Skunk, Long-tailed Texan.....	406	Small-footed Bat.....	502, 523
Skunk, Louisiana.....	406	Small-footed Hare.....	356
Skunk, Lower California.....	406	Small-footed Wood Rat.....	209
Skunk, Mearns'.....	412	Small-toothed Cotton Rat.....	178
Skunk, Merriam's Spotted.....	413	Small-toothed Fox.....	389
Skunk, Miller's.....	410	Small-toothed Spermophile.....	104
Skunk, Motzorongo.....	412	Small-winged Bat.....	478
Skunk, Narrow-headed Spotted.....	416	Smith's Coyote.....	378
Skunk, Northern Plains.....	406	Smoky Shrew.....	447
Skunk, Oaxaca Spotted.....	416	Snow-loving Red-backed Vole.....	233
Skunk, Olympic Mountains Spotted.....	415	Snowy Bat.....	519
Skunk, Pigmy Spotted.....	415	Soft-furred Chickaree.....	83
Skunk, Puget Sound.....	406	Soft-furred Lynx.....	370
Skunk, Rock Spotted.....	414	Solitary Wood Rat.....	215
Skunk, San Felipe.....	412	Sonora Peccary.....	36
Skunk, San Pedro Martir Spotted.....	414	Sonora Spiny Mouse.....	323
Skunk, Slender Spotted.....	413	Sonora White-footed Field Mouse.....	142
Skunk, Sonoran.....	411	Sonoran Beaver.....	128
Skunk, Texan.....	411	Sonoran Skunk.....	411
Skunk, White-backed.....	411	Sonoran Spermophile.....	103
Skunk, White-spotted.....	411, 413	Sonoyta Desert Mouse.....	165
Skunks.....	405	Sooty Mouse.....	136, 171
Skunks, Hooded.....	410	Sooty Pocket Gopher.....	267
Skunks, Large-striped.....	406	Sornborger's Black Bear.....	394

	PAGE.		PAGE.
South Dakota Chickaree	81	Spermophile, Neglected.....	103
Southern Otter.....	434	Spermophile, Nelson's.....	99
Southern Pocket Mouse.....	308	Spermophile, Nushagak.....	109
Southern Raccoon-fox	399	Spermophile, Olivaceous	104
Southern Shrews.....	459	Spermophile, Oregon.....	113
Southern Weasel.....	429	Spermophile, Osgood's.....	109, 110
Sowerby's Sperm Whale.....	23	Spermophile, Padre Island.....	100
Spear-nosed Bat	514	Spermophile, Pale.....	103
Specter Bat	511	Spermophile, Pallid.....	104
Sperm Whales.....	20	Spermophile, Park.....	100
Spermophile, Alberta.....	111	Spermophile, Parry's.....	108
Spermophile, Allen's.....	105	Spermophile, Perote.....	103
Spermophile, Armed.....	112	Spermophile, Plain-tailed.....	116
Spermophile, Ashy	106	Spermophile, Red-thighed.....	108
Spermophile, Bailey's.....	100	Spermophile, Richardson's	111
Spermophile, Barrow's.....	110	Spermophile, Ring-tailed.....	116
Spermophile, Bay.....	105	Spermophile, Rock.....	97, 115
Spermophile, Beechey's.....	114	Spermophile, Round-tailed	102
Spermophile, Belding's.....	113	Spermophile, San Bernardino	107
Spermophile, Bering's.....	110	Spermophile, Sierra Madre.....	106
Spermophile, Black-headed.....	114	Spermophile, Small-toothed.....	104
Spermophile, Buckley's	115	Spermophile, Sonoran.....	103
Spermophile, Bushy-tailed.....	114, 149	Spermophile, Spotted.....	99
Spermophile, Charming	98	Spermophile, Stephen's	112
Spermophile, Chestnut-tailed	106	Spermophile, Stone's.....	110
Spermophile, Cinnamon-hued.....	97	Spermophile, Striped	106
Spermophile, Columbian	111	Spermophile, Texas.....	105
Spermophile, Dark	106	Spermophile, 13-striped	104
Spermophile, Dark Spotted.....	100	Spermophile, Townsend's.....	112
Spermophile, Death Valley.....	101	Spermophile, Trader.....	98
Spermophile, Desert.....	101	Spermophile, Trinity Mountains...	107
Spermophile, Douglas'.....	115	Spermophile, Utah	115
Spermophile, Drab Gray	101	Spermophile, White-tailed.....	97
Spermophile, Elegant.....	113	Spermophile, Wortman's.....	107
Spermophile, Fisher's.....	114	Spermophile, Yakima.....	112
Spermophile, Franklin's	105	Spermophiles.....	96
Spermophile, Golden-neck.....	107	Sphagnum Moss Shrew.....	447
Spermophile, Goldman's.....	116	Spider Monkeys.....	534
Spermophile, Harris'.....	97	Spiny Pocket Mouse.....	316
Spermophile, Hoary.....	112	Spiny Pouched Rats.....	316
Spermophile, Island	115	Spiny Shrew.....	448
Spermophile, Kadiak Island	108	Spotted Agouti	336
Spermophile, Kennicott's	99, 112	Spotted Great-eared Bat.....	490
Spermophile, Languishing.....	101	Spotted Lynx.....	371
Spermophile, Large Spotted	100	Spotted Skunks.....	412
Spermophile, Little.....	105	Spotted Spermophile.....	99
Spermophile, Long-tailed.....	113	Spotted Spiny Rat	319
Spermophile, Lower California.....	98	Squirrel, Abert's	66
Spermophile, Mexican.....	103	Squirrel, Acapulco.....	74
Spermophile, Mohave Desert.....	103	Squirrel, Alfaro's Pigmy.....	60
Spermophile, Nagai Island.....	109	Squirrel, Allen's.....	63

	PAGE		PAGE.
Squirrel, Apache.....	64	Squirrel, Mexican Flying.....	126
Squirrel, Apazote.....	61	Squirrel, Michoacan.....	70
Squirrel, Arizona Gray.....	64	Squirrel, Minnesota Gray.....	79
Squirrel, Banded-back.....	76	Squirrel, Mimic.....	66
Squirrel, Bangs' Flying.....	124	Squirrel Monkeys.....	533
Squirrel, Barber's.....	68	Squirrel, Mountain.....	72
Squirrel, Black Fox.....	66	Squirrel, Nayarit.....	63
Squirrel, Black-backed.....	62	Squirrel, Nelson's.....	72
Squirrel, Blackfooted Gray.....	78	Squirrel, Nicaragua.....	75
Squirrel, Boreal Flying.....	124	Squirrel, Oak Woods.....	70
Squirrel, Brown's Pigmy.....	60	Squirrel, Oaxaca.....	69
Squirrel, California Flying.....	122	Squirrel, Olmypic Mountains Flying.....	123
Squirrel, California Gray.....	78	Squirrel, Oregon Flying.....	123
<i>Squirrel, Cat.</i>	397	Squirrel, Perote.....	69
Squirrel, Chiapas.....	74	Squirrel, Plain-backed.....	67
Squirrel, Chiriqui.....	62	Squirrel, Projecting-teeth.....	59
Squirrel, Chiriqui Pigmy.....	60	Squirrel, Puebla.....	70
Squirrel, Colima.....	72	Squirrel, Pug-nosed Flying.....	125
Squirrel, Colima Mountain.....	71	Squirrel, Richardson's Flying.....	122
Squirrel, Collie's.....	73	Squirrel, Richmond's.....	62
Squirrel, Common Flying.....	125	Squirrel, Rio Managua.....	77
Squirrel, Deppe's.....	60	Squirrel, Sinaloa.....	73
Squirrel, Durango.....	67	Squirrel, Stephen's Flying.....	124
Squirrel, Dusky Alaskan Flying.....	122	Squirrel, Swarthy.....	75
Squirrel, Dusky Flying.....	123	Squirrel, Tehuantepec.....	74
Squirrel, Dusky Gray.....	79	Squirrel, Texas Fox.....	65
Squirrel, Eastern Fox.....	65	Squirrel, Thomas'.....	75
Squirrel, Escondido River.....	77	Squirrel, Toluca.....	63
Squirrel, Fire-bellied.....	69	Squirrel, True's.....	73
Squirrel, Florida Flying.....	126	Squirrel, Variegated.....	77
Squirrel, Florida Gray.....	80	Squirrel, White-eared Gray.....	79
Squirrel, Fox.....	65	Squirrel, Yucatan.....	75
Squirrel, Frisky Flying.....	123	Squirrel, Yukon Flying.....	125
Squirrel, Golden-bellied.....	68	Squirrels.....	59
Squirrel, Goldman's.....	77	Squirrels, Flying.....	121
Squirrel, Gray's Black.....	76	Squirrels, Tree.....	59
Squirrel, Guatemala.....	74	Star-nosed Mole.....	467
Squirrel, Guerrero.....	72	Star-nosed Moles.....	467
Squirrel, Hoffmann's.....	61	Stejneger's Harbor Seal.....	443
Squirrel, Honduras.....	76	Stejneger's Sperm Whale.....	23
Squirrel, Huachuca.....	64	Steller's Sea Cow.....	14
Squirrel Inn Mouse.....	155	Steller's Sea-lion.....	437
Squirrel, Kaibab.....	67	Stephens' Flying Squirrel.....	124
Squirrel, Klamath Flying.....	123	Stephens' Mouse.....	153
Squirrel, La Cienaga.....	67	Stephens' Pocket Mouse.....	312
Squirrel, La Salado.....	70	Stephens' Spermophile.....	112
Squirrel, Large-eared Flying.....	124	Stephens' Wood Rat.....	216
Squirrel, Lion Hill.....	62	Stickeen Jumping Mouse.....	327
Squirrel, Little Gray.....	61	Stone's Caribou.....	39
Squirrel, Makkovik Flying.....	125	Stone's Lemming Vole.....	225
Squirrel, Manzanillo.....	73	Stone's Sheep.....	54

	PAGE.		PAGE.
Stone's Sperophile.....	110	Teapa Weasel.....	433
Straight-eared Mastiff Bat.....	497	Tehautepec Cotton-tail.....	344
Straight-headed Pocket Gophers ..	268	Tehuantepec Field Mouse	168
Straw-colored Bat.....	501	Tehuantepec Squirrel.....	74
Streator's Chickaree.....	82	<i>Tejon</i>	11
Streator's Kangaroo Rat.....	294	Temporal Dolphin	33
Streator's Shrew	447	Tennessee Cotton Mouse	143
Streator's Weasel.....	427	<i>Tepachiche del Cofre de Perote</i>	399
Streator's Wood Rat.....	206	<i>Tepichichi del Cofre de Perote</i>	399
Striped Chipmunk.....	95	Texan Cotton Rat.....	175
Striped Sperophile.....	106	Texan Peccary.....	35
Striped-face Rice Rat.....	187	Texan Skunk.....	411
Stupid Vole	227	Texan Spiny Mouse.....	323
Subarctic Field Mouse	146	Texan White-tailed Deer.....	45
Suerre Rice Rat.....	190	Texas Field Mouse	145
Sumichrast's Vesper Rat	172	Texas Fox Squirrel.....	65
Summit Arctic Fox.....	382	Texas Gray Fox.....	387
Sunny Mouse.....	154	Texas Jack Rabbit.....	355
Surber's Wood Rat.....	209	Texas Mole.....	471
Swamp Hares.....	340	Texas Opossum.....	6
Swamp Rice Rat.....	181	Texas Pocket Gopher.....	264, 281
Swarthy Mouse	139	Texas Sperophile.....	105
Swarthy Pocket Gopher.....	277	Texolo Wood Rat.....	207
Swarthy Squirrel.....	75	The Dalles Pocket Gopher.....	276
Swift Bat	481	Thick-set Kangaroo Rat	299
Swift Fox.....	384	Thick-set Shrew.....	462
Swift Gray Fox.....	388	Thievish Coyote.....	378
Swift Hare.....	360	13-Striped Sperophile	104
Swift Pocket Gopher.....	284	Thomas' Bat	493
Swift-footed Kit Fox	385	Thomas' Guerrero Field Mouse....	171
Swimming Rice Rat.	181	Thomas' Squirrel.....	75
Swimming Shrews.....	457	Thomas' White-tailed Deer.....	47
		Three-toed Anteater.....	11
Tabascan Rice Rat.....	187	Thurber's Field Mouse.....	146
Tabasco Opossum.....	6	Tiburon Island Kangaroo Rat.....	291
Tailless Bat	531	Timber Wolf.....	376
Tailless Hares.....	351	Tiny Mouse.....	136, 176
Talamanca Rice Rat.....	183	<i>Titi</i>	534
Tamaulipas Coyote.....	379	<i>Tlacuazin</i>	5
Tamaulipas Eyra.....	368	<i>Tlacuazin de Agua</i>	1
Taos Chickaree.....	84	<i>Tlacuazin Raton</i>	2
Tapir, Baird's.....	58	Tlalpam Field Mouse	164
Tapir, Dow's	58	Tlalpam Harvest Mouse.....	203
Tapirs.....	58	Tlalpam Shrew.....	464
Tatemales Pocket Gopher.....	283	Todos Santos Island Mouse.....	162
Tawny Field Mouse.....	143	Todos Santos Mouse.....	171
Tawny Raccoon Fox.....	398	Todos Santos Harvest Mouse	202
Taylor's Mouse.....	135	Todos Santos Wood Rat	214
Teapa Cotton Rat.....	178	Toltec Cotton Rat.....	177
Teapa Mouse	169	Toluca Squirrel	63
Teapa Rice Rat.....	185	Tomes' Long-eared Bat	508

	PAGE.		PAGE.
Tonalá Cotton Rat	177	Uinta Mountains Pocket Gopher ..	273
Tonila Rice Rat.....	184	Unalaska Lemming.....	260
Torrid Spiny Mouse	321	Unalaska Meadow Vole	242
Totontepec Field Mouse.....	169	Underwood's Bat.....	518
Tough-skinned Pocket Gopher	267	Ungava Arctic Fox.....	383
Townsend's Big-eared Bat.....	491	Ungava Red-backed Vole	232
Townsend's Chipmunk.....	86	Ursine Anteater.....	10
Townsend's Gray Fox.....	387	Uruapan Spiny Mouse	323
Townsend's Hare.....	363	Utah Chipmunk.....	86
Townsend's Meadow Vole	243	Utah Kangaroo Rat	296
Townsend's Mole.....	468	Utah Meadow Vole.....	239
Townsend's Pocket Gopher.....	275	Utah Prairie-dog.....	118
Townsend's Sperrophile	112	Utah Sperrophile.....	115
Trader Sperrophile	98		
Tree Squirrels	59	<i>Vacca de Agua</i>	14
Tres Marias Islands Bat.....	489, 517	Valley Meadow Vole.....	240
Tricky Coyote	377	Valparaiso Harvest Mouse.....	194
Trinity Mountains Sperrophile ..	107	Vampire Bats.....	503
Tropical Mole Mouse.....	133	Vancouver Island Chickaree	82
Tropical Pocket Gopher.....	270	Vancouver Island Shrew	449
Tropical Shrew.....	464	Vancouver Meadow Vole	243
Tropical Weasel.....	432	Variegated Squirrel.....	77
Tropical Wood Rat.....	220	<i>Venado</i>	45
Troublesome Field Mouse	149	Vera Cruz Cotton-tail.....	347
Troublesome Mouse.....	158	Vera Cruz Spiny Mouse	319
Troublesome Pocket Gopher.....	282	Villous Howler.....	333
Trowbridges's Shrew	452	Virginia Hare.....	360
True's Bat	479	Virginia Opossum	7
True's Lemming Vole.....	258	Virginian White-tailed Deer	43
True's Mole.....	469	Volcan de Chiriqui Cotton Rat....	180
True's Mouse	159	Volcan Toluca Harvest Mouse	199
True's Squirrel.....	73	Volcano of Irazu Mouse.....	204
True's Swamp Hare	341	Volcano of Irazu Shrew	464
True's White-tailed Deer.....	46	Vole, Acadian.....	235
Tucubaya Free-tailed Bat	499	Vole, Alaska Red-backed.....	230
Tulare Jack Rabbit.....	357	Vole, Albemarle Meadow.....	235
Tulare Kangaroo Rat.....	293	Vole, Anderson's Lemming	257
Tulare Mole Mouse	132	Vole, Arizona Meadow	238
Tule Meadow Vole.....	241	Vole, Aztec Meadow.....	237
<i>Tulomuco</i>	418	Vole, Baird's Meadow.....	250
Tumbala Rat.....	173	Vole, Barren Ground Meadow.....	237
Tumbala Rice Rat	187	Vole, Beach Meadow.....	238
Tundra Meadow Vole.....	241	Vole, Big-footed Meadow.....	251
Tundra Shrew	447	Vole, Bluegrass Meadow	249
Tunkas Brocket.....	51	Vole, California Meadow	240
Tunkas Rat	174	Vole, California Red-backed	233
Tuxtla Rat.....	173	Vole, Cantankerous Meadow.....	244
Two-toothed Sperm Whale	22	Vole, Carolina Red-backed	232
Two-toothed Whales.....	22	Vole, Cascade Mountain Meadow..	257
		Vole, Chapman's Lemming	257
Uinta Mountains Lynx.....	373	Vole, Coast Meadow	241, 244

	PAGE.		PAGE.
Vole, Colima or Dark Meadow	245	Vole, Nevada Meadow	240
Vole, Constable's	228	Vole, Newfoundland Meadow	238
Vole, Cooper's Lemming	255	Vole, Northern Meadow	247
Vole, Creeping Meadow	250	Vole, Northwestern Red-backed	229
Vole, Cunning Red-backed	233	Vole, Oaxaca Meadow	246
Vole, Dall's Lemming	258	Vole, Ochraceous Red-backed	230
Vole, Dawson's Red-backed	229	Vole, Olympic Meadow	244
Vole, Desert Meadow	241	Vole, Olympic Mountain	226
Vole, Drummond's Meadow	237	Vole, Orca Red-backed	230
Vole, Dusky Red-backed	232	Vole, Oregon Meadow	250
Vole, Dutcher's Meadow	240	Vole, Pallid Meadow	247
Vole, Dwarf Meadow	239	Vole, Peale's Meadow	238
Vole, Florida Water	252	Vole, Pennsylvania Meadow	235
Vole, Forest	236	Vole, Penobscot Field	235
Vole, Fulvous-bellied	246	Vole, Pigmy Meadow	247
Vole, Gale's Red-backed	231	Vole, Pine Meadow	249
Vole, Gapper's Red-backed	230	Vole, Popof Island Meadow	242
Vole, Goss' Lemming	256	Vole, Prairie Meadow	248
Vole, Graham Mountain Meadow	245	Vole, Preble's	227, 258
Vole, Gray Meadow	239	Vole, Reddish Meadow	240
Vole, Gray Rock Meadow	247	Vole, Rhoads' Red-backed	230
Vole, Gray-tailed Meadow	239	Vole, Richardson's Meadow	251
Vole, Guadalupe Mountains Mea- dow	245	Vole, Rich-coloured Red-backed	231
Vole, Guatemalan Meadow	252	Vole, Rock Meadow	246
Vole, Gull Island Meadow	238	Vole, Rufous Red-backed	229
Vole, Hall Island Meadow	243	Vole, St. Matthew Island Meadow	243
Vole, Hayden's Meadow	248	Vole, Sawatch Meadow	236
Vole, Idaho Red-backed	232	Vole, Secretive	227
Vole, Innuvit Lemming	256	Vole, Short-tailed Meadow	247
Vole, Innuvit Meadow	243	Vole, Short-tailed Red-backed	231
Vole, Intermediate	226	Vole, Silly Lemming	256
Vole, Jalapa Meadow	250	Vole, Sitka Meadow	242
Vole, Kadiak Meadow	242	Vole, Snow-loving Red-backed	233
Vole, Labrador Lemming	256	Vole, Stone's Lemming	255
Vole, Large Labrador Meadow	237	Vole, Stupid	227
Vole, Least Upland Meadow	248	Vole, Townsend's Meadow	243
Vole, Little Labrador Meadow	236	Vole, True's Lemming	258
Vole, Long-tailed	228	Vole, Tule Meadow	241
Vole, Long-tailed Meadow	244	Vole, Tundra Meadow	241
Vole, Loring's Red-backed	231	Vole, Unalaska Meadow	242
Vole, Louisiana Meadow	248	Vole, Ungava Red-backed	232
Vole, MacFarlane's Meadow	241	Vole, Utah Meadow	239
Vole, Mackenzie's	228	Vole, Valley Meadow	240
Vole, Mexican Meadow	245	Vole, Vancouver Meadow	243
Vole, Mogollon Mountains Meadow	246	Vole, Western Red-backed	233
Vole, Mole-like Meadow	249	Vole, White Mountains Lemming	256
Vole, Mt. Mazama Red-backed	232	Vole, White-footed	228
Vole, Mount Zempoaltepec Meadow	251	Vole, Woodland Meadow	249
Vole, Mountain	227	Vole, Wrangel Red-backed	229
Vole, Mountain Meadow	245	Vole, Wrangel's Lemming	257
		Vole, Yakutat Meadow	242

	PAGE.		PAGE.
Vole, Yellow-cheeked Meadow . . .	246	Weasel, Northern	429
Voies	129, 226	Weasel, Oregon	431
Voies, Lemming	255	Weasel, Pugnacious	428
Voies, Meadow	233	Weasel, Richardson's	426
Voies, Red-backed	229	Weasel, Richly-colored	430
		Weasel, Rio Grande Bridled	432
Wagner's Field Mouse	149	Weasel, Small-eared	426
Wallawalla Jack Rabbit	356	Weasel, Southern	429
Walrus	439	Weasel, Streater's	427
Walrus, Atlantic	439	Weasel, Teapa	433
Walrus, Pacific	439	Weasel, Tropical	432
Wandering Bermuda Bat	482	Weasel, Washington	429
Wandering Jack Rabbit	353	Weasel, Yellow-cheeked	431
Wandering Pocket Gopher	282	Weasels	404, 423
Wandering Rice Rat	184	West Indian Seal	445
Wandering Shrew	448, 465	Western Bat	480
Wandering Swamp Hare	341	Western Bushy-tailed Wood Rat	221
Wapiti	42	Western Desert Cotton Rat	170
Wapiti, Pacific Coast	42	Western Desert Jack Rabbit	356
Ward's Musk Ox	56	Western Hare	366
Warner Mountain Jumping Mouse	326	Western Moles	
Warree	37	Western Pocket Gopher	279
Warring Shrew	465	Western Porcupine	332
Washington Hare	361	Western Red-backed Vole	233
Washington Weasel	429	Western Wolf	376
Water Opossum	1	Whale, Arctic Baleen	17
Waterhouse's Large-eared Bat	508	Whale, Arctic Right	16
Watling's Island Bat	503	Whale, Baird's Sperm	24
Watson's Bat	524	Whale, Beaked Sperm	22
Watson's Rat	174	Whale, Blue Finback	20
Weasel, Alaskan	426	Whale, Cachalot Sperm	21
Weasel, Alleghany Mountains	428	Whale, Common Killer	26
Weasel, Allen's	431	Whale, Cope's Baleen	17
Weasel, Allied	433	Whale, Cope's Finback	19, 24
Weasel, Arctic	428	Whale, Cope's Killer	26
Weasel, Arizona	431	Whale, Davidson's Finback	19
Weasel, Bridled	432	Whale, Fighting Baleen	18
Weasel, Chestnut-brown	430	Whale, Gill's Bottle-nosed	30
Weasel, Common	426	Whale, Gray's Baleen	17
Weasel, Decorated	431	Whale, Gray's Bottle-nosed	29
Weasel, Eskimo	428	Whale, Gray's Japanese Baleen	16
Weasel, Feeble	427	Whale, Grebnitzky's Two-toothed Sperm	23
Weasel, Florida	430	Whale, Ice Baleen	15
Weasel, Goldman's Bridled	432	Whale, Large-toothed Killer	27
Weasel, Haida	429	Whale, Lesson's Finback	20
Weasel, Kadiak Island	428	Whale, Mobile Baleen	18
Weasel, Long-tailed	430	Whale, Northern Finback	20
Weasel, Michoacan Bridled	432	Whale, Pacific Finback	19
Weasel, Mountain	430	Whale, Pigmy Sperm	22
Weasel, Mousing	428	Whale, Pilot or Ca'ing	27
Weasel, Murderous	429		

	PAGE.		PAGE.
Whale, Sharp-beaked Bottle-nosed.	29	Wind River Mountains Chickaree..	81
Whale, Sharp-nosed Finback.....	19	Wolf, Arctic.....	376
Whale, Siebold's Baleen.....	16	Wolf, Autocrat Timber.....	374
Whale, Sowerby's Sperm.....	23	Wolf, Dusky.....	376
Whale, Stejneger's Sperm.....	23	Wolf, Mexican Timber.....	376
Whale, Two-toothed Sperm.....	22	Wolf, Timber.....	376
Whale, White.....	25	Wolf, Western.....	376
Whale, White-beaked Bottlenosed.	29	Wolverine.....	416
Whale, Yellow-bellied Finback....	20	Wolverine, Buff-colored.....	417
Whales, Baleen.....	15	Wolverine, Mount McKinley.....	417
Whales, Bottle-nosed.....	29	Wood Bison.....	57
Whales, Finback.....	18	Wood Rats.....	205
Whales, Killer.....	26	Wood Shrew.....	461
Whales, Sperm.....	20	Woodchuck.....	119
Whales, Two-toothed.....	22	Woodchuck, Canadian.....	120
White Bat.....	494	Woodchuck, Dakota.....	121
White Honduras Bat.....	528	Woodchuck, Engelhardt's.....	121
White Mountains Lemming Vole...	256	Woodchuck, Greedy.....	120
White Mountains Shrew.....	453	Woodchuck, Hoary.....	120
White Whale.....	25	Woodchuck, Labrador.....	120
White-backed Skunk.....	411	Woodchuck, Olympic Mountains..	121
White-banded Hutia.....	331	Woodchuck, Yellow-bellied.....	120
White-beaked Bottle-nosed Whale.	29	Woodchucks.....	119
White-bellied Mole Mouse.....	130	Woodland Jumping Mouse.....	328
White-bellied Mouse.....	138	Woodland Meadow Vole.....	249
White-bellied Rice Rat.....	182	Wooly Mouse.....	158
White-bellied Shrew.....	459	Working Pocket Gopher.....	273
White-eared Cotton Rat.....	179	Wortman's Spermophile.....	107
White-eared Gray Squirrel.....	79	Wrangel's Lemming Vole.....	257
White-eared Pocket Mouse.....	308	Wrangel Red-backed Vole.....	229
White-footed Deer Mouse.....	140	Wrinkled-face Bat.....	530
White-footed Raccoon Fox.....	399		
White-footed Vole.....	228	Xometla Mouse.....	162
White-lipped Harvest Mouse.....	202		
White-lipped Peccary.....	37	Yaki Mole Mouse.....	133
White-nosed Pocket Gopher.....	268	Yakima Spermophile.....	112
White-spot Field Mouse.....	142	Yakutat Bay Shrew.....	451
White-spotted Skunk.....	411, 413	Yakutat Meadow Vole.....	242
White-striped Bat.....	493, 525	Yapock.....	1
White-tailed Mouse.....	165	Yavapai Pocket Mouse.....	302
White-tailed Spermophile.....	97	Yellow Pocket Mouse.....	315
White-throated Brown Bat.....	485	Yellow Wood Rat.....	217
White-throated Capuchin.....	536	Yellow-bellied Finback Whale....	20
White-throated Wood Rat.....	217	Yellow-bellied Woodchuck.....	120
White-tipped Bushy-tailed Wood		Yellow-cheeked Meadow Vole.....	246
Rat.....	222	Yellow-cheeked Weasel.....	431
White-tipped Swimming Shrew...	458	Yellowish Lemming.....	259
White-toothed Pocket Gopher....	273	Yellowish-brown Bat.....	481
White-toothed Wood Rat.....	212	Yelm Prairie Pocket Gopher.....	277
Whitish Harvest Mouse.....	193	Yohaltun Field Mouse.....	147
Wild Cat.....	371	Ypanema Nose-leaf Bat.....	528

	PAGE.		PAGE.
Yucatan Bat	523	Yucatan Tree Porcupine	335
Yucatan Brown Bat	485	Yucatan White-tailed Deer	47
Yucatan Coati	403	Yukon Flying Squirrel	125
Yucatan Cotton-tail	345	Yukon Lemming	259
Yucatan Eyra	369		
Yucatan Free-tailed Bat	498	Zacatecas Pocket Mouse	311
Yucatan Mouse	163	Zacatecas Wood Rat	224
Yucatan Opossum	5	Zamora Mouse	166
Yucatan Peccary	36	<i>Zarro de Agua</i>	1
Yucatan Squirrel	75	Zygomata Pocket Gophers	271

g - F. C. Mueller