FOR THE PEOPLE
FOR EDUCATION
FOR SCIENCE

LIBRARY
OF
THE AMERICAN MUSEUM
OF
NATURAL HISTORY

TURNER ON BIRDS

London: C. J. CLAY AND SONS,
CAMBRIDGE UNIVERSITY PRESS WAREHOUSE,
AVE MARIA LANE.

Glasgow: 50, WELLINGTON STREET.

Lipsig: F. A. BROCKHAUS.
New York: THE MACMILLAN COMPANY.
Bombay and Calcutta: MACMILLAN AND CO., LTD.

TURNER ON BIRDS:

A SHORT AND SUCCINCT HISTORY

OF THE

PRINCIPAL BIRDS NOTICED BY PLINY AND ARISTOTLE,

59.82.102

FIRST PUBLISHED BY

DOCTOR WILLIAM TURNER,

1544.

EDITED,

WITH INTRODUCTION, TRANSLATION, NOTES, AND APPENDIX,

BY

A. H. EVANS, M.A.

CLARE COLLEGE, CAMBRIDGE.

CAMBRIDGE :

AT THE UNIVERSITY PRESS

1903

THE UNIVERSITY
OF CAMBRIDGE
LIBRARY

20-81081-Jan. 24
Cambridge:

PRINTED BY J. AND C. F. CLAY,
AT THE UNIVERSITY PRESS.

PREFACE.

IN translating this treatise it has been thought advisable to adhere as closely as possible to the original text of Turner, though in many places a more modern style of phraseology would doubtless have better pleased the ear of the non-scientific reader.

Not a few difficult points of interpretation have arisen in the course of the work, and the Editor welcomes this opportunity of expressing his gratitude to Professor Newton and to Mr R. D. Archer-Hind of Trinity College for the invaluable help that he has received from them in elucidating the hard passages and in revising the proofs. The Editor's thanks are also due to the Syndics of the University Press for undertaking the present publication.

9, HARVEY ROAD,
CAMBRIDGE.

July, 1903.

NOTE.

It may be remarked that the pages of Turner's work are not numbered in the original; but, for convenience of reference, the pagination is marked in the margin—the numbers being included in square brackets.

INTRODUCTION.

WILLIAM TURNER, author of the rare treatise here republished, was a native of Morpeth in Northumberland and is supposed to have been the son of a tanner of that town. By the aid of Thomas, the first Lord Wentworth, he was enabled to enter Pembroke Hall in the University of Cambridge, where he graduated B.A. and was elected a fellow of his College in 1530. At Pembroke he became acquainted with Ridley (who instructed him in Greek) and Latimer, two of the most earnest advocates of the Reformed doctrines, which he himself, both then and afterwards, strenuously embraced; but there is no need to dwell upon his theological views or the polemical works in which they were set forth.

While at Cambridge Turner was a zealous student of botany, and in 1538 published a *Libellus de re herbaria*. About two years later he left this University for Oxford, and soon after suffered imprisonment for preaching without a licence. On his release he quitted England, and travelled by way of the Netherlands and Germany to Italy, attending the botanical lectures of Luca Ghini at Bologna, where, or at Ferrara, he took the degree of M.D.

Thereafter he proceeded to Switzerland, forming a close friendship with the great naturalist Conrad Gesner of Zurich, Professor of Medicine and Philosophy in the School of that city, who held him in high esteem, and with whom he afterwards kept up a correspondence. He seems to have been at Basel in 1543, but early in 1544 he was at Cullen (Cologne), where he published not only the present work—dedicated

to Edward Prince of Wales (afterwards King Edward the Sixth)—but also edited the *Dialogus de Avibus* of his friend Gybertus Longolius of Utrecht, who died the preceding year. He pursued his botanical studies in several parts of Germany, as well as in the Netherlands, including East Friesland, for he became physician to the Count of Emden, and visited the islands of Juist and Norderney lying off the coast of that province.

On the death of King Henry the Eighth he returned to England, becoming chaplain and physician to Lord Protector the Duke of Somerset; but he lived at Kew, where he established a botanic garden. He was, moreover, incorporated M.D. of Oxford, and was appointed a prebendary of York. In 1550 the Privy Council sent letters for his election as Provost of Oriel College in Oxford, but the post had been already filled, and a similar disappointment awaited him in regard to the Presidency of Magdalen College in the same University. He then applied to Sir William Cecil for leave to return to Germany, but was soon after consoled by being appointed Dean of Wells, and, having in 1551 published the first part of his *New Herbal*, was during the next year ordained priest by his old friend Ridley, then Bishop of London.

On the accession of Queen Mary Turner had to vacate his deanery, and betook himself for safety once more to the Continent, visiting Rome and several places in Germany and Switzerland. When Queen Elizabeth ascended the throne he returned to England, and recovered his deanery, to which was attached the rectory of Wedmore in Somersetshire; but in 1564 he was suspended for nonconformity and seems to have come to live in London. In 1557 he had addressed a letter on English Fishes to Gesner, which was included in that naturalist's *Historia Animalium*; and in 1562 he published the second part of his *Herbal*, which he dedicated to Lord Wentworth, the son of his original benefactor. On the 7th of July 1568 Turner died at his house in Crutched Friars in the City of London, and was buried in the church of St Olave, Hart Street.

Turner married Jane, daughter of George Ander, Alderman of Cambridge, by whom he had issue Winifred, Peter and Elizabeth.

It must be understood that, his scientific work apart, nearly the whole of Turner's life was spent in religious controversy, and he published a considerable number of polemical works, the titles of which may be seen in the bibliography appended to the excellent 'Life' prefixed to Mr Jackson's facsimile reprint of the *Libellus de re herbaria*¹, whence all the particulars above given are taken. Other lists of Turner's works may be found in Cooper's *Athenae Cantabrigienses* (I. pp. 257—259) and the *Dictionary of National Biography* (LVII. pp. 365, 366).

Turner's object in writing the present treatise is fully set forth in his 'Epistola Nuncupatoria' prefixed to it. While attempting to determine the principal kinds of birds named by Aristotle and Pliny, he has added notes from his own experience on some species which had come under his observation, and in so doing he has produced the first book on Birds which treats them in anything like a modern scientific spirit and not from the medical point of view adopted by nearly all his predecessors; nor is it too much to say that almost every page bears witness to a personal knowledge of the subject, which would be distinctly creditable even to a modern ornithologist.

This knowledge is especially evident in his account of the habits of the Hobby (p. 19), Hen-Harrier (p. 19), Water-Ousel (p. 23), Moor-Buzzard (p. 33), Osprey (p. 37), Godwit (p. 45), Wheatear (p. 53), Sandpiper (p. 57), Fieldfare (p. 59), Cuckoo (p. 69), Black-headed Gull (p. 77), Black Tern (p. 79), Swallows (p. 101), Cormorant (p. 111), Shrike (p. 119), Redbreast and Redstart (p. 157); while his keen eye for distinctions is shown in his descriptions of the Black Cock and Grey Hen (p. 43), Godwit (p. 45), Tree-Creeper (p. 53),

¹ *Libellus de re herbaria novus*, by William Turner, originally published in 1538, reprinted in facsimile, with notes, modern names, and a Life of the Author, by Benjamin Daydon Jackson, F.L.S. *Privately Printed*. London: 1877.

Doves (p. 59), Lapwing (p. 77), Nutcracker (p. 95), Reed-Bunting (p. 103), Kites (p. 117), Bullfinch (p. 161) and others. He is most careful to tell us whether he observed the various species in England or abroad and their comparative abundance, and to note the breeding of the rarer species, such as the Spoonbill (p. 151), and Crane (p. 97) within our islands—that of the Crane being of special interest; the whole account of the Cuckoo (p. 69) is also most noticeable, as is that of the curious Walt-rapp (p. 95) of which Gesner writes as follows:

GESNER *De Corvo Sylvatico*¹ (p. 337).

AUIS, cuius hic effigies habetur, à nostris nominatur uulgo ein *Waldrapp*; id est coruus sylvaticus...Sunt qui phalacrocoracem hanc auem interpretentur, quoniam & magnitudine & colore ferè coruum refert: & caluescit, ut uidi, cum adultior est. Turnerus Aristotelis coruum aquaticum & Plinij phalacrocoracem, & coruum sylvaticum nostrum auem unam esse arbitratur, tertium genus graculi. Coruus sylvaticus Helvetiorum, inquit, auis est corpore longo et ciconia paulò minore, cruribus breuibus, sed crassis: rostro rutilo, parum adunco (curuo) & sex pollices longo: alba in capite macula, & ea nuda, si bene memini....Sic ille.

Doubtless Turner's work is not free from errors, as in the case of the very old story of the breeding of the Bernicle-Goose (which, however, he was most loth to credit even when assured of its truth by an Irish Divine), in his confounding of the *Onocrotalus* with the *Ardea stellaris* and the Cornish with the Alpine Chough; yet these are but small blots on a very excellent treatise, which compares most favourably with other writings of his time.

It is quite evident from various passages that Turner was acquainted with Aristotle's works in the original Greek, and especially with his *History of Animals*; but he preferred quoting that author from the Latin translation of Theodorus

¹ Conradi Gesneri Tigurini medici & Philosophiæ professoris in Schola Tigurina Historiæ Animalium Liber III. qui est de Auium natura. Tiguri apud Christoph. Froschovervm, Anno M.D.LV.

Gaza of Thessalonica, the most celebrated Scholar of his day, who, fleeing from the sack of Constantinople, played a conspicuous part in the rise of the "New Learning," and after a course of teaching in Rome, entered successively the service of the Popes Nicholas the Fifth and Sixtus the Fourth, eventually dying in poverty in Lucania about 1484.

Exact transcription of a text was considered by no means necessary in those days: consequently we find many observations and explanations inserted in the text of Aristotle and Pliny, which had no place in the original¹.

Besides referring to Gesner, Turner mentions other learned men by name and occasionally quotes from their works; while his pages also inform us of many places that he visited.

The following excerpts from Gesner not only give instances of correspondence between him and Turner, but also shew that the former was accustomed to correct the latter from his wider knowledge of Ornithology.

De Branta vel Bernicla... (p. 107).

Idē [Turnerus] post librum suum de avibus publicatum, in epistola ad me data, Berniclas siue Brantas (inquit) ex putridis nauis malis fungorum more nasci, minimè fabulosum esse doctorum & honestorum uirorum oculata fides mihi persuasit. Branta anserem palustrem ualde refert: his tamen notis ab eo differt. Branta breuior est, à collo quod rubescit nonnihil, ad medium usq̃ uentrem, qui candicat, nigra est. anserum more segetes populatur. In Vuallia (quæ pars est Angliæ) in Hibernia & Scotia aues istæ adhuc rudes & implumes in littore, sed non sine forma certa & propria auis passim inueniuntur. Et rursus, Præter brantam aut berniclam est alia auis, quæ originem suam arbori refert acceptam. Arbores sunt in Scotia ad littus maris crescentes, è quibus prodeunt ueluti fungi parui, primum informes, postea paulatim integram auis formam acquirunt, perfectæ tandem magnitudinis illæ, rostro aliquantisper pendent, paulò post in aquam decidunt, & tum demum uiuunt. Hoc tot tantæq̃ integritatis uiri affirmauerunt ut credere audeam, & aliis credere suadeã. Hæc ille. Eliota Anglus &c....

¹ The precise references to Aristotle and Pliny are now supplied, from the texts of Aubert and Wimmer, and Sillig respectively.

De Vulpansere (p. 156).

Chenalopex (inquit Turnerus Anglus) ab ansere & uulpe nomen habet...

Et rursus in epistola ad me, Vulpanserem Angli vocât a Bergander, nidulatur in cuniculorũ foueis more uulpium, anate maior, minor ansere, alis ruffis. Eliota Anglus...

De Pygargo (p. 199).

Pygargum (inquit Turnerus) literatores quidam ineptè trappum à Germanis dictum (tardam, uel bistardam) interpretantur. Sed pygargus Anglorum lingua, nisi fallar, ernam vocatur, an erne. Ego ernam audio dici genus aquilę quod apud Frisios ad Oceanum Germanicum per hyemem degat, colore nigro, quod cornices quædam ut ex escarum eius reliquijs uictitent sequantur. Pygargus est forte quam Anglicè dicimus ringetayle, Eliota. Sed Turnerus ringtalum Anglis dictum ab albo circulo caudam circumeunte, buteone minore, subbuteonem Aristotelis esse suspicatur. Quod si minor est buteone, non poterit esse pygargus.

De Caprimulgo (p. 235).

[Having quoted Turner] Idem postea in litteris ad me missis, caprimulgum se uidisse scribit prope Bonnam (Germaniæ ciuitatem ad ripam Rheni, supra Coloniam) ubi à uulgo appellatur *Naghtranen*, id est coruus nocturnus. Nos auis illius quæ Argëtinę vocatur *Nachtram*, corrupto forsân nomine, alibi *Nachtrap*, effigiem infrâ ponemus cum historia nycticoracis.

De Carduele (p. 235).

...Auis Aristoteli thraupis, *θραυπῖς*, dicta à Gaza carduelis conuertitur: quod & Hermolao probatur. Turnero quidem non assenserim, qui chloridem nostram (*Grünling* uulgo vocant) thraupin esse coniicit.

De Rala terrestri (pp. 481—482).

...Perdix rustica vel rusticula Plinij (inquit Turnerus in epistola ad me) ab Anglis vocatur rala. Est autem rala duplex, altera cibum è ripis fluminum petit, altera degit in ericeto in locis sylvestribus. Aquaticam illam Colonia diu alui, & male uolare deprehendi, & egregiè pugnacem. Rostrum & crura erāt rubra, plumę multis maculis resperse.

Montana verò illa & syluestris crura habet multò breuiora aquatili, & plumas undiq̃ magis cinereas, sed rubra interim crura habet & rostrum. auis utraq̃ apud nos regium epulum (real Itali regium vocant, Galli royal, & forte hinc ductum est ralæ vocabulū. à colore crurum forte erythropus fuerit) vocatur. Hæc ille.

De Cinclo (p. 592).

Turnerus cinculum auem Anglicè interpretatur a water swallow, (quasi dicas hirundinem aquaticam,) Germanicè *ein Steinbeisser*, (sed nostri aliam auē, coccothrausten nostrum, *Steinbeisser* appellant.) Auicula (inquit) quam ego cinculum esse puto, galerita paulo maior est, colore in tergo nigro, uentre albo, tibiis longis, & rostro neutiquam breui. Vêre circa ripas fluminum ualde clamosa est & querula, breues & crebros facit uolatus. ¶ Huic Turneri descriptioni cognata uideri potest merula aquatica nostra: magis uerò illa, quam circa Argentoratum *Lyssklicker* appellant: quam non similem modo Turneri cinclo, sed prorsus eandem esse conijcio, cuius figuram sequens pagina cōtinet.

De Motacilla quam nostri albam cognominant (p. 593).

Turnerus in libro de Auibus Cnipológon Aristotelis (id est culicilegā interprete Gaza) hãc auem esse putat...[quotation]....Sed postea in epistola ad me, Culicilegam Aristotelis (inquit) in terra Bergensi uidi, tota cinerei ferè coloris est, & speciem habet pici Martij, illa uerò quam culicilegam esse putabã, est uariola nisi fallor.

De Nycticorace (pp. 602—604).

TURNERUS in litteris ad me missis caprimulgum auem se uidisse scribit prope Bonnam (Germaniæ ciuitatem ad ripam Rheni, supra Coloniam,) ubi à uulgo appellatur *Naghtrauen*, id est coruus nocturnus. Nos in præcedente pagina effigiem adiecimus auis quæ circa Argentoratum, ut audio *Nachtram*, alibi *Nachtrab* nominatur. quæ tamen neq̃ caprimulgus neq̃ nycticorax mihi uidetur. [The figure is of the Night-Heron, unmistakably.]

De Onocratalo (pp. 607—608).

Onocrotalus Machliniensis, quæ *Vogelhain* à Brabantis uocatur, quinquaginta annis, ut ipsi ferunt, Machliniæ uixit, cygno maior est. pennę foris albicant, in fundo uerò rubrum

quiddam ostendunt. collum duas spithamas longum est, aut paulò longius, rostrum, quod rubrum habet, dodrantali longitudine est & quatuor uncias longius, & in fine hamì propemodum more incuruum & uersus finem latius latiusq, proturbatur. crura anserinis similia, breuia, nimirum pro magnitudine tanti corporis: in pectore magnum habet ueluti sacculum protuberantem. Alis est longissimis, & ipsis in summa extremitate nigris, Guil. Turnerus in epistola ad me....

Bononiæ uisus est mihi onocrotalus (uerba sunt ex epistola Angli cuiusdam amici ad me) plumis cinereis tectus, cygno maior, palmipes, capite mergi, rostro quatuor palmas ferè longo, & in fine adunco, collo deplumi, amplissimo, ut anatem deuorare posset. Captam aiebāt in lacu Benaco....

Solis uictitat piscibus, & bis tantum anno bibit. Turnerus...Pisces præsertim anguillas audissimè uorat botaurus auis, Turnerus. idem amicus quidam de onocrotalo ad me scripsit. ego onocrotalum quidem hoc facere non dubito: de botauro dubitari potest, præsertim cum multi etiã non indocti cum onocrotalo eum confundant.

De Perdice (p. 644).

...Quæ Aristoteles & Ouidius de perdice scribunt, omnia nostræ pernici uulgarì conueniunt, nempe uolandi nidulandi, ratio, astutia, circa prolem sollicitudo, corporis grauitas, & uocis stridor, à quo etiam nomen accepisse uidetur, Turnerus in epistola ad nos.

The 'Avium præcipuarum...Historia' was reprinted by Dr George Thackeray, Provost of King's College, Cambridge, in 1823; but the reprint is as rare as, if not rarer than, the original. Two copies are in the Library of King's College.

The following is a list of the Birds determined by Turner.

ALAUDIDÆ. <i>Alauda arvensis</i> .	ALCEDINIDÆ. <i>Alcedo ispida</i> .
Lerk or Laverock. German	Kynges fissher. G. <i>Eisssvogel</i> ,
<i>Lerch</i> , p. 80.	pp. 18—22.
<i>A. arborea</i> . Wodlerck, p. 80.	[Turner recognised two kinds
<i>A. sp.?</i> Wilde Lerc or Heth	of Kingfishers described by
Lerk. G. <i>Heid Lerch</i> , p. 80.	Aristotle and Pliny, but
<i>Galerita cristata</i> . [No English	does not state what they
name.] G. <i>Copera</i> , p. 80.	are].

- ANATIDÆ. *Anas boscas*. Duck, pp. 22, 48.
Anser (2 species.) Gose. G. Ganss, p. 22.
Bernicla leucopsis. Brant or Bernicle Gose, p. 26.
Cygnus olor. Swan. G. Swän, p. 120.
Mareca penelope. Wigene, p. 48.
Nyroca ferina. Pochard, p. 48.
Querquedula crecca. Tele, p. 48.
Tadorna cornuta. Bergander, p. 24.
- ARDEIDÆ. *Ardea* sp.? (white). Cryel or Dwarf Heron, p. 38.
A. cinerea. Heron. G. Reyger, p. 36.
Botaurus stellaris. Bittour, Buttor, Buttour, or Myre Dromble. G. Pittour, Rosdom, pp. 38, 40, 122.
- CAPRIMULGIDÆ. *Caprimulgus europæus*. [No English name], p. 48.
- CERTHIIDÆ. *Certhia familiaris*. Creper, p. 52.
- CHARADRIIDÆ. *Charadrius plumbealis*. Pluver. G. Pulver, p. 132.
Vanellus vulgaris. Lapwing. G. Kywit, pp. 76, 174.
- CICONIIDÆ. *Ciconia alba*. Stork. G. Storck, Sax. Ebeher, p. 54.
- CINCLIDÆ. *Cinclus aquaticus*. Water-Craw, p. 22.
- COLUMBIDÆ. *Columba* sp.? Dove. G. Taube, Sax. Duve, p. 59. [Venice Dove, p. 62.]
C. ænas. Stocdove. G. Holtztaube, p. 60.
C. palumbus. Coushot or Ringed Dove. G. Ringel Taube, p. 60.
Turtur communis. Turtel Duve, p. 60.
- CORVIDÆ. *Corvus corax*. Raven. G. Rabe, p. 64.
C. cornix. Winter Crow, p. 64.
C. corone. Crow. G. Krae, Kraeg, p. 64.
C. frugilegus. [No English name], p. 64.
C. monedula. Caddo, Chogh, Ka. G. Döl, Sax. Älke, p. 92.
Garrulus glandarius. Jay. G. Mercolphus, p. 144.
Nucifraga caryocatactes. [No English name.] G. Noubrecher, p. 94.
Pica rustica. Py, Piot. G. Elster, Atzel, pp. 142, 144.
Pyrrhocorax graculus. Cornish Choghe. G. Bergdöl, p. 90. [Confounded with *P. alpinus*.]
- CUCULIDÆ. *Cuculus canorus*. Cuckow, or Gouke. G. Kukkuck, p. 66.
- CYPSELIDÆ. *Cypselus apus*. Chirche Martnette. Rok Martinette. G. Kirch Swalbe, pp. 100, 102.
C. melba. Great Swallow. G. Geyr Swalbe, p. 102.
- FALCONIDÆ. *Accipiter nisus*. [No English name], p. 66.
Aquila sp.? Right Egle. G. Edel Ärn, p. 36.
Astur palumbarius [?] Sparhauc. G. Sperwer, p. 18.
Buteo vulgaris. Bushard, p. 16.
Circus æruginosus. Balbushard, p. 32.
C. cyaneus. Hen-Harroer, Ringtale, p. 18. [Turner calls the male Hen-Harroer, the female Ringtale, erroneously considering them two species.]
Falco æsalon. Merlin. G. Smerl, p. 16.

FALCONIDÆ. *F. subbuteo*. Hobby,
p. 18.

Gypaëtus barbatus [?]. [No English name], p. 128.

Haliaëtus albicilla. Egle or Erne. G. Ärn, Adler, p. 30.

Milvus ater. [No English name], p. 116.

M. iclinus. Glede, Puttok, or Kyte. G. Weye, p. 116.

Tinnunculus alaudarius. Kastrel, Kistrel, or Steingall, p. 166.

FRINGILLIDÆ. *Carduelis elegans*. Goldfinche. G. Distelfinck, Stigelitz, pp. 40, 50.

C. spinus. Siskin. G. Zeysich, Engelchen, p. 108 (cf. p. 50).

Emberiza citrinella. Yellowham, Yowlyng. G. Geelgorst, p. 106.

E. miliaria. Bunting. G. Gersthammer, pp. 134, 158.

E. schæniclus. Rede Sparrow. G. Reydt Müss, pp. 102, 134.

Fringilla cælebs. Chaffinche, Sheld-appel, Spink. G. Büchfink, p. 72.

F. montifringilla. Bramlyng. G. Rowerl, p. 72.

Ligurinus chloris. Grenefinche. G. Kirsfincke, pp. 104, 106.

Linota cannabina? Linot. G. Flasfinc, pp. 50, 158.

Passer domesticus. Sparrow. G. Mische, Spätz, Lüningk, Sax. Sperlingk, p. 132.

Pyrhula europæa. Bulfinche. G. Blödtfinck, p. 160.

Serinus canarius. Canary Bird, p. 108.

GRUIDÆ. *Grus communis*. Crane. G. Krän, Kränich, pp. 94, 96.

HIRUNDINIDÆ. *Cotile riparia*.

Bank Martnet. G. Über Swalbe, Speiven, p. 102.

Hirundo rustica. Swallowe. G. Schwalb, Sax. Swale, pp. 96, 100, 102.

IBIDIDÆ. *Comatibis cremita*. [Red-cheeked Ibis]. G. Waltrap, pp. 92, 94.

LANIIDÆ. *Lanius excubitor*. Schric, Shrike, or Nyn Murder. G. Nuin Mürder, Neun Mürder, pp. 116, 118, 168.

Lanius minor? [No English name], p. 168.

LARIDÆ. *Hydrochelidon nigra*. Stern, p. 78.

Larus sp.? [Grey Gull.] Se Cob or See Gell, p. 78.

L. sp.? [White Gull.] White Semaw. Se Cob or Seegell. G. Wyss mewwe, pp. 74, 78.

L. ridibundus. White Semaw with a blak cop, pp. 74, 76.

MEROPIDÆ. *Merops apiaster*. [No English name], p. 112.

MOTACILLIDÆ. *Anthus pratensis*. Titlyng, p. 68.

Motacilla lugubris or *M. alba*. Wagtale. G. Wasser Steltz, Quikstertz, p. 64.

ORIOOLIDÆ. *Oriolus galbula*. Witwol. G. Witwol, Weidwail, Kerscurife, pp. 148, 172, 174.

OTIDIDÆ. *Otis tarda*. Bistard or Bustard. G. Träp or Trap Ganss, pp. 130, 166.

PANDIONIDÆ. *Pandion haliaëtus*. Osprey. G. Vishärn, pp. 34, 36.

PARIDÆ. *Parus caruleus*. Non. p. 132.

- PARIDÆ. *P. major*. Great Titmouse, or Great Oxei. G. *Kölmeyse*, p. 130.
P. palustris or *P. ater*. Less Titmouse. G. *Meelmeyse*, p. 130.
- PHALACROCORACIDÆ. *Phalacrocorax carbo*. Cormorant. G. *Dücher*, p. 110.
P. graculus? Douker (pt), Loun, *Dücher* (pt.), p. 176.
- PHASIANIDÆ. *Attagen* [possibly *Bonasa sylvestris*, the Hazel Grouse], pp. 42, 44.
Gallus ferrugineus (domesticus). Cok, Hen. G. *Hän, Hen, Sax. Hön*, p. 82.
Numida meleagris. [No English name?] Kok of Inde? pp. 82, 86, 140.
Pavo cristatus. Pecok. G. *Pffaw, Sax. Pagelün*, p. 136.
Perdix cinerea. Pertridge. G. *Velt hön, Raphön*, p. 138.
Phasianus colchicus. Phesan. G. *Fasant, Fasian*, p. 140.
- PHENIX. [No English name], p. 140.
- PICIDÆ. *Dendrocopus major*? Specht, or Wodspecht. G. *Elsterspecht*, pp. 146, 148.
Gecinus viridis. Hewhole, Huhol, Raynbird? G. *Grünspecht*, pp. 88, 112, 114, 146, 148.
Iynx torquilla. [No English name], pp. 146, 148.
Picus martius. [No English name]. G. *Craspecht*, p. 148.
- PLATALEIDÆ. *Platalea leucorodia*. Shovelard. G. *Lefler, Löffel Ganss*, pp. 38, 150.
- PODICIPEDIDÆ. *Podiceps minor*, Douker (pt), *Dücher* (pt), p. 176.
- PROCELLARIIDÆ. *Puffinus* sp.? Bird of Diomedæ, p. 70.
- PSITTACIDÆ. Parrot. Popinjay. G. *Papegay*, p. 150.
- RALLIDÆ. *Crex pratensis*. Daker Hen, Rale (?) G. *Schryk, Scrica*, pp. 70, 128, 140.
Fulica nigra. Cout, pp. 32, 76.
Gallinula chloropus. Mot Hen or Water Hen. G. *Wasser Hen*, p. 170.
Porphyrio caelestis. [No English name], p. 152.
- SCOLOPACIDÆ. *Actitis hypoleuca*. Water Swallow. G. *Steynbisser*, pp. 54, 56.
Limosa belgica. Godwitt or Fedoa, p. 44.
Scolopax rusticula. Wodecok. G. *Holtz Sneppf*, pp. 42, 86.
Totanus calidris. Redshanc, p. 102.
- SITTIDÆ. *Sitta cæsia*. Nut-jobber. G. *Nushækker, Meyspecht*, p. 162.
- STRIGIDÆ. *Asio otus*. Horn Owl. G. *Ranseul, Schleier Eul*, p. 130.
Bubo ignavus. Lyke Foule. G. *Schuffauss, Schüffel, Kautz*, p. 46.
Strix stridula (?) Owl, Howlet. G. *Eul, Sax. Üle*, p. 120.
- STRUTHIONIDÆ. *Struthio camelus*. Oistris. G. *Strauss*, p. 164.
- STURNIDÆ. *Sturnus vulgaris*. Sterlyng. G. *Stär, Stör*, p. 164.
- SULIDÆ. *Sula bassana*. Solend Guse, p. 28.
- SYLVIIDÆ. *Accentor modularis* (?) Hedge-sparrow, or Dike Smouler. G. *Grassmusch, Koelmusch*, p. 136.
Daulias luscinia. Nyghtyngall. G. *Nachtgäl*, p. 108.

SYLVIIDÆ. *Erithacus rubecula*.
Robin Redbreaste. G. *Rötbrust*,
Rötkelchen, p. 154.

Pratincola rubicola. Stonchatter
or Mortetter. G. *Klein Brach-*
vogelchen, p. 158.

Regulus cristatus (?). [No English
name.] G. *Gold Hendlin*, pp.
154, 168.

Ruticilla phaniceus. Rede Tale.
G. *Röstertz*, p. 154.

Saxicola ananthe. Arlyng, Clot-
burd, Smatche or Steinchek.
G. *Brechvögel*, p. 52.

Sylvia atricapilla (?). [No Eng-
lish name.] G. *Grasmuklen*,
p. 44.

S. rufa? Lingett. G. *Graes-*
musch, *Grassmusch*, p. 111,
[cf. p. 136].

TETRAONIDÆ. *Coturnix commu-*
nis. Quale. G. *Wachtel*, p. 62.
Lagopus mutus. [No English
name], p. 104.

L. scoticus. (?) Morhen, p. 86.
Tetrao tetrix. [No English
name], p. 42.

TROGLODYTIDÆ. *Troglodytes par-*
vulus. Wren. G. *Kuningsgen*,
Zaunküningk, p. 152.

TURDIDÆ. *Turdus iliacus*, *mu-*
sicus, *viscivorus*. Thrusche,
Thrushe, Throssel, Mavis, or
Wynghrushe. G. *Drossel*,
Durstel, *Weingaerdsvoegel*, pp.
170, 172.

T. merula. Blak Osel or Blak-
byrd. G. *Merl*, *Amsel*, p. 114.

T. pilaris. Feldfare or Feldefare.
G. *Krammesvögel*, *Wachholter-*
vögel, pp. 58, 170. [Confounded
with Mistletoe Thrush?]

UPUPIDÆ. *Upupa epops*. Howpe.
G. *Houp*, *Widhopff*, p. 174.

VULTURIDÆ. *Vultur* sp. Geir.
G. *Geyr*, p. 176.

A V I V M
PRAECIPV
A R V M, Q V A R V M

APVD PLINIVM ET ARI-
stotelem mentio est, brevis &
succincta historia.

*Ex optimis quibusque scripto-
ribus contexta, scholio illu-
strata & aucta.*

*Adiectis nominibus Græcis, Germanicis &
Britannicis.*

*Per Dn. Guilielmum Turnerum, artium & Me-
dicinæ doctorem.*

*Coloniæ excudebat Ioan. Gymnicus,
Anno M. D. XLIIII.*

DE HISTORIA AVIUM.

[p. 3] Illustrissimo Valliæ principi, Eduuardo
filio hæredi, serenissimi & potentis-
simi Henrici VIII. regis Angliæ,
Franciæ, & Hiberniæ, Guilielmus
Turnerus S. P. D.

PRUDENS admodum, & si quid ego intel-
ligo (illustrissime princeps) necessaria im-
primis regij prophetæ fuit admonitio, qua reges,
principes & iudices terræ, ut intelligerent, &
eruditionem consequerentur, admonuit. Nam
ut summus ille rerū architectus Deus optimus
maximusq; caput super omnes reliquas corpo-
ris partes in homine, qui perfectissimè in se
[p. 4] Reipublicæ simulachrum gerit, collocavit, &
omnes quinque sensus simul in eo reposuit,
ut pro reliquis omnibus mēbris (quibus solus
tactus est concessus) uideret, audiret, gustaret &
odoret, & eorum salutem consuleret : ita principem
Reipublicę, corpori ex multis membris conflato,
ueluti caput præfecit, ut prudentia, eruditione,
& sensibus suis non tam exterioribus, quā inte-
rioribus, totius Reipublicæ commodis & salutem
prospiceret. In pedibus uisum, in tibijs audi-
tum, in manibus olfactum, in brachijs odoratum
nemo requirit: sed hæc omnia in capite requi-
runtur.

To the most illustrious Prince of Wales,
Edward, son and heir of the most serene
and mighty Henry VIII, King of Eng-
land, France and Ireland, William Turner
wishes long life and health.

EXCEEDING wise, and if I understand aright,
necessary above all things, most illustrious
Prince, was the warning of the royal prophet, in which
he admonished kings, princes, and judges of the earth
that they should have understanding and seek learn-
ing. For, as that architect supreme of the universe,
God most good and great, placed the head above all
the remaining parts of the body in man, who in him-
self shews forth the image of a most perfect State,
and stored up in it all the five senses at once, that it
should see, hear, taste, and smell for all the remaining
members (to which touch alone has been allowed),
and should consult for their well-being; so he hath
set the Prince, as it were a head, over the State,
a body welded together of many members, that he
should provide for the advantage and well-being of
the whole State by his wisdom, learning and senses,
not so much external as internal. No one demands
sight in the feet, hearing in the legs, smell¹ in the
hands, or smell in the arms; but all these things are
necessary in the head.

¹ This should probably be "taste" (gustum).

Quum igitur tot sensus in capite uni tantū corpori præfecto requirantur : quot sensus, quantum sapientiæ & eruditionis ab eo capite exi-
 [p. 5] guntur, cui plus quàm trecentorum milium corporum præfectura committitur? Quòd si quis forsan respondeat, non in principe, sed in ijs solis, qui illi à consilijs sunt, eruditionem & prudentiam requiri : hunc ego dignum cenferē, qui pro tali responso, omnibus sensibus, excepto tactu, orbatus in media sylua uepribus & spinis densa, caueis & fossis formidabili, quatuor ducibus comitatus statueretur, nobis dicturus, nū proprijs malit uti sensibus an alienis? & num tutius illi sit, ducum suorum incertorum sensibus, an proprijs duci? & qua ratione cæcus & surdus odoratu & gustu destitutus ipse, cæci'ne an uidētes sui sint duces, dignoscere possit?

In consiliarijs summam prudentiam & eruditionem non uulgarem requiri, non diffiteor :
 [p. 6] uerū non in ijs solis, nam si illi, qui principi sunt à consilijs, ad tempus bene consulant, & postea in ipsius perniciem malè suadeant, ut Absaloni Achitofelem fecisse legimus : quomodo perniciosum illorum consiliū ipse subodorabitur & depræhendet, nisi eruditione & prudentia consiliarios suos aut superet, aut faltem æquet? Quare nō in consiliarijs tantū, sed in principe ipso eruditio & sapiētia requiruntur. Non defunt, qui fati esse principi existimant, quo cæteris mortalibus præstet, si regio uestitu, diuitijs, copijs, scitè pulsando testudinem, & tela dextrè uibrando, subditis suis præluceat : uerū fortis-
 [p. 7] simi quiq; & sapientissimi reges longè diuersum

Inasmuch therefore as so many senses are requisite in the head, which is set over one body alone, how many senses and what a wealth of wisdom and learning are demanded from that head, to whom more than three hundred thousand bodies are given in charge? But if any should chance to answer that learning and wisdom are needed not in the Prince, but only in those who are his councillors, I should consider it fitting that he for such a reply should be set, accompanied by four guides, in the midst of a wood tangled with briers and thorns, and dangerous with its pits and ditches, deprived of all his senses, except that of touch, and should tell us whether he preferred to use his own senses or those of others: or whether it would be safer for him to be led by the senses of his doubting guides or by his own; and in what way he, being blind and deaf, and destitute of smell and taste, could determine whether his guides were blind or able to see.

I fail not to confess that the highest wisdom, and learning of no common sort, are requisite in councillors, but not in them alone; for if they who are the advisers of the Prince, give good counsel for the time, and afterwards prompt him ill to his destruction, as we read that Achitofel did in the case of Absalom, how shall he smell out and detect their fatal advice, unless he either excels or at least equals his councillors in learning and wisdom? Wherefore not only in councillors but in the Prince himself are learning and wisdom requisite. There are not wanting those who think it enough for a Prince, as matters in which he should surpass other mortals, if he outshines his subjects in royal garb, in riches, in resources, in cunningly striking the lyre, and in skilfully throwing the spear; but all the bravest and wisest kings have

fenferunt. Nã Mithridates rex Põti & Bithynië, se regnorum fuorum caput esse intelligēs, et tot corporibus, quot præerat, unicã uernaculã suam linguã minimè sufficere, uiginti duas linguas gētium, quas sub ditione sua habuit, ita perfectè didicit & percalluit, ut uiginti illarum gentium uiris sine interprete promptè responderit, & sua cuique lingua non fecus atque gentilis fuisset, locutus fuerit. Idē rerum abditas naturas ita peruestigauit, & in re medica ita foeliciter fuit uersatus, ut aduersus lethalia uenena antidotum, quod hodie etiamnum ab eo nomen fortitum, Mithridatium appellatur, suo Marte inuenerit. Alexander ille Macedonum rex, tam

[p. 8] naturæ quàm fortunæ dotibus iure suspiciendus, tanto bonarum artium & philosophiæ potissimum studio flagrauit, ut etiã in zelotypiam quandam literariam inciderit. Nam is cū omnem propè Asiam armis & exercitu teneret, ubi primū Aristotelem libros suos de auscultatione physica inuulgasse acceperat, in tantis negocijs cum Aristotele, missa statim epistola de editis libris, his uerbis expostulabat: Quòd disciplinas ἀκροαματικάς edidisti, non rectè fecisti. nam qua alia re cæteris præstare poterimus, si ea, quæ abs te accepimus, omnium prorsus fuerint communia? Quippe ego doctrina anteire malim, quàm copijs atq; opulentijs. Hæc Alexander.

Diuino approbatus oraculo rex ille Daud,

thought quite differently. For Mithridates, king of Pontus and Bithynia, understanding that he was the head of his domains, and that his native tongue alone was by no means sufficient for the numerous bodies, over which he reigned, learned so perfectly and understood so thoroughly the twenty-two tongues¹ of the nations, which he had under his sway, that he gave immediate answers to twenty men of those nations without an interpreter, and spoke to each in his own tongue just as if it had been native to him. He also so thoroughly traced out the hidden natures of things, and occupied himself to such good purpose in the science of medicine, that he discovered by his own exertions an antidote to deadly poisons, which even to-day is called Mithridatium, a name derived from him. The great Alexander, king of the Macedonians, rightly renowned as much for the gifts of nature as for those of fortune, burned with so great a zeal for the noble arts, and philosophy in particular, that he even descended to a sort of literary jealousy. For though he was holding almost all Asia by force of arms and his troops, when first he heard that Aristotle had made public his books '*De Auscultatione Physica*,' in the midst of such great concerns he expostulated with Aristotle in the following words, a letter having been at once sent off concerning the publication of the books: "In that you have published your teachings called ἀκροαματικά you have not done rightly; for in what other thing shall I be able to excel the rest, if those things, which I have heard from you, become henceforth the common property of all? For I should prefer to stand first in learning rather than in resources and wealth." Thus said Alexander.

The great king David, approved by the voice of

¹ 'Duas' is perhaps a misprint for 'duarum.'

- [p. 9] qui & propheta fuit diuino numine adflatus, atq; ideo qd regi maximè necessariũ foret, cognoscens, ante omnia literas, nempe sacras expetiuit, ut sibi tẽperare non potuerit, quin diceret, Benedictus es domine, doce me iustificationes tuas, in uia mandatorũ tuorũ delectatus sum, sicut in omnibus diuitijs : in mandatis tuis exercebor, & considerabo uias tuas. Reuela oculos meos, & considerabo mirabilia de lege tua. Bonitatẽ & disciplinã & scientiam doce me : ego autem in toto corde meo scrutabor mandata tua. Nisi quòd lex tua meditatio mea est, tunc fortè perissem in humilitate mea. Quàm dulcia faucibus meis eloquia tua, super mel ori meo. Bonum mihi lex oris tui super milia auri & argenti.
- [p. 10] Lucerna pedibus meis uerbum tuum, & lumen femitis meis. Declaratio sermonum tuorum illuminat, & intellectum dat paruulis. Hactenus rex Dauid, & pace & bello omnium regum illustrissimus.

Rex Solomon huius filius, omniũ, quos unquã terra genuit, sapientiss. cuius unius auctoritati plus tribuendum est, quàm sexcentis adulatoribus diuersum suadẽtibus, cùm totius orbis conditor & omnium bonorum largitor Deus pater, illi, quod sibi optimum, & ex usu suo maximè fore iudicaret, ultro offerret, & poscenti mox se daturũ promitteret, ad hunc modum, ut diuinæ literæ testantur, respondit. Nunc domine Deus, tu me regnare fecisti seruum tuum pro

[p. 11] Dauide patre meo, ego autem sum puer paruulus, & ignorans ingressũ, & introitũ meũ : & seruus tuus in medio est populi, quem elegisti,

God, who was moreover a prophet filled with divine inspiration, and therefore well aware of what was especially necessary for a king, sought before all things learning, and that of course divine, so that he was unable to restrain himself from saying "Blessed art thou, O Lord, teach me thy righteousness, I have delighted in the way of thy commandments, as in all riches: in thy statutes will I exercise myself, and I will consider thy ways. Open thou mine eyes, and I will consider the wonderful things of thy law. Teach me goodness and instruction and learning; but with my whole heart will I examine thy commandments. Unless thy law had been my meditation, then should I perchance have perished in my low estate. How sweet are thy sayings to my mouth, better than honey to my lips. The law of thy mouth is a good to me beyond thousands of gold and silver. Thy word is a lantern unto my feet, and a light unto my paths. The telling of thy discourses giveth light and understanding to babes." Thus far king David, the most illustrious of all kings both in peace and war.

King Solomon, his son, the wisest of all that earth ever bore, to whose single authority more weight is to be given than to six hundred flatterers persuading to a different course, when God the Father, maker of all the world and giver of all good things, of his own accord offered to him what he should judge to be best for himself and for his greatest advantage, and promised that he would grant it at once on his request, replied in this manner, as the Scriptures testify. "Now, O Lord God, thou hast made me, thy servant, to reign in the room of David my father, but I am a little child, and know not my coming in and entering; and thy servant is in the midst of the people whom thou hast chosen, an in-

populi infiniti, qui numerari & supputari non potest præ multitudine. Dabis ergò seruo tuo cor docile, ut populum tuum iudicare possit, & discernere inter bonum & malum: quis enim potest iudicare populum istum, populum tuum hunc multum? Huc usq; Solomon, qui in philosophia tam diuina quàm humana ita non multis pòst annis profecit, ut de stirpibus à cedro usq; ad hissopum disputauerit, & de bestiis, uolucris, reptilibus, & piscibus differuerit.

Quare, prudentissimi quique principes, & [p. 12] fortissimi, nō satis habebant, subditos suos diuitijs, honoribus, uestitu, incessu, & bellica gloria excellere, nisi literis, linguis, philosophia tam diuina quàm humana insuper multum superarent, & à tergo relinquerēt. Quod pater tuus omnium regum, qui hodie uiuunt, eruditissimus, satis ut regem tantum decet, intelligens, & cui Reipub. gubernaculum committitur, quàm necessaria literæ & philosophia sint, prudenter secum perpendens, liberos suos semper eruditissimis quibusque præceptoribus commisit. Duci enim Richmundiæ, piæ memoriæ, fratri tuo Georgium Folberium præceptorē olim meum, uirum insigniter doctum, et mirum rectè instituendæ iuuentutis artificem, & tibi nūc uirum longè [p. 13] doctissimum (uti audio) præfecit.

Qua de causa, illustrissime & optime princeps, sapiētissimorum & fortissimorum regum exempla secutus, atq; potentissimi & eruditissimi patris tui consilio obtemperans, qui te ad meliores imbibendas literas, nunquam non inuitat, incitat & hortatur, dum ætas tua adhuc tenera

numerable people, a people which cannot be numbered or counted for their multitude. Thou shalt give therefore to thy servant a heart that may be taught, that he may be able to judge thy people, and to discern between good and evil: for who is able to judge this people, this great people of thine?" Thus far spoke Solomon, who not many years afterwards so excelled in philosophy both divine and human that he disserted about plants from the cedar even to the hyssop, and discoursed of beasts, birds, reptiles, and fishes.

Wherefore all the most wise and brave Princes have not considered it sufficient to surpass their subjects in riches, honours, garb, gait, and warlike glory, unless beyond this they excelled them far in learning, tongues, and philosophy both divine and human, and left them in the rear. And this your father, the most learned of all the kings who are alive at the present day, well understanding, as becomes so great a king, and one to whom the helm of the State is entrusted, wisely pondering in his mind how necessary learning and philosophy are, always committed his children to the care of the most learned of instructors. For over your brother the Duke of Richmond, of pious memory, he set Georgius Folberius, once my tutor, a man of remarkable learning, and a wondrous handicraftsman for rightly instructing youth, and now over you (as I hear) a man by far the most learned of all.

Wherefore, most illustrious and worthy Prince, following the steps of the wisest and bravest kings, and yielding to the advice of your most powerful and learned father, who so constantly invites you to the draught of superior learning, spurs you on, and exhorts you, while your years are yet tender and

est, & literarũ capacissima, omne genus bonarum literarum obuijs ulnis amplectere, discere, & imbibe, & exantlati in bonas literas laboris olim te minimè pœnitebit. Sed ut ad propositam metam minori cum negotio possis peruenire, libellum De historia auium, in quo Latinis nominibus Græca, Germanica & Britànica in [p. 14] gratiam tuam apposui, ex Aristotele & Plinio, & optimis quibusq; scriptoribus contexui. Hunc ego nominis tui celebritati dedico, & dono: etiã atq; etiã te obtestans, ut hoc meum qualecunq; munusculum æqui boni'q; consulas. Quod si te facturum intellexero, & hunc libellum figuris & auium moribus, & medicinis auctum, & de herbis alium etiam librum, breui, uolente Deo, in lucem emittam. Vale. Dominus Iesus
te nobis sanctissimis moribus
institutum, & optimis literis imbutum, quàm diutissimè inculumem
conferuet.

Coloniæ 5. Idus Februarij,
Anno M.D.XLIIII.

most amenable to learning, embrace with open arms every kind of noble literature, learn and drink it in, and hereafter you will surely not repent of the labour expended upon this noble literature. But that you may be able to reach with less trouble the goal that is laid before you I have compiled from Aristotle and Pliny and all the best writers this little book on 'The History of Birds,' in which I have placed for your pleasure the Greek, German, and British names side by side with the Latin. This I dedicate and offer to the glory of your name: again and again praying you to receive this little gift, such as it is, with fair and favourable consideration. And if I understand that you will do this, I will shortly, God willing, bring to the light of day a further edition of this little book with figures of the birds, their habits, and curative properties, as well as another book on plants. Farewell. May the Lord Jesus preserve you as long as possible unharmed to us, trained in most holy ways and filled with the best of learning.

Cullen [Cologne]. February 9th, 1544.

ARISTOTELES¹.

Buteo.

Aefalo.
Circus.

Percæ
Fringillarij.
Rubetarij.

ACCIPITRUM genus præcipuum Buteo est, Triorchæ² à numero testium nuncupatus: secundum æfalo, tertium circus. Stellaris autem, palumbarius, & pernix³ differunt. Appellantur subuteones, qui latiores⁴ sunt: alij percæ & fringillarij uocantur: alij læues⁵ & rubetarij, qui abundè uiuunt⁶, atque humiuolę sunt. Genera non pauciora quàm decem esse accipitrum aliqui prodiderunt, quæ modo quoq; uenandi⁷ inter se dissident. Alij enim columbam humi confidentem, rapiunt, uolantem non appetunt: alij super arborem, aut tale quid conscendentem, uenantur: sìn humi est, aut uolat, non inuadunt. Alij neq; humi, neq; in sublimi manentē, adgrediūtur, sed uolantem capere conantur. Fertur etiam à columbis quodq; accipitrū genus cognosci. Itaq; cū accipiter prouolat, sī sublimipeta est, manent quo constiterunt loco: sed sī humipeta qui prouolat, est, non manēt, sed continuò auolant.

¹ *Hist. An.* Bk IX. 128—130.

² Aristotle has simply *κράτιστος μὲν ὁ τριόρχης*.

³ Other readings are *πτερνίς*, *πέρνης*, *πτέρνης*.

⁴ Instead of *πλατύτεροι*, some texts have *πλατύπτεροι*, which would make better sense and mean 'broad-winged.'

⁵ *λείοι*, or according to another text *ἐλειοί*.

⁶ The word *εὐβιώτατοι* here, and corresponding expressions throughout the passages quoted in this book, might possibly mean that the birds in question have no particular faults, or are of ordinary respectability. Gaza, however, followed as usual by Turner, seems to have interpreted the word rightly here.

⁷ These three words are not found in Aristotle.

Of the ten kinds of Accipitres.

ARISTOTLE.

THE chief kind of Accipitres is Buteo, which from the number of its testicles is named Triorcha, Æsalo is the second, Circus is the third. Again Stellaris, Palumbarius, and Pernix differ. Those which have more breadth are called Subuteones ; other kinds are named Percæ and Fringillarii ; others Læves and Rubetarii, which get their living most easily, and fly near to the ground. Some have asserted that there are no fewer than ten kinds of the Accipitres which differ from each other in their several modes of hunting. For some sorts seize a Dove when sitting on the ground, but do not touch one flying ; others seek their prey when perched upon a tree, or such like, but if it be on the ground or flying do not attack it. And others seize it neither on the ground, nor when resting aloft, but strive to catch it flying. Moreover it is said that each kind of Accipitres is recognised by Doves. So, when the Accipiter comes forth, if it be such as hunts on high, they stay where they have settled, but, if that which comes be such as takes them on the ground, they stay not, but forthwith fly off.

PLINIUS¹.

Accipitrum genera fedecim inuenimus. Ex
 Circus. ijs circon claudum altero pede, prosperrimi
 †pecuarię augurij nuptialibus negocijs, & †pecuniarię rei².
 Buteo. Triorchē à numero testiū, cui principatum
 in augurijs Phœmone dedit: buteonē hunc
 appellant Rom. Aefalona Græci uocant, qui
 solus omni tēpore apparet. Cæteri hyeme
 Cymindis. abeunt. Nocturnus accipiter cymindis uocatur,
 [p. 17] rarus etiam in syluis, interdiu minùs cernens:
 bellum internecinum cum aquila gerit: cohæ-
 rentesq; sæpè præhenduntur. Hæc Plinius.

Quauquam Aristoteles decem esse accipitrum genera tradat, & Plinius sedecim: neuter tamen horum hæc ita distinxit genera, & descripsit, ut procliue sit lectori suum cuique peculiare nomen ex illorum præscriptis imponere. Quare à me nemo horum exactam differentiam, & cuiusque nomen Britannicum aut Germanicum cum Latino & Græco coniunctum, iure poterit exigere. Ego tamen, quod nomen Britannicum, cuiq; Latino imponēdum esse censeo, lectorem minimè celabo.

Buteo. Buteo τριόρχης Græcè dictus, Anglorum busharda est, nisi fallar: nam miluo magnitudine æquiparatur, semperq; ipse cernitur, qualem Aristoteles octauo libro de historia animalium buteonem describit.

Aefalo. Αισάλων, quoniam iuxta Plinij sententiam omni tempore apparet, & inter minores accipitres sola merlina siue smerla, semper adpareat, mihi Anglorum merlina, & Germanor. smerla esse uidetur.

¹ Hist. Nat. Lib. x. cap. viii.

² If the reading pecuarię is accepted, the meaning would appear to be 'for cattle breeding.'

PLINY.

Of Accipitres we have found sixteen kinds. Circus among them, halting in a foot, of lucky omen in nuptial affairs and money business. Triorches next, to which Phœmone¹ gave the foremost place in auspices, named from the number of its testicles: the Romans call it Buteo and the Greeks Æsalon: it is the only kind which may be seen at every time. The rest leave us in winter. An Accipiter that flies by night is called Cymindis; it is rarely found in woodlands, in the day it scarce can see: it wages deadly warfare with the Aquila, and they are often captured clinging to each other. So far Pliny.

Though Aristotle may set forth that there are ten kinds of Accipitres, and Pliny that there are sixteen, yet neither of them has distinguished or described the kinds so that it may be easy for a reader to apply to each its proper name from their accounts. So no one can in fairness claim from me their exact difference, nor yet the British or the German name of each, together with the Latin or the Greek equivalent. I will, however, surely not conceal from you, my reader, what I think to be the British name, and to which Latin name it ought to be applied.

Buteo, called in Greek *τριόρχης*, if I do not err, is the Buzzard of the English, for it is compared with Milvus as to size; moreover it is seen at all times, and is such a bird as Aristotle makes his Buteo in the eighth book of the 'History of Animals.'

Αίσάλων, since in Pliny's judgment it appears at every season, and among the smaller Hawks the Merlin or the Smerl alone seems to appear² at all times, is, I think, the Merlin of the English and Smerl of the Germans.

¹ Phœmone, called 'Daughter of Apollo,' was a priestess at Delphi. (See Pliny *Hist. Nat.* ed. Hardouin: Lipsiæ, 1791, Index Auctorum, p. 340.)

² This seems to be the force of the subjunctive here, if it is not an oversight.

[p. 18] *Palumbarius.* *Accipitrem palumbarium ideo Anglorum sparhau-*
cam, & Germanorum speruuerum esse puto, quòd pa-
lumbes, columbos, perdices & grandiusculas aues inse-
quatur.

Fringillarius. *Fringillarium Anglorum hobbiã esse conijcio. Est*
autem hobbia accipiter minimus, coloris cæteris nigri-
oris. In capite duos habet nigerrimos in pallido neuos.
Galeritas & fringillas plerumq; captat, in excelsis ar-
boribus nidulatur, & hyeme nusquam cernitur.

Rubetarius. *Rubetarium esse credo accipitrem illum, quem Angli*
hen harroer nominant. Porro ille apud nostros à dila-
niandis gallinis nomen habet. Palumbarium magni-
tudine superat, & coloris est cinerij. Humi sedentes
aves in agris, & gallinas in oppidis & pagis repentè
adoritur. Præda frustratus, tacitus discedit, nec un-
quam secundum facit insultum. Hic per humum om-
nium uolat maximè.

Subbuteo. *Subbuteonem esse puto, quem Angli ringtalum appel-*
lant, ab albo circulo, qui caudam circuit. Colore est medio
inter fuluum & nigrũ, butcone paulò minor, sed multò
agilior. Prædam eodem modo, quo superior captat.

[p. 19] *DE ALCEDONE.*

Ἀλκυὼν, alcedo, Anglicè the kynges fyfsher, Germa-
nice eyn eifvogel.

ARISTOTELES¹.

Alcedo non multò amplior passere est, co-
 lore tum uiridi, tum cœruleo, tum etiam leuiter
 purpureo insignis: uidelicet non particulatim
 colore ita distincta, sed ex indiscreto uariè re-
 fulgens corpore toto & alis & collo, rostrum
 subuiride, longum & tenue. Alcedonum² quo-

¹ *Hist. An.* Bk IX. 85.

² *Hist. An.* Bk VIII. 47.

The *Accipiter palumbarius*¹ I take to be the Sparrow-Hawk of the English and the Sperwer of the Germans, since it preys on Doves, Pigeons, and Partridges and the bigger sorts of birds.

The *Fringillarius* I guess to be the Hobby of the English. Now the Hobby is a very little Hawk of darker colour than the other kinds. It has upon the head two spots of deep black on a lighter ground. It catches for the most part Larks and Finches, nests on lofty trees, and is not seen in winter anywhere.

The *Rubetarius* I think to be that Hawk which English people name Hen-Harrier. Further it gets this name among our countrymen from butchering their fowls. It exceeds the *Palumbarius* in size, and is in colour ashen. It suddenly strikes birds when sitting in the fields upon the ground, as well as fowls in towns and villages. Baulked of its prey it steals off silently, nor does it ever make a second swoop. It flies along the ground the most of all.

The *Subbuteo* I think to be that Hawk which Englishmen call Ringtail from the ring of white that reaches round the tail. In colour it is midway from fulvous to black; it is a little smaller than the *Buteo*, but much more active. It catches prey in the same manner as the bird above.

OF THE *ALCEDO*.

Ἄλκυων, *alcedo*, in English the kynges fisher, in German eyn eissvogel.

ARISTOTLE.

The *Alcedo*, not much larger than the *Passer*, is remarkable for being in its colour green and blue, and even slightly purple, not, that is to say, in separate parts, as if it had the colour perfectly distinct, but variably shining over every part alike of the whole body, with the wings and head. The beak is greenish, and is long and thin. The tribe of

¹ Later authors are probably more correct in applying this name to the Goshawk, which suits even Turner's account better.

que genus aquas adamat, quod duplex est : alterum uocale, harundinibus insidens, alterum mutum, quod ampliore corpore est . utrique dorsum cœruleum. Sed alcedo apud mare quoque uersatur.

PLINIUS ¹.

Ipse auis paulò amplior pascere², colore [p. 20] cyaneo, ex parte maiore, tantum purpureis & candidis admixtis pennis, collo gracili ac pro-cero. Alterum genus earum, magnitudine distinguatur, & cantu. Minores in harundinetis canunt. Halcyonem uidere rarissimū est, nec nisi Vergiliarum occasu, & circa solstitia, brumam'ue, naue aliquando circumuolata, statim in latebras abeuntem. Fœtificant bruma, qui dies Halcionides uocantur, placido mari per eos & nauigabili, Siculo maximè. In reliquis partibus est quidem mitius pelagus. Siculū utiq; tractabile. Faciunt autem septem ante brumam diebus nidos, & totidem sequentibus pariūt. Nidi earum admirationem habent, pilæ figura paulū eminente, ore perquàm angusto, grandium [p. 21] spongiarum similitudine, ferro intercidi non queunt, frangunturq; ictu ualido, ut spuma arida maris. Nec unde confingantur inuenitur. Putant ex spinis aculeatis, piscibus enim uiuunt. Subeunt & in amnes. Pariunt oua quina.

¹ *Hist. Nat. Lib. x. cap. xxxii.*

² Lege 'passere.'

Kingfishers, of which there are two sorts, is fond of watersides : one is a vocal bird, which sits on reeds, the other, which is of a larger size, is mute. The back is blue in both. The Kingfisher, however, also haunts the sea.

PLINY.

This bird is little bigger than the *Passer*, for the most part blue in colour, with the wings alone of purple mixed with white, and with a long and slender neck. Each of the two kinds may be distinguished by its size and voice. The lesser sing in reed-beds. It is very rare to see the *Halcyon*, and this occurs only towards the setting of the *Pleiades* and near the solstice or in winter-time, when, after circling round the ship awhile, it hurriedly departs again to its retreat. They breed in winter, at the season called the *Halcyon days*¹, wherein the sea is calm and fit for navigation, the Sicilian sea particularly so. Elsewhere indeed the ocean is less boisterous. The Sicilian is certainly gentle enough. Now these birds build their nests in the seven days before the winter solstice, and hatch out their young in the seven following. Their nests compel our wonder, of a ball-like shape, with a small jutting part and very narrow hole, like sponges of great size ; they cannot be cut open with an iron tool, but may be broken by a vigorous blow, as dry sea-foam² may be. It is not known of what these are composed. Some think of pointed bones, since the birds live on fish. They also dive in rivers, and lay five eggs each.

¹ For the origin of this ancient tradition, the reader may be referred to any work dealing with Greek mythology.

² By 'dry sea-foam' Pliny probably meant masses of whelks' eggs.

Præter hæc duo ab Aristotele & Plinio descripta genera, auem noui, quæ si alcedonum generibus non sit adscribenda, sub quo genere contineatur, prorsus nescio. Ea sturno paulò minor est, corpore toto nigro, excepto uentre albo. Caudam habet breuiusculam, rostrum alcedone paulò breuius. Ante uolatum, alcedonis more crebrò nutat, & in uolatu gemit: uoce alcedonē ita refert, ut, nisi uideas, alcedonem esse iurares: in ripis fluminum, non procul à mari uidi, aliàs nusquam. pisciculis uicētat ut superiora alcedonum genera. Nidum huius nunquam uidi. Morpetenses, apud quos auem uidi, cornicem uocant aquaticam.

a uuater
crauu.

DE ANATE.

Νῆττα, *anas*, Anglicè *a duck*, Germanicè *eyn endt*.

PLINIUS¹.

[p. 22] Anates solæ, quæque sunt eiusdem generis, in sublimē sese protinus tollunt, atq; è uestigio cælum petunt, & hoc etiam ex aqua.

DE ANSERE.

Χήν, *anser*, Anglicè *a goose*, Germanicè *eyn ganß*.

Aristoteles simul & Plinius duo præcipua anserum genera faciunt: hic anserem in maiorem & minorem, ille in domitū et ferū diuidens. Sed Plin. præter hæc duo anserum præcipua genera, Penelopes² et chenalopeces, ut unus textus habet, & ut alius habet, chenalopeces, & chenerotes anserini esse generis tradit. Prior lectio sic habet,

¹ *Hist. Nat. Lib. x. cap. xxxviii.*

² Judging from p. 148 of the original work the singular of this word is 'Penelops,' and it is probably by mistake that the Wigeon has been called *Mareca penelope*.

Besides the two kinds thus described by Aristotle and Pliny I know of a bird, of which, if it should not be properly ascribed to the Kingfisher tribe, I really cannot say under what head it ought to go. It is a little smaller than a Starling, with the body wholly black, except for a white belly, and it has the tail comparatively short, the beak a little shorter than the Kingfisher. Before a flight it dips repeatedly, after the manner of the Kingfisher, and cries out as it flies; it is so like the Kingfisher in voice that, if you did not see it, you would swear it was a Kingfisher. I have observed it on the banks of streams not far from the sea-side, but nowhere else. It lives on little fishes, like the aforesaid kinds of Kingfishers. I never saw its nest. The inhabitants of Morpeth, where I saw the bird, call it a water craw¹.

OF THE ANAS.

Νῆττα, anas, in English a duck, in German eyn endt.

PLINY.

Anates only, and birds of like kind, rise in the air at once, and make straight for the sky, and that even from the water.

OF THE ANSER.

Χήν, anser, in English a goose, in German eyn ganss.

Aristotle agrees with Pliny in making two chief kinds of Geese, the latter separating them into the greater and the less, the former into tame and wild. But Pliny tells us that besides these two chief kinds of Geese, there are of the Goose kind Penelopes and Chenalopeces, as one text has it, as another goes, Chenalopeces and Chenerotes. The first reading stands thus:—

¹ The bird meant is undoubtedly the Water Ousel or Dipper (*Cinclus aquaticus*), which still goes by the name of Water Craw in the north of England. It is curious that Turner should never have seen its nest when he was in Northumberland.

Anferũ generis funt Penelopes, & quibus lautiores epulas Britannia non nouit, chenalo-
peces, anfere ferè minores. *Altera sic habet:*
Anferini generis funt chenalopeces & quib⁹
lautiores epulas Britannia nō nouit, chenerotes¹.

[p. 23] *Posterior lectio mihi magis approbatur, nam & nos una aue locupletat, et penelopes anatini potiùs q̃ anferini generis eruditis esse uidentur. Sed quænam istæ aues, & quibus nominibus apud nostrates appellantur, dicere tentabo. Chenalopex, ab anfere et uulpe nomen habet, & Latinè à Gaza uulpanfer dicitur. Nostrates a bergander. hodie bergandrum nomināt, anate longior & grandior uulpanfer est, pectore rufescente, in aquis degit, & in cuniculorũ foueis. interdum & in excelsarum rupium cauernis (unde fortè nomen ab Anglofaxonibus, nostris patribus sortitus est) nidificat. Nusquam aliàs uulpanferem uidi, nisi in Tamisi fluuio. Aiunt tamen frequētem esse in insula Tenia uocata, & illic in scrobibus cuniculorum nidulari. Moribus admodum uulpinis est. nam dum teneri adhuc pulli sunt, si quis eos captare tentet, prouoluit sese uulpanfer ante pedes captantis, quasi iam capi possit, atq; ita allicit ad se capiendam hominem, eousq; dum pulli effugiant: tum ipse auolat & renocat prolem. Chenerotes quænam aues sint, puto paucissimos hodie esse, qui nouerunt. Neq;*

¹ *Hist. Nat. Lib. x. cap. xxii.*

“Of the Goose kind there are Penelopes and also Chenalopeces, the latter generally smaller than a Goose; and Britain knows no richer feast than these.”

The second runs:—

“Of the Goose kind are Chenalopeces and Cherotes, Britain knows no richer feast than these.”

To me the latter reading most approves itself, for it both makes us richer by one bird, and the Penelopes seem to our learned men to be of the Duck tribe rather than of the Goose. But I will try to say what these birds are and by what names they go among our countrymen. The Chenalopex¹ takes its name from the Goose and the Fox, while it is called by Gaza Vulpanser in Latin, though our people nowadays name it Bergander². It is longer than a Duck and bigger, with a ruddy breast. It lives upon the waters and in coney's burrows. At times it even nests in holes of lofty rocks (whence possibly the name was first allotted to it by our ancestors the Anglo-Saxons). I have nowhere else seen the Vulpanser save upon the river Thames. Nevertheless they say that it is plentiful upon the isle which is called Tenia³, and that it breeds in coney's burrows there. In habits it is very like a Fox, for, while the young are still of tender age, should any one attempt to capture them, the old Vulpanser rolls upon the ground before his very feet⁴, as if she could be taken there and then, and thus allures the man to follow her, until the young are able to escape; then she flies off and summons back her brood. I think that there are very few men now who know what sort of birds the

¹ Turner's bird was undoubtedly the Sheld-Drake (*Tadorna cornuta*), notwithstanding the fact that the name *Chenalopex* has been conferred on the so-called 'Fox-Goose' of Africa.

² The Sheld-Drake is still the Bargander or Bergander of some districts of England; possibly the correct spelling should be Burgander, i.e. Burrow Duck. The word seems to have nothing to do with Berg=a mountain.

³ Possibly St Mary's, or even Coquet Island.

⁴ The Sheld-Drake does not usually behave thus.

ego, licet Britannus, chenerotes nostros satis noui: nā præter duo Aristot. genera, anserū adhuc duo genera noui in Britānia, ad quorū neutrum si chenerotes pertineāt, chenerotes mihi penitus ignotos esse ingenuè fatebor. Prior anser à nostris hodie brāta & berniclæ uocatur, & fero ansere minor est, pectore aliquò usq[ue]

[p. 24] nigro. Cætero cinerio, anserum ferorū more uolat, strepit, paludes frequētat, & segetē depopulatur. Caro huius paulò insuauior est, & diuitibus minùs appetita. Nidum berniclæ, aut ouum nemo uidit: nec mirum, quum sine parentis opera berniclæ ad hunc modū spontaneam habeāt generationē. Quum ad certum tempus, malus nauis in mari cōputruit, aut tabulæ, aut antennæ abiegnæ, inde in principio ueluti fungi erumpūt: in quibus temporis progressu, manifestas auiū figuras cernere licebit, deinde pluma uestitas, postremò uiuas & uolantes. Hoc, ne cui fabulosum esse uideatur, præter cōmune omnīū gentiū littoralīū Angliæ, Hiberniæ & Scotiæ, testimoniū Gyraldus ille præclarus historiographus qui multò fælicius q̃ pro suo tempore Hiberniæ historiam conscripsit, nō aliam esse berniclarū generationē testatur. Sed, quum uulgo non satis tutū uideretur fidere, et Gyraldo ob rei raritatem non satis crederem, dum hæc, quæ nunc scribo, meditarer, uirum quendam, cuius mihi perspectissima integritas fidem merebatur, professione Theologum, natione Hibernum, nomine Octauianū, consului num Gyraldum hac in re fide dignum censeret? qui per ipsum iurans, quod profitebatur euangelium, respondit, uerissimum esse, quod de generatione huius auis Gy-

[p. 25] raldus tradidit, seq[ue] rudes adhuc aues oculis uidisse, & manibus contrectasse: breuiq[ue] si Londini mensem unum aut alterum manerem, aliquot rudes auiculas mihi aducētas curaturū. Ista berniclæ generatio nō usq[ue] adco

Chenerotes are. And, though I am a Briton, I am not quite sure about our Chenerotes; for as yet, apart from the two kinds that Aristotle gives, I know two sorts of Geese in Britain and will frankly own that, if the Chenerotes are not to belong to either of them, they are quite unknown to me. The first Goose by our people nowadays is called the Brant and Bernicle, and is a smaller bird than the Wild Goose, with the breast partly black. The rest is ashen grey. It flies, gabbles, haunts swamps, and devastates green crops, like the Wild Goose. Its flesh is somewhat strong, and is the less sought after by the rich. No one has seen the Bernicle's nest or egg, nor is this wonderful, since Bernicles without a parent's aid are said to have spontaneous generation in this way: When after a certain time the firwood masts or planks or yard-arms of a ship have rotted on the sea, then fungi, as it were, break out upon them first, in which in course of time one may discern evident forms of birds, which afterwards are clothed with feathers, and at last become alive and fly. Now lest this should seem fabulous to anyone, besides the common evidence of all the long-shore men of England, Ireland, and Scotland, that renowned historian Gyraldus¹, who composed a history of Ireland in much more happy style than could have been expected in his time, bears witness that the generation of the Bernicles is none other than this. But inasmuch as it seemed hardly safe to trust the vulgar and by reason of the rarity of the thing I did not quite credit Gyraldus, while I thought on this, of which I now am writing, I took counsel of a certain man, whose upright conduct, often proved by me, had justified my trust, a theologian by profession and an Irishman by birth, Octavian by name, whether he thought Gyraldus worthy of belief in this affair. Who, taking oath upon the very Gospel which he taught, answered that what Gyraldus had reported of the generation of this bird was absolutely true, and that with his own eyes he had beholden young, as yet but rudely formed, and also handled them, and, if I were to stay in London for a month or two, that he would take care that some growing chicks should be brought in to me. This curious generation of the Bernicle will not appear so very

¹ Giraldu Cambrensis, *Topographia Hibernica* Distinctio 1. cap. xv.

prodigiosa illis uidebitur, qui quod Aristoteles de uolucres ephemero scripsit, legerint.

De ephimero autem Aristoteles¹ libro quinto de historia animalium ita scribit. Hyppanis fluuius apud Cymerium Bosporum sub solstitio, desert ueluti folliculos acinis maiores, quibus quadripedes uolucres erumpunt: quod genus animalis in postmeridianum² usque diei tempus uiuit & uolat: mox descendente sole, macrescit & languet³: deinde occidente, moritur, uita non ultra unum diem protracta: unde ephemerum, id est, diarium⁴ appellatum est. Hęc Aristotel.

Quę si uera sunt, & tãto philosopho digna, superioris auis generationi non parum fidei adstruent.

a folend
guse.

[p. 26] *Alter anser, de quo promisi me dicturũ, marina auis est, ex uenatu piscium uictitans, magnitudine superiore anseris paulò minor: anserem tamen noce & forma per omnia refert, nidulatur in mari Scotico, in rupibus excelsis, insulę Bassi, per antiphrasim, opinor, dictę: nec alias uspiam in tota Britannia. Hic tanto amore suos pullos prosequitur, ut cum pueris per funes in corbibus ad auferēdos eos demissis, acerrimè non sine uitę periculo conflictetur. Nec silentio prætereundum est, ex adipe huius anseris (est enim insigniter adiposus) unguentum à Scotis ad multos morbos utilissimum fieri, quod cum commageno à Plinio⁵ celebrato, meritò bonitate & remediorum numero potest certare. Iam quũ anserum genera, licet diligentissimè inquirēs, apud Britannos plura inuenire non possim, chenerotes*

¹ Bk v. 107.

² μέχρη δειλῆς.

³ These two words are not in the original Greek.

⁴ This explanation is not given by Aristotle. We have here an instance of the insertions common in old authors, which will not be noticed hereafter in each case, as being too numerous. Another instance is found with regard to 'Albicilla' (p. 30).

⁵ *Hist. Nat. Lib. x. cap. xxii.*

marvellous to those who may have read what Aristotle wrote about the flying creature called Ephemerus. Now Aristotle writes thus of the Ephemerus in the fifth book of his History of Animals:—

“The river Hyppanis¹, near the Cymerian Bosphorus² when the solstice is nigh, brings down small pouches, as it were, each larger than a grape, from which four-footed flying creatures burst; a sort of animal which lives and flies until the afternoon of the same day, but presently at the sun’s going down withers and languishes, and finally, at the sun’s setting, dies, lasting no longer than a single day, whence it is called Ephemerus, that is, the creature of a day.” Thus Aristotle writes.

Now if these things are true, and worthy of the great philosopher, they will impart no little credibility as to the generation of the aforesaid bird.

The second Goose, of which I promised I would speak, is a sea-bird, which lives by hunting fishes, somewhat less in size than the Goose given above; and yet in voice and aspect it recalls the Goose in every way; it nests within the Scottish sea, upon the lofty cliffs of the Bass Isle—so called, as I opine, by an antiphrasis³—and nowhere else in all Britain. This bird looks to its young with so much loving care, that it will fight most gallantly with lads that are let down in baskets by a rope to carry them away, not without danger of its life. Nor must we fail to mention that a salve, most valuable for many a disease, is made by Scots from the fat of this Goose (for it is wonderfully full of fat) which may deservedly rival the Commagenum vaunted much by Pliny, in its virtue and the number of its cures.

Now since, though searching with the greatest care, I cannot find any more kinds of Geese among Britons,

¹ Now the Bog.

² Between the Sea of Azov and the Black Sea.

³ As if the derivation was from the French *bas*=low.

(qui ab amore mihi nomen habere videntur), aut berniclae aut Bassani anseres sunt, aut mihi prorsus ignoti.

DE AQVILA.

ἀετὸς, aquila, Anglicè an egle, Germanicè ein ärn, oder ein adler.

ARISTOTELES¹.

Aquilarum plura sunt genera. Vnum, quod [p. 27] pigargus ab albicante cauda dicitur, ac si albicillam nomines. gaudet hæc planis, & lucis et oppidis. Hinnularia² à nonnullis uocata cognomine est. montes etiam, fyluamq; suis freta uiribus, petit. reliqua genera rarò plana & lucos adeunt.

Pygargus, quum sit primum aquilarum genus, Germanorum literatores turpiter errant, qui pygargum suum trappum faciunt, qui apud Aristotelem tetrix, & Plinio tetrao est, ut postea docebo. Pygargus Anglorum lingua, nisi fallar, crna uocatur.

an erne.

DE PLANGA AUT CLANGA EX ARISTOTELE³.

Alterum genus magnitudine secundum & uiribus, clanga⁴ aut planga nomine, saltus & conualles, & lacus incolere solitum, cognomine anataria⁵, & morphna, à macula pennæ, quasi neuia⁶ dixeris, cuius etiam meminit Homer. in exitu Priami⁷.

[p. 28] PLINIUS⁸ DE MORPHNA SIVE PLANGA.

Morphnos, quam Homerus & percnon uocat, aliqui & plancum & anataria, secūda magnitu-

¹ *Hist. An.* Bk IX. 111.

² νεβροφόνον = fawn-slayer.

³ *Hist. An.* Bk IX. 112.

⁴ For πλάγγος some texts have πλάνος. The word 'clanga' does not seem to be represented in the Greek.

⁵ νητροφόνος = duck-slayer.

⁶ This explanation is not in Aristotle.

⁷ *Iliad*, Bk XXIV. l. 316.

⁸ *Hist. Nat.* Lib. x. cap. iii.

the Chencrotes (which seem to me to get their name from "love"¹) are either Bernicles, or the Geese of the Bass, or are decidedly unknown to me.

OF THE AQUILA.

ἀετὸς, aquila, in English an eagle, in German ein ärn, or ein adler.

ARISTOTLE.

Of Aquilæ there are several kinds. One which is called Pygargus from its whitish tail, as though you were to name it Albicilla, loves plains, groves, and towns. For by-name it is called by certain Hinnularia. It even seeks the mountains and the wood, relying on its might. The other kinds seldom approach the plains and groves.

Now, seeing that Pygargus is the first kind of the Aquilæ, the German scribblers err disgracefully, who reckon it their Trapp, which is the Tetrax in the works of Aristotle and the Tetrao of Pliny, as I shall shew afterwards. Pygargus, if I err not, in the English tongue is called an Erne.

OF THE PLANGA OR CLANGA, FROM ARISTOTLE.

Another kind, second in size and strength, by name Clanga or Planga, generally haunts glades and valleys and lakes. It has the by-name Anataria, and Morphna from the marking on the wing, as though you should say spotted. Of this Homer makes mention in the scene of Priam's death.

PLINY ON THE MORPHNA OR PLANGA.

Morphnos, which Homer also calls Percnos, some name Plancus and Anataria, second in size and

¹ A very doubtful derivation.

† in deic-
ctu.

† hostem
aliās abūdat.

[p. 29]

dine & ui, huicq; uita circa lacus. Ista circa stagna aquaticas aues appetit mergētes se subinde, donec sopitas lassatasq; rapiat. Spectanda dimicatio, aue ad perfugia littorum tendente, maximè si condensa harundo sit: aquila † inde ictu abigente ala, & cūm appetit in lacus cadente, umbramq; suam nanti sub aqua à littore ostendente: rursus aue in diuersa, & ubi minimè se credat expectari, emergente. Hæc causa est gregatim aibus natandi, quia plures simul non infestantur: resperfu pinnarū † hostē obcæcantes. Sæpè & aquilæ ipsæ non tollerantes pondus apprehensum, unà merguntur. Hæc Plinius.

Omnia, quæ Aristoteles & Plinius percno hætenus tribuerunt, Anglorum balbushardo conueniunt, si solam magnitudinem exceperis, quæ si alia adfuerint, hic fortassis non oberit. est autem illa, quam anatariam esse conijcio, auis buteone maior & longior, neuo albo in capite, colore fusco proximo, ad ripas fluminum, stagnorum et paludium semper degens, uiuit ex uenatu anatum et gallinarum nigrarum, quas Angli coutas nomināt. Venationem hanc, cuius meminit Plinius, inter aquilam istam (si aquila dicenda sit) & aues aquaticas, non solum ego sæpissimè uidi, sed infiniti apud Anglos quotidie uident. Si qua terræ portuncula super aquas inter arundincta emineat, in hac solet nidum facere, ut quoniam uolatu non admodum ualet, à præda non procul absit. Aues subitò adoritur, & sic capit. Cuniculos ista interdum etiam dilaniat. Nunc an ista anataria sit nec ne, doctis uiris iudicandum propono.

strength; it passes its life round lakes. By pools it chases water-birds, which dive from time to time, until it catches them sleepy and weary. The contest is a sight to see, the quarry seeking refuge on the shore, chiefly where reeds are thick, and thence the Aquila drives it away with a stroke of the wing and plunges in the lake as it swoops from above, shewing its shadow to the bird as it swims under water from the shore. Again the latter tries a different place and comes up where it thinks that it will least be marked. This is the cause of birds swimming in flocks, for they are not molested when in companies, and blind their enemy by splashing with their wings. The Aquilæ themselves, moreover, often are immersed, not being able to support the weight that they have clutched. Thus Pliny.

All things that Aristotle and Pliny have attributed to the bird Percnos so far well agree with the Balbushard of the English¹, if one may except its size alone, and if the rest be present, that perhaps should not stand in the way. Now the bird which I apprehend to be the Anataria, being bigger and longer than the Buteo, with a white patch upon the head, and nearly fuscous in colour, always haunts the banks of rivers, pools, and swamps; it lives by hunting Ducks and those black fowls which Englishmen call Coutts. The conflict of which Pliny makes mention above between this Eagle (if it should be called an Eagle) and the water-birds I have seen often, and not I alone, but countless Englishmen witness it daily. If anywhere a little space of ground rises among the reed-beds, there the bird is wont to make a nest, that, since in power of flight it is not very strong, it may not be far distant from its prey. It suddenly attacks birds, and thus takes them. It also sometimes butchers coney. Now whether this may be the Anataria or not I put it to the learned to decide.

¹ The Bald-Buzzard or Marsh-Harrier (*Circus æruginosus*).

DE TERTIO GENERE EX ARISTOTELE¹.

Tertium genus colore nigricās, unde nomen [p. 30] accepit, ut pulla & fulua² uocetur, magnitudine minima, sed uiribus omnium præstantissima. Hæc colit montes & fyluas & leporaria cognominatur.

PLINIUS³.

Melænaëtos.

Melænaëtos à Græcis dicta, eademq; ualeria, minima magnitudine, uiribus præcipua, colore nigricans: sola aquilarum fœtus suos alit, cæteræ fugant: sola sine clangore, sine murmuratione.

DE QUARTO GENERE EX ARISTOT.⁴percnop-
terus.

Quartum genus percnopterus ab alarum notis, capite albicante, corpore minore, quàm cæteræ adhuc dictæ, hæc est. Sed breuioribus alis, cauda longiore, uulturis speciem hæc refert. Subaquila⁵, & aquila montana cognominatur. In- [p. 31] colit lucos, degener, nec uicijs cæterarum caret, & bonorum, quæ illæ obtinent, expers est: quippe quæ à coruo, cæterisq; id genus auibus uerberetur, fugetur, capiatur. Grauis enim est, uictu iners: examinata⁶ fert corpora: famelica semper est, et querula, clamat, & clangit.

DE HALIÆETO.

Haliæetus Græcè & Latinè, Anglicè an *osprey*, Germanicè *eyn viſharn*.

¹ *Hist. An.* Bk IX. 113.² Aristotle has merely: καλεῖται δὲ μελανίετος καὶ λαγωφόνος.³ *Hist. Nat.* Lib. X. cap. iii.⁴ *Hist. An.* Bk IX. 114; a very free version.⁵ Some texts read γυπαίετος for ὑπαιετός.⁶ Lege 'exanimata.' Aristotle has τὰ τεθνεῶτα φέρων.

OF THE THIRD KIND FROM ARISTOTLE.

The third kind in colour is blackish, whence it has received its name, so that the bird is called Pulla and Fulvia, in size the least of all and yet chiefest in strength. It haunts mountains and woods, and is called also Leporaria.

PLINY.

The bird called Melænaetos among the Greeks, which is the same as the Valeria, is very small in size, but chief in strength, in colour blackish : of the Aquilæ this kind alone fosters its young, the others drive them off : it is the only one without a scream, without a softer note.

OF THE FOURTH KIND FROM ARISTOTLE.

The fourth kind, called Percnopterus, from having spots upon the wings, is whitish on the head ; it has a smaller body than the other sorts spoken of hitherto. But with its shorter wings and longer tail it has the aspect of a Vulture. It is called besides Subaquila and Mountain Aquila. It dwells in woodlands, an ignoble bird, not lacking the bad qualities of others, but void of the good that they possess. For it is beaten, put to flight, and caught by the Raven and by other birds like that. Further it is unwieldy, sluggish to get food, and carries off dead bodies ; it is always ravenous and querulous ; it cries continually and screams.

OF THE HALIÆTUS.

Haliaetus in Greek and Latin, in English an Osprey, in German eyn vishärn.

PLINIUS¹.

Supereſt Haliæetus, clariffima oculorum acie, librans ex alto ſeſe uiſoꝝ in mari piſce, præceps in eũ ruens, & diſcuſſis pectore aquis rapiens.

ARISTOTELES².

Haliæetos, hoc eſt, marina aquila, ceruice [p. 32] magna & craſſa, alis curuantibus, & cauda lata eſt. Moratur hæc in littoribus & oris. Accidit huic ſæpius, ut quum ferre, quod ceperit, nequeat, in gurgitem demergatur.

Haliæetos apud Anglos hodie notior eſt, quàm multi uelint, qui in uiuarijs piſces alunt: nam piſces omnes breui tempore aufert. Piſcatores noſtrates eſcis fallendis piſcibus deſtinatis, haliæcti adipem illinunt, aut immiſcent, putantes hoc argumento eſcam efficaciorẽ futuram, quòd haliæeto ſeſe in aëre librâte, piſces quotquot ſubſunt (natura aquilæ ad hoc cogente, ut creditur) ſeſe reſupinõt, & uẽtres albicantes, ut quem liberet, eligeret, exhibeãt.

DE AQUILA VERA EX ARISTOTELE³.

Sextum genus gneſium, id eſt, uerũ germanuꝝ appellant. Vnũ hoc ex omni aquilarum genere, ueri incorruptiꝝ ortus creditur. Maxima omnium aquilarum hæc eſt, maior etiam [p. 33] quàm offiſfraga⁴: ſed cæteras aquilas uel ſeſqui- altera portione excedit, colore ruſſa eſt, conſpectu rara.

Angli. a
right egle
German.
eyn edel arn.

DE ARDEA.

ἐρωδιός, ardea, Anglicè an heron. Germanicè eynerger.

¹ Hist. Nat. Lib. x. cap. iii.

² Hist. An. Bk IX. 115, a free version.

³ Hist. An. Bk IX. 116.

⁴ Gaza translates φήνη by ossifraga, but it is very doubtful what bird the ossifraga really was. Possibly it should be identified with the Lämmergeier (cf. Prof. Newton, Dict. Birds, p. 660).

PLINY.

The *Haliæetus* remains, with eyesight of the keenest, poising itself aloft when it spies fishes in the sea below, then dashing headlong on them and securing them, the waters being parted by its breast.

ARISTOTLE.

The *Haliæetos*, that is to say Sea Eagle, has the neck both big and thick, bowed wings, and a broad tail. It bides upon the sea-coast and the shores. It often happens, when it cannot lift what it has taken, that it is submerged beneath the tide.

The Osprey is a bird much better known to-day to Englishmen than many who keep fish in stews would wish; for within a short time it bears off every fish. Our anglers smear or mix their bait with Osprey's fat, arguing that thus the bait will prove more efficacious from the fact that, when the Osprey hovers in the air, whatever fishes be below turn up and shew their whitish bellies (as it is believed, the nature of the *Aquila* compelling them to this), that it may choose that one which it prefers.

OF THE TRUE AQUILA FROM ARISTOTLE.

The sixth kind men call Genuine, or true and thoroughbred. Of all the various kinds of *Aquilæ* this is the only one that is believed to be of true and unstained origin. This is the largest of all *Aquilæ* and bigger even than the *Ossifrage*, for it surpasses by one half as much the other *Aquilæ*; in colour it is reddish brown, but it is rarely seen.

OF THE ARDEA.

ἐρωδιός, *ardea*, in English a heron, in German *eyn reyger*.

ARISTOTELES¹.

Ardearum tria sunt genera, pella, alba, stellaris, piger cognomine. Pellæ coitus difficilis est: uociferatur enim, & sanguinem ex oculis (ut aiunt) emittit cum coit; parit etiam ægrè fummoq; cum dolore. Pella fagax² & cœnæ gerula est, & operosa³. Agere interdiu solet: colore tamen & prauo & aluo humida. Reliquarum duarum, alba colore est pulchro, & coit, & nidulatur & parit probè, pascitur paludibus, lacu, campis & pratis. Sed stellaris piger cognominata, (in fabula est, ut olim è feruo in auem transferit) atque, ut cognomē sonat, iners ociofaq; est. Phoici appellatæ⁴, peculiare præ cæteris est, ut oculos potissimū appetat⁵. Petit lacus & fluuios ardea⁶ & albardeola, quæ magnitudine minor est, rostro lato, porrectoq;.

Pella apud Anglos in excelsis arboribus, nō procul à ripis fluminum crescentibus nidum facit. Superior pars corporis cyanea est, inferior autem nōnihil candidat, uentris excrementis liquidioribus inuadentes se subito aquilas, aut accipitres abigit, & se ita defendit. Vidi & huius generis, licet raras, albas, quæ neque corporis magnitudine, neque figura, sed solo colore, à superiore distulerunt. Visa est etiam alba cum cyanea apud Anglos nidulari, & prolem gignere. Quare eiusdem esse speciei, satis constat. Albardeolam, quæ Græcè λευκερωδιός dicitur, semel tantum in Italia uidi, pella multo minor est, & hominis conspectū nō perinde atq; cærulea fugit. Hæc si nō uidissem, Anglorū shouedromble.

¹ *Hist. An.* Bk IX. 19.² *Hist. An.* Bk IX. 93.³ Aristotle's word is ἐπαγπος, which Sundevall renders by the Swedish equivalent of 'forages round the fields.'⁴ *Hist. An.* Bk IX. 94.⁵ This seems to mean that the φῶς cats other creatures' eyes, for Aristotle says: μάλιστα γάρ ἐστιν ὀφθαλμοβόρος τῶν ὀρνίθων.⁶ *Hist. An.* Bk VIII. 46.

ARISTOTLE.

Of Ardeæ there are three kinds, Pella, Alba, and Stellaris, but the last has the by-name of Piger. The coupling of Pella is difficult, for it screams while it couples and (they say) emits blood from its eyes: it also brings forth painfully and with extreme distress. The Pella is sagacious, quick at getting food¹, and always busy. It is wont to be astir by day; yet it is mean in colour, with the belly wet. Of the remaining two the Alba, fair of colour, couples, nests and brings forth well; it feeds in marshes, on a lake, in fields and meadow-ground. But the Stellaris, by-named Lazy (in the fable it is said of old to have been changed from a slave to a bird), as its by-name imports, is slow and indolent. The bird called Phoix has beyond all others this peculiarity that it especially attacks the eyes. The Ardea and the Albardeola, which is of smaller size and has a broad and elongated bill, seek lakes and rivers.

The Pella builds its nest in England on the lofty trees that grow not far from the banks of streams. The upper part of the body is blue, the lower is, however, somewhat white. It routs Eagles or Hawks, if they attack it suddenly, by very liquid mutings of the belly, and thereby defends itself. Of this kind I have seen some white, though they are rare, which differed from the aforesaid neither in their size nor shape of body, but in colour only. Furthermore the white has been observed in England to nest with the blue, and to bear offspring. Wherefore it is clear that they are of one species. I have only once seen—and that was in Italy—the Albardeola, which is called *λευκερωδιός* in Greek; it is much smaller than the Pella and by no means shuns the sight of man so much as does the blue. Had I not seen it, I should have declared the Albardeola to be the English

¹ That is, for its young.

[p. 35]
a buttour
ein pittour.
ein rofdō.

Phoix.

lardam albardcolā esse iudicassē. Stellaris est, quā Angli buttourum, aut bittourum, & Germani pittourum & rofdommum nominant: nam avis est toto corporis habitu ardeis reliquis similis, ex piscium uenatu ad ripas paludium & amnium uiuens, pigerrima & stolidissima, ut quæ in retia ab equo facticio agi potest facilimè. Colore est serè, quantum memini, phasian, rostro limo indito, aspinos ronchos uoce refert: oculos hominum audissimè omnium auium appetit. Quare si quid impediāt, quò minùs stellaris esse possit, (quod mihi nondum cernere datum est) phoica esse oportebit, quam Aristoteles oculos maximè appetere testatur, quam & cæteræ ardeæ idem facere sæpè uisè sunt.

DE AVRIVITTE.

Auriuittis.

Χρυσομήτρης, nō ut quidam codices habent, ῥυσομήτρης, auriuittis, Anglicè a gold finche, Germanicè cyn distelfinck, oder cyn stigeliß.

Auriuittis una est ex auiculis, quæ carduorum semine uictitant¹, & uermes etiam oblatos, non attingunt. Alij goldfincam, aut distelhuincam, spinum, alij carduellem esse uolunt. Sed si quis, ex spinuoris præter hanc aliam aurea uitta redimitam ostēderit, cui magis auriuittis nomen competat, quàm huic, opinionem meam faciliè patiar explodi, alioqui non uideo, quin digna sit, quæ probetur.

[p. 36]

DE ATTAGENE.

Ἀτταγὴν, ἀττάγας, attagen, attagena.

Attagen, ut scribit Aristoteles, gallinagini similis est colore. Attagenam uarijs distinctam esse maculis, Aristophanes² his uersibus testatur:

*Si quis ex nobis erit fugitiuus atq; uisus notis,
Attagen sanè apud nos uarijs appellabitur.*

¹ Aristotle's groups of Birds are as follows: (1) γαυψώνυχες (crooked-clawed); (2) σκωληκοφάγα (worm-eating); (3) ἀκανθοφάγα (eating thistle seeds); (4) σκιποφάγα (? grub-eating); (5) περιστεροειδῆ (dove-like); (6) σχιζόποδα (cleft-footed); (7) στεγανόποδα (wholly webbed); (8) βαρέα (heavy, i.e. ground kinds). A few Birds, however, can hardly be placed under any of these.

² *Aves*, ll. 761—762.

Shovelard¹. Stellaris is that kind which Englishmen denominate buttour or bittour, and the Germans call pittour or rosdorn. Now it is a bird like other Herons in its state of body generally, living by hunting fishes on the banks of swamps and rivers, very sluggish and most stupid, so that it can very easily be driven into nets by the use of a stalking horse. So far as I remember, it is nearly of the colour of a Pheasant, and the beak is smeared with mud; it utters brayings like those of an ass. Of all birds it aims at men's eyes most readily. Wherefore if anything hinders this kind from being the Stellaris (which is not yet given to me to see) it ought to be the Phœnix, inasmuch as Aristotle testifies that it aims chiefly at the eyes, though other Ardeæ also often seem to do the same.

OF THE AURIVITTIS.

Χρυσομήτις (not as some texts have it χρυσομήτης), aurivittis, in English a gold finche, in German cyn distelfinck or cyn stigelitz.

The Aurivittis is one of the small birds that feed on seeds of thistles, and do not touch worms even when offered to them. Some will have it that the Goldfinch or the Distelvinc is but the Spinus², some the Carduelis. But if anyone can shew another of the thistle-eating birds save this, girt with a golden band, to which the name of Aurivittis is more fitting than to this, I gladly will allow my opinion to be ignored, but otherwise, I do not see why it should not be worthy of approval.

OF THE ATTAGEN.

Ἀτταγὴν, ἀττάγας, attagen, attagena.

The Attagen, as Aristotle writes, is like the Gallinago in colour. And Aristophanes bears witness in these lines that the Attagena is marked with varied spots:—

“If any of you be a runaway and branded with the marks, he shall assuredly be called with us the spotted Attagen.”

¹ That is, the Spoonbill of modern books (*Platalea leucorodia*), while the buttour is of course the Bittern (*Botaurus stellaris*)

² Turner himself considered Spinus to be the Greenfinch (cf. p. 85 of the original).

PLINIUS¹ DE ATTAGENE.

Attagen maximè Ionius celebratur, uocalis aliàs, captus uerò obmutescit, quondam existimatus inter raras aues. Iam & in Gallia Hispaniaq; capitur, & per alpes.

PETRUS GYLLIUS².

Attagen, est perdice paulò maior, uersicoloribus picta plumis in dorso, & color ruffus est, uescitur grano, breuib; est alis, & puluerator est.

Falluntur igitur Britannici ludimagistri, qui suū Wodcoccum attagenem faciunt, qui solis uescitur uermibus, & grana nunquam attingit. An attagens apud
 [p. 37] *Anglos inueniantur nec'ne, multum sanè ambigo. nam qui attagenem describunt, marem à fœmina non separant. unde colligo eundem fuisse colorem, & candem figuram maris & fœminæ. Cæterum in hoc auium genere, quod apud nos ad attagenis formam proximè accedit, mas à fœmina ita differt ut duorum generū istiusmodi rerū inperito uideri possint. Vtranque tamen auem describam.*

Mas gallo domestico paulò minor, totus niger est, excepta ea parte caudæ, quæ podicem tegit, ea enim alba est. Cæterū nigredo huius nonnihil splendescit, ad eum ferè modum, quo columborum nigrorum torques circa colla splendescunt. ad uiriditatem igitur proximè accedit. in capite rubrum quendam habet, sed carneum cirrū, & circa genas duos habet ueluti lobos rubros, & eos carneos. Fœmina tota maculis distincta est, & à perdice, nisi maior esset, & ruffa magis, ægrè dignosci

¹ *Hist. Nat. Lib. x. cap. xlviii.*

² Petrus Gyllius was the author of the work *De vi et natura Animalium etc.* Lugd. Bat. 1533.

PLINY OF THE ATTAGEN.

The Attagen is most renowned as an Ionian bird ; it usually is noisy, in captivity however it is dumb. In former times it was considered rare, but now it is taken in Gaul, in Spain, and on the Alps.

PETRUS GYLLIUS.

The Attagen is rather larger than the Perdix, and is marked with particoloured feathers on the back, in colour it is reddish, and it feeds on grain. It has short wings, and rolls itself in dust.

Accordingly our British schoolmasters are wrong who make their Woodcock the Attagen, which lives only on worms and never touches grain. Indeed I seriously doubt whether Attagenes be found in England or not, for those who give descriptions of the Attagen, do not distinguish the male from the female, whence I infer that they have the same colours and are like in form. But in the kind of bird which with us comes the nearest to the Attagen in form, the male differs so greatly from the female that they might appear to be of separate kinds to the man inexperienced in things like this. Nevertheless I will describe each bird.

The male¹ is somewhat less than a domestic cock and is entirely black, save that part of the tail which overlies the vent, for that is white. Moreover the black colour of the bird is somewhat glossy, very nearly as the collar round the neck of our black pigeons is. So it approaches very near to green. Upon its head it has a red but fleshy sort of comb², and round its cheeks two red lobes as it were and those fleshy. The hen is wholly marked with spots, and, were she not a bigger bird and more rufous, could scarcely be distinguished from a Partridge. Both frequent

¹ Turner here undoubtedly refers to the Blackcock (*Tetrao tetrix*).

² The Blackcock has two erectile patches of red skin over the eyes, which in the breeding season even reach above the top of the head ; and the word 'cirrum' must be taken to mean such a patch here.

possit. In desertis locis & planis, erica potissimum confitis, ambo degunt. grano nescuntur, et summis ericæ germinibus. Breues habens alas, & breues faciunt uolatus. Hæc avis, si attagen nō sit, gallina uidetur esse Varronis rustica. Erasmus in Adagio, Attagenæ nonilunium, attagenam auctu palustrem facit, & uarijs maculis distinctam. quod si satis exploratum
 [p. 38] *mihi esset, Anglorum goduittam, siue fedoam, attagenam esse, indubitanter auderem adfirmare. Est autem ipsa gallinagini ita similis, ut nisi paulò maior esset, & pectoris color magis ad cinereū uergeret, altera ab altera difficulter possit distingui. uiuit in locis palustribus, et ad ripas fluminū. rostrum habet longum, sed capta triticum non secus atque columbi, comedit. triplo pluris quàm gallinago apud nos uenditur, tantopere eius caro magnatum palatis arridet. harum si neutra sit attagena, attagenam nusquam uidi.*

DE ATRICAPILLA.

Μελανκόρυφος, atricapilla, Germa. ut creditur, cyn graßmußten.

ARISTOTELES¹.

Atricapillam etiam plurima edere aliqui referunt, sed post Africam strutionem. Iam uel decem & septē oua atricapillæ reperta sunt. sed plura etiam quàm uiginti parit, & numero impari semper, ut narrant. Nidificat ea quoq; in arboribus, & uermiculis alitur. Proprium huius
 [p. 39] & lusciniae præter cæteras aues, ut linguae summæ acumine careant. Ficedulæ & atricapillæ² uicibus commutantur. Fit enim ineunte autumno ficedula, ab autumno protinus atricapilla,

¹ *Hist. An.* Bk IX. 88.

² *Hist. An.* Bk IX. 256—257.

waste open places, and especially those covered with heather. They feed on grain and on the topmost buds of heather. They have short wings and take short flights. This bird, if it be not the Attagen, appears to be Varro's *Gallina rustica*. Erasmus in his proverb of "the Attagena's new-moon" makes his Attagena a marsh-bird, marked with varied spots. If this approved itself sufficiently to me I confidently would venture to affirm that the Attagena was what the English call the Godwitt or Fedoa¹. Furthermore the bird is so much like the Woodcock, that, if it were not a little larger, and did not the breast verge upon ash-colour, the one of them could hardly be distinguished from the other. It is found in marshy places and on river banks. The beak is long; but in captivity it feeds on wheat, just as our Pigeons do. With us it sells for thrice as much again as any Woodcock, so much does its flesh tickle the palates of our magnates. Of these two, if neither be the Attagena, then I have nowhere seen the Attagena.

OF THE ATRICAPILLA.

Μελαγκόρυφος, atricapilla, in German, as is supposed, cyn grasmuklen.

ARISTOTLE.

The Atricapilla, as some report, lays the most eggs of all, next to the Struthio of Africa. No fewer than seventeen eggs of the Atricapilla have been found, but it lays even more than twenty and, as some narrate, in number always odd. It also nests in trees and feeds upon small worms. It is peculiar to this and the Luscinia beyond all other birds that they have no point to the tip of the tongue. Ficedulæ and Atricapillæ change in turn. For when autumn sets in the bird becomes a Ficedula, from autumn onwards it becomes an Atricapilla, nor is

¹ See Prof. Newton's *Dict. Birds*, p. 248.

nec inter eas discrimen aliquod, nisi coloris & uocis est. Auem esse eandem cōstat, quia dum immutaretur hoc genus, utrūque conspectum est, nondum absolutè mutatum, nec alterutrum adhuc proprium ullum habens appellationis¹. Hæc Aristoteles.

Atricapillam in Anglia nunquam uidi, neq. sæpius in uita quàm semel, idq̃, in Italia in domo Ducis Ferrariensis. Eamq̃, mihi uir utriusq̃, linguæ nō uulgariter doctus, D. Franciscus, duci à sacris concionibus exhibuit. Anglorū lingettæ, & Germanorum grafmuscho, quod ad corporis magnitudinem attinet, similis erat: sed atrum habebat caput, & reliquum corporis colorem magis ad cinerium uergentem.

[p. 40]

DE BVBONE.

Búas², bubo, Anglicè, alyke foule, Germanicè eyn
schuffauß / eyn schuffel / eyn fauß.

ARISTOTELES³.

Bubo è noctuarum genere est, & noctuæ specie quidem similis, sed magnitudine non minor quàm aquila.

PLINIUS⁴.

Bubo funebris, & maximè abominatus, publicis præcipuè auspicijs, deferta incolit, nec tantum desolata, sed etiam inaccessa: noctis monstrum, nec cantu aliquo uocalis, sed gemitu. Volat nūquam quò libuit, sed transfuersus auferitur. Hęc Plinius.

Hanc auem semel Venetijs iusta aquilæ magnitudine uidi, sed crura erant paulò breuiora quàm aquilæ crura solent esse. Cætera aquilæ similis erat.

¹ Aristotle has ἐκατέρω ἰδιόν τι ὑπῆρχεν οὐδέν according to one text. He has no word to represent 'appellationis.'

² Or Bpúas.

³ Hist. An. Bk VIII. 39.

⁴ Hist. Nat. Lib. X. cap. xii.

there any difference between the two, save that of colour and of voice. That the bird is the same is evident, since, while this kind is undergoing the change, each of the two is seen, not fully changed as yet, and neither having so far any proper name. Thus far Aristotle.

In England I have never seen the Atricapilla, nor yet but once in life, and that in Italy within the house of the Duke of Ferrara. And Don Francisco of the Holy Council of the Duke, a man uncommonly well versed in both the tongues, shewed it to me himself. The bird was like the English lingett and the German grasmuschen, so far as size of body went; but it had a black head, and the remaining colour of its body verging more to grey.

OF THE BUBO.

Βύας, bubo, in English alyke foule, in German eyn schuffaüss (eyn schüffel), eyn kautz.

ARISTOTLE.

The Bubo is of the race of Noctuæ, and is in aspect very like a Noctua, but not less than an Aquila in size.

PLINY.

The Bubo is a fatal bird, of evil omen beyond other sorts, especially at public auguries; it lives in desert places, and not merely those that are unpeopled, but those also hard of access: monster of the night it utters not a song, but only a groan. It never flies where it intends, but is borne off aslant. So Pliny says.

This bird I saw at Venice once, of a full Eagle's size¹, its legs, however, were a little shorter than an Eagle's legs are wont to be. But it was like an Eagle otherwise.

¹ Turner probably meant the Eagle Owl (*Bubo ignavus*).

[p. 41]

DE BOSCA.

Bosca, avis est aquatica, anati similis, sed minor. Quum multæ sint aves aquaticæ anati similes, sed minores, ut sunt, telæ uocatæ ab Anglis Vuigene & pochardæ, eam puto boscam esse, quæ proximè ad magnitudinem & similitudinem anatis accedit. Hoc quum pocharda faciat, illam Aristotelis esse boscam iudico.

DE CAPRIMVLGO.

Caprimulgus, αἰγιοθήλας.

ARISTOTELES¹.

Caprimulgus, avis est montana, magnitudine paulò maior, quàm merula, minor quàm cuculus, moribus mollior. Parit duo ova, aut tria cùm plurima. Sugit, caprarum ubera aduolans, unde nomen accepit. Cùm uber fuxerit, extingui, capramq; excæcari aiunt, parùm clarè interdiu uidet, sed noctu perspicax.

PLINIUS².

[p. 42] Caprimulgi appellantur grandiores merulæ aspectu, fures nocturni, interdiu etiam uifu carent.

Cùm essem apud Heluctios, senem quendam conspiciatus, capras pascentem in montibus, quos herbas quærendi gratia ascenderam, rogabam num auem nouisset merulæ magnitudine, interdiu cæcam, noctu perspicacem, quæ caprarum ubera noctu sugere solcat, unde capræ postea cæcæ euadunt? qui respōdit, se in Heluctiorum montibus ante quatuordecim annos, multos uidisse, & multas iniurias ab ipsis passum, ut qui semel

¹ *Hist. An.* Bk IX. 109.² *Hist. Nat.* Lib. X. cap. xl.

OF THE BOSCA.

The Bosca is a water fowl, like to a Duck¹, but smaller. Since there are many water fowls like to the Duck, but smaller (as there surely are), called Teles, Wigenes and Pochards by the English, I believe the Bosca to be that which comes nearest the size and likeness of a Duck. And since the Pochard does so, I decide that it is Aristotle's Bosca.

OF THE CAPRIMULGUS.

Caprimulgus, αἰγροθήλας.

ARISTOTLE.

The Caprimulgus is a mountain bird, in size a little bigger than the Merula, less than the Cuculus; in disposition it is milder. It lays two eggs or three at most. Flying to the udders of she-goats, it sucks them, and thus gets its name. They say that the udder withers² when it has sucked at it, and that the goat goes blind. By day the bird sees dimly, but quite well at night.

PLINY.

Caprimulgi, as they are called, look bigger than a Merula, and act as thieves by night; by day they even lack the power of sight.

When I was in Switzerland I saw an aged man, who fed his goats upon the mountains, which I had gone up intent on search of plants: I asked him whether he knew a bird of the size of a Merula, blind in the day-time, keen of sight at night, which in the dark is wont to suck goats' udders, so that afterwards the animals go blind. Now he replied that he himself had seen many in the Swiss mountains fourteen years before, that he had suffered many losses from those very

¹ That is, the Wild Duck (*Anas boscas*).

² This rendering appears much preferable to translating this word 'the goat dies,' as, judging from the punctuation, some would have it.

sex capellas à caprimulgo occæcatas habuerat. cæterū nūc omnes ad unum ab Heluetijs usque ad inferiores Germanos, ubi hodie non solum capras lacte priuant & occæcant, sed & oves insuper occidunt, auolasse. Nomen auis quærenti, paphum, id est, sacerdotem dici respōdit. Sed uctulus ille mecū fortè iocatus est. Ego uerò, siue iocatus fuerit, siue seriò locutus, aliud Germanicū caprimulgi nomen quā quod me docuit iste, non teneo. Si qui sint, qui melius & aptius nomen in prōptu habeant, proferant.

DE CARDVELE.

Carduelis, si Gazæ credimus, Græcè θραυπίς dicitur, [p. 43] & inter spiniuoras auiculas Aristoteles recenset. Nec plura de carduele apud Aristotelem lego. Plinius¹ scribit cardueles auium minimas imperata facere, nec uoce tantum, sed pedibus & ore pro manibus.

Præter auiculam illam spiniuoram aurea uitta redimitam, aliam noui spiniuoram colore uiridem, quæ non secus atq; auriuittis rostro è duabus situlis uicissim ascendentibus & descendentibus, cibum ex una, & potū ex altera desumit. Quin & hoc facit miliaria, quam linotam nostrates appellant. Eadē quoq; hominē quiduis cātantē, uoce imitatur. Quare nō sola illa, quæ Græcè θραυπίς, & Latinè Theodoro carduelis dicitur, imperata facit, & rostro & pedibus pro manibus utitur.

Dictū mihi difficile uidetur, quamnam è tribus, quum omnes illæ carduorum semine uescantur, Plinius carduelem fecerit, nū thraupin aut acanthin, aut chry-

¹ *Hist. Nat. Lib. x. cap. xlii.*

birds; so that he had once had six she-goats blinded by Caprimulgi, but that one and all they now had flown away from Switzerland to Lower Germany, where nowadays they did not only steal the milk of she-goats, making them go blind, but killed the sheep besides. And, on my asking the bird's name, he said that it was called the Paphus, otherwise the Priest. But possibly that aged man was jesting with me. Yet whether he was jesting, or spoke gravely, still I have no other German name than what he gave to me for Caprimulgus. If there be any then who have in readiness a better or a fitter name than this, let them produce it.

OF THE CARDUELIS.

The Carduelis, if we believe Gaza, is in Greek called *θραυπίς*. Aristotle also numbers it among small thistle-eaters¹. I find in Aristotle nothing more than this about the Carduelis. Pliny writes that Cardueles, smallest of all birds, perform set tasks, and not in song alone, but with their feet and beak in place of hands.

Besides that thistle-eating little bird² adorned with band of gold I know another thistle-eating sort, in colour green³, which with its beak takes up its food from one of two small buckets moving up and down alternately, its water from the other, as the Aurivittis does. The Miliaria moreover does the same, which bird our countrymen call Linot. Furthermore it mimics with its song a man when singing anything. And so it is not only that one kind, in Greek called *θραυπίς* and in Latin named by Theodorus⁴ Carduelis that performs the tasks that it is bid, and uses beak and feet in place of hands.

It seems to me then difficult to say, since all three birds feed upon thistle-seeds, which of them Pliny meant by Carduelis, whether it should be the Thraupis, or the Acanthis, or the Chrysomitris. And should it be the Thraupis, as

¹ See p. 35.

² The Aurivittis, p. 35.

³ Probably Turner means the Siskin (*Carduelis spinus*).

⁴ Theodorus Gaza.

somitř. Si thraupin, ut Gaza credit, avis illa aureis plumis carduelis non erit: nam illa apud Aristot. nō θραυπίς, sed Chrysomitris est. Quæ nā igitur avis carduelis sit, non audeo pronunciare.

DE CŒRVLEONE.

Κυανός, cœruleo, Anglicè, a clotburd, a smatche, an arlyng, a sleincheh, German. eyu brech uögel.

[p. 44]

ARISTOTELES¹.

Cœruleo maximè in Scyro² colit, faxa amans: magnitudine minor quàm merula, maior paulò quā fringilla: pede magno³ est, scanditq; faxa: colore cœruleo: rostro tenui & longo: crure breui, similiter ut pipo est.

Cœruleo, si ea sit avis, quam conijcio esse, in cuniculorum foveis, & sub lapidibus in Anglia nidulatur, & in hyeme non apparet.

DE CERTHIA.

ARISTOTELES⁴.

Certhia, auicula est exigua, cui mores audaces, domicilium apud arbores, uictus ex coffis, ingenium sagax in uitæ officijs.

Avis est quædam, quam Angli creperam, id est, reptitatricem nominant, quòd super arbores semper repat, quam certhiam esse credo. Ea regulo paulò maior, pectore pallido, cætera fusca & maculis nigris distincta

¹ *Hist. An.* Bk IX. 97.

² Another reading is Νισύρω.

³ μεγαλόπους, but some read μελανόπους.

⁴ *Hist. An.* Bk IX. 92.

Gaza believes, that bird with golden feathers will not be the *Carduelis*, for with Aristotle that is not the *θραυπίς*, but the *Chrysomitris*. Therefore I dare not pronounce what bird the *Carduelis* is.

OF THE CŒRULEO.

Κυανός, cœruleo, in English a clot-burd, a smatche, an arlyng, a steinchek, in German eyn brech vögel.

ARISTOTLE.

The Cœruleo chiefly dwells in Scyros and loves rocks; in size it is a little smaller than a *Merula*, a little larger than a *Fringilla*: it has large feet, and climbs on rocks: in colour it is blue: the beak is long and thin: the legs are short as in the *Pipo*.

The Cœruleo¹, if it be the bird which I conjecture, nests in rabbit holes and under stones in England, and does not appear in winter.

OF THE CERTHIA.

ARISTOTLE.

The *Certhia* is a very little bird of bold habits; its home is upon trees, its food is grubs; it shews wise instinct for the needs of life.

There is a certain bird which Englishmen call Creeper, that is Climber, for it always climbs about on trees: this I believe to be the *Certhia*. It is a little bigger than the *Regulus*, having a whitish breast, the other parts dull brown, but varied with black spots; its note is sharp, its

¹ Whatever bird Aristotle's may be, Turner's is certainly the Wheatear. Belon and Gesner seem to think that the former is the Blue Thrush, but Sundevall is certainly wrong in suggesting that it is the Wall-Creeper (*Tichodroma muraria*).

[p. 45] *est, uoce acuta est, & rostro' tenui, & leuiter in fine adunco, nunquam quiescit, sed semper per arbores picorum more scandit, & coffos è corticibus eruens, comedit.*

DE CICONIA.

Πελαργός, ciconia, Anglicè a stork, Germanicè eyn storck, & Saxonice eyn ebeher.

Ciconia, ut Germanis auis est notissima, ita Britannis meis plerisq: omnibus tam ignota est, quàm quæ omnium ignotissima. Nec mirum, quum nusquam in insula nostra nisi captiua Ciconia uideatur. Apud Germanos in summis tectis, aliquando in ipsis summis fumarijs nidulatur. Auis est mediæ magnitudinis inter gruem & ardeam, pennis albis & nigris distincta: crura longa habet, rostrum gruinam, sed rubrum & crassum: circa lacus & paludes degit, ranas, bufones, angues, & pisces comedens.

PLINIUS DE CICONIIS¹.

Ciconiæ, quó'nam è loco ueniant, aut quò se referant, incompertum adhuc est. E longinquo uenire, non dubium, eodem, quo grues modo, illas hyemis, istas æstatis aduenas. Abi-
[p. 46] turæ congregantur in loco certo: comitatæq; sic ut nulla sui generis relinquatur, nisi captiua & ferua, ceu lege prædicta die recedūt. Nemo uidit agmen discedentium, cùm discessurum appareat, nec uenire, sed uenisse cernimus. Vtrumque nocturnis fit temporibus. Ciconiæ nidos eosdem repetunt, & geneticum senectutem educant.

DE CINCLO.

Κίγκλος, ἡ σεισπορυγίς, Anglicè a uuater suuallouu, Germanicè eyn steynbijsjer.

¹ *Hist. Nat. Lib. x. cap. xxiii.*

beak is slender and is slightly hooked towards the tip; it never rests, but is for ever climbing up the trunks of trees after the manner of the Woodpeckers, and it eats grubs, picking them from the bark.

OF THE CICONIA.

Πελαργός, ciconia, in English a stork, in German eyn storck, and in Saxon eyn ebeher.

The Stork, though one of the best known of birds among the Germans, is to nearly all my countrymen of Britain as unknown as the most unknown bird. And little wonder since a Stork is nowhere to be seen, save as a captive, in our island. With the Germans the bird nests upon roofs or even chimney tops at times. It is a bird of middle size between a Crane and a Heron and distinguished by feathers of black and white: it has long legs and a beak like a Crane's, but red and stout; it lives near lakes and marshes, eating frogs, toads, snakes and fishes.

PLINY ON CICONIÆ.

From what parts the Ciconiæ may come, or whither they betake themselves, is not yet ascertained. It is indubitable that they come from far in the same manner as the Grues, but the former arrive in winter and the latter in summer. When ready to depart these birds collect at some fixed place, and after gathering, so that none of their tribe, unless a prisoner or a slave, is left behind, they disappear, on an appointed day, as if by law. No one has ever seen the whole array in very act to go, though it may haply shew itself when ready to depart; nor do we view it coming, but when it has come. Both these events take place at night. Ciconiæ seek the same nests again, and tend their parents in old age.

OF THE CINCLUS.

Κίγκλος, ἡ σεισοπυγίς, in English a water swallow, in German eyn steynbisser.

ARISTOTELES¹.

Cinclus ex mari & fluuijs uictū petit. astutus est, & captu difficilis: sed captus, omnium maxime mitescit. Lēsus hic est, incontinens enim parte sui posteriore.

Auicula, quā ego cinclū esse puto, galerita paulò maior est, colore in tergo nigro, uentre albo, tibijs longis, & rostro neutiquam breui: uere circa ripas
 [p. 47] *fluminum, ualde clamosa est & querula, breues & crebros facit uolatus.*

DE CHALCIDE.

ARISTOTELES².

Chalcis rarò apparet, montes etenim incolit. nigro colore est, magnitudine accipitris, quem palumbarium nominant: forma longa ac tenui, Iones cymindem appellant. cuius Homerus etiam meminit in Iliade cū dicit:

*Chalcida dij perhibens, homines dixêre cymindem*³.

Sunt, qui eandem hanc auẽ non aliam esse atque ptyngem uelint. Interdiu minùs apparet, quia non clarè uidet, sed noctu uenatur, more aquilæ: pugnat uerò cum aquila adeò acriter, ut fæpius ambæ implexæ, deferantur in terram⁴, & uiuẽ à pastoribus capiantur. Parit hæc oua duo, & faxis speluncisq; nidulatur.

¹ *Hist. An.* Bk IX. 76.

² *Hist. An.* Bk IX. 79—80.

³ *Iliad*, Bk XIV. l. 291. 'Perhibens' is a misprint for 'perhibent.'

⁴ Aristotle has not these five words.

ARISTOTLE.

The Cinclus seeks its food from the sea and from rivers. It is cunning and is hard to catch, but grows the gentlest of all birds when caught. It is moreover maimed, being without control over its hinder parts.

The little bird which I believe to be the Cinclus¹ is a little bigger than the Galerita, with black colour on the back, and a white belly; while it has long shanks and a bill by no means short: in spring it is exceeding clamorous and querulous about the banks of rivers, where it takes short and incessant flights.

OF THE CHALCIS.

ARISTOTLE.

The Chalcis is not often seen, for it haunts mountains. It is of a black colour, and of the size of the Accipiter which they name Palumbarius. Its form is long and slender; the Ionians call it Cymindis. Of it furthermore Homer makes mention in the Iliad, wherein he says:—

The Gods know it as Chalcis, men say Cymindis.

Some there are who would make this very bird none other than the Ptynx. By day it shews itself but little, since it does not see clearly, although it hunts by night, after the manner of the Aquila; yet with the Aquila so keenly does it strive, that frequently both birds fall to the ground clutching each other, and are caught alive by shepherds. It lays two eggs, and nests in cliffs and caves².

¹ Turner evidently means the Common Sandpiper (*Actitis hypoleucos*).

² Sundevall says that Klb thought that this bird was the Hawk Owl, but himself refers it doubtfully to the Glossy Ibis (*Plegadis falcinellus*), which however is not a night bird and does not live on hills. Belon and Gaza thought that it was an owl of some kind.

[p. 48]

DE COLLVRIONE.

Κολλυρίων, *collurio*, Anglicè, *a feldfare* aut *a feldfare*. Quibusdā German. *eyn frammesvögel*.

ARISTOTELES¹.

Collurio ijsdem, quibus merula uestitur, magnitudo eius eadem quæ superioribus, id est, uireoni & mollicipiti², capitur potius hyberno tempore.

Auis, quam collurionem esse puto, turdum magnitudine æquat, sed caudam habet longiorem, & magis mobilem, & pectus maculosum. In æstatē apud nos aut rarō aut nunquam uidetur: in hyeme uerō tanta copia est, ut nullius auis maior sit. baccis aquifoliæ arboris, sorbi minimæ, & similium arborum uestitur. gregatim uolat, & inter uolandum obstrepera est.

DE COLVMBIS.

Περιστερά, *columba*, Anglicè *a doue*, Germanicè *eyn taube*. Saxonice *eyn duue*.

ARISTOTELES³.

Columbacei uerò generis plures species sunt. Est enim liuia à liuore dicta, diuersum certè à columba genus, quippe minor quàm columba sit, & minùs patiens mansuescere: liuet enim plumis, & penè nigricat, & pedibus rubris scabrosisq; est. Quas ob res, nullus id genus cellare alit. Maximo inter hæc genera sunt corpore palumbes. Secundum magnitudinis locum uinago obtinet, paulò maior quàm columbus⁴ est. Minimum ex his turtur est. pariunt columbæ omnibus anni temporibus, pullosq; educant si locum apricum habeāt & cibum. Sin minùs,

Liuia.

[p. 49]

Palumbes.

Vinago.

Turtur.

¹ *Hist. An.* Bk ix. 99.

² These five words are not represented in Aristotle. Moreover the *πάρδαλος* is here omitted, and is placed later (p. 107 of the original work), but there seems to be some doubt as to the correct reading in the Greek.

³ *Hist. An.* Bk v. 43. The readings vary in places, but the rendering is decidedly free.

⁴ Gaza and Turner make indiscriminate use of 'Columbus' and 'Columba' for the same kind of bird.

OF THE COLLURIO.

Κολλυρίων, collurio, in English a feldfare or a feldefare. According to some Germans eyn krammesvogel.

ARISTOTLE.

The Collurio feeds on the same meat as the Merula. Its size is that of the aforesaid kinds—that is, the Vireo and the Mollicept—it is caught chiefly in the winter time.

The bird which I consider to be the Collurio equals a Thrush in size, but has a longer and more flirting tail, also a spotted breast. Rarely or never is it seen with us in summer: yet its plenty is so great in winter that of no kind is there more. It eats the berries of the Holly, the Least Service, and like trees. It flies in companies, and on its flight is very noisy.

OF DOVES.

Περιστερὰ, columba, in English a dove, in German eyn taube, in Saxon eyn duve.

ARISTOTLE.

Of the Dove-kind, however, there are many sorts. For first there is the Livia, named from its livid colour, which is certainly a different kind from the Columba, inasmuch as it is smaller and less ready to be tamed: and it is livid in its plumage, verging upon black, and has moreover red and roughened feet. Wherefore nobody keeps this kind in cotes. Of greatest size among the several sorts are the Palumbes; the Vinago holds the second place herein, a little bigger than the Columbus. The smallest of them is the Turtur. The Columbæ breed at all times of the year, and rear their young, if they have but a sunny place and food. If otherwise they breed

æstate tantummodo fœtant. Sed proles præstantior uere est, quàm autumnino, deterrima æstate, & omni tempore calidiore.

De ijsdem in alio libro¹ ad hunc modum scribit,

[p. 50] Alia frugibus uiuunt, ut palumbes, columbus, uinago, turtur. Visuntur semper columbi, atque palumbes, sed turtur æstate tantū, hyeme se condit, latitat² enim suo tempore. Vinago autumnino & conspicitur, & capitur, cui magnitudo maior, quàm columbo, minor quā palumbi est.

Liua. Πελειάς, quæ liuia Latinè dicitur, est syluestris illa columba, quam Angli a stocdoue, & Germani cyn holts taube nominant.

Palumbes. Φάττα, Latinè palumbes siue palumbus, dicta ab Anglis a couchot a ringged doue, & à Germanis cyn ringel taube appellatur. Hæc longè aliter atque liuia nidulatur. Nidificat autem liuia in cauis aliquando arboribus, interdum & in templorum muris. Palumbes uerò in condensa hedera, aut super ramum arboris ex pauculis ligniculis transuersim positis, tenuissimum nidum construit. Quòd si quis mihi parùm hac in re fidat, torquatos columbos esse palumbes ueterum, Aristotelem diligentius legat, & Martialem³ poetam de ijsdem ita scribentem audiat,

[p. 51] Inguina torquati, tardant hebetantq; palumbi
Non cdat hanc uolucrem, qui uolet esse salax.

Politianus⁴ de palumbis ita scribit :

Dum sua torquatæ repetunt dictata palumbes.

Τρυγών.
Turtur.

Turtures in Germania sunt multò frequentiores quā in Anglia. Turturem Angli & Saxones communi uocabulo turtel diuine nominant.

¹ *Hist. An.* Bk VIII. p. 45.

² The Greek is φωλεί γάρ. But how far Aristotle referred to birds 'hibernating' is very doubtful.

³ *Epigr.* Lib. XIII. lxvii.

⁴ A poet and scholar of the Renaissance.

only in summer. Yet in spring the young are better than in autumn, they are worst of all in summer and at every hotter season.

Of the same he writes in another book after this manner:—

Other birds live on crops, as the Palumbes, the Columbus, the Vinago and the Turtur. The Columbi and Palumbes may be always seen, the Turtur only in the summer. In the winter it lies hid, for it conceals itself at the due time. But the Vinago is both seen and caught in autumn, of which bird the size is greater than that of the Columbus, but less than that of the Palumbes.

Πελειῶς, which in Latin is called Livia, is that dove of the woodlands which the English name a stocdove, and the Germans *eyn holtztaube*.

Φάττα, in Latin Palumbes or Palumbus, is called by Englishmen a Coushot or a Ringged Dove, and by Germans named *eyn ringel taube*. It nests far otherwise than does the Livia, for that bird sometimes breeds in hollow trees and sometimes even in the walls of churches. But the Palumbes builds a nest of the frailest of a few small twigs laid crosswise in a mass of ivy or upon a bough. Now in this thing if there be anyone who places little confidence in my opinion that our collared doves are the Palumbes of the ancients, let him read with greater care his Aristotle and give ear as well to Martial the poet writing thus of the same birds:—

Ringed doves make a man's loins slow and dull;
Who would be lusty should not eat this bird.

Politian writes of the Palumbi thus:—

While ringed doves seek again their accustomed haunts.

Turtle Doves are much more plentiful in Germany than in England. English and Saxons in common call it *turtel duve*.

Vinago.

Oivás, quæ Latinè uinago dicitur, mihi nunquã hæcenus uisa est, nec quid habeat nominis apud nostros, aut apud Germanos compertum habeo. Vidi tamen Venetijs columbos hijs nostratibus sesquialtera portione maiores: sed hos non uinagines fuisse credo, sed columbos è Campania ad Venetos aduectos, ubi Plinius columbos scribit esse grandissimos.

DE COTVRNICE.

ὄρνις, coturnix, Anglicè a quale, Germanicè eyn machtel.

PLINIUS¹.

Coturnix parua auis, & cùm ad nos uenit, terrestres potiùs quàm sublimis. Aduolant & hæ simili modo, quo grues & ciconiæ, non sine [p. 52] periculo nauigantium, cùm appropinquauere terris. Quippe uelis sæpè insidunt, & hoc semper noctu, merguntq; nauigia. Coturnicibus, ueratri, siue ut alij legunt, ueneni: semen gratissimus cibus. quam ob causam eas damnaue mensæ, simulq; comitalẽ propter morbum despui fuetum, quem solę animalium sentiunt præter hominem.

Quæ, quum ita se habeant, demiror quis malus genius Britannis meis in mentem immisit, ut eas tantopere in deliciis habeant, quum tot malis, ueneno scilicet, et comitali morbo, illarum caro sit obnoxia. Coturnix perdici similis est: sed multis partibus minor. Coturnix, ut scribit Aristot. hoc sibi peculiare & proprium uindicat, ut & ingluuiem, & gulam propè uentriculum amplam & latam habeat.

¹ *Hist. Nat. Lib. x. cap. xxiii.*

Olvás, in Latin called *Vinago*, has never met my eye up to this time, nor have I yet found out what name it bears among our countrymen or among Germans. But I have seen doves in Venice half as big again as those of our own land, although I do not think that they could be *Vinagines*, but birds brought to those parts out of Campania, where Pliny notes the Doves to be exceeding large.

OF THE COTURNIX.

ὄρνις, coturnix, in English a quale, in German eyn wachtel.

PLINY.

The Coturnix is a little bird, and, when it comes to us, keeps on the ground more than aloft. Yet it flies hither just as *Grues* and *Ciconiæ*, not without danger to sea-faring men, when they approach the land. For these birds often settle on the sails, and that always at night, and so sink ships. The seed of *Veratrum*, or, as others read, *Venenum*, is a very grateful food to the Coturnices, and for this cause men have condemned them for the table; furthermore it is the custom for them to be spurned on account of the falling sickness, to which, they alone of animals, save man, are subject.

Now since these things are so, I marvel much what evil genius put it into the mind of my fellow Britons to esteem them thus among their delicacies, when their flesh is liable to ills so many, namely poison and the falling sickness. The Quail is like the Partridge, although many times smaller. As Aristotle writes, it claims a property peculiar to it of having both crop and gullet large and wide near to the stomach.

DE CORNICE.

Κορώνη, *cornix*, Anglicè *a crouu*, Germanicè *eyn frae*, & *eyn fraeg*. *Cornix* avis est omniuora, nam carnes, pisces, & grana interdum uorat, circa littora maris, & ripas fluminum multùm uersatur, ut ea animalia, quæ [p. 53] unda eiecit, tangat. *Cornix* tota nigra est. & media magnitudine inter monedulam & coruum.

Est & marina quædam cornix, quam aliqui *hybernæ* cornicem uocant, capite, cauda, & alis nigris, cætera cineria: an hanc aliquando uiderint *Aristoteles* & *Plinius*, dubito: nam de ea nusquam mentionem fecerunt. Supereſt adhuc & alia *cornix* graniuora, rostro albo, cætera nigra. Hæc σπερμολόγον, id est, frugilegam *Aristotelis* Longolius esse coniecit.

DE CORVO.

Κόραξ, *coruus*, Anglicè *a raucn*, Germanicè *eyn rabe*. *Coruus*, quum sit avis cornice maior, tota nigra & carniuora, omnibus fatis notus est. *Corui* locis arctioribus & ubi fatis pluribus non sit, duo tantùm incolunt, & suos pullos cum iam potestas uolandi est, primùm nido eijciunt, deinde regione tota expellunt. Parit coruus quatuor aut quinque.

DE CVLICILEGA.

Κνιπολόγος, *culicilega*, Anglicè *a uuagtale*, Germanicè *eyn wasser stels*, *eyn quiffsterß*.

ARISTOTELES¹.

Culicilega, magnitudine est quæta spinus, [p. 54] colore cinerea, distincta maculis, uoce parua, quæ & ipsa lignapeta² est.

Culicilegam esse iudico auiculam, quam aliqui *môtacillam* nuncupant. est autem illa albo & nigro uariè

¹ *Hist. An.* Bk VIII. 44.

² The Greek is ἔστι δὲ καὶ τοῦτο ξυλοκόπον.

OF THE CORNIX.

Κορώνη, cornix, in English a crow, in German eyn krae and eyn kraeg.

The Crow¹ is an omnivorous bird, for it eats flesh and fish and sometimes grain; it much frequents sea-coasts and river-banks, that it may there obtain those animals which the tide has thrown up. The Crow is wholly black and is midway in size between a Daw and a Raven.

There also is a certain Sea Crow, which some call the Winter Crow², with black head, tail, and wings and the remainder grey: but whether Aristotle or Pliny ever saw this bird I am uncertain, for they have not mentioned it in any place. There still remains another Crow³, a grain-eater, with white beak, but black otherwise. Longolius conjectured this to be Aristotle's σπερμολόγος, that is frugilega.

OF THE CORVUS.

Κόραξ, corvus, in English a raven, in German eyn rabe.

The Raven, inasmuch as it is bigger than the Crow, quite black, and a flesh-eater, is sufficiently well known to all. In places with less space, and where there is not room for many, Ravens dwell only in pairs, and, when their young have just gained power of flight, the parents first banish them from the nest, and later drive them out of the whole neighbourhood. The Raven has a brood of four or five.

OF THE CULICILEGA.

Κυπολόγος⁴, culicilega, in English a wagtail, in German eyn wasser steltz, eyn quikstertz.

ARISTOTLE.

The Culicilega is a bird of the same size as the Spinus, ash-coloured, and marked with spots: its voice is poor; moreover it pecks wood.

The Culicilega I judge to be that little bird, which some name Motacilla, inasmuch as it is variously marked with

¹ The Carrion Crow (*Corvus corone*).

² The Hooded or Grey Crow (*Corvus cornix*).

³ The Rook (*Corvus frugilegus*).

⁴ Sundevall thinks that this bird is *Certhia familiaris*.

distincta, cauda longa, quam semper motitat. degit plurimum ad ripas fluminum, ubi muscas captat & uermiculos, quin & aratrum uermium causa sequitur, quos uersat & exhibet cum gleba aratrum.

DE CVCVLO.

Κόκκυξ, *cuculus*, Anglicè *a cukkouu*, & *a gouke*, Germanicè *cyn fuffuf*.

ARISTOTELES¹.

Cuculus ex accipitre fieri, immutata figura, à nonnullis putatur: quoniam quo tempore is apparet, accipiter ille, cui similis est, non aspicitur. Sed ita ferè euenit, ut ne cæteri item accipitres cernantur cùm primam uocem emisit cuculus, nisi perquàm paucis diebus. Ipse autem breui tempore æstatis uisus, hyeme nō [p. 55] cernitur. Est hic neque aduncis unguibus, ut accipiter, neq; capite accipitri similis: sed ea utraque parte columbum potiùs quàm accipitrem repræsentat. Nec alio quàm colore imitatur accipitrem, nisi quòd maculis distinguitur, ceu lineis, cuculus uelut punctis. Magnitudo atq; uolatus similis accipitrũ minimo, qui magna ex parte non cernitur per id tempus, quo cuculus apparet. Nã uel ambo unà uisĩ aliquando sunt. Quin etiam ab accipitre interim² cuculus uisus est, quod nulla auis suo in genere solet facere. pullos cuculi nemo ait se uidisse. parit tamẽ, uerũ non in nido, quem ipse fecerit, sed

¹ *Hist. An.* Bk VI. 41—44.

² Aristotle has *καρσθιόμενος*.

black and white, and it has a long tail, which it is always jerking. It mostly haunts the banks of rivers, where it catches flies and little worms; moreover it follows the plough for the sake of the worms which are turned up and laid bare with the clod.

OF THE CUCULUS.

Κόκκυξ, cuculus, in English a cuckow, and a gouke, in German cyn kukkuck.

ARISTOTLE.

By some the Cuculus is thought to come by change of form from an Accipiter¹, since, at the season when the former appears, the Accipiter which it resembles is not seen. But commonly it so falls out that the other Accipitres are likewise absent when the Cuculus utters its earliest cry, save for a very few days. Further the bird itself is only seen for a short time in summer; it is not observed in winter. Nor has it the claws hooked as an Accipiter, nor yet a head like an Accipiter: but in both of these parts it counterfeits a Columbus rather than an Accipiter. In naught but colour does it imitate the Accipiter, except that in its marks, it is distinguished as it were by lines, the Cuculus by spots. The size and mode of flight are like those of the least of the Accipitres, which for the most part at the time wherein the Cuculus appears, is not to be observed. Yet on occasion both have been seen at once. The Cuculus, moreover, has been known to be struck down by the Accipiter, which thing no bird is ever wont to do to one of its own kind. Nobody says that he has seen young of the Cuculus, and yet it breeds, although not in a nest which it has made itself: but sometimes

¹ Such a tradition is still common in many parts of this country and on the Continent.

interdum in nidis minorum auium, & oua, quæ aliena reperit, edit: maximè uerò nidos palū-
 [p. 56] bium petit, quorum & ipforum oua esu absumit, sua relinquens: parit maiori ex parte singula oua, rarò bina. Curucæ quoque in nido parit, fouet illa & excudit & educat. Quo quidem præcipuè tempore¹ & pinguis & grati saporis pullus cuculi est. Genus eorum quoddam nidos facere procul in petris excelsis, præruptisq; affolet.

Curuca a
 titlyng.

Cuculum hîc nobis satis graphicè depinxit Aristoteles, si curucam eadem diligentia descripsisset, non fuisset hodie tam omnibus ferè incognita quàm nunc est. Ego suspicor Anglorum titlingam esse curucam Aristotelis. Nam nullam auem in uita frequentius cuculi pullum sequentem, & pro suo educantem, quam illam obseruari. Est autem illa luscinia minor, sed eadem corporis figura, colore subuiridi, culices & uermiculos in ramis arborum scclatur, rarò humi consistit, hyeme non cernitur.

DE CRECE EX ARISTOTELE².

[p. 57] Sed cū omnibus quaterni digiti sint, tres parte priore habentur, unus parte posteriore pro calce, ut tute sit, qui minutus inest ijs, quæ longa habent crura, ut in crece euenit³. Est⁴ autem crex moribus pugnacibus, ingenio ualens ad uictum, fed cætera infœlix.

Est auis quædam apud Anglos, longis cruribus, cætera coturnici, nisi quod maior est, similis, quæ in segete & lino, uere et in principio æstatis non aliam

¹ This apparently means 'when in the nest.' How then does Aristotle say that 'no one has ever seen the young'? The passage may be an interpolation, as may be another which follows referring to Hawks. If so, the fact of nesting on rocks may also refer to Hawks, and be a further interpolation.

² *Hist. An.* Bk II. 46.

³ 'ut...euenit.' These words are not in Aristotle.

⁴ *Hist. An.* Bk IX. 91.

in the nests of smaller birds, and it devours the eggs of the others that it finds. It mostly seeks the nests of the Palumbes and eats those birds' eggs, leaving its own behind. For the most part it lays a single egg or rarely two. It also lays in the Curuca's nest, and that bird sits upon the eggs, hatches and rears the young. And at that time indeed the offspring of the Cuculus is both particularly fat and of a grateful flavour. A certain kind of Cuculus is wont to make its nest far off on steep and very lofty rocks.

Here Aristotle has portrayed the Cuculus to us most graphically, and, had he described the Curuca in the same careful way, it would not at this day have been so little known to almost everyone as now it is. The Curuca of Aristotle I suspect to be the Titling¹ of the English. For I have observed no other bird in life more frequently than this following the Cuckow's young and rearing it, as though its own. Now it is less than the Luscinia, but with the same figure of body, and in colour somewhat green; it hunts for gnats and little worms among the boughs of trees. It seldom settles on the ground, and is not seen in winter.

OF THE CREX FROM ARISTOTLE.

But seeing that all birds have four toes each, three are directed forwards and one backwards by way of a heel, for safety's sake; the last is very small in such as have long legs, as happens with the Crex. Further the Crex is of pugnacious habit, clever in procuring food, but of bad omen otherwise.

There is a certain bird in England with long legs, otherwise like a Quail, except that it is bigger, which in spring as well as early summer makes no other cry among the corn and flax

¹ It is impossible to say with certainty what Turner's 'Titlyng' was; but probably he meant the Tree-Pipit, which he confounded with the Titlark.

habet uocem, quàm crex crex: hãc enim uocem semper ingeminat, quam ego Aristotelis creccm esse puto. Angli auem illam uocant a daker hen, Germani eyu schryf. nusquam in Anglia nisi in sola Northumbria uidi & audiui.

DE DIOMEDEIS AVIBUS EX PLINIO¹.

Nec Diomedæas præteribo aues. Iuba catarractas uocat, & eis esse dētes, oculosq; igneo colore, cætera cãdidos tradēs. Duos semper [p. 58] ijs duces, alterum ducere agmen, alterum cogere. Scrobes excauare rostro, inde crate cōsternere, & operire terra, quæ antè egesta fuit, in his fœtificare. Fores binas omniũ scrobibus, orientem spectare, quibus exeant in pascua, † occafum † occidentem, quibus redeant. Aluum exoneraturas subuolare semper, & contrario flatu. Vno hæ in loco totius orbis uisuntur, in insula, quam diximus nobilem Diomedis tumulo, atque delubro, contra Apulię oram, fulicarum similes. Aduenas barbaros clangore infestant. Græcis tantũ adulantur, miro discrimine, uelut generi Diomedis hoc tribuētes, ædemq; eius quotidie pleno gutture madētibus pennis perluunt.

DE FICEDVLA.

[p. 59] Συκαλὶς, *ficedula* Latinè dicta, non est Germanorum *sneppa*, quæ locis gaudet humidis, & solis uer-

¹ *Hist. Nat. Lib. x. cap. xliv.*

than *crex crex*, and moreover it repeats this sound incessantly; I think that it is Aristotle's *Crex*. This bird the English call a Daker Hen, and the Germans *eyn schryk*¹. I have not seen or heard it anywhere in England, save in Northumberland alone.

OF THE AVES DIOMEDEÆ FROM PLINY.

And I will not omit the birds of Diomedæ², which Juba calls *Cataractæ*, telling us that they have teeth and fire-coloured eyes, but otherwise are white. They always have two captains, one to lead the band, the other to bring up the rear. These birds dig furrows with the beak, then cover them with wattlework, and hide this with the earth thrown out at first; in these places they breed. Each furrow has two openings, one facing east, by which they may go out towards their feeding grounds, the other facing west, by which they may return. They always flutter out to disburden the belly, and against the wind. In one place only of the whole world are they to be seen, namely that island which we have set down as famous for the tomb and shrine of Diomedæ, over against the shore of Apulia. They are like *Fulicæ*. Strangers who come there they attack with clamour, only on the Greeks they fawn, with wonderful discernment, paying as it were this tribute to the race of Diomedæ, and every day they purify his shrine with brimming throats and water-laden wings.

OF THE FICEDULA.

Συκαλὶς, in Latin called *ficedula*³, is not the *sneppa* of the Germans, which delights in wet localities, and feeds only

¹ Schlegel (*Vog. Nederl.* 11. 60) says that the Dutch *schriek* is the Water Rail (*Rallus aquaticus*), but Turner evidently means the Corn Crake (*Crex pratensis*). Naumann (*Naturg. Vög. Deutschl.* ix. p. 496) gives *Schrecke* as a local name for the Corn Crake.

² Apparently Shearwaters of some species are meant. For the story see any work on Mythology.

³ For the supposed change of *Ficedula* into *Atricapilla*, see p. 39.

mibus uestitur: sed auicula Germanorum graspmuscho similis, ficubus & uuis uictitās, ut pulchrè his uersibus Martialis¹ testatur:

*Cùm me ficus alat, & pascar dulcibus uuis,
Cur potiùs nomen non dedit uua mihi?*

DE FRINGILLA.

Σπίτσα, *fringilla*, Anglicè *a chaffinche*, *a sheld* appel, *a spink*, Germanicè *eyn büchfink*.

Fringillæ, autore Aristotele, æstate tepidis locis, & hyeme, frigidis degunt, & inde puto apud Latinos nomen accepisse, quòd in frigore plures conuolantes apud nos cernantur, quàm æstate. *Pascere* magnitudine æquat, *uarijs coloribus*, albo nempe, uiridi & ruffo distincta est. *maris pectus rubescit, fæminæ pallefcit: cantat mas primo uere. Nidulatur fringilla in summis fruticum ramis, aut arborum infimis, nidumq; intus ex lana, forisq; ex musco facit.*

DE MONTIFRINGILLA.

Οροσπίζης, *mōtifringilla*, Anglicè *a bramlyng*, Germanicè *eyn rowert*.

ARISTOTELES².

[p. 60] *Montifringilla fringillæ similis est, & magnitudine proxima: sed collo cœruleo est, & in montibus degit, unde nomen accepit.*

Anicula, quam ego montifringillam esse credo, fringillæ magnitudine & corporis figura similis est: sed mas in collo plumas habet cœruleas, quas nō æquè promptè in fæmina depræhendas. Rostrum luteum est, & alæ uarijs coloribus, albo, nigro, & luteo nimirum distinguuntur, ut auriuittis. Vox illi insuanis & stridula est.

¹ *Epig. Lib. XIII. xlix.*

² *Hist. An. Bk VIII. 41.*

on worms¹; but is a little bird like the grasmusch of the Germans, living upon figs and grapes, as Martial prettily bears witness in these lines:—

Since the fig gives me nourishment, and I feed on sweet grapes,
Why has the grape not rather given me a name?

OF THE FRINGILLA.

Σπίζα, fringilla, in English a chaffinche, a sheld-appel², a spink, in German eyn büchfink.

Fringillæ—on Aristotle's authority—in summer haunt warm places, and in winter cold; and thence I think that they received their name among the Latins³, for when it is cold more are seen flocking round us than in summer time. In size the bird equals a Sparrow and is marked with various colours, namely, white and green, and russet. In the male the breast is ruddy, in the female pale. The male sings in the early spring. The Fringilla nests upon the highest boughs of shrubs or on the lowest boughs of trees, and fashions its nest inwardly of wool and outwardly of moss.

OF THE MONTIFRINGILLA.

Ὀροσπίζης, montifringilla, in English a bramlyng, in German eyn rowert.

ARISTOTLE.

The Montifringilla is like the Fringilla, and similar in size, but with a blue neck; and it lives in mountains, whence it has its name.

The little bird which I believe to be the Montifringilla, in size and shape of body is like the Fringilla, but the male has blue feathers upon the neck, which one cannot perceive so quickly in the hen. The beak is yellow, and the wings in truth are marked with various colours, yellow, black and white, as in the Aurivittis. Its note is unmelodious and grating.

¹ See p. 35.

² Shell-apple, or Apple-sheiler is still a Northumbrian name for the Chaffinch. The word 'sheld' may mean 'parti-coloured.'

³ Here Turner's mistaken etymology (*Fringilla a frigore*) is evident.

DE FLORO EX ARISTOTELE¹.

Anthos, siue florus, uermibus pascitur, & magnitudo illi, quanta fringillæ est. uicitat circa aquas & paludes, & ei color pulcher est, & uita commoda, odio equum habet, pellitur enim ab equo pabulo herbæ, qua uestitur. Nubeculans, nec ualens oculorum acie est, quippe qui uocem equi imitetur, atque aduolans [p. 61] equum fuget: sed interdum excipiat occidaturq; ab equo. In ægithum florus tantum odium gerit, ut ne mortuarum auiũ sanguis posse misceri dicatur.

DE FVLICA.

Κέπφος, *fulica*, Anglicè *a uuhite semauu uuith. a blak cop.* Germanicè, *eyu wyß mewe.*

Recentiores Græci, qui post Aristotelem scripserunt, larum & cepphum eandem auem fecerunt, ut Erasmus in Adagio, λάρος κέπφος, ex Aristophane² & eius interprete ostendit. Aristoteles uerò duas facit diuersas aues libro de historia animalium octauo his uerbis, ἔστι δὲ λάρος ὁ λευκός καὶ κέπφος. Iam quãnam ratione autores istos conciliẽ, nescio, nisi dicam poetæ rerum peculiars & proprias notas, & discrimina, philosophis multò negligentius obseruantes, aues corporis figura, natalibus, & uictus ratione similes, licet manifestis notis differentes, easdem aues fecisse, quas seueriores philosophi ad amussim omnia expedientes, in diuersas species distinxerunt.

Sed inter huius ætatis grammaticos, non minor est opinionum uarietas de fulica, quãnam illa sit, quàm [p. 62] inter Græcos de nomine controuersia fuit. Sunt enim

¹ *Hist. An.* Bk VIII. 41 and Bk IX. 18, 22, freely rendered.

² *Pax*, l. 1067. By 'interpreter' is evidently meant the Scholiast, who says that the Proverb is used of those who promise much, and perform little. .

OF THE FLORUS FROM ARISTOTLE.

The Anthos, that is Florus, feeds on worms ; its size is that of the Fringilla. It gets victual round waters and marshes ; its colour is fair, and its life easy to it. It holds the Horse in hatred, inasmuch as it is driven by the Horse from the grassy pastures where it feeds. It is purblind and nowise keen of eyesight, while it imitates the neighing of the Horse, and flying at it puts the Horse to flight, yet sometimes it is caught and then killed by the Horse. The Florus has so great a hatred of the Ægithus that it is stated that the blood of these two birds, even when dead, cannot be mixed.

OF THE FULICA.

Κέπφος, fulica, in English a white semaw, with a black cop, in German eyn wyss mewe.

The later Greeks, who have written after Aristotle, have made the Larus and the Cepphus the same bird, which fact Erasmus in his Proverb *λάρος κέπφος* shews, from Aristophanes and his interpreter. But Aristotle in the eighth book of his History of Animals keeps the two birds distinct, using the following words :—"There is the *λάρος* that is white, also the *κέπφος*." Now in what way to reconcile these authors I know not, unless I say that poets who observe more negligently than philosophers the peculiar properties of things, and their diversities, have made these birds the same, which are alike in form of body, breeding-time, and way of feeding, although differing in manifest respects, whereas philosophers, more strict than they, gauging all things exactly, have distinguished them as different kinds.

And yet there is not less diversity of opinion among the critics of our day about the Fulica, and what that bird may be, than there was controversy among the Greeks about its name. For there are teachers of a sort¹ in Lower

¹ 'Literatores' is here apparently used in a somewhat scornful sense.

Anglorū
lapuuinga.

in inferiori Germania literatores aliqui, qui fulicam kyuuittam suam esse uolunt, ex eo forsan opinionem suam adstruentes, quòd apud Plinium fulicæ cirrum tribui legerint. Est autem Germanorum kyuuitta cornice minor, plumis ferè uiridibus, et nigris per totum dorsum et caput et collū: uentre albo, longa, & semper erecta in capite: crista plumea, alis obtusioribus, & inter uolandum magnum strepitum edentibus, unde & uannellus a barbaris dicitur: aquis uerminum gratia, quibus solis uictitat, appropinquat, sed ipsas non ingreditur, in planis & in locis erica consitis, plurimū degit. Ad depopulandum uermes, nostrates in hortis sæpè alunt.

Sed hanc esse fulicam non patitur, quod Vergilius de fulica Georgicorum primo¹ ad hunc modum scribit:

*Iam sibi tum curuis malè temperat unda carinis,
Cum medio celeres reuolant ex æquore mergi,
Clamoremq; ferunt ad littora, cumq; marinæ
In sicco ludunt fulicæ.*

Hinc satis liquet kyuuittam non esse fulicam, quum non sit auis marina nec aquatica. Non desunt qui fulicam gallinam illam nigram aquaticam, alba in fronte macula, esse uolunt. Sed isti Vergil. et Aristotelis autoritate facilè erroris conuincuntur, quorum alter auem facit marinam, alter, nempe Aristoteles lib. octauo historicæ animalium, apud mare uiuere testatur. Quare quum palustris illa gallina neque auis sit marina, neq; apud mare uictum petat, sed in stagnis, paludibus, & recentibus aquis perpetuò degat: nec Vergilij fulica, nec Aristotelis κέπφος esse poterit. Sed iam restat, ut quam auem fulicam esse iudicem, ostendam.

Est auis marina, magnitudine monedulæ, sed alis acutioribus & longioribus, colore tota albo, excepto nigro, quem in capite gerit cirro: rostro etiam & pedibus puniccis. Hanc ego sæpè in mari nauigans, ex eo

¹ Lib. I. l. 360—3.

Germany, who will have it that the Fulica is their Kywit, possibly resting their opinion on what they have read in Pliny of a tuft being attributed to the bird Fulica. The Kywit of the Germans is, however, smaller than a Cornix, with the plumage almost green and black on the whole back and head and neck, the belly white, a long and always upright feathery crest upon the head, and somewhat rounded wings, which during flight make a great hurtling, whence it is even named by foreigners Vannellus. It approaches waters for the sake of worms, on which alone it feeds, but does not enter them. It mostly lives in open country, and in places overgrown with heather. Our people often keep this bird in gardens, to destroy the worms.

And yet what Vergil in this manner writes in the first book of his Georgics of the Fulica will not permit this bird to be his Fulica :—

“And now the waters scarce restrain themselves from the ships’ curving keels, while the swift Mergi wing their way once more out of the Ocean’s midst, bringing their noisy voices to the shore, and while the Fulicæ, frequenters of the sea, disport themselves on land.”

Hence it is clear enough that the Kywit is not the Fulica, since it is not a sea-bird nor a water-bird. There are not wanting those who would have that black Water Hen, with a white frontal patch, to be the Fulica¹. But on the strength of Vergil and Aristotle such are easily convicted of mistake, for one of these makes it a sea-bird, and the other, namely Aristotle in the eighth book of his History of Animals, bears witness that it lives about the sea. Wherefore, since that Marsh Hen is neither a sea-bird nor seeks its food about the sea, but constantly haunts pools, and marshes, and fresh waters, it can neither be the Fulica of Vergil nor the *κέπφος* of Aristotle. But it still remains that I should shew what bird I judge the Fulica to be.

There is a sea-bird², like a Daw in size, but with the wings sharper and longer, wholly white in colour, save for a black patch which it bears on the head, and with the beak and feet of purplish red. I often, journeying upon the sea, have had

¹ *I.e.* the Coot (*Fulica atra*).

² The Black-headed Gull (*Larus ridibundus*).

tempore, quo historiam animalium Aristotelis legeram, consideravi, tū præsertim, quando uel deficiente uento, uel flante contrario, emissa anchora, uentum sequundiorē quiescentes expectauimus. Hæc statim soluta anchora, gauijs comitata aduolat, ex purgamentis nauis eiectis, escæ nonnihil sibi promittens, diutino clangore defatigata, tandem kep̄ profert, ut lari cob. unde à nostris marini cobbi dicuntur. Ficri potest, ut in fulicarum genere quædam sint cinereæ, licet Plin. ubi ex autoritate Iubæ Diomedæas aues, fulicis similes albas esse tradit, uideatur fulicas omnes albas facere, nam [p. 64] non de eo, quod rarius, sed frequentius accidit, in genere loquuntur classici scriptores. Nidulātur lari & fulicæ in eisdem locis, in excelsis nempe petris, & marinis rupibus.

DE GAVIA.

Λάπος, gauia, a se cob or a seegell.

Gauiarum duo genera Aristot. facit: alterum album, quod apud mare, alterum cinerium, quod circa lacus & flumios uictum quærit. Gaviā albam à fulica parū differre arbitror, solo nimirum cirro et rostro. Gauia cinerea, quæ ad flumina & lacus ascendit, querula semper & clamosa est. pisciculos captat & uermes ad ripas lacuum. huius generis est & alia parua auis, nostrati lingua sterna appellata, quæ marinis laris ita similis est, ut sola magnitudine & colore ab illis differre uideatur: est enim iste larns, marinis minor & nigrrior. Tota æstate tam improbè clamosa est, quo tempore parurit, ut iuxta lacus & paludes degentes, immodico clamore tantum nō obtundat. hāc ego sanè auem esse credo, cuius improba garrulitas adagio, Larus parurit, locum fecit. uolat ferè perpetuò super lacus & paludes, nunquā quiescens, sed prædæ semper inhians. Nidulatur hæc in densis arundinetis. Marinæ gaviæ in petris & rupibus maritimis nidificant.

this bird in mind, from the time that I read Aristotle's History of Animals, and then especially when through the wind failing or blowing contrary, the anchor being dropped, we have been calmly awaiting a more favourable wind. The anchor being weighed this bird immediately flies to us in the company of Gulls, promising something to itself by way of food out of the refuse cast forth from the ship; at last exhausted by its constant cries it merely utters "keph," as Gulls cry "cob." And hence they are called Sea-Cobs by our countrymen. It may be that some of the race of Fulicæ are grey, though Pliny, when on the authority of Juba he relates that the birds known as Diomedæ's are white like Fulicæ, seems to put down all Fulicæ as white; for classical authorities speak not in any class of what more rarely, but of what more frequently occurs. Gulls nest in the same places as do Fulicæ, forsooth on lofty crags and rocks about the sea.

OF THE GAVIA.

Λάρος, gavia, a se cob or a see-gell.

Aristotle makes two kinds of Gaviæ, one white, which seeks its food about the sea, the other grey, which seeks it round the lakes and rivers. Now I think that the white Gavia differs but little from the Fulica, only indeed as to the hood and beak. The grey Gavia, which comes up to our rivers and lakes, is always querulous and full of noise. It catches little fishes and eats worms upon the banks of lakes. There is another small bird of this kind, called Stern¹ in local dialect, which is so like the sea Lari that it seems to differ from them only in its size and colour; for it is a Larus, though smaller than the sea Lari and blacker. Throughout the whole of summer, at which time it breeds, it makes such an unconscionable noise that by its unrestrained clamour it almost deafens those who live near lakes and marshes. This I certainly believe to be the bird whose vile garrulity gave rise to the old Proverb "Larus parturit." It is almost always flying over lakes and swamps, never at rest, but always open-mouthed for prey. This bird nests in thick reed-beds. The sea Gaviæ breed on crags and rocks about the sea.

¹ The Black Tern (*Sterna nigra*).

[p. 65]

DE GALERITA.

Κόρυδος, ἡ κορυδαλός, *Ang. a lerk or a laucrock, Germa. cyn lercf. Plinius naturalis historię undecimo libro, galeritam Gallico uocabulo, postea alaudam esse dictam ostendit. quare galeritę potius uocabulo, quàm alaudę Latinis utendum esse ex auctoritate Plinij censeo.*

ARISTOTELES DE GALERITIS¹.

Galeritarum duo sunt genera: alterum terrenum cristatum, alterū gregale. nec singulare more alterius, uerū colore simile, quāquam magnitudine minus, & galero carēs, cibo uerò idoneū, galeritę nunquā in arbore consistunt, sed humi².

Prius hoc Aristotelis genus uariam habet in uarijs regionibus cristam, alicubi semper apparentem: in alijs locis talem, ut pro arbitrato suo possit erigere aut deponere, quū una cademq; sit utriusq; auis magnitudo. & galeritam hanc maiorem, Angli proprię lercam nominant.

a uuilde
lerc or a heth
lerk.

[p. 66]

Alterum genus Aristotel. à nostris fera alauda, à Germanis heid lercf nominatur, in planis & locis erica confitis, & ad ripas lacuūm, uermium causa, quibus uiuitat, magna ex parte degit. Duplo ferē superiore minus est, & rostro tenui, & carne longē suauissima.

a uuodlerck.

Supereſt tertium galeritę genus, Germanis copera, à lōgiffima crista, ut arbitror, ita dictum, Aristoteli planē incognitum: nam prius Aristotelis genus esse non potest, quia minor est quàm ut illud esse possit: minūs autem illud genus esse non potest, quia galerum habet, qui Aristotelis posteriori generi deest. Quare galerita ista, Aristoteli fuit incognita. Et cūm Colonienſes aucupes coperam (quę medię est magnitudinis inter Aristotelis

¹ *Hist. An. Bk IX. 101.*

² *Hist. An. Bk IX. 66.*

OF THE GALERITA.

Κόρυδος, ἢ κορυδαλός, in English a lerk or a laverock, in German cyn lerch.

Pliny, in the eleventh book of his Natural History, has shown us that Galerita¹, taken from a Gaulish word, was called Alauda afterwards, wherefore I think that on Pliny's authority the name of Galerita should be used by those who write in Latin rather than Alauda.

ARISTOTLE, OF THE GALERITÆ.

Of Galeritæ there are two kinds, one is a crested ground-bird, but the other lives in flocks, not singly as the former. Yet in colour it is similar, though of a smaller size, and not having a crest. Moreover it is fit for food. The Galeritæ never sit upon a tree, but always on the ground.

The first kind given by Aristotle has in different lands a different crest, in one place always evident, elsewhere such that the bird can raise or lower it at will, although the size of either is one and the same. This larger Galerita Englishmen call the Lerc proper, while Aristotle's second sort is by our countrymen named a Wilde Lerc, and by the Germans a heid lerch; this for the most part lives in open country, and in places overgrown with heather, and on banks of lakes, for the sake of the worms on which it feeds. This bird is smaller by nearly one half than the aforesaid, with a slender beak, and flesh by far the sweetest.

There still remains a third kind of Galerita, the Copera of the Germans, thus named I believe from its very long crest, and certainly unknown to Aristotle, for it cannot be his first kind, inasmuch as it is smaller than that bird can be; likewise it cannot be the smaller sort, because it has a crest, which is not present in the latter kind. Wherefore this Galerita was unknown to Aristotle. And since the fowlers of Colonia [Cullen] with one accord assure us that the Copera (which is midway in size between Aristotle's crested Galerita

¹ Galerita is usually supposed to have some connexion with the Latin galea=a helmet.

galeritam cristatam, & non cristatam) concordibus adfirment suffragijs, hanc nullā habere peculiarem cantunculam, sed ineptè aliarum, quibuscum uiclitat, auium uoces referre, adducor planè ut credam hanc esse recentiorum Græcorum corydon, cuius in sequenti adagio mentio est ἐνάμουσις¹ καὶ κόρυδος φθέγγεται, & in hoc uersu :

εἰ κύνω δύναται κόρυδος παραπλήσιον ᾄδειν.

Nā galerita maior, pulchrè & suauiter cantat, & minorem cantu non minùs ualere tradunt aucupes. Hæc igitur quum uoce nihil possit, sed ineptè tantum aliarum uoces, suo garritu mentiri, recentiorum Græcorum erit corydos.

[p. 67]

DE GALLIS ET gallinis.

Αλέκτωρ, gallus, Anglicè a cok, Germanicè eyn hân. Αλέκτορις, gallina, Anglicè a hen, Germanicè, eyn hen. Saxones dicunt eyn hōn.

VARRO² DE RE RUSTICA LIBRO TERTIO.

Rusticæ
gallinæ.

Villaticæ.

Gallinæ rusticæ, sunt in urbe raræ, nec ferè manfuetę sine cauea uidentur Romæ, similes facie non his uillaticis gallinis nostris, sed Africanis aspectu, ac facie incontaminata. In ornatibus publicis solent poni cum psiticeis ac merulis albis, item id genus rebus inusitatis. Neque ferè in uillis oua ac pullos faciunt, sed in fyluis.

Africanę.

Meleagrides.

Gallinæ Africanæ, sunt grandes, uariæ, gibberæ, quas Meleagrides Græci appellant. Hæ nouissimæ in triclinium ganearium introierunt, [p. 68] è culina propter fastidium hominū. Venerunt³ propter penuriam magno. De tribus generibus, gallinæ •saginantur maximè uillaticę. Eas in-

¹ Undoubtedly the reading should be :—ἐν ἀμούσοις=among those with little voice. Both these proverbs are to be found in the *Adages* of Erasmus, Chil. II. Cent. ii. 92.

² Bk III. cap. ix.

³ No doubt a misprint for 'veneunt'=are sold.

and the non-crested) has no song of its own, but feebly imitates the notes of other birds with which it feeds, I am assuredly led to believe that it must be the Corydos of the later Greeks, of which mention is made in the proverb below:—

Ἐνάμουσις καὶ ὁ κόρυδος φθέγγεται,

and in this verse:—

εἰ κύκνω δύναται κόρυδος παραπλήσιον ᾄδειν.

For the larger *Galerita* sings fairly and sweetly, and the fowlers say that in its song the smaller kind is worth no less. Wherefore the third kind, since it has no power of voice except feebly to imitate the voices of the others by a twitter of its own, will be the *Corydos* of the later Greeks.

OF THE GALLI AND GALLINÆ.

Ἀλέκτωρ, gallus, in English a cok, in German eyn hān.

Ἀλέκτορις, gallina, in English a hen, in German eyn hen. The Saxons say eyn hōn.

VARRO, IN HIS THIRD BOOK DE RE RUSTICA.

The wild *Gallinæ* are rare in a city, and are scarcely seen at Rome tame, unless in a cage: they are not in appearance like the *Gallinæ* of our country-houses, but in look recall the African, and have the face unmarked¹. During public festivities these birds are wont to be exhibited with *Psitaci*, white *Merulæ*, and other unfamiliar kinds of that description. They do not usually lay their eggs or hatch their young at country-houses, but among the woods.

The African *Gallinæ*, which the Greeks call *Meleagrides*, are big, speckled, and hunch-backed. They have been the last to enter the dining room of eating-houses from the kitchen through people's fastidiousness. And from their rarity they have advanced to a great price. Of the three kinds those of the country-house are chiefly fattened. These they keep shut up in

¹ It is impossible to reconcile the statements of Varro and Columella as they stand. Various alterations of the text, which is possibly unsound, have been suggested.

Gallus medicus a
bauncok or
a cok of
kynde.

cludunt in locum tepidum & angustum & tenebrosum, quòd motus earum & lux pinguedini inimica, electis ad hanc rem maximis gallinis, nec continuò his, quas Melicas appellant falsò, quòd antiqui ut thetim thelim dicebāt, sic Medicā Melicā uocabant. Hæ primò Medicę dicebantur, quia ex Media propter magnitudinem erant allatæ.

COLUMELLA¹.

Cohortalis. Gallinarū alię sunt cohortales, alię rusticę,
Rustica. alię Africanę. Cohortalis est auis, quæ uulgò
per omnes ferè uillas conspicitur. Rustica,
quæ non dissimilis uillaticę, per aucupem deci-
[p. 69] pitur, eaq; plurima est in insula, quā nautæ in
Ligustico mari sitam, producto nomine alitis,
Africana. gallinariā uocitauerunt. Africana est, quam
plerique Numidicam dicunt, meleagridi similis,
nisi quòd rutilam galeam, & cristam capite gerit,
quæ utraque in Meleagride sunt cœrulea.

PLINIUS².

Africæ. Simili modo pugnant Meleagrides in Bæotia. Africæ, hoc est gallinarum genus gibberum, uarijs sparsum plumis, quę nouissimę sunt peregrinarum auium in mensis receptæ, propter ingratum uirus: uerùm Meleagri tumulus nobiles eas fecit.

ARISTOTELES³.

[p. 70] Item Hadrianæ paruo quidem sunt corpore, sed quotidie pariunt. Ferociunt tamen & pullos sæpè interimūt. Color his uarius. Oua⁴

¹ *De re rustica*, Lib. viii. cap. ii.

² *Hist. Nat.* Lib. x. cap. xxvi.

³ *Hist. An.* Bk vi. 1.

⁴ *Hist. An.* Bk vi. 5.

a warm, narrow, and dark place, for exercise and light hinder the fattening. The largest birds are chosen for this purpose, and not always those which men mistakenly call Melicæ, because the ancients, as they used to say Thelis for Thetis, also used to call Medica Melica. At first they were called Medicæ because on account of their size they were brought hither out of Media.

COLUMELLA.

Of Gallinæ some are court-yard birds, others again are wild, others are African. The court-yard bird is that which commonly is seen at nearly every country-house. The wild sort, which is not unlike that of the country-house, is trapped by bird-catchers. It is abundant in the island lying in the Ligurian sea, which sailors, lengthening the bird's name out, have called continuously Gallinaria. The kind from Africa, which many call Numidica, is like the Meleagris, save that on its head it bears a helmet and a crest of red, but in the Meleagris both of these are blue¹.

PLINY.

In a like way the Meleagrides fight in Bœotia. The Africæ, that is a hunch-backed kind of Gallinæ, are sprinkled here and there with variegated feathers; and they are the last of foreign birds to be received at table, on account of their unpleasant flavour: but the tomb of Meleager has ennobled them².

ARISTOTLE.

Likewise the Hadrianic birds are small indeed in body, but they lay their eggs daily. Yet they are fierce and often kill their chicks. They are of varied

¹ See Art. Guinea Fowl in Prof. Newton's *Dict. B.* p. 399.

² The reader may here be referred to any work relating to mythology.

alia candida sunt, ut columbarum & perdicum, alia pallida, ut palustrium, alia punctis distincta, ut Meleagridum & phasianorum.

In paucis istis, quos recensui autoribus, nō paucae sunt de rebus, quas tractauere, cōtrouersiae. Varro primū gallinas rusticas non uillicatis¹ sed Africanis similes esse scribit. Columella autē rusticam non dissimilem esse uillicae tradit. Varro Africanas, meleagrides facit, quod & Plinius etiam facere uidetur. Columella autem uarijs notis Africanas à meleagribus distinguit. Aristoteles Hadrianas gallinas facit uarias, ut Plinius itidem facit, & paruo corpore. Varro Africanas, quas non alias esse constat quàm Hadrianas, uarias & grandes facit. Verūm tanta autorum inter se dissidia componere, penes me non est. Sed quid de generibus istis sentiam, paucis aperiam. Gallina apud nos rustica nusquam reperitur, si gallina illa, quam morhennam uocant non sit, quam uarijs de causis

[p. 71] *antē attagenem esse conieci. Columellae meleagrides uidentur illae esse aues, quas nonnulli pauones Indicos appellant: nam illas paleis² & cristis caeruleis esse, in confesso est.*

a kok of
inde.

DE GALLINAGINE.

Ασκαλώπαξ, gallinago, Anglicè a uuod cok, Germanicè eyu holz jnepf.

ARISTOTELES³.

Gallinago per fepes⁴ hortorum capitur, magnitudine quanta gallina est, rostro longo, colore attagenae, currit celeriter, & hominem mirè diligit. Hæc in arbore nunquam⁵ sedet, & humi nidulatur.

¹ A misprint for 'uillaticis.'

² Perhaps the reading should be galeis, cf. p. 69, ll. 6—7.

³ *Hist. An.* Bk IX. 102. 66.

⁴ Aristotle has in addition—ἐρκεσι=in nets.

⁵ *Hist. An.* Bk IX. 66.

colouring. Of certain kinds of birds the eggs are white, as those of Columbæ and Perdices, others are pale, as those of marsh-birds; others marked with spots, as those of Meleagrides and Phasiani.

In those few authors, whose works I have scanned, not a few of the things which they have treated are disputable. First Varro tells us that the wild-bred Gallinæ are not like those of country-houses, but the African; while Columella states that the wild sort is not unlike that of the country-house. Varro makes the Africanæ to be the Meleagrides, which Pliny also seems to do. Yet Columella separates the Africanæ from the Meleagrides by various characters. And Aristotle makes his Hadrianic fowls of various colours, as does Pliny also, and of little size. But Varro makes the Africanæ big and of varied colours, though it is quite clear that they are nothing but the Hadrianic birds. But after all it is not in my power to adjust the mutual differences of authors, when so great; and yet in a few words I will disclose what I think of these kinds. The wild Gallina is not found with us in any part, if it be not that which they name Morhen, and this I formerly conjectured to be the Attagen for several reasons. The Meleagrides of Columella seem to be those birds which some call Indian peacocks¹, for they are admitted to have wattles and blue crests.

OF THE GALLINAGO.

Ἰσκαλώπαξ², gallinago, in English a wod cok, in German cyn holtz snepff.

ARISTOTLE.

The Gallinago is taken among the hedges of our gardens; it is of the size of a Gallina, but has a long bill, and the colour of the Attagenæ: it runs with speed, while it is wonderfully fond of man. This bird never sits on a tree and it nests on the ground.

¹ Turner was, of course, wrong in his conjecture.

² Turner makes σκολόπαξ the same as ἰσκαλώπαξ.

Gallinagines apud nostrates nunquam, nisi hyeme uidentur, quare de prole & modo nidulandi, nihil habeo, quod dicā. Capitur apud Anglos diluculo potissimum & crepusculo in syluis, retibus in loco arboribus uacuo, suspensis, & ueniente aue demissis.

DE COLIO, SIVE GALGULO¹, UT VERTIT GAZA,
EX ARISTOTELE.

[p. 72] Galgulo magnitudo quanta ferè turturi est: color luteus, lignipeta hic admodum est, magna'que ex parte macerie² pascitur, uocẽ emittit grandem, incola maximè Peloponesi hæc auis est.

Omnia, quæ Aristoteles ἁλτικνὸς colio, siue galgulo tribuit, Anglorum huhulo, & Germanorum grunspechto (si incolam maximè esse Peloponesi exceperis) conueniunt. Nam turturem ferè magnitudine æquat, lignipeta est: maceriam contundit, & uocem grandem emittit. Sed nihil hic definitio, sed inquirō tantum. Galgulus Plinio ἰστρος Græcè dicitur, & Aristoteli, si Theodoro fidimus, etiam κελεός. Quanquam mihi textum Græcum consulenti, alia auis κολῖός, & alia κελεός uidetur: nam κολῖός ἐστὶ ξυλοκόπος σφόδρα, καὶ νέμεται ἐπὶ τῶν ξύλων τὰ πολλὰ. Id est, colius est lignipeta ualde, & magna ex parte ad ligna pascitur, ὁ μὲν γὰρ κελεός παρὰ ποταμόν οἰκεῖ καὶ λόλμας³, quæ uerba Theodorus circa fruteta & nemora reddidit sed recte'ne an secus, doctis iudicandum relinquo. Vidi in Alpbis abieti insidentem auem, magnitudine turturis, uiridibus ueluti maculis in luteo distinctam, quæ tota corporis effigie

[p. 73] *picū Martium retulit, sed caput reliquo corpori (secus*

¹ *Hist. An.* Bk VIII. 44.

² Possibly a misprint for 'materie.'

³ *Hist. An.* Bk IX. 22. In his errata Turner alters λόλμας to χλόχμας, but evidently he means λόχμας.

Woodcocks are never seen with us save in the winter, wherefore I have naught to say about their young or mode of nesting. They are chiefly caught in England in the woods at daybreak and at dusk, by means of nets hung in some place devoid of trees, and dropped when the bird comes.

OF THE COLIUS, OR GALGULUS, AS GAZA RENDERS IT, FROM ARISTOTLE.

Of the Galgulus the size is almost that of the Turtur: it is yellowish in colour, and hacks timber very much, and for the most part feeds on trees: it utters a loud cry. This bird is mainly an inhabitant of the Peloponnese.

All that Aristotle has so far attributed to the Colius or Galgulus is in agreement with the Huhol of the English and the Grunspecht of the Germans (if one may except its being chiefly an inhabitant of the Peloponnese). For it is nearly equal to the Turtle-Dove in size; it hacks the timber, hammers rotten wood, and utters a loud cry. But I give no decision here, I only ask. The Galgulus of Pliny is said to be called the Icteros in Greek, and if we trust to Theodorus [Gaza] is also the *κελεός* of Aristotle. Though, on consulting the Greek text *κολιός* seemed to be one bird, and *κελεός*¹ another, for the reading was:—*κολιός ἐστι ξυλοκόπος σφόδρα, καὶ νέμεται ἐπὶ τῶν ξύλων τὰ πολλά*. That is, the Colius is especially a wood-hunter and for the most part feeds on wood, *ὁ μὲν γὰρ κελεός παρὰ ποταμὸν οἰκεῖ καὶ λόχμας*, which words Theodorus renders “around the thickets and the groves,” but whether rightly so or otherwise I leave to be decided by the learned. In the Alps I saw sitting upon a fir a bird of the size of a Turtle-Dove, marked as it seemed with green patches on yellow, which to me in the whole aspect of the body called to mind the *Picus Martius*, save that its head was like in colour to the rest of its body

¹ Turner appears to have had a text with the word *κελεός* in one place instead of *κολιός*.

atq; in pico sit) colore fuit simile, tibijs fuit breuibus, et capite erecto, & rostro longiusculo. An hæc galguli species fuerit, nihil statuo, sed fuisse suspicor.

DE GRACVLIS.

Erasmus in eruditissimo adagiorum opere, quoties κολοιὸς occurrit (occurrit aut non raro) graculum reddit. Theodorum Gazam hac in re, licet aliàs libenter, minimè secutus, qui κόλοιον semper monedulam uertit. Ego quoq; hac in re Erasmus potiùs quàm Gazam, uarijs de causis, imitari decreui:

ARISTOTELES SECUNDUM TRANSLATIONEM GAZÆ¹.

Monedularum tria sunt genera: unum, quod graculus uocatur, magnitudine quanta cornix, rostro rotundo, rutilo. Alterum, lupus cognominatum, paruū & scurra. Tertium, quod familiare, Lydiæ ac Phrygiæ terræ, idemq; palmipes est.

[p. 74] *Primum graculorum genus, quod Græci κορακλαν uocant, Plinio Pyrrhocorax est, Anglis a cornish choghe, Germanis eyn bergdöl, cornice paulò minor est, rostro luteo, paruo, & in fine nonnihil adunco; frequens est in alpibus, & apud Anglos in Cornubia, uocem habet monedula acutior, & magis querulani. Secundum genus λυκος καὶ βομολόχος, Græcè dictum, Latinis propriè monedula, quasi monetula à moneta dicitur, quam sola auium, ut inquit Plinius, furatur. Aurum non omnia tria genera furantur, sed secundum genus tantum, quare*

Monedula.
Plin. li. 10.

¹ *Hist. An. Bk IX. 100.*

(otherwise than it is in Picus), and the legs were short, the head was upright, the beak rather long. As to whether this may have been a kind of Galgulus, I do not certify, but I suspect it to have been.

OF THE GRACULI.

Erasmus in his very learned work on Proverbs, as often as *κολοιδς* occurs (and it occurs not seldom) renders it by Graculus, in this thing following by no means Theodorus Gaza—though at other times he does so freely—who in every case renders *κολοιδς* by Monedula. And in this thing I also have determined for divers reasons here to imitate Erasmus rather than Gaza.

ARISTOTLE ACCORDING TO THE TRANSLATION OF GAZA.

Of Monedulæ there are three sorts: the first, which is called Graculus, in size as big as Cornix with a curved red bill. The next, also named Lupus, small, and a mimic. The third, which is well known in Lydia and Phrygia, is web-footed.

Now the first kind of Graculi, which the Greeks call *κορακίας*, is the Pyrrhocorax of Pliny and the Cornish Choghe of Englishmen, *eyn bergdöl* of the Germans. It is a little smaller than the Cornix, with a yellow bill¹, not large, and somewhat hooked towards the tip, it is abundant in the Alps and in Cornwall in England. It has a sharper and more querulous cry than the Monedula. The second sort called *λύκος* and *βωμολόχος* in Greek, is by the Latins strictly named Monedula, as if it were Monetula, from the Moneta [money] which alone of birds, as Pliny says, it steals. The three kinds do not all steal gold—only the second does—

¹ Here there is an evident confusion between the Chough (*Pyrrhocorax graculus*) with its red bill, and the yellow-billed Alpine Chough (*P. alpinus*).

secundum genus solum erit monedula, de cuius furacitate pulchrè etiā his uersibus scribit Ouidius¹:

*Mutata est in auem, quæ nunc quoq; diligit aurum,
Nigra pedes, nigris uelata monedula pennis.*

Phalacro-
corax.

Monedulā Angli uocāt, a caddo, a chogh or, a ka. Germ. cyn döhl, & Saxon. cyn älf. Multò minor est pyrrhocorace monedula, & in syluis nidulatur, & in cauis arboribus, & in templorū turribus. Tertium genus, Aristoteles 8 libro historiæ animalium² ita describit: Palmipedum grauiore circa lacus & amnes uersantur, ut anas, phalaris, urinatrix. Ad hæc [p. 75] bosca similis anati, sed minor, & qui coruus appellatus est, cui magnitudo quanta ciconiæ, sed crura breuiora, palmipes natansq; est, colore niger, infidet arboribus & nidulatur in ijs. Hæc Aristoteles. Coruus iste, nisi fallar, Plinio phalacrocorax est, & Heluctiorum Vualtrapus, de quo Plin. ad hunc modum scribit³: Iam & in Gallia Hispaniaq; capitur attagen, & per alpes etiam, ubi & phalacrocoraces, Balearium insularum peculiare: sicut & alpium pyrrhocorax. Et alibi de eodem⁴: Quædā animalium naturaliter caluent, sicut struthiocameli, & corui aquatici, quibus apud Græcos nomen est inde.

Phalacro-
corax .i.
coruus cal-
uus.

¹ *Metamorph.* Lib. VII. ll. 467—8.

² Bk VIII. 48.

³ *Hist. Nat.* Lib. X. cap. xlviii.

⁴ *Op. cit.* XI. cap. xxxvii.

wherefore this second kind alone shall be Monedula ; moreover Ovid happily describes its thievish habits in the following lines :—

Was changed into a bird, which even now loves gold,
Monedula the black of foot, in plumage black arrayed.

The English call the Monedula a Caddo, Chogh, or Ka ; Germans eyn döl ; and Saxons eyn älke. The Monedula is much smaller than the Pyrrhocorax, and nests in woods and hollow trees and towers of churches. The third kind is thus described by Aristotle in the eighth book of his History of Animals :—

Of web-footed birds the heavier haunt lakes and rivers, as the Anas, Phalaris, and Urinatrix. Add to these the Bosca, which is like the Anas but smaller, and that which is called Corvus, whose size is that of a Ciconia, but it has shorter legs ; it is web-footed and a swimmer : black in colour, it perches on trees, and nests in them. So far Aristotle.

Unless I err, this Corvus is the Phalacrocorax of Pliny and the Swiss Waltrapus¹, of which Pliny writes after this fashion :—

Further, the Attagen is caught in Gaul and Spain, and even on the Alps, where Phalacrocoraces also are, proper to the Balearic isles, as the Pyrrhocorax is to the Alps.

And in another place of the same bird :—

Some animals are naturally bald, as Struthiocameli and Corvi Aquatici, whence is their name among the Greeks.

¹ Mr Rothschild identifies this bird with *Comatibis comata* = *C. eremita* (L.), no doubt rightly. See *Bull. Brit. Orn. Club*, xii. p. 56. *Novitates Zoologicae*, 1897, p. 371, and Pliny, Lib. x. cap. xlviii.

Iam ut sciatis qualis'nam avis sit Heluetiorum Vualtrapus, quam conijcio phalacrocoracem esse, & tertium genus graculi, avis est corpore longo, & ciconia paulò minore, cruribus breuibus, sed crassis, rostro [p. 76] *rutilo, parùm adunco, & sex pollices longo, albam quoque in capite maculam, & eam nudam, nisi malè memini, habuit. Si palmipes sit, & interdum natet, indubitanter tertium graculorum genus esse adfirmarem: uerùm licet auem in manibus habuerim, an palmipes fuerit nec'ne, & caluus, non memini: quare donec isthæc certiùs nonero, nihil statuam.*

Præter hæc tria graculorum genera ab Aristotele descripta, noui & quartum genus, quod in alpinis Rheticis uidi, Aristotelis lupo minus, nigrum & albis maculis per totum corpus, more sturni distinctum, garulitate superiora genera multùm superans, semper in syluis & montibus degens: cui Rheti nucifragæ nomen, à nucibus quas rostro frangit & comedit, indiderunt.

Graculus
nucifrag⁹
eyn nouf-
brecher.

DE GRUE.

Γεράνος, grus, Anglicè a crane, Germanicè eyu frän / oder eyu fränich.

ARISTOTELES¹.

Alia de ultimis propè ueniunt, ut grues faciunt, quæ Scythicis ad paludes Aegypto², unde Nilus profluit, ueniunt: quo in loco pugnare cū pygmeis dicuntur. Non enim id fabula est, sed certè genus tum hominum tum etiam equorum pusillum, ut dicitur est, deguntq; in cauernis, unde nomen troglodytæ, à fubeundis cauernis accepere. Grues³ etiam multa prudenter faciunt: loca enim longinqua petunt, fui commodi gratia, & altè uolant, ut procul prospici-

¹ *Hist. An.* Bk VIII. 75—76.

² A variant reading is τὰ ἄνω τῆς Αἰγύπτου.

³ *Hist. An.* Bk IX. 70.

And now, that you may know what sort of bird the Switzers' Waltrapus may be, which I conjecture is the Phalacrocorax, and the third kind of Graculus, it is a bird long in the body, which is rather less than that of the Ciconia, and the legs short but stout, the bill reddish, a little hooked, and six inches in length—further it had a white spot on the head, and that, unless my memory fails me, bare. If it be web-footed and swim at times, I should affirm that it undoubtedly was the third kind of Graculus; but, though I have myself had the bird in my hands, I do not now remember whether it was web-footed or not, nor whether it was bald. Wherefore I will determine nothing, until I shall have a surer knowledge of these things.

Besides the said three kinds of Graculi described by Aristotle I know a fourth, which I have seen upon the Rhætic Alps, smaller than Aristotle's Lupus, black and marked with spots of white on the whole body, as a Starling is; it far surpasses all the above-named kinds in chattering; it always lives in woods and mountains. Now to this the Rhetians have given the name of Nucifraga, from the nuts which it breaks with its bill and eats.

OF THE GRUS.

Γερανός, grus, in English a crane, in German eyn krän, or eyn kränich.

ARISTOTLE.

Others come almost from earth's utmost parts, as do the Grues, which come from the Scythians to the Egyptian marshes, whence the Nile flows forth: in which place they are said to fight with Pygmies. And this is no mere fable, but assuredly there is, as it is said, a dwarf race both of men and horses, and they live in caves, whence they have got the name of Troglodytæ, from dwelling in caves. The Grues furthermore do many things with prudence, for they seek for their convenience distant places, and fly high that they may look out far, and, if they shall

cere possint, & si nubes tempestatem'ue uiderint, conferunt se in terrā, & humi quiescūt. Ducem etiam habent, & eos, qui clament, dispositi¹ in extremo agmine, ut uox percipi possit. Cū consistunt, cæteri dormiunt, capite subter alam condito, alternis pedibus insistentes. Dux detecto capite, prospicit, & quod senserit, uoce significat.

Pipers.

Vipiones Plin. dicuntur minores grues & iuniores, [p. 78] ut pipiones iuniores dicuntur columbæ. Apud Anglos etiam nidulantur grues in locis palustribus, & earum pipiones sæpissimè uidi, quod quidam extra Angliam nati, falsum esse contendunt.

DE HIRVNDINE.

Χελιδών, *hirundo*, Anglicè à *suuallouuc*, Germanicè *eyn schwalb*. Saxonibus *est eyn swale*.

ARISTOTELES².

Hirundo carnibus uescitur, bis in anno parit, & tota hyeme latet. Omnino ratio brutorum, magnā refert uitæ humanæ similitudinem magisq; in minori genere, quàm in maiore. uideris intelligentiæ rationem, quod primum in auium genere hirundo in effigendo cōstituendoq; nido ostendit, confingit implicito luto, festucis ad normam lutariæ paleationis, & si quādo luti inopia est, se ipsa madefaciens, uolutat in pul-
[p. 79] uerem omnibus pennis. Stragulum etiam facit more hominum duriore primum subijciens, & modicè totum consternens, pro sui corporis magnitudine.

¹ 'dispositi' is here apparently attracted to 'qui.'

² Four passages are incorporated in Turner's selection:—*Hist. An.* Bk VIII. 39, Bk VI. 36, Bk VIII. 107 and Bk IX. 51, 52.

have seen clouds or a storm, betake themselves to earth, and take rest on the ground. They have a leader also and those who, disposed at each end of the band, may call out, that their voice may be perceived. The others sleep when they alight, with the head hidden underneath the wing, standing alternately on either foot. The leader gazes round him with uncovered head, and by his cry gives notice of whatever he perceives.

The smaller, that is younger, Cranes are called by Pliny Vipiones, as young Doves are known as Pipiones. Cranes, moreover, breed in England in marshy places, I myself have very often seen their pipers¹, though some people born away from England urge that this is false.

OF THE HIRUNDO.

Χελιδών, hirundo, in English a swallowe, in German eyn schwalb. Among the Saxons it is eyn swale.

ARISTOTLE.

The Hirundo feeds on flesh, and lays eggs twice a year, for the whole winter also it lies hid. The way of brutes upon the whole bears a marked likeness to the life of man, and more so in the smaller than the larger kinds. One may observe the understanding way which the Hirundo foremost in the ranks of birds shews in the constitution and construction of its nest. It builds it by applying mud to straws, after the rule of daub and wattle work, and if there ever be a scarcity of mud, it wets itself and rolls itself in dust with all its feathers. It moreover makes a bedding after the manner of men, first laying a foundation of the harder stuff below, and moderately covering the whole in proportion to its size.

¹ Young pigeons are still called Pipers in England.

PLINIUS ¹.

Hirundines luto construunt, stramento roborant. Si quando inopia est luti, madefactum multa aqua pennis puluerem spargunt. Ipsum uerò nidum mollibus plumis floccisque consternunt, tepefaciendis ouis, simul ne durus sit infantibus pullis.

Secundū. Alterum genus est hirundinum rusticarum & agrestium, quæ rarò in domibus, diuersos figura, sed eadē materia nidos confingunt, totos supinos, faucibus porrectis in angustum, utero [p. 80] capaci: mirum qua peritia occultandis habiles pullis, & substernendis molles.

Tertium. Tertium hirundinum genus est, quæ ripas excauant, atque ita †internidificant. Non faciunt hæ nidos, migratque multis diebus antè, si futurum est, ut auctus amnis attingat.

† in terra
foetificat.

DE APODIBUS, EX ARISTOTELE ².

Nonnullæ aues depedes ³ sunt, quæ ob eam rem apodes à paruitate pedum nuncupantur. quod genus auiculæ, pennis plurimum ualet, sicut & cætera quoque propè similia, ut pennis præualere, sic pedibus degenerare uidentur. ut hirundo & falcula siue riparia ⁴. Hæc enim omnia, & moribus, & uolatu, & specie proxima [p. 81] inter se conspiciuntur. Apparet apes omnibus anni temporibus: riparia æstate tantum cum imber incescit: tum enim & apparet & capitur.

Riparia siue
falcula

¹ *Hist. Nat.* Lib. x. cap. xxxiii.

² *Hist. An.* Bk I. 10.

³ κακόποδες in the Greek.

⁴ These two words are probably interpolated.

PLINY.

Hirundines build nests of mud, and strengthen them with straw. And if there ever be a scarcity of mud, they sprinkle a good store of water from their feathers on the dust, which is thus moistened. The nest itself they further line throughout with soft feathers and wool, to thus keep the eggs warm, and also that it may not be too hard for the young chicks.

There is another sort of the Hirundines of the country and the fields, which rarely build their nests in houses, different in shape, but of the same material, and facing wholly upwards, having entrances prolonged into a strait with a capacious belly¹; it is wonderful how skilfully they are adapted for concealing young, and soft for them to lie upon.

There is a third kind of Hirundines which bore in banks, and thus breed within holes. These make no nests, and migrate many days before, if it be likely that the stream in flood should reach them.

OF THE APODES, FROM ARISTOTLE.

Some birds are weak-footed, and for that reason from the smallness of their feet are known as Apodes. This kind of little bird is very strong upon the wing, just as some others that are nearly like it seem to lose in strength of foot proportionately as they gain in power of flight, as the *Hirundo* and the *Falcula*, in other words *Riparia*. For all these in their habits, flight, and look seem very near each other. The Apes may be seen at all times of the year, but the *Riparia* only in summer, when the rains begin: for then it is both noticed and is caught. In

¹ Pliny evidently refers to *Hirundo rufula*, which builds a flask-shaped nest against a cliff.

Apodes. deniq; rara hæc auicula est. Apodes¹, quos aliqui cypsellos uocant, similes esse hirūdinum, iam dictum est: haud enim ab hirundine discerni possunt, nisi quod tibijs sunt hirsutis. Nidum specie cistellæ² productæ longius fictæ ex luto, imò aditu dato arctissimo faciunt, idq; locis angustis³, intrā saxa & specus, ut & belluas, & homines possint deuitare.

PLINIUS⁴.

Apodes. Plurimū uolant, quæ apodes uocātur, quia carēt usu pedum. Ab alijs Cypseli appellantur, hirundinum specie. Nidificant in scopulis. Hæ [p. 82] sunt, quæ toto mari cernuntur: nec unquam tam longo naues, tamq; continuo cursu, recedunt à terra, ut non circumuolitēt eas apodes. Cætera genera residunt, & insistant: his quies nisi in nidis nulla: aut pendent, aut iacent. *Hactenus Plinius & Aristoteles.*

Aristoteles tria tantum hirundinum genera facit: domesticas, apodes, & falculas. Plinius autem quatuor genera facere uidetur: domesticas, rusticas, apodes & riparias. Quod si uerum sit, hirundines domesticæ, sanguinolento pectore nobiles, erunt primum genus. Secundum genus maximæ illæ & nigerrimæ hirundines gregatim plerumq; uolantes, facere uidentur. Tertium genus, hirundines quæ in summis turribus & altis templorum fenestris nidulantur, efficiunt. Quartum genus ripariæ siue falculæ erunt. Quod si ista diuisio parum arrideat, ad primum genus referantur hirundines illæ in domibus rusticorum semper nidificantes, quæ à reliquis generibus, duæ sanguinolentæ

Hirūdines
domesticæ.

¹ *Hist. An.* Bk IX. 108.

² κυψέλεσις is the word in Greek.

³ στενωπό; another reading is στεγνωπό=under cover.

⁴ *Hist. Nat.* Lib. X. cap. xxxix.

short this little bird is rare. The Apodes, which some call Cypseli, are like Hirundines, as has been said before, for they are not to be distinguished from the *Hirundo*, save by having hairy legs. The nest which they construct looks like a little basket formed of mud somewhat drawn out, an entrance of the straitest opening beneath; and this they place in cracks within the rocks and caves, that they may avoid both beasts and men.

PLINY.

The birds which, because they cannot use their feet, are called Apodes, live chiefly on the wing. They are named Cypseli by some, in aspect they are like Hirundines. They nest in crags. These are they which are seen on all parts of the sea, nor do ships ever leave the land upon so long or so continuous a course but that the Apodes still fly around them. The other kinds alight and settle, but for these there is no rest save in their nests alone, they either hang or lie. So far Pliny and Aristotle.

Now Aristotle makes only three kinds of Hirundines, those of the house, the Apodes and the *Falculæ*. Yet Pliny seems to make four kinds, those of the house, the *Rusticæ*, the Apodes, and the *Ripariæ*. If that be true, our House Swallows, well known from their blood-coloured breast, will be the first-named kind. Those very large and black Swallows, that mostly fly in flocks, appear to form the second kind. Those Swallows which make nests upon the tops of towers, and in lofty church windows, constitute the third. And the *Ripariæ* or *Falculæ* will be the fourth. But should the said division not approve itself, then to the first-named kind may be referred those Swallows which invariably build on the houses of the country people. Two patches of a blood colour, which one may see on each side

maculæ, quas utrinque in pectore uideas, distinguunt, quod Onidius¹ his uersibus pulchrè ostendit:

[p. 83] *Altera tecta subit, nec adhuc de pectore cædis
Excessère notæ, signataq; sanguine pluma est.*

*Hoc primum genus Angli a suuallouu nominât
& Germani eyn schwalb.*

Apodes.

*Secundum genus faciunt apodes tam maiores quàm
minores. maiores uoco maximas illas hirundines, gre-
gatum & altiùs cæteris uolantes, quæ in arbore, more
hirundinum aliarum nunquam consistere uisuntur. mi-
nores uoco, quæ in scopulis, templorum fenestris æditori-
bus & summis turribus nidos figunt. Maiores Germani
uocant geyr swalben, Angli the great suuallouues. Mi-
nores Angli uocant rok martinettes or chirche martnettes,
Germani uocant firdy swalben.*

Facula siue
riparia

*Tertium genus, quod in ripis nidulatur, Angli a
bank martuet², Germani eyn über³ swalbe, aut speiren
nominant.*

DE HÆMATOPODIBUS, EX PLINIO⁴.

Rostrum & prælonga crura rubra hæmoto-
podi⁵ sunt, multò Porphyryone minori: quâquam
[p. 84] eadem crurum altitudine. Nascitur in Aegypto.
Insistit ternis digitis, præcipuum ei pabulum
muscæ. Vita in Italia paucis diebus.

*Est apud Anglos in locis palustribus avis quædam
longis & rubris cruribus, nostra lingua redshanca dicta,
cui an descriptio hæmotopodis Pliniani conueniat nec-
ne, qui apud Anglos degunt, inuestigent & examinent.*

DE IVNCONE.

*Σχοίνικλος, iunco, Anglicè a rede sparrouu, Ger-
manicè eyn reydt mûß. Iunco, ut scribit Aristoteles octauo
historiæ animalium, & capite tertio, ad ripas lacuum &
fluminum uiclitat, & caudam frequenter motitat, & ex
eodem constat, auem esse paruum: nam turdo minorem*

¹ Metam. Lib. vi. ll. 669—70.

² A misprint for 'martnet.'

³ A misprint for 'über.'

⁴ Hist. Nat. Lib. x. cap. xlvii.

⁵ Another reading is 'Himantopus,' but the Stilt-Plover has not a red bill.

of the breast, distinguish these from the remaining sorts, as Ovid prettily sets forth in these verses:—

“The other haunts our roofs, nor have the marks of slaughter yet departed from its breast, and its plumage is stained with blood.”

The English call this first kind a Swallow, and the Germans cyn schwalb.

The Apodes, the greater and the less, compose the second kind. I call greater those very great Swallows that fly in flocks, and higher than the rest, which are never observed to settle on a tree, after the manner of our other Swallows. I call less those which fix their nests to rocks, lofty church windows and the tops of towers. The greater kind the Germans call geyr swalben, and the English the Great Swallows; but the less the English call rok martinets or chirche martnettes, the Germans kirch swalben.

The third kind, that which breeds in banks, the English name a bank martnet, the Germans cyn ufer swalbe or speiren.

OF THE HÆMATOPODES, FROM PLINY.

The Hæmatopus has its bill and very long legs red, and is much less than the Porphyrio, though of the same height of leg. It is native in Egypt. It stands on three toes to a foot; flies are its favourite food. It lives in Italy but a few days.

There is in marshy places in England a certain bird with long red legs, called Redshank in our tongue, but whether the description of the Hæmatopus of Pliny agrees with this or not let those who live in England seek out and enquire.

OF THE JUNCO.

Σχοίνικλος, junco, in English a rede sparrow, in German eyn reydt müss.

The Junco, as Aristotle writes in the eighth book of his History of Animals, and in the third chapter, lives on the banks of lakes and streams, and flirts its tail continually; and it is clear from him that it is a small bird, for he makes

facit. Ego igitur quum nullam aliam nouerim auiculam, iuncis & harundinibus insidentem, præter Anglorum passerem harundinarium, illum iunconem esse iudico. Avis est parua, passere paulò minor, cauda longiuscula & capite nigro. cætera fusca.

DE LINGULACA, EX ARISTOTELE¹.

[p. 85] Lingulaca, quæ Græcè γλώττις dicitur lin-
guam exerit longam, unde nomē habet, una
est è coturnicum ducibus, formam habet auium
lacustrium.

DE LAGOPODE EX PLINIO².

Præcipuo sapore lagopus est: pedes leporino uillo ei nomen hoc dedere. Cetero candidè³, columborum magnitudine, non extra terram⁴, in qua nascitur, eam uesci: quando nec uiua manfuescit, & corpus occisè, statim marcescit. Est & alia, nomine eodem, à coturnicibus magnitudine tantum differens, croceo tinctu cibus

† gratissima. †aptissima. *Huius hoc uersu Martialis⁵ meminit:*

Si meus aurita gaudet lagopede Flaccus.

DE LIGVRINO SIVE SPINO.

Ακανθίς, spinus, siue ligurinus, Anglicè, a grene finche, ut conijcio, Germanicè, cyn firšfinche.

ARISTOTELES⁶.

[p. 86] Ligurini, & uita & colore ignobiles sunt, fed ualent uocis amœnitate⁷, & ex auium albo sunt, quæ carduorum femine uescuntur⁸. Florus, spinus, & ægithus, odium inter se exercēt. Spinus etiam bellum cum asino gerit.

¹ *Hist. An.* Bk VIII. 83.

² *Hist. Nat.* Lib. X. cap. xlviii.

³ After 'dedere' substitute a comma for the full stop.

⁴ Pliny seems to have written 'facile' here, in addition.

⁵ *Epigr.* Lib. VII. lxxxvi.

⁶ Three passages are here combined:—*Hist. An.* VIII. 42, IX. 22, IX. 92.

⁷ The words of Aristotle are φωνὴν μέντοι λιγυρὰν ἔχουσιν.

⁸ See p. 35.

it less than a *Turdus*. Therefore, since I know no other little bird which sits upon the rushes and the reeds, save the Reed Sparrow of the English, I believe that kind to be the Junco. Now this bird is small, a little smaller than a Sparrow, with a longish tail, and a black head. The other parts are brown.

OF THE LINGULACA, FROM ARISTOTLE.

Lingulaca, in Greek called *γλωττίς*, puts forth a long tongue, whence comes its name; it is one of the leaders of Coturnices, it has the form of a lake-haunting bird.

OF THE LAGOPUS, FROM PLINY.

The Lagopus is in flavour excellent, its feet shaggy as in a hare have given it this name. Otherwise it is white, in size as the Columbi; it is not eaten except in the land of which it is a native, since it is not tameable while living, and when killed its flesh soon putrefies. There is another bird of the same name, differing but in size from the Coturnices, most excellent for food with yellow saffron sauce.

Of this Martial makes mention in the following verse:—

If my Flaccus delights in the eared lagopes.

OF THE LIGURINUS OR SPINUS.

Ἀκανθίς, spinus, or ligurinus, in English a grene finche, as I suppose, in German *eyn kirsfincke*.

ARISTOTLE.

The Ligurini, commonplace in mode of life and colour, yet excel in pleasantness of song. And they are of the list of birds which feed on thistle-seed. The Florus, the Spinus, and the *Ægithus* shew mutual dislike. The Spinus wages war moreover with the Ass.

Spinum Aristotelis grenefincam nostram esse arbitror: nam illa inter spinas plurimum degit, & ex herbarum feminibus uictitat. Avis, quam spinum esse iudico, magnitudine passerem æquat, tota uiridis est, præsertim mas in hoc genere, fœmina ferè pallida est. uescitur carduorum maiorum semine & lapparum, ut auriuittis minorum, nidulatur in ramis salicū aut prunorum syluestriū. cantat amænè, & cibum & potū è situlis haurire non recusat.

Sed obijciat mihi forsan quispiam, hanc colore uiridi adeoq; amæno, non posse spinum esse, quem Aristoteles colore ignobilem esse testatur. Sciat uelim, qui hoc mihi obijcit, eundem Aristotelem uiridem colorem damnare, etiam in aue, quæ tota uiridis est, & à uiriditate nomen accepit. Verba Aristotelis¹ hæc sunt: Vireo², qui totus uiridis est, docilis & ad uite munera [p. 87] ingeniosus notatur, sed malè uolat, nec grati est coloris. Hæc Aristoteles.

DE LVTEA.

Χλόρευς, luteus siue lutea, Anglicè a yellow ham, a youulryng. Germanicè cyn geelgerst.

ARISTOTELES.

Luteus à colore partis suæ inferioris pallido dictus, magnitudine alaudæ est. Parit oua quatuor aut quinque. Nidum sibi ex symphyto stirpitis euulso facit. Sed stragulum subijcit ex lana & uillo.

¹ *Hist. An.* Bk IX. 98, 89.

² *Χλόριων.*

I think that Aristotle's Spinus is our Grenefine, for it lives for the most part among thorns, and feeds upon the seeds of grasses. The bird which I believe to be the Spinus in its size equals a Sparrow, and is wholly green, and in this kind the male especially, the female being somewhat pale. It feeds upon the seeds of bigger thistles and of burdocks, as the Aurivittis does upon the smaller; and it nests on branches of the willow or wild plum. It is a pleasant songster, and does not refuse to draw its food and water up in little buckets.

But some one may perhaps object to me that this bird with its green and somewhat pretty colour cannot be the Spinus, inasmuch as Aristotle testifies that it is commonplace in colour. I should like the man who thus objects to me to know that Aristotle equally condemns green colour even in a bird which is entirely green, and from its greenness has received its name. These are the words of Aristotle:—

“The Vireo, which is entirely green, is singled out as easy to be taught, and clever for the business of life; but it flies badly and its colour is unpleasing.”

So far Aristotle.

OF THE LUTEA.

Χλωρεύς¹, luteus or lutea, in English a yellow ham, a yowlyng, in German eyn geelgorst.

ARISTOTLE.

The Luteus, so named from the pale colour of its lower parts, is of the size of an Alauda. And it lays four eggs, or even five. It builds itself a nest of comfrey torn up by the roots, but spreads within a covering of wool and hair.

¹ Aristotle in his *History of Animals* mentions three birds, χλωρίς, χλωρίων, and χλωρεύς: but Turner's quotation with regard to χλωρεύς is found in Aristotle under χλωρίς in a passage (*Hist. An.* Bk IX. 83) where there seems to be no alternative reading.

Auicula, quam luteum esse credo, passerc paulò maior est. Maris pectus & uenter lutea sunt: fæminæ uerò pectus luteum, & uenter pallidus est, in capite dorso & alis, pennis fuscis lutæ intermîscentur. Rostrum utrique firmum & breue, in quo tubercum quoddam dentem mentiens, reperias, præter uermes, hordeo & aucna libenter uestitur. Cauda huius auiculæ longiuscula est, & frequenter motitans.

DE LVTEOLA.

Χλώρις, *lutcola*, Anglicè *a fiskin*, Germanicè *eyn* [p. 88] *zeyfich*, quibusdam *eyn engelchen*.

Lutcola, lutea superiùs descripta, multò minor est, & colore ad uiriditatem magis tendente, pectore luteo est, & rostro longiusculo, tenui & acuto, auriuittis simili, duas habet maculas nigras: alteram in fronte, alteram sub mento, cantillat non insuauiter. Rara apud Anglos hæc est, nec uspiam ferè alibi quàm in caueis cernitur. Semel tamen in Cantabrigianis agris uidisse recordor. Huius generis sunt, quas Anglia aues canarias uocat.

DE LVSCINIA.

Αἰδών, *luscinia, philomela*, Anglicè *a nyghtyngall*, Germanicè *eyn nachtgâl*.

ARISTOTELES¹.

Parit luscinia æstate quinq; aut sex oua, conditur ab autumnno usq; ad uernos dies, luscinia² canere solet assiduè diebus ac noctibus quindecim, cùm fylua³ fronde incipit opacari. dein canit quidem, sed non assiduò, mox adulta æstate uocem mittit diuersam, nò insuper uar[ia]m, aut celerem⁴ modulatamq; sed simplicem,

¹ *Hist. An.* Bk. v. 31.

² *Hist. An.* Bk. ix. 255.

³ Aristotle has ὄρος (mountain) here.

⁴ Or τραχέα = harsh.

The little bird, which I believe to be the Luteus. is somewhat bigger than a Sparrow. It is yellow on the breast and belly in the cock; but in the hen the breast is yellow and the belly pale. Yellow are mixed with dark feathers upon the head, back, and wings. In each of them the beak is short and stout, and on it one may find a sort of knob that simulates a tooth. Apart from worms it eats barley and oats freely. The tail of this small bird is rather long and is in constant motion.

OF THE LUTEOLA.

Χλωρίς, luteola, in English a siskin, in German *eyn zey-sich*, or of some *eyn engelchen*.

The Luteola is much smaller than the Lutea above described, and with a colour tending more to green. It has a yellow breast, a longish, slender, pointed bill, like that in *Aurivittis*, and two spots of black, one on the forehead, one beneath the chin; it warbles with some sweetness. In England it is rare, and scarcely to be seen elsewhere than in cages. Yet I remember having seen it once among the fields of Cambridgeshire. Of this kind are those which England calls Canary birds¹.

OF THE LUSCINIA.

Ἀηδών, luscinia, philomela, in English a nyghtyngall, in German *eyn nachtgäll*.

ARISTOTLE.

In summer the Luscinia lays five or six eggs, but from autumn it lies hid continually until the days of spring. Now the Luscinia is wont to sing incessantly for fifteen days and nights, when woods begin to become dark with foliage. Later it sings indeed, but not incessantly, then in the height of summer it gives forth a different note, not varied over and above, or quick and modulated, but a simple

¹ Gesner, the first to describe the Canary-bird, states that Turner informed him of it.

colore etiam immutatur, & quidem in terra Italia per id tempus alio nomine appellatur, apparet non diu, abdit enim fese & latet.

Aristoteles præter unam notam nullam ostendit peculiarem, qua ab alijs auibus luscinia differret, ea autem est quòd linguæ summæ acumine careat. Quamquam & hoc etiam cum atricapilla commune habet. Colore luscinia, & corporis magnitudine aniculam illam proximè refert, quam Angli lingettam, & Germani Graefmufch. passerem gramineum nominant. Passere paulò minor est, & tenuior, & longiori corporis figura, color pectoris ferè cinereus est, cætera subfusca.

DE MERGO.

Αἰθρία, mergus, Anglicè a cormorant, German. cyn dücher.

ARISTOTELES¹.

Mergus marina auis est, ex piscium uenatu uictitans, subit tamè altius in fluuios. Mergus [p. 90] & gauia² faxis maritimis oua bina aut terna pariunt. Sed gauiæ æstate, mergi à bruma, ineunte uere. Incubant more cæterarum auium, sed neutra earum auium conditur.

Mergus.

Mergus, auis est magnitudine ferè anseris pulla, rostro longo & in fine adunco, palmipes est, & corpore graui, forma corporis aui sedenti, erecta est. Plinius in arboribus nidulari scribit, at Aristoteles in faxis maritimis. Quod uterq; aut uidit, aut à referentibus aucupibus didicit, scripto mandauit. Et ego utrumque obseruauì, nam in rupibus marinis iuxta hostiū Tinæ fluuij mergos nidulantes uidi, & in Northfolcia cum

¹ *Hist. An.* Bk I. 6; Bk VIII. 48, freely rendered.

² *Hist. An.* Bk v. 30.

sound. It also changes colour, and during that time is known, at least in the land of Italy, by another name; it is not seen for long, since it conceals itself, and so lies hidden.

Aristotle provides no special mark, save one, by which the *Luscinia* differs from the rest of birds, and that is that it lacks the point at the tip of the tongue. Though even this it has in common with the *Atricapilla*. In colour and in size of body the *Luscinia* comes nearest to that little bird which Englishmen call Lingett and the Germans Grass-Sparrow. It is a little smaller than a Sparrow and more slender, with a longer shape of body, and the colour of the breast is nearly grey; the other parts are brownish.

OF THE MERGUS.

Αἰθρία, *mergus*, in English a cormorant, in German *eyn dücher*.

ARISTOTLE.

The *Mergus* is a sea-bird, and it lives by hunting fishes, yet it makes its way somewhat far up the rivers. The *Mergus* and the *Gavia* lay two or three eggs each upon rocks in the sea, the *Gaviæ* in summer and the *Mergi* when the spring arrives after the solstice. They incubate like other birds, but neither of these birds conceals itself.

The *Mergus*, a sad-coloured bird, is nearly equal to a Goose in size, with the bill long and hooked at the end; it is web-footed, heavy in the body, and the attitude is upright in the sitting bird. Pliny writes that it nests on trees, but Aristotle says on sea-rocks. What each man saw or learnt from the reports of bird-catchers he has set down in writing. And I have observed both birds myself, for I have seen *Mergi* nesting on sea-cliffs about the mouth of the Tyne river, and on lofty trees in Norfolk with the

ardcis in excelsis arboribus. Qui in rupibus maritimis nidificant, ex præda marina ferè uiuūt, qui uerò in arboribus, amnes, lacus, & fluuios, uictus causa petunt.

DE MEROPE EX ARISTOTELE¹.

Merops.

Sunt, qui meropes genitorum suorum senectutem educare confirmant, uicemq; reddi, ut parentes non modò senescentes, uerùm etiã cùm iam datur facultas, alantur opera liberorum: nec matrem aut patrem exire, sed in cubili manentes, pasci labore eorum, quos ipsi genuerunt, enutrierunt, educarunt. Pennæ huius auis inferiores pallidæ sunt, superiores cœruleæ sunt ut halcyonis: postremæ pinnulæ rubræ habentur. Parit sex aut septem æstare in præcipitijs mollioribus, intrà uel ad quatuor cubita subiectis, terræ etiã cauernas subiens, cunabula facit.

PLINIUS².

Nec uerò ijs minor solertia, quæ cunabula in terra faciunt, corporis grauitate prohibente sublimē petere. Merops uocatur, genitores suos reconditos pascens, pallido intus colore pennarum, superne cyaneo, priori subbrutulo. Nidificat in specu, sex pedum defossa altitudine.

Meropem ingenuè fateor me nunquam uidisse, nec quēquam cōuenisse, qui aliquando uiderit. Tametsi non sum nescius apud Germanos, grammaticos non indoctos, esse, qui grunspæchtum suum, meropem esse doceant: sed Aristotele & Plin. reclamantibus. Picus uiridis nidum

¹ *Hist. An.* Bk IX. 82, freely rendered.

² *Hist. Nat.* Lib. X. cap. xxxiii.

Herons¹. Such as make their nests on sea-cliffs generally live on prey from the sea, but such as breed on trees seek rivers, lakes, and streams to get their food.

OF THE MEROPS, FROM ARISTOTLE.

There are some who insist that Meropes foster the old age of their parents and thus take their turn, so that the parents not in age alone are nourished by the labour of their offspring, but as soon as power is given to these: that neither does the mother-bird fare forth nor yet the father, but they stay within a resting place and are fed by the aid of those which they themselves have bred, nourished and reared. The plumage of this bird is pale beneath, but blue above like that in Halcyon: the pinnules at the end of the wings are reckoned red. It lays six or seven eggs in summer in the softer banks, and makes its nurseries by boring into these for quite four cubits, and it also uses hollows in the soil.

PLINY.

Nor truly is less skill shewn by those birds which make their nurseries in the soil, since the weight of their bodies hinders them from mounting to a height. The kind called Merops feeds its parents in retreat; the colour of its feathers underneath is pale, the upper surface blue, the former being somewhat red. It breeds within a hole, bored out six feet in depth.

In fairness I admit that I have never seen the Merops, nor have I met anyone who ever saw it. Still I am not unaware that there are not unlearned schoolmasters among the Germans, who would teach us that their grunspecht is the Merops, though against the sense of Aristotle and Pliny.

¹ Compare with this Sir T. Browne's *Notes and Letters on the Natural History of Norfolk* (ed. Southwell) p. 11 (1902).

sibi roſtro ſuo in arboribus facit: ubi enim picus arborem tundēs, illam ex ſono ſubcauam eſſe depræhendit, inſtante tempore partus, eam in qua poſtea nidulaturus eſt, roſtro perforat. Nulla uſpſiam arbor tam alta eſt, quam impediēte ulla corporis grauitate, non uolatu traicere poſſit. Pennæ huius quoque ſuperiores ſunt uirides, inferioresq; niſi malè memini, lutæ aut ſaltem pallidæ ſunt. quare quum merops prohibente corporis grauitate, in ſublime petere, atq; ideo in arboribus nidulari non poſſit, & ſuperne colore ſit cyaneo, Germanorum picus uiridis, quem Britanni à faciendis foraminibus, huholam nominant, merops Ariſtotelis & Plinij eſſe non poterit.

Meropem
non eſſe
Germano-
rum grun-
ſpechtum.

DE MERULA.

Κορτυφός¹, *merula*, Angliçè *a blak ofel*, *a blak byrd*, Germanicè *eyn merl*, aut *eyn amſel*.

[p. 93]

ARISTOTELES².

Merularum duo ſunt genera: alterum nigrum & uulgare: alterum candidum, magnitudine quidem compari, & uoce ſimili, ſed circa Cylenam Arcadię familiare, nec uſquã alibi naſcens. Eſt etiam ex hoc genere, quæ ſimilis nigræ eſt, ſed fuſca colore, & magnitudine paulò minor, uerſari hæc in faxis & tectis ſolita eſt, nec roſtrum rutilum, ut merula habet. Merula³ etiam & colore, & uoce per tempora immutatur. Nam ex nigra redditur ruſa, & uocem emittit diuerſam. Strepitat enim per hyemem, quum per æſtatem tumultuans cantet.

PLINIUS⁴.

Merula, ex nigra ruſſeſcit, canit æſtate, [p. 94] hyeme balbutit, circa ſolſtitium muta, roſtrum

¹ A misprint for κορτυφός.

² *Hist. An.* Bk IX. 95.

³ *Hist. An.* Bk IX. 254; freely rendered.

⁴ *Hist. Nat.* Lib. X. cap. xxix.

Now the Green Picus makes itself a nest with its own bill in trees: for when a Picus hammering on a tree discovers by the sound that it is hollow at the core, the breeding season being close at hand, it bores that with its bill in which it afterwards intends to nest. There is not anywhere a tree so tall which this bird cannot reach by means of flight, for any weight of body that it has. Its plumage is moreover green above and, if my memory serves me, yellow underneath, or pale at least. Since then the Merops, hindered by its weight of body is incapable of rising to a height, and thus of making nests in trees, and has blue upper parts, the grunspecht of the Germans, which the Britons from the holes it makes call huhol [that is, Hew-hole], cannot be the Merops known to Aristotle and Pliny.

OF THE MERULA.

Κόττυφος, merula, in English a blak osel, a blak byrd, in German cyn merl or cyn amsel.

ARISTOTLE.

Of Merulæ there are two sorts, one black and common, and the other white, of equal size indeed and having a like voice, but which is well-known round Cyllene in Arcadia, and not bred elsewhere. There is of this kind another also, which is like the black, but dull in colour and a little less in size. It usually haunts rocks and roofs, but has not the bill ruddy like the Merula. The Merula in colour and in voice moreover changes with the season, for it turns from black to rufous, and utters a different cry. For it chatters in winter, but sings lustily in summer¹.

PLINY.

From black the Merula turns rufous, in summer it sings, but in winter it babbles, and about the solstice

¹ The readings in Aristotle differ considerably. 'Sings lustily' may go with 'in winter.'

quoq; anniculis in ebur transfiguratur, dūtaxat maribus.

DE MILVO SIVE

miluio.

ἰκτινος, *miluus*, *Anglicè*, a *gledc*, a *puttok*, a *kyte*, *Germanicè* *eyn weye*.

PLINIUS¹.

Milui ex accipitrū genere sunt, magnitudine differentes. Idem uidentur artem gubernandi docuisse, caude flexibus, in cœlo monstrante natura, quod opus esset in profundo. Milui & ipsi hybernis mensibus latent, non tamen ante hirundines abeunt. Traduntur & solstitijs affici podagra.

ARISTOTELES².

Milui pariunt bina magna ex parte, interdum & terna, totidemq; excludunt pullos. Sed [p. 95] qui Aetolius³ nuncupatur, uel quaternos aliquando excludit.

Duo miluorum genera noui, maius & minus: maius colore propemodum ruffo est, apud Anglos frequens, & insigniter rapax. Pueris hoc genus cibum è manibus in urbibus & oppidis cripere solet. Alterum genus est minus, nigrius, & urbes rariùs frequentans. Hoc genus ut in Germania sæpissimè, ita in Anglia nunquam me uidisse recordor.

DE MOLLICIPITE.

Μαλακοκρανέυς, *molliceps*, *Anglicè* a *shrike*, a *nyn* murder, *Germanicè* *eyn muin mûrder*.

ARISTOTELES⁴.

Molliceps eodem in loco semper sibi fedem statuit, atque ibidem capitur. Grădi & cartila-

¹ *Hist. Nat. Lib. x. cap. x.*

² *Hist. An. Bk vi. 38.*

³ Other readings are αἰγώλιος and ἐγώλιος.

⁴ *Hist. An. Bk ix. 98.*

it is dumb. In yearlings furthermore the bill puts on a look of ivory, provided they are males.

OF THE MILVUS OR MILVIUS.

ἰκτινος, milvus, in English a glade, a puttok, a kyte, in German eyn weye.

PLINY.

Milvi are of the race of Accipitres, though differing in size. They seem, moreover, to have taught mankind the art of steering, by the turning of the tail, nature thus shewing in the sky what might be useful in the sea. Milvi lie hidden in the winter months, yet not until Hirundines depart. They are reported also to be affected with the gout about the solstice.

ARISTOTLE.

Milvi lay for the most part two eggs each, but sometimes three, and hatch as many young. But that kind which is named Ætolian at times lays even four.

I know two sorts of Kites, the greater and the less; the greater is in colour nearly rufous, and in England is abundant and remarkably rapacious. This kind is wont to snatch food out of children's hands, in our cities and towns. The other kind is smaller, blacker, and more rarely haunts cities. This I do not remember to have seen in England, though in Germany most frequently.

OF THE MOLLICEPS.

Μαλακοκρανέϋς, mollicept, in English a shrike, a nyn murder, in German eyn nuin mürder.

ARISTOTLE.

The Mollicept invariably takes its stand in the same place, and thereat it is caught. It has a big

gineo capite est, magnitudine paulò minor quã turdus, ore firmo, paruo, rotũdo, colore totus cinereo depes¹, & pēnis inualens est, capitur maximè noctua.

Mollicipitem esse arbitror auiculam, quam Germani nuinmurder non sine causa nominant. Porro ut omni-
 [p. 96] *bus perspicuum sit, quænam & qualis illa sit, formam auis & mores quanto licebit compendio perstringam. Magnitudine, minimum turdorum genus æquat, è longinquo contēplanti, tota apparet cinerea. Propius autem insipienti, mentum, pectus & uenter alba apparent, ab utroque oculo ad collum usque, longa & nigra macula, sed nonnihil obliqua porrigitur. Capite tam grandi est, ut aui triplo maiori (modò rostrum longius & maius esset) proportionem suam satis responderet. Rostrum nigro est, & mediocriter breui, & in fine adunco, sed omnium firmissimo & fortissimo est, utpote quo manum semel meam duplici chirotheca munitam, sauciauerit, & auium ossa & capita confringat & conterat quàm ocysimè. Ala utraque nigra tota est, nisi quòd alba linea grandiuscula, mediam utrinque alam transuersim distinguit. Caudam picæ similem habet, lōgiusculam nimirum, & uariam. Tibias & pedes pro ratione corporis omnium minimos, & eos nigros habet. Alas habet breues, & ueluti per saltus sursum atque deorsum uolitat. Viuit ex scarabeis, papilionibus, & grandioribus insectis: sed non solis istis, uerumetiam, more accipitris, auiibus. Occidit enim regulos, fringillas, & (quod ego semel uidi) turdos. Tradunt etiam aucupes hanc picas quasdam*
 [p. 97] *syluestres interdum iugulare, & cornices in fugam adigere. Aues, quas occidit, non unguibus, ut accipitres, uolando perniter adsequitur, sed ex insidijs adoritur, & mox (quod iam sæpius expertus sum) iugulum petit, & cranium rostro comprimit & confringit. Ossa comminuta & contusa deuorat: & quando esurit, tantos carnis bolos in gulam ingerit, quantos riētus oris angustia potest capere. Præter morem etiam reliquarum auium, quando uberior præda contigit, nonnihil in fu-*

Offisfraga
dici possit,
si eius illi
magnitudo
adeffet.
Nam nec
moribu⁹, nec

¹ ἄπovs. Another reading is εὔπovs.

and gristly head, and is a little smaller than a Thrush in size ; the bill is strong but small, and curved ; in colour it is wholly grey, while it is weak-footed and feeble on the wing, it is caught chiefly by the Noctua.

The Mollicept I think to be that little bird which Germans call *nuinmurder*, not without a cause. Further that it may be quite clear to all which and what sort of bird it really is, I will touch on its form and habits as compendiously as may be. In size it equals the least of the Thrushes, and to one observing from afar seems wholly grey. And yet, to one inspecting it more nearly, the chin, the breast and belly appear white, and from each eye there reaches to the neck, although somewhat oblique, a long black patch. It has so big a head that (were the bill longer and larger) it assuredly would answer in proportion for a bird of thrice its size. The bill is black and moderately short, and hooked at the tip, but is the stoutest and strongest of all, so much so that the bird once wounded my hand, although protected by a double glove, and very speedily it crushes and breaks up the bones and skulls of birds. Each wing is wholly black, except that a white line of some size marks transversely the middle of the wing on either side. The tail is like that of a Pie, that is to say, longish and particoloured. Of all it has the shortest legs and feet proportionately to its body, and these parts are black. It has short wings, and flies as if by bounds upwards and downwards. It lives on beetles, butterflies, and biggish insects, and not only these, but also birds after the manner of a Hawk. For it kills *Reguli* and Finches and (as once I saw) Thrushes ; and bird-catchers even report that it from time to time slays certain woodland Pies, and can put Crows to flight. It does not seize the birds it kills with its claws, after a swift flight, as Hawks do, but attacks them stealthily and soon (as I have often had experience) aims at the throat and with its beak squeezes and breaks the skull. Then it devours the crushed and bruised bones, and when anhungered crams into its gullet lumps of flesh as big as the gape's narrowness can take. Again, beyond the habit of the rest of birds, when prey happens to be more plentiful, it lays by some for future scarcity.

colore ab ea
multum
abludit.

turam penuriã reponit. Muscas enim grandiores & insecta iam capta in aculeis & spinis arborum figit & suspendit: omnium animi facilius cicuratur, & mansueta, carnibus alitur, quæ si fuerint sicciore, aut prorsus exangues, potum requirit. In Anglia sæpius quàm bis nunquam uidi, in Germania sæpissimè. Nomen huius apud nostros neminem inueni, qui nouerit, præter Dominum Franciscum Louellum, tam animi quàm corporis dotibus equitem auratum nobilissimum. Iam si cui mollicipitis Aristotelis descriptio huic non uidetur per omnia conuenire, tyrannorum albo adscribat, aut auem ostendat, cui descriptio melius competat.

DE NOCTVA.

Γλαύξ, noctua, Anglicè an ouul, or an houlet,
[p. 98] Germanicè eyn eul & eyn ùle Saxonice.

ARISTOTELES¹.

Noctuæ, cicuniæ², & reliqua, quæ interdiu nequeunt cernere, noctu uenando cibum sibi adquirunt: uerum non tota nocte id faciunt, sed tempore uespertino et matutino. Venantur autem mures, lacertas, uerticillos, & eiusmodi bestiolas. Noctuam³ cæteræ omnes aues circumuolant, quod mirari uocatur, aduolantesque percutiunt⁴. Qua propter aucupes ea constituta, auicularum genera multa & uaria capiunt.

DE OLORE.

Κυκνὸς, olor, Anglicè a fuuan, Germanicè eyn swān.

ARISTOTELES⁵.

Olor.
Olores palmides⁶ sunt, apud lacus & paludes uiuentes, qui nec probitate uictus, morum, pro-
[p. 99] lis, fenectutis uacant⁷. Aquilam si pugnam

¹ *Hist. An.* Bk IX. 122.

² Apparently a misprint for cicumæ, said to mean 'horned owls.' Aristotle has *νυκτικόρακες* which he identifies in Bk VIII. 84 with *ὥτοί*.

³ *Hist. An.* Bk IX. 11.

⁴ Aristotle has *τῶλονσι* = pluck it.

⁵ *Hist. An.* Bk IX. 78.

⁶ That is, palmipedes.

⁷ Aristotle has *εὐβίοτοι δὲ καὶ εὐήθεις καὶ εὐτεκνοὶ καὶ εὐγῆροι*.

For it impales and hangs the bigger flies and insects on the thorns and spines of shrubs, so soon as they are caught: of all birds it is tamed most easily, and when accustomed to the hand is fed on meat, and, should this happen to be somewhat dry or altogether bloodless, it requires drink. In England I have never seen it oftener than twice, although most frequently in Germany. Among our people I have found no one who knew its name, except Sir Francis Lovell, that most noble knight, endowed with equal gifts of mind and body. Now if Aristotle's description of the *Molliceps* does not appear to any one in all points to agree with this let him ascribe it to the list of the *Tyranni*, or shew us a bird, which the description fits better than this.

OF THE NOCTUA.

Γλαύξ, noctua, in English an owl or an howlet, in German cyn eul, and in Saxon eyn ũle.

ARISTOTLE.

The Noctuæ, Cicumæ and the rest, which cannot see by day, obtain their food by seeking it at night: and yet they do not do this all night long, only at eventide and dawn. They hunt moreover mice, lizards, and scorpions, and small beasts of the like kind. All other birds flock round the Noctua, or, as men say "admire," and flying at it buffet it. Wherefore this being its nature¹, fowlers catch with it many and different kinds of little birds.

OF THE OLOR.

Κύκνος, olor, in English a swan, in German cyn swän.

ARISTOTLE.

Olores are web-footed, and they live on lakes and marshes; they get food with ease, are peaceable, prolific and attain to a great age. They repulse the

¹ Or, possibly, 'the bird being set down on the ground.'

cœperit, repugnantes uincunt. Ipſi tamen nunquam, niſi prouocati, pugnam inferunt. Canere ſoliti ſunt, & iamiam morituri. Volant etiam in pelagus longius, & iam quidam cùm in mari Africo nauigarunt, multos canentes uoce flebili & mori nonnullos conſpexêre.

Si quis olorem nunquam uiderit, & ex hac Ariſtotelis deſcriptione non ſatis qualis ſit auis didicerit, ſciat auem eſſe albam, anſere multò maiorem, forma tamen & uictu ſimilem, pedibus nigris, & roſtro parùm turbinato, colore rutilo, in cuius ſumma parte, qua capiti committitur, nigerimum tuberculum, atque id rotundum, & in roſtrum ſeſe inflectens, exiſtit.

DE ONOCRATALO.

Onocrotalus. *Sunt hodie non parùm multi eruditione inter omnes conſpicui, qui grandifonam illam lacuſtrem auem, Anglis buttoram & Germanis pittourum, & roſdommam uocatam, Onocrotalum eſſe contendunt. Quorũ*
 [p. 100] *ego ſententiæ lubens ſubſcriberem, (pulchrè enim cum uoce auis nominis etymologia conuenit:) niſi Plinij autoritas de onocrotalo ad hunc modum ſcribentis, non diſſuaderet. Onocrotali, inquit, olorum ſimilitudinem habent, nec diſtare uidentur omnino, niſi faucibus ipſis ineſſet, alterius uteri genus. huc omnia inexplabile animal congerit, mira ut ſit capacitas, mox perfectâ rapina, ſenſim inde in os reddita, in ueram aluum ruminantis modo refert. Gallia hos ſeptentrionalis, proximè¹ oceano mittit. Hæc Plinius².*

¹ This ſhould apparently be ‘proxima,’ as ſome texts have it.

² *Hist. Nat. Lib. x. cap. xlvii.*

Aquila successfully, should he begin a fight ; and yet, unless provoked, never induce the fight. These birds are wont to sing even when just about to die. They also fly afar over the main, and men ere now, who have been sailing on the African sea have met with many singing mournfully and seen some of them die.

Should any one have never seen a Swan, nor learnt sufficiently what sort of bird it be from this account of Aristotle, let him know that it is a white bird, much bigger than a Goose, though like in form and feeding ; with black feet, and a bill hardly spindle-shaped¹, reddish in colour ; on the highest part of which, where it adjoins the head, stands forth a very black and rounded knob, sloping towards the bill.

OF THE ONOCROTALUS.

There are many to-day conspicuous among all for learning to no small degree who maintain that the loud-sounding lacustrine bird, called Buttor by the English, and Pittour or Rosdomm by the Germans, is the *Onocrotalus*. To whose opinion I would willingly subscribe, (the more so as the etymology of the bird's name agrees well with its voice,) did not the authority of Pliny writing of the *Onocrotalus* after this manner dissuade me therefrom.

The *Onocrotali*, he says, have a similitude to the *Olores*, and they do not seem to differ in any way, save that there is a kind of second belly in the very jaws. Herein the insatiable animal crams everything at once, so marvellous is its capacity, and presently, the plundering complete, it gradually returns all to the mouth, and thence transfers it to the real belly in the manner of a ruminant. Northern Gaul, where nearest to the ocean, sends us these. So far Pliny.

¹ This passage is not easily rendered, as it is difficult to see what Turner intended by 'turbinato.' Turbo is a conical shell, spindle and so forth ; but it is hard to say how a Swan's beak could be considered either conical or spindle-shaped.

Nunc paucis auem illam uobis depingam, quā onocrotalum esse affeuerant. Auīs est tota corporis figura Ardeæ similis. longis cruribus, sed ardeæ breuioribus. longo collo, & mirè plumoso, & rostro nec breui nec obtuso. caput pennæ tegunt nigerrimæ. reliquum uerò corpus, fusca & pallida maculis nigris densissimè respersæ. Pedes habet longissimos, nam inter extremos ungues medijs digiti pedis unius & calcis eiusdem, [p. 101] *spithames longitudo intercedit. Ungues habet longissimos, nam ille, qui calcis uicem in auibus gerit, longitudine sesquiunciam superat. quare ad fricandos dentes nostrates utuntur, & argento inferunt. Medius digitus utriusque pedis, qui cæteris longior est, unguem habet portentosum, nempe dentatum & ferratum, non secus atque pectunculorum testæ ferratæ sunt, ad lubricas anguillas, quas cepit¹, retinendas, à natura proculdubio ordinatum. Cauda illi breuissima est, et stomachus capacissimus, quo ingluuiei loco utitur. Ventriculum non cæterarum auium uentriculis, sed canino similem habet, & cum grandem & capacem.*

Sed ne cui falsa esse uideantur, quæ de hac aue iam scripsi, aut ex aliorum relatu potiùs quàm certa experientia didicisse uidear: dum prima huius libri folia adhuc sub prælo essent, auem mihi hanc contemplanti, secantiq;, & nū tales haberet uentriculum & stomachum, quales Plinius illi tribuit, inuestiganti: aderant uir eruditissimus, & abstrusorum naturæ arcanorum studiosissimus inuestigator Ioannes Echthius, Medicus apud Colonienses celeberrimus: Cornelius Sittardus, Medicinæ prima laurea decoratus. M. Lubertus Estius, artium liberalium professor, ambo simplicium medicamentorum pulchrè gnari, & ad miraculum usque studi- [p. 102] *osi: & Conradus Embeccanus uir non uulgariter doctus, et Gymnicinæ officinæ castigator insigniter diligens, cum alijs aliquot bonarum artium studiosis, qui me nihil de hac aue hîc scripsisse testari possunt & uolunt, quod cum illis omnibus non uiderim. Ad ripas lacuum & paludium desidet, ubi rostrum in aquas in-*

¹ A misprint for 'cepit.'

Now in a few words I will portray to you that bird which they assert to be the *Onocrotalus*. In general make of body it is like the Heron, with long legs, though shorter than that bird's. The neck is long and marvellously thick with plumes, the beak is neither short nor blunt. Very black feathers clothe the head, but on the body generally they are dusky and pale, and most thickly sprinkled with black spots. It has very long feet, indeed there is a span's length from the claw-tips of the middle toe of either foot to the heel of the same. It has very long claws, for that which serves in birds the purpose of a heel exceeds an inch and a half in length, on which account our countrymen use it to pick their teeth, and mount it in silver. The middle toe of either foot, which is longer than the rest, has a prodigious claw, that is to say, toothed and serrated, not unlike the shells of little scallops are, doubtless contrived by nature to retain the slippery eels, which the bird catches. The tail is very short, the gullet most capacious, and it uses it in the place of a crop. It has a belly not like that of other birds, but like that of a dog; it also is large and capacious. But lest what I have written thus far of this bird seem false to anyone, or lest I seem to have learnt the above from the reports of others rather than from sure experience: while the first pages of this book were still at press, and while I was examining the bird and was dissecting it, and taking note whether it really had a belly and a stomach such as Pliny had assigned to it, there were assisting me Joannes Echthius, a very learned man and a most zealous student of the more abstruse secrets of nature, a physician much renowned among the men of Cullen: Cornelius Sittardus decorated with the highest laurel-wreath of Medicine: Marcus Lubertus Estius, professor of the liberal arts, both excellently skilled in that of simpling, and wonderfully earnest, and as well as these Conradus Embecanus, a man well-informed in no common degree, and a remarkably careful corrector in the printing-house of Gymnicus, with certain others versed in learned arts, who can and will bear witness to the fact that I have written nothing here about this bird which I have not observed in company with all of them. It sits about the sides of lakes and marshes, where putting

ferens, tantos édit bombos, ut ad miliarium Italicum facile possit audiri. Pisces & præsertim anguillas uorat auidissimè, nec ulla auis est, excepto mergo, quæ ista uoracior est. Nunc quid simile habet ista cygno? Nihil planè, quod se oculis conspiciendum offerat. Et Moïses Leuit. undecimo capite, proximè cygnum inter immundas aues onocrotalum recenset. Vnde non immeritò suspicio quibusdam orta est in Gallia, aut Iudæa auem forma olori similem alicubi posse reperiri. Quòd si nusquam talis inueniatur: probabile est, aut Plin. à mendacibus relatoribus suis esse falsum, aut ea, quæ de similitudine inter onocrotalum & cygnū tradidit, non de corporis sed uocis similitudine intellexisse. Nam & olores interdum bombos emittunt ruditui asinuino non dissimiles: sed breues, & quæ longè audiri non possunt. Verùm si hanc meam interpretationē uariæ, reconditæq; eruditionis uiri, suis suffragijs minimè approbauerint, [p. 103] hanc saltem Aristotelis ardeam stellarem esse mecum consentient. Nam præter cætera, quæ superius attigi, Aristoteles in fabula fuisse ostendens, ardeam stellarem ex seruo auem fuisse factam, opinioni meæ multum patrociniatur. Vt fugitiuorum enim seruorū post fugam depræhensorum, cutis, loris, flagris, uirgis, & scorpionibus icta, uerberum uibicibus, tota maculosa redditur: ita huius auis plumæ nigris ubique maculis, sed potissimum in tergo, distinctæ & ucluti picturatæ, serui flagris cæsi cutem proximè referūt. Quam rem fabulæ occasionem dedisse ex hoc colligo, quòd fabularum uariarum autor Aristophanes¹, de attagene aue, quod ad plumarum colores attinet, huic similima, ad hunc modum scribat:

*Si quis ex uobis crit fugitiuus atq; uflis notis,
Attagen is sanè apud nos uariius appellabitur.*

DE ORTYGOMETRA EX ARISTOTELE².

Ortygometra, id est, coturnicū matrix, auis est forma perinde ac lacustres. Cruribus ideo

Onocrotalum quibusdam hodie dictū, ardeam, esse stellarem apud Aristotelem.

¹ See p. 36.

² *Hist. An.* Bk VIII. 83, freely rendered, and interpolated.

its beak into the water it gives utterance to such a booming as may easily be heard an Italian mile away. It gorges fishes and especially eels most greedily, nor is there any bird, except the Mergus, that devours more. Now what resemblance has it to a Swan? Distinctly none that brings itself in view before our eyes. Now in the eleventh chapter of Leviticus Moses enumerates the *Onocrotalus* next to the Swan among the unclean birds. And a suspicion has arisen thence, not undeservedly, within a certain class, that somewhere within Gaul or Judæa a bird of Swan-like form may possibly exist. If such, however, nowhere can be found, it seems likely that Pliny either was deceived by lying story-tellers or he understood that which he has related of the similarity between the *Onocrotalus* and *Cygnus* to refer to a resemblance not of body, but of voice. For even Swans utter at certain times booms not unlike the braying of an ass: but short, and which cannot be heard afar. However if men of deep and varied learning by their votes shall not approve this rendering of mine, at least they will agree with me that the said bird is Aristotle's *Ardea Stellaris*. For to omit the rest, which I have touched upon above, that author certainly gives countenance to my opinion when he shews a tale to have existed that the *Ardea Stellaris* from a slave was turned into a bird. For as the skin of an absconding slave, caught subsequent to flight, stricken with thongs, whips, rods and knotted ropes, becomes all mottled with the wales of stripes, so too the feathers of this bird are marked, and painted as it were, with mottlings of black in every part, though chiefly on the back, and thus may well recall to us the skin of slaves cut up with whips. And that this thing gave rise to the aforesaid tale, I gather from the fact that Aristophanes, author of various plays, writes of the *Attagen*, a bird very like ours so far as colour of the feathers goes, to this effect:—

“If any of you be a runaway, and branded with the marks, with us assuredly he shall be called the spotted *Attagen*.”

OF THE ORTYGOMETRA FROM ARISTOTLE.

The *Ortygometra*, that is, dam of the *Coturnices*, in form is much like marsh-birds. Certain birds are

longis aues quædam innituntur, quòd earum uita fit palustris.

[p. 104] *Ortygometram aliqui eandem esse auem cum crece et cychramo uolunt. Sed Aristoteles, peculiare caput crece donauit, et octauo libro historiæ animalium, cychramum a matrice, quam ertygometram uocat, his uerbis distinguit. Coturnices (inquit) cum hæc adeunt loca, sine ducibus pergunt: at cum hinc abeunt, ducibus lingulaca, oto, & matrice, proficiscuntur, atque etiam cychramo, à quo etiam reuocatur noctu, cuius uocem cum senserint aucupes, intelligunt parari discessum. Hæc ille.*

Fieri igitur non potest, ut matrix & cychramus eadem auis sit. Aliqui ertygometram esse uolunt Germanorum scricam, & Anglorum daker hennam, quorum ego sententiæ accederem, si crecem eandem cum ista, euincerent.

DE OSSIFRAGA.

ARISTOTELES¹.

Offifragæ magnitudo maior est quàm aquilæ, color ex cinere² albicans. Probè³ & sætificat, & [p. 105] uiuit, cænæ gerula & benigna est. Nutricat enim bene, & suos pullos & aquilæ. Cum enim illa suos nido eiecerit, hæc recipit eos, ac educat.

PLINIUS⁴.

Quidam adijciunt genus aquilæ, quam barbaram uocant Thusci offifragam.

¹ *Hist. An.* Bk VIII. 39.

² Apparently a misprint for 'cinereo.'

³ *Hist. An.* Bk IX. 123.

⁴ *Hist. Nat.* Lib. X. cap. iii.

perched upon long legs because their life is passed in marshes.

Some will have *Ortygometra* to be the same as *Crex* and *Cychramus*. But Aristotle has attributed a peculiar sort of head to *Crex*, and in the eighth book of his *History of Animals* distinguishes his *Cychramus* from *Matrix*, which he calls *Ortygometra*, in the following words :—

Coturnices (he says) when they come to these places travel without guides : but when they go away set out with the *Lingulaca*, the *Otus*, and the *Matrix* as their guides, and also with the *Cychramus*, by which they are moreover summoned back at night. And when the fowlers have heard its cry, they know the birds' departure is at hand.

Thus he writes.

Therefore it is impossible that the *Matrix* and the *Cychramus* should be the same. Others will have the *Ortygometra* to be the *Scrica* of the Germans and the *Daker Hen* of the English, and I should accede to their opinion, if they could but prove the *Crex* to be the same as this.

OF THE OSSIFRAGA.

ARISTOTLE

In size the *Ossifraga* is greater than the *Aquila*, its colour whitish grey¹. Both in breeding it is comely and in way of life, it brings food home and is kindly. For it rears its own young with care, besides those of the *Aquila*. For when the latter has cast its progeny out of the nest, the former takes them to itself, and brings them up.

PLINY.

Some there are who add that kind of *Aquila*, which the Tuscans call a bearded *Ossifrage*².

¹ See p. 36.

² Possibly Pliny means the Lämmergeier (*Gypaëtus barbatus*).

DE OTO.

ὠτός, *otus*, *Anglicè a horn owl*, *Germanicè* eyn ranieul / oder eyn schleier eul.

ARISTOTELES¹.

Otus noctuæ similis est, pinnulis circiter aures eminentibus, præditus, unde nomen accepit, quasi auritum dixeris. Nonnulli ululam eū appellant, alij asionem². Blatero hic est & hallucinator, & planipes: faltantes enim imitatur. Capitur intentus in altero aucupes³, altero circumeunte.

[p. 106]

DE OTIDE EX PLINIO⁴.

Tetraonibus proximæ sunt, quas Hispania aues tardas appellat, Græcia otidas, damnatas in cibis. Emissa enim ossibus medulla, odoris tædium extemplò sequitur.

DE PARIS.

ἀγριθολός, *parus*, *Anglicè a tit mouse*, *German.* eyn meysje.

ARISTOTELES⁵.

Parorum tria sunt genera: fringillago, quæ maior est, quippe quæ fringillam æquet. Alter monticola cognomine est: quoniam in montibus degat, cui cauda longior. Tertius magnitudine sui exigui corporis discrepat, quanquam cætera similis est. parus⁶ plura oua parit.

Fringillago.

Primum parū, Angli uocant the great titmouse or the great oxei, Germani eyn fōlmeysje.

Parus medius.

Parum secundum, Angli the lefs titmouse nominant. Germanici eyn meelmeysje.

¹ *Hist. An.* Bk VIII. 84. very freely rendered.

² Aristotle has 'νυκτικόρακα,' instead of 'ululam' and 'asionem.'

³ A misprint for 'aucupe.'

⁴ *Hist. Nat.* Lib. X. cap. xxii.

⁵ *Hist. An.* Bk VIII. 40.

⁶ *Hist. An.* Bk IX. 88.

OF THE OTUS.

ὠτός, otus, in English a horn owl, in German cyn ranseul or cyn schleier eul.

ARISTOTLE.

The Otus is like a Noctua, furnished with little tufts sticking out near the ears, whence it has got its name, as though one should say "eared." Some call it Ulula, and others Asio. It is a babbler and a mischievous rogue, and is a mimic too, for when men dance it imitates their ways. It is caught while intent upon one of two bird-catchers, the other circumventing it.

OF THE OTIS FROM PLINY.

Next to the Tetraones come those birds, which Spain calls "*Aves tardæ*" and Greece "*Otides*," condemned as food for man. For when the marrow issues from the bones, disgust at the smell follows there and then.

OF THE PARI.

αἰγιθαλος, parus, in English a titmouse, in German eyn meyse.

ARISTOTLE.

There are three kinds of Pari: *Fringillago* bigger than the rest, for it is equal to a *Fringilla*. The next *Monticola* by name, for it inhabits mountains, has a longer tail. The third kind differs in the size of its small body, though not otherwise unlike the rest. Parus lays many eggs,

The first Parus the English call the Great Titmouse or Great Oxeye, the Germans eyn kölmeyse.

The second Parus the English name the Less Titmouse, the Germans eyn meelmeyse.

[p. 107]
Parus mi-
nimus.

Parum tertium, Angli nonnam à similitudine quam cum uclata monacha habet, nominant.

Nidulantur pari in cauis arboribus, uescuntur non solùm uermibus, sed & canabino semine, & nucibus, quas rostris suis acutioribus solent perforare, & nucleos erucere. Sæuo duo priora genera multùm delectantur. Parus maximus ineunte statim uere cantiunculam quandam breuem, nec admodum iucundam exercet, aliàs mutus, huic pectus luteum est, intercurfante linea nigra maiuscula. Cæterorum corpora albo, nigro, pallido, & cyaneo coloribus distinguuntur.

DE PARDALO.

Pardalus, Angl. (ut creditur) a pluuer, Germa. eyn puluer.

ARISTOTELES¹.

Pardalus etiam auicula quædam perhibetur, quæ magna ex parte gregatim uolat, nec singularè hanc uideris, colore tota cinereo est, magnitudine proxima mollicipiti² est: sed pennis & pedibus bonis, uocem frequentem nec grauem emittit.

[p. 108] *Si auis illa pardalus sit quam esse suspicor, celerimè currit, & sibilum, quem pastores & aurigarum pueri labijs porrectis edunt, uoce imitatur. Pennas habet ad cinereum colorem proximè uergentes, quarum singule singulis flauis maculis sunt respersæ, & ea auicula, quam mollicipitè esse conijcio, multò maior est. Fieri potest, ut eius auis plures sint species.*

DE PASSERIBVS.

στρούθος, passer, Angli. a sparrow, German. eyn müſche oder eyn ipāß. Quidam eyn lüningf, Saxones eyn ipertling uocant.

Passer, authore Aristotele³, puluerat & lauat, et auis est omnium salacissima. Et quanquam Aristoteles unum tantùm passerum genus fecerit, tria tamè genera esse constat, quæ nunquam nouit, & quæ recentiores

¹ *Hist. An.* Bk IX. 99.

² Aristotle has ἐκείνους, referring to the χλωρίων and the μαλακοκρανέυς.

³ *Hist. An.* Bk IX. 260, Bk V. 8.

The third Parus the English name the Nun from the resemblance that it bears to a veiled sister.

The Pari nest in hollow trees, they feed not only on worms, but on hempseed and nuts, which they are wont to bore with their sharp-pointed beaks, and thence extract the kernels. The two former kinds are very fond of suet. The Greatest Parus, when the spring arrives, at once utters a sort of little song, short and not very pleasing, it is dumb at other times; its breast is yellow with a somewhat big black line running along the middle. Of the other kinds the bodies are diversified by white, black, grey, and blue.

OF THE PARDALUS.

Pardalus, in English (as is believed) a pluver, in German cyn pulver.

ARISTOTLE.

The Pardalus again is held to be a certain little bird, which for the most part flies about in flocks, and cannot be seen solitary; it is wholly grey in colour, and in size comes nearest to the Mollicept: but it has strong wings and feet, and utters a frequent but not deep-toned cry.

If that bird be the Pardalus which I suspect, it runs very swiftly, and by its cry mimics the whistle which shepherds and post-boys make with pouting lips. It has the feathers almost ash-colour, each sprinkled with one yellow spot, and is much bigger than the little bird which I suppose to be the Mollicept. It well may be that there are several kinds of this bird.

OF THE PASSERES.

στρουθός, passer, in English a sparrow, in German cyn müsche or cyn spätz. Some call it cyn lüningk, the Saxons eyr sperlingk.

The Passer, Aristotle says, both dusts itself and washes, and is of all birds most wanton. And though Aristotle has made only one kind of Passeres, yet it is clear that there are three kinds, which he never knew; but which the later

Passer tor-
quatus.

Passer mag-
nus.

Huic mol-
licipitis apud

[p. 109] Aristotelem
descriptio
magna ex
parte con-
uenit.
Passer tro-
glodites.

Græci inuenerunt. Primum horum trium est passer torquatus, à communi passere, nō solū torque albo, sed & uoce, & modo nidificandi differēs. Hoc genus in Germania frequens est, sed apud Anglos rarum. Secundus passer magnus Aucuario dicitur, & in summis arborum ramis plerumque solet sedere. hunc narijs de causis Anglorum buntingam, & Germanorum Gersthammeram esse suspicor. Tertius passer Aristoteli incognitus, est passer troglodites, apud Paulum Aeginetam, & Aetium celeberrimos medicos, multū celebratus. Qualis autem illa avis sit, ex Paulo & Aetio, quorum descriptiones mox subijciam, omnibus facili patebit.

PAULUS AEGINETA DE PASSERE

TROGLODITE.

His accedit laudatissimum remedium troglodites. est autem passerculus omnium auium minima, ea sola excepta, quæ regulus appellatur, hoc enim solo paulò maior est, eiq; similis: colore inter cinerium & uiride, tenui rostro, in muris maximè & in sepibus degens.

AETIUS.

Troglodites est passerculus minimus, iuxta sepes & muros uictum quæritans. Estq; hoc animalculum omnium auicularum minimum, ex [p. 110] cepta ea, quæ regulus appellatur, similis autem regulo in multis, præterquam quòd in fronte auricolores pennas non habet. Est autè troglodites passer regulo paulò maior & nigrior, caudamq; semper subrectam, & albo colore retrò interpunctam habet. Magis item garrulus

Greeks discovered. First of these three is *Passer torquatus*, differing from the common *Passer* not alone in its white collar, but also its note and mode of nesting. This kind is plentiful in Germany, but rare among the English. The second *Passer* is called in the Supplement¹ the Great, and for the most part it is wont to sit on the top boughs of trees. For several reasons I consider this to be the Bunting of the English and the Gersthammer of Germans. The third *Passer*, unknown to Aristotle, is the *Passer troglodytes*, fully recognised by Paulus Ægineta and Aëtius, doctors of great renown. And so what sort of bird it is will easily be seen by everyone from Paulus and Aëtius, and their descriptions I will forthwith add.

PAULUS ÆGINETA² OF THE PASSER TROGLODYTES.

There is a remedy most highly prized besides these, namely *Troglodytes*: this is nothing but a little Sparrow, the least of all birds, with the exception only of that kind which is called *Regulus*. It is a little bigger than that bird alone, and similar to it: in colour between grey and green, and with a slender bill. It lives chiefly in walls and hedges.

AËTIUS³.

The *Troglodytes* is the very least of Sparrows, seeking for its food near hedges and near walls. This little animal moreover is the smallest of all little birds, except that which is called *Regulus*, while it is like the *Regulus* in many ways, save that it has not golden-coloured feathers on the forehead. The *Passer troglodytes* is a little larger and blacker than the *Regulus*; it always has its tail cocked up, which is spotted behind with white. Likewise it is more noisy than the

¹ It seems impossible to ascertain what this Auctuarium was.

² A medical writer of Ægina, whose chief work was *De Re Medica Libri Septem*.

³ A Greek medical writer of Amida in Mesopotamia who wrote *Βιβλία Ἰατρικὰ Ἑκκαίδεκα*.

quàm regulus est, & fanè iuxta summum alæ lineamentum cinerij amplius coloris. Breues item facit uolatus, naturalem autem uim omnino admiratione dignam habet.

Nihil est in hac descriptione, quod non ad amussim auiculæ conueniat, quam Angli passerem sepriarium, Colonienses aucupes koelmusslum nominant. Sed quoniam tam in Germania quàm in Anglia uarijs nominibus appellatur, & non omnes eum ex uno nomine agnoscunt, omnia eius, quæ noui nomina, ut omnibus innotescat, subiiciam.

Vocatur apud Anglos an hedge sparrow, hoc est passer sepriarius, & a dike smouler, hoc est, in sepibus [p. 111] delitescens. Vulgus Coloniëse hunc passerem eyn gras-musch appellat. uerū peritiores quiq: aucupes eyn koelmusch, hoc est, passerem in foraminibus & cauernis degentem, nuncupant. Hic Germanos monitos uolo, quò duæ sint aues, grasmuschi, sua lingua uocatæ, illā solā esse trogloditen, quæ per totum annum regulo similis cernitur, & non illam, quæ circa fauces plumosa¹ inenunte statim hyeme discedit. Nidum huius passeris semel humi factum inter urticas uidi, & pullos antequam uolare possunt, relicto nido, inter herbas fruticesq: reptitantes, sæpius obseruaui: uermibus pascitur, & paulò ante uesperum solet impensius strepere, & omnium ferè auium postrema dormitum petit.

DE PAVONE.

Ταῶν, pavo, Anglicè a peccok, Germanicè eyn pffaw. Saxonice eyn pagelūn.

PLINIUS².

Pauo gemmantes laudatus expandit colores, aduerso maximè sole, quia sic fulgentius radiant, simul umbræ quosdam percussus cæteris, quæ [p. 112] in opaco clariùs micat, conchata quærit cauda, omnesq: in aceruum contrahit pēnarum, quos

¹ This is probably a misprint for some other word.

² *Hist. Nat. Lib. x. cap. xx.*

Regulus, and certainly towards the upper border of the wing the colour is more nearly grey. And though it takes short flights, its natural energy is worthy of all admiration.

In this description there is nothing that does not agree exactly with the little bird, which Englishmen name *Passer sepiarius*, the bird-catchers of Cullen the *koelmusch*. But inasmuch as both in Germany and England it is called by various names, and all men do not recognise it by the same, I will subjoin those of its appellations which I know, that so it may be known to all.

By the English it is called a Hedge-Sparrow, which is the same as *Passer sepiarius*, and also a Dike Smouler, one, that is, hiding itself in hedges. The common people of Cullen call it *cyn grassmusch*, but those who know better and fowlers name it *cyn koelmusch*, that is, a Sparrow dwelling within holes and caverns. Now here I wish the Germans to be warned, that since there are two birds called *grasmusch* in their tongue, the *Troglodytes* is that kind alone which throughout the year is noticed to be like the *Regulus* and not that which is feathered round the jaws, and goes away so soon as winter comes. Of this *Passer* I once met with a nest built on the ground among nettles and I have often seen young having left the nest before that they could fly, creeping among the grass and shrubs. It feeds on worms, and it is wont a little before evening to cry out with not a little vehemence; it goes to roost almost the last of all the birds.

OF THE PAVO.

Taóv, pavo, in English a peacock, in German *cyn pffaw*, in Saxon *cyn pagelün*.

PLINY.

The Peacock is admired for setting forth his jewelled colours, generally counter to the sun, since thus they shine the brighter, while with concave tail he gains certain reflexions of the shade for other feathers which shine brighter in the dark, and at the same

spectare¹ gaudet oculos. Idem cauda annuis uicibus amissa cum folijs arborum, donec renascatur iterum cum flore: pudibundus ac mœrens quærit latebram. Viuit annis uiginti quinque. Colores incipit fundere in trimatu. Ab auctoribus traditur non tantum gloriosum animal, sed maleuolum, sicut anser uerecundum. Pauones in capitibus suis ueluti crinita habent arbusculâ.

DE PERDICE.

Πέρδιξ, *perdix*, *Anglicè a pertridge*, *Germanicè* eyu velt hön/ *ader eyu raphôn*.

ARISTOTELES².

Perdix auis est pulueratrix, & non altiuola, & eadem non in nido sed in condensa frutice [p. 113] aut segete prolem suam munit. Aues enim grauiore nidos sibi non faciunt, ut coturnices & perdices, & reliquæ generis eiusdem. Quibus enim uolandi facultas deest, ijs nidus non prodest: sed facta in aprico³, area, (alibi enim nusquam pariunt) atque materia ut uepribus⁴ quibusdam congestis, quoad accipitrum & aquilarum iniuriam deuitare possint. Oua edunt, & incubant, mox cum excluderint, protinus pullos educunt. propterea quod nequeunt suo uolatu ijs cibum administrare. Resouent pullos suos sub se, ipsæ ducendo more gallinarum, & coturnices & perdices. Nec eodem loco pariunt & incubant, ne quis locum percipiat, longioris temporis mora.

¹ A misprint for 'spectari.'

² *Hist. An.* Bk IX. 260, 59—61.

³ There is a reading *λείω* besides *ήλίω*.

⁴ Aristotle has *ἀκανθών τινα καὶ ὕλην*, so no doubt 'ut' is a misprint for 'et.' Gaza has 'ut.'

time draws into a cluster all the eyes upon his feathers, which he is well pleased should be admired. The same bird, having lost his tail, when the trees shed their leaves by annual change, ashamed and sorrowing seeks a hiding place, until it once more grows together with the flowers. He lives for five and twenty years, and in the third begins to shew his colours. He is reported by authorities to be an animal not only proud but also ill-disposed, just as the Goose is bashful. Peacocks have on their heads as it were a bush of hair.

OF THE PERDIX.

Πέρδιξ, perdix, in English a pertridge, in German eyn velt hön, or eyn raphön.

ARISTOTLE.

The Perdix is a bird that dusts itself, and flies not high¹; moreover it finds safety for its young not in a nest, but in thick shrubs and corn. For birds of heavy body make no nest, such as Coturnices and Perdices, and others of like sort. For to those kinds in which facility of flight is wanting, there is small advantage in a nest; but in some sunny place (for they breed nowhere else) a space is cleared and sticks and a few briars are collected there sufficient for them to avoid attacks of Accipitres and Aquilæ. They lay their eggs and sit; so soon as these are hatched, they lead their young away forthwith because they cannot supply food to them by means of flight. Both Coturnices and Perdices cherish their chicks beneath them, themselves leading them in the same way as hens lead theirs. They do not lay and incubate in the same place (year after year), lest any one should find it through the length of time for which they sit. Should

¹ See p. 35.

Cùm ad nidum quis uenando accefferit, pro-
 [p. 114] uoluit se perdix ante pedes uenantis, quasi iam
 capi possit¹, atque ita ad se capiendam hominem
 allicit, eoque dum pulli effugiant, tum ipsa
 uolat, & reuocat prolem. parit oua non pauci-
 ora quàm decem.

*Est & alia avis, quæ perdix rustica dicitur, Anglis
 rala dicta, cuius his uersibus Martialis² meminit :*

Rustica sum perdix, quid refert si sapor idem ?

Charior est perdix, si sapit illa minus.

DE PHASIANO.

*Phasianus, Anglicè a pheasant, Germanicè eyu sasant/
 oder eyu sasian.*

PLINIUS³.

Phasiani geminas aures ex pluma submit-
 tunt, subriguntq̃. quæ ueluti cornicula appa-
 rent.

ARISTOTELES⁴.

Phasianorum oua punctis distincta sunt ut
 meleagridum, puluerant ut gallinæ & perdices.
 [p. 115] Phasiani à pediculis infestantur, & nisi inter-
 dum puluerent, eisdem interimuntur.

DE PHÆNICE.

PLINIUS⁵.

Aethiopes atque Indi discolores maximè &
 inenarrabiles ferunt aues, & ante omnes nobilem
 Arabiæ phœnicem : haud scio an fabulosè, unum
 in toto orbe, nec uisum magnoperè. Aquilæ
 narratur magnitudine, auri fulgore circa colla,
 cætera purpureus, cœruleam roseis caudam pen-

¹ Aristotle has *ὡς ἐπιληπτος οὖσα*, which means 'as if disabled.'

² *Epigr.* Lib. XIII. lxxvi.

³ *Hist. Nat.* Lib. X. cap. xlviii.

⁴ *Hist. An.* Bk VI. 5, Bk IX. 260, Bk V. 140.

⁵ *Hist. Nat.* Lib. X. cap. ii.

any man when hunting come up to the nest, the *Perdix* casts herself before the hunter's feet, as if she could be caught at once, and thus attracts the man to take her, till the chicks escape, whereon she flies off and recalls her brood. She lays no fewer than ten eggs.

And there is yet another bird, which is called *Perdix rustica*: it is called "ræle" by Englishmen, and of it Martial makes mention in the following lines:—

I am a country partridge, but what matters it so that the flavour be the same? The partridge is the dearer, if it tastes less well¹.

OF THE PHASIANUS.

Phasianus, in English a phesan, in German cyn fasant or cyn fasian.

PLINY.

The *Phasiani* lower and erect two ears of feathers, which look like small horns.

ARISTOTLE.

Of *Phasiani* the eggs are marked with spots, like those of *Meleagrides*; they dust themselves, just as *Gallinæ* and *Perdices* do. *Phasiani* are a prey to lice, and if they do not sometimes dust themselves, are killed by them.

OF THE PHŒNIX.

PLINY.

The *Æthiopians* and *Indians* tell of birds of very varied colouring and indescribable, and of the *Phœnix* of Arabia, most noteworthy of all: I know not whether falsely, that there is but one in the whole world, and this not often seen. It is declared to be of the size of an *Aquila*, with golden sheen around the neck, but purple otherwise, varied with roseate feathers on

¹ The text here given is probably corrupt, which makes the rendering uncertain.

nis distinguuntibus, cristis faciem caputq; plumo apice honestante. Primus atque diligentissimus togatorum de eo prodidit Manilius, senator ille maximus, nobilis, doctore nullo, autor est, neminem extitisse, qui uiderit uescētem. [p. 116] Sacrum in Arabia foli esse, uiuere annis 660. fenescētem casia thurisq; furculis construere nidum, replere odoribus, & superemori. Ex ossibus deinde ac medullis nasci primò ceu uermiculum, inde fieri pullum.

DE PICA.

Κίττα, *pica*, *Anglicè a py, or a piot, Germanicè eyn elster/ oder eyn aßel.*

ARISTOTELES¹.

Pica uoces plurimas commutat, singulis enim ferè diebus diuersam emittit uocem. Parit oua circiter nouem numero. Nidum in arboribus facit ex pilis & lana, glandes cùm deficiunt, colligit, & in repositoio abditas, referuat.

PLINIUS².

Minor nobilitas, quia non ex longinquo [p. 117] uenit, sed expressior loquacitas certo generi picarum est, quàm psittacis est. Nec discunt tantum, sed diligunt meditatesq; intrà semet, cura atq; cogitatione, intentionem non occultant. Constat emori uictas difficultate uerbi, ac nisi subinde eadem audiant, memoria falli, quærentes mirum in modum hilarari, si interim audierint id uerbum. Nec uulgaris ijs forma, quamuis non

¹ *Hist. An.* Bk IX. 81.

² *Hist. Nat.* Lib. X. cap. xlii.

a tail of blue, tufts beautifying the face, a feathery crown the head. First of our citizens and with great care Manilius wrote of it, that noted senator, of such high birth ; of his own knowledge he asserts that nobody exists who ever saw it eat. He says that in Arabia it is considered sacred to the sun, and lives for six hundred and sixty years. When it grows old it makes itself a nest with cassia and twigs of frankincense, this nest it stores with scents and on the top it dies. Then from its bones and marrow is produced what seems a little worm, but afterwards becomes a chick.

OF THE PICA.

Κίττα, pica, in English a py or a piot, in German cyn elster or cyn atzel.

ARISTOTLE.

The Pica oftentimes changes its notes, for almost every day it utters different cries. It lays about nine eggs. It makes a nest in trees, of hair and wool, and when acorns grow scarce, it gathers them and keeps them hidden in store.

PLINY.

Less fame, because it does not come from distant lands, though more distinct loquacity characterizes a certain sort of Picæ than the Psittaci. Not only do they learn, but they delight to talk, and meditating carefully and thoughtfully within themselves hide not their earnestness. They are known to have died when overcome by difficulty in a word, and, should they not hear the same things constantly, to have failed in their memory, and while recalling them to be cheered up in wondrous wise, if meanwhile they have heard that word. Nor¹ is their beauty of an

¹ Or perhaps, 'their form is not commonplace, though not showy to the eye.'

ſpectanda, fatis illis decoris in ſpecie ſermonis humani eſt. Verùm addiſcere alias negant poſſe, quàm quæ ex genere earum ſunt, quæ glande ueſcantur, & inter eas faciliùs, quibus quini ſunt digiti in pedibus: ac ne eas quidem ipſas, niſi primis duobus uitę annis. Nuper et adhuc tamen rara ab Appennino ad urbem [p. 118] uerſus cerni cœpere picarum genera, quæ longa inſignes cauda uarię appellantur, proprium his calueſcere omnibus annis cùm feratur rapa.

Plinius duo picarum genera facere uidetur: poſterius hoc genus Plinij, picarum genus eſſe uidetur, quod paſſim in Germania & Anglia longa cauda præditum, oua & pullos gallinarum populatur. Aliud genus picę, tam longa cauda ornatum, quàm hoc eſt, non noui. noſtra quoque pica uulgaris calueſcere quotannis ſolet. Alterum autem picę genus diu ſanè dubitari quod nam eſſet, & adhuc non ſatis teneo. Cùm eſſem in Italia ad ripam Padi, ambulantiſus mihi, & itineris mei comitiſus, auis quędam picę ſimilis, lingua Britannica iaia, & Germanica mercolphus appellata, conſpiciendam ſeſe commodùm obtulit, cuius nomen Italicum quum à monacho quodam, qui tum fortè aderat, percontarer, picam granatam dici reſpondit. Qua re cùm apud Italicum etiam uulguſ non ſolùm priſtinę lingue Romanę, ſed & rerum ſcientię, non obſcura ueſtigia adhuc ſupereſſe depræhenderem, ſuborta eſt mihi hinc ſuſpicio, auem hanc è generibus picarum eſſe, & quòd [p. 119] ſcirem eandem, altera uulgari pica, multò expreſſius humanas uoces imitari, ita ſuſpicionem meam auxit, ut parùm abſit, quin credam hanc eſſe alterius generis picam, nam & glandibus ueſcitur magis omnibus alijs auibus.

Iaia An-
 glorum,
 mercolphus
 Germanorū.
 Hanc meā
 opinionē
 Perottus¹
 cōfirmat,
 quem ſi ua-
 cat lege.

¹ Probably Reader in Greek to Edward VI. (cf. *Dict. National Biogr.* XLV. p. 21).

ordinary sort, though not considerable to the eye; for them it is enough honour to have a kind of human speech. -However people deny that others are able to learn, save those belonging to the group which lives on acorns—and of these again those with the greatest ease which have five toes upon each of their feet: nor even they except during the first two years of life. Of late, however, and as yet infrequently, towards the city from the Appennines there have begun to be observed some sorts of *Picæ* which being remarkable for the length of their tails have been called “*variæ*.” They have this special mark that they grow bald in every year when rape is sown.

Of *Picæ* Pliny seems to make two kinds: this latter kind of his would seem to be that Pie which here and there in Germany and England plunders both the eggs and chicks of fowls, possessing a long tail. I do not know another kind of Pie provided with a tail so long as this. And furthermore our common Pie is wont to grow bald every year. Now what the second sort of Pie might be I doubted very long, nor have I yet grasped it sufficiently. But when I was in Italy upon the banks of the Po, and while my fellow-travellers and I were walking out, a certain bird like a Pie, in English called a Jay, in German *mercolphus*, offered itself conveniently for observation. Thereupon I asked a certain monk, who then by chance was present, its Italian name, and he replied to me that it was called the Seed Pie¹. When therefore I perceived that with the common people of Italy not only patent traces of the old Roman tongue still actually existed, but also of things scientific, a suspicion rose within me that this bird was of the group of Pies; moreover, since I knew that the same imitated human tones much more correctly than the other Pie, which is the commoner, so much was I confirmed in my suspicion that I can scarcely refuse to credit that this Pie was Pliny's second kind, particularly as it lives on acorns more than any other bird.

¹ Ghiandaja is the modern Italian name, derived from ‘*glans*’=an acorn.

DE PICO MARTIO.

a rayn
byrde.

Δρυοκολάπτης, *picus martius*, *piſo*, *iyux*, *torquella*,
turbo, *Anglicè & Germanicè a ſpecht*, *eyn ſpecht*.

ARISTOTELES¹.

Galgulum
interpretatur
Gaza.

Id eſt, uiuit
ad ligna.

[p. 120]

Alia culicibus² gaudent, nec alio magis quàm uenatu culicum uiuūt, ut *piſo* tum maior tum minor, utrumque *picum martium* uocant. Similes inter ſe ſunt, uocemq; ſimilem emittunt, ſed maiorem, quæ maior eſt. Item *κόλιος*³, cui magnitudo quanta turturi ferè eſt, color luteus, lignipeta hic admodum eſt, magnaq; ex parte more *picorum* *νέμεται ἐπὶ τῶν ξύλων*, quod eſt, ut interpretatur Gaza, ex macerie uiuit: uocem emittit magnam, incola maximè Peloponeſi hic eſt.

Obſerua ubi Ariſtoteles duo tantùm picorum genera facit, ibidem illum galgulum deſcribere, & ubi tria facit, eundem omittere.

ARISTOTELES⁴ LIB. 9. CAP. 9. DE HISTORIA
ANIMALIUM.

Alauda gallinago, & coturnix nunquam in arbore conſiſtunt, ſed humi. Contrà atq; picus martius, qui nunquam humi conſiſtere patitur. Tundit hic quercus, uermium & culicū cauſa, quo exeant: recipit enim egreſſos lingua ſua, quam maiuſculam, & latiuſculam habet. Scādit per arborem omnibus modis: nam uel reſuſpinus, more ſtellionū, ingreditur. Vngues etiam habet commodiores quàm monedula⁵, ad tuti-
[p. 121] *orem arborum reptationem, his enim adfixis aſcendit. Sunt pici Martij cognomine, tria genera: unum minus quàm merula, cui rubidæ*

¹ *Hist. An.* Bk VIII. 43—44.

² Gaza translated Ariſtotle's *σκηίψ* by *culex* (=gnat). Moſt probably it may be uſed for various ſmall winged creatures.

³ There is another reading, *κελεύς*.

⁴ *Hist. An.* Bk IX. 66—69.

⁵ *κολοιὺς* may be a miſreading here and below for *κελεύς*, but this ſeems doubtful.

OF THE PICUS MARTIUS.

Δρυοκολάπτῃς, *picus martius*, *pipo*, *ilyn*x, *torquella*, *turbo*, in English and in German a *specht*, *eyn specht*.

ARISTOTLE.

Some birds delight in grubs, and as a rule live on no other prey, as do the great and little *Pipo*, both of which people call *Picus Martius*. Resembling one another they utter like cries, although the greater has the louder cry. Again there is the *κολιός*, the size of which is, nearly as may be, that of the *Turtur*, and its colour yellowish. It pecks wood freely, and, as the *Pici* do, lives for the most part on the trunks, that is, lives on the wood¹, as *Gaza* renders it: it utters a loud cry, and is especially a resident in the Peloponnese.

Note that, when Aristotle only makes two sorts of *Pici*, in that passage he describes the *Galgulus*, when he makes three, he does not mention it.

ARISTOTLE BOOK 9, CHAP. 9, OF THE HISTORY OF
ANIMALS.

Alauda, *Gallinago*, and *Coturnix* never alight on trees, but always on the ground. It is however otherwise with *Picus Martius*, which never can endure sitting upon the ground. It hammers oaks for worms and grubs, that they may shew themselves, and when they issue forth it takes them on its tongue, which it has somewhat long and broad. It climbs about a tree in every way, for it even walks upside down, after the way of *Lizards*. It has claws better formed for creeping safely on the trees than even the *Monedula*, and climbs with them stuck in. There are three sorts of birds that have the special name of *Picus Martius*, one less than a *Merula*, which has some

¹ Cf. p. 88.

aliquid plumæ inest. Alterû maius quàm mē-
rula: tertium non multò minus quàm gallina.
Nidulatur in arboribus tum alijs cum oleis.
Pascitur formicis & coſſis. Cùm coſſos uenatur,
tam uehementer excauare, ut ſternat arbores
dicitur. Iam uerò miteſcens, quidam amygdalū,
quod rimæ inferuiſſet ligni, ut fixū conſtanter
içtu reciperet, tertio içtu pertudit, & nucleum
edit. Paucis¹ quibuſdam utrinque bini ungues
ut auiculæ, quam Iyngem uocant. Hæc paulò
maior quàm fringilla eſt, corpore uario. Habet
ſibi propriam digitorum, quā modò dixi, diſpo-
[p. 122] ſitionem, & linguam ſerpentibus ſimilem: quippe
quam in longitudinem, menſura quatuor digi-
torum porrigat, rurfumq; contrahat intra ro-
ſtrum, collum etiam porrigit in auerſum reliquo
quieſcente corpore, modo ſerpentium, unde tor-
quilla uulgò appellata eſt: quanquam turbo ab
antiquis. Ungues ei grandes, & ſimiles, ut
monedulis exeunt, uoce autem ſtridet.

Picus mi-
nimus.
Medius.
Maximus.

*Primum pici genus Angli ſpechtam, & uuod-
ſpechtam, Germani elſterſpechtam nominant. Secundum
genus Angli huholam, hoc eſt, foraminum dolatorem,
Germani grunſprechtā nuncupant. Tertiū genus Angliā
nō nouit, Germani aut craspechtam .i. cornicinum
picum appellāt, quòd cornicem plumarū colorc & magni-
tudine etiam penè æquet.*

*Plinius præter hæc tria Ariſtotelis genera, quartum
pici genus facere uidetur, nam lib. 10. cap. 33 ſcribit.
picum aliquem ſuſpendere nidum in ſurculo primis
in ramis cyathi modo, ut nulla quadrupes poſſit ac-
cedere.*

*Præter uireonem ſolum, in Europa nullam aliā
[p. 123] auem ita nidulari noui. Quare nullam aliam, quàm
hanc, quæ pici quartum genus eſſe poſſit, inuenio.*

¹ *Hist. An.* Bk II. 46—47.

feathers red. The second bigger than a *Merula*, the third not much less than a barn-door hen. It breeds in various trees and olives in particular. It feeds on ants and grubs, and when hunting for grubs is said to excavate so vigorously as to fell trees. Indeed one that was tamed broke at the third attempt an almond which it had inserted in a chink of the wood, that being fixed it might more surely receive the stroke, and ate the kernel out. In some few birds there are two claws in front and two behind, as in the little bird which men call *Iynx*. This kind is not much larger than a *Fringilla*, and has the body mottled. It has moreover the peculiar arrangement of the toes, of which I have just spoken, and a tongue like that of serpents, for it shoots it out up to a distance of four fingers' length, and draws it back again within the beak; it twists its neck moreover backwards, with its body still, just as the serpents do, whence it is commonly called *Torquilla*, although it is the *Turbo* of old writers. It has claws of great size, which are like those that grow on the *Monedula*, it has a strident cry.

Of *Picus* the first kind the English call the *Specht* and *Wodspecht*, which the Germans name the *elsterspecht*. The second kind Englishmen term *Hewhole*, that is, hewer of holes, the Germans *grunspecht*. The third kind England knows not, but in Germany they call it *craspecht* or the *Crow-Picus*, for it is very nearly like a *Crow* in colour of the plumage and also in size. Besides these three sorts of *Aristotle* *Pliny* seems to make a fourth, for in Book 10 and chapter 33 he tells us that a certain *Picus* hangs its nest, in fashion like a cup, upon a twig among the outer branches of a tree, so that no quadruped is able to come nigh. Except the *Vireo* alone, I know no other bird in Europe which places its nest in such a way. Wherefore I find no other than the above which the fourth kind of *Picus* possibly can be.

DE PSITACO.

Psitacus, Anglicè *a popiniay*, Germanicè *eyn papegay*.

ARISTOTELES¹.

Nam & Indica auis, cui nomen psitacæ, quam loqui aiunt², talis est, & loquacior³ quum biberit uinum, redditur.

PLINIUS⁴.

Super omnia humanas uoces reddunt psitaci, & quidem fermocinantes. India auem hanc mittit. Psitacen uocant, uiridè toto corpore, torque tantū miniato in ceruice distinctā. Imperatores salutat, & quæ accipit uerba, pronunciat: in uino ꝑcipuè lasciua. Capiti eius duritia eadem quæ rostro. Hæc cū loqui discit, ferro uerberatur radio: nō sentit aliter ictus. [p. 124] Cū deuolat, rostro se excipit, illi innititur, leuioremq; se ita pedum infirmitati facit.

DE PLATEA.

Πέλεκας, *platea*, *platelea*, *pelcanus*, Anglicè *a shouclard*, Germa. *eyn lesler*/ od' *eyn löffel gāß*.

ARISTOTELES⁵.

Platea fluuiatilis, conchas maiusculas, leuesq; deuorat, quas ubi sua ingluuie coxerit, euomit, ut hiantibus testis exuens, legat atque edat.

PLINIUS⁶.

Platea nominatur aduolans ad eas, quæ se in mari mergunt, & capita illarum morfu corri-

¹ *Hist. An.* Bk VIII. 85.

² τὸ λεγόμενον ἀνθρωπόγλωττον.

³ ἀκολαστότερον = reckless.

⁴ *Hist. Nat.* Lib. X. cap. xli.

⁵ *Hist. An.* Bk IX. 71.

⁶ *Hist. Nat.* Lib. X. cap. xl.

OF THE PSITACUS.

Psitacus, in English a popinjay, in German eyn papegay.

ARISTOTLE.

An Indian bird indeed, the Psitace by name, which people say can speak, is such a one as this, and is reported as more talkative after it has drunk wine.

PLINY.

Beyond all Psitaci repeat men's words, and even talk connectedly. India sends this bird, which they call Psitace, with the whole body green marked only by a scarlet ring upon the nape. It will pronounce "Hail Emperor," and any words it hears; it is especially sportive after wine. The hardness of the head is the same as of the beak. And when the bird is being taught to speak, it is beaten with an iron rod, else it feels not the strokes. When it flies down it receives its weight upon its beak, and supports itself thereon; and thus lightens itself to remedy the weakness of its feet.

OF THE PLATEA.

Πελεκάν, platea, platelea, pelecamus, in English, a shovellard, in German eyn lefler or eyn löffel ganss.

ARISTOTLE.

The Platea, a river bird, devours biggish shellfish, if they be but smooth, and, after it has seethed them in its crop, it casts them up again, that stripping them off from their gaping shells, it so may pick and eat them.

PLINY.

The Platea, as it is called, flies at those birds which dive below the sea, and seizes their heads with a bite

piens, donec capturam extorqueat. Eadem cum se deuoratis impleuit conchis, calore uentris coctas, euomit, atque ita esculenta legit, testas excernens.

[p. 125]

HIERONYMUS.

Pelicanus cum suos à serpente filios occisos inueniunt, lugent, sed & sua latera percutiunt, & sanguine excusso, corpora mortuorum sic reuiuiscunt.

Couradus Gestnerus, cum Tiguri agerem, homo ut doctissimus, ita candidissimus, huius mihi auis cognitionem (ut fatear, per quem profeci) primus omnium communicauit, & ideo Germanis lesterā uocari docuit, quod rostrū cochleari simile haberet.

DE PORPHYRIONE EX PLINIO¹.

Porphyrio solus morfu bibit: idem ex proprio genere omnem cibū aqua subinde tingens, deinde pede ad rostrum ueluti manu adferēs, laudatissimi in comagene². Rostra ijs & prælonga crura rubent.

DE REGVLO.

Τροχίλος, πρέσβυς, βασιλεύς, trochilus, senator, regulus, Anglicè a uuren, Germanicè ein funningſgen/ oder ein zaunſinningſ.

[p. 126]

ARISTOTELES³.

Trochilus & fruteta incolit, & foramina, capi difficulter potest, fugax atque infirmis moribus est, sed uictus probitate, & ingenij solertia præditus. uocatur idem senator & rex, quam ob rem aquilam cum eo pugnare referunt.

¹ *Hist. Nat. Lib. x. cap. xlvi.*

² It almost seems as if 'Comagene' should be 'Commageno,' in which case we might translate 'They are highly prized for ointment.'

³ *Hist. An. Bk ix. 75.*

until it wrenches their prey from them. So too when it has filled itself with shell-fish that it has devoured, it casts them up, seethed by its belly's heat, and so picks out the eatable parts, sifting off the shells.

HIERONYMUS.

Pelecani, when they find their young killed by a serpent, mourn, and beat themselves upon their sides, and with the blood discharged, they thus bring back to life the bodies of the dead.

Conrad Gesner, a man most learned as he also was most truthful, first imparted to me while I was at Zurich knowledge of this bird (that I may own from whom I profited), and taught me that it was called lefler by Germans because it has a spoon-shaped beak.

OF THE PORPHYRIO FROM PLINY.

The Porphyrion alone drinks with a bite, it also is peculiar in dipping all its food from time to time in water, and then bearing it to its beak with its foot, as with a hand. The best are found in Comagene. Their beaks and very long legs are red.

OF THE REGULUS.

Τροχιλος, πρέσβυς, βασιλεύς, trochilus, senator, regulus, in English a wren, in German cyn kuningsgen or cyn zaunkünigk.

ARISTOTLE.

The Trochilus inhabits shrubberies and holes, and cannot easily be caught. Now it is shy and of a feeble habit, but endowed with great ability of getting food and knowledge of its craft. The same is called both senator and king, on which account the Aquila, they say, fights with it.

Trochilus, est avium omnium minima, cauda longa & semper erecta, rostro longiusculo, sed tenuissimo, colore ferè fuluo. nidum facit foris ex musco, intus ex plumis aut lana, aut floccis, sed plurimū ex plumis. Ovi erecti & in altero suo fine cōsistentis, formā nidus habet, in medio ucluti latere hostiolum est, per quod ingreditur & egreditur. In posticis ædibus & stabilis stramine tectis, interdū nidū construit, sed sæpius in sylvis. anis est etiam soliuaga, & gregatim nunquam uolat, imò quoties alium sui generis offendit, mox illi bellum indicit, & conflictatur. Quare anes illæ, quæ in Bauaria pennis auricoloribus, quas in capitibus ceu coronas aureas ferunt, in sylvis æstate degentes, & gregatim ad urbes hyeme aduolantes, reguli non sunt sed tyranni Aristotelis, ut postea docebo.

[p. 127]

DE RVBECVLA ET
ruticilla.

- Rubecula Ερίθακος, ἢ ἐρίθῆα, rubecula, Anglicè a robin red-breſte, Germanicè *eyn rōtbruſt/ oder eyn rōtfeldchen.*
- Ruticilla. Φοινικουρὸς, & ut alter textus habet, φοινικούργος, Plinio *phœnicurus*, Gazæ ruticilla, Anglicè a rede tale, Germanicè *eyn rōt ſterg.*

ARISTOTELES¹.

Rubecula & ruticilla uermibus aluntur. Rubeculæ, & quæ ruticillæ appellantur, inuicem tranſeunt, eſt'que rubecula hyberni temporis, ruticilla æſtiui, nec alio ferè inter ſe differunt, niſi pectoris colore & caudæ².

¹ *Hist. An.* Bk IX. 256.

² The three words 'pectoris & caudæ' are not in Aristotle.

The Trochilus is smallest of all birds, with a long tail¹ always cocked up, and a bill somewhat long but very slender, it is nearly reddish-brown in colour. The nest it makes is outwardly of moss and inwardly of feathers, wool, or down, but mainly of feathers. The nest has the form of an upright egg standing on one of its ends, while in the middle of one side there is a little postern as it were, by which the bird goes in and out. It sometimes builds its nest at the back of a house or in sheds thatched with straw, but usually in woods. It also is a bird that roves alone, and never flies in flocks; nay more, so often as it meets another of its kind it forthwith declares war, and fights. Wherefore the birds with plumes of gold-colour that they wear on their heads like golden crowns, which pass the summer in Bavaria in woods, and in the winter flock to towns, are not the Reguli of Aristotle, as I presently shall prove, but the Tyranni.

OF THE RUBECULA AND THE RUTICILLA.

Ἐπίθακος, ἢ ἐπιθέα, rubecula, in English a robin redbreast, in German eyn rötbrust or eyn rötkelchen.

Φοινίκουρος, and, as another text has it, φοινικουργος, in Pliny phœnicurus, the ruticilla of Gaza, in English a rede-tale, in German eyn rötstertz.

ARISTOTLE.

The Rubecula and the Ruticilla feed on worms. Rubeculæ and Ruticillæ, as the birds are called, change into one another, and what in winter is the Rubecula in summer is the Ruticilla², while they hardly differ from each other save in colour of the breast and tail.

¹ Turner evidently means the Wren (*Troglodytes parvulus*), but with this the 'long tail' does not agree; perhaps there is a misprint.

² As Sundevall remarks, Aristotle probably only meant that the Redstart was called φοινίκουρος in summer and ἐπίθακος in winter. Sundevall ascribes the misinterpretation to Gaza, whose work Turner admittedly used. This is the more likely as the section of Aristotle quoted concerns birds which change their plumage and note at different seasons.

Omnia, quæ hîc Aristoteles de duabus auibus istis conscripsit, Plinius ex ipso in opus suum transcripsit. Sed uterque hac in re, aucupum relatibus magis quàm sua experientia nixus, à ueritatis tramite longissimè aberrauit, nam utraque auis simul conspicitur, & rubeculæ domitæ, & in caueis alitæ, eandem perpetuò formam retinent. quin & eodem tempore nidulantes,

Rubeculæ
nidulatio

[p. 128] *sed modis longè diuersis sæpissimè in Anglia uidi. Rubecula, quæ non secus æstate quàm hyeme rubrû habet pectus, quàm possit longissimè ab oppidis et urbibus in densissimis uepretis, & fruticetis ad hunc modum nidulatur. Vbi multa querna reperit folia, aut quernis similia, ad radices ueprum, aut densiorum fruticum, inter ipsa folia nidum construit: & iam constructum, opere ueluti topiario folijs contegit. Nec ad nidum ubiq; patet aditus, sed una tantum uia ad nidum itur. ea quoque parte, qua nidum ingreditur, longum struit ex folijs ante hostium nidi uestibulum, cuius extremam partem pastum exiēs, folijs claudit. Hæc, quæ nunc scribo, admodum puer obseruaui, non tamen inficias inuerim, quin aliter nidulari possit. Si qui alium nidulandi modum obseruauerint, edant, & huiusmodi rerum studiosis, & mihi cum primis nō parum gratificabuntur. Ego, quod uidi, alijs candidè sum impertitus.*

Phœnicuri
nidulatio.

Phœnicurus, quem rubicillam¹ uocat, in excauatis arboribus & (quod sæpè expertus sum) in rimis & fissuris murorum, posticarum ædiū, in medijs urbibus, sed ubi hominum minor frequentia concursat, nidulatur.

Rubeculæ
æstate can-

[p. 129] *semper motitat uterque. Phœnicura fœmina, & proles adeo rubeculæ pullis similes sunt, ut uix ab oculatissimo tantè nunquam audiui.*

Phœnicurus mas nigro est capite, & cauda rubra, cætera fœminæ, nisi quod subinde cantillat, similis. Caudam

discerni possint. Verùm motu caudæ dignoscuntur. Rubeculæ licet caudam moueant, postquam tamen submiserint, statim erigunt, nec tremit bis aut ter more ruticillarū. Ruticillæ enim simul atque caudam mouere ceperint, non cessant donec ter aut quater simul

¹ No doubt a misprint for Ruticilla, cf. pp. 154, 160.

All that Aristotle here has written of these two birds Pliny has copied from him into his own work. But in this matter each of them, relying on the tales of fowlers more than on his own experience, has wandered greatly from the path of truth. For both the birds are seen at the same time; moreover tame *Rubeculæ*, when fed in cages, constantly retain the same appearance. Moreover I have very often seen the birds in England nesting at the same time, though in very different ways. The *Rubecula*, which has a ruddy breast no less in summer than in winter, nests as far as possible from towns and cities in the thickest briers and shrubs after this fashion. Where it finds oak leaves in plenty, or leaves like the oak, it builds its nest among the leaves themselves close to the roots of briers or the thicker shrubs: and when completed covers it with leaves as if with topiary work. Nor does access lie open to the nest on every side, but by one way alone is entrance gained. And at that place where it enters the nest the bird builds a long porch of leaves before the doorway and, on going forth to feed, closes the end with leaves. But, what I now describe, I first observed when quite a boy, nevertheless I am not going to deny that it may build otherwise. If any have observed another way of nesting, let them tell it, and they certainly will not a little gratify the students of such things, myself among the first. I have imparted truthfully to others what I saw.

The *Phœnicurus* which he [Gaza] calls the *Rubicilla* nests in hollow trees and (as I often have had experience) in chinks and cracks of walls and outhouses in the midst of our towns, though where the throng of men is not so great. The male has a black head, a red tail, but otherwise is like the female, save that he repeatedly utters a little song. Either sex flirts the tail continually. The female *Phœnicurus* and its brood are so much like young of *Rubecula* that they can scarcely be distinguished by the sharpest eye. But by the motion of the tail they may be recognised. For the *Rubeculæ*, although they move the tail, yet, after they have lowered it, at once raise it again, nor does it quiver twice or thrice as does that of the *Ruticilla*. For no sooner have the *Ruticillæ* once begun to move the tail than they go on till they have lightly moved it three or four times altogether

leuiter mouerint, ut alas, iuniores auiculæ cibum à matribus efflagitantes, motitant. Rubeculæ in æstate, ubi in syluis satis superq; alimenti suppetit, nec ullo infestantur frigore, (quæ res cogit illas in hyeme ad urbes, oppida & pagos confugere) cum prole ad desertissima quæq; loca secedunt. Quare, minùs mirandum est, rubeculas in æstate non passim occurrere. Ruticillas quid miri est in hyeme nō esse obuias, quum per totam hyemem delitefcant? Adhæc cūm rubeculæ pulli, in fine autumnii perfectam ferè in pectoribus rubedinem naçti, ad pagos & oppida propiùs accedunt, ruticillæ, quæ antea per totam æstatem cernebantur, disparent, nec amplius in proximum usque uer cernuntur. Quæ quum ita se habeant, quid Aristoteli aut illi hoc referentibus erroris ansam præbuerit, facilè quiuis potest colligere.

DE RVBETRA.

[p. 130] Βάτις, Latinè rubetra dicta, ab Aristotele inter auiculas uermiuoras numeratur. Porro, quænam auis sit, prorsus diuinare non possum. Gybertus Longolius linariam, siue miliariam esse rubetrā putabat, quod rubis crebrò insideat. Sed quum Anglorum buntinga in rubis tam frequēs sit, quid uetat quò minùs & ipsa quoq; batis dici possit. Nihil igitur certi habemus, quod nomen Britannicum aut Germanicum sit huic aui imponendum.

Sed quum auium suprà commemoratarum altera seminibus herbariū uescatur, & altera hordeo & tritico, & batis Arist. uermiuora sit, delegēda est auicula quæpiam, quæ solis uermibus pascitur, qualis est auicula Anglis stonchattera, aut mortettera dicta, & Germanorum klein brachuogelchen. Hæc si batis non sit, mihi prorsus ignota est. Porro illa, quam Angli linotam, & Germani flasfincam uocant, ueteribus, si Ruellio, credimus, erit miliaria.

just as young small birds flutter their wings soliciting meat from their mothers. In summer, when there is enough and more of food found in the woods, and they are not troubled by any cold (a thing which forces them in winter to resort to cities, towns, and villages), *Rubeculæ* retire to the most solitary places with their young. And so it is no marvel that *Rubeculæ* do not occur in summer everywhere. And what wonder is it that *Ruticillæ* are not met with in winter, since throughout the whole of winter they are hidden? And further, when the young *Rubeculæ*, having almost assumed the full red on their breasts at the end of autumn, come nearer to towns and villages, the *Ruticillæ*, which were hitherto seen during the whole summer, disappear and then are no more noticed till the following spring. Wherefore, things being thus, anyone may easily perceive what gave a handle to Aristotle or to those who reported this error to him.

OF THE RUBETRA.

Βατίς, in Latin called *Rubetra*, is by Aristotle classed among the little birds that feed on worms. Beyond this I cannot guess at all what sort the bird may be. Yet Gybertus Longolius¹ believed that the *Rubetra* was the *Linaria* or the *Miliaria*, because it often perched on brambles. But since the Bunting of the English sits so commonly on brambles, what forbids that bird from also being called the *Batis*? On this account we have no certainty as to what name, British or German, should be given to this bird.

But inasmuch as of the birds mentioned above the one eats seeds of grasses, and the other wheat and barley, and as Aristotle's *Batis* lives on worms, some small bird must be chosen which eats worms and nothing else. Now such a little bird is that called by the English *Stonchatter* or *Mortetter* and the *klein brachvogelchen* of the Germans. If this be not the *Batis*, it is quite unknown to me. Besides that which the English call the *Linot* and the Germans the *flasfinc* must be the *Miliaria* of older works, if we believe Ruellius².

¹ For this author see Introduction.

² Ruellius wrote *De natura stirpium libri tres* (1536) and edited one or more medical or other works.

DE RVBICILLA.

Πυρρούλας, *rubicilla*, Angli. a *bulfinche*, Germa. *eyn blödtfinck*.

Rubicillam Aristoteles inter eas aues connumerat, quæ uermibus uescuntur: sed pluribus uerbis cam non describit. Ego nominis etymologiam secutus, rubicillam [p. 131] *Anglorum bulfincam, & Germanorum blouduincam esse conijcio. Nam omnium, quas unquam uidi auium mas in hoc genere, pectore est longè rubidissimo: fœmina uerò pectore toto est cinereo, cætera mari similis. Sed ut facilius omnes intelligant, de qua aue scribam, magnitudine passeris est, rostro breuissimo, latissimo, et nigerrimo, lingua latiore multò quàm pro corporis magnitudine. Pars ea linguæ, quæ cibi sapores dijudicans, oris cælum tangit, carnea & nuda est, reliquæ partes cornea pellicula obducuntur. Supremam auis partem plumæ cyanæ contegunt. cauda nigra est, & capite etiam nigro, uescitur libentissimè primis illis gemmis ex arboribus ante folia & flores crumpentibus, & semine canabino. auis est imprimis docilis, & fistulam uoce sua proximè imitatur. nidulatur in sepibus, & oua quatuor excludit, ut plurima quinq;. eundem colorem per totum annum seruat, nec locum mutat. Quæ quum ita se habeant, non potest hæc atricapilla esse, ut quidam uolunt, utcumq; extremo linguæ acumine carere uideatur.*

DE SALO, QVI GRÆCE

ἄγριθος dicitur.

ARISTOTELES¹.

[p. 132] *Salus uitæ commoditate, & partus numero commendatur, sed alterius pedis clauditate cedit. Sali & flori sanguinè misceri negant, tâ ingens inter salum & florū feruet odiū. Salo etiam preliū cū asino est, propterea quòd asinus spi-*

¹ *Hist. An.* Bk IX. 89, 22, 14.

OF THE RUBICILLA.

Πυρρόυλας¹, rubicilla, in English a bulfinche, in German cyn blödtfinck.

Aristotle counts the Rubicilla among those birds which feed on worms: but he does not describe it in more words. I, following the derivation of the name, conjecture that it is the Bulfinch of the English and the bloudvinc of the Germans. For of all the birds I ever saw the male of this kind has by far the reddest breast: the female however has the breast wholly grey, though otherwise like the male. But, that all may understand more easily about which bird I write, it is the size of a Sparrow, with the beak particularly short and broad and black, the tongue much broader than is in proportion to its body. That part of the tongue which discriminates the flavour of the food and meets the palate of the mouth is flesh-coloured and naked, while the other parts are covered with a horny pellicle. Bluish grey feathers clothe the upper parts. The tail is black and the head also black. It feeds most greedily on those earliest buds, which burst out on the trees before the leaves and flowers, as well as hemp-seed. It is the readiest bird to learn, and imitates a pipe very closely with its voice. It nests in hedges where it lays four eggs or five at most. It keeps the same colour throughout the year, and does not change its home. And since these things are so, it cannot be the *Atricapilla*, as some will have it, though it may appear to lack the point at the tip of the tongue.

OF THE SALUS, WHICH IN GREEK IS CALLED αἶγιθος.

ARISTOTLE.

The Salus is well thought of for its skill in gaining a livelihood, and for the number of its young, although it suffers from lameness in one of its feet. And men deny that the blood of the Salus and the *Florus* ever mixes, for so great an enmity rages between the birds. There is war also between the Salus and the Ass,

¹ Sundevall thinks that the Πυρρόυλας is the Redbreast, but the description does not agree with his idea. Another reading is πυρρουράς.

netis sua ulcera scabendi causa atterat: tum igitur ob eam rem, tū etiam quòd si uocem rudentis audierit¹, oua abigat per abortum, pulli etiam metu in terram labantur. Itaque ob eam iniuriam aduolās, ulcera eius rostro excauat.

PLINIUS².

Aegithus auis minima cum asino pugnat, spinetis enim se scabendi causa atterens, nidos eius dissipat, quod adeò pauet, ut uoce audita omnino rudentis oua eijciat, & pulli ipsi metu [p. 133] cadant: igitur aduolans ulcera eius rostro excauat.

DE SITTA.

Sitta, Anglicè a nut iobber, Germanicè eyu nushäffer oder eyu meyspecht.

ARISTOTELES³.

Sunt & ei, quæ fitta dicitur, mores pugnaces, fed animus hilaris, cōcinnus, compos uitæ facilioris. Rē maleficam illi tribuunt, quia rerum callat cognitione, prolem numerosam faciliemq; progignit, cariosa ligna contundens, ex coffis, quos inde eruit, uiçtitat. Sitta⁴ aquilæ oua frangit, aquila tum ob eam rem, quum etiam quòd carniuora est, aduersatur.

Auicula, quam Angli nucipetam uocant, & Germani meyspechtum, paro maximo paulò maior est, pennis cyaneis, rostro longiusculo, & per arbores eodem modo, quo picus ascendit, & easdem uiçtus gratia contundit: nuces rostro etiam perforat, & nucleos commedit. nidu- [p. 134] latur in cauis arboribus more pici, uoce ualde acuta & sonora est.

¹ Aristotle has 'κᾶν ὀγκήσῃται, ἐκβάλλει τὰ ὠὰ καὶ τοὺς νεοττοὺς,' as if the bray of the ass shook the eggs and young out of the nest. Gaza seems to have had the reading ἐκτίκτουσι for ἐκπίπτουσι

² *Hist. Nat.* Lib. x. cap. lxxiv.

³ *Hist. An.* Bk IX. 91, very freely rendered.

⁴ *Hist. An.* Bk IX. 17

because the Ass is wont to rub its sores against the thorn-bushes to scratch them, therefore for this cause, and also because the bird has heard the brayer's voice it prematurely drops its eggs, while even nestlings fall down to the ground with fear. So for that injury (the bird) attacking it scoops out its sores.

PLINY.

The *Ægithus*, a very little bird, wages war with the Ass, because it, rubbing against thorn-bushes to scratch itself, destroys the nest, and this the bird dreads so much that, if it merely hears the brayer's voice, it drops its eggs, and the young also fall to the ground with fear. Accordingly attacking it the bird scoops out its sores.

OF THE SITTA.

Sitta, in English a nut jobber, in German *eyn nushäkker* or *eyn meyspecht*.

ARISTOTLE.

That bird which is called Sitta has pugnacious habits but a cheerful disposition ; it is elegant and well adapted to get food with ease. Yet men attribute witchcraft to it, since it is cunning in knowledge of affairs ; it produces numerous young with ease ; hammering on rotten trees, it lives upon the grubs which thence it digs. The Sitta breaks the eggs of the *Aquila*, on which account, and also since it is carnivorous, the *Aquila* is its enemy.

The small bird which the English call the Nut-seeker and Germans the *meyspecht* is somewhat bigger than the biggest *Parus*, with blue plumage and a longish beak. It climbs trees in the same way as the *Picus*, and hammers the same for food ; it also bores nuts with its beak, and eats the kernels. It nests in hollow trees, as does the *Picus*, while its note is very sharp and loud.

DE STRVTHIONE.

Στρουθός, λιβυκός, *struthio*, aut *struthiocamelus*,
Anglicè an oïstris, *Germanicè* cyn ſtrauß.

ARISTOTELES¹.

Struthio, etiam libicus, eodem modo partim auem, partim quadrupedem refert, quippe qui non ut quadrupes pennas habeat, ut non auis sublimis non uolet, nec pennas ad uolandum commodas gerit, sed pilis similes. Itē quasi quadrupes sit, pilos habet palbebrę superioris, & gibber² capite, parte colli superiore est. Itaq; cilia habet pilosiora, sed quasi auis sit, infrā pennis integitur. Bipes etiam tanquam auis, bifulcus tanquam quadrupes est. Nō enim digitos habet, sed ungulam bipartitam. quarum rerum [p. 135] causa est, quod magnitudine non auis, sed quadrupes est. Magnitudinem enim auiū minimam esse propè dixerim, necesse est. Corpus enim molem sublimem mouere, nequaquam facilē est.

DE STVRNO.

Ψάρος, *sturnus*, *Anglicè a sterlyng*, *Germanicè* cyn ſtär/ oder cyn ſtör.

ARISTOTELES³.

Sturnus niger est, albis maculis distinctus, magnitudine merulæ.

PLINIUS⁴.

Sturnorum generi proprium, cateruatim uolare, & quodam pilæ orbe circumagi, omnibus in medium agmen tendentibus.

¹ *De partibus Animalium*, IV. 14.

² No doubt a misprint for 'glaber.'

³ *Hist. An.* Bk IX. 102, freely rendered.

⁴ *Hist. Nat.* Lib. X, cap. xxiv.

OF THE STRUTHIO.

Στρουθός, λιβυκός, struthio or struthiocamelus, in English an oistris, in German eyn strauss.

ARISTOTLE.

The Struthio, or Libyan bird, in like manner partly recalls a bird, partly a quadruped, seeing that it, unlike a quadruped, has wings, and yet, unlike a bird, it does not fly aloft, nor has it feathers fit for flight, since they resemble hairs. Likewise as if it were a quadruped, it has hairs on the upper eyelid, while the head and upper portion of the neck are bare. So also it has somewhat hairy eyelashes, yet it is covered with feathers beneath, as if it were a bird. Moreover it is biped like a bird, but yet it is cloven-footed like a quadruped. That is, it has not toes but a divided hoof. The cause of these things is that in its size it is not a bird but a quadruped. For I would almost say that a bird's size must be extremely small, for it is by no means easy to move aloft a body when the mass is vast.

OF THE STURNUS.

Ψάρος, sturnus, in English a sterlyng, in German eyn stār or eyn stör.

ARISTOTLE.

The Sturnus is black, varied with white spots, and of the bigness of a Merula.

PLINY.

It is peculiar to Starlings in their kind to fly in crowds, and wheel about as it were in a ball, all tending to the middle of the band.

DE TINVNCVLO.

Κέγχρις, *tinunculus*, Anglicè a *kistrel*, or a *kastrel*, or a *steingall*.

ARISTOTELES¹.

[p. 136] Omnes, quibus ungues adunci, parcius generant, excepto tinunculo, qui plurima in adunco genere parit. Iam enim quatuor eius reperti sunt pulli, sed plures etiam procreari posse, apertum est. Tinunculo² uentriculus ingluuiei similis est, & solus in adunco genere bibit. Rubra sunt eius oua modo minij.

Tinunculus colore multò magis est fuluo quàm reliqui accipitres, & corpore paruo. Auiculas insequitur, & ut quidam mihi retulére, papiliones interdum. In cauis nidulatur arboribus, & in templorum muris, & æditioribus turribus, ut apud Germanos Argentorati & Coloniae, & apud Anglos Morpeti obseruauit. pullos etiam diu uolantes tantisper alit, dùm ipsi ex proprio uenatu uiuere possunt.

DE TETRAONE.

Τέτριξ, όupaξ, *tetrao*, Anglicè a *bustard*, or a *bistard*, Germanicè *eyn trāp*/ oder *eyn trap gānß*.

ARISTOTELES³.

[p. 137] Tetrix, quam Athenienses uragem uocant, nec terræ, nec arbori suum nidum committit, sed frutici⁴.

PLINIUS⁵.

Decet tetraones suus nitor, absolutaq; nigritia, in supercilijs cocci rubor. Alterum eorum genus uulturum magnitudinē excedit, quorum

¹ *Hist. An.* Bk VI. 2.

² *Hist. An.* Bk II. 88, Bk VIII. 50, Bk VI. 6.

³ *Hist. An.* Bk. VI. 4.

⁴ Aristotle has χαμαιζήλους φυτοίς.

⁵ *Hist. Nat.* Lib. X. cap. xxii.

OF THE TINNUNCULUS.

Κεγχρίς, tinnunculus, in English a kistrel or a kastrel, or a steingall.

ARISTOTLE.

All birds with crooked claws¹ breed somewhat sparingly, save the Tinnunculus, and it of all the crooked-claw kind lays the most eggs. For of this bird four young have been already found, while it is evident that more might be produced. The stomach in Tinnunculus is not unlike a crop, whereas it is the only one of all the crooked-claw kind that drinks. Its eggs are red—like scarlet.

The Tinnunculus is of a much more fulvous colour than are other Hawks, and small in body. It chases little birds, and, as some men have told me, butterflies at times. It nests in hollow trees, church walls, and lofty towers, as I have seen in Germany at Strassburg and at Cullen, and in England at Morpeth. It also feeds its young long after they can fly until such time as they can live apart by hunting for themselves.

OF THE TETRAO.

Τέτριξ, οὐραξ, tetrao, in English a bustard or a bistard, in German eyn tröp or eyn trap gänss.

ARISTOTLE.

The Tetrix, which Athenians call the Urax, trusts its nest not to the ground, nor to a tree, but to low-growing plants.

PLINY.

Their glossy plumage well becomes the Tetraones, as does furthermore their perfect blackness and the scarlet redness of their eyebrows. But one kind exceeds in size the vultures and recalls their colour-

¹ For Aristotle's groups of Birds see p. 35.

et colorem reddit. Nec ulla auis, excepto struthiocamelo, maius corpore implēs pondus, in tantum aucta, ut in terra quoq; præhendatur. gignunt eos alpes, & septentrionalis regio. In uiuarijs saporem perdunt. Moriuntur contumacia spiritu reuocato. Proximæ eis sunt, quas Hispani aues tardas appellāt, Græcia otidas dānatas cibis. Emissa enim ossibus medulla, odoris tedium extemplo sequitur.

[p. 138]

DE TYRANNO.

Tyrannus, Anglicè a nyn murder, Germanicè eyu neun mürder/ oder eyu gold hendlin.

ARISTOTELES¹.

Vescitur & uermibus tyrannus, cui corpus non multò amplius quàm locustæ, crista rutila ex pluma elatiuscula, & cætera elegans, cantuq; suauis hæc auicula est.

Quaquam Aristoteles unum tantum tyranni genus faciat, Colonienfes tamen aucupes tria genera esse contendunt. Primum uocant die grosse neun murder, quod Angli etiam schricum nominant: & ego Aristoteles mollicipitem esse conijcio: sturnum magnitudine æquat, color eius à cyanco ad cinerum uergit. Secundum genus eiusdem est coloris, cuius & superius, sed passerem magnitudine non excedit. Hoc genus etiam in aues sæuit. Tertium genus, quod Aristotelis tyrannus est, auicula est regulo paulò maior, crista rutila redimita, & cæteris generibus (si aucupibus credere phas² sit) cede³ & corporis effigie non dissimilis. Secundum & tertium tyranni genus apud Anglos hætenus nunquam uidere
 [p. 139] *contigit, & primum genus licet in Anglia sit, paucissimis tamen notum est: sunt tamen, qui norunt, & schricum uocant.*

¹ *Hist. An.* Bk VIII. 41.

² Of course a misprint for 'fas.'

³ No doubt this should be 'sede.'

ing. There is no other bird, except the *Struthio-camelus*, which attains so great a weight of body, growing to such a size that it may even be caught upon the ground. The Alps produce them, as do northern lands. In mews they lose their flavour. They die of stubbornness by holding back their breath. Very near them are those which Spaniards call "*Aves tardæ*" and Greece "*Otides*"; they are condemned as food. For when the marrow issues from the bones, disgust at the smell follows there and then¹.

OF THE TYRANNUS.

Tyrannus, in English a nyn murder, in German eyn neun mürder or eyn gold hendlin.

ARISTOTLE.

The Tyrannus also lives on worms². Its body is but little larger than a locust's is. This little bird moreover has a somewhat upright crest of reddish feathers, and is otherwise pretty; its song is sweet.

Though Aristotle makes but one kind of Tyrannus, yet the bird-catchers of Cullen state that there are three. The first they call die grosse neun mürder, which the English name the Schric for their part, this I take to be the *Molliceps* of Aristotle. In size it equals the *Sturnus*, while its colour verges from blue to grey. The second kind is of the same colour as the foregoing, but in size does not exceed a Sparrow. Furthermore this kind is cruel towards other birds. The third kind, which is Aristotle's Tyrannus, is a small bird which is little bigger than the *Regulus*, adorned with a red crest and not unlike the other kinds (if it be right to trust the fowlers) in its haunts and form of body. It has not ever happened to me hitherto in England to observe the second or the third kind of Tyrannus and, although the first exists in England, it is known to very few. Yet there are some who know it and who call it Schric.

¹ See p. 106.

² See p. 35.

DE TRYNGA.

Τρύγγας, trynga, Anglicè a uwater hen, or a mot hen, Germanicè eyn wasser hen.

ARISTOTELES¹.

Lacus & fluuios petunt iunco, cinclus, & trynga², quæ inter hæc minora, maiuscula est, turdo enim æquiparatur: omnibus his cauda motitat.

Iam, ut sciatis, quam auem tryngam esse putem, auis tota pulla est, excepta ea caudæ parte, quæ podicem tegit, ea enim candida est, & tum cernitur, cùm caudam erigit. alis parùm ualet, atque idco breues facit uolatus. In stagnis, quæ nobilium ædes obducunt, & in piscinis apud Anglos plerunq; degit. Si quando periclitatur, ad arundineta densiora solet confugere.

DE TVRDO.

Κίχλα, turdus, Anglicè à thrusche, Germanicè eyn frammesjügel/ oder eyn wachholternügel.

[p. 140]

ARISTOTELES³.

Turdorum tria sunt genera, unū uisciuorū, quod nisi uisco resinaq; non uescitur, & magnitudine picæ est. Alterum pilare, quod sonat acute. & magnitudine merulæ est. Tertium quod iliacum quidam uocāt, minimum inter hæc, minusque maculis distinctum est. Mutat⁴ & turdus colorē, quippe collo æstate uarius, hyeme distinctus spectetur, uoce tamē eadē est. Turdus⁵ nidos ex luto, ut hirundines, facit, in excelsis arboribus, ita deinceps continuato opere,

¹ *Hist. An.* Bk VIII. 47.

² Another reading is πύγαργος; but this word is elsewhere used of an Eagle, cf. p. 30.

³ *Hist. An.* Bk IX. 96.

⁴ *Hist. An.* Bk IX. 254.

⁵ *Hist. An.* Bk VI. 3.

OF THE TRYNGA.

Τρύγγας, trynga, in English a water hen or a mot hen, in German eyn wasser hen.

ARISTOTLE.

The Junco and the Cinclus live on lakes and streams, as does the Trynga, which among these little birds is somewhat largest, for it equals in its size a Turdus ; all these wag their tails.

And now, that you may know what bird I think the Trynga is, it is an altogether dusky bird, save that part of the tail which lies above the vent, for that is white and only visible when it erects its tail. It is weak on the wing, and therefore takes short flights. In England for the most part it haunts moats which surround the houses of the great, and fishponds. If danger ever threatens it is wont to flee to the thicker reed-beds.

OF THE TURDUS.

Κίχλα, turdus, in English a thrusche, in German eyn krammesvögel or eyn wachholtervögel.

ARISTOTLE.

There are three kinds of Turdi, one of which is called the Viscivorus, since it feeds on naught but mistletoe and gum, and is of the size of a Pica. The second, the Pilaris, which has a sharp note, is of the same size as a Merula. The third, which some call the Iliacus, is the least of them and less marked with spots. The Turdus changes colour also, since it may be seen mottled upon the neck in summer, while in winter it is spotted, though its voice continues similar. The Turdus makes its nests of mud, as do Hirundines, and places them in lofty trees, the building

ut quasi catena quædã nidorum contexta uideatur.

PLINIUS¹.

Turdi hyeme maximè in Germania cernuntur.

[p. 141] *Primum turdi genus Angli peculiariter nominant a thrushe, & Germani (nisi me fallant aucupes, qui me sic uocare docuerunt) cyn crammesuogel. Secundum genus Angli uocant a throssel, aut a mauis, Germani cyn drossel, aut cyn durtsel. Tertiũ genus ab Anglis a uuyngthrushe, & à Germanis cyn uueingaerdsuoegel nuncupatur. Hic turdus utrinque iuxta oculos, & in pectore & in ipso alæ flexu, intus & foris maculas habet latiusculas rubras. Huius nidum nunquam uidi: nec mirum, quum per æstatem apud nos nusquam uideatur. primum genus non nisi hyeme in Anglia cernitur, aut si uideatur, rarum est. Secundum genus per totum annum apparet maculoso ualde pectore, & catus sui gratia à multis in canebis alitur. Nidum intus ex luto aut lignorum carie liquore mixta, & artificiosè leuigata, foris ex musco in ramis arborum aut fruticum facit.*

DE VIREONE.

Χλωρόν, uirco, Anglicè a uuitunol, Germanicè cyn witwol/ oder cyn weidwail/ oder cyn ferjeruise.

ARISTOTELES².

Vireo docilis, & ad uitæ munia ingeniosus [p. 142] notatur, sed malè uolat, nec grati coloris est. Vireo³ totus uiridis ex obscuro⁴ est, hyeme hic non uidetur, sed æstiuo solstitio potissimùm uenit in conspectum. Discedit exortu arcturi fyderis, magnitudine turturis est.

¹ *Hist. Nat. Lib. x. cap. xxiv.*

² *Hist. An. Bk IX. 89.*

³ *Hist. An. Bk IX. 98.*

⁴ These two words are not in Aristotle.

being so continuous as to seem almost like a chain of nests woven together.

PLINY.

The Turdi are in winter chiefly seen in Germany.

The first kind of Turdus Englishmen particularly name a Thrushe, and Germans (if the bird-catchers, who taught me so to call it, lead me not astray) eyn crammesvogel. The second kind the English call a Throssel or a Mavis, but the Germans say eyn drossel or eyn durstel. The third is named a Wyngthrushe by the English and eyn weingaerdsvogel by the Germans. This Turdus has broadish red spots on each side near the eyes, as well as on the breast, and also both inside and outside at the bend of the wing. But I have never seen its nest, nor is that wonderful, since it is nowhere to be seen with us throughout the summer. The first kind is not observed in England save in winter, or, if it be seen, it is unusual. The second kind, with a much spotted breast appears throughout the year, and by many is kept in cages for its song. It builds a nest, moreover, inwardly of mud or else of rotten wood tempered with moisture and smoothed skilfully, and outwardly of moss, upon the boughs of trees or shrubs.

OF THE VIREO.

Χλωρίων, vireo, in English a witwol, in German eyn witwol or eyn weidwail, or eyn kersenrife.

ARISTOTLE.

The Vireo is teachable, and is remarkable for its capacity for the duties of life ; but it flies badly and is not of a pleasing colour¹. The Vireo is wholly of a dusky green ; it is not seen in winter here, but comes chiefly in view about the summer solstice, it departs at the rise of the star Arcturus, and is of the size of the Turtur.

¹ Cf. p. 86.

Virconem (quod scio) in Anglia nunquam uidi, sed in Germania sæpissimè. turture paulò minor est. Vocem fistulæ grandiusculæ, quæ infimam cantionis partem sustinet, similem emittit. Nidum in ramo quem in summa arbore suspendit, in formam rotundam construit, ne cui hominum aut ferarū ad eum pateret aditus.

DE VPVPA.

ἐποπς, upupa, Anglicè a houpe, Germanicè eyn houp/ oder eyn widhopff.

ARISTOTELES¹.

Vpupa potissimū nidum è stercore hominis facit. Mutat faciē tempore æstatis & hyemis, sicut & cæterarum quoque agrestium plurimæ.
[p. 143] Vpupa² una in suo genere non nidificat, sed stipites arborum subiens, parit sine ullo stramento, in cauis.

Anglorū
lapuuingā
non esse
upupam.

Literatores pleriq; omnes Britannici, upupam eam nominant auem, quam barbari ab alarum strepitu, uannellum nuncupant, & ipsi sua lingua lapuuingam uocant. Verū istorum crassus error facilè auctoritate Plinij³ de upupa ita scribentis, confutatur: Vpupa (inquit) obsœna aliàs pastu auis, crista uifenda plicatili, cōtrahens eam, subrigensq; per longitudinem capitis. Hæc ille. Sed Grammaticis nostris hic error est facilè condonandus, nam nusquam in tota Britannia upupa (quod ego scio) reperiri potest, apud Germanos tamen frequentissima. Ea est magnitudine turdi, alis per interualla fuscis, albis & nigris pennis distinctis, crista in capite ab ea parte rostri, qua capiti committitur, ad extremum usq; occiput in lōgitudinem porrigitur, quam pro adscētib; suis aut contrahit,

¹ *Hist. An.* Bk IX. 88.

² *Hist. An.* Bk VI. 4.

³ *Hist. Nat.* Lib. X. cap. xxix.

I have never seen the Vireo in England, so far as I know, but very often when in Germany. It is a little smaller than the Turtur. It gives forth a note like that of the large pipe which plays the bass part of a song. This bird suspends its nest upon a branch at the top of a tree, and fashions it in rounded form, that it should not afford access to any man or beast.

OF THE UPUPA.

ἔπωψ, upupa, in English a howpe, in German eyn houp or eyn widhopff.

ARISTOTLE.

The Upupa builds its nest chiefly of human dung. It changes its appearance in the summer season and in winter, as very many other wild birds do. The Upupa only of its kind builds not a nest, but entering the trunks of trees lays eggs in cavities, without any litter.

Nearly all British writers name that bird Upupa, which from the noise of its wings foreigners call Vannellus, though in their own tongue the former call it Lapwing. Yet their gross error may be easily refuted on the authority of Pliny, who thus writes of the Upupa.

The Upupa (he says) is a bird filthy otherwise as to its food, but to be noticed for its folding crest, which it contracts and then erects again along its head.

These are his very words. And yet our scholars may be well excused this their mistake, for nowhere in the whole of Britain is the Upupa to be found (so far as I know), though in Germany it is most plentiful. The bird is of the bigness of a Thrush, with wings barred here and there with brown, and marked with black and white feathers; the crest extends from the part of the bill which joins the head to the extremity of the occiput, along the length, this it contracts

aut dilatat, ut equus aures arrigit aut demittit. tibijs est ualde breuibus, alis obtusioribus, & lentè admodum uolat.

[p. 144]

DE VRINATRICE.

Κολυμβρίς, *urinatorix*, *Angl. a douker, Germa. eyn bücher.*

ARISTOTELES¹.

Alia degunt quidem in fluido, uictumq; inde petunt, sed aërem nõ humorẽ recipiunt, & foris pärere solent. Complura huius generis sunt, partim gressilia ut lutra, latæ & crocodilus: partim uolucres ut mergi & urinatrices.

Aristoteles urinatricis unum tantum genus commemorat, ego tamen tria urinatricum genera uidi. Horum primum totum nigrum est. & si cirrum, quẽ in capite gerit, exceperis, mergo, quo tamen triplo minor est: cætera, quod ad corporis attinet effigiẽ, non dissimile est. & hoc genus nautæ nostrates lounam nominant, alij doukeram. Secundum genus turdo non maius est, anati colore & corporis effigie simile, et hoc Angli mediam urinatricem nuncupât. Tertium genus adeo nuper ab ouo exclusum refert anserculum, ut nisi rostrum huius paulò tenuius esset, ægrè alterum ab altero dis-
[p. 145] *cerneres. Non enim pennas, sed lanuginem quandam, earum loco obtinet. Degunt hæc plerunque tria genera in aquis stagnantibus, aut fluuijs non admodum rapidis, in quorum ripis arundines & carices nascuntur.*

DE VULTVRE.

Γύψ, *uultur*, *Anglicè a geir, Germanicè eyn geir.*

ARISTOTELES².

Vultur nidificat in excelsissimis rupibus: unde fit ut rarò nidus & pulli cernantur.

¹ *Hist. An.* Bk I. 6, somewhat freely rendered.

² *Hist. An.* Bk VI. 35.

or spreads again according as it is disposed, as a horse pricks or droops its ears. It has very short legs and rounded wings, while it flies somewhat slowly.

OF THE URINATRIX.

Κολυμβίς, urinatrix, in English a douker, in German eyn dücher.

ARISTOTLE.

But other animals in truth live in the water and thence seek their food, yet they breathe air and not moisture, and they are wont to breed out of the water. Now there are many of this sort, in part going afoot, as are the Lutra, Latax, and Crocodilus; and in part winged, as the Mergi and the Urinatrices.

Aristotle makes mention only of one kind of Urinatrix, but I have observed three kinds of Urinatrices. Of these the first is wholly black, and, except for the tuft it bears upon its head, is not unlike the Mergus otherwise, so far as the outline of its body goes, though it is one-third less in size. This is the sort our sailors call the Loun, but others the Douker. The second kind, no bigger than a Thrush, is like a Duck in colour and in form of body; this the English call the middle Urinatrix. The third kind, when it is but newly hatched, recalls a Gosling, so that if its beak were not a little more slender you could scarce discern the one bird from the other. For it has no quills, but in place of these a sort of down. These three kinds for the most part live on stagnant waters or not very rapid rivers, on the banks of which grow reeds and sedges.

OF THE VULTUR.

Γύψ, vultur, in English a geir, in German eyn geyr.

ARISTOTLE.

The Vulture nests in very lofty rocks, and thus it chances that the nest and young are rarely to be seen.

quocirca Herodotus Brifonis rhetoris pater, uultures ex diuerso orbe nobis incognito aduolare putauit, argumēto quòd nidum nemo uidisset uulturis, & quòd multi exercitum sequētes, repentē appareant. Sed quanquam difficile nidum eius alitis uideris: tamen uifus aliquan-
 [p. 146] do est. Pariunt uultures oua bina. Cætera, quæ carne uescuntur, non plus quàm semel anno pârere exploratum est.

PLINIUS¹.

Vulturum præualent nigri, nidos nemo attigit, ideo etiam fuêre, qui putarent ex aduerso orbe aduolare falso. Nidificant enim in excellentissimis rupibus. Fœtus quidem sæpè cernuntur ferè bini. Vmbricius auruspiciū nostro æuo peritissimus, pârere tradit tria, uno ex his reliqua oua nidūq; lustrare, moxq; abijcere. Tri-duo aut antè aut biduo uolare eos, ubi cadauera futura sunt.

Perperã Grammatici quidam uulturẽ, gryphem nominant, uulturem & gryphem ineptè confundentes. quum gryps sit a gryphen, animal ut creditur uolatile & quadrupes.

¹ *Hist. Nat. Lib. X. cap. vi.*

Wherefore Herodotus, the father of the rhetorician Briso, thought that Vultures winged it from some other world unknown to us, his argument being that nobody had ever seen a Vulture's nest, although a multitude at once come into sight when following an army. And yet, however difficult it be to see the nest of this bird, still it has been seen at certain times. Vultures lay two eggs each. Besides it is a well-known fact that animals which feed on flesh do not breed more than once a year.

PLINY.

Of Vultures the black are most plentiful. No one has ever reached their nests and therefore there have been some who erroneously thought that they flew hither from another world. They really nest in very lofty rocks. Indeed the offspring, generally twins, are often seen. Umbricius, the most skilful augur of our age, asserts that they lay three eggs, with one of which they cleanse the others and purify the nest, and afterwards throw it away. And that they fly three or two days beforehand to a place where carcasses are likely to be found.

Quite wrongly certain scholars call the Vulture Gryps, confounding foolishly the Vulture and the Gryps, since the Gryps is a Gryphon, or an animal believed to be both winged and quadruped.

AVIVM LOCI COMMVNES ex Ariftotele.

*SI Plinium, Ariftotclē, Ariftophanē aut quemcūq;
alium idoneum ſcriptorem te legere contingat,
locos huiusmodi cōmunes, quales exempli tantū gratia
ſubijciam, in procinctu, libro inſcriptos chartacco ha-
bere expedit, ut ad eos aues omnes, de quibus apud iſtos
legis, certo referas, quod ſi feceris, nō dubito quin in
aviū cognitione multū breui ſis profeſſurus.*

Appendices habentes.

*Olor, anſer, anas, gallinaceus, perdix, ciconia, aſca-
laphus, tarda, noctua, paſſer.*

Ingluivcs habentes.

Gallinaceus, palumbes, perdix & columbus.

Gulas totas amplas habentes.

Anſer, anas, gauia, cataraſta, & tarda.

Gregales aues.

Olor, anſer minor, grus, & platca.

Frugibus uiſitantes.

Palumbes, columbus, turtur, & uinago.

Lacus frequentantes.

*Ardeola, albardeola, ciconia, gauia cineria, iunco,
cinclus, trynga, calidris, & alcedones.*

Mare frequentantes.

*Alcedo, carulus, gauia alba, fulica, mergus, rupes
& cataraſta.*

COMMON PLACES, REFERRING TO BIRDS,
FROM ARISTOTLE.

IF it should happen that you read Pliny, Aristotle, Aristophanes, or any other suitable writer, it is fitting to have ready for use such Common Places of this kind as I will add for the sake of example only, written in a note-book, that you may with certainty refer to all those birds, of which you read in their pages. And if you do this, I doubt not that in a short time you will make great progress in the knowledge of birds.

Those having appendices¹.

Olor, anser, anas, gallinaceus, perdix, ciconia, ascalaphus, tarda, noctua, passer.

Those having craves.

Gallinaceus, palumbes, perdix, and columbus.

Those having wide gullets.

Anser, anas, gavia, cataracta, and tarda.

Gregarious birds.

Olor, the smaller anser, grus, and platea.

Those living on crops.

Palumbes, columbus, turtur, and vinago.

Those frequenting lakes.

Ardeola, albardeola, ciconia, the grey gavia, junco, cinclus, trynga, calidris, and alcedones.

Those frequenting the sea.

Alcedo, carulus², the white gavia, fulica, mergus, rupex³, and cataracta.

¹ That is *cæca* or blind-guts.

² κύανος.

³ Perhaps χαρδριός.

Amnes & lacus frequentantes.

*Olor, anas, phalaris, urinatrix, bosca, coruus pal-
mipes, uterque anser, uulpauser, capella, penclops, aquila
marina.*

Spinas appetentes.

Spinus, carduelis, & auriuittis.

Culicibus uictitantes.

Pici Martij, galgulus, culicilega.

*Vermibus aut ex toto aut magna ex parte
uictitantes.*

*Fringilla, passer, rubetra, luteola, & pari omnes,
ficcdula, atricapilla, rubicilla, rubecula, syluia, curuca,
asylus, florus, montifringilla, regulus & frugilega.*

*Plures locos cuique licebit huius modi
excogitare.*

Those frequenting rivers and lakes.

Olor, anas, phalaris, urinatrix, bosca, the web-footed corvus, either kind of anser, vulpanser, capella, penelops, the sea aquila.

Those feeding on thistles.

Spinus, carduelis, and aurivittis.

Those feeding on insects.

Pici martii, galgulus, culicilega.

Those feeding on worms, either wholly, or for the most part.

Fringilla, passer, rubetra, luteola, and all the pari, ficedula, atricapilla, rubicilla, rubecula, sylvia, curuca, asylus, florus, montifringilla, regulus, and frugilega.

Any one may devise more Places of this sort.

NON deerunt forsan, qui mihi hoc uicio
 uerfuri funt, quòd libellus iste meus
 coniecturarum multò plus quàm certarum ad-
 fertionum in se contineat : quibus respondeo, in
 re ardua, & nondum fatis explorata mihi con-
 sultius & modestius uideri, hēſitanter & modeste
 coniectando uestigare, & ita inquirere, quàm
 temere & impudenter de rebus incompertis pro-
 nunciare. Quòd autem de moribus & medicinis
 auium nihil hic ſcripſerim, in cauſa fuerunt,
 temporis anni infœlicitas, & anguſtia (breuiori
 enim ſpacio quàm duobus menſibus totus liber
 [p. 150] conſcriptus eſt) & pecuniæ copia minor, quàm
 quæ huiuſmodi negotio abſoluendo ſufficeret.
 Nam quis ſine magna pecuniæ ui in longinquas
 regiones proficiſci poteſt, peregrinarum auium
 formas & mores contemplaturus, & illic diu ea
 de cauſa manſurus? Quis familia aut uoca-
 tione ſua, aut alijs negocijs foras prodire pro-
 hibitus, ſine maximis impenſis omnia auium
 genera ab aucupibus ad ſe ex uarijs mūdi plagis
 allata, curare poterit? et ià allata, quò mores
 ad plenum perueſtiget, in uiuarijs & caueis ſine
 maximis ſumptibus alere quis ſufficiet? Hoc

PERORATION TO THE READER.

THESE perhaps will not be wanting those who will attribute this to me as a fault, that this little book of mine contains within it many more conjectures than sure statements: to whom I reply, that it seemed to me much more prudent and becoming on a subject that is difficult and not yet sufficiently explored to tread doubtingly and modestly by conjecture, and so to enquire, than to pronounce rashly and immodestly on things undetermined. Moreover that I have written nothing here of the habits and medicinal nature of birds, I have for reasons the unsuitability of the time of year and its brief span (for the whole book was written in a space of less than two months) and a supply of money too slender to suffice for the perfection of a work of that kind. For who without great command of money can set off for distant regions, to observe the forms and habits of foreign birds, and there to stay a long time for that purpose? Who, hindered by his household or his calling or other business from going abroad could without vast expense give heed to all the kinds of birds brought to him by fowlers from the various quarters of the world? and when brought, who would be capable without vast expense of keeping them in vivaria and cages, that he might investigate their habits to the full?

Alexander ille omnium ethnicorum regum potentia, bellica gloria, & literarum studio maximus & nobilissimus fecū animo perpendens, [p. 151] Aristotelem iam de animalibus scripturū, quem priuatis sumptibus negocium illud absolueri non potuisse cognouerat, ad conducendos aucupes, & uenatores, & ad alenda in uiuarijs animalia iam capta, 480 milibus coronatorum donauit, & ita instruxit. Talis si hodie alicubi Alexander existeret, non dubitarem quin nouus nobis Aristoteles alicunde renasceretur, qui prioris illius Aristotelis animalia omnia, paucis exceptis, nō solum nobis cum moribus medicis suis facultatibus, & huius tēporis nominibus exhiberet, sed & multa plura animalia quā prior, & humano generi non minūs utilia nos doceret. Quōd igitur Aristoteles de tam multis animantibus [p. 152] tā fœliciter scripserit, Alexā dri potius liberalitati quā Aristotel. diligentē tribuendum erat: quamuis et ea summo opere laudāda erat. Nam si Alexandri munificentia Aristoteli animalia illa spectanda nō exhibuisset, historiam animalium nobis tam absolutam nunquam ædidisset. Mirari igitur definant scriptorum huius temporis ociosi cēsores, frigidius, indoctius, & minori cum diligentia stirpium, auium, piscium & quadrupedum historias hac ætate cōscribi, quā apud seculum prius tractabantur. Quum hodie quicquid præclari in lucem emittitur, priuatis tenuioris fortunæ studiosorū uirorum sumptibus & typographorum impenſis edatur. Sed ad te iam redeo, candidissime lector, quem etiam atq; etiam ob- [p. 153] testor, ut si qua tibi sese peregrina facie offerat

The well-known Alexander, the greatest and most renowned of all kings of the nations in power, war-like glory, and zeal for learning, weighing this in his mind, presented Aristotle with 480,000 crowns, when he was about to write on animals, since he knew that the philosopher could not carry out that task with his private means, for the purpose of hiring fowlers and hunters, and for keeping in vivaria the animals which had been already caught, and provided to that end. If such an Alexander existed anywhere to-day, I should not doubt that a new Aristotle would be born again for us from somewhere, who not only would display to us all the animals, with few exceptions, of that former Aristotle, with their habits, their medicinal properties, and their latter-day names, but would inform us of many more animals than the former, and those not less useful to the human race. That Aristotle therefore wrote so happily about so many living creatures is to be put down rather to the liberality of Alexander than to the diligence of Aristotle: though that too must be praised without stint. For if the bounty of Alexander had not supplied to Aristotle those animals to be examined, he never would have published so complete a History of Animals for us. Therefore let the ease-loving critics of the present day cease to wonder that the histories of plants, birds, fishes, and quadrupeds are written in this age with less spirit, less learning, and less diligence than that with which they were treated in a former age. To-day whatever of value is brought to light is published at the private expense of very zealous men of slender fortunes and printers. But now I return to you, most ingenuous reader, and beseech you once and again that, if any bird of foreign aspect meet

auis, paucis mihi illam, addito nomine gentis tuæ,
depingere non dedigneris, & mihi & omnium
bonarū literarum studiosis fcies te magnopere
gratificaturum, nam nec facti in fecunda huius
libri editione ero immemor, nec quicquā, quod ad
hoc disciplinæ genus pertinet,
mihi exploratum, te celabo.
Vale. Coloniæ Calend.
Martijs.

your eye, you will not disdain to depict it for me in a few words, with the addition of the name of your family, and you will know that you will greatly gratify both me and those who are zealous of all good learning, for I shall be neither unmindful of the act in a second edition of this book, nor conceal from you anything, which pertains to this kind of teaching, and is found out by me.

Farewell. At Cullen. 1st March.

ΑΛΒΕΡΤΟΣ Ὁ ΓΕΛΡΙΕΤΣ

τῷ τῆς βίβλου σπουδαίῳ ἀναγνώστῃ
 ξυπράττειν.

Ποίκιλα εἰ ἐθέλεις πτην' εἰδέναι ὧ φιλόμουσε
 Τοῦτο ποίημα βραχὺ τάχ' ἀνάγνωσον ἄγε.
 Οὐδεὶς ἐστ' ἀκριβῶς περὶ τούτων γράψεν ἱατρῶν,
 Ὡστε βλέπεις τῇ δ' ἐν πάντα γραφέντα βίβλῳ.
 Τοῦ ὄν σπουδαίου Τουρνήρου τὸν πόνον οὗτον
 Τπέλαβεν: τούτου καὶ ἀπόλαυε καλῶς.

Ἄλλο.

Φωνὰς μανθανέμεν πτηνῶν χιλίας ἀναγνωστά
 Βούλεις Τουρνήρου τὴν βίβλον ὠνέεο.
 Τοῖα γὰρ οὐκ ἱατρῶν μηδεὶς πρὶν ἐγράψατο πάντων,
 Ἐν βιβλίῳ τούτῳ, ποῖα δοθέντα βλέπεις.
 Εἰ ἄρ' ἀβρῶς τε σαφῶς, τῆς γραφθείσης ἀπολαύειν
 Βίβλου ἀνδάνει, κίνεε δεῦρο πόδον.

Aliud eiusdem ad candidum lectorem.

*Accipe quæ docti medici tibi cura paravit
 Turneri, notas quisquis auebis aues.
 Has tibi tam uarijs manus ingeniosa figuris
 Expressit, nosse ut quamlibet inde queas.
 Ne quoque non posses has pernouisse, Latinas,
 Anglas, Teutonicas, Argolicasq; facit.*

Liber ad lectorem.

*Quisquis aues uarias de nomine deq; figuris
 Nōsse cupis lector, me lege, doctus eris.
 Nec dabo, crede mihi, tibi munera parua laboris:
 Nam uolucrum res est maxima scire genus.*

Crebrò Grāmatici hīc hærēt, stāt κωφὰ πρόσωπα,

Nec facit officio stultula turba satis.

Hīc ipsos medicos errare miserrima res est,

Quos decet hēc animis nota tenere suis.

Seu Maro sit pueris, seu Naso pōta legendus,

Seu fuerint quævis scripta legenda tibi:

Dispeream, si non multò tibi maximus error

Occurret passim, nī bene nōris aues.

Autorum nimiam placet haud posuisse cateruam.

Vt tibi, quæ teneo, nostra probare queam.

Plinius hīc ille est, & Aristoteles, reliquiq̃,

Quotquot de uolucrum nos ratione docent.

Hoc scio, Turneri miraberis ipse laborem,

Doctrinam, summam cum pietate fidem.

[p. 156]

Αδελον.

En tibi, quos docti dedit hīc pia cura labores

Turneri medici, candidè lētor habe.

Inuenies nimium quæ te didicisse iuuabit,

Hinc uenient fructus in tua uota boni.

FINIS.

APPENDIX.

EXCERPTA EX OPERE IOANNIS CAIL BRI-
TANNI DE RARIORUM ANIMALIUM ATQUE
STIRPIUM HISTORIA, fol. 17—23.

De Haliaeto.

De Auibus.

HALIAETOS, id genus aquilæ est, quod ex mari lacubusq; prædam quærit, vnde nomen inuenit. Is magnitudine Milui est, capite albis & fuscis distincto [f. 17 b] lineis, vt melino: rostro aquilino: oculis in medio nigris, in ambitu aureis: lingua ferè humana, nisi quòd ad radicem vtrinq; habet appendicem: colore per summa asturis, per ima albo: gutture maculis notato ruffis vt & ventre, pectore medio pure candido: crure crasso & squamoso: pede vncungui & cæruleo: digitis quatuor, per superna ad dimidiam longitudinem etiam squamosis, ad reliquam incis, per inferna asperis & aculeatis tenacitatis causa: & his tam validis, vt flexos vix vlla vi extendas. Prædator is est piscium, discussis decidentis corporis impetu aquis, ex eisq; viuut. Et quamuis ex pisce viuat, fidipes tamen est vtroque pede, non altero palmipes, vt vulgus putat. Giraldus Cambrensis libro de Topographia Hiberniæ, vbi de auibus biformibus agit, hunc Aurisfrisiū vocat: & altero pede aperto & vnguibus armato esse, altero clauso cum vulgo scribit. Supra magnitudinem corporis alæ longitudo est, quæ ad pedes Romanos duos & digitos vndecim extenditur. Inoleuit opinio istic apud nostrum vulgus in Britannia, eam inesse vim naturalem huic aui, vt quem conspexerit piscem, eum se quàm mox resupinare & conuertere, atque ad sum-

APPENDIX.

EXTRACTS FROM THE WORK OF JOHN CAIUS
'DE RARIORUM ANIMALIUM ATQUE STIRPIUM
HISTORIA' (1570).

OF THE SEA EAGLE.

THE *Haliaetos* is that kind of Eagle, which seeks its prey from the sea and lakes, whence it takes its name. It is of the size of a Kite, having the head marked with white and dusky lines, as in a badger; an Eagle's beak; eyes black in the middle, golden in the outer circle; a tongue almost like that of man, except that at the root it has an appendage on either side; the colour above that of a Goshawk, white below; the throat marked with rufous spots, as is the belly; the middle of the breast pure white; the legs thick and scaly; the foot with curved claws and blue; four toes scaly above for quite half of their length, fissured for the rest, rough on the lower part and sharp for a firm hold; and these so strong that you can scarcely straighten them by any force when bent. This bird is a preyer upon fishes, the water being cleft by the shock of its body as it plunges, and on them it lives. And though it lives on fishes, yet it is cloven on each foot, not webbed on one as the vulgar think. Giraldus Cambrensis in his book on the Topography of Ireland, when he treats of unequally formed birds, calls this the *Aurifrisius*¹; and writes in common with the vulgar that it has one foot free-toed and armed with claws, the other closed (with webs). The length of the wing surpasses that of the body, for it extends to two Roman feet and eleven inches. In this affair an opinion has grown up among our common people in Britain, that such a natural power exists in this bird that any fish which it sees turns upwards on its back as soon as possible and rises to the surface

¹ *Aurifrisius* must be the Latinized form of the old French name 'Orfraie'—which, like Osprey, is a corruption of *Ossifraga*.

mam aquam ascendere, in eaque fluitare vt sopitum, quo facilior præda sit volanti. Ideoq, eius adeps studiosius asseruatur à nostris piscatoribus, quòd eandem vim habere creditur. Frequentes sunt apud nos in maritimis locis & Vecti insula. Nostri an Osprey vocant. Moribus placidus est & tractabilis, & famis [f. 18] patientissimus. Vixit enim septem dies apud me sine cibo, & in alta quiete: nisi si hoc non mos fecit sed fames, quæ omnia domat. Carnem oblatam recusauit: piscem non obtuli, quòd cum ex hoc viuere didici. Caro illi nigra est.

De anserẽ Brendino.

Anser Brendinus, auis est marina, palmipes, figura anseris, sed magnitudine paulo infra, capite albo exiguo & curto, sed rostro nigro, à quo linea nigra ad oculum vtrumque ducitur, collo fusco, à pectore ad caudam ex dimidio corpore inferiori albo, coxendice murino (vt est Columbæ vulgaris color aut gruis) ex superiori, ad collum fusco, vt & ad caudam medio inter vtrumq, murino: alis item murinis, cum cauda longitudine paribus, sed pennis ad extremum obfuscat. Cauda nigra est ex albo enata, pede nigro & palmato. Gregalis auis est & garrula. Ex pisce viuuit, frequens apud nos per littora in Britannia. Vulgus Britannorum quod ad littus habitat à coloris varietate a Brendgose nominat. Ornithopolæ Londinenses Bernaclum vocant, cum dicendum putem Berndclacum seu Brendclacum, quòd antiqui Britanni atque item Scoti, anseres marinos, palustres & lacustres omnes Clakes dicebant, cum tamen hodie corruptè dicimus Fenlakes & Fenlagges, cum dicendum fuit Fenclakes. Varium item colorem, Brend, seu per metathesim Bernd iidem appellabant. Vnde [f. 18 b] bernded seu brended id animal dicitur, quod in colore murino variegatum est albo, vt est hic anser. Non est

of the water, and therein floats as if stunned, in order that it may more readily become a prey to the winged creature. And therefore is its fat preserved with greater keenness by our fishermen because it is believed to have the same virtue. They are abundant with us on sea-coasts and in the Isle of Wight. Our people call it an Osprey. In its habits it is quiet and amenable, and most patient of hunger. For it has lived with me for seven days without food, in deep repose; unless indeed it was not habit that caused this, but hunger, which tames all things. It refused flesh when offered to it; I did not offer fish, because I understood that it lived on it. Its flesh is dark.

OF THE BRENT GOOSE.

The Brent Goose is a web-footed sea-bird, of the appearance of a Goose, but a little less in size, with the head small and short and white, but the beak black, from which a black line reaches to either eye; the neck dusky, the lower half of the body from the breast to the tail white; the flank mouse-coloured above (like the colour of a common Pigeon or Crane), dusky towards the neck, and also the intermediate parts towards the tail mouse-coloured; the wings also mouse-coloured, equal in length to the tail, but with the feathers darkened at the tip. The tail is black from a white base, the foot black and webbed. It is a gregarious and noisy bird. It lives on fish, and is common with us in Britain about the coasts. The common people of the Britons who live on the coast, name it a Brendgose from the varied nature of its colouring. The London bird-dealers call it Bernacle, yet I should think that it ought to be named Berndclac or Brendclac, because the Britons of old, as also the Scots, called all the Geese whether of the sea, marshes or lakes "Clakes," though at the present day we corruptly say Fenlakes and Fenlagges¹, when we should say Fenclakes. The same people, moreover, call anything variegated Brend or by metathesis Bernd. Whence an animal is said to be bernded or brended which is variegated with white upon a mouse-coloured ground², like this Goose. It is not, there-

¹ According to this notion 'Grey-Lag' would be for 'Grey-Clak.'

² Possibly, then, 'Bergander' is for 'Bernd-gander.'

Anfer
Baffanus.

ergo Scotorum Baffanus anfer, qui in Baffe Scotorum Infula nidum ponit atq, oua, à qua nomen habet. In hanc infulã rupem existentem, nec in fummo quantam Miluus oberret (vt Poëta dixit) fed exiguam, venturi ftato anni tempore anferes, quo prius fpeculatu, qua obferuatione præmiffis nuntijs vtantur quàm ingrediuntur: quo anni tempore hoc faciant, qua folitudine infulæ, concludentibus fe incolis ad aliquot dies, donec fe firmauerint anferes, ne abigant, quanta multitudine atq, denfitate inuolent, fic vt in ferenitate folem adimant, quot pifces afferant, quot oua pariant, & quantos fructus in annos ex eis anferumq, plumis atq, oleo percipiant infulani (nam Pupinorum pinguedinem habent atq, guftũ) longum effet recensere.

De Anate Indica.

Eft apud nos ex India anas, eadem planè corporis figura, eodem roftro & pede quo vulgaris, fed ex dimidio maior ea & grauior. Caput illi rubefcit vt fanguis, & bona pars coniuncti colli à posteriore parte. Id totum callofa caro eft, & incifuris diftincta: quaq, ad nares finit, carunculam demittit à reliqua carne figura feperatam, qualis cygnis eft, roftro coniunctam. Nudum plumis caput eft, & ea quoque colli pars quæ rubefcit, nifi quòd in fummo capite crifta eft plumea atque candida, per totam capitis longitudinem pro-
[f. 19] tenfa: quam, cum excandefcit, erigit. Sub oculis ad roftri initium per inferna, inordinatæ maculæ nigræ carni funt inductæ: & vna atq, altera à fummo oculo ad fuperna eleuatæ. Oculus flauefcit, feperatus à reliquo capite circulo nigro. Sub extremo oculo in auerfum macula eft fingularis, feperata à ceteris. Roftum totum eft cœruleum, nifi quòd in extremo macula nigrefcit vna. Pluma illi per totum colli proceffum reliquum, alba. Qua corpori collum iūgitur, circulus eft plumeus niger, rara pluma alba, maculofus & inæqualis, per ima anguftior, per fumma latior. Poft eum per totum imum ventrem pluma

fore, the Bass Goose of the Scots, which has its nest and eggs on the Bass, a Scottish Isle, and thence takes its name. Now when at a certain season of the year the Geese are about to return to this precipitous island rock—not so big on the top as a Kite could hover over (as the Poet has said), but very small—it would be too long to recount what spying, what circumspection (scouts having been sent ahead) they use before they alight: at what time of year they do this, the solitary state of the isle, when the inhabitants shut themselves up for several days, until the Geese have settled down, lest they should drive them off, in what numbers and in what a throng they fly to it, so that in clear weather they obscure the sun, how many fishes they bring home, how many eggs they lay, and what profit the dwellers in the isle make annually from the feathers and the oil of these Geese (for they possess the fatness and the taste of Pupins).

OF THE INDIAN DUCK.

There is among us, a Duck from India, with exactly the same form of body, the same beak and foot as the common bird, but bigger by half and heavier. Its head is red as blood, as is a good part of the adjoining neck behind. The whole of this is callous flesh and marked with fissures: and where it ends at the nostrils it makes a caruncle like that in Swans, separated in form from the rest of the flesh, which joins the beak. The head and red part of the neck are devoid of feathers, save that on the top of the head is a white feathery crest, extending over the whole length of the head; and this the bird erects, when it is excited. Under the eyes to the beginning of the beak at its lower part irregular black spots are arranged on the flesh: and one or two reach upward from the top of the eye to the parts above. The eye is yellowish, being separated by a black ring from the rest of the head. Close behind the eye is a solitary mark, apart from the rest. The whole beak is blue save that at the tip one spot shews black. The plumage over all the rest of the neck is white. Where the neck joins the body, there is a ring of black feathers spotted and irregular—with an occasional white one—narrower below, broader above. Behind this the plumage is white over the whole of the belly below:

alba est: per summum corpus, fusca, sed ab circulo illo nigro pluma alba in summo diuisa. Extremæ alæ atque cauda cum splendore virescunt, vt Cantharides. Tibiarum cutis fusca est, incisuris leuibus per circuitus ducta. Membrana per interualla digitorum pedis pallefcit magis, vna atque altera resperfa macula fusca, incerta lege disposita, nisi in interuallo sinistri pedis, vbi sex per digiti extremi longitudinē disponuntur. Tardo gradu incedit propter corporis grauitatem. Vox illi non qualis ceteris anatibus, sed rauca, qualis faucibus humanis catarrho obfessis. Mas maior est quàm fœmina. Ea similis mari est, nisi quòd non ita variegato corporis colore est. Viuit ex cœnosis aquis, & alijs quibus cetera vulgaris anas gaudet.

[f. 19 b]

De Anate Turcica siue Indica altera.

Anati quidem similis est quæ Turcica siue Indica dicitur, sed quantitate & magnitudine corporis anferem ferè diceres. Tota est candida, nisi quòd rostrum, tibiæ, atque pedes rubent, genæq; item callosa carne, & rostri tuber supra nares. Caro illi dulcis est, & vox sibilus. Sunt eius generis quedam, colore albo & nigro variegatæ. In aqua viuit, locisq; gaudet cœnosis vt ceteræ anates.

De Pica marina.

Pica marina (vt nostrum vulgus nominat) paulo maior est terrestri, coloris quidem varietate in corpore similis, sed colore pedis, digitorum numero, incisuris, cauda atque rostro admodum dissimilis. Nam pes rubet, & digito posteriori destituitur, nec incisuras habet is & tibia, sed impressiones quasdã, piscium squamis quàm simillimas. Fissus est, sed ita in digitis vtrinq; luxuriat cutis, quemadmodũ in fulicis penè, vt ad natandũ quoque pes factus videatur. Aus est Amphibios. Caudam curtam habet, rostrum longum atq; tenue, perpendiculariter latum, non teres, colore

and dusky over the upper surface of the body, but the white feathering above is divided by the black ring. The ends of the wings and the tail have a greenish gloss as in *Cantharides*. The skin of the legs is dusky, marked all round with slight fissures. The web in the spaces between the toes is paler, marked now and again with a dusky spot, arranged in no precise plan, except in those of the left foot, where six are distributed over the length of the outer toe. The bird moves with slow step on account of the weight of its body. Its voice is not like that of other Ducks, but hoarse; such (as comes) from the human throat when attacked by a catarrh. The male is larger than the female. She is like the male, save that she has not so varied a colouring of body. The bird gets its living from muddy waters, and those others wherein the other common Duck delights.

OF THE TURKISH OR SECOND INDIAN DUCK.

That which is called the Turkish or Indian is like a Duck, but (judging) from the bulk and bigness of its body you would almost call it a Goose. It is entirely white, except that the beak, legs and feet are red, while the cheeks also have callous skin, and there is a protuberance on the beak above the nostrils. Its flesh is sweet, and its voice whistling. There are some of this kind variegated with black and white. It lives upon the water, and delights in muddy places, as do other Ducks.

OF THE SEA PIE.

The Sea Pie (as our common people call it) is a little larger than the land Pie, yet like it in the varied colour of its body, while very dissimilar in the colour of the foot, the number of toes, their scutellations, the tail and the beak. For the foot is red and lacks a hind toe, nor has that member or the tibia scutellations, but merely marks, as like as may be to the scales of fishes. It is cloven-footed, but there is such an extent of skin on the toes on each side—almost as much as in Coots—that the foot would even seem to be formed for swimming. It is an amphibious bird. It has a short tail, a beak long and thin, vertically broad, not smooth, in colour

in fummo ad caput rubrum, per reliquum pallidum, nec est in fine acutum, sed obtusum id. In mensa grata auis est.

De gallina Getula domestica.

Gallina Getula domestica, paulo minor nostra est, [f. 20] colore in supernis obscurè ruffa, in infernis pallida, pluma capitis incompressa & erecta, crista ferrata, humili, simplici, carnea: gutturosa magis perpetuò, quàm nostræ cum gloriunt: tibijs pedibusque plumosis, maximè per exteriora & posteriora ut columbis, ne per interiores plumas gressus impedirentur: cetera ut vulgaris.

De Meleagride.

Meleagris pulcherrima auis est, magnitudine corporis, figura, rostro, & pede Phasianus similis: vertice corneo, in apicem corneum à posteriori parte precipitem, in anteriori leniter acclivem eleuato, armatus. Eum natura voluisse videtur inferiori capitis parti tribus veluti lacinijs se promittentibus committere atque deligare: inter oculum & aurem vtrinque vna, & in fronte media item vna: omnibus eiusdem cum vertice coloris: ita ut infideat capiti eo modo, quo ducalis pileus illustrissimo duci Veneto, si quod iam aduersum est, auersum fieret. Rugosus is est: inferius, per circuitum: qua se attolit, in directum. In summo collo ad occipitium nascuntur erecti quidam atque nigri pili (non plumæ) in contrarium versi. Oculi toti nigri, æquè & in orbem palpebræ atque cilia, si maculam in summa & posteriori parte supercilij vtriusque demas. Imum caput per longitudinem vtrinque caro quædam callosa colore sanguineo occupat, quæ ne propendeat veluti Galli gallinacei palea¹, ut [f. 20 b] replicaretur natura voluit, & auerso ductu in duos processus acutos à capite liberos finiret. Ex hac carne attollunt se vtrinque carunculæ, quibus nares in ambitu vestiuntur, & caput in anteriori parte à cetero rostro pallido separatur. Harum ad rostrum margines inferiores, replicantur etiam leuiter sub utroque nare.

¹ Possibly 'palea' is a misprint for 'galea,' comb.

red on the top near the head, pale on the remainder, nor is it sharp at the end, but blunt. It is an acceptable bird for the table.

OF THE DOMESTIC GETULIAN HEN.

The domestic Getulian Hen, is a little smaller than our own, in colour dull rufous above, pale below, with an erect crest of disintegrated feathers on the head, a serrated comb, low, simple and fleshy: more continuously noisy than ours are, when they cluck: with the legs and feet feathered, for the most part outwardly and behind as in Doves, that the progress should not be hindered by feathers on the inside, otherwise it is like the common kind.

OF THE MELEAGRIS.

The Meleagris is a very beautiful bird, like to a Pheasant in bigness of body, form, beak and foot: provided with a horny poll rising to an abrupt horny peak at the back, which slopes down gently in front. Nature seems to have designed to join and bind this to the lower part of the head by three hanging lappets as it were; one on each side between the eye and the ear, and also one on the middle of the forehead, all of the same colour as the poll, so that it sits on the head in the same way that the ducal cap does on that of the most noble Duke of Venice, if that part which is usually in front be turned behind. It is wrinkled round about below, but vertically where it rises above. From the top of the neck to the occiput spring certain erect black bristles (not feathers), turned backwards. The eyes are wholly black, and equally so are the eyelids and eyelashes around them, if you except a mark on the top and back of each eyebrow. A kind of callous flesh of a blood-red colour covers the lower part of the head along its length; nature has designed that it should be folded, and should not hang forward like the wattle of the Fowl, and being led backward end in two acute processes free from the head. From this flesh rise on either side caruncles, by which the nostrils are clothed round about, and by which the head is divided in front from the rest of the pale-coloured beak. The lower edges of these by the beak are also folded slightly under each nostril. What intervenes

Quod inter verticem & carnem est à dextra & sinistra parte, album deplume est, leui cœruleo mixtum. Color verticis atque apicis, idem prorsus est cū colore dactyli. Tibiæ nigræ sunt, & in anteriori parte, squamosa incisura duplici notatæ: in posteriori nulla, sed leues, & veluti punctis quibusdam sui coloris respersæ. Color illi sub faucibus exquisitè est purpureus: in collo obscurè purpureus: in cetero corpore per summa contuenti, qualis confurgit si album & nigrum pollinem vtcunque tenuiter tritum, colori fusco rarius aspergas, nec tamen commisceas. Tali colori maculæ albæ ouales aut rotundæ per totum corpus inesse visuntur, per summa minores, per ima maiores, comprehensæ interuallis linearum (vt apparet in plumarum compositione naturali) quæ se mutuo interfecant obliquo hinc inde ductu, per summa tantum corporis, non item per ima. Id non ex toto corpore solum deprehendes, sed ex singulis auulsis plumis. Superiores enim, obliquis lineis se mutuo interfecantibus, aut, si mauis, orbiculis quibusdam ex albo & nigro (vt dixi) polline confectis, & per extremitates coniunctis, vt in fauis aut rhetibus, maculas ouales aut [f. 21] rotundas albas in spacijs fuscis comprehendunt: inferiores non item. Vtręq; tamen simili lege positæ sunt. Nam in alijs plumis, ordine ita iunctæ sunt, vt ferè triangulos acutos faciant: in alijs, vt oualem figurā representent. Huius generis ordines tres aut quatuor in singulis suis plumis sunt, ita vt minores in maiorū complexu reponantur. In extremis alis & in cauda, rectis lineis equidistantibus procedunt per longitudinem maculæ. Inter gallum & gallinam vix discernes, tanta est similitudo, nisi quòd gallinæ caput totum nigrum est. Vox illi est diuisus sibilus, non fonorior, non maior voce coturnicis, sed similior voci perdicis, nisi quòd submissior ea est, nec ita clara. Cursu velox est.

De Morinello.

Morinellus, auis nobis cum Morinis communis, stulta admodum est, sed in cibis delicata, coque apud

between the poll and the flesh on the right and left is white and unfeathered, mixed with light blue. - The colour of the poll and peak of the head is exactly the same as that of the toes. The legs are black, marked in front with a double scaly fissure, on the back with none, but smooth, and sprinkled as it were with some spots, of a peculiar colour. The colour below the jaws is exquisite purple; on the neck dark purple; on the rest of the body, if you look from above, it stands out as if you sprinkled black and white flour, ground very fine with dusky colour at intervals, and yet did not mix them up. On such a ground oval or round white spots seem to be imposed throughout the whole body, smaller above and larger below, arranged in lines at intervals (as appears in the natural structure of feathers) which cut one another here and there in reciprocal oblique arrangement, only on the top of the body, however, and not below. This you will observe not only from the body as a whole, but from individual feathers if plucked out. For the upper feathers, with their oblique lines cutting one another reciprocally, or, if you will it, with certain rounds composed of black and white flour (as I have said), and meeting towards the tips, as in honey-combs or nets, enclose oval or round white spots within dusky spaces: but the lower do not. Both, however, are arranged by a similar law, for on some feathers they are so joined in rows as to almost make acute triangles; in others so as to present an oval form. Of this kind there are three or four rows on each feather, so that the smaller are contained in the compass of the larger. At the tips of the wings and on the tail, spots lie along the length in equidistant straight lines. You can hardly distinguish between the cock and the hen, so great is the likeness, save that the head of the hen is wholly black. The voice is a shrill double cry, not more sonorous or louder than that of the Quail, but like that of the Partridge, except that the latter is lower, and not so clear. In running it is swift.

OF THE MORINELLUS.

The Morinellus, a bird common to us and the Morini¹, is very foolish, but delicate to eat, and on that account is a

¹ The people of a district in Northern France.

nos in fummis delitijs atque pretio est. Imitatrix auis est. Ideo, vt Scops & Otus saltandi imitatione, ita hæc noctu ad lumen candelæ pro capientis gestu capitur. Nam si is expandit brachium, extendit & illa alam: si is tibiam, & illa itidem. Breuiter quicquid gerit auceps, idem facit & ales. Ita humanis gestibus intenta auis, ab aucipe decipitur, & rhete obuelatur. Auis parua est, magnitudine Sturni, tribus tantum digitis anterioribus, posteriori nullo, vertice nigro, genis candidis, coturnicis ferè colore, si cinericei [f. 21 b] parum admisceas, potissimum circa collum. Morinellum voco duplici de causa, & quòd auis est apud Morinos frequētissima, & quòd auis stolidia est, quæ stultitia græcis *μωρότης* dicitur. Eam ob rem nostri etiam Doterellū vocant, quasi stultitia delirantem dicant.

De Puphino siue Pupino.

Est auis quædam marina nostras, parui anatis (quàm *βόσκαδα* Græci vocant) magnitudine & figura corporis, pedibus palmatis & rubescentibus, ad posteriora magis positus quàm ceteris palmipedibus exceptis pygoscelibus: rostro tenuiore magis latitudine se demittente, quàm longiore processu se extendente, quatuor incisuris rubris à summa, duabus ab ima parte fulcato, in colore pallentis ochræ. Quod inter has & caput est, subcæruleum est, & ea figura qua luna est, cum exacti dies decem sunt à coitu. Per summa corporis totius nigrescit, nisi qua oculi sunt, qui in albo constituti sunt: per ima exalbescit tota, nisi summo pectore, qua nigricat. Viuit ex mari. Hunc nostri Puphinum dicunt, nos Pupinum à naturali voce pupin. Latitat in cauernis, vt charadrius. Eam ob rem educita è cuniculi cauea auis hæc est, loco non procul à mari posito, à venatore quodam immissa viuerra. In piscis vsu apud nos est in solenni ieiunio per [f. 22] quadragesimam: carne & gustu, Phocæ marinæ haud dissimilis. Gregale animal est, & sua habet latitandi tempora, vt Cuculus & Hyrundo. Oua parit in terræ

very great luxury with us, and of great price. It is a mimic. And so, as the Scops and Otus are taken by an imitation of dancing, this bird is caught at night by the light of a candle according to the motion of the captor. For if he stretches out an arm, the bird lifts a wing; if he stretches out a leg, it does likewise. In short, whatever part the fowler plays, the bird does the same. So being intent on the man's actions, it is fooled by the bird-catcher and caught in his net. It is a little bird, of the size of a Starling, with only three fore-toes and no hind-toe, a black crown, white cheeks, and colour almost that of a Quail, if you were to mix with it a little ash-colour, especially round the neck. I call it *Morinellus* for a double reason, both because it is a bird most abundant among the Morini, and because it is a foolish bird, foolishness being by the Greeks called *μωρότης*. On this account our people also call it *Doterell*, as if they were to say doating with folly.

OF THE PUPHIN OR PUPIN.

There is a certain sea-bird of our country, in size and form of body like a little Duck (which the Greeks call *βόσσκας*), with webbed and reddish feet, placed nearer to the hinder parts than in other web-footed kinds except the *Pygosceles*: with a somewhat thin beak, rather more extended in breadth vertically than stretching laterally to a very great length, furrowed by four red grooves above, and two below, pale ochre in colour. The part lying between these and the head is bluish, and of such a shape as is the moon, when ten days have elapsed from conjunction. The bird is black on the upper surface of the whole body, save where the eyes are set, which are enclosed in white: but it is wholly white below, save on the upper breast, where it is black. It gets its living from the sea. This bird our people call the Puphin, we say Pupin from its ordinary cry of "pupin." It hides in holes, as the *Charadrius* does. And so it is driven out from a rabbit's burrow by a ferret turned in by any hunter in a place situated not far from the sea. It is used as fish among us during the solemn fast of Lent: being in substance and taste not unlike a Seal. It is a gregarious animal, and has its proper time for lying hidden, as the Cuckoo and Swallow.

*Coruorans
seu Mergus.*

cuniculis bina magna ex parte. Alis non confidit nisi conspecto mari. Pigrum videtur animal, sed iniurię patiens. Vescitur carne lubentius quàm pisce, & cuniculi quàm alterius animalis, sed vtroque crudo: cocta & assa respuit. Cetera non attingit humana edulia. Aestate se lauat, sed nunquam, quod observatione deprehendi potest, bibit: an quod aqua marina carebat, nescio. Excrementum alui illi est quale accipitri. Cum non esset quod ederet, cibum voce naturali geminata & summissa, pupin, pupin clamitando, implorabat. Alebam domi meae ad menses octo. Mordebat lubenter ministrantes cibum, aut attingentes, sed benignius atque innocentius. Exiguo cibo fatiatur. Non enim vorax auis est vt Coruorans noster, quem tu (Gefnere charissime) coruum aquaticum & mergum recte nominas, nostri Cormorantem corruptè dicunt, nescientes ex vocis ætymo coruum vorantem appellari debere. Quod naturaliter facit, cum illi à natura vnicum tantum intestinum sine elice (vt aiunt) rectum sit concessum, propter vehementiam caloris naturalis, citissimè absorbentis quæ assumuntur omnia.

De Spermologo seu Frugilega.

Spermologus noster à cornice nigra nulla in re differt, nisi morū innocentia, *προλόβω* seu ingluvie, [f. 22 b] qua granum legendo continet, vt ad suos referat: (est enim gutturofus) albo callo, qui in summo rostro est ad caput: & voce, quam habet gutturalem & raucam. Vnde forsan nostris, quibus nomina rerum multa Latina sunt & Græca (vti libro nostro de symphonia vocum Britannicarum diximus) rouce dicitur, quasi rauce Anglis, raucus Latinis sit dicendus: victu quoque à coruo differt, quod frumento, hordeo, & cetero semine (vnde nomen *σπερμολόγου* inuenit apud Græcos) vescitur. Vescitur & vermibus, vbi frugis frumentique copia non est. Hinc rustici nostri dubitant

It lays for the most part two eggs in rabbit burrows in the earth. It does not trust to its wings save in sight of the sea. It seems a lazy animal, but patient of injury. It eats flesh more readily than fish, and that of a rabbit in preference to that of any other animal, but in either case raw: it throws up what is boiled or roasted. Other human victuals it does not touch. In summer it washes itself but never drinks, so far as can be ascertained by observation; whether this was because salt water was wanting, I know not. The droppings are like those of an *Accipiter*. When there was nothing to eat it begged for food with its ordinary cry repeated and lowered, by calling out "pupin, pupin." I kept one at my house for eight months. It bit with right good will those who supplied it with food or touched it, but in a mild and harmless way. It was satisfied with little food. For it is not a voracious bird, as our *Corvorant* is, which you (dearest Gesner) rightly name *Corvus aquaticus* and *Mergus*, while our people corruptly say *Cormorant*, not knowing from the derivation of the word that it ought to be called the Crow that devours. And this it does naturally, since it is endowed by nature with only one intestine straight and without a coil (as they say), on account of the vehemence of the natural heat, which very quickly consumes all that it swallows.

OF THE SPERMOLOGUS OR FRUGILEGA.

Our *Spermologus* differs in nothing from a black Crow, save in the harmlessness of its habits, in the *προλόβος* or crop, which holds the grain as it is picked up, that it may bear it to its young (for it is pouched): in the white callus, which extends from the base of the beak to the head; and in its voice, which is guttural and harsh. Whence possibly by our people, among whom there exist many Latin and Greek names for things (as we have said in our book on the Harmony of British words) it is called *Rouke*, as if it should be called *Rauce* by the English and *Raucus* by the Latins: also in food it differs from the Crow in that it eats wheat, barley, and other seeds (whence it got the name of *σπερμολόγος* among the Greeks. It also eats worms, when there is not plenty of corn or grain. Hence our country people doubt

maior ne sit ex eis vtilitas agris dum legunt vermen, frugis & fementis pestem, an inutilitas hominibus dum vorant granum, hominis nutrimentum. Tanta tamen multitudo est, vt legibus condemnentur: innocens alioqui auis atq; vtilis. Agricolis enim teneri adhuc ex nido, in cibo sunt. Non nisi excelsis arboribus, idq; societate quadam numeroq; nidificant, cohabitant, & confidunt.

De Sacropsittaco.

Psittacorum plura esse genera obseruauit. Quidam enim pusilli, magnitudine videlicet turdi, toto quidem corpore virides sunt, sed caudam longam atque gracilem, & eam aut flauam, viridem, aut puniceam habent. Quidam¹ rursus grandes sunt admodum, cornicis magnitudine, ex toto punicei seu rubri, nisi sub imo ventre, [f. 23] extremis alis, & extima cauda, quibus partibus cum cœruleo virescunt. His rostrum est magnū, cauum, pellucidum, & aduncum, medio tantum sui pallidum, vtroque extremo ex parte nigrum, vt & inferior maxilla tota nigra, cuius cauitatem lingua dura & nigra ceterorum Psittacorum modo & forma occupat. Vtrinque genæ, in cute rugosa, figura ferè triangulari obtusa, candicant, rubris mollium pennarum ordinibus equidistanti parallelo inductis, & alicubi etiam sine ordine. Oculus paruus, & in albo cilio cuticulari constitutus, pupillam habet nigram, quam circundat aureus circulus. Digitos habet quatuor ita efformatos, vt videatur natura voluisse omnes anteriores fecisse, retorxisse tamen duos in auersum firmandi corporis causa. Hos Brasilia mittit, quos propter insignem magnitudinem Sacropsittacos nominamus. Veteres enim quod præclarum magnumque erat, sacrum dicebant. Vt sacrum os, sacram anchoram, sacrum falconem, quem hierofalconem dicunt, sacrum piscem, *ἱερὸν μένος*, sacram famem, & sacrum morbum. Huius generis imaginem quam à nobis accepisti, subiunge. Ceteros inter hos magnitudine medios, aliæ regiones, vt insula Hispania, Aegyptus, & India ferunt, sed colore vario. Alij enim toto corpore cinereo, caudas habent fultas, breuiores & puniceas,

Sacrum.

¹ A misprint for Quidam.

whether their utility is greater in the fields, when they pick up the vermin, destructive to crops and seeds, or their harmfulness to men, when they devour grain, the food of man. However the number of them is so great, that they are condemned by the laws; harmless and useful birds in other respects. For, when still tender from the nest, they are used as food by country men. Only in lofty trees, and that as it were, in company and in numbers do they nest, abide and roost.

OF THE SACROPSITTACUS.

I have observed that there are many kinds of Parrots. For some that are small, namely of the size of a Thrush, have the whole body quite green, but the tail, which is long and slender, either yellow, green, or crimson. Again some are very large, of the size of a Crow, altogether crimson or red, except under the lower belly, on the ends of the wings, and on the tip of the tail, on which parts they shew greenish blue. These have a large beak, hollow, shining and hooked, pale only in its middle, partly black at each end, while the lower jaw is entirely black, the cavity being filled by the tongue, which is hard and black of the style and shape of that of other Parrots. The cheeks on both sides are whitish on the wrinkled skin, in shape almost obtusely triangular, the rows of soft red feathers being set in equidistant parallel lines, though in some places also without being in rows. The eye, small and set in a white ring of skin, has a black pupil, which is surrounded by a golden circle. It has four toes fashioned in such a way that nature seems to have intended to place all of them in front, but to have turned two of them back for the sake of supporting the body. Brasil sends us these birds, which we call Sacropsittaci on account of their remarkable size. For the ancients used to call that Sacrum, which was notable and large. As the *os sacrum*, *anchora sacra*, *falco sacer*—which they call *Hicrofalco*, *piscis sacer*, *ἱερὸν μένος*, *fames sacra*, and *morbus sacer*. Compare with these the picture of this kind, which you have received from us. Other regions, as the Spanish Isle, Egypt and India, produce other kinds which are midway in size, but with various colouring. For some with the whole body grey have strong tails, some-

in extremoq̃ue lunares. Alij ex toto virides sunt. Alij cum corpore vireant, cauda longa flauescunt. [f. 23 b] Mores habent omnes similes, & victum communem. nisi quòd Sacropsittacus, pane bera macerato, carne, & pisce etiam vescatur.

De Coruis albis.

Anno domini. 1548. Augusto mense, coruos duos candidos ex eodem nido vidi & contrectavi istic in Cumbria nostræ Britanniae, apud eiusdem prouinciæ comitem natiuos, atque ita ad aucupium factos vt accipitres. Nam & brachio falconarij quietè insidere, & soluti ad eius vocem atque signum vel è longinquo quàm celerrimè aduolare docti erant. Hos nihil est infaustum cōsequutū, vt albas illas hirundines, de quibus Alexander Myndius apud Ælianū. Nam qui *Vrfus albus.* coruū album notat, colorem notat: vt qui vrfum *Vulpes nigra.* album & vulpem nigram: quorum vtrumque vidimus ex Moscouia istic in Britannia. Quanquam vulpem nigram non queo dicere, etsi vulgus nigram vocet: sed potius fuscā, aut obscurè griseam reliquo corpore. Nam genas atque tibias tantum fuscas habet, clunes atque caudam.

Expletis iam quæ de volatilibus ad te scripsi mi Gefnere, suo ordine pisces consequuntur.

what short and crimson, and crescent-shaped at the tip. Others are entirely green. Others, though they are green on the body, are yellowish on the long tail. All have like habits, and the same kind of food, save that the *Sacropsittacus* eats bread soaked in beer, flesh, and even fish.

OF WHITE RAVENS.

In the year 1548, in the month of August, I saw two white Ravens from the same nest, and handled them at the very place in Cumberland of our Britain, bred on the property of a lord of that county, and trained for bird-catching just like hawks. For they had been taught both to sit quietly on the arm of the falconer, and when loosed to fly as quickly as possible to his call and sign even from a distance. Nothing unlucky followed them, as in the case of those white Swallows, about which Alexander Myndius wrote according to *Ælian*. For he who notes a white Raven notes the colour; as he does who notes a white Bear and a black Fox; both of which I have seen here in Britain from Muscovy. Yet I can hardly call a Fox black, although the common people call it black; but rather dusky or dull grey on the rest of the body. For it has only the cheeks and the legs, with the rump and the tail, dusky.

And now those things being finished which I have written to you of flying creatures, my *Gesner*, the fishes follow in due order.

INDEX.

- Acanthis*, 52, 53
Accentor modularis? xvii
Accipiter, 14-18, 57, 66, 67, 206, 207
A. nisus, xv
A. palumbarius, 14, 15, 18, 19, 56, 57
Accipitres, 14-17, 38, 66, 67, 116-118, 138, 139, 166
Actitis hypoleucos, xvii, 57
Adler, xvi, 30
Ægithus, 74, 104, 162
Æsalo, 14-16
Æsalou, 16, 17
Ἀετός, 30, 31
Africanæ or *Africæ*, see *Gallinæ*
Ἀιδών, 108
Ἀιγυθαλός, 130, 131
Ἀιγιθος, 160, 161
Ἀιγοθήλας, 48, 49
Ἀιτωλός, 116
Ἀισάλων, 16, 17
Ἀιθηα, 110, 111
Ἀκανθίς, 104, 105
Ἀκανθοφάγα, 40
Alauda, xiv, 80, 81, 106, 107, 146, 147
A. arborea, xiv
A. arvensis, xiv
Alaudidae, xiv
Albardeola, 38-40, 180, 181
Albicilla, 28, 30, 31
Alcedinidae, xiv
Alcedo, 18-20, 22, 180, 181
A. ispida, xiv
Ἀλέκτωρ, 82, 83
Ἀλέκτορις, 82, 83
Älke, xv, 92, 93
Ἀλκυών, 18, 19
Amsel, xviii, 114, 115
Anas, xii, xiv, 22, 23, 32, 48, 92, 93, 176, 180-183, 198, 204
A. boscas, xv, 47
A. indica, 196
A. turcica, 198
Anataria, 30-33
Anatidae, xv
Anser, xi, xii, xv, 22-24, 26, 28, 110, 122, 180-183, 194, 196
A. bassanus, 30, 196
A. brendinus, 194
A. minor (the smaller), 180, 181
A. palustris, xi
Anthus pratensis, xvi
Apes, 98, 99
Apodes, 98-103
Apple-sheiler, 73
Aquila, xii, xv, 16, 17, 30-38, 46, 47, 56, 57, 120, 123, 128, 129, 138-141, 152, 153, 162, 163, 192
A. marina, 182
A. montana, 34
Aquila, Sea, 183
Aquila vera, 36
Aquilæ, 30, 31
Ardea, xv, 36-40, 54, 112, 124
A. alba, 38, 39
A. cinerea, xv
A. pella, 38, 39
A. stellaris, x, 38-41, 126, 127
Ardeæ, 38, 39
Ardeideæ, xv
Ardeola, 180, 181
Arlyng, xviii, 52, 53
Ärn, xvi, 30, 31
Ärn, Edel, xv, 36
Ascalaphus, 180, 181
Asio, 130, 131
A. otus, xvii
Ἀσκαλώπας, 86, 87
Astur, 192
A. palumbarius, xv
Asylus, 182, 183
Atricapilla, 44-47, 71, 110, 111, 160, 161, 182, 183
Attagen, xvii, 40-45, 86, 92, 93, 126, 127
Attigena, 40, 41, 44, 45, 87
Atzel, xv, 142, 143
Aurifrisius, 192, 193
Aurimittis, 40, 41, 50, 51, 106-109, 182, 183
Aves Canariæ, 108

- Aves Diomedea*, 70, 71, 78
Aves tarda, 130, 131, 168, 169 (see also *Tarda*)
 Balbushard, xv, 33
Balbushardus, 32
 Bald-Buzzard, 33
Baptea, 40
 Bargander, 25
Βασίλειος, 152, 153
 Bass Goose, 31, 197
 Batis, 158, 159
Βάτις, 158, 159
 Bauncok, 84
 Bergander, xii, xv, 24, 25, 195
Bergandrus, 24
 Bergdöl, xv, 90, 91
 Bernacle, 195
Bernaclus, 194
 Berndclac, 195
Berndclacus, 194
 Berndgander, 195
Bernicla, xi, 26, 30
B. leucopsis, xv
 Bernicle (Goose), x, xv, 27, 31
 Bistard, xvi, 166, 167
Bistarda, xii
 Bittern, 41
 Bittour, xv, 40, 41
 Blackcock, ix, 43
 Blak byrd, xviii, 114, 115
 Blödtfinck, xvi, 160, 161
 Bloudvine, 161
Bloudvinca, 160
Βωμολόχος, 90, 91
Bonasa sylvestris, xvii
Bosca, 48, 49, 182, 183
Βόσκας, 204, 205
Botaurus, xiv
B. stellaris, xv, 41
 Brachvogelchen, xviii, 158, 159
 Bramlyng, xvi, 72, 73
 Brant, xv, 27
Branta, xi, 26
 Brech vogel, xviii, 52, 53
 Brendclac, 195
Brendclacus, 194
 Brend Gose, 194, 195
 Brent Goose, 195
Βύας (*Βρύας*), 46, 47
Bubo, 46, 47
B. ignavus, xvii, 47
 Bächfink, xvi, 72, 73
 Bulfinc, 161
Bulfınca, 160
 Bulfinch, x
 Bulfinche, xvi, 160, 161
 Bunting, xvi, 135, 159
 Bunting, Reed, x
Buntinga, 134, 158
 Burgander, 25
 Burrow Duck, 25
 Bushard, xv
Busharda, 16
 Bustard, xvi, 166
Butco, xii, 14-18, 32
B. vulgaris, xv
 Buttor, xv, 123
Buttora, 122
 Buttour, xv, 40, 41
 Buzzard, 17
 Buzzard, Moor-, ix
 Caddo, xv, 92, 93
Calidris, 180, 181
 Canary Bird, xvi, 109
Capella, 182, 183
Caprimulgide, xv
Caprimulgus, xii, xiii, 48-51
C. europæus, xv
Carduelis, xii, 40, 41, 50-53, 182, 183
C. elegans, xvi
C. spinus, xvi, 51
Carulus, 180, 181
Cataracta, 70, 71, 180, 181
Cephus, 74, 75
Certhia, 52, 53
C. familiaris, xv, 65
Certhiide, xv
 Chaffinche (Chaffinch), xvi, 72, 73
Chalcis, 56, 57
Charadriide, xv
Χαραδρίος, 181
Charadrius, 204, 205
C. pluvialis, xv
Χελιδών, 96, 97
Χήν, 22, 23
Chenalopez, xii, 22-25
Cheneros, 22-28, 31
Χλωρεύς, 106, 107
Χλωρίων, 106, 107, 132, 172, 173
Chloris, xii
Χλώρις, 107-109
 Chogh, xv, 92, 93
 Choghe, 90, 91
 Choghe, Cornish, xv, 90, 91
 Chough, 91
 Chough, Alpine, x, 91
 Chough, Cornish, x
Χρυσομίτρίς, 40, 41
Chrysomitris, 52, 53
Ciconia, x, 54, 55, 62, 63, 92, 93, 180, 181
C. alba, xv
Ciconiide, xv
Cicumæ, 120, 121
Cicumie, 120
Cinclide, xv
Cinclus, xiii, 54-57, 170, 171, 180, 181
C. aquaticus, xv, 23

- Circus*, 14-17
C. aeruginosus, xv, 33
C. cyaneus, xv
 Clake, 194, 195
Clanga, 30, 31
 Clothurd, xviii, 52, 53
Cnipolegus, xiii
 Cob, xvi, 79
Cobbi marini, 78
Coccothraustes, xiii
Ceruleo, 52, 53
 Cok, xvii, 82, 83
 Cok of Kynde, 84
Colius, 88, 89
Collurio, 58, 59
Columba, xv, 14, 58-60
C. anas, xv
C. palumbus, xv
C. vulgaris, 194
Columbæ, 58, 59
Columbi, 14, 18, 42, 44, 58-61
Columbidæ, xv
Columbus, 18, 58-62, 66, 67, 180, 181
Comatibis comata, 93
C. crenata, xvi, 93
 Coot, 77, 199
Copera, xiv, 80, 81
Cormorans, 206
 Cormorant, ix, xvii, 110, 111, 207
 Corn-Crake, 71
 Cornish Chloghe, xv, 90, 91
Cornix, xii, 64, 65, 90, 91, 118, 148, 208
C. aquatica, 22
C. hyberna, 64
Corvidæ, xv
Corvorans, 206
 Corvorant, 207
Corvus, x, 34, 64, 65, 92, 93, 206, (albus) 210
Corvus (web-footed), 183
C. aquaticus, x, 92, 93, 206, 207
C. calvus, 92
C. corax, xv
C. cornix, xv, 65
C. corone, xv, 65
C. frugilegus, xv, 65
C. monedula, xv
C. nocturnus, xii, xiii
C. palmipes, 182
C. sylvaticus, x
C. vorans, 206
Corydos, 82, 83
Cotile riparia, xvi
Coturnix, 62, 63, 68, 104, 105, 126-129, 138, 139, 146, 147, 202, 204
C. communis, xviii
 Coughshot, xv, 60, 61
 Cout, xvii, 33
Couta, 32
 Crane, Corn-, 71
 Crammiesvogel, 172, 173
 Crane, x, 55, 94, 95, 97, 195
 Craspecht, xvii, 149
Craspechtus, 148
 Creeper, 53
 Creeper, Tree-, ix
 Creeper, Wall-, 53
Crepera (Creper), xv, 52
Crex, 68-71, 128, 129
C. pratensis, xvii, 71
 Crow, xv, 64, 65, 119, 149, 207, 209
 Crow (that devours), 207
 Crow-Picus, 149
 Crow, Carrion, 65
 Crow, Hooded or Grey, 65
 Crow, Sea, 65
 Crow, Winter, xv, 68
 Cryel Heron, xv, 38
 Cuckoo, ix, x, 205
Cuculidæ, xv
Cuculus, 48, 49, 66-69, 204
C. canorus, xv
 Cukkow, xv, 66, 67, 69
Cuticilega, xiii, 64, 65, 182, 183
Curuca, 68, 69, 182, 183
Cychramus, 128, 129
Cygnus, xiii, xiv, 126, 127, 196
C. olor, xv
Cymindis, 16, 17, 56, 57
Cypselidæ, xv
Cypselus apus, xv
C. melba, xv
Cypsellus, 100, 101
 Daker Hen, xvii, 70, 71, 128, 129
Dautias lusciniæ, xvii
 Daw, 65, 77
Dendrocopus major? xvii
 Dike Smouler, xvii, 136, 137
 Diomedæ (birds of), xvii, 71, 79
 Dipper, 23
 Distelfinck, xvi, 40, 41
 Distelvine, 41
Distelvinca, 40
 Döl, xv, 92, 93
 Doterel, 205
Doterellus, 204
 Douker, xvii, 176, 177
 Dove, x, xv, 15, 58-61, 63, 97
 Dove, Ringged, xv, 60
 Dove, Venice, xv, 63
 Drossel, xviii, 172, 173
Δρυκολάπτης, 146, 147
 Dächer, xvii, 110, 111, 176, 177
 Duck, xv, 22, 23, 25, 33, 49, 177, 199, 205
 Duck, Indian, 197
 Duck, Turkish, 199
 Duck, Wild, 47

- Durstel, xviii, 172, 173
 Duve, xv, 58, 59
 Duve, Turtel, xv, 60
 Eagle, 31, 33, 37, 39, 47, 193
 Eagle, Sea, 193
 Ebeher, xv, 54, 55
 Egle, xvi, 30
 Egle, Right, xv, 36
 Ἐγῶλιος, 116
 Eissvogel, xiv, 18, 19
 Elster, xv, 142, 143
 Elsterspecht, xvii, 149
Elsterspecht, 148
Emberiza citrinella, xvi
E. miliaria, xvi
E. schaniensis, xvi
 Endt, 22, 23
 Engelchen, xvi, 108, 109
Ephemerus, 28, 29
Ephimerus, 28
 Ἐποπς, 174, 175
Erithacus rubecula, xvii
 Ἐριθακος, 154, 155
 Ἐριθέα, 154, 155
Erna, xii, 30
 Erne, xii, xvi, 31
 Ἐρωδιός, 36, 37
 Eul, xvii, 120, 121
 Eul, Schleier, xvii, 130, 131 (see Ransaul)
Falco aesalon, xv
F. sacer, 208, 209
F. subbuteo, xvi
Falconidae, xv, xvi
Falcula, 98-102
 Fasant, xvii, 140, 141
 Fasian, xvii, 140, 141
Fedoa, xvii, 44, 45
 Feldefare, Feldfare, xviii, 58, 59
 Fenclake, 194, 195
 Fenlagge, 194, 195
 Fenclake, 194, 195
Ficedula, 44, 45, 70, 71, 182, 183
 Fieldfare, ix
 Finch, 19, 119
 Flasfinc, xvi, 159
Flasfinka, 158
Florus, 74, 75, 104, 105, 160, 161, 182, 183
 Fowl, 201
 Fox Goose, 25
Fringilla, 18, 52, 53, 72-75, 118, 130, 131, 148, 149, 182, 183
F. caelebs, xvi
F. montifringilla, xvi
Fringillago, 130, 131
Fringillarius, 14, 15, 18, 19
Fringillide, xvi
Frugilga, 64, 65, 182, 183, 206, 207
Fulica, 70, 71, 74-79, 180, 181, 198
F. atra, 77
F. nigra, xvii
Fulvia, 34, 35
Galerita, xiii, 18, 56, 57, 80-83
G. cristata, xiv
Galgulus (Galgulus), 88-91, 146, 147, 182, 183
 Galli, 82, 83
Gallina, 86, 87, 138, 140, 148
G. cohortalis, 84
G. getula, 200
G. nigra, 32
G. nigra aquatica, 76
G. numidica, 84, 85
G. palustris, 76
G. rustica, 82, 84, 86
Gallinaceus, 180, 181
Galline, 18, 82-87
G. africana (africa), 82, 84, 86, 87
G. Hadrianae, 84-87
G. villatica, 82, 84, 86
Gallinago, 40, 41, 86-88, 146, 147
Gallinula chloropus, xvii
Gallus, 82, 83
G. domesticus, xvii, 42
G. ferrugineus, xvii
G. gallinaceus, 200
G. medicus, 84
 Γαμψώνυχες, 40
 Ganss, xv, 22, 23
 Ganss, Löffel, xvii, 150, 151
 Ganss, Trap, xvi, 166, 167
Garrulus glandarius, xv
Gavia, 78, 110, 111, 180, 181
G. alba (White Gavia), 78, 180, 181
G. cinerea (Grey Gavia), 78, 180, 181
Gavia marinae, 78
Gavinus viridis, xvii
 Geelgorst, xvi, 106, 107
 Geir, xviii, 176, 177
 Γερανός, 94, 95
 Gersthammer, xvi, 135
Gersthammera, 134
 Geyr, xviii, 176, 177
 Geyr Swalbe, xv, 102, 103
 Gliandaja, 145
 Γλαύξ, 120, 121
 Glede, xvi, 116, 117
 Γλώττις, 104, 105
 Godwit, ix
 Godwitt, xvii, 45
Godwitta, 44
 Goldfinc, 41
Goldfinka, 40
 Goldfinche, xvi, 40, 41
 Gold hendlin, xviii, 168, 169
 Goose, 23, 25, 27, 29, 111, 123, 139, 195, 197

- Goose, Bass, 31, 197
 Goose, Bernacle (or Bernicle), x, 27, 31, 195
 Goose, Brant (or Brent), xv, 27, 195
 Goose, Fox, 25
 Gose, xv, 22
 Goshawk, 19
 Gosling, 177
 Gouke, xv, 66, 67
Graculi, 90, 91, 94, 95
Graculus, x, 90, 91, 94, 95
 Graesmusch, xviii, 110
 Grasmuklen, xviii, 44, 45
 Grasmuschen, 47
Grasmuschus, 46, 72
 Grasmusch (Grasmusch), xvii, xviii, 73, 136, 137
 Grass-sparrow, 111
 Greenfinch, 41
 Grenefinc, 107
Grenefinca, 106
 Grenefinche, xvi, 104, 105
 Grey-clak, 195
 Grey Hen, ix
 Grey-Lag, 195
 Grouse, Hazel, xvii
Gruide, xvi
 Grünling, xii
 Grunspecht, xvii, 89, 113, 115, 149
Grunspechta, 148
Grunspechtus, 88, 112, 114
Grus, 54, 62, 63, 94-96, 180, 181, 195
G. communis, xvi
 Gryphen, 178
Gryps, 178, 179
 Guinea Fowl, 85
 Gull, 79
 Gull, Black-headed, ix, 77
 Gull, Grey, xvi
 Gull, White, xvi
Γυπαίετος, 34
Γύψ, 176, 177
 Guse, Solend, xvii, 28
Gypaëtus barbatus, xvi, 129
Hadriana, see *Gallina*
Hematopus, 102, 103
Halcyon, 20, 21
Halicetus, 36, 37, 192, 193
Halicetus, 34, 35
H. albicilla, xvi
 Hän, xvii, 82, 83
 Harrier, see Hen-Harrier and Marsh-Harrier
 Hawk, 18, 19, 39, 68, 119, 167
 Hedge-Sparrow, xvii, 136, 137
 Hen, xvii, 82, 83
 Hen, Getulian, 201
 Hen-Harrier, ix, 19
 Hen-Harroer, xv, 18
 Hen, Marsh, 77
 Hen, Mot, xvii, 170, 171
 Hen, Water, xvii, 77, 170, 171
 Heron, xv, 36, 37, 41, 55, 113, 125
 Heron, Blue, 38
 Heron, Cryel, xv, 38
 Heron, Dwarf, xv, 38
 Heron, Night, xiii
 Hewhole, xvi, 115, 149
Hierofalco, 208, 209
Himantopus, 102
Hinnularia, 30, 31
Hirundinide, xvi
Hirundo, 96-102, 116, 117, 170, 173
H. agrestis, 98
H. (alba), 210
H. aquatica, xiii
H. domestica, 100
H. riparia, 98-102
H. rufula, 99
H. rustica, xvi, 98, 100, 101
Hobbia, 18
 Hobby, ix, xvi, 19
 Holtzsneppf, xvii, 86, 87
 Holtztaube, xv, 60, 61
 Hön, xvii, 82, 83
 Horn Owl (Owl), xvii, 130, 131
 Houp, xviii, 174, 175
 Howlet, xvii, 120, 121
 Howpe, xviii, 174, 175
 Huhol, xvii, 89, 115
Huhola, 114, 148
Huholus, 88
Hydrochelidon nigra, xvi
Hyrundo, 204
Iaiä, 144
Ibidide, xvi
 Ibis, Glossy, 57
 Ibis, Red-cheeked, xvi
Icteros, 88, 89
Ἰκτινος, 116, 117
Iliacus, 170, 171
lynx, 146-149
I. torquilla, xvii
 Jay, xv, 145
Junco, 102, 103, 170, 171, 180, 181
 Ka, xv, 92, 93
 Kastrel, xvi, 166, 167
 Kautz, xvii, 46, 47
Κέγχις, 166, 167
Κελεός, 88, 89, 146
Κέπφος, 74-77
 Kersenrie, xvi, 172, 173
Κίχλα, 170, 171
Κίγκλος, 54, 55
 Kingfisher, xiv, 21, 23

- Kirsfincke, xvi, 104, 105
 Kistrel, xvi, 166, 167
 Kite, x, 117, 193, 197
 Kite (Blacker), 117
 Κίττα, 142, 143
 Κνιπολόγος, 64, 65
 Koelmussh (Koelmusch), xvii, 136, 137
Koelmusshus, 136
 Κόκκυξ, 66, 67
 Kok of Inde, xvii, 86
 Κολίος, 88, 89, 146, 147
 Κολλυρίων, 58, 59
 Kölmeyse, xvii, 130, 131
 Κολοίος, 90, 91, 146
 Κολυμβρίς (Κολυμβίς), 176, 177
 Κορακίας, 90, 91
 Κόραξ, 64, 65
 Κορώνη, 64, 65
 Κορυδαλός, 80, 81
 Κόρυδος, 80-83
 Κοπτιφός (Κοπτιφός), 114, 115
 Krae, xv, 64, 65
 Kraeg, xv, 64, 65
 Krammesvögel, xviii, 58, 59, 170, 171
 Krän, xvi, 94, 95
 Kränich, xvi, 94, 95
 Κρανός, 52, 53, 181
 Kukkuck, xv, 66, 67
 Κύκνος, 82, 83, 120, 121
 Kuningsgen, xviii, 152, 153
 Kyngesfissler, xiv, 18, 19
 Kyte, xvi, 116, 117
 Kywit, xv, 77
Kyvitla, 76

Læves, 14, 15
Lagopes (*Lagopus*), 104, 105
Λαγωφόνος, 34
Lagopus mutus, xviii
L. scoticus, xviii
 Lämmergeier, 36, 129
Laniide, xvi
Lanius excubitor, xvi
L. minor?, xvi
 Lapwing, x, xv, 175
Lapwinga, 76, 174
Laride, xvi
 Lark, 19
Λάρος, 74, 75, 78, 79
Larus, xvi, 74, 75, 78, 79
L. ridibundus, xvi, 77
 Laverock, xiv, 80, 81
 Lefler, xvii, 150, 151, 153
Leflera, 152
Leporaria, 34, 35
 Lerc, 81
 Lerc, Wilde, 81
Lerca, 80
 Lerch, xiv, 80, 81
 Lerch, Heid, xiv, 80, 81

 Lerck, Wod-, xiv, 80
 Lerk, xiv, 80, 81
 Lerk, Heth, xiv, 80
 Lerk, Wilde, xiv, 80
Λευκερωδός, 38, 39
Libicus, 164
Λιβικός, 164, 165
 Libyan Bird, 165
Ligurinus, 104, 105
L. chloris, xvi
Limosa belgica, xvii
Linaria, 158, 159
 Lingett, xviii, 47, 111
Lingetta, 46, 110
Lingulaca, 104, 105, 128, 129
 Limot, xvi, 51, 159
Linota, 50, 158
L. cannabina, xvi
Livia, 58-61
 Löffel Ganss, xvii, 150, 151
 Loun, xvii, 177
Louna, 176
Λύκος, 90, 91
 Lüningk, xvi, 132, 133
Lupus, 90, 91, 94, 95
Luscinia, 44, 45, 68, 69, 108-111
Lutea (*Luteus*), 106-108
Luteola, 108, 109, 182, 183
 Lyke Foule, xvii, 46, 47
 Lyssklicker, xiii

Μαλακοκρανέυς, 116, 117, 132
Mareca penelope, xv, 22
 Marsh-Harrier, 33
 Marsh-Hen, 77
 Martinette (or Martnette), Chirche, xv, 102, 103
 Martinette, Rok, xv, 102, 103
 Martnet, Bank, xvi, 102, 103
Matrix, 128, 129 (cp. 126)
 Mavis, xviii, 172, 173
Medicæ, 84, 85
 Meelmeyse, xvii, 130, 131
Melanaëtos, 34, 35
Μελαγκόρυφος, 44, 45
Μελαναέτος, 34
Melagris, 82-87, 140, 141, 200, 201
Melice, 84, 85
Mercolphus, xv, 144, 145
Mergus, xiv, 76, 77, 110, 111, 126, 127, 176, 177, 180, 181, 206, 207
 Merl, xviii, 114, 115
 Merlin, xv, 17
Merlina, 16
Meropide, xvi
Merops, 112-115
M. apiaster, xvi
Merula, 48, 49, 52, 53, 82, 83, 114, 115, 146-149, 164, 165, 170, 171
M. aquatica, xiii

- Mewe, Wyss, xvi, 74, 75
 Meyse, 130, 131
 Meyspecht, xvii, 162, 163
Meyspechtus, 162
Miliaria, 50, 51, 158, 159
Milvius (*Milvus*), 16, 17, 116, 117, 192, 196
Milvus ictinus, xvi
M. ater (*nigrior*), xvi, 116
 Mistletoe-Thrush, xviii
Molliceps, 58, 59, 116, 117, 120, 121, 132, 133, 168, 169
Monedula, 64, 76, 90-93, 146-149
Monetula, 90, 91
Monticola, 130, 131
Montifringilla, 72, 73, 182, 183
 Moor-Buzzard, ix, 33
 Moorhen, 171
 Morhen, xviii, 87, 171
Morhenna, 86
Morinellus, 202-205
Morphna (*Morphnos*), 30, 31
 Mortetter, xviii, 159
Mortettera, 158
Motacilla, 64, 65
M. alba, xiii, xvi
M. lugubris, xvi
Motacillidae, xvi
 Mot Hen, xvii, 170, 171
 Müsche, xvi, 132, 133
 Myre Dromble, xv, 38

 Nachtgäl, xvii, 108, 109
 Nachtra, xii, xiii
 Nachtrab, xiii
 Nachtrap, xii
 Naghtrauen, xii, xiii
Νεβροφόνος, 30
Νήττα, 22, 23
Νηττοφόνος, 30
 Neun mürder, xvi, 168, 169
 Neunmürder, Grosse, 168, 169
 Night-Heron, xiii
Noctua, 46, 47, 120, 121, 130, 131, 180, 181
Nonna, 132
 Noubrecher, xv, 94
Nucifraga, 94, 95
N. caryocatactes, xv
Nucifeta, 162
 Nuin mürder, xvi, 116, 117
Νυκτικόραξ, 120, 130
Numida melagris, xvii
Numidica, see *Gallina*
 Nun (Non), xvi, 133
 Nushäkker, xvii, 162, 163
 Nutcracker, x
 Nutjobber, xvii, 162, 163
 Nutseeker, 163
Nycticorax, xii, xiii

 Nyghtyngall, xvii, 108, 109
 Nyn murder, xvi, 116, 117, 168, 169
Nyroca firina, xv

Οἰβάς, 62, 63
Oistris, xvii, 164, 165
Olor, 120-123, 126, 180-183
Onocrotalus (*Onocratalus*), x, xiii, xiv, 122-127
 Orfraie, 193
Oriolide, xvi
Oriolus galbula, xvi
Ὄροσπίτης, 72, 73
Ὄρνυξ, 62, 63
Ortygometra, 126-129
 Osel, Black, xviii, 114, 115
 Osprey, ix, xvi, 34, 35, 37, 193-195
Ossifraga, 36, 118, 128, 129, 193
O. barbata, 128
 Ossifrage, 37, 129
 Ossifrage, Bearded, 129
Otididae, xvi
Otis, 130, 131, 168, 169
O. tarda, xvi
Ὠτὸς, 120, 130, 131
Otus, 128-131, 204, 205
Ὀυπαξ, 166, 167
 Ousel, Water, 23
 Owl (Oul), xvii, 120, 121
 Owl, Eagle, 47
 Owl, Hawk, 157
 Owl, Horn, xvii, 130, 131
 Owl, Horned, 120
 Oxei, Great, xvii, 130
 Oxeeye, Great, 131

 Pagelün, xvii, 136, 137
Palmipedes, 120
Palumbarius, 14, 15, 18, 19, 56, 57
Palumbes, 58-61, 68, 69, 180, 181
Palumbus, 18, 60, 61
Pandion haliaëtus, xvi
Pandionide, xvi
 Papegay, xvii, 150, 151
Paphus, 50, 51
Πάρδαλος, 58
Pardalus, 132, 133
Parida, xvi
 Parrot, xvii, 209
 Partridge, 43, 63, 141, 203
Parus, 130-133, 182, 183
P. ater, xvii
P. caeruleus, xvi
P. major, xvii
P. maximus, 132, 162
P. medius, 130
P. minimus, 132
P. palustris, xvii
Passer (*Paseer*), 18, 72, 104, 106, 108, 137-139, 160, 168, 180-183

- Passer communis*, 134
P. domesticus, xvi
P. gramineus, 110, 111
P. harundinaceus, 104, 105
P. magnus, 134
P. sepiarius, 136, 137
P. torquatus, 134, 135
P. troglodytes (troglodytes), 134-137
 Passer, Common, 135
 Passer, Great, 135
Passerculus, 134
 Παυο, 136-139
P. cristatus, xvii
P. indicus, 86
 Peacock, 137, 139
 Peacock, Indian, 87
 Pecok, xvii, 136, 137
 Πελεῖας, 60, 61
Pelecanus (Pelicanus), 150-153
 Πελεκάν, 150, 151
 Pella, 38, 39
Penelopes, 22-25
Penelops, 22, 182, 183
Pernæ, 14, 15
Pernopterus, 34, 35
Pernos, 30-33
Perdix, xiv, 18, 42-44, 138-141, 180, 181, 202
P. cinerea, xvii
P. rustica (rusticula), xii, 140, 141
P. vulgaris, xiv
 Πέρδιξ, 138, 139
 Περιστέρα, 58, 59
 Περιστερειδῆ, 40
 Πέρνης, 14
Pernix, 14, 15
 Petrige, xvii, 138, 139
 Pffaw, xvii, 136, 137
Phalacrocoracidae, xvii
Phalacrocorax, x, 92-95
P. carbo, xvii
P. graculus? xvii
Phalaris, 92, 93, 182, 183
Phasianidae, xvii
Phasianus, 40, 140, 141, 200
P. colchicus, xvii
 Φάττα, 60, 61
 Pheasant, 41, 201
 Φήνη, 36
 Phesan, xvii, 140, 141
Philomela, 108, 109
Phanicura, 157
Phoenicurus, 154-157
Phœnix, xvii, 140, 141
 Φοινικουρός (φοινικούργος), 154, 155
 Φοῖνιξ, 38-41
 Φῶξ, 38
Pica, 118, 142-145, 170, 171
P. granata, 144
P. marina, 198
Pica rustica, xv
P. vulgaris, 144
Picidae, xvii
Picus, 54, 114, 115, 146-149, 162, 163
P. cornicinus, 148
Picus, Crow, 149
Picus, Green, 115
Picus martius, xiii, xvii, 88, 89, 146, 147, 182, 183
P. maximus, 148
P. medius, 148
P. minimus, 148
P. viridis, 112
 Pie, 119, 145, 199
 Pie, Common, 145
 Pie, Sea, 199
 Pie, Seed, 145
Pigargus (Pygargus), xii, 30, 31
 Pigeon, 43, 45, 195
Piger, 38, 39
Pilaris (Turdus), 170, 171
 Piot, xv, 142, 143
 Pipers (of Crane or Pigeon), 96, 97
Pipiones, 96, 97
 Pipit, Tree-, 69
Pipilo, 146, 147
 Pittour, xv, 40, 41, 123
Pittourus, 40, 122
 Πλάγγος, 30
Plaucus, 30, 31
Planga, 30, 31
 Πλάνος, 30
Platalea leucorodia, xvii, 41
Plataleidae, xvii
Plataea, 150, 151, 180, 181
Platlea, 150, 151
Plegadis falcinellus, 57
 Plover, Stilt, 102
 Pluver, xv, 132, 133
 Pochard, xv, 49
Pochardæ, 48
Poicipedidae, xvii
Podiceps minor, xvii
 Popinjay, xvii, 150, 151
Porphyrio, 102, 103, 152, 153
P. caelestis, xvii
Pratincola rubicola, xviii
 Πρῶσπος, 152, 153
 "Priest," 51
Procellariidae, xvii
 Ψάρος, 164, 165
Psitaca, 150
Psitacæ, 150, 151
Psitacus (Psittacus), 82, 83, 142, 143, 150, 151
Psittacidae, xvii
Psittacus (Sacropsittacus), 208-211
 Πτέρνης, 14
 Πτερίς, 14
Ptynx, 56, 57

- Puffinus*, xvii
Πύγαργος, 170; see *Pigargus*
Pulla, 34, 35
Pulver, xv, 132, 133
Puphin, 205
Puphinus, 204
Pupin, 197, 205
Pupinus, 196, 204
Πυρρόυλας, 160, 161
Πυρρουνάς, 161
Puttok, xvi, 116, 117
Py, xv, 142, 143
Pygargus, see *Pigargus*
Pygosceles, 204, 205
Pyrrhocorax, 90-93
P. alpinus, xv
P. graculus, xv
Pyrrhula europæa, xvi

Quail, 63, 69, 203, 205
Quale, xviii, 62, 63
Querquedula crecca, xv
Quikstertz, xvi, 64, 65

Rabe, xv, 64, 65
Rail, Water, 71
Rala, xii, xiii, 140
R. aquatica, xii
R. aquatilis, xiii
R. montana, xiii
R. sylvestris, xiii
R. terrestris, xii
Rale, xvii, 141
Rallidæ, xvii
Rallus aquaticus, 71
Ranseul, xvii, 130, 131
Raphön, xvii, 138, 139
Rauce, 206, 207
Raucus, 206, 207
Raven, xv, 35, 64, 65
Raven, White, 211
Rayn byrde, xvii, 146
Redbreast, ix, 161
Redbreste, Robin, xvii, 154, 155
Rede Sparrow, xvi, 102, 103
Rede tale, xviii, 154, 155
Redshanc, xvii
Redshanca, 102
Redshank, 103
Redstart, ix, 155
Reed-Bunting, x
Reed-Sparrow, 105
Regulus, 52, 53, 118, 119, 134-137, 152-155, 168, 169, 182, 183
R. cristatus? xviii
Reydt Müss, xvi, 102, 103
Reyger, xv, 36, 37
Ringel Taube, xv, 60, 61
Ringged Dove, xv, 60, 61
Ringtail, 19

Ringtale, xv
Ringtalus, xii, 18
Ringtayle, xii
Riparia, 98-102
Robin Redbreste, xvii, 154, 155
Rook, 65
Rosdom, xv, 40, 41, 123
Rosdomma, 122
Rosdommus, 40
Rötbrust, xvii, 154, 155
Rötkelchen, xvii, 154, 155
Röstertz, xviii, 154, 155
Rouce (Rouke), 206, 207
Rowert, xvi, 72, 73
Rubecula, 154-159, 182, 183
Rubetarius, 14, 15, 18, 19
Rubetra, 158, 159, 182, 183
Rubicilla, 156, 157, 160, 161, 182, 183
Rupex, 180, 181
Ῥυσσητρης, 40, 41
Rusticula, xii
Ruticilla, 154-159
R. phænicurus, xviii

Sacropsittacus, 208-211
Salus, 160, 161
Sandpiper, ix
Sandpiper, Common, 57
Saxicola oenanthe, xviii
Σχιζόποδα, 40
Σχοίνικλος, 102, 103
Schrecke, 71
Schric, xvi, 169
Schrieus, 168
Schriek, 71
Schryk, xvii, 70, 71
Schuffauss, xvii, 46, 47
Schüffel, xvii, 46, 47
Schwallb, xvi, 96, 97, 102, 103
Scolopacidae, xvii
Scolopax rusticula, xvii
Scops, 204, 205
Scrica, xvii, 128, 129
Sea-Aquila, 183
Sea-Cob, 79
Sea-Eagle, 193
Sea-Pie, 199
Se Cob, xvi, 78-79
See Gell, xvi, 78-79
Σεισπονήγης, 54, 55
Semaw (white with a black cop), xvi, 74, 75
Senator, 152, 153
Serinus canarius, xvi
Shearwater, 71
Sheldappel, xvi, 72, 73
Sheld-Drake, 25
Shell-apple, 73
Shovelard, xvii, 41, 150, 151
Shovelarda, 38

- Shric, 169
Shricus, 168
 Shrike, ix, xvi, 116, 117
 Siskin, xvi, 51, 108, 109
Sitta, 162, 163
S. caesia, xvii
Sittidae, xvii
Σκνιποφάγα, 40
Σκολόπαξ, 87
Σκωληκοφάγα, 40
 Smatche, xviii, 52, 53
 Smerl, xv, 17
Smerla, 16
 Snepff, Holtz, xvii, 42, 86
Sneppa, 70, 71
 Solend Guse, xvii, 28
 Sparhauc, xv
Sparhauc, 18
 Sparrow, xvi, 73, 105, 107, 109, 111, 132, 133, 135, 161, 169
 Sparrow, Grass, 111
 Sparrow-Hawk, 19
 Sparrow, Hedge, xvii, 136, 137
 Sparrow, Rede, xvi, 102, 103
 Sparrow, Reed, 105
 Spatz, xvi, 132, 133
 Specht, xvii, 146, 147, 149
Spechta, 148
 Speiren, xvi, 102, 103
 Sperlingk, xvi, 132, 133
Σπερμολόγος, 64, 65, 206, 207
Spermologus, 206, 207
 Sperwer, xv, 18, 19
 Spink, xvi, 72, 73
Spinus, 40, 41, 64, 65, 104-107, 182, 183
Σπίτσα, 72, 73
 Spoonbill, x, 41
 Står, xvii, 164, 165
 Starling, 23, 95, 165, 205
Στεγανόποδα, 40
 Steinbeisser, xiii
 Steinchek, xviii, 52, 53
 Steingall, xvi, 166, 167
Stellaris, 14, 15, (*Ardea*) x, 38-41, 126, 127
 Sterlyng, xvii, 164, 165
 Stern, xvi, 79
Sterna, 78
S. nigra, 79
 Steynbisser, xvii, 54, 55
 Stigelitz, xvi, 40, 41
 Stilt-Plover, 102
 Stodove, xv, 60, 61
 Stonchatter, xviii, 159
Stonchattera, 158
 Stör, xvii, 164, 165
 Storck, xv, 54, 55
 Stork, xv, 54, 55
 Strauss, xvii, 164, 165
Strigidae, xvii
Strix stridula? xvii
Στρούθος, 132, 133
Στρουθός, 164, 165
Struthio, 44, 45, 164, 165
Struthio camelus, xvii, 92, 93, 164, 165, 168, 169
Struthionidae, xvii
Sturnidae, xvii
Sturnus, 22, 94, 164, 165, 168, 169, 204
S. vulgaris, xvii
Subaquila, 34, 35
Subulco (*Subbulco*), xii, 14, 15, 18, 19
Συκαλός, 70, 71
Sula bassana, xvii
Sulide, xvii
 Swalbe, Geyr, xv, 102, 103
 Swalbe, Kirch, xv, 102, 103
 Swalbe, Über, xvi, 102, 103
 Swale, xvi, 96, 97
 Swallow (*Swallowe*), ix, xvi, 96, 97, 101-103, 205
 Swallow, Black, 101
 Swallow, Great, xv, 102, 103
 Swallow, House, 101
 Swallow, Water, xiii, xvii, 54, 55
 Swan, xv, 120, 121, 123, 127, 197
 Swän, xv, 120, 121
Sylvia, 182, 183
S. atricapilla, xviii
S. rufa, xviii
Sylviidae, xvii, xviii
Tadorna cornuta, xv, 25
Ταών, 136, 137
Tarda, xii, 180, 181, (see also *Ardea tarda*)
 Taube, xv, 58, 59
 Taube, Ringel, xv, 60
Tela, 48
 Tele, xv, 49
 Tern, Black, ix, 79
Tetrao, 30, 31, 130, 131, 166, 167
T. tetrix, xviii
Tetraonidae, xviii
Tetrix, 30, 31, 166, 167
Τέτριξ, 166, 167
Thraupis, xii, 52, 53
Θραυπίς, xii, 50, 51
 Throssel, xviii, 172, 173
 Thrusche, xviii, 170, 171
 Thrush, 59, 119, 175, 177, 209
 Thrush, Blue, 53
 Thrush, Mistletoe, xviii
 Thrushe, xviii, 172, 173
Tichodroma muraria, 53
Tinnunculus (*Tinnunculus*), 166, 167
Tinnunculus alaudarius, xvi
 Titlark, 69

- Titling, 69
Titlinga, 68
 Titlyng, xvi, 68, 69
 Titmouse, 130, 131
 Titmouse[e], Great, xvii, 130, 131
 Titmouse, Greatest, 133
 Titmouse[e], Less, xvii, 130, 131
Torquella (Torquilla), 146-149
Totanus calidris, xvii
 Trapp (Träp), xvi, 31, 166, 167
Trappus, xii, 30
 Tree-Creeper, ix
 Tree-Pipit, 69
Triorchia, 14, 15
Triorches, 16, 17
 Τριόρχης, 14, 16, 17
 Τροχίλος, 152, 153
Trochilus, 152, 153
Troglodytes (Troglodites), 134-137
T. parvulus, xviii, 155
Troglodytidae, xviii
 Τρύγας, 170, 171
 Τρυών, 60
Trynga, 170, 171, 180, 181
Turbo, 146-149
Turdida, xviii
Turdus, 58, 102, 105, 118, 170-174, 208
T. iliacus, xviii, 170, 171
T. merula, xviii
T. musicus, xviii
T. pilaris, xviii, 170, 171
T. viscivorus, xviii, 170, 171
 Turtel Dove, xv, 60, 61
 Turtle-Dove, 89
Turtur, 58-61, 88, 89, 146, 147, 172-175, 180, 181
T. communis, xv
Tyrannus, 120, 121, 154, 155, 168, 169
 Üle, xvii, 120, 121
Ulula, 130, 131
 Ὑπαιερός, 34
Upupa, 174, 175
U. epops, xviii
Upupida, xviii
Urxax, 166, 167
Urinatrix, 92, 93, 176, 177, 182, 183

Valeria, 34, 35
Vannellus, 76, 77, 174, 175
V. vulgaris, xv
 Velt hön, xvii, 138, 139
 Venice Dove, xv, 63

Villaticæ, 82, 84, 86
Vinago, 58-63, 180, 181
Vipiones, 96, 97
Vireo, 58, 59, 106, 107, 148, 149, 172-175
Viscivorus (Turdus), 170, 171
 Vishärn, xvi, 34, 35
 Vogelhain, xiii
Vulpanser, xii, 24, 25, 182, 183
Vultur, xviii, 34, 166, 176-178
V. niger, 178
 Vulture, 35, 177, 179
 Vulture, Black, 179
Vulturide, xviii

 Wachholtervögel, xviii, 170, 171
 Wachtel, xviii, 62, 63
 Wagtale, xviii, 64, 65
 Waldrapp, x
 Wall-Creeper, 53
 Waltrap, xvi
 Walt-rapp, x
Waltrapus, 92-95
 Wasser Hen, xvii, 170, 171
 Wasser Steltz, xvi, 64, 65
 Water-Craw, xv, 22, 23
 Water-Hen, xvii, 170, 171
 Water-Hen (black with a white frontal patch), 77
 Water Ousel, ix, 23
 Water Rail, 71
 Water Swallow, xiii, xvii, 54, 55
 Weidwail, xvi, 172, 173
 Weingaerdsvogel, xviii, 172, 173
 Weye, xvi, 116, 117
 Wheatear, ix, 53
 Widhopff, xviii, 174, 175
 Wigene, xv, 48, 49
 Wigeon, 22
 Witwol, xvi, 172, 173
Wodcoccus, 42
 Wodcok, xvii, 86, 87
 Wodlerck, xiv, 80
Wodspechta, 148
 Woodcock, 42, 45, 89
 Woodpecker, 55
 Woodspecht (Wodspecht), xvii, 149
 Wren, xviii, 152, 153, 155
 Wyngthrushe, xviii, 172, 173

 Yelowham, xvi, 106, 107
 Yowlring, xvi, 106, 107

 Zaunküningk, xviii, 152, 153
 Zeysich, xvi, 108, 109

AMNH LIBRARY

100099726