**BACTERIAL
PYROGENS**

**PARTICULARLY
PYROGENIC
POLYSACCHARIDES
of
BACTERIAL
ORIGIN**

**An
Annotated
Bibliography**

**BAXTER
LABORATORIES,
INC.**

**Morton Grove,
Illinois**

589.75
G 43

B A C T E R I A L

P Y R O G E N S

(Particularly Pyrogenic Polysaccharides

of

Bacterial Origin)

An Annotated Bibliography

Edited

By

Leonard G. Ginger, Ph. D.
Assistant Scientific Director

William F. Windle, Ph.D., Sc. D.
Scientific Director

and

Ione E. Johnson, A. B.
Librarian

BAXTER LABORATORIES, INC.
Morton Grove - Illinois

MAY, 1952

67721

P R E F A C E

Although there has been a general awareness for many years that the inadvertent occurrence of fever following the parenteral administration of fluids is related to some product of microbial metabolism, it is only within the past several decades that studies have led to the isolation from microbial sources of pyrogenic materials of a polysaccharide nature possessing remarkable physiologic activity. It is now appreciated that the vague phenomena, which formerly were called "non-specific, foreign protein reactions," have a common etiology. Trace contamination of a protein is usually sufficient to result in the formation of enough of the pyrogenic polysaccharide complex to impart the observed type of activity. As a consequence it is likely that protein is not the primary causative agent.

The recent availability of pyrogenic polysaccharides has stimulated basic science investigations, leading to the realization that these extremely active substances are capable of eliciting a miscellany of complex physiological changes, even in sub-febrile doses, which appear to be of therapeutic importance.

This annotated bibliography was compiled to provide a selected list of references, particularly relating to pyrogenic polysaccharides of bacterial origin, to acquaint the worker in this field with the published investigations and to aid in orienting and focusing current researches. Out of necessity, the bibliography also includes many pertinent references on vaccines. It must be emphasized, however, that a very significant number of references relating to non-specific, foreign protein therapy have been deliberately omitted because of the conviction that it is practically impossible to duplicate many of such materials; hence such information is not of practical significance. Some references in this latter category have been included in selected instances where deemed desirable.

The period covered by this bibliography extends from 1785 through March, 1952.* It is grossly arranged in numbered chronological order but is alphabetical within each year. To facilitate usage, an author index and an extensive subject index are provided.

A great number of the references in very recent years is concerned with investigations involving the use of a purified pyrogenic substance called PYROMEN. The name of this preparation has been changed to PIROMEN. Particular attention is directed to this alteration in name to avoid the possible confusion that might otherwise arise with the appearance of both names in the published literature.

It is a particular pleasure to acknowledge the efforts of Miss Ella Salmonsens, Medical Librarian, John Crerar Library, and Miss Jean Reilley, Medical Reference Librarian, American Medical Association, in providing necessary reference materials and much valuable advice. Mrs. Maud Hinson prepared the extensive index, and several members of our own staff, in particular, Mrs. Helen Heartstedt, Mrs. Mae Swanson, and Mrs. Joyce Eckhart, contributed significantly in the compilation and recording of the bibliographic material.

The Editors

May 1, 1952

* Extended in The Addendum to August, 1952

ERRATA

<u>Abstract</u>	<u>Line</u>	<u>Should read</u>
18	3	"Riforma"
	8	"Metchnikoff"
39	10	"vacuolation"
74	6	"contractions"
79	4	"mechanism"
132	7	"appears"
137	14	"leukocytes"
163	2	"lueischen"
203	12	"markedly by"
207	2	"glutathione"
226	13	"gram-positive organisms"
230	2	"praecox"
254	5	"antichancroid"
340	11	"criterion"
354	3	"cotton"
404	5	"testing"
421	5	"H ₂ O ₂ "
430	10	"diastases"
491	2	"Handbook"
506	3	"sclerosis"
509	6	"Seitz"
511	6	"pyocyaneus"
561	4	"reviews"
617	14	"nucleoprotein"
619	11	"Electric"
	12	"decrease"
629	3	"aspecific"
631	10	"resumes"
661	32	"release"
674	6	"synthetic"
685	16	"volume"
712	4	"leukocyte"
728	2	"leukocyte"
736	1	"Chambers"
752	11	"lymphocytes"

- 1 7 8 5 -

1. DICKINSON, C.

An Inquiry into the Nature and Causes of Fever (with a review of the several opinions concerning its proximate cause as advanced by different authors and particularly as delivered from the practical chair in The University of Edinburgh, etc.)

Edinburgh, 1785

The Edinburgh theory on the etiology of fever is stated to be that a product of tissue destruction ("putrefaction") produces fever.

- 1 8 2 6 -

2. BLACK, J.

A comparative View of the More Intimate Nature of Fever

London, 1826

- 1 8 3 7 -

3. CLUTTERBUCK, H.

An Essay on Pyrexia or Symptomatic Fever as illustrative of the Nature of Fever in General

London, 1837

Black and Clutterbuck considered fever an overt manifestation of an awakening of the defensive powers of the cells against toxins in the blood stream.

- 1 8 5 5 -

4. PANSUM, P. L. P.

"Zur Lehre von der putriden oder septischen Infectionen." (On the study of the putrid or septic infections)

Schmidt's Jahrbücher (Leipzig) 1855

- 1 8 5 5 -

A toxic principle found in decomposing protein substances was soluble in water but not in alcohol. Sterilization did not alter its toxicity. In minute quantities it was capable of producing fever, and the lethal dose for the dog was only 0.012 Gm.

- 1 8 6 5 -

5. BILLROTH, TH.

Beobachtungstudien über Wundfieber und (accidentelle) Wundkrankheiten (Observational studies on wound fever and accidental wound infections)

Arch. f. klin. Chir. 6:372-495, 1865

Related findings were observed earlier. Fever appeared during the course of injections in patients. Chemical products in the medicaments were first considered to be the pyrexial agents. After a rise in temperature was noted when ordinary distilled water was injected, it was concluded that the solvents were probably contaminated.

- 1 8 6 6 -

6. FRESE, H.

Experimentelle Beiträge zur Aetiologie des Fiebers (Experimental contribution on the etiology of fevers)

Berlin, 1866

In decomposing tissue a soluble toxic material can be secured which resists boiling and passes through filters.

- 1 8 7 2 -

7. MANASSEIN, W.

Ueber die wasserigen und alkoholischen Extracte der Muskeln und der Leber von fieberenden und hungernden Thieren (On aqueous

- 1 8 7 2 -

and alcohol extract of muscle and liver of fasting, feverish animals)

Arch. Path. Anat. 56:220-247, 1872-1873

Glycogen depletion of organs during febrile processes was noted.

- 1 8 7 3 -

8. SENATOR, H.

Untersuchungen Über die fieberhaften Process und seine Behandlung (Research in the febrile process and its treatment)

Berlin, Hirschwald, 1873

- 1 8 7 6 -

9. BURDON-SANDERSON, J.

On the process of fever

Practitioner 16:257-337, 417, 1876

The term "pyrogens" was coined to cover all fever-producing agents.

10. SCHULEIN, W.

Über das Verhältniß der peripheren zu centralen Temperatur im Fieber (On the relationship of peripheral to internal temperature in fever)

Arch. Path. Anat. 66:109-110, 1876

A vasoconstrictive mechanism, produced by fever directly or indirectly, raises internal temperatures.

- 1 8 7 8 -

11. ILIFFE, W.

Remarkable oscillation of temperature without apparent cause

Lancet 2:728, 1878

A patient without other symptoms in whom an increase or decrease in body temperature appeared regularly was described. The cause was not found.

- 1 8 8 7 -

12. VON JAUREGG, J. WAGNER

Ueber die Einwirkung fieberhafter Erkrankungen auf Psychosen
(Action of fever-increasing illnesses on psychotic patients)

Jahrbuch fur Psychiat. Bd. 7:94-102, 1887-8

It was noted that progressive paralysis in long-established neurosyphilis appeared to improve under the stress of acute, accidental infections. The fever engendered was considered to be the curative agent. In attempting to reproduce these beneficial results, von Jauregg utilized both crude typhoid vaccine and injections of malarial parasites, in his later studies, (1917). In the years between 1887 and 1909, he secured favorable results through the use of Koch's old tuberculin.

- 1 8 8 9 -

13. ROUSSY, G.

Recherches experimentales. Substances calorigenes et frigorigenes d'origine microbienne; Pyretogenine et Frigorigenine
(Experimental researches. Calorigenic and frigorific substances of microbial origin; Pyretogenine and Frigorigenine.)

Gaz. d'Hopit. (Paris) 62:171, 1889

Fever and chills were caused by two separate principles residing in bacteria; these were released by lytic action of enzymes and diastases within the body. Roussy isolated these substances from gram-negative organisms.

- 1 8 8 9 -

14. ROUSSY, G.

Recherches cliniques et experimentales sur la pathogenie de la fievre: Theorie générale sur la nature et les rôles physiologiques et pathogene des diastases ou ferments solubles (Clinical and experimental research on the pathogeny of fever: General theory on the nature and physiologic and pathogenic roles of diastases or soluble ferments)

Gaz. med. de Liege 188-189, i, 289-293, 1889. See also: Gaz Hopit. (Paris) 62:286-289, 1889 and Arch. de physiol. norm et path. (Paris) 5:355-370, 1890. Abstract, J.A.M.A. 12:517, 1889

The work of Roussy on the enzymatic etiology of fevers was forgotten in the interest awakened by the school headed by Pasteur.

- 1 8 9 1 -

15. BUCHNER, H.

Tuberculin reaktion durch Proteine nicht-spezifischer bacterien (A tuberculin reaction through nonspecific bacterial protein)

Münch. med. Wchnschr. 38:841-842, 1891

16. RICHTER, P.

Experimentelle Untersuchungen über Antipyrese and Pyrese, nervöse and kunstliche Hyperthermie (Experimental research on anti-pyrese and pyrese, nervous and artificial hyperthermia)

Arch. f. path. Anat. 73:118-165, 1891

- 1 8 9 2 -

17. KANTHACK, A. A.

Acute leukocytosis produced by bacterial products

Brit. M. J. 1:1301, 1892

The author injected sterilized or filtered cultures of Vibrio Metchnikovi, B. pyocyaneus and yeast intravenously into the rabbit, producing immediate pyrexia, initial leukopenia which

- 1 8 9 2 -

was succeeded by marked leukocytosis. The latter was mainly an increase in eosinophils, and the neutrophils were only slightly increased. The increase in leukocytes often lasted over 72 hours.

- 1 8 9 3 -

18. CENTANNI, E.

Ueber infektiöses Fieber. Das Fiebergift der Bacterien (On infectious fever. Fever toxins of bacteria)

Deutsch. med. Wchnschr. 20:148-150; 176-178, 1893 Reforma Med. 256, 1893

A number of pathogens were investigated for the presence of the fever toxin: Diplococcus pneumoniae, B. typhi, B. coli, B. aerogenes and pyocyaneus, the pyrogenic Staphylococci and that of erysipelas, the bacilli of anthrax and tuberculosis, the vibrio (Metschnikoff, Finkler, and Deneke) and the organisms of tetanus, influenza and diphtheria. Activity for fever production appeared to be present in all these toxins and in extracts from the bacterial cells. Centanni found pyrogenic ability of bacterial toxins was not proportional to toxicity; all appeared to contain the same volume of Pyrotoxina bacterica, a water-soluble concentrate, non-albuminoid in character.

19. COLEY, W. B.

The treatment of malignant tumors by repeated inoculations of erysipelas: With a report of ten original cases

Am. J. Med. Sci. 105:487-489, 1893

Pure culture material of S. erysipelatis was injected in ten cases of inoperable or recurrent carcinoma or sarcoma. Even without production of frank erysipelas, response was favorable, particularly in the sarcomas. The curative action is believed to be a systemic rather than a localized one.

20. RUMPF, T.

Die Behandlung des Typhus abdominalis mit abgetödteten Culturen des Bacillus pyocyaneus (Treatment of typhus abdominalis with killed cultures of Bacillus pyocyaneus)

- 1 8 9 3 -

Deutsch. med. Wchnschr. 19:987, 1893

- 1 8 9 4 -

21. CENTANNI, E.

Ueber Infektious Fieber (On infectious fever)

Chem. Zentr. (4th Series) 6:597, 1894

Pyrotoxina bacterica, the fever-producing substance from bacterial cells, was extracted from the organisms following a period of heating (3 hours at 60° C.) and subsequently a more intense treatment at the boiling point. Filtration through a porous clay filter followed; the concentrate was precipitated with absolute alcohol, and the albuminous fraction removed. Re-precipitation and dialysis precedes drying in vacuo.

Administered in superficial vessels under the skin, a chemotactic reaction of the local variety is produced. No suppuration appears, but there is a powerful leukocyte attraction. Larger doses bring forth a flattened, edematous swelling and adjacent blood vessels become hemorrhagic. The chromatin portion of the bacterial cell may be the site of origin of the fever toxin.

22. TEISSIER, J.

Des Pyrexies apyretiques ((Atypical fevers)

Sem. med. (Paris) 14:197-199, 1894

Teissier noted the prevalence of febrile conditions without other symptoms of specific disease. 30 years later, he applied the knowledge gained through his experience (with Rivallier and Reilley) to the treatment of disease making use of the 'propidon' of Delbet, a preparation secured by the digestion of Ps. aeruginosa, streptococci and staphylococci.

- 1 8 9 5 -

23. COLEY, W. B.

Erysipelas toxins and erysipelas serum in the treatment of inoperable malignant tumors. Further observations

Med. Rec. 57:609-612, 1895

24. CZERNY, V.

Über Heilversuche bei malignant Geschwulsten mit Erysipeltoxinen
(On research of healing of malignant growths with erysipelt toxin)

Münch. med. Wchnschr. 185:42; 833-835; 1895

The use of bacterial toxins in the treatment of malignancies was introduced in conjunction with Emmerich and Schöll.

25. EMMERICH, R. and SCHOLL, H.

Klinische Erfahrungen über die Heilung des Krebses durch Krebsserum (Erysipelserum) (Clinical experiences on the healing of cancer with cancer serum (Erysipelserum))

Aerztl. Centr. A. (Wien) 7:209; 228; 242; 358-360; 378-381; 706-712, 1895

26. SWAIN, J.

The treatment of malignant tumors by the toxin of the Streptococcus erysipelatis and B. prodigiosus

Brit. M. J. 2:1415, 1895

27. COLEY, W. B.

Further observations upon the treatment of malignant tumors with the toxins of erysipelas and Bacillus prodigiosus, with a report of 160 cases

Bull. Johns Hopkins Hosp. 65:157, 1896 (Abstract)

In all cases reported, the tumor was either inoperable or had recurred after surgery; in nearly all, microscopic examination confirmed the diagnosis of malignancy. These patients, observed

- 1 8 9 6 -

for a period over four years, presented all varieties of carcinoma or sarcoma. Sarcoma was present in 93, carcinoma and epithelioma in 62 patients. "Of the cases of sarcoma nearly one-half showed more or less improvement; the variety that showed the greatest improvement was the spindle-celled, that which showed the least, the melanotic."

- 1 8 9 8 -

28. BATTLE, W. H.

A case of sarcoma treated with Coley's Fluid

Brit. Med. J. 2:1554, 1898

The complete disappearance of a large tumor of the clavicle, with axillary metastases, followed treatment with Coley's fluid.

29. COLEY, W. B.

The treatment of inoperable sarcoma with the mixed toxins of erysipelas and Bacillus prodigiosus and final results in 140 cases

J.A.M.A. 31:389; 456, 1898

The differences between tumor response to mixed bacterial toxins are pointed out. Results thus far would indicate that the most successful therapeutic results are secured in treatment of spindle-celled sarcoma, in which favorable response has been secured in 50% or more of the cases. Far more resistant are round-celled sarcomata. No improvement in melanotic sarcoma has become permanent. The action of the mixed toxins appears to be one of coagulation-necrosis with fatty degeneration, unlike inflammation or local escharotic effects. That the action is specific is shown by the regression which occurs when injections are made subcutaneously and in sites removed from the tumor.

The dangers attendant on this type of therapy are those of shock, with circulatory collapse, and pyemia of septic origin. The use of the toxins after surgical removal of the tumor is advocated. Mild doses are advocated for this purpose.

- 1 8 9 8 -

30. FOWLER, G. R.

The use of animal toxins in the treatment of inoperable malignant tumors

Am. J. Med. Sci. 116:161-162, 1898

The author presents a review of the use of bacterial toxins in the treatment of neoplasms. Among the reports discussed were those of regressions after accidental infection with erysipelas and after use of the septic products of the causative streptococcus. The value of mixed toxins of erysipelas and B. prodigiosus in the treatment of malignancies was assessed. The rationale for the action of Streptococcus erysipelatis and its toxin is presented. Possibility of new applications of this type of treatment lies in the suggested use of the serum of animals infected with Str. erysipelatis, of the venom of the cobra capello, and in induced suppuration in malignant growths.

- 1 9 0 3 -

31. MACFAYDEN, A.

Upon the immunizing effects of the intracellular contents of the typhoid bacillus as obtained by the disintegration of the organism at the temperature of liquid air

Proc. Roy. Soc. (London) 71:351, 1903

A method for the fractionation of bacterial cellular components is briefly described.

- 1 9 0 6 -

32. COLEY, W. B.

Late results of the treatment of inoperable sarcoma by the mixed toxins of erysipelas and Bacillus prodigiosus

Am. J. Med. Sci. 137:376-78, 1906

The author reviews his work on the treatment of inoperable neoplastic growths through use of living cultures of Str. erysipelatis and its further development in the combined or mixed toxins of

- 1 9 0 6 -

B. prodigiosus and Str. erysipelatis. The results obtained in 38 personal cases so treated are summarized.

- 1 9 0 7 -

33. BEEBE, S. P. and TRACY, M.

The treatment of experimental tumors with bacterial toxins

J.A.M.A. 49:1493-1495, 1907

Inoculations, intratumorally or at a distance of a number of bacterial toxins produced some regression in lymphosarcoma in the dog. Vaccines of B. prodigiosus alone or in combination with Str. pyogenes vaccine, produced some softening and sloughing of tumor tissue and, in one animal, complete regression. A suspension of a vaccine made from B. coli communis was likewise beneficial. Soluble prodigiosus toxins, and a vaccine from Staphylococcus pyogenes aureus, were valueless for this purpose.

34. COLEY, W. B.

Inoperable sarcoma. A further report of cases successfully treated with the mixed toxins of erysipelas and Bacillus prodigiosus

Med. Rec. 72:129-131, 1907

The mixed toxins of B. prodigiosus and Str. erysipelatis were administered to twelve patients with inoperable sarcoma of the long bones. No recurrence of these tumors could be traced for eight of these patients, observed for periods ranging from three to eight years. Four other patients were living, but had been under care for shorter intervals.

- 1 9 0 8 -

35. LANGE, F.

Experimentelle Untersuchungen über das Verhalten der Leukocyten nach Injektionen von Bakterienextrakten (Experimental investigations on the behavior of leukocytes after bacterial extract injections)

- 1 9 0 8 -

Deutsch. Arch. f. klin. Med. 94:552-554, 1908

The initial drop in leukocytes may indicate transient storage in sites of the most intense activity of bacterial toxins. Local and total increases in certain of the cell types are discussed.

- 1 9 0 9 -

36. COLEY, W. B.

The treatment of inoperable sarcoma by bacterial toxins (The mixed toxins of the Streptococcus erysipelatis and the Bacillus prodigiosus)

Proc. Roy. Soc. Med. (Surg. Lectures) Pt. III. 3:1-4, 1909

In summarizing the results obtained in all cases of inoperable sarcoma treated with Coley's mixed toxins, the theories on the modus operandi of these toxins, prophylactic indications, mode and length of administration are presented. The toxic side effects are enumerated.

- 1 9 1 0 -

37. LOEB, L.

The treatment of inoperable sarcoma by erysipelas and prodigiosus toxins

J.A.M.A. 54:262-264, 1910

The author states that approximately 4 to 9 per cent of cases of inoperable sarcoma treated with the toxins of Str. erysipelatis and B. prodigiosus are cured, but the modus operandi remains obscure.

- 1 9 1 1 -

38. BENDIX, B., and BERGMANN, J.

Über das sogenannte Kochsalzfeber (On the so-called 'Salt Fever')

Monatschr. f. Kinderheilk. 11:387-88, 1911

After injections of physiologic saline solution, unexpected temperature increases were noted. Careful preparation of the solution and care in injection appeared to prevent the reaction which was considered to be the result of bacterial contamination.

39. HORT, E. C. and PENFOLD, W. J.

The dangers of saline injections

Brit. M. J. 2:1589, 1911

In a study of the undesirable effects which frequently follow injection of saline solution in both humans and animals, the authors listed: the production of fever, rigors, subnormal temperature, diarrhea, intestinal hemorrhage, Cheyne-Stokes breathing, and, with hypertonic solutions or rapid injection, convulsions and sudden death. Pulmonary edema may follow the injection of large amounts of solution. Vaculation of hepatic cells and changes in heart muscle and renal tissue have been noted on histologic examination. Bacterial products in water, and not the saline, are responsible for these changes. After filtration and heating of water, 19% of the animals used in this study showed slight febrile reactions; only a few organisms were found on culture, and the protein content was very small. When freshly distilled water was used for solutions, few reactions occurred, but when water was used after standing in sterile vessels, fever was produced in direct proportion to the amount injected. A solution containing 73,000 organisms per cc. was used 10 days after distillation; it produced a far less intense elevation of temperature than a sample two months old which contained but 40 organisms. When the contaminated solution was centrifuged, the pyrogenic substance was found to be concentrated just below the surface of the supernatant fluid. This pyrogen was heat-stable and passed readily through a Berkefeld filter.

40. KILIANI, M.

Pharmakologische Wertbestimmung der technischen Fiebermittel (Pharmacologic evaluation of technical fever materials)

Arch. inter. d. pharmacodyn. et d. therap. 20:333-352, 1911

41. LOWENSTEIN, J.

Zur Behandlung die progressiven Paralysis mit nucleinsäuren Injektionen (On the treatment of progressive paralysis with nucleic acid injections)

Klin. Wehnschr. (Berlin) 48:714-716, 1911

The later work of Meyer and Fischer corroborated that of Lowenstein who found nucleic acid to be superior to tuberculin ('old' tuberculin of Koch) for arrest of progressive paralysis.

The mechanism by which results were achieved was unexplained and remissions were not permanent.

42. MÜLLER, P. T.

Ueber den Bakteriengehalt des in Apotheken erhältlichen distillierten Wassers (On the bacterial content of distilled waters stored in pharmacies)

Münch. med. Wehnschr. 58:2739, 1911

43. NOBEL, E.

Zur Kenntniss des temperaturensteigernden Giftes des Bakterium Coli communis (To the knowledge of information on temperature-increasing toxins of B. coli communis)

Arch. Exper. Path. u. Pharmacol. 68:371-382, 1911

An active factor which produced fever was removed from the cells of B. coli communis after hydrolysis, extraction and dialysis. Its characteristics resembled those of cleavage products of protein.

44. THOMSON, D.

A remarkable daily variation in leukocytes in several diseases (Malarial fever, Hodgkin's disease, cancer)

Brit. M. J. 1:1586, 1911

During the sporulation of malarial parasites, toxins are liberated in the blood stream, the mononucleated leukocytes decrease in number, and the total leukocyte count is lower as the temperature rises. When fever declines, the mononuclear cells increase. Even after quinine therapy, marked leukocytosis occurs at the same hour as that of the last sporulation of Plasmodia.

- 1 9 1 1 -

The reduction of the population of malarial parasites is accompanied by a change from severe leukopenia to a marked increase in white cells. Five to ten million dead parasites mixed with corpuscles will stimulate leukocytosis for one hour; this is a good diagnostic sign for malaria.

In Hodgkin's disease a daily variation in leukocytes from 6,000 to 75,000 per cm. may be noted. The mononuclear cells account for 50 to 80 per cent of the increase. A similar change is seen in cancer of the stomach, but fever is absent. The similarity between the leukocyte picture in malaria and that seen in malignancies leads one toward acceptance of the virus theory of etiology for the latter.

45. WECHSELMANN, W.

Neuere Erfahrungen über intravenöse Salvarsaninjektionen ohne Reaktions Erscheinungen (More recent experiences with intravenous salvarsan injections without appearance of reactions)

Münch. med. Wchnschr. 58:1510, 1911

Febrile reactions after salvarsan injection were frequently reported in the literature during the first five years after use of the Ehrlich synthetic began. German physicians sought to place blame upon the solvent (Wasserfehler) while French writers considered the inherent toxicity of arsephenamin responsible. Wechselmann found that temperature elevations persisted in some instances after special treatment and prompt use of the distilled water solvent. The toxic principle therein resisted both autoclaving and filtration.

- 1 9 1 2 -

46. GREENWOOD, H. H.

Note on a case of melanotic sarcoma treated by Coley's Fluids
Lancet 2:881, 1912

Coley reported that the mixed toxins of Str. erysipelatis and B. prodigiosus were of very limited benefit in melanotic sarcoma. A report is presented of the recurrence, in a rapidly spreading form, of melanotic sarcoma previously removed by surgical means. As a last resort, the mixed toxins were administered. The patient had a complete regression and regrowth had not occurred in one and one-half years.

47. HORT, E. and PENFOLD, W. J.

Microorganisms and their relation to fever

J. Hyg. 12:361-390, 1912

Basic arguments are presented for each of the two opposing theories on the origin of bacterial pyrogens. One school of thought, dating from the days of Roussy, would place origin within the tissues and fluids of the host, others consider alien sources (bacterial contamination) responsible for genesis. This discussion considers the possibilities that pyrogens are liberated from bacteria during disintegration within the body or when grown in vitro and that they may be the results of enzymatic breakdown of metabolizable materials, in body tissues or in synthetic media.

48. HORT, E. and PENFOLD, W. J.

A critical study of experimental fever

Proc. Roy. Soc. Med. (London) Series B. 85:174-186, 1912

The terms 'water fever', 'salt fever', 'sugar fever', 'ferment fever', and 'tissue fever' have been applied to pyrexial phenomena in the past. Such terms are misleading. A single entity may be responsible for the appearance of all fever. Its appearance in contaminated water would account for the universality of pyrexia. The present procedures for isolating such materials are insufficiently standardized for reliability in results.

49. HORT, E. C. and PENFOLD, W. J.

The relation of salvarsan fever to other forms of injection fever

Proc. Roy. Soc. Med. (Pt. III. Pathology) 5:131-139, 1912

On the basis of the theory presented by Wechselsman, who found gross contamination of medicinal solutions by bacterial products to be responsible for so-called 'salvarsan fever', the authors studied the pyrogenetic properties of undegraded bacterial protein as well as the efficacy of filtration for the production of nonpyrogenic solutions. They noted, on filtration, a fever-producing substance passed through ordinary bacterial filters from salvarsan solution and other fluids. "Salvarsan fever is (not) necessarily due to the injection of organisms grown on water or saline", they reported, but were unable to determine whether or not pyrogens were products of bacterial protein decomposition. Safeguards against pyrogenic contamination of distilled water were described.

- 1 9 1 2 -

50. LÜDKE, H.
 Ueber Partialgifte im baktierien Protoplasma (On Partial toxin in bacterial protoplasm)
 Deutsch. Arch. f. klin. Med. 105:411-432, 1912
51. MORESCHI, C. and GOLGI, A.
 Dei rapporti anafilassi e febbre (Pyrogenic action of typhoid anaphylatoxin)
 Policlin. (Med. Sect.) 19:431-441, 1912
 Subcutaneous injection of typhoid anaphylatoxin in rabbits produced elevations of body temperature initially but only mild hypersensitive reactions without pyrexia when the injection was repeated.
52. SAMELSON, S.
 Über das sogenannte Kochsalzfeber (On so-called 'Salt Fever')
 Monatschr. f. Kinderheilk. 11:125-133, 1912
53. SEBASTIANI, V.
 I varii tipi febrile riprodotti con un' piro tossino batterica (Varied types of fever reproduced by a bacterial pyrogen)
 Arch. di Biol. (Firenze) 66:137-154, 1912
 A single toxic principle is released by many bacterial specimens. When released in the blood stream of animals or humans it produces a shock phase, followed by fever, with changes in the blood picture. To the pyrogenic agent isolated from B. prodigiosus, the name "pyrotoxin" was applied.
54. VON JAUREGG, J. WAGNER-
 Ueber Behandlung der progressiven Paralyse mit Bacterientoxinen (On treatment of progressive paralysis with bacterial toxins)
 Wiener klin. Wchnschr. 25:61-63, 1912
 After long experience with the use of malarial therapy and the old tuberculin of Koch in the treatment of neurosyphilis, Von Jauregg, with Josef Halban, injected therapeutically killed

- 1 9 1 2 -

cultures of staphylococci and streptococci. The temperatures of patients with tabes andluetice involvement of the central nervous system were not elevated to the degree seen after old tuberculin, but it was noteworthy that relapse was less frequently seen after treatment.

- 1 9 1 3 -

55. CITRON, J. and LESCHKE, E.

Ueber den Einfluss der Ausschaltung des Zwischenhirns auf das infectiöse und nichtinfectiöse Fieber (On the influence of the transection of the midbrain on infectious and non-infectious fever)

Zeitschr. f. Exper. Path. u. Therap. 14:379-390, 1913

Febrile response could not be elicited in the rabbit after section of the medial hypothalamus at the anterior border of the midbrain. Biological pyrogenic agents (Trypanosoma nagana and staphylococci), colloids and chemicals were equally ineffective.

56. FERNANDO-TORRES, F. R.

Vacunotérapia o antigenotérapia antífica

Seman. med. 22:1557, 1913

Typhoid vaccines may be used as nonspecific therapy in conditions other than typhoid itself. Torres suggested that the action of such bacterial materials might be dual, the antigenic-immunizing factor affecting one mechanism for defense, while a second principle produces a more obscured cellular activity.

57. VAUGHAN, V. C., JR. (with J. W. Vaughan)

Protein split products in relation to immunity and disease

Philadelphia, Lea and Febiger, 1913

Proteins could be split into two parts, either by enzymatic action within the organism, or by chemical procedures. A soluble poisonous particle remained in the menstruum. This product played an important role in the genesis of disease; on the injection of very minute amounts immune body production was

- 1 9 1 3 -

stimulated. Vaughan linked infantile diarrhea (cholera infantum) with the presence of 'thyrotoxin', a biological split protein which he found in ice cream and milk.

- 1 9 1 4 -

58. CENTANNI, E.

Über den chemische Wirkung des Bakteriengifte (Chemical action of bacterial toxins)

Zentr. Biochem. Biophys. 16:415, 1914

A basic product of secondary albumin cleavage and not an amino acid or simple polypeptide carried the pyrogenic characteristics found in bacterial proteins. The substance was dialyzable.

59. JOACHIM, A.

Über zehn Fälle von geheilter Paralyse progressive nach Behandlung mit Tuberculin (Ten cases of progressive paralysis cured after treatment with tuberculin)

Wiener klin. Wchnschr. 27:1409-1411, 1914

Detailed case histories for ten patients who presented evidences of psychosis and asocial overt behavior are presented. Remission of the progressive paralysis was accompanied by subjective improvement which enabled them to resume normal activities. The manner by which tuberculin alters central nervous tissue in paresis is not clarified. Hyperpyrexia was not emphasized in this study.

- 1 9 1 5 -

60. HARMER, T. W.

A study of the efficiency of mixed toxins (Coley) in inoperable sarcoma. A critical analysis of 134 microscopically proven cases

Boston M. & S. J. 172:331;373; 411; 440, 1915

An analysis of 134 case reports is presented. Criteria met by

- 1 9 1 5 -

all cases included: Proof by microscopical examination; primary or recurrent inoperable sarcoma; treatment period three weeks or longer; no other therapy given during mixed toxin treatment. Cases were divided into six groups on the basis of response to therapy. Twelve patients received no benefits from therapy; 5 experienced some softening of the neoplasm but no regression; 20 sarcomas disappeared and regrew after some months; growths disappeared, but metastases were seen later in 10 patients. Some diminution of the size of the sarcoma was seen in 14 patients. The 73 patients for whom apparent cures were obtained justify the use of the toxins in these inoperable cases. The greatest percentage of cure or amelioration was noted in the small round cell type of sarcoma. In a few patients, striking relief from pain was achieved.

61. LÜDKE, H.

Die Behandlung des abdominal-typhus mit intravenösen Injektionen von Albumose (Treatment of abdominal typhus through intravenous injections of albumose)

Münch. med. Wchnschr. 62:321-323, 1915

Deutero-albumose produced a febrile response when injected intravenously in treatment of typhoid. Pyrexia might be caused by action of this primary cleavage product of albumin or outside contamination of the solution by pyrogen-yielding organisms.

- 1 9 1 6 -

62. BIELING, R.

Untersuchungen über die veränderte Agglutininbildung mit Ruhrbacillen vorbehandelter kaninchen (Research in the changing agglutination picture in animals previously treated with colon bacilli)

Ztschr. Immunitätsforschung. u. Exp. Therap. 28:246, 1916

The agglutinins increased when previous immunization with colon, dysentery or diphtheria bacillus preparations preceded infections, such as typhoid.

- 1 9 1 6 -

63. GAY, F. P., and CHICKERING, H. T.

Studies in Typhoid Immunization. VI. Treatment of typhoid fever by intravenous injections of polyvalent sensitized typhoid vaccine sediment

Arch. Int. Med. 17:303-328, 1916

64. HEKTOEN, L.

Vaccine treatment

J.A.M.A. 66:1591-1594, 1916

In subacute and chronic localized infections, specific vaccines have, when properly used, valuable stimulatory properties on leukocyte activity and on specific antibody mechanisms. In typhoid fever and, perhaps, in other infections, intravenous injection of specific vaccines and other substances may induce crisis by activity either superimposed upon or varying from the stimulation of antibody production. This the author sees as "a new and interesting development in the study and treatment of infectious diseases."

65. JONA, J. L.

A contribution to the experimental study of fever

J. Hyg. 15:169-194, 1916

A non-protein thermogenic factor was isolated from B. coli communis and E. typhosa. It was colloidal in form, thermo-stable in dry heat at 110° C. and resistant to boiling. Organic solvents were ineffective. In combination with calcium it formed inert salts. In minute quantities (4 gammas) it produced fever in the rabbit.

66. MILLER, J. L. and LUSK, F. B.

The treatment of arthritis by the intravenous injection of foreign protein

J.A.M.A. 66:1756, 1916. See also: J.A.M.A. 67:2010, 1916

Typhoid vaccine was given intravenously for the relief of arthritis. Relief of pain followed. The treatment was

- 1 9 1 6 -

considered primarily one of 'shock'. Unfavorable reactions occurred. In addition to the expected rise in temperature, marked chills, severe malaise and headache were encountered.

67. VAUGHAN, V. C., JR.

Poisonous Proteins, Protein Fever

Herter Lecture. J. Lab. Clin. Med. 2:15, 1916

The author quotes extensively from other authors in the support of his theory that fever is the result of internal destruction of proteins. Bacterial infection in its active phase is caused by liberation of 'protein poison' from bacterial cells. This 'protein poison' is the keystone of all protein molecules and is toxic only when set free. Once released, it possesses the power to destroy the structure of other proteins through the removal of one chemical group.

- 1 9 1 7 -

68. DAVIS, B. F. and PETERSEN, W. F.

A comparative study of lymph and serum ferments during protein shock reactions

J. Exper. Med. 26:699-706, 1917

A preparation obtained from cultures of B. coli produced shock, increased the flow of lymph and antiferments, and raised protease titers. The concentration of lymph fluid is not paralleled by serum changes. The peptidase titer is not increased in proportion to protease. Diastase was found to be unchanged. A rise in lipase was noted.

69. DOCHEZ, A. R. and AVERY, O. T.

The elaboration of specific soluble substance by pneumococcus during growth

J. Exper. Med. 26:447, 1917

During the growth of young cultures of pneumococci and particularly during the period of maximal multiplication, a specific substance is elaborated by the bacteria and is found in cell-free filtrates. Disintegration of the cell is not required for its liberation. It is stable to high temperatures and

- 1917 -

resists boiling. Tryptic digestion, alcohol, acetone or ether precipitation do not change its properties. In animals and in humans, experimentally or accidentally infected with the pneumococcus, a similar or identical substance appears early in the disease process. In blood and urine the specific precipitin reaction may have diagnostic importance.

70. DUPLAN

Les colloïdes et le colloïdotherapie (Colloids and colloid therapy)

Rev. Int. med. chir. 28:46-49, 1917

Administration of colloidal toxic bacterial substances during septicemia activates leukocytosis, both genesis and migration. The osmotic tension of the white cells is reduced after phagocytosis of these colloidal toxins; disintegration of the leukocyte is thus hastened. In the circulation the liberated nucleic acids initiate general systemic changes.

71. HOLLER, G.

Beobachtungen über die Wirkungsweise parenterale einverleibter Proteinkörper zu therapeutischen Zwecken (Observations on the mode of action of parenterally administered protein for purpose of therapy in infectious diseases)

Med. Klin. 13:1038-1041, 1917

The death rate in typhoid was reduced to 0.5% and the average duration of fever to 10 days when a 10% solution of deuterio-albumose was injected in very minute doses once or twice daily. Therapy should be continued until defervescence is well-established.

72. KIBLER, C. S. and McBRIDE, L. F.

Intravenous injection of typhoid vaccine

J. Infect. Dis. 21:13-20, 1917

Intravenous injections of typhoid vaccine were administered to six patients with typhoid fever and to one normal individual. Temperature changes, leukocyte counts, agglutinin and opsonin titers, blood pressure variations and coagulation time were recorded. In both ill and normal patients, chills followed immediately, accompanied by hyperpnea and, in some instances,

cyanosis. The second phase brought a rapid temperature rise which reached its peak in 3 to 5 hours. Defervescence extended over some 18 hours. Leukopenia was found during the first phase. The rise in polymorphonuclear cells followed temperature increases. Blood pressure was decreased during the entire period (10 to 15 mm. Hg.) and returned to pre-injection levels within a few days. Coagulation time was unchanged. Three patients were improved under therapy, and typhoid was apparently aborted in a second case. There was no improvement in two cases and one fatality followed.

73. KRAUS, R., PENNA, J. and BONORINO, C. J.

Ueber Heterobakterientherapie (On heterologous bacteria therapy)

Wiener klin. Wchnschr. 30:869-870, 1917

The therapeutic effects derived from a preparation made from the colon bacillus resembled those secured through the use of typhoid vaccine. The same nonspecific and generalized reaction follows foreign protein therapy. A single substance present in all three materials must be the active agent.

74. MEYER, J. and CARLSON, A. J.

Hunger and appetite in fever

Am. J. Physiol. 44:222-226, 1917

In an attempt to determine the effects of fever on hunger contractions and the mechanism through which such effects are mediated, hunger contractions in the stomach of gastric fistula dogs were observed during a 24 hour period of starvation. After observation of the normal fasting functions of the stomach, the effects of intravenous or subcutaneous injection of the following pyrogenic materials were measured: sodium nucleate, killed B. prodigiosus cultures, peptone, typhoid vaccine and caffeine citrate. The changes in hunger contractions produced by external heat were also recorded. Sodium nucleate, B. paratyphoid, tetanus and diphtheria were incubated with pyloric strips. Sectioning of splanchnic and vagus nerves and its effect on pyrogenic changes revealed the fact that lower vagus tone appeared to inhibit hunger contraction, but no association with the plexuses in the splanchnic area could be traced. Starvation of neonatal puppies did not induce fever. In vitro bacterial toxins did not alter activity of stomach muscle. Temperatures of 103° F. prevented hunger contractions.

- 1 9 1 7 -

75. SCULLY, F. J.

The reaction after intravenous injection of foreign protein

J.A.M.A. 69:20-23, 1917

Patients with articular arthritis experienced pronounced reactions after intravenous typhoid vaccine injections. A chill, accompanied by marked leukopenia, was first noted; rising temperature followed with maximal levels appearing within 3 or 4 hours. Leukocytosis, with the absolute increase largely a matter of neutrophil multiplication, marked the maximal point. The blood pressure mounted during the chill and declined thereafter. Hypotension persisted for 3 to 4 hours, and the return to normal levels was a gradual one. Relief from joint pain was reported by 40% of the patients after a single injection; over one-half were benefited by 2 or more injections.

- 1 9 1 8 -

76. AULD, A. G.

Pyrogenic therapy. With remarks on colloidal metals

Brit. M. J. 1:195, 1918

Various colloidal metals may exert a pyrogenic effect when parenterally introduced. In the course of his investigations the author found a marked reaction could be stimulated by the use of a colloidal platinum preparation over one year old. On injection of this material a sharp rigor followed, then pyrexia, headache, and perspiration. The blood pressure reached high levels during the chill period and fell in the febrile interval; the leukocytes began to rise as the fever increased.

A suspension of colloidal silver was markedly effective in producing the reaction. The investigator deduced from the identical effects elicited by these metals that the metal itself was not the pyrogenic agent. "If the pyrogen...was not colloidal platinum, it matters little therapeutically, as all pyrogenic agents in appropriate doses produce the same effects....Under present conditions typhoid vaccine is probably the best pyrogen to use..."

77. HASHIMOTO, M.

Fieberstudien "Über die spezifische ueberempfindlichkeit des Warmzentrums an sensibisierten Tieren (Fever studies on the excessive sensitivity of heat centers in sensitized animals)

Arch. Exper. Pathol. (Leipzig) 78:370-373, 1918

A primary injection of heterologous protein sensitizes certain cells of the temperature-regulating centers. When the specific antigen is again introduced, an immunologic reaction occurs and fever appears.

78. HASHIMOTO, M.

Fieberstudien. II. "Über die Einfluss unmittelbarer Erwärmung und Abkühlung des Warmzentrums auf die Temperaturwirkungen von verschiedenen pyrogenen und antipyretischen Substanzen (Fever studies. II. The influence of direct heating and cooling of the heat centers upon the temperature-activity of various pyrogenic and antipyretic substances)

Arch. Exper. Pathol. (Leipzig) 78:394-395, 1918

Pyrexia induced by typhoid vaccine and by anaphylactic shock can be modified by direct heat or cold applied to the heat regulatory centers. The action of some antipyretics is altered by these measures.

79. MEYER, J., COHEN, S. J., and CARLSON, A. J.

Gastric secretion during fever

Arch. Int. Med. 21:354-365, 1918

The underlying mechanism affecting gastric secretion during fever was studied further through observations of Pavlov pouch dogs in which fever was induced by intravenous injections of sodium nucleate, inactivated cultures of B. prodigiosus or external heat. Determinations were made of both free and total HCl, pepsin, chlorides and total secretion volume.

External heat (to increase body temperature 2 to 4° F.) or pyrogenic substances decreased total and free acid in gastric secretion, as well as the total volume. A relative increase in pepsin and slight chloride reduction were recorded. During fever, gastrin was ineffective. It was believed that toxins elaborated during fever directly depress the secreting mechanism in the stomach. In particular, the secreting powers of the cells are reduced to a point where nerve impulses can no longer stimulate activity.

- 1 9 1 8 -

80. WELLS, C. W.

Leukopenia and leukocytosis in splenectomized rabbits

J. Infect. Dis. 22:502-506, 1918

The spleen may exert an influence over the total number of leukocytes during the period of increase which follows the administration of killed bacteria. In initial leukopenia, the fate of the leukocytes in the blood from bone marrow and liver parenchyma was studied. Normal levels were maintained in one instance. The mechanism involved was not determined.

- 1 9 1 9 -

81. BALCAR, J. O., SANSUM, W. D. and WOODYATT, R. T.

Fever and the water reserve of the body

Arch. Int. Med. 24:116-128, 1919

The appearance of so-called 'thirst' or 'inanition' fevers (which include the dietary fevers of infants and 'sugar' and 'salt' fever in children and adults) is considered a result of (a) binding of water by electrolytes or organic molecules (b) inadequate amounts of water available for evaporation (c) increased peripheral temperatures. In fever the normal body reserves of water are bound by colloids to an abnormal degree. The marked diuresis and perspiration which follow the dissolution of fever by crisis are evidences of water liberation. Intake of additional fluid would, rationally, reduce fever. Glucose in solution may, when given intravenously at a rapid rate, encourage water loss through glycosuric diuresis and produce typical 'thirst' fever. Loss of water, which has a high specific heat, may depress the process of evaporation. Hypertonic glucose in a single large injection did not result in pyrexia except where the water reserve was previously depleted.

82. COWIE, D. M. and CALHOUN, H.

Nonspecific therapy in arthritis and infections. A study of the changes in the blood consequent to the intravenous injection of typhoid protein. A consideration of the analogy between the typhoid paroxysm and the malarial paroxysm.

Arch. Int. Med. 23:69-131, 1919

Ten case reports of patients suffering from arthritic disease or infections are presented. Intravenous typhoid vaccine was injected after prior determinations of both total and differential leukocyte counts. The usual picture of leukopenia followed injections. Leukocytosis appeared during the pyrexial response. Atypical lymphocytes, nucleated red cells and myelocytes were found in increasing numbers as the white count increased. There was no eosinophilia. Patients who had definite myelocytic response exhibited the best clinical improvement. During fever, swelling and heat increased in joints.

Malarial therapy brings responses similar to those observed after the administration of typhoid vaccine; the common etiological factor in the reaction is believed to be a protein. The reaction is not an anaphylactic one since eosinophilia is absent. In the leukopenic phase the neutrophils leave the blood stream and enter the tissues; the leukocytic phase is characterized by overproduction of cells by 'their respective mesenchymal fundaments.'

83. GOW, A. E.

A note on certain phenomena associated with the protein shock reaction and intravenous vaccine therapy

Quart. J. Med. 13:82, 1919

After a brief review of the literature describing the use of bacterial proteins, the writer presents 8 case histories illustrating the application of this type of therapy to varied disease entities. Coliform bacteria, streptococci, and organisms associated with respiratory disease were used in the preparation of vaccines. Symptomatic and clinical remissions were obtained in 5 patients. Only with the colon type vaccine was marked leukocytosis noted.

84. MÜLLER, E. F.

Die Bedeutung des Kasiens in der Milchtherapie (Significance of casein in milk therapy)

Münch med. Wchnschr. 66:1233, 1919

The protein in milk does not produce fever after intravenous injection. Milk, from which casein had been removed, still produced fever when injected. An endotoxin of bacterial origin is responsible for such pyrogenic activity.

- 1 9 1 9 -

85. MURPHY, J. B. and STURM, E.

Studies on Lymphoid Activity. III.

J. Exper. Med. 29:1, 1919

Animals subjected to dry heat, ranging between 45 and 65° C. for 5 to 15 minutes have sharply decreased total leukocyte counts; thereafter a sharp rise in multinuclear cells takes place. The lymphocytes rise sharply and the increase may reach 300 per cent. The lymphoid tissue stimulation persists 2 to 3 weeks.

86. MURPHY, J. B. and STURM, E.

The lymphocytes in natural and induced resistance to transplanted cancer

J. Exper. Med. 29:25, 1919

Stimulation of lymphoid elements through exposure to dry heat increases immunity to transplantable tumors. Immunity was judged by the percentage of inoculation failures in groups of mice.

87. MURPHY, J. B. and STURM, E.

Effect of stimulation of the lymphocytes on the rate of growth of spontaneous tumors in mice

J. Exper. Med. 29:31, 1919

Grafts were removed from spontaneously appearing tumors in mice and the animals were then exposed to high, dry heat. After temperatures were elevated, tumor tissue grafts were transplanted. In 36 of 61 test animals tumor growth ceased; all tumors were retarded in the remaining members of the group. Successful inoculation and consistent growth of tumor tissue was seen in 28 of 29 control animals.

88. RHYMER, P.

Ueber Milchinjektiontherapie (On milk injection therapy)

Bor Bl. f. Schweiz. Aerzt. 49:1337, 1919

Pyrogens, and not the proteins of milk, are responsible for the fever seen after milk injections. Fresh, sterile milk does not produce temperature elevations.

- 1 9 1 9 -

89. WALKER, I. C.

The treatment of bronchial asthma with vaccines

Arch. Int. Med. 23:220-234, 1919

A group of 178 patients received bacterial vaccines in the treatment of bronchial asthma. Of these, 28 were found to be sensitive to bacterial proteins and were treated with vaccines prepared from organisms to which they were sensitized. The remainder were treated with vaccines from organisms found in sputum or nasal secretions.

Sensitized patients treated with vaccines received much benefit from this type of therapy (75 per cent relieved, 21 per cent improved). Vaccines prepared from organisms grown on plain agar were somewhat less effective (46 per cent of the patients so treated were relieved; 16 per cent only improved). When dextrose bouillon was used for culture of streptococci, 37.5 per cent of the patients treated with a vaccine prepared from the cultures were freed from symptoms, and 25 per cent showed some improvement. Combinations of the two types of vaccines were least effective.

- 1 9 2 0 -

90. BARBOUR, H. G. and HOWARD, H. G.

Coli fever and blood volume (in dogs)

Proc. Soc. Exp. Biol. Med. 17:148-150, 1920

The antipyretic effect of intravenous dextrose is not noted in the healthy dog, but in febrile conditions produced by injections of E. coli organisms, it exerts a distinct fever-reducing effect. The mode of action of dextrose is not clear; it may give osmotic support through increasing the fluidity of the blood at a rapid rate. Such an alteration would aid in the elimination of heat.

When subcutaneous injections of killed colon bacillus preparations were given, sterile abscesses developed in the dog. The blood solids increased at a rate parallel to increases in body temperature; the hemoglobin was concentrated. Loss of blood fluidity in fever reduces blood volume and lowers the rate of heat elimination.

- 1 9 2 0 -

91. BARBOUR, H. G. and HOWARD, H. G.

Dextrose plethora and its antipyretic effect in coli fever

Proc. Soc. Exp. Biol. Med. 17:150-154, 1920

In infections produced by E. coli organisms, a higher percentage of water is held in the tissues. The action of sugars is a hygroscopic one. Withdrawal of water from the tissue spaces increases blood volume.

92. JOANNOVICS, G.

Zur Wirkung fermentativ gewonner Spaltungsprodukte aus Geweben und Bakterien (Action of split products obtained by digestion of tissues and bacteria)

Wiener klin. Wchnschr. 33:649, 1920

Human tumor tissue was digested and sterilized, and its products injected into patients bearing malignant neoplasms. There was proliferation of connective tissue and the area became fibrotic; smaller tumors decreased in size. Split products obtained by the digestion of bacterial cells possessed similar properties. Such products are heat stable (15 minutes at 100° C.) and do not pass through the Reichel filter.

93. SCHERBER, G.

Zur der Bakterienspaltungsproduket (On bacterial split products)

Wiener klin. Wchnschr. 33:654-55, 1920

Superficial epithelioma appeared to be improved by injections of Joannovic's tumor tissue and bacterial split products. The product was administered both intravenously and subcutaneously. For other types of neoplasms, surgical intervention is advised.

- 1 9 2 1 -

94. COWIE, D. M.

Non-specific protein therapy in arthritis

J.A.M.A. 76:310-311, 1921

Sterile albuminose, horse serum and bacterial proteins were used in the treatment of arthritis. The bacterial products were the most satisfactory and typhoid vaccine is the usual choice. No increase in antibody production followed such therapy. The question as to whether or not arthritis is an infectious process remains unanswered.

95. CULVER, H.

Protein therapy in dermatology

J.A.M.A. 76:311-13, 1921

Mixed results followed the use of foreign protein therapy in dermatologic conditions. Typhoid vaccine was used in one case of psoriasis. In certain chronic conditions, such therapy may be beneficial.

96. DUBOIS, E. F.

The basal metabolism in fever

J.A.M.A. 77:352-355, 1921

In fever the standard normal metabolism is much increased. With a reading of 40° C., the acceleration was plus 32 per cent in tuberculosis and plus 48 per cent in typhoid. In malaria and 'foreign protein infection' the increases in metabolic processes may reach plus 60 per cent. A straight line function could be demonstrated in accordance with van't Hoff's law. Reduced heat loss played no part in causing body heat elevation; this was caused solely by increased heat production through metabolic breakdown. When normal metabolism has been calculated by the surface area method, 13 per cent must be added for each degree of temperature above the normal (7.2 per cent per Fahrenheit degree). To this 10 per cent additional must be added in extreme cases of fever or when the food intake is larger than normal; for the restless patient the same increment should be added to calculations.

97. GOVAERTS, P.

Action du serum antiplaquetteique sur la elimination des microbes introduits dans la circulation. II. Effets de la injection de plaquettes (Action of antiplatelet serum on the elimination of bacteria introduced into the circulation. II. Effects of platelet injection

- 1 9 2 1 -

Compt. rend. Soc. de biol. 85:667-668; 745-747, 1921

Intraperitoneal injection of 1 or 2 cc. of washed antiplatelet serum was followed by an injection of either B. coli or E. typhosa organisms. Within 30 minutes the temperature had risen; there were fewer organisms present yet mortality was even higher than that of the control groups. Washed guinea pig platelets markedly reduced the number of organisms (E. typhosa) in circulation, and activity exceeded that seen after erythrocytes were injected.

98. GRAY, F. P.

Use of vaccines in prevention and treatment of influenza and its sequelae

J.A.M.A. 76:244, 1921

A single vaccine prepared from organisms believed to be causative or conjunctive in influenza, or typhoid protein preparations may be used with good effect. Many excellent reports have been made of good results from the use of a mixed vaccine containing those organisms most frequently found in fatal cases (Pneumococci, Streptococci, Staphylococci and B. influenzae).

99. HIRSCH, E. F.

Changes in the leukocytes and the alkali reserves of the blood in experimental infections

J. Infect. Dis. 28:275-278, 1921

Intravenous injection of bacteria caused leukopenia and alkali reserve depletion for about 2 hours afterwards. The normal alkali reserves paralleled the return of leukocytosis.

100. MILLER, J. L.

Foreign protein therapy in acute infections

J.A.M.A. 76:308, 1921

Foreign protein modified the course and mortality rate of typhoid fever. Approximately 40 per cent of patients recover after 2 or 3 injections; the disease terminates by crisis or by lysis. For at least 20 per cent of the patients, such injections shorten the period of illness. "In sepsis and septic endocarditis the blood became sterile within 24 hours after the febrile reaction,

but was not rendered free from bacterial multiplication. In acute alcoholics, foreign protein therapy may be dangerous.

101. PETERSEN, W. F.

Non-specific reaction

J.A.M.A. 75:312, 1921

Nonspecific protein therapy develops resistance to toxic or protein-split products, thereby increasing tolerance. The enzyme systems mobilize and leukocytes are increased. Antibody production may be increased. Treatment should begin early in the course of infection. At this time, cellular permeability is increased, and a local inflammation reaction appears. Later cellular resistance to intoxication is active, but cellular permeability decreases.

102. PETERSEN, W. F.

Protein therapy and nonspecific resistance

New York, 1922, p. 90

The author quotes Luke on the favorable effects which pyrogen injections afford in infectious processes. Stimulation of thermal centers in the brain produces a mobilization of antibody defenses which are progressively reduced in progressive infections

103. PRINZ, A.

Oral Reizetherapie (Oral Stimulative therapy)

Munch. med. Wchnschr. 68:1215-1217, 1921

The initial leukopenia and succeeding leukocytosis after parenteral injection of "foreign" proteins may be noted likewise when methylene blue (or yatren) is administered orally.

- 1 9 2 2 -

104. BARR, D. P., CECIL, R. L. and DUBOIS, E. F.

Clinical calorimetry, XXXIII. Temperature regulation after the intravenous injection of proteose and typhoid vaccine

Arch. Int. Med. 29:608-610, 1922

Heat production in human beings increased from 75 to 200 per cent during the chill produced by proteose or typhoid vaccine. Heat elimination did not increase. Equalization with the environmental heat followed, and elimination was from 20 to 40 per cent above the basal levels. After heat production ceases, elimination continues, producing lower values. That a rapid consumption of glycogen occurs during febrile phases may be ascertained from records of respiratory quotients made during the chill phase. Fever is not caused by failure in the elimination of water. The velocity of chemical reactions follows van't Hoff's law (30 to 60 per cent increase for each 3° C. rise between 37 and 40° C.).

105. PENFOLD, W. J. and ROBERTSON, D. G.

Haemolytic and water fevers

Med. J. Australia 1:129, 1922

The role of bacteria and bacterial products in the production of 'water' fever is discussed.

- 1 9 2 3 -

106. FURTH, O. and LANDSTEINER, K.

On precipitable substances derived from Bacillus typhosus and Bacillus paratyphosus B.

J. Exper. Med. 47:171-176, 1923

Extraction of a culture of E. typhosa through the use of boiling 75 per cent EtOH yielded an antigenic precipitate, insoluble in water, but soluble in weak alkali. It contained traces of phosphorus and 15.7 per cent nitrogen (ash-free preparation). After hydrolysis for five hours, there is little reduction of Fehling's solution. Tryptic digestion alters antigenic activity. A second fraction was obtained by extracting cells with saline and precipitating with 95% alcohol. The fraction had a high nitrogen content. Again tryptic digestion destroyed antigenicity.

From an alkaline extract of the cells held at room temperature, an alcoholic precipitate was secured and redissolved in alcohol. On acidification a final precipitate was obtained which contained little or no protein, a considerable amount of phosphorus, and 39.4 per cent reducing sugar (after 5 hours hydrolysis). This "C" fraction could also be isolated from paratyphosa B organisms; it contained 63.8 per cent reducing sugar. Elemental analysis showed 43.8 per cent carbon and only 1.85 per cent nitrogen.

107. SEIBERT, F. B. and MENDEL, L. B.

Protein fevers with special reference to casein

Am. J. Physiol., 67:105-110, 1923

A study of the fever-producing powers of pure native proteins was undertaken in an effort to determine whether or not bacterial contamination of such solutions produced febrile effects in the rabbit. Each protein was dissolved in freshly distilled water before injection; controls were given the solvent only. All of the 21 protein solutions gave positive reactions, except where the protein was soluble only in alcohol. Casein, on purification, was found to be nonpyrogenic, but could be rendered pyrogenic when it was kept under non-sterile conditions. Nearly sterile milk was pyrogenic in quantities greater than 1 cc. Sterile egg albumin did not produce fever; the egg proteins, edestin and hemoglobin were likewise nonpyrogenic when prepared under sterile conditions. "Protein per se may...not be the cause of fever in the so-called 'protein' fevers."

108. SEIBERT, F. B.

Fever-producing substance found in some distilled waters

Am. J. Physiol. 67:90-104, 1923

The author eliminated, in turn, all possible sources of fever production by parenteral infusion solutions. All experiments were repeated twice and in the same manner. Eliminated as possible causes of fever were: rate of injection; hypersensitivity or immunity (of the rabbit); hemolysis (in blood); pH of solutions; specific effect of ions; impurities from glass, cork or laboratory gases. Non-pyrogenic water could be produced by distillation through an all-Pyrex apparatus. The pyrogen was present in many distilled waters kept more than 4 to 5 days under unsterile conditions, and could be destroyed only by long and drastic heating. Small rod-like gram-negative organisms were isolated from several pyrogenic waters. "The pyrogen is a filterable product produced by a specific bacterium..."

- 1 9 2 5 -

109. HEIDELBERGER, M., GOEBEL, W. F. and AVERY, O. T.

The soluble specific substance of a strain of Friedlander's Bacillus

J. Exper. Med. 42:701-707, 1925

Isolation and chemical analysis of a nitrogen-free specific polysaccharide from the E strain of Friedlander's bacillus are described.

110. LING, C. Y.

The mechanism of reaction of nonspecific protein agents in the treatment of disease. I. The influence of various agents on temperature and leukocyte counts in normal persons and in rabbits. II. The influence of various agents on the mobilization of blood antibodies. III. The influence of various agents on the mobilization of blood enzymes in normal persons and in rabbits.

Arch. Int. Med. 35:598-608; 790-795, 1925

Therapy through induced fever is complicated by the fact that leukocytes do not play the principal role in body defenses, in spite of their phagocytic activity. These cells release endolysins, which, like bacteriolysins, are bactericidal, and elaborate protease and peptidase likewise. The rise in the leukocyte count is caused by an accelerated production of myeloid elements; it varies with the agent employed and the degree of infection present.

Seven foreign proteins and distilled water alone were administered to rabbits and to human beings. Typhoid vaccine (0.2 cc. per kg. of body weight), containing 100 million organisms per cc., and market milk were responsible for the production of more intense pyrexia than other proteins used. Bacterial products increased titer against E. typhosa and Staphylococcus aureus in unsensitized individuals. Complement in serum was depressed, but natural anti-sheep hemolysin was not affected. The slight trauma and bleeding seen in injection has, in itself, a slight nonspecific stimulating effect. The production of large amounts of proteolytic enzyme after foreign protein injection may be either an elaboration by tissue cells in defense or acceleration of the lymph flow from the intestinal tract, or both. Detoxification through the digestion and elimination of bacterial toxins may occur. Proteolytic activity was much increased when typhoid vaccine was injected; it decreased steadily after the initial dose. Peptone decreases proteolysis initially, but a rise takes place soon after. Lipolytic activity in serum was temporarily increased.

111. MENTEN, M. L. and MANNING, H. M.

Relationship of enteritidis - paratyphoid B infections to hyperglycemia in rabbits

J. Infect. Dis. 37:400-410, 1925

Rabbits were subjected to adverse environmental conditions for one year. During this period all blood sugar changes were recorded. Bacterial cultures were made of heart blood and organs of all animals dying or killed within stated periods. Agglutination tests with antisera obtained by rabbit inoculation with these isolated bacteria followed; all organs were examined. In addition the effects produced by both killed and living bacteria were observed in the rabbit and dog.

A general seasonal variation in temperature was noted; the blood sugar levels were found to be in reverse relationship to temperature. Hyperglycemia was seen in cold weather. Most marked hyperglycemic values were found after spontaneous infection with enteritidis-paratyphoid B groups. Hydropic degeneration of the islets of Langerhans was noted only in prolonged hyperglycemic conditions. Blood changes and histologic alterations were similar regardless of whether or not the injected organisms were viable. The dog exhibits a natural resistance to the organisms used in the study.

112. MULLER, E. F.

The involuntary nervous system: An important factor in the body's resistance

Arch. Int. Med. 35:796, 1925

The theory is presented that the route or site of injection and not the substance injected produces the phenomena seen in parenteral therapy. Dilution in peripheral vessels alters the leukocyte concentration. An overbalance of parasympathetic tonus follows with stimulation of myeloid tissue. This may be the result of kinetic response of the involuntary nervous system. Intradermal water, milk or salt solution (or air) caused rapid leukopenia and a high concentration of leukocytes in the liver. Peptone injection produced a migration of leukocytes away from the periphery toward organ storage centers.

113. PRIBRAM, B. O.

Proteintherapie und chirurgische Therapie des Magen-und-duodenalgeschwürs (Protein therapy and surgical treatment of stomach and duodenal ulcers)

- 1 9 2 5 -

Deutsch. med. Wchnschr. 51:141, 1925

Pain in peptic and duodenal ulcer was reduced, and some healing of the lesions was demonstrated roentgenically when surgical procedures and nonspecific protein therapy were used in conjunction.

114. SEIBERT, F. B.

The cause of many febrile reactions following intravenous injections

Am. J. Physiol. 71:621-651, 1925

Pyrogens in distilled waters were injected in the rabbit. After sensitizing doses, typical anaphylactoid reactions followed the provocative injection. Febrile reactions were most intense after intravenous injection but could be obtained by use of the subcutaneous and intraperitoneal routes. Non-chromogenic bacteria appear to produce the pyrexial substance, which is destroyed by heat, only after treatment with strong acid or alkalis. Some nitrogen is found, but the content appears to have little correlation with pyrogenic activity.

115. ZIMMER, A. and BUSCHMANN, P.

Vermeidung und Verhandlung von Schaden die durch Protein-Körpertherapie hervorgerufen werden (Avoidance and palliation of illness through so-called protein therapy)

Zeitschr. f. artztl. Fortbild. 22:513, 1925

For chronic conditions the best results with protein nonspecific therapy are those secured without the appearance of fever. In acute illness or infections a febrile reaction appears to be more satisfactory.

116. ZECKWER, I. T. and GOODELL, H.

Blood sugar studies. I. Rapid alterations in the blood sugar level of rabbits as the result of intravenous injections of killed bacteria of various types

J. Exper. Med. 42:43-56, 1925

Heat-killed, 18-hour cultures of B. proteus, B. coli, and paratyphosus A and B, produced changes in blood sugar levels after intravenous injection into the rabbit. The saline suspension contained 2 billion organisms. Marked hyperglycemia was

- 1 9 2 5 -

noted after injections of B. proteus, B. coli and paratyphosa B. Slight and irregular increases were seen after B. paratyphosa A and B. enteritidis solutions were injected. Leukopenia accompanied hyperglycemia and persisted after the blood sugar had returned to normal levels. No relationship could be traced between temperature fluctuation, blood sugar increases and the leukocyte count. B. faecalis alkaligenes, Str. hemolyticus or viridans, Staphylococcus aureus, B. pyocyaneus and B. typhosa, in similar preparations, had little effect on blood sugar levels.

- 1 9 2 6 -

117. BARBOUR, H. G.

Heat regulation and water exchange. XI. Blood and plasma changes in cocaine and B. coli fevers and aspirin antipyresis.

J. Pharm. & Exper. Therap. 29:427-440, 1926

When fever of a mild grade is produced in dogs through injection of B. coli, blood plasma is slightly reduced. Blood solids rise. Fluid loss may be the result of increased capillary permeability with escape of plasma into tissues from the general circulation. Increased vascularity may be seen. The alkaline content of the blood falls. When a synthetic antipyretic is administered, the fall of temperature is closely related to the return of fluid into the blood stream.

118. BOSNAME, R. K.

Thrombopenia of infective diseases and of protein injections

Folia Haemat. 33:105-121, 1926

Thrombocytes play an important role in immunity. Their normal activities include agglutination of bacteria and removal from the blood stream. When thrombocytes are destroyed, the bactericidal factor is liberated into the serum. Thrombopenia is the result of an exhaustion of activity and may be seen after the injection of parenteral protein, living or killed bacteria, toxins and in hemorrhage. Splenectomy prevents exhaustion.

119. CANNEYT, J. VAN

Action du Bacillus subtilis et de ses secretions sur le bacille de la tuberculose (Action of B. subtilis and its secretion on the tubercle bacillus)

- 1 9 2 6 -

Compt. rend. Soc. biol. 95:878-881, 1926

Inocula of preparations of *B. subtilis* overran the growth of the tubercle bacillus on potato cultures and exhibited a preferential selection between the mycobacterium and other contaminants. Cultures of *B. mesentericus vulgaris* had similar activity. The activity appears to be antibiotic in nature and not merely one of competition for growth elements.

120. CONSOULOFF, S.

Theoretical experiment in curing by increased temperature and relation to the cancer problem (Translated Title)

Ann. Univ. of Sofia Sci. 22:1-24, 1926

The spirochaeta of syphilis are sensitive to high temperatures; other parasitic organisms may be similarly affected. Human cellular tissue is passively stimulated by fever. These two facts explain many of the benefits derived from bacteriotherapy. Applications of these findings to cancer therapy are discussed.

121. DREYFUS, G. L. and HANAU, R.

Grundsatzliches über die verwendung des neuen Fiebermittels Saproviton in Neurologie (Fundamentals of the application of the new fever producer, Saproviton, in neurology)

Deutsch. med. Wchnschr. 52:1381-1383, 1926

122. FAHRAEUS, R.

Physikalische Gewebsveränderungen bei Fieber und Hyperpyretischen (Physical tissue alterations through fever and hyperpyretics)

Acta med. Scand. (Supplement) 16:608-610, 1926

A preliminary report on the alterations seen in tissues during fever in infections and that induced by hyperpyrexial agents.

123. FAHRAEUS, R.

Über Änderung des Aggregatzustandes lipoider Substantzen bei Fieber u. d. hyperpyretischen Temperaturen (Changes in the aggregate condition of lipid substances in fever and hyperpyretic temperatures)

Skand. Arch. Physiol. 49:125, 1926 (Abstract)

124. KRAUSS, E.

Untersuchungen über den minimalen eiweiss Gebrauch des Menschen unter gesunden und krankheiten Bedingungen (Minimal protein requirements of man in health and disease)

Deutsch. Arch. klin. Med. 150:13-59, 1926

Nitrogen excretion is increased after the injection of bacterial and other 'foreign' proteins only when fever is produced in the sensitized animal. In some types of tissue destruction urinary N remains near the minimal point because of the reuse of protein in metabolism. Minimal total N is placed at 1.0 to 1.1 Gm. per square meter of body surface.

125. OKURA, Y.

Influence of toxic substances on nitrogenous and purine metabolism in dogs: Influence of pyrogenous substances (Translated Title)

Jap. J. Med. Sci. (Tr. IV. Pharmacology_ 1:65-71, 1926

The acceleration of catabolic processes and the influence of the tissue destruction on toxicity must be considered when the effects of fever are evaluated.

126. REICHE

Toxisches oder dynamisches Eiweiss Fieber (Toxic or dynamic protein fever)

Ztschr. f. Kinderheit. 410:96, 1926

Addition of a protein concentrate to a milk formula increased body temperatures of healthy infants. Breast milk (or formula alone) were not pyretic agents. Split products of protein digestion may be absorbable in the digestive tract or intestine.

127. SOLOMON, H. C., BERK, A., THEILER, M. and CLAY, C. L.

Use of soduku in the treatment of general paralysis. A preliminary report

Arch. Int. Med. 38:391-404, 1926

The excessively high fever and distinct lymphaginitis which follows the experimental use of 'rat-bite fever' therapy appears to influence the central nervous system and produce favorable changes in progressive paralysis.

- 1 9 2 6 -

128. TOMCSIK, J.

On carbohydrate-like specific substance in the colon-Aerogenes group

Proc. Soc. Exper. Biol. & Med. 24:810, 1926

A method is given for separation and purification of the specific polysaccharides of B. lactis aerogenes, encapsulated B. coli strains and five other Aerogenes organisms. Dissolution of the capsule occurs through treatment with KOH; alcoholic precipitation follows. On repurification, the nitrogen content could not be reduced below 0.9 per cent. The polysaccharide was not antigenic for the rabbit, was specific for its own parent strain and gave no cross reactions with immune sera.

129. VON TUKACS, J.

Protein Körpertherapie und vegetatives Nervensystem (Non-specific protein therapy and the autonomic nervous system)

Wiener med. Wchnschr. 39:885-888, 1926

Nonspecific protein therapy develops a stimulatory effect in the sympathetic system, with the initial reaction one typical of sudden sympathetic response. Within a few hours, the opposing parasympathetic system is likewise stimulated; this effect is more prolonged and therapeutically important.

- 1 9 2 7 -

130. ANDREWS, E., PETERSEN, W. F. and BOIKAN, W.

Observations of eviscerated dogs (Effects of bacterial proteins)

Proc. Soc. Exp. Biol. Med. 25:109-110, 1927

The response of febrile agents may be conditioned by the effects of stimulation in splanchnic organs. When the proteins of B. coli organisms are administered to the eviscerated dog, neither chills nor fever follows.

131. BRANHAM, S. L. and HUMPHREYS, E. M.

Soluble antigens of B. enteritidis

J. Infect. Dis. 40:516-524, 1927

Bacteria-free filtrates from B. enteritidis were concentrated in vacuo and the properties of the opalescent supernatant fluid and grey precipitate observed. The precipitate was chiefly polysaccharide. Dialysates from the supernatant fluid contained less than 1/4,000,000 Gm. of protein, but were superior to living bacterial emulsions as antigens.

132. EVANS, C. LOVATT and ZECKWER, I. T.

On the nature of the hyperglycaemic response to injections of certain killed bacteria

Brit. J. Exp. Path. 8:280-288, 1927

A hyperglycemic response follows the injection of B. coli and B. proteus killed cells in rabbits. Splanchnic nerve stimulation appears to initiate glycogenolysis; removal of the thyroid decreases it, but thyroid preparations given orally increase the response. When the adrenals are inactive, hypoglycemia may reach lethal levels. Sectioning of the vagus nerve does not prevent reduction in blood sugar.

133. GOEBEL, W. F. and AVERY, O. T.

The soluble specific substance of Friedlander's bacillus. III. On the isolation and properties of the specific carbohydrates from types A and C Friedlander bacillus

J. Exper. Med. 46:601-607, 1927

Two specifically-reacting N-free polysaccharides were isolated from strains of types A and C Friedlander's bacillus. While chemically they possessed individual properties, the following similarities were noted: high specificity; non-reducing in pure form; reducing sugars found on hydrolysis; strongly acidic with glucuronic acid or its isomer within the molecule. The specific function and carbohydrate structure were apparently inseparable.

134. HEIDELBERGER, M.

The chemical nature of immune substances

Physiol. Rev. 7:107-128, 1927

The author presents a summary of the current information on the

- 1927 -

actual chemical nature of immune substances. He discusses in turn proteins and their derivatives, the lipoids and carbohydrates.

135. HOWARD, H. J.

Nonspecific protein therapy in eye inflammations with specific reference to the use of typhoid vaccine

China Med. J. 41:395-407, 1927

Typhoid vaccine arrests inflammation in uveal infections; benefits are proportional to the fever produced. Milk and protein split products may be used, but typhoid vaccine is the selection of choice.

136. LILIENTHAL, H. and COLEY, W. B.

Mediastinal sarcoma treated with Coley's fluid

Ann. Surg. 85:615, 1927

Three years after treatment with Coley's fluid, of a child for whom a diagnosis of inoperable sarcoma of the posterior mediastinum had been made, no residual, physical or roentgen evidence of the existence of the tumor could be demonstrated.

A series of six patients with inoperable sarcoma of the spine responded to mixed toxins with remarkable 'clinical remission.' Successful results may be expected in approximately 10 per cent of cases treated by this method.

137. MATSUNAMI, Y.

Experimentelle Untersuchungen über das Verhalten der Leukocyten (Experimental investigation on relations of leukocytes)

Okayama Igakkuai Zasshi 39:622-625, 1927

Protein, carbohydrates and fat have widely differing effects upon leukocytes. On intravenous injection, protein acts most intensely on the neutrophils, to a lesser degree on lymphocytes and affects eosinophils moderately. Carbohydrates stimulate neutrophil production, decrease the lymphocytes and infrequently alter the ratio of eosinophils. Only after the injection of fat is lymphocyte activity heightened; production continues for a considerable period.

Inorganic stimulants differ in activity from those of organic origin. The leukocytes are little influenced by the parenteral injection of metals. The salts have least activity and the alkalis the greatest. Eosinophils are increased in blood after injection of metallic colloids.

138. MUELLER, E. F.

Nonspecific protein therapy: Its action and its application
Wisconsin M. J. 26:287-288, 1927

In nonspecific protein therapy the vegetative nervous system alters and produces early changes through the splanchnic area. Major internal organs are mobilized into rhythmic action, followed by reflex action at the periphery which decreases capillary capacity.

139. SEIBERT, F. B.

"Welche substanzen erzeugen Fieber nach intravenöser Injektion"
(What substance produces fever after intravenous injection?)

Arch. Exp. Path. Pharmacol. 121:247-249, 1927

The author opposes the theory of Donath and his school, holding that fever after intravenous injection of distilled water is caused, not by nucleic acid, but by a metabolic product of the bacterial contaminants.

140. STOKES, J. H.

Modern Clinical Syphilology

Philadelphia, W. B. Saunders Co. 1st Edition, 1927, pp. 177;
346

The "water error" (Wasserfehler) is commonly found on large dermatologic services where large amounts of distilled water are prepared in advance. Showers of reactions, with chills and fevers were noted Mondays on the author's clinic. The 'tubing reaction', first described by Stokes and Busman, is characterized by the appearance, within 30 minutes after an injection, of malaise, pain in the lumbar region, nausea and diarrhea with tenesmus. The chill may be the first symptom. With repetition of the rigor, sharp rises in temperature may be seen. There is often prostration and emotional disturbance. The fever may persist beyond the usual 8 hours. Profuse outbreaks of herpes have followed the reaction. Leukocytosis (with 96 per cent polymorphonuclear cells) is present.

The use of nuclein in neurosyphilis, advocated by Fischer and by Meyer, is discussed at length. Fischer observed remissions

- 1 9 2 7 -

in 58 per cent of the patients treated. He found sodium nucleinate to be a useful substitute. The initial dosage used was 2.5 to 3 cc. of a 10 per cent solution; interrupted treatment was the most successful. The marked leukocytosis, not fever produced, received credit for therapeutic benefits. Meyer, in comparing nuclein with tuberculin and arsephenamine alone, reported successes in 42 to 50 per cent of cases treated with the former. Tuberculin produced remissions in but 25 per cent of the cases treated; arsephenamine, in 7.0 to 7.5 per cent.

141. TEISSIR, P., REILLY, J. and RIVALIER, E.

Recherches experimentales sur l'infection chancrelleuse. I. Preparation d'une antigene Streptobacillaire soluble (Ses Effets sur l'animal et l'homme) (Experimental investigations on chancroid infection. I. The preparation of a soluble antigen Streptobacillin. Effects in animal and man.)

J. Physiol. et Path. 25:268-282, 1927

"Streptobacillin" a homologous vaccine-type extract, was prepared from Ducrey's bacillus. A mixture of nucleoprotein-globulin complexes, it is non-toxic to mice, guinea pig or healthy humans in the ordinary dosages. On intravenous injection certain reactions associated with protein 'shock' were noted. The substance may be used by subcutaneous or intradermal injection, as well. The preparation appears to have sensitizing activity; marked reactions occurred when injections were given to persons already infected with the bacillus. There was no complement fixation or inactivating ability in serum after the use of the extract, the precipitin reaction was feeble, and no bactericidal activity could be demonstrated.

142. TEISSIR, P., REILLY, J. and RIVALIER, E.

Recherches experimentales sur l'infection chancrelleuse. II. Les principes généraux d'une vaccinothérapie spécifique (Experimental investigations on chancroid infection. II. General principles of specific vaccine therapy)

J. Physiol. et Path. 25:319-338, 1927

The respective values of specific vaccine therapy and nonspecific protein therapy for the treatment of chancroid were compared. A homologous vaccine "Streptobacillin" was administered to persons infected with Ducrey's bacillus and the therapeutic results obtained appeared to parallel the degree of sensitization, most marked in conditions of long standing. The results would suggest a provocation of inflammatory processes, favoring leukocytosis and phagocytosis.

- 1 9 2 7 -

143. VON EULER, U.

Über Hyperadrenalinämie bei Fieberzuständen (On the increased blood adrenalin in fever conditions)

Pflüger's Arch. ges. Physiol. 217:699-716, 1927

- 1 9 2 8 -

144. ALLEN, A. W. and SMITHWICK, R. H.

Use of foreign protein in the treatment of peripheral vascular disease

J.A.M.A. 91:1161, 1928

Typhoid vaccine, containing 125 to 300 million bacilli per dose, was used in the treatment of Buerger's disease, in gangrene of arteriosclerosis and simple vasomotor types of vascular occlusion. The result of this treatment was most promising. It is believed that the benefits are chiefly derived from vasodilatation, which brings relief of vasospasm and increases capillary circulation.

145. BORCHARDT, W.

Zur Physiologie des Fiebers Über die Bedeutung von Schilddrüse und Nebennieren für das Fieber (Physiology of Fever. On the importance of thyroid and adrenals in fever)

Arch. Exp. Path. u. Pharmacol. 137:45-70, 1928

In an exhaustive study of the fever produced by bacterial, chemical and mechanical stimulation of the midbrain, a primary action upon the heat centers was noted. This can be delayed by denervation of the liver. The presence of thyroid and adrenals is important. The adrenals are not necessary to fever production, but after removal, susceptibility to fever is increased. However, no rigor follows when adrenals are removed. When adrenals are intact, puncture of the heat center of the brain alone leads to marked leukocytosis.

- 1 9 2 8 -

146. BORCHARDT, W.

Neutrophile Leukocytosis und Nebennieren (Neutrophilia, leukocytosis and adrenals)

Klin. Wchnschr. 7:2440-2442, 1928

147. CENTANNI, E.

Die Reversibilitat in der antibakteriellen und antitumoralen Vaccinebehandlung (Reversibility in antibacterial and anti-tumor vaccine therapy)

Deutsch. med. Wchnschr. 7:519-530, 1928

148. CHRISTIAN, S. L. and PALMER, L. A.

An apparent recovery from multiple sarcoma, with involvement of both bone and soft parts, treated by toxins of erysipelas and Bacillus prodigiosus (Coley)

Am. J. Surg. 4:188-197, 1928

A case report is presented involving myelosarcoma of the femur, with wide-spread metastases and recurrence at the stump after mid-thigh amputation. The diagnosis was microscopically and roentgenologically confirmed. The neoplastic area regressed and finally disappeared after a series of treatments with Coley's mixed toxins. Two years later, the patient was reported to be in good health.

149. COLEY, W. B.

End results in Hodgkin's Disease and lymphosarcoma treated by the mixed toxins of erysipelas and Bacillus prodigiosus, alone or combined with radiation.

Trans. Am. Surg. 46:331-357, 1928

The author discusses the theories prevailing on the etiology of Hodgkin's disease and the manner in which differentiation may be made between this condition, lymphosarcoma and other members of the lymphoma group. An impressive number of cases of Hodgkin's disease and lymphosarcoma have been treated with the combined toxins of erysipelas and B. prodigiosus alone or in conjunction with radium and X-rays. Five-year survivals in this group have been frequently attained.

150. HESSE, E. and TAUBMANN, G.

Untersuchungen über das Fieber nach Heuimfus (Inquiry into fever after hay infusion)

Arch. Exp. Path. u. Pharmacol. 136:239-256, 1928

Extracts of decomposed hay, injected parenterally, caused temperature increases. Analyses of these 'pyrogens' revealed that humic acid was present. Like the humic acid from turf and peat, it produced fever and leukopenia.

151. KUTNEZOWSKY, N.

Über die Einfluss lokaler thermischer Reize auf den Speicherungsprozess in den Lymphknoten (The influence of local hyperemia upon the storage process in lymph nodes)

Ztschr. ges. Exp. Med. 62:44-61, 1928

In the stimulation of the reticuloendothelial system, the route of introduction of various substances may be of great importance. Storage of substances directly introduced into the blood stream takes place in the reticuloendothelial cells of the blood vessels; if the material is introduced into the lymph gland, storage occurs in the cells of the lymph nodes. When colloids are injected subcutaneously, storage is seen in macrophages of the connective tissue. It is possible to influence the character of distribution of colloids through local hyperemia; storage thereafter takes place much earlier and in larger amounts than is seen in unstimulated tissue. Heating of the lymph nodes before injection of a dye hastened the rate of storage.

152. SHWARTZMAN, G.

Studies on Bacillus typhosus toxic substance I. The phenomenon of local skin reactivity to B. typhosus culture filtrate

J. Exper. Med. 48:247, 1928. See also: Proc. Soc. Exper. Biol. Med. 25:560, 1928

Toxic filtrates were obtained from tryptic digest broth cultures of E. typhosa strains. In rabbits intracutaneous injections of a sensitizing dose, followed in 24 hours by a provocative injection, produced hemorrhagic-necrotic skin lesions at the site of the initial injection. From 78 to 79 per cent of normal rabbits were sensitive to this reaction, which was not dependent on the amount of filtrate used. No relationship was shown

- 1 9 2 8 -

between the degree of erythema after preparatory injection and the intensity of the hemorrhagic reaction produced by intravenous filtrate. Skin preparation with nonspecific irritants did not render tissues responsive to subsequent injections of filtrate. Neither did the injection of the bacterial filtrate into the skin provoke the hemorrhagic-necrosis phenomenon, which is not related closely to bacterial hypersensitivity. The filtrates themselves resisted inactivation by heat at a range of pH between 4.0 and 9.0.

153. TOMCSIK, J. and KUROTKHIN, T. J.

On the role of carbohydrate haptens in bacterial anaphylaxis

J. Exper. Med. 47:379-388, 1928

Carbohydrate haptens obtained from B. lactis aerogenes, the pneumobacillus and from a yeast were found capable of producing anaphylactic shock in vivo and in vitro in guinea pigs passively sensitized with immune rabbit sera. Toxic and minimal shock-producing doses were found to have a wide margin, corresponding to those noted in protein anaphylaxis. Active sensitization could not be produced with these substances.

- 1 9 2 9 -

154. BREUER, S.

Insulinartige Substanzen aus Bacterium coli und Rauschbrandbakterien (Insulin-like substance from B. coli and Blackleg bacteria)

Munch. med. Wchnschr. 76:741-742, 1929

Filtrates from two cultures of B. coli were injected into the rabbit. A severe hypoglycemia followed, similar to but less marked than that produced by insulin; it was relieved by the administration of sugar. The filtrates from cultures of Clostridia chauvoei were four times as active as those prepared from B. coli but were orally inactive. When concentrated through use of acid alcohol, the potent principle was thermostable.

155. FREMONT-SMITH, L., MORRISON, R. L., MAKEPEACE, A. W.

Capillary flow in man in typhoid vaccine fever

J. Clin. Invest. 7:489, 1929

Changes in the capillary blood flow appear in three stages. At the onset of fever complete stasis begins, with cyanosis appearing at the height of the reaction. As fever mounts, the blood flow is increased. The typical flush is seen near the peak of febrescence; it continues throughout the succeeding fall in temperature and, occasionally, for some hours thereafter.

Stasis is initiated by a rapid constriction of the terminal arterioles and is prevented by the local application of heat. It also can be abolished once it has begun by the same means. Constriction in renal blood vessels leads to water retention and a subsequent dilution of blood, producing diuresis in the later stages of fever.

156. GROLLMAN, A.

Variations in the cardiac output of man. V. The cardiac output of man during the malaise and pyrexia following the injection of typhoid vaccine

J. Clin. Invest. 8:25-32, 1929

The cardiac output, pulse rates, blood pressure, temperature and oxygen consumption of 8 individuals were studied after the intramuscular injection of typhoid vaccine. Soon after injection the cardiac output was increased to a significant degree, but at the height of malaise and fever, the output was somewhat reduced. Pulse rate and oxygen consumption paralleled the degree of malaise and of fever.

157. HOFF, F.

Ueber den Einfluss von Baktierieneiweiss auf das Blut (Influence of bacterial protein on the blood)

Verh. Deutsch. ges. inn. Med. 41st Kong. Inn. Med. 208-211, 1929

The influence of bacterial protein on cellular and chemical elements, volume and rate of flow of the blood is discussed.

- 1 9 2 9 -

158. URBANEK, J.

Die Bedeutung der Tuberculose für die entzündlichen Erkrankungen des Uvealtrakts (The importance of Tuberculose for inflammatory ailments of the uveal tract)

Berlin, S. Karger, 1929

Extracts from cultures of human tubercle bacilli and pure bacterial proteins from organisms with specific activity were combined and used in treating 500 patients with uveal infections seen on the services of the Meller Clinic, Vienna. After 0.1 mg. was administered as a sensitizing dose, injections were continued, with the range of dosage from 0.0001 to 0.1 mg. The toxicity associated with tuberculin was absent in this preparation.

159. WAGNER, W.

Untersuchung der Bakteriziden Bestandteile des B. pyocyaneus.
(Investigation of the bactericidal constituent of B. pyocyaneus.)

Zeitschr. Immunitäts. forschung., 63:483-491, 1929

A highly potent bactericidal and detoxifying solution was prepared from cultures of B. pyocyaneus and pyocyanase.

- 1 9 3 0 -

160. CARTER, E. B.

A proposed chemical test for pyrogen in distilled water for intravenous injections

J. Lab. Clin. Med. 16:289-290, 1930

If pyrogens are not present, the color of a sample of distilled water, to which H_2SO_4 (10 cc.) and 0.05 N $KMnO_4$ (0.1 cc.) have been added, remains unchanged after boiling.

161. HAMEL, J.

Die Behandlung des Trippers bei Frauen mit Pyrifer (Treatment of gonorrheal infections of women with pyrifer)

Dermat. Ztschr. 60:404-412, 1930

Specific vaccines were added to pyrifer and administered to 15 patients with gonococcal infections. The treatment was successful for 73.3 per cent of the group. Pyrifer alone, in dosages calculated to increase the body temperature to 40° C., brought relief to 34.8 per cent of the patients in a second group. The values of adjuvant therapy are discussed.

162. HARRISON, F. C., TARR, H. L. A. and HIBBERT, H.

Factors relating to carbohydrates and polysaccharides. XXXIII. Synthesis of polysaccharides by bacteria and enzymes

Canad. J. Res. 3:449-463, 1930

163. KRUMBRUCH, O.

Pyriefieberhandlung bei leuischen Nervenkrankheiten (Pyri(fer)-fever treatment in neurosyphilitic conditions)

Deutsch. Med. Wchnschr. 56:2170-2173, 1930

A preparation produced by Rosenberg of Freiburg (Pyrifer) has been used for therapeutic fever production in the Psychiatric and Neurologic disease Clinic of the University of Göttingen. As prepared from the killed bacteria of the colon group found in milk, pyrifer gives rapid elevation of temperature, occasionally chills, and a strong leukocytic response. It is utilized in seven different strengths containing from 50 to 500 million bacterial 'units' and is usually administered intravenously, although more protracted fever follows intramuscular injections. Leukocyte values rise to 20,000 or 25,000 total counts, with neutrophils of immature type accounting for most of the increase; lymphocytes are decreased in number.

Eight of 13 tabetic patients treated with pyrifer improved. Paralysis with tabes is less favorably influenced than pure paralysis with neurasthenic complications. Optic atrophy may be hastened by fever therapy. There is speculation as to whether or not heightened blood circulation may be credited for the benefits received from this type of therapy.

164. RADEMAKER, L.

The cause and elimination of reactions after intravenous infusions

Ann. Surg. 92:195-201, 1930

Two specimens of distilled water, one known to contain pyrogens, were carefully studied. In the pyrogenic solution, total and organic solids, ammonia and an ammonia fraction were increased. The residue produced fever in animals. The bacterial count was high in both pyrogenic and non-pyrogenic waters but the pyrogenic specimens contained only one-third as many organisms as the nonpyrogenic. The use of water traps in the distillation step and immediate sterilization and sealing of solutions were advocated as measures against contamination of products destined for intravenous injection.

165. SPINELLI, A.

Una sostanza ipertermica nella levodura (A substance producing hyperthermia contained in yeast)

Biochem. terap. sper. 17:397-401, 1930

Autolyzed yeast produces a pyrexial substance after standing at room temperature for 15 to 25 hours. On injection, this material, in amounts of 0.7 to 0.8 cc., increased body temperature in the rabbit to 41.8° within 5 or 6 hours.

166. STETNER, E.

Über die Wirkung von Bakterieneiweiss ein Spritzungen bei Kindern, mit besonderer Berücksichtigung des Geleimrheumatismus (Zur Frage der Herpes sentstehung) (The action of bacterial protein, etc. on the arthritis of childhood, with special attention to the question of the appearance of herpes)

Ztschr. f. Kinderh. 48:623-639, 1930

167. WESTPHAL, K. and WEGSCHEIDER, K.

Zur Pyriferbehandlung der Tabes und Taboparalyse (Pyrifer treatment of tabes and tabetic paralysis)

Deutsch. Med. Wchnschr. 56:1731-1733, 1930

- 1 9 3 0 -

Excellent results were obtained by the use of pyrifer in classic tabes, with lightning pains, gastric crises and "girdle pain". It was less effective when ataxic complications were present (5 of 13 patients improved). The improvement of psychic symptoms in tabes is most spectacular. In one instance, improvement was seen in sexual impotency. The authors review the literature on the use of pyrifer. They find 87 per cent of tabetic patients exhibited improvement, while in paralysis, 45 per cent in a large group (104 persons) were benefited.

- 1 9 3 1 -

168. ARNETH, J. and ALBACHT

Über das qualitative und quantitative (reaktive) Verhalten der Leukocyten bei der Fiebertherapie mit Neosaprovitan B (Qualitative and quantitative proportions of leukocytes in Neosaprovitan B fever therapy)

Ztschr. f. ges. Exper. Med. 77:641-688, 1931

Neosaprovitan B, a vaccine prepared from saprophytic organisms, exerts a profound effect upon blood-forming organs through the mediation of the vegetative nervous system. Leukocytosis appears early, while lymphocytes increase within 24 hours after injection. Suggestions are made for its application to tabes dorsalis and multiple sclerosis therapy.

169. DAVESNE, J., and HABER, P.

L'influence de l'hyperthermie, de l'acide et de l'alcalose sur la production des agglutinins (Influence of alkalosis and acidosis and the thermal reactions (produced by a vaccine) on the production of agglutinins)

Compt. rend. Soc. biol. 100:1053-1054, 1931

A preparation which contained the endotoxins of histolytic bacilli produced fever in rabbits upon intravenous injection. When acidosis or alkalosis appeared during pyrogenic fever, the production of agglutinins was unchanged.

- 1 9 3 1 -

170. DELAFIELD, M. E.

A comparison of the changes in the blood sugar and blood phosphorus in rabbits following the injection of suspensions of Bact. aertrycke

J. Path. Bact. 34:177-194, 1931

Hyperglycemia appeared in rabbits after injections intravenously of suspensions of B. aertrycke or filtrates from such suspensions. Hypoglycemia and death followed. Elevated blood sugar levels and inorganic phosphorus bore inverse relationships to each other; increases above initial values of P were seen during the hypoglycemic phase. Organic acid-soluble blood phosphorus rose and fell with the carbohydrate content. Diphtheria toxins did not alter blood sugar and phosphorus in the same way; the latter reached a high level two days after the injection of toxin. B. aertrycke cells were active after autoclaving, although their enzymic activity was reduced. Alcohol and acetone extraction did not alter their capacity for changing blood sugar and phosphorus ratios.

171. DIETEL, F.

Fieberbehandlung mit Pyrifer (Fever therapy with pyrifer)

Med. klin. 27:622-624, 1931

A bacterial preparation (pyrifer) had been used in over 600 cases of gonorrhea and in psoriasis. The dosage was not easily determined, since the febrile reaction was not a prerequisite for therapeutic response. Good results were also obtained by its use in progressive paralysis, tabes and vulvovaginitis.

172. EVANS, A. C.

The effect of bacterial by-products on leukocytes

J. Bact. 21:42-43, 1931

Certain by-products of bacterial culture growth were compared with known toxins (tetanus and diphtheria). The phagocytic activity of leukocytes was not impaired by washings from cultures of Streptococci scarlatinae. Organic acids and free H ions impaired leukocytic functions.

173. FINKELSTEIN, H. and WEIL, E.

Zur Kenntnis des alimentarien Fiebers. IV. Das Kochsalzfieber (Information on alimentary fevers. IV. Saline fever)

Ztschr. Kinderh. 50:259-287, 1931

Saline 'fever' may appear after a single dose of hypertonic solution given orally or after repeated oral administration of hypotonic or isotonic solutions. Amounts as small as 50 cc. per kilogram of body weight may bring temperature increases. Ringer's solution has similar hyperthermic effects. The pattern of reaction is one of defense against disturbance of normal tissue concentrations of salts; the reaction diminishes the pressure gradient with the production of fever. This last-named phenomenon may be the culmination of tissue activity directed toward dehydration.

174. FINKELSTEIN, H.

Zur Kenntniss des alimentarien Fiebers. VI. Hyperthermie oder Fieber? (Information on alimentary fevers. VI. Hyperthermia or fever?)

Ztschr. Kinderh. 50:637-650, 1931

When body cells are forced to yield intracellular water, fever appears. Protein and sodium chloride retain water and are additional factors in its diversion from its physical destination. Desiccation of the liver during periods of high body temperature produces profound alterations in body chemistry, and colloid chemical changes are seen in the body cells. "The...theory of pyrogenous substances or protein split products as basic causes of temperature elevations...has inadequate background. Presence of neither...essential for fever production..."

175. FINKELSTEIN, H. and WEIL, E.

Zur Kenntniss des alimentarien Fiebers V. Zur Pharmakologie des Eiweiss-und Kochsalzfiebers (Information on alimentary fevers. V Pharmacology of protein and saline fevers)

Ztschr. Kinderh. 50:288-291, 1931

No distinction between hyperthermia and 'true' fever was found.

176. GOLDIE, H.

Les substances antihemolyses (antihemolytic substances)

Compt. rend. Soc. biol. 109:7-9, 1932

Thermostable and filtrable disaggregations of the products of culture filtrates of Streptococci, Staphylococci, E. typhosa and E. coli exert a marked antihemolytic influence on the blood. While the substances appear to be related to antibiotic and immunologic factors in the bacteria, they are not present in very young or very old cultures. Purified products are crystalline NH_2 compounds.

- 1 9 3 1 -

177. HEIDELBERGER, M. and KENDALL, F. E.

Specific and nonspecific polysaccharides of type IV pneumococcus

J. Exper. Med. 53:625-639, 1931

Three N-containing polysaccharides were isolated from autolyzed cultures of Type IV. pneumococci. One was a type-specific carbohydrate, which was separated with difficulty and required more alcohol for precipitation than the specific polysaccharides of Types I, II and III. A serologically inactive substance allied in structure to chitin was found. The "C" substance of Tillett, Goebel and Avery was present.

The inactive fraction had the lowest optical rotation value and the highest carbon content. It was least soluble in alcohol or acetic acid and contained a small amount of nitrogen. On hydrolysis it yielded crystals which exhibited the optical rotation characteristics of glucosamine. The reducing sugars accounted for 55 per cent of all chemical constituents; total N was 5.9 per cent. Like chitin and chitosan, the inactive polysaccharide was partially hydrolyzed by snail juice.

178. KROMAYER, E.

Über Fieber Behandlung mit Pyrifer (Fever treatment with pyrifer)

Dermat. Wchnschr. 93:1547-1550, 1931

A preparation from the sterilized protein of killed bacteria from milk (pyrifer) produced fever of 39-40° C. in 1 to 2 hours after injection, and chills in 6 to 10 hours. Its use in buboes, gonorrhea, balanitis, psoriasis and staphylococcic infections

increased leukocytes in the blood and produced other changes indicative of an action upon heat-regulating nervous centers. Some benefits were noted in all ailments treated, with the exception of psoriasis. The preparation is ineffective when given intramuscularly. Headache and herpes labialis were listed among the side effects.

179. LEUDEL, E. and DUTT, J. N.

The influence of elevated temperature on the reaction curve of endocrine glands (Translated Title)

Fermentsforsch. 12:335-340, 1931

Febrile conditions, whether produced by infections or through the injection of pyrogens, increase the interferometer reading for the Abderhalden reaction. The functions of the suprarenals may be widely altered in fever.

180. LUSTIG, B. and LANGER, A.

Ueber die Bestimmung und den Gehalt der freien, Eiweiss und gebundenen nichteiweiss Kohlenwasserstoffs im norm und pathologischen Seren (Determination and content of free, protein-bound and non-protein carbohydrate in normal and pathologic serum)

Biochem. Ztschr. 242:320-337, 1931

High values were found for sugars associated with protein in the serum during febrile infections. The highest values were recorded in carcinoma, sarcoma and pneumonia. No increases were noted in chronic conditions in which fever did not appear.

181. NEUBERGER, J.

Einblick in die Wirkungsweise einer Pyriferkur (Glance at the manner of action of pyrifer therapy)

Med. klin. 27:954-956, 1931

The action of pyrifer on neurosyphilis was observed in patients treated on the services of the State Hospitals in Berlin. Temperature seldom rose above 40° C. and local reactions were infrequent. Pyrifer gave a stronger therapeutic reaction than either specific or nonspecific protein therapy. The leukocyte and differential counts were altered; the white cell count rose to 16,000 in many cases, occasionally as high as 33,000. The

- 1 9 3 1 -

decrease began 6 to 7 hours after administration. A sharp, initial leukopenia sometimes antedated leukocytosis; where this occurred, stab cells made up approximately 50 per cent of the total leukocyte count. Although it is an active hematopoietic agent, pyrifur is not a cure in the final sense. In neurosyphilis, it is a valuable adjunct in combined therapy.

182. SABIN, A. R. and WALLACE, G. B.

On the nature of the chill-producing principle in anti-pneumococcus serum

J. Exper. Med. 53:339-362, 1931

A chill may occur 30 to 60 minutes following the injection of anti-pneumococcus serum. It is accompanied and followed by a rise in temperature which is prolonged for several hours. Since the same reaction has been noted after intravenous injection of other solutions and in transfusion, a study of its properties and origin was made. The influence of drugs upon the chill-producing factor was ascertained.

No relationships between the chill, anaphylaxis or the toxic effects of drawn blood were noted. The principle appears to be formed in blood after standing; it is not removed from solution by anticoagulants, dialysis, filtration or moderate heat. It was independent of the formed elements, the lipids or fibrinogen. Fractionation methods, utilizing the changes in pH and NaCl concentration, removed acid and alkaline globulin fractions from antibody solutions. The acid globulin apparently acts as an adsorptive carrier for the chill-producing fraction, while the alkaline globulin serves as a vehicle for the antibody substance.

183. SHWARTZMAN, G.

Phenomenon of local skin reactivity to bacterial filtrates in the treatment of mouse sarcoma 180

Proc. Soc. Exper. Biol. Med. 29:737, 1931

Agar washings of Meningococcus 44D group I were filtrated and injected intravenously into mice bearing sarcoma 180. Doses ranging from 0.25 to 0.5 cc. were given singly, as repeated injections, and at varied time intervals during tumor growth. The majority of neoplasms responded with prompt hemorrhage followed by complete regression or regression succeeded by slow regrowth. The effect appears to be selective. The substance was highly toxic.

184. SOLARI, L. A.

Temperature et reactions thermiques chez les chiens hypophysoprives ou á tuber lésé (Temperature and thermal reactions in hypophysectomized or tuber-injured dogs)

Compt. rend. Soc. biol. 108:125, 1931

185. SPEIRER, C.

Die unspezifische Behandlung des Gonorrhoe mit Pyrifer (Non-specific treatment of gonorrhea with pyrifer)

Dermat. Wchnschr. 92:13-17, 1931

Immunologic mobilization, and not natural resistance, are cardinal features of all nonspecific 'fever' therapeutics. The authors found a death point for gonococci in vitro at 40 to 41° C.; others, particularly Koch, have noted no absolute destruction through heat in vivo. A bacterial protein material from apathogenic colon type bacilli (pyrifer) was used in treatment in all seven strengths (50 to 500 million organisms). Pyrifer fever resembled that of malaria. Most frequently encountered reactions were those of headache, muscle pain and herpes labialis. While pyrifer has a place in the therapeutic armamentarium, it is not a panacea.

186. WOLFF, L. K. and RAS, G.

"Über unspezifische Therapie II. (On nonspecific therapy. II.)

Ztschr. ges. exp. Med. 75:719-736, 1931

A nonspecific bactericidal effect was found to be common to organic compounds (31), inorganic salts (12), biologic proteins (12) and suspensions of dead bacteria (14), when these materials were injected intravenously into the rabbit. When tolerance to histamine developed, there was tolerance to all other materials injected. Appearance of a tolerance to gram-negative bacilli produced similar tolerance to all other gram-negative organisms, but none to gram-positive bacterial preparations. When the materials were injected in a reverse order, the tolerance exhibited likewise was in reverse. Subcutaneous injection itself is bactericidal. When leukocytes were destroyed by coagulation, the blood and serum of the rabbit exhibited bactericidal properties.

- 1 9 3 1 -

187. ZOZAYA, J.

Serological study of polysaccharides of meningococcus, B. anthracis, B. proteus, B. subtilis and B. mesentericus

J. Exper. Med. 54:725-732, 1931

- 1 9 3 2 -

188. BANUS, M. G. and GINSBURG, E.

Changes in blood associated with fever induced by killed B. coli

Am. J. Physiol. 101:106-114, 1932

Vaccines of B. coli or typhoid organisms, or the suspensions of B. coli, produced a very marked alteration in the blood of the dog. A series of injections of increasing strength was more effective in producing high temperature levels than was a single dose. The concentration of the blood increased; hemoglobin and oxygen capacity were increased together. The alkali reserve declined. The reactions observed included the appearance of bloody stools and diarrhea, nausea and vomiting.

189. HARTMAN, F. A. and SCOTT, W. J. M.

The protection of adrenalectomized animals against bacterial intoxication by extract of adrenal cortex

J. Exper. Med. 55:63-65, 1932

Adrenalectomized rats were given preliminary injections of cortin or equivalent volumes of saline solution until body weight appeared to be standardized. Surviving animals were divided into groups two weeks later; one group received standard typhoid vaccine followed by 0.5 cc. of cortin or saline, respectively; injections were continued until death ensued or animals recovered from the intoxication produced. A second group received injections of saline or cortin daily with increasing doses of killed Staphylococcus aureus organisms (2 to 16 billion).

Acute intoxication from killed E. typhosa and chronic intoxication after injection of Staphylococcus aureus were resisted by those animals protected with cortical extracts. Larger amounts were necessary for protection during the peak of the intoxication. The authors believe that inadequate production of cortin may be one factor in prolonging severe pyogenic infections which endure over two to three weeks.

190. HENCH, P. S.

Usual and unusual reactions to protein (fever) therapy

Arch. Int. Med. 49:1, 1932

Physiologic changes during the "protein" reaction include those in basal metabolism and blood pressure, peripheral and splanchnic vasomotor mechanisms and in size and permeability of the arterioles and capillaries. Fluctuations in renal function are seen. There may be pH, N, P, urea and uric acid, albumin and allantoin concentration alterations and the amounts excreted may vary. Immunological phenomena may be observed (antibody and complement fixation) and the flow of bile, lymph, saliva, breast milk or menses may be increased. The activity of liver, gastrointestinal tract and spleen may vary from the norm. Of physical changes observed in blood, those of volume alteration, specific gravity, freezing point and viscosity are most evident. There are certain cellular blood characteristics which reflect the results of temperature elevation; platelet fragility is increased and the content of fibrinogen may vary. Sedimentation rate may be affected. Chemically, CO₂ tension and combining power of plasma and total NPN of whole blood and serum may reflect the general state during fever. Sugar tolerance and the ratio of albumin to globulin have shown variation in hyperpyrexia.

191. MURRAY-LYON, R. M.

Nonspecific protein therapy in rheumatism

Edinburgh M. J. 39:619-627, 1932

In chronic arthritis, typhoid vaccine or protein preparations bring a temporary response only. In the subacute arthritic conditions, typhoid vaccine may be beneficial. Its action on collagenous tissue has not been explained.

192. NOUVELAS, P. A.

Pirogenos: Modo de producto (Pyrogen production)

Siglo med. 89:77; 1080; 163, 1932

- 1 9 3 2 -

193. VON PAP, Z.

Neosaprovitan

Arch. f. Psychiat. 97:450-460, 1932

194. ZOZAYA, J.

Carbohydrates adsorbed on colloids as antigens

J. Exper. Med. 55:325-351, 1932

Bacterial polysaccharides, including dextran, are non-antigenic when administered in small doses alone. When adsorbed on a colloid carrier, as haptens, their antigenicity appears. The possibility is considered that anticarbohydrate antibodies are produced.

- 1 9 3 3 -

195. CANZIANI, G. and LONGO, V.

Donaggio's 'inhibition phenomenon' in the urine and spinal fluid of neuro- and psychopathic patients (Translated Title)

Boll. soc. ital. biol. sper. 8:1476-1478, 1933

The Donaggio 'inhibition phenomenon' does not appear in the normal individual after induced fever unless mental symptoms appear; it is not seen in neuropsychopathic or schizophrenic subjects. During epileptic seizure, the reaction is positive and may be a useful diagnostic sign.

196. CARMINATI, V.

Influenza di culture di germi lattacidogeni sulla crescita del cancro del topo (Influence of lactic acid bacilli on mouse cancer)

Boll. d. Inst. sierotherap. 12:205-220, 1933

The author injected cultures of termobacterium bulgaricus in milk serum into mice bearing cancer implants, with inconclusive results.

197. DONAGGIO, A.

Curves showing the 'Inhibition phenomenon' in urine and spinal fluid under various febrile conditions (Translated Title)

Boll. soc. ital. biol. sper. 8:1459-1462, 1933

Hyperpyrexia induced by injections of pyrogenic substances (proteins and vaccines) does not cause immediate changes in the urine and spinal fluid. Resistance is exhibited in a latent period, extending 12 to 24 hours after the onset of ordinary fever, and to 36 hours after injection of pyrogens. No lag is seen when fever is induced by physical means. Fever provoked by the injection of malarial parasites is preceded by alterations in urine and spinal fluid before the rise in temperature. The inhibiting substance can be removed from fluids by dialysis or by adsorption with bone charcoal or kaolin, followed by filtration.

198. DURAN-REYNALS, F.

Reaction of transplantable and spontaneous tumors to blood-carried bacterial toxins in animals unsusceptible to the Shwartzman phenomenon

Proc. Soc. Exper. Biol. Med. 31:341, 1933

The tumor hemorrhage and necrosis which follows the intravenous injection of bacterial filtrates of *B. coli* was studied in rats and mice, animals normally unsusceptible to the Shwartzman phenomenon. The majority of animals carried rapidly growing, malignant tumors, a smaller proportion bore slower-growing spontaneous or transplanted malignant growths, embryomas or granulomas.

A strongly positive hemorrhagic reaction was seen in the group bearing rapidly growing transplanted tumors; the larger the tumor the greater, in general, was the reaction. Slower growing and less malignant tumors were not affected. Animals bearing regressed tumors were resistant to regrafted tumor tissue. Toxic effects of the filtrate appeared to be related to the size and age of the tumor. The investigators found only tumors showing malignancy and rapid growth were capable of response to bacterial toxins.

199. GIAJA, J. and DIMITRIJEVIC, I.

La thermoregularisation sur la fièvre (Thermoregulation during fever)

Arch. int. pharmacodyn. 45:342, 1933

- 1 9 3 3 -

Fever caused by induced Pyocyaneus infection in rats did not inhibit the temperature regulating apparatus of the brain centers. The temperature remains high during wide variations in environmental temperature.

200. GORDON, R. E. and STARK, C. N.

The toxicity of young cells of S. paratyphi A and B when lysed by bacteriophage.

J. Infect. Dis. 62:45-47, 1933

Young bacterial cells did not yield toxic fractions after autolysis and filtration, but a slight toxicity was shown by preparations made from S. paratyphi A and B cells 6 or more days old. The maximal toxicity was present between 8 and 14 days of age. Aging again reduced the toxic factor. At room temperature for 2 months toxicity was gradually lessened. Filtration, additional bacterial inoculations and low temperatures did not reduce activity.

201. HSAIO-CH' IEN, CHANG

Gastric secretion in fever and infectious diseases

J. Clin. Invest. 12:155-169, 1933

Gastric acidity falls during febrile conditions and finally reaches levels 66 per cent below the normal. Chloride ion concentration falls with the volume of secretion. The decreases are in direct proportion to the degree of fever.

202. HEUBNER, W. and SILBER, W.

Der Bestimmung der antipyretischen Stoffen (Testing of antipyretics)

Arch. Exp. Path. Pharmacol. 169:530-536, 1933

The most regular and enduring fever reproducible in rabbits is seen after infusion of hay extracts. If subcutaneous injections are used, no ill effects follow high temperature elevations. The method may be employed for testing antipyretic drugs.

203. LEVINE, P. and FRISCH, A. W.

Further observations on specific inhibition of bacteriophage action

Proc. Soc. Exper. Biol. Med. 31:46, 1933

Specific inhibition of bacteriophage by two extracts from organisms of the Salmonella group was reported (Proc. Soc. Exper. Biol. Med. 30:993, 1933). In addition, the specificity of the inhibition was determined through isolation of specific phages for B. paratyphosa B and B. suispestifer from chicken stool filtrates. The extract of B. suispestifer had only a feeble influence on anti-B. paratyphosa phage, while the latter was inhibited markedly paratyphosa B. extract. The latter was only slightly specific for anti-B. suispestifer phage; its homologous extract was far more potent as an inhibitor. The order of activity of extracts did not run parallel to living bacillus phage titer.

204. LEWISOHN, R. and ROSENTHAL, N.

Prevention of chills following transfusion of citrated blood

J.A.M.A. 100:466-469, 1933

Transfusion reaction rates were reduced 1 to 12 per cent by careful preparation of solution and rigorous antisepsis of apparatus used in transfusion. Pyrogen reactions are reduced to a minimum by such precautions.

205. RADEMAKER, L.

The cause and elimination of reactions after intravenous infusions

Surg. Gynec. and Obst. 56:956, 1933

Pyrogenic reactions are produced by toxic by-products elaborated by the bacteria and not by bacterial components. Careful distillation, immediate sterilization and cautious attention to the prevention of later contamination will reduce pyrogens in fluid.

- 1 9 3 3 -

206. THOMPSON, S. A.

Preparation of dextrose and saline solutions and apparatus for intravenous and subcutaneous use

Am. J. Surg. 22:127, 1933

In a committee report on the causes of reactions following intravenous infusions, the presence of bacterial pyrogens in improperly distilled water, is considered to be the most important cause of the twelve listed. The correct method of distillation is described, and a permanganate color test is recommended for use. Techniques for the preparation of glucose, normal and 2 per cent saline solution are presented.

207. WAELSCH, H. and WEINBERGER, E.

Das Gehalt des glutathiones im Blut (Glutathione content of blood)

Arch. exp. Path. Pharmacol. 169:625-632, 1933

Glutathione and ergothionine in human blood are increased during pyrexia produced by injections of pyrifin, milk protein or bacteria. Glutathione alone does not influence the temperature of the normal rabbit nor in animals after the removal of thermoregulatory organs.

- 1 9 3 4 -

208. BANKS, H. M.

A study of hyperpyrexia reaction following intravenous therapy

Am. J. Clin. Pathol. 4:260-291, 1934

Organisms isolated from various pyrogenic waters were identified as Pseudomonas scissa and Ps. ureae. A suspension of killed organisms elicits a fever of 3 to 4° F. in the rabbit, but no immunity or hypersensitivity was noted after repeated injection.

Bacterial suspensions of the two organisms were injected separately in patients with advanced neurosyphilis, dementia paralytica and tabes dorsalis. Central nervous system disturbances began to clear within three weeks; the psychiatric symptoms abated to a considerable extent.

209. BIERRY, H.

L'action spécifique et la structure chimique (Specificity and chemical structure)

Compt. rend. Soc. biol. 115:1168-1170, 1934

The role of carbohydrate and protein in determining specificity of reaction to living or killed bacteria is briefly discussed.

210. BLUM, D. N. and COLEY, W. B.

Large malignant tumor (sarcoma) of the uterus with multiple bone metastases (femur, pelvic bones and skull)

Am. J. Surg. 1:47, 1934

A patient with a malignant myoma and multiple skeletal metastases received a series of injections with 'Coley's toxins' after total hysterectomy. No return of malignancy was evident six years after treatment.

211. BURNET, F. M.

The phage-inactivating agent of bacterial extracts

J. Path. Bact. 38:285-299, 1934

The findings of Levine and Frisch that bacterial extracts may be capable of inhibiting activity of phages lysing parent strains are confirmed. Extracts from resistant strains fail to inhibit phages to which bacteria themselves are resistant. Homologous antibacterial serum destroys the phage inactivating power.

212. CECIL, R. L.

The medical treatment of chronic arthritis

J.A.M.A. 103:583, 1934

Small intravenous doses of streptococcus vaccine, given four to five days apart, and subpyrexial in dosage (50,000-100,000,000 organisms) brought some improvement in chronic arthritis. At the time of writing, the author preferred this type of treatment.

213. DARROW, K. E.

Review of causes of reactions following intravenous injections of glucose and normal saline (with special reference to pyrogen containing distilled water as a cause)

Journal-Lancet 54:65-66, 1934

214. DELAFIELD, M. E.

Blood sugar changes and toxic effects produced in rabbits by certain fractions derived from B. aertryke

Brit. J. Exper. Pathol. 15:130, 1934

Fractions prepared by the method outlined by Raistrick and Topley were capable of producing hyperglycemia and toxic effects in the rabbit. Antibacterial and antitoxic immunity production in mice by means of these fractions appeared to parallel their toxicity for the rabbit. The active fraction of B. aertryke is precipitable in alcohol (50-68% by weight), a second inactive fraction is obtained by use of excess alcohol. The alcohol-soluble material is also inactive. The hyperglycemic effect can be removed from the precipitable active fraction by acidic or alkaline hydrolysis. Repeated injections of this fraction produces tolerance to the chemical response.

215. GOUGH, G. A. C. and BURNET, F. M.

The chemical nature of the phage-inactivating agent in bacterial extracts

J. Path. Bact. 38:301-303, 1934

The phage-inactivating agent of bacterial extracts is a polysaccharide which can be converted into a hapten by alkaline treatment. The power to inactivate different phages is lost by degrees during degradation by either heat or alkali. The investigators consider the specific somatic antigen of dysentery and Salmonella bacilli to be a labile polysaccharide and to be responsible for serological properties and phage susceptibility of the bacteria.

216. HOOPER, F. E., RENFREW, A. G. and JOHNSON, T. B.

The chemical study of bacteria. Studies on the complex carbohydrates isolated from the culture medium and the ultrafiltrate after the growth of tubercle bacilli

Am. Rev. Tuberc. 29:66-72, 1934

A fraction secured by the hydrolysis of a culture of tubercle bacilli 6 weeks old contained d-galactose, d-mannose and d-arabinose with a small amount of uronic acid. The nitrogen-free ultrafiltrate contained a polysaccharide with identical specific reactivity. The polysaccharide did not contain phosphorus.

217. HOWIE, J. E.

Observations on the use of pyrifer

J. Ment. Sci. 80:521-525, 1934

Pyrifer is a preparation produced from killed colon bacilli and prepared in seven strengths, containing from 50 million to 500 million bacteria per cc. The initial dosage recommended ranges from 25 to 50 million bacteria. The intravenous route is preferred. The febrile reaction begins one-half to 1 hour after administration and reaches its height in 3 hours, falling by lysis. A secondary rise follows with lower maxima; there is complete recovery in 24 to 48 hours. A regular fever rhythm appears, usually after the 3rd treatment. Side effects are moderate in degree and, unlike malarial or cabinet therapy, the treatments can be given to patients with serious cardiac lesions without adverse effects. The successive phases of leukocytosis and lymphocytosis resemble those seen in other types of nonspecific therapy. Howie administered pyrifer to 15 patients with advanced general paralysis, all too feeble for malarial therapy. Six were improved, 2 discharged as cured. The average duration of treatment was between 3 and 6 weeks. In schizophrenia (15 cases) improvement was seen in 5. The treatment was ineffective for chronic manic states. Pyrifer reduced seizures in one epileptic patient from 32 in 3 weeks to 16 in the same length of time. No permanent remission could be obtained.

218. LANFRANCHI, F.

Ricerche sul pirogene batterico: influenza degli amino acid nei mezzi sintetici, sulla produzione del pirogene (Investigation on bacterial pyrogens: influence of synthesis of amino acids in media on pyrogen production)

- 1 9 3 4 -

Ann. igiene 44:448-469, 1934

219. NEDZEL, A. J.

A study of muscle temperature during bacterial chill

Proc. Soc. Exper. Biol. Med. 32:279, 1934

The muscular tremor observed during bacterial chill was not accompanied by an increase in temperature in the skeletal muscles. Such muscular activity differs from that of muscular work which is accompanied by increased heat production.

220. (COUNCIL ON) PHARMACY and CHEMISTRY, AMERICAN MEDICAL ASSOCIATION

Erysipelas and prodigiosus toxins (Coley)

J.A.M.A. 103:167, 1934

The Council briefly reviews the literature on the use of the mixed toxins of Coley. Its members conclude that "...the combined toxins of erysipelas and B. prodigiosus may sometimes play a significant role in preventing or retarding malignant recurrence or metastasis (and) occasionally they may be curative in hopelessly inoperable neoplasms. Probably their value is strictly limited to tumors of entodermal or mesodermal derivation, and more particularly in the case of bone tumors, to those exhibiting little or no osteoplasia." For these reasons, the Council retained the toxins of Coley in New and Non-official Remedies with a view to facilitating their further study, particularly for a prophylactic use in conjunction with conservative or radical surgery.

221. PIOTTI, M.

Terapia pirotogena in casi di ebefrenia iniziale (Pyrogen therapy in cases of early hebephrenia)

Schizofrenie 4:443-447, 1934

222. RAISTRICK, H. and TOPLEY, W. W. J.

Immunizing fractions isolated from B. aertrycke

Brit. J. Exp. Path. 15:113, 1934

The preparation of an immunizing fraction is described.

- 1 9 3 4 -

223. WRIGHT, I. S.

The modern medical treatment of diseases of the peripheral vascular system

Med. Clin. N. Amer. 17:1429, 1934

Leukocytosis or immobilization of immune bodies may account for a portion of the benefits secured through the use of 'fever therapy'. Vascular changes bring relief from spasm in partially occluded vessels; local capillary dilatation brings an improved blood supply to ulcerated areas and healing is promoted.

- 1 9 3 5 -

224. APITZ, K.

Studies on the chemical nature of Shwartzman-active substances

J. Immunol. 29:343-358, 1935

The nucleoprotein fraction of bacteria, obtained by autolysis or extraction, is Shwartzman-active. The potency is increased in alkaline solutions. A metabolic fraction can be isolated from agar washings through alcohol precipitation; it contains a highly unstable Shwartzman-active fraction, not identical with type-specific carbohydrates. It may be dialyzed slowly from filtered washings but resists tryptic digestion.

225. DURAN-REYNALS, F.

Reaction of spontaneous mouse carcinomas to blood-carried bacterial toxins

Proc. Soc. Exper. Biol. Med. 32:1517, 1935

Previous experiments had revealed the insusceptibility of slow-growing malignant carcinomas of the mouse to the tumor-necrotic action of certain bacterial filtrates. Highly toxic agar washings of human typhoid bacilli, of mouse typhoid II and the less toxic filtrates of two other types of mouse typhoid bacilli (I and III) were injected, separately and in combination, in mice bearing mammary carcinomas.

The highest incidence of local reaction, as well as the most intense effects, were seen in larger tumors after large doses

- 1 9 3 5 -

of the toxins. Mortality was higher. Partial or total inhibition of tumor growth was higher in the group bearing smaller tumors and receiving smaller doses of the toxins. It is believed that newly formed vessels are most susceptible to toxins.

226. CECIL, R. L.

Nonspecific protein therapy.

J.A.M.A. 105:1846-1854, 1935

The history of nonspecific protein therapy is reviewed. Included are the following sources of therapeutic preparations: native proteins; serums; protein-split products; enzymes; tissue extracts; bacterial extracts and vaccines. Differences between nonspecific reactions and focal processes are probed. The possibility that all types of 'protein' therapy may be specific in calling forth body defenses is presented. The therapeutic usefulness of leukocytosis is still unexplained. The author notes that gram-negative bacteria and their products cause febrile reactions more often than gram-negative organisms. Application of this type of treatment to specific disease conditions is evaluated.

227. GIEHM, G.

Die moderne Therapie der Schizophrenie (Modern therapy of schizophrenia)

Med. Klin. 31:547-549, 1935

A suspension of bacterial albumins (pyrifer) was used in the therapy of schizophrenia. Bacterial proteins from non-pathogens may prove efficacious in selected cases.

228. HEKTOEN, L.

Nonspecific proteins

J.A.M.A. 105:1765-1767, 1935

Foreign proteins united with body fluids are taken up by body cells, particularly those of the reticuloendothelial system; humoral and intracellular alterations follow. The products of these processes affect the entire organism in a complex fashion. Resultant phenomena are partially temporary in effect, partially permanent. The temporary effect is that exercised on the autonomic or sympathetic nervous system. An increase

in chemical activity accompanies fever. Capillary walls and cells of the reticuloendothelial system are important sites of chemical activity. Proteolytic and lipolytic enzyme output is accelerated. Fibrinogen and complement are decreased initially, then increased. The colloid stability of plasma is lessened. Altered metabolism is shown in the increased NPN of the blood. Histamine and choline are released as secretory stimulation follows dilatation in the splanchnic area. A local edema may be an outward manifestation of detoxification.

229. KOPP, I.

Metabolic rates in therapeutic fever

Am. J. Med. Sci. 190:491-494, 1935

Fever was induced by physical means and by mixed typhoid vaccine in 7 patients with neurosyphilis. One patient received fever therapy by all five methods employed. Various combinations of fever production were studied. Metabolic rates at fixed temperature levels were higher during periods of elevation than during periods of falling temperature. Rapidity of fever production did not affect the results; no differences in basal metabolism could be traced to the technique by which fever was produced.

230. MENZIES, A. W. C.

Pyretotherapy in Dementia Praecox

Lancet 2:994-996, 1935

Dmelcos (Ducrey bacillus atoxic vaccine) intramuscularly or sulfosin produced improvement in 26 schizophrenic patients and a complete remission for four of these. The recovery rate cannot be altered by this type of fever alone; accessibility of the patient to psychiatric treatment was increased and the psychic pattern was broken.

231. PINKSTON, J. O.

Peripheral circulation during experimental fever

Am. J. Physiol. 110:448, 1935

Sympathectomy delays initial vasoconstriction seen in peripheral vessels following the injection of typhoid-paratyphoid vaccine. In the normal, innervated ear of the rabbit, constriction begins

- 1 9 3 5 -

within 10 minutes after injection and persists until the fever maximum is attained. After sympathectomy, vasoconstriction is noted in 50 per cent of animals and, in the majority of these, delayed vessel constriction occurs. Inactivation of the adrenals may abolish the early responses of the vascular system. In the general peripheral vasodilation after febrile maximal points have been reached, the sympathectomized animals show a similar delay in vessel dilatation.

232. PINKSTON, J. O.

Experimental fever in sympathectomized animals

Am. J. Physiol. 111:539, 1935

Complete removal of sympathetic chains from the stellate to the pelvic ganglia brings a profound delay in both the initial febrile reaction and in the attainment of maximal temperature levels. The fever curves are flattened. With splanchnic section or adrenal inactivation only, the behavior of the curve in induced fever approaches that seen in the normal febrile animal.

233. SHWARTZMAN, G.

Hemorrhagic necrosis and regression of sarcoma 180

Science 82:201, 1935

The lethality of E. typhosa filtrates was much reduced when certain proportions of homologous antisera were combined with the material. Single intravenous injections of either antiserum-filtrate mixtures or filtrate alone were administered to sarcoma-bearing mice. The marked reduction in lethality which followed did not affect the hemorrhagic-necrotic action of the filtrates on tumors. Regression was prompt when a mixture of 200 neutralizing units of antityphoid serum was mixed with 300 reacting units of E. typhosa.

234. WALTER, C. W.

Economical intravenous therapy.

J.A.M.A. 104:1688-1790, 1935

Physiologic reactions ranging from fever and chills to cyanosis, diarrhea, nausea, vomiting and collapse have been seen after intravenous infusions of saline solutions contaminated from impurities in distilled water, chemicals rubber tubing and glassware. While individual susceptibility, velocity of injection,

- 1 9 3 5 -

its volume, and the temperature and pH of the solutions must be considered, pyrogens in water take first rank as causative agents in a reaction. The methods by which infusion materials may be rendered pyrogen-free are outlined in detail.

- 1 9 3 6 -

235. ANDERVONT, H. B.

The reaction of mice and of various mouse tumors to the injection of bacterial products

Am. J. Cancer 27:77-83, 1936

Bacterial filtrates obtained from meningococcus and B. coli cultures were used to determine their toxicity for mice, using changes in cutaneous tumors as a standard for judging activity of the filtrates. Toxicity was greater for tumor-bearing mice held at 37° C. than when room temperature was used; the same was true for the normal animal. Normal mice were more resistant to toxic filtrates than were the tumor-carrying animals. The route of injection was not important. Certain strain differences in susceptibility to toxic filtrates were shown. Spontaneous mammary carcinomas were resistant to B. coli filtrates while chemically-induced tumors responded with regression.

236. ANDERVONT, H. B. and SHEAR, M.J.

Production of Shwartzman reaction in rabbits with purified fraction of B. coli filtrate

Proc. Soc. Exper. Biol. Med. 34:673, 1936

The original filtrate and a purified fraction from B. coli were capable of eliciting the Shwartzman reaction and producing tumor hemorrhage in the mouse. The minimal effective tumor-hemorrhage dose of the original filtrate was 0.2 cc. of the 1:50 dilution. When the purified fraction was used, the minimal effective dose for tumor hemorrhage and that eliciting a positive Shwartzman were of the same magnitude.

237. CENTANNI, E.

Sulla presenza di un principio pirogene e antipirogene nella cultura batterica (On the presence of a pyrogenic and anti-pyrogenic principle in bacterial cultures)

- 1 9 3 6 -

Baglivi 2:241-253, 1936

238. CO TUI, McCLOSKEY, K. L., SCHRIFT, M. H. and YATES, A. L.

Filtration studies on reactive infusion fluids

Proc. Soc. Exper. Biol. Med. 35:297, 1936

The pyrogenic substance found in infusion fluids is of a particulate nature, with a diameter somewhat larger than 50 millimicrons.

239. FOGG, L. C.

Effect of certain bacterial products upon the growth of mouse tumor

Pub. Health Rept. 51:56, 1936

A small gram-negative bacillus was isolated as a contaminant of mouse sarcoma 180 tissue cultures. When injected intravenously into mice bearing sarcoma 180 growths, tumor regression appeared in 20 per cent of the animals, and hemorrhage and necrosis in 28 per cent. However, mortality reached 45 per cent and the use of immune sera did not alter the mortality rate. An alcohol precipitable fraction was prepared which, on injection, produced complete tumor regression in 42 per cent of the mice, and necrotic changes in 22 per cent. Mortality was reduced to 33 per cent. Addition of bacterial vaccine or this fraction inhibited in vitro tissue cultures of sarcoma 180 but did not injure normal tissues.

240. GIUFFRÉ, I.

Il trattamento ipertermico con 'pyrifer' nella terapia della schizofrenia e della demenza paralytica (Hyperthermic treatment with 'pyrifer' in therapy of schizophrenia and paralytic dementia)

Arch. gen. di. neurol., psichiat. e psicoanal. 17:223-246, 1936

Two patients were cured and one notably improved after injections of a bacterial protein secured from a nonpathogenic strain of B. coli. Temporary remissions of hebephrenic behavior followed therapy in two other instances. Three failures are reported. Malarial and sulfur injections were ineffective in these conditions.

241. JACOBI, M.

The effect of the Schwartzman reaction with bacterial filtrate on transplantable tumors in animals

Am. J. Cancer. 26:770-774, 1936

The Schwartzman phenomenon was provoked in experimental animals bearing mouse sarcoma 180, Flexner-Jobling rat carcinoma, Walker rat carcinosarcoma and Rous chicken sarcoma. Both preparatory and provocative injections of E. typhosa filtrate utilized the undiluted toxin. Preparatory injections were given intratumorally. All tumors responded with violent hemorrhage and necrosis after the provocative injection, complete sloughing and healing followed in some, slow regrowth of tumor tissue in others. When the injections were made only into tumor tissue or only intraperitoneally, no hemorrhage occurred.

242. MORGAN, W. T. J.

Studies in immunochemistry. I. The preparation and properties of a specific polysaccharide from B. dysenteriae (Shiga)

Biochem. J. 30:909-925, 1936

A polysaccharide containing 97 per cent reducing sugars on acid hydrolysis was isolated from the 'smooth' form of B. dysenteriae. Strongly dextrorotatory, it contained but 1 per cent of nitrogen, all in the form of acetylated amino groups. It accounts for the serological specificity of the bacterium. The occurrence and significance of certain nonspecific polysaccharides are discussed.

243. SHEAR, M. J.

Chemical treatment of tumors. IV. Properties of hemorrhage-producing fraction of B. coli filtrate

Proc. Soc. Exper. Biol. Med. 34:325-326, 1936

The hemorrhage-active fraction of B. coli filtrate previously described was further purified by methyl alcohol precipitation. The minimum dose for hemorrhagic activity was 0.2 cc. of a 1:500 dilution. After solution in water the fraction gave a strongly positive Molisch test and a negative biuret test; it exhibited no selective absorption peak. Activity was demonstrated after storage for 8 weeks in cold temperatures. In biologic tests, normal tissue capillaries were not rendered more permeable after injection of the fraction.

- 1 9 3 6 -

244. SHEAR, M. J. and ANDERVONT, H. B.

Chemical treatment of tumors. III. Separation of hemorrhage-producing fraction of B. coli filtrate

Proc. Soc. Exper. Biol. Med. 34:323-325, 1936

A method is described for the separation of the tumor-hemorrhage producing fraction of B. coli filtrate from toxic and inert contaminants. The method involves centrifugation, filtration, alkaline precipitation, and the removal of calcium by acidification with HCl and precipitation with alcohol. The final product was a water-soluble fraction, highly potent for the production of mouse tumor hemorrhage in amounts as low as 0.1 cc. of a 1:100 dilution of a stock solution containing 22 mg. per cc. of the active fraction.

245. SHWARTZMAN, G.

Reactivity of malignant neoplasms to bacterial filtrates. II. Relation of mortality to hemorrhagic necrosis and regression elicited by certain bacterial filtrates

Arch. Path. 21:509-524, 1936

Mice bearing 12-day-old sarcoma 180 implants were injected intravenously, intra-peritoneally or subcutaneously with one of the following bacterial filtrates: meningococcus "agar washings", B. enteritidis, B. coli, E. typhosa, Staphylococcus aureus, Streptococcus hemolyticus and the tubercle bacillus. The ability of the preparations to elicit the Shwartzman phenomenon in rabbits was determined shortly before parenteral injection.

E. typhosa and meningococcal filtrates were more toxic for the tumor-bearing mouse than for normal animals; filtrates of B. enteritidis were toxic for both normal and tumor-bearing test animals. All three were potent agents for tumor destruction. B. coli, Str. hemolyticus and tubercle bacillus filtrates produced only slight changes in tumor tissue. Staphylococcus aureus filtrate produced no changes. Since all filtrates were highly toxic, it was evident that the lethal effects and tumor-destructive activity exist independently. In certain given proportions, serum-E. typhosa mixtures protected animals against the lethal effects, while tumor hemorrhage, necrosis and regression proceeded unchanged.

- 1 9 3 6 -

246. WALTER, C. W.

Preparation of safe intravenous solutions

Surg., Gynec. & Obst. 63:643-646, 1936

Pyrogenic bacterial substances are the chief offenders in producing the reactions noted after intravenous therapy. Proper preparation of solutions and careful cleaning of transfusion or infusion equipment eliminates many of the hazards attending parenteral therapy.

247. WITEBSKY, E. and SALM, H.

Hemorrhagic necrotic skin lesions in rabbits produced by Hemophilus influenzae and Hemophilus pertussis

Proc. Soc. Exper. Biol. Med. 34:351, 1936

Skin areas of the rabbit, prepared with intradermal injections of living H. influenzae 24 hours before the intravenous injection of the same suspension or E. typhosa agar washings filtrate, responded with hemorrhage and necrosis within 3 to 6 hours. The response to intravenous H. influenzae exceeded the one noted after injection of H. pertussis. Sites prepared with H. pertussis were unchanged by intravenous challenging doses. Heat killed H. influenzae were likewise effective in producing hemorrhagic necrosis in sites previously prepared by the intradermal injection of the same organism or E. typhosa agar-washing filtrate.

- 1 9 3 7 -

248. ABERNETHY, T. J. and FRANCIS, T., JR.

Studies on the somatic C polysaccharide of Pneumococcus. I. Cutaneous and serological reactions in pneumonia

J. Exper. Med. 65:59-74, 1937

The intracutaneous injection of pneumococcal somatic polysaccharide produced a characteristic skin response in patients ill with pneumococcal lobar pneumonia. Certain patients ill with non-pneumococcal febrile diseases gave similar responses. The response was not present during convalescence. No skin reaction could be obtained in 7 fatal cases.

- 1 9 3 7 -

249. BOYLAND, E. and BOYLAND, M. E.

Studies in tissue metabolism. IX. The action of colchicine and B. typhosus extract

Biochem. J. 31:454-460, 1937

Agar washings of E. typhosa cultures were injected to determine the effect of tumor hemorrhage produced by such means upon tumor, liver and intestine ascorbic acid content. Colchicine was injected and its toxicity for normal and tumor-bearing mice, capacity for hemorrhage production in the transplanted tumor or the mouse and rat and tissue metabolic changes were determined, in vitro and in vivo. The effect of colchicine upon the ascorbic acid content of tumor, liver and intestine and its ability to produce the Schwartzman phenomenon in the rabbit were observed.

Both colchicine and E. typhosa agar washings were capable of producing tumor hemorrhage and reducing ascorbic acid content and metabolism in grafted tumors, but dosages necessary to initiate these effects were close to the toxic levels. Tumor-bearing animals were more sensitive to the toxic effects of colchicine than normal animals. The size of the tumor and the survival rate were significantly related. Ascorbic acid content of the rat and mouse intestine diminished after colchicine; liver metabolism was not inhibited. Colchicine did not elicit the Schwartzman phenomenon, nor did it produce hemorrhage in the liver.

250. CO TUI, McCLOSKEY, K. L., SCHRIFT, M. and YATES, A. L.

A new method of preparing nonpyrogenic intravenous infusion fluids, based on removal of pyrogen by absorptive filtration

Ann. Surg. 106:1089-1094, 1937

Adsorptive filtration, followed by ultrafiltration, is advocated for the preparation of pyrogen-free intravenous solutions. The sterile solution is passed through the compressed Seitz EK (serum) type asbestos filter and the 200-second Zsigmondy membrane filter. The Jena, Berkefeld, and 42-second Zsigmondy type filters are not effective for this purpose. It is believed that the pyrogen particle diameter lie between 50 millimicron and 1 micron.

251. CO TUI, SCHRIFF, M. H. and RUGGIERO, W. F.

A new method of preparing infusion fluids

J.A.M.A. 109:250-252, 1937

Filtration through adsorptive filters and a 200-second Zsigmondy filter will render saline and dextrose solutions pyrogen-free. The Seitz serum #3 was employed.

252. CO TUI, SCHRIFF, M. H. McCLOSKEY, AND YATES, A. L.

Filtration studies on pyrogenic inulin

Proc. Soc. Exper. Biol. Med. 36:227-230, 1937

Dahlia inulin, given intravenously, produced fever and leukopenia. The pyrogen particle resembles that found in infusion fluids and can be adsorbed by standard methods.

253. DENNIS, E. W. and SENEKJIAN, H.

Typhoid leucociden

Proc. Soc. Exper. Biol. Med. 36:61, 1937

The activity of a typhoid toxic fraction, leucociden, on normal rabbits' blood and non-immune human blood, was manifested by a reduction in the number of leukocytes present as the concentration of toxin was increased. Only the granulocytes were affected. Neutrophils which survived showed degenerative changes. The alcohol precipitate was stable for one year or more without loss of potency. An incomplete inactivation could be produced by heat. Specific immune globulin (Felix anti-typhoid serum) neutralized the toxin. One preparation of leucocidin contained 19.9 per cent of carbohydrate calculated as glucose.

254. FEVER THERAPY (A symposium)

New York, Paul Hoeber, Inc., 1937

Therapeutic fever production by physical and bacterial agents was considered by international authorities. A discussion of the antichancroid vaccine of Rimbaud was included.

255. GERBER, I. E. and BERNHEIM, A. I.

Histological studies of tumor reactivity to bacterial filtrates
Am. J. Pathol. 13:673, 1937 (Abstract)

A single intravenous injection of 20 to 100 reacting units of meningococcus 'agar-washing' filtrate produced severe degenerative changes in sarcoma 180 of the mouse. Vascular engorgement, edema and necrosis of the tumor cells preceded regression (or regrowth) of the tumor. Hemorrhage was seen in the early stages of change and vessel lesions and thrombi were absent. The filtrate appeared to affect tumor cells directly. The reaction "is analogous to and is elicited by the bacterial active principles of the Schwartzman phenomenon". No preparatory local injection was required.

256. HJORTH, P.

Investigations of a thermostable bactericidal substance in human serum, demonstrated particularly in the serum of fever patients

Acta Path. et microbiol. scand. 14:412-426, 1937

The serum of patients with febrile conditions contains the thermostable, bactericidal substance previously described by Wulff (J. Immunol. 27:451, 1934). Fever alone does not influence production of the bactericide. In undulant fever and influenzal pneumonia, the sera were not bactericidal, whereas inactivated sera from patients with cancer, cirrhosis and nephritis, in which temperatures were normal or only slightly increased, were high potent. This substance was active against meningococci, gonococci, and gram-negative cocci; Staphylococcus aureus growth was not inhibited. The serum did not lose potency completely until after exposure for one-half hour to 68° C. Minute amounts of active serum restored bactericidal effectiveness. The properties of the substance resemble those credited to beta-lysin by Petterson (Microben. (Jena), 1934).

257. MILLER, F. R.

The induced development and histogenesis of plasma cells

J. Exper. Med. 64:333-347, 1937

The mechanism by which the injection of tuberculo-protein stimulates plasma cells after intravenous or intraperitoneal injection was studied at length. Plasma cells were found in normal connective tissue. Genesis proceeds, it is believed, in lymphocytes or mesenchymal cells. The Russell body cells constitute

the end point in the evolution of these cells. The greatest concentration was found in those organs closely concerned with detoxification, as the omentum and lymph nodes, particularly after the injection of bacterial proteins. They are noted in spleen, liver, the body wall of the cecum. Polyblasts in the 'milkspots' of the omentum were precursors of these cells.

258. OLITZKI, L. and AVINERI-SHAPIRO, S.

The hypothermic factor of B. dysenteriae Shiga

J. Exper. Path. 18:316-321, 1937

In B. dysenteriae Shiga, and in other organisms, a hypothermic fraction may exist. Hyperthermic reactions may precede and follow the drop in temperature. The lipoid and carbohydrate fractions of the organism considered in this study produced temperature increases only; the hypothermic activity was closely associated with the protein fraction which may produce hyperglycemia and leucopenia.

259. OLITZKI, L., LEIBOWITZ, J. and BERMAN, M.

Further investigations on the chemistry, toxicity and other biological properties of different fractions of dysentery bacteria

Brit. J. Exper. Path. 18:305-316, 1937

Bacterial antigens, on injection, have the ability to produce hyperglycemia combined with a fall of inorganic blood phosphorus and followed by hypoglycemia and rising phosphorus index. The relationships of these alterations to the rise and fall of leukocytes were determined. Fractions obtained from B. dysenteriae Shiga, Flexner, and Y (fractions of Furth and Landsteiner) as well as the hapten were injected intravenously in test animals. Blood sugar and leukocytes were examined. Changes in body weight were recorded. Toxic symptoms were observed. A leucopenic factor was present in P1 and P2 fractions; a slight leucopenic effect was observed after administration of the C fraction. The hapten did not effect leukocytes. In P1, C and the hapten a glycemic factor occurs; it was present in unpurified polysaccharide 2 but not in the residue after alcohol extraction. P2 was strongly toxic, but fraction C and the hapten were non-toxic. The glycogen in liver tissue, galactose tolerance and diastase in blood were altered during the hyperglycemic reaction. Spleen and bone marrow changes took place during the period of leukopenia. Antileucopenic and anti-glycemic immunity existed independently.

260. PYRIFER

Gehees Codex, (Dresden), 1937, p. 1350

A temperature-increasing bacterial product from apathogenic colon bacteria grown in milk. It is marketed in seven strengths, ranging from 50 million to 500 million bacterial cells per cc. Three types of packaging are obtainable. It has been used extensively in paresis and other luetic conditions involving the central nervous system, but uses in other types of infection have been reported. The beginning dose is 0.5 to 1.0 cc. of strength I, with gradual increases to maintain desired reactions.

261. SHWARTZMAN, G., MORELL, S., and SOBOTKA, H.

Chemical investigations on the active principles of the phenomenon of local skin reactivity to bacterial filtrates I. Purification by dialysis and attempts at fractional precipitation

J. Exper. Med. 65:323-337, 1937

The active principles in the phenomenon of local skin reactivity (Shwartzman reaction) are retained by cellophane membranes. This method effects considerable purification, for culture filtrates of E. typhosa, B. coli and meningococcus, increasing reactive potency about three-fold. Fractional precipitation of concentrates of these filtrates, using acid, dilute alkali or alcohol, was not successful in this study.

262. WECHSLER, Z.

Der Einfluss der pyrifier fiebers auf die Blutliquorschranke im Tieversuche (Influence of pyrifier on plasma barriers in animal investigations)

Schweiz. med. Wochenschr. 67:1048, 1937

263. ALDER, A.

Fieberbehandlung der epidemischen Kinderlähmung mit Pyrifer
(Fever therapy with pyrifer in poliomyelitis)

Med. Klin. 34:1527, 1938

The development of pyrifer and of its application is discussed. The case histories appended would indicate some usefulness for such bacterial substances in the therapy of poliomyelitis.

264. CHASIS, H., RANGES, H. A., GOLDRING, W. and SMITH, H. W.

The control of renal blood flow and glomerular filtration in normal man

J. Clin. Invest. 17:683-697, 1938

The diodrast, inulin and phenol red clearance of the human renal tubule was studied under a variety of experimental conditions. The physiologic regulation of renal blood flow and glomerular filtration rate was examined during the pyrexial reaction induced by intravenously administered typhoid vaccine, during water diuresis, and when certain stimulants and diuretics were administered. The action of typhoid vaccine was initiated with a rise in blood pressure during the period of chill accompanied by efferent arteriolar constriction and decreased renal plasma flow. The renal ischemia gave way to hyperemia in the febrile period; the filtration fraction increased, indicating a dilatation of efferent structures. The filtration fraction varies inversely to the renal blood flow, while the filtration rate tends to be independently maintained.

265. CONWAY, E. A.

Reaction of lymphatic tissue in early stages of Bacterium monocytogenes infection

Arch. Path. 25:200, 1938

After infection with B. monocytogenes a rapid mobilization of the pre-existing lymphocytes takes place. Intensified production in lymphatic nodules of the mesenteric lymph node and splenic periarterial tissue begins. The free stem cell from which the monocyte develops in diffuse lymphatic tissue, the mesenteric node and spleen, is morphologically identical with lymphocytes found in these tissues.

- 1 9 3 8 -

266. GERBER, I. E. and BERNHEIM, A. I.

Morphologic study of the reactivity of mouse sarcoma 180 to bacterial filtrates

Arch. Path. 26:971-983, 1938

In the mouse bearing growing tumor implants, the injection of meningococcus filtrate, capable of producing local reactivity, produced hemorrhage, engorgement and edema in tumor tissue within 4 hours after administration. Cellular necrosis followed, but capillary rupture, thrombosis and other vascular changes were not seen. The amount of degeneration noted in tumor tissue appeared to be a function of the amount of filtrate injected. Large doses (100 rabbit-reacting units) brought complete sloughing of malignant tissue with healing. With smaller dosages, the undestroyed residuum contained peripheral neoplastic cells, and regrowth was more commonly seen. Carcinoma was unaffected, and some hemorrhage and necrosis took place in tumors produced by dibenzanthracene.

267. GORO, M.

Action of various substances on the secretion of gastric juice

Jap. J. Med. Sci. (IX) 5:341-342, 1938

The oral administration of a 10 per cent solution of either mono- or polysaccharide diminished gastric secretion in the dog. A weaker polysaccharide solution, averaging from 1 to 2 per cent, stimulated the secretory powers of the mucosa.

268. KOSTYAL, L.

Wirkungsweise der Eiweisskorper (Foreign protein action)

Ztschr. Immunitats. 92:318-335, 1938

The antitoxic action of the blood in diphtheria was increased by intramuscular injections of omnadin. Pyrifer and milk, similarly administered, reduced such action. The activity is related to an alteration in colloid stability and of the globulin fractions, particularly pseudoglobulin I and II. Euglobulin may enter into such activities.

269. PASCHKIS, K. and MARTOS, K.

(The central nervous regulation of the blood)

Arch. Int. pharmacodyn. 58:432-436, 1938

Phenobarbital, administered before pyrifer, prevents the expected leukocytosis. If given before nucleic acid, phenobarbital

- 1 9 3 8 -

depresses leukocyte genesis and produces a leukopenia. Both pyrififer and nucleic acid stimulate leukocyte production.

270. RANSOM, S. W. JR.

Fever induced by the intravenous injection of typhoid-paratyphoid vaccine

Arch. Int. Med. 61:285-296, 1938

The reaction of the normal cat to intravenous typhoid-paratyphoid vaccine, is one of cyclic stimulation, first, heat conservation, sympathetic stimulation and heat production and, secondly, a period of heat loss and parasympathetic stimulation. The cycle of these alternate reactions is repeated three times. The heat preservation mechanism predominates, and the end result appears to be sustained hyperthermia. Anaesthesia appears to affect either the mechanism of heat preservation and loss or a coordinating control of these phenomena; etherization of cats before vaccine injection results in hypothermia and parasympathetic response to the bacterial material. In typhoid fever, intravenous typhoid vaccine brings forward similar mechanisms.

271. ROBERTSON, R. C. and YU, H

Leukopenia and the toxic substances of B. typhosus

J. Hyg. 38:299, 1938

The intraperitoneal or intravenous injection of toxic filtrates from several strains of E. typhosa provoked, in the rabbit given sublethal doses, sharp leukopenia. The leukocytosis which followed persists two days. In vitro mixtures of the filtrate and blood produced no pronounced leukocyte changes. Histologic sections from lung, liver, spleen and bone marrow displayed an accumulation of neutrophils in these organs.

The constant liberation of toxic substances in human typhoid fever is believed to be responsible for the leukopenic response.

- 1 9 3 8 -

272. SEVAG, M. G., LACKMAN, D. B., and SMOLENS, J.

The isolation of the components of streptococcal nucleoproteins in serologically active form

J. Bio.. Chem. 124:425-436, 1938

From hemolytic streptococci of Lancefield group A, new antigenic substances, designated labile antigens, were isolated. Both nucleic acid and protein have been liberated from the complex structure, through hydrolysis, chloroform extraction, and a foam-reducing agent.

273. WIEN, K.

Glucose tolerance and lactate utilization during fever

Quart. J. Pharm. Pharmacol. 11:34-45, 1938

When fever was induced in rabbits through subcutaneous injections of a B. coli vaccine, oral glucose tolerance was lowered. Peak levels of blood sugar far exceeded those normally present. After glucose infusions, initially high levels appeared, but normal blood sugar values quickly returned. Lactate utilization was impaired in such febrile states, and the ability of the liver to form glycogen after the administration of sodium dl-lactate was inhibited. Insulin injections did not increase glycogen reserves in the liver.

274. ANDERSON, C. G., HAWORTH, W. N., RAISTRICK, H. and STACEY, M.

Polysaccharide synthesized by microorganisms. IV. The molecular content of luteose

Biochem. J. 33:272-279, 1939

Luteose, a neutral polysaccharide remaining after removal of the malonic acid content of luteic acid, is composed of beta-glucose units linked through the 1,6-positions. The molecular structure may be of the closed-chain type. Luteic acid is a metabolic product of penicillin luteum (Zukal).

275. ANDERVONT, H. B. and SHIMKIN, M. B.

The effect of ascorbic acid upon the hemorrhage produced by bacterial filtrate in transplanted tumors

Am. J. Cancer 36:451-459, 1939

Minute intraperitoneal injections of a bacterial filtrate of B. prodigiosus into tumor-bearing mice (Sarcoma #37) were followed by intracutaneous injections of neutralized or unneutralized ascorbic acid. Similar experiments tested the effects of ascorbic acid and bacterial filtrate in combination against Sarcoma 180 and Lung Tumor F transplants. Ascorbic acid prevented the appearance of hemorrhage and regression of tumors after injection of the filtrate from B. prodigiosus. Acidity was unrelated to hemorrhage prevention, but reducing power of ascorbic acid was apparently involved in the phenomenon. Hemorrhage inhibition was directly proportional to the amounts of filtrate and ascorbic acid used. The route of administration appeared to be unimportant. A theory is advanced that the bacterial filtrate may produce hemorrhage in tumor tissue through sudden lowering of tumor ascorbic acid content, thus weakening the capillaries and permitting hemorrhagic extravasation.

276. CO TUI, BENAGLIA, A. E., RUGGIERO, W. F. and YATES, A. L.

Study of the fever-producing principle in the typhoid vaccine

Proc. Soc. Exper. Biol. & Med. 41:272-275, 1939

There is a close relationship between the fever-producing substance found in vaccines from B. typhosus cultures and pyrogens isolated from contaminated waters and infusion solutions. Initial leukopenia, rise in temperature and chills are identical; either may produce transient gastrointestinal disturbances.

- 1 9 3 9 -

277. CO TUI and SCHRIFT, M. H.

Production of pyrogen in sera by bacteria

Proc. Soc. Exper. Biol. & Med. 42:549-552, 1939

Normal horse serum was inoculated with cultures of Pr. vulgaris, B. coli, Staphylococcus aureus, B. subtilis and two unidentified organisms isolated from water. Following incubation periods of varying duration, the culture fluids were filtered and injected intravenously in the dog. A pyrogenic symptom-complex which included fever, leucopenia, occasionally chills, and gastrointestinal disturbances, was noted. Pr. vulgaris, B. coli and Staphylococcus aureus filtrates produced severe reactions. Pyrexia was insufficient after B. subtilis injections, but leucopenia was present. "On the basis of these experiments it is suggested that the chill and fever-producing substances in sera may be pyrogens, caused either by direct bacterial growth, or by pyrogen already present in chemicals used in the purification and concentration of therapeutic sera."

278. CO TUI, SCHRIFT, M. H. and RUGGIERO, W. F.

Production of pyrogen in gum acacia by bacteria

Proc. Soc. Exper. Biol. & Med. 41:533-535, 1939

Pyrogen-producing bacteria (B. subtilis, two unidentified water organisms and Staphylococcus albus) grown in gum acacia solutions produce an "active" acacia capable of eliciting febrile responses after injection. Leucopenia and other changes seen with reactive inulin and typhoid vaccine follow.

279. DENNIS, E. W. and SENEKJIAN, H.

A leucocidal toxin extracted from typhoid bacilli

Am. J. Hygiene 30:103-11, 1939

A crude extract of typhoid bacilli presents two active factors: a leukocidal factor with selective lethal effect on the pseudo-eosinophilic granulocytes and a toxic somatic antigen. The former loses activity on purification by trichloroacetic acid, but the latter increases in potency during the same process. Leukocidin can be demonstrated in extracts of S. typhimurium, S. schottmulleri, S. suispestifer (voldagsen strain), S. enteritidis, S. gallinarum and S. paratyphi, listed in decreasing order of activity.

280. GARDNER, R. E., BAILEY, G. H. and HYDE, R. R.

Hemorrhagic activity of toxic carbohydrate complexes from bacteria on a transplantable rat tumor

Am. J. Hygiene 29B:1-14, 1939

The theory has been presented that generalized vascular shock following injection of bacterial filtrates or specific substances initiates tumor hemorrhagic processes. Such materials were injected into rats bearing transplanted Walker tumor No. 256, all cutaneous, and approximately 10 days old. Controls were used for comparison. Non-bacterial carbohydrates and a yeast were also injected.

The yeast carbohydrate produced acute systemic reactions and a hemorrhage in tumors which differed from that occurring after use of bacterial filtrates. A polysaccharide-free component of the soluble specific substance from S. enteritidis also had activity and produced a severe reaction. Most marked results were obtained with fractions isolated from the cultures of enteric organisms, particularly those endogenous in the rat. However, other bacteria, including K. pneumoniae, had distinct activity.

281. HARKAVY, J. and ROMANOFF, A.

Local hemorrhagic-necrotic skin reactions in man (Shwartzman phenomenon)

J. Allergy 10:566, 1939

A comparison is made between mechanisms of the Shwartzman and Arthus phenomena, and three case histories are presented for illustration. Each presented skin lesions fulfilling all criteria for Shwartzman reactions. The provocative factor which precipitated lesions may be either a parenterally introduced antigen from a bacterium or a product derived by an interaction between an injected foreign protein and a preformed homologous antibody.

282. IVANOVICS, G.

Chemische Untersuchungen über de Polysaccharide des Milzbrand-bacillus (Chemical studies in the polysaccharides of the anthrax bacillus)

Zentr. Bakt. parasitenk. I. Abt. 144:244-246, 1939

The water soluble precipitate produced after concentration and purification of a culture of avirulent anthrax bacilli contained

- 1939 -

0.6% nitrogen and 60% reducing substances.

283. IVANOVICS, G.

Das Schicksal der Kapselsubstanz der Milzbrandbacillen im Organismus (Fate of the anthrax bacillus capsular substance in the organism)

Ztschr. f. Immunitätsforschung, u. Exp. Therap. 96:408-412, 1939

284. IVY, A. C. and GRAY, J. S.

The application of recent contributions in basic medical sciences to surgical practice. Reactions to intravenous therapy

Inter. Abstr. Surg. 69:414-416, 1939

A brief review of major milestones in the literature of pyrogens since 1911.

285. MASUCCI, P.

Antigenic polysaccharide complex from cocci

U. S. Patent #2,166,963, July 25, 1939

A method of extracting and precipitating antigenic polysaccharides from cocci.

286. MIELNIK-KOSMIDERSKI, S.

Über Der Einflusung der Phagozytose durch die Bakterienkapsel und Bakterielle Polysaccharide (Influence of phagocytosis of bacterial capsule and bacterial polysaccharide)

Ztschr. f. Immunitätsforschung, u. Exp. Therap. 95:203-217, 1939

Free or somatic polysaccharide from the capsule of scleroma bacteria enhances phagocytosis of unencapsulated bacteria to a degree limited by the number of bacteria and of leukocytes. Very small amounts of polysaccharides decrease phagocytosis below normal levels. The amount of polysaccharide released around bacteria by antibody may increase (or decrease) phagocytosis.

287. MORGAN, H. R. and BECKWITH, T. D.

Immunological relationships of polysaccharides of mucoid organisms of typhoid-salmonella groups

J. Bact. 37:389-399, 1939

Protein-free polysaccharide extracts from E. typhosa, S. paratyphi, A, B. and C., S. enteritidis, aertrycke, sulpestifer or S. pullorum, gave rise to precipitins for polysaccharides and agglutinins for the organisms after testing with antisera produced in the rabbit. Both immunized and non-immune animals were inoculated intracutaneously with the polysaccharide, and sterile abscesses were produced in both.

288. NELSON, C. M.

The cause of chills following intravenous therapy

J.A.M.A. 112:1303-1306, 1939

Extremes in temperature of solutions intended for venoclysis, the amount of solution, rate of administration, concentrations, preparation of glass and tubing of equipment and pH were all eliminated as factors in producing rigors after intravenous administration. Contamination of distilled water, when immediate autoclaving is neglected, was the primary cause of reactions. Aging alone in properly autoclaved solutions does not produce reactive solutions.

289. RANSON, S. W., JR., CLARK, G. and MAGOUN, H. W.

The effect of hypothalamic lesions on fever induced by intravenous injection of typhoid-paratyphoid vaccine

J. Lab. & Clin. Med. 25:160-168, 1939

Injection of typhoid-paratyphoid vaccine in the cat stimulates both the mechanisms for heat conservation and those governing heat loss. The stimulation of the sympathetic system was expressed by febrile reactions in the normal cat; with chronic lesions of the hypothalamus prolonged temperature reduction, enduring as long as twenty hours, followed the injection of typhoid vaccine and this was accompanied by other signs of parasympathetic overactivity (defecation, urination and salivation). The portion of the heat loss mechanism which is stimulated by pyrogen lies caudal to the hypothalamus.

290. SMITH, E. V.

The effects of injections of toxic extracts of the typhoid bacillus on the blood picture in rabbits

Am. J. Hygiene 29:15-23, 1939

- 1 9 3 9 -

Extracts from the cappola strain of the typhoid bacillus were injected intravenously in the rabbit in dosages ranging from sublethal to highly toxic amounts. Single and repeated injections were administered; the extract was also prepared with fresh normal, inactivated normal and immune serums. The severe systemic reactions, extreme leucopenia and tendency toward prolonged coagulation time were observed after typhoid extract administration. The systemic reactions were dependent upon the dose and independent of the dose for blood changes. The systemic effects appear to become less intense after repeated injection; no complete immunity could be demonstrated, and the changes in blood cellular components were not effected. Mixtures of serum and toxic typhoid extract produced symptoms characteristic of small sublethal doses of the extract when given alone.

291. SMITH, H. W.

Physiology of the renal circulation

The Harvey Lecture Series 35:200-203, 1939-1940

In a review of known facts on renal circulation, it is stated that the only known method of inducing renal hyperemia is by the use of pyrogens "either as present in typhoid vaccine or in pyrogenic inulin." Hyperemia is manifest at the time of the chill phase and continues for several hours. The relative increase depends upon the control level and is "...in general... from 36 to 60 per cent above the control levels". Antipyretics have little influence. The dilatation is centered in the efferent arteriole; the filtration fraction is decreased, but the rate of filtration is essentially unchanged.

292. TERRY, G. G.

Fever and Psychoses

New York: Paul B. Hoeber, Inc., 1939

Biologically induced fever (through the use of organic, bacterial and "foreign" agents) produces some evidence of improvement in manic-depressive states, but only a few benefits in schizophrenics.

It has been stated that Coley's fluid has therapeutic usefulness in the manic-depressive states.

293. WALKER, E. L. and HANDMAN, M. S.

Biological reactions of malignant neoplasms. I. The reaction to bacterial fractions

J. Immunol. 37:507-524, 1939

After a brief review of the use of bacterial products in the therapy of malignant neoplasms and of methods for their preparation, details of a comprehensive study of the fractions obtained from Salmonella typhimurium are appended.

Crude fractions, nucleoproteins, nucleic acid and polysaccharides were tested as preparatory or provocative agents in the Shwartzman reaction; all were found suitable for either rôle. Several strains of transplanted sarcoma or carcinoma were allowed to multiply for 10 to 21 days in the rat; intra-abdominal injections of the bacterial fractions were then administered in 1 to 6 doses at the tolerated levels. There were intervals between doses which ranged from 3 to 7 days. Tumor hemorrhage, necrosis and regression were seen in Sarcoma 180 on injection of nucleoproteins polysaccharides or fractions containing one or both of these constituents. In a maximum percentage of rats sublethal doses of these active fractions produced total regression. Few carcinoma strains underwent complete regression without recurrence. The authors do not find a single special bacterial substance responsible for activity in the Shwartzman phenomenon and against malignant neoplasms, but find that the ordinary proteins and carbohydrates of the bacterial cell carry the hemorrhagic-necrotic functions.

- 1 9 4 0 -

294. ANDERVONT, H. B.

Effect of colchicine and bacterial products on transplantable and spontaneous tumors in mice

J. Nat. Cancer Inst., 1:361-6, 1940

This study was undertaken to compare the effects of bacterial products and colchicine in inducing tumor hemorrhage and affecting tumor growth. Young mice bearing either Sarcoma 37 or Carcinoma F as transplantable tumors or mammary carcinoma as a spontaneous tumor, were given colchicine solutions or B. coli filtrate, in varying dosages, intraperitoneally. The effective dose which elicited tumor-hemorrhage without killing the animals was determined by injection of single doses at various levels of potency. Injections of colchicine in varying doses were repeated at differing time intervals. In order to determine whether or not colchicine and filtrate have a supplementary effect on each other, separate injections were made of each substance at practically the same time.

The effective dose of colchicine, 0.1 mg. was close to the lethal dose and was no more effective than bacterial filtrate which has a much wider range of potency. Repeated injections of colchicine in doses too small to elicit tumor hemorrhage had no influence on tumor growth; while injections of hemorrhage-producing doses were toxic to the majority of animals. Both substances were supplementary to each other in eliciting hemorrhage in transplanted tumors, but their combined action was no more effective than either alone in treatment of spontaneous tumors. Tumor F, after the 17th serial passage, became susceptible to hemorrhage induction by bacterial filtrate.

295. ASHENBURG, N. J., SANDHOLZER, L. A., SCHERP, H. W. and BERRY, G. P.

The influence of bacterial and nonbacterial polysaccharide upon bacteriophagy

J. Bact., 39:71-72, 1940

In an investigation of the effect of capsular polysaccharide (from Aerobacter) upon the activity of a bacteriophage to which strains of Aerobacter are susceptible, inhibition of lysis was noted. Non-bacterial polysaccharide (starch glycogen and gum arabic), and polysaccharide from a bacteriophage-resistant strain inhibited lysis to an equivalent degree. "The results did not support the hypothesis that susceptibility to bacteriophagy depends upon immunological specificity."

296. BEESON, P. B. and HOAGLAND, C. L.

The use of calcium chloride in the treatment of chills

N. Y. State J. Med. 40:803-804, 1940

Intravenous administration of calcium chloride in humans aborts the chills due to malaria, blood transfusions, and typhoid vaccine. Relief of the typhoid chill appeared to have no effect on the subsequent elevation of the body temperature.

297. CAMERON, G. R., DELAFIELD, M. E. and WILSON, J.

Pathological changes produced in rats and mice by a toxic fraction derived from Bacterium typhimurium

J. Path. Bact. 51:223-233, 1940

A non-dialyzable protein-polysaccharide fraction from B. typhimurium was injected intraperitoneally, in doses ranging from 0.05-1.0 mg. into 95 mice and 54 albino rats. The animals were sacrificed at $\frac{1}{2}$ hour to 6 days following injection, and various tissues were stained for microscopic study. Forty mice were injected with living B. typhimurium and their livers examined microscopically at intervals varying from 1 to 22 days following the initial injection. Blood sugar and hepatic glycogen content were also determined in animals injected with the bacterial fraction.

A profound disturbance of the carbohydrate metabolism was shown by the histological and chemical results. The liver showed marked portal congestion associated with hemorrhage, thrombosis, and necrosis; the spleen and lymph glands exhibited necrosis of Malpighian bodies and lymphoid follicles. Infection of the animals with living B. typhimurium produced similar results of more severe degree.

Liver glycogen in rats and mice, and blood sugar in mice were rapidly decreased in amount following initial injection. Carbohydrate disturbances are primarily of hepatic origin.

298. CENTANNI, E.

Weitere Beitrage zur Kenntniss des pyrogenen Wirkstoffs des Fiebers (Further studies toward knowledge of active pyrogenic materials of fever)

Deutsch. med. Wchnschr. (Anal. Series) 10:263-265, 1940

The chemical character of the pyrogenic active material formed in the decomposition of bacteria was studied. It is reported that

- 1 9 4 0 -

the pyrogen is not a polypeptide or an organic base. It is insoluble in fat solvents and not reduced by common hydrolytic enzymes. Stability during autoclaving or treatment with strong acids has been confirmed. It is divisible by specific precipitants and adsorption agents, but is non-dialyzable.

299. DE RENZI, S. and GRASSELINI, A.

Blood sugar in provoked hyperthermia (Translated Title)

Boll. soc. ital. biol. sper. 15:802-803, 1940

Hyperthermia induced by injection of killed bacteria increased the blood sugar in both normal and diabetic individuals. Glucose or galactose administered during fever brought blood sugar to peak levels. During hyperthermia, adrenalin and insulin produced only insignificant changes in blood sugar.

300. EHELSON, L. M.

The inability of desoxycorticosterone acetate to protect the adrenalectomized rat against typhoid vaccine.

Endocrinology, 27:340, 1940

Since adrenalectomized animals are protected against typhoid vaccine by cortico-adrenal extracts, it was of interest to note whether synthetic corticoids were equally efficacious. Wistar rats were adrenalectomized, and on the day after operation were given corticosteroids; 20 rats were given 2 mg. of desoxycorticosterone acetate daily; 20 rats were given 2 cc. of adrenal cortex extract daily; 4 rats were given 4 mg. of the synthetic daily. After 5-10 days of such treatment, typhoid vaccine was given intraperitoneally in doses varying from 0.5 to 0.8 cc. Treatment with adrenal substance was continued in those animals that survived an observation period of 3 weeks. A few of the animals in each group were given second vaccine injections 24 to 48 hours after the first. The synthetic product is unable to protect the adrenalectomized rat against typhoid vaccine, while the natural substance increases the resistance of such animals.

301. FREEMAN, G. G., CHALLINOR, S. W. and WILSON, J.

The use of a synthetic medium in the isolation of the somatic antigens of Bact. typhimurium and Bact. typhosum.

Biochem. J. 34:307-324, 1940

Antigenic fractions from B. typhimurium and B. typhosum have been prepared by 3 methods: (a) a development of the tryptic digestion method of Raistrick and Topley, (b) Boivin's method of extraction with cold trichloroacetic acid, and (c) extraction with organic solvents such as ethyleneglycol and diethyleneglycol (Morgan, 1937) and the products have been compared by chemical analysis and determination of toxicity in mice. The purest products obtained by these methods show a similarity of properties. It may be concluded that, by any of these methods, the essential antigenic complex of the bacteria may be obtained in a state approaching chemical homogeneity.

On the grounds of yield and suitability for use in extracting large quantities of bacterial bodies, the tryptic digestion method and Boivin's method are to be recommended.

302. GRAY, J. S., CULMER, C. U., WIECZOROWSKI, E. and ADKISON, J. L:

Preparation of pyrogen-free urogastrone

Proc. Soc. Exper. Biol. and Med. 43:225-228, 1940

Extracts of normal human male urine were prepared by 3 methods: the benzoic acid adsorption procedure of Katzman and Dolisy; further purification procedures involving neutral acetone fractionation. Assay of the gastric inhibitory factor was performed in 6 Heidenhain pouch dogs injected subcutaneously with histamine; rectal temperatures and percentage inhibition of free acid secretion were recorded after injection of the extract. Pyrogen assays were also made by recording rectal temperature curves for a group of 5 rabbits.

The pyrogenic effect of the extracts was progressively lost with the use of the more purified materials, though gastric inhibition was not materially affected. Utilizing the acid acetone fractionation procedure, the gastric inhibitory factor was obtained free from pyrogen.

"The results indicate quite clearly that the gastric inhibitory factor is separate and distinct from the pyrogenic factor in extracts of normal male urine. Until such time as the gastric inhibitory factor is definitely identified, we propose to call it urogastrone."

303. GRAY, J. S., WIECZOROWSKI, E., CULMER, C. U. and ADKISON, J. L.

The presence of pyrogen in urine and its separation from urogastrone.

Amer. J. Physiol., 129:589-596, 1940

- 1940 -

This study was undertaken to determine whether the gastric inhibition produced by urine extracts was due to pyrogenic substances. Six Heidenhain pouch dogs were used for assay of the gastric inhibitory factor under controlled conditions; the potency of the injected material was measured by percentage decrease in mgm. of free acid secreted. Pyrogen was assayed by rise in rectal temperature of the pouch dogs and in rabbits. Crude urine extracts were prepared by several modifications of the benzoic acid adsorption method of Katzman and Doisy (methods given). Purification of the crude extracts was accomplished through fractionation with acetone.

It was found that the major portion of the pyrogen in the crude extracts originated in bacterial contamination; measures which minimize bacterial growth permit preparation of crude extracts containing no more pyrogen than equivalent amounts of fresh urine. Fresh, untreated urine from healthy human subjects was found to contain small and variable amounts of pyrogenic substance. Purification of crude extracts relatively free of pyrogen consistently yielded products exhibiting high urogastrone and little or no pyrogenic potency.

It was concluded that urogastrone is not pyrogen. In disease the quantity of pyrogen in the urine is increased.

304. LEES, J. C. and LEVY, G. A.

Emergency preparation of pyrogen-free water

Brit. Med. J., 1:430, 1940

A method is described for removing pyrogens from contaminated water, thus rendering it suitable for intravenous use. The method consists of shaking the water with powdered charcoal (1 part per 1,000) for fifteen minutes, then allowing to stand for several minutes after which the charcoal is readily removed by decanting through a filter paper.

305. MORGAN, H. R.

Preparation of antigenic material inducing leucopenia from Eberthella typhosa cultured in a synthetic medium

Proc. Soc. Exp. Biol. Med. 43:529-532, 1940

Small amounts of antigenic substance isolated from cultures of E. typhosa induce fever and leukopenia in the rabbit while larger amounts are lethal to both mice and rabbits. Leukopenia was the result of the disappearance of neutrophils from the blood. The toxic properties of the antigen remain after 30 minutes boiling.

306. MORGAN, W. T. J. and PARTRIDGE, S. M.

Studies in immunochemistry - the fractionation and nature of antigenic material isolated from Bact. dysenteriae (Shiga)

Biochem. J. 34:169-191, 1940

A polysaccharide, a phospholipin and a polypeptide-like component were isolated from the antigenic complex of "smooth" (Shiga) B. dysenteriae. Dissociation of the complex is described and a hypothetical structure suggested.

307. NISI, NAOTO

Physiology and pathology of the hypophysis temperature-regulatory function

J. Chosen Med. A. 30:1-21, 1940

In rabbits, completely hypophysectomized or with only the anterior lobe removed, central pyrogenic stimulants do not act, but peripheral stimulants like tetrahydro-beta-naphthylamine or adrenaline are effective, although the response gradually decreases. The abnormal responses return to normal if the anterior lobe is implanted or its extract injected.

308. OMITI, M.

The influence of artificial fever on the nonspecificity in the syphilis reaction. III. Experiment with pyrifer.

Mitt. med. ges. Okayama 52:2413-2432, 1940 (English version)

A comprehensive study of the effects of pyrifer on the globulin-albumin alterations in serum.

309. PALMER, J. W. and GERLOUGH, T. D.

A simple method for preparing antigenic substances from the typhoid bacillus

Science 92:155-156, 1940

The capsular polysaccharides of pneumococci may be separated from their associated proteins by extracting the latter with phenol, in which polysaccharides are insoluble. Fractional precipitation of polysaccharides by alcohol or glacial acetic acid may follow. After drying with acetone, a white powder is obtained (substance A). Phenol extraction may likewise be used for the product obtained by tryptic digestion; the product (substance B) resembles substance A in both appearance and antigenic properties.

- 1940 -

Both substances, obtained in yields of 10 to 15 per cent of the weight of dried typhoid organisms, give negative or faint biuret reactions and a strong Molisch reaction. Nitrogen content, corrected for ash and moisture, is from 3.4 to 4.6 per cent. Hexosamine, if present, accounts for less than 10 per cent of total N. Both substances give strong precipitin reactions with antityphoid rabbit serum.

310. PARTRIDGE, S. M. and MORGAN, W. T. J.

Immunization experiments with artificial complexes formed from substances isolated from the antigen of Bact. shigae

Brit. J. Exp. Path. 21:180-195, 1940

A specific polysaccharide component of the antigen of B. shigae failed to induce immune bodies in the rabbit. The polypeptide component of the same complex produced low titer homologous precipitins but no agglutinins. A synthetic 'complex' built from these components in combination induced the formation of potent immune bodies, agglutinated B. shigae at high titer and precipitated with the specific polysaccharide. The complex fixed complement when Shiga antibacterial sera was added.

311. SHAPIRO, C. J.

The effect of toxic carbohydrate complex from S. enteritides on transplantable rat tumors in tissue culture

Am. J. Hygiene, 31:Sec. B. 114-26, 1940

Tissue cultures of rat sarcoma #10 or Walker carcinoma #256 were maintained in vivo in healthy rats as intradermal, subcutaneous or intramuscular growths. On plasma growth medium they were exposed to a toxic carbohydrate complex derived from S. enteritides, capable of causing tumor hemorrhage and necrosis in doses as small as 0.0005 mg. Control tissue tumor cultures were maintained in the growth medium and saline. Doses of 2000 to 5000 provocative units of polysaccharide were used in all in vivo experiments. No effects were noted in vitro. It is believed that the plasma of a tumor-bearing host does not interact with the hemorrhage-producing agent to form a substance toxic to either young or sensitive cells of these neoplastic growths.

312. SHEAR, M. J. and TURNER, F. C.

Studies on the chemical treatment of tumors. V. Separation of the hemorrhage-producing fraction from Bacillus prodigiosus filtrates

J. Biol. Chem., 133:Lxxxvii, 1940 (Abstract)

Methods which had been used to separate hemorrhage-producing fractions from B. coli were applied to broth cultures of B. prodigiosus. An untreated broth culture filtrate contained 330 mouse tumor units per cc. of active principle. In a commercial sample of Coley's mixed toxins, 100 units per cc. were present. Further treatment of the above filtrate gave a concentration which contained 20,000 mouse units per cc. In studying growth on simple synthetic media, it was noted that filtrates had the same potent quality as those from broth cultures. On treatment with CHCl_3 , an emulsion was obtained which possessed most of the activity of the filtrate. A water-soluble precipitate remained after CHCl_3 was removed and $\text{C}_2\text{H}_5\text{OH}$ employed. A solution of this concentrate had a potency of 133,000 mouse units per cc. and contained 0.2 gamma per mouse tumor unit. Lethal dosages of these concentrates ranged from 100 to 1000 times the minimum effective dose. The active principle was nondialyzable through cellophane and was heat-stable. Large scale preparation techniques are given in detail.

313. SUTHERLAND, A. M. and WOLFF, H. G.

Experimental studies on headache: Further analysis of mechanism of headache in migraine, hypertension and fever

Arch. Neurol. and Psychiat. 44:929-949, 1940

In twelve patients with chorea or rheumatoid arthritis, fever therapy was induced by intravenous typhoid vaccine. A tambour was placed on the temporal artery and the pulsations recorded. Photographic records of the pulsations of the cerebrospinal fluid were made from a needle introduced into the lumbar sac and connected to a Frank capsule. Control records of temporal artery and cerebrospinal fluid pulsations were made simultaneously. Then an appropriate dose of typhoid vaccine was given intravenously, and estimates of the degree of headache, determinations of blood pressure, and pulsations of temporal artery and cerebrospinal fluid were made at frequent intervals.

Increased amplitude of pulsations of temporal artery and cerebrospinal fluid in all instances was found to be followed by onset of headache or sensation of head fullness. Exacerbations and remissions of the headache were found to closely parallel the increase or decrease in amplitude of pulsations. The headache ended with the ultimate decrease in amplitude. These findings are similar to those seen in histamine headache. Increasing the intracranial pressure decreased the intensity of the headache.

- 1 9 4 0 -

314. WALTER, C. W.

The relation of proper preparation of solutions for intravenous therapy to febrile reactions

Ann. Surg. 112:603-625, 1940

While the bacterial pyrogens of water, apparatus and rubber tubing are believed to be the principal causes of febrile reactions during intravenous therapy, in dextrose solution the dehydration products should also be eliminated if possible. Such products of molecular interaction include hydroxymethylfurfural and levulinic acid. The alkaline film following sterilization of soft glass may cause some polymerization of dextrose during sterilization. The intravenous kit itself should be carefully sterilized if re-use contemplated.

315. ABERNATHY, T. J. and AVERY, O. T.

The occurrence during acute infections of a protein not normally present in the blood. I. Distribution of the reactive protein in patients' sera, and the effect of calcium on flocculation reaction with C polysaccharide of Pneumococcus

J. Exper. Med. 73:173-182, 1941

Sera obtained from both humans and monkeys during acute infections contained a protein which was precipitated by the C polysaccharide of pneumococci. C-precipitable substance was inactivated by exposure to temperatures above 65° C., or the range of temperatures in which protein is denatured. After ammonium sulfate separation, the reactive material was found in the albumin fraction. Decalcification by additions of potassium oxalate and sodium citrate inhibited precipitation from reactive serum; the inactivation is reversible through the addition of minute amounts of Ca. The reactive protein is non-specific with respect to the inciting cause of the infection.

316. DELAUNAY, A., LEHOULT, Y., and SARCIRON, R.

Recherches sur le chimiotactisme leucocytaire. Pouvoir chimiotactique des antigenes glucido-lipidiques (Investigations on the chemotaxis of leukocytes. The chemotactic power of glucido-lipoid antigens)

Compt. rend. Soc. biol. 135:1529-1532, 1941

Autolyzed typhoid bacilli, in suspensions diluted from 1:10 to 1:1000, were injected in the guinea pig. Each of the 4 doses given contained 0.5 cc. of the diluted suspension. Microscopic examination of the injected tissue showed an invasion of neutrophils disseminated in the hypodermis. When the glucido-lipoid antigen of the typhoid bacillus was injected alone, it consistently produced the leukocyte migration, and antigens of a number of other bacteria (E. coli, the Flexner, para-typhoid B. and Shiga organisms) have identical activity. Maxima for leukocyte migration lie between 1:100 and 1:1000 dilutions and stronger doses have lower chemotactic power. The importance of the colloid structure in immunology is emphasized.

- 1 9 4 1 -

317. ELLIS, E. L. and SPIZIZEN, J.

The rate of bacteriophage inactivation by filtrates of Escherichia coli cultures

J. Gen. Physiol. 24:437-445, 1941

The anti-coli phage is inactivated by culture filtrates of E. coli in a pattern influenced by the temperature at which the reaction occurs. The rate of inactivation at 37° C. is proportional to phage concentration and filtrate concentration; at 0° C. the rate is proportional to the square root of the concentration of the filtrate. Coli-phage is inactivated by other colloids (starch, inulin, gum arabic); the inactivation is delayed by either very high or very low salt concentrations.

318. FELDMAN, J. and GELLHORN, E.

The influence of fever on the vago-insulin and sympathetico-adrenal systems

Endocrinol. 29:141-143, 1941

Typhoid-paratyphoid vaccine was injected intraperitoneally (0.7 cc. per kg.) into the normal rat and into adreno-demedullated or vagotomized animals. One group received vaccine after extirpation in toto of adrenals, and medulla, and sectioning of the vagus nerves. The vaccine produced hyperglycemia in the normal animal, hypoglycemia in the adreno-demedullated animals but no alterations in blood sugar after vagus section in adreno-demedullated rats. Hyperthermia appeared in the normal and vagotomized rat; removal of adrenals and medulla produced hypothermic reactions. In the vagotomized animal, both hyperthermia and high blood sugar levels persisted much longer than in the normal animal. The excitation of autonomic centers may affect discharge over the vago-insulin system as well as in the sympathetico-adrenal network.

319. FREEMAN, G. G. and ANDERSON, T. H.

The hydrolytic degradation of the antigenic complex of Bact. typhosum Ty 2

Biochem. J. 35:564-577, 1941

Dissociation by slow hydrolysis of the antigen of B. typhosa Ty 2 yielded the following components: a polysaccharide, constituting 50 to 60 per cent of the total; an insoluble polypeptide

fraction (16 per cent), a soluble nitrogenous material (from 10 to 20 per cent) and a small amount of lipid. The polysaccharide was highly dextro-rotatory, ketose and pentose-free, but contained d-galactose, d-mannose and d-glucose. There were no uronic acid residues. Identical polysaccharides have been prepared from the dried organisms directly. The polypeptide component was insoluble in water, soluble in dilute alkalis, and contained 8.2 per cent N, partly as tyrosine and arginine units. Approximately 50 per cent of the total nitrogen appeared as free amino-groups after acid hydrolysis. The nature of the soluble nitrogenous component was not determined.

320. GOLDRING, W., CHASIS, H., RANGES, H. A., and SMITH, H. W.

Effective renal blood flow in subjects with essential hypertension

J. Clin. Invest. 20:637-653, 1941

Filtration rates, diodrast clearance, and maximal rate of tubular excretion of diodrast were determined in 60 patients with essential hypertension. The results were compared with those obtained from a previous study of the normal kidney. The effective blood flow per unit of functional renal tubular tissue was usually below normal, the filtration fraction elevated, and the maximal rate of diodrast excretion far below normal. Small doses of pyrogenic inulin resulted in renal hyperemia and a fall in the filtration fraction. In some patients, the maximal rate of diodrast excretion in the tubule increased during hyperemia.

321. HARTWELL, J. L., SHEAR, M. J., KAHLER, H. and TURNER, F. C.

Properties of the fraction of B. prodigiosus which produced hemorrhage in mouse sarcomas

Cancer Res. 1:741, 1941

A summary is given of the methods now used to determine the properties of the hemorrhage-producing filtrate concentrate of Shear. Protein content was below 1 per cent; the biuret, ninhydrin, xanthoproteic and Millon's tests were negative. High molecular weight components were present as shown by dialysis of the filtrate. The Molisch reaction was positive, while iodine-iodide and Benedict tests gave negative results. Freezing of active solutions destroyed most of the potency. All fractions appeared to be equally toxic to tumor tissue when a bioassay was made on mice bearing growing sarcomas. Two active fractions were obtained by ultra-centrifugation and electrophoresis.

322. MACLEOD, C. M. and AVERY, O. T.

The occurrence during acute infections of protein not normally present in the blood. II. Isolation and properties of the reactive protein

J. Exper. Med. 73:183-190, 1941

The reactive protein present in the serum albumin fraction of blood and precipitable by the C polysaccharide of pneumococci, becomes soluble in tap-water after removal of the alcohol-ether soluble lipids and is no longer precipitable by calcium in small amounts. It still retains capacity for precipitation with the C polysaccharide.

323. MORGAN, H. R.

Immunologic properties of an antigenic material isolated from Eberthella typhosa

J. Immunol. 41:161-180, 1941

The pyrexial substance from E. typhosa is antigenic and elicits fever in the animal when circulating antibodies are present. Washed precipitates of antigen and excess antibody cause elevations in temperature and leucopenia. Leukocytosis follows. The factor is heat-stable. Activity remains in the acid-insoluble portion after acid hydrolysis.

324. MORGAN, W. T. J. and PARTRIDGE, S. M.

Studies in immunochemistry. VI. The use of phenol and of alkali in the degradation of antigenic material isolated from Bact. dysenteriae (Shiga)

Biochem. J. 35:1140-1163, 1941

The antigenic complex of B. dysenteriae Shiga is soluble in 90 per cent phenol solution, partially soluble in aqueous alkaline solution. The 'smooth' variant yields the antigenic complex and nonspecific water-insoluble simple proteins, the 'rough' variant yields no antigenic material. Polysaccharide components can be isolated in a free state by dialysis of the phenol solution. This protein-free carbohydrate fails to induce formation of specific 'Shiga' agglutinins or precipitins in the rabbit. The acidic (polypeptide-like) protein component appears to be a conjugated protein, non-antigenic when separated from the polysaccharide, but, on recombination with undegraded specific polysaccharide, a powerful antigen. A simple protein isolated

by phenol or alkaline dissociation was a higher N content and an optical rotation which differs from the conjugated protein. It does not form an antigenic complex when mixed with 'Shiga' polysaccharide in aqueous solution.

325. NISHISHITA, S.

Studies on the fluctuation of the lipoid content of the blood in the fever period

Jap. J. Exper. Med. 19:97-107, 1941

In normal rabbits, the total lipids are reduced after induced fever. The drop may be as much as 30 per cent below normal levels. After 24 hours, normal levels reappear. Cholesterol is markedly affected. In infectious disease, blood lipids may be slightly lower during the febrile phase and rise above normal during convalescence. The cholesterol content varies over a wide range. In starvation, blood lipids increase in the early phases, fall slightly below normal in ten days; when fever accompanies starvation, cholesterol rises with lecithin and fatty acids. A rise of cholesterol of 155 per cent above normal was seen on the 6th day of fever and starvation.

326. OLITZKI, L., AVINERI-SHAPIRO, S. and BENDERSKY, I.

The leucopenic action of different micro-organisms and the antileucopenic immunity

J. Immunol. 41:361, 1941

Heat-killed bacterial suspensions of various strains of Coccaeae, Bacteriaceae, Bacillaceae and the mycobacteria were injected into the guinea pig or rabbit to determine the effect of different antigens and of varying amounts of each, the development of antileukopenic immunity of both passive and active character, and its species or strain specificity. The leukopenic factor was present in certain groups of organisms only and qualitative differences due to their chemical composition were noted. Within the same species of any one group, quantitative variations for leukopenic activity could be found. Bacterial vaccines or normal rabbit sera produced a nonspecific antileukopenic resistance lasting from 2 to 3 days. Anti-leukopenic immunity could be invoked by those bacteria which contain a common heat-resistant O antigen, but passive immunity against leukopenia was conferred with difficulty.

- 1941 -

327. PASRICHA, C. L., MALIK, K. S. and PAUL, B. M.

'Pyrogen' found in salines for intravenous use

Indian Med. Gaz. 76:216-218, 1941

The soluble, heat-stable pyrogens which develop in parenteral solutions through bacterial protein splitting are not easily detected by permanganate titer. Only extreme care in production will prevent their appearance, but adsorption on charcoal will usually remove all traces.

328. SANDERS, T. E.

Non-specific protein therapy in ocular disease

J. Iowa Med. Soc. 31:51-54, 1941

Use of foreign protein therapy materially shortens the course of many ocular infections and reduces permanent damage. It is particularly effective in the treatment of ocular trauma. Foreign protein therapy has been used, with more or less success, in typhoid fever, pneumonia, puerperal fever, arthritis, pelvic inflammation, syphilis of the central nervous system, other nervous disorders, multiple sclerosis, peptic ulcer, dermatologic conditions, and in thrombo-angiitis obliterans.

329. SANDWEISS, D. J., SUGARMAN, M. H., FRIEDMAN, M. H. F., SALTZSTEIN, H. C. and FARBMAN, A. A.

The effect of urine extracts on peptic ulcer. An experimental and clinical study

Am. J. Diges. Dis. 8:371-382, 1941

Pregnancy urine and normal female urine apparently contain a principle capable of exercising a beneficial effect on peptic ulcer, as determined by the prolongation of life spans of dogs with experimental ulcers, by the fibroblastic proliferation and epithelization of ulcer craters. The action by which healing is initiated appears to be concerned with suppression of secretion, but healing appears through stimulation of fibroblastic and epithelial cells and formation of blood vessels, as well. When 14 patients with duodenal ulcer were given large doses (30 to 40 mg.) of normal female urine extract, subcutaneously or intramuscularly, after injection of stimulating amounts of histamine, total volume, but not acidity, of gastric secretion decreased. For chronic peptic ulcer, a high percentage of remissions and total improvement was obtained. Relapses within 6 months were less frequently seen than with other medicaments.

330. SHEAR, M. J.

Effect of a concentrate from B. prodigiosus filtrate on subcutaneous primary induced mouse tumors

Cancer Res. 1:731, 1941

A nitrogen-containing polysaccharide, obtained as a concentrate from broth cultures of B. prodigiosus, was injected in mice bearing primary chemically-induced tumors. Doses more than 500 times the minimum hemorrhage-producing dose were lethal for most of the tumor-bearing strains, but normal mice tolerated even larger doses. Severe hemorrhage was produced in a few hours in the primary spindle cell sarcoma, but regrowth was commonly seen. Mice with larger tumors usually succumbed to a given dose while those with small tumors survived. Repeated injections of gradually-increased small doses of the polysaccharide produced tolerance to the lethal effect, but lethality was always more pronounced for the tumor-bearing mouse.

331. WEINBERG, J. and GOLDSTEIN, H. H.

Modification of the use of typhoid vaccine in the production of hyperpyrexia

Illinois M. J. 79:139-140, 1941

The authors devised an expanding dosage method for the production of therapeutic hyperpyrexia. Typhoid vaccine was administered in a low initial dose until fever was produced; at the height of fever, a second larger dose was injected. Very high temperatures were secured by this method. A definite dosage schedule is appended to the article.

332. WIENER, A. S. DREMLAND, B. H., HYMAN, M. A. and SAMWICK, A. A.

Transfusion reactions. Experience with more than 3000 blood transfusions

Am. J. Clin. Path. 11:102-121, 1941

In 1936, the incidence of chills and fever appearing after blood transfusion was 12 per cent, on the average, regardless of the method employed in transfusion. When the saline solution used was exchanged for another type, reactions dropped to 3.8 per cent and has continued to be between 2 and 3 per cent.

- 1 9 4 1 -

333. WRIGHT, I. S.

The conservative treatment of occlusive arterial disease

New England J. Med. 225:805-810, 1941

Typhoid vaccine has been used intravenously in the treatment of thromboangiitis obliterans with highly satisfactory results. The 'H' antigen of the typhoid bacillus will produce a therapeutically useful fever without the distressing chill.

334. YAMAMOTO, K.

Studien über die Beziehung zwischen der Virulenz und den giftigen Leibessubstanzen der Maustyphusbazillen (Studies on the relationship between virulence and toxic endogenous substances in mouse typhoid bacilli)

Tohoku J. Exp. Med. 41:69-91, 1941

The less virulent strains of the typhoid bacilli have low percentages of polysaccharides and high protein content. Endogenous toxic substances are more readily isolated from more virulent strains, which yield larger amounts of the carbohydrate-protein complex.

- 1 9 4 2 -

335. BLOOM, N.

Treatment of peripheral vascular diseases

Virginia M. Month. 79:273, 1942

The action of typhoid vaccine on the collateral vessels supplying the lower limbs with blood is one of vasodilatation and improved tissue nutrition. Buerger's disease is greatly improved by this type of therapy.

336. CENTANNI, E.

Immunitäts Erscheinung im experimentellen Fieber mit besonderen Berücksichtigung des pyrogenen Stoffes aus Typhusbakterieren (Appearance of immunity in experimental fever with particular emphasis on the pyrogenic materials from the typhoid bacillus)

Klin. Wchnschr. 21:664-669, 1942. See also: Bull. Hyg. (London) 17:890, 1942

The pyrogenic decomposition product obtained from typhoid bacilli appears to exist as a prosthetic group on a large protein complex molecule. Agar cultures of the typhoid bacillus 1 to 2 days old are prepared by autolysis and either pepsin or papain digestion; trypsin is less effective and acidic or alkaline hydrolysis injures the pyrogen. The pyrogen can be boiled in neutral solution without damage. "It even appears that through such action its activity is increased, either through destruction of interfering material or through..liberation of..fever material." The large polypeptides and albumin may be removed through the use of trichloroacetic acid or ammonium sulphate. Histamine hydrolysis and dialysis will remove materials of smaller molecular mass. Centanni preferred fractional precipitation with alcohol and glycol for further purification before filtration. The fever unit was 0.05 mg.

A new substance appeared during a study of the method outlined above, a biological unknown related to the polysaccharides. It is found in combination with the 'klastins' or endotoxins which produce characteristic immunity and certain toxic symptoms, as fever and inflammation. The pyrogenic factor itself contains both a hapten and a toxic group. The former produces pure histologic immunity through its action as an anti-pyrogen, with a prevention of antibody formation. Through fractional adsorption on aluminum hydroxide, the hapten can be obtained in a fairly pure state and may be then used in immunization against the pyrogenic reaction. This, the 'Third Immunity' of Centanni, has important potentialities in the treatment of disease.

337. CHAPMAN, C. J.

Use of rabbits for the detection of pyrogenic substances in solutions for intravenous administration.

Quart. J. Pharm. Pharmacol. 15:361-366, 1942

Initial leukopenia, after the injection of pyrogen-containing infusion solutions, may be used as a simple indicator of the presence of pyrogen. In the test rabbit, a drop of 4000 leukocytes per cmm. of blood or more is considered evidence of pyrogen contamination in the injected solution.

338. CHASIS, H., GOLDRING, W. and SMITH, H.

Reduction of blood pressure associated with pyrogenic reaction in hypertensive subjects

J. Clin. Invest. 21:369-376, 1942

No inherent vasodilator or depressor activity is known to reside in pyrogens, and the vasomotor responses noted are believed to be complex. The initial pressor phase is characterized by vasoconstriction, probably neurogenic in origin; systolic and diastolic pressures are moderately reduced shortly thereafter. A dual activity in pyrogens is suggested by the fact that the temperature rise may be blocked without altering the hypotensive action and renal vasodilation. Premedication with amidopyrine allows the hypotensive effect to be applied in hypertensive states, without a corresponding increase in body temperature. The delayed activity of the second phase of the hypotensive cycle after administration of pyrogens may, if dosage is not controlled, produce alarming circulatory crises.

339. CO TUI, and SCHRIFF, M. H.

A tentative test for pyrogens in infusion fluid

Proc. Soc. Exper. Biol. Med. 49:320-323, 1942

These investigators used, with success, a procedure which eliminated some of the errors commonly encountered in pyrogenic testing. Since the rabbit is responsive to other substances and has a highly labile thermoregulatory mechanism, the lack of response to pyrogen injection has more significance than the positive test. The dog is less sensitive to pyrogenic materials; hence positive pyrogenic reactions in this animal are highly significant. Preliminary tests on the rabbit are confirmed through injection of a test dose in the dog. The test dose is graduated according to the sensitivity displayed. The rabbit is 3 times, the dog 6 times less sensitive than human beings. In 214 tests, clinical and laboratory data have been in agreement.

340. CO TUI and SCHRIFF, M. H.

Production of pyrogen by some bacteria

J. Lab. Clin. Med. 27:569-575, 1942

In a study of pyrogen production in 18 strains of micro-organisms, representative of different genera, families and species

pyrogen production was found to be common to many types of bacteria. It was not confined to organisms found in water, and was not associated with nucleic acid, with lack of affinity for the gram stain, pathogenicity, chromatin production, nor was it confined to lower biologic forms. Saccharomyces Pink wild yeast produces pyrogen. In this study, a new criterion was employed in the judgment of toxicity; the production of antibodies in experimental animals.

341. CO TUI and WRIGHT, A. M.

The preparation of non-pyrogenic infusion and other intravenous fluids by adsorptive filtration

Ann. Surg. 116:412-425, 1942

The adsorptive filtration method, and a standardized apparatus, have been adopted for the production of non-pyrogenic physiologic salt, dextrose and sodium citrate solutions. The use of calibrated asbestos filter pads is a feature of the method.

342. DELAUNAY, A.

Recherche des constituents bacteriens responsables de l'appel leucocytaire (Investigation of bacterial constituents responsible for the leukocyte response)

Ann. Inst. Pasteur, 68:72-74, 1942

Marked leukocytic response was produced by an autolysate of staphylococci. Dealbuminized autolysates yielded an active, specific polysaccharide, capable of affecting the leukocyte count in concentrations ranging between 0.001 and 1.0 mg. The physiologic carbohydrate content is directly concerned in the stimulation of leukocyte production. Chemotaxis may be involved but this was not verified by a comparison with glycogen. Similar activity is shown by preparations prepared from typhoid, colon, paratyphoid B and Shiga organisms. It is believed that the active principle is a specific glucido-lipid antigen, since local leukocyte response follows the injection of small quantities of the substance beneath the skin.

343. DUBOS, R. J., HOBERMAN, H. D. and PIERCE, C.

Some factors affecting the toxicity of cultures of Shigella dysenteriae

- 1 9 4 2 -

Proc. Nat. Acad. Sci. 28:453, 1942

Conditions which favor aerobic metabolism greatly increase the toxicity of the cells in S. dysenteriae cultures, as well as the yield of dry material. When 1.5 mg. of cells grown under anaerobic conditions were injected into mice, all survived; 0.1 mg. of cells grown on silica gel plates killed all mice. Glucose in silica gel medium decreased the toxicity of cells. 'Smooth' and 'rough' varieties were similarly affected by environmental growth conditions. The toxic principle was bound to the cell itself and released in solution through autolysis. It was water soluble, stable and resistant to proteolysis.

344. DUGGAN, W. F.

Choroiditis centralis serosa: Diagnosis, pathologic physiology and therapy

Arch. Ophthalmol. 27:123-138, 1942

Bacterial body cellular elements and toxins produce a local ischemic condition after injection. As this is overcome by pyrexial factors, oxygen is increased in the area with beneficial effects.

345. FAVORITE, G. O. and MORGAN, H. R.

Effects produced by the intravenous injection in man of a toxic antigenic material derived from Eberthella typhosa: Clinical, hematological, chemical and serological studies

J. Clin. Invest. 21:589-599, 1942

No marked changes in the total protein, urea nitrogen, creatinine, chloride, glucose and CO₂-combining power of blood appeared after a purified antigenic material obtained from E. typhosa was administered to human beings with neurosyphilis. Repeated injections produced antibodies in high titer. The leukopenia and leukocytosis usually noted after pyrogen injections were present. The sedimentation rate rose after each injection. In animals receiving intravascular injections of this substance, pathologic changes involving heart muscle, lung, liver and spleen tissue, the bone marrow and vascular bed were produced.

346. FEDOROV, N. A. and SHUR, E. L.

The role of the viscera in regulating the temperature of the body of an animal under physiological and pathological conditions

Am. J. Physiol. 137:30-38, 1942

The sites of thermogenesis were studied in normal animals artificially heated or cooled and febrile states after blood transfusion. All experiments were made on angiotomized dogs. In normal fasting dogs, lowest temperatures were found in the aorta, highest in the hepatic vein. Important thermogenesis occurred in the intestines. Hepatic heat production rose rapidly during cooling of the skin and the reverse took place during heating. In transfusion and anaphylactic shock, variations were much greater in temperature, particularly with respect to visceral thermogenesis. Both liver and intestines participate in determining febrile reactions to foreign protein substances.

347. GERBER, I. E. and GROSS, M.

Production of the Shwartzman phenomenon with a sulfonamide conjugate of a bacterial filtrate

Proc. Soc. Exper. Biol. Med. 51:237-238, 1942

In four experiments performed with the rabbit, the skin was prepared by an intradermal injection of meningococcal filtrate, and in four others, a conjugate of paraminobenzenesulfonylacetamide with bacterial filtrate was used for preliminary conditioning to the Shwartzman reaction. Both filtrate alone and the conjugate were injected 24 hours later. The filtrate was still able to elicit the reaction despite conjugation.

348. HIATT, E. P.

The effect of denervation on the filtration rate and blood flow in dog kidneys rendered hyperemic by administration of pyrogens

Am. J. Physiol. 136:38-41, 1942

In order to determine whether or not the delayed renal hyperemia induced by pyrogens was under neural or humoral control, the ureters of two dogs were explanted to the exterior, 100 mg. of pyrogenic inulin administered, and urine from each kidney separately collected. The determinations of glomerular filtration rate and diodrast clearance were repeated after

- 1 9 4 2 -

denervation of the left kidney in each animal.

Denervated kidneys functioned in the same manner as normal kidneys during the experiment. In response to pyrogen, a decreased filtration fraction and increased renal blood flow were obtained. The pyrogenic renal hyperemia is apparently humoral and not neural in nature.

349. HUTNER, S. H. and ZAHL, P. A.

The action of sulfanilamide compounds on the lethal factor of bacterial toxins

Science 96:563-564, 1942

Sulfanilamide exhibits a protective action on mice after administration of lethal doses of purified Salmonella endotoxin. The method may have useful application in the toxicity encountered after the induction of hemorrhage in tumors through bacterial fractions. The antitoxic effect is inhibited by para-amino-benzoic acid.

350. KNOPPERS, A.

The rat tail as a part for temperature regulation (Translated Title)

Arch. Neurl. physiol. 26:364, 1942

Since the temperature of the rat tail rises after injections of pyrogens the rise in temperature may be utilized in testing for fever production.

351. KOPP, I. I.

Plasma protein in therapeutic fever

J. Lab. Clin. Med. 27:1054-1062, 1942

Fever induced by malaria organisms produced plasma protein shifts in patients with general paresis. The globulin shifted, albumin was decreased. The fibrinogen content varied. Injections of typhoid vaccine and physically-induced fever brought smaller changes. Destruction of red cells, as well as the infectious process itself, may account for plasma globulin increases in malaria.

352. KOPP, I. I.

Relationship of the plasma protein to the corrected sedimentation rate

J. Lab. Clin. Med. 27:1072-1077, 1942

As globulin in plasma increases in malarial or typhoid vaccine fever the sedimentation rate is proportionately heightened.

353. MORGAN, W. T. J. and PARTRIDGE, S. M.

An examination of the O-antigenic complex of Bact. typhosum

Brit. J. Exper. Path. 23:151-165, 1942

A substance possessing the immunologic properties of the O antigen of E. typhosa was prepared by diethyleneglycol extraction. Three components were found, a protein, a phospholipin and a specific polysaccharide. The protein component was chemically and immunologically similar to the conjugated protein found in B. Shigae antigen. Toxicity of both antigenic preparations appears to reside in the undegraded polysaccharide component; degradation removes toxicity.

354. NEAL, P. A., SCHNEITER, R. and CAMINITA, B. H.

Report on acute illnesses among rural mattress makers using low-grade stained cotton

J.A.M.A. 119:1074-1082, 1942

An acute illness appeared among workers on a rural mattress project, in a cotton mill and in cottonseed-processing plants. High concentrations of stained cotton dust were found in the occupational environment. Samples of the cotton were given physical, chemical, mycologic serologic, biologic and bacteriologic examinations. A gram-negative bacillus was found in stained cotton and cotton dust; the disabling illness, characterized by acute intoxication appearing shortly after exposure, could be reproduced by inhalation of dust containing the bacterium. Aerosols of a sterile filtrate from cultures of the 'cotton' bacterium produced similar symptoms.

355. NECHELES, H., DOMMERS, P., WEINER, M., OLSON, W. H. and RYCHEL, W. M.

Depression of gastric motility without elevation of body temperature following the injection of pyrogen

Am. J. Physiol. 137:22-29, 1942

Pentnucleotide, yeast or thymus nucleic acid, peptone, B. coli vaccine or triple typhoid vaccine, a bacterial pyrogen, and tap water were administered in subfebrile doses into unanesthetized fasting dogs in which continuous gastric motility had been produced by prostigmine. Pentnucleotide inhibited gastric motility in all tests. The average duration of activity was 30 minutes. Yeast nucleic acid had deep depressive effects, thymic nucleic acid none. Peptone, 5 and 50 mg., neither affected temperature nor motility. B. coli vaccine (12-1/2 million organisms) depressed gastric motility. The pyrogenic preparation depressed gastric motility profoundly.

356. OLITZKI, L., AVINERI-SHAPIRO, S. and KOCH, P. K.

The hypothermic and adreno-hemorrhagic effects of bacterial vaccines

J. Immunol. 45:237-248, 1942

A large number of vaccines from a variety of bacterial species were tested for the following: hypothermic effects, adreno-hemorrhagic effects; nonspecific resistance through treatment with bacterial vaccines; nonspecific resistance through treatment with serum or plasma, heterologous or homologous; active and passive immunization.

Adreno-hemorrhagic, hypothermic and leukopenic reactions appeared to be associated and were produced only by vaccines from certain organisms, in the main, the gram-negative groups. Reactions were varied in intensity and occurrence and influenced by such nonspecific factors as amounts, age and weight of experimental animals, and the species. Normal serum or plasma produced a resistant period extending 2 to 3 days, vaccine, for 12 days. However, vaccine prevented leukopenia after injection of bacterial products for two days only. It did not protect against the adreno-hemorrhagic effects, nor did passive immunization.

357. RETES, F. P.

Algunas alteraciones hemáticas provados por el pyrifer en los esquisofrenicas (Some alterations in blood provoked by pyrifer in schizophrenics)

Rev. neuro-psiquiat. 5:381-413, 1942

No differences in the kind, degree or duration of leukocytosis was observed in schizophrenic patients who received injections of pyrifer; a small reduction in the hemoglobin content of the blood followed.

358. SCHRADE, W.

Effect of bacterial substances (pyrogens) on blood lipides (Translated Title)

Ztschr. f. ges. exper. Med. 110:518-534, 1942

Blood fats, phosphatides and cholesterol are decreased after the injection of bacterial pyrogens in the dog. Free and total cholesterol is less affected. Catabolism of fat may be stimulated. Normal levels are resumed after pyrogen activity ceases.

359. SOLOMON, H. A. and SOMKIN, E.

An improved method of obtaining controlled hyperpyrexia with triple typhoid vaccine

Am. J. Med. Sci. 203:736-740, 1942

Continuous infusion of triple typhoid vaccine permits prolonged hyperpyrexia to be maintained. Only one chill phase occurs. The method is applicable to the chemotherapeutic treatment of subacute bacterial endocarditis, pyogenic lung infections, chronic arthritis and ulcerative colitis. It may be used to augment the effects of sulfonamides. Calcium gluconate or chloride and morphine sulfate suppress hyperpyrexia.

360. WEINSTEIN, J. J.

Reactions from the transfusion of unpooled liquid human plasma. Analysis of 1500 transfusions

Med. Ann. Dis. Columbia 11:226-229, 1942

- 1 9 4 2 -

A total of 15 reactions (1 per cent indidence) in 1500 transfusions were observed; all were mild. From the standpoint of precipitating causes, they were divided into 3 groups (1) those caused by agglutination (2) allergic reactions and (3) pyrogenic contamination. The last-named was most frequently encountered. An outline is presented for the prevention of plasma reactions.

361. WINKELSTEIN, A. and SHWARTZMAN, G.

The use of concentrated and purified antitoxic B. coli serum in the treatment of indeterminate ulcerative colitis

Am. J. Diges. Dis. 9:133-136, 1942

Whole and concentrated horse serum, strongly antitoxic, was prepared from immunizing doses of B. coli. Since this organism is considered to be a secondary invader in chronic ulcerative colitis, intramuscular injections of the concentrated B. coli antitoxin (Shwartzman) were administered to 29 patients exhibiting chronic aspects of colitis. The results were encouraging, 22 in 29 patients were benefited by this type of therapy.

362. YOUNGNER, J. S. and NUNGESTER, W. J.

The effect of pneumococcus polysaccharide on the circulation of the blood in small blood vessels

J. Bact. 43:94, 1942 (Abstract)

Intravenous injection of various quantities of pneumococcus polysaccharide produced changes in the rate and uniformity of flow of erythrocytes. Increased sedimentation rates were observed.

363. ZAHL, P. A., HUTNER, S. H., SPITZ, S., SUGUIRA, K. and COOPER, F. S.

The action of bacterial toxins on tumors. I. Relationship of the tumor-hemorrhagic agent to the endotoxin antigens of gram-negative bacteria

Am. J. Hygiene 36:223-242, 1942

The antigenic endotoxin of Shigella dysenteriae, on extraction with cold ethylene glycol, yields three components: a phospholipid, non-toxic, non-antigenic and not a hapten; a polysaccharide, comprising approximately 50 per cent of the endotoxin,

- 1 9 4 2 -

and haptenic in structure but nontoxic; the toxic polypeptide fraction, non-antigenic. The last-named fraction gives intense color reactions for tyrosine and arginine. It may be this type of fraction which produces tumor hemorrhage.

364. ZAHL, P. A. and HUTNER, S. H.

The action of bacterial toxins on tumors. II. Effect of sulfanilamide on toxin-induced hemorrhage

Proc. Soc. Exper. Biol. Med. 51:285-287, 1942

The hemorrhagic factor found in Salmonella endotoxin is believed to be a component of the O antigen and to exhibit parallel lethal effects in a manner common to most negative bacteria. The lethal activity is reduced in vivo by sulfanilamides, but the mode of action of such protection is not understood. It may, like ascorbic acid, affect the threshold of vascular resistance or participate in a toxicity-reducing mechanism.

- 1 9 4 3 -

365. DELAUNAY, A.

Recherche sur la phagocytose. V. Chimotactisme leucocytaire et toxines microbiennes (Research in phagocytosis. V. Leukocyte chemotactism and bacterial toxins)

Rev. immunol. 8:30-45, 1943

Bacterial toxins with the characteristics of proteins were isolated from staphylococci, corynebacteria of the pseudotubercle type, and organisms of diphtheria and tetanus. On injection, they inhibited chemotaxis of the leukocytes, preventing phagocytosis.

366. DELAUNAY, A.

Le pouvoir leucopéneant de l'antigène glucido-lipidique du Bacille d'Eberth et la leucopenia dans la fièvre typhoïde (The leukopenic ability of the glucido-lipid antigen of E. typhosa and leukopenia in typhoid fever)

Compt. rend. Soc. biol. 137:540, 1943

The leukopenic factor has been sought in E. typhosa. It has been considered to be nonprotein (Morgan); the Molisch reaction is positive. Trichloroacetic extracts of bacterial preparations produce leukopenia on injection.

367. DELAUNEY, A.

Mise en evidence d'une nouvelle propriété des antigènes glucido-lipidiques; Leur pouvoir leucopéneant (A new property of glucido-lipid antigens: Their leukopenic ability)

Compt. rend. Soc. biol. 137:589-590, 1943

A new property of antigens, the leukopenia-producing ability, appears when lethal doses of glucido-lipid antigens from colon or typhoid organisms are administered. The leukocyte count is depressed within one hour (from 10,000 to 25,000 initially to 5,000 or less). This is an early sign of intoxication. The leukocyte index is balanced by the increased mononuclear cells. In about 3 hours, the white cells increase. With more moderate dosages, the return to normal levels begins in 24 hours. Minute (gamma) dosages bring an early leukocytosis; the animal so treated remains in perfect health. This stimulation is believed to be a chemotactic one. According to J. Jolly, leukopenia occurs when leukocytes are retained in viscera, liver, lung and spleen - only small numbers remain in tissues. Glucido-lipid antigens bring intravisceral mobilization as the leukocytes migrate inward from the peripheral areas.

368. EDELMANN, A., MAHANNA, D. L., LEWIS, L. A., THATCHER, J. S. and HARTMAN, F. A.

The use of adrenal extract in fever therapy

J. Clin. Endocrinol. 3:20-27, 1943

369. GUERRA, F. and BARBOUR, H. G.

The mechanism of aspirin antipyresis in monkeys

J. Pharm. & Exper. Therap. 79:55-61, 1943

In normal monkeys aspirin produces only a slight decrease in temperature, but in fever produced by the subcutaneous injection of yeast, it induces vasodilation and perspiration.

370. HARRISON, J. W., MYERS, R. J. and HART, D. S.

Purified mineral-free water for pharmaceutical purposes

J. Am. Pharm. A. 32:121-128, 1943

By the use of ion exchange resins, pyrogens in mineral-free water or in solutions intended for parenteral use may be reduced. The amount of pyrogenic material remaining does not produce temperature increases above 0.5° C.

371. HARTWELL, J. L., SHEAR, M. J. and ADAMS, J. R., JR.

Chemical treatment of tumors. VII. Nature of the hemorrhage-producing fraction from Serratia marcescens (Bacillus prodigiosus)

J. Nat. Cancer Inst. 4:107-122, 1943

The active material contained 47.5 per cent C, 7.1 per cent H, 2.2 per cent N, 1.1 per cent P and 2.2 per cent acetyl. Ash was 3.5 per cent. The minimal hemorrhage-producing dose contains about 0.1 gamma of total solids. Neither protein nor polypeptide can be demonstrated after hydrolysis; hexosamine, methylpentose and a firmly bound phospholipid are present. Although stable at room temperatures, the potency is lost at increasing rates as temperatures increase between 60° C. and 100° C.

372. HARTWELL, J. L., SHEAR, M. J. and ADAMS, J. R., JR.

Nature of the bacterial polysaccharide which produces hemorrhage in mouse sarcoma

Cancer Res. 3:122, 1943

"Further data were given on the material obtained from cultures of B. prodigiosus". The polysaccharide fraction had previously given negative results in tests for protein; in this study colorimetric tests for tyrosine, arginine and histidine gave positive results. The active agent could be separated from the proteinate material by tryptic digestion. Potency of the active portion disappeared when the acetyl content was lost during treatment with 0.1 N/NaOH. By use of 0.1 N/NaHCO₃, the acetyl was removed with injury to activity. Hydrolysis (1.5 N/HCl) produced 60 per cent reducing sugars and 16 per cent of a water-insoluble, alkali-soluble product containing N and P. The reducing sugar was approximately 30 per cent methylpentose.

- 1 9 4 3 -

Glucosamine was about 4 per cent. No ketose or desoxy- sugars were noted; galactose, mannose and inositol were not present. A small amount of uronic acid was measured. The end-point dose (producing hemorrhage in 50 per cent of tumors) was about 0.1 gamma.

373. HUTNER, S. H. and ZAHL, P. A.

Action of bacterial toxins on tumors. IV. Distribution of tumor-hemorrhage agents among bacterial species

Proc. Soc. Exper. Biol. Med. 52:364-368, 1943

When 30 gram-negative and 7 gram-positive killed organisms, pathogenic or non-pathogenic, were injected in doses approaching lethal limits into mice bearing mouse sarcoma 180, representatives of gram-negative species induced tumor hemorrhage (with few exceptions). But one gram-positive bacterium (Listeria monocytogenes) possessed the capacity to produce vascular stasis in tumors. It is argued that tumor-hemorrhaging toxins may be bound in the somatic O antigens, characteristic of gram-negative organisms.

374. KAHLER, H., SHEAR, M. J. and HARTWELL, J. L.

Chemical treatment of tumors. VIII. Ultracentrifugal and electrophoretic analysis of the hemorrhage-producing fraction from Serratia marcescens (Bacillus prodigiosus)

J. Nat. Cancer Inst. 4:123-129, 1943

Analyses of 3 preparations of the polysaccharide fraction from S. marcescens which produces tumor hemorrhage gave the following information: the non-dialyzable solids were approximately 1 per cent of weight. The partial volume of polysaccharide by ultracentrifugal analysis was 0.75; the sedimentation rates differed. No polysaccharide component appeared to carry potency for other activities than that of producing hemorrhage in neoplasms. This potency was about equally divided between sediment and supernatant fluid. The electron micrograph revealed that the active principle was a large symmetrical particle with molecular weight somewhere between 3 and 30 million. It appears to remain in the material most easily sedimentable. The potency is lost rapidly at ordinary temperatures and pressures.

375. KNIGHT, H. C., EMORY, M. L., and FLINT, L. D.

A method of inducing therapeutic fever with typhoid vaccine using the intravenous drip technique

Vener. Dis. Inform. 24:323, 1943

The infusion of a dilute solution of typhoid vaccine in 5 per cent glucose-saline permits control of fever levels without danger to the patient. Over 1400 patients with gonorrhea of a resistant type were treated by means of an initial intravenous 'shock' dose of concentrated material, followed in 1 or 2 hours by intravenous slow infusion.

376. MORGAN, H. R.

Pathologic changes produced in rabbits by a toxic somatic antigen derived from Eberthella typhosa

Am. J. Pathol. 19:135-143, 1943

Marked inflammatory changes with later necrosis and fibrotic 'walling' of the area followed the injection in the skin of somatic fever-producing antigen of E. typhosa. When injected intravenously, it caused vascular damage and subsequent thrombosis, necrosis of hepatic cells and changes in heart muscle, adrenals and bone marrow. The lymphoid tissue of the spleen may be injured. Hyperplasia of bone marrow follows injury to the hematopoietic tissues.

377. McCLOSKEY, W. T., PRICE, C. W., VAN WINDLE, W., JR., WELCH, H., and CALVERY, H. O.

Results of the first U.S.P. Collaborative Study of pyrogens

J. Am. Pharm. A. 32:69-73, 1943

Tolerance to the pyrogen from Ps. aeruginosa develops on repetition of injections over a considerable time; with an intervening rest period of three weeks, the response phase is restored. The selection of animals for the bio-assay is emphasized and a method outlined for shortening test periods.

- 1 9 4 3 -

378. NECHELES, H.

Depression of the stomach by nonspecific substances

Proc. Inst. Med. (Chicago) 14:345-346, 1943.

Prolonged depression of gastric motility and secretion was possible through the injection, in the dog, of pyrogenic substances. The rectal temperature was unchanged. Pepsin, enterogasterone and other depressants of gastric motility may owe their effects to pyrogens.

379. OTTENSOOSER, F.

Sobre o pirogenio (About pyrogen)

Arq. biol. (São Paulo) 27:1-?, 1943

Biological and chemical tests for the presence of pyrogens are evaluated. The production of gelation in certain media, of initial leukopenia and subsequent leukocytosis are characteristic of bacterial pyrogens and pentose nucleotides. Pyrogen may be produced in or related to nucleic cellular material.

380. SHEAR, M. J., PERRAULT, A. and ADAMS, J. R., JR.

Chemical treatment of tumors. VI. Method employed in determining the potency of hemorrhage-producing bacterial preparations

J. Nat. Cancer Inst. 4:99-105, 1943

The bioassay method for determining the potency of bacterial preparations for production of mouse tumor hemorrhage was found to be inadequate for study of chemical and physical properties of the active agent. A modified technique was adopted, sensitive enough to enable distinctions between potencies to be made. A standard material of determined potency was used for reference. All contaminated tumor mash was discarded. Examination of the interior of the tumor was performed. The mouse tumor unit was defined as 'the amount of activity contained in the end point dose'.

381. SHEAR, M. J. and TURNER, F. C.

Chemical treatment of tumors. V. Isolation of the hemorrhage-producing fraction from Serratia marcescens (Bacillus prodigiosus) culture filtrate

J. Nat. Cancer Inst. 4:81-97, 1943

Highly potent preparations of the active agent for tumor hemorrhage were prepared from filtrates of S. marcescens through precipitation with chloroform, ethyl alcohol and dialysis. The active fractions were tested for potency, chemical properties and toxicity. The carbohydrate components were in the majority (positive Molisch) and the protein very small or absent (negative biuret, trichloroacetic acid and sulfosalicylic acid). pH changes between 3.0 and 10.0 did not affect potency. Lyophilization produced no loss of activity. The most active concentrate obtained had a potency of 130,000 m.t.u. per cc. The lethal dose ranged from 100 to 1000 times the minimum hemorrhagic dose (0.1 gamma total solids). Fractions of low potency were eliminated through a bioassay technique. When tumors were treated early in growth, they did not respond to bacterial action; old tumors were prone to develop brown discolorations which obscured hemorrhage.

382. TAYLOR, R. D. and PAGE, I. H.

Severe renal irritation resulting from foreign protein (typhoid) fever therapy

J.A.M.A. 121:754-755, 1943

Severe renal irritation developed during treatment with intravenous typhoid vaccine of patients ill with malignant hypertension and Bright's disease with nephrosis. Caution must be employed in the use of this type of therapy where renal disease exists.

383. WELCH, H., CALVERY, H. O., McCLOSKEY, W. T. and PRICE, C. W.

A method of preparation and test for bacterial pyrogens

J. Am. Pharm. A. 32:65-69, 1943

Lack of antigenicity in Pseudomonas aeruginosa pyrogen was shown by the fact that, in a test in rabbits, 11 injections were made in each animal over a period of 9 weeks without tolerance to the dosage developing.

- 1 9 4 3 -

384. ZAHL, P. A. and BJERKNES, C.

Induction of decidua-placental hemorrhage in mice by the endotoxins of certain gram-negative bacteria

Proc. Soc. Exper. Biol. & Med. 54:329-332, 1943

Endotoxins of Shigella paradysenteriae Flexner, S. typhimurium, and Rhodospirillum rubrum were injected in mice intraperitoneally on the 12th to 18th day of gestation; one-fifth of the LD₅₀ dose was used. All mice responded with decidua-placental hemorrhage. No other tissues were hemorrhagic.

385. ZAHL, P. A. and HUTNER, S. H.

Action of bacterial toxins on tumors. III. Some biological properties of purified Salmonella typhimurium endotoxins

Proc. Soc. Exper. Biol. & Med. 52:116-118, 1943

The object of this study was the determination of the activity of the purified O antigen in respect to lethality, hemorrhage production in transplanted tumors and production of protective antibodies in mice.

In three stages of purification, fractions were active for all three of the above phenomena.

386. ZAHL, P. A. HUTNER, S. H. and COOPER, F. S.

Action of bacterial toxins on tumors. V. Immunological protection against tumor hemorrhage

Proc. Soc. Exper. Biol. & Med. 54:48-50, 1943

After immunizing doses of Shigella paradysenteriae (1/2 to 1 LD₅₀) were injected intraperitoneally in mice for 6 days; an interval of 10 to 15 days was allowed before the implantation of sarcoma 180. Seven days later hemorrhage-inducing doses of Flexner endotoxin were administered. Non-immunized mice were used as controls.

Immunized animals showed no hemorrhage at dosages of endotoxin below LD₅₀. Controls were found to have tumor hemorrhage at dosage ranging above 1/200 of the LD₅₀.

- 1 9 4 3 -

387. ZAHL, P. A., HUTNER, S. H. and COOPER, F. S.

Action of bacterial toxins on tumors. VI. Protection against tumor hemorrhage following heterologous immunization

Proc. Soc. Exper. Biol. & Med. 187-189, 1943

Heterologous immunization was as effective as homologous in the protection against induction of tumor hemorrhage. A common antigenic toxic component may characterize all gram-negative bacteria.

- 1 9 4 4 -

388. BIOCCA, E. and OTTENSOOSER, F.

Sobre uma possível relação entre a ação pirogenica e um fenómeno de gelificação apresentado por diversos germes (On a possible relation between pyrogenic activity and the phenomenon of gelification presented by various bacteria)

Arq. biol. (Aso Paulo) 28:36-39, 1944

Pyrogen formation is found to closely parallel the presence of colon bacteria of the Salmonella-Shigella-Proteus groups, S. hemophilus, Brucella and vibrios of threadlike structures. These threads appear during growth of the organisms in 3 per cent KOH. The phenomena has a close relationship to gelation. The possibility exists that nucleic acid may play a part in the fibril development; similarities to the agglutination of red cells are noted.

389. BOSE, B. C. and AHUJA, M. L.

Detection of pyrogens in fluids by biological methods

Indian J. Med. Res. 32:9-14, 1944

Hyperpyrexia as an index for the presence of pyrogens in water gives only moderately satisfactory results; the leucopenia index is still less accurate. In this series, distilled water and reservoir waters were compared for the presence of pyrogens. Care in handling of test rabbits was observed in this study. The accuracy of rectal temperature readings depended upon the position of the thermometer during reading. In the normal rabbit, allowances of 0.6° C. in temperature were made. Only slight temperature changes were noted after the injection of

- 1 9 4 4 -

redistilled water. From reservoir waters, two strains of organisms were isolated; one belonged to the Achromobacterium prodigiosus group, the other, from the same species, produced chromogenic cultures. Both were aerobic, non-sporulating. One was gram-positive. Culture suspensions produced fever, but only insignificant decreases in the leukocyte count.

390. BRUES, A. M. and SHEAR, M. J.

Chemical treatment of tumors X. Reactions of four patients with advanced malignant tumors to injections of a polysaccharide from Serratia marcescens culture

J. Nat. Cancer Inst. 5:195-201, 1944

Four patients with inoperable tumors (Ewing's sarcoma, carcinoma of the prostate, lymphosarcoma and myeloma) were given injections of the hemorrhage-producing filtrate fraction prepared from B. prodigiosus. Chills, transitory fever and leukocytosis followed. No alterations were seen in multiple myeloma. There was immediate pain in tumorous areas in Ewing's sarcoma and carcinoma of the prostate; symptoms were somewhat relieved for a short period thereafter. In lymphosarcoma (the patient was in extremis) widespread destruction of tumor tissue followed the injection, and a temporary remission occurred. All patients died from malignancies. Post-mortem examination showed evidences of hemorrhage in tumor tissue.

The two patients who were temporarily benefited by therapy developed high NPN and uric acid levels in the blood. One patient excreted, on a low purine diet, 2500 mg. of uric acid in the first 24 hours after the injection. Some evidences of congestive failure were noted in two patients. The toxic response to uneliminated waste products is a serious medical problem where rapid destruction of neoplastic tissue follows chemotherapy.

391. CAMPBELL, D. J.

Gonorrhea in North Africa and the Central Mediterranean

Brit. M. J. 1:44, 1944

Adjuvants to the standard treatment are necessary if results are to be obtained in the intractable type of gonorrhea found in North Africa and the Levant. The use of TAB vaccine has been helpful.

392. CO TUI, HOPE, D., SCHRIFT, M. H., POWERS, J., WALLEN, A, and SCHMIDT, L.

Purified pyrogen from Eberthella typhosa

J. Lab. Clin. Med. 29:58-62, 1944

A concentrate from a filtrate of E. typhosa was deproteinized. It contained 1 glucosamine unit for every 5 or 6 hexose units. Total hydrolysis gave a yield of 30 per cent for reducing sugars; activity was destroyed. The minimum pyrogenic dosage (MPD) was 0.06 micrograms per kg. for the rabbit, 0.24 for the dog and 0.02 for man. The approximate lethal dose (for the rabbit) lies between 175 and 190 micrograms per kg. 15 micrograms per kg. was a toxic dose for man.

393. DELAUNAY, A., SARCIRON, R. C., and PAGES, J.

Antigenes glucido-lipidiques et pouvoir phagocytaire des cellules reticuloendotheliales (Glucido-lipidic antigens and phagocytic ability of reticuloendothelial cells)

Compt. rend. Soc. biol. 138:345, 1944

Sublethal doses of bacterial glycolipids from staphylococci and the virulent bacillus of Charbon, provoked a sharp increase resistance to bacterial invasion in the guinea pig and mouse. This may be caused by a diminution in phagocyte mobility or the preparations may act as reticuloendothelial toxins. A 'smooth' E. typhosa organism, viable, was mixed with trypan blue or azoprotein and injected intravenously in the mouse. The dosages ranged from 0.5 to 0.005 mg. and was followed by the glycolipid antigen. The reticuloendothelial cells, particularly the Kupfer cells, either contained dyes or the bacteria. It was concluded that reticuloendothelial cellular elements have no phagocytic power; if the antigen possessed the power to lower resistance to bacterial invasion, it must have done so through polynuclear diapedesis.

394. FRANKE, F. E.

Action of toxic doses of the polysaccharide from Serratia marcescens (Bacillus prodigiosus) on the dog and guinea pig

J. Nat. Cancer Inst. 5:185-193, 1944

In this study a series of unanesthetized and anesthetized dogs, and paired rats were given intravenous, intraperitoneal or

- 1 9 4 4 -

subcutaneous doses of the polysaccharide of B. prodigiosus ranging from 1 to 8 mg. per kg. of body weight in the dog, and 5.0 to 10.0 mg. (subcutaneous administration only) per kg. in the rat. Adrenalin was administered to certain animals in an attempt to prevent blood pressure depression.

Death from circulatory failure occurred in the anesthetized dog 1 to 4 hours after the administration of 0.3 to 4.3 mg. per kg. of body weight of the polysaccharide. Adrenaline was ineffective for restoration or maintenance of the blood pressure. In the unanesthetized dog, death occurred in some 12 to 24 hours after the administration of amounts between 1 and 8 mg. Necropsy revealed extensive intestinal congestion and hemorrhage. Subendocardial hemorrhage was also noted. Capillary permeability in rats and fragility in dogs approached the normal. Rise in body temperature and increased NPN were recorded for all animals.

395. FRANKE, F. E.

Attempts to produce anaphylaxis in guinea pigs with the polysaccharide from Serratia marcescens (Bacillus prodigiosus)

J. Nat. Cancer Inst. 5:173-177, 1944

Sensitizing doses of a lyophilized preparation of the polysaccharide from S. marcescens (P-21) and constant shocking doses of the material did not produce a typical picture of anaphylaxis in the guinea pig.

396. FRANKE, F. E. and RICHERT, D.

Effects of sublethal doses of a polysaccharide from Serratia marcescens (Bacillus prodigiosus) on the electrocardiogram, blood ascorbic acid and non-protein nitrogen of the dog

J. Nat. Cancer Inst. 5:179-183, 1944

Electrocardiographic studies of the dog, following the injection, subcutaneously or into the peritoneum, of 3.0 to 5.7 mg. per kg. of body weight of the polysaccharide from S. marcescens, revealed no significant changes. Development of a sinus tachycardia was noted. Four of the animals exhibited a rise in blood ascorbic acid, ranging from 0.98 to 1.86 mg. per cent. The increase in non-protein nitrogen ranged from 44 to 61.7 mg. per cent. There was considerable gastrointestinal congestion and hemorrhage. Only mild intoxication was noted in two animals. Serum calcium was within normal limits.

397. GOLDRING, W. and CHASIS, H.

Hypertension and hypertensive diseases

New York. The Commonwealth Fund, 1944

The renal pattern of response to pyrogens is shown in the maintenance of the glomerular filtration rate, the increase in renal plasma flow and decrease in filtration fraction.

Inulin may contain pyrogens and produce fever.

398. HANNAN, M., WEAVER, R. H., and SCHERAGO, M.

The relationship of pyrogen to the Dolman test for enterotoxin
J. Bact. 47:213-214, 1944

The reactions noted in kittens after pyrogen administration and those invoked by injection of staphylococcal enterotoxin were compared. Filtrates from cultures of enterotoxigenic strains of S. aureus, pyrogenic and non-pyrogenic, and from cultures of the same organism, grown on pyrogen-free or pyrogenic media, but lacking toxic properties, were employed. It was found that Dolman test filtrates contain pyrogen but that this substance was not identical with enterotoxin, nor did it affect the Dolman test.

399. LILIENTHAL, H.

Coley's mixed toxins of erysipelas and prodigiosus. Report of two cases of inoperable sarcoma treated by Coley's method

J. Mt. Sinai Hosp. 10:623, 1944

Complete disappearance of a 'pigmented' sarcoma and of an extremely malignant embryonic tumor followed treatment with Coley's toxins.

400. LIVE, I., STUBBS, E. L. and MACKEY, W. L.

The influence of intracutaneous injection of sonic filtrate of Brucella abortus on the blood leukocyte picture in cattle positive to brucellosis

J. Infec. Dis., 75:170-174, 1944

- 1 9 4 4 -

Marked changes in leukocyte distribution in the blood of cows with brucellosis appear within 6 to 7 hours after the intracutaneous injection of the sonic filtrate of Brucella abortus. A rise in immature (single-lobed) neutrophils is the most significant feature, but the total white count remains elevated for 48 hours. Leukocytosis is not seen in uninfected animals.

401. MENKIN, V.

Chemical basis of fever

Science 100:337-338, 1944

Toxic pyrogens are released from cells after injury to their structure in fever. 'Pyrexin', from the euglobulin fraction of exudates in inflammatory processes, is a nitrogenous, thermostable substance, non-toxic to mice. A non-pyrogenic, proteolytic material, 'necrosin', is found in association with 'pyrexin'; the latter may be a by-product of proteolysis. When 'pyrexin' or crude euglobulin fractions are injected into the marginal ear veins of the rabbit, an average increase in body temperature of 2.7° F. follows within 1 or 2 hours.

402. RITTER, W. L. and NUSSBAUM, M. A.

Occupational illnesses in cotton industries. I. 'Cotton Fever'.

Mississippi Doct. 22:96, 1944

The disease known as 'cotton fever', 'cotton cold', or 'card room fever' is discussed with reference to its incidence in Mississippi cotton factories. The illness, with its characteristic symptoms, attacks new employees who work in enclosed rooms and handle untreated cotton. First symptoms appear within 24 hours after the employee begins work; fever, headache, nausea and epistaxis usually disappear within a few hours. The severity of the symptoms depends on the grade of cotton handled and its dust concentration; continuous handling within a given environment produces immunity to the infection. Immunity is lost during exposure to lower grades of cotton, prolonged absence from exposure and when the concentration of cotton dust is increased. A bacterium, Aerobacter cloacae, may be the causative organism.

403. ROBINSON, E. S. and FLUSSER, B. A.

Studies on pyrogens. The isolation of pyrogens from various microorganisms

J. Biol. Chem. 153:529-534, 1944

Precipitates from suspensions of typhoid vaccine, Proteus vulgaris and Pseudomonas aeruginosa cultures, after extraction with 95 per cent phenol, dialysis and reprecipitation with acetone, contain no nitrogen. Ash remains (4 to 12 per cent). Reducing sugars are found only after hydrolysis. The pyrogen from typhoid vaccine is homogeneous, with a molecular weight of approximately 62,000.

The minimal pyrogenic dose for the rabbit has been found to be 0.5 gamma per kg. for the preparation from E. typhosa, 2.0 gammas per kg. for that from Ps. aeruginosa, and 1.0 gamma for that from Proteus vulgaris. The maximal temperature increase for typhoid pyrogen occurred when 25 to 50 gammas per kg. were injected.

404. ROSA, M. C. and de ARAGAO, R. M.

Do pirogenio (Pyrogen)

Hospital (Rio de Janeiro) 25:201-211, 1944

The occurrence of pyrogens in aqueous solutions is discussed with specific reference to testing methods. Variations in body temperature in test animals may have spontaneous generation unrelated to pyrogens.

405. SHWARTZMAN, G.

Phenomenon of local skin reactivity to Serratia marcescens (Bacillus prodigiosus). Immunological relationships between Serratia marcescens culture filtrates and Shear polysaccharide

Cancer Res. 4:191, 1944

The Shear polysaccharide and S. marcescens culture filtrates are closely related in tumor hemorrhage and necrosis-inducing properties. Serologic cross-precipitation of rabbit immune sera with the two substances and cross neutralization of provocative factors showed close antigenic relationships exist between the two, with a somewhat heightened antigenicity residing in the culture filtrate.

- 1 9 4 4 -

406. SHEAR, M. J. and PERRAULT, A.

Chemical treatment of tumors. IX. Reactions of mice with primary subcutaneous tumors to injection of a hemorrhage-producing bacterial polysaccharide.

J. Nat. Cancer Inst. 4:461-476, 1944

Normal mice given small doses of a special preparation of the polysaccharide from *S. marcescens* containing 100,000 m.t.u. per cc. developed tolerance during the administration of gradually increasing doses and survived a single injection containing 16 times the LD₅₀ dose. Tumor-bearing mice also developed resistance to lethal effects of doses 13 times the size of the LD₅₀ for the tumor-bearing mouse. When tumors were small or of medium size, dosages were tolerated which proved fatal when injected into animals with large, extensive neoplastic growths. Where tumors were less than 15 mm. in diameter, hemorrhage or necrosis was not observed. The degree of hemorrhage appears to depend on the size of the dose of polysaccharide.

An adsorbent (aluminum hydroxide suspension) was precipitated and used in the injection of polysaccharide. Hemorrhage and necrosis were regularly produced in tumors.

407. TAYLOR, R. D. and PAGE, I. H.

Studies on the mechanism of the hypotensive effect of substances eliciting leukocytosis and fever

Am. J. Med. Sci. 208:281-290, 1944

With a view toward practical therapeutic application, the principles underlying relationships between leukocytosis, hypotension and pyrogenic agents were studied in the dog with experimental renal hypertension and in humans afflicted with malignant hypertension.

Leukocytosis was induced in hypertensive dogs by subcutaneous injection of turpentine, subcutaneous and intraperitoneal injection of sterile pus, intramuscular and intravenous injection of the leukocyte-producing factor of Menkin and by 70 per cent acetylmethylamine. Terminal malignant hypertension in humans was treated by an albumin fraction from pork serum containing the buffy layer of leukocytes and an extract of human

leukocytes precipitated by $(\text{NH}_4)_2\text{SO}_4$. Mean arterial pressure of the dog was not affected by sterile pus, acetylmethylamine, or by Menkin's leukocyte-producing factor. Consistent reduction of blood pressure was not secured in malignant hypertensive patients; allergic factors were the only mechanisms clearly involved in the lower pressures achieved. The authors suggest that 'non-specific' and inflammatory reactions impair that function of the adrenal cortex which maintains vascular reactivity and this may, in certain cases, bring a reduction in arterial pressure.

408. URBACH, E., GOLDBURGH, H. L. and GOTTLIEB, P. M.

General Sanarelli-Shwartzman phenomenon with fatal outcome following typhoid-vaccine therapy

Ann. Int. Med. 20:989-994, 1944

409. YOUNG, E. G. and RICE, F. A. H.

Leukocytosis as an index of pyrogenicity in fluids for intravenous use

J. Lab. Clin. Med. 29:735-741, 1944

Leukocytosis was the most sensitive index of response to pyrogens in the dog. Progressively less sensitive were neutrophilia, leucopenia and hyperpyrexia. Leukocytosis was proportional to the amount of pyrogen in more dilute solutions. The lowest amount of pyrogen from Ps. aeruginosa required to produce a positive leukocytic response in the dog contained 0.056 gamma of N.

410. YOUNG, E. G. and HAWKINS, W. W.

The role of pyrogens in the alleged leucocytic response to allantoin

J. Pharmacol. & Exper. Therap. 81:10-16, 1944

Allantoin in Ringer's solution produced leukocytosis and neutrophilia in man when given subcutaneously or by the intravenous route. Pyrogens, and not allantoin, were the responsible agents.

- 1 9 4 4 -

411. ZAHL, P. A. and BJERKNES, C.

Effect of the endotoxin of Shigella paradysenteriae on pregnancy in rabbits.

Proc. Soc. Exper. Biol. Med. 56:153-155, 1944

12. ZAHL, P. A. and HUTNER, S. H.

Protection against the endotoxins of some gram-negative bacteria conferred by immunization with heterologous organisms

Am. J. Hyg. 39:189-196, 1944

Homologous protection by the endotoxins of 3 gram-negative organisms (S. typhimurium, Sh. paradysenteriae and Rhodospirillum rubrum) was most effective for the mouse immunized one week before injection of 4 times the LD₅₀ dose. Trichloroacetic acid extracts of Shigella and S. typhimurium endotoxins produced similar results. Heterologous protection was less efficient but still significantly protective. Toxins of 'rough' variants of bacteria may form a link between toxic nucleoproteins and the complex O antigens.

413. ZAHL, P. A. and HUTNER, S. H.

Temperature factors in the action of certain bacterial endotoxins

Proc. Soc. Exper. Biol. & Med. 56:156-158, 1944

External heat and cold did not affect the production of tumor hemorrhage and vaginal bleeding found after the injection of bacterial endotoxins in gravid mice. Histopathologic effects and temperatures produced were unrelated. It is possible that gram-negative bacteria possess endotoxins which act as neural poisons.

414. ZAHL, P. A., HUTNER, S. H. and COOPER, F. S.

Effect of sulfonamides on toxic and antigenic actions of endotoxins of certain gram-negative bacteria

Proc. Soc. Exper. Biol. & Med. 55:4-7, 1944

Limited protection against the lethal action of bacterial endotoxins is afforded by orally administered sulfonamides.

415. ALTSCHULE, M. D. and FREEDBERG, A. S.

Circulation and respiration in fever

Medicine 24:403-411, 1945

The authors present a comprehensive summary of the mechanisms involved in the cardiovascular and pulmonary manifestations of endogenous infectious fever. Cardiorespiratory functions are altered during the chill, and flush phases of fever, and during defervescence. In the prodromal stage, no significant physiologic or chemical changes occur. Sympathetic discharge initiates the chill, producing generalized vasoconstriction; with circulatory stasis of varying degree. The cardiac output falls, venous blood becomes deoxygenated, hence O_2 consumption increases. Body temperature rises. There is a possibility that stasis in the respiratory center gives rise to hyperventilation and the blood changes characteristic of alkalosis. In the flush phase, marked vasodilatation and increased cardiac output precede the acceleration of circulation time. Arterio-venous shunts open to permit equalization of blood pressure; the sudden vasodilatation brings hypotension. Hyperventilation diminishes. The circulation and respiration return to normal limits during defervescence.

416. ALTSCHULE, M. D., FREEDBERG, A. S. and McMANUS, M. J.

Circulation and respiration during an episode of chill and fever in man

J. Clin. Invest. 24:878-889, 1945

Observations on the changes following the administration of a pyrogenic agent (typhoid vaccine) revealed the presence of 4 divisions in the febrile response: prodromal, chill phase, flush phase, and defervescence. The phenomena associated have been reviewed (Medicine 24:403, 1945). It was noted that lymphatic function was greatly increased during the flush phase.

417. BANDELIN, F. J.

A leucocyte response to pyrogen in rabbits

J. Am. Pharm. A. 34:48-51, 1945

In a search for a more reliable and sensitive method for pyrogen determination, the leucocyte response of the rabbit to varying dilutions of pyrogen was selected as a standard of reference. Similar changes occur after injection of distilled,

- 1 9 4 5 -

tap and river waters. The initial leucopenia may be considered evidence of pyrogenicity; the secondary leukocytosis is corroborative. When used together they are considered to be more sensitive indicators of the presence of pyrogen than are body temperature elevations. In the rabbit, the magnitude of the leukocyte response was unaffected by solution dilution until a ratio of 1:6500 was reached; the magnitude of fever was affected by dilutions exceeding 1:3500.

418. BINKLEY, F., GOEBEL, W. F. and PERLMAN, E.

Studies on the Flexner group of dysentery bacilli. II. The chemical degradation of the specific antigen of type Z Shigella paradysenteriae (Flexner)

J. Exper. Med. 81:331-347, 1945

The specific antigenic complexes of Sh. paradysenteriae were shown to be phospholipid-acetylated carbohydrate-protein complexes grossly similar to the specific antigens of other gram-negative microorganisms. Dissociation of the antigen complex by acid yields a phospholipid, a carbohydrate and a protein fraction. Only the latter is toxic. On the other hand, alkaline degradation produces the same fractions, with only a carbohydrate possessing toxic qualities. Further studies on the complex suggest that a purine (or pyrimidine-like) derivative is associated with toxicity.

419. BONETTI, E.

Elimination of histidine and total nitrogen during fever (Translated Title)

Boll. soc. ital. biol. sper. 20:222-223, 1945

After intravenous injection of the 'pyretamine' of Russo, elimination of total nitrogen and of histidine was accelerated in the early stages of fever; excretion fell below initial levels as temperatures fell. The normal levels of excretion were resumed about 4 hours after fever disappeared. In the 12 hours thereafter some diminution of the output of both total N and histidine was seen.

420. BRADLEY, S. E., CHASIS, H., GOLDRING, W. and SMITH, H. W.

Hemodynamic alterations in normotensive and hypertensive subjects during the pyrogenic reaction

J. Clin. Invest. 24:749-758, 1945

Eight subjects, 6 of whom had essential hypertension and 2 controls with normal blood pressure held at basal levels (bed rest), received premedication with amidopyrine during the 24 hours immediately preceding pyrogen administration. Contaminated inulin and typhoid vaccine were used as pyrogenic agents. Cardiac output, arterial pressure, renal plasma flow and glomerular filtration rates were determined.

In the normal subjects, pyrogen administration produced increased cardiac output, decreased peripheral resistance and hypotension. Two hypertensive subjects exhibited blood pressure reductions to the normal range; in a third, hypotension was extreme and required special measures (Trendelenburg position); for all hypertensive subjects, hypotension to shock levels was noted in extremities. All subjects had high degrees of renal vasodilatation.

421. CAMPBELL, D. H. and CHERKIN, A.

The destruction of pyrogens by hydrogen peroxide

Science, 102:535-536, 1945

Pyrogenic solutions of gelatin were rendered nonpyrogenic by heating with KMnO_4 or H_2O_2 . Where the oxidation does not affect other constituents of the solution, the method may be applied, provided no harmful excess remains.

422. CAMEJO, E. G.

Pirogenos en las soluciones inyectables (Pyrogens in injectable solutions)

Escuela farm. 8:11, 1945

Pyrogens may be caused by the presence of killed bacteria, by bacterial proteins or they may be 'microscopic particles'. Although they may be eliminated in most cases by activated charcoal filtration, only the biologic test gives conclusive proof of their presence or absence in pharmaceutical solutions.

- 1 9 4 5 -

423. DeMERRE, J. and PROBEY, T. F.

Les pyrogens (The pyrogens)

Brussels, 1945. Belgian-American Educational Fdn., Inc. 1945

424. HARTMANN, F. L.

The use of fever therapy in the treatment of the common leukemias
Clinics 4:81-86, 1945

The employment of typhoid vaccine to invoke fever appears to enhance the effectiveness of irradiation of the spleen, lymph nodes and long bones in treating leukemia. During the febrile period, a restoration of susceptibility to X-rays is noted in patients who have become roentgen-resistant through long-continued treatment. Colchicine also increases the effectiveness of X-ray treatment.

425. KLENDISHOJ, N. C. and WITEBSKY, E.

Transfusion of O blood conditioned by the addition of blood group specific substances. Further clinical investigations

J.A.M.A. 128:1901-1093, 1945

In 1830 blood transfusions of homologous blood, compared with 389 transfusions of O blood with added group A and B specific substances, 6.5 per cent of reactions followed transfusion of homologous blood. Of these, 4.4 per cent were pyrogenic reactions. With 'conditioned' blood, reactions occurred in 4.9 per cent of transfusions; again, the majority were caused by pyrogens.

426. LOVE, J. and DRISCOLL, R. H.

Anaphylactoid (Sanarelli-Shwartzman) reaction following therapeutic anti-typhoid injections

U. S. Naval Med. Bull. 45:1104-1110, 1945

In a subject, for whom non-specific therapy had been prescribed, the second intravenous injection of typhoid vaccine proved fatal. Autopsy findings revealed generalized petechiae and ecchymoses through the parenchyma and brain; there was massive hepatic and renal necrosis. The cause of death was believed to be a generalized anaphylactoid reaction, which can be avoided through allowing a lapse of 48 hours between injections of vaccine.

427. MENKIN, V.

Chemical basis of fever with inflammation

Arch. Path. 39:28-36, 1945

'Pyrexin' a temperature-elevating fraction, has been secured by enzyme action in 'necrosin' the toxic substance present in the euglobulin of inflammatory exudates. The exudate, its fractions and 'necrosin' were injected intravenously into rabbits and compared with changes noted in rectal temperatures after injection of physiologic saline solution. The whole exudative material produced increases, regardless of the amount administered, averaging 2.37° F. Neither pseudoglobulin or albumin fractions were pyrexial. Necrosin, the euglobulin fraction, produced maximal increases (to 4.4° F.) within 6 hours, or a 276 per cent increase. The active portion of necrosin alone produced a 290 per cent increase in temperature. Nonhemolyzed canine blood serum increased temperature 67 per cent; hemolyzed blood brought these figures to 209.5 per cent. The injury to blood cells may release a pyrogenic factor and this also explains the fever-activity of malarial therapy and trauma.

'Pyrexin' has no proteolytic activity but may be released through such processes. Purified necrosin brings rapid liquefaction of thromboses. It yields the fever-producing agent, stable to heat, inactive after ashing. The latter is insoluble in ether, 95 per cent alcohol and concentrated sulfuric acid, distilled water, sodium chloride and ammonium sulfate, but is soluble in weak alkalis. In various samples, the nitrogen content ranged between 10.23 and 12.93 per cent, phosphorus from 0.79 to 1.39 per cent. It was not hydrolyzed by trypsin. The Molisch reaction was negative, the ninhydrin positive. It may be a peptide linked to a nucleic acid derivative. The mode of action shown by the substance is one of stimulation of the fever control centers of the hypothalamus. Its activity is reduced by salicylates and barbiturates.

428. MEYER, S. R.

Indicios da existencia de uma substancia antineoplasica formade mas liquidos de cultura de Penicillium notatum (Evidence for the existence of an anti-neoplastic substance as a metabolite from Penicillium notatum culture fluids)

Arq. Inst. biol. (Sao Paulo) 16:307-314, 1945

- 1 9 4 5 -

429. PROBEY, T. F. and PITTMAN, M.

The pyrogenicity of bacterial contaminants found in biologic products

J. Bact. 50:397-411, 1945

In the preparation of human blood and plasma, gram-positive and gram-negative bacilli were isolated. Most pyrogenic of these were gram-negative cocci, B. subtilis, certain corynebacterium, colon and Pseudomonas groups. Gram-positive organisms were 100 to 1000 times less pyrogenic than gram-negative organisms; the more usual reaction to the injection of such organisms was a drop in temperature. Blood serum culture medium appeared to produce organisms yielding more pyrogen. Autoclaving for 15 minutes did not destroy pyrogenicity, but, in the gram-positive bacilli preparations, such drastic treatment markedly reduced it.

430. RODNEY, G. and WELCKE, M.

Studies on bacterial pyrogens (Abstract)

J. Bact. 50:129, 1945

Bacterial pyrogens from various strains of Ps. aeruginosa, E. typhosa and B. subtilis contained 2 per cent glucosamine from which a large proportion of reducing sugars (32 per cent) could be released. All were hexoses; galactose, fructose and uronic acid were not found. Thymus nucleic acid was closely bound to the hexose. Hydrolysis did not follow treatment with lysozyme, diastases and pancreatic enzymes. The pyrogens did not appear to increase cell oxidation, although they caused body temperature in animals to rise.

431. SMITH, D. C., SHAFER, J. C. and CRUTCHFIELD, A. J.

Fever therapy with intravenous foreign protein in neurosyphilis

Southern Med. J. 38:3; 194, 1945

'With the method here described, we have available a method for producing prolonged high fever, foreign protein reaction, freedom from the untoward reactions associated with malaria and an opportunity to intensify chemotherapy'.

432. WELCH, H., PRICE, C. W., CHANDLER, V. L. and HUNTER, A. C.

The: thermostability of pyrogens and their removal from penicillin

J. Am. Pharm. A. 34:114-118, 1945

Adsorptive filtration with asbestos filter pads will remove the thermogenic substance from growing penicillia, but temperatures of 250° C. for 30 minutes are required to destroy bacterial pyrogens. Ampoules may not be pyrogen-free; all other glassware used in tests should be subjected to 250° C. heating for 40 minutes.

433. ZAHL, P. A., STARR, M. P. and HUTNER, S. H.

Effect of bacterial toxins on tumors. VII. The tumor hemorrhage factor in bacteria

Am. J. Hygiene 41:41-56, 1945

Tumor hemorrhage production was the criterion used for 38 strains of 25 species of plant pathogens. Suspensions of cultures employed included those prepared from Agrobacteria, corynebacteria, Pseudomonas and xanthomona strains. Of the 32 types of streptococci in this study which were gram-positive, it was found that a factor bound in the O antigen of gram-negative organisms is lacking. This factor has particularly toxic affinity for vascular structures. Some connection exists between hemorrhage production and lethality and the 'smooth' type of Salmonella choleraesuis; potency of this type is 10 times that of 'rough' strains.

434. ZITTLE, C. A., DEVLIN, H. B., RODNEY, G. and WELCKE, M.

Removal of bacterial pyrogens from protein hydrolysates

J. Lab. Clin. Med. 30:75-79, 1945

Heating to a temperature of 250 deg. F. for 30 to 40 minutes will destroy pyrogens; or, in penicillin solutions, they may be removed by adsorptive filtration with asbestos pads.

- 1 9 4 6 -

435. AGUILAR, R. M., CORTINA, M. C. and VILLAPARD

Pirogenes y transfusiones de Sangre (Pyrogens and Blood transfusion. Preliminary report.)

Ann. med. Mexico 7:1-6, 1946

436. ALGIRE, G. H.

Effect of a bacterial polysaccharide and of tourniquet shock on peripheral capillary circulation in unanesthetized mice

Federation Proc. 5:217, 1946 (Abstract)

In vivo microscopic studies were made of the effects of a bacterial polysaccharide from B. prodigiosus and of tourniquet shock on peripheral capillary circulation in the unanesthetized mouse. A transparent chamber introduced into a dorsal skin flap permitted daily measurements and photographic studies to be made. In both shock and the reaction to polysaccharide the capillary blood supply decreased to striated muscle and was followed by stasis, rouleaux formation and occlusion. Capillary dilatation was not seen. After polysaccharide injection, edema in subcutaneous tissues was present, suggesting increased capillary permeability. Edema after tourniquet pressure apparently did not cause increased permeability in peripheral tissues.

437. ALGIRE, G. H.

The vascular reaction of normal and neoplastic tissues to a bacterial polysaccharide from Bacillus prodigiosus culture filtrate

Cancer Res. 6:491, 1946 (Abstract)

Changes in the vascular bed of normal striated muscle and of neoplastic tissue were measured through the transparent chamber method. A decrease in the capillary supply to the panniculus carnosus layer of striated muscle and to transplanted sarcoma tissue was accompanied by vascular occlusion. Maximal occlusion was seen in three to four hours after injection of B. prodigiosus polysaccharide; recovery of capillary function followed within 18 hours in survivors. Neither hemorrhage nor necrosis was observed in normal tissues. In sarcoma transplants, occlusion was followed, in three to four hours, by petechial hemorrhage. Necrosis developed. The few tumor cells which survived retained the capacity to obtain a new vascular supply and thereby resume growth.

438. BECK, L. V.

The Shear tumor-necrotizing bacterial polysaccharide as a pyrogen
Federation Proc. 5:6, 1946 (Abstract)

After a maximum rise in temperature of 2 to 3° C. was produced by the polysaccharide from S. marcescens, increased amounts of the substance did not bring about further increases. A weak fever reaction was shown with lethal and semi-lethal dosages (20 to 100 micrograms per kg.), and death usually occurred within 24 hours after administration. An extremely rapid and extensive decrease in ear temperature accompanied fever; skin temperatures appeared to parallel those of the rectum. When rabbits were tied down in the copper trough of an animal table, rectal temperature was held to near normal levels; ear temperatures remained below normal. The fever reaction was not altered appreciably when animals were restrained on cardboard. It required 500 mg. per kg. body weight of acetyl salicylic acid to counteract the fever reaction. Isopyrene was more effective (150 mg. per kg.) while dial, 40 mg. per kg. intravenously was the most active antipyretic agent.

439. BECK, L. V., and FISHER, M.

Physiological Studies on tumor-inhibiting agents. I. Effect of S. marcescens polysaccharide on rectal temperatures in normal and in tumor-bearing mice

Cancer Res. 6:489, 1946 (Abstract)

In the normal rabbit 0.005 micrograms per kg. of the polysaccharide from S. marcescens produces a measurable fever when injected intravenously; 0.5 micrograms per kg. provokes high fever. The largest amount of polysaccharide administered to the rabbit was 100 micrograms per kg. body weight. No other pyrogenic substance has been found to possess so intense a pyretogenic action for the rabbit.

In the mouse, relationships are not clearly defined between tumor-necrotizing activity of the polysaccharide and its pyrogenic action. In these experiments, both normal and tumor-bearing mice received a total dose of 0.10 to 10 micrograms (0.4 to 400 micrograms per Kg.) intravenously and 0.5 to 10 micrograms intraperitoneally. Larger amounts of polysaccharide produced no increases in rectal temperatures; subnormal temperatures were noted, which were more pronounced and enduring in the tumor-bearing mouse. The larger the tumor the greater were such temperature decreases and the reduction in the chance of survival.

440. BECK, L. V. and FISHER, M.

- 1 9 4 6 -

Physiological studies on tumor-inhibiting agents. II. Effect on rectal temperatures in normal rabbits of the Serratia marcescens tumor-necrotizing polysaccharide of Shear

Cancer Res. 6:410-420, 1946

Fever of moderate grade was usually produced by the injection of 0.5 micrograms per kg. body weight in the rabbit. Diarrhea generally accompanied the elevation of temperature. The LD₅₀ dose for the rabbit exceeds 20 micrograms per kg. but is probably not above 100 micrograms. For the guinea pig LD₅₀ dosage is from 100 to 200 micrograms per kg., and that for the dog appears to be of the same magnitude. The normal mouse tolerates 100 micrograms per animal, but tolerance is greatly reduced (to 15 micrograms per mouse) with subcutaneous tumors. The polysaccharide acts through the central nervous system and not the peripheral areas in the animal; this is shown by the minute amounts which will produce fever, counteraction of fever by antipyretics and extreme vasoconstriction noted after administration. The febrile reaction may be influenced by the environmental temperature. When the spinal cord is cut, the drop in temperature ~~advances~~ at a rate of over 2° C. per hour.

441. BEESON, P. B.

Development of tolerance to typhoid bacterial pyrogen and its abolition by reticuloendothelial blockade

Proc. Soc. Exper. Biol. & Med. 61:248-250, 1946

Tolerance develops when daily injection of typhoid pyrogen is continued for a considerable period of time. It is believed that an accelerated rate of disposal of the foreign substance may underlie the phenomenon. The reticuloendothelial system is presumed to be closely connected with this mechanism, since tolerance development is prevented by reticuloendothelial blockade.

442. BRINDLE, H. and RIGBY, G.

The preparation of non-pyrogenic water and infusion fluids, using activated charcoal

Qtly. J. Pharm. Pharmacol. 19:302-339, 1946. See also: Pharm. J. 157:85, 1946

Untreated waters from Manchester water mains or from a Cheshire stream were agitated at intervals for 15 minutes with charcoal (0.1 per cent) and filtered through sintered glass. Pyrogenic activity, as shown by production of fever in the rabbit, was much reduced. The average rise in temperature was at least 1.5° C. before treatment, and 0.3° C. after. Strongly polluted water could not be rendered pyrogen-free if storage preceded treatment. Heavily contaminated pharmaceutical solutions required more than 0.1 per cent charcoal. There was little adsorption of medicaments on charcoal; procaine HCl solution was an exception with adsorption of 27 to 35 per cent.

443. DILLER, I. C. and SHEAR, M. J.

Cytological effects of S. marcescens polysaccharide on tumors
Cancer Res. 6:488, 1946 (Abstract)

A single injection of the polysaccharide isolated from S. marcescens caused striking degenerative changes and damage to cell nuclei in mouse sarcoma 37. Alterations were evident within 2 hours after the injection; cells appear to be most strongly affected during the division stages. The impact of the polysaccharide was most clearly seen in the period of mitotic activity, and at four hours after administration of the agent no division could be detected. When undamaged cells were seen after 6 hrs., they were in the resting phase. Normal tissues which divide at a rapid rate (intestinal epithelium) were damaged by the polysaccharide. Spontaneous tumors of the rat and mouse and neoplastic tissue from the human patient were examined after polysaccharide therapy. The nuclear changes were similar in all material studied, but the degree and extent varied.

444. DUNN, T. B. and LEHMANN, S.

- 1 9 4 6 -

Cancer Res. 6:488, 1946 (Abstract)

Transplanted epithelial tumors of the mouse appeared to be unchanged on gross examination 24 hours after a single injection (0.1 mg.) of S. marcescens polysaccharide. Microscopic examination revealed extensive necrosis in the five types of epithelial neoplasms observed: adenocarcinoma of the small intestine; myoepithelioma of the salivary gland; pulmonary tumors and two squamous cell carcinomas. The effect of bacterial polysaccharides on carcinomas may be overlooked unless microscopic studies are made.

445. GLUCK, R. and WILSON, J. L.

Fever produced by the oral administration of amino acids

Am. J. Dis. Child. 71:601-610, 1946

Fever from an undetermined cause appeared in infants ill from excessive dietary intakes of monosaccharides and casein hydrolysates in sterile solutions. Six well infants had sudden elevations of temperature on the second day of feeding; normal levels were resumed within 24 hours. The fever was not associated with dehydration nor was it influenced by an antibiotic (sulfadiazine). The work of Reitschel and of Holt is recalled. A specific dynamic action of amino acids and not a pyrogenic effect may be responsible for "oral" fevers.

446. FAVORITE, G. O. and MORGAN, H. R.

Therapeutic induction of fever and leucocytosis, using a purified typhoid pyrogen

J. Lab. Clin. Med. 31:672-676, 1946

When a purified pyrogenic antigen isolated from E. typhosa is injected intravenously fever, profound leukocytosis, localized pain in the head, back and legs, and occasionally, nausea and vomiting follow. Symptoms disappear, leaving a sense of fatigue and exhaustion. The leukocytosis produced is characterized by the appearance of many granulocytes of the non-filamentous type; glucose, chloride and CO₂-combining power of the blood vary little from the norm. The leukocytosis was considered the most valuable aspect of pyrogen therapy as shown in the treatment of neurosyphilis.

447. KOCH, P. K. and OLITZKI, L.

The action of dysentery toxins on different laboratory animals

Exper. Med. & Surg. 4:54-68, 1946

Two variants of S. dysenteriae were examined, and both a neurotoxin and thermostable lethal factor were isolated. In the S-variant a thermostable factor, lethal for mice and small animals, is responsible for hemorrhage of the cecum (rabbit) when injected with the neurotoxic principle present in both S- and R-variants. Other thermostable, toxic materials may be involved in the changes seen after administration of S. dysenteriae; these changes include hypothermia, decreased liver glycogen output, leucopenia and decreases in ascorbic acid in the adrenals.

448. KUNA, S., EDISON, A. D. and BUTZ, C.

A method for large scale testing for pyrogen

J. Am. Pharm. A. (Sci. Ed.) 35:59-63, 1946

Temperature elevations may be observed in the rabbit after handling. A thermocouple box method has been devised which eliminates the possibility of error from such factors. Other procedures are outlined which may increase sensitivity in the rabbit. Thermogenic response is intensified when low initial temperatures exist.

449. LEWIS, L. A. and PAGE, I. H.

Method of assaying steroids and adrenal extracts for protective action against toxic material (typhoid vaccine).

J. Lab. Clin. Med. 31:1325-1329, 1946

Since some protective ability had been previously demonstrated for adrenal extracts against the toxic effects of typhoid vaccine, an assay method was devised to determine the factor or factors responsible for such protection. Adrenalectomized rats were given a test dose of typhoid vaccine which was predetermined to be the MLD for the species. Various adrenal extracts and other steroid compounds were administered once or twice daily for three days before the vaccine injection. Survivors were sacrificed after 21 days. The efficacy of 5% glucose in maintaining blood sugar levels and the protection afforded against the toxicity of typhoid vaccine were additional features of the study. Of eleven steroids tested, 11-dehydro-17-hydroxycorticosterone was the most powerful protective agent, but some capacity for protection was shown by all compounds with an O linkage in the 11-position. No differences in potency could be demonstrated between natural and synthetic Compound A acetates. Maintenance of blood glucose alone proved to be ineffective as a protective measure.

- 1946 -

450. MACPHERSON, C. F. C., HEIDELBERGER, M, ALEXANDER, H. E. and LEIDY, G.

The specific polysaccharides of types A, B, C, D and F, Hemophilus influenzae

J. Immunol. 52:207-219, 1946

The chemical, physical and immunological properties of preparations of specific carbohydrates of H. influenzae are described.

451. MILLER, C. F. and BOOR, A. K.

Protection of mice against the lethal action of gonococcal endotoxin by penicillin

Proc. Soc. Exp. Biol. & Med. 61:18-20, 1946

Crude gonococcal endotoxin was injected in graded doses into the peritoneum of the mouse. Subcutaneous doses of 1500 units each of penicillin were given at intervals ranging from 90 minutes before to 24 hours after administration of the endotoxin. Seventy-six per cent of all control mice died in comparison to 36.5% when penicillin was employed.

452. NAUTS, H. C., SWIFT, W. E. and COLEY, B. L.

The treatment of malignant tumors by bacterial toxins as developed by the late William B. Coley, M. D., reviewed in the light of modern research

Cancer Res. 6:205-216, 1946

The authors review the use of bacterial toxins in the treatment of malignancies, as reported in the literature of the past 50 years. The basic principles underlying the choice of these agents and their place in the modern therapeutic armamentarium against neoplastic disease are given in detail. A careful appraisal is made of 'Coley's toxins' and similar agents used in tumor therapy, and errors in the earlier application of these substances are judged in the light of recent developments. More scientific study is indicated. "...toxin therapy has clinical value and...further extensive research is warranted to produce better preparations and further refinements in the technic of administration..."

A bibliography containing 80 references is appended.

453. OLITZKI, L., CZACZKES, J. W. and KUZENEK, A.

Endotoxic factors of Rickettsia Prowazeki and their immunological relationship to the endotoxins of other gram-negative micro-organisms

J. Immunol. 53:365-370, 1946

Hypothermia and decreased liver glycogen follows the intrabdominal injection of heat-killed Rickettsial organisms. To determine whether or not immunologic relationships exist between these factors in R. Prowazeki and in certain other gram-negative bacteria (Proteus, S. dysenteriae) and two gram-positive bacteria, Corynebacterium eros and Mycobacterium butyricum. Any of the three gram-negative organisms elicited immunization against hypothermia and decreased liver glycogen produced by either of the other two. This is conceded to be "a specific immune reaction directed against a toxic product which is a common constituent of a large group of microorganisms." The gram-positive organisms used did not elicit hypothermia or immunity against temperature reduction.

454. OLITZKI, L., SHELUBSKY, M. and HESTRIN, S.

Pathogenizing effect of different carbohydrates on Eberthella typhosa

Proc. Soc. Exper. Biol. Med. 63:491-496, 1946

In determining the chemical basis for the pathogenizing activity of certain carbohydrates, mice were injected with the test organism (E. typhosa) intraperitoneally and with a sterile aqueous solution of the tested pathogenizing substance in varying concentrations. By this method 15 polysaccharides and 11 simple sugars and oligosaccharides were examined. Three classes of carbohydrates were distinguished: (1) powerful pathogenizing sugars, levan, dextran, mucin and agar; (2) moderate pathogenizing in two cellulose preparations, in gum acacia, glycogen, mannan, and inulin; (3) non-pathogenizing crystalline sugars and the starch and galacto-mannan of the polysaccharides. Pathogenizing ability was enhanced by addition of kaolin to the solution injected. The pathogenicity of levan, produced in vitro, may be an intrinsic property. It is suggested that specific colloidal patterns, rather than polar groupings or individual unit structure, may underlie pathogenizing activity.

455. PAINE, T. F.

Illness in man following inhalation of Serratia marcescens

- 1 9 4 6 -

J. Infect. Dis. 79:226-232, 1946

Four adult males, two of whom had histories of previous exposure, were subjected to an aerosol concentration of S. marcescens of approximately 2 million per cubic foot of air for two and one-half hours. Blood, nasal secretion and sputum cultures were made, serum agglutinins determined and blood changes recorded. Two and one-half to four hours after the exposure all subjects became acutely ill. Ocular and respiratory symptoms appeared. Chills and fever, tachycardia and generalized myalgias were present. Subjective improvement appeared after diaphoresis and defervescence. No significant changes in blood or urine components were noted, but sputum and nasal secretions produced positive cultures. Marked neutrophilia appeared at the onset of the reaction. Specific serum agglutinins were present in high titer. Previous exposure appears to be immunologically significant.

456. REIMANN, S. P., HOLLOMAN, A. L., OAKLEY, R., McCONNELL, J. R. and SHEAR, M. J.

Gross and microscopic pathology of human tumors following administration of S. marcescens polysaccharide

Cancer Res. 6:489, 1946 (Abstract)

Without previous sensitization, the injection of a polysaccharide prepared from B. prodigiosus produces necrosis and hemorrhage in both animal and human tumors. Intravenous injection of the polysaccharide in a single dose or in divided doses, was followed in a few hours by swelling in the tumor area. For several patients, a constant intravenous drip of the polysaccharide in solution was used and, for others, intraperitoneal injections. Diffuse ecchymosis appeared within 48 hours in tumors near the surface. The size of the tumor is believed to be a determining factor in judging the amount of necrosis and hemorrhage produced by a given dosage. Small neoplasms showed more extensive necrotic changes. Microscopically, the hemorrhages were seen to be focal in most cases, and the necrosis was of the liquefaction type.

The distinction between hemorrhage from polysaccharide and that occurring spontaneously in softer sarcomas is often a finely drawn one. Fibrosarcomas show little or no hemorrhage, hence presence of these changes after polysaccharide injection may be considered a positive confirmation of polysaccharide activity.

457. TODD, J. P., MILNE, G. R. and LAURIE, J. P.

Non-pyrogenic saline infusion fluids

Chem. and Druggist 145:536, 1946

The definitions of "pyrogen-free" as a term for solutions are reviewed and standards are discussed. Pyrogen-free water is easier to prepare than pyrogen-free saline solution, and the authors conclude from such findings that NaCl per se is pyrogenic.

458. TODD, J. P. and MILNE, G. R.

Tests on nonpyrogenic fluids

Pharm. J. 156:157-158, 1946

The skill of the operator is considered to be highly important in the production of pyrogen-free intravenous solutions. Pyrogen tests on tap water, distilled water, normal saline and dextrose saline solutions are reviewed.

459. TODD, J. P., MEARNS, J. T. and MILNE, G. R.

Preparation of nonpyrogenic saline infusion fluids

Pharm. J. 156:158-159, 1946

The pH of dextrose-saline solutions bears an important relationship to certain changes which lower standards set for such fluids.

460. TONNESEN, M. and VESTERDAL, J.

The pyrogenic effect of Danish penicillin (in English)

Acta Pharm. Toxicol. 2:47-68, 1946

These experiments were performed to determine whether the pyrogenic effects of Danish penicillin could be tested in a manner similar to that adopted in the U.S.P. XII. In rabbits, rectal temperature increased after injection were used as a measure of pyrogenicity; the same preparations were tested simultaneously in therapeutic doses on both normal humans and those with gonococcal infections. Removal of pyrogen from the preparations was attempted. With the test used in the United States, it was noted that penicillins meeting the requirements did not usually cause febrile reactions in humans. A marked reduction in pyrogen concentrations could be obtained by filtration through the Seitz equipment. Guinea pigs had no febrile responses to penicillin. A large number died within 4 days from toxic conditions which could not be correlated to the presence or absence of pyrogens.

- 1 9 4 6 -

461. TUCKER, J.

Typhoid shock therapy

Cleveland Clin. Quart. 13:67-72, 1946

Experiences with fever elicited by typhoid vaccine in several thousand patients during the past 15 years have been useful for determining correct applications and contraindications for typhoid shock therapy. Most successful results have been realized in the following conditions: central nervous system virus infections, chronic rheumatoid arthritis, chorea, and eye and skin diseases.

462. WARREN, M. R. and WERNER, H. W.

The evaluation of antipyretics against pyrogen-induced fever

J. Am. Pharm. A. 35:257-259, 1946

A purified pyrogen was prepared from Proteus vulgaris by cellular lysis, repeated phenol extraction, dialysis and acetone precipitation. The minimal pyrogenic dose was 0.5 cc. per kg. body weight in the rabbit when the pyrogen was injected subcutaneously. Temperature was reduced to normal levels by the oral administration of aminopyrene, acetanilid or acetylsalicylic acid; acetophenetidin arrested temperature increase but did not reduce fever.

463. WEENS, H. S. and HEYMAN, A.

Cardiac enlargement in fever therapy induced by intravenous injection of typhoid vaccine

Arch. Int. Med. 77:307-316, 1946

Enlargement of the heart during typhoid vaccine therapy was noted in 8 of 15 patients with neurosyphilis. In four other patients, cardiac changes persisted for six months following hyperpyrexia. No significant differences could be found between patients with normal cardiac size and those with enlarged structures. Blood volume, cardiac output, and protein and hemoglobin levels did not differ widely from the norm.

464. WESTFALL, B. B. and DUNN, T. B.

Blood uric acid and proteose, body temperature and glomerular clearance of rabbits implanted with the Brown-Pearce carcinoma and treated with the polysaccharide from Serratia marcescens

J. Nat. Cancer Inst. 7:115-121, 1946

Blood proteose and uric acid, creatinine and creatinine clearance were determined in adult male rabbits after injection of a single dose of S. marcescens polysaccharide. All animals bore Brown-Pearce carcinomas at least three weeks old. The range of dosage extended from 5.0 to 300 micrograms per kg. body weight and all animals tolerated 50 micrograms per kg. At this level, no significant rise in blood proteose occurred, There was a slight increase in blood uric acid with accompanying decrease in glomerular filtration. Hemorrhagic changes in tumor tissue were similar to those which occur spontaneously and were therapeutically unimportant. Larger dosages, in the presence of proliferating tumors, brought peripheral vascular collapse; at the 50 microgram per kg. level, tumor hemorrhage, ascites, decrease in renal clearance, collapse and death followed in sequence.

465. WRETLIND, K. A. J.

About the pyrogen activity of dialyzed casein digest

Acta Physiol. Scand. 12:385-388, 1946 (in English)

Temperatures of rabbits given aminosol (3.3% solution) by the intravenous route rose an average of 0.41° C. within one hour. The solution was used alone and with glucose additions. In the third hour, the increased temperature still existed to a lesser degree (0.28° C.).

466. ZINOV'EVA, M. T.

Mineral metabolism. Calcium level in arterial and venous blood in all kinds of fever (Translated Title)

Byull, Ekspt. Biol. Med. (Leningrad) 21:70-72, 1946

Fevers prolonged two days are accompanied by a drop in both the arterial and venous blood calcium content. In fever induced through the injection of metabolism-heightening chemicals (as dinitrophenol), calcium rises early and falls to normal with the temperature. In pyrexia produced by amine toxins (from putrified meat) the calcium of blood rises only after the reduction of temperature from fever levels.

- 1 9 4 7 -

ALGIRE, G. H., LEGALLAIS, F. Y. and PARK, H. D.

Vascular reactions of normal and malignant tissues in vivo.
 II. The vascular reactions of normal and neoplastic tissues
 of mice to a bacterial polysaccharide from Serratia marcescens
(Bacillus prodigiosus) culture

J. Nat. Cancer Inst. 8:53-62, 1947

The polysaccharide fraction P-3 was injected intraperitoneally (0.2 mg. and 0.02 mg.) into mice bearing transparent chambers in dorsal skin flaps. Daily quantitative measurements were made of the vascular bed in normal striated muscle and neoplastic tissues prior to and following tumor transplantation. The capillary blood supply was traced by photographs, as were the changes preceding hemorrhage and necrosis and the pattern of regrowth when it occurred.

In both normal and neoplastic tissue, a period of vascular stasis preceded hemorrhage. Occlusion of capillaries was observed in both tumor and normal striated muscle tissue; edema was present. Where the tumor host survived, the surviving tumor cells began new growth with capillary proliferation. The mechanism through which hemorrhage and necrosis takes place is closely related to tissue anoxia and capillary damage.

BARSI, I.

A new treatment of rheumatoid arthritis

Brit. M. J. 2:252-253, 1947

A febrile reaction was seen after the first infusion of 300 ml. of blood (from pregnant women) into male and female patients. Improvement of rheumatoid arthritis followed in 64% of the group; relief from symptoms was spectacular. The possibility that a mild "protein" shock initiates benefits in this therapy is discussed.

BECK, L. V., DILLER, I. C., BLAUCH, B. and FISHER, M.

Reduction in toxicity of Serratia marcescens polysaccharide to tumor-bearing mice produced by beef adrenal extract (Upjohn)

Cancer Res. 7:725, 1947 (Abstract)

When large doses of Shear's polysaccharide (S. marcescens) are required to produce hemorrhage and necrosis in extensive or resistant tumor transplants in mice, toxicity is the limiting factor. The preparation P-8 (National Cancer Institute) was lethal in

- 1 9 4 7 -

500 gamma doses for 64 of 75 mice within 48 hours. All mice bore 7-day transplants of sarcoma 37. A second group, bearing neoplasms of the same age, were given beef adrenal extract prior to administration of P-8 polysaccharide and the same amount (0.25 cc.) at the end of the working day. Of these, 22 in a group of 45 survived 48 hours or longer. The possibility of this difference being the result of chance was estimated to be about 1 in 10,000. A concentrated hog adrenal extract in oil was ineffective in counteracting the lethal effects of P-8 polysaccharide.

470. BEESON, P. B.

Tolerance to bacterial pyrogens. I. Factors influencing its development

J. Exper. Med. 86:29-38, 1947

Repeated daily intravenous injections (7 to 10 days) of pyrogens from E. typhosa, B. prodigiosus and Ps. aeruginosa produces a gradual diminution of febrile response. Tolerance develops and the increase in body temperature produced is slight and it remains constant. Tolerance is lost after a period of rest and pyrogens again produce higher febrile levels, on resumption of injections. No passive transfer of tolerance could be confirmed. Cross tolerance was noted. Antipyretics did not alter the course of development of tolerance.

Amidopyrine, administered through a stomach tube 1 hr. before injection of typhoid vaccine, prevented febrile response. Cabinet fever therapy did not reduce responsiveness to pyrogens.

471. BEESON, P. B.

Tolerance to bacterial pyrogens. II. Role of the reticuloendothelial system

J. Exper. Med. 86:39-44, 1947

Tolerance to pyrogens in rabbits may be the result of stimulation of the reticuloendothelial system. After tolerance was established, febrile response could be heightened through blockade of the reticuloendothelial system. Tolerance develops, according to the hypothesis presented, when ability to remove pyrogens (or other bacterial toxins) from the blood is enhanced through awakening of the defense mechanism. Protection is thus given against tissue injury.

472. BEESON, P. B.

- 1 9 4 7 -

Effect of reticuloendothelial blockade on immunity to the Shwartzman phenomenon

Proc. Soc. Exper. Biol. Med. 64:145-149, 1947

When Thorotrast injections followed typhoid vaccine, rabbits naturally immune (or rendered immune through previously invoked Shwartzman reactions) developed areas of hemorrhagic necrosis, and the toxic effects of the vaccine were enhanced.

473 BERNSTEIN, C. and KLOTZ, S. D.

Fever therapy in herpes zoster

J. Lab. & Clin. Med. 32:1544, 1947

Mixed typhoid vaccine injected intravenously brought prompt relief from true herpes zoster. There was an early disappearance of pain.

474 BERTRAND, I. and QUIVY, D.

Le probleme des pyrogenes. Son état actuel (Problem of the pyrogens. Actual state)

Rev. d'Heat. 2:234-240, 1947

The chemical structure of the pyrogens has not yet been fully explained. Carbohydrates are present in relatively large amounts but the nitrogen content is negligible. Polysaccharides are present. The pattern of molecular form should be studied further.

Methods of testing for pyrogenic contaminants and the role of equipment (glass and tubing) in harboring pyrogen-producing bacteria are discussed. In transfusion of blood or infusion of fluids, the rate of infusion may, if accelerated, produce dangerous pyrogenic reactions when contaminants are present in stored blood or solutions.

475 BLACK-SCHAFFER, B., HEIBERT, T. G. and KERBY, G. P.

Experimental study of purpuric meningococcemia in relation to the Shwartzman phenomenon

Arch. Path. 43:28-54, 1947

Twice-washed meningococci, living or dead, possess preparatory and provocatory substances with ability to induce the Shwartzman phenomenon. The toxic materials were present in small amounts only in the washings. Potency was highest in strains producing

purpuric meningococcemia. Bilateral renal necrosis, found after injection of living organisms, was believed to be the product of vasospasm and the thrombosis it engendered.

476. BOQUET, P., DELAUNEY, A., LEHOULT, Y. and LEBRUN, J.

Action d'un extrait du bacille typhique sur la pression artérielle du lapin (Action of an extract of Bacillus typhosa on the arterial pressure of the rabbit)

Compt. rend. Soc. biol. 141:269-271, 1947

Marked, rapid and sustained peripheral vasoconstriction followed injection of B. typhosa endotoxin in the rabbit. Two to 16 mg. of the toxic substance were injected by subcutaneous, intra-peritoneal or intravenous routes. The carotid pressure remained constant during the toxic period; in extremis, a reduction occurs, as would be expected.

477. BOQUET, P., DELAUNAY, A., LEHOULT, Y. and LEBRUN, J.

Observation directe des reactions vasculaires cutanées chez le lapin soumis a l'épreuve d'une endotoxine typhique (Direct observations of the cutaneous vascular reactions in the rabbit subjected to typhus endotoxin)

Compt. rend. Acad. Sci. 225:1193-1194, 1947

The reactions of cutaneous vessels in the ears of rabbits were studied by means of window-flaps and microscopic examinations. Contraction of a severe degree occurs 5 to 15 minutes after intravenous or intraperitoneal injection of 6 to 8 mg. of a B. typhosa polysaccharide-lipid-polypeptide antigen; the cycle of alternating dilatation and contraction which follows is confined to arterioles alone. Removal of the upper cervical ganglion did not alter the response. The initial response to purified B. typhosa antigen is believed to be independent of local sympathetic innervation and to resemble similar reactions observable in anaphylactic and traumatic shock. Leucopenia and inhibition of diapedesis accompany this manifestation of intoxication.

478. BRADLEY, S. E. and CONAN, N. J.

Estimated hepatic blood flow and bromsulfalein extraction in normal man during the pyrogenic reaction.

J. Clin. Invest. 26:1175, 1947 (Abstract)

The cardiac output and renal blood flow are increased after

- 1 9 4 7 -

pyrogen administration even when fever is prevented by anti-pyretic drugs. Hepatic circulation in normal human subjects before, and after, intravenous injections of typhoid vaccine was traced through bromsulfalein clearance and hepatic venous cateterization. The blood flow in the liver increased in all subjects; to 300 per cent increase was noted in one individual. Despite hyperemia and increased bromosulfalein loading, the latter was removed from blood at a rate 50% below normal levels. Hepatic venous pressure rose occasionally as hyperemia developed in the liver.

474. BRAGDON, J. H.

The hepatitis of hyperthermia

New England J. Med. 237:765-769, 1947

Liver damage may be seen after fever is induced by physical or biological methods. Hepatitis, believed to be of virus origin may be activated through increased body temperatures. The critical temperatures above which liver cells are irreversibly changed are probably varied, but are believed to be not far above those encountered frequently. Anoxia, or the destruction of functioning enzyme systems, may underlie liver cell necrosis; the same factors are destructive to certain central nervous system cellular components.

480. BUENO, P.

Reticuloendothelial cells reacting to toxic antigens and to infection

Arch. Path. 44:635-638, 1947

Nuclear tumefaction and disintegration in the undifferentiated mesenchymal cells, in germinal elements of lymphatic tissues and lymph nodes, and the Kupffer cells of the liver of the guinea pig followed subcutaneous or intracardiac injection of the antigens of paratyphoid A., brucellosis, anthrax, tetanus and diphtheria organisms.

Close relationships exist between sites of antigen activity and those concerned with antibody formation.

485. CHAMBERS, W. W. and WINDLE, W. F.

Site of action of a bacterial pyrogen in cats with central nervous system lesions

Federation Proc. 6:89, 1947 (Abstract)

In the normal cat injection of a purified pyrogenic extract obtained from Pseudomonas aeruginosa (PYROMEN) produced pilo-erection, chills and fever. Only fever occurred when the pyrogen was administered to cats with chronic hypothalamic lesions and no response was registered in the chronic spinal animal. The last-named show poor adjustment to environmental temperature changes.

482. COLLIER, H. O. J. and PARIS, S. K.

The decline of pyrogenic activity of intravenous solutions during storage

Quart. J. Pharm. Pharmacol. 20:376-380, 1947

It has been reported that strongly pyrogenic solutions may lose their activity during storage. To test the validity of this theory, solutions of ten compounds commonly used for intravenous therapy were stored at reduced pressures (508-635 mm. Hg.) and tested for pyrogen content at intervals. Solutions included saline, dextrose M/6 sodium lactate. Darrow's solution, Hartmann's and sodium sulphate solutions.

The average span of pyrogenic activity was 92 days or less; in one instance, pyrogens were present after 110 days storage. Temperature and light were relatively unimportant factors in the storage life of pyrogens; initial concentration of pyrogens in solution was less significant than the chemical nature of the pyrogen under consideration and the presence of stabilizing substances. Two solutions of the same type may contain pyrogens widely dissimilar. The rate of loss of pyrogenicity varied widely. In general, dilute solutions were less stable for pyrogen retention, but it was noted that such substances can exist in solid materials, eg. dextrose. It was also noted that the febrile activity of pyrogens wanes after long storage.

483. CONDINI, P.

Amora sulla bactererterapia dei tumori (Bacteriotherapy of tumors)
Terapia (Milan) 32:69-70, 1947

484. CREECH, H. J., HAMILTON, M. A., DILLER, I. C., NISHIMURA, E. T. and SHEAR, M. J.

Comparative studies of the immunological, toxic and tumor-necrotizing properties of S. marcescens polysaccharides

- 1 9 4 7 -

Fractionation of the polysaccharide from one strain of S. marcescens has yielded a product which appears to be less toxic and antigenic in the mouse. No significant alterations of the tumor-necrotizing property occurred. Studies have been made of the protection afforded by passive immunization of mice against the toxic fraction of the polysaccharide from the "GW" strain. The gamma globulin fraction of rabbit antisera, bearing antibodies, protected a large percentage of the animals when injected a few hours before the injection of a lethal dose of polysaccharide. Previous injection of the fraction into mice bearing sarcoma 37 prevented toxic effects of a large dose of S. marcescens polysaccharide but did not interfere with tumor necrosis. A strain of S. marcescens (724) has been found to be capable of producing less toxic polysaccharides.

485. DILLER, I. C.

Degenerative changes induced in tumor cells by Serratia marcescens polysaccharide

Cancer Res. 7:605-626, 1947

A detailed account of the cellular changes induced in tumors by the polysaccharide of S. marcescens is presented. Studies were carried out on mouse sarcoma 37, on spontaneously appearing rat and mouse carcinoma and on human biopsy material taken from 16 carcinoma patients. The combined effects of polysaccharides and roentgen therapy on neoplastic destruction were observed.

Two effects of polysaccharide activity were noted; one, a vascular change, the other, one of nuclear destruction. Maximal damage was seen within six hours. There was marked inhibition of resting cell division for three days after treatment. In some animals no viable cells were detected three days after treatment with a dose lethal for one-fifth to one-third of the mice. Roentgen therapy increased three-fold the number of tumors which showed regression. Injection directly into the tumor did not inhibit growth. Resistance to the necrotizing effects of polysaccharide developed in the tumor host after repeated injections. Human tumors underwent similar but less pronounced destructive changes.

486. DILLER, I. C.

The effect of simultaneous administration of bacterial polysaccharide and adrenal cortex extract on cells of mouse tumors and on the adrenal glands of the host

Cancer Res. 7:715, 1947 (Abstract)

The action of the polysaccharide of S. marcescens on sarcoma 37

in mice was not inhibited by simultaneous administration of adrenal cortical extract. The necrotizing effect was delayed with maximal effects appearing in 24 hours. Chemical as well as structural changes are suggested by the appearance of the cytoplasm of cells and their staining characteristics. Cellular changes in adrenals of the host follow administration of either substance alone; damage appears to be prevented by simultaneous injection.

487. EDSALL, G. and WETTERLOW, L. H.

Missed contaminations in biologic products: The role of cryophile bacteria

J. Bact. 54:31-32, 1947

Bacterial species which thrive below the incubation temperature ranges produce pyrogens; refrigerated serums, when once contaminated, are fertile sources of such pyrogenic substances. Gram-negative organisms which grow only at lower temperature ranges have been isolated; they resemble Pseudomonas and grow most actively at 20 to 25 degrees C. but will multiply at 0 degrees. Tests for pyrogens should be made at room temperature as well as at 35 to 37°C.

488. FELTON, L. D., PRESCOTT, B., KAUFMAN, G. and OHINGER, B.

Studies on immunizing substances in pneumococci. XIV. The distribution of specific polysaccharide in mouse tissues after injection of a "paralyzing" dose

Federation Proc. 6 (Pt. 2) 427, 1947 (Abstract)

In the mouse a large dose of the antigenic polysaccharide of pneumococci "paralyzes" the immunologic mechanism for about 15 months; the phenomenon is type-specific. The concentration of the antigen in the tissues was traced and comparisons made of histochemical preparations from normal animals. In order of decreasing precipitin titer the polysaccharide was found in these tissues: liver, spleen, kidney, skin, bone marrow. In some instances, it could be demonstrated in muscle and lung tissues and in urine and the intestines. It was present after 15 months in the liver but had decreased or was absent in other tissues. Mice injected with a paralyzing dose of whole cell vaccine showed similar distributions of polysaccharide.

489. FRANCE, O. and LOSDAN, M.

Pirètoterapia biológica en reumatismo inflamatoriós (Biological pyretotherapy in inflammatory rheumatism)

Ref. argent. reum. 12:72-75, 1947

As adjuvant therapy with antibiotics (sulfonamides or sulfonamide-penicillin injections) typhoid vaccine containing both antigenic and metabolic products of the bacteria was administered. Dosage ranged from 100 to 500 million organisms per ml. on an average, but injection of 2000 million bacterial cells daily proved to be innocuous. In a series of patients with rheumatism of septic origin, 92% received benefit; there was some response in cases of inflammatory rheumatism. Rheumatoid arthritis was not improved.

490. GEORGI, C. E.

Pyrogen testing with "Polish" rabbits

J. Am. Pharm. A. 36:95, 1947

The small size of the "Polish" rabbit makes it an economical and useful animal for the testing of pyrogens.

491. GIFFORD, S. R.

Ocular Therapeutics (A Handbood of)

Philadelphia, Lea and Febiger, Inc. 4th Edition, revised by D. Vail, 1947, P. 119.

A nonspecific action in ocular infections is credited to a "mixed toxin" derived from killed cultures of streptococcus erysipelatis, and B. prodigiosus, grown separately, mixed together and diluted to strengths designated by the percentage of protein contained therein. The adult dose is 3 minims intramuscularly. Within four or five hours slight chills are followed by mild temperature increases (100° - 102°) and a slight inflammation at the site of injection. Results have been rewarding in uveal infections, corneal ulcer and postoperative iridocyclitis.

492. GINSBERG, H. S., GOEBEL, W. F., and HORSFALL, F. L., JR.

Inhibition of mumps virus multiplication by a polysaccharide

Proc. Soc. Exper. Biol. Med. 66:99, 1947

- 1 9 4 7 -

Multiplication of the mumps virus in the chick embryo can be inhibited by the capsular polysaccharide of Friedlander's bacillus, type B. After adsorption of the polysaccharide chick erythrocytes do not adsorb the virus.

493. HARTLEY, F.

The decline of pyrogenic activity in intravenous solutions during storage

Quart. J. Pharm. Pharmacol. 20:444, 1947

Wide variations in stability between different types of solutions must be considered in reckoning the rate of disappearance of pyrogens in stored solutions. Rapid disappearance of pyrogens may be seen in one and long continuation of pyretic activity in another. The presence of pyrogenic activity can be demonstrated after three years' storage.

494. HOLLOMAN, A. L.

Reactions of patients and of tumors to injections of Serratia marcescens polysaccharide in eight cases of malignant disease

Approaches to Tumor Therapy, Amer. Assn. for the Advancement of Science, Washington, D. C. XI. pp. 273-276

Four patients with fibrosarcoma, two with Hodgkin's disease, one with malignant melanoma and some with chondrosarcoma were treated with the polysaccharide of S. marcescens. The initial dose was 0.005 mg.; this was increased later to 0.04 mg. in a single dose. Reactions were severe and resembled those of shock, although anti-shock therapy was ineffective in treatment. Elevation of the lower half of the body, and continuous intravenous drip of a solution of 5% glucose in saline, containing 1 mg. of thiamine and a low concentration of epinephrine were employed. The highest temperature elevation recorded was 108.4°F. A short intense rise in blood pressure was succeeded by hypotension.

495. HORSFALL, F. L., JR.

The modifying effects of certain substances of bacterial origin on the course of infection with pneumonia virus of mice (PVM)

J. Exper. Med. 85:623-646, 1947

Since it has been demonstrated that infection subsequent to inoculation with pneumonia virus could be modified in mice by injections of streptococcus MG, the effect of other bacterial species was determined after intranasal inoculations with PVM. Both living and dead bacteria in various concentrations and

- 1 9 4 7 -

bacterial polysaccharides were employed. In addition, non-bacterial polysaccharides were injected. Varied time intervals between inoculations were observed. Streptococcus MG, S. salivarius, type II and Group A hemolytic streptococcus were injected during the test periods. A second pneumococcus organism, E. coli and H. influenzae, type B., were included in the group. Virus titration end points and hemagglutination titers were determined in the infected lungs as well as the frequency and degree of pulmonary consolidation.

Both bacterial and non-bacterial substances modified the course of PVM infection, but were inactive when administered by other than the nasal route. They appeared to inhibit virus multiplication. Competition may exist between these active substances and PVM for some intracellular system required for multiplication of the virus. Chemical blockade of virus receptors in the cells of the respiratory tract did not appear to be involved in the mechanism of inhibition. The evidence, still unconfirmed, would indicate that the active component may be a purified polysaccharide.

496. JAGER, B. V. and NICKERSON, M.

The altered response of human beings to the intramuscular administration of typhoid vaccine during massive salicylate therapy.

Am. J. Med. 3:408-422, 1947

The effects of injection of typhoid vaccine in a control group of healthy adults and in patients suffering from arteriosclerotic heart disease, acute rheumatic fever and from multiple sclerosis were contrasted with those observed in 14 patients receiving massive salicylate therapy. A number of individuals in both groups had been previously immunized against typhoid. Salicylates as medication had been administered for periods ranging from 7 to 132 days before typhoid injections, and were continued for 16 to 70 days after.

While the control group exhibited typical reactions to typhoid vaccine - development of H and O antibodies, transient leukocytosis, lymphopenia, a rise in the red cell sedimentation rate and plasma fibrinogen levels and increases in gamma globulin - the patients treated with salicylates had only slight changes or none. The systemic and local reactions noted in individuals previously immune to typhoid were absent during massive salicylate therapy. Antibody production may be prevented through the action of salicylates by a reduction in toxicity of antigenic proteins or through selective liver damage which inhibited production of precursors. A direct action on lymphocytic antibody production may be the result of adrenal and pituitary stimulation or there may be rapid destruction of typhoid toxins in the circulation.

497. KUN, E.

Effect of meningococcal endotoxin on histamine content of blood and tissues of rabbits

Proc. Soc. Exp. Biol. Med. 66:197-201, 1947

Intravenous injection of meningococcal endotoxin reduces blood histamine and increases that of liver and muscle. Histamine levels in the blood could not be sustained by simultaneous administration of endotoxin and histamine; the latter procedure appeared to shorten the survival time of test animals.

498. McCONNELL, J. R., HALLETT, S. F. and SHEAR, M. J.

Effect on sarcoma 37 in tissue culture of two tumor-necrotizing agents

Cancer Res. 7:716, 1947 (Abstract)

Hanging drop cultures of sarcoma 37 grown for 18 hours were treated directly with concentrations of emetine hydrochloride ranging from 10 mg. per cc. to 0.00001 mg. The cells rapidly disintegrated, nuclei shriveled and became pyknotic. The rate of destruction was proportional to emetine concentration and toxicity appears to reside in a rapid surface reducing effect. Similar concentrations of S. marcescens polysaccharide produced no necrotizing effect in vitro. The method is informative but does not appear to be as reliable as a screening procedure as in vivo tests.

499. MONDOLFO, H. and HOUNIE, E.

Sobre el origen del pirogeno bacteriano (On the origin of bacterial pyrogens)

Dia Med. 10:1724-1725, 1947

In a review of the factors concerned with pyrogen production, the role of culture medium in increasing pyrogenicity of supernatant fluids is described. Blood serum increases the pyrogenicity of E. coli cultures.

500. MORGAN, H. R. and BENNETT, G. A.

Intra-articular changes induced in rabbits by injection of typhoid somatic antigen

Arch. Path. 44:609-620, 1947

The toxic somatic antigen of S. typhosa was injected directly into the sub-patellar joint of the rabbit in dosages ranging from 0.25 mg. to 0.5 mg. and the injections were continued at intervals of 3 to 7 days for varying periods of time. Concomitant injections intradermally were employed as conditioning procedures preparatory for the Schwartzman reaction and were followed by

- 1 9 4 7 -

provocative injections 18 to 24 hours later. Tissues were examined both macroscopically and microscopically.

Repeated intra-articular injection of the antigen produced acute arthritis which progressed to a chronic state. Synovial tissue hypertrophy, surface "lipping" and subsynovial lymphoid accumulations were seen. The dermal reaction passed through a self-limiting course. Vascular thrombi were extensive throughout heart, lungs and liver.

. NASMYTH, P. A.

The decline of pyrogenic activity in intravenous solutions during storage

Quart. J. Pharm. Pharmacol. 20:445, 1947

In the discussion during the symposium, Nasmyth noted that the work of Collier and Paris, just reported, had shown that a wide range of temperatures might be seen in the normal rabbit. In the seven employed by these investigators, the differences extended to 500% and the lower of the temperatures recorded in these rabbits was well within normal limits for the species.

. OAKEY, R.

Reactions of patients to injection of Serratia marcescens polysaccharide in nine further cases of malignant disease

Approaches to tumor therapy.

American Association for the Advancement of Science, 1947, pp. 277-278

Serratia marcescens polysaccharide may be given to patients repeatedly with the necessary safeguards against extreme toxic and shock states. No tumor was completely destroyed when 0.03 to 2.0 mg. of polysaccharide was given in 1 to 14 injections to members of a group of 9 patients bearing tumors of varied types; changes in the size and consistency of most of them could be seen. The systemic reactions were similar to those noted with other pyrogenic substances; the most alarming feature was the fall in blood pressure with its threat of complete peripheral circulatory collapse.

. OCHOA, A. G. and FIGUEROA, E. S.

Los Polisacáridos de Sporotrichum schenckii (Polysaccharides of Sporotrichum schenckii)

Rev. d. Inst. salub. y enferm. trop. (Mexico) 8:143-153, 1947

The method of Palmer and Gerlough (C. A. 34:7328¹) was applied to 2.25 Gm. of the dried fungus, Sp. schenckii. A small quantity (0.03 Gm.) of a compound extracted by this method gave a positive Molisch reaction and reduced Benedict's solution after hydrolysis. A carbohydrate was obtained by alcohol precipitation of the culture medium (0.10 Gm. in 230 cc. of medium) which was negative for Millon, biuret and xanthoproteic tests. Neither the compound or its carbohydrate component were antigenic or toxic; both were precipitable by homologous and heterologous immune rabbit serum and by serum from a patient with sporotrichosis.

504. PERRAULT, A. and SHEAR, M. J.

Large scale preparation of the tumor-necrotizing polysaccharide from S. marcescens

Cancer Res. 7:714, 1947 (Abstract)

Culture of S. marcescens in synthetic media may be adapted to large scale operations (up to 350 liters). No differences in potency between preparations so secured and those obtained in small scale can be seen. Active fractions from the organisms themselves may be subjected to purification.

505. RADEMAKER, L.

Reactions to intravenous administration to solution

J.A.M.A. 135:1140-1141, 1947

The majority of pyrogenic substances can be eliminated from solutions during manufacture by use of triple distilled water, immediate sterilization of water, solutions and glassware. Rigid cleansing of distilling apparatus must be done periodically. Pyrogenic materials retained in moisture films on glassware, in tubing and needles, can be eliminated by washing and sterilization. Adsorption filtration of solutions is possible and is an additional safeguard.

506. REUBI, F.

Le traitement de la sclérose en plaques (Treatment of multiple sclerosis)

Schweiz. Med. Wchnschr. 77:1177-1179, 1947

- 1 9 4 7 -

For multiple sclerosis, pyretotherapy, vaccines, antimyelotic substances, coagulation inhibitors, vasodilators and radio-active materials were ineffective. Some results were seen in a few patients after antihistamine administration. The brain tissue extract of V. Murali is said to activate nerve regeneration; a tentative trial of this substance is suggested by the author.

7. ROTENHEIM, C. A., SATKE, H. A. and LIMAYE, M. V.

"
Über Pyrogene (On pyrogens)

Pharm. Acta Helvet. 22:673-678, 1947

A review of pyrogens, from the standpoint of the interest of pharmacist and manufacturer, with methods for eliminating such substances from solutions.

- . SACK, T. and SELIGMAN, A. M.

Some effects of iodinated bacterial polysaccharide on patients with malignant tumors

Cancer Res. 7:715, 1947 (Abst.)

When iodinated polysaccharide from S. marcescens was administered intravenously to seven patients with metastatic carcinoma, the effects of massive dosage could be observed, since from 2 to 3000 gammas were administered as individual doses. Post-injection reactions followed a common pattern, namely, chill, a temperature increase sustained for one or more hours and a gradual fall to normal levels within 12 to 30 hours after injection. The blood pressure rose during the chill, fell when temperature reached maximal levels, with a hypotensive period occurring 8 to 15 hours after injection. In four instances antishock therapy was required. Asthmatic symptoms, albuminuria and mild congestive heart failure appeared. However, the reactions did not differ markedly from those which occur after administration of the untreated polysaccharide. Post-injection exhaustion appeared to be prevented or greatly reduced by adrenal cortical extract, given in combination with the iodinated polysaccharide.

34. SMITH, W. E. and PENNELL, R. B.

Reducing the pyrogenicity of concentrated protein solutions

J. Bact. 54:715-718, 1947

A method is described for elimination of pyrogens from concentrated protein solutions, such as serum, plasma and hemoglobin.

This is accomplished through the use of a new type of Setiz pad, special filters and an ion exchange agent "decalso".
 "Pyrogen removal is a function of the degree of pyrogenicity and the concentration of the protein solution and the proper relationship of the amount of 'decalso' and filter pad area to these variables."

510. STUTTGEN, G.

Die Beeinflussung des cholinesterase im Serum des menschlichen Blutes durch pyrifer und U. V. Bestrahlung, besonders im Hinblick auf die Therapie einzelner Dermatosen (Regulation of cholinesterase in human blood serum by means of pyrifer and U. V. light rays, particularly in view of therapy of certain dermatoses)

Klin. Wchnschr. 24:758-759, 1947

It has been postulated that the bacterial product, pyrifer, regulates the equilibrium maintained between the vagotonic and sympathotonic systems. In testing this theory, its effect on cholinesterase was ascertained. It was noted that, with increased cholinesterase activity, ultra violet light rays produced erythemic skin changes.

511. WEGER, P.

Über rein dargestellte hochwirksame Fieberstoffe (Highly potent pure pyrogens)

Naturwissenschaften 34:59-60, 1947

The effects of pyrogens from E. typhus, E. coli, proteus, pyocyanus and abortus equi organisms were observed in the rabbit, mouse, horse and in man. A response could be elicited in the rabbit through the administration of 0.25 micrograms, but higher doses gave a higher temperature level. During this period animals appear quite normal and maintain their weights. Tolerance does not appear on repeated daily injections, while, in human beings, increased doses must be administered in order to achieve the same responses.

Chemical analyses of the pyrogens used are reported.

- 1 9 4 8 -

512. ALBERTY, R. A. and HEIDELBERGER, M.

Fractionation and physical-chemical studies of a commercial preparation of the specific polysaccharide of Type I pneumococcus

J. Am. Chem. Soc. 70:211-213, 1948

A colloidal electrolyte was separated from Type I pneumococcus. Electrochemical and solubility characteristics resembled those of the proteins. The crude preparation showed an isoelectric point at pH 2.8. After fractionation with isopropyl alcohol, electrophoresis gave an isoelectric point for one component, that of nucleic acid, slightly below this figure (2.0).

513. ALEXANDER, L., LOMAN, J., LESSES, M. F. and GREEN, I.

Blood and plasma transfusions in multiple sclerosis

Multiple Sclerosis and the Demyelinating Diseases. Baltimore, The Williams and Wilkins Co., 1950. Chapter 13.

The high incidence of pyrogenic reactions in patients with multiple sclerosis is in contrast with those reported in the literature on transfusion reactions.

Fifty patients in a group of 111 with multiple sclerosis were selected for study. Whole blood and plasma transfusions were given on the theory that, if it be a virus disease, passive immunity might be conferred through active immunity existing in healthy donors. Saline infusions and immune globulin injections were utilized in an effort to obtain information on fractions of blood which might be therapeutically useful. Reactions occurred in 27% of the 298 transfusions given or in 50% of the patients. There were no hemolytic reactions. There were 30 pyrogenic reactions involving 18% of the patients. Allergic symptoms appeared after transfusion in 14%. 10% of these were seen during the first transfusion. Both allergic and neural symptoms were often present in combination. Fever, temporary paralysis or spasticity, diplopia and "blindness" were seen. In the group of 23 patients who were benefited by transfusion, 60% had reactions. Of these, 26% were allergic reactions. Among the patients who did not improve after transfusion, 30% had allergic reactions. Clinically significant improvement was seen in 34% of all patients treated; 6% had incomplete remissions, three complete remission. Improvement was seen in but one patient after injection of immune globulin.

514. BAYER, O. and HORSTER, J. A.

Morphologische Veränderung an Plasmacytomzellen unter Pyriferrwirkung (Morphologic alteration in plasma cells under pyriferr

Klin. Wchnschr. 26:623-625, 1948

A diagnosis of plasmacytoma was made for a patient whose blood exhibited numerous Russell body cells with increased globulin fraction. The total protein in the blood serum was 8.9 Gm. per cent; 5.8 Gm. per cent was globulin. Autoagglutination took place between the cells and serum. Of the leukocytes (6400) 35% were segmented. Forty-three per cent of the lymphocytes were plasma cells. On sternal puncture, marrow specimens revealed deeply basophilic cells with extruded nuclei and many vacuoles.

After injection of 25E pyrifur, the temperature rose to 39.5° C. Vacuoles within cells diminished and staining qualities approached the normal. 50E pyrifur was given 19 days later as the cell structure approached its normal appearance; the temperature elevation was less marked.

515. BECK, L. V., BERKOWITZ, D. and SELTZER, B.

Physiological studies on tumor-inhibiting agents. III. Effect on apparent systolic blood pressure in mice of the Serratia marcescens tumor-necrotizing polysaccharide of Shear.

Cancer Res. 8:162-168, 1948

The polysaccharide produced by S. marcescens was injected intraperitoneally in the mouse. Dosages ranged from 5.0 to 100 gammas per mouse. For mice bearing transplanted tumors (Sarcoma 37) a definite and profound decrease in apparent systolic blood pressure followed. A fall in the rectal temperature accompanied this phenomenon. The extent of tumor area bore direct relationship to the reductions in systolic pressure and temperature. Normal mice showed smaller changes even when larger doses of polysaccharide were administered. The degree of prostration and response to toxicity after injection of polysaccharide was increased in the mouse in direct proportion to the amount of tumor tissue present.

516. BELLACH, H.

Dangers of artificial hyperpyrexia: Report of serious reaction from intravenous injection of typhoid vaccine

Connecticut State M. J. 12:843, 1948

A shocklike and hypotensive reaction was observed after administration of typhoid vaccine. Adequate anti-shock therapy brought a prompt recovery. The dosage does not appear to be a major factor in determining the incidence of untoward reactions. Careful

- 1 9 4 8 -

study of patients before institution of therapy and hospitalization during treatment is advisable.

517. BENACCHIO, L.

Contributo clinico alla terapia vaccinica endovenosa nelle infezione tifo-partifione (Clinical contribution on intravenous vaccine therapy in typhoid-anti-typhoid infection)

Gior. batteriol. e Immunol. 39:337-358, 1948

Forty-eight of 50 patients serologically positive for E. typhosa received anti-typhoid vaccine intravenously, with recovery following therapy considerably accelerated. The roseola herpes disappeared soon after injection, the spleen diminished in size. Leukocytosis and eosinophilia were observed during the treatment.

518. BOQUET, P., DELAUNAY, A., LEHOULT, Y. and LEBRUN, J.

Observation direct des reactions cutanees chez le lapin soumis a l'epreuve d'une endotoxine typhique (Direct observations of vascular reactions of the skin in the rabbit injected with typhoid endotoxin)

Compt. rend. Acad. Sci. 225:1193-1194, 1948

Injection of 6 to 8 mg. of purified typhoid glucido-lipido-polypeptide antigen produces within 5 to 15 minutes constriction and dilatation of the arterioles. This is the first symptom of typhoid intoxication. Sympathetic innervation of blood vessels does not prevent vascular reactions.

519. CREECH, H. J., HAMILTON, M. A. and DILLER, I. C.

Comparative studies of the immunological and tumor-necrotizing properties of polysaccharides from Serratia marcescens (Bacillus prodigiosus)

Cancer Res. 8:318-329, 1948

Comparative studies of the immunological, toxic and tumor-necrotizing properties of purified polysaccharides obtained from culture filtrates of S. marcescens (P3, P5 and P10) and those of fractionation products from one strain of the same organism are presented. Immunization of rabbits followed the injection of 2 mg. of polysaccharide in divided doses over a period of two weeks; the antisera was fractionated and studied serologically. The lethal dosage was determined by the mortality which resulted from intraperitoneal injection of 100 to 2000 gammas of the materials into normal mice. Tumor-necrotizing potency was

evaluated by histologic examination of tumor tissue following administration of 10 or 20 gammas of the preparations into Swiss mice bearing transplanted Sarcoma 37.

Toxicity and antigenicity were more pronounced in P₃ type polysaccharides, while less variation between the three types was evident in their effect on hemorrhage and necrosis production in tumors. Serologically the polysaccharides from two strains of S. marcescens differed. Some fractionation products were but weakly antigenic and relatively nontoxic but retained tumor-necrotizing properties. Storage for a period of years reduced anti-neoplastic qualities but altered toxicity and antigenicity very little. Freezing appeared to enhance the tumor-necrotizing activity.

520. CREECH, H. J., HAMILTON, M. A., NISHIMURA, E. T. and HANKWITZ, R. F., JR.

The influence of antibody containing fractions on the lethal and tumor-necrotizing actions of polysaccharide from Serratia marcescens (Bacillus prodigiosus)

Cancer Res. 8:330-336, 1948

Normal mice and those bearing dorsal transplants of Sarcoma 37 were passively immunized with gamma globulin fractions isolated from rabbit serum after injection of polysaccharides produced by two strains of S. marcescens. Both antisera and gamma globulin fractions protected mice against lethal doses of polysaccharide from the same strain of S. marcescens but not against that of a heterologous strain. One type of antisera and gamma globulin gave no protection against homologous polysaccharide. Lethal activity of one polysaccharide was decreased by use of the specific gamma globulin fraction. The reduction of lethality did not affect tumor hemorrhage production in the mouse.

521. DELAUNAY, A., BOQUET, P., LEBRUN, J., LEHOULT, Y. and DELAUNAY, M.

The mode of action of bacterial endotoxins. IV. The vasomotor disturbances and their consequences in the intoxicated animals.

J. Physiol. 40:89, 1948

The mesenteric vessels of guinea pigs and mice and the cutaneous vessels of the rabbit ear were examined microscopically after a lethal dose of S. typhi antigen. A few minutes after injection of the antigen, irregular contractions of arterioles and medium-sized arteries were seen, with weaker contractions affecting the venous side. Dilatation of the vessels occurred in the terminal stages. Throughout intoxication, diapedesis of leukocytes was inhibited.

- 1 9 4 8 -

522. DILLER, I. C., BLAUCH, B. and BECK, L. V.

Histological changes in adrenal glands of tumor-bearing mice injected with Serratia marcescens polysaccharide alone and in combination with adrenal cortical extract

Cancer Res. 8:591-605, 1948

Young albino mice bearing 7 day-old implants of sarcoma 37 were injected with either 0.02 mg. P-3 polysaccharide intraperitoneally or 0.5 cc. of adrenal cortical extract (Upjohn) by the same route but in divided doses or the two substances in combination. Similar experiments were carried out with mice bearing methylcolanthrene induced tumors and also with litter mates. Both survival studies, tumor growth measurements and histologic examinations of tumor and adrenal tissues followed.

Either the polysaccharide or the adrenal cortical extract caused tumor hemorrhage and necrosis, and similar effects on the adrenal glands were noted. The adrenal medulla showed more extensive necrotic changes than did the cortex. In combination, polysaccharide and adrenal cortical extract delayed tumor necrosis and produced minimal glandular changes. In healthy mice or tumor bearing mice where the growth is still in the primary stages, no tumor necrosis and little adrenal damage followed the administration of these agents.

523. FARR, R. S., LEQUIRE, V. S., SCHORK, P. K. and GAYHART, C. H.

The augmentation of the pyrogenic and leukocytic effects of typhoid vaccine by homologous plasma in the rabbit

Project NM007039. Report No. 17, Naval Medical Res. Inst. 1948

This article is reviewed in detail in: Proc. Soc. Exper. Biol. & Med. 75:661-666, 1950

524. FRONTALI, N. and TOOCHI, G.

Gli inibitori enzimatici nel processo febbrile; (Enzymatic inhibitors and the febrile process)

Boll. Soc. ital. biol. sper. 24:245-247, 1948

"Piretamina" was administered by various routes and the effect of chemical inhibitors assessed.

525. GINSBERG, H. S., GOEBEL, W. F. and HORSFALL, F. L., JR.

- 1948 -

The inhibitory effect of polysaccharide on mumps virus multiplication

J. Exper. Med. 87:385-410, 1948

The viruses of mumps, influenza A and B and Newcastle disease were cultivated in allantoic fluid and inoculated into fertile seven-to-eleven-day-old eggs. The inhibitory properties of the capsular polysaccharides of Friedlander's bacillus, types A, B, and C, and streptococcus MG polysaccharide were tested, as were those of levan and dextran, blood group substance A, apple pectin and commercial corn starch. Hemagglutination was used as a measure of mumps virus concentration. It was found that the polysaccharides from type specific Friedlander bacilli inhibit the multiplication of mumps virus in the allantoic sac of the chick embryo. Concentrations as low as five gammas were effective. This polysaccharide had no effect on other viruses used in the studies, and no other polysaccharide inhibited mumps virus. The mechanism involved may be one of blockade of the metabolism of host cells and not one influencing the virus or its infectivity

526. GINSBERG, H. S., GOEBEL, W. F. and HORSFALL, F. L., JR.

The effect of polysaccharides on the reaction between erythrocytes and viruses, with particular reference to mumps virus

J. Exper. Med. 87:411-424, 1948

In an effort to determine whether or not a single mechanism is involved in the inhibition of hemagglutination and multiplication of viruses by the polysaccharides of Friedlander bacilli A, B, and C, or of other non-bacterial polysaccharides, the viruses of mumps, influenza A, B, and Newcastle disease or PVM were titrated against chicken, human group O and mouse erythrocytes. Polysaccharides active as inhibitors do not block adsorption of the virus by erythrocytes nor by cells of the living allantoic membrane in all instances. There is no correlation between effects of polysaccharides on hemagglutination by viruses and their inhibition of virus multiplication. Wide variation is shown between in vitro and in vivo inhibiting activity of polysaccharides.

527. HALL, V. E., FIELD, J. and GRANT, R.

The metabolic activity of the central nervous system as it affects the thermostatic behavior of the body.

Memorandum Report, MCREXD-686-113D. U.S. Air Force Materiel Command, 1948

- 1 9 4 8 -

528. HALL, V. E., GRANT, R. and FIELD, J.

The influence of substances affecting body temperature on thermal polypnea

Federation Proc. 7:48, 1948

The effect of substances which alter body temperature was determined on oxidative and glycolytic activity of cerebral cortex. In the same studies, the respiratory response to heat (thermal polypnea) was carefully measured. Thermal polypnea appears when the rectal temperature of the rabbit has been increased about 2°C. by administration of typhoid-paratyphoid vaccine. When panting is present before injection, it ceases within fifteen minutes, and inhalation of 95% oxygen (and 5% CO₂) is not depressed by vaccine. Normal respiration is observed in the vagotomized rabbit. Polypnea may be provoked in the rabbit by 3 mM per Kg. magnesium chloride at environmental temperature of 20° C. or higher.

529. HALL, V. E., FIELD, J. and GRANT, R.

The metabolic activity of the central nervous system as it affects the thermostatic behavior of the body

Memorandum Rpt. U. S. Air Force Air Materiel Command MCREXD-696-113D, 1948, 99 pp.

The effects of intravenous injections of typhoid-paratyphoid vaccine in the rabbit and the cat were carefully studied by collaborating investigators, and comparisons were made with those noted after injection of a number of chemical thermogenic and thermostatic substances. The site of action of pyrogenic substances in the central nervous system, reactions after "shock" doses and effects of T.P.T. vaccine on heat production, heat loss and cold defense were ascertained and evaluated.

While results were inconclusive and no proof of a thermal "trigger" reaction in the central nervous system was forthcoming, the report enumerates many interesting facts on calorogenesis after pyrogen administration and presents a series of methods well adapted for use in such studies.

530. HANSEN, J. L.

Hyperthermia in association with shock

Acta chir. Scand. 96:530-546, 1948

Although the body temperature during traumatic shock is usually

- 1948 -

below normal levels, postoperative hyperthermia has been reported. Rectal temperatures may reach maximal limits within 24 hours after operation while shock is pronounced and no infection appears to be present; decreases are rapid following treatment for shock. The phenomenon has been noted after gastric operations, particularly when cancer or ulcer have been found. The clinical picture is one of coldness of the extremities, marked blood pressure reductions, oliguria; the pathology represented included oedema of cerebrum, vasodilatation, stasis and intravascular space alterations. Serous exudate of the nervous tissue was noted in the brain stem. It is suggested that hyperadrenalemia would produce similar symptoms (arteriolar contractions, paleness, piloerection and perspiration). It has been shown that adrenalin itself may be pyrogenic. Impedance of blood flow at the portion near the vena cava may be a factor.

531. HATTA, S., AOYAMA, K., and TANJI, S.

Studies on the bacteria (and harmful substances)

Japanese J. Pub. Health, 5:24, 1948

532. HICKS, M., HOLT, H., GUERRAND, S. and LEAVELL, B. S.

The effect of spontaneous and artificially induced fever on liver function

J. Clin. Invest. 27:580-587, 1948

Diminished bromsulfalein clearance is seen in induced fever and may be the result of impaired function in hepatic tissues. Abnormal retention of bromsulfalein is seen during febrile periods in infections, and body temperature at the time of testing must be considered in evaluation.

533. KIRCHMAIER, H. and SCHOLTZ, W.

Zur Pyriferbehandlung des Typhus und Paratyphus im Kindesalter
(Pyrifer treatment of typhus and paratyphus in childhood)

Artzl. Wehnschr. 3:426-428, 1948

Elevated body temperatures fell in crisis, and diarrhea ceased after one to four injections of pyrifer were administered to children ill with typhoid or paratyphoid fever. For the entire group, the period of illness was not appreciably shortened over that occurring under conservative treatment.

- 1 9 4 8 -

534. KUN, E.

Effect of bacterial endotoxins on glycogen synthesis

Proc. Soc. Exper. Biol. & Med. 68:496-498, 1948

The influence of endotoxins from Salmonella and meningococcal organisms upon the synthesis, in vitro and in vivo, of glycogen was studied. In the rat diaphragm and rabbit liver slices there was inhibition of glycogenesis from pyruvate and glucose; insulin diminished this inhibition in vitro in the rat diaphragm. In vivo toxins completely prevented glycogenesis in the liver of the rat.

535. KUN, E.

Inhibition of the phosphorylation of glucose by meningococcal endotoxin

J. Biol. Chem. 174:761-762, 1948

Since the addition of Zn-free insulin to meningococcal endotoxin prevents the inhibition of glucose utilization in rat muscle extracts in vitro, it is believed that meningococcal endotoxin prevents phosphorylation of glucose through a reduction of hexokinase activity. "...The disturbances of the carbohydrate metabolism in meningococcal toxic rabbits and meningococcal meningitis in humans...may be explained by a direct effect of meningococcal endotoxin on enzymatic reactions in the host..."

536. KUN, E., and MILLER, C. P.

Effect of bacterial endotoxins on carbohydrate metabolism of rabbits

Proc. Soc. Exper. Biol. & Med. 67:221, 1948

The injection of lethal doses of sterile, unpurified meningococcal and Salmonella endotoxins in the rabbit, increased blood glucose, lactic acid and inorganic phosphorus. Liver glycogen decreased while lactic acid content of tissues increased; pyruvic acid in blood and tissues was much reduced. Succinic-dehydrogenase activity in both muscle and liver was inhibited; cytochrome oxidase activity was not affected. A state of tissue anoxia may exist with corresponding interference with carbohydrate metabolism.

537. LEWIS, L. A. and PAGE, I. H.

Changes in blood leukocyte level of adrenalectomized and normal rats following administration of typhoid vaccine

Am. J. Physiol. 153:148-152, 1948

Adrenalectomized rats were maintained on either adrenal extract, dehydrocorticosterone acetate or physiologic saline solution for three days, at which time 1.33 MLD of typhoid vaccine was administered intraperitoneally. Determination of blood leukocyte and differential levels and of hemoglobin followed. Results were compared with values found in normal animals and humans after immunizing doses of typhoid vaccine.

Blood leukocyte values changed little after injection of adrenal extract, compound A acetate or saline, but marked lymphopenia and neutrophilia were noted after typhoid vaccine injections. These were enhanced by previous treatment with the steroids. Within 24 hours, all animals receiving saline alone succumbed; pre-treatment with hormones prevented lethal effects of typhoid vaccine. The lymphopenia does not appear to be altered by adrenal cortical hormones, but the increase in neutrophils is markedly augmented.

538. MILLER, C. P., HAWK, W. D. and BOOR, A. K.

Protection against bacterial endotoxins by penicillin and its impurities

Science 107:118-119, 1948

The protective effects of penicillin against a series of endotoxins injected into the mouse was confirmed through observations of animals receiving both purified and impure penicillin before intraperitoneal administration of endotoxin. Preparations were derived from gram-negative organisms. Such protection was demonstrated against the endotoxins of S. paratyphi A., S. paratyphi B., Shigella paradysenteriae, S. enteritidis, S. aertrycke and Aerobacter. The LD₅₀ for mice pre-treated with penicillin was three times greater when the crystalline product was employed and nine times greater when mice were given the impure product. The impure penicillin preparation lost its protective activity when treated with penicillinase or heat sufficient to inactivate penicillin; addition of the original amount of crystalline penicillin restored its capacity for protection.

539. MORGAN, H. R.

Resistance to the endotoxins of enteric bacilli in man

- 1 9 4 8 -

J. Clin. Invest. 27:706-709, 1948

Seven patients, two with gonorrhea, two with asymptomatic serologic syphilis and three without infections, received either the somatic antigen of S. typhosa, that of S. schottmulleri or Sh. dysenteriae. Tolerance to the toxic effects of the antigens developed after repeated injections. Subsequent injections did not produce hyperpyrexia and other symptoms, and cross-tolerance developed between the heterologous antigens. However, specific "O" agglutinins were demonstrable only for immunologically related antigens.

540. MORGAN, H. R.

Tolerance to the toxic action of somatic antigens of enteric bacteria

J. Immunol. 59:129-134, 1948

Rabbits developed tolerance to the pyrogenic effect of the somatic antigens of S. typhosa, S. schottmulleri and Sh. dysenteriae within three to four days following initial intravenous injections. Such tolerance was not specific for antigenic structure and appears to bear little relationship to the presence of specific circulating antibody. After a rest period, the resistant state disappears at a time when the level of circulating antibody still remains specifically elevated. Animals develop specific antibodies without comparable decrease in febrile response to pyrogenic antigen, and the preparation from Sh. dysenteriae which is but feebly antigenic induces an active state of tolerance.

541. PINGERT, F. P. and FERRY, C. W.

U. S. Patent 2,432,970, 1948

A method for reducing the pyrogen contents of parenteral solutions sensitive to temperature changes and to usual acid and alkaline treatments is described.

542. RABINER, A. M., ROSENBERG, M. and FRIEDMAN, H.

Neuronitis and neuronopathy

Ann. Int. Med. 29:432-444, 1948

For 6 of a group of 7 patients presenting the classic symptoms of acute infectious neuronitis or neuronopathy, benefit was

seen after typhoid vaccine therapy. Relief of pain was the most dramatic feature of this treatment. Acute inflammation of the peripheral nerves, nerve roots, brain stem and cord subsided more slowly, and the recovery of motor power lagged behind other improvements. In two patients there was an immediate return of sensation after the first injection of vaccine; in others, a considerable delay was observed. In the areas involved, regional hyperesthesia and hyperalgesia were noted. The excellent responses to vaccine therapy which occurred without noteworthy temperature elevations would appear to challenge the current belief that the beneficial effects of vaccino-therapy depend on the creation of lethal environmental conditions for infecting organisms.

543. RAUSS, K.

A dysenteria vedoltas kerdese (The problem of immunization against dysentery)

Nepegeszsegugy, 29:449-454, 1948

Trichloroacetic acid extracts of S. dysenteriae protect mice against infection while type-specific activity is found in vaccines prepared from the same organism. It is believed that the toxic substance and the protective factor are identical.

544. SACK, T. and SELIGMAN, A. M.

Chemical alteration of polysaccharide from Serratia marcescens. II. Effects of iodopolysaccharide in patients with malignant tumors

J. Nat. Cancer Inst. 9:19-34, 1948

Intravenous injections of iodopolysaccharide, in dosages ranging from 2 to 3,000 gammas, were administered to seven patients with inoperable or recurrent Roentgen-resistant malignant tumors. The polysaccharide was given over varying periods ranging from 1 to 64 days, or 1 to 9 injections. The toxic reactions which followed resembled those observed after the administration of unmodified polysaccharide and consisted of a prodromal period, a chill 15 to 45 minutes after injection, a period of flush and maximal temperature and progressive hypotension and defervescence. Tolerance to iodopolysaccharide developed rapidly. Cross resistance to the effects of unmodified polysaccharide appeared. No ~~clinical~~ evidence of tumor destruction could be found.

- 1 9 4 8 -

545. SELIGMAN, A. M., SHEAR, M. J., LEITER, J. and SWEET, B.

Chemical alteration of polysaccharide from Serratia marcescens.
I. Tumor-necrotizing polysaccharide tagged with radioactive iodine

J. Nat. Cancer Inst. 9:13-18, 1948

Radioactive iodine was incorporated in the molecule of the Shear polysaccharide and its effects on normal and tumor-bearing animals and humans were observed. Saturation of the molecule occurred over a broad range of I concentrations; a stable iodine linkage could be demonstrated.

When injected intravenously in dosages ranging from 12.5 to 300 gammas, radioactivity disappeared rapidly from the blood stream, but there was a prolonged maintenance of a residum (about 20%). Greatest radioactivity was present in the liver, lung and kidney, decreasing in this order. Tumor-necrotizing properties were not affected, and little radioactivity existed in tumor tissue.

546. TAL, C. and OLITZKI, L.

The toxic and antigenic properties of fractions prepared from the complete antigen of Shigella dysenteriae

J. Immunol. 58:337-348, 1948

A series of fractions were prepared from the complete antigen of the S strain of Shigella dysenteriae. A phospholipid, a conjugated protein and a simple protein, undegraded and degraded polysaccharides were found on fractionation. The conjugated complexes contained varying amounts of phosphorus while the degraded substances were free of this element.

These fractions were tested for their ability to produce the following alterations in animals: lethal effects (mice); hypothermia (in mice, rats and guinea pigs); leukopenia (guinea pig and rabbit); reduction of liver glycogen and adrenal ascorbic acid (in rats); the Schwartzman phenomenon (rabbits) and hemorrhage production in adrenals (guinea-pig) or placentas of pregnant mice. Antigenicity of the fractions for the rabbit was determined.

The simple protein, phosphorus-free, did not elicit the endotoxic reactions produced by the conjugated complexes. The conjugates reduced glycogen and ascorbic acid, and produced leukopenia and hypothermia. Production of a type specific antigen followed administration of the protein. The undegraded polysaccharide produced the Schwartzman reaction, leukopenia and hypothermia and,

combined with small quantities of S. dysenteriae or E. typhosa (O form), was capable of definite pathogenizing action. Anti-genicity depended on the linkage of carbohydrate and protein.

547. TARDIEU, G., GUILLAUMONT, B. and SUTTER, J.

Encephalite choreique ou chorée prolongée. Evolution tardive vers le parkinsonisme. Success de la pyretotherapie par le TAB puis de la radiothrapie des noyaux gris centraux. (Choreic or prolonged encephalitis. Late evolution toward parkinsonism. Success of pyretotherapy by TAB and then radiotherapy of the central gray matter)

Bull. Soc. med. Hop. 64:334-335, 1948

Bilateral chorea stemming from rheumatism responded to TAB therapy within a short period, but relapse followed one year later. TAB therapy was unavailing at this time, but some remission followed roentgenotherapy of basal ganglia.

548. TAUB, A. and HART, F.

Detoxification of pyrogens by hydrogen peroxide in some U.S.P. injections

J. Am. Pharm. A. 37:246-250, 1948

Effective destruction of pyrogens in water, saline solution, or dextrose-saline solution intended for injection, may be carried out by the use of hydrogen peroxide and prolonged boiling (1 hr.) for the first two fluids and by treatment with 100 to 200 mg. acid-washed activated charcoal (per 100 ml.) and filtration through a Selas candle, for dextrose saline solution.

549. TAYLOR, R. D., CORCORAN, A. C., FERTIG, H. H. and PAGE, I. H.

The effects of bacterial pyrogens on malignant hypertensive persons

J. Lab. Clin. Med. 33:1469-1470, 1948

A group of 35 patients, each with clearly-defined symptoms of malignant hypertension, received repeated injections of bacterial pyrogens which produced both fever and leukocytosis. The results were evaluated from the standpoints of changes in arterial pressure, cardiac, cerebral and renal functions and fundoscopic signs.

Diastolic pressure was reduced in 16 of 35 patients. Fourteen

- 1 9 4 8 -

patients became free of retinal hemorrhages and, of 16 patients with congestive heart failure, five were relieved during therapy; electrocardiograms became normal in seven, and heart size decreased, for the entire group, from plus 18% to plus 5% during treatment. Tolerance to pyrogens appeared in five to nineteen weeks when arterial pressures rose. However, the malignant syndrome did not, in the majority of patients, reappear.

Remissions for periods of 24 to 30 months, on an average, were obtained by consistent treatment. Bacterial pyrogens are effective in the absence of severe renal damage.

550. WAKIM, K. G., KRUSEN, F. H. and ELKINS, E. C.

Effect of artificially induced fever on circulation in arthritic patients

Arch. Phys. Med. 29:274-284, 1948

In a discussion of fever production through the use of passive (physical) agents and active substances (colloids, bacterial proteins, living parasites), it is noted that prodromal and chill phases are absent in the former. Both increase leukocyte formation and accelerate cardiac output, blood flow and the size of capillaries. There is an increase in metabolic catabolism.

551. WELLS J. A. and RALL, D. P.

Mechanism of pyrogen-induced fever

Proc. Soc. Exper. Biol. & Med. 68:421-424, 1948

Pyrogen (PYROMEN) will induce fever under conditions where increased heat production from muscular activity is impossible. Under such conditions, the mechanism of the pyrogenic reaction is one of rapid reduction of heat losses.

552. ANDERSON, W. H. and BRODERSON, R.

Hypothermia in the mouse as a bioassay of endotoxin protection factor in impure penicillin

Proc. Soc. Exper. Biol. & Med. 70:322-325, 1949

Young adult mice conditioned in a constant temperature for two days before and during experiments received either (1) 0.5 ml. impure penicillin containing 10,000 micrograms crystalline penicillin per ml. intraperitoneally 20, 18 and 2 hours before injection of 0.05 ml. of endotoxin (2) impure penicillin alone or (3) no treatment. In further experiments hydrogen peroxide oxidation was used to destroy protective activity, and comparisons were made with controls protected by unoxidized materials. A constant dose of endotoxin was administered to a third group; it contained various dilutions of the protective factor. S. aertrycke endotoxin had the ability to reduce temperature markedly, while impure penicillin previously injected protected against lethal effects and hypothermia.

553. AUBERT, J. P.

Existence of a polysaccharide in Bacillus megatherium

Compt. rend. Soc. biol. 229:477-478, 1949

Methods for preparing polysaccharide concentrates from B. megatherium are described. Sporulation decreases the quantity obtainable. The age, the culture, and carbon sources in the media wield influence over the amounts produced.

- 1949 -

554. BAKER, E. E. and GOEBEL, W. F.

Studies on the enzymatic hydrolysis of the capsular polysaccharide of Klebsiella pneumoniae, Type B. *Federation Proc.* 8:400, 1949

An enzyme capable of degrading capsular polysaccharide was isolated from soil bacilli. A motile, gram-negative, spore-forming organism elaborates such an enzyme when it is grown in the presence of 0.1% capsular polysaccharide on a medium containing ample sources of N but no growth factor. The enzyme specifically hydrolyzes capsular polysaccharide as shown by a change in the viscosity of the carbohydrate. Loss of serological activity accompanies this drop in viscosity. More than one enzyme may be involved in the phenomenon.

555. BENVENUTI, M.

Su di un tentativo di cura associati vaccino-seroterapici della poliomielite anteriore acuta. (A trial of the treatment of acute anterior poliomyelitis with vaccine and serum therapy)

Riv. clin. Pediat. 47:137-141, 1949

A trial was made of the use of anti-rabies vaccine with maternal blood serum for five patients with severe or complete paralysis of the lower extremities. The average dose was 3 ml. of vaccine per day and total dosage was approximately 25 ml.

Arrest of the paralysis was secured in all five cases. Regression of the paralysis was succeeded by return of motility. The sequelae were insignificant.

556. BENNETT, I. L., WAGNER, R. R. and LEQUIRE, V. S.

The production of fever by influenzal viruses. II. Tolerance in rabbits to the pyrogenic effect of influenzal viruses

J. Exper. Med. 90:335-347, 1949

Prior injection of certain strains of viruses (PR8 strain of influenza A, Lee strain of influenza B and "B" strain of NDV) conferred tolerance, in the rabbit, to the pyrogenic effects of homologous virus as well as some degree of protection against the lymphopenic action of such virus. The period of tolerance extended, on the average, over 11 days. The protective effect of heated virus was lost when hemagglutinins were destroyed. It appears to be independent of specific antibody formation. No cross tolerance was demonstrable between bacterial pyrogens and viruses. Prevention of fever by antipyretic substances did not interfere with protective influence of previous conditioning. No modification of the reactions could be secured with reticuloendothelial blockade.

557. BENNETT, I. L., WAGNER, R., and LEQUIRE, V. S.

Pyrogenicity of influenza virus in rabbits

Proc. Soc. Exper. Biol. & Med. 71:132-133, 1949

One ml. of chorio-allantoic fluid from infected embryos produced a rise in temperature in the rabbit; doses as small as 0.25 ml. also produced fever of a low degree. The rise in temperature began about 1½ to 2 hours after injection of fluid secured from PR-8 infected embryos, reached a peak of 3 to 4 degrees F. above baselines in four hours, then fell to normal levels. When normal chorio-allantoic fluid or fluids freed from virus particles through centrifugation were injected, no fever resulted; the virus re-suspended in physiologic saline solution again produced temperature elevations. Homologous immune serum neutralized the virus responses. Exposure to heat destroyed hemagglutinin titers and rendered preparations non-pyrogenic. Unlike heat-stable pyrogens the fever element of both PR8 influenza virus and the Lee strain B were destroyed by temperatures above 62° C. The Lee strain required temperatures considerably higher for complete destruction of the pyrexial material. Tolerance to the substance was quickly attained but lost again in eleven days. Antibody titers increased in the period.

558. BHARUCHA, K.

Pyrogen and its prevention

Indian J. Med. Sci, 3:349-350, 1949

Techniques for production of nonpyrogenic solutions are outlined. Freshly distilled water should be used in preparing solutions. Adsorption of pyrogens by activated charcoal or asbestos is possible. Inactivation with 0.01 N/HCl with subsequent neutralization of the acid eliminates pyrogen because of its carbohydrate nature.

559. BOROFF, D. A.

Study on toxins and antigens of Shigella dysenteriae. I. Toxicity and antigenicity of whole organisms and various fractions of Shigella dysenteriae.

J. Bact. 57:617-632, 1949

Somatic fractions of S. dysenteriae or its "smooth" variant produce identical antibodies and cross reactions with antiserum while the "rough" variant fractionated products do not induce immunologic reactions. The toxic properties, present in all forms and fractions, are attributable to a single endotoxin, which may consist of a toxic protein, nonantigenic, a specific polysaccharide, inducing antibody formation, and a lipoid.

560. BOROFF, D. A. and MACRI, B. P.

Study on toxins and antigens of S. dysenteriae. II. Active protection of rabbits with whole organisms and fractions of Shigella dysenteriae

J. Bact. 58:387-394, 1949

Reciprocal protection against homologous strains of S. dysenteriae is afforded by either "smooth" or "rough" whole organisms or their respective toxins. Chemically purified toxins give similar protection. Nontoxic variants give as effective protection as toxic parent strains. The presence of antitoxin or the lethality of its toxin are not determining factors in inducing immunity.

561. BRUETSCH, W. L.

Why malaria cures general paralysis

J. Indiana M. A. 42:211-216, 1949

The author reveals the literature on malarial fever therapy and concludes that in the therapy of luetic general paralysis, fever per se is a minor factor. Beneficial results depend upon the kind of fever produced and upon activation of the reticuloendothelial system to stimulate production of macrophages. Both malarial and typhoid fevers are most active in producing intense macrophagic responses. Tissue immunity produced by reticuloendo-

thelial stimulation inhibits spirochetal growth, disintegration and disappearance. Regression and resolution of the inflammatory processes in the brain follow. "The killing mechanism of the spirochetes in general paralysis...must be looked for in antibody production by the R-E cells." Any phagocytosis of spirochetes by malaria organisms or ingestion of pallida by capillary endothelial cells in the brain must be considered unproven.

562. BUNTING, M. I., ROBONOW, C. F. and BUNTING, H.

Factor affecting elaboration of pigment and polysaccharide by Serratia marcescens.

J. Bact. 58:114-115, 1949

Capsule formation in A. aerogenes is encouraged when growth of the organism is restricted by excessive carbohydrate in the medium, but deficiencies of N and P. Cells assume a nuclear appearance as ribonucleotide leaves the cytoplasm. The accumulation of prodigiosin, a pyrrole derivative, parallels the elaboration of polysaccharide in the cell.

563. CHAMBERS, WW., KOENIG, H., KOENIG, R. and WINDLE, W. F.

Site of action in the central nervous system of a bacterial pyrogen

Am. J. Physiol. 159:209-216, 1949

Febrile response to a bacterial pyrogen from a Pseudomonas species (PYROMEN) appears to be mediated through the central nervous system, and the cord must be in functional communication with one or more brain centers. Such responses are prevented by transection of the spinal cord in the lower cervical or upper thoracic areas; side reactions were not inhibited. When transection was made below T-5, the development of fever was not prevented. Decortication, thalamic and caudal hypothalamic lesions do not prevent febrile reactions. Decerebration at a pontile or medullary level did not prevent fever; an integrating mechanism appears to exist in the medulla oblongata or the upper spinal cord; connections with the spinal cord below the cervical region renders the organism fully responsive to pyrexial agents.

- 1949 -

564. CHASIS, H., GOLDRING, W. and BALDWIN, D. S.

Effects of febrile plasma, typhoid vaccine and nitrogen mustard on renal manifestations of human glomerulonephritis

Proc. Soc. Exper. Biol. & Med. 71:565-567, 1949

No decreases in proteinuria were noted in patients with diffuse glomerulonephritis after repeated intravenous administration of plasma taken from febrile patients during pneumococcal and streptococcal infections. Febrile plasma did not induce diuresis. In one patient with marked renal function impairment, administration of typhoid vaccine reduced proteinuria and the filtration rate, but not inhibition of urine flow. HN_2 (0.2 mg. per kg. body weight) reversed the renal manifestations of glomerulonephritis, reducing proteinuria and, in one instance, increasing urine output.

565. CREECH, H. J., HANKWITZ, R. F., JR. and WHARTON, D. R. A.

Further studies of the immunological properties of polysaccharides from Serratia marcescens (Bacillus prodigiosus). I. The effects of passive and active immunization on the lethal activity of the polysaccharides

Cancer Res. 9:150-157, 1949

In mice bearing dorsal implants of Sarcoma 37, passive immunization with gamma-globulin fractions from serum rendered immune against strains of S. marcescens protected the animal to some extent against the lethal dose of the polysaccharide of the same organism. Active immunization in tumor-bearing mice was accomplished by the injection of small intraperitoneal doses of the polysaccharide followed by the dose ordinarily lethal to 90% of test animals. A prolonged tolerance toward lethal activity of the polysaccharide could be established in both normal and tumor bearing mice, by a single injection of 1 per cent of the 90% lethal dose of polysaccharide from any of the strains studied. This appears to be a true immunological response.

566. CREECH, H. J. and HANKWITZ, R. F., JR.

Further studies of the immunologic properties of polysaccharides from Serratia marcescens (Bacillus prodigiosus). III. Passive immunization against the lethal activity of the polysaccharides with fractions of mouse antiserum elicited by a single injection of polysaccharide

Cancer Res. 9:589-591, 1949

Protective antibodies multiplied rapidly in the serum of mice

after injection of globulin fractions of mouse antiserum; the latter was elicited by single injections of S. marcescens. At a low globulin ratio (16-25) one fraction from P-10 preparation afforded a high degree of protection against an LD₇₀ dose of S. marcescens polysaccharide; a second preparation was less effective.

567. CURRY, J. J. and SHAW, E. A.

Continuous intravenous injection of typhoid vaccine in treatment of certain ophthalmic diseases

Arch. Ophthalmol. 42:123-125, 1949

Controlled hyperpyrexia through slow continuous intravenous injection of a suspension of killed typhoid organisms (1 billion per cc. and 1 cc. per liter of physiologic saline solution) produced excellent results for 17 patients with ophthalmic conditions ranging from nonspecific iritis to syphilitic keratitis. The risk of reaction is believed to be reduced considerably by this method of administration.

568. DAGLEY, S. and DAWES, E. A.

Factors influencing the polysaccharide content of Escherichia coli

Biochem. J. 45:331-336, 1949

Glucose, lactose and galactose increase the growth of E. coli and increase likewise stored polysaccharide up to a maximum. Fructose and mannose do not produce such increases in the polysaccharide. At a fixed sugar concentration, pH affected such storage with glucose and galactose media but not with mannose or fructose. Strain differences as to polysaccharide storage were also noted. Polysaccharide content falls during the stationary phase of growth. The Sahayun method, when coupled with the Hagedorn-Jensen test for reducing sugars, was not satisfactory for the estimation of total bacterial polysaccharide.

569. DELAUNAY, A., DELAUNAY, M. and LEBRUN, J.

The mechanism of lymphocytic lesions provoked by epinephrine and Salmonella typhi

Compt. rend. Acad. Sci. (Paris) 288:139, 1949

Vascular disturbances, destructive to lymphocytes, and not directly toxic effects upon nuclear or cytoplasmic components characterize the effects of S. typhi endotoxin or epinephrine. Hypodermic injection of lymphocytes with these substances into the guinea pig provokes the same 'lesions' as shown by lympho-

- 1 9 4 9 -

cytes in serum after injection of S. typhi endotoxin or epinephrine only.

570. DOMBA, E. and GABOR, T.

Extraction of Hormones, Viruses, Toxins and Antitoxins (Trans. title)

Hung. Patent #139,965, Sept. 24, 1949

Pyrogens may be removed from bacterial and biological preparations by cyclic or heterocyclic acids. Removal with ETOH and 10% BzOH is followed by cooling and filtration when room temperatures are again attained.

571. DORCHE, J. and BOUTHIER, G.

Pyrogenes des solutions injectables (Pyrogens in injectable solutions)

Ann. pharm. franc. 7:267-276, 1949

The application and techniques of the pyrogen test of the U. S. Pharmacopeia are discussed and modifications suitable for small laboratories are suggested.

572. ELLIS, F. A. GRANT, R. and HALL, V. E.

Typhoid-paratyphoid vaccine and respiratory response to the inhalation of carbon dioxide

Am. J. Physiol. 158:16-20, 1949

The hypothesis is presented that pyrogenic substances may depress respiration, and other neuronally mediated processes connected with periodicity of breathing, at medullary or higher levels. The inhalation of CO₂ increased breathing, and no significantly important alterations were produced by subsequent injection of typhoid vaccine. Respiratory minute volume was increased in anaesthetized animals.

573. FELTON, L. D.

The significance of antigen in animal tissues

J. Immunol. 61:107-117, 1949

In mice injection of 0.5 mg. of specific polysaccharide antigen may produce immunologic paralysis extending through the normal

life of the animal; the carbohydrate may be isolated from body tissues for at least one year after immunization, and hypersensitivity is exhibited toward the type-specific organism. No antibody is produced either toward bacteria or immunizing doses of the polysaccharide. Human beings may become non-reactive toward immunizing doses of polysaccharides from an initial paralyzing reaction. It is postulated that a given antigen may become firmly fixed to cell surfaces of the reticuloendothelial tissues and lose capacity to act as an antigen.

574. FRANCKENSTEIN, R.

Über die typhösen Erkrankungen in Kindersalter und ihre Behandlung mit Pyrifer (Enteric fevers in older children: Treatment with pyrifer)

Medizin. klin. 44:199-203, 1949

Nerve regulatory centers affected by cyclic infection by E. typhi in children produced symptoms which were benefited by the injection of pyrifer, 20 to 50 units, intravenously. Temperatures rose in one-half to one hour thereafter, with mild rigors. Other symptoms of a psychotic nature associated with the usual clinical picture were seen in 80 children infected by E. typhosa or paratyphi B. Younger children gave most rapid response to pyrifer and very slight reactions were followed by excellent results, with very little relation between fever production, onset of chills and the therapeutic result being evident. Pyrifer reduced temperature elevation in a child, 18 months old, in whom encephalitis, stupor and a roseola appeared after paratyphus infection.

575. GERMAN, A.

Pyrogenes (Pyrogens)

Ann. pharm. franc. 6:464-479, 1949

After reviewing the literature of pyrogenic phenomena and making certain comparisons between the action of known pyrogens and that incidental to vaccinothrapy and "protein shock", the author concludes that pyrogenic reactions may be responsible for benefits derived from vaccine injections. It is difficult to explain why antibodies appear after injection of killed bacteria in infections in which little antibody formation is developed by the pathogens present. Shock may be responsible for some beneficial reactions to so-called "protein" substances, but it is noted that superior results followed the use of three strong producers of pyrogen, ie. streptococci, staphylococci and Ps. aeruginosa, in a bacterial preparation (prodidon). The initial leukopenia following the injection of pyrogens and the ensuing leukocytosis

- 1 9 4 9 -

may be brought about by a dual activity in pyrogen: (1) massive liberation of antibody by lymphopenia, and (2) defense reactions by the organism with the release of leukocytes in an effort to combat stress.

576. GINSBERG, H. S. and HORSFALL, F. L., JR.

A resistant variant of mumps virus. Multiplication of the variant in the presence of inhibitory quantities of Friedlander bacillus polysaccharide

J. Exper. Med. 90:393-407, 1949

A large single inoculum of mumps virus or serial passage of the virus in the presence of inhibitory quantities of capsular polysaccharide of Friedlander bacillus, type B may produce variant virus strains, with infective and agglutinative qualities identical with the parent strain, but slower to multiply and not inhibited by polysaccharide. In the absence of the polysaccharide, serial passage again produces the original sensitive strain.

577. GRANT, R.

Nature of pyrogen fever: Effect of environmental temperature on response to typhoid-paratyphoid vaccine

Am. J. Physiol. 159:511-523, 1949

Oxygen consumption was used as a measure of calorigenesis when intravenous injections of typhoid-paratyphoid vaccine or "Pyrexin" were given to rabbits kept under constant environmental conditions. The amount given was about 100 times the minimal pyrogenic dose. In moderate and warm environments, moderate doses of TPT brought initial temperature increases, a temporary fall in about 1 hour and renewed inhibition of heat loss mechanisms toward the end of the 2nd hour. With very large doses the early course of fever is the same, but pyrexia may be prolonged. With very small doses, initial temperature rises are identical with those produced by larger amounts, but the final rise in temperature begins early. Oxygen consumption during the first hour is not affected by temperature changes; after the first hour, hypothermia follows in the shorn cold animal. The scant evidence available suggests that pyrogens may interfere with thermoregulation at lower levels of the brain stem.

578. GRANT, R. and ROBBINS, M. E.

Effect of ethyl carbamate on temperature regulation

Federation Proc. 8:59, 1949

Prior or subsequent intravenous injections of urethane did not alter the inhibition of heat loss mechanisms by bacterial pyrogens; intraperitoneal injections of urethane may allow vaso dilation, with polypnea unaffected.

579. GRANT, R., ROBBINS, M. and HALL, V. E.

Effect of environmental temperature on the febrile reaction to typhoid paratyphoid vaccines and other pyrogens

Am. J. Med. 6:395, 1949 (Abstract)

After the intravenous injection of pyrogen, the oxygen production in the rabbit increases to a maximum during the first 20 minutes, and is approximately 15 per cent higher than that of controls for the first hour. It remains slightly above normal during the second and subsequent hours of fever if environmental temperatures remain moderate or warm; when exposed to severe cold, oxygen values show sharp reductions below control values during the second hour and return to normal levels during the third hour. In normal environmental temperatures the second hour after a pyrogen injection brings an inhibition of heat defense mechanisms, and fever increases are again recorded.

580. GROSSMAN, M. I. and BLICKENSTAFF, D.

Quantitative aspects of inhibition of gastric secretion by pyrogens

Am. J. Physiol. 159:572, 1949 (Abstract)

A lag period between the appearance of pyrexia after intravenous injection of pyrogen and inhibition of gastric secretion has been reported. Another gap exists later in the test between temperature relations and gastric secretion inhibition.

581. HALL, V. E., FISHGOLD, J. and GRANT, R.

Oxygen consumption of liver slices from febrile and normal rabbits

Federations Proc. 8:66, 1949 (Abstract)

If, as suggested, heat production of the liver is increased during fever produced by bacterial pyrogens, measurement of O_2 consumption in vitro of liver slices taken from rabbits 15 minutes after injection of typhoid-paratyphoid vaccine does not confirm such observations.

- 1 9 4 9 -

582. HATTA, S., AOYAMA, K., and TANJI, S.

The cause of fever by injection and its real condition (translated title)

Sogo Igaku 113:887, 1949

An article in the original Japanese text.

583. HATTA, S., AOYAMA, K. and TANJI, S.

Elimination of pyrogen substances (translated title)

Saishin Igaku 41:612, 1949

584. HATTA, S., AOYAMA, K., and TANJI, S.

Studies on the bacteria and its harmful substance of injection and internal medicine in general use II. (translated title)

Jap. J. Pub. Health 5:20-23, 1949

585. HEYMAN, A., and BEESON, P. B.

Influence of various disease states upon the febrile response to intravenous injection of typhoid bacterial pyrogen

J. Lab. & Clin. Med. 34:1400-1403, 1949

Malarial fever appears to render its victims resistant to later injections of typhoid bacterial pyrogen; normal febrile response was lacking. With cirrhosis of the liver, the response was intensified. In many infectious diseases, little or no deviation from the normal fever curve is seen.

586. HOLMAN, J. and SWINEFORD, O., JR.

Studies in bacterial allergy. II. Preparation of crude nucleoprotein and polysaccharide fractions of bacteria

J. Allergy, 20:418, 1949

After alcohol precipitation of crude polysaccharide from the colon bacillus, *Ps. pyocyaneus*, *Streptococci* (*hemolyticus viridans* or *fecalis*), and the causative organisms of respiratory infections, nucleoprotein could be separated by precipitation with dilute acetic acid.

587. HOSOYA, S., HOMMA, Y., EGAMI, F. and YAGI, Y.

On the isolation of antigenic substances from Pseudomonas aeruginosa (Translated title)

Jap. J. Exper. Med. 20:55-68, 1949

An antigenic nucleoprotein was prepared from the Tsuchijima strain of Ps. aeruginosa by precipitation with $ZnCl_2$, dialysis and acetone precipitation. It was heat stable ($100^\circ C.$, 1 hr.) and contained 5.8% carbohydrates and 11.5% N. The nucleoprotein was precipitable with immune rabbit sera in dilutions to 1:250,000. It conferred immunity to a homologous strain of living bacilli.

588. KORNERUP, T.

Damage to visual organs resulting from typhoid inoculations

Acta Ophthalmol. 27:311-454, 1949

After typhoid vaccine therapy, optic atrophy, neuritis or neuroretinitis, temporary color blindness or muscular paralysis appeared in certain patients. Allergy or activation of latent virus infection may play roles in such conditions.

589. LEESE, C. E. and GREENE, K. V.

Physiological effects of bacterial polysaccharides upon the cardiovascular and neuromuscular systems

Federation Proc. 8:93, 1949

In the rabbit injections of polysaccharide ranging from 2 to 25 gammas per kg. were administered. At higher dosage levels, disturbances of nervous reflexes, convulsions, urination, defecation and vomiting preceded death. In cats, arterial blood pressure fell while venous pressures increased. A loss in autonomic controls was seen; the vagus, splanchnic and cardiosympathetic nerves required increased stimuli after polysaccharide was administered. Decerebration was fatal when the drug was administered prior to operation. Intense cardiovascular changes with heart block and dilatation of chambers were seen in the turtle perfused with polysaccharide preparations.

590. LEHMANN, H.

Beginning of a chemotherapy of neoplastic ailments (Translated title)

- 1 9 4 9 -

Schweiz. Apoth. Zeit. 87:741-743, 1949

Pyrogens, cytostatic preparations, radioactive materials and the hormone therapeutic preparations are discussed.

591. LEQUIRE, V. S.

Augmentation of the thermogenic effects of pyrogens by homologous plasma in rabbits.

Naval Medical Research Inst. Project NM007047. Report #6, 1949
See: Lequire, J. Infect. Dis. 88:194-206, 1951

592. LIBERALI, C. H.

Metodos de Ensaio para pirogenio (Tests for and elimination of pyrogens)

Anais faculdade farm. odontol, Univ. Sao Paulo 7:129-170, 1949

Better results are secured when rabbits used in pyrogen testing are carefully checked for temperature changes 4 or 5 times a day for two to three consecutive days before the day of pyrogen administration. A group average for 5 rabbits should be computed. Pyrogens may be eliminated to varying extents by $Al(OH)_3$, $Fe(OH)_3$ and $Cu(OH)_2$.

593. LONSEN, W. and LIEBERT, E.

Studies on a new pyrogen fever treatment

Illinois Med. J. 96:186-190, 1949

PYROMEN is a nonprotein, bacteria-free product, free from toxic substances, whose action in the treatment of neurosyphilitic patients compares favorably with those secured through malarial and typhoid vaccine fever therapy. It appears to be less dangerous than malarial therapy, and has been used, in this study, for older patients. Because of the small dosages used, administration is not difficult. Its action in these cases appears to rest in a strong mobilization of body defense mechanisms.

594. MAHONEY, V. P. and HAMMERMAN, S.

Combined treatment of symptomatic paresis with penicillin and fever produced by continuous typhoid vaccine

J. Nerv. & Ment. Dis. 109:133-141, 1949

Although continuous intravenous typhoid vaccine therapy is practicable for adjuvant fever therapy with penicillin, some selection of patients would improve the results obtained in these investigations. The two criteria used were (1) no clinical evidence of cardiovascular disease (2) ages under 60 years.

595. MARTELLI, F. and NUROLONE, U.

La piretoterapia nel trattamento della paralisi facciale primitiva con particolare riferimento alla forma inveterata (Pyretotherapy in the treatment of primary facial paralysis, with particular reference to chronic cases)

Minerva med. (Turin) 2:403-406, 1949

In primary idiopathic facial paralysis, but not in secondary manifestations associated with more fundamental causes, benefits were secured through use of agents, physical or otherwise, which elevated the temperature. Results would support the hypothesis that primary facial paralysis may be of allergic etiology, with fever acting, in these improved cases, as anti-allergen.

596. MCGINTY, D. A., WILSON, M. L. and RODNEY, G.

The ulcer-inhibiting action of pyrogens

Proc. Soc. Exper. Biol. & Med. 70:334-336, 1949

Extracts from urine known to possess pyrogenic activity have inhibited ulcer formation in rats. These observations have been extended to the use of purified substances from cultures of B. prodigiosus, Ps. aeruginosa and E. typhi, given by vein in 0.5 ml. volume prior to pyloric ligation in the rat, or by stomach tube during the period of recovery from anaesthesia. Pyrogens from B. prodigiosus (3 to 10 gammas) protected 18 of 28 rats against ulcer (index 72); the index fell to 15 when dosage levels were raised to 35 or 50 gammas. Intraperitoneal injections were less effective and higher dosages were required (100-1000 gammas). Typhoid pyrogens inhibited ulcer formation in 75% of the rats under therapy. Ps. aeruginosa pyrogen was less effective than that secured from cultures of B. prodigiosus. Non-pyrogenic polysaccharides (pneumococcal carbohydrates) had no anti-ulcer activity. The mechanism of action by which results are obtained is not clear, but both volume and acidity of gastric secretions appear to be reduced.

597. MENKIN, V.

An additional thermostable component in the mechanism of leukocytosis with inflammation

- 1 9 4 9 -

Federation Proc. 8:362, 1949

The thermostable, pyrogenic factor, pyrexin, induces the initial leukopenia. It is found in the euglobulin fraction in the acid exudates of inflammatory processes. A second thermostable component appears to be in close association with pyrexin and to play a part in the induction of high levels of leukocyte formation.

598. MORGAN, H. R. and NEVA, F. A.

Tolerance to the toxic effects of somatic antigens of enteric bacilli in typhoid and paratyphoid fever convalescents.

J. Clin. Invest. 28:800, 1949 (Abstract)

Intravenous test doses of the endotoxin of S. typhosa (5 micrograms) produced no evidences of the marked febrile responses and reactions seen in normal individuals when it was administered to convalescent patients after infections with typhoid or paratyphoid organisms. Shigella dysenteriae (7 micrograms) somatic antigen was not productive of febrile reactions after resistance had developed following spontaneous infections.

599. MUETHER, R. O.

Pyrogens in intravenous solutions

Hospital Prog. 30:301, 1949

The author briefly reviews the causes of pyrogen reactions and their sources, symptoms and treatment, and the manner in which bioassay of parenteral fluids may be conducted.

600. NECHELES, H. (Editor)

21st Annual Report, Med. Res. Institute, Michael Reese Hospital 1949

The observation has been made that even minimal amounts of certain bacterial pyrogens may depress normal functions of the stomach, ie. secretion of acid and normal emptying time. The reduction of gastric spasm in ulcer conditions might aid healing processes. Doses of pyrogen too minute to produce fever depress secretory activity, and it is tentatively stated that these substances may be responsible for therapeutic results achieved by certain preparations commonly prescribed in ulcer therapy.

601. OLITZKI, L.

Anti-leukopenic tolerance of rabbits and antibody formation in the course of treatment with typhoid vaccine

Acta Med. Orient. 8:103, 1949

Tolerance to the leukopenic effects of typhoid vaccine developed in rabbits during treatment and was unrelated to development of O antibodies in the blood serum. The tolerance was noted most frequently after intra-abdominal administration. Leukocytes collected in pulmonary areas during the period of leukopenia and were released in the succeeding phase.

602. OTT, W. H.

A quantitative assay method for pyrogens

J. Am. Pharm. A. (Sci. Ed.) 38:179-184, 1949

When test animals were carefully selected for similarity of response to a standardized pyrogen (a standard error variation computed on the basis of the assay values in 4 rabbits was found to be minus or plus 60%), a method for reducing error was developed, which would greatly increase the number of animals used in assay. The standard error is inversely proportional to the square of the number of animals used in the assay. The correction can be aided further by standardization procedures adapted by each testing laboratory and preparation of dosage-response curves for groups of animals tested under these conditions.

603. PAGE, I. H. and CORCORAN, A. C.

"Treatment with Pyrogen." Arterial Hypertension. Chicago; The Year Book Publishers, Inc. 2nd Edition, 1949, pp. 328-330

Thirty-five patients were treated with the pyrogenic agent PYROMEN for periods extending from 5 to 19 weeks. A good clinical response was shown by 19 patients. The most marked difference between improved and unimproved groups appeared to be the presence of arteriolar sclerosis. Delayed response to pyrogens in both normal and hypertensive patients involve vasodilatation, particularly in the renal vascular bed. Increased renal blood flow usually persists for 24 to 36 hours after the pyrogenic reaction. In the above group, patients who improved averaged 62 mg. per minute per 1.73 sq. m. body surface as shown by TMPAH readings and only 26 mg. in unimproved patients. Blood pressure was reduced; there was regression of retinal hemorrhage and absorption of retinal exudates. Congestive heart failure resistant to therapy improved in five instances. Immediate hemodynamic

- 1 9 4 9 -

responses and adrenal plasma and tissue enzyme interrelationships may enter into the mechanism of response to pyrogens. For hypertension, pyrogen treatment must be given under careful supervision; hospitalization is required.

604. PAGE, I. H. and TAYLOR, R. D.

Pyrogens in the treatment of malignant hypertension

Mod. Concepts Cardiovasc. Dis. 18:51, 1949

PYROMEN is considered to be the most satisfactory agent in the treatment of malignant hypertension, according to the results evaluated by the authors. Twenty patients have been treated with this agent over a period of four years. All selections for this therapy were based on the presence of renal function adequate enough to excrete para-amino hippurate at 35 mg./min./1.73 sq. m. body surface, or to concentrate urine at specific gravity of 1.020 or above. PYROMEN was administered intravenously daily for 5 or 6 days per week. The initial dose was 50 micrograms, successive doses were increased to maintain a temperature rise to 103 or 104° F. each day. When tolerance developed, treatment was discontinued for 3 to 7 days. Of 20 patients so treated, ten survived. Of five who discontinued treatment before results were obtained, 4 died, four patients who exhibited early improvement developed essential hypertension later. Where cardiac decompensation existed PYROMEN therapy appeared to be of value.

605. PATTERSON, J. L., HEYMAN, A. and NICHOLS, F. T.

Effects of fever induced by bacterial pyrogen on cerebral circulation and oxygen consumption

Am. J. Physiol. 159:584, 1949 (Abstract)

Cerebral blood flow and oxygen consumption levels were determined in 16 patients with asymptomatic neurosyphilis after intravenous injections of typhoid vaccine. Comparisons were made with readings secured during the afebrile state. Cerebral vascular resistance was significantly decreased (average 25.4%) as the rectal temperature increased to a mean 3.8° F. above previous readings. The blood flow was unchanged. When vascular resistance was above the normal in the afebrile state, values were higher. Mean arterial pressure reductions of 2.4% accompanied cerebral vascular tension decreases. There was a high degree of correlation between oxygen consumption and cerebral flow during fever, but these levels were unrelated to each other during afebrile periods.

606. PEISS, C. N., FIELD, J., HALL, V. E. and GOLDSMITH, M.

Comparison of respiration and glycolysis in the brains of normal and febrile rabbits

Am. J. Physiol. 157:283-286, 1949

In comparing oxygen consumption and anaerobic glycolysis rates in both febrile cortical tissue and normal preparations, no statistically important differences could be ascertained, nor was there support for the thesis that the cellular metabolism of the central nervous system is a limiting factor on thermic regulation of the animal.

607. PHILLIPS, K.

The clinical response to vaccine in 125 cases of rheumatic disease

Rheumatism 5:53-57, 1949

Rheumatoid osteo-arthritis, myositis and fibrositis improved in a study covering 125 patients. A stock mixed vaccine was employed.

608. PINCHOT, G. B., CLOSE, V. P., and LONG, C. H. N.

Adrenal changes produced in rats by infection with E. coli

Endocrinology, 45:135-142, 1949

Rats were given large doses of living E. coli organisms intravenously. The adrenal cortical response was similar to that produced by a single injection of ACTH in its effect on adrenal size, ascorbic acid and cholesterol levels. Dosages of adrenal cortical hormone many times larger than are required for the maintenance of adrenalectomized animals failed to prevent a depletion in adrenal cholesterol or prolong survival time after infection with B. tularensis. Manifestations of stress were present.

609. ROBERTSON, H. F., KISSEN, M. D., FLOTHOW, M. W.

The inherent dangers of fever therapy in the treatment of paresis

Urol. & Cutan. Rev. 53:409-412, 1949

The literature on malarial therapy in paresis is reviewed, and results achieved with typhoid fever therapy assessed. With all types of therapy a 50% improvement may be expected, but typhoid

- 1 9 4 9 -

vaccine therapy was productive of results 10% better than those achieved with other forms employed. However, a decrease in plasma proteins, the onset of renal dysfunction and certain vascular changes are contraindications to further therapy.

610. ROSA, M. C.

Loss of pyrogenic activity by solutions prepared for intravenous injection (Translated Title)

Rev. brasil. farm. 31:121-127, 1949

"...pyrogenic activity disappeared from physiologic saline solution after 71 days. Isotonic glucose solutions prepared and sealed in an atmosphere of CO₂...pH 4.5-4.8, lost activity after 41 or 62 days respectively...(when) prepared without these precautions and with pH 6.8 retained activity 53 days.

pH between 4.5 and 6.8 did not influence the rate of loss of pyrogenic activity. Concentration may be a more important factor.

611. RZUCIDLO, L.

Odczyn opadania ciałek czerwonych jako metoda roznicowania wielocukrow bakteryjnych (Erythrocyte sedimentation rate in the differentiation of bacterial polysaccharide)

Med. Doswiad. i. Mikrobiol. (Warsaw) 1:40-51, 1949

The erythrocyte sedimentation rate is increased after administration of polysaccharide-lipid-protein complexes from bacteria of the Salmonella group, or their carbohydrate derivatives. Other polysaccharides in suspension fluids do not enter into such complexes and have no effect on sedimentation rates.

612. SCHEINMAN, L. J.

The use of pyrogens in the treatment of lower nephron nephrosis

J. Urol. 61:1003-1009, 1949

Inconclusive results during a study of the effect of increased renal flow on symptomatology of lower nephron nephrosis suggest that the rabbit may not be the best animal for use in the study of the problem because it possesses a most efficient reticuloendothelial system.

613. SELIGMAN, A. M. and SACK, T.

Chemical alteration of polysaccharide from Serratia marcescens; Reaction of polysaccharide with radioactive paraiodobenzene-diazonium chloride and use of product in experimental and chemical study

J. Nat. Cancer Inst. 10:105-118, 1949

The preparation of a radioactive iodobenzene azo product is described. Toxicity of the radioactive product was similar to the unmodified polysaccharide of S. marcescens as was shown by changes following the administration of the carbohydrate in three patients with malignant tumors.

614. SHETLAR, M. R., BRYAN, R. S., FOSTER, J. V., SHETLAR, C. L. and EVERETT, M. R.

Serum polysaccharide levels in experimental inflammation

Proc. Soc. Exper. Biol. & Med. 72:294-296, 1949

In experimental inflammation, bacterial, or, as in this study, produced by irritant agents such as turpentine or talc, a delayed but impressive increase in serum polysaccharide was recorded. Since maximum elevation does not appear until three to six days after injury, the role of tissue destruction is believed to be minor or non-existent. Regenerating tissue may be responsible for the elevation which is independent of temperature.

615. SHETLAR, M. R., FOSTER, J. V., KELLY, K. H., SHETLAR, C. L. BRYAN, R. S. and EVERETT, M. R.

The serum polysaccharide level in malignancy and in other pathological conditions

Cancer Res. 9:515-519, 1949

Serum polysaccharide levels in neoplastic disease are higher than the peak normal levels. In benign tumors, no production of abnormal levels was noted. Levels were lower in carcinoma of the skin and of the breast. After surgical removal blood levels increased. The increases may be connected with tissue proliferation and repair. The serum polysaccharide is increased in many chronic conditions and most infectious processes, hence levels are not specifically diagnostic for neoplastic disease.

- 1 9 4 9 -

616. WHARTON, D. R. A., and CREECH, H. J.

Further studies of the immunological properties of polysaccharides from Serratia marcescens (Bacillus prodigiosus). II. Nature of the antigenic action and the antibody response in mice.

J. Immunol. 67:135-153, 1949

A tolerance which reached its height on the third day after a single injection of P-10 polysaccharide-lipid complex of S. marcescens protected mice against a lethal dose of the polysaccharide. Heating did not diminish ability to produce antibodies but did reduce toxic properties. The agglutinating antibody reached high titer levels on the 10th day after administration of the substance. Anaphylaxis was not produced (in guinea pig). Two fractions may be present in the complex, a toxic and an immunizing component.

617. YAGI, Y., EGAMI, F., HOSOYA, S., NONOBE, K., HAYASHI, T. and HOMMA, Y.

Studies on the endotoxine of Salmonella enteritidis, Gartner. I. Studies on the chemical nature of the endotoxin. II. Studies of the amino acid content and the sugar component of the endotoxin

Jap. J. Exper. Med. 30:245-259, 1949

A crude endotoxin was prepared from S. enteritidis and subjected to fractional precipitation. The resulting concentrate contained 2.9-3.6% N, 1.1-1.58% P, 2.1% S and 43-55% reducing sugars. Positive reactions to the Bial and Dische tests for carbohydrates were noted. Absorption curves reached their peaks at 2400-2700 Angstrom units. Three fractions were secured by heating with acetic acid: a nucleoprotein (MLD 1 mg.), a polysaccharide, non-toxic, which contained methyl-pentose, pentose and desoxy-sugars. The methyl-pentose fraction exhibited two hydrazones and a phenylosazone which resembled those found in the breakdown of fucose.

618. ALBOT, G. and BERNARD, BRUKER, N.

Nouveau traitement des diarrhees par les emulsions de Bacillus Subtilis (New treatment of diarrhea by emulsions of B. subtilis)

Sem. hopit. (Paris) 3400-3403, 1950

619. ALTSCHULE, M. D., PARKHURST, B. H., and PROMISEL, E.

Effects of intravenous injections of typhoid vaccine on blood leukocytes and adrenal cortex

A.M.A. Arch. Int. Med. 86:505-518, 1950

When typhoid vaccine was injected intravenously into psychotic patients (6), cellular redistribution in blood was observed, with lymphocytes and eosinophils decreased by approximately 80%, and granulocytes increased from 50 to 200% after each treatment. The lymphocyte count returned to normal after three days. No correlation with the degree of fever produced could be measured. Electirc shock therapy did not produce such startling changes (eosinophils decreased 48%, lymphocytes but 16%). Changes in the urinary output of 17-ketosteroids were seen. Urine uric acid: creatinine ratios varied inversely to the degree of vaccine fever.

620. ARENDSHORST, W. and FALLS, H. F.

Role of the adrenal cortex in treatment of ocular diseases with pyrogenic substances

A.M.A. Arch. Ophth. 44:635-642, 1950

This study was planned to determine the effects of "protein shock" therapy in ocular disease and as an investigation of the possible release of 11, 17-oxysteroids from the adrenal cortex. The circulating eosinophil response to injections of the antigen was taken as the most sensitive index of cortical activity. Typhoid H antigen was used as a "foreign" protein, and in dosages ranging from 15 to 25 million units initially to higher concentrations, was administered to patients with various uveal diseases.

Correlation was noted between eosinopenia, fever and the general response to the pyretogenic agent. Six of 21 patients had poor eosinopenic responses and proved to be resistant to foreign protein "shock". Drop in eosinophils usually occurred about 18 hours following injections. Improvement in ocular tissues was seen in 14 patients. The effects of pyretotherapy were attributed by the authors to the initiation of an "alarm" reaction and succeeding stimulation of the adrenal cortex, but not satisfactory explanation for the mechanism involved could be produced.

- 1950 -

When foreign protein therapy induces a refractory condition, the use of ACTH or cortisone is advised; the adrenalcortical factors are useful when pyrogenic therapy no longer produces eosinopenia or must be prolonged for longer than one week.

621. ASHENBURG, N. J., SANDHOLZER, L. A., SCHERP, H. W., and BERRY, G.P.

The inhibition of bacteriophagy by bacterial and nonbacterial polysaccharides

J. Bact. 59:681-687, 1950

Relationships between polysaccharide and bacteriophage resemble those existing between antigen and antibody, the former existing as a surface antigen and inhibiting lytic action. Nonspecific inhibition has been credited to starches, glycogen and gums. In this study the effect of nonbacterial polysaccharides and those obtained from capsules of Aerobacter and Klebsiella was to inhibit the lytic action of Friedlander Bacillus bacteriophage. All exhibited this activity, but inhibition was not complete; it varied with the polysaccharide used and the host organism.

622. BECK, L. V. and VOLOSHIN, T.

Influence of adrenal hormones on toxic and tumor-damaging effects of certain substances

A. J. Physiol. 163:696-697, 1950 (Abstract)

Cortisone and adrenal extract may counteract the damage wrought by tumor-necrotizing substances on sarcoma #37. Mice injected either intraperitoneally or subcutaneously with cortisone exhibited increased tolerance for arsenite administered one hour later by either intravenous or subcutaneous routes. When adrenal extract was injected peritoneally, there was an increased tolerance for both arsenite and mapharsen, and some protection against arsenite could be secured by previous injection of ACTH. The increase in tolerance was greater in the male and at the L5 level. Cortisone or adrenal extract, when administered before intraperitoneal injections of the tumor-necrotizing polysaccharide of S. marcescens, increased tolerance, and cortisone also afforded some protection against the intravenously injected polysaccharide. Protective patterns differed from those invoked against arsenicals. The adrenocortical factors do not protect mice against lethal or sublethal amounts of alpha peltatin or acridines.

623. BENNETT, I. L., JR., and BEESON, P. B.

The properties and biologic effects of bacterial pyrogens

Medicine 29:365-400, 1950

In a comprehensive survey of the literature of pyrogenic substances extending from 1855 to the present day, the bacteriology, physical characteristics and chemical properties of these substances are outlined and discussed. In turn, the application of pyrogens to therapeutic uses and physiologic response to their administration are outlined in detail. "The role of pyrogens in disease is still incompletely understood." A bibliography containing 246 references is appended.

624. BERGSMAN, A. and HEDENIUS, P.

De pyrogena substansernas betydelse vid parenteralt bruk av vitskor (Problem of pyrogens in the use of parenteral solutions)

Nord Med. 43:672-676, 1950

625. BLICKENSTAFF, D. and GROSSMAN, M. I.

A quantitative study of the reduction of gastric acid secretion associated with pyrexia

Am. J. Physiol. 160:567-571, 1950

Pyrexia produced by intravenous pyrexin or bacterial pyrogen was not directly responsible for inhibition of secretory activity in the dog. PYROMEN, prepared from cultures of *Ps. aeruginosa*, in doses ranging from 0.5 microgram per kg. to 10.0, and pyrexin, from sterile pleural exudate, 2 to 30 mg. were administered to 'gastric pouch' dogs and secretions were collected for three or more hours thereafter. A preliminary injection of histamine, 0.0125 mg. subcutaneously, was repeated at frequent intervals throughout the test. In no test was there depression of secretion without a rise in body temperature. When only slight increases in temperature were registered, secretion was not inhibited. However, a lag of approximately one hour between onset of pyrexia and reduction of gastric secretion, as well as the extension of inhibition past the period of temperature reduction, would appear to rule out pyrexia as the primary agent in this phenomenon. The state of hydration did not affect results.

626. BLUMENTHAL, L. S. and FUCHS, M.

Headache Clinics. I. What is the headache problem?

Am. Pract. ? Digest Treat. 1:1012-1015, 1950

Work on ACTH and PYROMEN, which appears to act in a manner similar to adrenal cortical derivatives, is still in the experimental

- 1950 -

stages. Proper application of PYROMEN, the antihistamines, ACTH, potassium thiocyanate, banthine and dehydrogenated alkaloids of ergot will be described in later publications.

627. BOQUET, P. and IZARD, Y.

Effect of dibenamine on vascular response of rabbits to typhoid vaccine

Proc. Soc. Exper. Biol. & Med. 75:254-259, 1950

The intense initial peripheral vasoconstriction produced in rabbits by an injection of typhoid endotoxin was noted in earlier studies. It was believed to be caused by the release of adrenalin (or adrenalin-like substances) from tissues. To confirm these findings, the role of adrenolytic agents in inhibiting or modifying vascular response was tested by injecting 0.5 to 0.8 ml. per kg. of S. typhosa endotoxin into the rabbit and following this with intravenous dibenamine (6-10 mg. per kg.) Cutaneous circulation was observed by three methods, two visual and one involving the use of thermocouples. Dibenamine modified vasoconstriction when administered either before or after the endotoxin, but, for sustained effect, blocking was necessary before and after the vaccine.

628. BOVE, E.

Nogle problemer i. pyrogenbestemmelsen (Problems of pyrogen determination)

Sven. farm. tidsskr. 54:721-726, 1950

629. CARLETTI, B.

La vaccinazione aspecifica intradermica nella malattia di Heine-Medin. Parte I. Storia, tecnica e meccanismo d'azione (A specific intradermal vaccination in poliomyelitis. I. History, techniques and mechanism of action)

G. Mal. Infett. parassit (Florence) 2:180-187, 1950

Intracutaneous injection of 500 to 5000 million nonspecific organisms in a vaccine preparation was made every 4 to 7 days for 20 injections. The treatment was often repeated 2 to 3 months later. The site of injection was the cutis of the lower limbs, and administration was usually made at several points in the area. Among the organisms present in the killed state in the vaccine, E. coli, Pr. vulgaris, the pyogenic streptococci, M. pyogenes and catarrhalis and S. paratyphosa A and B are noted. Seventy of 242 patients treated improved; circulation increased and neuromuscular function was benefited.

630. CHARY, R. and ARDRY, R.

Note sur le controle pyrogenique des solutions citratées utiliseés dans le transfusion sanguine (Notes on the pyrogenic control of citrated solutions used in blood transfusion)

Ann. biol. clin. 8:504-506, 1950

631. CLEMENTE, C. D.

Leukocytosis in the rat upon oral administration of a bacterial pyrogen

Am. J. Physiol. 163:703, 1950

Significant leukocytosis was observed when PYROMEN, 1 gamma daily, was administered to the rat. White cell counts increased from normal levels (15,000 cells per cu. mm.) to 35,000 or 40,000 cells per cu. mm. after 1 to 4 gammas of PYROMEN, but tolerance became evident after sustained dosages at constant levels, and white counts resumed normal levels. While increased doses were ineffectual during the period when the leukocyte count was returning to normal levels, a secondary leukocytosis could be elicited after a rest period of 6 days; the response was of shorter duration and lower intensity than the first. The percentages of granular cells to the non-granular types did not alter with the rising count. In only two of four animals observed was leukopenia noted. In two animals, circulating eosinophils showed an initial decrease but increased during the leukocytosis along with other granular cells. Histologically, no pathologic changes could be demonstrated in endocrine, digestive, lymphatic or excretory organs after oral administration of PYROMEN.

632. CORELLI, F.

L'importanza del pirogeno nella pratica medica e trasfusionale (contributo alla profilassi delle reazioni da trasfusione, da ipodermoclisi, fleboclisi, da endovenose di calcio, ecc.) (Importance of pyrogen in medical transfusion practice; contribution to prophylaxis, reactions due to transfusion, to hypodermoclysis, to phleboclysis, to intravenous calcium, etc.)

Policlinico (sez. prat.) 57:345-347, 1950

Reactions may occur even when depyrogenated solutions are employed since glassware, tubes, needles, or residues of blood and plasma may produce pyrogens, which after sterilization, continue to act as "denatured" pyretogenic agents. A thorough cleaning must precede sterilization, for elimination of possible pyrexial materials.

- 1 9 5 0 -

633. DORCHE, J., and CASTAING, M.

Sur la recherche des substances pyrogenes dans les solutes injectables; les essais basés sur la leucocytose du lapin (Research on pyrogenic substances in injectable solutes; tests based on leukocytosis of the rabbit)

Ann. pharm. franc. 8:353-357, 1950

The leukocyte count is too variable in the rabbit for use as a test factor for pyrogens in solutions intended for parenteral injection. Injection of pyrogen-free solutions and observations of the increases in the polymorphonuclear cell count should precede the administration of pyrogenic solutions. Differences between the grades of polynucleosis should be recorded for use as a test control measure.

634. DORCHE, J. and CASTAING, M.

Sur la recherche des substances pyrogenes dans les solutes injectables; atténuation des propriétés pyrogenes au cours de la conservation des solutes (Research on pyrogenic substances in injectable solutes. Reduction of pyrogenic properties during the storage of solutions)

Ann. pharm. franc. 8:365-368, 1950

Prolonged storage of solutions reduces the pyrogenic factors in most solutions. Glucose solutions are exceptions.

635. DORCHE, J., BOUTHIER, G., ARDIET, M. T., and CASTAING, M.

Sur la recherche des substances pyrogenes dans les solutes injectables; Commentaires et application de l'essai de la Pharmacopée française (Research on pyrogenic substances in injectable solutes. Commentary on and application of the bio-assay according to the French Pharmacopoeia)

Ann. pharm. franc. 8:358-365, 1950

Factors which influence the bio-assay may be physiologic or chemic in nature. Hypotonic solutions may yield false positive readings for pyrogenic content. The sensitiveness of the rabbit to handling is noted; special precautions must be taken in placing the rectal thermometer during the test.

636. FARR, R. S. and LEQUIRE, V. S.

Leukocytic and pyrogenic effects of typhoid vaccine and augmentation by homologous plasma

Proc. Soc. Exp. Biol. Med. 75:661-666, 1950

These experiments were designed to study the correlation between the magnitude of leukocytic and thermogenic responses to pyrogen in the rabbit and the amount of pyrogen administered. Vaccines from 2 strains of *E. typhosa* were used. Fifty-eight male rabbits were divided into four groups; in each group one of the following was injected intravenously: physiologic saline solution; saline plus typhoid vaccine; normal rabbit plasma; normal plasma with typhoid vaccine. Animals were observed for six consecutive hours, during which total and differential leukocyte counts and rectal temperatures were frequently measured

Saline and homologous plasma alone were without effect. Plasma-typhoid injections produced high fever, initial heterophile leukopenia and secondary heterophile leukocytosis. Saline-typhoid solutions produced only initial heterophile leukocytosis. Both these groups developed mononuclear leukopenia with the plasma-typhoid combination producing a more pronounced drop. By increasing the number of organisms in the saline suspension tenfold, it was possible to produce a response similar to that seen after plasma-typhoid injections, and conversely, reduction of cells in plasma-typhoid suspensions brought an initial heterophile leukocytosis. Five variables must be considered when using the leukocyte count as an index for the pyrogen test: amount of pyrogen; time at which the observation is made; differential composition of the leukocyte population; the type of bacteria in the source; the media used for suspensions.

637. Fever therapy in asthma

Queries and Minor Notes. J.A.M.A. 144:1419, 1950

"The salutary effect of acute febrile disease on allergic manifestations has been known for centuries. Chronic and intractable asthma frequently becomes more completely relieved by intercurrent non-respiratory infection than by almost any other means." The mechanism by which benefits are achieved is not clearly defined. Typhoid vaccine, colloidal sulfur and malarial therapy have been employed as has high frequency current. "More recently a pure carbohydrate constituent obtained from certain bacteria and known as PYROMEN has been used to produce fever."

638. FLENSBORG, E. W.

General reactions following treatment of asthma in children with bacterial vaccine (English)

Ugeskr. f. Laeger 112:392-399, 1950

- 1 9 5 0 -

Reactions to administration of the common pathogenic respiratory bacteria in the treatment of asthma of childhood become more severe during extended treatment. These are considered to be late reactions occurring, on the average, after two days. They may continue from 6 hours to 6 days. Repeated overdosage may induce prolonged temperature increases or status asthmaticus. Best treatment is prophylactic application of a minute dose, careful increases in dosage with attention to even slight reactions, and frequent temperature recordings during the first four days.

639. FLENSBORG, E. W., NEERBORG, G. and SAMSE-JENSEN, T.

Probable bacterial-allergic asthma in children and its treatment (English)

Ugeskr. f. Laeger 193:387-392, 1950

Children, one to four years of age, with asthma, but giving a negative cutaneous reaction to usual allergens, may have a bacterial allergy as an aftermath of early infection in the upper respiratory tract. Clinically, symptoms are identical with those seen in patients who have positive cutaneous reactions. Eosinophilia is noted in both groups. The authors have used a special standard vaccine prepared by the State Serum Institute with better results than are seen after specific desensitization in children with positive cutaneous reactions. General reactions, however, may be marked even with small doses of vaccine. Treatment should be continued one year after disappearance of the clinical symptoms.

640. FRANZOLIN, C.

Sulle modificanzioni leucocitarie provocati della piretoterapia nelle m. infiammatorie acute e croniche (Leukocytic modifications induced by pyretotherapy in acute and chronic inflammatory diseases)

Acta med. (Patav.) 11:194-199, 1950

641. FREIMULLER, F., and RHOMAKO, M.

Über die Frubehandlung der poliomyelitis anterior acuta mit pyrogenen substanzen (Pyrogenic substances in the early treatment of acute anterior poliomyelitis)

Wiener Ztschr. inn. Med. 31:501-506, 1950

642. GOSSELIN, W.

Effets des polysaccharides sur le sarcome de la souris (Effects of polysaccharides upon the mouse sarcoma)

Rev. brasil Cancer 3:27-50, 1950

While mice tolerate dosages of the polysaccharides of S. marcescens 1000 times higher per unit of body weight than do human beings, 25 to 50% of the animals die with 24 hours. A method of reducing toxicity of the material is a prerequisite for its use in extensive cancer therapy. Antihistamines do not reduce toxicity, nor does vitamin B₆. Rutin (1.0 ml. of 0.1% to 50 gammas of the polysaccharide) brought about the survival of 11 out of 12 animals, and did not affect the hemorrhage-inducing and necrotic qualities, which suggests that the hemorrhagic action of the polysaccharides does not depend on capillary fragility.

643. GRANT, R.

Emotional hypothermia in rabbits

Am. J. Physiol. 160:285-290, 1950

A transient hypothermia may be induced in rabbits by light restraint as well as by any activity which may activate heat loss mechanisms. Cold, applied or environmental, may inhibit the phenomenon, as do pyrogen injections. When 5 gammas of PYROMEN, prepared from Ps. aeruginosa, were given intravenously to a restrained animal that had recently regained its normal temperature after a hypothermic period, interruption of the fever began about 1 hour after injections as shown by ear vasodilation and polypnoea. The temperature fell to 1.25° below normal then stabilized at this level. In the unrestrained animal an initial drop in temperature was succeeded by an intense febrile state with no subsequent hypothermia. Modification of cerebral cortical control upon thermoregulation is postulated. Pyrogens may act on the motor mechanism of thermoregulation and not on higher centers of control. Animals used in pyrogen testing should not be restrained.

644. GRANT, R. and HIRSCH, J. D.

Pyrogen fever in rabbits; Effects of adrenalectomy

Am. J. Physiol. 161:528-533, 1950

In a preliminary study, changes in rectal and ear temperature and

- 1 9 5 0 -

in the respiratory rate of the adult rabbit were recorded after the intravenous injection of 0.1 ml. per kg. of PYROMEN, containing 50 gammas per ml. of purified pyrogen. Seven days later a comparable amount of PYROMEN was injected and the effect on oxygen consumption observed in a part of the group. Bilateral adrenalectomy was then performed in two stages, 2 to 6 weeks apart and DCA implanted subcutaneously. After stabilization of rectal temperature and body weight, the effect of PYROMEN was again studied. When animals were maintained in a healthy state through use of DCA, adrenalectomy affected the course of pyrogen-induced fever but little. It is suggested that the removal, by adrenalectomy, of inhibition exerted on shivering by epinephrine may underlie a mechanism which leads to calorogenesis.

645. GRANT, R., HIRSCH, J. D. and HIRSCH, B. B.

Pyrogen fever in rabbits: Effects of adrenalectomy and thyroidectomy

Federation Proc. 9:50, 1950

Shivering appeared to increase oxygen consumption following adrenalectomy with subsequent maintenance on DCA. No other effect on the normal responses to pyrogen injections could be remarked. Thyroxine, after thyroidectomy, had no effect upon normal responses to pyrexial agents. While neither the adrenals nor thyroid are essential intermediaries in fever production, the mechanism by which heat is produced may be altered in the adrenalectomized animal; increased shivering with subsequent thermal response may be caused by shivering precipitated by the removal of the inhibition normally exerted by epinephrine.

646. HARKNESS, W. D., LOVING, W. L. and HODGES, F. A.

Pyrexia in rabbits following injection of filtrates of typical mold cultures

J. Am. Pharm. A. 39:502-504, 1950. Abstract, Federation Proc. 9:282, 1950

When injected intravenously, preparations from 16 out of 31 species of mold cultures brought elevations in the body temperature of the rabbit. A medium containing dextrose and peptone (Sabouraud) was required for production of pyrogenic characteristic inorganic salts and N sources in the medium restrained its development.

647. HARKNESS, W. D. and VOS, B. J.

Comparison of the effects of injection of pyrogenic solutions by

intravenous and intramuscular routes

J. Am. Pharm. A. (Sci. Ed.) 39:413-414, 1950

Rabbits were given intravenous injections of a pyrogenic solution prepared by the method of Welch in amounts ranging from 0.001 cc. per kg. to 0.1 cc., and the same dosages in more concentrated aliquots of solution were administered intramuscularly. The ratio of the intramuscular to the intravenous response to the same amount of pyrogen was computed. Six rabbits were used at each dosage level and a rise of 0.6° (U.S.P. XIII assay) was considered significant. All pyrogen dosages gave significant temperature rises when administered intravenously. At any given dose this route produces higher rise in temperature. Responses to intramuscular doses of pyrogen were graded and lower for any given dosage. "When the temperature response is plotted against the logarithm of the dose a linear relationship is seen to exist within certain limits."

648. HATTA, S., AOYAMA, K., and TANJI, S.

Studies on the bacterial pyrogenic substances

Jap. Med. J. 3:125-135, 1950

A simplified method for detecting pyrogens in solution is described. Pyrogens were extracted from pyrogen-positive strains of bacteria (*sarcinae* and *micrococcus* sp.); from three strains of fungi (*penicillium* sp., *fusarium* and *macrosporium*) and from *S. typhi* and *S. paratyphi*. Pyrogenic materials were found in both mycelia and metabolic products. When mycelia were crushed, extracted, and filtered and the filtrates condensed and crystallized, a yield of 80 to 100 mg. was secured from 50 Gm. of cultured mycelia. The crystals were colorless and had melting points of 163° . Tests for protein and for sugar (Fehling and Molisch) were negative. There was a C content of 37.7%, of H, 8.12%. Pyrogens fluoresced blue and were dialyzable through semi-permeable membranes. When injected into rabbits 15 mg. per body weight, they produced temperature increases ranging from 1.5 to $2.0^{\circ}\text{C}.$, but 30 mg. injected intravenously in the mouse did not prove lethal.

The bacterium, *Micrococcus* sp. G10 produced two types of substances one absorptive and one nonabsorptive on chromatographic partition. Both fluoresced blue and were dialyzable. Pyrexia in rabbits was produced by 15 mg. per kg. of the absorptive portion and by 10 mg. per kg. of the nonabsorptive. Thirty mg. was not a lethal dose for mice.

Phenol extraction of *S. typhi* and *S. paratyphi* cultures, water precipitation, followed by dialysis and precipitation of supernatant fluid, produced yields of 70 to 100 mg. of pyrogenic

- 1 9 5 0 -

material from 5000 cc. S. typhi cultures and 16 to 200 mg. from S. paratyphi cultures. Both Molisch and Fehling tests were positive. Both preparations fluoresced blue. Fifty gammas per kg. of S. typhi pyrogen and 75 gammas of the paratyphi preparation produced febrile reactions in rabbits.

The fluorescence of pyrogens in ultraviolet light was applied to a method for detecting these substances in solution. Solutions of dextrose, fructose, invert sugar, CaCl_2 , 10% sodium citrate, Salso-brocanon solution and Glynogen, were subjected to U.V. light. Three per-cent of the tubes exhibited fluorescing phenomena; only eight of twenty such tubes were pyrexia-producing for the rabbit. When compared with negative solutions, only 4% of the latter produced fever. The pyrogen zone appears to be around 2650 and absorption spectra at this wave length remain consistent.

649. HEIDELBERGER, M., MACLEOD, C. M., MARKOWITZ, H. and ROE, A. S.

Improved methods for the preparation of the specific polysaccharides of pneumococcus

Neutralization of acidity in developing bacterial cultures should be avoided if one wishes to isolate certain cell constituents in their native states. The specific polysaccharides of Type I and V pneumococcus may be damaged by alkaline environmental conditions and on precipitation yield smaller amounts of antibody-carrying materials. Degradative processes may also be initiated by bicarbonate disodium phosphate.

650. HEILBRUNN, L. V., and WILSON, W. L.

Effect of bacterial polysaccharide on cell division

Science, 112:56-57, 1950

Recent experiments with a polysaccharide extracted from S. marcescens (Shear's polysaccharide) would tend to prove that it exerts a heparin-like action on cell division. Both polysaccharides have similar inhibiting influences upon fertilization, although such inhibition may be overcome by additions of sperm in excess amounts. Both act to prevent mitotic gelation which may be preliminary to cell division. The bacterial polysaccharide also resembles heparin in preventing coagulation of human blood but high concentrations are necessary. Its potency is about 1/300 that of heparin.

In one experiment outlined, Chaetopterus eggs were immersed in dilute solutions of bacterial polysaccharide (0.058%-0.125%) before and after fertilization. The polysaccharide enters egg cells more rapidly than heparin. A solution of Shear's P-25,

- 1 9 5 0 -

0.15%, reduced cleavage from 100% (in seawater) to 0.5%. Excess sperm additions reduced the degree of inhibition. The increase in protoplasmic viscosity before mitosis was also prevented. Use of the polysaccharide or dicumarol in cancer therapy is discussed.

651. HEILMEYER, L.

Angänge einer Chemotherapie neoplastischer Erkrankungen
(Chemotherapy of neoplastic diseases)

Schweiz. med. Wchnschr. 79:530-547, 1949 and Naturwissensch. 37:58-65, 1950

In reviewing the literature dealing with the effects of chemical cytostatics upon the cells of tumors, other active substances are considered. In addition to cell nucleus toxins (colchicine, tryptaflavine, the urethanes, mustard gases, stilbamidine, aminopterin and metallic salts) other active materials, including the polysaccharides and pyrogens, hormonal substances and radioactive preparations, are effective under certain conditions.

652. HEISTØ, H.

Pyrogene reaksjoner, med saerlig henblikk pa deres forebyggelse
(Pyrogenic reactions with special considerations of their prevention)

Tidsskr. norsk. laegerfor. 70:264-266, 1950

In parenteral therapy dangers of unpleasant reactions exist when pyrogens are present. Prevention may rest in careful production of material destined for injection, in elimination of pyretic substances after production and the cautious approach to initial therapeutic application.

653. HEYMAN, A., PATTERSON, J. L., JR. and NICHOLS, F. T., JR.

The effects of induced fever on cerebral functions in neurosyphilis

J. Clin. Invest. 29:1335-1341, 1950

Fever was induced in 27 patients through use of typhoid bacterial pyrogens, malaria parasites or induced hyperthermia. Cerebral function changes in dementia paralytica or asymptomatic neurosyphilis were assessed. Pyrogenic material averaged 1 billion organisms per cc. or 0.1 cc. each injection. Rectal temperatures ranged between 101.0 and 104.0 degrees. At the height of pyrexial response, cerebral blood flow, oxygen consumption and arteriovenous

- 1 9 5 0 -

oxygen differences and vascular resistance were determined. Paretic patients with the greatest amount of mental deterioration had the lowest oxygen concentration in this area; with fever, the optimum increase expected was approached. There was no correlation between the degree of fever and O_2 consumption. Mean arterial blood pressure fell during fever. Patients with asymptomatic neurosyphilis had normal oxygen consumption in the afebrile state which was not altered during fever. In dementia paralytica subnormal oxygen consumption was evident before fever (2.1 cc per 100 Gm. per minute); during fever a 24% increase was registered (2.6 cc.). It is believed that, in the normal individual, cerebral blood flow and oxygen consumption are likewise altered during pyrogen hyperthermia.

654. HILL, A. G. S., DEANE, H. W. and COONS, A. H.

Localization of antigen in tissue cells. V. Capsular polysaccharide of Friedlander's Bacillus, type B. in the mouse

J. Exper. Med. 92:35-44, 1950

The fate of a capsular polysaccharide (that of Friedlander bacillus, Type B) was studied by means of homologous antibody labelled with fluorescein. When injected intravenously into the mouse, this acid polysaccharide is rapidly taken up by phagocytes in body tissues. It was seen in collagenous fibers of all organs, in capillary endothelium, in the hepatic epithelium and bile, the juxtaglomerular segment of the distal renal tubule and in a few casts. It had penetrated epithelium in uterine glands, the hormone-secreting tissues of the ovary, the suprarenal cortex, and lymph nodes and lymphocytes of the spleen. Synovial membranes contained large amounts. "The polysaccharide was found in high concentration in pulmonary macrophages" after inhalation.

655. HOWE, E. E. and TISHLER, M.

U. S. Patent #2,532,403, Dec. 5, 1950

A method for removing pyrogens from aqueous solutions of the acid salts of streptomycin.

656. HOWIE, J. E.

Pyrexial treatment of ulcerative colitis

The Lancet 25:827-828 (#6642), 1950

The author presents a brief case report of a patient suffering from ulcerative colitis for two years, who, after a course of

twelve intravenous injections of pyrifer, has been free from any clinical signs of chronic colitis despite two attacks of dysentery. In other similar cases it is said that results were somewhat complicated by the fact that this preparation produced less fever; wartime processes may have been responsible.

657. JACOBS, F. A.

Damage produced by Pseudomonas aeruginosa fraction in sarcoma 37

Canad. Res. 10:227, 1950

Cultures of Ps. aeruginosa R, grown on semi-synthetic media containing glycerol and fatty acids, contained a fraction capable of inducing damage to Sarcoma 37. Intraperitoneal administration of fractions from either culture fluid or cell pack of the organism produced macroscopic hemorrhage and succeeding necrosis of tumor tissue in mice. The tumor damage was confirmed microscopically. Fractionation of the culture fluid showed that the activity was resident in a water-soluble, alcohol-insoluble polysaccharide. The method of measuring tumor response to treatment is described.

658. KAPLAN, M. H., COONS, A. H. and DEANE, H. W.

Localization of antigen in tissue cells. III. Cellular distribution of pneumococcal polysaccharides, Types II and III in the mouse

J. Exper. Med. 91:15-30, 1950

Distribution of pneumococcal polysaccharides in the tissues of the mouse was followed histochemically by means of the fluorescein-antibody technique (Coons, et al.) or the periodic acid method (Schiff; McManus). Preparation of tissues was spaced at intervals ranging from 1 day to 6 months after a single dose of the polysaccharide under consideration.

Polysaccharides were found in most constant and striking concentrations in the cells of the reticuloendothelial system, the capillary endothelium and in fibroblasts throughout the body. They were detectable in monocytes and lymphocytes, hepatic, cardiac and smooth muscle cells, in uterine epithelium, the steroid-forming cells of the adrenal cortex, in testes and ovaries. They were persistent for 75 days in macrophages of the lymphoid organs, the Kupffer cells of the liver, interstitial macrophages of the myocardium, in lung septal cells, capillary endothelium and renal glomeruli. After six months, they were still present

- 1 9 5 0 -

in macrophages of the spleen, liver and heart and in peritubular capillary endothelium of the kidneys.

The smallest dose of polysaccharide to produce a positive precipitin reaction in serum and detectable quantities in cells in 24 hours was 0.03 mg. A comparison of the distributions of polysaccharide, acid dyes and suspensoids is made. Possible relationships between distribution, antigenicity and toxicity may exist.

659. KASS, E. H., and FINLAND, M.

Effect of ACTH on induced fever

New England J. Med. 243:693-695, 1950

Varying numbers of killed typhoid bacilli were injected intravenously in two patients with chronic rheumatoid arthritis, prior to or after administration of ACTH. The duration and intensity of the febrile response was reduced by administration of ACTH in advance of the pyrogen. That the antipyretic effect of ACTH is not to be credited to tolerance developed toward pyrogens has been confirmed in the rabbit. The mechanism by which ACTH exerts the antipyrexial effect is not known. There is no alteration of the fundamental pathologic process of the illness under treatment. It is necessary to differentiate between antipyretic and antipathologic results after the use of ACTH.

660. KIERLAND, R. R. and KULWIN, M. H.

Clinical evaluation of a new pyrogenic agent

A.M.A. Arch. Dermat. & Syph. 62:571-572, 1950

The pyrogenic agent, PYROMEN, was used clinically in 46 dermatologic patients suffering from a number of diseases. The usual initial dose was 25 micrograms given intravenously; subsequent dosages were based on the production of oral temperatures of 102.0 degrees F. The drug was administered on alternate days. The largest amount given in a single injection was 125 micrograms; the greatest number of injections given to one patient was ten.

Results showed PYROMEN produced a satisfactory "therapeutic" fever. Individual duration of temperature elevation was fairly "constant and...independent of the dose of the material injected." Untoward side reactions were few and mild. The authors conclude that PYROMEN "is a safe agent for use in producing therapeutic fever" since the results compared favorably with those obtained by the use of typhoid vaccine and lacked certain unpleasant side effects noted with the latter preparation.

661. KIRKENDALL, W. M., HODGES, R. E. and JANUARY, L. E.

The ACTH-like effect of fever in man

J. Lab. Clin. Med. 36:845-846, 1950

A marked drop in circulating eosinophiles regularly occurs four to eight hours after injection of a pyrogenic agent in patients with malignant hypertensive conditions. The phenomenon has been reported in humans after typhoid vaccine injections and in animals after induced fever. Since eosinopenia occurs also after administration of ACTH, the hypothesis that fever-producing agents may stimulate production of ACTH was tested. PYROMEN, typhoid vaccine and physically-induced fever were employed for 27 human patients, all with normally functioning pituitary and adrenal glands, but with hypertensive disorders. The circulating eosinophil and total leukocyte counts, and the ratio of multi-nucleated cells to mononuclear types and of uric acid to creatinine were determined; excretion of corticosteroids was measured in five patients.

In response to fever an initial leucopenia occurs; the decrease is mainly one of mononuclear cells. Eosinopenia follows fever in three to eight hours. Occasionally chills precede the marked leukocytosis (increased polymorphonuclear elements) which follows in 12 to 24 hours after injection. The administration of therapeutic doses of an antihistamine, of aspirin or dibenamine did not prevent such changes. No changes in the uric acid/creatinine ratio or in corticosteroid excretion followed fever produced by the three agents employed. It was believed that these would be altered if the leukocyte response reflected release of the adrenocorticotrophic hormone. However, the short period of stress occasioned by the administration of pyrogenic materials may be only long enough to alter the blood picture alone. It is concluded that "in persons with normal pituitary glands, fever causes the release of small amounts of ACTH."

662. LEESE, C. E., POEL, W. E. and BERMAN, H.

Effects of the Shear bacterial polysaccharides upon cardiovascular response

Fed. Proc. 9:76, 1950

When 100 to 1200 gammas per kg. body weight of the Shear polysaccharide was administered to the rabbit, the circulation time increased. A depressive action was seen with blood pressure falling to one-third to one-half the normal levels; a slow rise began within five to ten minutes. Once the blood pressure regained the normal level, subsequent injections had no influence. The depressive effects of the polysaccharide was not prevented by an antihistamine. The EKG was indicative of tachycardia with inversion of both P and T waves; there was an abnormally high S-T takeoff which disappeared within six hours.

- 1 9 5 0 -

663. MENKIN, V.

Further studies on thermostabile leukocytosis factor of exudates
Federation Proc. 9:338, 1950

A thermostable leukocytosis factor may be isolated from exudates which contain the pyrogenic (thermolabile) factor. The thermostable factor does not produce fever or leukopenia but induces leukocytosis through stimulation of the discharge of young granulocytes into the circulation. It is not a polypeptide but contains at least one carbohydrate group. It is believed that the thermolabile leukocytosis factor may be a polypeptide.

664. MUNK, F.

Zur Fieberbehandlung der akuten Poliomyelitis (Pyretotherapy of acute poliomyelitis)

Therap. Gegenwart 89:274-275, 1950

The author examines the results obtained in poliomyelitis through the use of temperature-elevating substances. In his own services, vaccine, pyrifur and nucleohexyl have been utilized, as reported by R. Muller and H. Arens (Therap. Gegenw. 1947-1948).

665. NESSET, N. M., MCLALLEN, J., ANTHONY, P. Z. and GINGER, L. G.

Bacterial pyrogens. I. A pyrogenic preparation from a Pseudomonas species

J. Am. Pharm. A. (Sci. ed.) 39:456-459, 1950

A relatively non-toxic bacterial pyrogen has been obtained from a culture of a Pseudomonas organism. After growth over a period ranging from one to ten days, cellular materials were digested by trypsin, centrifuged, washed and lyophilized. Digestion by the tryptic method employed was compared with proteolysis by papain, pepsin, polidase-S, pancreatin, a fish enzyme, duodenal powder, ictozyme, protease and an intestinal preparation; none yielded satisfactory results. Nine enzyme preparations with diastatic activity failed to produce detectable decreases in the pyrogenicity of cells. Other solvents and fractionation methods were studied.

Approximately 50 per cent of the concentrate obtained was pyrogenic polysaccharide. Seven per cent was hexosamine and 27 per cent desoxyribonucleic acid. Ribonucleic acid was present (8.1%). Protein content was negligible. The ash constituted

12.9% of the concentrate. For the rabbit the MPD was 0.3 gammas per kilogram of body weight. The LD₅₀ in the mouse was 75,000 gammas per kilogram of body weight.

666. NEVA, F. A. and MORGAN, H. R.

Tolerance to the action of endotoxins of enteric bacilli in patients convalescent from typhoid and paratyphoid fevers

J. Lab. Clin. Med. 35:911-922, 1950

Initial dosages of 5.0 micrograms of typhoid antigen (or alternate doses of S. dysenteriae antigen and typhoid antigen) were administered to convalescents recovering from typhoid or paratyphoid A fevers. The antigens had been prepared by the method of Morgan. When compared with healthy subjects who received the same amounts of the two antigens, a definite tolerance or resistance to the pyrogenic activity of toxic somatic antigen was demonstrated after convalescence; this was unrelated to the amount of circulating antibody. No tolerance was exhibited toward the toxins following recovery from other infections (bacillary dysentery, gonococcal arthritis, pneumococcal pneumonia and tularemia). Other investigators have advanced the theory that increased activity in the reticuloendothelial system may underlie the mechanism by which tolerance develops.

667. New light on endotoxins

Editorial Note. The Lancet I:722, 1950

The somatic antigen or endotoxin which may be extracted from members of the colon-typhoid-dysentery group (possibly excepting S. shigae) is attributable to a polysaccharide and nitrogen complex. It produces circulating antibodies, a terminal circulatory hypertension, hyperthermia, transient hyperglycaemia and succeeding hypoglycaemia. Its hemorrhagic action on the placenta is followed by abortion; similar activity occurs in transplanted tumors. Adrenal cortical extract appears to prevent certain toxic effects, which are more lethal for the guinea pig than for mice; the ratio was 17.5 to 1 when guinea pigs were injected subcutaneously with a suspension of heat-killed B. coli organisms. Mice received intravenous doses. The gross appearance of lesions suggests that capillary hemorrhages are produced by a terminal hypertensive action. Antibodies in serum from a mouse immunized with endotoxin were 30 times more potent than commercial B. coli antiserum in protecting others of the species against B. coli infection.

- 1 9 5 0 -

668. NOOJIN, R. O., PACE, B. F. and PRAYTOR, H. B.

Prolonged fever produced by three injections of typhoid vaccine

Am. J. Syphilis 34:153-160, 1950

The advantages inherent in typhoid vaccine fever therapy are: (1) easy administration, (2) controllability, (3) development of natural resistance, (4) simplicity, (5) inexpensiveness, (6) no after-treatment as with malarial therapy and (7) minimum side effect. Some of the disadvantages are: (1) hospitalization is required, (2) timing of fever is needed, and (3) some hypersensitivity may be exhibited. If the rise in fever is considered as proportional to the benefits derived, 20 million organisms should be injected intravenously, followed by subsequent injections at two-hour intervals for the first two doses, then three times per day for the duration of the course of treatment.

669. PIORKOWSKI, G.

Wirkungssteigerung fiebererzeugender Mittel bei gleichzeitiger intravenöser und intracutaner Anwendung (Increased efficacy of fever-producing drugs by using combined intravenous and intracutaneous application)

Med. Klin. 45:630-631, 1950

670. RALL, D. P., and WELLS, J. A.

Inhibition of thermal panting by bacterial pyrogen

J. Pharmacol. & Exper. Therap. 98:27, 1950

Failure of the dog to pant in response to pyrogenic fever is interpreted as being, in part, due to inhibition of the thermal panting mechanism. Chronic cord transected dogs responded to environmental temperature elevation with fever (103.5-105°F.) and panting. The latter was inhibited by introduction into a running venoclysis system of purified pyrogens from Pr. vulgaris or Ps. aeruginosa (PYROMEN). The degree of inhibition closely followed increases in dosage, and respiratory rate decreased sharply after injection of 0.04 to 1.2 mg. pyrogen; average reduction in 15 to 30 minutes ranged between 15 and 53 per cent.

671. RANDOLPH, T. G. and TOLLINS, J. P.

Pyromen in the treatment of perennial allergic symptoms

Ann. Allergy 8:626-640, 1950

A group of 150 patients has been treated with PYROMEN, intravenously, orally or by a combination of the two routes, over periods ranging from 10 days to 8 months. Subpyrexial doses of 0.5 to 10.0 gammas have been employed. The frequency of administration has been dependent on the allergic manifestation treated and the degree of control obtained. PYROMEN therapy has been found to be most efficacious in treatment of those patients for whom partial or complete specific diagnosis had been made; a combination of PYROMEN therapy with inhalant desensitization has resulted in excellent management of perennial allergic symptoms. The authors believe that beneficial effects obtained during this study may be the result of stimulation of the pituitary-adrenal system. Side effects with this therapeutic regimen have been infrequent and mild in nature.

672. REINHARD, E. H., GOOD, J. T. and MARTIN, E.

Chemotherapy of malignant neoplastic diseases

J.A.M.A. 142:383-390, 1950

In a lengthy discussion of chemotherapeutic agents used in the treatment of neoplasms, the authors discuss the potency of S. marcescens polysaccharide, the reticuloendothelial cytotoxic sera and other biologic antineoplastic agents. Toxicity is the limiting factor in application of the polysaccharide of S. marcescens.

673. RODBARD, S., SACI, H. and MALIN, A.

Body fluid redistribution in induced hypothermia and hyperthermia

Federation Proc. 9:107, 1950

Actual fluid volume and fluid shifts in hyperthermia and hypothermia were determined in both rabbits and chicks. The data suggest that there is a marked shift from plasma and interstitial spaces to the intracellular phase during hypothermia, but hematocrit and specific gravity do not show equivalent increases. It is suggested that both blood cellular elements and plasma proteins may be stored. This phenomenon is reversed during hyperthermia as water leaves the cells of tissue and releases plasma protein and red cells into the circulating blood.

- 1 9 5 0 -

674. RODNEY, G., and DEVLIN, H. B.

Chemical composition of some bacterial pyrogens

Federation Proc. 9:220, 1950

Pyrogens in a purified form were isolated from strains of B. prodigiosus, B. subtilis, Ps. aeruginosa and E. typhi grown on synthetic media. After digestion, nucleic acid and pyrogens were removed, and the latter was fractionated by means of organic solvents. Carbohydrates present in hydrolysates were quantitatively separated by filter paper partition chromatography. Analyses for nitrogenous materials and phosphorus were made. Pyrogen from different sources differed in composition both as to quality and quantity; activity also differed. The most active pyrogen was obtained from B. prodigiosus which initiated reactions in rabbits in concentrations of 0.01-0.005 micrograms per kg. body weight.

675. SCHMENGLER, R. D.

Effect of fever shock therapy in acute rheumatic and allergic diseases. Its significance for acute rheumatism (Translated Title)

Klin. Wchnschr. (Heidelberg) 27:1027, 1950

A vaccine produced from the colon bacillus (pyrifer) was injected in varying amounts into patients who presented the following pathologic entities: (1) acute rheumatic polyarthrititis; (2) acute rheumatic fever without joint involvement; (3) poly-symptomatic rheumatic conditions, often with cutaneous lesions; (4) acute allergic disorders with known antigens.

Following therapy prompt defervescence occurred and the local symptoms disappeared. A sympathetic reaction is postulated; a cellular anaphylaxis may set in motion a sympathetic response ending in pathologic processes. This latter response is reversed by pyrifer.

676. SCHOLTZ, H. G.

Physikalische Hyperthermie oder Pyriferfieber? (Physical hyperthermia or fever induced by pyrifer?)

Arch. phys. Therap. 2:53-56, 1950

677. SMITH, J. McC. and MIRICK, G. S.

The inhibitory effect of bacterial suspensions on pneumonia

- 1950 -

virus of mice

Bact. Proc. (1950) 69

Gram-negative bacillary cells had an inhibitory effect on the inflammatory and lethal activity of the pneumonia virus when applied intranasally to mice, before infection or two days after. Species employed included *S. typhosa* and *paratyphi*, A, B and C, *S. typhimurium*, *E. coli*, *S. dysenteriae* (Shiga and Flexner) and *Proteus* OX19. Preparations given intraperitoneally exhibited no activity. Filtrates of broth were inactive. The agent was not a polysaccharide, since a trypsin digest was likewise ineffective. Activity of cells was not affected by autoclaving, phenol or acetone extraction. A similarity to the mucinase from *V. comma* described by Burnet is discussed.

678. STUART, E. G.

Alterations in connective tissue mast cells induced by bacterial pyrogens

Am. J. Physiol. 163:754, 1950

In connective tissue mast cells in abdominal skin of the mouse, a concentration of granules around a central nucleus may be seen with a few cells exhibiting some dispersion of granules when cell boundaries are indefinite. After peritoneal and oral PYROMEN in dosages ranging between 5 and 50 gammas per kg. body weight, skin biopsies were taken at intervals varying from 15 minutes to 30 days after injection. A response to PYROMEN was elicited which reached its peak in seven to ten days after a single (5 gammas per kg.) dose. "Numerically, morphologically and spatially it followed a definite pattern dependent on the dosage, elapsed time and strain of experimental mice employed" Granules of many mast cells dispersed, the dispersion beginning in the vicinity of small venules and capillaries between 5 to 15 minutes after administration, according to the dosage given. Groups of mast cells some distance from blood vessels were less rapidly changed; there was an alteration in shape away from the ovoid; granules spread out from the cell center. Many of the pericapillary cells assumed a spindle shape.

679. STUART, E. G.

The mast cell response to administration of a bacterial pyrogen

J. Nat. Cancer Inst. 10:1375-1376, 1950

After 5 to 50 gammas of PYROMEN per kg. body weight had been administered to mice by intraperitoneal injection or the oral route, skin biopsies were taken at intervals between 15 minutes and 24

- 1 9 5 0 -

hours. Dispersion of granules of mast cells began between 2 and 15 minutes after administration of PYROMEN with most rapid alterations evident in areas near small blood vessels.

680. STUTTGEN, G.

Pyrifer als antigen im rahmen der bakterzellen autoallergie (Pyrifer as an antigen in the picture of autoallergy of the bacterial cell.)

Zeitschr. Haut u. Geschlechtskr. 9:17-20, 1950

681. SWINEFORD, O., JR.

Observations on the use of bacterial antigens in the treatment of asthma. A brief critical review

Am. Pract. 1:612-618, 1950

The author presents summarized data from twenty-eight authors who have reported on the treatment of asthma with bacterial antigens. Characteristics, clinical and laboratory, of atopic and tuberculin allergies and of the Arthus reaction are enumerated. The treatment by such means is empirical but of definite value.

682. TAKOS, M. J. and MOE, G. K.

Prevention of pyrogen-induced renal hyperemia in the dog by dihydroergocornine

Proc. Soc. Exper. Biol. & Med. 75:51-52, 1950

After control periods in which glomerular filtration rate and the effective renal plasma flow were determined, dihydroergocornine (DHO) was administered to female dogs (0.01 mg. per kg. intravenously and a like amount subcutaneously) and followed by a pyrogen derived from Shigella organism. Rectal temperatures were recorded. Comparisons were made with controls receiving pyrogens alone, and all experiments were repeated.

In animals treated with pyrogen alone, the renal plasma flow increased 27 to 79 per cent while the glomerular filtration rate remained within normal limits. DHO is an alkaloid known to interfere with certain central autonomic mechanisms. Premedication with this drug brought a great reduction in renal hyperemic response, with a range of -4 to plus 25%. It did not affect the glomerular filtration rate or inhibit fever. The antagonism between pyrogen and DHO is presumed to indicate a blocking action on the central nervous system by the alkaloid.

683. TAL, C. and GOEBEL, W. F.

- 1950 -

On the nature of the toxic component of the somatic antigen of Shigella paradysenteriae type 2 (Flaxner)

J. Exper. Med. 92:25-34, 1950

The toxic component of the somatic antigen of dysentery bacilli appears to be neither the lipid, protein or carbohydrate of the complete antigen. Fractions disassociated from the complete antigen yielded a protein component nearly as toxic as the original antigen; carbohydrates were still bound to the protein complex. A milder procedure yielded a product almost protein free; further treatment eliminated 50% of the total lipids but did not change serologic, antigenic or toxic properties. The U.V. absorption was approximately 2600⁰ for one component, primarily polysaccharide in nature. Further hydrolysis yielded a polysaccharide, serologically active but non-toxic, and an insoluble precipitate, not identifiable as protein or lipid.

684. WAGNER, R. R., and BENNETT, I. L.

The production of fever by influenzal viruses

J. Exper. Med. 91:135-145, 1950

Toxic bacterial filtrates which have an affinity for and destructive effect upon the cell receptors for influenzal viruses diminished the febrile response to PR8 and Lee strains of influenza virus and that of Newcastle disease. Both Cholera vibrio and Clostridium welchii filtrates had antipyrexial influence, but near lethal amounts were necessary to demonstrate a protective effect. However, the ability of animals to respond to the pyrogenic effect of typhoid vaccine was unimpaired by injection of receptor destroyers. "These findings give support to the hypothesis that union of virus and host receptor substance plays a part in the production of fever by these viruses." Chemical substances (ie; sodium periodate) may be neutralized by normal serum and allantoic fluid, since the cell receptor activity was not blocked by their use; glucose is also believed to neutralize the action of sodium periodate.

685. WALKER, L., OLSON, W. H. and NECHELES, H.

Depression of gastric secretion with pyrogens and antipyretics without fever

Am. J. Physiol. 163:758, 1950 (Abstract)

Increases in the temperature of the body may cause a partial or complete inhibition of hydrochloric acid secretion by the

- 1950 -

stomach. Injection of pyrogens can produce temperature elevations with inhibition of gastric secretion and, may be capable of depressing secretory ability when subpyrexial quantities are administered. After histamine stimulation, pyrogens in subpyretic dosages were injected into dogs with total gastric pouches. The inhibitory effect was not constant. When larger pyrogen increments were utilized, consistent inhibition of secretion was obtained even after the temperature elevation was abolished by antipyretics. Sixty per cent reduction in total volume of gastric juice and 72 per cent reduction in total mEq. of free HCl accompanied this alteration, produced by injection of 50 gammas of PYROMEN, and 10 grains of aspirin given orally. A latent period of 30 to 60 minutes preceded these findings.

686. WAUER, D.

Innere Abteilung des Stadtischen Behring-krankenhauses Berlin-Zehlendorf. Zur Pyrifertherapie akuter und chronischer Leberparenchymschaden (Treatment of acute and chronic involvements of the liver parenchyma with pyrifer)

Deutsch. Gesundheitsw. 5:489-492, 1950

Nonspecific desensitization with pyrifer as anti-allergic agent was beneficial in both acute and chronic hepatic parenchymatous conditions. Allergy may be etiological in many hepatopathies, and pyrifer activity is considered one producing a change in capillary permeability.

687. WINDLE, W. F.

Changes in the hypophysis and suprarenal glands induced by a bacterial pyrogen

Anat. Rec. 106:94-95, 1950

Histologic studies were made of the organs of 32 virgin female rabbits previously treated with a new purified bacterial pyrogen, P.V. PYROMEN, and these were compared with organs from six controls. The average dose of PYROMEN was either 1 or 5 gammas per kg. body weight. Six animals received 0.01 to 0.1 gammas per kg.; six others received very high dosages (500 to 20,000 gammas per kg.). The LD₅₀ is 80,000 gammas per kg. Tissues were prepared with perfusion formalin fixation and hematoxylin-eosin staining procedures.

PYROMEN appeared to act as a mild stress stimulus, but no animal exhibited lesions resembling those seen in exhaustion stages of the general adaptation syndrome. Changes in endocrine organs

were not related to dosage levels. Within 12 to 24 hours a panendocrine response was manifested with vacuolation of cell cytoplasm in the suprarenal medulla, hypophyseal basophil and chromophobe cells, all zones of the suprarenal cortex, the parathyroid, islets of the pancreas and ovarian stroma cells. Leucopenia and eosinopenia appeared within one hour and reached their peak levels in four hours. Lymphocytosis followed. In contrast to the action of ACTH, no suprarenal hypertrophic changes were seen; in the mouse mast cells show marked granulocytic dispersion after PYROMEN. Prolonged administration of PYROMEN stimulated growth of the suprarenal cortex, with increased width of the reticularis but decreases in other zones. Capsule cells appeared to be transformed to glomerulosa cells. Prolonged high doses led to a reduction of anterior lobe basophil population associated with involutional changes.

688. WINDLE, W. F.

Endocrine and connective tissue cellular responses to administration of a bacterial pyrogen

Abst. of Communications, XVIII Inter. Physiol. Congress (Copenhagen) Aug. 15-18, 1950, p. 518

Events following PYROMEN administration were studied in various species. One to 5 gammas per kg. induced leucopenia and eosinopenia within one hour in the rat, cat, dog and human. The low point was reached in 4 hours, and lymphocytosis followed. Increased output of urinary corticoids was not detected in rabbits and cats, but pronounced temporary increases in dogs and man have been reported by others. In the mouse, connective tissue mast cell granular dispersion was noted within one hour. A panendocrine reaction to PYROMEN was seen in all organs of rabbits. These changes began in 12 to 24 hours, and maximal effects were present for periods up to 10 days; vacuolation of cell cytoplasm in adrenal medulla and cortex, pituitary chromophobes and basophils and parathyroid and pancreatic cells were seen. Interstitial cells of the ovarian stroma became hypertrophied within 24 hours, with vacuolation. Prolonged administration of PYROMEN created hyperactivity of lymphatic organs, inducing hyperplasia and even metaphasia with granulocyte production in the spleen. In all organs there was increased vascularity. Proliferation of certain connective tissue cells followed. After spinal cord transection in dogs and cats, fibrous and glial scars appear to be inhibited while the cellular tissues developing between cut surfaces permit the penetration of regenerating central neurons.

689. WINDLE, W. F. and CHAMBERS, W. W.

Regeneration in the spinal cord of the cat and dog

- 1 9 5 0 -

J. Compar. Neurol. 93:241, 1950

In a series of experiments to determine the site of action of bacterial pyrogens, spinal cords of cats and dogs were transected at various levels. One group was used for control purposes. The rest received PYROMEN intravenously or intraperitoneally in cumulative dosages ranging from 300 to 11,000 gammas. In addition one control received 150 units of hyalase for 10 doses; one animal received 225 gammas of typhoid concentrate and 35 mg. "Pyrexin", and a second was given 3,856 gammas of typhoid concentrate alone. One animal had received old and inactive pyrogen.

The animals were sacrificed at intervals ranging from 8 to 139 days after administration of PYROMEN. Cords of animals living 30 to 50 days after transection revealed new growth of neurons across or into cut regions. There was some connective tissue scarring at severed ends, but no glial barrier. Rather a blending of spinal cord parenchyma with a loose tissue containing non-nervous tissue cells had occurred. Through such regions nerve fibers had in some cases penetrated into or through the scar along strands of reticulum-like cells. Serial sections and staining by the pyridine-silver method revealed that they were derived from severed or injured nerve roots, nerves to blood vessels and intrinsic spinal cord neurons. No evidences of functional regeneration were seen.

690. WINDLE, W. F.

Spinal cord regeneration associated with a cellular reaction induced by administration of a purified bacterial pyrogen

International Anatomical Congress, July 24-28, 1950. Abst., Proc. of the Congress, p. 196.

Transected spinal cords of dogs and cats, when given varying dosages of PYROMEN for periods of time ranging from 8 to 61 days, revealed clear-cut evidence of neuron growth into or across the cut ends. Some connective tissue scarring between severed ends and the usual amount of retrograde degeneration were observed. There was no glial barrier, but a blending of spinal cord parenchyma with connective tissue was noted; through such areas, nerve fibers could be seen growing into the scar. The nerve fibers were derived from severed or injured posterior and anterior nerve roots, vascular and meningeal nerves, and intrinsic spinal cord neurons. Numerous blood vessels were observed at scar-cord juncture and there were many macrophages at the interstices between strands of non-nerve cells and the nerve fibers.

- 1 9 5 0 -

691. WINDLE, W. F., CHAMBERS, W. W., RICKER, W. A., GINGER, L. G. and KOENIG, H.

Reactions of tissues to administration of pyrogenic preparation from a Pseudomonas species

Am. J. Med. Sci. 219:422-426, 1950

Sensitivity to PYROMEN was shown in test animals in the following descending order: dog, rabbit, cat, monkey, guinea pig and rat. Administration of massive doses induced histological changes in lymphoid and myeloid organs similar to those observed with other pyrogenic substances or pyrogen-free saline solution. There was increased macrophage and lymphocyte production in all species studied; where treatment was prolonged, myelocyte and granulocyte production in the spleen of the rabbit was noted.

692. WINDLE, W. F. and WILCOX, H. H.

Extramedullary hemopoiesis in rabbit and cat induced by bacterial pyrogens

Am. J. Physiol. 163:762, 1950

Histological changes in the spleen after PYROMEN are related to time of treatment and to some degree to the size of the dose. Most marked changes were observed in the red pulp, where proliferation of reticular cells and reduction of blood flow through sinuses were noted during the first 20 days of treatment; where treatment had been continued a month or more, phagocytosis of static blood by macrophages was noted. Groups of hemopoietic cells appeared in pulp cords. 'Islands' of hemopoietic tissue, similar to those found in embryonic states, were seen in sinuses of the spleen. When the dosage was massive or therapy prolonged the hematopoietic picture resembled, for the spleen, that of bone marrow. The greatest increases in spleen weight occurred in the cat, but hepatic hemopoiesis was noted in the rabbit after massive or prolonged therapy. Hepatic activity was not noted in controls nor with lower dosage levels of PYROMEN.

693. WINDLE, W. F., WILCOX, RHINES, R., and CLEMENTE, C.

Changes in endocrine organs induced by bacterial pyrogens

Federation Proc. 9:137, 1950

The organs of rabbits were examined histologically for signs of stress-linked changes after administration of Ps. PYROMEN

- 1 9 5 0 -

(100,000-200,000 gammas total dosage). Treatment extended from 19 to 303 days. Comparisons were made with the appearance of organs after 500-20,000 gammas per kg. body weight of P.V. PYROMEN had been administered for 3 to 6 weeks.

Neither group had organs typical of the exhaustion stage associated with the general adaptation syndrome. Ps. PYROMEN-treated rabbits exhibited hyperplasia, hyperactivity of lymphatic and reticuloendothelial tissues, while P.V. PYROMEN induced but slight changes in these tissues. Both preparations produced blood component alterations resembling those seen after stress or following the administration of ACTH. Changes suggestive of involution after marked growth stimulation of the adrenal cortex were noted.

Rabbits and rats given P.V. PYROMEN thrice weekly (to 122 days) in dosages of 1 or 5 gammas per kg. body weight exhibited increased cytoplasmic activity in endocrines. Vacuolization of anterior lobe basophils and cells of the pars intermedia of the hypophysis appeared within 12 hours. Similar changes appeared in adrenal cortex after prolonged administration; at higher dosage levels involutional changes were seen.

694. WIRTH, F.

Über Pyriferverhandlung bei schwerer Diphtherie (Fever therapy (pyrifer) in severe diphtheria)

Deutsch. med. Rsch. 4:204-206, 1950

Seventy-one of 133 patients treated with antitoxin and penicillin were given a bacterial pyrexial agent, with the result that recovery was more rapid and less difficult. Mortality appeared to be reduced by adjuvant therapy. Neurological complications were unchanged.

695. ALGIRE, G. H. and LEGALLAIS, F. Y.

Vascular reactions in normal and malignant tissues in vivo. IV. The effect of peripheral hypotension on transplanted tumors

J. Nat. Cancer Inst. 12:399-421, 1951

The transparent chamber technique was used in making microscopic observations of changes in the blood supply of transplanted tumors when peripheral blood pressure was reduced. Direct correlation between hypotension and reduction of circulation in capillaries of tumors and surrounding tissues in the host was secured. Reduction of tumor circulation was related to the duration and degree of peripheral hypotension. Necrotic changes in tumor tissue follow in prolonged reductions. A polysaccharide from Serratia marcescens furnishes a mechanism which may be related to this phenomenon: progressively increasing sluggishness of flow in the arteries; stasis of blood; occlusion of capillaries in striated muscle; necrosis in sarcoma tissue. In the latter instance, hemorrhage follows ischemia within a few hours and tumor damage may be apparent in 24 hrs. thereafter. Work is now in progress to determine whether hypotensive effects are responsible for the therapeutic benefits achieved.

696. ARCHER, B. J.

Remission in rheumatoid arthritis following fever therapy with liver damage

New York J. Med. 51:2657-2658, 1951

Cephalin flocculation tests became positive soon after initiation of therapy with injections of typhoid bacteria in a patient resistant to the effects of gold salts and salicylates. The elevated sedimentation rate returned to normal and remission of arthritic symptoms followed. The cephalin index returned to normal at the time of relapse, two weeks later.

697. BECK, L. V.

Reducing substances as indicators of protection afforded by cortisone against tumor-damaging effects of Serratia marcescens polysaccharide

Cancer Res. 11:4, 1951

Marked decrease in the content of reducing substances - ascorbic acid and glutathione - accompanied hemorrhagic necrosis induced in mouse sarcoma #37 by arsenite, a-peltatin and P25, the polysaccharide from S. marcescens. When 5 micrograms of cortisone per Gm. mouse weight were injected before P25, the minimal dose of the polysaccharide required for production of tumor necrosis

- 1 9 5 1 -

was doubled. It was shown that the minimum dose of P25 required to decrease ascorbic acid and glutathione content of tumors is twice that required when preceded by injections of cortisone.

698. BRIL' M T, Knokhitkin

O Metodke pirogennoi terappii sifilisa (Method of Pyrogenous Syphilis Therapy)

Vestnik vener. dermat. 1951(3) 28-31

699. CHARONNAT, R. and LECHAT, P.

Investigaciones sobre el ensayo de los pirogenos en los soluciones inyectables (Investigations on the test for pyrogens in injectable solutions)

Laboratorio, 10:545-556, 1950

700. CHARONNAT, R. and LECHAT, P.

Researches sur la nature des pyrogenes des solutes injectables. I. Substances definies d'origine biologique ayanat une action pyrogene (Investigations into the nature of pyrogens in injectable solutions)

Ann. pharm. franc. 9:17-22, 1951

The pyrogen of Robinson and Flusser appeared to be a polysaccharide approximating 62,000 molecular weight, containing 0.29-2.38% P but without azotes. Menkin's "pyrexine" has 11% N, but 1% P, gives no Molisch reaction and is considered to be a glucopeptide.

In a series of experiments, pyrexial substances, both bacterial and chemical, were injected intravenously into the rabbit. From an opotherapeutic product, a substance was extracted which was, when purified, insoluble in water but soluble in monosodium carbon solution. 0.1 mg./kg., in the rabbit, elevated temperature from 1 to 4 degrees. Nucleic acids, thymonucleic acid, and a 7% nucleotide solution were also investigated. Adenylic acid was not pyrogenic. Adenosine (three types) had specific hyperthermic action, but it was believed that an impurity was responsible.

701. CHARONNAT, R. and LECHAT, P.

Researches sur la nature des pyrogenes des solutes injectables. II. Essais de definition quantitative des substances pyrogenes. III. Indice pyrogeniques (II. Tests of quantitative definition of pyrogenic substances. III. Pyrogenic indices)

Ann. pharm. franc. 9:22-30, 1951

- 1 9 5 1 -

II. After extended studies it could not be confirmed that the log of hyperthermia provoked by the pyrogen of B. pyocyaneus has the relationship of a linear function to the logarithm of the dose.

III. An empiric formula is presented which may be employed to correlate the hyperthermic curve and amounts of pyrogens injected. The law involved is one far more complex than that of a straight line function. The pyrogen unit is not an index.

702. CLEMENTE, C. D.

Production of periods of leukocytosis and tolerance by consecutive oral administrations of different bacterial pyrogens

Federation Proc. 10:27, 1951

Tolerance to oral administration of several bacterial pyrogens (from a Pseudomonas species, Pr. vulgaris and B. subtilis) followed leukocytosis. Normal white cell levels reappeared. To differentiate between a generalized effect of PYROMEN and possible existence of species differences, 1 gamma Ps. PYROMEN was fed in drinking water daily. In 2 to 5 days the total white count had ascended from an average of 16,000 leukocytes per cu. mm. to levels between 29,000 and 35,000. Normal levels reappeared in 7 to 10 days. When the pyrogen employed was one obtained from either Pr. vulgaris or B. subtilis, a secondary leukocytosis was evoked and a second period of tolerance followed. In some animals the period of white cell elevation extended to 20 days. The most prolonged results were achieved with Pr. vulgaris pyrogen.

703. CLEMENTE, C. D., CHAMBERS, W. W., GREENE, L. C., MITCHELL, S. Q. and WINDLE, W. F.

Regeneration of the transected spinal cord of adult cats

Anat. Rec. 109:280, 1951

Certain bacterial pyrogens assist in inhibiting collagenous scarring after complete spinal cord resection. Loose reticular and macrophage matrices are substituted for scar tissue. The penetration of intraspinal nerve fibers on regeneration is thereby rendered possible. Regenerative activity was demonstrated histologically when 20 gamma of PYROMEN per Kg. body weight were administered initially to spinal cats and the amounts increased (to 200 gamma per Kg. body weight) for successive periods of approximately two weeks each. Treatment alternated with rest periods. When the treatment was discontinued prematurely, growth of the dense collagenous tissue was resumed.

- 1 9 5 1 -

704. CLEMENTE, C. D., CHAMBERS, W. W. and WINDLE, W. F.

Blending of regenerating peripheral fibers with cerebral cortex through glial barrier inhibition

Am. J. Physiol. 167:774, 1951

The published observations of Windle and Chambers (1950) and Clemente and co-workers (1951) on the anatomical regeneration of severed spinal cords of adult cats have been augmented through a report of the results obtained when peripheral nerves were implanted into the cerebrum. The proximal stump of the temporal branch of the right facial nerve was implanted into the cerebral cortex of subject and control. One animal received a bacterial polysaccharide intravenously (PYROMEN) for a period of 30 days. Sections of cerebral tissue stained with silver were examined. No sharp demarcation between peripheral nerve and brain parenchyma could be seen in tissue following PYROMEN therapy. The regenerating fibers grew radially and blended with the substance of the cerebrum. In untreated tissue, distinct glial barriers were seen and no blending was observed.

705. CLOUGH, P. W.

Pyrogens.

Editorial, Ann. Int. Med. 34:816-819, 1951

The literature on the therapeutic applications of pyrogenic substances is partially reviewed. Present use of pyrogens is, according to the author, confined to attempts to stimulate certain defense mechanisms. "This is most evident in such diseases as uveitis and other ocular infections...it might be used more frequently than it is in certain other types of infection for which no specific therapy is available and which are resistant to less drastic procedures..." Application of pyrogens in the therapy of (a) inoperable neoplasms (b) malignant hypertension appears to be promising but is limited by severity of the reaction. In summarizing, the editor states that "this form of treatment is.. still in the early experimental stage".

The characteristics of bacterial pyrogens, clinical manifestations following their administration, and the increases in tolerance seen after continued pyrogen therapy are briefly discussed. Abolition of pyrogenic reactions has been secured through use of antipyretics, ACTH and cortisone, or reticuloendothelial blocking agents.

706. FARIA, R.

Post-transfusional reactions. A clinical and statistical study

-1 9 5 1 -

(Translated Title)

Seara médica 6:359, 1951

The reactions characterized as pyrogenic may be controlled after transfusion of blood by (1) discontinuance of transfusion, administration of anti-febrile drugs, sedatives (2) substitution of plasma and other equipment for transfusion. Calcium intravenously has been advocated as efficacious for these reactions as has novaldin in one gram injections intravenously. "Prophylaxis of pyrogenic reactions includes...care in the collection and administration of blood and a completely closed technic". Pyrogenic reactions may be actually pseudopyrogenic reactions; the role of chilling and the use of frozen preserved blood are considered. The author observed an incidence of 3.9 per cent unfavorable after-effects after 2.069 transfusions, and these included all reactions recorded.

707. FISHGOLD, J. T., GRANT, R., FIELD, J. and HALL, V. E.

Oxygen consumption and glucose exchange in vitro of liver slices from febrile rabbits

Am. J. Physiol. 166:113-120, 1951

Changes in liver function in fever may be caused by (a) heat production accelerating metabolic process or (b) disturbance of the metabolism of carbohydrate resulting in hyperglycemia and the typical diabetic syndrome.

When oxygen consumption was measured in liver slices taken from animals (rabbits) sacrificed at the onset of the chill, it was found that lower values were present after pyrogen (typhoid paratyphoid vaccine) than were seen in untreated animals. The hydrogen ion concentration was a factor also, since incubation of slices at pH 7.2 brought oxygen consumption closer to normal levels. No release of epinephrine could be measured at the onset of fever and, in the absence of epinephrine, five times as much glucose was released from tissue suspensions after fever as from normal tissue. Addition of epinephrine increased glucose values both in the febrile tissue preparations and in those taken from normal animals. The suggestion is made that early hyperglycemia in fever may be caused in part by a sudden release of carbohydrate from the liver through sympathetic-adrenal stimulation, and that heat production may be mediated through nervous pathways and not through direct peripheral action when pyrogens are administered.

708. FITCH, E. M. J. and WASHBURN, A. C.

The effects of PYROMEN administered intravenously in subpyrexial

- 1 9 5 1 -

doses to depressed patients. A preliminary study

New England J. Med. 245:961-968, 1951

Among the favorable results reported after PYROMEN therapy has been the appearance of increased euphoria. To determine whether or not this response might be therapeutically useful in the treatment of the mentally depressed patient, and whether or not it was independent of fever, subpyrexial doses of PYROMEN were administered to 62 individuals, 31 of whom reported extreme depression. Of these, four were termed schizoid personalities and four manic depressive. Fifteen patients presented some symptoms of allergic disease. The average dose of PYROMEN ranged from initial injections of 0.5 to 1.0 gamma, to 2.0 gammas two or three times weekly for an average period of 79 days. Overdosage appeared to bring an exacerbation of symptoms but it was rarely necessary to reduce the amount given. Among the symptoms observed were headache, dizziness, nausea and aching. A tendency toward increased bleeding was noted; menstruation was induced in 1 patient, intra-menstrual bleeding in two. Dysmenorrhea was relieved. Caution is indicated in the use of PYROMEN where a hemorrhagic lesion may exist. Nocturia and frequency of urination were present in the early stage of treatment; a loss of weight may have been related to loss of edema fluid. Symptoms indicative of vitamin balance disturbance, ie. metallic taste, cheilosis, glossitis, and bleeding gums, appeared in a few patients and were relieved by administration of vitamins B and C in high dosages.

On the basis of 537 pairs of differential counts it was concluded that hematologic response could not be used for evaluation. The eosinophil count was not valid enough to be used as a gauge of response. There were no significant changes in hemoglobin, erythrocyte levels, platelets, reticulocytes, total serum protein, prothrombin time, vitamin C., calcium, phosphorus or bilirubin. A placebo (saline solution) occasionally brought a drop in eosinophils.

Allergic states yielded to PYROMEN therapy (asthma, severe acne and chronic eczematoid dermatitis), but seborrheic dermatitis was not improved. Depression, fatigue, anxiety and insomnia disappeared. One of the schizophrenic subjects improved but symptoms were exaggerated in the passive variety of schizoid involvement. The drug appears to bring the psychiatric patient toward an acceptance of psychotherapeutic techniques. The theory is advanced that central nervous symptoms of a functional nature may be the results of allergic states with cerebral edema, or that a shift from the parasympathetic to the sympathetic system has changed both psychic and somatic behavior. A common agent, opposed by PYROMEN, may be involved in these syndromes.

709. GINGER, L. G., NESSET, N. M., RIEGEL, B. and FITZSIMONS, E. J.

Bacterial Pyrogens. II. Pyrogenic preparations from various bacterial species

J. Am. Pharm. A. (Sci. Ed.) 40:421-424, 1951

Pyrogenic concentrates, prepared from the cell bodies of five bacterial species by means of tryptic digestion, were characterized by essential chemical and biological differences. Cellular materials from E. coli, Pr. vulgaris, S. typhi and B. subtilis were grown and collected by a method previously described (J. Am. Pharm. A. 39:456, 1950). Both cellular material and concentrates were analyzed for the following constituents: ribonucleic acid, desoxyribonucleic acid, hexosamine, pyrogenic polysaccharide and free lipid. Total nitrogen, protein and nitrogenous residue were determined. The smallest content of pyrogenic polysaccharide (35.8%) was found in the concentrate from Pr. vulgaris; the largest content was found in the concentrate from B. subtilis. The polysaccharide content bore little relationship to biological activity; the most active concentrate was that from S. typhi. It contained the smallest amounts of hexosamine and unclassified reducing sugars. Proteus vulgaris concentrate, with low pyrogenic polysaccharide content possessed high biological activity. On the other hand, the concentrates prepared from B. subtilis contained 63.2% pyrogenic polysaccharide but had the lowest biological activity.

Differences in pyrogenic polysaccharide composition may account for the variations in activity. "The possibility that biological activity may be related to differences in molecular structure cannot be overlooked..."

710. GREENE, L. C.

Healing of thermal burns in cats treated with PYROMEN

Am. J. Physiol. 167:789, 1951

Burns, 3 and 1.6 cm. in diameter, were produced on either side of the vertebral column of the cat by the method outlined by Peters (1943-1944). In a series of 12 animals, intravenous injection of a purified bacterial polysaccharide (PYROMEN) were administered at several dosage levels. Four animals received initially 20 gammas per kg. per day steadily increasing to 60 gammas per kg. per day in one week. Maintenance doses were 10 gammas per day for 85 days. Two animals received 0.2 gamma three times a day. Treatment was continued until healing was complete.

Marked differences were noted between scab appearance in the treated animals and controls. For the former, scab formation was rapid,

- 1 9 5 1 -

and sloughing took place in a fragmentary fashion, leaving smooth surfaces, small scars and textures resembling those of normal skin. Sloughing and healing were completed 15 to 25 days earlier in the treated animal than in the untreated.

711. HALLMAN, N., LAGUS, K.

Pyrogeenireaktioista eri tavoin valmistetuilla injektiooliuksilla (Pyrogen reactions to solutions prepared by Finnish method as prescribed by U.S.P.)

Duodecim. 67:335-343, 1951

712. HIRSCHMANN, J.

Konstitutionelle Unterschiede von Temperatur und Leukocytenkurven unter Pyriferbelastung (Constitutional differences in temperature and elukocyte variations during pyrifer administration)

Nervenarzt 22:260-264, 1951

713. GUTHRIE, T. C.

Effects of artificially-induced fever on the electroencephalograms of patients with multiple sclerosis

Trans. New York Neurol. Soc. and New York Acad. Med. (Sect. on Neurology and Psychiat.) Meeting, May 8, 1951. A.M.A. Arch. Neurol. and Psychiat. 66:660-661, 1951 (Abstract)

Under the experimental conditions outlined, increased heat appears to affect multiple sclerosis adversely. When the temperature was increased from 0.8 to 5.4° F. through the injection of 20 million killed typhoid organisms, electroencephalograms recorded for five patients were indicative of unfavorable activity; there was decreased strength of grip, a loss in visual acuity and increases in dysarthria. The permanence of these changes is discussed and a further consideration of such therapy is suggested.

714. HAYNES, B. W., DEBAKEY, M. C. and DENMAN, F. R.

Renal function studies of severely burned patients

Ann. Surg., 134:617-625, 1951

During a study of renal function in eight severely burned patients,

who were receiving adequate hemotherapy and fluids, the renal plasma flow, glomerular filtrate rate and fraction excretion were determined. An analysis of both arterial and venous blood was added. There were changes in plasma protein concentration. The possibility that there is increased thyroxine production in burns was suggested, and certain similarities to results obtained through DOCA were noted. In fevers, the maintenance of glomerular filtration is accompanied by effective increases in blood plasma flow. Reduction of filtration factors has been reported. Pyrogen, without actual production of fever, increases the renal plasma flow but does not affect glomerular filtration. Filtration fractions are decreased.

715. KIRKENDALL, W. M., HODGES, R. E. and JANUARY, L. E.

The ACTH-like effect of fever in man

J. Lab. & Clin. Med. 37:771-779, 1951

In studies planned to determine whether or not eosinopenia observed after induced fever is caused by liberation of ACTH, elevation of temperature was induced in 27 afebrile patients, all with normally-functioning adrenal and pituitary glands. Fever was induced by either (1) 15 to 100 million typhoid organisms in vaccine (2) moist air fever cabinet or (3) an endogenous polysaccharide of Proteus vulgaris (PV PYROMEN). The last-named was administered intravenously in 25 to 100 gamma doses.

The effects of aspirin, dibenamine and chlor-trimeton before and after injection of Pv PYROMEN were observed in four patients. Determinations of uric acid/creatinine ratios, circulating leukocytes, corticosteroid excretion and rectal temperatures were employed in evaluating the results obtained.

The blood leukocyte changes following fever induced by all of the methods employed simulated those produced by ACTH with three major differences (1) neutrophilia and lymphopenia were greater after induced fever (2) eosinopenia occurred later than when ACTH was administered, particularly when typhoid vaccine was the pyrogenic agent, (3) leukopenia produced by typhoid vaccine was prolonged to 48 hours. The leukocyte response following fever cabinet pyrexia most nearly resembled that seen in ACTH therapy. Aspirin minimized the pyretic effects of Pv PYROMEN but did not affect leukocyte changes. Neither dibenamine or chlor-trimeton had capacity to alter febrile or leukocytic responses. None of the agents provoked consistent increases in uric acid/creatinine ratios or corticosteroid excretion.

Pv PYROMEN gave a 210% increase in leukocytes within 8 hours; levels returned to normal in 24 hours. There was a sharp rise in

- 1 9 5 1 -

polymorphonuclear cells to levels above those secured with other hyperthermic agents. It is possible that both immunization and toxic effects (not mediated through the anterior pituitary) occur with Pv PYROMEN therapy. Pv PYROMEN does not increase urinary corticosteroid excretion, but this may merely imply that the renal threshold is high. "...with normal pituitary glands, fever and pyrogenic substances may cause the release of small amounts of ACTH..." the authors believe.

716. LEQUIRE, V. S.

The augmentation of the thermogenic effects of pyrogens by homologous plasma in rabbits

J. Infect. Dis., 88:194-206, 1951

Mechanisms involved in the augmentation of pyrogenic response by use of homologous plasma were demonstrated through studies of the minimal and moderate febrile reactions in the rabbit after injection of saline or plasma suspensions of pyrogens from E. typhosa, P. aeruginosa, Aerobacter cloacae and PR-8 Str. influenzae.

Thermogenic response and duration of reaction were heightened when plasma was the suspending fluid. The latent period before initial temperature elevation was shortened. An exception was noted when influenza virus A was injected; the febrile reaction proceeded to peak levels regardless of the material used for suspension, although time of onset and duration were extended through the use of plasma. The pyrogens from E. typhosa were completely removed from the blood stream within 15 minutes when a saline medium was used for the suspending fluid; with plasma, fever was prolonged for two hours. Heparin decreased the pyrogenic activity of E. typhosa and restored a latent period; heparin itself was pyrogenic. With plasma in combination with heparin a febrile response could be obtained.

When plasma pyrogens were pretreated through suspension in this medium, clearance time was prolonged. The activity of pyrogens appears to be a selective precipitation of fibrin and agglutination of the bacterial coat. It is believed that pyrogens do not act directly on the temperature regulation centers, when combined with plasma but alter the surfaces of bacteria or react with certain blood constituents. It was noted that influenza virus A immune serum abolishes leukocytosis and pyrogenic activity.

717. MARTIN, S. P.

Effect of a group of bacteria and their carbohydrates on human leukocytes

Am. J. Med. 11:246, 1951 (Abstract)

Since the somatic carbohydrates of S. typhosa appear to prevent diapedesis of leukocytes from blood vessels, inhibiting cell migration, the effects of bacteria (S. typhosa and P. aeruginosa), of their carbohydrates (purified) and of heparin upon the oriented migration of leukocytes were studied. The slide cell was employed.

718. MCDONALD, R. K., SOLOMON, D. H. and SHOCK, N. W.

Aging as a factor in the renal hemodynamic changes induced by a standardized pyrogen

J. Clin. Invest. 30:457-462, 1951

Changes in the function and anatomical structure appear in the kidney of the aging individual. Alterations in renal function are shown in decreased urea clearance and rate of filtration, the reduction of the effective plasma flow and tubular transfer. Some intimal thickening is seen in renal arteries and arterioles even in the nonhypertensive individual.

The standard pyrogen test was applied to 54 male subjects divided into three groups: young, middle-aged and old. Decreases in renal function were not significant until after the age of 50. When 0.05 cc. TAB vaccine was administered and analyses made of albumin, globulin, inulin, PAH, blood urea N, total serum protein, hematocrit and blood pressure, it was noted that the baseline C declined with advancing age, but no changes were attributable to the pyrogen test itself. Effective plasma flow mean baseline declined with age and, for all ages, pyrogen injections increased flow. Systolic pressure fell in middle-aged and aged subjects; diastolic pressure reductions were approximately the same for all age groups. Renal afferent resistance was highest in older groups, and the fall was more abrupt. Minimal efferent resistance was highest in older subjects. While the filtration factor declined at the peak of the pyrogen reaction in all groups without distinction, the largest absolute decline occurred in aged subjects. The responsiveness of the renal vascular system to pyrogen at older ages would indicate persistence of arteriolar dilatation under such conditions and a certain reversibility of the usual functional alterations.

The purified carbohydrates (PYROMEN-in a concentration of 0.25-5.0 gammas per ml.) and the organisms themselves produced similar inhibitions of human cell migration. The profound effects they exert may be basic to the phenomena of leukopenia and fever. "In addition to the shock these agents produce it would appear that they have a direct effect upon the leukocytes."

- 1 9 5 1 -

719. MENCZEL, E.

A note on the depyrogenization of infusion solutions by hydrogen peroxide

J. Am. Pharm. A. 40:175-176, 1951

The Taub and Hart method for depyrogenizing infusion solutions has been slightly modified by workers of the Government hospital, "Yarkon". One modification replaces the Soxhlet apparatus with large pyrex boiling flasks. After boiling, contents are readjusted to the original volume. The usual standard titration with permanganate is replaced by other methods; fair results were obtained with the spot test of Feigl. The reagent used, a mixture of 0.4% ferric chloride solution and 0.8% potassium ferricyanide solution in equal parts, is sensitive to 0.08 gamma of hydrogen peroxide in a drop. Sensitivity of the test may be further increased if a 10 drops per 100 cc. solution ratio is maintained. Dextrose causes no interference. Both oxidizable substances and pyrogenic contamination may be detected through use of the test. It may be applied to parenteral dextrose solutions. Elimination of detectable hydrogen peroxide by boiling with manganese dioxide for 15 minutes (0.1%) or the use of activated charcoal (1%) for saline and dextrose combinations is to be recommended.

720. MITCHELL, S. Q., and STUART, E. G.

Role of adrenal gland in mechanism of leucocyte changes upon administration of PYROMEN

Am. J. Physiol. 167:810, 1951

PYROMEN, a purified bacterial polysaccharide, was injected intravenously into cats in dosages ranging from 0.5 to 35.0 gammas per kg. body weight. The leucocyte reaction was studied in both normal and adrenalectomized animals. When adrenalectomy was unilateral, no significant changes were elicited in the leucocyte counts until the second or third day after the operation when the typical response was observed. The remaining gland was unable to compensate fully. After bilateral adrenalectomy with maintenance on DCA or cortisone, no leucocyte changes could be noted at any dosage level employed. When portions of the adrenal gland were transplanted into the adrenalectomized animal and desoxycorticosterone discontinued, PYROMEN again induced leucopenia on the 5th, 6th and 7th days; on the 8th and 9th days the initial rise in leucocytes was recorded.

721. MOST, S.

Effect of Shear's Polysaccharide on plasma clotting

Nature, 168:342, 1951

It has been previously suggested (Heilbrunn, Wilson and others) that the polysaccharide extracted from the bacterium, S. marcescens may have heparin-like structure and activity. The action was presumed to involve prevention of mitotic gelation or clotting before division. Like heparin, the substance gives a metachromatic reaction with toluidine blue.

A solution of polysaccharide P25, containing 36 mg. per cc., was diluted with distilled water and added to freshly oxalated human plasma containing 1 cc. of 0.1 M sodium oxalate to 9 cc. of blood. The clotting time at 37°C. of total volumes of 0.1 cc. plasma and equal amounts of polysaccharide and M calcium chloride (0.025) was computed from the time of addition of the last-named chemical until complete clotting had taken place. Shear's polysaccharide inhibited the clotting of recalcified plasma in a manner resembling that of heparin, but commercial heparin had an activity of 138 to 320 times more effective for the purpose stated.

722. PAGE, I. H.

Treatment of essential and malignant hypertension

J. A. M. A., 147:1311-1318, 1951

In a review of the present status of therapy and of the possible additions to the therapeutic armamentarium in essential and malignant hypertension the use of pyrogens is held to be "the treatment of choice" in the latter condition. The author considers it to be a "harsh" treatment as applied in malignant hypertension, but a life-saving procedure.

An infusion of PYROMEN is employed in amounts sufficient to raise the body temperature to 101.0 or 102.0° F. Saline solution is preferred as a diluent, and 200 cc. are infused daily, for six days a week over a period of three to four months. Since tolerance may develop, dosages may increase to very high levels and a rest period of 3 to 5 days is required. In the treatment of some patients it is possible to fix a maintenance level early in the treatment period. Reactions to PYROMEN may be diminished through preliminary intravenous injections of amobarbital (0.2 Gm.) or tolserol (0.5 Gm.) before infusion. Tolserol should be well-diluted to reduce dangers of venous thrombosis resulting from its use.

- 1 9 5 1 -

723. SAETHER, J.

Pyrogener (Pyrogens)

Nord. Med. 54:501-504, 1951

724. SCHOCH, A. G. (Editor)

Clinical experience with PYROMEN. PYROMEN for the treatment of chronic dermatoses

Current News in Dermat. and Syphilol., November, 1951

PYROMEN therapy was evaluated from the results obtained in a group of 52 patients presenting stubborn dermatoses. The most frequently given dosages were 0.5 or 1.0 gamma initially, increased as tolerance developed. Three or four intravenous injections were given per week. Improvement was seen in 39 patients. After interruption of therapy for one to two months, improvement was still evident in 25 patients. Six cases of psoriasis improved but exacerbations were noted. The importance of the psychotherapeutic features of the treatment was stressed by the reporting clinician, who found spontaneous major environmental changes coincident with the fluctuations in about one-half of the patients, and "psychic catharsis" where reactions to treatment occurred. Temporary favorable results from PYROMEN and psychotherapy in combination are reported. Two individuals became conditioned to pyrogenic injections and reacted similarly to injections of saline solution. Astonishing improvement was shown by certain patients. "Although the usefulness of the drug, per se, remains to be determined, it seems that the results would usually not have been as good without PYROMEN".

The Editor quotes Dr. Kierland (Mayo Clinic), "PYROMEN...is of value but not specifically so in the treatment of many..chronic dermatoses...unresponsive to the usual therapeutic measures. There is evidence that subfebrile doses are effective." He considers the action a nonspecific one, similar to that noted in foreign protein therapy, but more controllable than that of TAB vaccine or other pyrexia-producing agents.

725. SCOTT, D., JR. and CLEMENTE, C. D.

Conduction of nerve impulses in regenerated fibers of the spinal cord of the cat

Am. J. Physiol. 167:825, 1951 (Abstract)

The growth of severed nerve fibers and a return of their capacity to transmit nerve impulses were observed in the cat after resection of the spinal cord and treatment with a bacterial polysaccharide, PYROMEN. Responses to stimuli in rapidly conducting descending fibers to the site of the lesion and below the point of severance were recorded. In the animal receiving PYROMEN therapy, impulses conducted to the site of the lesion at 25 to 40 meters per second were registered at a distance some 20 mm. below the lesion. Neural activity below the site of resection was undiminished in velocity. No activity could be demonstrated below the site of the lesion in the control group. Approximately 15% of the total number of nerve fibers normally present appeared to be capable of conducting impulses after PYROMEN therapy.

726. SHTAKEL 'BERG, N. A.

Raziotie likhoradochnoi reaktsii u zhivotnykh pri razlichnoi lokalizatsii vvedeniia pirogennykh veshchestv. (Development of fever reaction in animals in relation to injection sites of pyrogenous substances)

Fiziol. Zh. SSSR (Moskva) 37:195-205, 1951

727. SOYLEMEZOGLU, B., and WELLS, J. A.

Studies on the mechanism of the leukocyte response to bacterial pyrogen

J. Pharmacol. & Exper. Therap. 101:33-34, 1951

The relationships between pituitary-adrenal discharge, appearance of the alarm reaction and the activity of bacterial pyrogens, ACTH and cortisone were examined. Intravenous and intramuscular injections of ACTH (10 mg.) or PYROMEN (500 micrograms) in the dog produced widely differing results. While changes after intravenous administration of pyrogen were essentially the same as those produced by ACTH given by the same route, intramuscularly ACTH induced leukocytosis and PYROMEN did not. The responses to ACTH were modified by adrenalectomy. It is concluded that pyrogen-induced changes take place through some mechanism differing wholly or in part from that of pituitary-adrenal discharge.

- 1 9 5 1 -

728. SOYLEMEZOGLU, B. and WELLS, J. A.

Comparison of leukocyte response to ACTH and bacterial pyrogen

Proc. Soc. Exper. Biol. and Med. 77:43-47, 1951

Removal of the adrenal gland substantially reduced the eosinophilic response in the dog to ACTH but did not affect the response to pyrogen.

Normal animals received either 500 micrograms of Ps. PYROMEN or Pv PYROMEN intravenously or 10 mg. ACTH. Bilaterally adrenalectomized animals received pyrogen-free saline to establish a non-febrile condition, then one or the other of the preparations by vein. Blood samples were examined in 5 minutes, after 15 minutes, 30 minutes and at hourly intervals thereafter for 6 hours. Differences between leukocyte responses were observed. In the normal animal the eosinopenia produced by pyrogens and ACTH was similar. When adrenals were removed eosinopenia was reduced in animals given ACTH, but not in pyrogen-treated dogs. Adrenalectomy reduced total leukocytosis following PYROMEN; however, other factors than pituitary adrenal discharge enter into leukocyte changes after pyrogen administration.

729. STARKA, J.

Pyrogeny (Pyrogens)

Vojenske Zdravotnicke Listy 20:26-28, 1951

730. STETSON, C. A., JR.

Studies on the mechanism of the Schwartzman phenomenon. Certain factors involved in the production of the local hemorrhagic necrosis

J. Exper. Med. 93:489-504, 1951

Many hematologic changes follow intravenous injection of materials found capable of eliciting the Schwartzman phenomenon in the rabbit. These changes include the appearance of thrombocytopenia to a marked degree, a rapid accumulation of platelets and leukocytes in capillaries of the lung, preliminary occlusion of the skin capillaries through thrombi of platelets and neutrophils, and complete interruption of the blood supply, necrosis and hemorrhage. When purified rabbit liver glycogen or agar were administered intravenously, leukopenia developed more rapidly and disappeared more quickly than is the case after bacterial toxins are injected. Glycogen produced agglutination or clumping of platelets and leukocytes. Thrombi bear a resemblance to those noted before severe hemorrhage in mouse sarcoma 180 after injection of either glycogen or meningococcal toxin.

731. STUART, E. G.

Accelerating effect of PYROMEN in contrast to inhibitory effect of cortisone on the Arthus phenomenon in rabbits.

Federation Proc. 10 (Pt. I.) 133, 1951. Abstr. Am. J. Med. Sci. 221:470, 1951

Twenty-four male rabbits were sensitized with crystalline egg albumin cutaneously; 1/3 of the group received daily injections of cortisone (10 mg.) intramuscularly and eight others were given PYROMEN intravenously (10 gammas daily). Precipitin tests for serum antibody were made at intervals. Controls (8 animals) exhibited Arthus reactions in one week, pronounced on the 10th day and appearing in intracutaneous sites throughout the term of experimentation. After PYROMEN injections, the Arthus reaction appeared earlier (4th or 5th days). The lesions passed rapidly through the usual course, but with more intensified inflammation, hemorrhage, suppuration and necrosis. The size of the area exceeded that seen in control animals. Scab formation was rapid and resorption occurred, usually without sloughing. Precipitin titers were noted earlier in PYROMEN-treated animals and were at least 50% above those seen in controls. "These effects of PYROMEN may be correlated with its known property of stimulating the leukopoietic, phagocytic and reticular elements of the body".

732. STUART, E. G.

Connective tissue mast cell responses to bacterial pyrogens, ovalbumin and cortisone

Anat. Rec. 109:351, 1951

Pyrogens extracted from a Pseudomonas species, PYROMEN, or from Proteus vulgaris and B. subtilis were injected intraperitoneally into C-57 mice in dosages of 10 gammas each. Type response of the connective tissue mast cells differed from that produced by substances chemically unrelated. Proteins and steroids, as ovalbumin and the adrenocortical factors or hormones, ACTH, DCA and cortisone, were also injected in 10 mg. single doses. Only slight responses were elicited by the first two; cortisone had a marked influence widely different from that exerted by the pyrogens. Ovalbumin, 1 mg. single injection, produced dissimilar effects; the pyrogenic concentrate from E. typhosa, (10 gammas) produced a histologic picture resembling that seen after ovalbumin injections. With the pyrogens used, alterations in granulation, pyknosis or expansion of cells were seen. Distribution of cells and cell groups were altered and cellular deterioration was observed.

- 1 9 5 1 -

733. TENNENT, D. M. and OTT, W. H.

Tolerance in rabbits to bacterial pyrogens

Federation Proc. 10:396, 1951

Studies have been made of the factors which influence the development of tolerance in rabbits injected with bacterial pyrogens. Special emphasis was placed upon the size of the dose and time required for recovery of the normal febrile response. Tolerance and recovery time were found to be functions of the amount of pyrogen injected. With levels of dosage sufficient to produce 0.6° C. increase in body temperature, a rest period extending beyond one week was necessary to restore the normal febrile response in some animals.

734. THOMAS, L. and GOOD, R. A.

Bilateral cortical necrosis of kidneys in cortisone-treated rabbits following injection of bacterial toxins

Proc. Soc. Exper. Biol. & Med. 76:604, 1951

Five groups of rabbits were studied after each group had received one of the following: (a) 25 mg. cortisone intramuscularly daily for four days with 2 cc. of meningococcal toxin intravenously (1:10 dilution) the third day, (b) meningococcal toxin alone by vein, (c) 25 mg. cortisone daily for four days, (d) 25 mg. cortisone per day and 0.4 mg. S. marcescens toxin intravenously on the third day, (5) P-25 toxin alone. Smaller doses of cortisone were administered also, and the effects of shortened periods of treatment were noted.

Bilateral renal cortical necrosis was evident in a majority of the animals treated with cortisone and bacterial toxin in combination. No changes were evident in control animals. Seven of eight animals treated with 10 mg. cortisone and 0.2 mg. P-25 toxin developed necrotic areas in the kidneys.

735. WELLS, J. A. and SOYLEMEZOGLU, B.

Suppression of leukocyte response to bacterial pyrogen by nitrogen mustard

J. Pharmacol. & Exper. Therap. 103:366, 1951

Since the preponderance of unsegmented granulocytes in the leukocyte picture following the injection of bacterial pyrogens suggests bone marrow stimulation, methyl bis (b-chloroethyl)-amine was administered for its anti-hematopoietic effect, the total and

differential leukocyte counts performed at hourly intervals thereafter, and comparisons made with normal levels. In the rabbit initial leukopenia was not succeeded by rises in total leukocytes. "It is assumed that the biphasic leukocyte response to pyrogen (PYROMEN) is due to irreversible removal of cells from the circulation, and that the subsequent leukopenia is normally obscured by a release of newly formed cells from the bone marrow."

736. WINDLE, W. F., and CHAMBERS, W. W.

Regeneration in the spinal cord of the cat and dog

A.M.A. Arch. Neurol. & Psychiat., 65:261, 1951 (Abstract)

In experiments planned to determine the possible site of action of a purified bacterial pyrogen (PYROMEN) the spinal cords of both cats and dogs were transected at different levels. The pyrogen was administered for periods extending from 8 to 61 days. The pyridine-silver method was employed for staining sections

Histologic evidence of new growth of neurons into or across the cut was secured from the cords of animals living 42 days or more. Some connective tissue scarring and the usual amount of retrograde degeneration were observed. There appeared to be no barrier of glial tissue, but a blending of the connective tissue with the parenchyma of the spinal cord. Through such areas, the nerve fibers could be seen growing into the scar. Numerous blood vessels were seen at the scar-cord junction; macrophages were numerous in the spaces between strands of non-nerve cellular elements and the nerve fibers in the cord. Many nerve fibers had penetrated the scar and there was invasion of the opposite cut end. Serial sections revealed that nerve fibers were outgrowths of severed or injured nerve roots, vascular and meningeal nerves and intrinsic cord neurons. No functional regeneration was obtained.

737. WITTICH, F. W.

PYROMEN in the treatment of perennial respiratory allergies

Ann. Allergy 9:502-407; 518, 1951

Fifty patients who presented symptoms of chronic, perennial asthma received PYROMEN when conservative procedures had not proved wholly successful. All received careful study as to offending allergens, and many were under immunization therapy or on eliminatory diets. Subfebrile dosages of 0.5 gamma initially were increased as indicated by response. Oral therapy gave variable results and was abandoned in favor of intravenous measures for the most part; a few instances occurred where oral

- 1 9 5 1 -

therapy appeared to sustain clinical improvement established previously by intravenous dosage.

Good results were obtained for 12 patients and 18 others reported lessened severity and frequency of attacks. The disappearance of fatigue was notable. There appeared to be a more complete response to bronchodilators and vasoconstrictors after PYROMEN therapy. Twenty patients were unbenefited. While therapeutic results were less rapid than those seen after ACTH or cortisone therapy, the side effects were relatively few and far less hazardous. The use of PYROMEN as adjuvant to other measures for the control of asthma is indicated.

738. ZINDLER, G. A.

Pyromen therapy in the treatment of food allergy.

Ann. Allergy 9:494-501, 1951

A large group of patients (125) with varied types of allergies to foodstuffs received subfebrile doses of PYROMEN intravenously. Initial amounts injected ranged from 2.0 to 3.0 gammas; graduated amounts were then administered until benefits were evident. Intradermal administration of PYROMEN was effective in a schedule of dosages over the same range.. Sublingual administration was effective when accompanied by booster doses, given by either the intradermal or intravenous routes. Oral dosage ranges were 50 to 100% above parenteral. Results were most satisfactory where control of the offending allergen accompanied the administration of PYROMEN. Tolerance was shown by the return of the ability to accept a general diet without untoward reactions. Increases in appetite and gains in weight were noted even where qualitatively restricted diets were prescribed.

739. ZYKINA, E. S.

Razvitie likhordochnoi reaktsii pri vvedenii pirogennykh veshchesh u normal' nuiu i defferentiro vannuiu konechnosti (Development of febrile reactions during injection of pyrogenous substances into the normal and deferentectomized extremities)

Fiziol. zh. SSSR (Moskva) 37:186-194, 1951

- 1 9 5 2 -

740. CLEMENTE, C. D.

Penetration of brain by regenerating peripheral nerve fibers

Anat. Rec. 112:319, 1952

When the temporal branch of the facial nerve of the cat was severed and the stump implanted into the temporal cortex, the regenerating nerve fibers in animals receiving PYROMEN* rapidly penetrated the cortex and blended with cerebral tissue. Such regrowth was seen as early as 30 days after operation. Control animals exhibited glial barriers which prevented blending of regenerating nerve fibers with central nervous parenchyma. No blending was shown when cortisone was administered to cats with peripheral nerve implantation in cerebral tissue. "The effects of other steroids are being studied."

741. CLEMENTE, C. D., SCOTT, D., JR., CHAMBERS, W. W. and WINDLE, W. F.

Structural and functional regeneration in the central nervous system

A.M.A. Arch. Neurol. Psychiat., 1952 (in press)

742. DOUGLAS, W. W., and PATON, W. D. M.

The hypothermic and antipyretic effect of preparations of ACTH

Lancet 1:342, 1952

Preparations of ACTH lower body temperature and delay the response to pyrogens in normal rabbits. Where pyrexia is established after pyrogen administration, it is reduced by subsequent administration of ACTH. This hypothermic effect has not been previously reported.

In these studies, pyrexia, induced through injection of 1 microgram per kilogram body weight of the cells of Proteus vulgaris, was measurable in 20 to 30 minutes. ACTH, pyrogen-free, was injected in amounts of 1 or 3 units per kilogram body weight and the temperature reduction measured. Each dose reduced the

*PIROMEN

- 1952 -

temperature from 0.24 to 1.97° C. at the single unit level of dosage, and as much as 2.5° C. at higher dosages. The onset of pyrexial response from pyrogen was delayed for 1-1/2 to 2 hours - the exact period of the hypothermic influence of ACTH. The possibility that oxytocic activity from posterior lobe pituitary hormone might be responsible was ruled out by careful testing.

The significance of hypothermic effects of ACTH in relation to routine testing of drugs is apparent. The ability of ACTH to delay the appearance of pyrogenic responses will make longer testing periods (to five hours, preferably) obligatory. It is believed that the activity of ACTH in a given disease condition may be wholly or in part due to nonspecific metabolic or vascular alterations.

743. FREEMAN, G.

The anticoagulant effect of bacterial polysaccharides in normal and thrombocytopenic plasma of leukemia

Blood, 7:235-242, 1952

Polysaccharides from extracts of cultures of Friedlander's bacillus, type A, of Serratia marcescens (P-25 of Shear) or type II pneumococcus were compared for respective anticoagulant activity with a heparin sodium solution. Recalcification and clotting times were computed for both normal and thrombocytopenic plasma.

An arbitrary amount of polysaccharide from Friedlander's bacillus, type A, inhibited retraction of the fibrin clot. Pneumococcal polysaccharide has less anticoagulant activity than either the Friedlander or S. marcescens somatic carbohydrate. These two materials were used in comparing recalcification time when amounts of polysaccharides were increased in blood platelet ranges remained normal. Comparisons with thrombocytopenic plasma were also made. Effective concentrations of the polysaccharides were as low as 0.05 mg. per 0.1 ml. normal plasma, but heparin was shown to be 400 to 1000 times more effective as an anticoagulant.

In leukemic blood the two bacterial extracts had an exaggerated anticoagulant action. The evidence that bacterial polysaccharides exist in combination with antibody in disease states is reviewed. Non-glucoseamine polysaccharide in serum may be increased to 300 mg. per cent in disease. Work in progress at this time may prove more conclusively a relationship between the presence of such polysaccharides, the bleeding phenomenon and fever in leukemia.

744. GRANT, R. and WHALEN, W. J.

Latency of response to bacterial pyrogens as affected by previous incubation with blood

Federation Proc. 11(Pt. I) 58, 1952 (Abstract)

When small amounts of PYROMEN* (from Pseudomonas aeruginosa) were incubated with whole blood, heparinized, and injected in animals, the onset of fever reactions appears to be hastened. There is vasoconstriction, inhibition of polypnea, and rise in rectal temperature initially, but no augmentation of fever is obtained. No such effects followed the use of plasma, either heparinized or citrated. No explanation for either acceleration or augmenting effects seen after injection of other pyrogens in citrated plasma (LeQuire, J. Infect. Dis. 88:194, 1951) can be offered at this time. It is believed that tissue "injury" products are not involved in the production of a latent period before temperature increases.

745. GUERRIERI, J. A.

Treatment of neurodermatitis with PYROMEN - A new therapeutic agent

Clin. Med. 59:29-35, 1942

Chronic and acute neurodermatitis, in varying manifestations and recalcitrant to treatment, has shown response to minute doses of a bacterial polysaccharide, PYROMEN.*

Fifty-two patients, in a group of 60 under treatment, had some lessening or complete remission of dermatologic symptoms; improvement was shown with 1 week in conditions present for 6 months or less, and in 2 weeks for chronic conditions (over 1 year). The initial dose administered was either 1.0 or 2.0 gammas, and the therapy was continued, weekly or semi-weekly, for periods ranging from 2 weeks to 7 months. Remission followed in 18 patients with acute forms of neurodermatitis and in 8 with chronic dermatoses. The remissions extended longer than when ACTH (or cortisone) was administered. Greater safety was evident; but one febrile reaction was reported and there were no undesirable side effects. No resistant exacerbations followed the discontinuance of therapy. PYROMEN* may achieve its beneficial action through increasing the natural defenses of the dermis against local trauma or toxic invasion.

*PYROMEN

- 1 9 5 2 -

746. NICHOLLES, P. S. and BUBEL, H. C.

Increased resistance to infection by the concomitant administration of bacterial pyrogen and vaccine

Bact. Proc. (Soc. Am. Bacteriologists) 1952, p. 74

Pyrogen from S. typhosa administered parenterally with B. tularensis significantly increased the resistance of test animals to a challenging dose of a highly virulent strain of B. tularensis. Rabbits treated simultaneously with pyrogen and vaccine (B. tularensis) produced antibodies to a greater degree than did rabbits which received pyrogen or vaccine only.

747. SCHOPP, A. C., FELLHAUER, C. M. and EIGEL, E. G.

Preliminary observations on a new drug in the treatment of acute poliomyelitis

Med. Bull. St. Louis Univ. 4:6-10, 1952

PYROMEN* was administered to every second patient admitted for the treatment of acute poliomyelitis during a period of six months. Clinical impressions of the results of this therapy for 53 patients were recorded, although random selection of this type does not afford opportunity for carefully controlled studies. Supportive treatment was employed for the majority of these patients. The usual dose of PYROMEN* ranged from 3.0 to 5.0 gammas, administered twice daily. More intensive therapy was employed in a few cases; in one, a child 17 months old, a single massive dose of 80 gammas was administered without harmful effects. A direct relationship could be noted between early reactions to the drug and the therapeutic benefits achieved. Reactions were varied; fever, chills and myalgia appeared most frequently; extreme lethargy was noted in a few patients. These symptoms of reaction disappeared early and no after effects were apparent.

While mortality was not appreciably altered by the administration of PYROMEN*, many beneficial results followed treatment. In spinal paralytic paralysis, spasm, and its accompanying pain, was relieved. Residual spasticity was less often observed. The most dramatic recoveries occurred in encephalitic involvement, where complete reversal of the clinical picture of acute illness was observed. While the 7 deaths reported were those of patients with bulbar poliomyelitis, the PYROMEN-treated group included patients with the most serious aspects of this

*PYROMEN

- 1952 -

form of the disease. The drug appears to exert its most pronounced effects where higher centers of the central nervous system are affected. The duration of acute illness is materially shortened. The drug appears to be safe for use and merits further study in the treatment of acute poliomyelitis.

748. SCOTT, D., JR. and CLEMENTE, C. D.

Mechanism of spinal cord regeneration in the cat

Federation Proc. 11(Pt. I) 143, 1952 (Abstract)

A second factor is involved in neuronal regeneration in the spinal cord of the cat following treatment with PYROMEN*, since a small number of animals have failed to show significant regeneration after treatment. When, at the time of cord resection, the distance separating proximal and distal stumps of the cord were measured and the proximation compared with results after PYROMEN* treatment, it was found that when the distance was 1 mm. or less between the two sections of the cord at operation, active regeneration followed with transmission of potentials to 25 to 30 per cent of normal. No regeneration was seen beyond the site of the lesion when the separation was 2 mm. or more. It is believed that both treatment with the bacterial polysaccharide PYROMEN* and close proximation of the two cut ends are necessary for successful regeneration of spinal cord neurons in adult cats.

749. SEIBERT, F. B.

Introduction. Symposium on bacterial pyrogens

Tr. New York Acad. Sc. 14:157-158, 1952

The nature of 'true' pyrogens has not yet been fully established. While it is believed that many of the pyrogens are of carbohydrate structure, the polysaccharides of certain organisms, e.g. the tubercle bacillus, do not produce temperature increases, while protein fractions are pyrogenic. It would appear that the purified polysaccharides of tuberculin and of the parent bacillus are not pyrogens. "The isolation of a pure and stable pyrogen, with well-defined chemical and biological properties which could be easily reproduced, would furnish a valuable standard for the evaluation of pyrogen in products for intravenous use."

*PIROMEN

- 1 9 5 2 -

The effect of pyrogens on capillaries, their relationships to the Schwartzman phenomenon and to tumor hemorrhagic and necrotizing changes in tumor tissue, are not, to date, fully explained. The alterations in blood component distribution and vascular pressures are well-established. It is not yet known whether or not the leucocyte response is a direct or indirect phenomenon; the time required for its production after pyrogen administration would suggest that some intermediary is required for its appearance.

In studies now in progress, tuberculin protein fractions have been shown to have more definite effects than the polysaccharides on the leukocytes. Fractions which produced clumping of leukocytes, swelling and inhibition of cellular migration also produced fever after intravenous injection.

750. SOKOLOFF, B., EDDY, W. H., CONE, G. and POWELLA, R.

Stress reactions and tumor-breaking property of bacterial polysaccharide

Federation Proc. 11 (Pt. I) 427, 1952

Increased capillary fragility appears to be the prime factor in the tumor-breaking property of bacterial polysaccharide. Rats bearing Crocker rat carcinoma were given 0.5 mg. per 100 Gm. body weight of P-25 (Shear bacterial polysaccharide) alone or with 10 mg. per 100 Gm. body weight of flavonoid compound. When the flavonoid accompanied the dose of bacterial polysaccharide 66 per cent survived and the remaining 34 per cent had an extension of life for, on the average, 46 hours. Controls succumbed in approximately 7 hours and 44 minutes. When the ascorbic acid level of the blood plasma was reduced to 0.01 mg. per 100 cc. plasma and 0.1 mg. per 100 Gm. body weight of P-25 injected, the mortality was increased to 25 per cent or a 10 per cent increase over control groups; tumor destruction, however, was very extensive, approximately 5.3 times greater in area than in controls.

751. TENNENT, D. M.

Isolation of bacterial pyrogens

Federation Proc. 11(Pt. I) 58, 1952 (Abstract)

In a comparison of methods for isolating active pyrogenic substances, a crude bacterial pyrogen concentrate, prepared by precipitation of harvest supernatants of coliform or Salmonella organisms, had higher activity than that reported for purified

fractions secured through phenol extraction or tryptic degradation. Analysis of the crude concentrate has been completed. The effects of filtration, lyophilization, and dialysis were studied. Alcohol precipitation with pH adjustment yielded a preliminary fractionation; solvent extraction and precipitation with heavy metals were not successful. A large proportion of the active substance was lost during Darco separation, Florex or acid-washed alumina adsorption.

752. THOMPSON, R. M.

A lipo-protein-nucleic acid complex in the treatment of radiation injury. A preliminary report

Mill. Surg. 110:51-58, 1952

The principal therapeutic problem in mass radiation injury lies in the high mortality rate found among victims who receive the approximate median lethal or sublethal exposures to ionizing rays. The most important factor increasing the death rate is the development of susceptibility to infection which follows pancytopenia and impaired antibody formation. Radiation damages circulating lymphocytes and lymphatic tissues.

The activation or stimulation of the reticuloendothelial system through non-specific therapy has been well-documented. Certain nucleic acid salts have been utilized in the treatment of agranulocytosis. On these bases, trial has been made of the protection against radiation after-effects afforded by the injection of a lipo-protein-nucleic acid complex (reticulose). Animals receiving 450 r. of radiation and intramuscular injections within 24 hours of 2 cc. of reticulose, showed an arrest in the sharp fall in total leukocytes. The leukocyte level was sustained at approximately 3000 cells per cu. mm. for several days, and then increased to 7000 cells, on the average, within 21 days after radiation. Leukocytes fell to dangerously low levels in control animals. Normal animals showed marked elevation in the total leukocyte counts when given reticulose (without exposure) for 30 days. Normal cellular components were noted in bone marrow, but the spleen exhibited striking hyperplasia of the reticuloendothelial cells in the perifollicular zone. Irradiated animals had fewer cellular elements in the bone marrow, but the production of granulocytes and erythrocytes was elevated. Megakaryocytes were present in smaller numbers than normally seen. A moderate depletion of Malpighian corpuscles and an increase in mature granulocytes were noted in the spleen.

- 1 9 5 2 -

753. TOPPER, Y. J. and LIPTON, M. M.

Biosynthesis of capsular streptococcal polysaccharide

Federation Proc. 11(Pt. I) 299, 1952 (Abstract)

The specific activities of the capsular polysaccharide of hemolytic streptococci, Group A, and of glucosamine derived from it were studied after culture on a medium containing glucose- $1C^{14}$. The isotope content of glucosamine and glucuronic acid was the same, indicating a similar source for both. Glucose appeared to be directly converted to glucosamine without structural changes. When identical quantities of glucose- $1C^{14}$ were added to equimolar, non-isotopic glucosamine, glucuronic acid and glucosone, the polysaccharide was ten times as radioactive as the glucosamine it contained. "These findings indicate that free glucosamine is efficiently incorporated into the bacterial polysaccharide whereas this is not the case for glucuronic acid."

754. WINDLE, W. F.

Activities of certain bacterial polysaccharides (Symposium on Bacterial pyrogens)

Tr. New York Acad. Sc. 14:159-161, 1952

The physiological activities and histologic changes which follow the administration of highly purified non-protein bacterial polysaccharides in humans and in animals are reviewed. Polysaccharide preparations derived from a Pseudomonas species (PYROMEN)* and from Proteus vulgaris were used in these studies. These polysaccharides are of low toxicity, the LD₅₀ in mice being about 60,000 micrograms.

The present measure of activity, or 'minimal PYROMEN* response' is a rise in the number of granular and nongranular circulating leukocytes. This can be elicited in most human subjects by less than 0.05 micrograms of PYROMEN* per kilogram of body weight, given intravenously. Human subjects are more variable in response to a given dose than laboratory animals; the latter require nearly 10 times the minimal human dosage. The total leukocyte count may rise 50 to 100 per cent above the initial level during the peak of a minimal PYROMEN* response and it is possible to obtain superimposed responses by administration of similar amounts of the polysaccharide shortly after each peak in leukocyte production has passed. A second type of response has been noted in both human subjects and in animals; this is characterized by a marked leukopenia initially, with a lesser

*PYROMEN

degree of leukocytosis thereafter. The leukocytosis is accompanied by a depression in the number of circulating lymphocytes and eosinophils and often, by an increased body temperature. In the cat, leukopenia is accompanied by a rise in the plasma ascorbic acid level.

Administration of the polysaccharide preparation produces histologic changes; hyperplasia of the thymus, lymphatic and splenic corpuscles; depletion of the lymphocyte population of peribronchial tissue and of the tunica propria of intestines; and reduction of cytoplasmic droplets in the fascicular zone of adrenals and follicular colloid of the thyroid gland. The bone marrow was not studied. In the cat and rabbit, treated daily or three times weekly for 10 to 120 days with a polysaccharide from Pr. vulgaris, certain changes reminiscent of rejuvenation or reversal to the fetal state appeared in lymphoid tissue, the reticuloendothelial system and the endocrines. There was no evidence of the type of pathology seen in true stress.

Endocrine mechanisms appear to be affected by PYROMEN*. Chronic bilateral adrenalectomy in cats abolished the minimal PYROMEN* response; transplantation of adrenal cortical tissue revived it. Toxicity was increased for the adrenalectomized animal. Hypophysectomy altered the degree of response, reduced the leukocytosis which follows. Thyroidectomy delayed the onset of leukopenia and leukocytosis and ovariectomy produced a similar picture. Estrus, pregnancy and lactation produce variations in the response to PYROMEN*.

The enhancing of the healing process in skin burns by PYROMEN* may be related to an increase in the permeability of connective tissue. The most dramatic activity of PYROMEN* to date has been its striking inhibition of glial barriers in the transected spinal cord and the regrowth of neural fibers across the approximated severed ends. Recent experiences have shown that glial barriers against the growth of peripheral nerves may be inhibited in brain tissue.

755. WINDLE, W. F.

Regeneration in the spinal cord

J. Paraplegia 2:3-5, 1952

In a discussion of the therapeutic implications inherent in the discovery that prevention of glial overgrowth at the site of resection of the spinal cord follows the use of PYROMEN*, the author lays emphasis upon the fact that the application of these

*PYROMEN

- 1 9 5 2 -

findings to human beings must await the outcome of further investigation. In progress at the present time are certain experiments to determine whether or not alterations can be achieved in scars already formed and the effects of surgical removal of such scars followed by treatment with PYROMEN*.

756. WINDLE, W. F., CLEMENTE, C. D. and CHAMBERS, W. W.

Inhibition of formation of a glial barrier as a means of permitting a peripheral nerve to grow in the brain

J. Compar. Neurol., 1952 (in press)

757. WINDLE, W. F., CLEMENTE, C. D., SCOTT, D., JR. and CHAMBERS, W. W.

Induction of neuronal regeneration in the Central Nervous System

Transactions, Am. Neurol. Assn., 1952 (in press)

After complete transection of the thoracic spinal cord of the cat, glia cells from the parenchyma, with fibroblasts from blood vessels and pia mater form membranes over the cut ends. These are well-developed within one week and firm scar tissue is seen within one month in untreated animals.

Administration of PYROMEN* or ACTH from the day of operation delays the early development of the pia-glial membrane over the cut ends of the cord. A loose type of "granulation tissue", consisting of a well-vascularized reticulum, filled with connective tissue cells poorly differentiated, permits regenerating intraspinal neurons to penetrate the newly-formed tissue within fifteen to thirty days after treatment is begun. Successful regeneration depends upon proximation of severed ends and continued therapy. Electrophysiological studies have confirmed histologic findings. When separation of the cord ends was not more than 1 mm. at operation, recorded electric potentials after regrowth were within 25 to 30 per cent of normal. With separation of 2 mm. or more scar tissue formed and no potentials were recorded. Unless other measures are taken to eliminate scars, it is doubtful if paraplegics with spinal cord lesions of long standing can be benefited by the therapy employed in our experiments.

*PYROMEN

758. ZAHL, P. A., and HUTNER, S. H.

Biology of pyrogens Symposium on bacterial pyrogens

Tr. New York Acad. Sc. 14:161-163, 1952

The most frequently found and most potent pyrogens appear to be those associated with the somatic antigens of gram-negative organisms. Certain toxic effects appear after intravenous administration of these substances: hemorrhage induction in transplanted sarcoma; interference with the homiothermic system; vascular damage in deciduo-placental tissues; induction of leukopenia, hyperglycemia and adrenal cortex damage. Immune bodies may be produced. "All immunochemical work to date indicates that pyrogenicity, at least as associated with the gram-negative bacteria, represents but one aspect of poisoning by endotoxins." "Immunologic aspects have been ignored by some workers while others have neglected the purely pyrogenic..." The relationships of pyrogens to shock and stress have not been fully studied.

The sources of gram-negative bacteria which produce pyrogens are multiple. The Pseudomonas, for example, thrive in moist locations, as do most hydrogen bacteria. Pyrogenic contaminants may be reduced in number by preventive measures or by the use of chemicals (chlorobenzene, dichloro-preparations or ethylene oxide) where heat may be damaging. In apparatus and in packaging, polyethylene offers advantages over glass, since it is an inert material ~~without~~ without nutrient salts in its composition.

Pyrogenic endotoxin may be dependent on the presence of phosphate groups attached to the polysaccharide or to a protein component. It may be an antimetabolite. In the organism, pyrogenic endotoxin does not injure isolated cells, but appears to attach most strongly the epithelial cells of newly proliferated capillaries. Another interesting line of study would lead the worker to consider the possibility of antagonism between pyrogenic endotoxin and anti-thyroid substances.

AUTHOR INDEX

Abernethy, T. J.	248	Banks, H. M.	208
315		Barbour, H. G.	90
Adams, J. R., Jr.	371	91, 117, 369	
372, 380		Barr, D. P.	104
Adkison, J. L.	302	Barsi, I.	468
303		Battle, W. H.	28
Aguilar, R. M.	435	Bayer, O.	514
Ahuja, M. L.	389	Beck, L. V.	438
Albacht,	168	439, 440, 469, 515	
Alberty, R. A.	512	522, 622, 697	
Albot, G.	618	Beckwith, T. D.	287
Alder, A.	263	Beebe, S. P.	33
Alexander, H. E.	450	Beeson, P. B.	296
Alexander, L.	513	441, 470, 471, 472	
Algire, G. H.	436	585, 623	
437, 467, 695		Bellach, H.	516
Allen, A. W.	144	Benacchio, L.	517
Altschule, M. D.	415	Benaglia, A. D.	276
416, 619		Bendersky, I.	326
American Med. Assn.	220	Bendix, B.	38
Anderson, C. G.	274	Bennett, I. L. Jr.	556
Anderson, T. H.	319	557, 623,	
Anderson, W. H.	552	684	
Andervont, H. B.	235	Bennett, G. A.	500
236, 244, 275,	294		
Andrews, E.	130	Benventi, M.	555
Anthony, P. Z.	665	Bergmann, J.	38
Aoyama, K.	531	Bergsman, A.	624
582, 583, 584,	648	Berk, A.	127
Apitz, K.	224	Berkowitz, D.	515
Archer, B. J.	696	Berman, H.	662
Ardiet, M. T.	635	Berman, M.	259
Ardry, R.	630	Bernard, B. N.	618
Arendshorst, W.	620	Bernheim, A. I.	255
Arneth, J.	168	266	
Ashenburg, N. J.	295	Bernstein, C.	473
621		Berry, G. P.	295
Aubert, J. P.	553	261	
Auld, A. G.	76	Bertrand, I.	474
Avery, O. T.	69	Bharucha, K.	558
109, 133, 315,	322	Bieling, R.	62
Avineri-Shapiro, S.	258	Bierry, H.	209
326, 356		Billroth, Th.	5
Bailey, G. H.	280	Binkley, F.	418
Baker, E. E.	554	Biocca, E.	388
Balcar, J. O.	81	Bjerknes, C.	384
Baldwin, D. S.	564	411	
Bandelin, F. J.	417	Black, J.	2
Banus, M. G.	188	Black-Schaffer, B.	475

Blauch, B.	469	Calhoun, H.	82
522		Calvery, H. O.	377
Blickenstaff, D.	580	383	
625		Camejo, E. G.	422
Bloom, N.	335	Cameron, G. R.	297
Blum, D. N.	210	Caminita, B. H.	354
Blumenthal, L. S.	626	Campbell, D. H.	421
Boikan, W.	130	Campbell, D. J.	391
Bonetti, E.	419	Canneyt, J. van,	119
Bonorino, C. J.	73	Canziani, G.	195
Boor, A. K.	451	Carletti, B.	629
538		Carlson, A. J.	74
Boquet, P.	476	79	
477, 518, 521,	627	Carminati, V.	196
Borchardt, W.	145	Carter, E. B.	160
146		Castaing, M.	633
Boroff, D. A.	559	634, 635	
560		Cecil, R. L.	104
Bose, B. C.	389	212, 226	
Boshame, R. K.	118	Centanni, E.	18
Bouthier, G.	571	21, 58, 147, 237,	
635		298, 336	
Bove, E.	628	Challinor, S. W.	301
Boyland, E.	249	Chambers, W. W.	481
Boyland, M. E.	249	563, 689, 691, 703,	
Bradley, S. E.	420	704, 736, 741, 756,	
478		757	
Bragdon, J. H.	479	Chandler, V. L.	432
Branham, S. L.	131	Chapman, C. J.	337
Breuer, S.	154	Charonnat, R.	699
Bril' M T, Knokhitkin	698	700, 701	
Brindle, H.	442	Chary, R.	630
Broderson, R.	552	Chasis, H.	264
Brues, A. M.	390	320, 338, 397, 420,	
Bruetsch, W. L.	561	564	
Bryan, R. S.	614	Cherkin, A.	421
615		Chickering, H. T.	63
Bubel, H. C.	746	Christian, S. L.	148
Buchner, H.	15	Citron, J.	55
Bueno, P.	480	Clark, G.	289
Bunting, H.	562	Clay, C. L.	127
Bunting, M. I.	562	Clemente, C. D.	631
Burdon-Sanderson, J.	9	693, 702, 703, 704,	
Burnet, F. M.	211	725, 740, 741, 748,	
215		756, 757	
Buschmann, P.	115	Close, V. P.	608
Butz, C.	448	Clough, P. W.	705
		Clutterbuck, H.	3

Cohen, S. J.	79	Delaunay, A.	316
Coley, B. L.	452	342, 365, 366, 367	
Coley, W. B.	19	393, 476, 477, 518,	
23, 27, 29, 32, 34,		521, 569	
36, 136, 149, 210		Delaunay, M.	521
Collier, H. O. J.	482	569	
Conan, N. J.	478	DeMerre, J.	423
Condini, P.	483	Denman, F. R.	714
Cone, G.	750	Dennis, E. W.	253
Consouloff, S.	120	279	
Conway, E. A.	265	DeRenzi, S.	299
Coons, A. H.	654,	Devlin, H. B.	434
658		674	
Cooper, F. S.	363	Dickenson, C.	1
386, 387, 414		Dietel, F.	171
Corcoran, A. C.	549	Diller, I. C.	443
603		469, 484, 485, 486	
Corelli, F.	632	519, 522	
Cortina, M. C.	435	Dimitrijevic, I.	199
Co Tui	238	Dochez, A. R.	69
250, 251, 252, 276,		Domba, E.	570
277, 278, 339, 340,		Dommers, P.	355
341, 392		Donaggio, A.	197
Council on Pharmacy and		Dorche, J.	571
Chemistry, Amer. Med.		633, 634, 635	
Association	220	Douglas, W. W.	742
Cowie, D. M.	82	Dremland, B. H.	332
94		Dreyfus, G. L.	121
Creech, H. J.	484	Driscoll, R. H.	426
519, 520, 565, 566,		DuBois, E. F.	96
616		104	
Crutchfield, A. J.	431	Dubos, R. J.	343
Culmer, C. U.	302	Duggan, W. F.	344
303		Dunn, T. B.	444
Culver, H.	95	464	
Curry, J. J.	567	Duplan	70
Czaczkas, J. W.	453	Duran-Reynals, F.	198
Czerny, V.	24	225	
Dagley, S.	568	Dutt, J. N.	179
Darrow, K. E.	213	Eddy, W. H.	750
Davesne, J.	169	Edelmann, A.	368
Davis, B. F.	68	Edison, A. D.	448
Dawes, E. A.	568	Editorial, <u>Lancet</u>	667
Deane, H. W.	654	Edsall, G.	487
658		Egami, F.	587
de Aragão, R. M.	404	617	
DeBakey, M. C.	714	Ehelson, L. M.	300
Delafield, M. E.	170	Eigel, E. G.	747
214, 297		Elkins, E. C.	550

Ellis, E. L.	317	Franke, F. E.	394
Ellis, F. A.	572	395, 396	
Emmerich, R.	25	Franzolin, C.	640
Emory, M. L.	375	Freedberg, A. S.	415
Evans, A. C.	172	416	
Evans, C. L.	132	Freeman, G. G.	301
Everett, M. R.	614	319, 743	
615		Freimuller, F.	641
Fahraeus, R.	122	Fremont-Smith, L.	155
123		Frese, H.	6
Falls, H. F.	620	Friedman, H.	542
Farbman, A. A.	329	Friedman, M. H. F.	329
Faria, R.	706	Frisch, A. W.	203
Farr, R. S.	523	Frontali, N.	524
636		Fuchs, M.	626
Favorite, G. O.	345	Furth, O.	106
446		Gabor, T.	570
Fedorov, N. A.	346	Gardner, R. E.	280
Feldman, J.	318	Gay, F. P.	62
Fellhauer, C. M.	747	Gayhart, C. H.	523
Felton, L. D.	488	Gehes Codex	260
573		Gellhorn, E.	318
Fernando-Torres, F. R.	56	Georgi, C. E.	490
Ferry, C. W.	541	Gerber, I. E.	255
Fertig, H. H.	549	266, 347	
<u>Fever Therapy</u>	254	Gerlough, T. D.	309
Field, J.	527	German, A.	575
508, 529, 606,	707	Giaja, J.	199
Figueroa, E. S.	503	Giehm, G.	227
Findelstein, H.	173	Gifford, S. R.	491
174, 175		Ginger, L. G.	665
Finland, M.	659	691, 709	
Fisher, M.	439	Ginsberg, H. S.	492
440, 469		525, 526, 576	
Fishgold, J. T.	581	Ginsburg, E.	188
707		Giuffre, I.	240
Fitch, E. M. J.	708	Gluck, R.	445
Fitzsimons, E. J.	709	Goebel, W. F.	109
Flensburg, E. W.	638	133, 418, 492,	525
639		526, 554, 683	
Flint, L. D.	375	Goldburgh, H. L.	408
Flothow, M. W.	609	Goldie, H.	176
Flusser, B. A.	403	Goldring, W.	264
Fogg, L. C.	239	320, 338, 397,	420
Foster, J. V.	614	Goldsmith, M.	606
615		Golgi, A.	51
Fowler, G. R.	30	Good, J. T.	672
France, O.	489	Goldstein, H. H.	331
Francis, T., Jr.	248		
Franckenstein, R.	574		

Good, R. A.	734	Harmer, T. W.	60
Goodell, H.	116	Harrison, F. C.	162
Gordon, R. E.	200	Harrison, J. W.	370
Goro, M.	267	Hart, D. S.	370
Gosselin, W.	642	Hart, F.	548
Gottlieb, P. M.	408	Hartley, F.	493
Gough, G. A. C.	215	Hartman, F. A.	189
Govaerts, P.	97		368
Gow, A. E.	83	Hartmann, F. L.	424
Grant, R.	527	Hartwell, J. L.	321
	528, 529, 572, 577,		371, 372, 374
	578, 579, 581, 643,	Hashimoto, M.	77
	644, 645, 704, 744		78
Grassellini, A.	299	Hatta, S.	531
Gray, F. P.	98		582, 583, 584, 648
Gray, J. S.	284	Hawk, W. D.	538
	302, 303	Hawkins, W. W.	410
Green, I.	513	Haworth, W. N.	274
Greene, K. V.	589	Haynes, B. W.	714
Greene, L. C.	703	Hayashi, T.	617
	711, 720	Hedenius, P.	624
Greenwood, H. H.	46	Heibert, T. G.	475
Grollman, A.	156	Heidelberger, M.	109
Gross, M.	347		134, 177, 450, 512
Grossman, M. I.	625		649
	580	Heilbrunn, L. V.	650
Guerra, F.	369	Heilmeyer, L.	651
Guerrand, S.	532	Heistø, H.	652
Guerrieri, J. A.	745	Hektoen, L.	64
Guillaumont, B.	547		228
Guthrie, T. C.	713	Hench, P. S.	190
		Hesse, E.	150
Haber, P.	169	Hestrin, S.	454
Hall, V. E.	527	Heubner, W.	202
	528, 529, 572, 579,	Heyman, A.	463,
	581, 606, 707		585, 605, 653
Hallett, S. F.	498	Hiatt, E. P.	348
Hallman, N.	711	Hibbert, H.	162
Hamel, J.	161	Hicks, M.	532
Hamilton, M. A.	484	Hill, A. G. S.	654
	519, 520	Hirsch, B. B.	645
Hammerman, S.	594	Hirsch, J. D.	644
Hanau, R.	121		645
Handman, M. S.	293	Hirsch, E. F.	99
Hankwitz, R. F. Jr.	520	Hirschmann, J.	712
	565, 566	Hjorth, P.	256
Hannan, M.	398	Hoagland, C. L.	296
Hansen, J. L.	530	Hoberman, H. D.	343
Harkavy, J.	281	Hodges, F. A.	646
Harkness, W. D.	646	Hodges, R. E.	661
	647		715

Hoff, F.	157	Kahler, H.	321
Holler, G.	71	374	
Holloman, A. L.	456	Kanthack, A. A.	17
494		Kaplan, M. H.	658
Holman, J.	586	Kass, E. H.	659
Holt, H.	532	Kaufman, G.	488
Homma, Y.	587	Kelly, K. H.	615
617		Kendall, F. E.	177
Hooper, F. E.	216	Kerby, G. P.	475
Hope, D.	392	Kibler, C. S.	72
Horsfall, F. L, Jr.	492	Kierland, R. P.	660
495, 525, 526,	576	Kiliani, M.	40
Horster, J. A.	514	Kirchmaier, H.	533
Hort, E. C.	39	Kirkendall, W. M.	661
47, 48, 49		715	
Hosoya, S.	587	Kissen, M. D.	609
617		Klendishoj, N. C.	425
Hounie, E.	499	Klotz, S. D.	473
Howard, H. G.	90	Knight, H. C.	375
91		Knoppers, A.	350
Howard, H. J.	135	Koch, P. K.	356
Howe, E. E.	655	447	
Howie, J. E.	217	Koenig, H.	563
656		691	
Hsaio-Ch'ien Chang	201	Koenig, R.	563
Humphreys, E. M.	131	Kopp, I. I.	229
Hutner, S. H.	349	351, 352	
363, 364, 373, 385,		Kornerup, T.	588
386, 387, 412, 413,		Kostyal, L.	268
414, 432, 433, 758		Kraus, R.	73
Hyde, R. R.	280	Krauss, E.	124
Hyman, M. A.	332	Kromayer, E.	178
		Krumbruch, O.	163
Iliffe, W.	11	Krusen, F. H.	550
Ivanovics, G.	282	Kulwin, M. H.	660
283		Kun, E.	497
Ivy, A. C.	284	534, 535, 536	
Izard, Y.	627	Kuna, S.	448
		Kurotchkin, T. J.	153
Jacobi, M.	241	Kutnezowsky, N.	151
Jacobs, F. A.	657	Kuzenek, A.	453
Jager, B. V.	496		
January, L. E.	661	Lackman, D. B.	272
715		Lagus, K.	711
Joachim, A.	59	Landsteiner, K.	106
Joannovics, G.	92	Lanfranchi, F.	218
Johnson, T. B.	216	Lange, F.	35
Jona, J. L.	65	Langer, A.	180
		Laurie, J. P.	457

Leavell, B. S.	532	Macfayden, A.	31
Lebrun, J.	476	Mackey, W. L.	400
477, 518, 521,	569	MacLeod, C. M.	322
Lechat, P.	699	649	
700, 701		MacPherson, C. F. C.	450
Lees, J. C.	304	Macri, B. P.	560
Leese, C. E.	589	Magoun, H. W.	289
662		Mahanna, D. L.	368
Legallais, F. Y.	467	Mahoney, V. P.	594
695		Makepeace, A. W.	155
Lehmann, H.	590	Malik, K. S.	327
Lehmann, S.	444	Malin, A.	673
Lehoult, Y.	316	Manassein, W.	7
476, 477, 518, 521		Manning, H. M.	111
Leibowitz, J.	259	Markowitz, H.	649
Leidy, G.	450	Martelli, F.	595
Leiter, J.	545	Martin, E.	672
Lequire, V. S.	523	Martin, S. P.	717
556, 557, 591,	636	Martos, K.	269
716		Masucci, P.	285
Leschke, E.	55	Matsunami, Y.	137
Lesses, M. F.	513	McBride, L. F.	72
Leudel, E.	179	McCloskey, K. L.	238
Levine, P.	203	250	
Levy, G. A.	304	McClosky, W. T.	377
Lewis, L. A.	368	383	
449, 537		McConnell, J. R.	456
Lewisohn, R.	204	498	
Liberalli, C. H.	592	McDonald, R. K.	718
Liebert, E.	593	McGinty, D. A.	596
Lilienthal, H.	136, 399	McLallen, J.	665
Limaye, M. V.	507	McManus, M. J.	416
Ling, C. Y.	110	Mearns, J. T.	459
Lipton, M. M.	753	Menczel, E.	719
Live, I.	400	Mendel, L. B.	107
Loeb, L.	37	Menkin, V.	401
Loman, J.	513	427, 597, 663	
Long, C. H. N.	608	Menten, M. L.	111
Longo, V.	195	Menzies, A. W. C.	230
Lonsen, W.	593	Meyer, J.	74
Losdan, M.	489	79	
Love, J.	426	Meyer, S. R.	428
Loving, W. L.	646	Mielnik-Kosmiderski, S.	286
Lowenstein, J.	41	Miller, C. P.	451
Ludke, H.	50	536, 538	
61		Miller, F. R.	257
Lusk, F. B.	66	Miller, J. L.	66
Lustig, B.	180	100	
		Milne, G. R.	457
		458, 459	

Mirick, G. S.	677	Nungester, W. J.	362
Mitchell, S. Q.	703	Nurolone, U.	595
720		Nussbaum, M. A.	402
Moe, G. K.	682		
Mondolfo, H.	499	Oakey, R.	456
Morell, S.	261	502	
Moreschi, C.	51	Ochoa, A. G.	503
Morgan, H. R.	287	Ohinger, B.	488
305, 323, 345, 376		Okura, Y.	125
446, 500, 539, 540		Olitzki, L.	258
598, 666		259, 326, 356, 447	
Morgan, W. T. J.	242	453, 454, 546, 601	
306, 310, 324, 353		Olson, W. H.	355
Morrison, R. L.	155	685	
Most, S.	721	Omiti, M.	308
Mueller, E. F.	138	Ott, W. H.	602
Muether, R. O.	599	733	
Muller, E. F.	84	Ottensooser, F.	379
112		388	
Müller, P. T.	42		
Munk, F.	664	Pace, B. F.	668
Murphy, J. B.	85	Page, I. H.	382
86, 87		407, 449, 537, 549	
Murray-Lyon, R. M.	191	603, 604, 722	
Myers, R. J.	370	Pages, J.	393
		Paine, T.	455
Nasmyth, P. A.	501	Palmer, J. W.	309
Nauts, H. C.	452	Palmer, L. A.	148
Neal, P. A.	354	Pansum, P. I. P.	4
Necheles, H.	355	Paris, S. K.	482
378, 600, 685		Park, H. D.	467
Nedzel, A. J.	219	Parkhurst, B. H.	619
Neerborg, G.	639	Partridge, S. M.	306
Nelson, C. M.	288	310, 324, 353	
Nesset, N. M.	665	Paschkis, K.	269
709		Pasricha, C. L.	327
Neuberger, J.	181	Paton, W. D. M.	742
Neva, F. A.	598	Patterson, J. L., Jr.	605
666		650, 653	
Nicholes, P. S.	746	Paul, B. M.	327
Nichols, F. T., Jr.	605	Peiss, C. N.	606
653		Penfold, W. J.	39
Nickerson, M.	496	47, 48, 49, 105	
Nishimura, E. T.	484	Penna, J.	73
520		Pennell, R. B.	509
Nishishita, S.	325	Perlman, E.	418
Nisi, Naoto,	307	Perrault, A.	380
Nobel, E.	43	406, 504	
Nonobe, K.	617	Petersen, W. F.	68
Noojin, R. O.	668	101, 102, 130	
Nouvelas, P. A.	192	Phillips, K.	607
		Pierce, C.	343

Pinchot, G. B.	608	Ricker, W. A.	691
Pingert, F. P.	541	Riegel, B.	709
Pinkston, J. O.	231	Rigby, G.	442
232		Ritter, W. L.	402
Piorkowski, G.	669	Rivalier, E.	141
Piotti, M.	221	142	
Pittman, M.	429	Robbins, M. E.	578
Poel, W. E.	662	579	
Powella, R.	750	Robertson, D. G.	105
Powers, J.	392	Robertson, H. F.	609
Praytor, H. B.	668	Robertson, R. C.	271
Prescott, B.	488	Robinson, E. S.	403
Pribram, B. O.	113	Robonow, C. F.	562
Price, C. W.	377,	Rodbard, S.	673
383, 432		Rodney, G.	430
Prinz, A.	103	434, 596, 674	
Probey, T. F.	423	Roe, A. S.	649
429		Rollins, J. P.	671
Promisel, E.	619	Romanoff, A.	281
Queries & Minor Notes		Rosa, M. C.	404,
J.A.M.A	637	610	
Quivy, D.	474	Rosenberg, M.	542
Rabiner, A. M.	542	Rosenthal, N.	204
Rademaker, L.	164	Rotenheim, C. A.	507
205, 505		Roussy, G.	13
Raistrick, H.	222	14	
274		Ruggiero, W. F.	251
Rall, D. P.	551	276, 278	
670		Rumpf, T.	20
Randolph, T. G.	671	Rychel, W. M.	355
Ranges, H. A.	264,	Rzucidlo, L.	611
Ranson, S. W. Jr.	270	Sabin, A. R.	182
289		Saci, H.	673
Ras, G.	186	Sack, T.	508
Rauss, K.	543	544, 613	
Reiche	126	Saether, J.	723
Reilly, J.	141	Salm, H.	247
142		Saltzstein, H. C.	329
Reimann, S. P.	456	Samelson, S.	52
Reinhard, E. H.	672	Samsøe-Jensen, T.	639
Renfrew, A. G.	216	Samwick, A. A.	332
Retes, F. P.	357	Sanders, T. E.	328
Reubi, F.	506	Sandholzer, L. A.	295
Rhines, R.	693	621	
Rhomako, M.	641	Sandweiss, D. J.	329
Rhymer, P.	88	Sansum, W. D.	81
Rice, F. A. H.	409	Sarciron, R. C.	316
Richert, D.	396	393	
Richter, P.	16	Satke, H. A.	507

Scheinman, L. J.	612	Shur, E. L.	346
Scherago, M.	398	Shwartzman, G.	152
Scherber, G.	93	183, 233, 245,	261
Scherp, H. W.	295	361, 405	
621		Silber, W.	202
Schmengler, R. D.	675	Smith, D. C.	431
Schmidt, L.	392	Smith, E. V.	290
Schneider, R.	354	Smith, H. W.	264
Schoch, A. G.	724	291, 320, 338,	420
Scholl, H.	25	Smith, J. McC.	677
Scholtz, H. G.	676	Smith, W. E.	519
Scholtz, W.	533	Smithwick, R. H.	144
Schork, P. K.	523	Smolens, J.	272
Schopp, A. C.	747	Sobotka, H.	261
Schrade, W.	358	Sokoloff, B.	750
Schrift, M. H.	238	Solari, L. A.	184
250, 251, 252,	277,	Solomon, D. H.	718
278, 339, 340,	392	Solomon, H. A.	359
Schulein, W.	10	Solomon, H. C.	127
Scott, D., Jr.	725	Somkin, E.	359
741, 748, 757		Soylemezoglu, B.	727
Scott, W. J. M.	189	728, 735	
Scully, F. J.	75	Speirer, C.	185
Sebastiani, V.	53	Spinelli, A.	165
Seibert, F. B.	107	Spitz, S.	363
108, 114, 139,	749	Spizizen, J.	317
Seligman, A. M.	508	Stacey, M.	274
544, 545, 613		Stark, C. N.	200
Seltzer, B.	515	Starka, J.	729
Senator, H.	8	Starr, M. P.	433
Senekjian, H.	253	Stetner, E.	166
279		Stetson, C. A., Jr.	730
Sevag, M. G.	272	Stokes, J. H.	140
Shafer, J. C.	431	Stuart, E. G.	678
Shapiro, C. J.	311	679, 720, 731,	732
Shaw, E. A.	567	Stubbs, E. L.	400
Shear, M. J.	236	Sturm, E.	85
243, 244, 312,	321	86, 87	
330, 371, 372,	374,	Stuttgen, G.	510
380, 381, 390,	406	680	
443, 456, 484,	498	Sugarman, M. H.	329
504, 545		Sugira, K.	363
Shelubsky, M.	454	Sutherland, A. M.	313
Shetlar, C. L.	614,	Sutter, J.	547
615		Swain, J.	26
Shetlar, M. R.	614	Sweet, B.	545
615		Swift, W. E.	452
Shimkin, M. B.	275	Swineford, O., Jr.	586
Shock, N. W.	718	681	
Shtakel' Berg, N.A.	726		

Takos, M. J.	682	von Tukacs, J.	129
Tal, C.	546	von Pap, Z.	193
683		Vos, B. J.	647
Tanji, S.	531		
582, 583, 584,	648	Waelsch, H.	207
Tardieu, G.	547	Wagner, R. R.	556
Tarr, H. L. A.	162	557, 684	
Taub, A.	548	Wagner, W.	159
Taubmann, G.	150	Wakim, K. G.	550
Taylor, R. D.	382	Walker, E. L.	293
407, 549, 604		Walker, I. C.	89
Teissier, J.	22	Walker, L.	685
Teissir, P.	141	Wallace, G. B.	182
142		Wallen, A.	392
Tennent, D. M.	733	Walter, C. W.	234
751		246, 314	
Terry, G. G.	292	Warren, M. R.	462
Thatcher, J. S.	368	Washburne, A. C.	708
Theiler, M.	127	Wauer, D.	686
Thomas, L.	734	Weaver, R. H.	398
Thompson, R. M.	752	Wechselmann, W.	45
Thompson, S. A.	206	Wechsler, Z.	262
Thomson, D.	44	Weens, H. S.	463
Tishler, M.	655	Weger, P.	511
Todd, J. P.	457	Wegscheider, K.	167
458, 459		Weil, E.	173
Tollins, J. P.	671	175	
Tomcsik, J.	128	Weinberg, J.	331
153		Weinberger, E.	207
Tonnesen, M.	460	Weiner, M.	355
Toochi, G.	524	Weinstein, J. J.	360
Topley, W. W. J.	222	Welch, H.	377
Topper, Y. J.	753	383, 432	
Tracy, M.	33	Welcke, M.	430
Tucker, J.	461	434	
Turner, F. C.	312	Wells, C. W.	80
321, 381		Wells, J. A.	551
		670, 727, 728,	735
Urbach, E.	408	Werner, H. W.	462
Urbanek, J.	158	Westfall, B. B.	464
		Westphal, K.	167
Van Winkle, W. Jr.	377	Wetterlow, L. H.	487
Vaughan, V. C. Jr.	57	Whalen, W. J.	744
67		Wharton, D. R. A.	565
Vesterdal, J.	460	616	
Villapard	435	Wieczorowski, E.	302
Voloshin, T.	622	303	
von Euler, U.	143	Wien, K.	273
von Jauregg, J. Wagner	12	Wiener, A. S.	332
54		Wilcox, H. H.	692,
		693	
		Wilson, J. L.	445

Wilson, J.	297
301	
Wilson, M. L.	596
Wilson, W. L.	650
Windle, W. F.	481
563, 687, 688, 689,	
690, 691, 692, 693,	
703, 704, 736, 741,	
754, 755, 756, 757	
Winkelstein, A.	361
Wirth, F.	694
Witebsky, E.	247
425	
Wittich, F. W.	737
Wolff, H. G.	313
Wolff, L. K.	186
Woodyatt, R. T.	81
Wretlind, K. A. J.	465
Wright, A. M.	341
Wright, I. S.	223
333	
Yagi, Y.	587
617	
Yamamoto, K.	334
Yates, A. L.	238
250, 252, 276	
Young, E. G.	409
410	
Youngner, J. S.	362
Yu, H.	271
Zahl, P. A.	349
363, 364, 373, 384	
385, 386, 387, 411	
412, 413, 414, 433,	
758	
Zeckwer, I. T.	116
132	
Zimmer, A.	115
Zindler, G. A.	738
Zinov'eva, M. T.	466
Zittle, C. A.	434
Zozaya, J.	187
194	
Zykina, E. S.	739

A D D E N D U M

1 9 5 1

759. BENNETT, I. L., JR.

Comparison of leukocyte changes produced by pyrogens and by anaphylaxis in the guinea pig

Proc. Soc. Exp. Biol. and Med. 77:772-775, 1951

Hematologic response to pyrogens differs from the leukopenia and leukocytosis which follows injection of an antigen into sensitized animals. Circulating eosinophils rise after anaphylaxis and decline uniformly in the reaction elicited by pyrogens. Pyrogen contamination of antigenic substances is not an important factor in the production of anaphylactic shock.

760. BUSCAINO, V. M.

L'iperpiretoterapia della schizofrenia (Fever therapy in schizophrenia)

J. brasil Psiquiat, 1:154-161, 1951

Typhoid-paratyphoid vaccine was administered to schizophrenic patients for prolonged periods, accompanied by medicaments which afforded protection to hepatic tissues. More promising results were obtained than could be elicited by 'shock' treatment. The hypothesis is presented that schizoid behavior follows the production of lesions in cerebral cortex and mesodermal tissues. Such lesions may be secondary to the release of toxins by the liver or mucosa, with failure of detoxification through the liver and reticuloendothelial system.

761. HIRSCHMANN, J.

Konstitutionelle Unterschiede von Temperatur-und Leukocytenkurven unter Pyriferbelastung (Constitutional differences in temperature and leucocyte responses to 'pyrifer')

Nervenarzt 22:260-264, 1951

When individuals with varying body structural characteristics received a small injection of pyrifer (10 units), those with high sympathetic tonus, but smaller potential differences (ie., the pyknic individual), showed small differences in temperature, while the athletic and the

1 9 5 1

asthenic exhibited a tendency toward greater variations in temperature response. Irrespective of body type, the allergic patient showed wide reactive variations. Hormonal imbalance also influenced the range of temperature variation; this was noted particularly at the climacteric period. Temperature curves after the administration of pyrifera were evaluated in the patient with multiple sclerosis.

762. KERBY, G. P., and BENNETT, I. L., Jr.

Splanchnic removal of bacteria from blood stream of pyrogen-tolerant rabbits

Proc. Soc. Exper. Biol. and Med. 78:48-50, 1951

Splanchnic removal of bacteria from the blood stream of animals with established tolerance to pyrogens differed little from that observed in control animals. No significant alteration in the rate of removal for M. aureus or Klebsiella, pneumoniae type B, could be noted when continuous infusion of the organisms had been preceded by injections of Pv. Piromen, the P25 polysaccharide of Shear, or triple typhoid vaccine.

763. LANKFORD, C. E., HOYO, H., and LUTTERINGER, J. T.

Intracellular polysaccharide of Enterobacteriaceae

J. Bact. 62:621-626, 1951

Deeply staining polar bodies were seen in the cytoplasm of "smooth" strains of Shigella, Salmonella, Escherichia, Aerobacter, Klebsiella and Serratia marcescens, but not in so-called "rough" varieties. Such polar bodies may be the sites of the polysaccharide haptens of the O antigen.

764. LASFARGUES, E. Y. L., WHARTON, D. R. A. and DIFINE, J. C.

Some effects of polysaccharide preparations from Serratia marcescens and Aerobacter aerogenes on cells in tissue culture

Cancer Res. 11:425-427, 1951

No evidences of damage to tissue cultures of chick heart fibroblasts or mouse sarcoma 37 cells could be noted when preparations of polysaccharide materials from S. marcescens and A. aerogenes were added to the nutrient solutions. Some evidences of a stimulating effect on tissue growth could be noted. This may be caused by an enzymatic breakdown of the polysaccharide into simple sugars which supply energy for tissue functions. A detoxifying mechanism may be involved. The hemorrhagic necrotic action of bacterial polysaccharides may be one of anoxia after vascular damage occurring during shock.

1951

765. ROSENZWEIG, A.

Bactifebrin therapy of acute encephalo-radiculo-neuritis and Landry-Guillain-Barre syndrome, poliomyelitis acuta anterior and herpes zoster

Confin. neurol. (Basle) 11:227-240, 1951 (In English)

Prompt improvement followed the use of bactifebrin in Guillain-Barre syndrome (22 cases), lymphocytic meningitis (17 cases) and herpes zoster (17 cases). In acute paralytic poliomyelitis improvement followed closely after the injection, enduring some 2 to 4 hours. Pain was localized in paralyzed muscles during the period of activity. The preparation is therapeutically effective only when the lesions are situated in the proximal intrameningeal portion of the peripheral neuron. Experimental infections with "Tesch disease" (meningo-encephalomyelitis ensoötica suum) and rabies in rabbit, guinea pig and mouse were not favorably altered by the use of bactifebrin. Hyperpyrexia production did not influence results obtained.

766. SCHELLONG, F., and SOESTMEYER, T.

Herdsuche durch Pyrififer (Search for infectious foci by use of pyrififer)

Deutsch. med. Wchnschr. 76:761-763, 1951

Marked pain in infected internal organs was elicited by the intravenous administration of 0.3 cc. of 15E pyrififer. Localized pain, of a "grinding" or "boring" nature, usually appeared within 15 hours after administration, coinciding with the maximal temperature peak. The reliability of the test was determined in suspected gallbladder and duodenal disease, in appendicitis, renal and gynecologic conditions and in laryngeal and dental infections. The localized pain elicited was not related to the Schwartzman-Sanarelli phenomenon, but was considered to be an intense neuro-vegetative stimulation in inflamed areas, with a marked reduction of the threshold for pain. While the method is contradicted in chronic inflammation (as in tuberculosis), its value for differential diagnosis in acute infection appears to be established.

767. STAUDER, K. H.

Über die Dekompensation alter Hirnherde durch Infektionskrankheiten
(On decompensation of old brain lesions through infections)

Arch. Psychiat. Nerven. 187:165-176, 1951

Infectious disease, but not nonspecific stimuli (pyrififer and other pyrogenic substances) reactivated latent neurological syndromes (extrapyramidal and jacksonian symptoms, hemianopsia, orbital muscle disorders).

768. VON WOLFFERSDORF, H.

Pyrifer-behandlung von verschiedenen Erkrankungen des Sehnerven von 1939 bis 1950 unter besonderer Beachtung von Makula-Erkrankungen bei akuter retrobulbärer Neuritis (Treatment of various diseases of the optic nerve with pyrifer from 1939 till 1950, with particular attention to macular diseases in acute retrobulbar neuritis)

Klin. Mbl. Augenheilk. 119:271-279, 1951

When pyrifer was employed in retrobulbar neuritis, papillitis, postneuritic atrophy, methyl alcohol poisoning and kindred conditions, function was restored in 80% of the cases treated (107 patients).

769. YUKIMURA, T. and TAKEDA, S.

Relation between the pyrogen test and paper partition chromatography of protein hydrolyzates

Igaku to Seibutsugaku 19:187-190, 1951

A common peptide was found in pyrogen-positive casein hydrolyzates when paper partition chromatographic methods were applied. Pyrogenous hydrolyzates may produce fevers which are not symptomatically identical.

1 9 5 2

770. ALGIRE, G. H., LEGALLAIS, F. Y. and ANDERSON, B. F.

Vascular reactions of normal and malignant tissues in vivo. V. The role of hypotension in the action of a bacterial polysaccharide on tumors

J. Nat. Cancer Inst. 12:1279-1291, 1952

Peripheral or intratumoral circulation restriction by mechanical means (occlusion) or biochemical substances (histamine, bacterial carbohydrate complexes), is followed by damage or regression in tumor tissue. Retarded blood flow and resultant stasis and tissue anoxia, and not a direct cytotoxic effect, underlie these tissue changes.

In mice bearing rapidly growing strain L and sarcoma 37 implants, hypotension was produced in neoplastic tissue by single or repeated injections of bacterial polysaccharides obtained from S. marcescens culture filtrates (P-10, P-20 and P-25 of Shear). Systolic pressure was observed microscopically in arteries adjacent to tumors and capillary changes within tumor tissue were noted. In vivo criteria of tumor damage were supplemented by histologic studies of excised tissue. Hypotension, in both degree and duration, correlates to the dosage administered to normal or tumor-bearing animals. No vascular changes were observed in the absence of hypotension. The functional capillaries within the tumor were reduced in extent after the initial dose of the bacterial polysaccharide (by approximately 80%). Circulation was retarded within the first hour following administration. White thrombi were observed after 5 hours. In 24 hours, the center of the tumor was opaque and necrosis followed within 3 days. The most pronounced and prolonged fall in blood pressure was produced by the initial injection of the polysaccharide; succeeding injections may provoke resistance to the tumor-necrotizing effects.

771. BAILEY, A. A., ROOKE, E. D. and RODIN, E. A.

Investigation of a bacterial pyrogen as a therapeutic agent in neurologic disorders

Proc. Staff Meet. Mayo Clinic 27:340-343, 1952

PIROMEN was employed as a therapeutic agent for 57 patients exhibiting disease entities in which it was desirable to inhibit the formation of scar tissue and to promote the regeneration of neurons and axones. The product was given in dosages ranging between 2 and 9 micrograms, for limited periods, in the treatment of amyotrophic and multiple sclerosis, traumatic and retrobulbar neuritis, neuronitis, traumatic paraplegia, arachnoiditis and central nervous system involvements of indeterminate types. Its effect on residual evidences of poliomyelitis was also determined.

1 9 5 2

Some improvement was noted in 4 of 22 patients with multiple sclerosis; spasticity appeared to be benefited in late stages of the disease; such improvement may have a subjective basis. The dosage of PIROMEN may be a determining factor in these conditions. Work, still unpublished, would indicate a possible therapeutic role for the drug in the treatment of encephalitis of allergic origin if results of experiments now in progress confirm early observations. At this time, much more study of the efficacy of this preparation is needed before its value in treatment of crippling neurologic disease can be evaluated.

772. FRACCHIA, A. A. and BRUNSCHWIG, A.

Hyperthermia (114°F Rectal) with recovery

J.A.M.A. 149:926-927, 1952

Intravenous injection of Coley's toxin (0.12 ml. to 1.48 ml. daily for a ten-day period) produced symptoms of mental confusion, facial congestion and cardiac arrhythmia, accompanying a temperature reading (rectal) of 114°F. Administration of antipyretics, 5% dextrose solution and other temperature-reducing measures brought a rapid drop to 104° F. within 15 minutes. The patient exhibited no ill effects from this brief period of hyperpyrexia. Inactivity of an extensive ovarian carcinoma for 17 years, after surgical procedures and Coley's toxins had been employed, is confirmed.

773. GINGER, L. G. and WINDLE, W. F.

Studies on the chemistry and physiological action of bacterial pyrogens

Second International Congress of Biochemistry, Paris, July 21-27, 1952.
Resumé of Communications, Sec. 28, pp. 445-446.

Significant differences may exist between pyrogens obtained from different bacterial species. Minimum pyrogenic doses from nonprotein concentrates prepared from B. subtilis, S. marcescens, a Pseudomonas species, E. coli, Proteus vulgaris and S. typhi, ranged, in the rabbit, between 7.5 and 0.22 gamma, per Kg. body weight, and were unrelated to the polysaccharide content. A new fraction from Pseudomonas species was a highly pyrogenic, electrophoretically homogeneous substance (PC-4) which may represent the pyrogenic factor. It appears to be a stable polysaccharide-amino acid-lipid complex, in which the polysaccharide component consists of hexosamine, an unclassified reducing sugar and a nonreducing carbohydrate. PIROMEN toxicity is slight (LD₅₀ 60,000 in the mouse) and dosages above 1000 gamma have been given to animals without toxic manifestations. At dosages, for man and animals, of 0.05 - 0.5 gammas/kg., early granulocytosis, hypertrophy of the thymus, and lymphatic system, release of thyroid colloid and adrenocortical secretions followed within 2 to 5 days. Chronic adrenalectomy abolished the leukocyte response and the latter could be revived by adrenal cortex transplantation. Burn healing was enhanced, with natural debridement and edema reduction. PIROMEN prevented formation of pia-glial scars in spinal cords after resection and permitted neuronal regeneration.

1952

774. NEWELL, F. W.

Fever therapy in ophthalmology today

Qtly. Bull. Northwestern U. M. School 26:151-156, 1952

Dramatic response of inflamed tissues of the eye follows hyperpyrexia, induced by either passive agents, or by parenteral injection of pyrogenic substances. The present status of fever therapy in ophthalmology is re-evaluated with a view to clarifying the roles of heat, immunologic response or "trigger" activity on the adrenocortical system, in the reduction of ocular inflammation. If the initiation of stress underlies the beneficial effects secured through the use of bacterial pyrogens, it is believed that such results can be most easily obtained by administration of ACTH and cortisone. The use of bacterial pyrogens is well-justified, however, if it is confirmed that other important activities follow their application in ophthalmology.

775. PEARLMAN, M. D.

Management of endogenous uveitis

Eye, Ear, Nose and Throat Month. 31:202-206, 1952

Nonspecific fever therapy has been effectively used in the past for the therapy of inflammatory ocular disease. Typhoid vaccine and the bacterial extract PIROMEN are the most effective pyrogenic agents now in use. "Milk and its protein derivatives, proteolac and omnadin, are less effective." The mode of action of these pyrogenic substances has not yet been definitely ascertained; the most recent theory presented would assign a stress-producing activity to these materials "thereby stimulating an ACTH and cortisone response."

776. POEL, W. E. and BELKIN, M.

Variations in eosinophil morphology and leukocyte counts in the rat following injection of a tumor-necrotizing bacterial polysaccharide

Am. J. Physiol. 169:453-459, 1952

Leukocytosis and lymphopenia which follow the injection of bacterial polysaccharides suggest adrenal cortical activation. To observe more fully the changes in circulating leukocytes, particularly alterations in the eosinophil ratios, the polysaccharide of Shear (P-25) was injected in the rat in dosages equivalent to 3 gammas per Gm. body weight. Changes in the hematologic picture were compared with those seen after the injection of saline solution. Leukopenia of both neutrophils and lymphocytes appeared in the first hour; leukocytosis followed. The elevation

1952

in leukocytes persisted approximately 24 hours. Eosinophil levels followed a "W" pattern when plotted graphically, peaks corresponding to 4, 6 and 12 hours after injection. Eosinophil structure was altered; fragmented, pyknotic and granulocytic cells were seen. At the 6 hour "peak" the transient recovery observed appeared to correspond either to an exhaustion or a mobilization phase; marked eosinopenia followed such recovery. Adrenocortical or unknown factors may initiate alterations.

777. SAMTER, M. and KOFOED, M. A.

On the rationale of treating allergic diseases with bacterial pyrogens

J. Allergy 23:327-334, 1952

In comparison with a placebo, the clinical effectiveness of PIROMEN, when given in a limited dosage range, for the therapy of anaphylactic allergies (rhinitis, bronchial asthma, and urticaria) was no greater than a corresponding amount of its solvent. The compound was found to have low antigenicity for man; it affords no demonstrable inhibiting effect on the formation and distribution of anaphylactic antibodies in the experimental animal.

778. SCOTT, D., Jr. and CLEMENTE, C.

Conduction of nerve impulses in the regenerated fibers of the spinal cord of the cat

A.M.A. Arch. Neurol. and Psychiat. 67:830, 1952

Adult cats were permitted to recover without treatment following resection of the spinal cord or were treated with PIROMEN for periods ranging from 7 to 17 months. Electrophysiological studies were confirmed by histologic examinations. Stimulation of the pyramidal tract fibers above the site of the lesion evoked nerve impulses below the site of resection after treatment with the bacterial polysaccharide; the two factors which affected the extent of regeneration were (1) administration of PIROMEN and (2) close proximation of the proximal and distal portions of the cord after section. Corticotrophin appears to have similar activity while cortisone blocks its effectiveness.

779. WALTON, C. H. A. and ELLIOTT, G. B.

Sudden death from bronchial asthma following injection of PIROMEN

J. Allergy 23:322-326, 1952

In an asthmatic female, age 31, with a stormy history of severe paroxysmal

1 9 5 2

asthma complicated by hypertension, toxemia and transient psychotic episodes, control of asthmatic symptoms had been possible through the use of epinephrine, isuprel, aminophyllin and ACTH. PIROMEN was administered for one week before death, producing only slight shivering in dosages ranging between 1 and 5 gammas. Temperature elevation was registered after the first 3 injections only. No immediate reaction occurred after the last injection of PIROMEN, but asthmatic symptoms were not benefited. Dyspnea increased, epinephrine in repeated dosage brought no relief. Deep cyanosis and death followed. On autopsy, intense bronchial and bronchiolar obstruction, due to mucus plugging and edema, was seen. Bronchopulmonary lymph nodes were enlarged, but all others appeared normal in size. A persistent thymus was found. The lungs were crepitant. Congestion of both brain and pituitary was seen. It was not possible to link the administration of the medication (PIROMEN) with the terminal asphyxia. "Asthmatic patients of this type have a notoriously bad prognosis and often require very little to upset their delicately balanced respiratory state..."

AUTHOR INDEX (TO ADDENDUM)

Algire, G. H.	770	Legallais, F. Y.	770
Anderson, B. F.	770	Lutteringer, J. T.	763
Bailey, A. A.	771	Newell, F. W.	774
Belkin, M.	776	Pearlman, M. D.	775
Bennett, I. L., Jr.	759	Poel, W. E.	776
762		Rodin, E. A.	771
Brunschwig, A.	772	Rooke, E. D.	771
Buscaino, V. M.	760	Rosenzweig, A.	765
Clemente, C.	778	Samter, M.	777
Difine, J. C.	764	Schellong, F.	766
Elliott, G. B.	779	Scott, D., Jr.	778
Fracchia, A. A.	772	Soestmeyer, T.	766
Ginger, L. G.	773	Stauder, K. H.	767
Hirschmann, J.	761	Takeda, S.	769
Hoyo, H.	763	Von Wolffersdorf, H.	768
Kerby, G. P.	762	Walton, C. H. A.	779
Kofoed, M. A.	777	Wharton, D. R. A.	764
Lankford, C. E.	763	Windle, W. F.	773
Lasfargues, E. Y. L.	764	Yukimura, T.	769

SUBJECT INDEX

- ABDERHALDEN REACTION, in fever, 179
- ABORTION, infectious, effect of filtrate of B. abortus on leukocyte picture in cattle, 400
- ACACIA, bacteria grown in, prodn. of pyrogen, 278
colloids, inactivation of coliphage, 317
polysaccharide, effect upon bacteriophagy, 295
- ACETANILID, evaluation against pyrogen-induced fever, 462
- ACETIC ACID, pptn. of nucleoprotein with, 586
- ACETONE, pptn. of antigenic nucleoprotein from Ps. aeruginosa with, 587
- ACETOPHENETIDIN, evaluation against pyrogen-induced fever, 462
- ACETYLMETHYLAMINE, leukocytosis in hypertensive dogs by, 407
- ACETYLSALICYLIC ACID, 661, 685
dosage, to counteract fever from bacterial polysaccharide, 438
effect, before and after inj. of Pv Pyromen, 715
on blood and plasma changes in fevers, 117
evaluation against pyrogen-induced fever, 462
mechanism of antipyresis in monkeys, 369
- ACID, treatment of parenteral solns. for reducing pyrogens, 541
- ACIDOSIS, influence on prodn. of agglutinins, 169
- ACRIDINE, toxicity, adrenocortical factors do not protect mice against, 622
- ACTH: See Pituitary Preparations
- ADRENALIN, effects, on blood pressure ineffective for respiration after toxic doses of polysaccharide, 394
effects, on blood sugar during induced hyperthermia, 299
on hypophysectomized rabbits, 307
in blood: See Blood, adrenalin
mechanism of lymphocyte lesions provoked by, 569
- ADRENALS, and neutrophil leukocytosis, 146
ascorbic acid, decrease from action of dysentery toxin, 447
changes in rats by infection with E.coli, 608
induced by bacterial pyrogen, 687
cortex, effect of typhoid vaccine on, 619
extract, abolition of pyrogenic reactions with cortisone, 705, 778
changes in tumor-bearing mice injected with, 486, 522
connective tissue mast cell responses to cortisone, 732
dehydrocorticosterone acetate, 537
dosages in rats infected with B. tularensis, 608
effect of cortisone on Arthus phenomenon in rabbits, 731
effect of cortisone on regeneration of peripheral nerve fibers, 740
necrosis of kidneys in cortisone-treated rabbits after inj. of toxins, 734
post-injection exhaustion in metastatic carcinoma, prevented or reduced with, 508
prevents toxic effect of endotoxins, 667
protection of adrenalectomized animals against bacterial intoxication, 189

- ADRENAL
cortex
extract
 reducing substances, indicators of protection by cortisone against tumor-damaging S. marcescens polysaccharide, 697
function, impaired by non-specific and inflammatory reactions, may reduce arterial pressure, 407
hormone (synthetic): See 11-Dehydro-17-hydroxy-corticosterone; Desoxycorticosterone
 role in therapy of diseases of eyes with pyrogens, 620
effect of bacterial polysaccharide and adrenal cortex ext.on, 486
 on blood sugar levels, 132
excision, changes in leukocyte level of rats after typhoid vaccine, 537
effects of fever from typhoid-paratyphoid vaccine in rats, 318
effects on pyrogen fever in rabbits, 644, 645
in animals, protection against bacterial intoxication by ext. of adrenal cortex, 189
in dog, comparison of leukocyte response to ACTH and bacterial pyrogen (Pyromen), 728
inability of desoxycorticosterone acetate to protect rat against typhoid vaccine, 300
protection of rats against bacterial intoxication by ext. of adrenal cortex, 189
extract: See also Adrenals, cortex, extract
 beef, (Upjohn) reduction in toxicity of S. marcescens polysaccharide in tumor-bearing mice, 469
 in fever therapy, 368
 method of assaying for protective action against toxic material (typhoid vaccine), 449
hemorrhagic effects of bacterial vaccines, 356
hormones, influence on toxic and tumor-damaging effects of certain substances, 622
importance in fever, 145
inactivation in experimental fever, 231, 232
influence of fever on reaction curve of, 179
mechanism of leukocyte response to bacterial pyrogen (Pyromen), 720, 727
physiology, altered in febrile conditions, Abderhalden reaction, 179
 effect of dibenamine on vascular response of rabbits to typhoid vaccine, 627
plasma, in mechanism of response to pyrogen therapy of arterial hypertension, 603
stimulation of pituitary-adrenal system in Pyromen therapy of allergy, 671
- ADSORPTION: See Filtration, adsorptive
- AEROBACTER cloacae: See Bacteria
- AFRICA, north, gonorrhea in, 391
- AGE, factor in renal hemodynamic changes induced by standardized pyrogen, 718
- AGGLUTINATION, autoagglutination, between plasma cells and serum in plasmacytoma, 514
bacterial, role of thrombocytes in, 118
hemagglutination, cause of reactions from transfusion of unpooled plasma, 360
research in changing picture in animals previously treated with colon bacilli, 62
- AGGLUTININS, "O", demonstrable only for immunologically related antigens, 539
production, influence of alkalosis, acidosis and thermal reaction (vaccine) on, 169

- AGRAULOCYTOSIS, therapy, protection against radiation by lipo-protein-nucleic acid complex (reticulose), 752
- AIR, injection, 112
- ALBUMINS, suspension used in therapy of schizophrenia (pyrifer), 227
cleavage product, albumose, in therapy of abdominal typhus, 61
carried pyrogenic characteristics of bacterial proteins, 58
connective tissue mast cell responses to ovalbumin, 732
production of fever with, in rabbits, 107
- ALBUMOSE, **deutero-**, in therapy of typhoid, 61, 71
therapy of arthritis, 94
- ALCOHOLISM, foreign protein therapy may be dangerous in acute alcoholics, 100
- ALKALI, conversion of polysaccharide into haptens by, 215
effect on leukocytes, 137
in degradation of antigenic material from *B. dysenteriae*, 324
precipitation, in method for sepn. of hemorrhage-prod. fraction of *B. coli*, 244
reserve, in blood: See Blood, alkali reserve
solutions, increase potency of Schwartzman-active nucleoprotein fraction of bacteria, 223
treatment of parenteral solns. for reducing pyrogens, 541
- ALKALOSIS, influence on prodn. of agglutinins, 169
- ALLANTOIC FLUID, viruses of mumps, influenza and Newcastle disease cultivated in, 525
- ALLANTOIN, in Ringer's soln., role of pyrogen in alleged leukocytic response, 410
- ALLERGY: See also Anaphylaxis; Asthma; Food, allergy
bacterial: See also under names of bacteria as, *Pneumococcus* prepn. of crude nucleoprotein and polysaccharide fractions of bacteria, 586
pyrifer as antigen in picture of autoallergy of bacterial cell, 680
Pyromen, for anaphylactoid, 777
cause, of hepatopathies, 686
of primary facial paralysis, 595
of reactions from transfusion of unpooled plasma, 360
damage to visual organs from typhoid vaccine therapy, 588
reactions to whole blood and plasma transfusions in multiple sclerosis, 513
salutary effect of febrile disease on, known for centuries, 637
therapy, effect of fever shock in, 675
of asthma, acne, and chronic eczematoid dermatitis with Pyromen, 708, 777
of perennial symptoms with Pyromen, 671
- ALUMINUM HYDROXIDE, elimination of pyrogens with, 592
suspension, in inj. of polysaccharide for mice tumors, 406
- AMIDOPYRINE, premedication, admin. through stomach tube prevented febrile response to typhoid vaccine, 470
premedication, in pyrogenic reaction in hypertension, 338, 420
- AMINES, toxins, (from purified meat) pyrexia by, 466
- AMINO ACIDS, content of endotoxin of *S. enteritidis*, 617

AMINO ACIDS

oral admin. of casein hydrolysates in infants produced fever, 445
synthesis in media, influence on pyrogen prodn., 218

AMINO (NH₂) COMPOUNDS, crystalline, purified products from bacterial filtrates yield, 176

AMINOPYRENE, evaluation against pyrogen-induced fever, 462

AMINOSOL: See Casein

AMMONIA, and fraction increased in pyrogenic distilled water, 164

AMOBARBITAL: See Barbitol Derivatives

AMPUTATION STUMPS, recurrence of sarcoma in, treated with mixed toxins (Coley's), 148

ANAPHYLAXIS: See also Allergy
attempts to produce, with polysaccharide from S. marcescens in guinea pigs, 395
bacterial, pyrogenic action of typhoid anaphylatoxin, 51
—— pyrogenic contamination not factor in, 759
role of carbohydrate haptens in, 153
typical anaphylactoid reactions after inj. of pyrogens in distilled water, 114
eosinophilia during, 759
fatal reaction (Sanarelli-Shwartzman) after therapeutic anti-typhoid injections, 426
therapy, Pyromen not effective for, 777

ANESTHESIA, ether, effect in cats administered typhoid-paratyphoid vaccine, 270

ANIMALS: See also Cats; Dogs, Guinea Pigs; Mice, Rabbits, Rats; etc.
experimental, action of dysentery toxins on, 447
fever in, 231, 232

fever reaction in relation to inj. sites of pyrogens, 726
hypothermic and adreno-hemorrhagic effects of bacterial vaccines, 356
infection, 62
influence of pyrifur fevers on plasma barriers, 262
prodn. of antibodies in, criterion for toxicity of pyrogen, 340
sensitivity to Pyromen, 691
variations in body temperatures, relation to pyrogens, 404
sarcoma, effect of Shwartzman reaction with E. typhosa filtrate on, 241
sensitized, excessive sensitivity of heat centers in, 77
toxins, in therapy of malignant tumors, 30
tumors, reaction to bacterial toxins in animals susceptible to Shwartzman phenomenon, 198
vasomotor disturbances in intoxicated animals, mode of action of endotoxins, 521

ANOXIA: See Oxygen, deficiency

ANTHRAX, bacilli, fate of capsular substances in organism, 283
bacilli, fever toxins of, 18
filtrates, insulin-like substances from, 154
polysaccharides of, 282

ANTIBIOTICS, action of B. subtilis and B. mesentericus and their secretions on tubercle bacillus, 119
filtrates of Streptococci, Staphylococci, E. typhosa, and B. coli related to, 176

ANTIBODIES, contg. fractions of S. marcescens polysaccharides, influence on lethal and tumor-necrotizing actions, 520
anaphylactic, not inhibited by Pyromen, 777
formation, carbohydrates adsorbed on colloids as antigens, 194

ANTIBODIES

formation

- close relationship between site of antigen activity and, 480
- impaired in radiation injury, 752
- in exptl. animals, criterion for toxicity of pyrogens, 340
- in mice, activity of purified O antigen, 385
- in protein therapy, 102
- in therapy with typhoid vaccine and anti-leukopenic tolerance of rabbits, 601
- increased in non-specific protein therapy, 101
- prevented through action of salicylates, 496
- homologous, labeled with fluorescein, fate of polysaccharide of Friedlander's bacillus, studied with, 654
- in blood, influence of various agents on mobilization, 110
- mechanism, stimulated by specific vaccines, 64
- nature of chill-producing principle in antipneumococcus serum, 182
- polysaccharide released around bacteria by, 286
- response and antigenic action of polysaccharides from S.marcescens, 616

ANTIGENS; See also Antibodies; complement, fixation; Immunity; Polysaccharides; Precipitin, reaction; etc.

- bacterial, fall and rise of phosphorus level of blood after injection of, 259
- heat-killed suspensions, leukopenic action, 326
- hyper- and hypoglycemia after inj., 259
- leukopenic action, 326
- pyrifer as antigen in picture of autoallergy of bacterial cell, 680
- Pyromen, non-antigenic, 777

therapy of asthma, critical review, 681

- B.enteritidis, soluble precipitate chiefly polysaccharide, 131
- B. shigae, polysaccharide failed to induce immune bodies, 310
- carbohydrates adsorbed on colloids as, 194
- cocci, polysaccharide complex from, 285
- colon-typhoid-dysentery group, attributable to polysaccharide and nitrogen complex, 667
- Ducrey's bacillus, Streptobacillin, soluble prepn. of, effects in man and animal, 141
- E. coli, Flexner, paratyphoid B and Shiga produced leukocyte migration, 316
- leukopenic ability, 367
- produced leukocyte migration, 316
- E. typhosa, acute arthritis produced by intra-articular injection of, 500
- and B. paratyphosus B, ext. of culture, 106
- cultured in synthetic medium, 305
- effects produced by inj., 345
- glucido-lipid chemotaxis of leukocytes, 316
- glucido-lipid, leukopenic ability, 366
- glucido-lipido-polypeptide, inj. into rabbit, vascular reactions of skin, 518
- hydrolysis of, 319
- hyperplasia after injury to hemopoietic tissue, 376
- immunologic properties of, 323
- induced fever and leukopenia in rabbit, 305
- injected intrav. produced fever and leukocytosis, 446
- injury to hemopoietic system by, 376

- leukopenic ability, 367
- O antigen complex of, 353
- pathologic changes in rabbit by 376
- polysaccharide, component of, 319, 334, 353
 - _____ toxicity, 353
- produced leukocyte migration, 316
- properties of, 323
- simple method for prep., 309
- synthetic medium in isolation of, 301
- tolerance to toxic effects, 539, 540, 598
- toxic antigens, effects of intrav. inj. on, 345
- glucido-lipid, and phagocytic ability of reticuloendothelial cells, 393
- chemotactic power of, 316
- leukopenic ability, 366, 367
- responsible for leukocyte response, 342
- glucido-lipido-polypeptide, (typhoid endotoxin) vascular reactions of skin after inj. in rabbit, 518
- gram-negative bacteria, common toxic component characterize all, 387
 - relationship of tumor-hemorrhagic agent to, 363
- labile, isolated from hemolytic streptococci, Lancefield group A, 272
- local hemorrhagic-necrotic skin reactions, 281
- localization in tissue cells, 654, 658
- mycobacteria, leukopenic action, 326
- nature of action and antibody response in mice of polysaccharides from S. marcescens, 616
- (nucleoprotein), isolation from Ps. aeruginosa, 587
- O, activity, lethality, hemorrhage-prodn. and antibodies prodn. in mice, 385
- complex, 412
- complex of E. typhosa, 353
- Salmonella endotoxin component of, 364
- tumor-hemorrhage toxins in characteristics of gram-negative organisms, 373, 763
- Pneumococcus, distr. of polysaccharides in tissue of mouse after inj. of large dose, 486
- localization in tissue cells of types II and III, 658
- polysaccharide complex from cocci, 285
 - from S. marcescens, nature of action and antibody response in mice, 616
 - specific, injected in mice produce immunologic paralysis extending through normal life, 573
- Pseudomonas, isolation, 578
- pyrifer as, in picture of autoallergy of bacterial cell, 680
- Salmonella, a labile polysaccharide, 215
 - paratyphosus, S. schottmülleri, tolerance to toxic effects, 539, 540
 - typhimurium, synthetic medium in isolation, 301
 - vasomotor disturbances in animals, 521
- Shigella dysenteriae, 559
 - E. typhosa antigenic protein similar to B. Shigae, 353
 - endotoxin, tumor-hemorrhagic agent, 363
 - fractionation and chemistry of material isolated from, 306
 - immunization experiments with artificial complexes from, 310
 - phenol and alkali in degradation of, 324
 - polysaccharide, component of, 306, 324, 363
 - _____ labile, in 215
 - produced leukocyte migration, 316
 - protection of rabbits with whole and fractions of, 560

Shigella dysenteriae,

- tolerance to toxic effects of, 539, 540, 598, 666
- toxic and antigenic properties of fractions from, 546
- type Z (Flexner), chemical degradation of, 418
- type Z (Flexner), nature of toxic component of, 683
- significance in animal tissues, 573
- specific, studies of fever in sensitized animals, 77
- toxic, reticuloendothelial cells reacting to, 480
- typhoid, antigenic factor of typhoid vaccine, 56
- tolerance to action of endotoxins after typhoid and paratyphoid fevers, 666
- tolerance to toxic effects, in convalescents, 598

ANTI-HISTAMINES, 661

- do not reduce toxicity of polysaccharides of S. marcescens in mice, 642

ANTI-HORMONE: See Enterogastrone

ANTI-PYRETICS: See also Fever; Pyrogens; under names of drugs as, Acetanilid; Acetophenetidin; Acetylsalicylic Acid; Aminopyrene

- abolition of pyrogenic reactions with, 705
- depression of gastric secretions with, without fever, 685
- did not alter course of tolerance to bacterial pyrogens, 470
- effect, modified by heat or cold application to heat regulating centers, 78
- testing, use of hay exts. infusions in rabbits, 202

APPARATUS, for intravenous and subcutaneous use, 206

- standardized, for prepn. of non-pyrogenic fluids, 341

APPETITE, in fever, 74

ARABINOSE, d-, in fraction from culture of tubercle bacilli, 216

ARGININE, toxic polypeptide gives color reaction for. 363

ARSEPHEMINAMINE, therapy in neuro-syphilis, 140

- toxicity, intravenous salvarsan inj. without reactions, 45
- relation to other forms of injection fever, 49

ARTERIES, disease: See Thromboangiitis obliterans

ARTERIOSCLEROSIS, altered response to typhoid vaccine during massive salicylate therapy, 496

- in arterial hypertension, therapy with pyrogen (Pyromen), 603
- therapy, with foreign protein (typhoid vaccine), 144

ARTHRITIS: See also Osteoarthritis; Rheumatism

- acute, produced by intra-articular inj. of typhoid antigen, 500
- rheumatic polyarthritis, fever shock therapy in, 675
- chronic, therapy, method for controlled hyperpyrexia, 359
- effect of artificially induced fever on circulation, 550
- in children, action of bacterial protein, question of herpes, 166
- therapy, by intravenous inj. of foreign protein (typhoid vaccine), 66
- subpyrexial doses of streptococcus vaccine, 212
- with nonspecific protein (albumose, horse serum, bacterial proteins), 94, 191
- with typhoid vaccine, 75, 82, 191

- ARTHRITIS, RHEUMATOID, chronic,
effect of ACTH on induced
fever, 659
therapy, febrile reaction after
infusion of blood from preg-
nant women, 468
induced fever by intravenous
typhoid vaccine, 313
remission after fever therapy
(typhoid bacteria), with li-
ver damage, 696
typhoid shock, 461
- ARTHUS PHENOMENON, 681
comparison of Shwartzman
phenomenon and, 281
effects of Pyrogen and cor-
tison in rabbit on, 731
- ASBESTOS, adsorption of pyrogens
by, 558
pads, 434
- ASCORBIC ACID, content of tumor,
liver and intestines, effect
of E. typhosa ext. and col-
chicine on, 249
effects, 364
upon hemorrhage, 275
in adrenals, 697
changes produced in rats by
infection with E. coli, 608
decrease, action of dysentery
toxins, 447
in blood: See Blood, ascorbic
acid
- ARSENITE, hemorrhagic necrosis in-
duced in mouse sarcoma #37
by, 697
- ASPIRIN: See Acetylsalicylic Acid
- ASTHMA
allergic, 737, 777
bronchial, 777, 779
pathology, gross, 779
childhood, reactions after vac-
cine therapy, 638
reactions, after therapy, 638,
779
therapy, antigens, critical
review, 681
fever, induced, 637
- Pyrogen, 637, 737, 777, 779
vaccines in, 89, 637, 638
- ATROPHY, muscular. pyifer ther-
apy of tabes, 167
- AUTOPSY, 394, 779
in fatal reaction from typ-
hoid vaccine, 426
- BACKACHE, from fever induced with
typhoid pyrogen, 446
- BACTEREMIA: See Meningococci
- BACTERIA: See also Coccus;
Dysentery, bacilli; Meningo-
cocci; Pneumococci; Polysac-
charides; Pyrogens; Serratia;
Staphylococci; Tubercle
Bacilli; Viruses; etc.
abortus, filtrates, effect on
leukocyte picture in cattle
positive to brucellosis, 400
aerobacter: See also Polysac-
charides, from aerobacter
(aerogenes) carbohydrate
haptens from, role in bacter-
ial anaphylaxis, 153
cloacae, may be causative organ-
ism of 'cotton fever', 402
aerobic, detection of pyrogens
in fluids by biological met-
hods, 389
aerogenes, fever toxins of, 18
aertrycke: See Salmonella
agrobacteria, corynebacteria,
pseudomonas and xanthomona,
plant pathogens, used in tumor
hemorrhage prodn., 433
anthrax: See Anthrax; Polysac-
charides
antigens: See Antigens
bulgaricus, effects of inj. of
cultures into mice with can-
cer implants, 196
cell, chromatin portion site of
origin of fever toxin, 21
elements and toxins produce is-
chemia after inj. in chor-
oiditis, 344

- Charbon, glycolipids from, provoked increase in resistance to infection, 393
- chill, muscular temperature during, 219
- clostridia, vaccine from endotoxin, effect of thermal reaction on prodn of agglutinins, 169
- welchii, filtrates have anti-pyrexial influence, 684
- cocci: See Coccus; and under names of cocci bacteria as Staphylococci; Streptococci; etc.
- coli: See also Polysaccharides
- adrenal changes produced in rats by infection with, 608
 - agglutination picture in animals previously treated with, 62
 - changes in blood and plasma during fever, and aspirin antipyresis, 117, 188
 - cultures, blood serum increases pyrogenicity, 499
 - endotoxin, attributable to polysaccharide and nitrogen complex, 667
 - fever, dextrose plethora and its antipyretic effects, 91
 - toxins of, 18
 - filtrate, activity in malignant neoplasms, 245
 - antihemolytic effect on blood, 176
 - effect on tumors in mice, 294
 - insulin-like substance from, 154
 - prodn. of Shwartzman reaction in rabbits, 236
 - properties of hemorrhage prodg, fraction, 243
 - purification of, 261
 - rate of bacteriophage inactivation by, 317
 - reaction of mice and tumors to inj. of, 235
 - sepn. of hemorrhage-prodg. fraction of, 244
 - induced fever in dog by inj. of, blood volume and, 90
 - injection, action of antiplatelet serum on elimination of bacteria, 97
 - injection in mice modified course of virus pneumonia, 495
 - isolation of bacterial pyrogens, 751
 - leukocyte response, 342
 - nature of hyperglycemic response to inj. of killed cells of, 132
 - prepn., shock reaction, effect on lymph, serum ferments and protease titer, 68
 - therapeutic effect, 73
 - protein, effect on eviscerated dogs, 130
 - in therapy of schizophrenia and paralytic dementia, 240
 - pyrifer from, injected in patients with acute rheumatic and allergic diseases, 675
 - pyrogen formation parallel presence of 388
 - serum, antitoxic, in therapy of ulcerative colitis, 361
 - temperature-increasing factor from, 43, 65; in milk, 260
 - toxin, temperature increasing factor, 43
 - therapy of experimental (in dog) tumors, 33
 - vaccine, depressed gastric motility in dog, 355
 - glucose tolerance lowered after inj., 273
 - therapy of varied diseases, 83
- coli-aerogenes group: See Polysaccharides
- contamination of protein solns. in prodn. of fevers, 107
- content of distilled water stored in pharmacies, 42
- Corynebacteria in tumor-hemorrhage production, 433
- Corynebacterium diphtheriae, agglutinins increased when previously immunized with, 62
- fever, toxins of, 18
- toxins inhibit chemotaxis of leukocytes, 365
- culture; See also under names of bacteria

BACTERIA,
culture

of Pr. vulgaris, B. coli, St. aureus, B. subtilis and two unidentified organisms from water inoculated in horse serum, prodn. of pyrogen, 277
pyrogenic and antipyrogenic principle in, 237
diphtheria: See Bacteria,
Corynebacterium diphtheriae
Diplococcus pneumoniae: See
Pneumococci
Ducrey's: See Antigens; Chancroid; Hemophilus, of Ducrey
dysentery: See Dysentery
Eberthella typhosa: See
Eberthella typhosa
endotoxins: See also Toxins,
bacterial; names of bacteria as Dysentery, bacilli;
Eberthella typhosa; Gonococci; Meningococci; Salmonella; Staphylococci; and under subheads of bacteria as Bacteria, gram-negative
mode of action, 521
new light on, 667
protection against by penicillin and its impurities, 538
by sulfonamides, 414
protection factor in impure penicillin, hypothermia
in mouse as bioassay of, 552
responsible for pyrogenic activity in milk therapy, 84
enteritidis: See Salmonella enteritidis
Escherichia: See Bacteria, coli
extracts, chemical nature of phage-inactivating agent in bacterial extracts, 215
phage-inactivating agent of, 211
filtrates: See also Meningococci, filtrate; Staphylococci; under subheads of Bacteria
chemical studies of skin reaction to, 261
hemorrhagic activity following injection, 280
reactivity of malignant neoplasms to, 245

Flexner's: See Dysentery,
bacilli

Friedländer's: See Bacteriophage, Friedlander's; Polysaccharides, from
Friedlander's bacillus
gram-negative: See also
Diplococci
causative agents of fever and chills released by enzymes, 13
cause febrile reactions more often than gram-positive, 226
contaminants in biologic products, pyrogenicity of, 429
role of cryophile organisms, 487
detection of pyrogens in, by biological methods, 389
endotoxins, effect of sulfonamides on, 414
immunological relationship of endotoxic factor of Rickettsia prowazekii to, 453
induction of decidua-placental hemorrhage in mice, 384
neural toxins in, 413
protection against by immunization with heterologous organisms, 412
tumor-hemorrhage agents, 373
found in stained cotton and dust, 354
inhibitory effect on pneumonia virus when applied intranasally to mice, 677
sources of prodn. of pyrogens, 758
tolerance to, 186
vaccine, hypothermic and adreno-hemorrhagic effects, 356
gram-positive, contaminants in biologic products, pyrogenicity of, 429
toxins, distribution of tumor-hemorrhage agents, 373
groups, and their carbohydrates, effects on leukocytes, 717
harmful substances, studies on, 531, 584

BACTERIA

hemophilus: See *Hemophilus*
histolyticus: See *Bacteria*,
clostridium
 injection, caused leukopenia
 and alkali reserve de-
 pletion, 99
 hyperthermia by, effect on
 blood sugar, 116, 299
 intoxication, protection of
 adrenalectomized animals with
 ext. of adrenal cortex, 189
 killed, change in blood sugar
 level after inj., 116, 299
Klebsiella: See *Bacteriophage*,
Friedlander's; *Polysac-*
charides, from *B. Friedländer*
lactis aerogenes: See *Bacteria*,
aerobacter
Listerella, capacity to produce
 vascular stasis in sarcoma,
 373
 living, role of carbohydrate and
 protein in reaction, 209
megatherium: See *Polysaccharides*
meningococcus: See *Meningococci*
mesentericus: See also
Polysaccharides
 action on tubercle bacillus,
 119
 metabolism, 343
 nucleoprotein fraction is
 Shwartzman-active, 223
 product, contaminant in dis-
 tilled water, cause of fever,
 139
micrococcus aureus, splanchnic
 removal of, 762
monocytogenes, infection, re-
 action of lymphatic tissue,
 265
 paratyphoid: See *Paratyphoid*,
 vaccine; *Salmonella*
paratyposus
 pneumococcus: See *Pneumococci*
 polysaccharides from: See
Polysaccharides
 preparations, bioassay method for
 detg. potency of hemorrhage-
 producing, 380
 of crude nucleoprotein and
 polysaccharide fractions
 (*coli*, *pyocyaneus*, *strep-*
tococci), 586

trichloroacetic exts. pro-
 duce leukopenia, 366
prodigiosus: See *Coley's*
Fluid; *Polysaccharides*;
Serratia marcescens
 products, effect on leukocytes,
 172, 717
 in water responsible for
 reactions from saline
 soln., 39
 role in prodn. of 'water
 fever', 105
 proteus group: See also
Polysaccharides
 ACTH-like effect of fever
 in man, 715
 killed cells inj. in
 rabbits, hyperglycemic
 response, 132
 nature of hyperglycemic res-
 ponse to inj., 132
 pyrexia from, hypothermic
 and antipyretic effect of
 prepn. of ACTH, 742
 pyrogen fever, evaluation
 of antipyretics against,
 462
 pyrogens in parenteral
 salines, 327
Pseudomonas aeruginosa, bac-
 tericidal soln. prepd.
 from cultures of, and
 pyocyanase, 159
 damage produced by fraction
 in sarcoma, 37, 657
 effect of bacteria and their
 carbohydrates on leuko-
 cytes, 717
 filtrates, produced py-
 rexia, leukopenia and
 leukocytosis, 17
 hyperthermia provoked by,
 701; 'propidon', (Delbet), 22
 infection in rats, thermo-
 regulation during fever,
 199
 killed cultures, therapy of
 typhus abdominalis, 20
 method of prepn. and test
 for pyrogens of, 383
 pyrogens from, 18, 430
 lowest amount required to
 produce leukocytic res-
 ponse in dog, 409

BACTERIA

Pseudomonas aeruginosa

pyrogen

site of action in
central nervous system,
481, 563

tolerance, acquired, to, 377,
383

Pseudomonas scissa and ureae,

suspensions used in ther-
apy of neurosyphilis,
dementia paralytica and
tabes dorsalis, 208

pseudotuberculosis, toxin

inhibit chemotaxis of leuko-
cytes, 365

pyocyaneus: See Bacteria,Pseudomonas aeruginosapyrogens: See Pyrogens,
bacterial

Salmonella: See Salmonella

Shigae: See Antigens, Shigelladysenteriae; Dysentery,

bacilli; Polysaccharides

spirillum, endotoxins of

Rhodospirillum rubrum, induc-
tion of decidua-placental
hemorrhage in mice, 384

split products, action of, 92

superficial epithelioma

improved by, 93

subtilis: See also Polysac-
charides

action of preparations of sec-
retion on tubercle bacillus,
119

emulsions, new therapy of
diarrhea by, 618

grown in gum acacia produce feb-
rile response, 278

pyrogens from, 430

synthesis of polysaccharides by,
162

tetanus: See Tetanus

therapy: See individual
organisms

reversibility in antibacterial
and antitumor vaccine therapy,
147

toxins: See under names of bac-

teria as, Diphtheria;

Eberthella typhosa; Toxins;

etc.

tularensis: See Tularemia

typhimurium: See Salmonella

typhoid: See Eberthella

typhosa; Polysaccharides

vibrio, Metchnikovi, filtrates

produced pyrexia, leuko-

penia and leukocytosis, 17

(Metchnikovi, Finkler and

Deneke), fever toxins of, 18

BACTERICIDES, effects of organic
compds., inorganic salts,
biologic proteins, and
suspensions of dead bac-
teria when inj. into
rabbit, 186

solution prepd. from cul-
tures of B. pyocyaneus
and pyocyanase, 159

thermostable substance in
serum from fever patients,
256

BACTERIOPHAGE, effect of poly-
saccharides (Aerobacter,
starch glycogen and gum
arabic) upon, 295

Friedländer's, inhibition by
bacterial and nonbac-
terial polysaccharides, 621

inactivating agent of bac-
terial exts., 211, 215

inactivation by filtrates of
E. coli, 317

inhibition by exts. of S.
suipestifer and para-
typhosa B, 203

toxicity of young cells of
S. paratyphi A and B when
lysed by, 200

BACTIFEBRIN, use in neural
intoxications, 765

BALANITIS: See Penis

BARBITAL DERIVATIVES, barbi-
turates reduce activity
of 'necrosin', 427

diminished reactions to

Pyromen by prelim. inj.

of amobarital, 722

effect of phenobarbital on
leukocytosis, 269

BENEDICT'S SOLUTION, reduced polysaccharides of Sporotrichum schenckii, 502

BENEDICT TEST, 321

BENZANTHRACENE, di-, produced tumors, 266

BENZOIC ACID, adsorption, in prepn. of pyrogen-free urogastrone, 302,303

BERLIN, State Hospitals, therapy of neurosyphilis, 181

BICARBONATE disodium phosphate, degradative processes of polysaccharides of pneumococcus initiated by, 649

BIOLOGICAL PRODUCTS: See Serums

BIURET TEST, 243

BLOOD, adrenalin, hyperadrenalemia may produce symptoms similar to postoperative hyperthermia, 530

adrenalin, increased in fever, 143

agglutination: See Agglutination

alkali content, 117

reserve, changes in leukocytes and, in exper. infections, 99
decline in fever induced by killed B. coli, 188

antibodies, influence of various agents on mobilization, 110

ascorbic acid, effects of sublethal doses of polysaccharide from S. marcescens, 396

bacterial contaminants in prepn., pyrogenicity of, 429

bromsulfalein, clearance during pyrogenic reaction (typhoid vaccine), 478

calcium, arterial and venous level in fevers, 466

carbohydrates; See Blood Sugar

cells: See also Erythrocytes; Leukocytes; Plasma, cells

changes after inj. of materials eliciting Shwartzman phenomenon in rabbit, 730
effects of typhoid vaccine on, 619

changes (increased conc., hemoglobin and oxygen and alkali reserve decrease) in fever induced by B. coli, 188

cholesterol, affected by fever, 325

cholinesterase, regulation by pyrififer and ultraviolet rays in therapy of dermatoses, 510

circulation: See also

Capillaries

accelerated during therapy (pyrififer), 163

in arthritis during fever (induced), 550

changes during chill and fever (typhoid vaccine), 416

collapse, dangers of therapy with mixed toxins (Coley's), 29

death from failure, after a admin. of polysaccharide, 394

peripheral, collapse threat after inj. S. marcescens polysaccharide in tumor therapy, 502

during exper. fever, 231

summary of mechanism involved in fever, 415

coagulation: See also Plasma, coagulation prevention by S. marcescens polysaccharide, 650

culture, after inhalation of S. marcescens, 455

diastase, altered during hyperglycemic reaction, 259

enzymes, effect of B. coli prepn. on, 68

influence of various agents on mobilization of, 110

esgothionone, increased during pyrexia induced by

pyrififer, milk protein or bacteria, 207

BLOOD

fats, phosphatides and cholesterol decreased after inj. of pyrogens in dog, 358

fibrinogen, in therapeutic fever, 351

formation, extramedullary hemopoiesis in rabbit and cat induced by pyrogens (Pyromen), 692

globulin, alteration of pseudoglobulin I and II after inj. of omnadin, pyrifer and milk in diphtheria, 268

glutathione, increased during pyrexia induced by pyrifer, milk protein or bacteria, 207

groups, specific substances (A and B) added to O blood for transfusion, pyrogenic reactions, 425

hemolysis: See Hemolysis

histamine, effect of meningococcal endotoxin on, 497

influence of bacterial proteins on (cells, chemistry, volume, rate), 157

lipoids, fluctuation during fever, 325

nitrogen, nonprotein, effect of sublethal doses of S. marcescens polysaccharide, 396

phosphorus, changes after inj. of suspensions of B. aertrycke in rabbit, 170

level, fall and rise after inj. of bacterial antigens, 259

picture, changes by bacterial pyrogen "pyrotoxin", 53

effect of typhoid exts. on, 290

plasma: See Serum

platelets, action of antiplatelet serum on elimination of bacteria injd. into circulation, 97

anticoagulant effect of bacterial polysaccharides in normal and thrombocytopenic plasma of leukemia, 743

thrombopenia of infective diseases and of protein injts., 118

pressure: See Blood Pressure

proteins: See also Blood, globulins

bound to carbohydrates in fevers, 180

during acute infections, 315, 322

effects of pyrifer on globulin-albumin alteration, 308

in therapeutic fever for general paresis, 351, 609

precipitable by C polysaccharide of pneumococci, 322

relation to corrected sedimentation rate, 352

proteose, of rabbits implanted with Brown-Pearce carcinoma, treated with S. marcescens polysaccharide, 464

sedimentation rate, effect of pneumococcus polysaccharide on capillary circulation, 362

relation of plasma proteins to, 352

serological study of polysaccharides of Meningococcus, B. anthracis, B. proteus, B. subtilis and B. mesentericus, 187

sugar, (free, protein-bound and non-protein), in normal and pathologic serum, 180

changes after inj. of suspensions of B. aertrycke, 170

effect of fever induced by inj. of killed bacteria, 299

fractions from B. aertrycke in rabbits, 214

inj. of protein-polysaccharide fraction from S. typhimurium, 297

hyperglycemia after intrav. inj. of killed bacteria, 116

BLOOD

sugar

hyperglycemia

nature of response to inj.
of killed bacteria (B. coli and proteus), 132
produced by inj. of bacterial
antigens, 259
produced by protein frac-
tion of B. dysenteriae,
258

relation of S. enteritidis-
paratyphosus B infections
in rabbits to, 111

level during induced fever
in rabbits, 273

maintenance with 5% glucose,
449

transfusion: See Blood Trans-
fusion

uric acid, of rabbits implant-
ed with Brown-Pearce carc-
inoma, treated with S.
marcescens polysaccharide
464

levels, after therapy with
S. marcescens polysaccha-
ride, 390

vessels: See Blood Vessels
volume, dextrose plethora and
its antipyretic effect
on coli fever, 91

(in dogs), in coli fever, 90

water content, in febrile dogs,
117

whole, latency of response to
pyrogens as affected by
previous incubation with,
744

BLOOD PRESSURE, arterial, action
of ext. of E. typhosa, 476

during pyrogenic therapy with
colloidal metals, 76

fall after inj. of S. marcescens
polysaccharide in tumor
patients, 502

high, analysis of mechanism,
313

arterial, therapy with pyro-
gen (Pyromen), 603

associated with pyrogenic re-
action, reduction, 338

effect of pyrogens on malig-
nant hypertension, 549,
604, 705

essential, effective renal
flow in subjects with, 320
inj. of pyrogenic agent,
ACTH-like effect of fever,
661

renal irritation from foreign
protein (typhoid) therapy,
382

response to pyrogens, 397
therapy by albumin fraction
from pork serum, 407
hypotensive effect of
fever-producing substances,
407

pyrogens (inulin, typhoid
vaccine) in; effect of
premedication with anti-
pyretics, 338, 420

Pyromen, in essential and
malignant, 722

in therapy of arthritis with
typhoid vaccine, 75

increase after typhoid vaccine,
72, 264

low, effect of peripheral
hypotension on activ. of
Shear polysaccharide, 770
on transplanted tumors, 695

systolic, effect of S.
marcescens tumor-nec-
rotizing polysaccharide
(in mice), 515

BLOOD TRANSFUSION

chills, use of calcium chloride
in, 296

citrated blood in, prevention
of chills after, 204

pyrogens, control in solutions
used in, 630

preliminary report on, 435

reactions caused by, 296,
332, 425, 513

after transfusion of group
specific conditioned blood,
425

prophylaxis against, 632

statistical study of
clinical use, 706

BLOOD TRANSFUSION

pyrogens

reactions

unpooled liquid human plasma,
analysis, 1500 transfusions, 360
whole blood and plasma, in
multiple sclerosis, 513

Type O blood conditioned by group-
specific substances, re-
actions, 425

BLOOD VESSELS: See also Capil-
laries; Kidneys, blood supply;
Skin, blood supply; Thrombo-
angiitis obliterans; Vena
cava; etc.

changes during therapy of
paresis with fever, 609
sarcoma with polysaccharide
(Shear), 770

constriction mechanism produced
by fever, raises internal
temperatures, 10

diseases, fever therapy in
peripheral diseases, 223

effect of drugs on, dilation in-
duced by aspirin, 369

engorgement, of sarcoma, 255

mechanism of lymphocytic lesions
provoked by epinephrine and
Salmonella typhi, 569

occlusion, maximum seen 3 to 4
hrs. after inj. of B.
prodigiosus polysaccharide,
437

therapy with foreign protein
(typhoid vaccine), 144

reaction of normal and neoplastic
tissues to S. marcescens
polysaccharide, 437, 467, 770

BODY, structure and induced fever,
712, 761

weight, changes after inj. of
dysentery fractions, 259

BONES, long, invoked fever (typhoid
vaccine) enhances effect
of irradiation in therapy
of leukemia, 424

marrow, accumulation of
neutrophils in, after inj.
of E. typhosa filtrate,
271

changes during leukopenia,
259

hyperplasia after injury to
hematopoietic tissue by
toxic antigen of E. typhosa,
376

sarcoma, therapy with toxins
(Coley's), 148

BOVIN'S METHOD of prepg. anti-
genic fractions from
B. typhimurium and B.
typhosum, 301

BRAIN: See also Hypothalamus;
Nervous System, central
blood circulation, effects of
fever induced by pyrogen,
605

cortex, blending of regenera-
ting peripheral fibers with,
through glial barrier in-
hibition by Pyrogen, 704

effect of substances affect-
ing body temperatures
determined on oxidative
and glycolytic activity of,
528

modification of control upon
thermoregulation in emo-
tional hypothermia in rab-
bits, 643

ecchymoses and petechiae of,
after anti-typhoid
therapy, 426

electroencephalography, ef-
fects of induced fever on
patients with multiple
sclerosis, 713

extracts, of V. Muralto, sug-
gested therapy of multi-
ple sclerosis, 506

fever produced by bacterial,
chemical and mechanical
stimulation of midbrain,
145

BRAIN.

- physiology, comparison of respiration and glycolysis in normal and febrile rabbits, 606
 - effects of induced fever in neurosyphilis, 653
 - pyrogens may interfere with thermoregulation at lower level of brain stem, 577
 - regeneration, induction (Pyromen or ACTH) of neuronal regeneration in, 757
 - inhibition of glial barrier as means of permitting peripheral nerve to grow in brain, 704, 740, 756, 757
 - penetration by regenerating peripheral nerve fibers, 704, 740, 756,
 - structural and functional, in central nervous system, 741
 - thermal centers, fever induced by B. pyocyaneus infection in rat did not inhibit, 199
 - stimulated in protein therapy, 102
 - transection of midbrain, influence on infectious and non-infectious fever, 55
- BREAST, cancer, diagnosis, serum polysaccharide level, 615
- cancer, reaction of spontaneous mouse carcinomas to toxins of E. typhosa, 225
 - spontaneous and induced tumor response to B. coli filtrate, 235
 - milk: See Milk, human
- BRIGHT'S DISEASE: See Nephritis
- BROMSULFALEIN: See Blood, bromsulfalein; Liver, function test
- BUBO, therapy with pyrifer, 178
- BUERGER'S DISEASE: See Thromboangiitis obliterans

- BURNS, healing, enhanced by Pyromen, 754
- healing, in cats treated with Pyromen, 710
- renal function in, pyrogen increases plasma flow, 714

- CALCIUM, chloride, 359, 721
- chloride, used for chill due to malaria, transfusions, and typhoid vaccine, 296
 - effect on flocculation reaction with C polysaccharide of Pneumococcus, 315
 - gluconate or chloride suppress hyperpyrexia, 359
 - in blood: See Blood, calcium
 - intravenously for post-transfusional reactions, 706
 - metabolism, in fevers, 466
 - non-protein thermogenic factor from B. coli and E. typhosa combined with, formed inert salt, 65

- CALORIMETRY, clinical, 104

- CANCER: See also Sarcoma; Tumors; and under names of organs or regions
- carcinoma, high value for sugar associated with protein in serum, 180
 - effect of bacterial polysaccharides may be overlooked, 444
 - effect of colchicine and B. coli filtrates, 294
 - daily variation in leukocytes, 44
 - epithelioma, inj. of Joannovic's tumor tissue and bacterial split products, 93
 - transplanted in mouse, unchanged after inj. of S. marcescens polysaccharide, 444
 - immunity, lymphocytes in natural and induced resistance to transplanted cancer in mice, 86

CANCER:

- metastases, effect of iodinated S. marcescens polysaccharide on, 508
- mouse, effect of cultures of B. bulgaricus on, 196
- reaction of spontaneous carcinomas to bacterial toxins, 225
- rat carcinoma, effect of toxic carbohydrate complex from S. enteritidis on, 311
- effect of Shear polysaccharide alone or with flavonoid compound on, 750
- therapy, by bacterial toxins as developed by late Wm. B. Coley, reviewed in light of modern research, 452
- effects of iodopolysaccharide in pateints, 544
- fever, theoretical experiments, 120
- of Brown-Pearce carcinoma implanted in rabbits with S. marcescens polysaccharide, 463
- of carcinoma with pure culture material of S. erysipelatis, 19
- of transplanted carcinoma, reactions to fractions from S. typhimurium, 293
- use of polysaccharide S. marcescens or dicumarol, 650
- with erysipelas serum, 25

CAPILLARIES: See also Blood

- Vessels; Skin, blood supply circulation, effect of bacterial polysaccharide on unanesthetized mice, 436
- effect of foreign protein therapy (typhoid vaccine), 144
- Pneumococcus polysaccharide, 362
- in typhoid vaccine fever, 155
- dilution in, alters leukocyte conc. in parenteral therapy, 112
- effect of nonspecific protein therapy upon, 138
- pyrogens on, 749

permeability, 117

after inj. of hermoohage-prodg. fraction of B. coli filtrate, 243

CARBOHYDRATES: See also Dextrose;

- Glycogen; Polysaccharides; Starch; Sugars; etc.
- acetylated, specific antigenic complexes of Sh. para-dysenteriae, 418
- adsorbed on colloids (as haptens) as antigens, 194
- chemical nature of immune substances, 134
- complex, isolated from culture medium and ultrafiltrate of tubercle bacilli, 216
- effects of S. typhosa and Ps. aeruginosa on leukocytes, 717
- effects upon leukocytes of inj. of, 137
- factors relating to polysaccharides and, 162
- fraction of B. dysenteriae produced hyperthermia, 258
- from culture medium of Sporotrichum schenckii, 503
- injection into tumor-bearing rats, 280
- metabolism, effect of bacterial endotoxins on, 536
- effect of inj. of protein-polysaccharide fraction from S. typhimurium on, 297
- pathogenizing effect on E. typhosa, 454
- role in specificity of reaction to living or killed bacteria, 209

CARBON, in ext. of culture of

- B. paratyphosus B, 106
- dioxide, respiratory response to inhalation of, 572

CARBUNCLE: See Anthrax

CARDIOVASCULAR DISEASES: See also
 Blood Vessels; Heart criteria
 for selection of patients for
 therapy symptomatic paresis
 with penicillin and typhoid
 vaccine, 594

effects of bacterial polysac-
 charides upon, 589
 ——— Shear bacterial
 polysaccharides upon, 662

CASEIN. digest, dialyzed
 (aminosol), pyrogen activity,
 465

hydrolysate, in sterile soln.
 and monosaccharides pro-
 duced fever in infants by
 oral admin., 445

glutathione and ergothione in
 blood increased during
 pyrexia, 207

in milk therapy, 84
 prodn. of fever in rabbits
 with, 107

CATABOLISM: See Metabolism

CATS, effect of hypothalamic
 lesions on induced fever, 289
 extramedullary hemopoiesis in-
 duced by bacterial pyrogens
 (Pyromen), 692

healing of burns treated with
 Pyromen, 710

intravenous admin. of Pyromen,
 role of adrenals in mech-
 anism of leukocyte changes,
 720

reaction after pyrogen admin.
 and inj. of S. aureus
 filtrates, 398

to typhoid-paratyphoid vaccine,
 270

regeneration, of peripheral
 nerves, penetration of brain
 by, 704, 740

of spinal cord, conduction of
 nerve impulses, 725, 778

————— site of action
 of bacterial pyrogens (Pyromen),
 689, 690

CATTLE, effect of filtrates of B.
abortus on leukocyte picture, 400

CECUM, hemorrhage from inj.
 of variant of S. dysenteriae,
 447

CELLOPHANE MEMBRANES, 261

CELLS: See also Bacteria, cells;
 Blood, cells; Leukocytes;
 Lymphocytes; Phagocytosis;
 Plasma, cells; Tissue; etc.
 body, mechanism of fever, 174
 division, effect of poly-
 saccharide S. marcescens
 on, 650

free stem, morphologically
 identical with lympho-
 cytes in lymphatic tissue,
 mesenteric node and spleen,
 265

localization of antigen in
 tissue, 654, 658

mast: See Tissue, connective
 permeability, in non-specific
 protein therapy, 101

CENTANNI, 'third immunity' of, 336

CENTRIFUGATION, separation of
 tumor-hemorrhage prodg.
 fraction of B. coli, 244
 ultracentrifugal analysis of
 hemorrhage-prodg. frac-
 tion of S. marcescens, 374

CEPHALIN FLOCCULATION TEST, in
 rheumatoid arthritis, 696

CHANCROID, experimental, 141, 142
 therapy, comparison of vaccine
 (Streptobacillin) and
 nonspecific protein, 142
 vaccines, 254

CHARCOAL, activated, adsorption
 of pyrogens, by, 558
 activated, detoxification of
 pyrogens, 548
 elimination of pyrogens in
 injectable solns., 327, 422
 in prepn. of non-pyrogenic
 water and infusion fluids,
 442

powdered, prepn. of pyrogen-
 free water for intravenous
 use, 304

CHARCOAL

removal of inhibiting substance
by, 197

CHEMOTHERAPY: See Tumors; under
names of diseases and chemo-
therapeutic agents

CHICKEN, body fluid redistribution
in induced hypothermia and
hyperthermia, 673

CHILDREN, diseases: See Asthma,
in children: Typhoid,
in children
'sugar' and 'salt' fever and
water reserve of body, 81

CHILLS, after typhoid vaccine,
66, 72, 264
after inj. of distilled water,
140
—— of proteose and
typhoid vaccine, 104
after pyrifur injection, 178
bacterial, muscular tremor and
temperature during, 219
circulation and respiration
during, 416
during pyrogenic therapy with
colloidal metals, 76
etiology, by principles in
bacteria, 13
following intravenous therapy,
288
fever-prodg. principle in
typhoid vaccine, 276
in therapy of arthritis with
typhoid vaccine, 66
incidence after transfusion, 332
nature of chill-prodg. principle
in antipneumococcus serum, 182
phase, occurs in method for
controlled hyperpyrexia, 359
prevention, after transfusion of
citrate blood, 204
produced in cat injd. with
bacterial pyrogen, 481
prodn. of pyrogen in sera by
bacteria, 277
summary of mechanisms in circula-
tion and respiration in fever,
415

therapy, calcium chloride, 296

CHINIOFON, therapy, oral
stimulative, 103

CHITIN, serologically inactive
substance allied to
chitin found in Pneumococcus
type IV polysaccharide, 177

CHLORIDE, ion conc. falls with
volume of gastric secre-
tion in fever, 201

CHLOROBENZENE: prevention of
pyrogenic contaminants, 758

CHLOROFORM, extraction, 272

CHLORO, di-, PREPARATIONS,
prevention of pyrogenic
contaminants, 758

CHLOR-TRIMETON, effects, before
and after inj of Pv
Pyromen, 715

CHOLERA, vibrio, filtrates have
antipyrexial influence,
684

CHOLESTEROL, adrenal changes in
rats by infection with
E. coli, 608
in blood: See Blood, choles-
terol

CHOLINESTERASE, in blood: See
Blood, cholinesterase

CHOREA, therapy, induced fever
by typhoid vaccine, 313
therapy, typhoid shock, 461

CHOROIDITIS, centralis serosa,
diagnosis, pathologic
physiology and therapy, 344

CHROMATIN, of bacterial cell,
site of origin of fever
toxin, 21

CITRATES: See Blood Transfusion,
citrate blood in

- CLAVICLE, tumor, with axillary metastases treated with Coley's fluid, 28
- CLINIC, Psychiatric and Neurologic Diseases, of University of Göttingen, 163
- CLOSTRIDIA chauvei: See Anthrax, bacilli
- COBRA, capello, venom, in therapy of malignant tumors, 30
- COCAINE, fever, blood and plasma changes, and aspirin anti-pyresis, 117
- COCCI, antigenic polysaccharide complex from, 285
thermostable bactericidal substance from serum of fever patients active against, 256
- COLCHICINE, effects, increases effectiveness of X-ray therapy in leukemia, 424
effect, on tissue metabolism, 249
——— tumors in mice, 294
toxicity, 249
- COLD, application to heat regulating center, modifies action of induced pyrexia and some anti-pyretics, 78
- COLEY'S FLUID (mixture of erysipelas-prodigiosus toxins), 312, 452
- AMA Council reviews literature on use, retained in NNR, 220
reactions after therapy, 772
therapy, of carcinoma, 27, 772
——— Hodgkin's disease and lymphosarcoma, alone or with radiation, 149
——— malignant tumors, 26, 27, 28, 29, 30, 32, 34, 36, 37, 46, 60, 136, 148, 452
sarcoma, 28, 29, 32, 34, 36, 37, 60, 399; melanotic, 46;
multiple, bone and soft tissue, 148; myo-, of uterus, 210
——— manic-depressive states, 292
——— ocular infections, 491
survival after, 772
- COLITIS, ulcerative, pyrexial therapy (pyrifer), 656
ulcerative, therapy, concd. and purified antitoxic B. coli serum, 361
therapy, controlled hyperpyrexia, 359
- COLLOIDS: See also Metals, colloidal; Starch; Sulfur and colloid therapy, 70
carbohydrates adsorbed on, as antigens, 194
chemical changes in body cells during fever, 174
electrolyte, separated from polysaccharide of type I pneumococcus, 512
in fever, normal water reserves bound by, 81
ineffective after section of medial hypothalamus in febrile response in rabbit, 55
injection, storage process, 151
specific patterns rather than polar group or unit structure may underlie pathogenicity of carbohydrates, 454
(starch, inulin, gum arabic) in-activation of coli-phage, 317
- COLOR BLINDNESS, temporary, after typhoid vaccine therapy, 588
- COLORIMETRY, tests for tyrosine, arginine and histidine in polysaccharide from B. prodigiosus, 372
- COMPLEMENT, fixation, 141
- CONSTITUTION, differences in temperature during pyrifer admin., 712
- CONVALESCENCE, tolerance to toxic effects of antigens of enteric bacilli in typhoid and paratyphoid, 598

- COPPER, hydroxide, elimination of pyrogens by, 592
- CORNEA, ulcers, therapy with mixed toxins (Coley's), 491
- CORTICOSTERONE: See 11-Dehydro-17-hydroxycorticosterone; Desoxycorticosterone
- CORTISONE: See Adrenals, cortex extract
- CORYNEBACTERIA: See Bacteria, corynebacteria
- COTTON, 'cotton fever', incidence in Miss. cotton factories, 402
low-grade stained, acute illness among mattress makers and cottonseed processors, 354
- COUNCIL on Pharmacy and Chemistry, AMA, reviews literature on use of mixed toxins of Coley, retains in NNR., 220
- CRANIUM, tumors, metastases of myosarcoma of uterus, treated with Coley's toxins after hysterectomy, 210
- CREATININE, in urine: See Urine, creatinine
- CULTURE: See also Bacteria, culture; and names of bacteria
evidence of antineoplastic substance as metabolite from culture fluids of Penicillium notatum, 428
- CULTURE MEDIA, blood serum, produce organisms yielding more pyrogen, 429
contg. dextrose and peptone (Sabouraud) required for prodn. of pyrogens in mold cultures, 646
cultures of staphylococci on pyrogenic or pyrogen-free media, 398
- gelation, by bacterial pyrogens, 379
glucose, lactose and galactose increase growth of E. coli and stored polysaccharide, 568
-1-C¹⁴, in culture of hemolytic streptococci group A, 753
influence of synthesis of amino acids in media on pyrogen prodn., 218
latency of response to bacterial pyrogens as affected by previous incubation with whole blood, 744
plasma growth of S. enteritidis, 311
potato, growth of tubercle bacilli on, 119
relation between pyrogenic activity and gelatination, 388
role in increasing pyrogenicity of supernatant fluids, 499
synthetic, and broth, culture of B. prodigiosus, 312
culture of E. typhosa in, 305
large scale prepn. of tumor-necrotizing polysaccharide from S. marcescens, 504
- CYANOSIS, after typhoid vaccine, 72
capillary flow in man in typhoid vaccine fever, 155
- CYTOSTATIC PREPARATIONS, in therapy of neoplastic ailments, 590
- DAHLIA, inulin, toxicity, 252
- DEATH, causes, anaphylactoid reaction after therapeutic anti-typhoid inj., 426
- "DECALSO": See Ions
- DECIDUA, -placental hemorrhage in mice, induced by endotoxins of gram-negative bacteria, 384

- DEGENERATION, fatty, action of mixed toxins (Coley's), 29
- DEHYDRATION, 173
fever and water reserve of body, 81
- 11-DEHYDRO-17-HYDROXYCORTICOSTERONE, most powerful protective agent tested against toxic material (typhoid vaccine), 449
- DEMENTIA, paralytica: See Paralysis General
- DEMENTIA PRAECOX, alterations in blood by pyrifer, 357
etiology, cerebral lesions as cause, 760; toxins, hepatic, 760
induced fever and, 292, 760
therapy, of early hebephrenia with pyrogen, 221
pyretotherapy (Dmelcos, Ducrey bacillus atoxic vaccine) or sulfosin, 230
with pyrifer, 217, 227, 240
with typhoid-paratyphoid vaccine, 760
- DEPRESSION, therapy, effects of Pyromen in subpyrexial doses, 708
- DERMATOSES: See Skin, diseases
- DESOXYCORTICOSTERONE, acetate, inability to protect rats against typhoid vaccine, 300
- DEUTEROALBUMOSE: See Albumose
- DEXTRAN: See Polysaccharides, bacterial
- DEXTROSE, exchange, in vitro of liver slices from febrile rabbit, 707
intravenous, antipyretic effect in febrile dog, 90
phosphorylation, inhibition by meningococcal endotoxin, 535
plethora and its antipyretic effect in coli fever, 91
solutions, dehydration products cause of febrile reaction, 314
isotonic, loss of pyrogenic activity, 610
pH in prepn. of non-pyrogenic fluids, 459
prepn. . and apparatus for intravenous and subcutaneous use, 206
——— of non-pyrogenic fluids, 251, 341
review of causes of reactions after intravenous inj., 213
tests on non-pyrogenic fluids, 458
tolerance, lowered after subcutaneous inj. of B. coli vaccine in rabbits, 273
- DIABETES MELLITUS, effect of fever induced by inj. of killed bacteria, 299
- DIAL, intravenously most active antipyretic agent, 438
- DIALYSIS, 321
purification of bacterial filtrates by, 261
removal of inhibiting substances by, 197
- DIAPYCNOSIS: See Skin, blood supply
- DIAPHRAGM, glycogen, effect of bacterial endotoxins on synthesis, 534
- DIARRHEA, in infants, linked with presence of 'thyrotoxicosis' (split protein), 57
reaction to fever induced by killed B. coli, 188
therapy, by emulsions of B. subtilis, 618
- DIASTASE: See also Enzymes
hydrolysis did not follow treatment with, 430
in blood: See Blood, diastase
in etiology of fever, 13, 14

- DIBENAMINE, effects before and after inj. of Pv Pyromen, 715
effect on peripheral vascular response of rabbits to S. typhosa endotoxin, 627
- DIETHYLENE GLYCOL: See Ethylene Glycol
- DIHYDROERGOCORNINE, prevention of pyrogen-induced renal hyperemia in dog, 682
- DINITROPHENOL, induced fever, calcium level in arterial and venous blood in, 466
- DIODRAST, clearance, filtration rates and maximal tubular excretion in essential hypertension, 320
clearance, of renal tubule under various exptl. conditions, 264
- DIPHTHERIA, bacilli: See Bacteria, Corynebacterium diphtheriae
blood in, antitoxic action increased by intramuscular inj. of omnadin, 268
therapy, with antitoxin, 694
with fever therapy (pyrifer), 694
toxins, effect on blood sugar and phosphorus, 170
effect on leukocytes, 172
inhibit chemotaxis of leukocytes, 365
- DISEASE: See also under names of diseases; Infections; Therapy; etc.
influence upon febrile response to typhoid pyrogen, 585
therapy, new substance, 'third immunity' of Centanni has potentialities in therapy, 336
protein, 115
- DIURESIS, after fever crisis
evidence of water liberation, 81
- control of renal blood flow and glomerular filtration during pyrexial reactions (typhoid vaccine), 264
febrile plasma did not induce, in glomerulonephritis, 564
- DOGS, action of mono- or polysaccharide solns. on gastric mucosa, 267
action of toxic doses of polysaccharide from S. marcescens, 394
assay of gastric inhibitory factor of urogastrone in, 302, 303
depression of gastric motility and secretion by pyrogens, 378
without elevation of temperature by pyrogen, 355
effect of bacterial pyrogens on blood lipides, 358
of sublethal doses of S. marcescens polysaccharide, 396
eviscerated, effects of bacterial proteins on, 130
experimental tumors, effects of bacterial toxins, 33
ulcers, effect of urine exts. on, 329
hypophysectomized or tuberinjured, temperature and thermal reactions in, 184
influence of toxic substances on nitrogenous and purine metabolism, 125
quantitative study of reduction of gastric secretion associated with pyrexia, 625
regeneration of spinal cord of, 736
of spinal cord, experiments to determine site of action of bacterial pyrogens (Pyromen), 689, 690
tentative test for pyrogens in infusion fluid, 339
- DOISY method for prepg. pyrogen-free urogastrone, 302, 303

DOLMAN TEST, for enterotoxin, relation of pyrogen to, 397, 398

DUODENUM, ulcers: See Peptic Ulcers

DUST, stained cotton, gram-negative bacillus in, acute illness among workers, 354

DYSENTERY, bacilli, agglutination picture in animals previously treated with, 62

bacilli, antigen: See Antigens, *Shigella dysenteriae* chemistry, toxicity and biological properties of fractions of, 259

cultures, factors affecting toxicity, 343

endotoxin attributable to polysaccharide and nitrogen complexes, 667

——— consists of toxic protein, polysaccharide and lipid, 559

——— effect on pregnancy in rabbits, 411

——— induced decidua-placental hemorrhage, 384

——— tolerance in typhoid and paratyphoid convalescents, 666

bacilli, fractions injected into test animals, 259

hypothermic factor of, 258

immunological properties

against tumor hemorrhage, 386

polysaccharide: See Polysaccharides

produced leukocyte response, 342

immunity, problems, 543

toxin, 559, 560

action on different laboratory animals, 447

EBERTHELLA typhosa, antigen: See Antigens

typhosa, endotoxin attributable to polysaccharide and nitrogen complex, 667

endotoxin, cutaneous vascular reactions in rabbit, 477

——— effect of dibenamine on peripheral vascular response in rabbits, 627

——— intravenous test doses of, 598

——— relation to virulence, 334

——— tolerance to action of, 666

extract, action on arterial pressure in rabbit, 476

——— effect on tissue metabolism, 249

——— toxic, on blood picture, 290

filtrate, activity in malignant neoplasms, 245

——— antihemolytic effect on blood, 176

——— hemorrhagic

necrosis and regression of sarcoma, 180, 233

——— purification, 261

——— toxic, provoked

leukopenia in rabbits, 271

immunological relationships of polysaccharides of, 287

injection, action of anti-platelet serum on elimination of bacteria, 97

——— in chronic

rheumatoid arthritis, 659

——— in rabbit, 247

——— in rheumatoid

arthritis, remission with liver damage, 696

intoxication in animals by, protection with ext. of adrenal cortex, 189

method of fractionation of cellular component at temperature of liquid air, 31

mixed with trypan blue or azoprotein injected into mouse, followed by glycolipid antigen, 393

EBERTHELLA

typhosa

- non-protein thermogenic factor isolated from, 65
 - pathogenizing effect of
 - different carbohydrates, 454
 - polysaccharide and protein content, 334
 - precipitable substances from, 106
 - produced leukocyte response, 342
 - pyrogen from, 430
 - development of tolerance to and its abolition by reticuloendothelial blockade, 441
 - immunity in exptl. fever, 336
 - influence of various diseases upon febrile response to, 585
 - purified, 392
 - titer, bacterial products increased, 110
 - toxic extracts, effect on blood picture in rabbits, 290
 - filtrates, provoked leukopenia in rabbit, 271
 - fraction, leucocidin, activity, 253
 - substance I, local skin reaction, 152
 - toxin, fever, 18
 - leukocidal, extd. from, 279
 - reaction of spontaneous mouse carcinomas to, 225
- EDEMA, after tourniquet pressure caused no increase in peripheral tissue permeability, 436
- after meningococcus filtrate injection in mouse sarcoma, 180, 266 of sarcoma, 255

EDESTIN, prodn. of fever in rabbit, 107

EDINBURGH THEORY: See Fever

EGGS, inoculated with cultures of viruses of mumps, influenza and Newcastle disease, 525

ELECTROPHORESIS: See Ions

EMETINE, hydrochloride, effect on sarcoma 37 in tissue culture, 498

EMOTIONS, disturbances, after inj. of distilled water, 140

emotional hypothermia in rabbits, 643

ENCEPHALITIS, allergic, Piromen in therapy, 771

choreic, evolution toward parkinsonism, TAB pyretotherapy, then radiotherapy, 547

ENDOCARDITIS, infectious, blood became sterile after febrile reaction with foreign protein, 100

infectious, method for controlled hyperpyrexia, 359

ENDOCRINES: See also Adrenals; Hormones; Pituitary Body; Thyroid; etc.

cellular response to bacterial pyrogen, 688, 693, 754

ENDOTOXIN: See Bacteria, endotoxins; Bacteria, gram-negative; Toxins, bacterial; names of bacteria, as Dysentery, bacilli; Eberthella typhosa; Gonococci; Meningococci; Salmonella; Staphylococci; and under subheads of Bacteria

ENTEROGASTRONE, depressant of stomach motility may owe effect to pyrogens, 378

ENTEROTOXIN: See Intestines, mucosa; Staphylococci, enterotoxin

ENZYMES: See also under names of enzymes, as Diastases; Pyocyanase

ENZYMES

- action in protein split products, 57
 - activity of B. aertrycke cells, 170
 - causative agents(in bacteria) of fever and chills released by, 13
 - hydrolysis of capsular polysaccharide of Klebsiella pneumoniae type B, 554
 - in blood: See Blood
 - influence of various agents on mobilization of, 110
 - inhibitors and febrile process ("piretamina"), 524
 - mechanism of reaction of non-specific protein therapy, 110
 - of response to pyrogen therapy of arterial hypertension, 603
 - pancreatic, hydrolysis did not follow treatment with, 430
 - 'pyrexin' secured by enzyme action in 'necrosin' of inflammatory exudates, 427
 - pyrogens from breakdown of metabolizable materials in body or synthetic media, 47
 - reactions, disturbance of carbohydrate metabolism may effect , 535
 - synthesis of polysaccharides by, 162
 - systems, destruction of, or anoxia may underlie liver cell necrosis, 479
 - mobilize in nonspecific protein therapy, 101
 - titer, (protease, peptidase, diastase and lipase) during protein shock reactions (B. coli), 68
- EOSINOPHILIA, after pyrogens, 759
- during anaphylaxis, 759
 - therapy of E. typhosa infection with typhoid vaccine, 517
- EOSINPHILS, eosinopenia follows fever (induced), 661 ; polysaccharide of Shear, 776

EPILEPSY, therapy with pyrifer, 217

EPINEPHRINE: See Adrenalin

EPITHELIOMA: See Cancer

EQUIPMENT, transfusion or infusion, cleaning of, 246

ERGOTHIONONE, in blood: See Blood, ergothionone

ERYSIPELAS, bacillus: See Streptococci, erysipelatis toxin: See Coley's Fluid

ERYTHEMA, produced by ultraviolet rays with increased cholinesterase activity, 510

ERYTHROCYTES: See also Blood, cells; Hemolysis destruction, in therapeutic fever, 351

- injection, action on elimination of bacteria in circulation, 97
- pneumococcus polysaccharide produced changes in rate and uniformity of flow, 362
- sedimentation rate in differentiation of bacterial polysaccharide, 611

ESCHERICHIA coli: See Bacteria, coli

ETHYL CARBAMATE, effect on temperature regulation, 578

ETHYLENE GLYCOL, extn. of antigenic endotoxin of S. dysenteriae, 363

- extn. of bacterial antigenic fractions, 301
- of substance from E. typhosa, 353

ETHYLENE OXIDE, use in prevention of pyrogenic contaminants, 758

EUGLOBULIN, in blood, 268
of inflammatory exudates, 401,
427

EXTREMITIES: See also Legs
blood supply: See also Extremities
diseases; Thromboangiitis
obliterans
diseases, typhoid vaccine therapy
of peripheral vascular diseases,
335
febrile reactions during inj.
of pyrogens into normal
and deafferented extremities,
739

EXUDATES: See also Edema; Pus
inflammatory, euglobin fraction
of (pyrexin), chemical basis
of fever, 401, 427
thermostabile leukocytosis
factor of, 663

EYES: See also Uvea
diseases, continuous intrav. inj.
of typhoid vaccine, 567
"mixed toxin" from St.
erysipelatis and B. pro-
digiosus, 491
non-specific protein therapy,
328
therapy, pyrogen, role of
adrenal cortex in, 620
—— stress, a factor in,
774, 775
—— with pyrifer, 768;
Piromen, 774, 775; typhoid
vaccine, 461
muscles, paralysis after typhoid
vaccine therapy, 588

FACE, paralysis: See Paralysis,
facial

FASTING, aqueous and alcoholic
ext. of muscle and liver of
feverish animals, 7

FATS: See Blood, fats; Lipoids

FATTY DEGENERATION: See Degeneration,
fatty

FELIX anti-typhoid serum: See
Globulin, immune

FEMUR, tumors, metastases of
myosarcoma of uterus,
treated with Coley's toxins
after hysterectomy, 210
tumors, myelosarcoma, with
metastases, treated with
Coley's mixed toxins, 148

FEVER: See also Malaria; Para-
typhoid; Pyrogens; Temper-
ature, Body; Typhoid
action of toxic doses of poly-
saccharide from S. marces-
cens on dog and guinea
pig, 394
active factor from cells of
B. coli communis resemble
cleavage products of prot-
ein, 43
after inj. of medicaments, 5
alimentary, 173, 174, 175
alteration in tissue during
infections, 122
and water reserve in
body, ('thirst', 'inanition',
'sugar' and 'salt' fever),
81
antigenic material from E.
typhosa elicits fever when
antibodies are present, 323
aqueous and alcoholic exts. of
muscle and liver of fast-
ing animals during, 7
atypical, 22
blister (cold sore): See Herpes
labialis
changes in circulation and
respiration after admin. of
typhoid vaccine, 416
changes in lipoids during, 123
cocaine and B. coli, in dogs,
blood and plasma changes
and aspirin antipyresis, 117
comparison of respiration and
glycolysis in brains of
normal and febrile rabbits,
606
effect on allergic manifesta-
tions known for centuries,
637

FEVER

effect

on reaction curve (Abderhalden reaction) of endocrines (adrenals), 179

of spontaneous and induced on liver function, 532

on vago-insulin and sympathetico adrenal systems, 318

etiology: See also Fever, pathogenesis

after intravenous inj., 139

by inj. and its real condition (in Japanese), 582

by principles in bacteria, 13

chemical basis, 401, 427

chemical character of pyrogens studied, 297, 769

cleavage products, 769

comparative view of nature of, 2

experimental contributions on, 6

nature and causes, Edinburgh theory on, 1

pyrexia or symptomatic fever as illus. nature of, 3

reactions after intravenous inj., cause, 114

excessive sensitivity of heat centers in sensitized animals, 77

experimental, 65

critical study, term 'water fever', 'salt fever', 'sugar fever', 'ferment fever' and 'tissue fever', 48

immunity, emphasis on pyrogenic materials from typhoid bacillus, 336

in sympathectomized animals, 232
peripheral circulation during, 231

from dahlia inulin, 252

filtrates of Vibrio Metchnikovi, B. pyocyaneus and yeast, 17

gastric secretions in, 201

hemolytic and water fevers, 105

high, after Coley's toxins, 772
— as index of pyrogens in water, 389

associated with shock, 530

improved method for con-

trolled hyperpyrexia with triple typhoid vaccine, 359

in rabbit, from substance in autolyzed yeast, 165

values for sugars associated with protein in serum during, 180

hunger and appetite in, 74

in animals, produced from residue of pyrogenic distilled water, 164

in rabbits after inj. of filtrates of mold cultures, 646

incidence after transfusion, 332

increased blood adrenalin, 143

induced: See also under names of bacteria, as Eberthella typhosa; Fever, therapeutic

after inj. of exts. of decomposed hay, 150

alteration of tissue during hyperpyretics, 122

analysis of mechanism, 313

body fluid redistribution in hypo- and hyperthermia, 673

by B. coli, dextrose plethora and its antipyretic effect, 91

by B. coli in dogs, blood volume and, 90

by B. pyocyaneus infection in rats, thermoregulation during, 199

by bacterial pyrogen, effect on cerebral circulation and oxygen consumption, 605

(by dinitrophenol) calcium level in arterial and venous blood in, 466

by Coley's toxins, 772

by influenzal viruses, 684

by influenzal viruses, tolerance in rabbits, 556

by infusion of hay exts. in rabbits for testing antipyretics, 202

by injd. killed bacteria, effect on blood sugar, 299

by inulin, 397

by oral admin. of amino acids in infants, 445

FEVER

induced

by physical or biological methods, 479
 by product from B. coli grown in milk, 260
 by proteins (milk, casein, albumin, edestin, hemoglobin) in rabbits, 107
 by pyripher, milk protein or bacteria, glutathione and ergothionone in blood increasing during, 207
 focal pain after pyripher, 766
 by sodium nucleate, B. prodigiosus or heat, effect on gastric secretion in dog, 79
 by subcutaneous inj. or B. coli vaccine in rabbits, 273
 by subcutaneous inj. of yeast, effect of aspirin on, 369
 by typhoid-paratyphoid vaccine, 270, 289
 by typhoid vaccine, 264
 by typhoid vaccine, and anaphylactic shock, modified by application of heat or cold to heat regulatory centers, 78
 by vaccine, effect on prodn. of agglutinins, 169
 by vaccines of B. coli or typhoid, or suspensions, changes in blood of dog, 188
 capillary flow in man in typhoid vaccine fever, 155
 cardiac output during malaise and pyrexia after inj. of typhoid vaccine, 156
 changes in lipoids in hyperpyretics, 123
 Donaggio 'inhibition phenomenon', diagnostic sign in mental symptoms, 195
 evaluation of antipyretics against, 462
 fluctuation of blood lipoids during, 325
 from E. typhosa, in rabbit, 305
 mechanism of pyrogen (Pyromen), 551
 "piretamina" admin. by various routes, effect of chemical inhibitors assessed, 524

(proteins and vaccines), 327
 'inhibition phenomenon' in urine and spinal fluid, 197
 'pyretamine' of Russo, elimination of histidine and total nitrogen during, 419
 pyripher in animal investigations, 262
 (Pyromen) quantitative study of reduction of gastric secretion during pyrexia, 625
 utilize temperature rise of rat tail in testing, 350
 varied types reproduced by bacterial pyrogen ("pyrotoxin" from B. prodigiosus), 53
 infectious and non-infectious, effect of transection of midbrain on, 55
Pyrotoxina bacterica extd. from organisms, 21
 intense, after Coley's toxins, 772
 latency of, overcome by incubation of pyrogen (Piromen) in blood, 744
 maximal temperature increase for typhoid pyrogen, 403
 mechanism of circulation and respiration in, summary, 415
 metabolism, basal, 96
 effect of toxic substances on nitrogenous and purine metabolism, 124
 nervous and artificial, research on 'anti-pyrese' and 'pyrese', 16
 oxygen consumption and glucose exchange in vitro of liver slices from febrile rabbits, 707
 pathogenesis, theory, 67, 174
 role of diastases or soluble ferments, 14
 pathology, process, 8, 9
 pharmacologic evaluation of technical materials, 40
 physiology, thyrioid and adrenals in, 145
 protein, pharmacology of, 175, 769

FEVER

reactions: See also Pyrogens,,
reactions

in hypophysectomized or tuber-
injured dogs, 184

relation of proper prepn. of
solns. for intravenous
therapy, 314

relation of peripheral to
internal temperature in, 10
saline, 38, 48, 52, 173

pharmacology of, 175

symptom-complex, leukopenia,
chills and gastro-intestinal
disturbances, 277

therapeutic: See also Para-
typhoid, vaccine; Pyrifer;
Pyrogens; under diseases, as
Arthritis; Neurosyphilis;
Paralysis, General, Tumors; etc.
ACTH-like effects in man, 661,
659, 715, 774, 775

————— in malignant
hypertensive conditions,
(Pyromen, typhoid vaccine and
physically induced), 661

alterations in blood provoked by
pyrifer in schizophrenics, 357

and leukocytosis, using purified
typhoid pyrogen, 446

and psychoses, 292

artificially induced, effect on
circulation in arthritics, 550

————— effect on
electroencephalograms in mul-
tiple sclerosis, 713

biological (typhoid vaccine) in
inflammatory rheumatism, 489

(cabinet) did not reduce
responsiveness to pyrogens, 470
colloidal metals, 76

continuous typhoid vaccine with
penicillin in symptomatic
paresis, 594

danger in therapy (malaria,
typhoid vaccine) of paresis, 609

dangers with typhoid vaccine, 516

effect of shock therapy in acute
rheumatic and allergic diseases,
675

effect on cerebral functions in
neurosyphilis, 653

effect on nonspecificity in
syphilis reaction, 308

————— plasma protein, 351

expanding dosage method
for prodn. with typhoid
vaccine, 331

from infusion of blood from
pregnant women, new
treatment of rheumatoid
arthritis, 465

hypotensive effect of sub-
stances eliciting, 407

in arthritis or infections,
analogy between typhoid
and malarial paroxysms, 82

in cancer, 120

in dementia praecox using
Dmelcos (Ducrey bacillus
atoxic vaccine) or
sulfosin, 230; or typhoid
vaccine, 760

in dermatology, clinical
evaluation of new pyrogenic
agent (Pyromen), 660

in early therapy of acute
anterior poliomyelitis, 641

in herpes zoster, (typhoid
vaccine), 473

in ophthalmology, 774, 775

in poliomyelitis, with
pyrifer, 263

in therapy of leukemias, 424

increased efficacy of drugs
by using combined intrav-
enous and intracutaneous
application, 669

leukocytic modifications in-
duced by, in acute and
chronic inflammatory
diseases, 640

(malaria or typhoid vaccine)
as globulin plasma increases
sedimentation rate heightens,
352

mechanisms of reactions of
nonspecific proteins, 110
metabolic rates in neuro-
syphilis, 229

method of inducing with
typhoid vaccine, intra-
venous drip technic, 375

method of 'pyrogenous' syphilis
therapy, 698

FEVER

therapeutic

- new pyrogen (Pyromen), 593
- nonspecific protein (chemical and physiological mechanisms), 228
 - _____ history reviewed, 226
 - _____ in ocular disease, 774, 775
 - _____ in rheumatism, 191
- of acute poliomyelitis, 664
- of disease of the eye, 775
- of diseases of peripheral vascular system, 223
- of gonorrhea using pyrifer, 185
- of neurosyphilis, (dementia paralytica and tabes dorsalis) using suspensions of killed Ps. scissa and ureae, 208
- of ulcerative colitis (pyrifer), 656
- physical hyperthermia or fever induced by pyrifer, 676
- physiologic changes during reactions to protein therapy, 190
- produced by physical and bacterial agents, 254
- prolonged fever produced by three injs. of typhoid vaccine, 668
- pyrifer, body structure and temp. variations, 712, 761
 - _____ in gonorrhea, psoriasis, progressive paralysis, tabes and vulvovaginitis, 171
 - _____ in paresis, syphilis and other infections, 260
 - _____ in schizophrenia and paralytic dementia, 240
 - _____ in severe diphtheria, 694
 - _____ in neurosyphilis, manner of action, 181
- pyrogen in early hebephrenia, 221
- remission in rheumatoid arthritis after fever therapy (typhoid bacteria) with liver damage, 696

- Saprovitan in neurology, 121
- soduku, rat-bite fever in general paralysis, 127
- TAB in choreic encephalitis, 547
- typhoid shock, 461
- with Neosaprovitan B in tabes dorsalis and multiple sclerosis, 168
- therapy, adrenals extract in, 368
- research in, 8
- with pyrifer, 178
- thermostable bactericidal substances in serum from febrile patients, 256
- toxin, chromatin of bacterial cell site of origin of fever, 21
- of bacteria, 18

FIBROSITIS: See Rheumatism

- FILTERS, adsorptive and 200-second Zsigmondy, 251
- asbestos, Seitz EK and 200-second Zsigmondy membrane filter, 250
- Berkefeld, 39
- Reichel, 92

FILTRATES: See under Bacteria and names of bacteria

- FILTRATION, adsorptive, 251, 252
- adsorptive, in prepn. of pyrogen-free solns., 250, 251, 341, 505
- with asbestos filter pads for removal of pyrogens from penicillin, 432, 434
- method of sepn. of hemorrhage-prodg. fraction of B. coli, 244
- Selas candle for detoxification of pyrogens in dextrose-saline solns., 548
- studies on reactive infusion fluids, 238
- use of Seitz filter pad and ion exchange agent "Decalso" for removal of pyrogens from protein solns., 509

- FINNISH METHOD, pyrogen reactions to solns. prepd. by, as prescribed by U. S. P., 711
- FLAVONOID COMPOUND, effect of Shear polysaccharide alone or with, on rat carcinoma, 750
- FLOCCULATION, reaction, effect of calcium on, with C polysaccharide of Pneumococcus, 315
- FLUIDS: See also Protein, solutions; Salt, solutions; Solutions
body, redistribution in induced hypothermia and hyperthermia, 673
citratd: See Blood Transfusion, citrated blood in
intake, reduces fever, 81
- FLUORESCENCE, fate of capsular polysaccharide of Friedländer's bacillus labelled with fluorescein, 654
method of detecting pyrogenic substances in soln., 648
- FOAM-REDUCING AGENT, 272
- FOOD, allergy, Pyromen therapy, 738
- FUNGI: See also Sporotrichum
cultures, filtrates of, pyrexia in rabbits after inj., 646
simplified method of detecting pyrogens in soln., 648
- GALACTOSE, d-, in fraction from tubercle bacilli, 216
tolerance, altered during hyperglycemic reaction, 259
- GANGLION, basal, radiotherapy in late evolution of choreic encephalitis toward parkinsonism, 547
- GASTRIC JUICE: See Stomach, secretion
- GASTRO-INTESTINAL TRACT: See also Intestines; Rectum; Stomach; etc.
disturbances, from typhoid vaccine, or pyrogens from water of infusion solns., 275
prodn. of pyrogen in sera by bacteria, 277
- GELATIN, pyrogenic solns. made nonpyrogenic by heating with KMnO_4 or H_2O_2 , 421
- GLIA: See Neuroglia
- GLOBULIN, immune, (Felix anti-typhoid serum) neutralized typhoid leucocidin, 253
in blood: See Blood, globulin
- GLUCIDES, glucido-lipid antigen of E. typhosa, leukopenic ability, 266, 267
- GLUCOSAMINE, from biosynthesis of capsular streptococcal polysaccharide, 753
- GLUCURONIC ACID, 133, 753
- GLUTATHIONE, in blood: See blood, glutathione
reducing substances as indicators of protection by cortisone against tumor-damaging S. marcescens polysaccharide, 697
- GLYCOGEN, depletion of organs during fever in animals, 7
in liver: See liver
lysis, splanchnic nerve stimulation initiates, thyroidectomy decreases it, 132
starch, (polysaccharide) effect upon bacteriophagy, 295
synthesis, effect of bacterial endotoxins on, 534
- GLYCOLYSIS: See Brain; Glycogen, lysis

GONOCOCCI, endotoxin, protection of mice against lethal action with penicillin, 451

GONORRHEA, resistance to endotoxins of enteric bacilli, 539
therapy, in women, with pyrififer, 161
method of inducing fever with typhoid vaccine using intravenous drip technic, 375
use of TAB vaccine in N. Africa and Central Mediterranean, 391
with pyrififer, 171, 1781 185

GRANULOCYTES, reduction by typhoid leucocidin, 253

GUINEA PIGS, action of toxic doses of polysaccharide from S. marcescens, 394
anaphylaxis in, 395, 616
carbohydrate haptens in bacterial anaphylaxis, 153

GUM ACACIA, (Arabic): See Acacia

HAGEDORN-JENSEN TEST: See polysaccharides

HAIR, erection in cat injd. with bacterial pyrogen, 481

HAPTEN, bacterial polysaccharides adsorbed on colloids (as haptens) as antigens, 194
carbohydrate, role in bacterial anaphylaxis, 153
intravenous inj. into test animal, 259
polysaccharide, phage-inactivating agent of bacterial, converted by alkali into, 215

HARVEY LECTURE, physiology of renal circulation, 291

HAY, infusion, inquiry into fever after, 150
infusion, of exts., induced fever in rabbits for testing antipyretics, 202

HEADACHE, after therapy of arthritis with typhoid vaccine, 66
during pyrogenic therapy with colloidal metals, 76
experimental studies, 313
work with ACTH and Pyromen, 626
reaction to pyrififer, 185
therapeutic fever (typhoid pyrogen) and leukocytosis, 446

HEART: See also Blood circulation; Cardiovascular System; Endocarditis; etc.

cardiac output during malaise and pyrexia after inj. of typhoid vaccine, 155
electrocardiogram, effect of sublethal doses of S. marcescens polysaccharide, 396
failure, congestive, improved in therapy of arterial hypertension with pyrogen (Pyromen), 603
congestive, relieved during therapy with bacterial pyrogens, 549
pyrogen (Pyromen) therapy in cardiac decompensation, 604
hypertrophy, in induced fever therapy (typhoid vaccine), 463
rate, sinus tachycardia after sublethal doses of S. marcescens polysaccharide, 396
size, decreased during therapy with bacterial pyrogens, 549

HEAT: See also Temperature
application to animals, effect on leukocyte count, lymphocytes and lymphoid tissue activity, 85
to dogs, effect on gastric secretion, 79
to heat regulating center modifies action of induced pyrexia and antipyretics, 78
dry, effect of stimulation of lymphocytes on rate of growth of tumors in mice, 87

HEAT

dry

stimulation of lymphocytes

increases immunity to trans-
plantable tumors, 86

regulation, and water exchange, 117

HEBEPHRENIA: See Dementia Praecox

HEMATOPOIESIS: See Blood, form-
ation; Hematopoietic SystemHEMATOPOIETIC SYSTEM: See also
Blood, formation injury, by
toxic antigen of E. typhosa,
376HEMOGLOBIN: See also Blood;
Erythrocytes; Hemolysis pro-
duction of fever with, in
rabbit, 107HEMOLYSIS, antihemolytic substances,
176
fever, 105HEMOPHILUS, influenzae, chemical,
physical and immunological
properties of polysaccharides
of, 450
influenzae, fever toxins of, 18
injection in mice modified course
of virus pneumonia, 495
production of hemorrhagic
necrotic skin lesions with,
247
vaccine, prevention and therapy
of influenza and sequels, 98
of Ducrey, atoxic vaccine
(Dmelcos) use in dementia
praecox, 230
pertussis, production of hemo-
rrhagic necrotic skin lesions
in rabbits with, 247HEMORRHAGE: See also Rectum,
hemorrhage; Shwartzman
Phenomenon; Tumor, hemorrhage
adreno-hemorrhagic effects of bac-
terial vaccines, 356
effect of ascorbic acid upon, 275
of bacterial filtrates on trans-
plantable rat tumor, 280of meningococcus filtrate in
mouse sarcoma, 180, 266

immunological protection

(B. paradysenteriae toxin)
against tumor hemorrhage,
386induction in mice, effect of
colchicine and B. coli
filtrate on tumors, 294in mouse tumors, use of
sulfonamide in toxicity
after, 349local skin reaction to B.
typhosus culture filtrate,
152

of sarcoma, 255

—producing bacterial prepns.,
bioassay method for detg.
potency, 380
fraction from B. pro-
digiosus filtrates, sepn.
of, 312fraction of S. marcescens,
analysis, 374polysaccharide, S. marces-
cens, inj. of mice with
subcutaneous tumors, 406—production in mouse sarcoma,
nature of polysaccharide, 372
production, tumors, effect
of bacterial toxins on, 433
chemical composition of
fraction from S. marcescens
(B. prodigiosus), 371toxin---induced, effect of sul-
fonamide on, 364tumor, action of endotoxin
antigens of gram-negative
bacteria, 363
—— agents, distribution
among bacterial species, 373
—— effect of colchicine
and B. typhosus exts. on, 249
vascular reactions of normal
and neoplastic tissue to
bacterial polysaccharide, 467HEMOSTASIS, in typhoid vaccine
fever, 155HEPARIN, 721
effects, 650

HERPES, 140

- labialis, reaction to pyrifery therapy, 185
- question of appearance in arthritis of childhood, 166
- roseola, disappearance after inj. of anti-typhoid vaccine in E. typhosa infection, 517
- zoster, therapy with bactifebrin, 765; typhoid vaccine, 473

HEXOSES, 32% reducing sugars contained in glucosamine from bacterial pyrogens, 430

HIGH FREQUENCY CURRENTS. fever therapy in asthma, 637

HISTAMINE: See also Antihistamines
in blood: See Blood, histamine
in liver: See Liver, histamine
in muscles: See Muscles, histamine

HODGKIN'S DISEASE, daily variation in leukocytes, 44
therapy, end results with mixed toxins (Coley's) and/or radiation, 149
reactions of patients and tumors to inj. of S. marcescens polysaccharide, 494

HORMONES: See also Adrenals; Endocrines; Pituitary Body; Pituitary Preparations; Thyroids; under names of synthetic hormones
• extraction, pyrogens removed by cyclic or heterocyclic acids, removal with EtOH and BzOH, cooling and filtration, 570
therapy of neoplastic ailments, 590

HUMIC ACID, in ext. of decomposed hay, 150

HUMORS: See also Fluids, body
control of pyrogenic renal hypermia, 348

HUNGER: See also Fasting; Starvation
in fever, 74

HYDROCHLORIC ACID: See Stomach, secretion

HYDROGEN, peroxide, depyrogenization of infusion solns., 421, 719
detoxification of pyrogens in some U.S.P. inj., 548

HYDROGEN ION CONCENTRATION, 152
effect on potency of hemorrhage-producing fraction from S. marcescens, 381
factor in oxygen consumption and glucose exchange in vitro of liver slice from febrile rabbits, 707
of dextrose-saline solns. in prepn. of nonpyrogenic fluids, 459

HYDROLYSIS, 272, 372
enzymatic, of capsular polysaccharide of Klebsiella pneumoniae, type B, 554
of antigen of E. typhosa, 319

HYPEREMIA, local, influence upon storage process in lymph nodes, 151

HYPERPNEA; See Respiration

HYPERPYREXIA: See Fever

HYPERTHERMIA: See Fever

HYPOPHYSIS: See Pituitary Body

HYPOTENSION: See Blood Pressure, low; Blood Pressure, high

HYPOTHALAMUS: See also Pituitary Body
lesions, effect on fever induced by typhoid-paratyphoid vaccines, 289
in cats, site of action of bacterial pyrogen, 481
medial, influence of transection on infectious and non-infectious fever, 55
stimulation of fever control center of, mode of action of 'necrosin', 427

HYPOOTHERMIA: See Temperature,
Body, low

HYSTERECTOMY: See Uterus, tumors

ICE CREAM, 'thyrotoxicon', bio-
logical split protein from,
57

IMMUNITY: See also Anaphylaxis;
Antibodies; Antigens; under
diseases, as Typhoid, immunity
action of S. paradysenteriae
toxin on tumors, 386
antileucopenic, 326
appearance in exptl. fever,
emphasis on pyrogenic materials
from typhoid bacillus, 336
chemical nature of immune sub-
stances, 134
effect of intracellular contents
of typhoid bacilli, 31
effect of passive and active
immunization on lethal act-
ivity of polysaccharides, 565
of reticuloendothelial blockade
on immunity to Shwartzman
phenomenon, 472
endotoxic factors of Rickettsia
Prowazeki and immunological
relationship to endotoxins
of other gram-negative bacilli,
453
filtrates of Streptococci,
Staphylococci, E. typhosa,
and B. coli related to immuno-
logic factors in bacteria, 176
heterologous, protection against
tumor hemorrhage, 387
illness in man after inhalation of
S. marcescens, 455
immune bodies in fever therapy of
diseases of peripheral vas-
cular system, 223
lymphocytes in natural and induced
resistance to transplanted
cancer in mice, 86
passive, against lethal activity
of polysaccharide (S. marces-
cens), 566

preparation of fraction from
B. aertrycke, 222
properties of S. marcescens
polysaccharides, 484, 519,
616
specific polysaccharides of
types A,B,C,D, and F H.
influenzae, 450
protection against endotoxins
of gram-negative bacteria,
412
protein split products in
relation to, 57
reactions to blood and plasma
transfusions in multiple
sclerosis, 513
relation between S. marcescens
culture filtrates and Shear
polysaccharide, 405
relationships of typhoid-
salmonella, 287
studies in immunochemistry,
242, 324
substances in pneumococci,
distribution of poly-
saccharide in mouse tissue
after inj., 488
susceptibility to bacteriophagy
depends upon specificity, 295
'third immunity' of Centanni
has potentialities in
therapy, 336
thrombocytes play important role
in, 118

IMPOTENCE, effect of pyrifur in
therapy of tabes, 167

INCUBATION, latency of response
to bacterial pyrogens as
affected by previous in-
cubation with whole blood,
744

INDUSTRY: diseases, acute ill-
nesses among rural mattress
makers using low-grade
stained cotton, 354
diseases, 'cotton fever', 402

INFANT, feeding, addition of
protein conc. to milk
formula increased body
temperature, 126

INFANT

feeding

fever produced by oral admin.

of amino acids, 445

so-called 'thirst' or 'inani-
tion' fevers and water reserve
of body, 81

INFECTIONS: See also various dis-
eases; under names of bacteria,
as Pneumococci; Staphylococci;
etc.

blood lipoids fluctuation during
fever, 325

polysaccharide level in, 615

protein in acute, 315, 322

thrombopenia, 118

experimental, changes in leuko-
cytes and alkali reserves of
blood, 99

gastric secretion in fever and, 201

pathogenesis, theory of fever
result of internal destruction
of proteins, 67

putrid or septic, 4

reactivation of neurologic syn-
dromes during, 767

reticuloendothelial cells reacting
to, 480

susceptibility, development in
radiation injury, 752

therapy with foreign protein in
acute, 100

pyrifer, product from B. coli
growth in milk, 260

with typhoid vaccine, 82

with vaccines, 64

INFLAMMATION: See also Neuritis ;

Uvea, inflammation;

Vulvovaginitis

chemical basis of fever with, 427

experimental, (bacteria, turpentine
or talc) serum polysaccharide
levels in, 614

leukocytic modifications induced
by pyretotherapy, 640

local reaction in non-specific
protein therapy, 101

therapy of, in uveal tissue, 774

thermostable component in
mechanism of leukocytosis,
597

INFLUENZA, bacilli: See

Hemophilus Influenzae

sequels, prevention and therapy

with mixed vaccines, 98

therapy with mixed vaccines, 98

virus, cultivation in allan-
toic fluid, 525

effect of polysaccharides on
reaction between erythrocytes
and viruses, 526

production of fever by, 556, 684

pyrogenicity in rabbits, 557

INFUSIONS: See Injections;
Solutions

INHIBITION PHENOMENON, Donaggio's,

in urine and spinal fluid

of neuro- and psychopathic
patients, diagnostic sign,
195

in urine and spinal fluid,
under febrile conditions,
197

INJECTIONS: See also Dextrose,
solutions; Fever, induced;
Fever, therapeutic; Salt
solution; etc.

bacterial cellular elements
and toxins produced
ischemia after, 344

fever after, 5

reaction in animals, relation
of inj. sites of pyrogens,
726

relation of salvarsan fever
to other forms, 49

intracutaneous, increased
efficacy of fever-producing
drugs by using combined
intravenous and, 669

of antigen of E. typhosa,
inflammation and necrosis
in rabbits after, 376

of aspecific vaccine in
poliomyelitis, 629

of B. abortus filtrate in
cattle, 400

of H. influenzae in rabbit,
247

of meningococcal filtrate,
prodn. of Shwartzman
phenomenon, 347

INJECTIONS

intracutaneous

- of pneumococcal somatic C polysaccharide, 248
- of toxic E. typhosa filtrates, 152
- of water, milk or salt soln. or air caused leukopenia and conc. of leukocytes in liver, 112

intramuscular, comparison of inj.

- of pyrogenic solns. by intravenous and, 647
- of concd. B. coli antitoxin in ulcerative colitis, 361
- of Dmelcos (Ducrey bacillus atoxic vaccine) in dementia praecox, 230
- of "mixed toxin" from St. erysipelatis and B. prodigiosus in ocular infections, 491
- of pyrifer in neurosyphilis, 163
——— ineffective when given by, 178

intranasal, of living and dead bacteria and bacterial polysaccharides, modified course of virus pneumonia, 495

intraperitoneal, minute doses of B. prodigiosus filtrate into mice with sarcoma #37, 275

- of bacterial pyrogen (Pyromen) in mice, 679
- of bacterial pyrogens and cortisone in mice, 732
- of ethyl carbamate (urethane), effect on temperature regulation, 578
- of gonococcal endotoxin in mouse, 451
- of S. marcescens polysaccharide, 396; alone and with adrenal cortical ext. in tumor-bearing mice, 522; effect on rectal temperatures in mice, 439; effect on sarcoma in mice. 37, 515; vascular reactions of normal and neoplastic tissues to, 467
- of toxic filtrates of E. typhosa in rabbit, 271
- of washed antiplatelet serum, from guinea pig, 97

intravenous, 252, 284

- allantoin in Ringer's soln, role of pyrogen in alleged leukocytic response, 410
- antityphoid vaccine, 517
- cause and elimination of reactions after, 164, 205
- calcium chloride, for chills, 296
- cause intense febrile reactions, 114
- compared with intramuscular route of, 647
- continuous, typhoid vaccine, in ophthalmologic disease, 567
- dangers, with typhoid vaccine, 516
- drip technic for, 375
- economy in use of, 234
- efficacy of fever production by combined intracutaneous and, 669
- ethyl carbamate (urethane) effect on temperature regulation, 578
- in shock produced by S. marcescens administration, 494
- index of pyrogenicity in, 409
- loss of pyrogenic activity in solutions prepared for, 610; in storage, 482, 493, 501, 634
- meningococcal endotoxin, effect on histamine content of blood and tissues, 497; and filtrate, 255
- method of preparing nonpyrogenic solutions, 241, 246, 250, 251
- pneumococci, polysaccharide from, 362
- protein, carbohydrate and fat, effects on blood components, 137
——— foreign. See also: Pyrogens, Vaccines
——— in articular arthritis, 75; in neurosyphilis, 431; tuberculo, 257
- proteose and typhoid vaccine, 104

INJECTIONS

- intravenous
 - Proteus vulg. polysaccharide (Pyromen), 715
 - 'pyretamine' of Russo, 419
 - pyrogens, elimination, 204, 206, 246, 314, 327, 370, 505, 599
 - for inhibition of gastric secretions, 580
 - from typhoid vaccines, febrile response to, 585; loss of activity, 482, 493, 501, 610, 634
 - Pyromen, in food allergies, 738; effect on adrenals and blood components, 720
 - S. marcescens, polysaccharide from, effect on renal temperatures (mice), 439
 - S. typhosa, antigen, and pathology after, 376
 - endotoxin, vascular response in rabbit, 627
 - S. typhosa, toxins, 271
 - S. typhosa, vaccine, blood picture after, 619; cardiac enlargement after, 463; fatality after, 426; neurosyphilis benefited by, 605; renal blood flow after, 264
 - typhoid-paratyphoid vaccine, and other pyrogens, 579; thermostatic behavior in body, 529; as "pyrexin", 577; temperature increases after, 270
 - use of rabbits for testing pyrogens in, 337
 - water for, nonpyrogenic, 204, 246, 505
- bacterial filtrates, vascular shock after, 280
- endotoxins of histolytic bacteria (in rabbits), 169
- ext. of E. typhosa in rabbit, action on arterial pressure, 476
- polysaccharide, reaction of patients with malignant tumors to, 390
- preparation, dextrose and saline solns. for, 206; safe sol., 246
- pyrogens in injectable solutes, 327, 505, 599; bio-assay according to French Pharmacop., 635
 - elimination, 204, 206, 246, 314, 327, 505, 599; by activated charcoal, 327, 422; by ion exchange resin, 370
 - nature of, 700
 - technics of U. S. Pharmacop. test, 571
 - tests based on leukocytosis of rabbit, 633
 - for, 699
 - pyrogens, prepn. of nonpyrogenic saline fluids, 457, 459
 - reactions and preventions, 505, 652
 - site or route, not substance, produce phenomena seen in parenteral therapy, 112
 - subcutaneous, allantoin in Ringer's soln., 410
 - ascorbic acid, 275
 - bactericidal activity following administration, 186
 - E. coli vaccine in rabbits, 273
 - hay exts. in rabbits for testing antipyretics, 202
 - Pyrotoxina bacterica, chemotactic reaction, 21
 - S. marcescens polysaccharides in dog, 296
 - typhoid anaphylatoxin in rabbits, 51
 - yeast to produce fever, 'aspirin' effects, 369
 - sub-patellar, of typhoid antigen in rabbits, changes induced by, 500

INSANITY: See also Dementia
 Praeccc; Mental Diseases;
 Paralysis, General
 manic-depressive, induced
 fever and, 292
 therapy with pyrifer, 217

INSULIN, diminished inhibition
 to glycogen synthesis, 534
 effect on blood sugar during
 induced hyperthermia, 299
 ——— on glycogen reserves in
 liver, 273

INSULIN

- influence of fevers (typhoid-paratyphoid vaccine) on vago-insulin and sympathetico-adrenal systems, 318
- like substances from B. coli and Blackleg bacteria (Clostridium chauvei), 154
- zinc-free, addition to meningococcal endotoxin prevents inhibition of glucose utilization, 535

INTERFEROMETRY, increase in reading for Abderhalden reaction in fever, 179

INTESTINES: See also Cecum; Colitis; Gastro-Intestinal Tract; Rectum; etc.

- adenocarcinoma, effect of S. marcescens polysaccharide, 444
- ascorbic acid content, effect of colchicine and E. typhosus ext. on, 249
- bacteria, resistance to endotoxins of enteric bacilli in man, 539
- tolerance to toxic action of antigens of enteric, 540
- epithelium, damaged by S. marcescens, 443
- mucosa, relation of pyrogen to Dolman test for enterotoxin, 397, 398

INULIN, clearance, pyrexial reaction to typhoid vaccine, 264

- colloid, inactivation of coliphage, 317
- dahlia, toxicity, 252
- and fever prodn., 397
- induced renal hyperemia, expt. to determine neural or humoral control, 348
- small doses resulted in renal hyperemia in essential hypertension, 320

IODINE, effects of iodinated S. marcescens polysaccharide on metastatic carcinoma, 508

-iodide test, 321

- radioactive, tagged tumor-necrotizing polysaccharide S. marcescens, 545

IODOBENZENE DIAZONIUM, para--, chloride, radioactive, reaction of polysaccharide with, 613

IODOPOLYSACCHARIDE: See under polysaccharides, from S. marcescens

IONS: See also Chloride; Hydrogen Ion Concentration

- "decalso" ion exchange agent for removal of pyrogens from protein solns., 509
- electrophoresis analysis of hemorrhage-producing fraction from S. marcescens, 374
- gave an iso-electric point for nucleic acid component of polysaccharide of pneumococcus after fractionation, 512
- ion exchange resins for reduction of pyrogens in mineral free water and parenteral solns., 370

IRIDOCYCLITIS, postoperative, therapy with "mixed toxin" from St. erysipelatis and B. prodigiosus, 491

IRON, hydroxide, elimination of pyrogens by, 592

ISCHEMIA, renal: See Kidneys, blood supply

ISOPYRENE, required dosage to counteract reaction from bacterial polysaccharide, 438

JOANNOVIC'S tumor tissue split products: See Cancer, epithelioma

- KAOLIN, pathogenizing ability of carbohydrates enhanced by, 454
removal of inhibiting substances by, 197
- KATZMAN-DOISY-method of preparing pyrogen-free urogastrone, 302, 303
- KIDNEYS, blood supply, changes during pyrogenic therapy of essential hypertension, 420
blood supply, control of flow, 264
effect of denervation on hyperemia, 348
effective flow in subjects with essential hypertension, 320
hemodynamic changes induced by standardized pyrogen, aging as factor in, 718
in therapy of arterial hypertension, 603
increase in plasma flow, response to pyrogens, 397
physiology of circulation, 291
pyrogen-induced hyperemia, prevention in dog by dihydroergocornine, 681
response to pyrogens, 397
ischemia, action of typhoid vaccine, 264
diseases, pyrogens in therapy of lower nephron nephrosis, 612
dysfunction, in typhoid vaccine therapy of paresis, 609
severe irritation from foreign protein (typhoid) therapy of hypertension and Bright's disease with nephrosis, 382
function test, filtration rate, diodrast clearance, in essential hypertension, 320
in essential hypertension treated with pyrogens, 420
innervation, control of pyrogenic hyperemia, 348
necrosis, after fatal reaction from therapeutic anti-typhoid inj., 426
in cortisone-treated rabbits after inj. of bacterial toxins (S. marcescens), 734
- physiology, control of blood flow and glomerular filtration 264
creatinine and creatinine clearance in therapy of carcinoma with S. marcescens polysaccharide, 464
decrease in filtration fraction, response to pyrogens, 397
excretion of para-amino hippurate, in selection of patients for therapy with pyrogens, 604
neural vs. humoral control of pyrogenic hyperemia, 348
of circulation, 291
response to pyrogens in hypertension and hypertensive diseases, 397
studies of severely burned patients, 714
- KLEBSIELLA pneumoniae: See Bacteriophage, Friedländer's; Polysaccharides, from B. Friedländer
removal from blood stream, after Piromen, 762
- LACTATE, utilization during induced fever in rabbit, 273
- LANDRY-GUILLAIN-BARRE Syndrome, effect of fever therapy, 765
- LANGERHANS' ISLANDS: See Pancreas
- LEGS: See also Extremities
pain, from induced fever with purified pyrogen, 446
- LEUCOCIDIN, activity of typhoid toxic fraction on rabbit's blood and non-immune human blood, 253
glucose, 19.9% in, 253
- LEUKEMIA, anticoagulant effect of bacterial polysaccharides in normal and thrombocytopenic plasma of, 743

LEUKEMIA

therapy, fever therapy (typhoid vaccine) or colchicine, to enhance effectiveness of X-ray therapy, 424

LEUKOCYTES: See also Blood, cells; Leukemia; Lymphocytes; Phagocytes; Phagocytosis; etc.

accumulation of neutrophils in lung, liver, spleen and bone marrow, 271

chemotaxis of, 316

inj. of bacterial toxins inhibit, 365

count, after intramuscular inj. of pyrif, 163

augmentation of effects of typhoid vaccine by homologous plasma in rabbit, 523, 636

bacterial constituents responsible for response, 342

changes in adrenalectomized and normal rats after admin. of typhoid vaccine, 537

—— induced by pyreto-therapy in acute and chronic inflammatory disease, 640

daily variation in malaria, Hodgkin's disease and cancer, 44

during pyrif therapy of neurosyphilis, 181

effect of dry heat applied to animals, 85

—— inj. of bacterial exts., 35

measure of Pyromen activity, 754

mechanism of reaction to non-specific protein therapy, 110

reduced by typhoid leucocidin, 253

rise and fall after inj. of bacterial antigens, 259

variations during pyrif admin., 712

degeneration of neutrophils by typhoid leucocidin, 253, 279

effect of bacterial by-products, 172; carbohydrates, 717

protein, carbohydrates and fats, 137

vaccine injections, 64

Polysaccharide (Shear) 776

Str. scarlatinae cultures, 172

tuberculin protein fractions and polysaccharides, 749

typhoid toxic fraction, 253

morphologic alteration by polysaccharide (Shear) 776; by pyrif, 514

picture, effect of inj. of filtrates of B. abortus in cattle, 400

--producing factor of Menkin, 407

qualitative and quantitative proportions of, in Neosaprovitan B. fever in therapy, 168

response, adrenals in mechanism of changes upon admin. of Pyromen, 620

comparison of ACTH and bacterial pyrogen (Pyromen), 728

to bacterial pyrogen (Pyromen), mechanism, 727; suppression by N mustard, 735

LEUKOCYTOSIS: See also Leukocytes, count

effect of phenobarbital admin. before pyrif and nucleic acid, 269

of pyrogens, characteristic, 379, 759

from admin. of colloidal toxic bacterial substances in septicemia, 70

allantoin in Ringer's soln., role of pyrogen, 410

antigens of E. typhosa, 323

from bacterial products, 17

bacterial pyrogens in hypertension, 549

induced fever with purified typhoid pyrogen, 446

inj. of distilled water, 140

oral admin. of bacterial pyrogen, 631, 702

pyrif in schizophrenics, 357

pyrogenic therapy with colloidal metals, 76

therapy with E. coli vaccine, 83

typhoid vaccine therapy of arthritis, 75, 82; infections, 82

LEUKOCYTOSIS, hypotensive effect
 of substances eliciting, 407
 in therapy, chancroid, 142
 ————— diseases of peripheral
 vascular system, 223
 ————— E. typhosa infection,
 with typhoid vaccine, 517
 ————— nonspecific protein,
 101
 index of pyrogenicity in fluids
 for intravenous use, 409
 neutrophilic and adrenals, 146,
 776
 normal alkali reserve parallels,
 in exper. infections, 98
 oral administration of methylene
 blue or yatren, produce, 103
 search for reliable and
 sensitive method for determin-
 ation of pyrogenicity in
 rabbits, 417
 test for pyrogens in injectable
 solutes based on, 633
 thermostabile factor of exudates,
 663, 597

LEUKOPENIA: See also Leukocytes,
 count
 ability to produce by glucido-
 lipid antigens of E. coli or
 typhoid, 367; of E. typhosa,
 366
 cutaneous vascular reactions in
 rabbit to typhus endotoxin, 477
 effect of bacterial pyrogens, 278,
 379, 759
 of dysentery toxins, 447
 of fractions from B. dysenteriae,
Shigae, Flexner and Y, 259
 experimental infections and, 99
 fever-producing principle in
 typhoid vaccine, 276
 from antigens from E. typhosa, 323,
 363, 366
 bacterial products, 17
 bacterial vaccines from various
 species, 356
 dahlia inulin, 252
 hay infusion fever, 150
 intravenous inj. of killed bacteria
 in rabbit, 116
 oral admin. of methylene blue
 or yatren, 103

protein fraction of B. 341
dysenteriae, 258
 pyrogenic factor (pyrexin), 597
 toxic filtrates of E.
typhosa in rabbit, 271
 typhoid vaccine, 72, 82; in
 arthritis, 75, 82
 immunity, effect of bacterial
 antigens on, 326
 in splenectomized rabbits, 80
 initial, as indicator of
 presence of pyrogen, 337
 production of pyrogen in sera
 by bacteria, 277
 tolerance to leukopenic effects
 of typhoid vaccine, 601

LIPASE: See Enzymes

LIPIDS: LIPINS: See Blood, fats;-
 Cholesterol; Phospholipin
 changes during fever and
 hyperpyretics, 123
 chemical nature of immune
 substances, 134
 content of antigen of E.
typhosa, 312
 effects upon leukocytes of inj.
 of, 137
 fractions of B. dysenteriae
 produced hyperthermia, 258
 glucido-lipid antigen, of E.
typhosa, leukopenic ability,
 366; of E. coli or typhoid,
 leukopenic ability, 367
 chemotactic power of antigens,
 316
 in blood: See Blood, lipoids
 lipo-protein-nucleic acid
 complex (reticulose) in
 therapy of radiation injury,
 752
 polysaccharide-lipid-protein
 complexes from Salmonella
 group, 611

LISTERELLA, monocytogenes: See
 Bacteria, Listerella

LIVER, accumulation of neutrophils
 in, after inj. of E. typhosa
 filtrate, 27

LIVER

acute and chronic involvement of parenchyma, therapy with pyrifer, 686

ascorbic acid content, effect of colchicine and E. typhosa ext. 249

blood supply, estimated flow and bromsulfalein extn. during pyrogenic reaction. 478

cirrhosis, influence upon febrile response to typhoid pyrogen, 585

damage, after fever therapy (typhoid bacteria) of rheumatoid arthritis, 696

denervation, effect on fever prodn., 145

detoxification, failure in dementia praecox, 760

dessication during fever, 174

extracts from fasting and feverish animals, 7

function test, abnormal retention of bromsulfalein during febrile periods, evaluations, 532

glycogen, altered in hyperglycemic reaction, 259

decrease, action of dysentery toxins, 447

decrease after intra-abdominal inj. of killed Rickettsial bacilli, 453

effect of bacterial endotoxins on carbohydrate metabolism, 534, 536

——— inj. of protein-polysaccharide fraction from S. typhimurium, 297

intravenous admin in rabbit, 730

reserves, effect of insulin inj. on, 273

heat production, oxygen consumption of liver slices (in vitro) from febrile and normal rabbits, 581, 707

hemopoiesis in rabbit induced by bacterial pyrogens (Pyromen) 692

histamine, effect of meningococcal endotoxin on, 497

inflammation of hyperthermia, 479

necrosis, after fatal reaction from therapeutic anti-typhoid inj., 426

anoxia or destruction of enzyme systems may underlie, 479

physiology, effect of spontaneous and induced fever, 532

venous catheterization, 478

LUNGS: See also Respiration; Respiratory Tract

accumulation of neutrophils after inj. of E. typhosa filtrate, 271

pyrogenic infections, method for controlled hyperpyrexia, 359

tumors, effect of S. marcescens polysaccharide, 444

LUTEIC ACID, metabolic product of penicillin luteum, 274

LUTEOSE, composed of beta-glucose unit linked through 1,6-positions, a neutral polysaccharide, 274

LYMPH, flow during protein shock reactions, 68

LYMPH NODES, induced development of plasma cells, 257

irradiation, invoked fever (typhoid vaccine) enhances effect of, in therapy of leukemia, 424

mesenteric. intensified prodn. of cells after infection with B. monocytogenes, 265

storage process in, influence of local hyperemia, 151

LYMPHANGITIS, use of soduki, therapeutic rat-bite fever, 127

LYMPHATIC SYSTEM: See also Lymph Nodes

function increased in flush phase of pyrogenic (typhoid vaccine) admin., 416

radiation damages circulating lymphocytes and tissues, 752

reaction to B. monocytogenes infection, 265

LYMPHOCYTES: See also Leukemia;
Leukocytes
atypical, in arthritis after
typhoid vaccine therapy, 82
effect of dry heat on increase
of, 85

—— stimulation by dry
heat on natural and induced
resistance to transplanted
cancer, 86, 87

—— inj. of proteins,
carbohydrates and fat upon,
137

mechanism of lesions provoked
by epinephrine and
Salmonella typhi, 569

radiation damages circulating
lymphocytes, 752

rapid mobilization after infections
with B. monocytogenes, 265

LYMPHOGRANULOMA: See Hodgkin's
Disease

LYMPHOID TISSUE: See also Lymphatic
system activity, effect of dry
heat on, 85, 86, 87

LYSINS: See also Bacteriophage;
Hemolysis

mechanism of reaction of non-
specific proteins in therapy
110

LYSIN, b-, bactericidal substance from
serum of fever patients resemble
properties of, 256

LYSIS: See also Dialysis; Glycogen,
lysis; Hemolysis; Hydrolysis
phage-inactivating agent of
bacterial extracts, 211

LYSOZYME, hydrolysis did not follow
treatment with, 430

MAGNESIUM, chloride, polypnea
provoked by, at environmental
temperature, 528

MALAISE, after therapy of arthritis
with typhoid vaccine, 66
cardiac output during, after
inj. of typhoid vaccine,
156

MALARIA, chills, calcium chloride
in, 296

daily variation in leukocytes,
44

induced, relation of plasma
proteins to sedimentation
rate, 352

parasites, inj. in progressive
paralysis (1917), 12

renders victims resistant to
inj. of typhoid pyrogen, 585

therapeutic, in asthma, 637;
in paresis, 561; dangers,
605; in schizophrenia and
paralytic dementia, 240

injury to blood cells may
release pyrogens, explains
fever activity, 427

plasma globulin increase in
general paresis, 351

MALONIC ACID, removal from luteic
acid, luteose remains, 274

MANNOSE, d-, in fraction from
culture of tubercle bacilli,
216

MEDIASTINUM. sarcoma, therapy with
Coley's fluid, 136

MELANOMA: See Tumors, melanoma

MELANOSARCOMA: See Sarcoma

MENINGITIS, bacteferbrin for, 765

MENINGOCOCCI, endotoxin. effect on
carbohydrate metabolism in
rabbits, 536

endotoxin, effect of glycogen
synthesis, 534

effect on histamine of blood
and tissues of rabbits, 497

MENINGOCOCCI,

endotoxin

- inhibition of phosphorylation of glucose by, 535
- filtrates, activity in malignant neoplasms, 245
- degenerative changes of sarcoma by, 255
- in therapy of mouse sarcoma #180; skin reactivity, 183
- prodn. of Shwartzman phenomenon with sulfonamide conjugate of, 347
- purification, 261
- reaction of mice and tumors to injs., 235; of mouse sarcoma #180, 266
- in blood, experimental study of purpuric meningococcemia in relation to Shwartzman phenomenon, 475
- polysaccharide: See Polysaccharides

MENOPAUSE, Temperature variations during, 712, 761

MENTAL DISEASES: See also Dementia

PraecccX; Insanity: Paralysis, General; etc.

- diagnosis, Donaggio's 'inhibition phenomenon' in urine and spinal fluid of neuro-and psychopathic patients after induced fever, 195
- therapy, effects of Pyromen in subpyrexial doses, 708; of typhoid vaccine on leukocytes and adrenal cortex, 619

MESENTERY, blood supply, mode of action of bacterial endotoxins in animals, 521

METABOLISM: See also Bacteria,

metabolism; Carbohydrates, metabolism; Oxygen, consumption; Protein, metabolism; etc.

- activity of central nervous system on thermostatic behavior of body, 527, 529
- anti-neoplastic substance from penicillin, as metabolite, 428
- basal, in fever, 96
- blockade, mechanism in inhibitory effect of polysaccharides on mumps virus, 525

- in nonspecific protein therapy, 228; in neurosyphilis, 229
- increase in catabolism from induced fever in arthritics, 550
- pyrogens from enzymatic breakdown material, 47
- tissue, action of colchicine and E. typhosa ext., 249

METALS: See also Ions; Iron; Potassium; etc.

- colloidal, eosinophils increased after inj. of, 137
- in pyrogenic therapy (platinum, silver), 76
- effect of inj. upon leukocytes, 137

METHYL bis (b-CHLOROETHYL)-AMINE:

See Mustard, nitrogen

METHYLENE BLUE, therapy, oral stimulative, 103

MICE, cancer in mice (carcinoma F), 86. 196, 225, 294

- distribution of polysaccharide in tissues after inj. of paralyzing dose, 488
- influence of adrenal hormones on toxic and tumor-damaging effects of certain substances, 622
- inhibitory effect of bacterial suspensions (gram-negative) applied intranasally to mice, 677
- intraperitoneal inj. of bacterial pyrogens and cortisone, response of connective tissue mast cells to, 732
- nature of antigenic and antibody response of polysaccharide from S. marcescens, 616
- normal and tumor-bearing, effect of S. marcescens polysaccharide on rectal temperature, 439
- passive immunization elicited by single inj. of polysaccharide (S. marcescens), 566

MICE

pathologic changes, from
 protein-polysaccharide fraction
 from S. typhimurium, 297
 protection against lethal action
 of gonococcal endotoxin by
 penicillin, 451
 relation of virulence and toxic
 substances in mouse typhoid
 bacilli, 334
 sarcoma, nature of bacterial poly-
 saccharides which produce
 hemorrhage, 372
 properties of hemorrhage-prodg.
 filtrate of B. prodigiosus, 321
 sarcoma #180, action of bacterial
 toxins on, 373
 effect of inj. of products from
 gram-negative bacillus upon
 growth, 239; of S. marcescens
 polysaccharide, 233, 642; of
 S. paradysenteriae toxin, 386
 relation of mortality to
 hemorrhagic necrosis and re-
 gression elicited by bacterial
 filtrates, 245
 skin reactivity to bacterial
 (Meningococcus) filtrate in
 therapy, 183
 sarcoma # 37, 275
 damage by Ps.aeruginosa fraction,
 657
 effect of bacterial polysaccharide,
 486; and adrenal cortex ext. on,
 486, 522; of colchicine, 294; of
 E. coli filtrate, 294; of S.
 marcescens polysaccharide, 515
 522, 565
 immunization, passive and active,
 on lethality of polysaccharides,
 565
 pneumonia, virus, modification by
 polysaccharides, 495
 reducing substances as indicators
 of protection afforded by
 cortisone against tumor-damaging
 effects of S. marcescens
 polysaccharide, 697
 toxicity of antigenic fraction from
 B. typhimurium and B. typhosum, 301
 tumors, action of S. typhimurium
 toxin on, 385

assay method for det.
 potency of hemorrhage-
 producing bacterial preps.,
 380
 degenerative changes induced
 in cells by S. marcescens
 polysaccharide, 485
 effect of B. prodigiosus
 filtrate on, 330
 hemorrhage, 244
 induction of hemorrhage by
 bacterial toxins, action of
 sulfonamide on lethal factor,
 349
 reaction to inj. of filtrates
 from Meningococcus and B.
 coli, 235
 reduction in toxicity of S.
 marcescens polysaccharide
 by beef adrenal ext. 469
 spontaneous, effect of
 stimulation of lymphocytes
 on rate of growth, 87
 unanesthetized, effect of
 bacterial polysaccharide and
 tourniquet shock on peri-
 pheral circulation, 436

MICROSCOPY, critical analysis of
 134 proven cases of sar-
 coma, 60
 electron, graphs reveal size
 and mol. wt. of active
 principle of polysaccharide
 from S. marcescens, 374

MIDBRAIN: See under Brain

MIGRAINE, analysis of mechanism,
 313

MILK, B. coli grown in, 260
 human, 126
 intradermal injection, 112;
 intramuscular injection,
 reduced antitoxic action of
 blood in diphtheria, 268
 therapy, of eye inflammations,
 135, 775
 effectiveness, compared to
 Pyromen, 775
 pyrogens, not protein, reason
 for fever, 88

- MILK
therapy
significance of casein, bacterial endotoxin responsible for pyrogenic activity, 84
'thyrotoxicosis', biological split protein from, 57
- MILLON'S TEST, 321
- MINERALS: See also Ions; Iron; Metals; etc.
metabolism, in fevers, 466
- MISSISSIPPI, cotton industries,
'cotton fever' incidence, 402
- MOLDS: See Fungi
- MOLISCH REACTION, 243, 321, 381, 700
chemical basis of fever with inflammation, 427
polysaccharides of Sporotrichum schenckii, 503
positive for leukopenic factor of E. typhosa, 366
- MONKEYS, mechanism of aspirin antipyresis, 369
- MONOSACCHARIDES, oral admin. of, and amino acids produced fever in infants, 445
gastric secretion in dog diminished by 10% soln. of, 267
- MORPHINE, sulfate, suppress hyperpyrexia, 359
- MORPHOLOGY, free stem cell identical with lymphocytes found in lymphatic tissue, mesenteric node and spleen, 265
reactivity of mouse sarcoma 180 to bacterial filtrates, 266
- MUCINASE, 677
- MUMPS: See Parotitis
- MUSCLES, animal, extracts, during fever, 7
histamine, effect of meningococcal endotoxin, 497
innervation; See Neuromuscular System
pain in, reaction to pyriperazine therapy, 185
temperature, during bacterial chill, 219
- MUSTARD, nitrogen (HN_2), effect on renal manifestations of glomerulonephritis, 564
nitrogen, (methyl bis (b-chloroethyl)-amine)
suppression of leukocyte to response to bacterial pyrogen by, 735
- MYCOBACTERIA: See Bacteria, Corynebacterium; Bacteria, mycobacteria
- MYELOCYTES, in arthritis after typhoid vaccine therapy, 82
- MYELOID, production, mechanism of reaction of nonspecific protein in therapy, 110
- MYOSITIS, therapy, clinical response to mixed vaccine, 607
- NATIONAL CANCER INSTITUTE, 469
- NAUSEA, reaction to fever induced by killed B. coli, 188
- NECROPSY: See Autopsy
- 'NECROSIN', non-pyrogenic, proteolytic material, 401
'pyrexin' secured by enzyme action in, 427
- NECROSIS: See also Kidneys, necrosis; Liver, necrosis; Schwartzman Phenomenon; Tumors
action of mixed toxin (Coley's Fluid) on, 29

NECROSIS

- hemorrhagic skin lesions, 247, 281; after cutaneous inj. of antigen of E. typhosa, 152 376
- of sarcoma cells, 255
- vascular reaction of normal and neoplastic tissue to bacterial polysaccharides, 467

NEOPLASMS: See Cancer; Sarcoma; Tumors

NEOSAPROVITAN B (vaccine from saprophytes): See Saprophytes

NEPHRITIS: See also under Kidneys
glomerular, effects of febrile plasma, typhoid vaccine and nitrogen mustard on, 564;
renal irritation from foreign protein (typhoid) therapy, 382

NEPHROSIS: See Kidneys, disease; Nephritis

NERVES: See also Nervous System; Neuritis; paralysis
facial, severed and stump implanted in cat, 740
impulses, conduction in regenerated spinal cord of cat, 725
optic: See also Neuritis, optic atrophy, after typhoid vaccine therapy, 588
atrophy, hastened by pyripher, 163
function restoration by pyripher, 768
peripheral, blending of regenerating fibers with cerebral cortex through glial barrier inhibition, 704, 740, 756
regeneration, brain tissue ext. of V. Muralt said to activate therapy of multiple sclerosis, 506
neuron, in disease, (Pyromen), 771
splanchnic, effect of febrile agents on, 130
section, effect on pyrogenic changes, 74

stimulation by killed bacteria, initiates glycogenolysis, 132

vagus, effect on blood sugar of sectioning, 132

regulation of cholinesterase in blood by pyripher and ultraviolet rays in therapy of dermatoses, 510

section, effect of typhoid-paratyphoid vaccine in rats, 318

section, effect on pyrogenic changes, 74

NERVOUS SYSTEM: See also Brain; Nerves; Neuroglia; Neurons; Spinal Cord; etc.

central: See also Brain

induction of neuronal regeneration in, 688, 689, 736, 748, 757, 773, 778

metabolic activity of, on thermostatic behavior of body, 527, 529

regulation of blood, 269

site of action in, of bacterial pyrogen (Pyromen), 563

diseases, therapy with Saprovitin, 121

typhoid vaccine therapy in virus infections of, 461

sympathetic, effect of non-specific protein therapy on, 129, 138; factor in body's resistance, 112; influence of typhoid-paratyphoid vaccine on, 318; stimulation by typhoid-paratyphoid vaccine, 270
regulation of cholinesterase in blood by pyripher and ultraviolet rays in therapy of dermatoses, 510

surgery: See Sympathectomy

NEURASTHENIA, complications of paralysis, treated with pyripher, 163

NEURITIS, optic, appeared after typhoid vaccine therapy, 588
retrobulbar, therapy with pyripher, 768, 771

- NEURODERMATITIS, therapy with
Pyromen, 745
- NEUROGLIA, blending of regenerating
peripheral fevers with cere-
bral cortex through glia
barrier inhibition, 704
inhibition of glial barriers in
transected spinal cord by
Pyromen, 754, 755, 756, 757
- NEUROMUSCULAR SYSTEM: effects of bac-
terial polysaccharides, 589
- NEURONITIS, and neuronopathy, therapy
with typhoid vaccine, 542 ; with
Pyromen, 771
- NEURONS, regeneration, induction
(Pyromen) in central nervous
system, 688, 689, 736, 748,
757, 773, 778
_____ after ACTH,
757
- NEUORETINITIS, after typhoid vaccine
therapy, 588
- NEUROSYPHILIS, asymptomatic, cerebral
blood flow and oxygen consump-
tion detn. after intravenous
typhoid vaccine, 605
therapy, cardiac enlargement in in-
duced fever (typhoid vaccine)
therapy, 463
effects of induced fever on
cerebral functions in, 653;
foreign protein, 431; pyrifery,
163, 181; Pyromen, 593; typhoid
antigens, 345, 446
metabolic rates in therapeutic
fever, 229
use of nuclein compared to tuber-
culin, arsphenamine, 140
with suspensions of Ps. scissa and
ureae, 208
- NEUTROPHILS: See Leukocytes
- NEW AND NON-OFFICIAL REMEDIES, retain
mixed toxins of Coley in, 220
- NINHYDRIN REACTIONS, 321
- chemical basis of fever with
inflammation, 427
- NITROGEN, chemical studies in poly-
saccharides of anthrax
bacillus, 282
content of antigen of B.
dysenteriae, 324
in blood: See Blood, nitrogen
in extract of culture of E.
typhosa and B. paratyphosus B,
106
mustard: See Mustard, nitrogen
nonprotein, increase after toxic
doses of polysaccharide, 394
soluble, constituent of antigen
of B. typhosa, 319
- NOSE, intranasal application of bac-
terial suspensions in
pneumonia virus of mice, 677
secretions, culture after in-
halation of S. marcescens,
455
vaccines from organisms found
in, used in therapy of
asthma, 89
- NOVALDIN, intravenously for post-
transfusional reactions, 706
- NUCLEIC ACID, cause of fever after
inj. of distilled water, 138
component of polysaccharide or
type I Pneumococcus, 512
liberated after admin. of
colloidal toxic bacterial
substances, 70
from streptococcal nucleo-
proteins, 272
lipo-protein-nucleic acid com-
plex (reticulose) in therapy
of radiation injury, 752
phenobarbital admin. depresses
leukocyte genesis, 269
relationship to gelation, 388
therapy of progressive paralysis
with inj. of, 41
thymic, 430
effect on gastric motility in
dog, 355
yeast, deep depressive effects
on gastric motility in dogs,
355

NUCLEIN, therapy, in neurosyphilis,
140

NUCLEOHEXYL, pyretotherapy of acute
poliomyelitis, 664

NUCLEOPROTEIN, antigenic, isolation
from Ps. aeruginosa by pptn.
with ZnCl₂, dialysis and
acetone pptn., 587
component of endotoxin of S.
enteritidis, 617
crude, prepn. from B. coli, Ps.
pyocyaneus, and Streptococci,
586
fraction of bacteria is Shwartzman-
active, 223
streptococcal, 272
toxic, 412

NUCLEOTIDES, pentose, inhibited
gastric motility in dogs, 355
pentose, prodn. of gelatin in
media, leukopenia and
leukocytosis, 379

OMENTUM, induced development of
plasma cells, 257

OMNADIN, intramuscular inj. increased
antitoxic action of blood in
diphtheria, 268

OSTEO-ARTHRITIS, therapy, clinical
response to mixed vaccine, 607

OVALBUMIN: See Albumin

OVARECTOMY, effect on Pyromen res-
ponse, 754

OXYGEN, consumption, comparison of
glycolysis and, in brains of
normal and febrile rabbits, 606
-----effects of fever
induced by bacterial pyrogen
on, 605
in vitro of liver slices from
febrile rabbit, 581, 707

paralleled degree of malaise
and fever after typhoid
vaccine inj., 156
deficiency, (anoxia) may under-
lie liver cell necrosis,
479

P25; See Polysaccharides, from
S. marcescens

PAIN: See also Muscles, pain in
relief, after therapy of
arthritis with typhoid
vaccine, 66, 75
in peptic ulcers with sur-
gical and protein therapy,
113
in sarcoma treated with mixed
toxins (Coley), 60

PANCREAS, islands of Langerhans,
hydropic degeneration in
hyperglycemia, 111

PANTING: See Respiration

PARA-AMINOBENZOIC ACID, anti-
toxic effect, 349

PARA-AMINOHIPPURATE: See
Kidneys, physiology

PARALYSIS: See also
Poliomyelitis
diplopia and "blindness: in
reactions to transfusions
in multiple sclerosis, 513
facial, allergic etiology,
pyretotherapy, 595
general: See Paralysis, Gen-
eral

PARALYSIS, GENERAL: See also
NEUROSYPHILLIS
plasma protein during, 351
therapy, fever, action, 12;
danger, 609

PARALYSIS, GENERAL

therapy

malaria as, 561
 with nucleic acid, 41; penicillin
 and typhoid vaccine, 594; with
 pyripher, 163, 171, 217, 240, 260;
 with soduka, 127; with Staphylo-
 cocci and Streptococci, 54; with
Ps. scissae and ureae, 208; with
 tuberculin, 59

PARAPLEGIA, therapy, Pyromen, 771

PARATYPHOID, bacillus: See Salmon-
ella paratyphosus

in children, therapy with
 pyripher, 533

infections, (B), in relation to
 hyperglycemia, 111

vaccines, effect of hypothalamic
 lesions on fever induced by,
 289

injection, intraperitoneal into
 rats, 318; intravenous, in cat,
 270

typhoid, effect of temperature
 on reaction, 577, 579

heat prodn. of liver during
 fever by, 581

in exptl. fever, 231

metabolic activity of central
 nervous system as it affects
 thermostatic behavior, 529

(Pyrexin), nitrogenous, ther-
 mostable substance, non-toxic
 to mice, 401; 'necrosin' as
 source, 427

respiratory response to in-
 halation of carbon dioxide and
 572

thermal polypnea after, 528

use of TAB in gonorrhea, 391

PARENTERAL THERAPY: See Injections;
 under various diseases and
 therapeutics

PARKINSONISM, late evolution of
 choreic encephalitis toward,
 TAB and radiotherapy, 547

PAROTITIS, virus, effect of poly-
 saccharide on reaction between
 erythrocytes and, 526

virus, inhibition of multi-
 plication by polysaccharide
 of Friedländer's bacillus,
 492, 576

inhibitory effect of poly-
 saccharides on, 525

PATHOLOGY: See Fever, pathology;
 Tumors, pathology; under
 diseases

PELTATIN, alpha-, hemorrhagic

necrosis induced in mouse
 sarcoma #37 by, 697

toxicity, adrenocortical factors
 do not protect mice against,
 622

PELVIS, tumors, metastases of
 myosarcoma of uterus, Coley's
 toxins in, 210

PENICILLIN, Danish, pyrogenic
 effect of, 460

impure, hypothermia in mouse
 as bioassay of endotoxin
 protection factor, 552

luteum, metabolism, luteic acid
 product of, 274

protection against bacterial
 endotoxins, and its im-
 purities, 538

against lethal action of
 gonococcal endotoxin, 451

pyrogens in, removal by fil-
 tration with asbestos pads,
 434

thermostability of, and re-
 moval from, 432

therapy, of symptomatic paresis
 with typhoid vaccine, 594;
 of diptheria, 694

PENICILLINASE, 538

PENICILLIUM notatum, metabolite
 from culture fluids, evidence
 of anti-neoplastic substance
 as, 428

PENIS, inflammation, therapy with
 pyripher, 178

PENTNUCLEOTIDE: See Nucleotides,
 pentose

- PENTOSE, nucleotide: See Nucleotides
- PEPSIN, depressant of gastric motility may owe effect to pyrogens, 378
- PEPTIC ULCER, effect of urine exts. on, exptl. and clinical study, 329
therapy, protein and surgical, 113
pyrogens, 596, 600
- PEPTIDASE: See Enzymes
- PEPTIDE: See Polypeptide
- PEPTONE, affected neither temperature nor gastric motility in dog, 355
effect on hunger and appetite in fever in dog, 74
migration of leukocytes from inj., 112
- PERMANGANATE, titer, detection of pyrogens in parenteral sols., 327
- PERSPIRATION, after fever crisis, evidence of water liberation, 81
during pyrogenic therapy with colloidal metals, 76
induced by aspirin in fever produced by yeast, 369
- PETTERSON'S beta-LYSIN, 256
- pH: See Hydrogen Ion Concentration
- PHAGOCYTES: See also Immunity; Leukocytes; Reticulo-endothelial System
mononuclear, polyblasts of omentum, precursors of stimulated plasma cells, 257
- PHAGOCYTOSIS: See also Leukocytes; Leukocytosis
glucido-lipidic antigens and phagocytic ability of reticuloendothelial cells, 393
in chancroid therapy, 142
leukocyte chemotactism and bacterial toxins, 365
of bacterial capsule and polysaccharide, influence, 286
- PHARMACIES, bacterial content of distilled water stored in, 42
- PHENOBARBITAL: See Barbitol Derivatives
- PHENOL, in degradation of antigenic material from B. dysenteriae, 324
red: See Phenosulphonphthalein
- PHENOSULPHONPHTHALEIN (phenol red), clearance during pyrexial reaction to typhoid vaccine, 264
- PHOSPHATIDE: Blood, Lipoids; Phospholipin
- PHOSPHOLIPIN, component of antigenic complexes of Sh. paradysenteriae, 418; of endotoxin in S. dysenteriae, 306, 363; of O antigen of E. typhosa, 353
- PHOSPHORUS: See also Phospholipin in blood: See Blood, phosphorus in ext. of culture of E. typhosa, 106
pyrogenic endotoxin dependent on presence of phosphate group attached to polysaccharide, 758
- PIRETAMINA: See Fever, induced
- PIROMEN (Pyromen)
acne, therapy of, 708, 777
adrenals, role in leukocyte changes after, 620, 720; removal, 644
allergy, food, therapy of, 738
antigenicity, lack of, 777
Arthus phenomenon, compared with cortisone administ., 731
asthma, perennial, therapy of, 637, 671, 708, 737, 779
bacteria, in blood stream, removal by, 762
B. subtilis, 702, 709, 732

PIROMEN(Pyromen)

blood, bacterial invasion, effect on removal, 762; plasma, effect of incubation with, on augmentation of Piromen, 716
 blood pressure, high, therapy of, 603
 brain, glial barrier inhibition by, 704, 740, 756, 757; regeneration, neuronal, induced by, 741, 757
 burns, healing of, influenced by, 704, 710, 754, 773
 cells, endocrinal, 688, mast, 678, 679, 688, 732; See also: Hematopoietic effect, 692, 691; Leukocytosis
 chemistry, 665, 709, 769
 chlor-trimeton, 715
 clinical studies, 604, 660, 661, 671, 708, 715, 722, 724, 737, 738, 745, 747, 771, 777, 779
 dermatology, therapy in, 660, 708, 724, 745, 777
 diluent, preferred, 722
 dosages, lethal, for mice, 754
E. coli as source of, 709
 endocrine system, effect on cells of, 688, 689, 693, 754
 eosinophils, 759
 gastric secretion depression after, 625, 685
 headache, therapy of, 626
 hematopoietic effect, 691, 692
 hypertension, therapy of, 603, 722; essential, 661, 722
 hypothermia, inhibition, 643
 lactation, variation after, 754
 leukocytosis, See : Leukocytosis
 liver, hematopoietic effect on, 692
 nerve, regeneration, in central nervous system, 563, 593, 688, 690, 703, 704, 725, 736, 741, 748, 754, 755, 756, 757, 771, 773, 778, 740, 756
 nitrogen mustard therapy, combined with, 735
 oral administration, 631
 ovariectomy, effect of, 754
 pituitary-adrenal system, stimulation by, 661, 687, 774, 778, 779
 pregnancy, effect of, 754
Proteus vulgaris, as source of, 670, 693, 702, 709, 715, 728, 732

Pseudomonas aeruginosa, as source of, 481, 563, 625, 643, 665, 670, 671, 689, 690, 691, 693, 703, 704, 709, 728, 732, 735, 736, 744, 749, 754, 762
 paraplegia, therapy for, 771
 renal system, hemodynamic changes after, 718
 response diminished by barbiturates, tolserol, 722
 retina, hemorrhage regression in, 603
 scar tissue prevention, 689, 690, 703, 710, 736, 740, 754, 755, 756, 757
 spinal cord. See: Nerve regeneration, central nervous system
 spleen, hematopoietic effect on, 691, 692
 stress factor in therapy by, 620, 774, 775
S. typhi as source of, 709
 thyroidectomy, effect of, 754
 therapy of acne, 708, 777
 allergy, 671, 708, 737, 777;
 anaphylactoid, 777; food, 738;
 perennial, 671, 737
 anaphylaxis, 777
 asthma, 637, 671, 705, 708, 737, 779
 burns, 704, 710, 754, 773
 cardiac decompensation, 604; failure, 603
 dermatoses, 708, 777; chronic, 724; neuro-, 745
 headache, 626
 hypertension, 603, 604, 722
 neuronitis, 771
 neurosyphilis, 593
 ophthalmologic conditions, 774, 775
 poliomyelitis, 747, 765
 retinal hemorrhage, 603
 rhinitis, 603
 sclerosis, multiple, 771
 tissues, effect on connective, 678, 679, 688, 732
 —————endocrine, 687, 688, 689, 693, 754
 —————nervous.
 See: Nerve, regeneration
 —————spleen, 691, 692.

PITUITARY BODY: See also Hypothalamus
changes induced by bacterial pyrogen, 687

-----pathologic, 779

excision, temperature and thermal
reactions in dogs, 184

physiology, ACTH-like effect of
fever in man, 661, 715

and pathology of temperature-
regulatory function, 307

stimulation of adrenal-pituitary
system in Pyromen therapy of
allergy, 671

PITUITARY PREPARATIONS, abolition
of pyrogenic reaction with
ACTH, 659, 705, 742; in
refractory condition after
protein therapy, 620

ACTH, for therapy of bronchial
asthma, 779; headache, 626
comparison of leukocyte res-
ponse with that of Pyromen, 727
ACTH-like effect of fever in man,
661, 715

PLACENTA, decidua-, hemorrhage in
mice induced by endotoxins of
gram-negative bacteria, 384
hemorrhagic action of endotoxin
from colon-typhoid dysentery
group followed by abortion, 667

PLANTS, pathogens, tumor hemorrhage
production, 433

PLASMA: See also Blood
barriers, influence of pyripher
fevers on, 262
cells, induced development and
histogenesis of, 257
plasmacytoma, morphological
changes by pyripher, 514
coagulation, effect of Shear's
polysaccharide, 721
from febrile patients, effect on
glomerulonephritis, 564
homologous, augmentation of
leukocytic and pyrogenic effects
of typhoid vaccine, 523, 636
-----of
thermogenic effects of pyrogens,
591, 716
pyrogenicity of contaminants of, 42

transfusion: See Blood Trans-
fusion

PNEUMOCOCCI, antiserum, nature of
chill-prod. principle, 182
cultures, specific soluble sub-
stance elaborated by, 69
infections, diagnosis by
precipitin reaction of
specific substances in blood
and urine, 69

plasma from, effect of inj. in
glomerulonephritis, 564

polysaccharide: See Poly-
saccharides

vaccines, in therapy and pre-
vention of influenza and
sequels, 98

PNEUMONIA, high values for sugar
associated with protein in
serum, 180

lobar, cutaneous and serological
reactions, polysaccharide of
Pneumococcus, 248

virus, bacterial and non-bacterial
substances modified course
in mice, 495

effect of polysaccharides on
reaction between erythro-
cytes and, in mice, 526

inhibitory effect of bacterial
suspensions in mice, 677

POLIOMYELITIS, acute anterior,
therapy with pyrogenic
substances, 641, 765; trial
therapy with vaccine (anti-
rabies) and serum (maternal
blood), 555

pyretotherapy (vaccine, pyripher
and nucleohexyl), 664

therapy, aspecific intradermal
vaccination, history,
technics and mechanism of
action, 629

(Pyromen), 747, 771

with pyripher, 263

POLYPEPTIDE, component of antigen
of B. dysenteriae, 306, 324
component of antigen of B.
typhosa, 319: of antigen of
B. shigae produced low titer
homologous precipitins, 310

POLYPEPTIDE

- thermolabile leukocytosis factor may be, 663
- toxic non-antigenic component of antigenic endotoxin of S. dysenteriae, 363

POLYPNEA: See Respiration

POLYSACCHARIDES: See also Acacia;

- Carbohydrates; Glycogen;
- Hapten; Inulin; Luteose; Starch; etc.
- antigenic; See Antigens
- bacterial, activities, (Symposium on bacterial pyrogens), 754
- anticoagulant effect in normal and thrombocytopenic plasma of leukemia, 743
- effects upon cardiovascular and neuromuscular systems, 589
- nerve tissue (Pyromen) 688, 689, 736, 748, 757, 773, 778
- (including dextran) adsorbed on colloids as antigens, 194
- influence of phagocytosis of, 286
- Sahayun method and Hagedorn-Jensen test not satisfactory for estimation of, 568
- effects, difference in pyrogenic composition may account for variations in, 709
- on reaction between erythrocytes and viruses (mumps), 526
- on virus pneumonia after intranasal inj. 495
- fractional precipitation by alcohol or glacial acetic acid, 309
- from B. aerobacter, detoxification mechanism, 764
- effect upon bacteriophagy, 295, 621; in polar bodies, 763; sarcoma tissue cultures, 764; role of B. lactis aerogenes in bacterial anaphylaxis, 153
- from B. anthracis, chemical studies, 282; serological study, 187
- from B. dysenteriae (Shiga), prepn. and properties, 242
- from B. enteritidis, 131

- from B. Friedländer, anti-coagulant effect in normal and thrombocytopenic plasma of leukemia, 743
- effect on reaction between erythrocytes and viruses, 526
- enzymatic hydrolysis of capsular polysaccharide of Klebsiella pneumoniae, of polar bodies, 763; loss of serologic activity, type B., 554
- fate studied by homologous antibody labelled with fluorescein, 654
- inhibition of bacteriophagy by, 621
- inhibitory effect on mumps virus, 492, 525
- isolation and chemical analysis, 109; of types A and C, 133
- (Klebsiella pneumonia), role in bacterial anaphylaxis, 153
- multiplication of variant of mumps virus in presence of, 576
- type Bin mouse, 654
- from B. megatherium, existence of, 553
- from B. mesentericus, serological study, 187
- from B. subtilis, chemical study, 773; physiological study, 773; serological study, 187
- from E. coli, 128
- chemical study, 773
- factors influencing content of, 568
- method for sepn. and purification, 128
- physiological study, 773
- Ps. pyocyaneus and Streptococci, prepn. of crude nucleoprotein and fractions of, bacterial allergy, 586
- from H. influenzae, types A, B, C, D, and F, chemical, physical and immunological properties, 450
- from Meningococcus, serological study, 187

POLYSACCHARIDES

- from Pneumococcus
 - C, cutaneous and serologic reactions, 248; effect of calcium on flocculation reaction with, 315; precipitation of blood protein with, 322
 - effect on capillary circulation, 362
 - separated from proteins by extg. with phenol; 309
 - type I, fractionation and physical-chemical studies, 512; type I (and V.), improved method for prepn., 649
 - type II and III, localization of antigen in tissue cells, 658
 - type IV, N-containing, specific and nonspecific, 177
- from Proteus vulgaris, 67, 693, 702, 709, 715, 728, 732, 754, 773
 - ACTH-like effect of fever in man (Pyromen), 715
 - chemical study, 773
 - physiological action, 773
 - serological study, 187
- from Pseudomonas species, 481, 563
 - 625, 643, 665, 670, 671, 690, 691, 693, 702, 703, 704, 709, 728, 732, 735, 736, 744, 749, 754, 762, 773
- from Salmonella species, erythrocyte sedimentation rate increased by, 611
 - immunological relationships, 287
 - lipid-protein complexes from, 611, 773
 - polar bodies of, 763
 - toxins, from S. enteritidis, 617, 311; from S. typhimurium, prodn. of pathologic changes, 297
- from Serratia marcescens, action of toxic doses on dog and guinea pig, 394
 - anaphylaxis not produced by, 395
 - anoxia in tumor tissue, prodn. by, 764
 - anticoagulant effect on normal and thrombocytopenic plasma, 721, 743
 - changes in adrenals of tumor-bearing mice after, 522; degenerative in tumor cells, 485
 - chemical alteration of, 613, 773; by iodopolysaccharide in tumors, 544; studied by additions of radioactive I, 545
 - chemotherapy of neoplastic diseases, 672; See also: Pyrogens, Sarcoma, Tumors
 - comparisons of immunological, toxic and tumor-necrotizing properties, 484
 - cytological effects on tumors, 443, 444, 498, 642, 650, 764
 - effect of detoxifying agents, on toxicity, 642; of adrenal cortex extracts combined with, 486; of immunization on lethality, 565; of storage and freezing on properties, 519; of sublethal dosages, 396
 - effect on blood pressure, in tumor tissue, 372, 436, 515, 770; systolic, 515; capillary circulation, 436; on plasma clotting, 721; rectal temperature, 439; sarcoma, mouse, 372, 498, 642, 650, 764,
 - fractions, antigenic and immunologic, 616
 - haptens, O antigen, site, 763
 - histologic appearance of tumor tissue after, 456
 - immunologic properties, 484, 519, 616; relationships, between filtrates and Shear polysaccharide, 405; of anti-body-containing fractions, 520; passive immunization against toxic activity, 566
 - iodinated, effects on carcinoma, 508
 - nitrogen content, effect on tumors, 330
 - pigment elaboration a factor in activity, 562
 - polar bodies as source of, 763
 - preparation, 504
 - pyrogen, active, from, 674; use as pyrogen, 438
 - reactions to injections of tumor-necrotizing preparations, mice, 406;

POLYSACCHARIDES

from S. marcescens

reactions to injections, by patients, 494, 502; stress, assoc. with tumor-destruction, 750

reducing substances as indicators of protection against tumor damage by, 697

removal, splanchnic, of bacteria, not altered by, 762

toxicity of, 345, 484, 569, 622, 642, 764; reduced by beef adrenal extract, 469; by cortisone, 697; by adrenal hormones, 622; by antihistamines, vitamin B12 and rutin, 642; not altered by storage, 519; enhanced by freezing, 519

vascular responses after, 437, 467, 622, 695, 764, 770

-(Shear); 275, 312, 321, 330, 371, 372, 374, 380, 381, 387, 390, 394, 395, 396, 405, 406, 436, 437, 438, 439, 440, 443, 456, 464, 467, 469, 484, 485, 486, 494, 498, 502, 504, 515, 519, 520, 522, 544, 545, 565, 566, 613, 616, 622, 650, 662, 695, 697, 721, 734, 743, 750, 762, 770

from Shigella dysenteriae, component of antigen, 306, 324, 363, 763

from Sporotrichum schenckii, assay, 503

from Staphylococcus, autolysate of, produced leukocyte response, 342

from starch glycogen and gum arabic, effect upon bacteriophagy, 295

from Streptococcus, biosynthesis of capsular, 753

inhibitory effect on mumps virus, 525

from tubercle bacillus. chemical analysis, 216

from tuberculin, 749

from typhoid bacilli, new substances related to, has potentialities in therapy, 336
method of sepn. and purification of, 128

from typhoid-salmonella groups, protein-free, immunological relationships of, 287

in serum: See Serum, Polysaccharide

nonbacterial, injected into mice, effects on cause of infection with pneumonia virus, 495

phage-inactivating agent of bacterial exts., can be converted into hapten by alkali, 215

solution (1 to 2%) stimulate secretion of gastric mucosa, 267; (10%) diminished gastric secretion in dog, 267

synthesis by bacteria and enzymes, 162; by microorganisms, 274

POTASSIUM, hydroxide, threads appear during growth of colon bacteria in, 388

permanganate, in test for pyrogen in distilled water, 160

PRECIPITATION, fractional, of bacterial filtrates, 261

PRECIPITIN, reaction, 141

reaction in diagnosis of pneumococcus infection, 69

PREGNANCY, blood, febrile reaction after first infusion of, new treatment of rheumatoid arthritis, 468

in mice, temperature factors in action of bacterial endotoxins, 413

in rabbits, effect of endotoxin S. paradysenteriae, 411

PROCAINE, hydrochloride soln., adsorption on activated charcoal, 442

PROPYL, iso-, ALCOHOL, fractionation of polysaccharide of type I pneumococcus, 512

PROSTIGMINE, prodn. of continuous gastric motility in dog, 355

PROTEASE: See Enzymes

PROTEINS: See also Albumin; Amino

Acids; Nucleoprotein; Polypeptide; Xanthoprotein; etc. and derivatives, chemical nature of immune substances, 134

bacterial: See also under names of bacteria, as *Bacteria*, coli action on arthritis of childhood, question of herpes, 166 certain phenomena associated with, 83

dialyzable product source of pyrogenic activity, 58 effects on eviscerated dogs, 130 extracts used in therapy of uveitis, 158

glutathione and ergothionine in blood increased during pyrexia, 207

in therapy of arthritis, 94; of dermatology, 95

influence of blood (cells, chemistry, volume, rate), 157

sensitivity in therapy of bronchial asthma with vaccines, 89

colorimetric tests for tyrosine, arginine and histidine in polysaccharide fraction of *B. prodigiosus* gave positive results, 372

component of O antigenic complex of *E. typhosum*, 353; of strain of typhoid bacilli, 334

decomposing substance, toxic principle found in, 4

effect upon leukocytes, 137

fevers, (milk, casein, albumin, edestin, hemoglobin), 107

pharmacology of, 175

theory of pathogenesis, 67

usual and unusual reactions to, 190

toxic or dynamic, 126

foreign; See Protein, therapy;

Toxins; names of bacteria;

Typhoid, vaccine; etc.

fraction of *B. dysenteriae* produced hyperglycemia and leukopenia, 258

heterologous, sensitizes certain cells of temperature-regulating centers, 77

hydrolysates, casein, peptides produce fever, 769

-----removal of bacterial pyrogens from 434

in blood: See Blood, proteins

in specific antigenic complexes of *Sh. paradysenteriae*, 418

injections, thrombopenia of, 118 metabolism, 124

influence of toxic substances on, 125

non-diazable, polysaccharide fraction from *S. typhimurium*, pathologic changes in rats and mice, 297

nonspecific, (chemical and physiological mechanisms), 228

bacterial, tuberculin reaction through, 15

poisonous, theory of pathogenesis of fever, 67

polysaccharide-lipid-protein complexes from *Salmonella* group, 611

reactions, lymph and serum ferments during shock, 68

requirements, minimal, of man in health and disease, 124

retains water in body, alimentary fevers, 174

role in specificity of reaction to living or killed bacteria, 209

solutions, conc. (serum, plasma, hemoglobin) reducing pyrogenicity, 509

split products, absorbable in digestive tract or intestines, 126

basic cause of fever theory, 174

in relation to immunity and disease, 57; pyrogen test, 767

in therapy of eye inflammation, 135

nonspecific protein therapy develops resistance to toxic or, 101, 102

PROTEIN

split products

'thyrotoxin' from ice cream
and milk, 57

therapy, effects of foreign
protein, 73, 100

in acute infections, 100; in
infectious diseases (typhoid),
mode of action, 71

intravenous foreign protein in
neurosyphilis, 431

lipo-protein-nucleic acid complex
(reticulose) in therapy of
radiation injury, 752

nonspecific, actions and appli-
cations, 138

———— avoidance and
palliation, 115

———— effect on autonomic
nervous system, 129

———— history reviewed, 226

———— in ocular disease,
135, 328

———— in rheumatism, 191

———— resistance, 101, 102

———— of chancroid, 142

———— of stomach and

duodenal ulcers, 113

———— reactions, 101, 110

PROTEOSE: See also Blood,
temperature regulation after intra-
venous inj., 104

PROTOPLASM, bacterial, partial toxin
in, 50

PSEUDOGLOBULIN, in blood, See Blood,
globulin

PSEUDOMONAS aeruginosa: See Bacteria,
Pseudomonas aeruginosa;
Pyocyanase

scissa and ureae: See Bacteria,
Pseudomonas

PSEUDOTUBERCULOSIS, bacilli: See
Bacteria, Corynebacterium

PSORIASIS, therapy with pyrif,er,
171, 178

therapy with typhoid vaccine,
95

PURINE, metabolism, influence of
toxic substances on, 125
toxicity, associated with
specific antigen of type Z
Sh. paradysenteriae, 418

PURPURA

meningococcic, experimental
study in relation to Shwartz-
man phenomenon, 475

PURILENCE: See: Pus, Staphylococci;
adrenal cortex extract as pro-
tection against, 189

PUS, sterile, leukocytosis induced
in hypertensive dogs by, 407

PUTREFACTION, a product of tissue
destruction produces fever, 1

PYOCYANASE, bactericidal soln.
from, 159

'PYRESE', and 'antipyrese', research,
16

'PYRETAMINE' of Russo, elimination
of histidine and total nit-
rogen during fever, 419

'PYREXIN" See Paratyphoid, vaccine;
typhoid, vaccine

PYRIFER: See also: Fever, thera-
peutic
antigenic characteristics, 680
antitoxic quality of blood
reduced by, 268
dosage, 260
effects of, focal pain produced,
useful as diagnostic aid, 766;
glutathione and ergothionine
increased, 207; globulin-
albumin alterations in serum,
308; morphologic alterations
in plasma cells, 514
effects on plasma barriers, 262;
neurologic syndromes, 767;
leukocytes and temperature
variations, 712
preparation, character, 260;
produced by Rosenberg,
Freiburg, 163

PYRIFER

therapy compared to physically-induced fever, 676; effect on allergy, 675; in balanitis, 178; in buboes, 178; colitis, 656; in dementia paralytica, 217, 240; diphtheria, 694; in dermatoses, 510; in enteric fevers, 574; in epilepsy, 217; in gonorrhea, 161, 171, 178; in neuritis, retrobulbar, 768; in neuro-syphilis, 163, 171, 181, 217; in poliomyelitis, 263, 664; in psoriasis, 171, 178; psychoses, 217; schizophrenia, 217, 227, 240; sclerosis, multiple, 712, 761; staphylococcic infections, 178; tabes, 167; typhus and paratyphus, 533; vulvovaginitis, 171

phenobarbital before administration prevents leukocytosis, 269

PYROGENS, 423, 723, 729, See also:

Piromen, Polysaccharides and specific bacteria

amino acid synthesis in media, effect on prodn., 218

antipyrogenic principle, in bacterial cultures, 237

bacterial, 403; assay for minimal dosage, 773; French bio-assay 635; for potency of hemorrhage-producing prepn., 380

biology of, symposium, 758

blood, lipids, changes after, 358

blood, for transfusion, report, 435; reactions from unpooled plasma, 360; incidence of, in multiple sclerosis, 513

chemistry of, 298, 379, 474, 511, 581, 674, 700

connective tissue mast cell response to, See: Piromen

contaminants in biologic products, 429, 499; in apparatus, 314

effects (of E. coli, E. typhosa, Proteus, B. pyocyaneus and B. abortus equi) on rabbit, mouse, horse, man, 511; of urethane on temperature, 578; changes in blood lipids, 358

E. typhosa, purified, effect of intravenous inj., 446; minimum, lethal and toxic dosages, 392; combination with B. tularensis increases tolerance to the latter alone, 746

gastric depression by, 355, 600 gelation, of media, 379

isolation. See Pyrogens, preparations; Bacteria (under specific names) from coliform of Salmonella organisms, 751

panting, thermal, inhibited by, 670

preparations from various species, 709; of safe infusion fluids, 246; method and test, 383; simple method for detecting, 648

properties and biologic effects-literature survey, 1855-; 623

Pseudomonas aeruginosa, preparations and test, 377, 383.

See Piromen, Polysaccharide reducing sugars (in Ps. aeruginosa, E. typhosa, and B. subtilis), 430

removal from protein hydrolysates, 434; from penicillin, 432 review, 284

symposium, 749, 754

thermostability, 432

tolerance to toxic activity, 470, 471, 540, 702, 733; abolition by reticuloendothelial blockade (typhoid pyrogen), 441

contaminants, cause of danger in saline fluids, 39; in medicinal solution, 49; prevention, by chlorobenzene, dichloro-prepn. of ethylene oxide, 758; in penicillin, 460

dose, minimal, for rabbits, 403; for man, 773

effects, ACTH-like effect, 661, 715, 742, 774, 775; augmentation of activity by plasma, 523, 591, 716; by whole blood, 744; of pyrexin, in inflammation, 597;

PYROGENS

- effects. on thermoregulatory centers, 577; of removal of adrenals and thyroid, 644, 645; of temperature on reactions to, 577, 579; inhibitory, 643; on hunger and appetite (dog), 74; on the lymphatic system, 773; on nitrogen and purine metabolism, 125; on gastric secretion, 580, 596
- elimination, 164, 583, 592; by adsorptive filtration, 238, 250; by hydrogen peroxide, 421, 548, 719; from solutions of streptomycin salts, 655; emergency, from water, 304; detailed methods for, 234; from serum, plasma, hemoglobin solutions, 509; from hormones, viruses, toxin-antitoxin preparations, 570; in solution manufacture, 558; review of methods, 505, 507, 541
- endotoxin, presence depends on phosphate attachment to hexose or protein component, 758
- exudates, thermostable leukocytosis factor, 663; euglobulin fraction in inflammation, 597
- humic acid (decomposed hay) as, 202
- hypothalamus, control over reaction, 289, 307; section, blocks pyrogenic activity, 403
- hypertension, reduction of blood pressure after, 338, 397
- inulin, dahlia, adsorption, 252; renal hyperemia not neurally controlled, 348
- literature, review, 284, 575, 705
- milk, source of, 84, 88. See also: Pyrifer
- nucleus, components, source, 379
- production, 278, 379; in gum acacia, 278; in protein solution, 107
- Proteus vulgaris, as source. See: Piromen, Polysaccharides and under name of bacterium
- Pseudomonas aeruginosa, as source. See: Piromen, Polysaccharides and under name of bacterium
- Pyrifer, as source. See Pyrifer reactions after administration, cause and elimination, 205; after deferenectionomy, 739; diminished by barbiturates, 722; at injection sites, 726; after transfusion. See Transfusions, blood, under Pyrogens; prevention, 205, 652
- renal changes, 348, 397, 682
- Serratia marcescens as source of, 438. See under; Polysaccharides and under names of bacterium
- solutions, as source, 107, 114, 164, 205, 206, 213, 234, 238, 246, 276, 370, 404, 482, 493, 501, 571, 599, 610, 624, 633, 635, 652, 700, 711
- decline in, during storage, 482, 493, 501, 571, 634
- depyrogenation, U.S.P., 548; preparation, 341, 442, 457, 459
- term, coined to cover all fever-producing agents, 9
- tests, 474, 592, 628
- animals used should not be restrained, 643
- based on leukocytosis of rabbit, for injectable solns., 633
- chemical, for distilled water, 160
- detection in fluids by biological methods, 389
- evaluation of biological and chemical tests for, 379
- for missed contaminations in biological products, role of cryophile bacteria, 487
- in aqueous solns. methods, 404
- investigations on tests for injectable solns., 699
- methods, 474
- methods for large scale testing in rabbit (thermocouple box), 448
- on non-pyrogenic fluids, 458
- quantitative assay method in rabbits, 602; def. of indices, 701
- search for reliable sensitive method for determination of pathogenicity in rabbits, leukocyte response, 417

PYROGENS

tests
 technic of U. S. Pharmacopeia
 test for injectable solns.,
 571
 tentative, in infusion fluid,
 339
 use of rabbits for detn.
 in solns. for intravenous
 use, 337
 with Polish rabbits, 490
 therapy, of allergy, 671, 737,
 738
 asthma (Pyromen), 637
 healing of burns in cats with
 (Pyromen), 704, 710, 754, 773
 injection in infectious
 processes, 102
 literature review, (editorial),
 705
 new fever therapy (Pyromen)
 of neurosyphilis, 593
 of arterial hypertension with
 (Pyromen), 603
 chronic dermatoses (Pyromen),
 724
 early hebephrenia, 221
 essential and malignant hyper-
 tension (Pyromen), 604, 722
 lower nephron nephrosis, 612
 neoplasms, 590
 neurodermatitis (Pyromen), 745
 poliomyelitis, acute anterior
 (Pyromen), 641, 747, 765
 uveitis, 774
 toxic, released from cell after
 injury by fever, chemical basis
 of fever, 401
 transfusion, blood, as source of
 reactions, clinical study, 706;
 comparison of homologous and
 conditioned O blood, 425;
 control of citrated solutions,
 204, 630; prophylaxis, intra-
 venous Ca., 631
 typhoid, anaphylatoxin effects, 51
 immunity in experimental fever,
 336
 influence of various diseases
 upon febrile response, 585
 typhoid vaccine, changes in cir-
 culation and respiration
 during chill and fever in man,
 416,

Augmentation of effects by
 homologous plasma, 523, 636,
 effects of fever on cerebral
 circulation and oxygen con-
 sumption, 605
 or imulin, only known method
 of inducing renal hyper-
 emia, 291
 --paratyphoid, and res-
 piratory response to inhala-
 tion of carbon dioxide, 572
 -----effect of temp-
 erature on response, 577, 579,
 -----heat prodn. of
 liver during fever by, 581
 -----in oxygen con-
 sumption in vitro of liver
 slices from febrile rabbits,
 707
 urine, content, separated from
 urogastrone, 303
 virus, influenza, pyrogenicity
 for rabbits, 557, tolerance
 to, 556
 water, 108, 160, 164, 276, 304,
 370, 442,
 yeast, as source of, 165

PYROMEN: See under Piromen

"PYROTOXIN:, name applied to
 agent isolated from B.
prodigiosus, 53

"PYROTOXINA bacterica", fever
 producing substance from
 bacterial cells, 18, 21,

RABBITS, action of typhoid
 bacillus extract on arterial
 pressure, 476; on tissue
 metabolism, 249
 antibody formation after
 typhoid vaccine, 310
 519, 601
 Arthus phenomenon in, action of
 cortisone and Pyromen on,
 731; skin lesions, 247, 518
 augmentation of pyrogenic acti-
 vity of typhoid vaccine by
 homologous plasma, 523
 591, 716

RABBITS

body fluid redistribution in thermal alterations, 673
changes, in blood, 116, 290, 325;
after B. aertrycke inj., 170, 214;
in cardiovascular response, 589,
662, in endocrines after pyrogens,
693; in carbohydrate metabolism,
116, 536; in oxygen consumption,
581, 707; in respirations and
glycolysis (brain) during fever, 606
dosage, minim. pyrogen, 773; lethal
S. marcescens, 440
effect, glutathione on temp. of, 207
hematopoeisis after bacterial pyro-
gens (Pyromen), 692
hyperthermia after bacterial fractions
and filtrates, 65, 152, 165, 169,
202, 271, 305, 440, 557; exudates
inflammatory, and necrosin, 427;
proteins, 107, 465; vaccines, 51,
273, 577, 579
hypothermia, emotional in, 643
necrosis, of kidneys, after bacterial
toxins, 734; effect of cortisone,
734
organs, pathologic changes after E.
typhosa antigens, 376
penicillin as pyrogen for, 460
polypnea in, 528
protective effect of S. dysenteriae
fractions, 560
sarcoma, 245
Shwartzman reaction, after E. coli
filtrate inj., 236
splenectomy in, effect on leuko-
cytosis, 80
temperature, normal range in, 501;
reading, 635
testing, for pyrogens, 602, 633, 635,
773; Polish rabbit for, 490
tolerance, for pyrogens, 339, 540,
556, 592, 733

RABIES, anti-rabies vaccine, trial of
acute anterior poliomyelitis, 555
therapy, bactifebrin not effective for,
765

RADIATION: See also Ultraviolet
Rays; X-Rays

tumor-necrotizing polysaccharide
S. marcescens tagged with radio-
active iodine, 545; poly-
saccharide with radioactive
iodobenzene, 613

RADIOTHERAPY, of choreic ence-
phalitis with late evolution
toward parkinsonism, 547
of Hodgkin's disease and lympho-
sarcoma, 149
of neoplasms, 590

RAT-BITE FEVER, therapeutic, soduku
general paralysis, 127

RATS, action of colchicine and E.
typhosa ext. on tissues, 249
adrenal changes after E. coli, 608
leukocytosis after pyrogens, 631
pathologic changes after S. typhi-
murium polysaccharide, 297
sarcoma and other neoplasms, after
adminis. of S. typhimur. fract.
293; after polysaccharide,
(Shear), 750; after bacterial
filtrates, 280
thermoregulation during B. pyocya-
neus induced fever, 199

RECTUM, temperature, biol. methods
for det. pyrogens, 389; effect
of S. marcescens polysaccharide
on normal rabbit, 440; fall,
during sarcoma, in mouse, 515;
not changed in dog after
pyrogens, 378

REDUCING SUBSTANCES (ascorbic acid
and glutathione) as indicators
of protection (by cortisone)
against tumor-damaging
effect of S. marcescens poly-
saccharide, 697
chemical studies of polysaccharide
of anthrax bacillus, 282

RESINS, ion exchange, to reduce
the pyrogens in parenteral
solutions, 370

RESISTANCE: See also Immunity
involuntary nervous system
important factor in, 112

RESPIRATION: See also Brain,
physiology
changes during chill and fever
(typhoid vaccine), 416
hyperpnea after typhoid vaccine, 72
inhibition of panting by bac-
terial pyrogen, in dog, 670
polypnea, effect of substances
affecting body temperature, 528
summary of mechanisms involved in
fever, 415
typhoid-paratyphoid vaccine and res-
ponse to inhalation of carbon
dioxide, 572

RESPIRATORY TRACT, disease, bacterial
allergy an aftermath of, 639;
bacterial allergy, prepn. of crude
nucleoprotein and polysaccharide
fractions of bacteria in, 586

RETICULOENDOTHELIAL SYSTEM: See also
Phagocytes: Phagocytosis: Spleen
blockade, abolition of tolerance to
typhoid pyrogen, 441; of pyro-
genic reactions, 705;
effect on immunity to Shwartzman
phenomenon, 472
cells reacting to toxic antigens
and to infection, 480
increased activity in, may underlie
mechanism of tolerance to endo-
toxins, 666
phagocytic ability and glucido-
lipidic antigens, 393
role in detoxification in dementia
praecox, 760; in tolerance to
bacterial pyrogens, 471
stimulation, route of inj. of sub-
stances important, 151

RETICULOSE: See Radiation, toxicity

RETINA, hemorrhage, reduced by bac-
terial pyrogen in malignant
hypertension, 549; regression
after Pyromen, 603

RHEUMATIC FEVER, acute, altered res-
ponse to typhoid vaccine

during massive salicylate
therapy, 496
therapy, effect of fever shock,
675

RHEUMATISM: See also Arthritis;
Osteo-arthritis
therapy, bilateral chorea res-
ponse to TAB, 547
inflammatory, typhoid vaccine
as adjuvant to antibiotics, 489
clinical response of fibrositis
tomixed vaccine, 607; non-
specific protein, 191

RHINITIS, therapy by Pyromen, 777

RHODOSPIRILLUM rubrum: See Bacteria,
spirillum

RICKETTSIA, bacilli, endotoxic
factors, immunological
relation to other gram-
negative bacilli endotoxins,
453

RIMBAUD, antichancroid vaccine, 254

RUSSELL BODIES, end point in
induced development and histo-
genesis of plasma cells, 257

RUTIN, effect on toxicity of poly-
saccharide S. marcescens in
mouse sarcoma, 642

SACCHARIDES See Carbohydrates;
Sugars; and names of
saccharides, as Mono-
saccharides; Polysaccharides

SACCHAROMYSES, pink: See yeast

SAHAYUN METHOD: See Polysa-
ccharides

SALICYLATES, reduces activity of
'necrosin', 427
therapy, massive, altered res-
ponse of humans to typhoid
vaccine during, 496

SALIVARY GLANDS: See also Parotitis
myoepithelioma, effect of S.
marcescens polysaccharide on,
444

SALMONELLA, aertryke, fractions
produced changes in blood sugar
and toxic effects in rabbits,
214; immunizing fractions
isolated from, 222; suspensions,
changes in blood sugar and
phosphorus after inj. into
rabbit, 170

antigen: See Antigens

endotoxin, component of O antigen,
364; tolerance to, 598

effect on carbohydrate metabolism
in rabbits, 536; on glycogen
synthesis, 534

sulfonamide, action on lethal factor,
349

enteritidis, carbohydrate complex from,
effect on rat sarcoma or carci-
noma, 311

filtrate, activity in malignant
neoplasms, 245

(Gärtner) endotoxin, chemical nature
amino acid and sugar component of,
617

infections in rabbits, relation to
hyperglycemia, 111

Gärtner: See Salmonella enteritidis
isolation of bacterial pyrogens, 751
paratyphosa, A and B, toxicity lysed
young cells, 200

B, infections in rabbits, relation
to hyperglycemia, 111; pre-
cipitable substances from, 106

B(schottmülleri), ext. inhibition of
bacteriophage action by, 203
endotoxins, tolerance to, 598, 666

polysaccharides: See Polysaccharides
schottmülleri (paratyphoid B): See
Salmonella paratyphosus

supestifer, exts., inhibition of
bacteriophage action, 203; cholera
types and potency for tumor
hemorrhage prodn., 433

typhimurium, effect of bacteria and
their carbohydrates on leukocytes,
717

endotoxins, biological properties,
action on tumors in mice, 385

endotoxin had ability to reduce
temperature markedly, 552;
induced decidua-placental
hemorrhage, 384; lymphocytic
lesions provoked by, 569
fractions, reactions of malig-
nant neoplasms to, 293

SALT, concentration, inactivation
of coliphage delayed by
very high or very low
concn., 317

fever, character, 38, 48, 52,
173; pharmacology of, 175

inert, formed by combination of
calcium with non-protein
thermogenic factor from
B. coli and E. typhosa, 65

solutions, intravenous, con-
taminated from distilled
water, 234; dangers from,
38, 39; pyrogens, 327

intradermal injection, effect
on leukocytes, 112

physiologic, 427; loss of pyro-
genic activity, 610

preferred diluent of Pyromen,
722

prepn. and apparatus for
intravenous and subcutaneous
use, 206, 251, 341, 457, 459

reactions after intravenous
injections, review, 213; in
transfusion, 332

tests on nonpyrogenic, 458

SALVARSAN: See Arsphenamines

SANARELLI REACTION, after thera-
peutic anti-typhoid inj.,
426

-Shwartzman phenomenon with fatal
outcome after typhoid
vaccine therapy, 408

SAPROPHYTES, vaccine, Neosaprovitan,
193; (Neosaprovitan B)
therapy, effect on hemato-
poietic system, 168

SAPROVITAN: See Fever, thera-
peutic; Nervous System,
diseases: Saprophytes

SARCOMA: See also Cancer; Tumors;
Organs and Regions involved;
specific therapy, as Coley's
Fluid; Polysaccharide (of
Shear)

animal, effect of Shwartzman re-
action on, 241

carbohydrates, in blood serum
during course of, 180

fibro-and chondrosarcoma, reactions
to S. marcescens polysaccharide,
494

lympho-end results with Coley's toxins
and/or radiation, 149

mediastinal, with Coley's fluid
therapy, 136; melanotic, therapy, 46

mouse, action of bacterial toxins
on 372, 373; of S. marcescens
polysaccharide, 321, 372, 414,
464, 489, 565, 697

(180) 373; E. typhosa produces re-
gression and necrosis in, 233;
gram-negative bacilli on growth,
239; meningococcus filtrate, 183,
255, 266; S. marcescens poly-
saccharide, 245, 231, 372, 435
520, 642; of S. paradysenteriae,
immunizing properties, 386;
typhoid, anti-typhoid serum, 233

(37) B. coli filtrate and colchicine
on, 294; hemorrhage, effect on, of
ascorbic acid, 275; Ps. aeruginosa
fraction, effect on, 657; S.
marcescens polysaccharide, 443,
489, 515, 565; effect of, on
blood pressure in, 515; of
immunization by, on activity, 565;
of cortisone protection against
activity, 697; reduction of toxicity
by adrenal extract, 469; vascular
reaction after, 437

myosarcoma, of uterus, with metastases
to bone, 210

therapy, See under specific headings,
as Coley's fluid, Polysaccharide
(of Shear)

SCHIZOPHRENIA: See Dementia praecox

SCLEROSIS: See also Arteriosclerosis
multiple, altered response to typhoid
vaccine during massive salicylate
therapy, 496

effects of artificially in-
duced fever on electro-
encephalograms in, 713
therapy (brain tissue ext. of
V. Muralt) suggested trial, 506
with Neosaprovitan B.,
168; with Piramen, 771;
with Pyrifer, 712, 761

SEITZ PAD: See Filtration

SELAS CANDLE: See Filtration

SEPTICEMIA, therapy, admin. of
colloidal toxic bacterial
substances, 70

SERRATIA marcescens, analysis of
hemorrhage-producing fractions,
374

marcescens, antigen: See Antigens
cultures, induced fever in dogs,
effect on gastric secretion,
79

----- (killed), effect on
hunger and appetite in fever,
in dogs, 74

filtrate, effect on mouse
tumors, 330

----- immunological re-
lationship between Shear poly-
saccharide and, 405

----- inj. into mice with
sarcoma #37, 275

----- sepn. of hemorrhage-
prodg. fraction, 312, 381
hemorrhage prodg. fraction,
lethal dose, 381

----- fraction,
nature of, 321, 371

illness in man after inhalation
of, 455

polysaccharides: See Poly-
saccharides

toxin: See also Coley's Fluid
toxin, necrosis of kidneys in
cortisone-treated rabbits
after inj., 734

----- pyrogenic agent from,
named "pyrotoxin", 53;
therapy of exptl. tumors in
dog, 33

SERUM: See also Blood; Typhoid,
antiserum

SERUM; See: Blood; Typhoid; Antiserum
blood and plasma changes in fevers,
117

increases pyrogenicity of E. coli
cultures, 499

globulin: See Blood proteins

homologous antibacterial, destroys
phage inactivating power, 211

horse, bacterial production of
pyrogen in, 277

in therapy of arthritis, 94

missed contaminations in, role of
cryophile bacteria, 487

of fever patients, thermostable
bactericidal substance in, 256

polysaccharide, level in exptl.
inflammation, 614

level in malignancy and in other
pathological conditions, 615

Seitz serum filter #3 employed in
prepg. infusion fluids, 251

therapy, trial of maternal blood in
acute anterior poliomyelitis,
555

SHEAR'S POLYSACCHARIDE: See Poly-
saccharides, from S. marcescens

SHIGELLA dysenteriae: See Antigens;
Dysentery, bacilli; Poly-
saccharides

SHOCK, from therapy with mixed toxins
(Coley's Fluid), 29

phase produced by "pyrotoxin" in
animals or humans, 53

surgical, hyperthermia with, 530

therapy, of arthritis with typhoid
vaccine, 66

tourniquet, effect on peripheral
capillary circulation in un-
anesthetized mice, 436

SHWARTZMAN PHENOMENON, action of col-
chicine and E. typhosus ext. on
tissue metabolism, 249

active principles in, 261

after therapeutic anti-typhoid inj.,
426

chemical nature, 224

comparison of Arthus phenomena and,
281

effect of inj. of products from
gram-negative bacillus upon
growth of mouse sarcoma #180,
239

fractions from antigen of Sh.
dysenteriae tested for, 546
General Sanarelli-, fatal outcome
after typhoid vaccine therapy,
408

immunity, effect of reticuloen-
dothelial blockade on, 472
intradermal inj., conditioning
procedure preparatory for, 500
mechanism in rabbit, 730

production, in rabbits with B.
coli filtrate, 236

with sulfonamide conjugate with
meningococcal filtrate, 347

reaction of meningococcus filtrate
analogous to, 255; tumors to
bacterial toxins in animals
unsusceptable to, 198; frac-
tions from Salmonella
typhimurium, 293

reactivity of malignant neo-
plasms to bacterial filtrates,
245

relation of purpuric menin-
gococcemia to exptl., 475

relationship of pyrogens to, 749
skin reaction to S. marcescens,
405

with E. typhosa filtrate, effect
effect on transplantable tumors,
241

SKIN, biopsies, 678, 679

blood supply, effect of dibena-
mine on vascular response to
typhoid vaccine, 627

mode of action of bacterial endo-
toxins in animals, 521

observations of vascular reactions
to inj. of typhoid endotoxin
in rabbit, 518

peripheral circulation during
experimental fever, 231

vascular reactions (leukopenia,
diapedesis) in rabbit to
typhus endotoxin, 477

SKIN

- cancer, diagnosis, serum polysaccharide level in, 615
 - diseases: See also Neurodermatitis
 - chronic dermatoses treated with Pyromen, 724
 - clinical evaluation of new pyrogenic agent (Pyromen), 660
 - protein therapy, 95
 - therapy, regulation of cholinesterase in blood by pyrifer and ultraviolet rays, 510
 - typhoid shock therapy, 461
 - reaction: See also Allergy
 - (hemorrhagic-necrotic) to B. typhosus filtrate, 152
 - to bacterial filtrates, 261
 - (Meningococcus) filtrates in therapy of mouse sarcoma #180, 183
 - temperature, relation of peripheral to internal temperature in fever, 10
- SNAIL, juice, hydrolysis of inactive polysaccharide of type IV pneumococcus, 177
- SODIUM, chloride: See also Salt
 - pyrogenic, 457
 - retains water in body, alimentary fever, 174
 - citrate, prepn. of non-pyrogenic fluids, 341
 - nucleate, effects on hunger and appetite in fever, 74
 - periodate, 684
- SODUKU: See Rat-Bite Fever
- SOIL, bacteria, enzyme capable of degrading capsular polysaccharide isolated from, 554
- SOLUTION: See also Alkali, solutions; Dextrose, solution; Injections; Salt, solutions
 - hypertonic, hypotonic or isotonic may cause 'saline fever' after oral dose, 173
 - parenteral: See injection, intravenous; Solutions

pyrogenic: See Fever; Pyrogens and specific headings

- SPINAL CORD: See also Neurons, Nervous System; Nervous System, Sympathetic; Polysaccharides; Pyrogens
- regeneration, associated with cellular reaction to purified bacterial pyrogen (Pyromen), 688, 689, 690, 703, 736, 754, 755, 757, 773, 778
 - conduction of impulses in fibers of cat, 725, 778
 - effect of Pyromen, 688, 689, 690, 703, 736, 754, 755, 757, 773, 778
 - mechanism in cat, 688, 689, 690, 725, 740, 757
 - of cat and dog, possible site of action of bacterial pyrogen (Pyromen), 688, 689, 736
 - proximation of, 703
 - transection, in dogs and cats, cellular responses to bacterial pyrogen (Pyromen), 688, 689, 690, 703, 736, 754, 757, 773, 778
- SPINAL FLUID, Donaggio's 'inhibition phenomenon' in, 195
- 'inhibition phenomenon' during induced fever (proteins and vaccines), 197
- SPINE, sarcoma, therapy with mixed toxins (Coley's), 136
- SPLEEN, accumulation of neutrophils in, after inj. of E. typhosa filtrate, 271
- changes during leukopenia, 259
 - excision, leukopenia and leukocytosis in rabbits, 80
 - prevent exhaustion of thrombocytes, 118
 - hemopoiesis in rabbit and cat induced by (Pyromen), 692
 - hypertrophy, diminished after inj. of anti-typhoid vaccine in E. typhosa infection, 517

SPLEEN

- irradiation, invoked fever (typhoid vaccine) in leukemia enhances effect of, 424
- myelocyte and granulocyte prodn. after admin. of prepn. from Pseudomonas (Pyromen), in rabbit, 691
- periarterial tissue, intensified prodn. of lymphatic nodules after infection with B. mono-cytogenes, 265

SPOROTRICHUM schneckii, polysaccharides of, 503

SPUTUM, cultures, after inhalation of S. marcescens, 455
vaccine from organisms found in, used in therapy of asthma, 89

STAPHYLOCOCCI, albus, grown in gum

- acacia solns. produce febrile response, 278
- aureus, filtrate, activity in malignant neoplasms, 245
- killed, protection of animals against bacterial intoxication by adrenal cortex exts., 189
- titer, bacterial products increased, 110
- toxin, therapy of exptl. (in dog) tumors, 33
- autolysate, produced marked leukocytic response, 342
- enterotoxin, inj. reactions compared with pyrogen admin., 398
- filtrates, antihemolytic effect on blood, 176
- glycolipids from, provoked sharp increase in resistance, 393
- infections, therapy with pyrifer, 178
- toxin, 18
- therapy of progressive paralysis with, 54
- with protein characteristics inhibit chemotaxis of leukocytes, 365
- vaccine prevention and therapy of influenza and sequels, 98

STARCH, colloids, inactivation of coli-phage, 317

STARVATION: See also Fasting, Hunger
blood lipids increase in early phase, 325

STATE SERUM INSTITUTE, vaccine prepared by, 639

STERILIZATION, cause and elimination of reactions after intravenous infusions, 205
elimination of pyrogenic substances in prepn. of intravenous solns., 505
of distilled water, 164
of transfusion and infusion equipment, 246, 632
relation of proper prepn. of solns. for intravenous therapy to febrile reactions, 314
removal of thermostabile pyrogens from penicillin, 432

STERIODS, in urine: See Urine, steroids
method of assaying for protective action against toxic material (typhoid vaccine), 449

STOMACH, motility, depression by pyrogenic substances, 378;
without elevation of temperature after inj. of pyrogen, 355
physiology, pyrogens depress normal function, use in ulcer therapy, 600
secretion, action of mono- or polysaccharides on dog, 267
both volume and acidity reduced by pyrogens, 596
depression by pyrogens, 378
——— with pyrogens and antipyretics without fever, 685
during fever (induced in dogs), 79
in fever and infectious diseases, 201
inhibition by pyrogens, 580
ulcers: See Peptic Ulcers

STOOLS, bloody, reaction to fever induced by killed B. coli, 188

STREPTOBACILLIN, Ducrey's B. ext. 141; therapy of chancroid, 142

STREPTOCOCCI, erysipelatis, culture material in therapy of carcinoma or sarcoma, 19

serum in therapy of cancer, 25

toxins, 18; mixed, of Coley: See Coley's Fluid

toxins and serums in therapy of malignant tumors, 23, 24

filtrates, anti-hemolytic effect on blood, 176

hemolyticus, biosynthesis of polysaccharide, 753

filtrates, activity in malignant neoplasms, 245

inj. in mice modified course of virus pneumonia, 495

labile antigens isolated from, 272

infections, effect of inj., in glomerulonephritis, of plasma from, 564

MG, injection in mice modified course of virus pneumonia, 495

polysaccharide inhibitory effect on mumps virus, 525

pyogenes, toxin, therapy of exptl. (in dog) tumors, 33

salivarius, inj. in mice modified course of virus pneumonia, 495

scarlatinae, culture washings, effect on phagocytic activity of leukocytes, 172

toxins, therapy of progressive paralysis with, 54

vaccine, in therapy of bronchial asthma, 89

prevention and therapy of influenza and sequels, 98

subpyrexial doses in chronic arthritis, 212

therapy of varied diseases, 83

STREPTOMYCIN, method for removing pyrogens from aqueous solns. of acid salts of, 655

STRESS, environmental, 111

adrenal changes in rat by infection with E. coli, 608

factor in uveal inflammation, 774, 775

reactions of Shear polysaccharide, 750

thrombopenia in, 118

SUGARS, glucose, lactose and galactose increase growth of E. coli and stored polysaccharide, 568

in blood: See Blood, sugar reducing, 133

component of endotoxin of S. enteritidis, 617

-----polysaccharide

fraction of B. prodigiosus, 372

-----polysaccharide of

type IV Pneumococcus, 177

in ext. of culture of E. typhosa 106

SULFADIAZINE, 445

SULFONAMIDES, effect on lethal factor of bacterial toxins, 349

effect on toxic and antigenic actions of endotoxins of gram-negative bacteria, 414

--on toxin-induced hemorrhage, 364

para-aminobenzenesulfonylacetylamide, conjugate with bacterial filtrate used for Shwartzman phenomenon, 347

SULFOSIN(1% suspension sulfur in olive oil), therapy of dementia praecox, 230

SULFUR: See also under names of compounds

colloidal, fever therapy in asthma, 637

therapy, hyperthermic therapy of schizophrenia and paralytic dementia, 240

SULFURIC ACID, in test for pyrogen in distilled water for intravenous inj., 160

SURGERY: See also under organs and diseases, as Adrenals, excision; Nerves, vagus; Uterus, tumors; under specific operations, as Sympathectomy; Thyroidectomy, etc.

SURGERY

contributions in basic medical sciences to, 284
therapy of stomach and duodenal ulcers, 113

SYMPATHECTOMY, in exptl. fever, 231, 232

SYNOVIAL MEMBRANE, hypertrophy, after intra-articular inj. of typhoid antigen in rabbits, 500

SYPHILIS: See also Neurosyphilis; Paralysis, General; Tabes Dorsalis; etc.

asymptomatic serologic, resistance to endotoxins of enteric bacilli in man, 539

therapy, influence of artificial fever on nonspecificity, 308

pyrifer, a product from B. coli grown in milk, 260

pyrogenous, 698

TAB (typhoid, paratyphoid A and B)VACCINE: See Fever, therapeutic; Paratyphoid, vaccine; Typhoid, vaccine

TABES DORSALIS: See also Neurosyphilis
therapy with Neosaprovitan B, 168; pyrifer, 167, 171; suspensions of killed Ps. scissa and ureae, 208

TACHYCARDIA: See Heart, rate

TALC: See Inflammation, experimental

TEMPERATURE: See also Cold; Heat; etc.

effects, factors in action of bacterial endotoxins, 413

on reactions to typhoid-paratyphoid vaccines and other pyrogens, 579

on response to typhoid-paratyphoid vaccine ("Pyrexin"), 577

on toxicity of bacterial filtrates for tumor-bearing mice, 235

environmental, 199, 200

effect on rate of bacteriophage inactivation by filtrates, 317

low, role of cryophile bacteria in missed contaminations in biologic products, 487

of liquid air, method of disintegrating typhoid bacilli, 31

of solns., cause of chills following intravenous therapy, 288

relation to inhibition of gastric secretion by pyrogens, 580

variations, relation of S. enteritidis-paratyphoid B infections to hyperglycemia in rabbits, 111

TEMPERATURE, BODY: See also Chills; Heat; Muscles, temperature; Rectum, temperature; Skin, temperature

augmentation of effects of pyrogens by homologous plasma in rabbits, 591

constitutional differences

during pyrifer admin., 712

effect of heat or cold applications to heat regulating center on action of induced and antipyretics, 78

heterologous protein sensitizes cells of heat-regulating centers, 77

high: See Fever

in hypophysectomized or tuberinjured dogs, 184

influence on thermal polypnea of substances affecting, 528

low: See also Chills

action of dysentery toxins, 447
after intra-abdominal inj. of heat-killed Rickettsial bacilli, 453

body fluid redistribution in induced hypo-and hyperthermia, 673

effects of typhoid-paratyphoid vaccine in rats, 318

emotional hypothermia in rabbits, 643

hypothermic and antipyretic effect on prepn. of ACTH, 742

————— effects of bacterial vaccines, 356

————— factor of B. dysenteriae, Shiga, 258

TEMPERATURE, BODY

low
 in mouse as bioassay of endotoxin
 protection factor in impure
 penicillin, 552
 mechanism, effect of typhoid-para-
 typhoid vaccine on, 289
 metabolic activity of central
 nervous system as it affects
 thermostatic behavior of body,
 527, 529
 method (thermocouple box) for large
 scale testing for pyrogen in
 in rabbit, 448
 oscillation without cause, 11
 physiology and pathology of hypo-
 physis temperature-regulatory
 function, 307
 rectal: See Rectum, temperature
 regulation, after inj. of proteose and
 typhoid vaccine, 104
 effects of ethyl carbamate
 (urethane) on, 578
 pyrogens may interfere at lower
 level of brain stem 577
 rat tail as a part for, 350
 role of viscera in, 346
 relief of typhoid chill has no effect
 on, 296
 wide range in normal rabbits, 501

"TESCHEN" DISEASE, therapy, bacte-
 febrin not effective for, 765

TETANUS, toxin, effect on leukocytes,
 172
 toxin, fever toxins, 18
 inhibit chemotaxis of leukocytes,
 365

TETRAHYDRO-BETA-NAPHTHYLAMINE, effect
 on hypophysectomized rabbits, 307

THERAPY: See also under names of
 diseases and therapeutic drugs
 intravenous: See injections,
 intravenous
 nonspecific, bactericidal effects
 found in organic compds.,
 inorganic salts, biologic pro-
 teins, and suspensions of dead
 bacteria when injd. into rabbits,
 186

use of typhoid vaccine, 56
 parenteral: See injections,
 intravenous

THERMOCOUPLE BOX, method for large
 scale testing for pyrogen in
 rabbit, 448

THOROTRAST, injections followed
 by typhoid vaccine in rabbits,
 areas of hemorrhagic necrosis
 developed and toxic effect of
 vaccine enhanced, 472

THROMBOANGIITIS, obliterans, therapy
 with typhoid vaccine, 144, 333,
 335

THROMBOCYTES: See Blood, platelets

THROMBOPENIA: See Blood, platelets

THROMBOSIS, extensive in heart,
 lungs and liver after intra-
 articular inj. of typhoid
 antigen in rabbits, 500
 purified necrosin brings rapid
 liquefaction of, 427

THYMUS, nucleic acid, 430

THYROID, importance in fever, 145
 preparation, increases hyper-
 glycemic response to *B. coli*
 and *B. proteus* killed cells
 and glycogenolysis, 132

THYROIDECTOMY, decreases glyco-
 genolysis, 132
 effect on pyrogen fever in
 rabbits, 645
 on Pyromen responses, 754

THYROTOXICON: See Protein, split
 products

THYROXINE, increased production in
 burns, 714

TILLET, Goebel and Avery, sub-
 stance "C" isolated from
 type IV pneumococci, 177

- TISSUE: See also Cells: Muscles; Nerves; etc.
 alteration during fever in infections and by hyperpyretics, 122
 anoxia, effect of bacterial endotoxins on carbohydrate metabolism, 536
 connective, endocrine and cellular response to admin. of bacterial pyrogen (Pyromen), 688
 mast cells, alteration induced by bacterial pyrogens, 678, 679; response to bacterial pyrogens, 732
 plasma cells found in, 257
 culture, sarcoma #37 in, effect of tumor-necrotizing agents (emetine HCl and polysaccharide S. marcescens), 498, 764
 destruction, influence on toxicity in fever, 125
 permeability, edema after tourniquet pressure caused no increase in, 436
 reaction to pyrogenic prepn. from Pseudomonas, 691
 scar, prevention: See Nerve, Piromen
- TOLSEROL, intravenous inj. diminished reaction to Pyromen, 722
- TOURNIQUET, shock effect on peripheral capillary circulation in un-anesthetized mice, 436
- TOXINS: See also Bacteria; under names of bacteria as, Eberthella typhosa; Serratia marcescens; Staphylococci; Typhoid; etc.
 bacterial: See also Diphtheria, toxins; Tetanus, toxins chemical action of, 58
 colloidal, admin. during septicemia, 70
 effect on hunger and appetite in fever, 74; on leukocytes, 35; on tumor hemorrhage prodn., 433
 endotoxins of gram-negative bacteria, effect of sulfonamides of, 414; temperature factors in action, 413; partial, in protoplasm, 50
 reactions of mouse carcinomas to, 225; tumors, 198
 Coley's (mixture of erysipelas and prodigious toxins): See Coley's Fluid
 extraction, pyrogens removed by cyclic or heterocyclic acids, with ETOH and BzOH, cooling and filtration, 570
 link between toxic nucleoproteins and complex O antigens and variants of bacteria, 412
- TRAUMA: See also Shock, surgical injury to blood cells may release pyrogens, explains fever activity, 427
- TRENDELENBURG POSITION, in extreme hypotension from pyrogens, 420
- TRICHLORACETIC ACID, extraction of bacterial antigenic fractions, 301
 extracts of bacterial preps. produce leukopenia on inj., 366
- TRYPANOSOMA nagana, toxin, effect after section of medial hypothalamus in rabbit, 55
- TRYPSIN, digestion, method for prepg. antigenic fractions from B. typhimurium and B. typhosum, 301; for antigenic substances from typhoid bacillus, 309
- TUBERCLE BACILLI, culture, anti-biotic action of B. subtilis and its secretion on; also cultures of B. mesentericus vulgaris, 119
 extracts from cultures combined with bacterial proteins, use in uveitis, 158
 fraction, chemical analysis, 216
 toxins, in hyperthermia, 18; filtrate activity, 245
- TUBERCULIN, allergy, 681
 polysaccharides, 749

TUBERCULIN

- protein fractions, effects on leukocytes, 749
- injection, stimulation of plasma cells by, 257
- reaction, through nonspecific bacterial protein, 15
- therapy of neurosyphilis, 140; of progressive paralysis, 12, 41, 54, 59; of uveitis with tuberculoase, 158

TUBERCULOSE: See Tuberculin

TULAREMIA, adrenal cortical hormone in prevention of depletion of cholesterol and prolongation of survival, 608

- therapy, vaccine, B. tularensis, combined S. typhosa pyrogens, 746

TUMORS: See also Cancer; Sarcoma:

- names of Organs or Regions
- adrenal hormones, influence on toxicity of tumor-damaging substances, 622
- antineoplastic substance in cultures of P. notatum, 428
- pathology, gross and microscopic, after S. marcescens, 456
- preparations, of E. coli filtrate, 243, 244; of S. marcescens polysaccharide, 504; potency determinations of, 380
- properties, of S. marcescens polysaccharide, 438, 484
- therapy, chemo-, hormones, with cytostatic, radioactive bacterial products, 590; in mice, 312, 406; with radioactive material, 590, 613; temperatures, high environmental, 87, 413; azobenzene product, resembling S. marcescens in activity, 613; cobra capello venom, 30; colchicine with S. marcescens, 249
- bacterio-, 433, 483; with erysipelas toxins, 23, 24, 30. See also: Coley's fluid; with hemorrhage producing fractions from E. coli filtrates, 235, 236, 243, 244; dosage, minimal, 236; properties, 243;

- meningo, filtrate, 235; separation, 244; with antigenic fraction from Salmonella, 349; Sh. dysenteriae, 363; with hemorrhage-producing fraction from S. marcescens, 312, 330, 371, 374, 390; cell changes after S. marcescens, 485, after other materials, 590; toxins of Coley, mixed, on prevention or regression, 220; on metastases, 220

therapy, protein, split products in, 92, 93

pyrogens, necrotizing activity, 749

reticuloendothelial cytotoxic substance, 672

reactions, to uneliminated waste products, 390; to toxic materials, 502

reversibility of therapy, 147

review of literature on, 651

serum polysaccharide levels after bacteriotherapy, 615

toxins, See Tumors, bacteriotherapy

vascular reaction to bacterial polysaccharides, 467; hypotension in malignant tissue, 695

venom, snake: See Chemotherapy

TURPENTINE: See also Inflammation, experimental leukocytosis induced in hypertensive dogs by, 407

TURTLES, cardiovascular changes with polysaccharide preps., 589

TYPHOID, antiserum hemorrhagic necrosis and regression of sarcoma #180, 233

bacilli: See Eberthella typhosa

antigens: See Antigens

immunity, agglutinins increased when previously immunized with colon bacilli, 62

studies, 63

mortality, foreign protein therapy modified rate, 100

therapy, by intravenous inj. of polyvalent sensitized typhoid vaccine sediment, 63

TYPHOID

- therapy
 - forty-eight patients positive for E. typhosa injected with anti-typhoid vaccine, 517; protein (deutero-albumose) parenteral admin., 71; with intravenous inj. of albumose, 61; with killed cultures of B. pyocyaneus, 20; with pyrifer, in children, 533, 574; with vaccine, 64, 72
 - tolerance to action of endotoxins of enteric bacilli in convalescents, 666; toxic effects of somatic antigens of enteric bacilli in convalescents, 598
 - toxin, pyrogenic action of anaphylatoxin, 51
 - vaccine, ACTH-like effect of fever in malignant hypertension, 661
 - adrenals, cortex after, 619; desoxycorticosterone no protection after, 300; effect of adrenalectomy on leukocyte level after, 537
 - amidopyrine, for prevention of fever, after, 470
 - anaphylaxis, fatal, after, 426
 - antibody formation after, 601
 - augmentation of effects by homologous plasma, 523, 636
 - bacteria, removal from splanchnic area after, 762
 - cardiac enlargement during therapy by, 463; output, during fever, 156
 - chills after therapy, use of CaCl for, 296
 - circulation, alterations, during chill and fever, 416; cerebral, 605; capillary flow during therapy, 155; hepatic blood flow, 478
 - continuous, combined with penicillin, in paresis, 594; in ophthalmic conditions, 567
 - effect on hunger and appetite, 74
 - fever after, 72, 156, 229, 264, 276. See also: Fever
 - fever, controlled, after triple vaccine adminis., 359; effect of expanding dose, 331; modified by direct heat or cold, 78; prolonged, 668
 - hemorrhagic-necrotic activity, toxicity, enhanced by thoro-trast, 472
 - hypothalamic lesions, effect on : fever after, 289
 - intramuscular, combined with massive salicylate therapy, 496
 - intraperitoneal, 318
 - intravenous, 72, 270, 473; drip, 375
 - leukocytes after, 523, 537, 601, 619, 636
 - metabolic rate after, in neurosyphilis, 229
 - method for assaying steroids and adrenal extract for protective activity against, 449
 - oxygen consumption after, 605
 - paratyphoid, 'necrosin' yields 'pyrexin' after enzymatic degradation, 427; 'pyrexin' thermostable, nontoxic substance, 401
 - renal function affected by, 382, 561
 - respiration during fever caused by, 416
 - sediment, polyvalent, in therapy of typhoid, 63
 - therapy in arthritis, 66, 75, 82, 94, 191, rheumatoid, 313; asthma, 637; chorea, 313; dermatoses, 95, 473; gonorrhea, 391; herpes zoster, 473; hypertension, 420, 661; infections, 81; neurologic disease, 542; neurosyphilis, 12, 229, 594, 609; nonspecific. See Protein, therapy, nonspecific; opthalmic, 135, 567; sarcoma, 180, 233; shock, 461; vascular disorders, peripheral, 144; thromboangiitis obliterans, 333, 335
 - therapy, mechanism, 110; preferred as agent, 76; X-Ray, enhancement by, in leukemia, 424
 - toxicity, 449; damage to visual organs, 588; dangers, 516, 609; enhanced by thorotrast, 472; fatal, 426

TYROSINE, toxic polypeptide gives color reaction for, 363

typhoid vaccine in psychotics, 619

ULTRAVIOLET RAYS, regulation of cholinesterase in blood by pyrifera and, in therapy of dermatoses, 510

UNDULANT FEVER: See also Bacteria, abortus
in animals, effect of sonic filtrates of B. abortus in cattle positive to brucellosis, 400

UNITED STATES PHARMACOPEIA, pre-description of Finnish method of prepg. solutions, 711
pyrogen test: See under Pyrogens

URETHANE: See Ethyl Carbamate

URIC ACID, in blood: See Blood, uric acid; Urine

URINE, creatine and creatinine, clearance in therapy of carcinoma with S. marcescens polysaccharide, 464

extracts, effect on peptic ulcer, 329

(male): See Urogastrone
possessing pyrogens, have inhibited ulcer formation, 596
histidine, elimination during fever ('pyretamine' of Russo), 419

'inhibition phenomenon', during induced fever (proteins and vaccines), 195, 197

nitrogen, elimination during fever ('pyretamine' of Russo), 419; increased by foreign proteins, 124

protein, in glomerulonephritis, effect of febrile plasma, typhoid vaccine and nitrogen mustard on, 564

pyrogen, separation from urogastrone, 302, 303

steroids, (17-ketosteroids), uric acid and creatinine output, changes during injns. of

UROGASTRONE, pyrogen-free, prepn. of, 302, 303
pyrogen in urine and its separation from, 303

URONIC ACID, in fraction from culture of tubercle bacilli, 216

URTICARIA, allergic, Pyromen in, 777

UTERUS, excision: See Uterus, tumors

tumors, myosarcoma with multiple bone metastases treated with Coley's toxins after total hysterectomy, 210

UVEA, infections, therapy with "mixed toxin" from St. erysipelatis and B. prodigiosus (Coley's Fluid), 491

inflammation, therapy with non-specific protein, typhoid vaccine, 135; with tuberculo, 158; with Pyromen, 774

VACCINES: See also Paratyphoid, vaccine; Typhoid, vaccine; etc.
antichancroid, Rimbaud, 254
aspecific, intradermal vaccination in poliomyelitis, history, technics and mechanism of action, 629

bacterial: See also Bacteria, clostridium

B. coli vaccine depressed gastric motility in dog, 355;
glucose tolerance lowered, 273
effects, hypothermic and adreno-hemorrhagic, 356
general reactions after therapy of asthma, 638
therapy of bronchial asthma, 89;
of experimental tumors, 33;
Neosaproviton B, from saprophytes, use in tabes dorsalis and multiple sclerosis, 168

VACCINES

- histolytic endotoxin, thermal reactions in rabbit from inj., influence on prodn. of agglutinins, 169
- mixed, clinical response in 125 cases of rheumatic disease, 607; in prevention and therapy of influenza and sequels, 98
- standard, prepd. by State Serum Inst. in therapy of asthma in children, 639
- therapy, phenomena associated with, 83
 - subacute and chronic localized infections, 64; of influenza, 98; of acute poliomyelitis, 664
 - reversibility in antibacterial and antitumor therapy, 147
 - specific, combined with pyripher in therapy of gonorrhea in women, 161

VAGINA, hemorrhage, after inj. of bacterial endotoxins in gravid mice, 413

VAN'T HOFF'S LAW, basal metabolism in fever, 96

velocity of chemical reaction during reaction to typhoid vaccine and proteose injcs., 104

VASOMOTOR SYSTEM: See also Blood Vessels; Capillaries; Nervous System, Sympathetic, Sympathectomy

disturbances, mode of action of bacterial endotoxins in, 521

foreign protein therapy (typhoid vaccine) in peripheral vascular diseases, 144, 333, 335

VENA CAVA, impedance of blood flow near, may be factor in post-operative hyperthermia, 530

VENOCLYSIS: See Injection, intravenous

VIBRIO: See Bacteria, vibrio

VIRUSES: See also Influenza, virus; Parotitis, virus; Pneumonia, virus;

extraction, pyrogens removed by cyclic or heterocyclic acids, removal with ETOH and BzOH, cooling and filtration, 570

infection, activation of latent infection by typhoid vaccine, 588

VISCERA, effects of bacterial proteins (B. coli) on, 130

effects of nonspecific protein therapy on, 138

removal, in dogs, effects of bacterial proteins, 130

role in regulating body temperature under physiological and pathological conditions, 346

VITAMINS, B₆, does not reduce toxicity of polysaccharide S. marcescens, 642

C: See Ascorbic Acid

VOMITING, reaction to fever induced by killed B. coli, 188

VULVOVAGINITIS, therapy with pyripher, 171

WATER, contaminated, 48, 442; dangers from saline infusions of, 39; from gum acacia solns, 278; distilled water, 39, 45, triple-distilled, 505; reactions, 108, 139, 140, 164, 205, 206, 288; review, 213; stored, bacterial content, 42; recontamination, 49

'fever', 48, 105

in blood; See Blood, water content

in body, during fever, 174.

See: Fever

intradermal injection, 112

WATER

- preparation, mineral-free, 370;
- pyrogen-free, 164, 204, 205,
- 206, 234, 288, 304,
- 558; correct method, 20;
- distillation, triple, for
- saline preparations, 505; safe-
- guards against later contamination,
- 49
- testing, biological, for pyrogens
- in, 114, 160, 389
- chemical, 160, 458
- tissues, water content, action
- of sugar on, 91; binding by coll-
- oids in 'thirst' and 'salt' fever,
- 81

WHOOPING COUGH, bacteria: See
Hemophilus pertussis

WOUNDS, infection, and fever, 5

XANTHOPROTEIN, 321

X-RAY, therapy of leukemia, fever
 therapy (typhoid vaccine)
 or colchicine enhances
 effectiveness, 424

YATREN: See Chiniofon

YEAST, autolyzed, produced pyrexial
 substance, 165
 carbohydrate haptens from, role
 in bacterial anaphylaxis, 153
 filtrates, produced pyrexia,
 leukopenia and leukocytosis, 17
 injection into tumor-bearing rats,
 280; subcutaneous, in the monkey,
 369
 wild, produces pyrogen, 340

ZINC, chloride, pptn. of anti-
 genic nucleoprotein from
Ps. aeruginosa with, 587