QL
596
P83M3
SUPPL.
ENT.

The
THOMAS LINCOLN
CASEY
LIBRARY
1925

Lieut. T. L. Casey U.S.A.

presented at the request of the late

Rev. A. Matthews.

TRICHOPTERYGIA

ILLUSTRATA ET DESCRIPTA

A MONOGRAPH OF THE TRICHOPTERYGIA.

Supplement

BY

THE REV. A. MATTHEWS, M.A.

EDITED BY

PHILIP B. MASON, M.R.C.S., F.L.S., F.Z.S., &c.

With Seven Plates.

LONDON :

O. E. JANSON & SON, 44, GREAT RUSSELL STREET, W.C.

1900.

AS A RECORD OF LONG AND INTIMATE FRIENDSHIP,
AND AS A TRIBUTE OF HEARTFELT GRATITUDE AND REGRET,

I DEDICATE THIS

S U P P L E M E N T

TO MY

'TRICHOPTERYGIA ILLUSTRATA'

TO THE MEMORY OF

George Robert Crotch.

FROM HIS PROFOUND BIBLIOGRAPHICAL KNOWLEDGE I DERIVED MUCH
VALUABLE HELP WHILE I WAS ENGAGED IN THE FORMER PART OF THIS WORK ;

AND BY HIS PERSISTENT EXERTIONS ON MY BEHALF

DURING THE LAST YEARS OF HIS LIFE,

ENTOMOLOGY HAS BEEN ENRICHED BY A LARGE PROPORTION
OF THE SPECIES DESCRIBED IN THESE PAGES.

EDITOR'S PREFACE.

As I HAVE already stated in my Preface to the 'Monograph of the Corylophidæ and Sphæriidæ,' the entire collections of Micro-Coleoptera belonging to the late Rev. ANDREW MATTHEWS passed into my possession at his decease, under an arrangement, made shortly before, that I should assist him in the publication of the before-mentioned and present works, and in issuing this Supplement to the 'Trichopterygia Illustrata et Descripta.' I am pleased to have fulfilled my obligation to my much lamented friend. It is to be regretted that the Author did not live to revise these works as they were passing through the press, as I fear errors have escaped notice which he, no doubt, would have rectified.

In preparing the manuscript of the present work for the press, I have only corrected any obvious errors that were detected, chiefly in the references, otherwise the whole of the matter and system of arrangement is the Author's own work.

In the systematic and descriptive part of the work the Author has only included those species that he had an opportunity of examining himself, and for the most part contained in his own collection; others that were only known to him by description will be found mentioned in the latter part of his Preface, with a translation of the original descriptions, or a short note of their leading characters. At the end of the Preface (page 7), I have given a list, in chronological order, of some species that he had apparently overlooked, or had been described since he completed his manuscript.

Mr. MATTHEWS's collections, which now form part of my Museum, contain the types of all the *Trichopterygia* described by him, except twelve of the species described in the present work, of which I have given a list at page 11, and stated the collections in which these may be found, for the information of anyone wishing to consult the types.

Of the Plates, IX. to XIV. have been prepared for me from the original drawings made by Mr. MATTHEWS with his well-known artistic skill and accuracy, the figures having been most carefully and satisfactorily transferred to zinc by Mr. J. COLLIN, of Newmarket, and I have to thank him for the trouble and pains he has taken to make them accurate copies of the originals. Plate XV. is a photo-lithographic reproduction of that published in Mr. MATTHEWS's '*Essay on Hydroscaphæ*,' and has been carefully prepared for me by Messrs. WEST, NEWMAN, AND Co., in order to complete the illustration of all the new genera discovered since the publication of the '*Trichopterygia Illustrata et Descripta*' in 1872. The Plates have unfortunately been numbered IX. to XV. consecutively, in continuation of those in the '*Monograph of the Corylophidæ and Sphæriidæ*,' as it was at first intended to issue the two works as one volume.

I must also express my thanks which are due to Mr. G. C. CHAMPION, Mr. C. O. WATERHOUSE, and Mr. GEORGE LEWIS, for the assistance they have given me, and more especially to Mr. O. E. JANSON, who has been of invaluable service to me in verifying and obtaining references to the literature, reading proofs, preparing the Index, and other matters in connection with the printing and publication.

PHILIP B. MASON.

BURTON-ON-TRENT :

July, 1900.

PREFACE.

THE publication of the 'Trichopterygia Illustrata' in 1872 aroused among entomologists an interest in that previously neglected family which has proved so fertile in its results, that about one hundred fresh species, including seven new genera, have already been brought to light. In the preface to that work I expressed my belief that the *Trichopterygidae* would eventually prove to be one of the most numerous families of Coleoptera, and subsequent events appear to have fully justified my anticipations. The collections sent from every recently explored part of the world contain on an average three-fourths of species new to science, and these in most cases exhibit very obvious distinctive characters. The genus *Trichopteryx* forms the principal if not the only exception, for its multitudinous species often so closely resemble each other that, without the aid of a powerful microscope, or the presence of types of allied species for comparison, it is difficult to perceive in what way they differ from those already known.

In the same preface I also endeavoured to explain my views on the affinities and systematic position of the *Trichopterygidae*, and a prolonged study of those insects, together with a more perfect knowledge of their anatomy, has tended to confirm my views with regard to their affinities, but slightly to modify them in respect of their systematic position. At that time, perceiving their close alliance to the *Staphylinidae*, I proposed that in a systematic arrangement the *Trichopterygidae* should be placed immediately before the *Staphylinidae*; but I am now inclined to think that that arrangement should be transposed, and that, instead of preceding the *Staphylinidae*, the *Trichopterygidae* should follow next in succession to that extensive group; by this change their alliance would be maintained, while the transition into other Clavicorn families would be more naturally accomplished. The chief cause which has inclined me to make this alteration consists in the formation of their anterior coxæ. In all the groups which have by general consent been hitherto placed before the *Staphylinidae*, and in most genera of the *Staphylinidae* themselves, these limbs are long and free, or, as they are sometimes termed, prominent—i.e., unconfined by any coxal cavity; but throughout the *Trichopterygidae* the anterior coxæ are placed in a cavity of the prosternum, which with very few exceptions is perfectly closed on both sides. This is an important anatomical character, and strongly indicative of affinity among groups in every section of the whole Order.

Within the last few years the importance of the skeleton of the under side in systematic arrangement has been brought into a prominent position, chiefly through the exertions of Dr. Le Conte and Dr. Horn. And, working upon their lines with the aid of more powerful instruments, I have discovered some few points in the skeleton of the *Trichopterygidae* which I had previously overlooked. The most noticeable of these exists in the size and form of their *mesosternal epimera*. In the *Trichopterygidae* this portion of the skeleton is unusually large, and extends from the outer margin of the body to the intermediate coxal cavity, inclosing the whole of its outer side. This enlargement of the mesosternal epimera prevails with very little variation through all the genera of the family, and is in truth one of its most distinctive characters. It is clearly shown in the plates of the new genera—any one of these may be regarded as a fair type of its older allies. The part which I had supposed to be this epimeron, and which in size and shape closely resembles that part of the skeleton in all other Clavicorn families, proves to be the inflexed apical angle of the *episternum* of the *metasternum*. The figure of the under side of *Nanosella fungi* in Pl. xiii. A, is an apt illustration of all that I have endeavoured to explain in the foregoing sentences.

I have placed *Limulodes* at the commencement of the *Trichopterygidae* on account of the similarity of its outline to *Hydroscapha*, and have also separated it from the other genera as a distinct tribe, which may be characterized thus: "Head, pronotum, and elytra together forming a compact cone, furnished at its apex with long setae." I have also good reason to believe that other species belonging to this Tribe will eventually be discovered.

I must here add a few remarks on an important essay on this Family by Dr. C. Flach, of Vienna, published in the 'Verhandlungen der Kaiserlich-königlichen zoologisch-botanischen Gesellschaft in Wien,' xxxix. 1889, pp. 481-532.* I am sorry that I cannot do this in a more satisfactory manner than by merely transcribing, or rather translating to the best of my ability, Dr. Flach's original descriptions. Some few, of which I have received types from Herr Reitter, will be found described by myself in the course of this work, and, so far as I can judge by those and by Dr. Flach's own characteristic figures of some others, I believe that most of his species will prove to be genuine:—

Ptenidium ovulum.—Thorax widest at the base; body egg-shaped; basal foveæ wanting; thorax with the elytra arched in one curve. Beetle of a fine chestnut brown, thorax finely margined; body large, upper side impunctate, transverse line on the scutellum without medial impressions. Hab. near the Caspian Sea.

Ptenidium Lederi.—Thorax with the basal and anterior foveæ small, but distinct; body oblong, pitchy black; thorax marked with a transverse depression at the base, surface very finely reticulate, very finely and sparingly punctured, thickly clothed with short hair; transverse line of the scutellum with a fovea on each side near the angles. Hab. Hamerat.

Ptenidium Brenskei.—Basal foveæ distinctly round, outer foveæ three times larger than the interior; thorax smooth; body pitchy black. Hab. Morea.

* Some notes and criticisms by the author on Herr Flach's essay will be found in the Ann. Mag. Nat. Hist. (6) ix. June, 1892, pp. 442-448.—ED.

Ptenidium turgidulum.—Thorax distinctly narrower than the elytra; elytra not much attenuated posteriorly; body narrow, rather inflated; as in *P. turgidum*, size small. Hab. Caucasus.

Ptenidium Penzigi.—Thoracic foveæ shallow; elytra sparsely and very finely punctured; basal and anterior foveæ of the thorax very slightly impressed; body castaneous; thorax as broad as the elytra, strongly rounded at the sides, together with the elytra indistinctly punctured, and sparingly covered with hairs; elytra attenuated posteriorly, apex obtuse; body large. Hab. Asia Minor.

Ptenidium Heydeni.—Less compressed, and less convex; thorax scarcely punctured, sides strongly rounded and broadly margined, margin castaneous, widest near the middle; elytra with seven rows of punctures on the shoulders; antennæ short, intermediate joints transverse, and yellow like the legs. Hab. Portugal.

Ptenidium obotrites.—Basal and anterior thoracic foveæ very distinct, interior much larger than the exterior; body large and broad, moderately long, covered with grey hairs. Hab. Germany, Mecklenburg.

Actidium Reitteri.—Elytra two and a half times longer than the thorax; body small and narrow; thorax slightly constricted, with the basal depressions indistinct; upper surface very closely and finely granulated, clothed with shining silver-grey hairs; thorax broad at the shoulders; antennæ rather slender, otherwise like *A. Boudierii*. Hab. Morocco.

Actidium Kraatzii.—Elytra rounded behind; thorax strongly constricted, nearly as broad as the elytra; body depressed and elongate; thorax finely reticulated, marked with setigerous punctures as in *Ptilium*, basal depression obsolete, strongly constricted; elytra with a shallow depression on each side of the suture, diffusely granulated and clothed with long hair, three times as long as the thorax; antennæ shorter than half the body, dark yellow. Hab. Hyères.

Actidium variolatum.—Thorax distinctly narrower than the elytra; elytra strongly convex, inflated towards the middle; antennæ and legs piceous brown; basal depression distinct; reticulation strong; head and thorax closely marked with large, umbilicated punctures, as in *A. coarctatum*; upper surface very shining; elytra granulated in fine diffuse curved rows; antennæ as long as half the body, with a distinct two-jointed club. Hab. Dalmatia.

Ptilium vexans.—The lateral channels of the thorax rather longer than the medial channel, straight and convergent anteriorly, the medial channel not extending to the basal or apical margins; elytra transversely and distinctly granulated; body small, convex, and thinly pubescent. Hab. Corsica.

(*Ptilium*) **Sahlbergi.**—Hinder angles of the thorax strongly rounded; upper surface very finely reticulated and pubescent, interspersed with larger setigerous tubercles; elytra margined, with the sutural angles rather acute; antennæ not half as long as the body, more piceous than the legs. Hab. Finland.

[*Obs.*—From the foregoing description, together with the excellent figure given by Dr. Flach, it is evident that this species cannot belong to *Ptilium*, therefore the subgeneric term *Trichoptilium*, proposed by Dr. Flach, will have to be used as its true generic name, and should take its place in the General List immediately before *Actidium*.]

Ptilium Hopffgarteni.—Large, black, head one quarter narrower than the thorax. In the male the fifth ventral segment slightly widened behind, the seventh with a transverse impression; hinder tibiae with a fascicle of setæ on the inner side above the apex. Hab. Croatia.

Ptilium Lederi.—More robust than the preceding; head one-third narrower than the thorax; thorax broadest before the base. This is the largest of the genus. Hab. Caucasus.

Ptilium Schwarzi.—Fuscous black, clothed with grey pubescence, strongly alutaceous, smaller than *P. Caledonicum*; thorax strongly constricted, with the hinder angles acute. Hab. Mecklenburg and Caucasus.

Ptilium Reitteri.—Eyes without colour, placed on an elevated excrescence, with 1-3 very minute facets; body moderately convex, faintly reticulate, strongly alutaceous;

thorax broadest in the middle, indistinctly margined, small, shining, bright reddish-yellow; thorax and elytra equidistantly tuberculated. Hab. Caucasus.

Ptilium Oertzeni.—Eyes black, moderately convex, with 8–10 distinct large facets; thorax widest near the middle, sides rounded, closely tuberculate in transverse rows, pitchy black; elytra separately rounded at the apex. Hab. Griechenland.

[Some of the foregoing species are described at length by Herr Flach, and figured, in the *Wien. Ent. Zeit.* vi. 1887, pp. 177–180, pl. ii.—ED.]

In *Wien. Ent. Zeit.* vii. 1888, p. 103, Dr. Flach described the following species of *Trichopteryx* :—

Trichopteryx rufilabris.—*T. lata*, Motsch. exacte fere statura, lata, fere quadrata, capite thoraceque nigropiceis, elytris atrombentibus, margine extrema, labro superiore, ore, antennis unicoloribus, pedibusque late rufescentibus; thorace toto et omni superficie transversim reticulatis, tuberculis modicis, ordinibus plus minusve arcuatis, transversis, haud dense obtectis, et minus dense fulvo-pilosis; caput oculis minoribus, ocellis c. e. a. 20 visibilibus; caetera omnia *T. lata* similia; thorace paulo ante angulos posticos latissimo; elytris latiore, flavo-marginato; elytris latitudini aequilongis; corp. long. 0·92, lat. 0·56 mm.

Patria.—Talyseh, Lenkoran.—Leder com. Reitter.

This species might be taken for *T. lata*, but can easily be distinguished by the conspicuous reddish yellow colour of its mouth, pale yellow antennae, more shining surface, and especially by the eyes, which are only half as large, and by the dark red colour of the elytra.

In the same paper (pp. 101, 102), this author described at length the following four species :—

Ptilium Sahlbergi.
 „ *Hopffgarteni.*
 „ *Lederi.*
 „ *Oertzeni.*

But these have been already noticed in this Preface.

In the *Deuts. Ent. Zeit.* xxx. 1886, p. 248, Dr. Flach described at length *Ptilium Œdipus*; but I found, from types kindly sent by Herr Reitter, that this species is identical with one described by myself in the *Ann. and Mag. Nat. Hist.* 1889, p. 195, under the name of *Ptilium obcaecatum*. Dr. Flach's name must therefore be used in future for this species.

L'Échange, *Revue Linnéenne* (Lyon), No. 50, Feb. 1889, p. 4.

In this periodical Mr. Rey very briefly notices many *Trichopterygida*, which he considers to be new species, but it is not possible to form any opinion of their validity from the short remarks appended to their names. All that I can do is to transcribe his remarks, and to leave the identification of the species to those who may have an opportunity of examining the original types.

Ptenidium obsoletum, Rey.—Nearly allied to *Pten. formicetorum*. It is rather larger and more convex; the punctuation of the elytra is almost obliterated, and without order or series.

Villié-Morgon; St. Raphael.

Ptenidium varicorne, Rey.—Appears to have the antennae rather longer and rather paler at the base than *P. fuscicorne*. It is probably a variety.

Bugey.

Ptilium nitidulum, Rey.—More depressed, more shining, and less pubescent than *P. kunzei*, to which it is very closely allied. The prothorax is more distinctly margined at the base, and the elytra more rugose, more parallel, and more obtuse at the extremity, etc.

Lyon: Villié-Morgon.

Trichopteryx pubescens, Rey.—Is smaller than *T. lata* (Motsch.), and slightly larger than *T. Marseullii* (Matth.), with the elytra less quadrate, with longer pubescence, etc. St. Raphael.

Trichopteryx castanea, Rey.—Is intermediate between *T. brevipennis* (Erich.) and *T. sericans* (Heer). The elytra are dark chestnut, very finely rugose, and very finely pubescent, etc.

Villié-Morgon; Cluny.

Trichopteryx spissata, Rey.—Is rather more oblong, and more convex than *T. sericans* (Heer), subsemicylindric, with the elytra more dilated, and more obtuse at the extremity; the posterior angles of the prothorax are slightly produced, giving the appearance of a true *Ptilium*.

Cluny.

Trichopteryx subnitens, Rey.—Generally more shining than *T. sericans*. It is probably a local variety.

Collioure.

Trichopteryx opacina, Rey.—Slightly smaller, slightly narrower, deeper and more dull black, and more finely rugose than *T. sericans*, with the antennæ altogether black, and the femora sometimes more or less dusky (var. *femorialis*, R.); the elytra are concolorous and have their sutural angle more straight, etc.

Lyon; St. Raphael.

Pteryx subtruncata, Rey.—This is perhaps a local variety of *P. suturalis* (Heer), but nevertheless the prothorax is more contracted behind, the elytra are narrower, more parallel at the sides, more abbreviated and more broadly truncate behind, etc.

In rotten stumps of pine, St. Raphael.

Ptinella elata, Rey.—Very similar to *P. aptera*, but the prothorax has two obsolete impressions on the disc, and the elytra are rather broader posteriorly; the eyes, which are very small, are altogether placed behind, as in *P. britannica* (Matth.).

Provence.

L'Abeille, xxii. 1884.

"Coléoptères de l'Ancien Monde publiées isolément, ou en langues étrangères."—Ptilidæ, pp. 60-64.

The following descriptions are copied from 'L'Abeille,' as I have never seen an authentic example of any of them, and have no means of consulting the original works:—

P. 60. **Hydroscapha granulum.**

Motschulsky, Étud. iv. 1855, p. 84.

"Nearly twice smaller than *Limnebius atomus*, and much more strongly attenuated behind; black, with the prothorax and elytra reddish-brown or flavescens; antennæ, legs, and (poitrine) pale. Hab. Smyrna."

Col. Motschulsky appears to have described this insect as a species of *Limnebius*, but, if it belongs to *Hydroscapha*, it is probably an immature form of *H. vrotchii*.

P. 60. **Nossidium scaphidiforme.**

Hoeh. Bull. Mosc. 1872, ii. p. 215.

"L. c. 0.5 mm.—Short, oval, rather convex; pitchy black, shining, slightly pubescent, very closely punctured; elytra abbreviate, subtruncate; apex of elytra, antennæ, and legs testaceous. Rather shorter and more convex than *N. pilosellum*, but as broad, rather more attenuated posteriorly, similar in punctuation, the pubescence thinner; prothorax as broad at the base as the elytra; elytra not covering the pygidium, slightly rounded at the apex, apical margin with the antennæ and legs flavescens. Hab. S. Russia, Kiew."

P. 61. *Ptenidium laticolle*.

Hoch. Bull. Mosc. 1872, ii. p. 214.

"L. c. 0.6 mm.—Allied to *P. fuscicorne*, but rather larger, and easily distinguished by the thorax strongly rounded at the sides behind, and distinctly broader than the elytra; very shining, black, and smooth: antennæ and legs entirely pale yellow; sometimes the apex of the elytra is also pale. The greatest width of the thorax is rather behind the middle, and is there one quarter broader than the elytra at the base, and also broader than they are at the middle; elytra suboval, widest before the middle. The foveolate punctures peculiar to this genus are sometimes distinct, sometimes indistinct, sometimes two in number, sometimes four, sometimes altogether wanting. Hab. The South of Russia, Kiev."

P. 61. *Ptenidium longicorne*.

Füss. Berl. Ent. Zeit. 1868, p. 357.

"L. c. 0.8 mm.—Black, shining, in shape much like *P. apicale*, but distinctly larger, elytra widest behind the shoulders, pubescence very short and scattered: antennæ longer than the head and thorax, reddish brown, the last joint furnished with very long grey hairs: head smooth: prothorax distinctly narrower than the elytra, with the anterior angles rounded, posterior very obtuse, sides faintly rounded at the middle, above convex, very faintly and sparingly punctured, with a distinct impression on each side at the posterior angles, and two small indistinct foveæ above the scutellum; scutellum large, triangular; elytra widest before the middle, gradually narrowed towards the apex, which is transparent brown and separately rounded, and ornamented with small punctures, which are more scattered than in *P. apicale*, and disposed in striæ: suture slightly elevated posteriorly; legs brownish yellow. Hab. Germany, Ahrweiler."

P. 62. *Ptenidium cæcum*.

Joseph, Berl. Ent. Zeit. 1882, p. 46.

"L. c. 0.6 mm.—Allied to *P. larigatum* (Gillm.). Without eyes, apterous, transparent yellow; antennæ half as long as the body; prothorax slightly broader than long; elytra flattened, separately rounded behind. Hab. Carniola, from the Grotte de Luëg."

P. 63. *Ptinella biimpressa*.

Reitter, Deuts. Ent. Zeit. 1878, p. 48.

"L. c. 0.8 mm.—Elongate, subparallel, depressed, pale yellow, rather thickly clothed with short pale hair; head large, obtuse in front: prothorax transverse, rather broader than long, rather longer but scarcely broader than the head, sides slightly dilated, rounded in front, contracted towards the base, posterior angles obtuse, disc very obsoletely alutaceous (*rapene*), flattened and very faintly marked with two impressions; elytra rather long, subparallel; legs thickly and very faintly asperate: abdomen broad, and obtusely rounded at the apex. Hab. N. Hungary, Csap."

"This species belongs to the group with the abdomen obtusely rounded at the apex, as *P. Britannica*, *P. testacea*, *P. denticollis*, and *P. Maria*. It may be distinguished by the obtuse angles and flattened disc of the thorax with two longitudinal impressions more or less distinct, but always visible: also by its long and almost parallel elytra, and short obtuse abdomen."

In the foregoing description all the characters agree entirely with *P. Britannica*, except that Herr Reitter does not mention the remarkable prolongation of the head, and the backward position of the eyes.

I may add here that in the collection of M. Jekel there are two specimens labelled "*N. scaphidiforme*," which are not in any way distinguishable from *N. pilosellum*.

LIST OF SPECIES.

THE following list contains the names of all the species with which I am acquainted, arranged according to their apparent affinity to each other.

Those which are marked * are described in the present volume. The species which I have not seen are omitted from this list, and mentioned only in the preface.

The following abbreviations indicate the region inhabited by each species, viz. :—

E.....	Europe.
As.	Asia.
Af.	Africa.
At.	Atlantic Islands.
Aust.	Australasia.
Am.	America, North.
Am. c.....	America, Central, and West Indian Islands.
Am. m.	America, South.
P.....	Polynesia.

Fam. HYDROSCAPHIDÆ.

GEN. *Hydroscapha*, Le Conte.

*natans, Le Conte	Am.
*Crotchii, Sharp	E.

Fam. TRICHOPTERYGIDÆ.

Tribe I. *LIMULODINA*.

GEN. I. *Limulodes*, Matth.

paradoxus, Matth.	Am.
-------------------	-----

Tribe II. *TRICHOPTERYGINA*.

{GEN. II. *Ptinella*, Mots.

*Championnæ, Matth.	Am. c.
*Lewisiæ, Matth.	As.
testacea, Heer	E., At.
*Fauvelii, Matth.	Aust.
Maria, Matth.	E.
denticollis, Fairm.	E.
Britannica, Matth.	E.
*Pacifica, Matth.	P.
*Braziliana, Matth.	Am. m.
flavidula, Fairm.	Am. m.
*Matthewsiana, Woll.	At.
aptera, Guérin	E., At.

pini, Le Conte	Am.
angustula, Gilm.	E., At.
quercus, Le Conte	Am.
*pellucida, Matth.	Am. c.
tenella, Erich.	E.
*rotundicollis, Mots.	E.
*incerta, Matth.	Am. c.

GEN. III. *Championella*, Matth.

*nubigena, Matth.	Am. c.
-------------------	--------

GEN. IV. *Pterycodes*, Matth.

*Salvinii, Matth.	Am. c.
-------------------	--------

GEN. V. *Dimorphella*, Matth.

*Reitteri, Matth.	Am. m.
-------------------	--------

GEN. VI. *Ptinellodes*, Matth.

Le Contei, Matth.	Am.
-------------------	-----

GEN. VII. *Pteryx*, Matth.

brunnea, Le Conte.	Am., Am. c.
suturalis, Heer	E.
balteata, Le Conte	Am.
Du Valii, Matth.	Am.

GEN. VIII. Mikado, Matth.

*Japonicus, Matth. As.

GEN. IX. Myrmicotrichis, Mots.

aequatorialis, Mots. Am. c.

GEN. X. Astatopteryx, Perris.

Perrisii, Matth. E.

GEN. XI. Actinopteryx, Matth.

fucicola, Allib. E., Af., At., Am., Am. c.

Australis, Matth. Aust.

GEN. XII. Trichopteryx, Kirby.

Schaumii, Matth. Am.

munda, Mots. E.

*Lewisii, Matth. As.

*Caucasica, Matth. As.

*laevicollis, Matth. Am. c.

*Hellenica, Matth. E.

*soror, Flach As.

*lucida, Matth. Am. c.

Haldemanni, Le Conte Am.

Sara, Matth. E.

Matthewsii, Woll. At.

glabricollis, Matth. Am.

orientalis, Mots. E.

*rufescens, Matth. Am. c.

cursitans, Neitn. As.

Dobrnii, Matth. Am.

*similior, Matth. Am. c.

discolor, Hald. Am. m.

Sallæi, Matth. Am. m.

Crotchii, Matth. At., Am., Am. c.

*occidentalis, Matth. Am. c.

thoracica, Waltl. E.

atomaria, De Geer E., Am., Am. c.

convexiuscula, Mots. E.

brevicornis, Mots. At.

anthracina, Matth. E., At.

*Godmanni, Matth. Am. c.

*grossa, Mots. Am., Am. c.

umbricola, Woll. At.

Ceylonica, Mots. As.

Alliberti, Matth. Am. c., Am. m.

laticollis, Mann. Am.

crassicollis, Matth. Am.

Henrici, Matth. Am.

*fratercula, Matth. E.

Taurica, Mots. E.

Sitkaensis, Mots. Am.

Marsculii, Matth. E.

Chilensis, Fairm. Am. m.

abrupta, Hald. Am.

*angustipennis, Matth. Am. c.

*truncatissima, Matth. Am. m.

grandicollis, Mann.

*minor, Matth. E., Am. c.

lata, Mots. E.

cantiana, Matth. E.

fascicularis, Herbst. Am.

*Lætitia, Matth. E.

Wollastoni, Matth. At.

*cognata, Matth. Am.

attenuata, Gillm. E.

*semititens, Matth. E.

Championis, Matth. E.

*angusta, Matth. E.

Brasiliensis, Matth. Am. m.

*inconspicua, Matth. Aust.

*glauca, Matth. Am. c.

*Darwini, Matth. Am. m.

atrata, Mots. E.

*reticulata, Matth. Am. m.

Josephi, Matth. Am.

*depressa, Matth. Am. c.

diffinis, Matth. Am.

*brevior, Matth. Am. c.

Hornii, Matth. Am.

Lethierryi, Reiche. Af.

*parallelopipeda, Matth. Am.

*xanthocera, Matth. Am.

*Californica, Matth. Am.

*castanea, Matth. Am.

carbonaria, Matth. E.

volans, Mots. As.

sericans, Heer E., At., Am.

brevis, Mots. E., As.

Canariensis, Matth. At.

bovina, Mots. E.

brevipennis, Erich. E.

Kirbii, Matth. E.

longicornis, Mann. E.

*dubitata, Matth. Am. c.

Edithia, Matth. E.

*fuliginosa, Matth. Am. c.

*bidilatata, Matth. Am. c.

*Grenadensis, Matth. Am. c.

*excelsa, Matth. Am. c.

*vicina, Matth. Am., Am. c.

longula, Matth. E.

picicornis, Mann. E.

nigricornis, Mots. At.

Jansonii, Matth. E.

parallela, Mots. Am.

Wenkeri, Matth. Am. m.

Montandoni, Allib. E., Am.

rivularis, Allib. E.

Guerinii, Allib. E.

*nigrita, Matth. Am. c.

obscæna, Woll. E., At.

elevata, Mots. E.

acuminata, Mots. E.

ovatula, Mots. At.

fuscula, Matth. E.

Waterhousii, Matth. E.

Chevrolati, Allib.	E.
suffocata, Halid.	E.
*moreus, Matth.	Am.
aspera, Halid.	Am.
tesserula, Matth.	Am.
Poweri, Matth.	E.
ambigua, Matth.	E., Am.
*vitrea, Matth.	Am. c.
Reichii, Matth.	Am. m.
dispar, Matth.	E.
*Japonica, Matth.	As.
variolosa, Muls.	E.
Motschulskii, Matth.	Am. m.
Aubæi, Matth.	Am., Am. c., Am. m.

GEN. XIII. *Smicrus*, Matth.

flicornis, Fairm.	E., Am., Am. c., Am. m.
-------------------	-------------------------

GEN. XIV. *Microptilium*, Matth.

pulchellum, Allib.	E.
--------------------	----

GEN. XV. *Ptiliodes*, Matth.

*Blackburni, Matth.	P.
---------------------	----

GEN. XVI. *Nephanes*, Thoms.

*pubescens, Matth.	Am. c.
lieviusculus, Matth.	Am.
meridionalis, Matth.	Am. c., Am. m.
Titan, Newm.	E., At.
flaviventris, Mots.	Am., Am. c.

Tribe III. *PTILIINA*.

GEN. XVII. *Ptilium*, Erich.

Collani, Mäkl.	Am.
*Smithsii, Matth.	Am. c.
Kunzei, Heer	E.
brevicollis, Matth.	E.
*Hornianum, Matth.	Am., Am. c.
rugulosum, Allib.	E.
Försteri, Matth.	E.
Spencei, Allib.	E.
Sharpi, Matth.	Am.
*obseurum, Matth.	Am.
*Columbianum, Matth.	Am.
*humile, Matth.	Am.
*fissicollis, Reitter	E.
cæsum, Erich.	E.
affine, Erich.	E.
*incognitum, Matth.	E.
exaratum, Allib.	E.
myrmecophilum, Allib.	E.
modestum, Wank.	E.
insigne, Matth.	E.
foveolatum, Allib.	E.

*planum, Matth.	Am. c.
*tropicum, Matth.	Am. c.
*Simsoni, Matth.	Aust.
angulicollis, Fairm.	E.
Halidai, Matth.	E.
Croaticum, Hampe	E.
Caledonicum, Sharp	E.
*Ædipus, Flach	E.
*impressum, Matth.	Am. c.
*rufo-testaceum, Matth.	Am. c.

GEN. XVIII. *Millidium*, Mots.

uriculatum, Aubé	E.
triramosum, Mots.	Am. c.

GEN. XIX. *Actidium*, Matth.

*Fowlerianum, Matth.	Am. c.
*lineare, Matth.	Aust.
coarctatum, Halid.	E., Af.
concolor, Sharp	E.
*politum, Matth.	Am.
*rotundicollis, Casey	Am.
*robustum, Casey	Am.
*granulosum, Casey	Am.
*Crotchianum, Matth.	Am.
Boudierii, Allib.	E.
Sharpianum, Matth.	P.
*attenuatum, Casey	Am.

GEN. XX. *Ptenidium*, Erich.

*strangulatum, Matth.	Am. c.
punctatum, Gyll.	E., At.
*Lawsoni, Matth.	Aust.
*Ulkei, Matth.	Am.
fuscescorne, Erich.	E.
nitidum, Heer	E.
Brisonti, Matth.	E.
*Reitteri, Flach	E.
lævigatum, Gillm.	E.
*foveatum, Matth.	Am. c.
foveicollis, Le Conte	Am.
macrocephalum, Neitn.	As.
Mannerheimii, Matth.	Am.
*concinnum, Matth.	Am. c.
*ignobile, Matth.	Am. c.
*impunctatum, Matth.	Am. c.
*nitens, Matth.	Am. c.
Bruckii, Matth.	At.
pullum, Mann.	Am.
*Caseianum, Matth.	Am.
*attenuatum, Matth.	E.
*Africanum, Matth.	Af.
*orientale, Flach	As.
atomaroides, Mots.	E., At.
Wankowiczii, Matth.	E.
evanesceus, Marsh.	E., At., Am.
Kraatzii, Matth.	E.

formicetorum, Kraatz.	E.
turgidum, Thoms.	E.
obesum, Matth.	Am. c.
*speculifer, Matth.	Am.
Gressneri, Erich.	E.

GEN. XXI. **Camptodium**, Mots.

*adustipenne, Mots.	As.
---------------------	-----

GEN. XXII. **Motschulskium**, Matth.

sinuato-colle, Matth.	Am.
-----------------------	-----

GEN. XXIII. **Micridium**, Mots.

vittatum, Mots.	E.
lineatum, Mots.	Am. c.

GEN. XXIV. Nanosella , Mots.	
fungi, Mots.	Am., Am. c.

GEN. XXV. **Throscidium**, Matth.

Germainii, Matth.	Am. m.
Fairmairii, Matth.	Am. m.
invisibile, Nietn.	At., As., Am. c.

GEN. XXVI. **Euryptilium**, Matth.

Saxonicum, Gillm.	E.
marginatum, Aubé	E.

GEN. XXVII. **Nossidium**, Erich.

pilosellum, Marsh.	E.
Americanum, Mots.	Am.
*posthumum, Matth.	Am.

Mr. Matthews unfortunately did not label the specimens he described as types; he only put the same number on all specimens of the same species. With the exception of the species enumerated below, all are to be found in his collection now in my possession.—ED.

Ptinella

incerta, Matth.	Godman & Salvin.
-----------------	------------------

Pteryx

brunnea, Le Conte	Mus. Brit.
Du Valii, Matth.	Dr. Le Conte.

Trichopteryx

dubitata, Matth.	Mus. Brit.
------------------	------------

Ptilium

impressum, Matth.	Mus. Brit.
rufo-testaceum, Matth.	Mus. Brit.

Actidium

attenuatum, Casey	Casey.
-------------------	--------

Ptenidium

Ulkei, Matth.	Ulke.
nitens, Matth.	Godman & Salvin.
obesum, Matth.	Godman & Salvin.

Camptodium

adustipenne Mots.	G. Lewis.
-------------------	-----------

Nanosella

fungi, Mots.	Godman & Salvin.
--------------	------------------

TRICHOPTERYGIA.

FAM. HYDROSCAPHIDÆ.

GENUS HYDROSCAPHA. PL. XV.

Hydroscapha, Le Conte, Trans. Am. Ent. Soc. v. p. 45, 1874; Matthews, Essay Gen. Hydrosc. 1876.

Corpus elongato-conicum, vel exemplis desiccatis ovatum.

Caput sessile, magnum, clypeo a vertice sutura diviso: oculis sat magnis, rotundatis.

Antennæ (Pl. XV., fig. 7) breves, 8- vel potius 10-articulatæ; 1 magno, longo, recurvato; 2 primo plus quam duplo breviori; 3 secundo breviori; 4-7 tertio similibus, gradatim latioribus; 8 longo, fusiformi, septimo quadruplo longiori, apice acuminato, atque in articulis duobus minutis, sæpe conspicuis, diviso.

Palpi maxillares (Pl. XV., fig. 5, s) robusti, 4-articulati; 1 parvo, apice excavato; 2 primo quadruplo longiori, pyriformi, apice excavato; 3 secundo fere pariter longo, multo crassiori, subovato, apice excavato; 4 minuto, ovali.

Palpi labiales (Pl. XV., fig. 6, m) breves, robusti, 3-articulati; 1 permagno, crasso, subpyriformi, apice excavato; 2 parvo, primo multo angustiori, et quadruplo breviori; 3 minuto aciculari.

Labrum (Pl. XV., fig. 3) corneum, antice rotundatum et profunde emarginatum, appendice mediali parva rotunda atque ad latera setis brevibus instructum.

Mandibule (Pl. XV., fig. 4) corneæ, falciformes, profunde bifidæ aut fere duplices. Stipes robustus, externe stylo brevi valido instructus; intus late extensus. Lamella membranacea tenuissima e basi stipitis usque ad mandibularum apices extenditum.

Maxillæ (Pl. XV., fig. 5) corneæ, unilobatæ, lobo valido cultriformi, acuti uncinato, dentibus sex longis, curvatis, acutissimis, ordinibus duobus dispositis, apices versus armatæ. Lobus hic totus ad basim fortiter detlexus deinde curvatim inflexus est. Stipes corneus, ad apicem profunde excavatus, ad basim dente obtuso externe armatus.

Mentum (Pl. XV., fig. 6, j) corneum, oblongum, lateribus concavis costa valida firmatis, antice triangulariter elongatum.

Labium (Pl. XV., fig. 6, k) corneum, breve, transversum, mento aliquantum latius, lateribus parum rotundatis et costa valida firmatis, margine apicali utrinque ad palpos accipiendos profunde excavata atque processu paraglossali instructa.

Lingua (Pl. XV., fig. 6, l) coriacea, magna, lata, subquadrata, margine anteriori leviter rotundata, indistincte tuberculata, et pilosa.

Pronotum transversum ad basim dilatatum.

Scutellum magnum, triangulare; subscutello magno, lato, longe producto.

Elytra truncata, postice attenuata.

Alæ (Pl. XV., fig. 11) magnæ, latæ, postice ad modum stipitis seu scapi abrupte attenuatæ, antice oblongo-ovaliter expansæ nervis duobus basalibus firmatæ, quorum superior prævalidus, e basi bipartitus, ad mediam alam in stigmatem

lato expansus est; *inferior*, etiam validus, ad mediam alam bipartitus, sinuatim inflectitur et dilatatur; membrana pellucida setis marginalibus prælongis instructa est.

Abdomen segmentis octo compositus, tribus basalibus, seu thoracicis, coriaceis; quinque apicalibus corneis.

Caput *subtus* magnum ore deflexo, parte gulari latâ.

Prothorax *prosterno* (Pl. XV., fig. 2, a) sat parvo; *episternis* minutis; *epimeris* parvis, receptacula coxarum postice cingentibus. Coxarum receptacula ovalia non contingentia.

Pectus mesosterno (Pl. XV., fig. 2, d) sat brevi, late carinato. margine anteriori profunde excavata et utrinque acuminatâ, carinâ latâ, antice sat longe et acute productâ; *episternis* humeralibus modicis; *epimeris* sat magnis ad receptacula coxarum extensis. Coxarum receptacula magna, ovalia, per carinam sat late divisa.

metasterno (Pl. XV., fig. 2, g) magno, subquadrato. ad latera corporis haud extenso, margine posteriori mediâ excavata et utrinque acuminatâ; *episternis* angustis, *metasterno* fere pariter longis; *epimeris* longis, minute triangulariter inflexis, reliquis sub elytra occultis.

Venter valde elongatus, retractilis, segmentis sex corneis compositus, 1-5 fere pariter longis, apicali cæteris longiori, per totum late et profunde conice impresso, apice profunde emarginato. Intra segmentum apicale, vaginâ corneâ inclusum et forsân ad natandum adhibendum, corpusculum oblongum corneum (Pl. XV., fig. 10) existit, ad angulum utrumque externum appendicibus sex laminatis instructum, corneis, recurvatis, ad apices tenuissimis attenuatis, internis prælongis, cæteris gradatim brevioribus atque musculorum serie propriâ agendis. Corpusculi hujusco latera intra abdomen spatio longo sunt extensa ad musculos affligendos, quibus ad modum caudæ piscinæ admoventi possint.

Pedes antici coxis magnis ovalibus; trochanteribus sat magnis, triangularibus; femoribus modicis; tibiis modicis ad apices dilatatis, calcaribus validis externis quatuor, interno singulo armatis; tarsis robustis, 3-articulatis, 1 et 2 brevibus, 3 antecedentibus ambobus duplo longiori, unguibus longis curvatis subtus ad basim validissime dilatatis atque dente magno acuto armatis.

intermedii anticis similes sed longiores et graciliores, coxis longioribus, atque tibiis calcaribus externis octo, interno singulo armatis.

postici (Pl. XV., fig. 8) coxis magnis, fere ad latera corporis extensis, per totas sat late laminatis; trochanteribus elongatis, ad apices oblique excavatis; femoribus simplicibus; tibiis elongatis vix dilatatis, calcaribus validis externis quinque interno singulo armatis, unguibus anticis similibus sed gracilioribus.

Coxæ anteriores modice distantes.

intermediæ per carinam divisæ.

posteriores sat remotæ.

In 1865 my lamented friend Mr. G. R. Crotch first discovered this extraordinary genus in Spain, and in 1873 he was fortunate enough to meet with it again in California. The American insect proved to belong to a species distinct from its European congener, and was described by Dr. Le Conte in 1874 under the name of *Hydroscapa natans*. The European species, *Hydroscapa Crotchii*, was shortly after described by Dr. Sharp in the Ent. Monthly Magazine. Specimens of both species were subsequently sent to me for dissection, and the result of my investigation convinced me that the genus was closely allied to the *Trichopterygide*. To the opinion which I then expressed as to the affinity of *Hydroscapa* to the *Trichopterygide* I still in the main adhere, but, on account of the unlobed maxillæ of *Hydroscapa*, consider that it should be regarded as forming a distinct family.

I do not at all agree with the idea, lately brought forward, that the specific name *Gyrinoides*, Aubé, can be properly applied to *H. Crotchii*. With *Limnebius*,

Hydroscapha has no anatomical connection, and it would seem almost impossible that so accurate and experienced an entomologist as Dr. Anbe could have overlooked the conspicuous maxillary palpi of the insect he was then describing; and, in corroboration of this conjecture, I have received from Dr. Sharp types of *Limnchius Gyrinoides*, taken at the same place and in company with *Hydroscapha*, which display their true generic characters beyond the possibility of a doubt.

In dried specimens, when the long abdomen of *Hydroscapha* has contracted, the resemblance between *L. Gyrinoides* and *H. Crotchii* is so close that they might easily be mistaken for each other; but the most casual glance at the palpi and antennæ is at once sufficient to distinguish them. And, again, if the term *Gyrinoides* be assigned to *H. Crotchii*, by what name can the *Limnchius* be known? On every account, therefore, it seems expedient to retain the name *Crotchii* for the species described by Dr. Sharp.

The elongate form and the number of the corneous segments of the abdomen in *Hydroscapha* indicate an affinity to the *Staphylinidæ*, and thus that genus presents a connecting link between the *Staphylinidæ* and the *Trichopterygidæ*, and such appears to be its proper place in any systematic arrangement.

Hydroscapha natans.

H. natans, Le Conte, Trans. Am. Ent. Soc. v. p. 46, 1874; Matthews, Essay Gen. Hydros. p. 16, 1876.

L. c. abd. retracto, $\frac{8}{16}$ l. = 1 mm.; abd. extenso, $\frac{14}{16}$ l. = 1.75 mm.—Conica, vel postice truncata, vel valde elongata, convexa, piceo-brunnea, nitidissima, pilis brevibus albidis sparsissime vestita; capite magno antice rotundato, oculis magnis, rotundatis; pronoto transverso, postice dilatato, distincte sat remote punctato; elytris ovalibus, apicibus abrupte truncatis, ordinibus transversis sinuatis leviter asperatis; abdomine elongato, ad apicem valde attenuato, segmentis quinque apertis; pedibus atque antennis pallide flavis, his ad apices infuscatis.

Body ovate, either truncate or elongate posteriorly.

Head large and broad, alutaceous, shining, remotely punctured, with the clypeus rounded and divided from the vertex by an arcuate suture; eyes large and rounded; antennæ short, pale yellow, with the club dusky.

Thorax short, transverse, rather longer and much broader than the head, widest at the base, shining, indistinctly alutaceous, remotely punctured, sides slightly margined and rounded, anterior angles produced, basal margin faintly sinuated, with the angles slightly acute.

Scutellum large, triangular.

Elytra very convex, longer and rather broader than the head and thorax, widest before the middle, faintly asperate in transverse rows, interstices alutaceous, suture elevated, sides slightly rounded, extremities abruptly truncate.

Abdomen convex, long and much attenuated posteriorly, nigro-piceous, alutaceous, with five segments exposed, shining and remotely punctured, basal segment densely pubescent, the apical deeply emarginate at the apex, and longitudinally impressed throughout its whole length.

Legs rather long, testaceous.

Under parts nigro-piceous, mouth yellow.

Differs from H. Crotchii in its larger size, broader form, sculpture of each part of the body, shorter antennæ, and nigro-piceous colour.

Habitat, North America. Found in a river near Los Angeles, in California, by Mr. G. R. Crotch.

Hydroscapha Crotchii.

H. Crotchii, Sharp. Ent. Mo. Mag. xi. p. 103, 1874; Matthews, Essay Gen. Hydros. p. 17, plate, 1876.

L. c. abd. retracto, $\frac{4}{15}$ lin. = 0.50 mm.; abd. extenso, $\frac{6}{16}$ lin. = 0.75 mm.—Conica, vel postice truncata, vel valde elongata, convexa, nitidissima, livido-brunnea, pilis brevissimis pallidis sparsissime vestita; capite magno, lato, antice rotundato, punctorum serie transversa impresso; pronoto brevi transverso, postice dilatato, angulis sat acutis, leviter remoteque punctato; elytris longioribus, ovalibus ad apices abrupte truncatis, ordinibus transversis sinuatis leviter asperatis; abdomine valde attenuato, segmentis quinque partim apertis; pedibus atque antennis pallidis, his ad apices infuscatis.

Body conic, either truncate or elongate posteriorly, convex, very shining, livid brown, sparingly clothed with very short pale hair.

Head very large and broad, alutaceous, shining, with a row of four punctures on each side near the eye, clypeus divided from the vertex by a suture and row of punctures. Eyes large, rounded. Antennæ pale yellow, dusky towards the apex.

Thorax rather short, transverse, broader but scarcely longer than the head, dilated at the base, alutaceous, remotely punctured, sides slightly rounded, basal margin sinuated, with the angles acute.

Scutellum moderate, acutely triangular.

Elytra rather long, very convex, rather broader and more than one-half longer than the head and thorax, widest before the middle, rather deeply asperate in transverse rows, suture not elevated, extremities broad and truncate.

Abdomen elongate, much attenuated, convex, with five segments exposed, basal segment alutaceous and densely pubescent, black, the other segments dusky castaneous, shining, remotely punctured, apical segment long, with a divergent longitudinal impression on each side near the middle, apical ventral segment deeply emarginate at the apex, with a longitudinal impression in the middle extending to the base.

Legs moderate, pale yellow.

Under parts nigro-piceous, with the mouth, coxæ, and epipleural folds of the elytra pale.

Differs from *H. natans* in the smaller size, narrower and more cylindric form, longer antennæ, depressed suture, and sculpture of every part.

Habitat, Europe. Found in the river Manzanares, in Spain, by Mr. G. R. Crotch.

FAM. TRICHOPTERYGIDÆ.

TRIBE Trichopterygina.

GEN. II.—PTINELLA.

Some entomologists have lately asserted that in *Ptinella* both sexes may possess wings: I cannot of course contradict this assertion without examination of the supposed winged males, but, until convinced to the contrary by ocular demonstration, I shall still adhere to my previously expressed opinion—that serviceable wings are confined to the females alone. Indeed, the law of what is called Nature strongly supports this idea, for, as it is well known in the history of Ants and their allies, the power of flight in the females is the main factor in the preservation and diffusion of the species. I have for many years enjoyed unusual facilities for investigating the habits and anatomy of insects of this genus; more than two thousand specimens have passed under my examination, and I have always found that the presence or absence of wings formed an unmistakable indication of the sex of the individual. I mean by “wings” not mere rudimentary apologies for those members, but fully developed serviceable wings, and I have many anatomical preparations which clearly prove that winged specimens are always females; in some of these the ova are seen fully formed, and ready for extrusion. I may go even farther than this, and say that in specimens in which the wings, although present, are more or less imperfect, the ovarium is in a similar condition, but in every such case male organs are wholly absent. I am fully aware that in some undoubted males minute appendages may be discovered arising from that part of the mesothorax on which the wings articulate, but these can by no stretch of imagination be regarded as wings.

Many interesting species have recently been added to this genus, and one long ago recorded by Col. Motschulsky from Finland, and described by that author under the name of *P. rotundicollis*, has again been found in its original locality: this is the smallest and most graceful of all these elegant insects.

In a note by Mr. Matthews he states that the genus *Ptinella* is furnished with fully developed eyes. He has carefully examined the species—*aptera*, *angustula*, and *denticollis* are particularly mentioned. He says that the only peculiarity about them is that they are concolorous with the other parts of the head. They are mainly situated on the lower surface, being scarcely visible from above.—ED.

SPECIERUM RECENTIORUM EPITOMA.

L. c. 1·12 mm.—Elongato-ovalis, depressa, flava, pronoto basim versus abrupte constricto, capite breviori, elytris brevibus, abdomine obtuso, ped. atque ant. modicis flavis	Championana.
L. c. 0·87 mm.—Elongata, convexa, obscure castanea, pronoto basim versus modice constricto, elytris sat longis, abdomine obtuso, ped. atque ant. longis, gracilibus, lute flavis	Fauvelii.
L. c. 0·80 mm.—Elongato-ovalis, sat depressa, castanea, pronoti lateribus rotundatis, elytris sat longis, ped. atque ant. lute flavis	Pacifica.
L. c. 0·80 mm.—Elongato-ovalis, sat depressa, mas. lute flava, fem. castanea, pronoto ad basim contracto, ped. atque ant. lute flavis	Braziliana.
L. c. 0·75 mm.—Elongato-ovalis, convexa, flava, pronoto brevi, haud constricto, elytris brevibus, ped. atque ant. flavis	Matthewsiana.

- L. c. 0.60 mm.—Late ovalis, sat depressa, mas. late flava, fem. castanea, pronoto parvo, lateribus rotundatis, elytris brevibus, ped. atque ant. robustis, flavis **pellucida.**
- L. c. 0.45 mm.—Elongata, depressa, translucide flava, abdomine obscurato, pronoto perbrevis, lateribus rotundatis, abdomine acuminato, ped. atque ant. pallide flavis **rotundicollis.**
- L. c. 0.85 mm.—Elongata, angusta, convexa, nitida, castanea, pronoto parvo, basim versus fortiter contracto, elytris longis, angustis, abdominis apice flavo, ped. atque ant. flavis **incerta.**
- L. c. 1 mm.—Elongata, depressa, translucide castanea, pronoto magno, haud constricto, elytris brevioribus, ped. atque ant. longis, robustis, late flavis **Lewisiana.**

Ptinella Championana.

P. Championana, Matthews, Trans. Am. Ent. Soc. xi. p. 116, 1884; Biol. Cent.-Amer. Col. ii. pt. i. p. 128, pl. 3, fig. 22, 1888.

L. c. $\frac{9}{16}$ lin. = 1.12 mm.—Elongato-ovalis, depressa, nitida, translucide flava, pilis brevissimis aureis vestita; *capite* magno, antice producto, minute et remote tuberculato; oculis parvis prominentibus; *pronoto* brevi, capite breviori, sed parum latiori, ad medium latissimo, ad basim fortiter et abrupte constricto, tuberculis minutis, remotis, interstitiisque glabris, nitidis, ornato, atque depressione lata in disco plerumque notato, lateribus marginatis, antice valde rotundatis, margine basali fere recta, angulis acutis, salientibus: *elytris* brevibus, capite atque pronoto multum brevioribus, haud latioribus, prope media latissimis, ordinibus transversis, remotis, modice asperatis, interstitiis glabris, nitidis, apicibus rotundatis; abdomine longo, obtuso, segmentis quinque apertis; pedibus atque antennis modicis, flavis.

Body elongate-oval, depressed, shining, translucent yellow, clothed with very short golden hairs.

Head large, produced in front, minutely and remotely tuberculate; eyes small and prominent; antennæ moderate, yellow.

Thorax short, shorter but rather broader than the head, widest at the middle, strongly and abruptly constricted at the base, ornamented with minute remote tubercles, interstices smooth and shining, generally with a broad depression on the disc, sides margined and much rounded anteriorly, basal margin nearly straight, with the angles acute and salient.

Scutellum large, triangular, and deeply asperated.

Elytra short, much shorter and not broader than head and thorax, widest near the middle, moderately asperated in remote transverse rows, interstices smooth and shining, sides moderately rounded, extremities rounded.

Abdomen long, obtuse, with five segments exposed.

Legs moderate, yellow.

Under parts entirely testaceous.

Differs from *P. testacea* in its short and abruptly constricted thorax, shorter legs and antennæ, and very short elytra.

Habitat, Central America. Found in El Reposo, Guatemala, by Mr. Champion.

Ptinella Lewisiana.

P. Lewisiana, Matthews, Cist. Ent. iii. p. 78, 1884.

L. c. $\frac{8}{16}$ lin. = 1 mm.—Elongata, sat depressa, nitidula, translucide castanea, pilis griseis parce vestita; *capite* magno, lato, antice valde rotundato; oculis parvis, prominulis; *pronoto* magno, capite latiori, haud longiori, ad medium latissimo,

leviter sed confertim tuberculato sive alutaceo, impressione levi, ovali, in medio disco utrinque notato, lateribus marginatis, antice valde rotundatis, postice contractis vix constrictis, margine basali fere recta, angulis acutis; *elytris* capite atque pronoto parum longioribus et latioribus, pone media latissimis, ordinibus obliquis remote sat profunde asperatis, lateribus parum rotundatis, apicibus valde rotundatis; abdomine sat obtuso, dilutiori, segmentis duobus ultimis late flavis; pedibus atque antennis longis, robustis, late flavis.

Body elongate, rather depressed, moderately shining, translucidly castaneous, sparingly clothed with griseous hairs.

Head large and broad, much rounded in front; eyes small and rather prominent; antennæ long and robust, bright yellow.

Thorax large, broader but not longer than the head, widest at the middle, lightly but closely tuberculate or alutaceous, with a faint oval impression on each side on the disc, sides margined, strongly rounded in front and contracted posteriorly, basal margin nearly straight with the angles acute.

Scutellum rather small, triangular, closely asperate.

Elytra rather longer and broader than the head and thorax, widest behind the middle, remotely and rather deeply asperated in oblique rows, sides slightly rounded, extremities much rounded.

Abdomen obtuse, pale, with the two apical segments bright yellow.

Legs long and robust, bright yellow.

Under parts castaneous, with the coxæ, mouth, and all the ventral segments bright yellow.

Differs from *P. Fauvelii* in its larger size, broader and not constricted thorax, shorter elytra, and robust legs and antennæ.

Habitat, Asia. A single specimen found by Mr. Lewis near Yokohama, in Japan.

Ptinella Fauvelii, sp. n.

L. c. $\frac{7}{16}$ lin. = 0.87 mm.—Elongata, convexa, nitida, obscure castanea, pilis sat longis griseis vestita; capite magno, lato, sat brevi, confertim tuberculato; oculis magnis, prominentibus; pronoto modico, capite vix longiori aut latiori, pone medium latissimo, basin versus constricto, convexo, nitido, profunde et confertim tuberculato, lateribus marginatis, antice rotundatis, margine basali fere recta, angulis acutis, prominentibus; *elytris* sat longis, ovalibus, capite atque pronoto sesqui longioribus, parum latioribus, pone media latissimis, ordinibus transversis, irregularibus confertim et rugose asperatis, lateribus marginatis et leviter rotundatis, apicibus latis, rotundatis, dilutioribus, extremis albidis; abdomine sat obtuso, pallide castaneo, segmentis quatuor apertis; pedibus atque antennis longis, gracilibus, late flavis.

Body elongate, convex, shining, dark castaneous, clothed with rather long grey hair.

Head large and broad, rather short, closely tuberculate: eyes large and prominent; antennæ long, slender, bright yellow.

Thorax moderate, scarcely longer or broader than the head, widest behind the middle, constricted towards the base, convex and shining, deeply and closely tuberculate, sides margined and rounded anteriorly, basal margin nearly straight, with the angles acute and prominent.

Scutellum moderate, triangular, deeply asperated.

Elytra rather long, oval, rather broader and one-half longer than the head and thorax, widest behind the middle, closely and rugosely asperated in irregular transverse rows, sides margined and slightly rounded, extremities broad, rounded, and paler, with the edge white.

Abdomen rather obtuse, pale castaneous, with four segments exposed.

Legs long and slender, bright yellow.

Under parts castaneous, with the mouth and terminal segments of the venter yellow.

Differs from others in its large size, longer elytra, and rugose sculpture.

Habitat, Australasia. A single specimen of the female found in New Zealand by Mr. Helms, and sent to me by M. Fauvel.

Ptinella Pacifica.

P. pacifica, Matthews, Cist. Ent. iii. p. 42, pl. ii. fig. 4, 1882.

L. c. $\frac{6}{16}$ — $\frac{7}{16}$ lin. = 0.75–0.87 mm.—Elongato-ovalis, sat depressa, sat nitida, plus minusve obscure castanea, pilis flavis sparse induta; *capite* sat brevi, antice valde rotundato, nitido, confertim reticulato, tuberculis minutis, longe remotis ornato; oculis feminarum parvis, prominentibus; *pronoto* modico, capite pariter longo, paulo latiori, ad medium latissimo, tuberculis minutis, equidistantibus, interstitiisque nitidis, leviter et confertim reticulatis, ornato, lateribus marginatis et valde rotundatis, margine basali fere recta, angulis obtusis; *elytris* sat longis, capite atque pronoto pariter longis et latis, pone media latissimis, ordinibus transversis, irregularibus, profunde asperatis, interstitiis nitidis, leviter reticulatis, apicibus valde rotundatis, extremis dilutioribus; abdomine dilutiori, segmentis quinque apertis, apicali obtuso; pedibus atque antennis longis, robustis, late flavis.

Body elongate-oval, rather depressed, rather shining, castaneous, more or less dark, thinly clothed with yellow hair.

Head rather short, much rounded in front, shining, closely reticulated, ornamented with minute very distant tubercles; eyes in the female small and prominent; antennæ long, robust, bright yellow.

Thorax moderate, not longer but rather broader than the head, widest at the middle, ornamented with minute equidistant tubercles, interstices shining, and finely and closely reticulated, sides margined and much rounded, basal margin nearly straight, with the angles obtuse.

Scutellum moderate, triangular, rather long, deeply asperated.

Elytra rather long, as long and as broad as the head and thorax, widest behind the middle, deeply asperated in irregular transverse rows, with the interstices shining and faintly reticulated, extremities much rounded, and narrowly pale.

Abdomen pale, with five segments exposed, apical segment obtuse.

Legs long, bright yellow.

Under parts castaneous, with the mouth and apex of the venter yellow.

Differs from *P. Matthewsiana* in its larger size, longer thorax, longer and broader elytra, and very different sculpture; from others in its broad thorax with its sides much rounded, and not contracted or constricted towards the base.

Habitat, Polynesia. Found by Mr. Blackburn under the bark of a dead tree near Honolulu, in the Sandwich Islands.

Ptinella Braziliانا.

P. Brasiliana, Matthews, Ann. Mag. Nat. Hist. 6, iii. p. 191, 1889.

L. c. $\frac{6}{16}$ — $\frac{7}{16}$ lin. = 0.75–0.87 mm.—Elongato-ovalis, sat depressa, nitidula, modice punctata, pilis aureis vestita; mascula late flava, feminea castanea, abdomine flavescenti atque alis nigris visis; *capite* magno, antice valde rotundato, minute punctato; oculis (♂) nullis, (♀) modicis, nigris; *pronoto* modico, capite parum longiori et latiori, ante medium latissimo, ad basim leviter contracto, minute sed distincte punctato, interstitiis glabris, in disco profunde impresso, lateribus marginatis, margine basali recta, angulis sat acutis; *elytris* capite atque pronoto parum longioribus et latioribus, ad media latissimis, profunde et confertim asperatis, interstitiis glabris, nitidis, lateribus ovaliter rotundatis et marginatis, apicibus contractis et valde rotundatis; abdomine flavescenti, segmentis quinque expositis, apicali obtuso; pedibus atque antennis longis, late flavis.

Body elongate-oval, rather depressed, moderately shining, moderately punctured, clothed with golden hair, bright yellow in the male, in the female castaneous, with the abdomen flavescent.

Head large, much rounded in front, minutely punctured; eyes (♂) wanting, (♀) moderate, black; antennae long, bright yellow.

Thorax moderate, rather longer and broader than the head, widest before the middle, contracted near the base, minutely but distinctly punctured, with the interstices smooth, deeply impressed on the disk, sides margined, basal margin straight, with the angles rather acute.

Scutellum large, triangular, deeply punctured.

Elytra rather longer and broader than the head and thorax, widest at the middle, deeply and closely asperate, interstices smooth and shining, sides ovally rounded and margined, extremities contracted and much rounded.

Abdomen with five segments exposed, apical obtuse.

Legs long, bright yellow.

Under parts ♂) bright yellow, (♀) castaneous.

Differs from *P. testacea* in its larger size, much broader form, longer legs and antennae, and especially in its punctate, not tuberculate, sculpture.

Habitat, South America. Found in Brazil near Blumenau. I am indebted to Herr Reitter for both sexes of this very distinct species.

Ptinella Matthewsiana.

P. Matthewsiana, Wollaston, Col. St. Helenæ, p. 21, 1877.

L. c. $\frac{6}{16}$ lin. = 0.75 mm.—Elongato-ovalis, convexa, sat nitida, confertim tuberculata, flava, pilis aureis vestita: capite magno, convexo, antice producto, leviter alutaceo; oculis maribus nullis; pronoto transverso, capite latiori, haud longiori, ante medium latissimo, tuberculis modicis, confertissime dispositis atque nonnullis aliis majoribus hic et illic dispersis, ornato, lateribus marginatis, valde rotundatis, haud constrictis, margine basali parum arenata, angulis obtusis; elytris brevibus, capite atque pronoto brevioribus, atque angustioribus, prope media latissimis, sat profunde asperatis, lateribus marginatis, fere rectis, apicibus latis, vix rotundatis; abdomine lato, obtuso, segmentis quatuor apertis; pedibus robustis, læte flavis; antennis sat longis, pallide flavis.

Body elongate-oval, convex, rather shining, closely tuberculate, yellow, clothed with golden hair.

Head large, convex, produced in front, moderately alutaceous; eyes wanting in the males; antennae rather long, pale yellow.

Thorax transverse, broader but not longer than the head, widest before the middle, closely covered with moderate tubercles, with some of a larger size scattered here and there, sides margined and much rounded, not constricted, basal margin slightly arcuate, with the angles obtuse.

Scutellum large, triangular, deeply asperated.

Elytra short, shorter and narrower than the head and thorax, widest near the middle, rather deeply asperated, sides margined and nearly straight, extremities broad, scarcely rounded.

Abdomen broad, obtuse, with four segments exposed.

Legs robust, bright yellow.

Under parts yellow, deeply reticulated.

Differs from others in size, broad thorax with its sides much rounded and not constricted, and in its narrow elytra.

Habitat, the Atlantic Islands. Found in St. Helena by Mr. Wollaston, all the specimens being males.

Ptinella pellucida.

P. pellucida, Matthews, Trans. Am. Ent. Soc. xi. p. 117, 1884; Biol. Cent.-Amer. Col. ii. pt. i. p. 129, 1888.

L. c. $\frac{5}{16}$ lin. = 0.63 mm.—Sat lata, et sat depressa, nitidula, pilis pallidis vestita, maribus late flava, fœmineis castanea; *capite* permagno, antice producto, confertim tuberculato: oculis fœminarum magnis, prominentibus, nigris: *pronoto* parvo, transverso, capite multo breviori, haud latiori, prope medium latissimo, tuberculis sat magnis, ordinibus transversis dispositis, interstitiisque sat profunde reticulatis, ornato, lateribus sat late marginatis, antice valde rotundatis, postice contractis, depressione longitudinali in disco notato, margine basali fere rectâ, angulis rectis; *elytris* brevibus, capite atque pronoto nec longioribus, neque latioribus, ad apices latissimis, ordinibus transversis, fere rectis, profunde asperatis, lateribus rectis et marginatis, apicibus latissimis, valde rotundatis; abdomine sat lato, apicem versus acuminato, segmentis quinque apertis; pedibus atque antennis robustis, sat longis, flavis.

Body rather broad and rather depressed, moderately shining, clothed with pale hair, bright yellow in the males, and castaneous in the females.

Head very large, produced in front, closely tuberculate: eyes in the females large and prominent, black: antennæ rather long, robust, yellow.

Thorax small, transverse, much shorter and not broader than the head, widest near the middle, ornamented with rather large tubercles in transverse rows; interstices rather deeply reticulated, with a longitudinal depression on the disc, sides much rounded anteriorly and contracted towards the base, basal margin nearly straight, with the angles right angles.

Scutellum large, triangular, deeply asperated.

Elytra short, not longer or broader than the head and thorax, widest at the apex, deeply asperated in nearly straight transverse rows, sides straight and margined, extremities very broad and much rounded. In the females the wings are seen as black vittæ.

Abdomen rather broad, obtuse, acuminate at the apex, with five segments exposed.

Legs yellow, rather long and robust.

Under parts yellow.

Differs from *P. flavidula* in its much smaller size, shorter thorax, and deeper sculpture; and from *P. quereus* in its larger size, more depressed form, and much deeper sculpture.

Habitat, Central America. Found in Guatemala by Mr. Champion.

Ptinella rotundicollis.

P. rotundicollis, Motschulsky, Bull. Mosc. xli. 2, p. 185, 1868.

L. c. vix $\frac{1}{6}$ lin. = 0.47 mm.—Elongata, sat depressa, nitidula, confertim sed leviter tuberculata, pilis brevissimis aureis vestita, pallide et translucide flava, abdominis basi castaneâ; *capite* permagno, antice rotundato, confertim tuberculato: oculis maribus nullis; *pronoto* perbrevis, capite breviori, haud latiori, ad medium latissimo, confertim et leviter tuberculato, sive alutaceo, lateribus valde rotundatis et leviter marginatis, margine basali fere rectâ, angulis valde obtusis; *elytris* capite atque pronoto parum latioribus, haud longioribus, ad apices latissimis, ordinibus transversis remotis leviter asperatis, interstitiis alutaceis, lateribus marginatis, vix rotundatis, apicibus latis, minime rotundatis; abdomine acuminato, segmentis quinque plus minusve exsertis, leviter asperatis, sive alutaceis; pedibus modicis, pallide flavis: antennis robustis, pallide flavis.

Body elongate, rather depressed and rather shining, closely and finely tuberculate, clothed with very short golden hair, transparent, pale yellow, with the base of the abdomen castaneous.

Head very large, rounded in front, closely tuberculate. eyes wanting in the males; antennæ robust, pale yellow.

Thorax very short, shorter and not broader than the head, widest at the middle, closely or finely tuberculate or alutaceous, sides much rounded and slightly margined, basal margin nearly straight, with the angles very obtuse.

Scutellum very large, triangular, closely and finely asperate.

Elytra rather broader, but not longer than the head and thorax, widest at the extremities, faintly asperated in remote transverse rows, with the interstices alutaceous, sides margined and very slightly rounded, extremities broad and scarcely rounded.

Abdomen acuminate, with five segments more or less exerted and slightly asperated, or alutaceous.

Legs moderate, pale yellow.

Under parts pale yellow, with the base of the venter castaneous.

Differs from others in its exceedingly minute size, large head, short thorax, and also in colour and sculpture.

Habitat, Europe. Found in Finland (Herr Reitter).

This pretty species was described by Col. Motschulsky more than twenty years ago from specimens received from Finland, and has lately been obtained by Herr Reitter in some numbers from the same country. This circumstance corroborates very strongly all that I have so often written as to the accuracy of that author's discrimination of the species of *Trichopterygidæ*, and of the genuineness of those which he has described. While I was engaged in preparing the former part of this work, Col. Motschulsky kindly sent me types of about fifty of his new species, and of these only three could possibly be considered as sexual or accidental varieties.

Ptinella incerta.

P. incerta, Matthews, Trans. Am. Ent. Soc. xi. p. 118, 1884; Biol. Cent.-Amer. Col. ii. pt. i. p. 129, pl. iii. fig. 21, 1888.

L. c. $\frac{7}{16}$ lin. = 0.87 mm.—Elongata, angusta, convexa, obscure castanea, nitida, pilis brevissimis flavis sparse induta; capite sat longo, obtuso, confertim et leviter tuberculato, sive alutaceo; oculis retropositis, sat magnis, prominentibus; pronoto parvo, capite parum latiori, haud longiori, prope medium latissimo, basin versus fortiter contracto, tuberculis sat magnis confertim obtecto, depressionibus duabus ovalibus scutellum versus notato, lateribus marginatis, antice rotundatis, postice contractis, margine basali fere recta, angulis acutis, prominentibus; elytris longis, angustis, subparallelis, capite atque pronoto longioribus, vix latioribus, convexis, nitidis, confertim et profunde asperatis, lateribus fere rectis et marginatis, apicibus concoloribus, valde rotundatis, extremis albidis; abdomine castaneo, segmentis quatuor apertis, apicali toto atque parte penultimi late flavis, apicali acuminato; pedibus atque antennis longis, flavis.

Body elongate, narrow, convex, dark castaneous, shining, sparingly clothed with very short yellow hair.

Head rather long, obtuse, closely and finely tuberculate or alutaceous; eyes placed far back, rather large and prominent; antennæ long, yellow.

Thorax small, rather broader, but not longer than the head, widest near the middle, strongly contracted towards the base, closely covered with rather large tubercles, and marked with two oval depressions near the scutellum, sides margined, slightly rounded anteriorly, and contracted posteriorly, basal margin nearly straight, with the angles acute and prominent.

Scutellum small, elongate triangular, deeply asperated.

Elytra long and narrow, subparallel, longer but scarcely broader than the head and thorax, convex, shining, closely and deeply asperated, sides nearly straight and margined, extremities concolorous and much rounded, with the edge white.

Abdomen with four segments exposed, castaneous with the apical segment and part of the penultimate bright yellow, apex acuminate.

Legs long, yellow.

Under parts castaneous, mouth, coxæ, and greater part of the venter bright yellow.

Differs from all other species in its narrow, convex, and subparallel form, strongly constricted thorax, dark castaneous colour, and deep sculpture.

Habitat, Central America. One specimen found near San Geronimo, Guatemala, by Mr. Champion.

GEN. III.—CHAMPIONELLA. PL. IX.

Championella, Matthews, Trans. Am. Ent. Soc. xi. p. 118, 1884; Biol. Cent.-Amer. Col. ii. pt. i. p. 130, 1888.

Corpus sat elongatum, convexum, capite atque pronoto angustis, elytris latioribus, brevibus, abrupte truncatis.

Caput prominens, sat magnum, parte gulari magnâ, convexâ; oculi parvi, retropositi; antennæ juxta oculos insertæ.

Antennæ (Pl. IX., fig. 7) 11-articulatæ; 1 magno, elongato-ovali, apice truncato; 2 primo majori, alioqui fere simili; 3–7 tenuibus, elongato-ovatis, sibiipsis paribus; 8 septimo parum majori; 9 octavo majori, leviter incrassato; 10 nono multum majori, ad basim rotundato, antice in collo brevi producto; 11 ovato, obtuso, præcedenti multum majori; tribus ultimis clavam moniliformem formantibus atque setis brevibus verticillatis instructis.

Palpi maxillares (Pl. IX., fig. 5, s) sat robusti, 4-articulati; 1 longo, gracili; 2 recurvato, pyriformi, incrassato; 3 magno, subovali, apice abrupte truncato; 4 longo, aciculato, atque sinuato.

Palpi labiales (Pl. IX., fig. 6, m) modici, 3-articulati; 1 robusto, suboblongo; 2 primo pariter longo, sed angustiori, antice parum dilatato, seta longâ, divergentis, ad apicem externe instructo; 3 cæteris minori, ovato.

Labrum (Pl. IX., fig. 3) magnum, semiorale, ante medium constrictum atque setis tribus longis instructum, antice rotundatum, in disco setarum ordine transverso, curvato ornatum.

Mandibule (Pl. IX., fig. 4) validæ, acie concavata, integrâ, dorso convexo, deplanato, minute costato. Lamella membranacea sat magna, rotundata. Stipes robustus in stylo valido, incurvato, acuto, externe productus, atque dente longo basali armatus.

Maxille (Pl. IX., fig. 5) validæ, quasi trilobatæ, lobo *exteriori* longo, leviter incurvato; *intermedio* sat brevi, articulo basali parvo, apicali multum longiori, setis longis ad apicem dense instructo; *interiori* valido, sat brevi, dentibus tribus validis, incurvatis, acutissimis ad apicem armato. Stipes latus, robustus.

Mentum (Pl. IX., fig. 6, j) oblongum, costis lateralibus firmatum atque setis nonnullis brevibus ad angulos anteriores instructum.

Labium (Pl. IX., fig. 6, k) oblongum, sub mento ferme occultum, processibus paraglossalibus rudimentariis.

Lingua (Pl. IX., fig. 6, l) modica, subconica, ad apicem truncata et quadridentata.

Pronotum modicum, transversum, lateribus rotundatis.

Scutellum modicum, triangulare.

Elytra brevissima, abrupte truncata, epipleuris latis, distinctis.

Alæ haud examinatæ.

Abdomen elongatum, segmentis quatuor, interdum quinque, corneis expositis, fere pariter longis, apicali integro, sat acuto; alioqui non examinatum.

Prothorax (Pl. IX., fig. 2, a) *prosterno* sat magno; *episternis* minutis; *epimeris* magnis, receptacula coxarum postice cingentibus; *coxarum receptaculis* rotundatis, fere contingentibus.

Pectus mesosterno (Pl. IX., fig. 2, d) magno, haud carinato, *episternis* magnis, suboblongis; *epimeris* permagnis, ad receptacula coxarum late extensis; *receptaculis coxarum* sat magnis, transverse ovalibus, fere contingentibus.

metasterno (Pl. IX., fig. 2, g) sat magno, transverse oblongo, ad latera corporis extenso, margine posteriori inter coxas recta; *episternis* totis sub elytris occultis; *epimeris* postice minute triangulariter inflexis.

Venter segmentis septem compositus, basali majori, apicali integro, acuminato.

Pedes antici graciles, trochanteribus parvis, trigonalibus; femoribus simplicibus; tibiis longis, gracilibus; tarsis brevibus, triarticulatis, 1 modico; 2 primo multum minori; 3 valde elongato, tenui, unguibus gracilibus, leviter curvatis.

intermedii anticis similes, tibiis apicem versus leviter dilatatis.

postici (Pl. IX., fig. 8) ceteris parum longioribus, trochanteribus modicis, claviformibus; femoribus sat robustis, simplicibus; tibiis apicem versus leviter dilatatis, margine interiori remote serrata; tarsis anticis similibus.

Coxae anteriores rotundatae, fere contingentēs.

intermediae ovales, fere contingentēs.

posteriores parvae, prominentes, haud laminatae, longe distantes.

Body rather elongate, convex, with the head and thorax narrow, elytra broader, short and abruptly truncate

Head prominent, rather large, gular portion large, convex; eyes small, placed far back; antennae inserted near the eyes.

Antennae (Pl. IX., fig. 7) eleven-jointed; 1 large, elongate-oval, truncate at the apex; 2 larger than the first, but almost similar; 3-7 slender, elongate-ovate, equal to each other; 8 rather larger than the seventh; 9 larger than the eighth, slightly incrassate; 10 much larger than nine, much incrassate and rounded at the base, and produced into a short neck at the apex; 11 ovate, obtuse, much larger than the preceding; the three last form a moniliform club, and are ornamented with short verticillate setae.

Maxillary palpi (Pl. IX., fig. 5, s) rather robust, four-jointed; 1 long and slender; 2 recurved, pyriform, incrassate; 3 large, suboval, abruptly truncate at the apex; 4 long, aciculate and sinuated.

Labial palpi (Pl. IX., fig. 6, m) moderate, three-jointed; 1 robust, suboblong; 2 equal in length to the first, but narrower, slightly widest in front, and furnished with a long divergent seta on the exterior apical angle; 3 ovate, much smaller than the preceding.

Labrum (Pl. IX., fig. 3) large, semi-oval, constricted before the middle and furnished with three long setae on each side, rounded anteriorly and ornamented with a curved row of short setae on the disc.

Mandibles (Pl. IX., fig. 4) strong, acute but not much hooked at the apex, edge concave and simple; convex and flat at back with small transverse costae. Lamella membranacea rather large, rounded. Stipes robust, produced into a strong, incurved and acute style at the apex, and armed with a long denticulation near the base.

Maxillae (Pl. IX., fig. 5) strong, apparently trilobed, with the *exterior* lobe long and slightly incurved; *intermediate* rather short with the basal joint small, the apical much longer, and densely furnished with long setae at the apex; *interior* robust, rather short, and armed at the apex with three strong and very sharp incurved teeth. Stipes broad and robust.

Mentum (Pl. IX., fig. 6, j) oblong, strengthened with lateral costa, and furnished with short setae at the anterior angles.

Labium (Pl. IX., fig. 6, k) oblong, almost concealed by the mentum, paraglossal processes rudimentary.

Lingua (Pl. IX., fig. 6, l) moderate, subconical, truncate and quadridentate at the apex.

Pronotum moderate, transverse, and rounded at the sides.

Elytra short, abruptly truncate, with the epipleura distinct and broad.

Wings not examined.

Abdomen elongate, with either four or five segments exposed and equal in length, apical segment entire and pointed.

Prothorax (Pl. IX., fig. 2, a) with the *prosternum* rather large; *episterna* minute; *epimera* large, enclosing the coxal cavities behind; *coxal cavities* rounded, and nearly contingent.

Mesosternum (Pl. IX., fig. 2, d) large, not carinated; *episterna* large, suboblong; *epimera* very large, extending broadly to the coxal cavities; *coxal cavities* rather large, transversely oval, nearly contingent.

Metasternum (Pl. IX., fig. 2, g) rather large, transversely oblong, extending to the sides of the body, with the posterior margin between the coxæ straight; *episterna* entirely covered by the elytra; *epimera* minutely and triangularly inflexed beyond the metasternum.

Venter composed of seven segments, of which the basal is rather the longest, and the apical entire and rather pointed,

Legs anterior slender, trochanters small, trigonal; femora simple; tibiæ long and slender; tarsi three-jointed; 1 moderate; 2 much smaller than the first; 3 very long and slender; claws feeble, slightly curved.

intermediate like the anterior, but with the tibiæ slightly dilated towards the apex.

posterior (Pl. IX., fig. 8) rather longer than the preceding, trochanters moderate, claviform; femora simple, rather robust; tibiæ slightly dilated towards the apex with the inner margin remotely serrated; tarsi like the anterior.

Coxæ anterior rounded, nearly contingent.

intermediate oval, nearly contingent.

posterior small, prominent, not laminated, and widely distant.

In the shortness of its elytra and length of its abdomen this genus resembles *Ptinella*, but in other respects is totally different. Its principal diagnostic characters consist in the hardness of its integument, seven ventral segments, and the absence of laminae on the posterior coxæ.

Championella nubigena.

C. nubigena, Matthews, Trans. Am. Ent. Soc. xi. p. 119, 1884; Biol. Cent.-Amer. Col. ii. pt. i. p. 130, pl. iii. fig. 24, 1888.

L. c. $\frac{5}{16}$ — $\frac{6}{16}$ lin. = 0.63–0.75 mm.—Elongato-ovata, valde convexa, nitida, obscure castanea, pilis argenteis, parce vestita; *capite* prominenti, sat longo; oculis parvis, haud prominentibus; *pronoto* capite paulum longiori et latiori, prope medium latissimo, nitido, leviter et confertissime tuberculato, sive alutaceo, lateribus rotundatis, leviter marginatis, margine basali fere recta, angulis obtusis; *elytris* brevibus, capite atque pronoto brevioribus, sed paulum latioribus, pone media latissimis, sat profunde et remote asperatis, interstitiis fortiter reticulatis, lateribus modice rotundatis, apicibus latis, fere recta truncatis; *abdomine* conico, sat acuto, segmentis quatuor totis apertis, apicem versus dilutiori; pedibus sat longis, læte flavis, femoribus posticis valde robustis. antennis modicis, obscure flavis.

Body elongate-ovate, very convex, shining, dusky castaneous, sparingly clothed with silver hair.

Head prominent, rather long; eyes small, not prominent; antennæ moderate, dusky yellow.

Thorax rather longer and broader than the head, widest near the middle, shining, finely and very closely tuberculate or alutaceous, sides rounded and slightly margined, basal margin nearly straight, with the angles obtuse.

Scutellum moderate, triangular, remotely asperate, with the interstices strongly reticulated.

Elytra short, shorter but rather broader than the head and thorax, widest beyond the middle, rather deeply and remotely asperate, with the interstices strongly reticulated, sides moderately rounded, extremities broad and nearly straight.

Abdomen conical, with four segments fully exposed, paler towards the apex.

Legs rather long, bright yellow, posterior femora very robust.

Under parts castaneous, with the mouth, coxæ, and two apical segments of the venter yellow.

Habitat. Central America. Found by Mr. Champion on Totonicapam, in Guatemala, at an elevation of 10,500 ft.

This species may be easily recognized by its dark colour, corneous integument, and abrupt truncation of the elytra.

GEN. IV.—PTERYCODES. PL. X.

Pterycodes, Matthews, Trans. Am. Ent. Soc. xi. p. 120, 1884; Biol. Cent.-Amer. Col. ii. pt. i. p. 131, 1888.

Corpus ovatum, valde convexum.

Caput magnum, prominens, parte gulari magnâ, valde convexâ; oculi modici; antennæ sat longi ante oculos insertæ.

Antennæ (Pl. X., fig. 7) longæ, graciles, 11-articulatæ: 1 permagno; ovali, apice truncato; 2 primo minori, antice contracto, apice abrupte truncato; 3-8 elongatis, gracillimis, bases atque apices versus contractis, setis brevissimis verticillatis ornatis; 9 precedenti formâ simili sed majori; 10 nono multum majori, ad basim valde incrassato, antice in collo sat brevi producto; 11 decimo majori, et magis incrassato, obtuse subconico; tribus ultimis clavam gracilem, moniliformem formantibus, atque setis longioribus verticillatis instructis.

Palpi maxillares (Pl. X., fig. 5, s) sat breves, 1-articulati: 1 sat longo, gracili; 2 modico, pyriformi; 3 ovali, modice incrassato, apice extremo truncato; 4 sat longo, aciculato et sinuato, setâ divergenti ante medium utrinque instructo.

Palpi labiales (Pl. X., fig. 6, m) magni, valde elongati, triarticulati; 1 longo, cylindrico; 2 cylindrico, primo multum breviori atque tenuiori; 3 precedenti multum minori, acute aciculato.

Labrum (Pl. X., fig. 3) magnum, semiovale, ante medium constrictum, setis tribus longis ad latus utrinque atque ordine curvato setarum breviarum in disco instructum.

Mandibulæ (Pl. X., fig. 4, t) validæ, acute uncinnatæ, acie simplici, concavata, dorso convexo, deplanato, costis parvis, transversis armato. Lamella membranacea magna, rotundata. Stipes robustus, in stylo longo, incurvato, acuto, externe productus, atque dente basali sat parvo armatus.

Marillæ (Pl. X., fig. 5) modicæ, quasi trilobatæ, lobo exteriori magno, robusto, ad basim dilatato, leviter incurvato; intermedio longo, articulo basali parvo, apicali elongato, setis longis, erectis ad apicem dense instructo; interiori modico, longe infra precedentem posito, dentibus aut quinque aut sex longis, acutissimis, curvatis ad apicem armato. Stipes sat robustus, setis duabus prælongis ad medium externe instructus.

Mentum (Pl. X., fig. 6, j) oblongum, costis lateralibus firmatum, atque setis duabus validis ad angulum anteriorem utrinque instructum.

Labium (Pl. X., fig. 6, k) sat magnum, suboblongum, sub mento ferne occultum,

- processu paraglossali brevi, acuto, atque seta divergenti ad apicem ejus instructo, utrinque armatum.
- Lingua* (Pl. X., fig. 6, l) elongata, angusta, ad apicem excavata, atque seta divergenti utrinque instructa.
- Pronotum* transversum, basim versus sat fortiter constrictum.
- Elytra* sat longa, truncata.
- Alæ* lanceolatæ, sub elytris visis, alioqui haud examinatæ.
- Abdomen* sat elongatum, segmentis tribus partim apertis, pariter longis, apicali integro, sat acuminato, reliquis haud examinatis.
- Prothorac* (Pl. X., fig. 2, a) *prosterno* sat magno, inter coxas anguste elevato atque pone eos triangulariter dilatata; *episternis* minutis; *epimeris* modicis, receptacula coxarum cingentibus; *corarum receptaculis* rotundatis.
- Pectus mesosterno* (Pl. X., fig. 2, d) magno, carinato; *episternis* elongatis, humeralibus; *epimeris* magnis, ad coxarum receptacula late extensis; *receptaculis corarum* rotundatis, per carinam divisis.
- metasterno* (Pl. X., fig. 2, g) transverse oblongo, a corporis lateribus sat late remoto, margine posteriori inter coxas fere recta; *episternis* latis, sat brevibus; *epimeris* magnis, postice sat late inflexis.
- Venter* segmentis sex compositus, basali atque apicali cæteris longioribus, apicali integro, conico.
- Pedes antici* coxis rotundatis; trochanteribus parvis, trigonalibus; femoribus longis, simplicibus; tibiis longis, gracilibus; tarsis triarticulatis: 1 longo; 2 primo duplo breviori; 3 elongato, gracillimo; unguibus gracilibus, leviter curvatis.
- intermedii* anticis longiores, alioqui similes.
- postici* (Pl. X., fig. 8) cæteris parum longiores, trochanteribus modicis, clavi-formibus; femoribus robustis, simplicibus; tibiis apices versus leviter dilatatis atque interne calcaribus brevibus armatis; tarsis anticis similibus.
- Coxæ anteriores* ovales, fere contingentes.
- intermediæ* rotundatæ, per carinam divisæ.
- posteriores* prominentes, sat angustæ, late triangulariter laminatæ, longe distantes.

Body ovate, very convex.

Head large and prominent; gular portion large and very convex; eyes moderate; antennæ inserted considerably in front of the eyes.

Antennæ (Pl. X., fig. 7) long and slender, 11-jointed; 1 very large, oval, with the apex truncate; 2 smaller than the first, contracted anteriorly, with the apex abruptly truncate; 3-8 elongate and very slender, equal to each other in size, contracted both towards the base and apex, ornamented near the middle with short verticillated setæ; 9 like the preceding in form, but larger; 10 much larger than the ninth, much incrassated towards the base, and produced into a short neck at the apex; 11 larger than the tenth, and more incrassated, obtusely subconical; the last three form a slender moniliform club, and are furnished with longer verticillated setæ.

Maxillary palpi (Pl. X., fig. 5, s) rather short, four-jointed; 1 rather long and slender; 2 moderate, pyriform; 3 oval, moderately incrassated, with extreme apex truncate; 4 rather long, aciculate and sinuated, furnished with a divergent seta on each side before the middle.

Labial palpi (Pl. X., fig. 6, m) large and very long, extending in front of the mouth, three-jointed; 1 long, cylindric; 2 cylindric, much shorter and more slender than first; 3 much smaller than the second, acutely aciculate.

Labrum (Pl. X., fig. 3) large, semi-oval, constricted before the middle and furnished on each side with three long setæ, and on the disc with a curved row of shorter setæ.

Mandibles (Pl. X., fig. 4, t) strong, sharply uncinuate, with the edge concave and simple, and the back convex and flat, armed with small transverse costæ.

- Lamella membranacea large and rounded. Stipes robust, produced anteriorly into a long incurved acute style, and armed at the base with a rather small tooth.
- Maxillæ* (Pl. X., fig. 5) moderate, apparently trilobed, with the *outer lobe* large and robust, dilated at the base, and slightly incurved; *intermediate* long, with the basal joint small, and the apical joint elongate and densely furnished with long erect setæ at the apex; *inner lobe* moderate, placed far below the preceding lobe, and armed at the apex with five or six long and very sharp curved teeth. Stipes rather robust, furnished with two very long setæ near the middle of the outer side.
- Mentum* (Pl. X., fig. 6, j) oblong, strengthened with lateral costæ, and furnished with two strong setæ at each anterior angle.
- Labium* (Pl. X., fig. 6, k) rather large, suboblong, almost hidden by the mentum, armed on each side at the anterior angles with a short acute paraglossal process bearing a divergent seta at its apex.
- Lingua* (Pl. X., fig. 6, l) elongate, narrow, with the apex concave, and furnished on each side with a divergent seta.
- Pronotum* transverse, rather strongly constricted near the base.
- Scutellum* moderate, triangular.
- Elytra* rather long, truncate.
- Wings* lanceolate, seen through the elytra; in other respects not examined.
- Abdomen* rather long, with three segments more or less exposed, equal to each other in length, apical segment entire and rather acuminate, in other respects not examined.
- Prothorax* (Pl. X., fig. 2, a) with the *prosternum* rather large, narrowly elevated between the coxæ, and triangularly dilated beyond them; *episterna* very small; *epimera* moderate, inclosing the coxal cavities; *coxal cavities* rounded.
- Mesosternum* (Pl. X., fig. 2, d) large, carinated; *episterna* elongate, humeral; *epimera* large, broadly extending to the coxal cavities; *coxal cavities* rounded, narrowly separated by the carina.
- Metasternum* (Pl. X., fig. 2, g) transversely oblong, not extending to the sides of the body, with the posterior margin between the coxæ straight; *episterna* broad and rather short; *epimera* large, broadly inflexed posteriorly.
- Venter* composed of six segments, with the basal and the apical segments rather the longest, apical segment conical and entire.
- Legs* long and slender, *anterior* with the coxæ oval, trochanters small, trigonal; femora long and simple; tibiæ long and slender; tarsi three-jointed, with the first joint long; second much shorter; third long and very slender; claws slender and faintly curved.
- intermediate* longer than the anterior, but in other respects similar.
- posterior* (Pl. X., fig. 8) rather longer than the preceding, trochanters moderate, claviform; femora robust and simple; tibiæ slightly dilated towards the apex, and armed with short spurs on the inner side; tarsi like the anterior.
- Coxæ* *anterior* oval, almost contingent.
- intermediate* rounded, separated by the carina.
- posterior* prominent, rather narrow, triangularly and broadly laminated, widely separated.

Pterycodes is allied to *Pteryx* and *Ptinella*, but differs conspicuously from them both in the smaller size of its metasternum, which does not extend to the sides of the body, in the length and form of the labial palpi, and in the diverse formation of its antennæ. From *Pteryx* it is also distinguished by the number of its ventral segments, and the shape of its thorax.

• *Pterycodes Salvini*.

P. Salvini, Matthews, Trans. Am. Ent. Soc. xi. p. 121, 1884; Biol. Cent.-Amer. Col. ii. pt. i. p. 132, 1888.

L. c. $\frac{7}{16}$ — $\frac{8}{15}$ lin.=0.87—1 mm.—Subovatus, valde convexus, nitidus, castaneus, pilis brevissimis aureis parce vestitus: *capite* prominenti, magno, lato; oculis magnis, sat prominentibus; *pronoto* modico, capite paulum latiori haud longiori, prope medium latissimo, basim versus constricto, punctis magnis foveolatis, interstitiisque alutaceis ornato, lateribus antice rotundatis, postice constrictis, latissime marginatis, margine basali leviter sinuata, angulis acutis, prominentibus; *elytris* ovatis, translucidis, alis subtus visis, capite atque pronoto parum longioribus, vix latioribus, prope media latissimis, punctis asperatis, profundis, remotis, ordinibus curvatis dispositis, interstitiis leviter alutaceis, ornatis, lateribus leviter rotundatis et leviter marginatis, apicibus latis valde rotundatis; abdomine conico segmentis tribus apertis, duobus ultimis flavis; pedibus atque antennis longis, gracilibus, lute flavis.

Body subovate, very convex and shining, castaneous, sparingly covered with very short golden hair.

Head prominent, large and broad, produced in front, alutaceous and shining, mouth yellow; eyes large and rather prominent; antennae very long and slender, bright yellow.

Thorax moderate, rather broader but not longer than the head, widest near the middle and constricted near the base, regularly marked with deep and rather remote foveolate punctures, with the interstices alutaceous, sides rounded in front and constricted behind, very broadly margined, basal margin slightly sinuated, with the angles acute and prominent.

Scutellum moderate, triangular, marked with remote foveolate punctures.

Elytra ovate, transparent, rather longer but scarcely broader than the head and thorax, widest near the middle, marked with deep asperate punctures in curved rows, with the interstices alutaceous, sides slightly rounded and slightly margined, extremities broad and much rounded.

Abdomen conical, rather acute, with three segments exposed, the last two yellow.

Legs long and slender, bright yellow.

Under parts castaneous, with the mouth, coxae, and apical segment of the venter bright yellow.

Habitat, Central America. This pretty insect was found on Cerro Zunil, in Guatemala, by Mr. Champion.

GEN. V.—*DIMORPHELLA*. PL. XI.

Dimorphella, Matthews, Ann. Mag. Nat. Hist. 6, iii. 1889, p. 190.

Corpus subovatum, valde convexus

Caput prominens, antice sat productum; parte gulari sat longa, convexa; oculi prominentes.

Antennae (Pl. XI., fig. 7) longae, graeciles, 11-articulatae; 1 magno, robusto; 2 primo fere pariter magno, ovato, apice truncato; 3-8 fere linearibus, sibiipsis paribus; 9 octavo parum longiori, leviter incrassato, ampullaeformi; 10 nono et longiori et crassiori, ampullaeformi; 11 decimo pariter longo, parum crassiori, anguste ovato; tribus ultimis clavam tenuem, elongatam formantibus, 7-11 setis verticillatis instructis.

Palpi maxillares (Pl. XI., fig. 5, s) sat robusti, 4-articulati; 1 longo, gracili; 2 magno, pyriformi; 3 permagno, ovato, apice truncato; 4 sat brevi, aciculari.

Palpi labiales (Pl. XI., fig. 6, m) perbreves, 3-articulati, omnibus fere paribus, transverse subquadratis, apicali setis nonnullis, erectis, longissimis, ad apicem externe instructo.

Labrum (Pl. XI., fig. 3) magnum, latum, transversum, subquadratum, margine anteriori minute denticulata, ad basim parum dilatatum atque setis longis utrinque instructum.

Mandibulae (Pl. XI., fig. 4, 1) magna, sat longa, obtuse uncinatae, dorso convexo, minute costato. Lamella membranacea ovalis. Stipes robustus in stylo magno, fortiter incurvato, externe productus.

Maxillae (Pl. XI., fig. 5) permagna, valde robustae, trilobatae, lobo exteriori robusto, lato, ad apicem acutissimo, ad basim fortiter constricto; intermedio magno, lato, ad apicem longe setoso; interiori magno, lato, ad apicem dentibus multis, longis, acutissimis, curvatis armato.

Mentum (Pl. XI., fig. 6, j) magnum, oblongum, costis lateralibus validis firmatum.

Labium (Pl. XI., fig. 6, k) oblongum, sub mento ferre occultum, in processu paraglossali brevi, acutissimo, divergenti, ad apicem utrinque productum.

Lingua (Pl. XI., fig. 6, l) perbrevis, lata, apice producto, profunde bifido.

Pronotum sat parvum, angustum, capite paulo longius et latius, pone medium latissimum, ad basim fortiter constrictum, in maribus prope mediis lateribus dente acuto, magnitudine variabili, armatum, margine basali recta, angulis salientibus.

Scutellum magnum, triangulare.

Elytra magna, tumide convexa, apicibus subtruncatis.

Alae lanceolatae, angustae, scapo longo, nervis atque setis marginalibus fuscis.

Abdomen segmentis novem compositum, tribus basalibus coriaceis, sex apicalibus plus minusve corneis.

Prothorax (Pl. XI., fig. 2, a) modicus; *prosterno* modico; *episternis* sat parvis; *epimeris* modicis, coxarum receptacula postice cingentibus; *receptaculis coxarum* fere circularibus, haud contingentibus.

Pectus mesosterno (Pl. XI., fig. 2, d) magno, valde et late elevato, atque antice et postice mucronato; *episternis* sat magnis, suboblongis; *epimeris* magnis, humeralibus, ad coxas late extensis; *receptaculis coxarum* magnis, circularibus, per carinam anguste divis.

metasterno (Pl. XI., fig. 2, f) sat magno, a lateribus corporis late remoto, margine posteriori inter coxas recta; *episternis* magnis, latis, ad coxas extensis; *epimeris* longis, postice sat late inflexis, atque ultra coxas productis; coxis prominentibus, sat late distantibus.

Venter segmentis sex compositus, omnibus fere pariter longis, apicali caeteris parum longiori, integro.

Pedes graciles, *antici* trochanteribus minutis, trigonalibus; femoribus simplicibus; tibiis gracilibus, ad apicem dilatatis, margine interiori serrata, calcaribus parvis armatis; tarsis triarticulatis; 1 et 2 perbrevibus; 3 valde elongato, gracillimo; unguibus gracillimis, leviter curvatis.

intermedii anticis similes, sed tibiis longioribus, leviter recurvatis.

postici (Pl. XI., fig. 8) trochanteribus claviformibus; femoribus simplicibus; tibiis margine interiori longe serrata; tarsis anticis similibus.

Coxae anteriores rotundatae, modice distantes.

intermediae rotundatae, per carinam angustam divisae.

posteriores magne, prominentes, per totas late laminatae, sat remotae; lamina ovata, ad mediam leviter acuminata.

Body subovate, very convex.

Head prominent, rather produced in front, gular portion large, very convex; eyes moderate, prominent.

Antennae (Pl. XI., fig. 7) long and slender, 11-jointed; 1 large and robust; 2 nearly as large as the first, ovate, with the apex truncate; 3-8 almost linear, very slender, equal in length; 9 rather longer than the eighth, slightly incrassate, ampulliform; 10 ampulliform, longer and more incrassate

- than the ninth: 11 narrowly ovate, as long, but more incrassate than the tenth; the last three forming an elongate club; 7-11 furnished with verticillate setæ.
- Maxillary palpi* (Pl. XI., fig. 5, s) rather robust, 4-jointed: 1 long and slender; 2 large, pyriform; 3 very large, ovate, truncate at the apex; 4 rather short, acicular.
- Labial palpi* (Pl. XI., fig. 6, m) very short, 3-jointed, all nearly equal in size, subquadrate, transverse, the apical joint furnished at its outer extremity with several very long erect setæ.
- Labrum* (Pl. XI., fig. 3) large and broad, subquadrate, with the anterior margin minutely denticulate, slightly dilated, and furnished with long setæ at the base on each side.
- Mandibles* (Pl. XI., fig. 4, t) large, rather long, and obtusely hooked at the apex, convex, flattened, and minutely costate at the back. Lamella membranacea oval. Stipes robust, and produced above into a large strongly curved style.
- Maxillæ* (Pl. XI., fig. 5) very large and very robust, apparently trilobed; *exterior* lobe robust and broad, very sharply pointed, and strongly constricted near the base; *intermediate* large and broad, densely furnished with long setæ at the apex; *interior* large and broad, armed with many long, very sharp, curved teeth at the apex.
- Mentum* (Pl. XI., fig. 6, j) large, oblong, with broad lateral costæ.
- Labium* (Pl. XI., fig. 6, k) oblong, almost entirely concealed by the mentum, produced on each side of the apex into a short, very acute, divergent paraglossal process.
- Lingua* (Pl. XI., fig. 6, l) very short and broad, produced and deeply bifid at the apex.
- Pronotum* rather small and narrow, rather longer and broader than the head, widest behind the middle, strongly constricted near the base, in the males produced on each side near the middle into an acute tooth variable in size, basal margin straight, with the angles acute and salient.
- Scutellum* large, triangular.
- Elytra* large and turgidly convex in the males, rather short and truncate in the females.
- Wings* lanceolate, narrow, scape long, nerves and marginal setæ brown.
- Abdomen* composed of nine segments, the three basal coriaceous, six apical more or less corneous.
- Prothorax* (Pl. XI., fig. 2, a) moderate; *prosternum* moderate; *episterna* rather small; *epimera* moderate, inclosing posteriorly the coxal cavities; *coxal cavities* almost circular, not contingent.
- Mesosternum* (Pl. XI., fig. 2, d) large, much and broadly elevated, and anteriorly and posteriorly acuminate; *episterna* rather large, suboblong; *epimera* large, humeral, broadly extending to the coxal cavities; *coxal cavities* large, circular, narrowly separated by the carina.
- Metasternum* (Pl. XI., fig. 2, g) rather large, widely remote from the sides of the body, posterior margin between the coxæ straight; *episterna* large and broad, reaching to the coxæ; *epimera* long, broadly inflected posteriorly, and extending beyond the coxæ; coxæ prominent.
- Venter* composed of six segments of nearly equal length, apical segment entire and rather the longest.
- Legs* slender, anterior with the trochanters minute, trigonal; femora simple; tibiæ slender, slightly dilated towards the apex, with the inner margin serrate, armed at the extremity with small spurs; tarsi three-jointed; 1 and 2 very short; 3 very long and very slender; claws slender, slightly curved.
- intermediate* like the anterior, but with the tibiæ rather longer and faintly recurved.
- posterior* (Pl. XI., fig. 8) with the trochanters claviform; femora simple; tibiæ with the inner margin very strongly serrate; tarsi like the anterior.

Coxæ anterior rounded, moderately distant.

intermediate rounded, separated by the narrow extremity of the carina.

posterior large and prominent, rather distant, broadly laminated for their whole length; lamina ovate, slightly acuminate at the middle.

It is not an easy matter to determine the tribe in which this elegant genus should be placed. As its name implies, a vast difference of form exists between the sexes; the males resembling *Ptenidium*, or more closely *Camptodium*, while the females might be readily supposed to belong to *Ptinella*. It is therefore with some hesitation that I have placed it among the *Trichopterygina*; the truncation of the elytra is, although evident, but very slight in the males; in the females it is strongly pronounced, leaving several segments of the abdomen uncovered; the abdomen has as many corneous segments as are found in *Ptinella*, and the antennæ in length and tenuity resemble *Pterycodes*.

Dimorphella Reitteri.

D. Reitteri, Matthews, Ann. Mag. Nat. Hist. 6, iii. p. 190, 1889.

L. e. $\frac{5}{16}$ – $\frac{6}{16}$ lin. = 0.63–0.75 mm.—Subovata, valde convexa, nitidissima, vel flava, vel rufocastanea, pilis aureis sparse vestita, elytris in maribus foveolate punctatis; *capite* modico, antice producto, modice punctato, oculis modicis, valde prominentibus: *pronoto* capite parum longiori, multum latiori, ad medium latissimo, minute et remote punctato, interstitiis glabris, nitidis, lateribus marginatis, atque in maribus dente acuto, magnitudine variabili, ad media armatis, basin versus fortiter constrictis, margine basali recta, angulis salientibus; *elytris* capite atque pronoto multum et longioribus et latioribus, ante media latissimis, in maribus tumide incrassatis, et punctis foveolatis profundis remote impressis, interstitiis nitidissimis, in feminis modice convexis et multum abbreviatis, lateribus fortiter marginatis, apicibus plus minusve truncatis; abdomine plus minusve exserto, apice sat obtuso, integro; pedibus atque antennis prælongis, gracillimis, late flavis.

Body subovate, very convex and very shining, either yellow or rufo-castaneous, sparingly clothed with golden hair, marked with foveolate punctures on the elytra of the males.

Head moderate, produced in front, moderately punctured; eyes moderate, very prominent; antennæ very long and slender, yellow.

Thorax rather longer and much broader than the head, widest at the middle, minutely and remotely punctured with the interstices smooth and shining, sides margined, and in the males armed with a sharp tooth of variable size near the middle, strongly constricted towards the base, basal margin straight with the angles acute and salient.

Scutellum very large, triangular, indistinctly punctured, and very shining.

Elytra in the males much longer and broader than the head and thorax, in the females shorter, widest before the middle, in the males exceedingly convex and much incrassated, remotely marked with deep foveolate punctures, with the interstices very shining, in the females moderately convex, translucent and much abbreviated with the extremities rounded, sides strongly margined, extremities more or less truncate.

Abdomen more or less exposed, with the apex rather obtuse and entire.

Legs long and slender, yellow.

Under parts yellow, with the metasternum distinctly punctured and a row of deep punctures on each ventral segment.

Habitat, South America. Found in Brazil, near Blumenau.

Four males and two females of this pretty insect were found near Blumenau, in Brazil, and kindly presented to me by Herr E. Reitter.

GEN. VIII.—MIKADO. Pl. XII.

Mikado, Matthews, Ann. Mag. Nat. Hist. 6, iii. p. 189, 1889.

Corpus obeonicum, convexum.

Caput sat magnum, antice valde elongatum, fere rostriforme, sessile, deflexum; oculis permagnis, valde prominentibus; antennis juxta oculos insertis.

Antennæ (Pl. XII., fig. 7) 11-articulatæ; 1 magno, incrassato; 2 primo pariter magno, ad apicem abrupte truncato; 3-7 gracilibus, sibiipsis fere paribus; 8 septimo duplo breviori; 9 octavo simili; 10 ampulliformi, magno, incrassato; 11 decimo duplo majori, ad basim valde incrassato, ultra medium contracto; duobus apicalibus clavam biarticulatam formantibus.

Palpi maxillares (Pl. XII., fig. 5, s) parvi, gracillimi, 4-articulati; 1 longo, gracili; 2 primo parum longiori, gracili, anguste pyriformi; 3 secundo fere pariter longo, modice incrassato, ad apicem abrupte truncato; 4 aciculato, valde elongato, tenuissimo, ad basim parum incrassato.

Palpi labiales (Pl. XII., fig. 6, m) sat magni, triarticulati, mouiliformes, apicali cæteris parum majori, ad apicem subtruncato, atque seta longâ externe instructo 2 et 3 ovaliter rotundatis, sibiipsis paribus.

Labrum (Pl. XII., fig. 3) magnum, suboblongum, basin versus abrupte dilatatum, ad apicem leviter emarginatum, et utrinque rotundatum, atque processu membranaceo utrinque instructum et setis sat longis fimbriatum.

Mandibulæ (Pl. XII., fig. 4, t) modicæ, sat angustæ, apicibus bidenticulatis; lamellâ membranaceâ ovali. Stipes modicus, in processu longo, gracili, leviter incurvato, et dente magno, acuto, ad basim armato externe productus.

Maxille (Pl. XII., fig. 5) magnæ, robustæ, quasi trilobatæ, lobo *externo* sat brevi, leviter incurvato, apice acutissimo; *intermedio* parvo, debili, setis ad apicem instructo; *interiori* elongato, sat gracili, ad apicem acute uncinato et dentibus tribus acutissimis armato.—Stipes magnus, robustus.

Mentum (Pl. XII., fig. 6, j) modicum, fere quadratum.

Labium (Pl. XII., fig. 6, k) modicum, suboblongum, antice parum dilatatum, atque in processu paraglossali longo, acuto, divergenti, utrinque productum.

Lingua (Pl. XII., fig. 6, l) sat parva, palpis parum brevior, apice bifido.

Pronotum sat magnum, ad basim latissimum, angulis posterioribus, maribus parum productis, femineis haud productis.

Scutellum modicum, late triangulare.

Elytra elongata, apicibus oblique truncatis.

Alæ modicæ, lanceolatæ, nervis solitis, tenuioribus, firmatæ, et setis marginalibus prælongis instructæ.

Abdomen segmentis septem compositum, quinque basalibus membranaceis, duobus apicalibus corneis.

Prothorax (Pl. XII., fig. 2, a) parvus, angustis; *prosterno* perparvo; *episternis* inconspicuis; *epimeris* modicis, receptacula coxarum postice circumdati; receptaculis coxarum sat magnis, ovalibus, fere contingentibus, et antice fere opertis.

Pectus mesosterno (Pl. XII., fig. 2, d) sat magno, antice valde deflexo, late carinato, carinâ antice in apice jaculiformi terminato; *episternis* modicis, humeralibus, antice deflexis; *epimeris* permagnis, ad coxarum receptacula extensis; receptaculis coxarum modicis, rotundatis, per carinam divisis.

metasterno (Pl. XII., fig. 2, g) permagno, ad latera corporis fere extenso, postice inter coxas minute bidentato; *episternis* longis, angustis, vix conspicuis; *epimeris* longis, angustis, sub clytris totis occultis.

Venter segmentis sex compositus, omnibus fere pariter longis, apicali integro.

Pedes omnes graciles et sat breves; *antiei* trochanteribus parvis, trigonalibus; femoribus simplicibus; tibiis apices versus leviter dilatatis; tarsis triarticulatis; 1 et 2 brevibus, sat crassis, et setis sat longis ad apices instructis; 2 primo breviori; 3 prælongo, tenuissimo, unguibus gracilibus, leviter curvatis.

intermedii anticis similes, parum longiores.

postici (Pl. XII., fig. 8) trochanteribus claviformibus; femoribus robustis; tibiis apices versus interne serratis et setis brevibus instructis; alioqui anticis similes.

Coxae anteriores sat magnæ, ovales, sat prominentes, contingentes.

intermediae rotundatæ, per carinam divisæ.

posteriores sat magnæ, prominentes, late subovaliter laminatæ, fere contingentes.

Body obconic, convex.

Head rather large, exceedingly produced in front, almost rostriform, sessile, and strongly deflexed; eyes large and very prominent; antennæ inserted very near the eyes.

Antennæ (Pl. XII., fig. 7) eleven-jointed; 1 large and robust; 2 as large as the first, abruptly truncate at the apex; 3-7 slender, equal to each other in size; 8 twice as short as the seventh; 9 like the eighth; 10 ampulliform, large and incrassate; 11 twice as large as the tenth, roundly incrassated at the base, and much contracted beyond the middle; the two last joints forming a biarticulate club.

Maxillary palpi (Pl. XII., fig. 5, s) small and very slender, four-jointed; 1 long and slender; 2 rather longer than the first, slender and narrowly pyriform; 3 as long as the second, moderately incrassate, abruptly truncate at the apex; 4 very long and slender, aciculate, slightly incrassate at the base.

Labial palpi (Pl. XII., fig. 6, m) rather large, three-jointed, moniliform, with the apical joint largest, subtruncate at the apex, and furnished on the outer extremity with a long incurved seta; the two basal joints oval and of equal size.

Labrum (Pl. XII., fig. 3) large, suboblong, abruptly dilated at the base, slightly emarginate in front, rounded on each side at the apex, and furnished with a membranaceous appendage fringed with setæ.

Mandibles (Pl. XII., fig. 4, t) moderate, rather narrow, bidentate at the apex; lamella membranacea oval. Stipes moderate, produced into a long, slender, incurved process behind the mandibles with a large acute tooth on the outer side of its base.

Maxille (Pl. XII., fig. 5) large, robust, apparently trilobed, *external* lobe short, slightly incurved, very acutely pointed, forming part of the stipes; *intermediate* lobe small and feeble, furnished with setæ at the apex; *interior* lobe long, rather slender, acutely hooked at the apex, and armed with three very acute teeth. Stipes large and robust.

Mentum (Pl. XII., fig. 6, j) moderate, nearly quadrate.

Labium (Pl. XII., fig. 6, k) moderate, suboblong, slightly dilated in front, and produced on each side into a long paraglossal process acutely pointed and slightly divergent.

Lingua (Pl. XII., fig. 6, l) rather small, shorter than the palpi, and bifid at the apex.

Pronotum rather large, widest at the base with the hinder angles acute and slightly produced in the males, but not produced in the females.

Scutellum moderate, broadly triangular.

Elytra elongate, obliquely truncate at the apex.

Wings moderate, lancolate, with rather feeble neuration, fringed on both sides with long, movable setæ.

Abdomen composed of seven segments, of which the two apical are corneous, and the rest membranaceous.

Prothoracæ (Pl. XII., fig. 2, a) small and narrow; *prosternum* very small; *episterna* inconspicuous; *epimera* moderate, inclosing the coxal cavities behind; coxal cavities large, oval, almost contingent, and partly open in front.

Pectus with the *mesosternum* (Pl. XII., fig. 2, d) rather large, strongly deflexed in front, broadly carinated, with the anterior extremity of the carina broadly

lanceiform; *episterna* moderate, humeral, anteriorly much deflexed; *epimera* very large, reaching to the coxal cavities; coxal cavities moderate, rounded, separated by the carina.

metasternum (Pl. XII., fig. 2, g) very large, almost extending to the sides of the body, with the hinder margin minutely bidentate between the coxæ; *episterna* long and narrow, scarcely visible; *epimera* long and narrow, covered by the elytra.

Venter composed of six segments of nearly equal size, apical segment entire.

Legs slender and rather short, *anterior* with the trochanters small, trigonal; femora simple; tibiæ slightly dilated towards the apex; tarsi three-jointed; 1 and 2 short and stout, with a long seta on each side at their apex, 2 shorter than 1; 3 very long and very slender; claws slender, faintly curved.

intermediate like the anterior, but rather shorter.

posterior (Pl. XII., fig. 8) with the trochanters large, claviform; femora robust, simple; tibiæ with the inner side serrate towards the apex and setose, in other respects like the anterior.

Coxæ anterior rather large, oval, rather prominent, and contingent

intermediate round, separated by the carina.

posterior rather large, prominent and contingent, broadly and subovally laminated for their whole length.

Mikado Japonicus.

M. Japonicus, Matthews, Ann. Mag. Nat. Hist. 6, iii. p. 189, 1889.

L. c. $\frac{3}{16}$ – $\frac{4}{16}$ lin. = 0.37–0.50 mm.—Obconicus, elongatus, postice validissime attenuatus, nitidissimus, flavus, pilis brevibus aureis vestitus; *capite* sat magno, antice elongato, fere rostriformi; oculis nigris, magnis, valde prominentibus; *pronoto* modico, glabro, nitidissimo, ad basim latissimo, lateribus valde rotundatis, margine basali maribus leviter arcuata, angulis productis, fæmineis fere recta, angulis haud productis; *elytris* pronoto duplo longioribus, haud latioribus, ad humeros latissimis, postice valde attenuatis, leviter et remote asperatis, lateribus fere rectis, leviter marginatis, apicibus oblique truncatis; abdomine convexo, segmentis duobus apertis; pedibus atque antennis brevibus, flavis.

Body obconic, elongate, strongly attenuated posteriorly, very shining, yellow, clothed with short golden hair.

Head rather large, elongate, and almost rostriform in front; eyes black, large, and very prominent; antennæ short, pale yellow.

Thorax moderate, smooth, and very shining, widest at the base, sides much rounded, basal margin in the males faintly arcuate, with the angles produced, in the females nearly straight, with the angles acute, not produced.

Scutellum moderate, broadly triangular.

Elytra twice longer, but not broader than the thorax, widest at the shoulders, much attenuated posteriorly, faintly and remotely asperated, sides nearly straight, slightly margined, extremities obliquely truncate.

Abdomen conical, with two segments exposed.

Legs short and slender, yellow.

Under parts wholly yellow; apex of the venter entire.

Habitat, Asia.

This very distinct and remarkable insect was found plentifully in a fungus near Yuyama, Higo, Japan, by Mr. Lewis. It bears no resemblance to any genus but *Myrmicotrichis*, from which it is abundantly distinguished by its elongate rostriform head, and conspicuous black eyes.

GEN. XI.—ACTINOPTERYX.

*Actinopteryx australis.**A. australis*, Matthews, Cist. Ent. i. p. 93, 1872.

L. c. $\frac{4}{16}$ – $\frac{7}{16}$ lin. = 0.75–0.87 mm. — Oblongo-ovalis, sat depressa, pilis nitide argenteis vestita; capite sat magno; pronoto maribus ad latera valde dilatato, ad medium latissimo, femineis leviter rotundato, punctis sat magnis, distinctis, interstitiis glabris, nitidis, atque depressione lata, longitudinali utrinque in disco impresso, margine basali fere recta, angulis acutis, vix productis; elytris ad apices parum attenuatis, confertim sat profunde asperatis, lateribus modice rotundatis, apicibus dilutioribus, parum rotundatis; abdomine attenuato, longius exserto; pedibus atque antennis longis, gracillimis, pallide flavis.

Body oblong-oval, rather depressed, clothed with shining silver hair.

Head large, much rounded in front, marked with rather large, distinct punctures; eyes small, not prominent; antennæ very long and slender, pale yellow, with the three apical joints slightly incrassate.

Thorax in the male large, transverse, much broader but scarcely longer than the head, widest at the middle, sides much rounded and distinctly margined; in the female slightly rounded at the sides, widest near the base; in both sexes marked with large and distinct punctures, with the interstices smooth and shining, with a broad shallow longitudinal impression on each side near the middle of the disc; basal margin nearly straight, with the angles acute, scarcely produced.

Scutellum small, triangular, moderately asperate.

Elytra a little longer than the head and thorax, slightly attenuated posteriorly, moderately asperated in very close wavy rows, with a broad longitudinal impression on each side of the suture, sides slightly rounded and margined, apex very little rounded, pale, with the extreme edge white.

Abdomen considerably exerted, attenuated towards the apex.

Legs long and slender, pale yellow, femora dusky

Under parts pitchy castaneous, with the mouth, coxæ, and apex of the venter pale.

Differs from *A. fuscicola* in shape and sculpture, and in the short posterior angles of the thorax.

Habitat, Australasia. Found on the shores of the Swan River by Mr. Brewer, and in New Zealand by Mr. Lawson.

This very distinct and remarkable species appears to have a peculiar habit of covering its upper surface with mud after the manner of *Georyssus*.

GEN. XII.—TRICHOPTERYX.

Like *Homalota* among the *Staphylinide*, *Trichopteryx* contains a vast number of species, which to the naked eye, especially of an inexperienced observer, present no obvious difference; although when examined under a powerful microscope they are found to differ from each other abundantly in outline or sculpture, and generally in both these characters. Some may possibly be climatic or local forms, but I believe the vast majority are true species of science at least, if not of nature. I say "of science," because wherever any reasonable doubt exists it is always better to separate, than injudiciously to amalgamate diverse forms. The fact also that the form which may be regarded as the original type is often found in the same region and under similar conditions as the conjectural climatic or local variety, is strong evidence that the latter is specifically distinct, and not the off-

spring of evolutionary progress. Thus, for instance, the species which I have called *T. minor*, found by Mr. Champion in Guatemala, bears so strong a resemblance to the European *T. grandicollis* that it might be considered a local form of that species, the difference between them consisting in the much smaller size of *T. minor*, and in the shape and sculpture of its thorax; if, however, climatic or other influences had in this case led to metamorphosis, their effect would have produced similar results through the whole race; but, in contradiction to such a theory, *T. grandicollis*, agreeing in every respect with the European type, was also taken by Mr. Champion in the same region as *T. minor*; to separate the two, therefore, seems the only way to avoid confusion. Instances of a similar kind have also occurred with other species of *Trichopteryx*, and notably in the case of *T. atomaria*, of which the original type form is found in almost every region together with allied species.

The species of *Trichopteryx* are probably more numerous than those of any other genus of Coleoptera, and owing to their minute size the difficulty of distinguishing them from each other is so great, that an incipient student feels often inclined to give them up in despair. And, since it is sometimes impossible to arrive at a satisfactory determination without actual comparison, collectors should always bear this fact in mind—they should collect plentifully whenever an opportunity occurs, and distribute authentic types with the utmost liberality.

Owing to the multitude of species now contained in this genus, no synoptic tabulation would prove of any use. I have, therefore, in the following pages, given a separate and full list of every species which has hitherto been found in each principal division of the world; and have added to every species a few of its leading diagnostic characters.

SPECIERUM OMNIUM EPITOMA,

REGIONIBUS PROPRIIS SEGREGATARUM.

SPECIES EUROPEÆ.

ELYTRIS RUFESCENTIBUS.

L. c. 1 mm.—Brevis, lata, valde convexa, nitida, minutissime tuberculata, pronoto valde dilatato, ad basim latissimo, angulis valde productis, elytris attenuatis, ped. atque ant. flavis	Sara.
L. c. 0.90 mm.—Oblonga, valde convexa, nitidissima, modico et remote tuberculata, pronoto ad basim latissimo, angulis latis, valde productis, elytris parum attenuatis, ped. atque ant. rufescentibus	Hellenica.
L. c. 0.80 mm.—Oblonga, subparallela, modice tuberculata, pronoto ad basim latissimo, angulis parum productis, elytris oblongis, ped. atque ant. longis, flavis	Guerinii.
L. c. 0.80 mm.—Oblonga, elongata, sat fortiter tuberculata, pronoto ad basim latissimo, angulis vix productis, elytris brevibus, ped. atque ant. brevibus, flavis	obscæna.
L. c. 0.75 mm.—Ovata, subelongata, fortiter tuberculata, pronoto ante basim latissimo, angulis parum productis, elytris ovalibus, abdomine rufescenti, ped. atque ant. læte flavis	acuminata.
L. c. 0.60 mm.—Oblonga, brevis, modice tuberculata, pronoto subquadrato, angulis vix productis, elytris brevibus, quadratis, abdomine nigro, ped. atque ant. læte flavis	elevata.
L. c. 0.63 mm.—Oblonga, subparallela, depressa, minute et distincte tuberculata, pronoto quadrato, angulis parum productis, elytris brevibus, quadratis, abdomine fusco, ped. atque ant. flavis	Waterhousii.

OMNINO CASTANEÆ.

- L. c. 0·80 mm.—Subovata, nitida, convexa, haud tuberculata, profunde punctata, pronoto lateribus rotundatis, angulis haud productis, ped. atque ant. flavescentibus **variolosa.**
- L. c. 1 mm.—Lata, oblonga, valde convexa, modice tuberculata, pronoto magno, ad basim latissimo, angulis vix productis, elytris quadratis, ped. atque ant. longis, læte flavis **Poweri.**
- L. c. 0·75 mm.—Oblonga, distincte tuberculata, pronoto ante basim latissimo, angulis parum productis, elytris oblongis, ped. læte flavis, ant. obscuris **ambigua.**
- L. c. 0·75 mm.—Subovalis, aliquantum depressa, minute tuberculata, pronoto postice parum contracto, angulis haud productis, elytris postice latissimis, ped. atque ant. longis, flavis **dispar.**
- L. c. 0·75 mm.—Elongata, suboblonga, pilosa, modice tuberculata, pronoto ad medium latissimo, angulis parum productis, elytris longis, ovalibus, ped. atque ant. longis, flavis **Edithia.**
- L. c. 0·65 mm.—Brevis, quadrata, fortiter tuberculata, pronoto ad basim latissimo, angulis productis, elytris quadratis, ped. atque ant. longis, læte flavis **fuscula.**
- L. c. 0·85 mm.—Angusta, elongata, nitida, obscure castanea, minute tuberculata, pronoto dilatato, ad basim latissimo, angulis valde productis, elytris parum attenuatis, ped. atque ant. læte flavis . . . **Championis.**

OMNINO NIGRE, VEL FUSCÆ.

PRONOTO MULTUM DILATATO.

- L. c. 0·80 mm.—Lata, ovata, postice attenuata, fuscescens, nitida, remote tuberculata, pronoto ad basim latissimo, angulis valde productis, dilutioribus, elytris brevibus, attenuatis, ped. atque ant. flavis **atomaria.**
- L. c. 0·90 mm.—Ovalis, postice vix attenuata, fuscescens, densius tuberculata, pronoto ad basim latissimo, angulis valde productis, dilutioribus, elytris subquadratis, ped. flavis, ant. brevibus, picco-testaceis **brevicornis.**
- L. c. 0·80 mm.—Ovalis, obtusa, nigra, valde convexa, haud attenuata, sat minute tuberculata, pronoto ad basim latissimo, angulis latis, valde productis, elytris quadratis, ped. atque ant. læte flavis . . . **convexiuscula.**
- L. c. 0·60 mm.—Brevis, lata, postice attenuata, convexa, nitida, fuscescens, modice tuberculata, pronoto magno, angulis valde productis, elytris brevibus, valde attenuatis, ped. atque ant. flavis **thoracica.**
- L. c. 0·70 mm.—Subovata, nigra, maribus postice attenuatis, fortiter et remote tuberculata, pronoto ad basim latissimo, angulis productis, elytris fœminicis vix attenuatis, ped. flavis, ant. plus minusve piccis **anthracina.**
- L. c. 0·80 mm.—Brevis, lata, nigra, postice attenuata, minute tuberculata, pronoto ante basim latissimo, angulis productis, valde dilatatis, elytris brevibus, ped. flavis, ant. picco-nigris **fratercula.**
- L. c. 1–1·10 mm.—Sat lata, convexa, nigro-aenea, setis erectis instructa, fortiter tuberculata, pronoto magno, ad medium latissimo, angulis productis, elytris subquadratis, ped. flavis, ant. picco-testaceis . . . **grandicollis.**
- L. c. 1·25 mm.—Lata, subquadrata, fuscescens, fortiter et confertim tuberculata, pronoto ante basim latissimo, angulis latis, productis, elytris quadratis, ped. læte flavis, ant. longis, flavis **lata.**
- L. c. 1·10 mm.—Subovalis, maribus postice attenuatis, nigra, modice et confertim tuberculata, pronoto ad basim latissimo, angulis acutis, productis, elytris fœminicis oblongis, ped. læte flavis, ant. vel læte vel obscure flavis **fascicularis.**
- L. c. 0·80 mm.—Oblongo-ovalis, postice attenuata, nigra, minute tuberculata, pronoto ad basim latissimo, angulis acutis, productis, elytris attenuatis, ped. obscure flavis, ant. picco-nigris **Lætitia.**

- L. c. 1 mm.—Oblongo-ovalis, valde convexa, postice attenuata, nigra, fortiter et remote tuberculata, pronoto nitido, ad basim latissimo, angulis valde productis, elytris attenuatis, ped. flavis, ant. longis, piceis **seminitens.**
- L. c. 0·75 mm.—Ovata, postice valde attenuata, sat depressa, nigra, fortiter tuberculata, pronoto magno, ad basim latissimo, foveâ elongatâ prope angulos impresso, angulis longe productis, elytris valde attenuatis, ped. læte flavis, ant. nigro-piceis **attenuata.**
- L. c. 0·80 mm.—Elongata, angusta, nitida, nigra, postice attenuata, indistincte tuberculata, pronoto brevi, ad basim latissimo, angulis productis, acutissimis, elytris brevibus, parum attenuatis, ped. læte flavis, ant. obscure flavis **angusta.**

PRONOTO VIX AUT NON DILATATO.

- L. c. 1 mm.—Oblonga, lata, convexa, nigra, minute et confertim tuberculata, pronoto vix dilatato, angulis productis, acutis, elytris quadratis, ped. læte flavis, ant. piceis **cantiana.**
- L. c. 0·90 mm.—Oblonga, convexa, nigra, modice tuberculata, pronoto brevi, haud dilatato, angulis vix productis, elytris oblongis, ped. flavis, ant. piceis **volans.**
- L. c. 0·80 mm.—Brevis lata, haud attenuata, fuscescens, haud nitida, fortiter tuberculata, pronoto ad basim latissimo, angulis vix productis, elytris quadratis, ped. flavis, ant. nigro-piceis **sericans.**
- L. c. 0·60 mm.—Perbrevis, convexa, haud attenuata, fuscescens, haud nitida, modice tuberculata, pronoto brevi, angulis acutis, parum productis, elytris brevibus, quadratis, ped. flavis, ant. piceo-testaceis **brevis.**
- L. c. 0·70 mm.—Ovalis, aterrima, haud nitida, convexa, modice et sat confertim tuberculata, pronoto brevi, ad basim latissimo, angulis acutis, vix productis, elytris ovalibus, ped. læte flavis, ant. brevibus, piceo-nigris **bovina.**
- L. c. 0·80 mm.—Brevis, lata, convexa, nigra, minute tuberculata, pronoto ante basim latissimo, angulis parum productis, elytris perbrevis, ped. piceis, tibiis atque tarsis anticis maribus dilatatis, ant. longis, piceo-testaceis **brevipennis.**
- L. c. 0·75 mm.—Ovata, convexa, nigra, modice tuberculata, pronoto magno, ante basim latissimo, angulis acutis, elytris ovalibus, postice contractis, ped. longis, flavis, tibiis atque tarsis anticis maribus dilatatis, ant. longis, piceis **Kirbii.**
- L. c. 0·80 mm.—Oblonga, parallela, sat depressa, nigra, haud nitida, minute et confertissime tuberculata, pronoto quadrato, angulis parum productis, elytris oblongis, ped. longis, flavis, ant. longis, piceis **longicornis.**
- L. c. 1 mm.—Elongata, angusta, convexa, nigra, modice tuberculata, pronoto brevi, ad medium latissimo, angulis parum productis, elytris oblongis, ped. flavis, ant. nigro-piceis **longula.**
- L. c. 0·75 mm.—Oblonga, convexa, nigra, minute tuberculata, pronoto ante basim latissimo, angulis acutis, productis, elytris brevibus, subquadratis, ped. atque ant. obscure flavis **carbonaria.**
- L. c. 0·75 mm.—Oblonga, convexa, nigra, nitida, modice tuberculata, pronoto ante basim latissimo, angulis latis, productis, ped. obscure flavis, ant. nigro-piceis **piceicornis.**
- L. c. 0·80 mm.—Oblonga, subcylindrica, nigra, modice tuberculata, pronoto brevi, subquadrato, angulis flavescens, parum productis, elytris oblongis, ped. læte flavis, ant. piceo-testaceis **Jansoni.**
- L. c. 0·80 mm.—Oblonga, convexa, nitida, fuscescens, remote tuberculata, pronoto ad basim latissimo, angulis parum productis, elytris oblongis, ant. longis atque ped. flavis **Montandoni.**
- L. c. 0·80 mm.—Oblongo-ovata, convexa, nitida, nigra, sat minute tuberculata, pronoto ad basim latissimo, angulis productis, elytris ovalibus, ped. læte flavis, ant. flavis **rivularis.**

- L. c. 0.50 mm.—Brevis, oblonga, parallela, nigra, vix tuberculata, pronoto brevi, quadrato, angulis rectis, elytris brevibus, quadratis, ped. lacte flavis, ant. longis, piccis Chevrolati.
- L. c. 0.80 mm.—Oblonga, parallela, nigra, modice et confertim tuberculata, pronoto brevi, ad basim latissimo, angulis parum productis, elytris oblongis, ped. atque ant. flavis Taurica.

SPECIES ASIATICÆ.

ELYTRIS RUFESCENTIBUS.

- L. c. 1.25 mm.—Lata, convexa, postice attenuata, pilosa, minute tuberculata, pronoto dilatato, ad basim latissimo, angulis rufescentibus, valde productis, elytris brevibus, attenuatis, ped. lacte flavis, ant. longis, obscure flavis Lewisii.
- L. c. 1 mm.—Lata, suboblonga, sat fortiter tuberculata, pronoto magno, ad basim latissimo, angulis latis, valde productis, elytris subquadratis, ped. atque ant. sat longis, lacte flavis Caucasica.
- L. c. 1.10 mm.—Brevis, lata, convexa, postice parum attenuata, minute tuberculata, pronoto magno, dilatato, ad basim latissimo, angulis latis, productis, elytris attenuatis, ped. atque ant. lacte flavis Ceylonica.
- L. c. 1 mm.—Ovata, convexa, postice attenuata, modice tuberculata, pronoto magno, ad basim latissimo, angulis latis, productis, elytris brevibus, attenuatis, ped. lacte flavis, ant. obscure flavis cursitans.
- L. c. 0.87 mm.—Omnino ovata, valde convexa, postice attenuata, pronoti margine basali arcuatâ, angulis valde productis; elytris valde attenuatis; ped. atque ant. flavis soror.

OMNINO NIGRÆ VEL FUSCÆ.

- L. c. 1.10 mm.—Suboblonga, lata, convexa, nigro-ænea, setis erectis instructa, modice tuberculata, pronoto magno, ad medium latissimo, angulis parum productis, elytris ovatis, ped. flavis, ant. picco-nigris grandicollis.
- L. c. 1.15 mm.—Oblonga, convexa, picco-nigra, setis erectis sæpe instructa, modice tuberculata, pronoto lato, ante basim latissimo, angulis parum productis, elytris longis, oblongis, ped. atque ant. longis, lacte flavis crassicollis.
- L. c. 1.10 mm.—Brevis, lata, postice parum attenuata, minute et dense tuberculata, pronoto magno, ad basim latissimo, angulis latis valde productis, elytris attenuatis, ped. atque ant. lacte flavis orientalis.
- L. c. 1 mm.—Oblonga, nigra, nitida, fortiter tuberculata, pronoto brevi, ad basim latissimo, angulis parum productis, elytris vix attenuatis, ped. obscure flavis, ant. longis, piccis atrata.
- L. c. 0.80 mm.—Oblonga, aterrima, haud nitida, modice tuberculata, pronoto brevi, subquadrato, angulis parum productis, elytris haud attenuatis, ped. flavis, ant. longioribus, piccis volans.
- L. c. 0.60 mm.—Brevis, quadrata, sat depressa, fusca, fortiter tuberculata, pronoto brevi, transversa, ad basim latissimo, angulis productis, elytris brevibus, quadratis, ped. atque ant. longis, gracilibus, lacte flavis fuscula.
- L. c. 0.65 mm.—Brevis, subquadrata, convexa, nigra, nitida, modice et distincte tuberculata, pronoto sat magno, ad basim dilatato, angulis productis, elytris brevibus, attenuatis, ped. lacte flavis, ant. piccis inconspicua.
- L. c. 0.65 mm.—Subovata, angusta, convexa, nitida, castanea, profunde et remote punctata, haud tuberculata, pronoto ad medium latissimo, angulis aentis, elytris ovatis, ped. atque ant. longis, flavis Japonica.

SPECIES AFRICANÆ ATQUE ATLANTICÆ.

ELYTRIS RUFESCENTIBUS.

- L. c. 0·80 mm.—Brevis, lata, valde convexa, nitida, postice attenuata, modice tuberculata, pronoto dilatato, ad basim latissimo, angulis latis, valde productis, elytris brevibus, attenuatis, ped. læte flavis, ant. flavis **Matthewsii.**
- L. c. 0·70 mm.—Brevis, convexa, tota obscure castanea, minute tuberculata, pronoto magno, dilatato, ad basim latissimo, angulis valde productis, elytris brevibus, valde attenuatis, ped. atque ant. læte flavis **Crotchii.**
- L. c. 1·10 mm.—Oblonga, subcylindrica, tota rufo-fusca, modice tuberculata, pronoto brevi, ad basim latissimo, angulis parum productis, elytris oblongis, haud attenuatis, ped. atque ant. flavis **Lethierryi.**
- L. c. 0·80 mm.—Oblonga, elongata, convexa, fortiter tuberculata, pronoto modico, ad basim latissimo, angulis vix productis, elytris brevibus, haud attenuatis, ped. flavis, ant. nigro-piceis **obsœna.**

ELYTRIS NIGRIS, VEL FUSCIS.

- L. c. 0·80 mm.—Lata, ovata, nigro-fusca, postice attenuata, nitidula, remote tuberculata, pronoto dilatato, ad basim latissimo, angulis valde productis, elytris brevibus, attenuatis, ped. atque ant. læte flavis **atomaria.**
- L. c. 0·90 mm.—Ovalis, postice vix attenuata, nigro-fusca, modice tuberculata, pronoto ad basim latissimo, angulis productis, elytris oblongis, ped. flavis, ant. brevibus, obscuris **brevicornis.**
- L. c. 0·80 mm.—Oblonga, convexa, nitida, haud attenuata, remote tuberculata, pronoto ad basim latissimo, angulis vix productis, elytris oblongis, ped. atque ant. flavis **Montandoni.**
- L. c. 0·80 mm.—Oblonga, convexa, nitidula, nigro-fusca, haud attenuata, modice tuberculata, pronoto subquadrato, angulis acutis, parum productis, elytris oblongis, apicibus atque suturâ rufescentibus, ped. flavis, ant. nigro-piceis **nigricornis.**
- L. c. 0·80.—Brevis, lata, haud attenuata, nigro-fusca, haud nitida, sat rugose tuberculata, pronoto ad basim latissimo, angulis parum productis, elytris quadratis, ped. flavis, ant. nigro-piceis **sericans.**
- L. c. 0·60 mm.—Ovalis, brevis, convexa, nigro-fusca, modice et confertim tuberculata, pronoto fere quadrato, angulis vix productis, elytris ovatis, ad media latissimis, ped. robustis, læte flavis, ant. obscure flavis **ovatula.**
- L. c. 0·70 mm.—Subovata, nigra, maribus postice attenuatis, fortiter et remote tuberculata, pronoto ad basim latissimo, angulis productis, elytris fœmineis vix attenuatis, ped. flavis, ant. plus minusve piceis **anthracina.**
- L. c. 0·90 mm.—Oblongo-ovata, fuscescens, nitidula, sat minute tuberculata, postice parum attenuata, pronoto ad basim latissimo, angulis valde productis, elytris parum attenuatis, ped. læte flavis, ant. longis, pallido flavis **Wollastoni.**
- L. c. 1·10 mm.—Lata, convexa, nigra, postice parum attenuata, fortiter tuberculata, pronoto valde dilatato, ad basim latissimo, angulis latis, valde productis, elytris brevibus, postice attenuatis, ped. atque ant. longis, læte flavis **umbricola.**
- L. c. 0·70 mm.—Oblonga, subparallela, aterrima, haud nitida, fortiter tuberculata, pronoto subquadrato, angulis parum productis, elytris sat longis, parallelis, ped. flavis, ant. brevibus, nigro-piceis **Canariensis.**

SPECIES AMERICANÆ.

ELYTRIS RUFESCENTIBUS.

- L. c. 1·10 mm.—Sat elongata, valde convexa, nitida, sat remote et distincte tuberculata, pronoto valde dilatato, angulis productis, elytris attenuatis, ped. atque ant. læte flavis **Haldemanni.**
- L. c. 0·80 mm.—Brevis, lata, valde convexa, nitida, haud tuberculata, pronoto magno, ante basim latissimo, angulis productis, pallidis, elytris brevibus, postice latissimis, ped. atque ant. flavis **glabricollis.**
- L. c. 0·80 mm.—Brevis, convexa, postice attenuata, modice et remote tuberculata, pronoto dilatato, ad basim latissimo, angulis productis, flavis, elytris postice attenuatis, ped. atque ant. læte flavis **Dohrnii.**
- L. c. 0·80 mm.—Subovata, nitida, postice parum attenuata, pronoto ad basim latissimo, fortiter tuberculato, angulis productis, pallidis, elytris testaceis, depressis, ped. atque ant. læte flavis **discolor.**
- L. c. 0·70 mm.—Brevis, lata, convexa, parum attenuata, interdum tota castanea, pronoto dilatato, confertim tuberculato, angulis valde productis, elytris brevibus, attenuatis, ped. atque ant. læte flavis. **Sallæi.**
- L. c. 0·75 mm.—Brevis, lata, convexa, valde attenuata, interdum tota castanea, pronoto dilatato, remote tuberculato, angulis productis, pallidis, elytris brevibus, valde attenuatis, ped. atque ant. læte flavis **Crotchii.**
- L. c. 0·80 mm.—Suboblonga, convexa, nitidula, haud attenuata, pronoto ad basim latissimo, modice et remote tuberculato, angulis parum productis, elytris subquadratis, ped. atque ant. obscure flavis **similior.**
- L. c. 0·80 mm.—Ovalis, elongata, convexa, nitidula, pronoto subquadrato, ad basim latissimo, remote tuberculato, angulis productis, elytris ovalibus, ped. atque ant. læte flavis **parallela.**
- L. c. 1 mm.—Oblonga, convexa, nitida, pronoto parum dilatato, minute tuberculato, angulis acutis, productis, flavis, elytris oblongis, ped. atque ant. longis, læte flavis **Wenkeri.**
- L. c. 0·75 mm.—Suboblonga, convexa, nitida, setis crectis instructa, pronoto ante basim latissimo, fortiter tuberculato, angulis productis, elytris brevibus, quadratis, ped. læte flavis, ant. obscure flavis **Brasiliensis.**
- L. c. 0·90 mm.—Brevis, lata, valde convexa, nitidissima, translucida, postice attenuata, pronoto magno, valde dilatato, minute tuberculato, angulis longe productis, elytris brevibus, valde attenuatis, ped. atque ant. flavis **lucida.**
- L. c. 0·90 mm.—Ovata, convexa, nitida, postice attenuata, pronoto magno, dilatato, distincte tuberculato, angulis valde productis, elytris brevibus, attenuatis, ped. læte flavis, ant. flavis **rufescens.**
- L. c. 1 mm.—Oblonga, lata, haud attenuata, pronoto ad basim latissimo, minute tuberculato, angulis productis, flavis, elytris quadratis, rufo-piceis, ped. atque ant. lætissime flavis **xanthocera.**
- L. c. 0·80 mm.—Subhemispherica, postice attenuata, pronoto nigro, nitidissimo, elytris brevibus, rufescentibus, ped. atque ant. flavis. **lævicollis.**
- L. c. 0·80 mm.—Ovata, latissima, pronoto nigro, elytris fuscis, dense tuberculata et pilosa, ant. longis atque ped. flavis **occidentalis.**

OMNINO CASTANEÆ.

- L. c. 1 mm.—Elongata, postice valde attenuata, nitidissima, haud tuberculata, pronoto dilatato, angulis validissime productis, elytris angustis, valde attenuatis, ped. atque ant. læte flavis **Schaumii.**
- L. c. 1 mm.—Oblonga, convexa, nitida, pronoto magno, ad basim latissimo, minute tuberculato, angulis acutis, valde productis, elytris oblongis, ped. atque ant. læte flavis **Californica.**
- L. c. 0·90 mm.—Oblonga, convexa, haud attenuata, pronoto ad basim latissimo, modice tuberculato, angulis valde productis, flavis, elytris oblongis, translucidis, ped. atque ant. flavis **castanea.**

- L. c. 0·80 mm.—Suboblonga, fuscescens, haud nitida, pronoto ad basim latissimo, minute tuberculato, angulis haud productis, elytris postice dilatatis, ped. longis, late flavis, ant. longis, flavis . . . Reichii.
- L. c. 0·70 mm.—Oblonga, sat convexa, haud nitida, pronoto ante basim latissimo, distincte tuberculato, angulis parum productis, elytris oblongis, ped. late flavis, ant. obscuris . . . ambigua.
- L. c. 0·75 mm.—Elongato-oblonga, pilosa, haud nitida, pronoto ante basim latissimo, confertim tuberculato, angulis haud productis, elytris subparallelis, ped. atque ant. longis, late flavis . . . Aubæi.
- L. c. 1 mm.—Suboblonga, depressa, haud nitida, pronoto minute tuberculato, ad basim abrupte contracto, angulis acutis, haud productis, elytris postice dilatatis, ped. atque ant. longis, late flavis . . . Motschulskii.
- L. c. 0·87 mm.—Elongata, angusta, pallide castanea, oculis nigris, pronoto brevi, ad medium latissimo, elytris brevibus, ad media latissimis, ant. longis atque ped. flavis . . . dubitata.

OMNINO NIGRÆ, AUT FUSCÆ.

PRONOTO DILATATO.

- L. c. 2 mm.—Lata, convexissima, nitida, postice attenuata, pronoto permagno, minute tuberculato, angulis productis, elytris sat brevibus, postice attenuatis, ped. late flavis, ant. obscuris . . . Godmanni.
- L. c. 1·10 mm.—Latissima, convexissima, nitida, setis erectis instructa, pronoto permagno, valde dilatato, minute et remote tuberculato, angulis productis, pallidis, elytris brevibus, valde attenuatis, ped. atque ant. obscure flavis . . . grossa.
- L. c. 1·10 mm.—Oblonga, convexa, sat nitida, pronoto magno, distincte tuberculato, ad basim latissimo, angulis latis, productis, elytris parum attenuatis, ped. atque ant. longis, flavis . . . laticollis.
- L. c. 1·10 mm.—Lata, convexa, nitidula, postice valde attenuata, setis erectis instructa, pronoto permagno, ante basim latissimo, fortiter tuberculato, angulis valde productis, elytris valde attenuatis, ped. atque ant. late flavis . . . Alliberti.
- L. c. 1·15 mm.—Oblonga, convexa, picco-nigra, haud attenuata, setis erectis sæpe instructa, pronoto lato, ante basim latissimo, modice tuberculato, angulis parum productis, elytris longis oblongis, ped. atque ant. longis, late flavis . . . crassicollis.
- L. c. 1·10 mm.—Suboblonga, lata, convexa, nigro-ænea, setis erectis instructa, pronoto magno, ad medium latissimo, sat fortiter tuberculato, angulis parum productis, elytris ovatis, ped. flavis, ant. nigro-piceis . . . grandicollis.
- L. c. 0·80 mm.—Oblongo-ovata, nitidula, setis erectis instructa, fuscescens, pronoto sat longo, fortiter tuberculato, ad basim latissimo, angulis productis, pallidis, elytris subquadratis, ped. atque ant. longis, flavis . . . minor.
- L. c. 1·10 mm.—Subovalis, maribus postice attenuatis, nigra, pronoto ad basim latissimo, modice et confertim tuberculato, angulis acutis, productis, elytris femineis oblongis, ped. late flavis, ant. vel late vel obscure flavis . . . fascicularis.
- L. c. 1·10 mm.—Ovalis, sat angusta, nitidula, pronoto modico, ad basim latissimo, sat minute tuberculato, angulis productis, pallidis, elytris angustis, postice attenuatis, ped. late flavis, ant. obscuris . . . cognata.
- L. c. 0·80 mm.—Brevis, convexa, nigra, pronoto valde dilatato, ante basim latissimo, minute et remote tuberculato, angulis latis, productis, elytris brevibus, quadratis, ped. atque ant. brevibus, piccis . . . Henrici.
- L. c. 0·75 mm.—Brevis, lata, convexa, pronoto lato, ante basim latissimo, fortiter tuberculato, angulis modice productis, elytris brevibus, postice contractis, ped. atque ant. flavis . . . abrupta.

- L. c. 0·80 mm.—Lata, ovata, postice attenuata, fuscescens, nitida, pronoto ad basim latissimo, remote tuberculato, angulis valde productis, dilutionibus, elytris brevibus, attenuatis, ped. atque ant. flavis **atomaria.**
- L. c. 0·70 mm.—Perbrevis, postice valde attenuata, convexa, nigra, pronoto magno, fortissime tuberculato, dilatato, ad basim latissimo, angulis latis, valde productis, elytris perbrevis, valde attenuatis, ped. læte flavis, ant. obscuris **truncatissima.**
- L. c. 0·75 mm.—Oblonga, fuscescens, nitida, setis brevibus erectis instructa, pronoto ante basim latissimo, fortiter tuberculato, angulis latis, productis, elytris quadratis, ped. læte flavis, ant. obscuris **Brasiliensis.**
- L. c. 1·25 mm.—Brevis, lata, convexa, nitida, setis erectis instructa, haud tuberculata, pronoto magno, ad basim latissimo, umbilicate punctato, angulis latis, productis, elytris subquadratis, strigose asperatis, ped. atque ant. læte flavis **reticulata.**
- L. c. 1 mm.—Suboblonga, thoracis lateribus valde dilatatis et rotundatis, elytris brevibus, angustis, ped. atque ant. longis, flavis **angustipennis.**

PRONOTO VIX AUT NON DILATATO.

- L. c. 0·85 mm.—Oblonga, convexa, nitidula, nigra, pronoto prope medium latissimo, modice et confertim tuberculato, angulis acutis valde productis, elytris haud attenuatis, prope media latissimis, ant. longis, atque ped. læte flavis **Hornii.**
- L. c. 1 mm.—Sat brevis, lata, nigra, pronoto brevi, ad basim latissimo, sat minute tuberculato, angulis parum productis, elytris postice parum attenuatis, ant. longis, atque ped. læte flavis **Sitkaensis.**
- L. c. 0·75 mm.—Oblonga, sat angusta, nigra, pronoto magno, ad basim latissimo, confertim et fortiter tuberculato, margine posteriori flavescenti, angulis productis, elytris oblongis, ped. læte flavis, ant. nigris **Josephi.**
- L. c. 1 mm.—Oblonga, convexa, nigra, parallela, pronoto ad basim latissimo, modice et confertim tuberculato, angulis pallidis, valde productis, elytris oblongis, ped. læte flavis, ant. piceis **parallelopipeda.**
- L. c. 0·80 mm.—Oblonga, sat lata, nigra, pronoto ante basim latissimo, modice tuberculato, angulis pallidis productis, elytris quadratis, ped. læte flavis, ant. piceis **diffinis.**
- L. c. 0·75 mm.—Oblonga, fuscescens, haud attenuata, pronoto parum dilatato, prope basim latissimo, remote et indistincte tuberculato, angulis valde productis, elytris quadratis, abdomine castaneo, ped. atque ant. læte flavis **brevior.**
- L. c. 0·75 mm.—Oblonga, nitida, aterritima, pronoto magno, ad basim latissimo, sat minute et remote tuberculato, angulis productis, pallidis, elytris brevibus, quadratis, ped. læte flavis, ant. nigris **vicina.**
- L. c. 0·60 mm.—Oblonga, nigra, haud nitida, pronoto brevi, parum dilatato, ad basim latissimo, modice tuberculato, angulis vix productis, elytris oblongis, ped. atque ant. flavis **nigrita.**
- L. c. 0·75 mm.—Oblonga, nigra, haud nitida, pronoto ad medium latissimo, fortiter tuberculato, angulis vix productis, elytris ovalibus, ad media latissimis, ped. longis, læte flavis, ant. obscuris **bidilatata.**
- L. c. 0·80 mm.—Brevis, lata, haud attenuata, fuscescens, haud nitida, pronoto ad basim latissimo, sat rugose tuberculato, angulis parum productis, elytris quadratis, ped. flavis ant. nigro-piceis **sericans.**
- L. c. 0·60 mm.—Perbrevis, sat depressa, fuscescens, haud nitida, pronoto brevi, ad basim latissimo, minute et remotissime tuberculato, angulis vix productis, elytris perbrevis, quadratis, ped. atque ant. læte flavis **tesserula.**
- L. c. 0·60 mm.—Suboblonga, sat depressa, nigra, haud nitida, pronoto brevi, ad basim latissimo, fortiter tuberculato, angulis vix productis, elytris ad apices latissimis, ped. flavis, ant. piceis **aspera.**

- L. c. 0.60 mm.—Brevis, lata, valde convexa, aterrima, nitidissima, glabra, pronoto prope medium latissimo, angulis rectis, elytris perbrevis, ad media latissimis, ant. longis, atque ped. lute flavis **mærens.**
- L. c. 0.80 mm.—Oblonga, convexa, nigra, setis erectis instructa, pronoto brevi, ante basim latissimo, modice tuberculato, angulis productis, elytris oblongis, ped. atque ant. piceis **Chilensis.**
- L. c. 1 mm.—Oblongo-ovalis, nigro-fusca, haud nitida, pilis nigro-fuscis dense vestita; pronoto ante basim latissimo; elytris brevibus, ante media latissimis; ant. piceis, ped. lute flavis **fuliginosa.**
- L. c. 0.90 mm.—Oblongo-ovalis, depressa, haud attenuata, piceo-nigra, elytris parallelis, ant. atque ped. flavis **depressa.**
- L. c. 0.60 mm.—Oblonga, brevis, lata, nigra, pronoto atque elytris vicissim prope media dilatatis, ped. atque ant. brevibus, robustis, flavis **Grenadensis.**

Trichopteryx Lewisii.

T. Lewisii, Matthews, Cist. Ent. iii. p. 79, 1884.

L. c. $\frac{9}{16}$ — $\frac{11}{16}$ lin. = 1.12–1.36 mm.—Perlata, convexissima, postice attenuata, capite atque pronoto nigris, elytris rufis, pilis longis flavis vestita; capite magno, antice rotundato, tuberculis minutis remote ornato; oculis modicis, haud prominentibus; pronoto permagno, valde dilatato, convexissimo, capite longiori et multo latiori, ad basim latissimo, tuberculis minutis, ordinibus sinuatis, confertissimis, interstitiisque sat nitidis, ornato, lateribus rotundatis et late marginatis, margine rufa, margine basali profunde sinuata, angulis rufis, acutis, latis et longe productis; elytris rufis, capite atque pronoto angustioribus, haud longioribus, ad humeros latissimis, in maribus postice valde attenuatis, ordinibus transversis confertissime, haud profunde, asperatis, lateribus fere rectis, apicibus latis, leviter rotundatis; abdomine longius exserto, apice tridentato; pedibus sat longis, lute flavis; antennis longis, gracillimis, obscure flavis.

Body very broad and very convex, attenuated posteriorly, head and thorax black, elytra rufous, clothed with long yellow hair.

Head large, rounded in front, ornamented with minute, remote tubercles; eyes moderate, not prominent; antennæ long and very slender, dusky yellow.

Thorax very large and very broad, very convex, longer and much broader than the head, widest at the base, closely ornamented with minute tubercles in wavy rows, interstices rather shining, sides rounded and broadly margined, margin rufous, basal margin deeply sinuated with the angles rufous, acute, broad, and much produced.

Scutellum large, triangular, closely but not deeply asperated.

Elytra rufous, narrower and not longer than the head and thorax, widest at the shoulders, much attenuated posteriorly in the males, very closely but not deeply asperated in transverse rows, sides nearly straight, extremities broad and slightly rounded.

Abdomen much exserted, tridentate at the apex.

Legs rather long, bright yellow.

Under parts castaneous, with the mouth, coxæ, and extremity of the venter pale.

Differs from T. umbricola in its larger size, broader and more depressed form, rufous elytra, and in sculpture.

Habitat, Asia. Found in various parts of Japan by Mr. Lewis.

Trichopteryx Caucasica.

T. Caucasica, Matthews, Ann. Mag. Nat. Hist. 6, iii. 1889, p. 191.

L. c. $\frac{8}{16}$ – $\frac{9}{16}$ lin. = 1.12 mm.—Late oblonga, valde convexa, nitidula, pilis aureis vestita, sat rugose tuberculata, rufo-picea, elytris rufescentibus; capite magno, nigro, leviter punctato, interstitiis glabris, nitidis; oculis parvis, haud prominentibus; pronoto magno, nigro-piceo, ad basin latissimo, capite multo latiori et longiori, tuberculis magnis confertim dispositis, interstitiisque leviter reticulatis ornato, lateribus marginatis et valde rotundatis, margine basali fere recta, angulis latis longe productis; elytris rufescentibus, fere quadratis, capite atque pronoto parum angustioribus, haud longioribus, profunde et confertim asperatis, lateribus rectis et marginatis, apicibus latis, vix rotundatis, dilutioribus; abdomine sat longe exserto, rufescenti, leviter asperato, apice bidentato; pedibus atque antennis sat longis, laete flavis.

Body broadly oblong, very convex, moderately shining, clothed with golden hair, coarsely tuberculate, rufo-piceous with the elytra rufescent.

Head large, black, indistinctly punctured, with the interstices smooth and shining; eyes small, not prominent; antennae rather long, bright yellow.

Thorax large, nigro-piceous, widest at the base, much broader and longer than the head, ornamented with large tubercles closely arranged, and slightly reticulated interstices, sides margined and much rounded, basal margin nearly straight with the angles broad and much produced.

Scutellum very large, triangular, deeply and closely asperated.

Elytra rufescent, almost quadrate, rather narrower and not longer than the head and thorax, deeply and closely asperated, sides straight and margined, extremities paler, broad, and very little rounded.

Abdomen considerably exserted, rufescent, faintly asperated, apex bidentate.

Legs rather long, bright yellow.

Under parts castaneous.

Differs from T. Matthewsii in its much larger size, coarser sculpture, large, dilated thorax, and quadrate elytra.

Habitat, Asia. Found in the Caucasian district (Herr Reitter).

Trichopteryx lævicollis.

T. lævicollis, Matthews, Ann. Mag. Nat. Hist. 6, xiii. 1891, p. 338.

L. c. $\frac{6}{16}$ – $\frac{7}{16}$ lin. = 0.75–0.85 mm.—Subhemispherica, postice attenuata, pilis aureis sat dense vestita, pronoto nigro, angulis rufescentibus, elytris rufescentibus; capite brevi, lato, nigro, nitido, oculis modicis, haud prominentibus, antennis sat brevibus, laete flavis; pronoto permagno, valde convexo, nitidissimo, ad basin latissimo, tuberculis minutis, remotis, interstitiis glabris, nitidissimis ornato, lateribus marginatis, margine basali concavata, angulis latis valde productis, rufescentibus; elytris rufescentibus, brevibus, postice attenuatis, capite atque pronoto multo brevioribus, sat angustioribus, ad humeros latissimis, ordinibus obliquis, transversis, profunde, asperatis, interstitiis nitidis, lateribus marginatis, apicibus leviter rotundatis; abdomine minime exserto, pygidio rufescenti; pedibus modicis, flavis.

Body subhemispheric, attenuated posteriorly, rather thickly clothed with golden hair, thorax black, with the angles rufescent, elytra rufescent.

Head black, short and broad, shining; eyes moderate, not prominent; antennae rather short, bright yellow.

Thorax very large, very convex, and very shining, widest at the base, ornamented with minute and remote tubercles with the interstices smooth and very shining, sides margined, basal margin concave with the angles broad and very much produced, rufous.

Scutellum large, triangular, rufescent, deeply asperate.

Elytra rufescent, short, attenuated posteriorly, much shorter and rather narrower than the head and thorax, widest at the shoulders, deeply asperated in transverse oblique rows with the interstices shining, sides margined, extremities slightly rounded.

Abdomen very little exerted, pygidium rufescent.

Legs robust, rather short, yellow.

Under parts black, coxæ yellow, meso- and metasterna deeply asperated, venter piceous with the apex rufescent.

Differs from other species in its short and exceedingly convex form, very shining thorax, rugosely sculptured and very short elytra.

Habitat, America. Found plentifully in the West Indian Islands by Mr. Smith.

Trichopteryx Hellenica.

T. Hellenica, Matthews, Ann. Mag. Nat. Hist. 6, iii. 1889, p. 191.

L. c. $\frac{7}{16}$ lin. = 0.87 mm.—Oblonga, valde convexa, nitidissima, pilis flavis vestita, rufo-picea, remote tuberculata; capite magno, modice tuberculato, interstitiis glabris, nitidis; oculis parvis, haud prominentibus; pronoto magno, valde convexo, nitidissimo, capite multum longiori et latiori, ad basin latissimo, tuberculis parvis, ordinibus remotis sinuatis dispositis, interstitiisque glabris, nitidissimis ornato, lateribus marginatis, valde rotundatis, margine basali leviter arcuata, angulis latissimis longe productis; elytris brevibus, rufo-piceis, postice parum attenuatis, capite atque pronoto parum brevioribus et angustioribus, ad humeros latissimis, ordinibus remotis, sinuatis modice asperatis, interstitiis fere glabris, lateribus marginatis et leviter rotundatis, apicibus latis, vix rotundatis, dilutioribus; abdomine castaneo, modice exserto; pedibus atque antennis sat brevibus, rufescentibus.

Body oblong, very convex and very shining, clothed with yellow hair, rufo-piceous, remotely tuberculate.

Head large, moderately tuberculate with the interstices smooth and shining; eyes small, not prominent; antennæ rather short, rufescent.

Thorax large, very convex, and very shining, much longer and broader than the head, widest at the base, ornamented with small tubercles in remote, sinuated rows, interstices smooth and very shining, sides much rounded and margined, basal margins faintly arcuate with the angles very broad and much produced.

Scutellum very large, triangular, and closely asperated.

Elytra rufo-piceous, short, slightly attenuated posteriorly, rather shorter and narrower than the head and thorax, widest at the shoulders, moderately asperated in remote wavy rows with the interstices almost smooth, sides margined and slightly rounded, extremities paler, broad, and scarcely rounded.

Abdomen castaneous, moderately exerted.

Legs rather short, rufescent.

Under parts castaneous.

Differs from *T. Caucasica* in its smaller size, very convex and shining thorax, and faint sculpture.

Habitat, Europe. Found in the Morea (Reitter).

Trichopteryx soror.

T. soror, Flach, Verh. Zool.-bot. Ges. Wien. xxxix. 1889, p. 520.

L. c. $\frac{7}{16}$ lin. = 0.87 mm.—Omnino ovata, valde convexa, haud nitida, pilis brevissimis nureis vestita, capite atque pronoto piceo-nigris, elytris rufescentibus; capite nigro, lato, antice rotundato, indistincte tuberculato, oculis parvis, haud prominenti-

bus; *pronoto* nitidulo, sat magno et lato, ad basim latissimo, nigro-piceo, angulis dilutioribus, tuberculis parvis, ordinibus transversis sinuatis dispositis, sparse ornato, interstitiis fortiter reticulatis, lateribus leviter marginatis, margine basali arcuata, angulis latis, valde productis; *elytris* brevibus, valde attenuatis, capite atque pronoto nec longioribus neque latioribus, ad humeros latissimis, confertissime sed indistincto asperatis, lateribus leviter marginatis, apicibus obtusis, angulis suturalibus haud rotundatis; *abdomine* longius exserto, rufo-piceo, alutaceo; pedibus atque antennis modicis, flavis.

Body perfectly ovate, very convex, not shining, clothed with very short golden hairs, head and thorax nigro-piceous, elytra rufescent.

Head black, broad, rounded in front, indistinctly tuberculate; eyes small, not prominent; antennae moderate, yellow.

Thorax moderately shining, rather large and broad, widest at the base, nigro-piceous, with the angles paler, ornamented with small tubercles in wavy transverse rows, interstices strongly reticulate, sides faintly margined, basal margin arcuate with the angles broad and much produced.

Scutellum large, triangular, closely asperate.

Elytra short, much attenuated posteriorly, not longer or broader than the head and thorax, widest at the shoulders, very closely but indistinctly asperate, sides faintly margined, extremities obtuse with the sutural angles not rounded.

Abdomen much exserted, rufo-piceous and alutaceous.

Legs moderate, yellow.

Under parts rufo-piceous, venter paler.

Differs from others in its perfectly oval form, and superficial sculpture; from *T. Hellenica*, to which it is closely allied, only in its darker colour and indistinct sculpture.

Habitat, Asia. Found in the Caucasus district by H. Leden (Reitter).

Trichopteryx lucida.

T. lucida, Matthews, Trans. Am. Ent. Soc. xi. p. 129, 1884; Biol. Cent.-Amer. Col. ii. pt. i. p. 135, 1888.

L. c. $\frac{7}{16}$ lin. = 0.87 mm.—Sat brevis et lata, postice valde attenuata, convexa, nitidissima, translucide castanea, pilis aureis, sat longis, vestita; *capite* sat magno, antice producto, minute et remote punctato; oculis magnis, valde prominentibus; *pronoto* sat magno, capite longiori et latiori, postice valde dilatato, ad basim latissimo, valde convexo, nitidissimo, tuberculis exiguis, remotis, interstitisque glabris, nitidissimis, ornato, lateribus leviter rotundatis, et late marginatis, margine basali parum arcuata, angulis acutis, valde productis, dilutioribus; *elytris* sat brevibus, postice valde attenuatis, capite atque pronoto angustioribus, haud longioribus, ad humeros latissimis, remote et irregulariter asperatis, lateribus fere rectis, apicibus latis, rotundatis, dilutioribus; abdomine parum exserto, testaceo; pedibus sat brevibus, lute flavis; antennis sat longis, gracillimis, pallide flavis.

Body rather short and broad, much attenuated posteriorly, convex, very shining, castaneous and translucent, clothed with rather long golden hair.

Head rather large, produced in front, minutely and remotely punctured; eyes large and very prominent; antennae rather long, very slender, pale yellow.

Thorax rather large, longer and broader than the head, much dilated posteriorly, widest at the base, very convex and very shining, ornamented with very minute, remote tubercles, interstices smooth and very shining, sides slightly rounded, and very broadly margined, basal margin rather arcuate with the angles acute and much produced, and pale.

Scutellum moderate, triangular, moderately asperated.

Elytra rather short, much attenuated posteriorly, narrower and not longer than the

- head and thorax, widest at the shoulders, remotely and irregularly asperated, sides nearly straight, extremities broad, rounded, and paler.
- Abdomen* not much exposed, testaceous.
- Legs* rather short, bright yellow.
- Under parts* entirely testaceous, metasternum dusky.
- Differs* from all other species in its very shining translucent surface, pale and slender antennæ, testaceous under parts, and in sculpture.
- Habitat*, Central America. Found in Cerro Zunil, Guatemala, at an elevation of four to five thousand feet, by Mr. Champion.

Trichopteryx rufescens.

T. rufescens, Matthews, Trans. Am. Ent. Soc. xi. p. 131, 1884; Biol. Cent.-Amer. Col. ii. pt. 1. p. 137, 1888.

L. c. $\frac{7}{16}$ lin. = 0.87 mm.—Sat lata et brevis, valde convexa, nitida, pilis brevibus flavis vestita, capite atque pronoto nigrescentibus, elytris rufescentibus; capite magno, tuberculis parvis, distinctis ornato; oculis magnis, haud prominentibus; pronoto permagno, postice valde dilatato, capite longiori et multo latiori, tuberculis distinctis, interstitiisque reticulatis ornato, ad basim latissimo, lateribus rotundatis et late marginatis, margine basali profunde sinuata, angulis latis, valde productis; elytris brevibus, postice valde attenuatis, rufescentibus, capite atque pronoto haud longioribus, sat angustioribus, ordinibus transversis confertim, sat leviter asperatis, lateribus rectis, apicibus valde rotundatis, late dilutioribus, extremis albidis; adomine modice exserto, rufo-piceo, apicem versus dilutiori; pedibus robustis, atque antennis gracilioribus, læte flavis.

Body short and rather broad, very convex, shining, clothed with short yellow hair, with the head and thorax blackish, and the elytra rufescent.

Head large, ornamented with small, distinct tubercles; eyes large and prominent; antennæ rather slender, bright yellow.

Thorax very large, much dilated at the base, longer and much broader than the head, widest at the base, ornamented with distinct tubercles, interstices reticulated, sides rounded and broadly margined, basal margin deeply sinuated with the angles broad and much produced.

Scutellum large, triangular, closely asperated.

Elytra short, much attenuated posteriorly, rufescent, not longer, but rather narrower than the head and thorax, closely but rather lightly asperated in transverse rows, sides straight, extremities broadly pale and much rounded, with the edge white.

Abdomen moderately exserted, rufo-piceous, paler towards the apex, apex faintly tridentate.

Legs robust, bright yellow, basal joints of all the tarsi dilated.

Under parts rufo-piceous, metasternum and gular region black, mouth and coxæ bright yellow.

Differs from others in its very convex form, posteriorly attenuated and rufescent elytra, bright yellow legs and antennæ, and in sculpture.

Habitat, Central America. Found in Guatemala and Panama by Mr. Champion.

Trichopteryx similior.

T. similior, Matthews, Trans. Am. Ent. Soc. xi. p. 130, 1884; Biol. Cent.-Amer. Col. ii. pt. i. p. 136, 1888.

L. c. $\frac{8}{16}$ lin. = 1 mm.—Oblonga, valde convexa, nitida, nigro-fusca, pilis flavis vestita; capite magno, antice rotundato, tuberculis remotis ornato; oculis magnis, haud prominentibus; pronoto magno, capite latiori et multo longiori, prope medium

latissimo, tuberculis sat magnis, ordinibus sinuatis, irregulariter dispositis, interstitiisque glabris, nitidis, ornato, lateribus sat late marginatis, margine flava, margine basali valde depressa, reflexa, et sinuata, angulis acutis, vix productis, flavis; *elytris* rufo-fuscis, brevibus, capite atque pronoto vix longioribus aut latioribus, prope media latissimis, postice parum contractis, ordinibus transversis modice asperatis, interstitiis nitidis, lateribus modice rotundatis, apicibus latis, rotundatis, extremis albidis; abdomine longius exserto, obtuso, apice tridentato; pedibus sat longis, late flavis; antennis longis, gracilibus, flavis, clavis fusciscentibus.

Body oblong, very convex, shining, brownish black, clothed with yellow hair.

Head large, rounded in front, remotely tuberculate; eyes large, not prominent: antennae long and slender, yellow with the club dusky.

Thorax large, broader and much longer than the head, widest near the middle, ornamented with rather large tubercles in irregular wavy rows, interstices smooth and shining, sides broadly margined, margin yellow, basal margin much depressed, reflexed and sinuated with the angles acute, scarcely produced, yellow.

Scutellum large, triangular, rugosely asperated.

Elytra rufo-fuscous, not longer or broader than the head and thorax, widest near the middle, slightly contracted posteriorly, moderately asperate in transverse rows, interstices shining, sides moderately rounded, extremities broad, rounded, with the edge white.

Abdomen rather long, obtuse, tridentate at the apex.

Legs long, bright yellow.

Under parts black, with the mouth and coxae yellow.

Differs from *T. discolor* in its smaller size, rufo-fuscous elytra, and in sculpture.

Habitat, Central America. Found in Guatemala by Mr. Champion.

Trichopteryx occidentalis.

T. occidentalis, Matthews, Ann. Mag. Nat. Hist. 6, xiii. 1894, p. 339.

L. e. $\frac{6}{16}$ — $\frac{7}{15}$ lin. = 0.76–0.87 mm. —Ovata, latissima, convexa, postice attenuata, capite atque pronoto nigris, elytris fuscis, pilis flavescentibus dense vestita; *capite* magno, lato, oculis modicis haud prominentibus, antennis longis, gracilibus, flavis; *pronoto* magno, ad basim valde dilatato, nigro, tuberculis modicis ordinibus irregularibus confertim dispositis, interstitiisque nitidis, leviter reticulatis ornato, lateribus leviter marginatis, margine basali leviter sinuata, angulis latis, valde productis, plus minusve rufescentibus; *elytris* fuscis, brevibus, capite atque pronoto sat angustioribus, multo brevioribus, ad humeros latissimis, profunde et confertim asperatis, interstitiis reticulatis, lateribus marginatis, fere rectis, apicibus leviter rotundatis; abdomine modice exserto, rufescenti; pedibus modicis, flavis.

Body ovate, very broad, convex, attenuated posteriorly, head and thorax black, elytra fuscous, thickly clothed with flavescent hair.

Head large and broad: eyes moderate, not prominent: antennae long and slender, yellow.

Thorax large, much dilated at the base, black, ornamented with moderate tubercles closely placed in irregular rows with the interstices shining and faintly reticulated, sides slightly margined, basal margin faintly sinuated with the angles broad, much produced, more or less rufescent.

Scutellum moderate, triangular, asperated.

Elytra fuscous, short, rather narrower and much shorter than the head and thorax, widest at the shoulders, attenuated towards the apex, deeply and closely asperated, with the interstices reticulated, sides margined, nearly straight, extremities slightly rounded.

Abdomen moderately exerted, rufescent.

Legs moderate, yellow.

Under parts castaneous, more or less dusky.

Differs from *T. levicollis* in its much more depressed form, less shining surface, deep and close sculpture, long antennæ, and thick pubescence.

Habitat, America. Found in profusion by Mr. Smith in the West Indian Islands.

Trichopteryx Godmanni.

T. Godmanni, Matthews, Trans. Am. Ent. Soc. xi. p. 128, 1884.

T. godmani, Matth., Biol. Cent.-Amer. Col. ii. pt. i. p. 134, pl. iii. fig. 20, 1888.

L. c. $\frac{1}{16}$ –1 lin. = 1.87–2 mm.—Latissima, convexissima, aterrima, nitidissima, pilis brevibus argenteis vestita; *capite* sat parvo, antice acuminato, fere glabrò; *oculis* parvis; *pronoto* permagno, valde convexo, ad basim latissimo, tuberculis minutis, ordinibus irregularibus dispositis, interstitiisque confertim reticulatis, ornato, lateribus modice rotundatis et leviter marginatis, marginibus atque angulis posterioribus dilutioribus, margine basali sinuata, angulis acutis, sat productis; *elytris* sat brevibus, postice attenuatis, capite atque pronoto parum angustioribus, fere pariter longis, ad humeros latissimis, densius pubescentibus, ordinibus transversis, fere rectis, leviter asperatis, lateribus fere rectis, apicibus latis, leviter rotundatis, dilutioribus, extremis albidis; abdomine sat longe exserto, apice bidentato; pedibus robustis, late flavis, tibiis anterioribus et articulo basali tarsorum leviter dilatatis; antennis modicis, pallide piceis.

Body very broad and very shining, deep black, very convex, sparingly clothed with silvery hairs.

Head rather small, acuminate in front, almost smooth; eyes small; antennæ moderate, pale piceous.

Thorax very large and very convex, widest at the base, ornamented with minute tubercles in close irregular rows, interstices closely and finely reticulated, sides moderately rounded and slightly margined, margin and posterior angles pale, basal margin sinuated with the angles acute and moderately produced.

Scutellum rather large, triangular, closely and finely asperate.

Elytra rather short, attenuated posteriorly, as long, but rather narrower than, the head and thorax, widest at the shoulders, more densely pubescent, finely asperated in almost straight transverse rows, sides nearly straight, extremities broad and slightly rounded with the edge white.

Abdomen considerably exposed, obtuse, and bidentate at the apex.

Legs robust, bright yellow, with the anterior tibiæ and basal joint of the tarsi slightly dilated.

Under parts black.

Differs from others in its great size, and very broad, convex form.

Habitat, Central America. Found near Ballhue in Vera Paz by Mr. Champion.

I have given a portrait of *T. Godmanni*, since the figure of this species in the 'Biologia Centrali-Americana' does not in any way express either the form or character of the insect.

Trichopteryx grossa.

Acratrichis grossa, Motschulsky, Bull. Mosc. xli. 2, p. 177.

T. grossa, Matthews, Trans. Am. Ent. Soc. xi. p. 128, 1884; Biol. Cent.-Amer. Col. ii. pt. i. p. 134, 1888.

L. c. $\frac{8}{16}$ — $\frac{9}{16}$ lin. = 1-1.12 mm.—Latissima, valde convexa, nigra, nitida, pilis griseis parce vestita; *capite* magno, lato, antice producto, fere glabro; *pronoto* permagno, valde convexo, nitidissimo, postice valde dilatato, punctis minutis, remotis, distincte impresso, interstitiis glabris, nitidis, lateribus rotundatis, sat late marginatis, atque seta erecta prope basim utrinque instructis, margine basali fere recta, angulis latis, valde productis, dilutioribus; *elytris* brevibus, postice valde attenuatis, fusciscentibus, capite atque pronoto brevioribus et parum angustioribus, ordinibus transversis, sinuatis, leviter et confertim asperatis, lateribus fere rectis, seta brevi, erecta, prope humeros atque altera basim versus utrinque instructis, apicibus latis, fere rectis, dilutioribus; abdomine modice exserto, apice profunde tridentato; pedibus robustis, obscure flavis; antennis longis, gracilibus, piceo-testaceis.

Body very broad and very convex, black, shining, sparingly clothed with grey hair. *Head* large and broad, produced in front, nearly smooth; eyes rather large, not prominent; antennae long and slender, pitchy-testaceous.

Thorax very large, very convex, and very shining, much dilated at the base, minutely, remotely, and distinctly punctured with the interstices smooth and shining, sides rounded, rather broadly margined, with an erect seta on each side near the base, basal margin nearly straight with the angles broad, much produced, and pale.

Scutellum very large and broad, finely asperated.

Elytra short, much attenuated posteriorly, fusciscent, shorter and rather narrower than the head and thorax, finely and closely asperated in transverse, wavy rows, sides nearly straight, with a short, erect seta on each side near the shoulders, and another towards the apex, extremities broad, nearly straight, and pale.

Abdomen moderately exserted, strongly tridentate at the apex.

Legs robust, dusky yellow.

Under parts black with the mouth and coxae piceous.

Differs from others in its large size, short, convex form, large, shining, and minutely punctured thorax, long antennae, and erect lateral setae.

Habitat, North and Central America. Found in Washington, and by Mr. Champion in Vera Paz, Guatemala.

Col. Motschulsky's description of *A. grossa* agrees with this species; his specimens were found in North America.

Trichopteryx fratercula.

T. fratercula, Matthews, Ent. Mo. Mag. xv. p. 65, 1878.

L. c. $\frac{6}{16}$ — $\frac{7}{16}$ lin. = 0.75-0.87 mm.—Sat brevis, lata, nigra, nitida, pilis brevibus fulvis vestita; *capite* magno, lato, tuberculis minutis, remotis, ordinibus fere rectis ornato, interstitiis nitidis, reticulatis; oculis magnis, sat prominentibus; *pronoto* magno, capite longiori, et multo latiori, ante basim latissimo, valde convexo, confertissime reticulato sive alutaceo, tuberculis minutis, remotis, ordinibus sinuatis, ornato, lateribus rotundatis et late marginatis, margine basali leviter sinuata, angulis latis, valde productis, atque ad modum *Fraterculæ arcticæ* rostri externe dilatatis; *elytris* brevibus, postice parum attenuatis, capite atque pronoto fere pariter longis, parum angustioribus, ordinibus transversis, sat remotis, modice

asperatis, apicibus latis, rotundatis, flavescens, extremis albidis; abdomine longe exserto, lato, obtuso, apice leviter tridentato; pedibus brevibus, robustis, flavis, femoribus obscuratis; antennis brevibus atque palpis nigris.

Body rather short and broad, black, shining, clothed with short fulvous hair.

Head large and broad, ornamented with minute remote tubercles in nearly straight rows, interstices reticulated and shining; eyes large, rather prominent; antennae short, pitchy black; palpi black.

Thorax large, longer and much broader than the head, widest before the base, very convex, very closely reticulate or alutaceous, ornamented with minute remote tubercles in distant wavy rows, sides rounded and broadly margined, basal margin slightly sinuated, with the angles much produced, very broad, and rounded on the outer side in the shape of the bill of the Puffin, *Fratercula arctica*.

Scutellum large, deeply asperated in transverse rows.

Elytra short, slightly attenuated posteriorly, as long as the head and thorax, but narrower, moderately asperated in remote transverse rows, sides slightly rounded, extremities broad, rounded, flavescens, with the edge white.

Abdomen much exserted, broad, obtuse, slightly tridentate at the apex.

Legs short, robust, yellow, with the femora dusky.

Under parts wholly black.

Differs from T. Henrici in its more convex and narrower form, yellow legs, and in sculpture; and from all others in the peculiar dilatation of the posterior angles of the thorax.

Habitat, Europe. Found by myself in Leicestershire.

Trichopteryx angustipennis.

T. angustipennis, Matthews, Trans. Am. Ent. Soc. xi. p. 132, 1884; Biol. Cent.-Amer. Col. ii. pt. i. p. 139, 1888.

L. c. $\frac{5}{16}$ lin. = 1 mm.—Suboblonga, thoracis lateribus valde dilatatis, nitida, valde convexa, piceo-nigra, pilis flavis vestita; capite magno, oculis magnis, haud prominentibus; pronoto magno, ad latera valde dilatato, prope medium latissimo, tuberculis magnis, ordinibus irregularibus dispositis, interstitiisque glabris nitidis ornato, lateribus valde rotundatis et fortiter marginatis, margine basali depressâ, sinuata et reflexâ, angulis flavis, acutis, vix productis; elytris fuscis, sat brevibus, capite atque pronoto haud longioribus, multo angustioribus, prope media latissimis, ordinibus transversis sat profunde asperatis, apicibus parum contractis et parum rotundatis, extremis albidis; abdomine longius exserto, obtuso; pedibus atque antennis flavis.

Body suboblong, with the sides of the thorax largely dilated, shining, very convex, nigro-piceous, clothed with yellow hair.

Head very large, moderately tuberculate, shining; eyes large, not prominent; antennae long and slender, yellow, with the apex dusky.

Thorax rather large, wider and much longer than the head, dilated at the sides, widest near the middle, ornamented with rather large tubercles in irregular rows, with the interstices smooth and shining, much depressed towards the base, sides strongly margined, with the margin yellow, basal margin sinuated and reflexed, with the angles acute, scarcely produced, yellow.

Scutellum large, triangular, rugosely asperate.

Elytra rufous brown, short, not longer but much narrower than the head and thorax, widest near the middle, rather contracted towards the apex, moderately asperate in transverse rows, with the interstices shining, extremities broad, not much rounded, with the edge white.

Abdomen much exposed, with the apex obtuse, faintly tridentate.

Legs rather long, bright yellow.

Under parts black, with the mouth and coxæ yellow.

Differs from *T. crassicollis* in its smaller size, depressed base of the thorax, short elytra, and general sculpture.

Habitat, Central America. Found in Guatemala by Mr. Champion.

Trichopteryx truncatissima.

T. truncatissima, Matthews, Ann. Mag. Nat. Hist. 6, iii. 1889, p. 192.

L. c. vix $\frac{6}{16}$ lin. = 0.75 mm.—Brevissima, postice validissime attenuata, convexa, rugose exsculpta, nigra, pilis fulvis dense vestita; *capite* magno, sat fortiter tuberculato; oculis modicis, haud prominentibus; *pronoto* magno, ad basim latissimo, ordinibus irregularibus rugose tuberculato, interstitiis reticulatis, lateribus rotundatis et marginatis, margine basali fere recta, angulis latis, longe productis; *elytris* perbrevibus, fuscescentibus, postice validissime attenuatis, capite atque pronoto multum angustioribus et multum brevioribus, ad humeros latissimis, ordinibus transversis profunde et confertim asperatis, lateribus rectis et marginatis, apicibus angustis, valde rotundatis, dilutioribus, extremis albidis; abdomine nigro, longe exserto; pedibus robustis, late flavis; antennis modicis, gracilibus, obscure flavis.

Body very short, exceedingly attenuated posteriorly, convex, rugosely sculptured, black, thickly clothed with fulvous hair.

Head large, strongly tuberculated; eyes moderate, not prominent; antennæ moderate, slender, dusky yellow.

Thorax very large, widest at the base, rugosely tuberculated in irregular rows, the interstices reticulated, sides rounded and margined, basal margin nearly straight, with the angles broad and much produced.

Scutellum large, triangular, and deeply asperated.

Elytra fuscescent, very short, exceedingly attenuated posteriorly, much shorter and much narrower than the head and thorax, widest at the shoulders, deeply and closely asperated in transverse rows, sides straight and margined, extremities narrow, much rounded and paler, with the edges white.

Abdomen much exserted, black.

Legs robust, bright yellow.

Under parts entirely black.

Differs from all other species in its very short form, large thorax, very small and excessively attenuated elytra, and in its very rugose sculpture.

Habitat, South America. Found in Brazil, near Blumenau.

This is a very distinct and remarkable species, easily recognised by its short and excessively attenuated elytra.

Trichopteryx minor.

T. minor, Matthews, Trans. Am. Ent. Soc. xi. p. 134, 1884; Biol. Cent.-Amer. Col. ii. pt. i. p. 140, 1888.

L. c. $\frac{7}{16}$ lin. = 0.87 mm.—Oblongo-ovalis, convexa, nitidula, nigra, pilis luteis dense vestita; *capite* modico, antice producto, tuberculis minutis remote ornato; oculis modicis, haud prominentibus; *pronoto* sat magno, prope medium latissimo, capite longiori et latiori, tuberculis magnis, ordinibus sinuatis dispositis, interstitiisque leviter reticulatis, ornato, lateribus marginatis, modice rotundatis, et seta erecta ante media utrinque instructis, margine basali parum reflexa, sinuata, angulis valde productis, dilutioribus; *elytris* sat brevibus, quadratis, fuscescentibus, capite atque pronoto nec longioribus, neque latioribus, ordinibus transversis confertim et

profunde asperatis, lateribus vix rotundatis, seta erecta prope humeros atque altera apices versus utrinque instructis, apicibus latis, valde rotundatis, extremis albidis; abdomine sat longo, fusciscenti, apice bidentato; pedibus robustis, late flavis; antennis longis, gracilibus, flaviscentibus.

Body oblong-oval, convex, rather shining, black, thickly clothed with luteous hair.

Head moderate, produced in front, ornamented with minute remote tubercles; eyes moderate, not prominent; antennae long and slender, flavescens.

Thorax rather large, widest near the middle, longer and broader than the head, ornamented with large tubercles in wavy rows, interstices faintly reticulated, sides margined, moderately rounded, and furnished with an erect seta before the middle, basal margin slightly reflexed and sinuated, with the angles much produced and pale.

Scutellum large, triangular, deeply and closely asperate.

Elytra rather short, quadrate, fusciscent, not longer or broader than the head and thorax, closely and deeply asperate in transverse rows, sides scarcely rounded, with an erect seta on each side near the shoulders, and another towards the apex, extremities broad, much rounded, with the edge white.

Abdomen rather long, fusciscent, apex bidentate.

Legs robust, bright yellow.

Under parts black, with the mouth and coxae bright yellow.

Differs from T. grandicollis in its much smaller size, oblong form, pale antennae, and deeper sculpture.

Habitat, Central America. Found plentifully in Guatemala by Mr. Champion.

Trichopteryx Lætitia.

T. Lætitia, Matthews, Ent. Mo. Mag. ix. p. 180, 1873.

L. c. $\frac{6}{16}$ – $\frac{7}{16}$ lin. = 0.75–0.87 mm.—Oblong-ovalis, modice convexa, nigra, haud nitida, pilis pallidis dense vestita; capite modico, antice producto, tuberculis minutis, interstitiisque nitidis, ornato; oculis magnis, prominentibus; pronoto sat brevi, leviter convexo, capite longiori et latiori, ad basim latissimo, tuberculis minutis, ordinibus transversis, sinuatis, confertim dispositis, interstitiisque sat profunde reticulatis ornato, lateribus marginatis et modice rotundatis, marginibus dilutioribus, margine basali sat profunde sinuata, angulis acutis, longe productis; elytris oblongis, postice plus minusve attenuatis, capite atque pronoto vix longioribus aut latioribus, ordinibus transversis modice et confertim asperatis, lateribus marginatis, fere rectis, apicibus valde rotundatis, dilutioribus, summis albidis; abdomine longius exserto, segmentorum marginibus dilutioribus, apice tridentato; pedibus modicis, obscure flavis, articulo basali tarsorum anticorum dilatato; antennis brevibus, piceo-nigris.

Body oblong-oval, moderately convex, black, not shining, thickly clothed with pale hair.

Head moderate, produced in front, ornamented with minute tubercles, interstices shining; eyes large and prominent; antennae short and slender, pitchy black; maxillary palpi black.

Thorax rather short, moderately convex, longer and broader than the head, widest at the base, ornamented with minute tubercles in transverse wavy rows, with the interstices rather deeply reticulated, sides margined and moderately rounded, with the margin pale, basal margin rather deeply sinuated, with the angles acute and much produced.

Scutellum large, acuminate, closely asperated.

Elytra oblong, more or less attenuated posteriorly, scarcely longer or broader than the head and thorax, moderately and closely asperated in transverse rows,

sides margined, nearly straight, extremities much rounded, paler, with the edge white.

Abdomen rather long, obtusely conical, black, with edges of the segments paler, apex obtusely tridentate.

Legs moderate, dusky yellow, with the basal joint of the anterior tarsi dilated; femora dusky.

Under parts black, with the mouth and coxæ piceous.

Differs from *T. fascicularis* in its smaller size, and smaller and less convex thorax, and shorter pitchy-black antennæ; from *T. Cantiana* in its larger size, attenuated form, and depressed thorax.

Habitat, Europe. Found in Belgium by Miss L. Matthews, and also in England.

Trichopteryx cognata.

T. cognata, Matthews, Cist. Ent. ii. p. 176, 1877.

L. c. $\frac{9}{16}$ lin. = 1.12 mm.—Ovalis, sat angusta, convexa, nigra, pilis brevissimis, pallidis vestita; *capite* sat parvo, tuberculis exiguis confertim ornato; oculis prominulis; *pronoto* parvo, valde convexo, capite parum longiori et latiori, ad basim latissimo, tuberculis minutis, ordinibus irregularibus confertim dispositis, interstitiisque dense reticulatis, ornato, lateribus leviter rotundatis, margine basali depressa, leviter trisinuata, angulis longe productis, extremis dilutioribus; *clytris* oblongis, attenuatis capite atque pronoto nec longioribus nec latioribus, ordinibus densis, leviter sinuatis confertim asperatis, lateribus leviter rotundatis, apicibus attenuatis, valde rotundatis, dilutioribus, extremis albidis; abdomine longe exserto, obtuso, apice minute tridentato; pedibus robustis, femoribus infuscatis, tibiis hete flavis; antennis gracilibus, piceo-testaceis.

Body oval, rather narrow, convex, black, clothed with very short pale hair.

Head rather small, thickly ornamented with minute tubercles; eyes rather prominent; antennæ slender, pitchy-testaceous.

Thorax small, very convex, rather longer and broader than the head, widest at the base, ornamented with minute tubercles in close irregular rows, with the interstices densely reticulated, sides slightly rounded, basal margin depressed, faintly trisinuated, with the angles much produced, pale at the apex.

Scutellum large, acuminate, closely asperated.

Elytra oblong, rather attenuated, not longer or broader than the head and thorax, asperated in close wavy rows, sides slightly rounded, extremities attenuated, much rounded, paler, with the edge white.

Abdomen much exserted, obtuse, minutely tridentate at the apex.

Legs robust, femora dusky, tibiæ bright yellow.

Under parts black, mouth fuscous, metasternum deeply reticulated.

Differs from *T. fascicularis* in its smaller size, narrower form, smaller head and thorax, and piceous antennæ.

Habitat, North America. Found in British Columbia by Mr. G. R. Crotch.

Trichopteryx seminitens.

T. seminitens, Matthews, Ent. Mo. Mag. xiv. p. 36, 1877.

L. c. $\frac{1}{8}$ lin. = 1 mm. Oblongo-ovalis, postice attenuata, valde convexa, nigra, nitidissima, pilis pallidis sparse induta; *capite* magno, antice producto, minute et remote tuberculato; oculis magnis, prominentibus; *pronoto* magno, valde convexo, nitidissimo, capite multo longiori et latiori, valde dilatato, ad basim latissimo, tuberculis sat magnis ordinibus remotis, sinuatis, dispositis, interstitiisque nitidissimis, levissime reticulatis, ornato, lateribus marginatis et leviter rotundatis, mar-

gine basali depressa, dilutiori, sat profunde sinuata, angulis latis, valde productis; *clytris* brevibus, postice valde attenuatis, capite atque pronoto multum brevioribus, sat angustioribus, ordinibus remotis, sinuatis, sat profunde asperatis, interstitiis nitidis, lateribus fere rectis, apicibus latis, leviter rotundatis, anguste dilutioribus, extremis albidis; abdomine longe exserto, sat obtuso, apice minute bidentato; pedibus robustis, sat brevibus, obscure flavis; antennis sat brevibus, piceis.

Body oblong-oval, attenuated posteriorly, very convex, black, very shining, sparingly clothed with pale hair.

Head large, produced in front, minutely and remotely tuberculate; eyes large and prominent; antennae rather short, piceous.

Thorax large, very convex and very shining, much longer and broader than the head, much dilated, widest at the base, ornamented with rather large tubercles in remote wavy rows, interstices very shining and faintly reticulated, sides margined and slightly rounded, basal margin depressed, pale, and deeply sinuated, with the angles broad, very much produced.

Scutellum large and broad, deeply and closely asperated.

Elytra short, much attenuated posteriorly, narrower and much shorter than the head and thorax, rather deeply asperated in remote wavy rows, interstices shining, sides nearly straight, extremities broad, slightly rounded, narrowly pale, with the edge white.

Abdomen long, rather obtuse, with the apex minutely bidentate.

Legs rather short, robust, dusky yellow.

Under parts black, with the mouth and coxae yellow.

Differs from *T. attenuata* in its larger size, much more convex form, shorter antennae, and the sculpture of the thorax; from *T. fascicularis* in its more convex form, and its very shining and remotely tuberculated thorax.

Habitat, Europe. Not common.

Trichopteryx Championis.

T. Championis, Matthews, Ent. Mo. Mag. xv. p. 64, 1878.

L. c. $\frac{7}{16}$ lin. = 0.87 mm.—Angusta, elongata, postice valde attenuata, convexa, obscure castanea, nitida, pilis fulvis parce vestita; capite sat parvo, antice rotundato, indistincte tuberculato; oculis parvis, haud prominentibus; pronoto modico, valde convexo, capite longiori et latiori, ad basim latissimo, tuberculis minutis, ordinibus sinuatis, indistinctis, dispositis, interstitisque nitidissimis, leviter reticulatis, ornato, lateribus rotundatis et late marginatis, margine basali arcuata, angulis longe productis, his atque margine tota flavis; *clytris* brevibus, angustis, valde attenuatis, capite atque pronoto fere pariter longis, parum angustioribus, ordinibus transversis, sinuatis, leviter asperatis, lateribus marginatis et leviter rotundatis, apicibus leviter rotundatis, dilutioribus; abdomine obtuso, longe exserto, apice minute tridentato; antennis longis atque pedibus hae flavis.

Body narrow, elongate, much attenuated posteriorly, convex, dark castaneous, shining, sparingly clothed with fulvous hair.

Head rather small, rounded in front, indistinctly tuberculate; eyes small, not prominent; antennae long, bright yellow.

Thorax moderate, very convex, longer and broader than the head, widest at the base, ornamented with minute tubercles in indistinct wavy rows, the interstices very shining and slightly reticulated, sides rounded and broadly margined, basal margin arcuate, with the angles much produced, the angles and whole margin flavescent.

Scutellum very large, rather deeply asperated.

Elytra short, narrow, much attenuated posteriorly, nearly as long but narrower

than the head and thorax, slightly asperated in transverse wavy rows, sides margined and slightly rounded, extremities slightly rounded and pale.

Abdomen elongate, obtuse, with five segments partly exposed, minutely tridentate at the apex.

Legs moderate, bright yellow.

Under parts dusky castaneous, with the margins of all the segments pale.

Differs from others in its narrow and very convex form, castaneous colour, and in sculpture.

Habitat, Europe. Found in Wicken fen, Cambridgeshire, by Mr. Downing.

Trichopteryx angusta.

T. angusta, Matthews, Ann. Mag. Nat. Hist. 6, iii. 1889, p. 192.

L. c. $\frac{6}{16}$ – $\frac{7}{16}$ lin. 0·75–0·87 mm.—Elongata, valde angusta, convexa, nitida, nigra, pilis fulvis parce vestita: *capite* magno, prominenti, fere glabro; oculis modicis, laud prominentibus; *pronoto* parvo, capite latiori et parum longiori, ad basin latissimo, modice dilatato, indistincte et confertim tuberculato, nitido, lateribus leviter marginatis et leviter rotundatis, margine basali fere recta, angulis productis, acutissimis; *elytris* fusciscentibus, perbrevibus, parum attenuatis, capite atque pronoto brevioribus atque angustioribus, ad humeros latissimis, ordinibus transversis, regularibus, sat profunde asperatis, lateribus ovaliter rotundatis, et leviter marginatis, apicibus obtusis, vix rotundatis; abdomine valde elongato, segmentis quinque expositis, nigro, leviter asperato, apice obtuso, integro; pedibus late, antennis obscure flavis.

Body elongate, very narrow, convex and shining, black, sparingly clothed with fulvous hair.

Head large and prominent, almost smooth; eyes moderate, not prominent; antennæ dusky yellow.

Thorax small, moderately dilated, shining, broader and rather longer than the head, widest at the base, indistinctly and closely tuberculate, sides slightly rounded and margined, basal margin nearly straight, with the angles produced and very acute.

Scutellum large, triangular, rather deeply asperate.

Elytra fusciscent, very short, slightly attenuated, shorter and narrower than the head and thorax, widest at the shoulders, rather deeply asperated in regular transverse rows, sides ovally rounded, and slightly margined, extremities obtuse, scarcely rounded.

Abdomen elongate, with five segments exposed, black and slightly asperated.

Legs moderate, bright yellow.

Under parts black, with the mouth and coxæ yellow.

Differs from other species in its intermediate size, long and narrow form, very short elytra, and long abdomen.

Habitat, Europe. Three examples of this species were found by myself near Gumley, Leicestershire.

Trichopteryx Brasiliensis.

T. Brasiliensis, Matthews, Ann. Mag. Nat. Hist. 6, iii. 1889, p. 192.

L. c. $\frac{5}{16}$ lin. = 0·75 mm. — Suboblonga, convexa, nitida, fortiter calata, fusciscent, pilis fulvis vestita: *capite* magno, lato, nigro, modice tuberculato; oculis parvis, laud prominentibus; *pronoto* sat magno, ante basin latissimo, nigrescenti, profunde et confertim tuberculato, interstitiis nitidis, lateribus marginatis, valde rotundatis, atque seta nigra, erecta pone media instructis, margine

basali sinuata, angulis latis, productis: *elytris* brevibus, quadratis, rufo-fuscis, capite atque pronoto multum brevioribus, et parum angustioribus, ordinibus transversis profunde et confertim asperatis, lateribus rectis, marginatis, seta brevi, nigra, erecta, prope humeros, atque altera prope apices instructis, apicibus fere rectis; abdomine sat longe exserto, rufo-fusco, et asperato, apice obtuso, integro; pedibus sat robustis, late flavis; antennis modicis, obscure flavis.

Body suboblong, convex, shining, deeply sculptured, more or less fuscous, clothed with fulvous hair.

Head large and broad, black, moderately tuberculate; eyes small, not prominent; antennæ moderate, dusky yellow.

Thorax rather large, widest before the base, nearly black, deeply and closely tuberculate, with the interstices shining, sides margined and much rounded, furnished with an erect black seta behind the middle, basal margin sinuated, with the angles broad and produced.

Scutellum very large, elongate-triangular, deeply asperated.

Elytra short, quadrate, rufo-fuscous, much shorter, and rather narrower than the head and thorax, deeply and closely asperate in transverse rows, sides straight, margined, and furnished with a short black erect seta near the shoulders, and another near the extremities, extremities nearly straight.

Abdomen considerably exposed, rufo-fuscous, and asperated, apex entire, obtuse.

Legs rather robust, bright yellow.

Under parts rufo-fuscous.

Differs from others in its short quadrate elytra, rufescent abdomen, and short erect lateral setæ, which are sometimes lost: it resembles *T. atomaria*, but is narrower and more deeply sculptured.

Habitat, South America. Found in Brazil, near Blumenau.

Trichopteryx inconspicua.

T. inconspicua, Matthews, Cist. Ent. i. p. 296, 1874.

L. c. $\frac{5}{16}$ — $\frac{6}{16}$ lin. = 0.63–0.75 mm.—Brevis, convexa, subquadrata, postice parum attenuata, nigra, nitida, pilis brevibus griseis sparse vestita; capite magno, lato, antice sat producto, tuberculis minutis, remotis ornato: oculis parvis, haud prominentibus; pronoto sat magno, capite parum longiori, multo latiori, postice in maribus valde dilatato, valde convexo, tuberculis parvis, ordinibus sinuatis, regulariter dispositis, interstitiisque nitidissimis, indistincte reticulatis ornato, lateribus rotundatis, leviter marginatis, margine basali fere recta, angulis acutis, dilutioribus, in maribus longe productis; *elytris* subquadratis, capite atque pronoto haud longioribus, parum angustioribus, in maribus postice attenuatis, ordinibus regulariter sinuatis, confertim, haud profunde asperatis, sutura sat elevata, lateribus leviter rotundatis, apicibus latis, parum rotundatis, dilutioribus; abdomine modice exserto, apice minute bidentato; pedibus late flavis, articulo basali tarsorum anticorum in maribus dilatato; antennis modicis, dilute piceis.

Body short, convex, subquadrata, rather attenuated posteriorly, black and shining, sparingly clothed with short grey hair.

Head large and broad, rather produced in front, ornamented with minute remote tubercles: eyes small, not prominent; antennæ moderate, pale piceous.

Thorax rather large, rather longer and much broader than the head, very much dilated at the base in the males, very convex, ornamented with small tubercles in regular sinuated rows, interstices very shining and indistinctly reticulated, sides slightly rounded and margined, basal margin nearly straight, with the angles acute, pale, and much produced in the males.

Scutellum large and broad, closely asperated.

Elytra subquadrate, rather narrower, not longer than the head and thorax, attenuated posteriorly in the males, closely but not deeply asperated in regular wavy rows, sides slightly rounded, suture elevated, extremities broad, slightly rounded and pale.

Abdomen moderately exerted, minutely bidentate at the apex.

Legs moderate, bright yellow, basal joint of the anterior tarsi dilated in the male.

Under parts black, with the coxæ yellow, and the mouth and extremity of the venter piecous, metasternum rather deeply asperated.

Differs from *T. anthracina* in its smaller size, closer and much smaller tubercles of the thorax, and paler antennæ.

Habitat, Australasia. Found plentifully in New Zealand by Mr. Lawson. The females are more oblong in form, and their thorax is scarcely dilated at the base.

Trichopteryx glauca.

T. glauca, Matthews, Trans. Am. Ent. Soc. xi. p. 131, 1884; Biol. Cent.-Amer. Col. ii. pt. i. p. 137, 1888.

L. c. $\frac{5}{16}$ lin. = 0.75 mm.—Suboblonga, sat brevis, convexa, maribus postice attenuata, fusco-castanea, pilis brevibus, glaucis, valde sericeis, dense vestita: *capite* magno, lato, antice rotundato, sat confertim tuberculato: oculis modicis, haud prominentibus: *pronoto* sat magno, capite longiori et latiori, maribus postice dilatato atque ad basim latissimo, femineis ante basim latissimo, tuberculis ordinibus sinuatis confertim dispositis, interstitiisque latis, nitidis ornato, lateribus leviter rotundatis, et leviter marginatis, margine basali plus minusve sinuata, angulis acutis, in maribus valde productis; *elytris* subquadratis, maribus postice attenuatis, capite atque pronoto parum angustioribus, fere pariter longis, confertim asperatis, lateribus fere rectis, apicibus latis, leviter rotundatis, dilutioribus, extremis albidis: abdomine longius exserto, ferrugineo, apice bidentato: pedibus robustis, flavis; antennis gracilibus, flavis.

Body suboblong, rather short, convex, attenuated posteriorly in the males, fusco-castaneous, thickly covered with very glossy short glaucous hair.

Head large and broad, rounded in front, rather closely tuberculate; eyes moderate, not prominent: antennæ slender, yellow.

Thorax rather large, longer and broader than the head, dilated posteriorly, and widest at the base in the males, widest before the base in the females, ornamented with close tubercles in sinuated rows, with the interstices broad and shining, sides slightly rounded and slightly margined, basal margin more or less sinuated, with the angles acute, and much produced in the males.

Scutellum large, triangular, deeply and closely asperated.

Elytra subquadrate, attenuated posteriorly in the males, rather narrower, not longer than the head and thorax, deeply and very closely asperated, sides nearly straight, extremities broad, slightly rounded, paler, with the edge white.

Abdomen rather long, ferruginous, bidentate at the apex.

Legs robust, yellow.

Under parts castaneous, with the mouth and coxæ yellow.

Differs from other species in size and form, rufescent colour, slender antennæ, close sculpture, and especially in its dense silky glaucous pubescence.

Habitat, Central America. Found near Bugaba, Panama, by Mr. Champion.

There is so much difference in the outline of the sexes of this very distinct species that the female, if seen alone, might be supposed to belong to the "*sericans*" group, in which the posterior angles of the thorax are not produced.

Trichopteryx Darwinii.

T. Darwinii, Matthews, Ann. Mag. Nat. Hist. 6, iii. 1889, p. 193.

L. c. $\frac{5}{16}$ lin. = 0.63 mm.—Oblonga, convexa, nitidula, modice tuberculata, tota castanea, pilis sat longis aureis vestita; *capite* antice rotundato, minute et remote tuberculato, interstitiis glabris, nitidis, oculis parvis, haud prominentibus; *pronoto* sat brevi, capite longiori et latiori, ad basim latissimo, tuberculis parvis, remotis, ordinibus transversis irregulariter dispositis, interstitiisque glabris, nitidis, ornato, lateribus sat late marginatis, margine basali fere rectâ, angulis acutis, vix productis; *elytris* sat brevibus, quadratis, capite atque pronoto vix latioribus aut longioribus, apices versus latissimis, ordinibus transversis, fere rectis, modice asperatis, interstitiis glabris, nitidis, lateribus marginatis et leviter rotundatis, apicibus fere rectis; abdomine modice exserto; pedibus flavis; antennis gracillimis, pallide flavis.

Body oblong, convex, rather shining, moderately tuberculate, entirely castaneous, clothed with rather long golden hair.

Head rounded in front, minutely and remotely tuberculate, with the interstices smooth and shining; eyes small, not prominent; antennæ very slender, pale yellow.

Thorax rather short, longer and broader than the head, widest at the base, ornamented with small remote tubercles irregularly arranged in transverse rows, interstices smooth and shining, sides strongly margined, basal margin nearly straight, with the angles acute, scarcely produced.

Scutellum very large, triangular, deeply asperate.

Elytra rather short, subquadrate, scarcely longer or broader than the head and thorax, widest near the apex, moderately asperate in nearly straight transverse rows, interstices smooth and shining, sides margined and slightly rounded, extremities nearly straight.

Abdomen moderately exposed.

Legs moderate, yellow.

Under parts castaneous, mouth and apical segments of the venter paler; metasternum deeply reticulated.

Differs from *T. brevis* in its broader form, shorter thorax, very slender pale yellow antennæ, and very different sculpture.

Habitat, South America. Found in a fungus by Mr. Darwin near Rio Janeiro.

Trichopteryx reticulata.

T. reticulata, Matthews, Ann. Mag. Nat. Hist. 6, iii. 1889, p. 193.

L. c. $\frac{10}{6}$ lin. = 1.25 mm.—Brevis et lata, valde convexa, nitida, elytris strigose asperatis, fusco-picea, pilis brevibus aureis vestita; *capite* magno, lato, umbilicatum punctato; oculis modicis, sat prominentibus; *pronoto* sat magno, capite longiori et multum latiori, ad basim latissimo, punctis umbilicatis sat confertim impresso, interstitiis confertim reticulatis, lateribus rotundatis, leviter marginatis, margine basali leviter arcuatâ, angulis latis, productis; *elytris* brevibus, subquadratis, postice parum attenuatis, capite atque pronoto angustioribus, haud longioribus, reticulatum et strigose asperatis, lateribus marginatis, apicibus latis, rotundatis; abdomine sat longe exserto, ordinibus transversis umbilicatum punctato, interstitiis confertim reticulatis; pedibus atque antennis modicis, late flavis.

Immatura tota flava.

Body short and broad, very convex, shining; elytra strigosely asperate, fusco-piceous, clothed with short golden hair: immature specimens are entirely yellow.

Head large and broad, marked with umbilicated punctures; eyes moderate, rather prominent; antennae bright yellow.

Thorax rather large, longer and much broader than the head, widest at the base, rather closely marked with umbilicated punctures, interstices closely reticulated, sides rounded and slightly margined, basal margin slightly arcuated, with the angles broad and produced.

Scutellum very large, triangular, strigosely asperated.

Elytra short, subquadrate, slightly attenuated posteriorly, narrower and not longer than the head and thorax, widest at the shoulders, reticulately and strigosely asperated, sides margined, extremities broad and rounded.

Abdomen considerably exserted, with a transverse row of umbilicated punctures on each segment, interstices closely reticulated, apex entire.

Legs moderate, bright yellow.

Under parts piceous, with the mouth and coxæ paler.

Differs from other species in its strigose and reticulated sculpture.

Habitat, South America. Found in Brazil, near Blumenau.

Trichopteryx depressa.

T. depressa, Matthews, Ann. Mag. Nat. Hist. 6, xiii. 1894, p. 339.

L. c. $\frac{7}{16}$ — $\frac{5}{16}$ lin. = 0·87–1 mm. — Oblongo-ovalis, sat depressa, postice haud attenuata, nitidula, nigra, pilis griseis parce vestita, elytris fere parallelis: capite magno, prominenti, in fronte rotundato, fere glabro, nitido; oculis magnis, prominentibus; antennis modicis, flavis; pronoto magno, ad basim latissimo, vix dilatato, tuberculis modicis irregulariter dispositis, interstitiisque nitidis. profunde reticulatis ornat, lateribus marginatis, margine basali leviter sinuata, angulis latis, modice productis; elytris fere parallelis, piceo-nigris, sat brevibus, capite atque pronoto nec longioribus neque latioribus, profunde et confertim asperatis, interstitiis profunde reticulatis, lateribus marginatis, fere rectis, apicibus modice rotundatis, extremis albidis; abdomine piceo, modice exserto; pedibus robustis, laete flavis.

Body oblong-oval, rather depressed, not attenuated posteriorly, rather shining, black, sparingly clothed with grey hairs; elytra nearly parallel.

Head large and prominent, rounded in front, nearly smooth and shining; eyes large and prominent; antennae moderate, yellow.

Thorax large, widest at the base, but scarcely dilated, ornamented with moderate tubercles irregularly distributed, with the interstices shining and deeply reticulated, sides margined, basal margin faintly sinuated, with the angles broad and moderately produced.

Scutellum large, triangular, deeply asperated.

Elytra nearly parallel, pitchy black, rather short, not longer or broader than the head and thorax, deeply and closely asperated, with the interstices deeply reticulated, sides margined, nearly straight, extremities moderately rounded, with the outer edge white.

Abdomen piceous, moderately exposed.

Legs robust, bright yellow.

Under parts piceous, deeply reticulated.

Differs from *T. occidentalis* in its narrower, more parallel, and depressed oblong form, reticulated sculpture, and black colour.

Habitat, America. Found sparingly by Mr. Smith in the West Indian Islands.

Trichopteryx brevior.

T. brevior, Matthews, Trans. Am. Ent. Soc. xi. p. 136, 1884; Biol. Cent.-Amer. Col. ii. pt. i. p. 140, 1888.

L. c. $\frac{9}{16}$ lin. = 0.75 mm.—Oblonga, modice convexa, nitida, pilis fulvis vestita; capite magno, lato, antice producto, tuberculis minutis ornato; oculis magnis, haud prominentibus; pronoto modico, fere quadrato, capite sat longiori, haud latiori, prope medium latissimo, tuberculis parvis, ordinibus sinuatis dispositis, interstitiisque nitidis, reticulatis, ornato, lateribus marginatis, margine basali leviter reflexâ et leviter sinuatâ, angulis acutis, valde productis, dilutioribus; elytris fusciscentibus, oblongis, capite atque pronoto haud longioribus aut latioribus, ordinibus transversis leviter asperatis, lateribus fere rectis, apicibus latis, rotundatis, dilutioribus, extremis albidis; abdomine modice exserto, piceo; pedibus robustis, atque antennis læte flavis.

Setis erectis prope medium pronoti, atque ad humeros et apices versus elytrorum utrinque instructa.

Body oblong, moderately convex, shining, clothed with fulvous hair.

Head large and broad, produced in front, ornamented with minute tubercles; eyes large, not prominent; antennæ moderate, bright yellow.

Thorax moderate, nearly quadrate, rather longer not broader than the head, widest near the middle, ornamented with small tubercles in transverse sinuated rows, interstices reticulated and shining, sides margined, basal margin slightly reflexed and slightly sinuated, with the angles acute, much produced, and pale, furnished with an erect seta on each side near the middle.

Scutellum large, triangular, closely asperate.

Elytra fusciscent, oblong, not longer or broader than the head and thorax, with an erect seta on each side near the shoulders, and another towards the apex, slightly asperated in transverse rows, sides nearly straight, extremities broad, rounded, and pale, with the edge white.

Abdomen moderately exserted, piceous.

Legs robust, bright yellow.

Under parts black, rather convex, with the posterior margin of the metasternum and apex of the venter yellow.

Differs from *T. diffinis* in its smaller size, much fainter sculpture, and shorter elytra.

Habitat, Central America. Found in Guatemala by Mr. Champion.

Trichopteryx parallelopipeda.

T. parallelopipeda, Matthews, Cist. Ent. ii. p. 175, 1877.

L. c. $\frac{8}{16}$ lin. = 1 mm.—Oblonga, convexa, sat nitida, nigra, pilis sat longis, flavis vestita; capite lato, tuberculis parvis, interstitiisque reticulatis, ornato; oculis modicis, haud prominentibus; pronoto modico, valde convexo, capite longiori et latiori, ad basim latissimo, tuberculis modicis, ordinibus irregularibus confertim dispositis, interstitiisque profunde reticulatis, ornato, lateribus parum rotundatis, leviter marginatis, margine basali depressa, leviter sinuatâ, angulis productis, dilutioribus; elytris oblongis, haud attenuatis, capite atque pronoto longioribus, haud latioribus, ordinibus sinuatis confertim asperatis, interstitiis profunde reticulatis, lateribus fere rectis, leviter marginatis, apicibus latis, rotundatis, angusto dilutioribus, abdomine parum exserto; pedibus modicis, læte flavis; antennis sat brevibus, piceo-nigris, articulo octavo parvo.

Body oblong, convex, rather shining, black, clothed with rather long yellow hair.

Head broad, ornamented with small tubercles, with the interstices reticulated; eyes

moderate, not prominent; antennæ rather short, pitchy-black, with the eighth joint small.

Thorax moderate, very convex, longer and broader than the head, widest at the base, ornamented with moderate tubercles closely arranged in irregular rows, with the interstices deeply reticulated, sides slightly rounded and margined, basal margin depressed and faintly sinuated, with the angles produced and pale.

Scutellum large, deeply and closely asperated, and reticulated.

Elytra oblong, not attenuated posteriorly, longer but not broader than the head and thorax, closely asperated in wavy rows, with the interstices deeply reticulated, sides nearly straight, slightly margined, extremities broad, rounded, and narrowly pale.

Abdomen slightly exerted.

Legs moderate, bright yellow.

Under parts black, with the mouth, coxæ, and posterior margin of the metasternum piceous, metasternum deeply asperated.

Differs from T. Hornii in its oblong parallel form, thorax widest at the base, and short black antennæ.

Habitat, North America. Found in British Columbia by Mr. G. R. Crotch.

Trichopteryx xanthocera.

T. xanthocera, Matthews, Cist. Ent. ii. p. 174, 1877.

L. c. $\frac{8}{16}$ lin. = 1 mm.—Oblonga, sat lata, convexa, piceo-nigra, elytris rufescentibus, pilis modicis, pallidis vestita; capite magno, lato, tuberculis exiguis, interstitiisque glabris ornato; oculis haud prominentibus; pronoto magno, convexo, postice dilatato, capite longiori et multo latiori, ad basim latissimo, tuberculis minutis, ordinibus transversis, fere rectis, dispositis, interstitiisque nitidis, indistincte reticulatis, ornato, lateribus parum rotundatis, flavo-marginatis, margine basali depressa, sinuata, angulis longe productis, flavescentibus; elytris brevibus, quadratis, capite atque pronoto haud longioribus aut latioribus, ordinibus distinctis, sinuatis, sat profunde asperatis, lateribus fere rectis, sutura rufescenti, apicibus latis, parum rotundatis, dilutioribus: abdomine sat exserto, obtuso, rufo-piceo, apice minute tridentato; pedibus sat robustis, læte flavis; antennis longis, lætissime flavis.

Body oblong, rather broad, convex, nigro-piceous, with the elytra rufescent, clothed with pale hair.

Head large and broad, ornamented with very small tubercles, with the interstices shining; eyes not prominent; antennæ long, very bright yellow.

Thorax large, convex, dilated posteriorly, longer and much broader than the head, widest at the base, ornamented with minute tubercles in nearly straight transverse rows, interstices shining and indistinctly reticulated, sides faintly rounded, with the margins yellow, basal margin depressed, sinuated, with the angles very much produced, flavescent.

Scutellum large, triangular, and closely asperated.

Elytra short, quadrate, not longer or broader than the head and thorax, rather deeply asperate in distinct wavy rows, sides nearly straight, suture rufescent, extremities broad, slightly rounded, and paler.

Abdomen rather long, obtuse, rufo-piceous, with the apex minutely tridentate.

Legs rather robust, bright yellow.

Under parts entirely rufo-piceous, metasternum dusky.

Differs from T. Hornii in its larger size, quadrate and broader form, rufescent elytra, fainter sculpture, and large, very bright yellow antennæ.

Habitat, North America. Found in British Columbia by Mr. G. R. Crotch.

Trichopteryx Californica.*T. Californica*, Matthews, Cist. Ent. ii. p. 174, 1877.

L. c. $\frac{7}{16}$ — $\frac{8}{16}$ lin. = 0.87–1 mm.—Oblonga, sat convexa, castanea, nitida, pilis brevibus aureis vestita; capite magno, lato, tuberculis minutis, interstitiisque nitidis, ornato; oculis haud prominentibus; pronoto magno, convexo, capite parum longiori, multum latiori, ad basim latissimo, piceo-castaneo, angulis late atque lateribus flavescentibus, tuberculis minutis, ordinibus irregularibus dispositis, interstitiisque nitidis, confertim reticulatis, ornato, lateribus marginatis et rotundatis, margine basali profunde sinuata, angulis acutissimis, longe productis; elytris oblongis, rufo-castaneis, sutura dilutiori, capite atque pronoto aliquantum et angustioribus, et longioribus, ordinibus sinuatis confertim asperatis, interstitiis nitidis, lateribus rectis, fere parallelis, sutura postice elevata, apicibus latis, parum rotundatis; abdomine sat exserto, obtuso, apice tridentato; pedibus gracilibus, late flavis; antennis longis, gracilibus, late flavis.

Body oblong, rather convex, shining, castaneous, clothed with short golden hair.

Head large and broad, ornamented with minute tubercles, with the interstices shining; eyes not prominent; antennae long and slender, bright yellow.

Thorax large, convex, rather longer and much broader than the head, widest at the base, piceo-castaneous, with the angles and sides flavescent, ornamented with minute tubercles in irregular rows, interstices shining and closely reticulated, sides margined and rounded, basal margin deeply sinuated, with the angles much produced and very acute.

Scutellum large, closely asperated.

Elytra oblong, rufo-castaneous, with the suture paler, slightly narrower and longer than the head and thorax, closely asperated in wavy rows, with the interstices shining, sides straight, nearly parallel, suture elevated posteriorly, extremities broad and slightly rounded.

Abdomen rather long, obtuse, tridentate at the apex.

Legs moderate, rather slender, bright yellow.

Under parts piceous, mouth, coxae, and posterior margin of the metasternum yellow, metasternum deeply reticulated.

Differs from *T. lethierryi* in its much smaller size, longer elytra, shorter legs, fainter sculpture, and shorter pubescence.

Habitat, North America. Found in California by Mr. G. R. Crotch.

Trichopteryx castanea.*T. castanea*, Matthews, Cist. Ent. ii. p. 173, 1877.

L. c. $\frac{7}{16}$ lin. = 0.87 mm.—Oblonga, convexa, rufo-castanea, sat nitida, pilis aureis dense vestita; capite modico, antice producto, tuberculis minutis, interstitiisque reticulatis ornato; oculis haud prominentibus; pronoto modico, capite longiori et latiori, ante basim latissimo, tuberculis parvis, ordinibus sinuatis dispositis, interstitiisque nitidis, profunde reticulatis, ornato, lateribus rotundatis et marginatis, margine basali depressa, sinuata, angulis acutis, longe productis, late flavis; elytris oblongis, convexis, translucidis, capite atque pronoto longioribus, haud latioribus, sat profunde asperatis, interstitiis reticulatis, lateribus marginatis, fere parallelis, apicibus latis, rotundatis; abdomine sat longe exserto, obtuso, apice minute bidentato; pedibus robustis, late flavis; antennis longis, sat robustis, articulis duobus basalibus permagnis, late flavis, reliquis dilutioribus.

Body oblong, convex, rufo-castaneous, rather shining, thickly covered with golden hair.

Head moderate, produced in front, ornamented with minute tubercles, with the

interstices reticulated; eyes rather large, not prominent. antennae long and rather robust, with the two basal joints very large, bright yellow, and the rest paler.

Thorax moderate, longer and broader than the head, widest before the base, ornamented with small tubercles in wavy rows, interstices shining and deeply reticulated, sides rounded and margined, basal margin depressed and sinuated, with the angles acute, much produced, and broadly yellow.

Scutellum large, elongate, and closely asperated.

Elytra oblong, convex, and translucent, longer but not broader than the head and thorax, rather deeply asperate, with the interstices reticulate, sides almost parallel and margined, extremities broad and rounded.

Abdomen rather long, obtuse, minutely bidentate at the apex.

Legs robust, bright yellow.

Under parts castaneous, with the mouth, coxae, posterior margin of the metasternum, and extremity of the venter flavescent.

Differs from *T. Edithia* in its larger size, parallel form, thorax widest near the base, and longer legs and antennae.

Habitat, North America. Found in British Columbia by Mr. G. R. Crotch.

Trichopteryx carbonaria.

T. carbonaria, Matthews, Ent. Mo. Mag. ix. p. 179, 1873.

L. c. $\frac{4}{16}$ lin. = 0.75 mm.—Oblonga, convexa, aterrima, pilis brevibus flavis sat dense vestita; capite permagno, lato, tuberculis minutis, ordinibus rectis dispositis, interstitiisque reticulatis ornato; oculis magnis, laud prominentibus; pronoto modico, capite latiori, vix longiori, pone medium latissimo, tuberculis minutis, distinctis, ordinibus sinuatis dispositis, interstitiisque reticulatis, nitidis, ornato, lateribus valde rotundatis, leviter marginatis, ad basim contractis, margine basali valde depressa, sinuata, angulis acutis, valde productis; elytris brevibus, quadratis, capite atque pronoto angustioribus, laud longioribus, prope media latissimis, profunde et confertissime asperatis, lateribus leviter rotundatis, apicibus latis, rectis, dilutioribus, summis albidis; abdomine sat longe exserto, obtuso, apice obsolete tridentato, pedibus atque antennis robustis, obscure flavis.

Body oblong, convex, deep black, rather thickly clothed with short yellow hair.

Head very large and broad, ornamented with minute tubercles in straight rows, with the interstices reticulated; eyes large, not prominent; antennae robust, dusky yellow.

Thorax moderate, broader but scarcely longer than the head, widest behind the middle, ornamented with minute distinct tubercles in wavy rows, interstices reticulated and shining, sides much rounded, slightly margined, and contracted at the base, basal margin much depressed, sinuated, with the angles acute and much produced.

Scutellum rather large, triangular, closely and deeply asperate.

Elytra short, quadrate, narrower, but not longer than the head and thorax, widest near the middle, deeply and very closely asperated, sides slightly rounded, extremities broad, straight, and paler, with the edge white.

Abdomen rather long, obtuse, with the apex obsoletely tridentate.

Legs robust, dusky yellow.

Under parts black, with the mouth and coxae piceous.

Differs from *T. picicornis* in its paler and more slender antennae, with the eighth joint linear, and in the very different sculpture of the thorax and elytra.

Habitat, Europe. Found by myself in Sherwood Forest in Nottinghamshire.

Trichopteryx dubitata.

T. dubitata, Matthews, Ann. Mag. Nat. Hist. 6, xiii. p. 340, 1894.

L. c. $\frac{7}{16}$ lin. = 0.87 mm.—Elongata, angusta, valde convexa, nitidula, pallide castanea (an immatura?), pilis crassis aureis vestita; *capite* magno, prominenti, confertim tuberculato; oculis nigris, magnis, prominentibus; antennis longis, gracilibus, flavis; *pronoto* brevi, capite paulo latiori, prope medium latissimo, tuberculis modicis ordinibus densis sinuatis dispositis, interstitiisque reticulatis ornato, lateribus valde rotundatis, fortiter marginatis, margine basali depressa, sinuata, angulis latis, modice productis; *elytris* perbrevibus, capite atque pronoto brevioribus, haud latioribus, prope media latissimis, confertim asperatis, interstitiis fortiter reticulatis, lateribus marginatis, apicibus latis, leviter rotundatis; abdomine longius exserto, leviter asperato; pedibus robustis, pallide flavis.

Body elongate, narrow, very convex, moderately shining, pale castaneous (perhaps immature), clothed with coarse golden hair.

Head large and prominent, closely tuberculate; eyes black, large and prominent; antennæ long and slender, yellow.

Thorax short, rather broader than the head, widest near the middle, ornamented with moderate tubercles in close wavy rows, interstices reticulate, sides much rounded and strongly margined, basal margin depressed and sinuated, with the angles broad, moderately produced.

Scutellum large, triangular, deeply asperated.

Elytra very short, shorter and not broader than the head and thorax, widest near the middle, closely asperated, with the interstices strongly reticulated, sides margined, extremities broad, slightly rounded.

Abdomen much exposed, faintly asperated.

Legs robust, pale yellow.

Under parts testaceous.

Differs from others in its narrow elongate form, small size, large and prominent eyes, short thorax and elytra, long slender antennæ, and coarse pubescence.

Habitat, America. One specimen found by Mr. Smith in the West Indian Island of St. Vincent.

Trichopteryx fuliginosa, sp. n.

L. c. $\frac{8}{16}$ lin. = 1 mm.—Oblongo-ovalis, convexa, nigro-fusca, haud nitida, pilis obscure fuscis dense vestita, sat fortiter confertissime tuberculata; *capite* modico, sat prominenti, dense tuberculato, oculis modicis; *pronoto* modico, capite latiori et longiori, ante basin latissimo, dense tuberculato, lateribus marginatis, margine basali sinuata, angulis acutis, modice productis; *elytris* brevibus, capite atque pronoto brevioribus, haud latioribus, ante media latissimis, confertim asperatis, lateribus vix rotundatis, apicibus valde rotundatis, anguste dilutioribus; abdomine longius exserto, obtuso, nigro; antennis sat longis, piceis; pedibus modicis, læte flavis.

Body oblong-oval, convex, fuscous black, not shining, thickly covered with dusky hair, strongly and very closely tuberculate.

Head moderate, rather prominent, closely tuberculate, eyes moderate, antennæ rather long, piceous.

Thorax moderate, longer and broader than the head, widest before the base, closely tuberculate, sides margined, basal margin sinuated, with the angles acute, moderately produced.

Scutellum very large, triangular, closely asperate.

Elytra short, shorter and not broader than the head and thorax, widest before the

middle, closely asperate, sides scarcely rounded, extremities much rounded, narrowly pale.

Abdomen much exerted, obtuse, black.

Legs moderate, bright yellow.

Under parts black, mouth and coxæ yellow.

Differs from other species in its dull sooty colour and close sculpture.

Habitat, Central America. Found in Mexico.

Trichopteryx bidilatata.

T. bidilatata, Matthews, Trans. Am. Ent. Soc. xi. p. 138, 1884; Biol. Cent.-Amer. Col. ii. pt. i. p. 141, 1888.

L. c. $\frac{5}{16}$ – $\frac{6}{16}$ lin. = 0.63–0.75 mm.—Oblonga, convexa, haud nitida, obscure castanea, pilis brevibus griseis dense vestita; *capite* magno, lato, tuberculis ordinibus transversis dispositis, interstitiisque reticulatis, ornato; oculis magnis, sat prominentibus; *pronoto* modico, transverso, capite parum latiori, haud longiori, ad medium latissimo, tuberculis sat magnis, ordinibus transversis, sinuatis dispositis, interstitiisque reticulatis, sat nitidis, ornato, lateribus rotundatis et marginatis margine basali fere recta, angulis minime productis; *elytris* oblongo-ovalibus, ad media latissimis, capite atque pronoto nec latioribus, neque longioribus, ordinibus transversis confertissime asperatis, apicibus latis, fere rectis, extremis albidis; abdomine longe exserto, obtuso, marginibus segmentorum dilutioribus; pedibus longis, lute flavis; antennis modicis, obscure flavis.

Body oblong, convex, not shining, dark castaneous, thickly clothed with short grey hair.

Head large and broad, ornamented with tubercles in transverse rows, interstices reticulated; eyes large, and rather prominent; antennæ moderate, dusky yellow.

Thorax moderate, transverse, rather broader but not longer than the head, widest at the middle, ornamented with rather large tubercles in transverse rows, interstices reticulated and rather shining, sides rounded and margined, basal margin nearly straight, with the angles slightly produced.

Scutellum large, triangular, and closely asperated.

Elytra oblong-oval, widest at the middle, not longer or broader than the head and thorax, very closely asperated in transverse rows, extremities broad, nearly straight, with the edge white.

Abdomen long, obtuse, with the margins of the segments paler.

Legs rather long, bright yellow.

Under parts black, with the coxæ and apex of the venter piceous.

Differs from *T. Edithia* in its smaller size, longer elytra, and dusky antennæ; and from all others in the doubly dilated outline of its body.

Habitat, Central America. Found in Guatemala by Mr. Champion.

Trichopteryx Grenadensis.

T. Grenadensis, Matthews, Ann. Mag. Nat. Hist. 6, xiii. p. 339. 1894.

L. c. $\frac{5}{16}$ lin. = 0.63 mm.—Oblonga, sat depressa, postice haud attenuata, atra, nitidula, pilis fusciscentibus parce vestita; *capite* permagno, lato, fere glabro, nitido; oculis modicis, haud prominentibus; antennis sat robustis, flavis; *pronoto* brevi, capite paullo latiori, ante basim latissimo, tuberculis modicis confertim sed irregulariter dispositis, interstitiisque reticulatis ornato, lateribus marginatis et rotundatis, margine basali fere recta, angulis rectis; *elytris* brevibus, capite atque pronoto nec longioribus, neque latioribus, prope medios latissimis, sat profunde

confertim ordinibus transversis asperatis, lateribus leviter rotundatis et marginatis, apicibus valde obtusis, leviter rotundatis; abdomine paullo exserto nigro; pedibus robustis, luteo flavis.

Body oblong, rather depressed, not attenuated posteriorly, deep black, rather shining, sparingly clothed with dusky hair.

Head very large and broad, nearly smooth; eyes moderate, not prominent; antennae rather robust, yellow.

Thorax short, rather broader than the head, widest before the base, closely ornamented with moderate tubercles irregularly arranged, with the interstices reticulated, sides margined and rounded, basal margin nearly straight, with the angles right angles.

Scutellum very large and broad, triangular, deeply asperated.

Elytra short, not longer or broader than the head and thorax, widest near the middle, rather deeply and closely asperated in transverse rows, sides slightly rounded and margined, extremities slightly rounded and very obtuse.

Abdomen not much exposed, black.

Legs robust, bright yellow.

Under parts piceous, with the mouth and coxae yellow.

Differs from *T. bidilatata* in its much smaller size, finer sculpture, and much shorter antennae; and from others in the peculiar dilatation of the thorax and elytra.

Habitat, America. Two specimens found by Mr. Smith in the West Indian Island of Grenada.

Trichopteryx excelsa.

T. excelsa, Matthews, Trans. Am. Ent. Soc. xi. p. 133, 1884; Biol. Cent.-Amer. Col. ii. pt. i. p. 139, 1888.

L. c. $\frac{7}{16}$ lin. = 0.87 mm.—Ovalis, sat brevis, valde convexa, nitidula, capite atque pronoto nigris, elytris rufo-fuscis, pilis fulvis sparse vestita; capite magno, antice rotundato, tuberculis minutis, remotis, ornato; oculis modicis, prominentibus; pronoto sat brevi, capite parum longiori, multum latiori, ante basim latissimo, postice valde depresso, tuberculis minutis, ordinibus sinuatis, remotis, dispositis, interstitiisque reticulatis, nitidis, ornato, lateribus valde rotundatis, leviter marginatis, margine basali leviter sinuata, angulis latis, valde productis, late flavescens; elytris brevibus, convexis, capite atque pronoto brevioribus, haud latioribus, ad media latissimis, ordinibus transversis, remotis, modice asperatis, lateribus valde rotundatis, apicibus valde rotundatis, anguste dilutioribus; abdomine longe exserto, obtuso; pedibus atque antennis modicis, obscure flavis.

Body oval, rather short, very convex, rather shining, head and thorax black, elytra rufo-fuscous, sparingly clothed with fulvous hair.

Head large, rounded in front, ornamented with minute remote tubercles; eyes moderate, prominent; antennae moderate, dusky yellow.

Thorax rather short, rather longer and much broader than the head, widest before the base, much depressed towards the base, ornamented with minute tubercles in remote wavy rows, interstices reticulated and shining, sides much rounded and slightly margined, basal margin faintly sinuated, with the angles broad, much produced, and broadly flavescens.

Scutellum very large and broad, triangular, moderately asperated.

Elytra short, convex, shorter and not broader than the head and thorax, widest at the middle, moderately asperated in remote transverse rows, sides much rounded, extremities much rounded and narrowly paler.

Abdomen much exposed, obtuse.

Legs moderate, dusky yellow.

Under parts castaneous, with the mouth and coxæ paler.

Differs from others in its short convex form, rounded sides, and depressed base of the thorax, short and rounded elytra, and in sculpture.

Habitat, Central America. Found by Mr. Champion on Totonicapam, Guatemala, at an elevation of 10,500 ft.

Trichopteryx vicina.

T. vicina, Matthews, Cist. Ent. ii. p. 172, 1877; Trans. Am. Ent. Soc. xi. p. 138, 1884; Biol. Cent.-Amer. Col. ii. pt. i. p. 138, 1888.

L. c. $\frac{1}{16}$ lin. = 0.75 mm.—Oblonga, parum convexa, nigerrima, pilis brevibus flavis parce vestita; *capite* permagno, lato, nitidissimo, tuberculis indistinctis impresso, oculis magnis, haud prominentibus; *pronoto* magno, capite longiori et latiori, ad basim latissimo, tuberculis parvis, ordinibus remotis, sinuatis, dispositis, interstitiisque nitidissimis, profunde reticulatis ornato, lateribus leviter rotundatis, late marginatis, margine dilutiori, margine basali fere rectâ, angulis latis, productis, dilutioribus; *elytris* brevibus, quadratis, lateribus parallelis, capite atque pronoto parum angustioribus, haud longioribus, ordinibus remotis, sinuatis, modice asperatis, interstitiis reticulatis, sutura postice elevata, apicibus vix rotundatis, anguste dilutioribus; abdomine parum exserto, apice tridentato; pedibus late flavis; antennis nigris.

Body oblong, slightly convex, deep black, sparingly clothed with yellow hair.

Head very large and broad, very shining, indistinctly tuberculate; eyes large, not prominent; antennæ moderate, black.

Thorax large, longer and broader than the head, widest at the base, ornamented with small tubercles in remote wavy rows, interstices deeply reticulated and very shining, sides slightly rounded and broadly margined, with the margin pale, basal margin nearly straight, with the angles broad and produced, pale.

Scutellum large, triangular, closely asperated.

Elytra short, quadrate, parallel at the sides, not longer but rather narrower than the head and thorax, moderately asperate in remote wavy rows, interstices reticulated, suture elevated posteriorly, extremities scarcely rounded, narrowly pale.

Abdomen slightly exposed, with the apex tridentate.

Legs bright yellow.

Under parts black, with the mouth and coxæ paler.

Differs from *T. picicornis* in its larger size, much fainter and very different sculpture of the thorax, and longer and more slender antennæ.

Habitat, North and Central America. Found in British Columbia by Mr. G. R. Croteh; and in Guatemala by Mr. Champion.

Trichopteryx nigrita.

T. nigrita, Matthews, Trans. Am. Ent. Soc. xi. p. 138, 1884; Biol. Cent.-Amer. Col. ii. pt. i. p. 141, 1888.

L. c. $\frac{1}{16}$ lin. = 0.63 mm.—Oblonga, modice depressa, haud nitida, fusco-nigra, pilis fulvis vestita; *capite* modico, antice sat producto, tuberculis parvis, remotis, interstitiisque reticulatis ornato, oculis sat prominentibus; *pronoto* modico, postice leviter dilatato, capite latiori, vix longiori, ad basim latissimo, tuberculis parvis, ordinibus sinuatis, remotis, dispositis, interstitiisque sat profunde reticulatis ornato, lateribus sat late marginatis, modice rotundatis, margine basali leviter sinuata, angulis latis, parum productis; *elytris* oblongis, fuscis, capite atque pronoto parum longioribus, haud latioribus, ordinibus transversis sat profunde et confertim aspe-

ratis, lateribus rectis, apicibus latis, valde rotundatis, dilutioribus; abdomine longius exserto, piceo, apice obtuse bidentato; pedibus atque antennis sat gracilibus, flavis, his obscurioribus.

Body oblong, moderately depressed, not shining, fuscous-black, clothed with fulvous hair.

Head moderate, rather produced in front, ornamented with small remote tubercles, with the interstices reticulated; eyes rather prominent; antennæ slender, dusky yellow.

Thorax moderate, slightly dilated posteriorly, broader but scarcely longer than the head, widest at the base, ornamented with small tubercles in remote wavy rows, with the interstices rather deeply reticulated, sides rather broadly margined and moderately rounded, basal margin faintly sinuated, with the angles broad and slightly produced.

Scutellum large, triangular, deeply and closely asperated.

Elytra oblong, brown, rather longer but not broader than the head and thorax, deeply and closely asperated in transverse rows, sides straight, extremities broad and much rounded, pale.

Abdomen considerably exposed, piceous, obtusely bidentate at the apex.

Legs rather slender, yellow.

Under parts black, with the coxæ and apex of the venter piceous.

Differs from *T. obscura* in its smaller size, more depressed form, and much deeper sculpture; and from *T. elevata* in its larger size, depressed form, and black colour.

Habitat, Central America. Found in Guatemala by Mr. Champion.

Trichopteryx mærens.

T. mærens, Matthews, Cist. Ent. i. p. 298, 1874.

L. c. $\frac{5}{16}$ lin. = 0.63 mm.—Brevis, lata, valde convexa, nitidissima, aterrima, pilis brevibus, pallidis, parce vestita; capite brevi, lato, fere glabro, punctis minutis, remotis, impresso; oculis parvis, haud prominentibus; pronoto brevi, transverso, valde convexo, capite paullo longiori, vix latiori, prope medium latissimo, glabro, nitidissimo, ad basim depresso, minutissime et remote punctato, lateribus modice rotundatis, late marginatis, margine translucida, margine basali recta, angulis rectis; elytris perbrevis, convexissimis, capite atque pronoto nec longioribus, nec latioribus, ad media latissimis, ad humeros pronoto angustioribus, ordinibus transversis levissime asperatis, interstitiis glabris, nitidissimis, apicibus latissimis, vix rotundatis, extremis flavis; abdomine minime exserto; pedibus atque antennis longis, gracilibus, hæte flavis, his sat obscurioribus.

Body short and broad, very convex and very shining, deep black, sparingly clothed with short pale hair.

Head short, very broad, nearly smooth, minutely and remotely punctured; eyes small, not prominent; antennæ very long, dusky yellow.

Thorax short, transverse, very convex, rather longer but scarcely broader than the head, widest near the middle, depressed at the base, smooth and very shining, very minutely and remotely punctured, sides moderately rounded and broadly margined, with the margin translucent, basal margin straight, with the angles right angles.

Scutellum large, very broad, triangular, indistinctly asperated.

Elytra very short and very convex, not longer or broader than the head and thorax, widest at the middle, narrower than the thorax at the shoulders, faintly asperated in transverse rows, interstices smooth and very shining, sides rounded and margined, extremities very broad and scarcely rounded, with the edge yellow.

Abdomen very little exposed, black.

Legs long and slender, bright yellow.

Under parts entirely black.

Differs from all other species in its short, rounded and very convex form, smooth and very shining surface, deep black colour, and long antennæ.

Habitat, North America. This remarkable species was one of the last discoveries of my much lamented friend Mr. G. R. Crotch, by whom it was found near Philadelphia. A second example has since then occurred in the same locality.

Trichopteryx tesserula.

T. tesserula, Matthews, Ann. Mag. Nat. Hist. 6, iii. p. 193, 1889.

L. c. $\frac{5}{16}$ lin. = 0.63 mm.—Brevissima, quadrata, sat depressa, nitidula, fusca, pilis pallidis vestita: *capite* permagno, antice producto, leviter et remote tuberculato, interstitiis leviter reticulatis; oculis parvis; *pronoto* brevi, capite parum latiori, haud longiori, ad basim latissimo, tuberculis parvis remotis, interstitiisque nitidis, reticulatis, ornato, lateribus marginatis, leviter rotundatis, margine basali fere rectâ, angulis minime productis; *elytris* perbrevibus, omnino quadratis, depressis, capite atque pronoto haud longioribus aut latioribus, modice asperatis, lateribus rectis, leviter marginatis, apicibus latis, vix rotundatis; abdomine modice exserto, rufo-piceo; pedibus atque antennis sat gracilibus, late flavis, parum articulo octavo parum incrassato.

Body very short, quadrate, rather depressed, moderately shining, clothed with pale hair, fuscous.

Head very large, produced in front, ornamented with small remote tubercles, interstices faintly reticulated; eyes small; antennæ moderate, slender, yellow, with the eighth joint incrassate.

Thorax short, rather broader but not longer than the head, widest at the base, ornamented with small remote tubercles, the interstices reticulated, sides margined and slightly rounded, basal margin nearly straight, with the angles scarcely produced.

Scutellum small and broad, triangular.

Elytra very short, perfectly quadrate, depressed, not longer or broader than the head and thorax, moderately asperated, sides straight, slightly margined, extremities very broad, scarcely rounded.

Abdomen moderately exerted, rufo-piceous.

Legs rather slender, bright yellow.

Under parts castaneous.

Differs from all other species in its very small size, and perfectly quadrate form.

Habitat, North America. Found in California by Mr. Casey.

Trichopteryx vitrea.

T. vitrea, Matthews, Trans. Am. Ent. Soc. xi. p. 139, 1884: Biol. Cent.-Amer. Col. ii. pt. i. p. 142, 1888.

L. c. $\frac{1}{16}$ – $\frac{5}{16}$ lin. = 0.50–0.63 mm.—Perbrevis, lata, valde convexa, transluide castanea, pilis brevibus, aurcis, sparsissime induta: *capite* magno, lato, antice producto, tuberculis exiguis remote ornato; oculis magnis, valde prominentibus; *pronoto* brevi, transverso, capite sat latiori, haud longiori, ante basim latissimo, tuberculis exiguis, remotis, interstitiisque glabris, nitidis, ornato, lateribus rotundatis, margine basali fere rectâ, angulis vix productis; *elytris* brevibus, subquadratis, capite atque pronoto fere pariter longis et latis, prope media latissimis, leviter sed confertim asperatis, lateribus rotundatis, apices versus contractis,

apicibus minime rotundatis: abdomine longe exserto, flavescenti; pedibus atque antennis robustis, læte flavis.

Body very short and broad, very convex, translucidly castaneous, very sparingly clothed with short golden hair.

Head large and broad, produced in front, ornamented with very minute remote tubercles: eyes large and very prominent; antennæ rather large, bright yellow.

Thorax short, transverse, broader but not longer than the head, widest before the base, ornamented with very minute remote tubercles, interstices smooth and shining, sides rounded, basal margin nearly straight, with the angles scarcely produced.

Scutellum very large, triangular, faintly but closely asperated.

Elytra short, subquadrate, not longer or broader than the head and thorax, widest near the middle, faintly but closely asperated, sides rounded, and contracted posteriorly, extremities very little rounded.

Abdomen long, flavescent.

Legs robust, bright yellow.

Under parts pale castaneous, metasternum dusky.

Differs from all other species in its very short and very convex form, and shining translucent surface.

Habitat, Central America. Found very plentifully near Bugaba, Panama, by Mr. Champion.

Trichopteryx Japonica.

T. Japonica, Matthews, Cist. Ent. iii. p. 81, 1884.

L. c. $\frac{5}{16}$ – $\frac{6}{16}$ lin. = 0.63–0.75 mm.—Subovata, sat angusta, convexa, nitida, pieco-castanea, pilis flavis vestita; *capite* permagno, lato, antice rotundato, remote et modice punctato; oculis sat magnis, prominentibus; *pronoto* modico, capite parum longiori et latiori, prope medium latissimo, punctis magnis, foveolatis sat remote impresso, interstitiis glabris, nitidis, lateribus valde rotundatis et late marginatis, margine basali recta, angulis acutis, sat prominentibus; *elytris* ovatis, rufo-castaneis, capite atque pronoto fere pariter longis et latis, pone media latissimis, punctis profundis, sat remotis, impressis, interstitiis glabris, nitidis, lateribus modice rotundatis, apicibus latis, minime rotundatis, extremis albidis; abdomine longius exserto, obtuso; pedibus sat magnis, læte flavis, antennis longis, obscure flavis.

Body subovate, rather narrow, convex, shining, dark-castaneous, clothed with yellow hair.

Head very large and broad, rounded in front, moderately and remotely punctured; eyes rather large and prominent; antennæ long, dusky yellow.

Thorax moderate, rather longer and broader than the head, widest near the middle, marked with deep foveolate remote punctures, interstices smooth and shining, sides much rounded and broadly margined, basal margin straight, with the angles acute and prominent.

Scutellum large, triangular, remotely punctured.

Elytra ovate, rufo-castaneous, not longer or broader than the head and thorax, widest behind the middle, marked with deep and rather remote punctures, interstices smooth and shining, sides moderately rounded, extremities broad and very little rounded, with the edge white.

Abdomen much exserted, obtuse.

Legs rather large, bright yellow.

Under parts castaneous, mouth and coxæ paler.

Differs from *T. variolosa* in its much smaller size and much narrower form; and

from all other species in the deeply punctured surface, and peculiar form of its body.

Habitat, Asia. Found not uncommonly in Japan by Mr. Lewis.

I am much inclined to regard this species as a local race of *T. variolosa*; but have thought it safer to separate them on account of the great and persistent difference in size and form which exists between Mr. Lewis's captures and the European species.

GEN. XV.—PTILIODES. PL. XIII. B.

Ptiliodes, Matthews, Cist. Ent. iii. p. 40, 1882.

Corpus suboblongum, sat convexum.

Caput magnum, latum, prominens; parte gulari magna, valde convexa; oculi sat magni, prominentes; antennæ sat longe ante oculos insertæ.

Antennæ (Pl. XIII. B. fig. 7) breves, robustæ, 11-articulatæ; 1 magno, obconico; 2 primo minori, ovato, apice abrupte truncato; 3 gracili, sublineari; 4 et 5 tertio fere similibus; 6-8-ovatis, sibiipsis fere paribus; 9 precedenti consimili sed majori; 10 nono multum majori, subrotundato, incrassato; 11 obtuse conico, decimo longiori, pariter crasso; duobus ultimis clavam moniliformem formantibus, atque setis perbrevibus verticillatis instructis.

Partes oris hand examinatæ.

Pronotum breve, transversum, ad basim leviter contractum, angulis posterioribus haud productis.

Scutellum sat parvum, triangulare.

Elytra oblonga, truncata.

Alæ lanceolatæ, sub clytris visæ.

Abdomen longius exsertum, sat obtusum, segmentis quatuor apertis, apicali obtuso, integro; alioqui non examinatum.

Prothorax (Pl. XIII. B. fig. 2, a) *prosterno* magno; *episternis* parvis, conspicuis; *epimeris* modicis, coxarum receptacula eicientibus; *coxarum receptaculis* magnis, ovatis, obliquis, contingentibus.

Pectus mesosterno (Pl. XIII. B. fig. 2, d) modico, breviter carinato; *episternis* modicis, oblongis; *epimeris* magnis, humeralibus, ad receptacula coxarum late extensis; *coxarum receptaculis* rotundatis, contingentibus.

metasterno (Pl. XIII. B. fig. 2, g) magno, subquadrato, ad latera corporis extenso, margine posteriori inter coxas recta; *episternis* atque *epimeris* sub clytris totis occultis.

Venter segmentis sex compositus, omnibus pariter longis, apicali leviter emarginato.

Pedes sat breves, sat robusti, *antici* trochanteribus parvis, trigonalibus; femoribus simplicibus; tibiis parum dilatatis; tarsis brevibus, triarticulatis, 1 sat longo; 2 primo multum minori; 3 elongato, gracillimo; unguibus gracilibus, leviter curvatis.

intermedii anticis similes, sed parum longiores.

postici (Pl. XIII. B. fig. 8) trochanteribus claviformibus, alioqui anticis similes, sed tibiis magis dilatatis.

Coxæ anteriores magnæ, ovatæ, contingentes.

intermediæ rotundatæ, contingentes.

posteriores sat magnæ, prominentes, ovaliter et late per totam longitudinem laminatæ, modice remotæ.

Body suboblong, rather convex.

Head large and broad, prominent, gular region large and very convex; eyes large and prominent; antennæ inserted considerably in front of the eyes.

Antennæ (Pl. XIII. B. fig. 7) short and robust, 11-jointed: 1 large, obconic; 2 smaller than the first, ovate, with the apex abruptly truncate; 3-5 slender,

sublinear, nearly equal to each other ; 6-8 ovate, nearly equal to each other ; 9 like the preceding in form, but larger ; 10 much larger than the ninth, subrotundate and incrassate ; 11 obtusely conical, equally incrassate, but longer than the ninth ; the two last form a moniliform club, and are ornamented with very short verticillated setæ.

Parts of the mouth not examined.

Thorax short, transverse, slightly contracted towards the base, posterior angles not produced.

Scutellum rather small, triangular.

Elytra oblong, truncate.

Wings lanceolate, seen through the elytra.

Abdomen long, rather obtuse, with four segments exposed, apical segment obtuse ; in other respects not examined.

Prothorax (Pl. XIII. B, fig. 2, a) with the *prosternum* large ; *episterna* small, but evident ; *epimera* moderate, enclosing the coxal cavities ; *coxal cavities* large, ovate, oblique, and contingent.

Mesosternum (Pl. XIII. B, fig. 2, d) moderate, with a short carina ; *episterna* moderate, oblong ; *epimera* large, humeral, broadly extending to the coxal cavities ; *coxal cavities* rounded and contingent.

Metasternum (Pl. XIII. B, fig. 2, g) large, subquadrate, extending to the sides of the body, with the posterior margin between the coxæ straight ; *episterna* and *epimera* entirely covered by the elytra.

Venter composed of six segments of equal length, apical segment faintly emarginate.

Legs rather short and rather stout, *anterior* with the trochanters small, trigonal ; femora simple ; tibiæ faintly dilated ; tarsi short, three-jointed ; 1 rather long ; 2 much smaller than the first ; 3 long and slender ; claws slender and faintly curved.

intermediate (Pl. XIII. B, fig. 8) like the anterior, but rather longer.

posterior with the trochanters claviform, in other respects like the anterior, but with the tibiæ more dilated.

Coxa anterior large, ovate, contingent.

intermediate moderate, rounded, contingent.

posterior rather large, prominent, ovally and broadly laminated on the whole outer margin, and rather widely distant.

It is difficult to determine the proper position of this genus : in the form of its antennæ it resembles *Ptilium* more than any of the present tribe ; in its short transparent elytra and long abdomen it is like *Ptinella* ; but the short and broad shape of its head and thorax, together with the comparative shortness of its legs and antennæ, incline me to place it next in succession to *Nephanes*.

As the genus is at present only represented by a single individual, I have not ventured to dissect the parts of its mouth ; although in such minute insects these organs must not be regarded as of primary importance, yet they often afford collateral evidence of affinity.

Ptiliodes Blackburni.

P. Blackburni, Matthews, Cist. Ent. iii. p. 41, pl. ii. fig. 3, 1882.

L. c. $\frac{5}{16}$ lin. = 0.63 mm.—Elongato-oblongus, sat depressus, haud nitidus, castaneus, pilis brevibus aureis vestitus ; capite permagno, lato, prominenti, confertissime tuberculato sive alutaceo ; oculis modicis, haud prominentibus ; pronoto capite parum latiori, vix longiori, pone medium latissimo, confertissime tuberculato sive alutaceo, lateribus leviter rotundatis, margine basali rectâ, angulis fere rectis ; elytris suboblongis, capite atque pronoto parum longioribus, haud latioribus, prope

humeros latissimis, translucidis, alis subtus quasi vittis nigris visis, confertissime asperatis, lateribus leviter marginatis, apicibus valde rotundatis, extremis albidis; abdomine sat elongato, segmentis tribus expositis, late flavo; pedibus atque antennis brevibus, late flavis.

Body elongate-oblong, rather depressed, not shining, castaneous, clothed with short golden hair.

Head prominent, very large and broad, closely tuberculate or alutaceous; eyes moderate, not prominent: antennæ short, bright yellow.

Thorax rather broader, not longer, than the head, widest behind the middle, very closely tuberculate or alutaceous, sides slightly rounded, basal margin straight, with the angles nearly right angles.

Scutellum moderate, triangular.

Elytra suboblong, rather longer but not broader than the head and thorax, widest near the shoulders, translucent, showing the wings like black vittæ below, very closely asperate, sides faintly margined, extremities much rounded, with the edges white.

Abdomen bright yellow, elongate, with three segments exposed.

Legs short, bright yellow.

Under parts rufo-castaneous, ventral segments bright yellow.

Habitat, Polynesia: Sandwich Islands. Found by Mr. Blackburn under the bark of a dead tree near Honolulu.

GEN. XVI.—NEPHANES.

Nephanes pubescens.

N. pubescens, Matthews, Trans. Am. Ent. Soc. xi. p. 141, 1884: Biol. Cent.-Amer. Col. ii. pt. i. p. 145, 1888.

L. c. $\frac{4}{16}$ — $\frac{5}{16}$ lin. = 0.50-0.63 mm. — Oblongus, sat depressus, haud nitidus, castaneus, pilis griseis dense vestitus; capite magno, antice sat producto, confertim tuberculato; oculis modicis, prominentibus: pronoto sat parvo, transverso, capite haud longiori, vix latiori, pone medium latissimo, tuberculis minutis, irregulariter dispositis, interstitiisque reticulatis, sat nitidis, ornato, lateribus leviter rotundatis, margine basali fere recta, angulis obtusis: elytris brevibus, capite atque pronoto nec longioribus neque latioribus, ad apices latissimis, ordinibus transversis leviter sed confertim asperatis, lateribus fere rectis, apicibus latis, fere rectis; abdomine longius exserto, obtuso, dilutiori: pedibus atque antennis longis, late flavis.

Body oblong, rather depressed, not shining, castaneous, thickly covered with grey hair.

Head large and broad, produced in front, closely and minutely tuberculate; eyes moderate and prominent; antennæ long, bright yellow.

Thorax rather small, transverse, not longer and scarcely broader than the head, widest behind the middle, ornamented with minute tubercles irregularly placed, interstices reticulated and rather shining, sides slightly rounded, basal margin nearly straight, with the angles obtuse.

Scutellum large, triangular, closely asperate.

Elytra short, not longer or broader than the head and thorax, widest at the apex, finely and closely asperated in transverse rows, sides nearly straight, extremities broad and nearly straight.

Abdomen considerably exposed, obtuse and pale.

Legs long, bright yellow.

Under parts dusky testaceous, rather deeply reticulated.

Differs from all other species of this genus in its elongate form and dense pubescence.

Habitat, Central America. Found near Capetillo in Guatemala by Mr. Champion.

TRIBE Ptiliina.

GEN. XVII.—PTILIUM.

The thirteen species now added to this genus comprise representatives of all the existing groups: *P. Simsoni* is allied to the beautiful *P. Halidayi* and *P. angulicollis*; *P. Hornianum* to *P. Kunzei*; *P. incognitum* and *P. fissicollis* belong to the group which exhibit longitudinal lines on the thorax, as *P. exaratum*; the remaining species, with one exception, may be referred to the group of which *P. Spencei* is the most common exponent. The exception I allude to exists in *P. obsoletum*; in this we meet with a very abnormal form, differing from all the other species both in general appearance, and in the almost total absence of eyes. It is represented by a single individual in the collection of Herr E. Reitter; and, since on that account dissection was unattainable, I have placed it in this somewhat heterogeneous genus.

I call *Ptilium* a heterogeneous genus on account of the dissimilarity of its component groups, but at the same time I cannot clearly see how these groups can be accurately defined as separate genera. Col. Motschulsky made an attempt to do this, and I would gladly adopt his divisions, if definition was possible. Let us take as an example the genus *Ptiliola*, originally named by Prof. Haldeman. Motschulsky mentions *P. Kunzei* as the type of *Ptiliola*, and could the genus be restricted to that species and *P. Collani* it would be sufficiently distinct; but it must also include *P. Spencei* with its numerous allies, many of which more closely resemble *P. exaratum*, and those with longitudinal lines on the thorax. And similar ambiguity exists even more palpably in some of his other divisions. On the whole I think that species whose affinities are so widely distributed and so closely interlaced should be comprised under a single genus, rather than divided into various indefinite genera.

The foregoing remarks were written a long time before I had seen Dr. Flach's 'Essay,' which I have noticed in the preface; and I do not consider it worth while to alter them to any great extent, as they still express my own ideas. As far as the genus *Ptilium* is concerned, Dr. Flach has, in the 'Essay' referred to, followed the lines laid down by Col. Motschulsky in the *Etud. Ent.* iv. p. 17, 1855, but has substituted fresh names for many of those originally given by that author, and has added some new genera for species which he had himself first described. Among the latter he created the genus *Typhloptilium* to contain an insect which he had previously described in the *Deuts. Ent. Zeits.* 1886, p. 248, under the name of *Ptilium (Edipus)*. But I do not perceive that this species presents any characters sufficient to justify generic separation. The absence of eyes cannot of itself be considered conclusive, for in the well-known genus *Ptinella* this is merely a sexual distinction and exists only in certain species. And, moreover, in *P. (Edipus)* minute eyes possessing visual facets can be discovered by close investigation. I am speaking now of my own specimen; it is quite possible that others may be found in which the eyes are totally absent. Nevertheless, some more obvious anatomical character would be required to constitute a new genus. But, under any circumstances, the name *(Edipus)*, by which this insect was described by Dr. Flach in 1886, must be retained for the species.

SPECIERUM RECENTIORUM EPITOMA.

L. c. 0.35 mm.—Oblongum, convexum, castaneum, confertim tuberculatum, ped. læte, ant. obscure flavis	Hornianum.
L. c. 0.75 mm.—Ovale, convexum, piceum, pilis griseis dense vestitum, ped. atque ant. flavis	obscurum.
L. c. 0.87 mm.—Elongato-ovale, convexum, ferrugineum, pronoto tuberculis sat magnis distinctis ornato, ped. atque ant. prælongis, flavis	Columbianum.
L. c. 0.70 mm.—Elongatum, modice convexum, ferrugineum, pilis flavis dense vestitum, pronoto confertim tuberculato, ped. atque ant. brevibus, læte flavis	humile.
L. c. 0.60 mm.—Elongatum, modice convexum, haud nitidum, castaneum, pronoto lineâ longitudinali in disco impresso, ped. atque ant. læte flavis	fissicolle.
L. c. 0.70 mm.—Elongatum, rufo-castaneum, alutaceum, pronoto lineis tribus parallelis impresso, ped. atque ant. læte flavis	affine.
L. c. 1 mm.—Elongato-ovatum, nigro-piceum, valde convexum, confertim tuberculatum, pronoto ad basin fortiter constricto, lineis tribus parallelis sat profunde impresso, ant. longis, piceis, ped. longis, læte flavis	incognitum.
L. c. 0.60 mm.—Elongatum, fere lineari, modice convexum, ferrugineum, pilis flavis dense vestitum, pronoto confertim tuberculato, elytris longis, angustis, ped. longis, atque ant. læte flavis	planum.
L. c. 0.50 mm.—Elongatum, convexum, rufo-castaneum, pilis flavis vestitum, pronoto leviter tuberculato, elytris ovalibus, ped. longis, atque ant. brevibus, robustis, læte flavis	tropicum.
L. c. 0.70 mm.—Oblongo-ovale, læte castaneum, nitidum, pilis aureis vestitum, pronoto eleganter annulatim impresso, ped. atque ant. robustis, læte flavis	Simsoni.
L. c. 0.85 mm.—Elongato-ovale, valde convexum, nigro-piceum, pronoto modice tuberculato, oculis fere nullis, ant. prælongis, gracillimis, atque ped. piceis	Œdipus.
L. c. 0.50 mm.—Subovatum, piceum, confertim tuberculatum, pronoto ad latera rotundato, ant. longis, piceis, ped. longis, flavis	Smithsii.
L. c. 0.50 mm.—Elongato-ovale, rufo-testaceum, confertim tuberculatum, pronoto sat magno, ad basin triimpresso, ped. atque ant. brevibus, læte flavis	impressum.
L. c. 0.50 mm.—Elongato-ovatum, nitidulum, confertim tuberculatum, rufo-testaceum, pronoto ad basin triangulariter impresso, ped. atque ant. longis, flavis	rufo-testaceum.

Ptilium Smithsii.

P. Smithsii, Matthews, Ann. Mag. Nat. Hist. 6, xiii. p. 341, 1894.

L. c. $\frac{1}{4}$ lin. = 0.50 mm.—Subovatum, modice convexum, vix nitidum, confertim tuberculatum, piceum, pilis brevibus obscuris vestitum; *capite* magno, confertim tuberculato; oculis magnis, prominentibus; antennis sat longis, piceis; *pronoto* parvo, capite nec longiori neque latiori, prope medium latissimo, antice atque postice contracto, confertim tuberculato, interstitiis nitidis, lateribus rotundatis, vix marginatis, margine basali leviter reflexa, fere recta, angulis obtusis; *elytris* modicis, capite atque pronoto longioribus, haud latioribus, ante media latissimis, postice parum contractis, confertim tuberculatis, interstitiis nitidis, lateribus leviter marginatis, apicibus rotundatis, dilutioribus; abdomine parum exserto, piceo; pedibus sat longis, flavis.

Body subovate, moderately convex, scarcely shining, closely tuberculate, piceous, clothed with short dusky hair.

Head large, closely tuberculate: eyes large and prominent; antennæ rather long, piceous.

Thorax small, not longer or broader than the head, widest near the middle, contracted in front and behind, closely tuberculate with the interstices shining, sides rounded, slightly margined, basal margin slightly reflexed, nearly straight, with the angles obtuse.

Scutellum large, triangular, closely tuberculate.

Elytra moderate, longer but not broader than the head and thorax, widest before the middle, slightly contracted behind, closely tuberculate with the interstices shining, sides faintly margined, extremities rounded and paler.

Abdomen slightly exposed, piceous.

Legs rather long, yellow.

Under parts piceous, metasternum closely asperated.

Differs from *P. Kunzei* in its rounded thorax and elytra, finer sculpture, and much longer antennæ and legs.

Habitat, America. Taken not uncommonly by Mr. Smith in the West Indian Island of Grenada.

Ptilium Hornianum.

P. Hornianum, Matthews, Trans. Am. Ent. Soc. xi. p. 143, 1884; Biol. Cent.-Amer. Col. ii. pt. i. p. 146, 1888.

L. c. $\frac{3}{16}$ lin. = 0.37 mm.—Oblongum, convexum, sat nitidum, castaneum, pilis griseis vestitum; *capite* magno, lato, antice producto, tuberculis minutis confertim obtecto: oculis modicis, prominentibus; *pronoto* parvo, transverso, capite breviori, vix latiori, ad medium latissimo, sat nitido, tuberculis minutis confertim obtecto, lateribus leviter rotundatis, et leviter marginatis, margine basali recta, et leviter reflexa, angulis fere rectis; *elytris* oblongis, convexis, capite atque pronoto parum longioribus, vix latioribus, ad media latissimis, profunde et reticulatim asperatis, lateribus minime rotundatis, apice lato, integro, valde rotundato; abdominis pygidio exserto, dilutiori; pedibus modicis, late flavis: antennis sat longis, obscure flavis.

Body oblong, convex, rather shining, castaneous, clothed with grey hair.

Head large and broad, elongate anteriorly, closely covered with minute tubercles; eyes moderate, prominent; antennæ rather long, dusky yellow.

Thorax small, transverse, shorter and scarcely broader than the head, widest at the middle, rather shining, closely covered with small tubercles, sides slightly rounded and slightly margined, basal margin straight and slightly reflexed, with the angles nearly right angles.

Scutellum small, triangular, closely asperated.

Elytra oblong, convex, rather longer, but scarcely broader than the head and thorax, widest at the middle, deeply and reticulately asperated, sides scarcely rounded, apex broad, entire, and much rounded.

Abdomen with the pygidium only exposed, pale.

Legs moderate, bright yellow.

Under parts pale castaneous, with the mouth, hinder part of the metasternum, and terminal segments of the venter flavescent.

Differs from *P. rugulosum* in its much smaller size, fainter sculpture, shining surface, and bright yellow legs; and from all others in its minute size.

Habitat, North and Central America. Found in Texas, near Columbus, by Mr. H. Ulke, and subsequently by Mr. Champion in Guatemala.

This interesting species is the smallest of all the *Trichopterygidae* hitherto discovered except *Nanosella fungi*.

Ptilium obscurum.*P. obscurum*, Matthews, Cist. Ent. ii. p. 171, 1877.

L. c. $\frac{1}{16}$ lin. = 0.75 mm. Ovale, convexum, piceum, pilis sat longis, griseis, sat dense vestitum; *capite* magno, antice obtuso, tuberculis parvis, ordinibus rectis, interstitiisque nitidissimis, leviter reticulatis, ornato; oculis sat magnis, haud prominentibus; *pronoto* magno, capite longiori et latiori, ad basin latissimo, tuberculis magnis, ordinibus fortiter sinuatis, interstitiisque nitidissimis, leviter reticulatis, ornato, lateribus leviter rotundatis, margine basali leviter rotundata, angulis obtusis; *elytris* sat brevibus, capite atque pronoto haud longioribus, vix latioribus, ad humeros pronoto angustioribus, pone media latissimis, ordinibus sinuatis, profunde asperatis, lateribus marginatis, leviter rotundatis, apicibus validissime rotundatis, late dilutioribus; abdominis pygidio solo exserto; pedibus late flavis; antennis modicis, flavis, articulo secundo incrassato.

Body oval, convex, piceous, rather thickly clothed with grey hair.

Head large, obtuse in front, ornamented with small tubercles in straight rows, interstices shining and lightly reticulated; eyes rather large, not prominent; antennæ moderate, yellow, with the second joint enlarged.

Thorax large, longer and broader than the head, widest at the base, ornamented with large tubercles in strongly sinuated rows, interstices very shining and lightly reticulated, sides slightly rounded, basal margin slightly rounded, with the angles obtuse.

Scutellum short, obtusely triangular, moderately asperated.

Elytra rather short, not longer and scarcely broader than the head and thorax, narrower than the thorax at the shoulders, widest behind the middle, deeply asperated in wavy rows, sides margined and slightly rounded, extremities very much rounded and broadly pale.

Abdomen with the pygidium only exposed.

Legs moderate, bright yellow.

Under parts pitchy castaneous, with the mouth, coxæ, margin of the metasternum, and terminal segments of the venter yellow.

Differs from *P. marginatum* in its smaller size, thorax without a margin, and piceous colour; and from *P. Columbianum* in its smaller size, broader form, broader and much shorter head and thorax, short obtuse scutellum, and piceous colour.

Habitat, North America. Found in a fungus by Mr. G. R. Crotch in British Columbia.

Ptilium Columbianum.*P. Columbianum*, Matthews, Cist. Ent. ii. p. 169, 1877.

L. c. $\frac{7}{16}$ lin. = 0.87 mm.—Elongato-ovale, convexum, ferrugineum, vel castaneum, pilis flavescens, sat dense vestitum; *capite* modico, antice triangulariter elongato, tuberculis ordinibus transversis ornato; oculis magnis, prominentibus; *pronoto* sat parvo, capite parum breviori et angustiori, ad basin aliquantum latissimo, tuberculis sat magnis, distinctis, irregulariter dispositis, interstitiisque nitidis, vix reticulatis ornato, lateribus marginatis et valde rotundatis, margine basali rotundata, leviter reflexa, nitida, angulis rotundatis; *elytris* longis, elongato-ovalibus, capite atque pronoto fere duplo longioribus, parum latioribus, pone media latissimis, ordinibus distinctis, fortiter sinuatis, asperatis, lateribus leviter marginatis et leviter rotundatis, apicibus latis, parum rotundatis; abdominis pygidio solo exserto; pedibus longis, late flavis; antennis praelongis, gracilibus, late flavis.

Body elongate-oval, convex, ferruginous or castaneous, rather thickly clothed with flavescent hair.

Head moderate, triangularly produced in front, ornamented with tubercles in transverse rows; eyes large and prominent; antennæ very long and slender, bright yellow; maxillary palpi large, yellow.

Thorax rather small, rather shorter and narrower than the head, slightly widest at the base, ornamented with rather large, distinct, and irregular tubercles, interstices shining and faintly reticulated, sides margined and much rounded, basal margin rounded, slightly reflexed, shining, with the angles rounded.

Scutellum elongate, acuminate, closely and deeply asperated.

Elytra long, elongate-oval, nearly twice longer, and rather broader than the head and thorax, widest behind the middle, asperated in distinct and very wavy rows, sides slightly margined and slightly rounded, extremities broad and slightly rounded.

Abdomen with the pygidium only exposed.

Legs long, bright yellow.

Under parts entirely castaneous.

Differs from *P. Sharpi* in its larger and broader thorax widest at the base, and much fainter sculpture; from *P. Spencei* in its much larger size, shape of its thorax, and very long antennæ.

Habitat, North America. Found by Mr. G. R. Crotch in British Columbia, and in California.

This species varies in the width of the elytra; probably a sexual distinction, as in *P. Spencei*.

Ptilium humile.

P. humile, Matthews, Cist. Ent. ii. p. 170, 1877.

L. c. $\frac{5}{16}$ – $\frac{6}{16}$ lin. = 0.63–0.75 mm.—Elongatum, modice convexum, ferrugineum, pilis flavis sat dense vestitum; capite sat magno, antice elongato, tuberculis parvis confertissime obtectum; oculis magnis, prominulis, nigrescentibus; pronoto parvo, brevi, capite breviori, vix latiori, fere ad medium latissimo, tuberculis sat magnis confertissime oblecto, lateribus valde rotundatis, late marginatis atque setis duabus brevibus pone media instructis, margine basali leviter sinuata, reflexa, nitida, angulis obtusis; elytris elongato-ovalibus, capite atque pronoto plus quam duplo longioribus, vix latioribus, ad media latissimis, ordinibus sinuatis, sat remotis, modice asperatis, apice ovali, angulis suturalibus fere rectis; abdominis pygidio solo exserto; pedibus atque antennis brevibus, robustis, læte flavis.

Body elongate, moderately convex, ferruginous, rather thickly clothed with yellow hair.

Head rather large, produced in front, very closely covered with small tubercles; eyes large and rather prominent, black; antennæ short, rather robust, bright yellow.

Thorax small and short, shorter and scarcely broader than the head, widest near the middle, very closely covered with rather large tubercles, sides much rounded and broadly margined, furnished with two short erect setæ behind the middle, basal margin slightly sinuated, reflexed, and shining, with the angles obtuse.

Scutellum moderate, elongate, acuminate, closely asperated.

Elytra elongate-oval, more than twice longer, but scarcely broader than the head and thorax, widest at the middle, moderately asperated in remote wavy rows, apex entire, oval, with the sutural angles nearly right angles.

Abdomen with the pygidium only exposed.

Legs short, robust, bright yellow.

Under parts entirely ferruginous.

Differs from *P. Columbianum* in its much smaller size, narrower form, short robust antennæ, and in the sculpture of its thorax; from *P. Spencei* in its narrower form, shorter antennæ, and in sculpture.

Habitat, North America. Found near San Diego, in California, by Mr. G. R. Crotch.

Ptilium fissicolle.

P. fissicolle, Reitter, Deuts. Ent. Zeit. 1884, p. 116.

L. c. $\frac{5}{16}$ lin. = 0.63 mm. — Elongatum, modice convexum, haud nitidum, leviter tuberculatum, castaneum, pilis flavescens vestitum, pronoto sulco unico mediali impresso; capite modico, confertim tuberculato; oculis modicis, sat prominentibus; pronoto modico, pone medium latissimo, confertissime tuberculato, sulco profundo mediali, nec ad marginem anteriorem neque basalem extenso, impresso, lateribus antice valde rotundatis, postice contractis, margine basali fere recta, angulis acutis; elytris ovatis, capite atque pronoto sesqui longioribus, parum latioribus, prope media latissimis, leviter et confertissime asperatis, lateribus leviter marginatis, apice attenuato; pygidio minime exserto; pedibus atque antennis modicis, late flavis, tibiis anterioribus ad apices dilatatis.

Body elongate, moderately convex, not shining, closely tuberculate, castaneous, clothed with yellow hair, with a single deep central channel on the thorax.

Head moderate, closely tuberculate; eyes moderate, rather prominent; antennae moderate, bright yellow.

Thorax moderate, widest behind the middle, very closely tuberculate, with a deep central channel not extending to the anterior or basal margins, sides much rounded in front and contracted behind, basal margin nearly straight, with the angles acute.

Scutellum rather large, triangular, closely asperated.

Elytra ovate, one-half longer and rather broader than the head and thorax, widest near the middle, finely and very closely asperate, sides slightly margined, apex attenuated.

Abdomen with the pygidium very little exserted, flavescens.

Legs moderate, bright yellow, anterior tibiae dilated at the apex.

Under parts castaneous, mouth, prothorax, and apex of the venter pale, metasternum closely tuberculate.

Differs from others in the single deep central channel on the thorax without any lateral impressions.

Habitat, Europe. Found in Corfu (Herr Reitter).

Ptilium affine.

P. affine, Erichson, Ins. Deutsch. iii. p. 27, 1848.

L. c. vix $\frac{6}{16}$ lin. = 0.70 mm. — Elongatum, rufo-castaneum, alutaceum, nitidum, pilis perbrevibus argenteis vestitum, lineis lateralibus pronoti parallelis; capite sat parvo, prominenti, obtuse triangulari, indistincte alutaceo, fere glabro; oculis parvis, haud prominentibus; pronoto sat parvo, capite latiori et parum longiori, pone medium latissimo, basin versus abrupte contracto, indistincte alutaceo, linea mediali lata et profunda, nec ad apicem, neque ad basin extensa, et utrinque linea lata, brevi, parallela, nec ad basin, neque ultra medium discum extensa, impresso, lateribus marginatis, antice rotundatis, postice contractis, margine basali recta, angulis rectis; elytris ovalibus, capite atque pronoto sat latioribus, plus quam sesqui longioribus, prope media latissimis, ordinibus transversis distincte sed levissime asperatis, interstitiis alutaceis, lateribus marginatis, apice integro, obtuso; pedibus atque antennis sat robustis, late flavis.

Body elongate, rufo-castaneous, alutaceous, moderately shining, clothed with silver hair, with the lines on the thorax all parallel.

Head rather small, prominent, obtusely triangular, indistinctly alutaceous, almost smooth; eyes small, not prominent; antennae rather robust, bright yellow.

Thorax rather small, broader and rather longer than the head, widest beyond the middle, abruptly contracted towards the base, indistinctly alutaceous, with a deep and broad medial line, not extending to the base or apex, and a shorter parallel line on each side extending from the middle of the disc nearly to the basal margin, sides margined, rounded in front and contracted behind, basal margin straight, with the angles right angles.

Scutellum small, triangular, with a faint longitudinal impression on the disc.

Elytra oval, rather broader and more than one-half longer than the head and thorax, widest near the middle, distinctly but faintly asperated in transverse rows, with the interstices alutaceous, sides marginod, apex entire, obtuse.

Legs rather robust, bright yellow.

Under parts rufo-castaneous, with the coxæ bright yellow.

Differs from *P. incognitum* in its much smaller size, rufo-castaneous colour, less convex form, totally diverse sculpture, as well as in the short and broad lateral lines on the thorax. From the other allied species it is distinguished by the lines on the thorax being parallel to each other.

Habitat, Europe. Found by Mr. Crotch in Cambridgeshire, and by Mr. Morris near Lewes, in Sussex.

This insect appears to be the true *Ptilium affine* of Erichson; it agrees in size, colour, and sculpture with his description in the "Ins. Deutsch." I had formerly received types from two continental sources labelled "*P. affine*," which were identical with the insect I found in Wicken Fen, and described as *P. affine* in my former work, the 'Trichopterygia Illustrata.'

I have now separated this species under the name *P. incognitum*. It is much larger than the *P. affine* of Dr. Erichson, nearly black, much more convex, and very diverse in sculpture.

The discrepancy of size and form existing between my own specimen and those subsequently taken in this country first drew my attention to the subject, and convinced me that they could not belong to the same species.

The change of nomenclature does not, however, affect any but my own specimen; all the other *Ptilia* with parallel lines on the thorax which I have hitherto seen belong to *Ptilium affine*, Erichs.

Ptilium incognitum.

P. incognitum, Matthews, Ann. Mag. Nat. Hist. 6, iii. p. 195, 1889.

P. affine, Matthews nec Erichson, Trich. Illust. p. 99, pl. xxii. fig. 5, 1872.

L. c. $\frac{7}{16}$ – $\frac{8}{16}$ lin. = 0.87–1 mm. Elongato-ovatum, nigro-piceum, valde convexum, leviter et confertim tuberculatum, pilis brevibus argenteis sparse vestitum, pronoto ad basin constricto, lineis tribus rectis, lateralibus parallelis, in disco impresso; capite parvo, convexo, antice elongato, leviter confertimque tuberculato; oculis parvis, haud prominentibus; pronoto modico, capite parum latiori, vix longiori, pone medium latissimo, basin versus abrupte fortiterque constricto, confertim tuberculato, lineis tribus rectis in disco distinctissime impresso, marginem utramque haud attingentibus, mediâ longissimâ, lateralibus parallelis atque ad extremum utrumque minute uncinnatis, margine basali fere rectâ, angulis prominulis, acutis, angulis anterioribus valde deflexis; clytris ovatis, convexissimis, capite atque pronoto multo latioribus, et duplo longioribus, pone mediâ latissimis, ad humeros valde attenuatis, leviter confertimque tuberculatis, atque tuberculis majoribus hic et illic ornatis, lateribus marginatis, apice obtuso, dilutiori; abdomine parum exserto, valde obtuso, pedibus longioribus, late flavis; antennis longis, piceis, ad bases dilutioribus.

Body elongate-ovate, nigro-piceous, very convex, finely and closely tuberculate, sparingly clothed with short silver hair, with three straight parallel lines on the thorax.

Head small, convex, elongate anteriorly, finely and closely tuberculate; eyes small, not prominent; antennæ long, piceous, becoming paler towards the base.

Thorax moderate, rather broader but scarcely longer than the head, widest behind the middle, abruptly and strongly constricted towards the base, finely and closely tuberculate, with three distinct straight parallel lines on the disc, not extending to either margin, the middle line being the longest, and the lateral lines parallel to the middle and faintly hooked at each extremity, basal margin nearly straight, with the angles prominent and acute, anterior angles much reflexed.

Scutellum large, elongate, triangular, elevated at the base, and marked with a short medial channel.

Elytra ovate, very convex, much broader and twice longer than the head and thorax, widest beyond the middle, very much attenuated at the shoulders, finely and closely tuberculate, ornamented with larger tubercles placed singly at different intervals, sides margined, apex obtuse and paler.

Abdomen slightly exerted, very obtuse.

Legs rather long, bright yellow.

Under parts castaneous, with the apex of the venter pale, and the coxæ yellow.

Differs from other species in its large size, dark colour, very convex elytra narrowed at the shoulders, and peculiar sculpture. In addition to the foregoing characters it is distinguished from *P. affine* by its much larger size, narrowness of the lines on the thorax, dark antennæ, and less dense pubescence.

Habitat, Europe. Found in England by myself in Wicken Fen, Cambridgeshire; also in Holland by M. Neervoort van de Poll.

This fine species is one of the largest of the present genus, and very distinct in appearance. It seems to be very rare, since among all the continental collections which I have examined it is represented only in those of M. Reiche and M. Neervoort van de Poll, and in both was labelled "*Pt. affine*;" under that name, therefore, I described the species in the '*Trichopterygia Illustrata*.'

Since then several specimens of the true *P. affine* of Erichson have been taken in various parts of this country by Mr. G. R. Crotch, and Mr. Morris of Lewes; the occurrence of these smaller, paler, and more pubescent insects induced me to compare them carefully with Erichson's description, with which I found that they entirely agreed. It consequently became necessary to distinguish our present species by another name, as I now have done.

Ptilium planum.

P. planum, Matthews, Trans. Am. Ent. Soc. xi. p. 144, 1884; Biol. Cent.-Amer. Col. ii. pt. i. p. 146, 1888.

L. c. $\frac{5}{16}$ lin. — 0.63 mm.—Elongatum, fere lineare, modice convexum, haud nitidum, ferrugineum, pilis flavis dense vestitum; capite magno, antice producto, confertim et leviter tuberculato; oculis magnis, prominentibus; pronoto parvo, transverso, capite breviori, haud latiori, ante medium latissimo, confertim et leviter tuberculato, lateribus basin versus fortiter contractis, modice marginatis, margine basali fere recta, angulis rectis; elytris longis, angustis, capite atque pronoto plus quam sesqui longioribus, vix latioribus, pone media latissimis, ordinibus transversis confertim et leviter asperatis, lateribus levissime rotundatis, apicibus latis, rotundatis; pedibus longis, gracilibus, flavis; antennis sat brevibus, lute flavis.

Body elongate, almost linear, moderately convex, not shining, ferruginous, densely clothed with yellow hair.

Head large, produced in front, finely and closely tuberculate; eyes large and very prominent; antennæ rather short, bright yellow.

Thorax small, transverse, shorter and not broader than the head, widest before the middle, closely and finely tuberculate, without impressions, sides much contracted towards the base, moderately margined, basal margin nearly straight, with the angles right angles.

Scutellum large, triangular, closely asperate.

Elytra elongate-oval, narrow, more than one-half longer but scarcely broader than the head and thorax, widest beyond the middle, finely and closely asperated in transverse rows, sides faintly rounded, apex broad and rounded.

Abdomen with the pygidium exposed, flavescent.

Legs long and slender, yellow.

Under parts entirely flavescent.

Differs from *P. foveolatum* in its larger size, broader form, plain thorax, and deeper general sculpture; and from all others in its narrow elongate form and small size.

Habitat, Central America. Found near Capetillo in Guatemala by Mr. Champion.

Ptilium tropicum.

P. tropicum, Matthews, Trans. Am. Ent. Soc. xi. p. 144, 1884; Biol. Cent.-Amer. Col. ii. pt. i. p. 147, pl. iii. fig. 23, 1888.

L. c. $\frac{5}{15}$ lin. = 0.63 mm.—Elongatum, modice convexum, haud nitidum, rufo-testaceum, pilis flavis dense vestitum; *capite* magno, antice prominenti, confertim et leviter tuberculato; oculis magnis, prominentibus; *pronoto* parvo, transverso, capite angustiori, haud longiori, ante medium latissimo, postice fortiter contracto, confertim et leviter tuberculato, lateribus modice marginatis, margine basali fere rectâ, angulis fere rectis; *elytris* ovalibus, elongatis, capite atque pronoto plus quam sesqui longioribus, parum latioribus, ordinibus transversis leviter sed confertim asperatis, apicibus latis, rotundatis; abdominis pygidio solo exserto; pedibus longis, gracilibus, flavis; antennis sat brevibus, læte flavis.

Body elongate, moderately convex, not shining, rufo-testaceous, thickly clothed with yellow hair.

Head large, prominent in front, closely but not deeply tuberculate; eyes large and prominent; antennæ rather short, robust, bright yellow.

Thorax small, transverse, narrower and not longer than the head, widest before the middle, strongly contracted posteriorly, closely but not deeply tuberculate; sides moderately margined, basal margin nearly straight, with the angles nearly right angles.

Scutellum large, triangular, closely asperated.

Elytra elongate-oval, more than one-half longer but broader than the head and thorax, widest beyond the middle, lightly but closely asperated in transverse rows, extremities broad and rounded.

Abdomen with the pygidium only exposed.

Legs long and slender, yellow.

Under parts entirely testaceous.

Differs from *P. foveolatum* in its larger size, broader form, thorax not foveolate, and rather deeper sculpture; from *P. planum* in its shorter and broader form, and short and very robust antennæ.

Habitat, Central America. Found by Mr. Champion at Bugaba in Panama.

Ptilium Simsoni.

P. Simsoni, Matthews, Cist. Ent. ii. p. 327, 1878.

L. c. $\frac{5}{16}$ – $\frac{6}{16}$ lin. = 0.63–0.75 mm.—Oblongo-ovale, læte castaneum, nitidum, pilis aureis vestitum; *capite* sat magno, foveis umbilicatis, interstitiisquo nitidis,

ornato; oculis magnis, haud prominentibus; *pronoto* modico, capite paulum longiori, vix latiori, ad medium latissimo, foveis magnis, profundo umbilicatis, sive annulis parvis, ordinibus transversis, sat remotis, catenulatum dispositis, interstitiisque glabris, nitidis, ornato, impressione lata, profunda, longitudinali, antice divergenti, a basi usque ad medium extensa, in disco utrinque notato, lateribus antice rotundatis, postice leviter constrictis, margine basali leviter rotundata et leviter reflexa, angulis fere rectis; *elytris* ovatis, integris, translucidis, capite atque *pronoto* longioribus et paulo latioribus, ad media latissimis, ordinibus distinctis, sinuatis, transversis, asperatis, apice lato, obtuso, dilutiori; abdominis pygidio solo exserto; pedibus atque antennis robustis, laete flavis.

Body oblong-oval, bright castaneous, shining, clothed with golden hair.

Head rather large and prominent, rounded in front, ornamented with umbilicated foveae, interstices shining; eyes large, not prominent; antennae rather long, robust, bright yellow.

Thorax moderate, rather longer but scarcely broader than the head, widest at the middle, ornamented with large and deep umbilicated foveae, or small impressed rings, arranged like a chain in remote transverse rows, interstices smooth and shining, with a broad deep longitudinal impression on each side of the disc, extending from the base to the middle and diverging anteriorly, sides rounded anteriorly and slightly constricted towards the base, basal margin slightly rounded and reflexed, with the angles nearly right angles.

Scutellum moderate, triangular, and deeply punctured.

Elytra entire, ovate, translucent, longer and slightly broader than the head and thorax, widest at the middle, asperated in distinct wavy transverse rows, apex broad, obtuse, and paler.

Wings seen through the elytra as narrow black vittae.

Abdomen with the pygidium slightly exposed.

Legs robust, bright yellow.

Under parts castaneous, with the terminal segments of the venter paler.

Differs from all other species in the remarkable and beautiful sculpture of the thorax.

Habitat, Australasia. Found in Tasmania by Mr. Simson.

This beautiful species belongs to the section of the genus which contains *P. Halidayi* and *P. angulicollis*, both of which are also remarkable for elegance and rarity.

Ptilium impressum.

P. impressum, Matthews, Ann. Mag. Nat. Hist. 6, xiii. p. 341, 1894.

L. c. $\frac{5}{16}$ lin. = 0.63 mm. — Elongato-ovale, modice convexum, vix nitidum, confertim tuberculatum, rufo-testaceum, pilis brevibus flavescens vestitum; capite magno, prominenti, confertim tuberculato; oculis modicis, sat prominentibus; antennis brevibus, robustis, laete flavis; *pronoto* parvo, capite nec longiori, neque latiori, confertim tuberculato, impressione sat profunda, ovali, ad medium, atque alia rotundata utrinque ad basin notato, lateribus leviter marginatis, antice rotundatis, postice sat contractis, ad media latissimis, margine basali recta, angulis fere rectis; *elytris* longis, capite atque *pronoto* plus quam sesqui longioribus, parum latioribus, prope media latissimis, sat leviter, confertissime asperatis, lateribus leviter marginatis, apice integro, obtuso; abdomine haud exserto; pedibus brevibus, laete flavis.

Body elongate-oval, moderately convex, slightly shining, closely tuberculate, clothed with short flavescent hair.

Head large and prominent, closely tuberculate; eyes rather prominent; antennae short, robust, bright yellow.

Thorax small, not longer or broader than the head, closely tuberculate, marked at the base with a rather deep oval impression in the middle, and a smaller round impression on each side near the hinder angles, sides faintly margined, widest near the middle, rounded in front and rather contracted behind, basal margin straight, with the angles nearly right angles.

Scutellum moderate, triangular, closely asperated.

Elytra long, more than one-half longer and rather broader than the head and thorax, widest near the middle, very closely and rather finely asperate, sides faintly margined, apex entire, obtuse.

Abdomen not exposed.

Legs short, bright yellow.

Under parts testaceous.

Differs from the following species in the greater size and peculiar sculpture of its thorax, and much shorter legs and antennæ.

Habitat, America. One specimen found by Mr. Smith in the West Indian Island of St. Vincent.

Ptilium rufo-testaceum.

P. rufo-testaceum, Matthews, Ann. Mag. Nat. Hist. 6, xiii. p. 341, 1894.

L. c. $\frac{3}{16}$ lin. = 0.50 mm.—Elongato-ovatum, nitidulum, rufo-testaceum, confertissime tuberculatum, fere alutaceum, pronoto ad basim triangulariter impresso, pilis brevissimis aureis sparse vestitum; *capite* magno, prominenti, antice producto, confertim tuberculato; oculis nigris, magnis, prominentibus; antennis longis, gracilibus, late flavis; *pronoto* sat parvo, capite nec longiori neque latiori, ante medium latissimo, confertim tuberculato, fere alutaceo, linea mediali abbreviata, ad basim triangulariter dilatata, impresso, lateribus antice valde rotundatis, postice fortiter contractis, margine basali recta, angulis acutis, prominentibus; *clytris* capite atque pronoto parum longioribus atque latioribus, prope media latissimis, leviter confertissime asperatis, lateribus marginatis, apice integro; abdomine haud exserto; pedibus sat longis, flavis.

Body elongate-ovate, moderately shining, rufo-testaceous, very closely tuberculate, almost alutaceous, marked at the base of the thorax with an elongate triangular impression, sparingly clothed with short golden hair.

Head large, very prominent, produced in front, closely tuberculate; eyes black, large, and prominent; antennæ long and slender, bright yellow.

Thorax rather small, not longer or broader than the head, widest before the middle, closely tuberculate, or alutaceous, with an abbreviated middle line triangularly dilated at the base, sides much rounded in front and strongly contracted towards the base, basal margin straight, with the angles acute and prominent.

Scutellum large, triangular, depressed in the middle, closely asperated.

Elytra rather longer and broader than the head and thorax, widest near the middle, finely and closely asperate, sides margined, apex entire.

Abdomen not exposed.

Legs robust, bright yellow.

Under parts testaceous.

Differs from *P. impressum* in its smaller size and narrower form, large triangular head, large and prominent eyes, smaller and strongly contracted thorax, with a triangular impression at the base and a short central line, long antennæ, and legs.

Habitat, America. One specimen found in the West Indian Island of Grenada by Mr. Smith.

Ptilium Œdipus, Flach.

P. Œdipus, Flach, Dent. Ent. Zeit. 1886, p. 248.

P. obcecatum, Matthews, Ann. Mag. Nat. Hist. 6, iii. 1889, p. 195.

L. c. $\frac{6}{15}$ – $\frac{7}{15}$ lin. = 0.75–0.87 mm.—Elongato-ovale, valde convexum, nitidulum, nigro-piceum, pilis pallidis sparsissime vestitum; capite magno, antice producto, minute et confertim tuberculato, oculis longe retropositis, valde minutis; pronoto modico, capite parum longiori et latiori, ante basin latissimo, tuberculis modicis, ordinibus irregularibus, transversis dispositis, interstitiisque leviter reticulatis ornato, lateribus rotundatis, leviter marginatis, margine basali fere recta, angulis obtusis; elytris ovalibus, capite atque pronoto longioribus, vix latioribus, ad media latissimis, ordinibus transversis, irregularibus sat leviter asperatis, interstitiis reticulatis, lateribus leviter marginatis, apicibus rotundatis, dilutioribus; pedibus modicis, piceis; antennis elongatis, gracillimis, piceo-testaceis.

Body elongate-oval, very convex, rather shining, pitchy black, sparingly clothed with pale hairs.

Head large, produced in front, minutely and closely tuberculate; eyes very minute, placed far back; antennæ very long and slender, pitchy testaceous.

Thorax rather longer and broader than the head, widest before the base, ornamented with moderate tubercles in irregular transverse rows, interstices faintly reticulated, sides slightly margined and rounded, basal margin nearly straight, with the angles obtuse.

Scutellum triangular, asperated.

Elytra oval, longer but scarcely broader than the head and thorax, widest at the middle, lightly asperated in irregular transverse rows, interstices reticulated, sides slightly margined, extremities rounded and paler.

Legs moderate, piceous.

Abdomen with the pygidium exposed, piceous.

Under parts pitchy black.

Differs from *P. Spencei* in its large size, much finer sculpture, rounded sides of the thorax, very long slender antennæ, and minute eyes.

Habitat, Europe. Found near Taygetus, in the Morea, by Herr Brenske (Reitter), and in Central Bosnia.

GEN. XIX.—ACTIDIUM.

Since 1872 the number of the species of Actidium has increased fourfold. Of the novelties now recorded, six are due to the researches of Messrs. Crotch and Casey in California, which appears to be the head-quarters of the genus. With one exception the Californian species belong to the convex and shining group, represented in Europe by *A. Boudierii*. *A. lineare*, from New Zealand, and *A. Fowlerianum*, from Central America, are narrow and pubescent like the European *A. coarctatum*. *A. Sharpianum*, from the Sandwich Islands, is so unlike any other species that it may possibly belong to a distinct genus; but, as the specimen is at present unique, I have not ventured to dissect it.

SPECIERUM RECENTIORUM EPITOMA.

- L. c. 0.40 mm.—Elongatum, valde angustum, fuscescens, pilis brevibus sericeis dense vestitum, pronoto brevi, transverso, elytris longis, valde attenuatis, ped. atque ant. flavis **Fowlerianum.**
- L. c. 0.50 mm.—Lineare, elongatum, convexum, nigrescens, pilis griseis dense vestitum, confertim et profunde punctatum, elytris linearibus, ped. atque ant. late flavis **lineare.**

- L. c. 0.50 mm.—Ovale, convexum, nigrum, nitidissimum, pilis brevissimis sparsissime indutum, levissime punctatum, elytris ovalibus, ped. atque ant. nigro-piceis . . . **politum.**
- L. c. 0.50 mm.—Elongato-ovatum, convexum, nitidulum, nigro-piceum, leviter alutaceum, pilis sparse indutum, pronoto basin versus fortissime contracto, elytris ovatis, ped. atque ant. flavis . . . **rotundicolle.**
- L. c. 0.50 mm.—Elongato-ovatum, convexum, haud nitidum, nigro-piceum, profunde alutaceum, pronoto sat magno, elytris ovatis, striâ suturali distinctâ, ped. atque ant. nigro-piceis . . . **robustulum.**
- L. c. 0.50 mm.—Elongato-ovatum, convexum, nitidum, nigro-piceum, granulose tuberculatum, pronoto parvo, brevi, elytris longis, angustis, pronoto haud latioribus, ped. atque ant. robustis, rufo-piceis . . . **granulosum.**
- L. c. 0.65 mm.—Ovatum, valde convexum, nigrum, haud nitidum, pronoto brevi, confertissime tuberculato, basin versus transverse impresso, elytris ovatis, inflatè convexis, ped. atque ant. longis, piceo-nigris . . . **Crotchianum.**
- L. c. 0.65 mm.—Elongatum, valde convexum, piceo-brunneum, capite elongato triangulari, oculis prominentibus, pronoto parvo, cordiformi, elytris longis, apice parum dilatato et recte truncato, ped. atque ant. piceis . . . **Sharpianum.**

Actidium Fowlerianum.

A. Fowlerianum, Matthews, Cist. Ent. iii. p. 47a, 1883; Trans. Am. Ent. Soc. xi. p. 146, 1884; Biol. Cent.-Amer. Col. ii. pt. i. p. 148, 1888.

L. c. $\frac{3}{16}$ – $\frac{4}{16}$ lin. = 0.37–0.50 mm.—Elongatum, valde angustum sat convexum, fuscescens, pilis brevibus, sericeis, argenteis, dense vestitum; capite magno, antice obtuso, confertim tuberculato; oculis parvis, prominentibus; pronoto parvo, transverso, capite nec longiori neque latiori, ante medium latissimo, confertim tuberculato, nitidiusculo, lateribus marginatis, antice rotundatis, postice sat abrupte contractis, margine basali rotundatâ; elytris elongatis, capite atque pronoto vix latioribus, fere duplo longioribus, ante media latissimis, convexis, confertim asperatis, postice attenuatis, lateribus marginatis, levissime rotundatis, apice rotundato; abdomine haud exserto; pedibus modicis, læte flavis; antennis robustis, obscure flavis.

Body elongate, very narrow, rather convex, fuscescens, densely covered with short, sericeous, silvery hair.

Head large, obtuse in front, closely tuberculate; eyes small and prominent; antennæ robust, dusky yellow.

Thorax small, transverse, not longer or broader than the head, widest before the middle, closely tuberculate, rather shining, sides margined, rounded in front, and rather abruptly contracted posteriorly, basal margin rounded.

Scutellum small, triangular, acuminate.

Elytra elongate, attenuated posteriorly, scarcely broader but nearly twice longer than the head and thorax, widest before the middle, convex, closely asperated, sides margined and faintly rounded, apex rounded.

Abdomen entirely covered by the elytra.

Legs moderate, bright yellow.

Under parts castaneous, with the mouth and coxæ yellow.

Differs from others in its minute size, elongate and attenuated form, close sculpture, and dense, sericeous, silvery pubescence.

Habitat, Central America. Found in Guatemala, near the city, by Mr. Champion.

Actidium lineare.*A. lineare*, Matthews, Cist. Ent. i. p. 296, 1874.

L. e. $\frac{1}{16}$ lin. = 0.50 mm.—Lineare, elongatum, valde convexum, nigrum, pilis griseis dense vestitum; *capite* magno, elongato, antice acuminato, punctis profundis, fere confluentibus impresso: oculis modicis, prominentibus; *pronoto* parvo, capite nec longiori neque latiori, punctis magnis, fere confluentibus, atque pone medium, discum impressione levissima, longitudinali, utrinque notato, lateribus fere rectis, leviter marginatis, margine basali valde rotundata; *elytris* elongatis, linearibus, capite atque pronoto sesqui longioribus, haud latioribus, ordinibus transversis, irregularibus confertim et sat profunde asperatis, lateribus fere rectis, leviter marginatis, apice lato, valde rotundato; pedibus modicis, late flavis, tibiis pone medias leviter dilatatis, tarsis brevibus; antennis sat robustis, late flavis.

Body linear, elongate, very convex, black, thickly clothed with griseous hair.

Head large, elongate, acuminate in front, covered with deep, nearly confluent punctures; eyes moderate, prominent; antennæ rather robust, bright yellow.

Thorax small, not longer or broader than the head, marked with deep and almost confluent punctures, and a faint longitudinal impression on each side behind the middle, sides nearly straight and faintly margined, basal margin much rounded.

Scutellum small, triangular, deeply punctured.

Elytra elongate, linear, one half longer, but not broader than the head and thorax, closely and rather deeply asperated in irregular, transverse rows, sides nearly straight and faintly margined, apex broad and much rounded.

Abdomen not exposed.

Legs moderate, bright yellow, tibiæ slightly dilated beyond the middle, tarsi short.

Under parts piceous; mouth, coxæ and venter paler.

Differs from *A. coarctatum* in its linear form, much smaller size, elongate and acuminate head, smaller and much narrower thorax, and fainter but closer sculpture.

Habitat, Australasia. Found in New Zealand by Mr. Lawson.

Actidium politum.*A. politum*, Matthews, Cist. Ent. ii. p. 168, 1877.

L. e. $\frac{1}{16}$ lin. = 0.50 mm.—Ovale, convexum, nigrum, nitidissimum, pilis brevissimis sparsissime indutum; *capite* sat magno, antice longius producto, et valde rotundato, indistinctissime punctato atque foveâ indistinctâ in fronte utrinque impresso; oculis sat parvis, prominulis; *pronoto* brevi, sat parvo, capite haud longiori, parum latiori, ad basim latissimo, nitidissimo, punctis remotis, indistinctis impresso, lateribus leviter rotundatis et leviter marginatis, margine basali valde rotundata; *elytris* ovalibus, capite atque pronoto parum latioribus et longioribus, ante media latissimis, nitidissimis, ordinibus distinctis, sinuatis, levissime asperatis, atque striatim punctatis, apice lato, valde rotundato, dilutiori; pedibus sat longis, atque antennis piceo-nigris.

Body oval, convex, black and very shining, very thinly covered with very short hair.

Head rather large, much produced and much rounded in front, very indistinctly punctured, and marked with an indistinct fovea on each side of the forehead; eyes small and rather prominent; antennæ pitchy black.

Thorax short and rather small, not longer but rather broader than the head, widest at the base, very shining, marked with remote, indistinct punctures, sides slightly rounded, and slightly margined, basal margin much rounded.

Scutellum moderate, marked with large punctures.

Elytra oval, rather longer and broader than the head and thorax, widest before the middle, very shining, faintly asperated in distinct, wavy rows, and punctured in striae, apex broad, much rounded, and paler.

Legs rather long, pitchy black.

Under parts piceous, mouth and metasternum paler.

Differs from other species in its minute size, very shining surface, and in sculpture.

Habitat, North America. A single example of this very distinct species was found by Mr. G. R. Crotch near San Diego in California.

Actidium rotundicolle.

A. rotundicolle, Casey, Bull. Calif. Ac. Sci. ii. p. 248, 1886.

L. c. $\frac{4}{16}$ lin. = 0.50 mm.—Elongato-ovatum, convexum, nitidulum, leviter alutaceum, nigro-piceum, pilis argenteis parce vestitum; *capite* magno, prominenti, antice obtuso, indistincte alutaceo; oculis modicis, prominulis, retropositis; *pronoto* parvo, capite latiori, haud longiori, prope medium latissimo, leviter alutaceo, nitidulo, lateribus marginatis, antice valde rotundatis, postice fortissime contractis, margine basali leviter rotundata, angulis obsoletis; *elytris* capite atque pronoto longioribus et parum latioribus, ante media latissimis, confertim et levissime asperatis, interstitiis alutaceis, lateribus ovate-rotundatis et marginatis, apice obtuso; pedibus robustis, lute flavis; antennis gracilibus, obscure flavis.

Body elongate-ovate, convex, rather shining, finely alutaceous, pitchy black, sparingly clothed with silvery hair.

Head large and prominent, obtuse in front, indistinctly alutaceous; eyes moderate, rather prominent, placed far back; antennae slender, dusky yellow.

Thorax small, broader but not longer than the head, widest near the middle, faintly alutaceous, rather shining, sides margined, much rounded in front, and very strongly contracted towards the base, basal margin slightly rounded with the angles obsolete.

Scutellum large, triangular, and alutaceous.

Elytra longer and rather broader than the head and thorax, widest before the middle, closely and very faintly asperate with the interstices alutaceous, sides ovately rounded and margined, apex obtuse.

Legs robust, bright yellow.

Under parts nigro-piceous, mouth and palpi bright yellow.

Differs from all others in the peculiar shape of its thorax and faint sculpture.

Habitat, North America. Found in Texas by Lieut. T. L. Casey, U.S.E.

Actidium robustulum.

A. robustulum, Casey, Bull. Calif. Ac. Sci. i. p. 324, 1885.

L. c. $\frac{4}{16}$ lin. = 0.50 mm.—Elongato-ovatum, modice convexum, haud nitidum, per totum alutaceum, nigro-piceum; *capite* magno, antice producto, fortiter alutaceo; oculis parvis, valde prominentibus; *pronoto* sat magno, capite latiori, haud longiori, pone medium latissimo, fortiter alutaceo, depressione levi ante basim impresso, lateribus marginatis atque una cum basi fere continuo rotundatis; *elytris* ovatis, capite atque pronoto parum longioribus, haud latioribus, prope media latissimis, fortiter alutaceis, stria snturali distincta, apice acuto; pedibus modicis, nigro-piceis; antennis sat gracilibus, nigro-piceis.

Body elongate-ovate, moderately convex, not shining, alutaceous throughout, nigro-piceous.

Head large, produced in front, strongly alutaceous; eyes small and very prominent; antennæ rather slender, nigro-piceous.

Thorax rather large, broader but not longer than the head, widest behind the middle, strongly alutaceous, with a faint depression near the base, sides margined, and almost continuously rounded with the base.

Scutellum elongate-triangular, alutaceous.

Elytra ovate, rather longer but not broader than the head and thorax, widest near the middle, strongly alutaceous, sutural stria distinct, apex acute.

Legs moderate, nigro-piceous.

Under parts pitchy black, metasternum deeply punctured.

Differs from *A. granulosum* in its broader form, larger thorax, more slender antennæ, and finer sculpture.

Habitat, North America. Found in California by Lieut. T. L. Casey, U.S.E.

Actidium granulosum.

A. granulosum, Casey, Bull. Calif. Ac. Sci. i. p. 325, 1885.

L. c. $\frac{1}{16}$ lin. = 0.50 mm.—Elongato-ovatum, convexum, nitidum, piceo-nigrum, pilis griseis sparse vestitum, totum confertissime tuberculatum sive alutaceum; capite permagno, granuloso; oculis parvis, prominentibus; pronoto parvo, brevi, capite breviori et haud latiori, antice latissimo, granuloso, pone medium discum utrinque depressione levi impresso, lateribus marginatis atque una cum basi continuo rotundatis; elytris longis, angustis, capite atque pronoto plus quam sesqui longioribus, haud latioribus, prope media latissimis, granulosi, lateribus marginatis, apice integro, attenuato; pedibus atque antennis sat magnis, rufo-piceis.

Body elongate-ovate, convex, shining, pitchy black, sparingly clothed with grey hair, closely tuberculate or coarsely alutaceous throughout.

Head very large; eyes small and prominent; antennæ rather large, rufo-piceous.

Thorax small and short, shorter and not broader than the head, widest in front, marked with a faint depression on each side behind the middle of the disc, sides margined and continuously rounded with the base.

Scutellum moderate, triangular and alutaceous.

Elytra long and narrow, more than one half longer, but not broader than the head and thorax, widest near the middle, sides margined, apex entire and attenuated.

Legs rather large, rufo-piceous.

Under parts rufo-piceous, metasternum deeply punctured.

Differs from others in its elongate-ovate form, small size, granulose sculpture, and rufo-piceous legs and antennæ.

Habitat, North America. Found in California by Lieut. T. L. Casey, U.S.E.

Actidium Crotchianum.

A. Crotchianum, Matthews, Cist. Ent. ii. p. 168, 1877.

L. c. $\frac{5}{16}$ lin. = 0.63 mm.—Ovatum, valde convexum, nigrum, haud nitidum, pilis brevissimis albidis vestitum; capite sat magno, antice producto, confertim tuberculato; oculis modicis, prominulis; pronoto brevi, transverso, capite vix longiori, parum latiori, ante medium latissimo, confertissime tuberculato, depressione transversa basim versus notato, lateribus antice rotundatis, postice contractis, margine basali fere recta; elytris ovatis, valde convexis, leviter inflatis, capite atque pronoto plus quam sesqui longioribus, parum latioribus, ad media latissimis, ordinibus transversis leviter, sed confertissime asperatis, lateribus modice rotundatis, leviter marginatis, apice attenuato, aliquantum dilutiori; pedibus atque antennis sat longis, piceo-nigris.

Body ovate, very convex, black, not shining, thinly clothed with very short whitish hair.

Head rather large, produced in front, closely tuberculate; eyes moderate, rather prominent; antennæ rather long, pitchy black, with the ninth joint scarcely incrassate.

Thorax short, transverse, scarcely longer but rather broader than the head, widest before the middle, very closely tuberculate, with a transverse depression near the base, sides rounded in front and contracted towards the base, basal margin nearly straight.

Scutellum moderate, closely asperated.

Elytra ovate, very convex, slightly inflated, more than one half longer and rather broader than the head and thorax, widest at the middle, lightly but very closely asperated in transverse rows, sides moderately rounded and slightly margined, apex attenuated, and rather paler.

Abdomen not exposed.

Legs rather long and slender, pitchy black.

Under parts entirely piceous.

Differs from *A. Boudierii* in its larger size, broader and more inflated form, fainter sculpture, and more robust antennæ.

Habitat, North America. Found in British Columbia by Mr. G. R. Crotch.

Actidium Sharpianum.

A. Sharpianum, Matthews, Cist. Ent. iii. p. 39, pl. ii. fig. 2, 1882.

L. c. $\frac{5}{16}$ lin. = 0.63 mm.—Elongatum, valde convexum, pieco-brunneum, pilis brevibus argenteis vestitum; capite modico, elongato, triangulari, confertim tuberculato; oculis magnis, prominentibus, longe retropositis; pronoto parvo, cordiformi, capite pariter longo, parum latiori, ante medium latissimo, postice attenuato, confertim tuberculato, impressione transversâ, ovali, basim versus leviter notato, lateribus leviter marginatis, margine basali rotundatâ; elytris sat longis, capite atque pronoto parum latioribus, fere duplo longioribus, prope media latissimis, convexis, confertim, sat leviter, asperatis, lateribus leviter marginatis, et leviter rotundatis, apice recte truncato et parum dilatato; abdominis pygidio exserto; pedibus piceis, robustis, tibiis omnibus prope medias dilatatis; antennis longis, piceis.

Body elongate, very convex, pitchy brown, clothed with short silvery hair.

Head elongate, triangular, closely tuberculate; eyes large, prominent, and placed far back; antennæ long and piceous.

Thorax small, cordiform, not longer but rather broader than the head, widest before the middle, attenuated posteriorly, closely tuberculate, with a faint, oval, transverse impression towards the base, sides slightly margined, basal margin rounded.

Scutellum elongate, triangular, closely asperated.

Elytra rather long, rather broader and nearly twice longer than the head and thorax, widest near the middle, convex, closely but not deeply asperated, sides slightly margined and slightly rounded, apex straightly truncate, and laterally dilated.

Abdomen with the pygidium exserted.

Legs robust, piceous, with all the tibiæ dilated near the middle, and the posterior coxæ laminated.

Under parts castaneous.

Differs from all other species in its elongate, convex form, small thorax, and especially in the abruptly truncate apex of its elytra.

Habitat, Polynesia. One specimen only, found by Mr. Blackburn, running on mud on the sea-shore near Honolulu in Hawaii.

Actidium attenuatum (mihi invisum).

A. attenuatum, Casey, Bull. Calif. Ac. Sci. i. p. 326, 1885.

L. c. 0·45 mm.—Angustissimum, subcylindricum, nitidulum, leviter alutaceum, aterrimum, pilis brevibus griseis modice vestitum; *capite* modico, antice sat producto; oculis modicis, prominentibus; *pronoto* capite parum latiori, vix longiori, convexo, lateribus leviter rotundatis, margine basali rotundata, angulis rotundatis; scutello triangulari; *elytris* capite atque pronoto laud latioribus, multum longioribus, valde convexis, modice asperatis, lateribus leviter rotundatis, apice sat obtuso; pedibus sat robustis, piceo-castaneis; antennis modicis, nigris.

The above is an analysis of Mr. Casey's elaborate description in his 'Essay on New Genera and Species of Californian Coleoptera.'

A. attenuatum seems to be a distinct species, allied to *A. granulorum* and *A. rotundicollis*, from both of which it apparently differs in its more narrow and elongate form, more dense pubescence, and finer sculpture.

It is at present represented by a single individual found by Mr. Casey in California.

GEN. XX.—PTENIDIUM.

Fifteen fresh species have been added to this genus. Of these, *P. Lawsoni* and *P. Ulkei* belong to the variolose group of which *P. punctatum* is the type; twelve are more or less allied to *P. evanescens*; and one, *P. stragulatum*, is of such an exceedingly abnormal character that, if I did not strongly object to subdividing on insufficient grounds, I should feel inclined to form a separate genus for its reception.

SPECIERUM RECENTIORUM EPITOMA.

L. c. 0·80 mm.—Elongatum, nitidissimum, nigrum, elytrorum humeris atque apicibus rufis, pronoto ad basin strangulatim contracto, ped. longis atque ant. flavis	stragulatum.
L. c. 0·90 mm.—Ovale, valde convexum, castaneum, nitidum, punctis foveolatis notatum, ped. atque ant. flavis	Lawsoni.
L. c. 0·75 mm.—Suboblongum, modice convexum, castaneum, nitidissimum, totum punctis foveolatis notatum, ped. atque ant. late flavis	Ulkei.
L. c. 0·75 mm.—Ovatum, piceo-nigrum, valde convexum, pronoto minute punctato, foveis basalibus profundis, scutello foveis magnis impresso; ped. atque ant. late flavis	foveatum.
L. c. 0·63 mm.—Ovale, piceo-nigrum, glabrum, nitidissimum, pronoti foveis basalibus magnis, profundis; ped. atque ant. late flavis	ignobile.
L. c. 0·80 mm.—Suboblongum, piceum, valde convexum, nitidissimum, totum impunctatum, ped. atque ant. late flavis	impunctatum.
L. c. 1 mm.—Elongato-ovale, aterrimum, glabrum, nitidissimum, pronoti foveis basalibus nullis, ped. atque ant. late flavis	nitens.
L. c. 0·75 mm.—Angustum, elongatum, modice convexum, nigrum; pronoto punctis magnis in disco notato, foveis basalibus minutis, ped. late, ant. obscure flavis	Caseianum.
L. c. 0·75 mm.—Elongatum, valde angustatum, nitidissimum, obscure castaneum, foveis basalibus pronoti permagnis; ped. atque ant. longis, late flavis	attenuatum.
L. c. 0·75 mm.—Anguste ovatum, convexum, alutaceum, nitidum, nigrum, pronoto fere impunctato; ped. atque ant. late flavis	Africanum.
L. c. 0·87 mm.—Breve, obesum, validissime convexum, glabrum, nitidissimum, pronoto glabro, punctis modicis quatuor basalibus impresso; ant. brevioribus, atque ped. obscure flavis	obesum.

- L. c. 1 mm.—Ovale, validissime convexum, nitidissimum, castaneum, elytris rufescentibus, pronoto minute punctato, et foveis duabus basalibus impresso, ant. longis, atque ped. laete flavis. **speculifer.**
- L. c. 0·87 mm.—Breve, valde convexum, nigrum, modice punctatum, pronoti foveis basalibus nullis; ped. brevibus, laete flavis; ant. obscure flavis **orientale.**
- L. c. 0·75 mm.—Elongato-ovatum, convexum, nigrum, leviter alutaceum, pronoti foveis basalibus quatuor profundis; ped. longis, laete flavis; ant. longis, gracillimis, obscure flavis **Reitteri.**
- L. c. 0·60 mm.—Ovatum, postice attenuatum, nigrum, nitidissimum, glabrum, capite atque pronoto magnis, elytris parvis, ant. longis, robustis, atque ped. ketissime flavis **concinnum.**

Ptenidium strangulatum.

P. strangulatum, Matthews, Trans. Am. Ent. Soc. xi. p. 148, 1886; Biol. Cent.-Amer. Col. ii. pt. i. p. 149, 1888.

L. c. $\frac{6}{16}$ — $\frac{7}{16}$ lin. = 0·75–0·87 mm.—Elongatum, ad humeros strangulatum, valde convexum, glabrum, nitidissimum, nigrum; capite magno, antice prominenti, punctis minutis, remote notato; oculis magnis, prominentibus; pronoto sat parvo, transverso, ad basim fortissime contracto, capite parum latiori, hand longiori, ante medium latissimo, glabro, nitidissimo, foveis quatuor valde indistinctis, equidistantibus, ad basim, atque duabus vel tribus, pariter indistinctis, ad angulum utrumque basalem impresso, lateribus late marginatis, et valde rotundatis, margine basali perbrevis, depressa et fere recta; elytris ovatis, capite atque pronoto multum latioribus, et fere duplo longioribus, prope media latissimis, glaberrimis, nitidissimis, humeris atque apicibus rufis, atque pariter et fortissime contractis, punctorum remotorum, striâ abbreviata, lineâ convergenti finita, ad suturam utrinque impressis, lateribus marginatis, et valde rotundatis, epipleuris præ modo latis, apice angusto, rotundato; pedibus sat longis, laete flavis; antennis modicis, flavis, clavis obscurioribus.

Body elongate, strangulated at the shoulders, very convex, smooth and very shining, black, with the shoulders and apex of the elytra red.

Head large and prominent, remotely and minutely punctured; eyes large and prominent; antennæ moderate, yellow, with the club dusky.

Thorax rather small, transverse, very strongly contracted towards the base, rather broader but not longer than the head, widest before the middle, smooth and very shining, with four indistinct, equidistant basal foveæ, and two or three indistinct foveæ near each posterior angle, sides much rounded and broadly margined, basal margin very short, depressed, and nearly straight.

Scutellum very long, triangular, sparingly punctured.

Elytra ovate, much broader and nearly twice longer than the head and thorax, widest near the middle, very smooth and shining, shoulders and apex red and equally contracted, with a row of remote punctures on each side of the posterior part of the suture terminating in a convergent impressed line, sides margined and much rounded, epipleura unusually broad, apex narrow and rounded.

Legs rather long, bright yellow, tarsi very small.

Under parts black, with the coxæ, apical segment of the venter entirely, and the posterior margin of the other segments flavescent.

Differs from all other species in the strangulated form of its body.

Habitat, Central America. Found in Guatemala by Mr. Champion.

I have carefully examined the anatomy of this remarkable species, and cannot find that it differs in any generic character from *Ptenidium*.

*Ptenidium Lawsoni.**P. Lawsoni*, Matthews, Cist. Ent. i. p. 123, 1873.

L. c. $\frac{6}{16}$ – $\frac{8}{16}$ lin. = 0·75–1 mm.—Ovale, convexum, castaneum, nitidum, pilis brevissimis flavis sparse vestitum, punctis foveolatis impressum; *capite* sat parvo, glabro, nitido, impunctato, pilis paucis longis remote instructo, clypei margine supra oculos atque antennis distincte reflexa; oculis magnis, prominentibus; *pronoto* modico, capite parum longiori atque latiori, pone medium latissimo, punctis foveolatis remotis per totum impresso, ad basim ordine transverso dispositis, interstitiis glabris, nitidis, lateribus marginatis, antice rotundatis, postice contractis, margine basali fere recta, angulis obtusis; *elytris* ovalibus, capite atque pronoto sat longioribus atque latioribus, ante media latissimis, punctis magnis, profundis confertim impressis, interstitiis glabris, nitidis, lateribus marginatis, apice lato, rotundato; pedibus sat longis, late flavis; antennis longis, gracilibus, flavis.

Body oval, convex, castaneous, shining, sparingly covered with very short yellow hair, marked with foveolate punctures.

Head rather small, smooth and shining, impunctate, furnished with some long distant hairs, with the margin of the clypeus above the eyes and antennae distinctly reflexed; eyes large and prominent; antennae long and slender, pale yellow.

Thorax moderate, rather longer and rather broader than the head, widest beyond the middle, marked throughout with remote foveolate punctures, arranged at the base in a transverse row, interstices smooth and shining, sides margined, rounded in front and contracted towards the base, basal margin nearly straight with the angles obtuse.

Scutellum large, triangular, acuminate, smooth and shining, with two foveolate punctures near the middle of the base.

Elytra oval, rather longer and rather broader than the head and thorax, widest before the middle, closely marked with large punctures, interstices smooth and shining, sides margined, apex broad and rounded, paler.

Legs rather long, bright yellow.

Under parts castaneous, with the mouth, prothorax, and terminal segments of the venter pale.

Differs from *P. punctatum* in its broader, more ovate and convex form, the punctured disc of the thorax, and in its bright castaneous colour; and from *P. Ulkei* in its larger size, more convex form, and closer sculpture.

Habitat, Australasia. Found near Auckland, in New Zealand, by Mr. Lawson.

*Ptenidium Ulkei.**P. Ulkei*, Matthews, Trans. Am. Ent. Soc. xi. p. 148, 1884.

L. c. $\frac{6}{16}$ lin. = 0·75 mm.—Oblongum, sat convexum, nitidissimum, punctis foveolatis per totum impressum, castaneum, pilis validis flavis sparse vestitum; *capite* modico, sat elongato, punctis profundis remote impresso; oculis magnis, prominulis; *pronoto* modico, capite parum latiori, haud longiori, pone medium latissimo, foveis profundis remotis per totum impresso, interstitiis nitidis, lateribus postice contractis, leviter marginatis, margine basali fere recta, angulis sat obtusis; *elytris* ovalibus, capite atque pronoto sat longioribus atque latioribus, ante media latissimis, convexis, punctis foveolatis, profundis, remote impressis, interstitiis nitidis, lateribus leviter marginatis, apice obtuso, dilutiori; pedibus atque antennis late flavis.

Body suboblong, moderately convex, very shining, marked throughout with foveolate punctures, castaneous, sparingly covered with stout yellow hair.

Head moderate, rather long, remotely marked with deep punctures, interstices shining; eyes large, rather prominent; antennæ moderate, bright yellow.

Thorax moderate, rather broader but not longer than the head, widest beyond the middle, remotely marked throughout with deep punctures, interstices shining, sides contracted towards the base, slightly margined, basal margin nearly straight with the angles rather obtuse.

Scutellum very large, triangular, remotely marked with deep punctures.

Elytra oval, convex, rather longer and broader than the head and thorax, widest before the middle, remotely marked with deep foveolate punctures, interstices shining, sides slightly margined, apex obtuse and paler.

Legs moderate, bright yellow with the femora dusky.

Under parts entirely castaneous.

Differs from *P. Lawsoni* in its smaller size, less convex form, and deeper and more remote sculpture.

Habitat, North America. Found in British Columbia by Mr. H. Ulke.

Ptenidium Reitteri.

P. Reitteri, Flach, Wien. Ent. Zeit. vi. 1887, p. 180; Verh. Zool.-Bot. Ges. Wien, xxxix. 1889, p. 498.

L. c. $\frac{6}{16}$ lin. = 0.75 mm.—Elongato-ovatum, nigrum, nitidum, convexum, leviter alutaceum, pronoti foveis basalibus quatuor; *capite* modico, antice producto, alutaceo, punctis foveolatis tribus, prope oculum, utrumque impresso; oculis modicis; *pronoto* modico, capite sat longiori et latiori, prope basim latissimo, alutaceo, punctis remotis in disco atque foveis quatuor profundis, mediis approximatis, ad basim impresso, lateribus marginatis, margine basali rectâ, angulis obtusis; *elytris* capite atque pronoto longioribus, hand latioribus, prope humeros latissimis, alutaceis, leviter et remote striatim punctatis, lateribus marginatis, apice obtuso, dilutiori; pedibus longis, læte flavis; antennis longis et gracilibus, obscure flavis.

Body ovate-elongate, black, convex, very shining, faintly alutaceous, with four thoracic foveæ.

Head moderate, produced in front, alutaceous, with three foveolate punctures on each side near the eye; eyes moderate, rather prominent; antennæ long and slender, dusky yellow.

Thorax moderate, rather longer and broader than the head, widest near the base, alutaceous, marked with a few remote punctures on the disc, and four deep foveæ, of which the interior are near together, at the base, sides margined, basal margin straight with the angles obtuse.

Scutellum large, triangular, with four large punctures at the base.

Elytra longer but not broader than the head and thorax, widest near the shoulders, alutaceous, faintly and remotely punctured in striæ, sides margined, apex obtuse, paler.

Legs long, bright yellow.

Under parts black, with the coxæ bright yellow.

Differs from *P. levigatum* in its smaller size, much narrower shape, alutaceous surface, smaller thoracic foveæ, and long slender antennæ.

Habitat, Europe. Found in Dalmatia.

Ptenidium foveatum.

P. foveatum, Matthews, Trans. Am. Ent. Soc. xi. p. 149, 1884; Biol. Cent.-Amer. Col. ii. pt. i. p. 150, 1888.

L. c. $\frac{5}{16}$ – $\frac{6}{16}$ lin. = 0.63–0.75 mm.—Obtuse ovatum, piceo-nigrum, nitidissimum, valde convexum, foveis basalibus pronoti profundis; *capite* magno, lato, punctis

tribus magnis utrinque notato; oculis magnis, sat prominentibus; *pronoto* modico, capite parum longiori et latiori, glabro, nitidissimo, punctis minutis in disco, foveis quatuor magnis, equidistantibus ad basim, atque duabus minoribus ad marginem anteriorem impresso, lateribus prope media latissimis, rotundatis et marginatis, margine basali recta, angulis obtusis; *scutello* magno, foveis quatuor magnis ad basim, duabus minoribus ad medium, atque singula ad apicem notato; *elytris* sat brevibus, capite atque pronoto parum longioribus atque latioribus, ante media latissimis, perglabris, nitidissimis, striâ suturali punctorum remotorum utrinque notatis; pedibus atque antennis læte flavis.

Body obtusely ovate, nigro-piceous, very shining and very convex, with the basal foveæ of the thorax deep.

Head large and broad, with three large punctures on each side near the eye; eyes large and rather prominent; antennæ moderate, bright yellow.

Thorax moderate, rather longer and rather broader than the head, widest near the middle, smooth and very shining, marked with minute punctures on the disc, and with four large equidistant foveæ near the base, and two smaller foveæ near the anterior margin, sides rounded and margined, basal margin straight with the angles obtuse.

Scutellum large, triangular, with four large foveæ at the base, two smaller on the middle, and one near the apex.

Elytra rather short, rather longer and rather broader than the head and thorax, widest before the middle, very smooth and very shining, with a sutural stria of remote punctures on each side, sides rounded and margined, apex slightly attenuated.

Legs rather long, very bright yellow, with the tibiæ slightly dilated towards the apex.

Under parts black, with the mouth and coxæ bright yellow, and the venter piceous.

Differs from other species in its short ovate form, and in the peculiar sculpture of the scutellum.

Habitat, Central America. Found near Aceytuno, in Guatemala, by Mr. Champion; and in Cordova, in Mexico, by M. Salle.

Ptenidium concinnum.

P. concinnum, Matthews, Ann. Mag. Nat. Hist. 6, xiii. 1894, p. 342.

L. c. $\frac{5}{16}$ lin. = 0.63 mm.—Subovatum, postice valde attenuatum, convexum, nitidissimum, pilorum expers, fere impunctatum, nigrum, pronoto puncturâ vix conspicuâ utrinque ad basim impresso; capite magno, antice rotundato; oculis magnis, prominentibus; antennis longis, robustis, latissime flavis; *pronoto* magno, capite latiori, haud longiori, ante medium latissimo, basim versus contracto, glabro, nitidissimo, punctura vix conspicua ad basim impresso, lateribus antice valde rotundatis, postice contractis, fortiter marginatis, margine basali recta, angulis sat acutis; *elytris* parvis, ovatis, capite atque pronoto multo angustioribus, vix longioribus, ante media latissimis, postice multum attenuatis, punctulis minutis, remotissimis impressis, apicibus obtuse rotundatis; abdomine haud exserto; pedibus robustis, læte flavis.

Body subovate, much attenuated posteriorly, convex, very shining, without hair, almost impunctate, black, with an indistinct puncture on each side of the base of the thorax.

Head large, rounded in front; eyes large and prominent; antennæ long, robust, very bright yellow.

Thorax large, broader but not longer than the head, widest before the middle, contracted towards the base, smooth and very shining, with a very indistinct puncture on each side of the base, sides much rounded in front, contracted

towards the base, strongly margined, basal margin straight with the angles rather acute.

Scutellum large, triangular, smooth and impunctate.

Elytra small, ovate, attenuated in front and behind, much narrower and scarcely longer than the head and thorax, widest before the middle, marked with minute and very remote punctures, sides margined, extremities obtusely rounded.

Abdomen not exposed.

Legs robust, bright yellow.

Under parts black, very shining, metasternum indistinctly punctured, mouth and coxæ bright yellow.

Differs from others in its small size, large head and thorax, small elytra, almost impunctate surface, shining black colour, large and very bright yellow antennæ and legs.

Habitat, America. Found rarely by Mr. Smith in the West Indian Islands of Grenada and St. Vincent.

Ptenidium ignobile.

P. ignobile, Matthews, Trans. Am. Ent. Soc. xi. p. 149, 1884; Biol. Cent.-Amer. Col. ii. pt. i. p. 151, 1888.

L. c. $\frac{5}{16}$ lin. = 0.63 mm.—Ovale, sat elongatum, piceo-nigrum, glabrum, nitidissimum, pronoti foveis basalibus magnis et profundis; *capite* magno, antice sat producto, impunctato; oculis magnis, prominentibus; *pronoto* modico, capite parum latiori, vix longiori, prope medium latissimo, glabro, nitidissimo, foveis quatuor magnis et profundis, equidistantibus, prope basim impresso, lateribus modice rotundatis et marginatis, margine basali recta, angulis obtusis; *elytris* ovalibus, capite atque pronoto longioribus, haud latioribus, ante media latissimis, impunctatis, nitidis, postice sat attenuatis, lateribus ovaliter rotundatis, leviter marginatis, apice dilutiori; pedibus atque antennis sat longis, lute flavis.

Body oval, rather elongate, nigro-piceous, smooth and very shining, with the thoracic foveæ large and very deep.

Head large, rather produced in front, impunctate; eyes large and prominent; antennæ rather long, bright yellow.

Thorax moderate, subquadrate, rather broader but not longer than the head, widest near the middle, smooth and very shining, with four large and very deep equidistant foveæ near the base, sides slightly rounded and slightly margined, basal margin straight with the angles obtuse.

Scutellum large, triangular, with four small foveæ at the base.

Elytra oval, rather narrow, longer but not broader than the head and thorax, widest before the middle, impunctate and shining, rather attenuated posteriorly, sides ovally rounded and slightly margined, apex paler.

Legs rather long, bright yellow.

Under parts piceous, with the mouth, coxæ, and apex of the venter yellow.

Differs from other species in its small size, narrow form, large and deep thoracic foveæ, and smooth impunctate surface.

Habitat, Central America Found in Guatemala, near the city, by Mr. Champion.

Ptenidium impunctatum.

P. impunctatum, Matthews, Trans. Am. Ent. Soc. xi. p. 150, 1884; Biol. Cent.-Amer. Col. ii. pt. i. p. 152, 1888.

L. c. $\frac{6}{16}$ – $\frac{7}{16}$ lin. = 0.75–0.87 mm.—Elongato-oblongum, piceum, valde convexum, nitidissimum, omnino impunctatum; *capite* magno, lato; oculis sat magnis

et sat prominentibus: *pronoto* sat magno, transverso, capite latiori, haud longiori, prope medium latissimo, impunctato, glabro, nitidissimo, lateribus rotundatis et marginatis, margine basali recta, angulis obtusis; *elytris* obtuse-ovatis, capite atque pronoto angustioribus, vix longioribus, prope media latissimis, lateribus marginatis, apice obtuso, late rufescenti; pedibus atque antennis modicis, late flavis.

Body elongate-oblong, piceous, very convex and very shining, impunctate.

Head large and broad; eyes rather large and rather prominent; antennæ moderate, bright yellow.

Thorax rather large, transverse, broader but not longer than the head, widest near the middle, impunctate, smooth and very shining, sides rounded and margined, basal margin straight with the angles obtuse.

Scutellum moderate, triangular.

Elytra obtusely ovate, narrower and scarcely longer than the head and thorax, widest near the middle, convex, very shining and impunctate, sides margined, apex obtuse and broadly rufescent.

Legs moderate, bright yellow.

Under parts piceous, with the head, mesosternum, coxæ, and venter flavescens.

Differs from others in its broad head and thorax, narrow elytra, and impunctate surface.

Habitat, Central America. Found in several parts of Guatemala by Mr. Champion.

Ptenidium nitens.

P. nitens, Matthews, Trans. Am. Ent. Soc. xi. p. 149, 1884; Biol. Cent.-Amer. Col. ii. pt. i. p. 151, 1888.

L. c. $\frac{5}{16}$ lin. = 1 mm.—Elongato-ovale, valde convexum, aterrimum, glabrum, nitidissimum, pronoti punctis basalibus nullis; *capite* magno, prominenti, impunctato; oculis magnis, prominulis; *pronoto* modico, capite latiori et parum longiori, ad basim latissimo, valde convexo, nitidissimo, punctis exiguis, remotis in disco notato, foveis basalibus nullis, lateribus modice rotundatis, vix marginatis, margine basali leviter rotundata, angulis obtusis; *elytris* ovatis, capite atque pronoto longioribus, haud latioribus, ante media latissimis, valde convexis, nitidissimis, fere glabris, punctis remotis indistinctissime impressis, lateribus leviter marginatis, apice obtuso, fusciscenti; pedibus atque antennis modicis, late flavis.

Body elongate-oval, very convex, deep black, smooth and very shining, without thoracic foveæ.

Head large and prominent, impunctato; eyes large and rather prominent; antennæ moderate, bright yellow.

Thorax moderate, broader and rather longer than the head, widest at the base, very convex and very shining, indistinctly marked with minute, remote punctures on the disc, without foveæ at the base, sides moderately rounded, scarcely margined, basal margin faintly rounded, with the angles obtuse.

Scutellum large, triangular, smooth and shining.

Elytra ovate, longer but not broader than the head and thorax, widest before the middle, very convex and very shining, almost smooth, very indistinctly marked with minute remote punctures, sides faintly margined, apex obtuse, fusciscent.

Legs moderate, bright yellow.

Under parts black, with the mouth and coxæ yellow.

Differs from *P. Bruckii* in the absence of the basal foveæ of the thorax; and from other species of this genus in its shining glabrous surface.

Habitat, Central America. Found in Chontales, Nicaragua, by Mr. E. M. Janson.

Ptenidium Caseianum.

P. Caseianum, Matthews, Ann. Mag. Nat. Hist. 6, iii. 1889, p. 194.

L. c. $\frac{6}{16}$ lin. = 0.75 mm.—Angustum, elongatum, modice convexum, sat profunde punctatum, nitidissimum, pilis griseis vestitum, nigrum, apice elytrorum rufo-piceo; *capite* magno, punctis sat magnis remote impresso, antice elongato, nigro, labro rufo-piceo; oculis modicis; *pronoto* sat parvo, capite latiori, haud longiori, ante basim latissimo, punctis sat magnis remote impresso, interstitiis glabris, nitidis, foveis quatuor basalibus minutis, inconspicuis, lateribus rotundatis, vix marginatis, margine basali fere rectâ, angulis rotundatis; *elytris* longis, angustis, capite atque pronoto parum angustioribus, plus quam sesqui longioribus, atque profundius et magis confertim punctatis, ante media latissimis, interstitiis leviter reticulatis, lateribus marginatis et leviter rotundatis, apice angusto, rufo-piceo; pedibus modicis, late flavis; antennis atque palpis obscure flavis.

Body long and narrow, moderately convex, rather deeply punctured, very shining, clothed with grey hairs, black, with the apex of the elytra rufo-piceous.

Head large, remotely marked with rather large punctures, labrum rufo-piceous; eyes moderate; antennæ and palpi dusky yellow.

Thorax rather small, broader, but not longer than the head, widest before the base, remotely marked with large punctures, the interstices smooth and shining, the four basal foveæ minute and inconspicuous, sides rounded and faintly margined, basal margin nearly straight, with the angles rounded.

Scutellum moderate, triangular.

Elytra long and narrow, rather narrower and more than one half longer than the head and thorax, and more deeply and more closely punctured, widest before the middle, the interstices faintly reticulated, sides margined and slightly rounded, apex narrow, rufo-piceous.

Legs moderate, bright yellow.

Under parts black.

Differs from other species in its small size, long and narrow form, and especially in sculpture.

Habitat, North America. Found in California by Mr. Casey.

Ptenidium attenuatum.

P. attenuatum, Matthews, Ann. Mag. Nat. Hist. 6, iii. 1889, p. 194.

L. c. $\frac{6}{16}$ lin. = 0.75 mm.—Elongatum, valde angustatum, convexum, nitidissimum, pilis brevibus, flavescens, sparse vestitum, obscure castaneum, foveis basalibus pronoti permagnis; *capite* magno, antice producto, indistincte punctato; oculis magnis, sat prominentibus; *pronoto* lato, capite longiori atque latiori, ante medium latissimo, indistincte punctato, foveis basalibus permagnis et profundis, lateribus leviter marginatis, margine basali fere rectâ, angulis sat obtusis; *elytris* valde angustatis, capite atque pronoto plus quam sesqui longioribus, haud latioribus, prope humeros latissimis, striis valde remotis minute punctatis, interstitiis alutaceis, castaneis, dimidio apicali flavescens, apice attenuato, lateribus leviter marginatis; pedibus atque antennis longis, flavis.

Body elongate, very narrow, convex and very shining, castaneous, sparingly clothed with yellow hair, thoracic foveæ very large and deep.

Head large, produced in front, indistinctly punctured: eyes large and rather prominent; antennæ long and slender, yellow.

Thorax broad, longer and broader than the head, widest before the middle, indistinctly punctured, basal foveæ very large and deep, sides faintly margined, basal margin nearly straight, with the angles rather obtuse.

Scutellum very large, triangular, acuminate at the apex.

Elytra much narrowed, more than one half longer but not broader than the head and thorax, widest near the shoulders, minutely punctured in very remote striae with the interstices alutaceous, castaneous with nearly the apical half flavescent, sides slightly margined, attenuated towards the apex.

Legs long, yellow.

Under parts castaneous, coxæ and apex of the venter yellow.

Differs from other species in its intermediate size, long, narrow form, very large thoracic foveæ, and peculiar sculpture.

Habitat, Europe. Found in Italy.

Ptenidium Africanum.

P. Africanum, Matthews, Ann. Mag. Nat. Hist. 6, iii. 1889, p. 194.

L. c. $\frac{6}{16}$ lin. = 0.75 mm.—Ovatum, convexum, nitidum, pilis brevibus flavis sparse vestitum, alutaceum, modice punctatum, nigrum; *capite* magno, alutaceo, fere impunctato; oculis modicis, laud prominentibus; *pronoto* magno, fere ad basin latissimo, alutaceo, fere impunctato, lateribus marginatis, minime rotundatis, margine basali fere rectâ, angulis obtusis; *elytris* capite atque pronoto sesqui longioribus, laud latioribus, ante media latissimis, alutaceis, modice striatim punctatis, depressione levi ad mediam suturam utrinque impressis, lateribus marginatis, modice rotundatis, apice obtuso, parum dilutiori; pedibus atque antennis læte flavis.

Body ovate, convex, shining, sparingly clothed with yellow hairs, alutaceous, moderately punctured, black.

Head large, alutaceous, almost impunctate; eyes moderate, not prominent; antennæ moderate, bright yellow.

Thorax large, widest almost at the base, alutaceous, almost impunctate, sides margined, very little rounded, basal margin nearly straight with the angles obtuse.

Scutellum large, triangular, acuminate at the apex, alutaceous, with a single small fovea near the middle of the base.

Elytra one half longer, but not broader than the head and thorax, widest before the middle, alutaceous, with moderate punctures disposed in striae, and a longitudinal depression on each side of the suture near the middle, sides margined, and moderately rounded, apex obtuse, rather paler.

Legs moderatæ, bright yellow.

Under parts black.

Differs from *P. evanescens* in its much smaller size, narrower form, thorax not contracted at the base, alutaceous surface and faint sculpture.

Habitat, Africa. Found on Mount Carrée, Algeria, by Mr. Lewis.

Ptenidium orientale.

P. orientale, Flach, Verh. Zool.-Bot. Ges. Wien, xxxix. p. 525, 1889.

L. c. $\frac{7}{16}$ lin. = 0.87 mm.—Breve, valde convexum, nitidissimum, modice punctatum, nigrum, pronoti foveis basalibus nullis; *capite* magno, antice producto, punctis duobus magnis in fronte impresso, oculis parvis, laud prominentibus; *pronoto* modico, capite sat longiori et latiori, ante basin latissimo, indistincte et remote punctato, interstitiis nitidissimis, lateribus marginatis, margine basali fere rectâ, angulis rotundatis; *elytris* sat parvis, capite atque pronoto longioribus et sat latioribus, ante media latissimis, modice et distincte punctatis, interstitiis nitidissimis, ad suturam utrinque longitudinaliter impressis, lateribus marginatis, apice sat fusciscenti; pedibus sat brevibus, læte flavis; antennis modicis, obscure flavis.

Body short, very convex and very shining, moderately punctured, black, without thoracic foveæ.

Head large, produced in front and marked with two large punctures; eyes small, not prominent; antennæ moderate, dusky yellow.

Thorax moderate, rather longer and broader than the head, widest before the base, indistinctly and remotely punctured with the interstices very shining, sides margined, basal margin nearly straight with the angles rounded.

Scutellum very large, triangular, impunctate and very shining.

Elytra rather small, longer and rather broader than the head and thorax, widest before the middle, moderately and distinctly punctured, interstices very shining, with a longitudinal impression on each side of the suture, sides margined, apex fuscous.

Legs rather short, bright yellow.

Under parts black, with the mouth and coxæ bright yellow.

Differs from *P. Africanum* in its much larger size, broader form, and smooth, not alutaceous surface; from *P. evanescens*, of which it may be a local form, in its smaller size, shorter form, and in the sculpture of the elytra.

Habitat, Asia. Found near Lenkoran by Herr Leder.

Ptenidium obesum.

P. obesum, Matthews, Trans. Am. Ent. Soc. xi. p. 151, 1884; Biol. Cent.-Amer. Col. ii. pt. i. p. 150, 1888

L. c. $\frac{7}{16}$ lin. = 0.87 mm.—Brevo, obesum, validissime convexum, glabrum, nitidissimum, nigro-castaneum; capite sat brevi, latissimo, glabro; oculis magnis, vix prominentibus; pronoto modico, transverso, capite parum longiori, et multo latiori, prope medium latissimo, validissime convexo, glaberrimo, nitidissimo, punctis quatuor parvis, equidistantibus, ad basim, atque duobus ad medium marginis anterioris notato, lateribus valde rotundatis et leviter marginatis, margine basali recta, angulis valde obtusis; elytris brevibus, ovatis, validissimo convexis, glaberrimis, nitidissimis, capite atque pronoto parum longioribus, vix latioribus, ante media latissimis, lateribus leviter marginatis, apice valde obtuso; pedibus atque antennis modicis, flavis, his obscurioribus.

Body short and tumid, exceedingly convex, smooth and very shining, dark castaneous.

Head short and very broad, smooth and very shining; eyes large, scarcely prominent; antennæ moderate, dusky yellow.

Thorax moderate, transverse, rather longer and much broader than the head, widest near the middle, exceedingly convex, very smooth and shining, with four small equidistant foveæ near the base, and two near the middle of the anterior margin, sides much rounded and slightly margined, basal margin straight with the angles very obtuse.

Scutellum short and broad, abruptly acuminate towards the apex.

Elytra short, ovate, exceedingly convex, very smooth and shining, rather longer, but scarcely broader than the head and thorax, widest before the middle, sides slightly margined, apex very obtuse.

Legs moderate, yellow.

Under parts black, with the mouth and coxæ piceous.

Differs from other species in its short and exceedingly convex form, shining and almost impunctate surface, and short antennæ; and from *P. speculifer* in its short dusky antennæ, impunctate surface, and especially in the disposition of the thoracic foveæ.

Habitat, Central America. Found near Ducnas in Guatemala by Mr. Champion.

Ptenidium speculifer.

P. speculifer, Matthews, Trans. Am. Ent. Soc. xi. p. 151, 1884.

L. c. $\frac{8}{16}$ lin. = 1 mm. Ovale, validissime convexum, glaberrimum, nitidissimum, pilorum expers, piceum, elytris rufescentibus; *capite* sat magno, lato, foveâ singula prope oculum utrumque impresso; oculis magnis, prominentibus; *pronoto* modico, capite sat longiori atque latiori, ad medium latissimo, punctis perexiguâ dense dispositis, et foveis duabus parvis prope marginem anteriorem atque etiam prope basim impresso, lateribus valde rotundatis et leviter marginatis, margine basali fere recta, angulis sat obtusis; *elytris* modicis, capite atque pronoto parum longioribus, vix latioribus, ante media latissimis, validissime convexis et nitidissimis, punctis perexiguâ dense impressis, lateribus rotundatis, leviter marginatis, apice obtuso, dilutiori; pedibus atque antennis sat gracilibus, læte flavis.

Body oval, exceedingly convex and very shining, without hair, piceous, with the elytra rufescent.

Head rather large and broad, with a small fovea on each side near the eye; eyes large and prominent; antennæ rather slender, bright yellow; palpi bright yellow.

Thorax moderate, rather longer and rather broader than the head, widest at the middle, closely marked with very minute punctures and with two small foveæ near the anterior margin and also near the basal margin, sides much rounded and slightly margined, apex obtuse and pale.

Scutellum large and broad, triangular, smooth and shining.

Elytra moderate, rather longer, but scarcely broader than the head and thorax, widest before the middle, exceedingly convex and very shining, closely marked with very minute punctures, sides rounded and slightly margined, apex obtuse and pale.

Abdomen with the apical segment pale.

Legs rather slender, bright yellow, with the anterior tibiæ dilated towards the apex.

Under parts rufo-piceous, mouth and coxæ yellow.

Differs from P. obesum in its larger size, longer antennæ, and peculiar arrangement of the thoracic foveæ.

Habitat, North America. Found by Mr. E. A. Schwarz near Cambridge, Massachusetts.

GEN. XXI.—**CAMPTODIUM**.—PL. XIV.

Camptodium, Motschulsky, Bull. Mose. xli. 2, p. 189, 1868; Matthews, Cist. Ent. iii. p. 82, pl. iv. 1884.

Corpus capite atque pronoto angustatis, elytris sat inflatis.

Caput deflexum, sat parvum, antice parum rotundatum; oculis magnis, valde prominentibus; antennis sub clypeo sat longe ante oculos insertis.

Antennæ (Pl. XIV., fig. 7) longæ, gracillimæ, 11-articulatæ: 1 et 2 permagnis, pariter longis; 2 ad apicem abrupte truncato; 3-8 elongato-ovalibus, tenuibus, sibi-ipsis paribus; 9 octavo longiori, modice incrassato; 10 nono multo majori, ad basim valde et ovaliter incrassato, antice in collo prælongo, tenuissimo producto; 11 magno, longo, ad basim valde incrassato, ad medium fortiter constricto, et antice multum attenuato; tribus hisce ultimis corona mediali setarum ornatis, atque clavam elongatam, submoniliformem formantibus.

Partes nullæ oris examinatæ.

Pronotum angustum, basim versus leviter constrictum.

Scutellum modicum, elongato-triangulare.

Elytra magna, integra, turgide inflata.

Alæ longæ, lanceolatæ, setis prælongis, nigris fimbriatæ, scapo basali longo exortæ, atque nervis solitis firmatæ.

Prothorace sat magno; *prosterno* (Pl. XIV., fig. a) magno; *episternis* minutis, triangularibus; *epimeris* magnis, coxarum receptacula postice cingentibus; receptaculis coxarum circularibus, fere contingentibus.

Pectus mesosterno (Pl. XIV., fig. d) sat magno, sat late carinato; *episternis* modicis, suboblongis, margine exteriori excavatâ; *epimeris* magnis, latis, ad coxas extensis; coxarum receptaculis modicis, circularibus, per carinam divisis.

metasterno (Pl. XIV., fig. g) permagno, corporis lateribus remoto, margine posteriori inter coxas rectâ; *episternis* magnis, longis, ad coxas extensis; *epimeris* magnis, longis, ad apices late inflexis, atque ultra metasternum productis.

Venter segmentis sex compositus, primo sat longo, apicali integro.

Pedes omnes longi, graciles; *tarsis* triarticulatis, 1 brevis; 2 primo multo longiori; 3 prælongo, gracili, ad medium superne leviter emarginato; unguibus longis gracilibus.

Coxæ anteriores rotundatæ, fere contingentes.

intermediæ rotundatæ, per carinam divisæ.

posteriores sat parvæ, prominentes, laminâ magnâ triangulari instructæ, latissime remotæ.

Body with the head and thorax narrow, elytra much inflated.

Head deflexed, moderate, rounded in front; eyes large and very prominent; antennæ inserted beneath the frontal plate at some distance in front of the eyes.

Antennæ (Pl. XIV., fig. 7) long and very slender, 11-jointed: 1 and 2 very large, equal in length, 2 abruptly truncate at the apex: 3-8 elongate-oval, very slender, equal to each other in size; 9 longer than the eighth, moderately incrassate; 10 much larger than the ninth, ovally and very strongly incrassated at the base, produced in front into a very long and very slender neck; 11 large and long, strongly constricted near the middle, and much attenuated from thence to the apex; the three apical joints furnished with a whorl of rather short erect setæ near the middle, and forming an elongate, submoniliform club.

Parts of the mouth not examined.

Thorax narrow, slightly constricted near the base.

Scutellum moderate, elongate-triangular.

Elytra large, entire, much inflated.

Wings long, lanceolate, with a long, slender basal scape, strengthened with the usual neurulation, and fringed on both sides with long black setæ.

Prothorax rather large, with the *prosternum* (Pl. XIV., fig. a) large; *episterna* minute, triangular; *epimera* large, inclosing the coxal cavities on the hinder side; coxal cavities circular, nearly contingent.

Pectus with the *mesosternum* (Pl. XIV., fig. d) rather large, broadly carinated; *episterna* moderate, suboblong, with the outer margin excavated; *epimera* large and broad, extending to the coxal cavities, with their outer margin excavated; coxal cavities moderate, circular, separated by the carina.

metasternum (Pl. XIV., fig. g) very large, not extending to the sides of the body, with the hinder margin between the coxæ straight; *episterna* large and long, extending to the coxæ; *epimera* large and long, broadly inflexed posteriorly, and produced beyond the metasternum.

Venter composed of six segments, the first rather long, the apical entire.

Legs long and slender; *tarsi* three-jointed, 1 short; 2 much longer than the first; 3 very long and slender, faintly emarginate on the upper side near the middle; claws long and slender.

Coxæ anterior rounded, almost contingent.

intermediate rounded, separated by the carina.

posterior rather small, prominent, furnished with a large triangular lamina, very widely separated.

This interesting genus was first described by Col. Motschulsky from examples found in India, and figured by him in the same work (Tab. viii. fig. 4). His description, like all his other descriptions, is very unsatisfactory, but, as far as it goes, agrees with the insect found in Japan by Mr. Lewis; his figure is much more expressive, and has convinced me that Mr. Lewis's insect is identical with *Camptodium adustipenne*.

Camptodium adustipenne.

C. adustipenne, Motschulsky, Bull. Mosc. xli. 2, p. 189, pl. 8, fig. 4, 1868; Matthews, Cist. Ent. iii. p. 83, pl. iv. 1884.

L. c. vix $\frac{1}{16}$ lin. = 0.70 mm.—Sat elongatum, capite atque pronoto angustatis, elytris turgide incrassatis, castaneum, nitidum, pilis flavescentibus sparse indutum; capite sat parvo, foveâ magnâ in fronte impresso; oculis magnis, valde prominentibus; pronoto angusto, capite parum longiori et latiori, ante medium latissimo, nitido, foveis quatuor magnis, profundis, et fere contingentibus, ad basim impresso, quarum exteriores duplices existunt, lateribus marginatis, et basim versus leviter constrictis, margine basali fere rectâ, angulis acutis, sat prominentibus; elytris ovatis, inflatis, capite atque pronoto multum latioribus et longioribus, prope media latissimis, punctis foveolatis remote impressis, lateribus valde rotundatis, apice obtuso; pedibus atque antennis praelongis, gracillimis, læte flavis.

Body elongate, with the head and thorax narrow, and the elytra inflated, castaneous, shining, sparingly covered with flavescent hairs.

Head rather small, with a large fovea on the forehead; eyes large and very prominent; antennæ very long and very slender, bright yellow.

Thorax narrow, rather longer and broader than the head, widest before the middle, shining, with four large, deep and almost contingent foveæ near the base, of which the exterior on each side is double, sides margined and slightly constricted near the base, basal margin nearly straight, with the angles acute and rather prominent.

Scutellum large, triangular, smooth and shining, with an impressed line within the lateral margins.

Elytra ovate, inflated, castaneous, becoming dusky towards the apex, much broader and longer than the head and thorax, widest near the middle, remotely marked with foveolate punctures, sides much rounded, apex obtuse.

Legs very long and slender, bright yellow.

Under parts castaneous, with the terminal segments of the venter bright yellow.

Habitat, Asia. Found in India by Col. Motschulsky; and in Japan by Mr. Lewis.

GEN. XXIV.—**NANOSELLA**.—Pl. XIII. A.

Nanosella, Motschulsky, Bull. Mosc. xli. 2, p. 187, pl. 8, fig. 3, 1868; Matthews, Trich. Illust. p. 68, 1872.

Corpus elongatum, lineare, convexum, elytris integris.

Caput magnum, prominens, haud deflexum, antice latum, obtusum, antennis sat longe ante oculos insertis; parti gulari permagnâ, valde convexâ; oculis parvis, retropositis.

Antennæ (Pl. XIII., A, fig. 7) breves, robustæ, 10- aut potius 11-articulatæ: 1 magno, elongato, subcylindrico; 2 primo plus quam duplo breviori, pariter crasso, ad apicem abrupte truncato; 3-8 secundo brevioribus et gracilioribus, sibiipsis fere pariter longis, gradatim sed leviter incrassatis; 9 brevi, dilatato, cyathiformi; 10 magno, subquadrato; 11 permagno, obtuse conico, duobus precedentibus parum latiori, multum longiori, apicem versus setis verticillatis ornato; tribus ultimis clavam quasi biarticulatam formantibus, nono atque

decimo fere in uno conjunctis, nihilominus facile distinguendis; articulis omnibus ad apices setis brevibus instructis.

Partes oris haud examinatae.

Pronotum sat longum, subquadratum, angulis posterioribus haud productis.

Scutellum sat parvum, triangulare.

Elytra valde elongata, linearia, apice integro.

Alae longae, angustae, lanceolatae, alioqui non examinatae.

Abdomen non examinatum, pygidio solo viso.

Prothorax (Pl. XIII., A, fig. 2, a) *prosterno* brevi; *episternis* sat inconspicuis; *epimeris* minutis, perbrevis; *receptaculis coxarum* ovalibus, contingentibus, postice semi-excisis.

Pectus mesosterno (Pl. XIII., A, fig. 2, d) modico, triangulariter carinato; *episternis* angustis, humeralibus; *epimeris* permagnis, ad coxarum receptacula late extensis; *receptaculis coxarum* obliquis, subovatis, per carinam anguste divis.

metasterno (Pl. XIII., A, fig. 2, g) quadrato, ad latera corporis extenso, postice inter coxas minute acuminato; *episternis* angulis anterioribus inflexis, reliquis una cum *epimeris* totis sub elytris occultis.

Venter segmentis sex compositus, apicali integro, caeteris longiori.

Pedes perbreves, *antici* trochanteribus minutis, trigonalibus; femoribus modicis, simplicibus; tibiis sat gracilibus; tarsis triarticulatis, 1 secundo duplo longiori; 3 valde elongato, gracillimo; unguibus parvis, leviter curvatis. *intermedii* anticis similes.

postici (Pl. XIII., A, fig. 8) trochanteribus claviformibus, tibiis levissime dilatatis, atque ad apices calcaribus minutis armatis, alioqui anticis similes.

Coxae anteriores ovales, contingentes.

intermediae ovatae, per carinam vix divisae.

posteriores magnae, prominentes, robustae, margine exteriori tota late laminata.

Body elongate, linear, convex, with the elytra entire.

Head large and prominent, not deflexed, broad and obtuse in front, gular portion large and very convex; eyes small and placed far back; antennae inserted considerably in front of the eyes.

Antennae (Pl. XIII., A, fig. 7) short and robust, 10, or more properly 11-jointed: 1 large, elongate, subcylindric; 2 more than twice shorter than the first, but equally broad, abruptly truncate at the apex; 3-8 shorter and more slender than the second, in length nearly equal to each other, but gradually and slightly incrassated: 9 cyathiform, very short and widely dilated; 10 large, subquadrate; 11 very large, obtusely conical, rather broader and much longer than the two preceding united, ornamented towards the apex with verticillated setae; the three last form an apparently biarticulate club, the ninth and tenth, although easily distinguished, being closely joined together; all the articulations are furnished with short setae at their apex.

Parts of the mouth not examined.

Pronotum rather long, subquadrate, hinder angles not produced.

Scutellum small, triangular.

Elytra entire, long and linear.

Wings apparently long and narrow, lanceolate; not thoroughly examined.

Abdomen not examined, the pygidium only visible.

Prothorax (Pl. XIII., A, fig. 2, a) with the *prosternum* short; *episterna* indistinct; *epimera* very small and short; *coxal cavities* oval, contingent, semi-excised.

Pectus with the *mesosternum* (Pl. XIII., A, fig. 2, d) moderate, triangularly carinated; *episterna* narrow, humeral; *epimera* very large, reaching broadly to the coxal cavities; *coxal cavities* oblique, subovate, very slightly separated by the carina.

metasternum (Pl. XIII., A, fig. 2, g) quadrate, extending to the sides of the body, the posterior margin produced into a minute point between the coxae; *episterna*

with the inner anterior angle inflected, their other parts, together with the whole *epimera*, concealed by the elytra.

Venter composed of six segments, the apical being the longest and entire.

Legs very short, *anterior* with the trochanters minute, trigonal; femora moderate, simple; tibiae rather slender; tarsi three-jointed, the first twice as long as the second, the third very long and very slender, with the claws small and faintly curved.

intermediate like the anterior.

posterior (Pl. XIII., A, fig. 8) with the trochanters moderate, claviform; tibiae faintly dilated and armed with minute spurs at the apex; in other respects like the anterior.

Coxæ anterior oval, contingent.

intermediate ovate, very slightly separated by the carina.

posterior large and prominent, robust and broadly laminated on the whole outer margin.

N. fungi, the only species contained in this genus, has been so often accurately described, and is so clearly represented in Pl. XIII. A of this work, that it is needless to say more than that it is bright yellow in colour and clothed with a short golden pubescence. Among the *Trichopterygidae* collected in Central America by Mr. Champion, I was very much pleased to find an example of this most minute of all the Coleoptera, more especially as it enabled me to complete my previously imperfect description by adding an account of its antennæ. In Dr. Le Conte's specimen these limbs had been broken off, but in Mr. Champion's the whole head had become detached from the thorax in the process of mounting the insect on card, and thus the antennæ could be examined without any trouble.

GEN. XXVII.—NOSSIDIUM.

Nossidium posthumum.

N. posthumum, Matthews, Cist. Ent. i. p. 298, 1874; Trans. Am. Ent. Soc. xi. p. 155, 1884.

L c. $\frac{1}{16}$ lin. = 0.75 mm.—Ovale, valde convexum, nitidum, testaceum, pilis brevibus flavis vestitum; *capite* sat breve, obtuse triangulare, punctis remotis impresso; oculis modicis, vix prominentibus; *pronoto* capite longiori et multo latiori, ad basim latissimo, glabro, nitido, remote et levissime punctato, lateribus late marginatis, vix rotundatis, margine basali haud depressa, leviter sinuata, angulis acutis; *elytris* semiovalibus, capite atque pronoto paulo longioribus, haud latioribus, ad humeros latissimis, ordinibus transversis, sinuatis, profunde punctato-asperatis, interstitiis glabris, nitidissimis, lateribus vix rotundatis, leviter marginatis, apice obtuso; pedibus atque antennis testaceis.

Body oval, very convex, shining, testaceous, clothed with short yellow hair.

Head rather short, obtusely triangular, remotely punctured; eyes moderate, scarcely prominent; antennæ with the two basal joints yellow (the rest were broken off).

Thorax longer and much broader than the head, widest at the base, smooth and shining, remotely and very faintly punctured, sides broadly margined and faintly rounded, basal margin not depressed, slightly sinuated, with the angles acute.

Scutellum small, triangular, marked with five punctures placed in the form of a triangle on the disc.

Elytra semioval, rather longer, but not broader than the head and thorax, widest at the shoulders, marked with deep asperated punctures in transverse sinuated rows, with the interstices smooth and very shining, sides slightly margined and scarcely rounded, apex obtuse.

Legs moderate, yellow.

Under parts entirely testaceous.

Differs from *N. pilosellum* in its much smaller size and narrower form, smooth head and thorax, and much more deeply sculptured elytra.

Habitat, North America. A single specimen found in the United States by Mr. G. R. Crotch; the precise locality not mentioned.

The single example of this very distinct species was sent to me by my friend, Mr. Crotch, shortly before his death, but did not arrive until that lamented event had taken place. His premature death proved to science a serious loss, and to his friends a perpetual grief; and to his cherished memory I now dedicate this second part of a work, to the former part of which he so largely contributed.

The following more precise localities have been omitted by the author:—

Ptinella pellucida (p. 22), San Geronimo and Zapote, Guatemala.

Trichopteryx laricollis (p. 47), Granada and St. Vincent, West Indies.

„ *rufescens* (p. 50), Balheu, San Gerónimo and Capetillo, Guatemala;
San Felix and Peña Blanca, Panama.

„ *similior* (p. 50), Rio Naranjo, El Tumbador and El Reposo, Guatemala.

„ *occidentalis* (p. 51), Granada and St. Vincent, West Indies.

„ *fratercula* (p. 53), near Gumley, Leicestershire.

„ *angustipennis* (p. 54), Capetillo and San Gerónimo, Guatemala.

„ *minor* (p. 55), El Tumbador and San Gerónimo, Guatemala.

„ *lætitia* (p. 56), near Spa, Belgium; Repton, near London, and near
Gloucester, England.

„ *seminitens* (p. 57), Snodland, Kent; Sherwood Forest; Scotland.

„ *depressa* (p. 63), Granada and St. Vincent, West Indies.

„ *brevis* (p. 64), Zapote and near the city, Guatemala.

„ *californica* (p. 66), Lake Tahoe, California.

„ *bidilatata* (p. 69), El Tumbador, Balheu, San Gerónimo and Senahue,
Guatemala.

„ *ricina* (p. 71), Quezaltenango, 7800 feet, Guatemala.

„ *nigrita* (p. 71), near the city, Guatemala.

„ *japonica* (p. 74), Kiushiu and South Yezo, Japan.

Ptilium hornianum (p. 80), El Reposo, Guatemala.

Ptenidium strangulatum (p. 96), Capetillo, Guatemala.

„ *impunctatum* (p. 100), Capetillo, San Gerónimo, and near the city,
Guatemala.

Camptodius adustipenne (p. 107), Nagasaki, Japan.

ERRATA.

Page 7, line 23. *for* Gilleb. *read* Guilleb.

„ 49, „ 26. *for* Leden *read* Leder.

INDEX.

Species unknown to the author, and synonyms, are printed in italics.

	PAGE		PAGE
<i>ACRATRICHIS</i>	53	<i>DIMORPHELLA</i>	30
<i>ACTIDIUM</i>	89	<i>dubitata</i> , Matth.	68
<i>ACTINOPTERYX</i>	37	<i>alta</i> , Rey	5
<i>adustipenne</i> , Mots.	107	<i>excelsa</i> , Matth.	70
<i>affine</i> , Erichs.	83	<i>fauvelii</i> , Matth.	19
<i>affine</i> , Matth.	84	<i>femorialis</i> , Rey	5
<i>africanum</i> , Matth.	103	<i>fissicolle</i> , Reitt.	83
<i>americanus</i> , Casey	7	<i>flachi</i> , Gangl.	7
<i>angusta</i> , Matth.	59	<i>foveatum</i> , Matth.	98
<i>angustipennis</i> , Matth.	54	<i>fowlerianum</i> , Matth.	90
<i>anophthalma</i> , Joseph	7	<i>fratercula</i> , Matth.	53
<i>attenuatum</i> , Casey	95	<i>fuliginosa</i> , Matth.	68
<i>attenuatum</i> , Matth.	102	<i>fungi</i> , Mots.	109
<i>australis</i> , Matth.	37	<i>fungicola</i> , Casey	7
<i>bicoloratus</i> , Friedr.	7	<i>funiginus</i> , Casey	7
<i>bidilatata</i> , Matth.	69	<i>glauca</i> , Matth.	61
<i>biimpressa</i> , Reitt.	6	<i>godmani</i> , Matth.	52
<i>blackburni</i> , Matth.	76	<i>godmanni</i> , Matth.	52
<i>brasiliensis</i> , Matth.	59	<i>granulosum</i> , Casey ...	93
<i>braziliana</i> , Matth.	20	<i>granulum</i> , Mots.	5
<i>brenskii</i> , Flach	2	<i>grenadensis</i> , Matth.	69
<i>brevior</i> , Matth.	64	<i>grossa</i> , Mots	53
<i>cæcum</i> , Joseph	6	<i>gyrinoides</i> , Aubé	14
<i>californica</i> , Matth.	66	<i>hellenica</i> , Matth.	48
<i>CAMPTODIUM</i>	105	<i>helleni</i> , Flach	3
<i>carbonaria</i> , Matth.	67	<i>hemparti</i> , Flach	3
<i>caseianum</i> , Matth.	102	<i>hurnianum</i> , Matth.	80
<i>castanea</i> , Matth.	66	<i>humile</i> , Matth.	82
<i>castanea</i> , Rey	5	<i>hungarica</i> , Reitt.	7
<i>caucasica</i> , Matth.	47	<i>HYDROSCAPHA</i>	13
<i>caralli</i> , Broun	7	<i>ignobile</i> , Matth.	100
<i>championana</i> , Matth.	18	<i>impressum</i> , Matth. ...	87
<i>CHAMPIONELLA</i>	24	<i>impunctatum</i> , Matth.	100
<i>championis</i> , Matth.	58	<i>incerta</i> , Matth.	23
<i>cognata</i> , Matth.	57	<i>incognitum</i> , Matth.	84
<i>columbianum</i> , Matth.	81	<i>inconspicua</i> , Matth.	60
<i>concinnum</i> , Matth.	99	<i>japonica</i> , Matth.	74
<i>crotchianum</i> , Matth.	93	<i>japonicus</i> , Matth.	36
<i>crotchii</i> , Sharp	16		
<i>darwinii</i> , Matth.	62		
<i>depressa</i> , Matth.	63		

	PAGE		PAGE
<i>kraatzi</i> , Flach	3	PTERYCODES	27
<i>latitia</i> , Matth.	56	PTILODES	75
<i>laevicollis</i> , Matth.	47	PTILIUM	78
<i>lancifer</i> , Fauv.	7	PTINELLA	17
<i>laticolle</i> , Hoch.	6	<i>pubescens</i> , Matth.	77
<i>lawsoni</i> , Matth.	97	<i>pubescens</i> , Rey	5
<i>lederi</i> , Flach	2	<i>radicola</i> , Casey	7
<i>lederi</i> , Flach	3	<i>reitteri</i> , Flach	3
<i>lewisiana</i> , Matth.	18	<i>reitteri</i> , Flach	98
<i>lewisii</i> , Matth.	46	<i>reitteri</i> , Matth.	33
<i>LIMNEBIUS</i>	5	<i>reticulata</i> , Matth.	62
<i>lineare</i> , Matth.	91	<i>robustulum</i> , Casey	92
<i>longicorne</i> , Füss.	6	<i>rotundicolle</i> , Casey	92
<i>longipennis</i> , Casey	7	<i>rotundicollis</i> , Mots.	22
<i>lucida</i> , Matth.	49	<i>rufescens</i> , Matth.	50
<i>madagascariensis</i> , Schauf.	7	<i>rufilabris</i> , Flach	4
<i>maerens</i> , Matth.	72	<i>rufo-testaceum</i> , Matth.	88
<i>matthewsiana</i> , Woll.	21	<i>sahlbergi</i> , Flach	3
<i>MICOPILAGUS</i>	7	<i>salvinii</i> , Matth.	30
<i>MIKADO</i>	34	<i>scaphidiforme</i> , Hoch.	5
<i>minor</i> , Matth.	55	<i>schuleri</i> , Gangl.	7
<i>montivaga</i> , Schauf.	7	<i>schwarzi</i> , Flach	3
<i>NANOSELLA</i>	107	<i>seutellare</i> , Guilleb.	7
<i>natans</i> , Lec.	15	<i>seminitens</i> , Matth.	57
<i>NEPHANES</i>	77	<i>sharpi</i> , Reitt.	7
<i>nigrita</i> , Matth.	71	<i>sharpianum</i> , Matth.	94
<i>nitens</i> , Matth.	101	<i>similior</i> , Matth.	50
<i>nitidulum</i> , Rey	4	<i>simsoni</i> , Matth.	86
<i>NOSSIDIUM</i>	109	<i>smithsii</i> , Matth.	79
<i>nubigena</i> , Matth.	26	<i>soror</i> , Flach	48
<i>obcaecatium</i> , Matth.	89	<i>speculifer</i> , Matth.	105
<i>obesum</i> , Matth.	104	<i>spissata</i> , Rey	5
<i>obotrites</i> , Flach	3	<i>strangulatum</i> , Matth.	96
<i>obscurum</i> , Matth.	81	<i>subnitens</i> , Rey	5
<i>obsoletum</i> , Rey	4	<i>subtruncata</i> , Rey	5
<i>occidentalis</i> , Matth.	51	<i>sulcatum</i> , Casey	7
<i>oedipus</i> , Flach	89	<i>tesserula</i> , Matth.	73
<i>oertzeni</i> , Flach	4	TRICHOPTERYX	37
<i>opacina</i> , Rey	5	TRICHOPTILIUM	3
<i>orientale</i> , Flach	103	<i>tropicum</i> , Matth.	86
<i>ovulum</i> , Flach	2	<i>truncatissima</i> , Matth.	55
<i>pacifica</i> , Matth.	20	<i>truncatum</i> , Casey	7
<i>parallelipeda</i> , Matth.	64	<i>turgidulum</i> , Flach	3
<i>pellucida</i> , Matth.	22	<i>ulkei</i> , Matth.	97
<i>penzigi</i> , Flach	3	<i>varicorne</i> , Rey	4
<i>planum</i> , Matth.	85	<i>variolum</i> , Flach	3
<i>politum</i> , Matth.	91	<i>rexans</i> , Flach	3
<i>posthumum</i> , Matth.	109	<i>vicina</i> , Matth.	71
<i>posticale</i> , Broun	7	<i>vitrea</i> , Matth.	73
<i>proprium</i> , Broun	7	<i>xanthocera</i> , Matth.	65
<i>PTENIDIUM</i>	95		

GENERAL EXPLANATION OF PLATES IX—XV.

TRICHOPTERYGIA.

1. Upper side.
2. Under side.
3. Labrum.
4. Mandible.
5. Maxilla.
6. Mentum, &c.
7. Antenna.
8. Hind leg.
9. Under side of elytron.
10. Caudal organ of *Hydroscapha*.
11. Wing of *Hydroscapha*.

- a. prosternum.
- b. prosternal episternum.
- c. „ epimeron.
- d. mesosternum.
- e. mesosternal episternum.
- f. „ epimeron.
- g. metasternum.
- h. metasternal episternum.
- i. „ epimeron.
- j. mentum.
- k. labium.
- l. lingua.
- m. labial palpi.
- n. paraglossæ.
- o. paraglossal processes.
- p. inner lobe of maxilla.
- q. outer lobe of maxilla.
- r. process of stipes.
- s. maxillary palpus.
- t. mandible.
- u. process of stipes.
- v. lamella membranacea.
- w. epipleural fold.
- x. epipleurum.

Championella.

Dimorpha 11.

Mikado.

Nanosella.

Ptiliodes.

Campodius.

A. Matthew. di.

B.

W. S. S. S.

Hymenoptera

3 9088 00356542 1
nhent QL596 P83M3 supp
Trichopterygia illustrata et descripta