QL
522.1
A1B21
1910
ENT

HEMIPTERA
HETEROPTERA

QL
522.1
A1 B21
1910
ENT

CATALOGUE

OF THE

NEARCTIC

HEMIPTERA-HETEROPTERA

BY

NATHAN BANKS

PHILADELPHIA :
AMERICAN ENTOMOLOGICAL SOCIETY.
1910.

PRESS OF
P. C. STOCKHAUSEN
PHILADELPHIA

PREFACE.

No catalogue of our species of Heteroptera has hitherto been published. In the three volumes published of Lethierry and Severin's Catalogue of the Hemiptera are about half of our species, but the large family of Capsidæ and the Cryptocerata have never been catalogued.

Dr. Uhler's "List" has been very useful, but it was only a list, included Mexican species, and is now twenty years behind hand. The difficulties that I met have been largely eased by the kindness of Mr. Otto Heidemann, and especially valuable has been his help in the Capsidæ, placing in my hands some of his manuscript notes.

No order of American insects has been so largely controlled by Europeans as the Heteroptera. Not only have they described most of our species, but these descriptions are often in societies little known to American entomologist, and almost none of them in entomological journals. Volume after volume of our entomological journals is issued without a single article on Heteroptera, and when one is found it is short, or only a list of names. It is hoped that this catalogue will encourage entomologists to devote more time to this order, so that our forms will be better known to us.

The species are arranged in 30 families; those commonly accepted by Hemipterists. I have not used any superfamilies or higher groups, not because I do not believe in them, but because there is no arrangement so far proposed of such exceptional merit as to win general acceptance. Moreover, a catalogue is not the place to exhibit philosophic classifications, but to give references to the genera and species.

Altogether there are 1268 species in the catalogue; the family Capsidæ leads with 348, the Lygæidæ with 160, the Pentatomidæ with 149, and the Coreidæ with 108.

Undoubtedly many species remain to be described, but it will take many years to double the number.

What is needed especially are synoptic tables of genera and species to be printed in our entomological journals. Revisions of groups will be very valuable, but monographs are at present impossible. So many types are in Europe that before much revisional work can be done, it will be necessary for some one to take over a set of our species and compare them with the types.

CATALOGUE

OF THE

NEARCTIC

Hemiptera-Heteroptera

CORIXIDÆ.

CORIXA.

Geoffroy, Hist. Ins. i, 477, 1762.

Sigara Fabricius, Syst. Ent. 691, 1775.

Macrocorisa Thomson, Opusc. Ent. i, 28, 1869.

Glænocorisa Thomson, *ibid.* 39.

abdominalis Say, Het. Hem. N. Harm. 38, 1832; Compl. Writ. i, 366, 1859.—Fla.

bimaculata Guérin, Icon. Reg. Anim. iii, Ins. 354, 1845.

acuminata Uhler, Trans. Md. Acad. Sci. i, 392, 1897.—Tex., Ill.

alternata Say, Journ. Acad. Nat. Sci. Phil. iv, 329, 1825; Compl. Writ. ii, 251, 1859.—N. States.

bilineata Prov., Nat. Can. iv, 108, 1872.—Can.

bivittata Prov., Pet. Faune Ent. Can. Hem. 203, 1887.

brimleyi Kirk., Can. Entom. 1908, 120 (*Arctocorixa*).—N. Car.

burmeisteri Fieber, Abh. böhm. Ges. Wiss. (5), vii, 236, 1851.—N. A.

calva Say, Het. Hem., N. Harm. 38, 1832; Compl. Writ. i, 366, 1859.—U. S.

convexa Fieber, Abh. böhm. Ges. Wiss. (5), vii, 249, 1851.—Labr.

decolor Uhler, Packard, Silliman's Amer. Journ. Sci. 1871, 106.—Cal.

dispersa Uhler, Rept. U. S. Geog. and Geol. Surv. Terr. v, 841, 1875.—W. States.

erichsoni Fieber, Abh. böhm. Ges. Wiss. (5), vii, 247, 1851.—Pa.

escheri Heer, Insektenfauna von Oeningen, iii, Rhynchoten, 87, 1853 (foot note); Uhler, Bull. U. S. Geol. and Geog. Surv. Terr. ii, No. 5, 341, 1876.—W. States.

expleta Uhler, Hem. Colo. 63, 1895.—Colo.

fossarum Leach, Trans. Linn. Soc. Lond. xii, 17, 1818.—Calif.

germari Fieber, Bull. Soc. Mosc. 1848 (2), 531; Kirkaldy, Journ. Quekett Micr. Club, 1901, pl. 4, fig. 29.—Alaska.

- harrisii** Uhler, Proc. Bost. Soc. Nat. Hist. xix, 444, 1878.—E. States.
hydatotrepes Kirk., Can. Ent. 1908, 119 (*Arctocorixa*).—N. Car.
inscripta Uhler, Proc. Calif. Acad. Sci. (2), iv, 294, 1894.—S. W. States.
interrupta Say, Journ. Acad. Phil. iv, 328, 1825; Compl. Writ. ii, 250, 1859.—E. States.
kennicotti Uhler, Trans. Md. Acad. Sci. i, 393, 1897.—E. States.
kirbyi Fieber, Abh. böhm. Ges. Wiss. (5), vii, 255, 1851.—Can.
carinata Kirby, Rich. Faun. Bor. Am. iv, 284, 1837; Can. Ent. 1878, 215.
lævigata Uhler, Proc. Ent. Soc. Wash. ii, 384, 1891.—W. States.
lineata Forster, Nov. Spec. Ins. 70, 1771 (*Notonecta*).—N. Am.
macroceps Kirk., Can. Ent. 1908, 119 (*Arctocorixa*).—N. Car.
mercenaria Say, Het. Hem. N. Harm. 39, 1832; Compl. Writ. i, 367, 1859; Champ., Biol. Cent.-Amer. Hem. Het. pl. 22, fig. 23, 1901.—S. W. States.
nitida Fieber, Abh. böhm. Ges. Wiss. (5), vii, 240, 1851.—N. Car.
noveboracensis Gmelin, Syst. Nat. xiii, 2119, 1789.—N. Y.
planifrons Kirby, Rich. Faun. Bor. Amer. iv, 284, 1837; Can. Entom. 1878, 216.—Can.
serrulata Uhler, Trans. Md. Acad. Sci. i, 391, 1897.—S. W. States.
signata Fieber, Abh. böhm. Ges. Wiss. (5), vii, 233, 1851.—Pa., Ill.
stigmatica Fieber, Abh. böhm. Ges. Wiss. (5), vii, 248, 1851.—N. Am.
sutulis Uhler, Bull. U. S. Geol. Surv. ii, 339, 1876.—Colo.
tarsalis Fieber, Abh. böhm. Ges. Wiss. (5), vii, 231, 1851.—Pa.
trilineata Prov., Nat. Can. iv, 108, 1872.—Can.
irivittata Prov., Pet. Faune. Ent. Can. Hem. 203, 1887.
tumida Uhler, Bull. U. S. Geol. and Geog. Surv. Terr. iii (2), 454, 1877.—W. States.
verticalis Fieber, Abh. böhm. Ges. Wiss. (5), vii, 236, 1851.—Pa.
vulnerata Uhler, Proc. Acad. Nat. Sci. Phil. 1861, 284.—W. States.
wallengreni Stål, Eugenes Resa. Ins. 268, 1859.—Calif.
zimmermanni Fieber, Abh. böhm. Ges. Wiss. (5), vii, 233, 1851.—Pa., Car.

TENAGOBIA.

- Bergroth, Ent. Mo. Mag. xxxv, 282, 1899.
socialis White, Trans. Ent. Soc. Lond. 1879, 274 (*Sigara*).—Calif.

CALLICORIXA.

- White, Ent. Mo. Mag. x, 62, 1873.
præusta Fieber, Bull. Soc. Nat. Moscou, 1848 (2), 521; Abh. böhm. Ges. Wiss. (5), vii, 229, 1851.—Alaska.
kollari Fieber, Abh. böhm. Ges. Wiss. (5), vii, 229, 1851 (*Corisa*).—U. S.
cubæ Uhler, Proc. Zool. Soc. Lond. 1894, 224 (not Guérin).

NOTONECTIDÆ.

BUENOA.

Kirkaldy, Wien. Ent. Zeit. 1904, 120.

Anisops Spinola, Essai, 58, 1837 (part).

albidus Champ., Biol. Cent.-Am. Rhynch. ii, 373, 1901.—Tex.

carinatus Champ., Biol. Cent.-Amer. Rhyn. ii, 372, 1901.—N. Mex.

elegans Fieber, Abh. böhm. Ges. Wiss. (5), vii, 484, 1852.—E. States.

margaritacea Bueno, Journ. N. Y. Ent. Soc. 1908, 238.—E. States.

platycnemis Uhler, et auct. var. (nec Fieb.).

platycnemis Fieber, Abh. böhm. Ges. Wiss. (5), vii, 485, 1852.—Pa.

NOTONECTA.

Linné, Syst. Nat. Ed. x, 439, 1758.

howardi Bueno, Journ. N. Y. Ent. Soc. 1905, 151.—Ariz.

indica Linné, Mant. Plantarum, i, 534, 1771; Bueno, Journ. N. Y. Ent. Soc. 1905, 150.—W. States.

americana Fabr., Syst. Ent. 690, 1775.

unifasciata Guérin, Bull. Soc. Zool. Acclim. iv, 581, 1858.

insulata Kirby, Fauna Bor. Amer. Ins. 289, 1837; Can. Entom. 1878, 216; Bueno, Journ. N. Y. Ent. Soc. 1905, 162.—U. S.

impressa Fieb., Abh. böhm. Ges. Wiss. (5), vii, 475, 1851.

rugosa Fieb., ibid., 476.

irrorata Uhler, Proc. Bost. Soc. Nat. Hist. xix, 443, 1878; Bueno, Journ. N. Y. Ent. Soc. 1905, 159.—E. States.

lutea Bueno, Journ. N. Y. Ent. Soc. 1905, 160; (?) Müller, Zool. Dan. Prod. i, 1776.—Can.

mexicana Amyot et Serville, Hém. 453, 1843; Bueno, Journ. N. Y. Ent. Soc. 1905, 158.—Cal., Ariz.

klugeri Fieber, Abh. böhm. Ges. Wiss. (5), vii, 474, 1851.

montezuma Kirkaldy, Trans. Ent. Soc. Lond. 1897, 402; Champ., Biol. Cent.-Amer. Rhynch. ii, pl. 22, fig. 8, 1901; Bueno, Journ. N. Y. Ent. Soc. 1905, 162.—Calif.

raleighi Bueno, Can. Ent. 1907, 207.—N. Car.

shooterii Uhler, Proc. Cal. Acad. Sci. (2), iv, 292, 1894; Bueno, Jour. N. Y. Ent. Soc. 1905, 161.—Cal.

uhleri Kirkaldy, Ann. Mag. Nat. Hist. (6), xx, 58, 1897; Bueno, Journ. N. Y. Ent. Soc. 1905, 157.—E. States.

undulata Say, Het. N. Harm. 39, 1832; Compl. Writ. i, 368, 1859; Kirkaldy, Trans. Ent. Soc. Lond. 1897, 410; Champ., Biol. Cent.-Am. Rhynch. ii, 370, 1901; Bueno, Journ. N. Y. Ent. Soc. 1905, 152.—U. S.

americana Her. Schfr., Wanz. Ins. ix, 44, 1853 (nec Fabr.).

punctata Fieb., Abh. böhm. Ges. Wiss. (5), vii, 476.

variabilis Fieb. Abh. böhm. Ges. Wiss. (5), vii, 477; Bueno, Journ. N. Y. Ent. Soc. 1905, 165.—E. States.

PLEA.

Leach, Trans. Linn. Soc. London, xii, 11, 1817.

striola Fieber, Abh. böhm. Ges. Wiss. (5), iii, 296, 1845.—U. S.

NEPIDÆ.**NEPA.**

Linné, Syst. Nat. Ed. x, 440, 1758.

apiculata Harris, Ins. Inj. Veg. p. 12, pl. 1, fig. 1, 1862; Uhler, Proc. Bost. Soc. Nat. Hist. xix, 140, 1878; Montd., Bull. Soc. Sci. Bucarest, viii, 511, 1898; *ibid.* xii, 101, 1903.—E. States.

RANATRA.

Fabricius, Skrift. Nat. Selsk. i, 213, 1790.

fabricii Guérin, Sagra's Hist. de Cuba, vii, Ins. 176, 1857.—Tex.

fusca Beauv., Ins. Afric. et Amér. 235, 1805; Bueno, Can. Ent. 1905, 188.—U. S.

nigra Her.-Schfr., Wanz. Ins. iv, 32, 1838.

kirkaldyi Bueno, Can. Ent. 1905, 188.—E. States.

quadridentata Stål, Cfv. Kgl. Vet. Akad. Förh. 1864, 204; Bueno, Can. Ent. 1905, 188.—W. States.

See additions page 102.

BELOSTOMATIDÆ.**ABEDUS.**

Stål, Stett. Ent. Zeit. 1862, 461.

Stenoscytus Mayr., Verh. zool.-bot. Ges. Wien. xiii, 341, 1863.

ovatus Stål, Stett. Ent. Zeit. 1862, 461.—W. States.

mexicanus Mayr., Verh. zool.-bot. Ges. Wien. 1863, 347 (*Stenoscytus*).

PEDINOCORIS.

Mayr., Verh. zool.-bot. Ges. Wien. 1863, 341.

identata Hald., Proc. Ac. Phila. vi, 364, 1884 (*Zaittha*).—W. States.

brachonyx Mayr, Verh. zool.-bot. Ges. Wien, 1863, 351; Champ., Biol. Cent. Amer. Rhynch. ii, pl. xxi, fig. 22, 1901.

macronyx Mayr, Verh. zool.-bot. Ges. Wien, 1863, 350.—W. States.

DEINSTOMA.

Kirkaldy, Entom. xxx, 258, 1897.

Serphus Stål, Stett. Ent. Zeit. 1862, 462 (preoc).

dilatata Say, Het. N. Harm. 38, 1832; (*Belostoma*), Comp. Writ. i, 366, 1859; Stål, Stett. Ent. Zeit. 1861, 462 (*Serphus*).—W. States.

stollii Her. Schfr., Wanz. Ins. ix, 35, 1853 (*Zaittha*).

BELOSTOMA.

Latreille, Gen. Crust. Ins. iii, 144, 1807.

Amorgius Stål, Hem. Afric. iii, 179, 1865.

americanum Leidy, Journ. Acad. Phila. (2), i, 58, 1847.—U. S.

litigiosum Duf., Ann. Soc. Ent. France, 1863, 383.

angustipes Mayr, Verh. zool.-bot. Ges. Wien. 1871, 427; Champ., Biol. Cent. Amer. Rhynch. ii, pl. 22, fig. 5—Calif.

annulipes Her.-Schfr., Wanz. Ins. viii, 28, 1848; Mayr, Verh. zool.-bot. Ges. Wien, 1871, 424; Montandon, Ann. Soc. Ent. Belg. 1896, 514.—W. States.

ruficeps Dufour, Ann. Soc. Ent. France, 1863, 382.

signoreti Dufour, ibid.

obscurum Dufour, Ann. Soc. Ent. France, 1863, 383.—E. States.

grisea Mayr, Verh. zool.-bot. Ges. Wien. 1871, 427.

uhleri Mont., Ann. Soc. Ent. Belg., 1896, 513 (1897).—U. S.

americanum Howard, Insect Book, pl. xxix, fig. 25, 1902.

BENACUS.

Stål, Öfr. Vet. Akad. Förh. xix, 203, 1862.

griseus Say, Het. New Harm. 38, 1832 (*Belostoma*); Compl. Writ. i, 365, 1859; Riley, Proc. Ent. Soc. Wash iii, 86, 1895.—U. S.

haldemanum Leidy, Jour. Ac. Phila. (2) i, 1847, 66 (*Belostoma*).

impressus Hald., Pr. Ac. Nat. Sci. Phil., 1852, 364. (*Belostoma*).

harpax Stål, Öfr. Vet. Akad. Förh. xi, 240, 1854.

distinctum Dufour, Ann. Soc. Ent. France, 1863, 382.

ZAITHA.

Amyot et Serville, Hémipt. 430, 1843.

Perthostoma Leidy, Jour. Acad. Phila. (2) i, 59, 1847.

anurus Her. Schf., Wanz. Ins. viii, 26, 1848 (*Diplonychus*).—W. S.

cupreomicans Stål, Öfr. Vet. Akad. Förh., 1854, 240.

subspinosa Duf., Ann. Soc. Ent. France, 1863, 38.

aurantiacum Leidy, Jour. Acad. Phila. (2), i, 60, 1847 (*Perthostoma*).—E. S.

lutaria Stål, Öfr. Vet. Akad. Förh. xii, 190, 1855.

bifoveata Hald., Stanbury's Expt. Great Salt Lake, 370, 1852; pl. 10, fig. 1.—Utah.

fluminea Say, Het. Hem. N. Harm. 38, 1832; Compl. Writ. i, 364, 1859 (*Belostoma*); Dufour, Ann. Soc. Ent. France, 1863, 388.—U. S.

fusciventris Dufour, Ann. Soc. Ent. France, 1863, 389; Mayr, Verh. zool.-bot. Ges. Wien. 1871, 417; Champ. Biol. C. Am. Rhynch. ii, 365, pl. 21, fig. 23, 1901.—S. W. States.

micantula Stål, Svensk. Vet. Ak. Handl. (2), (No. 7), 84, 1860.—S. W. S.

minuscula Uhler, Stand. Nat. Hist. ii, 258, 1885.

minor Dufour, Ann. Soc. Ent. France, 1863, 391.—Calif.

testacea Leidy, Jour. Ac. Phila. (2), i, 66, 1847, (*Perthostoma*).—U. S.
reticulata Hald., Stanbury's Exped. Great Salt Lake, 370, 1852.

NAUCORIDÆ.

AMBRYsus.

Stål, Stett. Ent. Zeit., 1862, 439.

guttapennis Stål, Enum. Hem. v, 143, 1876; Mont., Verh. zool.-bot. Ges. Wien. 1897, 22.—Calif.

melanopterus Stål, Stett. Ent. Zeit., 1862, 460; Enum. Hem. v, 143, 1876; Mont., Verh. zool.-bot. Ges. Wien. 1897, 19; Champ. Biol. Cent. Amer. Rhynch.-Het. ii, pl. 21, fig. 7.—Tex., Ariz.

mormon Montd., Bull. Soc. Sci. Bucarest xviii, 48, 1909.—Utah.

oblongulus Mont., Verh. zool.-bot. Ges. Wien. 1897, 11; Champion, Biol. C. Am. Rhynch. Het. ii, pl. xxi, fig. 3.—Tex.

pudicus Stål, Stett. Ent. Zeit. 1862, p. 460. Champion, Biol. C. Am. Rhynch. Het. ii, pl. xxi, fig. 6.—Cal.

puncticollis Stål, Enum. Hem. v, 143, 1876; Mont., Verh. zool.-bot. Ges. Wien. 1897, 20.—Tex., Ariz.

signoreti Stål, Stett. Ent. Zeit. 1862, 460; Mont., Verh. zool.-bot. Ges. Wien. 1897, 23.—W. States.

PELOCORIS.

Stål, Enum. Hem. v, 144, 1876.

carolinensis Bueno, Can. Entom. 1907, 227.—S. States.

poeyi Guérin, Iconog. Règne Anim. iii, Ins. 352, 1845; Sagra's Hist. de Cuba viii, Ins. 175, 1857 (*Naucoris*).—E. States.

GELASTOCORIDÆ.

(*Galgulidæ*.)

PELOGONUS.

Latreille, Gen. Crust. Ins. iv, 384, 1809.

americanus Uhler, Bull U. S. Geol. and Geog. Surv. Terr. ii, No. 5, 335, 1876.—U. S.

GELASTOCORIS.

Kirkaldy, Entom. 1897, 258.

Galgulus Latreille, Hist Nat. Ins. iii, 253, 1802 (preoc).

oculatus Fabr., Ent. Syst. Suppl. 525, 1798 (*Naucoris*); Syst. Rhyng. 111, 1803; Latr., Hist. Nat. Ins. xii, 386, 1809. Lugger, Sixth Ann. Rept. Ent. Minn. Exp. Sta. 23, 1900, fig. 16.—U. S.

variegatus Guérin, Iconog. Règne Anim. iii, Ins. 352, 1845 (*Galgulus*); Champ., Biol. C. Am. Rhynch. ii, 349, 1901.

pulcher Stål, Cefv. Kgl. Vet. Akad. Förh. 1854, 239.—Fla., Tex.

vicinus Champion, Biol. Cent. Amer. Rhynch. ii, 349, 1901.—Calif.

MONONYX.

Laporte, Essai, 15, 1832.

Phintius Stål, Cefv. Vet. Akad. Förh. xviii, 201, 1861.

fuscipes Guérin, Rev. Zool. 1843, 114.—Calif.

badius Her. Schf., Wanz. Ins. ix, 27, 1853.

obscurus Stål, Cefv. Vet. Akad. Förh. xi, 239, 1854.

nepæformis Fabr., Syst. Ent. 693, 1775 (*Naucoris*),—S. States.

bipunctatus Stål, Cefv. Vet. Akad. Förh. xi, 239, 1854; Champ.,

Biol. Cent. Amer. Rhynch. ii, pl. 20, figs. 27, 27a, 1901.

NERTHRA.

Say, Descr. Het. Hem. 37, 1832; Compl. Writ. i, 364, 1859.

stygica Say, Descr. Het. Hem. New Harm. 37, 1832 (*Naucoris*).

Compl. Writ. i, 364, 1859; Bueno, Ohio Nat. v, 288, 1905.—Ga., Fla.

SALDIDÆ.

SALDA.

Fabricius, Syst. Rhyng. 113, 1803.

Acanthia, of various writers.

anthracina Uhler, Bull. U. S. Geol. and Geog. Surv. Terr. iii (2), 438, 1877.—E. States.

bouchervillei Prov., Nat. Canad. iv, 106, 1872 (*Sciadopterus*); Faune Ent. Can. Hem. 192, 1886.—Can.

confluens Say, Het. N. Harm. 35, 1832 (*Acanthia*); Comp. Writ. i, 361, 1859. Uhler, Bull. U. S. Geol. and Geog. Surv. Terr. iii (2), 433, 1877.—E. States.

coriacea Uhler, Fifth Ann. Rept. U. S. Geog. Surv. f. 1871, 1872, p. 421, 1872; Bull. U. S. Geol. and Geog. Surv. Terr. iii (2), 436, 1877; Proc. Bost. Soc. Nat. Hist. xix, 433, 1878; Bull. 69 Minn. Sta., fig. 18.—N. States.

crassicornis Uhler, Bull. U. S. Geol. and Geog. Surv. Terr. iii (2), 438, 1877.—Can.

deplanata Uhler, Bull. U. S. Geol. and Geog. Surv. Terr. iii (2), 442, 1877.—U. S.

dispersa Uhler, Proc. Ent. Soc. Wash. ii, 383, 1893.—Ut. Colo.

elongata Uhler, Bull. U. S. Geol. and Geog. Surv. Terr. iii (2), 448, 1877.—Can.

explanata Uhler, Proc. Ent. Soc. Wash. ii, 383, 1893.—Ut.

- hirta** Say, Het. N. Harm. 34, 1832 (*Acanthia*); Comp. Writ. i, 359, 1859; Uhler, Bull. U. S. Geol. and Geog. Surv. Terr. iii (2), 436, 1877.—U. S.
- humilis** Say, Het. N. Harm. 35, 1832 (*Acanthia*); Comp. Writ. i, 360, 1859; Uhler, Bull. U. S. Geol. and Geog. Surv. Terr. iii (2), 451, 1877.—U. S.
- interstitialis** Say, Jour. Acad. Phil. iv, 324, 1825 (*Acanthia*); Comp. Writ. ii, 248, 1859; Uhler, Bull. U. S. Geol. and Geog. Surv. Terr. iii (2), 444, 1877.—U. S.
- laticollis** Reuter, Pet. Nouv. Entom. 1875, 544; Öfv. Finsk. Vet. Förh. xxx, 61, 1879 (*Acanthia*).—Alaska.
- ligata** Say, Het. N. Harm. 34, 1832 (*Acanthia*); Comp. Writ. i, 359, 1859; Uhler, Bull. U. S. Geol. and Geog. Surv. Terr. iii (2), 432, 1877.—U. S.
- variegata* Prov., Nat. Can. iv, 107, 1872.
- littoralis** Linn., Syst. Nat. x, 442, 1758 (*Cimex*); Uhler, Bull. U. S. Geol. and Geog. Surv. Terr. iii (2), 439, 1877; Reuter, Acta. Soc. Sci. Fenn. xxi (2), 8, 1895.—N. States, Can.
- luctuosa** Stål, Eugenes Resa, Hemipt. 263, 1859; Uhler, Bull. U. S. Geol. and Geog. Surv. Terr. iii (2), 445, 1877.—Cal.
- lugubris** Say, Het. N. Harm. 34, 1832; (*Acanthia*), Comp. Writ. i, 360, 1859; Uhler, Bull. U. S. Geol. and Geog. Surv. Terr. iii, 442, 1877.—U. S.
- major** Prov., Nat. Canad. iv, 107, 1872; Pet. Faune Ent. Can. Hem. 190, 1886.—Can.
- obscura** Prov., Nat. Canad. iv, 107, 1872; Pet. Faune Ent. Can. Hem. 190, 1886.—Can.
- opacula** Zett., Ins. Lapp. 268, 1838; Reuter, Acta Soc. Sci. Fenn. xxi, 19, 1895.—Mass.
- orbiculata** Uhler, Bull. U. S. Geol. and Geog. Surv. Terr. iii (2), 450, 1877.—U. S.
- pallipes** Fabr., Syst. Rhyng. 115, 1803 (*Acanthia*); Her. Schfr., Wanz. Ins. vi, fig. 600, 1842; Uhler, Bull. U. S. Geol. and Geog. Surv. Terr. iii (2), 446, 1877; Reuter, Acta Soc. Sci. Fenn. xxi (2), 21, 1895.—U. S.
- pellita** Uhler, Bull. U. S. Geol. and Geog. Surv. Terr. iii (2), 433, 1877.—Mass.
- polita** Uhler, Bull. U. S. Geol. and Geog. Surv. Terr. iii (2), 441, 1877.—Colo., Can.
- reperta** Uhler, Bull. U. S. Geol. and Geog. Surv. Terr. iii (2), 447, 1877.—N. States.
- saltatoria** Linn., Syst. Nat. (x), 448, 1758 (*Cimex*); Reuter, Acta Soc. Sci. Fenn. xxi (2), 15, 1895.—N. States, Can.
- scotica** Curtis, Brit. Entom. xii, pl. 548, 1835; Reuter, Acta Soc. Sci. Fenn. xxi (2), 12, 1895.—Ill., Utah.

- separata** Uhler, Proc. Bost. Soc. Nat. Hist. xix, 432, 1878.—N. E. States.
- signoretii** Guérin, Sagra's Hist. Nat. de Cuba viii, Ins. 167, 1857; Stål, Enum. Hemipt. iii, 148, 1873; Uhler, Bull. U. S. Geol. and Geog. Surv. Terr. iii. (2), 431, 1877; Champ., Biol. C. Am. Rhynch. ii, 339, 1900.—E. States.
- ornata* Stål, Stett. Ent. Zeit. 1862, 458.
- sphacelata** Uhler, Bull. U. S. Geol. and Geog. Surv. Terr. iii (2), 434, 1877.—E. States.
- stellata** Curtis, Ross's Second Voyage, Arctic Regions, app. lxxv. 1835 (*Acanthia*); Uhler, Bull. U. S. Geol. and Geog. Surv. Terr. iii (2), 442, 1877.—B. Am.
- xanthochila** Fieber var. *limbosa* Horvath, Rev. d'Ent. x, 80, 1891; Ann. Mus. Nat. Hung. vi, 566, 1908.—E. States.

SALDOIDA.

Osborn, Can. Entom. 1901, 181.

cornuta Osborn, Can. Entom. 1901, 182.—Fla.

slossoni Osborn, Can. Entom. 1901, 181.—Fla.

REDUVIIDÆ.

Reduviini.

ACHOLLA.

Stål, Stett. Ent. Zeit. 1862, 445.

Ascra Stål, *ibid*, 445.

ampliata Stål, Enum. Hemipt. ii, 72, 1872.—N. Mex.

multispinosa De Geer, Mém. Ins. iii, 23, 1773 (*Cimex*).—U. S.

sexspinosus Wolff, Icones. Cim. iii, 124, 1802 (*Reduvius*); Stål, Stett. Ent. Zeit. 1862, 445.

subarmatus Her. Schf., Wanz. Ins. viii, 83, 1848.

tabida Stål, Stett. Ent. Zeit. 1862, 446 (*Ascra*); Enum. Hemipt. ii, 72, 1872.—Calif.

ARILUS.

Hahn, Wanz. Ins. i, 33, 1831.

Prionotus Laporte, Essai, 8, 1832.

Prionotus Uhler, Check-list, 23, 1886.

cristatus Linn., Cen. Ins. Rar. 16, No. 42, 1763 (*Cimex*); Syst. Nat. (XII. Ed.), 723, 1767; Stål, Enum. Hem. ii, 72, 1872.—U. S.

novenarius Say, Amer. Ent. ii, 31, 1825 (*Reduvius*); Compl. Writ. i, 71, 1859.

denticulatus Westw., Drury's Illust. (new ed.), ii, 73, 1842.

ATRACHELUS.

Amyot at Serville, Hem. 374, 1843.

cinereus Fabr., Ent. Syst. Supp. 545, 1794 (*Reduvius*); Syst. Rhyng. 287, 1803.—S. Sates.

heterogeneus Am. and Serv. Hémipt. 374, 1843.

FITCHIA.

Stål, Öfv. Vet. Akad. Förh. 1859, 370.

* **aptera** Stål, Öfv. Vet. Akad. Förh. 1859, 371.—E. States.
nigrovittata Stål, *ibid*, 1866, 296.

spinosula Stål, Enum. Hemipt. ii, 79, 1872.—S. States.

HEZA.

Amyot et Serville, Hem. 374, 1843.

annulicornis Stål, Enum. Hem. ii, 77, 1872.—S. States.

PSELLIOPUS.

Bergroth, Rev. d'Entom. 1905, 112.

Milyas Stål, Rio Hem. ii, 62, 1858 (preoc).

cinctus Fabr., Gen. Insect. 302, 1776 (*Reduvius*); Ent. Syst. iv, 199, 1794; Her.-Sch., Wanz. Ins. viii, 83, 1848 (*Harpactor*);
Stål, Hem. Fabric. i, 106, 1868.—U. S. 5. 5†.

præcinctus Gmelin, Syst. Nat. i, 2198, 1788.

spinicollis Champ., Biol. Cent. Amer. Rhynch. ii, p. 245, 1899, pl. xv, fig. 2 (*Milyas*).—Calif.

zebra Stål, Stett. Ent. Zeit. 1862, 448; Enum. Hem. ii, 87, 1872 (*Milyas*).—W. S.

REDUVIUS.

Lamarck, Syst. Nat. 294, 1801.

Opsicætus Klug, Symb. Phys. ii, pl. ix, 1830.

personatus Linn., Fauna Suec. 942, 1761 (*Cimex*); Fabr., Syst. Rhyng. 267, 1803; Hahn, Wanz. Ins. ii, 17, 1834; Reuter, Act. Soc. Sci. Fenn. xix (15), 6, 1892; Bull. 22 Bur. Entom., fig. 18.—U. S.

quisquilius De Geer, Mém. Ins. iii, 281, 1773 (*Cimex*).

albosignatus Prov., Nat. Can. iv. 105, 1872.

senilis Van Duzee, Ent. News, 1906, 390.—Ariz.

SINEA.

Amyot et Serville, Hém. 375, 1843.

complexa Caudell, Can. Entom. 1900, 67; Jour. N. Y. Ent. Soc. 1901, 7.—Calif., Ariz.

- confusa** Caudell, Jour. N. Y. Ent. Soc. 1901, 6.—Ariz.
multispinosa Stål, Stett. Ent. Zeit. 1862, 443 (part, not De Geer).
diadema Stål, Enum. Hemipt. ii, 70, 1872 (part, not Fabr.).
undulata Cham., Biol. Cent.-Amer. Rhyn. ii, 292, 1899 (not Uhler).
coronata Stål, Stett. Ent. Zeit. 1862, 444; Caudell, Jour. N. Y. Ent. Soc. 1901, 7. —W. States.
diadema Fabr., Gen. Insect. 302, 1776 (*Reduvius*); Stål, Enum. Hem. ii, 70, 1872; Caudell, J. N. Y. Ent. Soc. 1901, 4.—U. S.
hispidus Thunberg, Nov. Ins. Sp. ii, 33, 1783 (*Cimex*); Stål, Œfv. Akad. Förh. 1866, 264 (*Irantha*).
celosus Gmelin, Syst. Nat. I, iv, 21, 1788.
raptatorius Say, Jour. Acad. Nat. Sci. Phil. iv, 327, 1825; Compl. Writ. i, 72, 1859.
multispinosa Am. et Serv. Hémipt. 375, 1843 (not De Geer); Stål, Stett. Ent. Zeit. xxii, 139, 1861; Walker, B. M. Cat. Hem. viii, 138, 1873.
raptoria Stål, Stett. Ent. Zeit. 1862, 444; Uhler, Bull. U. S. Geol. and Geog. Surv. Terr. No. 5, 1876, 326; Caudell, Jour. N. Y. Ent. Soc. 1901, 9.—U. S.
rileyi Mont., Proc. U. S. Nat. Mus. xvi, 51, 1893; Caudell, Jour. N. Y. Ent. Soc., 1901, 10.—S. States.
sanguisuga Stål, Stett. Ent. Zeit. 1862, 444; Enum. Hem. ii, 71, 1872; Candell, Journ. N. Y. Ent. Soc. 1901, 9.—S. W. S.
spinipes Herr. Schfr., Wanz. Ins. viii, 82, 1848 (*Harpactor*); Stål, Enum. Hem. ii, 71, 1872; Caudell, Jour. N. Y. Ent. Soc. 1901, 10.—W. States.
undulata Uhler, Proc. Calif. Acad. Sci. iv, 282, 1894; Caudell, Jour. N. Y. Ent. Soc. 1901, 6.—W. States.

Zelini.

REPITA.

Stål, Œfv. Vet. Akad. Förh. 1859, 369.

- taurus** Fabr., Syst. Rhyng. 291, 1803 (*Zelus*); Stål, Stett. Ent. Zeit. 1862, 446; Enum. Hem. II, 80, 1872; Champ., Biol. C. Am. Rynch. Het. II, 269, 1899, pl. xvi, fig. 13.—S. States.
lineatus Amyot et Serv. Hémipt. 373, 1843 (*Zelus*).

ROCCONOTA.

Stål, Œfv. Vet. Akad. Förh. 1859, 370.

- annulicornis** Stål, Enum. Hemipt. ii, 77, 1872 (*Heza*); Champ., Biol. C. Am. Rhynch. ii, 273, 1899, pl. xvi, fig. 21.—S. States.

ZELUS.

Fabricius, Syst. Rhyng. 281, 1803.

- Diplodus* Am. et Serv. Hém. 370, 1843.
Pindus Stål, Stett. Ent. Zeit. 1862, 454.
Diplacodus Kirkaldy, Entom. 1900, 242.

- bilobus** Say, Insects, Louis. 12, 1832; Compl. Writ. i, 306, 1859; Stål, Enum. Hem. ii, 88, 1872.—S. States.
- cervicalis** Stål, Enum. Hemipt. ii, 90, 1872.—W. States.
- **exsanguis** Stål, Stett. Ent. Zeit. 1862, 452; Enum. Hem. ii, 91, 1872; Champ. Biol. C.-Am. Rhynch. Het. ii, 259, 1899, pl. xv, fig. 23.—U. S.
- **luridus** Stål, Stett. Ent. Zeit. 1862, 452; Enum. Hem. ii, 91, 1872.
- georgiæ** Prov., Nat. Can. iv, 106, 1872 (*Darbanus*).—Ga.
- lævicollis** Champ. Biol. Cent.-Amer. Rhynch. Het. ii, 260, pl. xv, fig. 24, 1899.—Tex.
- marginatus** Prov., Faun. Ent. Can. Hem. 182, 1887 (*Euagorus*).—Can.
- palliatu**s Prov., Faun. Ent. Can. Hem. 182, 1887 (*Darbanus*).—Can.
- renardii** Kolenati, Melet. Entom. vi, 460, 1857 (*Zelus*); Stål, Enum. Hem. ii, 91, 1872.—W. States.
- socius** Uhler, Hayden's Surv. Mont., 420, 1872 (*Pindus*).—W. States.

*Apiomerini.***APIOMERUS.**

Hahn, Wanz. Ins. i, 29, 1831.

Herega Amyot et Serv., Hém. 354, 1843.

- crassipes** Fabr., Syst. Rhyng. 273, 1803 (*Reduvius*); Say, Amer. Entom. ii, pl. 31, fig. 4, 1825; Stål, Hemipt. Fabric. i, 117, 1868; Bull. 69 Minn. Sta., fig. 27.—U. S.
- linitaris* Say, Het. N. Harm. 31, 1832; Compl. Writ. i, 355; 1859.
- rubrolimbata* Am. et Serv., Hémipt. 354, 1843.
- flaviventris** Her.-Schr., Wanz. Ins. viii, 77, 1848; Stål, Enum. Hemipt. ii, 98, 1872.—W. States.
- longispinis** Champ., Biol. Cent.-Amer. Rhynch.-Het. ii, p. 239, 1899, pl. xiv. fig. 18.—Ariz.
- pictipes** Her.-Schfr., Wanz. Ins. viii, 75, 1848; Stål, Enum. Hem. ii, 98, 1872; Champ., Biol. C.-Am. Rhynch. Het. ii, pl. xiv, figs. 28, 29.—S. W. States.
- repletus** Uhler, Bull. U. S. Geol. and Geog. Surv. Terr. ii, No. 5, 329, 1876.—Calif.
- spissipes** Say, Jour. Acad. Phil. iv, 328, 1825 (*Reduvius*); Compl. Writ. i, 72, 1859; Stål, Enum. Hemipt. ii, 98, 1872; Champion, Biol. C.-Am. Rhynch. ii, 241, 1899, pl. xiv, fig. 25; Bull. 63 pt. iv, Bur. Entom., Dept. Agric.—W. States.
- subpiceus** Stål, Stett. Ent. Zeit. 1862, p. 454; Champ., Biol. Cent.-Amer. Rhynch. Het. ii, 237, 1899, pl. xiv, fig. 12.—Ariz.
- ventralis** Say, Het. N. Harm. 31, 1832 (*Ruduvius*); Compl. Writ. i, 355, 1859.—W. States.

HARPACTOR.

Laporte, Essai, 8, 1832.

americanus Bergroth, Ent. News, 1897, 96.—Calif., Wash.

Hammatocerini.

HAMMATOCERUS.

Burmeister, Handb. ii, 235, 1835.

Hammacerus Laporte, Essai. 79, 1832.

luctuosus Stål, Cefv. Vet. Akad. Förh. 1854, 237 ; Enum. Hemipt. ii, 100, 1872 ; Champion, Biol. Cent.-Am. Rhynch. ii, 227, 1899.—S. W. States.

purcis Drury, Illust. iii, 63, 1782 (*Cimex*) ; Blanchard, Hist. Ins. iii, 105, 1840 (as *furcis*).—S. States.

nycthemerus Burm., Handb. ii, 236, 1835.

HOMALOCORIS.

Perty, Delect. Anim. 216, 1834.

maculicollis Stål, Enum. Hem. ii, 101, 1872.—Ariz.

Ectrichodini.

ECTRICHODIA.

Lepell. et Serv., Encyc. Meth. x, 279, 1825.

Rhiginia Stål, Cefv. Vet. Ak. Förh. 1859, 176.

cinctiventris Stål, Enum. Hemipt. ii, 103, 1872.—Tex., N. Mex.

cruciata Say, Het. N. Harm. 33, 1832 (*Petalocheirus*) ; Compl. Writ. i, 358, 1859 ; Stål, Enum. Hemipt. ii, 103, 1872.—U. S.

bicolor Her.-Schfr., Wanz. Ins. viii, 53, 1848 (*Ectrychotes*).

crudelis Stål, Stett. Ent. Zeit. 1862, 455 (*Rhiginia*).

media Walk., B. M. Cat. Hem. viii, 62, 1873.

POTHEA.

Amyot et Serville, Hém. 344, 1843.

æneo-nitens Stål, Ann. Soc. Ent. France, 1864, 59 ; Enum. Hem. ii, 104, 1872.—W. States.

Conorhinini.

CONORHINUS.

Laporte, Essai. 78, 1832.

gerstæckeri Stål, Berl. Ent. Zeit. iii, 111, 1859 ; Hem. Fabr. i, 124, 1868.—Tex.

protractus Uhler, Proc. Calif. Acad. Sci. (2) iv, 284, 1894.—Calif.

rubrofasciatus De Geer, Mém. Ins. iii, 349, 1773 (*Cimex*) ; Am. et Serv. Hémipt. 384, 1843 ; Stål, Berl. Ent. Zeit. iii, 106, 1859.—S. States.

- gigas* Fabr., Syst. Ent. 729, 1775 (*Reduvius*); Her.-Schfr., Wanz. Ins. viii, 72, 1848.
erythrozonias Gmelin, Syst. Nat. i, 2181, 1788 (*Cimex*).
stalii Sign., Ann. Soc. Ent. France (3) viii, 967, 1860.
sanguisugus Le Conte, Proc. Acad. Nat. Sci. Phil. vii, 404, 1855;
 Insect Life ii, fig. 17; Bull. 22, Div. Entom., fig. 22.—
 S. States.
lateralis Stål, Berl. Ent. Zeit. iii, 107, 1859; Hem. Fabr. i, 124,
 1868.
variegatus Drury, Illust. i, 109, 1772 (*Cimex*).—S. States.
claviger Gmelin, Syst. Nat. i, 2179, 1788 (*Cimex*).
lecticularis Stål, Berl. Ent. Zeit. iii, 107, 1859; Hemipt. Fabric.
 i, 124, 1868.

MECCUS.

- Stål, Berl. Ent. Zeitschr. iii, 105, 1859.
phyllosomus Burm., Handb. ii, 246, 1835 (*Conorhinus*); Stål, Berl.
 Ent. Zeit. iii, 105, 1859.—Calif.

SPINIGER.

- Burmeister, Handb. ii, 234, 1835.
Acrocoris Hahn, Wanz. Ins. iii, 22, 1835.
bicolor Stål, Stett. Ent. Zeit. 1859, 396; Berl. Ent. Zeitschr. xiii, 234,
 1869.—Tex., Ariz.

Piratini.

MELANOLESTES.

- Stål, Cefv. Vet. Ak. Förh. 1866, 231.
abdominalis Her.-Schfr., Wanz. Ins. viii, 63, 1848 (*Pirates*), fig. 832;
 Bull. 22 Div. Entom., fig. 19.—U. S.
picipes Her.-Schfr., Wanz. Ins. viii, 62, 1848 (*Pirates*); Stål, Enum.
 Hemipt. ii, 107, 1872; Bull. 69 Minn. Sta., fig. 28.—U. S.
pungens Le Conte, Proc. Acad. Nat. Sci. Phil., 1855, 404.

RASAHUS.

- Amyot et Serville, Hém. 325, 1843.
Macrosandalus Stål, Cefv. Vet. Ak. Förh. 1866, 251.
biguttatus Say, Ins. Louis. 13, 1832 (*Petalochirus*); Comp. Writ. i,
 307, 1859; Stål, Stett. Ent. Zeit. 1862, 457; Enum. Hem. ii,
 106, 1872; Bull. 22, Bur. Ent., fig. 21.—S. States.
mutillarius Guérin, Sagra's Hist. de Cuba, Ins. 172, 1857 (*Pirates*).
hamatus Fabr., Spec. Ins. ii, 381, 1781 (*Reduvius*); Stål, Enum.
 Hemipt. ii, 106, 1872; Champion, Biol. C.-Amer. Rhynch.
 ii, 217, 1899.—S. W. States.
thoracicus Stål, Enum. Hem. ii, 106, 1872 (*Macrosandalus*).—S.
 States.

SIRTHENEA.

Spinola, Essai, 100, 1840.

carinata Fabr., Ent. Syst. Supp. 545, 1794 (*Reduvius*); Serv., Ann. Sci. Nat. xxiii, 221, 1830 (*Peirates*); Amyot et Serv., Hémipt. 326, 1843 (*Rasahus*); Stål, Hem. Fabric. i, 120, 1868.—S. States.

roseus Her. Schf., Wanz. Ins. viii, 62, 1848 (*Pirates*).

Saicini.

ONCEROTRACHELUS.

Stål, Hem. Fabr. i, 130, 1868.

acuminatus Say, Het. N. Harm. 31, 1832 (*Reduvius*); Compl. Writ. i, 356, 1859; Stål, Enum. Hemipt. ii, 124, 1872; Champ., Biol. C.-Am. Rhynch. Het. ii, pl. x, fig. 8.—U. S.

Stenopodini.

DIADITUS.

Stål, Œfv. Vet. Akad. Förh. 1859, 383.

pictipes Champion, Biol. C.-Am. Rhynch. ii, 189, 1898, pl. xi, fig. 21.—Tex.

GNATHOBLEDA.

Stål, Œfv. Vet. Akad. Förh. 1859, 380.

tumidula Stål, Enum. Hemipt. ii, 121, 1872.—S. States.

NARVESUS.

Stål, Œfv. Vet. Akad. Förh. 1859, 384.

carolinensis Stål, Œfv. Vet. Akad. Förh. 1859, 385; Enum. Hemipt. ii, 124, 1872; Champ., Biol. C.-Am. Rhynch. Het. ii, pl. xi, fig. 19.—E. States.

ONCOCEPHALUS.

Klug, Symb. Phys. 2, 1830.

Spilaloni Stål, Enum. Hem. ii, 123, 1872.

apiculatus Reuter, Act. Soc. Fenn. xii, 728, 1882.—Missouri.

geniculatus Stål, Enum. Hemipt. ii, 123, 1872 (*Spilaloni*); Reuter, Act. Soc. Fenn. xii, 727, 1882.—S. States.

PNIRONTIS.

Stål, Œfv. Vet. Akad. Förh. 1859, 381.

Centromelus Fieber, Europ. Hem. 42, 1861.

infirm Stål, Œfv. Vet. Akad. Förh., 1859, 382; Champion, Biol. C.-Am. Rhynch. ii, 183, 1898, pl. xi, fig. 11.—S. States.

languida Stål, Œfv. Vet. Akad. Förh. 1859, 382; Enum. Hemipt. ii, 120, 1872 (*Centromelus*); Champion, Biol. Cent.-Amer. Rhynch. ii, 183, 1898, pl. xi, fig. 13.—S. States.

PYGOLAMPIS.

Germar, Faun. Insect. Europ. 8 fasc, 1817 (?), 1827 (?).

Ochetopus Hahn, Wanz. Ins. i, 176, 1831.

• **pectoralis** Say, Insects Louis. 11, 1832 (*Reduvius*); Compl. Writ. i, 306, 1859; Uhler, Proc. Bost. Soc. N. H., xiv, 15, 1871.—E. States.

fuscipennis Stål, Cefv. Vet. Akad. Förh. 1859, 379; Enum. Hem. iv, 85, 1874.

serica Stål, Cefv. Vet. Akad. Förh. 1859, 380.—S. States.

SCHUMANNIA.

Champion, Biol. C.-Am. Rynch. Het. ii, 185, 1898.

mexicana Champ., Biol. C.-Am. Rhynch. Het. ii, 185, 1898, pl. xi, fig. 18.—N. Car.

STENOPODA.

Laporte, Essai, 26, 1832.

culiciformis Fabr., Syst. Ent. 728, 1775 (*Cimex*); Spec. Ins. ii, 376, 1781; Ent. Syst. iv. 189, 1794; Stål, Hemipt. Fabric. i, 129, 1868.—U. S.

cinerea Lap., Essai. Hem. 26, 1832; Amyot et Serv., Hem. 390, 1843.

EMESIDÆ.**BARCE.**

Stål, Hem. Afric. iii, 163, 1865.

• **annulipes** Stål, Berl. Ent. Zeits. 1866, 168; Enum. Hem. ii, 127, 1872; Banks, Psyche, 1909, 47.—E. States.

fraterna Say, Hem. N. Harm. 33, 1832 (*Ploiaria*) Compl. Writ. i, 358, 1859; Dohrn, Linn. Ent. xiv. 232, 1860; Banks, Psyche, 1909, 47.—E. States.

simplicipes Uhler, Proc. Bost. Soc. Nat. Hist. xix, 430, 1878 (*Emesodema*).

uhleri Banks, Psyche, 1909, 47.—E. States.

uhleri var. **brunnea** Banks, Psyche, 1909, 47.—N. Y.

EMESA.

Fabr., Syst. Rhyng, 263, 1803.

brevicoxa Banks, Psyche, 1909, 48.—Calif.

longipes De Geer, Mém. Ins. iii. 352, 1773 (*Cimex*); Dohrn, Linn. Ent. xiv, 221, 1860; Uhler, Proc. Bost. Soc. N. H. xiv, 15, 1871; Banks, Psyche, 1909, 48.—U. S.

brevipennis Say, Amer. Ent. iii, pl. 47, 1828; Compl. Writ. i, 106, 1859.

pia Am. et Serv., Hémipt. 394, 1843.

filum G. R. Gray, Griffith's Anim. Kingd. xv. 244, 1832.

affinis Dohrn, Linn. Ent. xiv, 222, 1860.

GARDENA.

Dohrn, Linn. Ent. xiv, 214, 1860.

americana Champ., Biol. C.-Am. Rhynch. Het. ii, 167, 1898.—Tex.

LUTEVA.

Dohrn, Linn. Ent. xiv, 214, 1860.

arizonensis Banks, Psyche, 1909, 45.—Ariz.

PLOIARIA.

Scopoli, Del. Faun. Insubr. i, 60, 1786.

carolina Her.-Schfr., Wanz. Ins. ix, 115, 1853; Dohrn, Linn. Ent. xiv, 250, 1860 (*Luteva*); Banks, Psyche, 1909, 44.—N. Car.

texana Banks, Psyche, 1909, 44.—Tex.

PLOIARIODES.

White, Ann. Nat. Hist. (5) vii, 58, 1881.

Cerascopus of Uhler's list.

californica Banks, Psyche, 1909, 46.—Calif.

errabunda Say, Hem. N. Harm. 34, 1832; Compl. Writ. i, 359 1859; Uhler, Proc. Bost. Soc. Nat. Hist. xiv, 197, 1871; Banks, Psyche, 1909, 46.—E. States.

maculata Hald., Proc. Acad. Phil. iii, 151, 1847 (*Ploiaria*).

tuberculata Banks, Psyche, 1909, 46.—E. States.

PLOIARIOPSIS.

Champion, Biol. C.-Am. Rhynch. Het. ii, 173, 1898.

hirticornis Banks, Psyche, 1909, 44.—N. Car.

ENICOCEPHALIDÆ.**ENICOCEPHALUS.**

Westwood, Trans. Ent. Soc. London, ii, 1837, 22.

Hymenocoris Uhler, Trans. Md. Acad. Sci. 180, 1892.

Hymenodectes Uhler, *ibid*, 179.

culicis Uhler, Trans. Md. Acad. Sci. 180, 1892 (*Hymenodectes*); Ashm., Proc. Ent. Soc. Wash. ii, 329, 1892 (*Schwarzi* MSS.); Johannsen, Psyche, 1909, 3.—U. S.

formicina Uhler, Trans. Md. Acad. Sci. 1892, 181 (*Hymenocoris*); Johannsen, Psyche, 1909, 4,—Calif.

NABIDÆ.**CARTHASIS.**

Champion, Biol. C.-Am. Rhynch. ii, 305, 1900.

Orthometrops Uhler, Proc. Ent. Soc. Wash. iv, 508, 1901.

decorata Uhler, Proc. Ent. Soc. Wash. iv, 509, 1901.—Md., D, C.

contrarius Reut., Mém. Soc. Ent. Belg. XV, 97, 1908.

METATROPIPHORUS.

Reuter, *Œfv. Kgl. Vet. Akad. Förh.* 1872, 93.

belfragii Reut., *Œfv. Kgl. Vet. Akad. Förh.*, 1872, 94.—Tex.

PAGASA.

Stål, *Rio Hem.* ii, 60, 1862.

fusca Stein, *Berl. Ent. Zeit.* i, 90, 1857 (*Prostemma*); Reuter, *Rev.*

d'Ent. ix, 291, 1890; *Bull.* 69 *Minn. Sta.*, fig. 31.—N. States.

nitida Stål, *Enum. Hem.* iii, 108, 1873.—N. States.

pallipes Stål, *Enum. Hemipt.* iii, 108, 1873.—S. States.

PHORTICUS.

Stål, *Rio Hem.* i, 69, 1858.

collaris Stål, *Enum. Hem.* iii, 109, 1873.—Tex.

REDUVIOLUS.

Kirby, *Fauna Bor. Amer.* iv, 279, 1837.

Nabicula Kirby, *ibid.*

Nabis, authors, not of Latr.

Coriscus, authors, not of Schrank.

Hoplistoscelis, Reuter, *Rev. d'Ent.* ix, 295, 1890.

annulatus Reut., *Œfv. Kgl. Vet. Akad. Förh.* 1872, 86.—E. States.

capsiformis Germ., *Silberm. Rev.* v, 132, 1837 (*Nabis*); Reut., *Œfv.*

Vet. Akad. Förh., 1872 (6), 88; *Mém. Soc. Ent. Belg.* xv,

114, 1908.—Car., Tex.

chenkeanus Kirkaldy, *Wien. Ent. Zeit.* 1901, 224.—N. Car.

elongata Hart, *Bull. Ill. State Lab. Nat. Hist.* vii, 262, 1907.—Ill.

ferus Linn., *Faun. Suec.* 256, 1761 (*Cimex*); Fieber, *Eur. Hem.* 161,

1861; Stål, *Enum. Hemipt.* iii, 113, 1873.—E. States.

heidemanni Reut., *Mém. Soc. Ent. Belg.* xv, 100, 1908.—Calif.

inscriptus Kirby, *Fauna Bor. Amer.* iv, 280, 1837; *Can Entom.* 1878,

213.—N. States, Can.

boreellus Reut., *Medd. Soc. Fauna Flora Fenn.* xi, 165, 1884

(*Nabis*).

kalmii Reut., *Œfv. Kgl. Vet. Akad. Förh.* 1872, 91 (*Nabis*).—Wisc.

limbatus Dahlb., *Vet. Akad. Handl.* 1850, 227; *Flor. Rhyn. Livl.* i,

695, 1860; Reut., *Œfv. Vet. Akad. Förh.* 1872 (6), 70.—

Can., Colo.

nigriventris Stål, *Stett. Ent. Zeit.* 1862, 458 (*Nabis*).—Tex.

propinquus Reut., *Œfv. Kgl. Vet. Akad. Förh.* 1872, 87.—E. States.

vicarius Reut., *ibid.*

roseipennis Reut., *Œfv. Kgl. Vet. Akad. Förh.* 1872, 89; *Rev. d'Ent.*

1890, 308; *Mém. Soc. Ent. Belg.* xv, 118, 1908.—E. States.

punctipes Reut., *Œfv. Kgl. Vet. Akad. Förh.* 1872, 89.

- **rufusculiss** Reut., Cefv. Kgl. Vet. Akad. Förh. 1872, 92.—E. States.
assimilis Uhler, Proc. Bost. Soc. Nat. Hist. xix, 422, 1878
(*Coriscus*).
- sericans** Reut., Cefv. Kgl. Vet. Akad. Förh. 1872, 83 (*Nabis*).—Tex.
- **sordidus** Reut., Cefv. Kgl. Vet. Akad. Förh. 1872, 85; Rev. d'Ent. ix,
299, 1890; Champ., Biol. C.-Am. Rhynch. ii, 303, 1899.—E.
States.
pallescens Reut., Cefv. Kgl. Vet. Akad. Förh. 1872, 85.
- subcoleoptratus** Kirby, Fauna Bor.-Amer. iv. 282, 1837; Can Entom.
1878, 214; Reuter, Cefv. Kgl. Vet. Akad. Förh. 1872, 81;
Stål, Enum. Hem. iii, 112, 1873; Kirkaldy, Wien. Ent. Zeit.
1901, 222; Bull. 22 Div. Entom. Dept. Agric., fig. 20.—E.
States.
canadensis Prov., Nat. Can. i, 1869, 211.
- vanduzeei** Kirkaldy, Wien. Ent. Zeit. 1901, 223.—Colo.
var. of *R. flavomarginatus* Scholtz, Reuter, Mém. Soc. Ent.
Belg. xv, 111, 1908.

ANTHOCORIDÆ.

ANTHOCORIS.

Fallen, Hem. Suec. 65, 1829.

Rhynarius Hahn, Wanz. Ins. i, 104, 1831.

- **antevolens** White, Ent. Mo. Mag. xvi, 146, 1879; Reut., Monog.
Anthoc. 77, 1884.—Cal., Missouri.
- melanocerus** Reuter, Monog. Anthoc. 81, 1884.—Colo.
- musculus** Say, Het. N. Harm. 32, 1832 (*Reduvius*); Compl. Writ. i,
357, 1859.—U. S.
- nigripes** Reut., Monog. Anthoc. 69, 1884.—S. States.
- fulvipennis** Reut., Monog. Anthoc. 69, 1884.—S. States.
- whitei** Reuter, Monog. Anthoc. 74, 1884.—Calif.

ASTHENIDEA.

Reuter, Monog. Anthoc. 48, 1884.

- temnostethoides** Reuter, Monog. Anthoc. 51, 1884.—U. S.

CARDIASTETHUS.

Fieber, Ent. Monatsch. iv, 266, 1860.

- luridellus** Fieber, Wien. Ent. Monatschr. iv, 271, 1860; Reut. Monog.
Anthoc. 142, 1884.—Pa.
- pergandei** Reuter, Monog. Anthoc. 141, 1884.—U. S.

DASYPTERUS.

Reuter, Cefv. Kgl. Vet. Akad. Förh. 1871, 564.

- assimilis** Reut., Cefv. Kgl. Vet. Akad. Förh. 1871, 564, Monog. Anthoc.
139, 1884; Champ., Biol. C.-Am. Rhynch. Het. ii, 332, 1900.
Tex., S. Car.

DILASIA.

Reuter, *Œfv. Kgl. Vet. Akad. Förh.* 1871, 563.

fuscula Reut., *Œfv. Kgl. Vet. Akad. Förh.* 1871, 563; *Monog. Anth.* 22, 1884.—*Tex., S. Car.*

DOLICHOMERUS.

Reuter, *Œfv. Kgl. Vet. Akad. Förh.* 1871, 557.

stälîi Reut., *Œfv. Kgl. Vet. Akad. Förh.* 1871, 558; *Monog. Anthoc.* 10, 1884.—*Tex., S. Car.*

elongatus Reut., *Œfv. Kgl. Vet. Akad. Förh.* 1871, 558; *Monog. Anthoc.* 11, 1884.—*S. Car.*

LASIOCHILUS.

Reuter, *Œfv. Kgl. Vet. Akad. Förh.* 1871, 562.

pallidulus Reut., *Œfv. Kgl. Vet. Akad. Förh.* 1871, 562; *Monog. Anthoc.* 17, 1884; *Champ., Biol. C.-Am. Rhynch. Het.* ii, 309, 1900.—*S. Car.*

LYCTOCORIS.

Hahn, *Wanz. Ins.* iii, 19, 1835.

fitchii Reut., *Œfv. Kgl. Vet. Akad. Förh.* 1871, 557; *Monog. Anthoc.* 7, 1884 (sub *campestris*); *Bull.* 69, *Minn. Sta.*, fig. 46.—*N. Y.*

PIEZOSTETHUS.

Fieber, *Ent. Monatschr.* iv, 265, 1860.

californicus Reut., *Monog. Anthoc.* 46, 1884; *Bull.* 69, *Minn. Sta.*, fig. 45.—*Cal.*

galactinus Fieber, *Weit. Beitr.* 107, 1836; *Europ. Hem.* 139, 1861; Reut., *Monog. Anthoc.* 36, 1884; *Champ., Biol. C.-Am. Rhynch. Het.* ii, 315, 1900.—*S. States.*

albipennis Her.-Schr., *Wanz. Ins.* ix, 231, 1853 (*Xylocoris*).

sordidus Reut., *Œfv. Kgl. Vet. Akad. Förh.* 1871, 560; *Monog. Anthoc.* 37, 1884; *Champ., Biol. C.-Am. Rhynch. Het.* ii, 316, 1900.—*Tex., S. Car.*

binotatus Reut., *Œfv. Kgl. Vet. Akad. Förh.* 1871, 560.

SCOLOPOSCELIS.

Fieber, *Ent. Monatschr.* viii, 66, 1864.

flavicornis Reut., *Œfv. Kgl. Vet. Akad. Förh.* 1871, 561; *Monog. Anth.* 154, 1884; *Champ., Biol. C.-Am. Rhynch. Het.* ii, 335, pl. xix, fig. 32, 1900.—*Tex.*

TETRAPHLEPS.

Fieber, *Ent. Monatschr.* iv, 266, 1860.

canadensis Prov., *Faun. Canad. Hem.* 90, 1886.—*Can.*

TRIPHLEPS.

Fieber, Ent. Monatschr. iv, 266, 1860.

insidiosus Say, Het. N. Harm. 32, 1832 (*Reduvius*); Compl. Writ. i, 357, 1859; Reuter, Monog. Anth. 90, 1884; Bull. 50 Bur. Entom., fig. 13.—U. S.

pseudo-chinche Fitch, Second Rept. Ins. New York, 295, 1856 (*Anthocoris*).

latulus Reut., Cefv. Kgl. Vet. Akad. Förh. 1871, 565.

rugicollis Reuter, *ibid*, 565.

tristicolor White, Ent. Mo. Mag. xvi, 145, 1879; Reut., Monog. Anth. 98, 1884; Champ., Biol. C.-Am. Rhynch. Het. ii, 327, 1900. Cal., Tex.

CIMICIDÆ.

CIMEX.

Linné, Syst. Nat. x, 441, 1758.

Acanthia Fabr., Syst. Ent. 693, 1775.

Clinocoris Fallen, Hem. Suec. 141, 1829.

Klinophilos Kirk., Entom. 1899, 219.

hirundinis Jenyns, Ann. Nat. Hist. iii, 242, 1839; Osborn, Bull. No. 5 (n. s.) Div. Ent. U. S. Dept. Agric. 161, fig. 91, 1896; Lugger, Bull. 69 Minn. Agric. Exp. Station 51, fig. 42, 1900.—U. S.

lectularia Linn., Fauna Suec. 909, 1761 (*Cimex*); De Geer, Mém. Ins. iii, 195, 1775 (*Cimex*); Am. et Serv., Hémipt. 311, 1843; Fieb., Eur. Hem. 135, 1861; Insect Life ii, fig. 16; Bull. 4, Div. Entom., fig 10; Cir. 47, Bur. Entom. Dept. Agric.—U. S.

pipistrelli Jenyns, Ann. Nat. Hist. iii, 243, 1839.—U. S.

HÆMATOSIPHON.

Champion, Biol. C.-Am. Rhynch. Het. ii, 337, 1900.

inodora Dugès, La Naturelleza (2) ii, 169, 1892 (*Acanthia*); Townsend, Proc. Ent. Soc. Wash. iii, 40, 1894; Champ., Biol. C.-Am. Rhynch. Het. ii, 337, 1900, pl. xx, fig. 1; Osborn, Bull. No. 5 (n. s.) Div. Ent. U. S. Dept. Agric. 160, fig. 90, 1896.—S. W. States.

CERATOCOMBIDÆ.

CERATOCOMBUS.

Signoret, Ann. Soc. Ent. France, 1852, 542.

Lichenobia Bærensp., Berl. Ent. Zeitschr. i, 165, 1857.

brasiliensis Reut., Monog. Ceratocomb. 1871, 7; Uhler, Proc. U. S. Nat. Mus. xxvii, 361, 1904.—N. Mex.

latipennis Uhler, Proc. U. S. Nat. Mus. xxvii, 362, 1904.—N. Mex.

niger Uhler, Proc. U. S. Nat. Mus. xxvii, 361, 1904.—N. Mex.

GLYPTOCOMBUS.

- Heidemann, Proc. Ent. Soc. Wash. vii, 192, 1905 (*Hypselosoma* acc. to Horvath).
saltator Heid., Proc. Ent. Soc. Wash. vii, 194, 1905; Horvath, Ann. Mus. Nat. Hung. vi, 565, 1908.—Md.

GERRIDÆ.**GERRIS.**

- Fabricius, Ent. Syst. iv, 188, 1794.
Limnotrechus Stål, Cefv. Vet. Akad. Förh. xxv, 395, 1868.
Hydrotrechus Stål, *ibid.*
canaliculatus Say, Hem. New Harm. 36, 1832; Compl. Writ. i, 363, 1859.—U. S.
conformis Uhler, Proc. Bost. Soc. Nat. Hist. xix, 435, 1878 (*Hygrotrechus*).—E. States.
gillettei Leth. and Severin., Cat. Hemipt. iii, 60, 1896.—Colo.
producta Uhler, Hem. Colo. 61, 1895 (*Limnotrechus*) (preoc.).
marginatus Say, Het. Hem. N. Harm. 36, 1832; Compl. Writ. i, 362, 1859.—U. S.
orba Stål, Eugenie's Résa. Ins. 264, 1866.—Calif.
remigis Say, Het. N. Harm. 35, 1832; Compl. Writ. i, 362, 1859; Uhler, Proc. Bost. Soc. N. H. xix, 435, 1878; Bull. 69 Minn. Sta., fig. 21.—U. S.
robusta Uhler, Stillman's Amer. Journ. Sci. 1871, 105.—W. States.
aptera Walk., Cat. Hem. viii, 165, 1873.

HALOBATES.

- Eschscholtz, Nat. Abh. Dorpat. i, 163, 1823.
sericeus Eschscholtz, Nat. Abh. Dorpat. i, 163, 1823.—Calif.
wuellerstorffi Frauenfeld, Verh. zool-bot. Ges. Wien. 1867, 458; White, Rept. Chall. Voy. Zool. vii, 40, 1883.—Fla.

HALOBATOPSIS.

- Bianchi, Ann. Mus. Zool. Acad. Imp. Sci. St. Petersburg. 1896, 70.
beginii Ashmead, Can. Entom. 1897, 56.—Can.

LIMNOGONUS.

- Stål, Hem. Fabric. i, 132, 1868.
franciscana Stål, Eugenie's Resa. Ins. 265, 1866 (*Gerris*).—Calif.
hesione Kirkaldy, Entomologist, 1902, 137 (*Gerris*).—Fla.

LIMNOPORUS.

- Stål, Cefv. Vet. Akad. Förh. xxv, 385, 1868.
rufoscutellatus Latr., Gen. Ins. iii, 134, 1807 (*Gerris*); Stål, Cefv. Vet. Akad. Förh. 1868, 396.—U. S.

METROBATES.

Uhler, Proc. Bost. Soc. Nat. Hist. xiv, 108, 1871.

hesperius Uhler, Proc. Bost. Soc. Nat. Hist. xiv, 109, 1871; xix, 438, 1878.—E. States.

RHEUMATOBATES.

Bergroth, Insect Life, iv, 321, 1892.

rileyi Bergr., Ins. Life, iv, 321, 1892; Insect Life, v, figs. 18, 19, 20.—E. States.

tenuipes Meinert, Ent. Medd. v, 17, 1895.—E. States.

TREPOBATES.

Uhler, Proc. Zool. Soc. Lond. 1894, 213.

Stephania White, Rept. Chall. Voy. Zool. vii, 79, 1883 (preoc.).

picta Her.-Schfr., Wanz. Ins. viii, 111, 1848 (*Halobates*).—E. States.

VELIIDÆ.**MACOVELIA.**

Uhler, Hayden's Geol. Surv. Mont. 422, 1872.

horni Uhler, Hayden's Surv. Montana, 422, 1872.—W. States.

MESOVELIA.

Mulsant et Rey, Ann. Soc. Linn. Lyon, 1852, 138.

mulsanti Buch. White, Trans. Ent. Soc. Lond. 1879, 268; Champion, Biol. C.-Am. Rhynch. ii, 123, 1898.—E. States.

bisignata Uhler, Stand. Nat. Hist. ii, 273, fig. 324, 1884.

MICROVELIA.

Westwood, Ann. Soc. Ent. France, 1834, 647.

americana Uhler, Hem. Colo. 61, 1895.—U. S.

capitata Guérin, Sagra's Hist. Nat. Cuba, vii, Ins. 174, 1857.—E. States.

marginata Uhler, Proc. Zool. Soc. Lond. 1893, 719.—U. S.

pulchella Westwood, Ann. Soc. Ent. France 1834, 647.—N. Y., N. J.

RHAGOVELIA.

Mayr, Verh. zool-bot. Ges. Wien. 1865, 445.

Neovelvia White, Journ. Linn. Soc. Lond. Zool. xiv, 487, 1879.

Trochopus Carpenter, Ent. Mo. Mag. 1898, 78.

armata Burm., Handb. ii, 212, 1835 (*Velia*); Champ., Biol. Cent.-Amer. Rhynch. Het. ii, pl. viii, fig. 24, 1898.—Tex.

collaris Burm., Handb. ii, 212, 1835 (*Velia*); Mayr, Novara Reise, Hemipt. 180, 1866.—U. S.

distincta Champion, Biol. Centr.-Amer. Rhynch. Het. ii, 135, 1898.—U. S.

distincta Walk., Cat. Hem, viii, 161, 1873 (*Velia*) (sine descript).

- obesa* Uhler, Proc. Bost. Soc. Nat. Hist. xiv, 107, 1871; xix, 434, 1878.
—E. States.
plumbea Uhler, Proc. Zool. Soc. Lond. 1894, 217.—Fla.
marinus Carpenter, Ent. Mo. Mag. 1898, 78 (*Trochopus*).

NÆOGEUS.

- Laporte, Essai, 32, 1832.
Hebrus Curtis, Ent. Mag. i, 198, 1833.
burmeisteri Lethierry et Severin, Cat. Hem. iii, 51, 1896.—U. S.
pusillus Burm., Handb. ii, 214, 1835 (nec Fallen).
concinus Uhler, Proc. Zool. Soc. London, 1894, 221.—E. States.
sobrinus Uhler, Rept. Chief Engineers, Geol. Surv. Appendix N. N.,
1330, 1877; Bull. U. S. Geol. and Geog. Surv. Terr. iii (2),
452, 1877 (*Hebrus*).—N. Mex.

HYDROMETRIDÆ.

HYDROMETRA.

- Latreille, Hist. Crust. Ins. iii, 250, 1802.
Limnobates Burm., Handb. ii (1) 210, 1835.
australis Say, Het. Hem. N. Harm. 35, 1832; Compl. Writ. i, 361,
1859; Bueno, Can. Entom. 1905, 14, 264.—Ga., Fla.
martini Kirkaldy, Entomol. 1900, 176.—U. S.
lineata Say, Het. Hem. N. Harm. 35, 1832 (not of Eschscholtz);
Compl. Writ. i, 361, 1859; Martin, Can. Entom. 1900, 70.

ISOMETOPIDÆ.

ISOMETOPUS.

- Fieber, Wien. Ent. Monatsschr. iv, 259, 1860.
Cephalocoris Stein, Berl. Ent. Zeit. iv, 79, 1860.
pulchellus Heid., Proc. Ent. Soc. Wash. ix, 128, 1908.—E. States.
signatus Heid., Proc. Ent. Soc. Wash. ix, 129, 1908.—Tex.
unicolor Heid., Proc. Ent. Soc. Wash. ix, 130, 1908.—Ariz.

HEIDEMANNIA.

- Uhler, Proc. Ent. Soc. Wash. ii, 119, 1891.
cixiiformis Uhler, Proc. Ent. Soc. Wash. ii, 121, 1891; Heid., Proc.
Ent. Soc. Wash. ix, 127, 1908.—D. C., Va.

CAPSIDÆ.

Plagiognathini.

AGALLIASTES.

- Fieber, Wien. Ent. Monatsschr. ii, 321, 1858.
apiatus Uhler, Hem. Colo. 53, 1895.—Colo., Kans.
bakeri Berg., Wien. Ent. Zeit. 1898, 35.—Colo.
signatus Uhler, Hem. Colo. 55, 1895 (preoc.).

- fumidus** Uhler, Hem. Colo. 54, 1895.—Colo.
pulicarius Fall., Mon. Cim. 95, 1807 (*Phytocoris*); Fieber, Eur. Hem. 312, 1861; Reut., Hem. Gym. Eur. i, 60, 1878.—Can.
uniformis Uhler, Proc. Ent. Soc. Wash. ii, 379, 1893.—Utah.

ATOMOSCELIS.

- Reuter, Bih. Vet. Akad. Handl. iii (1), 57, 1875.
pilosulus Uhler, Proc. Ent. Soc. Wash. ii, 377, 1893.—Utah.
seriatus Reut., Öfv. Vet. Akad. Förh. 1875 (9), 91.—Tex.

ATRACTOTOMUS.

- Fieber, Wien. Ent. Monatssch. ii, 317, 1858.
albidicoxis Reut., Act. Soc. Sci. Fenn. xxxvi (2), 79, 1908.—Ariz.
hesperius Reut., Act. Soc. Sci. Fenn. xxxvi (2), 78, 1909.—Calif.

CAMPYLOMMA.

- Reuter, Act. Soc. Sci. Fenn. xiii (1), 52, 1878.
verbasci Meyer, Rhyn. Schw. 70, 1843 (*Capsus*); Fieber, Eur. Hem. 311, 1861; Reut., Hem. Gym. Eur. i, 53, 1878.—N. Y.

CHLAMYDATUS.

- Curtis, Ent. Mag. i, 19, 1833.
Agalliastes Fieber, Wien. Ent. Monatsschr. ii, 321, 1858 (part).
associatus Uhler, Hayden Surv. Mont. 419, 1872 (*Agalliastus*); Reut., Act. Soc. Sci. Fenn. xxxvi (2) 82, 1909; 21 Rept. Ill. Entom. p. 88, pl. iii, fig. 3.—Pa., Colo.
luctuosus Prov., Pet. Faun Can, Hem. 137, 1887.—Can.
obliquus Uhler, Proc. Ent. Soc. Wash. ii, 378, 1893 (*Agalliastes*).—Utah.
suavis Reut., Öfv. Vet. Akad. Förh. 1875 (9), 92 (*Agalliastes*); Reut., Act. Soc. Sci. Fenn. xxxvi (2), 83, 1909.—Tex.

EUROPIELLA.

- Reuter, Act. Soc. Sci. Fenn. xxxvi (2), 83, 1909.
concinna Reut., Act. Soc. Sci. Fenn. xxxvi (2), 84, 1909.—Calif.
rubidus Uhler, Hem. Colo. 41, 1895 (*Stenarus*); Reut., Act. Soc. Sci. Fenn. xxxvi (2), 85, 1909.—U. S.
stigmaticus Uhler, Proc. Ent. Soc. Wash. ii, 379, 1893 (*Agalliastes*); Reut., Act. Soc. Sci. Fenn. xxxvi (2) 84, 1909.—W. States.
umbrina Reut., Act. Soc. Sci. Fenn. xxxvi (2), 85, 1909.—Calif.

LAODAMINA.

- Laodamina* Kirkaldy, Wien. Ent. Zeit. 1903, 13 (preoc. Ragonot, 1888).
Strongylotes Reut., Öfv. Vet. Akad. Förh. 1875, 88 (preoc.).
saliens Reut., Öfv. Vet. Akad. Förh. 1875 (9), 88; Act. Soc. Sci. Fenn. xxxvi (2) 79, 1909 (*Strongylotes*).—Tex.

LAMPETHUSA.

Distant, Biol. C.-Am. Rhynch. Het. i, 303, 1884.

anatina Dist., Biol. Cent.-Amer. Rhynch. Het., i, 303, 451, pl. xxix, fig. 13, 1884.—Tex.

LEUCOPŒCILA.

Reuter, Œfv. Finska Vet. Soc. Förh. xlix. (5), 24, 1907.

albofasciata Reut., Œfv. Finska Vet. Soc. Förh. xlix (5), 26, 1907 ;
Act. Soc. Sci. Fenn. xxxvi (2), 83, 1909.—Pa.

MAURODACTYLUS.

Reuter, Act. Soc. Sci. Fenn. xiii (1), 27, 1878.

angulatus Uhler, Hem. Colo. 53, 1895,—Colo.

consors Uhler, Hem. Colo. 53. 1895.—Colo.

MICROPHYLELLUS.

Reuter, Act. Soc. Sci. Fenn. xxxvi (2), 76, 1909.

Type *Phylus modestus* Uhler, which is not described.

MYOCHROOCORIS.

Reut., Act. Soc. Sci. Fenn. xxxvi (2), 76, 1909.

griseolus Reut., Act. Soc. Sci. Fenn. xxxvi (2), 77, 1909.—Tex.

NEOCORIS.

Douglas and Scott, Brit. Hem. 424, 1865.

bohemani Fall., Hem. Suec. 106, 1829 (*Phytocoris*) ; Fieber, Eur. Hem. 303, 1861 ; Reut., Hem. Gym. Eur. i, 57, 1878.—Can., N. Y.

PHENICOCORIS.

Reuter, Bih. Vet. Akad. Hand. iii (1), 55, 1875.

pubescens Reut., Œfv. Vet. Akad. Förh. 1875 (9), 90.—Tex.

PLAGIOGNATHUS.

Fieber, Wien. Ent. Monatsschr. ii, 320, 1858.

annulatus Uhler, Hem. Colo. 51, 1895.—U. S.

confusus Reut., Act. Soc. Sci. Fenn. xxxvi (2), 80, 1909.—Calif., Nev.

decolor Uhler, Proc. Ent. Soc. Wash. ii, 380, 1893 (*Agalliastes*) ;
Reut., Act. Soc. Sci. Fenn. xxxvi (2), 81, 1909.—W. States,

fraternus Uhler, Hem. Colo. 51, 1895.—Colo.

fuscus Prov., Nat. Can. iv, 105, 1872 (*Lygus*) ; Pet. Faun. Can. Hem 153, 1887.—U. S.

obscurus Uhler, Hayden Surv. Mont. 418, 1872 ; Proc. Bost. Soc. Nat. Hist. xix, 409, 1878 ; 23 Rept. Ill. Entom. p. 201, 1905.

grandis Reut., Œfv. Vet. Akad. Förh. 1875 (9), 91,—Tex.

mœrens Reut., Act. Soc. Sci. Fenn. xxxvi (2), 80, 1909.—Calif., Wyo.

politus Uhler, Hem. Colo. 52, 1895.—Colo., N. Y.

rubricans Prov., Pet. Faun. Can. Hem. 154, 1887.—Can.

PSALLUS.

Fieber, Wien. Ent. Monatsschr. ii, 321, 1858.

Apocremnus Fieber, *ibid.*

ancorifer Fieber, Eur. Hem. 304, 1861; Reut., Hem. Gym. Eur. i, 104, 1878; Act. Soc. Sci. Fenn. xxxvi (2), 77, 1909.—D. C.

breviceps Reut., Act. Soc. Sci. Fenn. xxxvi (2), 78, 1909.—Tex.

delicatus Uhler, Entom. Amer. iii, 34, 1888.—Ga.

guttulosus Reut., Cefv. Vet. Akad. Förh. 1875 (9), 89.—Tex.

juniperi Heideman, Jour. N. Y. Ent. Soc. 1905, 49.—N. Y.

sulphureus Reut., Cefv. Finsk. Vet. Soc. Förh. xlix (5), 23, 1907.—Fla.

variabilis Fall., Hem. Suec. 98, 1829; Reut., Hem. Gym. Eur. i, 116, 1878.—Can.

REUTEROSCOPIUS.

Kirkaldy, Wien. Ent. Zeit. 1905, 268.

Episcopus Reuter, Cefv. Vet. Akad. Förh., 1875, 90 (preoc.).

Aristoreuteria Kirk., Trans. Amer. Ent. Soc. xxxii, 124, 1906.

ornatus Reut., Cefv. Vetensk. Akad. Förh. 1875, 90 (*Episcopus*).—U. S.

avidus Dist., Biol. Cent.-Amer. Rhynch. Het. i, 433, 1886 (*Lygus*).

RHINACLOA.

Reuter, Cefv. Vet. Akad. Förh. 1875, 88.

forticornis Reut., Cefv. Vet. Akad. Förh. 1875 (9), 89; Act. Soc. Sci. Fenn. xxxvi (2), 82, 1909.—Tex.

TUPONIA.

Reuter, Bih. Vet. Akad. Handl. iii, 53, 1875.

subnitida Uhler, Hem. Colo. 45, 1895.—Colo.

Oncotylini.

AMBLTYLUS.

Fieber, Wien. Ent. Monatsschr. ii, 325, 1858.

sexguttatus Prov., Pet. Faun. Can. Hem. 150, 1887.—Can.

CYLLOCEPS.

Uhler, Proc. Zool. Soc. London, 1893, 711.

pellicia Uhler, Proc. Zool. Soc. London, 1893, 712.—S. Fla.

HOPLOMACHIDEA.

Reuter, Act. Soc. Sci. Fenn. xxxvi (2), 75, 1909.

consors Uhler, N. Amer. Fauna, 1893, 264 (*Hopломachus*).—Calif.

LOPUS.

Hahn, Wanz. Ins. i, 10, 1831.

Onychumenus Reuter, Act. Soc. Sci. Fenn. xiii (2), 286, 1879.

decolor Fall., Hem. Suec. 123, 1829 (*Capsus*); Fieber, Eur. Hem. 298, 1861; Reut., Gym. Eur. ii, 287, 1879.—Can.
chrysanthemi Hahn., Wanz. Ins. i, 10, 1831.

MACROTYLUS.

Fieber, Wien. Ent. Monatsschr. ii, 325, 1858.

Mermimerus Sahlb., Not. Faun. Fenn. xi, 293, 1870.

affiguratus Uhler, Hem. Colo. 50, 1895.—Colo.

amœnus Reut., Act. Soc. Sci. Fenn. xxxvi. (2), 75, 1909.—Conn.

angularis Uhler, Proc. Calif. Acad. Sci. (2) iv, 272, 1894.—Calif.

lineolatus Uhler, Proc. Calif. Acad. Sci. (2) iv, 270, 1894.—Calif.

regalis Uhler, Trans. Md. Acad. Sci. 1890, 86.—Calif.

tristis Uhler, Trans. Md. Acad. Sci. 1890, 87.—Calif.

verticalis Uhler, Proc. Calif. Acad. Sci. (2) iv, 272, 1894.—Calif.

vestitus Uhler, Trans. Md. Acad. Sci. 1890, 88.—Calif.

MEGALOCOLEUS.

Reuter, Rev. d'Entom. 1890, 254.

Macrocoleus Fieber, Wien. Ent. Monatsschr. ii, 325, 1858 (preoc.).

coagulatus Uhler, Bull. U. S. Geol. Surv. iii, 417, 1877 (*Macrocoleus*).
—Colo.

ONCOTYLUS.

Fieber, Wien. Ent. Monatsschr. ii, 318, 1858.

Anoterops Fieber, Eur. Hem, 72. 1861.

Cylindromelus Fieber, ibid. 383, 1861.

longipennis Uhler, Hem. Colo. 48, 1895.—Colo.

puberus Uhler, Proc. Calif. Acad. Sci. (2) iv, 270, 1894.—Calif.

repertus Uhler, Hem. Colo., 49, 1895.—Colo., Calif.

unicolor Prov. Nat. Can. ix, 105, 1872 (*Lygus*).—Can.

TINICEPHALUS.

Fieber, Wien. Ent. Monatsschr. ii, 318, 1858.

simplex Uhler, Hayden Surv. Mont. 417, 1872.—Colo.

Cyllocorini.

CERATOCAPSUS.

Reuter, Öfv. Vet. Akad. Förh. 1875, 87.

Melinna Uhler, Ent. Amer. iii, 69, 1888.

Hyperoides Kirk., Wien. Ent. Zeit. 1903, 14.

elongata Uhler, Proc. Calif. Acad. Sci. (2) iv, 257, 1894 (*Melinna*).—
Tex., Ariz., Calif.

fasciatum Uhler, Bull. U. S. Geol. Surv. Terr. viii, 421, 1877 (*Mega-*
cœlum).—Tex., Mass.

- lutescens** Reut., Cefv. Vet. Akad. Förh. 1875 (9), 87.—Tex.
 * **modesta** Uhler, Entom. Amer. iii, 69, 1888 (*Melinna*).—E. States.
pumila Uhler, Entom. Amer. iii, 69, 1888 (*Melinna*).—E. States.
setosus Reut., Act. Soc. Sci. Fenn. xxxvi (2) 70, 1909.—D. C., Pa.
pteridis (*Bryocoris*) (Auct. Amer. nec Fallen).

CYLLOCORIS.

- Hahn., Wanzen. Ins. ii, 97, 1834.
Kelidocoris Kol., Bull. Soc. Nat. Moscou, 1845, 101.
histrionicus Linn., Syst. Nat. (12) 728, 1767 (*Cimex*); Fieber, Eur. Hem. 282, 1861; Reut., Hem. Gym. Eur. iii, 406, 1883.—E. States.
agilis Fabr., Spec. Ins. 374, 1781.

CYRTORRHINUS.

- Fieber, Wien. Ent. Monatsschr. ii, 313, 1858.
Tytthus Fieber, ibid. viii, 82, 1864.
Breddiniessa Kirk., Wien. Ent. Zeit. xxii, 13, 1903.
marginatus Uhler, Hem. Colo. 43, 1895; Reut., Act. Soc. Sci. Fenn. xxxvi (2) 67, 1909.—Can., Colo.

DIAPHNIDIA.

- Uhler, Hem. Colo. 43, 1895.
debilis Uhler, Hem. Colo. 43, 1895.—Colo.
pellucida Uhler, Hem. Colo. 44, 1895.—Colo., D. C., Can.

GLOBICEPS.

- Lep. Serv., Ency. Méth. x, 326, 1825.
angustata Uhler, Hem. Colo. 42, 1895.—Colo.
flavomaculatus Fabr., Entom. Syst. iv, 182, 1794 (*Lygæus*); Reut., Hem. Gym. Eur. iii, 395, 1883.—Can.
selectus Fieber, Eur. Hem. 284, 1861.

HETEROCORDYLUS.

- Fieber, Wien. Ent. Monatsschr. ii, 316, 1858.
malinus Reut., Act. Soc. Sci. Fenn. xxxvi (2), 71, 1909.—N. Y., Ill.

ILNACORA.

- Reuter, Cefv. Vet. Akad. Förh. 1875, 85.
chloris Uhler, Bull. U. S. Geol. Surv. iii, 418, 1877 (*Stenarops*).—Colo.
divisa Reut., Cefv. Kgl. Vet. Akad. Förh. 1875 (9), 86.—Tex.
 * **malina** Uhler, Bull. U. S. Geol. Surv. iii, 419, 1877 (*Stenarops*).—E. States.
 * **stalii** Reut., Cefv. Kgl. Vet. Akad. Förh. 1875 (9), 86.—Tex., N. Y.
viridis Uhler, Hem. Colo., 41, 1895.—Colo.

MALACOCORIS.

Fieber, Wien. Ent. Monatssch. ii, 318, 1858.

provancheri Burque, Pet. Faun. Can. Hem. 144, 1887.—Can.

MECOMMA.

Fieber, Wien. Ent. Monatsschr. ii, 313, 1858.

Sphyracephalus Douglas and Scott, Brit. Hem. 349, 1865.

† **ambulans** Fall., Hem. Suec. 126, 1829 (*Cimex*); Reut., Hem. Gym. Eur. iii, 384, 1883.—N. States.

ORTHOTYLUS.

Fieber, Wien. Ent. Monatssch. ii, 314, 1858.

Diommatus Uhler, Entom. Amer. iii, 32, 1888.

angulatus Uhler, Hem. Colo. 44, 1895 (*Diommatus*).—Colo.

biguttatus Reut., Act. Soc. Sci. Fenn. xxxvi (2), 69, 1909.—Colo.

† **chlorionis** Say, Het. New Harm. 24, 1832; Compl. Writ. i, 346, 1859 (*Capsus*); 21 Rept. Ill. Entom., p. 90, pl. iv, fig. 1.—E. States.

congrex Uhler, Entom. Amer. iii, 33, 1888 (*Diommatus*); Reut., Act. Soc. Sci. Fenn. xxxvi (2), 67, 1909.—E. States.

dorsalis Prov., Nat. Can. iv, 104, 1872 (*Lygus*); Pet. Faun. Can. Hem. 138, 1887.—Can.

† **flavosparsus** Sahlb., Act. Soc. Sci. Fenn. i, 411, 1842; Reut., ibid xxxvi (2), 68, 1909.—N. States.

inconspicua Uhler, Proc. Ent. Soc. Wash. ii, 376, 1893 (*Asciodema*); Reut., Act. Soc. Sci. Fenn. xxxvi (2), 69, 1909.—Utah.

nigripes Reuter, Act. Soc. Sci. Fenn. xxxvi (2), 68, 1909.—Nev.

sericatus Uhler, Hem. Colo. 49, 1895 (*Oncotylus*).—Colo.

translucens Tucker, Kans. Univ. Sci. Bull. iv, p. 58, 1907.—Colo.

viridicatus Uhler, Hem. Colo. 48, 1895.—Colo., N. Mex.

PARTHENICUS.

Reuter, Öfv. Vet. Akad. Förh. 1875, 84.

psalliodes Reut., Öfv. Vet. Akad. Förh. 1875 (9), 85.—Tex.

PSEUDOXENETUS.

Reuter, Act. Soc. Sci. Fenn. xxxvi (2), 66, 1909.

Xenetus Uhler, Trans. Md. Acad. 1890, 80 (nec Dist.).

regalis Uhler, Trans. Md. Acad. Sci. 1890, 80 (*Xenetus*).—E. States.

scutellatus Uhler, Trans. Md. Acad. Sci. 1890, 81 (*Xenetus*).—E. States.

REUTERIA.

Puton, Pet. Nouv. Entom. i, 519, 1875.

irroratus Say, Het. New Harm. 241, 1832; Compl. Writ. i, 346, 1859 (*Capsus*); Reut. Act. Soc. Sci. Fenn. xxxvi (2), 70, 1909.—E. States.

RHINOCAPSUS.

Uhler, Trans. Md. Acad. Sci. 1890, 81.

vanduzeei Uhler, Trans. Md. Acad. Sci. 1890, 82.—N. States, Can.

TELEORHINUS.

Uhler, Trans. Md. Acad. Sci. 1890, 74.

cyaneus Uhler, Trans. Md. Acad. Sci. 1890, 75.—Calif.

TIRYAS.

Kirk., Wien. Ent. Zeit. 1903, 14.

Trichia Reut., Cefv. Vet. Akad. Förh. 1875 (9), 81 (preoc.).

punctulata Reut., Cefv. Vet. Akad. Förh. 1875 (9), 82 (*Trichia*); Act.

Soc. Sci. Fenn. xxxvi (2), 71, 1909.—Tex.

Cremnocephalini.

CLOSTEROCORIS.

Uhler, Trans. Md. Acad. Sci. 1890, 76.

ornata Uhler, Trans. Md. Acad. Sci. 1890, 77.—Calif.

COQUILLETIA.

Uhler, Trans. Md. Acad. Sci. 1890, 79.

insignis Uhler, Trans. Md. Acad. Sci. 1890, 79.—W. States.

mimetica Osborn, Proc. Iowa Acad. Sci. v, 236, 1898.—Iowa, Fla.

CYRTOPELTOCORIS.

Reuter, Cefv. Vet. Akad. Förh. 1875, 81.

albofasciatus Reut., Cefv. Vet. Akad. Förh. 1875 (9), 81.—Tex.

DACERLA.

Bergroth, Entom. News, 1897, 95.

Myrmecopsis Uhler, Proc. Calif. Acad. Sci. (2) iv, 276, 1894 (preoccupied).

inflatus Uhler, Proc. Calif. Acad. Sci. (2) iv, 276, 1894 (*Myrmecopsis*);

Bergr., Ent. News 1897, 95; Reut., Act. Soc. Sci. Fenn.

xxxvi (2) 8, 1909.—Calif.

ECTOPIOCERUS.

Uhler, Trans. Md. Acad. Sci. 1890, 73.

anthracinus Uhler, Trans. Md. Acad. Sci. 1890, 74.—Calif.

ORECTODERUS.

Uhler, Bull. 5. U. S. Geol. Surv. Terr. ii, 319, 1876.

amoenus Uhler, Bull. U. S. Geol. Surv. iii, 426, 1877.—Colo.

longicollis Uhler, Hem. Colo. 47, 1895.—Colo.

obliquus Uhler, Bull. 5 U. S. Geol. Surv. Terr. ii, 320, 1876; Kirk.,

Trans. Ent. Soc. Lond. 1902, 249, pl. vi, fig. 1, 2, 5.—

W. States.

SERICOPHANES.

Reuter, *Œfv. Vet. Akad. Förh.* 1875, 79.

ocellatus Reut., *Œfv. Vetensk. Akad. Förh.* 1875 (9), 79; Osborn, *Proc. Iowa Acad. Sci.* v, 238; 1898, figs.—Iowa, Tex.

Pilophorini.

ALEPIDIA.

Reuter, *Act. Soc. Sci. Fenn.* xxxvi (2), 75, 1909.

gracilis Uhler, *Hem. Colo.* 42, 1895 (*Pilophorus*).—E. States, Colo.

PAMILLIA.

Uhler, *Entom. Amer.* iii, 31, 1888.

behrensii Uhler, *Entom. Amer.* iii, 31, 1888.—Calif.

PILOPHORUS.

Westwood, *Syn. Brit. Ins.* 121, 1839.

amœnus Uhler, *Entom. Amer.* iii, 30, 1888.—E. States.

cinnamopterus Kirsch., *Rhyn. Wiesb.* 232, 1855 (*Leptomerocoris*); Fieber, *Eur. Hem.* 314, 1861; Reut., *Hem. Gym. Eur.* iv, 113, 1891.—E. States.

bifasciatus Fabr., *Syst. Ent.* 725, 1775 (preoc.).

schwarzi Reut., *Act. Soc. Sci. Fenn.* xxxvi (2), 74, 1909.—Calif.

confusus (*Pilophorus*) Auct. Amer. (nec Kirschb.); Uhler, *Ent. Amer.* iii, 29, 1888.—Calif.

walshii Uhler, *Entom. Amer.* iii, 30, 1888.—Ill.

Fulviini.

FULVIUS.

Stål, *Stett. Ent. Zeit.* 1862, 322.

Pamerocoris Uhler, *Bull. U. S. Geol. Surv.* iii, 425, 1877.

atratus Dist., *Biol. Cent.-Amer. Hem. Het.* i, 282, 1883.—S. W. States.

brunneus Prov., *Nat. Can.* iv, 104, 1872 (*Capsus*); *Pet. Faun. Can.* Hem. iii, 127, 1890; Reuter, *Entom. Tidskr.* 1895, 140.—U. S.

anthocoroides Uhler, *Bull. U. S. Geol. Surv.* iii, 425, 1877 (*Pamerocoris*); *Proc. Bost. Soc. N. H.* xix, 413, 1878.

heidemanni Reut., *Entom. Tidskr.* 1895, 142.—Md., Va.

PERITROPIS.

Uhler, *Proc. Ent. Soc. Wash.* ii, 121, 1891.

saldæformis Uhler, *Proc. Ent. Soc. Wash.* ii, 122, 1891; Reut., *Act. Soc. Sci. Fenn.* xxxvi (2), 66, 1909.—D. C., Ill.

*Dicyphini.***CYRTOPELTIS.**

Fieber, Europ. Hem. 76, 1861.

Engytatus Reut., Cefv. Vet. Akad. Förh. 1875 (9), 82.

Neoproba Dist., Biol. C.-Am. Rhynch. Het. i, 270, 1883.

melanocephalus Reut., Act. Soc. Sci. Fenn. xxxvi (2), 63, 1909.—Tex.

simplex Reut., Act. Soc. Sci. Fenn. xxxvi (2), 63, 1909.—Calif.

varians Dist., Biol. C.-Amer. Rhynch. Het. i, 271, 1883, pl. xxvi, fig. 7 (*Neoproba*); Reut., Act. Soc. Sci. Fenn. xxxvi (2), 62, 1909.—S. States.

geniculatus Reut., Cefv. Vet. Akad. Förh. 1875 (9), 83 (*Engytatus*) (preoc.).

DICYPHUS.

Fieber, Wien. Ent. Monatsschr. ii, 327, 1858.

californicus Stål, Freg. Eugenies Resa, Hem. 259, 1859 (*Capsus*).—Calif.

minimus Uhler, Ent. News 1899, 59; Farm. Bull. 120 Dept. Agric., fig. 12.—S. States.

vestitus Uhler, Hem. Colo. 46, 1895.—N. States.

HYALIODES.

Reuter, Cefv. Vet. Akad. Förh. 1875, 83.

vitripennis Say, Het. New Harm. 24, 1832 (*Capsus*); Compl. Writ. i, 345, 1859; Reut., Act. Soc. Sci. Fenn. xxxvi (2), 61, 1909.—U. S.

var. *discoidalis* Reut., ibid.

IDOLOCORIS.

Douglas and Scott, Brit. Hem. 380, 1865.

agilis Uhler, Bull. U. S. Geol. Surv. iii, 425, 1878.—U. S.

famelicus Uhler, Proc. Bost. Soc. Nat. Hist. xix, 413, 1878.—N. H.

MACROLOPHUS.

Fieber, Wien. Ent. Monatsschr. ii, 326, 1858.

separatus Uhler, Proc. Zool. Soc. London 1894, 194 (*Dicyphus*).—U. S.

*Cylapini.***CYLAPUS.**

Say, Het. New Harm. 25, 1832.

Valdasus Stål, Kgl. Svensk. Vet. Akad. Handl. vii (2), 56, 1860.

tenuicornis Say, Het. New Harm. 25, 1832; Compl. Writ. i, 347, 1859; Heid., Proc. Ent. Soc. Wash. ii, 68, 1891.—E. States.

SYSINAS.

Distant, Biol. C.-Amer. Rhynch. Het. i, 248, 1883.

linearis Dist., Biol. C.-Am. Rhynch. Het. i, 248, 1883, pl. xxiii, fig. 21.—N. Y.

Labopini.

HALTICUS.

Hahn, Wanzen. Ins. i, 113, 1832.

Halticocoris Douglas and Scott, Brit. Hem. 478, 1865.

apterus Linn., Faun. Suec. 894, 1761 (*Cicada*); Reut., Hem. Gym. Eur. iv, 18, 1891.—U. S.

pallicornis Fabr., Ent. Syst. iv, 69, 1794 (*Acanthia*); Fieber, Eur. Hem. 282, 1861.

bractatus Say, Het. New Harm. 25, 1832; Compl. Writ. i, 348, 1859 (*Capsus*).

citri Ashmead, Ent. Amer. iii, 155, 1887 (*Rhinacloa*); Ent. News, 1897, pl. x.—U. S.

? *canus* Dist., Biol. C.-Amer. Rhynch. Het. i, 430, 1884 (*Calacoris*).

minutus Uhler, Ann. Rept. Kans. Exp. Sta. f. 1889, 212 (preoc.).

uhleri Giard, C. R. Soc. Biol. iv, 81, 1892; Reut., Act. Soc. Sci. Fenn. xxxvi (2), 71, 1909; Bull. 33 Bur. Entom. Dept. Agric., fig. 25.

intermedius Uhler, Proc. U. S. Nat. Mus. xxvii, 360, 1904.—Ariz.

HYOIDEA.

Reut., Pet. nouv. Entom. ii, 34, 1876.

Bolteria Uhler, Entom. Amer. iii, 33, 1888.

amicta Uhler, Entom. Amer. iii, 34, 1888 (*Bolteria*).—N. Mex.

grisea Reut., Act. Soc. Sci. Fenn. xxxvi (2), 73, 1909.—Cal., Wyo.

picta Uhler, Proc. Ent. Soc. Wash. ii, 373, 1893 (*Bolteria*); Reut., Act. Soc. Sci. Fenn. xxxvi (2), 72, 1909.—Utah.

LABOPIDEA.

Uhler, Bull. U. S. Geol. Surv. iii, 415, 1877.

chloriza Uhler, Bull. U. S. Geol. Surv. iii, 415, 1877.—Utah.

LABOPS.

Burmeister, Handb. ii, 279, 1835.

hesperius Uhler, Hayden Surv. Mont. 416, 1872.—Colo., Can.

LOMATOPLEURA.

Reuter, Öfv. Vet. Akad. Förh. 1875, 67.

cæsar Reut., Öfv. Vet. Akad. Förh. 1875 (9), 67; Act. Soc. Sci. Fenn. xxxvi (2), 72, 1909.—Pa.

hesperus Kirk, Trans Ent. Soc. Lond. 1902, 252.—Fla.

coccineus Walk., B. M. Cat. Hem. Het. vi, 93, 1873 (*Capsus*)
(not of Meyer 1843).

instabile Reut., Act. Soc. Sci. Fenn. xxxvi (2), 72, 1909.—E. States.

ONCEROMETOPUS.

Reuter, Cefv. Vet. Akad. Förh. 1875, 65.

nigriclavus Reut., Cefv. Vet. Akad. Förh. 1875 (9), 66.—Tex.

ruber Reut., Cefv. Vet. Akad. Förh. 1875 (9), 66.—Tex.

ORTHOCEPHALUS.

Fieber, Wien. Ent. Monatsschr. ii, 316, 1858.

saltator Hahn, Wanz. Ins. iii, 11, 1835 (*Capsus*); Fieber, Eur. Hem.
293, 1861; Prov., Pet. Faun. Can. Hem. 136, 1887.—N.
States.

SEMIUM.

Reuter, Cefv. Vet. Akad. Förh. 1875, 80.

hirtum Reut., Cefv. Vet. Akad. Förh. 1875 (9), 80; Act. Soc. Sci.
Fenn. xxxvi (2), 73, 1909.

elegans Uhler, Entom. Amer. iii, 149, 1887 (*Eccritotarsus*).—
S. States.

STIPHROSOMA.

Fieber, Wien. Ent. Monatsschr. ii, 312, 1858.

atrata Uhler, Proc. Calif. Acad. Sci. (2), iv, 268, 1894.—Calif.

croceipes Uhler, Proc. Ent. Soc. Wash. ii, 373, 1893.—Utah.

robusta Uhler, Hem. Colo. 45, 1895.—W. States.

stygica Say, Het. New Harm. 23, 1832; Compl. Writ. i, 344, 1859
(*Capsus*).—E. States.

Resthenini.

HADRONEMA.

Uhler, Hayden's Surv. Mont. 412, 1872.

militaris Uhler, Hayden's Surv. Mont. 412, 1872.—Colo., Utah, Calif.

picta Uhler, Hem. Colo. 31, 1895.—Colo., Dak.

princeps Uhler, Proc. Calif. Acad. Sci. (2) iv, 251, 1894.—W. States.

robusta Uhler, Proc. Calif. Acad. Sci. (2) iv, 250, 1894.—Calif.

LOPIDEA.

Uhler, Hayden's Surv. Mont. 411, 1872.

confluens Say, Het. New Harm. 23, 1832 (*Capsus*); Compl. Writ. i,
343, 1859.—E. States.

robinæ Uhler, Proc. Ent. Soc. Phila. i, 24, 1861 (*Capsus*).

floridana Walk., B. M. Cat. Hem. Het. vi, 97, 1873 (*Capsus*).—Fla.

- marginata** Uhler, Proc. Calif. Acad. Sci. (2) iv, 249, 1894 (equal *L. floridana* acc. Distant).—U. S.
media Say, Het. New Harm. 22, 1832; Compl. Writ. i, 341, 1859 (*Capsus*); Bull. 69 Minn. Sta., fig. 56.—U. S.
nigridia Uhler, Hem. Colo. 30, 1895.—Colo.

RESTHENIA.

Spinola, Essai, 184, 1837.

- Platytylus* Fieber, Wien. Ent. Monatsschr. ii, 306, 1858.
atripennis Reut., Cefv. Vet. Akad. Förh. 1875 (9), 65.—Tex.
circumcincta Say, Het. New Harm. 22, 1832 (*Capsus*); Compl. Writ. i, 343, 1859.
confraterna Uhler, Hayden's Surv. Mont. 411, 1872; Proc. Bost. Soc. Nat. Hist. xix, 399, 1878.—W. States.
maculicollis Reut., Cefv. Vet. Akad. Förh. 1875 (9), 65.
divisa Her.-Schf., Wanz. Ins. ix, 166, 1853 (*Capsus*); Stål, Stett. Ent. Zeit. 1862, 317.—Tex.
tetrastigma Her.-Schf., Wanz. Ins. ix, 166, 1853 (*Capsus*).
incisus Walk., B. M. Cat. Hem. Het. vi, 92, 1873 (*Capsus*).—Fla.
insignis Say, Het. New Harm. 22, 1832 (*Capsus*); Compl. Writ. i, 342, 1859.—U. S.
insitiva Say, Het. New Harm. 21, 1832 (*Capsus*); Compl. Writ. i, 340, 1859.—U. S.
xanthomelas Walk., B. M. Cat. Hem. Het. vi, 192, 1873 (*Capsus*).
nigricollis Reut., Cefv. Vet. Akad. Förh. 1875 (9), 65.—N. Jer.
ornaticollis Stål, Stett. Ent. Zeit. 1862, 317; Dist., Biol. Cent.-Amer. Rhynch. Het. i, 253, 424, pl. xxii, fig. 5, 1883.—Tex.
bicinctus Walk., Cat. Hem. B. M. vi, 100, 1873.
rubrovittata Stål, Stett. Ent. Zeit., 1862, 318.—Tex.

Mirini.

COLLARIA.

- Provancher, Nat. Can. iv, 79, 1872.
Nabidea Uhler, Proc. Bost. Soc. Nat. Hist. xix, 397, 1878.
Trachelomiris Reuter, Cefv. Vet. Akad. Förh. 1875, 61.
meilleurii Prov., Nat. Can. iv, 79, 1872.—E. States.
coracina Uhler, Proc. Bost. Soc. Nat. Hist. xix, 398, 1878 (*Nabidea*).
oculatus Reut., Cefv. Vet. Akad. Förh. 1875 (9), 61.—N. Y., Tex.

CREONTIADES.

- Distant, Biol. C.-Am. Rhynch. Het. i, 237, 1883.
filicorne Walk., B. M. Cat. Hem. Het. vi, 96, 1873 (*Capsus*).—Fla.
rubrinervis Stål, Stett. Ent. Zeit. 1862, 321 (*Megacoelum*); Dist., Biol. C.-Am. Rhynch. Het. i, 237, 1883, pl. xxiii, fig. 12.—Tex.

MEGALOCERCEA.

Fieber, Wien. Ent. Monatsschr. ii, 301, 1858.

debilis Uhler, Hayden's Surv. Mont. 408, 1872; Reuter, Act. Soc. Sci. Fenn. xxxvi (2), 5, 1909.—W. States.

rubicunda Uhler, Hayden's Surv. Mont. 409, 1872.—Colo.

MESOMIRIS.

Reuter, Act. Soc. Sci. Fenn. xxxvi (2), 4, 1909.

curtulus Reut., Act. Soc. Sci. Fenn. xxxvi (2), 5, 1909.—Md.

MIRIS.

Fabricius, Ent. Syst. iv, 183, 1794.

Leptopterna Fieber, Wien. Ent. Monatsschr. ii, 304, 1858.

amœna Uhler, Hayden's Surv. Mont. 409, 1872.—Idaho, Dak.

dolobrata Linn., Syst. Nat. (10), 449, 1758 (*Cimex*); Fieber, Eur. Hem. 245, 1861.—U. S.

belangeri Prov., Nat. Can. iv, 78, 1872 (*Miris*).

dorsalis Say, Het. New Harm. 25, 1832; Compl. Writ. i, 348, 1859.—E. States.

vicina Prov., Nat. Can. iv, 77, 1872.—U. S.

instabilis Uhler, Rept. Geog. and Geol. Surv. 1875, v, 836; Bull. 5 U. S. Geol. Surv. Terr. ii, 316, 1876; Proc. Bost. Soc. Nat. Hist. xix, 395, 1878.

affinis Reut., Cefv. Vet. Akad. Förh. 1875 (9), 59; Bull. 69 Minn. Sta. fig. 57.

STENODEMA.

Laporte, Essai, 34, 1832.

Lobostethus Fieber, Wien. Ent. Monatsschr. ii, 301, 1858.

Brachytropis Fieber, *ibid.*

trispinosum Reut., Cefv. Finska Vet. Soc. Förh., Vol. 46 (No. 15), 8, 1904; Act. Soc. Sci. Fenn. xxxvi (2), 4, 1909.—E. States.

calcaratus (*Brachytropis*) of American authors, not of Fallen.

TERATOCORIS.

Fieber, Wien. Ent. Monatsschr. ii, 302, 1858.

discolor Uhler, Entom. Amer. iii, 68, 1888.—E. States.

herbaticus Uhler, Entom. Amer. iii, 67, 1888; Reut., Act. Soc. Sci. Fenn. xxxvi (2), 7, 1909.—Labr.

miraria Atk., Cat. Capsidæ, 38, 1889.

saundersi Douglas and Scott, Ent. Mo. Mag. v, 260, 1869; Reut., Act. Soc. Sci. Fenn. xxxvi (2), 7, 1909.—Colo.

longicornis Uhler, Hem. Colo. 29, 1895.

TRIGONOTYLUS.

Fieber, Wien. Ent. Monatsschr. ii, 302, 1858.

Callimiris Reuter, Öfv. Vet. Akad. Förh. 1875, (9), 60.

brevipes Jakovec, Hor. Soc. Ent. Ross. xi, 63, 1880; Reuter, Ann. Mus. Zool. Acad. Imp. Sci. St. Petersburg. ix, 5, 1904; Act. Soc. Sci. Fenn. xxxvi (2), 6, 1909.—N. States.

pallidicornis Reut., Öfv. Finska Vet. Soc. Förh. xlii, 161, 1899.

tenuis Reut., *ibid*, xlix, p. 1, 1907.

confusus Reut., Act. Soc. Sci. Fenn. xxxvi (2), 6, 1909.—Md.

ruficornis Fourc., Ent. Paris, 209, 1785 (*Cimex*); Fall., Hem. Suec. 133, 1829 (*Miris*); Fieber, Europ. Hem. 243, 1861.—E. States.
viridis Prov., Nat. Can. iv, 78, 1872 (*Miris*).

pulcher Reut., Öfv. Vet. Akad. Förh. 1875 (9), 59.—Tex.

uhleri Reut., Öfv. Vet. Akad. Förh. 1875 (9), 60 (*Callimiris*).—U. S.

tarsalis Reut., Öfv. Vet. Akad. Förh. 1875 (9), 60 (*Callimiris*); Act. Soc. Sci. Fenn. xxxvi (2), 6, 1909.—Wisc., Tex.

Capsini.

ADELPHOCORIS.

Reuter, Öfv. Finska Vet. Förh. xxxviii, 168, 1896.

rapidus Say, Het. Hem. N. Harm. 20, 1832 (*Capsus*); Compl. Writ. i, 339, 1859; Bull. 57 Bur. Entom., fig. 26; 23 Ill. Rept. p. 199.—U. S.

multicolor H. Schf., Wanz. Ins. viiii 19, 1848 (*Capsus*).

superbus Uhler, Wheeler Surv. 100 Merid. v, 838, 1875 (*Calocoris*).—W. States.

CALOCORIS.

Fieber, Wien. Ent. Monatsschr. ii, 305, 1858.

Closterotomus Fieber, *ibid*, 306.

Lophyrus Kol., Mel. Entom. ii, 106, 1845 (preoc.).

fulvomaculatus De Geer, Mém. Ins. iii, 294, 1773 (*Cimex*); Fieber, Eur. Hem. 253, 1861; Reut. Gymn. Eur. v, 184, 1896.—Can.

norvegicus Gmelin, Syst. Nat. iv, 2176, 1788 (*Cimex*); Reut., Gym. Eur. v, 204, 1896.—U. S.

bipunctatus Fabr., Syst. Rhyng. 235, 1803 (*Cimex*).

contiguus Walk., B. M. Cat. Hem. Het. vi, 95, 1873 (*Capsus*).

stramineus Walk., *ibid*, 96.

palmeri Uhler, Hayden's Surv. Mont., 410, 1872.—W. States.

tinctus Uhler, Hem. Colo, 34, 1895.—Colo.

CAMPTOBROCHIS.

Fieber, Wien. Ent. Monatsschr. ii, 304, 1858.

Callicapsus Reut., Öfv. Vet. Akad. Förh. 1875 (9), 75.

Euarmosus Reut., *ibid*, 76.

Mycterocoris Uhler, Proc. U. S. Nat. Mus. xxvii, 358, 1904.

- brevis** Uhler, Proc. U. S. Nat. Mus. xxvii, 359, 1904; Reut., Act. Soc. Sci. Fenn. xxxvi (2), 59, 1909.—Nev.
- cerachates** Uhler, Proc. Calif. Acad. Sci. (2) iv, 265, 1894 (*Deræocoris*); Reut. Act. Soc. Sci. Fenn. xxxvi (2), 57, 1909.—Calif., Ariz.
- fulvescens** Reut., Act. Soc. Sci. Fenn. xxxvi (2), 56, 1909.—Colo.
- grandis** Uhler, Entom. Amer. ii, 230, 1887; Reut., Act. Soc. Sci. Fenn. xxxvi (2), 56, 1909.—E. States.
- histrion** Reut., Öfv. Vet. Akad. Förh. 1875 (9), 75 (*Callicapsus*); Reut., Act. Soc. Sci. Fenn., xxxvi (2), 58, 1909.—Car., Tex.
- nebulosus** Uhler, Hayden's Surv. Mont. 417, 1872; Proc. Bost. Soc. Nat. Hist. xix, 408, 1878; Reut., Act. Soc. Sci. Fenn. xxxvi (2), 59, 1909.—Dak., Colo.
- nigrita** Reut., Act. Soc. Sci. Fenn. xxxvi (2), 55, 1909.—D. C.
- nitens** Reut., Act. Soc. Sci. Fenn. xxxvi (2), 56, 1909.—N. Y., D. C., Colo.
- robustus** Uhler, Hem. Colo. 39, 1895.—Colo.
- sayi** Reut., Öfv. Vet. Akad. Förh. 1875 (9), 76 (*Euarmonus*); Act. Soc. Sci. Fenn. xxxvi (2), 55, 1909.—Pa., Tex.
- schwarzi** Uhler, Proc. Ent. Soc. Wash. ii, 375, 1898; Reut., Act. Soc. Sci. Fenn. xxxvi (2), 58, 1909.—Ut., Wash., Can.
- validus** Reut., Act. Soc. Sci. Fenn. xxxvi (2), 58, 1909.—U. S.

CAPSUS.

Fabricius, Syst. Hyng. 241, 1803.

Rhopalotomus Fieber, Wien. Ent. Monatsschr. ii, 307, 1858.

- ater** Linn., Syst. Nat. (10), 447, 1758 (*Cimex*); Fieber, Eur. Hem. 264, 1861; Reut., Gym. Eur. V, 14, 1896.—U. S.

COCCOBAPHES.

Uhler, Proc. Bost. Soc. Nat. Hist., xix, 401, 1878.

- sanguinarus** Uhler, Proc. Bost. Soc. Nat. Hist. xix, 401, 1878; Reut., Act. Soc. Sci. Fenn. xxxvi (2), 37, 1909.—E. States.

DERÆOCORIS.

Kirschbaum, Jahrb. Ver. Nat. Nassau, x, 208, 1855.

Macrocapsus Reuter, Öfv. Finsk. Vet. Förh. xxi, 55; 1879.

- approximatus** Stål, Stett. Ent. Zeit. 1858, 185.—Alaska.

- seguisinus** Müll., Mant. Ins. Taur. 191, 1766 (*Cimex*); Reut., Gym. Eur. v, 32, 1896.—N. States.

capillaris Fabr., Syst. Ent. 725, 1775 (*Cimex*); Fieber, Eur. Hem. 266, 1861.

DICHROOSCYTUS.

Fieber, Wien. Ent. Monatsschr. ii, 309, 1858.

- elegans** Uhler, Proc. U. S. Nat. Mus. xxvii, 356, 1904.—Ariz.

- suspectus** Reut., Act. Soc. Sci. Fenn. xxxvi (2), 37, 1909.—Mass., D. C., N. Y.

rufipennis Auct. Amer. (nec Fall.).

ECERTOBIA.

Reuter, Act. Soc. Sci. Fenn. xxxvi (2), 36, 1909.

decora Reut., Act. Soc. Sci. Fenn. xxxvi (2), 36, 1909.—Colo.

EURYCHLOPTERELLA.

Reuter, Act. Soc. Sci. Fenn. xxxvi (2), 59, 1909.

luridula Reut., Act. Soc. Sci. Fenn. xxxvi (2), 60, 1909.—N. Y., D. C.

EUSTICTUS.

Reuter, Act. Soc. Sci. Fenn. xxxvi (2), 35, 1909.

Megacælum Uhler, Check list, 1887.

catulus Uhler, Proc. Calif. Acad. Sci. (2), iv, 237, 1894 (*Megacælum*).
—Tex., Pa.

pusillus Uhler, Entom. Amer. iii, 71, 1887 (*Megacælum*).—Ariz.

mundus Uhler, Entom. Amer. iii, 71, 1887 (*Megacælum*).—Ga., Fla.

grossus Uhler, Entom. Amer. iii, 70, 1887 (*Megacælum*); Reut., Act.
Soc. Sci. Fenn. xxxvi (2), 35, 1909.—E. States.

GARGANUS.

Stål, Stett. Ent. Zeit. 1862, 321.

albidivittis Stål, Stett. Ent. Zeit. 1852, 322; Dist., Biol. C.-Amer.

Rhynch. Het. I, pl. xxv, fig. 23.—Tex.

fusiformis Say, Het. New Harm. 23, 1832; Compl. Writ. i, 344, 1859
(*Capsus*).—U. S.

croceipes H. Schf., Wanz. Ins. viii, 16, 1848 (*Capsus*).

HORCIAS.

Distant, Biol. C.-Am. Rhynch. Het. i, 117, 1881.

Metriorrhynchus Reuter, Cefv. Vet. Ak. Förh. 1875 (9), 74 (pre-
occupied).

Metriorrhynchomiris Kirk., Trans. Amer. Ent. Soc. xxxii, 140,
1906.

fallax Reut., Act. Soc. Sci. Fenn. xxxvi (2), 42, 1909.—Pa.

limbatellus Walk., B. M. Cat. Hem. Het. vi, 93, 1873 (*Capsus*).—N. Y.

affinis Reuter, Cefv. Vet. Akad. Förh. 1875 (9), 74 (*Pæcilocap-*
sus).

marginalis Reut., Cefv. Vet. Akad. Förh. 1875 (9), 75 (*Pæcilocapsus*).
—N. Y.

nigrita Reut. (var.), Act. Soc. Sci. Fenn. xxxvi (2), 41, 1909.

sexmaculatus Barber, Bull. Brookl. Ins. Arts. Sci. i (9), 280, 1906
(*Pæcilocapsus*).—Tex.

IRBISIA.

Reuter, Cefv. Finsk. Vet. Förh xxi, 57, 1879.

Thyrilus Uhler, Proc. Cal. Acad. Sci. (2), iv, 267, 1894.

Leptomerocoris Stål, Stett. Ent. Zeit. 1858, 188 (nec Kirschb.).

- brachycerus** Uhler, Hayden's Surv. Mont. 416, 1872 (*Rhopalotomus*).
—Calif., Colo.
- gilvipes** Stål, Stett. Ent. Zeit. 1858, 187 (*Leptomerocoris*).—Alaska.
- pacificus** Uhler, Hayden's Surv. Mont. 415, 1872 (*Rhopalotomus*).—
W. States.
- sericans** Stål, Stett. Ent. Zeit. 1858, 188 (*Leptomerocoris*); Reut.,
Gym. Eur. v, 12, 1896.—Alaska.

LYGIDEA.

- Reuter, Öfv. Finska Vet. Förh. xxi, 54, 1879.
- mendax** Reut., Act. Soc. Sci. Fenn. xxxvi (2), 47, 1909. N. Y.
- morio** Reut., Act. Soc. Sci. Fenn. xxxvi (2), 47, 1909.—Calif.
- rubeculus** Uhler, Hem. Colo. 37, 1895 (*Neoborus*); Reuter, Act. Soc.
Sci. Fenn. xxxvi (2), 45, 1909.—N. States.
- var. *lurida*, *rosacea*, *infuscata* et *obscura* Reut., ibid 46.

LYGUS.

- Hahn, Wanzen Ins. i, 147, 1833.
- apicalis** Fieber, Eur. Hem. 275, 1861; Reuter, Hem. Gymn. Eur. v,
110, 1896; Act. Soc. Sci. Fenn. xxxvi, (2), 43, 1909.—Tex.,
Me.
- prasinus* Reut., Öfv. Vet. Akad. Förh. 1875 (9), 72.
- annexus** Uhler, Hayden's Surv. Mont. 413, 1872.—Colo., Minn.
- approximatus** Stål, Stett. Ent. Zeit. xix, 185, 1859; Reuter, Hem.
Gymn. Eur. v, 91, 1896; Act. Soc. Sci. Fenn. xxxvi (2), 43,
1909.—N. States.
- belfragei** Reut., Öfv. Vet. Akad. Förh. 1875 (9), 71.—N. Y., Tex.
- campestris** Linn., Syst. Nat. (10), 448, 1758 (*Cimex*); Reuter, Act.
Soc. Sci. Fenn. xxxvi (2), 44, 1909.—N. States.
- pastinacæ* Fall., Mon. Cim. 86, 1807; Fieber, Europ. Hem. 278,
1861.
- scutellatus* Uhler, Bull. U. S, Geol. and Geog. Surv. Terr. iii,
420, 1877.
- carolinæ** Reut., Öfv. Vet. Akad. Förh. 1875 (9), 71.—Car.
- contaminatus** Fall., Hem. Suec. 97, 1829 (*Phytocoris*); Fieber, Eur.
Hem. 274, 1861; Reut., Hem. Gymn. Eur. v. 112, 1896.
- convexicollis** Reut., Öfv. Vet. Akad. Förh. 1875 (9), 72; Act. Soc.
Sci. Fenn. xxxvi (2), 43, 1909.—N. States.
- distinguendus** Reut., Pet. Nouv. Entom. 1875, 541; Hem. Gymn.
Eur. v, 74, 1896; Act. Soc. Sci. Fenn. xxxvi (2), 45, 1909.
—Can.
- fasciatus** Reut., Öfv. Vet. Akad. Förh. 1875 (9), 72.—S. Car.
- flavonotatum** Prov., Nat. Can. iv, 103, 1872 (*Capsus*); Pet. Faun. Can.
Hem. 120, 1887.—Can.
- strigulatus* Walk., B. M. Cat. Hem. Het. vi, 94, 1873 (*Capsus*).
- guttatipes** Uhler, Hem. Colo. 35, 1895.—Colo.

- invitus** Say, Het. New Harm. 24, 1832; Compl. Writ. i, 345, 1859 (*Capsus*); Uhler, Proc. Bost. Soc. Nat. Hist. xix, 407, 1878.—U. S.
- lucorum** Meyer, Rhyn. Schweiz. 46, 1843 (*Capsus*); Fieber, Eur. Hem. 275, 1861; Reut., Hem. Gymn. Eur. v, 107, 1896.—N. States.
- olivaceus** Reuter, Æfv. Finsk. Vet. Soc. Förh. xlix (5), 6, 1907.—Fla.
- pabulinus** Linn., Faun. Suec. 253, 1761 (*Cimex*); Fieber, Eur. Hem. 276, 1861; Reut., Hem. Gymn. Eur. v, 114, 1896.—N. States.
- pabulinus signifer** Reut., Act. Soc. Sci. Fenn. xxxvi (2), 42, 1909.—Md
- plagiatus** Uhler, Hem. Colo. 35, 1895.—U. S.
- pratensis** Linn., Syst. Nat. (10), 448, 1758 (*Cimex*); Fieber, Eur. Hem. 273, 1861; Reut., Hem. Gymn. Eur. v, 98, 1896; Bull. 43, Bur. Entom., fig. 51.—U. S.
- lineolaris* Pal. Beauv., Ins. Afr. Amer., 187, 1805.
- oblineatus* Say, Het. New Harm. 21, 1832; Compl. Writ. i, 340, 1859 (*Capsus*).
- diffusus* Uhler, Hayden's Surv. Mont., 413, 1872.
- redimitus* Uhler, *ibid*.
- rubicundus** Fall., Hem. Suec. 92, 1829; Reut., Hem. Gymn. Eur. v, 72, 1896; Act. Soc. Sci. Fenn. xxxvi (2), 45, 1909.—W. States.
- pulverulenta* Uhler, Trans. Md. Acad. Sci. 1892, 183; Proc., U. S. Natl. Mus. xxvii, 357, 1904 (*Hadrodema*).
- sallei** Stål, Stett. Ent. Zeit. 1862, 321.—W. States.
- scutellatus** Dist., Biol. Cent.-Amer. Rhynch. Het. i, 274, pl. xxiii, fig. 9, 18, 1884.—Tex.
- viridis** Fall., Monog. Cim. 85, 1807 (*Cimex*); Reut., Hem. Gymn. Eur. v, 113, 1896; Act. Soc. Sci. Fenn. xxxvi (2), 42, 1909.—N. H., Mass.
- vitticollis** Reut., Æfv. Vet. Akad. Förh. 1875 (9), 71; Reut., Act. Soc. Sci. Fenn. xxxvi (2), 43, 1909.—E. States.
- monachus* Uhler, Can. Entom. 1886, 208; Bull. 13, Div. Entom. U. Dept. Agric. 1887, 63.
- vividus** Uhler, Proc. Calif. Acad. Sci. (2), iv, 260, 1894.—Calif.

NEOBOROPS.

Uhler, Hem. Colo. 36, 1895.

vigilax Uhler, Hem. Colo. 36, 1895.—Colo., Ariz.

NEOCAPSUS.

Distant, Biol. C.-Am. Rhynch. Het. i, 277, 1884.

cuneatus Distant, Biol. C.-Am. Rhynch. Het. i, 438, 1893, pl. xxxviii, fig. 6.—Tex.

NEUROCOLPUS.

Reuter, *Œfv. Vet. Akad. Förh.* 1875 (9), 69.

inops Uhler, Bull. U. S. Geol. Surv. iii, 413, 1877 (*Phytocoris*); Proc. Bost. Soc. N. H. xix, 402, 1878 (*Phytocoris eximius?*).—E. States.

mexicanus Distant, Biol. Cent.-Am. Rhynch. Het. i, 262, pl. xxiii, fig. 5, 1883; Reut., Act. Soc. Sci. Fenn. xxxvi (2), 13, 1909. N. Mex.

nubilus Say, Het. Hem. N. Harm. 22, 1832 (*Capsus*); Compl. Writ. i, 341, 1859; Reuter, *Œfv. Vet. Akad. Förh.* 1875 (9), 70.—U. S.

hirsutulus Walk., B. M. Cat. Hem. Het. vi, 95, 1873 (*Capsus*).

affinis Dist., Biol. C.-Amer. Rhynch. Het. i, 263, 1883.

PALLACOCORIS.

Reuter, *Œfv. Vet. Akad. Förh.* 1875 (9), 62.

suavis Reut., *Œfv. Vet. Akad. Förh.* 1875 (9), 62.—Tex.

PARACALOCORIS.

Distant, Biol. C.-Am. Rhynch. Het. i, 263, 1883.

colon Say, Het. New Harm. 25, 1832 (*Capsus*); Compl. Writ. i, 346, 1859.—E. States.

deleticus Reut., Act. Soc. Sci. Fenn. xxxvi (2), 39, 1909.—Colo.

heidemanni Reut., Act. Soc. Sci. Fenn. xxxvi (2), 40, 1909.—W. Va.

jurgiosus Stål, Stett. Ent. Zeit. 1862, 320 (*Calocoris*); Distant, Biol. C.-Am. Rhynch. Het. i, 263, 1883, pl. xxv, fig. 17.—Tex.

multisignatus Reut., Act. Soc. Sci. Fenn. xxxvi (2), 40, 1909.—D. C.

scrupeus Say, Het. New Harm. 22, 1832 (*Capsus*); Compl. Writ. i, 342, 1859; Reut., Act. Soc. Sci. Fenn. xxxvi (2), 39, 1909.—U. S.

tetrastigma H. Schf., Wanz. Ins. ix, 166, 1853.

externus H. Schf., *ibid.* viii, 16, 1848.

PARAXENETUS.

Reuter, Act. Soc. Sci. Fenn. xxxvi (2), 13, 1909.

guttulatus Uhler, Entom. Amer. iii, 150, 1888 (*Eucercocoris*); Reut., Act. Soc. Sci. Fenn. xxxvi (2), 13, 1909.—E. States.

PHYTOCORIS.

Fallen, Spec. Nov. Hem. 10, 1814.

Compsocercocoris Reut., *Œfv. Vet. Akad. Förh.* 1875 (9), 70.

Callodemus Uhler, Hem. Colo. 33, 1895.

augustulus Reut., Act. Soc. Sci. Fenn. xxxvi (2), 29, 1909.—W. Va.

annulicornis Reut., *Œfv. Vet. Akad. Förh.* 1875 (9), 70 (*Compsocercocoris*); Reut., Act. Soc. Sci. Fenn. xxxvi (2), 33, 1909.—Tex.

- antennalis** Reut., Act. Soc. Sci. Fenn. xxxvi (2), 32, 1909.—D. C., Fla.
- bakeri** Reut., Act. Soc. Sci. Fenn. xxxvi (2), 28, 1909.—Calif.
- breviusculus** Reut., Œfv. Vet. Akad. Förh. 1875 (9), 68; Act. Soc. Sci. Fenn. xxxvi (2), 21, 1909.—Tex.
- canescens** Reut., Act. Soc. Sci. Fenn. xxxvi (2), 30, 1909.—Calif.
- conspersipes** Reut., Act. Soc. Sci. Fenn. xxxvi (1), 22, 1909.—D. C.
- eximius** Reut., Œfv. Vet. Akad. Förh. 1875 (9), 67; Act. Soc. Sci. Fenn. xxxvi (2), 23, 1909.—Tex.
- fenestratus** Reut., Act. Soc. Sci. Fenn. xxxvi (2), 24, 1909.—D. C., N. Car.
- fumatus** Reut., Act. Soc. Sci. Fenn. xxxvi (2), 25, 1909.—D. C.
- heidemanni** Reut., Act. Soc. Sci. Fenn. xxxvi (2), 27, 1909.—N. Mex.
- interspersus** Uhler, Hem. Colo. 32, 1895; Reut., Act. Soc. Sci. Fenn. xxxvi (2), 19, 1909.—Colo.
- lævis** Uhler, Hem. Colo. 33, 1895 (*Callodemas*).—Colo., N. Mex.
- lasiomerus** Reut., Act. Soc. Sci. Fenn. xxxvii (2), 34, 1909.—Can.
- lineatus** Reut., Act. Soc. Sci. Fenn. xxxvi (2), 30, 1909.—Can.
- minutulus** Reut., Act. Soc. Sci. Fenn. xxxvi (2), 24, 1909.—Md.
- mundus** Reut., Act. Soc. Sci. Fenn. xxxvi (2), 18, 1909.—Md., D. C.
- pallidicornis** Reut., Œfv. Vet. Akad. Förh. 1875 (9), 69; Act. Soc. Sci. Fenn. xxxvi (2), 33, 1909.—Wisc.
- palmeri** Reut., Act. Soc. Sci. Fenn. xxxvi (2), 32, 1909.—Can.
- politus** Reut., Act. Soc. Sci. Fenn. xxxvi (2), 21, 1909.—Nev.
- puella** Reut., Œfv. Vet. Akad. Förh. 1875 (9), 69; Act. Soc. Sci. Fenn. xxxvi (2), 20, 1909.—E. States.
var. *confluens* et *infuscata* Reut., *ibid*, 20.
- roseus** Uhler, Proc. Cal. Acad. Sci. (2), iv, 253, 1894; (*Compsocero-coris*); Reut., Act. Soc. Sci. Fenn. xxxvi (2), 27, 1909.—Cal.
- subnitidus** Reut., Act. Soc. Sci. Fenn. xxxvi (2), 26, 1909.—D. C.
- tibialis** Reut., Œfv. Vet. Akad. Förh. 1875 (9), 68; Act. Soc. Sci. Fenn. xxxvi (2), 20, 1909.—E. States.
vilis Dist., Biol. C.-Am. Rhynch. Het. i, 260, pl. xxv, fig. 13, 1883.
- validus** Reut., Act. Soc. Sci. Fenn. xxxvi (2), 31, 1909.—Colo.
- vittatus** Reut., Act. Soc. Sci. Fenn. xxxvi (2), 28, 1909.—N. Y.

POECILOCAPSUS.

- Reuter, Œfv. Vet. Akad. Förh. 1875, 73.
- dislocatus** Say, Het. New Harm. 21, 1832 (*Capsus*); Compl. Writ. i, 339, 1859; Uhler, Proc. Bost. Soc. Nat. Hist. xix, 406, 1878.—E. States.
- goniphorus** Say, Het. New Harm. 21, 1832 (*Capsus*); Compl. Writ. i, 341, 1859.—U. S.
melaxanthus H. Schf., Wanz. Ins. viii, 18, 1848, fig. 794 (*Capsus*).

- lineatus** Fabr., Entom. Syst. Supp. 541, 1798 (*Lygæus*); Slingerland, Bull. 58 Cornell Agric. Exp. Sta. 1903,—U. S.
4-vittatus Say, Het. New Harm. 20, 1832 (*Capsus*); Compl. Writ. i, 339, 1859.
bellus Emmons, Nat. Hist. N. Y. Agric. v, pl. 30, fig. 1, 1854.
marginatus Walk., B. M. Cat. Hem. Het. vi, 96, 1873 (*Capsus*).—N. Y.
marmoratus Uhler, Proc. Cal. Acad. Sci. (2), iv, 263, 1894.—Tex., Md.
nigriger Stål, Stett. Ent. Zeit. 1862, 339 (*Brachycoleus*).—Fla., Tex.

PÆCILOSCYTUS.

- Fieber, Wien. Ent. Monatsschr. ii, 311, 1858.
americanus Reut., Œfv. Vet. Akad. Förh. 1875 (9), 73 (*Systratiotus*); Act. Soc. Sci. Fenn. xxxvi (2), 61, 1909.—Tex.
 * **basalis** Reut., Œfv. Vet. Akad. Förh. 1875 (9), 73.—U. S.
sericeus Uhler, Bull. U. S. Geol. Surv. iii, 422, 1877.
basivittis Reut., Act. Soc. Sci. Fenn. xxxvi (2), 61, 1909.—Colo.
cuneatus Dist., Biol. Cent.-Am. Rhynch. Het. i, 435, 1885.—Tex., Fla., Va.
diffusus Uhler, Hayden's Surv. Mont. 415, 1872.—Colo., Vt., Id.
elegans Reut., Act. Soc. Sci. Fenn. xxxvi (2), 60, 1909.—Calif.
 * **intermedius** Uhler, Proc. Calif. Acad. Sci. (2) iv, 261, 1894.—Calif., Ariz.
obscurus Uhler, Proc. Zool. Soc. London, 1893, 715.—S. Fla.
unifasciatus Fabr., Entom. Syst. iv, 178, 1794 (*Lygæus*), Fieber, Eur. Hem. 276, 1861.—U. S.
venaticus Uhler, Hayden's Surv. Mont. 414, 1872 (*Systratiotus*).—U. S.
rubronotatus Prov., Nat. Can. iv, 105, 1872 (*Rhopalotomus*).

PYCNOPTERNA.

- Fieber, Wien. Ent. Monatsschr. ii, 307, 1858.
amœna Prov., Pet. Faun. Can. Hem., 114, 1887.—Can.

STENOTUS.

- Jakolev, Bull. Nat. Mosc. lii (1), 288, 1877.
Oncognathus Fieber, Wien. Ent. Monatsschr. ii, 303, 1858 (pre-occupied).
binotatus Fabr., Entom. Syst. iv, 172, 1794 (*Lygæus*); Fieber, Europ. Hem. 247, 1861 (*Oncognathus*); Insect Life, v, figs. 8, 9.—E. States.

TROPIDOSTEPTES.

- Uhler, Proc. Bost. Soc. Nat. Hist. xix, 404, 1878.
amœnus Reut., Act. Soc. Sci. Fenn. xxxvi (2), 48, 1909.—E. States.
 var. *palmeri*, *plagiata*, *signata* et *scutellaris* Reut., ibid, 49.
saxeus (*Neoborus*) Auct. Amer. (nec Dist.).

- cardinalis** Uhler, Proc. Bost. Soc. Nat. Hist. xix, 404, 1878; Reut., Act. Soc. Sci. Fenn. xxxvi (2), 50, 1909.—E. States.
commissuralis Reut., Act. Soc. Sci. Fenn. xxx (2), 51, 1909.—Can.
pettiti Reuter, Act. Soc. Sci. Fenn. xxxvi (2), 50, 1909.—Pa.
plagifer Reut., Act. Soc. Sci. Fenn. xxxvi (2), 51, 1909.—Can.

*Myrmecorini.***MIMOCEPS.**

- Uhler, Trans. Md. Acad. Sci. 1890, 83.
gracilis Uhler, Trans. Md. Acad. Sci. 1890, 85.—Wisc., N. Y., Can.
insignis Uhler, Trans. Md. Acad. Sci. 1890, 84.—Ill.

*Clivinemini.***CLIVINEMA.**

- Reuter, Œfv. Vet. Akad. Förh 1875, 63.
rubida Uhler, Proc. U. S. Nat. Mus. xxvii, 355, 1904.—Ariz.
villosa Reut., Œfv. Vet. Akad. Förh. 1875 (9), 63.—Tex.

*Bryocorini.***CAULOTOPS.**

- Bergroth, Wien. Ent. Zeit. 1898, 33.
agavis Reut., Act. Soc. Sci. Fenn. xxxvi (2), 1, 1909.—Ariz.

CYRTOCAPSUS.

- Reuter, Œfv. Vet. Akad. Förh. 1875, 78.
caligineus Stål, Freg. Eugenies Resa. Ins. 258, 1859 (*Capsus*).—Fla., Calif.
pallipes Dist, Biol. Cent.-Amer. Rhynch. Het. i, 303, 1884 (*Pirithous*).

DACOTA.

- Uhler, Hayden's Surv. Mont. 413, 1872.
hesperia Uhler, Hayden's Surv. Mont. 414, 1872.—W. States.

ECCRITOTARSUS.

- Stål, Kgl. Svensk. Vet. Handl. ii (7), 57, 1861.
atratus Dist., Biol. Cent.-Amer. Rhynch. Het. i, 285, 1883.—Cal., Tex.
incurvus Dist., Biol. Cent. Amer. Rhynch. Het. i, 285, 1883.—S. W. States.
scabrosus Uhler, Hem. Colo., 40, 1895.—Colo.

MONALOCORIS.

- Dahlbom, Vetensk. Acad. Handl. 1850, 209.
filicis Linn., Syst. Nat. (10) 443, 1758 (*Cimex*); Fieber, Europ. Hem. 238, 1861.—E. States.

PYCNODERES.

Guérin, Sagra's Hist. Cuba, vii, Ins. 168, 1857.

dilatatus Reut., Act. Soc. Sci. Fenn. xxxvi (2), 3, 1909.—Md., D. C.

quadrимaculatus Guérin, Sagra's Hist. phys. Cuba, vii, Ins. 168, 1857; Reuter, Act. Soc. Sci. Fenn. xxxvi (2), 2, 1909.—Tex., Ariz.

SIXEONOTUS.

Reuter, Cefv. Vet. Akad. Förh. 1875 (9), 77.

insignis Reut., Cefv. Vet. Akad. Förh. 1875 (9), 78.—Tex., N. J., N. Y.

luteiceps Reut., Act. Soc. Sci. Fenn. xxxvi (2), 4, 1909.—Tex.

tenebrosus Distant, Biol. C.-Am. Rhynch. Het. i, 441, 1885 (*Eccritotarsus*); Reuter, Act. Soc. Sci. Fenn. xxxvi (2), 3, 1909.—Fla., Kans., D. C.

CAPSUS. (Unplaced.)

flavipes Prov., Nat. Can. iv, 104, 1872.—Can.

frontifer Walk., B. M. Cat. Hem. Het. vi, 94, 1873 (type lost).—N. America.

geminus Say, Het. New Harm. 23, 1832; Compl. Writ. i, 344, 1859.—Ind.

imbecillus Say, Het. New Harm. 24, 1832; Compl. Writ. i, 345, 1859.—Ind.

obscurus Walk., B. M. Cat. Hem. Het. vi, 93, 1873 (type broken).—Can.

ocreatus Say, Hem. New Harm. 19, 1832; Compl. Writ. i, 338, 1859.—Ga.

pallens Walk., B. M. Cat. Hem. Het. vi, 94, 1873 (type lost).—Can.

submarginatus Say, Hem. New Harm. 23, 1832; Compl. Writ. i, 344, 1859.—Mo., Ind.

PHYMATIDÆ.

Macrocephalini.

MACROCEPHALUS.

Swederus, Vet. Akad. Nya Handl. viii, 183, 1787.

cimicoides Swed., Vet. Akad. Nya Handl. viii, 183, 1787; Handl., Ann. Nat. Hofm. Wien, xii, 186, 1897.—Ga., Fla.

manicatus Fabr., Syst. Rhynch. 123, 1803 (*Syrtsis*); H.-Schf., Wanz. Ins. viii, 107, 1848, fig. 878; Handl., Ann. Nat. Hofm. Wien, xii, 187, 1897.—Tex., Car.

notatus Westw., Trans. Ent. Soc. Lond. iii, 24, 1843; Handl., Ann. Nat. Hofm. Wien, xii, 188, 1897; Champion, Biol. C.-Am. Rhynch. ii, 56, 1898.—Tex.

incisus Stål, Stett. Ent. Zeit. 1862, 440.

cliens Stål, *ibid.*

- prehensilis** Fabr., Syst. Rhyng. 123, 1803 (*Syrtis*); Westw. Trans. Ent. Soc. Lond. iii, 26, 1843; Handl., Ann. Nat. Hofm. Wien. xii, 194, 1897.—Fla.
pallidus Westw., Trans. Ent. Soc. Lond. iii, 27, 1843.
uhleri Handl., Verh. zool.-bot. Ges. Wien. 1898, 383.—Ariz.
arizonicus Ckll., Entom. 1900, 66.

*Phymatini.***PHYMATA.**

- Latreille, Hist. Ins. iii, 247, 1802.
Syrtis Fabr., Syst. Rhyng. 121, 1803.
Discomerus Laporte, Essai, 14, 1832.
albopicta Handl., Ann. Nat. Hofm. Wien. xii, 151, 1897; Champ., Biol. C.-Am. Rhynch. Het. ii, 54, pl. iv, fig. 9, 1898.—Ga.
annulipes Stål, Stett. Ent. Zeit. 1862, 439; Enum. Hem. v, 136, 1876; Handl., Ann. Nat. Hofm. Wien. xii, 151, 1897.—Calif.
fasciata Gray, Griffith's Anim. Kingd. Ins. ii, 242, pl. 93, fig. 3, 1832; Handl., Ann. Nat. Hofm. Wien. xii, 161, 1897.—E. States.
wolffi Stål, Enum. Hem. v, 133, 1876.
erosa Wolff, Icon. Cim. iii, 89, 1802 (nec Linné).
noualhieri Handl., Ann. Nat. Hofm. Wien. xii, 153, 1897; Champ., Biol. C.-Am. Rhynch. Het. ii, 54, pl. iv, fig. 8. 1898.—Pa., Ga.
pennsylvanica Handl., Ann. Nat. Hofm. Wien. xii, 163, 1897.—E. States.
salicis Cockerell, Entom. 1900, 66.—Ariz.
vicina Handl., Ann. Nat. Hofm. Wien. xii, 150, 1897.—Fla.

ARADIDÆ.

*Aradini.***ARADUS.**

- Fabricius, Syst. Rhyng. 116, 1803.
Piestosoma Laporte, Essai, 35, 1832.
Quilnus Stål, Enum. Hem. iii, 137, 1873.
abbas Bergr., Bull. Soc. Ent. Belg. 1889, clxxx.—Can., N. Y.
acutus Say, Het. New Harm. 28, 1832, Compl. Writ. i, 351, 1859.—E. States.
americanus Her.-Schf., Wans. Ins. viii, 115, 1848.
æqualis Say, Het. New Harm. 29, 1832; Compl. Writ. i, 352, 1859; Osborn, Ohio Nat. iv, 36, 1903.—E. States.
ampliatus Uhler, Bull. U. S. Geol. Surv. ii (5), 321, 1876.—Calif.
behrensi Bergr., Wien. Ent. Zeit. v, 97, 1886.—Calif.
borealis Heid., Proc. Ent. Soc. Wash. xi, 190, 1910.—N. H., Mich.
breviatus Bergr., Rev. d'Ent. 1887, 245.—D. C., Fla.

- caliginosus** Walk., B. M. Cat. Hem. Het. vii, 36, 1873.—B. Am.
- cincticornis** Bergr., Can. Entom. 1906, 198.—Ala.
- cinnamoneus** Panz., Fauna Ins. Germ. 100, 20, 1794; Fieber, Eur. Hem. 111, 1861; Bull. 69 Minn. Sta., fig. 37.—E. States.
- coarctatus** Heid., Proc. Ent. Soc. Wash. viii, 69, 1906.—Calif.
- compressus** Heid., Proc. Ent. Soc. Wash. viii, 70, 1906.—Wash.
- concinus** Bergr., Proc. Ent. Soc. Wash. ii, 337, 1892; Can. Ent. 1906, 198.—Calif.
- crenatus** Say, Het. New Harm. 28, 1832; Compl. Writ. i, 350, 1859; Osborn, Ohio Nat. iv, 37, 1903.—N. States.
- dilatatus* Dufour, Ann. Soc. Ent. France, 1844, 452.
- debilis** Uhler, Bull. U. S. Geol. Surv. ii (5), 322, 1876.—B. Col., Vanc.
- duryi** Osborn, Ohio Nat. iv, 39, 1903.—Ohio.
- duzei** Bergr., Proc. Ent. Soc. Wash. ii, 333, 1892; Osborn, Ohio Nat. iv, 38, 1903.—Can., Pa., Ohio.
- falleni** Stål, Rio. Hem. i, 68, 1858; Enum. Hem. iii, 136, 1873; Champ., Biol. Cent.-Amer. Rhynch. Het. ii, 66, 1898, pl. v, fig. 1.—S. States.
- pallidicornis* Stål, Enum. Hem. iii, 136, 1873.
- fasciicornis** Walk., B. M. Cat. Hem. Het. vii, 36, 1873.—Nova Scotia.
- fenestratus** Walk., B. M. Cat. Hem. Het. vii, 36, 1873.—B. Am.
- fuscoannulatus** Stål, Freg. Eugeniès Resa, Hem. 260, 1859.—Calif.
- gracilicornis** Stål, Enum. Hem. iii, 136, 1873; Bergroth, Rev. d'Entom. 1887, 246.—Fla.
- heidemanni** Bergr., Can. Entom. 1906, 199.—Oreg.
- hubbardi** Heid., Proc. Ent. Soc. Wash. vi, 232, 1904.—W. States.
- inornatus** Uhler, Bull. U. S. Geol. Surv., ii (5), 323, 1876.—U. S.
- lugubris** Fall., Hem. Suec. 139, 1829; Fieber, Eur. Hem. 114, 1861.—U. S.
- rectus* Say, Het. N. Harm. 29, 1832; Compl. Writ. i, 352, 1859.
- affinis* Kirby, Fauna Bor.-Amer. iv, 279, 1837; Can. Entom. 1878, 213.
- niger** Stål, Enum. Hem. iii, 137, 1873.—S. States.
- ornatus** Say, Het. New Harm. 29, 1832; Compl. Writ., 352, 1859; Osborn, Ohio Nat. iv, 38, 1903.—E. States.
- proboscideus** Walk., B. M. Cat. Hem. Het. vii, 35, 1873.—B. Am.
- quadrilineatus** Say, Journ. Acad. Phila. iv, 326, 1825; Compl. Writ. ii, 249, 1859.—E. States.
- robustus** Uhler, Proc. Bost. Soc. Nat. Hist. xiv, 104, 1871; xix, 419, 1878; Bull. 69 Minn. Sta., figs. 35, 36; Osborn, Ohio Nat. iv, 37, 1903.—E. States.
- shermani** Heid., Proc. Ent. Soc. Wash. viii, 68, 1906.—N. Car., Pa.
- similis** Say, Het. New Harm. 28, 1832; Compl. Writ. i, 351, 1859.—U. S.
- centriguttatus* Bergr., Rev. d'Entom. 1887, 246.

tuberculifer Kirby, Fauna Bor.-Amer. iv, 278, 1837; Canad. Entom. 1878, 213.—N. States, Can.

uniformis Heid., Proc. Ent. Soc. Wash. vi, 231, 1904.—Va., Mass., Pa.

Mezirini.

ANEURUS.

Curtis, Brit. Ent. pl. 86, 1825.

fiskei Heid., Proc. Ent. Soc. Wash. vi, 164, 1904.—E. States.

inconstans Uhler, Proc. Bost. Soc. Nat. Hist. xiv, 105, 1871; xix, 420, 1878.—Can., Mass.

minutus Bergr., Verh. zool.-bot. Ges. Wien. 1886, 58.—S. States.

politus Say, Het. New Harm. 30, 1832; Compl. Writ. i, 354, 1859 (*Aradus*).—Fla.

septentrionalis Walk., B. M. Cat. Hem. Het. vii, 30, 1873.—B. Am.

simplex Uhler, Proc. Bost. Soc. Nat. Hist. xix, 421, 1878.—U. S.

CALISIUS.

Stål, Rio Hem. i, 67, 1858.

pallipes Stål, Rio Jan. Hem. i, 67, 1858; Heid. Proc. Ent. Soc. Wash., vi, 229, 1904.—Fla.

MEZIRA.

Amyot et Serv. Hém. 305, 1843.

Brachyrhynchus Laporte, Essai, 54, 1832 (preoc.).

granulatus Say, Het. New Harm. 30, 1832; Compl. Writ. i, 353, 1859 (*Aradus*); Bull. 15 Bur. Ent. Dept. Agric. fig. 16.—U. S.

parvulus Her.-Schf. Wanz. Ins. ix, 139, fig. 956, 1853 (*Dysodius*).

mæstus Stål, Stett. Ent. Zeit. 1862, 438; Bergr., Proc. Ent. Soc. Wash. ii, 336, 1892; Champ., Biol. C.-Am. Rhynch. Het. ii, 102, 1898.—Calif., Ariz.

lobatus Say, Het. New Harm. 30, 1832; Compl. Writ. i, 354, 1859 (*Aradus*); Champ., Biol. C.-Am. Rhynch. Het. ii, 96, pl. vi, fig. 28, 1898; Osborn, Ohio Nat. iv, 40, 1903.—U. S.

NANNIUM.

Bergroth, Ent. Mo. Mag. xxxiv, 100, 1898.

pusio Heid., Proc. Ent. Soc. Wash. xi, 189, 1910.—Ohio.

NEUROCTENUS.

Fieber, Europ. Hem., 34, 1861.

elongatus Osborn, Ohio Nat. iv, 41, 1903; Heid., Proc. Ent. Soc. Wash. vi, 163, 1904.—Ohio, Pa.

hopkinsi Heid., Proc. Ent. Soc. Wash. vi, 163, 1904.—N. Car.

ovatus Stål, Stett. Ent. Zeit. 1862, 439; Bergr., Öfv. Finsk Vet. Soc. Förh. xxix, 183, 1887; Osborn, Ohio Nat. iv, 41, 1903.—E. States.

- pseudonymus** Bergr., Wien. Ent. Zeit. 1898, 27.—E. States.
simplex Uhler, Bull. U. S. Geol. and Geog. Surv. Terr. ii, No. 5, 323,
 1876 (*Brachyrhynchus*); Bergroth, Öfv. Finsk. Vet. Soc.
 Förh. xxix, 182, 1887; Osborn, Ohio Nat. iv, 40, 1903.—U. S.

PICTINUS.

- Stål, Enum. Hem. iii, 140, 1873.
aurivillii Bergr., Rev. d'Entom. 1887, 247.—S. States.

PROXIUS.

- Stål, Enum. Hem. iii, 139, 1873.
schwarzii Heid., Proc. Ent. Soc. Wash. vi, 230, 1904.—Fla.

TINGITIDÆ.*Tingitini.***ACALYPTA.**

- Westwood, Mod. Class. Ins. ii, Syn. Gen. 121, 1840.
thomsonii Stål, Enum. Hemipt. iii, 122, 1873.—S. Car.

AGRAMMODES.

- Uhler, Hem. Colo. 56, 1895.
costatus Uhler, Hem. Colo. 56, 1895.—Colo.

ACANTHOCHILA.

- Stål, Rio Hem. i, 61, 1858.
exquisita Uhler, Proc. Ent. Soc. Wash. i, 143, 1889.—Fla.

ACYSTA.

- Champ., Biol. C.-Am. Rhynch. Het. ii, 46, 1898.
perseæ Heid., Proc. Ent. Soc. Wash. x, 103, 1908.—Fla.

ATHEAS.

- Champion, Biol. C.-Am. Rhynch. Het. ii, 44, 1898.
austroriparius Heid., Bull. Buffalo Soc. Nat. Sci. ix, 235, 1909.—
 Tex., Fla.
exiguus Heid., Bull. Buffalo Soc. Nat. Sci. ix, 233, 1909.—Fla.
insignis, Heid., Bull. Buffalo Soc. Nat. Sci. ix, 232, 1909.—D. C., Md.
mimeticus Heid., Bull. Buffalo Soc. Nat. Sci. ix, 234, 1909.—Wisc.,
 Minn., Kans.

CORYTHUCA.

- Stål, Enum. Hem. iii, 119, 1873.
arcuata Say, Hem. New Harm. 27, 1832; Compl. Writ. i, 350, 1859
 (*Tingis*).—E. States.
cæolata Uhler, Proc. Calif. Acad. Sci. (2) iv, 279, 1894.—Calif.
ciliata Say, Hem. New Harm. 26, 1832; Compl. Writ. i, 348, 1859
 (*Tingis*); Morrill, Psyche x, 133, 1903.—E. States.

- cratægi** Morrill, Psyche x, 132, 1903.—N. States.
cydoniæ Fitch, Count. Gentl., 14 Febr., 1861, 25.—N. Y.
decnes Stål, Stett. Ent. Zeit. 1862, 324.—W. States.
floridana Heid., Bull. Buffalo Soc. Nat. Sci. ix, 236, 1909.—Fla.
fuscigera Stål, Stett. Ent. Zeit. 1862, 323 (*Tingis*); Enum. Hem. iii, 122, 1873.—W. States.
gossypii Fabr., Ent. Syst. iv, 78, 1794 (*Acanthia*); Syst. Rhyng. 126, 1803; Stål, Hem. Fabr. i, 93, 1868; Enum. Hem. iii, 123, 1873; Her.-Schfr., Wanz. Ins. v, 85, 1839, fig. 534.—S. States.
hispida Uhler, Proc. Calif. Acad. Sci. (2) iv, 279, 1894.—Calif.
hyalina Her.-Schfr., Wanz. Ins., v, 84, 1839, fig. 532 (*Tingis*); Fieber, Ent. Monog. 103, 1844.—S. States.
incurvata Uhler, Proc. Calif. Acad. Sci. (2) iv, 280, 1894.—Ariz., Calif.
juglandis Fitch, Third Rept. Ins. N. Y., 466, 1856 (*Tingis*).—E. States.
marmorata Uhler, Proc. Bost. Soc. Nat. Hist. xix, 415, 1878 (*Tingis*).—E. States.
pergandei Heid., Proc. Ent. Soc. Wash. viii, 10, 1906.—E. States.
polygraphia Lintn., Fourth N. Y. Rep. 109, 1888.—N. Y., Md.

DICHROCYSTA.

- Champion, Biol. C.-Am. Rhynch. Het. ii, 33, 1898.
pictipes Cham., Biol. C.-Am. Rhynch. Het. ii, 34, pl. iii, fig. 1, 1898.—Ariz.

GALEATUS.

- Curtis, Entom. Mag. 196, 1833.
peckhami Ashm., Ent. Amer. iii, 156, 1888 (*Sphærocista*); Van Duzee, Canad. Ent. 1889, 5; Uhler, Proc. U. S. Nat. Mus. xxvii, 362, 1904.—N. States.

GARGAPHIA.

- Stål, Stett. Ent. Zeit. 1862, 324.
amorphæ Walsh, Proc. Amer. Ent. Soc. iii, 409, 1864 (*Tingis*).—Ill.
angulata Heidemann, Can. Entom. 1899, 301; Bull. 23 Div. Entom., fig. 9.—E. States.
fasciata Stål, Enum. Hem. iii, 125, 1873.—Ill.
iridescens Champ., Biol. C.-Am. Rhynch. Het. ii, 10, 1898, pl. ii, fig. 1.—Tex.
opacula Uhler, Death Valley Exp., Amer. Fauna ii, 263, 1893.—Calif.
tiliæ Walsh, Proc. Amer. Ent. Soc. ii, 408, 1864 (*Tingis*).—E. States.

LEPTOBYRSA.

- Stål, Enum. Hem. iii, 119, 1873.
explanata Heid., Proc. Ent. Soc. Wash. x, 105, 1908.—E. States.

LEPTOPHYA.

Stål, Enum. Hem. iii, 121, 1873.

mutica Say, Hem. New Harm. 26, 1832; Compl. Writ. i, 349, 1859 (*Tingis*).—U. S.

LEPTOSTYLA.

Stål, Enum. Hem. iii, 120, 1873.

oblonga Say, Journ. Acad. Phil. iv, 325, 1825; Compl. Writ. ii, 248, 1859 (*Tingis*).—U. S.

MONANTHIA.

Lepelletier et Serville, Encycl. Meth. x, 653, 1825.

labeculata Uhler, Death Valley Expd., Amer. Fauna, ii, 264, 1893.—Calif.

tabida Her.-Schfr., Wanz. Ins. v, 86, 1839, fig. 535; Fieber, Ent. Monog. 70, 1844.—S. States.

PHYSATOCHILA.

Fieber, Europ. Hem. 120, 1861.

plexa Say, Hem. New Harm. 26, 1832; Compl. Writ. i, 349, 1856 (*Tingis*); Stål, Enum. Hem. iii, 129, 1873.—E. States.

TELEONEMIA.

Costa, Ann. Mus. Zool. ii, 144, 1864.

Amaurosterphus Stål, Hem. Fabric. i, 92, 1868.

belfragei Stål, Enum. Hem. iii, 132, 1873; Van Duzee, Bull. Buffalo Soc. Nat. Hfst. ix, 173, 1909.—S. States.

nigrina Champ., Biol. Cent.-Amer. Rhynch. Het. ii, 41, 1898.—Tex.

sacchari Fabricius, Ent. Syst. iv, 77, 1794 (*Acanthia*); Syst. Rhynch. 126, 1803; Fieber, Ent. Monog. 76, 1844; Stål, Hem. Fabric. i, 92, 1868; Enum. Hem. iii, 132, 1873.—S. States.

scrupulosa Stål, Enum. Hemipt. iii, 132, 1873; Champion, Biol. C.-Am. Rhynch. ii, 35, 1898.—Tex.

TINGIS.

Fabricius, Syst. Rhynch. 124, 1803.

Lasiacantha Stål, Enum. Hem. iii, 130, 1873.

Melanorhopala Stål, *ibid.*

clavata Stål, Enum. Hem. iii, 130, 1873 (*Melanorhopala*).—E. States.
henshawi Ashm., Can. Ent. 1888, 20 (*Cantacader*).

lurida Stål, Enum. Hem. iii, 130, 1873 (*Melanorhopala*).—N. States.

uniformis Stål, Enum. Hem. iii, 130, 1873 (*Melanorhopala*).—N. States.

Piesmini.

PIESMA.

Lepelletier et Serville, Encycl. Meth. x, 653, 1825.

Zosmenus Laporte, Essai, 49, 1832.

Aspidotoma Curtis, Ent. Mag. i, 196, 1833.

cinerea Say, Hem. New Harm. 26, 1832 (*Tingis*); Compl. Writ. i, 349; 1859; Stål, Enum. Hem. iii, 116, 1873.—U. S.

LYGÆIDÆ.

Blissini.

BLISSUS.

Burmeister, Handb. ii, 290, 1835.

hirtus Mont., Ann. Soc. Ent. Belg. xxxvii, 405, 1893.—Pa.

leucopterus Say, Het. New Harm. 14, 1831; Compl. Writ. i, 329, 1859 (*Lygæus*); Signoret, Ann. Soc. Ent. France, 1857, 31 (*Micropus*); Webster, Bull. 15, Div. Entom. Dept. Agric.; ibid, Bull. 69, Bur. Entom.—U. S.

ISCHNODEMUS.

Fieber, Europ. Hemipt. 44, 1861.

Micropus Spinola, Essai, 218, 1840 (preoc.).

badius Van Duzee, Bull. Buffalo Soc. Nat. Hist. ix, 168, 1909.—Fla.

conicus Van Duzee, Ent. News, 1909, 234.—Tex.

falicus Say, Het. New Harm. 15, 1831; Compl. Writ. i, 331, 1859 (*Lygæus*); Signoret, Ann. Soc. Ent. France, 1857, 27; Stål, Enum. Hem. iv, 131, 1874.—U. S.

punctatus Prov., Nat. Can. iv, 76, 1874 (*Rhyparochromus*).

lobatus Van Duzee, Bull. Buffalo Soc. Nat. Hist. ix, 169, 1909.—Fla.

rufipes Van Duzee, Bull. Buffalo Soc. Nat. Hist. ix, 167, 1909.—Fla.

slossoni Van Duzee, Ent. News, 1909, 233.—Fla.

Cymini.

ARPHNUS.

Stål, Enum. Hem. iv, 125, 1874.

coriaceipennis Stål, Freg. Eug. Resa, Hem. 247, 1859; Enum. Hem. iv, 126, 1874.—W. States, Calif.

CYMODEMA.

Spinola, Essai Hemipt, 213, 1840.

exiguum Horvath, Ann. Mus. Nat. Hung. vi, 559, 1908.—D. C.

tabida Spinola Essai Hem. 213, 1840; Fieber, Europ. Hemipt. 204, 1861.—U. S.

CYMUS.

Hahn., Wanz. Ins. i, 76, 1831.

angustatus Stål, Enum. Hem. iv, 126, 1874.—U. S.

bellus Van Duzee, Bull. Buffalo Soc. Nat. Hist. ix, 167, 1909.—Fla.

breviceps Stål, Enum. Hem. iv, 127, 1874.—U. S.

discors Horvath, Ann. Mus. Nat. Hung. vi, 559, 1908.—N. States.

tabidus Prov., Faun. Can. Hem. 72, 1890 (nec Spin.).

luridus Stål, Enum. Hem. iv, 126, 1874.—U. S.

ISCHNORHYNCHUS.

Fieber, Europ. Hemipt. 51, 1861.

championi Dist., Biol. C.-Am. Rhynch. Het. i, 193, 1882; pl. xix, fig. 3.—Tex.

geminatus Say, Het. N. Harm. 14, 1831; Compl. Writ. i, 330, 1859 (*Lygæus*).—U. S.

franciscanus Stål, Freg. Eug. Resa, Hem. 252, 1859 (*Cymus*).

didymus Prov., Faun. Ent. Can. Hem. 74, 1890 (nec Zett.).

resedæ Panzer, Faun. Germ. xl, 20, 1797 (*Cimex*).—N. States.

didymus Zett., Vet. Akad. Handl. 1819, 71; Stål, Enum. iv, 124, 1874.

salvini Dist., Biol. C.-Am. Rhynch. Het. i, 194, 1882; pl. xix, fig. 2.—Tex.

NINUS.

Stål, Freg. Eugenies Resa, Hem. 252, 1859.

notabilis Distant, Biol. C.-Am. Rhynch. Het. i, 191, 1882, pl. xix, fig. 4.—Fla.

Oxycarenini.

CROPHIUS.

Stål, Enum. Hem. iv, 141, 1874.

angustatus Van Duzee, Bull. Buffalo Soc. Nat. Hist. ix, 395, 1910.—Calif., Utah.

bohemani Stål, Freg. Eug. Resa, Hem. 251, 1859 (*Cymus*); Enum. Hem. iv, 142, 1874; Van Duzee, Bull. Buffalo Soc. Nat. Hist. ix, 394, 1910.—W. States.

disconotus Say, Het. New Harm. 14, 1831 (*Lygæus*); Compl. Writ. i, 330, 1859; Van Duzee, Bull. Buffalo Soc. Nat. Hist. ix, 392, 1910.—U. S.

heidemanni Van Duzee, Bull. Buffalo Soc. Nat. Hist. ix, 393, 1910.—Ariz.

impressus Van Duzee, Bull. Buffalo Soc. Nat. Hist. ix, 396, 1910.—Calif.

schwarzi Van Duzee, Bull. Buffalo Soc. Nat. Hist. ix, 392, 1910.—Ariz.

OXYCARENUS.

Fieber, Europ. Hem. 53, 1861.

scabrosus Uhler, Proc. U. S. Nat. Mus. xxvii, 353, 1904; Van Duzee, Bull. Buffalo Soc. Nat. Hist. ix, 397, 1910.—N. Mex.

Pachygronthini.

GEDANCALA.

Amyot et Serville, Hemipt. 248, 1843.

crassimana Fabr., Syst. Rhyng. 233, 1803; Stål, Enum. iv, 139, 1874.—S. States.

cubana Stål, Enum. Hem. iv, 139, 1874.—Tex.

dorsalis Say, Het. New Harm. 17, 1831; Compl. Writ. i, 335, 1859 (*Pamera*).—U. S.

dorsilinea Amyot et Serv., Hem. 258, 1843; Stål, Enum. Hem. iv, 139, 1874; Van Duzee, Bull. Buffalo Soc. Nat. Hist. ix, 170, 1909.

PACHYGRONTHA.

Germar, Silb. Rev. Ent. v, 152, 1837.

Actractophora Stål, Cefv. Vet. Akad. Förh. 1853, 260.

Peliosoma Uhler, Proc. Acad. Nat. Sci. Phil., 1860, 229.

bimaculata Distant, Biol. C.-Am. Rhynch. Het. i, 393, pl. xxxiv, fig. 23, 1893.—Tex.

Heterogastrini.

HETEROGASTER.

Schilling, Beitr. Ent. 84, 1829.

Phrygadicus Fieb., Europ. Hem. 201, 1861.

behrensi Uhler, Bull. Geol. and Geog. Surv. Terr. ii, 312, 1876.—W. States.

PHLEGYAS.

Stål, Hemipt. Afr. ii, 145, 1865.

Helenotus Uhler, Bull. U. S. Geol. Surv. ii, 312, 1876.

Peliopelta Uhler, Check List Hem. 15, 1886.

annulicornis Stål, Berl. Ent. Zeit. 1869, 230; Enum. iv, 138, 1874.—U. S.

abbreviata Uhler, Bull. Geol. and Geog. Surv. Terr. ii, 313, 1876 (*Helenotus*).

tropicalis Distant, Biol. C.-Am. Rhynch. Het. i, 411, 1893, pl. xxxvi, fig. 5.—Tex.

Lygæini.

BELONCHILUS.

Uhler, Proc. Bost. Soc. Nat. Hist. xiv, 104, 1871.

koreshanus Van Duzee, Bull. Buffalo Soc. Nat. Hist. ix, 165, 1909.—Fla.

numenius Say, Het. New. Harm. 15, 1831 (*Lygæus*); Compl. Writ. i, 331, 1859.—U. S.

LYGÆOSOMA.

Spinola, Essai Hem. 254, 1840.

solida Uhler, Death Valley Expd., N. Am. Fauna, ii, 262, 1893.—Calif.

LYGÆUS.

Fabricius, Ent. Syst. iv, 133, 1794.

Melanocoryphus Stål, Cefv. Vet. Akad. Förh. 1872, 41.

Melanopleurus Stål, Enum. Hem. iv, 105, 1874.

Ochrostomus Stål, *ibid*.

Graptolomus Stål, Hem. Fabric. i, 73, 1868.

- admirabilis** Uhler, Hayden's Surv. Mont. 405, 1872.—Colo.
- albulus** Dist., Biol. C.-Am. Rhynch. Het. i, 380, 1893, pl. xxxiv, fig. 2.
—Tex.
- belfragei** Stål, Enum. Hem. iv, 109, 1874.—Tex.
- bicolor** H. Schf., Wanz. Ins. 195, 1853; Stål, Enum. Hem. iv, 109, 1874; Dist., Biol. C.-Am. Rhynch. Het. i, 180, pl. xvi, fig. 17, 1882.—W. States.
- bipartitus* Sign., Ann. Soc. Ent. France, 1861, 947.
- dimidiatus* Walk., B. M. Cat. Het. v, 45, 1872.
- bicrucis** Say, Journ. Acad. Philad. iv, 322, 1825; Compl. Writ. ii, 246, 1859; Stål, Enum. Hem. iv, 113, 1874.—U. S.
- flavomarginellus* Stål, Freg. Eug. Resa, Hem. 241, 1859.
- bitransversus* Sign., Ann. Soc. Ent. France, 1860, 947.
- bistriangularis** Say, Het. New Harm. 13, 1831; Compl. Writ. i, 329, 1859; Uhler, Hayden's Surv. Mont. 405, 1872; Stål, Enum. Hem. iv, 109, 1874.—W. States.
- marginellus* Dallas, List Hem. B. M. ii, 548, 1852.
- vicinus* Dallas, *ibid*, 549.
- californicus** Walk., B. M. Cat. Het. v. 42, 1872.—Calif.
- circumlitus** Stål, Stett. Ent. Zeit. 1862, 309; Enum. Hem. iv, 113, 1874; Biol. C.-Am. Rhynch. Het. i, 186, pl. xvi, fig. 13, 1882.—W. States,
- circumcinctus* Uhler, Hem. Colo. 27, 1895.
- costalis** Herr.-Schæffer, Wanz. Ins. vii, 22, 1844; *ibid*, ix, 195, 1853; Stål, Enum. iv, 107, 1874.—W. States.
- facetus** Say, Het. New Harm. 13, 1831; Compl. Writ. i, 328, 1859; Stål, Enum. iv, 113, 1874.—Fla., Tex.
- kalmii** Stål, Enum. iv, 107, 1874; Uhler, Bull. Geol. and Geogr. Surv. Terr. ii, 302, 1876; Montd., Ann. Soc. Ent. Belg. 1893, 399; Dist., Biol. C.-Am. Rhynch. Het. i, pl. xvii, fig. 2.—U. S.
- var. *melanodemus* Montd., *ibid*, 400, 1893.
- lateralis** Dallas, List, ii, 550, 1852; Stål, Enum. iv, 113, 1874; Dist., Biol. C.-Amer. Rhynch. Het. i, pl. xviii, fig. 1.—Tex., Colo., Ariz.
- lineola** Dallas, List, ii, 549, 1852; Stål, Enum. iv, 110, 1874.—Tex., Fla.
- melanopleurus** Uhler, Death Valley Exped., N. Am. Fauna, ii, 262, 1893.—W. States.
- mimulus** Stål, Enum. Hem. iv, 113, 1874.—Tex.
- obscuripennis** Stål, Enum. Hem. iv, 112, 1874.—Tex.
- pusio** Stål, Enum. Hem. iv, 112, 1874.—Tex.
- pyrrhopterus** Stål, Enum. Hem. iv, 110, 1874; Dist., Biol. Cent.-Am. Rhynch. Het. i, pl. xvii, fig. 6, 1882.—Tex.
- reclivatus** Say, Journ. Acad. Phila. iv, 321, 1825; Compl. Writ., i, 329, 1859; Uhler, Bull. Geol. and Geogr. Surv. Terr. ii, 302, 1876; Stål, Enum. iv, 107, 1874.—U. S.
- var. *enotus* Say, Het. New Harm. 13, 1831; Compl. Writ. i, 329, 1859.

rubricollis Uhler, Proc. Calif. Acad. Sci. (2) iv, 244, 1894.—Calif.

rubriger Stål, Stett. Ent. Zeit. 1862, 309; Enum. Hem. iv, 113, 1874; Dist., Biol. C.-Amer. Rhynch. Het. i, pl. xvii, fig. 12.—S. States.

truculentus Stål, Stett. Ent. Zeit. 1862, 308; Enum. iv, 108, 1874; Dist., Biol. C.-Am. Rhynch. Het. i, pl. xvii, fig. 4.

turcicus Fabr., Syst. Rhyng. 218, 1803; Stål, Enum. iv, 107, 1874; Bull. 69 Minn. Sta., fig. 61.—U. S.

uhleri Stål, Enum. Hem. iv, 109, 1874; Barber, Bull. Brookl. Ins. i, 274, 1906; Dist., Biol. C.-Am. Rhynch. Het. i, pl. xvi, fig. 12.—Tex., Cal.

NYSIUS.

Dallas, List, ii, 331, 1852.

Ortholomus Stål, Cefv. Vet. Akad. Förh. 1872, 43.

arphnoides Baker, Invert. Pacifica, i, 140, 1906 (*Ortholomus*).—Calif.

californicus Stål, Freg. Eug. Resa, Hem. 242, 1859; Enum. iv, 120, 1874; Baker, Invert. Pac. i, 135, 1906.—Tex., Calif.

coloradensis Baker, Invert. Pacifica, i, 136, 1906.—Colo.

destructor Riley, Fifth Rept. Ins. Missouri, 113, 1873, fig. 41.—Mo.

ericæ Schill., Beit. Ent. i, 86, 1829; Flor. Rhyn. Livl. i, 292, 1860; Horvath, Ann. Mus. Nat. Hung. vi, 558, 1908.—U. S.

angustatus Uhler, Hayden's Surv. Mont. 406, 1872; Baker, Invert. Pac. i, 135, 1906.

raphanus Howard, Can. Ent. iv, 219, 1872.

grønlandicus Zett., Ins. Lapp. 262, 1840 (*Lygæus*); Stål, Enum. Hem. iv, 121, 1874.—Labr., Alaska.

inæqualis Uhler, Proc. Zool. Soc. Lond. 1894, 183.—Fla.

longiceps Stål, Enum. iv, 120, 1874; Baker, Invert. Pac. i, 139, 1906.—U. S.

var. *cookii* Baker, *ibid.*

minutus Uhler, Hem. Colo. 22, 1895 (var. of *angustatus* acc. to Baker).—Colo.

monticola Dist., Biol., C.-Am. Rhynch. Het. i, 385, 1893, pl. xxxiv, fig. 8.

nevadensis Baker, Invert. Pacifica, i, 139, 1906 (*Ortholomus*).—Nev.

strigosus Uhler, Proc. Calif. Acad. Sci. (2), liv, 239, 1894.—W. States.
senecionis Baker, Invert. Pac. i, 135, 1906 (nec Schilling).

thymi Wolff, Icon. Cim. 147, pl. xv, fig. 143, 1804 (*Lygæus*); Fieber, Eur. Hem. 169, 1861—N. States.

saint-cyri Prov., Nat. Can. iv, 77, 1872.

uhleri Baker, Invert. Pacifica i, 139, 1906 (*Ortholomus*).—Wisc.

ONCOPELTUS.

Stål, Hemipt. Fabr. i, 75, 1868.

Erythriscius Stål, Enum. Hem. iv, 102, 1874.

fasciatus Dallas, List Hem. B. M. ii, 538, 1852; Stål, Enum. iv, 103, 1874.—U. S.

aulicus Her.-Schf., Wanz. Ins. vi, 76, 1842 (nec Fabr.).

gutta Her.-Schf., Wanz. Ins. vii, 20, 1844; *ibid*, ix, 200, 1853; Stål, Enum. iv, 101, 1874; Dist., Biol. C.-Am. Rhynch. Het. i, pl. xvi, fig. 23.—Calif.

sandarachatus Say, Het. New Harm. 13, 1831 (*Lygæus*); Compl. Writ. i, 328, 1859; Stål, Enum. iv, 103, 1874.—Calif.

varicolor Fabr., Ent. Syst. iv, 149, 1794 (*Lygæus*); Syst. Rhynch. 219, 1803; Dallas, List Hem. B. M. ii, 538, 1852; Stål, Enum. iv, 102, 1874; Dist., Biol. C.-Am. Rhynch. Het. i, pl. xvi, fig. 21.—Calif.

ORSILLUS.

Dallas, List Hem. B. M. ii, 551, 1852.

Mecorhamphus Fieb., Europ. Hem. 173, 1861.

scolopax Say, Het. New Harm. 14, 1831 (*Lygæus*); Compl. Writ. i, 380, 1859.—W. States.

Myodochini.

CARPILIS.

Stål, Enum. Hem. iv, 153, 1874.

ferruginea Stål, Enum. Hem. iv, 153, 1874.—Tex.

CISTALIA.

Stål, Enum. Hem. iv, 164, 1874.

signoretii Guérin, Sagra's Hist. Cuba, Ins. vii, 165, 1857.—Tex.

CLIGENES.

Distant, Biol. C.-Am. Rhynch. Het. i, 405, 1893.

Pygæus Uhler, Proc. Zool. Soc. Lond. 1894, 187.

Salacia Stål, Enum. iv, 156, 1874 (preoc.).

Antillocoris Kirk., Entom. 1904, 280.

Botocudo Kirk., *ibid*.

pallidus Uhler, Proc. Zool. Soc. Lond. 1894, 187 (*Pygæus*).—E. States.
minutus Bergr., Rev. d'Ent. 1895, 144.

pilosula Stål, Enum. iv, 158, 1874 (*Salacia*).—Tex.

punctata Dist., Biol. C.-Am. Rhynch. Het. i, 406, 1893, pl. xxxv, fig. 23 (*Salacia*).—Tex.

CNEMODUS.

Herr.-Schæffer, Wanz. Ins. ix, 184, 1853.

mavortius Say, Het. New Harm. 19, 1831; Compl. Writ. i, 337, 1859 (*Astemma*).—U. S.

brevipennis Herr.-Schæffer, Wanz. Ins., ix, 184, 1853.

alacris Walk., B. M. Cat. Het. v, 147, 1872 (*Thaumastopus*?).

sobrius Uhler, Proc. Calif. Acad. Sci. (2), iv, 241, 1894.—Calif.

CRYPHULA.

Stål, Enum. Hem. iv, 164, 1874.

parallelogramma Stål, Enum. Hem. iv, 165, 1874.—Tex.

DELOCHILOCORIS.

Bergroth, Rev. d'Entom. xii, 154, 1893.

Dorachosa Dist., Biol. C.-Am. Rhynch. Het. i, 409, 1893 (pre-occupied).

illuminatus Dist., Biol. Cent.-Amer. Rhyng. i, 409, 1893 (*Dorachosa*), pl. xxxvi, fig. 1.—W. States.

umbrosus Dist., Biol. C.-Am. Rhynch. Het. i, 409, 1893 (as var. *Dorachosa illuminata*); Horvath, Ann. Mus. Nat. Hung. vi, 561, 1908 (*Aphanus*).—W. States.

DRYMUS.

Fieber, Europ. Hem. 46, 1861.

Lampronotus Dougl. and Scott, Ent. Mo. Mag. iv, 243, 1868.

Lamproplex Dougl. and Scott, ibid, 265.

unus Say, Het. New Harm. 16, 1831; Compl. Writ. i, 332, 1859 (*Pamera*);—U. S.

DYCODERUS.

Uhler, Proc. Ent. Soc. Wash. iv, 507, 1901.

picturatus Uhler, Proc. Ent. Soc. Wash. iv, 508, 1901.—Ariz.

EMBLETHIS.

Fieber, Europ. Hemipt. 51, 1861.

griseus Wolff, Icon. Cim. 113, fig. 107, 1802.

arenarius Fieber, Europ. Hemipt. 198, 1861; Bull. 69 Minn. Sta., fig. 63.—N. States.

vicarius Horvath, Ann. Mus. Nat. Hung. vi, 563, 1908.—N. States.

arenarius Prov., Faun. Ent. Can. Hem. 77, 1890 (nec Fieber).

EREMOCORIS.

Fieber, Europ. Hemipt. 49, 1861.

ferus Say, Het. New. Harm. 16, 1831 (*Pamera*); Compl. Writ. i, 333, 1859; Horvath, Rev. d'Ent. ii, 9, 1883.—U. S.

borealis Dallas, List, ii, 565, 1852 (*Rhyparochromus*).

obscurus Van Duzee, Ent. News, 1906, 388.—Vanc.

tropicus Distant, Biol. C.-Am. Rhynch. Het. i, 218, pl. xx, fig. 7, 1882.—W. States.

GASTRODES.

Westwood, Mod. Class. Ins. ii, Syn. gen. 122, 1840.

Platygaster Schill., Beitr. Ent. i, 82, 1829 (preoc.).

ferrugineus Linné, Syst. Nat. (X), 730, 1758; Fieber, Europ. Hemipt. 187, 1861.—N. States.

pacificus Prov., Faun. Ent. Can. Hem. 204, 1890 (*Platygaster*).—Vanc.

GONIANOTUS.

Fieber, Europ. Hemipt. 51, 1861.

marginepunctatus Wolff, Icones Cim., 148, fig. 144, 1804; Fieber, Europ. Hemipt. 197, 1861; Uhler, Proc. Boston Soc. Nat. Hist. xix, 392, 1878.—N. States.

HERÆUS.

Stål, Stett. Ent. Zeit. xxiii, 314, 1862.

plebejus Stål, Enum. iv, 147, 1874.—U. S.

triguttatus Guérin, Sagra's Hist. de Cuba, vii, Ins. 167, 1857 (*Lygæus*)—Fla.

LIGYROCORIS.

Stål, Öfv. Vet. Akad. Förh. 1872, 51.

abdominalis Guérin, Sagra's Hist. de Cuba, vii, Ins., 165, 1857 (*Lygæus*); Stål, Enum. Hem. iv, 146, 1874; Dist., Biol. C.-Am. Rhynch. Het. i, pl. xvii, fig. 21.—Fla.

piliger Stål, Stett. Ent. Zeit. 1862, 312.

balteatus Stål, Enum. Hem. iv, 145, 1874; Dist., Biol. C.-Am. Rhynch. Het. i, 202, pl. xvii, fig. 20, 1882.—Calif.

constrictus Say, Het. New Harm. 15, 1831 (*Pamera*); Compl. Writ. i, 332, 1859; Stål, Enum. iv, 146, 1874; Heid., Proc. Wash. Ent. Soc. v, 156, 1903.—E. States.

contractus Say, Het. New Harm. 15, 1831 (*Pamera*); Compl. Writ. i, 332, 1859.—E. States.

sylvestris Fieber, Eur. Hem. 171, 1861.

costalis Van Duzee, Can. Ent. 1909, 373 (*Perigenes*).—E. States.

diffusus Uhler, Proc. Bost. Soc. Nat. Hist. xiv, 101, 1871 (*Plociomera*).—U. S.

sylvestris Stål, Enum. Hem. iv, 145, 1874 (nec Fieber).

litigiosus Stål, Stett. Ent. Zeit. 1862, 313 (*Plociomera*); Enum. Hem. iv, 145, 1874; Dist., Biol. C.-Am. Rhynch. Het. i, pl. xvii, fig. 19, 1882—Fla.

multispinus Stål, Enum. Hem. iv, 145, 1874; Dist., Biol. C.-Am. Rhynch. Het. i, pl. xvii, fig. 18, 1882; Van Duzee, Bull. Buffaló Soc. Nat. Sci. Hist. ix, 171, 1909.—Fla.

nitidula Uhler, Death Valley Exped., N. Am. Fauna, ii, 262, 1893 (*Pamera*).—Calif.

pseudoheræus Barber, Bull. Brookl. Inst. i, 275, 1906.—Tex.

MICROTOMA.

Laporte, Essai Hemipt. 33, 1832.

atrata Goeze, Beytr. ii, 268, 1778.—U. S.

carbonaria Rossi, Faun. Etr., No. 1330, 1790; Fieber, Europ. Hemipt. 193, 1861.

MYODOCHA.

Latreille, Gen. Ins. iii, 126, 1807.

Chiroleptes Kirby, Fauna Bor. Amer. iv, 281, 1837.

- serripes** Oliv., Ency. Meth. viii, 106, 1811 (*Lygæus*); Her.-Schf., Wanz. Ins. ix, 213, 1853, fig. 966; Bull. 69 Minn. Sta., fig. 64.—U. S. *petiolata* Say, Het. New Harm. 19, 1831; Compl. Writ. i, 337, 1859. *raptor* Kirby, Faun. Bor. Amer. iv, 281, 1837; Can. Entom. 1878, 214 (*Chiroleptes*).

OZOPHORA.

Uhler, Proc. Bost. Soc. Nat. Hist. xiv, 102, 1871.

burmeisteri Guérin, Sagra's Hist. de Cuba, vii, Ins., 165, 1857; Stål, Enum. Hem. iv, 156, 1874.—Tex.

- picturata** Uhler, Proc. Bost. Soc. Nat. Hist. xiv, 102, 1871; ibid, xix, 390, 1878.—U. S.

unicolor Uhler, Proc. Calif. Acad. Sci. (2), iv, 242, 1894.—Calif.

PAMERA.

Say, Het. New Harm. 15, 1831.

Orthæa Dall., List, Hem. ii, 532, 1852.

Diplonotus Stål, Cefv. Vet. Akad. Förh. 1872, 51.

- basalis** Dallas, List Hem. ii, 575, 1852 (*Rhyparochromus*); Stål, Enum. Hem. iv, 152, 1874.—U. S.

bilobata Say, Het. New Harm. 17, 1831; Compl. Writ. i, 334, 1859; Stål, Enum. Hem. iv, 150, 1874; Distant, Biol. C.-Am. Rhynch. Het. i, pl. xvii, fig. 25.—U. S.

scutellata Dallas, B. M. List Hem. ii, 575, 1852.

curvipes Stål, Enum. Hem. iv, 148, 1874.—Tex., S. Car.

lineata Dallas, B. M. List, Hem. ii, 575, 1852 (*Rhyparochromus*).—S. States.

nitidicollis Stål, Enum. Hem. iv, 150, 1874.—Tex.

setosa Stål, Enum. Hem. iv, 150, 1874.—Tex.

vicina Dallas, B. M. List Hem. ii, 576, 1852 (*Rhyparochromus*).—S. States.

vincta Say, Het. New Harm. 16, 1831; Compl. Writ. i, 333, 1859.—U. S.

parvula Dallas, List, Hem. B. M. ii, 576, 1852 (*Rhyparochromus*); Dist., Biol. C.-Am. Rhynch. Het. i, 206, pl. xix, fig. 12, 1882.

amyoti Guérin, Sagra's Hist. de Cuba, vii, Ins. 166, 1857.

PAROMIUS.

Fieber, Europ. Hemipt. 45, 1861.

- longulus** Dallas, B. M. List, Hem. ii, 578, 1852; Stål, Enum. iv, 148, 1874; Dist., Biol. C.-Am. Rhynch. Het. i, 206, pl. xvii, fig. 26, 1882.—S. States.

PERIGENES.

Distant, Biol. Cent.-Amer. Rhynch. Het. i, 396, 1893.

fallax Heid., Proc. Ent. Soc. Wash. v, 156, 1903.—E. States.

similis Barber, Bull. Brookl. Inst. i, 276, 1906.—Tex.

PERITRECHUS.

Fieber, Europ. Hemipt. 48, 1861.

fraternus Uhler, Proc. Bost. Soc. Nat. Hist. xiv, 103, 1871; *ibid*, xix, 391, 1878.—U. S.

tristis Van Duzee, Ent. News, 1906, 388.—Vanc.

PETISSIUS.

Distant, Biol. C.-Am. Rhynch. Het. i, 407, 1893.

diversus Dist., Biol. C.-Am. Rhynch. Het. i, 407, pl. xxxv, fig. 22, 1893.—Tex.

PSEUDOPAMERA.

Distant, Biol. C.-Am. Rhynch. Het. i, 209, 1882.

ferrerei Distant, Biol. C.-Am. Rhynch. Het. i, 399, pl. xxxv, fig. 5, 1893.—Ariz.

PTOCHIOMERA.

Say, Het. New Harm. 18, 1831.

Plociomera of various authors.

antennata Van Duzee, Bull. Buffalo Soc. Nat. Hist. ix, 172, 1909.—Florida.

clavigera Uhler, Hem. Colo. 24, 1895.—Tex., Colo.

formosa Distant, Biol. C.-Am. Rhynch. Het. i, 210, pl. xix, fig. 25, 1882 (*Plociomera*).—Tex.

fuscicornis Stål, Enum. Hem. iv, 152, 1874.—Tex.

minima Guér., Sagra's Hist. de Cuba, vii, Ins. 166, 1857 (*Beosus*); Stål, Enum. Hem. iv, 153, 1874.—S. States.

nodosa Say, Het. New Harm. 18, 1831; Compl. Writ. i, 335, 1859; Stål, Enum. Hem. iv, 153, 1874.—U. S.

clavata Dallas, List, ii, 560, 1852 (*Aphanus*).

puberula Stål, Enum. Hem. iv, 153, 1874.—Tex.

PTEROMETUS.

Amyot et Serville, Hemipt. 256, 1843.

canadensis Prov., Faun. Can. Hemipt. 84, 1886.—Can.

RHAPTUS.

Stål, Enum. Hem. iv, 163, 1874.

collina Dist., Biol. C.-Am. Rhynch. Het. i, 410, pl. xxxvi, fig. 3, 1893.—S. States.

RHYPAROCHROMUS.

Curtis, Brit. Entom. xiii, pl. 612, 1836.

compactus Uhler, Proc. U. S. Nat. Mus. xxvii, 354, 1904.—Ariz.

sodalicius Uhler, Rept. Geol. and Geog. Surv. Terr. v, 835, 1877
(*Megalonotus*).—W. States.

tripunctatus Dallas, B. M. List Hem, ii, 559, 1852 (*Aphanus*).—Fla.

SCOLOPOSTETHUS.

Fieber, Europ. Hemipt. 49, 1861.

atlanticus Horvath, Rev. d'Entom. xii, 239, 1893.—U. S.

diffidens Horvath, Rev. d'Entom. xii, 240, 1893.—N. States.

thomsoni Reuter, Finl. Skand. Hem. i, 53, 1875.—U. S.

decoratus Thom., Opusc. Ent. ii, 201, 1870 (nec Hahn.).

SPHÆROBIUS.

Uhler, Proc. Zool. Soc. Lond. 1893, 710.

insignis Uhler, Geol. Surv. Mont. 407, 1872 (*Heræus*); Stål, Enum.
Hem. iv, 147, 1874.—Colo.

SPHRAGISTICUS.

Stål, Cefv. Vet. Akad. Förh. 1872, 55.

nebulosus Fallen, Mon. Cim., 65, 1807; Fieber, Europ. Hemipt. 190,
1861; Bull. 69 Minn. Sta., fig. 62.—U. S.

fallax Say, Het. New Harm. 16, 1831 (*Pamera*); Compl. Writ.
i, 334, 1859.

TEMPYRA.

Stål, Enum. Hem. iv, 155, 1874.

biguttula Stål, Enum. Hem. iv, 157, 1874.—Tex.

TRAPEZONOTUS.

Fieber, Europ. Hemipt. 50, 1861.

caliginosus Distant, Biol. C.-Am. Rhynch. Het. i, 216, pl. xx, fig. 3,
1882.—S. W. States.

rufipes Stål, Enum. Hem. iv, 179, 1874.—Tex.

UHLERIOLA.

Horvath, Ann. Mus. Nat. Hung. vi, 562, 1908.

floralis Uhler, Hem. Colo. 26, 1895 (*Rhyparochromus*).—Colo.

XESTOCORIS.

Van Duzee, Ent. News, 1906, 389.

nitens Van Duzee, Ent. News, 1906, 390 (= *Rhaptus collina* ?).—N. Y.

Geocorini.

GEOCORIS.

Fallen, Spec. Hemipt. disp. 10, 1814.

Ophthalmicus Schilling, Beit. zur Entom. i, 62, 1829.

atricolor Montd., Bull. Soc. Sci. Bucarest, xvi, 228, 1908.—W. States.

- bullatus** Say, Het. New Harm. 18, 1831; Compl. Writ. i, 336, 1859 (*Salda*); Montd., Bull. Soc. Sci. Bucarest, xvi, 218, 1908; Bull. 69 Minn. Sta., fig. 65.—U. S.
borealis Dall., List, ii, 585, 1852.
griseus Dall., ibid.
pallens Stål, Cefv. Vet. Akad. Förh. 1854, 236; Enum. Hem. iv, 136, 1874.
- bullatus** var. **obscuratus** Montd., Bull. Soc. Sci. Bucarest, xvi, 220, 1908.—Colo.
- decoratus** Uhler, Bull. Geol. and Geogr. Surv. Terr. iii, 110, 1877; Montd., Bull. Soc. Sci. Bucarest, xvi, 221, 1908.—U. S.
- decoratus** var. **solutus** Montd., Bull. Soc. Sci. Bucarest, xvi, 223, 1908.—Colo.
- discopterus** Stål, Enum. Hem. iv, 136, 1874.—U. S.
- duzei** Montd., Bull. Soc. Sci. Bucarest, xvi, 231, 1908.—Colo.
- howardi** Montd., Bull. Soc. Sci. Bucarest, xvi, 229, 1908.—Mich.
- imperialis** Distant, Biol. C.-Am. Rhynch. Het. i, 197, 1882, pl. xviii, fig. 18; Barber, Bull. Brooklyn Inst. i, 275, 1906.—Cal., Tex.
- lividipennis** Stål, Stett. Ent. Zeit. 1862, 311; Enum. Hem. iv, 137, 1874; Dist., Biol. C.-Am. Rhynch. Het. i, 198, pl. xvii, fig. 27, 1882.—N. Mex.
- piceus** Say, Het. New Harm. 18, 1831; Compl. Writ. i, 336, 1859 (*Salda*).—U. S.
- punctipes** Say, Het. New Harm. 18, 1831 (*Salda*); Compl. Writ. i, 336, 1859; Montd., Bull. Soc. Sci. Bucarest, xvi, 224, 1908.—U. S.
luniger Fieber, Wien. Ent. Monatsch. v, 269, 1861.
- scudderi** Stål, Enum. Hem. iv, 135, 1874.—Tex.
- tristis** Stål, Cefv. Vet. Akad. Förh. 1854, 236; Freg. Eug. Resa, Ins. 249, 1859; Enum. Hem. iv, 136, 1874.—Calif.
- ♂ **uliginosus** Say, Het. New Harm. 18, 1831; Compl. Writ. i, 337 1859 (*Salda*); Montd., Bull. Soc. Sci. Bucarest, xvi, 226, 1908.—U. S.
niger Dallas, List, Hem. ii, 586, 1852 (*Ophthalmicus*).
lateralis Fieber, Wien. Ent. Monatsch. v, 271, 1861.
limbatus Stål, Enum. Hem. iv, 136, 1874.
- uliginosus** var. **speculator** Montd., Bull. Soc. Sci. Bucarest, xvi, 227, 1908.—Mo., Mass.

PYRRHOCORIDÆ.

Larginæ.

ACINOCORIS.

Hahn., Wanz. Ins. ii, 113, 1834.

lunatus Fabr., Mant. Ins. ii, 302, 1787 (*Cimex*); Stål, Enum. Hem. i, 92, 1870.—Calif.

calidus Fabr., Syst. Rhyng. 230, 1803.

interruptus H. Schf., Wanz. Ins. ix, 181, 1853, fig. 978.

ARHAPHE.

Herr. Schæffer, Wanz. Ins. ix, 183, 1853.

carolina Her.-Schf., Wanz. Ins. ix, 183, 1853.—Car.

cicindeloides Walk., Cat. Het. B. M. vi, 36, 1873; Dist., Biol. Cent.-Am. Rhynch. Het. i, pl. xxi, fig. 2, 1883.—Ariz.

LARGUS.

Hahn., Wanz. Ins. i, 13, 1831.

Euryophthalmus Laporte, Essai, 38, 1832.

Lecadra Sign., Ann. Soc. Ent. France, 1862, 582.

cinctus Her.-Schf., Wanz. Ins. vii, 6, 1844; Dist., Biol. Cent.-Am. Rhynch. Het. i, 223, pl. xx, fig. 16, 1882.—S. States.

varians Stål, Enum. Hem. i, 94, 1871.

convivus Stål, Cefv. Vet. Akad. Förh. 1861, 198; Enum. Hem. i, 121, 1870; Dist., Biol. Cent.-Amer. Rhynch. Het. i, 231, pl. xxi, figs. 11, 12, 15, 1882—Calif., Tex.

obtus Walk., Cat. Hem. B. M. vi, 33, 1873.

succinctus Linn., Cent. Ins. Rar. 17, 1763 (*Cimex*); Syst. Nat. xii, 727, 1767; Fabr., Syst. Rhynch. 233, 1803; H. Schf., Wanz. Ins. vi, 78, 1842, fig. 648; Stål, Hem. Fabric. i, 80, 1868.—S. States.

rubrocinctus De Geer, Mém. Ins. iii, 339, 1773 (*Cimex*).

STENOMACRA.

Stål, Enum. Hem. i, 90, 1870.

marginella Her. Schf., Wanz. Ins. ix, 182, 1853 (*Largus*); Stål, Enum. Hem. i, 98, 1870; Dist., Biol. C.-Am. Rhynch. Het. i, pl. xx, fig. 23.—Ariz.

*Pyrrhocorinæ.***DYSDERCUS.**

Amyot et Serville, Hem. 272, 1843.

Piezodera Sign., Ann. Soc. Ent. France, 1861, 953.

albidiventris Stål, Cefv. Vet. Akad. Förh. 1854, 236; Enum. Hem. i, 121, 1870; Dist., Biol. C.-Am. Rhynch. Het. i, 229, pl. xxi, figs. 5, 10, 1883.—Tex.

lunulatus Uhler, Proc. Ent. Soc. Phila. i, 24, 1861.

concinus Stål, Cefv. Vet. Akad. Förh. 1861, 198; Enum. Hem. i, 121, 1870; Dist., Biol. Cent.-Am. Rhynch. Het. i, 231, pl. xxi, figs. 11, 12, 1882.—S. States.

mundus Walk., Cat. Het. B. M. v, 181, 1872.

mimus Say, Het. N. Harm. 20, 1832 (*Capsus*); Compl. Writ. i, 338, 1859; Stål, Stett. Ent. Zeit. 1862, 316; Dist., Biol. Cent.-Am. Rhynch. Het. i, 229, pl. xxi, fig. 4, 1883.—Calif., Tex., Ariz.

obliquus Her. Schf., Wanz. Ins. vii, 19, 1844, fig. 701 (*Pyrrhocoris*); Stål, Enum. Hem. i, 121, 1870; Dist., Biol. C.-Am. Rhynch, Het. i, 232, pl. xxi, figs. 16, 17, 18, 1883.—Colo.

bimaculatus Stål, Cefv. Vet. Akad. Förh. 1854, 236.

suturellus Her. Schf., Wanz. Ins. vi, 76, 1842, fig. 645 (*Pyrrhocoris*); Insect Life, i, fig. 52.—Fla., Tex.

BERYTIDÆ.

ACANTHOPHYSA.

Uhler, North Amer. Fauna, vii, 261, 1893.

echinata Uhler, North Amer. Fauna, vii, 261, 1893.—Ariz.

JALYSUS.

Stål, Rio Jan. Hem. ii, 59, 1862.

multispinus Ashm., Ent. Amer. iii, 1887, 155 (*Hoplinus*).—S. States.

perclavatus Van Duzee, Bull. Buffalo Soc. Nat. Hist. ix, 163, 1909.

spinosus Say, Amer. Entom. i, 28, 1824 (*Berytus*); Compl. Writ. i, 28, 1859; Bull. 69 Minn. Sta., fig. 69.—U. S.

trispinosus Westw., Hope, Cat. Hemipt. ii, 24, 1842 (*Neides*).

wickhami Van Duzee, Ent. News, 1906, 387.—Ariz.

METACANTHUS.

Costa, Atti Acad. Napoli, 1848, 258.

Protocantha Uhler, Proc. Zool. Soc. London, 1893, 708.

decorus Uhler, Proc. Zool. Soc. London, 1893, 708 (*Protocantha*).—Fla.

NEIDES.

Latreille, Hist. Nat. iii, 246, 1802.

muticus Say, Het. New Harm. 13, 1831 (*Berytus*); Compl. Writ. i, 328, 1859.—U. S.

decurvatus Uhler, Fifth Ann. Rept. U. S. Geol. Surv. f. 1871 (1872), 402.

gracilipes Stål, Freg. Eug. Resa, Hem. 235, 1859.

PRONOTACANTHA.

Uhler, North Amer. Fauna, vii, 260, 1893.

annulata Uhler, North Amer. Fauna, vii, 260, 1893.—Calif.

COREIDÆ.

Rhopalini.

AUFEIS.

Stål, Enum. Hemipt. i, 221, 1870.

impressicollis Stål, Enum. Hemipt. i, 222, 1870; Dist., Biol. C.-Am. Rhynch. Het. i, pl. xv, fig. 20.—W. States.

CORIZUS.

Fallen, Spec. nov. Hem. 8, 1814.

Rhopalus Fieber, Europ. Hem. 232. 1861.

bohemani Sign., Ann. Soc. Ent. France, 1859, 86.—U. S.

nigristernum Sign., ibid, 100; Stål, Enum. Hem. i, 225, 1870; Hambleton, Ann. Ent. Soc. Amer. i, 143, 1908.

hyalinus Fabr., Ent. Syst. iv, 168, 1794 (*Lygæus*); Stål, Hem. Fabr. i, 68, 1868; Enum. Hem. i, 222, 1870; Hambleton, Ann. Ent. Soc. Amer. i, 136, 1908.—U. S.

indentatus Hambleton, Ann. Ent. Soc. Amer. i, 139, 1908.—W. States.

lateralis Say, Journ. Acad. Phil. iv, 320, 1825 (*Coreus*); Compl. Writ. ii, 245, 1859; Sign., Ann. Soc. Ent. France (3), vii, 97, 1859; Hambleton, Ann. Ent. Soc. Amer. i, 140, 1908.—U. S.

punctipennis Dallas, B. M. List, Hem. ii, 526, 1852.

parvicornis Sign., Ann. Soc. Ent. France, 1859, 301; Hambleton, Ann. Ent. Soc. Amer. i, 143, 1908.—Calif., Tex.

punctatus Sign., Ann. Soc. Ent. France, 1859, 81; Hambleton, Ann. Ent. Soc. Amer. i, 144, 1908.—Tex.

punctiventris Dallas, Brit. Mus. List Hemipt. ii, 526, 1852; Stål, Enum. Hemipt. i, 223. 1870—N. States.

noveboracensis Sign., Ann. Soc. Ent. France, 1859, 97.

borealis Uhler, Proc. Acad. Nat. Sci. Phil., 1861, 284.

crassicornis Linn., Hambleton, Ann. Ent. Soc. Amer. i, 137, 1908.

sidæ Fabr., Ent. Syst., iv, 169, 1794 (*Lygæus*); Sign., Ann. Soc. Ent. France, 1859, 95; Stål, Hem. Fabric. i, 69, 1868; Hambleton, Ann. Ent. Soc. Amer. i, 142, 1908.—W. States.

pictipes Stål, Freg. Eugeniès Resa, Hem. 239, 1859.

mexicanus, *proximus*, *nebulosus*, *anticus* et *ventralis* Sign., Ann. Soc. Ent. France, 1859, 92, 95, 96, 98, 99.

scutatus Stål, Freg. Eugeniès Resa, Hem. 239, 1859 (*Rhopalus*); Hambleton, Ann. Ent. Soc. Amer. 1, 139, 1908.—W. States.

jactatus Sign., Ann. Soc. Ent. France, 1859, 81.

tuberculatus Hambleton, Ann. Ent. Soc. Amer. i, 140, 1908.—Wash.

validus Uhler, Proc. Ent. Soc. Wash. i, 370, 1893; Hambleton, Ann. Ent. Soc. Amer. i, 141, 1908.—W. States.

viridicatus Uhler, Hayden's Surv. Mont. 404, 1872; Hambleton, Ann. Ent. Soc. Amer. i, 138, 1908.—W. States.

hyalinus Uhler, Bull. U. S. Geol. Surv. ii (5), 34, 1875 (not Fabr.).

HARMOSTES.

Burmeister, Handb., ii (1), 307, 1835.

Syromastes Say, Het. N. Harm. 10, 1831 (not Latreille).

affinis Dallas, B. M. List, Hem. ii, 522, 1852; Van Duzee, Bull. Buffalo Soc. Nat. Hist. ix, 161, 1909.—Fla.

- fraterculus** Say, Het. N. Harm. 10, 1831 (*Syromastes*); Compl. Writ. i, 324, 1859; Stål, Enum. Hemipt. i, 221, 1870; Distant, Biol. C.-Am. Rhynch. Het. i, 168, 1881, pl. xv, fig. 18.—U. S.
- obliquus** Say, Hem. New Harm. 10, 1831; Compl. Writ. i, 324, 1859 (*Syromastes*).—U. S.
- propinquus** Distant, Biol. C.-Am. Rhynch. Het. i, 168, 1881, pl. xv, fig. 19; Barber, Bull. Brooklyn Inst. i (No. 9), 271, 1906.—U. S.
- reflexulus** Say, Het. N. Harm. 10, 1831 (*Syromastes*); Compl. Writ. i, 323, 1859; Stål, Enum. Hemipt. i, 220, 1870.—U. S.
- costalis* Her. Schfr., Wanz. Ins. ix, 270, 1853.
- virescens* Dallas, Brit. Mus. List, Hemipt. ii, 520, 1852.
- serratus** Fabr., Ent. Syst. iv, 75, 1794 (*Acanthia*); Stål, Hemipt. Fabric. i, 67, 1870.—Fla.
- perpunctatus* Dallas, Brit. Mus. List, Hemipt. ii, 524, 1852.

LEPTOCORIS.

Hahn., Wanz. Ins. i, 200, 1831.

Serinetha Spinola, Essai, 247, 1840.

- trivittatus** Say, Journ. Acad. Phila. iv, 322, 1825 (*Lygæus*); Compl. Writ. ii, 246, 1859; Stål, Enum. Hemipt. i, 226, 1870; Howard, Cir. 28 Div. Entom. Dept. Agric.—W. States.

PYRRHOTES.

Westwood, Hope Cat. Hem. ii, 6, 1842.

Jaddera Stål, Rio Jan. Hem. i, 50, 1860.

- æola** Dallas, Brit. Mus. List, Hemipt. ii, 463, 1852 (*Serinetha*); Stål, Enum. Hemipt. i, 227, 1870; Distant, Biol. C.-Am. Rhynch. Het. i, 173, 1882.—S. States.
- hæmatoloma** Her. Schfr., Wanz. Ins. viii, 103, 1848 (*Leptocoris*); Dallas, Brit. Mus. List, Hemipt. ii, 463, 1852 (*Serinetha*); Stål, Enum. Hemipt. i, 226, 1870; Distant, Biol. C.-Am. Rhynch. Het. i, 173, 1882, pl. xvii, fig. 5.—W. States.
- marginalis* Walk., Cat. B. M. Het. v, 45, 1872 (*Lygæus*).

Pseudophloeini.

CERALEPTUS.

Costa, Cim. Reg. Neap. (Cent. 2), 11, 1847.

- americanus** Stål, Enum. Hemipt. i, 219, 1870—W. States.

DASYCORIS.

Dallas, List Hem. B. M., ii, 487, 1852.

- humilis** Uhler, Fifth Ann. Rept. U. S. Geol. Surv. f. 1871, 1872, 403, 1872.—Colo.
- nigricornis** Stål, Enum. Hem. i, 279, 1870; Dist., Biol. Cent.-Amer. Rhynch. Het. i, 164, pl. xv, fig. 12, 1881.—Colo.
- pilicornis** Burm., Handb. ii, 309, 1835 (*Coreus*); Fieber, Europ. Hem. 221, 1861.—W. States.

SCOLOPOCERUS.

Uhler, Wheeler's Surv. Ariz. 832, 1875.

secundarius Uhler, Wheeler's Surv. Ariz. 833, 1875; Distant, Biol. C.-Am. Rhynch. Het. i, 164, 1881; Barber, Sci. Bull. Brooklyn Mus. i, No. 9, 268, 1906.—W. States.

uhleri Dist., Biol. C.-Am. Rhynch. Het. i, 164, 1881, pl. xvi, fig. 6.—Ariz.

Leptocorisini.

LEPTOCORISA.

Latreille, Fam. Nat. 421, 1825.

tipuloides De Geer, Mém. Ins. iii, 354, 1773 (*Cimex*); Amy. et Serv. Hem, 229, 1843; Stål, Enum. Hem. i, 218, 1870.—S. States.

crudelis Westw., Hope Cat. Hem. ii, 18, 1842.

Alydini.

ALYDUS.

Fabricius, Syst. Rhyng, 248, 1803.

conspersus Mont., Proc. U. S. Nat. Mus. xvi, 49, 1893.—N. States.

eurinus Say, Journ. Acad. Phil. iv, 324, 1825 (*Lygæus*); Compl. Writ. ii, 247, 1859; Mont., Proc. U. S. Nat. Mus. xvi, 50, 1893.—U. S.

ater Dallas, Brit. Mus. List, Hemipt. ii, 478, 1852.

pilosulus Her. Schfr., Wanz. Ins. viii, 101, 1848; Dallas, List, Hem. B. M. ii, 478, 1852; Mont., Proc. U. S. Nat. Mus. xvi, 50, 1893.—U. S.

pluto Uhler, Hayden's Surv. Mont. 404, 1872.—W. States.

setosus Van Duzee, Ent. News, 1906, 386.—Ariz.

DARMISTUS.

Stål, Œfv. Vet. Akad. Förh. 1859, 469.

subvittatus Stål, Œfv. Vet. Akad. Förh. 1859, 469; Dist., Biol. Cent.-Amer. Rhynch. Het. i, pl. xv, fig. 10, 1881.—W. States.

ESPERANZA.

Barber, Sci. Bull. Brooklyn Mus. i, 269, 1906.

texana Barber, Sci. Bull. Brooklyn Mus. i, No. 9, 270, 1906.—Tex.

HYALMENUS.

Amyot et Serville, Hem. 224, 1843.

longispinus Stål, Enum. Hem. i, 213, 1870.—Fla.

sinuatus Guérin, Sagra's Hist. de Cuba, vii, Ins. 162, 1857 (*Campulopus*) (nec Fabr.).

pulcher, Stål, Enum. Hem. i, 211, 1870; Dist., Biol. C.-Am. Rhynch. Het. i, 156, pl. xv, fig. 9, 1881.—Tex.

tarsatus Fabr., Syst. Rhyng. 250, 1803 (*Alydus*); Stål, Enum. Hemipt. i, 212, 1870; Distant, Biol. C.-Am. Rhynch. Het. i, 156, 1881.—S. States.

recurvus Her. Schfr., Wanz. Ins. viii, 98, 1848 (*Alydus*).

pallens Dallas, Brit. Mus. List Hemipt. ii, 476, 1852 (*Alydus*).

diversipes Westw., Hope Cat. ii, 19, 1842.

affinis Westw., ibid.

sinuatus Her. Schf., Wanz. Ins. viii, 98, 1848 (nec Fabr.).

pectoralis Stål, Rio Jan. Hem. i, 34, 1858.

formicarius Dist., Biol. C.-Am. Rhynch. Het. i, 459, 1893 (*Galeottus*), young.

MEGALOTOMUS.

Fieber, Europ. Hem. 58, 1861.

quinquespinosus Say, Journ. Acad. Phila. iv, 323, 1825 (*Lygæus*); Compl. Writ., ii, 247, 1859; Stål, Enum. Hemipt. i, 214, 1870.—U. S.

cruentus Her. Schfr., Wanz. Ins. viii, 100, 1848.

PROTENTOR.

Stål, Cefv. Vet. Akad. Förh. i 1867, 543.

Tetrarhinus Prov., Nat. Can. iv, 76, 1872.

belfragei Haglund, Stett. Ent. Zeit. 1868, 162; Stål, Enum. Hemipt. i, 217, 1870.—N. States.

quebecensis Prov., Nat. Can. iv, 76, 1872 (*Tetrarhinus*).

STACHYOCNEMIS.

Stål, Enum. Hem. i, 215, 1870.

apicalis Dallas, Brit. Mus. List Hemipt. ii, 479, 1852 (*Alydus*).—U. S.

TOLLIUS.

Stål, Enum. Hem. i, 213, 1870.

curtulus Stål, Eugeniès Resa, Ins. 234, 1859 (*Alydus*); Enum. Hemipt. i, 213, 1870.—W. States.

Coreini.

ALTHOS.

Kirkaldy, Entom. 1904, 280.

Margus Dallas, List Hem. B. M. ii, 523, 1852 (preoc.).

inconspicuus Her. Schfr., Wanz. Ins. vi, 14, 1842 (*Syromastes*), Stål, Stett. Ent. Zeit. 1862, 303; Enum. Hemipt. i, 184, 1870.—W. States.

obscurator Fabr., Syst. Rhyng. 200, 1803 (*Coreus*); Stål, Hem. Fabric. i, 57, 1868,—Calif.

impudens Stål, Rio Hem. i, 37, 1858.

ANASA.

Amyot et Serville, Hém. 209, 1843.

andresii Guérin, Sagra's Hist. de Cuba, vii, Ins. 159, 1857 (*Coreus*); Stål, Enum. Hemipt. i, 190, 1870.—Fla.

lugens Stål, Stett. Ent. Zeit. 1862, 301.

armigera Say, Journ. Acad. Phil. iv, 319, 1825 (*Coreus*); Compl. Writ. ii, 244, 1859; Stål, Enum. i, 192, 1870; Bull. 19, Bur. Entom., fig. 6.—S. States.

terminalis Dallas, B. M. List, Hem. ii, 506, 1852.

obliquus Uhler, Proc. Ent. Soc. Phil., i, 23, 1861 (*Gonocerus*).—Calif.

repetita Heid., Proc. Ent. Soc. Wash. vii, 11, 1905.—D. C., Md.

scorbutica Fabr., Syst. Ent. 706, 1775 (*Cimex*); Ent. Syst. iv, 129, 1794 (*Coreus*); Stål, Hemipt. Fabric. i, 56, 1868; Enum. Hemipt. i, 190, 1870.—S. States.

nebulosus Pal. Beauv., Ins. Afric. et Amer. 205, 1805 (*Acanthocerus*).

mæsta Dallas, Brit. Mus. List Hemipt. ii, 505, 1852.

spiniceps Stål, Stett. Ent. Zeit. 1862, 300.

tristis De Geer, Mém. Ins. iii, 340, 1773 (*Cimex*); Stål, Hemipt. Fabric. i, 56, 1868; Bull. 19 Bur. Entom., fig. 5.—U. S.

ordinatus Say, Journ. Acad. Phil. iv, 318, 1825 (*Coreus*); Compl. Writ. ii, 224, 1859.

destructor Hahn, Wanz. Ins. i, 8, 1831.

rugator Burm., Handb. ii, 311, 1835 (*Gonocerus*).

uhleri Stål, Hem. Fabric. i, 57, 1868; Dist., Biol. C.-Am. Rhynch. Het. i, pl. xiii, fig. 23, 1881.—W. States.

CATORHINTHA.

Stål, Cefv. Vet. Akad. Förh. 1859, 470.

guttula Fabr., Ent. Syst. iv, 162, 1794 (*Lygæus*); Stål, Hemipt. Fabric. i, 58, 1868; Enum. Hem. i, 187, 1870.—S. States.

dorsiger Westw., Hope Cat. ii, 25, 1842 (*Gonocerus*); Dallas, Brit. Mus. List, Hemipt. ii, 504, 1852 (*Anasa*).

mendica Stål, Enum. Hem. i, 187, 1870; Dist. Biol. C.-Am. Rhynch. Het. i, pl. xiii, fig. 21.—West. States.

selector Stål, Cefv. Kgl. Vet. Förh. 1859, 471; Enum. Hemipt. i, 188, 1870; Dist., Biol. C.-Am. Rhynch. Het. i, pl. xiii, fig. 19.—W. States.

texana Stål, Enum. Hemipt. i, 188, 1870; Barber, Sci. Bull. Brooklyn Mus. i, No. 9, 267, 1906.—Tex.

CHELINIDEA.

Uhler, Proc. Ent. Soc. Phil. ii, 365, 1863.

tabulata Burm., Handb. ii, 311, 1835 (*Gonocerus*); Stål, Enum. Hemipt. i, 181, 1870; Distant, Biol. C.-Am. Rhynch. Het. i, 136, 1881, pl. xiii, fig. 17.—W. States.

vittigera Uhler, Proc. Ent. Soc. Phil., ii, 366, 1863; Stål, Enum. Hemipt. i, 180, 1870.—U. S.

CIMOLUS.

Stål, Stett. Ent. Zeit. 1862, 302.

obscurus Stål, Enum. Hem. i, 189, 1870.—S. States.

CLETUS.

Stål, Freg. Eugeniès Resa, Hem. 236, 1859.

obsoletus Her. Schfr., Wanz. Ins. vi, 10, 1842 (*Gonocerus*); Mayr, Reise Novara, Hem. 119, 1866.—Amer. Bor. (?)

FICANA.

Stål, Stett. Ent. Zeit. 1862, 303.

apicalis Dallas, Brit. Mus. List, Hemipt. ii, 499, 1852 (*Gonocerus*); Stål, Enum. Hemipt. i, 188, 1870; Dist., Biol. C.-Am. Rhynch. Het. i, pl. xiv, fig. 1.—W. States.

HYPSELONOTUS.

Hahn, Wanz. Ins. i, 186, 1831.

fulvus De Geer, Mém. Ins. iii, 341, 1773 (*Cimex*); Stål, Enum. Hem. i, 201, 1870; Dist., Biol. C.-Am. Rhynch. Het. i, 152, 1881.—Tex.

striatulus Fabr., Syst. Ent. 721, 1775.

venosus Fabr., Ent. Syst. iv, 142, 1794.

punctiventris Stål, Stett. Ent. Zeit. 1862, 297; Enum. Hemipt. i, 201, 1870; Distant, Biol. C.-Am. Rhynch. Het. i, 150, 1881; Barber, Sci. Bull. Brookl. Mus. i, No. 9, 268, 1906.—Tex.

MADURA.

Stål, Rio Jan. Hem. i, 35, 1858.

perfidia Stål, Stett. Ent. Zeit. 1862, 304; Distant, Biol. C.-Am. Rhynch. i, 135, 1881, pl. xiii, fig. 16.—Tex.

PARYPHES.

Burmeister, Handb. ii (1), 335, 1835.

rufo-scutellatus G. R. Gray, Griffith's Anim. Kingd. xv, 241, 1833 (*Nematopus*); Stål, Enum. Hem. i, 205, 1870.—Calif.

Anisoscelini.

AMBLYOMIA.

Stål, Enum. Hem. i, 171, 1870.

bifasciata Stål, Enum. Hem. i, 172, 1870.—W. States.

CHONDROCERA.

Laporte, Essai, 44, 1832.

laticornis Laporte, Essai, 45, 1832.—Fla.

unicolor Guérin, Sagra's Hist. de Cuba, vii, Ins., 162, 1857 (*Petalotoma*).—Fla.

LEPTOGLOSSUS.

- Guérin, Voyage Coquille, Ins., 174, 1838.
Anisoscelis Spinola, Essai, 200, 1837 (not Latreille).
Theognis Stål, Stett. Ent. Zeit. 1862, 294.
ashmeadi Heid., Bull. Buffalo Soc. Nat. Hist. ix, 237, 1909.—Fla.
corculus Say, Het. New Harm. 12, 1831 (*Anisoscelis*); Compl. Writ. i, 326, 1859.—S. States.
excellens Mayr, Verh. zool.-bot. Ges. Wien. 1865, 434.
gonagra Fabr., Syst. Ent. 708, 1775 (*Cimex*); Syst Rhyng. 210, 1803; Stål, Hem. Fabr. i, 51, 1868.—Fla.
grallator Herbst, Gem. Naturg. vi, 259, 1784 (*Cimex*).
anticus H. Schf., Wanz. Ins. iii, 92, 1835, fig. 316 (*Anisoscelis*).
oppositus Say, Het. N. Harm. 12, 1831 (*Anisoscelis*); Compl. Writ. i, 327, 1859; Stål, Enum. Hem. i, 163, 1870; Bull. 19 Bur. Entom., fig. 9.—U. S.
tibialis Her. Schf., Wanz. Ins. vii, 12, 1844 (*Anisoscelis*).
phyllopus Linn., Syst. Nat., Ed. 12, 731, 1767 (*Cimex*); Burm., Handb. ii, 332, 1835; Stål, Enum. Hemipt. i, 161, 1870.—S. States.
albicinctus Say, Het. N. Harm. 12, 1831 (*Anisoscelis*); Compl. Writ. i, 326, 1859; Stål, Hemipt. Fabric. i, 52, 1868.
confusa Dallas, Brit. Mus. List, Hemipt. ii, 453, 1852 (*Anisoscelis*).
fraternus Westw., Hope Cat. ii, 16, 1842 (*Anisoscelis*).
zonatus Dallas, Brit. Mus. List, Hemipt. ii, 452, 1852 (*Anisoscelis*); Stål, Enum. Hemipt. i, 162, 1870; Dist., Biol. C.-Am. Rhynch. Het. i, pl. xii, fig. 16.—Fla., Tex., Cal.
occidentalis, Heid.

NARNIA.

- Stål, Stett. Ent. Zeit. 1862, 294.
femorata Stål, Stett. Ent. Zeit. 1862, 296; Dist., Biol. Cent.-Amer. Rhynch. Het. i, pl. xiii, fig. 4, 1881.—S. W. States.
inornata Distant, Biol. C.-Am. Rhynch. Het. i, 361, pl. xxxiii, fig. 10, 1892.—Calif.
pallidicornis Stål, Enum. Hemipt. i, 166, 1870.—S. W. States.
snowi Van Duzee, Ent. News, 1906, 384.—Ariz.
wilsoni Van Duzee, Ent. News 1906, 385.—Calif.

PACHYLIS.

- Lepelletier et Serville, Ency. Meth. x, 62, 1825.
Thasus Stål, Ann. Soc. Ent. France, 1865, 174.
gigas Burm., Handb. ii, 338, 1835; Dallas, Brit. Mus. List, Hemipt. ii, 383, 1852; Uhler, Bull. U. S. Geol. and Geog. Surv. Terr. ii, No. 5, 295, 1876; Dist., Biol. C.-Am. Rhynch. Het. i, 108, pl. x, fig. 18, 19, 1881.—Ariz., N. Mex.

PHTHIA.

Stål, Stett. Ent. Zeit. 1862, 294.

- picta** Drury, Illust. i, 107, 1770 (*Cimex*); Westw., Hope Cat. ii, 17, 1842 (*Leptoscelis*); Dallas, Brit. Mus. List, Hemipt. i, 457, 1852; Stål, Hemipt. Fabric. i, 53, 1868.—Fla., Tex.
ciliatus Fabr., Syst. Ent. 706, 1775 (*Cimex*).
leprosus Fabr., Syst. Ent. 719, 1775 (*Cimex*).
crenulatus Fabr., Ent. Syst. iv, 144, 1794 (*Cimex*).
dispar Fabr., Syst. Rhyng. 214, 1803 (*Lygæus*).
divisus Her. Schfr., Wanz. Ins. vii, 9, 1844 (*Anisoscelis*).
pulverulentus Her. Schfr., ibid.
obscura Dallas, Brit. Mus. List, Hem. ii, 458, 1852.

Acanthocephalini.

ACANTHOCEPHALA.

Laporte, Essai, 29, 1832.

- declivis** Say, Insects, Louis. ii, 1832 (*Rhinuchus*); Compl. Writ. i, 305, 1859; Stål, Enum. Hem. i, 150, 1870; Dist., Biol. C.-Amer. Rhynch. Het. i, pl. xi, figs. 19, 21, 24.—S. States.
alatus Burm., Handb. ii (1), 334, 1835 (*Diactor*).
thoracicus Dallas, B. M. List, Hem. ii, 428, 1852 (*Metapodius*).

METAPODIUS.

Westwood, Hope Cat. Hem. ii, 4, 1842.

- confraternus** Uhler, Proc. Bost. Soc. Nat. Hist. xiv, 90, 1871; xix, 382, 1878.—S. States
femoratus Fabr., Syst. Entom. 708, 1775 (*Cimex*); Syst. Rhyng. 205, 1803; Dall. B. M. List, Hem. ii, 430, 1852; Bull. 69 Minn. Exp. Sta., fig. 82.—U. S.
nasulus Say, Insects, Louis. 10, 1832 (*Rhinuchus*); Compl. Writ. i, 305, 1859.
obscurus Westw., Hope Cat. Hem. ii, 15, 1842.
granulosus Dallas, B. M. List, Hem. ii, 430, 1852; Dist., Biol. C.-Am. Rhynch. Het. i, pl. xi, fig. 18.—S. States.
alatus Her. Schf., Wanz. Ins. vi, 53, 1842 (*Diactor*) (nec Burm.).
thomasi Uhler, Hayden's Surv. Mont. 399, 1872.
instabilis Uhler, Proc. Bost. Soc. Nat. Hist. xiv, 98, 1871; xix, 381, 1878.—Pa., N. Car.
terminalis Dallas, Brit. Mus. List, Hemipt. ii, 431, 1852; Stål, Enum. Hem. i, 151, 1870; Uhler, Bull. U. S. Geol. and Geog. Surv. Terr. ii, No. 5, 298, 1876.—S. States.

Mictini.

ARCHIMERUS.

Burmeister, Handb. ii (1), 321, 1835.

- ashmeadi** Mont., Bull. Soc. Bucarest, viii, 194, 1899.—Fla.

- calcarator** Fabr., Syst. Rhyng. 192, 1803 (*Coreus*); Stål, Hem. Fabr., i, 47, 1868.—U. S.
alternatus Say, Journ. Acad. Philad., iv, 317, 1825; Compl. Writ. ii, 243, 1859.
squalus Burm., Handb. ii (1), 321, 1835.
albonotatus A. and S., Hém. 197, 1843.
rubiginosus Her. Schf., Wanz. Ins. vi, 83, 1842.
muticus Her. Schf., ibid, 52.
indecorus Walker, Cat. Heter. iv, 64, 1871; Dist., Biol. C.-Am. Rhynch. Het. i, pl. xi, figs. 4, 5.—Ariz.

CAPANUS.

- Stål, Stett. Ent. Zeit, 1862, 277.
Xuthus Stål, Enum. Hem. i, 136, 1870.
auriculatus Stål, Stett. Ent. Zeit. 1862, 290; Enum. Hemipt. i, 136, 1870; Dist., Biol. Cent.-Amer. Rhynch. Het. i, pl. xi, fig. 8, 1881.—S. States.
spurcus Stål, Stett. Ent. Zeit. 1862, 291; Enum. Hem. i, 137, 1870; Dist., Biol. C.-Am. Rhynch. Het. i, pl. xii, fig. 3, 1881.—Ariz.

EUTHOCTHA.

- Mayr, Verh. zool-bot. Ges. Wien. xv, 431, 1865.
galeator Fabr., Syst. Rhyng. 191, 1803 (*Coreus*); Stål, Hemipt. Fabric. i, 49, 1868.—U. S.
tibialis Her. Schfr., Wanz. Ins. vi, 21, 1842 (*Crinocerus*).

MOZENA.

- Amyot et Serville, Hém. 192, 1843.
brunneicornis H. Schf., Wanz. Ins. vi, 26, 1842 (*Archimerus*); Stål, Enum. Hem. ii, 134, 1872.—Ariz.
spinicrus Amyot et Serv., Hém. 192, 1843.
lineolata Her. Schf., Wanz. Ins. vi, 25, 1842 (*Archimerus*); Stål, Enum. Hemipt. i, 134, 1870.—S. States.
lunata Burm., Handb. ii (1), 322, 1835 (*Archimerus*); Her. Schf., Wanz. Ins. vi, 24, 1842 (*Archimerus*); Stål, Enum. Hemipt. i, 134, 1870.—Tex., N. Mex.
lurida Dallas, Brit. Mus. List, Hemipt. ii, 417, 1852 (*Archimerus*); Stål, Enum. Hemipt. i, 135, 1870; Distant, Biol. C.-Am. Rhynch. Het. i, pl. xii, fig. 1, 1881.—Tex.
obesa Mont., Bull. Soc. Bucarest, viii, 190, 1899.—Fla.
obtusa Uhler, Bull. U. S. Geol. and Geog. Surv. Terr. ii, No. 5, 296, 1876.—S. W. States.

SAGOTYLUS.

- Mayr, Verh. zool.-bot. Ges. Wien. xv, 431, 1865.
triguttatus Her. Schfr., Wanz. Ins. vi, 86, 1842 (*Crinocerus*); Dallas, B. M. List, Hem. ii, 402, 1852 (*Mictis*).—Ariz., Calif.

*Merocorini.***CORYNOCORIS.**

Mayr, Reise Freg. Novara, Hem. 95, 1866.

typhæus Fabr., Ent. Syst. Suppl. 537, 1798 (*Lygæus*); Dallas, List Hem. i, 419, 1852 (*Merocoris*).—U. S.

rugosus Amyot et Serville, Hém. 214, 1843.

acridioides Fabr., Syst. Rhyng. 200, 1803.

distinctus Dallas, Brit. Mus. List, Hemipt. ii, 419, 1852.—N. States.

*Chariesterini.***CHARIESTERUS.**

Laporte, Essai, 44, 1832.

antennator Fabr., Syst. Rhyng. 198, 1803 (*Coreus*); Dallas, B. M. List ii, 510, 1852; Stål, Enum. Hem. i, 178, 1870.—U. S.

dubius Say, Het. N. Harm. 10, 1831 (*Gonocerus*); Compl. Writ. i, 323, 1859.

mæstus Her. Schf., Wanz. Ins. vii, 3, 1844.

cuspidatus Distant, Biol. C.-Am. Rhynch. Het. i, 364, 1892; Barber, Sci. Bull. Brooklyn Mus. i, No. 9, 266, 1906.—S. W. States.

*Spartocerini.***SEPHINA.**

Amyot et Serville, Hém. 185, 1843.

grayi Van Duzee, Ent. News, 1909, 232.—Fla.

gundlachi Guér., Sagra's Hist. de Cuba, vii, Ins. 156, pl. xiii, fig. 6, 1857.—Fla.

limbata Stål, Stett. Ent. Zeit. 1862, 273; Dist., Biol. Cent.-Amer. Rhynch. Het. i, pl. xii, fig. 22, 1881,—Calif.

atra Walk., Cat. Het. B. M. iv, 4, 1871.

SPARTOCERA.

Laporte, Essai, 43, 1832.

Coreocoris Hahn, Wanz. Ins. ii, 14, 1834.

cinnamonea Hahn, Wanz. Ins. ii, 15, 1834 (*Coreocoris*); Dallas, Brit. Mus. List, Hemipt. ii, 375, 1852; Stål, Enum. Hem. i, 174, 1870.—S. States.

subfulvus Westw., Hope. Cat. ii, 8, 1842.

diffusus Say, Het. N. Harm. ii, 1831 (*Coreus*); Compl. Writ. i, 325, 1859.

confluenta Say, Het. New Harm. 11, 1831; Compl. Writ. i, 325, 1859 (*Coreus*); Van Duzee, Bull. Buffalo Soc. Nat. Hist. ix, 159, 1909.—Fla.

PENTATOMIDÆ.

ACANTHOSOMINÆ.

ELASMOTETHUS.

Fieber, Europ. Hem. 78, 1861.

Acanthosoma, of American authors.**atricornis** Van Duzee, Trans. Amer. Ent. Soc. xxx, 75, 1904 (*Acanthosoma*).—N. States.**cooleyi** Van Duzee, Trans. Amer. Ent. Soc. xxx, 74, 1904 (as var. of *cruciata*).—N. States.**cruciata** Say, Het. N. Harm. 3, 1831; Compl. Writ. i, 311, 1859 (*Edessa*); Bull. 69 Minn. Sta., fig. 83; Van Duzee, Trans. Amer. Ent. Soc. xxx, 73, 1904.—N. States.*borealis* Westw., Hope Cat. Hem. i, 31, 1837; Dist., Proc. Zool. Soc. Lond. 1890, 818.**ELASMUCHA.**

Stål, Ann. Soc. Ent. France, 1864, 54.

Clinocoris Hahn, Wanz. Ins. ii, 70, 1834 (not Fallen).**lateralis** Say, Het. N. Harm. 3, 1831 (*Edessa*); Compl. Writ. i, 312, 1859.—U. S.*nebulosa* Kirby, Fauna Bor. Amer, Ins. 277, 1837 (*Edessa*); Can. Ent. 1878, 138.*affinis* Westw., Hope Cat. Hem. i, 30, 1837.*picolor* Westw., ibid.

PENTATOMINÆ.

ÆLIA.

Fabricius, Syst. Rhyng. 188, 1803.

americana Dallas, Brit. Mus. List, Hemipt. i, 223, 1851.—N. States.**ARVELIUS.**

Spinola, Essai, 344, 1837.

albopunctatus De Geer, Mém. Ins. iii, 331, 1773 (*Cimex*); Am. et Serv., Hem. 150, 1843.—Tex.*gladiator* Fabr., Syst. Ent. 705, 1775; Syst. Rhyng. 162, 1803.*leucostictus* Gmel., Syst. Nat. i, 2148, 1788.*luteicornis* Westw., Hope Cat. i, 30, 1837.**BANASA.**

Stål, Rio Jan. Hein. i, 24, 1858.

Atomosira Uhler, Proc. Bost. Soc. Nat. Hist. xiv, 97, 1871.**calva** Say, Het. N. Harm. 7, 1831; Compl. Writ. i, 318, 1859 (*Pentatoma*); Van Duzee, Trans. Amer. Ent. Soc. xxx, 59, 1904.—U. S.*catinus* Dallas, B. M. List, Hem. i, 282, 1851.

- dimidiata** Say, Het. N. Harm. 7, 1831; Compl. Writ. i, 318, 1859 (*Pentatoma*); Van Duzee, Trans. Amer. Ent. Soc. xxx, 60, 1904.—N. States.
- euchlora** Stål, Enum. Hem. ii, 44, 1872; Van Duzee, Trans. Amer. Ent. Soc. xxx, 61, 1904.—S. States.
- imbuta** Walk., Cat. Het. B. M. ii, 358, 1867 (*Raphigaster*); Distant, Biol. C.-Am. Rhynch. Het. i, 80, 1880.—Tex.
- lenticularis** Uhler, Proc. Zool. Soc. London, 1894, 174.—Fla.
- packardi** Stål, Enum. Hem. ii, 43, 1872; Van Duzee, Trans. Amer. Ent. Soc. xxx, 61, 1904.—Atl. States.
- sordida** Uhler, Proc. Bost. Soc. Nat. Hist. xiv, 98, 1871 (*Atomosira*); Stål, Enum. Hem. ii, 44, 1872; Van Duzee, Trans. Amer. Ent. Soc. xxx, 60, 1904.—U. S.
- varians** Stål, Hem. Rio Jan. i, 22, 1858 (*Rhaphigaster*); Van Duzee, Trans. Amer. Ent. Soc. xxx, 60, 1904.—N. Mex.

BREPHOLOXA.

Van Duzee, Trans. Amer. Ent. Soc. xxx, 78, 1904.

- heidemanni** Van Duzee, Trans. Amer. Ent. Soc. xxx, 78, 1904.—Fla.

CARPOCORIS.

Kolenati, Melet. Entom. iv, 45, 1846.

- remotus** Horv., Ann. Mus. Nat. Hung. v, 296, 1907.—W. States.
- lynx* Auctors Amer. (nec. Fabr.).

CHLOROCORIS.

Spinola, Essai, 288, 1837.

- hebetatus** Distant, Biol. Cent.-Am. Rhynch. Het. i, Suppl. 335, 1889.—Ariz.

ŒENUS.

Dallas, List Hem. B. M. i, 194, 1851.

- delius** Say, Het. N. Harm. 8, 1831 (*Pentatoma*); Compl. Writ. i, 320, 1859.—U. S.
- æruginea* A. and S., Hémipt. 125, 1843.
- tarsalis* Dallas, Brit. Mus. List, Hemipt. i, 230, 1851.
- punctatissimus* Voll., Versl. Akad. Amst. Nat. (2), ii 183, 1868.

COSMOPEPLA.

Stål, Œfv. Vet. Akad. Förh. 1867, 525.

- binotata** Distant, Biol. Cent.-Amer. Rhynch. Het. i, Suppl. 327, 1889.—Ariz.
- carnifex** Fabr., Ent. Syst. Suppl. 535, 1798 (*Cimex*); Hahn., Wanz. Ins. ii, 117, 1834 (*Eysarcoris*); Stål, Hem. Fabr. i, 28, 1868; Mont., Proc. U. S. Nat. Mus. xvi, 46, 1893; Bull. 69 Minn. Sta., fig. 89.—U. S.

- conspicillaris** Dallas, Brit. Mus. List, Hemipt. i, 225, 1852; Mont., Proc. U. S. Nat. Mus. xvi, 46, 1893.—W. States.
decorata Hahn, Wanz. Ins. ii, 117, 1834 (*Eysarcoris*); Her. Schf., Wanz. Ins. vii, 96, 1844; Mont., Proc. U. S. Nat. Mus. xvi, 46, 1893.—S. W. States.
uhleri Mont., Proc. U. S. Nat. Mus. xvi, 48, 1893.—Calif., Nev.

DENDROCORIS.

Bergroth, Rev. d'Ent. x, 228, 1891.

- Liotropis* Uhler, Bull. Geog. Geol. Surv. iii, 399, 1877.
contaminatus Uhler, Trans. Md. Acad. Sci. 1897, 390 (*Liotropis*); Van Duzee, Trans. Amer. Ent. Soc. xxx, 62, 1904.—Ariz.
fruticicola Bergroth, Rev. d'Ent. x, 228, 1891; Van Duzee, Trans. Amer. Ent. Soc. xxx, 62, 1904.—Fla.
humeralis Uhler, Bull. Geogr. Geol. Surv. iii, 400, 1877 (*Liotropis*). U. S.
pini Mont., Proc. U. S. Nat. Mus. xvi, 51, 1893; Van Duzee, Trans. Amer. Ent. Soc. xxx, 63, 1904.—Calif.
schæfferi Barber, Sci. Bull. Brooklyn Museum, i, No. 9, 262, 1906.—Tex.

EDESSA.

Fabricius, Syst. Rhyng. 145, 1803.

- Ascra* Say, Ins. Louis. 8, 1832; Compl. Writ. i, 304, 1859.
bifida Say, Ins. Louis. 7, 1832; Compl. Writ. i, 303, 1859 (*Pentatoma*); Stål, Enum. Hem. ii, 58, 1872.—S. States.
albirenis Her. Schf., Wanz. Ins. vii, 127, 1844; Dallas, List, Hem. B. M. i, 336, 1851.
cornuta Guérin, Sagra's Hist. de Cuba, vii, Ins. 154, 1857.

EUSCHISTUS.

Dallas, List Hem. B. M. i, 193, 1851.

- bifibulus** Pal. Beauv., Ins. Afr. et Amer., 148, 1805; Stål, Enum. Hem. ii, 27, 1872; Van Duzee, Trans. Amer. Ent. Soc. xxx, 48, 1904.—Fla., Tex.
pallipes Dallas, Brit. Mus. List Hem. 204, 1851.
biformis Stål, Stett. Ent. Zeit. 1862, 100; Enum. Hem. ii, 28, 1872.—Ariz., Tex.
comptus Walk., B. M. Cat. Het. iii, 550, 1868; Stål, Enum. Hem. ii, 28, 1872; Distant, Biol.-Cent. Amer. Rhynch. Het. i, pl. i, fig. 11, 1881.—Tex.
conspersus Uhler, Trans. Md. Acad. Sci. 1897, 388; Van Duzee, Trans. Amer. Ent. Soc. xxx, 46, 1904.—Calif., Wash.
crassus Dallas, Brit. Mus. List, Hem. i, 205, 1851; Van Duzee, Trans. Amer. Ent. Soc. xxx, 48, 1904.—Ga., Fla.; N. Car.

- crenator** Fabr., Ent. Syst. iv, 101, 1794 (*Cimex*); Stål, Hemipt. Fabric. i, 26, 1868; Enum. Hem. ii, 27, 1872; Van Duzee, Trans. Amer. Ent. Soc. xxx, 48, 1904.—S. States.
- obscura* Palisot-Beauv., Ins. Afr. et Amer. 149, 1805; Dallas, B. M. List Hem. i, 208, 1851.
- pustulata* Palisot-Beauv., ibid.
- conterminus* Walk., B. M. Cat. Het. ii, 248, 1867.
- melanocera* Walk., ibid, iii, 444, 1868.
- fissilis** Uhler, Proc. Bost. Soc. Nat. Hist. xiv, 96, 1871; ibid, xix, 374, 1878; Hayden's Surv. Montana, 396, 1872; Stål, Enum. Hem. ii, 26, 1872; Van Duzee, Trans. Amer. Ent. Soc. xxx, 44, 1904.—U. S.
- euschistoides* Voll., Versl. Akad. Amst. Nat. (2) ii, 180, 1868 (*Diceræus*).
- jugalis* Prov., Faun. Ent. Can. Hem. 204, 1887.
- ictericus** Linn., Cent. Ins. 16, 1763; Syst. Nat. xii, 719, 1767 (*Cimex*); Fabr., Syst. Ent. 702, 1775; Stål, Enum. Hemipt. ii, 26, 1872; Van Duzee, Trans. Amer. Ent. Soc. xxx, 47, 1904.—S. States. *New England Also. — U. S.*
- rubro-fusca* Palisot-Beauv., Ins. Afr. et Amer. 185, 1805.
- cognatus* Dallas, B. M. List Hem. i, 204, 1851.
- impictiventris** Stål, Enum. Hem. ii, 26, 1872; Van Duzee, Trans. Amer. Ent. Soc. xxx, 45, 1904.—N. Mex.
- inflatus** Van Duzee, Trans. Amer. Ent. Soc. xxix, 107, 1903; ibid, xxx, 45, 1904.—Colo., N. Mex.
- politus** Uhler, Can. Entom. 1897, 117; Van Duzee, Trans. Amer. Ent. Soc. xxx, 46, 1904.—E. States.
- servus** Say, Het. N. Harm. 4, 1831 (*Pentatoma*); Compl. Writ. i, 314, 1859; Stål, Enum. Hemipt. ii, 26, 1872; Van Duzee, Trans. Amer. Ent. Soc. xxx, 45, 1904.—U. S.
- spilota* Westw., Hope Cat. i, 42, 1837.
- tristigmus** Say, Het. New Harm. 4, 1831 (*Pentatoma*); Compl. Writ. i, 314, 1859; Her. Schf., Wanz. Ins. vii, 101, 1844; Stål, Enum. Hem. ii, 26, 1872; Van Duzee, Trans. Amer. Ent. Soc. xxx, 47, 1904.—U. S.
- pyrrhocerus* Her. Schf., Wanz. Ins. vii, 42, 1844.
- luridus* Dallas, B. M. List Hem. i, 207, 1851.
- inconspicua* Westw., Hope Cat. i, 42, 1837.
- variolarius** Pal. Beauv., Ins. Afr. et Amer. 149, 1805 (*Pentatoma*); Stål, Enum. Hem. ii, 26, 1872; Van Duzee, Trans. Amer. Ent. Soc. xxx, 47, 1904; Farm. Bull. 120, Dept. Agric., fig. 13.—U. S.
- punctipes* Say, Journ. Acad. Phil. iv, 314, 1825; Compl. Writ. ii, 241, 1859.
- ictericus* Her. Schfr., Wanz. Ins. vi, 71, 1842 (nec Linné.).
- sordidus* Her. Schfr., ibid, 70, 1842.

zophilotensis Distant, Biol. Cent.-Am. Rhynch. Het. i, Suppl. 330, 1889; Van Duzee, Trans. Amer. Ent. Soc. xxx, 48, 1904.—Tex.

EYSARCORIS.

Hahn, Wanz. Ins. ii, 66, 1834.

intergressus Uhler, Proc. Ent. Soc. Wash. ii, 368, 1893.—W. States.

punctifer Walk., Cat. Het. B. M. ii, 274, 1867.—Calif.

HYMENARCYS.

Amyot et Serville, Hém. 124, 1843.

æqualis Say, Het. N. Harm. 7, 1831 (*Pentatoma*); Compl. Writ. i, 319, 1859; Stål, Enum. Hem. ii, 30, 1872.—U. S.

dentatus Her. Schf., Wanz. Ins. v, 64, 1839.

boxura Dallas, Brit. Mus. List, Hemipt. i, 244, 1851.

crassa Uhler, Trans. Md. Acad. Sci. 1897, 387.—Ariz.

nervosa Say, Het. N. Harm. 9, 1831 (*Pentatoma*); Compl. Writ. i, 321, 1859.—U. S.

pennsylvanica Westw., Hope Cat. i, 35, 1837.

perpunctata Am. et Serv., Hémipt. 124, 1843.

reticulata Stål, Enum. Hem. ii, 30, 1872; Dist., Biol. C.-Am. Rhynch. Het. i, 63, pl. vi, fig. 17, 1880.—Ariz.

LOXA.

Amyot et Serville, Hém. 137, 1843.

flavicollis Drury, Ill. ii, 67, 1773 (*Cimex*); Amyot et Serv., Hist. Nat. Hém. 137, 1843.—Fla.

albicollis Fabr., Spec. Ins. ii, 347, 1781.

viridis Pal. Beauv., Ins. Afric. et Amer. 111, 1805.

virescens Amyot et Serv., Hist. Nat. Hém. 137, 1843.

deducta Walk., B. M. Cat. Het. ii, 242, 1867.

florida Van Duzee, Bull. Buffalo Soc. Nat. Hist. ix, 156, 1909.—Fla.

MECIDEA.

Dallas, List Hem. B. M. i, 132, 1864.

longula Stål, Cefv. Vet. Akad. Förh. 233, 1854; *ibid*, 1856, 57; Van Duzee, Trans. Amer. Ent. Soc. xxx, 26, 1904.—W. States.

MENECLIS.

Stål, Cefv. Vet. Akad. Förh. 1867, 527.

insertus Say, Het. N. Harm. 6, 1831; Compl. Writ. i, 317, 1859 (*Pentatoma*); Stål, Cefv. Vet. Akad. Förh. 1867, 527.—N. States.

MORMIDEA.

Amyot et Serville, Hém. 134, 1843.

lugens Fabr., Syst. Ent. 716, 1775 (*Cimex*); Syst. Rhyng. 187, 1803; Her. Schf., Wanz. Ins. v, 66, 1839; Stål, Stett. Ent. Zeit. xxiii, 103, 1862.—U. S.

albipes Fabr., Ent. Syst. Supp. 535, 1798.

gamma Fabr., Syst. Rhyng. Index 7, 1803.

punctipes Pal. Beauv., Ins. Afric. et Amer. 113, 1805; Say, Journ. Acad. Phil. iv, 313, 1825.

pictiventris Stål, Stett. Ent. Zeit. 1862, 103; Distant, Biol. C.-Am. Rhynch. Het. i, 55, 1880, pl. v, fig. 11.—Tex.

sordidula Stål, Enum. Hem. ii, 21, 1872; Van Duzee, Trans. Amer. Ent. Soc. xxx, 43, 1904.—Tex., N. Mex.

tetra Walk., Cat. Het. B. M. iii, 551, 1868; Van Duzee, Trans. Amer. Ent. Soc. xxx, 43, 1904.—Tex., Ariz.

MURGANTIA.

Stål, Stett. Ent. Zeit, 1862, 105.

histrionica Hahn, Wanz. Ins. ii, 116, 1834, fig. 196 (*Strachia*); Stål, Enum. ii, 37, 1872; Cir. 103, Bur. Entom. Dept. Agric.—U. S.

varicolor Westw., Hope Cat. i, 37, 1837 (*Pentatoma*).—W. States.

violacens Westw., Hope Cat. i, 34, 1837 (*Pentatoma*); Dallas, List Hem. B. M. i, 263, 1851; Van Duzee, Trans. Amer. Ent. Soc. xxx, 57, 1904.—Fla.

concinna Dallas, List Hem. B. M. i, 265, 1851.

— *munda*, Stål. W. States not noticed.

NEOTTIGLOSSA.

Kirby, Rich. Fauna Bor. Amer. iv. 276, 1837.

Melanostoma Stål, Enum. Hem. ii, 18, 1872.

cavifrons Stål, Enum. Hemipt. ii, 18, 1872 (*Melanostoma*).—W. States.

sulcifrons Stål, Enum. Hemipt. ii, 18, 1872 (*Melanostoma*).—S. States.

trilineata Kirby, Fauna Bor. Am. iv, 276, 1837; Uhler, Bull. U. S. Geol. and Geog. Surv. Terr. iii, 401, 1877.—N. States.

undata Say, Het. N. Harm. 8, 1831 (*Pentatoma*); Compl. Writ. i, 319, 1859; Stål, Enum. Hem. ii, 18, 1872; Van Duzee, Trans. Amer. Ent. Soc. xxx, 50, 1904.—U. S.

NEZARA.

Amyot et Serville, Hém. 133, 1843.

Rhaphigaster Dallas, List Hem. B. M. i, 274, 1851 (not Laporte).

hilaris Say, Het. New Harm. 5, 1831; Compl. Writ. i, 304, 316, 1859 (*Pentatoma*); Van Duzee, Trans. Amer. Ent. Soc. xxx, 58, 1904.—U. S.

sarpinus Dallas, List Hem. B. M. i, 276, 1851.

marginata Pal. Beauv., Ins. Afric. et Amer., 147, 1805 (*Pentatoma*); Amyot et Serv., Hist. Nat. Hém., 145, 1843; Van Duzee, Trans. Amer. Ent. Soc. xxx, 58, 1904.—Fla.

spirans Dallas, List Hem. B. M. i, 280, 1851 (not Fabr.).

jurgiosa Stål, Stett. Ent. Zeit. 1862, 106.

- pennsylvanica** De Geer, Mém. Ins. iii, 330, 1773 (*Cimex*); Pal. Beauv., Ins. Afric. et Amer. 186. 1805; Van Duzee, Trans. Amer. Ent. Soc. xxx, 58, 1904; Bull. 69 Minn. Sta., fig. 84.—E. States.
- abrupta* Say, Het. N. Harm. 6, 1831; Compl. Writ. i, 317, 1859.
- viridula** Linn., Syst. Nat. x, i, 444, 1758 (*Cimex*); Fabr., Syst. Rhyng. 166, 1803; Van Duzee, Trans. Amer. Ent. Soc. xxx, 58, 1904.—S. States.
- smaragdulus* Fabr., Syst. Ent. 711, 1775.
- unicolor* Westw., Hope Cat. i, 38, 1837.
- torquatus* Fabr., Syst. Ent. 710, 1775.
- spirans* Fabr., Ent. Syst. Supp. 533, 1798.
- viridissimus* Wolff, Icones Cim. ii, 55, 1801.
- subsericea* Westw., Hope Cat. i, 38, 1837.
- tripunctigera* Westw., ibid.
- prasinus* Dall., B. M. List, Hem. i, 274, 1851.
- orbis* Stål, Öfv. Vet. Akad. Förh. 1853, 221.

PADÆUS.

- Stål, Stett. Ent. Zeit. 1862, 101.
- irroratus** Her. Schfr., Wanz. Ins. iv, 19, 1839 (*Cimex*); vii, 94, 1844; Stål, Stett. Ent. Zeit. 1862, 101.—Fla.

PENTATOMA.

- Olivier, Ency. Meth. iv, 25, 1790.
- Lioderma* Uhler, Proc. Bost. Soc. Nat. Hist. xiv, 97, 1871.
- Chlorochroa* Stål, Enum. Hem. ii, 33, 1872.
- Rhytidolomia* Stål, Enum. Hem. ii, 33, 1872.
- belfragei** Stål, Enum. Hem. ii, 33, 1872; Van Duzee, Trans. Amer. Ent. Soc. xxx, 37, 1904.—W. States.
- congrua** Uhler, Bull. U. S. Geol. Geog. Surv. Terr. ii, 288, 1876; Van Duzee, Trans. Amer. Ent. Soc. xxx, 40, 1904.—W. States.
- faceta** Say, Journ. Acad. Phila. iv, 315, 1825; Compl. Writ., ii, 242, 1859; Van Duzee, Trans. Amer. Ent. Soc. xxx, 38, 1904.—W. States.
- granulosa** Uhler, Hayden's Surv. Mont. 398, 1872.—Colo.
- harrisii** Westw., Hope Cat. i, 42, 1837; Distant, Proc. Zool. Soc. Lond. 1900, pl. 53, fig. 2.—U. S.
- ligata** Say, Het. N. Harm. 5, 1831; Compl. Writ. i, 315, 1859; Van Duzee, Trans. Amer. Ent. Soc. xxx, 41, 1904; Bull. 69, pl. 1, Bur. Entom.—U. S.
- rufocinctus* Her. Schf., Wanz. Ins. iv, 94, 1839, fig. 436.
- marginalis* Walk., B. M. Cat. Het. ii, 288, 1867.
- rubromarginatus* Kouch., Hor. Soc. Ent. Ross. iv, 98, 1867.
- osborni** Van Duzee, Trans. Amer. Ent. Soc. xxx, 37, 1904.—Colo., Tex.

- persimilis** Horvath, Ann. Mus. Nat. Hung. vi, 555, 1908 (*Chlorochroa*).
—N. States.
- juniperina* Van Duzee, Trans. Amer. Ent. Soc. xxx, 39, 1904
(nec Linn.).
- saucia** Say, Het. N. Harm. 6, 1831; Compl. Writ. i, 318, 1859; Van
Duzee, Trans. Amer. Ent. Soc. xxx, 36, 1904.—Atl. States.
- sayi** Stål, Enum. Hemipt. ii, 33, 1872; Van Duzee, Trans. Amer. Ent.
Soc. xxx, 41, 1904.—W. States.
- senilis** Say, Het. N. Harm. 5, 1831; Compl. Writ. i, 316, 1859; Van
Duzee, Trans. Amer. Ent. Soc. xxx, 37, 1904.—Atl. States.
- oblonga* Westw., Hope Cat. i, 37, 1837.
- grisea* Dallas, B. M. List Hem. i, 246, 1851.
- ovalis* Westw., Hope Cat. i, 39, 1837.
- uhleri** Stål, Enum. Hem. ii, 33, 1872; Van Duzee, Trans. Amer. Ent.
Soc. xxx, 39, 1904; 23 Ill. Rept. 116, 1905.—S. W. States.
- (?) *flavomarginatus* Kouch., Hor. Soc. Ent. Ross. iv, 99, 1867
(*Cimex*).
- viridicata** Walk., Cat. Het. B. M. ii, 283, 1867 (*Hymenarcys*); Uhler,
Rept. Geog. and Geol. Surv. v, 830, 1875; Van Duzee, Trans.
Amer. Ent. Soc. xxx, 36, 1904.—W. States.

PERIBALUS.

- Mulsant et Rey, Pun. Pentat. 262, 1866.
- Holcostethus* Fieber, Eur. Hem. 334, 1861 (part).
- abbreviatus** Uhler, Hayden's Surv. Mont. 397, 1872 (*Holcostethus*);
Van Duzee, Trans. Amer. Ent. Soc. xxx, 33, 1904.—W.
States.
- dubia** Dallas, B. M. List, Hem. i, 237, 1851.—N. Am.
- limbolarius** Stål, Enum. Hem. ii, 34, 1872; Van Duzee, Trans. Amer.
Ent. Soc. xxx, 32, 1904.—U. S.
- modestus* Uhler, Hayden's Surv. Mont. 396, 1872.
- piceus** Dallas, B. M. List, Hem. i, 236, 1851; Van Duzee, Trans.
Amer. Ent. Soc. xxx, 34, 1904.—Mont., Can.
- tristis** Van Duzee, Trans. Amer. Ent. Soc. xxx, 33, 1904.—B. Col.

PIEZODORUS.

- Fieber, Eur. Hem. 78, 1861.
- guildingi** Westw., Hope Cat. Hem. i, 31, 1837 (*Raphigaster*); Dallas,
List, Hem. B. M. i, 283, 1851; Stål, Enum. Hemipt. ii, 45,
1872.—Fla., N. Mex.

PRIONOSOMA.

- Uhler, Proc. Ent. Soc. Phil. ii, 363, 1863.
- podopioides** Uhler, Proc. Ent. Soc. Phil. ii, 364, 1863.—W. States.
- villosum* Prov., Faun. Ent. Can. Hem. 204, 1887.

PROXYS.

Spinola, Essai, 325, 1837.

punctulatus Pal. Beauv., Ins. Afr. et Amer. Hem. 188, 1805 (*Halys*);
Stål, Enum. Hemipt. ii, 29, 1872.—S. States.

tenebrosa Say, Het. N. Harm. 10, 1832; Compl. Writ. i, 304, 1859.

geniculata Stål, Stett. Ent. Zeit. 1862, 102.

SCIOCORIS.

Fallen, Hem. Suec. 20, 1829.

microphthalmus Flor, Rhyng. Livlands, i, 114, 1860.—N. H.

umbrinus Fieber, Rhynch. 16, 1851; Europ. Hem. 358, 1861.

SOLUBEA.

Bergroth, Rev. d'Entom. 1891, 235.

Æbalus Stål, Stett. Ent. Zeit. 1862, 102 (preoc.).

pugnax Fabr., Syst. Ent. 704, 1775 (*Cimex*); Syst. Rhyng. 161, 1803;
Stål, Enum. Hemipt. ii, 22, 1872; 23 Ill. Rept. 194, 1905.

—U. S.

typhæus Fabr., Syst. Rhyng. 162, 1803.

orthacantha Palisot-Beauv., Ins. Afr. et Amer. 130, 1805.

augur Say, Het. N. Harm. 3, 1831; Compl. Writ. i, 313, 1859.

THYANTA.

Stål, Rio Jan. Hem. ii, 58, 1862.

antiguensis Westw., Hope Cat. i, 36, 1837; Van Duzee, Trans. Amer.
Ent. Soc. xxx, 54, 1904.—Ariz., Calif.

tæniola Dallas, List Hem. B. M. i, 250, 1851; Dist., Biol. Cent.-
Am. Rhynch. Het. i, pl. vii, fig. 4, 1880.

brevis Van Duzee, Trans. Amer. Ent. Soc. xxx, 56, 1904.—Colo.

casta Stål, Stett. Ent. Zeit. 1862, 104; Van Duzee, Trans. Amer. Ent.
Soc. xxx, 54, 1904.—Calif., Ariz.

custator Fabr., Syst. Rhyng. 164, 1803 (*Cimex*); Her. Schf., Wanz.
Ins. vii, 96, 1844; Van Duzee, Trans. Amer. Ent. Soc. xxx,
53, 1904.—U. S.

calceata Say, Het. N. Harm., 8, 1831.

pallido-virens Stål, Hem. Eug. Resa, Ins. 227, 1859 (*Pentatoma*);
Van Duzee, Trans. Amer. Ent. Soc. xxx, 53, 1904.—Utah,
Calif.

perditor Fabr., Ent. Syst. iv, 102, 1794 (*Cimex*); Syst. Rhyng. 163,
1803; Stål, Hem. Fabr. i, 29, 1868; Enum. Hem. ii, 34,
1872; Van Duzee, Trans. Amer. Ent. Soc. xxx, 53, 1904; 23,
Ill. Rept. 117, 1905.—U. S.

fascifera Pal. Beauv., Ins. Afr. et Amer. 150, 1805.

collaris Westw., Hope Cat. i, 40, 1837.

transversalis Her. Schfr., Wanz. Ins. vi, 66, 1842.

fasciatus Walk., Cat. Het. B. M. ii, 245, 1867.

punctiventris Van Duzee, Trans. Amer. Ent. Soc. xxx, 55, 1904.—W. States.

rugulosa Say, Het. N. Harm. 8, 1831; Compl. Writ. i, 319, 1859 (*Pentatoma*); Van Duzee, Trans. Amer. Ent. Soc. xxx, 54, 1904.—W. States.

TRICHOPEPLA.

Stål, Öfv. Vet. Akad. Förh. 1867, 528.

atricornis Stål, Enum. Hem. ii, 34, 1872; Van Duzee, Trans. Amer. Ent. Soc. xxx, 35, 1904.—N. States

semivittata Say, Het. N. Harm. 9, 1831 (*Pentatoma*); Compl. Writ. i, 322, 1859; Her. Schf., Wanz. Ins. vii, 93, 1844; Stål, Enum. Hem. ii, 34, 1872; Van Duzee, Trans. Amer. Ent. Soc. xxx, 34, 1904.—U. S.

pilipes Dallas, B. M. List Hem. i, 247, 1851.

VULSIREA.

Spinola, Essai, 350, 1837.

violacea Fabr., Syst. Rhyng. 167, 1803 (*Cimex*); Her. Schf., Wanz. Ins. iv, 89, 1839; Stål, Hem. Fabr. i, 31, 1868.—Fla.

abdominalis Westw., Hope Cat. i, 32, 1837.

tau Voll., Versl. Akad. Amst. Nat. (2), ii, 186, 1868.

ancora Spinola, Essai, 351, 1837.

nigro-rubra Spinola, ibid.

ponderosa Stål, Öfv. Vet. Akad. Förh. 1854, 233.

HALYDINÆ.

BROCHYMENA.

Amyot et Serville, Hém. 106, 1843.

affinis Van Duzee, Trans. Amer. Ent. Soc. xxx, 29, 1904.—W. States.

annulata Fabr., Syst. Ent. 704, 1775 (*Cimex*); Syst. Rhyng. 182, 1803; Stål, Enum. Hem. ii, 16, 1872; Van Duzee, Trans. Amer. Ent. Soc. xxx, 30, 1904; Insect Life, vii, fig. 17.—E. States.

serrota Wolff, Icones Cim. v, 184, 1811.

carolinensis Westw., Hope Cat. Hem. i, 22, 1837.

harrisii Uhler, Proc. Bost. Soc. Nat. Hist. xiv, 95, 1871; ibid, xix, 373, 1878.

arborea Say, Journ. Acad. Phil. iv, 311, 1825; Compl. Writ. ii, 239, 1859; Van Duzee, Trans. Amer. Ent. Soc. xxx, 27, 1904.—U. S. *erosa* Her. Schfr., Wanz. Ins. v, 70, 1839.

cariosa Stål, Enum. Hem. ii, 17, 1872; Van Duzee, Trans. Amer. Ent. Soc. xxx, 30, 1904.—S. States.

laticornis Say, Het. N. Harm. 4, 1831; Compl. Writ. i, 315, 1859.—U. S.

marginella Stål, Enum. Hem. ii, 16, 1872; Van Duzee, Trans. Amer. Ent. Soc. xxx, 31, 1904.—Car., Fla., Tex.

myops Stål, Enum. Hem., ii, 16, 1872.—S. States.

quadripustulata Her. Schfr., Wanz. Ins. vii, 57, 1844 (nec Fabr.).

- obscura** Her. Schf., Wanz. Ins. v, 68, 1839 (*Halys*); vii, 59, 1844; Van Duzee, Trans. Amer. Ent. Soc. xxx, 29, 1904.—W. States.
tenebrosa Walk., B. M. Cat. Het. i, 231, 1867.
- poeyi** Guérin, Sagra's Hist. de Cuba, Ins. 151, 1857; Stål, Berl. Ent. Zeit. 1866, 156; Enum. Hem. ii, 17, 1872; Van Duzee, Trans. Amer. Ent. Soc. xxx, 28, 1904.—Fla.
- punctata** Van Duzee, Can. Entom. 1909, 369.—Ga., Va.
harrisii Van Duzee, Trans. Amer. Ent. Soc. xxx, 31, 1904 (not Uhler).
- quadripustulata** Fabr., Syst. Ent. 704, 1775 (*Cimex*); Syst. Rhynch., 182, 1803; Stål, Enum. Hem. ii, 16, 1872; Van Duzee, Trans. Amer. Ent. Soc. xxx, 28, 1904.—U. S.
serrata Pal.-Beauv., Ins. Afr. et Amer., 187, 1805.
pupillata Her. Schfr., Wanz. Ins. iv, 104, 1839.

GRAPHOSOMINÆ.

ONCOZYGIA.

- Stål, Enum. Hem. ii, 15, 1872.
- clavicornis** Stål, Enum. Hem. ii, 16, 1872.—Tex.

PODOPS.

- Laporte, Essai, 72, 1832.
- Amaurochrous* Stål, Enum. Hem. ii, 15, 1872.
- cinctipes** Say, Amer. Ent., pl. 43, 1828 (*Tetyra*); Comp. Writ. 1, 94, 1859; Van Duzee, Trans. Amer. Ent. Soc. xxx, 22, 1904.—E. States.
- dubia* Germar, Zeitschr. Ent. i, 64, 1839 (not of Pal. Beauv.); Her. Schfr., Wanz. Ins. v, 44, 1839.
- dubius** Pal. Beauv., Ins. Afr. et Am. 33, 1805 (*Scutellera*); Stål, Enum. Hem. ii, 15, 1872; Van Duzee, Trans. Amer. Ent. Soc. xxx, 77, 1904.—Va.
- parvulus** Van Duzee, Trans. Amer. Ent. Soc. xxx, 22, 1904.—N. States.

WEDA.

- Schouteden, Ann. Soc. Ent. Belg. 1905, 145.
- horvathi** Schouteden, Ann. Soc. Ent. Belg. 1905, 145.—Colo.

DISCOCEPHALINÆ.

DISCOCEPHALA.

- Laporte, Essai, 57, 1832.
- clypeata** Stål, Stett. Ent. Zeit. 1862, 96; Enum. Hemipt. ii, 6, 1872; Distant, Biol. C.-Am. Rhynch. Het. i, 45, 1880.—Tex.
- inobtrusa* Walk., Cat. Hem. B. M. i, 183, 1867.
- notulata** Stål, Stett. Ent. Zeit. 1862, 96; Distant, Biol. Cent.-Amer. Rhynch. Het. i, 45, 1880.—Ariz.

CYRTOCORINÆ.

CYRTOCORIS.

White, Trans. Ent. Soc. Lond. 1842, 89.

trigonus Germar, Zeitschr. Ent. i, 44, 1839.—Calif.

ASOPINÆ.

APETETICUS.

Dallas, List Hem. B. M. i, 105, 1851.

Podisus Her. Schf., Wanz. Ins. ix, 296, 1853.

Tylospilus Stål, Enum. Hem. i, 52, 1870.

Eupodisus Schout., Gen. Insect. Fasc: 52, 71, 1907.

acutissimus Stål, Enum. Hem. i, 53, 1870; Distant, Biol. Cent.-Amer. Rhynch. Het. i, 40, 1880; Van Duzee, Trans. Amer. Ent. Soc. xxx, 72, 1904.—S. States.

bracteatus Fitch, Third Rept. Nox. Ins. 18, 1856 (*Arma*); Heid., Proc. Ent. Soc. Wash. vi, 10, 1904; Van Duzee, Trans. Amer. Ent. Soc. xxx, 70, 1904.—N. States.

crocatu Uhler, Trans. Md. Acad. Sci. 1897, 384.

cynicus Say, Het. N. Harm. 3, 1831 (*Pentatoma*); Compl. Writ. i, 312, 1859; Heid., Proc. Ent. Soc. Wash. vi, 9, 1904; Van Duzee, Trans. Amer. Ent. Soc. xxx, 70, 1904.—N. States.

grandis Dallas, List Hem. B. M. i, 96, 1851 (*Arma*).

fuscescens Dallas, Brit. Mus. List, Hemipt. i, 102, 1851 (*Arma*) Distant, Biol. C.-Am. Rhynch. Het. i, 38, 1880.—Tex.

gillettei Uhler, Bull. 31, Colo. Exp. Sta., 12, 1896; Van Duzee, Trans. Amer. Ent. Soc. xxx, 69, 1904.—Colo.

lineolatus Her. Schfr., Wanz. Ins. v, 69, 1839 (*Halys*); Stål, Enum. Hemipt. ii, 129, 1872; Distant, Biol. C.-Am. Rhynch. Het. i, 36, 1880.—Tex.

halys Dallas, Brit. Mus. List, Hemipt. i, 105, 1851; Stål, Enum. Hemipt. i, 49, 1870.

punctipennis Her. Schfr., Wanz. Ins. ix, 338, 1853.

maculiventris Say, Descr. Amer. Ins. La. 11, 1831 (*Pentatoma*); Psyche, viii, 307, 1899; Van Duzee, Trans. Amer. Ent. Soc. xxx, 71, 1904.—U. S.

spinosus Dallas, List Hem. B. M. i, 98, 1851; Cir. 87 Bur. Entom. fig. 6.

modestus Dallas, List Hem. B. M. i, 101, 1851 (*Arma*); Dist., Biol. Cent.-Amer. Rhynch. Het. i, 38, 1880; Van Duzee, Trans. Amer. Ent. Soc. xxx, 71, 1904.—N. States.

aggressor Walk., Cat. Het. B. M. ii, 359, 1867 (*Rhaphigaster*).

mucronatus Uhler, Trans. Maryl. Acad. Sci. 1897, 386; Van Duzee, Trans. Amer. Ent. Soc. xxx, 72, 1904.—Fla. .

pallens Stål, Freg. Eugen. Resa, Ins. 222, 1859 (*Arma*); Van Duzee, Trans. Amer. Ent. Soc. xxx, 72, 1904.—Calif.

Acc.
To Parshley

Apeticus

Apeticus

Podisus

Podisus

- placidus** Uhler, Amer. Ent. ii, 203, 1870; Can. Ent. xxix, 116, 1897; Van Duzee, Trans. Amer. Ent. Soc. xxx, 71, 1904.—N. States.
- sagitta** Fabr., Ent. Syst. iv, 99, 1794 (*Cimex*); Syst. Rhyng., 161, 1803; Distant, Biol. Cent.-Amer. Rhynch. Het. i, 37, 1880; Van Duzee, Trans. Amer. Ent. Soc. xxx, 72, 1904.—Tex.
- didymus* Pal. Beauv., Ins. 128, 1805.
- monospilus* Walk., Cat. Het. B. M. i, 136, 1867.
- sereiventris** Uhler, Proc. Bost. Soc. Nat. Hist. xiv, 94, 1871; Van Duzee, Trans. Amer. Ent. Soc. xxx, 71, 1904.—U. S.

Podisus

EUTHYRHYNCHUS.

- Dallas, List Hem. B. M. i, 77, 1851.
- floridanus** Linn., Syst. Nat. xii, 719, 1767 (*Cimex*); Fabr., Syst. Rhyng. 158, 1803; Her. Schf., Wanz. Ins. iv, 102, 1839; Stål, Enum. Hem. i, 54, 1870; Dist., Biol. Cent.-Amer. Rhynch. Het. i, 21, 1880.—S. States.
- emarginatus* Say, Het. New Harm. 3, 1831; Compl. Writ. i, 313, 1859.
- trivittatus* Her. Schf., Wanz. Ins. iv, 103, 1839.
- coloratus* Walk., Cat. Het. B. M. i, 136, 1867.

MINEUS.

- Stål, Öfv. Vet. Akad. Förh. 1867, 498.
- strigipes** Her. Schf., Wanz. Ins. ix, 338, 1853 (*Asopus*); Van Duzee, Trans. Amer. Ent. Soc. xxx, 67, 1904.—S. States.

MUTYCA.

- Stål, Rio Jan. Hem. ii, 58, 1862.
- grandis** Dallas, List Hem. B. M. i, 96, 1851 (*Arma*); Fitch, 3d Rept. N. Y. 17, 1856; Van Duzee, Trans. Amer. Ent. Soc. xxx, 72, 1904.—Fla.
- phymatophora** Pal. Beauv., Ins. Afr. et Amer. 112, 1805; Guérin, Sagra's Hist. de Cuba, Ins. 364, 1857.—Fla
- grandis* Dallas, List Hem. B. M. i, 91, 1851 (*Canthecona*).

OPLOMUS.

- Spinola, Essai, 355, 1837.
- dichrous** Her. Schf., Wanz. Ins. iv, 89, 1839; Stål, Enum. Hem. i, 26, 1870; Distant, Biol. Cent.-Amer. Rhynch. Het. i, 29, 1880; Van Duzee, Trans. Amer. Ent. Soc. xxx, 64, 1904.—Ariz.
- mundus** Stål, Stett. Ent. Zeit. 1862, 87; Enum. Hemipt. i, 30, 1870; Distant, Biol. C.-Am. Rhynch. Het. i, 32, 1880.—Tex.

PERILLUS.

Stål, Stett. Ent. Zeit. 1862, 88.

Perilloides Schouteden, Gen. Ins. fasc. 52, 37, 1907.

bioculatus Fabr., Syst. Ent. 715, 1775 (*Cimex*); Syst. Rhyng. 175, 1803; Stål, Enum. Hem. ii, 129, 1872; Van Duzee, Trans. Amer. Ent. Soc. xxx, 66, 1904.—W. States.

clauda Say, Journ. Acad. Phil. iv, 313, 1825; Compl. Writ. ii, 240, 1859.

circumcinctus Stål, Stett. Ent. Zeit. xxiii, 89, 1862; Dist., Biol. Cent.-Amer. Rhynch. Het. i, 34, 1880; Van Duzee, Trans. Amer. Ent. Soc. xxx, 67, 1904.—N. States.

marginatus Prov., Nat. Can. iv, 74, 1872.

confluens Her. Schf., Wanz. Ins. v, 77, 1839 (*Asopus*); Stål, Stett. Ent. Zeit. 1862, 88; Distant, Biol. Cent.-Amer. Rhynch. Het. i, 33, 1880; Van Duzee, Trans. Amer. Ent. Soc. xxx, 65, 1904.—W. States.

exaptus Say, Journ. Acad. Phil. iv, 313, 1825 (*Pentatoma*); Compl. Writ. ii, 240, 1859; Uhler, Proc. Ent. Soc. Phil. i, 23, 1861; Van Duzee, Trans. Amer. Ent. Soc. xxx, 65, 1904.—N. States.

marginella Dallas, List, Hem. B. M. i, 109, 1851.

variegata Kirby, Rich. Faun. Bor. Amer. iv, 276, 1837.

splendidus Uhler, Proc. Ent. Soc. Phil. i, 22, 1861 (*Zicrona*); Van Duzee, Trans. Amer. Ent. Soc. xxx, 66, 1904.—W. States.

RHACOGNATHUS.

Fieber, Europ. Hem. 347, 1861.

americanus Stål, Enum. Hem. i, 33, 1870; Van Duzee, Trans. Amer. Ent. Soc. xxx, 68, 1904.—Ohio, Can.

STIRETRUS.

Laporte, Essai, 75, 1832.

Oncogaster Stål, Enum. Hem. i, 24, 1870.

anchorago Fabr., Ent. Syst. 86, 1794 (*Cimex*); Syst. Rhyng. 137, 1803; Distant, Biol. Cent.-Amer. Rhynch. Het. i, 28, 1880; Van Duzee, Trans. Amer. Ent. Soc. xxx, 63, 1904; Cir., 102 Bur. Ent., fig. 5.—S. States.

diana Fabr., Syst. Rhyng. 137, 1803.

finbriatus Say, Amer. Ent., pl. 43, 1828; Compl. Writ. i, 93, 1859.

personata Germ., Zeitschr. Ent. i, 16, 1839.

ornatus Dallas, List Hem. B. M. i, 80, 1851.

pulchellus Westw., Hope Cat. i, 42, 1837.

variegatus Her. Schf., Wanz. Ins. iv. 90, 1839.

violaceus Say, Amer. Ent., pl. 43, 1828; Compl. Writ. i, 94, 1859.

ZIRCONA.

Amyot et Serville, Hém. 86, 1843.

cærulea Linn., Syst. Nat. x, 445, 1758 (*Cimex*); Hahn, Wanz. Ins. ii, 65, 1834; Van Duzee, Trans. Amer. Ent. Soc. xxx, 68, 1904.

—W. States. *New England* also, — C. M. P.

concinna Westw., Hope Cat. i, 39, 1837.

violacea Westw., *ibid.*

cuprea Dallas, List Hem. B. M. i, 108, 1851.

illustris A. and S., Hémipt., 87, 1843.

SCUTELLERIDÆ.**ACANTHOLOMA.**

Stål, Cefv. Vet. Akad. Förh. 1867, 491.

denticulata Stål, Enum. Hem. i, 17, 1870; Van Duzee, Trans. Amer. Ent. Soc. xxx, 77, 1904.—Ill., Kans.

CAMIRUS.

Stål, Rio Jan. Hem. (2) 57, 1862.

Zophæssa Dallas, List Hem, i, 43, 1851 (preoc.).

consocius Uhler, Bull. U. S. Geol. and Geog. Surv. Terr. ii, 274, 1876 (*Zophæssa*).—Ariz., Tex., Cal.

porosus Germar, Zeitschr. Ent. i, 108, 1839; Van Duzee, Trans. Amer. Ent. Soc. xxx, 16, 1904.—S. States.

CHELYSOMA.

Bergroth, Rev. d'Entom. 1891, 235.

Orsilochus Stål, Cefv. Vet. Akad. Förh. 1867, 493 (preoc.).

guttatus Her. Schf., Wanz. Ins. iv, 4, 1839 (*Pachycoris*); Stål, Enum. Hem. i, 9, 1870.—S. C., Fla.

DIOLCUS.

Mayr, Verh. zool.-bot. Ges. Wien. 1864, 904.

chrysorrhæus Fabr., Syst. Rhyng. 138. 1803 (*Tetyra*); Her. Schfr., Wanz. Ins. v, 6, 1839; Van Duzee, Trans. Amer. Ent. Soc. xxx, 12, 1904.—S. States.

viridipunctatus Say, Hem. N. Harm. i, 1831 (*Scutellera*); Compl. Writ. i, 310, 1859.

irroratus Fabr., Syst. Ent. 699, 1775; Syst. Rhyng. 136, 1803 (*Scutellera*); Her. Schf., Wanz. Ins. iv, 55, 1839; Stål, Hem. Fabric. i, 13, 1868; Van Duzee, Trans. Amer. Ent. Soc. xxx, 12, 1904.—Fla.

nebulosus Pal. Beauv., Ins. Afric. et Amer. 32, 1805.

hebraicus Pal. Beauv., *ibid.* 81.

cordigera Pal. Beauv., *ibid.* 234.

flavescens Westw., Hope Cat. i, 12, 1837.

EUPTYCHODERA.

Bergroth, Mém. Soc. Ent. Belg. xv, 143, 1908.

Ptychodera Reut. Act. Soc. Sci. Fenn. xxxiii (8), 44, 1906 (preoc.).

corrugata Van Duzee, Trans. Amer. Ent. Soc. xxx, 16, 1904 (*Phinodera*); Reut., Act. Soc. Sci. Fenn. xxxiii (8), 45, 1906.—Colo.

EURYGASTER.

Laporte, Essai, 69, 1832.

alternatus Say, Amer. Ent., pl. 43, 1828 (*Tetyra*); Compl. Writ. i. 94, 1859; Stål, Enum. Hem. i, 18, 1870; Van Duzee, Trans. Amer. Ent. Soc. xxx, 18, 1904; Bull. 69 Minn. Sta., fig. 96.—U. S.

nicoletanensis Prov., Nat. Can. iv, 73, 1872.

carinatus Van Duzee, Trans. Amer. Ent. Soc. xxx, 18, 1904.—W. States.

HOMCEMUS.

Dallas, List Hem. B. M. i, 36, 1851.

ænifrons Say, Long's Exped. App. 299, 1824 (*Scutellera*); Compl. Writ. i, 199, 1859; Van Duzee, Trans. Amer. Ent. Soc. xxx, 13, 1904.—U. S.

exilis Her. Schfr., Wanz. Ins. iv, 5, 1839, fig. 346.

bijugis Uhler, Hayden's Geol. Surv. Montana, 393, 1872; Van Duzee, Trans. Amer. Ent. Soc. xxx, 13, 1904.—W. States.

consors Uhler, Bull. U. S. Geol. and Geogr. Surv. Terr. ii, 272, 1876.

grammicus Wolff, Icones Cim. 172, 1811 (*Cimex*); Van Duzee, Trans. Amer. Ent. Soc. xxx, 14, 1904.—S. States.

parvulus Germar, Zeitschr. Ent. i, 107, 1839; Her. Schfr., Wanz. Ins. v, 54, 1839.

proteus Stål, Stett. Ent. Zeit. 1862, 82; Distant, Biol. C.-Am. Rhynch. Het. i, 20, 1880; Van Duzee, Trans. Amer. Ent. Soc. xxx, 15, 1904.—Tex., Calif.

ODONTOSCELIS.

Germar, Entom, Zeitschr. i, 36, 1839.

Fokkeria Schout., Ann. Soc. Ent. Belg. 1904, 301.

Vanduzeeina Schout., Gen. Ins. fasc. 24, 85, 1904.

balli Van Duzee, Trans. Amer. Ent. Soc. xxx, 19, 1904.—Colo., Wyo.

producta Van Duzee, Trans. Amer. Ent. Soc. xxx, 20, 1904.—Colo.

crassa Schout., Ann. Soc. Ent. Belg. 1904, 301 (*Fokkeria*).

PACHYCORIS.

Burmeister, Handb. ii (1), 391, 1835.

torridus Scop., Ann. V., Hist. Nat. 110, 1772 (*Cimex*); Distant, Biol. C.-Am. Rhynch. Het. i, 14, 1880.—Calif.

linnæi Westw., Hope Cat. Hem. i, 12, 1837.

stållicii Uhler, Proc. Ent. Soc. Phila. ii, 159, 1863.

PHIMODERA.

Germar, Entom. Zeitschr., i, 60, 1839.

binotata Say, Long's Exped. App. 298, 1824 (*Scutellera*); Compl. Writ. i, 198, 1859; Reut., Act. Soc. Sci. Fenn. xxxiii (8), 42, 1906.—W. States.

torpida Walker, B. M. Cat. Het. i, 75, 1867; Uhler, Bull. U. S. Geol. and Geogr. Surv. Terr. ii, 274, 1876; Van Duzee, Trans. Amer. Ent. Soc. xxx, 16, 1904; Reut., Act. Soc. Sci., Fenn. xxxiii (8), 22, 1906.—Colo., Can.

var. *fusco-testacea* Reut., ibid, 23.

var. *obscura* Reut., ibid, 23.

torrida Reut., Act. Soc. Sci. Fenn. xxxiii (8), 17, 1906.—Nev.

SPHYROCORIS.

Mayr., Verh. zool-bot. Ges. Wien. 1864, 904.

obliquus Germar, Zeitschr. Ent. i, 94, 1839 (*Pachycoris*); Her. Schfr., Wanz. Ins. v, 54, 1839; Van Duzee, Trans. Amer. Ent. Soc. xxx, 15, 1904.—Ariz., Fla.

STETHAULAX.

Bergroth, Rev. d'Entom. 1891, 235.

Aulacostethus Uhler, Proc. Bost. Soc. Nat. Hist. xiv, 93, 1871 (preoc.).

marmoratus Say, Hem. N. Harm. 2, 1831 (*Tetyra*); Compl. Writ. i, 310, 1859; Van Duzee, Trans. Amer. Ent. Soc. xxx, 12, 1904.—S. States.

simulans Uhler, Bull. U. S. Geol. and Geogr. Surv. Terr. ii, 272, 1876; Van Duzee, Trans. Amer. Ent. Soc., xxx, 76, 1904.—Calif.

TETYRA.

Fabricius, Syst. Rhyng. 128, 1803.

Macraulax Dallas, List Hem. B. M. i, 36, 1851.

arcuata Fabr., Syst. iv, 83, 1794 (*Cimex*); Syst. Rhyng. 134, 1803; Stål, Hem. Fabric. i, 13, 1868; Van Duzee, Trans. Amer. Ent. Soc. xxx, 11, 1904.—Fla.

bipunctata Her. Schfr., Wanz. Ins. iv, 87, 1839; Dallas, B. M. List, Hem. i, 36, 1851.—S. States.

tristis Uhler, Proc. Ent. Soc. Phil. i, 21, 1861.

robusta Uhler, Trans. Md. Acad. Sci. 1897, 383.—Ariz.

CYDNIDÆ.**AMNESTUS.**

Dallas, List Hem. B. M. i, 110, 1851.

pusillus Uhler, Bull. U. S. Geol. Geogr. Surv. Terr. ii, 278, 1876; Sign., Ann. Soc. Ent. France, 1883, 372.—E. States.

spinifrons Say, Journ. Acad. Phil. iv, 316, 1825 (*Cydnus*); Compl. Writ. ii, 242, 1859; Uhler, Bull. U. S. Geol. and Geogr. Surv. Terr. iii, 370, 1877; Sign., Ann. Soc. Ent. France, 1883, 368.—U. S.

CYDNUS.

Fabricius, Syst. Rhyng. 184, 1803.

Microporus Uhler, Geol. Geogr. Surv. 1876, 275.

s. g. *Trichocoris* Uhler, *ibid*, 277.

s. g. *Rhytidoporus* Uhler, *ibid*, 1877, 380.

s. g. *Cryptoporus* Uhler, *ibid*, 381.

Æthus Dallas, List Hem. B. M. i, 110, 1851.

communis Uhler, Bull. U. S. Geol. and Geogr. Surv. Terr. iii, 379, 1877 (*Æthus*); Sign., Ann. Soc. Ent. France, 1882, 35.—Fla., Tex.

compactus Uhler, Bull. U. S. Geol. and Geogr. Surv. Terr. iii, 382, 1877; Sign., Ann. Soc. Ent. France, 1882, 41.—Tex.

conformis Uhler, Bull. U. S. Geol. and Geogr. Surv. Terr. ii, 277, 1876; *ibid*, iii, 372, 1877; Sign., Ann. Soc. Ent. France, 1881, 425.—Calif.

indentatus Uhler, Bull. U. S. Geol. and Geogr. Surv. Terr. iii, 380, 1877; Sign., Ann. Soc. Ent. France, 1882, 38.—Fla.

obliquus Uhler, Hayden's Surv. Mont. 1872, 394; Bull. U. S. Geol. and Geogr. Surv. Terr. iii, 373, 1877; Sign., Ann. Soc. Ent. France, 1882, 161.—W. States.

politus Sign., Ann. Soc. Ent. France, 1882, 36.—Calif.

testudinatus Uhler, Bull. U. S. Geol. and Geogr. Surv. Terr. ii, 276, 1876; *ibid*, iii, 374, 1877; Sign., Ann. Soc. Ent. France, 1881, 424.—Calif.

CYRTOMENUS.

Amyot et Serville, Hém. 90, 1843.

castaneus Am. et Serv., Hémipt. 91, 1843.—Tex., Ariz.

crassus Walk., B. M. Cat. Het. i, 147, 1867.

obtusus Uhler, Bull. U. S. Geol. and Geogr. Surv. Terr. iii, 369, 1877.

mirabilis Perty, Delect. Ins. 166, 1834 (*Cydnus*); Sign., Ann. Soc. Ent. France, 1881, 199.—U. S.

mutabilis Uhler, Bull. U. S. Geol. and Geogr. Surv. Terr. iii, 367, 1877 (not of Perty).

teter Spinola, Essai, 332, 1837 (*Cydnus*); Distant, Biol. C.-Am. Rhynch. Het. i, 2, 1880; Sign., Ann. Soc. Ent. France, 1881, 197.—Calif.

æthiops Am. et Serv., Hémipt. 92, 1843.

GEOTOMUS.

Mulsant et Rey, Pun. Pentat. 34, 1866.

Melanæthus Uhler, Geol. Geogr. Surv. iii, 390, 1877.

parvulus Sign., Ann. Soc. Ent. France, 1883, 208.—W. States.

elongatus Uhler, Bull. U. S. Geol. and Geogr. Surv. Terr. 1872, 14; *ibid*, iii, 393, 1877 (not of Her. Schf.).

pennsylvanicus Sign., Ann. Soc. Ent. France, 1883, 207.—E. States.

picinus Uhler, Bull. U. S. Geol. and Geogr. Surv. Terr. iii, 391, 1877 (not of Stål, 1853).

punctatissimus Sign., Ann. Soc. Ent. France. 1883. 216.—Alaska.

robustus Uhler, Bull. U. S. Geol. and Geogr. Surv. Terr. iii, 390, 1877; Sign., Ann. Soc. Ent. France, 1883, 59.—E. States.

subglaber Walk., B. M. Cat. Het. i, 150, 1867.—N. Am.

uhleri Sign., Ann. Soc. Ent. France, 1883, 211.—S. States.

HOMALOPORUS.

Uhler, Geol. Geogr. Surv. iii, 377, 1877.

congruus Uhler, Bull. U. S. Geol. and Geogr. Surv. Terr. iii, 377, 1877; Sign., Ann. Soc. Ent. France, 1881, 330.—Colo., Tex.

LOBONOTUS.

Uhler, Geol. Geogr. Surv. iii, 395, 1877.

anthracinus Uhler, Bull. U. S. Geol. and Geogr. Surv. Terr. iii, 395, 1877; Distant, Biol. C.-Am. Rhynch. Het. i, 9, 1880; Sign., Ann. Soc. Ent. France, 1883, 529.—Tex.

MACROPORUS.

Uhler, Geol. Geogr. Surv. ii, 1876, 278.

repetitus Uhler, Bull. U. S. Geol. and Geogr. Surv. Terr. ii, 279, 1876; *ibid*, iii, 379, 1877; Sign., Ann. Soc. Ent. France, 1881, 329.—Calif.

PANGÆUS.

Stål, Stett. Ent. Zeit. 1862, 95.

bilineatus Say, Journ. Acad. Phil. iv, 315, 1825 (*Cydnus*); Compl. Writ., ii, 242, 1859; Uhler, Bull. U. S. Geol. and Geogr. Surv. Terr. iii, 383, 1877; Distant, Biol. C.-Am. Rhynch. Het. i, 6, 1880; Sign., Ann. Soc. Ent. France, 1882, 254.—U. S.

femoralis Her. Schf., Wanz. Ins. v, 98, 1839 (*Cydnus*).

robertsoni Sign., Ann. Soc. Ent. France, 1882, 254.

discrepans Uhler, Bull. U. S. Geol. and Geogr. Surv. Terr. iii, 386, 1877; Sign., Ann. Soc. Ent. France, 1882, 249.—W. States.

margo Dallas, B. M. List Hem. i, 116, 1851; Stål, Stett. Ent. Zeit. 1862, 95; Uhler, Bull. U. S. Geol. and Geogr. Surv. Terr. iii, 388, 1877; Sign., Ann. Soc. Ent. France, 1882, 248.—Ariz.

- piceatus** Stål, Stett. Ent. Zeit. 1862, 96; Uhler, Bull. U. S. Geol. and Geogr. Surv. Terr. iii, 388, 1877; Sign., Ann. Soc. Ent. France, 1882, 262.—Calif.
insularis Dallas, B. M. List Hem. i, 120, 1851.
tenuis Walk., B. M. Cat. Het. i, 151, 1867.
spanbergi Sign., Ann. Soc. Ent. France, 1882, 259.—Tex.
uhleri Sign., Ann. Soc. Ent. France, 1882, 253.—Ga., S. Car.
rufifrons Uhler, Bull. U. S. Geol. and Geogr. Surv. Terr. iii, 384, 1877 (not of Her. Schfr.).

SEHIRUS.

- Amyot et Serville, Hém., 96, 1843.
Canthophorus Mul. et Rey, Pentat. 54, 1866.
cinctus Pal.-Beauv., Ins. Afr. et Amer., 114, 1805 (*Cydnus*); Uhler, Bull. U. S. Geol. and Geogr. Surv. Terr. iii, 397, 1877; Sign., Ann. Soc. Ent. France, 1884, 60; Bull. 69 Minn. Sta., fig. 94.—U. S.
ligatus Say, Het. N. Harm. 10, 1831; Compl. Writ. i, 322, 1859 (*Cydnus*).
albonotatus Dallas, B. M. List Hem. i, 127, 1851.

THYREOCORIDÆ.

THYREOCORIS.

- Schrank, Fauna Boica, ii, 46, 1801.
Corimelæna White, Mag. Nat. Hist. 1839, 3.
Coreomelas Amyot et Serv., Hém. 67, 1843.
Galgupha ibid, 68.
anthracina Uhler, Bull. U. S. Geol. and Geogr. Surv. Terr. ii, 270, 1876; Van Duzee, Trans. Amer. Ent. Soc. xxx, 6, 1904.—W. States.
cærulescens Stål, Stett. Ent. Zeit. 1862, 94; Distant, Biol. C.-Am. Rhynch. Het. i, 10, 1880; Van Duzee, Trans. Amer. Ent. Soc. xxx, 7, 1904.—W. States.
cyaneonigra Walk., B. M. Cat. Het. i, 78, 1867.
cyanea Uhler, Bull. U. S. Geol. and Geogr. Surv. Terr. ii, 270, 1876.
ciliata Uhler, Proc. Ent. Soc. Phil. ii, 156, 1863; Bull. U. S. Geol. and Geogr. Surv. Terr. ii, 270, 1876; Van Duzee, Trans. Amer. Ent. Soc. xxx, 6, 1904.—W. States.
denudata Uhler, Proc. Ent. Soc. Phil. ii, 155, 1863; Van Duzee, Trans. Amer. Ent. Soc. xxx, 7, 1904.—La.
extensa Uhler, Proc. Ent. Soc. Phil. ii, 155, 1863; Van Duzee, Trans. Amer. Ent. Soc. xxx, 9, 1904.—W. States.
guttiger Stål, Stett. Ent. Zeit. 1862, 94; Distant, Biol. C.-Am. Rhynch. Het. i, 11, 1880.—Tex.

- lateralis** Fabr., Syst. Rhynch. 142, 1803 (*Cimex*); Germar, Zeitschr. Ent. i, 39, 1839.—E. States.
gillettei Van Duzee, Trans. Amer. Ent. Soc. xxx, 8, 1904.
- marginella** Dallas, B. M. List Hem. i, 59, 1851; Van Duzee, Trans. Amer. Ent. Soc. xxx, 9, 1904.—Can.
- minuta** Uhler, Proc. Ent. Soc. Phil. ii, 155, 1863; Van Duzee, Trans. Amer. Ent. Soc. xxx, 76, 1904.—Fla.
- montanus** Van Duzee, Ent. News. 1909, 231.—W. States.
lateralis Van Duzee, Trans. Amer. Ent. Soc. xxx, 8, 1904 (nec Fabr.).
- nigra** Dallas, B. M. List Hemipt. i, 57, 1851; Van Duzee Trans. Amer. Ent. Soc. xxx, 6, 1904.—N. States.
- nitiduloides** Wolff, Icones. Cim. iii, 98, 1802 (*Cimex*); Germar, Zeitschr. Ent. i, 37, 1839; Distant, Biol. C.-Am. Rhynch. Het. i, 10, 1880; Van Duzee, Trans. Amer. Ent. Soc. xxx, 5, 1904.—U. S.
histeroides Say, Het. New Harm. 2, 1831; Comp. Writ. i, 311, 1859.
- pulicaria** Germar, Zeitschr. Ent. i, 39, 1839; Dallas, List B. M. Hem. i, 59, 1851; Van Duzee, Trans. Amer. Ent. Soc. xxx, 9, 1904.—U. S.
flavomarginata Thomas, Trans. Ill. St. Agric. Soc. v, 455, 1865.
- renormata** Uhler, Hem. Colo. 11, 1895; Van Duzee, Trans. Amer. Ent. Soc. xxx, 9, 1904.—W. States.
- sayi** Van Duzee, Trans. Amer. Ent. Soc. xxx, 10, 1904 (*Corimelaena*) —W. States.
albipennis Say, Het. New Harm. 2, 1831; Compl. Writ. i, 311, 1859 (preoc.).
- unicolor** Pal. Beauv., Ins. Afr. et Am. 33, 1805 (*Cimex*); Germar, Zeitschr. Ent. i, 37, 1839; Van Duzee, Trans. Amer. Ent. Soc. xxx, 4, 1904.—U. S.
helopioides Wolff, Icones. Cim. v, 174, 1811.
atra Am. et Serv., Hémipt. 68, 1843.

ADDITIONS.

Family **NEPIDÆ**, p. 8.

Ranatra brevicollis Montd., Bull. Soc. Sci. Bucarest. xviii, 184, 1910.
 —Calif.

Ranatra protensa Montd., *ibid*, 185.—N. Y.

Curicta howardi Montd., *ibid*, 181.—Texas.

Later Montandon (Bull. Soc. Sci. Bucarest xix, 62, 1910) puts *Ranatra kirkaldyi* Bueno as a synonym of *R. fusca*, and gives a new name, *R. americana*, p. 65, for *R. quadridentata* Bueno (nec Stål), and describes variety *edentula*, p. 66, of *R. americana* from Pennsylvania and Texas.

Family **NAUCORIDÆ**, p. 10.

Ambrysus heidemanni Montd., Bull. Soc. Sci. Bucarest, xviii, 188, 1910.—Wyo.

Ambrysus pulchellus var. **pallidus** Montd., ibid, 189.—Texas.

ERRATA.

Page 8, for *Deinstoma* read *Deinostoma*.

Page 13, under *Arilus*, for *Prionotus* Uhler, read *Prionidus* Uhler.

Page 15, for *Repita* read *Repipta*.

Page 23, top, for *rufusculiss* read *rufusculus*.

Page 27, for *Macovelia* read *Macrovelia*.

Page 29, bottom, at side, for *Laodamina* Kirkaldy read *Laodamia* ;
Laodamina, centered, is a new name.

Page 47, bottom, for *angustulus* read *angustulus*.

Page 50, near top, for *decnes* read *decens*.

Page 9, for *Ofr.* read *Æfv.*

INDEX TO GENERA.

Abedus	8	<i>Arctocouixa</i>	5, 6
Acalypta	55	Arhappe	70
Acanthia	11, 25	Arilus	13
Acanthocephala	79	<i>Aristoreuteria</i>	31
Acanthochila	55	Arphnus	58
Acantholoma	96	Arvelius	82
Acanthophysa	71	<i>Asciodema</i>	34
Acanthosoma	82	<i>Askra</i>	13, 84
Acholla	13	<i>Aspidotoma</i>	57
Acinocoris	69	<i>Astemma</i>	63
<i>Acrocoris</i>	18	Asthenidea	23
<i>Actractophora</i>	60	Atheas	55
Acysta	55	Atomoscelis	29
Adelphocoris	42	Atomosira	82
Ælia	82	Atrachelus	14
<i>Æthus</i>	99	Atractotomus	29
Agalliaestes	28, 29	Aufeis	71
Agrammodes	55	<i>Aulacostethus</i>	98
Alepidia	36		
Althos	75	Banasa	82
Alydus	74	Barce	20
<i>Amaurochrous</i>	92	Belonochilus	60
<i>Amaurosterphus</i>	57	Belostoma	9
Amblyomia	77	Benacus	9
Amblytylus	31	<i>Beosus</i>	67
Ambrysus	10, 103	<i>Berytus</i>	71
Amnestus	98	Blissus	58
<i>Amorgius</i>	9	<i>Bolteria</i>	38
Anasa	76	<i>Botocudo</i>	63
Aneurus	54	<i>Brachycoleus</i>	49
<i>Anisops</i>	7	<i>Brachyrhynchus</i>	54
<i>Anisoscelis</i>	78	<i>Brachytropis</i>	41
<i>Anoterops</i>	32	<i>Breddiniessa</i>	33
Anthocoris	23	Brepholoxa	83
<i>Antillocoris</i>	63	Brochymena	91
Apeteticus	93	Buenoa	7
<i>Aphanus</i>	64, 67		
Apiomerus	16	<i>Callicapsus</i>	42
<i>Apocremnus</i>	31	Callicorixa	6
Aradus	52	<i>Callimiris</i>	42
Archimerus	79		

<i>Callodemas</i>	47	<i>Coreocoris</i>	81
<i>Calisius</i>	54	<i>Coreomelas</i>	101
<i>Calocoris</i>	42	<i>Coreus</i>	73, 75, 76, 80, 81
<i>Camirus</i>	96	<i>Corimelæna</i>	101
<i>Campptobrochis</i>	42	<i>Coriscus</i>	22
<i>Camptopus</i>	74	<i>Corisa</i>	6
<i>Campylomma</i>	29	<i>Corixa</i>	5
<i>Canthophorus</i>	101	<i>Corizus</i>	72
<i>Capanus</i>	80	<i>Corynocoris</i>	81
<i>Capsus</i>	43, 51	<i>Corythuca</i>	55
<i>Cardiastethus</i>	23	<i>Cosmopepla</i>	83
<i>Carpilis</i>	63	<i>Creontiades</i>	40
<i>Carpocoris</i>	83	<i>Crinocerus</i>	80
<i>Carthasis</i>	21	<i>Crophius</i>	59
<i>Catorhintha</i>	76	<i>Cryphula</i>	64
<i>Caulatops</i>	50	<i>Cryptoporus</i>	99
<i>Centromelus</i>	19	<i>Curicta</i>	102
<i>Cephalocoris</i>	28	<i>Cydnus</i>	99
<i>Ceraleptus</i>	73	<i>Cylapus</i>	37
<i>Cerascopus</i>	21	<i>Cylindromelus</i>	32
<i>Ceratocapsus</i>	32	<i>Cylloceps</i>	31
<i>Ceratocombus</i>	25	<i>Cyllocoris</i>	33
<i>Chariesterus</i>	81	<i>Cymodema</i>	58
<i>Chelinidea</i>	76	<i>Cymus</i>	58
<i>Chelysoma</i>	96	<i>Cyrtocapsus</i>	50
<i>Chiroleptes</i>	66	<i>Cyrtocoris</i>	93
<i>Chlamydatus</i>	29	<i>Cyrtomenus</i>	99
<i>Chlorochroa</i>	88	<i>Cyrtopeltis</i>	37
<i>Chlorocoris</i>	83	<i>Cyrtopeltocoris</i>	35
<i>Chondrocera</i>	77	<i>Cyrtorrhinus</i>	33
<i>Cimex</i>	25	<i>Dacerla</i>	35
<i>Cimolus</i>	77	<i>Dacota</i>	50
<i>Cistalia</i>	63	<i>Darbanus</i>	16
<i>Cletus</i>	77	<i>Darmistus</i>	74
<i>Cligenes</i>	63	<i>Dasycoris</i>	73
<i>Clinocoris</i>	25, 82	<i>Dasypterus</i>	23
<i>Clivinema</i>	50	<i>Deinostoma</i>	8
<i>Closterocoris</i>	35	<i>Delochilochoris</i>	64
<i>Closterotomus</i>	42	<i>Dendrocoris</i>	84
<i>Cnemodus</i>	63	<i>Deræocoris</i>	43
<i>Coccobaphes</i>	43	<i>Diactor</i>	79
<i>Cœnus</i>	83	<i>Diaditus</i>	19
<i>Collaria</i>	40	<i>Diaphnidia</i>	33
<i>Compsocerochoris</i>	47	<i>Dichrocysta</i>	56
<i>Conorhinus</i>	17	<i>Dichroosecytus</i>	43
<i>Coquilletia</i>	35		

Dicyphus	37	Ficana	77
Dilasia	24	Fitchia	14
Diolcus	96	<i>Fokkeria</i>	97
<i>Diommatus</i>	34	Fulvius	36
<i>Diplacodus</i>	15	Galeatus	56
<i>Diplodus</i>	15	<i>Galeothus</i>	75
<i>Diplonotus</i>	66	<i>Galgulus</i>	10
Discocephala	92	<i>Galgupha</i>	101
<i>Discomerus</i>	52	Gardena	21
Dolichomerus	24	Garganus	44
<i>Dorachosa</i>	64	Gargaphia	56
Drymus	64	Gastrodes	64
Dycoderus	64	Gelastocoris	10
Dysdercus	70	Geocoris	68
<i>Dysodius</i>	54	Geotomus	100
Ecclitotarsus	50	Gerris	26
Ecertobia	44	Globiceps	33
Ectopiocerus	35	<i>Glænocorisa</i>	5
Ectrichodia	17	Glyptocombus	26
<i>Ectrychotes</i>	17	Gnathobleda	19
Edessa	84	Gonianotus	65
Elasmostethus	82	<i>Gonocerus</i>	76, 77, 81
Elasmucha	82	<i>Graptolomus</i>	60
Emblethis	64	<i>Hadrodema</i>	46
Emesa	20	Hadronema	39
<i>Engytatus</i>	37	Hæmatosiphon	25
Enicocephalus	21	Halobates	26
<i>Episcopus</i>	31	Halobatopsis	26
Eremocoris	64	<i>Halticocoris</i>	38
<i>Erythrischius</i>	62	Halticus	38
Esperanza	74	<i>Halys</i>	90
<i>Euagorus</i>	16	<i>Hammacerus</i>	17
<i>Euarmosus</i>	42	Hammatocerus	17
<i>Eucrocoris</i>	47	Harmostes	72
<i>Eupodisus</i>	93	Harpactor	17
Euptychodera	97	<i>Hebrus</i>	28
Europiella	29	Heidemannia	28
Eurychilopterella	44	<i>Helenotus</i>	60
Eurygaster	97	Heræus	65
<i>Euryophthalmus</i>	70	<i>Herega</i>	16
Euschistus	84	Heterocordylus	33
Eustictus	44	Heterogaster	60
Euthoctha	80	Heza	14
Euthyrhynchus	94	<i>Holcostethus</i>	89
Eysarcoris	86		

Homalocoris	17	Leptocoris	73
Homaloporus	100	Leptocoris	74
Homœmus	97	Leptoglossus	78
<i>Hoplinus</i>	71	<i>Leptomerocoris</i>	44
Hoplistoscelis	22	Leptophya	57
Hoplomachidea	31	<i>Leptopterna</i>	41
<i>Hoplomachus</i>	31	<i>Leptoscelis</i>	79
Horcias	44	Leptostyla	57
Hyaliodes	37	Leucopœcila	30
Hyalmenus	74	<i>Lichenobia</i>	25
Hydrometra	28	Ligyrocoris	65
<i>Hydrotrechus</i>	26	<i>Limnobates</i>	28
Hymenarcys	86	Limnogonus	26
<i>Hymenocoris</i>	21	Limnopus	26
<i>Hymenodectes</i>	21	<i>Limnotrechus</i>	26
Hyoidea	38	<i>Lioderma</i>	88
<i>Hypereides</i>	32	<i>Liotropis</i>	84
Hypselonotus	77	Lobonotus	100
<i>Hypselosoma</i>	26	<i>Lobostethus</i>	41
		Lomatopleura	38
		<i>Lophyrus</i>	42
Idolocoris	37	Lopidea	39
Ilnacora	33	Lopus	32
Irbisia	44	Loxa	86
Ischnodemus	58	Luteva	21
Ischnorhynchus	59	Lyctocoris	24
Isometopus	28	Lygæsuma	60
		Lygæus	60
<i>Jadera</i>	73	Lygidea	45
Jalysus	71	Lygus	45
<i>Kelidocoris</i>	33	<i>Macraulax</i>	98
<i>Klinophilos</i>	25	<i>Macrocapsus</i>	43
		Macrocephalus	51
		<i>Macrocoleus</i>	32
Labopidea	38	<i>Macrocorisa</i>	5
Labops	38	Macrolophus	37
Lampethusa	30	Macroporus	100
<i>Lampronotus</i>	64	<i>Macrosandalus</i>	18
<i>Lamproplex</i>	64	Macrotylus	32
Laodamia	29	Macrovelia	27
Largus	70	Madura	77
<i>Lasiacantha</i>	57	Malacocoris	34
Lasiochilus	24	<i>Margus</i>	75
<i>Lecadra</i>	70	Maurodactylus	30
Leptobyrsa	56	Meccus	18

INDEX.

v

<i>Mecidea</i>	86	<i>Nabicula</i>	22
<i>Mecomma</i>	34	<i>Nabidea</i>	40
<i>Mecorhamphus</i>	63	<i>Nabis</i>	22
<i>Megacælum</i>	44	<i>Næogeus</i>	28
<i>Megaloceroea</i>	41	<i>Nannium</i>	54
<i>Megalocoleus</i>	32	<i>Narnia</i>	78
<i>Megalotomus</i>	75	<i>Narvesus</i>	19
<i>Melanæthus</i>	100	<i>Neides</i>	71
<i>Melanocoryphus</i>	60	<i>Nematopus</i>	77
<i>Melanolestes</i>	18	<i>Neoborops</i>	46
<i>Melanopleurus</i>	60	<i>Neocapsus</i>	46
<i>Melanorhopala</i>	57	<i>Neoboris</i>	45, 49
<i>Melanostoma</i>	87	<i>Neocoris</i>	30
<i>Melinna</i>	32	<i>Neopropa</i>	37
<i>Menecles</i>	86	<i>Neottiglossa</i>	87
<i>Mermimerus</i>	32	<i>Neovelina</i>	27
<i>Merocoris</i>	81	<i>Nepa</i>	8
<i>Mesomiris</i>	41	<i>Nerthra</i>	11
<i>Mesovelina</i>	27	<i>Neurocolpus</i>	47
<i>Metacanthus</i>	71	<i>Neuroctenus</i>	54
<i>Metapodius</i>	79	<i>Nezara</i>	87
<i>Metatropiphorus</i>	22	<i>Ninus</i>	59
<i>Metriorrhynchomiris</i>	44	<i>Notonecta</i>	7
<i>Metriorrhynchus</i>	44	<i>Nysius</i>	62
<i>Metrobates</i>	27		
<i>Mezira</i>	54	<i>Ochetopus</i>	20
<i>Microphyllellus</i>	30	<i>Ochrostomus</i>	60
<i>Microporus</i>	99	<i>Odontoscelis</i>	97
<i>Microtoma</i>	65	<i>Œbalus</i>	90
<i>Micropus</i>	58	<i>Œdancala</i>	59
<i>Microvelia</i>	27	<i>Oncerometopus</i>	39
<i>Milyas</i>	14	<i>Oncerotrachelus</i>	19
<i>Mimoceps</i>	50	<i>Oncocephalus</i>	19
<i>Mineus</i>	94	<i>Oncogaster</i>	95
<i>Miris</i>	41	<i>Oncognathus</i>	49
<i>Monalocoris</i>	50	<i>Oncopeltus</i>	62
<i>Monanthia</i>	57	<i>Oncotylus</i>	32
<i>Mononyx</i>	11	<i>Oncozygia</i>	92
<i>Mormidea</i>	86	<i>Onychumenus</i>	32
<i>Mozena</i>	80	<i>Ophthalmicus</i>	68
<i>Murgantia</i>	87	<i>Oplomus</i>	94
<i>Mutya</i>	94	<i>Opsicætus</i>	14
<i>Mycterocoris</i>	42	<i>Orectoderus</i>	35
<i>Myochrocoris</i>	30	<i>Orsillus</i>	63
<i>Myodocha</i>	66	<i>Orsilochus</i>	96
<i>Myrmecopsis</i>	35	<i>Orthæa</i>	66

<i>Orthocephalus</i>	39	<i>Physatochila</i>	57
<i>Ortholomus</i>	62	<i>Phytocoris</i>	47
<i>Orthometrops</i>	21	<i>Pictinus</i>	55
<i>Orthotylus</i>	34	<i>Piesma</i>	57
<i>Oxycarenus</i>	59	<i>Piestosoma</i>	52
<i>Ozophora</i>	66	<i>Piezodera</i>	70
		<i>Piezodorus</i>	89
<i>Pachycoris</i>	97	<i>Piezostethus</i>	24
<i>Pachygrontha</i>	60	<i>Piezostoma</i>	52
<i>Pachylis</i>	78	<i>Pilophorus</i>	36
<i>Padæus</i>	88	<i>Pindus</i>	15
<i>Pagasa</i>	22	<i>Pirates</i>	18, 19
<i>Pallacocoris</i>	47	<i>Pirithous</i>	50
<i>Pamera</i>	66	<i>Plagiognathus</i>	30
<i>Pamerocoris</i>	36	<i>Platygaster</i>	64
<i>Pamillia</i>	36	<i>Platytylus</i>	40
<i>Pangæus</i>	100	<i>Plea</i>	8
<i>Paracalocoris</i>	47	<i>Plociomeræ</i>	65, 67
<i>Paraxenetus</i>	47	<i>Ploiaria</i>	21
<i>Paromius</i>	66	<i>Ploiariodes</i>	21
<i>Parthenicus</i>	34	<i>Ploiariopsis</i>	21
<i>Paryphes</i>	77	<i>Pnirontis</i>	19
<i>Pedinocoris</i>	8	<i>Podisus</i>	93
<i>Peliopelta</i>	60	<i>Podops</i>	92
<i>Peliosoma</i>	60	<i>Pœcilocapsus</i>	48
<i>Pelocoris</i>	10	<i>Pœciloscytus</i>	49
<i>Pelogonus</i>	10	<i>Pothea</i>	17
<i>Pentatoma</i>	88	<i>Prionidus</i>	13
<i>Peribalus</i>	89	<i>Prionosoma</i>	89
<i>Perigenes</i>	67	<i>Prionotus</i>	13
<i>Perilloides</i>	95	<i>Pronotacantha</i>	71
<i>Perillus</i>	95	<i>Prostemma</i>	22
<i>Peritrechus</i>	67	<i>Protentor</i>	75
<i>Peritropis</i>	36	<i>Protocantha</i>	71
<i>Perthostoma</i>	9	<i>Proxius</i>	55
<i>Petalocheirus</i>	17	<i>Proxys</i>	90
<i>Petalotoma</i>	77	<i>Psallus</i>	31
<i>Petissius</i>	67	<i>Pselliopus</i>	14
<i>Phenicocoris</i>	30	<i>Pseudopamera</i>	67
<i>Phimodera</i>	98	<i>Pseudoxenetus</i>	34
<i>Phintius</i>	11	<i>Pterotmetus</i>	67
<i>Phlegyas</i>	60	<i>Ptochiomera</i>	67
<i>Phorticus</i>	22	<i>Ptychodera</i>	97
<i>Phrygadicus</i>	60	<i>Pycnoderes</i>	51
<i>Phthia</i>	79	<i>Pycnopterna</i>	49
<i>Phymata</i>	52	<i>Pygæus</i>	63

Pygolampis	20	<i>Serinettha</i>	73
<i>Pyrrhocoris</i>	71	<i>Serphus</i>	8
Pyrrhotes	73	<i>Sigara</i>	5
<i>Quilnus</i>	52	Sinea	14
Ranatra	8, 102	Sirthenea	19
Rasahus	18	Sixeonotus	51
Reduviolus	22	Solubea	90
Reduvius	14	Spartocera	81
Repipta	15	Sphærobilus	68
Resthenia	40	<i>Sphærocista</i>	56
Reuteria	34	Sphragisticus	68
Reuteroscopus	31	<i>Sphyracephalus</i>	34
Rhacognathus	95	Sphyrocoris	98
Rhagovelia	27	<i>Spilaloniis</i>	19
<i>Rhaphigaster</i>	87	Spiniger	18
Rhaptus	67	Stachyocnemis	75
Rheumatobates	27	<i>Stenarops</i>	33
<i>Rhiginia</i>	17	<i>Stenarus</i>	29
Rhinacloa	31	Stenodema	44
Rhinocapsus	35	Stenomacra	70
<i>Rhinuchus</i>	79	Stenopoda	20
<i>Rhopalotomus</i>	43, 49	<i>Stenoscytus</i>	8
<i>Rhopalus</i>	72	Stenotus	49
<i>Rhynarius</i>	23	<i>Stephania</i>	27
Rhyparochromus	68	Stethaulax	98
<i>Rhytidolomia</i>	88	Stiphrosoma	39
<i>Rhytidoporus</i>	99	Stiretrus	95
Rocconota	15	<i>Strachia</i>	87
Sagotylus	80	<i>Strongylotes</i>	29
<i>Salacia</i>	63	<i>Syromastes</i>	72, 73, 75
Salda	11	<i>Syrtis</i>	52
Saldoida	13	Sysinas	38
Schumannia	20	<i>Systratiotus</i>	49
Sciocoris	90	Teleonemia	57
Scolopocerus	74	Teleorhinus	35
Scoloposcelis	24	Tempyra	68
Scolopostethus	68	Tenagobia	6
<i>Scutellera</i>	92	Teratocoris	41
Sehirus	101	Tetraphleps	24
Semium	39	<i>Tetrarhinus</i>	75
Sephina	81	Tetyra	92, 98
Sericophanes	36	<i>Thasus</i>	78
		<i>Theognis</i>	78
		Thyanta	90

Thyreocoris	101	Uhleriola	68
<i>Thyrilus</i>	44	<i>Valdasus</i>	37
Tingis	57	<i>Vanduzeeina</i>	97
Tinicephalus	32	<i>Velia</i>	27
Tiryas	35	Vulsirea	91
Tollius	75		
<i>Trachelomiris</i>	40	Weda	92
Trapezonotus	68	<i>Xenetes</i>	34
Trepobates	27	Xestocoris	68
<i>Trichia</i>	35	<i>Xuthus</i>	80
<i>Trichocoris</i>	99	<i>Xylocoris</i>	24
Trichopepla	91		
Trigonotylus	42	Zaitha	9
Triphleps	25	Zelus	15
<i>Trochopus</i>	27	Zircrona	96
Tropidosteptes	49	<i>Zophæssa</i>	96
Tuponia	31	<i>Zosmenus</i>	57
<i>Tylospilus</i>	93		
<i>Tytthus</i>	33

BOUND BY THE MD. BOOKBINDERY. HAGERSTOWN MD.

SMITHSONIAN INSTITUTION LIBRARIES

3 9088 00726 2314