1902

USNM

EX LIBRIS

William Healey Dall

Division of Mollusks
Sectional Library

Division of Mollusks
Sectional Library

423
A1K92
1864
Mo
John A. Verrill
from the Father Hy. Krebs
J. S. Thomas. U.S.
The West-Indian Marine Shells
H. M. Anders

with some remarks.

Presented to Smithsonian Institution
by John H. Redfield
Dec. 1884

a manuscript printed for circulation between
collectors.

Division of Mollusks
Sectional Library

by

Henrik Johannsen, Krebs

*

*

*

1864.

Index.

	Page.
Acmaea Eschh.	75.
Anomia L.	137.
Aplustrum Schm	92.
Aplysia L.	91.
Arca L.	124.
Artemis Poli	99.
Astralium Link.	82.
Atys Mtf.	94.
Avicula Brug.	131.
Buccinum L.	32.
Bulla L.	92.
Bullia Gray	33.
Bullinula Beck	92.
Calcar Mtf.	81.
Cancellaria Lmk.	33.
Capsa Brug.	105.
Capulus Mtf.	70.
Cardita Brug.	123.
Cardium L.	115.
Cassis Lmk.	34.
Cerithium Adans.	47.
Chama L.	117.
Chiton L.	89.
Cinulia Gray	71.
Coecum Flem.	75.
Columbella Lmk.	28.
Conopleura Hinds	15.
Conus L.	3.
Corbula Brug.	109.
Crassatella Lmk.	122.
Crepidula Lmk.	69.
Crucibulum Schm.	69.
Cumingia Sowb.	107.
Cyclas Brug.	107.
Cypraea L.	41.
Cypræacassis St.	35.
Daphnella Hinds	9-12.
Delphinula Lmk.	80.
Dentalium L.	91.
Diplodonta Brown	122.

	Page.
Dolium Lmk.	35.
Dreisena Beneden	130.
Eburnea Lmk.	36.
Emarginula Lmk.	85.
Erato Risso	78. 42
Eucharis Recl.	111.
Eulima Risso	73.
Euomphalus Sowb.	62.
Fasciolaria Lmk.	15.
Fossarus Adans.	58.
Fimbria M. de M.	122.
Fissurella Brug.	86.
Fusus Brug.	15.
Gastrochaena Sp.	114.
Glauconome Gray	107.
Globulus Schm.	82.
Ianthina Lmk.	65.
Lentilaria Schm.	121.
Lima Brug.	132.
Liotia Gray.	80.
Litiopa Rang.	52.
Littorina Ferr.	58.
Lobiger Krohn.	91.
Lophocercus Krohn.	91.
Lophorus Peli	89.
Lucina Brug.	128, 100-6.
Mactra L.	105.
Magdala Leach.	109.
Magnelia Risso	9-13.
Marginella Lmk.	43.
Melina Retz	132.
Mitra Lmk.	36.
Mitrularia Schm.	69.
Modiolus Lmk.	128.
Modulus Gray	62.
Monostygma Gray	71.
Murex L.	18.
Narica Recl.	68.
Nassa Lmk.	31.
Natica Adans.	66.

	Page.
Neaera Gray	111.
Nerita L.	76.
Neritina Lmk.	77.
Odontostoma Flem.	71.
Oliva Lmk.	38.
Oniscia Lmk.	35.
Ostrea L.	136.
Ovula Brug.	43.
Patella L.	88-90.
Pecten O. F. M.	133.
Pectunculus Lmk.	126.
Periploma Schm.	108.
Petricola Lmk.	108.
Phasionella Lmk.	79.
Pholadomya Br.	111.
Pholas L.	113.
Pinna L.	130.
Pisania Biv.	31.
Placunanomia Brod.	136.
Planaxis Lmk.	52.
Pleurotoma Lmk.	7.
Pleurotomaria Def.	85.
Plicatula Lmk.	135.
Psammobia Lmk.	104.
Psammosolea Risso	113.
Purpura Brug.	25.
Pyramidella Lmk.	71.
Pyrula Lmk.	18.
Ranella Lmk.	25.
Ricinula Lmk.	27.
Rissoa Flem.	53.
Rissoina d'Orb.	57.
Sanguinolaria Lmk.	104.
Saxicava Fl. de B.	107.
Scalaria Lmk.	63.
Serobicularia Schm.	107.
Semele Schm.	106.
Semicassis Klein.	34.
Semimarginula	86.
Sigaretus Lmk.	68.
Siliqvaria Brug.	75.

	Page.
Siphonaria Sowb	75.
Solarium Lmk.	62.
Solemya Lmk.	123.
Solen L.	112.
Spondylus L.	135.
Stomatella Lmk.	85.
Stomatia Helb.	85.
Strombus L.	1.
Stylina Flem.	66.
Tellina L.	99.
Terebra Adanson.	33.
Teredo L.	114.
Thracia Leach.	108.
Trapezium M. de M.	123.
Triforis Desh.	51.
Tritonium Cav.	22.
Trochita Risso	72.
Trochus L.	83.
Turbinella Lmk.	16.
Turbo L.	79.
Turbonilla Risso	72.
Turritella Lmk.	46.
Venus L.	95.
Vermetus Adanson	74.
Vitrinella Adams	80.
Volvaria Lmk	46.
Voluta L.	36.
Xenophora F. v. W.	68.

Strombaceæ.

Strombus Linné.

- S. accipitrinus* Mart. (syn. *S. costatus* Gml., *S. inermis* Sowb. Queen shell, spacy quen-shell). Station: 5 to 7 fathoms water on bluish sand in Spratbay St. Thomas; eats sea-weed; lives in floks. Habitat: — ? Lamarck! West-Indies; Reeve! Antillae; Mörch*)! Haity; D'Orb! Tortola; St. Thomas; St. Croix; Krebs! Guadeloupe; Beau**)! on the authority of C. B. Adams: St. Thomas; J. H. Newton! St. Johns; E. Hartwig! Florida; C. B. Adams! Bermuda; J. Redfield!
- S. alatus* Gml. (syn. *S. pyrolatus* Lamarck, *S. pugilis* Delw.) Station: — ? — ; habitat: — ? Lamarck! Gulf of Mexico; Reeve! West-Indies & Florida; C. B. Adams! Guadeloupe; Beau! Remark. only a variety of *S. pugilis* Linné!
- S. costoso-muricatus* Mart. (syn. *S. raninus* Gml., *S. affinis* Gml., *S. curruca* Bolt., *S. lobatus* Sw., *S. bitubercularis* Lam.). Station: 5 to 7 fathoms water on bluish sand; eats sea-weed; lives in floks together with *S. accipitrinus*. Habitat: l'océan des Antilles; Lamarck! Antillae; Mörch! Island of Grenada; Reeve! Guadeloupe; Beau! Anguilla; St. Barth; St. Martin; St. Thomas; Trinidad; Barbados; Tortola; Vera-Cruz & Carthagera

*) O. A. L. Mörch: catalogus cœnonchyliorum quæ reliquit comes de Yoldi. Hafniæ 1852.

**) M. Beau: catalogue des coquilles recueillies à la Guadeloupe & ses dépendances. Paris 1858.

in N. Gr.; Krebs! and according to C. B. Adams: Tortola; E. Hartwig! Jamaica; C. B. Ad! Carthagera in N. Gr. & Bermuda; J. Redfield!

- S. dubius* Sow. station: — ? ; habitat: Jamaica & Florida; C. B. Adams! Jamaica; John C. Jay's Cat. 4te Ed. Remark: not in the collections at St. Thomas.
- S. gallus* L. station: on gravel in 1 to 2 fathoms water. Habitat: Les mers d'Asie & d'Amerique; Lamarck! Antillae; Mörch! Cuba; Martinique; St. Lucy; d'Orb.! the island of St. Johns & at Longpoint on Tortola; Krebs! Guadeloupe; Beau!
- S. gigas* L. King-shell.-station: in 3 to 4 fathoms water on bluish sand; eats species of *sphaerococcus* (*Sphaerococcus confervoides* Ag.) habitat: Cuba; d'Orb.! Guadeloupe; Beau! Jamaica; C. B. Adams! Bermuda; J. Redfield! St. Thomas; St. Johns; St. Croix; Tortola; St. Barth and Carthagera N. Gr.; Krebs!
- S. Goliath* Chem. station: — ? ; habitat: Guadeloupe; Beau! Tortola; Krebs! remark: two specimens from Tortola, but by us until further considered a mere monstrosity of *S. gigas* Linné, though the two specimens have several characters alike, by which they differ from the *S. gigas*.
- S. granulatus* Sow. Antillae; Mörch!
- S. lentiginosus* L. Cuba; d'Orb.!
- S. pugilis* L. station: two fathoms water on mud-habitat: Gulf of Mexico; Reeve! Cuba; Martinique; Guadeloupe; St. Lucy; Carthagera. N. G.; New-Orleans; d'Orb.! Guadeloupe; Beau! Florida & Jamaica; C. B. Ad! St. Thomas; St. Croix; Tortola; Carthagera in N. G.; Vera-Cruz; Krebs!

Concae.

Conus Linné.

- C. armillatus* Ad. station: — ? ; habitat: St. Thomas (from Krebs) and Jamaica; C. B. Ad! Remark: we have not seen this species, but we suggest it to be *Conus Mindanus* Hwass, which C. B. Ad. has described as a new species.
- C. aurantius* Hwass. station: — ? ; habitat: Antillae; Mörch!
- C. barbadensis* Hwass. station: — ? ; habitat: West-Indies; Brugiére! Guadeloupe; Duchassain! Hamsbluff at St. Croix and Just van Dyck's island. near St. Thomas; Krebs! Remark: We do not consider this species to be anything else, than the full grown *Conus leucostictus* Hwass, which the specimens in our collection fully prove.
- C. baeticus* Reeve? station: — ? ; habitat: Antillae; Mörch!
- C. castus* Reeve. station: — ? ; habitat: St. Thomas; Krebs! Remark: our specimen answers perfectly to the drawing and the description in Reeve's icon. chon., but we do not consider it to be anything else than a young *Conus daucus* Hwass, which species does not get the impressed lines before they approach the full size.
- C. cedo-nulli* L. station: in deep water according to Reeve; habitat: West-Indies; Reeve! St. Thomas; St. Croix; Tortola; Krebs!
- C. centurio* Born. station: — ? ; habitat: les mers des Antilles; Lamarek! Island of Margarita; R. Swift! Hamsbluff at St. Croix; Krebs!
- C. characteristicus* Chem. station: — ? ; habitat:

ocean americaine; Lamarek! West-Indies; Reeve! Remark: in none of the collections at St. Thomas.

C. columba Hwass. station: probably in very deep water as Mr. Beau, who is an excellent collector, says: „I never found it without being more or less broken by rolling“; — habitat: common at the island of Marie galante near Guadeloupe; Beau! West-Indies; Reeve! Jamaica; C. B. Ad!

C. cretaceus Knr. station: — ? , habitat: West-Indies; S. Hovey! Jamaica; C. B. Ad!

C. daucus L. station: in very deep water; habitat: Martinique & St. Lucy; de Candé! Hotessier! Guadeloupe; Beau! de Candé! Hotessier! Cuba; de la Sagra! Auber! St. Martin; Tortola; Just van Duck's island near St. Thomas; Krebs! St. Thomas; R. Swift! Jamaica; C. B. Ad!

C. daucus v. *luteus* together with the orange-coloured species.

C. echinulatus Kien. station: — ?; habitat: Guadeloupe; Beau!

C. fumigatus Hw. station: — ?; habitat: seas of America (?); Reeve! Lmk.!

C. granulatus L. station: — ?; habitat: Guadeloupe; Beau! Ind. orient; Mörch!

C. leoninus Hw. station: see *conus spurius*; habitat: Antillae; Mörch! Coast of Mexico & West-Indies; Reeve! Guadeloupe; Beau! Remark: it does not deserve to stand at a species, scarcely as a variety of *Conus spurius* Gml.; *C. leoninus* stands as one of the strongest proofs for the eagerness of collectors to increase the numbers in their catalogues; the difference only consisting in the colour.

- C. leucostictus* Hw. (*C. nebulosus* Sol.) station: — ?; habitat: Antillae; Mörch! West-Indies; Reeve! Cuba; de la Sagra! Martinique & St. Lucy; de Candé! St. Johns; E. Hartwig! Jamaica; C. B. Ad! Guadeloupe; Beau! St. Thomas; St. Croix; St. Martin; Trinidad; Krebs!
- C. maculiferus* Brod. station: — ?; habitat: Guadeloupe; Beau!
- C. mercator* L. station: on sand; Reeve! habitat: Nevis; by Cap. Power according to Reeve! Antillae; Mörch! one very large, but poor, specimen found at Smithsbay, St. Thomas; Krebs!
- C. minutus* Reeve. station: — ?; habitat: Island of St. Vincent; Guilding!
- C. mindanus* Hw. station: at certain seasons, between loose stones close to the shore on 1 to 2 feet water; habitat: Philippines; Lamarek! Reeve! Antillae; Mörch! Curacao; R. Swift! St. Thomas; St. Barth; Anguilla; St. Martin; Krebs! remark: the old, full grown specimens are very different from the young, which have on granulation at all; see *Conus pusio*.
- C. mus* Hw. (*C. Barbadosensis* Hw.). station: young shells found in two to three feet water, between stones overgrown with seaweed. habitat: l'océan des Antilles sur les cotes de la Guadeloupe; Lamarek! Antillae; Mörch! Vera-Cruz; Cuba; St. Lucy; Martinique & Guadeloupe; d'Orb! Jamaica; C. B. Ad! St. Johns; E. Hartwig! St. Thomas; St. Croix; St. Johns; St. Barth; St. Martin; and at St. Martha & Carthagen in Nw. Gr.; Krebs! Arecibo Puer-toricco; J. H. Newton!
- C. musicus* Hw. station: — ?; habitat: West-Indies; by Capt. Power according to Reeve! West-Indies; by S. Hovey according to C. B. Adams!

- C. nodiferus* Kien. station: — ? — ; habitat: Guadeloupe; Beau!
- C. puertoricanus* Hw. station: — ? — ; habitat: Puertorico; Lamarek! Reeve!
- C. punctatus* Hw. (*C. piperatus* Delw., *C. biliosus* Bolt.) station: — ? — ; habitat: Carthagen. New Grenada; according to C. B. Ad! St. Martha in N. Gr.; Krebs!
- C. purpurascens* Brod. station: — ? — ; habitat: Guadeloupe; Beau!
- C. pusio* Brug. station: — ? — ; habitat: Antilles; Lamarek! remark: we consider the name of *c. pusio*-syn. with *conus Mindanus*; the *pusio* being described after young specimens, which do not possess the characters of the full grown shell.
- C. spurius* Gml. (*C. proteus* Hw. and according to Beau *C. pseudo Thomae* Chem.) station: on bluish mud, together with *Strombus accipitrinus* Mart. in 5 to 7 fathoms water; habitat: l'océan d'Atlantique et celui d'Amerique; Lamarek! Antillae; Mörch! St. Thomas; Reeve! d'Orb.! St. Johns; E. Hartwig! Guadeloupe; Beau! St. Thomas; St. Croix; St. Johns; Tortola; Krebs!
- C. pygmaeus* Reeve. station: — ? — ; habitat: ? Reeve! Antillae; Mörch! St. Thomas; Tortola; Krebs!
- C. radiatus* Gml. (*C. Martinianus* R., *C. gubba*. Ki.?). station: — ? — ; Antillae? Mörch!
- C. roseus* Lamarek. station: — ? — ; habitat: Les mers des Antilles; Lamarek! West-Indies; Reeve! St. Thomas; Krebs! remark: we consider this a variety of *Conus mus* Hw.; we possess a specimen which is rose-colored, but they do else correspond in all other respects with *C. mus*.

- C. testudinarius* Martini. station: — ? — ; habitat West-Indies; Lamarck! Reeve! Guadeloupe; Beau!
- C. varius* L. (*C. granulosus*. Bolt.), station: supposed to be in very deep water. Habitat: West-Indies; Surinam; Brasils; Lamarck! Antillae; Mörch! St. Croix; R. Swift! Antigua; Reeve! St. Thomas; Ziock! Krebs! Jamaica; C. B. Adams!
- C. verrucosus* Hw. (*C. granulatus*. Mart. non L.) station: — ? — ; habitat: Antillae; Mörch!
- C. Villepinii* Fish. & Bern. station: — ? — , habitat: Marie Galante; Beau!

Pleurotomaceæ.

Pleurotoma Lamarck.

- P. affinis* Grays Man. station: — ? , habitat: St. Vincent; Guilding! St. Thomas; Knox! Guadeloupe; Beau!
- P. albella* Ad. station: — ? — ; habitat: Jamaica; C. B. Adams!
- P. albida* Ad. station: — ? ; habitat: Jamaica; C. B. Adams!
- P. alabaster* Reeve. station: — ? — ; habitat: Carthagera in Nw. Gr.; John Redfield according to C. C. Adams!
- P. albocincta* Ad. station: — ? — ; habitat; St. Thomas; St. Martin & Waterisland; Krebs! Jamaica; C. B. Adams! Guadeloupe; Petit!
- P. albomaculata* Ad. station: — ? ; habitat: Jamaica; C. B. Ad.!
- P. albomaculata* d'Orb. Moll. de Cuba, pt. 2 pg. 176 pl. 24 fig. 15—18. station: — ? — ; habitat: Cuba; d'Orb.! Guadeloupe; Beau!

- P. angulifera* Reeve. station: very deep water, d'Orb.!
habitat: West-Indies; d'Orb.!
- P. Antillarum* d'Orb. Moll. de Cuba, pt. 2 173 pl. 24
fig. 1—3. station: — ? ; habitat: Cuba; Martinique;
d'Orb.! Guadeloupe; Beau!
- P. augustae* Ad. station: — ? — ; habitat: Jamaica;
C. B. Ad.!
- P. Auberiana* d'Orb. Moll. de Cuba, pt. 2 pg. 174 pl.
24 fig. 4, 6. station: — ? — ; habitat: Martinique;
Cuba; d'Orb.! Guadeloupe; Beau!
- P. balteata* Reeve. station: ? ; habitat: St. Thomas;
R. Swift!
- P. cancellata* Gray. station: — ? — ; habitat: St.
Vincent; Guilding! St. Thomas; Krebs!
- P. Candearia* d'Orb. Moll. de Cuba, pt. 2 pg. 175 pl.
24 fig. 10—12. station: — ? — ; habitat: Mar-
tinique; Guadeloupe; d'Orb.! Guadeloupe; Beau!
- P. candidula* Reeve. station: very deep water, d'Orb.!
habitat: West-Indies; d'Orb.!
- P. caribaea* d'Orb. Moll. de Cuba, pt. 2 pg. 173 pl. 23
fig. 32—34; station: — ? — ; habitat: Cuba; Gua-
deloupe; Martinique; d'Orb.! Guadeloupe; Beau!
- P. clathratus* Reeve. station: very deep water, d'Orb.;
habitat: West-Indies; d'Orb.!
- P. coccinata* Reeve. station: — ? — ; Saba; R. Swift!
St. Thomas; St. Johns; Crabisland; St. Barth; St. Mar-
tin; Anguilla; Krebs! St. Thomas; C. B. Ad.!
- P. collaris* Reeve. station: — ? — ; habitat: Guade-
loupe; Beau!
- P. costata* Gray. station: ? ; habitat: Guadeloupe
& Martinique; d'Orb.! Guadeloupe; Beau! St. Vincent,
Reeve! St. Thomas; St. Johns; St. Barth; St. Martin;

Anguilla; Krebs! remark: it is surely synonyme with *Pl. trifasciata*. Gray; we have specimens without lines and specimens with 1 and 2 and 3 lines on the upper whorls.

P. decorata Ad. is a *Daphnella*.

P. deminuta see *Mangelia quadrata* Reeve!

P. d'Orbignii Reeve. station: dredged from deep water d'Orbigny!; in bluish mud in 6 to 8 feet water R. Swift! Knox! habitat: St. Thomas; R. Swift! Knox! Lillien-skjold! Krebs!

P. Dorvillae Gray. station: ?; habitat: St. Vincent; Guilding! Guadeloupe; Beau!

P. elatior Ad. station: ? , habitat: Jamaica; C. B. Adams!

P. elatior d'Orb. Moll. de Cuba, pt. 2 pg. 173 pl. 23 fig. 35—37; station: ?; habitat: St. Thomas; de Candé! Guadeloupe; Beau!

P. flavo-cincta Ad. station: — ? — ; habitat: Jamaica; C. B. Ad.!

P. fucata Reeve. station: — ? — ; habitat: — ? Reeve! Antillae; Mörch! Guadeloupe; Duchassain! St. Thomas; St. Barth; Anguilla; St. Martin; Krebs!

P. fuscescens Gray. station: — ? — , habitat: Mauritius; Kien! Guadeloupe; Beau! St. Thomas; St. Croix; Krebs! Antillae; Mörch! remark: should this not prove to be synonyme with *Pleurotoma nigrescens* Gray?

P. fusco-cincta Ad. station: — ? — ; habitat: Jamaica; C. B. Ad.!

P. fusiformis Ad. station: — ? — ; habitat: Jamaica; C. B. Ad.!

P. Guildingii Reeve. station: — ? — ; habitat: St.

Vincent; Guilding! St. Thomas (?); Krebs! Guadeloupe; Beau!

- P. harpula* Val. station: — ? — ; habitat: New-Holland; Kien! Antillae; Mörch! St. Thomas; R. Swift! John Knox! St. Croir; Krebs! remark: stands very near *Pl. fuscescens* and *Pl. nigrescens* Gray and is perhaps synonyme.
- P. hastula* Reeve. station: — ? — ; habitat: St. Thomas; Krebs!
- P. Jayana* Ad. station: — ? — ; habitat: Jamaica; C. B. Ad.! St. Thomas; Krebs! remark: a very doubtful species, likely synonyme with. *Pl. zebra* Kiener!
- P. Lavelleana* d'Orb., pt. 2 pg. 174 pl. 24 fig. 7, 9. station: — ? — ; habitat: Cuba; Jamaica; Guadeloupe; Martinique; d'Orb.! Guadeloupe; Beau!
- P. lagueata* Reeve. station: in very deep water d'Orb.! habitat: West-Indies; d'Orb.!
- P. luctuosa* d'Orb. station: — ? — ; habitat: Cuba; d'Orb.! pt. 2 pg. 172 pl. 23, fig. 29—31. Guadeloupe; Beau!
- P. maculata* Ad. station: — ? ; habitat: St. Thomas; and Jamaica; C. B. Ad.!
- P. minor* Ad. station: — ? — ; habitat: Jamaica; C. B. Ad.!
- P. monilifera* Sowb. station: — ? — ; habitat: Jamaica; C. B. Ad.!
- P. nigrescens* Ad. station: — ? — ; habitat: Jamaica; C. B. Ad.! remark: is this and the *P. semigranosa* Reeve not synonyme?
- P. nigrescens* Gray. station: under flat pieces of madrepores in 1 to 2 feet water; habitat: St. Vincent; Guilding! Guadeloupe: Beau! St. Thomas; St. Croix;

Anguilla; & St. Martin; Krebs! remark: see *P. fuscescens* Gray & *P. harpula* Val. & *P. semigranosa* Reeve.

P. nodata Ad. station: — ? — ; habitat: Jamaica; C. B. Ad.!

P. obesicostata Reeve. station: very deep water d'Orb.! habitat: West-Indies; d'Orb.!

P. ornata d'Orb. pt. 2 pg. 171 pl. 23 fig. 26, 28; habitat: Cuba; d'Orb.!

P. pagodus Reeve. station: — ? — ; habitat: — ? Reeve! St. Thomas?; Krebs!

P. paria Reeve. station: — ? — ; habitat: ? Reeve! St. Thomas; John Knox!

P. pellis-phocæ Reeve. station: — ? — ; habitat: — ? Reeve! St. Thomas; R. Swift!

P. paxillus Reeve. station: — ? — ; habitat: St. Jan; St. Thomas; Krebs! remark: perhaps synonyme with *P. fuscescens* Gray, *P. harpula* Val. and *P. nigrescens* Gray.

P. pentagonalis Gray. station: — ? — ; habitat: St. Vincent; Guilding! Guadeloupe; Beau!

P. pulchella Reeve. station: — ? — ; habitat: — ? Reeve! Antillæ; Mörch!

P. pulchra Gray. station: — ? — ; habitat: St. Vincent; Guilding! St. Thomas; Knox! Krebs!

P. pygmaea Ad. station: — ? — ; habitat: Jamaica; C. B. Ad.!

P. pura Reeve. station: — ? — ; habitat: St. Martin; Krebs!

P. quadrifasciata Reeve. station: — ? — ; habitat: Jamaica; C. B. Ad.!

P. quadrilineata Ad. station: — ? — ; habitat: Jamaica; C. B. Ad.! remark: we have not seen authentic

- specimens of this species, but we are inclined to think it synonyme with *P. costata* Gray and *P. trifasciata* Gray.
- P. rubicata* Reeve. station: very deep water; habitat: West-Indies; d'Orb.!
- P. semigranosa* Reeve. station: — ? — ; habitat: — ? — ; Reeve! St. Thomas; Krebs! remark: see *P. nigrescens*. Ad.!
- P. solida* Ad. station: — ? — ; habitat: Jamaica; C. B. Ad.! West-Indies; S. Hovey!
- P. trifasciata* Gray. (see *P. costata* Gray.)
- P. turbinelloides* Reeve. station: very deep water; habitat: West-Indies; d'Orb.!
- P. urnula* Reeve (?). station: — ? — ; habitat: St. Thomas; R. Swift!
- P. Vespuciana* d'Orb. pt. 2 pg. 175 pl. 24 fig. 13, 15. station: — ? — ; habitat: Guadeloupe; Martinique; Cuba; d'Orb.! Guadeloupe; Beau!
- P. Virgo* Lmk. station: — ? — ; habitat: Cartagena. Nw-Gr., John Redfield, accordding to C. B. Ad.!
- P. zebra* Lmk. station: — ? — ; habitat: Mauritius; Kiener! Antillae; Mörch! St. Thomas; Knox; St. Thomas; St. Johns; Water-Island; Krebs!

Daphnella Hinds.

- D. lymnaeiformis* Kiener. station: — ? — ; habitat: Philippine Islands; Cuming! Antillae; Mörch! St. Thomas; Bland! Guadeloupe; Duchassain! Beau! Petit! St. Thomas; St. Johns; Jamaica; Ad.! Just van Dyck's Island and Water-Island near St. Thomas; St. Thomas; St. Johns; St. Barth; St. Martin; Anguilla; Krebs! remark: the localities of C. B. Adams refer to *Daphnella decorata*, described by him; it is certainly synonyme

with *D. lymnaeiformis* K. from which C. B. Adams, once in a conversation, acknowledged not to be able to distinguish it without by the locality; we do not acknowledge this to be a sufficient argument to establish species on and we really regret that such doctrines have been able to enter, to multiply and to confuse the nomenclature and the science.

Magnelia Risso.

- M. antillarum* Reeve. station: — ? — ; habitat: West-Indies; Reeve!
- M. angulata* Reeve. station: — ? — ; habitat: — ? — ; Reeve! remark: we have found a species at St. Thomas, which approaches the drawing in Reeve's *Ic. conch.* very much.
- M. badia* Reeve station: dead specimens between stones and species of corals in 2 feet water; habitat: — ? Reeve! St. Thomas; Krebs! remark: see *M. crassicosata* Ad.!
- M. balteata* Reeve. station: in mud from 6 to 8 feet water; habitat: — ? — ; Reeve! St. Thomas; St. Barth; St. Martin; Anguilla; Krebs!
- M. biconica* Ad. station: — ? — ; habitat: Jamaica; C. B. Adams!
- M. biconica* var. *alba* Ad. station: — ? — ; habitat: Jamaica; C. B. Adams!
- M. brevis* Ad. station: — ? — ; habitat: Jamaica; C. B. Adams!
- M. candidissima* Ad. station: — ? — ; habitat: Jamaica; C. B. Adams! St. Thomas; Bland!
- M. crassicosata* Ad. station: — ? — ; habitat:

- Jamaica; C. B. Ad.! remark: we consider this and *M. badia* Reeve as synonymes.
- M. densestriata* Ad. station: — ? — ; habitat: Jamaica; C. B. Adams! remark: we have until further laid this species together with *M. balteata* Reeve in our collection.
- M. Hornbecki* Reeve. station: — ? — ; habitat: St. Thomas; Hornbeck! Krebs! remark: Reeve's drawing of this shell is very bad.
- M. lanceolata* Ad. station: — ? — ; habitat: Jamaica; C. B. Adams!
- M. luteofasciata* Reeve. station: — ? — ; habitat: — ? — ; Reeve! St. Thomas; Knox!
- M. multilineata* Ad. station: — ? — ; habitat: St. Thomas (from Bland); Jamaica; C. B. Ad!
- M. muricoides* Ad. station: — ? — ; habitat: Jamaica; C. B. Ad.!
- M. occidentalis* Reeve. station: dredged from very deep water; habitat: West-Indies; Reeve! St. Thomas; Krebs! remark: *Pleurotoma occ.* Reeve.
- M. quadrata* Reeve. station: — ? — ; habitat: — ? — ; Reeve! St. Thomas; R. Swift! Krebs! St. Martin; Krebs! Jamaica; C. B. Adams! remark: it is a *Pleurotoma* of Reeve and we take it to be synonyme, with C. B. Adams' *Pleurotoma deminuta*.
- M. astricta* Reeve. station and habitat: — ? — ; remark: we have found a species at St. Thomas, which approaches the drawing and the description of *M. ast.* in Reeve's *lc. conch.* as much as many shells described and drawn in said work, answer to the descriptions and the drawings intended to represent them.

M. tineta Reeve. station: — ? — ; habitat: St. Thomas; Lillienskjold! remark: *Pleurotoma* Reeve.

M. triticea Reeve (*Pleurotoma*). station: — ? — ; habitat: St. Thomas; Knox! Anguilla; received twice from there; Krebs!

Conopleura Hinds.

C. cancellata d'Orb. (*Sinusigera*). station: — ? — ; habitat: Jamaica; de Candé!

Fusus Bruguière.

F. distans Lmk. station: — ? — ; habitat: — ? — ; Mörch! Philippines; Reeve! Cartagena, Nw. Gr.; John Redfield! Bay of Campechy; C. B. Adams! South-Caroline; Say!

F. gradatus Reeve. station: dredged from 15 fathoms water; habitat: St. Thomas; Krebs!

F. lineatus Chem. (*F. ansatus* Gml.). station: — ? — ; habitat: Antillae; Mörch!

F. muricoides Ad. station: — ? — ; habitat: Jamaica; C. B. Adams!

F. nitens Ad. station: — ? — ; habitat: Jamaica; C. B. Adams!

F. reuma Mart. station: — ? — ; habitat: Antillae Mörch!

Fasciolaria Lamarck.

F. gigantea d'Orb. station: — ? — ; habitat: Cuba; d'Orb.! Moll. de Cuba, pt. 2 pg. 169 pl. 23 fig. 25.

F. Tulipa Lmk. station: on 2 a 6 feet water on sand and small stones, where Algae are growing and where the sea is calm; it is in very large quantity in the creeks around the town of Carthagena in N. Gr.; habitat: Antillae; Mörch! Guadeloupe; Beau; Cuba; Guade-

loupe; St. Thomas; Columbia; d'Orb.! Honduras; Dyson!
 St. Thomas; St. Johns; Tortola; Trinidad; St. Martin;
 Carthagen, Nw. Gr.; Krebs! Florida; South-Caroline;
 Jamaica; Chagres; C. B. Ad.!

Turbinella Lamarck.

- T. ananas* Chem. (*M. infundibulum*. Gml., *B. sulcatum* Msch.) station: in deep water, but at certain seasons between stones close to the shore; habitat: West-Indies; Reeve! Antillae; Mörch! Martinique; Guadeloupe; St. Lucy; d'Orb.! Guadeloupe; Beau! Jamaica; C. B. Adams! St. Martin; St. Croix; St. Thomas; Krebs!
- T. annulata* Bolt. station: — ? — ; habitat: Antillae; Mörch! St. Johns and Just van Dyck's Island near St. Thomas; Krebs!
- T. attenuata* Reeve. station: — ? — ; habitat: — ? — ; Reeve! Antillae; (?) Mörch!
- T. brevicaudata* Reeve. station: — ? — ; habitat: — ? — ; Reeve! Tortola; St. Thomas; St. Johns; C. B. Adams! remark: should this not be synonyme with *P. filamentosa* Kock?
- T. capitellum* L. station: — ? — ; habitat: Antillae; Mörch! Martinique; Guadeloupe; St. Lucy; d'Orb.! Guadeloupe; Beau! St. Thomas; R. Swift! Krebs!
- T. carinifera* Lmk. station: — ? — ; habitat: St. Thomas; Krebs!
- T. cingulifera* Lmk. station: — ? — ; habitat: Guadeloupe; Beau!
- T. filimentosa* Koch. station: — ? — ; habitat: Antillae; Mörch! St. Thomas; Tortola; Krebs!
- T. fuscata* Gml. (*Turbinella angularis* Reeve). station: — ? — ; habitat: Antillae; Mörch!

- T. gibbula* Gml. (*T. filusus* Lmk.) station: — ? — ; habitat: Antillae; Mörch!
- T. globulus* Lmk. station: — ? — ; habitat: Antigua?; Krebs! remark: Does the figure in Reeve's Ic. conch. really represent this species?
- T. gracilis* Reeve. station: — ? — ; habitat: St. Thomas; Krebs!
- T. leucozonalis* Lamk. station: — ? — ; habitat: St. Thomas; St. Croix; Tortola; Krebs! Honduras; Reeve! Jamaica; Carthagenae; Nw. Gr. (acc. to Redfield!) C. B. Adams!
- T. lineata* Lmk. station: — ? — ; habitat: St. Thomas; Krebs!
- T. muricata* Born (*T. pugillaris* Lmk.) station: — ? — ; habitat: Antillae; Mörch! Tortola; Krebs! Martinique; d'Orb.! Guadeloupe; Beau! St. Johns; Jamaica; C. B. Adams!
- T. nana* Reeve. station: — ? — ; habitat: St. Thomas & St. Croix; Krebs!
- T. nassa* Gml. station: — ? — ; habitat: Cuba; St. Lucy; Martinique; Guadeloupe; St. Thomas; d'Orb.! Jamaica; St. Thomas; C. B. Adams!
- T. nigella* Chem. (*P. ocellata* Gml.) station: — ? — ; habitat: Antillae; Mörch! Guadeloupe; Beau! St. Thomas; Anguilla; St. Martin; Krebs!
- T. polygona* Lmk. station: — ? — ; habitat: Cuba; Bahia; d'Orb.!
- T. recurvirostra* Wagn. station: — ? — ; habitat: Philippines; Reeve! Antillae; Mörch!
- T. rhinoceros* Lmk. station: — ? — ; habitat: Zanzibar; Reeve! Carthagenae; Nw. Gr. (according to J. Redfield); C. B. Adams!

- T. rudis* Reeve. station: — ? — ; habitat: St. Thomas; St. Johns and Guadeloupe; Krebs!

Pyrula Lamarck.

- P. aruana* L. (*P. carica* Gml.) station: — ? — ; habitat: Antillae; Mörch!
- P. melongena* L. station: — ? — ; habitat: Antillae; Mörch! Cuba; Martinique; Guadeloupe; Carthagenae N. Gr.; Florida; d'Orb.! Guadeloupe; Beau! Carthagenae New-Gr.; Redfield! Jamaica; C. B. Adams! St. Juan de Nicaragua; Tortola; Krebs!
- P. morio* L. (*Fusus coronatus* Lmk.) station: on mud; habitat: Brasils; Martinique; Cuba; d'Orb.! Trinidad; Reeve! Guadeloupe; Beau! Duchassin!
- P. perversa* Lmk. station: — ? — ; habitat: Cuba; Mississippi; d'Orb.!

Murex Linné.

- M. alveatus* Kien. station: on 1 or 2 feet water, between pieces of Madrepores; habitat: Antillae; Mörch! Jamaica; C. B. Adams! Guadeloupe; Beau! St. Thomas; St. Croix; St. Martin; Krebs!
- M. Beauii* Fisch. & Bern. station: — ? — ; habitat: Marie Galante; Beau!
- M. bellus* Reeve. station: a dead specimen dredged from 10 fathoms water on bluish mud; habitat: Antillae?; Mörch! Tortola; A. H. Riise! St. Thomas; Krebs!
- M. bilineatus* Reeve. station: — ? — ; habitat: Goyave; Beau!
- M. brevifrons* Lmk. (*M. calcitrapa* Lmk.) station: — ? — ; habitat: Antillae; Mörch! Martinique; Guadeloupe; Cuba; St. Thomas; Brasils; d'Orb.! Jamaica; Carthagenae

Nw. Gr.; C. B. Adams! remark: this will surely later prove identical with *Murex cornu cervi* Martini.

M. Cailleti Petit. station: — ? — ; habitat: Guadeloupe; Beau!

M. cancellatus Sowb. station: — ? — ; habitat: St. Johns; Knox!

M. chrysostoma Gray. station: — ? — ; habitat: Island of Margarita; Mörch! R. Swift!

M. cornu-cervi Mart. (*M. breviformis* Lmk., *M. calcitrapa* Lmk., *M. purpuratus* Reeve) station: on stinking bluish mud 10 to 12 feet below the surface of the water in a creek with smooth water depositing eggs; habitat: Antillae; Mörch! Guadeloupe; Beau! St. Thomas; Krebs! remark: we possess specimens, which have been taken out of the eggs and every step of development until the size of *Murex purpuratus* Reeve, drawn in his *lc. conch.* and they fully prove the synonymes mentioned above, to be correct. It is proper to remark: that some specimens have one and others have two nodules between each varix, although taken out of one cluster of eggs, but all the old and full-grown specimens have only one nodule, some with a very faint indication of the second, between each varix, young specimens have no sculpture before on the fourth or fifth whorl.

M. crassivaricosa Reeve. station: — ? — ; habitat: Guadeloupe; Beau! St. Thomas; Krebs! remark: we have laid four specimens, with this name, in our collection, but we are inclined to think that the *Murex crassivaricosa* Reeve is only a *Murex cornu-cervi* Mart.

M. elegans Bech. station: — ? — ; habitat: St. Domingo; St. Martin; d'Orb.! Port-Louise in the island of Fajau near Guadeloupe; Beau! remark: Beau says:

„it is only found at certain seasons“. It is a very remarkable fact how common certain species of marine-shells appear at certain localities for a length of time and thereafter disappear for many years; perhaps some species, which are living in the deepest parts of the sea, come, at certain seasons, to the shores to breed; Rvd. Ziöck has at least once found 6 or 8 full-grown life specimens of *Turbinella ananas* Chem. on one to two feet water, close to the shore; Reeve says that the *Turbinella ananas* Chem. lives in deep water and we have never found it close to the shore, though we in 17 years have made excursions around the W. Ind - Islands; we have likewise once found several specimens of *Murex hexagonus* Lmk. in one place close to the shore and since then not found it for many years.

M. elongatus Lmk. station: — ? — ; habitat: Gulf of Mexico; Reeve! remark: Perhaps a variety of *Murex cornu-cervi* Mart; Reeve makes the same mistake by the drawing of this species as by Nr. 5, 13, 14, 20, 23, 24, 25 and many more in the Ic. conch. in representing young specimens, where the characteristic enameled lip is not yet enveloped.

M. funiculatus Reeve. station: on stones in one or two feet water in the creeks around Carthagera; habitat: Jamaica; C. B. Adams! Carthagera in New-Granada; Krebs!

M. hexagonus Lmk. station: young specimens found at St. Thomas between pieces of corals and stones on 1 to 2 feet water; habitat: Antillae; Mörch! Guadeloupe; Beau! St. Thomas; St. Johns; Jamaica; C. B. Adams! St. Thomas; Tortola; Krebs! remark: we found this shell quite commonly at the shores of St. Thomas in

1851—52; it has been rare before and since that period.

M. intermedius Ad. (*M. alveatus*. Kiener).

M. megacerus Sowb. station: — ? — ; habitat: West-Indies; Reeve! Guadeloupe; Beau!

M. Messorius Sowb. station: on stones in 1 or 2 feet water; habitat: — ? Reeve! Jamaica; C. B. Adams! Martinique; de Candé! Guadeloupe; Beau!

M. motacilla Chem. station: — ? — ; habitat: Senegal; Reeve! Guadeloupe; Beau!

M. nodatus Reeve. station: — ? — ; habitat: Guadeloupe; Beau!

M. nuceus Mörch. station: at St. Thomas, under stones and pieces of Modrepores on one or two feet water; habitat: Antillae; Mörch! St. Croix; St. Thomas; St. Martin; Krebs!

M. oculatus Reeve. station: — ? — ; habitat: Jamaica; C. B. Adams! We do not know this species, but it seems to stand *Murex pomiformis* Mart. very near.

M. pleurotomoides Reeve. station: — ? — ; habitat: St. Martin; Krebs!

M. pauperculus Ad. (*Triton Cantrainei* Recl.) station: between rocks and pieces of corals on 1 to 2 feet water; habitat: Jamaica; C. B. Adams! Guadeloupe; Beau! St. Thomas; St. Johns; Krebs!

M. pomiformis Mart. station: on coral-reefs; Reeve! dredged from four fathoms water, on bluish sandy mud and on sand in 3 to 4 feet water between sea-weed, where it feeds on *Lucina pensylvanica*; habitat: Antillae; Mörch! Rio-Janeiro; Martinique; Guadeloupe; St. Thomas; d'Orb.! St. Vincent; Guldin! Jamaica; C. B. Adams! St. Thomas; Tortola; St. Martin; Caru-

- pano in Venezuela; remark: young specimens have two and three nodules between each varix, but the full-grown specimens have only one nodule between each varix.
- M. pudicus* Reeve. station: — ? — ; habitat: Island of Domingo; Reeve!
- M. pudoricolor* Reeve. station: St. Thomas on a few feet water between stones; Knox! habitat: St. Thomas; Robert Swift! Guadeloupe; Beau!
- M. purpuratus* Reeve. see *M. cornu-cervi* Mart.
- M. rarispina* Lmk. station: — ? — ; habitat: St. Domingo; Lmk! Jamaica; C. B. Adams!
- M. rubidentatus* Reeve. station: — ? — ; Guadeloupe; Duchassain! Beau!
- M. spectrum* Reeve. station: — ? — ; habitat: Guadeloupe; Beau!
- M. similis* Sowb. station: — ? — ; habitat: Guadeloupe; Beau! Duchassain!
- M. trigonulus* Lmk. station: — ? — ; Marie Galante; Terre-de-Bas des Saintes; Beau!
- M. tetragonus?* Brod. (*M. breviculus* Sow.) station: — ? — ; habitat: Guadeloupe; Beau!

Tritonium Cuvier.

- T. americanum* d'Orb. station: — ? — ; habitat: Cuba; Brasils; d'Orb.!
- T. antillarum* d'Orb. (*T. tuberosum* Reeve station: on rocks in one to two feet water; Krebs! habitat: Antillae; Mörch! Cuba; Martinique; Guadeloupe; d'Orb.!
- Guadeloupe; Beau! St. Johns; Jamaica; Bermuda; C. B. Adams! St. Thomas; Tortola; Krebs!
- T. aequale* Reeve. station: — ? — ; habitat: Antillae; Mörch!

- T. caribaeum* d'Orb. station: — ? — ; habitat: Pernambuco; Cuba; d'Orb.!
- T. chlorostomum* Lmk. station: — ? — ; habitat: Cuba; St. Lucy; Martinique; Guadeloupe; d'Orb.! Jamaica; St. Johns; Bermuda; C. B. Ad.! St. Thomas; Krebs!
- T. Cantrainii* Recluz see *Murex pauperculus* Ad.
- T. costatum* Born. (*T. succinctum* Lmk.) station: — ? — ; habitat: Guadeloupe; Beau! Tortola; Krebs!
- T. cynocephalum* Lmk. station: — ? — ; habitat: Guadeloupe; Beau! St. Johns; Jamaica; Bermuda; C. B. Adams! St. Croix; St. Martin; Trinidad; Krebs!
- T. femorale* L. station: on bluish, sandy mud together with *Strombus gallus* Linne! habitat: Cuba; St. Lucy; d'Orb.! Guadeloupe; Beau! St. Johns; Jamaica; C. B. Adams! St. Thomas; St. Croix; St. Martin; Tortola; Trinidad; Krebs!
- T. labiosum* Wood. (Tr. Loroisis Pt.) station: between stones and coral-pieces on 2 feet water at St. Thomas; Krebs! habitat: Antillae; Mörch! Guadeloupe; Duchassain! Beau! St. Thomas; St. Croix; St. Martin; Tortola; Krebs!
- T. lanceolatum* Mk. station: under stones and pieces of corals in one and two feet water; Krebs! habitat: Antillae; Mörch! Guadeloupe; Duchassain! Beau! St. Thomas; St. Croix; Crabisland; Bermuda; Trinidad; St. Martin; Anguilla; Krebs!
- T. nobile* Con. station: — ? — ; habitat: Jamaica; St. Johns; Cartagena! Nw.-Gr.; Bermuda; C. B. Ad.!
- T. obscurum* Reeve. station: — ? — ; habitat: Jamaica; C. B. Adams! Guadeloupe; Beau! St. Croix; Krebs!

- T. martinianum* d'Orb. station: — ? — ; habitat: Cuba; Guadeloupe; St. Lucy; St. Thomas; d'Orb.! see *T. piliare* Linné & Lmk.
- T. parvum* Ad. station: Between stones and pieces of Madreporos on two feet water; Krebs! habitat: St. Thomas; Anguilla; St. Barth; Krebs! St. Thomas; Jamaica; Rum-key, Turks-Island; C. B. Adams!
- T. piliare* L. & Lmk. (*T. martinianum* d'Orb.) Antillae; Mörch! Guadeloupe; Beau! Jamaica; Carthagera in Nw. Gr.; St. Johns; Bermuda; Sandwich Isl. C. B. Adams! St. Thomas; Tortola; St. Martin; Barbadoes; Krebs!
- T. pulchellum* Ad. station: — ? ; habitat: Jamaica; C. B. Adams!
- T. reticulare* L. (*T. clathratum* Lmk.) Antillae; Mörch! Guadeloupe; Duchassain! Beau! Jamaica; Carthagera in Nw. Gren.; C. B. Adams! Tortola; Krebs!
- T. rostratum* Mart. (*Fusus cutaceus* L., *Cassidaria cingulata* Lmk., *Tr. undosum* Kiener.) station: — ? ; habitat: Antillae; Mörch!
- T. testaceum* Mörch. station: — ? — ; habitat: Antillae; Mörch! St. Thomas; St. Croix; St. Martin; Krebs! remark: we have cause to believe that the localities mentioned by C. B. Adams under the *T. obscurus* Reeve are relative to *Trit. testaceum*. Mörch.
- T. tigrinum* Brod. station: — ? — ; habitat: Guadeloupe; Beau!
- T. variegatum* Lmk. (*T. marmoratum* Lmk.) station: in two fathoms water on seaweed; habitat: Antillae; Mörch! Guadeloupe; Duchassain! Beau! Vera-Cruz; St. Thomas; Krebs!
- T. vestitum* Hinds. station: — ? ; habitat: Guadeloupe; Beau!

Ranella Lamarck.

- R. crassa** Desh. (*R. granulata* Lmk.) station: — ? ; habitat: Guadeloupe; Beau! Hotessier!
- R. cubaniana** d'Orb. station: — ? — ; habitat: Cuba; St. Lucy; d'Orb.! Guadeloupe; Beau!
- R. livida** Reeve station: — ? ; habitat: Jamaica; C. B. Adams!
- R. rhodostoma**: — ? ; station: — ? ; habitat: St. Croix according to a label in the Academy of natural sciences in Philadelphia; remark: the specimen in said Academy appears very like *R. Thomae* d'Orb.
- R. pondorosa** Reeve. station: — ? ; habitat: Guadeloupe; Duchassain! Jamaica; C. B. Adams! Hamsbluff at St. Croix; Krebs! remark: should Reeve's *R. livida* not be young specimens of this shell? ; the figure and the description in his *Ic. conch.* does not prove anything to the contrary.
- R. Thomae** d'Orb. station: — ? ; habitat: St. Thomas; d'Orb.!
- R. I sp. ind.** We have further two specimens of *Ranella* from St. Thomas, but we have not been able to find any information about them.

Purpura Bruguère.

- P. aberrans** Ad. station: — ? ; habitat: Jamaica; St. Thomas; C. B. Adams!
- P. bicostalis** Lmk. station: on stones above low-water mark; Krebs! habitat: Antillae; Mörch! Guadeloupe; Beau; St. Thomas; Krebs!
- P. fasciata** Reeve. station: on stones above low-water mark; Krebs! habitat: Guadeloupe; Petit! Beau! St.

- Johns; Jamaica; St. Thomas; C. B. Adams! St. Martha and Carthagera in Nw. Gr.; Krebs!
- P. floridana* Con. station: Florida Reeve! Guadeloupe; Petit! St. Thomas; Texas; Jamaica; C. B. Adams! Carthagera in Nw. Gr.; J. Redfield!
- P. galea* Chem. (*P. abbreviata* Bly.) station: on perpendicular rocks in 6 to 8 feet water: Krebs! habitat: Antillae; Mörch! Cuba; d'Orb.! Guadeloupe; Beau! St. Thomas; St. Johns; Jamaica; C. B. Adams! St. Thomas! St. Johns; Krebs!
- P. gibbosa* Reeve. station: — ? — ; habitat: Antillae; Mörch! St. Thomas; Krebs! Jamaica; C. B. Adams! remark: Mörch thinks it may be a *Rhizochilus* of Steenstrup.
- P. haemastoma* Lmk. station: — ? — ; habitat: Guadeloupe; d'Orb.! St. Thomas; Krebs! remark: it does not exactly live in the West-Indies, but is often found at the bottom of vessels which arrive from the coast of Africa at the port of St. Thomas.
- P. Kieneri* Desh. station: — ? ; habitat: Martinique Lamarek! Philippines; Reeve! remark: we think the last habitat to be correct as we never have seen a piece of one between those thousand of West-Indian shells which have passed through our hands.
- P. lineata* Lam. (*B. bulbus* Wood) station: — ? ; habitat: Antillae; Mörch!
- P. patula* Linné. station: on rocks at high water mark; Krebs! habitat: Antillae; Mörch! Cuba; Martinique; Guadeloupe; d'Orb.! St. Thomas; St. Martin; Carthagera and St. Martha in New. Gr.; Krebs!
- P. plicata* Mart. station: on and in the roots of certain species of *Gorgonia* in 2 to 3 fathoms water; Krebs!

habitat: Antillae; Mörch! St. Thomas; St. Johns; Anguilla; Krebs!

P. rustica Lam. station: — ? — ; habitat: Luzon (one of the Philippines) Reeve! Antillae; Mörch! St. Thomas; St. Martin; Krebs!

P. subdeltoidea Blainv. station: — ? — ; habitat: Cuba; Martinique; d'Orb.!

P. trapa Bolt. (*P. deltoidea* Lmk., *M. nodus* L., Meusch, *Hippocastanum fasciatum* Mart.) station: on rocks and stones at high-water mark; Krebs! habitat: Antillae; Mörch! Guadeloupe; Beau! Carthagera in Nw.-Gr.; Bermuda; Jamaica; C. B. Adams! St. Thomas; Krebs!

P. undata Lmk. station: — ? — ; habitat: Pernambuco; Cuba; d'Orb.! Guadeloupe; Beau!

P. vexillum Lmk. station: — ? — ; habitat: Antilles; d'Orb.!

Recinula Lamarck.

R. ferruginosa Reeve. station: — ? — ; habitat: Antillae; Mörch! remark: we think it is synonyme with *R. nodulosa* Ad.

R. nodulosa Ad. station: on stones in one to two feet water; Krebs! habitat: Guadeloupe; Beau! St. Thomas; St. Johns; St. Martha in New-Gr.; C. B. Adams! St. Thomas; St. Johns; St. Barth; St. Martin; Anguilla; Carthagera in New-Gr.; Krebs!

R. rosea Reeve. station: — ? — ; habitat: Island of Masbate; Reeve! St. Thomas; Krebs!

R. turbinella Kien. station: on stones in one to two feet water; Krebs! habitat: Guadeloupe; Beau! Duchasain! Petit! Jamaica; C. B. Adams! St. Thomas; St. Barth; Anguilla; St. Martin; Krebs!

Columbella Lamarck.

- C. argus* d'Orb. station: — ? — ; habitat: Guadeloupe; Hotessier! Beau!
- C. atomella* Ducl. station: — ? — ; habitat: West-Indies; Guilding! according to a note of C. B. Adams!
- C. barbadensis* d'Orb. station: — ? — ; habitat: Antilles; d'Orb.!
- C. Broderipii* Sow. station: — ? — ; habitat: Guadeloupe; Beau!
- C. catenata* Sowb. (*Buccinum mitrula* Dk.) station: under pieces of Madreporos on 1 à 2 feet water Krebs! habitat: Guadeloupe; Beau! Jamaica; C. B. Adams! St. Thomas; St. Johns; St. Martin; Krebs!
- C. corniculata* Ad. station: — ? ; habitat: Jamaica; C. B. Adams!
- C. decipiens* Ad. (*Bucc. concinnum* Ad.) station: — ? — ; habitat: Jamaica; C. B. Adams! St. Thomas; Krebs! remark: perhaps synonyme with *C. subcostata* Ad.
- C. dermestoides* Kiener. station: — ? — ; habitat: West-Indies; Sowerby! Guadeloupe; Beau!
- C. dichroa* Sowb. station: — ? — ; habitat: St. Vincent; Guilding! Jamaica; C. B. Adams!
- C. dormitor* Sowb. (*Col. purpurascens* Ad.) station: — ? — ; habitat: St. Vincent; Guilding! Jamaica; C. B. Ad.!
- C. Ducloziana* d'Orb. station: — ? — ; habitat: St. Thomas; Jamaica; d'Orb.!
- C. fastigata* Kien. station: — ? — ; habitat: les iles saintes Kien. according to C. B. Adams!
- C. fenestrata* Ad. station: — ? — ; habitat: Jamaica! C. B. Adams! St. Thomas; Krebs!
- C. fusiformis* d'Orb. station: — ? — ; habitat: Martinique; Jamaica; d'Orb.!

- C. Guildingii* Sowb. station: — ? — ; habitat: St. Vincent; Guilding!
- C. Hotessieri* d'Orb.! station: — ? — ; habitat: Guadeloupe; Hotessier! Beau!
- C. lactea* Sowb. station: — ? — ; habitat: Antillae; Mörch!
- C. laevigata* Lmk. (*Col. concinna* Sowb.) station: — ? — ; Antillae; Mörch! Jamaica; St. Johns; Arecibo; C. B. Adams! Guadeloupe; Beau! Duchassain! Carthagenae in New-Gr.; Redfield! St. Thomas; St. Croix; St. Martin; Trinidad; St. Martha in New-Gr.; Krebs!
- C. la Fresnayi*. Fisch & Bern. station: — ? — ; habitat: Marie Galante; Beau!
- C. mercatoria* L. station: between sea-weed on sandy ground on 2 to 4 feet water; habitat: Cuba; Martinique; Guadeloupe; St. Lucy; Jamaica; d'Orb.! St. Johns; Jamaica; Bermuda; Turksisland; Arecibo; C. B. Adams! St. Thomas; St. Croix; St. Barth; St. Martin; Anguilla; Barbadoes; Trinidad; Carthagenae in New-Gr.; Krebs!
- C. nitida* Lmk. (*S. nitidula* (L.?) Sowb.) station: under pieces of corals on 2 to 3 feet water; habitat: Guadeloupe; Martinique; St. Lucy; Cuba; d'Orb.! St. Johns! Arecibo; C. B. Adams! St. Thomas; St. Johns; St. Croix; St. Barth; Anguilla; St. Martin; Barbadoes; Krebs! remark: d'Orbigny makes the remark that this species diminishes in size from Guadeloupe toward Cuba; we have seen very large specimens said to come from Margarita.
- C. obesa* Ad. station: — ? — ; habitat: Jamaica; C. B. Adams! Guadeloupe; Beau! St. Thomas; Carthagenae; New-Gr.; Krebs!

- C. monolifera* Sowb. station: — ? — ; habitat: West-Indies; Jay's Catalogue!
- C. ocellata* d'Orb. (pt. 2 pg. 135) station: — ? — ; habitat: Cuba; St. Lucy; d'Orb.!
- C. ocellata* Gml. station: under stones on 1 to 2 feet water; Krebs! habitat: Jamaica; Bermuda; C. B. Adams! Guadeloupe; Beau! Duchassain! St. Thomas; St. Johns; St. Croix; St. Barth; Anguilla; St. Martin; Krebs! remark: in a manuscript of C. B. Adams stands *C. parvula* Dunker (Philippi?) and *C. guttata* Sowb. as synonymes of this species, but the firts mentioned species is surely destinct.
- C. ovulata* Lmk. station: under pieces of madrepores on stony ground in 1 to 2 feet water; Krebs! habitat: Cuba; Guadeloupe; St. Lucy; d'Orb.! Jamaica; C. B. Adams! St. Thomas; St. Croix; St. Barth; Anguilla; St. Martin; Krebs!
- C. ovuloides* Ad. station: — ? — ; habitat: Turks Island; C. B. Adams! remark: surely synonyme with *C. ovulata*. Lmk.
- G. Peleei* Kiener. (*C. rudis* Sowb.) station: — ? — ; habitat: Martinique; Kiener! Nevis; Sowerby!
- C. parvula* Ph. station: — ? — ; habitat: St. Thomas; St. Johns; St. Martin; Krebs!
- C. pulchella* Kiener (*Buc. triticeum* Sol.) habitat: Guadeloupe; Beau! St. Thomas; St. Johns; St. Croix; St. Martin; Anguilla; Krebs!
- C. pusilla* Sowb. station: — ? — ; habitat: St. Vincent; Guilding! Jamaica; C. B. Adams!
- C. rudis* Sowb. (*C. Peleei* Kien.).
- C. Sagra* d'Orb.! station: — ? — ; habitat: Guadeloupe; d'Orb.! Beau!

- C. *Schrammii* Petit. station: — ? — ; habitat: Guadeloupe; Beau!
- C. *subcostata* Ad. station: under pieces of madrepores on 1 foot water; Krebs! habitat: Guadeloupe; Beau! Duchassain! Petit! Jamaica; Turks-Island; Bahama; C. B. Adams! St. Thomas! St. Johns; St. Croix; St. Barth; St. Martin; Anguilla; Trinidad; Krebs!
- C. *therpsichore* Leathes. station: — ? — ; habitat: Antillae; Mörch! Nevis; Cuming! West-Indies; Lmk.! South sea; Kiener!

Pisania Bivona.

- P. *pennata* Brown. (B. *plumatum* Gm.; *Fusus articulatus* Lmk.; *Murex accinctus* Born; *Purpura accincta* d'Orb.) station: — ? — ; habitat: Cuba; St. Lucy; Guadeloupe; Jamaica; d'Orb.! Honduras; Reeve! St. Johns! E. Hartwig! Tortola?; Krebs! remark: it is rare, if at all found at the Virgin Islands.
- P. *tritones* Reeve station: — ? — ; habitat: Antillae; Mörch!

Nassa Lamarck.

- N. *albescens* Dunk. station: — ? — ; habitat: West-Indies; Reeve!
- N. *antillarum* d'Orb. station: — ? — ; habitat: Guadeloupe; Beau!
- N. (Phos) *antillarum* Phil. (B. *zonale* Brug) station: Antillae; Mörch! Cuba; Reeve! Guadeloupe; Beau! St. Thomas and Carthagera in New-Gr.; Krebs!
- N. *ambigua* Pult. station: — ? — ; habitat: Guadeloupe; Beau! Jamaica; Turks-Island; C. B. Adams! St. Thomas; St. Croix; St. Barth; Anguilla; Krebs!

- N. (Phos) *Beauui* Fisch. & Bern. station: — ? — ; habitat: Marie Galante; Beau!
- N. *Candei* d'Orb. station: — ? — ; Guadeloupe; Beau! de Cande!
- N. *candidissima* Ad. station: — ? — ; habitat: Jamaica; C. B. Adams!
- N. (Phos) *guadeloupensis* Petit. station: Between stones on 1 foot water at Carthagenia; Krebs! habitat: Guadeloupe; Petit! Beau! St. Johns; Cartagena in New-Granada; C. B. Adams! St. Thomas; St. Martha in New-Gran.; Krebs!
- N. *Hotessieri* d'Orb. station: — ? — ; habitat: Jamaica; Guadeloupe; Martinique; d'Orb.! (Moll. de Cuba pt. 2 pg. 142 pl. 21 fig. 40, 42).
- N. *olivacea* Desh. station: — ? — ; habitat: West-Indies; Jay's Catalogue! Reeve! Guadeloupe; Lamarck!
- N. *polygona* Lmk. station: — ? — ; habitat: Cuba; Martinique; Guadeloupe; Rio-Janeiro; d'Orb.! Moll. de Cuba pt. 2 140.
- N. (Strongylocera) *textilina* Mörch. station: — ? — ; habitat: Antillae; Mörch!
- N. *vibex* Say. station: — ? — ; habitat: Guadeloupe; Beau!

Buccinum Linné.

- B. *auriculatum* Lmk. (B. *coromandelinum* Lmk., B. *Caribaeorum* d'Orb.) station: between large rocks and in excavations of rocks & madrepores near the shore; habitat: Jamaica; St. Martha; Carthagenia; C. B. Adams! St. Thomas; St. Croix: St. Martha; Krebs! remark: on a table of C. B. Adams stands: „B. corom. Lmk. (Pollia tineta Conr.?)“

- B. pulchellum* Ad. station: — ? — ; habitat: Jamaica; St. Thomas; C. B. Adams!

Bullia Gray.

- B. aciculata* Lmk. (Bucc. monile Linné? Mörch) station: — ? — ; habitat: Antillae?; Mörch! Guadeloupe; Beau! Duchassain!
- B. — ?* station: — ? — ; habitat: St. Barth and Anguilla; Krebs!

Terebra Adanson.

- T. castanea* Kien. station: — ? — ; habitat: Guadeloupe; Beau!
- T. cinerea* Born. station: — ? — ; habitat: Antillae; (*T. accicularis* Lmk.) Mörch! Guadeloupe; St. Lucy; Cuba; d'Orb.!
- T. hastata* Gml. (syn: *T. lancinula* Gml. acc. to Mörch; *T. costata* acc. to C. B. Adams) station: — ? — ; habitat: Antillae; Mörch! St. Johns; Jamaica; C. B. Adams! Guadeloupe; Beau! Anguilla; Krebs!
- T. jamaicensis* Ad. station: it buries itself in white sand at low-water mark; habitat: St. Johns; Jamaica; Arecibo; C. B. Adams! Guadeloupe; Beau! St. Thomas; Krebs!
- T. rudis* Gray. station: — ? — ; habitat: St. Lucy; d'Orb.!
- T. sp. ind.* station: — ? — ; habitat: St. Martha, New-Gr. in the collection of H. Krebs!

Cancellaria Lamarck.

- C. Candeï* d'Orb. station: — ? — ; habitat: Martinique; de Cande!

- C. reticulata* L. station: — ? — ; habitat: Cuba; Guadeloupe; d'Orb.! Marie-Galante; Beau! St. Thomas; Tortola; St. Martin; Krebs!
- C. rugosa* Lmk. station: — ? — ; habitat: Guadeloupe; St. Thomas; Tortola; C. B. Adams! St. Thomas; Anguilla; St. Martin; Krebs! Marie-Galante; Beau!

Cassis Lamarck.

1. Semicassis Kleiner.

- C. abbreviata* Lmk. (*C. lactea* Kiener) station: — ? — ; habitat: Antillae; Mörch! Guadeloupe; Beau! St. Thomas; Krebs!
- C. bisulcata* Wag? station: — ? — ; habitat: Antillae; Mörch!
- C. cicatricosa* Meusch. (*C. recurvirostris* Gml. non Reeve; *C. granulosa* jun. Kien.) station: Antillae; Mörch! Guadeloupe; Beau! Duchassain! St. Thomas; St. Johns; Jamaica; Carthagenae; North-Caroline; C. B. Adams! Tortola; St. Croix; Krebs!
- C. cornuta* L. (*C. labiata* Chem.) station: — ? — ; habitat: Guadeloupe; Beau!
- C. gibba* Gml. (*C. globulus* Bolt.; *C. abbreviata* Lmk.; acc. to Mörch!) station: Antillae; Mörch! Carthagenae; St. Martin; Krebs!
- C. ventricosa* Mart. (*Cass. granulosa* Lmk.; *B. granulatatum* Born according to Mörch! *Bucc. tessellatum* Chem. according to Beau!) station: at certain seasons of the year in seaweed on 2 to 3 feet water and on the white sand on the beach; Krebs! habitat: Antillae; Mörch! Guadeloupe; Beau! St. Thomas; Krebs!

2. Cassis Browne.

- C. flammea* L. station: in 3 fathoms water on seaweed;

Krebs! habitat: Antillae; Mörch! Cuba; d'Orb.! Jamaica; C. B. Adams! Guadeloupe; Beau! Barbadoes; Antigua; Tortola; Krebs!

C. Madegascariensis Lmk. station: — ? — ; habitat: Antillae; Mörch! Jamaica; C. B. Adams! St. Thomas; St. Croix; Krebs!

C. tuberosa L. station: — ? — ; habitat: Antillae; Mörch! Cuba; d'Orb.! Jamaica; Bermuda; C. B. Adams! Guadeloupe; Beau! Duchassain! Antigua; St. Johns; Tortola; St. Martin; Barbadoes; Krebs!

3. Cypræcassis Stutchb.

C. testiculus L. (C. erumena Brug. accord. to C. B. Ad.!) station: — ? — ; habitat: St. Johns; Jamaica; Arecibo; Carthagenae; Bermuda; C. B. Adams! Guadeloupe; Beau! St. Croix; Tortola; St. Barth; Barbadoes; Trinidad; Carthagenae; Krebs!

Oniscia Lamarck.

O. Denisonii Reeve. station: — ? — ; habitat: Guadeloupe; Beau!

O. Oniscus L. station: — ? — ; habitat: Guadeloupe; Beau! St. Johns; St. Thomas; Jamaica; Arecibo; C. B. Adams! St. Thomas; Tortola; St. Barth; Anguilla; St. Martin; Trinidad; Krebs!

Dolium Lamarck.

D. galea L.? station: — ? — ; habitat: St. Thomas; R. Swift! Jamaica; C. B. Adams!

D. pennatum Mart. station: — ? — ; habitat: St. Johns; Jamaica; C. B. Adams! Guadeloupe; Beau! St. Johns; Tortola; St. Martin; St. Barth; Barbadoes; Trinidad; Krebs! remark: C. B. Adams above mentioned

habitats are marked by *D. perdix* L. which is an East-India shell.

Eburnea Lamarek.

- E. balteata** Sow. station: — ? — ; habitat: Antillae; Mörch!
E. glabrata Lmk. station: — ? — ; habitat: Curaçao; R. Swift! Jamaica; Curaçao; de Candé!
E. Tankervillii Swains. station: — ? — ; Island of Margarita; Swift!

Volutacea.

Voluta Linné.

- V. Beauii** Fischer & Bern. station: — ? — ; habitat: Marie Galante; Beau!
V. musica L. station: — ? — ; habitat: Cuba; Guadeloupe; St. Lucy; d'Orb.! Guadeloupe; Beau! Margarita; R. Swift! Barbadoes; Krebs! remark: not hitherto found at the Virgin-Islands; the largest ever seen were from the island of Margarita.
V. Guildingii Sowb. station: — ? — ; habitat: St. Vincent; Reeve! remark: Reeve's drawing of this shell the *lc. conch.* looks very much like *Mitra dermestina* Lamarek.
V. Vespertilio Rumphius. station: — ? — ; habitat: Antilles; d'Orb.!

Mitra Lamarek.

- M. alreolata** Reeve. station: — ? — ; habitat: Antillae; Mörch!
M. barbadensis Gmelin. (*M. striatula* Lmk.) station: on coral-reefs according to Guilding! habitat: St. Vincent;

Reeve! Cuba; de la Sagra! Auber! Martinique; Candé!
Guadeloupe; Beau! Duchassain! Hotessier! St. Thomas;
St. Johns; St. Barth; St. Martin; Barbadoes; Krebs!

M. candida Reeve. station: — ? — ; habitat: St.
Thomas; Lillienkjöld! Mörch! Krebs! St. Croix;
Krebs! remark: It is pale yellow or pink when very
fresh and perfect.

M. caelagaena Reeve. station: — ? — ; habitat:
Crab Island; A. H. Riise! St. Thomas; Krebs!

M. cavea Reeve. station: — ? — ; habitat: Guade-
loupe; Beau! remark: see *M. microzonias* var. *bifasciata*.

M. dermestina Lmk. (*M. speciosa* Reeve? Mörch!) sta-
tion: — ? — ; habitat: Antillae; Mörch! St. Thomas;
St. Johns; Passage-Island; St. Martin; Anguilla; Krebs!

M. microzonias Reeve? station: — ? — ; habitat:
Antillae; Mörch!

M. microzonias var. *bifasciata* (*M. cavea* Reeve; *M.*
microzonias Kiener.)

M. nodulosa Gml. (*M. granulosa* Lmk.) station: on rocks
Guilding! habitat: St. Vincent; Reeve! Cuba; de la
Sagra! Auber! St. Lucy; Martinique; Guadeloupe; Candé!
Hotessier! St. Thomas; St. Johns; Krebs!

M. nucleola Reeve. Does the drawing in Reeve's Ic. conch.
not represent a *M. nodulosa* Gml.?

M. pulchella Reeve. station: — ? — ; habitat: Bar-
badoes; Humphreys! Guadeloupe; Beau! remark: is it
not synonyme with *M. derme*. Lmk.?

M. semicostata Anton. station: — ? — ; habitat: St.
Croix; St. Martin; Anguilla; Krebs!

M. sulcata Gml. (*M. microzonias* Lmk.) station: — ? — ;
habitat: Antillae; Mörch!

M. sulcata bifasciata (*M. cavea* R.! *M. microz.* K.)

- M. puella* Reeve. station: — ? — ; habitat: St. Thomas; Gruner! Guadeloupe; Beau! St. Thomas; St. Croix; St. Johns; St. Martin; Krebs!
- M. tiarella* Lmk. station: — ? — ; habitat: West-Indies; Jay's Catalogue!

Oliva Brugière.

- O. angulata* Lmk. (*V. incrassata* Sol., *V. corticata* & *striatula* Mart.) station: — ? — ; habitat: Antillae; Mörch!
- O. attenuata* Reeve. station: — ? — ; habitat: Cuba; C. B. Adams!
- O. bullula* Reeve. station: — ? — ; habitat: Guadeloupe; Beau! St. Thomas; St. Barth; Krebs!
- O. dealbata* Reeve. station: — ? — ; habitat: Guadeloupe; Beau!
- O. Esther* Ducl. station: — ? — ; habitat: Antillae, Mörch!
- O. exigua* Mart. (*O. conoidalis* Lmk.; *Voluta jaspidea* Gml.; *Olivina* d'Orb.) station: — ? — ; habitat: Curaçao; Candé! Cuba; de la Sagra! Guadeloupe; Hostessier! St. Johns; Jamaica; Carthagera; Tortola; C. B. Adams! Barbadoes; Curaçao; St. Thomas; Krebs!
- O. flammulata* Lmk. (*Mica marmorata* Mart.) station: — ? — ; habitat: Antillae; Mörch!
- O. fimbriata* Reeve. station: — ? — ; habitat: Guadeloupe; Beau! St. Johns; Barbadoes; Jamaica; South-Caroline; Brasils; Marecaibo; Carthagera; C. B. Adams!
- O. fulgida* Reeve. station: — ? — ; habitat: St. Thomas; St. Barth; Anguilla; Krebs! Guadeloupe; Beau!
- O. Guildingii* Reeve. station: — ? — ; habitat: St. Vincent; Reeve! Jamaica; C. B. Adams!

- O. ispida* Link. (*O. fusiformis* Lmk.) station: — ?
— ; habitat: Guadeloupe; Beau! Jamaica; C. B. Adams! Great North-Side-Bay at St. Thomas; Krebs!
- O. leucostoma* Ducl. (*O. ancillaria* d'Orb.) station: — ?
— ; habitat: Antillae; d'Orb.!
- O. ligneola* Reeve. station: — ? — ; habitat: Guaranao in Venezuela; Swift!
- O. litterata* Lmk. station: — ? — ; habitat: Antillae; Mörch! Florida; R. Swift! Bermuda; C. B. Adams! Vera-Cruz; Krebs!
- O. mica* Ducl. station: — ? — ; habitat: Jamaica; de Candé!
- O. miliola* d'Orb. station: — ? — ; habitat: Jamaica; Martinique; de Candé!
- O. millepunctata* Ducl. station: — ? — ; West-Indies; Reeve! Antillae; Mörch!
- O. minuta* Link. (*O. ziczac* Ducl.) station: — ? — ; habitat: Antillae; Mörch!
- O. mitrola* Ducl. station: — ? — ; habitat: Antillae; Mörch!
- O. myriadina* Ducl. station: — ? — ; habitat: Cuba; de la Sagra! St. Thomas & Jamaica; de Candé! Jamaica; Bahamas; C. B. Adams!
- O. nana* Lmk. (*O. micans* Sol.) station: — ? — ; habitat: West-Indies; Reeve! Jamaica; de Candé! Antillae; Mörch! remark: should this not be synonyme with *O. strigata* Reeve?
- O. nitidula* Link. (*O. mutica* R.) station: — ? — ; habitat: Guadeloupe; Beau! South-Caroline; Jamaica; Portocabello; C. B. Adams! St. Thomas; St. Barth; Bahama; Krebs!
- O. nivea* d'Orb. station: — ? — ; habitat: St. Barth

de Candé! Guadeloupe; Hotessier! Cuba; de la Sagra! Auber! Florida; d'Orb.!

- O. oryza* Lmk. station: — ? — ; habitat: Guadeloupe; Beau! Bermuda; Bahama; Jamaica; C. B. Adams! St. Barth; St. Martin; Krebs!
- O. parvula* Mart. (*Voluta nivea* Gml., *Olivina* d'Orb., *Oliva eburnea* Lmk.) station: — ? — ; habitat: St. Barth; de Candé; Guadeloupe; Hotessier! Beau! Cuba; de la Sagra! Auber! Florida; d'Orbig.! Jamaica; C. B. Adams! St. Thomas; St. Croix; St. Barth; Anguilla; Barbadoes; Krebs! remark: we have specimens with gray, yellow, orange and rose ornaments; is it not synonyme with *Oliva nana* Lmk.?
- O. pulchella* Duclos. station: — ? — ; habitat: Florida; Jamaica; C. B. Adams! St. Thomas; Krebs! remark: very closely allied with *O. lanceolata* R.; we have seen specimens of *O. pulch. Duc.* with spots close to the suture.
- O. pura* Reeve. station: — ? — ; habitat: St. Martin?; Krebs!
- O. reticularis* Lmk. (*O. olivaceus* Meusch) station: we suppose it buries itself in sand; a very young life specimen has been dredged from blue sandy mud in 10 to 12 fathoms water; habitat: Cuba; Auber! St. Lucy; Guadeloupe; Hotessier! Florida; Alphen! Jamaica; Carthagera; Curaçao; Cumana; Bermuda; C. B. Adams! Guadeloupe; Beau! St. Thomas; St. Johns; St. Croix; St. Barth; Tortola; St. Martin; Anguilla; Trinidad; Sta. Martha in New-Granada; Krebs! remark: Beau gives as synonymes *O. araneosa* Lmk., *O. timoria* Ducl., *O. venulata* Ducl., *O. obesina* Ducl., *O. pindarina* Ducl.

- O. rosalina* Duclos. station: — ? — ; habitat: Guadeloupe; Duchassain! Jamaica; Turks-Island; C. B. Adams!
- O. rufifasciata* Reeve. station: — ? — ; habitat: St. Thomas; Krebs! remark: nothing else than a variety of *O. nitidula* Link.
- O. scripta* Lmk. station: — ? — ; habitat: Guadeloupe; Beau! Jamaica; C. B. Adams! Curaçao; St. Lucy; Cuba; d'Orbigny!
- O. strigata* Reeve. station: — ? — ; habitat: Hatfield, Jamaica; E. Chitty!

Cypracaceae.

Cypraea Linné.

- C. bicallosa* Gray. station: — ? — ; habitat: Marie Galante; Beau!
- C. cervus* L. (*C. cervina* Lmk.) station: — ? — ; habitat: Antillae; Mörch!
- C. cornea* Gray (*C. rosea* Kien.) station: — ? — ; habitat: Guadeloupe; Beau!
- C. exanthema* L. station: — ? — ; habitat: Cuba; de la Sagra! Auber! Martinique; Guadeloupe; St. Lucy! de Candé! Hotessier! Duchassain! Beau! St. Thomas; St. Barth; St. Martin; Krebs!
- C. globosa* Sow. (*C. pilula* Kiener.) station: — ? — ; habitat: Cuba; Auber! New-Orleans; d'Orbigny! Guadeloupe; Beau! St. Thomas; St. Croix; St. Johns; St. Barth; Anguilla; Krebs!
- C. mus* L. station: — ? — ; habitat: Cuba; de la Sagra! Martinique; de Candé! Curaçao; R. Swift!
- C. nivea* Gray. station: — ? — ; habitat: Guadeloupe; Beau! Cuba; Auber! New-Orleans; d'Orbigny!
- C. pediculus* L. station: — ? — ; habitat: Cuba; de

- la Sagra! Martinique; de Candé! Guadeloupe; Hotessier!
 Beau! St. Thomas; St. Croix; St. Johns; Tortola; St.
 Martin; St. Barth; Anguilla; Barbadoes; Trinidad; Krebs!
- C. rotunda* Kiener (*C. quadripunctata* Gray) station: — ?
 — ; habitat: Cuba; St. Lucy; New-Orleans; d'Orb.!
 St. Thomas; St. Croix; St. Johns; St. Barth; St. Mar-
 tin; Anguilla; Barbadoes; Trinidad; Jamaica; Krebs!
- C. spurca* Linné (*C. flavcola* Reeve; *C. lota* Born) sta-
 tion: — ? — ; habitat: Martinique; St. Lucy; Cuba;
 d'Orbigny! Guadeloupe; Beau! St. Thomas; St. Croix;
 St. Martin; Barbadoes; Krebs!
- C. stercoraria* L. station: — ? — ; habitat: Guade-
 loupe; Beau!
- C. succincta* L. (*C. cinerea* Mart.; *C. sordida* Link.) sta-
 tion: — ? — ; habitat: St. Thomas; Tortola; St.
 Barth; St. Martin; Anguilla; Barbadoes; St. Domingo;
 Krebs! Cuba; Martinique; Jamaica; d'Orbigny! Guade-
 loupe; Beau! Duchassain!
- C. suffusa* Gray (*C. armindina* Ducl.) station: — ? — ;
 habitat: Guadeloupe; Beau! St. Thomas; St. Croix;
 St. Barth; St. Martin; Anguilla; Barbadoes; Krebs!
- C. subrostrata* Gray. station: — ? — ; habitat:
 Guadeloupe; Beau! St. Thomas; St. Johns; St. Croix;
 St. Barth; Anguilla; Barbadoes; Krebs! remark: we
 have beside a white variety.
- C. tremeza* Ducl. station: — ? — ; habitat: West-
 Indies; Reeve! *as found in the West Indies*

Erato Risso.

- E. Maugeriae* Gray. station: dredged from blue sandy
 mud in 10 to 12 fathoms water; habitat: St. Thomas;
 St. Barth; Krebs! Guadeloupe; Beau!

Ovula Brugière.

- O. acicularis* Lmk. station: — ? — ; habitat: Cuba; d'Orbigny! St. Martin; Krebs! Guadeloupe; Beau!
- O. gibbosa* Lmk. station: — ? — ; habitat: Cuba; de la Sagra! Auber! Martinique; Guadeloupe; St. Lucy; Hotessier! de Candé! Guadeloupe; Beau! St. Thomas; St. Johns; Anguilla; Trinidad; Krebs!

Marginella Lamarck.

- M. albolineata* d'Orb. station: — ? — ; habitat: Cuba; Auber! Jamaica; C. B. Adams!
- M. apicina* Mke. (*M. flavida* Redf.) station: — ? — ; habitat: Bahama; Redfield! St. Thomas; Swift! Krebs!
- M. avena* Kien. station: — ? — ; habitat: Cuba; Auber! de Candé!
- M. candida* Redf. station: — ? — ; habitat: St. Thomas; St. Croix; St. Barth; Anguilla; Krebs!
- M. caribaea* d'Orb. station: — ? — ; habitat: Cuba; de la Sagra! Auber! St. Thomas; de Candé! remark: described in d'Orbigny's Moll. de Cuba pt. 2 pg. 97.
- M. catenata* Mont. station: — ? — ; habitat: Spanish-town; St. Martin; St. Barth; Guadeloupe; Krebs!
- M. chrysomelina* Redf. station: — ? — ; habitat: West-Indies; Krebs!
- M. conoidalis* Kiener. station: — ? — ; habitat: Antillae; Mörch! Havana; Jay! remark: is this not synonyme with *M. apicina* Mencke?
- M. crassilabrum* Sowb. station: — ? — ; habitat: West-Indies; Jay!
- M. Delessertiana* Recluz. station: — ? — ; habitat: Guadeloupe; Beau!
- M. interrupta-lineata* Muhl. (*M. interrupta* Lmk.) sta-

tion: — ? — ; habitat: Antillae; Mörch! Carthagenae; St. Martha; Redfield! St. Thomas; d'Orbigny! St. Martha; Carthagenae in New-Granada; Krebs! remark: d'Orbigny's locality „St. Thomas“ is surely a mistake; he may have bought it or received it from a collector in said place, but we doubt that it has been found at the island of St. Thomas, where we have collected shells in 17 years and never seen a trace of it; Mr. J. Redfield has told us that he always received *M. interrupta-lineata* Mühlf. and *M. obesa* Redf. intermixed when the shells came from Carthagenae, but only *M. interrupta*-l. when they came from Sta. Martha; Mr. Cotoy has told R. Swift Esq. that he *M. int-st.* was always found along the shore and *M. ob.* in 8 to 10 fathoms of water.

- M. frumentum* Sowb. station: — ? — ; habitat: West-Indies; Jay!
- M. fusca* Sowb. station: — ? — ; habitat: West-Indies; Jay!
- M. glans* Meneke. station: — ? — ; habitat: St. Domingo; Jay!
- M. lactea* Kiener (Pers. parva Mart.) station: — ? — ; habitat: Antillae; Mörch! Guadeloupe; Beau!
- M. Lavelleana* d'Orb. (*M. minima* Guild.) station: — ? — ; habitat: Jamaica; Martinique; de Candé! St. Thomas; R. Swift! Krebs!
- M. longivaricosa* Lmk. station: — ? — ; habitat: Cuba; Sagra!
- M. maculosa* Kien. station: — ? — ; habitat: Curaçao; R. Swift!
- M. marginata* Born. (*M. bivaricosa* Lmk.) station: — ? — ; habitat: Antillae; de Candé! Mörch! Curaçao; R. Swift! Sta. Martha; Krebs!

- M. muralis* Hinds. station: — ? — ; habitat: Curaçao; Mörch! Jamaica; Jay!
- M. nivea* Ad. station: — ? — ; habitat: Jamaica; Jay!
- M. obesa* Redfield (*M. similis* Sowb.) station: in 8 to 10 fathoms water along the coast of Venezuela and New-Granada; habitat: Sta. Martha; Redfield! Carthage; Jay!
- M. ovuliformis* d'Orb. station: — ? — ; habitat: Martinique; Guadeloupe; St. Thomas; de Candé! Guadeloupe; Beau!
- M. pallida* L. (*M. avenacea* Desh.) station: — ? — ; habitat: St. Thomas; St. Johns; St. Martin; St. Barth; Barbadoes; Krebs! Honduras; Jay! St. Domingo; Guadeloupe; de Candé! Cuba; Auber! Guadeloupe; Beau! remark: (Beau has as synonymes: *M. avena.* Kien. & *M. varia.* Sow. *Entirely different*.)
- M. pellucida* Pfr. (*M. diaphana* Ki.) station: — ? — ; habitat: Antillae; Mörch! St. Thomas; R. Swift! Krebs!
- M. phrygia* Sowb. station: — ? — ; habitat: Antillae; Mörch!
- M. pulcherrima* Ad. station: — ? — ; habitat: Bahama; Redf! remark: very like *M. catenata* Mont., but only ornamented with white spots, not with lines.
- M. pulchra* Gray. station: — ? — ; habitat: St. Domingo; Jay!
- M. rubella* Ad. station: -- ? — ; habitat: Jamaica; Adams! Jay! St. Thomas; Krebs!
- M. subcoerulea* Mart. (*V. prunum* Gm., *V. coerulescens* Lmk.) station: Antillae; Mörch! Curaçao; Candé! Swift! Guadeloupe; Beau! Sta. Martha; Krebs!
- M. sulcata* d'Orb. station: in mud on 4 to 6 fathoms

water; habitat: Martinique; Candé! St. Thomas; A. H. Riise!

M. subtriplicata d'Orb. station: — ? — ; habitat: St. Barth; St. Thomas; Candé! Guadeloupe; Beau! Guadeloupe; St. Martin; Anguilla; St. Johns; Krebs!

M. tessellata Lmk. station: — ? — ; habitat: Venezuela; Jay!

M. zonata Kien. station: — ? — ; habitat: West-Indies; Jay!

M. xanhostoma Mörch station: — ? — ; habitat: Curaçao; Mörch!

Volvaria Lamarek.

V. pellucida Schm. (*V. pallida* Lmk. non L.) station: — ? — ; habitat: Guadeloupe; d'Orb.! Beau! Duchassin! St. Thomas; St. Johns; St. Barth; Anguilla; St. Martin; Barbadoes; Krebs!

Turritella Lamarek.

T. bicingulata Lmk. (*Xylohelix* *T. marmoreus* minor Chem.) station: — ? — ; habitat: Antillae; Mörch!

T. caribaea d'Orb. station: — ? — ; habitat: Cuba; Sagra!

T. exoleta L. (*Haustator* Mtf.; *Terebellum* Browne; *T. torcularis* Born according to Beau) station: — ? — ; habitat: Jamaica; C. B. Adams! Guadeloupe; Beau! Porto-Plata; St. Martin; Trinidad; Krebs!

T. imbricata L. station: — ? — ; habitat: Antillae; Lamarek! Mörch! Jamaica; St. Lucy; d'Orbigny!

T. Meta Reeve. station: — ? — ; habitat: St. Martha; Carthagera; Krebs! remark: perhaps a variety of *T. imbr.* L.

T. variegata (*T. marmoratus* Ch.; *Elob. marmoreum* Bolt., *T. imbricata* Lmk.) station: — ? — ; habitat: Antillae; Mörch! Jamaica; St. Lucy; d'Orbigny! Carthageria; Redfield! Cumana; J. H. Newton! Island of Margarita and at Mayaguez in Puertorico; R. Swift! St. Thomas; Krebs!

Cerithium Adanson.

C. albovittatum Ad. station: — ? — ; habitat: Jamaica; C. B. Adams! remark: described in „Contributions“ No 7 page 122.

C. algicola Ad. station: — ? — ; habitat: Rum-Key, Turks-Island; Bahama; Jamaica; C. B. Adams! St. Thomas; Porto-Plata; Barbadoes; Krebs!

C. ambiguum Ad. (synonymes according to Beau: *C. costatum* Wood, *C. servile* C. B. Ad.) station: — ? — ; habitat: Jamaica; C. B. Adams! Guadeloupe; Beau! remark: described in „Proceedings of the Boston Soc. of Nat. History“. Jan. 1st. 1845 page 4.

C. atratum Born. station: on rocks close to the shore between the marks of high and low water; at other seasons on blue sandy mud in 2 fathoms water; habitat: Guadeloupe; Lamarck! Beau! Rio-Janeiro; Jamaica; C. B. Adams! Key West; J. H. Newton! St. Thomas; Krebs!

C. bicolor Ad. station: — ? — ; habitat: Jamaica; C. B. Adams! St. Thomas; Th. Bland! Arecibo; J. H. Newton! remark: described in the „Proceedings of the Boston. S. of Nat. Sc.“ 1st January 1845 page 5.

C. columellare d'Orb. station: — ? — ; habitat: St. Thomas; Guadeloupe; St. Domingo; Jamaica; Cuba; Candé! Guadeloupe; Beau!

- C. dissimile* Yoldi station: — ? — ; habitat: Antillae; Mörch!
- C. eburneum* Brug. station: — ? — ; habitat: Antilles; Lamarek! Guadeloupe; Martinique; Cuba; d'Orb.! Guadeloupe; Beau! St. Thomas; Krebs!
- C. ericense* Kien. station: — ? — ; habitat: Guadeloupe; Beau!
- C. exile* Ad. station: — ? — ; habitat: Jamaica; C. B. Adams! remark: described in „Contributions“ No 7 April 1850 page 120.
- C. ferrugineum* Say. station: — ? — ; habitat: Guadeloupe; Beau!
- C. flavum* Ad. station: — ? — ; habitat: Jamaica; C. B. Adams! remark: described in „Contributions“ No 7 page 122 April 1850.
- C. fusiforme* Ad. station: — ? — ; habitat: Jamaica; C. B. Adams! remark: described in „Contributions“ No 7 page 120 April 1850.
- C. gemmulosum* Ad. station: — ? — ; habitat: Jamaica; C. B. Adams! St. Thomas; Krebs! remark: described in „Contributions“ No 7 page 120 April 1850.
- C. gibberulum* Ad. station: in myriades on mud in 8 to 15 fathoms water; habitat: Jamaica; C. B. Adams! St. Thomas; St. Johns; St. Martin; Porto-Plata; Krebs! remark: described in the „Proc. of the Boston S. of Nat. History“ 1st January 1845 page 5.
- C. interruptum* Mke.? station: — ? — ; habitat: Saba; R. Swift! St. Thomas; St. Martin; Porto-Plata; Krebs!
- C. iota* Ad. station: — ? — ; habitat: Jamaica; Ad.! remark: described in „Proc. of the Boston S. of Nat. History“ January 1st 1845.

- C. *Lafondii* Mich. station: — ? — ; habitat: Mediterranean sea; Lmk.! Jamaica; Ad.!
- C. *Lavalleanum* d'Orb. station: — ? — ; habitat: Cuba; d'Orb.!
- C. *lima* Brug. station: — ? — ; habitat: Guadeloupe; Lmk.! d'Orb.! Beau! Cuba; Martinique; d'Orb.!
- C. *litteratum* Brug. station: young specimens dredged from 9 fathoms water on blue sandy mud; habitat: Cuba; Sagra! Auber! Martinique; Guadeloupe; Candé! Carthagera; Bermuda; Redfield! Jamaica; Ad.! St. Thomas; Tortola; St. Martin; Barbados; Trinidad; Krebs!
- C. *megasoma* Ad. station: — ? — ; habitat: Jamaica; Ad.! remark: described in „Contributions“ No 7. April 1850 page 119.
- C. *mirabile* Ad. station: — ? — ; habitat: Jamaica; Adams! remark: described in the 7te „contribution“ page 118.
- C. *muscarum* Say. station: Jamaica; Ad.! Cuba; Florida; d'Orb.!
- C. *mutabile* Ad. station: — ? — ; habitat: Jamaica; Ad.! remark: described in „Proc. of the Boston Soc. of N. Hist.“ 1st January 1845 page 5.
- C. *minimum* Gml. (C. *zonale* Brug.) station: on rocks in the surface of the water close to the shore and on the beach where there is coarse gravel; only where the sea is calm; habitat: Curaçao; Mörch! St. Thomas; Tortola; Carthagera; Krebs! remark: it varies very much and several shells described as separate species will in the future be shown to be synonymes with this.
- C. *nigrescens* Mk. station: on rocks in the surface of the water along the shore; habitat: Guadeloupe; Petit! Beau! St. Thomas; St. Johns; Jamaica; Florida; Ad.!

- St. Thomas; St. Martha; Carthagera; Krebs! remark: it is surely synonyme with *C. septemstriatum* Say. and perhaps with *C. minimum* Gml.
- C. ornatum* Duk. station: — ? — ; habitat: Antillae; Mörch!
- C. Petitii* Kien. station: — ? — ; habitat: St. Croix; Saba; R. Swift! Guadeloupe; Duchassain! Porto-Plata; St. Martin; Krebs!
- C. punctatum* L. (according to Beau: *C. subulatum* Montf. *C. Emersonii* C. B. Ad.) station: — ? — ; habitat: Guadeloupe; Beau! Duchassain! St. Thomas; Porto-Plata; Krebs!
- C. rugulosum* Ad. station: — ? — ; habitat: Jamaica; Ad.! St. Thomas; St. Johns; St. Martin; Krebs!
- C. scalariforme* Say. station: — ? — ; habitat: Jamaica; E. Chitty! Florida; Ad.!
- C. Sagra d'Orb.* station: — ? — ; habitat: Cuba; Auber!
- C. semiferrugineum* Lnk. station: — ? — ; habitat: Guadeloupe; Beau! St. Thomas; St. Martin; Krebs! remark: a variety of *C. litteratum* Brug.
- C. septemstriatum* Say. station: — ? — ; habitat: Bahama; Ad.! remark: see *C. nigrescens* Mnk.
- C. terebellum* Ad. station: — ? — ; habitat: Jamaica; Ad.! St. Thomas; St. Martin; Pto.-Plata; Krebs!
- C. versicolor* Ad. station: — ? — ; habitat: Jamaica; Ad.! Guadeloupe; Beau! St. Thomas; Bland! remark: described in C. B. Ad. „Contribution“ No 7 page 119. April 1850.
- C. variabile* Ad. station: — ? — ; habitat: Jamaica; Ad.! remark: described in „Proc. of the Boston Soc. of N. Hist.“ 1st January 1854.

C. vicinum Ad. station: — ? — ; habitat: Jamaica;
Ad! remark: described in 7te „Contribution“ of C. B.
Adams page 122.

Triforis Deshayes.

- T. dealbatus* Ad. station: — ? — ; habitat: Jamaica;
Ad! remark! described in „Contribution“ No 7 page
117. April 1850.
- T. decoratus* Ad. station: — ? — ; habitat: Jamaica;
Ad! remark: described in „Contribution“ No 7 page
117. April 1850.
- T. exiguus* Ad. station: — ? — ; habitat: Jamaica;
Ad! remark: described in „Contribution“ No 7 page
118. April 1850.
- T. intermedius* Ad. station: — ? — ; habitat: Ja-
maica; Ad! remark: described in „Contribution“ No
7 page 119. April 1850.
- T. melanura* Ad. station: — ? — ; habitat: Jamaica;
Ad! remark: described in „Contribution“ No 7 page
117. April 1850.
- T. mirabilis* Ad. station: — ? — ; habitat: St. Johns;
Jamaica; Ad! remark: described in „Contribution“ No
7 page 118. April 1850.
- T. modestus* Ad. station: — ? — ; habitat: Jamaica;
Ad! remark: described in „Contribution“ No 7 page
117. April 1850.
- T. nanus* Ad. station: — ? — ; habitat: Jamaica;
Ad! remark: described in „Contribution“ No 7 page
117. April 1850.
- T. ornatus* Desh. (Hinds according to Beau) station: — ?
— ; habitat: Jamaica; Ad! Guadeloupe; Beau! St.
Thomas; St. Croix; Krebs!

T. pulchellus Ad. station: — ? — ; habitat: Jamaica; St. Thomas; Ad.! remark: described in „Contribution“ No 7 page 121. April 1850.

T. turris Thomae d'Orb. station: — ? — ; habitat: Cuba; Guadeloupe; d'Orb.! Guadeloupe; Beau!

Paludinacea.

Litiopa Rang.

L. effusa Ad. station: — ? — ; habitat: Jamaica; Ad.! remark: described in „Contribution“ No 5 page 71.

L. maculata Rang. station: On seaweed in the gulfstream Ad.! habitat: Jamaica; Cuba; d'Orbig.! St. Thomas; Krebs!

L. obesa Ad. station: — ? — ; habitat: Jamaica; Ad.! St. Thomas; Krebs! remark: described in the „Contribution“ No 5 page 71; the nucleus of this species is strongly costulated like that of *L. mac. R.*

L. striata Pf. station: on sea-weed; Ad.! habitat: in the gulfstream; Ad.!

Planaxis Lamarck.

P. nucleus Lmk. station: on rocks between high and low water mark; habitat: Martinique; Guadeloupe; St. Lucy; Cuba; Auber! Candé! Beau! Jamaica; Ad.! St. Thomas; Sta. Martha; St. Johns; St. Martin; Anguilla; Krebs!

P. lineatus Da Costa (*P. pedicularis* Lmk.) station: on rocks and in hollows between high and low water mark; habitat: Antillae; Mörch! England; Da Costa! Java; Lmk.! St. Vincent (Canary-Islands); Schmidt! Jamaica; Ad.! Guadeloupe; Petit! Beau! St. Johns; St. Martin; St. Thomas; Krebs!

Rissoa Flemingville (Cingula; Alvania).

- R. aberrans** Ad. station: — ? — ; habitat: Jamaica; Ad.! Vieques; A. H. Riise! remark: described in „Contribution“ No 7 page 113.
- R. affinis** Ad. station: — ? — ; habitat: Jamaica; Ad.! remark: described in „proceedings of the Boston Soc. of Nat. hist.“ 1st January 1845 page 6.
- R. albida** Ad. (Rissoina acc. to von Möhrenberg) station: — ? — ; habitat: Jamaica; Ad.! St. Thomas; A. H. Riise! remark: described in „Proceed. of the Boston Soc. of Nat. History.“ 1st January 1845 page 6.
- R. Auberiana** d'Orb. station: dead in the sand of the beach; habitat: Cuba; Sagra! St. Thomas; Jamaica; Candé! St. Thomas; Krebs! remark: described in „Mollusques de Cuba“. 2d. part, page 22.
- R. canaliculata** (Odostomia) Ad. station: — ? — ; habitat: Jamaica; Ad.!
- R. caribaea** d'Orb. station: dead in the sand of the beach; habitat: Jamaica; de Candé! Cuba; Sagra! remark: described in „Mollusques de Cuba“ part 2 page 21.
- R. carinata** Recluz. station: — ? — ; habitat: St. Thomas! A. H. Riise!
- R. chesnellii** v. *major* Mich. station: — ? — ; habitat: Cuba; A. H. Riise!
- R. concinna** (Cingula) Ad. station: — ? — ; habitat: Jamaica; Ad.! remark: described in the „Contribution“ No 5 page 70.
- R. conica** (Cingula?) Ad. station: — ? — ; habitat: Jamaica; Ad.! remark: described in „Contribution“ No 5 page 70.
- R. crassicosta** Ad. station: — ? — ; habitat: Ja-

- maica; Ad.! remark: described in „Proc. of the Boston Soc. of Nat. History“. 1st January 1845 page 6.
- R. *dubiosa* Ad. station: — ? — ; habitat: Jamaica; Ad.! St. Thomas; Th. Bland! remark: described in „Contributions“ No 7 page 114.
- R. *eulimoides* Ad. station: — ? — ; habitat: Jamaica; Ad.! remark: described in „Contribution“ No 7 page 115.
- R. *gemmulosa* (Odostomia) Ad. station: — ? — ; habitat: Jamaica; Ad!
- R. *gradata* d'Orb. station: in the sand of the beach; habitat: Jamaica; d'Orb.! remark: described in „Mollusque de Cuba“ part 2d. page 23.
- R. *laevigata* Ad. (Rissoina; Möhrenstern) station: — ? — ; habitat: Jamaica; Ad.! St. Thomas; A. H. Riise! remark: described in „Contribution“ No 7 page 114.
- R. *laevissima* Ad. (Rissoina; Möhrenstern) station: — ? — ; habitat: Jamaica; Ad.! St. Thomas; Bland! A. H. Riise! remark: described in „Contribution“ No 5 page 115.
- R. *melanura* Ad. station: — ? — ; habitat: Jamaica; Ad.! St. Thomas; A. H. Riise! Guadeloupe; Trinidad; St. Martin; St. Croix; St. Thomas; Krebs! remark: we agree with C. B. Adams, who has described this species in „Contribution“ No 8 page 116, that it is not a Rissoa and we take it to be the shell described by d'Orbigny as *Eulina incerta*, but he himself doubts it is an *Eulina*; we doubt it to be either a Rissoa or an *Eulina*; we have specimens with two and three strongly enveloped varices on each whorl and in some few specimens are the varices forming two or three continuous

rows from the, in the most cases, black nucleus until the last whorl.

R. *multicostata* Ad. station: — ? — ; habitat: Jamaica; Ad.! remark: described in the „Contribution“ No 7 page 114, in which Ad. refers to *Rissoina elegantissima* d'Orb., with which we consider it synonyme — Professor C. B. Adams, who has contributed so much to the development of the natural history of the West-Indies and who, by his practical way of collecting and arranging as he collected, has done more for the knowledge of the geographical distribution of the marine shells on both sides of the american continent, than any man of science before him, had quite a sickly passion for describing „supposed new shells“, by which he, as well as many of his contemporaries, has confused and burdened the science with innumerable synonymes, which men of science in future days will have trouble to find through.

R. *ovuloides* (*Odostomia*) Ad. station: — ? — ; habitat: Jamaica; Ad.!

R. *princeps* Ad. station: — ? — ; habitat: Jamaica; Ad.! remark: described in his „Contribution“ No 7 page 116. where he says it is allied to *R. albida* Ad.

R. *pulchra* Ad. (*Rissoina* acc. to Möhrenberg) station: — ? — ; habitat: Jamaica; Ad.! St. Thomas; Th. Bland! A. H. Riise! St. Barth; Krebs! remark: described by C. B. Adams in „Contribution“ No 7 page 114; very near allied to *Rissoina Sagraia* d'Orb.! and *Rissoa cancellata* Phil.

R. *pusilla* Desh. station: — ? — ; habitat: Cuba; A. H. Riise!

R. *solida* (*Odostomia*) Ad. station: — ? — ; habitat:

Jamaica; Ad.! remark: he describes it in „Contribution“ No 5 page 70 as *Cingula solida*.

- R. *scalarella* Ad. (*Rissoina Möhrenberg*) station: — ? — ; habitat: Jamaica; Ad.! St. Thomas; Th. Bland! A. H. Riise! remark: described in „Proc. of the Boston S. of Natural History.“ 1st January 1845. see *Rissoina Catesbya d'Orbigny* and „Contribution“ No 7 page 113.
- R. *scalaroides* Ad. (*Rissoina Möhrenberg*) station: — ? — ; habitat: Jamaica; Ad.! St. Thomas; A. H. Riise! remark: described in „Contribution“ No 7 page 113; by Philippi considered a *Rissoina*; see *Zeitschrift* 1848 page 13.
- R. *scalaroides minor* Ad. station: — ? — ; habitat: Jamaica; Ad.! remark: described in „Contribution“ No 7 page 113.
- R. *striosa* Ad. station: in sand on the beach; Ad.! habitat: Jamaica; Ad.! St. Thomas; Th. Bland! remark: described in „Contribution“ No 7 page 116.
- R. *subangulata* Ad. station: — ? — ; habitat: Jamaica; Ad.! remark: described in „Contribution“ No 7 page 112.
- R. *trivariosa* Ad. station: — ? — ; habitat: Jamaica; Ad.! St. Croix; A. H. Riise! remark: described in „Proc. of Boston Soc. of Natural History.“ 1st January 1845 page 6. see *R. melanura* Ad.
- R. *umbilicata* Ph. station: — ? — ; habitat: St. Thomas; A. H. Riise!
- R. *vitrea* Ad. station: — ? — ; habitat: Jamaica; Ad.! remark: described in „Contribution“ No 7 page 113.

Rissoina d'Orbigny.

- R. albida* Ad. station: — ? — ; habitat: St. Thomas; A. H. Riise!
- R. Browniana* d'Orb. station: In sand on the beach; habitat: St. Thomas; Haity; Martinique; de Candé! St. Thomas & St. Johns; Krebs! remark: described in „Mollusques de Cuba“ 2 part, page 28.
- R. cancellata* Ph. station: dead in the sand on the beach; habitat: Cuba; Pfeiffer! remark: described in Zeitschrift 1847 page 127.
- R. Catesbya* d'Orb. station: dead in the white sand on the beach and life on dead madrepores laying on the beach between high and low water mark; habitat: Cuba; Sagra! St. Thomas; R. Swift! Martinique; St. Thomas; Haity; Candé! Portoplata; St. Thomas; St. Croix; St. Johns; St. Martin; Krebs! Guadeloupe; Beau! St. Thomas; A. H. Riise! remark: This *R. Cat. d'Orb.* is certainly the same as the shell A. H. Riise Esq. has in his Collection by the name of *Rissoa Dunkeri* Pf. and we consider it the same shell as *C. B. Adams Rissoa scalaroides*, *R. scalaroides minor* and *R. scalarella*. The number of ribs and their proportion are very variable; we have a large number of specimens before us and we have come to the conclusion that they are one species.
- R. decussata* Montg. station: — ? — ; habitat: St. Thomas; A. H. Riise!
- R. dubiosa* C. B. Ad. station: — ? — ; habitat: St. Thomas; A. H. Riise!
- R. elegantissima* d'Orb. station: only found dead in the sand on the beach; habitat: Haity; d'Orbig.! St. Martin; Anguilla; Krebs! remark: Described in „Mollusques de Cuba“ part 2, page 26.

- R. *Phillippiana* Pfeiff. station: — ? — ; habitat: Viegues; Cuba; A. H. Riise!
- R. *reticulata* Sowb. station: — ? — ; habitat: St. Thomas; A. H. Riise!
- R. *Sagraina* d'Orb. station: — ? — ; habitat: Martinique; St. Thomas; Candé! remark: described in „Mollusque de Cuba“ part 2d. page 25. See *Rissoa pulchra* Ad.
- R. *scalariformis* Ad. station: — ? — ; habitat: St. Thomas; A. H. Riise!
- R. *scalaroides* see *Rissoa scalaroides* Ad. and *Rissoina Catesbya* d'Orb.!
- R. *Sloania* d'Orb. station: only found dead in the sand of the beach; habitat: Jamaica; Candé! remark: described in „Mollusques de Cuba“ part 2 page 28.
- R. *striato-costata* d'Orb. station: dead in the sand of the beach; habitat: Cuba; Sagra! Haity; Candé! St. Barth; Passage-Island near St. Thomas; Porto-Plata; Krebs! remark: described in „Mollusques de Cuba“ part 2d page 27.

Fossarus Adans.

- F. *sulcatus* d'Orb. station: — ? — ; habitat: Guadeloupe; Beau! remark: we are of opinion it is a young *Narica*.

Littorina Ferussac.

- L. *angulifera* Lmk. (*Littorina scabra* d'Orb.) station: Always on Mangletrees; d'Orb.! On bulwarks and old wood close to the shore; Krebs! habitat: Brasils; Mörch! All the Antilles; Jamaica; d'Orb.! Cuba; Sagra! Martinique; Guadeloupe; Rang! Candé! Saint Cur. Ho-

tessier! St. Thomas; St. Johns; St. Croix; Portorico; Jamaica; Krebs!

L. Antonii Phil. station: — ? — ; habitat: Jamaica; Ad.! St. Thomas; Th. Bland! Antilles; Mörch!

L. carinata d'Orb. Moll. de Cuba 1. 209; station: — ? — ; habitat: Havana; Sagra! Auber! Martinique; Candé! Rang! Antilles; Mörch! remark: d'Orbigny observes that those at Martinique are smaller than those at Havana. It would be very interesting to study if a species has a centre where the individuals envelope themselves to the greatest size and to the highest degree of perfection and from there toward the radius decrease in size and perfection. Several series of shells in our collection above us to cherish this idea.

L. columellaris d'Orb. station: On rocks; Beau! habitat: Pernambuco; d'Orb.! Martinique; Candé! Point à Pitre; Guadeloupe; Beau! Cat. pg. 12; Havana; K. Prescott! remark: Described in „Mollusques de Cuba“ p. 1 p. 213.

L. Gundlachi Ph. station: — ? — ; habitat: Cuba; Gundlach! remark: Described in „Zeitschrift“ 1848—150.

L. guttata Ph. (*L. punctata* Pf.) station: Found on poles in quiet water, close to the beach; habitat: Antilles; Mörch! Guadeloupe; Petit! Beau! Duchassain! St. Thomas; Bland! Jamaica; Ad.! St. Thomas; St. Johns; St. Martin; Porto-Plata and Carthagera (N. Gr.); Krebs!

L. dilatata d'Orb. station: — ? — ; habitat: Havana; Sagra! Auber! K. Prescott! Florida; Alphen! Porto-Plata; Krebs! Jamaica; Ad.! remark: Described in „Mollusques de Cuba“ 1—207.

L. flava Brod. (*L. irrorata* var. Petit) station: — ? — ; habitat: Brasils (?); Ad.! Guadeloupe; Petit!

- L. lineata* d'Orb. station: On rocks; Beau! Especially in small caves on the rocks between high and low-water mark; Krebs! habitat: Havana; Sagra! Auber! Martinique; Candé! Rio Janeiro; d'Orbigny! Guadeloupe; Beau! Duchassain! St. Thomas; Th. Bland! Krebs! St. Johns; E. Hartwig! Krebs! Jamaica; Ad.! Arecibo; J. H. Newton! St. Martin; Porto-Plata; Chagres; Krebs! remark: Described in „Mollusque de Cuba“ 1—208. Those d'Orbigny had received from Martinique were the largest, those next were from Havana and the smallest from Rio Janeiro; the largest we have are from St. Thomas the smallest from Porto-Plata.
- L. jamaicensis* Ad. station: — ? — ; habitat: Jamaica; Ad.! remark: Described in „Contribution“ No 5 page 71.
- L. minima* Wood. station: Very common on the poles of the wharfs; Beau! habitat: West-Indies; S. Hovey! Antilles; Mörch! Guadeloupe; Beau! St. Martin; Krebs! St. Croix; Riise!
- L. muricata* Linné (*Tectus muricatus* Ch.? *gothicus* Bolt?) station: On rocks always over high water mark; habitat: Antillae; Mörch! Cuba; Sagra! Auber! Martinique; Candé! St. Lucy and Florida; d'Orbigny! Guadeloupe; Beau! St. Martin; St. Thomas; Krebs!
- L. mespilum* Mode. (*L. fusca* Pfr. according to C. B. Adams) station: — ? — ; habitat: Jamaica & Florida; Ad.! Havana; K. Prescott!
- L. naticoides* d'Orbig. station: — ? — ; habitat: Havana; Sagra! Auber! remark: Described in „Moll. de Cuba“ 1—214.
- L. nodulosa* Gml. station: — ? — ; habitat: Martinique; Candé! St. Lucy! d'Orbig.! Jamaica; Ad.! Arecibo;

J. H. Newton! Havana; K. Prescott! remark: Described in „Moll. de Cuba“ 1—205.

- L. *nodulosa* Ph. station: — ? — ; habitat: Guadeloupe; Beau!
- L. *d'Orbigniana* Phil. station: — ? — ; habitat: Guadeloupe; Beau! Cat. pg. 12; Carthagenä (New Gr.); Krebs!
- L. *scabra* Linné. station: — ? — ; habitat: Guadeloupe; Beau! Mörch! 46. Ind. orient. d'Orbig. 1 Moll. de Cuba pg. 212.
- L. *tesselata* Phil. station: — ? — ; habitat: Jamaica; Ad.! St. Johns; E. Hartwig! Guadeloupe; Beau! Cat. pg. 12.
- L. *tigrina* d'Orb. station: In clusters on wood laying on the beach between high & low water mark; habitat: Havana; Sagra! Auber! Guadeloupe; Saint Cyr Hotessier! Chagres; Krebs! remark: Those we found at Chagres were, with exception of one specimen, of a uniform pale colour.
- L. *trochiformis* Dillw. (L. *nodulosa* d'Orb. acc. to Beau) station: — ? — ; habitat: Guadeloupe; Beau! St. Martin; Krebs! Jamaica; Ad.! Havana; K. Prescott! Tortola; E. Hartwig!
- L. *tuberculata* d'Orb. (L. *nodulosa* Ph. acc. to Beau) station: On rocks; d'Orb.! habitat: Cuba; Sagra! Auber! Porto-Plata; Krebs! Guadeloupe; Beau! remark: Described in „Moll. de Cuba“ 1. 206.
- L. *undulata* d'Orb. station: — ? — ; habitat: Martinique; Candé! remark: Described in „Moll. de Cuba“ 1—212.
- L. *ziczac* Chem. station: In crevices of the rocks between high and low water mark; Krebs! on the coast;

Beau! habitat: Antilles; Mörch. pag. 46! Havana; Sagra! Auber! Martinique; Candé! Guadeloupe; Beau. Cat. pg. 12! St. Thomas; Krebs! Th. Bland! E. Hartwig; Barbados; Chemnitz! Lmk.! Jamaica; Ad.! Arecibo; J. H. Newton! Carthagen (N. Gr.); Redfield!

Modulus Gray.

- M. lenticularis* Ch. (Trochus unidens List d'Orb.) station: On sandy mud about 2 to 4 feet water, especially on sea-weed; habitat: St. Thomas; St. Martin; St. Barth; Guadeloupe; St. Martha; Carthagen; Krebs! Cuba; Sagra! Auber! Martinique; Candé! Guadeloupe; Hotessier! Petit! Beau! Florida; d'Orb.! Ad.! St. Johns; E. Hartwig! St. Thomas; Th. Bland! Carthagen; Redfield; Jamaica; Ad.! Antillae; Mörch pag. 45!
- M. perlatus*. station: On plants close to the coast; Krebs! habitat: Carthagen N. G.; Krebs! remark: Decidedly different from the former.
- M. — ?* station: — ? — ; habitat: Virgin-Islands; Krebs!

Solarium Lmk.

- S. aethiops* Menke. station: — ? — ; habitat: Puer-torico; Philippi! remark: Described in „Zeitschrift“ 1848 page 167.
- S. bisulcatum* d'Orb. (Euomphalus Sowb.?) station: Dead in sand on the beach; habitat: Jamaica; Martinique; d'Orb.! Jamaica; Ad.! Arecibo; J. H. Newton! remark: Described in „Moll. de Cuba“ 2 part 66.
- S. delphinuloides* d'Orb. station: Found in sand; habitat: Jamaica; d'Orb.! remark: Described in „Moll. de Cuba“ 2—67.

- S. granulatum* Lmk. station: — ? — ; habitat: Guadeloupe; Hotessier! Cuba; d'Orb.! Haity; Mazatlan; (?Kr.) Menke! Jamaica; Ad.! Carthageria; St. Martha; Porto-Plata; Krebs! remark: The young specimens are perfectly smooth on the surface.
- S. Herbertii* Dh. (*Heliacus* d'Orb. *Tr. cylindraceus* Chem. *Tr. cylindricus* Gml.) station; Inserted in slimy zoophytes on rocks about low-water mark; habitat: Cuba; Auber! Jamaica; Ad.! St. Thomas; Krebs! Guadeloupe; Beau! Hotessier! Antillae; Mörch pag. 47!
- S. infundibuliformis* Chem. (*Tr. Chemnitzii* Ki.) station: — ? — ; habitat: Porto-Plata; St. Thomas; St. Martin; Krebs! Antillae Mörch pag. 47!
- S. inornatum* d'Orb. station: In sand; habitat: St. Thomas; Candé! remark: Described in „Moll. de Cuba“ 2—67.
- S. nubilum* Menke. station: — ? — ; habitat: Cap. Haity; Philippi!

Scalariacea.

Scalaria Lamarck.

- S. albida* d'Orb. (*Sc. fragilis* Hanl. acc. to Beau) station: — ? — ; habitat: Cuba; Auber! Guadeloupe; Beau! remark: Should this not be Sowerby's *Scalaria tenuis*?
- S. Candearia* d'Orbig. station: — ? — ; habitat: Jamaica & St. Thomas; Candé! remark: The specimens described by d'Orbigny were found dead in the sand on the beach.
- S. coronata* Lmk. station: — ? — ; habitat: Guadeloupe! Beau pg. 12!
- S. crenata* Linné. station: — ? — ; habitat: Jamaica; Ad.! Arecibo; J. H. Newton! St. Thomas; St. Martin;

- Anguilla; Krebs! remark: See our remark on *Sc. Hotessieriana* d'Orbig.
- S. denticulata* Sowb. station: — ? — ; habitat: West-Indies; Jay's cat! Guadeloupe; Beau!
- S. echinaticosta* d'Orb. station: The specimen described found in the sand on the beach; habitat: St. Thomas; d'Orbig! Krebs! R. Swift!
- S. foliaceicostata* d'Orb. station: — ? — ; habitat: Guadeloupe; Beau! Martinique; Guadeloupe; St. Thomas; Candé! Anguilla; St. Thomas; Porto-Plata; Krebs! remark: Described in „Moll. de Cuba“ 2—12.
- S. fragilis* Hanley. station: — ? — ; habitat: Guadeloupe; Petit! Jamaica; Ad.!
- S. Hotessieriana* d'Orb. station: — ? — ; habitat: Guadeloupe; Hotessier! Beau pg. 12! St. Martin; Porto-Plata; Krebs! remark: Take d'Orbigny's drawing of *S. Hot.* and Lamareks description of *Scalaria crenata* Linné; they approach surely each other very much.
- S. lamellosa* Lmk. (*Sc. clathrus* Linné acc. to Beau) station: Young specimens found on pieces of dead madrepores in a depth of water of 3 feet; habitat: Seas of Europe; Lamarek! St. Thomas; Th. Bland! St. Johns; J. Mac-Murry! Jamaica; Ad.! Cuba; Auber! Guadeloupe; St. Lucy; Hotessier! Candé! St. Thomas; St. Johns; Passage-Island; St. Martin; Anguilla; Guadeloupe; Krebs! Guadeloupe; Beau pag. 12!
- S. lineata* Say. station: — ? — ; habitat: West-Indies; according to a note in Amherst Museum! Guadeloupe; Beau pag. 12!
- S. ligata* Ad. station: — ? — ; habitat: Jamaica; Ad.! remark: Described in Contribution No 4 page 67. January 1850.

- S. Martini* Wood? station: — ? — ; habitat: St. Barth; St. Martin; Krebs!
- S. modesta* Ad. station: — ? — ; habitat: Jamaica; Ad.! Porto Cabello; R. Swift! remark: *S. venosa* Sowb. is according to Jay's Catalogue synonyme with this species.
- S. muricata* Kur. station: — ? — ; habitat: Florida; Ad.! St. Thomas; Th. Bland! Guadeloupe; Petit!
- S. pernobilis* Fisch. & Bern. station: — ? — ; habitat: Marie Galante; Beau!
- S. tenuis* Sowb. station: — ? — ; habitat: St. Thomas; Th. Bland! St. Thomas; St. Johns; St. Croix; St. Martin; Barbados; Trinidad; Porto-Plata; Krebs! Guadeloupe; Beau pg. 12!
- S. uncinaticosta* d'Orb. station: — ? — ; habitat: Guadeloupe; Candé! Beau pg. 12! St. Thomas; Krebs! R. Swift! remark: Described in „Moll. de Cuba“ 2—19.
- S. varicosa* Lmk. station: In mud on 1 to 2 feet water; habitat: Tortola; R. Swift! St. Croix; H. Krebs!

Ianthinacea.

Ianthina Lamareck.

- I. exigua* Lmk. station: — ? — ; habitat: Guadeloupe; Beau!
- I. fragilis* Ad. station: — ? — ; habitat: Jamaica; Ad.!
- I. globosa* Sow. station: — ? — ; habitat: Jamaica; Ad.!
- I. planispira* Reeve. station: floating on shore on sandy beaches; habitat: St. Thomas; A. H. Riise! Guadeloupe; Beau! St. Thomas & St. Johns; Krebs!

- I. prolongata* Blainville. station: — ? — ; habitat: Cuba; d'Orb.! Jamaica; Ad.!
- I. umbilicata* d'Orb. station: — ? — ; habitat: Cuba; d'Orb.! remark: Described in 2d. part page 85 in mollusques de Cuba.

Stylina Fleming.

- S. subulata* Brod. station: Found dead in holes bored by Littophages in madrepores: habitat: St. Thomas; Th. Bland! R. Swift! J. Knox! St. Thomas; St. Croix; St. Martin; Krebs!

Naticacea.

Natica Adanson.

- N. canrena* L. station: On sand 4 to 6 feet under water at a certain season of the year (Nov. De?) habitat: West-Indies Mörch 1—134! Brasils; Cuba; Martinique; St. Lucy; Guadeloupe; Florida; d'Orb.! Jamaica; Turks-Island; Bermuda; Ad.! St. Barth; St. Thomas; St. Johns; Tortola; Anguilla; St. Martin; Krebs! Guadeloupe; Beau!
- N. fuscata* Chem. (*N. brunnea* Link. *N. mamilaris*) station: In the clear sand 4 to 5 feet from the shore i Great Northside Bay at St. Thomas; habitat: West-Indies; Mörch pag. 132! Cuba; St. Lucy; d'Orb.! Jamaica; Navy-Bay; Carthagera; Ad.! Guadeloupe; Beau! Trinidad; Jamaica; St. Thomas; St. Martin; Porto-Plata; Carthagera; Krebs!
- N. Lacernula* d'Orb. station: — ? — ; habitat: Martinique; Candé! Cuba; de la Sagra! remark: see Moll. de Cuba 2. 35.

- N. lactea* Ph. station: lives in sand; habitat: St. Thomas; Krebs! remark: the epidermis is gray.
- N. marochiensis* ? Lmk. station: — ? — ; habitat Guadeloupe; Beau!
- N. Menkeana* Pfr. station: -- ? — ; habitat: Puer-torico; Zeitsch. 1851 pg. 46! West-Indies; Krebs!
- N. nitida* Donovan (*N. mamilla* L. *Naticina lactea* Guild.) station: — ? — ; habitat: West-Indies; Mörch pag. 132! St. Thomas; Tortola; Barbados; Krebs! Cuba; Martinique; Guadeloupe; St. Lucy; d'Orb. 2. 32! St. Thomas; Tortola; Jamaica; Cuba; Carthagen; Ad.!
- N. ochrostoma* Recluz. station: — ? — ; habitat: West-Indies? remark: Journal de Coch. 1850 pg. 391.
- N. Pfeiferi* Ph. station: — ? — ; habitat: St. Thomas. ?; Krebs! remark: Very like *Natica lactea* Ph. but the epidermis is yellowish-brown; perhaps only a variety.
- N. proxima* Ad. station: — ? — ; habitat: Jamaica; Tortola; Arecibo; St. Croix; C. B. Ad. St. Thomas; St. Johns; Barbados; Porto-Plata; Krebs! remark: By us considered young *Nat. canrena* L.
- N. pulchella* Pfr. (*N. jamaicensis* Ad. *N. Sagriana* d'Orb.) station: — ? — ; habitat: Jamaica; Turks-Island; Ad. St. Thomas; Anguilla; Trinidad; Krebs! Cuba; Sagra! d'Orb. 2. 34. of Moll. de Cuba.
- N. pusilla* Say. station: — ? — ; habitat: Guadeloupe; Beau!
- N. rugosa* Chem. (*N. cancellata* Gml.) station: — ? — ; habitat: Guadeloupe; Beau! West-Indies; Mörch pag. 133! St. Johns; Ad. St. Johns; Porto-Plata; St. Croix; Trinidad; Krebs!
- N. sulcata* Born. (*Natica rugosa* Gml. *N. costata* Mke.)

station: — ? — ; habitat: West-Indies; Mörch pag. 133! Cuba; Auber! remark: we believe it is synonyme with *N. rugosa* Chem.

Sigaretus Lamarck.

- S. depressus* Philippi (*S. antillarum* Recluz) station: — ? — ; habitat: West-Indies; Mörch pag. 131! Cuba; St. Lucy; d'Orb.! Guadeloupe; Beau! St. Thomas; Tortola; Krebs!
- S. haliotideus*. station: — ? — ; habitat: Jamaica; C. B. Adams!
- S. maculatus* Say. station: — ? — ; habitat: Jamaica & St. Johns; C. B. Adams!
- S. zonatus* d'Orb. (*S. martinianus* Ph.) station: — ? — ; habitat: West-Indies; Mörch pg. 131! Guadeloupe; Hotessier! Beau! St. Thomas; St. Martin; Krebs!

Naricaceæ.

Narica Recluz.

- N. granulosa* Recluz. station: — ? — ; habitat: St. Thomas; Krebs!
- N. lamellosa* d'Orb. station: — ? — ; habitat: Cuba; Martinique; Guadeloupe; Candé! Guadeloupe; Beau! St. Johns; St. Barth; Anguilla; Porto-Plata; Krebs!
- N. striata* d'Orb. station: — ? — ; habitat: Cuba; Auber!
- N. sulcata* d'Orb. station: — ? — ; habitat: Havana; Auber! St. Lucy; Jamaica; de Candé!

Xenophorea.

Xenophora Fischer von Waldheim.

- X. trochiformis* Born. (*Onustus* Humph. *Phorus* Mtf.)

station: in 8 to 10 feet water on sand; habitat: St. Thomas; Mörch pg. 49! Martinique; Candé! Guadeloupe; Beau! St. Thomas; Tortola; Porto-Plata; St. Croix; St. Johns; Krebs!

X. *caribæa* Petit. station: — ? — ; habitat: Marie-Galante; Beau!

Calyptraeacea.

Crucibulum Schumacher.

(Calypeopsis Les. Dispotea Say).

C. *auriculatum* Ch. (*P. auriculata* Gml.) station: — ? — ; habitat: West-Indies; Mörch pag. 146! St. Thomas; St. Croix; Anguilla; St. Martin; St. Barth; Guadeloupe; Barbados; Puertorico; Porto-Plata; Krebs!

C. *radiata* Brod. station: — ? — ; habitat: Caracas; Mörch pag. 146! (La Guyra?) St. Thomas; Tortola; St. Croix; Krebs!

Mitrularia Schumacher.

M. *esquestris* Lmk. station: — ? — ; habitat: Africa; l'île de Prince; Cuba; Florida; d'Orb.! East-Indies; Mörch! remark: Is confounded by d'Orbigny and others with *Crucibulum auriculatum* Chem.

Trochita Schumacher. (Infundibulum d'Orb.)

T. *candæana* d'Orb. station: — ? — ; habitat: St. Domingo; Jamaica; St. Thomas; Martinique; de Candé! Cuba; Sagra! Porto-Plata; St. Thomas; St. Martin; Anguilla; Krebs! St. Thomas; R. Swift!

Crepidula Lamarck.

C. *aculeata* Chem. station: On dead Madrepores laying

in 2 feet water; habitat: West-Indies; Mörch pg. 147! Cuba; Patagonia; Brasils; Guadeloupe; d'Orb.! Guadeloupe; Beau! Jamaica; Ad.! St. Thomas; St. Johns; St. Croix; Krebs!

C. fornicata Lmk. station: — ? — ; habitat: Barbadoes; Lamarek!

C. protea d'Orb. station: Found on *Strombus pugillis*; habitat: Patagonia; Brasils; Cuba; d'Orbig.! Guadeloupe; Beau! St. Thomas; Porto-Plata; Krebs!

C. Riisei Dnk. station: — ? — ; habitat: St. Juan de Puertorico; Riise!

C. unguiformis Lmk. (*C. plana* Say acc. to C. B. Ad.!) station: — ? — ; habitat: Guadeloupe; Beau! Jamaica & Carthageria; C. B. Ad.! remark: We think the *C. fornicata*, *C. protea* and *C. auguiformis* are synonyms; they appear under very different shape and colours.

Capulus D. Montfort (*Hipponyx*).

C. antiquatus L. (*P. mitrula* Gm. — *P. cernua* Spr.) station: — ? — ; habitat: West-Indies; Mörch! Guadeloupe; Beau! Cuba; de la Sagra! Florida; d'Orb.! Martinique; Guadeloupe; St. Lucy; Candé! Hotessier!

C. incurvus Gm. (*P. militaris* L., Dw., *P. intorta* Lmk.) station: — ? — ; habitat: West-Indies? Mörch pg. 145! Cuba; Guadeloupe; d'Orb. 2. 187! Guadeloupe; Beau! St. Thomas; St. Croix; St. Martin; St. Barth; Guadeloupe; Barbadoes; Porto-Plata; Krebs!

C. pilosa Desh. station: — ? — ; habitat: Guadeloupe; Beau!

C. subrufa Mart. station: — ? — ; habitat: West-Indies; Mörch pg. 145! Cuba; Sagra! Guadeloupe; Beau!

Porto-Plata; Guadeloupe; Krebs! remark: Moll. de Cuba; d'Orb. p. 2. pg. 187 pl. 24. fig. 24—25.

C. trigona Gml. — ? — ; habitat: West-Indies; Mörch pag. 45!

Ringiculacea.

Cinulia Gray.

C. semistriata d'Orb. station: — ? — ; habitat: Jamaica; Candé! remark: Described in d'Orbigny's Moll. de Cuba 2. 103. pl. 31— $\frac{17}{19}$.

Pyramidellacea.

Pyramidella Lamarck.

P. dolabrata L. station: — ? — ; habitat: West-Indies; Mörch pg. 45! Cuba; Sagra! Guadeloupe & St. Lucy; Hotessier! Guadeloupe; Beau! St. Thomas; St. Jan; Crabisland; Portoplata; St. Martin; Anguilla; Krebs!
P. terebellum Lmk. station: — ? — ; habitat: Jamaica; E. Chitty! St. Johns; Jamaica; Bermuda; Ad.!

Odontostoma Fleming.

O. depressa d'Orb. station: — ? — ; habitat: Cuba; Sagra! remark: Described in d'Orb. Moll. de Cuba 1—238.
O. globosa d'Orb. station: — ? — ; habitat: Cuba; d'Orb.! remark: Described in d'Orb. Moll. de Cuba 1—239.
O. pusilla (*Pyramidella*) Pfef. station: — ? — ; habitat: Cuba; Zeitschrift 1851. pg. 88.

Monostygma Gray.

1 spec-indet. St. Thomas; Krebs!

Turbonilla Risso (Chemnitzia).

- T. babylonia** Ad. station: — ? — ; habitat: Jamaica; Ad.!
- T. cancellata** d'Orb. station: — ? — ; habitat: Cuba; Sagra! remark: Described by d'Orb. in Moll. de Cuba 1—225. pl. 17. fig. 1—3.
- T. dubia** d'Orb. station: — ? — ; habitat: Jamaica; Guadeloupe; Martinique; St. Thomas; Cuba; Rio Janeiro; d'Orb.! Guadeloupe; Beau! remark: Described by d'Orb. in „Moll. de Cuba“ 1. 226. pl. 17 fig. 4—6.
- T. elegans** d'Orb. station: — ? — ; habitat: Guadeloupe; de Candé! Beau! remark: Described by d'Orb. in „Moll. de Cuba“ 1—223. pl. 16. fig. 25. 27.
- T. exilis** Ad. station: — ? — ; habitat: Jamaica; Ad.!
- T. flavocincta** Ad. station: — ? — ; habitat: Jamaica; Ad.!
- T. laevigata** d'Orb. station: — ? — ; habitat: St. Thomas; de Candé! remark: Described by d'Orb. in „moll. de Cuba“ 1—227 pl. 17 fig. 7—9.
- T. laevis** Ad. station: — ? — ; habitat: Jamaica; Ad.! St. Thomas; Krebs!
- T. latior** Ad. station: — ? — ; habitat: Jamaica; Ad.! St. Thomas; Krebs!
- T. modesta** d'Orb. station: — ? — ; habitat: Jamaica; Candé! remark: Described by d'Orb. „moll. de Cuba“ 1—222 pl. 16 fig. 22. 24.
- T. multicostata** Ad. station: — ? — ; habitat: Jamaica; Ad.!
- T. obeliscus** Ad. station: — ? — ; habitat: Jamaica; Ad.!
- T. ornata** d'Orb. station: — ? — ; habitat: Martinique; Guadeloupe; Jamaica; Candé! Jamaica; Ad.! Gua-

deloupe; Beau! remark: Described in d'Orb. „moll. de Cuba“ 1—221 pl. 16 fig. 18. 21.

T. pulchella d'Orb. station: — ? — ; habitat: Martinique; St. Thomas; Guadeloupe; Candé! Cuba; Sagra! Guadeloupe; Beau! remark: Described by d'Orbig. in „Moll. de Cuba“ 1. 220 pl. 16 fig. 14—17.

T. punctata Ad. station: — ? — ; habitat: Jamaica; Ad.!

T. pupoides d'Orb. station: — ? — ; habitat: Cuba; St. Thomas; d'Orb.! Guadeloupe; Beau! remark: Described by d'Orb. in „Moll. de Cuba“ 1. 224. pl. 16. fig. 32. 36.

T. pusilla Ad. station: — ? — ; habitat: Jamaica; Ad.! St. Thomas; Krebs!

T. reticulata Ad. station: — ? — ; habitat: Jamaica; Ad.! St. Thomas; Krebs!

T. simplex d'Orb. station: — ? — ; habitat: Jamaica; Candé! remark: Described by d'Orb. in „Moll. de Cuba“ 1—224 pl. 16. 28—31.

T. substriata Ad. station: — ? — ; habitat: Jamaica; Ad.! St. Thomas; Krebs!

T. subulata Ad. station: — ? — ; habitat: Jamaica; Ad.! St. Thomas; Knox!

T. turris d'Orb. station: — ? — ; habitat: St. Thomas; d'Orb.! R. Swift! remark: Described by d'Orb. in „moll. de Cuba“ 1—219. pl. 16. fig. 10—13.

Eulimacea.

Eulima Risso.

E. affinis Ad. station: — ? — ; habitat: Jamaica; Ad.!

E. arcuata Ad. station: — ? — ; habitat: Jamaica; Ad.!

- E. bifasciata* d'Orb. station: — ? — ; habitat: Guadeloupe; St. Thomas; de Candé! Guadeloupe; Beau! St. Thomas; R. Swift! remark: Described in „Moll. de Cuba“ 1. 216. pl. 16 fig. 1—3; is this not *E. fulvo-cincta* Ad.?
- E. conica* Ad. station: — ? — ; habitat: Jamaica; Ad.!
- E. fulvo-cincta* Ad. station: — ? — ; habitat: Jamaica; Ad. St. Thomas; Knox!
- E. gracilis* Ad. station: — ? — ; habitat: Jamaica; Ad.!
- E. incerta* d'Orb. station: — ? — ; habitat: Jamaica; de Candé! remark: Described in „Moll. de Cuba“ 1—218 pl. 16 fig. 7—9.
- E. jamaicensis* Ad. station: — ? — ; habitat: Jamaica; St. Thomas; St. Johns; Ad. St. Martin; St. Barth; St. Thomas; St. Johns; Krebs!
- E. subearinata* d'Orb. station: — ? — ; habitat: Guadeloupe; de Candé! Beau! remark: Described in „Moll. de Cuba“ 1—217 pl. 16 fig. 4—6.

Vermetacea.

Vermetus Adanson.

- V. corrodeus* d'Orb. station: on rocks close to the surface of the sea; habitat: Cuba; Martinique; d'Orb.!
- V. decussata* Gml. station: — ? — ; habitat: West-Indies; Mörch pg. 53.
- V. irregularis* d'Orb. station: — ? — ; habitat: Cuba; Martinique; de Candé!
- V. lumbricalis* L. (*V. Knorri* D.) station: on rocks in the surface of the sea; habitat: Cuba; Auber! St. Thomas; Krebs!

Siliquaria Brugière.

- S. muricata* Born. (*S. echinata* Gml.) station: found on sandy beaches; habitat: West-Indies; Mörch pg. 53!
 1 spec. ind. station: between small shells picked on the beach; habitat: Porto-Plata; Krebs!

Caecacea.

- Caecum* Fleming. 3 spec. ind.: St. Thomas; Krebs!

Siphonariacea.

Siphonaria Sowerby.

- S. lineata* d'Orb. station: — ? — ; habitat: Cuba; Sagra! Guadeloupe; Beau!
S. picta d'Orb. station: — ? — ; habitat: Cuba; Auber! Rio Janeiro; d'Orb.!

Acmaeacea.

Acmaea Eschholtz.

- A. fungus* Meush. (*P. fungoides* Bolt., *P. octoradiata* var. Gm.) station: — ? — ; habitat: West-Indies; Mörch!
A. Hamillei Fisch. station: — ? — ; habitat: Guadeloupe; Beau!
A. leucopleura Gm. (*C. extensorium*? Sowb. *C. rugosa* Less) station: — ? — ; habitat: West-Indies; Mörch! Cuba; St. Lucy; Martinique; d'Orb.! Jamaica; St. Thomas; Ad.!
A. melaleuca Gm. station: — ? — ; habitat: West-Indies; Mörch!
A. melanosticta Gm. (*L. antillarum* Sow. Pat. *palescens* Ph.) station: — ? — ; habitat: West-Indies; Mörch! St. Thomas; Krebs!
A. pustula Helb. (*P. punctulata* Gm., *P. puncturata* . . .)

station: on flat pieces of Madrepores close to the shore;
 habitat: West-Indies; Mörch pg. 144! Turks-Island;
 St. Johns; Jamaica; St. Thomas; Ad.! Guadeloupe; Beau!
 Barbadoes; Trinidad; St. Martin; Anguilla; St. Barth;
 St. Johns; St. Thomas; St. Croix; Puertoricco; Porto-
 Plata; Krebs!

Neritacea.

Nerita Linné.

- N. antillarum* Gm. station: —? —; habitat: West-Indies; d'Orb.! St. Thomas? Carthagenae? St. Martha? Krebs! remark: Yellow columella, one tooth, spire flat; we are in doubt if those we have found, at the three places noted, are *N. ant.* Gm.
- N. nigerrima* Chem. station: on wood along the shore in creeks with smooth water; habitat: West-Indies; Mörch pg. 168! Porto-Plata? Krebs!
- N. peloronta* L. (*Pila* Kl. *N. erythron.* Recluz.) station: On rocks from 1 foot below low water and until 2 to 3 feet above highwater; habitat: West-Indies; Mörch pg. 169! Jamaica; Havana; Bermuda; Carthagenae; Ad.! St. Thomas; St. Johns; Guadeloupe; Barbadoes; St. Martin; Carthagenae; Krebs! Cuba; Sagra! Auber! Candé! Martinique; d'Orb.!
- N. præcognita* C. B. Ad.! station: On rocks from 1 foot below low-water until 2 to 3 feet above highwater; habitat: Jamaica; Ad.! Guadeloupe; Beau! St. Thomas; St. Johns; St. Croix; Carthagenae; Krebs! remark: A doubtful species.
- N. varia* Meuschen. (*Theliostyla* Mörch, *N. tessellata* Gml.) station: On rocks where the waves washe over; West-Indies; Mörch pg. 168! Cuba; Sagra! Auber! Martini-

que; Candé! Guadeloupe; Beau! St. Thomas; St. Johns; Sta. Martha; Carthagera; Krebs!

- N. variegata* Chem. (*N. versicolor* Gml., *Neritina striata* Chem. acc. to d'Orb.) station: On rocks where the waves wash over; habitat: Cuba; Martinique; d'Orb.! Florida; Alphen! Ad.! West-Indies; Mörch pg. 169! Guadeloupe; Beau! Duchassain! St. Thomas; St. Johns; Porto-Plata; Carthagera; Krebs! Turks-Island; Havana; Jamaica; Arecibo; Carthagera; Ad.!

Neritina Lamarck.

- N. brasiliana* Recluz. (*N. virginea* Lmk. acc. to Beau.) station: — ? — ; habitat: Guadeloupe; Beau!
- N. chlorina* Link. (*N. meleagris* Lmk.) station: In sea water in small ponds in the rocks along the shore and on poles where the sea is calm and smooth; habitat: West-Indies; Mörch pg. 167! Cuba; Guadeloupe; d'Orb.! Guadeloupe; Beau! St. Domingo; Lmk.! Jamaica; Carthagera; Ad.! St. Thomas; Carthagera; Krebs!
- N. Domingensis* Lmk. station: — ? — ; habitat: St. Domingo; Lmk.!
- N. jamaicensis* Ad. station: — ? — ; habitat: Jamaica; E. Chitty!
- N. microstoma* d'Orb. station: — ? — ; habitat: Martinique; de Candé!
- N. ornata* Ad. station: — ? — ; habitat: Jamaica; Ad!
- N. pulchella* Gray. station: — ? — ; West-Indies; Mörch pg. 167!
- N. punctulata* Lmk. (*N. cassicula* Sow. acc. to Beau! *N. fluviatilis*? acc. to C. B. Adams) station: — ? — ; habitat: Martinique; Mörch! Guadeloupe; Beau! Duchassing! d'Orb.! Jamaica! Ad.!

- N. pupa* L. station: In brackish water even a good distance from the sea; habitat: Guadeloupe; Beau! Duchassain! Cuba; Sagra! Auber! West-Indies; Mörch pg. 167! St. Croix; A. H. Riise! Arecibo; Jamaica; Ad.! Porto-Plata; Krebs!
- N. pusilla* Ad. station: — ? — ; habitat: Jamaica; Adams!
- N. pygmæa* Ad. station: — ? — ; habitat: Jamaica; Ad.!
- N. succinea* Recluz. station: — ? — ; habitat: Guadeloupe; Beau! Crabisland; A. H. Riise! St. Thomas; H. Haagensen! St. Johns (Fishbay); Krebs!
- N. tenebrosa* Ad. (*N. fluviatilis*? according to C. B. Ad.) station: — ? — ; habitat: Jamaica; Ad.!
- N. trabalis* Meusch. station: — ? — ; habitat: West-Indies; Mörch pg. 167!
- N. tristis* d'Orb. (syn. *N. pupa* L. accord. to Jay's Catal.) station: — ? — ; habitat: Guadeloupe; Beau! Cuba; Sagra! Jamaica; Ad.! St. Croix; A. H. Riise!
- N. turriculata* Mke. (*N. virginea* Sow.) station: — ? — ; West-Indies; Mörch pg. 166!
- N. virginea* L. station: In brackish water, sometime in a considerable distance from the shore (at North-Star. St. Croix) in salt water in the wester-most part of the harbour of St. Thomas; habitat: West-Indies; Mörch pg. 166! Florida; d'Orbig.! Cuba; Sagra! Auber! Martinique; Guadeloupe; Candé! Hotessier! Jamaica; St. Domingo; Arecibo; Ad.! St. Thomas; St. Croix; Krebs!
- N. viridis* L. station: In salt and brackish water; on poles in the western part of the harbour of St. Thomas, where the sea is smooth; habitat: West-Indies; Mörch! Martinique; St. Lucy; Guadeloupe; Candé! Cuba; d'Orb.!

Guadeloupe; Beau! Jamaica; St. Johns; Arecibo; Bermuda; Ad.! St. Thomas; St. Croix; St. Johns; St. Barth; Anguilla; St. Martin; Trinidad; Porto-Plata; Krebs!

Trochacea.

Phasianella Lamarck.

- P. affinis* Ad. station: Below Madrepores in 1 or 2 feet water; habitat: St. Thomas; St. Johns; Jamaica; Ad.! St. Thomas; St. Barth; Krebs!
- P. brevis* Ad. station: — ? — ; habitat: Jamaica; St. Thomas; Ad.!
- P. brevis* d'Orb. station: — ? — ; habitat: Cuba; Sagra! Martinique; Candé! remark: Described in „moll. de Cuba“ 2—79 pl. 20 fig. 19—21.
- P. concinna* Ad. station: — ? — ; Jamaica; Ad.! St. Thomas; St. Martin; St. Barth; Krebs! St. Thomas; Crab-island; Coro; A. H. Riise!
- P. concolor* Ad. station: — ? — ; habitat: Jamaica; Ad.!
- P. tessellata* Ad. (*P. minuta* Ant., *P. lipeata* Dkr. according to specimens in the collection of A. H. Riise) station: — ? — ; habitat: Jamaica; Ad.! Guadeloupe; Beau! Hotessier! St. Johns; Anguilla; Trinidad; Krebs!
- P. umbilicata* d'Orb. station: dredged May 1860 from 10 to 12 fathoms water, bluish sandy mud, 2 specimens; habitat: Martinique; Guadeloupe; Jamaica; Cuba; Candé! Sagra! Guadeloupe; Beau! St. Thomas; Krebs! remark: Described in „moll. de Cuba“ 2—78 pl. 19 fig. 35—37.

Turbo Linné.

- T. Cailletii* Fischer & Bern. station: Taken in a fish; habitat: Guadeloupe; Beau!

- T. castaneus* Chem. (*T. hippocastanum* Lmk.) station: Between *Zostera* (*maritima*?) on 2 to 3 feet water in the western part of the harbour of St. Thomas; habitat: West-Indies; Mörch pg. 162! Margarita; R. Swift! St. Thomas; Tortola; St. Martin; Guadeloupe; Trinidad; Porto - Plata; Krebs! Martinique; Jamaica; Cuba; Florida; d'Orb.!
- T. crenulatus* Chem. station: — ? — ; habitat: West-Indies; Mörch pg. 162! Guadeloupe; Beau!
- T. pica* L. station: on rocks where the waves are washing over; habitat: West-Indies; Mörch pg. 154! Guadeloupe; Beau! Cuba; Martinique; Guadeloupe; Sagra! Auber! Candé! Jamaica; Carthagenia; Ad.! St. Thomas; Bermuda; St. Croix; St. Johns; Porto-Plata; Krebs! remark: common food between the lower classes.

Liotia Gray.

- L. cruentata* Mühlf. (*Delphinula radiata* Eien.) station: Under pieces of Madrepores in 2 to 3 feet water, where the sea is pretty rough; habitat: West-Indies; Mörch! Guadeloupe; Beau! Duchassain! Jamaica; Ad.! St. Thomas; Anguilla; Krebs!

Delphinula Lamk.

- D. tuberculosa* d'Orb. (*Trochus Schrammii* Fischer accor. to Beau) station: — ? — ; habitat: Jamaica; St. Thomas; Candé! Guadeloupe; Beau! remark: Described in „moll. de Cuba“ 2—66.

Vitrinella Adams (*Skoenia* acc. to Beau).

- V. Adamsii* Fischer. station: found dead in the fine white sand of the beach; habitat: Guadeloupe; Beau!

- V. *Beauui* Fischer. station: — ? — ; habitat: Guadeloupe; Beau!
- V. *helicoidea* Ad. station: in the fine white sand of the beach; habitat: Jamaica; Ad.! St. Thomas; Krebs!
- V. *hyalina* Ad. station: — ? — ; habitat: Jamaica; Ad.!
- V. *interrupta* Ad. station: — ? — ; habitat: Jamaica; Ad.!
- V. *megastoma* Ad. station: — ? — ; habitat: Jamaica; Ad.! Guadeloupe; Beau!
- V. *Petitii* Fisch. station: — ? — ; habitat: Guadeloupe; Beau!
- V. *Schrammii* Fisch. station: — ? — ; habitat: Jamaica; Ad.!
- V. *tincta* Ad. station: — ? — ; habitat: Jamaica; Ad.!

Calcar D. Montfort (Stella Kl.)

- C. *caelatus* Chem. station: on coarse gravel in 4 to 5 feet water; habitat: West-Indies; Mörch pg. 160! Cuba; Sagra! Auber! Martinique; Guadeloupe; St. Lucy; Candé! Guadeloupe; Beau! Bermuda; St. Johns; St. Thomas; Jamaica; Ad.! St. Thomas; Tortola; Krebs!
- C. *inermis* Gml. station: — ? — ; habitat: Cuba; Sagra! Martinique; St. Croix; Candé! St. Thomas; Kr.! remark: This is certainly nothing else than the young Calcar Tuber L.; we have compared it with the drawing of Chemnitz.
- C. *ramosus* Meusch. (*T. imbricatus* Gml.) station: — ? — ; habitat: West-Indies; Mörch pg. 160! Curaçao; A. H. Riise! St. Thomas; St. Croix; Tortola; Guadeloupe; Kr.!
- C. *Spenglerianus* Ch. (*Tuber aculeatus* Gml.) station:

Said to live in very deep water; habitat: Ind. orient; Mörch pg. 160! Curaçao; R. Swift! Guadeloupe; Beau! Jamaica; Ad.! Porto-Plata; St. Martin; Krebs! remark: perfect specimens are very rare.

- C. Tuber L. station: On rocks near the shore in 1 to 8 feet water; habitat: West-Indies; Mörch pg. 160! Guadeloupe; Beau! Jamaica; Arecibo; Bermuda; Ad.! Trinidad; St. Martin; St. Johns; Kr.!

Astraliu Link. (Imperator Mtf.).

- A. brevispinum Lmk. station: — ? — ; habitat: Cuba; Martinique; Candé!
- A. calcar L. (T. inermis Gml.? T. spinulosa Lmk. Ki.?) station: — ? — ; habitat: West-Indies; Mörch pag. 160! Cuba; Sagra! Martinique; Candé! Guadeloupe; Beau! remark: see A. phoebia Bolt. & Calcar inermis Gml.
- A. costulatum (Tr.) Lmk. station: — ? — ; habitat: West-Indies; Lamarek! St. Croix; Kr.!
- A. phoebia Bolt. (A. deplanatum Link. P. longispina Lmk.) station: In Marsch 58, found on sand about $\frac{1}{2}$ foot below the level of the sea; habitat: West-Indies; Mörch pg. 160! St. Thomas; Candé! Guadeloupe; Beau! St. Thomas; St. Croix; Tortola; Carthagena; Kr.! remark: We have no doubt that the A. calcar L. and A. phoebia B. are synonymes; 1721—22 in Chemnitz are certainly young specimens of A. ph.
- A. radians Lmk. station: — ? — ; habitat: Near Guadeloupe; Lmk.! St. Croix; Krebs!
- A. rhodostomum Lmk. station: on a rock about the level of the sea. June 52; habitat: St. Martha-New-Granada; Kr.! remark: those found at St. Martha

are the only specimens seen by the collectors at St. Thomas.

Globulus Schum. (Rotella Lmk. d'Orb.)

- G. *anomalis* d'Orb. station: — ? — ; habitat: St. Thomas; Candé! remark: Described in „Moll. de Cuba“ 2—64.
- G. *carinatus* d'Orb. station: — ? — ; habitat: St. Thomas; Candé! remark: Described in „moll. de Cuba“ 2—62.
- G. *diaphanus* d'Orb. station: — ? — ; habitat: St. Thomas; Candé! remark: Described in „moll. de Cuba“ 2—62.
- G. *semistriatus* d'Orb. station: — ? — ; habitat: Cuba; Sagra! remark: Described in „moll. de Cuba“ 2—61.
- G. *strictus* d'Orb. station: — ? — ; habitat: Jamaica Candé!

Trochus Linné.

- T. *asperulus* Lamarck station: — ? — habitat: St. Domingo; Lmk.!
- T. *aster* Ph. station: — ? — ; habitat: Guadeloupe; Beau!
- T. *ater* Less. station: — ? — ; habitat: West-Indies; Mörch pg. 154!
- T. *concavus* Gml. station: — ? — ; habitat: Martinique; d'Orb.!
- T. *crenulatus* Wood. station: — ? — ; habitat: St. Johns; St. Thomas; Jamaica; Ad.! Carthagera; Kr.!
- T. *cubanus* Ph. station: — ? — ; habitat: Guadeloupe; Beau!

- T. excavatus* Lmk. (*Tr. umbilicaris* Ch.) station: On rocks near the shore at the level of the sea; habitat: West-Indies; Mörch pg. 153! Martinique; Guadeloupe; Cande! Hotessier! Guadeloupe; Beau! Jamaica; Ad.! St. Thomas; St. Croix; St. Johns; Crabisland; Porto-Plata; Krebs!
- T. fasciatus* Born. (*C. lun. laevis* Chem., *Tb. dentatus* Gm., *Tb. carneolus* Lmk., *Tr. livido-maculatus* Ad.) station: Under pieces of Madrepores in 1 foot water; habitat: Jamaica; Ad.! West-Indies; Mörch! Martinique; Guadeloupe; St. Lucy; Hotessier! Candé! Cuba; Sagra! St. Thomas; St. Johns; St. Martin; Anguilla; Trinidad; Barbadoes; Porto-Plata; Krebs! Guadeloupe; Beau!
- T. flammulatus* Lmk. station: — ? — ; habitat: St. Domingo; Lmk.
- T. heliacus* Ph. station: — ? — ; habitat: Guadeloupe; Beau!
- T. Hotessierianus* d'Orb. (*Tr. occultus* Phil., *Tr. nasaviensis* Ch., *Tr. interrupta striatus* C. B. Ad.) station: In small ponds of seawater in the rocks, left when it is low-water; habitat: West-Indies; Mörch pg. 153! Florida; d'Orb.! Guadeloupe; St. Lucy; Candé! Hotessier! Cuba; Auber; Sagra! St. Thomas; St. Martin; Porto-Plata; Kr.! Guadeloupe; Beau!
- T. javanicus* Lmk. station: Said to be taken on corals in 3 fathoms water; habitat: Jamaica; Ad.! Carupano; Kr.!
- T. indusii* Chem. (*Omphalius* Ph., *Tr. carneus* Gml., *Tr. canaliculatus* d'Orb., *Tr. scalaris* Anton, *Tr. crenulatus* Bolt.); station: On pieces of madrepores in 1 or 2 feet of water; habitat: West-Indies; Mörch pg. 153! Cuba; Auber! All the Virgin-Islands; Krebs!

- T. jujubinus* Gml. (*Tr. lunatus* Bolt.) station: On pieces of Madreporos in 2 feet water; habitat: West-Indies; Mörch pg. 157! Jamaica; St. Thomas; Carthageria; Ad.! All the Virgin Islands; Kr.!
- T. microstomus* d'Orb. (*Tr. bidentatus* Mk.) station: — ? — ; habitat: West-Indies; Mörch pg. 154!
- T. pulcher* C. B. Adams. station: On pieces of madreporos in a few feet water and dredged from 6 and from 12 feet water on bluish sandy mud; habitat: Jamaica; Ad.! St. Thomas; St. Croix; Porto-Plata; Krebs! remark: Described in C. B. Adams „Contributions“ page 69— ; should this species not be Woods *Tr. interruptus*? We have specimens of *pulcher* with 9 whorls.
- T. Schrammii* Fisch. see *Delph. tuberculosa* d'Orb.

Stomatia Helbling.

- S. picta* d'Orb. station: On pieces of Madreporos in a few feet water; habitat: Cuba; d'Orbig.! St. Martin; Guadeloupe; Beau! St. Thomas; St. Barth; Pto.-Plata; Anguilla; Krebs! remark: Described in „Moll. de Cuba“ 2—184 pl. 24 fig. 19, 21.

Stomatella Lamarck.

- S. coccinea* A. Ad. station: — ? — ; habitat: St. Thomas; Mörch pg. 152! St. Thomas; St. Croix; Kr.!

Pleurotomaria Defrance.

- P. Gouayana* Fisch. & Bern. station: In very deep water; habitat: Marie Galante; Beau!

Fissurellacea.

Emarginula Lamarck.

1. Emarginula.

- E. notata* L. (*P. manipula* Humphr) station: — ? — ; habitat: West-Indies; Mörch pg. 148!

- E. Rollandii* Fisch. station: — ? —; habitat: Guadeloupe; Beau!

2. Subemarginula.

- E. depressa* Blainw. station: — ? —; habitat: Guadeloupe; Beau! remark: C. B. Adams gave us a specimen of *E. octoradiata* Gml. from St. Thomas with this name.
- E. octoradiata* Gml. (*E. tricarinata* Sow., *P. laqueara* Gray, *E. Listeri* Ant.) station: Common on rocks between high and low-water mark; habitat: West-Indies; Mörch pg. 149! Jamaica; St. Johns; Ad.! St. Thomas; St. Johns; St. Barth; St. Martin; Anguilla; Barbadoes; Porto-Plata; Kr.!

Fissurella Bruguière.

- F. alternata* Say. station: — ? —; habitat: Guadeloupe; Beau! Jamaica; Carthagera; Ad.!
- F. Antillarum* d'Orb. station: — ? —; habitat: Florida; Cuba; d'Orb.! remark: Described in „moll. de Cuba“ 2—198 pl. 24 fig. 40—42.
- F. atricapilla* Dw. (*F. viridula* Lmk.) station: — ? —; habitat: West-Indies; Mörch pg. 149!
- F. barbadensis* Gml. (*P. porphyrozonias* Gml., *P. angusta* Gml., *F. radiata* Lmk.) station: On rocks from high-water-mark until 4 to 5 feet below the level of the sea; habitat: Florida; Cuba; St. Lucy; Martinique; d'Orb.! West-Indies; Mörch pg. 149! Jamaica; Ad.! Guadeloupe; Beau! St. Thomas; St. Croix; Crabisland; St. Johns; St. Martin; Anguilla; Puertorico; Pto.-Plata; Kr.!
- F. cancellata* Sal. (*F. Sowerbyi* Guild.) station: — ? —; habitat: Guadeloupe; Beau! St. Thomas; St. Martin; St. Barth; Kr.!

- F. sayennensis* Lmk. station: — ? — ; habitat: Guadeloupe; Beau!
- F. Dysonii* Reeve. station: — ? — ; habitat: St. Johns; Ad.! Guadeloupe; Beau!
- F. elongata* Reeve. station: — ? — ; habitat: Guadeloupe; Beau! St. Thomas; St. Martin; St. Barth; Kr.!
- F. fascicularis* Lmk. (*Clypidella* Sow.) station: — ? — ; habitat: Jamaica; Ad.! Arecibo; Newton! West-Indies; Mörch pg. 150!
- F. gemmulata* Reeve. (*F. minuta* Sowb., *F. elongata* Ad.) station: On pieces of Madreporas in a few feet water; habitat: Guadeloupe; Beau! Jamaica; Ad.! St. Thomas; St. Johns; St. Croix; Crab Island; St. Barth; St. Martin; Anguilla; Kr.! remark: We have in a note by C. B. Adams seen that he considered *F. gemmulata* Reeve synonyme with *F. elongata* Ad.
- F. graeca* L. (*F. Listeri* d'Orb.) station: — ? — ; habitat: West-Indies; Mörch pg. 149! St. Thomas; St. Johns; St. Martin; Porto-Plata; Kr.! St. Lucey; Martinique; d'Orb.! St. Thomas; St. Johns; Jamaica; Arecibo; Carthagena; Ad.! remark: Moll. de Cuba 2—197 pl. 24 fig. 37—39.
- F. larva* Reeve. station: — ? — ; habitat: Guadeloupe; Beau! St. Martin; Guadeloupe; Kr.! St. Johns; Adams!
- F. nimbosea* Lmk. station: At Carthagena on rocks where the waves broke over; habitat: Martinique; Candé! Guadeloupe; Beau! Carthagena, Kr.!
- F. nodosa* Born. (*F. spinosa* Gml. acc. to Mörch, *F. jamaicensis* Gml. acc. to C. B. Ad.) station: On rocks from highwater mark to 3 to 6 feet under the level of the sea, especially between seaweed; habitat: West-

Indies; Mörch pg. 149! Jamaica; Arecibo; Ad.! Cuba; St. Lucy; Martinique; d'Orb.! Guadeloupe; Beau! St. Thomas; St. Croix; Puertoricco; Pto.-Plata; Kr.!

F. pustula L. station: — ? — ; habitat: West-Indies; Mörch pg. 150! Cuba; Florida; d'Orb.! Guadeloupe; Beau! St. Thomas; St. Croix; St. Martin; Crabisland; Anguilla; Puertoricco; Pto.-Plata; Kr.! Guadeloupe; St. Johns; Ad.!

F. radiata Lmk. (*F. balanoides*? Reeve acc. to Beau) station: — ? — ; habitat: Guadeloupe; Beau!

F. reticulata Say. station: — ? — ; habitat: Jamaica; Ad.!

F. rosea Lmk. station: — ? — ; habitat: Jamaica; Arecibo; Ad.! remark: „Compare it with *F. barbadosis*“ says C. B. Adams in a note.

F. viridula Lmk. station: — ? — ; habitat: Guadeloupe; Beau! St. Thomas; Jamaica; Arecibo; Ad.!

Patellacea.

Patella Linné.

P. albicosta C. B. Ad. station: — ? — ; habitat: Jamaica; Ad.! Guadeloupe; Beau!

P. confusa Guild. (*P. notata* Lmk.) station: — ? — ; habitat: Guadeloupe; Beau!

P. cubaniana d'Orb. station: — ? — ; habitat: Cuba; Florida; d'Orb.!

P. cubensis Reeve. station: — ? — ; habitat: Guadeloupe; Beau!

P. elegans Ph. station: — ? — ; habitat: St. Thomas; Kr.!

P. notata Chem. station: — ? — ; habitat: St. Thomas; Jamaica; Arecibo; Ad.!

- P. pulcherrima* Guild. (*P. Candearia* d'Orb.) station: — ? — ; habitat: Guadeloupe; Beau! Cuba; St. Lucy; Martinique; Guadeloupe; d'Orb.!

Chitonacea.

Chiton Linné.

- C. acutiliratus* Reeve. (*Ch. pectinatus* Sow. acc. to Beau) station: — ? — ; habitat: West-Indies; Mörch pg. 139! Guadeloupe; Beau!
- C. asper* Shut. station: — ? — ; habitat: Guadeloupe; Beau!
- C. assimilis* Reeve. station: On rocks near the surface of the sea; habitat: Guadeloupe; Beau! St. Thomas; Krebs!
- C. candisatus* Shut. station: — ? — ; habitat: Guadeloupe; Beau!
- C. costatus* Ad. station: — ? — ; habitat: Jamaica; Ad.!
- C. Erythronatus* C. B. Ad. station: — ? — ; habitat: Jamaica; Ad.! Guadeloupe; Beau!
- C. granulatus* Gml. station: — ? — ; habitat: Cuba; Martinique; d'Orb.!
- C. marmoratus* Ch. (*Lophorus* Poli, *C. marmoreus* Ch., *C. punctatus* L.?) station: On rocks along the shores; habitat: Barbadoes; Mörch pg. 138! Guadeloupe; Beau! Jamaica; Arecibo; Ad.! St. Croix; Carthagera; Kr.!
- C. marmoratus* Reeve. station: — ? — ; habitat: Guadeloupe; Beau!
- C. multicostatus* Ad. station: — ? — ; habitat: Jamaica; Ad.!
- C. papillosus* Ad. station: — ? — ; habitat: Jamaica; Ad.! Guadeloupe; Beau!

- C. piceus* Gml. station: On rocks at the surface of the sea at St. Thomas; Carthagera June 52; Bermuda 25 April 1858; habitat: West-Indies; Mörch pg. 139! Guadeloupe; Beau! St. Johns; Jamaica; Bermuda; Ad.! Carthagera; Bermuda; St. Thomas; Kr.! St. Thomas; R. Swift!
- C. purpurascens* Ad. station: — ? — ; habitat: Jamaica; Ad.! Guadeloupe; Beau!
- C. rugosus* Gray. (*C. squalidus* Ad. acc. to Ad.) station: — ? — ; habitat: Jamaica; Ad.!
- C. salamandra* Spl (*C. piceus* var. Gml., *C. occidentalis* Reeve acc. to Mörch, *C. marmoreus* Reeve syn. with *C. occidentalis* Reeve acc. to Beau) station: On rocks at the shore of Carthagera; habitat: West-Indies; Mörch pg. 139! Carthagera; Kr.!
- C. Schrammii* Shut. station: — ? — ; habitat: Guadeloupe; Beau!
- C. spiculosus* Gray. station: — ? — ; habitat: Guadeloupe; Beau!
- C. squamosus* L. (*C. scaber variegatus* Ch., *Scutigerulus angulatus* Meusch., *C. marmoreus* Reeve) station: on rocks close to the shore; habitat: West-Indies; Mörch pg. 138! Guadeloupe; Beau! St. Thomas; R. Swift! Jamaica; Porto-Cabello; Ad.! Carthagera; Kr.!
- C. squamulosus* C. B. Ad. station: — ? — ; habitat: Guadeloupe; Beau! Jamaica; Ad.!
- C. strigatus* (*Chitonella*) Shut. (*C. laevis* Reeve. acc. to Beau) station: — ? — ; habitat: Guadeloupe; Beau!
- C. undatus* Spgl. (*C. squamosus* Born, Ch., Lmk., *C. tuberculatus* L.?) station: — ? — ; habitat: West-Indies; Mörch pg. 138!
- C. viridis* Spgl., Wood (*C. squamosus denticulatus* Ch.,

C. foveolatus Sowb. accord. to Mörch, *C. excavatus* Gray acc. to Beau) station: — ? — ; habitat: Guadeloupe; Beau! West-Indies; Mörch pg. 138!

Dentalium Linné.

- D. antillarum* d'Orb. station: — ? — ; habitat: St. Thomas; d'Orb.! Guadeloupe; Beau!
- D. disparile* d'Orb. station: — ? — ; habitat: Martinique; d'Orb.! Guadeloupe; Beau!
- D. domingense* d'Orb. station: — ? — ; habitat: St. Domingo; Martinique; St. Thomas; Cuba; d'Orb.!
- D. semistriatum* Guild. station: — ? — ; habitat: Guadeloupe; Beau!

Aplysiacea.

Aplysia Linné.

- A. protea* Rang. station: feeds on species of *Ulvae*, close to the shore in smooth water; habitat: Guadeloupe; Beau! St. Thomas; Carthagen; Kr.! remark: we have observed it swimming but not as George Johnston says in *Einl. in die Conchyphiologie*. Stuttgard 1853 pag. 127 „with the foot upward“, but with the matel extended as mentioned pg. 129; it swims quick when alarmed and takes then a shape very different from when it crawls about.

Lophocercus Krohn.

- 1 species ind. at St. Thomas; Krebs!

Lobiger Krohn.

- L. Sowerbii* Fischer. station: — ? — ; habitat: Guadeloupe; Beau!

Bullacea.**Aplustrum Schumacher.**

- A. physis** L. & d'Orb. (*Hydalina filosa* Sch.) station — ? — ; habitat: West-Indies; Mörch pg. 136! Cuba; St. Lucy; d'Orb.! Guadeloupe; Hotessier! Beau! St. Thomas; Krebs!

Bullinula Beck.

- B. undata** Brug. (*B. nitidula* Sol., *B. elegans* Mke.) station: — ? — ; habitat: West-Indies; Mörch pg. 136! Cuba; Sagra! St. Vincent; Guilding! St. Johns; Ad.! Guadeloupe; Beau! St. Thomas; St. Croix; St. Martin; Kr.!

Bulla Linné.

- B. acuta** d'Orb. station: — ? — ; habitat: Cuba; Sagra! Jamaica; St. Thomas; Guadeloupe; Martinique; Candé! Guadeloupe; Beau!
- B. antillarum** d'Orb. station: — ? — ; habitat: St. Thomas; d'Orb.! Cuadeloupe; Beau!
- B. Auberii** d'Orb. station: — ? — ; habitat: Cuba; d'Orbig.!
- B. bidentata** d'Orb. station: — ? — ; habitat: Cuba; Sagra! Jamaica; St. Thomas; Guadeloupe; Martinique; d'Orb.! Guadeloupe; Beau! Porto-Plata; St. Thomas; Krebs!
- B. canaliculata** d'Orb. (see *Actaeon*) station: — ? — ; habitat: Cuba; Sagra! St. Thomas; Porto-Plata; Kr.!
- B. Candéana** d'Orb. (see *Philine*) station: — ? — ; habitat: Cuba; Sagra! Guadeloupe; Beau! Guadeloupe; Martinique; St. Thomas; Jamaica; Candé!
- B. caribaea** d'Orb. station: May 60, 3 specimens in

10 to 12 fathoms water on bluish sandy mud; habitat: Jamaica; St. Thomas; Martinique; Guadeloupe; Candé! Cuba; d'Orb.! Guadeloupe; Beau! Porto-Plata; Krebs! remark: 1 specimen is double the size as the shell described by d'Orb.

B. fragilis Lmk. station: — ? — ; habitat: Guadeloupe; Beau!

B. maculosa Mart. (*B. ibyx* Meusch., *B. umbilicata* Bolt., *B. striata* Brug. non Ph., *B. media* Ph.) station: — ? — ; habitat: West-Indies; Mörch pg. 136! Guadeloupe; Beau! Jamaica; Arecibo; Turksisland; Ad.! St. Thomas; St. Croix; St. Johns; Tortola; St. Martin; St. Barth; Barbadoes; Trinidad; Krebs!

B. Petitii d'Orb. station: on bluish mud in 10—12 fathoms water; May 1860 dredged 4 dead and 3 life specimens; habitat: Cuba; Auber! St. Thomas; Krebs! remark: the animal greenish and according to the three strong canaliculated semilunar teeth an *Atys*.

B. recta d'Orb. station: — ? — ; habitat: St. Thomas; Guadeloupe; Candé! Guadeloupe; Beau! Jamaica; Ad.! St. Thomas; Krebs!

B. physis Linné. station: — ? — ; habitat: St. Thomas; Krebs!

B. Sagrae d'Orb. station: — ? — ; habitat: Martinique; Candé!

B. solida Gml. (*B. ampulla* d'Orb.) station: — ? — ; habitat: West-Indies; Mörch pg. 136! Cuba; Sagra! Guadeloupe; Hotessier! Martinique; de Candé!

B. sulcata d'Orb. station: — ? — ; habitat: St. Thomas; Candé! Guadeloupe; Candé! Beau! Porto-Plata; Krebs!

B. undata Brug. is a *Bullinula*.

Atys de Montfort.

- A. *Guildingii* Sow. station: — ? — ; habitat: West-Indies; Mörch pg. 135! Jamaica; Ad.! St. Thomas; St. Martin; Kr.!

Philine Ascanias (*Bullaea* Lmk.).

- P. *candeana* (*Bullaea*) d'Orb. station: — ? — ; habitat: Guadeloupe; Candé! Beau!

Actaeon de Montfort (*Tornatella*).

- A. *canaliculata* (Bulla) d'Orb. (*B. olivula* C. B. Ad., *T. bullata* Ki.?) station: — ? — ; habitat: Guadeloupe; Beau! West-Indies; Mörch pg. 136!

Dimyaria.

1. Tellinacea.

Venus Linné.

- V. *affinis* Gm. (*C. laeta* Lmk. non Linné) station: — ? — ; habitat: Antillae; Mörch pg. 27!
- V. *albida* Gm. (*Cyth. albida* Gray acc. to C. B. Adams) station: — ? — ; habitat: Carthagera (from Krebs!); St. Thomas; Jamaica; C. B. Ad.!
- V. *antillarum* d'Orb. station: — ? — ; habitat: Florida; Cuba; St. Thomas; Jamaica; Martinique; d'Orb.! remark: Moll. de Cuba 2—278 pl. 26 fig. 41—43.
- V. *Auberiana* d'Orb. station: — ? — ; habitat: Cuba; Auber! St. Thomas; R. Swift! remark: Moll. de Cuba 2—227 pl. 26 fig. 35, 37.
- V. *Beauii* Recl. (see V. *subrostrata* Lmk.) station: — ? — ; habitat: Guadeloupe; Beau! (p. 24).
- V. *cardioides* Lmk. station: — ? — ; habitat: Cayenne; Jamaica; Lmk.!
- V. *circinata* Born. (*Dione*; *Cyth. guineensis* Lmk.) station: in white fine sand on 2 to 4 feet water in creeks where the water is smooth; habitat: Brasilia; Mörch pg. 27! Guadeloupe; Beau pg. 24! Jamaica; Ad.! Virgin-Island; R. Swift!*) St. Thomas; Trinidad; Carthagera; Krebs!
- V. *crenulata* Ch. (*V. crenata* Gray, *V. subrostrata* Lmk? *V. eximia* Ph., *Omphalocladium*) station: — ? — ; habitat: Antillae; Mörch pg. 24! Guadeloupe; Beau

*) List of Marine-Shells of the Virgin-Islands. March 63.

pg. 24! Jamaica; Ad.! Virgin-Islands; Swift! St. Thomas; Krebs!

V. *cubani* d'Orb. station: — ? — ; habitat: Cuba; Florida; Martinique; d'Orb.! St. Thomas? Swift! remark: Moll. de Cuba 2—278 pl. 26, fig. 44—46.

V. *Dione* L. (*Dione* Gray) station: — ? — ; habitat: Antillae; Mörch pg. 27! Trinidad but not at any of the Antilles; d'Orb.! Guadeloupe; Beau pg. 24! Carthage & Puerto Rico; Krebs! Carthage (from Redfield!); Jamaica; Navy-Bay; Ad.!

V. *dysera* — ? station: — ? — ; habitat: Carthage (from Redfield!); Ad.! remark: see V. *ziczac*.

V. *flexuosa* Linné. station: in sandy mud in creeks with smooth water; habitat: until 39° south; Martinique; Rio Janeiro; La Plata; d'Orb.! Guadeloupe; Beau pag. 24! Carthage; Jamaica; St. Thomas; Krebs! Jamaica; Ad.! Virgin-Islands; R. Swift! remark: R. Swift in his cat. of Marine shells of the Virgin Isl. places a sign of interrogation after „*flexuosa*“, in which he may be correct as the West-Indian shell does not answer in every respect to the description and drawings; he further gives a note of Woodward, according to which the W.-I. shell should be V. *punctifera* Gray of which Mr. S. has however not been able to find any account.

V. *granulata* Gm. (V. *marica* Ch., Cy. *lavacrum* Bolt.) station: in gravelly sand, in creeks with smooth water, close to the surface of the sea and to the shore; habitat: Antillae; Mörch pg. 23! Virgin-Isl.; Swift! Cuba; de la Sagra! Auber! Martinique; Guadeloupe; Candé! Guadeloupe; Beau pg. 24! St. Thomas (Th. Bland); St. Johns (E. Hartwig); Jamaica; Ad.! St. Thomas; St. Johns; Tortola; St. Croix; Carthage; Krebs!

- V. hebraea* Lmk. (*C. rubiginosa* Phil., *C. varians* Haul. by Swift, *C. varians* Ph. by Beau) station: 7 life specimens from bluish sandy mud in 10 to 12 fathoms water; habitat: Brasils; Mörch pg. 27! Rio-Janeiro; d'Orb.! Martinique; Guadeloupe; Candé! Beau pg. 24! St. Thomas; Krebs! Virgin-Isl.; Swift!
- V. inaequalvis* d'Orb. station: — ? — ; habitat: Cuba; Auber! Martinique; Florida; d'Orb.! Virgin-Isl.; Swift!
- V. Listeri* Gray. (Hanley by Swift; Dosina; *V. cancellata* Ch., *Chama persica* Meusch., *V. puerpera* var. 2 Lam. Desh.) station: — ? — ; habitat: Antillae; Mörch pg. 24! Virgin-Island; Swift!
- V. lucinalis* Lmk. station: — ? — ; habitat: St. Thomas; Lmk.!
- V. mactroides* Born. (*C. corbicula* Gml. Lmk.) station: — ? — ; habitat: Antillae; Mörch pg. 28! Guadeloupe; Cuba; d'Orb.! Guadeloupe; Curaçao; Jamaica; Ad.! St. Juan del Norte de Nicaragua; Chagres; Carthagen; Krebs!
- V. maculata* L. (*Card. tigrinum* Mart.) station: — ? — ; habitat: Antillae; Mörch pg. 28! Rio Janeiro; St. Lucia; Martinique; Guadeloupe; Cuba; Jamaica; d'Orb.! Guadeloupe; Beau pg. 24! St. Thomas; Krebs! Tortola; Marecaibo; Jamaica; Ad.! Virgin-Isl.; Swift!
- V. minuta* Koch. station: — ? — ; habitat: Virgin-Island; Swift!
- V. paphia* L. (*Chione*) station: in coarse sand in 1 foot water in creeks with smooth water; habitat: Antillae; Mörch pg. 24! Cuba; de la Sagra! St. Lucy; Martinique; Jamaica; Candé! Guadeloupe; Beau pg. 24! St.

- Thomas; Tortola; St. Johns; St. Croix; Carthagera;
Krebs! Jamaica; Ad.! Virgin-Island; Swift!
- V. *pectorina* Lmk. (V. *elegans* Gray; Timoclea; Leach.)
Brasilia; Mörch pg. 23! West-Indies? Krebs!
- V. *Philippi* d'Orb. (Cyther. *concentrica* Lmk., C. *patagonica* Phil.) station: — ? — ; habitat: Martinique;
Rio-Janeiro; d'Orb.! remark: Moll. de Cuba 2—270.
- V. *pygmæa* Lmk. station: — ? — ; habitat: Antillae;
Mörch pg. 24! Guadeloupe; Beau pg. 24! Jamaica;
Ad.! Virgin-Isl.; Swift! St. Johns; St. Thomas; Tortola;
St. Croix; St. Barth; Krebs!
- V. *rugosa* Ch. Gm. (non L., V. *rugosa* ind. orien. Ch., V. *rigida* Sol.; Dillw.) station: — ? — ; habitat: Antillae;
Mörch pg. 25! Rio Janeiro; d'Orb.! Martinique;
Candé! Guadeloupe; Beau pg. 24! St. Thomas; Tortola;
Jamaica; Ad.! St. Croix; Tortola; St. Johns; Krebs!
Virgin-Islands; Swift!
- V. *subrostrata* Lmk. (V. *Bouii* Recluz; see V. *crenul.*)
station: — ? — ; habitat: Isl. of St. Johns; Lmk.!
- V. *trigonella* Lmk. station: — ? — ; habitat: Virgin-Isl.; Swift!
- V. *trippla* Linne. station: — ? — ; habitat: Antillae;
Mörch pg. 28! Carthagera; New-Gr.; Krebs!
- V. *ziczæ* L. (V. *cancellata* L., V. *cingenda* Dw., V. *dysera* d'Orb.; V. *Lamarekii* Gray (Beau) —) station: in localities like those of V. *paphia*; habitat: Antillae;
Mörch pg. 24! Brasils; d'Orb.! Martinique & Guadeloupe;
Candé! Beau pg. 24! Jamaica; Ad.! Cuba; de la Sagra!
St. Thomas; Jamaica; Ad.! Virgin-Isl.; Swift! St. Croix;
St. Thomas; St. Johns; Tortola; Puertorico; Carthagera;
Krebs!

Artemis Poli.

- A. *concentrica* Born. (Dosina; Cy. *concentrica* Lmk.) station: — ? — ; habitat: Brasils; Mörch pg. 2! South-Caroline; Virginia; d'Orb.! Guadeloupe; Beau pg. 24! Virgin-Islands; Swift! St. Croix; St. Thomas; Krebs!
- A. *elegans* Con. station: — ? — ; habitat: St. Thomas (from Krebs!); Jamaica; South-Caroline; Ad.!
- A. *tenuis* Recluz. (Dosina & Cyclina Recluz, Lycinopsis Petit) station: in sand, white and fine; about 12 feet water in the harbour of St. Thomas; habitat: Virgin-Islands; Swift! St. Thomas; Krebs! Guadeloupe; Beau! remark: see Journ. de Conch. 1853 pg. 250, 1856 pg. 155. 2

Donax Linné.

- D. *cayennensis* Lmk. station: — ? — ; habitat: Bahia; St. Lucy; d'Orb.! Guayana; Lmk.!
- D. *denticulata* L. station: in white, fine sand in 2—3 feet water; habitat: Antillae; Mörch pg. 17! Guadeloupe; Beau pg. 25! Virgin-Isl.; Swift! St. Thomas; St. Croix; Crabisland; St. Johns; Tortola; St. Barth; Trinidad; St. Martha; Carthagenia; St. Juan del Norte; Chagres; Krebs!
- D. *martinicensis* Lmk. see Tell. *punicea* Born.
- D. *rugosa* d'Orb. station: — ? — ; habitat: Martini-que; Cuba; d'Orb.! Jamaica; Ad.!
- D — ? station: — ? — ; habitat: Cumana by E. Hartwig; Ad.! Chagres; St. Juan del Norte; Krebs!

Tellina Linné.

- T. *antillarum* d'Orb. station: — ? -- ; habitat:

- Cuba; Auber! Guadeloupe; Beau pg. 25! remark: Moll. de Cuba 2—250 pl. 30 fig. 45—46.
- T. Antoni* Ph. (Tellinella Gray) station: — ? — ; habitat: Guadeloupe; Beau pg. 25! Antillae; Mörch pg. 13! Virgin-Islands; Swift! remark: a doubtful species.
- T. bimaculata* L. (*T. sexradiata* Lmk.; *Heterodonax* Mörch; *Arcopagia* d'Orb.) station: in sand close to the shore; habitat: Antillae; Mörch pg. 17! Virgin-Islands; Swift! Cuba; Florida; d'Orb.! Bermuda from Redfield; Jamaica; Ad.! Tortola; St. Croix; St. Thomas; Trinidad; Curaçao; St. Martha; Krebs!
- T. brevifrons* Say? station: — ? — ; habitat: Jamaica; Ad.!
- T. Candearia* d'Orb. station: — ? — ; habitat: Guadeloupe; Beau pg. 25! Martinique; Candé! Rumkey, Turks-Island (Mrs. A. Holland); Ad.! remark: Moll. de Cuba 2—251 pl. 25 fig. 50—52.
- T. caribæa* d'Orb. station: — ? — ; habitat: Guadeloupe; Beau pg. 25! Petit! Guadeloupe; Cuba; d'Orb.! remark: Moll. de Cuba 2—250 pl. 30 fig. 47—49.
- T. carnaria* L. (*Strigilla* Turton; *Lucina carnaria* Lmk.) station: in sand close to the shore; habitat: Antillae; Mörch pg. 15! Guadeloupe; Beau pg. 25! Virgin-Is.; Swift! Brasils; Martinique; Guadeloupe; Cuba; d'Orb.! Jamaica; Ad.! St. Thomas; St. Johns; Tortola; St. Barth; St. Croix; Trinidad; Carthagera; Chagres; Krebs!
- T. cayennensis* Lmk. (*Phammobia*) see *T. constricta* Ph.
- T. cerena* Ad. station: — ? — ; habitat: St. Johns by E. Hartwig; Jamaica; Ad.!
- T. consobrina* d'Orb. station: — ? — ; habitat: Martinique; d'Orb.! remark: Moll. de Cuba 2—254 pl.

26 fig. 12—14; see Swifts remark to *T. subradiata* Sch.

- T. constricta* Phil. (*Solen Brug*; *Phammobia Lmk.*) station: — ? — ; habitat: Rio Janeiro; Cuba; d'Orb.! Carthagera by Redfield; Jamaica; Ad.! Nicaragua; Kr.!
- T. crystallina* Chem. (*T. Schrammii Recluz*) station: in fine white sand in bays with a strong surf; habitat: Carthagera; Krebs! Redfield! St. Thomas; R. Swift! H. Haagensen!
- T. cuneata* d'Orb. station: — ? — ; habitat: Guadeloupe; Beau pg. 25! d'Orb.! Cuba; Florida; d'Orb.! remark: Moll. de Cuba 2—256 p. 26 fig. 21—23.
- T. decussata* Ad. station: — ? — ; habitat: Carthagera (from Krebs!); Jamaica; Ad.!
- T. digitalina* Lmk. station: supposed to be in rather deep water; habitat: Guadeloupe; Beau pg. 25! Virgin-Islands; R. Swift!
- T. exilis* Lmk. station: — ? — ; habitat: Guadeloupe; Beau pg. 25! d'Orb.! Petit! Jamaica; Ad.!
- T. fausta* Soldr. (*Arcopagia Leach*; *T. remies Born non L.*; *T. laevis Wood*); station: in clayish sand close to the shore; habitat: Antillae; Mörch pg. 13! Guadeloupe; Beau! Martinique; St. Lucy; Candé! Cuba; Auber! New-Orleans; Alphen! Virgin-Isl.; Swift! Carthagera (by Redfield); Jamaica; Ad.! St. Thomas; Tortola; Carthagera; Krebs!
- T. flexuosa* Say (et Sow., *T. mirabilis Ph.*) station: — ? — ; habitat: Guadeloupe; Beau pg. 25! Guadeloupe; Jamaica; Florida; d'Orb.
- T. Gruneri* Ph. (*T. interstriata Say*; *T. inornata Ad. acc. to C. B. Ad.*) station: in sand in rather deep water in the harbour of St. Thomas; habitat: Virgin-Islands;

- Swift! St. Thomas; St. Johns; St. Croix; Krebs! Guadeloupe; Beau pg. 25! remark: Swift says in his „list of marine-shells from the Virgin-Islands“: „Mr. Dietz suggests that *T. int.* Say may be the same; see description in Binneys reprint of Say pg. 125“. After the List has been printed, we find a note to the same effect by C. B. Adams, who gives the above synonymes.
- T. Guadeloupensis* d'Orb. station: — ? — ; habitat: Guadeloupe; Beau pg. 25! d'Orb.! St. Thomas; Swift! remark: Moll. de Cuba 2—252 pag. 26 fig. 1—3.
- T. interrupta* Solander (*T. Listeri* Bolt., *T. maculosa* Lmk.) station: — ? — ; habitat: Guadeloupe; Beau pg. 25! Martinique; Candé! Cuba; de la Sagra! Virgin-Isl.; Swift! Bermuda (by Redfield); Tortola; Jamaica; Ad.! St. Thomas; St. Johns; Tortola; Carthagen; Kr.!
- T. laevis* Ch. (*T. laevigata* L.) station: — ? — ; habitat: Antillae; Mörch pg. 15! Virgin-Isl.; Swift! Bermuda (by Redfield); Tortola; Ad.!
- T. lineata* Hanl. (*T. lineata* Turt., *T. brasiliana* Lmk.) station: — ? — ; habitat: Rio Janeiro; d'Orbigny! Cuba; Martinique; Candé!
- T. magna* Spgl. (*T. acuta* Wood; *T. elliptica* Lmk.) station: — ? — ; habitat: Antillae; Mörch pg. 15! Bermuda; Ad.! Virgin-Islands; Krebs!
- T. Martinicensis* d'Orb. station: — ? — ; habitat: Guadeloupe; Beau pg. 25! d'Orb.! remark: Moll. de Cuba 2—255 p. 26 fig. 15—17.
- T. nitens* Ad. station: — ? — ; habitat: Jamaica; Adams!
- T. pfauperata* d'Orb. station: — ? — ; habitat: Guadeloupe; Martinique; d'Orb.! Beau pg. 25! remark: Moll. de Cuba 2—255 p. 26 fig. 18—20.

- T. pellucida* Ph. station: — ? — ; habitat: Guadeloupe; Beau pg. 25!
- T. pisiformis* L. (d'Orb.) station: — ? — ; habitat: Puertorico; d'Orb.! Guadeloupe; Beau pg. 25! Jamaica; Chagres (by Krebs); Ad.!
- T. pudica* Hanl. station: — ? — ; habitat: Guadeloupe; Beau pg. 25!
- T. punicea* Born. (*Peronaeoderma* Mörch pg. 12, *T. angulosa* Gm. accord. to Swift, *Donax martinisensis* Lmk., *T. alternata* Say & Sowb. acc. to d'Orb., *T. striata* Ch. acc. to C. B. Ad.) station: in sand outside the surf.; habitat: North-America; Rio-Janeiro; Martinique; Guadeloupe; d'Orb.! Virgin-Isl.; Swift! Surinam; Jamaica; Ad.! Carthagen; Redfield! Krebs! remark: see *T. striata* Hanl.
- T. radiata* L. (*Musculus*; Mörch pg. 13! *T. unimaculata* Lmk.; *T. nivea* Wood) station: in sand on 4 to 6 feet water; habitat: Guadeloupe; Beau pg. 25! Cuba; Martinique; Jamaica; New - Orleans; d'Orb.! Bermuda (by Redfield); Jamaica; Ad.! St. Thomas; St. Croix; St. Johns; Tortola; Trinidad; Carthagen; Krebs!
- T. sincera* Hanl. station: — ? — ; habitat: Guadeloupe; Beau pg. 25!
- T. similis* Sow. station: — ? — ; habitat: Antillae; Mörch pg. 14! Martinique; Candé! Virgin-Isl.; Swift!
- T. striata* Hanl. (non Chem. nec Philippi; *Don. martinicensis* Lmk. acc. to Swift!) station: — ? — ; habitat: Virgin-Islands; Swift! remark: Hanleys R. S. pl. 13 fig. 10 suppl.; Swift says: Hanley in his ill. & desc. cat. of recent shells plate 13 fig. 10 app. page 347 refers *Don. mart.* to *Tellina* as *T. striata* Ch. but the figure vol. x. tab. 170 fig. 1654—55 represents a variety of

- T. punicea* Born. *T. striata* H. stands in Beau's Cat. of Guadeloupe shells, but specimens of *T. subradiata* Schum. have been sent from Guadeloupe (by Beau) as *T. striata*. See *T. punicea* Born.
- T. Sauleyetiana* Recluz. station: — ? — ; habitat: Guadeloupe; Beau pg. 25! Virgin-Isl.; St. Thomas in Longbay; Krebs!
- T. subradiata* Schm. station: — ? — ; habitat: St. Thomas (by Krebs); St. Johns (by Hartwig); Ad.! Virgin-Isl.; Swift! remark: Philippi makes *T. subradiata* Schum. synonyme with *T. alternata* Say, but Swift thinks them distinct according to the specimens of his collection; we do not see any difference though we have procured authentic north-american specimens of *T. alt.* Say to compare with West-Indian specimens.
- T. vespuciana* d'Orb. station: — ? — ; Martinique; Jamaica; d'Orb.! remark: Moll. de Cuba 2—254 pl. 26 fig. 12—14.
- T. vitrea* d'Orb. station: — ? — ; habitat: Guadeloupe; Beau pg. 25! d'Orb.! remark: Moll. de Cuba 2—253 pl. 26 fig. 4—5.

Psammobia Lmk.

- P. squamosa* Lmk. station: — ? — ; habitat: Virgin-Islands; Swift! remark: very rare.
- P. spec. ind.* Virgin-Islands; Swift!

Sanguinolaria Lmk.

- S. nivea* (Solen) Chem. (*S. rosea* Lmk., Tell. *rosea* Gm., Sol. *sanguinolentus* Gm., Sol. *fucatus* Spgl., Tell. *rubicunda* Bol., *Lobaria rosacea* Sch.) station: in sand outside the surf or in the surf; habitat: Cuba; Martini-

que; Guadeloupe; d'Orb.! Marie Galante; Beau pg. 24! Virgin-Islands; Swift! Jamaica; Ad.! St. Thomas; St. Croix; Tortola; Carthagera; Trinidad; Krebs!

Capsa Bruguière.

- C. coccinea* (Cardium) Martyn. (*C. efflora* in d'Orb., Tell. anomale Ch. Bolt., *Corbula radiata* Bolt., *Caps. rugosa* var. Lmk., Sang. *rugosa* Blv., *S. rug.* v. 2 Anton., Asaphis) station: in coarse sand in creeks where the water is smooth; habitat: Florida; Cuba; Martinique; Guadeloupe; St. Lucy; d'Orb.! Carthagera & Bermuda (by Redfield); Jamaica; Ad.! Virgin-Islands; Swift! St. Thomas; St. Johns; St. Croix; Tortola; Carthagera; Krebs! remark: Mörch pg. 9!; Beau pg. 24 says: it is not eatable. But this is a mistake; it is often in the market of St. Thomas.

Mactra L.

- M. alata* Wood. (*M. carinata* Lmk., *M. striatula* L.⁴ Hanley) station: in sand outside a strong surf at Carthagera; habitat: Brasils; Mörch pg. 4! Virgin-Islands; Puertorico; Venezuela; Swift! Jamaica; Ad.! Carthagera; Krebs!
- M. donacaeformis* Gray. (*Mulinia*) station: — ? — ; habitat: Nevis; Gray! remark: we think it synonyme with *M. guad.* Recl.
- M. guadeloupensis* Recl. station: — ? — ; habitat: Guadeloupe; Beau pg. 26! Guayvia on the coast of Venezuela in Swifts collection; Blume!

Donacilla Lmk. (*Mesodesma* Desh.)

- D. rosea* d'Orb. station: — ? — ; habitat: Martini-

que; Candé! remark: described in Moll. de Cuba 2—238 pl. 25 fig. 39—41.

Semele Sch. (*Amphidesma* Lmk.)

- S. cancellata* d'Orb. (A. cane. Sow.) station: — ? — ; habitat: Guadeloupe; Martinique; d'Orb.! Guadeloupe; Marie Galante; Beau! Virgin-Islands; Swift! St. Thomas; Krebs! remark: Beau pg. 25; Moll. de Cuba 2—241 pl. 25 fig. 42—44.
- S. decussata* Wood. (*Tellina* Wood, A. Jayanum Ad.) station: — ? — ; habitat: Virgin-Isl.; Swift! Guadeloupe; Beau! St. Johns (by E. Hartwig); St. Thomas (by Bland) & Jamaica; Ad.! remark: Wood in G. C. 160 pl. 43 fig. 2—3 and Ind. test. p. 28 pl. 4 fig. 81; Ad. in proc. Bost. Soc. N. H. 2 pg. 10.
- S. formosum* Sowb. station: — ? — ; habitat: St. Johns (by E. Hartwig); Jamaica; Ad.!
- S. purpurascens* Gm. (*Venus* Gm.; A. *variegata* Lmk.; *Tellina obliqua* Wood.) station: — ? — ; habitat: Antillae; Mörch! Virgin-Isl.; Swift! Rio-Janeiro; Cuba; Martinique; Guadeloupe; d'Orb.! Guadeloupe; Beau! St. Thomas; Krebs! remark: Wood in Ind. test. pg. 22 Hanley's ed.; Mörch pg. 16; Beau pg. 25.
- S. reticulata* Lmk. (*Tellina* Lmk.; *Lucina reticulata* Lmk.) station: — ? — ; habitat: Antillae; Mörch! Guadeloupe; Beau! Virgin-Islands; Swift! Rio Janeiro; Cuba; Martinique; d'Orb.! St. Thomas; Portoplata; Krebs! remark: Chem. vol. 6 fig. 118; Mörch pg. 15; Beau pag. 25.
- S. subtruncata* Sow. station: — ? — ; habitat: Guadeloupe; Beau! Virgin-Isl.; Swift!

Cumingia Sowb.

- C. *Antillarum* d'Orb. station: in crevices on the lower part of madrepores in 8 to 10 feet water; habitat: Virgin-Islands; Swift! Jamaica; Ad.! St. Thomas; Swift! Bland! J. Knox! Krebs! Guadeloupe; Beau! St. Domingo; d'Orb.! remark: Moll. de Cuba Lavignon 2—236 pl. fig. 36—38.
- C. *lineata* d'Orb. station — ? — ; habitat: North-America until Brasils; d'Orb.! *C. lineata* d'Orb., Say
- C. *sinuata* A. Adams. station: — ? — ; habitat: St. Thomas; A. H. Riise! remark: is this not synonyme with C. *Ant.* d'Orb.?
- C. *Petitiana* d'Orb. station: — ? — ; habitat: Guadeloupe; d'Orb.! Beau pg. 25! remark: see *Scrobicularia*; Moll. de Cuba 2—236 pl. 25 fig. 33—35.

Scrobicularia Schumacher. (Lavignon; d'Orbigny!
Cumingia; Sowb.)

Cyclas Bruguière.

- C. *maritima* d'Orb. station: in muddy lagoons; habitat: Cuba; Auber! St. Thomas; Swift! remark: Moll. de Cuba 2—280 pl. 26 fig. 47—60.

Glaucanome Gray.

- G. sp. ind. station & habitat: in the creeks and lagoons by Carthagera, N. Gr.; Krebs!

Saxicava Fleurian de Bellevue.

- S. *minuta* Guild. station: in a crevice of a dead madre-pore; habitat: St. Croix; A. H. Riise! St. Thomas; Krebs!

Petricula Lmk.

- P. divaricata* Ch. station: in crevices on madrepores; St. Thomas; Virgin-Isl.; Swift! remark: Swift says it distinguishes itself by purple syphons from *P. lapicida* Ch., which has yellow syphons; see *P. lapicida* Ch. vol. 10 tab. 172 fig. 1666—67, Woods Ind. test. pl. 8 fig. 73, Sowb. Tes. pl. 146 fig. 24—25.
- P. lapicida* Chem. (*Venus* Chem., *P. costata* Gray & Lmk., *P. divaricata* Chem., *Ven. divergens* Gm., *Pr. lucinalis* Lmk.) station: in crevices on dead madrepores; habitat: Australia; Sowb.! St. Thomas; Swift! Krebs! St. Domingo; Cuba; Martinique; d'Orb.! Antillae; Mörch! remark: we have a large number of specimens, but are not able to distinguish the shells of *P. div.* & *lap.*; Swifts remark about the syphons (see *P. div.*) are very interesting, but perhaps the colour of the syphons varies by the same species.
- P. pholadiformis* Lmk. station: — ? — ; habitat: United States; Cuba; d'Orb.!
- P. rocellaria* Lmk. station: — ? — ; habitat: St. Thomas (by J. H. Newton); Jamaica; Ad.!
- P. typica* (*Chloristodon*) Jonas. station: in crevices and holes in Madrepores on 2 to 10 feet water; habitat: Antillae; Mörch pg. 20! Virgin-Islands; Swift!

Anatinacea.**Periploma Schum.**

- P. inæquivalvis* Schum. station: — ? — ; habitat: Marie Galante; Beau pg. 25!

Thracia Leach. (Blainv. 1824).

- T. rugosa* Conrad. station: — ? — ; habitat: Rio Janeiro; d'Orb.!

- T. plicata*? Desch. (*T. undata* or *undulata* in Cumings Collection) station: in sand on about 8 to 20 feet water; habitat: Virgin-Islands; Swift! St. Thomas; Tortola; Krebs! remark: Encycl. meth. vol. 3, pg. 1039.
- T. sp. ind.* station: — ? — ; habitat: Virgin-Islands; Swift! remark: like *Phaseolina*.

Magdala Leach. (*Lyonsia* Turton; *Osteodesma* Desh.)

- M. Beauiania* d'Orb. station: — ? — ; habitat: Antillae; d'Orb.! Guadeloupe; Beau! remark: Moll. de Cuba 2—225 pl. 25 fig. 26—28; Beau pg. 26! Deshayes 1827.
- M. sp. ind.* station: — ? — ; habitat: Virgin-Islands; Swift!

Corbulacea.

***Corbula* Bruguière.**

- C. Barrattiana* Ad. station: — ? — ; habitat: Jamaica; Ad.! remark: desc. in Contr.
- C. Blandiana* Ad. station: — ? — ; habitat: Jamaica; Ad.! remark: desc. in Contr.
- C. caribæa* d'Orb. station: numerous in mud on 6 fathoms water together with a few specimens of *C. Kjærana*; habitat: Cuba; Auber! St. Domingo; Candé! St. Thomas; Krebs! remark: Moll. de Cuba 2—284 pl. 27 fig. 5—8; Encycl. meth. pl. 230, 1792.
- C. cubaniana* d'Orb. station: — ? — ; habitat: Cuba; Auber! Guadeloupe; Beau! remark: moll. de Cuba 2—283 pl. 26 fig. 51—54.
- C. Chyttiana* Ad. station: — ? — ; habitat: Jamaica; Ad.! remark: Described in Contr.

- C. Dietziana* Ad. station: — ? — ; habitat: Jamaica; Ad.! remark: Descr. in Contr.
- C. disparalis* d'Orb. station: — ? — ; habitat: Jamaica; Martinique; Guadeloupe; Cuba; d'Orb.! remark: Moll. de Cuba 2—283 pl. 27 fig. 1—4.
- C. duplex* Ad. station: — ? — ; habitat: Jamaica; Ad.! remark: desc. in Contr.
- C. elliptica* Recl. station: — ? — ; habitat: Guadeloupe; Beau pg. 26!
- C. Kjærana* Ad. station: — ? — ; habitat: Jamaica; St. Juan de Nicaragua (by Krebs); Ad.! St. Thomas; Krebs! remark: desc. in contr.
- C. Knoxiana* Ad. station: — ? — ; habitat: Jamaica; Ad.! remark: descr. in contr.
- C. Krebsiana* Ad. station: numerous on 10 to 12 fathoms water together with a few *C. caribæa* d'Orb.; habitat: Jamaica; Ad.! St. Thomas; Krebs! remark: descr. in contr.
- C. Lavelleana* d'Orb. station: in mud on 12 feet water; habitat: Jamaica; Martinique; Guadeloupe; Cuba; d'Orb.! Virgin-Islands; Swift! St. Thomas; Krebs! remark: describsd in moll. de Cuba 2—284 pl. 27 fig. 9—12.
- C. nitens* Turton. (*Ervilia*, *Mya nitens* Laskey, *Amphidesma purpurascens* Lmk. No. 14) station: — ? — ; habitat: Antillae; Mörch! St. Thomas; Tortola; Krebs! Virgin-Islands; Swift! remark: descr. Turtons B. B. pg. 55 pl. 19 fig. 4.
- C. Newtoniana* Ad. station: — ? — ; habitat: Jamaica; Ad.! St. Thomas; Th. Bland! remark: descr. in Contr.
- C. operculata* Phil. station: one specimen dredged in

May 1860 in bluish sandy mud on 10 to 12 fathoms water; habitat: St. Croix; Mörch pg. 30! Virgin-Isl.; Swift! St. Thomas; Krebs!

- C. *qvadrata* Hinds (Eucharis Recl.) station: — ? — ; habitat: Antilles; d'Orb.! Guadeloupe; Beau pag. 26! St. Thomas; Th. Bland! Krebs! Porto - Plata; Krebs! Virgin - Islands; Swift! remark: Hinds Z. P. 1843, Reeve Conch. icon. sp. 40, Jour. de Conch. 1850 pg. 168.
- C. *Swiftiana* Ad. station: — ? — ; habitat: Jamaica; Ad.! St. Thomas; Bland! remark: described in Cont.; several of the Jamaica species are identic with d'Orbigny's species.

Neaera Gray. (*Sphena* Turton acc. to d'Orb.)

- N. *alternata* d'Orb. station: — ? — ; habitat: Martinique; d'Orb.! remark: desc. in Moll. de Cuba 2—286 pl. 27 fig. 17—20.
- N. *cleryana* d'Orb. station: — ? — ; habitat: Jamaica; Cuba; St. Thomas; Guadeloupe; d'Orb.! remark: desc. Moll. de Cuba 2—285.
- N. *ornatissima* d'Orb. station: dredged 4 specimens in May 1860 in bluish-sandy mud from 10 to 12 fathoms water; habitat: Guadeloupe; Martinique; Cuba; St. Thomas; d'Orb.! St. Thomas; Swift! Krebs! remark: desc. in Moll. de Cuba 2—286 pl. 27 fig. 13—16.

Myacea.

Pholadomya G. B. Sowerby 1823.

- P. *candida* Sowb. station: St. Thomas harbour in sand on 10 to 20 feet water about 40 to 60 feet from the shore; habitat: Tortola; Mörch pg. 5! Guadeloupe; Beau pag. 26! St. Thomas; E. Hartwig! Dr. Ravn! Vir-

gin-Islands; Swift! Tortola; St. Thomas; Krebs! remark: desc. in Gen. of shells No 19 fig. 184.

Solenacea.

Solen Linné 1758.

- S. bidentatus* Spgl. (*Tagelus* Gray, *Siliqvaria* Sch., *Macha* Gray, *S. bidens* Ch., *S. divisus* Fabr, *S. fragilis* Sol., Pult, *S. tenuis* Wood Sup.?) station: lives in sand close to the shore even at places where the sand remained dry when low-water and in mud on 10 to 12 fathoms water; habitat: Antillae; Mörch pg. 8! Martinique; Candé! Guadeloupe; Beau pg. 26! Jamaica; Ad.! Virgin-Islands; Swift! St. Thomas; St. Croix; St. Johns; Trinidad; Carthagera; St. Juan de Nicaragua; Krebs!
- S. lucidus* Spgl. (*Siliqua* Meg., *Machaera* Gould, *Aulus* Oken, *Liguminaria* Sch., *Solecurtoides* Desmoulins) station: — ? — ; habitat: America centralis; Mörch pag. 7!
- S. obliquus* Spgl. (*S. ambiguus* Lmk.; *Solena* Browne) station: in sand close to the shore; habitat; Cuba; Sagra! Puertorico; Mörch! Martinique; Candé! Guadeloupe; Beau! Jamaica; Ad.! Virgin-Isl.; Swift! Porto-Plata; St. Thomas; Puertorico; Krebs! remark: Lmk. No 7, Hanleys R. S. pg. 13, Woods Ind. test. sup. pl. 11 fig. 17.
- S. gibbus* Spgl. (*S. caribæus* Lmk., *Siliqvaria notata* Sch., *Baphia coerulescens* Meusch?) station: — ? — ; habitat: Antillae; Mörch pg. 8! Cuba; d'Orb.! North-Bedford in Mass.; Florida; Cuba; Ad.! Arecibo; J. H. Newton! Virgin-Islands; Swift! St. Thomas; Carthagera; St. Juan de Nicaragua; Krebs! remark: Mollusques de Cuba 2—231.

Psammosolea Risso. (Solecurtis Desh.)

- P. Sanctae Marthae* Ch. (Solen Ch.) station: in sand not very distant from the shore; habitat: Antilles; d'Orb.! Guadeloupe; Beau pg. 26! Virgin-Islands; Swift! St. Thomas; Tortola; Krebs! remark: Moll. de Cuba 2—232 pl. 25 fig. 31—32.

Pholadea.**Pholas Linné.**

- P. campechensis* Gml. station: found dead on the sandy beach; habitat: Carthagera; Krebs!
- P. candeana* d'Orb. station: — ? — ; habitat: Martinique; Candé! Havana; Sagra! Florida; d'Orb.! remark: Moll. de Cuba 2—215 pl. 25 fig. 18—19.
- P. caribæa* d'Orb. station: — ? — ; habitat: Cuba; Sagra! Mexico; Pet. de Saussaye! Virgin-Islands; Swift! St. Thomas; Krebs! remark: Moll. de Cuba 2—216 pl. 25 fig. 20—22.
- P. corticaria* Sowb. station: — ? — ; habitat: Jamaica; Ad.!
- P. costata* (L.) Gray. station: — ? — ; habitat: Antillae; Mörch pg. 3! Cuba; New-Orleans; d'Orb.! North-Bedford in Mass; Florida; Ad. Mayaguez; Swift!
- P. Hornbecki* d'Orb. (*P. Beauiana* Recl.) station: — ? habitat: St. Thomas; Hornbeck! Guadeloupe; Beau pg. 27! Virgin-Islands; Swift! remark: Moll. de Cuba 2—217 pl. 25 fig. 23—25, Journ. de Conch. 1853 pg. 49 pl. 2 fig. 1—3.
- P. Krebsii* Ad. station: — ? — ; habitat: St. Thomas (by Krebs); Ad. remark: by us supposed to be *P. striata* L.
- P. pusilla* L. station: in wood close to the shore; ha-

bitat: Rio-Janeiro; Cuba; d'Orb.! St. Thomas?; Kr.!
 remark: see *P. striata* L.

- P. striata* L. (*Martesia* Leach, *Ph. pusilla* L., *Ph. clavata* Lmk.) station: in wood in the surface of the water; Antillae; Mörch pg. 2! Virgin-Isl.; Swift! Guadeloupe; Beau pg. 27! St. Thomas; Charthagena; Krebs! remark: Woods Ind. test. tab. 2 fig. 7.

Teredo L.

- T. navalis* L. station: in wood in the surface of the sea; habitat: Cuba; Auber! St. Thomas; Krebs!
T. norvegica Spgl. station: in wood; habitat: Guadeloupe; Beau pg. 27!

Gastrochaenacea.

Gastrochaena Spengler 1783.

- G. hians* (Pholas) Ch. (*Rocellaria* Fl. de Bellv, *Trapezium* Blv., *G. cuneiformis* Phil. non Spgl. fide Mörch, *Fistulana rupestris* Bosc., *G. truncata*? Sowb.) station: in life madrepores from the surface of the sea until on 20 feet water; habitat: Antillae; Mörch pg. 1! Guadeloupe; Beau pg. 27! Virgin-Isl.; Swift! St. Domingo; Cuba; Martinique; St. Thomas; d'Orb.! St. Thomas; St. Croix; Krebs! remark: its tubes sometimes 8 to 10 inches long.; Chem. vol. 10 tab. 172 fig. 1678—79; Wood's Ind. test. pl. 2 fig. 11, Moll. de Cuba vol. 2 pg. 228, Hanley's Ed. fig. Woods Ind. test. pl. 2 fig. 114 text., Wood's Ind. test. suppl. pl. 9 fig. 42.
G. ovata? Sowb. Z. P. 1834. station: — ? — ; habitat: Virgin-Isl.; Swift! remark: Woods Ind. test. Suppl. pl. 9 fig. 42.
G. rostrata Spgl. (*Ph. hians* Ch., *G. callosa* Phil., *G.*

Chemniziana d'Orb., *Rocellaria rostrata* Spgl. fide Mörch) station: in life madrepores until on 20 feet water; we have never seen the tubes over one inch long; they are general in pieces of madrepores in which *Gast. hians* and several species of the boring lithophagi are living; habitat: Antillae; Mörch pg. 1! Guadeloupe; Beau pg. 27! St. Thomas by Dr. Hornbeck acc. to d'Orb.! Virgin-Islands; Swift! St. Thomas; Tortola; St. Croix; Kr.! remark: Moll. de Cuba 2—229 pl. 25 fig. 29—30, Chem. vol. 10 tab. 172 fig. 1680—81, Jonas Zeits. für Mal. 1844 pg. 136.

Cardiacea.

Cardium Linné 1758.

- C. antillarum* d'Orb. (*C. graniferum* Brod.) station: in sand between the roots of sea-weed; habitat: Virgin-Islands; Swift! Guadeloupe; Beau pg. 23! Cuba; Guadeloupe; Martinique; Jamaica; d'Orb.! remark: Moll. de Cuba 1846 2—309 pl. 27 fig. 53—55, Sowb. 2 pg. vol. 4—346, Reeve Conch. icon 1844; Swift says: Broderips name was first published & should have precedence.
- C. citrinum* Ch. (*Liocardium* Sws., *C. triste* L.?, *C. serratum* Ch. non L., *C. hiatus* Meuschen, *C. lineatum* Gm., *O. laevigatum* Lmk.) station: in sand between sea-weed; habitat: Antillae; Mörch pg. 35! Guadeloupe; Beau pg. 23! Rio Janeiro; Cuba; Martinique; St. Lucy; Guadeloupe; d'Orb.! Bermuda (by Redfield); Jamaica; Ad.! Virgin-Islands; Swift! St. Thomas; Tortola; five varieties from Hamsbluff at St. Croix; Trinidad; Krebs!
- C. isocardia* L. (*Trachycardium* Mörch pg. 34, *Pectunculus* Mart.) station: — ? — ; habitat: Antillae; Mörch! Guadeloupe; Beau pg. 23! Cuba; Sagra! Mar-

tinique; St. Lucy; d'Orb.! Jamaica; Carthagera (by Redfield); Ad.! Virgin-Islands; Swift! St. Thomas; St. Croix; Tortola; St. Johns; Trinidad; Krebs!

C. laevigatum L. (Reeve, *C. pristis* Valene) station: — ? — ; habitat: St. Thomas; Mörch pg. 35! Guadeloupe; Beau! Margarita; d'Orb.! St. Thomas; Tortola; Ad.! Virgin-Islands; Swift! St. Thomas; St. Croix; Tortola; Krebs!

C. leucostomum Born. (*C. marmoreum* Lmk.) station: — ? — ; habitat: Antillae; Mörch pg. 34!

C. medium L. (*Fragum* Bolt., *Hemicardium* Sws.) station: — ? — ; habitat: Antillae; Mörch pg. 36! Guadeloupe; Beau pg. 23! Florida; Cuba; Martinique; St. Lucy; d'Orb.! Jamaica; St. Thomas; Carthagera; Ad.! Virgin-Isl.; Swift! St. Thomas; Tortola; St. Croix; Crab-island; St. Johns; Krebs!

C. muricatum L. station: in sand between *Zostera* in 1 to 4 feet water; habitat: Antillae; Mörch pg. 34! Guadeloupe; Beau pg. 23! Rio Janeiro; Cuba; Martinique; Guadeloupe; Jamaica; d'Orb.! Carthagera; Jamaica; South-Caroline; Ad.! Puertorico; Virgin-Islands; Swift! Krebs!

C. oviputamen Reeve. station: — ? — ; habitat: Carthagera (by Krebs); Ad.!

C. Petitianum d'Orb. (*C. ringiculum* Sowb.) station: in sand between *Zostera*; habitat: Guadeloupe; d'Orb.! Beau pg. 23! St. Vincent; Guilding; Virgin-Isl.; Swift! St. Thomas; St. Johns; Tortola; Krebs! remark: moll. de Cuba 2—309 pl. 27 fig. 50—52, Sowb. Z. L. 1840, Conch. icon. spg. 115; Swift says: Sowerby's name should be retained as it was published long before d'Orbignys.

C. spinosum Meusch. (*Papyridea* Sws., *Solen bullatus* Ch.

- non Linné, *C. soleniforme* Brug., *C. bul. pars* Lmk.) station: in sand on shallow water; habitat: Antillae; Mörch pg. 33! Guadeloupe; Beau pg. 23! Cuba; Martinique; St. Lucy; Guadeloupe; d'Orb.! Tortola; Jamaica; Ad.! Virgin-Islands; Swift! Trinidad; Carthagenia; Kr!
- C. subelongatum* Sow. station: — ? — ; habitat: Guadeloupe; Beau pg. 23! St. Thomas; Cuba; Martinique; d'Orb.! Virgin-Isl.; Swift! Hamsbluff at St. Croix; Krebs!

Chamaea.

Chama L.

- Ch. arcinella* L. (*Arcinella spinosa* Schm.) station: often on *Strombus* pug. between seaweed on 8—10 feet water; habitat: Antillae; Mörch pg. 37! Guadeloupe; Beau pg. 23! Guadeloupe; Martinique; Cuba; d'Orb.! Jamaica; Ad.; Virgin-Islands; Swift! St. Thomas; St. Croix; Tortola; Krebs! remark: when young moving about.
- Ch. ferruginea* Reeve. station: in mud about 2 feet water in creeks; habitat: Antillae; Mörch pg. 37! Virgin-Islands; Swift! St. Thomas; St. Johns; Tortola; Trinidad; Krebs!
- Ch. florida* Lmk. station: on rocks; habitat: Guadeloupe; Beau pg. 23! St. Domingo; Lmk! Virgin-Isl.; Swift! St. Thomas; St. Croix; St. Barth; Crab Island; Kr!
- Ch. frondosa* Brod. station: — ? — ; habitat: Guadeloupe; Beau pg. 23!
- Ch. macrophylla* Ch. (*Ch. bicornis* L., *Ch. imbricata* Lmk.) station: in mud close to the shore; habitat: Antillae; Mörch pg. 36! Cuba; Martinique; Guadeloupe; St. Domingo; St. Lucy; d'Orb.! Jamaica; Puertorico; Bermuda; Ad.! Virgin-Islands; Swift! Krebs!

- Ch. radians* Lmk. station: — ? — ; habitat: Guadeloupe; Beau pg. 23!
- Ch. sarda* Reeve. station: fastened on rocks and accord. to Beau on *Strombus gigas*; habitat: Antillae; Mörch pg. 37! Guadeloupe; Beau pg. 23! Virgin-Isl.; Swift! St. Thomas; St. Johns; St. Barth; Trinidad; Krebs!
- Ch. sordida* Reeve? station: — ? — ; habitat: Central-America; Mörch pg. 37!
- Ch. variegata* Reeve. station: — ? — ; habitat: Virgin-Islands; Swift!
- Ch. venosa* Reeve. station: — ? — ; habitat: — ? — ; Guadeloupe; Beau pg. 23!
- Ch. 3 sp.* not determ. in the collection of R. Swift Esq.; they are from the Virgin-Islands.

Lucinacea.

Lucina Bruguière 1792.

- L. americana* Ad. station: — ? — ; habitat: St. Thomas (by J. H. Newton); Jamaica; Ad.! remark: see contributions. = *Diplommatina*
- L. aurantia* Desh. (V. pens. var. Ch.) station: in sand on 4 to 6 feet water, where the wind plays freely with the waves; habitat: Antillae; Mörch pg. 32! Guadeloupe; Beau pg. 23! Virgin-Islands; Swift! common at Smithsbay, St. Thomas; St. Croix; Krebs! remark: Chem. vol. 7 fig. 396.
- L. Candearia* d'Orb. station: — ? — ; habitat: Guadeloupe; Martinique; d'Orb.! Guadeloupe; Beau pg. 23! Virgin-Isl.; Swift! remark: Moll. de Cuba 2—299 pl. 27 fig. 43—45.
- L. caribaea* d'Orb. station: — ? — ; habitat: Guadeloupe; Beau pg. 23! *sp. inedita?*

- C. chrysostoma* Meuschen (*Loripes* Poli, *V. edentula* Ch., *Anodonta alba* Link., *L. Philippiana* Reeve) station: in sand on shallow water in creeks; habitat: Florida; Cuba; Martinique; d'Orb.! Puertorico; Jamaica; Ad.! Carthagenae; St. Thomas; St. Croix; Tortola; Puertorico; Krebs! Virgin-Isl.; Swift! remark: Chem. 7—427—29; Phill. Abb. pt. 3 pl. 1 fig. 2; Swift observes: the shell Phill. refers to *L. edent.* is in *Conch. icon. sp.* 23 *L. Philippiana* Reeve; d'Orbigny draws the attention to the marvelous hurry with which this spec. & they *L. jamaicensis* breed; it is used as food.
- L. costata* d'Orb. (*L. textilis* Ph., *L. antillarum* Reeve, *L. ornata* Ad.) station: — ? — ; habitat: Rio Janeiro; Havana; St. Thomas; Guadeloupe; Jamaica; d'Orb.! Jamaica; Ad.! St. Thomas; St. Croix; Krebs! Virgin-Isl.; Swift! remark: see *pilula* Ad.
- L. digitalis* Lmk. (*T. pisiformis* L.) station: — ? — ; habitat: Virgin-Islands; Swift! *Strophodonta*
- L. flexuosa* d'Orb. station: — ? — ; habitat: Guadeloupe; Beau pg. 23! *L. flexuosa* d'Orb.
- L. granulosa* C. B. Ad. (*L. semireticulata* d'Orb.) station: — ? — ; habitat: Jamaica; St. Thomas; Ad.! remark: Moll. de Cuba 2—297; is this not *L. antillarum* Reeve? No. = *Diplodonta semireticulata* Ph.
- L. janeiroensis*? Reeve. station: — ? — ; habitat: Jamaica; St. Thomas (by Krebs); Ad.! *Diplodonta*
- L. jamaicensis* Spgl. (*Codakia* Scopoli) station: in sand in smooth water; habitat: Antillae; Mörch pg. 32! Guadeloupe; Beau pg. 23! Rio - Janeiro; Martinique; Cuba; Jamaica; d'Orb.! St. Thomas; St. Croix; St. Johns; Carthagenae; St. J. Nicaragua; Tortola; Krebs! Jamaica;

- Ad.! Virgin-Isl.; Swift! remark: Chem. 7 pg. 409; used as food; d'Orbigny remarks: it breeds very fast.
- L. obliqua* Reeve non Phill. (*L. pectinata* C. B. Ad., *L. pecten* Reeve, *L. occidentalis* Reeve) station: — ? —; habitat: Virgin-Islands; Swift! remark: Swift says: Adams name has the precedence if it is synonyme and *L. obl.* Ph. is quite distinct from *L. obl.* R.; Contributions pg. 245 Nov. 1851, Catalogue of N. | A. shells January 1847, Conch. Icon. June 1850 sp. 42 & Aug. 1850.
- L. pecten* Lmk. (*L. imbricatula* C. B. Ad. acc. to Beau pg. 23, *L.*) station: in coarse sand on one foot water in a creek; habitat: Jamaica; Ad.! Antillae; Mörch pg. 33! Guadeloupe; Beau pg. 23! St. Thomas; Tortola; Krebs! remark: Contributions.
- L. pensylvanica* (Venus) *L.* (*Codakia* Scop.) station: in sand close to the shore in shallow water; May 60 dredged 7 specimens of one line diameter in 6 fathoms water on bluish sandy mud; habitat: Antillae; Mörch pg. 32! Guadeloupe; Beau pg. 23! Virgin-Isl.; Swift! Jamaica; Yucatan; Ad.! New-Orleans; Martinique; Cuba; St. Thomas; d'Orb.! St. Thomas; St. Croix; St. Johns; Tortola; Krebs!
- L. pilula* Ad. (*L. costata* d'Orb.?) station: — ? —; habitat: St. Thomas (by Krebs); Ad.! *obliqua*
- L. punctata* (Venus) *L.* station: — ? —; habitat: Antillae; Mörch pg. 33! = *orbicularis*
- L. quadrisulcata* d'Orb. (*Cyclas* Klein, *Strigilla* Gray, *T. divaricata* Ch.) station: — ? —; habitat: Antillae; Mörch pg. 32! Guadeloupe; Beau pg. 32! Virgin-Isl.; Swift! Krebs! remark! Beau gives as synonyms *L. div.* *L.* & *L. dentata* Wood, but *L. quadrisul*

d'Orb. as a separate species; Moll. de Cuba 2—292 pl. 27 fig. 34—36, Chem. vol. 6—129 fide Mörch.

L. scabra Lmk. (*L. scobinata* Recl., *Tell. muricata* Spgl., *Cyrachæa* Leach, *Myrtea* Turt.?, *T. imbricata* Chemn.) station: dredged from 6 to 12 fathoms water in bluish sandy mud; habitat: Antillae; Mörch pg. 32! Guadeloupe; Beau pg. 23! Martinique; Cuba; Jamaica; St. Thomas; d'Orb.! Jamaica; Ad.! Virgin-Islands; R. Swift! St. Thomas; Tortola; St. Croix; Krebs!

L. semireflecta d'Orb. station: — ? — ; habitat: San Blas in Patagonia; Rio-Janeiro; Cuba; d'Orb.! remark: Moll. de Cuba 2—297. *Diploporina*

L. serrata d'Orb. (*L. Chemnitzii* Ph., *L. divaricata* Rv. non L.) station: — ? — ; habitat: St. Thomas (by J. H. Newton); Ad.! Virgin-Islands; Swift! Cuba; St. Martin; d'Orb.! Guadeloupe; Beau pg. 23! remark: Moll. de Cuba 2—295 pl. 27 fig. 37—39, Zeitschrift 1848, Conch. Icon. spec. 47. *denticulata* (Perrinitz)

L. tigerina L. (*Codakia* Scop., *Chama* Mart., *Lentilaria* Sch., *Cyth. tigerina* Lmk) station: in coarse sand between zostera on one foot water in bays and creeks; habitat: Antillae; Mörch pg. 33! Guadeloupe; Beau pag. 23! New-Orleans; Cuba; Martinique; Guadeloupe; St. Thomas; d'Orb.! Bermuda; Carthageria; Redfield! Virgin-Isl.; Swift! Jamaica; Ad.! Puertorico; St. Thomas; St. Croix; St. Johns; Carthageria; Trinidad; Krebs! remark: commonly eaten by poor people.

L. trisinuata d'Orb. station: — ? — ; habitat: Guadeloupe; Beau pg. 23! Martinique; Guadeloupe; d'Orb.! remark: Moll. de Cuba 2—300 pl. 27 fig. 46—49.

L. virgo? Reeve. station: — ? — ; habitat: Guade-

loupe; Beau pg. 23! Jamaica; Ad.! Virgin-Islands in the cabinet of H. Krebs; Swift! — *perforata*

- 5 species indetermined from the Virgin-Islands in the cabinet of R. Swift Esq. St. Thomas.
- 3 species indetermined from the West-Indies in the cabinet of H. Krebs Esq. St. Thomas.

Fimbria Mergele v. Muehlfeld.

- F. magna** Meg. (*Venus fimbriata* L., *Gafrarium* Bolton, *Corbis* Cuv., *Idotea perforata* Sch.) station: — ? — ; habitat: Antillae; Mörch pg. 33!

Diplodonto Brown 1831.

- D. semiaspera?** Ph. station: — ? — ; habitat: Virgin-Islands; Swift!
- D. species indetermined.** Virgin-Islands; Swift!

Astarteacea.

Crassatella Lmk. 1799 in prodr.

- C. antillarum** Reeve. (*C. rostrata* Lmk.?) station: — ? — ; habitat: Island of Margarita; Mörch pg. 39!
- C. martinicensis** d'Orb. station: — ? — ; habitat: Jamaica; Martinique; St. Domingo; d'Orb.! Virgin-Isl.; Swift! Krebs! remark: Moll. de Cuba 2—288 pl. 27 fig. 21—23.
- C. guadaloupensis** d'Orb. station: living on madrepores; Beau pg. 22!, under pieces of dead madrepores in gravelly sand on one foot water and dredged in bluish sandy mud from 10 to 12 fathoms water; habitat: Cuba; Guadeloupe; St. Domingo; St. Thomas; d'Orb.! Virgin-Islands; Swift! Krebs!
- C. rostrata** Lmk. station: — ? — ; habitat: West-Indies; South America; Lmk.!

Carditacea.

Cardita Bruguière 1798.

- C. domingensis* d'Orb. station: — ? — ; habitat: St. Domingo; Cuba; d'Orb.! remark: Moll. de Cuba 2—291 pl. 27 fig. 27—29.
- C. gibbosa* Rv. station: — ? — ; habitat: Florida; Yucatan; Ad.! = *floridana* (Lamarck)
- C. ovata* Ad. station: — ? — ; habitat: Jamaica; Adams! = *Venus pygmaea* St.
- C. spec. ind.* in the collection of R. Swift Esq. at St. Thomas; he observes in his list of the marine-shells of the Virgin-Islands: „may be *C. gracilis* Shuttl. who received specimens from Puertorico“, but Swift has not the diagnosis; see *Trapezium*. -3

Trapezium Megerle v. Muehlfeld.

(*Coralliophaga* Blv., *Cardita* Lmk.)

- T. dactylus* Brug. (*Cardita* Brug., *Chama coralliophaga* Ch., *C. carditoidea* Blv., *Cypricardia Hornbeckiana* d'Orb.?) station: on the lower side of madrepores; habitat: Antillae; Mörch pg. 19! St. Thomas; d'Orb.! Guadeloupe; Bean pg. 23! Virgin-Islands; Krebs! St. Thomas; Jamaica; Ad.! remark: Moll. de Cuba 2—266 pl. 26 fig. 33—34.
- T. gracilis* (*Cypricardia*) Shuttl. station: — ? — ; habitat: Guadeloupe; Beau pg. 23! remark: we have received specimens from Guadeloupe, which we are not able to distinguish from the former.

Solemyacea.

Solemya Lmk.

- S. occidentalis* D. station: dredged with bluish sandy

mud on 10 to 12 fathoms water; habitat: Crab Island; St. Thomas; Krebs! Guadeloupe; Beau! remark: Jeon. de Conch.

Arcacea.

Arca L. 1758.

- A. *Adamsi* Shuttl. station: in crevices of madrepores; habitat: Guadeloupe; Beau pg. 22! Virgin-Isl.; Swift! Tortola; St. Thomas; Porto-Plata; Krebs!
- A. *americana* Gray. (*A. pexata* Say) station: — ? — ; habitat: Guadeloupe; Beau pg. 22! Florida; Cuba; d'Orbigny!
- A. *antiquata* L. (*Anodara* Gray, *A. Deshayesii* Ph., *A. hemidermos* Ph.) station: small specimens fastened on madrepores and old specimens loose in blue sandy mud on 10 to 12 fathoms water; habitat: Cuba; Martinique; d'Orb.! Jamaica; Cuba; Carthagenia; Ad.! Virgin-Islands; Swift! Krebs!
- A. *auriculata* Lmk. station: — ? — ; habitat: St. Thomas; Cuba; d'Orb.!
- A. *barbadensis* Petiver station: — ? — ; habitat: Cuba; St. Lucy; Martinique; St. Thomas; d'Orb.!
- A. *brasiliensis* Lmk. (*Scapharca* Gray) station: — ? — ; habitat: St. Croix; Mörch pg. 41! Guadeloupe; Beau pg. 22! Rio Janeiro; Martinique; Guadeloupe; Cuba; d'Orbigny!
- A. *candida* Helbling. (*Lithoarca* Gray, *A. ovata* Gmelin., *A. trapizina* Lmk., *A. Helbingii* Brug.) station: fastened on the lower side of madrepores; habitat: Antillae; Mörch pg. 40! Guadeloupe; Beau pg. 22! Florida; Martinique; Cuba; d'Orb.! Jamaica; St. Thomas; Ad.! Virgin-Isl.; Swift! Carthagenia; Jamaica; St. Thomas; Tor-

tola; Krebs! remark: Deshayes says in a note in Lmk. an. sans vert. vol. 6 pg. 469: this species should be retained as *A. nivea* Chem. vol. 7 pg. 538.

- A. incongrua* Say. station: — ? — ; habitat: Navy-Bay; Florida; South Carolina; Carthagenae; Ad.! Carthagenae?; Krebs!
- A. indica* Gml. station: — ? — ; habitat: Guadeloupe; Beau pg. 22! Virgin-Isl.; Swift!
- A. labiata* Sowb. station: — ? — ; habitat: Guadeloupe; Beau pg. 22!
- A. lactea* L. station: — ? — ; habitat: Guadeloupe; Beau pg. 22! Natal Coast; Jamaica; Sicily; St. Thomas; Ad.!
- A. Listeri* Ph. (*Barbatia* Gray, *A. fusca* Brug, *A. granulata* Meusch.) station: — ? — ; habitat: Antillae; Mörch pg. 40! Cuba; Martinique; Guadeloupe; d'Orb.! St. Thomas; Jamaica; Ad.! Virgin-Islands; Swift! Kr.!
- A. noae* L. station: in 10 fathoms water in mud, 6—7 attached together and in crevices of large pieces of madrepores; habitat: Guadeloupe; Beau pg. 22! Tortola; St. Thomas; Jamaica; Carthagenae; Bermuda; Ad.! Virgin-Islands very common; Krebs! remark: C. B. Ads says it is synonyme with *A. occidentalis* Phil.
- A. notabilis* Bolt. (*Anodara* Gray, *A. Deshayesii* Hanley) station: — ? — ; habitat: Antillae; Mörch pg. 41! Guadeloupe; Beau pg. 22! Virgin-Islands; Swift!
- A. occidentalis* Ph. (*Cilota* Browne, *Daphne* Poli, *Navicula* Blv.) station: as *A. noae*; habitat: Virgin-Isl.; Swift! Antillae; Mörch pg. 39! remark: see *A. noae* L.
- A. rhombea* Born. (*A. Chemnitzii* Phil., *Scapharca*) station: in crevices of corals, at the root of gorgoniae and in mud on 10 fathoms water; habitat: Antillae; Mörch

pg. 41! Virgin-Islands; Swift! remark: Chem. vol. 7 fig. 553—56, Zeitsch. 1851 pg. 50.

- A. *sqvamosa* Lmk. (A. *gradata* Brod., A. *donaciformis*, Rv., A. *divaricata* Sowb., A. *domingensis* Lmk., *Daphnoderma* Mörch, A. *angulata* Meusch.) station: below madrepores; habitat: Antillae; Mörch! Guadeloupe; Beau pg. 22! St. Domingo; Lmk.! St. Thomas; St. Johns; Ad.! Virgin-Islands; Swift! Krebs!
- A. *umbonata* Lmk. (Cilota Brown, A. *ventricosa* Lmk. according to C. B. Ad.! A. *imbricata* Brug; A. *mutabilis* Reeve, A. *americana* d'Orb. non Gray) station: fastened to madrepores and in the roots of gorgoniae; habitat: Antillae; Mörch pg. 40! Guadeloupe; Beau pg. 22! Ad.! Virgin-Islands; Swift! Chagres; St. Thomas; Antigua; Jamaica; Carthagenae; Ad.! St. Thomas; St. Croix; Tortola; Krebs! remark: Adams separates A. *umb.* from A. *imbricata*; Moll. de Cuba 2—317 pl. 28 fig. 121

Pectunculus Lmk. (*Axinaea* Poli, *Glycimeris* da Costa)
station: we are not able to give the station of any of the *Pectunculi*.

- P. *angulatus* Lmk. spec. 8 (*Area angulosa* Gml.) habitat: Tortola; Swift!
- P. *castaneus* Lmk. non Reeve. (P. *variegatus* Chem.) habitat: Guadeloupe; Beau pg. 22! St. Johns; Jamaica; Ad.!
- P. *castaneus* Reeve non Lmk.; habitat: Tortola; Swift! Krebs! remark: perhaps Adams habitat refers to this species.
- P. *decussatus* L. (P. *pennaceus* Lmk. spec. No 6) habitat: Virgin-Islands; Swift! Tortola; Krebs!

- P. hirtus* Phil. habitat: Cumana; Philippi Zeitsch. 1846 pg. 191! Isl. of Margaritha; Swift! West-Indies; Krebs!
- P. minimus* Gml. habitat: St. Thomas; Riise! remark: the name derives from Mörch but the specimens are by us considered the young of *P. pectinatus* Gm.
- P. oculatus* Reeve. spec. 38 (*P. pectiniformis* Lmk. acc. to d'Orbigny) habitat: West-Indies; Reeve! Tortola; Swift! Krebs! Martinique; d'Orb.! Guadeloupe; Beau pg. 22! remark: Reeve's description does not satisfy us.
- P. pectenoides* Desh. habitat: Tortola Swift!
- P. pectinatus* Gml. (*Area* Gmel., *A. costata* Meusch., *A. pectunculus minor* Ch., *P. minor* d'Orb.) habitat: Isl. la Plata; Reeve! St. Thomas; Riise! Krebs! Tortola; Carthagera (by Krebs); Ad.! Guadeloupe; Beau pg. 22! Tortola; Swift! Krebs! remark: Riise has d'Orb.'s name from Mörch.
- P. scriptus* Lmk. habitat: St. Domingo; Lmk.
- P. sericatus* Reeve spec. 49; habitat: Tortola; Reeve! Swift! Krebs! St. Thomas; Krebs! St. Johns; Ad.! Guadeloupe; Beau pg. 22!
- P. spadiceus* Reeve. habitat: — ? — ; Reeve! St. Juan del Norte de Nicaragua; Krebs!
- P. undatus* L. (*P. undulatus* Lmk. spec. 3, *P. lineatus* Reeve fide Mörch pg. 42) habitat: Virgin-Isl.; Swift! St. Thomas; Tortola; Krebs! Guadeloupe; Beau pg. 22!
- P. spec. ind* No 4. habitat: Riise has 2 specimens from Crabisland; Krebs has one valve from St. Martin; remark: it is inequilateral and approaching *P. spadiceus*, but mostly white with a few brown specks; diameter from 4 to 6 lines.
- P. spec. ind.* No 2. habitat: Porto-Plata; St. Thomes;

Krebs! remark: two perfect specimens and two odd valves; perhaps young undatus or decussatus or hirtus; remark: are *P. hirtus* Phil. and *P. undatus* L. and *P. decussatus* L. & *angulatus* L. not synonymes?

Heteromyaria.

Mytilacea.

Mytilus L.

- M. exustus* L. (*M. Domingensis* Lmk., d'Orb., *Hormomya* Mörch) station: on rocks which are over low-water mark; habitat: Antillae; Mörch pg. 53! Guadeloupe; Beau pg. 22! Rio Janeiro; Cuba; Martinique; d'Orb.! Jamaica; St. Thomas; Ad.! Virgin-Islands; Swift; Carthage; St. Thomas; St. Johns; Krebs! remark: it envelopes itself very differently in places where the sea is smooth and in places where the sea is rough.
- M. Lavelleanus* d'Orb. station: — ? — ; } habitat: Guadeloupe? d'Orb.! remark: Moll. de Cuba 2—328 pl. 28 fig. 3—5.
- M. viator* d'Orb. station: — ? -- ; habitat: Patagonia; Cuba; d'Orb.!

Modiolus Lmk.

- M. americana* Tarvart d'Herbigny (*M. amer.* Leach?, *M. tulipa* Lmk., *Perna* Adans., *Volsella* Scop., *Pholas* Kl.) station: between seaweed on gravel in a few feet water in creeks; habitat: Antillae; Mörch pg. 53! Guadeloupe; Beau pg. 22! Cuba; Martinique; Guadeloupe; St. Lucy; d'Orb.! Jamaica; Bermuda; Ad.! Virgin-Isl.; Swift! all the West-Indian islands from Trinidad to St. Domingo; Krebs!

- M. barbata*? L. station: — ? — ; habitat: North-Bedforth; Jamaica; St. Thomas; d'Orb.!
- M. castanea* Say. station: between sea-weed in coarse sand close to the shore; habitat: South-Carolina; Jamaica; St. Thomas; Ad. Florida; on a specimen in the Academy of N. Sc. in Philadelphia! Virgin-Isl.; Swift! St. Thomas; St. Croix; Tortola; Krebs!
- M. litoralis*? Say. station: — ? — ; habitat: Virgin-Islands; Swift!
- M. modiolus* L. (*Arca* L., *M. citrinus polydentalus* Ch., *Tamarindiformis striatus* Meusch., *M. citrinus* Bolt., *M. sulcatus* Lmk., *M. exustus* Gml., d'Orb. non L., *M. Chemnitzii* Pot. & Mich., *Brachydontes* Sws.) station: on rocks above low-water-mark; habitat: Antillae; Mörch pg. 54! Guadeloupe; Beau pg. 22! Cuba; d'Orb.! Virgin-Islands; Swift! Krebs!

Lithophagus Megerle v. Muchfeld.

- L. bisulcatus* d'Orb. (*Mod. appendiculatus* Ph.) station: boring in life madrepores; habitat: Antillae; Mörch pg. 56! Guadeloupe; Beau pg. 22! Cuba; Jamaica; Martinique; Guadeloupe; St. Domingo; d'Orb.! Virgin-Islands; Swift! Krebs! remark: Moll. de Cuba 2—333 pl. 28 fig. 14—16; we are in possession of several very interesting specimens, where the animal has been obliged 5—6 times to form concave bases to rest on to be able to keep communication with the sea and to prevent to be overgrown by the madrepores; they are consequently able to loose the byssus and to refasten it as they are moving upward; it will be understood that thereby is formed a wedgeformed hole from the centre of the madre-pore toward the surface as the animal of the Litho-

phagus increases in size and the hole is intersected at intervals by the above mentioned, on the upper side concave, on the lower side convex, bases.

- L. caribæus* Ph. station: — ? — ; habitat: Antillae; Mörch pg. 56!
- L. dactylus* Sow. (*Lithodomus* Cuv., *L. antillarum* d'Orb., *L. corrugata* Phil.) station: in life madrepores; habitat: Antillae; Mörch pg. 55! Virgin-Isl.; Swift! Cuba; Martinique; Guadeloupe; d'Orb.! remark: Moll. de Cuba 2—332 pl. 28 fig. 12—13.
- L. fuscus* Gml. (*Mytilus brunneus* Soldr., *Mod. cinnamomea* var. Lmk., *Mod. Favannii* Pot. & Mich., *Botula* Mörch) station: in life madrepores; habitat: Antillae; Mörch pg. 55! Guadeloupe; Beau pg. 22! Virgin-Islands; Swift! Cuba; Martinique; St. Domingo; Jamaica; d'Orb.! St. Thomas; Jamaica; Lmk.! St. Thomas; St. Croix; Tortola; Krebs!
- L. niger* Lister. (*Pholas* Lister, *L. antillarum* Phil.) station: in madrepores; habitat: Antillae; Mörch pg. 56! Guadeloupe; Beau pg. 22! Cuba; Martinique; St. Domingo; St. Lucy; d'Orbig.! Virgin-Islands; Swift! Krebs! remark: Moll. de Cuba 2—332 pl. 28 fig. 10—11; the *Lethophagi* are only found alive in the madrepores as long as these are alive.

Dreisenia Beneden.

- D. Rissei* ? ; station: in brackish water between the roots of *Avicennia*; habitat: Crabisland; Riise! Porto-Plata; Krebs!

Pinnacea.

Pinna L.

- P. muricata* L. (*P. nobilis* Ch., *P. seminuda* Lmk., Pen-

naria Browne, *P. rigida* Sold.) station: in mud and sand on 1 to 3 feet water in smooth water; habitat: Cuba; Martinique; Guadeloupe; d'Orb.! Virgin-Islands; Swift! Jamaica; St. Thomas; Carthageua; Ad.! Krebs!

P. pernula Ch. (*P. haud ignobilis* Ch., *P. degenera* Link., *P. carnea* Gm., *P. flabellum* Lmk., *P. varicosa* (Lmk.) d'Orb., *Chimaera* Poli) station: as the former; habitat: Antillae; Mörch pg. 51! Guadeloupe; Beau pg. 22! Cuba; Martinique; Guadeloupe; Trinidad; St. Croix; d'Orb.! Jamaica; Tortola; St. Thomas; Ad.! Krebs! Virgin-Islands; Swift!

Monomyaria.

Maleacea.

Avicula Bruguière.

- A. Candeara* d'Orb. (*Malleus* Link.) station: — ? — ; habitat: Cuba; de Candé! Guadeloupe; Beau pg. 21! remark: Moll. de Cuba 2—343 pl. 28 fig. 25—27.
- A. colymbus* Bolton. (*A. Pteria* Scop., *Glaucus* Poli, *Anonica* Oken, *M. Hirundo* Ch., *A. atlandica* Lmk., *A. communis* Lmk.) station: at the root of *Gorgoniae*; habitat: Antillae; Mörch pg. 51! Jamaica; St. Thomas; Tortola; Ad.! Virgin-Islands; Swift!
- A. crocata* Swains. station: — ? — ; habitat: Jamaica; St. Johns; St. Thomas; Carthageua; Tortola; Ad.!
- A. guadelupensis* d'Orb. station: — ? — ; habitat: Guadeloupe; Beau pg. 21! d'Orb.! remark: Moll. de Cuba 2—343 pl. 28 fig. 23—24.
- A. heteroptera* Lmk. station: — ? — ; habitat: Antillae; Mörch pg. 51!
- A. longisquamosa* Dunk. station: — ? — ; habitat: Guadeloupe; Beau pg. 21!

A. macroptera Lmk. station: — ? — ; habitat: Guadeloupe; Beau pg. 21! Virgin-Islands; Swift!

A. radiata Leach. (*Margaritifera* Browne; *Meleagrina* Lmk., *A. guadeloupensis* d'Orb. according to Beau pg. 21) station: — ? — ; habitat: Antillae; Mörch pg. 50!

Melina Retz. (*Isognomum* Kl., *Isogonum* Bolt., *Perna* Lmk.).

M. alata Gm. (*Ostrea* Gm., *Sutura* Meg., *Ala corvi pendula* Ch., *Riparia isogoalata* Meuschen., *Perna obliqua* Lmk.) station: in large clusters in creeks with stones and smooth water; habitat: Antillae; Mörch pg. 49! Guadeloupe; Beau pg. 22! Cuba; Martinique; d'Orb.! Jamaica; Ad.! Virgin-Islands; Swift! Krebs! Carthagenia; Krebs! remark: is eatable.

M. bicolor Ad. station: — ? — ; habitat: Jamaica a St. Johns; St. Thomas; Ad.!

M. perna L. (*Concha semiaurita* Ch., *O. canterinata* Meusch., *M. perniformis* Retz, *Perna sulcata* Lmk., *P. vulcella* b Lmk., *P. Linnaei* Pfr., *P. Lamarekiana* d'Orb.) station: Antillae; Mörch pg. 49 — 50! Guadeloupe; Beau pg. 22! Martinique; de Candé!

M. vulsella Lmk. *P. Lamarekiana* d'Orb. accor. to Swift, *Concha semiaurita* var. Ch., *P. Chemnitziana* d'Orbigny) station: on rocks at the level of the sea; habitat: Cuba; Martinique; St. Croix; d'Orb.! Virgin-Isl.; Swift! Krebs! Carthagenia; Krebs!

Pectinea.

Lima Bruguière 1792.

L. bullata Born. (*Limatula* S. Wood) station: — ? — ; habitat: Antillae?; Mörch pg. 57! remark: see *L. scabra* Born.

- L. caribæa* d'Orb. (*Radula* Kl.) station: — ? — ; habitat: Antillae; Mörch pg. 57! Cuba; Auber! remark: Moll. de Cuba 2—337 pl. 28 fig. 17—19.
- L. cubaniana* d'Orb. station: — ? — ; habitat: Cuba; Auber! remark: Moll. de Cuba 2—337 pl. 28 fig. 20—22.
- L. inflata* Lmk. (*L. pellucida* Ad.) station: in small caves in the sand between stones on 1 to 2 feet water; this species, as well as *L. tenera* & *L. scabra*, moves quick about by opening and shutting the valves; habitat: Jamaica; St. Thomas; Guadeloupe; Ad.! St. Thomas; St. Martin; St. Barth; Krebs!
- L. multicostata* Sowb. station: — ? — ; habitat: Florida; Jamaica; St. Thomas; St. Johns; Ad.! remark: Adams says: generally confounded with the mediterranean species, which is the *L. squamosa* Lmk.
- L. scabra* Born (*Ctenoides* Klein, *L. aspera* Ch., *L. glacialis* Gm. & Lmk., *Limaria asperula* Link., Swift gives as synonymes: *O. bullata* Born. & *L. fragilis* Lmk.) station: see *L. inflata*; habitat: Antillae; Mörch pg. 57! Guadeloupe; Beau pg. 21! Cuba; Martinique; St. Lucy; d'Orb.! Jamaica; St. Thomas; Porto-Cabello; St. Johns; Ad.! Virgin-Islands; Swift! Krebs!
- L. squamosa*? Lmk. (see *L. multicostata* Sowb.) station: — ? — ; habitat: Guadeloupe; Beau pg. 41!
- L. tenera* Ch. (*Ctenoides* Klein, *Ost. glacialis* Solis, *Limaria glacialis* Lmk.) station: as *L. inflata* Lmk.; habitat: Antillae; Mörch pg. 57! Carthagen; Jamaica; Ad.! Virgin-Islands; Krebs!
- Pecten O. Fried. Mueller (Rondelet 1606.)
- P. antillarum* d'Orb. station: — ? — ; habitat: Guadeloupe; Beau pg. 21!

- P. exasperatus* Sowb. station: — ? — ; habitat: Antillae; Mörch pg. 59! Virgin-Islands; Swift!
- P. fucatus* Reeve. station: — ? — ; habitat: Virgin-Isl.; Swift!
- P. gibbus* Lmk. station: — ? — ; habitat: Cuba; Martinique; Guadeloupe; d'Orb.!
- P. imbricatus* Gm. (*Ostrea* Gm., *Pera venatoria imbricata* Ch.) station: — ? — ; habitat: Antillae; Mörch pg. 58! St. Thomas; St. Johns; Ad.! Virgin-Islands; Swift! St. Croix; Trinidad; Tortola; Krebs!
- P. nodosus* L. (*C. coralinus* Ch. acc. to Beau) station: — ? — ; habitat: Antillae; Mörch pg. 58! Guadeloupe; Beau pg. 21! Cuba; Guadeloupe; Martinique; d'Orb.! St. Thomas; Bermuda; St. Johns; Ad.! Virgin-Islands; Swift! Krebs!
- P. nucleus* Born. (*P. turgidus* Lmk., Argus Gray) station: — ? — ; habitat: Antillae; Mörch pg. 59! Guadeloupe; Beau pg. 21! Virgin-Islands; Swift!
- P. ornatus* Lmk (*Pectinium* Link., *Chlamys* Bolton) station: — ? — ; habitat: Antillae; Mörch pg. 57! Guadeloupe; Beau pg. 21! Florida; Cuba; St. Lucy; d'Orb.! Jamaica; St. Johns; Ad.! St. Thomas; Carupana; Tortola; Krebs! Virgin-Islands; Swift!
- P. Sowerbii* Guild. station: — ? — ; habitat: Florida; Jamaica; St. Thomas; Carthagera; Ad.!
- P. sulcatus* Lmk. No 21 (*P. antillarum* Recluz) station: — ? — ; habitat: Virgin-Islands; Swift!
- P. ziczac* L. (*Vola* Klein, *Pandora* Meg., *Janeira* Sch.) station: — ? — ; habitat: Antillae; Mörch pg. 59! Guadeloupe; Beau pg. 21! Cuba; Martinique; Guadeloupe; St. Lucy; St. Domingo; St. Thomas; Jamaica;

d'Orb.! Tortola; Jamaica; Carthageria; Ad.! Virgin-Isl.;
Swift! Krebs!

Spondylus L. (Lister 1686.)

- S. americanus* Lmk. No 2 (*S. folia brassicae* Ch., *S. gaederopus* Ch. acc. to d'Orb.) station: — ? — ; habitat: St. Domingo; Lmk.! Guadeloupe; Beau pg. 21! St. Domingo; Cuba; Martinique; d'Orb.! Virgin-Islands; Swift!
- S. coccineus* Lmk. station: — ? — ; habitat: Guadeloupe; Beau pg. 21! remark: perhaps synonyme with *S. fimbr.* M.
- S. echinatus* d'Orb. station: — ? — ; habitat: Cuba; Martinique; Guadeloupe; d'Orb.! Virgin-Islands; Swift!
- S. fimbriatus* Meusch. (*Ostrea spondyloides* Gm., *S. spatuliferus* Lmk. No 13, *S. divaricatus* Schrit.) station: on rocks in 7 to 10 feet water where the surf is strong; habitat: Bermuda; St. Thomas; Carupano; Ad.! Virgin-Islands; Swift! Krebs!
- S. radians* Lmk. No 19 station: — ? — ; habitat: Virgin-Isl.; Swift!
- S. striato-spinosus* Ch. station: — ? — ; habitat: Jamaica; St. Thomas; Ad.! remark: is this not our *S. fimb.* M.?

Plicatula Lmk.

- P. depressa?* Lmk. station: — ? — ; habitat: Virgin-Islands; Swift!
- P. ramosa* Lmk. station: — ? — ; habitat: Guadeloupe; Beau pg. 21!
- P. spondyloidea* Meusch. (*Ostrea* Meusch., *Sp. plicatus* Ch., *P. cristata* Lmk., *P. reniformis* Linkn., *P. barbades*

sis (Petiver) d'Orb.) station: on rocks in 7—10 feet water; habitat: Guadeloupe; Beau pg. 21! Antilles Rio-Janeiro; Patagonia d'Orb.! Jamaica; Florida; St. Thomas; Ad.! Virgin-Islands; Swift! Krebs!

Ostreacea.

Ostrea L. 1768.

- O. arborea Ch. (Mulleria Fer., O. radicum Ch., O. rhizophorae Ch., O. parasitica Gm.) station: in lagoons and creeks on the root of Avicennia and Rhizophora; habitat: Antillae; Mörch pg. 62! Cuba; Jamaica; Ad.! Guadeloupe; Beau pg. 21! Virgin-Islands; Swift!
- O. cristata Born. station: — ? — ; habitat: Guadeloupe; Beau pg. 21!
- O. frons L. Ch. (O. transtra Meusch., rubella Lmk., limacella Lmk., erucella Lmk., folium (L.) d'Orb.) station: on the roots of trees in lagoons and creeks; habitat: Cuba; Martinique; Guadeloupe; St. Lucy; St. Domingo; New-Orleans; d'Orb.! Guadeloupe; Beau pg. 21! Jamaica; Tortola; Ad.! Virgin-Islands; Swift! Krebs!
- O. rostralis Lmk. station: — ? — ; habitat: Guadeloupe; Beau pg. 21!
- O. spreta d'Orb. station: — ? — ; habitat: Rio-Janeiro; Cuba; d'Orb.! remark: moll. de Cuba 2—365 pl. 28 fig. 30.
- O. spec. ind. No 1 from St. Croix and Guadeloupe, in the cabinets of R. Swift & H. Krebs; Swift says: it approaches O. cucullata Born in Lmk. an. sans vert. No 34.

Placunanomia Broderip 1832.

- P. echinata? Brod. (P. abdominalis Gray) station: between the branches of corals; habitat: Guadeloupe; Beau pg. 21! Virgin-Islands; Swift!

P. rudis Brod. station: nests especially in the dead shells of *Pinna muricata*; habitat: Virgin-Islands; Swift! Krebs!

Anomia L. 1767.

A. simplex d'Orb. station: — ? — ; habitat: Guadeloupe; Beau pg. 21! Martinique; Cuba; d'Orb.! Virgin-Islands; Swift: remark: Moll. de Cuba 2—367 pl. 28 pg. 31—33.

Copy of a letter, from the Author.

Copenhagen.

Ole Fuhr's Gade 24-1

1st December, 1884.

Messrs. R. Friedländer & Sohn,
Berlin.

Dear Sirs,

In reply of your favor dated 29th last month I beg to inform you, that "the West-Indian Marine Shells, with some remarks", 1864, was only printed in 26 copies, of which 3 were - according to the law - delivered to the public Libraries, 7 were lost in transmitting them to St. Thomas, 3 went to the Universities of Sweden and Norway, and a few given to friends.

The work is of very little value, and it was only printed with the view to get it supplied with notes from those friends to whom it was circulated.

I have a copy, with many notes in, and a copy, which consists of the proof-sheets from

the printer. These last are very little
manipulated and as good as a clean copy.

Consequently there are none for
sale; my friends tease me that the book
is the costliest they know; on account of
a copy has been sold in Altona at
auction for 10 R_s, it is only 137 pages;
but if it is a scientific gentleman, who
wishes the use of it for some months,
I might be willing to lend him it against
receipt.

At the same time I have this
opportunity to write I beg to ask if you
are acquainted with a gentleman, who
interests himself in Genealogy, and who
would permit me to address him about
some German families.

Dear Sir!

Your obedient servant,
H. Krebs.

SMITHSONIAN INSTITUTION LIBRARIES

3 9088 00604 3236