QL
684
I 3M4X
Birds

WILD BIRDS
IN
CITY PARKS

WALTER

EIGHTEENTH THOUSAND

allensis

.35

Wild Birds in City Parks

QL

684

I 3W4X

BIRDS

Being hints on identifying 200 birds,
prepared primarily for the spring
migration in Lincoln Park, Chicago,
but adapted to other localities

BY

HERBERT EUGENE WALTER

AND

ALICE HALL WALTER

FOURTH ENLARGED REVISION

With Chart and Key

AUTHORS' EDITION

1910

Copyright 1901, 1902, 1904, 1910.
BY ALICE HALL WALTER.

Contents.

	Page
Note to Fourth Revision.....	5
Preface.....	7
General Hints	9
Particular Hints	12
Water and Shore Birds	13
Grebes and Loons	13
Gulls and Terns	14
Ducks, Geese and Swans.....	15
Hérons and Bitterns	19
Rails and Coots	20
Plover and Sandpipers	22
Land Birds	25
Bobwhite and Grouse	25
Doves	25
Birds of Prey	26
Cuckoos and Kingfishers.....	29
Woodpeckers.....	29
Swifts, Hummingbirds, etc.....	30
Perching Birds	31
Key to Land Birds	53
Table of Occurrence in Lincoln Park, Chicago	60
Table of Arrival in Lincoln Park, Chicago	64
Table of Distribution and Occurrence	71
Index to Common Names.....	89

“The first sparrow of spring ! The year beginning with younger hope than ever ! The faint silvery warblings heard over the partially bare and moist fields from the bluebird, the song sparrow and the red-wing, as if the last flakes of winter tinkled as they fell.”

—*Thoreau's Walden.*

Note to the Fourth Edition.

A springtime has not yet passed without the arrival of the migrating birds. Upon this attractive subject an increasing number of new books appear every year, yet it seems probable from the way in which successive printings of this little book have been exhausted that there still exists a need for a simple, compact field manual at a minimum price.

The present edition is considerably enlarged by descriptions of over fifty additional birds, a revised key and a new table of comparative distribution. The entire subject matter also has undergone a thorough revision with the end in view of making the book useful in all localities in north-eastern United States from the Mississippi to the Atlantic coast.

The authors wish to thank all those who have joined hands with them in the past and hope in the name of the birds that the circle may grow still larger in the days to come.

"All nature is so full that that district produces the greatest variety which is most examined."

—*Gilbert White*, 1768.

Preface.

The object of this little book is to furnish those who may be interested in making the acquaintance of wild birds with a simple letter of introduction to 200 birds, the majority of which are commonly seen during the spring migration.

Complete descriptions have been avoided, in the belief that the student should rely upon his own observation for the discovery of minor details. *The living bird* is the one important fact which will make the brief hints offered of value.

Anyone caring to make use of these hints may be assured that during the migrations of the birds city dwellers have one of the keenest delights of country life brought to their very doors, because many birds, migrating largely at night, are attracted by the lights of the city and stop off in their long journey to feed, so that a city park often contains a greater variety of feathered visitors than an equal area in the country.

We wish to remind those of our friends who have asked for pictures in a future edition that every springtime this book is copiously illustrated in our parks and around our homes by hundreds of living birds and that these illustrations are all life size, absolutely accurate in detail and colored true to nature.

“As for the birds * * * they add immeasurably to the wholesome beauty of life.”

—*Theodore Roosevelt, in Bird Lore, Vol. II, p. 98.*

General Hints.

"A good observer is quick to take a hint and follow it up."—*John Burroughs*.

The identification of birds depends quite as much upon accurate observation of their size, motions, flight, characteristic attitudes, manner of feeding, company, song, call-notes and haunts, as upon details of form and color. Especial care is necessary to insure correct estimates of size for the reason that living birds often appear smaller to the unpracticed eye than they actually are. The familiar *English Sparrow* is a convenient standard of size because it is usually at hand either in city or country for instant reference. Remembering that it is 6 inches long a practical though rough division of wild birds may be made as follows:—

1. Birds smaller than the *English Sparrow*.
2. Birds about the size of the *English Sparrow*.
3. Birds decidedly larger than the *English Sparrow*.

If a few general characteristics of the common bird-families be kept in mind, and these are quickly and almost unconsciously learned, the identity of a strange bird may usually be narrowed down to a few possibilities. For example:

Woodpeckers climb up and down the trunks of trees bracing with their tails and tapping the bark vigorously;

Nuthatches are smaller than woodpeckers and have much the same habit of climbing up and down tree-trunks but with a freer wig-wagging motion, often descending head downward;

Flycatchers sit erect with drooping tails, watching alertly for insect prey upon which they pounce in mid air, afterwards returning to their perch;

Swallows skim through the air in graceful and long sustained flights;

Sparrows have stout seed-cracking bills, feed upon the ground, seldom fly high or far at a time and are for the most part fine songsters;

Warblers are tiny, tireless, gaily-colored explorers of the twigs of trees and bushes;

Kinglets are smaller than warblers and quite as restless in their motions, but arrive earlier in the migration;

Wrens, with tails erect, slip mouse-like about brush heaps, crevices and bushes, though often perching in sight while singing;

Thrushes, who with the exception of the *Robin* and *Bluebird* are very plainly dressed, run about on the ground stopping suddenly in a listening attitude. When singing they fly up to some perch, although many of the unrivaled singers of this family are silent during their brief sojourn in city parks;

Vireos are most at home on the boughs of trees and sing freely as they glide in and out among the leaves to feed.

Female birds can usually be identified after the adult males have been seen, although the females and young of many species are obscurely marked or quite different from the adult males, a fact especially true of the warblers. Immature birds are not considered in the present scheme of

study since they are a source of confusion to the beginner and occur in any considerable number only during the fall migration.

The time of arrival, that is, when a bird may be expected during the migration, is a point worthy of particular attention. Many wild guesses may be avoided by simply noticing the dates when a bird has been known to arrive in any given locality from year to year.

By eliminating all birds whose arrival is decidedly earlier or later than that of the one in question, the matter is usually narrowed down to a few species whose identification is thus made much easier.

The safest way, however, is to make careful notes about a stranger on the spot. Memory is never more treacherous than in the case of the description of a doubtful bird when one appeals to some book of reference, a museum collection or the judgment of others. It is not advisable to attempt extensive "field-notes" at first but *it is essential* to keep a daily, dated list of all identified birds. A blank chart for this purpose is inserted in the back part of the book.

Never *chase* birds. Have patience, stand still a great deal and use your common sense all the time.

Do not make yourself believe that you see a certain bird because it has been reported by others. Do your own looking and listening and do it well.

If you cannot go birding alone take along as few friends as possible because birds are suspicious of human beings in flocks, especially when they move about much.

A pair of field or opera-glasses is a valuable aid, although practice is necessary in learning to adjust them rapidly and to fix them instantly upon the bird. Do not feel obliged to use the glasses when you are near enough to see well without them.

Keep the sun at your back, otherwise colors will deceive you.

Remember that birds do not stay "indoors" on account of rain, clouds or unfavorable weather. Warm spring rains literally fill the parks with feathered travelers who often tarry but a day.

Particular Hints.

For the convenience of those desiring access to more detailed sources of information the scientific name of each bird is placed in parenthesis below its common name, while prefixed to the common name is its number in the official list of the American Ornithologist's Union as revised to date.

Following the hints given about each bird are the names (in **black-faced type**) of any bird or birds for which it might easily be mistaken.

If information is desired as to the probable date of arrival of any bird mentioned, reference should be made to the table on page 65.

The table beginning on page 74 indicates the birds which may be expected in given localities in northeastern United States.

Part 1.

WATER AND SHORE BIRDS.

3. HORNED GREBE. 14 in.

(Colymbus auritus.)

Sooty black, extending in narrow line up *back of neck*; rest of neck, sides and upper breast rich chestnut; glossy white below; head and chin black; buffy "horns" or crests, slanting backward from the eyes; slender black bill, tipped with yellow; eye, red; no tail. WINTER: black parts sooty; brown replaced by white; grayer below; crests and ruff less conspicuous. **Pied-billed Grebe.**

6. PIED-BILLED GREBE. 13½ in.

(Podilymbus podiceps.)

Brownish-black, showing gray on head and neck; belly, dusky white, otherwise brownish below; black throat-patch; a conspicuous black band across bill. WINTER: throat whitish; browner below; no band on bill. **Horned Grebe.**

7. LOON. 33 in.

(Gavia imber.)

Black, showing greenish on head and neck, spotted with square or oval patches of white on back and wings and striped with white in front of wings; white below; two conspicuous white-striped bars across the black neck giving the effect of a *broad black collar*; tail, very short; eye, red; large black bill, long and pointed; dives and sinks like the *Grebes*; note, an eerie, prolonged cry.

51. HERRING GULL. 24 in.

(*Larus argentatus*.)

White; wings and back, pearly blue-gray; bill mostly yellow; outer wing feathers partly black, marked and *tipped with white*; the closed wing shows black towards the end, marked with a regular line of white spots and tipped with white at the extreme point; looks large and heavy in flight. WINTER: Streaked on head and neck with gray. IMMATURE: Dark, much streaked with brownish; bill darker. **Ring-billed Gull. Kittiwake.**

54. RING-BILLED GULL. 19 in.

(*Larus delawarensis*.)

Similar to 51, but *smaller*; bill, greenish-yellow, bright at tip, *banded with black around the middle*; wing tipped at extreme point with black; feet greenish-yellow; eyelids vermilion-red. **Herring Gull.**

60. BONAPARTE GULL. 13 in.

(*Larus philadelphia*.)

Back and wings, light pearl-gray; head and throat, dark slate color; back of head, neck, underparts and square tail, white; wings tipped, and *bordered narrowly on the outside edge with black*; feet and legs, red; bill black. WINTER: hood, grayish white. **Common Tern.**

70. COMMON TERN. 15 in.

(*Sterna hirundo*.)

Pearly gray back and wings, whiter on rump; *top of head, shining black*; pure white on throat; dusky-white below; tail forked, *outer edge darkest*; bill long and red, blackening towards tip, held at right angles to the body in flight; feet reddish. WINTER: front of head and under-parts, white; bill nearly black. **Bonaparte Gull.**

129. AMERICAN MERGANSER. 25 in.

(*Mergus americanus*.)

Black; rump and tail, ashy; head, throat and upper part of neck, greenish; wing, *largely white*, edged with black and crossed by a black bar; white below, showing salmon tinge in flight; narrow white collar; long red bill tipped with black; eye, red. FEMALE: ashy-gray; *head and neck, brown*; salmon or brownish tinge across upper breast; *throat, white*; wings dark with a white patch; small crest on back of neck; feet orange; eye yellow. **American Golden-eye. Red-breasted Merganser.**

130. RED-BREASTED MERGANSER. 22 in.

(*Mergus serrator*.)

Long ragged crest; head and neck, black; broad white collar; broad cinnamon band streaked with black across breast, otherwise white below, showing salmon tinge in flight; wing largely white, edged and barred twice with black; eye and bill, red. FEMALE: back and wings dark gray turning to brown on head and neck; much paler on sides of neck and throat; gray across breast; crest small; white wing-patch. **American Merganser.**

131. HOODED MERGANSER. 18 in.

(*Lophodytes cucullatus*.)

Black, including throat and neck; *large circular crest, white bordered with black*; white below running up in front of the wings in two points; sides brownish, finely lined with black; white wing-patch, crossed by black bar; lengthwise white streaks on end of wings; *short black bill*; eye yellow. FEMALE: grayish-brown; throat white; crest small; sides unmarked. **Buffle-head.**

132. MALLARD. 23 in.

(*Anas platyrhynchos*.)

Gray-brown head, throat and neck glossy green to blue; narrow white collar; breast, rich brown; belly, gray lined with black; tail white, set off by black feathers curling up from either side of black rump; wing-patch purple, bordered on either side with a black and then a white bar. FEMALE: buffy-brown and black; much streaked and speckled; lighter on throat; shows wing-patch as in male.

133. BLACK DUCK. 22 in.

(*Anas rubripes*.)

Dark brown mottled with rusty, lighter below; wing-bar violet-purple edged with black; throat with few markings; legs brownish; bill greenish-black; *in flight white wing lining shows*; a suspicious pond and swamp duck often active at night; call, "quack". **Female Mallard.**

139. GREEN-WINGED TEAL. 14½ in.

(*Nettion carolinensis*.)

Gray, sides and shoulders finely lined with black; *white bar in front of wing*; head and throat brown with a broad green stripe from eye to back of head ending in a small tuft; wings, gray-brown with brilliant green and black wing-patch, buff-bordered; breast, pale chestnut speckled with black; buffy patches on sides of tail; white belly. **FEMALE**: mottled brown; top of head and back of neck dark brown; sides of head and neck buff, finely streaked; throat buff, unmarked; no green on head nor white bar in front of wings; wing-patch as in male but smaller. **Blue-winged Teal.**

140. BLUE-WINGED TEAL. 15 in.

(*Querquedula discors*.)

Back and underparts thickly mottled brownish and black, lightest below; head slaty with purplish gloss; *crescent-shaped white stripe in front of eye*; *shows blue* on bend of wing, followed by white bar and bright green wing-patch; white patch on sides of tail; bill black. **FEMALE**: dusky-brown; black on top of head; streaked and whitish on neck and sides of head; throat and about base of bill, *white*; back and underparts mottled and spotted; wing shows blue but no green; no white stripes on head. **Green-winged Teal.**

143. PINTAIL. 27 in.

(*Dafila acuta*.)

Head and throat brown, darkest on top; long, swan-like neck, black above, finely waved white and dusky on back and sides; *long middle tail-feathers*, black; white below with a conspicuous curved white stripe running up sides of neck to head and ending in a point; bronzy patch on wings. **FEMALE**: tail shorter but pointed; dusky, everywhere streaked; no white stripe on neck; whitish wing-bars; smaller than male. **Old-squaw.**

144. WOOD DUCK. 18 in.

(*Aix sponsa*.)

Highly variegated; long, glossy *crest* showing green and purple, marked by two narrow white parallel lines curving from bill and behind eye almost to end of crest; throat white extending irregularly in two stripes, one up behind eye, the other nearly around neck; breast glossy brown spotted with white and set off on either side by a conspicuous white stripe bordered with black; wings highly iridescent, marked by green patch bordered with white. FEMALE: duller, less conspicuously marked; head brownish-gray, slightly crested, greenish; throat and line extending from it around base of bill, with space about eye, white.

146. REDHEAD. 20 in.

(*Marila americana*.)

Puffy head and *upper* part of neck and throat, bright reddish-brown; breast and upper part of back with rump, *black*; belly white; middle back and *sides* evenly and finely waved black and white, *which shows gray at a distance*. FEMALE: grayish-brown, almost white on throat; head brown; lighter on neck. **Canvas-back.**

148. AMERICAN SCAUP DUCK. 18½ in.

(*Marila marila*.)

Similar to 149, but greenish on head; sides faintly barred; guttural call; less frequent on fresh water than 149. **Lesser Scaup.**

149. LESSER SCAUP DUCK. 16 in.

(*Marila affinis*.)

Head, neck and breast black, showing *purple* on head; back black and white, narrowly barred; upper part of wing dark, finely mottled with white; lower part of wing brownish-black with small white patch; lower breast and belly white, *sides barred*; bill, bluish tipped with black. FEMALE: black replaced by brown, lighter on head; white around bill. **Ring-necked Duck.** Female, easily confused with female **Lesser Scaup.**

151. AMERICAN GOLDEN-EYE. 19 in.

(*Clangula clangula americana*.)

Black, glossed with green; white below, extending around the neck in a collar; head bunchy, slightly crested and marked *below* and just back of eye by a nearly oval *white spot*; wings largely white; short bill. FEMALE: brown, snuff-colored on head; white collar and wing patch; band of bluish-gray across breast and down sides. **American Merganser.**

153. BUFFLE-HEAD. 15 in.

(*Charitonetta albeola*.)

Head, throat and upper neck, iridescent black; *black crest broadly banded with white over head* from eye to eye; back black, ashy on tail; wings black with large white patch; *collar and entire under parts, white*. FEMALE: smaller; brownish, lighter below; no crest; white patch on cheeks and small white wing-patch. **Hooded Merganser.**

154. OLD-SQUAW. 21 in.

(*Harelda hyemalis*)

WINTER PLUMAGE: Back, long tail-feathers and breast with *patch on sides of neck* black; throat, upper breast, top of head, neck and belly white; cheeks grayish-brown; wings black and gray; bill crossed by yellowish band. FEMALE: dark brown, lighter on head, grayish on tail; throat, breast and region about the eye gray-white; below white, brown on lower part of throat; no long tail-feathers; smaller than male. SUMMER PLUMAGE: head and neck, dusky black; grayish-white on sides of head; female shows more brown. **Pintail.**

165. WHITE-WINGED SCOTER. 19—20 in.

(*Oidemia deglandi*.)

Black, brownish below; *patch on wing* and *spot under eye, white* (nearly absent in winter); iris almost white; upper bill, red-orange on sides, *swollen on top*; fly high except in stormy weather. FEMALE: grayish-brown; whitish around upper bill and ears; white wing-patch; iris black. **American and Surf Scoters,**

167. RUDDY DUCK. 15—16 in.

(*Erismatura jamaicensis*.)

Upper half of head black, cheeks and chin white; elsewhere chestnut-red turning grayish; brown on rump and wings; belly white; tail blackish, *stiff* (often held erect by male while swimming); bill *broad*, short; feet large. WINTER PLUMAGE AND FEMALE: brown, somewhat mottled; *white stripe below eye*; swift, whirring flight; sink without diving or dive and swim below water like the grebes.

172. CANADA GOOSE. 40 in.

(*Branta canadensis*.)

Head and neck, *black with white throat-patch* extending up to and just behind eye; brownish-gray marked with lighter, darkest on back; rump and tail, black above and white below; fly after a leader in harrow-shaped ranks, often high up, crying "honk, honk, honk."

190. AMERICAN BITTERN. 30 in.

(*Botaurus lentiginosus*.)

Streaky, tawny and dark-brown, mixed with white and yellowish below; conspicuous black streak on sides of neck; short brown tail; *breast feathers loose and baggy*; yellow eye; call, loud resounding squawk, or booming notes which have given it the common name of "stake-driver."

191. LEAST BITTERN. 13 in.

(*Ixobrychus exilis*.)

Crown with back, tail and upper half of wings, shining greenish-black; rest of wing, buff and bright chestnut; throat and baggy breast, whitish shading into yellowish-brown on sides of neck and chestnut on back of neck; buffy-white below; tufts of dark feathers on either side of breast; yellow eye. **Green Heron.**

194. GREAT BLUE HERON. 45 in.

(*Ardea herodias*.)

Slaty-gray; top of head and *throat white*; long bill yellow beneath; legs *black*; *bend of wing reddish-brown*; blackish crest; feathers on breast long and streaky; stands on one leg; flies slowly with neck drawn in and legs extended behind; hoarse call. **Cranes.** (Cranes are rare; fly with necks extended; call, loud and resounding.)

201. GREEN HERON. 17 in.

(*Butorides virescens.*)

Wings and back dark green, latter tinged with blue-gray; crown greenish-black; neck and breast reddish-brown; throat and line down middle of neck to breast whitish, ending in light streaks on breast; the smallest heron; carries neck curved in flight; squawking note, "scow." **Night Heron.**

202. BLACK CROWNED NIGHT HERON. 24 in.

(*Nycticorax nycticorax naevius.*)

Green-black above; *sides of head*, throat and lower parts white; *ashy on lower half of wing* and tail; *neck short and thick*; pointed black bill; legs yellow; nests in flocks; feeds by night except in nesting season; call, "quawk." **IMMATURE:** brown, much streaked and mottled; appear slightly larger than adult. **Little Green Heron.**

212. VIRGINIA RAIL. 9½ in.

(*Rallus virginianus.*)

Streaked, blackish and olive-brown; back of neck and top of head plain brown; elsewhere *cinnamon-red* except *flanks* which are *heavily barred cross-wise*; long feet; short upturned tail; in fresh or salt marshes; grunting and guttural calls; one note of female much like sharp "click" of the Flicker. **King Rail. Clapper Rail.**

214. SORA. 8½ in.

(*Porzana carolina.*)

Olive-brown with lengthwise markings of black and some white; short tail turned up, showing buffy white underside; sides *barred* posteriorly with white; *front of head and throat black*; slaty line over eye; sides of neck and breast grayish-slate; bill *short and yellowish*; legs long; very stealthy; skulks and crouches in grass. **Young:** no black; throat whitish, brown on breast. **Virginia Rail.**

219. FLORIDA GALLINULE. 13½ in.

(*Gallinula galeata*.)

Bluish-slate, washed with olive-brown on back, rump, wings and tail; grayish on belly with *white streaks along flanks*; under tail-coverts *white*; brilliant *red bill* and *plate* across forehead; legs green to red; call, "chuck" or clucking notes; walks and *swims*. **American Coot.**

221. AMERICAN COOT. 15 in.

(*Fulica americana*.)

Smooth blackish slate color, lighter below, black on head and neck; white bill shows *dark marks near the tip*; turned up tail; white underneath; wing edged with white; eye red; young show white below and no marks on bill; toes with scalloped edges. May be mistaken for a small duck. **Florida Gallinule.**

228. AMERICAN WOODCOCK. 10½ in.

(*Philohela minor*.)

Back, mixed black and rusty slate; below reddish-brown, no markings; *large* head with short neck and *very long straight bill*; eye set *high up and far back*; crown *barred crosswise* with black and rusty; short tail; burrows in soft mud for earth-worms; nocturnal. **Wilson Snipe.**

230. WILSON SNIPE. 10½—11½ in.

(*Gallinago delicata*.)

Black and buff, much streaked; wings, dark brown marked with white; below white, streaked; sides and *lining of wings barred with black*; throat white; rusty tinge on breast and black-barred tail; *long, probing bill*; usually in fresh-water marshes or open swamps; *flight swift and crooked*; varied notes; call, harsh "scaipe"; makes a "bleating" sound (not vocal.) **Woodcock.**

239. PECTORAL SANDPIPER. 9 in.

(*Pisobia maculata*.)

Black mottled with rusty; white below, thickly streaked on breast; upper side of tail black, middle tail-feathers long and pointed; wet meadows; harsh creaky, whistling note; mating call musical, repeated eight times by male with throat greatly inflated. **Wilson Snipe. White-rumped Sandpiper.**

242. LEAST SANDPIPER. 5½ — 6 in.

(*Pisobia minutilla*.)

Blackish edged with reddish brown; light line over eye; shows whitish wing-bar; outer tail-feathers ash-gray; below white, finely streaked across breast; call sweet, "peep-peep"; in flocks along fresh and salt water; the *smallest* sandpiper. **Semipalmated Sandpiper.**

246. SEMIPALMATED SANDPIPER. 6 in.

(*Ereunetes pusillus*.)

Grayish-brown, marked with black and buffy; rump very dark; tail tapering; underparts *pure white*, slightly tinged and streaked across breast; white line over eye and dusky line beneath; note "peep-peep". **Other Sandpipers.**

248. SANDERLING. 8 in.

(*Calidris leucophaea*.)

SUMMER: mottled black, white and *rusty*; *white wing-bar*; below white, marked with black and tinged with *rusty* on throat and upper breast; *lacks the hind toe*; feet and bill black. **WINTER:** Grayish above, pure white below; call, a shrill whistle; mostly frequents beaches along the coast; runs rapidly, at times lifting its wings as if about to fly. **Spotted Sandpiper.**

254. GREATER YELLOW-LEGS. 15—16 in.

(*Totanus melanoleucus*.)

Black, much speckled with white; *head, neck and lower parts* white, streaked and marked with black except on middle of belly; bill *long, black*; legs and feet *yellow*; tail shows white barred with black; utters whistling call frequently; found near fresh and salt water. **Yellow-legs. Bartramian Sandpiper.**

255. YELLOW-LEGS. 11 in.

(Totanus flavipes.)

Similar in plumage and habits to 254, but *smaller* and more common; in flocks; most abundant in fall migration; call, three whistling notes resembling that of 254. **Greater Yellow-legs.**

256. SOLITARY SANDPIPER. 8½ in.

(Helodromas solitarius.)

Back, dusky olive-brown *finely spotted with white*, crown and back of neck showing dark streaks; white below, throat unmarked; sides of head and neck with breast slightly buffy and *distinctly streaked*; sides lightly barred; wings dark brown with one narrow white wing-bar; outer tail-feathers *white barred with black*; note, a soft whistle; frequents fresh water mostly. **Spotted Sandpiper.**

261. BARTRAMIAN SANDPIPER. 12 in.

(Bartramia longicauda.)

Black edged *regularly with buff*; neck *slender, buff, finely streaked*; tail *barred with black*; outer wing-feathers show *white* barring; below pale buff with arrow-marks of black; belly lighter with few markings; flute-like call or soft, mournful note; walks slowly, *nodding* head; flies high; when alighting *raises wings to full height before* folding them; grassy plains and pastures. **Golden Plover.**

263. SPOTTED SANDPIPER. 7½ in.

(Actitis macularia)

Greenish-ash marked lightly with black; *long white line over eye*; pure white below, *adults with dark round spots*; wings brownish, marked broadly with white bar; flies close to the water, wings full-spread, showing white bar; note, penetrating "pee-weet, weet"; walks with tilting motion. **Solitary Sandpiper.**

272. AMERICAN GOLDEN PLOVER. 10½ in.

(Charadrius dominicus.)

SUMMER: black, mottled with yellow; sides of breast white, otherwise black below; *forehead and line over eye, white*; below eye, *black*; lining of *wings ashy*. WINTER: whitish below. Plains and sandy hills or fields; runs swiftly; melodious whistle; in flocks. **Black-bellied Plover.**

273. KILLDEER. 10 in.

(Oxyechus vociferus.)

Gray-brown; rump rusty-brown; tail rather long; white below; white collar followed by a black collar; *black band across breast*; forehead, line over eye and wing-bar white; bill black; runs very swiftly; note, loud and persistent "kill-dee," often heard high overhead; found mostly inland.

274. SEMI-PALMATED PLOVER. 6 $\frac{3}{4}$ in.

(Aegialitis semipalmata.)

Forehead, throat and collar around neck white, followed by a black collar broadening across breast; below white; gray-brown above; in small scattering flocks along beaches or water-courses; call, very sweet, two or three notes, ascending; rather silent but not shy. **Piping Plover.**

Part 11.

LAND BIRDS.

289. BOB-WHITE. 10 in.

(Colinus virginianus.)

MALE: reddish-chestnut, much variegated with black and buff; tail ashy; *throat and line over eye white*; forehead, line below eye and band across breast, black; belly white, finely barred with wavy black lines; spring call of male, "bob-white"; varied notes; in flocks except during breeding season; *highly beneficial*. FEMALE: white markings buffy, little black on breast.

300. RUFFED GROUSE. 17 in.

(Bonasa umbellus.)

Black and reddish-buff, *much barred*; below barred with grayish-white; tail mottled showing broad black bar; *black feather-tufts* on side of neck (longest on male) which form a ruff; woodland and shrubby pastures; *call* of male, a remarkable whirring of the wings described as "*drumming*;" call of female "cluck" or a note of distress when young are disturbed. Canada Grouse. Prairie Hen.

316. MOURNING DOVE. 12 in.

(Zenaidura macroura carolinensis.)

Gray-blue, tinged with brownish; tail broadly tipped with white, *tapering to a narrow point*; sides of neck iridescent; marked with black on wings and sides of head; rapid whistling flight; note, "coo-oo." Pigeon. Cuckoos.

331. MARSH HAWK. 22 in.

(*Circus hudsonius*.)

Bluish-gray, showing light streaks; gray below, belly white marked with brown; tail barred; *white rump*; (female brown, barred and streaked more heavily); note shrill.

332. SHARP-SHINNED HAWK. 12 in.

(*Accipiter velox*.)

Slate-gray, crown darker; edge of wing barred with black; *tail long, square*, barred broadly, white-tipped; below white slightly tinged with brown, streaked or faintly barred; *usually silent* except at nesting period; soars in narrow circles; flies swiftly; attacks poultry. **Cooper Hawk.**

333. COOPER HAWK. 17 in.

(*Accipiter cooperi*.)

Resembles closely 332, but is *larger*, crown almost black; tail *rounded* and showing more white at tip; flies *swiftly*; attacks poultry; cackling note. **Sharp-shinned, Red-shouldered and Red-tailed Hawks.**

337. RED-TAILED HAWK. 19—22 in.

(*Buteo borealis*.)

Much like 339 above but without shoulder-patch; *tail rich, red-brown*, barred *once* with black (more or less completely barred in female and immature); below white; upper breast and throat streaked (*adults show red-brown tinge* on sides); call, a squealing whistle; woodland; beneficial. **Red-shouldered and Broad-winged Hawks.**

339. RED-SHOULDERED HAWK. 18—20 in.

(*Buteo lineatus*.)

Dark brown varied with reddish-brown, buffy or grayish; edge of wing barred black and white, showing *four notches* in *flight*; red-brown shoulder-patch; tail blackish, barred and tipped with white; below reddish-buff, *barred* with whitish; (immature *streaked* below or *heavily spotted*; shoulder-patch less showy); note, "kee-you"; beneficial, although commonly known as a hen-hawk. **Red-tailed and Broad-winged Hawks.**

343. BROADWINGED HAWK. 14 - 18 in.

(*Buteo platypterus*.)

Dark brown, slightly mottled with reddish-buff; tail, broadly barred; *edge of wing shows three notches in flight*; below white, *thickly barred with reddish-brown* and finely lined with blackish; heavy flight; notes, high pitched (one resembles that of the Wood Pewee). **Red-shouldered and Red-tailed Hawks.**

352. BALD EAGLE. 33 in.

(*Haliaetus leucocephalus*.)

Dark brown, *tail, head and neck white*; yellow bill; edge of wing black; (immature resembles Golden Eagle; adult plumage appears the third year); usually near water; eats fish, carrion, and at times water-birds and mammals; call-notes harsh. **Fish Hawk. Golden Eagle.**

360. AMERICAN SPARROW HAWK. 10½ in.

(*Falco sparverius*.)

Reddish-brown, usually barred or spotted with black; head bluish, *marked on sides with black*; tail white-tipped with one broad black bar; below white, more or less spotted; narrow pointed wings. FEMALE: much more barred, darker and *streaked below. The smallest hawk.*

364. AMERICAN OSPREY. 23 in.

(*Pandion haliaetus carolinensis*.)

Brown-black, feathers slightly tipped with whitish; *head mottled with white; white below*, sometimes faintly spotted; in flight shows dark above, head and lower parts white; nests in colonies along coast; fish-eating; dives for prey; call, a high wailing whistle. **Bald Eagle.**

336. AMERICAN LONG-EARED OWL. 14½ in.

(*Asio wilsonianus*.)

Very dark brown, lightly mottled with yellowish and white; wings and tail barred; below, streaked on breast, thickly mottled and barred elsewhere; *long, blackish ear-tufts* with light edges; yellow eyes; barking, whirring call; wooded swamp land; beneficial. **Short-eared Owl. Barred Owl.**

367. SHORT-EARED OWL. 14½—15½ in.

(*Asio flammeus*.)

Streaked and mottled dark brown with yellowish; tail broadly barred, also edge of wing; yellowish-white below streaked heavily across breast but *without bars*; *car-tufts very short*; eyes yellow; grassy marshes and rank undergrowth; sometimes feeds by day in cloudy weather; call shrill; beneficial. **Long-eared Owl. Barred Owl.**

368. BARRED OWL. 18—20 in.

(*Strix varia*.)

Conspicuously and regularly barred (more finely on head) brown and white with slight yellowish tinge; below white *barred* across breast, *streaked broadly* on belly; *no car-tufts*; bill *yellow*; eyes almost *black*; woodland; call, a resonant hoot; beneficial. **Short-eared Owl.**

372. SAW-WHET OWL. 7½—8 in.

(*Glaux acadica*.)

Dark chocolate brown, finely lined on head, spotted on back; tail slightly barred *with white*; below white, thickly streaked; eyes *yellow*; bill black; *no car-tufts*; the smallest owl common to northeastern America; call resembles "filing a saw"; flies like a woodcock; thickly wooded areas; beneficial. **Screech Owl.**

373. SCREECH OWL. 8—10 in.

(*Otus asio*.)

Gray to reddish-brown, streaked with black and somewhat mottled; below white, streaked and irregularly *barred*; *car-tufts*; call, *ceerie*, *wailing tremolo*; frequents orchards; beneficial. **Saw-whet Owl.**

375. GREAT-HORNED OWL. 22 in.

(*Bubo virginianus*.)

Mottled buff, black and yellow; *long car-tufts*; eyes *yellow*; below finely barred, *patch on throat white*; wild forested country; very loud, varied calls, "whoo", a scream or bark; attacks poultry and smaller birds but also destroys many mice and rabbits.

387. YELLOW-BILLED CUCKOO. 12¼ in.

(*Coccyzus americanus*.)

Bronzy olive-gray; white below; *lower half of curved bill yellow* tipped with black; outer tail-feathers black tipped broadly with white; wings show a *reddish-brown tinge in flight*; glides stealthily from tree to tree and keeps concealed among the leaves; slowly droops and raises tail when perching; song, loud "kuk-kuk, kuk-kuk". **Black-billed Cuckoo. Mourning Dove.**

388. BLACK-BILLED CUCKOO. 12 in.

(*Coccyzus erythrophthalmus*.)

Bronzy olive-gray; white below; bill *entirely* black; *only a little white* on tail; movements like those of 387; song, soft "coo-coo, coo-coo". **Yellow-billed Cuckoo. Mourning Dove.**

390. BELTED KINGFISHER. 14 in.

(*Ceryle alcyon*.)

Blue-gray; white collar; *big, crested head*; sits erect watching for fish and flies near the surface of the water, uttering a *rattling* call.

393. HAIRY WOODPECKER. 9 in.

(*Dryobates villosus*.)

White band down the middle of black back; wings black spotted with white; beneath white; outer tail-feathers white *not barred*; red band on neck of male; large bill; note, sharp resonant "plink". **Downy Woodpecker. Yellow-bellied Sapsucker.**

394c. DOWNY WOODPECKER. 6¼ in.

(*Dryobates pubescens medianus*.)

Bill small; closely resembles the *Hairy Woodpecker* but is *scarcely larger* than the *English Sparrow* and has outer tail-feathers *barred with black*. **Hairy Woodpecker. Yellow-bellied Sapsucker.**

402. YELLOW-BELLIED SAPSUCKER. $8\frac{1}{4}$ in.

(Sphyrapicus varius.)

Mixed black and white; *yellowish-white underneath*, streaked on sides; white bar *lengthwise* the wing; scarlet patch, bordered with black on crown and throat (throat of female white); harsh, squealing note. Downy Woodpecker. Hairy Woodpecker.

406. RED-HEADED WOODPECKER. $9\frac{1}{2}$ in.

(Melanerpes erythrocephalus.)

Black and white, not streaked; *entire head, neck and upper part of breast crimson*; pure white belly; note, a loud "kr-r-r-r-r". Red-bellied Woodpecker.

409. RED-BELLIED WOODPECKER. $9\frac{1}{2}$ in.

(Centurus carolinus.)

Black and white, back, wings and tail *barred transversely* and very evenly; below grayish-white, *tinged with red*; top of head and back of neck crimson. FEMALE: crown gray. Red-headed Woodpecker. Hairy Woodpecker.

412a. NORTHERN FLICKER. $12\frac{1}{2}$ in.

(Colaptes auratus luteus.)

Brown, barred with black; head grayish; back of neck marked with bright red; black crescent on cinnamon-brown spotted breast; in flight shows *white rump* and yellow lining of wings and tail; feeds much *on ground*, unlike other woodpeckers; sharp, nasal call-note; mating call, "flee-ker", repeated. Meadowlark.

417. WHIP-POOR-WILL. $9\frac{3}{4}$ in.

(Antrostomus vociferus.)

Mottled, strongly resembling the bark of certain trees; narrow white band across upper part of breast; big head; in daytime lies *motionless, lengthwise a branch*. Nighthawk.

420. NIGHTHAWK. $9\frac{1}{2}$ in.

(Chordeiles virginianus.)

On the wing overhead after insects just before or at dusk; long, pointed wings *show a white patch*; flight resembles that of *Chimney Swift*; sharp note, "skeep"; often nests on the flat roofs of city buildings. Whip-poor-will.

423. CHIMNEY SWIFT. $5\frac{1}{4}$ in.

(*Chaetura pelagica*.)

Dark sooty-gray; *bob-tail*; long, pointed wings which often vibrate rapidly in flight; twittering note; never seen perching like other birds. **Swallows.**

428. RUBY-THROATED HUMMING-BIRD. $3\frac{1}{4}$ in.

(*Trochilus colubris*.)

Iridescent green and purple; throat of male ruby-red; frequents flowering shrubs and plants; darts here and there and poises in mid air with wings vibrating so rapidly that they make a humming noise; "squeaky note".

444. KINGBIRD. $8\frac{1}{2}$ in.

(*Tyrannus tyrannus*.)

Dusky black; below white; *tail tipped with broad band of white*; fire-red patch on crown, usually concealed; spreads tail when about to alight; when flying high, may be mistaken for a **Tree Swallow**.

452. CRESTED FLYCATCHER. 9 in.

(*Myiarchus crinitus*.)

Olive, showing rusty brown on wings and tail; *crested head*; throat and breast, ashy-gray; wing-bars; belly sulphur-yellow; usually perches high; note, hoarse whistle. **Blue Jay.**

456. PHOEBE. 7 in.

(*Sayornis phoebe*.)

Dull olive-brown, darker on head and tail; whitish below; *bill* and feet black; tail drooping but jerking constantly; note, "phoe-be." **Wood Pewee. Least Flycatcher.**

459. OLIVE-SIDED FLYCATCHER. $7\frac{1}{2}$ in.

(*Nuttallornis borealis*.)

Olive-gray, wings and tail darker; throat and a narrow line along middle of belly *white*; *large, flattened head*; bill *heavy*, black above, light below; *long wings*; tail *notched*; whitish tufts on flanks; bold, conspicuous; plaintive call or loud whistle. **Crested Flycatcher.**

461. WOOD PEWEE. $6\frac{1}{4}$ in.

(*Myiochanes virens*.)

Dark olive-brown, brownish-black on wings and tail; frowzy head; below white, olive tinge on sides; two narrow white wing-bars; plaintive note repeated slowly and often, "pee-to-way, pee-to-way, hee-rue." **Phœbe. Least Flycatcher.**

463. YELLOW-BELLIED FLYCATCHER. $5\frac{1}{2}$ in.

(*Empidonax flaviventris*.)

Bright olive-green, tail and wings dark brown; yellowish wing-bars; sulphur-yellow below, *brightest between legs*; tinged elsewhere with pale olive; yellow eye-ring; call, a soft, slow whistle, "pee-a." **Yellow-throated Vireo. Least Flycatcher. Pine Warbler.**

465. GREEN-CRESTED FLYCATCHER. $5\frac{3}{4}$ in.

(*Empidonax virescens*.)

Olive-green, wings and tail dark; two *showy* wing-bars; head *broad*; below white, pure on throat and middle of belly, elsewhere palely tinged with greenish-yellow; (immature brighter); open woods near water; common notes, "cheep" or a two-syllabled squeak resembling that made by a rubber doll. **Traill, Least and Yellow-bellied Flycatchers.**

466. TRAILL FLYCATCHER. $5\frac{1}{2}$ —6 in.

(*Empidonax traillii*.)

Olive-green, *tinged with brown*; whitish below, sides grayish; belly slightly yellow; throat *white*; dull, *narrow* wing-bars; pale *eye-ring*; bill, black above, light below; prefers alder growth; a short, soft note, also a longer, sharp, rasping call, accented on second syllable. **Acadian Flycatcher. Least Flycatcher.**

467. LEAST FLYCATCHER. $5\frac{1}{4}$ in.

(*Empidonax minimus*.)

Olive-gray, wings, tail and head darker; two narrow wing-bars; *grayish below*, whiter on throat; *jerks its tail* and utters a sharp, energetic call. "che-bec." **Wood Pewee. Phoebe.**

474. HORNED LARK. $7\frac{3}{4}$ in.

(*Otocoris alpestris*.)

Streaked grayish-brown, somewhat *tinged* with *chocolate*; line *over eye* and *throat yellow* (paler in summer); *breast-patch* and line from bill to and *below eye black*; feathers on sides of crown long, *black*, *erectile*; tail *black*, *white-edged*; ground feeder; *runs*; whistling call; in flocks. **Vesper Sparrow. Pipit. Prairie Horned Lark.**

476b. PRAIRIE HORNED LARK. $7\frac{1}{4}$ in.

(*Otocoris alpestris praticola*.)

Similar to 474 but smaller; yellow markings almost white; formerly confined to prairies, now spreading eastward. **Horned Lark. Vesper Sparrow. Pipit.**

477. BLUE-JAY. $11\frac{1}{2}$ in.

(*Cyanocitta cristata*.)

Narrow black collar; *crested*; wings and tail deep blue, white-tipped and black-barred; bold; harsh, noisy notes. **Loggerhead Shrike. Kingfisher. Bluebird.**

488. AMERICAN CROW. 19 in.

(*Corvus brachyrhynchos*.)

Entirely black, including bill and feet; often seen in flocks; wings appear *frayed* and *ragged in flight*; note, a lusty "caw." **Bronzed Grackle.**

490. FISH CROW. 16 in.

(*Corvus ossifragus*.)

Most easily distinguished from the common crow by smaller size and *very nasal note*, "car-car-car;" found on the Atlantic coast northward to Connecticut. **Crow.**

494. BOBOLINK. $7\frac{1}{2}$ in.

(*Dolichonyx orizivorus*.)

Black with light back, creamy buff on neck; long merry musical song; open fields; sings on the wing; FEMALE: sparrow-colored, streaked above.

495. COWBIRD. 8 in.

(*Molothrus ater.*)

Black, head and neck glossy brown; *sparrow-like bill*; fly in flocks, uttering a greasy, squeaking note. FEMALE: brownish. **Bronzed Grackle. Rusty Blackbird.**

497. YELLOW-HEADED BLACKBIRD. 10¼ in.

(*Xanthocephalus xanthocephalus.*)

Black; head, throat and breast *yellow*; white wing patch. FEMALE: dark brown, yellow markings duller; no wing patch; smaller than male.

498. RED-WINGED BLACKBIRD. 9¼ in.

(*Agelaius phoeniceus.*)

Black showing scarlet shoulder-straps edged with yellow. Likes marshy places; notes "quonk-a-ree," metallic whistle and chattering. FEMALE: much streaked, marked with rusty. **Cowbird. Rusty Blackbird.**

501. MEADOWLARK. 10½ in.

(*Sturnella magna.*)

Streaked; black crescent on a *bright yellow* breast; *outer tail feathers white*; flies straight, *hovering* as it reaches the ground; sweet, plaintive song; nasal call-note. **Flicker. Dickcissel.**

506. ORCHARD ORIOLE. 7 in.

(*Icterus spurius.*)

Rich *chestnut-brown*, head, wings and tail black; narrow white wing-bar; high trees; loud, clear song. FEMALE: olivaceous above, greenish-yellow below with two white wing-bars; young male like female but with black throat. **Baltimore Oriole.**

507. BALTIMORE ORIOLE. 7½ in..

(*Icterus galbula.*)

Bright orange, head, throat, back and wings black; two white wing-bars; loud call, "co-weet, co-weet, co-weet;" song noteworthy. FEMALE: smaller, brownish-yellow showing little black. **Orchard Oriole. Scarlet Tanager.**

509. RUSTY BLACKBIRD. 9½ in.

(*Euphagus carolinus.*)

Black, uniformly iridescent and more or less rusty; tail nearly even. FEMALE: smaller and rustier with pale line over eye. **Bronzed Grackle. Red-wing Blackbird.**

511. PURPLE GRACKLE. 13 in.

(*Quiscalus quiscula.*)

Back purple to greenish otherwise similar to 511b except that feathers *on back, belly and rump show iridescent bars.* **Bronzed Grackle. Rusty Grackle. Crow.**

511b. BRONZED GRACKLE. 13 in.

(*Quiscalus quiscula æneus.*)

Iridescent black; head steely blue-green or purple; *body* distinctly bronzy; tail carried rudderwise in flight; pompous walk; rusty, grating call. **Crow.**

517. PURPLE FINCH. 6¼ in.

(*Carpodacus purpureus.*)

Not "purple," but faded crimson-red, somewhat streaky; belly whitish; *large, heavy bill*; noteworthy song. FEMALE: mottled olive-brown, *resembling sparrows.*

521. AMERICAN CROSSBILL. 6 in.

(*Loxia curvirostra minor.*)

Male brickish to bright-red, wings and tail dark brown; *bill crossed and curved at tip*; feeds on seeds of cone-bearing trees; movements *sluggish* and *parrot-like*; flight, undulating; go in small flocks, singing on the wing. FEMALE: dull olive-green; yellowish rump; belly whitish. **White-winged Crossbill.**

528. REDPOLL. 5½ in.

(*Acanthis linaria.*)

Streaked dark brown and white; belly, also bars and edges of wing white; sides streaked; *rump, crown, throat, breast* and sides suffused with rosy-red; blackish throat-patch; bill sharply pointed; canary-like song; in flocks about weedy patches. FEMALE: shows red only on crown; no throat-patch. **Purple Finch. Pine Siskin. Tree Sparrows.**

529. AMERICAN GOLDFINCH. $5\frac{1}{4}$ in.

(Astragalinus tristis).

Canary-yellow, crown, tail and wings black; white wing-bars; canary-like song; also sings in its undulating flight, "per-*chic*-o-ree, per-*chic*-o-ree"; in flocks. FEMALE: olive-tinged, without pure black or yellow. **Yellow Warbler.**

533. PINE SISKIN. 5 in.

(Spinus pinus.)

Black and *buffy-olive*, much streaked; below whitish, streaked; *tail notched*; slender, pointed bill; wing and tail-feathers dark-brownish showing *yellow* edges; whitish wing-bars; prefers evergreen or in spring, budding trees; pleasing song. **Redpoll. Female Goldfinch.**

534. SNOWFLAKE. 7 in.

(Plectrophenax nivalis.)

SUMMER PLUMAGE: male, *white*, back, part of wing and inner tail-feathers *black*; female, entirely streaked with black above, showing rusty tinge on wings. WINTER PLUMAGE: similar but male streaked with black on back, showing rusty brown tinge; a ground-feeder; *walker*; goes in flocks; song, sweet and rapid. **Horned Lark. Lapland Longspur.**

540. VESPER SPARROW. $6\frac{1}{4}$ in.

(Poœcetes gramineus.)

Streaked above and below; underparts whitish, tinged with buff across breast and on sides; shows *two conspicuous white tail feathers in flight*; noteworthy song. **Female Purple Finch.**

542a. SAVANNA SPARROW. $5\frac{1}{4}$ in.

(Passerculus sandwichensis savanna.)

Much streaked above and below; line over eye and edge of wing, yellowish; cheek sometimes suffused with yellow tinge; *tail short*, feathers pointed; movements stealthy; song, a weak trill. **Henslow Sparrow.**

546. GRASSHOPPER SPARROW. 5¼ in.

(*Coturniculus savannarum australis.*)

Crown black, divided and bordered by buff lines; back broadly, neck finely streaked, black, buff and gray; *tail even, feathers pointed*; below *bright buff*; belly whitish; before eye orange-buff; edge of wing yellow; weak, grasshopper-like song, "chir-r-r-r"; old fences about dry pastures; skulking, mouse-like; flies low and not far. **Savanna and Henslow Sparrows.**

547. HENSLOW SPARROW. 5 in.

(*Coturniculus henslowii.*)

Back conspicuously marked and streaked, showing much chestnut; head and neck a peculiar greenish tinge, *finely lined with black*; below tinged with buff and finely streaked; dark stripes and spots on sides of head; *sharp tail*; hides in the grass and can be carefully observed only with patience. **Grasshopper Sparrow.**

554. WHITE-CROWNED SPARROW. 7 in.

(*Zonotrichia leucophrys.*)

Ashy-gray; rump, tail and wings brown streaked with black; *crown conspicuously striped* with black and white; no yellow; two white wing-bars; in flocks, often with *White-throated Sparrows*; scratches with both feet; song notable. **White-throated Sparrow.**

558. WHITE-THROATED SPARROW. 7 in.

(*Zonotrichia albicollis.*)

White throat and wing-bars; unmarked ashy breast; whitish on belly; crown striped with black and white; broad white stripe over the eye shading to yellow in front; yellow and white markings obscure; in flocks about shrubbery; clear, sweet whistle, "pee-bod-dee-dee-dee." **IMMATURE: White-crowned Sparrow.**

559. TREE SPARROW. 6¼ in.

(*Spizella monticola.*)

Streaked, showing much grayish; two *showy white wing-bars*; chestnut-brown cap; breast whitish, *shading to a dark spot in the middle*; in flocks, often with *Juncos*; a winter visitor; pleasing song. **Chipping Sparrow. Field Sparrow.**

560. CHIPPING SPARROW. $5\frac{3}{4}$ in.

(*Spizella passerina*.)

Streaked; *plain ashy breast*; two narrow white wing-bars; *chestnut-brown cap*; whitish line over eye; song, a strong clear trill. **Field Sparrow. Tree Sparrow.**

563. FIELD SPARROW. $5\frac{3}{4}$ in.

(*Spizella pusilla*.)

Streaked; crown and *bill* reddish-brown; neck ashy; brownish-white *unspotted* breast; two indistinct white wing-bars; song, "fe-u, fe-u, fu, fee, fee, fee," beginning slowly and ending almost in a trill. **Chipping Sparrow. Tree Sparrow.**

567 SLATE-COLORED JUNCO $6\frac{1}{4}$ in.

(*Junco hyemalis*.)

Slate color; blackish bib (female brownish) over a white belly; *ivory bill*; *two white tail feathers*; feeds in flocks on ground, often in company with other birds; "sucks its teeth" for a call-note; song, a melodious trill.

581. SONG SPARROW. $6\frac{1}{2}$ in.

(*Melospiza melodia*.)

Reddish-brown, showing black streaks; *ashy* line over eye; whitish below with dark brown streaks which form a *spot in the middle of the breast*; cheery song, varied. **Swamp Sparrow. Lincoln Sparrow.**

583. LINCOLN SPARROW. $5\frac{3}{4}$ in.

(*Melospiza lincolni*.)

Finely streaked with black and brown; white beneath, with *broad buff band* across the breast and down the sides; narrow streaks on throat, upper breast and sides; dark stripe on each side of throat from corner of mouth; very shy. **Song and Swamp Sparrows.**

584. SWAMP SPARROW. $5\frac{3}{4}$ in.

(*Melospiza georgiana*.)

Very darkly colored and heavily streaked; chestnut cap, sometimes indistinctly streaked; *black forehead*; plain ashy breast; shy; frequents bushes along the water's edge. **Song Sparrow. Lincoln Sparrow.**

585. FOX SPARROW. $7\frac{1}{2}$ in.

(*Passarella iliaca.*)

Fox-red back and tail; ashy about neck and head; white breast *thickly* streaked with dark spots; scratches like a hen; alert; noteworthy song. **Brown Thrasher. Hermit Thrush.**

587. TOWHEE. $8\frac{3}{4}$ in.

(*Pipilo erythrophthalmus.*)

Black above and black bib (female rich brown); reddish-brown sides; white underneath; *outer tail-feathers white*; scratches about under bushes; brilliant notes, "tow-hee" and "che-wink." **Robin.**

593. CARDINAL. $8\frac{1}{2}$ in.

(*Cardinalis cardinalis.*)

Bright red, including large blunt bill; *crested*; black markings on face and throat; fine songster. FEMALE: olive-gray, tinged lightly with red and obscurely marked. **Tanagers.**

595. ROSE-BREASTED GROSBEAK. $8\frac{1}{4}$ in.

(*Zamelodia ludoviciana.*)

Black, including throat; white rump and wing-patches; *breast white with a brilliant rose-red blotch*; lining of wings also rose-red; *large beak*; sluggish; song, a rich carol. FEMALE: yellowish-brown darkly mottled; no red.

598. INDIGO BUNTING. $5\frac{3}{4}$ in.

(*Passerina cyanea.*)

Indigo blue, wings and tail blackish; in flocks; often feeds in the grass; noteworthy song. FEMALE: brown. **Bluebird.**

604. DICKCISSEL. 6 in.

(*Spiza americana.*)

Streaked; ashy on head and neck; white chin; *black throat-patch*; *yellow breast*; white belly; *line over eye, yellow*; wings show chestnut-brown. FEMALE: less yellow; no throat-patch. **Yellow-breasted Chat. Meadowlark.**

608. SCARLET TANAGER. 7¼ in.

(*Piranga erythromelas.*)

Scarlet; black wings and tail; rather sluggish; note, "chip, chree;" song resembles that of robin. FEMALE: olive-green; usually near by. Cardinal. Summer Tanager.

610. SUMMER TANAGER. 7½ in.

(*Piranga rubra.*)

Vermilion all over; no black. FEMALE: yellowish-olive; usually near the male. Cardinal.

611. PURPLE MARTIN. 7½ in.

(*Progne subis.*)

Blue-black all over; often nests in bird houses; twittering song. FEMALE: lighter, streaked with grayish-white below. Tree Swallow. Chimney Swift.

612. CLIFF SWALLOW. 6 in.

(*Petrochelidon lunifrons.*)

Steel-blue back and crown; rump, throat and sides of head chestnut; forehead whitish; belly white, breast brown-gray marked by steel-blue patch; tail square; in flocks; nests in eaves of buildings or under cliffs; twittering call. Barn and Tree Swallows.

613. BARN SWALLOW. 7 in.

(*Hirundo erythrogaster.*)

Steel-blue; chestnut shading to white below; forehead chestnut; deeply forked tail. IMMATURE: tail shorter; colors duller. Cliff Swallow.

614. TREE SWALLOW. 6¼ in.

(*Iridoprocne bicolor.*)

Metallic blue-green; shining white below; tail appears nearly square in flight; often seen skimming near the surface of still water. Barn Swallow. Purple Martin.

616. BANK SWALLOW. 5¼ in.

(*Riparia riparia*.)

White throat and belly; brownish band across breast, otherwise dull brown-gray; tail notched; nests in holes near top of soft banks(usually sand); twittering call; in flocks. Rough-winged Swallow. Swift.

619. CEDAR WAXWING. 7 in.

(*Bombycilla cedrorum*.)

Smooth snuff-brown; *crested*; chin and line from bill to crest, black; end of tail banded with yellow; in small flocks; "monotonous lisping note."

621. NORTHERN SHRIKE. 10½ in.

(*Lanius borealis*.)

Gray, very light above eyes and on forehead; below hoary; breast, throat and sides show dusky, *wavy, cross-wise bars*; line from bill to eye, continued broadly *below eye* back to nape, *black*; wings and tail black, mostly white-tipped, former marked by white patch; *hooked bill*; straight even flight; seeks commanding perch; song, strong, varied, sometimes mimetic. **Loggerhead Shrike. Blue Jay.**

622. LOGGERHEAD SHRIKE. 9 in.

(*Lanius ludovicianus*.)

Slaty-gray; commonly white beneath, *unmarked*; wings and tail black, marked with white; *hooked bill*; bar through eye and *between forehead and bill*, black; imitates notes of other birds. **Blue Jay. Catbird.**

624. RED-EYED VIREO. 6 in.

(*Vireosylva olivaceus*.)

Olive-green; crown slaty; white below, lightly tinged on sides; *dark line above a white one over eye*; clear, persistent song. **Warbling Vireo.**

627. WARBLING VIREO. $5\frac{1}{4}$ in.

(*Vireosylva gilva*.)

Mouse-gray; olive rump; white below tinged with greenish yellow; creamy-white around eye; song, rippling warble. **Red-eyed Vireo. Least Flycatcher.**

628. YELLOW-THROATED VIREO. $5\frac{1}{2}$ in.

(*Lanivireo flavifrons*.)

Bright olive-green shading to ashy-blue on rump; *yellow throat* and breast sharply defined against white belly; line around eye and to bill, yellow; *two white wing-bars*; slow, distinct song. **Yellow-bellied Flycatcher. Blue-headed Vireo.**

629. BLUE-HEADED VIREO. $5\frac{1}{2}$ in.

(*Lanivireo solitarius*.)

Olive-green; *ashy-blue head*; pure white below; sides yellowish; line around eye and to bill, white; *two white wing-bars*; song, noteworthy. **Yellow-throated Vireo.**

631. WHITE-EYED VIREO. $5\frac{1}{2}$ in.

(*Vireo griseus*.)

Rich olive-green; *two broad yellowish-white wing-bars*; below white, *sides and lower belly bright yellow*; white iris; *line from bill to and around eye yellow*; thickets and road-side undergrowth; emphatic call, "whip-tee-you;" mimics notes of other birds. **Yellow-throated and Blue-headed Vireos.**

636. BLACK AND WHITE CREEPER. $5\frac{1}{4}$ in.

(*Mniotilta varia*.)

Black and white, *striped*; white belly (female less streaked below); creeps up and down trees like the *Nuthatches*; song rapid, rasping, "he-ho, he-ho, he-ho." **Blackpoll Warbler.**

637. PROTHONOTARY WARBLER. $5\frac{1}{4}$ in.

(*Prothonotaria citrea*.)

Entire head, neck and breast, *bright orange-yellow*; blue-gray wings, rump and tail; prefers willows near water; strong, ringing song. FEMALE: duller. **Blackburnian Warbler. Summer Warbler.**

641. BLUE-WINGED WARBLER. $4\frac{3}{4}$ in.

(*Helminthophila pinus*.)

Bright olive-green; tail and wings ashy-blue, the latter with two *white* bars; *top of head, forehead and under parts bright yellow, unmarked*; white tail-spots; *narrow black line through eye*; edge of woods and second growth; two characteristic songs. **Golden-winged and Brewster Warblers.**

642. GOLDEN-WINGED WARBLER. 5 in.

(*Helminthophila chrysoptera*.)

Blue-gray; crown and *patch on wing*, golden yellow; *throat and stripe through eye, black*; white below; song, drawling "zee, zee, zee, zee." **Blue-winged and Parula Warblers.**

645. NASHVILLE WARBLER. $4\frac{3}{4}$ in.

(*Helminthophila rubricapilla*.)

Olive-green; head and neck bluish gray; indistinct brown patch on crown; *bright yellow below*; no wing-bars; *white eye-ring*; song, not loud, "wee-see, wee see, wit-a-wit." **Tennessee Warbler.**

647. TENNESSEE WARBLER. $4\frac{1}{2}$ in.

(*Helminthophila peregrina*.)

Olive-green, brightest on rump; head and neck, blue-gray; *dull white below*; faint white line over eye; no wing-bars; song, shrill and feeble. **Nashville Warbler.**

648a. NORTHERN PARULA WARBLER. $4\frac{1}{2}$ in.

(*Compsothlypis americana usneae*.)

Grayish-blue, *yellowish across the back*; throat and breast yellow, latter crossed by brownish band; wing-bars; peculiar song. **Golden-winged Warbler.**

650. CAPE MAY WARBLER. $5\frac{1}{4}$ in.

(*Dendroica tigrina*.)

Yellowish-olive spotted with black on back; crown very dark; cheeks yellow with an *orange-brown ear-patch*; yellow below, black streaks on breast; wing-bars; song all on one note. **Black-throated Green Warbler. Magnolia Warbler.**

652. YELLOW WARBLER. 5 in.

(*Dendroica aestiva*.)

Yellow all over shading to olive on back; wings and tail darker, showing yellow edgings; orange-brown streaks on breast; song, penetrating and unceasing; nests often in city trees. FEMALE: not so yellow, less streaked. **American Goldfinch.**

654. BLACK-THROATED BLUE WARBLER. 5¼ in.

(*Dendroica caerulescens*.)

Slaty-blue; below white; black throat and sides; *white patch on wing*; hoarse note, "kree, kree." FEMALE: grayish; *no black* but may be recognized by *wing patch*. **Blue-gray Gnatcatcher.**

655. MYRTLE WARBLER. 5½ in.

(*Dendroica coronata*.)

Blue-gray, streaked; crown, *rump* and spot on each side of breast *yellow*; white below, marked on breast and sides with black; usually in small flocks. **Magnolia Warbler.**

657. MAGNOLIA WARBLER. 5 in.

(*Dendroica magnolia*.)

Back and tail black; crown blue-gray; yellow rump; *yellow* below, streaked with black on breast and sides; white patch on wings; broad white band across tail, *seen from below*. **Myrtle Warbler. Cape May Warbler.**

658. CERULEAN WARBLER. 4½ in.

(*Dendroica caerulea*.)

Azure-blue, marked with black on back and sides of head; white below, streaked with blue; two white wing-bars; frequents tree-tops; song resembles that of *Parula*. **Blue-gray Gnatcatcher.**

659. CHESTNUT-SIDED WARBLER. 5 in.

(*Dendroica pensylvanica*.)

Back shows black streaks mixed with olive-green; yellow cap; white below; *chestnut markings* on sides of breast; wing-bars; song, chee, chee, chee, *chee-ar*." **Bay-Breasted Warbler.**

660. BAY-BREASTED WARBLER. $5\frac{1}{2}$ in.

(*Dendroica castanea*.)

Ashy-brown, black-streaked; black cheeks and forehead; throat, upper part of breast, sides and crown, *rich chestnut-brown*; white wing-bars; song, soft, liquid warble. **Chestnut-sided Warbler.**

661. BLACK-POLL WARBLER. $5\frac{1}{2}$ in.

(*Dendroica striata*.)

Streaked black and white; *black crown*; white cheeks; below white with line of black spots down sides; tree-tops; song, a staccato musical "chink" repeated rapidly. **Black and White Creeper.**

662. BLACKBURNIAN WARBLER. $5\frac{1}{4}$ in.

(*Dendroica blackburniæ*.)

Black, streaked with white; middle of crown, throat and sides of head *flaming orange*; black patch behind eye; white wing-bars; usually, high up in evergreen trees; shrill song; FEMALE: duller. **Prothonotary Warbler.**

667. BLACK-THROATED GREEN WARBLER. $5\frac{1}{4}$ in.

(*Dendroica virens*.)

Olive-green; *bright yellow cheeks*; throat and upper part of breast black; tail shows much white; white wing-bars; song, repeated often and rapidly, "chee, chee, chee, char, chee." **Cape May Warbler. Hooded Warbler.**

671. PINE WARBLER. $5\frac{3}{4}$ in.

(*Dendroica vigorsii*.)

Olive-green; yellow breast; sides of head suffused with yellowish, sometimes forming a *yellow line over eye*; two dull white wing-bars; song, a loud, penetrating trill. **Yellow-bellied Flycatcher. Female Maryland Yellow-throat. Female Wilson Black-cap.**

672. PALM WARBLER. 5 in.

(*Dendroica palmarum*.)

Streaky olive-gray; *chestnut cap*; yellow line over eye; beneath yellowish (brightest on throat and under tail) streaked with brown; feeds on ground in flocks; *tilts its tail* constantly. **Water-Thrushes. Yellow Palm Warbler.**

672a. YELLOW PALM WARBLER. 5½ in.

(*Dendroica palmarum hypochrysea*.)

Similar to 672 but *uniform* yellow beneath and more strongly olive above; song, soft, broken trill, "tzee-tzee". **Palm Warbler.**

673. PRAIRIE WARBLER. 4¾ in.

(*Dendroica discolor*.)

Olive-green, *middle of back spotted with chestnut*; sides of head, broad line over eye and under parts, bright yellow; *lines through and below eye with chain of streaks down sides black*; white *tail-spots* and yellowish wing-bars; in wild pastures; song, thin, "zee", repeated on ascending scale. **Magnolia, Canadian, Cape May and Female Black-throated Green Warblers.**

674. OVEN-BIRD. 6¼ in.

(*Seiurus aurocapillus*.)

Olive-green showing yellow tinge; below pure white, thickly spotted; *orange-brown cap* bordered with black stripes; scratches like a hen; *tilts tail*; under shrubbery; note, "teacher" repeated rapidly with increasing emphasis. **Water-Thrushes. Wood Thrush.**

675. WATER-THRUSH. 6 in.

(*Seiurus noveboracensis*.)

Plain olive-brown; below white, tinged with sulphur-yellow and thickly streaked, finely *on chin and throat*; belly plain; *short bill*, yellowish line over eye; *wags tail*; noticeable song, high, ending low. **Oven-bird. Other Water-Thrushes.**

675a. GRINNELL WATER-THRUSH. $5\frac{1}{4}$ in.

(*Seiurus noveboracensis notabilis*.)

Very dark olive-brown; below white, palely tinged with yellowish, thickly streaked; *pale* stripe over the eye; skulks along the very edge of the water; *tilts tail*; brilliant song. Louisiana Water-Thrush. Oven-bird. Palm Warbler.

676. LOUISIANA WATER-THRUSH. $6\frac{1}{4}$ in.

(*Seiurus motacilla*.)

Olive; white below, buff-tinged and streaked with black; *throat and middle of belly unmarked*; *white line over eye*; loud, ringing song. Grinnell Water-Thrush. Oven-bird. Palm Warbler.

678. CONNECTICUT WARBLER. $5\frac{1}{2}$ in.

(*Oporornis agilis*.)

Olive-green; head ashy, throat darker; *white ring* around eye; belly yellow; wings long and pointed; low bushes and swampy places; penetrating song. Mourning Warbler.

679. MOURNING WARBLER. $5\frac{1}{2}$ in.

(*Oporornis philadelphia*.)

Bright olive-green; head, neck, throat and upper part of breast *bluish-ash*; yellow belly; black band on breast; about bushes and in the grass; noteworthy song. Connecticut Warbler.

681. MARYLAND YELLOW-THROAT. $5\frac{1}{4}$ in.

(*Geothlypis trichas*.)

Olive-green showing brownish tinge; *broad, black stripe across forehead and through eye*, bordered with grayish-white; yellow beneath, lighter on belly; wings short; in low bushes; song, "witch-i-ty, witch-i-ty, witch-i-ty." FEMALE: duller, *without black stripe* and easily confused with Female Mourning, Connecticut and Pine Warblers. Kentucky and Wilson Warblers.

683. YELLOW-BREASTED CHAT. $7\frac{1}{4}$ in.

(*Icteria virens*.)

Olive-green; bright yellow below, belly *white*; region about eye blackish, marked with a white line above eye; white spot and shorter line below eye; the largest warbler; remarkable song. **Dickcissel. Maryland Yellow-throat. Yellow-Throated Vireo.**

684. HOODED WARBLER. $5\frac{1}{2}$ in.

(*Wilsonia citrina*.)

Olive-green; yellow below; head, neck and upper breast black; yellow "*mask*" across forehead and cheeks; white spots on tail; FEMALE: less black. **Maryland Yellow-throat. Wilson Warbler. Canadian Warbler.**

685. WILSON WARBLER. $4\frac{3}{4}$ in.

(*Wilsonia pusilla*.)

Bright olive-green; yellow below; *black cap*; in low bushes. **Pine Warbler. Female Maryland Yellow-throat.**

686. CANADIAN WARBLER. $5\frac{1}{2}$ in.

(*Wilsonia canadensis*)

Bluish-lead color; yellow below with a *necklace of black spots*; no wing-bars; note-worthy song. **Magnolia Warbler.**

687. AMERICAN REDSTART. $5\frac{1}{4}$ in.

(*Setophaga ruticilla*.)

Male black, with *six flame-colored spots* (female olive, with *six yellow spots*); spreads tail like a fan; song, "chee, chee chee, chew;" darting motions.

697. AMERICAN PIPIT. 6 in.

(*Anthus rubescens*.)

Streaked grayish-brown; below buff, breast and sides darkly streaked; tip of wing and *outer edge of tail whitish*; light line above and ring around eye; *long hind toe-nail*; *wagging walk*; undulating flight; in flocks; song, soft "dee-dee". **Larks. Water-thrushes.**

703. MOCKINGBIRD. $10\frac{1}{2}$ in.

(*Mimus polyglottus*.)

Ashy-gray, wings and tail darker; whitish below; *wing shows white patch*; outer tail-feathers white; famous for its song. **Catbird. Loggerhead Shrike. Cuckoos.**

704. CATBIRD. $8\frac{3}{4}$ in.

(*Dumatella carolinensis*.)

Slate-gray; bill, crown and tail black; *chestnut patch under tail*; about shrubbery; willing to be looked at; calls like a cat; song, greatly varied.

705. BROWN THRASHER. $11\frac{1}{4}$ in.

(*Toxostoma rufum*.)

Bright rusty-red; white below, much streaked; long bill and *long, sweeping tail*; wing-bars; song, loud and varied; about bushes. **Wood Thrush.**

718. CAROLINA WREN. $5\frac{1}{2}$ in.

(*Thryothorus ludovicianus*.)

Reddish-brown, brightest on rump; below rusty-buff; throat and *line over eye*, whitish; wings and tail barred, the latter long and carried erect; "quick as a flash," flight jerky; very loud, ringing song and striking call-notes; near wooded streams in wild pasture undergrowth. **Bewick and Long-billed Marsh Wrens.**

721. HOUSE WREN. 5 in.

(*Troglodytes aedon*.)

Brown, barred with darker; below grayish, tinged with brown, also barred; rusty-colored tail, *two inches long*; remarkable song. **Other Wrens.**

722. WINTER WREN. $4\frac{1}{4}$ in.

(*Nannus hiemalis*.)

Dark tawny-brown, barred on wings and tail; lighter below, barred except on throat and breast; pale line over eye; *stub tail, carried erect*; fine songster. **Other Wrens.**

725. LONG-BILLED MARSH WREN. $5\frac{1}{4}$ in.

(*Telmatodytes palustris*.)

Brown, almost black on head, tail and between shoulders, *the latter streaked with white*; barred only on wings and tail; *line over eye* and middle of breast white; *long bill*; spluttering song. **Other Wrens.**

726. BROWN CREEPER. $5\frac{1}{2}$ in.

(*Certhia familiaris americana*.)

Brown finely streaked with white; silky white below; long curved bill; *flies to the base of tree-trunks and works up in spirals* bracing with its long pointed tail; calls softly, "screep, screep." **Nuthatches.**

727. WHITE-BREASTED NUTHATCH. $5\frac{3}{4}$ in.

(*Sitta carolinensis*.)

Ashy-blue; cap and back of the neck, black; white below; constantly exploring the bark of trees; does not brace, woodpecker fashion, with its tail; note, nasal "yank, yank, yank." **Red-breasted Nuthatch. Brown Creeper.**

728. RED-BREASTED NUTHATCH. $4\frac{3}{4}$ in.

(*Sitta canadensis*.)

Bluish-gray; crown and *stripe through eye*, black with a white line over eye; *below rusty-red*, throat white; climbs up and down and around trees; note, "cark, cark, cark." **White-breasted Nuthatch.**

731. TUFTED TITMOUSE. $6\frac{1}{4}$ in.

(*Baeolophus bicolor*.)

Ashy-gray; below whitish; sides reddish-brown; *head crested*; bill and band above black; loud emphatic song; whistling call "peto." **Chickadee. Catbird.**

735. CHICKADEE. $5\frac{1}{4}$ in.

(*Penthestes atricapillus*.)

Ashy-gray; *black throat* and cap; white cheeks; frowzy; lively; often swings head downward from the tips of twigs; call-note "chick-a-dee-dee"; song musical, "pewee," confused with note of *Phæbe*.

748. GOLDEN-CROWNED KINGLET. 4 in.

(*Regulus satrapa*.)

Olivaceous; two whitish wing-bars; *orange-yellow stripe edged with black on crown*; restless; often seen feeding in evergreens; note, rapid "tzee, tzee, tzee." **Ruby-crowned Kinglet.**

749. RUBY-CROWNED KINGLET. 4 $\frac{1}{4}$ in.

(*Regulus calendula*.)

Dark greenish-olive; light underneath; *ring about eye* and wing-bars white; male shows at will a brilliant ruby crown-patch; remarkable song. **Golden-crowned Kinglet. Warblers.**

751. BLUE-GRAY GNATCATCHER. 5 in.

(*Poliioptila caerulea*.)

Blue-gray; white beneath; *long tail* edged with white; male has a black line over eye; fidgety, never long in a place; floating flight; wheezy song. **Black-throated Blue Warbler. Cerulean Warbler.**

755. WOOD THRUSH. 7 $\frac{3}{4}$ in.

(*Hylocichla mustelina*.)

Bright brown on head shading to olive-brown on rump and tail; white beneath with conspicuous *round* black spots except on throat and middle of belly; remarkable song. **Other Thrushes. Fox Sparrow.**

756. WILSON THRUSH. 7 $\frac{1}{4}$ in.

(*Hylocichla fuscescens*.)

Uniform tawny brown; no ring about eye; below white, faintly tinged and lightly spotted; shy, about bushes; notable song. **Other Thrushes.**

757. GRAY-CHEEKED THRUSH. 7 $\frac{1}{2}$ in.

(*Hylocichla aliciae*.)

Uniform olive; below white, *very palcely* suffused with buff; cheeks *gray tinged*; breast and sides of throat spotted; **Olive-backed and other Thrushes,**

758_a. OLIVE-BACKED THRUSH. $7\frac{1}{4}$ in.

(*Hylocichla ustulata*.)

Uniform dark olive; below white, darker on sides; cheeks, throat, breast and ring about eye buff; spotted on breast; song loud and ringing. **Gray-cheeked Thrush. Hermit Thrush. Wilson Thrush.**

759b. HERMIT THRUSH. $7\frac{1}{4}$ in.

(*Hylocichla guttata pallasii*.)

Olive-brown; *tail reddish-brown*; below dull white tinged with buff and thickly spotted across the breast; shy, hiding in bushes; famous for its song but silent during migration. **Other Thrushes.**

761. ROBIN. 10 in.

(*Planesticus migratorius*.)

Dark slate color; black head; *yellow bill*; throat white streaked with black; underparts bright chestnut-red; sings "cheer-i-ly, cheer-i-ly, cheer-up." **IMMATURE:** spotted with black. **Towhee.**

766. BLUEBIRD. $6\frac{1}{2}$ in.

(*Sialia sialis*.)

Sky-blue; brownish-red below; female shows less blue; usually in pairs; call, "pu-ri-ty, pu-ri-ty;" often heard before seen. **Blue Jay.**

Note to Key.

Since no key can make up for inaccurate observation, it is essential that at least one or two points about an unknown bird shall have been *clearly seen* before turning to any key for aid in identification.

Keep in mind (1) the *relative size* and (2) see accurately some one or more distinguishing markings of any bird in question.

The following key *includes the land birds* described in the foregoing pages. These are arranged by number, first in three groups according to size (using the Robin and English Sparrow as 10 and 6 inch measures of length respectively) and second, according to distinguishing colors or markings.

Suppose a bird at least as large as a Robin, i. e., 10 inches or more in length, is seen, showing white outer tail-feathers and some yellow beneath. Upon referring to the key only one number, namely 502, will be found which answers this description. When more than one number fulfils the conditions of the observation it becomes necessary to compare descriptions before coming to a decision.

Blackface figures indicate female birds that are noticeably unlike the corresponding males.

BRIGHT COLORS. STREAKED, BARRED or SPOTTED	Showing BLUE Showing RED Showing YELLOW Showing BLACK Showing BROWN
BRIGHT COLORS. <i>NEVER</i> STREAKED, BARRED or SPOTTED	Showing BLUE Showing RED Showing YELLOW Showing BLACK Showing BROWN
IRIDESCENT COLORS	
DULL COLORS	FEW MARKINGS Sharply defined..... Obscure No MARKINGS.....
BACK	STREAKED Entirely streaked or mottled, no 'distinct cap.... Rump plain, cap or striped crown Gray or ashy about head or neck..... BARRED AND SPOTTED Barred, including wings..... Black and white, wings barred or spotted..... No MARKINGS Back and wings alike..... DISTINCTIVE MARKINGS Rump patch white, yellow or brown On back of neck, between shoulders or middle of back.....

10 inches or more	Between 6 and 10 inches.	Less than 6 inches.
390, 477.....		658.....
412a, 761.....	406, 409, 517, 521, 595.....	428, 528, 687.....
412a, 501.....	604.....	650, 652, 655, 657, 662, 667, 671 672a, 681 686, 687.....
360, 412a, 477, 501.....	406, 409, 493, 494, 534, 604.....	533, 650, 655, 657, 659, 660, 662, 667, 686, 687.....
289, 331, 339, 343.....	585, 755, 756, 759b.....	598, 659, 660, 718 721, 722, 725 726.....
360, 705.....		
.....	766.....	598, 629, 642, 648, 654, 727, 728 751.....
761.....	498, 593, 608, 610, 766.....	
497.....	498, 507, 619, 683.....	529, 628, 637, 642, 648a, 671, 678 679, 681, 684, 685.....
488, 490, 497.....	444, 495, 498, 509, 611, 613.....	428, 654.....
.....	614.....	
.....	495, 506, 587, 619, 766.....	728.....
316, 511, 511b.....	493, 495, 509, 611, 612, 613.....	428.....
.....	614.....	
316, 703.....	417, 444, 567, 622, 704.....	463, 683, 735, 748, 749.....
364, 387, 388, 497.....	456, 459, 461, 495, 509, 593.....	466, 467, 529, 598, 627, 647, 749..
621.....	608, 611, 624.....	
488.....	456, 495, 509, 610, 611.....	423, 598, 627, 681.....
501.....	373, 417, 420, 494, 517, 540.....	533, 542a, 546, 547, 583, 636, 650, 581, 595.....
.....	558.....	661.....
.....	554, 559, 585, 604.....	560, 662, 726.....
289, 300, 360, 368.....	372, 409.....	563, 584.....
412a.....		721, 722.....
.....	393, 394c, 402.....	636.....
316, 387, 388, 488.....	456, 495, 567, 611, 624, 674.....	423, 627, 645, 647, 675a, 678, 679, 681, 684, 685, 686, 728, 735.....
511b, 761.....	676, 683, 756, 757, 758a...	
331, 412a.....	595, 612.....	650, 655, 657, 725.....
412a.....	393, 394c, 402, 409.....	547, 648, 673, 725.....

UNDER PARTS

No MARKINGS—Pure white, yellow or ashy.....

Dull colors, indistinctly shaded

FEW, IF ANY MARKINGS

Distinctive red, yellow or brown breast.....

BELLY white or yellow, sharply contrasting with
breast.....

DISTINCTIVE MARKINGS

A line of streaks down sides or across breast.....

Spotted or finely streaked with throat and middle
of belly plain.

Bib, collar or band across breast.....

Spot or patch on throat, sides, breast or sides of
neck.....

TINGED

Noticeably with yellow, red, buff or brownish ...

HEAD

TOP

Crested or red crown-patch displayed at will

Crown conspicuously striped.....

Distinct cap, no line over eye.....

Distinct cap, with line over, through or back of
eye.....

Forehead showing black, sometimes with bar
through eye

Hood

SIDES

Eartufts.....

Conspicuous line over or through eye.....

Inconspicuous but distinctive line over eye.....

Conspicuous eye-ring or, black or yellow mask...

Distinctly marked or, colored on cheeks or sides
of head.....

Neck band.....

BILL.

Hooked, or noticeably long, sometimes curved...

Large and stout, or noticeably short and thick....

Used for hammering.....

10 inches or more.	Between 6 and 10 inches.	Less than 6 inches.
387, 388,	393, 394a, 406, 444, 461, 463	316, 466, 546, 563, 584, 627, 629,
.....	554, 614, 624, 683.....	641, 642, 645, 658, 678, 726, 727,
.....		751.....
703.....	456, 461, 506, 507, 509, 622,	467, 563, 584, 647, 725, 735, 748,
.....		749, 751.....
501, 761.....	613, 683, 766.....	628, 637, 645, 671, 681, 684, 685,
.....	452, 567, 587, 604, 612, 683,	628, 631, 654, 678, 679
.....	402, 585.....	655, 657, 658, 659, 661, 662, 673
.....		686.....
360.....	417, 420, 567, 587, 655, 676,	547, 583, 648a, 654, 658, 660, 667,
.....		684.....
390, 412a.....	417, 587, 616.....	583, 667, 679.....
375, 412a, 501.....	402, 474, 558, 559, 581, 595,	428, 528, 642, 679, 689, 735
.....	604, 612.....	
.....	402, 409, 452, 517, 675, 757	463, 528, 547, 563, 583, 629, 671,
.....	758a.....	672, 672a, 722, 728.....
289, 390, 477.....	402, 444, 452, 593, 619, 731,	749.....
.....	331, 554, 558, 674.....	546, 636, 748.....
.....	409, 704.....	529, 641, 657, 660, 661, 685, 729
.....		735.....
.....	559	560, 563, 642, 655, 659, 672, 728
.....	593, 619.....	584, 660, 681, 751.....
352, 497.....	406, 495, 506, 507.....	629, 637, 679, 684.....
300, 366, 375.....	373, 474.....	
289, 501.....	393, 394c, 474, 474b, 558,	628, 641, 642, 650, 657, 667, 671,
.....	604, 622, 675, 676, 683.....	673, 675a, 718, 725, 751.....
497.....	509, 624, 675a.....	542a, 647, 658, 672, 697.....
.....	683, 758a.....	463, 749, 681, 628, 645, 629, 684
.....		678, 631.....
316, 360, 621.....	409, 619, 731, 757.....	547, 583, 642, 650, 658, 660, 661,
.....		662, 667, 735.....
390, 477.....	393, 494.....	
331, 360, 387, 388,	521, 622.....	428, 725, 726.....
621, 705.....		
.....	495, 517, 593, 595.....	529, 598.....
412a.....	393, 394c, 402, 406, 409, 412a	

WINGS	Showing notches in flight..... Long and pointed..... One or two conspicuous wing-bars..... Spots, patches or bright markings.....
TAIL	<div>MARKINGS</div> Outer feathers conspicuously white or spotted.... Reddish brown or, tipped with white or yellow.. Barred or, banded with black or white..... <div>SIZE</div> Very long, sometimes keeled or forked..... Very short..... <div>MOVEMENTS</div> Used for bracing..... Jerking or tilting motions..... <div>SHAPE</div> Square Rounded..... Notched..... Narrow and pointed.....

10 inches or more.	Between 6 and 10 inches.	Less than 6 inches.
339, 343, 488.....	417, 420, 611, 613, 614.....	423.....
705.....	452, 461, 506, 507, 554, 559.....	463, 465, 467, 528, 529, 628, 629.....
		631, 641, 648a, 658, 659, 660, 661,.....
		667, 671, 673, 726, 749.....
393, 497, 621, 703,.....	402, 406, 420, 498, 587, 595,.....	642, 654, 655, 657, 687.....
	622.....	
387, 390, 501, 703,.....	393, 394c, 474, 540, 567, 587,.....	655, 658, 660, 667, 671, 672, 687..
761.....	622.....	697, 751.....
316, 332, 333, 337,.....	444, 452, 585, 619, 759b.....	
339, 360, 387, 477.....		
300, 331, 332, 360,.....	394c, 402, 409.....	657, 721, 722, 725.....
366, 477.....		
316, 387, 388, 511b,.....	613.....	726, 751.....
705.....		
	412a.....	423, 428, 721, 722, 727, 728.....
412a.....	393, 394c, 402, 406, 409.....	726.....
	456, 674, 675, 675a, 676.....	467, 672, 672a, 675a, 697.....
332.....	612, 614.....	546.....
333.....		
	459, 613, 616.....	533.....
316.....		542a, 547.....

Table of Occurrence.

In this table the birds observed by the authors in Lincoln Park, Chicago, are arranged according to the order of their comparative frequency. The abundance of the birds, that is, the number of individual birds of different species, is in no wise shown but simply the relative representation of each species. For instance, out of a total of 454 mornings covering observations during six years, the *Robin* was seen on 383 different mornings. The record for 1902, during the absence of the authors, was made by Mr. H. V. Bozell.

NUMBER OF MORNINGS SEEN
IN LINCOLN PARK, CHICAGO

	1898	1899	1900	1901	1902	1903	Total
1 Robin.....	62	52	57	69	61	82	383
2 Bronzed Grackle.....	57	58	57	60	57	72	361
3 Blue Jay.....	49	58	61	65	40	51	324
4 Cowbird.....	41	34	44	44	39	54	256
5 Junco.....	33	35	32	32	51	42	225
6 Towhee.....	35	18	24	44	45	48	214
7 Song Sparrow.....	27	33	19	38	37	36	190
8 Flicker.....	29	18	21	33	37	46	184
9 Ruby-crowned Kinglet.....	31	28	23	30	22	39	173
10 White-throated Sparrow....	30	23	23	31	26	35	168
11 Bluebird.....	26	1	7	31	35	50	150
12 Wilson Thrush.....	34	35	23	21	11	20	144
13 Yellow-bellied Sapsucker..	23	12	15	30	27	34	141
14 Yellow Warbler.....	27	30	24	18	13	27	139
15 Belted Kingfisher.....	25	21	24	27	21	19	137
16 Redheaded Woodpecker..	23	23	16	26	16	26	130
17 Myrtle Warbler.....	19	25	11	21	22	31	129
18 Brown Thrasher....	27	8	20	21	16	29	121
19 Brown Creeper.....	18	15	19	14	28	27	121
20 Catbird.....	22	27	16	17	10	17	109
21 Golden-crowned Kinglet....	18	14	13	26	15	23	109
22 Chipping Sparrow.....	25	11	11	12	17	32	108
23 American Goldfinch.....	21	19	12	20	5	18	95
24 Tree Swallow.....	13	11	16	19	13	22	94
25 American Redstart.....	18	16	19	8	10	20	91
26 Barn Swallow.....	15	14	10	19	14	19	91
27 Fox Sparrow.....	11	9	14	9	27	21	91
28 Chimney Swift.....	16	11	14	15	11	22	89
29 Blue-gray Gnatcatcher....	18	16	11	10	16	17	88
30 White-crowned Sparrow....	9	5	18	22	16	17	87
31 Palm Warbler.....	8	21	7	11	19	20	86
32 Hermit Thrush.....	13	9	10	18	9	27	86
33 Black and White Creeper..	15	15	11	13	12	18	84
34 Least Flycatcher.....	18	18	8	14	7	18	83
35 Magnolia Warbler.....	13	13	13	5	16	20	80
36 B'k-throated Blue Warbler	12	14	9	13	12	16	76
37 Olive-backed Thrush.....	8	4	14	16	11	23	76
38 Phoebe.....	15	4	5	16	15	19	74
39 Baltimore Oriole.....	18	6	10	15	11	13	73
40 Grinnell Water-Thrush....	17	6	9	14	8	17	71
41 Field Sparrow.....		2	12	17	23	17	71
42 Swamp Sparrow.....	4	4	14	12	17	17	68
43 Crow.....	4	11	7	11	16	18	67

NUMBER OF MORNINGS SEEN
IN LINCOLN PARK, CHICAGO

	1898	1899	1900	1901	1902	1903	Total
44 Chestnut-sided Warbler....	11	6	11	13	9	16	66
45 Oven-bird.....	9	11	11	12	11	11	65
46 White-breasted Nuthatch...	12	6	17	21	6	2	64
47 Maryland Yellow-throat....	18	17	10	8	3	15	61
48 Wood Pewee.....	12	14	9	6	7	13	61
49 Downy Woodpecker.....	14	7	4	8	17	10	60
50 House Wren.....	1	...	10	11	13	24	59
51 Red-eyed Vireo.....	13	10	5	6	6	16	56
52 Loggerhead Shrike.....	5	1	16	13	7	13	55
53 Indigo Bunting.....	12	17	2	6	5	11	53
54 Bl'k-throated Green Warbler	4	9	4	9	6	12	44
55 Blackburnian Warbler....	11	4	9	7	5	6	42
56 Black-poll Warbler.....	6	7	8	3	3	15	42
57 Kingbird.....	8	1	5	3	6	17	40
58 Scarlet Tanager.....	7	3	7	4	3	10	34
59 Ruby-thr. Humming-bird..	7	9	4	3	1	10	34
60 Mourning Dove.....	5	3	6	10	2	8	34
61 Rose-breasted Grosbeak....	5	2	6	8	6	6	33
62 Tree Sparrow.....	...	...	10	3	12	8	33
63 Winter Wren.....	3	5	11	...	3	10	32
64 Meadowlark.....	3	...	7	13	5	4	32
65 Purple Martin.....	1	...	...	6	4	20	31
66 Cape May Warbler.....	3	2	9	2	6	7	29
67 Warbling Vireo.....	7	1	4	3	4	10	29
68 Mourning Warbler.....	5	8	2	4	4	5	28
69 Wilson Black-cap Warbler.	9	1	3	2	5	8	28
70 Yellow-billed Cuckoo.....	8	7	1	3	1	7	27
71 Crested Flycatcher.....	5	2	...	1	10	9	27
72 Bobolink.....	2	...	9	10	1	4	26
73 Canadian Warbler.....	8	3	6	4	1	4	26
74 Red-winged Blackbird....	4	...	2	8	3	7	24
75 Wood Thrush.....	8	3	...	5	3	3	22
76 Orchard Oriole.....	4	3	5	1	1	6	20
77 Red-breasted Nuthatch....	...	...	10	...	3	7	20
78 Marsh Hawk.....	...	2	1	8	5	4	20
79 Hairy Woodpecker.....	3	1	4	...	4	7	19
80 Bay-breasted Warbler.....	1	1	2	1	4	10	19
81 Parula Warbler.....	6	...	...	2	9	2	19
82 Chickadee.....	3	...	7	8	...	...	18
83 Pine Warbler.....	...	...	3	1	5	9	18
84 Henslow Sparrow.....	3	2	1	2	3	6	17
85 Nashville Warbler.....	1	1	4	1	...	9	16
86 Yellow-throated Vireo.....	2	...	2	2	3	7	16

NUMBER OF MORNINGS SEEN
IN LINCOLN PARK, CHICAGO

	1898	89	1900	1901	1902	1903	Total
87 Cedar Waxwing.....	6	5	...	1	2	1	15
88 Purple Finch.....	4	...	4	...	1	5	14
89 Vesper Sparrow.....	...	...	6	...	4	3	13
90 Blue-headed Vireo.....	...	...	2	1	6	4	13
91 Gray-cheeked Thrush.....	...	...	...	...	3	10	13
92 American Sparrow Hawk....	1	1	...	4	2	4	12
93 Yellow-bellied Flycatcher...	...	...	5	...	2	4	11
94 Nighthawk.....	...	2	1	1	...	7	11
95 Lincoln Sparrow.....	...	...	1	...	1	8	10
96 Black-billed Cuckoo.....	...	1	2	2	2	2	9
97 Rusty Blackbird.....	...	2	...	3	...	2	7
98 Yellow-breasted Chat.....	...	...	...	...	1	6	7
99 Long-billed Marsh Wren.....	1	2	...	...	...	3	6
100 Louisiana Water-Thrush....	...	...	1	...	3	2	6
101 Prothonotary Warbler.....	...	1	...	...	3	1	5
102 Savanna Sparrow.....	...	...	...	...	...	5	5
103 Blue Golden-winged Warb..	1	...	1	1	1	..	4
104 Whip-poor-will.....	1	1	...	1	...	1	4
105 Connecticut Warbler.....	...	...	...	...	...	3	3
106 Dickcissel.....	...	...	...	1	2	...	3
107 Tennessee Warbler.....	2	...	...	...	...	...	2
108 Mockingbird.....	...	...	...	2	...	...	2
109 Cerulean Warbler.....	...	...	1	...	1	...	2
110 Summer Tanager.....	...	1	...	...	...	...	1
111 Cardinal.....	...	...	1	...	...	...	1
112 Red-bellied Woodpecker....	1	...	...	...	...	...	1
113 Hooded Warbler.....	...	...	...	...	...	1	1
114 Yellow-headed Blackbird...	...	...	...	1	...	...	1
Total number of days.....	66	77	74	78	72	87	454
Av'ge No. of kinds seen daily	18.83	12.75	14.59	16.23	16.45	19.53	16.34

Table of Arríbal.

NOTE:—The following table gives the average date of the first arrival of each bird together with the data from which it was reckoned, for the seven years from 1897 to 1903 inclusive, *as observed in Lincoln Park*, Chicago, by the authors. The record for 1902, in the absence of the authors, was kept by Mr. Harold V. Bozell.

Attention is called to the fact that many birds which are common in the neighboring country are rare in the Park. Among these may be mentioned the *Red-winged Blackbird*, *Bobolink*, *Warbling Vireo*, *Prairie Horned Lark* and *Tufted Titmouse*.

	Average First Appearance	MARCH					APRIL					MAY				
		1897	1898	1899	1900	1901	1902	1903	1897	1898	1899	1900	1901	1902	1903	
Family COLUMBIDAE																
Mourning Dove.....	April 15.			25							6	10	26	1	9	
Family FALCONIDAE.																
Marsh Hawk.....	April 12.				25						25	19	21	1		
Sparrow Hawk.....	April 17.				24	21					25	18		30	18	
Family CUCULIDAE																
Yellow-billed Cuckoo.....	May 14														19	1 16 17 17 12
Black-billed Cuckoo.....	May 16														27 12 15 16 12	
Family ALCEDINIDAE...																
Belted Kingfisher.....	April 1.	28	25	22	26				10		11	10				
Family PICIDAE																
Hairy Woodpecker.....	April 11.				25	6					11	19			24	
Downy Woodpecker.....	March 30			27	10	6				7	8	25	5			
Yellow-bellied Sapsucker.....	April 3			30	19	27				25		8	6	1		
Red-bellied Woodpecker.....	May 22														22	
Red-headed Woodpecker.....	April 29									26	29	28	27	27	28	6
Flicker.....	March 31		25		23	24	18	15	11	7						
Family CAPRIMULGIDAE																
Whip-poor-will.....	May 12														21	9
Nighthawk.....	May 15										30					10 13 18 18 18
Family MICROPODIDAE																
Chimney Swift.....	April 29										30		30	17	22	1 11
Family TROCHILIDAE																
Ruby-throated Hummingbird.....	May 15														10	9 13 16 17 13

Family TYRANNIDAE

Kingbird.....	May 3	29 30	21	10	6 10	5
Crested Flycatcher.....	May 3	30	21 29	3	11	..
Phoebe.....	March 27	18 15 17	17 1	..	..	..
Wood Pewee.....	May 9	27	28	19 10 11 16	12	..
Yellow-bellied Flycatcher.....	May 4	..	29	..	1 12	..
Least Flycatcher.....	May 4	30 20 30	21 12	..	6	9

Family CORVIDAE

Blue Jay.....	Resident	..	..	..	..	..
Crow.....	March 11	9 26 12	3 6 8	..	..	..

Family ICTERIDAE

Bobolink.....	May 9	..	..	14	8	4 12 8
Cowbird.....	March 23	16	19 1 14	4 10 1	..	..
Yellow-headed Blackbird.....	May 17	..	..	..	17	..
Red-winged Blackbird.....	April 5	..	22 26 17	18	7	..
Meadowlark.....	March 17	14	27 19 10 13	..	..	..
Orchard Oriole.....	May 8	..	..	..	15	6 11 10 2 5
Baltimore Oriole.....	May 3	..	..	..	2	6 4 2 5
Rusty Blackbird.....	March 31	..	30	..	..	..
Bronzed Grackle.....	March 20	29 14 20 24 23 10 14	17	19 5	..	..

Family FRINGILLIDAE

Purple Finch.....	April 16	..	24	15	11 15	..
American Goldfinch.....	May 2	..	25	28	30	6 7 2 7
Vesper Sparrow.....	April 14	..	..	22	18 1	..
Savanna Sparrow.....	March 19	..	19	..	..	..
Henslow Sparrow.....	April 24	..	29 26 21 30 24 15	..	..	..
White-crowned Sparrow.....	April 23	..	11	29	20 29	5 9 2
White-throated Sparrow.....	April 13	..	24 21 26 13 19 21 17	..	..	..
Tree Sparrow.....	March 23	..	7 13	6 5	..	..
Chipping Sparrow.....	April 12	..	15 13 18 8	8 13 12	..	..
Field Sparrow.....	April 3	..	11 19	23 17 5	..	..
Junco.....	March 16	28 9 14 23 18	8 13	..	..	..
Song Sparrow.....	March 20	15 19 27 18	8 12	8	..	..
Lincoln Sparrow.....	May 14	..	..	..	10 23 12	10
Swamp Sparrow.....	April 14	..	11	20 21 18	12	7

	Average First Appearance	MARCH			APRIL			MAY		
		1896	1898	1901	1902	1907	1896	1898	1901	1907
Family FRINGILLIDAE—Continued										
Fox Sparrow.....	March 23	..	30 14	18 10 17	8 ..	5 ..	..	..	..	..
Towhee.....	March 27	28	17 ..	25 10 18	7 .. 11	7 ..	..	..	..	..
Cardinal.....	April 11	..	..	..	..	25 ..	..	..	..	..
Rose-breasted Grosbeak.....	May 7	..	..	..	..	28 ..	..	..	..	..
Indigo Bunting.....	May 11	..	..	..	..	..	10 11	4 ..	9 ..	..
Dickcissel.....	May 17	..	..	..	..	..	14 10 21	10 2 6	16 18	..
Family TANAGRIDAE										
Scarlet Tanager.....	May 5	..	..	..	..	..	14 4	2 5	3 5	..
Summer Tanager.....	May 4	..	..	..	..	..	..	4 ..	..	..
Family HIRUNDINIDAE										
Purple Martin.....	May 6	..	..	..	..	27 23 17	25 30	..	..	..
Barn Swallow.....	April 21	..	..	..	..	18 ..	23 26 10 18 24	1 ..	..	..
Tree Swallow.....	April 11	..	..	11 ..	..	12 18 12 ..	1 ..	..	..	..
Family BOMBYCILLIDAE										
Cedar Waxwing.....	April 16	..	27 ..	..	..	11 ..	24 5	16 ..	..	..
Family LANIIDAE										
Loggerhead Shrike.....	March 13	..	14 ..	9 3 9 1	..	11 ..	..	..	..	..
Family VIREONIDAE										
Red-eyed Vireo.....	May 5	..	..	..	..	29 ..	..	10 ..	12 4 2 4	..
Warbling Vireo.....	May 8	..	..	..	..	27 ..	..	15 12 6 11	7 5	..
Yellow-throated Vireo.....	May 3	..	..	..	..	..	28 ..	1 ..	12 5 12	..
Blue-headed Vireo.....	May 8	..	..	..	..	..	..	17 5	2 8	..
Family MNIOTILTIDAE										
Black and White Creeper.....	April 28.	..	..	..	..	28 30 21 26	28 27	..	4 ..	..
Prothonotary Warbler.....	May 11	..	..	..	..	..	27 ..	16 ..	..	19

	Average First Appearance	MARCH					APRIL					MAY					
		1897	1898	1899	1900	1901	1897	1898	1899	1900	1901	1897	1898	1899	1900	1901	1902
Family PARIDAE																	
White-breasted Nuthatch.....	April 1	29			18 31	7											
Red-breasted Nuthatch.....	April 28																
Chickadee.....	April 1			19				1		13					3		
Family SYLVIIDAE																	
Golden-crowned Kinglet.....	March 31	14			28 19	7			11	6 5							
Ruby-crowned Kinglet.....	April 8				19 14	11 3	7 17 13										
Blue-gray Gnatcatcher.....	April 23					25 22	18 28 22 25										
Family TURDIDAE																	
Wood Thrush.....	May 1					18				29			3	1	4		12
Wilson Thrush.....	April 16																
Gray-checked Thrush.....	May 16						10 12	7 11 26	8 11								
Olive-backed Thrush.....	May 1															21 11	
Hermit Thrush.....	April 4	31				19				30 22 29			1	4	8		
Robin.....	March 15	28	9	14 24	14 10	7				7 5 13							
Bluebird.....	March 18	28	8	31 17	7	7			4								

Notes.

Table of Distribution and Occurrence.

The following table has been compiled in the hope that wherever the bird student may be in the northern states, east of the Mississippi river, he may have at hand for comparison the results of some reliable observer in the same or a nearby area. No better method of gaining a clear idea of the movements of birds can be suggested than to study carefully their distribution and occurrence within related areas. In order to do this understandingly, the main topographic features, taken as a whole or state by state, should be considered. The bird population of any region will be found to be closely related to its mountains, river basins, prairies, forests and wet or dry areas.

Attention is called to the meeting and intermingling of the Carolinian and Alleghenian faunal areas in these northeastern states. The teacher of geography as well as the bird student will find great interest in following out the limits of life conditions in these areas.

The table contains in addition to the 200 species already described, 148 species of frequent but rarer occurrence throughout the area covered. It should be said that considerable difficulty lies in the way of reducing to a few arbitrary symbols the notes of several ornithologists whose data are not equally complete and who may not use the same descriptive terms, or who may attach a different significance to the same terms. The authors nevertheless have earnestly tried to interpret correctly each authority cited and trust that the useful features of the result will outweigh its shortcomings.

The *varying dates* of the published lists for the different localities given should be kept constantly in mind in making

comparisons with new lists, because the bird population of any locality is as truly changeable, although for different reasons, as its human population.

The data given have been compiled from the following published lists:—

1. **Maine.** Knight, O. W., *Birds of Maine.* 1908.
2. **New Hampshire.** Allen, Glover M., *A List of the Birds of New Hampshire.* 1903.
3. **Vermont.** Perkins, G. H., and Howe, C. D., *A Preliminary List of the Birds found in Vermont.* 1901.
4. **Massachusetts.** Brewster, Wm., *The Birds of the Cambridge Region of Massachusetts.* 1906.
5. **Massachusetts.** Clarke, H. L., *The Birds of Amherst and Vicinity*, 2nd edition. 1906.
6. **Rhode Island.** Howe, R. H. Jr., and Sturtevant E., *The Birds of Rhode Island.* 1899. Supplement, 1903.
7. **Long Island.** Braislin, W. C., *A List of the Birds of Long Island, N. Y.* 1907.
8. **New York City.** Chapman, F. M., *The Birds of the Vicinity of New York City.* 1906.
9. **Pennsylvania and New Jersey.** Stone, Witmer, *The Birds of Eastern Pennsylvania and New Jersey.* 1894.
10. **Ohio.** Jones, Lynds., *The Birds of Ohio.* 1903.
11. **Indiana.** Butler, A. W., *The Birds of Indiana.* 1897.
12. **Illinois.** Woodruff, F. M., *The Birds of the Chicago Area.* 1907.
13. **Missouri.** Widmann, Otto, *A Preliminary Catalog of the Birds of Missouri.* 1907.

KEY TO TABLE OF DISTRIBUTION AND OCCURRENCE.

Column I Number

- A.....abundant.
- a.....common or very common.
- F.....frequent, rather common, not uncommon.
- f.....infrequent, not common, rather rare, uncommon.
-rare.
- =.....very rare or casual.
- x.....accidental, out of normal range.
- i.....irregular, from year to year common, rare or absent.
- (blank)..no record.
- d.....doubtful, records scanty or incomplete.

Column II Status

- R.....Permanent Resident. (May be partially migratory, increasing in number during the spring or fall or both; or within the limits of a single state withdrawing largely to the north in summer and to the south in winter in search of special breeding and feeding areas.)
- R.....Permanent Resident most abundant in winter.
- r.....Winter Resident.
- P.....Winter Resident with a few in summer.
- s.....Summer Resident.
- S.....Summer Resident with a few in winter.
- V.....Winter Visitant.
- V.....Winter Visitant with a few in summer.
- v.....Summer Visitant.
- T.....Regular Transient, a spring and fall migrant.
- t.....Regular Transient, most abundant in fall.
- 3.....Regular Transient most abundant in spring.
- I.....Regular Fall Transient, not recorded in Spring.
- I.....Regular Transient with some wintering locally.
- II.....Regular Transient with some summering, or breeding locally.
- III.....Regular Transient with a few winter and summer.
- IV.....Regular Transient frequent or common in winter, infrequent or rare in summer.
- X.....Regular Transient frequent or common in summer, rare in winter.
- d.....doubtful, may occur or probably occurs.
- ?.....no record within recent years. (usually indicates species formerly common).
- D.....an introduced species.

Column III Distribution

- g.....general throughout area cited.
- L.....not general, confined mostly or entirely to particular regions within area cited.
- l.....general, but restricted to particular breeding environments. (r. g. swamps dry pasture, second growth, etc.) or, general, but for no apparent reason absent from many localities throughout area cited, hence local.
- c.....found on the coast.
- C.....found largely on the coast, a few inland.
- o.....found largely inland, a few on the coast.
- n.....inland.
- m.....confined mostly to mountains (High altitudes are equivalent to northern latitudes).
- ?.....distribution doubtful.
- i.....irregular, or erratic, likely to occur at any point. (Usually applies to species going in flocks).

Figures in any column indicate number of recorded instances of rare species.

Symbols are occasionally placed without reference to the column-title in order to express the data given.

117. 4 cm. x 10 cm.
 122. South-western Cormorant
 123. American White Pelican
 129. American M.
 130. Red-breasted
 131. Hatched M.

1
 2
 3
 4
 5
 6
 7
 8
 9
 10
 11
 12
 13
 14
 15
 16
 17
 18
 19
 20
 21
 22
 23
 24
 25
 26
 27
 28
 29
 30
 31
 32
 33
 34
 35
 36
 37
 38
 39
 40
 41
 42
 43
 44
 45
 46
 47
 48
 49
 50
 51
 52
 53
 54
 55
 56
 57
 58
 59
 60
 61
 62
 63
 64
 65
 66
 67
 68
 69
 70
 71
 72
 73
 74
 75
 76
 77
 78
 79
 80
 81
 82
 83
 84
 85
 86
 87
 88
 89
 90
 91
 92
 93
 94
 95
 96
 97
 98
 99
 100

hard			
mill			
mill			
at the			hard
at the			
the			

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80																				

1. Name of the person or organization
 2. Address
 3. City
 4. State
 5. Zip

6. Date
 7. Time
 8. Location

9. Description of the event
 10. Details of the event

11. Name of the person or organization
 12. Address

13. City
 14. State

15. Zip
 16. Date

17. Time
 18. Location

19. Description of the event
 20. Details of the event

21. Name of the person or organization
 22. Address

23. City
 24. State

25. Zip
 26. Date

27. Time
 28. Location

29. Description of the event
 30. Details of the event

31. Name of the person or organization
 32. Address

33. City
 34. State

Western Pa.
 477
 478 Black-throated Blue Warbler
 479 Monte Warbler

H P
 7 11
 8 12
 9 13
 10 14
 11 15
 12 16
 13 17
 14 18
 15 19
 16 20
 17 21
 18 22
 19 23
 20 24
 21 25
 22 26
 23 27
 24 28
 25 29
 26 30
 27 31
 28 32
 29 33
 30 34
 31 35
 32 36
 33 37
 34 38
 35 39
 36 40
 37 41
 38 42
 39 43
 40 44
 41 45
 42 46
 43 47
 44 48
 45 49
 46 50
 47 51
 48 52
 49 53
 50 54
 51 55
 52 56
 53 57
 54 58
 55 59
 56 60
 57 61
 58 62
 59 63
 60 64
 61 65
 62 66
 63 67
 64 68
 65 69
 66 70
 67 71
 68 72
 69 73
 70 74
 71 75
 72 76
 73 77
 74 78
 75 79
 76 80
 77 81
 78 82
 79 83
 80 84
 81 85
 82 86
 83 87
 84 88
 85 89
 86 90
 87 91
 88 92
 89 93
 90 94
 91 95
 92 96
 93 97
 94 98
 95 99
 96 100
 97 101
 98 102
 99 103
 100 104
 101 105
 102 106
 103 107
 104 108
 105 109
 106 110
 107 111
 108 112
 109 113
 110 114
 111 115
 112 116
 113 117
 114 118
 115 119
 116 120
 117 121
 118 122
 119 123
 120 124
 121 125
 122 126
 123 127
 124 128
 125 129
 126 130
 127 131
 128 132
 129 133
 130 134
 131 135
 132 136
 133 137
 134 138
 135 139
 136 140
 137 141
 138 142
 139 143
 140 144
 141 145
 142 146
 143 147
 144 148
 145 149
 146 150
 147 151
 148 152
 149 153
 150 154
 151 155
 152 156
 153 157
 154 158
 155 159
 156 160
 157 161
 158 162
 159 163
 160 164
 161 165
 162 166
 163 167
 164 168
 165 169
 166 170
 167 171
 168 172
 169 173
 170 174
 171 175
 172 176
 173 177
 174 178
 175 179
 176 180
 177 181
 178 182
 179 183
 180 184
 181 185
 182 186
 183 187
 184 188
 185 189
 186 190
 187 191
 188 192
 189 193
 190 194
 191 195
 192 196
 193 197
 194 198
 195 199
 196 200
 197 201
 198 202
 199 203
 200 204
 201 205
 202 206
 203 207
 204 208
 205 209
 206 210
 207 211
 208 212
 209 213
 210 214
 211 215
 212 216
 213 217
 214 218
 215 219
 216 220
 217 221
 218 222
 219 223
 220 224
 221 225
 222 226
 223 227
 224 228
 225 229
 226 230
 227 231
 228 232
 229 233
 230 234
 231 235
 232 236
 233 237
 234 238
 235 239
 236 240
 237 241
 238 242
 239 243
 240 244
 241 245
 242 246
 243 247
 244 248
 245 249
 246 250
 247 251
 248 252
 249 253
 250 254
 251 255
 252 256
 253 257
 254 258
 255 259
 256 260
 257 261
 258 262
 259 263
 260 264
 261 265
 262 266
 263 267
 264 268
 265 269
 266 270
 267 271
 268 272
 269 273
 270 274
 271 275
 272 276
 273 277
 274 278
 275 279
 276 280
 277 281
 278 282
 279 283
 280 284
 281 285
 282 286
 283 287
 284 288
 285 289
 286 290
 287 291
 288 292
 289 293
 290 294
 291 295
 292 296
 293 297
 294 298
 295 299
 296 300
 297 301
 298 302
 299 303
 300 304
 301 305
 302 306
 303 307
 304 308
 305 309
 306 310
 307 311
 308 312
 309 313
 310 314
 311 315
 312 316
 313 317
 314 318
 315 319
 316 320
 317 321
 318 322
 319 323
 320 324
 321 325
 322 326
 323 327
 324 328
 325 329
 326 330
 327 331
 328 332
 329 333
 330 334
 331 335
 332 336
 333 337
 334 338
 335 339
 336 340
 337 341
 338 342
 339 343
 340 344
 341 345
 342 346
 343 347
 344 348
 345 349
 346 350
 347 351
 348 352
 349 353
 350 354
 351 355
 352 356
 353 357
 354 358
 355 359
 356 360
 357 361
 358 362
 359 363
 360 364
 361 365
 362 366
 363 367
 364 368
 365 369
 366 370
 367 371
 368 372
 369 373
 370 374
 371 375
 372 376
 373 377
 374 378
 375 379
 376 380
 377 381
 378 382
 379 383
 380 384
 381 385
 382 386
 383 387
 384 388
 385 389
 386 390
 387 391
 388 392
 389 393
 390 394
 391 395
 392 396
 393 397
 394 398
 395 399
 396 400
 397 401
 398 402
 399 403
 400 404
 401 405
 402 406
 403 407
 404 408
 405 409
 406 410
 407 411
 408 412
 409 413
 410 414
 411 415
 412 416
 413 417
 414 418
 415 419
 416 420
 417 421
 418 422
 419 423
 420 424
 421 425
 422 426
 423 427
 424 428
 425 429
 426 430
 427 431
 428 432
 429 433
 430 434
 431 435
 432 436
 433 437
 434 438
 435 439
 436 440
 437 441
 438 442
 439 443
 440 444
 441 445
 442 446
 443 447
 444 448
 445 449
 446 450
 447 451
 448 452
 449 453
 450 454
 451 455
 452 456
 453 457
 454 458
 455 459
 456 460
 457 461
 458 462
 459 463
 460 464
 461 465
 462 466
 463 467
 464 468
 465 469
 466 470
 467 471
 468 472
 469 473
 470 474
 471 475
 472 476
 473 477
 474 478
 475 479
 476 480
 477 481
 478 482
 479 483
 480 484
 481 485
 482 486
 483 487
 484 488
 485 489
 486 490
 487 491
 488 492
 489 493
 490 494
 491 495
 492 496
 493 497
 494 498
 495 499
 496 500
 497 501
 498 502
 499 503
 500 504
 501 505
 502 506
 503 507
 504 508
 505 509
 506 510
 507 511
 508 512
 509 513
 510 514
 511 515
 512 516
 513 517
 514 518
 515 519
 516 520
 517 521
 518 522
 519 523
 520 524
 521 525
 522 526
 523 527
 524 528
 525 529
 526 530
 527 531
 528 532
 529 533
 530 534
 531 535
 532 536
 533 537
 534 538
 535 539
 536 540
 537 541
 538 542
 539 543
 540 544
 541 545
 542 546
 543 547
 544 548
 545 549
 546 550
 547 551
 548 552
 549 553
 550 554
 551 555
 552 556
 553 557
 554 558
 555 559
 556 560
 557 561
 558 562
 559 563
 560 564
 561 565
 562 566
 563 567
 564 568
 565 569
 566 570
 567 571
 568 572
 569 573
 570 574
 571 575
 572 576
 573 577
 574 578
 575 579
 576 580
 577 581
 578 582
 579 583
 580 584
 581 585
 582 586
 583 587
 584 588
 585 589
 586 590
 587 591
 588 592
 589 593
 590 594
 591 595
 592 596
 593 597
 594 598
 595 599
 596 600
 597 601
 598 602
 599 603
 600 604
 601 605
 602 606
 603 607
 604 608
 605 609
 606 610
 607 611
 608 612
 609 613
 610 614
 611 615
 612 616
 613 617
 614 618
 615 619
 616 620
 617 621
 618 622
 619 623
 620 624
 621 625
 622 626
 623 627
 624 628
 625 629
 626 630
 627 631
 628 632
 629 633
 630 634
 631 635
 632 636
 633 637
 634 638
 635 639
 636 640
 637 641
 638 642
 639 643
 640 644
 641 645
 642 646
 643 647
 644 648
 645 649
 646 650
 647 651
 648 652
 649 653
 650 654
 651 655
 652 656
 653 657
 654 658
 655 659
 656 660
 657 661
 658 662
 659 663
 660 664
 661 665
 662 666
 663 667
 664 668
 665 669
 666 670
 667 671
 668 672
 669 673
 670 674
 671 675
 672 676
 673 677
 674 678
 675 679
 676 680
 677 681
 678 682
 679 683
 680 684
 681 685
 682 686
 683 687
 684 688
 685 689
 686 690
 687 691
 688 692
 689 693
 690 694
 691 695
 692 696
 693 697
 694 698
 695 699
 696 700
 697 701
 698 702
 699 703
 700 704
 701 705
 702 706
 703 707
 704 708
 705 709
 706 710
 707 711
 708 712
 709 713
 710 714
 711 715
 712 716
 713 717
 714 718
 715 719
 716 720
 717 721
 718 722
 719 723
 720 724
 721 725
 722 726
 723 727
 724 728
 725 729
 726 730
 727 731
 728 732
 729 733
 730 734
 731 735
 732 736
 733 737
 734 738
 735 739
 736 740
 737 741
 738 742
 739 743
 740 744
 741 745
 742 746
 743 747
 744 748
 745 749
 746 750
 747 751
 748 752
 749 753
 750 754
 751 755
 752 756
 753 757
 754 758
 755 759
 756 760
 757 761
 758 762
 759 763
 760 764
 761 765
 762 766
 763 767
 764 768
 765 769
 766 770
 767 771
 768 772
 769 773
 770 774
 771 775
 772 776
 773 777
 774 778
 775 779
 776 780
 777 781
 778 782
 779 783
 780 784
 781 785
 782 786
 783 787
 784 788
 785 789
 786 790
 787 791
 788 792
 789 793
 790 794
 791 795
 792 796
 793 797
 794 798
 795 799
 796 800
 797 801
 798 802
 799 803
 800 804
 801 805
 802 806
 803 807
 804 808
 805 809
 806 810
 807 811
 808 812
 809 813
 810 814
 811 815
 812 816
 813 817
 814 818
 815 819
 816 820
 817 821
 818 822
 819 823
 820 824
 821 825
 822 826
 823 827
 824 828
 825 829
 826 830
 827 831
 828 832
 829 833
 830 834
 831 835
 832 836
 833 837
 834 838
 835 839
 836 840
 837 841
 838 842
 839 843
 840 844
 841 845
 842 846
 843 847
 844 848
 845 849
 846 850
 847 851
 848 852
 849 853
 850 854
 851 855
 852 856
 853 857
 854 858
 855 859
 856 860
 857 861
 858 862
 859 863
 860 864
 861 865
 862 866
 863 867
 864 868
 865 869
 866 870
 867 871
 868 872
 869 873
 870 874
 871 875
 872 876
 873 877
 874 878
 875 879
 876 880
 877 881
 878 882
 879 883
 880 884
 881 885
 882 886
 883 887
 884 888
 885 889
 886 890
 887 891
 888 892
 889 893
 890 894
 891 895
 892 896
 893 897
 894 898
 895 899
 896 900
 897 901
 898 902
 899 903
 900 904
 901 905
 902 906
 903 907
 904 908
 905 909
 906 910
 907 911
 908 912
 909 913
 910 914
 911 915
 912 916
 913 917
 914 918
 915 919
 916 920
 917 921
 918 922
 919 923
 920 924
 921 925
 922 926
 923 927
 924 928
 925 929
 926 930
 927 931
 928 932
 929 933
 930 934
 931 935
 932 936
 933 937
 934 938
 935 939
 936 940
 937 941
 938 942
 939 943
 940 944
 941 945
 942 946
 943 947
 944 948
 945 949
 946 950
 947 951
 948 952
 949 953
 950 954
 951 955
 952 956
 953 957
 954 958
 955 959
 956 960
 957 961
 958 962
 959 963
 960 964
 961 965
 962 966
 963 967
 964 968
 965 969
 966 970
 967 971
 968 972
 969 973
 970 974
 971 975
 972 976
 973 977
 974 978
 975 979
 976 980
 977 981
 978 982
 979 983
 980 984
 981 985
 982 986
 983 987
 984 988
 985 989
 986 990
 987 991
 988 992
 989 993
 990 994
 991 995
 992 996
 993 997
 994 998
 995 999
 996 1000
 997 1001
 998 1002
 999 1003
 1000 1004
 1001 1005
 1002 1006
 1003 1007
 1004 1008
 1005 1009
 1006 1010
 1007 1011
 1008 1012
 1009 1013
 1010 1014
 1011 1015
 1012 1016
 1013 1017
 1014 1018
 1015 1019
 1016 1020
 1017 1021
 1018 1022
 1019 1023
 1020 1024
 1021 1025
 1022 1026
 1023 1027
 1024 1028
 1025 1029
 1026 1030
 1027 1031
 1028 1032
 1029 1033
 1030 1034
 1031 1035
 1032 1036
 1033 1037
 1034 1038
 1035 1039
 1036 1040
 1037 1041
 1038 1042

1. *Hydrolysis of the polymer*
 2. *Hydrolysis of the polymer*

Fre. Henry.

Tri White-browed Nuthatch

116 Red-breasted Nuthatch

Memo.

Index of Common Names

Bittern, American.....	19	Purple.....	35
Least.....	19	Grebe, Horned.....	13
Blackbird, Red-winged.....	34	Pied-billed.....	13
Rusty.....	35	Grouse, Ruffed.....	25
Yellow-headed... ..	34	Grosbeak, Rose-breasted.....	39
Bluebird.....	52	Gull, Herring.....	14
Bobolink.....	33	Bonaparte.....	14
Bob-white.....	52	Ring-billed.....	14
Buffle-head.....	18	Hawk, American Sparrow.....	27
Bunting, Indigo.....	39	Broad-winged.....	27
Cardinal.....	39	Cooper.....	26
Catbird.....	49	Marsh.....	26
Chat, Yellow-breasted.....	48	Red-tailed.....	26
Chickadee.....	50	Red-shouldered.....	26
Coot, American.....	21	Sharp-shinned.....	26
Cowbird.....	34	Heron, Black-crowned Night..	20
Creepers, Black and White.....	42	Great Blue.....	19
Brown.....	50	Green.....	20
Crossbill, American.....	35	Hummingbird, Ruby-throated..	31
Crow.....	33	Jay, Blue.....	33
Fish.....	33	Junco.....	38
Cuckoo, Black-billed.....	29	Killdeer.....	24
Yellow-billed.....	29	Kingbird.....	31
Dickcissel.....	39	Kingfisher, Belted.....	29
Dove, Mourning.....	25	Kinglet, Golden-crowned.....	51
Duck, American Scaup.....	17	Ruby-crowned.....	51
Black.....	16	Lark, Horned.....	33
Lesser Scaup.....	17	Prairie Horned.....	33
Ruddy.....	19	Loon.....	13
Wood.....	17	Mallard.....	15
Eagle, Bald.....	27	Martin, Purple.....	40
Finch, Purple.....	35	Meadowlark.....	34
Flicker.....	30	Merganser, American.....	15
Flycatcher, Crested.....	31	Hooded.....	15
Green-crested.....	32	Red-breasted.....	15
Least.....	32	Mockingbird.....	49
Olive-sided.....	31	Nighthawk.....	30
Traill.....	32	Nuthatch, Red-breasted.....	50
Yellow-bellied.....	32	White-breasted.....	50
Gallinule, Florida.....	21	Old-Squaw.....	18
Gnat-catcher, Blue-gray.....	51	Oriole, Baltimore.....	34
Golden-eye, American.....	17	Orchard.....	34
Goldfinch, American.....	36	Osprey, American.....	27
Goose, Canada.....	19	Oven-bird.....	46
Grackle, Bronzed.....	35		

Index of Common Names—Continued

Owl, American Long-eared.....	27	Cliff.....	40
Barred.....	28	Tree.....	40
Great Horned.....	28	Swift, Chimney.....	31
Saw-whet.....	28	Tanager, Scarlet.....	40
Screech.....	28	Summer.....	40
Short-eared.....	28	Teal, Blue-winged.....	16
Pewee, Wood.....	32	Green-winged.....	16
Phoebe.....	31	Tern, Common.....	14
Pintail.....	16	Thrasher, Brown.....	49
Plover, American Golden.....	23	Thrush, Gray-cheeked.....	51
Semi-palmated.....	24	Hermit.....	52
Pipit, American.....	48	Olive-backed.....	52
Rail, Virginia.....	20	Wilson.....	51
Red-head.....	17	Wood.....	51
Redpoll.....	35	Titmouse, Tufted.....	50
Redstart, American.....	48	Towhee.....	39
Robin.....	52	Vireo, Blue-headed.....	42
Sanderling.....	22	Red-eyed.....	41
Sandpiper, Bartramian.....	23	Warbling.....	42
Least.....	22	White-eyed.....	42
Pectoral.....	22	Yellow-throated.....	42
Semipalmated.....	22	Warbler, Bay-breasted.....	45
Solitary.....	23	Blackburnian.....	45
Spotted.....	23	Black-poll.....	45
Sapsucker, Yellow-bellied.....	30	Black-throated, Blue.....	44
Scoter, White-winged.....	18	Black-throated, Green.....	45
Shrike, Loggerhead.....	41	Blue Golden-winged.....	43
Northern.....	41	Blue-winged.....	43
Siskin, Pine.....	36	Canadian.....	48
Sora.....	20	Cape May.....	43
Snipe, Wilson.....	21	Cerulean.....	44
Snowflake.....	36	Chestnut-sided.....	44
Sparrow, Chipping.....	38	Connecticut.....	47
Field.....	38	Hooded.....	48
Fox.....	39	Magnolia.....	44
Grasshopper.....	37	Mourning.....	47
Henslow.....	37	Myrtle.....	44
Lincoln.....	38	Nashville.....	43
Savanna.....	36	Palm.....	46
Song.....	38	Northern Parula.....	43
Swamp.....	38	Prairie.....	46
Tree.....	37	Pine.....	45
Vesper.....	36	Prothonotary.....	42
White-crowned.....	37	Tennessee.....	43
White-throated.....	37	Wilson, Black-cap.....	48
Swallow, Bank.....	41	Yellow.....	44
Barn.....	40	Yellow Palm.....	46

Index of Common Names—Continued

Water-Thrush.....	46	Red-headed	30
Grinnell.....	47	Wren, Carolina.....	49
Louisiana.....	47	House.....	49
Waxwing, Cedar.....	41	Long-billed Marsh.....	50
Whip-poor-will.....	30	Winter.....	49
Woodcock, American.....	21	Yellow-legs.....	23
Woodpecker, Downy.....	29	Greater.....	22
Hairy.....	29	Yellow-throat, Maryland.....	47
Red-bellied.....	30		

Migration Record

[illegible]

Weather Signs

W = Warm
C = Cold
F = Fair
Cl = Cloudy
R = Rainy

Wind Signs

W = West
E = East
S = South
N = North

Record Sign

○ = good
X = many
* = several
+ = two
- = one

MARCH

Weather Record

Direction of Wind

19

- Bittern, American
- Least
- Blackbird, Red-winged
- Busty
- Yellow-headed
- Bluebird
- Bobolink
- Bob-white
- Bull-head
- Bunting, Indigo
- Cardinal
- Catbird
- Chaff, Yellow-bellied
- Chickadee
- Coot, American
- Cowbird
- Creeper, Black and White
- Brown
- Cross-bill, American
- Crow, Common
- Fish
- Cuckoo, Black-billed
- Yellow-billed
- Dickcissel
- Dove, Mourning
- Duck, American
- Black
- Lesser Scaup
- Ruddy
- Wood
- Eagle, Bald
- Finch, Purple
- Flicker
- Mocker, Crested
- Green, Crested
- Grass
- Olive-backed
- Tall
- Yellow-bellied
- Gallinule, Florida
- Guillemot, Blue-gray
- Golden-eye, American
- Golden-eye, American
- Goose, Canada
- Grackle, Common

SMITHSONIAN INSTITUTION LIBRARIES

3 9088 00055 5730