UNIVERSITY OF
ILLINOIS LIBRARY
AT URBANA-CHAMPAIGN
NATURAL HIST SURVEY

A Bibliography of the Northern and Western Corn Rootworm: An Update 1977 through 1988

Eli Levine
Siu Yau Chan

Illinois Natural History Survey, Lorin I. Nevling, Chief
A Division of the Illinois Department of Energy and Natural Resources

Printed by authority of the State of Illinois
X03207-2.5M-2-90
US ISSN 0073-490X

Cover: Molly Hardin Scott
Editors: Audrey S. Hodgins and Eva L. Steger

A catalog of the publications of the Illinois Natural History Survey is available without charge from the address below.
A price list and an order blank are included with the catalog.
Illinois Natural History Survey
Distribution Center
Natural Resources Building
607 East Peabody Drive
Champaign, Illinois 61820

Manuscripts of high quality dealing with any aspect of natural history will be considered for publication in one of the Illinois Natural History Survey series: *Bulletin*, *Biological Notes*, *Circular*, and *Special Publications*. Authors who are not employees of the Survey are required to pay printing costs. Manuscripts should follow the recommendations of the third edition of the *Council of Biology Editors Style Manual* except that journal names in literature cited are to be spelled in full. The Survey expects to publish only one or two manuscripts by non-Survey authors yearly. Send three copies of manuscripts to be considered for publication to Office of the Chief, Illinois Natural History Survey, 607 East Peabody Drive, Champaign, Illinois 61820. Before a manuscript is accepted for publication in the *Bulletin* or in *Biological Notes*, two or more outside referees must recommend it.

Citation: Levine, E., and S.Y. Chan. 1990. A bibliography of the northern and western corn rootworm: an update 1977 through 1988. Illinois Natural History Survey Biological Notes 135. 29 p.

Introduction

In 1974 the Illinois Natural History Survey published a bibliography of the northern and western corn rootworm. An update through 1976 was published in 1977. In the twelve years since that publication, the literature on these two insects has grown immensely. This bibliography supplements those earlier publications and covers the period from January 1977 through December 1988. Like its predecessors, this bibliography is generally limited to literature dealing with the northern corn rootworm, *Diabrotica barberi* (referred to as *D. longicornis barberi* before the subspecies was raised to species level in 1983), and the western corn rootworm, *D. virgifera virgifera*. Literature on the Mexican corn rootworm, *D. virgifera zea*, is now also included since this pest is present in the southwestern part of the United States. The 434 titles in the present list were obtained from standard reference sources (e.g., AGRI-COLA, BIOSIS Previews, CAB Abstracts, Dissertation Abstracts International, OCLC, and Science Citation Index) and from the personal files of field crop entomologists in the United States and Canada.

Entries are listed alphabetically by author and then by date of publication. Only materials published in English were selected, including journal articles, books and portions of books, M.S. and Ph.D. theses, agricultural extension publications, manuals, and fact sheets. Only the latest revisions are listed for agricultural extension publications (1989 revisions were included when 1988 versions were out of print). Literature that dealt solely with insecticide evaluations or recommendations and contained no discussion of pest biology was excluded.

Dr. Eli Levine is an Associate Entomologist in the Center for Economic Entomology at the Illinois Natural History Survey and an Associate Professor of Agricultural Entomology at the University of Illinois at Urbana-Champaign. Mr. Siu Yau Chan is a student in the Graduate School of Library and Information Science at the University.

Except for citations preceded by a ♦ symbol, copies were obtained for each document (abstracts in the case of Ph.D. theses). Key words to be included in the subject index were then assigned on the basis of content and are given within parentheses at the end of each entry. Key words taken directly from titles were listed in the index but are not repeated in the parenthetical listings. Citations preceded by a ♦ were assigned key words solely on the basis of their titles. The subject index and a list of periodicals cited appear at the end of this publication.

This bibliography (with periodic updates) will be available on diskette (5.25 or 3.5 inch) in Wordperfect or ASCII text file format for IBM-compatible computers. These diskettes are available at a cost of \$5.00 by contacting the senior author at the Natural History Survey.

Acknowledgments

The search of literature was assisted by Carla G. Heister, technical librarian of the Illinois Natural History Survey Library, and David M. Bott, graduate research assistant, University of Illinois at Urbana-Champaign. Valuable advice was obtained from Jenny Kogan, librarian for the Soybean Insect Research Information Center, Illinois Natural History Survey and University of Illinois at Urbana-Champaign, and from professors Lawrence W. Auld and Donald W. Krummel of the Graduate School of Library and Information Science, University of Illinois at Urbana-Champaign. Their assistance is gratefully acknowledged. Thanks are also extended to the many individual researchers who sent us copies of the publications of their respective states or provinces and titles of master's theses. Finally, we especially wish to recognize Audrey S. Hodgins for superb editorial assistance. This publication is a contribution of the Illinois Natural History Survey and the Illinois Agricultural Experiment Station, College of Agriculture, University of Illinois at Urbana-Champaign. The literature search was supported in part by funding from USDA Hatch Project ILLU-12-0310.

Citations

1. Agee, H.R., V.M. Kirk, and J.C. Davis. 1983. Comparative spectral sensitivity and some observations on vision related behavior of northern and western corn rootworm adults (*Diabrotica*: Coleoptera). J. Georgia Entomol. Soc. 18(2): 240–245. (*D. barberi*, *D. virgifera virgifera*)
2. Agriculture Canada. 1981. Northern corn rootworm, *Diabrotica longicornis* (Say). Agric. Canada. Insect Identification Sheet 54. 2 pp. (*D. barberi* as *D. longicornis*, biology)
3. Ahmad, N., D.D. Walgenbach, and G.R. Sutter. 1979. Comparative disappearance of fonofos, phorate and terbufos soil residues under similar South Dakota field conditions. Bull. Environ. Contam. Toxicol. 23: 423–429. (*Diabrotica* spp., insecticide degradation, insecticide persistence)
4. Ahmad, N., D.D. Walgenbach, and G.R. Sutter. 1979. Degradation rates of technical carbofuran and a granular formulation in four soils with known insecticide use history. Bull. Environ. Contam. Toxicol. 23: 572–574. (*Diabrotica* spp., insecticide persistence, insecticide degradation, insecticide history)
5. Ahmad, N., D.D. Walgenbach, and G.R. Sutter. 1979. Differential degradation rates of technical and formulated carbofuran in soil. Proc. N. D. Acad. Sci. 33: 75. (*Diabrotica* spp., insecticide degradation, insecticide persistence)
6. Ahmad, N., D.D. Walgenbach, and G.R. Sutter. 1979. Some properties of microsomal mixed function oxidase in southern and western corn rootworm larvae (abstract). Proc. N. Cent. Br. Entomol. Soc. Am. 1978, 33: 48. (*D. virgifera virgifera*)
7. Andersen, J.F. 1984. The role of attractants and feeding stimulants in the selection of *Cucurbita* flowers by Diabrotic beetle beetles (Coleoptera: Chrysomelidae). Ph.D. Thesis, University of Illinois at Urbana-Champaign. (*D. virgifera virgifera*, plant volatiles, kairomones, semiochemicals, indole, electroantennogram, phenology, sex ratio, cucurbitacins)
8. Andersen, J.F., and R.L. Metcalf. 1986. Identification of a volatile attractant for *Diabrotica* and *Acalymma* spp. from blossoms of *Cucurbita maxima* Duchesne. J. Chem. Ecol. 12(3): 687–699. (*D. virgifera virgifera*, plant volatiles, indole, electroantennogram, phenology, sex ratio, cucurbitacins)
9. Andersen, J.F., and R.L. Metcalf. 1987. Factors influencing distribution of *Diabrotica* spp. in blossoms of cultivated *Cucurbita* spp. J. Chem. Ecol. 13(4): 681–699. (*D. virgifera virgifera*, *D. barberi*, pollen, cucurbitacins, host plant selection, plant volatiles)
10. Andersen, J.F., R.D. Plattner, and D. Weisleder. 1988. Metabolic transformations of cucurbitacins by *Diabrotica virgifera virgifera* LeConte and *D. undecimpunctata howardi* Barber. Insect Biochem. 18(1): 71–77. (metabolism, detoxification)
11. Andersen, J.F., and P.J. Wilkin. 1986. Methods for the study of pheromones and kairomones. Pages 57–82 in J. L. Krysan and T. A. Miller, eds. Methods for the study of pest *Diabrotica*. Springer-Verlag, New York. (*D. barberi*, *D. virgifera virgifera*, *D. virgifera zea*, bioassay, electroantennogram, cucurbitacins, *Cucurbita* spp., plant volatiles, indole)
12. Andow, D., and R. Wetzler. 1982. A new method for mass marking insects. Entomologist's Monthly Mag. 118: 211–212. (*D. barberi* as *D. longicornis*, mark and recapture)
13. Anonymous. 1987. An alternate school of thought. Crops Soils Mag. 39(8): 25–26. (*D. barberi*, *D. virgifera virgifera*, extended diapause, environmental factors, temperature, crop rotation)
14. Apple, J.W., H.C. Chiang, L.M. English, L.K. French, A.J. Keaster, G.F. Krause, Z.B. Mayo, J.D. Munson, G.J. Musick, J.C. Owens, E.E. Rasmussen, R.E. Sechriest, J.J. Tollefson, and J.L. Wedberg. 1977. Impact of northern and western corn rootworm larvae on field corn. N. Cent. Reg. Res. Publ. 239. Research Division, University of Wisconsin, Madison. 10 pp. (*D. virgifera virgifera*, *D. barberi* as *D. longicornis*, yield losses, damage [larval], host plant resistance, economic thresholds, sampling [egg], economics, root damage ratings)
- ♦ 15. Arnold, F.J. 1981. Residual activity and efficacy of permethrin and fenvalerate for control of corn soil pests. M.S. Thesis, University of Missouri, Columbia. (*Diabrotica* spp., pyrethroids)
16. Bach, C.E. 1977. Distribution of *Acalymma vittata* and *Diabrotica virgifera* (Coleoptera: Chrysomelidae) on cucurbits. Great Lakes Entomol. 10(3): 123–125. (*D. virgifera virgifera*, *Cucurbita* spp., competition)
17. Bach, C.E. 1988. Effects of host plant patch size on herbivore density: patterns. Ecology 69(4): 1090–1102. (*D. virgifera virgifera*, population dynamics, Cucurbitaceae)
18. Bach, C.E. 1988. Effects of host plant patch size on herbivore density: underlying mechanisms. Ecology 69(4): 1103–1117. (*D. virgifera virgifera*, Cucurbitaceae, movement [adult], population dynamics)
19. Ball, H.J. 1981. Larval and adult control recommendations and insecticide resistance data for corn rootworms in Nebraska (1948–1981). Report 3 (rev.). Agric. Exp. Sta., University of Nebraska, Lincoln. 16 pp. (*D. virgifera virgifera*, *D. barberi* as *D. longicornis*, insecticide recommendations, insecticide history, pest management)
20. Ball, H.J. 1982. Spectral response of the adult western corn rootworm (Coleoptera: Chrysomelidae) to selected wavelengths. J. Econ. Entomol. 75(5): 932–933. (*D. virgifera virgifera*, trap color, traps [adult], vision)
21. Ball, H.J. 1983. Aldrin and dieldrin residues in Nebraska soils as related to decreasing LD₅₀ values for the adult western corn rootworm—1962–1981. J. Environ. Sci. Health B 18(6): 735–744. (*D. virgifera virgifera*, insecticide residues)

22. Ball, H.J., and P.P. Su. 1979. Effect of sublethal dosages of carbofuran and carbaryl on fecundity and longevity of the female western corn rootworm. *J. Econ. Entomol.* 72(6): 873–876. (*D. virgifera virgifera*, oviposition, insecticides [sublethal dosage])
23. Ballard, J.B., and Z.B. Mayo. 1979. Predatory potential of selected ant species on eggs of western corn rootworm. *Environ. Entomol.* 8(3): 575–576. (*D. virgifera virgifera*, predation, biological control)
24. Bartelt, R.J., and H.C. Chiang. 1977. Field studies involving the sex-attractant pheromones of the western and northern corn rootworm beetles. *Environ. Entomol.* 6(6): 853–861. (*D. virgifera virgifera*, *D. barberi* as *D. longicornis*, trap capture, environmental factors, temperature, wind, trap height, mating status, periodicity, pheromone traps)
- ♦25. Bereza, K. 1983. Corn rootworms. Fact sheet. Ontario Ministry of Agriculture and Food. Order no. 83-052 (rev. of 1981 ed.). 3 pp. (*Diabrotica* spp., biology)
- ♦26. Bergman, M.K. 1980. Optimizing sampling for corn rootworm larvae. M.S. Thesis, Iowa State University, Ames. (*D. virgifera virgifera*, *D. barberi*, sampling [larval])
27. Bergman, M.K. 1983. Field studies on the population dynamics and management of corn rootworms. Ph.D. Thesis, Purdue University, West Lafayette, IN. (*D. barberi*, *D. virgifera virgifera*, economic thresholds, sampling [larval], cultivation time applications, planting date, heat units, phenology, adult emergence)
28. Bergman, M. 1987. Corn rootworm control: do we have any new solutions? Pages 41–49 in Illinois Agricultural Pesticides Conference '87, Thirty-ninth Spray School, summaries of presentations. Coop. Ext. Serv., University of Illinois at Urbana-Champaign. (*Diabrotica* spp., pest management, cultivation time applications, adult control, dispersal, movement [adult])
29. Bergman, M.K., J.J. Tollefson, and P.N. Hinz. 1981. Sampling scheme for estimating populations of corn rootworm larvae. *Environ. Entomol.* 10(6): 986–990. (*D. virgifera virgifera*, *D. barberi*, sampling [larval], flotation [larval], soil washing apparatus)
30. Bergman, M.K., J.J. Tollefson, and P.N. Hinz. 1983. Spatial dispersion of corn rootworm larvae (Coleoptera: Chrysomelidae) in Iowa cornfields. *Environ. Entomol.* 12(5): 1443–1446. (*D. virgifera virgifera*, *D. barberi*, sampling [larval], dispersion [larval])
31. Bergman, M.K., and F.T. Turpin. 1982. Cultivator versus planting treatment for rootworms. Pages 39–42 in Thirty-fourth Illinois Custom Spray Operators Training School, summaries of presentations. Coop. Ext. Serv., University of Illinois at Urbana-Champaign. (*Diabrotica* spp., cultivation time applications)
32. Bergman, M.K., and F.T. Turpin. 1984. Impact of corn planting date on the population dynamics of corn rootworms (Coleoptera: Chrysomelidae). *Environ. Entomol.* 13(3): 898–901. (*D. virgifera virgifera*, *D. barberi*)
33. Bergman, M.K., and F.T. Turpin. 1986. Phenology of field populations of corn rootworms (Coleoptera: Chrysomelidae) relative to calendar date and heat units. *Environ. Entomol.* 15(1): 109–112. (*D. virgifera virgifera*, *D. barberi*)
34. Bollinger, E.K., and J.W. Caslick. 1985. Factors influencing blackbird damage to field corn. *J. Wildlife Management* 49(4): 1109–1115. (*D. barberi* as *D. longicornis*, *Agelaius phoeniceus*, damage by birds)
35. Bollinger, E.K., and J.W. Caslick. 1985. Northern corn rootworm beetle densities near a red-winged blackbird roost. *Can. J. Zool.* 63(3): 502–505. (*D. barberi* as *D. longicornis*, *Agelaius phoeniceus*, predation)
36. Bollinger, E.K., and J.W. Caslick. 1985. Red-winged blackbird predation on northern corn rootworm beetles in field corn. *J. Applied Ecol.* 22(1): 39–48. (*Agelaius phoeniceus*, *D. barberi* as *D. longicornis*)
37. Branson, T.F. 1978. Optimum temperature for long-term storage of eggs of *Diabrotica virgifera* (Coleoptera: Chrysomelidae). *Ent. Exp. Appl.* 24(2): 199–200. (*D. virgifera virgifera*, rearing)
38. Branson, T.F. 1982. Olfactory response of larvae of *Diabrotica virgifera virgifera* to plant roots. *Ent. Exp. Appl.* 31(3): 303–307. (movement [larval], host location, olfaction)
39. Branson, T.F. 1986. Larval feeding behavior and host-plant resistance in maize. Pages 159–182 in J. L. Krysan and T. A. Miller, eds. *Methods for the study of pest Diabrotica*. Springer-Verlag, New York. (*D. barberi*, *D. virgifera virgifera*, movement [larval], host plant selection, artificial infestation, root damage ratings, lodging, tolerance, antibiosis, antixenosis, host location)
40. Branson, T.F. 1987. The contribution of pre-hatch and post-hatch development to protandry in the chrysomelid, *Diabrotica virgifera virgifera*. *Ent. Exp. Appl.* 43(3): 205–208. (egg hatch duration, sex ratio, adult emergence, rearing)
41. Branson, T.F., and P.L. Guss. 1983. Olfactory responses of *Diabrotica* spp. (Coleoptera: Chrysomelidae) to cut fruits of bitter and nonbitter *Cucurbita* spp. *Environ. Entomol.* 12(3): 700–702. (*D. virgifera virgifera*, cucurbitacins, olfaction, arrestants [feeding])
42. Branson, T.F., and P.L. Guss. 1984. Sterilization of adult male western corn rootworms by gamma radiation in an inert atmosphere. *J. Kansas Entomol. Soc.* 57(4): 717–719. (*D. virgifera virgifera*, sterile insect release)
43. Branson, T.F., P.L. Guss, and J.J. Jackson. 1977. Mating frequency of the western corn rootworm. *Ann. Entomol. Soc. Am.* 70(4): 506–508. (*D. virgifera virgifera*, reproductive status, oviposition, mating status)
44. Branson, T.F., P.L. Guss, and J.J. Jackson. 1978. Sterilizing male western corn rootworms with gamma radiation (Coleoptera: Chrysomelidae). *J. Kansas Entomol. Soc.* 51(2): 270–273. (*D. virgifera virgifera*, sterile insect release)
45. Branson, T.F., P.L. Guss, and J.L. Krysan. 1978. Winter populations of some *Diabrotica* in Central Mexico: voltinism and pheromone response. *Ann. Entomol. Soc. Am.* 71(2): 165–166. (*D. virgifera virgifera*, *D. virgifera zcae*, diapause)
46. Branson, T.F., and J.J. Jackson. 1988. An improved diet for adult *Diabrotica virgifera virgifera* (Coleoptera: Chrysomelidae). *J. Kansas Entomol. Soc.* 61(3): 353–355. (rearing, diet [adult])

47. Branson, T.F., J.J. Jackson, and G.R. Sutter. 1988. Improved method for rearing *Diabrotica virgifera virgifera* (Coleoptera: Chrysomelidae). J. Econ. Entomol. 81(1): 410-414.
48. Branson, T.F., and J.L. Krysan. 1981. Feeding and oviposition behavior and life cycle strategies of *Diabrotica*: an evolutionary view with implications for pest management. Environ. Entomol. 10(6): 826-831. (*D. barberi*, *D. virgifera virgifera*, *D. virgifera zeae*, voltinism, feeding behavior, ovipositional preference, host plant resistance, antibiosis, antixenosis, tolerance)
49. Branson, T.F., and J. Reyes R. 1983. The association of *Diabrotica* spp. with *Zea diploperennis*. J. Kansas Entomol. Soc. 56(1): 97-99. (*D. barberi*, *D. virgifera virgifera*, *D. virgifera zeae*, host plant resistance)
50. Branson, T.F., J. Reyes R., and H. Valdés M. 1982. Field biology of Mexican corn rootworm, *Diabrotica virgifera zeae* (Coleoptera: Chrysomelidae), in central Mexico. Environ. Entomol. 11(5): 1078-1083. (*D. virgifera virgifera*, diapause, soil moisture, population dynamics, predation, pathogens [insect], irrigation, rainfall, environmental factors, egg hatch, developmental time, oviposition, weeds)
51. Branson, T.F., and G.R. Sutter. 1985. Influence of population density of immatures on size, longevity, and fecundity of adult *Diabrotica virgifera virgifera* (Coleoptera: Chrysomelidae). Environ. Entomol. 14(6): 687-690. (root damage ratings, oviposition, size [adult])
52. Branson, T.F., and G. R. Sutter. 1986. Influence of application date on damage resulting from controlled infestations with eggs of *Diabrotica virgifera virgifera* (Coleoptera: Chrysomelidae). J. Econ. Entomol. 79(3): 838-839. (artificial infestation, egg hatch duration, root damage ratings)
53. Branson, T.F., G.R. Sutter, and J.R. Fisher. 1980. Plant response to stress induced by artificial infestations of western corn rootworm. Environ. Entomol. 9(2): 253-257. (*D. virgifera virgifera*, root damage ratings, yield losses, plant stress)
54. Branson, T.F., G.R. Sutter, and J.R. Fisher. 1981. The independence of rootworm damage ratings and the size of maize root systems. J. Kansas Entomol. Soc. 54(4): 694-696. (*D. virgifera virgifera*, root size, root damage ratings)
55. Branson, T.F., G.R. Sutter, and J.R. Fisher. 1981. Non-diapause eggs of western corn rootworms in artificial field infestations. Environ. Entomol. 10(1): 94-96. (*D. virgifera virgifera*, artificial infestation, non-diapause strain)
56. Branson, T.F., G.R. Sutter, and J.R. Fisher. 1982. Comparison of a tolerant and a susceptible maize inbred under artificial infestations of *Diabrotica virgifera virgifera*: yield and adult emergence. Environ. Entomol. 11(2): 371-372. (root damage ratings, tolerance, host plant resistance, yield losses)
57. Branson, T.F., H. Valdés M., and A.L. Kahler. 1986. Evaluation of Mexican populations of maize for resistance to larvae of *Diabrotica virgifera virgifera* (Coleoptera: Chrysomelidae). Maydica 31(2): 173-177. (*D. virgifera virgifera*, *D. virgifera zeae*, host plant resistance, root damage ratings)
58. Branson, T.F., V.A. Welch, G.R. Sutter, and J.R. Fisher. 1983. Resistance to larvae of *Diabrotica virgifera virgifera* in three experimental maize hybrids. Environ. Entomol. 12(5): 1509-1512. (host plant resistance, root damage ratings, yield losses, antixenosis)
59. Briggs, S.P. 1987. Using traps for insect management. Pages 15-19 in 1987 Thirteenth Annual Illinois Crop Protection Workshop proceedings. Coop. Ext. Serv., University of Illinois at Urbana-Champaign. (*Diabrotica* spp., traps [adult], emergence traps, cucurbitacin vial traps, pest management)
60. Briggs, S.P., and J.T. Shaw. 1983. Monitoring for black cutworms and corn rootworms. Pages 135-138 in Thirty-fifth Illinois Custom Spray Operators Training School, summaries of presentations. Coop. Ext. Serv., University of Illinois at Urbana-Champaign. (*D. barberi*, *D. virgifera virgifera*, cucurbitacin vial traps, root damage ratings, visual counts, population measurement, sampling [adult])
61. Bruss, R.G. 1981. Intrafield dispersal of adult corn rootworms. Ph.D. Thesis, Iowa State University, Ames. (*D. barberi*, *D. virgifera virgifera*, movement [adult], temperature, population density, pheromone)
62. Burkhardt, C.C., and G.J. Michels, Jr. 1981. Western corn rootworm: research on distribution, damage and control in Wyoming. Agric. Exp. Sta. Bull. RJ169. University of Wyoming, Laramie. 18 pp. (*D. virgifera virgifera*, biology, distribution [geographical], crop rotation, root damage ratings, economics)
63. Bush, M.R. 1988. Impact of selected cultural practices on soil invertebrates in continuous corn and a new serological technique for identifying predators of corn rootworms. M.S. Thesis, Purdue University, West Lafayette, IN. (*Diabrotica* spp., predation, biological control)
64. Buttery, R.G., and L.C. Ling. 1985. Volatile components of corn roots: possible insect attractants. J. Agric. Food Chem. 33: 772-774. (*D. barberi*, plant volatiles, olfaction, host plant selection)
65. Call, D.J., D.D. Walgenbach, and G.R. Sutter. 1977. Corn rootworm cholinesterase activity as influenced by species, sex, age, and cold storage. J. Econ. Entomol. 70(5): 533-535. (*D. virgifera virgifera*, sex differences)
66. Calvin, D. 1986. Western and northern corn rootworm management in Pennsylvania. Special Circ. 333. Coop. Ext. Serv., Pennsylvania State University, State College. 8 pp. (*D. barberi*, *D. virgifera virgifera*, pest management, biology, sampling [adult], economic thresholds, crop rotation)
67. Calvin, D.D., and F.L. Poston. 1984. Managing arthropods in corn with an emphasis on biological control. Pages 87-97 in Proc. Intl. Symp.: The Role of Biological Control in Pest Management. IOBC/VHRS, November 16-20, 1981. Santiago, Chile. University Press, Ottawa, Canada. (*D. barberi* as *D. longicornis*, *D. virgifera virgifera*, crop rotation)
68. Cameron, K. 1982. Effect of ammonia on post-diapause western corn rootworm eggs and larvae. M.S. Thesis, South Dakota State University, Brookings. (*D. virgifera virgifera*, fertilizer)
69. Capinera, J.L., N.D. Epsky, and D.C. Thompson. 1986. Effect of adult western corn rootworm (Coleoptera: Chrysomelidae) ear feeding on irrigated field corn in Colorado. J. Econ. Entomol. 79(6): 1609-1612. (*D. virgifera virgifera*, silk feeding, yield losses, irrigation)

70. Capinera, J.L., T.J. Weissling, and E.E. Schweizer. 1985. Compatibility of intercropping with mechanized agriculture: effects of strip intercropping of pinto beans and sweet corn on insect abundance in Colorado. *J. Econ. Entomol.* 78(2): 354–357. (*D. virgifera virgifera*)
71. Chapman, R.A., and C.R. Harris. 1980. Insecticidal activity and persistence of terbufos, terbufos sulfoxide and terbufos sulfone in soil. *J. Econ. Entomol.* 73(4): 536–543. (*D. barberi* as *D. longicornis*, soil type, soil moisture, temperature, environmental factors, bioassay, toxicity, insecticide persistence, insecticide degradation, phorate, carbofuran, insecticide volatility, insecticide residues)
72. Chiang, H.C. 1978. Pest management in corn. *Ann. Rev. Entomol.* 23: 101–123. (*D. barberi* as *D. longicornis*, *D. virgifera virgifera*)
73. Chiang, H.C. 1982. Factors to be considered in refining a general model of economic threshold. *Entomophaga* 27 (Special Issue): 99–103. (*D. virgifera virgifera*, plant stress, host plant resistance, phenology)
74. Chiang, H.C., and L.K. French. 1980. Host tolerance, a short-term pest management tool—maize and corn rootworm as a model. *Plant Prot. Bull., F.A.O.* 28(4): 137–138. (*D. virgifera virgifera*, host plant resistance, survival, root damage ratings)
75. Chiang, H.C., L.K. French, and D.E. Rasmussen. 1980. Quantitative relationship between western corn rootworm population and corn yield. *J. Econ. Entomol.* 73(5): 665–666. (*D. virgifera virgifera*, artificial infestation, yield losses, economic thresholds, rainfall, population density)
76. Chio, H. 1977. Comparative mechanisms for resistance to aldrin, carbofuran, fonofos, terbufos, and phorate in four species of *Diabrotica* beetles. Ph.D. Thesis, University of Illinois at Urbana-Champaign. (*D. barberi* as *D. longicornis*, *D. virgifera virgifera*, insecticide resistance)
77. Chio, H., C.S. Chang, R.L. Metcalf, and J. Shaw. 1978. Susceptibility of four species of *Diabrotica* to insecticides. *J. Econ. Entomol.* 71(3): 389–393. (*D. barberi* as *D. longicornis*, *D. virgifera virgifera*, insecticide resistance, bioassay, aldrin, heptachlor, DDT, fensulfothion, fonofos, malathion, diazinon, methoxychlor, carbaryl, carbofuran, chlorpyrifos, phorate, terbufos, permethrin)
78. Chio, H., and R.L. Metcalf. 1979. Detoxification mechanisms for aldrin, carbofuran, fonofos, phorate, and terbufos in four species of *Diabrotica*. *J. Econ. Entomol.* 72(5): 732–738. (*D. barberi* as *D. longicornis*, *D. virgifera virgifera*, insecticide resistance, insecticide degradation)
79. Clement, S.L., D.A. McCartney, and M.A. Casey. 1987. Adult corn rootworm (*Diabrotica*, Coleoptera: Chrysomelidae) populations in field corn in northeastern Ohio. *J. Kansas Entomol. Soc.* 60(4): 535–540. (*D. barberi* as *D. longicornis*, *D. virgifera virgifera*, oviposition, movement [adult], population density)
80. Clement, S.L., G.Szatmari-Goodman, and B.M. Drees. 1979. The status and control of Ohio corn rootworms. *Ohio Report* 64(1): 8–10. (*D. barberi* as *D. longicornis*, *D. virgifera virgifera*, distribution [geographical], insecticide control, movement [adult])
81. Coats, J.R. 1986. Toxicological methods and laboratory testing of insecticides. Pages 205–226 in J. L. Krysan and T. A. Miller, eds. *Methods for the study of pest Diabrotica*. Springer-Verlag, New York. (*D. virgifera virgifera*, bioassay, insecticide resistance, insecticide degradation, insecticide bioactivity, biodegradation [enhanced])
82. Coats, S.A.O. 1986. Migratory flight potential and the role of juvenile hormone in flight regulation in the western corn rootworm (*Diabrotica virgifera virgifera*). Ph.D. Thesis, Iowa State University, Ames. (movement [adult], periodicity, dispersal, reproductive status, mating status, ovarian development)
83. Coats, S.A., J.R. Coats, and C.R. Ellis. 1979. Selective toxicity of three synthetic pyrethroids to eight coccinellids, a eulophid parasitoid, and two pest chrysomelids. *Environ. Entomol.* 8(4): 720–722. (*D. barberi* as *D. longicornis*, bioassay, permethrin, cypermethrin, fenvalerate)
84. Coats, S.A., J.A. Mutchmor, and J.J. Tollefson. 1987. Regulation of migratory flight by juvenile hormone mimic and inhibitor in western corn rootworm (Coleoptera: Chrysomelidae). *Ann. Entomol. Soc. Am.* 80(5): 697–708. (*D. virgifera virgifera*, movement [adult], ovarian development, reproductive status, mating status)
85. Coats, S.A., J.J. Tollefson, and J.A. Mutchmor. 1986. Study of migratory flight in the western corn rootworm (Coleoptera: Chrysomelidae). *Environ. Entomol.* 15(3): 620–625. (*D. virgifera virgifera*, movement [adult], periodicity)
86. Culy, M.D. 1987. The impact of western corn rootworm (Coleoptera: Chrysomelidae) on pollination success in seed corn production fields. Ph.D. Thesis, Purdue University, West Lafayette, IN. (*D. virgifera virgifera*, economic thresholds, silk feeding, environmental factors)
87. Danielson, S.D., F.P. Baxendale, R.J. Wright, J.F. Witkowski, J.B. Campbell, L.L. Peters, G.L. Hein, A.F. Hagen, K.J. Jarvi, R.C. Seymour, and J.A. Kalisch. 1989. 1989 insect management guide. Publ. EC 89–1509. Coop. Ext. Serv., University of Nebraska, Lincoln. 18 pp. (*D. barberi*, *D. virgifera virgifera*, extended diapause, crop rotation, pest management, insecticide recommendations, adult control)
- ♦88. Day, E. 1985. Studies on the susceptibility of corn rootworms to infection by *Beauveria bassiana*. M.S. Thesis, University of Illinois at Urbana-Champaign. (*Diabrotica* spp., fungi, biological control, pathogens [insect])
- ♦89. Deng, Y. 1987. Electroantennogram responses of four *Diabrotica* species and *Acalymna vittatum* (Coleoptera: Chrysomelidae) to a number of benzenoid semiochemicals. M.S. Thesis, University of Illinois at Urbana-Champaign.
90. Dicke, F.F., and W.D. Guthrie. 1978. Genetics of insect resistance in maize. Pages 299–317 in D. B. Walden, ed. *Maize breeding and genetics*. John Wiley and Sons, New York. (*D. barberi*, *D. virgifera virgifera*, host plant resistance, tolerance, antibiosis, root regrowth)
- ♦91. Dobson, I.D. 1985. Pheromone-mediated behaviors of *Diabrotica barberi* Smith and Lawrence and *D. virgifera* LeConte (Coleoptera: Chrysomelidae). M.S. Thesis, University of Guelph, Guelph, Ontario, Canada. (*D. virgifera virgifera*)
92. Dobson, I.D., and P.E.A. Teal. 1986. Field studies of the temporal response patterns of male *Diabrotica virgifera virgifera* LeConte and *D. barberi* Smith and Lawrence to 8R-methyl-2R-decyl propanoate. *Physiol. Entomol.* 11(4): 405–410. (pheromone, temperature, environmental factors, periodicity)

93. **Dobson, I.D., and P.E.A. Teal.** 1987. Analysis of long-range reproductive behavior of male *Diabrotica virgifera virgifera* LeConte and *D. barberi* Smith and Lawrence to stereoisomers of 8-methyl-2-decyl propanoate under laboratory conditions. *J. Chem. Ecol.* 13(6): 1331-1341. (pheromone)
94. **Doersch, R.E., J.D. Doll, R.A. Flashinski, C.R. Grau, R.G. Harvey, J.L. Wedberg, and G.L. Worf.** 1989. Corn pest management in Wisconsin. Publ. A1684, University of Wisconsin Extension, Madison. 64 pp. (*D. barberi*, *D. virgifera virgifera*, insecticide recommendations, crop rotation, extended diapause)
95. **Dominique, C.R.M.** 1983. Bionomics of the northern corn rootworm *Diabrotica longicornis* (Say) (Coleoptera: Chrysomelidae) in Quebec. Ph.D. Thesis, McGill University, Quebec, Canada. (*D. barberi* as *D. longicornis*, threshold temperature, heat units, adult emergence, oviposition, ovipositional preference, temperature, environmental factors, mortality, rearing, soil moisture, soil type, biology, phenology)
96. **Dominique, C.R., and W.N. Yule.** 1983. Laboratory rearing technique for the northern corn rootworm, *Diabrotica longicornis* (Coleoptera: Chrysomelidae). *Can. Entomol.* 115(5): 569-571. (*D. barberi* as *D. longicornis*, oviposition)
97. **Dominique, C.R., and W.N. Yule.** 1983. Threshold temperature and thermal constant for egg development of northern corn rootworm, *Diabrotica longicornis* (Coleoptera: Chrysomelidae). *Can. Entomol.* 115(8): 1051-1052. (*D. barberi* as *D. longicornis*, heat units)
98. **Dominique, C.R., and W.N. Yule.** 1984. Bionomics of the northern corn rootworm, *Diabrotica longicornis* (Say) (Coleoptera: Chrysomelidae), in Quebec. *Revue d'Entomologie du Québec* 29(1): 12-16. (*D. barberi* as *D. longicornis*, biology, phenology)
99. **Dominique, C.R., W.N. Yule, and P. Martel.** 1983. Influence of soil type, soil moisture, and soil surface conditions on oviposition preference of the northern corn rootworm, *Diabrotica longicornis* (Coleoptera: Chrysomelidae). *Can. Entomol.* 115(8): 1043-1046. (*D. barberi* as *D. longicornis*)
100. **Edwards, C.R., and F.T. Turpin.** 1988. Managing corn rootworms—1989. Publ. E-49, Coop. Ext. Serv., Purdue University, West Lafayette, IN. 5 pp. (*D. barberi*, *D. virgifera virgifera*, pest management, sampling [adult], sampling [larval], extended diapause, insecticide recommendations)
101. **Ellis, C.R.** 1982. A survey of granular application equipment and insecticide rates used for control of corn rootworms (Coleoptera: Chrysomelidae) in southern Ontario. *Proc. Entomol. Soc. Ont.* 113: 29-34. (*D. barberi*, *D. virgifera virgifera*, insecticide application equipment, insecticide control, insecticide calibration)
102. **Ellis, C.R., and B. Beattie.** 1984. Effect of banding and incorporation on the efficacy of granular insecticides for control of corn rootworms (Coleoptera: Chrysomelidae) in grain corn. *Proc. Entomol. Soc. Ont.* 115: 31-36. (*D. barberi*, *D. virgifera virgifera*, insecticide application equipment, insecticide control)
103. **Felsot, A.S.** 1981. Factors affecting the performance of Furadan against corn rootworms. Pages 79-80 in *Thirty-third Illinois Custom Spray Operators Training School*, summaries of presentations. Coop. Ext. Serv., University of Illinois at Urbana-Champaign. (*D. virgifera virgifera*, carbofuran, root damage ratings, insecticide resistance, insecticide persistence)
104. **Felsot, A.S.** 1982. Insecticide degradation: above and below ground. Pages 97-107 in *Eighth Annual Illinois Crop Protection Workshop*. Coop. Ext. Serv., University of Illinois at Urbana-Champaign. (*Diabrotica* spp., insecticide persistence)
105. **Felsot, A.** 1984. Persistence of isofenphos (Amaze) soil insecticide under laboratory and field conditions and tentative identification of a stable oxygen analog metabolite by gas chromatography. *J. Environ. Sci. Health B* 19(1): 13-27. (*D. barberi*, *D. virgifera virgifera*, insecticide persistence, insecticide degradation)
106. **Felsot, A.** 1985. Factors affecting the bioactivity of soil insecticides. Pages 134-138 in *Thirty-seventh Illinois Custom Spray Operators Training School manual*. Coop. Ext. Serv., University of Illinois at Urbana-Champaign. (*D. barberi*, *D. virgifera virgifera*, insecticide bioactivity, insecticide persistence, toxicity, environmental factors, insecticide resistance)
107. **Felsot, A.** 1987. Fate and interactions of pesticides in conservation tillage systems. Pages 35-43 in G. J. House and B. R. Stinner, eds. *Arthropods in conservation tillage systems*. Misc. Publ. 65, Entomol. Soc. Am., College Park, MD. (*D. virgifera virgifera*, *D. barberi*, no-till, chlorpyrifos, phorate, terbufos, carbaryl, carbofuran, DDT, aldrin, environmental chemodynamics, insecticide degradation, biodegradation [enhanced], insecticide persistence, pesticide interactions)
108. **Felsot, A.S., W.N. Bruce, and K.S. Steffey.** 1987. Degradation of terbufos (Counter) soil insecticide in corn fields under conservation tillage practices. *Bull. Environ. Contam. Toxicol.* 38: 369-376. (*Diabrotica* spp., no-till, insecticide degradation)
109. **Felsot, A., E. Levine, K.L. Steffey, and M.A. Foster.** 1984. Soil insecticide behavior and efficacy under conservation tillage systems. Pages 146-154 in *Thirty-sixth Illinois Custom Spray Operators Training School manual*. Coop. Ext. Serv., University of Illinois at Urbana-Champaign. (*Diabrotica* spp., no-till, adult emergence, insecticide persistence, root damage ratings)
110. **Felsot, A.S., J.V. Maddox, and W. Bruce.** 1981. Enhanced microbial degradation of carbofuran in soils with histories of Furadan use. *Bull. Environ. Contam. Toxicol.* 26: 781-788. (*D. barberi*, *D. virgifera virgifera*, insecticide persistence, insecticide degradation, biodegradation [enhanced], insecticide history)
111. **Felsot, A.S., K.L. Steffey, and E. Levine.** 1986. Conservation tillage and pesticides. *Ill. Nat. Hist. Sur. Reports* 253: [3-4]. (*D. barberi*, *D. virgifera virgifera*, no-till, yield losses)
112. **Felsot, A.S., K.L. Steffey, E. Levine, and J.G. Wilson.** 1985. Carbofuran persistence in soil and adult corn rootworm (Coleoptera: Chrysomelidae) susceptibility: relationship to the control of damage by larvae. *J. Econ. Entomol.* 78(1): 45-52. (*D. barberi*, *D. virgifera virgifera*, insecticide degradation, insecticide persistence, insecticide resistance, root damage ratings)
113. **Felsot, A., L. Wei, and J. Wilson.** 1982. Environmental chemodynamic studies with terbufos (Counter) insecticide in soil under laboratory and field conditions. *J. Environ. Sci. Health B* 17(6): 649-673. (*D. barberi*, *D. virgifera virgifera*, insecticide persistence, environmental factors, soil type, insecticide degradation)

114. Felsot, A.S., J.G. Wilson, D.E. Kuhlman, and K.L. Steffey. 1982. Rapid dissipation of carbofuran as a limiting factor in corn rootworm (Coleoptera: Chrysomelidae) control in fields with histories of continuous carbofuran use. *J. Econ. Entomol.* 75(6): 1098–1103. (*D. barberi* as *D. longicornis*, *D. virgifera virgifera*, insecticide persistence, insecticide degradation, biodegradation [enhanced], insecticide history)
115. Ferguson, J.E. 1985. Disposition of cucurbitacin as a factor in host plant selection by Diabrotic beetle and electrophoretic analysis of the coevolved genus *Cucurbita*. Ph.D. Thesis, University of Illinois at Urbana-Champaign. (*D. virgifera virgifera*, longevity, diet [adult], electrophoresis)
116. Ferguson, J.E., E.R. Metcalf, R.L. Metcalf, and A.M. Rhodes. 1983. Influence of cucurbitacin content in cotyledons of Cucurbitaceae cultivars upon feeding behavior of Diabrotic beetles (Coleoptera: Chrysomelidae). *J. Econ. Entomol.* 76(1): 47–51. (*D. virgifera virgifera*, *Cucurbita* spp.)
117. Ferguson, J.E., and R.L. Metcalf. 1985. Cucurbitacins: plant-derived defense compounds for Diabrotic beetles (Coleoptera: Chrysomelidae). *J. Chem. Ecol.* 11(3): 311–318. (*D. virgifera virgifera*, sequestration)
118. Ferguson, J.E., R.L. Metcalf, and D.C. Fischer. 1985. Disposition and fate of cucurbitacin b on five species of Diabrotic beetles. *J. Chem. Ecol.* 11(9): 1307–1321. (*D. virgifera virgifera*, *Cucurbita* spp., sequestration, metabolism, detoxification)
119. Fielding, D.J. 1985. Influence of varying amounts of bait on numbers of *Diabrotica virgifera* and *D. barberi* caught in cucurbitacin traps. M.S. Thesis, University of Illinois at Urbana-Champaign. (*D. virgifera virgifera*, cucurbitacin vial traps, sex ratio)
120. Fielding, D.J., and W.G. Ruesink. 1985. Varying amounts of bait influences numbers of western and northern corn rootworms (Coleoptera: Chrysomelidae) caught in cucurbitacin traps. *J. Econ. Entomol.* 78(5): 1138–1144. (*D. barberi*, *D. virgifera virgifera*, cucurbitacin vial traps, sex ratio)
- ♦ 121. Fischer, D.C. 1981. Tachinid parasitoids of *Acalymma vittata*, *Diabrotica undecimpunctata* and *Diabrotica virgifera*. M.S. Thesis, University of Illinois at Urbana-Champaign. (*D. virgifera virgifera*, parasites, biological control)
122. Fischer, D.C. 1983. *Celatoria diabroticae* Shimer and *Celatoria setosa* Coquillett: tachinid parasitoids of the diabrotic beetle. Ph.D. Thesis, University of Illinois at Urbana-Champaign. (*D. barberi* as *D. longicornis*, parasites, biological control, rearing, developmental time, kairomones, threshold temperature)
123. Fisher, J.R. 1980. A modified emergence trap for quantitative adult corn rootworm studies (Coleoptera: Chrysomelidae). *J. Kansas Entomol. Soc.* 53(2): 363–366. (*Diabrotica* spp., adult emergence, sampling [adult])
124. Fisher, J.R. 1981. System for extracting corn rootworm larvae from soil samples. *J. Econ. Entomol.* 74(1): 103–105. (*D. barberi*, *D. virgifera virgifera*, soil washing apparatus, flotation [larval])
125. Fisher, J.R. 1984. Comparison of emergence of *Diabrotica virgifera virgifera* (Coleoptera: Chrysomelidae) from cut and uncut corn plants in artificial and natural infestations. *J. Kansas Entomol. Soc.* 57(3): 405–408. (adult emergence, artificial infestation, emergence traps)
126. Fisher, J.R. 1985. Comparison of controlled infestations of *Diabrotica virgifera virgifera* and *Diabrotica barberi* (Coleoptera: Chrysomelidae) on corn. *J. Econ. Entomol.* 78(6): 1406–1408. (root damage ratings, artificial infestation)
127. Fisher, J.R. 1986. Development and survival of pupae of *Diabrotica virgifera virgifera* and *D. undecimpunctata howardi* (Coleoptera: Chrysomelidae) at constant temperatures and humidities. *Environ. Entomol.* 15(3): 626–630. (environmental factors)
128. Fisher, J.R. 1987. Greenhouse method for studying development and survival of *Diabrotica virgifera virgifera* (Coleoptera: Chrysomelidae). *J. Econ. Entomol.* 80(1): 286–289. (environmental factors, soil moisture, heat units)
129. Fisher, J.R., and M.K. Bergman. 1986. Field sampling of larvae and pupae. Pages 101–121 in J. L. Krysan and T. A. Miller, eds. *Methods for the study of pest Diabrotica*. Springer-Verlag, New York. (*D. barberi*, *D. virgifera virgifera*, sampling [larval], sampling [pupal], flotation [larval], sampling statistics)
130. Fisher, J.R., T.F. Branson, and G.R. Sutter. 1984. Use of common squash cultivars, *Cucurbita* spp., for mass collection of corn rootworm beetles, *Diabrotica* spp. (Coleoptera: Chrysomelidae). *J. Kansas Entomol. Soc.* 57(3): 409–412. (*D. barberi*, *D. virgifera virgifera*, sex ratio)
131. Fisher, J.R., J.J. Jackson, G.R. Sutter, and T.F. Branson. 1986. Influence of adult diet on damage to corn by larvae of the western corn rootworm, *Diabrotica virgifera virgifera* (Coleoptera: Chrysomelidae). *J. Econ. Entomol.* 79(1): 114–115. (artificial infestation, diet [adult], rearing)
132. Foott, W.H., and P.R. Timmins. 1977. Observations on new insect pests of grain corn in Essex County, Ontario. *Proc. Entomol. Soc. Ont.*, 1977, publ. 1978, 108: 49–52. (*D. virgifera virgifera*, distribution [geographical])
133. Foster, R.E. 1977. Corn rootworm egg sampling. M.S. Thesis, University of Illinois at Urbana-Champaign. (*D. barberi* as *D. longicornis*, *D. virgifera virgifera*, sampling [egg], distribution [spatial], oviposition)
134. Foster, R.E. 1983. The development and use of sampling and scouting procedures for corn insects and the relationship of corn rootworm densities to corn yield. Ph.D. Thesis, Iowa State University, Ames. (*D. barberi*, *D. virgifera virgifera*, sampling [adult], sampling [sequential], sampling statistics, sampling [egg], damage [larval], yield losses, economic thresholds, model, root damage ratings, planting date, fertilizer, plant density, rainfall)
135. Foster, R.E., W.G. Ruesink, and W.H. Luckmann. 1978. Northern corn rootworm egg sampling (abstract). *Proc. N. Cent. Br. Entomol. Soc. Am.*, 1978, publ. 1979, 33: 60. (*D. barberi* as *D. longicornis*, oviposition, sampling [egg], distribution [spatial])
136. Foster, R.E., W.G. Ruesink, and W.H. Luckmann. 1979. Northern corn rootworm egg sampling. *J. Econ. Entomol.* 72(5): 659–663. (*D. barberi* as *D. longicornis*, oviposition, sampling [egg], distribution [spatial])

137. Foster, R.E., and J.J. Tollefson. 1986. Frequency and severity of attack of several pest insects on corn in Iowa. J. Kansas Entomol. Soc. 59(2): 269-274. (*D. barberi*, *D. virgifera virgifera*, economic thresholds, sampling [adult], root damage ratings)
138. Foster, R.E., J.J. Tollefson, J.P. Nyrop, and G.L. Hein. 1986. Value of adult corn rootworm (Coleoptera: Chrysomelidae) population estimates in pest management decision making. J. Econ. Entomol. 79(2): 303-310. (*D. barberi*, *D. virgifera virgifera*, sampling [adult], yield losses, economic thresholds)
139. Foster, R.E., J.J. Tollefson, and K.L. Steffey. 1982. Sequential sampling plans for adult corn rootworms (Coleoptera: Chrysomelidae). J. Econ. Entomol. 75(5): 791-793. (*D. barberi* as *D. longicornis*, *D. virgifera virgifera*, sampling [adult], sampling [sequential])
140. Gergerich, R.C., H.A. Scott, and J.P. Fulton. 1986. Evaluation of *Diabrotica* beetles as vectors of plant viruses. Pages 227-249 in J. L. Krysan and T. A. Miller, eds. Methods for the study of pest *Diabrotica*. Springer-Verlag, New York. (*D. barberi*, *D. virgifera virgifera*, vector [disease], virus transmission, disease transmission, pathogens [plant])
141. Gilbertson, R.L., III. 1985. Ecology and epidemiology of corn stalk rot in Colorado. Ph.D. Thesis, Colorado State University, Fort Collins. (*D. virgifera virgifera*, *Fusarium* spp., fungi, vector [disease], pathogens [plant])
142. Gilbertson, R.L., W.M. Brown, and E.G. Ruppel. 1984. Association of stalk rot fungi and western corn rootworm beetles in Colorado (abstract). Annual Meeting of the Am. Phytopathol. Soc. (Pacific Div.), June 25-27, 1984. Phytopathology 74(9): 1138. (*D. virgifera virgifera*, *Fusarium* spp., vector [disease], pathogens [plant])
143. Gilbertson, R.L., W.M. Brown, Jr., E.G. Ruppel, and J.L. Capinera. 1986. Association of corn stalk rot *Fusarium* spp. and western corn rootworm beetles in Colorado. Phytopathology 76(12): 1309-1314. (*D. virgifera virgifera*, fungi, vector [disease], pathogens [plant])
- ♦ 144. Godfrey, L.D. 1980. Population dynamics of western corn rootworm beetles in first-year and continuous corn. M.S. Thesis, Purdue University, West Lafayette, IN. (*D. virgifera virgifera*, first-year cornfields, continuous cornfields)
145. Godfrey, L.D., and F.T. Turpin. 1983. Comparison of western corn rootworm (Coleoptera: Chrysomelidae) adult populations and economic thresholds in first-year and continuous corn fields. J. Econ. Entomol. 76(5): 1028-1032. (*D. virgifera virgifera*, population dynamics, sex ratio, first-year cornfields, continuous cornfields, crop rotation, reproductive status, root damage ratings, movement [adult], dispersal, ovarian development, population density)
146. Goodwin, D. 1985. The ecology of two species of corn rootworm (Coleoptera: Chrysomelidae) on volunteer corn in soybean fields in northern Illinois (abstract). Bull. Ecol. Soc. Am. 66(2): 179. (*D. barberi*, *D. virgifera virgifera*, crop rotation, oviposition, weeds, phenology, adult emergence)
147. Goodwin, D.M. 1987. The ecology of northern corn rootworm (*Diabrotica barberi*) and western corn rootworm (*Diabrotica virgifera*) on volunteer corn in soybean fields in northern Illinois. Ph.D. Thesis, University of Illinois, Chicago. (*D. virgifera virgifera*, crop rotation, sampling [adult], first-year cornfields, sampling [egg], oviposition, foxtail, *Setaria* spp.)
148. Gorder, G.W., P.A. Dahm, and J.J. Tollefson. 1982. Carbofuran persistence in cornfield soils. J. Econ. Entomol. 75(4): 637-642. (*D. barberi* as *D. longicornis*, *D. virgifera virgifera*, insecticide persistence, insecticide degradation, soil moisture, biodegradation [enhanced])
149. Gorder, G.W., J.J. Tollefson, and P.A. Dahm. 1980. Carbofuran residue analysis and control of corn rootworm larval damage. Iowa State J. Res. 55(1): 25-33. (*D. barberi* as *D. longicornis*, *D. virgifera virgifera*, root damage ratings, insecticide history, insecticide residues, insecticide degradation, insecticide persistence)
150. Gray, M.E. 1986. The population dynamics of the western and northern corn rootworms in four tillage systems. Ph.D. Thesis, Iowa State University, Ames. (*D. barberi*, *D. virgifera virgifera*, ovipositional preference, survival, no-till, adult emergence, root damage ratings, sampling [egg])
151. Gray, M.E., J.R. Coats, and J.J. Tollefson. 1984. Effect of an insecticide and a herbicide combination on corn rootworm (Coleoptera: Chrysomelidae) damage. J. Econ. Entomol. 77(2): 465-467. (*D. virgifera virgifera*, root damage ratings, carbofuran, alachlor, cyanazine)
152. Gray, M.E., and J.J. Tollefson. 1987. Influence of tillage and western and northern corn rootworm (Coleoptera: Chrysomelidae) egg populations on larval populations and root damage. J. Econ. Entomol. 80(4): 911-915. (*D. barberi*, *D. virgifera virgifera*, no-till, root damage ratings, population dynamics)
153. Gray, M.E., and J.J. Tollefson. 1988. Emergence of the western and northern corn rootworms (Coleoptera: Chrysomelidae) from four tillage systems. J. Econ. Entomol. 81(5): 1398-1403. (*D. barberi*, *D. virgifera virgifera*, adult emergence, no-till)
154. Gray, M.E., and J.J. Tollefson. 1988. Influence of tillage systems on egg populations of western and northern corn rootworms (Coleoptera: Chrysomelidae). J. Kansas Entomol. Soc. 61(2): 186-194. (*D. barberi*, *D. virgifera virgifera*, no-till, ovipositional preference, sampling [egg], population dynamics)
155. Gray, M.E., and J.J. Tollefson. 1988. Survival of the western and northern corn rootworms (Coleoptera: Chrysomelidae) in different tillage systems throughout the growing season of corn. J. Econ. Entomol. 81(1): 178-183. (*D. barberi*, *D. virgifera virgifera*, no-till)
- ♦ 156. Grogan, J.E. 1979. Impact of rootworm larval damage on yield and quality of corn grain. M.S. Thesis, Purdue University, West Lafayette, IN. (*Diabrotica* spp., yield losses)
157. Guse, C.A. 1985. The effects of weather on visual counts and cucurbitacin trap catches of adult western corn rootworms. *Diabrotica virgifera*. M.S. Thesis, University of Illinois at Urbana-Champaign. (*D. virgifera virgifera*, environmental factors, cucurbitacin vial traps, temperature, rainfall)
158. Guss, P.L., P.E. Sonnet, R.L. Carney, T.F. Branson, and J.H. Tumlinson. 1984. Response of *Diabrotica virgifera virgifera*, *Diabrotica virgifera zea*, and *Diabrotica porracea* to stereoisomers of 8-methyl-2-decyl propanoate. J. Chem. Ecol. 10(7): 1123-1131. (pheromone)

159. Guss, P.L., P.E. Sonnet, R.L. Carney, J.H. Tumlinson, and P.J. Wilkin. 1985. Response of northern corn rootworm, *Diabrotica barberi* Smith and Lawrence, to stereoisomers of 8-methyl-2-decyl propanoate. J. Chem. Ecol. 11(1): 21-26. (pheromone)
160. Guss, P.L., J.H. Tumlinson, P.E. Sonnet, and A.T. Proveaux. 1982. Identification of a female-produced sex pheromone of the western corn rootworm. J. Chem. Ecol. 8(2): 545-556. (*D. barberi*, *D. virgifera virgifera*, *D. virgifera zea*)
161. Gustin, R.D. 1979. Effect of two moisture and population levels on oviposition of the western corn rootworm. Environ. Entomol. 8(3): 406-407. (*D. virgifera virgifera*, population density, soil moisture, ovipositional preference)
162. Gustin, R.D. 1981. Soil temperature environment of overwintering western corn rootworm eggs. Environ Entomol. 10(4): 483-487. (*D. virgifera virgifera*, survival)
163. Gustin, R.D. 1983. *Diabrotica longicornis barberi* (Coleoptera: Chrysomelidae): cold hardness of eggs. Environ. Entomol. 12(3): 633-634. (*D. virgifera virgifera*, survival, temperature)
164. Gustin, R.D. 1984. Effect of crop cover on oviposition of the northern corn rootworm, *Diabrotica longicornis barberi* Smith and Lawrence. J. Kansas Entomol. Soc. 57(3): 515-516. (weeds, ovipositional preference)
165. Gustin, R.D. 1986. Effect of intermittent low temperatures on hatch of western corn rootworm eggs (Coleoptera: Chrysomelidae). J. Kansas Entomol. Soc. 59(3): 569-570. (*D. virgifera virgifera*, survival, cold hardness, egg hatch)
166. Gustin, R.D., and G.E. Wilde. 1984. Cold hardness of western corn rootworm (Coleoptera: Chrysomelidae) eggs from northern and southern zones of the corn belt. J. Kansas Entomol. Soc. 57(4): 722-725. (*D. virgifera virgifera*, survival, temperature)
167. Hansen S.K., I.C. McDonald, D.A. Gabrielson, O.A. Johnson. 1986. Characterization studies of sex-limited esterases among beetles of the genus *Diabrotica* (Coleoptera: Chrysomelidae). Proc. N. D. Acad. Sci. 40: 65. (*D. virgifera virgifera*, electrophoresis)
168. Harris, C.R. 1977. Biological activity of chlorpyrifos, chlorpyrifos-methyl, phorate, and Counter in soil. Can. Entomol. 109(8): 1115-1120. (*D. barberi* as *D. longicornis*, terbufos, soil moisture, soil type, temperature, insecticide persistence, bioassay)
169. Harris, C.R., and R.A. Chapman. 1980. Insecticidal activity and persistence of phorate, phorate sulfoxide, and phorate sulfone in soils. Can. Entomol. 112(7): 641-653. (*D. barberi* as *D. longicornis*, insecticide persistence, insecticide volatility, soil type, soil moisture, temperature)
170. Harris, C.R., R.A. Chapman, C. Harris, and C.M. Tu. 1984. Biodegradation of pesticides in soil: rapid induction of carbamate degrading factors after carbofuran treatment. J. Environ. Sci. Health B 19(1): 1-11. (*Diabrotica* spp., carbaryl, bufencarb, insecticide persistence, insecticide degradation, biodegradation [enhanced], insecticide history, phorate, isofenphos)
171. Harris, C.R., R.A. Chapman, J.H. Tolman, P. Moy, K. Henning, and C. Harris. 1988. A comparison of the persistence in a clay loam of single and repeated annual applications of seven granular insecticides used for corn rootworm control. J. Environ. Sci. Health B 23(1): 1-32. (*D. virgifera virgifera*, insecticide persistence, insecticide degradation, biodegradation [enhanced], soil moisture, insecticide history, carbofuran, chlorpyrifos, disulfoton, fonofos, isofenphos, phorate, terbufos)
172. Harris, C.R., and S.A. Turnbull. 1977. Laboratory assessment of the potential of 16 experimental compounds as soil insecticides. Can. Entomol. 109(8): 1109-1114. (*D. barberi* as *D. longicornis*, bioassay, chlorpyrifos, carbofuran, dieldrin, soil type, soil moisture, temperature, insecticide volatility, insecticide persistence, terbufos)
173. Harris, C.R., and S.A. Turnbull. 1978. Laboratory studies on the contact toxicity and activity in soil of four pyrethroid insecticides. Can. Entomol. 110(3): 285-288. (*D. barberi* as *D. longicornis*, chlorpyrifos, carbofuran, bioassay, soil type, temperature, soil moisture)
- ♦ 174. Harvey, J.P. 1978. Biosystematics and hybridization in five species of diabroticines. M.S. Thesis, University of Illinois at Urbana-Champaign. (*Diabrotica* spp., taxonomy)
- ♦ 175. Hass, M.J. 1987. Soil degradation comparisons of corn rootworm insecticides in the field and laboratory. M.S. Thesis, South Dakota State University, Brookings. (*Diabrotica* spp., insecticide degradation)
- ♦ 176. Hedrick, T.W. 1978. First year corn damage—northern corn rootworm ovipositional studies. M.S. Thesis, South Dakota State University, Brookings. (*D. barberi*, first-year cornfields)
177. Hedrick, T.W., and D.D. Walgenbach. 1978. First year corn damage; northern corn rootworm ovipositional studies (abstract). Proc. N. Cent. Br. Entomol. Soc. Am., 1978, publ. 1979, 33: 62. (*D. barberi*, first-year cornfields, continuous cornfields)
- ♦ 178. Hein, G.L. 1981. Comparison of adult corn rootworm sampling techniques as population estimators and as predictors of subsequent larval damage. M.S. Thesis, Iowa State University, Ames. (*Diabrotica* spp., sampling [adult], population measurement, prediction of damage)
179. Hein, G.L. 1984. Adult population dynamics and oviposition of northern and western corn rootworms in continuous cornfields under commercial production. Ph.D. Thesis, Iowa State University, Ames. (*D. barberi*, *D. virgifera virgifera*, emergence traps, sampling statistics, sex ratio, flight, adult emergence, survival, sampling [egg], model, fecundity)
180. Hein, G.L., M.K. Bergman, R.G. Bruss, and J.J. Tollefson. 1985. Absolute sampling technique for corn rootworm (Coleoptera: Chrysomelidae) adult emergence that adjusts to fit common-row spacing. J. Econ. Entomol. 78(6): 1503-1506. (*D. barberi*, *D. virgifera virgifera*, emergence traps, sampling [adult], population measurement)
181. Hein, G.L., and D.E. Foster. 1986. Corn rootworm management. Pm-670 (rev.), Coop. Ext. Serv., Iowa State University, Ames. Unpagd pamphlet. (*D. barberi*, *D. virgifera virgifera*, biology, pest management, sampling [adult], crop rotation, extended diapause, insecticide recommendations, silk feeding, sampling [sequential])
182. Hein, G.L., and J.J. Tollefson. 1984. Comparison of adult corn rootworm (Coleoptera: Chrysomelidae) trapping techniques as population estimators. Environ. Entomol. 13(1):

266–271. (*D. barberi*, *D. virgifera virgifera*, sampling [adult], traps [adult], sex ratio, population measurement)

183. **Hein, G.L., and J.J. Tollefson.** 1985. Seasonal oviposition of northern and western corn rootworms (Coleoptera: Chrysomelidae) in continuous cornfields. *J. Econ. Entomol.* 78(6): 1238–1241. (*D. barberi*, *D. virgifera virgifera*, temperature, heat units)

184. **Hein, G.L., and J.J. Tollefson.** 1985. Use of the Pherocon AM trap as a scouting tool for predicting damage by corn rootworm (Coleoptera: Chrysomelidae) larvae. *J. Econ. Entomol.* 78(1): 200–203. (*D. barberi*, *D. virgifera virgifera*, sampling [adult], root damage ratings, traps [adult], prediction of damage)

185. **Hein, G.L., and J.J. Tollefson.** 1987. Model of the biotic potential of western corn rootworm (Coleoptera: Chrysomelidae) adult populations, and its use in studying population dynamics. *Environ. Entomol.* 16(2): 446–452. (*D. virgifera virgifera*, adult emergence, sampling [adult], sampling [egg], mortality, dispersal, movement [adult])

186. **Hein, G.L., J.J. Tollefson, and R.E. Foster.** 1988. Adult northern and western corn rootworm (Coleoptera: Chrysomelidae) population dynamics and oviposition. *J. Kansas Entomol. Soc.* 61(2): 214–223. (*D. barberi*, *D. virgifera virgifera*, adult emergence, sampling [adult], sampling [egg], sex ratio, survival, mortality, population density, weather, movement [adult], dispersal)

187. **Hein, G.L., J.J. Tollefson, and P.N. Hinz.** 1985. Design and cost considerations in the sampling of northern and western corn rootworm (Coleoptera: Chrysomelidae) eggs. *J. Econ. Entomol.* 78(6): 1495–1499. (*D. barberi*, *D. virgifera virgifera*, sampling [egg], oviposition)

188. **Hellman, J.L., G.P. Dively, A. Brown, J.M. Luna, and M. Graustein.** 1988. Pest control recommendations for field crops. Bull. 237. Coop. Ext. Serv., University of Maryland, College Park, and University of Delaware, Newark. (*D. barberi*, *D. virgifera virgifera*, pest management, insecticide recommendations, sampling [adult])

♦ 189. **Henn, R.A.** 1981. Western corn rootworm larvae and lesion nematodes on corn roots. M.S. Thesis, Iowa State University, Ames. (*D. virgifera virgifera*)

190. **Hibbard, B.E., and L.B. Bjostad.** 1988. Behavioral responses of western corn rootworm larvae to volatile semiochemicals from corn seedlings. *J. Chem. Ecol.* 14(6): 1523–1539. (*D. virgifera virgifera*, plant volatiles, carbon dioxide, kairomones, attractants)

191. **Higgins, R.A., H.L. Brooks, and P.E. Sloderbeck.** 1989. Insect management for corn in 1989. Publ. MF-810 (rev., Entomology 126). Coop. Ext. Serv., Kansas State University, Manhattan. 16 pp. (*D. barberi*, *D. virgifera virgifera*, crop rotation, pest management, insecticide recommendations, sampling [adult], adult control)

192. **Higgins, R.A., T.J. Gibb, and G.E. Wilde.** 1988. Corn rootworm management in Kansas field corn. Publ. MF-845 (rev., Entomology 128). Coop. Ext. Serv., Kansas State University, Manhattan. 4 pp. (*D. barberi*, *D. virgifera virgifera*, pest management, biology, sampling [adult], adult control)

193. **Hill, R.E., and Z.B. Mayo.** 1980. Distribution and abundance of corn rootworm species as influenced by topography and crop rotation in eastern Nebraska. *Environ. Entomol.* 9(1): 122–127. (*D. barberi* as *D. longicornis*, *D. virgifera virgifera*, behavior, continuous cornfields, oviposition, movement [adult], dispersal, distribution [geographical], population density)

194. **Hintz, A.M., and B.W. George.** 1979. Successful laboratory hybridization of *Diabrotica virgifera* (western corn rootworm) and *Diabrotica longicornis* (northern corn rootworm) (Coleoptera: Chrysomelidae). *J. Kansas Entomol. Soc.* 52(2): 324–330. (*D. barberi* as *D. longicornis*, *D. virgifera virgifera*, mating)

195. **Homan, H.W., C.R. Baird, G.W. Bishop, J. Blackmer, and D.G. Bolz.** 1984. The western corn rootworm. Current information series 533 (rev.). Coop. Ext. Serv., University of Idaho, Moscow. Unpaginated pamphlet. (*D. virgifera virgifera*, biology, pest management, adult control, insecticide recommendations)

♦ 196. **Hooten, R.S.** 1979. Egg hatching patterns for individual western corn rootworm females. M.S. Thesis, South Dakota State University, Brookings. (*D. virgifera virgifera*, egg hatch)

♦ 197. **Huckla, D.L.** 1981. The integrated corn pest management program in Missouri: implementation, pest problems, pesticide usage, and grower acceptance. M.S. Thesis, University of Missouri, Columbia. (*Diabrotica* spp.)

198. **Hudon, M., and I. Ogilvie.** 1984. Corn pest management. Pages 98–102 in Proc. Intl. Symp.: The Role of Biological Control in Pest Management. IOBC/WHRS. November 16–20, 1981. Santiago, Chile. University Press, Ottawa, Canada. (*D. barberi* as *D. longicornis*, host plant resistance)

199. **Irwin, B.J.** 1977. A bibliography of the northern corn rootworm and the western corn rootworm: an updating through 1976. Ill. Nat. Hist. Sur. Biol. Notes 101. 8 pp. (*D. barberi*, *D. virgifera virgifera*)

200. **Jackson, J.J.** 1985. *Diabrotica* spp. Pages 237–254 in P. Singh and R. F. Moore, eds. Handbook of insect rearing. Vol. 1. Elsevier Science Publ., Amsterdam. (*D. virgifera virgifera*, diet [adult], pathogens [insect])

201. **Jackson, J.J.** 1985. Parasitism of the western corn rootworm with the nematode, *Steinernema feltiae*. Ph.D. Thesis, University of Minnesota, St. Paul. (*D. virgifera virgifera*, parasites, temperature, soil moisture, environmental factors, biological control)

202. **Jackson, J.J.** 1986. Rearing and handling of *Diabrotica virgifera* and *Diabrotica undecimpunctata howardi*. Pages 25–47 in J. L. Krysan and T. A. Miller, eds. Methods for the study of pest *Diabrotica*. Springer-Verlag, New York. (*D. virgifera virgifera*, pathogens [insect])

203. **Jackson, J.J., and D.G. Davis.** 1978. Rearing western corn rootworm larvae on seedling corn (Coleoptera: Chrysomelidae). *J. Kansas Entomol. Soc.* 51(3): 353–355. (*D. virgifera virgifera*)

204. **Jackson, J.J., and N.C. Elliott.** 1988. Temperature dependent development of immature stages of the western corn rootworm, *Diabrotica virgifera virgifera* (Coleoptera: Chrysomelidae). *Environ. Entomol.* 17(2): 166–171. (threshold temperature, heat units)

205. **Jensen, S.G.** 1979. Laboratory transmission of maize chlorotic mottle virus by corn rootworms (abstract). Annual Meeting of the Am. Phytopathol. Soc., August 5–11, 1979. Phytopathology 69(9): 1033. (*D. barberi* as *D. longicornis*, *D. virgifera virgifera*, disease transmission, vector [disease], virus transmission, pathogens [plant])
206. **Jensen, S.G.** 1985. Laboratory transmission of maize chlorotic mottle virus by three species of corn rootworms. Plant Dis. 69(10): 864–868. (*D. barberi*, *D. virgifera virgifera*, disease transmission, vector [disease], virus transmission)
207. **Johnson, T.B., D.R. Raemisch, A.S. Las, and F.T. Turpin.** 1985. Response of western corn rootworm (Coleoptera: Chrysomelidae) adults to several semiochemicals in cornfields. J. Econ. Entomol. 78(1): 85–88. (*D. virgifera virgifera*, cucurbitacins, pheromone, *Cucurbita* spp.)
208. **Johnson, T.B., and F.T. Turpin.** 1985. Northern and western corn rootworm (Coleoptera: Chrysomelidae) oviposition in corn as influenced by foxtail populations and tillage systems. J. Econ. Entomol. 78(1): 57–60. (*D. barberi*, *D. virgifera virgifera*, no-till, *Setaria* spp., weeds, environmental factors, soil moisture, root damage ratings, survival, ovipositional preference)
209. **Johnson, T.B., F.T. Turpin, and M.K. Bergman.** 1984. Effect of foxtail infestation on corn rootworm larvae (Coleoptera: Chrysomelidae) under two corn-planting dates. Environ. Entomol. 13(5): 1245–1248. (*D. barberi*, *D. virgifera virgifera*, weeds, alternate food source, alternate host, *Setaria* spp., weed competition, population dynamics)
210. **Jolivet P., E. Petitpierre, and T.H. Hsiao, eds.** 1988. Biology of Chrysomelidae. Series Entomologica, Vol. 42. Kluwer Academic Publ., Boston, MA. (*D. barberi*, *D. virgifera virgifera*, feeding stimulant, cucurbitacins, metabolism, cucurbits, electrophoresis, genetics, taxonomy, nematodes, *Neoplectana carpo-capsae*, biological control)
211. **Kahler, A.L., A.E. Olness, G.R. Sutter, C.D. Dybing, and O.J. Devine.** 1985. Root damage by western corn rootworm and nutrient content in maize. Agron. J. 77(5): 769–774. (*D. virgifera virgifera*, nutrient uptake, compensatory root growth, yield losses, root damage ratings, mineral nutrition, root regrowth)
212. **Kahler, A.L., R.E. Telkamp, L.H. Penny, T.F. Brandon, and P.J. Fitzgerald.** 1985. Registration of NGSDCRW1(S2)C4 maize germplasm. Crop Sci. 25(1): 202. (*D. virgifera virgifera*, host plant resistance, root damage ratings, compensatory root growth, tolerance)
213. **Kantack, B.H.** 1988. South Dakota insecticide recommendations handbook. Pages 16–18 in Publ. EC 683. Coop. Ext. Serv., South Dakota State University, Brookings. (*D. barberi*, *D. virgifera virgifera*, sampling [adult], pest management)
- ♦214. **Karr, L.L.** 1984. Temporal efficiency of the Pherocon AM trap for adult corn rootworm sampling. M.S. Thesis, Iowa State University, Ames. (*Diabrotica* spp., sampling [adult], traps [adult], periodicity)
- ♦215. **Karr, L.L., and J.R. Coats.** 1985. Insecticidal activity of dextro-limonene, a monoterpenoid isolated from ethereal oils of citrus (abstract). 190th Am. Chemical Soc. National Meeting, Chicago. September 8–13, 1985.
216. **Karr, L.L., and J.R. Coats.** 1988. Insecticidal properties of d-limonene. J. Pestic. Sci. 13(2): 287–290. (*D. virgifera virgifera*, bioassay, ovicides)
217. **Karr, L.L., and J.J. Tollefson.** 1987. Durability of the Pherocon AM trap for adult western and northern corn rootworm (Coleoptera: Chrysomelidae) sampling. J. Econ. Entomol. 80(4): 891–896. (*D. barberi*, *D. virgifera virgifera*, environmental factors, temperature, trap color, sampling [adult], traps [adult], trap capture)
218. **Karren, J.B.** 1986. Corn rootworm control. Insect control series 21 (rev.). Coop. Ext. Serv., Utah State University, Logan. 6 pp. (*D. virgifera virgifera*, biology, insecticide recommendations, pest management, adult control)
219. **Kirk, V.M.** 1978. Drought cracks as oviposition sites for western and northern corn rootworms (abstract). Proc. N. Cent. Br. Entomol. Soc. Am., 1978, publ. 1979, 33: 25. (*D. barberi*, *D. virgifera virgifera*)
220. **Kirk, V.M.** 1979. Drought cracks as oviposition sites for western and northern corn rootworms (*Diabrotica*: Coleoptera). J. Kansas Entomol. Soc. 52(4): 769–776. (*D. barberi*, *D. virgifera virgifera*)
221. **Kirk, V.M.** 1981. Base of corn stalks as oviposition sites for western and northern corn rootworms (*Diabrotica*: Coleoptera). J. Kansas Entomol. Soc. 54(2): 255–262. (*D. barberi* as *D. longicornis*, *D. virgifera virgifera*, soil moisture, ovipositional preference)
222. **Kirk, V.M.** 1981. Corn rootworm: population reduction associated with the ant, *Lasius neoniger*. Environ. Entomol. 10(6): 966–967. (*D. barberi* as *D. longicornis*, *D. virgifera virgifera*, predation, biological control, ants)
223. **Kirk, V.M.** 1981. Earthworm burrows as oviposition sites for western and northern corn rootworms (*Diabrotica*: Coleoptera). J. Kansas Entomol. Soc. 54(1): 68–74. (*D. barberi* as *D. longicornis*, *D. virgifera virgifera*)
224. **Kirk, V.M.** 1982. Carabids: minimal role in pest management of corn rootworms. Environ. Entomol. 11(1): 5–8. (*D. barberi* as *D. longicornis*, *D. virgifera virgifera*, predation, biological control)
225. **Krysan, J.L.** 1978. Diapause, quiescence, and moisture in the egg of the western corn rootworm, *Diabrotica virgifera*. J. Insect Physiol. 24(6/7): 535–540. (*D. virgifera virgifera*, humidity, soil moisture, egg hatch)
226. **Krysan, J.L.** 1978. Egg dormancy in *Diabrotica*: embryological, geographical and taxonomic aspects (abstract). Proc. N. Cent. Br. Entomol. Soc. Am. 1978, publ. 1979, 33: 60. (*D. virgifera virgifera*, diapause, genetics, taxonomy, embryonic development, distribution [geographical])
227. **Krysan, J.L.** 1982. Diapause in the Nearctic species of the *virgifera* group of *Diabrotica*: evidence for tropical origin and temperature adaptations. Ann. Entomol. Soc. Am. 75(2): 136–142. (*D. barberi*, *D. virgifera virgifera*, *D. virgifera zaeae*, voltinism, evolution)
228. **Krysan, J.L.** 1986. Introduction: biology, distribution, and identification of pest *Diabrotica*. Pages 1–23 in J. L. Krysan and T. A. Miller, eds. Methods for the study of pest *Diabrotica*. Springer-Verlag, New York. (*D. barberi*, *D. virgifera virgifera*,

- D. virgifera zea*, distribution [geographical], taxonomy, identification [egg], identification [larval], identification [adult], sex determination, sex differences)
229. Krysan, J.L. 1987. A phenocline in the sculpturing of the egg chorion in the *virgifera* species group of *Diabrotica* (Coleoptera: Chrysomelidae). *Coleopterists Bull.* 41(4): 323-326. (*D. barberi*, *D. virgifera virgifera*, taxonomy)
  230. Krysan, J.L., and T.F. Branson. 1977. Inheritance of diapause intensity in *Diabrotica virgifera*. *J. Heredity* 68(6): 415-417. (*D. virgifera virgifera*, genetics, egg hatch)
  231. Krysan, J.L., and T.F. Branson. 1983. Biology, ecology, and distribution of *Diabrotica*. Pages 144-150 in D.T. Gordon, J.K. Knoke, L.R. Nault, and R.M. Ritter, eds. *Proc. Intl. Maize Virus Dis. Colloq. and Workshop*, August 2-6, 1982. Wooster, OH. (*D. barberi*, *D. virgifera virgifera*, *D. virgifera zea*, distribution [geographical], evolution, dispersal, host plant selection, movement [adult])
  232. Krysan, J.L., T.F. Branson, and G.D. Castro. 1977. Diapause in *Diabrotica virgifera* (Coleoptera: Chrysomelidae): a comparison of eggs from temperate and subtropical climates. *Ent. Exp. Appl.* 22(1): 81-89. (*D. virgifera virgifera*, evolution, humidity, soil moisture, egg hatch, temperature)
  233. Krysan, J.L., T.F. Branson, P.L. Guss, and R.D. Gustin. 1982. Corn rootworm biology. Pages 59-64 in Eighth Annual Illinois Crop Protection Workshop. Coop. Ext. Serv., University of Illinois at Urbana-Champaign. (*D. barberi*, *D. virgifera virgifera*, *D. virgifera zea*, distribution [geographical])
  234. Krysan, J.L., D.E. Foster, T.F. Branson, K.R. Ostlie, and W.S. Cranshaw. 1986. Two years before the hatch: rootworms adapt to crop rotation. *Bull. Entomol. Soc. Am.* 32(4): 250-253. (*D. barberi*, *D. virgifera virgifera*, extended diapause)
  235. Krysan, J.L., and P.L. Guss. 1978. Barriers to hybridization between *Diabrotica virgifera* and *D. longicornis barberi* (Coleoptera: Chrysomelidae). *Ann. Entomol. Soc. Am.* 71(6): 931-934. (*D. virgifera virgifera*, mating, egg hatch, reproductive isolation, fecundity, egg viability, insemination, fertility)
  236. Krysan, J.L., J.J. Jackson, and A.C. Lew. 1984. Field termination of egg diapause in *Diabrotica* with new evidence of extended diapause in *D. barberi* (Coleoptera: Chrysomelidae). *Environ. Entomol.* 13(5): 1237-1240. (*D. virgifera virgifera*, egg hatch duration, crop rotation, egg hatch)
  237. Krysan, J.L., and T.A. Miller, eds. 1986. *Methods for the study of pest Diabrotica*. Springer-Verlag, New York. (*D. barberi*, *D. virgifera virgifera*, *D. virgifera zea*, biology, distribution [geographical], taxonomy, rearing, pheromone, kairomones, sampling [egg], sampling [larval], sampling [adult], artificial infestation, host plant resistance, insecticide evaluation, bioassay, vector [disease])
  238. Krysan, J.L., and R.F. Smith. 1987. Systematics of the *virgifera* species group of *Diabrotica* (Coleoptera: Chrysomelidae: Galerucinae). *Entomography* 5: 375-484. (*D. barberi*, *D. virgifera virgifera*, *D. virgifera zea*, taxonomy, distribution [geographical], biology, host plant selection)
  239. Krysan, J.L., R.F. Smith, T.F. Branson, and P.L. Guss. 1980. A new subspecies of *Diabrotica virgifera* (Coleoptera: Chrysomelidae): description, distribution, and sexual compatibility. *Ann. Entomol. Soc. Am.* 73(2): 123-130. (*D. barberi*, *D. virgifera virgifera*, *D. virgifera zea*, distribution [geographical], mating, pheromone, taxonomy)
  240. Krysan, J.L., R.F. Smith, and P.L. Guss. 1983. *Diabrotica barberi* (Coleoptera: Chrysomelidae) elevated to species rank based on behavior, habitat choice, morphometrics, and geographical variation of color. *Ann. Entomol. Soc. Am.* 76(2): 197-204. (taxonomy, pheromone, mating, distribution [geographical], morphology, coloration [insect])
  241. Krysan, J.L., and G.R. Sutter. 1986. Aldrin susceptibility as an indicator of geographic variability in the northern corn rootworm, *Diabrotica barberi* (Coleoptera: Chrysomelidae). *Environ. Entomol.* 15(2): 427-430. (insecticide resistance, extended diapause, bioassay)
  242. Krysan, J.L., P.H. Wilkin, J.H. Tumlinson, P.E. Sonnet, R.L. Carney, and P.L. Guss. 1986. Responses of *Diabrotica lemniscata* and *D. longicornis* (Coleoptera: Chrysomelidae) to stereoisomers of 8-methyl-2-decyl-propanoate and studies on the pheromone of *D. longicornis*. *Ann. Entomol. Soc. Am.* 79(4): 742-746. (*D. barberi*, *D. virgifera virgifera*)
  243. Kuhlman, D.E. 1982. Silk feeding insects and economic thresholds. Pages 83-88 in Eighth Annual Illinois Crop Protection Workshop. Coop. Ext. Serv., University of Illinois at Urbana-Champaign. (*D. barberi* as *D. longicornis*, *D. virgifera virgifera*)
  244. Kuhlman, D. 1988. Bugs, corn, and set-aside acres. Pages 122-128 in Illinois Agricultural Pesticides Conf. 1988, Fortieth Spray School, summaries of presentations. Coop. Ext. Serv., University of Illinois at Urbana-Champaign. (*Diabrotica* spp., crop rotation, oviposition, cover crops)
  245. Kuhlman, D.E., and J.T. Shaw. 1982. Influence of crop rotation on corn rootworms. Pages 22-24 in Thirty-fourth Illinois Custom Spray Operators Training School, summaries of presentations. Coop. Ext. Serv., University of Illinois at Urbana-Champaign. (*D. barberi*, *D. virgifera virgifera*, oviposition, root damage ratings, soybeans)
  246. Kuhlman, D., and K. Steffey. 1987. Rootworms in corn after soybeans? Survey results and management suggestions. Pages 15-27 in Illinois Agricultural Pesticides Conf. '87, Thirty-ninth Spray School, summaries of presentations. Coop. Ext. Serv., University of Illinois at Urbana-Champaign. (*D. barberi*, *D. virgifera virgifera*, extended diapause, root damage ratings)
  247. Kuhlman, D.E., K.L. Steffey, and M.E. Gray. 1988. 1989 insect pest management guide: field and forage crops. Pages 6-9 in Circular 899 (rev.). Coop. Ext. Serv., University of Illinois at Urbana-Champaign. (*D. barberi*, *D. virgifera virgifera*, extended diapause, sampling [adult], insecticide recommendations, crop rotation, adult control, biology)
  248. Ladd, T.L., Jr. 1984. Eugenol-related attractants for the northern corn rootworm (Coleoptera: Chrysomelidae). *J. Econ. Entomol.* 77(2): 339-341. (*D. barberi*, traps [adult], semiochemicals, isoeugenol)
  249. Ladd, T.L., Jr., J.L. Krysan, and P.L. Guss. 1985. Corn rootworms (Coleoptera: Chrysomelidae): responses to eugenol and 8R-methyl-2R-decyl propanoate. *J. Econ. Entomol.* 78(4): 844-847. (*D. barberi*, *D. virgifera virgifera*, traps [adult], pheromone, attractants, semiochemicals)

250. Ladd, T.L., B.R. Stinner, and H.R. Krueger. 1983. Eugenol, a new attractant for the northern corn rootworm (Coleoptera: Chrysomelidae). J. Econ. Entomol. 76(5): 1049-1051. (*D. barberi*, *D. virgifera virgifera*, semiochemicals, traps [adult], movement [adult], dispersal)
251. Ladd, T.L., B.R. Stinner, and H.R. Krueger. 1983. Find new attractant for corn rootworm. Ohio Report 68(5): 67-69. (*D. barberi*, *D. virgifera virgifera*, eugenol, traps [adult], semiochemicals, movement [adult], dispersal)
252. Ladd, T.L., Jr., B.R. Stinner, and H.R. Krueger. 1984. Influence of color and height of eugenol-baited sticky traps on attractiveness to northern corn rootworm beetles (Coleoptera: Chrysomelidae). J. Econ. Entomol. 77(3): 652-654. (*D. barberi*, trap height, traps [adult], attractants, semiochemicals, trap color)
253. Lampman, R.L. 1986. Chemical ecology of defense compounds in *Anasa* species (Heteroptera: Coreidae) and volatile attractants for *Diabrotica* species (Coleoptera: Chrysomelidae). Ph.D. Thesis, University of Illinois at Urbana-Champaign. (*D. barberi*, *D. virgifera virgifera*, estragole, indole, eugenol, isoeugenol, phenology, sex ratio, semiochemicals, kairomones, plant volatiles)
254. Lampman, R.L., and R.L. Metcalf. 1987. Multi-component kairomonal lures for southern and western corn rootworms (Coleoptera: Chrysomelidae: *Diabrotica* spp.). J. Econ. Entomol. 80(6): 1137-1142. (*D. virgifera virgifera*, kairomones, attractants, semiochemicals, traps [adult], indole, eugenol)
255. Lampman, R.L., and R.L. Metcalf. 1988. The comparative response of *Diabrotica* species (Coleoptera: Chrysomelidae) to volatile attractants. Environ. Entomol. 17(4): 644-648. (*D. barberi*, *D. virgifera virgifera*, host plant selection, traps [adult], semiochemicals, kairomones, estragole, eugenol, indole, isoeugenol, plant volatiles)
256. Lampman, R.L., R.L. Metcalf, and J.F. Andersen. 1987. Semiochemical attractants of *Diabrotica undecimpunctata howardi* Barber, southern corn rootworm, and *Diabrotica virgifera virgifera* LeConte, the western corn rootworm (Coleoptera: Chrysomelidae). J. Chem. Ecol. 13(4): 959-975. (*D. barberi*, host plant selection, traps [adult], indole, eugenol, estragole, kairomones, isoeugenol)
257. Lance, D.R. 1988. Potential of 8-methyl-2-decyl propanoate and plant-derived volatiles for attracting corn rootworm beetles (Coleoptera: Chrysomelidae) to toxic bait. J. Econ. Entomol. 81(5): 1359-1362. (*D. barberi*, *D. virgifera virgifera*, semiochemicals, traps [adult], attractants, estragole, eugenol, pheromone, repellents, cucurbitacin vial traps, kairomones, adult control, plant volatiles)
258. Lance, D.R. 1988. Responses of northern and western corn rootworms to semiochemical attractants in corn fields. J. Chem. Ecol. 14(4): 1177-1185. (*D. barberi*, *D. virgifera virgifera*, pheromone, eugenol, repellents, disruption of mating, traps [adult], kairomones)
259. Lance, D.R., and J.R. Fisher. 1987. Food quality of various plant tissues for adults of the northern corn rootworm (Coleoptera: Chrysomelidae). J. Kansas Entomol. Soc. 60(3): 462-466. (*D. barberi*, oviposition, longevity, fecundity, rearing)
- ♦ 260. Lawson, E.C. 1986. Influence of tillage and depth in the soil on soil temperature and survival of overwintering western corn rootworm eggs. M.S. Thesis, Iowa State University, Ames. (*D. virgifera virgifera*)
261. Lazarus, W.F. 1981. Optimal management of a common property pest under risk: an application to the corn rootworm. Ph.D. Thesis, University of Illinois at Urbana-Champaign. (*D. barberi* as *D. longicornis*, *D. virgifera virgifera*, risk management, economics, economic thresholds, model, crop rotation, crop insurance, sampling [adult], insecticide resistance, movement [adult], genetics, soybeans)
262. Lazarus, W.F., and B.L. Dixon. 1984. Agricultural pests as common property: control of the corn rootworm. Am. J. Agric. Econ. 66(4): 456-465. (*D. barberi* as *D. longicornis*, *D. virgifera virgifera*, insecticide resistance, crop rotation, soybeans, economics)
263. Lazarus, W.F., and E.R. Swanson. 1983. Insecticide use and crop rotation under risk: rootworm control in corn. Am. J. Agric. Econ. 65(4): 738-747. (*D. barberi* as *D. longicornis*, *D. virgifera virgifera*, economic thresholds, crop insurance, soybeans, economics, risk management)
264. Levine, E. 1985. Corn insect research results, 1984. Pages 49-57 in Thirty-seventh Illinois Custom Spray Operators Training School manual. Coop. Ext. Serv., University of Illinois at Urbana-Champaign. (*D. barberi*, *D. virgifera virgifera*, oviposition, tillage, adult emergence, no-till)
265. Levine, E. 1987. Extended diapause and egg-hatching patterns in corn rootworms. Pages 11-14 in Illinois Agricultural Pesticides Conf. '87, Thirty-ninth Spray School, summaries of presentations. Coop. Ext. Serv., University of Illinois at Urbana-Champaign. (*D. barberi*, *D. virgifera virgifera*, egg hatch, egg hatch duration)
266. Levine, E., D. Kuhlman, K. Steffey, and H. Oloumi-Sadeghi. 1988. New developments regarding extended diapause in northern corn rootworms: research and survey results. Pages 145-153 in Illinois Agricultural Pesticides Conf. '88, Fortieth Spray School, summaries of presentations. Coop. Ext. Serv., University of Illinois at Urbana-Champaign. (*D. barberi*, *D. virgifera virgifera*, crop rotation, soybeans, root damage ratings, egg hatch)
267. Levine, E., and R.L. Metcalf. 1988. Sticky attractant traps for monitoring corn rootworm beetles. Ill. Nat. Hist. Sur. Reports 279: [1-2]. (*D. barberi*, *D. virgifera virgifera*, traps [adult], semiochemicals, kairomones)
268. Levine, E., H. Oloumi-Sadeghi, R.L. Metcalf, and R. Lampman. 1988. Dry cucurbitacin-containing baits for controlling adult western corn rootworms, *Diabrotica virgifera virgifera* (Coleoptera: Chrysomelidae), in field corn. Cucurbit Genetics Coop. 11: 79-82. (oviposition, adult control, mortality)
269. Lew, A.C. 1979. Pheromone secretory cells and the behavioral aspect of mating in the western corn rootworm, *Diabrotica virgifera* LeConte. Ph.D. Thesis, University of Nebraska, Lincoln. (*D. virgifera virgifera*, spermatophore, reproductive behavior)
270. Lew, A.C., and H.J. Ball. 1978. Pheromone secretory cells of *Diabrotica virgifera* LeConte (abstract). Proc. N. Cent. Br. Entomol. Soc. Am., 1978, publ. 1979, 33: 59-60. (*D. virgifera virgifera*)

271. **Lew, A.C., and H.J. Ball.** 1978. The structure of the apparent pheromone-secreting cells in female *Diabrotica virgifera*. *Ann. Entomol. Soc. Am.* 71(5): 685-688. (*D. virgifera virgifera*, morphology)
272. **Lew, A.C., and H.J. Ball.** 1979. The mating behavior of the western corn rootworm *Diabrotica virgifera* (Coleoptera: Chrysomelidae). *Ann. Entomol. Soc. Am.* 72(3): 391-393. (*D. virgifera virgifera*, reproductive behavior)
273. **Lew, A.C., and H.J. Ball.** 1980. Effect of copulation time on spermatozoan transfer of *Diabrotica virgifera* (Coleoptera: Chrysomelidae). *Ann. Entomol. Soc. Am.* 73(4): 360-361. (*D. virgifera virgifera*, mating, spermatophore, reproductive status, mating status)
274. **Lew, A.C., and G.R. Sutter.** 1985. Toxicity of insecticides to northern corn rootworm (Coleoptera: Chrysomelidae) larvae. *J. Kansas Entomol. Soc.* 58(3): 547-549. (*D. barberi*, *D. virgifera virgifera*, rearing, bioassay, terbufos, carbofuran, insecticide resistance, phorate, chlorpyrifos, isofenphos, ethoprop, fonofos)
- ♦275. **Lockwood, M.J.** 1978. Western corn rootworm sex pheromone; field trapping studies. M.S. Thesis, South Dakota State University, Brookings. (*D. virgifera virgifera*, pheromone traps)
276. **Lockwood, M.J., and D.D. Walgenbach.** 1978. Western corn rootworm pheromone; field studies (abstract). *Proc. N. Cent. Br. Entomol. Soc. Am.*, 1978, publ. 1979, 33: 60. (*D. virgifera virgifera*, sampling [adult], pheromone traps)
277. **Lummus, P.F., J.C. Smith, and N.L. Powell.** 1983. Soil moisture and texture effects on survival of immature southern corn rootworms, *Diabrotica undecimpunctata howardi* Barber (Coleoptera: Chrysomelidae). *Environ. Entomol.* 12(5): 1529-1531. (*D. virgifera virgifera*, soil type)
- ♦278. **MacDonald, P.J.** 1988. Mortality of overwintering eggs and longevity and movement of first instars of corn rootworm (Coleoptera: Chrysomelidae). M.S. Thesis, University of Guelph, Guelph, Ontario, Canada. (*Diabrotica* spp., movement [larval])
279. **Madder, D.J., and M. Stemeroff.** 1988. The economics of insect control on wheat, corn and canola in Canada, 1980-1985. *Bull. Entomol. Soc. Can.*, 20 (2, supplement), 22 pp. (*D. barberi*, *D. virgifera virgifera*)
280. **Matin, M.A.** 1983. Development and application of population sampling methods for the stages of northern corn rootworm, *Diabrotica longicornis* (Say) (Coleoptera: Chrysomelidae) in Quebec corn fields. Ph.D. Thesis, McGill University, Ottawa, Ontario, Canada. (*D. barberi* as *D. longicornis*, sampling [egg], sampling [larval], sampling [adult], sampling statistics, distribution [spatial])
281. **Matin, M. A., and W.N. Yule.** 1984. Effects of corn phenology on spatial distribution and population estimation of adult northern corn rootworms (Coleoptera: Chrysomelidae). *J. Econ. Entomol.* 77(5): 1176-1179. (*D. barberi*, sampling [adult], dispersion [adult], distribution [spatial], population measurement)
282. **Matin, M.A., and W.N. Yule.** 1984. Population fluctuations of northern corn rootworm, (Coleoptera: Chrysomelidae) in a Quebec cornfield. *Revue d'Entomologie du Québec* 29(2): 74-82. (*D. barberi*, population dynamics, mortality, tillage, environmental factors)
283. **Matin, M.A., and W.N. Yule.** 1986. Techniques for extracting northern corn rootworm eggs from soil. *Pedobiologia* 29(1): 21-26. (*D. barberi*, sampling [egg], egg separation)
284. **Matin, M.A., W.N. Yule, and P. Martel.** 1984. Comparison of three methods for sampling northern corn rootworm (Coleoptera: Chrysomelidae) adult populations in two Quebec cornfields. *J. Econ. Entomol.* 77(5): 1344-1348. (*D. barberi*, emergence traps, sampling [adult], adult emergence)
285. **Mayo, Z. B.** 1977. Control of larvae of the western and northern corn rootworm with liquid starter fertilizer-insecticide combinations and the influence of depth of placement. *J. Econ. Entomol.* 70(2): 234-236. (*D. barberi* as *D. longicornis*, *D. virgifera virgifera*, insecticide placement, root damage ratings, carbofuran, ethoprop, fonofos, fensulfothion, insecticide evaluation)
286. **Mayo, Z. B.** 1980. Influence of planting dates on the efficacy of soil insecticides applied to control larvae of the western and northern corn rootworm. *J. Econ. Entomol.* 73(2): 211-212. (*D. barberi* as *D. longicornis*, *D. virgifera virgifera*, insecticide evaluation, root damage ratings, carbofuran, fensulfothion, diazinon, phorate, fonofos, terbufos)
287. **Mayo, Z. B.** 1984. Influences of rainfall and sprinkler irrigation on the residual activity of insecticides applied to corn for control of adult western corn rootworm (Coleoptera: Chrysomelidae). *J. Econ. Entomol.* 77(1): 190-193. (*D. barberi* as *D. longicornis*, *D. virgifera virgifera*, bioassay, carbaryl, phosmet, PennCap-M, diazinon, malathion, adult control, insecticide residues, insecticide evaluation)
288. **Mayo, Z. B., Jr.** 1986. Field evaluation of insecticides for control of larvae of corn rootworms. Pages 183-203 in J. L. Krysan and T. A. Miller, eds. *Methods for the study of pest Diabrotica*. Springer-Verlag, New York. (*D. barberi*, *D. virgifera virgifera*, insecticide evaluation, root damage ratings, lodging, yield losses, insecticide application equipment)
289. **Mayo, Z. B., and L.L. Peters.** 1978. Planting vs. cultivation time applications of granular soil insecticides to control larvae of corn rootworms in Nebraska. *J. Econ. Entomol.* 71(5): 801-803. (*D. barberi* as *D. longicornis*, *D. virgifera virgifera*, insecticide evaluation, carbofuran, terbufos, ethoprop, fonofos, diazinon, phorate, fensulfothion, bufencarb, root damage ratings)
- ♦290. **McAuslane, H.J.** 1986. Monitoring for corn rootworms (Coleoptera: Chrysomelidae) in field corn in southern Ontario. M.S. Thesis, University of Guelph, Guelph, Ontario, Canada. (*Diabrotica* spp., sampling [adult])
291. **McAuslane, H.J., C.R. Ellis, and O.B. Allen.** 1987. Sequential sampling of adult northern and western corn rootworms (Coleoptera: Chrysomelidae) in southern Ontario. *Can. Entomol.* 119(6): 577-585. (*D. barberi*, *D. virgifera virgifera*, sampling [adult], economic thresholds, continuous cornfields, first-year cornfields, sampling [sequential])
292. **McAuslane, H.J., C.R. Ellis, and P.E.A. Teal.** 1986. Chemical attractants for monitoring for adult northern and western corn rootworms (Coleoptera: Chrysomelidae) in Ontario. *Proc. Entomol. Soc. Ont.* 117: 49-57. (*D. barberi*, *D. virgifera virgifera*, traps [adult], trap color, pheromone, semiochemicals, kairomones, sampling [adult])

293. McDonald, I.C., J.L. Krysan, and O.A. Johnson. 1982. Genetics of *Diabrotica* (Coleoptera: Chrysomelidae): inheritance of xanthine dehydrogenase, hexokinase, malate dehydrogenase, and esterase allozymes in two subspecies of *D. longicornis*. Ann. Entomol. Soc. Am. 75(4): 460–464. (*D. barberi*)
294. McDonald, I.C., J.L. Krysan, and O.A. Johnson. 1985. Genetic variation within and among geographic populations of *Diabrotica barberi* (Coleoptera: Chrysomelidae). Ann. Entomol. Soc. Am. 78(3): 271–278. (genetics)
295. McGovern, T.P., and T.L. Ladd, Jr. 1988. 2-methoxyphenols, 1, 2-dimethoxybenzenes, and 1, 3-benzodioxoles as attractants for the northern corn rootworm (Coleoptera: Chrysomelidae). J. Econ. Entomol. 81(3): 826–829. (*D. barberi*, semiochemicals, kairomones, eugenol, isoeugenol)
296. Metcalf, R.L. 1979. Plants, chemicals, and insects: some aspects of coevolution. Bull. Entomol. Soc. Am. 25(1): 30–35. (*D. barberi* as *D. longicornis*, *D. virgifera virgifera*, cucurbitacins, host plant selection, kairomones, semiochemicals, *Cucurbita* spp., Cucurbitaceae, evolution)
297. Metcalf, R.L. 1986. Coevolutionary adaptations of rootworm beetles (Coleoptera: Chrysomelidae) to cucurbitacins. J. Chem. Ecol. 12(5): 1109–1124. (*D. barberi*, *D. virgifera virgifera*, kairomones, semiochemicals, host plant selection, *Cucurbita* spp., Cucurbitaceae, evolution)
298. Metcalf, R.L. 1986. Foreword. Pages vii–xv in J. L. Krysan and T. A. Miller, eds. Methods for the study of pest *Diabrotica*. Springer-Verlag, New York. (*D. barberi*, *D. virgifera virgifera*, yield losses, insecticide resistance, semiochemicals, plant volatiles, cucurbitacins, pest management)
299. Metcalf, R.L. 1987. Plant volatiles as insect attractants. CRC Critical Reviews in Plant Sci. 5(3): 251–301. (*D. barberi*, *D. virgifera virgifera*, kairomones, semiochemicals, estragole, eugenol, isoeugenol, indole, *Cucurbita* spp., Cucurbitaceae, host plant selection)
- ♦300. Metcalf, R.L., J.E. Ferguson, D. Fischer, R. Lampman, and J. Andersen. 1983. Controlling cucumber beetles and corn rootworm beetles with baits of bitter cucurbit fruits and root. Cucurbit Genetics Coop. 6: 79–81. (*Diabrotica* spp.)
301. Metcalf, R.L., J.E. Ferguson, R. Lampman, and J.F. Andersen. 1987. Dry cucurbitacin-containing baits for controlling diabrotic beetles (Coleoptera: Chrysomelidae). J. Econ. Entomol. 80(4): 870–875. (*D. virgifera virgifera*, arrestants [feeding], eugenol, *Cucurbita* spp., Cucurbitaceae, adult control)
302. Metcalf, R.L., R.A. Metcalf, and A.M. Rhodes. 1980. Cucurbitacins as kairomones for diabrotic beetles. Proc. National Acad. Sci. 77(7): 3769–3772. (*D. barberi* as *D. longicornis*, *D. virgifera virgifera*, host plant selection, evolution, arrestants [feeding])
303. Metcalf, R.L., and A.M. Rhodes. 1982. Cucurbitacins and rootworm control. Pages 65–67 in Eighth Annual Illinois Crop Protection Workshop. Coop. Ext. Serv., University of Illinois at Urbana-Champaign. (*D. barberi* as *D. longicornis*, *D. virgifera virgifera*, baits, kairomones, semiochemicals, host plant selection, *Cucurbita* spp., Cucurbitaceae, adult control)
304. Metcalf, R.L., A.M. Rhodes, R.A. Metcalf, J. Ferguson, E.R. Metcalf, and P-Y. Lu. 1982. Cucurbitacin contents and diabroticite (Coleoptera: Chrysomelidae) feeding upon *Cucurbita* spp. Environ. Entomol. 11(4): 931–937. (*D. virgifera virgifera*, kairomones, semiochemicals, host plant selection, Cucurbitaceae)
- ♦305. Miller, K.G. 1985. Root rating–corn yield interactions. M.S. Thesis, South Dakota State University, Brookings. (*Diabrotica* spp., root damage ratings, yield losses)
306. Mitchell, B.K. 1988. Adult leaf beetles as models for exploring the chemical basis of host-plant recognition. J. Insect Physiol. 34(3): 213–225. (*D. virgifera virgifera*, indole, attractants, host plant selection)
307. Montgomery, M.E., G.J. Musick, J.B. Polivka, and D.G. Nielsen. 1979. Modifiable washing-flotation method for separation of insect eggs and larvae from soil. J. Econ. Entomol. 72(1): 67–69. (*D. barberi* as *D. longicornis*, flotation [egg], flotation [larval], egg separation)
- ♦308. Mori, K., and H. Watanabe. 1984. Synthesis of the propionate of (2R, 8R)- and (2S, 8R)-8-methyl-2-decanol, the pheromones of western corn rootworms (*Diabrotica virgifera virgifera*), employing chiral compounds of microbial origin as starting materials. Tetrahedron 40(2): 299–304.
309. Munson, G. 1989. Insect pests of corn. Pages 56–76 in 1989 Missouri insect control recommendations. Publ. MX-3. Coop. Ext. Serv., University of Missouri and Lincoln University, Columbia. (*D. barberi*, *D. virgifera virgifera*, insecticide recommendations)
310. Munson G., A.J. Keaster, J.A. Grundler. 1985. Soil insect control in reduced tillage cropping systems. Agric. Guide 4906 (rev.). Coop. Ext. Serv., University of Missouri, Columbia. Unpagged pamphlet. (*D. barberi*, *D. virgifera virgifera*, no-till, pest management, insecticide recommendations, crop rotation)
311. Munson, G., A.J. Keaster, J.A. Grundler. 1986. Corn rootworm control. Agric. Guide 4151 (rev.). Coop. Ext. Serv., University of Missouri, Columbia. Unpagged pamphlet. (*D. barberi*, *D. virgifera virgifera*, biology, pest management, crop rotation, insecticide recommendations)
312. Musick, G.J. 1987. History, perspective, and overview of entomological research in conservation tillage systems. Pages 1–10 in G. J. House and B. R. Stinner, eds. Arthropods in conservation tillage systems. Misc. Publ. 65, Entomol. Soc. Am., College Park, MD. (*D. barberi* as *D. longicornis*, *D. virgifera virgifera*, oviposition, no-till, mortality)
313. Musick, G.J., H.C. Chiang, W.H. Luckmann, Z B Mayo, and F.T. Turpin. 1980. Impact of planting dates of field corn on beetle emergence and damage by the western and the northern corn rootworms in the corn belt. Ann. Entomol. Soc. Am. 73(2): 207–215. (*D. barberi* as *D. longicornis*, *D. virgifera virgifera*, adult emergence, emergence traps, root damage ratings)
314. Naranjo, S.E. 1987. Population dynamics and oviposition of the northern corn rootworm, *Diabrotica barberi*: significance of host plant phenology. Ph.D. Thesis, Cornell University, Ithaca, NY. (model, temperature, reproductive development, rearing, diet [adult], planting date, adult emergence, sex ratio, mortality, flight, reproductive status, dispersal, movement [adult])
315. Naranjo, S.E., and A.J. Sawyer. 1987. Reproductive biology and survival of *Diabrotica barberi* (Coleoptera:

- Chrysomelidae): effect of temperature, food, and seasonal time of emergence. *Ann. Entomol. Soc. Am.* 80(6): 841-848. (*D. virgifera virgifera*, longevity, fecundity, food quality, pollen, reproductive development, reproductive status, rearing, oviposition)
316. Naranjo, S.E., and A.J. Sawyer. 1988. Impact of host plant phenology on the population dynamics and oviposition of northern corn rootworms, *Diabrotica barberi* (Coleoptera: Chrysomelidae), in field corn. *Environ. Entomol.* 17(3): 508-521. (adult emergence, mortality, dispersal, emergence traps, sex ratio, planting date)
317. Naranjo, S.E., and A.J. Sawyer. 1988. A temperature- and age-dependent simulation model of reproduction for the northern corn rootworm, *Diabrotica barberi* Smith and Lawrence (Coleoptera: Chrysomelidae). *Can. Entomol.* 120(1): 1-17. (oviposition, reproductive development, reproductive status)
318. Naranjo, S.E., and D.C. Steinkraus. 1988. Discovery of an entomophthoralean fungus (Zygomycetes: Entomophthorales) infecting adult northern corn rootworm, *Diabrotica barberi* (Coleoptera: Chrysomelidae). *J. Invert. Pathol.* 51(3): 298-300. (biological control, pathogens [insect])
319. Nault, L.R., W.E. Styer, M.E. Coffey, D.T. Gordon, L.S. Negi, and C.L. Niblett. 1978. Transmission of maize chlorotic mottle virus by chrysomelid beetles. *Phytopathology* 68(7): 1071-1074. (*D. barberi* as *D. longicornis*, *D. virgifera virgifera*, vector [disease], virus transmission, pathogens [plant])
- ♦320. Newton, J.P. 1978. Microbial degradation of carbofuran in a Nebraska corn field. M.S. Thesis, University of Nebraska, Lincoln. (*Diabrotica* spp., insecticide degradation, biodegradation [enhanced])
321. Ostlie, K.R. 1987. Extended diapause: northern corn rootworm adapts to crop rotation. *Crops Soils Mag.* 39(8): 23-25. (*D. barberi*, root damage ratings)
322. Oyervides-Garcia, M. 1984. Selection-induced differences among Iowa stiff stalk synthetic strains. Ph.D. Thesis, Iowa State University, Ames. (*D. virgifera virgifera*, genetics, host plant resistance)
323. Oyervides-Garcia, M., and A.R. Hallauer. 1986. Selection-induced differences among strains of Iowa stiff stalk synthetic maize. *Crop Sci.* 26(3): 506-511. (*D. virgifera virgifera*, host plant resistance, yield losses, lodging, genetics)
324. Palmer, D.F. 1983. Embryologic studies of the western corn rootworm, *Diabrotica virgifera virgifera* LeConte: developmental patterns, embryonic stages, viability and diapause of eggs of different specific gravities. Ph.D. Thesis, University of Minnesota, St. Paul. (embryonic development, egg hatch, egg viability)
325. Palmer, D.F., and H.C. Chiang. 1978. Post-diapause embryonic development of western corn rootworm eggs of varying specific gravities (abstract). *Proc. N. Cent. Br. Entomol. Soc. Am.* 1978, publ. 1979, 33: 60. (*D. virgifera virgifera*)
326. Palmer, D.F., L.K. French, and H.C. Chiang. 1977. Effects of chill temperatures on developing western corn rootworm eggs. *Environ. Entomol.* 6(6): 862-864. (*D. virgifera virgifera*, survival, egg viability, embryonic development, egg hatch duration, cold hardness)
327. Palmer, D.F., M.B. Windels, and H.C. Chiang. 1977. Artificial infestation of corn with western corn rootworm eggs in agar-water. *J. Econ. Entomol.* 70(3): 277-278. (*D. virgifera virgifera*, root damage ratings)
328. Peairs, F.B., and S.D. Pilcher. 1988. Western corn rootworm management. Service in Action 5,570. Coop. Ext. Serv., Colorado State University, Ft. Collins. 2 pp. (*D. virgifera virgifera*, crop rotation, biology, insecticide recommendations, pest management)
329. Peters, L.L., L.J. Meinke, and J.F. Witkowski. 1987. Corn rootworm control. NebGuide G87-839. Coop. Ext. Serv., University of Nebraska, Lincoln. 4 pp. (*D. barberi*, *D. virgifera virgifera*, biology, pest management, crop rotation, adult control)
- ♦330. Piedrahita, O.H. 1984. Electrophoretic identification of larvae of two *Diabrotica* sp. (Coleoptera: Chrysomelidae) and their competition and movement. M.S. Thesis, University of Guelph, Guelph, Ontario, Canada. (taxonomy, electrophoresis, identification [larval], movement [larval])
331. Piedrahita, O., C.R. Ellis, and O.B. Allen. 1985. Effect of spacing and clumping of corn plants on density of corn rootworm larvae (Coleoptera: Chrysomelidae) per root system. *Can. Entomol.* 117(2): 139-142. (*Diabrotica* spp., distribution [spatial], sampling [larval], density [larval])
332. Piedrahita, O., C.R. Ellis, and J.P. Bogart. 1985. Electrophoretic identification of the larvae of *Diabrotica barberi* and *D. virgifera virgifera* (Coleoptera: Chrysomelidae). *Ann. Entomol. Soc. Am.* 78(4): 537-540. (electrophoresis, identification [larval], taxonomy)
333. Piedrahita, O., C.R. Ellis, and J.P. Bogart. 1985. Interaction of northern and western corn rootworm larvae (Coleoptera: Chrysomelidae) in a controlled environment. *Environ. Entomol.* 14(2): 138-141. (*D. barberi*, *D. virgifera virgifera*, competition, survival, distribution [spatial])
334. Pimentel, D., C. Shoemaker, and R.J. Whitman. 1978. Systems management program for corn pest control in New York State. *Search Agric.* 8(1): 1-16. (*D. barberi*, pest management, sampling [larval], sampling [pupal], yield losses, crop rotation, economics)
335. Poinar, G.O., Jr., J.S. Evans, and E. Schuster. 1982. Use of *Neoplectana carpocapsae* for control of corn rootworm larvae (*Diabrotica* sp. Coleoptera). *IRCS Medical Sci.: Biochem.: Environ. Biol. and Medicine: Microbiology, Parasitology and Infectious Diseases* 10: 1041. (biological control, nematodes)
336. Poinar, G.O., Jr., J.S. Evans, and E. Schuster. 1983. Field test of the entomogenous nematode, *Neoplectana carpocapsae*, for control of corn rootworm larvae (*Diabrotica* sp., Coleoptera). *Prot. Ecol.* 5: 337-342. (biological control)
337. Prystupa, B., C.R. Ellis, and P.E.A. Teal. 1988. Attraction of adult *Diabrotica* (Coleoptera: Chrysomelidae) to corn silks and analysis of the host-finding response. *J. Chem. Ecol.* 14(2): 635-651. (*D. barberi*, *D. virgifera virgifera*, host plant selection, attractants, bioassay)
338. Raemisch, D.R. 1985. The influence of bitter cucurbit fruit on the biology and behavior of western corn rootworm adults. Ph.D. Thesis, Purdue University, West Lafayette, IN. (*D. virgifera virgifera*, *Cucurbita* spp., attractants, semiochemicals, kairomones, aggregation pheromone, fecundity, longevity)

339. **Raemisch, D.R., and F.T. Turpin.** 1984. Field tests for an adult western corn rootworm aggregation pheromone associated with the phagostimulatory characteristic of bitter *Cucurbita* spp. J. Agric. Entomol. 1(4): 339–344. (*D. virgifera virgifera*, cucurbitacins)
340. **Rhodes, A.M., R.L. Metcalf, and E.R. Metcalf.** 1980. Diabrotic beetle responses to cucurbitacin kairomones in *Cucurbita* hybrids. J. Amer. Soc. Hort. Sci. 105(6): 838–842. (*D. virgifera virgifera*, semiochemicals)
341. **Rosen, P.P., D.A. Gabrielson, and I.C. McDonald.** 1988. Identification of a sex-limited esterase gene from *Diabrotica virgifera virgifera*. Proc. N. D. Acad. Sci. 42: 50.
342. **Ruesink, W.G.** 1986. Egg sampling techniques. Pages 83–99 in J. L. Krysan and T. A. Miller, eds. Methods for the study of pest *Diabrotica*. Springer-Verlag, New York. (*D. barberi*, *D. virgifera virgifera*, distribution [spatial], sampling [egg], egg separation, flotation [egg], sampling statistics)
343. **Ruesink, W.G., and J.T. Shaw.** 1983. Evaluation of a trench method for sampling eggs of the northern and western corn rootworms (Coleoptera: Chrysomelidae). J. Econ. Entomol. 76(5): 1195–1198. (*D. barberi*, *D. virgifera virgifera*)
- ♦344. **Ruppel, R.F.** 1977. Control of field corn insects. Extension Bull. E-828. Coop. Ext. Serv., Michigan State University, East Lansing. 6 pp. (*Diabrotica* spp., pest management)
345. **Ruppel, R.F.** 1983. Protecting corn from corn rootworms. Extension Bull. E-736. Coop. Ext. Serv., Michigan State University, East Lansing. 4 pp. (*D. barberi*, *D. virgifera virgifera*, biology, insecticide recommendations, crop rotation, pest management, adult control)
346. **Ruppel, R.F., H.L. Russell, and S.J. Jennings.** 1978. Indices for projecting emergence of corn rootworm adults in Michigan. J. Econ. Entomol. 71(6): 947–949. (*D. barberi* as *D. longicornis*, *D. virgifera virgifera*, heat units, temperature, adult emergence, emergence traps)
347. **Ruud, S.K., I.C. McDonald, D.A. Gabrielson, and O.A. Johnson.** 1988. Species distribution and characterization studies of sex-limited esterases within the *virgifera* group of *Diabrotica* (Coleoptera: Chrysomelidae). Ann. Entomol. Soc. Am. 81(6): 954–963. (*D. barberi*, *D. virgifera virgifera*, *D. virgifera zaeae*, genetics, electrophoresis, taxonomy)
348. **Sawyer, A.J.** 1985. Efficient monitoring of the density of adult northern corn rootworm (Coleoptera: Chrysomelidae) in field corn. Can. Entomol. 117(2): 171–183. (*D. barberi*, sampling statistics, sampling [adult], sampling [sequential])
349. **Sechriest, R.E., L. Crain, C. Johnson, and R. Foster.** 1977. Control of corn rootworms by adult suppression. Pages 72–76 in Twenty-ninth Illinois Custom Spray Operators Training School, summaries of presentations. Coop. Ext. Serv., University of Illinois at Urbana-Champaign. (*D. barberi*, *D. virgifera virgifera*, adult control, oviposition, root damage ratings, yield losses, sampling [adult], sampling [egg])
350. **Sechriest, R.E., D.E. Kuhlman, and J.L. Wedberg.** 1978. Aerial applications of insecticides to prevent rootworm egg-laying. Pages 103–108 in Thirtieth Illinois Custom Spray Operators Training School, summaries of presentations. Coop. Ext. Serv., University of Illinois at Urbana-Champaign. (*D. barberi*, *D. virgifera virgifera*, adult control, sampling [adult], root damage ratings)
351. **Shaw, J.T.** 1982. Cucurbitacins and rootworm adult sampling. Page 68 in Eighth Annual Illinois Crop Protection Workshop. Coop. Ext. Serv., University of Illinois at Urbana-Champaign. (*Diabrotica* spp., sampling [adult], cucurbitacin vial traps)
352. **Shaw, J.T., and D.E. Kuhlman.** 1978. Managing corn rootworm populations by crop rotation. Pages 98–100 in Thirtieth Illinois Custom Spray Operators Training School, summaries of presentations. Coop. Ext. Serv., University of Illinois at Urbana-Champaign. (*D. barberi*, *D. virgifera virgifera*, pest management, oviposition, root damage ratings, weeds, volunteer corn, sampling [adult], sampling [egg], soybeans)
353. **Shaw, J.T., J.H. Paullus, and W.H. Luckmann.** 1978. Corn rootworm oviposition in soybeans. J. Econ. Entomol. 71(2): 189–191. (*D. barberi* as *D. longicornis*, *D. virgifera virgifera*, crop rotation, volunteer corn, root damage ratings, weeds, sampling [adult], ovipositional preference)
354. **Shaw, J.T., W.G. Ruesink, S.P. Briggs, and W.H. Luckmann.** 1984. Monitoring populations of corn rootworm beetles (Coleoptera: Chrysomelidae) with a trap baited with cucurbitacins. J. Econ. Entomol. 77(6): 1495–1499. (*D. barberi*, *D. virgifera virgifera*, sampling [adult], sex ratio, cucurbitacin vial traps, *Cucurbita* spp., root damage ratings, oviposition, movement [adult], dispersal, sampling statistics)
- ♦355. **Shufuran, K.A.** 1986. The accuracy of integrated pest management scouting observations and factors affecting the performance of scouts employed in Kentucky. M.S. Thesis. University of Kentucky, Lexington. (*Diabrotica* spp., sampling [adult])
356. **Siegfried, B.D., and C.A. Mullin.** 1988. Properties of a cytochrome P-450-dependent epoxidase in aldrin-resistant western corn rootworms, *Diabrotica virgifera virgifera* LeConte. Pestic. Biochem. Physiol. 31(3): 261–268. (insecticide resistance, detoxification, metabolism)
357. **Smith, B.C.** 1979. Population changes of the northern corn rootworm (Coleoptera: Chrysomelidae) and corn yield losses in southwestern Ontario. Proc. Entomol. Soc. Ont. 1979, publ. 1980, 110: 85–91. (*D. barberi* as *D. longicornis*, sampling [egg], sampling [larval], carbofuran, phorate, population density)
358. **Smith B.C.** 1981. Effect of soil applications of carbofuran on egg populations of the northern corn rootworm (Coleoptera: Chrysomelidae). Can. Entomol. 113(1): 73. (*D. barberi* as *D. longicornis*, predation, fecundity, oviposition)
359. **Smith, M.V.** 1983. Northern (*Diabrotica longicornis*) and western (*Diabrotica virgifera*) corn rootworm beetles as competitors of foraging honey bees, *Apis mellifera*. Can. Beekeeping 10(8): 173–174. (*D. barberi* as *D. longicornis*, *D. virgifera virgifera*, goldenrod, competition, pollen)
360. **Solheim, B.A.** 1982. Toxicity and acetylcholinesterase inhibition by carbofuran and terbufos insecticides on *Diabrotica* species (Insecta: Coleoptera: Chrysomelidae). Ph.D. Thesis, Iowa State University, Ames. (*D. barberi*, *D. virgifera virgifera*, bioassay, cholinesterase activity)

361. **Sorensen, A.A.** 1988. Integrated pest management in the Midwest: issues, constraints, and solutions. Pages 59–73 in 1988 Fourteenth Annual Illinois Crop Protection Workshop proceedings. Coop. Ext. Serv., University of Illinois at Urbana-Champaign. (*D. barberi*, *D. virgifera virgifera*)
- ♦ 362. **Spackman, E.W., and C.C. Burkhardt.** 1979. Corn rootworm control. Bull. 688. Coop. Ext. Serv., University of Wyoming, Laramie. Unpaginated sheet. (*Diabrotica* spp., pest management)
- ♦ 363. **Spike, B.P.** 1986. Effect of corn rootworm injury, nitrogen and plant density on the growth and yield of corn. M.S. Thesis, Iowa State University, Ames. (*Diabrotica* spp., fertility, yield losses, damage [larval])
364. **Spike, B.P., and J.J. Tollefson.** 1988. Western corn rootworm (Coleoptera: Chrysomelidae) larval survival and damage potential to corn subjected to nitrogen and plant density treatments. J. Econ. Entomol. 81(5): 1450–1455. (*D. virgifera virgifera*, adult emergence, root damage ratings, root regrowth, soil moisture, lodging, artificial infestation, emergence traps, rainfall, fertilizer)
365. **Sridodo.** 1980. Selection of management variables for regressing Iowa corn yields on management, climatic, and soil variables. Ph.D. Thesis, Iowa State University, Ames. (*Diabrotica* spp., yield losses, environmental factors, tillage, crop rotation, fertilizer, weeds)
- ♦ 366. **Stamm, D.E.** 1981. Implementation of a western corn rootworm beetle scouting procedure to predict larval damage in Nebraska corn fields. M.S. Thesis, University of Nebraska, Lincoln. (*D. virgifera virgifera*, sampling [adult], prediction of damage, damage [larval])
367. **Stamm, D.E., Z.B. Mayo, J.B. Campbell, J.F. Witkowski, L.W. Andersen, and R. Kozub.** 1985. Western corn rootworm (Coleoptera: Chrysomelidae) beetle counts as a means of making larval control recommendations in Nebraska. J. Econ. Entomol. 78(4): 794–798. (*D. virgifera virgifera*, sampling [adult], root damage ratings, economic thresholds)
368. **Steffey, K.L.** 1979. Spatial dispersion and population estimation of northern and western corn rootworm adults in Iowa cornfields. Ph.D. Thesis, Iowa State University, Ames. (*D. barberi* as *D. longicornis*, *D. virgifera virgifera*, sampling [adult], distribution [spatial], dispersion [adult], mark and recapture, sampling statistics, population measurement)
369. **Steffey, K.L.** 1982. Sequential sampling for rootworm beetles. Pages 69–72 in Eighth Annual Illinois Crop Protection Workshop. Coop. Ext. Serv., University of Illinois at Urbana-Champaign. (*D. barberi* as *D. longicornis*, *D. virgifera virgifera*, sampling [adult], sampling [sequential])
370. **Steffey, K.L.** 1983. Corn rootworm biology, damage, and scouting procedures. Pages 131–142 in Ninth Annual Illinois Crop Protection Workshop. Coop. Ext. Serv., University of Illinois at Urbana-Champaign. (*D. barberi*, *D. virgifera virgifera*, tillage, soil moisture, root damage ratings, sampling [egg], sampling [larval], sampling [adult], damage [larval])
371. **Steffey, K., and K. Kinney.** 1988. Corn rootworm control: do root ratings tell the whole story? Pages 80–96 in Illinois Agricultural Pesticides Conf. '88, Fortieth Spray School, summaries of presentations. Coop. Ext. Serv., University of Illinois at Urbana-Champaign. (*Diabrotica* spp., root damage ratings, root regrowth, yield losses)
372. **Steffey, K.L., and D.E. Kuhlman.** 1985. Corn rootworm control: 1984 and beyond. Pages 152–165 in Thirty-seventh Illinois Custom Spray Operators Training School manual. Coop. Ext. Serv., University of Illinois at Urbana-Champaign. (*Diabrotica* spp., tillage, insecticide placement, pest management)
373. **Steffey, K.L., D.E. Kuhlman, and S.P. Briggs.** 1985. Should soil insecticides be used on first-year corn after soybeans? Pages 91–95 in Thirty-seventh Illinois Custom Spray Operators Training School manual. Coop. Ext. Serv., University of Illinois at Urbana-Champaign. (*Diabrotica* spp., first-year cornfields, crop rotation)
374. **Steffey, K.L., D.E. Kuhlman, and A.S. Felsot.** 1986. Corn rootworm management. Pages 108–116 in Thirty-eighth Illinois Custom Spray Operators Training manual. Coop. Ext. Serv., University of Illinois at Urbana-Champaign. (*D. barberi*, *D. virgifera virgifera*, pest management)
375. **Steffey, K., D. Kuhlman, and E. Levine.** 1986. Corn rootworm control data—1985. Pages 37–49 in Thirty-eighth Illinois Custom Spray Operators Training manual. Coop. Ext. Serv., University of Illinois at Urbana-Champaign. (*D. barberi*, *D. virgifera virgifera*, insecticide evaluation, insecticide placement, tillage, cultivation time applications, insecticide incorporation, pest management)
376. **Steffey, K.L., and J.J. Tollefson.** 1982. Spatial dispersion patterns of northern and western corn rootworm adults in Iowa cornfields. Environ. Entomol. 11(2): 283–286. (*D. barberi* as *D. longicornis*, *D. virgifera virgifera*, sampling [adult], sampling statistics, distribution [spatial], dispersion [adult])
377. **Steffey, K.L., J.J. Tollefson, and P.N. Hinz.** 1982. Sampling plan for population estimation of northern and western corn rootworm adults in Iowa cornfields. Environ. Entomol. 11(2): 287–291. (*D. barberi* as *D. longicornis*, *D. virgifera virgifera*, sampling [adult], sampling statistics, population measurement)
378. **Stinner, B.R., D.A. McCartney, and D.M. Van Doren, Jr.** 1988. Soil and foliage arthropod communities in conventional, reduced and no-tillage corn (maize, *Zea mays* L.) systems: a comparison after 20 years of continuous cropping. Soil and Tillage Res. 11(2): 147–158. (*D. barberi* as *D. longicornis*, *D. virgifera virgifera*, terribus, predation, root damage ratings)
379. **Storch, R.H., and J.L. Krysan.** 1980. Embryology of *Diabrotica undecimpunctata howardi* (Coleoptera: Chrysomelidae) from germ band formation to hatching. Ann. Entomol. Soc. Am. 73(4): 362–366. (*D. barberi*, *D. virgifera virgifera*, diapause, embryonic development)
380. **Strnad, S.P.** 1986. Host finding and feeding behavior of western corn rootworm larvae. Ph.D. Thesis, Purdue University, West Lafayette, IN. (*D. virgifera virgifera*, carbon dioxide, host location, olfaction, chemo-orientation, movement [larval], soil type, survival, starvation, infestation [larval])
381. **Strnad, S.P., and M.K. Bergman.** 1987. Distribution and orientation of western corn rootworm (Coleoptera: Chrysomelidae) larvae in corn roots. Environ. Entomol. 16(5): 1193–1198. (*D. virgifera virgifera*, infestation [larval], host location, distribution [spatial])

382. **Strnad, S.P., and M.K. Bergman.** 1987. Movement of first-instar western corn rootworms (Coleoptera: Chrysomelidae) in soil. *Environ. Entomol.* 16(4): 975-978. (*D. virgifera virgifera*, chemo-orientation, host location, movement [larval], carbon dioxide, soil type, survival, starvation)
383. **Strnad, S.P., M.K. Bergman, and W.C. Fulton.** 1986. First-instar western corn rootworm (Coleoptera: Chrysomelidae) response to carbon dioxide. *Environ. Entomol.* 15(4): 839-842. (*D. virgifera virgifera*, host location, olfaction, chemo-orientation)
384. **Sutherland, T.A.** 1986. Development of field and laboratory techniques to evaluate the relative attractiveness of corn plants to corn rootworm beetles (*Diabrotica virgifera virgifera* and *D. barberi*). Ph.D. Thesis, University of Nebraska, Lincoln. (attractants)
385. **Sutter, G.R.** 1982. Comparative toxicity of insecticides for corn rootworm (Coleoptera: Chrysomelidae) larvae in a soil bioassay. *J. Econ. Entomol.* 75(3): 489-491. (*D. virgifera virgifera*, terbufos, phorate, chlorpyrifos, fonofos, bufencarb, isofenphos, ethoprop, carbofuran)
386. **Sutter, G.R., and T.F. Branson.** 1980. A procedure for artificially infesting field plots with corn rootworm eggs. *J. Econ. Entomol.* 73(1): 135-137. (*D. barberi*, *D. virgifera virgifera*, artificial infestation, root damage ratings)
387. **Sutter, G.R., and T.F. Branson.** 1986. Artificial infestation of field plots. Pages 147-157 in J. L. Krysan and T. A. Miller, eds. *Methods for the study of pest Diabrotica*. Springer-Verlag, New York. (*D. virgifera virgifera*, rearing)
388. **Sutter, G.R., T.F. Branson, and J.R. Fisher.** 1981. The effect of varying populations of corn rootworm larvae on corn production. Pages 58-62 in *Thirty-third Illinois Custom Spray Operators Training School, summaries of presentations*. Coop. Ext. Serv., University of Illinois at Urbana-Champaign. (*D. virgifera virgifera*, yield losses, root damage ratings, artificial infestation, adult emergence, host plant resistance, tolerance, population density)
389. **Sutter, G.R., J.R. Fisher, and T.F. Branson.** 1981. Performance of insecticides in field plots artificially infested with western corn rootworm eggs. Pages 68-72 in *Thirty-third Illinois Custom Spray Operators Training School, summaries of presentations*. Coop. Ext. Serv., University of Illinois at Urbana-Champaign. (*D. virgifera virgifera*, artificial infestation, root damage ratings, yield losses, adult emergence, lodging, insecticide evaluation)
390. **Tauber, M.J., C.A. Tauber, and S. Masaki.** 1986. Variability and genetics of seasonal adaptations. Pages 192-217 in *Seasonal adaptations of insects*. Oxford University Press, New York. (*D. virgifera virgifera*, *D. virgifera zea*, diapause)
391. **Tollefson, J.J.** 1980. Suppressing adult corn rootworm populations to reduce larval damage. Pages 115-121 in *Thirty-second Illinois Custom Spray Operators Training School, summaries of presentations*. Coop. Ext. Serv., University of Illinois at Urbana-Champaign. (*Diabrotica* spp., adult control, oviposition, sampling [egg], sampling [adult], root damage ratings)
392. **Tollefson, J.J.** 1986. Field sampling of adult populations. Pages 123-146 in J. L. Krysan and T. A. Miller, eds. *Methods for the study of pest Diabrotica*. Springer-Verlag, New York. (*Diabrotica* spp., sampling [adult], sampling statistics, sampling [sequential], traps [adult], emergence traps)
393. **Tollefson, J.J.** 1986. Why soil insecticidal immunity. *Solutions Mag.* 30(1): 48-50. (*D. barberi*, *D. virgifera virgifera*, root damage ratings, carbofuran, bufencarb, phorate, ethoprop, insecticide degradation, biodegradation [enhanced], isofenphos)
394. **Tollefson, J.J.** 1988. A pest adapts to the cultural control of crop rotation. *Proc. British Crop Prot. Conf.* 3: 1029-1033. British Crop Protection Council, Brighton Center, England. (*D. barberi*, *D. virgifera virgifera*, extended diapause)
395. **Tollefson, J.J., J.F. Witkowski, J.C. Owens, and P.N. Hinz.** 1979. Influence of sampler variation on adult corn rootworm population estimates. *Environ. Entomol.* 8(2): 215-217. (*D. barberi* as *D. longicornis*, *D. virgifera virgifera*, sampling [adult], population measurement)
396. **Townsend, L.** 1984. Corn rootworm beetles. *Publ. ENT-45*. Coop. Ext. Serv., University of Kentucky, Lexington. 4 pp. (*D. barberi*, *D. virgifera virgifera*, biology, crop rotation, sampling [adult], pest management)
397. **Townsend, L., and M. Bitzer.** 1985. Silk clipping insects on corn. *Publ. ENT-48*. Coop. Ext. Serv., University of Kentucky, Lexington. 3 pp. (*D. barberi*, *D. virgifera virgifera*, silk feeding, pollination, insecticide recommendations, pest management)
398. **Turpin, F.T.** 1977. Insect insurance: potential management tool for corn insects. *Bull. Entomol. Soc. Am.* 23(3): 181-184. (*Diabrotica* spp., pest management, economics, yield losses, crop rotation)
399. **Turpin, F.T., and M.K. Bergman.** 1984. Rootworm control—past, present, and future. Pages 145-149 in *1984 Tenth Annual Illinois Crop Protection Workshop*. Coop. Ext. Serv., University of Illinois at Urbana-Champaign. (*D. barberi*, *D. virgifera virgifera*, pest management)
400. **Turpin, F.T., M.K. Bergman, D.R. Raemisch, and T.B. Johnson.** 1984. Rootworm biology: new developments. Pages 121-124 in *1984 Tenth Annual Illinois Crop Protection Workshop*. Coop. Ext. Serv., University of Illinois at Urbana-Champaign. (*Diabrotica* spp., pheromone, movement [adult], cucurbitacins)
401. **Turpin, F.T., and L.D. Godfrey.** 1982. Interfield movement of western corn rootworms. Pages 128-129 in *Thirty-fourth Illinois Custom Spray Operators Training School, summaries of presentations*. Coop. Ext. Serv., University of Illinois at Urbana-Champaign. (*D. virgifera virgifera*, movement [adult], economic thresholds, first-year cornfields, continuous cornfields)
402. **Turpin, F.T., D. Leva, and D. Freeman.** 1978. Impact of silk feeding by western corn rootworm beetles on corn grain yield. Pages 101-102 in *Thirtieth Illinois Custom Spray Operators Training School, summaries of presentations*. Coop. Ext. Serv., University of Illinois at Urbana-Champaign. (*D. virgifera virgifera*, pollination, yield losses)
403. **Turpin, F.T., and J.M. Thieme.** 1978. Impact of soil insecticide usage on corn production in Indiana: 1972-1974. *J. Econ. Entomol.* 71(1): 83-86. (*Diabrotica* spp., yield losses, economics, insecticide control)
404. **Turpin, F.T., and A.C. York.** 1981. Insect management and the pesticide syndrome. *Environ. Entomol.* 10(5): 567-572. (*Diabrotica* spp., pest management)

405. Tyler, B.M.J. 1979. The northern corn rootworm, *Diabrotica longicornis* (Say) (Coleoptera: Chrysomelidae), and other selected arthropods in three tillage systems in Ontario. Ph.D. Thesis, University of Guelph, Guelph, Ontario, Canada. (*D. barberi* as *D. longicornis*, no-till, adult emergence, temperature, soil moisture, oviposition, predation, ovipositional preference)
406. Tyler, B.M.J., and C.R. Ellis. 1979. Ground beetles in three tillage plots in Ontario and observations on their importance as predators of the northern corn rootworm, *Diabrotica longicornis* (Coleoptera: Chrysomelidae). Proc. Entomol. Soc. Ont. 110: 65-73. (*D. barberi* as *D. longicornis*, biological control, Carabidae, no-till, predation)
407. Ullstrup, A.J. 1978. Evolution and dynamics of corn diseases and insect problems since the advent of hybrid corn. Pages 283-297 in D. B. Walden, ed. Maize breeding and genetics. John Wiley and Sons, New York. (*D. barberi*, *D. virgifera virgifera*, host plant resistance, root regrowth, antibiosis)
408. University of Illinois at Urbana-Champaign, Coop. Ext. Serv. 1983. Corn rootworm. Fact sheet NHE-26 (rev.). 5 pp. (*D. barberi*, *D. virgifera virgifera*, biology, pollination, silk feeding, sampling [larval], sampling [adult], crop rotation)
409. VanWoerkom, G. 1977. Seasonal patterns of adult emergence and flight of the western corn rootworm, *Diabrotica virgifera*. Proc. Indiana Acad. Sci. 86: 230. (*D. virgifera virgifera*, emergence traps, sex ratio, phenology)
- ♦410. VanWoerkom, G.J. 1977. Vertical flight distribution and influence of environmental factors on locomotor behavior of adult western corn rootworms. M.S. Thesis, Purdue University, West Lafayette, IN. (*D. virgifera virgifera*, temperature, wind, movement [adult], sampling [adult])
411. VanWoerkom, G.J., F.T. Turpin, and J.R. Barrett, Jr. 1980. Influence of constant and changing temperatures on locomotor activity of adult western corn rootworms (*Diabrotica virgifera*) in the laboratory. Environ. Entomol. 9(1): 32-34. (*D. virgifera virgifera*, environmental factors, sampling [adult], movement [adult])
412. VanWoerkom, G.J., F.T. Turpin, and J.R. Barrett, Jr. 1983. Wind effect on western corn rootworm (Coleoptera: Chrysomelidae) flight behavior. Environ. Entomol. 12(1): 196-200. (*D. virgifera virgifera*, sampling [adult], environmental factors, movement [adult])
- ♦413. Vassalotti, P.M. 1982. Bionomics of the northern corn rootworm on first year corn in eastern South Dakota. M.S. Thesis, South Dakota State University, Brookings. (*D. barberi*, biology, first-year cornfields)
414. Walgenbach, D.D., and G.R. Sutter. 1977. Corn rootworm susceptibility to insecticides. Pages 68-71 in Twenty-ninth Illinois Custom Spray Operators Training School, summaries of presentations. Coop. Ext. Serv., University of Illinois at Urbana-Champaign. (*D. virgifera virgifera*, insecticide resistance, toxicity, bioassay)
415. Warnes, D.D., and J.A. Lofgren. 1981. A search for herbicide effects on rootworm control in corn with carbamate and phosphate insecticides. Proc. N. Cent. Weed Control Conf., 36: 128. (*D. barberi* as *D. longicornis*, *D. virgifera virgifera*)
416. Wedberg, J.L. 1985. Corn rootworms. Extension publ. A3328 (UWEXnote 4). Coop. Ext. Serv., University of Wisconsin, Madison. 4 pp. (*D. barberi*, *D. virgifera virgifera*, biology, sampling [adult], crop rotation)
417. Weiss, M.J. 1983. Influence of corn plant density on population estimates and population dynamics of western (*Diabrotica virgifera virgifera* LeConte) and northern (*D. longicornis barberi* Smith and Lawrence) corn rootworm (Coleoptera: Chrysomelidae). Ph.D. Thesis, University of Nebraska, Lincoln. (phenology, sex ratio, oviposition, sampling [adult], heat units, model, population measurement)
418. Weiss, M.J., and G. Jensen. 1985. The western corn rootworm. Publ. 8409 (rev.). Coop. Ext. Serv., Montana State University, Bozeman. 2 pp. (*D. virgifera virgifera*, biology, sampling [adult], pest management, insecticide recommendations)
419. Weiss, M.J., and Z B Mayo. 1983. Potential of corn rootworm (Coleoptera: Chrysomelidae) larval counts to estimate larval populations to make control decisions. J. Econ. Entomol. 76(1): 158-161. (*D. barberi*, *D. virgifera virgifera*, heat units, sampling [larval], population measurement)
420. Weiss, M.J., and Z B Mayo. 1985. Influence of corn plant density on corn rootworm (Coleoptera: Chrysomelidae) population estimates. Environ. Entomol. 14(6): 701-704. (*D. barberi*, *D. virgifera virgifera*, phenology, oviposition, sampling [adult], population measurement)
421. Weiss, M.J., Z B Mayo, and J.P. Newton. 1983. Influence of irrigation practices on the spatial distribution of corn rootworm (Coleoptera: Chrysomelidae) eggs in the soil. Environ. Entomol. 12(4): 1293-1295. (*D. barberi*, *D. virgifera virgifera*, distribution [spatial], sampling [egg], soil moisture, oviposition, rainfall, ovipositional preference)
422. Weiss, M.J., K.P. Seevers, and Z B Mayo. 1985. Influence of western corn rootworm larval densities and damage on corn rootworm survival, developmental time, size and sex ratio (Coleoptera: Chrysomelidae). J. Kansas Entomol. Soc. 58(3): 397-402. (*D. virgifera virgifera*, damage [larval], density [larval], size [adult])
423. White, R. 1977. Sexual characters of species of *Diabrotica* (Chrysomelidae: Coleoptera). Ann. Entomol. Soc. Am. 70(2): 168. (*D. barberi*, *D. virgifera virgifera*, sex differences, sex determination)
424. Wilde, G. 1978. Corn rootworm control in Kansas. Agric. Exp. Sta. Bull. 616. Kansas State University, Manhattan. 15 pp. (*D. barberi*, *D. virgifera virgifera*, biology, root damage ratings, crop rotation, insecticide recommendations, pest management, adult control)
425. Wilkin, P.J., J.L. Krysan, and P.L. Guss. 1986. Electroantennogram responses of corn rootworms and other closely related *Diabrotica* (Coleoptera: Chrysomelidae) to sex pheromones and synthetic attractants. J. Kansas Entomol. Soc. 59(4): 571-579. (*D. barberi* as *D. longicornis*, *D. virgifera virgifera*)
426. Willson, H.R., and B. Easley. 1987. Insect pests of field crops. Bull. 545. Coop. Ext. Serv., Ohio State University, Columbus. 21 pp. (*D. barberi*, *D. virgifera virgifera*, biology, crop rotation, insecticide recommendations, pest management)

427. **Wisniewski, M.N.** 1984. The western corn rootworm synthetic sex pheromone: optimal concentration and source, area of influence, and male response disruption. Ph.D. Thesis, Iowa State University, Ames. (*D. virgifera virgifera*, pheromone traps, mark and recapture, disruption of mating)
428. **Witkowski, J.F., D.L. Keith, and Z B Mayo.** 1982. Evaluating corn rootworm soil insecticide performance. NebGuide G82-597. Coop. Ext. Serv., University of Nebraska, Lincoln. 2 pp. (*Diabrotica* spp., insecticide evaluation, root damage ratings)
429. **Witkowski, J.F., D.L. Keith, and Z B Mayo.** 1986. Western corn rootworm soil insecticide treatment decisions based on beetle numbers. NebGuide G86-774. Coop. Ext. Serv., University of Nebraska, Lincoln. 4 pp. (*D. virgifera virgifera*, pest management, sampling [adult], economic thresholds)
430. **Witkowski, J.F., and J.C. Owens.** 1979. Corn rootworm behavior in response to trap corn. Iowa State J. Res. 53(4): 317–324. (*D. barberi* as *D. longicornis*, *D. virgifera virgifera*, movement [adult], trap crop, oviposition, adult emergence)
431. **Yaro, N.D., J.L. Krysan, and T.L. Ladd, Jr.** 1987. *Diabrotica cristata* (Coleoptera: Chrysomelidae): attraction to eugenol and related compounds compared with *D. barberi* and *D. virgifera virgifera*. Environ. Entomol. 16(1): 126–128. (attractants, evolution, kairomones, semiochemicals)
432. **Yule, W.N., C.R. Dominique, and M.A. Matin.** 1983. Bionomics and sampling methods for the northern corn rootworm in Quebec. Proc. of the Tenth Intl. Congress of Plant Protection, 20–25 November 1983, Brighton, England: Plant Protection for Human Welfare. Vol. 1: 165. (*D. barberi* as *D. longicornis*, threshold temperature, heat units, oviposition, sampling [egg], sampling [larval], sampling [adult], biology)
433. **Zacharias, T.P.** 1984. Economically optimal integrated pest management strategies for control of corn rootworm and soybean cyst nematode. Ph.D. Thesis, University of Illinois at Urbana-Champaign. (*D. barberi* as *D. longicornis*, *D. virgifera virgifera*, economics, model, risk management)
434. **Zacharias, T.P., and A.H. Grube.** 1986. Integrated pest management strategies for approximately optimal control of corn rootworm and soybean cyst nematode. Am. J. Agric. Econ. 68(3): 704–715. (*D. barberi* as *D. longicornis*, *D. virgifera virgifera*, crop rotation, model, economics)

Periodicals Cited

- Agric. Canada** — Agriculture Canada
- Agric. Exp. Sta.** — Agricultural Experiment Station (various states)
- Agron. J.** — Agronomy Journal
- Am. J. Agric. Econ.** — American Journal of Agricultural Economics
- Ann. Entomol. Soc. Am.** — Annals of the Entomological Society of America
- Ann. Rev. Entomol.** — Annual Review of Entomology
- Bull. Ecol. Soc. Am.** — Bulletin of the Ecological Society of America
- Bull. Entomol. Soc. Am.** — Bulletin of the Entomological Society of America
- Bull. Entomol. Soc. Can.** — Bulletin of the Entomological Society of Canada
- Bull. Environ. Contam. Toxicol.** — Bulletin of Environmental Contamination and Toxicology
- Can. Beekeeping** — Canadian Beekeeping
- Can. Entomol.** — Canadian Entomologist
- Can. J. Zool.** — Canadian Journal of Zoology
- Coleopterists Bull.** — Coleopterists Bulletin
- Coop. Ext. Serv.** — Cooperative Extension Service (various universities)
- CRC Critical Reviews in Plant Sci.** — CRC Critical Reviews in Plant Sciences
- Crop Sci.** — Crop Science
- Crops Soils Mag.** — Crops and Soils Magazine
- Cucurbit Genetics Coop.** — Cucurbit Genetics Cooperative
- Ecology**
- Ent. Exp. Appl.** — Entomologia Experimentalis et Applicata
- Entomography**
- Entomologist's Monthly Mag.** — Entomologist's Monthly Magazine
- Entomophaga**
- Environ. Entomol.** — Environmental Entomology
- Great Lakes Entomol.** — Great Lakes Entomologist
- Ill. Nat. Hist. Sur. Biol. Notes** — Illinois Natural History Survey Biological Notes
- Ill. Nat. Hist. Sur. Reports** — Illinois Natural History Survey Reports
- Insect Biochem.** — Insect Biochemistry
- Iowa State J. Res.** — Iowa State Journal of Research
- IRCS Medical Sci.** — International Research Communications System Medical Science
- J. Agric. Entomol.** — Journal of Agricultural Entomology
- J. Agric. Food Chem.** — Journal of Agricultural and Food Chemistry
- J. Amer. Soc. Hort. Sci.** — Journal of the American Society for Horticultural Science
- J. Applied Ecol.** — Journal of Applied Ecology
- J. Chem. Ecol.** — Journal of Chemical Ecology
- J. Econ. Entomol.** — Journal of Economic Entomology
- J. Environ. Sci. Health B** — Journal of Environmental Science and Health, Part B
- J. Georgia Entomol. Soc.** — Journal of the Georgia Entomological Society (now Journal of Entomological Science)
- J. Heredity** — Journal of Heredity
- J. Insect Physiol.** — Journal of Insect Physiology
- J. Invert. Pathol.** — Journal of Invertebrate Pathology
- J. Kansas Entomol. Soc.** — Journal of the Kansas Entomological Society
- J. Pestic. Sci.** — Journal of Pesticide Science
- J. Wildlife Management** — Journal of Wildlife Management
- Maydica**
- Misc. Publ., Entomol. Soc. Am.** — Miscellaneous Publications, Entomological Society of America
- N. Cent. Reg. Res. Publ.** — North Central Regional Research Publication
- Ohio Report**
- Pedobiologia**
- Pestic. Biochem. Physiol.** — Pesticide Biochemistry and Physiology
- Physiol. Entomol.** — Physiological Entomology
- Phytopathology**
- Plant Dis.** — Plant Disease
- Plant Prot. Bull., F.A.O.** — Plant Protection Bulletin, Food and Agricultural Organization
- Proc. British Crop Prot. Conf.** — Proceedings of the British Crop Protection Conference
- Proc. Entomol. Soc. Ont.** — Proceedings of the Entomological Society of Ontario
- Proc. Indiana Acad. Sci.** — Proceedings of the Indiana Academy of Science

Proc. Intl. Maize Virus Dis. Colloq. and Workshop — Proceedings of the International Maize Virus and Disease Colloquium and Workshop

Proc. Intl. Symp. — Proceedings of the International Symposium.

Proc. National Acad. Sci. — Proceedings of the National Academy of Sciences USA

Proc. N. Cent. Br. Entomol. Soc. Am. — Proceedings of the North Central Branch of the Entomological Society of America

Proc. N. Cent. Weed Control Conf. — Proceedings of the North Central Weed Control Conference

Proc. N. D. Acad. Sci. — Proceedings of the North Dakota Academy of Sciences

Prot. Ecol. — Protection Ecology

Revue d'Entomologie du Québec

Search Agric. — Search Agriculture

Soil and Tillage Res. — Soil and Tillage Research

Solutions Mag. — Solutions Magazine

Tetrahedron

Index

Adult control 19 28 87 191 192 195 218 247 257 268
287 301 303 329 345 349 350 391 424

Adult emergence 27 40 56 95 109 123 125 146 150 153
179 180 185 186 264 284 313 314 316 346 364 388 389
405 409 430

Age 65 317

***Agelaius phoeniceus* (red-winged blackbird)** 34 35 36

Aggregation pheromone 338 339

Alachlor 151

Aldrin 21 76 77 78 107 241 356

Allozymes 293

Alternate food source 209

Alternate host 209

Ammonia (*see also* fertilizer) 68

Antibiosis 39 48 90 407

Antixenosis 39 48 58

Ants 222

***Apis mellifera* (honey bee)** 359

Arrestants, feeding 41 301 302

Artificial infestation 39 52 53 55 56 75 125 126 131
237 327 364 386 387 388 389

Attractants 7 8 24 64 190 248 249 250 251 252 253
254 255 256 257 258 267 292 295 299 306 337 338 384
425 431

Baits 119 120 257 268 300 301 303

Beauveria bassiana 88

Bees 359

Behavior (insect) (*see also* ovipositional preference) 1 39 48
91 93 116 190 193 221 240 269 272 338 380 410 412
430

Bibliography 199

Bioassay 11 71 77 81 83 168 172 173 216 237 241
274 287 337 360 385 414

Biodegradation, enhanced (*see also* insecticide degradation;
insecticide persistence) 81 107 110 114 148 170 171 320
393

Biological control 23 63 67 88 121 122 201 210 222
224 318 335 336 406

Biology 2 25 50 62 66 95 98 181 192 195 210 218
228 231 233 237 238 247 311 315 328 329 338 345 370
396 400 408 413 416 418 424 426 432

Bionomics (*see* biology)

Biosystematics (*see* taxonomy)

Biotic potential 185

Bufencarb 170 289 385 393

Carabidae, carabids 224 406

Carbaryl 22 77 107 170 287

Carbofuran 4 5 22 71 76 77 78 103 107 110 112
114 148 149 151 170 171 172 173 274 285 286 289 320
357 358 360 385 393

Carbon dioxide 190 380 382 383

***Celatoria* spp.** 122

Chemo-orientation 380 382 383

Chlorpyrifos 77 107 168 171 172 173 274 385

Cholinesterase activity 65 360

Cold hardiness (of eggs) 163 165 166 326

Coloration, insect 240

Compensatory root growth (*see also* root regrowth) 211 212

Competition 16 330 333 359

Conservation tillage (*see also* no-till) 107 108 109 111 312

Continuous cornfields 144 145 177 179 183 193 291 401

Control (*see* adult control; biological control; insecticide
control; insecticide evaluation; pest management)

Cover crops 244

Crop insurance 261 263

Crop rotation 13 62 66 67 87 94 145 146 147 181
191 193 234 236 244 245 247 261 262 263 266 310 311
321 328 329 334 345 352 353 365 373 394 396 398 408
416 424 426 434

***Cucurbita* spp.** 7 8 9 11 16 41 115 116 118 130
207 296 297 299 301 303 304 338 339 340 354

Cucurbitaceae, cucurbits (*see also* *Cucurbita* spp.) 16 17 18
116 210 296 297 299 300 301 303 304 338

Cucurbitacin vial traps 59 60 119 120 157 257 351 354

Cucurbitacins (*see also* cucurbitacin vial traps) 7 9 10
11 41 115 116 117 118 207 210 268 296 297 298 301
302 303 304 339 340 351 354 400

Cultivation time applications 27 28 31 289 375

Cyanazine 151

Cypermethrin 83

Damage by birds 34

Damage, larval (*see also* root damage ratings) 14 134 363 366
370 422

DDT 77 107

Defense compounds 117 253

Degradation (*see insecticide degradation*)**Density, larval** 331 422**Detoxification** 10 78 118 356**Development** (*see also embryonic development; ovarian development; reproductive development*) 40 97 127 128 204**Developmental time** 50 122 422

***Diabrotica* spp.** 3 4 5 9 15 25 28 31 41 49
 59 63 88 104 108 109 123 130 156 170 174 175 178
 129 130 134 137 138 140 146 147 150 152 153 154 155
 197 200 214 244 254 278 290 300 305 320 331 344 351
 355 362 363 365 371 372 373 391 392 398 400 403 404
 428

D. barberi 1 9 11 13 26 27 29 30 32 33 39
 48 49 60 61 64 66 87 90 91 92 93 94 100
 101 102 105 106 107 110 111 112 113 119 120 124 126
 127 228 229 231 233 234 235 236 237 238 239 240 241
 159 160 163 164 176 177 179 180 181 182 183 184 186
 187 188 191 192 199 206 208 209 210 213 217 219 220
 227 228 229 231 233 234 235 236 237 238 239 240 241
 242 245 246 247 248 249 250 251 252 253 255 256 257
 258 259 264 265 266 267 274 279 281 282 283 284 288
 291 292 293 294 295 297 298 299 309 310 311 314 315
 316 317 318 321 329 332 333 334 337 342 343 345 347
 348 349 350 352 354 360 361 370 374 375 379 384 386
 393 394 396 397 399 407 408 413 416 417 419 420 421
 423 424 426 431

D. barberi* as *D. longicornis 2 12 14 19 24 34 35 36
 67 71 72 76 77 78 79 80 83 95 96 97 98
 99 114 122 133 135 136 139 148 149 168 169 172 173
 193 194 198 205 221 222 223 224 243 261 262 263 280
 285 286 287 289 296 302 303 307 312 313 319 346 353
 357 358 359 368 369 376 377 378 395 405 406 415 425
 430 432 433 434

D. longicornis (*see D. barberi* as *D. longicornis*)

D. virgifera virgifera 1 6 7 8 9 11 13 14 16
 17 18 19 20 21 22 23 24 26 27 29 30 32
 33 37 38 39 40 41 42 43 44 45 46 47 48
 49 50 51 52 53 54 55 56 57 58 60 61 62
 65 66 67 68 69 70 72 73 74 75 76 77 78
 79 80 81 82 84 85 86 87 90 91 92 93 94
 100 101 102 103 105 106 107 110 111 112 113 114 115
 116 117 118 119 120 121 124 125 126 127 128 129 130
 131 132 133 134 137 138 139 140 141 142 143 144 145
 146 147 148 149 150 151 152 153 154 155 157 158 160
 161 162 163 165 166 167 171 179 180 181 182 183 184
 185 186 187 188 189 190 191 192 193 194 195 196 199
 200 201 202 203 204 205 206 207 208 209 210 211 212
 213 216 217 218 219 220 221 222 223 224 225 226 227
 228 229 230 231 232 233 234 235 236 237 238 239 242
 243 245 246 247 249 250 251 253 254 255 256 257 258
 260 261 262 263 264 265 266 267 268 269 270 271 272
 273 274 275 276 277 279 285 286 287 288 289 291 292
 296 297 298 299 301 302 303 304 306 308 309 310 311
 312 313 315 319 322 323 324 325 326 327 328 329 332
 333 337 338 339 340 341 342 343 345 346 347 349 350
 352 353 354 356 359 360 361 364 366 367 368 369 370
 374 375 376 377 378 379 380 381 382 383 384 385 386
 387 388 389 390 393 394 395 396 397 399 401 402 407

408 409 410 411 412 414 415 416 417 418 419 420 421
 422 423 424 425 426 427 429 430 431 433 434

D. virgifera zea 11 45 48 49 50 57 158 160 227 228
 231 233 237 238 239 347 390

Diapause (*see also extended diapause; non-diapause strain*) 45
 50 68 225 226 227 230 232 236 324 325 379 390

Diazinon 77 286 287 289**Dieldrin** 21 172**Diet, adult** 46 115 131 200 314**Disease transmission** 140 205 206

Dispersal 28 61 82 145 185 186 193 231 250 251 314
 316 354

Dispersion, adult 281 368 376**Dispersion, larval** 30**Disruption of mating** 258 427

Distribution, geographical 62 80 132 193 226 228 231 233
 237 238 239 240

Distribution, spatial 133 135 136 280 281 331 333 342 368
 376 381 421

Disulfoton 171

Economic thresholds 14 27 66 73 75 86 134 137 138
 145 243 261 263 291 367 401 429

Economics 14 62 261 262 263 279 334 398 403 433 434**Egg chorion** 229

Egg hatch 40 50 165 196 225 230 232 235 236 265 266
 324

Egg hatch duration 40 52 236 265 326**Egg production** (*see fecundity*)**Egg separation (extraction)** 283 307 342**Egg viability** 235 324 326**Electroantennogram** 7 8 11 89 425**Electrophoresis** 115 167 210 330 332 347**Embryonic development** 226 324 325 326 379**Emergence** (*see adult emergence*)

Emergence traps 59 123 125 179 180 284 313 316 346 364
 392 409

Environmental chemodynamics 107

Environmental factors (*see also humidity; rainfall; tempera-
 ture; weather*) 13 24 50 71 86 92 95 106 113 127
 128 157 201 208 217 282 365 410 411 412

Esterase 167 293 341 347**Estragole** 253 255 256 257 299**Ethoprop** 274 285 289 385 393

Eugenol 248 249 250 251 252 253 254 255 256 257 258
 295 299 301 431

Evolution 48 227 231 232 296 297 302 407 431

Extended diapause (*see also diapause*) 13 87 94 100 181
 234 236 241 246 247 265 266 321 394

- Fecundity** 22 51 179 235 259 315 338 358
- Feeding behavior** 39 48 116 380
- Feeding stimulant** 7 210
- Fensulfothion** 77 285 286 289
- Fenvalerate** 15 83
- Fertility (insect)** 235
- Fertilizer** 68 134 285 363 364 365
- First-year cornfields** 144 145 147 176 177 291 373 401 413
- Flight** 82 84 85 179 314 409 410 412
- Flotation, egg** (*see also* egg separation) 307 342
- Flotation, larval** 29 124 129 307
- Fonofos** 3 76 77 78 171 274 285 286 289 385
- Food quality** 259 315
- Foxtail** 147 208 209
- Fungi, fungus** 88 141 142 143 318
- Fusarium* spp.** 141 142 143
- Genetics** 90 210 226 230 261 293 294 322 323 347 390
- Goldenrod** 359
- Hatch** (*see* egg hatch)
- Heat units** 27 33 95 97 128 183 204 346 417 419 432
- Heptachlor** 77
- Herbicide** 151 415
- Host finding** (*see* host location)
- Host location** 38 39 380 381 382 383
- Host plant resistance** 14 48 49 56 57 58 73 74 90 198 212 237 322 323 388 407
- Host plant selection** 9 39 64 115 231 238 255 256 296 297 299 302 303 304 306 337
- Humidity** 127 225 232
- Hybridization** 174 194 235
- Identification, adult** 228
- Identification, egg** 228
- Identification, larval** 228 330 332
- Indole** 7 8 11 253 254 255 256 299 306
- Infestation, larval** 380 381
- Insecticide application equipment** 101 102 288
- Insecticide bioactivity** 81 106
- Insecticide calibration** 101
- Insecticide control** (*see also* adult control; insecticide evaluation; pest management) 80 101 102 403
- Insecticide degradation** (*see also* insecticide persistence) 3 4 5 71 78 81 104 105 107 108 110 112 113 114 148 149 170 171 175 320 393
- Insecticide evaluation** 237 285 286 287 288 289 375 389 428
- Insecticide history** 4 19 110 114 149 170 171
- Insecticide incorporation** 375
- Insecticide persistence** (*see also* biodegradation, enhanced; insecticide degradation) 3 4 5 71 103 104 105 106 107 109 110 112 113 114 148 149 168 169 170 171 172
- Insecticide placement** 285 372 375
- Insecticide recommendations** 19 87 94 100 181 188 191 195 213 218 247 309 310 311 328 345 397 418 424 426
- Insecticide residues** 21 71 149 287
- Insecticide resistance** 19 76 77 78 81 103 106 112 241 261 262 274 298 356 414
- Insecticide volatility** 71 169 172
- Insecticides, sublethal dosage** 22
- Insemination** 235
- Integrated pest management** (*see* pest management)
- Intercropping** 70
- Irrigation** 50 69 287 421
- Isoeugenol** 248 253 255 256 295 299
- Isofenphos** 105 170 171 274 385 393
- Juvenile hormone** 82 84
- Kairomones** (*see also* semiochemicals) 7 11 122 190 237 253 254 255 256 257 258 267 292 295 296 297 299 302 303 304 338 340 431
- Larval control** (*see* biological control; insecticide evaluation; pest management)
- Lasius neoniger*** 222
- Limonene** 215 216
- Lodging** 39 288 323 364 389
- Longevity (insect)** 22 51 115 259 278 315 338
- Loss in yield** (*see* yield losses)
- Malathion** 77 287
- Mark and recapture** 12 368 427
- Mating** (*see also* mating status) 194 235 239 240 258 269 272 273 427
- Mating status** 24 43 82 84 273
- Metabolism** 10 118 210 356
- Methoxychlor** 77
- Mineral nutrition** 211
- Mixed function oxidase** 6
- Model, modeling** 73 134 179 185 261 314 317 417 433 434
- Morphology** 240 271
- Mortality** 95 185 186 268 278 282 312 314 316
- Movement, adult** 18 28 61 79 80 82 84 85 145 185 186 193 231 250 251 261 314 354 400 401 410 411 412 430
- Movement, larval** 38 39 278 330 380 382
- Nematodes** 189 201 210 335 336 433 434
- Neoapectana carpocapsae*** 210 335 336

Nitrogen 363 364

No-till (*see also* conservation tillage) 109 111 150 152 153
154 155 208 264 310 405 406

Non-diapause strain 55

Nutrient uptake 211

Olfaction 38 41 64 380 383

Ovarian development (*see also* reproductive development;
reproductive status) 82 84 145

Ovicides 216

Oviposition 22 43 48 50 51 79 95 96 99 133 135
136 146 147 150 154 161 164 176 177 179 183 186 187
193 208 219 220 221 223 244 245 259 264 268 312 314
315 316 317 349 352 353 354 358 391 405 417 420 421
430 432

Oviposition behavior (*see* ovipositional preference)

Ovipositional preference 48 95 150 154 161 164 221 353
405 421

Parasites 121 122 201

Pathogens, insect 50 88 200 202 318

Pathogens, plant 140 141 142 143 205 319

Pennac-M 287

Periodicity 24 82 85 92 214

Permethrin 15 77 83

Pest management 19 28 48 59 66 67 72 74 87
94 100 138 181 188 191 192 195 197 198 213 218 224
247 298 310 311 328 329 334 344 345 352 355 361 362
372 374 375 396 397 398 399 404 418 424 426 429 433
434

Pesticide interactions 107

Phenology 7 8 27 33 73 95 98 146 253 281 314
316 409 417 420

Pheromone (*see also* aggregation pheromone) 11 24 45 61
91 92 93 158 159 160 207 237 239 240 242 249 257
258 269 270 271 275 276 292 308 400 425 427

Pheromone traps 24 275 276 427

Phorate 3 71 76 77 78 107 168 169 170 171 274 286
289 357 385 393

Phosmet 287

Plant density 134 363 364 417 420

Plant stress 53 73

Plant volatiles 7 8 9 11 64 190 253 255 257 298
299

Planting date 27 32 134 209 286 313 314 316

Pollen 9 315 359

Pollination 86 397 402 408

Population density 51 61 75 79 145 161 186 193 357
388

Population dynamics 17 18 27 32 50 144 145 150 152
154 179 185 186 209 282 314 316 417

Population estimates (*see* population measurement)

Population measurement (*see also* sampling) 60 178 180 182
281 368 377 395 417 419 420

Predation 23 35 36 50 63 222 224 358 378 405 406

Prediction of damage 178 184 366

Protandry 40

Pyrethroids 15 83 173

Rainfall 50 75 134 157 287 364 421

Rearing 37 40 46 47 95 96 122 131 200 202 203
237 259 274 314 315 387

Reduced tillage (*see* conservation tillage; no-till)

Repellents 257 258

Reproductive behavior 93 269 272

Reproductive development (*see also* reproductive status) 314
315 317

Reproductive isolation 235

Reproductive status 43 82 84 145 273 314 315 317

Risk management 261 263 433

Root damage ratings 14 39 51 52 53 54 56 57 58
60 62 74 103 109 112 126 134 137 145 149 150 151
152 184 208 211 212 245 246 266 285 286 288 289 305
313 321 327 349 350 352 353 354 364 367 370 371 378
386 388 389 391 393 424 428

Root regrowth (*see also* compensatory root growth) 90 211
364 371 407

Root size 54

Sampling, adult 60 66 100 123 134 137 138 139 147 178
180 181 182 184 185 186 188 191 192 213 214 217 237
247 261 276 280 281 284 290 291 292 348 349 350 351
352 353 354 355 366 367 368 369 370 376 377 391 392
395 396 408 410 411 412 416 417 418 420 429 432

Sampling, egg 14 133 134 135 136 147 150 154 179 185
186 187 237 280 283 342 349 352 357 370 391 421 432

Sampling, larval 26 27 29 30 100 129 237 280 331 334
357 370 408 419 432

Sampling, pupal 129 334

Sampling, sequential 134 139 181 291 348 369 392

Sampling statistics 129 134 179 280 342 348 354 368 376
377 392

Seed corn 86

Semiochemicals (*see also* kairomones) 7 89 190 207 248
249 250 251 252 253 254 255 256 257 258 267 292 295
296 297 298 299 303 304 338 340 431

Sequestration 117 118

Setaria spp. (foxtail) 147 208 209

Sex determination 228 423

Sex differences 65 228 423

Sex pheromone (*see* pheromone)

- Sex ratio** 7 8 40 119 120 130 145 179 182 186 253
314 316 354 409 417 422
- Silk feeding** 69 86 181 243 397 402 408
- Size, adult** 51 422
- Soil moisture** 50 71 95 99 128 148 161 168 169 171
172 173 201 208 221 225 232 277 364 370 405 421
- Soil temperature** (*see also* temperature) 162 260
- Soil type** 71 95 99 113 168 169 172 173 277 380 382
- Soil washing apparatus** 29 124
- Soybeans** 146 147 245 246 261 262 263 266 352 353 373
433 434
- Spatial distribution** (*see* distribution, spatial)
- Spermatophore** 269 273
- Stalk rot** 141 142 143
- Starvation** 380 382
- Steinernema feltiae*** 201
- Sterile insect release** 42 44
- Stress** (*see* plant stress)
- Survival (insect)** 74 127 128 150 155 162 163 165 166 179
186 208 260 277 315 326 333 364 380 382 422
- Synthetic pyrethroids** (*see* pyrethroids)
- Systematics** (*see* taxonomy)
- Taxonomy** 174 210 226 228 229 237 238 239 240 330 332
347
- Temperature** (*see also* soil temperature; threshold tempera-
ture) 13 24 37 61 71 92 95 127 157 163 165 166
168 169 172 173 183 201 204 217 227 232 314 315 317
326 346 405 410 411
- Terbufos** 3 71 76 77 78 107 108 113 168 171 172
274 286 289 360 378 385
- Threshold temperature** 95 97 122 204 432
- Tillage** (*see also* conservation tillage; no-till) 150 152 153 154
155 208 260 264 282 310 365 370 372 375 378 405 406
- Tolerance (host plant)** 39 48 56 74 90 212 388
- Topography** 193
- Toxicity** 71 83 106 173 274 360 385 414
- Trap capture** 24 217
- Trap color** 20 217 252 292
- Trap crop** 430
- Trap height** 24 252
- Traps, adult** (*see also* cucurbitacin vial traps; emergence traps;
pheromone traps) 20 59 182 184 214 217 248 249 250
251 252 254 255 256 257 258 267 292 392
- Vector, disease** 140 141 142 143 205 206 237 319
- Virus transmission** 140 205 206 319
- Vision** 1 20
- Visual counts** (*see also* sampling, adult) 60 157
- Voltinism** 45 48 227
- Volunteer corn** 146 147 352 353
- Weather** (*see also* environmental factors) 157 186
- Weed competition** 209
- Weeds** 50 146 164 208 209 352 353 365
- Wind** 24 410 412
- Yield losses** 14 53 56 58 69 75 111 134 138 156 211
288 298 305 323 334 349 357 363 365 371 388 389 398
402 403
- Zea diploperennis*** 49

Illinois Natural History Survey
Natural Resources Building
607 East Peabody Drive
Champaign, Illinois 61820

A Division of the Illinois Department of Energy and Natural Resources