Chama Anthon T. 1952

UNITED STATES DEPARTMENT OF AGRICULTURE

MISCELLANEOUS PUBLICATION No. 74

WASHINGTON, D. C.

MAY, 1930

AN ANNOTATED LIST OF THE IMPORTANT NORTH AMERICAN FOREST INSECTS

Compiled by F. C. CRAIGHEAD, *Principal Entomologist in Charge*
and WILLIAM MIDDLETON, *Associate Entomologist*
Division of Forest Insects, Bureau of Entomology

CONTENTS

	Page		Page
Government projects relating to forest insects.....	1	The most important forest insects—Contd.....	
Tree-killing bark beetles.....	1	Insects affecting forest products.....	11
Insects injurious to forest products.....	2	Insects affecting seeds, forest reproduc-	
Insects affecting forest management.....	3	tion, and nursery plants.....	15
Insects injurious to shade trees and		Defoliating insects.....	18
ornamental shrubs.....	3	Literature cited.....	23
The most important forest insects.....	4	Appendix.....	27
Tree-killing bark beetles.....	4	Index.....	28
Insects boring in living trees.....	6		

GOVERNMENT PROJECTS RELATING TO FOREST INSECTS

The investigations by the United States Government of the insects affecting forest trees, forest products, and shade trees are centered in the division of forest insects of the Bureau of Entomology. The investigations carried on by the division are grouped for administrative convenience into projects, the most important of which are briefly described in the following paragraphs.

TREE-KILLING BARK BEETLES

Several species of tree-killing bark beetles of the genus *Dendroctonus* destroy annually over 6,000,000,000 feet of timber valued at from \$15,000,000 to \$20,000,000. Biological investigations of these beetles and the associated cooperation with Federal and private agencies in their control constitute one of the major activities of the division. The biological studies relate not only to the seasonal history and habits of the species, but also to the environmental factors governing the behavior of the insects in all stages of development, such as the part played by predators, parasites, and other associated insects; the effects of climate, forest type, and site; the characteristics of the trees selected for attack; and the reaction of the tree itself during and after attack. All these factors, to which

much attention is given, have an important bearing on brood development, and their study is throwing considerable light on the causes underlying the rise and fall of bark-beetle epidemics. Results have already been obtained which will have an application in timber-sale regulations, logging operations, and slash disposal. In certain cases, it has been possible, as a result of these researches, to predict the decline of epidemics and thus to save considerable money that might otherwise have been expended needlessly in direct control. These studies are centered on the national forests in California, Oregon, Montana, Arizona, Colorado, and North Carolina.

As the stumpage value of timber increases in the Western States, the protection of the great stands of mature timber from these bark-beetle losses constitutes an ever-growing problem. A considerable part of the funds appropriated is utilized in rendering service to agencies cooperating in protection from this class of insects. This service consists largely in the direction and administration of the technical features of control projects against tree-killing bark beetles as well as in the analysis of the results. In this work close cooperation is maintained with the Forest Service, the National Park Service, the Office of Indian Affairs, and private owners. At the request of these organizations or individuals surveys are made of beetle-infested areas and recommendations given as to the advisability of control measures and as to the methods to be employed. Occasionally these projects cover enormous areas and involve very complex administrative details.

Field laboratories for the regional administration of these investigations are maintained at Coeur d'Alene, Idaho, and at Palo Alto, Calif. From four to six additional temporary bases are in use during the field season.

INSECTS INJURIOUS TO FOREST PRODUCTS

From a monetary standpoint, because of the higher values involved, insects destructive to forest products take an even greater annual toll than do the bark beetles. There are a great number of insects destructive to all forms of wood products from the green felled tree to the finished seasoned article. Much time and labor have been devoted to the development of methods for preventing damage to material of this character. In many cases all that is necessary for adequate protection is to make simple alterations in the methods of handling the material in the woods, or at the mill or warehouse. Suggested modifications of building codes to insure insect-proof construction are being widely adopted (87). Again, repellent sprays or dips are effectively employed, or the wood is impregnated with chemicals that ward off insect attack. Two experimental grounds are maintained—one at East Falls Church, Va., and the other in the Canal Zone, Panama—where chemicals and wood preservatives are tested for efficiency against wood-boring insects. The Forest Products Laboratory of the Forest Service, at Madison, Wis., treats wood samples with preservatives to be tested at these experimental grounds. Cooperation is maintained, also, with many organizations interested in wood preservation.

INSECTS AFFECTING FOREST MANAGEMENT

Cooperation with the Forest Service experiment stations forms another important line of activities. At present entomologists are stationed at, or are in close touch with, five of these experiment stations, namely, the Lake States Forest Experiment Station, St. Paul, Minn., the Northeastern Forest Experiment Station, Amherst, Mass., the Appalachian Forest Experiment Station, Asheville, N. C., the California Forest Experiment Station, Berkeley, Calif., and the North Pacific Forest Experiment Station at Portland, Oreg. It is planned for the future to have entomologists assigned to each station.

The investigations carried on under this form of cooperation have to do with the insect problems involved in the management of the more important forest types of the regions. Through studies of the major forest insect pests a better understanding of the conditions favoring or retarding serious insect damage is gained, and this knowledge is made available to foresters so that it can be applied in developing sound practices in timber culture. Some species of trees and some types of forests are little affected by insects, but in others insects may be the limiting factor in successful timber production. As better cultural practices for growing timber are attempted, the complex relations existing in the forests must be more fully understood to avoid fatal mistakes. Insects constitute one of the most important of the biotic factors. Although greatest recognition is given to their rôle as destroyers of green standing timber, other important activities that bring about changes in the composition of the forest can not be disregarded. Insects often increase the percentage of undesirable species of trees, inhibit the reproduction of certain desirable species, affect the rate of growth, and thus lengthen the rotation period and augment the ill effects of fire. Investigations dealing with these varied problems are under way.

Some other problems of a more specific character are under investigation, such as the entomological aspects of slash disposal (72), the interrelation of insects and forest fires (78, 79), turpentine practices and insect damage, insects affecting nursery practices, and insect injury to the seed of forest trees.

INSECTS INJURIOUS TO SHADE TREES AND ORNAMENTAL SHRUBS

Little in the way of investigation of insects affecting shade and ornamental trees and hardy shrubs is done at the present time, and most of the effort in this work is expended in service. There is a great and increasing demand from municipal authorities, estate owners and managers, small property owners, tree surgeons, and nurserymen for advice on methods of handling their entomological problems. The division of forest insects attempts to meet this demand by correspondence, publications, and occasional surveys and studies. Whenever possible, observations are made on important species and, where practical, experiments in methods of control and improvements in such methods are conducted.

THE MOST IMPORTANT FOREST INSECTS

This list includes those species of forest insects which are of the greatest economic importance at the present time. In the discussion of each species the common and technical names are given. There is added some information on the occurrence of the insect, the character of damage or notable epidemics, and the most important or available references to it in literature.

For the convenience of those not acquainted with the systematic classification of insects, the species listed have been arranged in groups according to the type of damage, such as bark beetles, leaf feeders, etc., and these groups are further subdivided as necessary.

TREE-KILLING BARK BEETLES

THE SOUTHERN PINE BEETLE

Dendroctonus frontalis Zimm.

In the southeastern part of the United States the southern pine beetle stands out as the most important tree-killing bark beetle. This species becomes exceedingly abundant at irregular intervals; and for several years, during one of these outbreaks, thousands of pines may be attacked and killed. In 1910 and 1911 timber valued at over \$2,000,000 was destroyed (1, 40, 42)¹.

THE EASTERN SPRUCE BEETLE

Dendroctonus piceaperda Hopk.

In past years, before the virgin spruce forests of the Northeast were cut, this bark beetle was responsible for serious depredations. Of late years little timber has been killed by this beetle in the United States, though serious losses are reported from Canada (42).

THE TURPENTINE BEETLES

Dendroctonus valens Lec.

Dendroctonus terebrans Oliv.

The red turpentine beetle (*Dendroctonus valens*) and the black turpentine beetle (*D. terebrans*) are widely distributed in North America. They attack the bases and roots of practically all species of pine and spruce, causing the exudation of large masses of pitch. They rarely kill trees and are of importance more as a result of the attention they attract than because of the damage inflicted (1, 30, 42, 46).

THE MOUNTAIN PINE BEETLE

Dendroctonus monticolae Hopk.

The mountain pine beetle has a very wide range, extending from the Sierras of central California through the northern Rockies into Canada. It attacks lodgepole, sugar, western white, and yellow pines. Nearly every year severe outbreaks are in progress in some part of this area. One notable infestation has been progressing since 1909 and now bids fair to destroy practically all the lodgepole pine on the Bitter Root and Beaverhead National Forests (42, 46).

¹ Reference is made by italic numbers in parentheses to Literature Cited, p. 23.

THE BLACK HILLS BEETLE

Dendroctonus ponderosae Hopk.

The Black Hills beetle is the most destructive enemy of yellow pine in the Rocky Mountain region. It has a marked tendency to increase suddenly to extraordinary numbers and, after destroying great quantities of timber, to subside quickly. Notable epidemics occurred in the Black Hills from 1898 to 1905 and in the Kaibab National Forest from 1920 to 1925 (42).

THE DOUGLAS FIR BEETLE

Dendroctonus pseudotsugae Hopk.

The Douglas fir beetle is most destructive in the northern Rockies and Canada. It seldom kills Douglas fir west of the Cascade Range. Outbreaks are local and sporadic, frequently associated with windthrows, defoliations, or logging operations (42).

THE WESTERN PINE BEETLE

Dendroctonus brevicornis Lec.

The western pine beetle occurs over practically the same territory as the mountain pine beetle and extends to the southern limit of the yellow pine type in California. This species rarely causes spectacular outbreaks but takes an annual toll of from 0.1 per cent to occasionally 5 per cent of the stand. It is of economic importance only in the western yellow pine forests (42, 46).

THE SOUTHWESTERN PINE BEETLE

Dendroctonus barberi Hopk.

In habits this species resembles its near kin, the western pine beetle, though it is not so destructive. Occasional outbreaks in southern Colorado, Arizona, and New Mexico take a considerable toll of timber but are short-lived. The activities of the beetle increase in years of drought (42).

THE JEFFREY PINE BEETLE

Dendroctonus jeffreyi Hopk.

The Jeffrey pine beetle restricts its activities entirely to the tree for which it is named. It is most injurious in the forests east of the Sierras, where it normally takes a small annual toll. Following a severe windstorm in the Inyo National Forest, Calif., some 6,000,000 feet of timber were killed by this species in 1924 and 1925 (42, 46).

THE ENGRAVER BEETLES

Ips spp.

The engraver beetles of the genus *Ips* are represented by numerous species. They are usually associated with dying or recently felled softwoods. Occasionally they contribute to the death of timber weakened from other causes and may even kill outright healthy timber when they are present in great numbers. Sporadic out-

breaks, in which large groups of young trees and occasionally mature trees are killed, often follow drought, windfalls, and slashings. Epidemics will not continue in healthy timber because the broods fail to develop, and such outbreaks as occur are short-lived. In the southern part of the United States the three most important species of Ips are *grandicollis* Eichh., *calligraphus* Germ., and *avulsus* Eichh. In the Western States *confusus* Lec., *emarginatus* Lec., *integer* Eichh., *oregoni* Eichh., and *radiatae* Hopk. are the most destructive species. *I. oregoni* very commonly kills the tops of mature western yellow pines, rendering the trees susceptible to the attack of Dendroctonus beetles (1, 46, 58, 63, 66).

MONTEREY CYPRESS BARK BEETLE

Phloeosinus cristatus Lec.

Several species of *Phloeosinus*, a genus of bark beetles, apparently are capable of killing trees belonging to those genera allied to Cupressus. This species seems to be particularly aggressive in Monterey cypress planted in the vicinity of San Francisco. It is usually associated, however, with scale insects and fungi, and its specific role has not been accurately determined.

THE HICKORY BARK BEETLE

Scolytus quadrispinosus Say

The hickory bark beetle is an important enemy of hickory in the Eastern States. Every few years local outbreaks of the hickory bark beetle destroy considerable timber in the natural range of the tree from Massachusetts southward to Georgia and westward into the Mississippi Valley. Recent studies indicate that these outbreaks are correlated with droughts or conditions unfavorably affecting the trees (2, 43).

THE FIR BARK BEETLE

Scolytus ventralis Lec.

This species, widely distributed throughout the Western States, attacks standing dead, dying, or living firs, frequently killing them but more often destroying only the tops of the trees or patches of bark on the trunk, which results in defects. In California, following the drought of 1924, very extensive losses were reported (40, 46).

INSECTS BORING IN LIVING TREES

THE SMALL METALLIC WOOD AND BARK BORERS

Agrilus anxius Gory

Agrilus bilineatus Web.

The bronze birch borer (*Agrilus anxius* Gory) is very destructive to several species of birch (*Betula*) in the Northeast. Its control is becoming an important problem in the management of hardwood stands. The 2-lined chestnut borer (*A. bilineatus* Web.) is of secondary importance throughout the eastern part of the United States. It occasions the death of oaks weakened by other causes, though in Minnesota it is reported to be more of a primary pest (8, 14, 57, 59).

THE TURPENTINE BORER

Buprestis apicans Hbst.

The turpentine borer attacks the exposed wood resulting from fire scars, blazes, and turpented faces of several species of southern pines, making the trees susceptible to windthrow, and the butt log unfit for lumber. In recent years, with the wider adoption of conservative turpentine practices and forest management, injury by this insect is receiving greater recognition (8, 19).

THE LOCUST BORER

Cyllene robiniae Forst.

The locust borer frequently prevents the growth of a valuable tree, the black locust, in some regions. It occurs throughout the Eastern States and has been introduced with the host into Colorado and adjacent States (17, 29, 41).

THE OAK TWIG PRUNER

Elaphidion villosum Fab.

The oak twig pruner, occurring throughout the eastern half of the United States, locally causes serious damage to various hardwoods by cutting off the branches. The pruning in this case is done by the larvae (48).

THE SUGAR MAPLE BORER

Glycobius speciosus Say

The sugar-maple borer is an especially injurious pest of the sugar maple tree in the Northeastern States. This borer attacks trees apparently in full vigor, especially those growing in the open, and kills limbs and sometimes the entire tree. This insect is probably the most serious enemy of the sugar maple tree (25).

THE LIVING HICKORY BORER

Goes pulcher Hald.

THE LIVING BEECH BORER

Goes pulverulenta Hald.

THE WHITE OAK BORER

Goes tigrina DeGeer

THE OAK SAPLING BORER

Goes tessellata Hald.

The four species of roundheaded borers listed above attack the trunks of a variety of hardwoods, including oak, hickory, beech, elm, sycamore, blue beech, and ironwood, throughout the eastern part of the United States. The larvae bore deeply into the wood, causing large, unsightly defects and culls in the lumber. In younger trees these defects frequently cause breakage under the strain of wind or ice storms (18).

THE PINE AND HEMLOCK FLAT-HEAD BORERS

Melanophila spp.

Two species of the genus *Melanophila*, the eastern hemlock bark borer (*M. fulvoguttata* Ham.) and the western flat-headed borer (*M. gentilis* Lec.), are of considerable economic importance. The eastern form kills hemlock weakened by defoliation and other causes. The western species is frequently an important pest during epidemics of the western-pine bark beetle in western yellow pine and to trees affected by drought (8, 46).

THE TWIG TUNNELERS

Oberea spp.

Several species of *Oberea* bore down through the pith in the twigs of several species of trees and shrubs, causing the death of the infested portion. The work can always be distinguished from that of other twig girdlers by the linear series of small holes cut through the bark. *O. myops* Hald. in rhododendron and azalea, *O. schaumii* Lec. in poplar, and *O. tripunctata* Fab. and *O. bimaculata* Oliv. in plum, apple, peach, and dogwood are the most important (18).

THE LONG-HORNED GIRDLERS

Oncideres cingulata Say*Oncideres putator* Thom.

The hickory twig girdler (*Oncideres cingulata*) and the huisache girdler (*O. putator*) cause considerable damage to a great variety of hardwood trees in the Eastern and Gulf States by cutting off the branches. The girdling is done by the adult beetle (18, 37).

THE COTTONWOOD BORER

Plectrodera scalator Fab.

In the central part of the United States the cottonwood borer causes serious injury to the base and roots of several species of cottonwood and willow. It is sometimes abundant enough in certain localities to kill these trees (52).

THE PRIONID ROOT BORERS

Prionus spp.

Several species of these large roundheaded borers—*Prionus latifrons* Drury and *P. imbricornis* L. in the East, and *P. californicus* Motschulsky in the Western States—bore in the roots of living hardwoods, thus lowering the vitality of the trees. The attack is frequently followed by root rots such as *Armellaria* and attacks of secondary bark borers which ultimately kill the trees (16).

THE RED OAK BORER

Romaleum rufulum Hald.

The red oak borer is of considerable economic importance throughout its range in the central and eastern part of the United States and Canada. The habits of this species somewhat resemble those of

Prionoxystus and Goes. The borers attack living oak trees, mining deeply into the sapwood and heartwood (18).

THE ELM BORER

Saperda tridentata Oliv.

THE LINDEN BORER

Saperda vestita Say

THE POPLAR BORER

Saperda calcarata Say

The larvae of the poplar borer riddle the heartwood of several species of poplar, opening it to decay and making the trees subject to windfall. It is widely distributed throughout the range of the host plants. Several other species of this genus, among which *Saperda tridentata* in elm and *S. vestita* in linden are important, are injurious to living trees (18, 27, 38).

THE WESTERN FIR BORER

Tetropium abietis Fall

In the Pacific coast region the western fir borer occasionally kills large numbers of true firs by mining beneath the bark. Trees weakened by defoliation or fire are particularly susceptible to attack (18).

THE WESTERN LARCH BARK BORER

Tetropium velutinum Lec.

The western larch bark borer is occasionally a pest of considerable importance in stands of western larch and hemlock in the Rocky Mountain and Pacific coast regions of the United States. The larvae mine in the inner bark (18, 68).

THE CEDAR POLE BORER

Trachykele blondeli Mars.

The larvae of the western cedar pole borer riddle the heart and sapwood of living western red cedar, causing serious defects and the consequent rejection of the damaged material for lumber, shingles, and poles. This borer occurs in Oregon, Washington, and California (9).

THE PITCH MOTHS

Dioryctria abietella D. and S.

Dioryctria xanthaenobarea Dyar

Dioryctria ponderosae Dyar

Dioryctria amatella Hulst

The inner bark of the trunks, the new growth of the leaders, and the cones, of pine, Douglas fir, spruce, and fir are often seriously injured by the caterpillars of the pitch moths. Death of parts or even the entire tree and the loss of seed of the infested cones frequently follow the injury. These insects are important, but little work has been done on them.

THE ZIMMERMAN PINE MOTH

Pinipestis zimmermani Grote

The Zimmerman pine moth is a serious insect pest of pine, attacking western yellow pine, white pines, Scotch pine, and Austrian pine. The caterpillar bores into the bark of the trunk, usually avoiding the base and top, and causes the malformation known as spike top or sometimes kills the entire tree. The injury it produces also brings about a considerable depreciation in the value of the timber. The species is especially abundant in the northern and western part of the United States (6).

THE CARPENTER WORM

Prionoxystus robiniae Peck

The larvae of this large moth bore in a great variety of eastern hardwoods, especially oak and locust, causing large wormholes and resulting defects in the lumber (44, 48).

THE DOUGLAS FIR PITCH MOTH

Synanthedon novaeocensis Hy. Edw.

The Douglas fir pitch moth is an important enemy of Douglas fir over the northern Rocky Mountain and Pacific coast regions. The insect causes pitch seams, gum check, and other defects which seriously depreciate the value of the timber. Its attack appears to be restricted to that portion of the trunk which later clears itself of branches (5).

THE SEQUOIA PITCH MOTH

Vespamima sequoiae Hy. Edw.

The Sequoia pitch moth attacks almost all conifers of the Northwest. It is particularly abundant and destructive to lodgepole pine in certain regions of Montana. The insect infests the extreme base of the trees, causing a heavy pitch flow over the base of the trunk and surrounding ground. The trees attacked are frequently killed and jeopardize other trees in the neighborhood through increased fire hazard (4).

THE LEOPARD MOTH

Zeuzera pyrina Fab.

The leopard moth kills the limbs of hardwood trees and is locally serious in the New England and the Northeastern States. The species was introduced from Europe and is spreading slowly to the South and West (45).

THE CAMBIUM MINERS

Agromyza aceris Greene*Agromyza amelanchieris* Greene

Birch, cherry, maple, oak, poplar, and many other less commonly used hardwood trees have small, yellowish-brown spots or streaks called "pith flecks" made by the larvae of several species of flies, the names of two species being given above. In the Eastern States this injury sometimes lowers the grade of the product for certain uses (34).

THE HEMLOCK BARK MAGGOT
Cheilosia alaskensis Hunter

The hemlock bark maggot in the northern Rocky Mountain and Pacific coast regions causes defects in western hemlock known as "black check." The injury is caused by the larvae working in the growing tissue of the tree. Similar defects in fir, spruce, and pine are caused by related species (?).

INSECTS AFFECTING FOREST PRODUCTS
AMBROSIA BEETLES OR PINHOLE BORERS

A number of species of beetles included in the genera *Corthylus*, *Gnathotrichus*, *Pterocyclon*, *Xyleborus*, and *Platypus* are important because of the injury they do to fire-scarred trees, green logs, and green lumber. These insects, while working in the wood, not only make holes but also introduce wood-staining fungi. In some localities a considerable quantity of wood otherwise sound and useful is discarded because by the work of these beetles it is rendered unfit for certain special uses (39, 40, 62).

THE COLUMBIAN TIMBER BEETLE
Corthylus columbianus Hopk.

The Columbian timber beetle attacks living white oak, chestnut, and yellow poplar, entering directly through sound bark. In some localities from 15 to 25 per cent of the white oak lumber may be affected, and where tight barrel staves are being manufactured whole trees are discarded because of injury by this insect. The wood may still be used for such purposes as base for veneer and for the invisible parts of various structures, as its strength is little impaired (39, 40, 62).

THE OAK TIMBER WORM
Eupsalis minuta Drury

The oak timber worm, entering through wounds, attacks the wood of living trees and also frequently causes much damage to green logs and produces defects in the lumber. It occurs throughout the eastern part of the United States (39, 40, 62).

THE CHESTNUT TIMBER WORM
(*Lymericylon*) *Melittomma sericeum* Harris

The chestnut-timber worm attacks the wood of living chestnut. Entering through a wound, it bores deeply into the center of the tree. In the recent past, when chestnut was a more abundant forest tree, this insect presented a serious problem, as from 75 to 90 per cent of all the chestnut timber produced was defective (39, 40, 62).

THE SAPWOOD TIMBER WORM
Hylecoetus lugubris Say

The sapwood timber worm attacks dying trees and green saw logs of basswood, buckeye, chestnut, black walnut, cottonwood, yellow poplar, and birch, causing considerable damage. Often a 5 to 10 per cent loss occurs in logs on which the bark has been left (40, 62).

THE BLACK-HORNED PINE BORER

Callidium antennatum Newm.

The black-horned pine borer attacks a variety of softwoods and, together with other closely allied species, is widely distributed over the United States. Considerable damage is caused to rustic houses and other structures where these woods with the bark still on are used for support or ornamentation. The presence of bark is necessary for attack by this species (18, 68).

THE BELTED CHION

Chion cinctus Drury

The belted chion, a long-horned borer, attacks several species of partly seasoned hardwoods, causing severe damage to the sapwood of oak and hickory, often making holes from one-half to three-fourths of an inch wide and 2 feet long and rendering it unfit for construction purposes. This insect will attack wood even after the bark has been removed from the log. Small stock used in the manufacture of rustic furniture or in making small products, such as mallets and mauls, is subject to severe damage by this insect (18, 62, 68).

THE PAINTED HICKORY BORER

Cyllene caryae Gahan

The painted hickory borer causes serious injury to green saw logs. The larvae make large oval holes in the sapwood. This part of the log, which should furnish the best grades of material, is often ruined by the resulting defects in the lumber. Besides hickory this borer attacks walnut, honeylocust, mulberry, and osage orange (18, 22).

THE WESTERN CEDAR BARK BORER

(Hylotrupes) Hewicallidium amethystinum Lec.

The western cedar bark borer is of considerable economic importance in the Pacific coast region of the United States. It injures wood of recently felled giant arborvitae and incense cedar (18, 68).

THE OLD HOUSE BORER

Hylotrupes bajulus L.

The old-house borer attacks a variety of dry, seasoned, softwood products. This species, introduced from Europe, is of considerable economic importance and is becoming more abundant and serious each year. It confines its activities to such materials as joists, studding, and rafters (18, 62).

THE CEDAR-TREE BORER

Hylotrupes ligneus Fab.

The cedar-tree borer attacks a variety of softwoods (firs, spruces, hemlocks, cedars, arborvitae, etc.), damaging the wood for commercial purposes and causing shedding of the bark of material used in rustic work. The larvae mine first between the bark and wood and later in the sapwood, occasionally going into the heartwood (18, 68).

THE PINE SAWYERS

Monochamus spp.

Several species of pine sawyers are of economic importance, namely, *Monochamus titillator* Fab. in the Southeast, *M. scutellatus* Say and *M. confusor* Kirby in the Northeast, and *M. maculosus* Hald. in the West. They attack dying and recently felled pine, spruce, and fir, their work resulting in a high percentage of culled lumber. Following windfalls and burns these species are especially destructive, causing rapid deterioration of the timber before it can be salvaged. *M. marmorator* Kirby attacks and kills living fir (*Abies balsamea*) in the Northeastern States and Canada (18, 67).

THE BANDED ASH BORER

Neoclytus caprea Say

The banded ash borer is of considerable economic importance in the eastern part of the United States, causing extensive damage to the sapwood and outer heartwood of recently felled and partly seasoned ash logs. In some large operations the grub of this beetle has been known to have completely destroyed the entire cut of one season (18, 62, 68).

THE GRAY ASH BORER

Neoclytus conjunctus LeConte

This beetle has habits similar to those of the banded ash borer. The grub mines the sapwood and heartwood of ash and oak. It is a common species in the Rocky Mountain region and in the Pacific Coast States (18).

THE RED-HEADED ASH BORER

Neoclytus erythrocephalus Fab.

The red-headed ash borer attacks recently felled hardwoods of a number of varieties, often causing from 20 to 90 per cent loss in the lowering of the grade of the lumber. Ash and hickory particularly are selected for attack. Green logs, felled during the summer in the Gulf coast and Mississippi Delta areas and left in the woods for several weeks before being removed, often suffer a total loss as the result of the activities of this insect (18, 68).

THE POLE BORER

Parandra brunnea Fab.

The pole borer is a fairly large wood borer which causes considerable damage to the bases of telephone poles and to shade trees throughout the eastern part of the United States. Telephone poles are sometimes completely severed, and the entire heart is eaten out of shade trees causing large limbs to fall or the entire tree to break over.

THE TANBARK BORER

Phymatodes variabilis Fab.

The tanbark borer is of economic importance in the tanbark industries. The grub of this beetle mines chiefly in the bark of oak (18, 39, 68).

THE FLAT OAK BORER
Smolicum cucujiforme Say

The flat oak borer is a typical heartwood feeder that attacks seasoned oak and hickory products. Its work is also often found in old fire scars on oak trees (18).

THE POWDER-POST BEETLES
Lyctus planicollis Lec. and other species of *Lyctus*

Several species of the so-called lyctus powder-post beetles are extremely destructive to seasoned hardwoods, particularly oak, hickory, ash, persimmon, etc. The annual loss to such stored products as billets, lumber, handles, and spokes runs into millions of dollars (61).

THE LEAD-CABLE BORER
Scobicia declivis Lec.

The lead-cable borer is a serious pest of oak stave wood on the west coast and has an interesting and expensive habit of boring into suspended lead-sheathed telephone cables (10).

THE RED-SHOULDERED POWDER-POST BEETLE
Xylobiops basilaris Say

The red-shouldered powder-post beetle is of considerable economic importance, causing large losses to partly seasoned wood products used in the manufacture of shuttle blocks, mallets, mauls, etc., as well as to small poles and saplings used for rustic furniture. Hickory and persimmon blocks and poles are destroyed by both adults and grubs and literally reduced to powder in a few months' time (40, 53).

OTHER BOSTRICHID POWDER-POST BEETLES
Apatides fortis Lec.
Dendrobiella aspera Lec.

Other powder-post beetles occur in the Southwest and attack mesquite, reducing the wood to powder within a short time. The large form, *Apatides fortis*, prefers wood which has dried for several months, but the smaller form, *Dendrobiella aspera*, shows a decided preference for freshly cut wood. Fence posts, poles, and cordwood are severely damaged (20).

THE CARPENTER ANT
Camponotus herculeanus pennsylvanicus DeG.

The carpenter ant is widespread and abundant in the United States. It attacks living trees through wounds, dead trees, logs, and forest products, frequently entering the wooden parts of buildings.

TERMITES (NONSUBTERRANEAN)
Kalotermes spp.

In certain sections of the United States, namely, the South Atlantic, Gulf, Southwestern, and Pacific Coast States, nonsubter-

ranean or dry-wood termites (*Kaloterms* spp.) seriously damage the woodwork of buildings and furniture, and telegraph, telephone, power, and electric-light poles. These termites are able to live in dry wood and do not need to maintain contact with the ground as they do not require much moisture (60).

TERMITES (SUBTERRANEAN)

Reticulitermes spp.

The most injurious termites throughout the entire United States are ground-nesting or subterranean forms (*Reticulitermes* spp.). The regions where damage is most serious are portions of the country with warm or temperate climates, including the Eastern States, the central West, and the Pacific coast, as well as all areas south of these regions. These termites are chiefly injurious to wood either in contact with the ground or insufficiently protected, insulated, or removed from the ground. The annual damage caused by these termites in the United States is many millions of dollars, these losses being due not only to the value of the wood destroyed but also to the cost of labor of removal, reconstruction, or replacement (60, 87).

INSECTS AFFECTING SEEDS, FOREST REPRODUCTION, AND NURSERY PLANTS

THE SPRUCE TWIG GALL LICE

(*Chermes* or *Gillettea*) *Adelges cooleyi* Gill.

(*Chermes*) *Adelges abietis* L.

These insects and other related species are important forest, nursery, and ornamental-tree pests, producing characteristic twig-deforming growths which frequently kill the parts attacked. They occur throughout the natural range of the spruces and firs.

THE NUT WEEVILS

Curculio spp.

Weevils of this group are important destroyers of the seeds of certain hardwood trees. Hickory nuts, acorns, chestnuts, and hazelnuts are heavily infested by these snout beetles which make the seed unviable with a resultant loss in reproduction. Infestation in seeds collected for nursery planting occasionally exceeds 20 per cent (3, 25, 47).

THE CONE BEETLES

Conophthorus spp.

The cone beetles bore into the base of the developing cones of many species of pine. Drying and falling, or prolonged retention of the cones, may follow the attack. In some years, in restricted localities, the entire seed crop is destroyed. The more important species are *Conophthorus ponderosae* Hopk., the yellow pine cone beetle; *C. lambertianae* Hopk., the sugar pine cone beetle; and *C. coniperda* Sz., the eastern white pine cone beetle (46, 50, 51).

ACORN GALL MAKERS

Cynipidae

A number of 4-winged gallflies destroy large quantities of oak seed by producing gall formations in the cotyledons. They lower the natural reproduction of oak and are also responsible for a heavy loss in collected seed (26).

SEED CHALCIDS

Megastigmus albifrons Walk.

Megastigmus milleri Rohwer

Megastigmus piceae Rohwer

Megastigmus pinus Parfitt

Megastigmus spermatophus Wachtl

The seeds of many trees are attacked by certain chalcids that feed on plant tissue. These small, 4-winged flies destroy the seeds of pine, spruce, fir, juniper, and many other trees. The Douglas fir chalcid (*Megastigmus spermatophus*) is at times particularly destructive (50).

THE PALES WEEVIL

Hylobius pales Boh.

The pales weevil destroys young pine seedlings. The larvae breed in green pine stumps and logs, and the adults girdle pine seedlings from 1 to 5 years old by feeding at the base. The inroads of this beetle frequently make several plantings necessary in order that a fully stocked stand may be secured.

The species is injurious chiefly in the New England States (56).

THE PINE BARK LOUSE

(*Chermes*) *Pineus pinicorticis* Fitch

The pine bark louse, a relative of the spruce twig gall lice, occurs on the bark and needles of white pine producing a waxy secretion which appears as a whitish gray mold. Trees heavily infested for several years show the effect in scant, poor-colored foliage and stunted new growth.

MAY BEETLES OR WHITE GRUBS

Phyllophaga spp.

The control of the larvae or white grubs of a number of species of May beetles constitutes an important problem in nurseries of both forest and ornamental stock. These insects kill large numbers of both evergreen and deciduous seedling trees and shrubs by feeding on the roots. The beetles when abundant defoliate hardwood trees (21).

THE WHITE PINE WEEVIL

Pissodes strobi Peck

The white pine weevil is the most serious pest of white pine in the Northeastern States. Successive killing of the terminal shoots

of saplings and young trees results in stunting and malformation and ultimately in low-grade timber. Over 75 per cent of the white pines in this region are weeviled, resulting in a reduction in value of from 20 to 25 per cent below that of trees of normal development (32).

THE EUROPEAN PINE-SHOOT MOTH

Rhyacionia buoliana Schiff.

This imported pine-shoot moth, first found in the United States in 1914, is known in Europe as a very serious pest. The species is confined to pine but is recorded as attacking "all species" of this genus. It has been found in this country on Scotch pine, Austrian pine, mugho pine, white pine, red pine, and longleaf pine. This moth has become established in Illinois, Ohio, West Virginia, Pennsylvania, New Jersey, New York, Massachusetts, Connecticut, Rhode Island, and Florida. At present it seems to be largely a nursery pest.

THE NANTUCKET PINE MOTH

Rhyacionia frustrana Comst.

The Nantucket pine moth is common on young pine throughout the Eastern and Southern States. Periodically it becomes of great importance locally. Some years ago it was introduced into the Forest Service plantations at Halsey, Nebr., where it has become established and is proving very injurious, especially to western yellow pine, the most desirable tree species (23, 53, 65).

THE PINE NEEDLE SCALE

Chionaspis pinifoliae Fitch

The pine needle scale is one of the few scale insects that are serious forest pests in the United States. It occasionally becomes sufficiently abundant in the forests of California and Minnesota to weaken the trees and make them susceptible to bark-beetle attack. It is also an important pest of ornamental pine and spruce (23, 44).

THE CYPRESS BARK SCALE

Ehrhornia cupressi Ehrhorn

Monterey cypress in California, especially trees in thickly planted hedgerows or windbreaks, are severely injured by this scale insect (23, 35).

THE PINE SOFT SCALES

Toumeyella spp.

Toumeyella numismaticum (P. and McD.), *T. parvicorne* Ckll., *T. pini* King, and unidentified species of this genus have been reported as injurious to pines east of the Rocky Mountains. *Pinus sylvestris*, *P. divaricata*, *P. radiata*, *P. virginiana*, and *P. mugho* are recorded as hosts from Louisiana, Michigan, Minnesota, Nebraska, North Carolina, and Pennsylvania. *T. numismaticum* promises to be an important pine nursery problem, as it has appeared in injurious numbers in forest-tree nurseries of the South and has killed many

trees in the plantations of the Nebraska National Forest at Halsey, Nebr., and in Minnesota.

DEFOLIATING INSECTS

THE LOCUST LEAF MINER

Chalepus dorsalis Thunb.

The locust leaf miner causes widespread defoliation of black locust. It is abundant practically every year over considerable areas of its range from Massachusetts through the southern Appalachians and the Ohio Valley. The feeding causes the leaves of the trees to turn brown in the late summer. No study has been made of its effect on the growth of the trees (48, 53).

THE GRAPE COLASPIS

Colaspis brunnea Fab.

Widespread defoliation of pines in the Gulf States in 1925 and 1926 was attributed to this small beetle. The species is not often so abundant. Severest injury occurred in southern Louisiana and eastern Texas. Young trees were more severely infested than older ones.

THE ORANGE-STRIPED OAK WORM

Anisota senatoria A. and S.

THE SPINY OAK WORM

Anisota stigma Hbn.

THE GREEN-STRIPED MAPLE WORM

Anisota rubicunda Fab.

Throughout the eastern hardwood belt oaks are frequently defoliated by the orange-striped oak worm, the spiny oak worm, and *Anisota virginiensis* Drury, and maples by the green-striped maple worm. The first-mentioned species is the most important and injurious (25, 44, 48).

THE BIRCH LEAF SKELETONIZER

Bucculatrix canadensisella Chambers

This insect is a frequent defoliator of birch throughout the northern part of the United States and Canada and feeds to some extent on alder and oak.

THE SPRUCE BUD WORM

Cucoccia fumiferana Clem.

The spruce bud worm is periodically one of the most destructive forest insects in the United States. A recent epidemic in the North-eastern States and Canada destroyed from 50 to 75 per cent of the spruce and fir over great areas. In the Lake States and locally throughout the Rocky Mountains notable losses have been reported. The caterpillars feed on fir, spruce, Douglas fir, pine, larch, and hemlock (64).

THE CATALPA SPHINX

Ceratonia catalpae Bdv.

Catalpas are frequently defoliated by the caterpillar of the catalpa sphinx moth. When the trees are grown in plantations this insect occasionally becomes extremely abundant and destructive. The most serious injury has occurred in the Ohio Valley (44, 48).

THE LARCH CASE-BEARER

Coleophora laricella Hbn.

The larch case-bearer, introduced from Europe, has become widely distributed in the northeastern part of the United States, the Lake States, and Canada and is causing considerable injury to larch. An extensive outbreak of this insect occurred in New England from 1923 to 1925 and in Maine thousands of acres of larch were defoliated. The trees which were stripped annually began to die in 1926.

THE PANDORA MOTH

Coloradia pandora Blake

This species causes serious defoliation of western yellow pine and lodgepole pine growing on the volcanic soils east of the Sierra Nevada and Cascade Ranges in California and Oregon. Trees are defoliated only in alternate years and, although few die as a direct result of defoliation, they are rendered more susceptible to bark-beetle attack (55).

THE YELLOW-NECKED CATERPILLAR

Datana ministra Walk.

THE WALNUT CATERPILLAR

Datana integerrima G. & R.

These two species of *Datana* are occasionally important. The yellow-necked caterpillar is a general feeder and is sometimes abundant in hardwood forests. The walnut caterpillar is a frequent defoliator of walnut trees throughout the eastern hardwood belt (25, 44, 48).

THE HEMLOCK LOOPERS

Ellopiia spp.

Hemlock, spruce, and Douglas fir in the Lake States and on the Pacific coast are sometimes very severely defoliated and killed by several species of caterpillars belonging to the genus *Ellopiia*. *E. fiscellaria* Gn. in the Lake States, *E. somniaria* Hulst in the northern Pacific States, and an undetermined species in California are the most important (28).

THE DOUGLAS FIR LOOPER

Galenara sp.

At higher elevations in the spruce-fir type of the southern Rocky Mountain region, particularly in New Mexico, the Douglas fir looper

has periodically destroyed the timber over considerable areas. Several recent outbreaks have occurred in the Sacramento Mountains. Douglas fir is the favored host but spruce foliage is eaten as well. *Dendroctonus engelmanni* frequently follows and destroys not only weakened trees but vigorous stands outside the area of defoliation.

THE LODGEPOLE NEEDLE TIER

Argyrotaenia pinatubana Kearf.

The lodgepole needle tier, supposed to be the same as the defoliator of eastern white pine, became very abundant in 1920 to 1925 on the headwaters of the Madison River in Yellowstone National Park and the Madison National Forest. In conjunction with a sawfly (*Neodiprion* sp.) these insects killed most of the lodgepole pine over an area of 75 square miles.

THE BROWN-TAIL MOTH

Nygmia phaeorrhoea Donovan

The brown-tail moth is an insect which has been introduced and become established in the New England States. It is locally and periodically abundant, defoliating apple, pear, plum, oak, willow, elm, maple, and other common trees and shrubs. It does not attack conifers. The hairs from the caterpillars are a source of annoyance, causing a rash and irritation of the skin (12).

THE DOUGLAS FIR TUSSOCK MOTH

Hemerocampa pseudotsugata McD.

This defoliator has come into prominence as an insect of economic importance within the last ten years in the northern Rocky Mountain regions, Washington, and British Columbia. Locally through this region it has killed a great deal of Douglas fir, but the outbreaks usually disappear in about three years. The female is wingless. The caterpillars shed poisonous hairs which cause a rash on susceptible skins.

OAK AND MAPLE DEFOLIATORS

Heterocampa spp.

Caterpillars belonging to the genus *Heterocampa* occasionally become exceedingly abundant in hardwood forests and severely defoliate trees over large areas. In 1918 and 1919 maple and birch were severely defoliated in New Hampshire by *Heterocampa bilineata* Pack. and *H. biundata* Walk., and elm in Essex County, N. Y., by *H. bilineata* in 1918. *H. guttivitta* Walk. defoliated maples, birch, and beech in New York, Vermont, and New Hampshire in 1910 and 1919. *H. manteo* Dbdy. defoliated forest trees, oak, beech, maple, and hickory, in 1922 over an extensive area involving Virginia and North Carolina. Many oaks were completely stripped. In most of these defoliations caterpillars of more than one genus were reported present although species of *Heterocampa* played the dominant rôle (15, 25).

THE TENT CATERPILLARS

Malacosoma disstria Hübn.*Malacosoma americana* Fab.*Malacosoma californica* Pack.*Malacosoma constricta* Stretch*Malacosoma pluvialis* Dyar*Malacosoma fragilis* Stretch

The tent caterpillars are frequently important forest defoliators as well as farm and orchard pests. They have a wide range of host plants. Some species are widely distributed over the United States, whereas others are rather restricted in distribution. A recent infestation on the Carson National Forest in New Mexico spread from 10 sections in 1926 to approximately 225 sections in 1929 (23, 25, 48).

THE PINE BUTTERFLY

Ncophasia menapia Felder

At irregular periods large areas of western yellow pine, western white pine, and lodgepole pine are defoliated in Idaho, Montana, and the Pacific Coast States by caterpillars of the pine butterfly. Death of the trees through loss of the needles or attack by bark beetles follows severe defoliation (24).

THE HEMLOCK BUDWORM

Peronea variana Fern.

The hemlock budworm, as recognized at present, occurs in the northeastern part of the United States and Canada as well as in the Pacific Northwest, British Columbia, and Alaska. At times it is extremely destructive over wide areas, feeding chiefly on hemlock, although the spruces and firs are also attacked. At present active outbreaks occur along the St. Lawrence River in the eastern part of Canada and on the Olympic peninsula on the Pacific Coast. Little is known of its habits.

THE GIPSY MOTH

Porthetria dispar L.

The control of the gipsy moth is a problem in the care of shade, park, and forest trees in the New England States. The nearly omnivorous feeding habits of the larvae of this insect and the methods of pupation and egg laying have rendered it a pest exceptionally difficult to control. It has killed many trees, particularly oaks, in the infested area (12).

THE CALIFORNIA OAK WORM

Phryganidia californica Pack.

Considerable areas of oaks on the Pacific coast are frequently defoliated by the caterpillars of the California oak moth. This species is at times sufficiently abundant and injurious to demand control measures (11).

THE PINE NEEDLE MINER AND OTHER CONIFEROUS LEAF MINERS

Recurvaria spp.

These leaf miners are often important pests of coniferous trees, especially pines, in the forests of the Pacific coast region. The genus is widespread throughout the United States and contains many species attacking other genera of coniferous trees besides pines. In the Yosemite National Park, Calif., great areas of lodgepole pine were killed by *Recurvaria milleri* Busck (54).

THE SATIN MOTH

Stilpnotia salicis L.

The satin moth, a recently introduced insect, has become established in the New England States. It has increased in abundance and spread with great rapidity. The caterpillars prefer poplar and willow, but will feed to a certain extent on other species of deciduous trees, chiefly oaks. The principal damage caused by the insect is the complete defoliation of many poplars and willows (13).

THE LARCH SAWFLY

Nematus erichsonii Htg.

Larch in the Northeastern States and in the Lake States is periodically defoliated by the larvae of the larch sawfly. Several of these great outbreaks occurring since 1880 have practically wiped out all the mature larch in the northeastern part of the United States and southeastern Canada.

The western larch in Idaho is similarly attacked by two related sawflies (36, 53).

THE LODGEPOLE SAWFLY

Neodiprion sp.

The lodgepole sawfly occasionally kills lodgepole pine in Idaho, Montana, and Wyoming. The most recent destructive outbreak occurred in connection with the outbreak of the lodgepole needle tier mentioned under that species.

OTHER SAWFLIES

Neodiprion spp.

Jack pine in Minnesota was severely defoliated in 1923 and 1924 by sawfly larvae of this genus. *Neodiprion banksianae* Roh. and another species near *N. scutellaris* Roh. were found attacking western yellow pine in Idaho in 1922 and young lodgepole pine in West Yellowstone, Mont., in 1924 and 1925.

During a widespread outbreak in 1926, which extended from Massachusetts to Louisiana, pines in Mecklenburg County, southern Virginia, suffered a severe defoliation, a large percentage of the pines in the county being attacked.

An outbreak of *N. excitans* Roh. severely defoliated pines in Montgomery County, eastern Texas, in 1919 and was reported as doing serious damage to pine timber.

The red-headed pine sawfly, *N. lecontei* Fitch, attacks young pines in the Eastern States and as far west as Wisconsin. Not infrequently the trees infested are completely defoliated and, as a result, are killed or severely stunted (31, 49).

THE BIRCH LEAF MINER

Phyllotoma nemorata Fallen

This sawfly was described as a new species native to America but was later found to be a known species that had been introduced from Europe. It was first discovered in North America in Nova Scotia, about 1905. Since then it has spread rapidly south and west in the birch forests of Maine, Massachusetts, and New Hampshire. The insect mines the leaves of birch, showing a preference for white birch. The defoliation is most complete late in the growing season.

LITERATURE CITED

- (1) BLACKMAN, M. W.
1922. MISSISSIPPI BARK BEETLES. Miss. Agr. Expt. Sta. Tech. Bul. 11, 130 p., illus.
- (2) ———
1924. THE EFFECT OF DEFICIENCY AND EXCESS IN RAINFALL UPON THE HICKORY BARK BEETLE (*ECCOPTOGASTER QUADRISPINOSUS* SAY). Jour. Econ. Ent. 17: 460-470, illus.
- (3) BROOKS, F. E.
1922. CURCULIOS THAT ATTACK THE YOUNG FRUITS AND SHOOTS OF WALNUT AND HICKORY. U. S. Dept. Agr. Bul. 1066, 16 p., illus.
- (4) BRUNNER, J.
1914. THE SEQUOIA PITCH MOTH, A MENACE TO PINE IN WESTERN MONTANA. U. S. Dept. Agr. Dept. Bul. 111, 11 p., illus.
- (5) ———
1915. DOUGLAS FIR PITCH MOTH. U. S. Dept. Agr. Bul. 255, 23 p., illus.
- (6) ———
1915. THE ZIMMERMAN PINE MOTH. U. S. Dept. Agr. Bul. 295, 12 p., illus.
- (7) BURKE, H. E.
1905. BLACK CHECK IN WESTERN HEMLOCK. U. S. Dept. Agr., Bur. Ent. Circ. 61, 10 p., illus.
- (8) ———
1917. FLAT-HEADED BORERS AFFECTING FOREST TREES IN THE UNITED STATES. U. S. Dept. Agr. Bul. 437, 8 p., illus.
- (9) ———
1928. THE WESTERN CEDAR POLE BORER OR POWDER WORM. U. S. Dept. Agr. Tech. Bul. 48, 16 p., illus.
- (10) ——— HARTMAN, R. D., and SNYDER, T. E.
1922. THE LEAD-CABLE BORER OR "SHORT-CIRCUIT" BEETLE IN CALIFORNIA. U. S. Dept. Agr. Bul. 1107, 56 p., illus.
- (11) ——— and HERBERT, F. B.
1920. CALIFORNIA OAK WORM. U. S. Dept. Agr. Farmers' Bul. 1076, 14 p., illus.
- (12) BURGESS, A. F.
1923. CONTROLLING THE GIPSY MOTH AND THE BROWN-TAIL MOTH. U. S. Dept. Agr. Farmers' Bul. 1335, 28 p., illus.
- (13) ——— and CROSSMAN, S. S.
1927. THE SATIN MOTH, A RECENTLY INTRODUCED PEST. U. S. Dept. Agr. Dept. Bul. 1469, 23 p., illus.
- (14) CHAPMAN, R. N.
1915. OBSERVATIONS ON THE LIFE HISTORY OF *AGRILUS BILINEATUS*. Jour. Agr. Research 3: 283-294, illus.

- (15) COLLINS, C. W.
1926. OBSERVATIONS ON A RECURRING OUTBREAK OF *HETEROCAMPA GUTTIVITTA* WALKER AND NATURAL ENEMIES CONTROLLING IT. Jour. Agr. Research 32: 689-699.
- (16) CRAIGHEAD, F. C.
1915. CONTRIBUTIONS TOWARD A CLASSIFICATION AND BIOLOGY OF THE NORTH AMERICAN CERAMBYCIDAE, LARVÆ OF THE PRIONINÆ. U. S. Dept. Agr., Off. Sec. Rpt. 107, 24 p., illus.
- (17) ———
1919. PROTECTION FROM THE LOCUST BORER. U. S. Dept. Agr. Bul. 787, 12 p., illus.
- (18) ———
1923. NORTH AMERICAN CERAMBYCID LARVÆ. A CLASSIFICATION AND THE BIOLOGY OF NORTH AMERICAN CERAMBYCID LARVÆ. Canada Dept. Agr. Ent. Branch Bul. (n. s.) 27, 238 p., illus.
- (19) ———
1927. THE TURPENTINE BORER ON THE FLORIDA NATIONAL FOREST. Forest Worker 3 (4): 11-12.
- (20) ——— and HOFFER, G.
1921. PROTECTION OF MESQUITE CORDWOOD AND POSTS FROM BORERS. U. S. Dept. Agr. Farmers' Bul. 1197, 12 p., illus.
- (21) DAVIS, J. J.
1929. COMMON WHITE GRUBS. U. S. Dept. Agr. Farmers' Bul. 940, 27 p., illus.
- (22) DUSHAM, E. H.
1921. THE PAINTED HICKORY BORER. N. Y. Cornell Agr. Expt. Sta. Bul. 407, p. 173-203, illus.
- (23) ESSIG, E. O.
1926. INSECTS OF WESTERN NORTH AMERICA. . . 1035 p., illus., New York.
- (24) EVENDEN, J. C.
1926. THE PINE BUTTERFLY, *NEOPHASIA MENAPHIA* FELDER. Jour. Agr. Research 33: 339-344, illus.
- (25) FELT, E. P.
1905. INSECTS AFFECTING PARK AND WOODLAND TREES. N. Y. State Mus. Mem. 8, 2 v., illus.
- (26) ———
1917. KEY TO AMERICAN INSECT GALLS. N. Y. State Mus. Bul. 200, 310 p., illus.
- (27) ——— and JOUTEL, L. H.
1904. MONOGRAPH OF THE GENUS *SAPERDA*. N. Y. State Mus. Bul. 74. Ent. 20, 86 p., illus.
- (28) FRACKER, S. B., and GRANOVSKY, A. A.
1928. AIRPLANE DUSTING TO CONTROL THE HEMLOCK SPANWORM. Jour. Forestry 26: 12-33, illus.
- (29) GARMAN, H.
1915. THE LOCUST BORER (*CYLLENE ROBINIAE*) AND OTHER INSECT ENEMIES OF THE BLACK LOCUST. Ky. State Forester Bien. Rpt. 2: [32]-63. [Also Ky. Agr. Expt. Sta. Bul. 200, p. 99-135, illus. 1915.]
- (30) GRAHAM, S. A.
1922. THE RED TURPENTINE BEETLE IN ITASCA PARK. Minn. State. Ent. Rpt. 19: [15]-21.
- (31) ———
1925. TWO DANGEROUS DEFOLIATORS OF JACK PINE. Jour. Econ. Ent. 18: 337-345.
- (32) ———
1926. BIOLOGY AND CONTROL OF THE WHITE-PINE WEEVIL, *PISSODES STROBI* PECK. N. Y. Cornell Agr. Expt. Sta. Bul. 449, 32 p., illus.
- (33) ——— and BAUMHOFER, L. G.
1927. THE PINE-TIP MOTH IN THE NEBRASKA NATIONAL FOREST. Jour. Agr. Research 35: 323-333, illus.
- (34) GREENE, C. T.
1917. TWO NEW CAMBIUM MINERS (DIPTERA). Jour. Agr. Research 10: 313-318, illus.
- (35) HERBERT, F. B.
1920. CYPRESS BARK SCALE. U. S. Dept. Agr. Bul. 838, 22 p., illus.

- (36) HEWITT, C. G.
1912. THE LARGE LARCH SAWFLY [NEMATUS ERICHSONII] WITH AN ACCOUNT OF ITS PARASITES, OTHER NATURAL ENEMIES AND MEANS OF CONTROL. Canada Dept. Agr. Expt. Farms Div. Ent. Bul. 10, 2d ser., Ent. Bul. 5, 42 p., illus.
- (37) HIGH, M. M.
1915. THE HUISACHE GIRDLER. U. S. Dept. Agr. Bul. 184, 9 p., illus.
- (38) HOFER, G.
1920. THE ASPEN BORER AND HOW TO CONTROL IT. U. S. Dept. Agr. Farmers' Bul. 1154, 11 p., illus.
- (39) HOPKINS, A. D.
1904. INSECT INJURIES TO HARDWOOD FOREST TREES. U. S. Dept. Agr. Yearbook 1903: 313-328, illus.
- (40) ———
1904. CATALOGUE OF EXHIBITS OF INSECT ENEMIES OF FORESTS AND FOREST PRODUCTS AT THE LOUISIANA PURCHASE EXPOSITION, ST. LOUIS., MO., 1904. U. S. Dept. Agr., Div. Ent. Bul. 48, 56 p., illus.
- (41) ———
1907. THE LOCUST BORER AND METHODS FOR ITS CONTROL. U. S. Dept. Agr., Bur. Ent. Circ. 83, 8 p., illus.
- (42) ———
1909. PRACTICAL INFORMATION ON THE SCOLYTID BEETLES OF NORTH AMERICAN FORESTS. I. BARKBEETLES OF THE GENUS DENDROCTONUS. U. S. Dept. Agr., Bur. Ent. Bul. 83, pt. 1, 169 p., illus.
- (43) ———
1912. THE DYING HICKORY TREES: CAUSE AND REMEDY. U. S. Dept. Agr., Bur. Ent. Circ. 144, 5 p., illus.
- (44) HOUSER, J. S.
1918. DESTRUCTIVE INSECTS AFFECTING OHIO SHADE AND FOREST TREES. Ohio Agr. Expt. Sta. Bul. 332, p. 161-487, illus.
- (45) HOWARD, L. O., and CHITTENDEN, F. H.
1916. THE LEOPARD MOTH: A DANGEROUS IMPORTED INSECT ENEMY OF SHADE TREES. U. S. Dept. Agr. Farmers' Bul. 708, 12 p., illus.
- (46) KEEN, F. P.
1929. INSECT ENEMIES OF CALIFORNIA PINES AND THEIR CONTROL. Calif. State Dept. Nat. Resources, Forestry Div. Bul. 7, 113 p., illus.
- (47) KORSTIAN, C. F.
1927. FACTORS CONTROLLING GERMINATION AND EARLY SURVIVAL IN OAKS. Yale Univ. School Forestry Bul. 19, 115 p., illus.
- (48) KOTINSKY, J.
1921. INSECTS INJURIOUS TO DECIDUOUS SHADE TREES AND THEIR CONTROL. U. S. Dept. Agr. Farmers' Bul. 1169, 100 p., illus.
- (49) MIDDLETON, W.
1921. LECONTE'S SAWFLY, AN ENEMY OF YOUNG PINES. Jour. Agr. Research 20: 741-760, illus.
- (50) MILLER, J. M.
1914. INSECT DAMAGE TO THE CONES AND SEEDS OF PACIFIC COAST CONIFERS. U. S. Dept. Agr. Bul. 95, 7 p., illus.
- (51) ———
1915. CONE BEETLES: INJURY TO SUGAR PINE AND WESTERN YELLOW PINE. U. S. Dept. Agr. Bul. 243, 12 p., illus.
- (52) MILLIKEN, F. B.
1916. THE COTTONWOOD BORER. U. S. Dept. Agr. Bul. 424, 7 p., illus.
- (53) PACKARD, A. S.
1890. THE FIFTH REPORT OF THE UNITED STATES ENTOMOLOGICAL COMMISSION. . . 957 p., illus. Washington, [D. C.]
- (54) PATTERSON, J. E.
1921. THE LIFE HISTORY OF RECURVARIA MILLERI, THE LODGEPOLE PINE NEEDLE-MINER, IN THE YOSEMITE NATIONAL PARK, CALIFORNIA. Jour. Agr. Research 21: 127-142, illus.
- (55) ———
1923. PINE DEFOLIATION ON THE KLAMATH RESERVATION. Timberman 24 (8): 39, illus.

- (56) PEIRSON, H. B.
1921. THE LIFE HISTORY AND CONTROL OF THE PALES WEEVIL (*HYLOBIUS PALES*). Harvard Forest Bul. 3. 33 p., illus.
- (57) ———
1927. CONTROL OF THE BRONZE BIRCH BORER BY FOREST MANAGEMENT. Jour. Forestry 25: 68-72.
- (58) ST. GEORGE, R. A.
1925. THE RECENT DEATH OF LARGE QUANTITIES OF SOUTHERN PINES. Amer. Lumberman (whole No.) 2607: 50-51, illus.
- (59) SLINGERLAND, M. V.
1906. THE BRONZE BIRCH BORER: AN INSECT DESTROYING THE WHITE BIRCH. N. Y. Cornell Agr. Expt. Sta. Bul. 234, p. [65]-78, illus.
- (60) SNYDER, T. E.
1915. INSECTS INJURIOUS TO FORESTS AND FOREST PRODUCTS. BIOLOGY OF THE TERMITES OF THE EASTERN UNITED STATES, WITH PREVENTIVE AND REMEDIAL MEASURES. U. S. Dept. Agr., Bur. Ent. Bul. 94. pt. 2. 95 p., illus.
- (61) ———
1926. PREVENTING DAMAGE BY LYCTUS POWDER-POST BEETLES. U. S. Dept. Agr. Farmers' Bul. 1477, 12 p., illus.
- (62) ———
1927. DEFECTS IN TIMBER CAUSED BY INSECTS. U. S. Dept. Agr. Bul. 1490. 47 p., illus.
- (63) SWAINE, J. M.
1918. CANADIAN BARK BEETLES. II. A PRELIMINARY CLASSIFICATION WITH AN ACCOUNT OF THE HABITS AND MEANS OF CONTROL. Canada Dept. Agr. Ent. Branch Bul. 14 (pt. 2), 143 p., illus.
- (64) ——— CRAIGHEAD, F. C., and BAILEY, I. W.
1924. STUDIES ON THE SPRUCE BUDWORM. (*CACOEZIA FUMIFERANA CLEM.*). Canada Dept. Agr. Ent. Branch Bul. (n. s.) 37, 91 p., illus.
- (65) SWENK, M. H.
1927. THE PINE TIP MOTH IN THE NEBRASKA NATIONAL FOREST. Nebr. Agr. Expt. Sta. Research Bul. 40. 50 p., illus.
- (66) TRIMBLE, F. M.
1924. LIFE HISTORY AND HABITS OF TWO PACIFIC COAST BARK BEETLES. Ann. Ent. Soc. Amer. 17: 382-[391], illus.
- (67) WEBB, J. L.
1909. SOME INSECTS INJURIOUS TO FORESTS. IV. THE SOUTHERN PINE SAWYER. U. S. Dept. Agr., Bur. Ent. Bul. 58. pt. 4, p. 41-56, illus.
- (68) ———
1911. INJURIES TO FORESTS AND FOREST PRODUCTS BY ROUNDHEADED BOREES. U. S. Dept. Agr. Yearbook 1910: 341-358, illus.

ADDENDA

Most of the papers cited below were published subsequent to the preparation of this one.

- (69) ALLISON, J. H., and ORR, L. W.
1929. A NEW MENACE TO SCOTCH AND JACK PINE. Jour. For. 27: 821-824.
- (70) ANNAND, P. N.
1928. A CONTRIBUTION TOWARD A MONOGRAPH OF THE ADELGINAE (PHYLLOXERIDAE) OF NORTH AMERICA. Stanford Univ. Pubs., Biol. Sci. vol. 6, no. 1, 146 p.
- (71) BUSCK, A.
1915. THE EUROPEAN PINE-SHOOT MOTH; A SERIOUS MENACE TO PINE TIMBER IN AMERICA. U. S. Dept. Agr. Bul. 170, 11 p., illus.
- (72) DIVISION OF FOREST INSECT INVESTIGATIONS, BUREAU OF ENTOMOLOGY.
1927. THE RELATION OF INSECTS TO SLASH DISPOSAL. U. S. Dept. Agr. Dept. Circ. 411, 12 p.
- (73) FRIEND, R. B.
1926. THE SPRUCE GALL APHID (*ADELGES ABIETIS* L.) AND ITS CONTROL. Conn. Agr. Expt. Sta. Rpt. Bul. 285, p. 223-228.
- (74) ———
1927. THE BIOLOGY OF THE BIRCH LEAF SKELETONIZER *BUCCALATRIX CANADENSIS*ELLA, CHAMBERS. Conn. Agr. Expt. Sta. Bul. 288: [395]-486, illus.

- (75) KEEN, F. P.
1929. TUSsock Moth Menace. *Timberman*, 30 (11) : 108.
- (76) MACALONEY, H. J.
1930. THE WHITE PINE WEEVIL (*PISSODES STROBI* PECK)—ITS BIOLOGY AND CONTROL. *N. Y. State Col. For. Bul.*, vol. 3, no. 1, 87 p., illus.
- (77) MARLATT, C. L.
1928. HOUSE ANTS: KINDS AND METHODS OF CONTROL. *U. S. Dept. Agr. Farmers' Bul.* 740 (Rev.) 14 p., illus.
- (78) MILLER, J. M.
1929. THE RELATION OF WINDFALLS TO BARK-BEETLE EPIDEMICS. *Fourth Intern. Cong. Ent.* 1928, *Trans.* vol. 2, p. [992]-1002.
- (79) ——— and PATTERSON, J. E.
1927. PRELIMINARY STUDIES ON THE RELATION OF FIRE INJURY TO BARK-BEETLE ATTACK IN WESTERN YELLOW PINE. *Jour. Agr. Research*, 34 : 597-613, illus.
- (80) PATCH, E. M.
1909. CHERMES OF MAINE CONIFERS. *Maine Agr. Expt. Sta. Bul.* 173 : [277]-308, illus.
- (81) PATTERSON, J. E.
1929. THE PANDORA Moth, A PERIODIC PEST OF WESTERN PINE FORESTS. *U. S. Dept. Agr. Tech. Bul.* 137, 19 p., illus.
- (82) PEIRSON, H. B.
1927. MANUAL OF FOREST INSECTS. *Maine For. Ser. Bul.* 5, 130 p., illus.
- (83) ———
1929. OBSERVATIONS ON THE BIRCH LEAF MINING SAWFLY. *Jour. Econ. Ent.* 22 : 588-594.
- (84) ST. GEORGE, R. A.
1929. PROTECTION OF LOG CABINS, RUSTIC WORK, AND UNSEASONED WOOD FROM INJURIOUS INSECTS. *U. S. Dept. Agr. Farmers' Bul.* 1582, 19 p., illus.
- (85) ——— and BEAL, J. A.
1929. THE SOUTHERN PINE BEETLE: A SERIOUS ENEMY OF PINES IN THE SOUTH. *U. S. Dept. Agr. Farmers' Bul.* 1586, 18 p., illus.
- (86) SNYDER, T. E.
1910. DAMAGE TO CHESTNUT TELEPHONE AND TELEGRAPH POLES BY WOOD-BORING INSECTS. *U. S. Dept. Agr. Bur. Ent. Bul.* 94, pt. 1, 12 p., illus.
- (87) ———
1929. TERMITES IN BUILDINGS. *U. S. Dept. Agr. Leaflet* 31, 5 p., illus.
- (88) ——— and ST. GEORGE, R. A.
1924. DETERMINATION OF TEMPERATURES FATAL TO THE POWDER-POST BEETLE, *LYCTUS PLANICOLLIS* LECONTE, BY STEAMING INFESTED ASH AND OAK LUMBER IN A KILN. *Jour. Agr. Research*, 28 : 1033-1038, illus.
- (89) SWAINE, J. M.
1924. THE CONTROL OF THE DESTRUCTIVE SPRUCE BARK BEETLE IN EASTERN CANADA. *Canada Dept. Agr. Ent. Br. Pamphlet* 48 n. s., 20 p., illus.

APPENDIX

These additional references should be consulted under the species mentioned.

The southern pine beetle (85); the eastern spruce beetle (89); the engraver beetles (85); the Jeffrey pine beetle (78); the black horned pine borer (84); the belted chion (84); the cedar tree borer (84); the pole borer (86); the powder post beetle (88); the red shouldered powder post beetle (84); termites (87); the spruce twig gall lice (70, 73, 80); the pine bark louse (80); the white pine weevil (76); the European pine shoot moth (71); the pine soft scales (69); the birch leaf skeletonizer (74); the larch case bearer (82); the Pandora moth (81); the Douglas fir tussock moth (75); the birch leaf miner (83).

INDEX

	Page		Page
Acorn gall makers.....	16	Douglas fir beetle.....	5
Adelges abietis.....	15	Douglas fir chalcid.....	16
Adelges cooleyi.....	15	Douglas fir looper.....	19-20
Agrilus anxius.....	6	Douglas fir pitch moth.....	10
Agrilus bilineatus.....	6	Douglas fir tussock moth.....	20
Agromyza aceris.....	10	Eastern hemlock bark borer.....	8
Agromyza amelanchieris.....	10	Eastern spruce beetle.....	4
Ambrosia beetles.....	11	Eastern white pine cone beetle.....	15
Anisota rubicunda.....	18	Ehrhornia cupressi.....	17
Anisota senatoria.....	18	Elaphidion villosum.....	7
Anisota stigma.....	18	Elloplia fuscicollis.....	19
Anisota virginiana.....	18	Elloplia somnaria.....	19
Apatides fortis.....	14	Elloplia spp.....	19
Argyrotaenia pinatubana.....	20	Elm borer.....	9
Banded ash borer.....	13	Engelmann spruce beetle.....	20
Belted chion.....	12	Engraver beetles.....	5-6
Birch leaf miner.....	23	Eupsalis minuta.....	11
Birch leaf skeletonizer.....	18	European pine shoot moth.....	17
Black Hills beetle.....	5	Fir bark beetle.....	6
Black-horned pine borer.....	12	Flat oak borer.....	14
Black turpentine beetle.....	4	Galenara sp.....	19-20
Bronze birch borer.....	6	(Gillettea) Adelges cooleyi.....	15
Brown-tail moth.....	20	Gipsy moth.....	21
Bucculatrix canadensisella.....	18	Glycobius speciosus.....	7
Buprestis apricans.....	7	Goes pulcher.....	7
Cacoecia fumiferana.....	18	Goes pulverulenta.....	7
California oak worm.....	21	Goes tessellata.....	7
Callidium antennatum.....	12	Goes tigrina.....	7
Cambium miners.....	10	Grape colaspis.....	18
Camponotus herculeanus pennsylvanicus.....	14	Gray ash borer.....	13
Carpenter ant.....	18	Green-striped maple worm.....	14
Carpenter worm.....	10	Hemerocampa pseudotsugata.....	20
Catalpa sphinx.....	19	Hemicallidium amethystinum.....	12
Cedar pole borer.....	9	Hemlock bark maggot.....	11
Cedar-tree borer.....	12	Hemlock budworm.....	21
Ceratomia catalpae.....	19	Hemlock loopers.....	19
Chalepus dorsalis.....	18	Heterocampa bilineata.....	20
Cheilosia alaskensis.....	11	Heterocampa biundata.....	20
(Chermes) Adelges abietis.....	15	Heterocampa guttivitta.....	20
(Chermes) Adelges cooleyi.....	15	Heterocampa mantee.....	20
(Chermes) Pinus pinicorticis.....	16	Heterocampa spp.....	20
Chestnut timber worm.....	11	Hickory bark beetle.....	6
Chionaspis pinifoliae.....	17	Hickory twig girdler.....	8
Chion cinctus.....	12	Huisache girdler.....	8
Colaspis brunnea.....	18	Hylecoetus lugubris.....	11
Coleophora laricella.....	19	Hyllobius pales.....	16
Coloradia pandora.....	19	Hylotrupes bajulus.....	12
Columbian timber beetle.....	11	(Hylotrupes) Hemicallidium amethystinum.....	12
Cone beetles.....	15	Hylotrupes ligneus.....	12
Conophthorus coniperda.....	15	Ips avulsus.....	6
Conophthorus lambertianae.....	15	Ips calligraphus.....	6
Conophthorus ponderosae.....	15	Ips confusus.....	6
Conophthorus spp.....	15	Ips emarginatus.....	6
Corythilus columbianus.....	11	Ips grandicollis.....	6
Cottonwood borer.....	8	Ips integer.....	6
Curculio spp.....	15	Ips spp.....	5-6
Cyllene caryae.....	12	Jeffrey pine beetle.....	5
Cyllene robiniae.....	7	Kaloterms spp.....	14-15
Cynipidae.....	16	Larch case-bearer.....	19
Cypress bark scale.....	17	Larch sawfly.....	22
Datana integerrima.....	19	Lead-cable borer.....	14
Datana ministra.....	19	Leopard moth.....	10
Dendrobiella aspera.....	14	Linden borer.....	9
Dendroctonus barberi.....	5	Living beech borer.....	7
Dendroctonus brevicornis.....	5	Living hickory borer.....	7
Dendroctonus engelmanni.....	20	Locust borer.....	7
Dendroctonus frontalis.....	4	Locust leaf miner.....	18
Dendroctonus jeffreyi.....	5	Lodgepole needle tier.....	20
Dendroctonus monticolae.....	4	Lodgepole sawfly.....	20, 22
Dendroctonus piceaperda.....	4	Long-horned girdlers.....	8
Dendroctonus ponderosae.....	5	Lycetus planicollis.....	14
Dendroctonus pseudotsugae.....	5	Lycetus spp.....	14
Dendroctonus terebrans.....	4	(Lymexylon) Melittomma sericeum.....	11
Dendroctonus valens.....	4	Malacosoma americana.....	21
Dioryctria abietella.....	9	Malacosoma californica.....	21
Dioryctria amatella.....	9	Malacosoma constricta.....	21
Dioryctria ponderosae.....	9	Malacosoma dissita.....	21
Dioryctria xanthanobarea.....	9	Malacosoma fragilis.....	21

	Page		Page
Malacosoma pluvialis	21	Powder-post beetles (see also red-shouldered powder-post beetle)	14
Maple defoliators	20	Prionid root borers	8
May beetles	16	Prionoxystus robiniae	10
Megastigmus albifrons	16	Prionus californicus	8
Megastigmus milleri	16	Prionus imbricornis	8
Megastigmus piceae	16	Prionus laticollis	8
Megastigmus pinus	16	Prionus spp.	8
Megastigmus spermatophus	16	Recurvaria milleri	22
Melanophila fulvoguttata	8	Red-headed ash borer	13
Melanophila gentilis	8	Red-headed pine sawfly	23
Melittotomma sericeum	11	Red oak borer	8
Monochamus confusus	13	Red-shouldered powder-post beetle	14
Monochamus maculosus	13	Red turpentine beetle	4
Monochamus marmorator	13	Reticulitermes spp.	15
Monochamus scutellatus	13	Rhyacionia buoliana	17
Monochamus spp.	13	Rhyacionia frustrana	17
Monochamus titillator	13	Romaleum rufulum	8
Monterey cypress bark beetle	6	Saperda calcarata	9
Mountain pine beetle	4	Saperda tridentata	9
Nantucket pine moth	17	Saperda vestita	9
Nematus erichsonii	22	Sapwood timber worm	11
Neoclytus capraea	13	Satin moth	22
Neoclytus conjunctus	13	Sawflies	22
Neoclytus erythrocephalus	13	Scobicia declivis	14
Neodiprion banksianae	22	Scolytus quadrispinosus	6
Neodiprion excitans	22	Scolytus ventralis	6
Neodiprion lecontei	22	Seed chalcids	16
Neodiprion scutellaris	22	Sequoia pitch moth	10
Neodiprion spp.	20, 22	Small metallic wood and bark borers	6
Neophasia menapia	21	Smidicium cucujiforme	14
Nut weevils	15	Southern pine beetle	4
Nygmia phaeorrhoea	20	Southwestern pine beetle	5
Oak and maple defoliators	20	Spiny oak worm	18
Oak sapling borer	7	Spruce bud worm	18
Oak timber worm	11	Spruce twig gall lice	15
Oak twig pruner	7	Stilpnotia salicis	22
Oberea bimaculata	8	Sugar maple borer	7
Oberea myops	8	Sugar pine cone beetle	15
Oberea schaumii	8	Synanthedon novaeorleansis	0
Oberea spp.	8	Tanbark borer	13
Oberea tripunctata	8	Tent caterpillars	21
Old-house borer	12	Termites (nonsubterranean)	14-15
Oncideres cingulata	8	Termites (subterranean)	15
Oncideres putator	8	Tetropium abietis	9
Orange-striped oak worm	18	Tetropium velutinum	9
Painted hickory borer	12	Toumeyella numismaticum	17-18
Pales weevil	16	Toumeyella parvicorne	17
Pandora moth	19	Toumeyella pini	17
Parandra brunnea	13	Toumeyella spp.	17
Peronea variana	21	Trachykele blondell	9
Phloeosinus cristatus	6	Turpentine beetles	4
Phryganidia californica	21	Turpentine borer	7
Phyllophaga spp.	16	Twig tunnelers	8
Phyllotoma nemorata	23	Two-lined chestnut borer	6
Phymatodes variabilis	13	Vespa mima sequoiae	10
Pine bark louse	16	Walnut caterpillar	19
Pine butterfly	21	Western cedar bark borer	12
Pine needle miner	22	Western fir borer	9
Pine needle scale	17	Western flat-headed borer	8
Pine sawyers	13	Western larch bark borer	9
Pine soft scales	17-18	Western pine beetle	5
Pineus pinicorticis	16	White grubs	16
Pinhole borers	11	White oak borer	7
Pinipestis zimmermani	10	White pine weevil	16-17
Pissodes strobi	16-17	Xylobiops basilaris	14
Pitch moths. (See Dioryctria spp., Douglas fir pitch moth, and sequoia, pitch moth.)		Yellow-necked caterpillar	19
Plectrodera scalator	8	Yellow pine cone beetle	15
Pole borer	13	Zeuzera pyrina	10
Poplar borer	9	Zimmerman pine moth	10
Porthetria dispar	21		

ORGANIZATION OF THE UNITED STATES DEPARTMENT OF AGRICULTURE

May 1, 1930

<i>Secretary of Agriculture</i>	ARTHUR M. HYDE.
<i>Assistant Secretary</i>	R. W. DUNLAP.
<i>Director of Scientific Work</i>	A. F. WOODS.
<i>Director of Regulatory Work</i>	WALTER G. CAMPBELL.
<i>Director of Extension Work</i>	C. W. WARBURTON.
<i>Director of Personnel and Business Administration.</i>	W. W. STOCKBERGER.
<i>Director of Information</i>	M. S. EISENHOWER.
<i>Solicitor</i>	E. L. MARSHALL.
<i>Weather Bureau</i>	CHARLES F. MARVIN, <i>Chief.</i>
<i>Bureau of Animal Industry</i>	JOHN R. MOHLER, <i>Chief.</i>
<i>Bureau of Dairy Industry</i>	O. E. REED, <i>Chief.</i>
<i>Bureau of Plant Industry</i>	WILLIAM A. TAYLOR, <i>Chief.</i>
<i>Forest Service</i>	R. Y. STUART, <i>Chief.</i>
<i>Bureau of Chemistry and Soils</i>	H. G. KNIGHT, <i>Chief.</i>
<i>Bureau of Entomology</i>	C. L. MARLATT, <i>Chief.</i>
<i>Bureau of Biological Survey</i>	PAUL G. REDINGTON, <i>Chief.</i>
<i>Bureau of Public Roads</i>	THOMAS H. MACDONALD, <i>Chief.</i>
<i>Bureau of Agricultural Economics</i>	NILS A. OLSEN, <i>Chief.</i>
<i>Bureau of Home Economics</i>	LOUISE STANLEY, <i>Chief.</i>
<i>Plant Quarantine and Control Administration</i> ..	LEE A. STRONG, <i>Chief.</i>
<i>Grain Futures Administration</i>	J. W. T. DUVEL, <i>Chief.</i>
<i>Food, Drug, and Insecticide Administration</i> ..	WALTER G. CAMPBELL, <i>Director of Regulatory Work, in Charge.</i>
<i>Office of Experiment Stations</i>	_____, <i>Chief.</i>
<i>Office of Cooperative Extension Work</i>	C. B. SMITH, <i>Chief.</i>
<i>Library</i>	CLARIBEL R. BARNETT, <i>Librarian.</i>

This publication is a contribution from

<i>Bureau of Entomology</i>	C. L. MARLATT, <i>Chief.</i>
<i>Division of Forest Insects</i>	F. C. CRAIGHEAD, <i>Principal Entomologist, in Charge.</i>

U. S. DEPARTMENT OF AGRICULTURE,

BUREAU OF ENTOMOLOGY—BULLETIN No. 58, Part V.

L. O. HOWARD, Entomologist and Chief of Bureau.

SOME INSECTS INJURIOUS TO FORESTS.

INSECT DEPREDACTIONS IN NORTH AMERICAN FORESTS

AND PRACTICAL METHODS OF PREVENTION
AND CONTROL.

BY

A. D. HOPKINS, PH. D.,

In Charge of Forest Insect Investigations.

ISSUED DECEMBER 4, 1909.

WASHINGTON:

GOVERNMENT PRINTING OFFICE.

1909.

BUREAU OF ENTOMOLOGY.^a

L. O. HOWARD, *Entomologist and Chief of Bureau.*

C. L. MARLATT, *Assistant Entomologist and Acting Chief in Absence of Chief.*

R. S. CLIFTON, *Executive Assistant.*

CHAS. J. GILLISS, *Chief Clerk.*

F. H. CHITTENDEN, *in charge of truck crop and stored product insect investigations.*

A. D. HOPKINS, *in charge of forest insect investigations.*

W. D. HUNTER, *in charge of southern field crop insect investigations.*

F. M. WEBSTER, *in charge of cereal and forage insect investigations.*

A. L. QUAINANCE, *in charge of deciduous fruit insect investigations.*

E. F. PHILLIPS, *in charge of bee culture.*

D. M. ROGERS, *in charge of preventing spread of moths, field work.*

ROLLA P. CURRIE, *in charge of editorial work.*

MABEL COLCORD, *librarian.*

FOREST INSECT INVESTIGATIONS.

A. D. HOPKINS, *in charge.*

H. E. BURKE, J. L. WEBB, JOSEF BRUNNER, S. A. ROHWER, T. E. SNYDER, W. N. DOYENER, *agents and experts.*

MARY E. FAUNCE, *preparator.*

WILLIAM MIDDLETON, *student assistant.*

W. D. EDMONSTON, *forest ranger, detailed from Forest Service.*

^a Organization of the Bureau on November 1, 1909.

LETTER OF TRANSMITTAL.

U. S. DEPARTMENT OF AGRICULTURE,
BUREAU OF ENTOMOLOGY,
Washington, D. C., August 23, 1909.

SIR: I have the honor to transmit herewith, and to recommend for publication as Bulletin No. 58, Part V, of the Bureau of Entomology, manuscript of a paper by Dr. A. D. Hopkins, in charge of forest insect investigations of this Bureau, entitled "Insect Depredations in North American Forests and Practical Methods of Prevention and Control."

The paper is based largely on the observations and experience of the author and of the assistants engaged in forest insect investigations, and contains a list of publications relating to forest insects.

Respectfully,

L. O. HOWARD,
Entomologist and Chief of Bureau.

HON. JAMES WILSON,
Secretary of Agriculture.

CONTENTS.

	Page.
Introduction	57
Insect depredations in North American forests.....	57
Character and extent of depredations.....	57
Insects cause the death of trees.....	58
The southern pine beetle.....	58
The eastern spruce beetle.....	58
The Engelmann spruce beetle.....	59
The Black Hills beetle.....	59
The mountain pine beetle and the western pine beetle.....	59
The Douglas fir beetle.....	59
The hickory barkbeetle.....	60
The larch worm.....	60
Insect injuries to the wood of living trees.....	60
The oak timber worm.....	60
The chestnut timber worm.....	60
Carpenter worms.....	61
Ambrosia beetles.....	61
The locust borer.....	61
Turpentine beetles and turpentine borers.....	61
The white pine weevil.....	62
Insect injuries to the wood of dying and dead trees.....	62
Coniferous trees.....	63
Sawyers.....	63
Ambrosia beetles.....	63
Pinhole borers in cypress.....	63
Hardwood trees.....	64
Round-headed borers, timber worms, and ambrosia beetles..	64
Insect injuries to forest products.....	64
Crude products.....	64
Round-headed borers, timber worms, and ambrosia beetles..	64
Manufactured unseasoned products.....	65
Ambrosia beetles and other wood borers.....	65
Seasoned products in yards and storehouses.....	66
Powder-post beetles.....	66
Finished products.....	66
Utilized products.....	66
Powder-post beetles, white ants, and other wood-boring in-	
sects.....	66
Insects in their relation to the reduction of future supplies of timber....	67
Interrelations of forest insects and forest fires.....	67
Losses from forest insects.....	67
Insect-killed timber as fuel for fires.....	68
Fire-killed timber injured by insects.....	68

Insect depredations in North American forests—Continued.

Interrelations of forest insects and forest fires—Continued.	Page.
Destruction of insects by fire	68
Durability of insect-killed timber.....	69
Interrelation of forest insects and forest fungi	69
Decay following injury by insects	69
Summary and estimates relating to character and extent of insect damage.	69
Standing timber killed and damaged by insects.....	70
Reduction in the Nation's wealth.....	71
Reduction in cash revenue.....	71
Reduction in value of finished and commercial products	71
Methods of prevention and control	71
General principles of control	72
Control of barkbeetles which kill trees.....	73
Maintaining control of barkbeetles.....	74
Examples of successful control of barkbeetles.....	75
Control of insects which cause defects in living timber.....	78
Prevention of injury to dying and dead trees	78
Prevention of loss from insect injuries to natural and artificial reproduction.	79
Utilization of immune and resistant varieties and races of trees.....	79
Prevention of insect injuries to forest products.....	79
Crude products.....	79
Manufactured products	81
Unseasoned products	81
Seasoned products.....	81
Past and present conditions of powder-post injury.....	82
Tan bark	83
Utilized products.....	83
Timbers and woodwork in structures	83
Utilization of natural enemies and factors in the control of injurious insects.....	85
Beneficial insects.....	86
Beneficial diseases of insects	86
Beneficial birds	86
Beneficial climatic conditions	87
Utilization of waste caused by insects.....	87
Present conditions and opportunities	88
Forest entomology as applied to American forests	88
Present knowledge	88
Needs	89
Elementary and technical knowledge of forest entomology for the forester.....	91
Present requirements of instruction.....	91
Conclusion.....	91
General estimates of amount of damage caused by forest insects.....	92
How losses can be prevented.....	93
Publications relating to forest insects	96
Historical and general	96
Prepared by officials of the branch of Forest Insect Investigations, Bureau of Entomology, and of the West Virginia Agricultural Experiment Station	97
Publications relating to forest statistics.....	100
Forest resources.....	100
Forest fires	101

SOME INSECTS INJURIOUS TO FORESTS.

INSECT DEPREDACTIONS IN NORTH AMERICAN FORESTS AND PRACTICAL METHODS OF PREVENTION AND CONTROL.

INTRODUCTION.

It is the purpose of this part of the bulletin to give a summary of facts, conclusions, and estimates relating to the forest-insect problem as applied to North American conditions and to call attention to its importance in the future management of private and public forests.

The matter is presented in as brief and concise a form as possible, in order that the information may be readily available to the general reader, as well as to the forester and student, and references are made to publications in which more detailed accounts may be found.

The statements and conclusions relating to the insects and their work and to methods for their control are based almost entirely on investigations and observations by the writer and by assistants in the Bureau of Entomology working under his direction, carried on in all of the principal forest areas of the United States. The historical data have been gathered from the publications listed on pages 96-97. The estimates of the amount and value of standing timber killed by insects and of the forest products destroyed and reduced in value through insect injuries to the crude and seasoned products are based on our published results of investigations (pp. 97-100), on unpublished notes, on technical and practical knowledge of the subject, and on the published forest statistics relating to the amount and value of timber, fire losses, etc. (pp. 100-101).

INSECT DEPREDACTIONS IN NORTH AMERICAN FORESTS.

CHARACTER AND EXTENT OF DEPREDACTIONS.

The records of notable depredations by insects on the timber supply of Europe during the past four hundred years, on that of the

United States during the past century, and especially those that have come under the observation of the writer and assistants engaged in forest insect investigations during the past eighteen years, furnish conclusive evidence that this class of enemies has been, and is now, an important factor in the destruction and waste of forest resources.

INSECTS CAUSE THE DEATH OF TREES.

It has been conclusively demonstrated that certain species of insects are the direct or primary cause of the death of forest trees of all ages, and that from time to time they multiply to such an alarming extent that their depredations assume the character of a destructive invasion, which results in the death of a large percentage of the best timber over thousands of square miles.

There are many species of barkbeetles which prefer to attack matured and healthy trees, and there are many examples of whole forests of century-old trees having perished from the girdling effect of the mines of the beetles, which are extended in all directions through the inner living bark on the main trunks of the trees. Indeed, we find among these bark-boring beetles the most destructive insect enemies of North American forests. Some notable examples of the depredations of these barkbeetles are given below.

The southern pine beetle.—In 1890–1892 a destructive invasion of the southern pine beetle (*Dendroctonus frontalis* Zimm.) extended from the western border of West Virginia through Maryland and Virginia into the District of Columbia, northward into southern Pennsylvania, and southward into North Carolina. In this area, aggregating over 75,000 square miles, a very large percentage of the mature and small trees of the various species of pine and spruce was killed by this beetle. In many places in West Virginia and Virginia nearly all the pine trees of all sizes on thousands of acres were killed, while shade and ornamental trees within the same area suffered the same as those in the forest. Since 1902 this barkbeetle has been more or less active in the Southern States from Virginia to Texas, and in some localities and during certain years it has killed a large amount of timber. Records of extensive destruction of timber in the Southern States are found dating back to the early part of the nineteenth century (Wilson, 1831). This species may be considered one of the most dangerous insect enemies of southeastern conifers and, therefore, a constant menace to the pine forests of the Southern States. (Hopkins, 1899*b*, 1903*b*, 1909*b*.)

The eastern spruce beetle.—During the period between 1818 and 1900 there were several outbreaks of the eastern spruce beetle (*Dendroctonus piceaperda* Hopk.) in the spruce forests of New York, New England, and southeastern Canada (Peck, 1876, 1878; Hough, 1882; Packard, 1890; Pinchot, 1899, p. 74; Hopkins, 1901*a*, 1909*b*).

This species caused the death of a very large percentage of the mature spruce over an area of thousands of square miles. In the aggregate many billions of feet of the best timber were destroyed. The larger areas of this dead timber furnished fuel for devastating forest fires, with the result that in most cases there was a total loss.

The Engelmann spruce beetle.—Another barkbeetle (*Dendroctonus engelmanni* Hopk.), similar in habits to *piceaperda*, has from time to time during the past fifty years caused widespread devastations in the Rocky Mountains region to forests of Engelmann spruce, in some sections killing from 75 to 90 per cent of the timber of merchantable size. (Hopkins, 1908*a*, p. 161; 1909*b*, pp. 126–132.)

The Black Hills beetle.—One of the most striking examples of the destructive powers of an insect enemy of forest trees is found in the Black Hills National Forest of South Dakota, where during the past ten years a large percentage of the merchantable timber of the entire forest has been killed by the Black Hills beetle (*Dendroctonus ponderosæ* Hopk.). It is estimated that more than a billion feet of timber have been destroyed in this forest as the direct result of the work of this beetle. This destructive enemy of the western pine is distributed throughout the forests of the middle and southern Rocky Mountains region, where, within recent years, it has been found that in areas of greater or less extent from 10 to 80 per cent of the trees have been killed by it. (Hopkins, 1902*a*, 1903*b*, 1905*b*, 1908*a*, and 1909*b*, pp. 90–101.)

The mountain pine beetle and the western pine beetle.—The sugar pine, silver pine, western yellow pine, and lodgepole pine of the region north of Colorado and Utah, westward to the Cascades, and southward through the Sierra Nevadas are attacked by the mountain pine beetle (*Dendroctonus monticolæ* Hopk.) and the western pine beetle (*Dendroctonus brevicomis* Lec.), and, as a direct consequence, billions of feet of the timber have died. In one locality in north-eastern Oregon it is estimated that 90 to 95 per cent of the timber in a dense stand of lodgepole pine covering an area of 100,000 acres has been killed within the past three years by the mountain pine beetle. Throughout the sugar-pine districts of Oregon and California, as the result of attacks by this same destructive barkbeetle, a considerable percentage of the largest and best trees is dead. (Webb, 1906; Hopkins, 1908*a*, 1909*b*, pp. 80–90.)

The Douglas fir beetle.—The Douglas fir throughout the region of the Rocky Mountains from southern New Mexico to British Columbia has suffered severely from the ravages of the Douglas fir beetle (*Dendroctonus pseudotsugæ* Hopk.), with the result that a large percentage of dead timber is found, much of which will be a total loss. (Hopkins, 1909*b*, pp. 106–114.)

Three other species of beetles, having destructive habits similar to those above mentioned, depredate on the pines of New Mexico

and Arizona, and still another has contributed greatly to the destruction of the larch throughout the northeastern United States and southeastern Canada. (Hopkins, 1909*b*, pp. 49, 53, and 77.)

The hickory barkbeetle.—Within the past ten years the hickory barkbeetle (*Scolytus quadrispinosus* Say) has caused the destruction of an enormous amount of hickory timber throughout the northern tier of States from Wisconsin to Vermont and southward through the eastern Atlantic States and into the Southern States as far as central Georgia. (Hopkins, 1904*b*, pp. 314–317.)

The larch worm.—There are also many examples of widespread depredations chargeable to insects which defoliate the trees, thus contributing to their death. Notable among these are the depredations by the larch worm (*Nematus erichsonii* Hartig), which, during several extensive outbreaks since 1880, has killed from 50 to 100 per cent of the mature larch over vast areas in the northeastern United States and southeastern Canada. It is evident that the amount of merchantable-sized timber that has died as the result of defoliation by this insect will aggregate many billions of feet. (Packard, 1890, pp. 879–890; Pinchot, 1899; Hopkins, 1908*a*.)

INSECT INJURIES TO THE WOOD OF LIVING TREES.

It has been determined that insects of a certain class attack the wood and bark of living timber and that, while they do not contribute materially to the death of the trees or give much external evidence of their presence, they produce wounds in the bark and wormhole and pinhole defects in the wood which result in a depreciation in commercial value amounting to from 5 to 50 per cent. These defects in the wood are not detected until after the trees have been felled and the logs transported to the mill and converted into lumber. Thus to the actual damage to the lumber is added the expense of logging and manufacture of the defective, low-grade material, much of which must be discarded as worthless culls. (Hopkins, 1894*a*, 1894*b*, 1904*b*, 1905*a*, 1906*b*.)

The oak timber worm.—One of the most destructive of the class of depredators just mentioned is the oak timber worm (*Eupsalis minuta* Drn.). It enters the wood of the trunks of living trees through wounds in the bark and at the base of broken or dead branches and extends its "pinhole" burrows in all directions through the solid heartwood. The losses occasioned by this insect in the hardwood forests of the eastern United States are enormous and usually affect the wood of the finest examples of old trees. (Hopkins, 1894*a*, 1904*b*.)

The chestnut timber worm.—The chestnut throughout its range is damaged in a like manner by the chestnut timber worm (*Lymexylon sericeum* Harr.). Practically every tree of merchantable size is

more or less affected, and a large percentage is so seriously damaged that the product is reduced to that of the lowest grade. It is estimated that the reduction in value of the average lumber product at any given time is not far from 30 per cent, thus involving extensive waste and an increased drain on the forest to supply the requirements for clear lumber. This insect also attacks the oaks, and especially the red oak, the older trees of which are often as seriously damaged as are the chestnut. (Hopkins, 1894a, 1904b.)

Carpenter worms.—The oaks, especially the white oak and the red oak, are seriously damaged by carpenter worms of the genus *Prionoxystus*. The holes made by these insects through the heartwood of the best part of the trunks are sometimes 1.5 inches in diameter one way by 0.75 inch the other, thus causing serious damage to the wood. These, with other large wood-boring beetle larvæ, sometimes infest the top part of the trunk and the larger branches of oak trees, where their continued work results first in the dead and so-called "stag-horn" top and subsequently in broken, decayed, and worthless trunks. (Hopkins, 1894a, 1904b.)

Ambrosia beetles.—One of the commonest defects in white oak, rock oak, beech, whitewood or yellow poplar, elm, etc., is that known to the lumber trade as "grease spots," "patch worm," and "black holes." This defect is caused by one of the timber beetles or ambrosia beetles, *Corthylus columbianus* Hopk., which makes successive attacks in the living healthy sapwood from the time the trees are 20 or 30 years old until they reach the maximum age. Thus the black-hole and stained-wood defect is scattered all through the wood of the best part of the trunks of the trees. The average reduction in value of otherwise best-grade lumber amounts, in many localities, to from 25 to 75 per cent. The defect is commonly found in oak and elm furniture and in interior hardwood finish in dwellings and other buildings. (Hopkins, 1893g, 1894b, 1904b.)

The locust borer.—The locust, as is well known, suffers to such an extent from the ravages of the locust borer (*Cyrtene robinæ* Forst.) that in many localities the trees are rendered worthless for commercial purposes or they are reduced in value below the point of profitable growth as a forest tree, otherwise this would be one of the most profitable trees in the natural forest or artificial plantation and would contribute greatly to an increased timber supply. (Hopkins, 1906b, 1907a, 1907c.)

Turpentine beetles and turpentine borers.—While the softwood trees, or conifers, suffer far less than the hardwoods from the class of enemies which cause defects in the living timber there are a few notable examples of serious damage. There is a common trouble affecting the various species of pine throughout the country known as basal wounds or basal fire wounds. It has been found that a large

percentage of this injury to the pine in the States north and west of the Gulf States and in the Middle and South Atlantic States is caused by the red turpentine beetle (*Dendroctonus valens* Lec.) and in the Southern States by the black turpentine beetle (*Dendroctonus terebrans* Oliv.). These beetles attack the healthy living bark at and toward the base of the trunks of medium to large trees and kill areas varying in size from 1 to 10 square feet. These dead areas are subsequently burned off by surface fires and are then generally referred to as fire-wounds. The further damage to the exposed wood by successive fires, decay, and insects often results in a total loss of the best portion of the tree, or a reduction in value of the lower section of the trunk of from 10 to 50 per cent. (Hopkins, 1904*a*, 1909*b*.) These and similar wounds in the bark of trees, including those caused by lightning and by the uncovering and exposure of the wood in turpentineing, offer favorable conditions for the attack of the turpentine borer (*Buprestis apricans* Hbst.), the work of which, together with that of two or three others with similar habits, is very extensive, and causes losses amounting to from 10 to 50 per cent of the value of the wood of the best part of the trees thus affected. (Hopkins 1904*a*.)

The white pine weevil.—The abnormal development of white pine trees as the result of successive attacks on the terminals of the saplings and young trees by the white pine weevil (*Pissodes strobi* Peck) is an element of loss of considerable importance, especially in mixed stands and in open pure stands of this timber. The value of such trees is reduced from 20 to 50 per cent below those of normal development, and there is an additional loss from the effect of their spreading branches or crowns in the suppression or crowding out of trees which would otherwise occupy the space thus usurped. (Hopkins, 1906*c*, 1907*d*.)

There are many other examples of insects which damage the wood and bark of living trees, but those mentioned should be sufficient to demonstrate the importance of insects in this relation.

INSECT INJURIES TO THE WOOD OF DYING AND DEAD TREES.

Timber dying from insect attack and other causes, including fire, disease, storms, etc., is attacked by certain wood-boring insects which extend their burrows through the sound sapwood and heartwood, and thus contribute to the rapid deterioration and decay of a commodity which otherwise would be available commercially during periods of from one to twenty years or more after the death of the trees, depending on the species of trees and on the character of the product desired. This loss often amounts to from 25 to 100 per cent during the period in which the dead timber would otherwise be almost as valuable as if living. (Hopkins, 1894*a*, 1901*d*, 1904*a*, 1905*a*; Webb, 1909.)

CONIFEROUS TREES.

Sawyers.—One of the most striking examples of the destruction or deterioration of the wood of dying and dead timber, familiar to all lumbermen, is the injury to fire-killed and storm-felled pine, fir, spruce, etc., caused by boring larvæ known as “sawyers.” These borers hatch from eggs deposited by the adult beetles in the bark of the dying trees, and after feeding on the inner bark for a time they enter the solid wood and extend their large burrows deep into the heartwood. Fire-killed white pine is especially liable to this injury, and is often so seriously damaged within three or four months during the warm season as to reduce the value of the timber 30 to 50 per cent. The shortleaf, loblolly, and longleaf pines of the Southern States are damaged to a somewhat less extent, but instances are known in which more than one billion feet of storm-felled timber within limited areas were reduced in value 25 to 35 per cent within three months after the storm. (Webb, 1909.) The fire-killed and insect-killed sugar pine, silver pine, and yellow pine of the western forests are also damaged in a similar manner and the value of the product greatly reduced within a few months after the trees die. The aggregate losses from this secondary source in the coniferous forests of the entire country contribute largely to the annual waste of millions of dollars’ worth of forest products which otherwise might be utilized. (Hopkins, 1905*a*, p. 385; Webb, 1909.)

Ambrosia beetles.—Wood-boring insects of another class, known as timber beetles or ambrosia beetles, cause pinhole defects, principally in the sapwood, although some of them extend their burrows into the heartwood. These insects make their attack in the early stage of the declining or dying of the tree, or before the sapwood has materially changed from the normal healthy condition, and often in such numbers as to perforate every square inch of wood. Thus the wood is not only rendered defective on account of the presence of pinholes, but the holes give entrance to a wood-staining fungus which causes a rapid discoloration and produces still further deterioration of the product.

The sapwood of trees dying from the attack of other insects or from fire, storm, or other causes is often reduced in value 50 per cent or more, and in some cases the value of the heartwood is reduced in a like manner from 5 to 10 per cent. (Hopkins, 1894*a*, 1895*c*, 1898*b*, 1904*a*, 1905*a*.)

Pinhole borers in cypress.—An example of the destructive work of insects which attack dying and dead trees is found in the cypress in the Gulf States, where these trees are deadened by the lumbermen and left standing several months, or until the timber is sufficiently dry to be floated. Upon investigation it was found that trees dead-

ened at certain seasons of the year were attacked by the ambrosia beetles or pinhole borers, and that in some cases millions of feet of timber had been reduced 10 to 25 per cent or more in value. (Hopkins, 1907*b*.)

HARDWOOD TREES.

Roundheaded borers, timber worms, and ambrosia beetles.—The principal damage to dying and dead hardwood trees is caused by certain roundheaded wood-borers (Cerambycidae) with habits similar to the sawyer, by the timber worms mentioned as damaging living timber, and by ambrosia beetles having habits similar to those that attack the sapwood and heartwood of conifers. All of the hardwoods suffer more or less, but the greatest damage is done to the wood of hickory, ash, oak, and chestnut, which are often reduced in value 10 to 25 per cent or more within the period in which it would otherwise remain sound and available for commercial purposes. (Hopkins, 1894*a*, 1904*a*, 1905*a*.)

INSECT INJURIES TO FOREST PRODUCTS.

Damage is caused by various species of insects which are attracted by the varying conditions prevailing at different stages during the process of utilizing the forest resources, from the time the trees are felled until the logs are converted into the crude and finished product and until the latter reaches the final consumer, or even after it is placed in the finished article or structure. As a result, additional drains are made on the timber to meet the demand for the higher grades of lumber and for other supplies to replace those injured or destroyed. From the writer's personal investigations of this subject in different sections of the country it is evident that the damage to forest products of various kinds from this cause is far more extensive than is generally recognized. This loss differs from that resulting from insect damage to standing timber in that it represents more directly a loss of money invested in material and labor. (Hopkins, 1894*a*, 1903*c*, 1904*a*, 1905*a*.)

CRUDE PRODUCTS.

Roundheaded borers, timber worms, and ambrosia beetles.—Round timber with the bark on, such as poles, posts, mine props, sawlogs, etc., is subject to serious damage by the same class of insects as those mentioned under injury to the wood of dying and dead trees. The damage is especially severe when material is handled in such a manner as to offer favorable conditions for attack (Hopkins, 1905*a*), as when the logs are left in the woods on skidways or in mill yards for a month or more after they have been cut from the living trees. Under such conditions there is often a reduction in value of from 5 to 30 per cent or more, due to wormhole and pinhole defects caused by

roundheaded and flatheaded wood-borers and timber beetles. Frequently the insects continue the work in the unseasoned and even dry lumber cut from logs which had been previously infested. They also continue to work in mine props after they have been placed in the mine, and in logs and other material used for the construction of cabins, rustic houses, etc., and in round timbers generally.

The products from saplings, such as hickory hoop-poles and like material, are often seriously injured or rendered worthless by roundheaded and flatheaded borers and wood-boring beetles, sometimes resulting in a loss of from 50 to 100 per cent of the merchantable product. (Hopkins, 1905*a*.)

Stave and shingle bolts left in moist, shady places in the woods or in close piles during the summer months are often attacked by ambrosia beetles and timber beetles. The value of the product is often reduced, as a consequence, from 10 to 50 per cent or more. (Hopkins, 1894*a*, 1905*a*.)

Handle and wagon stock in the rough is especially liable to injury by ambrosia beetles and roundheaded borers. Hickory and ash bolts from which the bark is not removed are almost certain to be greatly damaged if the logs and bolts cut from living trees during the winter and spring are held over for a few weeks after the middle of March or first of April. (Hopkins, 1905*a*.)

Pulpwood, and cordwood for fuel and other purposes, cut during the winter and spring and left in the woods for a few weeks or months or in close piles after the beginning of the warm weather, are sometimes riddled with wormholes or converted into sawdust borings, causing a loss of from 10 to 100 per cent. One example reported from near Munising, Mich., represents a loss of \$5,000 from injury to spruce and fir pulpwood cut in the winter and kept in piles over summer.

MANUFACTURED UNSEASONED PRODUCTS.

Ambrosia beetles and other wood borers.—Freshly sawed hardwood placed in close piles during warm, damp weather during the period from June to September is often seriously injured by ambrosia beetles. Heavy 2-inch to 3-inch stuff is also liable to attack by the same insects, even in loose piles. An example of this was found in some thousands of feet of mahogany lumber of the highest grade, which had been sawed from imported round logs and piled with lumber sticks between the tiers of plank. Native species of ambrosia beetles, principally *Pterocyclon mali* Fitch, had entered the wood to such an extent as to have reduced the value 50 per cent or more within a few weeks. Oak, poplar, gum, and similar woods often suffer severely from this class of injury, causing losses varying from 5 to 50 per cent. (Hopkins, 1905*a*.)

Lumber and square timbers of both soft and hard woods with the bark left on the edges are frequently damaged by flatheaded and roundheaded wood borers, which hatch from eggs deposited in the bark before or after the lumber is sawed. There are examples of losses from this character of injury amounting to from 20 to 50 per cent or more.

Telegraph and telephone poles, posts, mine props, etc., are frequently injured before they are set in the ground, especially if the bark remains on them during a few weeks after the middle of March.

SEASONED PRODUCTS IN YARDS AND STOREHOUSES.

Powder-post beetles.—Hardwood lumber of all kinds, rough handles, wagon stock, etc., made partially or entirely of sapwood, are often reduced in value from 10 to 90 per cent by a class of insects known as powder-post beetles. The sapwood of hickory, ash, and oak is most liable to attack. The reported losses from this source during the past five or six years indicate that there has been an average reduction in values of from 5 to 10 per cent or more. (Hopkins, 1903*c*, 1905*a*.)

Old hemlock and oak tanbark is often so badly damaged by various insects which infest dead and dry bark that in some tanyards as much as 50 to 75 per cent of the bark that is over three years old is destroyed. In one tannery in West Virginia it is estimated that more than \$30,000 worth of hemlock bark was thus destroyed. (Hopkins, 1905*a*.)

FINISHED PRODUCTS.

The greatest loss of finished hardwood products, such as handle, wagon, carriage, and machinery stock, is caused by powder-post beetles. This is especially true of hickory and ash handles and like products in the large and small storehouses of the country, including the vast amount of material held in storage for the army and navy. When material of this kind is once attacked it is usually worthless for the purposes indicated, and therefore must be replaced with new material. In some cases losses have amounted to from 10 to 50 per cent, and it is estimated that the average losses have been as much as 10 per cent on nearly all sapwood material that has been in storage more than one year. (Hopkins, 1903*c*, 1905*a*.)

UTILIZED PRODUCTS.

Powder-post beetles, white ants, and other wood-boring insects.—The finished woodwork in implements, machinery, wagons, furniture, and the inside finish in private and public buildings are often seriously damaged by powder-post beetles, thus requiring increased demands for new material. (Hopkins, 1903*c*, 1905*a*.)

Construction timbers and other woodwork in new and old buildings are often so seriously damaged by powder-post beetles, white ants, and other wood-boring insects that the affected material has to be removed and replaced by new, or the entire structure torn down and rebuilt. (Hopkins, 1905*a*.)

Construction timbers in bridges and like structures, railroad ties, telephone and telegraph poles, mine props, fence posts, etc., are sometimes seriously injured by wood-boring larvæ, termites, black ants, carpenter bees, and powder-post beetles, and sometimes reduced in efficiency from 10 to 100 per cent.

INSECTS IN THEIR RELATION TO THE REDUCTION OF FUTURE SUPPLIES OF TIMBER.

Insects not only reduce future supplies by killing the mature trees and destroying the wood of timber that is inaccessible for utilization, but through injuries inflicted upon trees during the flowering, fruiting, germinating, seedling, and sapling periods of early growth they prevent normal reproduction and development. (Hopkins, 1904*a*, 1906*c*.)

INTERRELATIONS OF FOREST INSECTS AND FOREST FIRES.

Investigations conducted by the writer and assistants in all sections of the country during the past ten years indicate to them quite conclusively that the average percentage of loss of merchantable timber in the forests of the entire country to be charged to insects during a five or ten year period is infinitely greater than most people realize. (Hopkins, 1906*a*, pp. 4-5, 1908*b*, p. 345, 1909*b*, pp. 5, 24; Forbes, 1909, pp. 51-52.)

Losses from forest insects.—The writer estimates (p. 70) that for a ten-year period the average amount of timber in the forests of the entire country killed and reduced in value by insects would represent an average loss of \$62,500,000 annually.^a

It has been estimated (Hopkins, 1905*b*, p. 5; 1908*a*, p. 162) that the Black Hills beetle killed approximately 1,000,000,000 feet b. m. of timber during a period of ten years, which at \$2.50 per thousand would amount to an average of \$250,000 annually. This is merely one example of very destructive depredations by a single species of barkbeetle in a single national forest.^b (See also p. 70.)

Prof. Lawrence Bruner, state entomologist of Nebraska, at a meeting of the American Association of Economic Entomologists, held at

^a *Losses from forest fires.*—It has been estimated that "on the average, since 1870, forest fires have yearly cost \$50,000,000 in timber." (Cleveland, T., jr., 1909, p. 3.)

^b It has been estimated that the losses of timber from forest fires on all of the National Forests of the United States from 1905 to 1908, inclusive, average only \$165,062 annually. (Cleveland, T., jr., 1908, p. 541.)

Baltimore, Md., in December, 1908, spoke as follows: "I can agree with Doctor Hopkins that the insects are far more important in destroying our forests than fires." (Bruner, 1909, p. 53.)

Insect-killed timber as fuel for fires.—It has often happened that after insects have killed the timber over extensive areas the standing and fallen dead trees furnished fuel for great forest fires which have not only destroyed or charred the dead timber but killed the living timber and reproduction and swept on into adjacent areas of healthy timber. Indeed, abundant evidence has been found during recent investigations to indicate that some of the vast denuded areas in the Rocky Mountains and other sections of the country are primarily due to widespread devastation by insects, and that subsequent fires destroyed the timber and prevented reproduction. (Hopkins, 1906*a*.)

It is also evident that a considerable percentage of dead timber, and especially that found in coniferous forest regions, which has generally been believed to have been fire-killed is a result of primary attack by insects. This has been demonstrated in many cases by the pitch-marked galleries of the destructive barkbeetles on the surface of the wood of the old dead trees which had escaped subsequent fires.

Fire-killed timber injured by insects.—It is true that a vast amount of timber has been killed outright or has died as the direct result of forest fires, but in almost every case observed insects have contributed to a greater or less extent to the death of recently fire-injured trees which might otherwise have recovered, and especially to the rapid deterioration of the wood of a large percentage of the injured and killed trees. It is evident that in some cases fire-scorched and fire-killed timber has contributed to the multiplication of one or more of the insect enemies destructive to living timber, and thus the injury started by the fire may have resulted in a destructive outbreak of beetles. However, it is evident that this has happened only when the destructive beetle was already present in abnormal numbers in the forest surrounding the fire-swept area. Therefore, *it is believed that injuries by fire are not as a rule an important factor in contributing to subsequent depredations by barkbeetles.* Such fires, however, contribute to the multiplication of the insects which depredate on the bark and wood of dying and dead trees, so that in forested areas where fires are frequent the damage to the wood of such trees is more severe, and fewer injured trees recover on account of the abundance of secondary barkbeetle enemies which do not, as a rule, attack and kill living timber.

Destruction of insects by fire.—There is another important feature in the relation of insects and fire, in which the fire contributes to the destruction of the principal barkbeetle enemies of the living timber. This happens when the fire burns the timber while it is infested, thus effectually destroying the broods of the insects. It is perfectly plain that the dying and dead foliage of the beetle-infested trees and the

dead bark on the trunks would contribute to the spreading of crown fires and thus the bark on the entire infested trunks would be sufficiently scorched to kill the insects. Therefore, complete fire control may easily contribute to more extended depredations by insects on the living timber, thus increasing, rather than diminishing, the need for insect control. However, the setting of fires or permitting them to burn for the purpose of combating insects should never be undertaken or permitted.

Durability of insect-killed timber.—Some of the matured larch trees which evidently died as a result of defoliation by the larch worm between 1881 and 1885, and which had escaped subsequent depredations by fire and wood-boring insects, were found by the writer in 1908 to be standing and sound enough to be utilized for railroad ties and many other purposes. Under similar conditions the heartwood of red spruce and white pine in the East, of Engelmann spruce in the Rocky Mountains, and of Douglas fir in the Northwest coast region have been found by the writer to be sound enough for profitable utilization for pulp wood, lumber, fuel, and other purposes from twenty to thirty years after it had been killed by insects or fire. Thus it is shown that timber killed by insects and fire would be available for utilization for many years were it not for injuries through the secondary attacks of wood-boring insects and the destruction of insect-killed timber by forest fires.

INTERRELATION OF FOREST INSECTS AND FOREST FUNGI.

Decay following injury by insects.—It is well known that the burrows in the bark and wood of living and dead trees and in the crude and finished products often contribute to the entrance of bark and wood decaying fungi. Deterioration and decay are thus far more rapid than would otherwise be possible. It is also known that trees injured and dying from primary attack by parasitic fungi are attractive to certain insects which breed in the bark and wood of sickly and dying trees, and that certain other complicated troubles affecting forest trees are the result of an intimate interrelation and interdependence of insects and fungi. There can be no doubt, however, that certain species and groups of both insects and fungi are independently capable of attacking and killing perfectly vigorous and healthy trees.

SUMMARY AND ESTIMATES RELATING TO CHARACTER AND EXTENT OF INSECT DAMAGE.

The killing of trees by insects; the damage by them to the wood of living, dying, and dead timber; the destruction of insect-killed timber by subsequent forest fires; the damage to fire-killed timber by insects; and the damage from decay resulting from insect injuries to the wood, have all been more or less continuous for centuries and are still going on in the forest and woodland areas of this country.

While these depredations are not always evident or important in all forests or localities, yet almost every year, somewhere in the forests of the country, there are widespread depredations.

In every forest and woodland there is an ever present but inconspicuous army of insects which require the bark, wood, foliage, and seeds of the various tree species for their breeding places or food. Thus, the accumulated but inconspicuous injuries wrought during the period required for the growth of a tree to commercial size go far toward reducing the average annual increment below the point of profitable investments.

The accumulated damage to crude, finished, and utilized products reduces the profits of the manufacturer, increases the price of the higher grades to the consumer, and results in an increased drain on the natural resources.

In any attempt to estimate in *dollars* or *feet, board measure*, the extent of losses or waste of timber supplies caused by insects there are many conflicting factors which contribute to the difficulty of arriving at accurate conclusions. The published information concerning the amount in board feet of standing timber in the country is admittedly only an estimate, as are also the published data relating to average stumpage value. The published statistics relating to the amount and value of forest products are of course more accurate, but until more complete data can be furnished by the forest experts on the various complicated phases of forest statistics, any figures given by the forest entomologist relating to the value of timber and commercial products destroyed or reduced in value by insects must be considered on the same basis as the other estimates, and as the best that can be presented on available evidence.

Standing timber killed and damaged by insects.—When we consider the amount of standing merchantable timber killed by insects and the amount of standing timber, living, dying, and dead, which has been reduced in quantity and value through their agency during a ten-year period, we would estimate that such timber represents an equivalent of more than 10 per cent of the quantity and stumpage value of the total stand of merchantable timber in the United States at any given time.^a A certain percentage of such timber is a total

^a The estimate of the area and stand of the present forests of the United States, as given in Circular 166 of the Forest Service, page 6, is two trillion five hundred billion feet (2,500,000,000,000) board measure. The average stumpage value has been given as \$2.50 per one thousand feet b. m., making a total value of the standing merchantable timber of \$6,250,000,000. Ten per cent of this amount would be \$625,000,000, as the amount to be charged to insects for a 10-year period, or an average of \$62,500,000 annually. As an example, it has been estimated that over 1,000,000,000 feet b. m. of timber was killed by the Black Hills beetle in the Black Hills National Forest within a period of ten years. This, at \$2.50 per one thousand feet stumpage, would be an average of \$250,000 annually in a single forest of 1,294,440 acres.

loss because of the impossibility of utilization; but in some cases a greater or less percentage can be, and in some cases is, utilized within the period in which it is of sufficient value to yield a profitable return on the cost of logging and manufacture, although its value is greatly reduced.

Reduction in the Nation's wealth.—When we consider the forest resources both in merchantable timber and young growth as an important asset of the Nation's wealth; as representing a given value to the people for direct utilization; as a cover to the soil for protection of the land from erosion; as protection of headwater streams and of game; and as contributing to the æsthetic value of health and pleasure resorts, it would be difficult indeed to estimate the amount or percentage of loss of timber or the reduction in the land values, in each case, chargeable to insects. It is plain, however, that in the aggregate it is considerably greater than when estimated on stumpage values alone.

Reduction in cash revenue.—When we consider the problem from the standpoint of direct utilization we can estimate the annual loss on a basis of mill values; but here again we meet with complications, since much of the damaged material is left standing or is discarded in the woods or at the mill without measurement. Therefore we are left to judge from our observations and knowledge of the general conditions as regards dead and damaged timber found in the forests of the country, and the information from lumbermen in different sections, as to the percentage of loss from defective timber. On this basis we can estimate that the amount of insect-killed and damaged timber left in the woods, plus the reduction in value of that utilized, to be charged to insects is not far from an equivalent of 10 per cent of the value of the annual output of forest products of all kinds, in the rough. The total value of the forest products of the United States in 1907 is given as \$1,280,000,000; the losses from insect depredations would therefore represent an annual loss in a cash value of more than \$100,000,000. (Hopkins, 1895*c*, 1904*a*.)

Reduction in value of finished and commercial products.—When we consider the aggregate loss to the manufacturers of the finished products, to the trade, and to the consumer from insect injuries to the wood, it is evident that it amounts to many millions of dollars in addition to the estimated loss of crude products, or at least 3 per cent of the mill value.

METHODS OF PREVENTION AND CONTROL.

The results of extensive investigations and of practical applications of the knowledge gained during recent years have demonstrated that some of the most destructive insect enemies of American forests and of the manufactured and utilized products can be controlled, and

serious damage prevented, with little or no ultimate cost over that involved in forest management and business methods. (Hopkins, 1904*b*, 1905*a*, 1908*a*, 1909*b*.)

There are, of course, certain insects and certain injuries which, under present conditions and available information, can not be controlled or prevented, but it is very evident that if the information now available through the publications of the Department of Agriculture and through direct correspondence with its experts is properly utilized in the future it would result in the prevention of at least 30 per cent of the estimated annual waste of forest resources that has been caused by insects within recent years, and thus contribute greatly to the conservation of forest resources.

GENERAL PRINCIPLES OF CONTROL.

The ordinary spraying and similar methods employed in dealing with fruit and shade tree insects are, of course, not available for practical application in the case of forest trees. But there are other and less expensive methods of accomplishing the desired results.

In all efforts to control an outbreak or prevent excessive loss from forest insects it should be remembered that as a rule it is useless to attempt the complete extermination of a given insect enemy of a forest tree or forest product. Experience has demonstrated that it is only necessary to reduce and weaken its forces 75 per cent or more. It can not then continue an aggressive attack, but must occupy a defensive position against its own enemies until conditions resulting from avoidable negligence and mismanagement by the owners of the forests and manufacturers of forest products favor its again becoming destructive. Forest insects can thus be easily kept under control by good management.

The desired control or prevention of loss can often be brought about by the adoption or adjustment of those requisite details in forest management and in lumbering and manufacturing operations, storing, transportation, and utilization of the products which at the least expenditure will cause the necessary reduction of the injurious insects and establish unfavorable conditions for their future multiplication or continuance of destructive work.

It is, however, of the utmost importance that any adjustment or modification in management or business methods should be based on expert technical knowledge or advice relating to the species, habits, life history, and natural enemies of the insects involved and the essential features of the methods for their control. This should be supplemented by expert knowledge or advice on the principles of technical and applied forestry in the proper management, care, and utilization of the forest and its resources, and still further supplemented by practical knowledge and experience relating to local con-

ditions and facilities favorable and unfavorable for success in practical applications according to the recommended method or policy of control.

As has been shown, the mature or merchantable timber is the most susceptible to injury or death from the ravages of insects. Therefore, considered from the standpoint of insect control and the prevention of one of the greatest items of loss, it is important that such matured timber should be utilized before it begins to deteriorate, or before it reaches the stage of unprofitable growth.

For the greatest success in dealing with forest insects, it must be recognized that there are certain features in the habits and seasonal history of each species which differ to a greater or less extent from those of all other species, even of the same genus; that there are certain features in the characteristics of the various species of trees which differ from those of all other species; and that as a rule it is the technical knowledge of these peculiar features or characteristics of the trees and their enemies which furnishes the clew to successful methods of control.

There are also many peculiar features in the prevailing conditions in different localities, some of them favorable, others unfavorable, for the practical application according to a given method, so that while certain general advice may apply in a broad sense and be available for utilization by the practical man, whether owner, manager, or forester, without further advice, it is often necessary to diagnose a given case before specific expert advice can be given as to the exact cause and the most effective method or policy to be adopted, just as a physician must diagnose a case of illness or injury before prescribing the required treatment for his patient.

Therefore, in a consideration of the problem as to how far the waste of forest resources caused by insects can be prevented and how far the damaged timber can be utilized, we will attempt to give only general statements based on the results of our observations relating to some of the principal kinds of loss discussed in the first part of this paper, namely, by insects which (1) kill the trees, (2) cause injuries to the wood of living timber, (3) reduce future supplies, and (4) cause injuries to the manufactured products. In addition, we will consider the utilization of natural enemies of injurious insects, the utilization of damaged timber, and the present conditions and opportunities for success in the general control of forest insects.

CONTROL OF BARKBEETLES WHICH KILL TREES.

The barkbeetles which kill trees attack the bark on the trunk and destroy the life of the tree by extending their burrows or galleries in all directions through the inner living bark. The broods of young grubs or larvæ develop within the inner bark, on which they feed.

Those of some species develop to the adult stage within the inner bark and are exposed when the bark is removed, while those of other species transform to the adults in the outer corky bark and the larvae are not exposed when the bark is removed. Some species have two or more generations in a season or annually, while others have but one, and in a few species it requires two years for a single generation to develop. (Hopkins, 1909*b*.)

The barkbeetles of the genus *Dendroctonus* represent the most destructive enemies of the principal coniferous tree species of American forests, and at the same time are among the easiest of control. The general requisites for success are embodied in the following rules:

(*a*) Give prompt attention to the first evidence of a destructive outbreak, as indicated by an abnormal percentage of yellow or red topped dying trees, and especially when such trees occur in groups of ten or more or cover large areas; (*b*) secure authentic determination of the particular species of insect responsible for the trouble; and (*c*) take prompt action toward its control according to specific expert advice, published or otherwise, on the best method for the destruction of the necessary 75 per cent or more of the insects in the infested trees.

Some of the methods to be adopted to meet the requirements of various local conditions are as follows:

(1) Utilize the infested timber and burn the slabs during the period in which the broods of the destructive beetles are in the immature stages or before the developed broods emerge from the bark; or

(2) Fell the infested trees and remove the bark from the main trunk and burn the bark if necessary;^a or

(3) Remove the infested bark from the standing timber and burn the bark when necessary;^a or

(4) Immerse the unbarked logs in ponds, lakes, or streams, where the bark will remain soaked long enough to kill the insects; or

(5) Remove the unbarked logs or products to a locality where there are no trees liable to attack within a radius of 20 miles or more.

MAINTAINING CONTROL OF BARKBEETLES.

Future trouble of a serious nature from barkbeetles which kill trees can be prevented within a given forest or area of greater or less extent if an insect-control policy is adopted in connection with, or independent of, a fire-control policy by which groups of dying trees will receive similar prompt attention as that required for the prevention or control of forest fires.

In state and national forests.—In all forest reserves in which there is an organized force of rangers and fire wardens or patrols each

^a If the broods develop to adults in the outer bark, it must be burned; if they develop in the inner bark and are exposed when the bark is removed, burning is not necessary. As a rule the burning of the tops to destroy the insects is not necessary.

officer should be furnished with instructions for the location of beetle-infested trees, and with equipment and directions for taking the necessary action whenever the conditions demand or warrant it.

In private forests.—Private forests should receive the same attention as public forests, but this is often far more difficult on account of intervening forests, where the owners either can not or will not give the matter the required attention. While it may be advisable to have some laws to govern the treatment of timber infested with a dangerous pest, when the owner refuses to take any action such a law should apply only to the more extreme cases or as a last resort on authoritative advice. It is probable that in most cases legislation will not be necessary, and more ultimate good will result without than with strict laws, especially when it can be made clear to the owner that his personal interests demand that he take the proper action and that, when necessary, his neighbors will render assistance, as is done in the case of a forest fire.

Inaccessible areas.—There are yet large inaccessible areas in the East and West where it is not practicable or possible at present to control the depredations by these beetles and which must therefore be left to the same natural adjustment that has been going on in all forests from their beginning. While under such natural control much of the older matured timber will be lost it will usually be replaced by young growth, either of the same species of trees or of a different species, so that under normal conditions the forest will be perpetuated; but under exceptional conditions and combinations of detrimental influences, such as secondary insect enemies, fire, drought, etc., extensive areas may be completely denuded, never to be reforested under natural conditions. Therefore it will evidently not be very long before it will pay to adopt insect-control policies even in the areas that are inaccessible for profitable lumbering.

EXAMPLES OF SUCCESSFUL CONTROL OF BARKBEETLES.

We have a sufficient number of examples of successful control of depredations by the destructive barkbeetles to demonstrate the practicability of the advice based on the results of recent entomological investigations.

Control of the eastern spruce beetle.—The control of an alarming outbreak of the eastern spruce beetle (*Dendroctonus piceaperda* Hopk.) in northeastern Maine in 1900 and 1901 was effected by the concentration of regular logging operations into the areas of infested timber and placing the logs in lakes and streams and driving them to the mills on the Androscoggin River. Thus, with little or no additional expense, there was a saving to one firm, according to its estimates, of more than \$100,000.

Control of the Black Hills beetle.—An extensive outbreak of the Black Hills beetle (*Dendroctonus ponderosæ* Hopk.) in the vicinity of Colorado Springs, Colo., in 1905-6, which was threatening the living pine timber of the entire section, was brought under control through the efforts of the private owners of forests and those of forest officials in the adjoining National Forests. It was accomplished by cutting and barking about 1,000 beetle-infested and beetle-killed pine trees. The cost of the operations was largely, if not entirely, covered by the utilized felled timber, although there was considerable unnecessary expense involved through the felling and barking of trees from which the beetles had emerged and from the unnecessary burning of the bark and tops.

The successful control of another serious outbreak of the Black Hills beetle, in 1906, on an extensive private estate in southern Colorado, was effected through the efforts of the owners, who had some 500 infested trees felled and barked within the necessary period to destroy the broods. A large percentage, but not all, of the infested timber was thus treated. These operations were so successful that not a single infested and dying tree could be found when the area was inspected in 1908. In this, as in the other case, considerable unnecessary expense was involved in the burning of the bark and tops, but the value of utilizable timber was evidently more than enough to pay all expenses. It is evident that in this case a destructive invasion was prevented.

The most striking example of success relates to a large estate near Idaho Springs, Colo., and in the adjoining Pike National Forest. In May, 1907, it was found that some 63,000 feet of standing timber on the estate was infested by the Black Hills beetle, and the owner was advised by the Bureau of Entomology that unless the ravages were checked at once the beetle would kill the timber not only on this estate but that on the adjoining estates and National Forest, and that therefore radical action should be taken according to the recommendations and detailed instructions given relating to a necessary control policy. No action was taken, however, before the first of the following July, and therefore not in time to prevent the broods of beetles from swarming from the infested trees and extending their ravages. In December, 1907, another examination of the timber was made, and it was found that instead of 65,000 feet of timber in the old infestation there was nearly four times as much timber involved in the new, or over 250,000 feet. The owner was again notified of the serious character of the outbreak, and further suggestion made that if the logs from the infested trees were converted into lumber and the slabs burned before May, 1908, it would result in the protection of the remaining living timber. Immediate steps were then taken by the owners to carry out the original

recommendations. An expert in locating infested timber, working under instructions from this Bureau, gave instructions to the manager of the estate in locating and marking the infested trees and in the essential features in the methods of utilization to destroy the necessary number of beetles; he also marked infested timber on an adjoining estate and on the National Forest. Five months later, in May, 1908, this expert reported that the larger clumps of infested trees on the estate had been converted into lumber and the slabs burned, and that the marked trees on the adjoining estate and National Forest had been cut and barked. In November, 1908, another inspection of the forest on the estate and surrounding area was made by the expert, and on December 1 he reported as follows:

Nothing could be more satisfactory than the results obtained by the cutting of the infested timber on the estate. Your recommendations and instructions submitted to the owner, and carefully followed by the manager of the estate, have clearly demonstrated that insect infestation can be controlled, and at no expense to the owner of the timber involved: in fact a very satisfactory price was realized, resulting in a net profit, I understand, of over \$5 per thousand feet, board measure, on the 240,000 feet cut. This, of course, does not include the profit of the milling operations, but for the logs sold at the mill, after deducting the expenses of cutting and logging. The sawmill was owned and operated by an Idaho Springs firm, and the manufactured article sold in that town. I spent six days on the estate, November 18 to 23. After a very thorough examination of the timber, I found only three infested trees, isolated individuals, and over a mile from where the large clumps of infested trees were cut. With the exception of those three trees, there is no new infestation on the estate. I also examined the adjoining lands, but no new infestation was observed. The infested trees which I marked in December, 1907, had been cut and barked. On the Pike National Forest, contiguous to the first mentioned estate, where you will remember I marked some clumps of infested trees, no new infestation was found, not one tree.

This most gratifying result demonstrated two important things: One, that an extensive outbreak by the most destructive barkbeetle enemy of the pine timber of the central Rocky Mountain forests, involving in this case more than 1,000 infested trees, can be controlled without expense, and even at a profit, whenever the conditions are favorable for the utilization of the infested timber; the other, that the essential details of the recommendations and expert advice, based on the results of scientific research, can be successfully applied by a manager of a private forest or by the rangers of national and state forests. It also indicates quite conclusively that the widespread depredations in the Black Hills National Forest could have been prevented with very little expense to the Government if the matter had received prompt attention in 1901, when the first investigations were made and essentially the same recommendations submitted. But, through the lack of public appreciation of the importance of the problem at the time, and the lack of sufficient authority and funds later, the outbreak was allowed to extend beyond practical control, and in conse-

quence a large percentage of the timber of the entire National Forest has been killed. There were, then, no forcible examples of the practical value of such recommendations based on scientific research, and no other argument was effective in arousing public interest in the threatening character of the outbreak or confidence in scientific advice or methods of control. Now we have several examples demonstrating the practicability of forest-insect control in America which should lead to confidence in the results of scientific research as a basis for success in practical application.

Control of the hickory barkbeetle.—The complete control of the hickory barkbeetle (*Scolytus quadrispinosus* Say), which threatened the total destruction of the hickory trees on Belle Isle Park, at Detroit, Mich., in 1903, was effected by felling and removing the infested trees and converting them into merchantable products, all without cost to the park commission. (Hopkins, 1904b.)

CONTROL OF INSECTS WHICH CAUSE DEFECTS IN LIVING TIMBER.

The class of insects which causes defects in the wood of living timber can be controlled to a greater or less extent, depending upon local conditions, and a large percentage of the losses prevented through the adoption of certain requisite details in forest management. Of these the following are especially important:

(1) The utilization of all of the defective and infested timber that will pay expenses for manufacture into merchantable products, such as lumber, cordwood, etc.

(2) The burning of infested timber and waste material not available for use, including dead standing and fallen timber, to remove the breeding places of insects like the oak timber worm and the chestnut timber worm, which go from the dead to the living timber.

(3) The prevention of wounds of any kind in the bark of living trees.

(4) The prevention of future losses by the practice of improved forestry methods which will eliminate favorable conditions for injury and contribute to a perpetual supply of vigorous, healthy timber to be utilized before it passes the stage of profitable increment.

PREVENTION OF INJURY TO DYING AND DEAD TREES.

A large percentage of the injury to the wood of insect, fire, and lightning killed trees and those killed or dying from injuries by storms, disease, etc., can be prevented as follows:

(1) By the prompt utilization of such timber within a few weeks or months after it is dead or found to be past recovery.

(2) By removing the bark from the merchantable portions of the trunks within a few weeks after the trees are dead (the work to be done either before or after the trees are felled).

(3) By felling the trees and placing the unbarked logs in water.

(4) By the adoption of a system of forest management which will provide for the prompt utilization of all trees which die from any cause.

PREVENTION OF LOSS FROM INSECT INJURIES TO NATURAL AND ARTIFICIAL REPRODUCTION.

The successful control of the insects which destroy or prevent the normal development of natural reproduction is a far more difficult problem than that presented by other classes of insect injuries, but in this as in the others a great deal can be accomplished toward preventing the reduction of future supplies.

Much can be accomplished in nurseries and small plantations by the adoption of the ordinary methods of controlling farm and orchard insects, but in the natural forests reliance must be placed largely on systems of forest management which will bring about unfavorable conditions for the work of the more important enemies. (Hopkins, 1906*c*.)

UTILIZATION OF IMMUNE AND RESISTANT VARIETIES AND RACES OF TREES.

Certain individuals representing varietal or racial forms of trees of a given species are sometimes found to be either immune or decidedly more resistant to the insects which are destructive or seriously injurious to the life or wood of other individuals or varieties of the same species. This fact suggests the importance of recognizing the well-known principle of improvement by selection. Thus, selecting seed or cuttings from such immune and resistant trees for artificial propagation, or taking great pains to leave such trees in commercial or selection cuttings for natural reproduction, will undoubtedly be an important step toward providing against damage and loss in the future. (Hopkins, 1906*b*, 1907*a*, 1907*c*, 1907*d*.)

PREVENTION OF INSECT INJURIES TO FOREST PRODUCTS.

The problem of artificial control and prevention of insect injuries to forest products offers less difficulties perhaps than that relating to many other branches of the general subject of forest-insect control. In most cases the principle of prevention is the only one to be considered, since the damage is done soon after the insects enter the wood, and therefore it can not be repaired by destroying the enemy.

CRUDE PRODUCTS.

The proper degree of moisture found in the bark and wood of newly felled trees, saw logs, telegraph poles, posts, and like material, cut in the fall and winter and left on the ground or in close piles

during a few weeks or months in the spring and summer, or during the period when the particular species of injurious insects are flying, are some of the conditions most favorable to attack. The period of danger varies with the kind of timber and the time of the year it is felled. Those felled in late fall and winter will generally remain attractive to ambrosia beetles and adults of round and flat headed borers during March, April, and May. Those felled during the period between April and September may be attacked in a few days after they are felled, but the period of danger from a given species of insect may not extend over more than a few weeks. Thus certain kinds of trees felled during certain seasons are never attacked, while if they are felled at other times and seasons the conditions for attack may be most favorable when the insects are active, and then the wood will be thickly infested and ruined. The presence of bark is absolutely necessary for successful infestation by most of the wood-boring grubs, because the eggs and young stages must occupy the inner and outer portions before the latter can enter the wood. Some ambrosia beetles and timber worms will, however, attack barked logs, especially those in close piles or otherwise shaded or protected from rapid drying. A large percentage of the injury to this class of products can be prevented, as follows:

(1) Provide for as little delay as possible between the felling of the tree and its manufacture into rough products. This is especially necessary with trees felled from April to September in the region north of the Gulf States and from March to November in the latter, while the late fall and winter cuttings should all be worked up by March or April.

(2) Do not leave the round timbers in the woods or on the skidways during the danger period, or, if this is unavoidable, take every precaution to facilitate the rapid drying of the inner bark by keeping the logs off the ground, in the sun, or in loose piles, or else, if possible, the opposite extreme should be adopted and the logs kept in water.

(3) Remove the bark within a few days after the trees are felled, from poles, posts, and other material which will not be injured by checking or season cracks.

(4) Take advantage of the proper months or seasons in which to fell or girdle different kinds of trees to avoid danger.

Damage to products cut from saplings and left with the bark on can be prevented by transporting the material from the woods soon after it is cut, so that it will not be left in piles or bundles in or near the forest during the season of insect activity. Damage may also be prevented by care not to leave the material stored in one place for several months.

Pinhole damage to stave and shingle bolts cut during a warm season can be prevented by removing the bark from the timber as

soon as it is felled and by converting the bolts into the smallest practicable dimensions and piling them in such a manner as to facilitate rapid drying.

Damage to unseasoned handle and wagon stock in the rough can be prevented by taking special precautions to provide against the same favorable conditions for attack as mentioned in connection with round timbers. This is especially necessary with hickory and ash if cut during the winter and spring.

Damage to pulpwood and cordwood can be prevented to a great extent by placing the sticks of wood in triangular or crib piles immediately after they are cut from the trees, especially if the timber is cut during the danger period or must be held for a few months during the warm season. Peeling or splitting the wood, or both, before it is piled will also provide against damage from insects.

MANUFACTURED PRODUCTS.

UNSEASONED PRODUCTS.

Freshly sawed hardwood lumber placed in close piles during warm, damp weather in the period from July to September, inclusive, presents the most favorable conditions for injury by ambrosia beetles. In all cases it is the moist condition and retarded drying of the lumber which induces attack. Therefore any method which will provide for the rapid drying of the lumber before or after piling will tend to prevent loss. It is important, also, that heavy lumber should, as far as possible, be cut only in the winter and piled so that it will be well dried out before the middle of March.

The damage to lumber and square timber when the bark is left on the edges or sides can be prevented by removing the bark before or immediately after the lumber is sawed, or by sawing and piling the material during the winter, or if sawed at other times it should be piled so that rapid drying will be facilitated.

SEASONED PRODUCTS.

Unfinished seasoned products.—Injury by powder-post beetles to dry hardwood lumber and other material in stacks or storehouses can be prevented as follows:

(1) Have a general inspection of the material in the yards and storehouses at least once a year, preferably during November or February, for the purpose of (a) sorting out and destroying or otherwise disposing of any material that shows the slightest evidence of injury, as indicated by the presence of fine powdery boring dust, and (b) sorting out and destroying all old and useless sapwood material of any kind that will offer favorable breeding places for the insects.

(2) Prevent the introduction into the lumber yards or store-houses of any infested material, remembering that the insect may be thus distributed to or from all parts of the world.

(3) Adopt a system of classification of all dry or seasoned hardwood stock which will provide for (a) the separation of the pure heartwood material from the pure and part sapwood material; (b) classification of all kinds of wood most liable to attack, such as hickory, ash, oak; (c) the successive utilization or sale of the older material (remembering that material one year old or over is far more liable to injury); (d) providing against the accumulation of refuse material in which the insects could breed; and (e) treating the best material with linseed oil or kerosene to prevent attack.

Finished seasoned products.—Damage to finished handles, oars, spokes, rims, hubs, wheels, and other unpainted wagon, carriage, machinery, and implement stock in factories, wholesale and retail storehouses, and army and navy stores can be prevented by the adoption of the same general rules as those given under rough products. In addition, damage can be controlled and prevented in the following manner:

Sort out and (a) destroy all articles showing the slightest evidence of powder-post injury, or (b) treat with kerosene oil such infested and slightly injured articles as may be tested for required strength and found to be of sufficient value for retention, placing the same in quarantine for a sufficient time to determine whether the treatment is successful.

Damage by powder-post insects to many kinds of articles can be prevented and at the same time the material otherwise benefited by treating the sapwood with linseed oil or kerosene, either by immersing it in the oil or by applying the oil with a brush, the application to be made as soon as possible after the articles are finished from recently seasoned, uninjured stock.

PAST AND PRESENT CONDITIONS OF POWDER-POST INJURY.

Up to 1906 there were a great many reports of extensive losses of valuable material from the ravages of powder-post beetles which were seriously affecting all industries involved in the manufacture, sale, and utilization of the classes of hardwood products affected by them. In response to these reports and accompanying appeals to the Department of Agriculture for information on causes and remedies, the problem was thoroughly investigated and specific advice and instructions relating to practical methods of control and prevention have been widely disseminated, both through publications of the Department and special correspondence.

Reports of present conditions from our principal correspondents, together with the less frequent requests for advice, indicate that

the disseminated information has been extensively utilized and that it has been worth many millions of dollars toward eliminating the losses and reducing the drain on the limited supply of the kinds of timber required to replenish the damaged and destroyed material.

The army and navy stores of handles, tent poles, wheelbarrows, oars, and many other hardwood articles have suffered severely from powder-post damage, involving an enormous loss, but the carrying out of the information already supplied has evidently contributed greatly toward the elimination of this source of loss to the Government.

TAN BARK.

Damage to hemlock and oak tan bark by the class of insects which in some cases has been so destructive to these products in the past can be easily prevented without cost, as follows:

- (1) Utilize the bark within three years from the time it is taken from the trees.

- (2) Prevent the accumulation in the yards and store-sheds of old bark and waste material in which the insects can breed.

These simple methods have been extensively adopted since their recommendation in correspondence and publications between about 1894 and 1904, and afford one of the most striking examples of the value of expert information on the peculiar habits of insects and of how millions of dollars can be saved without cost through a simple adjustment in methods of utilization.

UTILIZED PRODUCTS.

Damage and loss from insect injuries to timber and other woodwork in structures of various kinds, to telephone and telegraph poles, posts, railroad ties, mine props, etc., can be prevented to a large extent through the adoption of the proper methods of management or of treating the material with preservatives before and after it is utilized.

TIMBERS AND WOODWORK IN STRUCTURES.

Injuries to timbers and woodwork in dwellings, outbuildings, bridges, etc., by powder-post insects can be prevented as follows:

- (1) Use nothing but heartwood for the concealed parts most liable to damage.

- (2) If it is necessary to use all or part sapwood material, attack can be prevented by treating the sap portions with kerosene, coal tar, creosote, or linseed oil. Facilities for future treatment can be provided wherever the rough or finished woodwork is exposed, as in outbuildings, bridges, etc., if care is taken to expose the sapwood portions.

(3) If the untreated timbers and woodwork in old buildings show evidence of attack, the affected portions should be given a liberal application of kerosene.

Damage by white ants, or termites, can often be prevented in the following ways:

(1) By the use of nothing but sound wood for underpinning and foundation timbers and the removal of decaying timbers from old structures.

(2) By preventing moist conditions of the wood in any part of the structure and especially that in foundation timbers.

(3) By the treatment of timbers necessarily exposed to moist conditions with creosote, zinc chlorid, corrosive sublimate, etc.

(4) If the timbers become infested, further progress of insect damage can be prevented by removing the badly damaged parts and soaking the remainder with kerosene, fumigating with bisulphid of carbon, and by removing any adjacent decaying or other wood in which the insects have been breeding or may breed, such as logs, stumps, etc.

Log cabins and rustic work.—Damage by bark and wood boring insects to the unbarked logs and poles used in rustic cabins, summer houses, fences, etc., can be largely prevented by cutting the material in October and November and utilizing it at once, or by piling it off the ground or under cover in such a manner as to offer the best facilities for the rapid and thorough drying of the inner bark before the middle of March or the 1st of April following. If these necessary precautions are not taken, and there is evidence that insects are at work in the bark and wood, the damage can be checked by injecting bisulphid of carbon through natural or artificial openings in the affected bark, and immediately stopping these and other openings with putty or a similar substance.

Poles, posts, piles, ties, mine props, and similar products.—Insect damage to poles, posts, and similar products can be prevented to a greater or less extent by the preservative treatments which have been tested and recommended by the Forest Service for the prevention of decay. These should be applied before the material is utilized for the purposes intended, or, if it be attacked after it has been utilized, further damage can be checked to a certain extent by the use of the same substances.

It is often of prime importance to prevent injury from wood-boring insects, for the reason that such injuries contribute to more rapid decay. Therefore anything that will prevent insect injury, either before or after the utilization of such products, will contribute to the prevention of premature deterioration and decay.

UTILIZATION OF NATURAL ENEMIES AND FACTORS IN THE CONTROL OF INJURIOUS INSECTS.

Were it not for the natural checks and natural factors of control of some of the more destructive insect enemies of forest trees and forest products, artificial control would in many cases be impossible, and the depredations would evidently be far more continuous and complete. These natural factors in the control of the depredating insects consist of parasitic and predatory insects, diseases of insects, birds, adverse climatic conditions, etc. While one or more of these beneficial factors exert a continuous and powerful influence toward the prevention of a more extensive waste of forest resources, it has been repeatedly demonstrated that they can not be depended on to prevent widespread devastations or to otherwise work for the best interests of the private or public owner by protecting the best trees and the best tree species. The insects and birds which prey upon the depredating insects also have factors to contend against, consisting of insects, birds, diseases, and climatic conditions. Therefore under normal conditions the tendency is toward the preservation of a balance between the warring factors, but frequently the enemies of the trees get the ascendancy and take on the character of an invasion, which may continue for two or three or even ten years before the balance is again adjusted through the influence of the natural enemies or diminished food supply. Thus a vast amount of timber or of a given forest product may be destroyed before the factors of natural control can prevail.

It is evident that the most effective utilization of the factors of natural control will be through the alliance with them of the owner of the forest in the artificial reduction of the enemies of the trees rather than by efforts to make the natural enemies of the injurious insects his allies through artificial introduction or dissemination. The former is accomplished by the adoption of methods of combating the invaders which will reduce and weaken their forces below their power of prosecuting aggressive movements and attacks, or, as previously stated, to reduce their numbers to the point where they must occupy a defensive position against their natural enemies and be dependent for their supplies of food and breeding places upon that furnished through avoidable mismanagement of the forests and manufacturing operations. Thus the owner of the forest can contribute greatly toward the preservation of a balance which will be to his material benefit. On the other hand, he may in the future, as in the past, contribute greatly to the multiplication of the depredating insects and to greatly increased losses caused by them, through neglect or a disregard of available information on the fundamental prin-

ciples of insect control in the management of forests and manufacturing enterprises.

BENEFICIAL INSECTS.

The beneficial insects comprise those which are internal or external parasites of the immature or mature stages of the injurious insects, and predators which feed on the young or adults of insects either before or after they make their attack on the trees or products. These two beneficial factors are doubtless far more effective in the long run than any other agencies of natural control. Yet they, in combination with all other factors, can not be always relied upon to render continued and efficient control. They can, however, be relied upon to respond to artificial assistance in reducing the numbers of the depredators.

BENEFICIAL DISEASES OF INSECTS.

It is very evident that the parasitic fungi and bacteria which sometimes cause epidemics among injurious insects often exert a powerful influence toward the control of extensive outbreaks or invasions of insect enemies of forests. Indeed, it appears that the greatest service rendered by this class of natural enemies is in the frequent sudden appearance of an epidemic which kills off a destructive species of insect after the latter has increased to such numbers and extended its depredations over such vast areas as to be far beyond the control of man or his insect and bird allies. Numerous examples of this kind of natural control are found in the sudden ending of widespread depredations by various species of caterpillars and sawfly larvæ which defoliate deciduous and coniferous trees. As a rule, however, the beneficial effects of the diseases of insects prevail only after the injurious insects have increased to excessive numbers. Therefore this factor of insect control can not be depended upon to hold the insects in check or prevent outbreaks. The fact, however, that it operates on a class of insect enemies of the forest (defoliators) which at present can not be controlled by any known artificial methods renders the services of the diseases all the more valuable.

It is believed that with further knowledge of nature's method of propagating, perpetuating, and disseminating the diseases which cause epidemics among insects, they may be utilized more or less successfully through artificial propagation and dissemination to prevent threatened invasions of defoliating insects.

BENEFICIAL BIRDS.

It is very evident that certain kinds of birds, such as woodpeckers, render valuable service toward the natural control of destructive bark and wood boring insects. They appear to render the greatest

service, however, where but few trees are being killed or injured, because their concentrated work on such trees may contribute toward the prevention of an abnormal increase of the insects. They also render some service as allies of the other beneficial factors which assist in artificial control. It is evident, however, that where many hundreds or thousands of trees are being killed the comparatively limited number of birds in any forest under the most favorable conditions could have little or no beneficial effect. Therefore, while the birds should be classed among the valuable friends of the forest, and should be protected, it is plain that they can not, even with the utmost protection, always be relied upon to protect the forest against destructive ravages of insects.

We must remember, in this connection, that there are complicated interrelations between birds, injurious insects, and beneficial insects which do not necessarily always operate to the benefit of the forest. In fact, it may sometimes be quite the reverse. Therefore, in order to derive the greatest benefit from the conflict between the birds, the insect enemies of the trees, and the insect friends of the trees, we must utilize our knowledge of the factors which are contributing toward the preservation of a balance, so that whenever the enemies of the forest threaten to get beyond natural control we may enter the field through artificial means and endeavor to force them back to their normal defensive position.

BENEFICIAL CLIMATIC CONDITIONS.

The benefits to be derived from climatic conditions which are detrimental or destructive to insect enemies of the forest, while sometimes very great, are necessarily unreliable, and thus can not be depended upon to assist in artificial control. In fact, the very condition which may contribute to the destruction of one depredator may favor the multiplication of another.

UTILIZATION OF WASTE CAUSED BY INSECTS.

When we come to consider the vast amount of standing timber in the forests of the country which has been injured or killed by insects, and will go to waste if it is not utilized within a limited period, we realize that there are great possibilities in its utilization as a means of preventing the reduction of future supplies of living healthy timber. It is all the more important that the insect-infested timber should be utilized, because in so doing we can contribute more perhaps than in any other way to the reduction of the insects to or below their normal numbers, and thus provide against serious injury in the future, as well as to the maintenance of control.

PRESENT CONDITIONS AND OPPORTUNITIES.

Unfortunately, the examples of management, or rather mismanagement, which are contributing to an extension or increase of waste are far in excess of those which under proper management are contributing to the reverse. This is due in a large part to the conditions that have prevailed in American forests in the past that have rendered it impracticable to adopt improved policies of forest management, but at present it is largely due to a lack of appreciation of the importance of the subject, and of the opportunities to prevent such losses, when it is not only practicable but possible to do so, and when it will, at the same time, yield large returns on the necessary additional expense. This is especially true in thousands of farmers' woodlots and private holdings in the States east of the Mississippi River, in which from 25 to 90 per cent of all of the serious injury of the past can be prevented in the future with little or no additional expense over that required for ordinary good forest management.

FOREST ENTOMOLOGY AS APPLIED TO AMERICAN FORESTS.

It is only within the past eleven years that any attempt has been made toward a systematic investigation of the insect enemies of the forest trees and forest products of the entire country. The state of knowledge of the subject previous to that time can be judged by the fact that a number of the most destructive enemies of forest trees are found to be new to science, and that nothing whatever was known of the habits and seasonal history of a large number of the more important known species which are common enemies of forest trees and forest products, while scarcely anything was known in regard to practical methods of controlling the principal enemies of the forest and its products, or of preventing losses from their ravages as applied to conditions in this country.

PRESENT KNOWLEDGE.

Within the past ten years forest-insect investigations have been conducted by the Bureau of Entomology of the Department of Agriculture in all of the principal forest regions of the United States, and have led to the following results:

Results of investigations.—Satisfactory progress has been made toward the attainment of some of the fundamental objects of the investigations, one of which has been the laying of a substantial foundation for forest entomology in this country on which future progress can be made along the lines of acquiring, disseminating, and applying information of immediate practical value in the protection of our forest resources.

Acquired and new information.—The principal insect enemies of the forests and forest products of North America have become

known, and the general character and extent of their depredations have been ascertained.

The more important facts in the life history, habits, and practical methods of control relating to some of the more destructive insects have been determined.

A mass of original data has been collected relating to forest insects in general, including not only those which are destructive or injurious, but also those which are beneficial or neutral in their relation to the forest, as represented by a collection of more than a million specimens of insects and their work.

As a direct result of the investigations of forest insects during the past six years, at a cost of less than \$53,000, there has been accumulated a reserve fund of information now available through publications, correspondence, and field demonstrations which, if properly utilized for practical application, would evidently prevent from 10 to 30 per cent of the annual losses at a very small cost.

Disseminated information.—In addition to information disseminated in all sections of the country through correspondence, lectures, demonstrations, exhibitions, etc., the published information, based on results of investigations conducted by the West Virginia Agricultural Experiment Station and by the Bureau of Entomology of the U. S. Department of Agriculture during the past eighteen years, is represented by over 1,300 pages, 99 plates, and 340 figures. (See list of publications.)

NEEDS.

The work that has been done is only a beginning in the vast field of forest entomology. There is need of more systematic work (or so-called pure science) on the different stages of the thousands of species of injurious and beneficial insects involved. *This is absolutely necessary in order to have the further scientific basis of facts on which to build the structure of complex details necessary to success in practical application in its broadest sense.*

There is need of further detailed study of habits and seasonal history of the species of injurious and beneficial insects, as well as of the local and other conditions favorable and unfavorable for their multiplication and work.

Further information is desirable on the principal factors of natural control of injurious insects, in order that it may be better utilized to facilitate artificial control.

There is special need of more general information and public interest in the subject of losses from insect depredations on standing timber and timber products, and a better realization of the possibility and practicability of preventing losses.

Looking to this end, there is need of further demonstration and educational work along the lines which will bring the matter to the

attention of the man in the woods, sawmill, factory, or trade who is in direct touch with the local conditions and business methods.

There is a special need of more experts in forest entomology, and there will be an increasing demand for such experts in the future, to organize and take charge of insect-control policies in state forests, in public parks, in the more extensive private or commercial forests, and in extensive manufacturing enterprises, and to give instructions to students in forestry schools and forestry departments in state and other institutions of learning.

The need of trained and experienced experts in forest entomology for all of this class of work can not be too strongly urged. There is perhaps no branch of economic or applied science which requires more technical knowledge and practical experience as a basis for proper investigations and authentic instructions and advice than forest entomology, and there is perhaps no other feature in the science and practice of forestry in which advice and application based on insufficient knowledge is so dangerous.

There is quite a general recognition of the importance of guarding against contagious diseases, of the necessity of consulting a physician in cases of serious illness, and of relying on authorized pharmacists to fill the prescription, and then administering the prescribed treatment according to directions, but it is a notable fact that there are comparatively few persons who, even when deeply interested in preserving the health of the forest, have heretofore recognized the importance of guarding against insect epidemics or of consulting an expert forest entomologist in case of a threatened or existing outbreak. It has often happened that when such advice has been sought and received, the treatment was not administered according to the recommendations but changed to suit the ideas of some one entirely ignorant of the facts and principles upon which it was based. This has often resulted not only in failure to accomplish the desired end, but has contributed to an aggravation of the trouble and increased loss. The determination of the cause of specific troubles affecting the individuals of a given species of forest tree or an injury to a given type of product is just as complicated and requires the same elements of experience, training, and skill as that required for the determination of the cause of a given disease or the character of a given injury affecting man. It is just as important to know the cause or character of injury in the former as it is in the latter, in order to prescribe the specific treatment which will yield the desired results.

Therefore, in order to make the best progress toward preventing future waste of our forest resources from depredations by insects, every one interested in the subject, and especially those in authority in the public and private institutions of investigation and learning, should see to it that the instructions to students and the information

given out to the public is not only the best available but that it is limited to the range of expert knowledge of the subject possessed by the instructor or investigator.

ELEMENTARY AND TECHNICAL KNOWLEDGE OF FOREST ENTOMOLOGY FOR THE FORESTER.

While it may be desirable that every professional forester should have an expert knowledge of forest entomology, it is rarely possible, even under exceptionally favorable conditions, for him to acquire more than the necessary elementary knowledge, and even this has not been possible under the conditions which have necessarily prevailed in the forest schools, and in the practice of forestry, in this country. Little or no time has been available for acquiring the necessary information from subsequent study and practical experience. Therefore this feature in the education of the American forester has been practically neglected.

PRESENT REQUIREMENTS OF INSTRUCTION.

As long as expert forest entomologists and authentic text-books based on American insects and conditions are not available for giving a complete course in technical and applied forest entomology the requirements of such a course should be limited to instruction in elementary entomology, and in elementary principles of applied forest entomology, which will give the necessary foundation for intelligent observations and utilization of available information as required in future practice.

CONCLUSION.

There is conclusive evidence that insects have been in the past, and are now, important factors in the waste and reduction of timber supplies, and will continue to be such in the future (pp. 57-58).

They attack perfectly healthy trees of all ages and kill them (p. 58).

They have at times killed a large percentage of the best timber over thousands of square miles of heavily forested lands (pp. 58-60).

They reduce the value of living timber and that of crude and finished products (pp. 60-66).

The accumulated evidence through many years of investigation and observation in the principal forest areas of the entire country by the writer and the field assistants in forest insect investigations furnishes the basis for the following summarized statements and estimates:

A large percentage of pine and spruce timber was killed by the southern pine beetle in 1890-1892 over an area of 75,000 square miles in West Virginia, Maryland, Pennsylvania, Virginia, and North Carolina (p. 58).

Billions of feet of matured spruce have been killed by the eastern spruce beetle during the past half century in the northeastern United States and southeastern Canada (p. 58).

A large percentage of the matured Engelmann spruce of the Rocky Mountains region has been killed by the Engelmann spruce beetle within the past fifty years (p. 59).

A large percentage of the pine timber of merchantable size in the Black Hills National Forest and other national and private forests of the central Rocky Mountains region has been killed during the past ten years by the Black Hills beetle (p. 59).

A large percentage of the best matured pine timber of the region north and west of Colorado and Utah has been killed within the past twenty years by the mountain pine beetle and the western pine beetle (p. 59).

A large percentage of the matured Douglas fir, or red fir, of the Rocky Mountains region has been killed by the Douglas fir beetle (p. 59).

The supply of hickory timber in the forests and woodlots of the States east of the Mississippi River has been greatly reduced by the ravages of the hickory barkbeetle (p. 60).

Practically all of the matured eastern larch has been killed over vast areas in the northeastern United States and southeastern Canada by the larch worm and eastern larch beetle (p. 60).

The wood of living timber has been rendered defective by wood-boring insects to such an extent as to reduce the value of a vast amount of standing timber from 50 to 75 per cent (p. 60).

Rapid deterioration of the wood of dying and dead trees has been caused by wood-boring insects, often amounting to from 25 to 100 per cent during the period in which it would otherwise be available for utilization (p. 62).

Crude manufactured and finished forest products have been damaged by insects to such an extent as to cause an estimated average annual loss of 10 per cent of its mill value (pp. 64-67).

Insects are the cause of greatly reducing our forest resources by killing the inaccessible timber; by reducing the quantity through injuries to the wood of living and dying timber; by preventing normal reproduction and development of future supplies, and through destroying forest products.

GENERAL ESTIMATES OF AMOUNT OF DAMAGE CAUSED BY FOREST INSECTS.

The results of extensive observations during the past ten years in the principal forested areas of all sections of the country, and during an additional eight years in West Virginia, indicate to the writer that the amount of standing timber killed by insects, together with

the standing living, dying, and dead timber reduced in quality and value by them in the forests of the country, to be found at any given time, has been not far from 10 per cent of the total stand of merchantable-sized timber (pp. 70-71).

Considering the forest in its broadest sense as a source of national wealth of a given value for all purposes, including direct utilization, protection of land from erosion, protection of headwater streams, protection of game, and as contributing to the real and æsthetic value of health and pleasure resorts, it is evident to the writer that the total damage caused by insects has been equivalent to an average additional 5 per cent of the value of the merchantable-sized timber of the entire country (p. 71).

Considering the problem of insect damage to standing timber and crude products on the basis of direct utilization of the forest resources, it is evident that the reduction in value below that of healthy timber or sound products at the time of utilization, including losses from handling defective material, has amounted to an equivalent of at least 10 per cent of the average annual mill value of the aggregate output of forest products of all kinds. This, of course, includes the killed and damaged merchantable-sized timber considered under the estimate relating to standing timber, given above. Since the killed and damaged standing timber is involved in any given annual output, this estimate on a basis of utilization represents more nearly a direct reduction in cash values (p. 71).

The writer estimates that the annual loss caused by insects injurious to finished and utilized products, including the consequent increased drain on the forest resources to replace that prematurely destroyed by insects, is equivalent to at least 3 per cent of the original or mill value.

HOW LOSSES CAN BE PREVENTED.

The results of extensive investigations and of practical applications during recent years have demonstrated that some of the most destructive insect enemies of American forests and of manufactured and utilized products can be controlled and serious damage prevented with little or no ultimate cost over that involved in good forest management and business methods.

It is evident that if the information now available through publications of the Department of Agriculture and through direct correspondence with its experts is properly utilized in the future it will result in the prevention of an equivalent of at least 30 per cent of the estimated annual waste of forest resources that has been caused by insects within recent years and thus contribute greatly to the conservation of the forest resources. This can be accomplished as follows:

(1) By the adoption or adjustment of certain requisite details in forest management, in lumbering and manufacturing operations, and in storing, transporting, and utilizing the products which, at the least expense, will bring about the necessary reduction of the injurious insect and unfavorable conditions for their future multiplication or destructive work.

(2) By the adoption of policies of control, based upon expert technical knowledge or advice relating to the species, habits, life history, and natural enemies of the insects involved, and methods for their control, supplemented by expert knowledge or advice on the principles of technical and applied forestry in the proper management, care, and utilization of the forest and its resources and still further supplemented by practical knowledge and experience relating to local conditions and facilities favorable and unfavorable for successful application according to a given method or policy of control.

(3) By reliance on technical advice furnished by recognized experts in forest entomology and forestry as a basis for success in practical application by the owner or forester.

(4) By utilization of so-called matured timber, and especially dense or pure stands of such timber, thus removing one of the favorable conditions for rapid deterioration through attacks by wood-boring insects or death through the attack of destructive bark-boring or defoliating insects.

(5) By the utilization of a knowledge of the principles of natural control as a means of contributing to the efficiency of artificial control.

(6) By prompt recognition of the first evidences of the work or destructive outbreaks of the principal insect depredators, by authentic identification of the species involved, and by prompt action in adopting the proper method or methods of control for the prevention of losses.

It should be remembered that as a rule it is useless to attempt the extermination of an insect enemy of the forest or its products. It is only necessary to reduce and weaken its forces at least 75 per cent, so that it can not continue an aggressive invasion, but must occupy a defensive position against its own enemies and become dependent upon favorable conditions resulting from avoidable negligence and mismanagement by the owners of the forests and the manufacturers of forest products.

While beneficial insects, beneficial birds, and beneficial diseases exert a continuous and powerful influence toward the prevention of a more extensive waste of forest resources, it has been repeatedly demonstrated that they can not be depended upon always to prevent widespread devastations or otherwise to work for the best interests of the private or public owner by protecting the best trees and the best tree species.

The best way to utilize the factors of natural control is to become their allies and assist in the reduction of the enemy, rather than to try to make them our allies through artificial introduction or dissemination.

A large percentage of the waste caused by insects can be prevented by the utilization of infested material, and at the same time, without additional expense, this will contribute greatly to the control of insects which cause such waste and also prevent injuries and depredations in the future.

Under past conditions the poor management or neglect of the average forest has contributed to the increase of depredations by insects.

Under present conditions of better management of local forests and of the more progressive manufacturing enterprises much is being accomplished toward the reduction of losses. In the average forest, and in the average business enterprise dealing with forest products, present conditions are little better than in the past. This is largely due to a lack of appreciation of the importance of the subject and failure to realize the opportunity and practicability of preventing a large percentage of the loss.

PUBLICATIONS RELATING TO FOREST INSECTS.

HISTORICAL AND GENERAL.

1831. WILSON, ALEXANDER, and CHARLES LUCIEN BONAPARTE.—American Ornithology. (Robert Jameson.) Edinburgh. Vol. I, pp. 133-134.
1876. PECK, C. H.—The black spruce. <Trans. Albany Inst., VIII, pp. 283-301.
1878. PECK, C. H.—30th Rep., N. Y. State Museum.
1882. HOUGH, F. B.—Report on Forestry, submitted to Congress by the Commissioner of Agriculture, Part VIII. Insect Ravages, pp. 259-274.
1. The injuries done to spruce and other coniferous timber by insects.
1882. STRETCH, R. H.—Notes on *Pieris menapia* Felder. <Papilio, II, pp. 103-110, Pl. II.
1883. SCUDDER, S. H.—Mass. Soc. for the Prom. of Agr. Boston. (Pamphlet.)
1884. FYLES, THOMAS W.—*Nematus crichsonii*. <14th Rep. Ent. Soc. Ont. for 1883, p. 17.
1884. FLETCHER, JAMES.—Forest trees. <Can. Dept. Agr., Rep. Ent. for 1884, p. 7.
1885. FLETCHER, JAMES.—The larch saw-fly. <15th Ann. Rep. Ent. Soc. Ont., pp. 72-77.
1887. JACK, JOHN G.—Notes of 1885 on some injurious and other common insects. <17th Ann. Rep. Ent. Soc. Ont., p. 17.
1888. FLETCHER, JAMES.—Larch saw-fly. <Exp. Farms Canada, Rep. Ent. and Bot. for 1887, p. 40.
1889. LINTNER, J. A.—*Nematus crichsonii* Hartig. <5th Rep. Inj. and other Ins., State of N. Y., pp. 164-173, fig. 4.
1890. PACKARD, ALPHEUS S.—Insects injurious to forest and shade trees. Revised and enlarged edition of Bul. 7. <5th Rep. U. S. Ent. Comm., U. S. Dept. Agr.
1891. FYLES, THOMAS W.—*Nematus crichsonii*. A retrospect. <22d Ann. Rep. Ent. Soc. Ont., pp. 28-30.
1893. LINTNER, J. A.—*Nematus crichsonii* Hartig. <8th Rep. Inj. and other Ins., State of N. Y. for 1891, pp. 168-169, fig. 22.
1897. HOWARD, L. O.—Damage by the white pine butterfly. <Bul. 7, n. s., Div. Ent., U. S. Dept. Agr., pp. 77-78.
1898. FISKE, W. F.—Bul. 17, n. s., Div. Ent., U. S. Dept. Agr., pp. 67-69.
1899. HUTT, H. L.—The larch saw-fly. <29th Ann. Rep. Ent. Soc. Ont., pp. 94-95.
1899. PINCHOT, GIFFORD.—A primer of forestry. <Bul. 24, Part I, Div. For., U. S. Dept. Agr., pp. 73-74.
1900. CARY, AUSTIN.—Insect damage to spruce timber in Maine and New Hampshire. <The Forester, VI, No. 3, pp. 52-54.
1901. FLETCHER, JAMES.—Pine butterfly. <Proc. Ent. Soc. Wash., IV, pp. 30-31.
1904. JARVIS, F. D.—Larch saw-fly. <34th Ann. Rep. Ent. Soc. Ont. for 1903, p. 100.

1906. FLETCHER, JAMES.—Forest and shade trees. The larch saw-fly. <Can. Dept. Agr., Centr. Exp. Farm, Rep. Ent. and Bot. for 1905, pp. 189, 190, 191.
1906. FYLES, THOMAS W.—The tamarack growth in the eastern townships of the Province of Quebec. <Can. For. Journ., II, pp. 132-134.
1907. FYLES, THOMAS W.—In the tracks of *Nematus erichsonii* Hartig. <37th Ann. Rep. Ent. Soc. Ont. for 1906, pp. 105-106.
1908. COOLEY, R. A.—The Douglas spruce cone moth (*Cydia pseudotsugana* Kearfott). <Proc. Am. Assn. Econ. Ent., pp. 125-130.
1909. BRUNER, LAWRENCE.—Journ. Econ. Ent., II, No. 1, pp. 52-53, February.
1909. FORBES, S. A.—Journ. Econ. Ent., II, No. 1, pp. 51-52, February.

PREPARED BY OFFICIALS OF THE BRANCH OF FOREST INSECT INVESTIGATIONS, BUREAU OF ENTOMOLOGY, AND OF THE WEST VIRGINIA AGRICULTURAL EXPERIMENT STATION.

1890. HOPKINS, A. D.—First report of the entomologist. <3rd Ann. Rep. W. Va. Agr. Exp. Sta., pp. 145-170.
- 1891a. HOPKINS, A. D.—Second report of the entomologist. <4th Ann. Rep. W. Va. Agr. Exp. Sta., pp. 59-64.
- 1891b. HOPKINS, A. D.—Forest and shade tree insects. I. Yellow locust. <Bul. 16, W. Va. Agr. Exp. Sta., pp. 85-91, 1 pl., April.
- 1891c. HOPKINS, A. D.—Forest and shade tree insects. II. Black spruce. <Bul. 17, W. Va. Agr. Exp. Sta., pp. 93-102, 1 pl., May.
- 1892a. HOPKINS, A. D.—Notes on a destructive forest-tree scolytid. <Science, XX, pp. 64-65, July 29.
- 1892b. HOPKINS, A. D.—The pine beetle of the Virginias. <Hardwood, II, p. 7, November 25.
- 1892c. HOPKINS, A. D.—The first announcement of the importation of the European bark-beetle destroyer into America. <Proc. Ent. Soc. Wash., II, p. 353.
- 1892d. HOPKINS, A. D.—Third report of the entomologist. <5th Ann. Rep. W. Va. Agr. Exp. Sta., pp. 40-46.
- 1893a. HOPKINS, A. D.—Fourth report of the entomologist. <6th Ann. Rep. W. Va. Agr. Exp. Sta., pp. 29-48.
- 1893b. HOPKINS, A. D.—Damage to forests by the destructive pine barkbeetle. <Insect Life, V, pp. 187-189, January.
- 1893c. HOPKINS, A. D.—Pinholes in chestnut. <Hardwood, III, p. 5, 1 fig., February.
- 1893d. HOPKINS, A. D.—Catalogue of West Virginia Scolytidae and their enemies. <Bul. 31, W. Va. Agr. Exp. Sta., pp. 121-168, April.
- 1893e. HOPKINS, A. D.—Catalogue of West Virginia forest and shade tree insects. <Bul. 32, W. Va. Agr. Exp. Sta., pp. 171-251, May.
- 1893f. HOPKINS, A. D.—Notes on timber worms and bark borers. <Proc. Ent. Soc. Wash., III, p. 82.
- 1893g. HOPKINS, A. D.—Notes on food habits of *Corthylus punctatissimus*. <Proc. Ent. Soc. Wash., III, pp. 104, 107.
- 1893h. HOPKINS, A. D.—Short notes on list of pine-infesting insects. <Proc. Ent. Soc., Wash., III, pp. 192-193.
- 1893i. HOPKINS, A. D.—Destructive scolytids and their imported enemy. <Insect Life, VI, pp. 123-129, December; also 24th Ann. Rep. Ent. Soc. Ont., p. 71.
- 1894a. HOPKINS, A. D.—Defects in wood caused by insects. <Bul. 35, W. Va. Agr. Exp. Sta., pp. 291-306, 26 figs., January.

- 1894b. HOPKINS, A. D.—Black holes in wood. <Bul. 36, W. Va. Agr. Exp. Sta., pp. 311-336, 43 figs., 2 pls., February.
- 1894c. HOPKINS, A. D.—The relations of insects and birds to present forest conditions. <Garden and Forest, p. 348.
- 1894d. HOPKINS, A. D.—Insect enemies of the yellow poplar. <The Timberman, pp. 8, figs. 11, July.
- 1894e. HOPKINS, A. D.—Notes on some discoveries and observations of the year in West Virginia. <Insect Life, VII, pp. 145-151, October.
- 1894f. HOPKINS, A. D.—Serious trouble over. <Southern Lumberman; Hardwood, VI, pp. 270-271, November; Timberman.
- 1894g. HOPKINS, A. D.—Some interesting conditions in wood resulting from the attack of woodpeckers and insects. Abstract. <Proc. Amer. Assn. Adv. Sci., p. 252.
- 1894h. HOPKINS, A. D.—Fifth report of the entomologist. <7th Ann. Rep. W. Va. Agr. Exp. Sta., pp. 34-44.
- 1895a. HOPKINS, A. D.—Sixth report of the entomologist. <8th Ann. Rep. W. Va. Agr. Exp. Sta., pp. 27-35.
- 1895b. HOPKINS, A. D.—The northern timberworm of the grub species. <Southern Lumberman, p. 7, 5 figs., January.
- 1895c. HOPKINS, A. D.—On the study of forest-tree insects. <Proc. Assn. Econ. Ent. In Bul. 2, n. s., Div. Ent., U. S. Dept. Agr., pp. 75-79; 26th Ann. Rep. Ent. Soc. Ont., pp. 80-83.
- 1896a. HOPKINS, A. D.—Seventh report of the entomologist. <9th Ann. Rep. W. Va. Agr. Exp. Sta., pp. 66-164, pls. 5, figs. 2, maps 3.
- 1896b. HOPKINS, A. D.—Some notes on insect enemies of trees. <Can. Ent., XXVIII, pp. 243-250, October.
1897. HOPKINS, A. D.—Eighth report of the entomologist. <10th Ann. Rep. W. Va. Agr. Exp. Sta., pp. 42-57.
- 1898a. HOPKINS, A. D.—The periodical cicada in West Virginia. <Bul. 50, W. Va. Agr. Exp. Sta., pp. 46, figs. 23, map 1, January.
- 1898b. HOPKINS, A. D.—On the history and habits of the "wood engraver" ambrosia beetle, *Xyleborus xylographus* Say (*Xyleborus sareseni* Ratz.), with brief description of different stages. <Can. Ent., XXX, pp. 21-29, pls. 2, February. (= *Xyleborus sareseni* Ratz., A. D. H., 1909.)
- 1898c. HOPKINS, A. D.—Insect enemies of the locust tree. <W. Va. Farm Review, pp. 88-93, figs. 6, March.
- 1898d. HOPKINS, A. D.—Insects detrimental and destructive to timber products. <Proc. 19th Ann. Meet. Soc. Prom. Agr. Sci., pp. 103-108.
- 1898e. HOPKINS, A. D.—Ninth report of the entomologist. <11th Ann. Rep. W. Va. Agr. Exp. Sta., pp. 32-40.
- 1899a. HOPKINS, A. D.—Tenth report of the entomologist. <12th Ann. Rep. W. Va. Agr. Exp. Sta., pp. 25-36.
- 1899b. HOPKINS, A. D.—Report on investigations to determine the cause of unhealthy conditions of the spruce and pine from 1880 to 1893. <Bul. 56, W. Va. Agr. Exp. Sta., pp. 197-461, figs. 99, April.
- 1899c. HOPKINS, A. D.—Preliminary report on the insect enemies of forests in the Northwest. <Bul. 21, n. s., Div. Ent., U. S. Dept. Agr., pp. 27, December.
- 1900a. HOPKINS, A. D.—Eleventh report of the entomologist. <13th Ann. Rep. W. Va. Agr. Exp. Sta., pp. 15-17.
- 1900b. HOPKINS, A. D.—The pin-hole borers and how to prevent losses from the ravages of the oak timber worm. <Southern Lumberman, XXVII, No. 427, p. 425.

- 1900c. HOPKINS, A. D.—The periodical cicada or seventeen-year locust in West Virginia. A revision of Bul. 50, with additional data relating to the 1898 and 1900 swarm. <Bul. 68, W. Va. Agr. Exp. Sta., pp. 267–330, figs. 14, pls. 3, maps 9, September.
- 1901a. HOPKINS, A. D.—Insect enemies of the spruce in the Northeast. <Bul. 28, n. s., Div. Ent., U. S. Dept. Agr., pp. 48, pls. 16, 56 orig. figs., 5 text figs.
- 1901b. HOPKINS, A. D.—On the development and evolution of the scolytid gallery. [Paper read before Section F, Amer. Assn. Adv. Sci., Denver, Colo. (August).] Abstract published in Science.
- 1901c. HOPKINS, A. D.—On insects detrimental and destructive to forest products used for construction material. [Paper read at Denver Meet. Amer. Assn. Econ. Ent. (August).] <Bul. 31, n. s., Div. Ent., U. S. Dept. Agr., pp. 60–62.
- 1901d. HOPKINS, A. D.—Insect enemies of forests and forest products. [Paper read at Denver Meet. Amer. Forestry Assn.] <The Forester, VIII, pp. 250–254, 5 figs.
- 1902a. HOPKINS, A. D.—Insect enemies of the pine in the Black Hills Forest Reserve. <Bul. 32, n. s., Div. Ent., U. S. Dept. Agr., pp. 24, pls 7, figs. 18, 5 text figs.
- 1902b. HOPKINS, A. D.—On the study of forest entomology in America. <Bul. 37, n. s., Div. Ent., U. S. Dept. Agr. (Proc. 14th Ann. Meet. Assn. Econ. Ent., pp. 5–28. Reprint, *ibid*).
- 1902c. HOPKINS, A. D.—Some insect enemies of living trees. <Proc. 22d Ann. Meet. Soc. Prom. Agr. Sci., pp. 66–69.
- 1902d. HOPKINS, A. D.—The powder-post beetle. <Amer. Agriculturist, November 15; New England Homestead, November 1.
- 1903a. HOPKINS, A. D.—Some of the principal insect enemies of coniferous forests in the United States. <Yearbook U. S. Dept. Agr. for 1902, pp. 265–282, figs. 10, pls. 2. (Separate No. 268, Office of Secy. of Agr.)
- 1903b. HOPKINS, A. D.—Forest insect explorations in the summer of 1902. <Can. Ent., XXXV, pp. 59–61, March.
- 1903c. HOPKINS, A. D.—Powder-post injury to seasoned wood products. <Cir. 55, Bur. Ent., U. S. Dept. Agr., pp. 5, fig. 1.
- 1903d. HOPKINS, A. D.—Methods of work and some results in forest insect investigations. <Proc. Assn. Agr. Colls. and Exp. Stas., pp. 180–182.
- 1903e. HOPKINS, A. D.—Insect enemies of the redwood. <Bul. 38, Bur. Forestry, U. S. Dept. Agr., pp. 32–40, figs. 4.
- 1904a. HOPKINS, A. D.—Catalogue of exhibits of insect enemies of forests and forest products at the Louisiana Purchase Exposition. <Bul. 48, n. s., Div. Ent., U. S. Dept. Agr., pp. 56, pls. 22.
- 1904b. HOPKINS, A. D.—Insect injuries to hardwood forest trees. <Yearbook U. S. Dept. Agr. for 1903, pp. 313–328, figs. 17, pl. 1. (Separate No. 327, Office of Secy. of Agr.)
1905. BURKE, H. E.—Black check in western hemlock. <Cir. 61, Bur. Ent., U. S. Dept. Agr., pp. 10, figs. 5.
- 1905a. HOPKINS, A. D.—Insect injuries to forest products. <Yearbook U. S. Dept. Agr. for 1904, pp. 381–398, figs. 14. (Separate No. 355, Office of Secy. of Agr.)
- 1905b. HOPKINS, A. D.—The Black Hills beetle, with further notes on its distribution, life history, and methods of control. <Bul. 56, Bur. Ent., U. S. Dept. Agr., pp. 24, figs. 6.

1906. FISKE, W. F.—A new species of the curculionid genus *Paraplinthus*. <Proc. Ent. Soc. Wash., VIII, pp. 31–32.
- 1906a. HOPKINS, A. D.—Barkbeetle depredations of some fifty years ago in the Pike's Peak region of Colorado. <Proc. Ent. Soc. Wash., VIII, pp. 4–5.
- 1906b. HOPKINS, A. D.—Some insects injurious to forests. The locust borer. <Bul. 58, Part I, Bur. Ent., U. S. Dept. Agr., pp. 16, figs. 6, pl. 1.
- 1906c. HOPKINS, A. D.—Insect enemies of forest reproduction. <Yearbook U. S. Dept. Agr. for 1905, pp. 249–256, figs. 9. (Separate 381, Office of Secy. of Agr.)
1906. WEBB, J. L.—Some insects injurious to forests. The western pine-destroying barkbeetle. <Bul. 58, Part II, Bur. Ent., U. S. Dept. Agr., pp. 17–30, figs. 7–12, Pls. II, III.
1907. BURKE, H. E.—Notes on the larva of *Calopus angustus* Lec. <Proc. Ent. Soc. Wash., VIII, pp. 64–66.
- 1907a. HOPKINS, A. D.—Some insects injurious to forests. Additional data on the locust borer. <Bul. 58, Part III, Bur. Ent., U. S. Dept. Agr., pp. 31–40.
- 1907b. HOPKINS, A. D.—Pinhole injury to girdled cypress in the South Atlantic and Gulf States. <Cir. 82, Bur. Ent., U. S. Dept. Agr., pp. 4, fig. 1.
- 1907c. HOPKINS, A. D.—The locust borer and methods for its control. <Cir. 83, Bur. Ent., U. S. Dept. Agr., pp. 8, figs. 4.
- 1907d. HOPKINS, A. D.—The white pine weevil (*Pissodes strobi* Peck). <Cir. 90, Bur. Ent., U. S. Dept. Agr., pp. 8, figs. 6.
1908. BURKE, H. E.—A new buprestid enemy of *Pinus edulis*. <Proc. Ent. Soc. Wash., IX, pp. 117–118, fig. 6.
1908. FISKE, W. F.—Notes on insect enemies of wood-boring Coleoptera. <Proc. Ent. Soc. Wash., IX, pp. 23–27.
- 1908a. HOPKINS, A. D.—Notable depredations by forest insects. <Yearbook U. S. Dept. Agr. for 1907, pp. 149–164. (Separate 442, Office of Secy. of Agr.)
- 1908b. HOPKINS, A. D.—Work of the Bureau of Entomology against forest insects. <Journ. Econ. Ent., I, No. 6, pp. 343–348, December.
- 1909a. HOPKINS, A. D.—Contributions toward a monograph of the scolytid beetles. I. The genus *Dendroctonus*. <Tech. Ser. 17, Part. I, Bur. Ent., U. S. Dept. Agr., pp. xiii+164, pls. 8, text figs. 95.
- 1909b. HOPKINS, A. D.—Practical information on the scolytid beetles of North American Forests. I. Barkbeetles of the genus *Dendroctonus*. <Bul. 58, Part I, Bur. Ent., U. S. Dept. Agr., pp. 169, pls. II, text figs. 102.
- 1909c. HOPKINS, A. D.—Journ. Econ. Ent., II, No. 1, pp. 49–53, February.
1909. WEBB, J. L.—Some insects injurious to forests. The southern pine sawyer. <Bul. 58, Part IV, Bur. Ent., U. S. Dept. Agr., pp. 41–56, figs. 13–24.

PUBLICATIONS RELATING TO FOREST STATISTICS.

FOREST RESOURCES.

1907. NORTH, S. N. D., AND PINCHOT, GIFFORD.—The lumber cut of the United States, 1907. <Forest Products, No. 2, Dept. Commerce and Labor, Bur. Census. Compiled in cooperation with the Dept. of Agr., For. Serv.
1909. KELLOGG, R. S.—The timber supply of the United States. <Cir. 166, For. Serv., U. S. Dept. Agr., pp. 6 and 8.

FOREST FIRES.

1903. PINCHOT, GIFFORD.—Rep. of the Forester for 1903. <From Annual Reports, U. S. Dept. Agr.
1903. SUTER, H. M.—Forest fires in the Adirondacks in 1903. <Cir. 26, For. Serv., U. S. Dept. Agr.
1904. STERLING, E. A.—Attitude of lumbermen toward forest fires. <Yearbook U. S. Dept. Agr. for 1904, p. 134.
1907. CRAFT, Q. R.—The progress of forestry in 1907. <Yearbook U. S. Dept. Agr. for 1907, p. 568.
1908. CLEVELAND, T., JR.—The progress of forestry in 1908. <Yearbook U. S. Dept. Agr. for 1908, p. 541.
1908. WILSON, JAMES.—Rep. Secretary of Agr. for 1908, p. 75.
1909. CLEVELAND, T., JR.—The status of forestry in the United States. <Cir. 167, For. Serv., U. S. Dept. Agr., p. 23.

O

VOL. III

FEBRUARY, 1930

NO. 1

BULLETIN
OF
THE NEW YORK
STATE COLLEGE OF FORESTRY
AT
SYRACUSE UNIVERSITY

Nelson C. Brown, Acting Dean

Technical Publication No. 28

The White Pine Weevil (*Pissodes
strobi* Peck)—Its Biology
and Control

by

Harvey J. MacAloney

TRUSTEES OF THE NEW YORK STATE COLLEGE OF FORESTRY

Ex-Officio

HON. HERBERT H. LEHMAN.....Albany, N. Y.
Lieutenant Governor

DR. FRANK P. GRAVES.....Albany, N. Y.
Commissioner of Education

DR. CHARLES WESLEY FLINT.....Syracuse, N. Y.
Chancellor

HON. ALEXANDER MACDONALDAlbany, N. Y.
Conservation Commissioner

Appointed by Governor

HON. ALFRED E. SMITHNew York City

HON. EDMUND H. LEWIS.....Syracuse, N. Y.

HON. JOHN R. CLANCY.....Syracuse, N. Y.

HON. HAROLD D. CORNWALL.....Glenfield, N. Y.

HON. GEORGE W. DRISCOLL.....Syracuse, N. Y.

HON. WILLIAM KELLEY.....Syracuse, N. Y.

HON. EDWARD H. O'HARA.....Syracuse, N. Y.

HON. J. HENRY WALTERS.....New York City

HON. CHARLES A. UPSONLockport, N. Y.

President

ALFRED E. SMITH

Vice-President

JOHN R. CLANCY

CONTENTS

	PAGE
Introduction.....	5
General Position in Classification.....	10
Historical References.....	10
Description of the Insect.....	10
Adult.....	10
Egg.....	12
Larva.....	12
Pupa.....	12
Life History and Habits.....	12
Hibernation.....	12
Resumption of Activity.....	14
Egg Laying.....	16
Incubation Period and Larval Activity.....	18
Pupal Stage.....	20
Habits of the New Generation.....	20
Feeding.....	21
Copulation.....	21
Flight Habits.....	23
Host Trees.....	25
Indications of Injury and Character of the Damage.....	27
Effect on Axial and Radial Growth.....	28
Loss in Axial Growth.....	28
Effect on Radial Growth.....	29
Effect on Quality of Timber.....	31
Injuries often Mistaken for Weeviling.....	33
Summarization of Plot Data.....	34
Manner in Which Study was Carried Out.....	34
Data on Pure Stands.....	34
Data on Mixed Stands.....	36
Data on Other Species Liable to Attack.....	39
Effect of Silvical and Habitat Factors.....	41
Composition of the Stand.....	41
Age, Height and Vigor.....	43
Site Factors.....	45
Exposure.....	45
Climatic Factors.....	47
Scarcity of Food Supply as a Limiting Factor.....	48
Control Measures.....	49
Natural Control.....	50
The Climate.....	50

Control Measures — <i>Continued</i> :	PAGE
Parasitic and Predatory Insects.....	50
Parasites.....	50
Hyperparasites.....	50
Predacious Insects.....	50
Associated Species.....	51
Birds and Other Animals.....	55
Prevention of Attack by the Insect.....	57
Sprays and Repellents.....	57
Banding Materials.....	58
Sticky tree-banding material.....	58
Raw Wool.....	58
Direct Control of the Insect.....	60
Picking the Beetles by Hand.....	60
Jarring the Insect into a Net.....	60
Removal of the Infested Leaders.....	61
Repair of the Injury by Silvicultural Methods.....	63
Density of Stocking.....	63
Mixed Stands.....	65
Application of Control Measures to all Species Liable to Attack.....	67
Summary.....	67
Bibliography.....	70
Graphs and Plates.....	78

THE WHITE PINE WEEVIL (*PISSODES STROBI* PECK), ITS BIOLOGY AND CONTROL

Harvey J. MacAloney

Introduction

Forest Entomology is taking its place as one of the most important branches of scientific forestry. Every forester or manager in charge of woodlands should possess at least a general knowledge of the most important injurious forest insects of his region and the means of controlling them, or at least of minimizing the damage done by them. This information is fully as essential to the forester as is a knowledge of the best species of tree to plant, according to the soil and other silvicultural conditions prevailing in the particular region. In managing a forest all the biological factors, including soil-composition and moisture, humidity, rainfall, aspect, exposure and altitude, and liability to insect and fungus injury should be considered.

Owing to economic conditions, brought about by increasing population and higher standards of living, we have ever-increasing land values, and therefore it is essential that these lands be managed properly. The correct management of forest lands is necessary to the economic life of the nation. Public sentiment, if it be properly directed, should prevent abuses in timber production just as in any other business. By helpful and intelligent suggestions the forester can aid in solving the problem of timber depletion, and the growth now on the land can, by careful handling, be cut in such a way as to insure its replacement by young growth. The really essential problem is that a forest be maintained on the land, and that those areas which are better suited for growing trees than for other purposes be afforested. The people of the nation are becoming insistent that a supply of timber which will be adequate for the nation's use be assured. The proper management of the forest is one means of insurance that this demand be satisfied. The accomplishment of this desired end involves not only correction of certain practices of destructive lumbering but also protection against excessive destruction by windfall, fire, disease and insect attack.

Until recently the importance of the destructive actions of forest insects has been under-emphasized. Few people have realized that these pests do more damage each year, not only by the actual killing of living timber, but also by the destruction or reduction of timber value, than is done by any other single agency. The injury usually is not so spectacular as that caused, for example, by a forest fire. In consequence the popular fallacy has been prevalent, and still exists, that there is no necessity for control work unless an epidemic is actually under way.

The annual loss due to insect attacks upon living trees and upon rough, finished or utilized forest products cannot be accurately estimated. In 1910 Dr. A. D. Hopkins, at that time chief of the Division of Forest Insects, Bureau of Entomology, estimated the annual loss due to insect depredations upon forests and forest products as some \$100,000,000. This would be a very low estimate for the present time, because insect injuries have increased rather than diminished and at the same time the value of timber has increased greatly. Since 1910 there have been severe outbreaks of the Spruce Budworm in Maine, the Lake States, the Pacific States and in Canada from the East to the West. The Gipsy Moth still takes its toll in New England; the *Dendroctonus* beetles do immense damage in almost all parts of the South and West; and wood borers, powder-post beetles, termites and numerous others contribute to the growing total of damage.

The following statement by Jaenicke (1921) will show exact figures for one species alone; "Perhaps a single instance of the tremendous loss which the western pine beetle (*Dendroctonus brevicomis* Lec.) can inflict on the yellow pine of this region will suffice. One of the finest bodies of western yellow pine in the West is located in southern Oregon in the vicinity of Klamath Falls. It is a privately owned body of timber covering an acreage of about 600,000 acres, and a volume of at least ten billion feet. Thirty million dollars is a conservative valuation of the property. During the past ten years the forest fire loss on this area amounted to only \$15,000, because of the efficacy of the fire protection plan. During this same ten-year period there has been a \$3,000,000 loss for which the western pine beetle has been entirely responsible. In other words, during the past ten years the western pine beetle has killed ten per cent of the stand. This killing is still going on."

The Pales weevil (*Hylobius pales* Herbst) often does great damage to young white pine seedlings in regions where the white pine is common. The loss cannot be even approximately estimated, as the value of a small seedling cannot be considered in terms of the mature forest, except by the manipulation of complicated formulæ and even then it is theoretical.

Thus it can be seen that even an approximation, or a guess, of the loss caused annually by forest insects will not be of any great value. It has, however, been estimated by different authorities as somewhere between \$100,000,000 and \$250,000,000.

The Bureau of Entomology during the past twenty years has made noteworthy progress in determining the life-histories of destructive forest insects; in devising means of combatting them; and in meeting emergencies by educational work and systematic control measures. Tree-killing insects, insects causing reduced growth and others causing injuries resulting in losses in the timber value of the trees take a large toll of the forests, and in their more serious outbreaks often threaten extensive forest regions.

The subject of the ecological relations of forest insects is an extremely complicated one. It covers a field involving many fundamentally different conditions of site and of tree associations, any one or more members of which may be subject to attack by insects. The presence of any one condition of site or of host or of insect, or a combination of various conditions and insects, may so modify the forest that the ultimate forest association may be entirely different from that which would prevail under another set of conditions. There are some defoliating insects such as the spruce budworm (*Cacoecia fumiferana* Clem.) which, because of an abnormal amount of food material, become epidemic, and by repeated defoliations kill the trees. Such attacks are in turn followed by invasions of secondary insects, the natural function of which is that of merely assisting in converting the dead wood to humus. These are the wood borers which often do great damage to manufactured lumber, rendering it unfit for utilization. In addition to these, there are some insects such as the White Pine Weevil (*Pissodes strobi* Peck) which normally do not kill the tree itself, but kill back a portion of it so as to greatly reduce its ultimate lumber value.

The White Pine Weevil is the most serious insect pest affecting the white pine, and in some localities it is the most serious of all pests. This weevil is a native insect and has always been of importance throughout the range of the white pine, although, as Butterick (1916) states, "Many people thought the white pine in its younger stages had almost no enemies." The first known mention of its economic importance was made by Peck in 1817 when he described the insect. The great white pines were probably among the first green things seen by the Pilgrim Fathers when they landed at Plymouth Rock in 1620, and doubtless some of these trees had been weeviled. At the time Peck (1817) wrote, the manufacture of wooden ships was at its height and one of the factors limiting the suitability of white pine for use as masts was the injury and consequent crook caused by white pine weevil attack. Peck stated that, "The species which is so injurious to the white pine is, I believe, pretty generally diffused over our country and is undescribed." He goes on further to say that if it were not for the parasites which prey upon this insect, "Our forests would scarce produce a single mast." Many of these trees were two hundred years old when they were taken out of the forests for masts, and in the same stands were other trees of the same age, rejected because of their crookedness.

Although Peck commented on its numerical abundance when he described the weevil, a half century later conditions became still more favorable for rapid multiplication. This is especially true in sections where abandonment of farms was a common occurrence. About 1850, in the New England States and New York, the farmers

began to abandon their lands and the peak was reached between the end of the Civil War and 1880. Many of the abandoned fields reverted to the forest types and were seeded in 'almost' wholly by white pine. These old field pine stands were in many cases poorly stocked, and the trees were weeviled repeatedly. Added to this acreage of old pine are the many acres of plantation which have been planted in the past thirty years. This gave abundant food for the weevil and all recent observers agree that the insect has been more abundant and the percentage of infested pine much greater in the past half century than before.

Various systems of artificial or natural regeneration have been used in the past thirty years. There are pure and mixed plantations in densities varying from 12' x 12' to 3' x 3'. There are natural stands which are pure or mixed in composition. There are stands which are essentially natural but in which regeneration is aided by silvicultural practice with or without the aid of additional planting. It has been found, however, in some regions that the growing of the white pine is almost an impossibility under the generally accepted ideas for handling this tree, owing to the ravages of two native insects—the White Pine Weevil and the Pales weevil. However, in course of time satisfactory methods have been developed for bringing young stands through to maturity in spite of these insects, and at the present time these practices give every promise of producing a crop that will go through a full rotation without appreciable injury, and at the same time will improve the forest environment and insure a stand of high grade material more adjustable to future markets.

The range of the White Pine Weevil is very similar to that of the white pine itself, but the insect is not so important in the more westerly or southerly portions as it is in the northeastern part of the United States and southeastern Canada and the northern portions of the Lake States.

This study was begun at the New York State College of Forestry in 1923 and carried on there until May, 1925. During this time the prosecution of the project was largely made possible through the cooperation given and the interest shown by Dean Frederick Franklin Moon. Continuance of the work was the result of the cooperation of certain timberland owners in Massachusetts who made available to the Bureau of Entomology of the United States Department of Agriculture a contribution of five thousand dollars for this work. Much of the life history and many control experiments were worked out at the Harvard Forest, Petersham, Massachusetts, and its vicinity, as there is a greater variety of silvical conditions in this locality than is usually found where weevil injury

is prevalent. The Agricultural Experiment Stations of the states of Connecticut and Vermont supplied temporary assistants to lay out sample plots in different conditions in their respective states, and the Entomological Branch of the Dominion of Canada Department of Agriculture cooperated by paying the salary and expenses of the writer, thereby making possible a study of three months' duration in eastern Canada in 1927. Permission to study permanent plots on the Yale Forest at Keene, New Hampshire, made available for observation forest conditions different from those which are found at the Harvard Forest forty miles to the south. During the autumn of 1928, a survey of conditions in eastern Pennsylvania was made possible through the aid of the Pennsylvania Department of Forests and Waters and the Pennsylvania Forest School at Mont Alto. The Biltmore Plantations at Asheville, North Carolina, and some natural stands near the Bent Creek

Figure I. Life History Stages of the White Pine Weevil.
(a) Adult (smaller figure natural size); (b) Pupa;
(c) Larva. A. D. Hopkins, Bureau of Entomology, U. S.
D. A.

Laboratory of the U. S. Bureau of Entomology were also studied at this time.

The author wishes to express his appreciation to all who made this investigation possible, and to convey his sincere thanks to Dr. F. C. Craighead, U. S. Bureau of Entomology; Dr. M. W. Blackman, New York State College of Forestry; Director R. T. Fisher and Assistant Director A. C. Cline, Harvard Forest, for their criticisms and suggestions in the prosecution of the investigation and the preparation of the manuscript. The determinations of the parasitic and predacious insects were made by Dr. J. M. Aldrich,

of the Division of Insects, U. S. National Museum, and by Dr. A. G. Boving and Messrs. A. Busck, A. B. Gahan, C. Heinrich, and S. A. Rohwer of the Division of Taxonomy, Bureau of Entomology, U. S. Department of Agriculture. The author is very grateful for their excellent cooperation. Thanks are also due Field Assistants D. DeLeon, R. C. Hall, L. Rintel and R. O. Hall who aided the writer in the taking of data and the study of the plots during the course of the field work.

GENERAL POSITION IN CLASSIFICATION

The White Pine Weevil (*Pissodes strobi* Peck) belongs to the order *Coleoptera*, suborder *Rhyncophora*, family *Curculionidae*, tribe *Pissodini*, genus *Pissodes*, species *strobi* Peck. Originally the genus *Pissodes* was included in the tribe, *Hylobiini* but Champion (1902) lists it as a separate tribe, *Pissodini*, with the single genus *Pissodes*. This tribe differs from the tribe *Hylobiini* in that the antennae are inserted at or just behind the middle of the beak instead of near the tip, and the antennal club has the first joint smooth and glabrous and not ringed or wholly pubescent. The front coxae also are separated and the prosternum is not emarginate or produced toward the sides, while in the *Hylobiini* the front coxae are contiguous and the prosternum is emarginate and produced toward the sides.

Historical References

The name *Pissodes* (pitch-colored) was proposed for this genus in 1871 by Germar to include certain old species. The genus itself was not described, however, until 1824, when Germar published a complete generic diagnosis together with a short description of *P. nemorensis*. The White Pine Weevil, a species distinct from *P. nemorensis* which is a southern form, was described by Peck in 1817 under the name *Rhynchaenus strobi*. He says, "In technical language it may be called *Rhynchaenus strobi*; femoribus muticis, rufogriseus; thoracis punctis duobus, scutello maculaque versus elytrorum apiceni albis." In 1831, Say wrote, "Dr. Harris sent me this insect as the *P. strobi*, or White pine Weevil of Professor Peck, whose name having the priority must be adopted." Since then it has been known as *Pissodes strobi* Peck.

The species name is derived from the specific name of the white pine (*Pinus strobus* L.) and from the fact that this tree is the favorite host. It was on this tree that the injury was first noticed and commented upon by Peck.

Description of the Insect

Adult:—An oblong-oval beetle, from 4 to 6 millimeters in length; the color ranges from dark to light brown with spots of white with two or more varying shades on each elytron. The

body covering is made up of small scales which vary to a considerable degree in size, density and color. The brown scales, which generally have a distinct-reddish cast, predominate but the white scales are found in irregular areas all over the ventral surface of the abdomen, thorax and legs. Across the posterior third of the elytra is a broad band composed of brownish-yellow scales at the margin and changing into white ones toward the suture. This band may or may not have a dark border.

There is considerable variation in the shape of the pronotum, and the head and prothorax together are slightly shorter than the elytra. In some specimens the pronotum is distinctly broader than long while in others it is subequal in breadth and length. The anterior margin may be entire or it may be slightly emarginate in the middle. The constriction of the pronotum varies from gradual to quite sharp. The pronotal punctures are quite irregular in size and are sometimes almost completely obscured by the flat or slightly, elevated areas surrounding them. The elytra are slightly broader than the pronotum, with the sides subparallel and the apex rounded. The third and fifth interspaces are more prominent than the others, and the third and ninth are joined around a distinct impression at the apex of the fourth, seventh and eighth interspaces. The fifth interspace is strongly elevated at the apex. The very distinct striae punctures may be irregular or regular in size. The head is distinctly narrower than the prothorax and is characterized by a long slender beak. This beak has the antennae inserted at or just behind the middle; the eyes are at the base of the beak and are widely separated. The legs are strong, almost equal in length and each one has an incurved spine on the inner apex of the tibia. The mouth parts are characteristically those generally found in the *Rhyncophora* in that the labrum and clypeus are absent and the epistoma is only represented externally by a smooth area faintly defined by an obscure line and lateral bristles.

The distinction between males and females of the genus is not easily visible to the naked eye, and the use of a hand lens is generally necessary to recognize the sex. In the *females* there are but seven visible abdominal tergites, the eighth being completely covered by the seventh which forms the pygidium and which is the last dorsal segment left exposed by the elytra. In order to see these segments it is necessary to spread the elytra and true wings, and this is difficult with living specimens. The beak is longer, smoother and more slender than in the males. The apical or seventh sternite is usually shorter than the two preceding sternites together which are generally less convex and more evenly punctured. The inner apical tooth of the tibiae is also smaller.

In the *males* there are eight visible abdominal tergites; the seventh is distinguished by the emarginate posterior margin, while the eighth is prominent, with the apex broadly rounded, and forms the pygidium. The abdominal sternites three and four are more convex than in the female, less evenly punctured and more shining toward the middle. The beak is stout, shorter, less shining and more distinctly punctured. The inner apical tooth of the tibiae is usually more prominent.

Egg: The egg is pearly white, subopaque, slightly oblong, equally rounded at both ends and about one millimeter in length.

Larva: The larva is a small, yellowish-white, footless grub, about seven millimeters long when full grown. There are three thoracic segments and ten abdominal segments (nine true segments and the anal lobes which represent the tenth). The head is small, light brown, with the mandibles and anterior margin dark brown; there are distinct eye spots. The row of yellowish dots on each side of the abdomen are spiracles. The antennae are very small, jointed and situated at the anterior extremity of the frontal suture. The larvae are characterized by having a distinct clypeus and labrum, which are absent in the adult.

The legitimacy of the separation, by Hopkins (1911), of *P. approximatus* as a distinct species from *P. strobi* has often been questioned. The writer was able to obtain authentic specimens of the larvae of each of these, and Dr. A. G. Böving, of the U. S. Bureau of Entomology, has very kindly called attention to the following main differences:

1. The head in *strobi* is shorter and more rounded, and the frons anteriorly narrower than in *approximatus*.

2. The spiracles in *strobi* are surrounded by a strong and dark not fully closed peritremal ring, but in *approximatus* the peritremal ring is weak and light.

3. The setae of the abdominal segments, and particularly on the ninth abdominal segment, are much shorter in *strobi* than in *approximatus*.

Pupa: The pupa is creamy white in general color; with the mandibles and eyes brown. As the pupa approaches maturity the beak and legs begin to turn brown and the general color of the adult is gradually assumed. The abdominal tip is square, armed with two spines, one on each side of the median line. The pupa is of the same length as the adult, while the larva is usually slightly longer.

LIFE HISTORY AND HABITS

Hibernation

The White Pine Weevil usually hibernates in the adult stage in the lowermost, well decomposed layer of the litter or duff and occasionally in the uppermost layer of the soil beneath trees fed

upon during the preceding autumn. These trees may have been weeviled that season or they may be trees to which one or more individuals of the new generation had migrated after leaving the larval host. The adults may also hibernate in the litter at the bases of trees cut late in the fall, probably attracted there by the odor of the escaping pitch.

Harris (1862) states that "The beetles are found in great numbers in April and May on fences, buildings and pine trees; they probably secrete themselves during the winter in the crevices of the bark, or about the roots of trees, and deposit their eggs in the spring; or they may not usually leave the trees until spring." Hopkins (1907), nearly fifty years later, wrote that "The winter is passed in the adult stage, evidently in the ground." Graham (1918) stated that the weevil hibernated in the litter beneath the trees. In the same year, in the Report of the Crown Lands Department of the Province of New Brunswick, it was anonymously reported that the adults hibernate in the litter beneath the trees. This was further corroborated in 1926 by Graham and by MacAloney. Graham reported that during the winter of 1914-1915 he succeeded in collecting the weevils from litter obtained from beneath pine trees. He used a modification of the Berlese trap. The litter was placed in the cone and heated slowly and on two occasions he obtained living weevils. The writer has never obtained weevils in this manner or by sifting material thru different meshes of screening, although many hours were spent on the latter operation in infested plantations, without success. However, in the spring, just after the frost was out of the ground and before any weevils were found on the leaders, little difficulty has been experienced in finding adults by digging away the upper layers of needles and examining the lower layers at the bases of trees fed on the previous fall. If this is done at the time root elongation has begun and before the beginning of axial elongation, the weevils can be found just at the surface of the mineral soil. They almost immediately become active on exposure to heat.

Experiments begun in 1925 and continued through the field season of 1927 show that there is a great mortality during the hibernation period. About 250 weevils were placed in stands and in cages which were as near natural conditions as possible. Only one weevil was found in the spring in all the places where they were "planted" in the stands. This one was in the duff beneath a tree which showed unmistakable signs of fall feeding. Only one living weevil was found in the cages, and this one eventually died from starvation after emergence from hibernation, as no fresh food was available. In one pure plantation 25 pairs of weevils were placed in a cage, which was sunk in the ground well below the surface of the mineral soil. All of these died.

A large number of trees were examined during the springs of 1927 and 1928, and a search made in the litter under all trees which showed injury from fall feeding. In most cases only one weevil was found under a tree, but in some cases two or more. Seven were found under one tree, and all were alive and became very active on exposure to the warm air. A few of the weevils were found in the upper layers of the litter, but the majority were at the surface of the mineral soil. Those which were above that were very probably making their way to the surface of the litter preparatory to ascending the trees. In the cases of the weevils which were "planted" in the stands it must be remembered that it was possible for them to fly from the area. This, however, was not the case in the cages, and, as stated previously, the mortality was very high.

The weather conditions preceding and during the hibernation period have a distinct bearing on the success of the hibernation. An early winter with an extended period during which the ground is covered with a blanket of snow is favorable, while an open winter, especially one which was preceded by an open fall such as was the case in New England in 1927, causes the death of many of the weevils while in hibernation. The reduction in weeviling, as a whole, over the entire New England area in 1928 is evidence of this. The condition which will do most to reduce the numbers during hibernation is a combination of sudden changes in temperature and an abnormal rainfall.

Hopkins (1907) has stated that, in common with some of the other species of *Pissodes* and allied forms, "It appears that an individual adult may live two or three years and will deposit eggs each year." In 1927 an attempt was made to determine whether this was the case. A cage about ten feet high and six feet square was built around an open growing tree which was about six feet high. Six pairs of weevils were liberated in the cage on June 4. The last date on which feeding was noticed was July 4. At that time the leader was dead, having been weeviled. It was removed so that none of the new generation would develop and be present in the cage. No evidence of fall feeding was found and no weevils had resumed activity the following spring nor up to the time the cage was removed the next July. The top of the cage was removed so that snow could fall around the bottom of the tree and approximate natural winter conditions. The fact that no weevils were found does not necessarily prove that they will not live over a second winter because the mortality is so great that even a hundred weevils might have been placed in the cage and all might have died.

Resumption of Activity

Resumption of activity by adults begins in the spring about the time the buds on the white pines begin to swell, and apparently

depends entirely on the weather. Egg laying does not usually begin until several days have elapsed. This gives rise to the belief that the females must feed for a time before they are physiologically able to oviposit. A very early spring and consequently an early beginning of growth will result in an early beginning of the oviposition period. Examples of the dependence on the weather were observed in 1926 and 1927. In the spring of 1926 the weather conditions were adverse and general weevil activity, as usual, did not begin until the buds began to swell. This was about two weeks later than in the previous year. At the Rainbow Forest Plantations, 12 miles north of Hartford, Connecticut, weevils were found for the first time that season on April 23. No pairs were previously seen and there had been no flight. Seventy miles to the north, at the Harvard Forest, Petersham, Massachusetts, the first weevils were observed on May 3, and the first eggs several days later.

In the same region in 1927 there was a warm period in late March and early April and bud swelling began about April 13. Adult weevils were found on April 20 and eggs the following day, so it is probable that some adults were out earlier than the date noted. About this time a cold period set in, with the temperature to a minimum of 16° F. This period lasted for several days, and the weevils descended to the litter beneath the trees and did not reappear until another warm period began about May 1. This drop in temperature caused a temporary cessation in the height growth of the trees and the new beginning of height growth and the activities of the weevils coincided, both being resumed on May 2.

In Canada, in the range of the Northern white pine, the springs are usually cold and wet. This would preclude early emergence from hibernation and would result in later egg-laying. The contrary would be the case in the southern portion of the range of the insect. In the Carolinas, although the pines are in the higher altitudes, there is an early spring, and consequent early emergence of the weevil from hibernation. Thus the probabilities are that in Canada the egg-laying would not begin until the first of May and in the Carolinas not until about the end of March or the first week in April. In New York, around Syracuse, the beetles emerge from hibernation about the middle of April in an average season.

During the feeding and oviposition period, the gregarious habit of the adult weevils is especially noticeable. Usually there are several weevils on a leader which has been attacked. There is a distinct odor from the pitch which is escaping from the punctures, and this doubtless acts as an attraction to other weevils, thus explaining the presence of several on each leader attacked. The males at this time are very attentive to the females and at times

there will be three or four of them to each female found, although at other times the females may be alone.

Direct observations in the field show that the weevils are likely to be more numerous on leaders already weeviled than on those just being attacked, because the former give forth more odor. In one area where white pine slash was being caged under closely meshed wire screening, specimens of *Pissodes strobi* and *Pissodes approximatus* were found in the cage before there was time enough to close it. These weevils came from a pile of slash close by and undoubtedly were attracted by the odor.

If a weevil is intent on making a puncture, or if copulation is in progress, little difficulty is experienced in making observations. Care need only be taken not to breathe in the direction of the leader or to stand between the tree and the wind movement or the source of light. Breathing on the weevils causes them to release their grasp of the stem and to drop to the ground or lodge in the needle clusters. If the weevil is simply clinging to the stem, the moment it detects a foreign object it will "play possum."

In unseasonable weather refuge is taken in the bud clusters and from such positions it is usually extremely difficult to remove the weevils and a considerable amount of shaking is usually necessary before they can be dislodged.

Egg Laying

After resumption of activity, the first few days are spent in feeding and copulation. During this time punctures are to be found in the upper third of the leader and also in the apical buds. Then the females begin to lay eggs. The following observations made by Peck (1817) are rather inaccurate but are included here merely for the historical reference. He states, "The insect is so small that it is impossible to detect it in the act of depositing its eggs, which are probably placed under the thin skin or epidermis of the shoot. I suspect that the larvae remain in the wood more than one year, and that the shoot dies the second year after the eggs are placed in it." Fitch (1853) states that the weevil deposits her eggs in the bark of the topmost shoot and drops one egg in each puncture at irregular intervals through the whole length. The writer has found that, as a rule, it is only the upper third of the leader in which the eggs are laid, and there may be two or more in each cavity.

The manner of depositing the eggs, which has been reported previously by Blackman (1919) and Graham (1926), is as follows: The excavation is carefully made in the bark not quite as far as the wood; then the female reverses herself and feels about with her ovipositor until she finds the excavation. In this excavation she deposits one, two and occasionally three eggs. Reversing

herself again she pushes them farther into the cavity with her beak. The whole operation may take nearly a half hour, depending apparently on the thickness of the bark and probably on the temperature of the insect itself. The writer examined a large number of leaders which had just been weeviled and found that in almost every case the eggs were deposited in punctures which had not encountered a resin duct. The resin exudes from the punctures where the ducts were encountered and flows over the bark, eventually filling in any egg punctures which have been made and covering the eggs contained therein. Exposure to the air for a short time causes it to harden and it acts as a sealing agent.

Peirson (1922) is authority for the statement that a single weevil will lay as many as one hundred eggs. Graham (1929) records an average of one hundred and fifteen eggs per leader, but he reported earlier (1926) that only one female had been observed in the caged tree from which he obtained this figure.

During the summer of 1928 the writer caged a number of paired weevils on leaders of white pine, simulating natural conditions as closely as possible. These pairs were removed from trees a day or two after oviposition was apparent, and in each case copulation was actually in progress. There is a possibility that each female had laid a few eggs previous to the time she was taken, but examination of the leaders, from which each was removed, showed comparatively few eggs. The cages were removed and placed on new leaders four times, and in two cases five times, so each female actually was placed on five or six leaders. These cages were of galvanized wire cloth, sixteen mesh, and were cylindrical in form (Plate I). The upper end was closed tight with metal elips, and the lower end when placed over the leader was closed in a similar way. This made a tight cage, although strong wind movement eventually loosened several of them to such an extent that the weevils were able to squeeze themselves out around the leader and escape. These cages were fastened to a stake driven in the ground alongside of the tree, and thus made rigid enough to withstand ordinary wind movement.

Each time a cage was transferred the old leader was removed and taken to the laboratory for careful examination. The number of punctures was counted and then the bark was shaved off in small strips and the eggs and larvae counted. This caging experiment began May 10, when the first larvae were caged, and ran until August 13. The fourth collection of leaders, on July 10, was the last in which eggs or larvae were found. By this time some of the weevils had escaped, and on July 30 only four of the cages had weevils in them. Six cages had pairs in them for nearly a month, so each female may have been fertilized more than once. The leaders, at this time, had several punctures

in them but no eggs. Only one living weevil was found on August 13, and two leaders had punctures in them, so the weevils had died between July 30 and that date.

The greatest number of eggs laid in the cages by any one female was 82, with an average of 41 for six series of cages, which were carried through the oviposition period. The number of eggs laid by these six individuals was 49, 31, 82, 27, 10, and 45. It seems probable that an average of 50 eggs per female is low, as it is quite probable that some of the females had oviposited previous to the time they were removed from the trees.

There is the additional fact that in the majority of cases each female oviposits in more than one leader, and also that several females may oviposit in one leader. The writer has found from 20 to 350 eggs in one leader, and often several females are found on a single terminal shoot. In a heavily infested pure white pine plantation several hundred weeviled leaders were examined during the course of an experimental control project by removal of the infested leaders. The average number of eggs and larvae removed from these leaders was about one hundred and twenty-five. Even though many of the egg punctures contained two and sometimes three eggs, the average number of punctures totalled one hundred and seventy, showing that a large number were for feeding purposes or were flooded by an exudation of resin before the eggs were deposited. In a less severely infested stand or in certain mixtures where the infestation is generally low these averages would be considerably less.

If the beginning of oviposition is late because of unseasonable weather the cessation will usually be correspondingly late, as there is evidence to show that this period can be correlated rather closely with the period of accelerated height growth. Generally speaking, however, the peak of the oviposition is reached within two weeks after accelerated height growth starts, then begins to decrease gradually and after another month has passed has practically ceased.

Incubation Period and Larval Activity

The eggs hatch in from six to twenty days depending on the weather conditions. Immediately after hatching, the larvae, which are voracious feeders, begin to feed down the stem on the phloem tissues, thereby depriving the shoot, wholly or in part, of the food necessary for elongation. As they grow larger, they feed also on the outer portion of the wood. If there are a large number of larvae the living tissue will be quickly eaten and the growth may be completely stopped.

The larvae follow down the stem side by side, often packed in closely in a ring, and many of those at the rear are undoubtedly starved to death. The final survival of larvae represents only

a fraction of the number of eggs laid, and the eventual development to adults depends on a number of factors. These are the number of eggs laid, the vigor of the leader, the time of oviposition, and the presence of parasitic and predacious insects and other predators—such as insectivorous birds. As a larva which is a little further developed than its companions attains full size it drops behind, bores through the wood into the pith and pupates. The remaining larvae work down the stem and if they are not eaten by birds, a fate which befalls many, they pupate in the pith or in the wood. If there is not sufficient room for all to pupate in the pith, and this is usually the case, the remainder will pupate in the wood but as near the pith as possible. The head is usually upward during pupation, just the opposite to that usually maintained during the larval period, but quite often, as may be seen in Plate II-1, the head may be downward.

There is a striking difference in the character of the pupation chambers in the pith and those in the wood. Those in the pith, as Graham (1926) points out, have no "chip cocoon" or lining of frass as has usually been reported, and those in the wood have it only on the side toward the bark. The main portion of the chamber is smooth and with no lining whatsoever, only the marks of the mandibles being visible. (Plate II—2 and 3)

Occasionally some of the larvae when they come to a node will bore into a lateral, feeding and pupating there. Several cases were also observed during the season of 1926 where larvae from eggs laid late in the season had gone upward into the 1926 growth and pupated there. This occurs, however, only when much of the axial growth of the tree has been completed at the time the larvae hatch, and when there would probably be sufficient food available to induce the larvae to go up instead of down. In rare cases leaders of the current season are weeviled, but this only occurs when the eggs are laid very late in the season and the shoots are nearly full grown.

During the progress down the shoot it quite often happens that one larva may get ahead of the others, and almost invariably in this case it will be smothered by the resin flow. It has also been found that weevil mortality in infested leaders in mixed and in dense pure stands is exceedingly high, often being one hundred per cent. In such cases the leader very often is not killed and, if the growing point of the leader is still above that of the laterals, the stem will remain straight at that point, although there may be a slight decrease in the growth of that year due to the attack. All of this shows that the concerted feeding of a considerable number of larvae is necessary to overcome the resistance of the leader.

The length of the larval period varies according to the weather conditions, but, on the average, approximately four to six weeks elapses from the time the eggs hatch until pupation commences.

In 1928, at Petersham, Mass., the first larvae were found on May 28. They were very small, and apparently were in the first stage. The first pupae were observed on July 12 and they apparently had transformed on or about that date, as larvae in the prepupal stage were very numerous. Apparently it is not easily possible to determine definitely the number of larval stages, although it is quite certain that there are either five or six. The instars representing the different stages appear to be similar, and the number of stages can not be counted from the different sizes of the instars as the length and width of the male and female larvae of the same stage may vary.

Pupal Stage

The pupal stage lasts about two weeks and the young adults remain in the pupal chambers two weeks or longer before they finally bore out from the leader and emerge. Nearly three months is the minimum time for development from the first eggs to the first adults in the northeastern states. In warmer regions this time is considerably less; and larvae from eggs laid in warm weather in the northeast will also have a shorter developmental period. In 1928, also a late season, the total period for the earliest adults noted was 100 days. It must, however, be remembered, that in both instances these were the first adults.

Blackman (1916), Pierson (1922) and Graham (1926) all refer rather definitely to the beginning and cessation of the oviposition period. Weather conditions being considered, it is shown in Graph I that these dates, and those of the larval and pupal periods, can be correlated with definite periods of the height growth. In 1928 the accelerated height growth at Petersham began in the week ending May 24, and the first larvae were found a few days after this. This accelerated height growth began to decrease in the week previous to the time that the first pupae were observed, and was very definitely retarded at that time.

Habits of the New Generation

After a short time spent in the pupal chamber, the first young adults emerge from the leader and very often can be found feeding on the living twigs of the tree from which they apparently emerged. Practically all the members of the new generation emerge in the latter part of the summer or the early fall, but there is such an overlapping of the life stages that, as stated previously, the larvae from eggs laid very late in the season may winter over as full grown larvae or immature adults. Blackman (1916) has reported that the larvae may pass the winter in the leaders. The writer on two occasions has obtained weevils from cages containing infested leaders in April, and none were found when the cages were examined in January and February. The weevils were dead

at the time they were found, and it is assumed that they emerged and died from starvation as there was no green material in the cages. In one case the cages were placed in a barn where no heat was available and, with the exception of the sunlight and precipitation factors, the conditions approximated those which would have been found in a plantation or stand. In the other case, however, the leaders were collected in December and placed in a warm room, resulting in early emergence during the winter.

Feeding

Fall feeding has been observed in cages and on trees in young plantations and natural stands. The young adults feed on the buds and the tender bark of the leaders, and also on any portion of the branches which has tender bark and is exposed to the sun. On warm days almost every tree which has a weeviled leader will, if examined, be found to have its population of weevils busily engaged in feeding. In some cases the injury, when a number of weevils have been feeding on the same twig, presents the appearance of that caused by the Pales weevil, in that much of the bark appears to be gnawed away. On close inspection, however, many small punctures, characteristic of the White Pine Weevil, will be found. The twigs are not often killed by fall feeding, but will be calloused when the injuries heal. Sometimes the cause of an aborted leader can almost certainly be traced to fall feeding by the White Pine Weevil during the preceding season.

Feeding may continue at intervals, even though snow may have fallen in the intervening time until settled cold weather prevails and then the weevils go into hibernation under the trees upon which they have been feeding. In 1925 feeding was observed in central Massachusetts as late as the middle of November even though a rather severe snow storm had occurred in the same area on October 10 and the snow had remained for some days.

Copulation

It has long been known that the females are fertilized in the spring during the short feeding and copulation period immediately after the resumption of activity. But it is possible that copulation may also occur in the fall. To determine this possibility the writer made extensive observations on a large number of the new generation of adults in cages. A quantity of litter and dead needles was placed in the bottoms of the cages so that the weevils could take shelter at any time or could feed on the fresh twigs which were also in the cages. So long as a temperature above 70° F. was maintained (usually 75° and 80°) and the weevils were hungry, they were quite active and copulation was frequently observed. About fifty pairs were observed in copulation and several of these more than once. As they were observed for

the first time they were removed to another cage, where the same conditions for shelter and food prevailed, in the hope that the females would remain alive over winter and could be observed from time to time. The conditions in the cages, probably owing to the drying out of the litter, were evidently unfavorable, for all the weevils eventually died.

Little difficulty was met with in collecting the coupled pairs if they were actually copulating. Several times, however, in attempting to remove pairs just after the male had mounted the female, he immediately feigned death and dropped to the bottom of the cage. During the course of the writer's observations many punctured twigs were removed, the bark stripped and each cavity carefully examined. Although there was abundant proof that copulation had actually taken place, and the females placed in the cages with the twigs, no eggs were found in any of the punctures.

The time involved in a single copulation does not appear to have any definite limits and may be several hours. If the temperature is favorable it can take place at any time during the day or night. Several pairs were observed coupled for over two hours. A few pairs were observed as late as twelve o'clock midnight, and had been in the dark for some time, but it is not known whether or not copulation actually began in the dark. The application of light had no apparent effect on these pairs but in the same cage other individuals were stimulated and became active in a short time.

Graham (1926) mentions a case where a female emerged from hibernation in a cage which had been placed over a tree the previous fall. No other weevil was observed in the cage during the whole period of oviposition. She laid 115 eggs most of which were fertile and hatched. This brought to Graham's mind the possibility of fertilization the fall previously or reproduction parthenogenetically. The next fall he placed a large number of weevils in a cage and watched them. He observed but one pair in copulation, and that only once. He explains, however, that there was no cover in the cage except green twigs for food and this may have made the conditions unfavorable for any unusual activity. Hopkins (1907) found that the ovaries in the newly emerged females were undeveloped and they could not possibly lay eggs in less than a month.

On May 1, 1928, one female was found in the lower layer of the litter next the mineral soil beneath a pine tree. A few days later three more were found under the same conditions. All four were placed in separate cages which were in an open field. These cages were sunk to the mineral soil and had the sides banked with earth. The litter, placed at the bottoms of the cages, was screened and examined so as to be certain no weevils were present. Eggs

were found in the leaders in all cages, and larvae were found later in the season in three of them. The facts that the spring of 1928 was very backward and that conditions in the field were not favorable for an early resumption of weevil activity make it appear that fertilization can take place in the fall and will last over until the following spring.

The caged specimens observed during the fall of 1926 were most active when hungry and, if heat was applied at this time, many would attempt to fly about the cage. Striking the ends of the cage or the glass front with apparent force did not deter them from trying again and again. When they had appeased their hunger or their sexual desires or when the cage was allowed to cool the great majority of them would take cover in the litter.

Flight Habits

If there is a favorable air temperature these young adults will fly readily. Migration to new areas by flight is possible if the air temperature rises to approximately 75° F. This often occurs in the fall especially in the middle of the day when the sun is bright. The weevils are relatively strong fliers and, when carried high in the air by the wind currents, may cover a considerable distance, a flight of several miles being possible.

The manner in which the weevils reach the leaders has for some time been a matter of doubt, and experiments were made in the spring of 1926 to determine how this was accomplished. It was found that the weevils may reach the leaders by one of three ways. In the early spring, before the temperature is high enough to enable them to fly, they have been seen crawling up the stem. As soon as the temperature rises sufficiently to permit flight they will fly directly to the leaders. Several were caught and it was found that both males and females will fly in the spring as well as in the fall. From observation and experiments conducted after the new generation had emerged in 1926 the writer found that ordinarily an air temperature of at least 70° F. is necessary before the weevils will resort to flight. The temperature range most conducive to flight is between 70° and 85° with the optimum from 75° to 80°. Above 85° the weevils will seek shade, especially if the temperature remains above that limit for any considerable time.

The way the leaders are reached was determined by banding them and the bases of the trees with sticky tree banding material. In the spring of 1926 two series of sample plots were laid out before the snow and ice were off the ground and some time before the weevils had resumed activity. The banding material was inspected at least once each week thereafter, until late in the summer, in order to be sure that it was remaining sticky, and any applications which showed signs of hardening or glazing over from rainy

weather were combed or renewed. Each series consisted of a central plot, with a check plot on each of the four sides. Each plot was one-tenth acre in area.

One series of plots was in a severely infested plantation. The leaders of all the trees in this central plot were banded just above the lateral growth of the previous year, and the border trees were also banded at the bottom. The branches below the basal bands of these border trees and any above which had their tips resting on the ground, were removed. This was to prevent any weevils from getting to the trees in the plot except by flying directly to the leader. The sample plot records show that there was a substantial decrease in percentage of weeviling in the central plot over the average of the four check plots. The percentage of trees weeviled in this plot decreased from 51 in 1925 to 12 in 1926, a decrease of 39 per cent. In the check plots there was an average infestation in 1925 of 37 per cent and in 1926 of 57 per cent, an increase of 20 per cent.

The second series of plots was in a plantation less seriously infested than that in which the first series was laid out. All the trees in this central plot were banded at the base. The border trees had an additional band on the leaders, just below the lateral growth of the previous year, to prevent weevils from getting to the leaders from neighboring trees in the check plots by crawling along the interlocked branches and on up the stems. As in the first treated plot, and for the same reason, all the trees banded at the base had the branches below the bands and those above which had their tips resting on the ground removed. As in the former case there was a decrease in the percentage of trees weeviled. There were 27 per cent weeviled in 1925, and 15 per cent in 1926. In the check plots around this central plot there was an average infestation of 21 per cent in 1925 and 35 per cent in 1926, or an increase of 14 per cent.

Infestation in the check plots thus materially increased in both series at the same time that it decreased materially in the treated plots. Both treated plots showed a substantial decrease in percentage of weeviling over that of their respective series of check plots, and also over the previous year's infestation. This was particularly striking in the plot in Series 1, where all the trees were banded at the leader.

The fact that trees banded at the bottom were weeviled shows that the weevils do not necessarily have to crawl up the trees. The data from the other plot, with all trees banded at the top, show that a certain percentage of the weevils will fly directly to the leaders, but the fact remains, because of the much higher infestation in the check plots, that many trees must have been reached in some other way. This must therefore have been by flying to

some other part of the tree than the leader, and crawling up the stem until the leader was reached.

From this information it can definitely be stated that the weevils reach the leader in any one of the three ways enumerated above, but probably more often by flying to the tree, striking the lower branches and then crawling the rest of the way. In both treated plots a number of weevils were caught in the bands at the base, and only one in a band at the leader. In the central plot of Series 1, with only the border trees banded at the bottom, five weevils were found on four bands. In the other central plot thirteen weevils were found on nine bands, some of which were on trees in the center of the plot.

HOST TREES

Quite a number of species of trees have been found to serve as breeding hosts for the White Pine Weevil. The white pine, *Pinus strobus* L., is the favorite host and the injury is found in varying degrees in almost every locality where this species is common. The Norway spruce, *Picea abies* (L.) Karst., is also severely attacked and at times may be injured to as great an extent as the white pine. All of the native pines and spruces of the northern region are liable to attack, although the white spruce, *Picea glauca* (Moench) Voss, and the red pine, *Pinus resinosa* Ait., are rarely affected. In addition a number of exotic coniferous species have been reported as hosts. The species of western soft pines listed would probably be weeviled to a greater extent if the numerical abundance were great. This would also probably apply to the sugar pine, *Pinus lambertiana* Dougl., and the other soft pines if planted in this region.

Injury due to the White Pine Weevil has never been observed by the writer in Canada hemlock, *Tsuga canadensis* (L.) Carr., or balsam fir, *Abies balsamea* (L.) Mill., but such is reported by Packard (1890). He states: "The grub and pupa occurred . . . at the end of July in the fir; on the 30 of July I took five mature beetles from under the bark of a hemlock tree." From this description of the location of the beetles in the hemlock it appears that Packard confused *Pissodes strobi* with some other weevil, and probably the same applies to his statement regarding the fir. It is quite probable that the species reported by Hopkins (1904) in Deodar, *Cedrus deodara* (Roxb.) Loud., is *Pissodes deodarae* Hopk., is it infests the same portion of the stem in Deodar cedar as the White Pine Weevil infests in the species it attacks. The White Pine Weevil has also been reported in southeastern Texas on shortleaf pine, *Pinus echinata* Mill., and in northeastern Texas on loblolly pine, *Pinus taeda* L. Apparently this is a mistake, as *Pissodes nemorensis* is the only species likely

to be found in this region, and this species works in stumps and bases of felled or dying trees.

It must be borne in mind, however, that in localities within the range of the weevil where the insect is not abundant many of the species listed would not be attacked except in rare cases. At times where there is a mixed stand of conifers the other species in the stand may be weeviled to a greater extent than the white pines. This has been observed in New England and New York in areas where Norway spruce and white pine are either in the same plantation or are near each other. This brings up the Host Selection Theory and the possibility of a difference in the food materials causing the production of a strain of *Pissodes strobi* which prefers some other species than the white pine. This would explain why the other species were attacked to a greater degree.

Mr. Ralph C. Hall, School of Forestry, Michigan, has also reported greater infestation of jack pine in Northern Michigan than of white pine. In this region the white pine had been removed and the jack pine, which formerly grew mostly where the other species would not grow, was able to seed in these logged areas more quickly than the white pine. There were so few white pines available to attack that the jack pine was infested.

A list of the trees known to be attacked by the weevil in the general range of the insect, in the order of their susceptibility, follows:

Severely: White Pine, *Pinus strobus* L.

*Norway Spruce, *Picea abies* (L.) Karst.

Commonly: Pitch Pine, *Pinus rigida* Mill.

Jack Pine, *Pinus banksiana* Lamb.

*Japanese Red Pine, *Pinus densiflora* Sieb. & Zucc.

*Western White Pine, *Pinus monticola* Dougl.

*Limber Pine, *Pinus flexilis* James.

*Foxtail Pine, *Pinus balfouriana* Murray.

Red Spruce, *Picea rubra* Link.

Occasionally: *Scotch Pine, *Pinus sylvestris* L.

*Western Yellow Pine, *Pinus ponderosa* Dougl.

*Mugho Pine, *Pinus montana mughus* (Scop.) Willk.

Black Spruce, *Picea mariana* (Mill.) B. S. P.

Rarely: *Blue Spruce, *Picea pungens* Engelm.

White Spruce, *Picea glauca* (Moench) Voss.

*Douglas Fir, *Pseudotsuga taxifolia* (La Marek) Britton.

Red Pine, *Pinus resinosa* Ait.

*Himalayan Pine, *Pinus excelsa* Wall.

* Species not native to the region in which *Pissodes strobi* occurs.

INDICATIONS OF INJURY AND CHARACTER OF THE DAMAGE

The first indication of attack by weevils is the presence of tiny glistening droplets of pitch on the upper third of the growth of the previous year. Closer examination will show tiny holes under these droplets, and also in the apical buds. These are due to the boring activity of both sexes in search of food and to the work of the females in preparing cavities in the stem for the deposition of their eggs. The punctures in the buds may indicate a serious injury, as the inner portion of the bud is often almost entirely consumed. In a leader heavily infested the droplets of pitch are very numerous, as many of the cavities tap resin canals, although eggs are seldom placed in cavities causing any considerable flow of pitch. By merging together, these droplets may completely cover the bark of that portion of the stem where the punctures are. The whitish color which this pitch assumes after it hardens is very noticeable against the darker background of the needles and bark, as shown in Plate II-4. It is the first notable indication that the shoot has been weeviled. This covering may to some extent act as a protection to the eggs and young larvae against predators.

Height growth starts normally in the infested leaders; but soon after the eggs hatch and the larvae begin to feed, the leader is girdled and assumes a wilted appearance characteristic of this injury. There is considerable variation in the growth of the new leaders, depending on the time of attack and the number of eggs deposited. If the stem is girdled almost immediately, as is the case when a large number of eggs are laid just as the buds are swelling, there will be little or no elongation. If only a few eggs are laid in the shoot, or if it is not attacked until later in the oviposition period, the elongation may vary from a very small amount to nearly full length. Weeviled leaders are very conspicuous in a plantation or natural stand and especially in young open stands in old fields where the infestation is severe. As a rule the new growth withers, and the tip droops and turns brown (Plate III). If the shoot is killed just after bud swelling begins there will be only a brown stub sticking up straight, altho the death of the growing tip due to feeding on the buds may also be the cause of this.

A leader killed by weeviling will be replaced by a lateral or laterals. The inherent tendency of the white pine to axial growth makes this possible. After the weeviled leader has died, at least one lateral, and usually several of the whorl immediately below the killed portion, begins to show an upward tendency. In course of time a new leader, or several, will become established and in many cases there is a fork or noticeable crook. The deformed trees that thus result from this injury are known as "bayonet"

(Plate IV) or "staghorn" (Plate V-2) pines. The first occurs when one lateral finally gains supremacy over the others and becomes the leading shoot. The second occurs when two or more laterals survive and form a forked tree. It is not uncommon to find several of each of these in a single tree. Such trees may have 20, 40, 60 or 70 leaders. These are called "cabbage" pines, and the shape instead of being conical becomes globular (Plate VI-2). In such cases the trunks are practically worthless for lumber. Peck (1817) points out that, "The lofty stature of this tree depends upon the constant health of the leading shoot for a long succession of years. If the leading shoot is destroyed, the trunk can rise no higher, although it may happen that in the circle of branches immediately below it, one more vigorous than the rest may be forced by those on each side of it into a more ascending direction and become a kind of leading shoot—yet the tree is deformed." One, and occasionally two laterals may eventually gain dominance and the others left behind will die. When these stubs—often some inches in diameter—break, a jagged wound is left which can not heal and which permits the entrance of wood rots, and in many of the trees so injured the heart is defective.

Effect on Axial and Radial Growth

The injury to the trees is more apparent when the attack occurs on the young than on the old trees. It is very difficult to estimate this damage exactly, but it is an unquestionable fact that there is loss in both height and radial growth. The loss in height growth is caused by the killing back of at least two years' growth each time the tree is weeviled. Usually at least one lateral will gain dominance but the resulting leader will be shorter at any given time than the natural leader would have been, and there will also be a crook in the trunk. However, once the new leading shoot has definitely gained supremacy it will have an annual elongation similar to the non-weeviled trees, providing it is not weeviled again.

Loss in Axial Growth

Graham (1926) records the loss in height growth for each weeviling as "the difference between the length of the main stem above the topmost living whorl and the length of the lateral that assumes the lead." Graham, however, does not include the difference in amount of growth that actually occurred in the lateral that assumed the lead and that which would have occurred in the real leading shoot the season of weeviling had it not been weeviled. Nor does he take into account the loss for several seasons following until the new leader is firmly established as such, as the death of two or more years' growth is undeniably a great physiological shock to the tree. This loss, over a period of years, can be approximately computed.

In Graph II is shown the approximate loss which can be expected in trees, killed back two years, up to several years after weeviling. This approximation was obtained by measuring the growth of the year before weeviling, of the year of weeviling and of the four years after weeviling, and comparing these figures with the growths of the corresponding years of non-weeviled trees growing under similar conditions. The stimulation of the new leader over the corresponding lateral is also shown. Sixty trees of each class were measured.

Quite often three years' growth and occasionally four or five years' may be killed. When this occurs there is even greater loss. The number of years' growth killed back depends on two conditions—(1) the vigor of the tree and specifically the height growth of the preceding year, and (2) the relation of the diameter and length of the leader to the number of eggs laid in the stem. The continued loss in the height growth for some years after will also depend on these conditions. The fact that, as a rule, it is only the trees with the most vigorous leaders which are attacked, even though the trees themselves are not actually dominant, makes this at once a very serious injury.

Occasionally, as Walker (1911) has also reported, a very vigorous leader will recover, but this seems to be only when a small number of eggs have been laid, more particularly late in the season, and the larvae are smothered by the resin. In many cases recovery actually occurs in attacked leaders of pines in mixed stands, but often when this happens the leader is permanently retarded, as a lateral has attained supremacy while the injured leader was recovering from the attack. If a lateral, in the course of its struggle for leadership, once gets its tip above that of the real leader, it immediately becomes the leader itself and remains so even though the original leader is not killed.

Effect on Radial Growth

Aside from the loss in axial growth there is also an appreciable decrease in the diameter growth in certain parts of the tree attacked and an increase in other parts of the tree. In the portions of the tree where for several years there is a continued decrease, the loss is greater the year after attack than the year of attack. This may be caused by three factors. The effect of the shock due to the injury is noticeable in the diameter growth as well as in the axial growth. The energy required to counteract, or attempt to counteract, the injury is so great that the tree can not carry on its normal functions of making diameter growth and laying up stored food for use in the early part of the next season. There is less food manufactured because of the death of the leader and its growth of laterals and there is more food used because the stimulus for height growth of the lateral or laterals, striving for leadership,

uses up materials that normally are converted into stored food and utilized the following season. The total amount of these losses will be governed by the amount of new growth that was actually made before the previous season's growth was girdled and the new growth wilted. A leader that is not girdled until late in the season may have transported some materials down the stem to be stored, before the effect of weeviling was felt. Also the growth of laterals, immediately below the killed portion, which are striving for the leadership, will not be so far advanced in the additional growth which is made than would be the case with those which had an earlier stimulation.

If one tree alone is considered this loss in growth will be negligible, but if all affected trees, over a wide area, be taken into account it can readily be seen that the ultimate loss of the stored materials, and subsequently the loss of diameter increment, will be considerable. This continued decrease in radial growth for several seasons will be found at the base, at breast height and at the internode below the killed portion. In the newly developing leader there is a decrease in radial growth the year of attack corresponding to that in the lower parts of the tree, but the year after attack there is an increase, followed by a noticeably larger one the next year. In the second internode of this new leader, which grows the year of weeviling, this increased diameter growth does not become apparent until the second year after weeviling, and is accentuated the third year, so the increases here correspond to those in the lowermost internode of the new leader.

The reason that there is less stored food in the parts of the tree below the weeviling is due in part to the cutting off of the conducting system between the leader and the main stem when the weeviled leader dies. If more than two years' growth is killed the loss will be even greater. Theoretically, there will be a loss each year until a new conducting system is completed and the new leader is firmly established. If, however, the tree becomes overtopped this loss will persist even longer than that. The temporary loss in annual diameter growth in the new leader is caused partly by the shock and partly by the diversion of the food materials in order that height growth may be increased. The increase in this new growth after the first season following weeviling is probably due to the fact that more energy is poured into the new leader, due to the inherent tendency for axial growth which is found in white pine. This additional energy manufactures a supply of food greater than is ordinarily produced. There is a very limited downward conducting system available at this time, so storage of food in the stem of the tree is severely curtailed, and additional axial growth results.

TABLE I.—To show the loss in diameter growth, in millimeters, in different parts of weeviled trees. Explanation in text.

No. of trees	PORTION OF TREE	Condition of tree	NUMBER OF YEARS AFTER WEEVILING					Year of wee- viling	NUMBER OF YEARS BEFORE WEEVILING				
			5	4	3	2	1		1	2	3	4	5
20	Annual Diameter Growth at Base	non- weeviled	5.14	5.02	4.88	5.36	5.58	5.84	6.74	7.70	7.90	7.60	7.28
48		weeviled	2.56	3.16	3.66	4.62	4.70	5.32	7.22	7.84	8.06	8.26	7.94
20	Annual Diameter Growth at D. B. H.	non- weeviled	5.52	5.64	5.98	6.68	7.16	7.82	8.24	7.68	7.40	6.82	5.74
48		weeviled	4.96	4.32	4.58	5.64	6.42	7.24	9.46	9.48	7.40	7.00	3.62
20	Annual Diameter Growth of Internode below Weeviling	non- weeviled	5.52	5.76	7.18	8.24	7.86	7.78	7.00	2.90			
48		weeviled	5.04	5.10	6.24	7.10	6.64	6.94	7.60	3.26			
20	Annual Diameter Growth First Internode of New Leader	non- weeviled		7.82	7.98	7.40	7.46	6.28	2.66				
48		weeviled		7.02	7.02	6.98	5.78	4.50	2.58				
20	Annual Diameter Growth Second Internode of New Leader	non- weeviled	7.38	7.76	7.28	7.32	6.18	2.62					
48		weeviled	7.04	7.56	6.74	6.40	4.98	2.52					

Table I shows these decreases or increases in the diameter increment in millimeters. The material for this table was obtained from trees removed from plantations at the Rockwell Springs Experiment Station, Syracuse, N. Y., and is based on 48 weeviled trees and 20 non-weeviled trees. Sections were removed from the base, breast high, one internode below weeviling, the internode of the new leader the year of weeviling and the internode of the new leader the year after weeviling. The corresponding internodes were taken in the non-weeviled trees. In all cases this average diameter growth was obtained by taking the average radial increment and doubling it, and the measurements of the three upper portions were taken halfway between the nodes.

Effect on Quality of Timber

The first noticeable result of the killing back of the leader or leaders is the crook or fork (Plate V-I) which must invariably occur. This decreases the value of the trees materially, especially if they are in a pure stand which has not reasonable density or vigor. Usually the lumber sawn from such trees is usable only for boxboards or the cheaper grades.

The quality of the lumber in trees weeviled once or even several times may not be seriously impaired, provided the trees have nearly straightened at the point of attack (Plate V-3). There

are, however, many knots in a board from a weeviled tree. Although the tree may have straightened entirely there will always be a larger knot at the node from which the new leader came than at an uninjured node. If the weeviled leader persists for many years, as is often the case, there will be a dead, and often a loose knot extending to the center of the tree (Plate VII-1, 2, 4), and this will show in all boards sawn from that side of the tree. There is also uneven grain and boards with crooked grain will be common. The boards will also show clear lumber on the side of the stem from which the successful leader arose, and the opposite side will have several knots of varying sizes (Plates VII-3, VIII-1.) If this condition would remain constant—that is, having the clear portions always on the same arc of the stem cylinder—that side would produce longer lengths of clear lumber than is usually the case. But, unfortunately, this is seldom true, as following a second weeviling, even immediately above the first, the new leader will usually arise from another arc of the stem cylinder, and a greater crook at the new place of attack often results.

In the older trees there is often a large dead snag, traceable to weeviling. The breaking off of this dead portion, usually through mechanical means, allows the entrance of fungal spores, thus resulting in decayed sections near the crook or crotch, causing defective lumber (Plate VIII, 2, 3). In specific reference to *Trametes pini* Fries (which is the basic cause of the heart rot in white pines in the northeastern states) Hartig (1894) states that if the spores gain a footing on a fresh branch wound which is not protected by a covering of resin they push their germ tubes into the stem and the mycelium spreads upward and downward. However, Hartig was referring to the wounds in dead branches of older trees and also states that younger trees enjoy immunity from infection because in their cases wounds on living portions are quickly protected by an exudation of resin. It is the older trees in this region which show the injury, and in many cases it is very probable that ingress to the heartwood was gained through wounds such as could be caused by the breaking of a dead stub.

However, there is also every possibility that spores can gain entrance through the dead leader, as no exudation of resin occurs in the dead weeviled leader if it is left on the tree. This leaves a wound which is not covered by resin, as it is only the living sapwood that carries it, and there would be a large enough space uncovered to allow the entrance of the spores. The white pine on the old field rocky soil which is so common in the Northeast is subject to heart rot, and it must be remembered that entrance may be had by other means than the dead stubs caused by weevil attack. However, it is also quite probable that much of the infection in the stem would not occur if dead stubs, which can be traced to weevil injury, were not so abundant. Pruning the shoots in the summer,

where it is a practicable undertaking, would be effective in reducing the possibility of this injury, as the exuding pitch would glaze over the cut surface and prevent entrance of the spores.

Injuries Often Mistaken for Weeviling

After four or five years it is often difficult to designate the year of injury merely by examining the weeviled leader, and after a lapse of several years it may be impossible to tell if a crook or fork was caused by weeviling or some other injury. There are at least four agencies each of which causes an injury that might be mistaken for weeviling.

Injury caused by bud freezing is perhaps the one which looks most like weeviling and an untrained observer often mistakes it for the latter. It is more prevalent in some strains of Scotch Pine planted in the Northeast than in any other of the species liable to weevil attack. Under certain conditions of temperature and moisture the terminal bud will be killed back and the leader dies to the lateral whorl of the previous year. A dead leader very often results, but absence of emergence holes should show at a glance that weevils are not guilty of the injury (Plate IX-1). Often a paint brush effect (Plate XIII-1) may be produced due to the growth of adventitious buds and needles. Belyea and MacAloney (1926) found that these dead leaders persisted for years and could very easily be mistaken for weeviling.

When overtopped by broad-leaved trees, or occasionally by coniferous trees, white pine and other species affected by the weevil are often injured by whipping (Plate IX-2). The leader is killed and a lateral from below forms the new leader. This may happen for several years in succession and in many cases the tree will be completely overtopped. If the whipped leader has been broken off it may be impossible to tell whether the resulting crook was caused by that or by weeviling.

A black aphid, *Dilachnus strobi* Fitch, also attacks the leader of the previous season. The bark is often severely injured and the leader sometimes dies. The absence of white pitch in the early summer, the presence of a black sootlike material on the bark, and the absence of emergence holes later in the summer will enable the observer to distinguish this injury from weeviling. If the leader is killed there will be a crook and in the course of a few years it may be difficult to tell one injury from the other when the leader has been removed.

In some localities the injury done by the red squirrel, and reported by Hosley (1928), is the most serious of all these (Plate IX-3). At times in small areas it may be of even more importance than the weevil. The squirrel may injure the tree by removing

either the central bud or the entire apical cluster, or it may remove a considerable part of the leader as well. In any case the buds are eaten by the squirrel. The result is an aborted bud or killed leader and eventually a crook or fork. As a rule no emergence holes are found, but the odor of the fresh resin oozing from the wound may attract the weevils and the shoots will be weeviled and development of the new generation completed.

SUMMARIZATION OF PLOT DATA

Manner in Which Study was Carried Out

Because of the probability indicated by previous studies that silvicultural measures will in the long run prove to be the cheapest and most effective means of minimizing the weevil injury, much stress was placed on this phase of the investigation. These observations and the information gained from studying silvicultural methods practiced in such forests as the Harvard Forest at Petersham, Massachusetts, have shown that stands of a certain character or growing under certain conditions, such as a groupwise mixture of white pine and hardwood, are relatively immune to weevil injury. Special attention was given to determining the prevalence of the weevil and the amount of damage done by it in stands of different compositions and densities and on different sites.

For this purpose thirty-seven permanent sample plots and two hundred and sixty-five semi-permanent or temporary sample plots were laid out and studied in various parts of the states of Massachusetts, Connecticut, Vermont, New Hampshire, New York, Pennsylvania, North Carolina, and the Canadian Provinces of New Brunswick and Nova Scotia. These plots vary in size from one-fortieth acre to one-half acre depending on the site and forest conditions, with most of them one-tenth acre or one-fifth acre. Attempts were made in all cases to lay out the plot in what appeared to be the average condition in that stand unless some particular condition was to be studied. The permanent sample plots will be examined and studied periodically for a considerable length of time to determine the ultimate effect of various mixtures and the seasonal variation and abundance of the weevil. These plots are located near Syracuse and Cranberry Lake, New York; Petersham, Winchendon and Amherst, Massachusetts; Rainbow, Connecticut; Keene, New Hampshire; and Fredericton, New Brunswick; and cover a variety of conditions.

Data on Pure Stands

Examination of one hundred and forty-four sample plots, laid out in pure plantations under thirty years of age and in natural stands of all ages, shows that, disregarding other factors and

within certain limits of spacing, the density of the stand has a distinct relation to the amount of injury. Also, if the percentage of trees weeviled be considered, the amount of weeviling decreases with an increase in density. Generally, in regions of severe infestation, the wider the spacing the more severe will be the injury, but no distinct reduction will be observed until a spacing of 4 x 4 is reached. The wider spacing allows more light to reach the lower branches and there is more likelihood of forking than there is in the narrower spacings. There may be as many trees weeviled per acre in a dense stand but as the number of trees increases so will the number of unweeviled trees, and those which are weeviled will have more chance to straighten out because of the keen competition for height growth which is going on all the time. The actual percentage of trees weeviled will, of course, be less as the density increases, but, as is the case with the injury, this is not particularly noticeable until a 4 x 4 spacing is reached. In dense plantations this reduction is striking. Here the trees are even-aged and of approximately the same height. In a natural stand the density may be high, but if the trees are uneven-aged, and therefore probably of uneven height, a condition may be found which, in so far as the relation of the tops is concerned, may approximate a widely spaced stand and the taller trees may be weeviled and severely injured.

Practically all the plots were taken in stands where the site quality was either II or III. A few plots in plantations were undoubtedly on site quality I. As a rule there is less recovery from weevil injury and more infestation in stands where the site quality is a little poorer than II than in better sites. Usually in such a site the density of stocking is below normal and the trees which are in the stand are vigorous. In a stand on the poorest site, also, the few trees which are vigorous may be severely injured, but as there are not so many of them as there are on a better site, the ultimate damage is not of so much financial importance. The plots in the older stands were all in natural growth and these showed very plainly the influence of density on the injury. Very few plantations over thirty years of age were found, and those which were studied had been thinned to such an extent that no accurate data were available except for the few trees remaining and therefore no plots were laid out. In addition to this the number of plots would be too few to allow for a fair study. Table II shows the amount of weeviling in the different pure stands of white pine.

TABLE II.—To show the Decrease in Weeviling with the Increase in Density in Young and Old Pure Stands, on Acre Basis.

No. of plots	Kind of stand	Approx. age	Density	Average density	Density wh. pine	Wh. pine weeviled	Percent weeviled
1	Natural.....	15 or less...	0- 500	480	470	190	40.4
2	Natural.....	15 or less...	501-1000	838	838	333	39.7
1	Natural.....	15 or less...	2001-3000	2820	2820	180	5.7
2	Natural.....	15 or less...	3001-4000	3310	3290	230	7.0
2	Planted.....	15 or less...	0- 500	357	357	242	67.8
11	Planted.....	15 or less...	501-1000	738	738	363	49.2
13	Planted.....	15 or less...	1001-1500	1239	1212	447	36.9
5	Planted.....	15 or less...	1501-2000	1860	1860	468	25.2
1	Planted.....	15 or less...	2001-2500	2448	2148	696	32.4
3	Planted.....	15 or less...	4000 plus	4860	4860	67	1.4
3	Natural.....	16-30.....	0- 500	397	393	182	46.3
7	Natural.....	16-30.....	501-1000	728	705	421	59.7
3	Natural.....	16-30.....	1001-1500	1188	1162	623	53.6
1	Natural.....	16-30.....	1501-2000	1632	1592	856	53.8
2	Natural.....	16-30.....	2001-3000	2645	2600	915	34.4
2	Natural.....	16-30.....	3001-4000	3470	3460	735	21.2
2	Natural.....	16-30.....	5000 plus	6490	6404	134	2.1
1	Natural.....	16-30.....	9000 plus	9040	9040	280	3.1
8	Planted.....	16-30.....	0- 500	284	272	220	80.9
14	Planted.....	16-30.....	501-1000	764	763	392	51.4
14	Planted.....	16-30.....	1001-1500	1207	1207	529	43.8
6	Planted.....	16-30.....	1501-2000	1764	1760	710	40.2
3	Planted.....	16-30.....	2001-2500	2342	2342	835	35.6
1	Planted.....	16-30.....	4000 plus	4840	4840	1160	24.0
26	Natural.....	30 or more	0- 500	288	271	207	76.4
6	Natural.....	30 or more	501-1000	732	661	401	60.7
2	Natural.....	30 or more	1001-1500	1368	1265	683	54.3
1	Natural.....	30 or more	1501-2000	1840	1820	660	36.3
1	Natural.....	30 or more	2001-3000	2840	2760	180	6.7

Data on Mixed Stands

Study of one hundred and twenty-two plots in the mixed stands must be approached from a rather different angle from that used in examination of the plots in the pure stands. There are, as before, the artificial stands and the natural stands, but in addition there is also the question of the compositions and the groupings. If it were possible to go back one hundred and fifty years, it would be found that there was practically no pure pine on the heavy soils. White pine formerly grew with hardwoods and to a lesser extent with softwoods—either singly or in groups.

If the mixture be with another conifer, there may be adequate protection in one locality and very little in another. Whether or not the white pine will be protected will depend on the height of the accompanying conifer, and also on its liability to weeviling. The rate of growth of each species must be considered, and if the white pine will grow as fast as the other species the protection value may be very slight; whereas, if the white pine is younger, or grows more slowly, so that it may be shorter in its earlier years at

least, it may receive enough protection to enable it to escape severe attacks.

The choice of the other species of a mixture should be determined by the economic conditions in any given locality. For example, in central New England a white pine—hardwood mixture, groupwise in make-up if possible, offers excellent protection but care must be taken in regions where the gipsy moth is abundant to avoid the use of hardwoods which may be subject to excessive defoliation by this insect. Ash, sugar maple and yellow birch are some of the better hardwoods, the foliage of which is not favorable food for the larvae of the gipsy moth; whereas, the red and white oaks are particularly subject to attack. In Maine and the Maritime Provinces, there is more likelihood of a market for pulp wood, from both broadleaved and coniferous species, and consequently a mixture with a species valuable for its pulp properties would be suitable provided it furnished the desired protection. Except where fuel wood is desired and has a ready market, it is doubtful if it would be advisable to allow the poorer hardwoods, such as gray birch and fire cherry, to grow.

In Graph III is shown the difference in amount of weeviling which can be expected in individual trees in pure old field stands of low density and individual trees from mixtures of white pine and the better hardwoods. The data for these curves were obtained from a large number of trees. The trees, in the pure stands, were climbed and the height above ground at which each weeviling occurred was measured. This was arbitrarily taken as one foot above the crook or fork, on the assumption that the eggs were laid at that point. The trees in the mixed stands were not climbed because of the lack of branches, but the weevilings as high as 40 feet were noted. This graph does not take into account the percentage of trees weeviled per acre, but the sample plot records show that there is a distinct decrease in percentage of pine trees weeviled in the mixed stands.

White pine growing in mixture with other conifers is also a common occurrence throughout the range. When growing with older hemlock, which therefore was taller in the younger stages of the stand, it is usually of good form and quality. However, hemlocks of the same age in the younger stages will be shorter and therefore will not offer adequate protection. In parts of Maine and the Maritime Provinces white pine grows naturally with the spruces and balsam fir. In most of the stands studied, the pines are now dominant and of good form, but the number of white pines per acre is low. Weeviling has occurred only occasionally. The annual growth of the spruces is usually less than that of the white pine and unless they are older there will not be adequate protection. There is no danger of whipping

in these stands, however, and a group-wise system of planting might be made in localities where these species are already present in groups in large numbers.

It appears that white pines growing with red pine, pitch pine or larch, whether naturally or in plantations, are generally weeviled considerably unless the white pines are in the understory. In Nova Scotia a number of stands of a red pine-white pine mixture were found where the red pines were much older. Here protection seems adequate, but in plantations where the two species have been planted in alternate rows and at the same time, weeviling of the white pine is considerable. In all white pine-larch mixtures observed or studied, the pine was considerably damaged by the weevil.

At Great Bear Springs near Syracuse, New York, a Scotch pine-white pine mixture, in alternate rows, gives good protection. Examination of data obtained from sample plots laid out in this mixture and from an adjacent pure white pine stand gives conclusive evidence of the protection afforded by the Scotch pine. Three plots (paralleling one another), two in the mixed and one in the pure stand of the same size and under apparently similar site conditions, were taken. In 1928, the percentage of trees weeviled in the plot in the pure stand was 83.5. The percentage of white pine weeviled in the plot in the mixed stand immediately adjacent to the pure stand was 56.4, while in the other plot it was 28.0. In addition to the fact that there is a smaller percentage of infestation in the second plot, there is less weeviling in the rows of trees farthest from the pure stand. Also the percentage of weeviled trees in the pure stand weeviled more than once is 47.1. In the first plot in the mixed stand this percentage is 24.2, and in the second 28.6. Three other plots in the mixed stand, one in the center of the area and the other two on a western slope at the end of the plantation opposite Plots I and II, showed still further decreases in percentages. The plot in the center of the area has a percentage of trees weeviled of 23.0, while the two on the western exposure have 20.4 per cent and 9.7 per cent of infestation.

The Scotch pine, in the earlier stages of its life, grows faster than the white pine, and its branches tend to spread laterally to a great degree, thus possibly giving a certain amount of protection after the neighboring white pines have been outstripped. Most of the white pines which are in the upper canopy have been weeviled, while those at least two feet below the Scotch pines are not weeviled to any extent. The difference in height growth in these three classes of trees up to 1924 is shown in Graph IV, which shows that at all times after 1914 (five years or more after planting) the Scotch pines were at least two feet above the non-weeviled

white pines and were never more than two feet above the weeviled white pines. Furthermore it will be seen that the weeviled white pines were always taller than the non-weeviled ones.

There are two ways in which white pine is found naturally in mixture with other species, both affording good protection against weevil attack, so long as the white pine is in the lower canopy. These mixtures are either stemwise or groupwise in composition. A stemwise mixture has the species present mixed in singly over the area, and, especially in a mixture with hardwoods, the white pine is in danger of being whipped and overtopped. A groupwise mixture will have groups of pine surrounded by groups of the other species and this composition often occurs after logging, as well as in a virgin forest. The maximum protection is obtained in such a stand where the pines are allowed to dominate the groups in which they are most numerous and vigorous. There will be adequate protection by the species in the surrounding groups and in addition the danger from whipping is minimized. Cline and Lockard (1925) found that, owing to the competition, the size of the white pine group necessary for one survival depends both on the site and on the accompanying species. Ordinarily the groups of white pine should not be larger than one-tenth acre and generally even smaller. Larger groups tend to have the same status as a pure stand, and if the weevils get established they may do as much damage as in a pure stand. Table III shows the amount of weeviling in the different stands of mixed white pine and hardwoods, and mixed white pine and softwoods.

Data on Other Species Liable to Attack

Because of its widespread use in forest planting, Norway spruce is the next most important species subject to weevil attack. Even in widely spaced stands this species tends to straighten much sooner than the white pine, and often will be almost entirely straight by the end of the first summer after weeviling. Very often the new leader apparently comes from an adventitious bud which arises just below the killed portion, instead of from the first living whorl of laterals, as is always the case in white pine. As a result of this, forked leaders are not so common. With the exception of white pine, the percentage of weeviling in Norway spruce is higher than in any of the other species which are more or less commonly planted. No plantations older than thirty years were studied, but in all those between sixteen and thirty years the percentage of infestation varied from 29.2 at a density of 2000 trees per acre to 35.5 at a density of 600 trees per acre. In some localities where both species are found, either together or in separate stands, the Norway spruce is sometimes even more seriously affected than the white pine.

TABLE III. To show the Decrease in Weeviling with Increase in Density in Young and Old Mixed Stands, on Acre Basis.

No. of plots	Kind of stand	Approximate age (years)	Total Density	Class density	Density white pine	White pine weeviled	Per cent weeviled
4	Mixed pine and hardwood stemwise	15 or less..	1027	0- 500	401	78	19.4
5		15 or less..	1915	501-1000	698	140	20.1
4		15 or less..	2320	1001-1500	1133	276	24.4
2		15 or less..	3500	1501-2000	1600	40	2.5
4		16-30.....	746	0- 500	252	70	27.8
2		16-30.....	1475	501-1000	615	252	41.0
10		30 or more	706	0- 500	261	103	39.5
3		30 or more	1170	501-1000	658	313	46.7
2		15 or less..		0- 500	356	32	9.0
3		15 or less..	1800	501-1000	779	181	23.2
2	Mixed pine and hardwood groupwise	15 or less..		1001-1500	1469	387	26.4
1		15 or less..		2001-3000	2690	370	13.8
2		15 or less..		4000-5000	4475	40	.9
1		15 or less..		7000 plus	7420	70	.9
1		16-30.....		0- 500	421	88	20.9
3		16-30.....		501-1000	710	280	39.4
1		16-30.....		1001-1500	1170	370	31.6
6		30 or more	713	0- 500	322	127	39.5
2		30 or more	1105	501-1000	662	65	9.8
1		30 or more	1370	1001-1500	1100	100	9.0
1	Mixed pine and hemlock	30 or more	1800	1501-2000	1690	265	15.7
6		30 or more	873	0- 500	177	48	27.1
1	White pine and pitch pine	30 or more	900	501-1000	620	10	1.6
1		16-30.....	2510	1001-1500	1470	550	37.4
1	White pine and white yellow pine	30 or more	660	0- 500	480	50	10.4
2		16-30.....	1092	0- 500	232	210	90.5
2	White pine and Scotch pine	16-30.....	1200	501-1000	641	488	76.1
4		16-30.....	1041	0- 500	377	154	40.8
4	White pine and Scotch pine	16-30.....	1206	501-1000	627	261	41.6
1		16-30.....	1420	1001-1500	1260	480	38.1
12	White pine and red pine	15 or less..	1020	0- 500	382	160	41.9
1			805	501-1000	566	152	26.9
1	White pine and red pine	16-30.....	1200	0- 500	85	40	47.0
9		30 or more	585	0- 500	261	165	63.2
1	White pine and cedar	30 or more	635	0- 500	270	215	79.6
1		16-30.....	1005	501-1000	625	385	61.6
1	White pine and J. red pine	15	1138	0- 500	416	242	58.2
2		16-30.....	672	0- 500	278	224	80.6
1	White pine and Norway spruce	16-30.....	1590	501-1000	800	440	55.0
3		16-30.....	1090	0- 500	240	117	48.3
1	White pine and mixed conifers	16-30.....	2080	501-1000	520	50	9.6
1		16-30.....	2880	1001-1500	1400	280	20.0
2		16-30.....	3200	2001-2500	2210	630	28.5
4		30 or more	2023	0- 500	205	90	43.9

Study of the plots taken in Scotch pine plantations showed that there is a greater percentage of weeviling in pure stands of this species, just as in white pine stands, than when it is in a mixture. Belyea (1923) reported almost as much weeviling in stands of this species as in white pine at the Great Bear Springs plantations, and more weeviling of Scotch pine than of white pine in a plantation where these species were planted in alternate rows. Belyea and MacAloney (1926) found, however, that nearly all the injury in the Scotch pine at Great Bear Springs, previously attributed to the weevil, was in reality a weather injury. The percentage of trees weeviled in a pure stand of this species is not so high as in a pure white pine stand of equal density, 34.3 per cent

being the highest recorded in the sample plot data, and this was in a density of 102 trees per acre. The lowest density, of white pine of the same age, is 480 and the percentage of infestation is 40.4. In a mixed plantation at Great Bear Springs the average percentage of weeviling in five plots was 0.31 for the Scotch pine and 28.1 for the white pine. There is a considerable amount of Scotch pine planted in the Northeastern States, especially in New York, but the damage due to weather injury is of much greater consequence than that caused by weeviling.

Western yellow pine is not planted extensively in the Northeastern States, but the few plots taken show that the liability to weeviling is not great. An average infestation of 8.0 per cent was found in three mixed stands of a density of 600, and in two pure stands of about 1,200 trees per acre an average infestation of 2.2 per cent was found. All the plantations were under thirty years of age.

One plot of Japanese red pine showed an infestation of 45.4 per cent, and one plot in natural red spruce had 30.6 per cent of its trees weeviled. Both stands were under fifteen years of age. The red pine and the white spruce are so rarely weeviled that it was considered unnecessary to lay out plots.

TABLE IV. To show the Decrease in Weeviling, in Species other than white pine, with Increase in Density on Acre Basis.

No. of plots	Kind of stand	Species	Approximate age (years)	Density class	Total density	Density species affected	Number of weevilings	Percent weeviled
1	Pure...	Norway spruce	16-30	501-1000	620	620	220	35.5
1	Pure...	Norway spruce	16-30	1001-1500	1220	1220	380	31.1
1	Pure...	Norway spruce	16-30	1501-2000	1990	1990	580	29.2
3	Mixed...	Norway spruce	16-30	0- 500	754	416	163	39.2
2	Mixed...	Norway spruce	16-30	501-1000	1407	688	213	31.1
1	Pure...	Scotch pine...	15	0- 500	102	102	35	34.3
1	Pure...	Scotch pine...	15	1001-1500	1220	1210	120	9.9
2	Mixed...	Scotch pine...	16-30	0- 500	974	243	39	16.2
7	Mixed...	Scotch pine...	16-30	501-1000	1208	664	21	3.2
2	Pure...	White yel. pine	16-30	1001-1500	1200	1112	25	2.2
3	Mixed...	White yel. pine	16-30	501-1000	1061	578	45	8.0
1	Pure...	Red spruce...	15	4000 plus	4920	3920	1200	30.6
1	Mixed	Jap. red pine..	15	501-1000	1138	722	328	45.4

In Table IV is shown the amount of weeviling in species other than white pine, where there was a sufficient number of trees present in the stand to warrant laying out the plot.

Effect of Silvical and Habitat Factors

Composition of the Stand

Within a few years after the inception of management policies on the Harvard Forest it was found that wherever white pine occurred with hardwoods or hemlock the amount of weeviling was

small. Fisher and Terry (1920) stated that "The weevil, however, is less numerous in the mixed growth of cut-over lands than in the pure plantation." Previous to this, however, both Hopkins (1907) and Blackman (1916 and 1919) wrote that an unfavorable condition for injury was mixed stands of pine and hardwood. It has generally been considered that this is because the pines are shaded from the sun and the surrounding trees act as a barrier against flight. This barrier action would, of course, prevail in the early spring when the hardwood trees are bare as well as in the fall before the leaves have fallen. This is during the time when the air temperature is sufficiently high to permit general flight, and undoubtedly this is a very important means of keeping the weevil from such stands. Pyle (1927) in Pennsylvania found, by experimentally classing different degrees of shade from 0.1 to 1.0, that below 0.3 shade there was a much greater amount of weeviling than above, with the optimum for protection against weevil attack between 0.4 and 0.7. Above this latter degree there was whipping which, in its action on the leaders, is as damaging as weeviling in pure stands. Pyle has come to the conclusion that a correlation of facts seems to indicate that shade density, and thus the character of the growth in the pine, plays an important role in the severity of weevil attack.

These reasons partly explain the small amount of infestation, but the fact that groups of pines surrounded by hardwood are not weeviled to any extent, even though pure pine stands nearby are weeviled repeatedly, indicates that there are other reasons. It was found, in experimental cages, that adult weevils will feed more readily on shoots cut from pine growing in mixture with hardwoods. Also the females will almost invariably go to the most vigorous leaders to oviposit. Additional experiments, similar to those just stated, were conducted on pine shoots from pure open growing stands and from pure dense stands to determine if possible whether the same reaction prevailed on the part of the weevils. As in the former case, it was found that they fed more readily on the shoots from the open growing stands and the females oviposited in them first. Indeed as a general rule the shoots from the dense stands were not attacked to any extent until the others were practically riddled with holes.

The reason why the leaders from pure stands are most favorable for feeding and oviposition is a complicated one. Above all other things the importance of the odor is probably paramount, and other factors are also probably concerned. The leaders in such stands are, on the average, larger and more vigorous than those on trees in either a dense pure stand or a mixed stand. The needles are more numerous, longer and thicker, and the buds are bigger; there is more bark area and consequently a greater number of lenticels.

The inner bark is thicker, and thus has more resin in it. Therefore there are more outlets available for the emanation of the odor from this larger amount of resin. The moisture content in the average sized leader from an open stand is greater than in the average sized leader from either of the other stands. This was ascertained by removing a number of leaders from the different stands and weighing them. Then they were placed in an oven and completely dried, after which they were weighed again. The loss of weight in each case represented the amount of moisture which was in each bundle of leaders. Then the average was taken. The greater amount of moisture in the leaders from these open growing trees is because of the average larger size, and this also makes them more suitable for attack. All these conditions tend to make these more vigorous leaders in the open stands more attractive to the ovipositing beetles.

Age, Height and Vigor

Weeviling usually begins when the trees are two or three feet high and five or six years old, but sometimes earlier. When trees two feet high are weeviled they usually are killed at least to the lowest whorl of laterals, thus often resulting in a scrubby tree. Sometimes the trees may be killed back under the ground and in such cases they usually die. Such an injury is very similar to that caused by *Pissodes approximatus* Hopk. or *Pissodes nemorensis* Germ. and may very easily be mistaken for either. Beginning with a height of two or three feet the rate of infestation in a widely spaced old field pure stand becomes increasingly greater, reaching the peak when the trees are twenty to twenty-five feet high. From this point it gradually decreases until at a height of sixty feet it has practically ceased.

From observations made during the field seasons of 1925 and 1926 it appears that in old field stands about one-half of the total weevilings in individual trees occurs before the trees are thirty feet high. Also it is apparent that there is a greater percentage of weeviling in the present 25- and 35-foot classes than in the basal 25 and 35 feet of the other height classes, because the widespread planting of white pine (the favorite host of the white pine weevil) has created a condition conducive to rapid multiplication. It will be seen in Graph V that there has been a marked increase in the percentage of weeviling in the lower height classes in the last thirty years in low density old-field stands on the medium soils of the abandoned farms. For this graph the weevilings in dominant, co-dominant and intermediate trees below 25 and 35 feet were taken in all of the height classes from 25 to 85 feet. The trees in the higher height classes came in on the farm land when the abandonment began after the Civil War. With the increase in reproduc-

tion from decade to decade and the added food supply caused by the widespread planting from the beginning of the present century, it was inevitable that there would be an increase in the amount of weeviling. In many plantations, where conditions for attack are especially favorable, the infestation is even greater than in these old field stands. Owing to the difficulty which would necessarily arise in attempting to ascertain the year and age when the weeviling occurred in the older trees the height classes were arbitrarily chosen as the means of collecting the data.

While the weeviling above thirty feet is considered of little importance under the present silvicultural practice of growing white pine in short rotations, it will be of importance in longer rotations if clear lengths above this height are desired. If the present slack market persists in the boxboard industry clear lumber must be produced if stumpage is to be sold at a profit. In any case it would be worth more, as clear lumber is of more value than lumber which has crooked grain and large and sometimes loose knots.

As stated previously one factor limiting attack is the vigor of the tree. Those leaders which are attacked are usually those which have been most vigorous. Peirson (1922) states that the weevil in flying over the stand will choose only those trees which are tallest, while Graham (1926) points out that he has observed that the trees with the most vigorous leaders, and not necessarily the tallest ones, are most susceptible to infestation. Fitch in 1858 stated that "young thrifty-growing pines are its favorite resort, and among those it selects those that are the most vigorous and whose topmost shoots have made the greatest advance the preceding year. But I have seen it so numerous that not only the topmost shoots of every tree in the grove, but many of the lateral ones also were invaded and destroyed by it." From the writer's observations it appears that Fitch's account is most nearly correct. A large number of weeviled trees have been observed, where the trees were higher than those surrounding them, and where their leaders were not as long as those on some of the co-dominant trees nearby which were not attacked. That is, as Fitch stated, the weeviling is most apt to occur on those leaders which made the greatest growth the year previous. In heavily infested plantations in central Massachusetts, where the annual infestation is often over 50 per cent, it is apparent that both the vigorous and some of the less vigorous leaders must be attacked. The trees most often attacked are those in the upper canopy, or the dominant and co-dominant trees, and these are generally the ones with the most vigorous leaders. This holds in pure stands of varying densities, but the pine trees in mixed stands of pine with certain hardwoods, such as maple, ash, birch, oak, aspen, etc., are normally not in these two height classes until they have reached the height of thirty feet or

more. This partial suppression by the accompanying species is one of the reasons why pine trees in these stands are relatively immune to weevil attack until the leaders are above the crown canopy of the other species.

Site Factors

Site is of great importance as a factor determining the rate of infestation. It is generally understood that the white pine will grow on almost any site, but the fact remains that it will not thrive equally on all of them. From observations that has been made during the course of this investigation it is also apparent that weeviling is more prevalent in stands growing on certain sites and soils. In a previous paper (1926) the writer rather obscurely stated that stands on light sandy soils generally show the greatest percentage of infestation. Other conditions being equal, such a stand may have as many trees weeviled per acre, but there will not be as much permanent damage done as in a stand on the medium agricultural soils where the typical old field stands are usually found, and where many of the pure plantations have been set out. Weeviled trees on such a soil will be weeviled more frequently than those on a light sandy soil, because the growth is more vigorous, and consequently the leaders are more attractive to the beetles.

A stand on a wet, cold soil has little weeviling, but the growth conditions are also poor. In this case, the trees when weeviled are more apt to be killed back several years, so the injury is more severe in the individual cases than where the leaders are vigorous. Mixed pine and hardwoods are more prevalent on the richer soils. In such a stand where the pine has a chance to develop and is not suppressed, the growth is generally good and when the trees are weeviled the recovery from the injury is much quicker than in the trees weeviled in pure stands. As a general rule most of the old field stands and plantations in the Northeast can be classed as being on Site II, or a little better than Site III, and it is in such stands that the most severe permanent injury is found.

Exposure

The exposure also has a direct bearing on the amount of weeviling. As a rule stands on southern and eastern exposures (those directly exposed to the morning sun where the air temperature is higher earlier in the day than on the other exposures) are attacked with greater intensity. This is especially noticeable where there are different exposures in the same stand. When a stand on a western or northern exposure is weeviled to any extent, it generally can be explained by the fact that there is little pine nearby on other slopes, or the weevils have become established there and have not migrated to other places.

On the protected exposures the steeper the slope the less will be the weeviling, and most of what does occur will be at the top of the slope. That the weevils might also fly over trees on a protected slope to get to those at the top of the slope or on the other side, where the air temperature is higher earlier in the day, is shown by the following instance: Near Barre, Massachusetts, there is a pine plantation of several acres which is partly on a medium to steep west slope, and partly on a gentle southwest slope. Site conditions appear to be similar. The nearest stand of pine is across the road at the bottom of the west slope and approximately two hundred yards distant. It is probable that most of the weevils which caused the weeviling in the plantation came from this stand, as there is no other within one mile. The first weeviling in the plot on the southeastern slope occurred in 1919, and by 1922 there were 18 weevilings in 17 trees. In 1922 the first trees in the plot on the western slope were weeviled—six being attacked. The weevils necessarily had to fly up and over the west slope and toward the morning sun in order to get to the trees on the southeast slope. Up to and including the year 1926, the percentage of trees weeviled at least once in a tenth-acre plot on the west slope was 38.3 and in a plot of the same size on the southeast slope the percentage was 60.0. The total number of trees in each plot was not the same—on the western slope there were 107 trees, and on the southeastern slope there were 75 trees. In the first named plot, 45 trees, and in the second 41 trees, were weeviled. The total number of weevilings on the southeastern slope was 66, as compared with 51 on the western slope; an excess of 15 weevilings or 29.4 per cent.

In 1924, the feeding and oviposition, in a plantation at the Rockwell Springs Experiment Station of the New York State College of Forestry, were heavier on a southeast slope and the top of a small knoll facing that slope than on the northeast and northwest slopes of the knoll. One leader on the southeast slope had sixteen weevils on it at one time during the feeding period, and the average per leader occupied was six. On the northwest slope the most found on any one leader was six, and the average per leader occupied was about three. The trees in the plantation were of the same age, practically the same height, and the other site conditions seemed to be uniform. Another thing which must be considered is that these weevils came in from outside the plantation that season, as the leaders weeviled the previous season had been removed and burned before the new generation had matured and emerged. The percentage of trees weeviled on the southeast slope was also higher than on the others.

It is apparent that a difference in the temperature on the different slopes offers an explanation of the importance of this factor; that is, the morning sun will cause a higher temperature

on the slopes exposed first, and general flight would therefore be to that slope first, as the weevil is a heat-loving insect. On the steep eastern and southern slopes there may be more weeviling than on the gentle ones, as there tends to be a "terraced" effect that allows more light and sunshine for each leader. On the protected exposures the steeper the slope the lesser will be the weeviling with most of what does occur at the top of the slope.

Climatic Factors

As stated in the section on Life History, the weather is a factor affecting the development of the weevil from the time the eggs are laid and hatched as well as in the resumption of activity. This limits the time of activity in each locality but the local weather factors as a whole also have much to do with the degree of infestation. A wet, cold spring, or an abnormal amount of precipitation during the feeding and oviposition periods, undoubtedly limits the infestation. Lack of sunlight and warm weather materially affect these periods, as the white pine weevil prefers to work in the sunlight. Likewise such a poor growing season results in a decreased amount of the stored food necessary for the early part of the next season's height growth, and also in a decrease in the numbers of the new generation of weevils, because the leaders will not be as vigorous as if the growing season were good, and they are therefore not so suitable as a breeding host. This possibly may be one factor to explain the rise and fall in the intensity of infestation from time to time.

There is probably no other time during the life of the weevils that is as critical as the hibernation period. The mortality of hibernating beetles must be enormous. Thus, when there is an open winter, preceded by a wet autumn, as happened in New England in 1927, the great and sudden changes in temperature and rainfall are sufficient to kill the beetles. It has long been known that it is not necessarily excessive cold that kills insects in hibernation but rather the rapid changes from one extreme to the other. If local climatic influences peculiar to the various parts of New England where the permanent plots are established are left out of consideration, the plot data show that there was a decrease in 1928 in the young stands where the weevil infestation had been increasing for a number of years. This decrease, while slight in some localities, was heavy in others and seemed to be general over the entire New England region affected by the heavy rains in the fall of 1927 and the open winter which followed.

The climatic difference in certain portions of the range of the white pine also limits the degree of infestation. In Vermont, for example, the pine is found mostly in the rich valley bottoms, where conditions for development of the weevil, and subsequent hiberna-

tion, are rather unfavorable. This can be explained by a variety of factors, chief among which are, (1) the late spring and early fall frosts followed by a rise in temperature often accompanied by rain, (2) the protection afforded by the slopes, and (3) the site conditions, which are such that when weeviling does occur the injury is not so serious as in other regions. The growth of the pine under these conditions is sufficiently vigorous to enable the injured individual to recover relatively early. On the other hand, scattered stands on sandy soils in Vermont are weeviled to nearly as great an extent as those in central New England. One reason for this is that the conditions for hibernation are favorable, and more weevils may survive.

Scarcity of Natural Food Supply as a Limiting Factor

In Vermont, in addition to the influences stated under climatic factors, the scarcity of natural breeding places must also be considered. In some localities no natural white pine can be found for miles, and plantations set out in these areas have seldom or never been attacked. The Bass River plantations in southern New Jersey, the plantations of the York Water Company near Jacobus, Pennsylvania, and the Biltmore Plantations near Asheville, North Carolina, are examples where scarcity of breeding places over an area of several miles has either prevented attack or has kept it at a minimum.

Toward the more westerly and southerly portions of the range of the white pine, weeviling is not severe. Pine in these sections is found in scattered clumps. The nearer the limits of the range the lesser will be the number of trees and consequently a decreased amount of food material. Thus we will find the infestation falling off as food material decreases and at the limits of the range of the white pine there will be little, if any, weeviling.

Quite often where there has been no white pine on islands in lakes or reservoirs, natural pine seedlings have come in or plantations have been set out. Such a case is found at the Cross River Reservoir of the New York City Water Supply Commission. A white pine plantation set out there on an island where there was no natural pine escaped injury for twelve years. At that time one or two trees were weeviled and in four years' time the infestation was firmly established. It is probable that the first weevils in that particular plantation were wafted across the water, about a quarter-mile, by a favorable wind, and at the present time more than half of the trees have been weeviled, and an adjoining Norway spruce plantation is also infested.

The white pine has also been set out in plantations extending over 2,000 acres in western Montana and northern Idaho. Up to the present time no weeviling has been reported and the growth in one

plantation set out in 1911 is nearly double that of western white pine of the same age growing naturally within the bounds of the plantation.

Weymouth pine, as the white pine is known in Europe, has been used extensively in forest plantations in different countries there. While there are insect pests of lesser importance affecting this pine in Europe it is fortunate that no insect as important as the White Pine Weevil is present. It is doubtful if the white pine weevil will ever obtain a foothold in Europe, for two reasons. In the first place the white pine was taken over as seed and germinated there, and in the second place, if the insect did occur European Forest Managers would in all probability institute direct control measures which, under the existing circumstances are impractical in wide spread areas here.

CONTROL MEASURES

In all efforts to control or prevent excessive loss from depredations of forest insects it should be remembered that as a rule it is useless to attempt the complete extermination of any species. In the case of tree killing bark borers, such as members of the genus *Dendroctonus*, it is now considered sufficient if enough of the brood is destroyed to prevent it from continuing an aggressive attack. The remaining ones must then occupy a defensive position against their own enemies until conditions, often due to avoidable negligence and mismanagement by the owners of the forest, favor its again becoming epidemic. Control of the white pine weevil presents a different problem. The trees themselves are not often killed, but the leaders are. The result is a tree which may be crooked or forked, or both, depending on the number of times the leading shoots are killed, with a subsequent loss in timber value. In the case of the tree killing bark borers the progeny of many females is necessary in order that a tree may be girdled and killed and the brood successfully complete its life cycle, while one fertilized female weevil may lay in one leader, and possibly in more than one, enough eggs to cause its death once the larvae start their work. But the numbers of larvae necessary to kill a leader is comparatively small, while many hundreds of the bark beetle larvae must be present in order to kill a tree.

The problem of keeping the White Pine Weevile in check and controlling the injury resulting from weeviling must be considered from a number of angles. Among the most important points requiring consideration are the cost and the completeness of the control measures. For convenience these measures may be classed as Natural Control, Prevention of Attack, Direct Control of the Beetles, and Repair of the Injury.

Natural Control

The Climate

The most important natural control factor is the climate. It has already been discussed fully and, although it can not be manipulated by man, is mentioned here to emphasize its importance.

Parasites and Predatory Insects

Control of the White Pine Weevil by parasites and predatory insects is at best a very uncertain method. This measure can not be left out of consideration, however, as these classes of insects are undoubtedly of great value. In some areas the parasites and predatory insects undeniably have been the most important means of lessening the amount of weevil emergence from the leaders. In the main, however, control by this means is in the nature of a remedial or curative measure rather than a preventive one. A large laboratory, especially fitted for the breeding and liberation of these insects, would be necessary in order that this method could be given a fair trial. The following is the complete list of insects, collected or bred from weeviled leaders by the author, or reported from other sources:

PARASITES

Hymenoptera

<i>Eurytoma pissodis</i> Gir.	<i>Rogas aciculatus</i> Cress.
<i>Eurytoma</i> sp.	<i>Eupelmus pini</i> Taylor.
<i>Microbracon pini</i> Mues.	<i>Rhopalicus pulchripennis</i> Cwfd.
<i>Doryctes</i> sp.	<i>Calliephialtes nubilipennis</i> Vier.
<i>Coeloides pissodis</i> Ashm.	<i>Calliephialtes comstockii</i> Cress.
<i>Coeloides</i> sp.	<i>Schenkia</i> sp.
<i>Spathius canadensis</i> Ashm.	<i>Phygadeuon nitidulus</i> Prov.

Diptera

<i>Lonchaea corticis</i> Taylor.	<i>Muscina stabulans</i> Fall.
<i>Gaurax apicalis</i> Mall.	<i>Compsilura concinnata</i> Mg.

Lepidoptera

Undetermined Pyralid.

HYPERPARASITES

Hymenoptera

<i>Deleter undulatus</i> Davis (on Diptera).	<i>Homoporus</i> sp. (on Diptera).
<i>Pleurotropis</i> sp. (on Diptera).	<i>Berecynthus</i> sp. (on Lepidoptera).

PREDOCIOUS INSECTS

Formicoidea

Pheidole sp.

Coleoptera

<i>Tenebrionidae.</i>	<i>Thanasimus dubius</i> Fabr.
<i>Elasmocerus (Monophylla) terminatus</i> Say.	<i>Enoclerus</i> sp.

ASSOCIATED SPECIES (Leaf Feeders, Scavengers, etc.)

Hymenoptera

<i>Hylaeus</i> sp.	<i>Megaspilus</i> sp.
<i>Hemitaxonus dubitatus</i> Nort.	<i>Prosynacra</i> sp.

Lepidoptera

<i>Dioryctria abietella</i> D. & S.	<i>Pinipestis zimmermanni</i> Grote.
<i>Eucordylea atrupictella</i> Dietz.	

Diptera

<i>Scaptomyza graminum</i> Fall.	<i>Leptocera</i> sp.
<i>Forcipomyia specularis</i> Coq.	<i>Hippelates</i> sp.
<i>Oscinella cocendix</i> Fitch.	<i>Tachydromia</i> sp.
<i>Chymomyza amoena</i> Lw.	<i>Aphiochaeta</i> sp.
<i>Madiza glabra</i> Fall.	

Parasites. The most important primary parasites are *Lonchaea corticis* Taylor; *Eurytoma pissodis* Gir., *Microbracon pini* Mues., *Doryctes* sp., and *Coeloides pissodis* Ashm. The first two are external feeders, while the last three feed internally. Graham (1926) reported 50 per cent parasitism by *Eurytoma pissodis* in several pine clumps near Ithaca, N. Y., and states that in this general locality it was by far the most important parasite of this weevil. During the summer of 1927 the author found several areas in New Brunswick and Nova Scotia where weeviled leaders were heavily parasitised, many of them showing nearly 100 per cent parasitism. The species most common seemed to be *Eurytoma pissodis*, followed closely by *Lonchaea corticis*. The latter, which is the species called *laticornis* in a previous article by the writer, seems to be almost as common as the former over the entire range of the weevil. Graham (1926) reported *L. rufitarsus* from Ithaca, but it seems probable that it is the same species which the writer has taken in New York, as well as in Massachusetts, Connecticut and the Maritime Provinces.

Microbracon pini is very common all over Massachusetts and Connecticut, and is nearly as numerous as the two preceding. Neither *Doryctes* sp., nor *Coeloides pissodis* are as numerous as the others, but apparently have approximately the same range. An undertermined species of *Coeloides* and one of *Eurytoma*, both of which may be varieties of the ones already mentioned, have also been found in Massachusetts. *Spathius canadensis* has previously been reported from other Coleopterous hosts and there seems no doubt but that it is a primary parasite. Several species

of *Rhogas* (*Rogas*) have been reported as primary parasites on the larvae of the army worm (Britton, 1914) oriental peach moth (Stearns, 1919), fall webworm (Dustan, 1923), canker worm (Porter & Alden, 1924) and the green clover worm (Hill, 1925), in North America, and also on the larvae of various insects in France, Russia and India. *R. aciculatus* can therefore apparently be considered as primary on the white pine weevil.

Eupelmus pini has already been described and reported by Taylor (1927). There is some doubt whether or not *Phygadeuon nitidulus* Prov. is primary on the white pine weevil, as in this genus we find both primary parasites and hyperparasites. In all the literature available there are more of this genus considered as primary than as secondary species, and at least one species, not determined, has been reported by Franklin (1921) from Massachusetts as primary on the larvae of the Cranberry Girdler. No definite evidence was obtained by the writer, but it has been placed as a primary parasite because of the number of other species of this genus being primary.

According to Klimesch (1923) *Ips typographus* in Europe is at times heavily parasited by *Rhopalicus suspensus* (Latz). The species of this genus bred from weeviled leaders, *R. pulchripennis*, is also primary. The two species of *Calliephialtes nubilipennis* (Vier.) and *comstockii* (Cress), are undoubtedly primary, although *comstockii* has generally been considered as primary on a Lepidopteron. Cushman (1915) described a species of this genus, *thurberiae*, which was reared from larvae of the Thurberia Cotton Boll Weevil. Leiby (1922) refers to a species of this genus as primary on the Goldenrod Gall Maker, and Gill (1925) also found a species primary on the Pecan Nut Casebearer. The Ichneumon *Schenkia* is probably a primary parasite in common with the vast majority of this family.

Among the Dipterous forms primary on the white pine weevil *Lonchaea corticis* (Taylor) is by far the most important. Properly speaking this species is a predator, as the larvae, after devouring the weevil larvae, may go along the tunnels and attack others. During the summer of 1926 the life stages of this species were studied. The adults were present in large numbers almost from the time the female weevils began ovipositing and continued so until the middle of July. Mating was not observed, so it is not known when fertilization takes place. Several eggs are laid either in the egg cavity or in the larval tunnel of the weevil. These eggs are one millimeter long and a half-millimeter thick and are sculptured with fine lines along the long axis and are pearly white in color. They hatch in a week to ten days, and the larvae, if they are not already beside a weevil larva, crawl along the tunnel until they find one. They attack it, fastening

themselves by means of their mouthparts, and devour it in a short time (Plate X-1, 2). Usually several attack one weevil larva so that it is devoured completely with only the chitinous covering and the head capsule left as evidence of their work. They then proceed along the tunnels looking for more food. Pupae are also attacked by this species. The fly larva goes into the pupal stage about the same time as those weevil larvae which escape death, and the new generations of both fly and weevil are ready to emerge at about the same time. Many of the flies, however, winter in the leader as pupae and emerge the following spring. No information is available on the hibernation quarters of the adults which emerge during the summer or early fall, but it is believed that, like closely related Muscid flies, they may hibernate in almost any place that will give adequate protection.

No mention has been made in the literature of the genus *Gaurax* as to whether or not the species *apicalis* Mall. is a primary parasite, and the author has not been able to determine this definitely, although it apparently is parasitic at times.

Muscina stabulans Fallen is generally considered as a scavenger, but there is excellent evidence to support the writer in placing this species as a primary external parasite, although, like *L. corticis*, it really is a predator. A considerable number of small larvae, looking not unlike those of *Lonchaea corticis*, were removed from weeviled leaders with weevil larvae with the intention of studying their feeding habits. Indeed it was not until about a week later that their size attracted the attention of the writer. At the time they were removed they were about the size of half-grown larvae of *L. corticis*. In a short time, however, they began to grow much larger than the larvae of this species and when they had pupated it was fully realized that they belonged to another genus. When they emerged they were determined as *Muscina stabulans* Fallen. Britton (1914) reported it as unmistakably a parasite on the larvae of the army worm in Connecticut, and Smith (1921) reared it from larvae of the fall army worm. It has also been reported by Herold (1923) from Germany bred from the larvae of the cutworm *Euxoa segetum* Schiff.

Webber and Schaffner (1926) report that Mr. C. W. Johnson, of the Boston Society of Natural History, recovered one specimen of *Compsilura concinnata* Meigen from the white pine weevil. This specimen was found in a mount containing a weeviled white pine shoot and obviously must have emerged after the mount was prepared. There was no puparium in sight and the maggot had apparently pupated within the burrow of the host.

Hyperparasites. At least four of the species bred from the weeviled leaders are hyperparasites. Three of them, *Deleter un-*

dulatus Davis, *Pleurotropis* sp. and *Homoporus* sp., are parasitic on Dipterous insects and the fourth, *Berecynthus* sp., is parasitic on Lepidoptera.

Predacious Insects. A predacious insect differs from a true parasite in that it may feed upon a number of victims, whereas the true parasite completes its development on one host. Aside from the parasitic insects affecting the white pine weevil there are numerous predatory insects which do their share in decreasing the numbers of this pest.

Packard (1885) reported that the larvae of Tenebrionid beetles are very commonly predacious on the white pine weevil. Hunter and Pierce (1912) reported that the Mexican cotton boll weevil is attacked by an ant of the genus *Pheidole*. Specimens of the same genus have been bred from weeviled leaders by the author. Wheeler (1913) stated that this is one of the Harvesting Ants, and that pieces of other insects are carried to the nests to be used as food. No mention is made by him that this genus is predatory, but it is quite possible that the white pine weevil may be attacked in the same manner as the cotton boll weevil.

Graham (1926) reports the rearing of a Clerid, *Elasmocerus* (*Monophylla*) *terminatus* Say, from a weeviled leader, and that larvae have been found in the terminals evidently feeding on the weevil larvae. The writer has taken adults of *Thanasimus dubius* Fabr. in the act of running up and down leaders which were already weeviled or in which eggs were being laid, but they were not actually seen feeding on the weevil adults or larvae. A species of *Enoclerus*, in the larval form, was taken from a weeviled leader in the Cumberland Valley region in southeastern Pennsylvania, and the presence of partly devoured weevil larvae is good evidence of its predatory nature. However valuable these Clerids may be, it must not be forgotten that they will also feed on larvae of the parasites present as well as on the weevil larvae, so that they may be killing off species which would otherwise help in reducing the numbers of the weevil.

Plate X-3 shows a larva of an undetermined Pyralid moth in a pupal chamber of the white pine weevil. These are often found when the pupal chambers are constructed in the pith. In the particular case photographed the larva was removed from the elongated cavity which was formed by the weevil larva when it bored through the wood into the pith and made its pupal chamber. A number of fragments of a young adult of the weevil were found in this cavity. Microscopic examination of the material showed that the body tissue was fresh when it was taken, as it showed a clear white color and the cellular structure was firm, and the light brown coloration of the body wall indicates that the weevil was a young adult. Had the weevil been dead

when the moth larva began to feed upon it, there would have been a discoloration in the muscular tissue and the cell structure would have been altered.

As stated previously, parasites and predators have indubitably been the cause of decreasing the numbers of weevils in certain areas. However, in certain other areas, in the Northeastern States and the Maritime Provinces, examined by the author, there was very little parasitism, which bears out the contentions advanced by many entomologists that control by parasites and predators is uncertain. Particularly is this true if the pest is native, because in so far as parasites and predators are concerned, the natural or biotic balance is more firmly established than it would be with a recently imported pest. The breeding conditions may also be so favorable for the pest that it can multiply at a greater rate than its enemies.

Associated Species. Sixteen associated species, belonging to the orders *Hymenoptera*, *Diptera* and *Lepidoptera*, have been bred from the cages containing weeviled leaders, taken in the field, or reported by other investigators. One of the *Lepidoptera*, *Eucordylea atrupictella* Dietz, is a leaf feeder. Britton (1916) reports that injury by the Pine Tip Moth (*Pinipestis zimmermanni* Grote) was particularly apparent in connection with the tunnels of the weevil. At the same time he reported breeding the moth *Dioryctria abietella* D. & S. from white pine leaders. It apparently has been considered as a scavenger at times and a wood borer at others, although it is generally known as a cone borer. The writer has also bred this moth from weeviled leaders.

Certain flies of the family *Phoridae*, of which the genus *Aphiochaeta* forms a part, are primary parasites according to Brues (1903). Hunter and Pierce (1912) also list three species as parasitic on the cotton boll weevil, as well as being scavengers. However the probability is that the species bred from the weeviled leaders is primarily a scavenger, in common with practically every other species of the genus, but it may lay eggs on dead material and the larvae feed upon it.

All the rest of the associated species are primarily scavengers in the tunnels of the weevil larvae, with the possible exception of *Hylaeus* sp. and *Hemitaxonus dubitatus* Nort. Both these species were taken as adults from the leaders in the field.

Birds and Other Animals

The value of birds as a means of preventing the increase of the white pine weevil through destruction of the larvae was shown very strongly during the summers of 1926 and 1927. In every place visited it was found that a large number of leaders had had the bark stripped from the wood by birds (Plate X, 4, 5), and the larvae contained therein had been eaten by them. In practically

all the leaders examined (several hundred) all the larvae which had still been feeding at the time the bark was stripped by the birds had been eaten by them or had dropped to the ground where they would die or be picked up by the various ground-feeding species of birds and rodents, which are numerous in such plantations. Many of the pupae, in addition, had been picked out of their chambers. In one heavily infested plantation in Petersham, Massachusetts, covering over three acres, 30 per cent of the weeviled leaders had been stripped of the bark and the larvae had been destroyed. The birds actually observed in various localities comprise the white-breasted nuthatch, downy woodpecker, chickadee, rose-breasted grosbeak and certain warblers which were not identified.

Hopkins (1907) stated that "Woodpeckers also feed on the matured larvae, pupae and adults." McAtee (1926) reports that the Biological Survey has records of the yellow billed cuckoo, English sparrow and bluebird feeding on this pest. It is probable that other species present in the plantations were also feeding on the larvae, but we have no record of such. Fitch (1858) reports "Newly perforated holes larger than the mouths of the burrows—work of small birds which are very efficient in ferreting out and devouring the larvae of this weevil." Felt (1913) reported that woodpeckers preyed on this species, and Forbush (1913) states that "The downy woodpecker is one of the most useful birds to the lumberman, for it feeds on such destructive insects as the . . . white pine weevil. I have seen many shoots from which this insect had been removed by birds, and most of them showed the characteristic work of this woodpecker." The chickadee is also mentioned by this author—"While working among the pine trees I saw several chickadees go to infested shoots, peck them and apparently extract the larvae. These dying shoots seem to be the principal attraction which brought them to the pines."

Graham (1926) also reports the partial check of the weevil in one plantation in New York State by allowing chickens to run among the trees, and the possibility of destruction of the weevils in the fall by the birds which feed along the ground is suggested. In the spring towhees have been observed and heard, by the author, scratching at the bases of pine trees in natural stands, but no positive proof that they were finding weevils was obtained. Observations in the summer suggest the possibility that they may have been searching for larvae which had fallen from leaders stripped by other birds.

The various field mice, shrews and wood mice, are insect eaters, and the fact that runways can be found in the litter at the depth where the weevils hibernate is a good indication that they very probably take their toll of the weevils during the hibernation period.

Prevention of Attack by the Insect

Prevention of attack by the white pine weevil means the saving of the leader and consequently a straight stem at all times. The two methods tried by the writer are (1) spraying with repellents and (2) banding the leader with some material which will prevent the weevil from reaching it except by flying directly to it.

Sprays and Repellents

Experiments have been carried on by at least three other investigators in different parts of the weevil's range to determine the feasibility of stomach poisons and repellents as measures of control. Good results, as repellents, have been obtained by the writer with lime sulphur in a proportion of ten ounces to eight quarts of water, and with lead arsenate in a proportion of two ounces to eight quarts of water. In each case calcium caseinate was added as a sticker and spreader. The cost of spraying has not been given by Walden (1915) or Graham (1926), but the writer found that for lead arsenate it was \$1.45 and for lime sulphur it was \$1.70 per acre. The cost is mostly for labor as both materials are relatively inexpensive. The use of a good adhesive agent in the spray mixture is necessary so that the film of spray will stick to the leader for the entire period of oviposition. The area treated by the writer was about three acres in extent, and was divided into strips. In all the sprayed strips the infestation was reduced about one-half, although the infestation in this particular area was very low—3 per cent on the check strips, 1.7 per cent on the strips sprayed with lime sulphur, and 1.1 per cent on the strips sprayed with the lead arsenate. Up to the time of a heavy rainstorm the percentage of infestation was negligible and no feeding punctures were found. This rain washed off the material and practically all the weeviling happened after this storm. Considerable waste of material was unavoidable as there were no spray nozzles obtainable that would make a "mist-spray" and still concentrate the material in a small stream.

Britton and Walden (1911), in addition to the work done with lime sulphur and lead arsenate, experimented with other materials, but on a small scale. Of these, whale-oil soap, in a proportion of eight ounces to one gallon of water, gave good results with no injury to the leaders, but is more expensive to use than either of the other materials mentioned above. Graham (1916 and 1926) reported experiments with sixteen different materials or mixtures and found only carbolineum and creosote, sprayed on pure, to be completely effective. He did not get good results with lime sulphur, as did Walden (1915) in Connecticut or the writer in Massachusetts. Walden sprayed over eight hundred trees and the writer over eleven hundred, while Graham sprayed only ten, and this may explain the difference in results.

If an effective repellant can be used economically it will mean the saving of the leader and consequently straight stems. At the present time the labor cost per acre per year is too great to allow widespread use of this method in plantations or natural stands. It will be an efficient and desirable means of control of ornamental or shade trees, or in small plantations where the aesthetic value is paramount.

Banding Materials

Banding the leaders with some material which will prevent the weevils from getting to them, except on the wing, is very effective, but the bands must be renewed each season. The rate of weeviling is reduced in severe infestations very noticeably. Two materials, sticky banding material and raw wool, have been used in this work, the bands in all cases being placed above the previous season's growth of laterals.

Sticky tree-banding material. As previously stated in the section on Flight Habits, the central plot of one series in a severely infested plantation had all trees banded with sticky tree-banding material just above the lateral growth of the previous year. In this plot the percentage of trees weeviled decreased from 51 in 1925 to 12 in 1926, while the average for the four check plots increased from 37 in 1925 to 57 in 1926. Thus, while the infestation was very materially decreased in the treated area it was increasing in the check areas. The disadvantages encountered in using the sticky tree-banding material are that it lasts but one year, is very apt to become hardened by the weather and must therefore be inspected frequently, and the cost is prohibitive. The material is expensive and the labor charge will be at least \$2.50 per acre per year.

Raw Wool. Four series of plots were laid out in 1927 in connection with the experiments with the wool bands. Two of these were at the Rainbow Plantations of the Connecticut Agricultural Experiment Station and the others were on the Choate Plantations at Petersham, Massachusetts.

Each of the series of plots at Rainbow consisted of a central plot and four check plots, and all of the plots were one-twentieth acre in area. In one central plot the infestation, the year before the experiment was 3.4 per cent, while the average for the check plots was 2.5 per cent. In 1927, the infestation in the treated plot decreased to 2.2 per cent, while the average for the check plots rose to 10.8 per cent. In 1928, the treated plot again had 2.2 per cent and the check plot average dropped to 4.1 per cent. In the other plot results were somewhat similar. The treated plot had 7.0 per cent infestation the year before the experiment, and the average for the four check plots was 3.3 per cent. No weeviling occurred in the treated plot in 1927, but, as in the previous

series of plots, the average infestation in the checks rose to 15.3 per cent. In 1928, the treated plot had 12.1 per cent infestation while that in the check plots was 8.1 per cent. These figures show a decrease in the treated plots of both series the year the leaders were banded (in one case to zero), and a distinct increase in the check plots. In 1928, the effectiveness of the bands was lost, as practically every band had been partially or wholly removed. One treated plot had the same percentage of infestation and the other had a very noticeable increase, while in both series of checks the infestation had decreased.

The first series of plots at Petersham consisted of a central plot and four check plots, each one-tenth acre in size. The amount of weeviling the year before the experiment was not noted. In the treated plot the infestation in 1927 was 3.5 per cent, while the average for the checks was 17.5 per cent. In 1928, the infestation in the plot treated the season before rose to 18.6 per cent, and dropped to 14.6 per cent for the check plots. In the other series, seventy trees were taken in each plot. In 1927, the banded trees showed an infestation of 13.1 per cent, and that in the check trees was 55.1 per cent. In 1928, the infestation in the treated area was 18.8 per cent and was 21.0 per cent in the untreated area.

It can be seen that the results obtained at Petersham were very similar to those obtained at Rainbow, in that there was a distinct decrease the year the bands were applied and the following year the infestation increased again in the treated plots, even though it decreased in the check areas. The disadvantages found in the wool bands are that the wool is removed by birds for nesting purposes, and it is almost imperative that it be tied to the leader to prevent the wind blowing it away. It was also found that it offers an excellent place for the deposition of eggs of the pine bark louse (*Adelges pinicorticis* Fitch) as each leader, around which a band remained over the winter, had a considerable number of the egg masses between the band and the bark. The cost of application, in this method, was about the same as with the tanglefoot.

In both methods the taller the trees, the greater was the cost. If the material is not properly applied, it will not give protection—in other words the entire surface round each leader must be covered for at least an inch, and, in the case of the sticky banding material, it must be inspected periodically to see that it is retaining its sticky quality.

This method is practical for ornamental and shade trees, but the raw wool in particular gives the leader an unsightly appearance. In extensive forest plantations, the cost of application will make this method prohibitive.

Direct Control of the Insect

As Graham (1926) points out "The direct methods of control are in their very nature rather expensive and are therefore limited in application to trees that have some value in addition to the wood produced. . . . Direct control is applicable to shade and ornamental trees, to windbreaks along highways, on watersheds, and in some private forests where the aesthetic value is an important consideration." In young plantations it would appear from observations and data collected that direct measures are feasible if the work is begun before weeviling becomes prevalent. The cost of such an operation over a period of years must be considered and it is doubtful if this increased financial burden, necessary to obtain minimum injury, would be considered by the average timberland owner, even though the operation, carried over year by year, in a lightly infested stand might not cost more than twenty-five cents per acre per year.

Three direct control methods have been tried, and all are, within the limits mentioned in the detailed account of each, effective. These methods are: (1) picking the beetles by hand, (2) jarring the insects into a net, and (3) removal of the infested leaders.

Picking the Beetles by Hand

One instance of picking the beetles from the leaders has been reported to the writer. Many of the trees in a plantation of about five acres on the Choate Estate at Petersham were being attacked severely in the spring of 1923. These trees were about six to eight feet high, and the leaders could be reached quite easily. The caretaker and one other man spent a considerable time going from tree to tree in each row and picking off the weevils which they saw. Their work was very carefully done and the attack was noticeably checked for two years. The cost was not considered for this operation, but it is estimated, on a basis of 50 cents per man hour to have been about \$2.50 per acre. This method is limited to trees within reach of a man's hand, and, as it must be very carefully done in order that the beetles will not drop off, is not practical economically.

Jarring the Insect into a Net

The collections of the adults from the leaders during the period preceding egg-laying in the spring has often been advanced as a means of control. Walden (1914) and Felt (1916) have reported that a considerable reduction of weevils in plantations is possible. In Connecticut, Walden found that six collections, about a week apart, cost between \$1.50 and \$2.00 per acre. Felt gives the cost in an operation in New York as \$1.28 per acre. Both these cases, however, were at a period when the labor cost

was not as high as at present. The feeding period for an individual female is very short—not over a week—and great care must be exercised to get all the weevils. Several collections must necessarily be made in each stand or plantation, as the weevils do not emerge from hibernation simultaneously. There is also the probability of immigration from outside infested areas when flying is generally possible. As the adults feed upon and take refuge among the buds during unseasonable weather, these must be examined and any weevils found removed. Jarring the leader on one side with a fairly heavy stick so as to knock the weevils off into a net held on the opposite side is the only practicable way to collect those on the leader itself. However, when the weevils are in the bud clusters, they cannot be dislodged in this manner, but must be picked out. This method is practical with ornamentals and possibly with young plantations isolated from native pine which is infested, or of value aesthetically. The cost in an extensive plantation or over a wide area, however, is prohibitive. In addition to this, once the trees get above seven or eight feet in height, the leaders are too high to be reached without the aid of a step-ladder or box.

Removal of the Infested Leaders

The removal of the infested leaders after they have wilted has been recommended for many years as an excellent means of control. In areas where there is natural pine in the surrounding country, from which the treated areas could be reinfested, this is financially impractical in either forest plantations or natural stands. Where there is a scarcity of pine nearby or where the plantations have aesthetic value, removal of the leaders is practical only so long as they can be reached by hand. It will be necessary to go over the stand at least twice each season; the second time to remove those leaders which were missed the first time as well as those which wilted after the first operation.

The leaders as they are removed must either be burned or placed in tight receptacles. Burning is the one certain means of killing all the larvae and pupae of the weevil, but if the leaders are placed in tight receptacles, screened on at least one side, and left in the plantation, the smaller parasites will not be killed. These receptacles should be so placed that water will not collect in them and drown the parasites. They should be left in the plantation until the following spring. By this time all the weevils will have died, the screen can be removed and those parasites which have hibernated in the tunnels of the weeviled leaders will be able to escape into the plantation. The use of number twelve or fourteen mesh screening is advisable, as these meshes will allow the escape of practically all the parasites, some of which

are too large to escape through the number sixteen mesh, which is the size which has heretofore been used.

This method is not practical in forest plantations, where there is a liability of reinfestation from surrounding areas, as the cost of removal often exceeds a dollar per acre per year. In order to have any appreciable effect, it must be carried on over a period of years. Below eight feet, the cost should not be more than twenty-five cents per acre per year to be of economic value. Above this height, the cost will be greater, depending on the height of the trees and the amount of the infestation. The total cost of the operation up to the time the trees are thirty feet high would amount to a considerable sum at compound interest by the end of even a short rotation; but the increasing value of the stand may, in some cases, warrant the expenditure.

Results obtained by the removal of infested leaders from a number of plantations in the Petersham area for two or three years have yielded some interesting data. In five heavily infested areas, treated for three years, two showed an average decrease in number of weeviling of 38.9 per cent the second year and 50.7 per cent the third year. Check plots in plantations adjacent to the treated ones showing decreases, showed an average infestation the first year the leaders were removed of 27.4 per cent. The second year this rose to 44.2 per cent and the third year it was 38.0 per cent. The infestation, therefore, was greater by 61.3 per cent the second year and by 38.6 per cent the third year, while it was decreasing in the treated plots. In the remaining three there was an average increase of 57.2 per cent the second year and of 41.7 per cent the third year. The average decrease for all five plantations for the two years was 17.9 per cent and 30.5 per cent respectively.

In stands which are isolated and which have a small amount of infestation, treating the leaders is practical. The cost should not be excessive in such isolated areas if carried on from year to year from the time the trees are liable to weeviling. If not begun until several years after the weevils first get into the stand, as happened in the first five plantations treated, the weevils become established and the cost will be much greater than if begun the first year. In South Carolina and in parts of Ontario and Quebec infestations were completely controlled by the removal of the wilted leaders before the adults emerged from them. At Rockwell Springs Experiment Station, near Syracuse, New York, the infestation in a plantation spaced 3' x 3' has been kept down to a total of less than two per cent over a period of ten years through the annual removal of the infested leaders.

The importance of the second inspection each year is borne out by the following incident. On the Swann State Forest in

Western Massachusetts, the removal of leaders has been practiced extensively for a period of years. Over an area of eighty-five acres of young white pine, the infestation, the second year after the removal operation began, was only 1 per cent. The cost of removal of the weeviled leaders for the third year was only eleven cents per acre. However, a number of the leaders were missed the next year and the cost had risen to twenty-six cents per acre. No check areas were available for comparison.

The amount of crook which will necessarily occur can be largely overcome by an additional operation which can be performed at the time the leaders are removed, with very little additional labor. The bud clusters on all but one lateral should be removed by pinching or clipping. This seems to result in the remaining lateral absorbing all the available nutrient materials which would have gone to the other laterals as well as the amount it would normally have used itself, thus giving an extra stimulus for growth. Removing the bud clusters does not impair the growth efficiency as much as if the entire length of each lateral was removed. In other words, the amount of leaf surface is not reduced to the extent that it would be if each entire lateral were removed. The laterals so treated do not grow at all in length and not appreciably in diameter; and there is no wound such as would occur if they were removed at their bases. The effect on the future quality of the timber will also be better in those trees where the laterals are left—there will be small knots, but they will probably be tight.

Repair of the Injury by Silvicultural Methods

The information gained in studying the sample plots is in accord with that secured from previous studies by Hopkins (1907), Blackman (1916, 1919), Fisher and Terry (1920), Pierson (1922), and Graham (1926), in indicating clearly that under certain forest conditions the injury, as well as the severity of attack, may be substantially decreased. Permanent injury is found most frequently in stands which are below par in site and density. It can not be expected that a stand will be free from insect or other injury unless it is maintained at a reasonable standard of density and vigor. Repair of the injury will be considered from two angles—(1) density of stocking in pure stands, and (2) production of mixed stands.

Density of Stocking

Under optimum site conditions, density of stocking will tend to keep permanent injury by the weevil at a minimum. Such a stand, however, is financially practical only when it can be procured naturally, and at a high enough density to offset any weeviling

which may occur (Plate XI-1) Pierson (1922) and Graham (1926) state that this minimum amount of damage can be expected at the end of the rotation, if a fully-stocked stand of from 1,200 to 1,500 trees per acre is maintained throughout the early years and, thereafter, a fully-stocked stand according to age class requirements. The writer's observations show that this will hold only where there is not a very severe infestation. Where there is an average infestation every year of nearly 50 per cent, as is often the case in the "weevil country" of central Massachusetts, it would appear that a greater number of trees per acre is necessary, even when the trees are fifteen years old in plantations of those densities. In the main, however, if this density is not maintained, it will be below the standard necessary to produce a good crop.

Throughout the range where it can reproduce itself in pure stands on abandoned lands, there are many thousands of acres of old field pine which are in good condition. In these stands the site conditions and stockings are such that the trees are vigorous. Sparsely-spaced stands, such as those which often come in on medium soil in abandoned pastures, will generally be weeviled quite heavily. In a closely-spaced stand only a relatively small number of trees will be in the dominant or codominant height classes at fifteen years of age; and, after a stand has reached a height of thirty feet, it will be necessary to remove the dead and suppressed trees. This will improve the condition of the stand and allow the remaining trees to develop more rapidly.

The denser the stand, the greater will be the tendency to produce relatively straight stems when weeviling does occur in such stands. In Graphs VI and VII, it will also be seen that in pure stands, both natural and artificial, growth conditions being similar, the greater the number of trees per acre the less will be the percentage of weeviling. Graph VI shows the cumulative increase in number of trees weeviled from year to year in four plots in adjacent pure plantations where the spacings are 6' x 6', 5' x 5', 4' x 4', and 3' x 3'. The same number of trees were examined in all four plots and the site conditions are practically equal. In addition to the marked decrease in weeviling, the 3' x 3' plantation has only 12.1 per cent of the weeviled trees attacked more than once; while the 4' x 4' has 27.7 per cent; the 5' x 5' has 30.1 per cent; and the 6' x 6' has 40.9 per cent. The material for this graph was obtained from an acre plantation on the Harvard Forest. The plantation is divided into blocks of one-quarter acre each, with a different density, as previously stated, in each.

Graph VII shows the amount of weeviling in white pine in pure stands and mixed stands of the same age, and under similar site conditions. The white pine was planted in densities of 8' x 8',

6' x 6', and 4' x 4' under gray birch, aspen and fire cherry in 1916. Half of each area was released in 1922, so for six years there has been a condition approximating a pure stand while in the remainder the mixed condition has persisted. Each plot is one-quarter acre in size and all trees were taken in each one. By comparing the results as shown in the curves it is apparent that there is considerably less weeviling in the mixed stand than in the pure stand.

The possibility of controlling the weevil injury by dense plantings is impractical from a financial standpoint. The cost of planting in closer spacing than 6' x 6' makes plantations denser than that prohibitive. At the present time, in the majority of localities, a spacing which will give more than 1,500 trees per acre is not considered in planning for a future forest, unless it can be secured naturally. However, it appears that on light sandy soil, where the ultimate weevil injury is usually not so severe as on the old field, medium soils, a densely stocked stand for ornamental or aesthetic purposes would be practical.

There are disadvantages with which the forest owner will have to contend if a pure stand is to be produced. When trees in the intermediate and suppressed crown classes are weeviled they are generally left behind and will eventually die. Lack of natural pruning in a pure stand is also a disadvantage. The lower branches will die in such a stand but they will persist for years instead of falling off. The knots in the lumber produced will be small, but there will be very little clear lumber at the end of a sixty year rotation. While these disadvantages may not be considered as important as the high cost of the original planting they are factors which must be considered if high grade lumber is expected at the end of the rotation.

Mixed Stands

The plots laid out in the different parts of the Northeast have yielded abundant proof that the most advantageous way to protect white pine from the weevil is to grow it in mixture, preferably with species that will be of value in the final crop. These species may be either coniferous or broadleaved trees, the latter in any case depending almost entirely on the economic conditions in that particular locality. There will not be so many white pines in a mature mixed stand as in a pure white pine stand, but a larger percentage of those surviving will be clean-boled and straight. In a region heavily infested with weevils it is very probable that comparatively little of the pure pasture pine or the trees in widely-spaced pure plantations will be straight and free enough of limbs to produce high grade timber.

The maximum amount of protection, and subsequently the maximum amount of good lumber, will be found where there is a

groupwise mixture of white pine and the better hardwoods (Plate XI-2). In a stemwise mixture, there is danger of whipping, which, in its damage to the leading shoots, may be almost or quite as injurious as the white pine weevil. Previous investigators have suggested that the reason for the protection afforded by these mixed stands is the shading of the species liable to attack and the mechanical barrier against flight caused by the accompanying species. The leaves of the broadleaved species are still on the trees when the fall migration takes place and this barrier action is one of the reasons why the species liable to attack are not infested to any extent even though pure white pine stands nearby may be severely injured. The odor from the smaller number of pine trees in a mixed stand will not attract as many weevils as will the odor from a greater number of trees in a pure stand concentrated in an area of the same size.

The preference, shown by the weevils, for feeding and ovipositing in leaders from pure open stands is also a factor which must be considered, as we have already seen. There is also evidence to show that in many cases the white pine leaders, although attacked, may recover, and very few weevils, if any, develop in those which are killed. This is undoubtedly due to the fact that there are usually fewer eggs laid in such leaders, and the small number of larvae are more likely to be smothered by the resin. If accelerated height growth has begun this is especially likely to occur.

Weedings in the early stages, if the mixture be with hardwoods, and thinnings as the stand becomes older, will be necessary so that the white pine will not be crowded out. By judicious weeding the groups of the species liable to attack, such as the white pine, can be kept fairly well segregated and the final stand will contain groups of much better quality than can be grown in the pure stands. Aside from the protection offered, an advantage in growing white pine in mixture with other species is that it cleans itself relatively early in life, and the boles will be clear for some distance.

As stated in the Summarization of Plot Data, an alternate row mixture of the white pine and the Scotch pine has given maximum protection against weevil attack and also favors the repair of the injury when attack does occur.

A natural mixture of the Canada hemlock and the white pine gives good protection to the latter species if the hemlocks are several years older and therefore several feet high at the time the white pines begin to grow (Plate XII).

In fact this protection can be obtained in any coniferous mixture, whether it be an artificial planting in alternate rows or a natural regeneration, groupwise or stemwise in composition. The accompanying species, however, must be immune from

weeviling and faster growing in the earlier years, if both are the same age, or if slower growing it must be considerably older, at the time the white pines become liable to attack, so that it will be taller than the latter. There will be little danger of whipping in such stands if the leaders of the white pines are high enough to be above the spreading branches in the lower part of the crowns and still be at least two or three feet below the leaders of the accompanying species. The disadvantage met with in the regeneration of a coniferous mixture is that there will be no protection offered if the white pines are as tall as the other species.

Application of Control Measures to All Species Liable to Attack

Protection of the white pine has been emphasized in these control measures because this species is the favored host of the white pine weevil and is the most widely planted or naturally regenerated species, in the range of this insect, of those which are susceptible to attack. Norway spruce, which is damaged to almost as great an extent as the white pine (Plate XIII-2), is grown in large plantations, and the injury can be minimized by the same measures. All the other species which are attacked by the White Pine Weevil can be protected against infestation. Those measures given for ornamental and shade trees are particularly applicable for the exotic species which are planted for these purposes.

SUMMARY

The White Pine Weevil (*Pissodes strobi* Peck) is the most serious insect pest attacking the white pine (*Pinus strobus* L.). The leading shoot is killed, resulting in loss in height, loss in radial increment and often poor form. The timber value is also materially affected; lumber sawn from weeviled trees will usually show large knots at the points of weeviling, and very often decay will be present. Short lengths will often be necessary because of the many crooks in the stems due to weevil injury.

The adult weevils hibernate mostly in the lowest layer of the litter, just at the surface of the mineral soil under trees fed on during the fall. Occasionally they hibernate at the bases of trees cut during the late summer or fall. The trees fed upon during the fall may or may not have been weeviled that season. Young adults and even full-grown larvae from eggs laid late in the season may, under very favorable conditions, hibernate in the leaders.

The weevils emerge from hibernation in the spring about the time the buds begin to swell, and activity apparently ceases with the cessation of accelerated aerial elongation, as no weevils are found from this time until the new generation emerges. This

period depends on the growing season and weather conditions. The females may be fertilized in the fall, as well as in the spring. Oviposition begins soon after activity is resumed, reaches the peak in two to three weeks and then gradually diminishes, ending with the cessation of activity of the old adults. Development from the egg to the adult covers a period of approximately three months. The new generation emerges in the late summer and feeds on the tender inner bark of the tips of branches and leading shoots, but does little damage as compared to that done in the spring. Feeding ordinarily goes on at intervals until settled cold weather prevails, and then the weevils go into hibernation.

The adult weevils, in both spring and fall, are relatively strong fliers. They can cover considerable distances when wafted by the wind, and can thus reinfest an area or infest a new area in a short time. The air temperature necessary for general flight appears to be between 70° and 85° Fahrenheit. Above and below this range, comparatively little flight has been observed. Both sexes will fly directly to the leading shoot, but the majority strike the trees lower down and crawl up the stem. In the early spring, before warm weather prevails, many of the weevils reach leading shoots by crawling up from the ground.

The degree of intensity of infestation depends primarily on the number of host trees available, weather factors (chiefly temperature), soil conditions and exposure. Adults feed and oviposit more readily on trees in pure, widely-spaced stands than on trees in dense, pure stands or in mixed stands of pine and other species, such as hemlock or hardwoods. Stands on medium agricultural (sandy loam) soils will generally show the most injury. Stands on sandy soils may have as many trees weeviled per acre, but the ultimate injury is not so severe. Plantations or stands on sunny exposures are more liable to weeviling than those on exposures protected from the morning sun.

Direct control measures are generally too expensive for common practice. In isolated stands, where the infestation is low and the danger of reinfestation is slight, or in stands which are being preserved for their aesthetic value, such control measures may be used advantageously.

Weeviling, in stands which are subject to attack from surrounding areas year after year, can sometimes be checked, but not controlled, by the removal of the infested leaders, collection of the adults during the spring feeding period, spraying with repellents or banding the leaders. The cost of these treatments, often more than a dollar per acre per year makes these measures prohibitive over wide forested areas.

The breeding and liberation of parasites and predatory insects in infested areas is an uncertain measure of control. The cost, necessary for breeding and liberating a number sufficient to have

an appreciable effect, would be too great to make such a project financially practical over a wide area at the present time.

Insectivorous birds are very valuable natural control agencies and they should be protected and encouraged in pine woodlands.

In pure stands, both natural and artificial, the greater the number of trees per acre the smaller will be the percentage of weevil-ing. A dense stand is financially practical only when it can be procured naturally at a high enough density to offset any weevil-ing which may occur. Where site conditions and the stocking are favorable for vigorous growth good stands of merchantable old field pine are common.

The most advantageous and cheapest way to protect white pine from the weevil and to control the injury is to grow it in mixture, preferably with species that will be of value in the final crop. The choice of the other species will be determined by the economic conditions in any given locality. Weedings in the early stages, if the mixture be with hardwoods, and thinnings as the stand becomes older, will be necessary so that the pines will not be crowded out. There will not be so many white pines in a mature mixed stand, as in a mature pure stand of normal stocking, but those present will be clean boled and straight, and the value of the whole crop will be greater.

BIBLIOGRAPHY

- Anon., 1921.**
White Pine Weevil and Reforestation. The Conservationist. Vol. IV, pp. 189-190.
- Anon., 1925.**
White Pine Weevil. Cons. Comm. N. Y. XV Ann. Rept., pp. 221-222.
- Anon., 1926.**
White Pine Weevil. Cons. Comm. N. Y. XVI Ann. Rept., p. 223.
- Anon., 1927.**
White Pine Weevil Control. Cons. Comm. N. Y. Ann. Rept. to the Legislature. Legis. Doc. 38, pp. 251-252.
- Bailey, H. L., 1926.**
Important Tree Insects of Vermont. Vermont Dept. Agri. Bull. 35, p. 49.
- Bazeley, W. A. L., 1921.**
Ann. Rept. of the Commissioner of Conservation and State Forester of Mass. Pub. Doc. 73, p. 41.
- Belyea, H. C., 1923.**
The Control of White Pine Weevil (*Pissodes strobi*) by Mixed Planting. Jour. For., IV, pp. 384-390.
- Belyea, H. C., and MacAloney, H. J., 1926.**
Weather Injury to Terminal Buds of Scotch Pine and Other Conifers. Jour. For., IV, pp. 685-691.
- Blackman, M. W., 1919.**
Report on the White Pine Weevil. Maine For. Dept.
- Blackman, M. W., and Ellis, W. O., 1916.**
Some Insect Enemies of Shade Trees and Ornamental Shrubs. N. Y. State College of Forestry at Syracuse University. Vol. XVI, No. 26, pp. 72-78.
- Blatchley, W. S., and Leng, C. W., 1916.**
Rhyncophora or Weevils of Northeastern America. pp. 178-180.
- Britton, W. E., 1914.**
Reference to *Rhogas terminalis* and *Muscina stabulans* Fall., in "Outbreak of Army Worms." XIV Rept. of the State Entomologist. Conn. Agri. Expt. Sta., pp. 132-133.
- Britton, W. E., 1916.**
The Pine Tip Moth. XVI Rept. of the State Entomologist. Conn. Agri. Expt. Sta., p. 122.
- Britton, W. E., 1919.**
XIX Rept. of the State Entomologist. Conn. Agri. Expt. Sta. Bull. 218, pp. 144-155.
- Britton, W. E., and Walden, B. H., 1911.**
XI Rept. of the State Entomologist. Conn. Agri. Expt. Sta. Rept. 35, pp. 307-309.
- Brues, C. T., 1903.**
Reference to Genus *Aphiochaeta* in "Monograph of the North American Phoridae." Trans. Amer. Ent. Soc., XXIX, p. 337.

Brues, C. T., 1920.

Reference to the White Pine Weevil in—Insects and Human Welfare, pp. 80–81.

Brues, C. T., 1923.

Choice of Food and Numerical Abundance among Insects. Jour. Econ. Ent. XVI No. 1, pp. 46–51.

Butterick, P. L., 1916.

Refers to the White Pine Weevil during the course of the article. Another Insect Enemy of the White Pine (Imported Pine Sawfly). Am. For., Vol. XXII, July, pp. 395–396.

Champion, G. C., 1902.

Insecta, Coleoptera, Biologia Centrali Americana, Vol. 4, part 4, p. 119.

Chittenden, F. H., 1899.

The White Pine. U. S. D. A. For. Div. Bull. 22, pp. 58–59.

Cline, A. C., and Fletcher, E. D., 1928.

References to weeviling in "Pruning for Profit." Harvard Forest and Mass. For. Assn.

Cline, A. C., and Lockard, C. R., 1925.

Mixed White Pine and Hardwood. Harvard Forest Bull. 8.

Colville, p., 1923.

A Few Hypotheses on White Pine Losses from Weevil Damage. The Ames Forester, pp. 72–80.

Cushman, R. A., 1915.

Reference to *Calliephialtes thurberiae* Cress. in "Descriptions of New Ichneumonidae and Taxonomic Notes." Proc. Ent. Soc. Wash. XVII, pp. 132–133.

Dustan, A. G., 1923.

Reference to *Rhogas hyphantriae* Gahan in "A Histological Account of Three Parasites of the Fall Webworm (*Hyphantria cunea* Drury)." Proc. Acad. Ent. Soc. No. 8, pp. 73–94.

Emmons, E., 1854.

Genus *Pissodes* (Germ.) Agriculture of New York, p. 108.

Felt, E. P., 1906.

Insects Affecting Park and Woodland Trees. Memoir 8, part 2, N. Y. State Museum, pp. 397–401.

Felt, E. P., 1913.

XXIX Rept. of the State Entomologist on Injuries and Other Insects of the State of New York. Bull. 175, pp. 30–33.

Felt, E. P., 1916.

White Pine Weevil. XXXI Rept. of the State Entomologist. N. Y. State Museum Bull. 186, p. 84.

Felt, E. P., 1924.

The White Pine Weevil, in "Manual of Tree and Shrub Insects," pp. 227–228.

Felt, E. P., 1928.

White Pine Weevil, in "Observations and Notes on Injurious and Other Insects of New York State." N. Y. State Museum Bull. 274, pp. 164–166.

Fernald, H. T., 1908.

White Pine Weevil, in 'Shade Trees,' by E. A. Start, G. E. Stone, and H. T. Fernald. Mass. Agri. Expt. Sta. Bull. 125, p. 58.

Fernald, H. T., 1921.

The White Pine Weevil (*Pissodes strobi* Peck). App. Ent., pp. 142-143.

Fisher, R. T., and Terry, E. I., 1920.

The Management of Second Growth White Pine in Central New England. Jour. For. 1V, pp. 358-367.

Fitch, A., 1858.

IV Rept. on the Noxious and Other Insects of the State of New York, pp. 732-736.

Forbush, E. N., 1913.

Useful Birds and Their Protection. Mass. State Bd. of Agri., pp. 168, 254-256.

Franklin, H. J., 1921.

Reference to species of *Phygadeuon* on the Cranberry Girdler in Report of the Cranberry Station for 1919. Mass. Expt. Sta. Bull. 206, pp. 149-168.

Frothingham, E. H., 1914.

White Pine Under Forest Management. U. S. D. A. Bull. 13, pp. 62-63.

Fuller, A. S., 1870.

White Pine Weevil. Am. Ent. Vol. II, p. 26.

Fuller, A. S., 1880.

A Good Word for the White Pine Weevil. Am. Ent. and Bot. Vol. III. (2nd Series 1), pp. 5-6.

Germar, E. F., 1817.

Magazin der Entomologie. Vol. II, p. 340.

Gevorkiantz, S. R., and Hosley, N. W., 1929.

Form and Development of White Pine Stands in Relation to Growing Space. Bull. 13. Harvard Forest, p. 44.

Gill, J. B., 1925.

Reference to species of *Calliephialtes* in "The Pecan Nut Case-bearer." U. S. D. A. Dept. Bull. 1303.

Girault, A. A., 1917.

New Chalcid Flies with Notes. Bull. Brooklyn Ent. Soc. Vol. XXI, pp. 86-89.

Gorham, R. P., 1922.

Insect Pests of the Year 1922 in New Brunswick. Proc. Acad. Ent. Soc., p. 21.

Graham, S. A., 1916.

Notes on the Control of the White Pine Weevil. Jour. Econ. Ent. Vol. IX, pp. 549-551.

Graham, S. A., 1918.

The White Pine Weevil and Its Relation to Second Growth White Pine. Jour. For. II, pp. 192-202.

Graham, S. A., 1926.

The Biology and Control of the White Pine Weevil, *Pissodes strobi* Peck. Cornell Univ. Agr. Expt. Sta. Bull. 449.

Graham, S. A., 1929.

Miscellaneous References to the White Pine Weevil, in "Principles of Forest Entomology.

Hawley, R. C., 1924.

Early Development of White and Red Pine Plantations. Jour. For. III, pp. 275-281.

Harrington, W. H., 1881.

Rhyncophora—Weevils. XI Rept. Ent. Soc. Ontario, 1880, pp. 49-57.

Harris, T. W., 1841.

A report on the Insects of Massachusetts Injurious to Vegetation, pp. 63-65.

Harris, T. W., 1862.

A Treatise on Some of the Insects Injurious to Vegetation, pp. 72-73.

Hartig, R., 1894.

Diseases of Trees. Translated from the German by William Somerville, Professor of Agriculture and Forestry, Durham College of Science, Newcastle-on-Tyne, England.

Herold, W., 1923.

Reference to *Muscina stabulans* Fall. in "Zur Kenntniss von *Euroa segetum* Schiff. (Saateule) III Feinde und Krankheiten." Zeitschr. angew. Ent. IX, No. 2, pp. 306-332. (Review Appl. Ent.)

Hicock, H. W., 1924.

Rainbow Forest Plantations. Conn. Agri. Expt. Sta. Bull. 262, For. Pub. 15.

Hill, C. C., 1925.

Reference to species of *Rhogas* in "Biological Studies of the Green Clover Worm." U. S. Dept. Agri. Dept. Bull. 1336.

Hopkins, A. D., 1893.

Catalogue of West Virginia Forest and Shade Trees Insects. W. Va. Agri. Expt. Sta. Bull. 32, p. 205.

Hopkins, A. D., 1899.

Report on Investigations to Determine the Cause of Unhealthy Conditions of the Spruce and Pine from 1880-1893. W. Va. Agri. Expt. Sta. Bull. 56, pp. 259-260, 345, 429, 441.

Hopkins, A. D., 1904.

Catalogue of Insect Enemies of Forests and Forest Products at the Louisiana Purchase Exposition, St. Louis, Mo. 1904. U. S. D. A. Div. Ent. Bull. 48. "Catalogue of exhibit," pp. 15 and ff., reprinted as part of U. S. D. A. Bur. Ent. Bull. 53. 1905.)

Hopkins, A. D., 1905.

Insect Enemies of Forest Reproduction. U. S. D. A. Year Book, 1905, pp. 249-256.

Hopkins, A. D., 1907.

The White Pine Weevil. U. S. D. A. Bur. Ent. Circ. 90.

Hopkins, A. D., 1911.

Contributions toward a Monograph of the Bark Weevils of the Genus *Pissodes*. U. S. D. A. Bur. Ent. Tech. Series 20.

Hosley, N. W., 1928.

Red Squirrel Damage to Coniferous Plantations and its Relation to Changing Food Habits. Ecology. Vol. IX. No. 1, pp. 43-49.

Houser, J. S., 1918.

Destructive Insects Affecting Ohio Shade and Forest Trees. Ohio Agri. Expt. Sta. Bull. 332, pp. 323-325.

Hunter, W. D., and Pierce, W. D., 1912.

Reference to species of *Pheidole* in "Mexican Cotton Boll Weevil." U. S. D. A. Bur. Ent. Bull. 114, p. 140.

Jaenicke, A. J., 1921.

Forest Insect Problems of the Pacific Coast. Jour. Econ. Ent. Vol. XIV, p. 449.

Klimesch, J., 1923.

Reference to *Rhopalicus suspensus* Katz on *Ips typographus* L. Wiener Allg. Forst U. Jagdzeitung XLI. Nos. 28 and 31. (Review Appl. Ent.)

LeConte, J. L., and Horn, G. H., 1876

The Rhyncophora of America North of Mexico. Proc. Amer. Phil. Soc. Vol. XV. No. 96, pp. 142-144.

Leiby, R. W., 1922.

Reference to species of *Calliephialtes* in "Biology of the Golden Rod Gall-maker." Jour. N. Y. Ent. Soc. XXX. No. 2, pp. 81-94.

Lintner, J. A., 1888.

IV Rept. on the Injurious and Other Insects of the State of New York, p. 24.

Lintner, J. A., 1892.

IX Rept. on the Injurious and Other Insects of the State of New York for the year 1892, pp. 344-345.

MacAloney, H. J., 1925.

Preliminary Observations on the Control of the White Pine Weevil Injury by Forest Management. Unpublished Thesis for the Degree of Master of Forestry at the N. Y. State College of Forestry.

MacAloney, H. J., 1926.

The White Pine Weevil in the New England States. Papers presented at the Forest Protection Conference, N. Y. State College of Forestry.

MacAloney, H. J., 1928.

Mixed Stands the Best Protection Against White Pine Weevil. Forest Worker. March. p. 13.

MacAloney, H. J., 1929.

The White Pine Weevil as a Shade-Tree Pest. Proceedings 5th National Shade-Tree Conference, Brooklyn Botanical Gardens.

McAtee, W. L., 1926.

Relation of Birds to Woodlots. Roosevelt Wild Life Bulletin. Vol. IV, No. 1, p. 115.

Muesebeck, C. F. W., 1925.

Microbracon pini—new species. A Revision of Parasitic Wasps of the Genus *Microbracon* occurring in America North of Mexico. Proc. U. S. Nat. Mus., Vol. LXVII, Art. 8, pp. 52-53.

- New Brunswick Crown Land Report for 1918, 1919.**
The White Pine Weevil, p. 114.
- Packard, A. S., 1881.**
Insects Injurious to Forest and Shade Trees. U. S. Ent. Comm. Bull. 7, pp. 185-188, 228-229, 236, 241.
- Packard, A. S., 1885.**
The White Pine Weevil and Its Injury to Shade and Forest Trees. Rept. Comm. Agri., pp. 322-325.
- Packard, A. S., 1890.**
The White Pine Weevil. U. S. Ent. Comm. Rept. No. 5, pp. 734-741, 829-830, 861, 872.
- Peck, W. D., 1817.**
On the Insects which Destroy the Young Branches of the Pear Tree and the Leading Shoot of the Weymouth-pine. Mass Agri. Repository and Journal, Vol. IV, No. 1, pp. 205-211.
- Peirson, H. B., 1922.**
Control of the White Pine Weevil by Forest Management. Harvard Forest Bull. 5.
- Peirson, H. B., 1823, 1923**
White Pine Weevil. Insects Attacking Forest and Shade Trees. Maine For. Ser. Bull. 1, pp. 35-37.
- Pierce, W. D., 1907.**
On the Biologies of the Rhyncophora of North America. Nebraska State Bd. of Agri. Ann. Rept. 1906-1907, p. 259.
- Pierce, W. D., 1908.**
A list of Parasites Known to Attack American Rhyncophora. Jour. Econ. Ent. Vol. 1, pp. 389-396.
- Porter, B. A., and Alden, C. H., 1924.**
Reference to species of *Rhogas* on Cankerworm in "The Cankerworms." U. S. D. A. Dept. Bull. 1238.
- Pyle, E. C., 1927.**
Weevil Attack in White Pine Underplantings. Extract from Penn. Dept. Forests and Waters "Service Letter," Dec. 29.
- Rane, F. W., 1915.**
The White Pine Weevil. Rept. Mass. State Forester Pub. Doc. 73, p. 53.
- Rane, F. W., 1917.**
Recent Troubles with the Forest Trees. Trans. Mass. Hort. Soc., p. 68.
- Report of the Entomologist, 1926.**
U. S. D. A. Bur. Ent. Bull., p. 27.
- Riley, C. V., and Howard, L. O., 1890.**
Some of the Bred Parasitic Hymenoptera in the National Museum Collection. Ins. Life, Vol. II, p. 348.
- Riley, C. V., and Howard, L. O., 1891.**
The White Pine Weevil. Ins. Life, Vol III, p. 468.

Say, T., 1831.

Descriptions of North American Curculionides, and an Arrangement of Some of our own Species Agreeably to the Method of Schonherr. New Harmony, Indiana. The Complete Writings of Thomas Say on the Entomology of North America. *Pissodes (Rhynchaenus) strobi*, Peck.

Saunders, W., 1883.

Insects Injurious to the White Pine—*Pinus strobus*. Ent. Soc. Ont., p. 55.

Simmons, J. R., 1916.

Three Common Dangers to White Pine. New England Homestead, Vol. LXXII, No. 2, p. 5.

Smith, R. C., 1921.

Reference to *Muscina stabulans* Fall in "Observations on the Fall Army Worm and Some Control Experiments." Jour. Econ. Ent. XIV, No. 3, pp. 300-305.

Spaeth, J. N., 1922.

Notes on Release of White Pine on Harvard Forest. Jour. For. Vol. XX, No. 2, pp. 117-122.

Spring, S. N., 1905.

The Natural Replacement of White Pine on Old Fields in New England. U. S. D. A. Bur. of For. Bull. 63, pp. 14-15.

Stearns, L. A., 1919.

Reference to *Rhogas* species in "Some Recently Recorded Parasites of the Oriental Peach Moth." Jour. Econ. Ent. XII, No. 4, pp. 347-348.

Swaine, J. M., 1925.

The Destruction of Canadian Forests by Insects. Pulp and Paper Magazine, June 25, p. 725.

Swaine, J. M., 1928.

Forest Entomology and its Development in Canada. Dom. Can. Dept. Agri. Pamphlet 97, pp. 4, 16, 19.

Tarbox, E. E., 1924.

Quality and Growth of White Pine as Influenced by Density, Site, and Associated Species. Harvard Forest Bull. 7.

Taylor, R. L., 1927.

A New Species of Parasitic *Hymenoptera* (*Chalcidoidea*, *Eupelmidea*). Brooklyn Ent. Soc. Vol. XXII, No. 4, pp. 205-207.

Taylor, R. L., 1928.

A New Species of *Lonchaea* Fallen (*Lonchacidae* *Diptera*). Bull. Brooklyn Ent. Soc. October, p. 191.

Taylor, R. L., 1929.

The Arthropod Fauna of Coniferous Leaders Weeviled by *Pissodes strobi* (Peck). Contribution from the Entomological Laboratory of the Bussey Institution, Harvard University, No. 301 Psyche, Vol. XXXV, No. 4, pp. 217-225.

Thomas, C., 1874.

VI Rept. of the State Entomologist on the Noxious and Beneficial Insects of the State of Illinois, pp. 133-134.

Vierick, H. L., et al., 1916.

The Hymenoptera, or Wasp-like insects of Connecticut. State Geol. and Nat. Hist. Survey of Conn. Bull. 22, pp. 210, 765.

Walden, B. H., 1914.

XIV Rept. State Entomologist. Conn. Agri. Expt. Sta. Bull., pp. 173-176.

Walden, B. H., 1914.

Experiments in Controlling the White Pine Weevil in 1915. XV Rept. State Entomologist. Conn. Agri. Expt. Sta. Bull., pp. 134-136.

Walker, E. M., 1911.

Some Injurious Insects at De Grassi Point, Lake Simcoe. Rept. Ent. Soc. Ont., p. 58.

Webber, R. T., and Schaffner, J. V. Jr., 1926.

Host Relations of *Compsilura concinnata* Meigen, an Important Tachinid Parasite of the Gipsy Moth and Brown-Tail Moth. U. S. D. A. Bull. 1363, p. 27.

Weiss, H. B., 1919.

White Pine Weevil (*Pissodes strobi* Peck). State Bd. of Agri., State of New Jersey, Cir. 5.

Wells, A. B., 1926.

Notes on *Hylobius pales* Herbst and *Pissodes strobi* Peck as Nursery Pests. Jour. Econ. Ent. Vol. XIX, No. 2, pp. 412-413.

Wheeler, W. M., 1913.

Reference to Pheidole species in "Ants—Their Structure, Development and Behavior," p. 268.

EXPLANATION OF GRAPHS

Graph I. To show the Correlation between the Periods of Development of the White Pine Weevil and Definite Periods in the Annual Height Growth of the Northern White Pine.

Graph II. To show the Approximate Loss in Height Growth of Trees killed back two years, each time a Tree is Weeviled. Based on 60 Weeviled Trees and 60 Non-weeviled Trees.

Graph III. To show the Contrast between the Average Number of Weevils per Tree Weeviled in the Different Height Classes in White Pine from Stands of low density pure old field pine and from Stands of mixed white pine and hardwoods. The broken lines indicate trees from mixed stands.

Graph IV. The Average Annual Heights of Scotch Pine, weeviled and non-weeviled White Pine, in alternate row even-aged mixture. Explanation in text.

Graph V. To show the Average Number of Weevilings per Tree below 25 and 35 Feet in all Height Classes. Pure old field pine, low density, all aged. Explanation in Text.

Graph VI. Comparison of the Cumulative Increase in Percentage of Trees Weeviled in Pure stands of Different Densities. Based on 125 Trees from each plot.

Graph VII. Comparison of the Percentage of Trees Weeviled in Stands of Different Densities. Explanation in Text.

A—White Pine from Pure Stand.

B—White Pine from Stemwise Mixture.

EXPLANATION OF PLATES

PLATE I

Type of Cage used in Experiments to Determine the Average Number of Eggs laid by each Female.

PLATE I

PLATE II

1. Showing the Positions the Pupae may assume while in their Pupal Chambers.
- 2 and 3. Showing the Position and Character of the Pupal Chambers in the Pith and in the Wood.
4. The Dried Pitch Masses on a Weeviled Leader.

PLATE II

PLATE III

A Weeviled Leader.

PLATE III

PLATE IV

A Plantation showing the Bayonnetted Effect on Weeviled Trees. (Courtesy Harvard Forest.)

PLATE IV

PLATE V

1. A Badly Forked Tree.
2. Staghorn Pine, showing several Weevilings in each Tree and the Resulting Staghorn Effect. (Courtesy Harvard Forest.)
3. A Weeviled Tip which has persisted for Many Years in a Log which was 12 Inches in Diameter at the Weeviled Point at Time of Cutting. Exit Holes are Plainly Evident.

PLATE V

PLATE VI

1. An Uninjured Tree. (Courtesy G. H. Lentz.)
2. A Typical Cabbage Pine. (Courtesy Harvard Forest.)

PLATE VI

PLATE VII

- 1 and 2. Boards with Knots to the Center. The Result of Weeviling.
3. Board showing Clear Portion on One Side; Knots on Opposite Side are due to Weeviling.
4. Board showing Large Knot and Persisting Leader after Weeviling.

PLATE VII

PLATE VIII

1. Showing Knots on One Side of Board due to Weeviling; Clear Portion on Opposite Side.

2 and 3. Showing Decay following Weeviling.

PLATE VIII

1

2

3

PLATE IX

Injuries Often Mistaken for Weeviling.

1. A Dead Leader in Scotch Pine caused by Weather Injury. The Absence of Exit Holes Distinguishes this Injury from Weeviling. (Courtesy G. A. Whipple.)
2. White Pine Trees overtopped and whipped by Gray Birch.
3. Squirrel Damage. A Crook or Fork results from this Injury and may often be mistaken for Weeviling. (Courtesy Harvard Forest.)

PLATE IX

PLATE X

- 1 and 2. Larvae of *Louhaca corticis* T. in the Larval Tunnels and Pupal Chambers of *Pissodes strobi* Peck. (See also Plate II-1.)
3. A Lepidopterous Larva in the Pupal Chamber of *Pissodes strobi* Peck.
- 4 and 5. Showing the Bark Stripped from Weeviled Leaders by Birds in order that the Larvae and Pupae may be eaten.

PLATE X

PLATE XI

1. Showing Density of Tree in a Pure Stand, Resulting in Minimum Injury. (Courtesy Harvard Forest.)
2. Mixed White Pine and Hemlock showing Straight Pines without Weevil Injury. (Courtesy Harvard Forest.)

PLATE XI

2

1

PLATE XIII

1. A Paint Brush Top in Scotch Pine caused by Weather Injury. (Courtesy G. A. Whipple.)
2. Weeviling in Norway Spruce.

PLATE XIII

2

1

Biology and Control of the White-Pine Weevil, *Pissodes strobi* Peck

Samuel A. Graham

AN ADULT WEEVIL ON A BUD CLUSTER OF WHITE PINE

The feeding punctures show on the buds

Published by the
Cornell University
Agricultural Experiment Station
Ithaca, New York

Received for publication January 11, 1926

CONTENTS

	PAGE
Injurious habits.....	4
Character of the injury.....	4
Host trees.....	7
Relation of injury to age and height of tree.....	7
Relation of injury to crown class and rate of growth.....	8
Life history and habits.....	10
Activities of the old adults.....	10
Developmental stages.....	13
Activities of the young adults.....	15
Control.....	17
Direct control.....	17
Chemical methods.....	18
Mechanical methods.....	18
Indirect control.....	21
Silvicultural methods.....	21
Biological methods.....	26
Summary.....	28
Bibliography.....	30

BIOLOGY AND CONTROL OF THE WHITE-PINE WEEVIL, PISSODES STROBI PECK

SAMUEL A. GRAHAM

The white-pine weevil is one of the most destructive pests of the eastern white pine. In some localities the planting of this tree has been practically abandoned because of its ravages, and throughout the range of its host tree it has been a large factor in discouraging private reforestation of waste lands. It is therefore an important forest insect, not only because of the actual damage that it does, but also because of the bad psychological effect which its presence produces upon owners of forest properties where white pine is an important tree.

This weevil was early brought to the attention of entomologists, and evidently it has always been an important forest insect throughout the range of the eastern white pine. The earliest observers of the insect found that to a certain degree it was controlled by several parasites. Peck (1817) states that "our forests would scarce produce a single mast" were it not for this controlling factor. Today, although the demand for masts is a small item in the white-pine trade, the merchantable value of the timber still depends on the straightness of the main stem, and any crook or deformity such as that caused by the weevil reduces the value of the timber.

Peck described this weevil in 1817 and made a few notes on its life history. Since that time it has been studied by a number of entomologists, each of whom has contributed more or less to the knowledge of the species. Unfortunately, the white-pine weevil has been confused with some of its close relatives. This has led to many misleading, and even incorrect, statements concerning its life history and habits. One of the purposes, therefore, of the experiments upon which this paper is based, was to determine the true life cycle and habits of the insect as distinguished from those of its close relatives. The second and more important purpose was to find, if possible, some practical methods of controlling the weevil.

The study was begun in 1915 as a thesis problem under the direction of Professor Glenn W. Herrick, of Cornell University, and the greater part of the biological data included in this bulletin were collected at Ithaca during the summer of 1915. During 1916, most of these observations were checked and further work on the control problem was conducted at Ithaca and also on the Luther Forest Preserve near Saratoga Springs, New York, for the Department of Forestry at Cornell University. In 1917 and 1918, further studies were carried on in Isanti County and other localities in Minnesota, for the Minnesota Agricultural Experiment Station. Some of the results of the Minnesota studies have appeared in an article in the *Journal of Forestry* (Graham, 1918). Thus there has been opportunity to check results in several widely separated localities. In this bulletin an attempt is made to collect all of these data into a comprehensive whole.

INJURIOUS HABITS

CHARACTER OF THE INJURY

The injury done by the white-pine weevil is confined to the terminal part of sapling pines. The lateral branches are seldom injured. From a point just below the terminal bud cluster, the larvae work downward in the inner bark and girdle the shoot as they progress. The injury

FIGURE 1. LEADING SHOOT KILLED BY WHITE-PINE WEEVILS

usually extends to a point below the topmost whorl of branches only, but many instances have been observed in which the tree has been killed below the second, or even the third, whorl from the top (figure 1). When the leader is destroyed in this way, the laterals in the topmost living whorl of branches compete for supremacy. In some instances one lateral succeeds in overtopping the others and takes the lead, and in such a case the tree either outgrows the injury or a bayonet trunk results. Sometimes, however, two or more laterals may take the lead, and produce a forked tree. In some cases weevil injury may render the trees commercially valueless.

Even where a single lateral takes the place of the killed leader and the tree outgrows completely the weevil injury, there is always a loss in height growth. If the leader is not killed below the first whorl, this loss is only the difference between the length of the terminal and the length of the lateral that takes its place. This difference varies in proportion

FIGURE 2. RELATION BETWEEN THE LENGTH OF THE LEADING SHOOT (TERMINAL), AND THE AVERAGE EXCESS OF THE LEADER OVER THE CORRESPONDING LATERAL

to the rate of height growth. The relation between the length of the leading shoot and the average excess of the length of the leader over that of the corresponding lateral, is shown in figure 2. If, however, the weevil girdles the tree below the topmost whorl, the loss in height is equal to the difference between the length of the main stem above the topmost living whorl and the length of the lateral that assumes the lead. The average loss per tree, in feet, at different densities of the stand, is shown

in figure 3. The number of trees per acre as indicated on this graph, is the original number before any thinning or crowding-out has taken place.

FIGURE 3. AVERAGE LOSS IN HEIGHT GROWTH PER TREE, IN FEET, RESULTING FROM WEEVIL ATTACK AT DIFFERENT DENSITIES

It has been estimated that from 70 to 90 per cent of all the white pines in New York and New England have been weeviled before they have reached the age of fifteen years. In Minnesota this percentage is somewhat lower. Unless steps are taken either to reduce this injury to the minimum or to assist the trees in outgrowing the resultant deformity, a crop of crooked, forked trees, of very little commercial value, will result.

In addition to the direct damage caused by weevil attack, there are other secondary injuries that are of considerable importance. One of these is snow breakage. The irregular, dense growth of weeviled trees holds the snow to a much greater degree than does the normal straight pine. An extreme example of combined weevil and snow injury was observed not far from the Cornell University campus, in a group of trees located in a hollow where heavy snows collect. These trees were broken beyond any hope of recovery, and were little more than sprawling shrubs, barely able to keep alive.

HOST TREES

The eastern white pine (*Pinus strobus*) is the favorite host of the white-pine weevil. However, the insect sometimes attacks other species of pine and some species of spruce. Scotch pine (*Pinus sylvestris*), pitch pine (*Pinus rigida*), and jack pine (*Pinus Banksiana*) are frequently attacked when growing in mixture with, or near, young white pines. Red pine (*Pinus resinosa*) is much more resistant but is occasionally injured. Norway spruce (*Picea excelsa*) is very susceptible and is second to white pine as a favored host of this insect. Other species that have been reported as somewhat susceptible are Japanese pine (*Pinus densiflora*) (Peirson, 1922), Himalayan cedar (*Cedrus deodara*) (Currie, 1905), red spruce (*Picea rubra*) (Hopkins, 1911), balsam fir (*Abies balsamea*) (Packard, 1890), and eastern hemlock (*Tsuga canadensis*) (Packard, 1890). It is possible that some of these earlier records may have resulted from the confusion of other species of *Pissodes* with the white-pine weevil. Packard, in his writings, evidently confused several species of *Pissodes*, and, as his records of hemlock and balsam fir as host trees of the white-pine weevil have not been verified by recent workers, it is possible that these two species should not be included as host trees of this insect.

Since white pine is not only the favorite host of the white-pine weevil but is also the most important of the susceptible species as a timber tree, the experiments in both New York and Minnesota have been confined almost entirely to the activities of the weevil on this tree.

RELATION OF INJURY TO AGE AND HEIGHT OF TREE

The first attack of the white-pine weevil usually occurs when the trees are from five to seven years old and from two to three feet in height. The infestation is generally rather light during the first year or so, but it steadily increases, reaching the maximum when the trees are from twelve to eighteen years old. From then on, the infestation declines rapidly in intensity, and it practically ceases to be economically important when the trees are from twenty-five to thirty years old.

The rate of infestation appears to be correlated with the rate of growth of the trees, as is shown later in this bulletin; but there seems to be also some connection between the height of the trees and weevil injury (figure 4). The weevil ordinarily ceases to work on the trees after they have reached a height of from twenty to thirty feet. No entirely satisfactory explanation of this has been offered. It has often been assumed that insects which work only on low trees are habitually low fliers, and that when a tree attains a certain height it is beyond their reach. This may be the true explanation; but it is equally probable that the changing eco-

FIGURE 4. INCREASE OF THE INTENSITY OF WEEVIL INFESTATION WITH HEIGHT GROWTH, FROM 1910 TO 1915

This graph was constructed from data collected on a series of sapling plots on the Luther Forest Preserve. In 1910 the trees on these plots averaged seven years of age, two years after weeviling began

logical conditions which take place conjointly with the development of the stand, may offer a more adequate explanation.

In this study it was impossible to follow through, from year to year, the development of the young white-pine stands that were examined. At the conclusion of the investigation, the trees on the plots had not yet passed beyond the period of greatest susceptibility, and therefore the curve represented in figure 4 is incomplete. This figure does, however, show that the rate of infestation increased rapidly during the period from 1910 to 1915, as the trees grew from two and one-half to seven feet in height.

RELATION OF INJURY TO CROWN CLASS AND RATE OF GROWTH

From the foregoing section it may be inferred that the weevil has a decided preference for trees of certain ages and heights. Observations have shown that even during the highly susceptible age the weevil shows a preference for certain trees. It seems to choose the most thrifty and rapidly growing trees in the stand.

In even-aged stands growing normally, the most rapidly growing trees are in the dominant and co-dominant crown classes. The distribution, according to crown class, of trees weeviled in 1916, is shown in table 1, wherein it is seen that the weevil injury is confined for the most part to these two classes. Although these data were collected from sample plots in New York, they are typical of the conditions existing in all localities where the white-pine weevil has been studied.

TABLE 1. DISTRIBUTION OF TREES WEEVILED PER ACRE IN 1916, ACCORDING TO CROWN CLASS

Plot no.	Average height growth in 1915 (inches)	Number of trees weeviled in 1916 in the several crown classes			
		Dominant	Co-dominant	Intermediate	Suppressed
2.....	20	128	48	16	0
3.....	20	176	64	0	0
4.....	22	176	48	0	0
5.....	24	192	0	0	0
6.....	15	112	0	0	0
7.....	15	528	64	0	0
8.....	14	240	32	0	0
9.....	18	384	96	0	0
10.....	14	160	0	0	0
11.....	15	144	0	0	0
12.....	16	192	16	0	0
Average.....	17.5	221	33.5	1.5	0

The crown class of a tree is indicative of two things: first, it indicates the position and size of the crown, relative to other trees in the stand; and secondly, in even-aged stands it is an indication of the rate of growth. The plots used as a basis for table 1 were all located in even-aged pure stands of young white pine, in which the dominant trees were, for the most part, rapidly growing trees. The data prove, therefore, that the dominant rapidly growing trees are the ones most attractive to the weevil. However, since the dominant trees were also the most vigorous, there is nothing to show whether it was the superior height or the rapid growth of the trees which determined their susceptibility.

It has been assumed by Peirson (1922) that the height of the tree is the more important factor. This hypothesis is in accord with Peirson's assumption that the beetles probably fly to the terminal shoots of the pines, and, in flying over a stand, select the tallest trees. In the work on which this paper is based, however, no evidence has been found that the beetles reach the terminals by flying. If they creep up the trees from the ground — and there is direct evidence that this does occur (pages 11-12) — it is difficult to see how superior height alone could be an attraction to the weevils.

In the field observations in New York, nothing was found which would prove that the weevil prefers the tallest trees if those trees are not also the most rapidly growing trees in the stand. There was, on the other

hand, no small amount of evidence in proof of the contention that the weevil chooses the most rapidly growing trees, irrespective of their height. In stands where the trees were of unequal ages and heights, the weevil chose for attack the younger and shorter trees whenever these were growing more rapidly than the older and taller trees.

Field observations as to this preference of the weevil were further supported by an experiment at Ithaca in which four saplings were inclosed in a large cage. One of the trees was shorter than the others but was growing more rapidly. Six pairs of weevils were introduced into this cage on May 3, 1915, and left free to select the tree best suited to their needs. On the day following their introduction, four of the six females began ovipositing on the small, rapidly growing tree. No other tree in the cage was attacked during the entire season.

From the evidence at hand, it would appear that, in determining the trees suitable for weevil attack, rate of growth is more important than superior height. But, since rapid growth and superior height go together, the dominant trees in the stand are found to be the ones most often attacked.

This habit of stunting the best trees causes more damage than the mere loss of so much height growth in the infested trees. It interferes with the normal development of the whole stand: for, by its reducing the height growth of the most vigorous trees, the natural process of thinning is retarded; backward trees, instead of being crowded out, overtake the leaders; the rate of growth for the stand, as a whole, is reduced; and, unless artificial thinnings are made, stagnation is likely to result.

LIFE HISTORY AND HABITS

ACTIVITIES OF THE OLD ADULTS

In 1915, the first adults were observed on April 26 and the next day they were seen to be ovipositing. Because of seasonal variations from year to year and because of the variations between different latitudes, a description of the general condition of arboreal vegetation at the time of emergence of the weevils from hibernation gives a more reliable indication of the emergence time than does a date alone. At the time when the weevils first appeared in 1915, the pine buds had just begun to swell. Of the deciduous trees, only the cherries were beginning to put forth leaves. The American elms had begun to set fruit, and the shadbush and the aspens were in full bloom. The oaks and the hickories showed scarcely any sign of activity.

The weevils, as soon as they appear, begin feeding on the buds of the white pine. It is usually the terminal cluster of the leading shoot that is thus attacked. The weevils chew holes in the buds, and sometimes make cavities of considerable size. This injury, in itself, is important; but it is overshadowed by the later and much more serious damage caused by the larvae. Sometimes, to vary their diet a little, the adults feed on the inner bark of the terminal shoot.

There is still some question as to the manner in which the adult weevils reach the leading shoots of the trees. Peirson (1922) assumes that they fly over the young pines and alight on the tallest terminals. This assumption appears to be open to some question, inasmuch as the weevil has

seldom been observed in flight. Only once in the course of these investigations has it been observed on the wing. This one occasion was in mid-afternoon of a warm day in the early spring of 1916. On that day, many weevils were flying. They were strong fliers, and when in the air their movements were similar to those of such bark beetles as *Hylurgops*. This unusual occurrence suggests the possibility of a short period of flight during early spring, whereby the weevil becomes widely disseminated, followed by a period when it seldom if ever takes wing.

FIGURE 5. ADULT WHITE-PINE WEEVIL

During the spring of 1916 an experiment was conducted to determine, if possible, how the adult weevils reach the leading terminal. Two bands of tree tanglefoot, one just above the ground and the other at the base of the leading shoot, were placed around the stems of forty trees in a stand of young white pine that had been heavily infested by the weevil in 1915. These bands were applied just before the adults began to emerge from hibernation. As soon as the weevils began to emerge, about two hundred were collected and liberated on the ground in the center of the plot. This number of weevils, added to the number already present, insured a heavy infestation.

With this arrangement of tanglefoot bands, it was certain that any weevils found on the terminal of a banded tree had not crawled up the tree. Any that were caught in the upper band must have alighted on a side branch, and any that were caught in or found below the lower band must have crawled up from the ground. During the entire season, no weevils were caught in any of the bands. Many weevils, however, were

found collected below the lower bands. Apparently they were attempting to crawl up the trunks in order to reach the terminals.

Up to June 15, none of the banded trees were infested. By July 21, however, three weeviled shoots were found on the plot. It is possible that the weevils may have flown to the tips; but it is more likely that, during a cold rainy period in the latter part of June, the tanglefoot may have become glazed sufficiently to permit its being crossed by the weevils.

Of the thirty unbanded trees on this plot, eleven were weeviled — an unusually large proportion for a single season. From this experiment it appears probable that, although the white-pine weevil is, under certain conditions, a good flier, it usually reaches the terminal shoot by creeping up the trunk of the tree.

In the spring, after having fed on the trees for a day or two, the beetles mate. During the entire oviposition period they are usually found in pairs, a male nearly always being found wherever a female is ovipositing. It appears, however, that they occasionally mate in the autumn previous to the period of oviposition. The possibility of this was first indicated by an occurrence in one of the experimental cages placed over uninfested white-pine saplings near Ithaca. In the spring, before any weevils had been introduced, a female appeared, presumably from the litter at the base of the trees, and promptly began to oviposit. She laid 115 eggs in one shoot and did not move to any other during the entire period. The eggs proved fertile and nearly all of them hatched. No other weevil was in the cage during any part of the season; therefore, the only possible explanation is that either the weevils can reproduce parthenogenetically, or fertilization in this instance had taken place during the preceding year. The latter supposition seems, on the face of it, the more probable. Further evidence on this point is presented later in this paper.

FIGURE 6. EGGS AND YOUNG LARVAE OF THE WHITE-PINE WEEVIL

After the brief period of feeding and mating, the females enter upon the task of oviposition. In ovipositing, the female makes with her beak a cavity in the bark of the terminal shoot, and excavates in the inner bark a chamber somewhat larger than the eggs she intends to lay. After carefully digging out and smoothing off the interior of this chamber, she turns around and feels about with her ovipositor until she finds the opening she has made. She deposits one, two, or occasionally three, of her small, pearly white, translucent eggs in the chamber. The operation of preparing the chamber and depositing the egg or eggs in it, often requires a half hour or more. The weevils continue ovipositing until sometime

in early July. The latest that oviposition has been observed in New York was on July 6. Eggs laid after July 1 are so few as to be a negligible quantity.

DEVELOPMENTAL STAGES

After a short period, which, depending on the temperature, varies from six to fourteen days, the eggs hatch and the tiny larvae burrow downward, first in the inner bark and then between the wood and the bark. They eat out all the soft inner bark and cambium as they advance. Many of these small larvae are smothered in the resin which is exuded by the tissues of the tree. The larva that works ahead of his brothers is at a distinct disadvantage, and is usually overwhelmed by resin and killed.

FIGURE 7. LARVAE OF THE WHITE-PINE WEEVIL, ALMOST FULL-GROWN

This illustrates how the larvae work side by side down the shoot

When, however, the larvae work in close formation, they kill the tissues of the tree so quickly that little resin is exuded. As a result of this, the larvae usually form a ring entirely around the stem and work downward, side by side.

When a larva that is a little older than his companions attains full size, he drops out of the line, burrows into the pith, turns upward, and forms a pupal cell. The remaining larvae work down the stem until each is fully grown. Sometimes, as has been mentioned previously, they pass below the second, or even the third, whorl of branches, thus making the total length of a single tunnel from eight to thirty inches. Occasionally, when they are crowded at a whorl, some of the larvae may work up the laterals and form in them their pupal cells. Frequently one finds more larvae attaining full growth at a given point than can be accommodated in the pith, and then the pupal chambers are formed in the solid wood or between the bark and the wood.

The pupal chambers made in the pith or the wood are merely cells hollowed out of the tissues, without any lining whatever. Those made beneath the bark, however, are of a different type. A hollow is made in the wood, and this is arched over with shreds of wood matted together, forming the chip cocoon which is described by Packard in his fifth report. In none of the infested shoots that have been cut open and examined

FIGURE 8. LARVAE AND PUPAE OF THE WHITE-PINE WEEVIL IN THEIR PUPAL CELLS

FIGURE 9. LARVAL TUNNELS OF THE WHITE-PINE WEEVIL

(and these number several thousand) has any lining been observed in the pupal chambers. Even those chambers that are roofed over with the chip cocoon have no lining on the side toward the wood.

The larva may remain in the pupal cell for two weeks before transforming to the pupal stage. This stage lasts about twelve days, when the transformation to the adult occurs. In 1915 the first pupae were observed on June 17, in shoots collected on May 9. If twelve days are added for the pupal period, fifty-one days were required from egg to adult under

conditions existing at Ithaca during the summer of 1915. This length of time would vary somewhat from season to season, since the length of the developmental period is influenced greatly by weather conditions.

ACTIVITIES OF THE YOUNG ADULTS

After the beetles transform, they remain in the pupal cells for a period of from two weeks to a month. Finally they gnaw their way to the outside. Emergence of the young adults continues from the middle of July through September and into October. After they emerge, the weevils feed for a short time on the newly formed buds or the young growth of pines. After this feeding period they seek a place in which to hibernate during the winter, although the weather at the time may still be warm.

Mention has already been made of an instance in which mating probably took place in the fall before hibernation. In order to determine whether or not this is possible, more than a hundred newly emerged weevils were placed in a cage in the laboratory on August 3. They were fed on fresh pine tips, which they ate voraciously at first. After a few days they became restless and began wandering around as if looking for a place in which to hibernate, but, since there was no litter in the cage, they were forced to keep in sight. They were kept in cages under observation until the last one died, in November. During this period but one pair copulated, and they only once. This indicates that occasionally the weevils may mate during the fall

in which they emerge, but that such mating is rare. Hopkins (1911) states that when the female weevils emerge their ovaries are undeveloped and they could not possibly produce eggs in less than a month after appearing as adults. If they laid eggs so late in the season as this would necessitate, the larvae would not be half grown by cold weather and could be found in the shoots during the winter. Never have living eggs, larvae, or pupae been found during the winter, although hundreds of infested shoots have been examined. One larva and one pupa, both in the pupal chamber, were found in the winter of 1914-15, but they were

FIGURE 10. WHITE-PINE WEEVIL ON A NEW SHOOT

both dead. Apparently they could not survive the cold weather. The larva of the parasite *Eurotoma pissodis* Gir. winters in the pupal cell of its host, and the confusion of this parasitic larva with the weevil may account for the statement repeatedly found in literature, that the weevil may pass the winter as a larva.

It seems possible that not all the adult weevils on the trees in mid-summer are adults that matured the preceding season. Possibly the beetles may live for two or three years and deposit eggs each year. From recorded observations of other species of the genus *Pissodes* in Europe and the United States, this fact has been definitely established (Hopkins, 1911). If this holds true for the white-pine weevil as well as for other species of the genus, then, if every immature individual should be wiped out of existence for one year or possibly two years in succession, there might still be enough adults the third year to reproduce the species and give it a fresh start. It may be due, in part, to its long life as adult, that the weevil is able to withstand the attacks of its enemies so well, winning out by persistence where other means might fail.

During the winter of 1914-15 the weevil was sought in hibernation, but without success. The method used in this search was to sift onto a white oilcloth the trash and litter that was taken from beneath the pines, and examine it carefully. The failure to find the weevil in this way was due, doubtless, to its close resemblance to the bits of trash in which it was hidden. In November, 1915, by the use of a modification of the Berlese trap, six adult weevils were found in the litter taken from under infested trees. On February 1, more litter from beneath the infested pines was examined by the same method and five more weevils were found. Therefore the statement can no longer be questioned, that the beetles hibernate in the litter beneath the infested trees.¹

A description of the trap, or extractor, used in finding these insects in hibernation, may be of interest. It consisted of an inverted cone, made of galvanized iron, with a one-inch opening at the bottom, where a detachable glass bottle was fastened. This cone was surrounded by a water jacket one inch thick. The water was kept warm by the application of heat to a tank at one side, the dimensions of which were 8 by 6 by 2 inches. The tank was connected to the water jacket by a half-inch pipe at the bottom and another at the top, thus providing for the circulation of the water. The legs were of wood and were attached to the water jacket by riveted hinges, so that they might be folded against the cone. A screen was placed about two-thirds of the way down the cone so that the material under examination might not drop to the bottom and clog the opening. The cone was then filled with litter.

In using this apparatus, the water in the jacket should be kept warm but not hot, so that the insects in the litter may be heated slowly. As soon as the insects become warm enough to come out of hibernation, they begin to work their way through the trash. Some burrow downward and fall into the bottle at the base of the cone. Others work their way upward and appear at the surface; these may be prevented from escaping by fastening a piece of cheesecloth over the top of the cone. So

¹ At the conclusion of this study of the life history of the white-pine weevil, it is evident that, of all the earlier workers, Hopkins (1907) has most accurately described the life cycle.

far, this apparatus has proved very useful in collecting all sorts of small and inconspicuous insects in hibernation which are easily overlooked in sifting litter.

CONTROL

The white-pine weevil, like other insects, can be controlled in several ways. The method to be selected depends on the conditions in each individual case. Control is largely an economic problem. The amount of labor and material used in these operations must be in proper proportion to the value of the trees protected. Naturally, the cost in all cases must be reduced to a minimum for the method used; but in some instances the value of the trees justifies the application of expensive control measures, whereas in other instances a comparatively low value of the trees will permit the application of only the indirect or preventive type of control. For example, white pines used in ornamental plantings have a high esthetic value that cannot be measured in dollars and cents. On the other hand, a forest whose only purpose is the production of wood can never be valued at more than the market price of the stumpage. Between these two extremes there are all gradations. In the first instance, a large expenditure would be justified; but in the second, only a small amount of labor and material can profitably be spent. In a timber forest, even a small, injudicious expenditure in the early years of the rotation may consume entirely the profit which the timber might have produced. In the following discussion of control, various methods are considered and the conditions under which each is applicable are pointed out.

Methods applicable in the control of the white-pine weevil may be divided into more or less arbitrary groups and subgroups, as follows:

- A. Direct control: Operations that deal directly with one or more stages of the insect, or with trees or parts of trees already infested. (Curative methods.)
 - 1. Chemical methods: Sprays; washes; repellents.
 - 2. Mechanical methods: Collecting adults; pruning shoots.
- B. Indirect control: Operations designed to modify environmental factors so that conditions become unfavorable for the rapid increase of the insect. (Preventive methods.)
 - 1. Silvicultural methods: Shading; dense planting.
 - 2. Biological methods: Parasites; predatory agents.

These groups are taken up in their proper sequence.

DIRECT CONTROL

The direct methods of control are in their very nature rather expensive, and are therefore limited in application to trees that have some value in addition to the wood produced. Only under unusually favorable conditions will growth be sufficiently rapid and stumpage prices sufficiently high to justify the application of these control methods in a timber forest.

Direct control is applicable to shade and ornamental trees in commercial nurseries, to windbreaks, along highways, on watersheds, and in some private forests where the esthetic value is an important consideration.

It is to be expected that not all methods of control are equally effective. In order to justify their use under any circumstances, direct methods must

be highly efficient in checking or preventing weevil injury. They must surpass in efficiency the indirect methods in proportion to the increased expense, or their use cannot be justified.

Chemical methods

It is doubtful whether the chemical method of applying sprays or washes will prove sufficiently effective in controlling this pest to make its use economically practical. Britton and Walden (1912), at the Connecticut Agricultural Experiment Station, have conducted a series of experiments testing various materials. They found lime-sulfur, 1 to 8, to be an effective repellent under certain conditions. Experiments with this material at Ithaca failed to give satisfactory control.

The results of insecticide tests conducted in the spring of 1916 are summarized in table 2. No safe and effective spray or wash has as yet been developed, with the possible exception of a spray of creosote or carbolineum.

TABLE 2. CHEMICAL EXPERIMENTS IN THE CONTROL OF THE WHITE-PINE WEEVIL, 1916

Material	Date of application	Method of application	Strength	Chemical injury	Number of trees treated	Number of trees weeviled
Kerosene emulsion.....	April 26	Painted	1 to 3.....	Considerable....	10	5
Creosote emulsion.....	April 13	Sprayed	1 to 2.....	None.....	10	3
Creosote.....	April 13	Painted	Pure.....	Trees badly injured.....	10	0
Carbolineum.....	April 13	Painted	Pure.....	Growth stopped.	10	0
Creosote.....	April 13	Sprayed	Pure.....	Very slight....	10	0
Carbolineum.....	April 13	Sprayed	Pure.....	None.....	10	0
Soluble sulfur.....	April 26	Sprayed	4 lbs. to 100 gal.	None.....	10	4
Lime-sulfur.....	April 26	Sprayed	1 to 8.....	None.....	10	5
Scalecide.....	April 26	Sprayed	1 to 25.....	None.....	10	8
Powdered lead arsenate.....	May 6	Sprayed	3 lbs. to 50 gal.	None.....	10	3
Powdered lead arsenate.....	May 6	Sprayed	2 lbs. to 50 gal.	None.....	10	2
Paste lead arsenate.....	May 6	Sprayed	5 lbs. to 50 gal.	None.....	10	4
Paste lead arsenate.....	May 6	Sprayed	10 lbs. to 50 gal.	None.....	10	2
Calite.....	May 6	Sprayed	1 to 50.....	None.....	10	3
Calite.....	May 6	Sprayed	2 to 50.....	None.....	10	3
Carbolic emulsion.....	May 26	Sprayed	1 to 3.....	None.....	10	4

Check trees

Group no.	Number of trees	Number weeviled
1.....	30	8
2.....	6	2
3.....	10	3
4.....	10	5
5.....	10	4

Mechanical methods

Collecting adults

In Connecticut, Walden (1915) conducted a series of experiments to prove whether it is possible to control the white-pine weevil by collecting the adults during the period of oviposition. In those experiments a large net was held on one side of the terminal and a sharp blow was

administered on the other side, the beetles being thus knocked into the net. They were then removed from the net and killed. Undoubtedly an immense number of adult weevils can be captured in this way, but the question arises as to whether any appreciable impression on the numbers of the weevil can be made by this method in a badly infested plantation. Furthermore, the expense is too great for use in commercial plantations, since it costs from \$1.50 to \$2 per acre to make six collections. Felt (1906) also recommends this method of weevil control, and Peirson (1922) obtained satisfactory results by collecting adults in this way.

A method employed on the Clark estate near Cooperstown, New York, has combined the collecting of adults with the cutting and burning of the infested shoots. There are on this estate about sixty-five acres of white pines, which were planted in 1910. These trees were attacked when they were smaller than is ordinarily the case. At the time of the first attack, in 1915, and for a number of years thereafter, a fight was waged against the weevil. Two boys were employed at \$1.50 a day for practically three months of each year, barring rainy days. This made the cost of protection a little more than \$3 per acre; which would be prohibitive on a strictly commercial plantation. However, it was the belief of the forester in charge that this method was effective.

From experiments near Ithaca it was found that during the early part of the breeding season it was impossible to collect more than 75 per cent of the weevils by the net method, and at this time oviposition was going on at a most rapid rate. Small, badly infested plots were selected and were gone over with the net and stick. At this time the terminal cluster of buds had not separated. The weevils hide in the terminal bud cluster, and, when shaken, wedge themselves between the buds. Those that were resting on the stem could easily be knocked off into the net, but those hidden among the buds wedged themselves in so tightly that, in spite of the hardest shaking and knocking, they were not dislodged. After a plot had been covered with the net, each shoot was examined very carefully, and in every plot a considerable number of weevils were found hidden among the buds. In one case there were more weevils left in these bud clusters than were caught in the net. Undoubtedly a number of weevils, when shaken off, missed the net and afterward found their way back to another terminal. After the bud cluster had separated, however, exceedingly good results were obtained by this method. It was impossible to tell just what percentage of the weevils were captured, but, judging from the small number to be found in the plots a day after the collecting, the proportion caught was high. From this it would appear that during the first of the season, collecting with a net is of little value though after the separation of the bud clusters it is satisfactory so far as the number of insects caught is concerned; but unless it is used in combination with some other method, such as pruning the weeviled shoots, its efficiency is doubtful.

Pruning shoots

The most practical and economical of the mechanical methods brought to the writer's attention, seems to be that of removing and destroying infested shoots before the emergence of the weevils. Even this method is too expensive and uncertain for general use in timber plantations. How-

ever, the practice has been advised by almost all the investigators who have studied the white-pine weevil, and, although it is rather expensive, it is reasonably effective if the work is done carefully and on a sufficiently extensive scale. The best time for such pruning, if it is possible to go over the plantation only once during the season, is the first half of the month of July. At this time practically all of the infested shoots have begun at least to wilt, and most of them have turned brown. They are then very conspicuous. The work can be carried on rapidly in a plantation where the trees are set out regularly. In irregular natural reproduction, it goes forward much more slowly and is correspondingly more expensive. If possible, it is better to remove the infested leaders twice during the year: once in the month of June, and again in August. Before July few of the weevils will have worked below the first whorl of branches; and if the shoots are removed before that time, the leader will come from this whorl and a whole year of height growth will not be lost. Also, the laterals in this topmost whorl, because of their small diameter, are capable of straightening up much more easily than are the larger branches of the lower whorls. In July many larvae pass below the first whorl, and sometimes by the end of that month, the second, and in a few cases the third, whorl has been passed. This results in a much more serious loss in height growth, and causes a crook that may never grow out. In June it is difficult to get all of the infested shoots, as some, in which the eggs have been recently laid, are rather inconspicuous; and so, in order to get those that have been missed, a second cleaning in August is advisable.

The infested shoots should be removed every year as long as the weevils are in sufficient numbers to endanger the final stand. This is especially true if the infestation is concentrated in spots, as is often the case. The weevils, as has already been mentioned, may perhaps live for as long as three years and deposit eggs each year. If this is the case, it will be necessary to prune for at least three years in succession, in order to suppress an outbreak. If there are any other infested trees near by, they should be treated in order to reduce the number of weevils flying into the plantation from outside. By this method the weevil can be kept within reasonable limits.

All infested shoots pruned from the trees should be disposed of in such a way as to make certain that the weevils do not get back to the trees. If a shoot is cut off and left lying on the ground, the weevils will go through their transformations and emerge just as if the shoot had been left on the tree. Therefore such shoots must be disposed of in some way. The method usually employed is that of burning the shoots. This certainly kills the weevils, but it also destroys the parasites and some of the predacious insects that prey upon the pest. Naturally the destruction of these beneficial forms of life is not to be desired. A much better plan is to place the infested shoots in receptacles tightly covered with wire mosquito netting. The wire netting should be no coarser than sixteen meshes to the inch. The weevils will be unable to escape, but most of the parasites can easily make their way through the meshes. Any tight receptacle will answer. It should not be stood upright, as in that position it would collect rain. The containers should not be opened, but should be left in the plantation until the following spring, when all the parasites will have emerged. By that time the weevils should all be dead, since the

containers offer neither food nor a suitable place in which they could hibernate.

Removal of the infested shoots suffices to keep the weevil in check, but it does not help the trees to recover from the attack. The laterals compete for the leadership, just as if the dead leader were still on the tree, and the characteristic forks and crooks result. If the trees are pruned in such a way as to favor one of these laterals as leader, they will be able to outgrow the weevil injury much more rapidly than would otherwise be the case. Experiments made in the course of the work show that, where the expenditure is possible, the pruning of all but one lateral in the topmost living whorl gives the best results.

INDIRECT CONTROL

In commercial forests, where the value of the trees must be measured in terms of the wood produced, it is essential that the cost of controlling tree insects be kept at the minimum. This is particularly true during the early years of the rotation, when even a small injudicious expenditure means a considerable loss. It is under such conditions that one must rely almost entirely on the indirect methods of control.

Silvicultural methods

It has been the observation of all who have made any study of the injury done by the white-pine weevil that this insect shows a decided preference for trees growing under certain conditions. If this holds consistently true, then one method of indirect control would be to secure for the growing of white pine those silvicultural conditions which are least conducive to weevil injury. In order to ascertain just what would be the best conditions to maintain, a series of experiments was carried on in New York and in Minnesota by means of sample plots laid out under as wide a variety of conditions as could be found.

These sample plots were, for the most part, one-tenth acre in area, although in some cases they contained only one-sixteenth of an acre. In a few instances larger plots were used. By using small plots it was possible to so place them that conditions were fairly uniform throughout a plot, which with plots of larger size would have been difficult if not impossible. Practically all the plots were in stands resulting from natural reproduction.

In these plots the relation of weevil injury to density, rate of growth, age, forest composition, and other factors, was studied with the object of determining what conditions would secure for white pine the greatest immunity from the weevil. The purpose of the experiments was to obtain data sufficiently conclusive to warrant the recommendation of certain definite silvicultural methods that would eliminate, or at least materially reduce, the weevil injury. This aim has been attained; and the correctness of the conclusions drawn from the data collected in this phase of the work have been verified by later investigators.

It has already been noted that under certain conditions the weevil is particularly abundant — for example, on young trees from five to twenty years old, and trees that are growing unusually rapidly. These two conditions are not of particular interest in the control phase of the work,

because their very nature precludes their being changed. It is obvious that all trees must pass through the susceptible age before being marketed, and that rapid growth is in itself a commercial asset. These investigations have, however, brought to light other silvicultural conditions favorable to the weevil, which can be so changed as to reduce the amount of injury, in some cases even to the point of almost complete elimination.

Shading

All students of the white-pine weevil have noticed the insect's preference for trees exposed to the sunlight, and the present experiments have confirmed this observation. Weevil injury is invariably greatest in open pure stands of white pine growing in full sunlight. In mixed stands where hardwoods shade the pine, the injury is much less, and it decreases with increasing intensity of shade until it reaches the zero point under a shade such as that cast by an average stand of oak or maple.

In view of these facts, which have been proved beyond a doubt, it would seem that some system of silviculture should be used which would provide shade during the susceptible period: that is, until the trees are about fifteen to twenty-five years of age or somewhere between twenty and thirty feet in height. The density of shade tolerated by white pine will vary with soil and moisture conditions, and therefore must be determined separately for each site. On the better soils in New York, and to a lesser degree in the Lake States, white pine reproduces itself well under the shade of mixed pine and hardwoods. It would, however, be necessary to liberate the young pines by means of thinning, in order to prevent their succumbing to the competition of the overtopping trees. On poorer soils it would be difficult, if not impossible, to grow pines under hardwoods.

Pearson (1922), as a result of experiments conducted in Massachusetts, also advises the growth of white pine in mixture with hardwoods in such a way that the pine will be slightly overtopped until it has reached the age of about fifteen years. In view of the fact that his experiments were carried on without knowledge of the work done around Ithaca or in Minnesota, it would appear that shading has received double recommendation as a silvicultural method of control.

Dense planting

The question next arises as to whether it is possible to protect pine from the weevil where shading is not feasible, such as in pure stands of volunteer young pine in old pastures, and in white-pine plantations. The answer to this question was found in data gathered from the sample plots of experiments regarding the influence on weevil injury of density of stand. From these experiments it appeared that in open stands where the crowns are all free, the trees are all equally subject to attack, but in the denser stands the percentage of trees that are attractive to the weevil is much less. In very dense stands the injury caused by this insect may be reduced to almost nothing.

The effect of the original density of the stand upon its susceptibility to weevil attack is illustrated in figure 11. The curve indicating the percentage of trees that show the effect of weevil injury at thirty years is a summary of data collected in Minnesota, and represents an average of all plots where the original density could be determined with reasonable

accuracy. The other curves in the figure are from data collected on the Luther Forest Preserve in New York State.

From these curves it is evident that the percentage of trees attacked by the weevil decreases with an increase of density. But, important as

FIGURE 11. EFFECT OF ORIGINAL DENSITY ON SUSCEPTIBILITY TO WEEVIL ATTACK

this fact may be, it is overshadowed by the influence of density on the ability of the trees to outgrow weevil injury. In the open, a tree has no competition from the sides and therefore the stimulation to straight growth is slight. White pines growing in the open, practically never outgrow weevil injury. They almost invariably become forked and

crooked. On the other hand, when a tree is growing in a closed stand, competition for light and space stimulates it to straight growth. The denser the stand, the greater is the competition and the greater is this stimulus to straight growth. As a result of this competition, trees growing in a dense stand are much better able to straighten and to outgrow the injury caused by the weevil attacks, and thereby to become merchantable. The percentage of unmerchantable trees bears a direct relation to the original stand density, as is indicated in figure 12, which shows the pro-

FIGURE 12. PER CENT OF UNMERCHANTABLE TREES PER ACRE WITH DIFFERENT ORIGINAL DENSITIES

portion of unmerchantable trees that may be expected at the age of thirty years at different densities. It is interesting to note that the curves for New York and Minnesota are fairly similar in spite of the climatic differences existing between the two localities.

On comparing figures 11 and 12, it will be noted that in fully stocked stands the proportion of weeviled trees that actually become unmerchantable is comparatively small. Where there is a strong stimulus to straight growth, almost all the weeviled trees succeed in outgrowing the injury. At a density of 1200 trees per acre, only a few trees are likely to be unmerchantable. At an original density of 1500 trees per acre, a fully stocked stand of straight merchantable trees is practically assured.

FIGURE 13. NUMBER OF TREES PER ACRE IN A FULLY STOCKED STAND OF WHITE PINE AT DIFFERENT AGES

The reduction in number is due to the natural thinning that results from competition
(After Hawley, *Practice of Silviculture*)

From these data it is apparent that if a density of from 1200 to 1500 trees per acre is maintained throughout the early years of the rotation, and a fully stocked stand thereafter (figure 13), then the loss resulting from the attack of the white-pine weevil will be at a minimum. On the other hand, if a fully stocked stand is not maintained, a serious loss may be expected. This conclusion has been independently verified by Peirson (1922), so that the facts of the case appear to be well established.

From the foregoing discussion it is evident that there are two possible courses of silvicultural procedure that may be followed for protection of white pine from the weevil. In mixed forests on good soil, this tree can be grown

FIGURE 14. TREE IN DENSE STAND OUTGROWING WHITE-PINE-WEEVIL INJURY

The leader was killed by weevils, and the lateral gained the lead and formed a nearly straight trunk. The remnant of the leader still projects from the side of the trunk

as an "under-story" until a height of from twenty to thirty feet has been attained. The shade should, however, be kept sufficiently light to permit good growth. In pure stands, the keynote of weevil control is a fully stocked stand. Open natural stands should have at least 1200 trees per acre, and, wherever possible, 1500 per acre. A density of at least 1200 trees per acre should be maintained until the trees are somewhere between twenty and twenty-five years old.

Biological methods

Control of insect pests by means of parasites and predatory agents is always an attractive possibility. Too often, however, closer investigation brings disappointment, particularly with native pests such as the white-pine weevil.

If it were possible to breed and liberate parasites in such large numbers as to be an effective control for this insect, or if the activities of predatory agents could be regulated, the problem of weevil control might be much simplified. Such regulation, however, is impossible, at least at the present time. Nevertheless these biological elements play an important part in reducing the numbers of the weevil. If it were not for parasites, predatory agents, and occasionally adverse weather conditions, this insect would be a much more serious menace to our white-pine plantations than it is. Therefore they cannot be disregarded.

Parasites

During the course of these investigations a record was kept of the activities of insect parasites of the white-pine weevil. The most abundant, both in New York and in the Lake States, is *Eurytoma pissodis* Gir. In addition to this parasite, three unidentified chalcids and two braconids have been reared. These, however, have been found in such small numbers as to be an almost negligible quantity so far as their value in controlling the weevil is concerned. It seems probable that they are more or less accidental parasites, and ordinarily find their host in some other insect; or that they are more or less general parasites, and therefore attack a number of other species of insects.

Peirson (1922) reared parasites of the white-pine weevil belonging to the genera *Epirus*, *Eurytoma*, *Microbracon*, and "apparently *Paracrias*." A number of other occasional parasites have been reared by various workers.

In the vicinity of Ithaca, *Eurytoma pissodis* Gir. is by far the most important parasite of the weevil. In several pine clumps the weevils were found to be parasitized to the extent of 50 per cent. A similar degree of parasitism by this species seems to hold consistently wherever the white-pine weevil is abundant. During the month of July the adult parasites were abundant among the infested pines, ovipositing on the larvae of the weevil. The eggs were not observed, but apparently they are deposited in the pupal cells of the weevil since the larvae feed externally upon their host and have been found only in these cells. When fully grown, the larvae of this parasite are only a little smaller than the weevil pupa. They pass the winter in the pupal cells of their host, and in the early spring they pupate and transform to the adult. The whereabouts

of this parasite during May and June is unknown. Perhaps another generation develops on an alternate host.

Predatory agents

In spite of its parasites, the weevil succeeds in maintaining itself. But these are not the only enemies it must face, for there are a number of predacious insects that prey upon it. Packard (1885) states that the young beetles of the family Tenebrionidae are very commonly predacious on the larvae of the white-pine weevil. In the course of this study none of these beetles have been observed at this work, but the adults of this family are very common about young pines. Larvae of the clerid beetles, however, have been found in the weeviled shoots, evidently feeding on the larvae of *Pissodes*, and one of these was successfully reared and identified as *Elasmoserus terminatus*.

The most important predacious insect that has been observed in this study is the larva of a fly which is not ordinarily considered as predacious, *Lonchaea rufatarsus*. The larvae of this species are universally present in shoots infested by the white-pine weevil, and, when confined with the weevil larvae, have been observed to feed upon them. Usually several will attack one larva at the same time and will devour it quickly. They feed upon the pupae also, to some extent.

In addition to insects, there are predacious mammals and birds which feed to a greater or less extent upon the weevil in its various stages. Undoubtedly, as compared with the birds, the mammals are of little account in this work; but in the pine thickets there are found several small mammals, the food of which consists to a greater or less degree of insects. The short-tailed shrew is very common in such places, and lives entirely on insects. During the months when the weevils are in or on the ground, particularly at the seasons when they are emerging from or entering hibernation, the shrews have an excellent opportunity to feed upon them. Practically all of our field and wood mice are very fond of insects of any kind — several of these animals, which were being kept in a cage, preferred them to grain — and so it seems likely that some weevils fall victims to these small rodents.

Of all the vertebrates, however, the birds are undoubtedly the greatest enemies of the weevil. It is more or less common, in infested patches, to find some of the infested shoots cleaned out by birds, although no birds have been observed in the act. Of our common birds, the most likely to do this work are the nuthatches, the chickadees, and the downy woodpeckers. Unfortunately, these birds continue to work on the dead shoots throughout the winter, when only the parasites are present, and thus undo a part of the good work that was begun in the summer. The pine thickets are a favorite place throughout the entire year for many of our birds, and undoubtedly they feed upon the adult weevils to some extent at least. In the spring and fall, when the weevils are entering and leaving the ground, they lay themselves open to the attack of the ground feeders, such as the chewinks, the thrashers, the grouse, and the quail. The chipping sparrows and the wrens are abundant in all white-pine thickets, and probably eat some of the weevils.

Domestic fowl may be of value in controlling the weevil in small plantations. Chickens and turkeys have been used in the South, with marked

success in some cases, to aid in the control of the closely related cotton-boll weevil; and in one instance in New York, the white-pine weevil has been at least partially checked by allowing chickens to run in the plantation.

SUMMARY

The white-pine weevil is one of the most dangerous and destructive pests of young white pine and Norway spruce. It attacks and kills the leading shoots of large seedlings and small saplings. Not only does this injury result in a loss in the rate of height growth, but in many cases it is the cause of serious crooks and forks in the main stem of the tree. In many instances, trees are rendered commercially valueless by the injury.

This weevil has but one generation a year. The adults appear on the trees in the early spring, after having hibernated over winter in the litter beneath the trees. They deposit their eggs in the leading shoots, and the larvae, when hatched, work downward as they feed upon the inner bark. Thus the shoot is girdled and killed. The injury frequently extends below the topmost whorl of laterals, and sometimes below even the third whorl from the top. After completing their growth, the larvae excavate pupal cells in the pith or the wood, transform to the pupal stage, and emerge as adults in August and September. After feeding for a short time on the new growth and buds of the pines, the young adults retire to their winter quarters in the litter beneath the trees.

A number of effective methods for the control of the white-pine weevil have been developed. They may be divided into two main groups, direct methods and indirect methods. In the first group are included all of those methods that deal directly with one or more stages of the insect, or with trees or parts of trees already infested. In the second group are included all operations designed to modify environmental factors so that conditions become unfavorable for the rapid increase of the weevil. The direct methods are expensive, and are applicable only in shade and ornamental plantings, along highways, on some watersheds, and in some private forests where the esthetic value is an important consideration. In the timber forest, only the indirect methods can be used except under very unusual circumstances.

Of the direct methods that have been developed for the control of this insect, the most effective is a combination of collecting adults in the spring, and pruning injured leaders in the summer. The best method of collecting adults is to jar the beetles from the trees into a net held beneath them. Pruning of injured leaders should be done in June and again in August, or, if only one treatment is to be given, in July. Pruning all but one lateral tends to stimulate recovery and is a desirable operation when possible.

Sprays and washes have been recommended for the control of the white-pine weevil, but they have not proved sufficiently effective to warrant their use.

In timber forests, one of two practical methods of indirect control can be used. In established forests on good soil, white pine should be grown under a form of the shelter-wood system that will provide a light shade for the young trees until they have reached a height of from twenty to thirty feet. The shade should not, however, be sufficiently heavy to seriously interfere with the growth of the trees.

On the poorer soils, in plantations, and in volunteer growth in old fields where even-aged pure stands cannot be avoided, another method for weevil control is effective. It has already been shown that density is an important factor in determining both the abundance of the weevil and the amount of injury resulting from its attack upon the trees. With an original density of 1500 trees per acre, very little weevil injury will be evident at the end of the rotation. Even at the standard density of 1200 trees per acre, comparatively little injury will result if a fully stocked stand is maintained throughout the first thirty years after planting. This is due in part to a reduction in the amount of weeviling, but in a much greater measure to the effect of competition in stimulating straight growth.

Parasites and predatory agents play an important part in the natural control of this insect. The most abundant and effective parasite throughout the range of the weevil is *Eurytoma pissodis* Gir. In some cases more than 50 per cent of the weevil larvae are parasitized by this species. Predatory insects and vertebrates are important aids in reducing the numbers of the weevil.

In view of the facts brought to light by the experiments in both New York and Minnesota, it seems safe to say that the young white pine needs no longer to be seriously menaced by the white-pine weevil, because the insect can be controlled under any condition in which a fully stocked stand can be maintained throughout the first twenty-five or thirty years of the rotation, or where white pine can be grown under the shelter-wood system or some other system that will provide shade for the sapling pines. This means that where good silvicultural methods are in force, this pest can be effectively and economically controlled.

BIBLIOGRAPHY

- BELYEA, HAROLD CAHILL. The control of white pine weevil (*Pissodes strobi*) by mixed planting. Journ. forestry 21:384-390. 1923.
- BLACKMAN, M. W. Report on the white pine weevil. Maine Forestry Dept. 1919.
- BLACKMAN, M. W., AND ELLIS, W. O. The white pine weevil. In Some insect enemies of shade trees and ornamental shrubs. Bulletin of the New York State College of Forestry at Syracuse University, 16²⁶:72-78. 1916.
- BRITTON, W. E. The white pine weevil. In Nineteenth report of the State Entomologist of Connecticut for the year 1919. Connecticut Agr. Exp. Sta. Bul. 218:144-155. 1920.
- BRITTON, W. E., AND WALDEN, B. H. Record of preliminary tests to prevent damage by the white pine weevil. Connecticut Agr. Exp. Sta. Ann. rept. 35(1911):307-309. 1912.
- CHITTENDEN, F. H. The white-pine weevil. In The white pine. U. S. Agr. Dept., Forestry Div. Bul. 22:58-59. 1899.
- CURRIE, ROLLA P. The white-pine weevil. In Catalogue of the exhibit of economic entomology at the Lewis and Clark Centennial Exposition, Portland, Oregon, 1905. U. S. Ent. Bur. Bul. 53:91. 1905.
- EMMONS, E. Genus *Pissodes* (Germ.). In Agriculture of New York, 5:108. 1854.
- FELT, EPHRAIM PORTER. White pine weevil. In Insects affecting park and woodland trees. New York State Mus. Mem. 8²:397-401. 1906.
- White pine weevil. In Twenty-ninth report of the State Entomologist on injurious and other insects of the State of New York, 1913, p. 30-33. 1915.
- FERNALD, H. T. White pine weevil. In Shade trees, by E. A. Start, G. E. Stone, and H. T. Fernald. Massachusetts Agr. Exp. Sta. Bul. 125:58. 1908.
- FITCH, ASA. White-pine weevil, *Pissodes Strobi*, Peck. In Fourth report on the noxious and other insects of the State of New York, p. 732-736. 1858.
- FROTHINGHAM, E. H. White pine weevil. In White pine under forest management. U. S. Agr. Dept. Bul. 13:62-63. 1914.
- FULLER, A. S. White pine weevil. Amer. ent. and bot. 2:26. 1870.
- A good word for the white pine weevil. Amer. ent. 3(2d ser., 1): 5-6. 1880.
- GIRAULT, A. A. New chalcid flies, with notes. Bul. Brooklyn Ent. Soc. 12:86-89. 1917.

- GRAHAM, S. A. Notes on the control of the white pine weevil. Journ. econ. ent. 9:549-551. 1916.
- The white-pine weevil and its relation to second-growth white pine. Journ. forestry 16:192-202. 1918.
- HARRINGTON, W. HAGUE. Rhynchophora — weevils. Ent. Soc. Ontario. Ann. rept. 1880:49-57. 1881.
- HARRIS, THADDEUS WILLIAM. A report on the insects of Massachusetts injurious to vegetation, p. 1-459. (Reference on p. 63-65.) 1841.
- A treatise on some of the insects injurious to vegetation, p. 1-640. (Reference on p. 72-73.) 1862.
- HOPKINS, A. D. Catalogue of West Virginia forest and shade tree insects. West Virginia Agr. Exp. Sta. Bul. 32:169-251. (Reference on p. 205.) 1893.
- Report on investigations to determine the cause of unhealthy conditions of the spruce and pine from 1880-1893. West Virginia Agr. Exp. Sta. Bul. 56:197-461. (Reference on p. 259-260, 345, 429, 441.) 1899.
- Insect enemies of forest reproduction. U. S. Agr. Dept. Year-book 1905:249-256. 1906.
- The white-pine weevil. U. S. Ent. Bur. Circ. 90:1-8. 1907.
- Practical information on the scolytid beetles of North American forests. I. Barkbeetles of the genus *Dendroctonus*. U. S. Ent. Bur. Bul. 83¹:1-169. 1909.
- Contributions toward a monograph of the bark-weevils of the genus *Pissodes*. U. S. Ent. Bur. Tech. ser., no. 20¹:1-68. 1911.
- HOUSER, J. S. The white pine weevil. In Destructive insects affecting Ohio shade and forest trees. Ohio Agr. Exp. Sta. Bul. 332:323-325. 1918.
- LINTNER, J. A. Fourth report on the injurious and other insects of the State of New York, p. 1-237. (Reference on p. 24.) 1888.
- *Pissodes strobi* Peck. In Ninth report on the injurious and other insects of the State of New York for the year 1892, p. 344-345. 1893.
- PACKARD, A. S. Insects injurious to forest and shade trees. U. S. Ent. Com. Bul. 7:1-275. (Reference on p. 185-188, 228-229, 236, 241.) 1881.
- The white pine weevil, and its injury to shade and forest trees. In Report of the Commissioner of Agriculture, 1885:322-325. 1885.
- The white-pine weevil. U. S. Ent. Com. Rept. 5:734-741, 829-830, 861, 872. 1890.

- PAUL, B. H. An ecological study of typical forest plantations in New York State. Thesis for degree of master in forestry, Cornell University. 1915.
- PECK, W. D. On the insects which destroy the young branches of the pear-tree, and the leading shoot of the Weymouth-pine. Massachusetts agr. journ. 4:205-211. 1817.
- PEIRSON, H. B. Control of the white pine weevil by forest management. Harvard Forest. Bul. 5:1-42. 1922.
- PIERCE, W. DWIGHT. On the biologies of the Rhynchophora of North America. Nebraska State Bd. Agr. Ann. rept. 1906-07:249-319. (Reference on p. 259.) 1907.
- RILEY, C. V., AND HOWARD, L. O. Some of the bred parasitic Hymenoptera in the national collection. Insect life 2:348-353. (Reference on p. 348.) 1890.
- The white pine weevil. Insect life 3:468. 1891.
- SANDERSON, E. DWIGHT. The white pine weevil. In Insects mistaken for the Mexican cotton boll weevil. Texas Agr. Exp. Sta. Bul. 74:5. 1904.
- SAUNDERS, WM. The white pine weevil — *Pissodes strobi*. In Insects injurious to the white pine — *Pinus strobus*. Ent. Soc. Ontario. Rept. 1883:55. 1884.
- THOMAS, CYRUS. *Pissodes strobi* — Peck. (White-pine weevil.) In Sixth report of the State Entomologist on the noxious and beneficial insects of the State of Illinois, p. 133-134. 1876.
- WALDEN, B. H. Experiments in controlling the white pine weevil. Connecticut Agr. Exp. Sta. Ann. rept. 38(1914):173-176. 1915.
- Experiments in controlling the white pine weevil in 1915. Connecticut Agr. Exp. Sta. Ann. rept. 39(1915):134-136. 1916.

Insects Attacking Forest and Shade Trees

BY
H. B. PEIRSON

BULLETIN NO. I
MAINE FOREST SERVICE,
AUGUSTA, MAINE
1923

In the absence of state appropriations available for the purpose, the work on which this publication is based has been made possible by contributions from persons interested in the establishment of forest entomology. A large part of the notes on white pine insects were collected by the writer on the Harvard Forest at Petersham, Massachusetts.

Contents

	Page
Introduction	5
Spruce and Fir Insects	8
Spruce Bud Worm	8
Introduction	8
History	9
Distribution	9
Nature of Damage	11
Food Plants	14
Seasonal History and Habits	15
Description of Stages	18
Natural Enemies	19
Control by Forest Management	20
Salvaging of Budworm Killed Timber	23
Bark Beetles	25
Eastern Spruce Beetle	25
Spruce Bark Beetle	26
Balsam Fir Bark Beetle	26
Secondary Bark Beetles	26
Control	26
Wood Borers	28
Round-Headed Borers	28
Flat-Headed Borers	28
Control	29
White Pine Insects	30
Introduction	30
Insects Attacking Seedlings	32
White Grubs.	32
Pales Weevil	33
Strawberry Crown Girdler	33

Insects Attacking Sapling Pine	35
White Pine Weevil	35
Mound Building Ants	37
White Pine Aphid	39
Gipsy Moth	39
Pine Tube Moth	40
Miscellaneous Species	40
Saw-Flies	41
European Pine Shoot Moth	41
Spittle Bugs	41
Pine Leaf Miner	41
Pine Leaf Scale	41
Insects Attacking Mature Trees	41
Bark Beetles	42
Pine Borers	42
Carpenter Ants	43
Control	44
Wooly Aphid	44
Summary	45
Larch Insects	47
Larch Saw Fly	47
Larch Case Bearer	47
Hardwood Insects	49
Poplar Borer	49
Maple Borer	51
Bronze Birch Borer	51
Forest Tent Caterpillar	51
Powder Post Beetles	52
Shade Tree Insects	53
Spruce Gall Aphid	53
Spruce Cone Worm	53
Oak and Maple Twig Pruner	53
Elm Leaf Beetle	54
Borers	54
Leaf Spots	55
Economic References	55

Insects Attacking Forest and Shade Trees

INTRODUCTION

The work in forest entomology was started in September, 1921. A survey of the field left no doubt that the principal problem to be solved was a practical control for the spruce bud worm which already had destroyed fully forty per cent of the merchantable spruce and fir in the State. Consequently most of the time has been devoted to a study of this problem, a report of which follows. The situation was aggravated by the fact that the State had failed to take any steps whatever to stop the wholesale destruction of its forests until the outbreaks had largely subsided. At the present time all efforts are being bent to prevent a reoccurrence of the devastation which these relentless insect enemies have wrought. New outbreaks of the bud worm are continually starting up and requiring attention.

Owing to the importance of the white pine crop in the southern portion of the State considerable time has been spent in a study of insect conditions there, and a few suggestions on control of the more important pests of the pine are included in this report. In like manner a large number of reports and inquiries on outbreaks of shade tree insects makes it seem wise to include a short resume of the more important of these.

As much time as possible has been given to field work and a large part of the forest district of the State has been covered, detailed field notes on forest conditions having been taken throughout the territory. With the

help of foresters, wardens, and others a forest type map is being made which will be used in connection with the entomology work.

There is a growing interest in the hardwoods of the state and altho time has not permitted the study in any detail of their condition, it has been very evident that there has been considerable damage from insects in several sections of the State. Hardwoods are, however, far more resistant to death from insect attack than are the softwoods. The greatest damage occurs in areas where the trees are either overmature or else are present in pure stands over wide areas, as is quite common in the case of poplar and white birch.

An insect outbreak is very much like a fire in that if located in its early stages it is easily stamped out, but if allowed to run and assume great proportions the chances of stopping it under present forest conditions are very slight. The time to fight an insect outbreak is not after the damage has been done, but immediately after the location of an incipient outbreak. Everything depends upon preparedness. It is time that the word "control" be dropped and the word "prevention" substituted, for there is no more excuse for wide spread insect outbreaks than there is for forest fires. Both can be prevented.

PLATE I. TYPICAL GROUP OF SPRUCE AND FIR DESTROYED BY
BUDWORM

The man is standing by the only live tree. Hairy moss clinging to the branches of dead trees often gives the appearance of foliage at a distance. Note particularly the manner in which the dead fir break off. (Courtesy of R. G. Stubbs).

Spruce and Fir Insects

SPRUCE BUDWORM

INTRODUCTION

Never in the history of forestry has such tremendous damage been done to the forests of the Northeast as that which has taken place in the last ten years following wide-spread outbreaks of the spruce budworm.* The loss that has occurred is beyond all belief and if it were not for actual cruising reports covering large areas of forest land, statements made might well be open to doubt. In Maine reports on township after township show all the merchantable fir and fifty or more per cent of the spruce dead, and in many cases the trees are still dying. There is no question but that fully two-fifths of the merchantable spruce and fir in the state is already dead. In the Provinces of Quebec and New Brunswick the loss is also stupendous and is variously estimated in the billions of dollars.

The fact that heavy infestations of the insect have repeatedly occurred in the past makes it plain that unless definite control measures are taken other and probably even more serious outbreaks are sure to occur in the future. Everywhere the percentage of fir on budworm killed areas is increasing, and as the foliage of this tree is the principal food of the budworm, and is the primary cause of outbreaks, some idea of the seriousness of the situation can be obtained.

*This insect, formerly known as *Tortrix fumiferana*, has just recently been placed in the genus *Cacoecia*.

HISTORY

The history of past outbreaks of the budworm and bark-beetles in the spruce and fir forests of the Northeast is of especial interest in that it throws much light on what may be expected in the present and future outbreaks. The earliest record in regard to the dying of spruce in the forests of the Northeast is found in a letter quoted by Packard. This letter written in 1818 mentions great destruction of spruce east of the Penobscot River. Very little was written on this outbreak due to the fact that at this time very little spruce and no fir was being cut, pine being the principal timber tree lumbered.

The next great outbreak occurred about 1880. The destruction wrought at this time brought forth much more attention, for spruce was being cut in large quantities. Such reports as:—"One billion feet of spruce killed along Allegash and tributaries of the St. John River", "great destruction of spruce in North Somerset County", and "the slump in the amount of spruce coming down the river after the outbreak was very noticeable",—are common. There is no question in the writer's mind but that each of these past outbreaks started with widespread budworm epidemics followed by bark-beetle outbreaks.

The present series of outbreaks of the budworm started about 1910 and are just now on the wane, altho local outbreaks are continually starting up and threatening the remaining green stands of spruce and fir. These are being followed by swarms of bark-beetles which in many localities are attacking the green spruce, and proving a serious menace to the remaining timber. Much interesting evidence relative to the older outbreaks has been found in old reports, obtained from lumbermen, and traced in the retarded ring growth of old spruce.

DISTRIBUTION

The distribution of the spruce budworm (*Cacoecia umiferana*) is restricted to North America, where it has

PLATE II. WORK OF SPRUCE, FIR, AND PINE INSECTS.

A. Cross section of spruce showing ring retardation, due to defoliation, on tree that has recovered from Budworm. B. Ring retardation on tree that has not recovered. C. Egg tunnels of *Pityokleines sparsus* in fir. (Slightly enlarged). D. Tunnels of *Ips perturbatus* in white spruce. ($\frac{1}{2}$ natural size). E. Tunnels of *Ips pini* in white pine. ($\frac{1}{2}$ natural size). F. Egg tunnel and larval mines of *Dendroctonus piceaperda* in spruce. G. Tunnels of Ambrosia Beetles in wood of spruce.

been reported from Maine, Vermont, New Hampshire, New York, New Jersey, Minnesota, Nova Scotia, Prince Edward Island, New Brunswick, Quebec, Ontario, and British Columbia.

NATURE OF DAMAGE

The spruce budworm is a defoliating insect which starts feeding in the spring within the new buds of the fir or spruce, hollowing them out. The larva then constructs a protecting shelter for itself by loosely binding together needles that have been bitten off. It later starts feeding on the foliage itself rapidly stripping the trees. Those trees that are completely stripped are usually killed outright. This is due in the first place to the interference with the normal functioning of the vital organs of the tree,—the lack of foliage bringing about the rapid drying out or death of the root hairs thru which water and essential minerals are absorbed, thus shutting off the food supply of the tree. In the second place conifers, unlike hardwoods, store up comparatively little reserve food so that there is nothing to fall back upon. This is the reason why hardwoods may be defoliated several seasons in succession and still survive, whereas a single defoliation usually kills a conifer.

Retarded Growth

Particularly in the hardwood and mixed softwood-hardwood type where the spruce and fir have not been severely defoliated the growth is often retarded. Much valuable data was collected throughout the State in regard to dates in which the different areas were swept by the budworm, by making a study of the ring growth. Following defoliation the annual growth is greatly reduced, so that it is possible to make an accurate estimate as to date of outbreak by cutting into green trees and counting the rings back to the point where retardation started. To obtain accurate results the count must be

made near the top of the tree. The first year's feeding is scarcely shown in that year's growth, as much of this ring is made up from food stored the previous year. The second year there is a decided reduction, so that the age of the outbreak is found by adding one to the number of rings between the inner bark and the ring in which decided retardation shows. A few rings may show an increase in growth during the second and succeeding years after feeding stops. Near the base of the tree, the retarded growth may not show up until four or five years after the first feeding.

Death from Secondary Causes

Large numbers of trees which have been only partially defoliated are so weakened that secondary enemies easily destroy them. It is, of course, the suppressed or over-mature trees that go first, due to their naturally weak condition. The nature of this dying, which usually extends over a period of several years, has been likened "to a greatly accelerated natural thinning that takes place in the normal forest", correlated with this, it has been very apparent that trees growing on poor or shallow soils are more likely to succumb. Due to the extreme importance of these secondary agencies in bringing about the death of the trees it is well to consider them separately.

1. *Winter Killing.* Large numbers of defoliated trees succumb the winter following defoliation due to winter killing. It has been shown that the ability of plants to withstand freezing depends in large measure on the concentration of the cell sap. Depletion of foliage prevents the formation of stored food thus lowering the concentration of the cell sap and making the trees subject to freezing.

2. *Drying.* In sample plots taken throughout Maine it has been very apparent that many of the trees that were only partially defoliated were unable to live through a dry season such as that of 1921. This is due largely

to the fact that there is little or no reserve food stored up in the trees. An interesting incident occurred in this connection. Of two ornamental firs that had been severely defoliated, one which was kept watered recovered; while the other died. Death from drying is particularly noticeable on mountain tops and other localities where the soil is shallow and moisture drains off quickly. Schierbeck suggests that balsam growing in mixture with hardwoods is better able to recover from serious defoliation due to the protection from evaporation of the soil moisture afforded by the hardwoods.

3. *Bark-beetles*. Considerable loss is caused by the swarms of bark-beetles which come into these areas of weakened trees. It is a common belief that these beetles never attack green healthy trees. This is a complete fallacy. In the normal forest they live and breed largely in sickly trees or green windfalls. Under these conditions birds and parasites are able to hold the beetles in check. Large areas of weakened trees such as are found after a budworm outbreak form ideal feeding and breeding places for bark-beetles so that they are able to increase rapidly in numbers and assume epidemic form. It is at this time that green trees are attacked and very often killed. Ordinarily the outflow of sap or resin brought about by the feeding of the bark-beetles drowns them, thus preventing their increase in green trees. It can readily be seen, however, that any great outflow of sap such as is caused by the attack of myriads of the beetles will so weaken the trees that they will become suitable breeding and feeding places. This is exactly what takes place during a bark-beetle epidemic and the trees are killed just as surely as if they were felled, the galleries made by the bark-beetles cutting off the flow of sap.

4. *Fungi*. Fungus diseases, particularly the shoestring root rot caused by *Armillaria mellea*, account for the death of large numbers of weakened trees. This fungus affects the bark and wood of the roots and destroys the living tissues. The decay very commonly extends up

into the bark and sapwood of the lower part of the trunk. Large amounts of resin commonly exude from the base of effected trees, accumulating as hard cakes. The fungus gets its name from the black strands resembling shoestrings which can be found attached to bark at the base of the tree and running thru the soil. The thin white sheets of mycelium spread out thru the cambium destroying the tissue. The fruiting bodies appear on the sides of the trunk or on exposed roots as clumps of honey-colored mushrooms.

Large numbers of sample plots taken by Craighead in Canada showed that over seventy-five per cent of the dying balsams had one or more roots affected by this disease. In Maine a very similar condition prevails, the disease being common throughout the State.

5. *Excess fruiting.* Following severe defoliation fir, in particular, seeds very heavily. This takes a great deal of strength from the tree at a time when the little reserve food present in the tree might do much to start it on the road to recovery. The trees apparently follow some natural law which causes them to bear a heavy crop of fruit when under adverse conditions in order to insure the continuity of the species.

FOOD PLANTS

The name spruce budworm has been very deceiving for the principal food of the budworm is the balsam fir (*Abies balsamea*). Counts made on hundreds of thousands of trees in Maine show that the relation between dead spruce and dead fir is seldom greater than two to three. In other words it is rarely over wide areas that more than two dead spruce will be found for every three dead fir. The first year's feeding during an epidemic is almost entirely on fir and it is only when the foliage from this tree is largely gone that spruce is attacked. Of the three species of spruce attacked red spruce is by all means the most severely injured. Black spruce is killed

to some extent when growing in mixture with fir. White spruce is rarely killed due to the fact that only the current year's foliage is suitable as food. In Maine this species is usually found in favorable growing conditions, particularly in abandoned pastures and is, as a rule, relatively hardy. In New Brunswick Craighead reports considerable damage to slow growing white spruce, particularly trees having low vitality.

At times hemlock, larch, and white pine are attacked but these are seldom killed. The most severe damage usually occurs in localities where the caterpillars have either dropped onto these trees from spruce and fir or else have been blown on to them.

The following is a list of known food plants:

<i>Pinus strobus</i> Lamb	. . .	White Pine.
<i>Larix laricina</i> (Du Roi) Koch	. . .	American Larch.
<i>Picea mariana</i> (Mill) B., S. & P.	. . .	Black Spruce.
<i>Picea rubra</i> Dietre	. . .	Red Spruce.
<i>Picea canadensis</i> (Mill) B., S. & P.	. . .	White Spruce.
<i>Picea Abies</i> (L) Karst	. . .	Norway Spruce.
<i>Tsuga canadensis</i> (L) Can.	. . .	Hemlock.
<i>Abies balsamea</i> (L) Mill.	. . .	Balsam Fir.

SEASONAL HISTORY AND HABITS

It is well worth while to review in a few words an active outbreak. In the first place observation has shown that the budworm is present at all times in the spruce and fir stands of northern Maine. It is, however, present in such small numbers that little or no attention is attracted by its feeding, which under non-epidemic conditions is limited almost entirely to the tips of fir receiving an abundance of sunlight. When conditions become favorable, the budworm again finding an abundance of food in the sunlight, such as young growth coming up in openings or older growth pushing up through the hardwoods, eggs are laid by the moths on the needles

at the tops of the fir trees that are in the sunlight. These eggs, which are laid from the first to the fifteenth of July, hatch in about ten days and the small caterpillars crawl into crevices in the bark, beneath lichens, or even into the cones, where they spin a small cocoon and thus pass the winter. They come out in the spring just before the balsam buds open and feed for three weeks or more on the foliage. An abundance of food and a favorable season mean that most of the caterpillars will mature and later turn into moths which again lay large numbers of eggs. Thus it is that an epidemic starts.

The moths in flying over the forest seek out and alight on the tops of fir trees. The first year's feeding is almost exclusively on fir, spruce being practically free from injury. The second season again finds the moths in flight. Some of them, however, remain and lay their eggs on the spruce. This accounts for the fact that the greatest injury is to the fir, and also emphasizes the fact that the intensity of an outbreak over wide areas depends largely upon the amount of fir that is present in a region. This habit of the moths seeking fir that is growing in the sunlight also accounts for the fact that spruce and fir found in mixture with hardwoods is less liable to damage from budworm attack than that growing in pure stands. It also accounts for the fact that spruce and fir coming up under the protection of hardwoods or budworm killed timber, is seldom damaged.

Wind undoubtedly plays an important part in the spread of an epidemic. It has been possible to trace the movement of the series of outbreaks occurring during the last ten years by a study of ring retardation in trees that have recovered from budworm injury. Notes were taken throughout the Maine Forestry District and dates of outbreaks plotted on a map. It is quite apparent that as a rule the outbreaks follow the valleys, spreading up on to the ridges. In some sections of the State the spread of the moths was north, in other sections south, depending apparently on the direction of the prevailing

PLATE III. SPRUCE AND FIR INSECTS.

A. Fir tip on which feeding has started. B. Egg mass of Budworm on fir needle (greatly enlarged). C. Larva of Budworm. (Enlarged $1\frac{1}{2}$ x). D. Pupa of Budworm. (Enlarged $1\frac{1}{2}$ x). E. Moth of Budworm. (Enlarged $1\frac{1}{2}$ x). F. Typical barkbeetle tunnel beneath bark, showing large adult tunnel and small larval tunnels (Natural size). G. Parasitic Hymenoptera laying egg in wood boring larvae. (Natural size).

winds during the period of flight. The moths are frequently blown on to high mountain tops where the spruce and fir are severely damaged.

Many reasons have been given for the somewhat periodic appearance of wide-spread outbreaks in the past, but few of these take into account the fact that during an active outbreak the caterpillars are present in such vast numbers and over such a great territory that it is utterly inconceivable that either birds or parasites could check their advance. The one great factor is lack of suitable food. When the majority of the fir has been defoliated the outbreak soon dwindles down.

DESCRIPTION OF STAGES

The Moth

The moth is robust with a wing expanse of slightly less than one inch. The fore wings which are approximately one-half inch in length, are large, broad, and oblong in shape. In color the fore wings are umber brown, the brown sometimes being replaced by a rust red. The ground color of the wing is a bluish slate. There are numerous fine, transverse, short brown lines which divide the wing into checks or squares. Numerous brown spots and patches occur on the wings and near the middle of the wing is a broad brown band crossing it in a zig zag fashion. The umber brown markings which show considerable variation are sometimes replaced by a bright rust red color and the clay-blue by a ferruginous brown. The legs, body, and hind wings are of a glistening umber brown. The length of the body is approximately one-half inch, and in the female is very stout. That of the male ends in a long distinct hairy tuft.

The Egg

The egg is pale green, broad, oval, scale-like, flat beneath and slightly convex above. The shell is thin. The egg is about one-sixteenth inch in diameter. The

eggs are laid in shingle-like patches over-lapping each other irregularly. These patches may be composed of as many as thirty eggs.

The Larva

When first hatched the larva is pale pea-green with a yellowish tinge. The head is dark brown. The mature larva is of a rich umber-brown color, diffused with olive green markings near the sutures or body segments. Very conspicuous yellowish-white warts occur along the sides. The head and fore legs are of a dark brown color. The full grown larva is approximately three-fourths inch in length.

The Pupa

The pupa is approximately one-half inch in length. The body is comparatively thick, tapering at both ends and is of a tan brown color. The antennae and legs are dark horn colored, the wings pale with dark veins, the thorax spotted and with three irregular, dark stripes, abdomen with two rows of stout spines above, and a lateral row of dark spots.

NATURAL ENEMIES

It is unquestionably true that birds and parasites play an important role in holding in check the budworm and subsequent outbreaks of bark-beetles. These, however, cannot be depended upon to prevent epidemics. Several very interesting incidents of the work of both parasites and birds are on record. It is claimed that the budworm gained unusual headway during seasons when the smoke from forest fires drove the birds out of the forests. Several observations show that spiders destroy large numbers of the young larvae.

Very interesting data were collected by the writer on the effects of heavy storms in holding in check incipient outbreaks of the budworm. The young larvae are

washed off from the boughs in large numbers and drowned, and during continued heavy showers an outbreak may be almost entirely stamped out.

CONTROL BY FOREST MANAGEMENT

It is the purpose of this paper to emphasize the possibilities of controlling and preventing wide-spread outbreaks of the spruce budworm by means of forest management. Work has progressed far enough to prove that this can economically be done. It is essential, however, that plans be put into operation at once, for the time to fight an insect outbreak is immediately after it starts. The belief held by some that the danger from the budworm has passed is entirely groundless for new outbreaks are continually starting up. One of the most startling results showing up from the field work of the past season is the ever increasing amount of fir that is coming in to the forests. This only emphasizes the need of putting control and preventive measures into effect immediately, if the remaining green stands of spruce are to be saved.

The following methods of control are already being put into operation in Maine by the writer and if continued and carried to completion should insure the timberlands of the State against future serious damage by the budworm. Already several outbreaks have been stamped out.

(1) *Forest Type Map*

It has been very apparent when studying spruce budworm outbreaks that the type of the forest growth is the controlling factor both as to determining where an outbreak may start and the extent of the damage likely to be done. Sample plots taken throughout the spruce region in Maine have emphasized this point, as have plots taken by Craighead in the Provinces. Outbreaks start in pure softwood areas where fir makes up a high percentage of the stand. Once started the intensity of the

outbreak depends largely upon the percentage of spruce and fir in an area. Spruce and fir occurring in mixture with hardwoods receives a reasonably high degree of immunity due to the protection or shade of the hardwoods. In like manner small isolated bodies of spruce and fir very commonly escape injury.

With these points in view a general forest type map of the entire forested region of the State is being made. Already over one-half of the Maine Forestry District has been typed and the remainder is rapidly being completed. This work is being carried on largely by the Chief Forest Fire Wardens who are in most cases men who are very familiar with their territory. A great deal of help has been received from foresters, cruisers, and timberland owners. For the present only eight types are being considered.

1. Spruce and Fir.	5. Coniferous Swamp.
2. Northern Hardwoods.	6. Aspen or Poplar.
3. Mixed Softwoods and Hard-
woods.	7. White or Paper
Birch.
4. White Pine.	8. Gray Birch.

The final type map will be of utmost importance in locating danger points and in planning control measures once an outbreak starts.

(2) *Patrol of Area*

The second phase of the problem of preventing future outbreaks takes the form of scouting and education. Fortunately the Maine Forestry District, in which the majority of the spruce and fir occurs, is fairly well patrolled by wardens, patrolmen, foresters, and timber cruisers. These men have, for the most part, been made acquainted with the nature of insect outbreaks thru talks and letters and have been requested to notify the office as soon as any are located. In this way the State already has several hundred men constantly on the lookout, and it is hoped to locate outbreaks long

before they reach serious proportions. Budworm and bark-beetle outbreaks often show up from the fire lookout stations, for the reddening of the trees is visible for long distances. In connection with this part of the work the type map will show the areas which should be visited once a year and every effort will be made to have this done. Reports of outbreaks are carefully checked and if deemed advisable the area is visited by the entomologist.

(3) Control of Outbreak

The third and final phase of the problem comes after the outbreak has been located and visited by the entomologist. If the infestation proves serious and gives promise of doing considerable damage, the infested area is marked on a map and the owner notified with the request that he plan his winter's operation, if possible, so as to clean cut the spruce and fir in this area. This request will not be made unless it is absolutely necessary in order to save the surrounding forest, and from the attitude shown so far by the land-owners it is believed that in the majority of cases the request will be granted. If the infestation is located in time it may not be necessary to operate in this region until the second year.

This method of stamping out new infestations, which has already been tried with success, is based upon the fact that the budworm passes the winter as small larvae which have just emerged from the eggs and are so weak that unless suitable food is present when they emerge in the spring the caterpillars will soon die from starvation. Clean cutting the area in the winter means that when the larvae emerge in the spring there is nothing in the area but dry tops, the food supply of the insect having been destroyed. Under average conditions the necessary size of the cut will be comparatively small and could well be handled by a small jobber. The operation should, if properly planned, practically, if not entirely, pay for itself. The protection to the surrounding timber

•

would more than offset any financial loss due to the operation. In the case of owners who do not do their own operating, a small stumpage sale could be planned for the area.

Improvement of Forest Crop

Considerable progress can be made in improving the forest crop so that the likelihood of future wide-spread outbreaks will be much lessened. Serious insect epidemics almost invariably occur in pure stands. In the present budworm outbreak the greatest amount of damage took place in the pure spruce and fir stands, the mixed soft and hardwood stands were far less seriously injured and it is questionable if any outbreak at all would have occurred if it had not been for the large areas of practically pure spruce and fir.

In the future every effort should be made to cull out the fir from the forest and to favor a mixed spruce-hardwood forest. In order to bring this condition about it will probably be necessary to leave a certain number of spruce seed trees and possibly girdle some of the over mature hardwoods in case they are making up too high a percentage of the stand. In areas in which fir does make up a high percentage of the stand a short rotation should be adopted. In the mixed softwood-hardwood type emphasis should be laid upon the culling out of all fir and the cutting of the softwoods as soon as economically possible after they out-top the hardwoods wherever the softwoods make up a high percentage of the stand. In many areas it would be decidedly worth while to merely girdle all fir in the winter in case they cannot be cut and utilized at the time of the operation.

SALVAGING OF BUDWORM KILLED TIMBER

One of the greatest problems which the timberland owners in Maine have had to face has been the salvaging of the dead wood left as a result of the budworm epi-

demics during the last ten years. Many small owners have been practically bankrupted, being unable to harvest or sell their budworm killed timber due to the flooded condition of the market with this dead wood. Some of the large companies which have been salvaging this insect-killed timber during the last five years already have from three to five years' supply cut.

The length of time that the timber is worth salvaging, after having been killed by the budworm varies for both spruce and fir depending largely upon the growing conditions. In general, it has been found that for pulp purposes fir must be salvaged within three years from time of death, and spruce within five and in some cases seven years. Rots and wood-boring insects attack the trees the summer following defoliation by the budworm and soon ruin the wood for lumber purposes, so it is necessary to salvage wood for this use immediately. The fact that the wood will not peel only adds to the difficulties and cost of salvage.

Apparently the more rapid the growth of the spruce and fir the more rapid it deteriorates after death. This seems to be general throughout Maine. Spruce and fir growing in swamps or on extremely shallow ground, such as is found on rocky ridges or mountain tops, soon dry out due to the small amount of sap in the wood, whereas that growing on the more favorable sites contains a considerable amount of sap and remains favorable ground for insects and fungi much longer, besides being more suitable in the first place.

Great savings could be made if these factors were taken into consideration when planning cutting operations. Unfortunately some operators are going at it in just the reverse manner, cutting the wood which will remain sound for the longest time first and leaving the wood which will rapidly deteriorate for future operations. The fact is that this is going to mean that much of the wood will never be salvaged that otherwise might have been. There is a strong reluctance on the part of many operators, and upon individuals as a whole, to face the real situation. The full effects of the budworm outbreak are yet to be felt.

BARK-BEETLES

Several species of bark-beetles at times destroy large areas of spruce and fir. These small beetles bore through the outer bark and lay their small whitish eggs along tunnels made in the inner bark, from which minute whitish grubs hatch and start feeding in the cambium, finally girdling the tree. The attack first shows up on the trees in the nature of fine reddish, sawdust-like borings deposited on the rough bark or lichens. The exuding of pitch mixed with the borings is also very common. The foliage of effected fir turns pale green, later taking on a yellowish and finally a reddish hue, the trees assuming a mahogany-like color. The bark finally loosens and is easily broken off. In the case of effected spruce the old growth needles do not turn color but drop off green leaving the tree with scanty foliage.

EASTERN SPRUCE BEETLE

Dendroctonus piceaperda is by all means the most injurious of the bark-beetles likely to kill spruce in Maine, and has in the past destroyed the red, black, and white spruce over wide areas. This is a reddish brown to black beetle approximately three-sixteenths inch in length. Its body is sparsely clothed with long hairs. The insect passes the winter in the inner bark. The beetles emerge from June to October, the majority coming out in July, and immediately attack trees excavating galleries. According to Hopkins, "The beetles enter the bark of healthy trees at a point from six to ten feet from the base, and that of weakened trees from near the base to the larger branches. In the living trees the entrance burrow is gradually extended obliquely upward, thence in a longitudinal direction upward through the inner bark, often grooving the surface of the wood. Along the sides of this gallery, which is usually about three times as broad as the beetle, the eggs are placed singly in small cavities or in groups in an elongated

cavity. The eggs are then protected by a mass of borings, closely packed and cemented with gum." As this gallery is being built numerous openings to the outside are excavated and then filled with borings as the gallery advances.

SPRUCE BARK-BEETLE

Polygraphus rufipennis is very abundant in the bark of dying spruce and may at times become of primary importance and attack green healthy trees. This beetle is stout, jet black, and slightly over one-sixteenth inch in length.

BALSAM FIR BARK-BEETLE

Pityokteines sparsus at times destroys large amounts of balsam fir in the Northeastern United States. This minute beetle, which is approximately one-sixteenth inch in length, is particularly active in destroying over-mature and weakened trees.

SECONDARY BARK-BEETLES

There are large numbers of species of bark-beetles which attack only weakened, over-mature, or dead trees. It is rarely that an injured conifer is not sooner or later attacked by these beetles, which do much to bring about the rapid death and deterioration of the trees by loosening the bark and allowing moisture to seep in. These beetles are admirably covered by Swaine in a paper entitled "Canadian Bark-Beetles".

Control. In general it can be stated that if active measures are taken to stamp out and prevent epidemics of the budworm, the danger from bark-beetles will be materially lessened, as these almost invariably start up in areas of weakened trees. Outbreaks of bark-beetles are easily located by the scanty foliage or by the yellowing and then reddening of the foliage

on infested trees which shows up for long distances. These so-called "beetle trees" often occur in clumps. Outbreaks can usually be controlled without great difficulty as the habits of bark-beetles render them readily vulnerable by methods that lumbermen can economically employ. The removal and utilization, peeling, or burning of the infested trees is the most satisfactory control, and an infestation can be checked in a single season by destroying these "beetle-trees". Floating the logs in the early spring, as is the usual custom, also destroys the beetles. When only a portion of an infestation can be treated, control measures should be concentrated against the centers of infestation, cutting the trees and either floating them or else peeling off and burning the bark. These methods of control do not apply to the old "red tops" from which the primary bark-beetles have emerged. They do apply though to freshly infested trees on which the foliage may be scanty, green, yellowish, or of a slightly reddish tinge. As in the case of budworm control it is highly desirable that as complete a patrol as possible of the forest region be maintained in order to locate incipient outbreaks.

There are two factors other than defoliation which may be the primary cause of bark-beetle outbreaks starting up. The first of these is a ground fire which scorches the base of the trees. The second is the accumulation of large amounts of green slash whether due to logging operations or windfall. Under certain conditions this accumulation of green slash may be a decided menace to the surrounding timber, not only due to the fact that it may serve as a breeding place for the beetles, but also due to the fact that such an area serves to attract large numbers of the beetles.

Climatic conditions play an important role in influencing the increase or decrease of bark-beetles. It has been found that the beetles breed most rapidly in warm weather when there is a moderate supply of moisture, but that their increase is retarded by periods of cold wet weather. The beetles are easily dried out by sunlight and wind striking the trunks of the brood trees.

WOOD BORERS

Considerable damage is done to unpeeled long lumber and pulp by wood-boring insects which attack the wood during the summer months. In like manner trees that have been injured by fire are severely injured by these borers. There are two principal types attacking spruce and fir.

ROUND-HEADED BORERS

Four species of *Monochamus* are commonly found boring in spruce and fir. These are large beetles with long antennae from which they get the common name "long horned beetles". Eggs are laid during the warm months in the bark of trees or logs which are in the sunlight. These eggs hatch into white colored grubs which begin boring beneath the bark and later enter the sapwood making tunnels of about the size of a lead pencil.

Monochamus scutellatus is black and approximately one inch in length with antennae of male about twice the length of the body. It is common in June and July.

Monochamus notatus is brown and of approximately the same size as *M. scutellatus*. It is found in June and July.

Monochamus titillator is a large brown mottled gray beetle with a body length of about one and one-half inches. It is most commonly found in the early fall.

Monochamus marmorator is a mottled gray beetle of about the size of *M. scutellatus*. This species attacks both green and dying fir. The adult is found active in the evening. It is common in June and July.

FLAT-HEADED BORERS

There are numerous species of flat-headed borers belonging to the family *Buprestidae* which cause considerable damage to dying and cut timber or pulp. As in the

case of the round-headed borers these beetles lay their eggs in the bark. The larvae, which are elongate and somewhat flattened, start boring beneath the bark, later going into the sapwood.

Control. Summer logging should be planned so that the logs are either peeled or else got out of the woods as rapidly as possible. Floating them will prevent injury. The round-headed borers attack principally logs lying in the sunlight so that some success has been obtained by keeping the logs in the shade. Craighead has made considerable headway in developing a cheap repellant spray which could be used on piled pulp or logs. The work, however, has not been completed. Dilutions of creosote oil containing up to eight parts of kerosene proved most effective. It is planned this summer to carry this work to completion in Maine. The normal loss from borers to piled unpeeled pulp amounts to nearly ten per cent the first year. The loss to logs cut for lumber purposes is much greater, and very commonly logs left lying in the open during the summer months are entirely ruined due to the worm holes.

White Pine Insects

INTRODUCTION

The changes in composition which have taken place in our New England forests during the last one hundred years are portentous. White pine which at one time seldom occurred in pure stands of any size now occurs in large bodies throughout central New England. This means that what at one time was a mixed forest crop has now become a specialized crop. Insects which at one time had only individual trees of the same species scattered here and there through the forest to feed upon, now have large forests of the same species to satisfy their greed. This fact alone accounts for the great increase of certain obnoxious forest insects. We find a striking analogy in the agriculture of the country. As man specializes more and more, insects which at one time had only small individual plots of a given food to feed upon, now find this same food in great abundance over large areas. No one factor so influences the rise and fall in abundance of an insect pest as the food supply.

The increase in abundance of white pine has hardly been able to keep up with the increased demand for it. From the earliest days of lumbering, the white pine has been the favorite tree in the North-east. In the great spruce regions of Maine it was the first tree to be cut and in these areas was so thoroughly culled out that it has not returned as a species of any importance in the new crop, the lumbermen falling back first upon the spruce, then the fir.

In central New England, however, the pine has seeded into the many abandoned pastures and at present forms the principal forest crop of this region. This so-called second growth white pine is by no means of as good

PLATE IV. WHITE PINE INSECTS.

A. Adult Roundheaded Borer. (Monochamus). (Natural size). B. Larva Roundheaded Borer. (Natural size). C. White Grub. (Natural size). D. Adult Flatheaded Borer. (Buprestid). (Natural size). E. Larva Flatheaded Borer. (Natural size). F. Adult White Pine Weevil. (Natural size). G. Adult Barkbeetle. (Greatly enlarged).

quality as that occurring in the original forest, but an ever increasing demand for this low grade pine has made it a tree of very great value. The realization that the pine will not last forever has resulted in large numbers of plantations of white pine being set out in an effort to keep the crop on a sustained yield basis. With the increase of the areas set out to white pine, has come a like increase in the prevalence of the insects attacking the pine. In some localities the apparent damage from insects has been so great that a decided damper has been put upon the interest in planting this species. This is decidedly unfortunate in view of the large program of reforestation which has so carefully been built up. It is highly important then to know in advance the extent to which existing planting plans and methods of management should be modified to meet these enemies. Enough white pine has been planted now in different localities throughout this region to make it reasonably safe to predict just what is to be expected in the way of insect damage.

INSECTS ATTACKING SEEDLINGS

WHITE GRUBS

Practically all injury likely to result from insect attacks on the seedlings can be circumscribed by proper planting methods. First in importance and probably the most common injury is that resulting from white grubs (*Phyllophaga*) which are well known to most nurserymen and farmers. These grubs are in reality the young of June bugs. Normally they feed upon the roots of grass and it is seldom that they become a serious pest to the agriculturist except when grass land is turned over and planted. As this is a very common practice the grubs are consequently often a serious menace.

The possible damage can usually be greatly minimized by cultural methods such as fall plowing or by planting the area to corn or clover the first year. Pasturing hogs

in the area is often resorted to. J. F. Illingsworth, an Australian entomologist, has recently obtained remarkable results by the use of crude white arsenic, applying eighty pounds per acre. This method of control is based upon the fact that in feeding a large amount of soil passes through the intestinal tract of the grub. By mixing the arsenic with the soil, when preparing the seed bed, almost perfect control may be expected. No detrimental effect upon plant growth was noticed even when the arsenic was used in excessive amounts.

PALES WEEVIL

It is practically useless to start a coniferous nursery on a white pine area that has just been cut or on a plot that is contiguous to such an area. The pales weevil, (*Hylobius pales*) which is a snout beetle, is attracted in large numbers to areas where freshly cut pine stumps, logs, boards, or even slash occur. These beetles are ravenous feeders, eating the tender bark off from pine seedlings and invariably proving a serious pest on such areas, often entirely destroying the nursery. It is not safe to plant in such an area until the third year after cutting. This statement also holds for planting of pine on cut-over pine lands during the first two years following the cutting. Such plantations are invariably a failure. The beetles breed in the fresh pine stumps and logs.

The damage may be materially mitigated by burning the slash over the stumps and utilizing the logs in the early spring, but this will not entirely prevent injury. The age of seedlings killed ranges from one to five years. Year old seedlings are seldom attacked, whereas those over five years of age are usually of sufficient size to overcome the damage.

STRAWBERRY CROWN GIRDLER

In 1915 the strawberry crown girdler (*Brachyrhinus [Otiorrhynchus] ovatus L.*) caused a loss of over \$15,000 in one Massachusetts nursery to white pine seedlings. The

PLATE V. WHITE PINE INSECTS.

A. Leader killed by White Pine Weevil showing characteristic drooping of current year's growth, and pupal chambers with exit holes. ($\frac{1}{3}$ natural size). B. Larva of Leconte's sawfly. C. Work of Pine Tube Moth. ($\frac{1}{3}$ natural size). D. Adult Pales Weevil. (Enlarged 2 x). E. Work of Pales Weevil on pine seedling, showing girdling effect. ($\frac{1}{3}$ natural size).

larvae, or small white grubs, girdle the stems and the roots at from one to three inches below the surface of the ground. There is a large range of food plants including red and Scotch pine and several species of spruce. The larvae pupate in the ground during early June at a depth of from one and one-half to three inches and may be destroyed at this time by plowing and harrowing. The adults can be trapped under boards, weed piles, burlap, etc., particularly during hot dry weather. They are active only at night. The larvae live for from ten months to one year. Fortunately it is seldom that this insect proves troublesome. At times the seedlings may be attacked by foliage eating insects, but these are easily controlled by using a lead arsenate spray.

INSECTS ATTACKING SAPLING PINE

WHITE PINE WEEVIL

Undoubtedly the most serious enemy of the white pine is the white pine weevil (*Pissodes strobi*), but extensive experiments carried on by the writer have shown that even the ravages of this destructive pest are easily overcome by proper forest management. This beetle lays its eggs in the terminal shoots of the white pine. These eggs hatch into small white grubs which immediately begin feeding just beneath the bark on the wood of the shoot. As the grubs increase in size they feed deeper into the wood, all of the time working downwards. Needless to say the leader is soon girdled and immediately begins to wither up and die. Often the grubs will continue feeding in the main stem until they have passed the first, and at times even the second and third, whorl of laterals.

The results of this destruction of the main shoot are three fold. In the first place it results in crooked, forked, many branched trees. Secondly it materially lengthens the time of rotation for the crop. In the third place injured trees are very apt to have large knots

PLATE VI. WHITE PINE INSECTS.

A. & B. Injury to pine caused by Mound Building Ants, showing constriction on main stem. C. Queen of *Formica exsectoides* (greatly enlarged). D. Pupal case of Mound Building Ant from which adult has emerged. (Greatly enlarged). E. Larva of Mound Building Ant. (Greatly enlarged).

and short internodes which lower the grade and value of the lumber. The method of control is based upon the fact that in a reasonably dense stand the trees do not have the opportunity to fork and branch out. The natural closing in of the stand does not allow room for more than one terminal shoot, and any others are soon crowded and shaded out. If then a stand of young pine can be started with a density of from 1200 to 2000 trees per acre, on reasonably good pine soil, the trees, although badly weeviled, will overcome the injury. As the majority of white pine plantations are set out using a six by six foot spacing, or 1210 trees per acre, the resulting stand will be of good quality in so far as lack of weevil injury is concerned.

Much of the very serious damage resulting from weevil injury in plantations is due to poor planting methods. The weevil attacks trees between the height of two and one-half and fifteen feet, seldom doing great damage above this height. It is patent then that the sooner the trees reach this height the less damaged they will be, for the trees may be attacked by the weevil each year. Trees planted on poor soil such as sandy ridges or swamp lands are naturally slow growing and may be weeviled ten or fifteen times before they even reach a height of ten feet. The internodes, or distances between the whorls of branches, are so short that in order to mature the grubs may destroy several years' growth during a single season. Such trees are almost invariably knotty and crooked regardless of the density of the stand. Poor stock used in planting or careless planting methods which result in a number of the trees dying leaving open spaces in the stand, allow the weeviled trees to fork and branch out. Such spaces should be replanted. On the average quality II site the mature stand of from fifty to sixty years should contain from 250 to 300 trees per acre.

MOUND BUILDING ANTS

Another insect of considerable importance which attacks sapling pine is the mound building ant, (*Formica*

exsectoides). These ants destroy all the vegetation around their nests for distances up to a twenty foot radius and kill young pine trees until they reach a height of six feet. The ants use a unique method in killing the trees. Formic

PLATE VII. Mounds of *Formica exsectoides* showing barren area around them.

acid is injected by them into the tissues of the main stem a short distance above the ground. The acid coagulates the cell contents, thus preventing the downward flow of the sap. The lesion appears superficially like a fungus canker. It is useless to plant pine within fifteen feet of these mounds, which are often three feet in height and six or more feet in diameter.

These ant colonies are not easily destroyed as the queens, or mothers, live deep down in the nest often five or six feet below the surface of the ground, and unless these are killed the colony will continue to live. The use of carbon bisulfide has, however, proved very satisfactory in destroying the colonies. From one to one and one-half pounds of the bisulfide should be poured into a shallow dish which is then placed on top of the nest. A few holes should then be punched in the mound with a stick and a large pan inverted over the nest in order to prevent the fumes from escaping. A little fresh earth or a few sods thrown around the base of the

pan will aid materially in keeping the gas within the nest. Carbon bisulfide is sold on the market in liquid form. On exposure to the air the liquid soon volatilizes to a gas of disagreeable odor. This gas, which is 2.68 times as heavy as air, penetrates deep into the recesses of the nest, killing the ants. The smaller the nests, the less bisulfide is necessary, and the easier it is to destroy the colony.

WHITE PINE APHID

Often times large colonies of black aphids, or plant lice, are found on the needles and small twigs of pine. These are commonly known as the white pine aphid (*Lachnus strobi*) and are often the cause of the browning of the needles and the final death of the trees. These little plant lice occurring as they do in large numbers take an immense amount of vitality from the trees attacked. They increase in numbers very rapidly during the year, there being several generations during the season, so that the resulting off-spring from a single female is enormous, it being estimated at one billion as the least possible number if all should live.

Fortunately these aphids are very frail, weak-bodied insects and are destroyed in large numbers by rains, and other agencies. It is quite common to find them in company with the mound building ants. In plantations they may be destroyed by spraying the trees with 40 per cent nicotine sulfate, used at a dilution of 1 part nicotine to 800 parts of water. It is seldom that large trees are damaged and even in the plantations they usually occur only on individual trees so that it is seldom necessary to spray extensively.

GIPSY MOTH

A large plantation of white pine in southern Maine was attacked this last summer by first stage gipsy moth larvae (*Porthetria dispar*) and would have been entirely

destroyed had it not been immediately sprayed. The caterpillars, which had been blown in great numbers on to the plantation from surrounding birches, were feeding on the epidermis of the main stem and laterals of the young pine, entirely girdling them. The small brown caterpillars are covered with both long and short hairs which aid materially in making them buoyant. The spinning of a long thread, as is the case with spiders, adds to their buoyancy. Records taken by Collins show that the larvae may be blown for several miles. The feeding of the larvae on the epidermis of the stems of seedlings is, however, unusual. The plantation was sprayed with hand sprayers using arsenate of lead at the rate of three pounds to fifty gallons of water. The more mature stages of the gipsy moth larvae feed to a considerable extent on the foliage of the white pine.

PINE TUBE MOTH

An interesting and rather unusual insect outbreak occurred this last summer in a large white pine plantation in New Haven, Conn. The insect causing the damage was the pine tube moth (*Eulia pinatubana*) which is quite common in Maine, but rarely causes any serious damage. In the fall the pale green larvae gather a bundle of about fifteen needles together into a tube within which the caterpillar lives, feeding on the ends of the needles. In October 1922, when the above plantation was visited, the tops of a large number of the trees had been entirely defoliated. Fall spraying is the only known method of controlling an outbreak, but fortunately this is seldom necessary.

MISCELLANEOUS SPECIES

There are several other insects, usually of minor importance, which attack sapling pines. These should be mentioned, although it is seldom that they will be found in large numbers. First in importance are the

larvae of saw-flies. Superficially these look like hairless caterpillars and are usually of a uniform color, either yellow or green, feeding entirely upon the foliage or needles. They can consequently be controlled by using a stomach poison such as arsenate of lead, two pounds being used to fifty gallons of water. In controlling such an outbreak trees should be sprayed, if possible, before the caterpillars become of any size. The European pine-shoot moth (*Evetria buoliana*) at times does considerable damage to sapling pines. The larvae work in the new buds and shoots, resulting in crooked trees. This insect is controlled by pruning off all infested shoots. Several species of spittle insects are found on pine. The most common species belong to the genus *Aphrophora*. These insects are found under masses of froth-like spittle which appear on the new growth twigs. Whatever damage which may occur is localized to the twig on which the insect is feeding and as these are sucking insects the injury is further localized to the cells within the twig. The pine leaf miner (*Paralechia pinifoliella*) which is the minute larvae of a small grayish brown moth, whose wing expanse is but three-eighths inch, at times becomes numerous enough to cause trees to turn brown. There are supposed to be several generations a year. The pine leaf scale (*Chionaspis pinifoliae*) at times becomes extremely numerous so that small trees may even appear white due to the large number of whitish scales occurring on the needles. This insect is best controlled by spraying the trees with Black Leaf Forty. It would, of course, only be in connection with transplants or ornamental trees that spraying would be advisable.

INSECTS ATTACKING MATURE TREES

Healthy mature white pine trees are seldom killed outright by insects. It is usually trees that have first been weakened by drought, fire, smoke, or over-maturity, which are seriously attacked and finally killed by insects. Weakened trees attract large numbers of in-

jurious insects which usually soon destroy them so that they are practically of no use for lumber purposes. The gipsy moth at times does considerable damage to healthy trees growing in mixture with hardwoods or in close proximity to them. At times saw-fly larvae may defoliate and kill large pine.

BARK BEETLES

Usually the first insects to attack the weakened pine are bark-beetles. These are small, oblong, hard-shelled insects usually from one-eighth to three-sixteenths inch in length. These beetles make tortuous galleries beneath the bark in which the eggs are laid. These eggs hatch into minute white grubs which burrow in all directions beneath the bark loosening it from the trees. There are several species commonly found attacking the pine in New England. *Ips pini* is probably the most common bark beetle found in pine. Three species of *Tomicus* are also commonly found attacking pine. These beetles multiply rapidly in spite of the fact that there is usually but one generation a year.

PINE BORERS

The greatest damage to the trees comes from wood-boring larvae. These are of two general types—the round-headed borers and the flat-headed borers. The first type are usually the larvae of *Monochamus* or the so-called round-headed borers. There are three species commonly found attacking the pine—one brown (*M. notatus*), one black (*M. scutellatus*), and one mottled gray (*M. titillator*), which are described under spruce and fir wood borers. These beetles particularly attack logs piled in the sun, or dying trees, and can be heard boring in the wood for quite a distance. The ribbed ragium (*Rhagium lineatum*) is also frequently found burrowing beneath the bark of white pine. The adults of the second type, or flat-headed borers, are usually broad,

elongate, rather flat bodied beetles, generally of a bronzed or metallic color. The larvae, or grubs, are much elongated and somewhat flattened. The head is small, although the body segments directly behind it are very broad, apparently forming part of the head. This gives them the name flat-headed or hammer-head borers. The most common species attacking pine are from one-half to one and one-half inches in length. The larvae start by tunneling directly beneath the bark, later boring down into the wood. They make broad, rather shallow burrows or tunnels which are filled with soft shaving like chips.

In the woods these borers are held in check somewhat by birds and parasitic hymenoptera. These latter are fly-like insects which have long ovipositors, or tail-like appendages, which they thrust through the bark of infested logs or trees in order to lay their eggs in the burrows of the wood-boring larvae. These eggs hatch into small grubs, which follow down the burrows and attack the larvae, boring into and eventually killing them. The parasites can usually tell fairly accurately just where the borers are working, probably due to the very perceptible rasping sound, caused by the chewing of the larvae. The eggs are consequently laid fairly close to the places where the borers are working, if not at times directly in the borers.

CARPENTER ANTS

Next in importance are the large black, or black and red, carpenter ants. These seldom attack a healthy tree, usually entering through decayed places or attacking felled trees. They are tireless workers and due to their great numbers soon honeycomb out a tree. It is quite common to find pine trees in the woods which have been hollowed out by these ants.

Control

All of the above wood-boring insects, including the bark beetles and carpenter ants, are best controlled by immediately felling those trees which show signs of dying, and utilizing the logs before the larvae have an opportunity to do their damage. It is bad policy to leave pine logs lying around for any length of time during the warm season before sawing. Conifers which show signs of dying seldom recover, due to the small amount of reserve food present. They are therefore very susceptible to adverse conditions. Dying trees left standing serve merely as breeding grounds for insects which are likely to spread into the surrounding healthy trees. If for any reason the dying trees can not be utilized, the mere felling and barking of the tree, combined if possible with burning of the slash, will aid materially in preventing possible outbreaks.

WOOLY APHID

There are several insects, usually of minor importance, which at times may become numerous enough to be a menace to the incipient forest. Among these the pine bark aphid, or wooly aphid (*Chermes pinicorticis*) is probably the most commonly found in the forest. This aphid appears as white tufts at the base of the needles on twigs, and as white flocculent patches on the bark of the trunks and branches. This insect undoubtedly checks the growth and lessens the vitality of the trees and at times may even kill them. In the forest it is questionable if it is ever necessary to carry on extensive control measures against this particular insect. The writer has, however, seen local infestations where it would have been advisable to cut out the infested trees and destroy them by burning, or else by utilizing the cut timber. Individual trees are at times sprayed with kerosene emulsion, which is very effective in controlling the aphids. Other insects which might be mentioned are white ants, several foliage feeders, and a few which feed upon the roots.

In spite of the somewhat formidable array of insect enemies no apprehension is felt by the writer as to the possible danger from planting white pine. It is true, however, that the planting of this species to the exclusion of other forest trees is highly undesirable from an entomological standpoint. A more extensive planting of red pine or spruce is worthy of serious consideration. There is a certain farming region in Massachusetts where very little white pine occurred. Twenty years ago large plantations were set out, and for years these escaped serious insect damage. Since that time an ever increasing number of plantations has been set out in this locality so that at the present time insects attacking the white pine have been attracted to this most favorable area in large numbers. The white pine weevil is particularly injurious, and has begun to attack even the older plantations.

SUMMARY

In concluding it might be well to summarize briefly just how plantations should be handled in order to prevent serious injury from insects. The seed beds should not be situated in close proximity to recently cut-over pine lands, that is within 200 yards of the nearest stumps, logs, or slash. If possible freshly plowed-under grass land should be avoided. It is best to use land that has previously been cultivated.

When setting out the plantation, cut-over pine lands should be avoided until the third year after cutting. Too much emphasis can not be laid upon the need of care in choosing the planting site and in the planting itself. There is no question but that weak trees are more susceptible to insect injury than vigorous growing ones. If a person is at all in doubt as to the proper site for planting, valuable information can be obtained by merely noting under what growing conditions the best stands of pine are found in that or similar localities. The fact that a white pine tree can be found in a swamp does

not signify that this is a favorable location for the tree, as some planters apparently believe. There is no question but that careful planting more than pays for the extra labor involved. Planting around ant mounds should either be avoided or else the colonies destroyed.

It is the writer's belief that in order to overcome weevil damage, the trees should not be spaced any further apart than six by six feet. The increased value of the resulting stand should more than pay for any increase in cost of planting. The six by six foot spacing is almost universally used now. In the maturing stand, dead or moribund trees should be cut out and destroyed, as these merely act as centers of infestations. It was previously suggested that some emphasis should be placed upon the planting of other species of conifers. This suggestion does not, however, mean that the planting of white pine should in any way be curtailed, for insects attacking red pine or spruce would also become obnoxious if these trees alone were planted in any quantity. As near as possible the natural balance found in nature should be kept. This does not mean the planting of mixed plantations, for these are not so economically handled. It does mean, however, that large areas should not be given over to a single species, as this merely invites insect damage.

Larch Insects

LARCH SAW-FLY

Between the years 1882-1885 the larch fly (*Nematus erichsonii*) destroyed practically all of the mature larch in the State. The destruction of the larch at this time is said to have been one of the prime factors bringing about the exodus of caribou from Maine into Canada, as these animals lived largely in the larch swamps feeding on the twigs and foliage of this tree. The saw-fly was again found active this summer in several localities where the larch is again becoming a dominant species making up, as it does in some parts of the State, fully one-quarter of the stand over wide areas. Several small outbreaks occurred during the summer of 1922.

The large black adult saw-fly lays its eggs during the latter part of June and the early part of July in the terminal shoots of larch. The larvae are found active during July and early August.

The full grown larvae are from one-half to three-quarters of an inch in length. They are easily recognized by the round jet-black head and the dull light green bodies. There are three pairs of black thoracic legs and seven pairs of short abdominal legs. As is the case with other saw-fly larvae they have the habit of curling up the head and rear end of the body into a "U" shape when disturbed.

LARCH CASE BEARER

Considerable attention was called to the browning of the larch, or tamarack, in Maine during 1922. In many localities large stands of larch appeared in July as if a fire had swept thru, whereas a few weeks previous they

had been green. In places where the larch occurred in large stands making up a high percentage of the forest whole hillsides and valleys looked brown.

Examination of the stands of larch have shown that the trees are being attacked by a comparatively new insect to this region—the larch case bearer (*Coleophora laricella*). This insect is very common throughout Europe where it is a serious pest. It was probably brought into this country on European larch, a tree that is quite widely planted in the northeastern United States.

There is but one generation of the insect a year. The small delicate gray moths started to emerge from the small cigar shaped cocoons about June twentieth. The moths, which are only about three-sixteenths inch in length, lay their eggs on the new needles and the young larvae upon hatching immediately begin to tunnel or mine within the needles, causing them to turn yellow and then brown starting from the tips. When very numerous practically all of the needles on a tree may be injured and in many cases destroyed, as in the present outbreak. As soon as the mine is long enough to form a case, a portion of the needle is cut off and the larvae proceed with this protection to another leaf. When attacking a needle the larva first cuts a circular hole thru which its head and later its body may extend for feeding purposes. The case, which at first is whitish, gradually becomes gray with age, and is lined by the larva with silk. Upon the approach of cold weather the larvae carrying their cases retire to a sheltered place on the branches or beneath bark and lichens becoming active early in the spring when they continue feeding.

There is at present every indication that considerable damage may be done to the larch, for the insect is becoming extremely abundant. It is hoped that this coming spring something can be done to check the outbreak.

Hardwood Insects

POPLAR OR ASPEN BORER

Trees of the genus *Populus*, which includes the aspen, popple, cottonwoods, poplars, and Balm of Gilead, are all subject to serious injury by wood-boring insects. As poplar is becoming an important pulp tree in Maine, even to the extent of being set out in plantations, some attention must be paid to holding these borers in check, for plantations are often completely destroyed.

Probably the most injurious of these borers is the so-called aspen borer (*Saperda calcarata*), which is very common in Maine. The adult is a large, robust, grayish green beetle, approximately one inch in length with light yellowish spots on its wing covers. During July and August the beetles gnaw oblong scars in the bark in which they lay their eggs. These hatch in about twenty days and the young larvae begin mining beneath the bark where they remain over winter. In the spring the larvae bore into the sap and heart wood. Here the larvae remain boring until the second season following egg laying. During June or July of the third year the adults emerge.

CONTROL

The insect is best controlled in the forest by locating and then cutting the so-called brood trees which have been attacked for a number of years in succession and usually contain several generations of the insects. These trees are easily distinguished by the dead tops or limbs, the deformed trunks, or by the frass and brown sap exuding from the egg scars. These brood trees should be cut and burned or else the main trunks split and exposed to the sun, which will dry out the wood thus killing the brood.

PLATE VIII. SHADE TREE INSECTS.

A. Larch Case Bearer, showing cases on needles. ($\frac{3}{4}$ natural size). B. Larva of Forest Tent Caterpillar. (Natural size). C. Larva of Elm Leaf Beetle. (Slightly enlarged). D. Elm Leaf Beetle. (Slightly enlarged). E. Adult Poplar Borer. (Natural size). F. Work of Oak Twig Pruner. Twig has been split so as to show burrow and end of twig that was severed. ($\frac{1}{2}$ Natural size). G. Adult Oak Twig Pruner. (Natural size). H. Adult Maple Borer. (Natural size). I. Work of spruce Cone Worm showing castings around base of cones.

MAPLE BORERS

Several wood boring insects do considerable damage to maples, particularly over mature trees. These trees should be cut and burned or utilized whenever possible, as should all trees heavily infested with borers.

BRONZE BIRCH BORER

There are large areas of white and yellow birch in the northern part of the State in which injury from the bronze birch borer (*Agilus anxius*) is everywhere very apparent. The somewhat flattened, bronze-colored beetles lay their eggs in June in crevices on rough surfaces of the bark. These hatch into flat creamy white grubs which bore thru the bark into the sap wood where zig zag tunnels are made. Damage shows up in the way of dead tops and in some areas of pure white birch hardly a tree has escaped injury. Poplar is also attacked. Trees occurring in mixture with softwoods are not so liable to injury.

FOREST TENT CATERPILLAR

Several thousand acres of poplar and white birch were stripped by the forest tent caterpillar (*Malacosoma disstria*) in 1922 in the region of Jackman, Lobster Lake, Chesuncook Lake, and Eagle Lake (T. 8, R. 12). Examinations of the stands in the fall showed that the insects had been heavily parasitised and it is believed that the outbreaks will not prove serious. In one locality the severity of the outbreak made it seem advisable to cut the stand and this was done as a protection to surrounding timber. The location of the stand made it possible to utilize the wood for pulp.

The brown eggs are laid in masses encircling the twigs. The caterpillars hatch out during April. When full grown they are from one and one-half to two inches in length. The head is blue and there is a row of white, diamond-shaped spots down the middle of the back. The larvae turn into moths during July. Unlike the apple tree tent caterpillar this species forms no tent.

POWDER POST BEETLES

Very frequently considerable damage is done by powder post beetles to hardwood material in storage. There are several injurious species but in Maine the European *Lyctus* (*Lyctus linearies*) is apparently the most common. This is a slender, somewhat flattened, rusty red brown beetle, from one-tenth to one-fifth inch in length. There are single rows of shallow punctures on each wing cover. The beetles lay their eggs in the sap wood of seasoned hardwood material of all kinds. Hickory, oak, and ash are particularly subject to injury but other woods are also attacked. The small, yellowish white larvae, upon hatching, immediately begin boring through the wood in all directions. When full grown the larvae are from one-eighth to one-fifth inch in length with three pairs of legs. The adults emerge in the early summer.

Frequently the inside of the wood is reduced to powder before exit holes are apparent. Usually damage first shows up in the way of small piles of fine powder around the stored products. At times furniture and even buildings have been known to crumble before the presence of the beetles has been realized.

It is well to bear in mind that heart wood material is never attacked. Sap wood material which has been seasoned for less than eight months is also immune from injury. Material in storage should be classified, keeping that which is most subject to injury by itself. This should be inspected at least once a year, preferably in the winter. Material which has been attacked by the beetles should be immediately destroyed. The oldest stock in storage should be utilized and sold first. Care should also be taken to inspect all new stock being brought into storage.

Shade Tree Insects

SPRUCE GALL APHID

The spruce gall aphid (*Chermes abietis*) causes considerable injury to Norway and white spruce used for ornamental purposes such as on lawns or as hedges. The galls usually occur near the base of the terminals and appear very much like small cones. In the fall these turn brown. The main injury comes from disfigurement to the trees, as the galls remain on the trees year after year. Infested trees should be sprayed in April, using a whale oil soap solution at a dilution of one pound of soap to two gallons of water.

SPRUCE CONE WORM

The spruce cone worm (*Dioryctria reniculella*) was very common in some sections of Maine in 1922. The amber brown larvae tunnel thru the new cones, often passing from one cone to another, and leaving unsightly masses of frass and castings around the cones. The larvae also feed on the foliage of terminal twigs, leaving masses of excrement and pitch on the sides of the twigs. When full grown the larvae are slightly over one-half inch in length. The moths are of a stone gray color. Infested cones should be collected and burned.

OAK TWIG PRUNER

Numerous inquiries have been received in regard to the damage caused by the oak and maple twig pruner, (*Elaphidion villosum*). This insect in its larval stage bores thru the twig working towards the base of the small limb or twig in which it is boring. Upon reaching matur-

ity in the fall, it gnaws a circular groove extending to the bark and then retreats back into the burrow plugging it with sawdust. The twig is usually broken off by the first high wind and if examined will give every appearance of having been severed with a knife. The larva transforms into a pupa and beetle within the fallen twig, emerging about the first of June. At times the insect may do considerable damage to shade trees, but is easily controlled by collecting and burning the fallen twigs either in the late fall or better still in the spring.

ELM LEAF BEETLE

The elm leaf beetle (*Galerucella luteola*) was apparently very prevalent in the cities and towns of southern and central Maine in 1922. The small beetles emerge from hibernation at about the time that the leaf buds are beginning to open. They immediately begin feeding on the tender leaves making small, irregular holes. Eggs are then laid on the lower surfaces of the leaves and the young hatching out in about six days begin skeletonizing the leaves. A second generation occurs in July. Most of the difficulty found in controlling this insect has been due to the fact that trees have not been sprayed at the right time. A difference of one week means the difference between success and failure. The trees should be sprayed just as the adults first begin feeding on the leaves, then again as the larvae are hatching, and a third spray may be necessary at the time that the second generation larvae are hatching. A fairly strong spray should be used, using four pounds of lead arsenate to fifty gallons of water. On the second and third sprays emphasis should be placed on spraying the under side of the leaves.

BORERS

A large number of inquiries have been received in regard to borers in shade trees—particularly maple, poplar, and elm. In general the treatment is similar for all. Wherever possible pruning and burning of

infested limbs is advisable, either in the late fall, early spring, or as soon as the injury shows up. In cases where the borers have entered the main stem or trunk, carbon bisulphid should be injected into the burrows, using from one to two teaspoonfuls to a hole and then plugging the hole in order to keep the gases from escaping. In some cases the larvae may be destroyed by forcing a flexible wire into the burrow. Painting freshly made egg scars with carbolineum is very effective.

LEAF SPOTS

Many different fungi attack the foliage of shade trees causing small brown spots, or at times the entire leaves to turn brown and fall off prematurely. As most of these fungi pass the winter on the fallen leaves, the diseases are easily checked by raking these up and burning.

Economic References

- Blackman, M. W. 1919. Report on the Spruce Bud Worm, Maine Forestry Dept.
- Craighead, F. C. 1922. Fighting Bark Beetles in Balsam. *Illus. Can. For. Mag.*, Vol. 18, No. 6, pp. 820-821.
- Craighead, F. C. 1922. Combatting Budworm Attack. *Illus. Can. For. Mag.*, Vol. 18, No. 8, p. 942.
- Felt, E. P. 1906. Insects Affecting Park and Woodland Trees. *Mem. New York State Mus.*, No. 8, Vol. I, pp. 59, 170-171, 189.
- Hofer, G. 1920. The Aspen Borer and How to Control It. *U. S. D. A. Farmers' Bul.* 1154.
- Hopkins, A. D. 1909. Bark Beetles of the Genus *Dendroctonus*. *U. S. D. A. Bur. of Ent. Bul.* 83. Part 1.
- Hopkins, A. D. & Snyder, T. E., 1917. Powder Post Damage by *Lyctus* Beetles to Seasoned Hardwood. *U. S. D. A. Farmers' Bul.* 778.

- Johannsen, D. A. 1913. Spruce Budworm & Spruce Leaf Miners. Maine Agr. Exp. Sta. Bul. No. 210.
- Packard, A. S. 1890. Insects Injurious to Forest and Shade Trees. Fifth Rept. U. S. Ent. Comm., pp. 791-792, 811-822, 830-839, 854-856, 879-890.
- Peirson, H. B. 1920. An Infestation of the White Pine Aphid. *Psyche*, Vol. 27, Nos. 2-3, pp. 62-63.
- Peirson, H. B. 1921. The Life History and Control of the Pales Weevil. Harvard Forest Bul. No. 3, Petersham, Mass.
- Peirson, H. B. 1922. Control of the White Pine Weevil by Forest Management. Harvard Forest Bul. No. 5, Petersham, Mass.
- Peirson, H. B. 1922. Mound Building Ants In Forest Plantations. *Journ. of Forestry*, Vol. 20, No. 4, April.
- Peirson, H. B. 1922. Forest Insects in the Northeast, *Amer. For. Mag.*, Vol. 28, No. 346, pp. 628-634.
- Peirson, H. B. 1922. Larch in Maine Threatened By Insect Depredation, *Amer. For. Mag.*, Vol. 28, No. 347, p. 666.
- Schierbeck, O. 1922. Treatise on the Spruce Bud Worm, Bark-Beetle, and Borer, Montreal.
- Swaine, J. M. 1918. Canadian Bark-Beetles. Part 2, Ent. Branch Dept. of Agr., Tech. Bul. No. 14, Ottawa, Can.
- Swaine, J. M. 1922. Insect Armies that Follow in the Wake of Fire. *Illus. Can. For. Mag.*, p. 666.

CONNECTICUT AGRICULTURAL EXPERIMENT STATION

NEW HAVEN, CONN.

BULLETIN 155, MAY, 1907.

ENTOMOLOGICAL SERIES, No. 14.

The Elm Leaf Beetle

CONTENTS.

	Page.
The elm leaf beetle	3
History and distribution in America	3
Life history and habits	5
Description	7
Effect upon the trees	8
Food plants	8
Number of generations	9
Natural enemies	9
Remedies	10
Spraying with poison	10
Destroying the pupae	12
Destroying the adult beetles	12
Outfit for spraying elm trees	12
Summary	13

The Bulletins of this Station are mailed free to citizens of Connecticut who apply for them, and to others as far as the editions permit.

CONNECTICUT AGRICULTURAL EXPERIMENT STATION.

OFFICERS AND STAFF.

BOARD OF CONTROL.

His Excellency, ROLLIN S. WOODRUFF, <i>Ex officio</i> , President.	
PROF. H. W. CONN	Middletown.
PROF. W. H. BREWER, <i>Secretary</i>	New Haven.
B. W. COLLINS	Meriden.
CHARLES M. JARVIS	Berlin.
EDWIN HOYT	New Canaan.
J. H. WEBB	Hamden.
E. H. JENKINS, <i>Director and Treasurer</i>	New Haven.

STATION STAFF.

Chemists.

Analytical Laboratory.

JOHN P. STREET, M.Sc., *Chemist in charge.*

E. MONROE BAILEY, M.S.

E. J. SHANLEY, Ph.B.

KATE G. BARBER, Ph.D., *Microscopist.*

Laboratory for the Study of Proteids.

T. B. OSBORNE, Ph.D., *Chemist in Charge.* C. A. BRAUTLECHT, Ph.B.

Botanist.

G. P. CLINTON, S.D.

Entomologist.

W. E. BRITTON, Ph.D.

Assistant in Entomology.

B. H. WALDEN, B.Agr.

Forester.

AUSTIN F. HAWES, M.F.

Agronomist.

EDWARD M. EAST, M.S.

Stenographers and Clerks.

MISS V. E. COLE.

MISS L. M. BRAUTLECHT.

MISS E. B. WHITTLESEY.

In charge of Buildings and Grounds.

WILLIAM VEITCH.

Laboratory Helper.

HUGO LANGE.

Sampling Agent.

V. L. CHURCHILL, New Haven.

The Elm Leaf Beetle.

BY

W. E. BRITTON,

State Entomologist.

Twelve years ago an account of this insect was published by this Station in Bulletin No. 121, which has for some time been out of print. Since then only a few scattered notes have appeared in the bulletins and reports. The object of the present paper is to place in the hands of the people of Connecticut a fairly complete and comprehensive account of the elm leaf beetle with up-to-date remedies and methods of treatment. There is a constant demand for such information from public school teachers, members of village improvement societies, and men employed in the street and park departments of our cities and towns, as well as from private individuals who desire to give the best care to trees on their home grounds.

HISTORY AND DISTRIBUTION IN AMERICA.

The elm leaf beetle was introduced into this country probably more than seventy years ago. In its native country, Europe, where it had long been known, it had from time to time caused serious injury in Italy, Austria, and the southern portions of France and Germany. In Northern Europe the insect occurs sparingly, but can hardly be called a pest. Harris states* that the elm leaf beetle attacked and seriously injured the elm trees of Baltimore, Md. in 1838 and 1839. The beetle seemed to spread chiefly northward, though slowly, until Southern New England was reached in the early nineties, and much damage done. In the coast towns of Connecticut many fine old elms, including some historic trees, were killed by its depredations.

* Insects Injurious to Vegetation, page 124.

Stamford, Norwalk, Bridgeport, Stratford, Milford and New Haven especially lost many noble trees. Later the inland cities were invaded, and the elm trees ravaged. In New Haven the pest was perhaps at its worst in 1895 and 1896. In 1896 many of the trees on the older streets about the center of the city were sprayed with poison by the street department. The following season the pest was less serious, and continued to subside until 1901, when it was again comparatively destructive. From

FIG. 1.—Elm leaves showing larvae and the damage which they do by feeding on the under surface, natural size.

1902 it diminished in abundance until 1906, when considerable damage was done to the trees.

So far as is known the distribution of this insect in America is confined chiefly to the lower altitudes of Southern New England and the Alleghanian region. From Charlotte, N. C., its southern limit, the elm leaf beetle now extends as far north as North Conway, New Hampshire. Up to this time, however, the beetle has done no particular damage in New Hampshire. Regions generally infested include the whole of Massachusetts,

Rhode Island, Connecticut, southeastern New York, New Jersey, eastern Pennsylvania, Delaware, Maryland and a portion of Kentucky, though isolated outbreaks have occurred in western New York, Pennsylvania, West Virginia, Ohio and North Carolina. It is of course found in Virginia, Vermont, New Hampshire, and probably in Maine. Kentucky, therefore, contains the western limit of the distribution of this insect, though we may expect that soon adjoining states may become infested. The insect exhibits a marked tendency to spread farther along river valleys than over mountains, and is distinctly a pest of city and village trees rather than of trees in the open fields and roadsides of the country.

FIG. 2.—Cluster of eggs, greatly enlarged.

LIFE HISTORY AND HABITS.

The overwintering beetles come out of their winter quarters during the warm days of early spring, mate, and as soon as the leaves unfold they begin to eat small round or oval-shaped holes through them. Many leaves are thus riddled as though shot had been sent through them, and appear like the illustration on the front page of this bulletin.

During the latter part of May and early in June the females deposit small clusters of yellow eggs on the under sides of the leaves. The period of ovipositing extends over about four weeks, and each female may lay five or six hundred eggs. In

about a week the eggs hatch and the young larvae or grubs feed upon the under surface of the leaves, eating off the green tissue between the veins and leaving the veins and the upper epidermis, as is shown in Fig. 1.

In about three weeks the larvae or grubs are fully grown, and crawl down the trunks of the trees or drop from the ends of the branches to the ground and transform to the naked pupa stage. The great proportion of the pupae are found close around the base of the tree or lodged in the crevices of the rough bark of the trunk and larger branches. Except for being

FIG. 3.—Larvae and pupae, twice natural size.

in crevices, they are unprotected. The writer has seen trees in New Haven where it would be possible to gather several quarts of these pupae at the base of a single tree.

The pupa stage lasts about ten days, then the adult beetles appear, and lay eggs for the second generation, which seldom does much harm in Connecticut. Those emerging late probably do not lay eggs for a second brood, but may be seen crawling and flying about for a time, feeding more or less, but early going into winter quarters, usually in church belfries, attics of houses, barns, sheds or other out-buildings. They also pass the winter in cracks of fences, telephone poles, or under the edges of the loose bark of the trees. In some of the cities worst infested the adults sometimes gather in church belfries in such numbers that they can be swept up by the half

bushel. The elm leaf beetle often occurs with the two-spotted lady beetle in dwellings, simply because they both seek the same kind of a place for hibernation. Correspondents frequently send both species to the writer and desire to know if they are not in some way responsible for the injuries to their carpets. There is, of course, no relationship or similarity in food habits of the three species. The lady beetle is predatory, and in the larval stage destroys numbers of plant lice, and therefore should never be destroyed. The elm leaf beetles should, of course, be killed wherever they are found.

DESCRIPTION.

The *eggs* are bright yellow in color, bottle-shaped, and resemble the eggs of the Colorado potato beetle, but are smaller. They are fastened vertically to the under side of the leaf in clusters of from five to twenty-five arranged in two or three irregular rows.

When first hatched, the *larva* is dark or nearly black, covered with tubercles bearing black hairs. As the larva increases in size it molts several times and on becoming full-grown is about one-half inch long, dull yellow in color, with a pair of longitudinal black stripes along the back. Head, legs, lateral tubercles and two rows of small tubercles between the dorsal stripes are black. The tubercles also bear black hairs.

The *pupa* is about one-fourth inch in length and bright orange yellow in color, with black hairs or spines. It is not enclosed in an earthen shell to protect it, but is found at the base of the tree perfectly naked and wholly unprotected.

FIG. 4.—Adult beetles, twice natural size.

The *adult beetle* is light yellow in color when it first emerges, but soon takes on a duller hue, and finally becomes a dull olive green. An indistinct black stripe extends from the base to the extremity of each wing cover just inside of the margin. Small black spots

or markings on the pronotum of the thorax vary greatly in size and shape. Legs and antennae are yellow.

EFFECT UPON THE TREES.

It has previously been mentioned that the adult beetles do more or less feeding, always eating holes entirely through the leaves, as is shown on cover of this bulletin. This of course injures the tree, but is much less serious than the damage caused by the larvae, which eat away the under surface of the leaves. The larvae are always more abundant than the adult beetles, and are more voracious in their feeding habits. The worst infested trees usually drop their leaves in Connecticut about the middle of July. If this happens, and is followed by a rainy season, new leaves will be put out, but in a season of a protracted drought the trees may fail to put forth new leaves. In either case the tree is undoubtedly weakened, and often seriously so. Sometimes the second crop of leaves is devoured by the second generation of beetles, but in Connecticut the white fungus mentioned in another part of this bulletin is apt to serve as an important check to the beetle in a wet season. Two complete defoliations, one succeeding the other, usually kill a tree. Usually, however, the defoliation is not quite complete, and the trees continue to exist in a greatly weakened and devitalized condition. In the cities and larger towns, on account of further injuries* by horses, by leaky gas pipes in the ground, and pavements which cut off the supply of moisture, many trees have died. In 1901 the writer was called to Norwich, where nearly every elm had died for a distance of about one and one-half miles on a prominent residential street. These trees had been weakened by the constant attacks of the beetle year after year, and a leaky gas main finally destroyed what little vitality remained. Most of the maple trees along the street survived.

In many cases young or newly planted trees seem to be especially subject to attack, and therefore should receive extra attention.

FOOD PLANTS.

Elms constitute the only food plants known for this insect, and the European species suffer more than the American ones. The English elm (*Ulmus campestris*) and its weeping variety

* For a discussion of these injuries the reader should consult Bulletin 131 of this Station.

known as the Camperdown elm are favorites of the beetle. The writer has seen these trees entirely defoliated in New Haven when the common white, or American elms, were uninjured. The American elm is, however, the next choice, followed by the Scotch elm (*U. montana*), and though no variety is wholly exempt from attack, the winged elm (*U. alata*), the slippery elm (*U. fulva*), the cork elm (*U. suberosa*) and the rock elm (*U. racemosa*) are much less frequently attacked.

NUMBER OF GENERATIONS.

According to Burgess,* "in New Jersey, Professor J. E. Smith has recorded only a single brood and sometimes a partial second brood, while in the latitude of Washington, D. C., according to the observations of Messrs. Riley, Howard and Marlatt, of the Division of Entomology, two annual broods and sometimes a partial third brood have been found."

Dr. E. P. Felt, State entomologist, of Albany, N. Y., finds† two well marked broods and a partial third brood at Albany and Troy, N. Y.

The writer has not followed out this matter carefully in Connecticut, but all stages are found on the trees during the first half of September, so presumably there are at least two broods, but the egg-laying period of the adults is so prolonged that the lines of demarcation are nearly obliterated. In Connecticut the leaves of the trees severely attacked by the first brood generally turn brown and drop about the middle of July, when the larvae are descending the trees to pupate.

NATURAL ENEMIES.

One of the most important natural enemies of the elm leaf beetle in Connecticut is a fungus known to botanists as *Sporotrichum globuliferum* Speg. (*S. entomophilum* Peck), which attacks the pupae and adults in late summer, especially in a moist season. In 1902 this fungus was prevalent, and the following season the elm leaf beetle did little damage to the trees. In 1906 the beetles were abundant, but as the month of July was

* Bulletin No. 4, page 17, Ohio Dept. of Agriculture, Div. of Nursery and Orchard Inspection. 1905.

† Bulletin No. 57, N. Y. State Museum, p. 14, 1902.

wet, the fungus developed and killed a great many beetles. Beetles attacked by this fungus are covered with white mold, as shown in Fig. 5.

Predatory bugs of at least three species of the genus *Podisus* feed upon the larvae and pupae, and Riley* records three species of beetles that also devour full-grown larvae and pupae. The praying mantis (*Stagmomantis carolina* Linn.) is also an enemy of the elm leaf beetle in the southern portion of its range.

FIG. 5.—Pupae and adults killed by fungus. Healthy specimens at the left, natural size.

REMEDIES.

Spraying with poison. Covering the foliage with some arsenical poison is the only sure means of preventing injury to the trees, and for this purpose arsenate of lead is unquestionably the most satisfactory of these poisons. It remains better in suspension and adheres to the foliage longer than Paris green or London purple, and is less liable to injure it. As there are now several brands of good arsenate of lead on the market, it

* Div. of Entomology, U. S. Dept. of Agriculture, Bulletins No. 6, p. 10 and No. 10, p. 13.

will no longer pay to make it up each time from lead acetate and arsenate of soda. The arsenate of lead made by the Merri-mac Chemical Co., Boston, Mass.; Schonmaker & Son, Cedar Hill-on-Hudson, N. Y., and the Bowker Insecticide Co., Boston, Mass., (Disparene) have all been used in our tests and have given satisfaction. Arsenate of lead should be used in the following proportions:—

Formula	Arsenate of lead	3-5 lbs.
	Water	50 gals.

There are two methods of spraying elms: (1) To spray as soon as the leaves unfold, the treatment being aimed especially at the parent beetles, and to forestall all injury. As growth takes place, new leaves are constantly appearing, and these will not carry poison unless the application is from time to time repeated. This should be done often enough to keep the foliage well coated with poison until July 1st, when growth usually ceases, and both sides of the leaves should be coated.

(2) The other method is to spray the under sides of the leaves very thoroughly about June 1st, or soon after the eggs begin to hatch. This treatment is aimed at the larvae, and sometimes a single spraying is sufficient, as the poison will remain throughout the season. By it the great bulk of damage will be prevented, but the leaves will show the small holes made by the parent beetles before the poison was applied.

If arsenate of lead cannot be procured and it seems desirable to use other poisons, Paris green can be substituted.

This should be used at the following rate:

Formula	Paris Green.....	1 lb.
	Fresh lime.....	3 lbs.
	Water	50 gals.

Paris green is quicker in its action upon the insects than arsenate of lead, but will not remain as long upon the trees. Unless the lime is added, there is danger of "burning" the leaves. Lime is not needed with arsenate of lead.

The cost of spraying elm trees will vary from ten cents each in case of small trees to five dollars or more for the largest trees, according to the price of labor and the efficiency of the outfit.

Destroying the pupae. If the trees have not been protected by spraying, and have been attacked and injured by a horde of beetles and their larvae, it is always advisable to destroy the insects in the pupa stage at the base of the trees, in order to reduce the crop for next year as much as possible. These can often be swept up in large quantities; they can be killed by sprinkling them with hot water; but best of all is to spray the ground not only close to the tree but as far away as the branches reach, and also spray the bark of the trunk and large branches, with kerosene emulsion or some other contact insecticide. The pupae are easy to kill, but as they remain in the pupa stage only about ten days, it is essential that this work be done at the right time, and it is only when we observe that most of the larvae descending the trunk have transformed to bright yellow pupae that we can know when is the proper time to act.

Destroying the adult beetles. The beetles should of course be destroyed in attics, belfries and other places where they hibernate. This can best be done by sweeping them up before they become very active in spring and dropping them into the fire, hot water or kerosene to kill them.

OUTFIT FOR SPRAYING ELM TREES.

The barrel hand-power pump can be used for spraying small trees or for a limited number of large trees, but if one expects to make a business of spraying street trees, it will pay to procure a power sprayer; this may be a steam or gasoline engine with pump, or what is perhaps better, the "Niagara Gas Sprayer," which utilizes cylinders of carbonic acid gas to furnish pressure, thus doing away with a pump. Each large village and city should have at least one power sprayer that can be put to immediate use for spraying street or park trees. Strong one-half inch hose should be provided in long lines of from fifty to two hundred feet, and from four to six of these can be attached to each power sprayer. Even a larger number might be attached, but while in operation some workmen would be in the way of others, so that nothing would be gained. Mr. H. L. Frost of Boston, who is in the spraying business, informed the writer that four lines of hose is about the most economical number for each outfit. Each hose is furnished with an extension rod with closing valve at the lower end

and a cluster of nozzles at the upper end. These rods should be of different lengths, from four to twelve or fourteen feet for the different kinds of work. Any of the standard nozzles such as "Vermorel," "Mistry," or "Spraymotor" will be found satisfactory, and if several are used together in a cluster one is able to cover more leaf surface in a given time.

For getting about in tall trees it will be necessary for the men to be provided with climbing irons and extension ladders.

SUMMARY.

The elm leaf beetle was introduced into this country from Europe about seventy years ago, and caused serious injury to trees at Baltimore, Md., in 1838 and 1839. From this point it spread slowly, chiefly to the northward, reaching Connecticut in the early nineties, and injuring and killing many fine old trees in the coast towns. Later, inland towns were attacked, and the trees ravaged. The insect is now found from Charlotte, N. C., as far north as North Conway, N. H., and as far west as central Kentucky, but has not proven destructive to elm trees north of Massachusetts. It is preëminently a pest of shade trees in cities and towns, and seldom injures trees in the open fields. Since 1896 the attacks have diminished, but the pest was again serious in 1906.

The winter is passed by the adult beetles in attics, belfries and cracks in fences, and they come forth in April, and later feed and lay their yellow eggs upon the unfolding leaves. Beetles eat holes through the leaves, eggs hatch in a week, and the larvae eat off the green tissue from the under surface, causing the leaves to turn brown and fall about the middle of July, at which time the larvae are about full grown. Then they descend to the base of the tree and transform to naked pupae; ten days later the adult beetles emerge and lay eggs for the second brood or go early into winter quarters.

Two complete defoliations in succession will kill a tree.

How to Fight the Elm Leaf Beetle.

(1) Search all attics, church belfries and cupolas for the dormant beetles in winter and early spring. Sweep them up and burn them.

(2) Spray the leaves with poison as soon as they have opened, if their shot-hole appearance shows that the beetles are there in abundance, and the under sides of the leaves should be coated about June 1st to destroy the larvae or grubs.

(3) When, later in the season, the yellow pupae appear on the trunks of trees and the ground beneath, kill them with a spray of kerosene emulsion or by sweeping them up and burning or soaking with kerosene.

The first and third measures should be taken by each householder, church or social organization, at individual expense. The spraying, which is difficult and expensive if the trees are large, can only be done by concerted action of the town or borough authorities. For extensive spraying work power sprayers are desirable, but small elms or a few large trees can be treated successfully by means of a hand pump of barrel size. The "Niagara gas sprayer" in operation is shown in Fig. 6.

Fig. 6.—"Niagara gas sprayer" and outfit for spraying street trees. (After Smith, Report New Jersey Expt. Station for 1905.)

The Spruce Gall-Aphid

Glenn W. Herrick and T. Tanaka

A GROUP OF GALLS ON SPRUCE IN JULY

Published by the
Cornell University
Agricultural Experiment Station
Ithaca, New York

Received for publication April 10, 1926

THE SPRUCE GALL-APHID

(*Adelges abietis* Kaltenbach)

Order, *Homoptera*

Family, *Phylloxeridae*

GLENN W. HERRICK AND T. TANAKA

Certain of the native spruce trees in this country, and one introduced species, the Norway spruce, are infested with a species of aphid, *Adelges abietis*, which causes galls of a familiar pine-apple type on the branches. The Norway spruce is especially subject to the attacks of this pest. All of the spruces, because of their evergreen character and their dense and shapely growth, are attractive trees in the plantings of both small and large grounds, and are useful for windbreaks in various situations. The Norway spruce is often used for hedges, but various factors contribute to its undesirability as a hedge plant, notably the injuries produced by the gall-aphid.

HISTORY, DISTRIBUTION, AND INJURY

It is not possible to give a detailed history of the spruce gall-aphid in America because the insect has been confused in the literature with other closely related species and one cannot be sure of the species under discussion. It was certainly introduced from Europe probably many years ago. It is widely distributed in the northeastern United States and in southern Canada. Just how far west it extends, we have no accurate means of knowing. It has a wide distribution in New York, for the typical galls have been received from many different localities. It appears to be more common and injurious in New York, Pennsylvania, and the New England States, than in any other part of the United States.

In this country the insect attacks particularly the Norway spruce (*Picea Abies*) and the white spruce (*Picea canadensis*). Britton (1924) records it as attacking also the black spruce (*Picea mariana*), the red spruce (*Picea rubra*), and sometimes the hemlock (*Tsuga canadensis*). The Colorado blue spruce (*Picea pungens*) is infested with a closely related species of gall-aphid (*Chermes cooleyi* Gillette), which causes galls on the branches similar in appearance to those of the spruce gall-aphid but longer and nearly always at the ends of branches, as shown by Gillette (1907-08).

The spruce gall-aphid is seriously injurious to Norway spruces especially, and is particularly harmful to young trees. There may be hundreds of the galls on a single small tree. Miss Patch (1909) counted 990 fresh galls on a white spruce three feet tall. The writers have seen as many on Norway spruce. In cases of such severe infestation, many of the branches are usually killed and often single young trees in hedges are killed outright, although the crowding and pruning undoubtedly contribute to the unhealthy condition of the plants. When branches are severely infested with the galls, they begin to shed their leaves and the tree becomes

ragged and unsightly in appearance. The work of the insect is a distinct handicap to the nurseryman, because customers do not like to purchase trees already infested with the unsightly galls, and they really should not accept infested trees unless they feel inclined to take vigorous measures of control after the trees have been set. The effect of the galls is to cause trees to become deformed and misshapen, thus detracting from their beauty and attractiveness.

SCIENTIFIC AND COMMON NAMES

: This aphid, together with other closely related species, has been placed heretofore in the genus *Chermes*, and this particular species has been known as *Chermes abietis*. It is found under this name in the references cited herein, in all except one instance. Comstock (1924) says it has been determined that the name *Chermes* should be applied to certain jumping plant-lice of the family Chermidae formerly known as the Psyllidae, and that those insects found on conifers which have been "quite generally known" under the generic name *Chermes* should be placed in the subfamily Adelginae. Therefore the forms on conifers heretofore known as *Chermes* will fall in the genus *Adelges*, and the species under discussion will be known as *Adelges abietis*.

The insect has been given several common names—*spruce gall-louse*, *European spruce bud-louse*, *yellow chermes*, *green-winged chermes*, and *green-winged adelges*. It seems to the authors that the common name *spruce gall-aphid* comes nearer to expressing the common and distinctive characteristics of the insect than any of the other names mentioned. The insect is an aphid, and its peculiar, distinctive, and commonly known injury to its food plant is in the form of a gall. The name *spruce gall-aphid*, therefore, indicates at once its food plant, its character and position among insects, and the form of its injury.

DESCRIPTION OF FORMS

The wingless oviparous female

The body of the adult wingless oviparous female is grayish green and is covered with a white waxy pulverulence which increases with age. The groups of pores are dark, and the mouth parts, antennae, and legs are dark brown. The body is more or less globular and oval as viewed from the dorsum. It is from 1.5 to 2 millimeters long and from 1 to 1.5 millimeters wide.

The head and the thorax are closely joined, and there are three transverse rows of groups of dorsal pores in this region. The mesothorax, the metathorax, and each segment of the abdomen, bear a transverse dorsal row of groups of wax pores. The groups of pores are arranged on the body in six longitudinal rows, four rows along the dorsum proper and one additional row along each side of the body. Near the caudal end of the body the two side rows disappear. The number of wax pores in each group is large, but is not the same in all of the groups. The wax pores are not so numerous on the ventral side of the body, although there are from two to four groups on most of the segments. There are two

pairs of spiracles on the ventral side of the thorax, and at least four pairs on the abdomen.

The antennae are short but conspicuous. The first segment is short and thick, the second is slightly longer but somewhat slenderer, while the third and last is as long as or longer than the first two. The stylets of the mouth parts are very long. The legs are short and are wholly concealed beneath the body. The tarsi are two-segmented, with the first segment very short.

The eggs of the wingless hibernating female

The eggs of the wingless hibernating female are laid in clusters of from 150 to 200, or even more, beneath the body of the female. They are oblong elliptical in shape, and greenish yellow in color when first laid. Later they turn darker, but they become yellowish again just before hatching. Each egg is furnished with a slender white waxy thread at least twice as long as the egg itself. This thread seems to be entangled among and connected with other longer threads underlying the batch of eggs and holding the mass fast to the branches.

The actual measurements of different eggs are as follows:

0.30×0.15 mm.	0.32×0.165 mm.
0.30×0.155 mm.	0.34×0.165 mm.
0.30×0.16 mm.	0.34×0.170 mm.

The first-instar nymph

The newly hatched nymph is yellow in color and the appendages are more or less transparent. The eyes are dark and each one seems to possess three facets. The antennae are three-segmented, the first two segments being short and the third one being longer than both of these together. There are no wax pores on the body.

Different individuals measure as follows:

0.42×0.20 mm.	0.47×0.22 mm.
0.42×0.22 mm.	0.49×0.23 mm.
0.45×0.21 mm.	

The second-instar nymph

The general color of the body and appendages in the second-instar nymph is very similar to that of the first-instar nymph, except that the body is now covered with a fine whitish waxy powder. The wax pores are therefore present, but they are not highly developed except on the head and the prothorax, where they are conspicuous.

The measurements of individuals are as follows:

0.27×0.51 mm.	0.32×0.64 mm.
0.28×0.56 mm.	0.34×0.64 mm.
0.29×0.61 mm.	0.36×0.67 mm.

The third-instar nymph

The body of the third-instar nymph is yellowish green and is covered with a white powdery secretion of wax. The appendages and the mouth

parts are light brown. The wax pores are well developed on the anterior part of the body. The nymph usually carries a cast skin, and sometimes two of them, attached to the posterior end of the body. These skins are filled with liquid and are more or less globular in shape.

Individuals of this stage measure as follows:

0.37×0.80 mm.	0.48×0.96 mm.
0.40×0.80 mm.	0.50×0.99 mm.
0.40×0.82 mm.	0.53×0.99 mm.
0.40×0.88 mm.	0.58×1.09 mm.
0.44×0.88 mm.	

The fourth-instar nymph, or pupa

In the nymph of the fourth instar, the body is stout and the wing-pads are well developed. The nymph is yellow in color and noticeably pulverulent. The eyes are black and prominent, and the legs are dark. This stage, which is of long duration, enables the nymph to grow considerably, and the individuals vary greatly in size. The following measurements show the variation in size:

0.68×1.32 mm.	0.90×1.99 mm.
0.78×1.61 mm.	0.90×2.02 mm.
0.80×1.75 mm.	0.94×2.10 mm.
0.85×1.90 mm.	1.03×2.21 mm.

The antennae are similar to those of the preceding instar, and the prothorax is fairly well differentiated from the fused meso-metathorax. The wing-pads of the mesothorax are large and extend to the middle of the second abdominal segment, while the hind wing-pads are smaller and are usually concealed beneath the front ones. The situation of the wax pores are well described by Miss Patch (1909) as follows:

Head with an anterior group of pores near base of each antenna and two median groups at posterior margin; prothorax with lateral gland area, an anterior median pair of groups, a small group (4 pores) between the lateral and median ones, and four groups near posterior margin of prothorax; mesothorax with an anterior lateral area; metathorax with two separated median groups; abdomen with small and widely separated median groups on segments I-VI, those on I being composed of about 6 pores, those on II of 3 and those of III-VI of 2 each, a very small group of lateral pores occur on segments II-IX (merged to a caudal group on IX), and about half way between the median group and lateral margin an additional group of about 3 pores occurs on segments I and II. The smallness of these inconspicuous groups is correlated aptly with the fact that this is not a flocculent species, the wax secretions not exceeding the pulverulent condition.

The winged oviparous female

In the winged oviparous female, the wings have an expanse of from 5.4 to 6.3 millimeters. The head is dark brown and about three times as wide as its length. The compound eyes are prominent and black. The antennae are smoky yellow and five-segmented; the first segment is short and stout, the second is longer than the first but less in diameter, the third is longer than the second, and the fourth and the fifth are sub-equal in length but are longer than the third (figure 1). Miss Patch (1909) says:

FIGURE 1. THE SPRUCE GALL-APHID

115, The winged female, which emerges from the gall in August. 116, Antenna of the winged female. 117, The pupa, which develops in the gall

(From Bulletin 173, Maine Agricultural Experiment Station)

"The sensorium of each of antennal joints III-V is confined to the distal half of the joint and extends not more than half way around the joint." There is a group of smaller sensoria on V.

The prothorax is smoky greenish yellow, with a dark triangular area in the middle of the dorsum; the mesothorax is pale yellow, with a dark triangular area near the front margin in the middle. Near the caudal border of the metathorax is a subquadrangular dark transverse plate. In the normal position the wings are folded along the abdomen in a roof-like manner. The fore wing has a wide green costal margin and a green stigma. The legs are slender and yellowish. The tarsi are two-segmented, with the basal segment very short and small and often concealed beneath the tip of the tibia. The abdomen is smoky yellow, with segments fairly distinct. The groups of wax pores "are much as in the pupa except for the abdomen which is as follows: I with median groups comparatively large and merged, II median groups separated and composed of 2 pores each, V with two small groups between median ones referred to. I-VIII with distinct lateral groups on darkened area, III-VI with small groups of 2 or 3 pores each midway between median and lateral groups" (Patch, 1909).

The measurements vary as follows:

Length of body

1.58 mm.
1.73 mm.
1.88 mm.
1.91 mm.
1.92 mm.
1.98 mm.
2.07 mm.
2.10 mm.

Expanse of wings

5.4 mm.
5.7 mm.
5.8 mm.
6.0 mm.
6.2 mm.
.....
6.3 mm.
.....

The eggs of the winged oviparous female

The eggs of the winged oviparous female are laid beneath the body of the female, while she sits lengthwise of a leaf, in a cluster of from 40 to 50. They remain covered with the dead body of the female, together with what waxy threads she may have secreted. They are dull yellow at first, and later become darker. Each egg is furnished with a waxy thread at one end.

The eggs measure as follows:

0.18×0.34 mm.
 0.18×0.36 mm.
 0.18×0.38 mm.

0.20×0.36 mm.
 0.20×0.38 mm.
 0.22×0.34 mm.

The first-instar nymph of the hibernating oviparous female

FIGURE 2. FIRST-INSTAR NYMPH OF THE OVER-WINTERING FEMALE, THE FORM WHICH PASSES THE WINTER ON THE SPRUCE

In the first-instar nymph of the hibernating oviparous female, the body is slender and the appendages are rather long. The body is light yellow when first hatched, but later becomes darker and somewhat green. The antennae are long and are composed of three segments, the first and second of which are short while the third is comparatively very long. The groups of wax pores are conspicuous and abundant. In general, they form six longitudinal rows as seen from above, four rows situated medianly and one row on each lateral edge of the body. There are usually four pores in each group of the median rows on the head, the thorax, and the first segments of the abdomen, but on the posterior segments of the abdomen there are more often but three pores in a group. Each of the lateral

groups tends to have numerous pores. This form (figure 2), which passes the winter on the branches of the spruce at the bases of buds, secretes more or less wax in the form of white threads that usually cover the body.

LIFE HISTORY

The spruce gall-aphids pass the winter as first-stage female nymphs clustered in crevices of the stem about the bases of the buds. On severely infested spruces the overwintering forms are abundant, there being as many as twenty to thirty about one bud. When examined with a lens they are seen to be purplish gray, but to the unaided eye they resemble small grains of gunpowder. By May 5, in 1923, the spruce buds had begun to grow and many of the nymphs had molted. So far as the writers' observations have gone, but one molt has been found in this generation, but some molts may have been missed. The females increase in size, and begin to secrete long, crinkly, curly, white, waxy hairs, and become bluish green in color. They all settle now at the bases of the developing branch buds. The weather continued cold after the above-named date, and the females remained rather stationary in development for a week. By May 16, however, females were found which had just begun to lay eggs. The individuals develop fairly in unison, although there are belated ones here and there that are sometimes a week behind the earlier ones in depositing eggs. By the end of May, in 1923, nearly if not quite all were depositing eggs. There is some evidence to indicate that egg laying is influenced by the development and activity of the tree. The buds on some trees start much earlier than do those on others, and on those trees which start early the aphids are found to be depositing eggs, while on the spruces, which are then still dormant, the aphids are inactive. At least this proved to be the case in some instances which the writers were fortunate enough to observe.

Each female deposits from 150 to 200 or more eggs beneath her body. On May 18 a female under observation was observed to exude from the caudal end of the abdomen a drop of liquid which appeared to be evidence of a beginning of activity in egg laying. On the 19th she became covered with a secretion of waxy threads. On the 21st, eggs could be seen beneath the body. These increased in number until, by the 29th, 198 eggs had been laid, some of which had already hatched. It is difficult to determine the exact length of the egg-laying period, because the first

FIGURE 3. OVERWINTERING FEMALES BEGINNING TO LAY EGGS, MAY 22

The insects are covered with white waxy threads

eggs are completely hidden by the body of the female and it is not easy to determine when fresh eggs have been added to the mass already present. The period appears to range from eight to fifteen days. For the reasons just given, it is difficult to determine the total number of eggs laid by a single female, but the number certainly ranges from 150 to more than 200, and probably as high as 250. The eggs and the body of the female are covered by a white mass of waxy threads and are more or less completely hidden from view (figure 3).

The eggs hatch in about one week. A female was removed from a batch of eggs on May 23. The eggs had been deposited on the 21st and 22d. On the 25th the light yellow eggs had turned green, and on the 26th the eyes of the embryo could be seen in some. On the 28th two eggs hatched, and on the 30th nearly all had hatched. The period of hatching, in this case, varied from seven to ten days, which is probably about the average period.

Formation of the galls and development of the nymphs

In 1923 the earliest nymph observed emerged from the egg on May 19. Others kept emerging up to June 5, by which time, in normal years, nearly if not quite all of the eggs will have hatched. The nymphs are active and they very soon crawl down behind the leaves of the buds, which have already begun to enlarge at their bases.

Formation of the galls

Cooley (1897) says:

A very young shoot broken out from the bud scales before any of the eggs have hatched shows the basal leaves on the side where the female passed the winter, distinctly swollen at the base. The starting of the gall must therefore be attributed to the female rather than to the nymphs. It is reasonable to suppose, however, that the nymphs settling in the partially formed gall have some influence on its further development.

All of our observations lead us to agree with this statement. In the case of a developing bud, at the base of which one or more females are situated with their beaks inserted in the tissues, we find more or less of the outer leaves of the bud enlarged and deformed even before the eggs hatch. The bases of these affected leaves are wider and thicker than the normal leaves. As the nymphs appear and crawl in between the bases of the leaves and insert their beaks into the tissues of the leaves, the irritation and stimulation are maintained and increased and the basal part of the affected leaves enlarges rapidly until a characteristic gall is formed. A normal bud, when it starts in the spring, is larger in diameter at the tip than at the base. On the contrary, a bud at the base of which one or more female aphids are hibernating, is as large in diameter at the base, and more often larger, than at the tip, when it begins to grow. The aphid unquestionably imparts to the leaves of the bud a stimulation that begins the formation of the gall.

The newly hatched nymphs are fairly active, and crawl up the young shoot which is now growing rapidly. They soon seek the spaces behind the modified leaves and crawl down among the bases of them. The modification of the leaves is apparently accelerated by the influence of the nymphs, for each leaf covering the aphids enlarges at the basal part while the distal part remains of normal size. Since the basal parts of many leaves become enlarged, a bulbous or gall-like enlargement of the whole

growing branch is formed and the branch itself is checked in growth. If the aphids are abundant enough to occupy spaces between all of the leaves, then the gall is formed the whole way around the branch (see cover illustration). If the aphids are few, the enlargement may take place on one side only of the growing branch. In such an event the branch is usually bent in the direction of the side on which the half-gall is formed. In some cases the gall so completely involves the whole bud that the growth of the branch is entirely checked. In most cases, however, the branch grows more or less beyond the gall, although never probably to what its full normal length would have been if no gall had been formed. Since the galls are deformations of the branch buds, they are always at the bases of branches.

The gall is more or less elongated, about 1 inch long, and shaped much like a pineapple, the sharp distal ends of the leaves answering to the spines and thus increasing the resemblance. The name *pineapple gall* is a very appropriate one. The gall is of a beautiful green color, somewhat yellowish at first but becoming deeper green later. After it has completely formed, reddish pubescent lines appear along the lines of junction of the leaves, and these add to its attractive coloration. The cells of the gall may be from 40 to 50 or even more in number, and each cell, containing from eight to twelve nymphs, is entirely closed. The red lines disappear as the gall grows older, and before the gall opens it turns yellowish in color as though it were ripening. During the latter half of August the leaves begin to separate and openings are formed leading to the cavities in which the aphids are situated. Through these openings the fourth-stage nymphs emerge, and the galls quickly turn dark and dry and harden; but they remain on the branches indefinitely, giving the trees an untidy appearance.

Development of the nymphs in the galls

Several nymphs are ensconced in the cavity behind each leaf, and their development can be traced by opening the galls at varying intervals during their growth. In 1923 the first nymphs to molt were found on June 4, and by June 7 many nymphs in their second instar were noted. On June 18 the first third-instar nymphs were observed, but most of the nymphs were still in the second instar. By July 2 most of the nymphs had transformed to the third instar, and on July 13 the first fourth-instar nymph was found. The fourth-instar nymphs are the so-called *pupae*, for in this stage the wing-pads are developed and are conspicuous. By August 9 most of the aphids had progressed to the fourth instar, and on August 15 the first winged adults were observed. By August 26 to 29, many of the pupae had emerged from the galls and had transformed to winged adults. It was not, however, until from September 5 to 10 that the majority of the galls were open.

The first eggs of the winged females were found on August 22. They were laid beneath the body of the female, on the leaf, to the number of 40 to 50. Many females were depositing eggs by September 11, and the first nymphs to hatch were discovered on that date. Many nymphs were present by September 29.

Emergence and habits of the winged female

When the different cells in the gall open in August, the pupae (fourth-instar nymphs) crawl out of the openings and settle on the leaves. In from

two to three days the nymphs molt and transform to the winged adult oviparous females. These females are certainly very inactive, and in the great majority of cases do not move far from the places in which the respective nymphs have cast their skins. The writers have never seen any indications of a migratory movement among these females. One wonders how this insect spreads from one plant to another, for it certainly does do so although the dispersal from a common center is gradual and very slow. The females settle rather quickly after the body has hardened, and, for the most part, go to the undersides of the branches. In from two to five days each one begins laying its quota of eggs, and the egg-laying continues for about ten days, when the full number, from 40 to 50, have been laid. The female dies when her eggs are all deposited, but her dead body covers the eggs and clings to the leaves long after the young nymphs have settled for the winter. Indeed, the shriveled bodies of the females may be found late in the winter. The eggs, which begin to be deposited during the first week of September, hatch through a period of two weeks, so that some of them do not hatch until the latter part of the month and probably some not until into October.

The young nymphs are active, and crawl about over the leaves apparently inserting their beaks now and then to feed. By the latter part of October most of the nymphs have migrated from the leaves to the twigs and have settled for the winter. They insert their beaks into the tender bark of the twigs, and many of them secrete waxy threads over their bodies. The nymphs tend to gather in groups mostly near the distal ends of the twigs, where they may be found during the winter.

Summary of life history

The insects overwinter as small nymphs of the oviparous, wingless, agamic females ensconced in crevices about the bases of the buds. In the spring these nymphs rapidly mature, and in May each adult lays from 150 to 200 or more greenish yellow eggs in clusters beneath her body (figure 4). These eggs hatch in about one week, and the young nymphs crawl down behind the already enlarged bases of the leaves of the spruce buds. The presence of the nymphs causes the leaves to enlarge further and the branch to become checked in its growth, thus forming a large gall resembling a miniature pine-

FIGURE 4. AN OVERWINTERING FEMALE JUST BEGINNING TO LAY EGGS, MAY 22

The insect is covered with white waxy threads

apple. The nymphs soon become entirely inclosed in the gall, and remain so until August. In the latter half of August (in New York) the galls open, and the nymphs crawl out of their cells, molt, and transform to winged, agamic, oviparous females. These females remain on the spruce, and in August and September each lays a cluster of from 40 to 50 eggs on the leaves. The eggs hatch in about one week, and the nymphs soon settle down in crevices about the buds, there to pass the winter. Thus the life cycle is completed, there being two parthenogenetic generations on the spruce each year.

PARTHENOGENETIC DEVELOPMENT

The species of the genus *Chermes* have been investigated with great detail in Europe through a period of many years, and considerable controversy has arisen regarding the life history and biology of the species *Adelges* (*Chermes*) *abietis*. Blochmann (1890) published a paper in which he held that *Chermes abietis* had an alternate or intermediate host plant, the larch tree. This immediately aroused the curiosity of Cholodkovsky as to how the insect lived in northern Europe, where the larch is not present. As a result of Cholodkovsky's careful and extended study (1910, 1911, 1915), he believed that there were two species of *Chermes*: one which he called *Chermes abietis* Kalt., and which passed through a cycle of two generations and lived permanently and parthenogenetically on the spruce; and another which he called *Chermes viridis* Ratz., with a cycle of five generations (one sexual) and living on both the spruce and the larch, the latter tree being the intermediate host. The accompanying diagram (figure 5) shows graphically the life cycle of these two species in Europe according to Cholodkovsky. Börner (1908, a and b), adopting the "parallel-row theory" of Dreyfus (1890), maintains that *Chermes abietis* develops two

FIGURE 5. GRAPHICAL REPRESENTATION OF THE LIFE HISTORY OF *CHERMES VIRIDIS* (A) AND OF *CHERMES ABIETIS* (B), ACCORDING TO CHOLODKOVSKY (AFTER STEVEN)

rices or strains or branches as it were, one of which lives on the spruce while the other migrates to the larch and passes through a complex cycle on that plant. In other words, he claims that the sexes (Sexuales) give rise to the hibernating nymphs (Fundatrix) which mature in the spring and lay the eggs that produce the gall-making forms (Fundatrigenia), part of which, when they mature, remain on the spruce and part of which migrate to the larch, thus living in parallel rows, as it were, on the spruce and the larch at the same time. Börner therefore holds that there is but one species of this form *Chermes abietis*, and that the parthenogenetic form of Cholodkovsky does not exist as a distinct species.

Our investigations in the vicinity of Ithaca do not show any sexual forms of this aphid. We have not found any evidence that there is a specific or varietal form or race that lives on the larch, there developing a sexual cycle. We should not like to say that further careful investigation might not disclose such a form. It is a problem that we hope to investigate at greater length. It is of interest in this connection to note that Miss Patch (1909) finds no evidence of *Adelges abietis* going to the larch. She says: "A different species of host plant is never sought by this *Chermes*."

Steven (1918) finds that in Scotland there are two separate cycles of the *Chermes* under consideration. One is a cycle of two parthenogenetic generations on the spruce, as we have found in Ithaca. This Steven calls *Chermes abietis* Kalt. of Cholodkovsky. He finds also a *Chermes* with a cycle of five generations, which passes one phase of its life history on the spruce and one phase on the larch. This, he believes, corresponds to *Chermes viridis* Ratz. of Cholodkovsky. This species or race we have not as yet found at Ithaca.

CONTROL

The senior author has been spraying Norway spruces for some years with different materials in an endeavor to control the spruce gall-aphid. Most of these trials have been isolated attempts to control the insect on two or three trees, but in the spring of 1924 an opportunity was offered to make a test on a larger scale. The results (Herrick, 1925) corroborated those of the earlier isolated sprayings and were very satisfactory.

A hedge of Norway spruce trees about five feet high, which runs along one end and about half the length of one side of the grounds of Prudence Risley Hall at Cornell University, has been abundantly infested with the gall-aphid, and the infestation has been gradually increasing until some of the young trees have nearly succumbed to the injuries inflicted by the insect. This hedge, which has a total length of about 550 feet, offered a fine opportunity for trying some control measures.

On April 9, 1924, the hedge was thoroughly sprayed with powdered lime-sulfur at the rate of $16\frac{1}{2}$ pounds to 50 gallons of water. The trees were sprayed from both sides, and care was taken to coat the branches thoroughly with the liquid. The senior author had this hedge under observation throughout the season, and was not able to find a single developing fresh gall. The caretaker of the grounds, who is interested in the problem and aided in the spraying, pruned the hedge in July, and, while doing so, kept a sharp lookout for galls. He failed to find a single one,

and, since he pruned each tree in a rather careful manner, it is likely that he would have seen galls if any had been present.

The senior author has along the front border of his own grounds a hedge of Norway spruce the trees of which are about five feet high. These small trees also have been severely infected with the gall-aphid. On April 21, 1924, this hedge, which is about 75 feet long, was sprayed with Sunoco oil at the rate of 1 part to 20 parts of water. Warm water was used and the oil emulsified readily, forming a very satisfactory milky emulsion. The hedge was sprayed from both sides in a very thorough manner with an abundance of the liquid. During the summer, frequent examinations were made of the trees in search of developing galls. The writer was unable to find any signs of the presence of the aphids and no fresh galls were discovered. It should be remarked that rain began to fall at about three o'clock in the afternoon of the day on which the trees were sprayed, and a drizzle continued through most of the afternoon.

On an individual tree of larger size, sprayed on April 5 with Sunoco oil at the rate of 1 gallon to 15 gallons of water, five freshly developing galls were found, one on June 10 at the tip end of a side branch, and four at the topmost end of the leader. Unquestionably the tip ends of these branches were left untouched by the oil. The lower branches of this tree were covered with old galls and the hibernating nymphs were abundant on nearly all of the branches at the time of spraying. No new galls, other than those mentioned, appeared on the tree.

All of the trees sprayed with oil were watched during the summer and fall for possible signs of injury, but no evidence of the slightest ill effect was observed until early in the following March, nearly a year after the application. Then it was noted that the snow underneath the trees was covered with spruce leaves, and on examination it was found that there appeared to be an abnormal shedding of the foliage. The ends of the branches of nearly all the young trees in the hedgerow were bare of leaves, making the effect very noticeable. The defoliation did not seem to be progressive, and did not appear to be serious although it did give the hedge a somewhat unthrifty and untidy appearance. Indeed, one cannot be positive that the effect was due to the oil, although there was no evidence of abnormal shedding of foliage on unsprayed spruce trees, many of which were examined. Later, on July 25, 1925, the trees were in excellent condition and the hedge showed no injury. On November 10, 1925, the hedge was in a perfectly normal condition.

In the spring of 1925 the attention of the writer was called to two Norway spruces about twenty feet high that were literally covered with the galls of this aphid. The trees had made little growth during the preceding year, and looked ragged and unhealthy. They were two among a half dozen fine trees standing in a row and close together. All of the trees had been making excellent growth, but these two, standing at the end of the row, had become infested, probably from two other infested trees some distance beyond them.

On April 6 these trees were sprayed with Scalecide, 1 quart to 20 quarts of water. The oil was fresh and emulsified very satisfactorily. The trees were sprayed very thoroughly and every part of them was covered. The weather during the next few days was propitious, and the oil had a

chance to do its work. April 7, 8, 9, and 10 were fine days without rain but with rather heavy frosts during the nights.

The results of this spraying with Scalecide seemed, up to November 10, to have been very satisfactory. Observations of the trees during the summer and fall failed to disclose a single new gall. All of the evidence indicated that this miscible oil is nearly a specific in destroying the spruce gall-aphid. There was no evidence of any injury to the tree by the oil, when applied during the spring of the year at the dilution of 1 part to 20 parts of water.

Britton (1924) says, regarding the control of this gall-aphid, that spruce trees have been effectively sprayed for several years with miscible oil, at a dilution of 1 to 20, in the fall and in April, and that there has never been any injury to the trees except once, and that was probably due to the contents of the original package not having been thoroughly mixed before dilution with water. It seems apparent that oil emulsions, miscible oils, and lime-sulfur either dry or liquid, are effective materials for the control of the spruce gall-aphid. The senior author has obtained effective results on a few trees at different times with the concentrated solution of lime-sulfur at winter strength, 1 gallon to 8 gallons of water, when the solution tested 32° Baumé. The lime-sulfur may be somewhat objectionable because it tends to stain the foliage of spruce trees; at least, it gives the trees a whitewashed appearance which may persist for some time and to which some owners might object. The oils do not produce any discoloration of the foliage, and if they do not produce marked injury to the trees, such as may prohibit their use on spruces, they are probably preferable.

REFERENCES CITED

- BLOCHMANN, F. Ueber die regelmässigen Wanderungen der Blattläuse, speziell über den Generationszyklus von *Chermes abietis* L. Biol. Centbl. 9 (1889-90): 271-284. 1890.
- BÖRNER, CARL. Systematik und Biologie der Chermiden. Zool. Anz. 32 (1907-08): 413-428. 1908 a.
- Über Chermesiden. II. Experimenteller Nachweis der Entstehung diöcischer aus monöcischen Cellaren. Zool. Anz. 33: 612-616. 1908 b.
- BRITTON, W. E. Spruce gall aphid. In Twenty-third report of the State Entomologist of Connecticut, 1923. Connecticut (New Haven) Agr. Exp. Sta. Bul. 256: 240-241. 1924.
- CHOLODKOVSKY, N. Aphidologische Mitteilungen. 26. Zur Kenntniss der westeuropäischen *Chermes*-Arten. Zool. Anz. 35 (1909-10): 279-284. 1910.
- Aphidologische Mitteilungen. 27. Über *Chermes abietis* Kalt. und *Ch. viridis* Ratz. Zool. Anz. 37: 172-174. 1911.
- [Russian title.] Chermes injurious to conifers. (Abstract.) Rev. appl. ent., Ser. A (Agricultural), 3: 592-599. 1915.
- COMSTOCK, JOHN HENRY. An introduction to entomology, p. 1-1044. (Reference on p. 429-432.) 1924.
- COOLEY, R. A. The spruce gall-louse. Massachusetts Agr. Coll. Rept. 34: 1-12. 1897.
- DREYFUS, L. Zu Prof. Blochmann's Aufsatz, "Ueber die regelmässigen Wanderungen der Blattläuse, speziell über den Generationszyklus von *Chermes abietis*." Biol. Centbl. 9 (1889-90): 363-376. 1890.
- GILLETTE, C. P. Chermes of Colorado conifers. Acad. Nat. Sci. Philadelphia. Proc. 59: 3-22. 1907-08.
- HERRICK, GLENN W. Methods of control for two shade-tree pests. Journ. econ. ent. 18: 630-632. 1925.
- PATCH, EDITH M. *Chermes abietis* Chol. In Chermes of Maine conifers. Maine Agr. Exp. Sta. Bul. 173: 290-294. 1909.
- STEVEN, H. M. Contributions to the knowledge of the family Chermesidae. No. I: The biology of the Chermes of spruce and larch and their relation to forestry. Roy. Soc. Edinburgh. Proc. 37 (1916-17): 356-381. 1918.

United States Department of Agriculture,

BUREAU OF ENTOMOLOGY,

L. O. HOWARD, Entomologist.

THE COTTONY MAPLE SCALE.

(Pulvinaria innumerabilis Rathvon.)

By J. G. SANDERS, Assistant.

The cottony maple scale (*Pulvinaria innumerabilis* Rathv.) at maturity is the most conspicuous scale insect indigenous to the United States, and has received much attention from entomologists and horticulturists on account of its occasional abundance. It was described in 1854 by Dr. S. S. Rathvon, of Lancaster, Pa., who gave the pest a very expressive and significant specific name. Since then over 60 articles and references concerning this species have appeared in entomological literature, besides hosts of others in newspapers and other periodicals. Of special worth is a monograph of the species, worked out in elaborate form by J. D. Putnam, which was published in the Proceedings of the Davenport (Iowa) Academy of Natural Sciences, p. 339 (1879).

FIG. 1.—*Pulvinaria innumerabilis*: adult females in position on twigs, with egg sacs—natural size (from Howard).

More recently in Bulletin 22, new series, Division of Entomology, Dr. L. O. Howard published a thorough treatise on "The Two Most Abundant Pulvinarias on Maple," giving in detail the distribution, food plants, and life history of the cottony maple scale and the nearly related maple-leaf scale, illustrated by numerous figures of the various stages of each species.

Sporadic outbreaks of this scale insect in various parts of the United States have been reported from time to time for many years past, but in each case natural enemies have subsequently increased with sufficient rapidity to effectually check its progress. Recently many reports of its extreme abundance have come to us from the city of Chicago and

vicinity, and from various sections of Illinois, Indiana, Wisconsin, and Iowa. It is interesting to note that most of the reports of serious injury by this insect come from those States or portions of States north of the fortieth parallel, where it seems to be attacked by fewer parasites than in the southern regions. A map of its distribution prepared by the author from locality cards shows that it is distinctively an Upper Austral zone species but occasionally reaches the Transition zone.

The cottony maple scale is classified in the soft, unarmored group (Non-Diaspinæ) of the family Coccidæ, and does not differ superficially from the Lecaniums, except in the formation of an ovisac by the female of the Pulvinaria. The male insect is a very small and delicate two-winged fly which develops its latter stage under a glassy test or covering and emerges late in August or early in September.

FOOD PLANTS.

The various species of maples, particularly the soft maple (*Acer saccharinum*), including the box-elder (*Acer negundo*), are the favorite food plants of this species. The writer has, however, found it on 47 different species of trees, shrubs, and vines, including various species of maple, oak, linden, elm, willow, poplar, beech, hawthorn, sycamore, locust, hackberry, osage-orange, mulberry, grape, poison-ivy, apple, pear, plum, peach, currant, gooseberry, rose, and Virginia creeper. It is possible that a very careful study will prove that all these infestations are not referable to one species, but that other species may be separated just as the maple-leaf Pulvinaria [*P. acericola* (Walsh and Riley)] and the osage-orange Pulvinaria [*P. macluræ* ("Kenn.," Fitch)], once considered identical, were found to be distinct from the cottony maple scale.

HABITS AND LIFE HISTORY.

The large, white, flocculent masses, resembling popcorn fastened to the twigs, which appear on infested trees during the month of June, are the cottony (really waxy) ovisacs of the female, provided for the retention of the eggs after oviposition (fig. 1). These may contain as many as 1,500 minute, oval, pale reddish-yellow eggs.

The larvæ hatch at various dates from June 1 to August 15, depending on the latitude and exposure to the sun's rays. After remaining in the ovisac for a day or two, they swarm over the twigs, instinctively migrating toward the light, and settle on the leaves along the midribs and veins, always preferring the under surface (fig. 2, *e*). The larvæ on box-elder become active somewhat in advance of those on maple. This difference may be due to the food, but it must be remembered that the shade of the box-elder is less dense than that of the maple, and consequently the greater amount of heat and light may be the controlling factor. The male larvæ, when fully grown, assume a propupal stage from which they pass to the true pupal stage having a pinkish

hue. In a few days the winged males appear, but remain beneath the scale for two or three days before emerging. The females at this time have a few dorsal brownish markings and have undergone two molts. The males and females copulate in September, and soon afterward a small proportion of the females migrate to the twigs, where they insert their probosces and become fixed for the remainder of their existence, meanwhile changing from a greenish color to buff and finally brown, with a slightly tessellated waxy covering. In this condition the winter is passed (fig. 3).

When the sap begins to flow in the early spring the female grows rapidly, mainly because of the rapid development of great numbers of

FIG. 2.—*Pulvinaria innumerabilis*: a, newly-hatched young; b, female, third stage, from above; c, same, from side; d, male, third stage; e, same, natural size, on leaf and petiole; f, same, enlarged, on leaf petiole showing two specimens parasitized—all greatly enlarged except e (from Howard).

eggs within the body. In May or early June the ovisac begins to form from the fine waxy threads exuded from the posterior spinnerets, pushing backward and upward until the body of the female is almost vertical to the twig. Meanwhile the ovisac is gradually being filled with eggs, which hatch and undergo all the changes mentioned above. Thus, fortunately, there is but a single brood each year.

PARASITES AND PREDACEOUS ENEMIES.

Notwithstanding the extreme prolificacy of this insect, it is usually held in check by its many natural enemies. Doctor Howard has observed the English sparrow apparently feeding upon the full-grown scales. The twice-stabbed ladybird (*Chilocorus bivulnerus* Muls.) is a common enemy of this as of other scale insects, particularly in its early stages. The smaller similarly marked *Hyperaspis binotata* Say (fig. 4) and *H. signata* Melsh. are also valuable enemies of this species. The writer found in the severely infested parks and cemeteries

in Chicago that the egg-contents of 80 to 85 per cent of the cottony ovisacs had been destroyed by the white mealy larvæ of *Hyperaspis*. These larvæ enter and devour the entire contents of one ovisac after another without destroying its form, so that only an examination will reveal the empty ones.

The predaceous caterpillar of *Latilia coccidivora* Comst. has been an efficient ally in controlling the pest in the vicinity of Washington, D. C. It is particularly effective where the cottony masses are in close proximity, because of its habit of eating its way through the masses, spinning a strong web about itself, and forming a gallery as it progresses. The caterpillar is very active and moves about freely within its web.

Several specimens of *Leucopis nigricornis* Egger, a common dipterous parasite of aphides, were reared from cottony maple scale collected in Chicago, Ill., on July 21, 1905, by the writer, and from other specimens sent to the Bureau from Crystal Lake, Ill. This parasite is considered to be a European species, and although aphides seem to be its favorite host, it is occasionally reared from nondiaspine Coccidæ.

FIG. 3.—*Pultrinaria innu-mcrabilis*: gravid female, greatly enlarged, before commencing to secrete egg sac in the spring (from Howard).

Several species of chalcid flies (Chalcidoidea) are truly parasitic upon the cottony maple scale as well as upon others of the Lecaniinæ. There is no doubt that many more Pulvinarias are killed by these interesting little fellows than by all other enemies. *Coccophagus lecanii* Fitch, *C. flavoscutellum* Ashm., *Atropates collinsi* How., *Aphycus pulvinariae* How., *Comys fusca* How., and *Eunotus lividus* Ashm. have been reared from the cottony maple scale.

REMEDIES.

In dealing with an infestation by the cottony maple scale, the most important matter to be considered in each case is the advisability of artificial means of control under the existing conditions. The natural enemies of this pest have done and are still doing such effective work in its control, that it is a question whether remedies should be applied when the infestation is not serious. Nature always maintains a balance, and this pest can not be in the ascendancy for any great length of time. On the contrary its parasites will eventually multiply so rapidly as to completely check its progress, and it may even narrowly escape extermination.

Insecticides applied for a pest always kill its parasites and oftentimes predaceous enemies are also destroyed. Hundreds of larvæ of *Hyperaspis binotata* were found to have been destroyed by a summer treatment with kerosene emulsion for the cottony maple scale in parks

of Chicago, and no doubt thousands of the minute chalcids were killed, as they were found quite plentifully on unsprayed trees.

If it is necessary to resort to artificial means of control, as seems to be the case in some sections, the spraying should be delayed until fall or winter, when the *Hyperaspis* is hibernating at the bases of the trees among lichens, moss or dried grass. Care should be exercised in the use of spray, and the bases of trees should be covered with canvas or other suitable material to prevent the spray from running down the tree-trunks and collecting at their bases.

WINTER TREATMENT.

When the trees have become dormant, after the falling of the leaves, they can be trimmed and thoroughly sprayed with a strong kerosene emulsion, which will kill every scale reached by the spray, without injury to the trees. Scarcely one-fourth the quantity of emulsion is required to spray a tree in winter condition that is necessary when a tree is covered with dense foliage and both sides of the leaves must be reached with the spray. Some excellent results have been obtained by Mr. S. Arthur Johnson¹ in experiments against this pest in Denver, Colo. He found that kerosene emulsion 25 per cent or more in strength or whale-oil soap at the rate of 1 pound to 1 gallon of water was very effective, apparently killing all scales which received the spray.

FIG. 4.—*Hyperaspis binotata*: a, adult; b, antenna; c, palpus; d, larva—enlarged (original).

SUMMER TREATMENT.

Kerosene emulsion of not more than 10 to 12 per cent of oil can be thoroughly applied with safety to maple trees, but the tips and margins of the leaves may be injured even at this strength. Box-elder will be almost defoliated with a 12 to 15 per cent solution. Less than a 10 per cent emulsion will be of little value against the young unless they have recently hatched. Since the hatching period may extend over six or eight weeks, it will be seen that more than one spraying will be necessary to insure success, and, coupled with the fact that it is a very difficult and disagreeable task to thoroughly spray a tree in foliage, the winter treatment will be found more satisfactory in every way. And at that season the larvæ of predaceous beetles would not be destroyed.

¹ 1905: Bul. 52, U. S. Dept. Agr., Bureau of Entomology, pp. 85-88.

KEROSENE EMULSION.

Stock solution (66 per cent oil).

Kerosene (coal-oil, lamp-oil) -----	gallons..	2
Whale-oil or laundry soap (or 1 quart soft soap) -----	pound..	$\frac{1}{2}$
Water -----	gallon..	1

Dissolve the soap in boiling water, *then remove from the fire*, add the kerosene immediately and thoroughly agitate the mixture until a creamy solution is obtained. This can be done by pouring the mixture into the tank of a spray-pump and pumping the liquid through the nozzle back into the tank. This is a stock solution, which must be diluted before using. In order to make a 10 per cent emulsion, add to each gallon of the stock solution about 6 gallons of water and agitate thoroughly before using. For a 30 per cent solution add to each gallon of the stock solution $1\frac{1}{3}$ gallons of water and agitate thoroughly. This strength will kill a large percentage of the hibernating females, without injury to the trees.

If a good naphtha soap can be obtained, the preparation of the emulsion will be simplified. It will be unnecessary to heat the solution, since the kerosene will combine readily with the naphtha and soap and form a perfect, cold, milky-white emulsion when the mixture is thoroughly agitated. If naphtha soap is used, double the amount called for by the formula, and emulsify in soft (rain) water.

Approved:

JAMES WILSON,

Secretary of Agriculture.

WASHINGTON, D. C., *September 7, 1905.*

U. S. DEPARTMENT OF AGRICULTURE,
BUREAU OF ENTOMOLOGY—BULLETIN NO. 58, Part III.

L. O. HOWARD, Entomologist and Chief of Bureau.

SOME INSECTS INJURIOUS TO FORESTS.

ADDITIONAL DATA ON THE
LOCUST BORER.

BY

A. D. HOPKINS,

In Charge of Forest Insect Investigations.

ISSUED MARCH 5 1907.

WASHINGTON:
GOVERNMENT PRINTING OFFICE.
1907.

LETTER OF TRANSMITTAL

U. S. DEPARTMENT OF AGRICULTURE,
BUREAU OF ENTOMOLOGY,
Washington, D. C., February 5, 1907.

SIR: I have the honor to transmit herewith a manuscript entitled "Additional Data on the Locust Borer," prepared by Dr. A. D. Hopkins, in charge of forest insect investigations of this Bureau. It comprises a partial revision of Part I of Bulletin No. 58, and additional information based on the results of investigations carried on since that part was issued. I recommend that it be published as Bulletin No. 58, Part III, of the Bureau of Entomology.

Respectfully,

L. O. HOWARD,
Entomologist and Chief of Bureau.

Hon. JAMES WILSON,
Secretary of Agriculture.

CONTENTS.

	Page.
Seasonal history.....	31
Hibernation.....	31
Activity of the overwintered larvæ.....	31
Activity of the adults.....	32
Variation in seasonal history between different latitudes and altitudes ...	32
Habits of larvæ and adults.....	33
Economic features.....	34
Destructive character of the work.....	34
Evidences of attack.....	34
Favorable and unfavorable conditions for destructive work.....	35
Natural enemies.....	35
Insects.....	35
Disease and sap flow.....	36
Methods of control.....	36
Time to cut locust to destroy the young larvæ.....	36
Destruction of infested trees and wood.....	37
Spraying the trunks and branches to kill the young larvæ.....	37
Poison bait.....	38
Damage to cut wood and danger of introducing the insect into new localities.....	38
Selection of localities for extensive plantings.....	38
Management of plantations to prevent injury.....	39
Selecting and breeding borer-resistant trees.....	39

SOME INSECTS INJURIOUS TO FORESTS.

ADDITIONAL DATA ON THE LOCUST BORER.

(*Cyrtene robiniae* Forst.)^a

By A. D. HOPKINS,
In Charge of Forest Insect Investigations.

This part of Bulletin 58 contains a partial revision of Part I, with additional information based on the results of subsequent investigations by the writer and one of his assistants, Mr. W. F. Fiske.

SEASONAL HISTORY.

The data under this head refer to the District of Columbia and vicinity, latitude 39°, altitude 10 to 90 feet above tide.

HIBERNATION.

Hibernation begins soon after the larvæ hatch from eggs deposited at various times from August to October, and the period is passed as minute larvæ, scarcely longer than the eggs from which they hatch, in small individual hibernating cells excavated by them just beneath the corky bark and in the outer layers of the living bark on the main trunk of the larger to small trees or small saplings, and larger to small branches.

ACTIVITY OF THE OVERWINTERED LARVÆ.

Activity of the overwintered larvæ begins in April, or with the beginning of the movement of the sap in the bark and just before the leaf buds open. In 1906 activity began April 11; on April 13 the more advanced individuals had entered to the wood, on the 16th were grooving the surface, and on the 25th some of them had entered the wood. By May 11 nearly all of them had entered the sapwood and some of them had extended their burrows into the heartwood and were rapidly

^a Order Coleoptera, Family Cerambycidae.

increasing in size and very active, so that by May 20 some of them were more than half grown. They continued actively feeding and growing until after the middle of July, when they began to transform to pupæ and continued transforming during August until all had transformed, probably by the 1st of September. The pupæ begin transforming to adults about the 1st of August and continue transforming probably into September, although the principal transformation is in August.

ACTIVITY OF THE ADULTS.

The adults begin to emerge as early as the 7th of August, and continue emerging until the last of September, the greater number coming out during the last part of August and the first half of September. Evidently all beetles are out by the first week in October.

The females begin to deposit eggs within a few hours after they emerge. The principal period of oviposition appears to be between the middle of August and middle of September, but oviposition continues until in October. The eggs hatch within eight or ten days after they are deposited, and the young larvæ excavate their hibernating cells and remain dormant until the following spring.

VARIATION IN SEASONAL HISTORY BETWEEN DIFFERENT LATITUDES AND ALTITUDES.

Phenological investigations of plants and insects by the writer^a during the past ten years indicate that the average difference in the dates of occurrence of the different stages of *Cyllene robinix* at different latitudes and altitudes in the eastern United States will not be far from four days later for each degree north of latitude 39° and for each 400 feet of altitude above Washington at the same latitude, or four days earlier for each degree south of latitude 39° at the same altitude.

Thus, at latitude 43° in central New York, or central Michigan, with altitude the same as at Washington, the dates would be about sixteen days later, and at altitudes of 1,000 feet at latitude 43° they would be about twenty-six days later; at the same altitude as that of Washington at latitude 35° in southern North Carolina and Tennessee they would be about sixteen days earlier or at 1,600 feet elevation about the same. Thus we would have about thirty-two days' difference between localities at the same altitude in central New York and southern North Carolina. We would also have thirty-two days' difference between Washington and localities at latitude 39° and altitudes of 3,200 feet in the mountains of Virginia and West Virginia.

^a Bull. 50, W. Va. Agric. Exp. Sta., 1898, pp. 17, 18, and Bul. 67, 1900, pp. 241-248, with map.

HABITS OF LARVÆ AND ADULTS.

When a larva begins activity in the spring it molts and proceeds to excavate an independent food and entrance burrow through the dead area of bark surrounding the hibernating cell or through the living bark immediately surrounding the dead area, until it reaches the cambium. It then excavates an irregular groove or cavity in the outer sapwood, returning frequently to the outer cell or opening to push out the borings and apparently to get relief from the exuding sap. A large per cent of the larvæ die before any further progress is made, but survivors grow rapidly and soon succeed in overcoming the many obstacles, including natural enemies, resistance of the tree, etc., and enter the sapwood. From this stage on until the larvæ have attained their full growth they are very active and destructive. Their food consists principally of the nutritious substances of the bark and wood, and probably of the liquids flowing into the burrow, but they do not hesitate to kill and feed upon each other when two or more come in contact within the same burrow. The fact that the entire development often takes place in a burrow scarcely more than twice the length of a matured larva indicates that food must be obtained from some source other than the wood and bark. Throughout its active life the larva frequently returns to the inner and outer bark to enlarge the burrow, and push out its borings, so that the burrow when completed is of a diameter throughout sufficient to allow the passage back and forth of the full-grown larva. When full grown the larva enlarges the inner end of the burrow, plugs the outer portion with boring chips, and in due time transforms in succession to the pupa and adult. When the adult is fully matured it escapes through the exit prepared by the larva. Immediately after a female emerges she is joined by one or more males, and within a few hours, or within twenty-four hours, she proceeds to deposit eggs. She runs about over the bark investigating the crevices, by means of her ovipositor, to locate those most suitable for an egg. Sometimes as many as twenty places are critically examined before one is selected, and it appears that but one egg is deposited in a place by the same female, but other females may find the same place and each deposit an egg, so that sometimes several eggs are found in one crevice. As a rule, however, there is but one. The faculty of the female in locating the most suitable place for an egg by means of the sensitive palpi on the tip of the ovipositor is remarkable.

The beetles feed principally on pollen from the flowers of golden-rod, but are very fond of any sweet liquid, such as sugar sirup placed on the trunks of the trees. They are found during the day on the trunks,

branches, and foliage of the locust, and during their principal period of activity, from toward the last of August to the middle of September, they are especially common on the golden-rod flowers. Mr. Fiske determined that they were also actively copulating and depositing eggs as late as 10 o'clock at night.

The attack of this insect is apparently confined to the black or yellow locust (*Robinia pseudacacia*).

ECONOMIC FEATURES.

DESTRUCTIVE CHARACTER OF THE WORK.

The destructive character of the work of the locust borer is a matter of great economic importance. This insect attacks the otherwise perfectly healthy trees, and in addition to causing the detrimental wormhole defects in the wood it often kills the trees or renders an otherwise valuable product worthless except for fuel. It is much more destructive in some localities and sections than in others, and also much more destructive to some trees in the same grove than it is to others. It is more destructive also to young saplings and the branches of medium-sized trees than to the larger trees.

The death of a tree is caused principally by injuries to the inner bark and cambium, resulting from repeated attacks. Injuries to the wood alone do not result in the death of trees except when all of the wood is practically destroyed or sufficiently injured to cause the tree to fall or be broken down by the wind.

The commercial value of the wood product is diminished or destroyed by the wormhole defects, but for certain purposes, as, for instance, fence posts, a limited number of such defects are not detrimental, except so far as they may contribute to decay.

EVIDENCES OF ATTACK.

The first evidence of attack is fine brownish boring dust and wet spots on the bark, first observed in April, when the overwintered larvæ begin to enter the inner bark. As soon as the larvæ begin to groove the surface of the wood and enter the sapwood, their presence, in addition to the wet spots, is indicated by yellowish boring dust mixed with liquids and the gum-like exudations. After all of the larvæ have entered the wood their presence is plainly shown by the quantities of yellowish boring dust lodged in the loose bark on the trunk, in the forks of the tree or branches, and around the base. At this stage, usually about the middle of May, the badly infested trees which will die are plainly indicated by the failure of the leaf buds to open, or by the dwarfed or faded and sickly appearance of the foliage, and toward the last of the month and until the larvæ have completed their work in July, by the breaking down of the branches and small trees.

FAVORABLE AND UNFAVORABLE CONDITIONS FOR DESTRUCTIVE WORK.

Favorable conditions for the destructive work of the borer appear to be found in the presence of isolated trees and groves in the open in localities where golden-rod is present or abundant; also, where less resistant varieties of the tree prevail.

Unfavorable conditions are found in forest growth or large areas of pure stands, or mixed stands where the locust predominates; also, in plantations and groves where resistant varieties prevail, and where there is no golden-rod or other favorite food for the beetles. It is also found that coarse, thick bark is less favorable than the thinner bark on old and young trees and saplings.

NATURAL ENEMIES.

INSECTS.

Several predaceous insect enemies of the larvæ have been observed, but so far no true parasites have been discovered. A large elaterid larva (*Hemirhipus fascicularis* Fab.) appears to be the principal enemy of the borer after the latter has entered the wood. It resembles the borer somewhat, but is easily distinguished by the more flattened and shiny body, long prothoracic legs and two curved spines on the last abdominal segment. This predaceous larva is frequently found in the empty mines of the Cyllene larvæ, therefore it is evidently an enemy of considerable importance.

A slender, cylindrical, whitish, footless dipterous larva of an undetermined species is sometimes found in the mines in the wood, and, according to an observation made by Mr. Pergande, it may attack and kill the borers.

Whitish, flattened larvæ of the nitidulid genus *Ips*, with prominent branched hooks on the last abdominal segment, are common in the sap at the entrance of the mines and in the burrows made by the young borers in the inner bark and outer wood. They are supposed to be sap feeders, but the writer found they would attack and devour young Cyllene larvæ when confined together in a bottle. Therefore it is possible that they kill a great many of the young borers before these enter the wood, which may account, in part, for the disappearance of such a large number of the young borers while in the bark-boring stage.

It was also demonstrated that if several young Cyllene larvæ of various sizes were placed together in a small vial, the larger ones would kill and eat the smaller ones. It is probable, therefore, that when several larvæ hatch from a cluster of eggs and but one survives—which is usually the case—the larger or stronger one has killed the weaker ones.

DISEASE AND SAP FLOW.

Dead larvæ are frequently found in the mines, covered with white flour-like spores, and sometimes these spores are so common that a perceptible cloud rises from the wood when it is split open. Experiments in placing some of the spores with healthy uninjured larvæ in bottles, as well as with those in the normal position in the wood, resulted in the death of the larvæ and the development of apparently the same disease, while the duplicate larvæ kept under the same conditions, but without contact with the spores, remained normal and healthy. Therefore this is a fungus which will kill the borers and one which is evidently of considerable importance.

The profuse flow of sap together with a gummy substance in the wounds made in the living bark and cambium is evidently detrimental to the normal progress of the young larvæ and apparently many of the latter are thus destroyed.

METHODS OF CONTROL.

It should be remembered that all the holes found in a tree and all other damage by the borer are not the work of one generation, but usually that of repeated annual attack during the life of the tree; also, that a burrow in the sapwood of a young tree remains the same burrow in the heartwood of the old tree, without change, as long as the tree exists, except in the healing of the original entrance. The number of borers and the annual amount of damage is not so great, therefore, as might appear, and, while each female is capable of depositing a hundred eggs, only a small percentage of the larvæ hatching from them survive the bark-infesting stage or complete their development to adults. This suggests that any method of management which will insure the destruction of a large per cent of the surviving larvæ and beetles each year will reduce the damage to a point where there will be practically no loss.

With our knowledge of the life history and habits of the insect it is now possible to make definite recommendations and suggestions for its control. Some of those of immediate practical importance are as follows:

TIME TO CUT LOCUST TO DESTROY THE YOUNG LARVÆ.

The cutting of locust for all purposes, including thinning operations and for private or commercial use, should be done during the period between the 1st of October and the 1st of April, the bark removed from the crude product, such as posts, poles, and the like, and the tops and thinnings burned. The removal of the bark from all desirable portions of the trunks of the trees felled during this period is important and necessary in order to destroy the larvæ before they

enter the wood. The work in all cases should be completed before the leaf buds begin to swell on the living trees in the spring.

DESTRUCTION OF INFESTED TREES AND WOOD.

When it is desirable to simply remove and destroy, by burning or otherwise, the badly infested and damaged trees to kill the broods of larvæ, the work should be done in May and June, when all such trees can be easily recognized by the boring dust, fading leaves, broken branches, etc., and must be completed before the beetles begin to emerge. Perhaps the best rule, applicable to all localities, latitudes, and elevations, is to complete the work by the time the flowers have all fallen from the trees, which will vary between different altitudes and latitudes from about the middle of May to the last of June. Another rule would be to complete the work before the earliest varieties of golden-rod begin to show evidences of flowering. This, however, would be the latest that the work should be done, because the beetles begin to emerge by the time the first golden-rod flowers appear.

SPRAYING THE TRUNKS AND BRANCHES TO KILL THE YOUNG LARVÆ.

Experiments have demonstrated that the hibernating larvæ may be killed by spraying the trunks and branches with a strong solution of kerosene emulsion. Therefore, when it is practicable or more desirable to adopt this method for the protection of small plantations, groves, or shade trees, the spraying should be done in the fall or winter, not earlier than November 1, and not later than April 1—in other words, during the dormant period of the tree. The following paragraphs, relative to the preparation of kerosene emulsion, are taken from Farmers' Bulletin No. 127, by C. L. Marlatt:

Kerosene emulsion (soap formula)—

Kerosene	gallons..	2
Whale-oil soap (or 1 quart soft soap)	pound..	$\frac{1}{2}$
Water	gallon..	1

The soap, first finely divided, is dissolved in the water by boiling and immediately added, boiling hot, away from the fire, to the kerosene. The whole mixture is then agitated violently while hot by being pumped back upon itself with a force pump and direct-discharge nozzle throwing a strong stream, preferably one-eighth inch in diameter. After from three to five minutes' pumping the emulsion should be perfect, and the mixture will have increased from one-third to one-half in bulk and assumed the consistency of cream. Well made, the emulsion will keep indefinitely, and should be diluted only as wanted for use.

For the treatment of large orchards or in municipal work requiring large quantities of the emulsion, it will be advisable to manufacture it with the aid of a steam or gasoline engine, as has been very successfully and economically done in several instances, all the work of heating, churning, etc., being accomplished by this means.

The use of whale-oil soap, especially if the emulsion is to be kept for any length of time, is strongly recommended, not only because the soap possesses considerable

insecticide value itself, but because the emulsion made with it is more permanent, and does not lose its creamy consistency, and is always easily diluted, whereas with most of the other common soaps the mixture becomes cheesy after a few days and needs reheating to mix with water. Soft soap answers very well, and 1 quart of it may be taken in lieu of the hard soaps.

In limestone regions, or where the water is very hard, some of the soap will combine with the lime or magnesia in the water and more or less of the oil will be freed, especially when the emulsion is diluted. Before use, such water should be broken with lye, or rain water employed. * * *

For use on locust trees dilute 1 gallon of emulsion with 2 gallons of soft water.

Pure kerosene and pure petroleum will effectually kill the insects, but may do some damage to the bark of the trees.

Experiments with carbolic emulsion indicate that this preparation is of no value to kill the young larvæ.

POISON BAIT.

Experiments showed that the beetles would feed readily on poisoned bait, such as sugar, sirup, or molasses with some arsenical mixed in, when this was smeared on the trees. Such baits are fatal to the beetles, but the danger of killing honeybees is so great that their use is not recommended in localities where honeybees are kept.

DAMAGE TO CUT WOOD AND DANGER OF INTRODUCING THE INSECT INTO NEW LOCALITIES.

We have determined that after the borers have once entered the wood they may complete their development in the cut and dry branches; they will evidently do so, therefore, in posts or other material manufactured from trees cut between the 1st of May and the middle of September; from this it is plain that locust should not be cut during this period for any purpose except to destroy the borers, or, if it should be necessary to cut it, the tops should be burned and the logs submerged in ponds or streams for a few days before they are shipped or manufactured. This is very important, both to prevent damage to the manufactured material and the introduction of the insect into the far West and other sections of the country which are at present free from it.

SELECTION OF LOCATIONS FOR EXTENSIVE PLANTINGS.

The fact that there are many sections and localities of greater or less extent within the natural home of the locust and its insect enemies where, from some unknown cause, the tree grows to large size and old age without perceptible injury from borers and other insects, suggests the importance of selecting such localities for any proposed extensive operations in the line of artificial planting, or utilization of

natural growth. It will be found, however, that no area of considerable extent, even in such localities, is entirely free from this and other destructive insect enemies, and that certain precautions and well-planned methods of management with reference to the control of the latter will be necessary.

MANAGEMENT OF PLANTATIONS TO PREVENT INJURY.

In the first place it is necessary, in order to provide against future losses from the borer, that a thorough survey be made in May and June, not only of the area to be utilized but of the entire neighborhood for a radius of a mile or more from its borders, for the purpose of locating and destroying scattering trees and groves which are more or less seriously infested or damaged by the borer. It would seem that the control of such large areas, by purchase or under a plan of cooperation between the owners of the land or trees, is one of the most important requisites for success in preventing future losses from the ravages of this and other insects in small as well as large plantations. In fact, it is the writer's opinion that, with this precaution properly and continuously carried out, locust may be successfully protected from the borer in any locality.

In the subsequent management of plantations and of natural forest and sprout growth it is important each year to locate and destroy the worst infested trees for the purpose of killing the borers in the wood, and to conduct the thinning and commercial cutting operations during the period between October of one year and April of the next, in order to destroy the young borers before they enter the wood.

Worthless, scrubby, borer-infested trees should be killed outright by stripping the bark from 4 or 5 feet of the lower stem during August to prevent sprouts and seed production from them, and at the same time to destroy the eggs and young borers. Trees deadened in this manner will usually be so completely killed that not a single root sprout will appear. Therefore this method is of special value in preventing sprout reproduction from inferior individual trees.

SELECTING AND BREEDING BORER-RESISTANT TREES.

The fact that some trees are more or less resistant to attack or injury by the borer, while adjacent ones in the same grove are attacked year after year and seriously damaged, suggests breeding races and varieties of the species which would be permanently resistant.

Breeding experiments have been begun in cooperation with the Bureau of Plant Industry and the Forest Service, but it will require several years to get definite results. In the meantime, however, it is important that seed and cuttings for commercial planting should be selected, as far as possible, from trees which show least damage from

the borer and are otherwise vigorous and healthy. From a well-established principle in the heredity of plants and animals, this practice of propagating from the best examples must certainly yield better results than would follow a disregard of the character of the trees from which seed or root propagations are made.

For reference to literature, and other information not included in this paper, the reader is referred to Part I of this bulletin, pages 1 to 16.

O

United States Department of Agriculture,

BUREAU OF ENTOMOLOGY,

L. O. HOWARD, Entomologist and Chief of Bureau.

THE LOCUST BORER¹ AND METHODS FOR ITS CONTROL.

By A. D. HOPKINS,

In Charge of Forest Insect Investigations.

The locust borer and its relation to detrimental and destructive injuries to the black or yellow locust² in the eastern United States have been subjects of special investigations by the Bureau of Entomology during the past two years, which have resulted in the determination of practical methods of control.

The locust borer is a whitish, elongate, so-called "round headed" grub or larva (fig. 1), which hatches from an egg (fig. 2) deposited by a black or brown and yellow striped long-horned winged beetle (fig. 3) found on the trees and on the flowers of golden-rod from August to October. The eggs are deposited in the crevices of the bark of living, growing trees from August to October, and the young borers (fig. 2, *b*, *c*) hatching therefrom excavate individual cells in the outer layers of the inner living

FIG. 1.—The locust borer (*Cyllene robiniae*): *a*, larva, dorsal view; *b*, same, lateral view. Line at right represents natural length (author's illustration). The larva in profile should show minute prothoracic feet.

¹ *Cyllene robiniae* Forst; Order Coleoptera, Family Cerambycidae.

² *Robinia pseudacacia*.

bark (fig. 4), where they pass the winter, and in the spring bore through the bark into the sapwood and heartwood. Here they transform in July and August to pupæ and in August and September to adult beetles, which soon emerge from the trees and deposit eggs for the next annual generation of borers and beetles.

The injury to the trees consists of wounds in the bark and sapwood which, if sufficiently severe or repeated year after year, result in either a stunted, worthless growth or the death of young and old trees, while the numerous worm holes in the wood reduce its commercial value or render it worthless.

The presence of the insects in injurious numbers is indicated (1) by the frequency of the adults on the golden-rod flowers and on the trees from the last of August until in October or later; (2) by the slight flow of sap and by the brownish borings where the young larvæ are at work in the bark during April; (3) by the yellowish borings lodged in the rough bark, in the forks of the tree, and on the ground around the base of the trunk, by the breaking down of the branches and young trees, and by the sickly appearance of the young twigs and leaves during May, June, and July.

This insect appears to be present and more or less injurious in all of that part of the United States which is east of the Great Plains and north of the Gulf States. Published information and reports of forest officials and others indicate that in Oklahoma and Indian Territory and west of the Great Plains the locust is now quite free from injury by the borer, but that these regions will remain exempt is by no means certain.

EXTENT OF DAMAGE AND LOSS.

So extensive is the damage to natural growth, artificial plantations, and shade trees that in some sections within the natural range of the tree in the Eastern States, but particularly in the Middle West, where both the tree and the insect have been introduced, it is considered unprofitable to grow the tree for shade or timber, and in such sections the natural sprout growth is often considered a pest rather than otherwise.

The loss resulting from defective timber, stunted growth, and the death of trees is represented by the difference in value between the damaged growth, or product, and the same if uninjured and healthy. This, if expressed in dollars, would represent a large sum.

CHARACTER OF THE INJURIES.

The destructive nature of the work of the locust borer is a matter of great economic importance. The borer attacks the otherwise perfectly healthy trees, and in addition to causing the detrimental worm-hole defects in the wood, it often kills the trees or renders an otherwise valuable product worthless except for fuel. It is much more destructive in some localities and sections than in others, and also much more

destructive to some trees in the same grove than it is to others. It is more destructive, also, to young saplings and the branches of medium-sized trees than to the larger trees.

The death of a tree is caused principally by injuries to the inner bark and cambium resulting from repeated attacks. Injuries to the wood alone do not result in the death of trees, except when all of the wood is practically destroyed or sufficiently injured to cause the tree to fall or be broken down by the wind.

The commercial value of the wood product is diminished or destroyed by the worm-hole defects, but for certain purposes, as, for instance, fence posts, a limited number of such defects are not detrimental, except so far as they may contribute to decay.

EVIDENCES OF ATTACK.

The first evidence of attack is fine brownish boring dust and wet spots on the bark, first observed in April, when the overwintered larvæ begin to enter the inner bark. As soon as the larvæ begin to groove the surface of the wood and enter the sapwood, their presence, in addition to the wet spots, is indicated by yellowish boring dust mixed with liquids and the gum-like exudations. After all of the larvæ have entered the wood their presence is plainly indicated by the quantities of yellowish boring dust lodged in the loose bark on the trunk, in the forks of the tree or branches, and around the base. At this stage, usually about the middle of May, the badly infested trees which will die are plainly indicated by the failure of the leaf buds to open or by the dwarfed or faded and sickly appearance of the foliage and, toward the last of the month until the larvæ have completed their work in July, by the breaking down of the branches and small trees.

FIG. 2.—The locust borer (*Cyllene robiniae*): a, eggs; b, c, larvæ from hibernation cells. Much enlarged (author's illustration).

FAVORABLE AND UNFAVORABLE CONDITIONS FOR DESTRUCTIVE WORK.

Favorable conditions for the destructive work of the borer appear to consist in the presence of isolated trees and groves in the open, in localities where golden-rod is present or abundant and where less resistant varieties of the tree prevail.

Unfavorable conditions are found in forest growth or large areas of pure stands, or mixed stands where the locust predominates; also in plantations and groves where resistant varieties prevail and where there is no golden-rod or other favorite food for the beetles. It is also found that coarse, thick bark is less favorable than the thinner bark on old and young trees and saplings.

METHODS OF CONTROL.

It should be remembered that all the holes found in a tree and all other damage by the borer are not the work of one generation, but usually that of repeated annual attack during the life of the tree; also that a burrow in the sapwood of a young tree remains the same burrow in the heartwood of the old tree, without change, as long as the tree exists, except in the healing of the original entrance; therefore the number of borers and the annual amount of damage are not so great as they might appear, and, while the females are capable of depositing a hundred eggs each, only a small percentage of the larvæ hatching from them survive the bark-infesting stage or complete their development to adults. This suggests that any method of management which will insure the destruction of a large percentage of the surviving larvæ and beetles each year will reduce the damage to a point where there will be practically no loss.

With our knowledge of the life history and habits of the insect, it is now possible to make definite recommendations and suggestions for its control. Some of those of immediate practical importance are as follows:

TIME TO CUT LOCUST TO DESTROY THE YOUNG LARVÆ.

The cutting of locust for all purposes, including thinning operations and for private or commercial use, should be done during the period beginning with the 1st of October and ending with the 1st of April, the bark removed from the crude product, such as posts, poles, and the like, and the tops and thinnings burned. The removal of the bark from all desirable portions of the trunks of the trees felled during this period is important and necessary, in order to destroy the larvæ before they enter the wood. The work in all cases should be completed before the leaf buds begin to swell on the living trees in the spring.

DESTRUCTION OF INFESTED TREES AND WOOD.

When it is desirable simply to remove and destroy, by burning or otherwise, the badly infested and damaged trees to kill the broods of larvæ, the work should be done in May and June, when all such trees can be easily recognized by the boring dust, fading leaves, broken branches, etc., but the work must be completed before the beetles begin to emerge. Perhaps the best rule applicable to all localities, latitudes, and elevations is to complete the work by the time the flowers have all fallen from the trees, which will vary between different altitudes and latitudes from about the middle of May to the last of June. Another rule would be to complete the work before the earliest varieties of golden-rod begin to show evidences of flowering. This, however, would be the latest that the work should be done, because the beetles begin to emerge by the time the first golden-rod flowers appear.

SPRAYING THE TRUNKS AND BRANCHES TO KILL THE YOUNG LARVÆ.

Experiments have demonstrated that the hibernating larvæ may be killed by spraying the trunks and branches with a strong solution of kerosene emulsion. Therefore, when it is practicable or more desirable to adopt this method for the protection of small plantations, groves, or shade trees, the spraying should be done in the fall or winter, not earlier than November 1 and not later than April 1, or, in other words, during the dormant period of the tree. The following paragraphs explaining the preparation of the kerosene emulsion are taken from Farmers' Bulletin No. 127 of this Department, by Mr. C. L. Marlatt:

Kerosene emulsion (soap formula)—

Kerosene.....	gallons..	2
Whale-oil soap (or 1 quart soft soap).....	pound..	$\frac{1}{2}$
Water.....	gallon..	1

FIG. 3.—The locust borer (*Cyllene robinia*): a, male beetle; b, female beetle. Much enlarged (author's illustration).

The soap, first finely divided, is dissolved in the water by boiling and immediately added boiling hot, away from the fire, to the kerosene. The whole mixture is then agitated violently while hot by being pumped back upon itself with a force pump and direct discharge nozzle throwing a strong stream, preferably one-eighth inch in diameter. After from three to five minutes' pumping the emulsion should be perfect, and the mixture will have increased from one-third to one-half in bulk and assumed the consistency of cream. Well made, emulsion will keep indefinitely and should be diluted only as wanted for use.

For the treatment of large orchards or in municipal work requiring large quantities of the emulsion, it will be advisable to manufacture it with the aid of a steam or gasoline engine, as has been very successfully and economically

done in several instances, all the work of heating, churning, etc., being accomplished by this means.

The use of whale-oil soap, especially if the emulsion is to be kept for any length of time, is strongly recommended, not only because the soap possesses considerable insecticide value itself, but because the emulsion made with it is more permanent, does not lose its creamy consistency, and is always easily diluted, whereas with most of the other common soaps the mixture becomes cheesy after a few days and needs reheating to mix with water. Soft soap answers very well, and 1 quart of it may be taken in lieu of the hard soaps.

In limestone regions or where the water is very hard some of the soap will combine with the lime or magnesia in the water, and more or less of the oil will be freed, especially when the emulsion is diluted. Before use, such water should be broken with lye or rain water employed. * * *

For use on locust trees dilute 1 gallon of emulsion with 2 gallons of soft water.

Pure kerosene and pure petroleum will effectually kill the insects, but may do some damage to the bark of the trees.

Experiments with carbolic emulsion indicate that this preparation is of no value to kill the young larvæ.

DAMAGE TO CUT WOOD AND DANGER OF INTRODUCTION INTO NEW LOCALITIES.

We have determined that after the borers have once entered the wood they may complete their development in the cut and dry branches. They will evidently do so, therefore, in posts or other material manufactured from trees cut between the first of May and the middle of September. From this it is plain that locust should not be cut during that period for any purpose except to destroy the borers, or, if it should be necessary to cut it, the tops should be burned and the logs submerged in ponds or streams for a few days before they are shipped or manufactured. This is very important, both to prevent further damage to the manufactured material by the borers remaining in the wood and also to prevent the introduction of the insect into the Far West and other sections of the country which are at present free from it.

SELECTION OF LOCATIONS FOR EXTENSIVE PLANTINGS.

The fact that there are many sections and localities of greater or less extent within the natural home of the locust and its insect enemies where, from some unknown cause, the tree grows to large size and old age without perceptible injury from borers and other insects suggests the importance of selecting such localities for any proposed extensive operations in the line of artificial planting or utilization of natural growth. It will be found, however, that no area of considerable extent, even in such localities, is entirely free from this and other destructive insect enemies, and that certain precautions and well-planned methods of management with reference to their control will be necessary.

MANAGEMENT OF PLANTATIONS TO PREVENT INJURY.

In the first place it is necessary, in order to provide against future losses from the borer, that a thorough survey be made in May and June, not only of the area to be utilized, but of the entire neighborhood for a radius of a mile or more from its borders, for the purpose of locating and destroying scattering trees and groves which are more or less seriously infested or damaged by the borer. It would seem that the control of such large areas, by purchase or under a plan of cooperation between the owners of the land or trees, is one of the most important requisites for success in preventing future losses from the ravages of this and other insects in small as well as large plantations. In fact, it is the writer's opinion that with this precaution properly and continuously carried out, locust may be successfully protected from the borer in any locality.

In the subsequent management of plantations and of natural forest and sprout growth it is important each year to locate and destroy the worst infested trees for the purpose of killing the borers in the wood and to conduct the thinning and commercial cutting operations during the period between October of one year and April of the next, in order to destroy the young borers before they enter the wood.

Worthless, scrubby, borer-infested trees should be killed outright by stripping the bark from 4 or 5 feet of the lower stem during August to prevent sprouts and seed production from them and at the same time to destroy the eggs and young borers. Trees deadened in this manner will usually be so completely killed that not a single root sprout will appear. Therefore this method is of special value in preventing sprout reproduction from inferior individual trees.

BREEDING AND PROPAGATING BORER-RESISTANT TREES.

The fact that some trees are to a greater or less extent immune from attack or injury by the borer, while adjacent ones in the same grove are attacked year after year and seriously damaged, suggests the idea of

FIG. 4.—The locust borer (*Cyllene robinia*): Hibernation or larval cells in outer portion of living inner bark. About natural size (author's illustration).

breeding races and varieties of the species which would be permanently immune.

Breeding experiments have been begun in cooperation with the Bureau of Plant Industry and the Forest Service, but it will require several years to get definite results. In the meantime, however, it is important that seed and cuttings for commercial planting should be selected as far as possible from trees which show least damage from the borer and are otherwise vigorous and healthy. From a well-established principle in the heredity of plants and animals this practice of propagating from the best examples must certainly yield better results than would follow a disregard of the character of the trees from which seed or root propagations are made.

Approved :

JAMES WILSON,

Secretary of Agriculture.

WASHINGTON, D. C., *February 1, 1907.*

O

PROTECTION FROM THE LOCUST BORER.

By F. C. CRAIGHEAD,

Specialist in Forest Entomology, Forest Insect Investigations.

CONTENTS.

	Page.		Page.
Introduction-----	1	How to recognize trees containing	
Historical-----	2	no borer defects-----	9
Observations by the writer-----	2	Character of growth of uninfested	
Investigations and experiments-----	3	stands-----	9
Natural growth in the vicinity		Conditions under which locust can	
of Falls Church, Va.-----	3	be grown-----	10
Plantations along the Pennsyl-		Control-----	11
vania Railroad-----	5	A method of handling severely	
Condition of tree necessary for		damaged plantations-----	11
borer attack-----	8	Treatment of shade trees-----	11
		Literature cited-----	12

INTRODUCTION.

The increasing value of black or yellow locust¹ for many purposes, and especially the recent demand for sound locust pins or tree-nails in the construction of wooden ships, render its protection from insect damage important.

In addition to the natural growth of locust in forests, farmers' woodlots, and abandoned fields, many attempts have been made to grow it on a commercial scale in plantations. These attempts in most cases have resulted in failure, owing to the serious damage to the wood and frequently the destruction of the trees caused by the locust borer.² The fact, however, that in natural growth and occasional plantations practically no injuries from the borer are found, while in other cases the trees are ruined for commercial purposes or killed outright, has led the writer to make a thorough investigation of the problem to ascertain the cause of the occasional immunity from borer injury.

As a result of these investigations, it appears practically certain that *plantations of this tree can be protected successfully from*

¹ *Robinia pseudacacia* L.² *Cyllene robiniae* Forst.

the borer and grown profitably on a commercial scale if the locusts are planted in thick stands or mixed with other trees so as to produce a densely shaded condition and natural pruning during the first 10 to 15 years of growth.

HISTORICAL.

The great variation in the extent of injury by borers to both planted and natural stands of black locust has been noted by many writers. In fact, as early as 1821, Pickering (2)¹ stated that trees of natural growth in groves were much less liable to injury than were transplanted trees. Schwarz (3) in 1890 observed that the insect lives in large colonies affecting all trees of small groves, while long hillsides full of locust are not infested. Cotton (5) observed that in Ohio injury was greater in single trees and plantations of considerable size than in natural forests. Hopkins (6) remarked that "Favorable conditions for the destructive work of the borer appear to be found in the presence of isolated trees and groves in the open. . . . Unfavorable conditions are found in forest growth or large areas of pure stands, or mixed stands where the locust predominates; also, in plantations and groves where resistant varieties prevail, and where there is no goldenrod or other favorite food for the beetles." Dearborn (1), Kellogg (4), and Garman (7) also call attention to this fact.

OBSERVATIONS BY THE WRITER.

In examining locust plantations during the last few years, the writer was greatly impressed with the absolute destruction of some tracts, while others, or parts of the same tracts, were thrifty and unmarred by borers. This was convincing evidence that the trees could be grown so as not to be injured by the locust borer. Many tracts, therefore, both planted and natural, were studied with the idea of securing evidence that might be applicable in a practical way. As far as possible the accurate history of many locust stands, both pure and mixed, was obtained and factors that might be responsible for the presence or absence of injury were weighed. As a result of such study, it was found that the amount of destruction was unquestionably greater in those tracts which had been pruned occasionally, closely grazed, or in which fire had gone through from time to time, killing out the underbrush and destroying the natural shade produced by weeds or shrubbery. The denser the growth, particularly weeds and undergrowth about the stem of the tree, the less was the amount of borer work and vice versa. Pure stands in open fields, where the trees were growing from 2 to 3 feet apart, were seldom injured, while near-by isolated trees were riddled by the borer.

¹ Reference is made by number in parenthesis to "Literature cited," p. 12.

Possible reasons for such immunity in dense stands were discussed with Dr. A. D. Hopkins in several letters during May, 1910. In a letter of May 11 Dr. Hopkins wrote:

Perhaps with the unpruned trees there is more shade and, since the beetles are sun-loving and active during the day, the more open and light conditions found in the pruned groves may serve as the attractive influence. This is the only thing I can think of that would make the difference. Observations should be made during the period of flight to determine the relative number of beetles found on trees under shaded and sunny exposures.

If this should lead to a solution of the problem and shaded conditions are favorable, then mixed planting with some quick growing tree like catalpa might be advisable.

This seems a very plausible theory to account for the immunity of unpruned and natural growths, and it is substantiated in the experiments conducted. No other factors considered seem to cover the conditions in the many localities where trees were examined.

INVESTIGATIONS AND EXPERIMENTS.

Natural growths of locust in the vicinity of Falls Church, Va., and plantations along the Pennsylvania Railroad right of ways were selected and used as experimental plats. Some of these near Falls Church were pruned while others were left as they were found. The history of plantations along the Pennsylvania Railroad as to planting and subsequent management has been furnished by the office of the forester of the railroad, which also provided facilities for the study of these plantations.

NATURAL GROWTH IN THE VICINITY OF FALLS CHURCH, VA.

1. On Miners Hill there is a stand of mixed growth of reproduction averaging 3 to 5 inches in diameter. The aspect is a westerly slope, the trees extending from the crest more than half way down the hill. It contains about an acre and is composed chiefly of oak, chestnut, tulip, persimmon, hickory, sassafras, maple, and locust. It is thickly undergrown with weeds and briars which now are dying out. The locust composed about 5 per cent of the stand. In the fall of 1915 these locusts were from 2 to 3 inches in diameter and contained no borers. In January, 1916, one end of this plot, constituting about one-tenth of the area, was pruned; all trees, except the locust, were cut out. The following fall adult beetles were observed ovipositing on the trees thus isolated, and examination in the spring of 1917 showed from 3 to 10 borers in each tree. Again in 1918 these trees were found to be heavily infested. In the remainder of the stand (that which is still growing naturally) about 1 tree in 10 can be found containing one or rarely two defects made by the borer.

2. Just across a road from the tract described above is an old orchard which is used as a pasture. About two dozen locusts of the

same age are growing here. These trees are gnarly, some broken off, and all heavily infested. In the spring of 1918, 10 to 35 larvæ were found in each of these trees. The cattle have kept down the weeds and underbrush which would otherwise protect the trunks.

3. In the same general locality, about one-half mile north of these tracts, lies an abandoned field covered with an almost pure stand of virgin scrub pine 12 to 14 years old. Mixed in this pine are quite a few locusts. During the winter of 1915-16 about an acre of this pine was cleared. The locusts, then averaging 4 inches in diameter, were left standing. None of the trees were infested, as were none of those in the midst of the remaining pine. All were straight, naturally well-pruned, thrifty trees. In the fall of 1916 these trees were attacked by the beetles, and many larvæ were found in them in the spring of 1917. The land has not been cultivated, so that a dense growth of weeds conceals the trunks for 6 to 8 feet above the ground. Scarcely any adults were found on these trees in the fall of 1918, and very few larvæ are expected in the spring of 1919.

In January, 1918, another part of this pine wood was cleared, leaving 15 trees exposed. In July the trees were examined and no borers found nor any evidence of injury in the past. During August and September, 1918, adults were observed ovipositing on these trees.

4. Another abandoned field on an easterly slope contains a clump of about a hundred small locusts, 2 to 6 inches in diameter, grouped about several large trees. These are closely spaced, averaging 2 feet apart. They are well mixed with sumac, tulip, and sassafras. In the surrounding field are many isolated trees. All these isolated trees are scrubby and badly infested by repeated borer attack, while those in the dense clump are tall, thrifty, and contain no borers or defects from past injury.

5. Another abandoned field one-half mile north of Green Gables station on the Washington and Virginia electric line contains several groups of locust from 6 to 18 years of age and many isolated trees. This field contains about 100 acres. Clumps of sassafras and pine mixed with locust and persimmon occupy much of the remaining area. Broom sedge and goldenrod cover the ground on all open spots. All isolated locusts are heavily infested and have been damaged severely by the borers. Many of these trees have fallen over (Pl. I, fig. 1).¹ Three excellent stands of locust occur here; one group of an almost pure stand is composed of trees 25 to 30 feet high and 6 to 8 inches in diameter. They grew very close together and have now thinned out to an average of 10 feet apart. There are more than 500 trees in this plot; all are thrifty and no evidence of borers can be found. They represent what can be grown in 18 years' time on

¹ The photographs were taken by Mr. R. A. St. George, scientific assistant, Forest Insect Investigations, Bureau of Entomology.

FIG. 1.—NATURAL GROWTH OF LOCUST IN AN OLD ABANDONED FIELD OF SOME 60 ACRES IN EXTENT AT FALLS CHURCH, VA.

Trees badly infested by the locust borer. Several killed and others blown over. Note scattered position. Diameter near base, 2 to 6 inches.

FIG. 2.—NATURAL GROWTH OF LOCUST IN AN OLD ABANDONED FIELD OF SOME 60 ACRES IN EXTENT AT FALLS CHURCH, VA.

Several old seed trees and surrounding root sprouts and seedlings. Note thick growth. None of these trees are infested. Diameter near base, 1 to 8 inches.

PROTECTION FROM THE LOCUST BORER.

FIG. 1.—NATURAL GROWTH OF LOCUST IN AN OLD ABANDONED FIELD OF SOME 60 ACRES IN EXTENT AT FALLS CHURCH, VA.

An isolated clump of some 30 trees, 2 to 4 inches in diameter. These were growing close together and mixed with tall weeds and sassafras bushes. Illustration after clearing the trunks in July when not a single borer was present. In September these were heavily infested.

FIG. 2.—ILLUSTRATION OF A FENCE ROW AT FALLS CHURCH, VA., SHOWING THICK LOCUST GROWTH OF 2 TO 5 INCHES IN DIAMETER NEAR BASE.

None of these trees contained borers in July, 1918, when the underbrush and small branches were cleared from the trees in the foreground. These trees were infested in September.

PROTECTION FROM THE LOCUST BORER.

FIG. 1.—A DENSE GROWTH OF ROOT SUCKERS FROM A FEW OLD TREES THAT WERE REMOVED, ON MINERS HILL NEAR FALLS CHURCH, VA.

This illustrates the type of growth that will not be attacked. In the background to the right of the road are many trees containing 5 to 20 borers each. It illustrates the pruned portion of example 1 (see p. 3).

FIG. 2.—LARGE MATURED LOCUSTS AT VIENNA, VA., SHOWING WHAT CAN BE GROWN IN A LOCALITY WHERE THE BORERS ARE VERY ABUNDANT.

These trees show no evidence of borer injury.

PROTECTION FROM THE LOCUST BORER.

old waste land, the soil of which has been enriched by a thick bed of humus beneath the trees.

Another group (Pl. I, fig. 2) near by contains about 30 trees similar in respect to size and conditions. These have seeded the surrounding soil so that a large clump is formed of dense growth, the outermost being some 6 to 8 feet high. None of these trees have been injured by the borer. A third group (Pl. II, fig. 1), forming a small oval clump, contains 38 trees from 2 to 4 inches in diameter and 15 feet high. They were closely set, averaging 4 feet apart. A few sassafras were intermixed. In July, 1918, not one of these trees contained a borer. August 20, 1918, the plot was thinned, all branches trimmed to 8 or 10 feet, and all weeds and briars removed. The trees were closely watched and many adults were observed ovipositing on them during September.¹ Adults were likewise observed on the isolated trees near by, but none in the dense plats described above.

6. A fence row opposite the eastern field station at Falls Church, Va., is densely matted with wild cherry, honeysuckle, and goldenrod along the sides. Through this grow a dozen locusts (Pl. II, fig. 2) 2 to 4 inches in diameter. None of these trees were infested on or previous to August 15, 1918, on which date all the surrounding growth was cleared from four trees. During September adults were observed ovipositing on these four. Many such fence rows exist wherever locust is grown and explain why so little trouble is experienced on the average farm.

7. Along the south bank of the Potomac River between Difficult Run and Scotts Run many locusts are growing on the wooded slope. This stand is a hardwood mixture composed chiefly of oak, chestnut, hickory, tulip tree, basswood, and butternut. All the locusts are tall, straight poles, reaching to the top of the stand, the trees averaging from 6 to 18 inches in diameter. All show a thrifty condition of growth and no borer defects. A low meadow, sometimes cultivated but now pastured, lies between this river terrace and the water. A few occasional locusts grow here, nearly all of which are infested and badly deformed. No goldenrod occurs in the meadow, and the adult beetles were observed to feed on several species of Eupatorium.

PLANTATIONS ALONG THE PENNSYLVANIA RAILROAD.

From the office of the forester of the Pennsylvania Railroad records have been secured of 42 plantations, comprising nearly 2,000 acres, on which over 1,000,000 trees were planted. These trees were about 2 years old when set out and usually were spaced 8 by 8 or 8 by 10 feet. Many of these tracts were personally examined by the writer, accompanied by Mr. I. T. Worthley, assistant forester. The history of each plantation was ascertained as accurately as possible.

¹ Examination of this plot on Apr. 8, 1919, showed an average of 15 borers per tree.

Some of these tracts are in excellent condition, others almost totally destroyed. It is necessary to describe only a sufficient number of these to give all conditions of culture and past history.

8. Along the line of the railroad, between Philadelphia and Harrisburg, Pa., near Kinzers, are two locust plantations, one on the south side planted in 1904, one on the north planted in 1905. Together they comprise about 25 acres, containing 44,000 trees. As far as can be ascertained, nothing was done in these plots until 1909 and 1910, when thorough prunings took place. The entire tracts were gone over and all the trees trimmed to a single straight stem. All natural growth of other trees or shrubs was thinned out. The writer personally assisted in one of these thinnings in 1910 and at that time noted a few borers in the larger trees. Nothing more was done, but an examination in 1912 showed the borer very abundant and many trees breaking off. At present it is difficult to find a good tree. A few have not been injured sufficiently to cause them to break, but the great majority form a broken tangle. Many root sprouts and suckers are present. These are now meeting the same fate.

9. Near New Brunswick, at Stelton, N. J., are several more plantations in a long strip on the old roadbed comprising some 8 or 10 acres of 13,000 trees. One of these lies along the present roadbed near the station. These trees have been pruned from time to time by the section crews. At present they are badly infested (in the spring of 1918 averaging 10 living borers to a tree) but few broke off until the summer of 1918, when a severe wind destroyed about 5 per cent. These trees average 2 to 5 inches in diameter.

Farther west, some distance from the present roadbed, is a similar stand planted at the same time. No care was taken of it and the trees grew up in thick weeds and other natural, shrubby growth. They are scarcely close enough to produce much natural pruning and all have several large branches. An examination in 1917 and 1918 showed that only a very few trees were infested or ever had been infested. The average diameter is larger than that of those nearer the station.

10. One mile east of Metuchen, N. J., are 10,000 trees planted in 1909, divided into about equal stands, one on the south and one on the north side of the roadbed. The north plot was pruned at two different times, but the years could not be determined. In the spring of 1918 all these trees were badly infested and many were broken off. Many root sprouts and suckers have grown out. The trees on the south side never were pruned or thinned. In the spring of 1918 it was almost impossible to find a borer in the tract; however, a certain amount of old work was present, though not sufficient to mar the trees greatly. These trees were not planted close enough to cause natural pruning and consequently are considerably branched. Very

few root sprouts occur in this tract. They are now producing a more dense shade and natural pruning is taking place. At an earlier date, when the trees were less dense, borers attacked, but as the density increased they probably were repelled before the trees were seriously injured.

The diameter of the stand on the south side averages several inches more than that of the standing trees on the north side. If these trees on the south side had been planted several feet closer, say 6 by 6 instead of 8 by 10 feet, it is the writer's opinion that they would now have unbranched stems with practically no borer defects.

11. On the low-grade freight line between Martic Forge and Columbia, Pa., are five plantations set out in 1906 containing about 150,000 trees. None of these have ever received any attention in the way of pruning. They present conditions varying from such as described at Kinzers (example 8) to almost perfect stands free from borer injury. In no single plantation are the trees all destroyed or all in good condition, but the extremes are found in different parts of each tract. Another factor has been responsible here. A definite correlation exists between those parts of the plantation that have been run over by fire and those parts which were by nature of their position less subject to fire. Where fire has burned over repeatedly, killing the undergrowth, the worst destruction by borers is found. In many places the locusts themselves have been killed. One tract near Shenks Ferry attracted particular attention. It extends from the roadbed across a bottom and up over a hillside. Fire no doubt has gone through the part near the tracks repeatedly, as evidenced by the different ages of scars on the standing trees. Here there is little undergrowth; the locusts are scattered (many have been fire-killed), and all are severely infested. On the hillside and over the crest, fires, for some reason, have not gone through. The locusts have grown up in a dense stand mixed with much underbrush, and oak and chestnut sprouts from the original stumpage. These trees are now in excellent condition; they are tall, straight, thrifty, not branched, and free from borer defects. In some parts the shade of the mixed stand has become so dense that all weeds, briars, and underbrush have been shaded out.

12. Along the railroad between Harrisburg and Huntingdon, Pa., much locust has been planted, and many natural stands occur. The condition of the locust in this region is generally so much better and so much more thrifty that the first examination gave the impression that it is an exceptionally favorable situation for the growth of the tree. It is no doubt true that certain localities are better adapted to the growth of black locust, but the essential factor in this location is considered to be more than purely a favorable situation. The locusts are growing in a narrow belt of river terrace from a few

yards to a mile wide between the foot of the mountain and the Susquehanna and Juniata Rivers. There is much humidity, and all vegetation is vigorous.

It is believed that the good condition of these locusts and the decided absence of injury by borers are due to the fact that the situation is conducive to a rapid growth of underbrush and plants characteristic of river shores which have afforded protection to the locusts. This is further emphasized by the fact that in other localities are found trees of equal size and good appearance, as described at Shenks Ferry (example 11), of the same age. Also in the same locality we find trees deformed and aborted by borer injury, as described in the following paragraph.

13. Along the Juniata River between Newport and Old Ferry, Pa., a continuous plantation of locust set out in 1904 extends for a distance of 2 miles. These trees, especially that part near Newport, show the best stand of all the plantations. After planting no attention was given them until 1914, when they were thinned to afford a view of the river from the trains. At this time most of the stems had reached a sufficient size to be immune from borer attack. The trees nearest Newport now average 5 to 8 inches in diameter. They are straight, free from branches, and about as tall as the telegraph poles near by. The bark is ridged naturally, showing absence of borer injury in the past. Nothing definite as to how these trees grew in the period intervening between planting and first pruning could be ascertained, but from a study of many plots of all ages along the river it is certain that they were intermixed with a dense growth of other shrubs and weeds. Natural pruning and thinning have taken place to such an extent that few living branches are found below the crown, and many trees have been suppressed and killed. Farther back from the railroad, where no pruning at all was done, the dead, suppressed trees and shaded-out branches give a good idea as to how rapidly this process takes place in the tree.

In this 2-mile strip of locusts the borers are serious in two places and have caused many trees to break off or have stunted the growth badly. In one of these places fire has gone through; the other was evidently pruned (as shown by scars) to afford a view of a pond just behind. These last trees were also isolated in the sense that the rows were only two trees deep. These trees average scarcely half the diameter of those in the better parts of the grove.

CONDITION OF TREE NECESSARY FOR BORER ATTACK.

All trees and all parts of the tree are not subject to attack by the borer to the same degree.

Moderately rough bark seems to be an essential condition, since it provides the necessary crevices in which the adults deposit their eggs.

Likewise, until they become $1\frac{1}{2}$ to 2 inches in diameter at the base, trees are not subject to attack unless the bark is rough. On younger trees the borers are found concentrated at the base and near crotches.

For some unknown reason trunks of trees reaching 5 to 6 inches in diameter and over (excepting old brood trees) rarely are found to contain borers. On such trees the larger branches frequently are infested, but such injury is seldom common enough to do much harm or even attract attention. It can be said, therefore, that protection from borer injury is necessary for only a comparatively short period during the tree's growth. Under good growing conditions this time should not exceed 10 years.

In every locust grove that has borers present, certain trees will be found on which they have concentrated. These are called brood trees. The thick, irregularly barked, gnarled appearance and stunted growth will distinguish such individuals. They are often continuously infested until they reach an old age, or 12 to 18 inches in diameter.

HOW TO RECOGNIZE TREES CONTAINING NO BORER DEFECTS.

The larval mine made by the locust borer destroys a certain amount of the growing tissue or cambium and makes a serious defect in the wood. This injury to the cambium accelerates growth to heal it over and produces a swollen or gnarly appearance. Many such defects give the entire stem a roughened, distorted shape. The bark is irregular and scaly. On the other hand, trees that have not been injured by the borer are characterized by very regular bark, which is grooved longitudinally between thick, dark ridges. With a little experience these features can be quickly recognized and until the tree reaches 10 to 12 inches in diameter it is possible to determine accurately whether the borer defect will be found in the wood.

CHARACTER OF GROWTH OF UNINFESTED STANDS.

Not only is the appearance of uninjured individual trees characteristic but pure stands of such trees have a different appearance from those that are damaged. The tops of isolated natural stands have a domelike outline, the innermost trees growing taller and straighter, while root sprouts continuously coming up around the borders form smaller and younger trees which give additional protection to those within. These younger trees are at first too small for infestation, and when they have reached a susceptible size are protected in their turn. The crowns are uniformly shaped and no branches project to break the contours. Planted stands, where the trees are of the same age, are uniform in height, the tops forming a flat outline. No large branches are found on the trunks, but many

small, naturally pruned dead branches are seen and many trees are thinned out naturally as the others increase in size. Few root sprouts appear. Infested plantations are very irregular in outline; broken tops, trees of irregular size, and many root sprouts and suckers are characteristic indications of borer damage.

CONDITIONS UNDER WHICH LOCUST CAN BE GROWN.

From the history of the foregoing tracts it is evident that black locust can be grown profitably under circumstances that require little care, or, in fact, better results are obtained without too much attention. After comparing all the data available it seems that *provision for sufficient shade during the period of growth subject to borer attack* is all that is necessary in order that this tree may be grown successfully.

This can be achieved by some system of close and mixed planting. Experiments of such a character should be tried. In nature it is accomplished by close reproduction coming up around the seed tree, by root sprouts from older trees, or often by the mixture of other plants growing with the locusts. Weeds and vines often form the needed shade, as illustrated by trees in old fence rows. It is very essential that this shade be present after the trees reach $1\frac{1}{2}$ to 2 inches in diameter, and that it be continued until they attain 5 or 6 inches. After this time thinning and pruning can be done with little or no subsequent injury by the borers.

Close planting or thick growth of these trees also is necessary to produce straight, unbranched boles. Trees in the open are always much branched and rather crooked, but those grown in forests are tall, straight, and naturally pruned while the branches are quite small.

That difference in site or locality is not the influencing factor in the growing of uninjured trees is evident from the fact that in every locality examined it was possible to find examples of borer-free and destroyed trees growing 100 yards apart. It is also evident that goldenrod is not necessarily associated with greater damage by the borer, for in the same abandoned field, massed with this plant, were found plots of trees absolutely free from injury and near-by isolated trees badly infested. Again, localities where no goldenrod is growing may have borer-infested trees, the adults feeding on other composites, as illustrated by example 7.

The idea has been advanced that the borers are more abundant in some localities than in others and that this will account for the difference in infestation. This difference can not be sustained, as the beetles are present everywhere within the natural range of this tree. Side by side we find stands of badly infested trees and trees containing no borers. It is rather to be believed that in localities where locust

was planted the beetles present either remained in about the same numbers or increased enormously, according as the condition of the trees retarded or favored their increase. In no new locality where plantations were put out would there be enough beetles present to infest all the trees. They only attack all the trees as they become sufficiently numerous.

CONTROL.

A METHOD OF HANDLING SEVERELY DAMAGED PLANTATIONS.

Many locust plantations have been abandoned and all hope of ever realizing any commercial product given up because of the severe devastation produced by the borers. Such tracts look hopeless with the greater percentage of the trees broken off or killed, but it is believed that they can be reclaimed after several seasons' care by virtue of the sprouting ability of this tree.

It is recommended that all such plantations be gone over and the broken-down and infested trees removed and burned during the winter. Unless otherwise desired it would be necessary to cut out only the living infested trees, because no beetles will breed in the dead ones. Especial attention should be given to the seriously damaged or so-called brood trees. If the cutting out of the infested trees can be done early in November it is not necessary to destroy or burn the wood. The larvæ require living wood for their early development and will not mature in dead material. This not only will reduce the numbers of the insects, but before the sprouts become large enough to be attacked a sufficiently dense stand will have been developed to provide natural protection, as illustrated in Plate III, figure 1.

TREATMENT OF SHADE TREES.

The locust is widely planted for ornamental and shade purposes. It is very desirable for such planting, because of its ability to succeed well in a variety of soils and situations and its rapid growth and good form of crown in the open. We often hear complaints of serious injury by the borer to locust shade trees; this is because such trees are usually grown in isolated situations most favorable for attack.

It has been found that the young borers can be killed readily by the use of an arsenical spray, applied to the bark when the new growth begins to open at the tips of the twigs in the spring. It is necessary to apply this mixture so thoroughly as to cover all parts of the trunk and reach every spot where a larva is working.¹

A thorough application will probably be necessary only every two years unless there are badly infested trees near by which are not treated and form centers of reinfestation. As a rule, spraying will

¹ The presence of a young borer can be determined by the oozing of sap and boring dust from a small hole through the bark. This hole is enlarged as the larva grows.

United States Department of Agriculture,

BUREAU OF ENTOMOLOGY,

L. O. HOWARD, Entomologist and Chief of Bureau.

THE BAGWORM.

(*Thyridopteryx ephemeraformis* Haw.).

By L. O. HOWARD and F. H. CHITTENDEN.

GENERAL APPEARANCE AND NATURE OF ATTACK.

Shade trees, shrubs, and hedges, and in particular evergreens, are much subject to injury by a medium-sized caterpillar which has a curious habit of crawling about on the infested trees in a bag-like case, whence its common name of bagworm or basket worm. In the shelter of these cases the insects undergo all their transformations, after which the bags remain attached to the plants for some time and are conspicuous objects on leafless trees and shrubs in late autumn and in winter. Like the tussock moth^a and the fall webworm^b this species is preeminently a pest on the streets and in parks and private grounds of cities and towns and is even more subject to fluctuation in numbers. It is, however, more limited in distribution and not found as a rule north of southern New York and the central portions of Pennsylvania and Ohio. South of these points it is in certain years very troublesome and the subject of much complaint. Such a year was 1907, when the bagworm attracted greater attention than any other tree defoliator. Numerous complaints were received of injuries in the region mentioned, and especially from the States of New Jersey, Pennsylvania, Maryland, Virginia, West Virginia, Ohio, Indiana, and Illinois. The natural enemies of this insect (see p. 6) were comparatively scarce, and there is a strong possibility of a recurrence of injuries in the years to come.

The general appearance of the bagworm is shown in figure 1, which illustrates the caterpillar when nearly full grown, in its characteristic bag. When removed from its bag it looks as shown in figure 2, *a*, which represents the larva at maturity. At this period in its develop-

FIG. 1.—Bag worm (*Thyridopteryx ephemeraformis*). Natural size (after Riley).

^a *Hemerocampa leucostigma* S. & A.

^b *Hyphantria cunea* Dru.

ment it may attain a length of about three-fourths of an inch. The body is soft in texture and dull brownish or blackish, while the head and thoracic segments are horny and whitish, mottled with dark brown.

ORIGINAL HOME AND PRESENT DISTRIBUTION.

The bagworm is unquestionably native to North America. It abounds in the Southern States, and its proper home—that is to say, the part of the country where it reaches its maximum—is in the Lower Austral life zone. It extends through a considerable portion of the Upper Austral zone, but there are indications that it has gradually spread into this territory from more southern regions." The shade trees of Baltimore, Washington, St. Louis, and other more southern cities are frequently defoliated by this insect. Northward it occurs

FIG. 2.—Bagworm (*Thyridopteryx ephemeraeformis*): a, Full-grown larva; b, head of same; c, male pupa; d, female pupa; e, adult female; f, adult male. All enlarged (from Howard).

through New Jersey and at many points in Pennsylvania, including the cities of Harrisburg, Ellittsburg, Allentown, and Swarthmore. Farther west it has been found at Columbus, Marietta, South Salem, and other localities in Ohio, at several points in West Virginia, at Brooklyn, Ind., in Pecatonica, Thompsonville, and Allendale, Ill., and so on west to Oklahoma. Everywhere south of these localities, except in the immediate Gulf region, it abounds. In the East it is commonly found in New York City and Brooklyn, and at several points on Long Island. In the Hudson River Valley region it has been recorded by Felt at Yonkers and Mount Vernon, and has been collected at New

^a In the main the bagworm is one of those characteristic forms like the wheel-bug (*Arilus cristatus* L.), the Carolina mantis (*Stagmomantis carolina* L.), and the larger digger wasp (*Sphecius speciosus* Dru.), true southern forms which are gradually extending their northward range by following the seacoast or valleys, or, if carried accidentally northward upon railroad trains, establishing themselves at points beyond their former habitat.

FIG. 3.—Arborvitae infested by bagworms. (Original.)

Haven, Conn., but it is not known to be injurious in these localities. It has been sent to the Bureau of Entomology from Springfield, Mass., but probably does not breed there.

FOOD PLANTS.

The bagworm, although a very general feeder, displays a particular fondness for evergreens of all kinds and especially for arborvitæ, hence it seems probable that one or the other of these was its original or normal food plant (fig. 3). The species becomes exceedingly abundant every few years, and at such times it may be found on shade, orchard, and forest trees of nearly all kinds. It is fond of the maples, particularly the cut-leaved and silver maples, and the related box elder; also of the willows, poplars, and mulberry, less so of the elms, and apparently still less so of the oaks. It feeds more or less freely, however, on most other trees and shrubs, and even on many low-growing semi-woody plants, such as elder, mallows (*Hibiscus*), and ragweed (*Ambrosia trifida*). Thus, in the absence of its choice food plants, it is able to subsist on the foliage of almost any of the plants of the character enumerated and which may be available, but it does not seem to live on grasses and herbaceous plants generally.

HABITS AND LIFE HISTORY.

The bagworm overwinters in the egg stage within the old female bags, and for this reason hand-picking in winter time is an efficacious remedy. In the late spring the young hatch from the eggs, crawl out upon the twigs, make their way to the nearest leaf, and immediately begin to feed and to construct cases or bags for themselves. They spin a large amount of silk and attach to it, for additional strength and protection, bits of leaf or of twig, evidently attempting to disguise the nature of the case as well as to strengthen it. The larva is remarkably soft-bodied, except for its head and strong thoracic plates, and it is necessary that the soft abdomen should have some protection. The construction of the bag of an allied species was carefully studied by H. G. Hubbard, and it is a very interesting performance.

The young larva (fig. 4, *a*) cuts off with its jaws a small fragment of leaf which it places between its front legs, gradually forming a pile fastened loosely with silk. When the pile becomes a transverse tangle about as long as the body it is fastened at each end loosely to the surface upon which it rests; then the caterpillar, after placing itself at right angles, dives under the mass, turning a complete somersault, so that it lies on its back, bound down by the fillet. It then twists around and stands upon its feet, having its neck under a sort of yoke (fig. 4, *b*). It makes the yoke into a complete collar, adding bits to each end until

the circle is complete. Then row after row of fragments is added until the case becomes a hollow cylinder (fig. 4, *c*). One end is then closed up and the inside lined with a tough coating of silk, the case being then extended upright and fastened at one end. When it is fully completed the larva crawls away, with the case carried upright like a cap on the upturned end of its body. In the illustration *d* shows a completed bag made by the young larva, tightly appressed to the flat surface, the larva being concealed within. Such bags may frequently

be found on leaves, and are quite puzzling to the uninitiated until the larva pokes out its head and slowly walks off. As the caterpillar grows the case is constantly enlarged, bits of twigs and any other small objects being used to ornament the outside, and these objects will vary with the kind of tree upon which the caterpillar is feeding. While the larva is small it carries its case erect, but when it is larger the case hangs down (fig. 1). The larval skin is cast four times, and during the molting the mouth of the bag is kept closed with silk. There is a small opening in the extremity of the bag through which excrement and cast skins are pushed.

The male bags are smaller than those of the females, reaching a length of about an inch, while those of the females are much larger. Toward the end of August, about Washington, D. C., the larva completes its growth, attaches its bag firmly by a silken band to a twig, strengthens it inside with an additional layer of silk, and within this retreat, which now becomes its cocoon, transforms to pupa with its head downward. The pupal period lasts about three weeks, and then the imago emerges. The male chrysalis works its way out of the lower opening, and the winged moth issues

FIG. 4.—*a*, Newly hatched bagworm before making its case; *b*, same just beginning case; *c*, showing case nearly completed; *d*, completed case, insect concealed within; *e*, larva after first molt. Highly magnified (original).

through the cracking skin, leaving the chrysalis hanging from the bag, as shown at *c*, figure 5. The chrysalis of the female does not push its way at all out of the bag, but the skin cracks and the female gradually works her way partly out, her head reaching the lower end of the bag, (fig. 5, *d*). The males fly about, seeking the bags of the females, and when one is found in which the head of the female is near the end, showing that she has emerged from her chrysalis skin, the male pushes his enormously protrusive and, in fact, telescopic genital apparatus up into the bag to the anal end of the female and fertilizes her. The female then works her way back into the chrysalis skin, gradually filling it with eggs until more than half of it is filled, scattering in among the eggs some of the sparse hairs from her body. Having done this she

forces her shriveled body out of the opening, falls to the ground, and dies. The eggs remain in this way until the following spring, when they hatch, as previously described.

There is thus only one generation annually.

NOTE.—There is a possibility that the bags of this extremely common insect might be made commercially useful. Its silk, from a practical standpoint, has always been ignored, but it is firmer and stronger and more easily spun as carded silk than that of most other native silk cocoons.

FIG. 5.—Bagworm at (*a*, *b*, *c*) successive stages of growth. *c*, Male bag; *d*, female bag. Natural size (from Howard).

NATURAL ENEMIES.

Although apparently well protected from the attacks of birds by its tough case, the bagworm is somewhat extensively parasitized by several forms of ichneumon and chalcids flies, most of them species which affect also similar tree-feeding caterpillars. Prominent among these is the common *Pimpla inquisitor* Say (fig. 6), which, however, more commonly parasitizes the tussock moth and tent caterpillars.^a The related *P. conquisitor* Say is also a parasite of the bagworm and a third species of ichneumon, *Allocota (Hemiteles) thyridopterigis* Riley (fig. 7), is usually the most abundant of all. Four or five individuals of this species commonly infest a single bagworm, spinning for themselves white silken cocoons within the bag.

^a *Malacosoma* spp.

The species last mentioned was for many years credited with being a primary parasite of the bagworm, although recent observations would indicate that it is a secondary parasite when infesting other

FIG. 6.—*Pinpta inquisitor*: Female, from side. Enlarged (from Howard).

FIG. 7.—*Allocota (Hemiteles) thyridopterigis*. Much enlarged (original).

caterpillars. This might be explained by the hypothesis that this parasite oviposits only in cocoons or cases of firm texture, and therefore can not be the primary parasite of an insect which is not provided with a case of that character.

FIG. 8.—*Spilochalcis marix*. Much enlarged (after Riley).

FIG. 9.—*Chalcis ovata*: Adult. Enlarged (from Howard).

Certain chalcis flies also breed in the bags of this insect. These include *Spilochalcis marix* Riley (fig. 8), *Chalcis ovata* Say (fig. 9), and the common little *Dibrachys boucheanus* Ratz. (fig. 10). This last is a hyperparasite, in the case of the bagworm probably secondary, and in the case of other caterpillars a

tertiary parasite. *Habrocytus thyridopterigis* Ashm. (fig. 11) is assumed to be a tertiary parasite on *Allocota thyridopterigis* Riley, when the latter is a secondary parasite of the tussock moth, but may be secondary when breeding in the cases of the bagworm.

REMEDIES.

When the bagworm occurs upon deciduous trees it can be controlled by hand-picking the bags in the winter, but when it affects evergreen trees it is practically impossible to apply this remedy with profit unless the plants are badly defoliated. Therefore for the treatment of evergreens spraying is a necessity.

The methods of controlling shade-tree pests in cities and towns, as outlined in Farmers'

FIG. 10.—*Dibrachys boucheanus*: Adult female and antenna of male. Much enlarged (from Howard).

Bulletin No. 99,^a on pages 25–29, are in part applicable to this species. This bulletin should be read by persons who have experienced trouble from the depredations of the bagworm.

Collecting the bags.—

One of the most important remedies consists in gathering the bags with the contained insect by hand and either burning them or preserving them to liberate the useful parasites which have been previously mentioned. This work may be facilitated by the use of a 12-foot pole pruner or similar appliance, and can be intrusted to those ordinarily unemployed, such as children and aged persons. Where the trees are very tall it will be necessary to use a long ladder. For best results the cooperation of neighbors who are troubled with the same pest should be secured.

FIG. 11.—*Habrocytus thyridopterigis*. Greatly enlarged (from Howard).

^a This publication may be had gratis on application to the Department of Agriculture.

Very considerable immunity from future injury will result by care in the employment of this method. It is particularly useful where only a few trees are infested. The bags are such conspicuous objects on defoliated or bare trees in winter that it is not at all difficult to detect them, but in cases where comparatively few insects are present on evergreen trees they are not so easily seen.

Encouraging the parasites.—When many trees are infested it is advisable to keep the hand-picked bags for a considerable time in receptacles, such as barrels covered with netting, preferably of wire, so that the numerous beneficial parasites of the pest will be able to issue in the spring and assist in the control of the bagworm the following year. One or two holes bored in the bottom of the barrel or box will prevent water from accumulating and drowning the insects. Where the bags can be placed in piles in an open space or inclosure distant from trees and free from disturbance, the young insects, having very limited powers of locomotion, will soon perish of starvation, as they will not be able to reach the trees or shrubs after they hatch.

Spraying with arsenicals.—On evergreen, where the bags are more or less difficult to find, hand-picking can not be advised. A striking instance of the futility of this method under such circumstances was given by Prof. C. V. Riley in his testimony at a conference on the gipsy moth in 1891. He said that he once tried to protect a cedar tree not more than 6 feet high, upon his own grounds at Washington, by hand-picking. He worked for two consecutive months picking off small bags from that tree, the progeny of not more than two females. Almost daily he went to the tree and found fresh specimens which he had overlooked the day before. For evergreen trees, therefore, an arsenical spray is the best remedy. In connection with the story of his experience just related, Professor Riley stated that he had absolutely stopped injury by the bagworm on large trees in the Smithsonian grounds by spraying, and in the summer of 1895 we had a similar experience on the grounds of the Agricultural Department at Washington. Trees sprayed with Paris green at the rate of 1 pound to 150 gallons of water were completely rid of larvæ of the bagworms. It is easier to reach the bagworms on evergreen than on large-leaved deciduous shade trees, such as maple and elm, but if carefully carried out spraying will result in the destruction of the bagworms, so that the collection of the bags in winter will not be necessary. Arsenate of lead at the rate of 1 pound of the prepared paste form to from 25 to 50 gallons of water will be found even more useful than the Paris green, as its greater adhesiveness renders it less likely to be washed off by rains, which in some seasons frequently occur almost daily at the time when the larvæ are beginning to work.

The best time to apply the arsenical is when the eggs hatch, or shortly afterwards, and the best methods of spraying shade trees are discussed in Farmers' Bulletin No. 99.

Approved:

JAMES WILSON.

Secretary of Agriculture.

WASHINGTON, D. C., *December 2, 1907.*

[Cir. 97]

O

THE PERIODIC widespread devastations of the southern pine beetle have been responsible for the destruction of more merchantable pine timber in the Southern States than has any other agency. During a single epidemic it has destroyed timber valued at \$2,000,000, and the present stumpage value of pine killed since 1891 is placed at no less than \$50,000,000.

This tiny beetle kills healthy, vigorous pines of practically all ages and of all species which occur within its range. To do this, it attacks the middle to upper portions of trees and destroys the soft inner bark through which the trees get their nourishment.

Recent investigations indicate that weather conditions have much to do with the abundance and destructiveness or the scarcity of this beetle. Abnormally dry weather seems to be followed by outbreaks, but when the drought is broken by normal or heavy rainfall the beetle may almost disappear. Low winter temperatures also effectively check these beetle outbreaks.

This bulletin describes the insect and its work, its life and habits, and the methods that have been employed to prevent and suppress outbreaks. It supersedes Farmers' Bulletin 1188, The Southern Pine Beetle: A Menace to the Pine Timber of the Southern States.

THE SOUTHERN PINE BEETLE: A SERIOUS ENEMY OF PINES IN THE SOUTH

By R. A. ST. GEORGE, *Associate Entomologist*, and J. A. BEAL, *Assistant Entomologist, Division of Forest Insects, Bureau of Entomology*

CONTENTS

	Page		Page
Beetle outbreaks and their importance-----	1	Natural enemies of the southern pine beetle-----	15
Evidence of beetle attack-----	3	Preventive measures by forestry methods-----	15
The beetle: Its work and its life throughout the year-----	4	Control measures-----	16
Bark beetles likely to be mistaken for the southern pine beetle-----	8	Investigations of outbreaks-----	18
Other beetles associated with the southern pine beetle-----	13		

BEETLE OUTBREAKS AND THEIR IMPORTANCE

EVERY FEW YEARS patches of dying and dead pines may be seen throughout the southern forests. These group killings can usually be attributed to a bark beetle known as the southern pine beetle.¹ Ordinarily it is a rather rare insect, breeding in small numbers in weakened and dying trees. Periodically, however, favorable conditions occur for development of its broods, enabling it to increase in a very short time from a few individuals to countless hordes. Results of recent investigations of the Bureau of Entomology indicate that abnormally dry spells bring about favorable conditions for outbreaks, both by weakening the trees and by creating a more favorable situation in the inner bark for the rapid development of the broods of the beetle. The resumption of normal or heavy precipitation effectively checks such development and very nearly eliminates the beetle from the forest. Low winter temperatures also effectively check the increase of the beetle.

Other agencies which affect the trees, making it easier for the beetles to overcome their resistance, are fire, wind, and lightning. Trees weakened by these agencies, or by mechanical means, constitute a menace to the surrounding forest. Many infestations owe their origin to the attractiveness of a single weakened tree; therefore such a tree should be removed immediately. During periods when the beetles are present in small numbers, abnormal trees enable the bark beetles to establish themselves and breed up to normal numbers, whereupon they again can become destructive, killing groups of surrounding healthy trees. Results of recent studies suggest that trees

¹ *Dendroctonus frontalis* Zimm.; order Coleoptera, family Scolytidae.

on burned areas serve as an attraction to the beetles. This is an additional reason for adequate fire protection. "Blow downs" occurring during the summer, as well as the large quantities of timber normally cut at this time, also create conditions which are a menace to the surrounding healthy trees. This freshly cut or damaged timber is frequently attractive to the beetles.

There is probably no more serious enemy of pines in the Southern States than this beetle. It attacks and kills healthy, vigorous pines

FIGURE 1.—Map showing distribution of the southern pine beetle. The dots represent known centers of infestation

of all species occurring within its range, including those of practically all ages. Extended observations in the Southern States since 1891 have led forest entomologists to place the present stumpage value of pine killed during this time at no less than \$50,000,000. Hopkins² places the value of timber killed by this beetle, during a single epidemic in the South Atlantic and Gulf States in the years

² HOPKINS, A. D. THE SOUTHERN PINE BEETLE: A MENACE TO THE PINE TIMBER OF THE SOUTHERN STATES. U. S. Dept. Agr. Farmers' Bul. 1188, 15 p., illus. 1921.

1908 to 1911, inclusive, at \$2,000,000. In 1922 and 1923 extensive outbreaks of this beetle were again reported or observed, and a vast quantity of timber was known to have been killed. The latest outbreaks took place in 1925, 1926, and 1927. It is well known that only a comparatively small percentage of the destroyed pine was ever salvaged after these epidemics. There is constant danger that the beetle status may again rise to the danger point.

No part of the South is free from this bark beetle. Its outbreaks occur periodically throughout the States of Georgia, Alabama, Mississippi, Louisiana, Texas, Arkansas, Tennessee, Kentucky, South Carolina, North Carolina, Virginia, West Virginia, Maryland, and Delaware. The beetle is also known to occur in the southernmost

FIGURE 2.—A group killing by the southern pine beetle. Appearance of forest area several years after the trees were killed. Note the new growth

portions of New Jersey, Pennsylvania, Ohio, Indiana, Illinois, Missouri, eastern Oklahoma, and northern Florida. Centers of infestation and distribution are shown in Figure 1.

EVIDENCE OF BEETLE ATTACK

Wherever patches of dying or dead pine are found (figs. 2 and 10) which show no evidence of fire or other destructive injury, there need be little doubt that they have been attacked by the southern pine beetle. These trees can be discovered most easily after they have been infested from 10 days to two weeks, when they display a yellowish-green foliage easily seen from long distances. The color gradually turns to a brownish green, and finally, about the time the beetles emerge, to a reddish brown. Pitch tubes often occur along the

middle and upper trunk (fig. 3), but they should not be confused with the larger tubes of the turpentine beetles³ which are found near the ground. (Fig. 15.) Reddish sawdustlike borings can often be found sprinkled along the trunk and on the ground around the base of the tree. A blue-stain fungus which results in lowering the grade of salvaged lumber is always associated with trees killed by this beetle.

FIGURE 3.—Pitch tubes made by the southern pine beetle along the middle trunk. About one-half natural size

(Fig. 4.) The stain quickly extends to the heartwood. (Fig. 5.)

Where any doubt exists as to the species attacking the trees, close examination of the inner bark from the middle of the trunk of the dying trees will positively determine the insect. Here the characteristic winding, S-shaped galleries in the inner bark (fig. 6) and on the surface of the wood (fig. 7) can be found, and by removing a portion of the outer bark the developing broods can be seen. The burrows of the grubs, or larvæ, extend only about half an inch from the parent gallery. Here the larvæ make oval cells in the outer bark and go into a resting stage—that is, become pupæ, before transforming into beetles.

THE BEETLE: ITS WORK AND ITS LIFE THROUGHOUT THE YEAR

Like most insects, the southern pine beetle passes through four distinct stages during its life history. These are the egg, the larva, the pupa, and the adult. The adult (fig. 8, at right) is a small brownish or blackish beetle about one-eighth inch long. It flies throughout the year except during the cooler winter months, when it remains dormant within its galleries beneath the bark. It usually attacks the

³ The red turpentine beetle (*Dendroctonus valens* Lec.) and the black turpentine beetle (*D. terebrans* Oliv.)

FIGURE 4.—Blue-stained, dead trees. The blue stain is always associated with trees killed by the southern pine beetle

FIGURE 5.—Cross section of pine tree killed by the southern pine beetle, showing the associated blue stain extending to heartwood, causing a lowering in grade of merchantable timber

central and upper portions of healthy pine trees (see title-page illustration) as well as weakened trees and occasionally girdled trees and felled logs. Sometimes the attack on standing trees extends almost to the ground. The beetles bore in through the bark and excavate long, winding, S-shaped galleries (fig. 7) which extend through the inner layers of the living bark and mark the surface of the wood.

FIGURE 6.—S-shaped galleries made by adults of the southern pine beetle in the inner, soft bark of pine. Note the oval larval or grub cells about one-fourth to one-half inch on each side of the egg tunnels. Slightly reduced

They deposit their eggs in small niches along the sides of the galleries. From

these eggs young grubs, or larvæ (fig. 8, at left), hatch and feed on the inner bark (fig. 6) until full grown, when they mine into the outer bark, excavate cells, and transform to the pupal or resting stage. In from 7 to 10 days they change to adult beetles and emerge from the tree through tiny exit holes. (Fig. 9.) The beetles then fly to other living trees or occasionally to suitable logs, where they lay eggs to start the next brood. They pass the winter in all stages of development, between the bark and the wood or in the bark itself. The trees that harbor them may be living, dying, or even dead. The more advanced beetles begin to emerge and fly in the spring about the time the redbud⁴

blooms, and the later ones continue to develop and emerge until midsummer when the blackberries begin to ripen. Trees attacked during the previous fall are dead and abandoned by the beetles before the following July. (Fig. 10.) Those attacked during the early part of the summer are dead and abandoned by midsummer of the same year.

⁴ *Cercis canadensis* L.

FIGURE 7.—S-shaped adult-beetle galleries marking the surface of the wood. The larger ones are those of the southern pine sawyer, which often destroys many of the broods of the southern pine beetle. Slightly reduced

From three to five generations occur annually, the number of generations depending on the section of the country in which the beetle occurs, the length of the summer season, the climatic conditions, and the vigor of the beetles. Flight begins in the spring and early summer. The first generation begins with the eggs of the overwintered beetles. When the larvæ from these eggs have developed into adult beetles the first generation is complete. The second generation begins with the eggs laid by the newly formed beetles, and so the process continues until checked by cold weather.

Under favorable conditions a brood can develop from egg to adult in from 30 to 40 days, and the number of beetles can increase as much as 1,000 per cent in a single generation. With the possibility of producing five generations a year, the beetles' capacity for increase and for starting sudden outbreaks is very great.

FIGURE 8.—A, mature larva and, B, adult of the southern pine beetle. Greatly enlarged. (The vertical line indicates the natural size of the beetle.) (Hopkins)

**BARK BEETLES
LIKELY TO BE
MISTAKEN FOR
THE SOUTHERN
PINE BEETLE**

A great many other bark beetles are often associated with the southern pine beetle, but in this bulletin only those most likely to be confused with our most serious enemy of southern pines will be mentioned, namely, the turpentine beetles and the Ips bark beetles.

The Ips bark beetles can be distinguished from the southern pine beetle by the form of the body and the type of tunnel they make. The chief difference between them is seen in the form of the posterior end of the abdomen. In the southern pine beetle this is gradually rounded out in a convex manner (fig. 11, A), whereas in the Ips bark beetles it is scooped out and armed on both sides of this depression with minute toothlike projections. (Fig. 11, C and D.) The egg galleries of Ips are straight or star shaped (figs. 12 and 13), whereas those of the southern pine beetle have a characteristic serpentine, S-shaped appearance. (Figs. 6 and 7.) Furthermore, the grubs of the southern pine beetle bore for a distance of about one-half inch from the parent tunnel and mature in the outer bark, whereas most of the species of the Ips grubs work from 2 to 3 inches from the adult gallery and mature between the inner bark and wood. One very small species of Ips⁵ works near the top of the tree and makes a larval mine about one-half inch long like that of the southern pine beetle, but this work can be readily distinguished because the adult gallery is comparatively straight and not S-shaped. (Fig. 14.)

⁵ *Ips avulsus* Eich.

FIGURE 9.—Exit holes made by new southern pine beetles emerging from the bark.
About two-thirds natural size

FIGURE 10.—Trees abandoned by the southern pine beetle soon after being killed

This difference in habits is made use of in control of beetle outbreaks. The southern pine beetle attacks and kills healthy, vigorous green pines of practically all species and sizes within its range, Shortleaf pine⁶ is probably its most favored host, while longleaf pine⁷ is least favored and rarely attacked. The only trees not attacked by it are the smaller pines below 2 inches in diameter (breast height) and less than 20 years old. The Ips beetles attack dying trees, recently killed trees, and trees weakened by some other agency. Although insects of this latter group may hasten the death of some trees, they seldom are the cause of it except under unnatural conditions. Trees seriously affected by drought, fire, girdling, or primary insects may be successfully attacked by Ips.

The turpentine beetles (fig. 11, B) are similar in form to the southern pine beetle but are about twice as large, being about one-fourth inch in length. They often attack the bases of pines surrounding newly constructed buildings or camps, or anywhere where the cutting or wounding of pines has taken place. They are rarely able to complete their life cycle in green trees and are usually drowned out by the flow of pitch. Normally they breed in stumps and felled logs and are not capable of killing trees. Occasionally, however, they attack weakened, dying, and girdled trees and develop healthy broods in them. In girdled trees they attack below the griddle, where growth has ceased. Trunks of standing trees are seldom attacked by this beetle at points more than 6 feet from the ground. Their work is readily detected, because there are large pitch tubes near the base of attacked trees (fig. 15), and when the bark is removed their long galleries, about a half inch wide and filled with pitch and borings, are exposed. The grubs, working

FIGURE 11.—Adult bark beetles: A. The southern pine beetle (note end of abdomen gradually rounded out in a convex manner); B. one of the turpentine beetles similar in shape to A but larger; C and D, parts of Ips beetles showing posterior end of abdomen scooped out and armed with toothlike projections. The short lines indicate the natural lengths of the beetles. (B, C, and D after Swain)

and develop healthy broods in them. In girdled trees they attack below the griddle, where growth has ceased. Trunks of standing trees are seldom attacked by this beetle at points more than 6 feet from the ground. Their work is readily detected, because there are large pitch tubes near the base of attacked trees (fig. 15), and when the bark is removed their long galleries, about a half inch wide and filled with pitch and borings, are exposed. The grubs, working

⁶ *Pinus echinata* Mill.

⁷ *Pinus palustris* Mill.

FIGURE 12.—Ips bark beetle gallery in center with long radiating larval tunnels, more or less at right angles to the adult egg tunnel. About two-thirds natural size

together, often completely excavate the soft inner bark around a portion of the base of the tree, thereby causing the outer bark to loosen and fall off. Hopkins⁸ has pointed out that this large area, where the wood has been exposed, is often mistaken for a fire scar. These beetles should cause little alarm, as they seldom, if ever, kill the trees.

OTHER BEETLES ASSOCIATED WITH THE SOUTHERN PINE BEETLE

A beetle⁹ commonly associated with trees attacked by the southern pine beetle is one of the long-horned beetles, the larvæ of which are known as round-headed borers or, locally, more commonly called sawyers or fish bait. The adult is an elongated beetle about 1 inch in length, of a mottled-gray color. (Fig. 16.) The male has long feelers or horns (antennæ), which are often two or three times as long as the body. The larvæ may be recognized as elongated, footless grubs of a grayish-white color, which vary in size according to the age of the individuals. When fully mature they are about 2 inches long. (Fig. 16, *i*.) Although this beetle is in no way responsible for the death of trees, it quickly reduces the value of the logs for salvage purposes, since it reduces the lumber from the best grades to No. 1 common. The injury is caused by the grubs, which often destroy some of the broods of the destructive pine beetle while working under the bark, but later ruin the sapwood of the logs if it is not salvaged during midsummer within two weeks after the trees begin to die.

FIGURE 13.—Types of galleries of the southern pine beetle and the associated Ips bark beetles in a 10-inch log. Note the star-shaped and straight galleries of Ips bark beetles on left and the serpentine S-shaped galleries of the southern pine beetle on right, as indicated by arrows

⁸ HOPKINS, A. D. CONTRIBUTIONS TOWARD A MONOGRAPH OF THE SCOLYTID BEETLES. I. THE GENUS DENDROCTONUS. U. S. Dept. Agr., Bur. Ent. Tech. Ser. 17 (1): 153. 1909.
⁹ *Monochamus titillator* Fab.

FIGURE 14.—Work of *Ips avulsus* in bark removed from upper part of tree. Note the comparatively straight adult gallery. About natural size

Other grubs, similar in form to the sawyer grubs and often mistaken for them, attack pines already infested by the southern pine beetle, or some other destructive agency, but these are not serious enemies of the southern pine and therefore are not discussed here. Many other bark beetles, most of them smaller species, work in dying and dead pine, but these are not likely to be confused with our most serious enemy of southern pines and therefore need not be mentioned further.

NATURAL ENEMIES OF THE SOUTH- ERN PINE BEETLE

Birds and parasites are agencies which help to preserve the balance of nature. Woodpeckers, especially, have been repeatedly observed destroying the broods of this beetle. (Fig. 17.) Unfortunately these agencies are not present in our forests in large enough numbers to prevent or suppress outbreaks.

PREVENTIVE MEAS- URES BY FOREST- RY METHODS

Outbreaks of the southern pine beetle usually occur in pure, even-aged pine stands, in which the trees are often attacked when only 20 years old. They rarely occur, however,

FIGURE 15.—Tubes of the black turpentine beetle within 5 feet of ground. Note that they are much larger than those of the southern pine beetle (fig. 3) usually found farther up the tree. About one-half natural size

among pines growing in mixed hardwood stands. General observations throughout the Piedmont region indicate that pure pine stands eventually are transformed to a mixed pine-hardwood composition, because the hardwoods come in in the openings where the pines have been killed by this bark beetle. Wherever possible, the better hardwoods should be encouraged throughout the pure stands of pine so that a mixed type of forest may be developed which will not only be immune to further bark-beetle outbreaks but which also will produce a better grade of softwood lumber. Summer cutting of pines should be avoided whenever possible. Fires in pine areas increase the danger from pine-beetle attack and also destroy the hardwoods within the area, thus tending to create pure pine stands which are always subject to attack.

FIGURE 16.—Work of the southern pine sawyer. Section of trunk showing: *a*, Egg pit in bark; *b*, entrance hole of larva into wood; *c*, pupal cell; *d*, emergence hole; *e*, *g*, *h*, sections of larval mines; *f*, scored surface of wood, scoring done by larva preparatory to entering wood; *i*, larva; *j*, adult. Insect one-half natural size. (Webb)

During years of drought pine areas should be carefully watched for the first evidence of the destructive southern pine beetle, so that measures may be taken to prevent any outbreaks from assuming epidemic proportions.

Since we know that single weakened trees serve as centers of attraction to the southern pine beetle, it is advisable to remove all such trees in order that they may not attract the beetles.

CONTROL MEASURES

Owing to the influence of weather conditions on the southern pine beetle, direct methods of control are recommended with considerable hesitation. It is evident that these beetles increase from a status of rarity to that of destructive abundance in a remarkably short time. On

the other hand, if heavy rains come they disappear quickly and the effort and money spent in control are wasted. It seems reasonable, however, to believe that if the situation were taken in hand at the outset and the insects thoroughly and persistently destroyed, it would be impossible for them to increase as rapidly as if nothing had been done, and considerable timber would thus be saved.

In control work, cooperation between the owners is necessary. It will do little good to clean up one area if timber in areas close to it serves as a source of infestation.

In such control the main objective is to destroy the broods developing in the bark of the tree, thus preventing them from emerging as beetles and destroying other living trees. This can be accomplished by one or another of the following methods:

(1) Felling and logging infested trees and removing the logs to a sawmill, the logs being immediately sawed and the slabs burned. This method is especially recommended where practical, since it permits salvage of the lumber which otherwise would be rapidly destroyed by borers and timber beetles.

(2) Felling and limbing infested trees in a north-and-south direction and, after they have lain in this position a few days, turning them completely over. This measure is effective only in a well exposed sunny situation during midsummer. The insects under the bark on the exposed side of the log are all killed after about three days of hot sun. The logs are turned so that the

FIGURE 17.—Tree attacked by the southern pine beetle. Note holes made by woodpeckers, indicating the high mortality of the brood. Portion of bark removed to show beetle galleries. About three-eighths natural size.

broods under the bark on the opposite side will be exposed to the sun and killed.

(3) Felling and barking infested trees and burning the bark and tops. This method is dangerous in the summer unless extreme precaution be taken to avoid fire. It is, however, a practical control measure in winter.

Whatever the method, it is essential that all the trees containing the living insects be treated. Some of these will still have green foliage or foliage that is only slightly faded, in which case boring dust or pitch tubes on the bark should be looked for as evidence of attack. When the foliage has turned brown, except where it has turned color late in the fall, the insects have already abandoned the trees.

All portions of the main stem up to a minimum of 4 inches in diameter should be treated. The limbs should either be scattered about so that they will be exposed to the sun and quickly dry out, or else they should be burned.

Do not treat trees, except for the purpose of salvage, if the bark shows innumerable small holes such as might be produced by a load of fine shot. The beetles have left the bark of such trees.

Prompt action is necessary. The sooner and the more effectively and persistently the work is carried out, the better will be the results.

Adequate fire protection will do much to reduce the ravages of the beetles, since they are attracted to scorched trees.

INVESTIGATIONS OF OUTBREAKS

The periodic widespread devastations of the southern pine beetle have been responsible for the destruction of more merchantable pine timber in the Southern States than has any other agency. For this reason, extensive investigations have been conducted from time to time to devise measures of control. The most recent facts contributing to a knowledge of the habits of this beetle have been gained as a result of projects carried on at a field station established during 1925, in cooperation with the Forest Service, in the Bent Creek section of the Pisgah National Forest, near Asheville, N. C. Insect studies pertinent to the forest districts of the Piedmont and coastal plains sections of the Southern Atlantic and Gulf States are now being conducted.¹⁰

¹⁰ Information relating to forest-insect work can be obtained by addressing either the Division of Forest Insects of the Bureau of Entomology, U. S. Department of Agriculture, Washington, D. C., or the Forest Entomological Laboratory, Appalachian Experiment Station, Asheville, N. C.

UNITED STATES DEPARTMENT OF AGRICULTURE

DEPARTMENT BULLETIN No. 1182

Washington, D. C.

December 12, 1923

THE IMPORTED PINE SAWFLY.¹

By WILLIAM MIDDLETON,

Scientific Assistant, Forest Insect Investigations.

CONTENTS.

Page.	Page.
Introduction.....	1
Descriptions.....	1
Life history and seasonal history.....	7
Effect of meteorological conditions.....	14
Parthenogenesis.....	15
Parasites.....	16
Hosts.....	17
Distribution in the United States.....	20
Economic importance.....	20
Control.....	21

INTRODUCTION.

A European insect, the imported pine sawfly,² has recently been found defoliating young pine trees in nurseries and on estates in certain of the New England and North Atlantic States.

This bulletin contains descriptions of the principal stages of the species and some information on its life history, host plants, distribution, and importance, together with a brief discussion of its control. The data presented are based on investigations of the sawfly conducted at the eastern field station of the Forest Insect Investigations, Bureau of Entomology, at East Falls Church, Va. The studies began with the receipt of material in August, 1915, and continued until the spring of 1919, when the last adults emerged in the cages.

Inasmuch as the species was an introduced and dangerous one which had not become well established in this country, great care was exercised in the cage work, in order to prevent its escape. A special double-walled, screen-wire insectary was constructed to which entrance was had only through a vestibule, and all the experiments were performed on young trees transplanted into this inclosure.

DESCRIPTIONS.

The following descriptions have been prepared, so that this species can be recognized in its principal stages. Those of the adult are by S. A. Rohwer.

¹ This bulletin was prepared under the direction of S. A. Rohwer, to whom the author is indebted for many helpful suggestions in conducting the investigations. The descriptions of the adult insects are by Mr. Rohwer. The majority of the material from which these studies were made was obtained from Dr. W. E. Britton, State entomologist of Connecticut.

² *Diprion simile* Hartig, order Hymenoptera; suborder Chalcidostogastra; superfamily Tenthredinoidea; family Tenthredinidae; subfamily Diprioninae.

ADULT.

This species is the only North American representative of the genus *Diprion* and may be readily distinguished from the other Nearctic species belonging to the subfamily Diprioninae by the large and densely punctured metascutellum. The specimens which have been examined show very little variation, and all seem to represent the typical form of the species.

Female (Fig. 1, a).—Length 7 to 9 millimeters. Clypeus truncate; head with large, rather close punctures; postocellar area convex and more than twice as wide as long; postocellar furrow well defined; antenna distinctly tapering, the joints much broader than long and with very short rami (apical joints practically without rami), third joint distinctly longer than the fourth; scutum shining, with large, distinct punctures; scutellum and metascutellum opaque and with the punctures closer; tibial spurs simple; pad-like part of sheath elliptical in outline and close to the median line; apical sternite deeply angulately emarginate medially, lancet with nine rows of regular teeth. Black with yellow markings; head black except yellow clypeus, supraclypeal area, and scape; thorax yellow with the following parts black: Sternum, mesepimeron, large spots on prescutum and scutum, lateral part of scutellar area, and metascutellum; abdomen yellow with tergites 3 to 6 inclusive and median spots on following two black; legs, except the somewhat dusky femora, yellow; wings hyaline, venation pale brown with stigma somewhat darker and the costa somewhat paler.

Male (Fig. 1, b).—Length 5 to 7 millimeters. Agreeing with female in general structure; antenna long, the basal rami slightly less than half as long as half the flagellum; hypandrium distinctly punctured, broadly rounded posteriorly; head and thorax more coarsely punctured than in the female; valves of penis, when seen from the side, broad and the ventral margin armed with small, widely separated teeth. Black; venter of abdomen and legs beyond trochanters dark rufous; wings hyaline, venation pale brown, costa paler.

FIG. 1.—*Diprion simile*: a, Female; b, male.

EGG AND EGG SLIT.

The eggs of the imported pine sawfly are about 1.5 millimeters long, with straight sides and bluntly rounded ends. They are oval in cross section, being 0.5 millimeter on the greater transverse diameter and 0.25 millimeter on the lesser transverse diameter. When first laid the eggs are pale whitish blue, translucent, shining, much like gelatin, and somewhat smaller than the above dimensions. After several days, development becomes marked by swelling and the color appears bluish green. When about to hatch they are quite swollen, sometimes as long as 1.75 millimeters, and are dark green. (Fig. 2, b.)

The eggs are laid in slits in the needles (Fig. 2, *a*). Adults emerging early lay in last year's needles, but those emerging when the present year's needles are of sufficient size appear to prefer these. The number of eggs deposited in a single needle varies. In the needles observed, from 2 to 16 eggs were laid, with an average of 8. Where more than one egg is laid the egg pockets adjoin one another, with merely a wall between. The pocket or slit is cut into the convex portion of the needle from an edge at an angle and the tissue composing the walls of the slit is of two textures (Fig. 2, *c*). The basal portions of these walls are the rather tough outer coating of the needle and the apical portions are delicate membranes, apparently made of torn needle pulp, which are pressed together following the laying of the egg, seal it in place, and somewhat conceal it. As the egg increases in size, it parts these lips and becomes exposed. When freshly made the cuts are yellowish green and the needle dark green, but as the eggs mature the cuts become somewhat brownish and the needles yellowish.

FIG. 2.—*Diprion simile*: *a*, Eggs in position in needle; *b*, egg, showing embryo; *c*, needle, showing eggs in pockets.

LARVA.

The following description ³ was made from a full-grown, sixth-instar larva of *Diprion simile* which had been preserved in alcohol. (Fig. 3.)

Length 20 millimeters; maximum breadth, across the metathorax, 4.5 millimeters. The head is circular in outline viewed from in front and the front plane slightly convex

FIG. 3.—*Diprion simile*: Full-grown larva.

viewed from the side; frons and epicranium with a few fine hairs; epistoma with 4 hairs; labrum with 4 hairs; eye disks not elevated, eyes large, lenses convex; antennæ between and slightly closer to pleurostomata than eyes, composed of cone and two disks or partial ring joints; maxillæ with cardo, stipes, palpifer and 4-jointed palpi, galea, and lacinia; lacinia not flattened, rather triangular in outline viewed from apex, armed with large hornlike process or seta on the side toward labium and with rather large setæ forming an apical transverse row on the side of the lacinia parallel with the interior surface of the trophi; labium with palpiger and 2-jointed palpi. Thorax with the tergum composed of areas A, B, C, and D; A excepting in the prothorax, B,

³ The terminology used in describing the larva is that used by the writer in the descriptions of *Neodiprion lecontei* (Fitch) (Jour. Agr. Research, v. 20, no. 10, p. 741-760, 1921), with the changes adopted in "Some suggested homologies between larvæ and adults in sawflies" (Proc. Ent. Soc. Wash., v. 28, no. 8, p. 173-192, 1921).

and C with a few setæ or spines; A of the prothorax bare and constricted and D bare; alar area wanting or incorporated in the spiracular area of the prothorax, large in the mesothorax and metathorax; in the pleurum, preepipleurite large, well defined, and possessing setæ or spines; postepipleurite large, well defined, spined in prothorax but bare in mesothorax and metathorax; prehypopleurite large, triangular, of heavy chitin and sparsely, finely haired; posthypopleurite large with a prominent spined lobe; legs with 4 joints and an apical claw, joint 3 with small soft pad on inner side at apex. Abdomen with the tergum of urites 1 to 9 inclusive composed of areas A, B, C¹, C², C³ and D; A, B, and C² spined, C¹, C³, and D bare; urite 10 consists tergally of the epiproct, which is rather thickly spined and without pseudocerci; spiracular area present on urites 1 to 8, wanting on urites 9 and 10; alar area present on urites 1 to 9 (reduced on urite 9), absent or indistinct on urite 10; the pleurum with preepipleurite and postepipleurite distinct, rather large and spined on urites 1 to 8, reduced, indistinct, but spined on urite 9, wanting or indistinguishable on urite 10; hypopleurite distinct but not divided into prehypopleurite and posthypopleurite, unspined and bearing the uropods on urites 2 to 8 inclusive, indistinct or wanting on urites 1, 9, and 10; uropods well developed on urites 2 to 8 inclusive, not developed on urites 1 and 9, and developed as postpedes on urite 10. The head is black and shiny, with the eyes yellowish and the membranes about antennæ, mandibles, trophi, and between labrum and epistoma whitish. The thorax is yellow, much darkened with velvety black which becomes greenish with growth and age; legs black. The abdomen is yellow, marked as thorax; uropods, postpedes, ventral region, and anus undarkened. Most of the body, the tergum and pleurum of the thorax and abdomen is a mottled black and yellow of rather regular pattern. The depressed portions of the body wall are black and the raised portions are yellow, giving much the appearance of a yellow larva dipped in some adhesive black coloring matter and then allowed to wear off the material darkening its more prominent areas.

LARVAL INSTARS.

In the study of *Diprion simile* the experiments on the larvæ offered an excellent opportunity to determine the number of larval instars and the size of the larvæ during each instar. For convenience, this information is arranged in tabular form below. It will be noted that the species has six feeding instars in larvæ producing female adults and five feeding instars in larvæ producing males. This feature was remarked upon by the writer in treating *Cladius isomerus* Norton in a recent paper on the subfamily Cladiinae⁴ and is probably common to sawflies, since H. C. Severin⁵ records a similar difference in the number of instars between larvæ producing male and female adults in his account of *Neurotoma inconspicua* Norton.

The prepupa of *Diprion simile* is rather well defined and differs from the larva sufficiently in general appearance to be readily recognized.

⁴ Rohwer, S. A., and Middleton, William. North American sawflies of the subfamily Cladiinae. With notes on habits and descriptions of larvæ by William Middleton. In Proc. U. S. Nat. Mus., v. 60, art. 1 (no. 2396), 1922. See p. 19.

⁵ Severin, H. C. The plum web-spinning sawfly. State Ent. S. Dak., Tech. Bul. 1. 1920. See p. 11, 26, tables 2, 3. Severin finds six instars for larvæ becoming males and seven instars for those becoming females. This larval period, however, covers the entire time spent by the larvæ above ground and therefore includes the stage recognized in the present paper as the prepupa.

TABLE. 1.—Average size of larval instars and prepupa of *Diprion simile*, in millimeters.

Instar or stage.	Head.		Body length.		Number of larvae used to obtain averages. ¹
	Height.	Width.	Young.	Full grown.	
	Milli- meters.	Milli- meters.	Milli- meters.	Milli- meters.	
I.....	0.75	0.50	2.75	6.0	11
II.....	.95	.75	4.00	8.0	11
III.....	1.20	1.00	7.00	12.0	8
IV.....	1.60	1.40	8.00	15.0	7
V (male penultimate).....	2.00	1.80	14.00	20.0	5
VI (male prepupa).....	2.00	1.80	² 14.00	³ 13.0	2
V (female antepenultimate).....	1.95	1.65	11.00	18.0	3
VI (female penultimate).....	2.20	2.00	15.00	25.5	3
VII (female prepupa).....	2.20	2.00	² 19.00	³ 12.0	2

¹ These larvæ were reared in isolation and measurements were made at frequent intervals. The dimensions obtained from them are perhaps few in number but accurate and from an instar or stage positively known. The averages obtained from these larvæ were found to apply to larvæ in the general rearing cages.

² Largest. Prepupæ do not feed and grow. They contract with development and the spinning of the cocoon.

³ Smallest.

The different instars of the larvæ of *Diprion simile* are colored as follows:

INSTAR I.—Before feeding: Head whitish, eye spots blackish. Body dull gray. After feeding: Head and legs become blackish.

INSTAR II.—Much the same as the first instar. The dorsum darkens somewhat at the approach of the time to shed.

INSTAR III.—Head black. Body green to bluish gray. The day before molting to the fourth instar the larva begins to show dark dorsal, supraalar, and epipleural lines. This change previous to shedding is doubtless caused by the darker coloration of the approaching instar showing through the skin about to be shed.

INSTAR IV.—Head black. Body considerably darkened, greenish black and yellow, pattern of mottling similar to that of full-grown larva. The larva pales somewhat with age or increased size.

INSTAR V.—Head black. Body velvety black and yellow. The black becomes greenish with age or increased size.

INSTAR VI.—Head black. Body velvety black and yellow.

Each instar seems to have two poorly marked phases; first, a rapid growing period, usually covering the greater part of the instar, during which the body increases in length; and, second, a preparatory period preceding molting during which there is a retardation in growth or even occasionally a contraction. The accompanying diagram (Fig. 4) shows the records of growth of a male and a female larva in body lengths, instars and days, and illustrates this feature.

PREPUPA.

The prepupa of the imported pine sawfly does not feed and its development is accompanied by contraction rather than increase in size. The cocoon is spun in this stage and, protected within, the prepupa changes gradually as it develops into the pupa.

The following description is from a prepupa preserved in alcohol which had not begun the spinning of its cocoon. Prepupæ after spinning do not differ structurally from those which have not spun but are considerably more contracted. A prepupa that is to produce a female adult will measure about 19 millimeters in length before spinning its cocoon, while, after spinning, it will be about 12 millimeters long.

The size of the prepupa is discussed under "Larval instars."

The prepupa of *Diprion simile* is similar to the larva, in structure, with the armature or spines reduced to indistinctness. The head is grayish white, eye disks and apices of mandibles, only, black. The thorax and abdomen are both pale greenish white, with narrow green, dorsal longitudinal stripes and broader green supraalar to alar longitudinal lines which are broken by yellowish spots on B, C¹, C², the pale spiracular area, and the yellowish alar area. Pleurum pale but for a pair of greenish pleural spots. Venter, uropods, postpedes, anus, and epiproct pale, the latter faintly grayish.

COCOON.

The cocoons (Fig. 5) of *Diprion simile* are cylindrical, with hemispherical ends. They are single-walled, of fine texture, rather glossy, fairly stiff, but thin. Cocoons from which males emerge are somewhat smaller than those from which females issue, 7 by 3.5 millimeters being the size of the former, while the latter are 9.3 by 4 millimeters. The cocoons are usually dark brown; sometimes they are pale tan when made but become dark brown within a week or 10 days, and in one instance a whitish cocoon was spun which failed to become colored.

FIG. 4.—*Diprion simile*: Larval growth charts. Stage of larva indicated by large figures. Female has six stages, male five. Growth records are shown for a single female and a single male larva.

In captivity the cocoons were often spun on the needles, and while this may be the rule in nature the writer is inclined to believe that many of the prepupæ of this species, like those of *Neodiprion lecontei*, will make their cocoons in the ground for the sake of the protection thus afforded in winter. This view is supported by a European observation⁶ which records the summer brood cocooning on the branches of the tree while the winter brood makes its cocoons beneath the tree.

The cocoon is spun of a "silken" thread from the vicinity of the mouth.⁷ The end of the cocoon inclosing the caudal extremities of the prepupa is made first, then the anterior end is inclosed by a sidewise motion of the head. After the entire cocoon is formed the prepupa works over the inside, plastering a sticky substance to it. This coating makes the glossy inner surface.

⁶ Britton, W. E. A destructive pine sawfly introduced from Europe. In Jour. Econ. Ent., v. 8, no. 3, p. 379-382. 1915. See p. 381.

⁷ This thread doubtless comes from the apex of the latium, where the writer has observed a similar substance protruding in some prepupæ of *Cimber americana* Leach in his possession.

LIFE HISTORY AND SEASONAL HISTORY.

The experiments to obtain data on the life history and seasonal history of *Diprion simile* were distinct from those pertaining to the choice of host plant and also from those pertaining to the ability of the insect to reproduce parthenogenetically. *Pinus sylvestris* was chosen as a host for this work because it seemed to be favored by the sawfly and because it had been shown to be a successful host in this locality by previous tests. In no instance was copulation observed, but opportunity for fertilization was provided by keeping constantly with each female at least two males in excellent condition. These experiments were made with small cylinder cages of screen wire on a steel frame, 14 inches in diameter by 24 inches high, the top of which was removable to permit easy access to the interior for observation or handling the material. These cages were placed over young trees of *Pinus sylvestris*, and adults of *Diprion simile* were then liberated within. There were also certain special life-history and seasonal-history studies, particularly some in which observations were made on larvæ of this sawfly reared in isolation to determine the number and duration of the instars and the influence of the sex of the individual upon this phase of development.

FIG. 5.—*Diprion simile*: Cocoons. *a*, Occupied by female, unopened; *b*, occupied by male, unopened; *c*, after emergence of female; *d*, male cocoon, showing emergence hole of a chalcid parasite; *e*, female cocoon, showing emergence hole of a dipterous parasite.

The data presented in the following account of *Diprion simile* show some singular likenesses and contrasts with the life of the related *Neodiprion lecontei*.⁸ It would be well to emphasize that the imported pine sawfly has the same apparent division of the emergence period into Broods A and B. It differs somewhat from Leconte's sawfly in the course of its issuance, a fact which may be explained either as inherent in the species or as caused by change of environment. The climate and other natural factors in the United States may be so different from those to which it is accustomed that it is unstabilized here. Should this prove true, the situation with regard to *Diprion simile* may become very different within a few years, and the insect may even become a more serious problem than it now promises to be. The imported pine sawfly has important life-history and seasonal-history advantages, and if it can become a recognized depredator while unstabilized by the conditions of this new environment, the destructive possibilities of the species when it is properly adjusted will be necessarily greater. In that case it may possibly present a new life history and seasonal history.

ADULT EMERGENCE.

The emergence of adults from the cocoons of a single colony of the larvæ of *Diprion simile* occurs over a considerable period, approximating a year. This emergence is not continuous, but divides itself

⁸ Middleton, William. Leconte's sawfly, an enemy of young pines. In Jour. Agr. Research, v. 20, no. 10, p. 741-760. 1921.

into a period early in the year and a period late in the year. The adults are separated into Brood A, those issuing in the first emergence period following cocooning, and Brood B, those issuing in the second emergence period following cocooning.

The early emergence period occurs from April to June and is not distinctly separated from the later emergence period, which occurs during July and August. The rearing cages, however, show a marked decrease in the number of adults coming from cocoons during June.

The following summary of records of the issuance of adults shows the duration of this period and its tendency to divide into two parts, the months of maximum emergence being April, May, and July.

Larvæ and pupæ collected in August, 1915:

1 adult emerged in August, 1915.

77 adults emerged in May, 1916.

5 adults emerged in June, 1916.

1 adult emerged in July, 1916.

Cocoons collected in April, 1917:

16 adults emerged in April, 1917.

95 adults emerged in May, 1917.

2 adults emerged late in June, 1917.

2 adults emerged in July, 1917.

1 adult emerged in April, 1918.

Larvæ cocooning in June of various years:

2 adults emerged in June of the same year in which they cocooned.

24 adults emerged in July of the same year in which they cocooned.

1 adult emerged in August of the same year in which it cocooned.

7 adults emerged in April of the year following that in which they cocooned.

8 adults emerged in May of the year following that in which they cocooned.

LONGEVITY OF ADULTS.

The average length of life for adults of *Diprion simile* was 6.6 days, with a range from 3 to 11 days. The life of the female was slightly longer than that of the male, the former averaging 7.5 days, with a range from 5.5 to 11 days, while the latter averaged 5.8 days, with a range from 3 to 9 days. The life of female adults known to have laid eggs was somewhat longer than the life of those not known to have laid eggs. Females laying eggs averaged 7.75 days, those not known either to have laid eggs or not to have laid eggs averaged 7.66 days, while those known not to have laid eggs averaged 7 days. The two shortest-lived males were the progeny of parthenogenetic females. Only two of these were available for this study, and these lived, one 3 and the other 4 days.

PROPORTION OF SEXES.

The males seem somewhat to exceed the females in number, and from a series of 227 adults issuing from cocoons in certain of the cages, 134, or 59 per cent, were males, while 93, or approximately 41 per cent, were females.

MATING.

Mating was not observed. At different times three males were caged in glass vials with a single unfertilized female without their paying her the slightest attention. This, however, was probably due to the close confinement of the sawflies under observation, which tended to distract them and prevent or discourage natural behavior. The fact that from the life-history and seasonal-history

experiments both sexes were obtained in the progeny, while, as will be stated farther on, in parthenogenetic experiments only male adults were obtained, indicates that mating is a regular occurrence.

OVIPOSITION.

The eggs are laid in the needles of the pine and the adults emerging from their cocoons early in the year oviposit in the old or last year's needles, while the adults emerging later in the year lay their eggs in the needles of the current year. The following note, made in observations on the habits and activities of the adults in the life-history and seasonal-history cages, shows the increase in attractiveness of the current year's needles.

All the females, except the one emerging at the latest date, i. e., June 23, 1916, laid their eggs in the old or grown needles, but this female, possibly because of the advance of the younger needles, delayed her oviposition several days and then apparently first attempted egg laying in these new leaves. After two slits or thrusts, possibly having succeeded in laying an egg in one of these, she began on the old needles and finished her work on them.

This note seems to indicate a preference for the young needles which were still too small for this female to oviposit in them.

The following account, taken from cage notes made during these experiments, gives an accurate description of the method of oviposition:

The female begins laying eggs near the base of the needle and works toward the tip, making a new incision for each egg. In oviposition the mandibles usually close upon the needle; the antennæ extend forward and are usually quiet and the wings are folded, at rest on the back. The ovipositor (Fig. 6, *b*, *c*) is exerted from between the right and left halves of the sheath, forming an angle of about 60° with the needle. It is worked into the needle and straightened until as it disappears it forms a right angle with the needle, saw edge forward (Fig. 6, *a*).

The posterior margin of the seventh sternite is membranous and is produced medially to form a sort of trough (*tr*) for conveying the eggs into the slit prepared for them. The sides of the trough are supported by a pair of diverging chitinous pieces or more rigid areas. This trough during oviposition is immediately in front of the slit that is being cut and serves as a support to the abdomen, maintaining a constant distance between body and needle and contributing to a rigidity and firmness of the base of the working parts in their relation to the needle. Posterior to the trough and basad of the sheath proper there is a pair of rather small rectangular plates or trowels (*t*) (Fig. 6, *d*), one plate at each side, attached to the base of the basal portion of the sheath (*sh*¹) and appearing similarly attached to the base of the lance (*le*). These plates were observed for the first time in this insect and are not found in *Neodiprion lecontei*, which belongs to the same subfamily, nor has the writer been able to discover similar developments in sawflies outside of the Diprioninae. The function of these plates or trowels is apparently to support and mold the torn needle fiber into walls and cover for the egg pocket. The sheath (*sh*¹ and *sh*²) is posterior to these plates (*t*) and its right and left halves cover respectively the right and left sides of the needle at the point of incision. During the sawing there are alternate contractions and expansions of the pleurum of the ninth urite which cause the sheath to flex or bend at the juncture of the basal (*sh*¹) and apical (*sh*²) portions, and a side-to-side oblique movement of the trough (*tr*). The saw is worked or dragged until the trough (*tr*) reaches the mesothoracic legs. The puncture is then complete and the abdomen is slightly elevated. The trough (*tr*) and the trowels (*t*) are now pushed back together, smoothing off the torn fiber at the mouth of the pocket and bringing together the walls of the cavity at its opening. The saw is withdrawn at the same time. It is not visible except through the needle tissue, being hidden by the trowel (*t*), and its withdrawal is accomplished or accompanied by a backward, upward folding movement. The entire egg-laying process consumes about two minutes.

Oviposition occurs over several days, the female going from needle to needle. Table 2 indicates the number of eggs laid and gives an idea of the number of needles upon a tree that are attacked.

FIG. 6.—*Diprion simile*: Ovipositor. *a*, Apex of abdomen of female adult in position over nearly completed egg slit (enlarged); *b*, three-quarters ventral view of apex of abdomen of female adult (enlarged); *c*, ventral view of portion of egg-laying apparatus of female adult (much enlarged); *d*, side view of trowel, from ovipositor of adult female (greatly enlarged).

TABLE 2.—Oviposition records of *Diprion simile*.

No.	Number of needles oviposited in and number of eggs in each.										Total number of eggs.	Host plant.
	1	2	3	4	5	6	7	8	9	10		
1...	9	0	3	9	5	-----	-----	-----	-----	-----	26	<i>Pinus austriaca</i> .
2...	9	-----	-----	-----	-----	-----	-----	-----	-----	-----	9	<i>P. laricio</i> .
3...	11	16	11	6	9	-----	-----	-----	-----	-----	53	<i>P. sylvestris</i> .
4...	12	2	7	6	-----	-----	-----	-----	-----	-----	27	Do.
5...	12	2	3	5	12	11	8	2	5	8	68	Do.
6...	5	10	8	6	12	4	13	14	7	2	81	Do.

Of these experiments the first and second were failures owing to the host plant chosen, and the eggs in the third and fourth were laid by virgin females, and therefore will be treated farther on, in considering the experiments on parthenogenetic reproduction in this species.

Table 3 shows the length of time elapsing between the emergence of the adult and oviposition.

TABLE 3.—*Preoviposition period of adults of Diprion simile.*

No.	Date adult emerged.	Date oviposition began.	Time in days from emergence to oviposition.	Host plant.	Remarks.
1...	May 8...	May 9...	1	<i>Pinus sylvestris</i>	
2...	May 11...	May 11...	0	do.....	
3...	May 12...	May 16...	4	<i>Pinus austriaca</i>	Host plant unfavorable.
4...	May 8...	May 10...	2	<i>Pinus sylvestris</i>	Unfertilized female.
5...	May 9...	May 11...	2	do.....	Do.
6...	June 20...	June 23...	3	do.....	New needles attractive.

From the consideration of all the available data the writer is of the opinion that a fertilized female will begin laying eggs soon after fertilization, provided material suitable for oviposition can be found. Such was the case with the first and second females in Table 3. Unfertilized females will begin laying after a two-day period during which they await mating, provided that suitable material is convenient, as shown by the fourth and fifth females, which were used in experiments on parthenogenetic reproduction. A period of longer than two days presumably indicates an unfavorable host, unfavorable condition of host, or other disturbing factor. In the instance of female No. 3, the four-day delay is believed to indicate that *Pinus austriaca* is an unfavorable host, a conclusion borne out by other observations discussed under "Hosts." In the instance of female No. 6, the three-day delay is believed to have been caused by the advanced condition of the young needles, which, though not of sufficient size for successful oviposition, were yet large enough to attract her and thus lengthen the time between emergence and oviposition.

FERTILITY AND INCUBATION.

In only one of the experiments was an accurate record of the eggs from laying to hatching kept, and here it was found that 48 eggs out of 69, or 69.5 per cent, hatched. The period of time elapsing between oviposition and hatching is called the incubation period. Table 4 records the results of investigations to determine the length of this period.

TABLE 4.—*Incubation period of Diprion simile.*

Date laid.	Date hatched.	Length of period. ¹	Host.	Remarks.
		<i>Days.</i>		
May 9 to 15.....	May 22 to 26.....	11-13	<i>Pinus sylvestris</i>	Unfertilized female. Do.
May 10.....	May 22.....	12	do.....	
May 11 to 15.....	May 25 to 29.....	14	do.....	
Do.....	do.....	14	do.....	
May 16.....	May 29.....	13	<i>Pinus austriaca</i>	
June 23 to 24.....	July 1 to 2.....	8	<i>Pinus sylvestris</i>	

¹ In counting the number of days in the incubation period, it was assumed that the first eggs laid were the first to hatch and that the latest laid were the latest hatched.

For eggs laid about the middle of May (May 9 to 19), from 11 to 14 days were required for development to the point of hatching, averaging 12.8 days. For eggs laid in late June, 8 days sufficed for development. Fertilized and unfertilized eggs, apparently, did not differ in the duration of the incubation period.

LARVAL DEVELOPMENT.

The larvæ of the imported pine sawfly shed their skins as they develop. The periods between moltings of the larva are called instars. The number of instars, the difference in the number between larvæ producing female adults and those producing males, and the size and appearance of the larvæ in each of the various instars have been treated under "Larval instars."

Table 5 shows for a number of larvæ the time spent in each instar, the average length of the instar, and the total length of the larval or feeding period. This table is arranged to show sex and date of hatching of each individual or group of individuals, because the sex makes a difference in the number of instars, and the rate of development of the larvæ appears to be accelerated as the season advances.

TABLE 5.—*Number and length of larval instars of Diprion simile, showing individual's date of hatching and sex.*

Date egg hatched.	Sex.	Length of instar.						Total.
		Instar I.	Instar II.	Instar III.	Instar IV.	Instar V.	Instar VI.	
		<i>Days.</i>	<i>Days.</i>	<i>Days.</i>	<i>Days.</i>	<i>Days.</i>	<i>Days.</i>	<i>Days.</i>
May 8 ¹	Male.....	6	5	3	6	12		32
Do. ¹		6	5	4	5	11		31
May 9 ¹		7	5	4	4	9		29
May 12 ¹		6	4	4	4	11		29
Do. ¹		6	4	4	4	10		28
May 22 ²		7	4	4	6	2		23
May 25 ²		6	5	5	4	6		26
Do. ²		4	7	7	5	3		26
Extremes.....		4 to 7	4 to 7	3 to 7	4 to 6	2 to 12		23 to 32
Average early.....		6	4½	4½	4½	8		28
July 1.....		4	5	2	3	2		16
Average mid-year.....		4	5	2	3	2		16
May 9.....	Female..	9	4	5	3	6	10	37
May 12.....		6	4	4	2	5	9	30
Do.....		6	4	4	2	4	9	29
Extremes.....		6 to 9	4 to 4	4 to 5	2 to 3	4 to 6	9 to 10	29 to 37
Average early.....		7	4	4½	2½	5	9½	32

¹ Isolated for stages.

² Cage work.

LARVAL HABITS.

The larvæ from the time of hatching to the latter part of the second instar or the beginning of the third feed only on the exterior of the needles, especially from the angles or sides. They then begin to consume the entire needle. At about the fourth or fifth instar they attack the young shoots, feeding on them basally to an extent sufficient frequently to cause the shoots to die and fall. This habit of feeding on the shoots is not essential to the development of the

larvæ, since caged individuals supplied with needles complete their growth successfully and become adults; nor does it, among those larvæ practicing it, supplant needle eating.

PREPUPAL AND PUPAL PERIOD.

When the larvæ have finished feeding they evacuate their alimentary tracts and shed their skins, becoming prepupæ. They then crawl about until each finds a suitable place and spins its cocoon. The length of time spent in the prepupa stage varies considerably. When adults emerge the same year this stage may be as brief as 10 days, when they emerge the following year it may be as long as 341 days.

Table 6 records the results of some experiments to determine the length of the prepupal period and the time spent within the cocoon. The prepupal period as recorded in this table is not separated from the pupal period. It is difficult to determine the exact duration of the pupal period. It seems to be brief and the pupa is seldom obtained by cutting open cocoons, the insects being nearly always found in the prepupal or in the unemerged or immature adult stage.

TABLE 6.—*Dates of becoming prepupa, of cocooning, and of emergence.*

ADULTS EMERGING THE SAME YEAR IN WHICH THEY COCOON.

Individual No.	Host plant.	Date individual became prepupa.	Date cocooned.	Days between time insect became a prepupa and cocooned.	Date adult emerged.	Days in cocoon.	Days between time insect became prepupa and emerged as adult.
1	<i>Pinus cembra</i>	June 29, 1917	July 2, 1917	3	July 9, 1917	7	10
2	<i>P. virginiana</i>		June 29, 1917		July 7, 1917	8	
3	<i>P. strobus</i>		June 18, 1917		June 27, 1917	9	
4	<i>P. monticola</i>		June 29, 1917		July 9, 1917	10	
5	<i>P. divaricata</i>		do.....		do.....	10	
6	<i>P. sylvestris</i>		June 22, 1917		July 5, 1917	13	
7	do.....	July 3, 1916	July 7, 1916	4	Aug. 1, 1916	25	29
Average.....				3.5		11.7	19.5

ADULTS EMERGING THE FOLLOWING YEAR.

8	<i>P. flexilis</i>		July 9, 1917		Apr. 15, 1918	280	
9	<i>P. taeda</i>	July 11, 1917	July 12, 1917	1	May 2, 1918	294	295
10	<i>P. monticola</i>		June 29, 1917		Apr. 24, 1918	299	
11	<i>P. divaricata</i>		do.....		May 5, 1918	310	
12	<i>P. virginiana</i>		do.....		May 6, 1918	311	
13	<i>P. sylvestris</i>		June 22, 1917		May 3, 1918	315	
14	do.....	June 15, 1918	June 17, 1918	2	May 22, 1919	339	341
Average.....				1.5		306.8	318

SUMMARY OF LIFE HISTORY AND SEASONAL HISTORY.

Variations in the length of the several stages of *Diprion simile*, due to known factors such as season, sex, etc., or to unknown factors such as what determines the prepupa to emerge the same or the next year, cause variations in the total length of the insect's life cycle. Table 7 indicates the various lengths of life and periods to be expected.

TABLE 7.—Life chart (average) of *Diprion simile*.

Period.	Progeny of adults issuing in the spring.				Progeny of adults issuing in summer.			
	Adults emerging same year as coo- cooning.		Adults emerging the next year after cocooning.		Adults emerging same year as co- cooning.		Adults emerging next year after cocooning.	
	Male.	Female.	Male.	Female.	Male.	Female.	Male.	Female.
	Days. 12.8	Days. 12.8	Days. 12.8	Days. 12.8	Days. 8	Days. 8	Days. 8	Days. 8
Incubation period....								
Larval period:								
First stage.....	6	7	6	7	4		4	
Second stage.....	4 $\frac{1}{2}$	4	4 $\frac{1}{2}$	4	5		5	
Third stage.....	4 $\frac{3}{8}$	4 $\frac{1}{2}$	4 $\frac{3}{8}$	4 $\frac{1}{2}$	2		2	
Fourth stage.....	4 $\frac{1}{2}$	2 $\frac{3}{8}$	4 $\frac{1}{2}$	2 $\frac{3}{8}$	3		3	
Fifth stage.....	8	5	8	5	2		2	
Sixth stage.....		9 $\frac{1}{2}$		9 $\frac{1}{2}$				
Total.....	28	32	28	32	16	1 18	16	1 18
Prepupa (including pupal period).....	10	10	341	341	29	29	295	295
Adult period.....	5.8	7.5	5.8	7.5	5.8	7.5	5.8	7.5
Total.....	56.6	62.3	357.6	393.3	58.8	62.5	324.8	328.5

¹ Obtained by rule of proportion, as no isolation for length of larval stages gave female adults.

The accompanying chart (Fig. 7) gives a general view of the activities of this species covering a period of about four years, especially the relations existing between stages, broods, generations, and colony periods, and shows the possible opportunities of the species to inbreed and crossbreed among the various colonies, broods, and generations.

EFFECT OF METEOROLOGICAL CONDITIONS.

The notes on the effects of meteorological conditions on this species are incomplete and do not lead to any definite conclusions with regard to the part climate will play in its establishment or in the restriction of its range in America. The small number of adults obtained in the rearing experiments undertaken under the life-history, seasonal-history, and host-plant work gives the impression that the climate of the section of the country where this work was undertaken, East Falls Church, Va., near Washington, D. C., is rather unfavorable for *Diprion simile*, while the apparent establishment of the species in New England, New York, and New Jersey points to the suitability of a somewhat cooler climate.

In the cage work, where the rapidity of development was noted, it was observed to be accelerated by increased temperature.

Table 8 records the duration of stages and instars in days for two colonies of *Diprion simile* between which there was a difference of 46 days in the date of egg laying, and shows the average temperature during each of the stages or instars.

TABLE 8.—*Effect of time of year and temperature on rate of development of the egg and larval instars of Diprion simile.*

Stage.	Earlier colony.			Later colony.		
	Date stage or instar began.	Time in stage or instar.	Average temperature in stage or instar.	Date stage or instar began.	Time in stage or instar.	Average temperature in stage or instar.
Egg.....	1916 May 9	Days. 13	° F. 61.61	1916 June 24	Days. 7	° F. 72.85
Larval instar:						
I.....	22	7	68.78	July 1	4	71.25
II.....	29	4	64.50	5	5	72.70
III.....	June 2	4	66.25	10	2	77.75
IV.....	6	6	64.33	12	3	79.00
V.....	12	5	67.90	15	2	75.25
VI.....	17	12	70.46	17	7	76.28
Prepupa.....	29			24		
Total.....		51	66.19		30	74.51

Some scattered observations record the larvæ feeding close to the trunk on a cool and rainy day (temperature 60° F., humidity 87 per cent), feeding vigorously and spread out on a warmer and less humid day (temperature 69° F., and humidity 68 per cent), and clustered on needles close to the bole on a still warmer and drier day (temperature 85° F., humidity 40 per cent). These observations would seem to indicate a rather low optimum temperature, somewhere between 69° and 85° F., with a rather delicate responsiveness to any considerable variation in temperature, and the fact that the larvæ were observed to be rather slow in developing during a damp period and inactive during a cool, rainy spell (temperature 56° to 71° F. and humidity 85 per cent) tends to show a similar sensitiveness to humidity. The observations, however, are too few and were made in too restricted an area to be of much value.

PARTHENOGENESIS.

Diprion simile can reproduce parthenogenetically; that is, eggs laid by virgin females are fertile and hatch, producing larvæ which grow, become prepupæ, spin cocoons, and finally emerge as adults. All the adults thus far obtained from eggs of unfertilized females, however, are males, a result believed usual for parthenogenetic reproduction in sawflies.

Females that have not mated appear to wait two days before commencing oviposition, as shown in Table 3, but after this period passes they begin to lay unfertilized eggs. Two of the experiments performed to obtain these unfertilized eggs were successful, one female laying 53 eggs, the other 27. A comparison of the number of eggs laid parthenogenetically with the number laid by a normal fertilized female indicates that the virgin female lays only half as many eggs as the fertilized female. In the two parthenogenetic experiments in which eggs were deposited about 72½ per cent of the eggs hatched, which is approximately the same as the percentage of hatching obtained from eggs of fertilized females. The mortality among the

FIG. 7.—*Diprion similis*: Chart showing in a general way the life history and seasonal history data obtained in studies of two separate lots of *Diprion similis* material, one received in August, 1915, which was under observation during 1915 and 1916, and the other received in April, 1917, and used in experiments during 1917, 1918, and 1919. It shows the division of the adult emergence period into at least two broods, A and B. Two adults, one emerging from the cocoon in July, 1916, and the other in April, 1918, seem to have been somewhat retarded, and these are indicated by a superimposed asterisk.

larvæ, however, was higher in the progeny of virgin females, and few survived to become adults. It may be that larvæ wanting male parents are lacking in vitality and when developing to the adult produce an insect with less viability. This is indicated by the death of a great percentage of the larvæ under apparently favorable conditions and the shorter life of adults from eggs laid by unfertilized females.

PARASITES.

Only two species of natural enemies were obtained from the experiments performed at the eastern field station, East Falls Church, Va. These were *Dibrachys nigrocyaneus* Norton and a species of *Eurytoma*. Britton and Zappe⁹ record the following: Hymenoptera: *Dibrachys nigrocyaneus* Norton, *Monodontomerus dentipes* Boheman, *Dibrachoides verditer* Norton, *Delomerista* n. sp., *Cerambycobius* sp. (probably new), *Eurytoma* sp., *Hemiteles utilis* Norton. Diptera: *Exorista petiolata* Coquillett.

All parasites specifically identified are native species, with the exception of *Monodontomerus dentipes* Boh., which is a European species already recorded from the United States.

Dibrachys nigrocyaneus was rather abundant, much more so than any of the other species, and Britton and Zappe state that only the first three species recorded in their list given above were reared in sufficient numbers to indicate that they are at all effective in holding the pest in check.

Table 9, from Britton and Zappe, gives the general results of an examination of cocoons collected during the winter of 1916-17.

A short experiment on the length of life of *Dibrachys nigrocyaneus* Norton was performed. From 65 adults caged without food all the males died within three days and all females

⁹ Britton, W. E., and Zappe, M. P. The imported pine sawfly. In Conn. Agr. Exp. Sta. Bul. 203, p. 273-290, 1917. See p. 283.

within eight days; and from 40 adults supplied with cotton soaked in sugar water all the males died within three days, but all the females, except 1, were living on the thirteenth day, and all were not dead until the seventeenth.

TABLE 9.—Cocoons of *Diprion simile* collected during the winter of 1916-17 and examined for parasites.

Condition.	Number.	Per cent.
Adult sawflies had emerged.....	1,321	41
Parasites had emerged.....	1,210	37
Torn open and eaten.....	191	6
Dead.....	518	16
Total.....	3,240	100

Several experiments were made with *D. nigrocyanus* Norton, with the object of observing oviposition and studying the relation of the parasite to this host. All resulted in failure, however, for no parasitism was obtained, although the chalcids endeavored to oviposit, usually more than once, in the cocoons supplied. Possibly *D. nigrocyanus* does not attack *Diprion simile* in the cocoon stage.

The presence and activity of these parasites must not lead to the conclusion that they will bring about an effective natural control of *D. simile*. If the permanent establishment of this pest is to be prevented, or if its multiplication and work are to be checked, artificial control measures, such as those recommended, must be applied.

HOSTS.

If the imported pine sawfly is able to feed on a number of species of pine this will be an important factor in the establishment of the species in this country, and it is extremely desirable to know if it can and will use our native pines successfully. Restricted to introduced ornamental species of pines, *Diprion simile* would at worst only become a nursery and estate problem, but if it will attack our American pines it may become a serious forest problem. At present the sawfly seems to be confined to young pines of imported species and has only been recorded from nurseries and estates.

To determine the range of host plants which this insect can successfully utilize, two series of host-tree experiments were conducted. One was to determine the ability of *Diprion simile* to use a given host successfully from oviposition to the production of adults. The other was to determine the preferred host plants of the sawfly, if there are any. In the first series of experiments a number of adult males and females of *D. simile* were confined upon a particular species of young pine to observe its suitability. In the experiments on choice of host species a quantity of small native and introduced pines were transplanted into the large screen-wire insectary previously mentioned and a number of males and females liberated within the inclosure.

Table 10 gives the results of the experiment to determine the imported pine sawfly's choice of host plants. Fifty-four male adults and fifty-nine female adults of *D. simile* were liberated in the insectary,

into which the following species of young pines had been transplanted.

Species.	Number of trees placed in insectary.	Species.	Number of trees placed in insectary.
<i>Pinus austriaca</i>	2	<i>Pinus ponderosa</i>	5
<i>Pinus cembra</i>	2	<i>Pinus resinosa</i>	3
<i>Pinus divaricata</i>	2	<i>Pinus strobus</i>	5
<i>Pinus flexilis</i>	3	<i>Pinus sylvestris</i>	15
<i>Pinus laricio</i>	4	<i>Pinus taeda</i>	1
<i>Pinus monticola</i>	3	<i>Pinus virginiana</i>	3

Twelve species and a total of forty-eight trees.

In Table 10 the host trees are arranged according to the number of trees of each species that were chosen by the females for oviposition and the suitability of these pines as hosts is indicated by the number of individuals of the sawfly that were able to develop through the various stages from egg to adult.

Frequency of selection for oviposition has been considered a somewhat better guide to the appeal of the various pines to the sawfly than the numbers of the different stages of the insect obtained. This is especially true where the number of adults obtained exceeds a third of the total number of larvæ, since the larvæ after hatching were cage-reared, and, although protected from parasites, doubtless experienced a number of difficulties due to handling and to their artificial surroundings. The species of pines believed most favorable for *D. simile* are therefore placed at the beginning of the table.

TABLE 10.—Experiment to determine the host-plant preference exhibited by *Diprion simile*.

Species and serial number of host tree.	Number of trees of species chosen.	Eggs.	Larvæ.	Cocoons.	Adults.
<i>Pinus sylvestris</i> , trees 1, 20, and 29.....	3	¹ x	13	11	5
<i>strobus</i> , trees 5 and 6.....	2	x	36	19	7
<i>cembra</i> , tree 2.....	1	x	12	11	8
<i>virginiana</i> , tree 14.....	1	x	21	20	10
<i>divaricata</i> , tree 1.....	1	x	14	13	4
<i>monticola</i> , tree 2.....	1	x	10	6	3
<i>flexilis</i> , tree 2.....	1	x	5	2	1
<i>taeda</i> , tree 9.....	1	8	5	3	1
<i>resinosa</i> , trees 16 and 18.....	2 ?	0	2		
<i>austriaca</i> , tree 2.....	1 ?	0	1		

¹ x=number not counted.

² 27 larvæ killed—rain.

³ 2=trees examined and no eggs found. Therefore, while it is possible that eggs may have been overlooked, it is also possible that a larva dropping or knocked from another tree may have reached these; consequently these species can not be given full credit for choice.

Table 11 gives a summary of experiments in which adults of *Diprion simile* were confined in cages on single young trees of certain species of pine. It shows in some degree the greater success attending the use of *Pinus sylvestris* as a host for the sawfly.

TABLE 11.—Experiments showing the results accompanying the forced use of certain species of pine as hosts by *Diprion simile*.

Species and serial number of host tree.	Parent females.	Eggs.	Larvæ.	Cocoons.	Adults.
<i>Pinus sylvestris</i> :					
Tree 1.....	1				
Tree 2.....	1	9	3	2	
Trees 4 and 5.....	1	¹ N 9	21		
Trees 6 and 3.....	1	79	48	9	
Trees 15 and 16.....	1	53	43		
Trees 17 and 2.....	1	27	18	2	2
Trees 18 and 19.....	1	68	² x	3	1
Tree 20.....	1				
Tree 21.....	1	9	3	2	1
Tree 29.....	1				
Tree 30.....	2	6	5	4	
<i>Pinus virginiana</i> , tree 10.....	3				
<i>Pinus austriaca</i> :					
Tree 1.....	1				
Tree 2.....	1	26	5		
<i>Pinus cembra</i> , tree 1.....	1				
<i>Pinus divaricata</i> , tree 36.....	2				
<i>Pinus flexilis</i> , tree 1.....	1				
<i>Pinus laricio</i> , tree 1.....	3	9			
<i>Pinus strobus</i> :					
Tree 16.....	2	13	2		
Tree 17.....	1	29	6		

¹ N=needles in which eggs have been deposited but the number of eggs not counted.² x=number not counted.

Britton and Zappe,¹⁰ in their paper on this species, give the following list, in which the pines are arranged "according to the preference shown by the sawflies in nature and to the largest number of sawflies reaching maturity" in their host-plant experiments:

<i>Pinus excelsa</i> Wall.	Bhotan pine	} Five-needled pines.
<i>Pinus cembra</i> Linn.	Stone pine	
<i>Pinus flexilis</i> James.	Limber pine	
<i>Pinus strobus</i> Linn.	White pine	
<i>Pinus koraiensis</i> Sieb. & Zucc.	Korean pine	

<i>Pinus montana</i> Du Roi.	Mugho pine	} Two-needled pines.
<i>Pinus densiflora</i> Sieb. & Zucc.	Japanese red pine	
<i>Pinus resinosa</i> Ait.	Red pine	
<i>Pinus sylvestris</i> Linn.	Scotch pine	
<i>Pinus ponderosa</i> Dougl.	Bull pine	
<i>Pinus laricio</i> Poir. var. <i>austriaca</i> Endl.	Austrian pine	

<i>Pinus rigida</i> Mill.	Pitch pine	Three-needled pine.
---------------------------	------------	---------------------

This list shows that *Diprion simile* has a decided preference for the five-needled and the softer two-needled pines. A comparison of this list with the writer's experiments shows an agreement as to the preference for the five-needled pines and the apparent lack of suitability and attraction possessed by *P. laricio*, *P. austriaca*, and *P. ponderosa*, but a disagreement regarding *P. sylvestris*, which was a favored tree and capable of carrying the species from egg to adult in the writer's cages, while, along with *P. ponderosa* and *P. laricio* variety *austriaca*, it was almost immune to infestation in the experiments of Britton and Zappe.

From these data it seems reasonable to suppose that *Diprion simile* is capable of utilizing several of our American pines as hosts and therefore may succeed in establishing itself in the forests of the United

¹⁰ Britton, W. E., and Zappe, M. P. *Op. cit.*, p. 278.

States. Furthermore, the species of pines represented above are from sufficiently different sections of the country to make available a possible host in most of our pine-growing regions and in such a variety of climates that some of them must be favorable for the insect's development.

DISTRIBUTION IN THE UNITED STATES.

FIG. 8.—*Diprion simile*: Distribution in the United States. A round black spot indicates a report from a definite locality; a square spot indicates occurrence in the State, but without definite locality.

Diprion simile is known to occur in nurseries in Connecticut, New York, Pennsylvania, and New Jersey. Specimens have been received from Massachusetts, and the species is recorded from Indiana.¹¹

The history of the sawfly in the United States suggests that it was introduced with imported nursery stock, a view which is supported by the fact that in Connecticut, New Jersey, and Massachusetts it has been intercepted at quarantine inspection ports.

The following list gives the localities from which the species is recorded (see also Fig. 8):

Connecticut: Derby, Greenwich, Hartford, New Canaan, and New Haven.¹¹

Indiana.

Massachusetts.

New Jersey: Elizabeth, Rutherford, and South Orange.¹²

New York: Flushing, L. I.

Pennsylvania: Chestnut Hill¹³ and Penbrook.¹³

ECONOMIC IMPORTANCE.

The imported pine sawfly is to be regarded as an extremely dangerous visitor in the United States, because in Europe, where it is associated with *Diprion pini* L., the two are responsible for an immense amount of damage. *D. pini* has usually been considered the chief cause of this injury, but the two species are so closely allied¹⁴ and so frequently confused that the presence of *D. simile* in this country must be regarded with apprehension. In Europe *D. pini*¹⁵ is accredited with serious damage to pine in southwestern Russia, in 1914 defoliating many acres of pine in the districts of Achtyr and Iziun; in Germany, in Prussia, and particularly in Brandenburg and Silesia, in

¹¹ Britton, W. E., and Zappe, M. P. *Op. cit.*, p. 275.

¹² From a letter from Harry B. Weiss, dated Aug. 12, 1916.

¹³ Localities for specimens determined by S. A. Rohwer.

¹⁴ *Diprion simile* Hartig has been considered a synonym of *D. pini* L.

¹⁵ Britton, W. E., and Zappe, M. P. *Op. cit.*, p. 276.

1914; in France, in 1906; in Sweden; in Norway, in 1914, when the larvæ nearly defoliated 10-year old pines on one plantation; and in England, where they are said to injure Scotch fir as well as pine. In many of these infestations *D. simile* doubtless took a prominent part and in some instances may have been the chief depredator.

Should *D. simile* become permanently established in this country, it will be capable of great destruction. At present it is probably confined to nurseries and estates, where it is doing sufficient damage to attract considerable attention, and nurserymen and gardeners are endeavoring to exterminate it.

CONTROL.

In the United States the seeming preference of *Diprion simile* for young trees makes this sawfly primarily a nursery pest and a nursery problem rather than a forest problem. This confinement to trees in nurseries and on estates is of considerable advantage in an attempt to eradicate or control the species, since in such situations the best opportunities for combating it are found. The infestation is more easily observed early in its course, and methods of control can be employed which, despite their effectiveness, could hardly be recommended to check a forest invasion, owing to the inaccessibility and vastness of the area. Because infestations of the sawfly are at present limited to young trees in nurseries and under similar conditions, it is doubly important to combat it now, first, to check its ravages, and, second, in the hope of exterminating it or at least preventing it from ever becoming established in our forests.

In the effort to control the species its life-history and seasonal-history advantages, already referred to, should be taken into consideration and strict watch maintained against its appearance or reappearance.

Infested trees should be sprayed with some stomach poison, such as a mixture of lead arsenate and water in the proportion of 2 pounds of powdered arsenate of lead to 50 gallons of water, or, in smaller quantities, 6 teaspoonfuls of the powder to 1 gallon of water. A knapsack, barrel, or tank spraying outfit, whichever is available and convenient to use, should give good results.

Hand picking and dropping or shaking the larvæ into pails partially filled with kerosene, crushing them with the hands protected by leather gloves, and knocking or jarring them from the trees and treading on them, are all possible methods of control, especially where labor can be obtained cheaply. Such methods as these have been used successfully in Europe.

ORGANIZATION OF THE UNITED STATES DEPARTMENT OF AGRICULTURE.

November 12, 1923.

<i>Secretary of Agriculture</i>	HENRY C. WALLACE.
<i>Assistant Secretary</i>	HOWARD M. GORE.
<i>Director of Scientific Work</i>	E. D. BALL.
<i>Director of Regulatory Work</i>	WALTER G. CAMPBELL.
<i>Director of Extension Work</i>	C. W. WARBURTON.
<i>Weather Bureau</i>	CHARLES F. MARVIN, <i>Chief</i> .
<i>Bureau of Agricultural Economics</i>	HENRY C. TAYLOR, <i>Chief</i> .
<i>Bureau of Animal Industry</i>	JOHN R. MOHLER, <i>Chief</i> .
<i>Bureau of Plant Industry</i>	WILLIAM A. TAYLOR, <i>Chief</i> .
<i>Forest Service</i>	W. B. GREELEY, <i>Chief</i> .
<i>Bureau of Chemistry</i>	C. A. BROWNE, <i>Chief</i> .
<i>Bureau of Soils</i>	MILTON WHITNEY, <i>Chief</i> .
<i>Bureau of Entomology</i>	L. O. HOWARD, <i>Chief</i> .
<i>Bureau of Biological Survey</i>	E. W. NELSON, <i>Chief</i> .
<i>Bureau of Public Roads</i>	THOMAS H. MACDONALD, <i>Chief</i> .
<i>Bureau of Home Economics</i>	LOUISE STANLEY, <i>Chief</i> .
<i>Fixed Nitrogen Research Laboratory</i>	F. G. COTTRELL, <i>Director</i> .
<i>Division of Accounts and Disbursements</i>	A. ZAPPONE, <i>Chief</i> .
<i>Library</i>	CLARIBEL R. BARNETT, <i>Librarian</i> .
<i>Federal Horticultural Board</i>	C. L. MARLATT, <i>Chairman</i> .
<i>Insecticide and Fungicide Board</i>	J. K. HAYWOOD, <i>Chairman</i> .
<i>Packers and Stockyards Administration</i>	] CHESTER MORRILL, <i>Assistant to the</i> <i>Secretary</i> .
<i>Grain Future Trading Act Administration</i>	
<i>Office of the Solicitor</i>	R. W. WILLIAMS, <i>Solicitor</i> .

This bulletin is a contribution from

<i>Bureau of Entomology</i>	L. O. HOWARD, <i>Chief</i> .
<i>Forest Insect Investigations</i>	F. C. CRAIGHEAD, <i>Entomologist in Charge</i> .

22

ADDITIONAL COPIES
OF THIS PUBLICATION MAY BE PROCURED FROM
THE SUPERINTENDENT OF DOCUMENTS
GOVERNMENT PRINTING OFFICE
WASHINGTON, D. C.
AT
5 CENTS PER COPY

PURCHASER AGREES NOT TO RESELL OR DISTRIBUTE THIS
COPY FOR PROFIT.—PUB. RES. 57, APPROVED MAY 11, 1922

Farmers' Bulletin 1259
United States Department of Agriculture

H. C. Clepper

A SAWFLY INJURIOUS *to* YOUNG PINES

YOUNG PINES, both nursery stock and natural reproduction, are often defoliated by larvæ of Leconte's sawfly. Defoliation of young pines is usually severe in its effects, killing, misshaping, or weakening the trees to such an extent that attacks of secondary enemies will kill them.

This insect is common on the scrub pines throughout the eastern United States and is a constant menace to the better species in nurseries or reforesting areas. It can be controlled in nurseries and parks, when the infestation is heavy, by spraying with lead arsenate at the rate of 2 pounds of the powder to 50 gallons of water; or, if scatteringly present, by hand methods such as knocking the larvæ from the trees and crushing them.

In larger and less accessible areas, where it would be impractical to attempt control by spraying, rangers and lumbermen should make a practice of destroying the colonies of these larvæ whenever they are found.

Contribution from the Bureau of Entomology

L. O. HOWARD, Chief

Washington, D. C.

January, 1922

A SAWFLY INJURIOUS TO YOUNG PINES.

WILLIAM MIDDLETON,

Scientific Assistant, Forest Insect Investigations.

The larva, or false-caterpillar, of an insect known as Leconte's sawfly¹ is a serious enemy of pine in nurseries, parks, and reforesting areas in the eastern United States. This insect shows a decided

FIG. 1.—Young silver pine in nursery, completely defoliated by larvæ of Leconte's sawfly.

preference for young trees and has demonstrated its ability to leave its local host and attack other species of pines. The effects of defoliation on young pines (figs. 1, 2), especially before late summer,

¹ *Neodiprion lecontei* Fitch; order Hymenoptera, suborder Chalastogastra, family Tenthredinidae, subfamily Diprioninae.

are usually severe, the part denuded often being killed. Trees not killed are frequently infested by secondary enemies which com-

FIG. 2.—Young scrub pine on edge of woods, completely defoliated by larvæ of Leconte's sawfly.

plete the work; and, if by chance the trees recover, they are stunted or misshapen and of little or no commercial or ornamental value.

APPEARANCE, LIFE, AND HABITS OF THE INSECT.

The needles are eaten by small larvæ one-eighth of an inch long when young to three-fourths or seven-eighths of an inch in length when full grown, which feed in colonies or groups. The young larvæ (fig. 3) are pale, whitish to leaden white, and unspotted, with the head brownish; but they change in appearance through a series of molts (five for the male and six for the female), becoming yellowish-white with a number of rows of black spots on the body (fig. 4) and with the head dark brown to orange. These larvæ have three pairs of legs on the thorax and eight pairs of smaller legs on the abdomen, and the head has only a single pair of eyes, situated one at each side in a rather large, circular, blackish spot.

FIG. 3.—Young larvæ of Leconte's sawfly, second and third stages. Four times natural size.

FIG. 4.—Larvæ of Leconte's sawfly, fourth and fifth stages. Four times natural size.

The larvæ are found clustered in colonies feeding on the pine needles and occasionally on the tender bark (fig. 5) of the young twigs.

When full grown the larvæ spin cocoons from which they emerge later as adults. Adult emergence is divided into two periods, a first issuance, brood A, and a second issuance, brood B. When the eggs are laid and hatch in the late spring or early summer, adults of brood A issue from the cocoons in the late summer and early fall

of the same year; but adults of brood B from this batch of eggs do not issue from their cocoons until the late summer and early fall of

the following year, a complete colony developmental period of 14 months. If the eggs are laid and hatch in the late summer, adults of brood A of these eggs issue from their cocoons in the spring and early

FIG. 5.—Terminal of young scrub pine showing defoliation and feeding work of larvæ of Leconte's sawfly on bark. Three-fourths natural size.

summer of the following year, and adults of brood B of the same batch of eggs emerge from their cocoons in the late summer and early fall of the same year as brood A, making a complete colony developmental period of 12 months. (See diagram, fig. 6.)

The cocoons (fig. 7) have been found several inches underground and only in this location; but it is quite possible that some larvæ, es-

FIG. 6.—Chart showing life and seasonal history of Leconte's sawfly through the active period of three years (November to March omitted, the insect being in the cocoon during this period).

pecially those the adults of which will issue before winter, may spin their cocoons in more exposed places, even on the pine twigs. The

FIG. 7.—Cocoons of Leconte's sawfly: At left, unopened ones containing the insects; at right, those from which adults have issued (emergence holes shown). The females issue from the larger cocoons and the males from the smaller ones. Approximately natural size.

cocoons are tough, papery, red-brown, capsule-shaped cases, five-sixteenths to seven-sixteenths of an inch long.

The adults (fig. 8) are rather robust, four-winged insects. The male is black with reddish yellow legs and has beautiful feathery feelers, or antennæ. The female has the head and first two thoracic segments reddish brown and the third thoracic segment and abdomen black. Her feelers are rather slender and not feathery.

A *B*

FIG. 8.—Leconte's sawfly : *A*, Adult female ; *B*, adult male. About eight times natural size.

The sawflies get their name from the complicated egg-laying organ of the female. It is well named "the saw," having blades and teeth and being used exactly as a saw when tearing into the leaves in cutting a pocket for the eggs.

The eggs are laid in shoe-shaped pockets, or slits, in the needles. Usually, attacked needles occur in a cluster, each needle bearing a

FIG. 9.—Needles of scrub pine showing eggs of Leconte's sawfly embedded in them. Much enlarged.

number of eggs. The egg scars, or pockets, are rather easily seen, being yellowish on the dark green needles. (Fig. 9.)

DISTRIBUTION.

Leconte's sawfly occurs throughout the eastern United States, the accompanying map (fig. 10) indicating localities from which the species has been recorded or specimens received.

TREES ATTACKED.

This species seems to have a preference for jack pine,² red pine,³ and scrub pine,⁴ each of which furnishes a local host in some part of the eastern United States throughout which the insect occurs. Its attack is not limited to these species, however, for it is also found on or recorded from white pine,⁵ scotch pine,⁶ loblolly pine,⁷ shore pine,⁸ silver pine,⁹ mugho pine,¹⁰ western yellow pine,¹¹ longleaf pine,¹² Austrian pine,¹³ and American larch.¹⁴

FIG. 10.—Map showing localities where Leconte's sawfly is known to occur.

PERIODICAL OUTBREAKS.

Leconte's pine sawfly, like most insect enemies of forest trees, appears and disappears periodically. For several years this species will be very abundant, then for a few years it will be rare. The cause for this periodic disappearance has not been determined, but the low records of parasitism indicate that some factor other than parasitism plays an important rôle.

NATURAL ENEMIES.

Eight species of insect parasites and an infectious disease have been found killing this sawfly. None of these enemies, however, has been found widely enough distributed or abundant enough to account for the periodic disappearance of the species, and it is certain that neither any nor all of these natural checks are sufficiently numerous or effective to justify disregard of the artificial control measures suggested.

² *Pinus banksiana*.

³ *P. resinosa*.

⁴ *P. virginiana*.

⁵ *P. strobus*.

⁶ *P. sylvestris*.

⁷ *P. taeda*.

⁸ *P. contorta*.

⁹ *P. monticola*.

¹⁰ *P. mughus*.

¹¹ *P. ponderosa*.

¹² *P. palustris*.

¹³ *P. austriaca*.

¹⁴ *Larix americana*.

CONTROL.

The control of Leconte's sawfly depends largely upon the extent and location of the infestation. In nurseries and parks, when the infestation is heavy, a thorough spraying with lead arsenate, 2 pounds of powdered lead arsenate to 50 gallons of water (or 6 teaspoonfuls to the gallon), will give good results. Spraying should begin when the larvæ are first discovered. In a scattered infestation, hand picking or knocking the larvæ from the trees and crushing them will be found more economical and at least as effective.

In large areas of either natural or artificial reproduction control can not be generally practiced because of its expense; but rangers and lumbermen should make it a habit to destroy the colonies of these larvæ whenever found.

Whenever these insects are observed in any locality and control measures are practiced against them, it is important that the territory be carefully surveyed for the following 14 months, since it is possible that some larvæ may have escaped the treatment and have spun cocoons. This possibility makes watchfulness necessary over the entire colony period of the species in order that an emergence of adults from these cocoons may not reestablish the infestation.

U. S. DEPARTMENT OF
AGRICULTURE

FARMERS' BULLETIN No. 1623

THE
GIPSY MOTH
AND THE
BROWN-TAIL MOTH

THE GIPSY MOTH and the brown-tail moth are two very destructive enemies of tree foliage which have gained entry into this country from Europe. The gipsy moth has spread over a large part of New England, and isolated colonies have been found in New York State east of the Hudson River, on Long Island, and in New Jersey. The brown-tail moth has become established in many localities, embracing about one-third of the area of the New England States.

Substantial progress has been made in exterminating the gipsy moth in New Jersey, and many isolated infestations have been cleaned up in other States.

During the last 10 years the brown-tail moth has been far less abundant than was previously the case, except in certain sections in the eastern part of the infested territory. The decrease has been due to such factors as heavy mortality during the winter, increase in the abundance of imported parasites, a fungous disease that attacks the larvae, and field control work including spraying operations.

The States and the Federal Government (the latter through research conducted by the Bureau of Entomology and quarantine and control work conducted by the Plant Quarantine and Control Administration) are actively engaged in destroying these insects by the use of appropriate measures. The Federal Government is chiefly concerned in preventing their spread, while the States aim to reduce the infestations and prevent the defoliation of valuable trees in woodlands, orchards, and parks.

The methods which have been found effective in destroying the insects, together with a brief statement of the progress that has been made, are set forth in this bulletin.

This bulletin is a revision of and supersedes Farmers' Bulletin 1335, Controlling the Gipsy Moth and the Brown-Tail Moth.

THE GIPSY MOTH AND THE BROWN-TAIL MOTH

By A. F. BURGESS, *Principal Entomologist in Charge of Moth Work, Plant Quarantine and Control Administration*

CONTENTS

	Page		Page
The gipsy moth.....	1	The brown-tail moth—Continued.....	
Introduction and spread.....	1	Native enemies of the brown-tail moth.....	27
Record of clean-up of outlying gipsy-moth colonies.....	7	Introduced parasites and enemies.....	28
Gipsy-moth extermination project in New Jersey.....	8	Methods of controlling the brown-tail moth.....	28
Life history of the gipsy moth.....	10	Organization and status of work against the gipsy moth and the brown-tail moth.....	29
Means by which the gipsy moth is spread.....	12	Maine.....	29
Food plants of the gipsy moth.....	13	New Hampshire.....	29
Injury caused by the gipsy moth.....	13	Vermont.....	30
Effect of climatic conditions on the gipsy moth.....	15	Massachusetts.....	30
Native enemies of the gipsy moth.....	15	Rhode Island.....	30
Introduced parasites and enemies.....	15	Connecticut.....	30
Methods of controlling the gipsy moth.....	16	New York.....	30
The brown-tail moth.....	24	New Jersey.....	30
Introduction and spread.....	24	Pennsylvania and Ohio.....	31
Life history of the brown-tail moth.....	24	Work of the Dominion of Canada.....	31
Food plants of the brown-tail moth.....	26	Work of the United States Department of Agriculture.....	31
Injury caused by the brown-tail moth.....	26	Cooperative work.....	32

THE GIPSY MOTH

INTRODUCTION AND SPREAD

IN 1869 a number of egg clusters of the gipsy moth (*Porthetria dispar* L.) were brought from France to Medford, Mass., by a French mathematician and astronomer, who seemed to have had a fanciful idea that he could cross this insect with silkworm moths and thus develop a hardy race of silk-producing insects. In the course of his rearing experiments some of the eggs were accidentally lost or some of the caterpillars escaped, and he made at that time public acknowledgment of this fact, evidently somewhat appreciating the danger.

It is clear now that the insect increased slowly and after 10 years seemed to have been noticed by local residents but was believed to be some native caterpillar. It was not until some 20 years after the introduction, namely, in the summer of 1889, that this insect became so abundant and destructive as to bring it into general public notice. At that time fruit and shade trees in the immediate region were completely defoliated and the caterpillars swarming into the houses became a very grave nuisance. This situation was so serious that the State of Massachusetts appropriated funds and delegated to the State department of agriculture the task of exterminating this pest.

The area that was found to be infested covered about 359 square miles and the trees in many towns surrounding Boston were com-

pletely defoliated each season for a number of years. As the work continued, more effective methods of treatment were adopted and better results were obtained; consequently, during the summers of 1898 and 1899 little defoliation could be found in the entire area and few specimens of the moth were located throughout the entire residential sections of the infested district. Careful examination indicated that the insect had been exterminated in some of the towns around the outside border of the originally infested district. In February, 1900, the work was discontinued by the general court because of the popular belief that the danger had passed, in spite of the advice of experts that the insect had not been stamped out.

Within the next five years the insect increased enormously. Many of the towns and cities in the old infested district were overrun with caterpillars which completely defoliated the trees in many of the residential sections and thousands of acres of woodland were stripped of leaves during the summer. The situation became so serious and intolerable that in 1905 control work was resumed by the State of Massachusetts. In the meantime the insect had spread far beyond the original limits of infestation, over 2,224 square miles being involved in Massachusetts as well as a considerable number of isolated areas in Maine, New Hampshire, and Rhode Island.

In 1906 an appropriation was made by Congress and the Secretary of Agriculture was authorized to take all possible measures, in cooperation with the States concerned, to prevent the spread of this pest. The insect had increased to such enormous numbers and had spread so rapidly that the utmost efforts of the Federal and State forces were only able to apply relief measures in the badly infested residential sections and partially retard the continued spread of the pest. Efforts were made to prevent the shipment of the insect to uninfested localities by inspecting products that were likely to carry the insect. This phase of the work was greatly strengthened as a result of the enactment of the plant quarantine law by Congress on August 20, 1912.¹ Since October of that year shipments from the infested district have been regulated by Federal quarantine.

On account of the serious situation that was resulting from the continued spread and increasing damage caused by this insect, efforts were constantly being made to develop improved means of control. The process of manufacturing lead arsenate, which was first made and used in the gipsy-moth work in 1893, was perfected so that the cost of production was reduced and the product could be more effectively applied. Spraying machinery and equipment were developed to a high point of efficiency. Advantage was taken of the results of experimental work in carrying on the field operations, and the details of field management were constantly improved in order that the greatest possible volume of effective work could be done with the funds available.

This pest is distributed by natural means, principally by the newly hatched caterpillars being blown by strong winds when accompanied

¹ This act created the Federal Horticultural Board and gave authority for the establishment of Federal quarantines to prevent the spread of dangerous introduced insect pests and plant diseases. The act making appropriations for the Department of Agriculture for the fiscal year ended June 30, 1929, transferred the functions of this board, together with insect-control activities, to the Plant Quarantine and Control Administration.

by high temperatures in the spring. Under favorable conditions such caterpillars may drift a distance of 20 miles.

In spite of the efforts that were made by the Federal Government and all the States concerned, the insect continued to spread. By 1914 it had covered the southern half of New Hampshire and extended as far east as Bangor, Me. On the west it had crossed the Connecticut River in Vermont and Massachusetts. Rhode Island and many towns in eastern Connecticut were found to be infested.

During the war period conditions were most unfavorable for preventing spread. The loss of efficient personnel and constant turnover of men, together with extraordinary increase in costs, made progress difficult. During this period weather conditions were especially favorable for spread toward the west. By the fall of 1922, colonies were found in many towns in Vermont and west of the Connecticut River in Massachusetts and Connecticut, and an infestation was located in New York State adjoining the Massachusetts State line. Every indication pointed to the continued spread of the insect unless more intensive work could be done.

As a result of this serious situation a conference was held in the office of the commissioner of farms and markets in Albany, December 26, 1922, which was attended by representatives from all the infested States, the Dominion of Canada, and the United States Department of Agriculture.² The conference considered the entire subject of the prevention of spread of the gipsy moth and its control, and adopted a resolution urging that sufficient funds be obtained by the States interested and the Federal Government, for the purpose of continuing and strengthening present control methods in the infested area, to do necessary scouting for the discovery and destruction of border infestations, to determine the location of the most practical place for a control zone, to take necessary steps to make control therein effective, and for the destruction of all infestations in and west of said zone.

In order to carry out this project legislation was passed by the State of New York in April, 1923, carrying an appropriation of \$150,000 to be administered by the department of conservation of that State. Federal funds were also provided for the fiscal year beginning July 1, 1923, in order to bring about effective cooperation.

The plan finally adopted, in cooperation with the New York Department of Conservation, was to locate a zone east of the Hudson River where clean-up operations would be centered in order to prevent westward spread of this pest. (Fig. 1.) The territory east of this zone was to be treated by the States concerned as far as their funds would permit, and their work was to be supplemented by liberation of imported parasites and other natural enemies of the insect by the Bureau of Entomology. Work in the New York portion of the area was to be financed by the State with such assistance as could be given by the bureau after covering the eastern part of the zone. This zone embraced an area of over 8,000 square miles extending from Long Island Sound east of the Hudson River (excluding

² THE GIPSY MOTH, AN IMMINENT MENACE TO THE FOREST AND SHADE TREES OF THE STATE OF NEW YORK. New York State Dept. of Farms and Markets, Agr. Bul. 148, 58 p., illus. 1922.

Westchester County, N. Y.) to the Canadian border, a distance of over 250 miles, and ranging in width from 25 to 30 miles.

While much of the extremely rugged country in the Adirondack and Green Mountains was avoided in northern New York and in Vermont, as well as the Catskills and some of the rougher country west of the Connecticut River in Connecticut, there are areas, embracing the Berkshire Hills in western Massachusetts and some of the

FIGURE 1.—Barrier zone in New England and New York

territory directly south and southwest of them in Connecticut and New York, where the terrain is extremely difficult.

In 1924, owing to the number of infestations found in Massachusetts and Vermont, and a vigorous colony that was located at Henrysburg, Quebec, by the inspection force of the Dominion entomologists of Canada, the quarantine line was moved westward to embrace the whole of the State of Vermont and additional towns in northwestern Connecticut. A foreign quarantine regulating the movement of Christmas trees was made effective covering the southern tier of towns in the Province of Quebec. (Fig. 2.) Since that time, owing to the effectiveness of the clean-up work in the barrier zone, sup-

plemented by a limited amount of scouting work carried on west of the zone as a precautionary measure, it has been possible to revise the area under quarantine and eliminate the area included in the zone. The Canadian quarantine was withdrawn July 1, 1928. (Fig. 3.)

The danger of reinfestation of the barrier zone depends to a very large extent on the density of infestation east of the zone. The

FIGURE 2.—Quarantined areas, 1924

following tabulation indicates the acreage defoliated by the gipsy moth east of the zone from 1924 to 1929, inclusive:

Acreage defoliated by the gipsy moth, 1924 to 1929, inclusive

Year	Acre	Year	Acre
1924	825	1927	140,920
1925	48,560	1928	262,514
1926	80,822	1929	551,133

It will be noted that defoliation has been increasing rapidly in the area east of the barrier zone, and during 1928 and 1929 the insect

was sufficiently abundant to cause some defoliation between the Connecticut River and the zone area. During the period covered by the tabulation, parasites and natural enemies of the gipsy moth have been scarce in most of the infested area, but in the summer of 1929 an increase was noted. On account of these conditions the control work done by the States and the cities and towns has been confined

FIGURE 3.—Quarantined areas, 1928

chiefly to residential sections and to parks and street trees. Very little work has been done in woodland areas. More infestations were found in the barrier zone in 1928 after July 1 than during the entire previous fiscal year, which made the problem of preventing the increase of the insect far more difficult and emphasized the importance of carrying on work in the territory east of the zone in order to make it possible to keep the zone free from infestations. As a result of the increase in infestation directly east of the zone, 7 towns in Vermont, 12 in Massachusetts, and 31 in Connecticut, embracing an area of 1,581 square miles, were added to the generally infested area under quarantine.

During the latter half of 1929 and the first half of 1930 the number of infested locations continued to increase in the barrier zone and additional funds were made available by Congress to carry on clean-up work. No action was taken, however, to provide for the examination of territory immediately east of the zone from which most of these infestations originated and which furnish a source of constant supply of the insect so long as they remain untreated. The scouting work during the fall of 1929 and spring of 1930 resulted in the discovery of an increasing number of infested locations, although practically all of those found in previous years had been exterminated. Serious infestations were found in Branford, Middlefield, and Meriden, Conn., and Chester, Mass., towns immediately adjoining or close to the barrier zone line to the east. These colonies will be treated by the States concerned. Three of these colonies are in heavy woodland, and as no scouting work is done in woodland in most of the towns which border the barrier zone on the east, there is every probability that more infestations of this nature are present which, if allowed to remain untreated, will cause constant reinfestation of the areas that are being cleaned up.

Since the barrier zone was established, large areas immediately adjoining the zone on the west have been examined and two small infestations found and eradicated. Thus far the work in the barrier zone has accomplished its purpose, namely, the prevention of spread of the insect to the west and the extermination of the colonies as rapidly as they are located within the zone area.

RECORD OF CLEAN-UP OF OUTLYING GIPSY-MOTH COLONIES

In 1912 a small area was found heavily infested with the gipsy moth in Geneva, N. Y. Prompt action was taken by the State department of agriculture, and the Federal Bureau of Entomology cooperated. Clean-up measures were applied. Several very thorough annual examinations have been made but no trace of the insect has been discovered. In February, 1914, a colony was discovered at Bratenahl, a suburb of Cleveland, Ohio. Immediate attention was given this colony by the Bureau of Entomology in cooperation with the Ohio Department of Agriculture. As a result of thorough inspection and treatment work for three seasons, the colony was exterminated. In May, 1914, a colony was found by State inspectors in North Castle, Westchester County, N. Y. The infestation was heavy at the center and scouting determined that it extended over about 1 square mile. Intensive work was done throughout this area and the surroundings by the State and Federal forces, and although the locality was a very difficult one to treat thoroughly, the moth was exterminated. In the same year a small colony was exterminated at Rutherford N. J., by the Bureau of Entomology. In June, 1921, a small infestation was found at Greenport, near the east end of Long Island, N. Y., and during July another was reported at Patchogue on the south side of the island. The Bureau of Entomology, in cooperation with the State department of farms and markets, exterminated the insect in these colonies. Since that time several small infestations have been found on Long Island and have been exterminated by the New York State Department of Conservation. (Fig. 4.)

GIPSY-MOTH EXTERMINATION PROJECT IN NEW JERSEY

Early in July, 1920, the gipsy moth was found on a large estate near Somerville, N. J., by one of the State inspectors. The infestation centered in a large plantation of blue spruce trees, several acres of which were defoliated. Dead trees in the worst infested portion indicated that they had been killed as a result of defoliation and that the infestation was one of long standing. (Fig. 5.)

These trees had been imported from Holland about 10 years before, and there is no doubt that the infestation came with this shipment, which was received prior to the enactment of the plant quarantine act. This shows clearly the pressing need for precautionary measures to keep out dangerous pests.

Scouting work carried on during the fall of 1920 and spring of 1921, financed by the State of New Jersey and the Federal Government, revealed infestations of this insect in scattered localities in an area of over 400 square miles surrounding Somerville.

Previous successes in cleaning up local infestations led to the adoption of a cooperative plan to clean up this large, newly discovered infestation. State and Federal funds were appropriated, and the work was organized under the direction of the field office for gipsy moth control of the Bureau of Entomology. When the limits of infestation had been determined, the area was placed under State quarantine, which required, as a condition of movement from the infested area, certification as to freedom from infestation of all products likely to carry any stage of the gipsy moth. Thorough inspection which permitted certification was done in cooperation with the Federal Government and by Federal inspectors. The area

FIGURE 4.—Map showing locations, outside of the large New Jersey infestation and the barrier zone, where the gipsy moth has been exterminated: 1, Cleveland (Bratenahl), Ohio; 2, Loretto, Pa.; 3, Geneva, N. Y.; 4, Schenectady, N. Y.; 5, North Castle and Garrison, N. Y.; 6, Brooklyn, Roslyn, Kew Gardens, and Greenport, on Long Island, N. Y.; 7, Deal Beach, Wyckoff, South Orange, Scotch Plains, Paterson, Madison, Glen Rock, Elizabeth, and Rutherford, N. J.; 8, Henrysburg, Quebec, Canada.

under regulation has been gradually reduced as the results of the clean-up work have made this possible.

As soon as active work was under way in New Jersey it was discovered that trees had been shipped during the preceding years from the estate where the insect was first located and that the danger of the establishment of the insect in many localities was very great. Fortunately a record of all these shipments was available. It was found that 261 shipments had been sent to the District of Columbia and to 15 States outside of New Jersey as follows: Connecticut, Delaware, Florida, Illinois, Indiana, Kentucky, Maryland, Minnesota, Missouri, New York, North Carolina, Ohio, Pennsylvania, Virginia, and Wisconsin. These shipments were traced and inspections made of the planted stock by State or Federal officials. In

addition to this, 318 shipments had been distributed in New Jersey, and these were followed up in the same way.

As a result of this work small infestations were found at Loretto, Pa., Garrison, Roslyn, and Kew Gardens, N. Y., and Deal Beach, Wyckoff, South Orange, Scotch Plains, Paterson, Madison, Glen Rock, and Elizabeth, N. J. Thorough work cleaned up these small infestations during the following year, and repeated scouting has failed to indicate the presence of the pest.

In the meantime intensive work was carried on in the central part of the infested area, and more than 3,000,000 egg clusters were treated the first season. The trees on several acres were not only completely defoliated at the time the colony was discovered, but many were dead and others were in a dying condition. This intensive work pre-

FIGURE 5.—Blue spruce defoliated and killed by the gipsy moth

vented defoliation the following season; in fact, no trees have suffered from gipsy-moth feeding in New Jersey since that time. During the next four years the scouting and clean-up operations were continued throughout the entire area known to be infested, particular attention being paid to a series of extensive woodland areas north of Somerville, known as the Watchung Ridges. These areas were by far the most difficult and expensive part of the territory to handle properly, as many square miles of woodland were involved. Substantial progress was made as the work proceeded. The density of the infestation in 1921 is indicated on the map in Figure 6.

Since 1925 continuous progress has been made in cleaning up and exterminating the gipsy moth in New Jersey. It has been necessary to operate in an area embracing 2,369 square miles, and the towns in the area where colonies have been found cover 924 square miles.

The area in which intensive scouting is necessary has been reduced to 197 square miles. (Fig. 7.)

LIFE HISTORY OF THE GIPSY MOTH

In the course of its life the gipsy moth passes through four stages—the egg, the larva or caterpillar, the pupa, and the adult or moth. (Fig. 8.)

FIGURE 6.—Map showing area in New Jersey infested by the gipsy moth in 1921, indicating the density of infestation and location of isolated colonies

THE EGGS

The female gipsy moth deposits a cluster containing 400 eggs or more, which she covers with buff-colored hairs. Most of the egg clusters are laid during July and hatch about the time the leaves begin to appear the following spring. They are deposited on the under side of branches of trees, on tree trunks, under loose bark, or in cavities in the trunks or branches, and are sometimes placed on stones or rubbish and in a variety of situations where they are concealed from view. As the female moths do not fly, egg clusters are

seldom found far from the food plant upon which the caterpillars developed.

THE LARVAE

The newly hatched larvae feed on the opening leaves, making small perforations. They grow rapidly and become full fed early in July. During this period they molt five or six times, and as they increase in size they eat a larger proportion of the foliage, so that, if the infesta-

FIGURE 7.—Map showing conditions of infested area in New Jersey in 1928, indicating colonies and reduced area requiring intensive work

tion is severe, trees may be stripped completely of foliage before the end of June.

THE PUPAE

When full grown, the caterpillars shed their skins and transform to pupae, which are chestnut brown and provided with tufts of yellow hairs. They remain in this stage about 10 days, after which the adult insects emerge.

FIGURE 8.—Life stages of the gipsy moth: A, Female moth; B and F, pupae; C, larvae or caterpillars; D, male moth; E, egg mass. All about three-fourths natural size

THE ADULTS

The male moth is dark brown, with black wing markings, and flies well. The female is nearly white, with black markings on the wings, and because of the weight of the abdomen she does not fly. After mating, the females begin depositing eggs.

The time of year when the different forms of the insect may be found in the field is shown in Figure 9.

FIGURE 9.—Time of year when the different stages of the gipsy moth are present in the field

MEANS BY WHICH THE GIPSY MOTH IS SPREAD

Egg clusters of the gipsy moth deposited on trees, lumber, stone, or other products that are likely to be shipped, may be carried long distances and cause the establishment of new colonies of the insect. Spread in this

way is prevented by inspection of such products before they are shipped. Caterpillars of this insect may be spread for limited dis-

tances by being carried on moving objects, such as trains, horse-drawn vehicles, or automobiles. The danger of such spread is very slight if the roadways are kept free from severe infestation. New colonies are started principally by the spread of newly hatched caterpillars. Experiments have shown that these caterpillars may be blown by the wind, if the temperature is high enough for the caterpillars to be active, and the stronger the wind the greater the probability of their being carried long distances. Caterpillars have been carried more than 20 miles in this way, and specimens have been caught in the air 50 feet above the ground, although probably they are carried much higher in the air. The temperature must range above 65° F. and the wind velocity must be 8 miles or more per hour, in order that spread of small caterpillars by wind may result.

In recent years a new avenue for spread of the gipsy moth has developed from the extensive use of the motor truck for long-distance hauling. Fortunately the material hauled is only occasionally of a dangerous nature, but chance of spread in this way is more or less constant.

The popularity of automobile camping parties presents another source of danger, particularly during the vacation season. There is a growing tendency, however, to restrict such camps to municipal or semipublic grounds. Under these conditions the danger can be minimized by proper inspection. Warning posters are being used in sections where parties are likely to camp, instructing them to examine their effects carefully so that no caterpillars or egg clusters of the insect will be carried away. Many camps in the heavily infested area are inspected annually.

FOOD PLANTS OF THE GIPSY MOTH

The food plants most favored by the gipsy moth are apple, the different species of oak, gray birch, alder, and willow. In cases of bad infestation nearly all deciduous trees are injured to a greater or less extent, with the exception of ash. Hickory is not a favored food plant, although the foliage occasionally shows severe feeding. Chestnut will not support the gipsy-moth caterpillars when in the first stage, and pine will not support them in the first two stages; but if other food plants are present severe injury may result from feeding by the larger caterpillars. Beech and poplar are sometimes fed upon freely, and occasionally the trees are defoliated. The different species of spruce are more susceptible to attack than the pines.

INJURY CAUSED BY THE GIPSY MOTH

Unless reduced in numbers by natural enemies, unfavorable climatic conditions, or the application of control measures, the gipsy moth is capable of causing enormous injury to tree growth. In the area of New England which has suffered most from this insect thousands of trees have been killed as a result of defoliation. (Fig. 10.) On many areas the trees were cut before they were mature and the wood sold at a loss on account of damage caused by the insect. Apple and oak have been injured most, but pine and other coniferous trees mixed with deciduous growth have suffered severely. (Fig. 11.)

FIGURE 10.—Deciduous trees defoliated by the gipsy moth

FIGURE 11.—White pine, in a mixed growth, defoliated by the gipsy moth

Many oak trees which have been weakened severely as a result of defoliation by the gipsy moth and the brown-tail moth have failed to recover because of the attacks of certain wood-boring insects. The species which has caused most damage in this way is a beetle (*Agrilus bilineatus* Weber) known as the two-lined chestnut borer, the larva of which feeds beneath the bark of injured trees.

EFFECT OF CLIMATIC CONDITIONS ON THE GIPSY MOTH

Investigations have shown that extremely low temperature during the winter is fatal to eggs of this insect that have been deposited in exposed situations. From -20° to -25° F. is usually required to injure them sufficiently to prevent hatching, and if the clusters are protected by snow, ice, or other material, the eggs are not killed even at this temperature. After extremely cold winters many eggs fail to hatch. Evidence also indicates that abnormally early frosts in the fall injure freshly deposited egg clusters.

During May, when the caterpillars are hatching, there are sometimes long periods of cold weather accompanied by heavy rains. This condition, or the occurrence of severe frosts in June, occasionally causes the death of the least hardy caterpillars.

NATIVE ENEMIES OF THE GIPSY MOTH

Few insect enemies of the gipsy moth native to New England cause any noticeable reduction in its numbers. This is shown by the fact that from 1900 to 1905, when no systematic effort was made to suppress the insect, alarming injury resulted, and native insect enemies did not increase to any marked degree. The same is true of the work of native insect-eating birds. While they undoubtedly feed to some extent on gipsy-moth caterpillars, there is no case on record where they have been able to control the insect. The "wilt," a disease which attacks and kills the caterpillars and pupae, has probably occurred in this country for many years. During some seasons, particularly in badly infested areas, it causes the death of an enormous number of the caterpillars and is often an important factor in reducing the infestation locally. As a rule, this disease is more common in heavy infestations, where overpopulation of caterpillars causes a scarcity of food.

INTRODUCED PARASITES AND ENEMIES

In 1905 an effort was made by the State of Massachusetts, in cooperation with the Bureau of Entomology, United States Department of Agriculture, to introduce the parasites and natural enemies of the gipsy moth from its native home in Europe and Japan. Since that time a large amount of parasitized material has been received nearly every year, and as a result some promising natural enemies have become established in this country and are assisting in bringing about the control of the species. The enemies which have become established and are at present destroying the largest number of gipsy-moth caterpillars and pupae are the *Calosoma* beetle (*Calosoma sycophanta* L.), two species of parasitic flies (*Sturmia scutellata* Desv. and *Compsilura concinnata* Meig.), one

of which attacks the brown-tail moth as well as caterpillars of many native insects, and two species of small wasplike flies (*Apanteles melanoscelus* Ratz. and *A. lacteicolor* Vier.), one of which likewise attacks the brown-tail moth. Two tiny parasites of the gipsy-moth eggs have also been introduced and colonized, one (*Oencyrtus kuvanae* How.) from Japan and the other (*Anastatus disparis* Ruschka) from Europe.

Several other parasites have been introduced and are established in this country; however, they have not increased sufficiently to be considered important factors in gipsy-moth control. Some additional parasites have been imported during the last three years, but their value has not yet been fully demonstrated.

The work of the natural enemies of the gipsy moth, including the imported parasites, the Calosoma beetle, and the wilt disease, has greatly reduced the numbers of the insect in many localities that are badly infested.

It is hoped that when the enemies of the moth are present in larger numbers over the entire infested territory the insect will become much less destructive than at present.

The records at the gipsy-moth laboratory at Melrose Highlands, Mass., indicate that the species of parasites introduced prior to 1923 increased gradually in the field until 1923, when they reached their greatest abundance, and the results of their work were very noticeable over much of the infested area.

In 1924 the gipsy moth was rather scarce, and parasitism was low; the following year it was still low and did not increase much in 1926. In 1925 the gipsy moth began to increase rapidly in the older infested area, and although some of the parasites gained slightly in 1927 and 1928 they did not prevent the rapid increase of the moth.

Until the natural enemies demonstrate greater effectiveness throughout the entire infested area it will be necessary to employ the best mechanical methods for restricting the spread and curtailing the increase of the gipsy moth. In the meantime further work is being done in foreign lands, particularly in central Europe and the Mediterranean countries, to determine the principal factors that are responsible for the periodical scarcity of the insect there and to take advantage of this knowledge for the purpose of developing more effective natural control in this country.

METHODS OF CONTROLLING THE GIPSY MOTH

GENERAL METHODS

Creosote.—One of the best methods of controlling the gipsy moth is to treat the egg clusters of the insect between August 1 and April 1 with creosote to which a small quantity of lampblack has been added. The material is sold by dealers in the infested region under the name of gipsy-moth creosote. It is applied with a brush and leaves a black residue on the clusters treated. Creosote may be obtained from nearly all the large hardware or seed stores in the infested district.

Burlap bands.—Gipsy-moth caterpillars usually seek shelter during hot, sunny days, and if a band of burlap is attached to a tree (fig. 12), large numbers of them will crawl beneath it, where they may

be crushed each day. Ordinarily a strip of burlap about 8 inches wide is placed loosely around a tree trunk and a piece of twine is passed around the center and tied to hold it in place. After this is done the top part of the burlap is folded down so that a double shelter is made beneath it. The extensive use of these bands has been discontinued during the last few years, owing to the expense

FIGURE 12.—Burlap band on tree with caterpillars beneath it

involved and because of the fact that if the burlap bands are applied early in the season, before the brown-tail moth caterpillars have pupated, an excellent place is furnished for these poisonous caterpillars to make their cocoons, and severe poisoning results to the workmen. If this method is to be used in areas where the brown-tail moth is abundant, the burlap should not be attached to the trees until after June 15, when most of the brown-tail moth caterpillars will have pupated.

FIGURE 13.—Caterpillars on tree trunk below sticky band

Sticky bands.—Bands of a widely known sticky substance for tree banding, which may be obtained on the market, may be used on tree trunks after the bark has been scraped so that the material can be applied evenly in a thin layer with a paddle. The purpose of this band is to prevent caterpillars from ascending the trees, and if the egg clusters have previously been treated, this is a very effective measure. Every week or 10 days during the caterpillar season a comb or other similar implement should be run over the band in order to prevent hardening of the surface and to bring up fresh, sticky material from the part of the band near the bark. (Fig. 13.) Placing these bands on the trees prevents the caterpillars from reaching the foliage; and as they usually mass in large numbers beneath the bands, conditions are favorable for wilt disease to develop, and the caterpillars often die in large numbers from this disease and from starvation.

Gipsy moth tree-banding material.³—This is a black, greasy substance which was prepared and tested by the Bureau of Entomology in cooperation with the Bureau of Chemistry. It is similar to the product known as “Raupeleim” that has been used in the German forests for many years to prevent injury to the trees by caterpillars. It is applied with a special “gun” designed for the purpose. (Fig. 14.) The thick, narrow band left on the tree trunk prevents the cater-

pillars from reaching the foliage. This material is cheap and effective and is easily applied because it is not necessary to scrape the trees before putting it on.

³ For full information concerning the preparation and use of this tree-banding material, apply to the gipsy-moth laboratory of the U. S. Department of Agriculture at Melrose Highlands, Mass.

Spraying.—The most effective spray material for the gipsy moth is lead arsenate. It can be used either in the paste or powder form. More effective control is possible if a sticker is used in the spray. Fish oil or raw linseed oil is excellent for that purpose, as it binds the insecticide firmly to the foliage so that it is not removed by rain or heavy dew.

If either of these stickers is used the quantity of poison can be reduced from 12½ pounds to 10 pounds of paste lead arsenate or from 6¼ pounds to 5 pounds of powdered lead arsenate to each 100 gallons of water. Fish oil is cheaper than linseed oil and is therefore to be preferred. Four fluid ounces of the oil is required for each pound of poison used; in other words, 1¼ pints of oil for 5 pounds of powder or 10 pounds of paste.

In preparing the spray mixture, add the oil after the poison is well mixed with water in the tank and while the whole is being agitated. Continue the agitation while the spray is being applied. The best grade of fish oil, known as "light pressed," should be used, as the cheaper grades are not so satisfactory.

Warning.—It is not safe to allow cattle to graze beneath trees after the spray has been applied. When there is danger of the spray drifting on buildings they should be drenched with water before and after the trees are treated. A garden hose can be used for this purpose. If the spray is allowed to dry, it will disfigure the painted surface. Spraying fruit trees with this mixture is not recommended if the fruit is more than half grown, as it is difficult to remove the spray residue.

For small operations the ordinary orchard sprayer may be used with one or more lines of hose, depending on the pressure that is available. Nozzles of the Vermorel or Bordeaux type are satisfactory.

In case large shade trees or valuable park or woodland trees are to be treated, the use of a high-power sprayer is more economical. The type that has given the most satisfactory results in the gipsy-moth work develops sufficient power to throw a solid stream of spray so it will drift through the trees. (See illustration on front cover.) The nozzle is constructed so that the stream will break into a fine mist high in the air, and this results in very satisfactory and rapid work. With such a sprayer it is unnecessary to use a small hose and climb trees, which is a slow and expensive operation.

A satisfactory high-power sprayer for this work is equipped with not less than a 10-horsepower gasoline engine and a triplex pump capable of delivering at least 35 gallons of liquid per minute at a

FIGURE 14.—Applying tree-banding material to protect the foliage from gipsy-moth caterpillars

pressure of at least 300 pounds. This machinery, together with a 400-gallon tank, should be mounted on a well-built truck. (Fig. 15.) High-grade 1-inch spray hose is required, and a nozzle pressure of 300 pounds or more is necessary, depending on the length of the hose and the elevation of the nozzle above the level of the sprayer. Machines that will develop 1,000 pounds working pressure are necessary in rough country where extremely long lines of hose are required. A machine of this capacity equipped with high-pressure hose will supply a hose line 5,000 to 6,000 feet in length and spray satisfactorily at an elevation of 600 feet above the level of the sprayer.

By using a small device, which is attached to the nozzle and is known as a spreader, it is possible to spray low growth very satisfac-

FIGURE 15.—High-power truck sprayer

torily. Fruit trees can be treated very rapidly in this way. (A nozzle equipped with spreader is shown in operation at the left of the picture on front cover.)

Tests have been made for several seasons with airships of both the heavier-than-air and lighter-than-air type, to determine whether infested woodland areas could be dusted satisfactorily and economically by these means. This method has not proved entirely satisfactory for use on forest growth. The cost is high, and the dust does not adhere as well as liquid spray when an adhesive like fish oil is used. For certain low-growing crops that have a high annual value this method has given excellent results in some sections of the country.

METHODS TO BE USED IN ORCHARDS

The methods to be used for controlling the gipsy moth in orchards depend largely on the severity of the infestation. If only a few egg clusters are present in the orchard, the early spraying which is

applied for the codling moth after the blossoms fall will be found useful, provided the quantity of lead arsenate used is increased to 10 pounds of paste, or 5 pounds of powder, to 100 gallons of water. This spray should be especially effective if fish oil is added. If the infestation is more serious, a second spraying early in June, with a similar quantity of poison, but without fish oil, will be found very satisfactory. Where the infestation is severe, it will probably be necessary to creosote egg clusters in the winter and spray in the spring. Thoroughness is a prime essential if good results are to be obtained. All poor or hollow trees should be removed; and if badly infested woodland is near by, the orchard trees should be banded.

Orchard infestations can be managed by following these methods, and it will not require much additional expense or a great deal of extra work to protect the trees. In making this statement it is assumed that the orchard is being cared for by up-to-date methods to protect it from the codling moth and other injurious insects and diseases, and it is not likely that these results can be brought about in neglected orchards or where the owners do not practice the best horticultural methods in handling their growing trees.

METHODS ADVISABLE IN CITIES AND TOWNS

The same methods that are used in orchards are applicable in cities and towns and for the treatment of park and shade trees. In certain instances it would probably be advisable to use tree-banding material or burlap, preferably the former, and to discontinue spraying in cases where the infestation is very light. If the infestation is bad, creosoting, banding, and spraying should all be used in their season, in order to bring the insect under control and reduce the numbers present to a minimum.

In town or city work careful attention should be given to near-by woodlands or isolated trees, particularly if they are located on high elevations immediately outside the residential area, as colonies in such locations may furnish a supply of caterpillars which will be distributed by wind throughout the town after it has been cleaned. In lightly infested woodland areas spraying may be used exclusively for control, but when heavy infestations exist, a combination of the most effective measures is usually necessary.

The method of handling the gipsy moth in any town, city, or park, or on private estates, should be based on the degree of infestation as determined by some one who is familiar with gipsy-moth work, if the best results are to be secured at a minimum expense. Much energy and money may be wasted in applying remedies unless their application is based on a thorough knowledge of existing conditions. An owner of an infested estate should have an examination made by some qualified person who can give reliable recommendations as to treatments. It should be borne in mind that conditions as to infestation vary from year to year, and this should be considered when plans for treatment are being made.

CONTROLLING THE GIPSY MOTH IN WOODLAND

Satisfactory control of the gipsy moth in woodland by the employment of hand methods such as have already been mentioned is im-

practicable under present conditions, unless the tree growth is particularly valued for purposes other than lumber. If the woodland is situated near a large city and occupies space that is likely to be utilized in a few years for building lots, considerable money may be expended to advantage in protecting the trees, as these will make the property much more valuable when the land is subdivided. Limited areas of woodland on private estates may be of sufficient value to the owners to justify a considerable expenditure for moth destruction. In all cases, however, the species of trees involved should be carefully studied before a plan of work is adopted, in order that the expense may be reduced as much as possible. Unfortunately the difficulty of treating the woodlands in the infested area of New England is considerably increased by the fact that they are for the most part composed of numerous species in mixture.

FIGURE 16.—Mixed deciduous and coniferous woodland before thinning

Experiments have shown that most coniferous trees are not injured by the gipsy moth if grown in isolated pure stands; and if the growth is such that the trees can be thinned to a stand of immune species, no hand suppressive measures are necessary in order to prevent injury by this insect. Such lots will not be attacked by the brown-tail moth, as the larvae of this insect do not feed on conifers.

If mixtures containing a large percentage of deciduous trees are to be protected from moth injury, the species involved should be carefully considered before a decision is reached as to the best methods of treatment. Sometimes practical methods of thinning (figs. 16 and 17) can be adopted so that species will be left that are only slightly subject to injury by these insects. Experiments have shown that mixtures of chestnut, pine, red maple, ash, and hickory, regardless of the proportion of each species, are seldom injured by the gipsy moth.

In woodlands the oaks are the most favored food plant of this insect, and, unfortunately, the infested region abounds in large areas where these species predominate. At present there seems to be no method aside from hand treatment which will prevent serious injury to oak woodland, but as a large part of such land consists of poor sprout growth, the amount of damage sustained is not always so great as it might at first appear. The greatest injury likely to be caused in such areas where oaks and gray birch abound is the dying of small pines or other valuable species which have been denuded by the caterpillars after the oaks and birches have been defoliated. This reduces greatly the chance that the sprout growth will be replaced by any species of value that can withstand gipsy-moth attack. Consideration is being given to devising some economical method for protecting and improving wood lots of this character at

FIGURE 17.—Same woodland after growth favored by the gipsy moth has been removed by thinning

moderate expense. It is true that in considerable areas of oak woodland the trees, although not mature, could be utilized for small timber, railroad ties, or cordwood, and in cases of bad infestation such woodland should be promptly cut if the wood can be sold to advantage. On cheap cut-over or infested lands in many sections of the territory white or red pine might be planted to advantage, but as this involves considerable expense and as the crop can not be harvested for a period of years, the question of the desirability of managing any wood lot in this way must, in the end, be decided by the owner of the property.

If the practice common in some European countries of maintaining municipal or state forests were well developed in the New England States, it would be possible in a period of years to transform considerable areas of land which are now destined to be worthless, and which form a favorable feeding ground for the gipsy moth, into well-managed forests of valuable growth.

THE BROWN-TAIL MOTH

INTRODUCTION AND SPREAD

The brown-tail moth (*Nygmia phaeorrhoea* Don.) was first found in the United States in Somerville, Mass., in the summer of 1897, and was undoubtedly introduced several seasons previous to that time on imported nursery stock. The insect increased enormously, and as the caterpillars were particularly fond of the foliage of fruit and ornamental trees and shrubs, they became an unbearable nuisance, particularly in residential sections. Not only was complete defoliation common in early summer, but as the hairs from the caterpillars caused serious poisoning to human beings, the presence of this pest became a veritable scourge in densely populated sections. The insect extended its range very rapidly because the moths of both sexes fly freely. This species occurs in many sections of Europe and is frequently seriously injurious.

The State of Massachusetts applied suppressive measures from the winter of 1897 until February, 1900, when this work was discontinued as was the case with that against the gipsy moth. By 1905, when work on the gipsy moth was resumed, the brown-tail moth was extremely abundant in eastern Massachusetts. It was also present in enormous numbers in Rhode Island, southern New Hampshire, and southeastern Maine. Not only did fruit and shade trees suffer defoliation, but large areas of oak woodland, particularly sprout growth, was completely defoliated.

The insect continued to spread until 1915, when most of the area east of the Connecticut River, with the exception of a portion of northern New Hampshire and Maine, was heavily infested. Some infestation also existed in Vermont and west of the Connecticut River in Massachusetts and Connecticut. Suppressive work was carried on by the States and the Federal Government and the residential sections were kept fairly free from this pest.

The area now infested by the brown-tail moth is shown in Figure 3, and is much less extensive than the maximum limits to which it had previously spread. Unfavorable winter conditions, particularly in the northern part of the territory, coupled with the work of natural enemies and disease and the continuous repressive measures used in the residential sections, have caused a remarkable decrease in the abundance of this pest. The insect has been found in Nova Scotia and New Brunswick, but the area of the infested territory has been greatly reduced there and it is only moderately abundant at the present time in sections of Nova Scotia.

This pest can easily be shipped on woody plants, but general spread in this manner has been prevented by the strict enforcement of inspection and quarantine regulations.

LIFE HISTORY OF THE BROWN-TAIL MOTH

The different stages in the development of the brown-tail moth are shown in Figure 18.

THE EGGS

The female moth deposits a small cluster of eggs on the underside of a leaf. They are usually laid in July, and are covered with brown

FIGURE 18.—Different stages of the brown-tail moth: A, Winter nest; B, male pupa; C, female pupa; D, cocoon in leaves; E, young caterpillars on leaf; F, full-grown caterpillar; G, female depositing eggs on a leaf, and egg mass also on leaf; H, egg mass removed from leaf and with some of the eggs exposed; I, male moth; J, female moth. All about three-fourths natural size

hair taken from the body of the female. Hatching begins about August 15.

THE LARVAE

The newly hatched larvae or caterpillars feed on the epidermis of the leaf. After molting once or twice they begin to construct a winter web. This is made by drawing together several terminal leaves and securely fastening them by silk which is secreted by the

caterpillars. The larvae from one or more egg clusters live and feed in common, and as cold weather approaches they retire to the web, in which they remain during the winter. In the spring these larvae leave the web as soon as the buds begin to develop and feed upon the bud scales and small leaflets. They become full grown about the middle of June.

THE PUPAE

After the caterpillars finish feeding they spin loose silken cocoons and pupate within them. These cocoons are sometimes constructed separately, but in many cases large numbers of them are spun in a single mass. About two weeks are spent in the pupal stage.

THE ADULTS

Emergence of the moth usually begins the first week in July. The adult brown-tail moth is pure white. The abdomen of the female is much larger than that of the male, but in both sexes the tip of the abdomen is covered with dark-brown hairs. These moths are attracted to strong light, such as electric arc lights; and as they fly at night, it is often possible to secure many specimens around the arc lights in cities and towns during the first half of July.

FOOD PLANTS OF THE BROWN-TAIL MOTH

The caterpillars of the brown-tail moth commonly feed on the leaves of apple, pear, cherry, oak, and willow, and they are sometimes found in considerable numbers on other common deciduous trees and shrubs. They never attack conifers and are seldom found on hickory, ash, chestnut, or birch.

Oak foliage is one of the favored foods of the brown-tail moth caterpillars in Europe and was severely injured in New England for a number of years after this insect became established there. It has not suffered in this way in recent years except in 1926 and 1927, when large numbers of webs were found on oak trees and the caterpillars caused considerable defoliation in some localities.

INJURY CAUSED BY THE BROWN-TAIL MOTH

The principal injury caused by the brown-tail moth is due to the feeding of the larvae in the spring. If the infestation is bad the caterpillars are often numerous enough to devour the leaves as fast as the trees are able to develop them. As the webs are made on the terminals, the growth of the trees is often severely checked. In severe infestations trees may be completely stripped (fig. 19), but as the larvae become full grown during the first part of June, there is usually an opportunity for the trees to produce new leaves before midsummer. The young larvae that hatch in August frequently skeletonize the leaves to a considerable extent. This does not damage the trees seriously, as the growing period for the season is nearly completed.

The bodies of the caterpillars of the brown-tail moth are provided with poisonous hairs. A microscopic examination of these hairs shows that the edges are barbed in such a way that when they come in contact with the human skin and are pressed into the flesh, intense

irritation is caused. These hairs are hollow and contain a poisonous substance which acts on the blood corpuscles. This causes serious poisoning and severe irritation, accompanied by external swelling, which is known as the brown-tail rash. Persons differ considerably in their susceptibility to this poison, but many cases are reported each year in the infested region, most of which are more serious than those of ivy poisoning. Many camps and summer cottages, particularly in wooded areas, can not be occupied with any comfort during the early summer if the caterpillars are abundant, on account of the poisoning due to these caterpillars. If clothing is hung on the line near badly infested trees the hairs frequently find lodgment and are brought into the houses, and severe poisoning may result later.

FIGURE 19.—Apple trees defoliated by the brown-tail moth. Note the hibernating webs on the twigs

The brown-tail moth has been less abundant during the last three years than heretofore. There have been local infestations, however, where serious injury resulted, particularly in northeastern Massachusetts, southeastern New Hampshire, and southwestern Maine, and in some of the river valleys in New Hampshire.

NATIVE ENEMIES OF THE BROWN-TAIL MOTH

One of the important native enemies of the brown-tail moth is a fungous disease (*Entomophthora aulicae* Reich.) which attacks the caterpillars, particularly in the spring. It was first reported in this country by Roland Thaxter in 1888. Like all diseases of this nature, the benefit derived from it is regulated largely by favorable or unfavorable weather conditions. This fungus sometimes works to a slight degree on the small caterpillars in the fall, and is found occasionally in the winter webs. As a rule, however, the greatest

mortality of caterpillars takes place in the spring, when they are nearly full grown, and the pupae may, under the most favorable conditions, be almost completely exterminated.

Native parasites and predacious insects have done very little to check the increase of the brown-tail moth.

INTRODUCED PARASITES AND ENEMIES

Some of the parasites (*Apanteles lacteicolor* Vier. and *Compsilura concinnata* Meig.) that were introduced as enemies of the gipsy-moth caterpillars also attack those of the brown-tail moth freely, and one additional introduced species (*Meteorus versicolor* Wesm.) is an important enemy of the insect. The Calosoma beetle (*Calosoma sycophanta* L.), both in the adult and larval stages, destroys many brown-tail moth larvae and pupae.

The parasites and natural enemies of this insect are more effective than is the case with the gipsy moth, and extremely low temperature in the winter often proves fatal to a large proportion of the small caterpillars in the webs.

These factors, together with the enormous amount of hand suppression that has been employed in many parts of the infested area, have resulted in a pronounced decrease in the abundance of the insect during the last few years.

METHODS OF CONTROLLING THE BROWN-TAIL MOTH

The brown-tail moth can be controlled by cutting off the winter webs and burning them before the caterpillars begin to emerge in April. These webs should be destroyed by fire, for if they are simply cut from the trees and left on the ground, the caterpillars will emerge, and no benefit will result from the work which has been done.

In orchard practice it is sometimes inadvisable to cut the winter webs, for where an infestation is bad the cutting is likely to leave poorly shaped trees. Spraying in the spring is not a satisfactory remedy unless the infestation is very light, because the caterpillars, when they occur in large numbers, do not allow the tree to put out sufficient foliage to hold the spray material.

The most effective method in orchards is to spray the trees before the middle of August, using 3 pounds of lead-arsenate powder to 100 gallons of water. Before spraying operations of this sort are attempted the orchardist should determine to what extent the trees are infested with egg masses of the brown-tail moth. If the infestation is very slight, it will be more satisfactory to cut and destroy the webs. If the infestation warrants, both shade, ornamental, and fruit trees may be sprayed to advantage at this time. Caution should be used, however, in spraying fruit trees, particularly if early-fall varieties are to be treated. The foliage should be treated, particularly the terminal shoots, and as much care as possible should be exercised not to cover the fruit. Late fall or winter varieties of fruit may be sprayed with lead arsenate in August, and although an occasional spot may be found on the fruit at the time of picking, no injury will result from it. In cases where only a few choice fruit trees are sprayed it is practicable to wipe the fruit before packing for sale;

but this will not be necessary if care is taken to spray only the terminal growth of the trees, as this is where the bulk of the egg clusters are deposited.

The damage caused by the brown-tail moth is ordinarily not so severe as that resulting from gipsy-moth infestation, because the brown-tail moth does not have so wide a range of food plants and, further, because the bulk of the feeding by the larvae is done early in the season, so that the trees have an opportunity to recover before midsummer. In the territory where both insects exist the caterpillars of the gipsy moth supplement the feeding which has been done by those of the brown-tail moth, and the injury is, therefore, greatly increased.

During the last few years this insect has seldom been found feeding on oak foliage, except in the worst infested localities, and elm, maple, and rose are not as badly infested as was the case about 15 years ago.

Thorough destruction of brown-tail moth webs in residential sections and in orchards has resulted in materially decreasing the insect in thickly settled sections. Elimination of worthless apple and wild cherry trees would help greatly in reducing the pest.

ORGANIZATION AND STATUS OF WORK AGAINST THE GIPSY MOTH AND THE BROWN-TAIL MOTH

Each State infested with these insects is carrying on control work, and many of the towns and cities are similarly engaged.

The Dominion of Canada and several of the Provinces have taken up control and eradication work.

MAINE

In Maine the work is in charge of the commissioner of agriculture, who has authority to appoint assistants to take charge of the operations. In the southwestern part of the State the gipsy-moth infestation is general and in places severe, but the greater part of the known infested area is only slightly infested. The severity of infestation increased in 1926 and 1927, and many heavily infested sections were found in 1928. The brown-tail moth is not now seriously abundant except along the seacoast in the southwestern part of the State. Over 12,000 square miles of territory are infested with the gipsy moth and about 7,000 with the brown-tail moth. The former has gradually spread to new territory, but the area infested by the latter has decreased considerably since 1919 and has not changed very greatly for several years.

NEW HAMPSHIRE

The moth work in New Hampshire is in charge of the State entomologist. Over 8,000 square miles are infested with the gipsy moth and about 4,000 with the brown-tail moth. Most of the territory south of Lake Winnepesaukee has suffered severe and repeated defoliation. In 1927 the largest defoliated areas centered around the lake and toward the east and northeast. In 1928 severe defoliation was general throughout the section east of the Merrimac River and to the foothills of the White Mountains. Serious stripping also occurred west of the river in the southern part of the State.

VERMONT

In Vermont the moth work is in charge of the commissioner of agriculture, who appoints an entomologist to carry on the field work. About 6,000 square miles are infested by the gipsy moth, the worst localities being along the Connecticut River. An occasional web of the brown-tail moth has been reported in recent years, but this insect does not appear to maintain itself in numbers in the State.

MASSACHUSETTS

The commissioner of conservation has charge of the moth work in Massachusetts. Each infested town is required by law to select a local superintendent, whose appointment must be approved by the commissioner. Owners are required by law to keep their property free from these pests, but can not be compelled to expend for this purpose more than \$5 per year on each \$1,000 assessed valuation. Towns and cities must expend funds for proper treatment of the street trees and those in parks and on public grounds. After the amounts fixed by law are expended, financial aid may be supplied by the State. The gipsy moth at present occurs to a greater or less extent in every town in the State east of Berkshire County. It has increased in abundance over much of the infested area during the last three years. The area infested by the brown-tail moth has been reduced so that at present it amounts to only a little over 4,000 square miles.

RHODE ISLAND

In Rhode Island the commissioner of agriculture has charge of the moth work, which is done under the supervision of the State entomologist. The gipsy moth occurs in all towns in the State, and several heavy infestations were treated during 1927 and 1928. The brown-tail moth infestation has decreased rapidly in recent years, and at present none of the towns are known to be infested.

CONNECTICUT

In Connecticut the work is in charge of the State entomologist. The brown-tail moth has not been found in this State for several years. The gipsy-moth area now occupies about 3,000 square miles. Control work has heretofore been effective in controlling the insect, but during the past two years the infestation has been increasing somewhat in the eastern part of the State.

NEW YORK

In New York State the field work against the small number of scattered colonies that had been found up to March, 1923, was supervised by the director of the bureau of plant industry, of the department of farms and markets. At that time the work was transferred to the department of conservation, which has worked co-operatively with the Bureau of Entomology in the New York section of the barrier zone.

NEW JERSEY

In New Jersey the State work is under the control of the State department of agriculture. The funds appropriated have been used in conjunction with Federal funds, and the direct supervision of the work has been in the hands of the United States Bureau of

Entomology. The results, up to the present time, have been very satisfactory.

PENNSYLVANIA AND OHIO

Work on one small colony in Pennsylvania and another in Ohio was done by arrangement with the State officials—in Pennsylvania with the bureau of plant industry of the department of agriculture of that State and in Ohio with the State nursery inspector. The results were satisfactory, as both colonies of the insects have been exterminated.

WORK OF THE DOMINION OF CANADA

Work on the gipsy moth and the brown-tail moth in Canada is conducted by the division of foreign pests suppression of the office of the Government entomologist. Cooperation with the Provincial governments of New Brunswick, Nova Scotia, and Quebec has been maintained. The brown-tail moth is not now known to occur in the Province of New Brunswick but is found in decreasing numbers in the Annapolis Valley in Nova Scotia. The gipsy-moth colony found at Henrysburg, Quebec, has been exterminated through the efforts of the Dominion and Provincial officials, and the insect is not known to occur in Canada at the present time.

WORK OF THE UNITED STATES DEPARTMENT OF AGRICULTURE

The gipsy-moth project has for many years been conducted by the Bureau of Entomology. On July 1, 1928, the quarantine and control work was transferred to the Plant Quarantine and Control Administration, and the research work was assigned to the Division of Forest Insects of the Bureau of Entomology.

The object is to control and prevent the spread of these insects. For a number of years it was believed to be quite impossible to restrict the spread of the brown-tail moth on account of the heavy migration of the adults of this species. In recent years, owing principally to the effectiveness of parasites introduced by the bureau, winter conditions unfavorable to the insect, and in some sections the effectiveness of a fungous disease which attacks the caterpillars, in addition to a large amount of hand suppression work, it has been possible not only to prevent spread but to record a marked reduction in the infested area. In certain sections near the seacoast in New Hampshire and northern Massachusetts and near some of the rivers, however, rather heavy infestations exist. One of the dangers in connection with the brown-tail moth at present is that favorable climatic conditions or temporary scarcity of natural enemies may enable it to increase rapidly and spread proportionately.

The gipsy moth was held in check for several years and did not spread beyond the Connecticut River. During the war period, when extreme difficulty was experienced in maintaining a field force on an effective basis, and funds were inadequate to meet the increased cost of operation, infestations developed rapidly, and strong winds immediately following the hatching period caused an extensive spread of the insect to the west and northwest. Field work indicated that many colonies were established during that period, as scattered infestations were found in the Berkshire Hills section of Massachusetts and Connecticut and in the Green Mountain region in Vermont.

This territory is extremely rugged and very difficult for field operations.

The establishment of the barrier zone has already been discussed in some detail, and the results secured thus far have justified the effort expended.

A large amount of experimental work has been done to devise better methods of controlling these insects. Natural enemies have been introduced from Europe, northern Africa, and Japan and colonized throughout most of the infested area. Specialists have spent considerable time studying the insects in their native homes and collecting parasites for shipment to this country. This phase of the work is being pressed as rapidly as possible. Investigations looking toward the planting and growing of forest trees that are not subject to attack by the gipsy moth are being continued.

The entire area known to be infested by either of these insects is under quarantine, and shipments of nursery stock, lumber, cordwood, and other forest products, including Christmas trees and greenery, and stone and quarry products are not permitted to leave the territory unless they are inspected and accompanied by a certificate stating that they are free from infestation.

COOPERATIVE WORK

Since the gipsy moth and brown-tail moth work was begun by the Bureau of Entomology more or less work has been done in cooperation with the States concerned. The introduction of parasites and natural enemies of the gipsy moth was conducted in cooperation with the State of Massachusetts for several years. After the infestation had covered large areas in other States an arrangement was made for this work to be carried on by the Bureau of Entomology. The general plan of field work in New England is for the States to manage the clean-up east of the barrier zone while the Federal forces work in the zone and cooperate in making the entire work effective.

In New York the barrier zone work is handled in cooperation with the State authorities.

The work in New Jersey is planned with the purpose of exterminating this large infestation, and close cooperation with the State has been maintained.

The gipsy moth and brown-tail moth quarantine covers the entire infested area, and its enforcement has thus far prevented long-distance spread of these pests.

ORGANIZATION OF THE UNITED STATES DEPARTMENT OF AGRICULTURE WHEN THIS PUBLICATION WAS LAST PRINTED

<i>Secretary of Agriculture</i> -----	ARTHUR M. HYDE.
<i>Assistant Secretary</i> -----	R. W. DUNLAP.
<i>Director of Scientific Work</i> -----	A. F. WOODS.
<i>Director of Regulatory Work</i> -----	WALTER G. CAMPBELL.
<i>Director of Extension Work</i> -----	C. W. Warburton.
<i>Director of Personnel and Business Administration.</i>	W. W. STOCKBERGER.
<i>Director of Information</i> -----	M. S. EISENHOWER.
<i>Solicitor</i> -----	E. L. MARSHALL.
<i>Weather Bureau</i> -----	CHARLES F. MARVIN, <i>Chief.</i>
<i>Bureau of Animal Industry</i> -----	JOHN R. MOHLER, <i>Chief.</i>
<i>Bureau of Dairy Industry</i> -----	O. E. REED, <i>Chief.</i>
<i>Bureau of Plant Industry</i> -----	WILLIAM A. TAYLOR, <i>Chief.</i>
<i>Forest Service</i> -----	R. Y. STUART, <i>Chief.</i>
<i>Bureau of Chemistry and Soils</i> -----	H. G. KNIGHT, <i>Chief.</i>
<i>Bureau of Entomology</i> -----	C. L. MARLATT, <i>Chief.</i>
<i>Bureau of Biological Survey</i> -----	PAUL G. REDINGTON, <i>Chief.</i>
<i>Bureau of Public Roads</i> -----	THOMAS H. MACDONALD, <i>Chief.</i>
<i>Bureau of Agricultural Economics</i> -----	NILS A. OLSEN, <i>Chief.</i>
<i>Bureau of Home Economics</i> -----	LOUISE STANLEY, <i>Chief.</i>
<i>Plant Quarantine and Control Administration</i> -----	LEE A. STRONG, <i>Chief.</i>
<i>Grain Futures Administration</i> -----	J. W. T. DUVEL, <i>Chief.</i>
<i>Food and Drug Administration</i> -----	WALTER G. CAMPBELL, <i>Director of Regulatory Work, in Charge.</i>
<i>Office of Experiment Stations</i> -----	-----, <i>Chief.</i>
<i>Office of Cooperative Extension Work</i> -----	C. B. SMITH, <i>Chief.</i>
<i>Library</i> -----	CLARIBEL R. BARNETT, <i>Librarian.</i>

BULLETIN OF THE U.S. DEPARTMENT OF AGRICULTURE

No. 170

Contribution from the Bureau of Entomology, L. O. Howard, Chief.
February 9, 1915.

THE EUROPEAN PINE-SHOOT MOTH; A SERIOUS MENACE TO PINE TIMBER IN AMERICA.

By AUGUST BUSCK,
Entomological Assistant, Forest Insect Investigations.

INTRODUCTION.

One of the most injurious insects to pine forests in Europe is a small orange-red moth, the larva of which eats out the new buds and kills or deforms the young twigs of pine trees, so as seriously and permanently to lower their timber value. This European pine-shoot moth, which is known under the scientific name *Evetria buoliana* Schiffermiller, has within very recent years been accidentally introduced into America on imported European pine seedlings and has unfortunately become established in several widely separated localities in the eastern and middle western States.

Early last summer (1914), a correspondent of the Bureau of Entomology complained of a serious insect injury to European pines under his surveillance on Long Island, and sent examples of the injury and of the larvæ causing it; the latter could not be identified as those of any of our known American pine pests, and the writer was therefore authorized to visit the affected localities in order to ascertain the extent of the injury and to obtain sufficient live material for study and rearing. From this material a large number of moths emerged during the latter part of June and these were at once recognized as the famous European pine-shoot moth.

Subsequent surveys, undertaken by the bureau through Mr. Carl Heinrich and the writer, established the fact that the species has been repeatedly introduced on European nursery stock, and that it has become established in nurseries and parks in several localities scattered over nine States.

In view of the experience with other introduced European insects, and considering the very serious financial loss caused abroad annually by this insect, its introduction into this country gives just cause for alarm, because incalculable injury may result to the vast American forest interests if this insect is permitted to become generally established on our native pines.

Some idea of the extent and permanent character of the injury which this insect is capable of inflicting may be gained from the illustration (Pl. I) of a European pine forest which has been infested by it for several years in succession, with the result that a majority of the tree trunks are so twisted and crooked that their value as timber is materially lessened.

HISTORY OF THE SPECIES IN EUROPE.

The species is a constant menace to pine forests in Europe and annually causes serious depredations, especially to young plantations of pine, in spite of continual preventive work against it. It has been the subject of much study and of an extensive literature from the time it was first described by Schiffermiller in 1776 to the present day. The species was named in honor of a Vienna entomologist, Baron Buol, who studied its injurious work during the latter part of the eighteenth century; since then numerous accounts have appeared of particularly severe outbreaks in many parts of Europe, from England to Russia, and from Scandinavia to southern France. It also occurs in Siberia.

One such outbreak in Denmark, in 1805–1807, is recorded by Niemann (1809).¹ This was so serious as nearly to cause pine culture to be abandoned in that country as hopeless. It is interesting to note that at that time the same preventive means were resorted to as are now employed against the insect, namely, the wholesale pruning and burning of all infested twigs.

The German forest entomologist, Ratzeburg, counted *Evetria buoliana* one of the most injurious forest insects and gave a detailed account (1840) of the life history, structure, and economic importance of the species. He mentioned especially an unusual outbreak in 1836–1838, which covered many parts of Europe. In the province of Furstenau the Rochesberg Mountain, which was covered with pines, became so seriously infested that it was under consideration by the authorities to burn it off and plant new trees. Other localities were only saved by strenuous systematic collecting of the infested twigs; thus, in the small province of Kesternich alone, 150,000 larvæ were gathered and destroyed.

Judeich and Nitsche (1895) state that the injury caused by *Evetria buoliana* is often fatal to the pine plantations. To quote from these authors, "If the attack is slight, it results in the branching of the tree, but if the attack is more severe and continued for several years, as we have seen it, then hardly any bud is spared and the pines become stunted into miserable small bushes from which numerous

¹ Dates in parentheses refer to "Literature," pp. 10–11.

branched shoots and large needle tufts stick out." These authors record many severe outbreaks and mention especially one in 1883-1885, in the Royal Forest Reserve, Pillnitz in Saxony, where nearly 75 acres of young pines planted in 1878 became infested to such an extent that hardly a shoot was spared, and in 1884 the entire plantation presented a pitiful, crippled appearance.

J. E. V. Boas (1898), who has made original investigations of the insect in Denmark, considers it one of the most injurious insects affecting forest trees. Among other outbreaks he mentions one in Jutland, Denmark, extending through several years around 1870, which "threatened the total destruction of the pine plantations."

The Belgian authority on forest insects, G. Severin (1901), regards *Evetria buoliana* as the most injurious insect to pines in Europe, and emphasizes the lasting injury to the timber resulting from even slight attacks of this insect.

All other European handbooks on entomology or on forestry contain similar accounts of this insect and express the same opinion as to its destructiveness to pine.

FOOD PLANTS.

Evetria buoliana is confined to pine and does not attack other coniferous trees, as spruce or larch, even though these grow alongside of the infested pines. While the species is most often mentioned on the yellow pine, or Scotch pine,¹ in Europe, because this is preeminently the forest tree of importance there, it attacks all species of *Pinus* indiscriminately, according to Ratzeburg and other authorities, and the American infestations have come in on European seedlings of the Austrian pine² and on mughus pine³ quite as often as on Scotch pine.

According to Ratzeburg and Severin, it also attacks and is equally injurious to American white pine,⁴ which is cultivated in Europe; and Mr. Carl Heinrich found the species on a small lot of another native American pine,⁵ which was growing immediately surrounded by infested European pine seedlings.

These latter records are particularly significant, as they prove beyond question that the pest will spread to our native American pines if not prevented.

The species attacks mainly young trees between 6 and 15 years of age, but it is often excessively destructive to younger plantings and seedlings and injurious also to older trees, though trees of 30 years or older are rarely seriously affected.

¹ *Pinus sylvestris*.

² *Pinus laricis* var. *austriaca*.

³ *Pinus montana* var. *mughus*.

⁴ *Pinus strobus*.

⁵ *Pinus resinosa*.

INTRODUCTION AND DISTRIBUTION IN AMERICA.

American nurseries have imported many thousands of pine seedlings annually from Europe, especially from France, Belgium, Holland, Germany, and England. Importations normally take place in the fall, winter, and early spring. At this time of the year the young larvæ of the pine moth lie dormant within the buds, so that an infestation is easily overlooked. It is evident that the pest has been present in a number of shipments of late years and that it thus has been introduced repeatedly into American nurseries. In a great majority of these cases, however, the species has been unable to establish itself and has died out during the first year. Many of the larvæ die from overheating en route, or from various other unfavorable circumstances incident to the handling and transplanting of the seedlings under different climatic conditions. Only by a combination of favorable conditions would the few surviving larvæ have been able to develop into moths and succeed in establishing the species in this country. This is probably the reason why the species as yet has become established in comparatively few American localities. It appears that such established infestation has taken place only in very recent years and especially within the last two years, or since the demand for European pines has become general.

Up to the present time the European pine moth has been discovered in only 32 nurseries and private estates, representing 20 localities in 9 States, namely:

State.	Locality.	Discovered in—
Illinois.....	Chicago.....	Private grounds.
Do.....	Glenview.....	One nursery.
Do.....	Dundee.....	Do.
Do.....	Western Springs.....	Do.
Do.....	Deerfield.....	Do.
Do.....	Kenilworth.....	Two private grounds.
Do.....	Bloomington.....	One nursery.
Ohio.....	Tippecanoe City.....	Do.
West Virginia.....	Elm Grove.....	Do.
Pennsylvania.....	Pittsburgh.....	Private grounds.
Do.....	Philadelphia.....	One nursery.
New Jersey.....	Somerville.....	One estate.
New York.....	Long Island.....	Nine nurseries and estates.
Do.....	Tarrytown.....	One nursery and one estate.
Do.....	Elmsport.....	One estate.
Massachusetts.....	Dedham.....	One nursery.
Do.....	North Abington.....	Do.
Do.....	Worcester.....	Do.
Connecticut.....	New Canaan.....	Do.
Rhode Island.....	Newport.....	Two nurseries and one estate.

In none of these localities, except on Long Island, has the species existed for more than the last two years, and in most of them it has become established only within the last year.

But the survey for this insect has so far covered only about 60 localities, which could be reasonably suspected to harbor the pest because it was known that importations of European seedlings had

WORK OF THE EUROPEAN PINE-SHOOT MOTH (EVETRIA BUOLIANA).

Section of European pine forest showing deformations in the trunk of *Pinus sylvestris* resulting from several consecutive years' injury. (After G. Severin.)

STAGES OF THE EUROPEAN PINE-SHOOT MOTH.

Moth and full-grown larva; both greatly enlarged. (Original.)

[Drawings by Miss Mary Carmody.]

WORK OF THE EUROPEAN PINE-SHOOT MOTH.
Fall work of the young larvae in the buds of *Pinus sylvestris*. (Original.)

WORK OF THE EUROPEAN PINE-SHOOT MOTH.

Malformations in pine resulting from injury by this pest. (Original.)

WORK OF THE EUROPEAN PINE-SHOOT MOTH.

Twisted growth of European pines caused by the work of this insect. (Original.)

WORK OF THE EUROPEAN PINE-SHOOT MOTH.

Typical first, second, and third years' growth of pine, after the primary injury by this insect, known as "baionnette" or "posthorn." (Original.)

[Photographs by Mr. H. C. Sands, of the New York Department of Agriculture.]

taken place, and the indications are very strong that the pest has become established in several other widely distributed localities, either by direct importation from Europe or by distribution from infested American nurseries. This is particularly to be suspected of localities where large importations and plantings of European pines have been made.

As yet the pest has been found only in nurseries and private parks supplied by these infested nurseries. In no case has it yet been found on forest trees in America. The species is therefore at present mainly a nursery problem in this country and consequently may yet be controlled and possibly even eliminated by proper measures under Federal and State supervision. That this condition can not long endure and that the pest, if not checked, will soon multiply and spread to native pines outside of nurseries and pass beyond the possibility of elimination is clearly indicated by all the evidence on hand.

LIFE HISTORY.

In Europe the moths (Pl. II, upper figure) issue in July, sometimes as early as the end of June, and in the warm evenings they swarm around the pines in large numbers. During the day they sit quietly on the branches, as can be ascertained by giving the tree a sharp jolt, which will cause the moths to fly out. When the insect sits still on the food plant it is not easily discovered, for the apparently striking orange-red color blends well with the natural surroundings and therefore must be classed as a protective coloration. Early in August the eggs are laid singly on the new buds for next year's growth, the terminal cluster of buds being nearly always chosen for oviposition. The young larva soon hatches and eats its way into the bud, making itself a roomy cell by devouring the live inside part. It attains a length of only a few millimeters during the fall months, and overwinters within the hollow bud. At this stage its presence is easily overlooked, though a trained eye will discover a small exudation of pitch over the entrance hole differing from the normal exudation of the buds. (See Pl. III.)

In May, as soon as the sap begins to rise in the trees, the larva the buds. (See Pl. III.)

leaves its winter quarters and bores into the bud next thereto, in turn destroying this and as many others as it needs for food. As the remaining buds adjoining begin to grow into young shoots the larva attacks them. It eats the entire inside of the youngest shoots and these consequently die. The more developed shoots are injured only on one side, and these sometimes continue to grow, but are bent downward at the injured spot. The larva (Pl. II, lower figure) feeds only on the soft growth on which the needles have not yet appeared, and by the time the needles have developed all, or nearly all, of the shoots in the infested cluster have become dead or injured. The

larva then makes a silk-lined chamber within one of the hollow shoots and here it pupates. After about three weeks the spiny pupa pushes itself half way out through the dry wall of its chamber and the moth, or adult, issues.

The full life history of the species in America has not been ascertained, because a full year has not elapsed since it was first discovered here. While in the main it is the same as in Europe, a very distinct difference has already been noticed, due to the longer and warmer summer and fall in this country. In Europe the young larva attacks only one bud and attains very little growth before it enters the dormant winter season, but in the warmer climate of America the larva eats out two, three, or more buds and attains nearly half of its growth before winter. This, of course, tends to make the species even more injurious here than it is in Europe.

While it is altogether probable that the species has here only one generation annually, as in Europe, the possibility is not absolutely excluded that on account of the longer season it may eventually develop two generations annually like the allied native species. This, of course, would greatly increase its power for injury.

CHARACTER OF INJURY.

During the entire spring the infested twigs are very noticeable by reason of the dead and injured buds and young shoots, and the empty pupa skin sticking out of the destroyed shoot is also a familiar and easily noticed sight during the summer months; but the extent of the injury caused by this insect is only realized later in the season, when the new growth is found to be either quite destroyed or permanently injured.

As may be gathered from the foregoing account of the life history, each one of these insects does very considerable damage, not only by destroying a large number of buds and young shoots but by injuring the adjoining shoots which remain and which normally should supplant the destroyed leaders; thus the trees are permanently disfigured. These injured shoots bend downward and outward and afterwards grow upward again in a curve, in the attempt to continue the normal upward growth of the tree. This results in a characteristic malformation (Pls. IV, V, VI), so familiar in European pine forests that it has a popular name in each country—as “posthorn” and “waldhorn” in Germany and Holland and “baionnette” in France, while the few examples which have so far occurred in America have suggested the name “Dutch pipe” to those who have noticed it. This injury does straighten out somewhat during the successive years’ growth, but never can be fully remedied and will always be noticeable and a serious detriment to the timber (Pl. I). Injury of this character is the result even when the species is present in only small numbers, the

repeated infestation of the leading twigs during several consecutive seasons producing additional malformations which result in a much distorted tree of little commercial value. If the pest becomes more abundant, then the trees are transformed by the effect of the injury into unsightly crippled bushes with no commercial value.

DESCRIPTION.

THE ADULT.

(Pl. II, upper figure.)

The European pine-shoot moth is a small, gayly colored moth, about one-half inch long and measuring about three-fourths of an inch across with the wings extended. The head and its appendages and the thorax are light orange-yellow, and the abdomen is dark gray. The forewings are bright ferruginous orange, suffused with dark red, especially toward the tips, and with several irregular, forked anastomizing, silvery crosslines and costal strigulae; the hindwings are dark blackish brown. The legs are whitish, the anterior ones reddish in front.

THE EGG.

The egg is very small, flat, whitish in color, and is laid singly at the base of a bud. Dissection of a female abdomen proves that upwards of a hundred eggs are laid by each female; this is a rather greater fecundity than is normal in this group of insects.

THE LARVA.

(Pl. II, lower figure.)

The young larva is dark brown with deep black head and thoracic shield, the latter divided by a narrow central line. The body of the older larva becomes somewhat lighter, but is still much darker than the larva of any of our allied native species. The full-grown larva is two-thirds of an inch long.

THE PUPA.

The pupa is stout, robust, light chestnut brown with darker head and back. The wing covers reach to the end of the fourth abdominal segment. The abdominal segments are armed with rings of short, sharp, blackish-brown spines.

ALLIED AMERICAN SPECIES.

There are in this country several indigenous species closely allied to *Evetria buoliana*, and like it confined to pine. Some of these already constitute a serious problem and periodically do considerable

damage to pine forests and more often to pine nurseries. They are the more capable of injury because there are two generations annually and they thus have two chances each year to accomplish their damaging work. None of these native species can, however, even with this advantage, be compared in destructiveness to the European species just introduced. This is partly due to the larger size of the introduced species and to the greater voracity of the larva, but is mainly due to the difference in the attack, which causes a different reaction of the tree.

The larva of the native species of the genus confines itself to a single twig and finds its food within this or within a single bud, or at most a few buds. This bud or twig dies, but the tree responds with the natural growth of the next set of buds and very often recovers from the injury without permanent disfigurement. The resulting injury to the trees is serious only when these native species are present in unusually large numbers. Moreover, each of the native American species is more or less confined to a single or a few species of *Pinus*, but the European pine-moth thrives indiscriminately on all species of *Pinus* and has consequently a greater chance to become excessively abundant. While several of the native species are continually of some economic importance and periodically become a serious menace even to larger trees, it is mainly when they occur in large numbers in nurseries that they become really troublesome. Large trees become checked in their growth by the loss of terminal twigs, but are not necessarily seriously deformed in their future growth, although an undesirable forking of the tree top is a quite common result.

On the other hand, the larva of the European pine-shoot moth is very voracious and not only destroys a number of buds and young sprouting shoots by eating their interior, but it invariably damages the remaining shoots in the cluster by nibbling their bases on the inner side. The subsequent growth of these injured shoots, in the effort to supplant the destroyed leader, causes greater permanent injury to the value of the tree than if they were entirely removed.

NATURAL ENEMIES.

Evetria buoliana in Europe is, to some extent, kept in check by a large number of parasitic enemies. As early as 1838 Hartig¹ recorded 14 ichneumonid wasps and 1 tachinid fly² which he had reared from pupæ of the pine-shoot moth. It has since been ascertained that there are several other parasites; among the ichneumonids Ratzeburg considered the following three, which he himself had reared, as the more important: *Pristomerus vulnerator* Panz., *Cremastus interruptor* Grav., and *Orgilus obscurator* Hald.

¹ See "Literature," p. 10.

² *Actia pinipennis* Fallen.

To promote the good work of these parasites specially constructed rearing houses have been erected in Europe during bad outbreaks of the pine moth. The infested twigs are collected in these small houses, which permit the escape of the parasites but not of the moths.

It is reasonable to suppose that some of the native parasites on some of the native species of *Evetria* will in time also attack *Evetria buoliana* in this country—in fact, parasitized larvæ have already been observed—but these native parasites can not be relied upon to keep in check their natural hosts, the American pine moths, which sporadically become very abundant and injurious in spite of the parasites, and presumably will be less effective in controlling the newly introduced host.

METHOD OF CONTROL.

The larva of the European pine-moth is so effectively protected within the buds that it can not be reached by any insecticide, and the only method of combating it is that used in Europe for more than a hundred years, namely, the pruning and destruction of the infested buds and twigs together with the larvæ they contain. Such hand picking is practiced every year in the government-controlled forest reserves of Europe.

This pruning must be done while the insect is within the twigs, and while it may be done throughout the entire year, except during the midsummer months when the insect is in the adult stage, it can be most profitably done in the fall and winter months while the young larvæ are yet within the undeveloped buds, because the pruning at this time will enable the secondary set of buds to develop in the spring without delay. The only drawback to the collecting of the larvæ in the fall and winter is that the infested buds are then less noticeable than in the spring when the injury is further developed. A little practice, however, soon enables instant recognition of the infested buds, even by an unskilled laborer; the slight exudation of pitch at the base of the bud covering the entrance hole of the larva (Pl. III) is very characteristic and easily recognized when once known.

In the spring, when the buds develop into young shoots, the injury is very much more apparent, and anybody can then distinguish the infested twigs at a glance. For this reason it is advisable to have the trees gone over again in the spring, so as to remove any infestation which has been overlooked in the fall. In America the work of the larva in the fall (September, October, and November) has progressed far more and is much more easily discovered than is the case in Europe, where the larvæ have attained very small proportions and

have attacked only one or two buds before the winter resting period intervenes.

The fact that this species is stationary during the greater part of the year and only found within definite parts of certain kinds of trees, namely, in the next year's buds of pines, makes effective control work much easier than is the case with insect pests which are general feeders and which are not confined to definite parts of the food plant, as, for example, the gipsy moth or the brown-tail moth. While the European pine-shoot moth is confined to nurseries and private parks and has not spread to the native pines, it should prove a comparatively easy task to eradicate the species absolutely within any limited area. At the present time it would even seem possible completely to stamp out this dangerous pest in America, and forestall the infestation of our native pine forests, provided that the danger of new infestation is removed. But when once the species has multiplied sufficiently to become generally distributed on the native pines the possibility of eradication will be past.

SYNONYMY OF *EVETRIA BUOLIANA* SCHIFFERMILLER.

- Tortrix buoliana* Schiffermiller, Syst. Verz. d. Schmett., p. 128, 1776.
Coccyx buoliana Treitschke, Schmetterlinge von Europa, vol. 8, p. 140, 1830.
Tortrix (Coccyx) buoliana Ratzeburg, Die Forst-Insecten, vol. 2, p. 202, 1840.
Retinia buoliana Guénée, Europaeorum Microlepidopterorum index methodicus, p. 46, 1845.
Coccyx buoliana Herrich-Schäffer. Bearb. d. Schmetterlinge von Europa, vol. 4, p. 221, 1849.
Evetria buoliana Meyrick, Handbook of British Lepidoptera, p. 470, 1895.
Evetria buoliana Rebel, Catalog der Lepidopteren des palaearctischen Faunengebietes, T. II, No. 1851, 1901.

LITERATURE.¹

1776. Schiffermiller, I. Systematisches Verzeichniss der Schmetterlinge der Wiener Gegend. Wien.
 Original description of *Evetria buoliana*.
 1809. Niemann, E. Forststatistik der Danischen Staaten, Altona.
 Describes outbreak in Denmark in 1805-1807, and the collecting of larvæ in the effort to control the species.
 1838. Hartig, T. *Tortrix buoliana*. In Jahresberichte über die Fortschritte der Forstwissenschaft und forstlichen Naturkunde. Jahrg. 1, Heft 2, p. 267-268, Berlin.
 Records the rearing of 15 species of parasites from *Evetria buoliana*.
 1840. Ratzeburg, J. T. C. Die Forst-Insecten, T. 2, p. 202-207, Taf. XIV, fig. 4. Berlin.
 Detailed account with illustrations of the life history, work, economic importance, remedies, natural enemies, and literature of the species, with notes of severe outbreaks in Germany, 1835-1838.

¹This is not intended to be a complete bibliography of *Evetria buoliana*; a large number of special articles have appeared in various publications in Europe, and every handbook on insects or forestry contains more or less exhaustive accounts of this pest.

1895. Judeich, J. F., and Nitsche, H. Lehrbuch der mitteleuropaischen Forstinsektenkunde, Bd. 2, p. 1004-1008. Wien.
Condensed (5 pages), life-history and economic importance with original figure of the injury done by the species.
1897. Lovink, H. J., and Ritzema Bos, J. Schade in jonge dennen bosschen tewegg gebracht door rupsen uit het bladrollergeslacht *Retinia* Gn. ("dennenknoprupen" "dennenlotrupen" "harsbuilrupen"). In Tijdschr. Plantenziekten, Jahrg. 3, Afl. 4, p. 83-133, figs. 6, pls. V-VII, Oct.
Detailed account of the species and its injury, with colored plates.
1897. Severin, G. Insectes. Extrait du Catalogue détaillé et illustré du Pavillon des eaux et forêts à l'Exposition internationale de Bruxelles-Tervueren, p. 46-49, pl. X. Bruxelles.
Contains short illustrated account of *Tortrix (Retinia) buoliana* Schiffmiller and its injury: Plate I of the present paper has been copied from this article.
1898. Boas, J. E. V. Dansk Forstzoologi. Copenhagen.
Condensed life history, injury, and references, with original observations and figures.
1898. Hess, R. A. Der Forstschutz, ed. 3 enl., v. 1, p. 492-494, figs. 174-175. Leipzig.
Condensed handbook information on *Tortrix (Retinia) buoliana* Schiff.
1901. Severin, G. Le genre *Retinia*, Pyrale des pommes, des bourgeons, de la résine. In Bul. Soc. Cent. Forest. Belg., t. 8, p. 598-605, 674-685, 2 pls., 7 figs.
Monographic account of the three most important injurious species of the genus *Evetria* in Europe, with text figure and colored plate of *Evetria buoliana*. It should be noted that the larva figured under and credited to *Evetria buoliana* belongs to *Evetria resinella*, figured on the next colored plate, and vice versa.
1912. Gillanders, A. D. Forest Entomology, ed. 2. Edinburgh and London.
Condensed handbook information.
1913. Nüsslin, O. Leitfaden der Forstinsektenkunde, 2. neubearb. und verm. Aufl., p. 417-418, figs. 350, 352. Berlin.
Condensed handbook information on *Grapholitha (Evetria) buoliana* Schiff.
1914. Busck, August. A destructive pine-moth introduced from Europe (*Evetria buoliana* Schiffmiller). In Jour. Econ. Ent., v. 7, no. 4, p. 340-341, pl. IX, August.
First notice of the pest in America.

ADDITIONAL COPIES

OF THIS PUBLICATION MAY BE PROCURED FROM
THE SUPERINTENDENT OF DOCUMENTS
GOVERNMENT PRINTING OFFICE
WASHINGTON, D. C.

AT

5 CENTS PER COPY

▽

UNITED STATES DEPARTMENT OF AGRICULTURE

DEPARTMENT BULLETIN No. 1490

Washington, D. C.

July, 1927

DEFECTS IN TIMBER CAUSED BY INSECTS

By THOMAS E. SNYDER, *Entomologist, Forest Insect Investigations, Bureau of Entomology*

CONTENTS

	Page		Page
Introduction-----	1	Defects classed as powder post—Con.	
Types of insect defects; definitions—	4	Powder post caused by Ptinidae	
Pinholes-----	6	and Anobiidae-----	32
Grub holes-----	6	Powder post caused by Bostrich-	
Powder post-----	7	idae-----	32
Defects classed as pinholes-----	7	Powder post caused by round-	
Pinhole defects formed in living		headed borers (Cerambycidae)---	33
trees; a nonpreventable loss---	10	Powder post caused by flat-	
Pinholes in living trees		headed borers (Buprestidae)---	34
caused by ambrosia beetles---	10	Other types of defects-----	36
Pinholes in living trees		Defects classed as ring distor-	
caused by timber worms---	14	tions-----	36
Pinhole defects formed in wood		Ring distortions caused by	
after the trees are felled; a		defoliation by the bud	
preventable loss-----	15	worm-----	36
Pinholes in felled trees		Defects classed as pitch pockets	
caused by ambrosia beetles---	15	and pitchy timber-----	37
Pinholes in felled trees		Defects classed as gum spots or	
caused by timber worms---	16	streaks-----	38
Pinhole injury to imported logs---	22	Defects classed as black check-	
Defects classed as grub holes or		Defects classed as bluing or	
wormholes-----	23	staining-----	40
Grub-hole injury to living trees;		Defects classed as pith flecks---	43
a nonpreventable loss-----	25	Pith-fleck injury caused by	
Grub-hole injury to green saw		the larvae of flies-----	43
logs and lumber; a prevent-		Pith-fleck injury caused by	
able loss-----	26	the feeding of adult wee-	
Defects classed as powder post-----	27	vils-----	43
Powder post caused by Lyctus		Summary-----	43
beetles-----	28	Literature cited-----	44

INTRODUCTION

It is extremely difficult to estimate the losses due to defects in timber caused by insects. Where the grade of lumber is lowered, the loss may be ascertained by the reduction in the grade, because of the number or size of the holes caused by insects. However, where entire large oak or spruce trees are left to rot in the forest,

because the wood has pinholes in it and hence is not suitable for high-grade cooperage or airplane stock, the loss is greater, but perhaps more intangible. Where such timber is accessible and there are suitable markets, it need not be an entire loss, but could be used for lower-grade products. Closer utilization is of prime importance in the prevention of waste in the conservation of our forest resources.

Better methods of manufacture and the use of forest products, and the elimination of waste, as advocated by the Federal Forest Products Laboratory at Madison, Wis., and the Bureau of Standards of the Department of Commerce, will go far in helping to relieve the great timber shortage, which is especially serious in the case of the hardwoods.

The purpose of this bulletin is briefly to describe and illustrate, from the viewpoint of the entomologist, for the benefit of graders, inspectors, manufacturers, or utilizers of timber products, the principal types of defects in timber caused by insects, the causes of these defects, and, where possible, the mode of applying recommended methods to prevent the damage and loss. The Forest Service has already published a circular on grading lumber (39).¹

Wood-boring insects not only destroy a considerable quantity of forest products, but also cause the loss of the labor expended during the process of their manufacture. The trees from which these products were cut are a loss, and additional trees must be taken from the forest to replace them. To this loss must be added percentages of the cost and upkeep of lumber camps, machinery, equipment, logging railroads, wages and keep of men and animals in the woods, storage in the mill pond, sawing, drying, finishing, and piling at the mill.

The direct money loss caused by insects to cut timber and lumber assumes an enormous aggregate—greater proportionately than that caused by insects to living timber. To the money loss of production costs must be added the loss of time necessary to properly season the wood.

Of course, damage to seasoned finished wood products causes relatively greater loss than does damage to crude forest products. Where the products are damaged after being put in place, the cost of replacement involves additional loss of labor and time, as well as the cost of the original and replaced products, a loss far greater than the value of the raw products. Often such replacement charges should be charged to both wood-destroying fungi and wood-boring insects and not to one agency alone, as frequently there is a close relationship between these forms of life in the destruction of timber. However, in many cases wood-destroying fungi alone are responsible for the destruction.

It has been demonstrated in practice that a large percentage of the \$45,000,000 annual loss (45) caused by wood-boring insects in the past can be prevented in the future. The increased prices of lumber and all other forest products make it even more essential that all avoidable waste caused by insect defects should be eliminated in the interest of conservation.

¹ Reference is made by italic numbers in parentheses to "Literature cited," page 44.

In order to accomplish this saving it is necessary for manufacturers of wood products to utilize all available information that has been obtained from experiments carried on for many years by the Bureau of Entomology, especially the results of scientific research on the specific causes of the different types of insect defects and methods of preventing them. Data contained in earlier bulletins by experts of this bureau, as well as new and unpublished data, have been used freely in this bulletin. The published articles by Hopkins (23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35), Burke (6, 7), Webb (50, 51), Craighead (9, 10, 11, 12, 13, 14), Snyder (43, 44, 45, 46, 47), and St. George (41) were the source of much information. The pioneer investigations and publications of Doctor Hopkins, former forest entomologist of the Department of Agriculture, in reality form the basis for this bulletin. Doctor Hopkins's investigations have done much to prevent waste and losses due to insects. References to these and many other publications are to be found in Chamberlin (8).

Timber inspectors and graders should be able to determine from the defect in the wood whether it was caused by insects working in (1) the living tree; (2) the freshly felled, green saw log or bolt, with or without the bark on; (3) the green, unseasoned lumber; or (4) seasoned rough or finished product.

FIG. 1.—Black holes in white oak made by *Corythylus columbianus*. A, slightly enlarged; B, slightly reduced

The rejection of timber with insect defects caused in the tree, log, or green lumber before the wood is dry or seasoned is often an avoidable loss, because the insects are no longer working in the wood. This type of insect defect is analogous to "pecky cypress" caused by a fungus. Even though the defects are such that they materially affect the strength of the wood or otherwise unfit it for the special

FIG. 2.—"Calico poplar," a defect caused by an ambrosia beetle (*Corthylus columbianus*). (25)

use intended, there are still many purposes for which it can be used. The grade may be merely lowered, according to the number and size of the holes; or the defect may unfit the wood for special uses, owing to unsightly appearance, likelihood to cause leaks (as for barrel staves and heads for tight cooperage), or weakening (as for high-grade airplane stock). Much waste can be avoided by utilizing such defective timber for other uses and in lower grades than originally intended.

Damaged implement or vehicle stock and other material in which great structural strength is required can be used for less exacting purposes, the defect being plugged and painted over; while low-grade lumber with wormholes but no living worms or decay can be used as the base for veneer if the holes are not large enough to cause depressions in the face veneer. On the other hand, the acceptance of material infested by powder-post beetles may lead to serious results through a break at a critical moment.

Special reference is made in this bulletin to the protection

and conservation of both crude and finished wood products, such as lumber, bolts, cooperage stock (finished and unfinished), agricultural-implement handles, tent poles, vehicle parts, timber for shipbuilding, oars, airplane stock, and other high-grade products.

TYPES OF INSECT DEFECTS; DEFINITIONS

The principal types of defects caused by insects may be classed in three definite groups, namely, pinholes, grub holes, and powder post. (Table 1.) The terms for these defects are those generally used by the loggers in the woods and the sawyers and graders at the mill.

TABLE 1.—*Classification of the more common defects in timber caused by insects*

Type of defect	Description	How and when made	Condition of defective ¹ timber	Preventable
Pinholes	Holes with dark streak in surrounding wood: Hardwoods— Stained area 1 inch or more long	By ambrosia beetles in living tree	Wormholes, no living worms or decay.	No.
	Softwoods— Stained area less than 1 inch long	By ambrosia beetles in recently felled trees and green logs. By ambrosia beetles in sapwood of felled trees, logs, or green lumber.	do	Yes.
Grub holes	Holes alone darkly stained: Larger than one-eighth inch in diameter	By wood-boring grubs in living trees	do	No.
	Smaller than one-eighth inch in diameter	By ambrosia beetles in felled trees, green logs, or green lumber.	do	Yes.
Powder-posted	Holes unstained: Holes circular, open— Holes less than one-eighth inch in diameter	By ambrosia beetles in recently felled green logs or green lumber.	do	Yes.
	Holes oval, with powderlike boring dust or shreds (frass); usually more than one-fourth inch in diameter	By timber worms in living trees.	do	No.
	Holes unstained, filled with granular or powdery boring dust: Hardwoods—	By round-headed borers in recently felled softwoods and hardwoods.	do	Yes.
	Softwoods— Boring dust tightly packed (pellets of digested and excreted wood). Boring dust shredded, loose	By round-headed borers and powder-post beetles in green or seasoned timber.	Wormholes, no living worms or decay, or powder-posted.	Yes.
Pitch pocket		By flat-headed borers in living or green felled trees.	do	No or yes.
Black check Bluing		By round-headed borers in seasoned wood.	do	Yes.
		By various insects in living trees	Wormholes, no living worms or decay.	No.
Pith flock	Stained area over 1 inch long	By the grubs of various insects in living trees	do	No.
		By fungus following insect wounds in living trees and recently felled sawlogs.	do	No or yes.
Gum spot Ring distortions		By the maggots of flies or adult weevils in living trees	do	No.
		By the grubs of various insects in living trees	do	No.
		By debilitating larvae	do	No.

¹ Wormholes, no living worms or decay, is divided into two types: Injury which, from the lumberman's standpoint, can not be prevented, and injury which can be prevented by proper and prompt handling of felled trees. Timber with these defects can be utilized for certain purposes. (See pp. 11, 14, 17, 20, 22, 24, 25.)

Powder-posted or living wormy is a continuous injury, hence timber having this defect can not be used with safety before treatment with chemicals, kiln-drying, or steaming in a kiln. This defect can be prevented by proper handling of the stock. (See pp. 30, 31, 33, 34, 35.)

PINHOLES

Pinholes are small, round, usually open holes ranging from one one-hundredth to one-fourth of an inch in diameter; they are made either by ambrosia beetles or timber worms. Pinholes caused by ambrosia beetles are of two types: (1) Pinholes caused by adult beetles boring into the trunks of growing trees for the purpose of laying eggs and rearing their young, or by larvae, which also may burrow in the wood; (2) pinholes made by adult beetles or larvae in freshly felled green saw logs (with or without the bark on), bolts, and green or partly seasoned lumber.

In commercial grading rules for various species of timber pinholes constitute a standard defect (considering only wormholes), but are sometimes recognized as "equivalent defects," that is, equivalent

FIG. 3.—Work of ambrosia beetles in both sapwood and heartwood of southern yellow pine

to other defects, such as knots. This grade in certain hardwoods is termed "sound wormy" and is of that class of defects in timber or lumber in which the insects are no longer present and hence no further damage results to the timber.

GRUB HOLES

Grub holes or the larger wormholes are oval, circular, or irregular holes three-eighths of an inch to 1 inch in diameter, produced by adult insects (1) boring into or laying eggs in the trunks of living trees, or (2) boring into green, recently felled logs for the purpose of laying eggs or rearing their young; the young or larvae cause most of the injury to the wood, which serves as both food and shelter.

Grub holes may also be classified as standard defects.

POWDER POST

Powder post is that class of defects in which the larvae of insects reduce the wood fibers of seasoned or partially seasoned wood to a powderlike condition by boring through the wood, which is both their shelter and their food.

Powder post occurs only in the seasoned or partially seasoned sapwood or heartwood of both hardwoods and softwoods. Logs, bolts, timbers, lumber, and crude or finished products are attacked. The infested wood is always more or less filled with fine or coarse powdery or granulated boring dust and is called powder-posted. This type of injury is dangerous, since the grubs continue their destructive work in the wood and also infest other timber near by.

DEFECTS CLASSED
AS PINHOLES

The term "pinholes" undoubtedly originated with stave makers and coopers, from the fact that such holes are often plugged with small wooden pins. Pinholes are small, round holes one one-hundredth to one-fourth of an inch in diameter in both the heartwood and sapwood of hard (broad-leaved) and soft (coniferous) living trees, green, moist saw logs, bolts, green timbers, or green piled lumber. These holes are made by either ambrosia

FIG. 4.—Pinholes caused by the pine-wood stainer (*Gnathotrichus matricarius*)

FIG. 5.—Pinholes in sapwood Douglas fir made by *Xyloterus* sp.

beetles or timber worms. In the former case (where caused by ambrosia beetles) the holes are made by both the small, slender, cylindrical, adult beetles, and, after hatching within the wood, by their

FIG. 6.—Pinholes made by the spruce timber beetle (*Xyloterus bivittatus*) in red spruce, West Virginia

larvae or young. In the latter case the injury is caused entirely by the larvae or grubs (young) of beetles or the so-called timber worms.

The larvae of some types of ambrosia beetles excavate side galleries at right angles to the gallery made by the adult, which tunnels into the wood to deposit eggs; other types excavate no side galleries. Some holes are clear; others are stained black by the action of fungi, some of which the beetles cultivate for food in their

FIG. 7.—Pinholes caused by ambrosia beetles in heading and tight cooperage in South Carolina

FIG. 8.—Pinholes in green hickory lumber caused by an ambrosia beetle (*Xyleborus xylographus*)

galleries. Injury to green heartwood stock and to partly seasoned stock of such woods as hickory and cypress in many cases does not produce the stain.

These holes are always open (never filled with dust), and are either clear or black and associated with discolored streaks or stains

in the surrounding wood. In the latter case, the injury is caused by the larvae or grubs (young) of the beetles.

PINHOLE DEFECTS FORMED IN LIVING TREES; A NONPREVENTABLE LOSS

When pinhole defects occur in the living tree, it is of course impossible for the lumberman to prevent the injury, since the holes

FIG. 9.—Pinholes with short stain streaks made in green ash lumber by *Platypus compositus*

have been made periodically, and often many years before the tree was cut. The dates when the injury was done can be determined by counting the annual rings of tree growth. Usually the wood surrounding pinholes of this type is stained.

PINHOLES IN LIVING TREES CAUSED BY AMBROSIA BEETLES

Pinholes are uniform in size, one-eighth of an inch in diameter or smaller (to one twenty-fifth inch), either darkly stained or un-

stained, or with short streaks in the surrounding wood; these holes run deeply in every direction in an irregular manner through the wood, and it is difficult to determine their extent from the surface. They are termed "pinholes," "shot holes," or "black holes" (36). Such holes occur in both the sapwood and heartwood of softwoods and hardwoods, but are more common in the sapwood.

One of the commonest defects (fig. 1) is caused in living trees by an ambrosia beetle,² and the holes are known as "black holes," "spot worm," "steamboats," "grease spots" (West Virginia), and "flag worm" (Arkansas). The holes usually occur in the best part of the wood, either one in a place, or two or more in a row. This defect is so prevalent in mature white oak and other oaks that it is often difficult to find a tree the wood of which is entirely free from it; in yellow poplar the defect is not so common. When the defect does occur in whitewood or yellow poplar the accompanying long, black, greenish or bluish streaks give a calico effect, and the wood is called "calico poplar" (fig. 2) (23, 27). The same defect occurs in beech, birch, basswood, maple, and elm in West Virginia, elm and oak in Michigan, oak in Arkansas, and chestnut in Tennessee.

Calico poplar and oak might be used, when available in sufficient quantity, as a special grade for interior natural-wood finish, as the effect is pleasing. Thus the apparently limited quantity could be utilized without having its value depreciated much below that of the first and second grades.

This defect causes a serious loss to stave and shingle stock. In Arkansas from 30 to 40 per cent of white oak Bourbon whisky-barrel-stave stock and staves has often been left to rot in the woods on account of injury by the oak timber worms and ambrosia beetles.

FIG. 10.—Pinholes made by *Xyleborus* sp. in green chestnut board, after piling

² *Corthythus columbianus* Hopkins.

FIG. 11.—Pinholes in the sapwood of high-grade cypress blocks; end view of logs in freight car; wood completely riddled by ambrosia beetles (*Platypus compositus* and *Xyleborus* sp.)

FIG. 12.—Pinholes caused by ambrosia beetles (*Xyleborus* sp.) in mahogany flitches

A loss of \$5 to \$20 per thousand feet of timber is a conservative estimate. This defect is classed as "wormholes, no living worms or decay," and can not be prevented in the tree, but of course similar defects in the green stock can be prevented by proper handling.

Pinholes one twenty-fifth to one-eighth of an inch in diameter in both heartwood and sapwood of hardwoods and softwoods (figs. 1, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, and 14) are caused by similar beetles.³

The defects caused by these pinholes and stains (discolored streaks and patches) (figs. 1, 2, 3, 4, 6, 9, and 13) reduce the grade of timber and its full strength, unfitting it

for structural timber, wagons, agricultural implements, tight cooperage, and shingles. A 25 per cent loss of elm cooperage stock in

FIG. 13.—Pinholes caused by ambrosia beetles (*Xyleborus* sp.) in imported greenheart (*Nectandra rodioei*)

FIG. 14.—Pinhole damage by ambrosia beetles (*Xyleborus grenadensis*) to Hura wood (Rakuda) imported from Central America

logs at the mills in Illinois was due to such defects. In some cases millions of feet of timber have been reduced 10 to 25 per cent or more in value by pinhole defects. In deadened standing cypress in the Gulf States, pinhole injury can be prevented by girdling the trees in March, April, October, and November (28), but from August to September is apparently the most effective season.

³ Platypus, Monarthrum, Xyleborus.

PINHOLES IN LIVING TREES CAUSED BY TIMBER WORMS

Pinholes or wormholes in the heartwood of chestnut and red oak, one one-hundredth to one-fourth of an inch in diameter, open, clear, not stained, but lined with a substance about the color of the wood, a large number of holes in a given space, are made by timber worms hatching from eggs laid in or near scars. The holes made by the grubs are not of uniform size; they may extend several feet through the wood. (Fig. 15.)

This defect, due to the chestnut timber worm,⁴ is one of the most common and serious defects in chestnut timber throughout its range. It is rare to find chestnut trees, logs, or telephone or telegraph poles free from this defect, and practically every tree of merchantable

FIG. 15.—Holes caused by the chestnut timber worm (*Melittomma sericeum*), which causes the defect classed in the grade "sound wormy chestnut"

size is more or less affected. There is a very low percentage of "clear" chestnut, the remainder being "wormy" and reduced to the lower grades. This defect is a "sound wormy" injury and can not be prevented. Unfortunately the chestnut is rapidly becoming extinct as a commercial tree, owing to the chestnut blight fungus.

Wormy chestnut and oak wood can be used wherever structural strength is not necessary. They are especially suitable as the base for veneer in manufacturing pianos, caskets, coffins, automobile running boards, and the lower grades of building lumber, because of the comparatively reasonable price at which this grade of lumber can be bought. Much timber that would otherwise be wasted can be thus utilized. There is, however, a limit to the size of the holes admitted.

Similar pinholes or wormholes one one-hundredth to one-eighth of an inch in diameter in oak timber (fig. 16) are caused by the oak timber worm.⁵ This defect is especially serious in large mature white oak trees. The holes occur in large numbers in a given space and extend in all directions through the solid heartwood. This insect usually affects the wood of the finest old, mature, or over-mature trees, sometimes causing defects which result in the discarding of entire trees for such uses as tight barrel staves. This defect is classified under the term "wormholes, no living worms or decay," and can not be prevented when occurring in living trees, although losses can be lessened by clean forest management; that is, removal

⁴ *Melittomma sericeum* Harris.

⁵ *Eupsalis minuta* Drury.

of old dead and dying snags, stag-headed trees, and trees badly fire-scarred. From 20 to 25 per cent of oak lumber may be wormy from this cause. A loss of 15 to 20 per cent of the product is considered low.

Pinholes or wormholes one-eighth to three-sixteenths of an inch in diameter, few to many grouped in a given space, accompanied by a staining of the wood, or all more or less connected by irregular blackened streaks, in oak, maple, tupelo gum, beech, and other hardwoods, are similar to those caused by the oak timber worm. This defect is caused by tenebrionid timber worms.⁶ The insect lays its eggs in the living tree near a scar or wound, and many larvæ working together excavate irregular cavities and longitudinal burrows sometimes a foot in length. This defect is fairly common throughout Pennsylvania, Maryland, and Virginia and is classed as "wormholes, no living worms or decay." Although the losses can be greatly lessened by clean forest management, they are not preventable from the lumberman's standpoint.

FIG. 16.—Pinholes caused by the oak timber worm (*Eupsalis minuta*), which causes the defect classed in the grade "sound wormy oak"

PINHOLE DEFECTS FORMED IN WOOD AFTER THE TREES ARE FELLED; A PREVENTABLE LOSS

Pinholes in felled timber are similar to those caused in living trees by ambrosia beetles and timber worms. Such holes are less than one-eighth of an inch in diameter and open, either darkly stained or unstained, or with dark streaks in the surrounding wood.

PINHOLES IN FELLED TREE⁷ CAUSED BY AMBROSIA BEETLES

In sapwood or heartwood of white oak, chestnut, yellow poplar,⁷ birch, beech, and sweet gum the pinholes caused by ambrosia beetles are about one-twelfth of an inch in diameter, open, usually isolated, and always accompanied by long, discolored streaks in the surrounding wood. Lumber with such defects is rejected for the higher grades.

⁶ *Strongyllum* spp.

⁷ The adult of *Corthylus columbianus* Hopkins causes pinholes in hardwoods with which discolored streaks are associated.

In Arkansas a 2 per cent loss of the product of green or newly manufactured white-oak Bourbon whisky-barrel staves occurred at one operation.

Pinholes one twenty-fifth to one-eighth of an inch in diameter occur in the sapwood of both hardwoods and softwoods (figs. 3, 4, 5, 7, 8, 9, 10, 12, 13, 14); in hardwoods usually the wood is not stained; but in freshly cut green hardwoods the wood may be stained. A few to a large number of holes occur in a given space; often every

FIG. 17.—Pinholes in yellow birch caused by the sapwood timber worm (*Hylcoctus lugubris*). (Drake)

square inch of wood is penetrated. These common pinhole defects in recently felled green logs or bolts (with or without the bark) and closely piled green timber and lumber are caused by ambrosia beetles⁸ (figs. 4 and 5). There are two types of galleries made by these beetles—one, a long, winding main gallery and usually no staining of the wood, and the other, a series of short side galleries at right angles to this main gallery (figs. 1 and 6), the gallery usually being accompanied by a staining of the wood. Unlike holes made by the timber worms, which

increase with the growth of the worms, these holes are more or less uniform in size.

PINHOLES IN FELLED TREES CAUSED BY TIMBER WORMS

Pinholes caused by timber worms in the heartwood of chestnut⁹ and oak¹⁰ can be recognized as follows: Holes from one one-hundredth to one-fourth of an inch in diameter; open and not stained; a large number to a given space.

Such holes may be a serious defect in square timbers used in structures the woodwork of which is exposed to the weather. Care should be exercised in utilizing wormy oak in structural work where strength is required, or in vehicle, ladder, or implement stock. It should be

⁸ *Gnathotrichus* and *Xyloterus*.

⁹ The larvae of *Melittomma sericeum* Harris cause wormy chestnut and oak; the eggs are laid under bark.

¹⁰ The larvae of *Eupsalis minuta* Drury cause wormy oak; the eggs are laid under bark.

rejected for barrel staves and heads of tight cooperage, unless the holes are few and can be plugged. Such stock can be used where the holes can be plugged, puttied, and painted over, or as a base for veneer.

Pinholes or wormholes one one-hundredth to three-sixteenths of an inch in diameter in green saw logs with the bark on, of basswood, buckeye, chestnut, black walnut, cottonwood, yellow poplar, and birch (fig. 17), are caused by the sapwood timber worm,¹¹ which may cause a 5 to 10 per cent loss of a log. This defect is described as "wormholes, no living worms or decay," and can be prevented.

The insect lays its eggs in dying trees and green saw logs which are allowed to lie in the woods with the bark on from April to July, in the States north of the Gulf. The eggs are deposited in crevices in the bark. The beetle will not lay its eggs in barked logs or logs which are floating in log ponds.

In the sapwood or heartwood of oak, maple, tupelo gum, and beech, grouped holes from one-eighth to three-sixteenths of an inch in diameter and all more or less connected by irregular blackened streaks are caused by tenebrionid timber worms.¹²

FIG. 18.—Holes caused by termites (*Coptotermes niger*) in log imported from Honduras; damage to the living tree caused this defect in the board

¹¹ *Hylecoetus lugubris* Say.

¹² The larvae of *Strongylus* spp. cause wormholes in hardwoods.

Pinholes of these and all other kinds made in timber after felling are preventable by prompt handling and rapid utilization of felled timber. Green logs or bolts, either with or without the bark on, should be placed in the mill pond as soon as possible after being cut, especially during the warm months of spring, summer, or fall, or in damp weather. In the mill pond the logs should be floated in loose booms, so that they can be rolled over and turned to make sure that they are periodically submerged on all sides. The damage can also

FIG. 19.—Grub holes and burrow caused by carpenter bee (*Xylocopa* sp.) in juniper from Arizona

be prevented by sawing the logs into timber and lumber as soon as possible.

If it is not possible to get the logs into the mill pond, they can be sun cured or dried rapidly (10, 11, 15, 16, 17, 18, 19, 20) to bring about conditions unfavorable to insect attack, which is accomplished by peeling them and placing them on "browse" (limbs) on knolls in open places in the sun. Other preventive measures consist of girdling before cutting, peeling, cutting, and leaving the tops on for a time before logging or after cutting. Barking the logs will prevent

FIG. 20.—A, enlarged holes made by carpenter ants (*Camponotus* sp.) in red cedar in Washington; B, redwood damaged by carpenter ants in California

FIG. 21.—Work of a horntail (*Urocerus albicornis*) in a decaying softwood log

damage by timber worms, as will also coating or spraying with preservatives.

Repellent, poisonous, or sticky chemical sprays, to prevent attack by ambrosia beetles on more valuable species and products, can be used where the bark is on the logs and the wood will not be discolored (12). The preservative coating known as hardened gloss oil has the advantage of preventing checking.

Green lumber should be kiln-dried (49); or, where this is not practicable, piled on stickers (1), to insure rapid air seasoning. Air

FIG. 22.—Grub holes made by the oak carpenter worm (*Prionoxystus robiniae*)

seasoning of green freshly sawn timber or lumber by open or loose piling is a preventive. In cribbing or proper piling of green sawn stock to facilitate rapid drying, care should be taken to insure against severe checking from too-rapid drying. Heavy dimension timber should be stacked in loose piles. If it were possible to kiln-dry the material, insect damage, of course, would be prevented.

In the case of damage to stave stock, by special methods of sawing the waste can be reduced to a minimum and a very considerable saving of material effected. Where the holes are not too numerous, they can be plugged with wooden pins and the wood used for casks.

FIG. 23.—A, grub holes in overcup oak, made by the Parandra borer (*Parandra brunnea*) near Vicksburg, Miss.; B, grub holes in soft maple made by the Parandra borer in Maryland

The defect does not deteriorate furniture and inside-finish stock. This defect is considered under "wormholes, no living worms or decay," and in the interest of conservation and closer utilization, timber with these defects should be used.

FIG. 24.—A, grub holes made by a flat-headed borer (*Buprestis apicans*) in the pitchy "fatwood" of the basal log of long-leaf pine; B, the same in a turpentine tree

PINHOLE INJURY TO IMPORTED LOGS

Large quantities of tropical woods in the rough, round, or squared log are yearly shipped into the United States from Central and South America, the West Indies, Africa, and the Philippines. (Figs. 12, 13, and 14.) Almost invariably such timber, if received within a year after it is cut, contains many species of pinhole-boring beetles, but these never live over the winter, except perhaps in the Gulf States.

Under present methods of lumbering, such imported logs usually are infested by pinhole-boring beetles of many species, which attack the logs after they are cut, and continue to live in them for several months and in some cases a year or more. If such logs are shipped into another country with a similar climate, the insects may survive and attack logs of native timber; if, on the other hand, logs are exported to a much colder or hotter country, the insects will rarely, if ever, become established in the country of import.

There is, therefore, some danger of introducing destructive species through the commercial interchange of timber in the form of logs.

FIG. 25.—Wormholes caused by "sawyers" (*Monochamus* spp.) in pine. A and B, wormholes in southern yellow pine; C, wormholes in white pine

For example, many insect species are widely distributed over tropical America that do not occur in tropical Africa, Asia, Australasia, and the Pacific islands. Tropical Australia, the Philippine Islands, and tropical Japan doubtless have a considerable number of the same species of pinhole borers. One country may have a few that do not occur in the others.

DEFECTS CLASSED AS GRUB HOLES OR WORMHOLES

Grub holes are medium to large, circular, oval, or irregular holes from three-eighths to 1 inch in diameter, in both sapwood and heart-

wood of all kinds of timber. They may either be stained black inside or be of the same color as the surrounding wood, and they may be free and open or filled with tightly-lodged boring dust, depending on the kind of insect making them. This boring dust does not fall out when the wood is jarred. Grub holes are made in the living tree, in the saw log, or in piled green lumber. In nearly all cases the injury is caused by the young borers, sawyers, or grubs, but occasionally also by adult termites or white ants (fig. 18), adult

FIG. 26.—Wormhole defects in cedar, caused by the round-headed borer *Callidium antennatum*

carpenter bees¹³ (fig. 19), or carpenter ants¹⁴ (fig. 20), or by the larvae of horntails¹⁵ (fig. 21). Usually this type of injury is considered as "wormholes, no living worms or decay," especially if the holes are stained black, and no further damage will result, except in rare cases.

Grub holes constitute a standard defect and are also included under "equivalent defects." They are often of considerable size

¹³ *Xylocopa* spp.

¹⁴ *Camponotus* spp.

¹⁵ *Tremex columba* L., or *Sirex* spp.; they usually damage only dead wood or trees or logs that have been left in the woods too long.

and consequently reduce the structural strength of the wood, especially when cut into smaller dimensions; therefore timber having such borer holes should be used in as large dimensions as possible.

FIG. 27.—Western red cedar shingles badly damaged by grub holes and burrows of flat-headed borers (*Trachykele blondeti*). (1). *a*, Quarter-sawn shingle showing both cross and longitudinal sections, *c*, *d*, of the larval mines; *b*, bastard-sawn shingle showing larval mines

GRUB-HOLE INJURY TO LIVING TREES; A NONPREVENTABLE LOSS

Grub-hole defects in living trees can not be prevented by lumbermen, since often these defects occurred many years before the tree was cut. Sometimes the insect holes are partially healed over by

new growth of the tree, or they may have been enlarged by carpenter ants. This type of injury can be recognized as follows:

(1) In the sapwood and heartwood of hardwoods very large circular holes, one-half to 1 inch in diameter, are open and but slightly stained and are usually lined with a silky yellowish-brown web.¹⁶ (Fig. 22.)

(2) In sapwood or heartwood of hardwood trees the holes are darkly stained and are open, containing little or no sawdust or frass.¹⁷

FIG. 28.—Powder-posted white ash shipbuilding lumber showing larval burrows and exit holes of adults of *Lyctus planicollis*; board from a closely piled stack of lumber throughout which larvae had burrowed

(3) In the heartwood of butt logs of gum and a few other hardwoods, unstained holes contain mined granular and fibrous frass¹⁸ (fig. 23).

(4) In the heartwood of butt logs of softwoods, chiefly pine, very irregular holes are narrowly oval, without stain, but surrounded by much pitchy "lightwood," or "fatwood," and are very tightly packed with fine granular frass¹⁹ (fig. 24) (?).

(5) Pinholes or wormholes in softwood logs and lumber in the Northern States are caused by a melandryid timber worm.²⁰ These larvae

gain entrance to the living sapwood through scars or blazes and also attack recently felled timber. The holes are filled with fine sawdust.

GRUB-HOLE INJURY TO GREEN SAW LOGS AND LUMBER; A PREVENTABLE LOSS

All types of borer holes other than those mentioned above are made after the trees have been felled, and are preventable by proper methods of handling the logs. These borer holes occur in the sap-

¹⁶ This injury is found principally in oak, chestnut, locust, and cherry and is caused by the carpenter worm, *Prionoxystus robiniae* Peck.

¹⁷ These holes are found principally in oaks and are caused by the round-headed borers (Cerambycidae) *Gocis* spp. and *Romuleum* sp.; in hickory, they are caused by *Gocis* sp.; in hickory in the South, small darkly stained holes are caused by a wood-boring caterpillar, *Cossula magnifica* Bailey; and in hard maple they are caused by the maple-tree borer, *Glycobius speciosus* Say.

¹⁸ This injury is caused by *Parandra brunnea* Fab.

¹⁹ This injury is caused by *Buprestis apicans* Hbst. and occurs especially in fire-scarred long-leaf pines and trees boxed for turpentine. This borer causes an excess of pitchy wood near the injury. The injury often amounts to the reduction of 5 to 10 per cent of the lumber to lower grades and the wind-throw of much second growth on turpentine operations.

²⁰ *Serropalpus barbatus* Schall.

wood of hardwoods or softwoods as unstained irregular holes from one-fourth to 1 inch in diameter, which may be open, loosely or tightly filled with powder, with granular or fibrous frass, or with pellets.²¹

These and similar types of borer injury can be prevented by prompt handling of the logs after they are felled. Logs, bolts, and sawn or squared timbers should never be allowed to lie where cut in the woods, after the 1st of February in the Gulf States, or after the 1st of April farther north. Other preventive measures include: Rapid utilization; submergence of logs in water, where they will not be attacked, and working them up as soon as removed from the water; sun-curing, with or without the bark on (care should be taken to provide against excessive checking); and removal of bark strips from freshly sawn material. The damage can often be prevented by peeling the bark (both outer and inner) from the logs or bolts, timbers, or edges of lumber, as the bark offers a favorable place under which the insects can lay eggs. Both the outer and inner bark of sawn timber should always be carefully removed and the timbers placed where they will season rapidly (1).

FIG. 29.—Powder-posted sapwood oak veneer laid on a core of chestnut (door stock); work of *Lyctus planicollis*. Note that the heartwood oak and the chestnut have not been attacked

DEFECTS CLASSED AS POWDER POST

Powder post is indicated by holes from one-sixteenth to one-fourth of an inch in diameter, in the surface of the wood, in the sapwood and

²¹ The principal injury of this character found in pines, spruces, and firs is caused by the pine sawyers, *Monochamus* spp. (fig. 25) (50); the loss due to pine sawyers in green logs and storm-felled timber is often as high as 35 per cent. In ash the defects are caused by the banded and red-headed ash borers, *Neoclytus caprea* Say and *erythrocephalus* Fabricius; in hickory by the banded hickory borer, *Cyllene pictus* Drury, and the red-headed ash borer; in locust by the locust borer, *Cyllene robiniae* Forster; in cedar by the round-headed borer *Callidium antennatum* Newm. (fig. 26); in cypress, western redwoods, and cedars by flat-headed borers, *Trachykele* spp. Burrows made by *Trachykele* are tightly packed with pellets of excrement, and shingle stock is full of holes (fig. 27) (7).

heartwood of both hardwoods and softwoods, from which the powder will fall when moved or jarred.²² The interior is honeycombed by irregular burrows made by the larvae and when badly damaged is converted into a mass of closely packed material, which readily crumbles into fine flourlike powder or coarser pellets of excreted wood. This is held together by an outer thin shell and intervening

fibers of sound wood. These defects will be discussed in the order of the size of the holes caused by the various types of insects. All powder-post damage can be prevented.

FIG. 30.—Powder-post defect in pine made by *Xyletinus peltatus*

POWDER POST CAUSED BY LYCTUS BEETLES

The injury caused by Lyctus beetles is confined to the white-wood or sapwood of hardwoods (34). It consists of small holes one-sixteenth to one-twelfth of an inch in diameter, with irregular burrows filled with flourlike powder. Air-dry or kiln-dry sapwood material, and sapwood which has been stored or piled in one place for two, three, or more years, especially second-growth ash, hickory, and oak, are principally

affected; but other hardwoods, such as walnut, maple, persimmon, cherry, elm, poplar, and sycamore, are also damaged.

Seasoned shipbuilding and airplane lumber and gunstock blanks, stored in large quantities, and finished stores, such as wheelbarrows, tent poles, oars, airplane parts, shovel and pick handles, and many other hardwood articles used in the military services are subject to serious damage by powder-post beetles. (Fig. 28.)

Hickory, ash, and oak furniture, interior woodwork of buildings (fig. 29), and the woodwork of farming machinery and implement handles; ladder stock, such as rungs; vehicle stock, such as hubs, spokes, felloes, rims, singletrees, poles, and shafts; and cooperage stock (barrel-stave bolts) are also injured.

²² Insects which have this peculiar habit of reducing wood fiber to a powderlike condition belong chiefly to the families Lyctidae, Ptinidae, Anobiidae, Bostrichidae, and Cerambycidae. By far the larger part of the injury is caused by species of the genera Lyctus and Neoclytus.

The loss to seasoned hardwood products ranges from 10 to 50 per cent, sometimes representing a loss of thousands of dollars to a single manufacturer or dealer who neglects to adopt the proper preventive measures. The affected articles are not only reduced in value, but

FIG. 31.—Powder-posted seasoned hickory stock in the rough damaged by *Xylobius basilaris*. A, end of bolt; B, end of section of bolt; C, planed section of damaged bolt

frequently are rendered worthless for the purposes for which they are intended. In the aggregate the direct financial loss that has been caused by these beetles in this country has amounted to hundreds of thousands of dollars. The loss increases with the length of time the infested stock is held in storage; the wood may be reinfested by many

generations over a period of 20 years or more. In certain cases powder-post injury may be a menace to human life, as in the weakened woodwork of buildings, vehicle stock, or ladders.

Losses due to *Lyctus* beetles can be prevented by proper methods of classification and piling of stock by kinds; by keeping heartwood

FIG. 32.—A, Duali plywood, imported from the Philippine Islands, powder-posted by *Bostrychopsis parallela*; B, bostrychid powder-post injury to myrtle in Oregon

and sapwood stock separate; by periodical inspection, utilizing the older stock first (34); and by using only heartwood piling sticks; or by submergence in water for four months, which renders the wood immune from attack, even after removal from the water.

In the case of finished wood products, it may often be practicable to treat the wood with substances to prevent attack. Creosotes are

effective preventives, but they stain the wood; hence, where they can not be used, in the light of the discovery of the place and manner of the laying of the eggs in the pores of the wood, any substances that will close the pores will prevent oviposition in wood not previously infested. In wood from which beetles have emerged, however, eggs might be laid within the exit holes. Paraffin wax, varnish, linseed oil, or other fillers effectively close the pores of wood. A certain varnish known as hardened gloss oil is commonly used. Wood that has been seasoned less than 8 to 10 months will not be attacked by *Lyctus* beetles; therefore, in applying chemical preventives, only sapwood that has been seasoned for 8 to 10 months and longer need be treated. The seasonal history of these beetles indicates that preventives should be applied before March 1.

The great and recurring expense of treating infested wood can be avoided by prevention of attack by proper methods of management. Since only the sapwood or whitewood is attacked by *Lyctus* powder-post beetles, it is recommended that more heartwood be used to replace sapwood. Although the demand of the trade is for white-wood handles, etc., the prejudice against heartwood is not warranted

FIG. 33.—Holes made by the banded ash borer (*Neoclytus caprea*), one of the powder-post beetles

FIG. 34.—Powder-posted ash shipbuilding lumber, showing coarse powdery frass in the burrows made by *Neoclytus caprea*

and should be overcome by educational propaganda. Weight for weight, red or heartwood hickory is as strong as white or sapwood. Closer utilization can be effected by larger use of heartwood. Some manufacturers paint their stock, to overcome this trade prejudice.

Effective remedies include kiln-drying at high temperatures (180° F. and over); steaming at 130° F. (this will not insure against future attack) (46, 47); and treatment with orthodichlorobenzene or a mixture of kerosene and coal-tar creosote, after which the material should be kept in quarantine a sufficient length of time to deter-

mine whether a second treatment is required.

Partially damaged material which is too valuable to be destroyed should be salvaged, when practicable, by trimming off and burning the sap edges and other damaged and infested parts.

POWDER POST CAUSED BY PTINIDAE AND ANOBIIDAE

This damage consists of small holes one-sixteenth to one-eighth of an inch in diameter and irregular burrows in the wood of both softwoods and hardwoods (fig. 30). The damage is similar to that caused by *Lyctus* beetles, except that softwoods are also attacked and the defect occurs²³ in the heartwood. Most injury by Ptinidae is caused to seasoned wood, or logs that have been left lying in the woods too long.

FIG. 35.—Powder-posted southern yellow pine boards damaged by *Hylotrupes bajulus*

These insects²⁴ attack freshly felled logs with the bark on. The eggs are laid within the log near holes made by the adult beetles. Submerging the logs in the mill pond and prompt utilization will prevent much loss. In the case of vehicle, handle, and similar

POWDER POST CAUSED BY BOSTRICHIDAE

This damage consists of circular holes one-eighth to three-eighths of an inch in diameter and irregular longitudinal burrows filled with frass, or with coarser dust in the sapwood and heartwood of hardwoods, which does not fall out so readily. (Figs. 31 and 32.)

These insects²⁴ attack freshly felled logs with the bark on. The eggs are laid within the log near holes made by the adult beetles.

Submerging the logs in the mill pond and prompt utilization will prevent much loss. In the case of vehicle, handle, and similar

²³ This defect is caused by *Xyletinus peltatus* Harris, which attacks both hardwoods and softwoods.

²⁴ Bostrichidae: *Scobicia*, *Xylobiops*, etc.

stock, all the bark and edgings should be removed; treatment with shellac or wax will also prevent the adult beetles from boring into the wood to lay eggs.

POWDER POST CAUSED BY ROUND-HEADED BORERS (CERAMBYCIDAE)

Holes about one-eighth of an inch in diameter, tightly packed with finer frass, in the sapwood or heartwood of oak and hickory. are made by the flat powder-post beetle.²⁵ It attacks both freshly cut and seasoned timbers and continues to work for a number of years. Damage can be prevented by prompt handling of the logs, removal of the bark, and disposal of infested stock.

Oval holes about one-fourth of an inch in diameter or irregular burrows tightly packed with coarse, powdery frass, in the heartwood and sapwood of ash, are caused by a round-headed borer.²⁶ (Figs. 33 and 34.) This insect attacks only freshly cut timbers, but when infested logs are stored the borers continue to work for several years (51).

Prompt utilization, submerging the logs in the mill pond, rapid seasoning, or removal of the bark will prevent this defect. Timber should be felled in the late fall or winter, so that the bark may dry somewhat and be less attractive to the beetles when they are flying and depositing their eggs early in the spring. In the Gulf States logs should not be allowed to lie in the woods at any time for more than two to three weeks, or after the 1st of April farther north, nor should they be

FIG. 36.—Ring distortions in balsam caused by the spruce bud worm. Base of tree attacked in 1911 showed accelerated growth for two years, followed by retardation and incomplete ring in 1918 (X), and rapid recovery later. (48)

²⁵ *Smolicum cucujiforme* Say.

²⁶ *Neoclytus caprea* Say.

stored in closely packed piles for long periods. After the 1st of February logs should be cut up into sizes as small as commercially practicable and seasoned as promptly as possible; or sun-cured; or placed in the mill pond. Narrow strips of bark left on the edges of boards and planks or timber cut from green logs in February and March serve to attract the insects to such places to lay their eggs; therefore the logs should be trimmed off and the trimmings burned.

Irregular oval holes from one-fourth to one-half inch in diameter, filled with a mixture of coarse granular and fine powdery frass, are caused by the "old house borer."²⁷ (Fig. 35.)

FIG. 37.—Pitch pockets in western yellow pine, caused by unsuccessful attack, years ago, by the Black Hills beetle (*Dendroctonus ponderosae*). A, pitch pockets in sawn board; B, end of log showing pitch pockets

It attacks only well-seasoned wood which has been stored for more than a year. This borer continues to work for years, until the wood is thoroughly perforated.

All bark edges should be removed from lumber and timber; the stock should be inspected frequently for evidences of falling dust, which indicates the presence of these borers, and all such wood should be promptly burned; or the stock may be dipped in orthodichlorobenzene, if only slightly injured.

POWDER POST CAUSED BY FLAT-HEADED BORERS (BUPRESTIDAE)

In bald cypress in the Eastern States and in western redwoods and cedars, powder-post injury is caused by flat-headed borers.²⁸

²⁷ *Hylotrupes bajulus* L.

²⁸ *Trachykele* spp.

The burrows are tightly packed with pellets of excrement, and shingle stock is full of holes. (Fig. 27.)

The injury occurs in living, dying, or dead trees, and consists of a flattened, oval, gradually enlarging, more or less tortuously winding mine or wormhole, which, when completed, widens out into an elongate-oval pupal cell. This cell connects with the outer surface by a short, oval exit hole. The mine has its surface marked by fine transverse, crescentic lines, and is usually tightly packed with sawdustlike borings and pellets of woody excrement.

To prevent such injury, the forest should be kept clear of dead and dying trees and of felled trees which afford ideal breeding spots. Such trees might be used for fuel, or they could be piled with the limbs and tops and burned. If trees must be deadened in the lumbering operations, the "deadening" should be done at a time of the year when the sap is not actively flowing. October, November, and December would probably be the best months for this. If the timber must be felled and left in the woods for a time, the felling should be done during the same months, and the logs should be barked and left so that they will dry quickly and thus become distasteful to these borers.

If the timber is found to be newly infested while standing, or on felling, the most practical remedy is to cut it into logs at once and place the logs in a pond or stream so that the larvae will be destroyed and further damage prevented. If the damage has been done before the lumberman has noticed the injury, which is usually the case, much loss can often be prevented by utilizing the damaged stock to the best advantage. It may be used for poles, posts, plank-ing, sills, small construction timbers, or where the wormholes are not particularly detrimental; it should not be used for cooperage, shipbuilding, shingles, doors, finishing, cabinetmaking, or furniture, in which clear stock is desired: otherwise the loss is apt to be severe, both because of the poor quality of the product and because of the extra labor necessary to produce it (7).

FIG. 38.—Longitudinal section of yellow pine sapling, showing damage to wood from attack of *Pinipestis zimmermanni*. (5)

OTHER TYPES OF DEFECTS

Other common defects—ring distortions, pitch pockets, gum spots, black check, staining, and pith flecks—are often caused by insects or associated with insect attack. Such defects, however, are not always necessarily caused by insects.

DEFECTS CLASSED AS RING DISTORTIONS

Characteristic distortions and abnormalities in the annual rings of trees result from defoliation by leaf-feeding insects, as well as from other causes, as pointed out by Hartig, Harper, and Craighead (48). Destruction of the leaves at certain periods causes a reduction in the normal amount of food manufactured by the tree, with consequent loss in the amount of wood laid on and the formation of incomplete annual layers or double rings (false rings). Many species of trees are subject to defoliation, though little is known of the resulting effects except in a few cases.

Larch, fir, spruce, jack pine, lodgepole pine, western yellow and white pines, hemlock, oak, hickory, catalpa, birch, and cherry are some of the trees subject to frequent or periodic defoliation.

FIG. 29.—A and B, two views showing small "pitch pocket" or "bird's eye," caused by larva of a fly in western yellow pine

RING DISTORTIONS CAUSED BY DEFOLIATION BY THE BUD WORM

Defoliation by the bud worm²⁹ on spruce and fir produces the first year a decided decrease in wood on the upper stem, while an abnormally larger ring is laid on at the base. In subsequent years a gradual reduction occurs, the narrowest ring being laid on some four or five years later throughout the tree. Gradual increase then

²⁹ *Cacoecia (Harmoloba) fumiferana* Clemens.

follows in recovering trees. In fir trees one, two, or three rings may fail to form completely around the stem, particularly at the base, and are represented only by partial arcs (fig. 36) (48).

Rings of traumatic resin ducts are frequently deposited in the terminal portions of defoliated fir trees during the first to third years of feeding.

Complete defoliation of certain hardwoods in the early spring produces a double ring the same year. These defects result in a great loss of increment, increasing the rotation from 3 to 10 years, and cause confusion in growth studies based on the annual rings.

Similar distortions of rings are found in Douglas fir, due to defoliation by the spruce bud worm, and in western fir, spruce, yellow pine, and particularly lodgepole pine, due to defoliation by the needle miner.³⁰

Such losses and defects can be prevented only by the prevention of defoliation, a problem of forest management.

DEFECTS CLASSED AS PITCH POCKETS AND PITCHY TIMBER

Pitch pockets are openings between the grain of the wood which contain more or less pitch or bark. They are graded as small, standard, and large pitch pockets. These pockets range from one-eighth inch to 2 inches in width and from 3 to 12 inches in length. Sometimes, instead of pockets, there are merely pitch streaks. These defects may be caused by insects as well as other agents.

The pockets are small, usually about one-half to 1 inch in length, and about one-half inch in width, full of pitch; in pine, they are parallel to the grain. Sometimes they are as large as 2 inches in diameter and contain a large quantity of pitch. This injury may be caused by unsuccessful attacks of bark beetles³¹ which have failed to kill the tree and have been drowned out by the flow of pitch. (Fig. 37.) It is a sound defect which can be prevented only by controlling the bark beetles that attack the living trees.

Excessive pitchy streaks in yellow and other pine lumber are sometimes caused by the larvae of a moth³² working in the cambium (fig. 38), especially in mature trees.

FIG. 40.—Gum spot in Douglas fir, caused by a flat-headed borer (*Melanophila drummondi*)

³⁰ *Recurvaria milleri* Busck.

³¹ *Dendroctonus* spp. The term "pitch pocket" is here used to define what is apparently the tree's further means of defense from attack by bark beetles, after the pitch running out of the entrance hole, forming a pitch tube, has failed to drown out the beetles.

³² *Pinipestis zimmermanni* Grote, the Zimmerman pine moth.

The wood of trees that have been infested by larvae of the moth is invariably so permeated with pitch that the lumber cut from such logs is either materially reduced in value or is rendered wholly unfit for commercial use. This moth is especially abundant; in south-eastern Montana a large percentage of the trees are pitch-soaked. For this reason the lumber is utilized only as rough lumber for sheds and similar structures where shrinkage can be discounted (5).

Serious damage by this moth can be lessened by clean forest management.

"Pitch seams," "gum check," or whatever these defects may be termed locally, have always been recognized as a serious depreciating factor in the utilization of Douglas fir (4).

FIG. 41.—Black check in western hemlock, radial section. (6)

The loss is occasioned by the work of the Douglas fir "pitch moth,"³³ which causes the difference in price between absolutely clear lumber and the lower grades or dimension stuff. These insects work in the portions of the trunks of living trees which later clear themselves of branches; hence only those logs are affected which, were it not for previous infestation by the moths, would yield the better grades of lumber.

Sawyers of Douglas fir estimate a general loss in the entire Douglas fir product of between 7.5 and 15 per cent due to this defect of pitch seams in the logs used (4). The depreciation is lowest in the Rocky Mountain region and heaviest toward the Pacific coast. Such pitch pockets are not always caused by insects.

Another pitch moth, the Sequoia pitch moth,³⁴ greatly retards the growth of lodgepole pine in western Montana (3).

Under present forestry conditions in this country it is impracticable to control these insects over large areas.

Similar defects due to the working of the larvae of moths and flies³⁵ (fig. 39) in pitch near insect wounds or other wounds, occur in all sections of the country.

DEFECTS CLASSED AS GUM SPOTS OR STREAKS

"Gum spots" in western hemlock³⁶ and Douglas fir³⁷ are caused by insect attack.

Gum spots entail practically no loss to mills, nevertheless they cause losses to the builder and consumer. Gum-spot defects are also due to causes other than insects.

³³ *Sesia novaeocensis* Hy. Edw.

³⁴ *Vespa minima sequoia* Hy. Edw.

³⁵ *Cheilosia* spp.

³⁶ *Melanophila drummondi* Kirby causes this defect in both hemlock and Douglas fir (fig. 40).

³⁷ The pitch moth (*Pinipestis zimmermanni* Grote) is responsible for not more than 10 per cent, *Dendroctonus pseudotsugae* Hopkins for not less than 70 per cent, and all other causes for about 20 per cent of the damage.

DEFECTS CLASSED AS BLACK CHECK

"Black check" is the lumberman's name for a common defect consisting of a dark stain in the heartwood and sapwood, surrounded by thickened, curled, or abnormal layers of wood (fig. 41) (6). It should not be confused with stains associated in the wood with other boring insects (as ambrosia beetles), where the wood is rarely distorted. Although also caused by other agents, it particularly applies to the defect in western softwoods (hemlock, fir, spruce, and yellow pine) produced by the maggots of several small flies³⁸ or moths,³⁹ but the term is also used for a somewhat similar defect in oaks produced by the larvae of large beetles.⁴⁰ (Figs. 42 and 43.)

FIG. 42.—Pile of logs with grub holes made by round-headed borer (*Romaleum rufulum*). This defect is considered as "wormholes, no living worms or decay"

These defects are caused by the insects injuring the growing tissue of the tree and killing a small area of the outer layers of wood. The subsequent growth of the tree finally heals over this injury, but many annual layers are stained and distorted. The original small pocket or cavity where the insect was working remains, and in coniferous trees it fills up with pitch.

Western hemlock contains more of this injury than any other softwood. In quarter-sawn (vertical-grain) wood the checks appear as small seams one-half to 1 inch long, with one side curly, while in bastard-sawn (flat-grain) boards they appear as oval or rounded spots from one-half to 1 inch in diameter. The defects in other conifers are similar but not so numerous.

³⁸ *Chcitosia* spp.

³⁹ *Parharmonia*.

⁴⁰ *Romaleum* sp. and *Gocis* sp.

In oak the black checks are larger, often 6 inches long over an area of several square inches. In quarter-sawn boards they appear as dark distorted or curly wood and run through many layers of wood, while in bastard-sawn boards they form black scars several inches long by 1 or 2 wide. In the Ozark Mountains of Arkansas this is a very common defect.

Timber from softwoods or hardwoods is not seriously reduced in strength by such defects, but the wood is lowered in grade and rendered useless for finishing, turning, staves, and woodenware. It can, however, be used for structural material or for purposes for which the marred appearance will not be detrimental. Since black

check is always the result of injury to the living tree, it is not preventable, from the lumberman's standpoint.

**DEFECTS CLASSED AS
BLUING OR STAINING**

FIG. 43.—Black check in oak, caused by round-headed borer (*Goes* sp.) in the living tree. The wound has healed over

bluish-black streaks or stains in the sapwood of pine, especially in southern yellow, western yellow, and sugar pines, as well as in certain hardwoods, such as red gum or sap gum, are classed as bluing. This defect occurs either in the standing tree or in recently cut green logs or green lumber and is caused by fungi developing from spores which are no doubt carried by insects. In pines the bluing directly follows in-

festation by tree-killing bark beetles⁴¹ in the standing trees (fig. 44) (29). This defect can be prevented only by the control of tree-killing bark beetles. Similar staining defects are caused by many ambrosia beetles⁴² in both the living trees and green saw logs and green, freshly sawn lumber. (Figs. 4, 9, and 13.) This staining can be prevented in green saw logs by prompt utilization of the green logs; by placing them in the mill pond soon after cutting; or by sun-curing. When the lumber is sawn it should be either kiln-dried or air-seasoned rapidly. Bluing is not always dependent on bark beetles (42).

⁴¹ *Dendroctonus* spp.

⁴² *Gnathotrichus*, *Xyleborus*, *Platypus*, etc.

FIG. 44.—Section of short-leaf pine, showing “blue stain” of sapwood after attack by the southern pine beetle (*Dendroctonus frontalis*). A, side view; B, cross section showing stain extending to the heartwood. Serious bluing in the log sometimes appears to take place without the aid of insects

FIG. 45.—A, Pith flecks in river birch, transverse section, caused by *Agromyza pruinosa* (2); B, pith flecks in silver maple, tangential section, caused by *Agromyza accris*. Natural size

More general and excessive bluing by stain fungi in green lumber after it has been sawn causes a sound defect and is also called "sap stain." Rapid handling of the logs, kiln-drying the lumber, steaming it in a kiln, and air-seasoning are preventive measures. Dipping the lumber in chemical solutions is effective under certain conditions, but this method of prevention has not been entirely satisfactory (37, 38, 40).

DEFECTS CLASSED AS PITH FLECKS

PITH-FLECK INJURY CAUSED BY THE LARVAE OF FLIES

Birch, cherry, maple, oak, poplar, and many other less commonly used hardwood trees have small "pith flecks," i. e., yellow-brown spots or narrow streaks in the sapwood and heartwood, due to the burrows made by larvae of flies⁴³ in the cambium of living trees (fig. 45) (2, 21, 22). This is a very common and widespread defect, more common in soft than in hard maples, and more common in river birch (*Betula nigra*) than in the other birches. Where clear birch is used, as for spool stock, it throws out considerable material. However, this is merely a trade prejudice and should be overcome by educational propaganda, in accordance with the move for closer utilization. This is a sound defect caused in the living tree, the flies laying their eggs in young branches and burrowing down the trunk through the cambium long distances to the roots, where they emerge and pupate. There is no known remedy to prevent the insect injury to the living tree, and hence there is greater need for closer utilization of the product.

PITH-FLECK INJURY CAUSED BY THE FEEDING OF ADULT WEEVILS

Another pith-fleck defect is that caused by the feeding of adult weevils.⁴⁴ This defect has been described (48) as occurring in fir. The adults puncture the bark to feed on the inner phloem, removing a small circular or oval disk one-fourth inch in diameter to one-fourth by 1 inch. This cavity fills with scar tissue and some gum and later heals over. It is very common in fir, but is not of much commercial importance. No doubt this defect occurs in other trees, as pine and spruce, which are attacked by species of this genus of weevils.

SUMMARY

Defects in timber caused by wood-boring beetles and grubs cause serious losses either by rendering the affected material unfit for use or by reducing it to lower grades.

Much of this loss can be prevented through proper methods of lumbering and management, usually involving but slight changes or modifications in present handling of the timber.

Some of the damage can not be avoided, but in such cases considerable loss can be prevented by periodical inspection and proper classification, piling, and handling of stock. More rapid and closer utilization of the material can often be attained.

⁴³ *Agromyza* spp.

⁴⁴ *Pissodes dubius* Rand.

This bulletin presents facts which will aid in determining the nature and cause of defects caused by insects, gives recommendations for avoiding preventable injury, and discusses the types of defects which are not preventable. Of course it is not intended to supersede previous classifications of timber and lumber, based on defects other than those due to insects.

The principal damage comes under two types of defects, designated as wormholes, with no living worms or decay, and powder post.

Sound wood can be utilized where the lower price offsets the lower grade and the defect is not objectionable. There will be no extension of the damage after the wood is dry or seasoned. Sound, wormy chestnut even can be exported with no fear of further damage from the insects which caused this defect to the wood of the tree when living.

Powder-posted stock can not be safely utilized because the damage is continuous in the seasoned and finished product and will not only cause further damage but will be a menace to other hardwood stock stored near by.

LITERATURE CITED

- (1) BETTS, H. S.
1917. THE SEASONING OF WOOD. U. S. Dept. Agr. Bul. 552, 28 p., illus.
- (2) BROWN, H. P.
1913. PITH-RAY FLECKS IN WOOD. U. S. Dept. Agr., Forest Serv. Circ. 215, 15 p., illus.
- (3) BRUNNER, J.
1914. THE SEQUOIA PITCH MOTH, A MENACE TO PINE IN WESTERN MONTANA. U. S. Dept. Agr. Bul. 111, 11 p., illus.
- (4) ———
1915. DOUGLAS FIR PITCH MOTH. U. S. Dept. Agr. Bul. 255, 23 p., illus.
- (5) ———
1915. THE ZIMMERMAN PINE MOTH. U. S. Dept. Agr. Bul. 295, 12 p., illus.
- (6) BURKE, H. E.
1905. BLACK CHECK IN WESTERN HEMLOCK. U. S. Dept. Agr., Bur. Ent. Circ. 61, 10 p., illus.
- (7) ———
1910. INJURIES TO FOREST TREES BY FLAT-HEADED BORERS. U. S. Dept. Agr. Yearbook 1909: 399-415, illus.
- (8) CHAMBERLIN, W. J.
[1924]. FOREST ENTOMOLOGY. AN ACCOUNT OF THE INJURIOUS AND BENEFICIAL INSECTS WHICH AFFECT FOREST AND SHADE TREES. 2 v. Ann Arbor, Mich. [Mimeographed.]
- (9) CRAIGHEAD, F. C.
1919. PROTECTION FROM THE LOCUST BORER. U. S. Dept. Agr. Bul. 787, 12 p., illus.
- (10) ———
1920. DIRECT SUNLIGHT AS A FACTOR IN FOREST INSECT CONTROL. Ent. Soc. Wash. Proc. 22: 106-108.
- (11) ———
1921. PROTECTION OF MESQUITE CORDWOOD AND POSTS FROM BORERS. U. S. Dept. Agr. Farmers' Bul. 1197, 12 p., illus.
- (12) ———
1922. EXPERIMENTS WITH SPRAY SOLUTIONS FOR PREVENTING INSECT INJURY TO GREEN LOGS. U. S. Dept. Agr. Bul. 1079, 11 p.
- (13) ———
1923. A BRIEF SUMMARY OF THE BUDWORM INVESTIGATIONS IN CANADA. Jour. Forestry 21: 134-138.

- (14) CRAIGHEAD, F. C., and LOUGHBOROUGH, W. K.
1921. TEMPERATURES FATAL TO LARVÆ OF THE RED-HEADED ASH BORER AS APPLICABLE TO COMMERCIAL KILN DRYING. Jour. Forestry 19: 250-254.
- (15) GRAHAM, S. A.
1918. THE CARPENTER ANT AS A DESTROYER OF SOUND WOOD. Minn. State Ent. Rpt. 17: 32-40, illus.
- (16) ———
1920. FACTORS INFLUENCING THE SUBCORTICAL TEMPERATURES OF LOGS. Minn. State Ent. Rpt. 18: 26-42, illus.
- (17) ———
1921. CONTROLLING INSECTS IN LOGS BY EXPOSURE TO DIRECT SUNLIGHT. Jour. Forestry 19: 512-514.
- (18) ———
1922. EFFECT OF PHYSICAL FACTORS IN THE ECOLOGY OF CERTAIN INSECTS IN LOGS. Minn. State Ent. Rpt. 19: 22-40.
- (19) ———
1924. TEMPERATURE AS A LIMITING FACTOR IN THE LIFE OF SUBCORTICAL INSECTS. Jour. Econ. Ent. 17: 377-383.
- (20) ———
1925. THE FELLED TREE TRUNK AS AN ECOLOGICAL UNIT. Ecology 6: 397-411, illus.
- (21) GREENE, C. T.
1914. THE CAMBIUM MINER IN RIVER BIRCH. Jour. Agr. Research 1: 471-474, illus.
- (22) ———
1917. TWO NEW CAMBIUM MINERS (DIPTERA). Jour. Agr. Research 10: 313-318, illus.
- (23) HOPKINS, A. D.
1894. DEFECTS IN WOOD CAUSED BY INSECTS. W. Va. Agr. Expt. Sta. Bul. 35, p. 291-306, illus.
- (24) ———
1894. BLACK HOLES IN WOOD. W. Va. Agr. Expt. Sta. Bul. 36, p. 313-336, illus.
- (25) ———
1904. INSECT INJURIES TO HARDWOOD FOREST TREES. U. S. Dept. Agr. Yearbook 1903: 313-328, illus.
- (26) ———
1904. CATALOGUE OF EXHIBITS OF INSECT ENEMIES OF FORESTS AND FOREST PRODUCTS AT THE LOUISIANA PURCHASE EXPOSITION, ST. LOUIS, MO., 1904. U. S. Dept. Agr., Div. Ent. Bul. 48, 56 p., illus.
- (27) ———
1905. INSECT INJURIES TO FOREST PRODUCTS. U. S. Dept. Agr. Yearbook 1904: 381-398, illus.
- (28) ———
1907. PINHOLE INJURY TO GIRDLED CYPRESS IN THE SOUTH ATLANTIC AND GULF STATES. U. S. Dept. Agr., Bur. Ent. Circ. 82, 4 p., illus.
- (29) ———
1909. PRACTICAL INFORMATION ON THE SCOLYTID BEETLES OF NORTH AMERICAN FORESTS. I. BARKBEETLES OF THE GENUS DENDROCTONUS. U. S. Dept. Agr., Bur. Ent. Bul. 83, pt. 1, 169 p., illus.
- (30) ———
1909. INSECT DEPREDATIONS IN NORTH AMERICAN FORESTS AND PRACTICAL METHODS OF PREVENTION AND CONTROL. U. S. Dept. Agr., Bur. Ent. Bul. 58, pt. 5, p. 57-101.
- (31) ———
1910. INSECT INJURIES TO THE WOOD OF DYING AND DEAD TREES. U. S. Dept. Agr., Bur. Ent. Circ. 127, 3 p.
- (32) ———
1910. INSECT INJURIES TO FOREST PRODUCTS. U. S. Dept. Agr., Bur. Ent. Circ. 128, 9 p.
- (33) ———
1912. DAMAGE TO THE WOOD OF FIRE-KILLED DOUGLAS FIR, AND METHODS OF PREVENTING LOSSES, IN WESTERN WASHINGTON AND OREGON. U. S. Dept. Agr., Bur. Ent. Circ. 159, 4 p.

- (34) HOPKINS, A. D., and SNYDER, T. E.
1917. POWDER-POST DAMAGE BY LYCTUS BEETLES TO SEASONED HARDWOOD.
U. S. Dept. Agr. Farmers' Bul. 778, 20 p., illus.
- (35) ——— and SNYDER, T. E.
1921. POWDER-POST DAMAGE TO TIMBER AND WOOD PRODUCTS. Engin. News-
Rec. 87: 269-271, illus.
- (36) HUBBARD, H. G.
1897. THE AMBROSIA BEETLES OF THE UNITED STATES. U. S. Dept. Agr.,
Div. Ent. Bul. (n. s.) 7: 9-30, illus.
- (37) HUBERT, E. E.
1921. NOTES ON SAP STAIN FUNGI. Phytopathology 11: 214-224, illus.
- (38) ———
1922. SOME WOOD STAINS AND THEIR CAUSES. Hardwood Rec. 52 (11):
17-19, illus.
- (39) IVORY, E. P., WHITE, D. G., and UPSON, A. T.
1923. STANDARD GRADING SPECIFICATIONS FOR YARD LUMBER AS RECOM-
MENDED BY THE DEPARTMENT OF AGRICULTURE. U. S. Dept. Agr.
Circ. 296, 75 p., illus.
- (40) LOUGHBOROUGH, W. C., and HUBERT, E. E.
1924. PROBLEMS IN THE SEASONING OF SOUTHERN HARDWOODS. South.
Lumberman 117 (1525): 170-174, illus.
- (41) ST. GEORGE, R. A.
1924. SOUTHERN PINE BEETLE AND OTHER INSECT ENEMIES OF SOUTHERN
FORESTS. Lumber Trade Jour. 86 (9): 37-38.
- (42) SCHRENK, H. VON.
1903. THE "BLUING" AND THE "RED ROT" OF THE WESTERN YELLOW PINE,
WITH SPECIAL REFERENCE TO THE BLACK HILLS FOREST RESERVE.
U. S. Dept. Agr., Bur. Plant Indus. Bul. 36, 40 p., illus.
- (43) SNYDER, T. E.
1916. EGG AND MANNER OF OVIPOSITION OF LYCTUS PLANICOLLIS. Jour.
Agr. Research 6: 273-276, illus.
- (44) ———
1922. INSECT INJURY TO GREEN LOGS AND LUMBER AND METHODS OF PRE-
VENTING THIS LOSS. South. Lumberman 108 (1422): 133-135,
illus.
- (45) ———
1923. CLOSER UTILIZATION HELPS CONSERVE OUR FORESTS—INSECT DEFECTS.
South. Lumberman 113 (1473): 131-134, illus.
- (46) ———
1923. HIGH TEMPERATURES AS A REMEDY FOR LYCTUS POWDER-POST BEETLES.
Jour. Forestry 21: 810-814.
- (47) ——— and ST. GEORGE, R. A.
1924. DETERMINATION OF TEMPERATURES FATAL TO THE POWDER-POST
BEETLE, LYCTUS PLANICOLLIS LECONTE, BY STEAMING INFESTED
ASH AND OAK LUMBER IN A KILN. Jour. Agr. Research 28:
1033-1038, illus.
- (48) SWAINE, J. M., CRAIGHEAD, F. C., and BAILEY, I. W.
1924. STUDIES ON THE SPRUCE BUDWORM [CACOECIA FUMIFERANA CLEM.]
Canada Dept. Agr. Bul. (n. s.) 37, 91 p., illus.
- (49) THELEN, R.
1923. KILN DRYING HANDBOOK. U. S. Dept. Agr. Bul. 1136, 64 p., illus.
- (50) WEBB, J. L.
1909. THE SOUTHERN PINE SAWYER. U. S. Dept. Agr., Bur. Ent. Bul. 58,
pt. 4, p. 41-56, illus.
- (51) ———
1911. INJURIES TO FORESTS AND FOREST PRODUCTS BY ROUNDHEADED BORERS.
U. S. Dept. Agr. Yearbook 1910: 341-358, illus.

ORGANIZATION OF THE UNITED STATES DEPARTMENT OF AGRICULTURE

July 8, 1927

<i>Secretary of Agriculture</i>	W. M. JARDINE.
<i>Assistant Secretary</i>	R. W. DUNLAP.
<i>Director of Scientific Work</i>	A. F. WOODS.
<i>Director of Regulatory Work</i>	WALTER G. CAMPBELL.
<i>Director of Extension</i>	C. W. WARBURTON.
<i>Director of Personnel and Business Administration</i>	W. W. STOCKBERGER.
<i>Director of Information</i>	NELSON ANTRIM CRAWFORD.
<i>Solicitor</i>	R. W. WILLIAMS.
<i>Weather Bureau</i>	CHARLES F. MARVIN, <i>Chief</i> .
<i>Bureau of Animal Industry</i>	JOHN R. MOHLER, <i>Chief</i> .
<i>Bureau of Dairy Industry</i>	C. W. LARSON, <i>Chief</i> .
<i>Bureau of Plant Industry</i>	WILLIAM A. TAYLOR, <i>Chief</i> .
<i>Forest Service</i>	W. B. GREELEY, <i>Chief</i> .
<i>Bureau of Chemistry and Soil</i>	, <i>Chief</i> .
<i>Bureau of Entomology</i>	L. O. HOWARD, <i>Chief</i> .
<i>Bureau of Biological Survey</i>	PAUL G. REDINGTON, <i>Chief</i> .
<i>Bureau of Public Roads</i>	THOMAS H. MACDONALD, <i>Chief</i> .
<i>Bureau of Agricultural Economics</i>	LLOYD S. TENNY, <i>Chief</i> .
<i>Bureau of Home Economics</i>	LOUISE STANLEY, <i>Chief</i> .
<i>Federal Horticultural Board</i>	C. L. MARLATT, <i>Chairman</i> .
<i>Grain Futures Administration</i>	J. W. T. DUVEL, <i>Chief</i> .
<i>Food, Drug, and Insecticide Administration</i>	WALTER G. CAMPBELL, <i>Director of Regulatory Work, in Charge</i> .
<i>Office of Experiment Stations</i>	E. W. ALLEN, <i>Chief</i> .
<i>Office of Cooperative Extension Work</i>	C. B. SMITH, <i>Chief</i> .
<i>Library</i>	CLARIBEL R. BARNETT, <i>Librarian</i> .

This bulletin is a contribution from

<i>Bureau of Entomology</i>	L. O. HOWARD, <i>Chief</i> .
<i>Forest Insect Investigations</i>	F. C. CRAIGHEAD, <i>Senior Entomologist, in Charge</i> .

47

ADDITIONAL COPIES
OF THIS PUBLICATION MAY BE PROCURED FROM
THE SUPERINTENDENT OF DOCUMENTS
GOVERNMENT PRINTING OFFICE
WASHINGTON, D. C.

AT
15 CENTS PER COPY

POWDER-POST DAMAGE BY LYCTUS BEETLES TO SEASONED HARDWOOD

A. D. HOPKINS

Forest Entomologist, and

T. E. SNYDER

Specialist in Forest Entomology

FARMERS' BULLETIN 778

UNITED STATES DEPARTMENT OF AGRICULTURE

Contribution from the Bureau of Entomology

L. O. HOWARD, Chief

POWDER-POST BEETLES often ruin stored hardwoods of the finest quality by turning them into a flourlike powder. They work in whitewood or sapwood, especially second-growth hickory, ash, and oak, which has been stored or piled in one place for two or three years or longer.

Sapwood seasoned for less than eight or ten months will not be attacked, and *heartwood is never attacked*.

In the manufacture of hickory implement handles, when feasible substitute heartwood for sapwood. See footnote 2, page 15.

By the adoption of the following system of inspection, classification, and the proper disposal of the seasoned sapwood of hardwood stock, loss by powder-post beetles can be prevented:

Inspect material in yards and storehouses annually, preferably in November and February, and sort out and burn material showing evidence of powder post. Burn all refuse and useless sapwood material.

Classify seasoned hardwood stock into (a) hickory, ash, oak, etc.; (b) heartwood, pure sapwood, and part sapwood; and (c) according to the number of years seasoned.

Utilize or sell oldest stock first.

Inspect all new stock to prevent introduction of powder-posted material.

Use only heartwood piling sticks in lumber piles.

To prevent attack treat the more valuable material, between October and March, with boiled linseed oil. Varnish or paraffin the sapwood portions of backs and interior surfaces of cabinet work, inside finish, and furniture.

Material once attacked is usually damaged beyond repair. If the injury is not too far advanced further damage can be stopped by saturating the wood with kerosene.

While there are a number of effective chemical treatments to prevent attack, and to kill the insects in the wood, the great objection to all of them is the great and recurring expense of treating material that never would be attacked if, by the proper methods of management, the premises were kept free from the pest.

POWDER-POST DAMAGE BY LYCTUS BEETLES ¹ TO SEASONED HARDWOOD.

CONTENTS.

	Page.		Page.
Evidence of attack and character of powder-post injury.....	4	Habits and seasonal history of Lyctus beetles.....	8
Classes of forest products damaged.....	4	The four species responsible for most of the losses.....	12
Extent of losses, and those upon whom they fall.....	7	Conditions favorable for attack.....	15
Character of the insects which cause the damage.....	8	Conditions unfavorable for attack.....	15
		Insect enemies of Lyctus beetles.....	17
		Methods of control.....	18

The sapwood of seasoned hardwood material of all kinds, both finished and unfinished, especially of hickory, ash, and oak, is often ruined by yellowish-white grubs from one-eighth to one-fifth inch in length which burrow through the solid wood in all directions and convert it into powder. These grubs are the young, or larvæ, of small, slender, somewhat flattened, reddish-brown to nearly black beetles, known as powder-post beetles. Wood that has been seasoned a year or longer is especially liable to this peculiar type of insect injury, which sometimes amounts to from 10 to 50 per cent of neglected material in storage. Powder-post damage by these beetles is widely distributed over the world and the consequent losses have been severe.

Under the general class of powder-post injury the work of insects representing several species and groups of species of beetles may be found, but by far the larger part and the most serious damage and losses are caused by different species of *Lyctus*.

¹ The *Lyctus* beetles which are responsible for the principal losses belong to four species, viz, *Lyctus linearis* Goeze, *L. planicollis* Le Conte, *L. parallelipedus* Melsheimer, and *L. caricollis* Le Conte. These are discussed separately on pages 12-15.

NOTE.—Systems of control which adapt certain features in business management to the habits and seasonal history of *Lyctus* powder-post beetles have been perfected by the Bureau of Entomology and have been adopted by many manufacturers and dealers in seasoned hardwood products, and there has been a marked decrease in powder-post losses since 1906. The object of this bulletin is to describe the methods which have been found effective in preventing these losses and to induce a more general adoption of them throughout the United States as well as to show the character and extent of the damage.

EVIDENCE OF ATTACK AND CHARACTER OF POWDER-POST INJURY.

Infested wood may be detected by the fine, flourlike powder found on or beneath piled or stored hickory, ash, oak, and other seasoned hardwood products. During the first year of infestation the powder comes from exceedingly minute holes in the wood, but after the second year the small holes from which the beetles have emerged are more or less conspicuous, and from these the powder will fall when the infested material is moved or jarred. When the wood is cut or split the interior is often found converted into a mass of closely packed powdery material which has been held together by an outer thin shell and intervening fibers of sound wood. The grubs, burrowing through the solid wood in all directions, have pulverized the wood fiber and have packed their burrows with this powdered wood. The injury by *Lyctus* beetles is always confined to the white wood or sapwood, although the heartwood is sometimes penetrated when the matured beetles are emerging from the wood.

CLASSES OF FOREST PRODUCTS DAMAGED.

A great variety of seasoned hardwood products is subject to powder-post damage, especially hickory, ash, and oak woodwork of farming machinery and implement handles (see illustration on title-page, and figs. 1, 2, and 3); ladder stock, as rungs, etc.; and vehicle stock, as hubs, spokes, felloes, rims, singletrees, poles and shafts. Other products affected include woodwork of electric street cars; shipbuilding lumber (fig. 4); the Army and Navy stores of handles, tent poles, wheelbarrows, oars, and many other hardwood articles; interior finish or trim and ornamental woodwork, as panels, mantels, doors, doorposts, staircases, wainscoting, flooring, etc.; construction timber, including beams, joists, roof framing, etc.; furniture, including tables, chairs, bureaus, cabinets, refrigerators (before use), filing cases (fig. 5), piano stock, bookcases, cabinet-work, etc.; inside rustic work; wood specimens and curios in museums; cooperage stock (barrel-stave bolts); shoe-last blocks; walking sticks, umbrella handles, measuring rules, and blocks to be converted into golf-stick heads; fish-net hoops; ornamental bamboo (fig. 6); Japanese fans; shuttle blocks, and "picker" sticks (for driving shuttles in looms), etc.

Hickory, ash, and oak are the kinds of wood most liable to injury, but persimmon, osage orange, black walnut, butternut, maple, elm, wild cherry, locust, poplar, sycamore, eucalyptus, sassafras, orange wood, fig, bamboo, and other woods are also attacked.

FIG. 1.—Powder-posted ash shovel handle, showing exit holes of adult beetles; the work of *Lyctus planicollis*. (Original.)

FIG. 2.—Powder-posted ash shovel handle with surface wood cut away to show burrows of larvæ within; work of *Lyctus planicollis*. (Snyder.)

EXTENT OF LOSSES AND THOSE UPON WHOM THEY FALL.

Powder-post causes a loss which falls alike on the dealer, the manufacturer or owner, and the consumer of finished products. The producer of the unseasoned or crude product is not affected, because it is only after the wood is seasoned a year or more that it is attacked and damaged by these insects. Second-growth sapwood of the best

FIG. 3.—Powder-posted hickory pickaxe handles, showing exit holes of adult beetles; work of *Lyctus linearis*. (Original.)

quality is particularly liable to attack and serious injury, especially when it has been stored or piled in one place for two or three years or more.

The loss to seasoned hardwood products ranges from 1 to 50 per cent, sometimes representing thousands of dollars to a single manufacturer or dealer who neglects to adopt the proper preventive measures. The affected articles are not only reduced in value, but frequently are rendered worthless for the purposes for which they are intended. In the aggregate the direct financial loss that has been

caused by these beetles in this country has amounted to hundreds of thousands of dollars annually. The loss increases with the length of time the infested stock is held in storage. In certain cases powder-post injury may be a menace to human life as in the weakened wood material of vehicles, ladders, etc.

CHARACTER OF THE INSECTS WHICH CAUSE THE DAMAGE.

The *Lyctus* beetles are small, slender, somewhat flattened, brownish to nearly black beetles, which, upon emerging from the wood where

FIG. 4.—Powder-posted white ash shipbuilding lumber, showing burrows of larvæ and exit holes of adult beetles; board from closely piled stack of lumber throughout which larvæ had burrowed; work of *Lyctus planicollis*. (Original.)

they breed and pass the winter as larvæ, fly or crawl about in search of suitable wood material in which to deposit their eggs.

HABITS AND SEASONAL HISTORY OF LYCTUS BEETLES.

There are four stages in the life history of these insects—namely, the egg, the larva or grub, the pupa or resting stage, and finally the adult or beetle.

The egg (fig. 7) is deposited in the pores of the wood by the female beetle soon after it emerges from the wood. The minute larva which hatches from the egg proceeds to burrow in and through the wood in all directions, feeding and growing as it proceeds, until it has attained its full size. The full-grown larva (fig. 8) is a yellowish-white grub ranging in length from one-eighth to one-fifth of an inch,

with three pairs of legs, and with the tip of the body curved under toward the head. Upon reaching maturity as a larva it excavates a cell at the end of its burrow (fig. 9), and in this transforms to the pupa (fig. 10). Later the pupa changes to the adult beetle (fig. 11), which can fly in search of suitable places to lay its eggs.

FIG. 5.—Powder-posted oak file case, showing exit holes of adult beetles; work of *Lyctus planicollis*. (Original.)

Seasoned wood is attacked by these beetles because the chemical changes which take place in the process of seasoning render the nutritive substances in the wood, such as sugar and starch, especially suitable as food for the development of the young or larval stage.

Each female beetle deposits many eggs,¹ and many females oviposit in a single piece of wood, so that the combined work of their nu-

¹ A large female of *Lyctus planicollis* Lec., about one-fourth inch in length, that was dissected contained ovaries as two separate, elongate ovate clusters one-tenth inch in length, one of which was about one-third the size of the other. These ovaries took up most of the space of the abdomen. The eggs were arranged in rows so as to have the appearance of being braided. The mature ovules, or fully formed eggs, are pointed at one end and rounded at the other, with the rudimentary strand present at the rounded end. The egg surface is apparently granular. At least 40 eggs were in the larger mass.

merous progeny, burrowing through the wood in quest of food for their development, results in the complete destruction of the interior wood fiber and its conversion into a mass of fine powder. If the first attack and the first generation do not accomplish this destruction, subsequent generations will follow in the same wood until nothing of the solid fiber is left but a thin outer shell.

FIG. 6.—Powder-posted ornamental section of bamboo, showing exterior and interior of split bamboo; work of *Lyctus parallelopipedus*. (Original.)

The different kinds of *Lyctus* beetles vary somewhat in their habits and seasonal history, but there is a general similarity. They pass the winter as larvæ in the wood, change to pupæ in the early spring, and during late spring and early summer the adult beetles emerge from the wood and fly about. Under natural out-of-door conditions the eggs are laid in the pores of the wood soon after activity commences in the spring, but in storehouses, sheds, or buildings kept warm and dry, the development may take place and the eggs may be deposited much earlier.

The larvæ remain dormant or active in the wood, according to the temperature of the place where stored, and in consequence infestation in wood stored in cold places may pass unnoticed. If the wood is re-

FIG. 7.—Egg of a powder-post beetle, *Lyctus planicollis*, in pore of wood on radial section of ash ladder-rung stock; pore opened to show egg. Highly magnified. (Snyder.)

moved from outdoors, open storage sheds, or other places which are cold in winter, to the dry kiln, the dormant larvæ start to work in the wood, as shown by the ejection of boring dust or powder. Handling or jarring infested wood appears temporarily to stop the activity of larvæ contained in it.

In general, the adults or beetles are active between the first of March and the last of July at Washington, D. C. (about latitude 39° and longitude 77° , and practically at sea level), but the period varies according to the species, the temperature of the storerooms, and the geographical location. According to the senior author, the variation due to latitude, longitude, and altitude from a date at Washington, D. C., will be approximately four days for each degree of latitude, 5° of longitude, and 400 feet of altitude, being earlier in the season southward, westward, and descending. The number of adult beetles that might be active and deposit eggs after the last of July at Washington, D. C., is so insignificant that practically no damage is done by them. Therefore the danger of attack would be over relatively earlier at parts south of Washington and westward, and later northward. The approximate dates of flying and

FIG. 8.—Larva of a powder-post beetle, *Lyctus planicollis*. Enlarged. (Chittenden.)

egg laying by each species under outdoor conditions at Washington, D. C., are given on pages 13, 14, and 15.

So far as known there is only one generation annually.

THE FOUR SPECIES RESPONSIBLE FOR MOST OF THE LOSSES.

While, as previously stated, there are many species of *Lyctus* beetles in the United States, the European *Lyctus*, the southern *Lyctus*, the velvety *Lyctus*, and the western *Lyctus* are the four species responsible for practically all the losses.

THE EUROPEAN LYCTUS.¹

The European *Lyctus* beetle is rusty red-brown, slender, somewhat flattened, elongate, and from one-tenth to one-fifth inch in length. The wing covers have single rows of large, rounded, very shallow punctures.

This species is commonly met with in northern Europe under natural conditions as well as in commercial products, especially in England, France, and Germany. Therefore it is evident that it was introduced into this country from Europe. In addition to the wide distribution of this species in the temperate zones of the world, it is especially common and destructive in the States north of North Carolina, Tennessee, and Arkansas. There are a great many records of damage to

FIG. 9.—Pupal cells of *Lyctus planicollis* in powder-posted white-ash shovel handle. (Snyder.)

FIG. 10.—Pupa of a powder-post beetle, *Lyctus planicollis*. (Chittenden.)

seasoned hardwood products by this beetle from this area, including Maine, Massachusetts, New York, Pennsylvania, Virginia,

¹ *Lyctus linearis* Goeze.

West Virginia, Ohio, Indiana, Illinois, Missouri, Iowa, Michigan, and Minnesota. The great number of specimens in the collection of the United States National Museum show but a single record from the South Atlantic and Gulf States, and this is evidently a temporary introduction. Undoubtedly the species is frequently carried into the southern region, but evidently it does not survive there. This beetle is common in commercial products of seasoned hickory, oak, walnut, and ash, and occasionally in poplar, wild cherry, and locust. It is not recorded from natural growth in the open in this country, but is said to infest such growth, as well as commercial products, in Europe.

Adults of the European species, which has been shown to have a relatively northern distribution in the United States, begin activity as early as the first part of March in its southern distribution and are still active during the first part of August in its northern distribution, general emergence of the adult beetles from infested wood occurring at Washington, D. C., from April until June. The eggs are deposited soon after the adult beetles emerge and are inserted into the pores of the wood by means of a long, slender, flexible ovipositor. The winter is passed in the larval stage in the wood. General pupation occurs from about the middle of April to the first of May in the District of Columbia.

THE SOUTHERN LYCTUS.¹

The southern *Lyctus* beetle is pitchy black, slender, somewhat flattened, elongate, and from one-tenth to one-fifth inch in length. The wing covers have smaller, finer, and deeper punctures in more or less distinct double rows. Individuals of this species vary extremely in size, and there is marked difference in size between the sexes, the males often being much smaller than the females. As the result of continued breeding in the same wood for several generations, in confinement the beetles were found to decrease in size.

This species is injurious to hardwood products in and from the South Atlantic and Gulf States to California, Arizona, and Nevada. This range probably represents the natural distribution of this southern species. It is frequently introduced into the Northern, Central, and Western States and transported to other countries, but this apparently never results in its permanent establishment except in warm storehouses. There are definite records of its occurrence at Cape

FIG. 11.—Adult powder-post beetle, *Lyctus planicollis*. (Chittenden.)

¹ *Lyctus planicollis* Lec.

Town, South Africa, in lumber received from the southern United States, and at Buenos Aires, Argentina, in the ash wood in refrigerators and shovel handles from the United States.

The southern *Lyctus* has been recorded as breeding under natural conditions in this country and is common in the Southern States in commercial products, such as seasoned ash, oak, hickory, persimmon, and sycamore. Rearing records show that it may breed continuously in the same wood during a period of at least six years, or until all the wood tissue has been converted into powder. The beetles emerge from infested wood of commercial products in heated buildings much earlier than where exposed to outdoor conditions. Owing to the character of the class of commercial products infested by *Lyctus* beetles, which are often stored indoors, there apparently are no great differences in the periods of activity of adults of species of northern and southern distribution.

Adults of this southern species are active from the middle of February till the last of September in the South Atlantic and Gulf States. General emergence of the adult beetles and mating occur from the middle of April to June, and very few beetles emerge at Washington, D. C., after the first part of July. Egg laying occurs a few days after emergence, and the eggs hatch after a period of not longer than 10 days. The winter is passed as larvæ in the wood, and full-grown larvæ are in the pupal cells at Washington by the first part of February. General pupation of the larvæ in the infested wood in rearing experiments at Falls Church, Va., occurred from about the middle of March to the first part of April.

THE VELVETY LYCTUS.¹

The velvety *Lyctus* is rusty red-brown to black, slender, flattened, and elongate, and from one-twelfth to one-sixth inch in length. The punctures on the wing covers are very fine and obscure and not placed in rows; the fine, dense, yellowish hairs on the wing covers are prominent, which gives the beetle a velvety appearance.

Extensive observations have been made on the habits of this species, which has been recorded as injurious in Texas, Louisiana, Florida, Georgia, South Carolina, Virginia, West Virginia, District of Columbia, Long Island, New York, Ohio, and Missouri.

The natural distribution of the species is evidently in the South Atlantic and Gulf States and in part of the Mississippi and Ohio River Valleys, from which it has been temporarily introduced into other States.

This beetle infests the seasoned sapwood of commercial products made from persimmon, hickory, ash, oak, and bamboo; it also lives

¹ *Lyctus parallelipipedus* Melsh.

in the dead wood of natural growth of osage orange, sassafras, and fig.

Adults of the velvety *Lyctus* are active from the middle of March till the last of August at Washington, D. C. General emergence occurs from June to the last of July.

THE WESTERN LYCTUS.¹

The western *Lyctus* is a rusty red-brown, slender beetle, somewhat flattened and elongate, and from about one-seventh to about one-fifth inch in length; the punctures on the wing covers are very fine and placed in more or less distinct double rows.

This species is recorded from California in commercial products, seasoned orange wood and hickory, tanbark oak, and cordwood of live oak.

Adults of this species are active from April till the middle of September.

CONDITIONS FAVORABLE FOR ATTACK.

Second-growth white wood or sapwood of the finest quality of hardwoods, especially hickory, ash, and oak, which has been stored or piled in one place for two or three years or longer is especially liable to damage by *Lyctus* beetles. This is particularly the case if the material has not been handled or moved and if the old stock is allowed to accumulate. Refuse and useless sapwood material and infested stock piled about the sheds or yards, sapwood piling sticks, etc., are sources of infestation. If the different species of hardwoods are not placed in separate piles, the species of woods not so liable to attack often become infested.

CONDITIONS UNFAVORABLE FOR ATTACK.

Material of species other than hickory, ash, and oak is less liable to attack. Sapwood seasoned for less than 8 to 10 months will not be attacked, and *heartwood is never attacked*² (fig. 12).

The conditions will be rendered very unfavorable for attack by powder-post beetles wherever hardwood stock in yards and store-houses is handled as follows: Material inspected and rehandled annually, preferably in November and February, and that showing evidence of powder-post attack sorted out and burned; all refuse and useless sapwood material burned; as far as practicable, all dry or seasoned hardwood stock separated into heartwood, pure sapwood, and part sapwood, and into hickory, ash, oak, etc., and classified according to the number of years it has been seasoned, *and the oldest*

¹*Lyctus carvicollis* Lec.

² Exhaustive strength tests conducted by the Forest Service show that the heaviest, and consequently the strongest, hickory averages below 10 rings per inch in rate of growth, and that, weight for weight, red hickory (heartwood) is as strong as white hickory (sapwood).

stock utilized and sold first; accumulations of refuse material in which the insects could breed prevented; the introduction into lum-

FIG. 12.—Powder-posted sapwood oak veneer laid on a core of chestnut (door stock); work of *Lycus planicollis*. Note that the heartwood oak and the chestnut have not been attacked. (Original.)

ber yards and storehouses of material infested with powder post prevented; and, finally, heartwood instead of sapwood piling sticks used in the piles.

INSECT ENEMIES OF LYCTUS BEETLES.

There are many insect enemies of powder-post beetles. Among the beetles, species of the family Cleridae are often beneficial in checking the multiplication of *Lyctus* beetles, the larvæ preying on the *Lyctus* larvæ and the adults preying on the *Lyctus* adults. Other beetles which commonly prey upon the powder-post beetles belong to two other families.¹

FIG. 13.—Exit holes of small, four-winged, wasplike parasites of *Lyctus* in powder-posted red oak interior finish. (Original.)

Many species of small, four-winged, wasplike parasites have the *Lyctus* beetles as their hosts. The emergence or exit holes of the adult parasites (fig. 13) are much smaller than those made by the beetles, but are often mistaken for those made by young *Lyctus* larvæ.

¹ Histeridae and Cucujidae.

In general, these insect enemies, although beneficial, can not be depended upon to keep the destructive beetles under control, and therefore usually can be disregarded when practical control measures are being instituted.

METHODS OF CONTROL.

The methods of combating this class of insects and of preventing losses from their ravages come under two primary heads: (1) The destruction of the insects, or remedies, and (2) prevention of attack.

REMEDIES.

Infested material, including infested sap edges of lumber, etc., and all refuse sapwood in which the insects might breed should be eliminated by sorting it out and disposing of it by burning or otherwise. All material showing the slightest evidence of powder-post damage should be discarded and destroyed, except possibly such articles as may be tested for required strength and found to be of sufficient value for retention after treatment.

Material slightly infested and damaged should be treated with kerosene oil, after which it should be kept in quarantine a sufficient length of time to determine whether a second treatment is required.

All partially damaged material, such as the sap edges of lumber and parts of other material too valuable to be destroyed, should, when practicable, have the damaged and infested parts cut away and burned.

The work should be done between October and the first of March in storehouses, and before the first of April in the open. If this is thoroughly done and, by annual inspection thereafter, infested material is disposed of as soon as found, there will soon be no trouble from this source, unless there is a continued introduction in lumber and material received from other yards and localities where methods of control are neglected. To avoid this, all material should be carefully inspected before shipment, or at least before it is placed in the yards or storehouses.

Where it is not practicable to remove the infested parts, or in case of the more valuable stock, the wood should be subjected to methods of treatment for the destruction of the insects between October and the first of March. Of the following remedies, that most suitable in each case should be selected, remembering that the treatment must not be detrimental to the wood for subsequent uses.

(1) Liberal applications of pure kerosene oil with a brush, or immersion of infested wood in vats of kerosene. The only objection to kerosene is the fire risk. However, kerosene soon evaporates, so that the treated material is not long near the danger point. Kero-

sene oil does not affect the application of shellac or varnish; the only effect on finishing is to make it more difficult to stain kerosene-treated sapwood to match the rest.

(2) Mixtures of 3 parts creosote and 1 part kerosene oil; 3 parts kerosene oil and 1 part creosote (to obtain a deeper penetration); and 1 part creosote and 3 parts naphtha have been used successfully. The wood should be dipped in vats of preservative, preferably heated by coils of steam pipe, or the preservative may be applied hot with a brush. *Of course, mixtures containing kerosene oil should not be heated over a direct fire.*

(3) Thorough steaming of the infested wood in a tight room or under pressure. Steaming under pressure weakens and discolors the wood and should not be applied to wood to be used for fine finish or where strength is essential.

(4) Subjecting seasoned wood to temperatures over 200° F. in dry kilns.

(5) Fumigation of infested wood in tightly closed drying rooms with the fumes of sulphur at the time of the emergence of the adult beetles. This has been recommended for killing the beetles and preventing egg laying.

The objection to treating infested material lies in the fact that if it is once infested it may be damaged beyond repair and not safe for future use. A break might result in a serious accident and reflect on the reputation of the manufacturer or distributor.

PREVENTION OF DAMAGE.

By the adoption of the following system of inspection, classification, and the proper disposal of the seasoned sapwood of hardwood stock, loss by powder-post beetles can be prevented:

(1) Inspect material in yards and storehouses annually, especially stock two or more years old, preferably in November and February. Then (a) sort out and burn material showing evidence of powder-post attack and (b) burn all useless sapwood material.

(2) Classify all dry or seasoned hardwood stock as (a) hickory, ash, oak, etc.; (b) heartwood, pure sapwood, and part sapwood; and (c) according to age, that is, the number of years it has been seasoned.

(3) *Utilize or sell oldest stock on hand first. Prevent the accumulation of old stock; in other words, keep the stock moving.*

(4) Prevent the accumulation of refuse material in which the insects can breed.

(5) Use only heartwood piling sticks in lumber piles.

(6) Inspect all new stock to prevent the introduction into lumber yards and storehouses of powder-posted material.

(7) If the beetles have not been eliminated from the yards and storehouses, stock that has been seasoned longer than eight months,

and which is to be held in storage, may be rendered immune by treating it with two coats of boiled linseed oil applied hot, or it may be immersed in vats of hot oil. The boiled oil will dry more rapidly than the raw linseed oil, especially if kerosene is added to the hot oil. The wood should be treated between October and the 1st of March.

Linseed oil has an advantage over other substances in that it can also be profitably applied to unseasoned timber, since it will prevent checking in seasoning. This oil stains the wood slightly yellow. In case of wagon stock, however, it can be used without prejudice to the trade and is an effective preventive.

Creosotes can be effectively used in case of stock to which the brown stain will not be detrimental. The wood should be dipped in vats of hot creosote, or the preservative should be applied hot with a brush.

In case of finished products or more valuable material, any substance which closes the pores of the wood may be effectively applied. For example, paraffin wax, varnish, etc., effectively close the pores of the wood and prevent the beetles from depositing the eggs, which are laid in these pores. Thus the sapwood portions of backs and interior surfaces of cabinet work, inside finish, and furniture, etc., should also be treated to prevent attack.

In conclusion it may be stated that while there are a number of effective chemical treatments to prevent attack, and to kill the insects after they get into the wood, the great objection to all of them is the great and recurring expense of treating material that never would be attacked if, by the proper methods of management, the premises were kept free from the pest.

U. S. DEPARTMENT OF AGRICULTURE

FARMERS' BULLETIN No. 1582

PROTECTION OF LOG CABINS, RUSTIC WORK AND UNSEASONED WOOD *from* INJURIOUS INSECTS

INSECTS cause considerable damage to the unbarked logs of the principal woods used in the construction of log cabins, rustic woodwork, and certain other unseasoned products. This damage, which varies from the making of numerous holes in the bark to the complete destruction of the sapwood and heartwood, causes annoyance and unsightly condition of the material as well as large financial loss. Some of the insects which cause the damage are active nearly every month of the year except December, January, and February. It has been found that by cutting the trees at certain seasons and by treating the wood with preventive and remedial substances practically all such insect damage can be prevented or checked. This bulletin tells how it can be done.

Although the information contained herein applies specifically to the insects which occur in the eastern and southern parts of the United States, the principles set forth can be applied equally well to problems of this nature which occur in the country as a whole, provided due allowance is made for individual differences in insect and tree species and also in the time of flight of the beetles because of differences in climate.

PROTECTION OF LOG CABINS, RUSTIC WORK, AND UNSEASONED WOOD FROM INJURIOUS INSECTS

By R. A. ST. GEORGE, *Associate Entomologist, Division of Forest Insects, Bureau of Entomology*

	Page		Page
How beetles and grubs damage unseasoned wood.....	1	Conditions favorable and unfavorable for attack.....	14
Woods principally used for cabins, rustic work, and certain wood products.....	1	Conditions favorable for bark beetles and ambrosia beetles.....	14
Classes of insects responsible and the damage they do.....	2	Conditions favorable for powder-post beetles.....	14
Bark beetles.....	2	Conditions favorable for roundheaded borers and flat-headed borers.....	15
Ambrosia beetles.....	4	Preventive and control measures.....	15
Wood borers.....	4	Prevention of attack.....	15
Seasonal activity of the insects.....	11	Treatment after attack.....	18

DURING the last few years there has been an increasing demand for information on the protection of log cabins and rustic work from injurious insects. This is largely due to the popularizing of National and State forests and parks for recreational purposes, as a result of which many rustic bridges, benches, and log cabins have been constructed in such places. But it is also due to the increased building of rustic summer homes and arbors on private woodlands and estates. Wood with the bark still on is subject to injury by many kinds of beetles; therefore manufacturers, as well as the users, of rustic furniture constantly request advice. In addition, concerns requiring small raw poles from which to manufacture certain finished products, such as shuttle blocks, mallets, and mauls, often suffer severe losses and call for advice.

HOW BEETLES AND GRUBS DAMAGE UNSEASONED WOOD

Woods cut at certain seasons of the year are subject to attack by beetles which fill the bark with holes, thus causing sawdustlike borings to fall out and lodge on the wood. The larvae, or grubs, of wood-boring beetles mine the inner bark, causing the bark to loosen and fall off, and they bore into the sapwood and sometimes the heartwood of logs, making large holes and often reducing them almost to dust within a few months, before the wood becomes well seasoned.

WOODS PRINCIPALLY USED FOR CABINS, RUSTIC WORK, AND CERTAIN WOOD PRODUCTS

The woods mainly used in the construction of log cabins and the rustic work in summer homes, furniture, and fences are spruce, fir, hemlock, tamarack, cedar, juniper, pine, birch, poplar, hickory, and

oak. The selection of the wood to be used for these purposes is often determined to a large extent by the local supply. Hickory, ash, dogwood, and persimmon, which are used in the manufacture of shuttle blocks, mallets, mauls, etc., are cut under very similar conditions and thus are subject to similar attack, and the protective measures described in this bulletin are equally applicable to stock cut for such purposes. Those woods which are probably most subject to insect attack and subsequent damage, if cut during certain seasons of the year, as explained on pages 14 and 15, are hickory, pine, persimmon, spruce, ash, and dogwood, although the others mentioned are by no means immune. The damage can be largely prevented if proper measures are taken.

From the standpoint of durability cedar, juniper, and white oak are probably the most desirable of the woods listed above, especially where the wood comes in contact with the ground.¹

FIGURE 1.—An adult bark beetle, *Ips avulsus* Eich.
Enlarged 18 diameters

CLASSES OF INSECTS RESPONSIBLE AND THE DAMAGE THEY DO

The insects responsible for damage, in cases where the bark remains on logs used in rustic work, can be conveniently grouped into three main types or classes, according to their structure and the character of their work, namely, bark beetles,² ambrosia beetles,² and wood borers.³ The bark beetles confine their activities to the bark and burrow between it and the sapwood, often scoring the latter, the ambrosia beetles bore directly through the bark into the sapwood and sometimes the heartwood, and the wood borers may penetrate all three parts of the log.

All three classes of these insects have four distinct stages, namely, the egg; the larva, also called worm or grub; the pupa, or transforming stage; and the adult, or beetle stage. Only the larval and adult stages are responsible for damage to woodwork.

In all instances the larvae hatch from eggs laid by the beetles. After the larva becomes mature it makes a cell in which to rest, called the pupal cell, and there becomes a beetle.

BARK BEETLES

The adult insects of the first class, the bark beetles, are short, cylindrical, reddish-brown to black insects, varying in length from about one-sixteenth to one-fourth inch. (Fig. 1.) They bore

¹ The United States Forest Service has conducted extensive tests and made detailed studies of the durability and chemical preservation of wood and should be consulted if further information of this character is desired.

² Order Coleoptera, family Scolytidae.

³ Order Coleoptera, families Cerambycidae, Bostrichidae, and Buprestidae.

through the outer bark to its soft inner portion, called the phloem, where they make tunnels of various types, along the sides of which they lay their eggs. (Fig. 2.) It is while the beetles are boring through the bark and constructing their tunnels that they push out to the entrances the fine brownish-white sawdustlike particles, or frass, which falls on the wood below and attracts attention by its unsightly appearance, and it is this tunneling that causes the bark

FIGURE 2.—Tunnels of adults and larval galleries of a bark beetle, *Leperisinus aculeatus* Say, beneath the bark of ash. Natural size. The frass that was in the galleries of this species adhered to the bark when it was removed

to loosen and fall off. The piles of frass on a log enable one readily to detect the presence of bark beetles within it. The larvae or grubs are tiny, whitish, cylindrical, slightly curved, legless creatures. (Fig. 3.) Upon hatching from the eggs, they extend their mines in all directions, quite often at right angles to the original (parent) tunnel, thereby aiding in the loosening of the bark. The galleries often contain borings or frass. This depends upon the species of bark beetle.

AMBROSIA BEETLES

The ambrosia beetles (fig. 4) or pinhole borers, which are the adults of the second class, closely resemble the bark beetles in general appearance. Their work differs considerably, however, and can easily be distinguished. After the ambrosia beetles enter the bark they bore immediately into the sapwood and sometimes into the

FIGURE 3.—Larva or grub of a pine bark beetle (*Ips* sp.). Enlarged 10 diameters

heartwood, where they extend their galleries in all directions, each making a hole about the size of a pinhead. (Fig. 5.) While boring these tunnels they push out sawdustlike particles, which either fall out loosely in piles (fig. 6), or come out in stringlike masses (fig. 7) as if being squeezed from a tube. These borings are white, whereas those of the bark beetles are usually brownish and fall loosely from the holes in the bark. In cer-

tain cases much of it, however, remains in the tunnels. The galleries of the ambrosia beetles are round, always free from borings, and quite often their walls are stained black.

Their food is not the wood, but a substance called "ambrosia," which is a coating formed of a minute fungus that is propagated by the beetles themselves. It is this that stains the walls of their galleries.⁴ The damage by these borers is caused almost entirely by the adult beetles, as the larvae, until mature, stay either in the original gallery or in separate cells where they are in most cases cared for and fed by the beetles.

Beetles of this group do not often cause much damage to wood after it has been put into log cabins or rustic work if the trees are cut in the fall of the year, because then it has seasoned by the time the beetles are flying and is too dry to be in a suitable condition for them to work in it. If the wood is cut green, however, and is used during the period of insect activity, it may be attacked immediately by the beetles, in which case a considerable quantity of boring dust will be exuded.

FIGURE 4.—An adult ambrosia beetle (*Xyleborus* sp.). Enlarged about 14 diameters

WOOD BORERS

The adults of the wood borers do not as a group present so uniform an appearance as do the bark beetles and ambrosia beetles. The

⁴ HUBBARD, H. G. AMBROSIA BEETLES. U. S. Dept. Agr. Yearbook 1896: 421-430, illus. 1897.

FIGURE 5.—Work of adult ambrosia beetles (*Xylotcrus* sp.) in pine. Two-thirds natural size

FIGURE 6.—Piles of boring dust or frass being pushed out as ambrosia beetles (*Gnathotrichus* sp.) are entering a hickory log. About one-third natural size

wood borers may be divided into three subgroups, namely, powder-post beetles, roundheaded borers, and flat-headed borers.

POWDER-POST BEETLES

FIGURE 7.—Boring dust or frass which certain species of ambrosia beetles (probably a species of *Xyloterus*) push out of the wood in cylindrical casts. About natural size

about $1\frac{1}{2}$ inches on each side of the entrance hole, although in some cases it may extend in one direction only. The injury can readily be detected by the loose, whitish, dustlike borings which may be found on the bark below the circular entrance hole of the parent gallery.

The larva or grub of this beetle has a curved form, somewhat similar in shape to that of the cutworms found in the garden. When it is doubled up the head of the grub is close to the end of the abdomen. (Fig. 10.) This beetle, in marked contrast to the ambrosia beetles, is exceedingly destructive to the wood, both in its adult and larval stages. In pieces of wood of small diameter it often extends its gallery or tunnel completely around the section, thus

The powder-post beetles are short, cylindrical, reddish-brown to black, hard-shelled insects, ranging in length from one-eighth to nearly one-half inch.

In the eastern section of the United States there is only one species⁵ that causes much damage to the class of materials under consideration. This beetle has reddish-brown markings and is about one-fourth of an inch long. (Fig. 8.) It is commonly known as the "red-headed shot-hole borer" and was mentioned years ago by C. V. Riley as doing considerable damage. It works principally in the wood of the hickory and persimmon, although it has been found in other woods. The adult bores through the bark and into the sapwood, making a cylindrical tunnel around the log just under the surface of the wood and at right angles to the grain. (Fig. 9.) The pores or cells of the wood are opened up, and the female beetle can then insert her eggs into them. The original or egg tunnel is usually bored

FIGURE 8.—An adult powder-post beetle, *Xylobiops basillare*. Enlarged 12 diameters

⁵ *Xylobiops basillare* Say.

greatly weakening it and causing it to break off readily. The larvae also extend their work to the pith in such pieces, completely destroying it and often literally reducing it to powder. In the larger pieces of wood this insect confines its work more to the sapwood, and by the time the larvae are full grown they have destroyed nearly all of the sapwood. (Fig. 11.) The work of the larval borers is entirely in the interior of the log and can not be detected without chopping into the wood. The borings are of the same texture as those of the adult ambrosia beetles but are packed tightly in the mines behind the larvae. When the borings are loosened they break up into cakes. Occasionally the injury is not noticed until the wood has been worked up into the finished product (fig. 12) and the emerging beetles leave holes that indicate the destruction which has been going on unnoticed inside of the manufactured article.

Hickory and persimmon woods used in the manufacture of shuttle blocks, mallets, mauls, etc., are liable to severe injury by this insect, sometimes suffering a 50 per cent loss. (Figs. 11 and 12.)

ROUNDHEADED BORERS

The roundheaded borers have a remarkable variation in general appearance.

(Fig. 13.) The beetles range in length from less than one-fourth inch to more than $1\frac{1}{2}$ inches. They also vary considerably in the place and the manner of laying their eggs. The adult beetles lay their eggs (1) on top of the bark⁶ (fig. 14), or (2), if the bark is

FIGURE 9.—Piece of hickory with the bark and a little of the wood removed, revealing tunnel of an adult powder-post beetle, *Xylobiops basilaris*. Entrance hole is shown at tip of arrow, and the gallery is seen extending on each side. Larvae, pupae, and emerging adults are shown in their cells. Natural size

⁶ *Chion cinctus* Drury.

removed, occasionally on the sapwood, over which they secrete a covering, or (3) in crevices under the bark⁷ (fig. 15), or (4) through slits or pits⁸ which they gnaw in it. (Fig. 16.)

The larvae (fig. 17) which hatch from these eggs are entirely responsible for the damage caused to the wood. They are elongate, fleshy, yellowish-white grubs, usually slightly tapering toward the tail end. Upon hatching, they bore into the soft layers of the inner bark, or phloem, which they begin to mine. Some species⁹ confine much of their activity to mining under the bark, thereby loosening it and causing it to fall off, and they also mine in the outer sapwood. (Fig. 18.) Others make large oval mines which extend deeper into the sapwood¹⁰ and heartwood.¹¹ Sometimes the grubs completely riddle the wood within a few months, although one species¹¹ has entirely different habits and may continue its destructive work in the

seasoned wood over a period of several years. As these galleries are often one-half inch wide and as much as 2 feet long, they weaken the material as well as cause an unsightly condition because of the large quantity of boring dust which many of the grubs push to the outside during the process of excavation. (Fig. 19.)

FIGURE 10.—Larva or grub of the powder-post beetle *Xylobius basillare*. It is in this stage that it does most of its injurious boring. Enlarged 9 diameters

Each species differs somewhat from others in the method of preparing its gallery.

Some grubs pack the borings, or frass, tightly behind them (fig. 11), while others push it out through the entrance hole (fig. 19). The composition of these borings varies from fine, white, and powdery material (fig. 11) to coarse, brownish particles (fig. 19) or shreds of wood fiber (fig. 20). These characters, combined with the species of wood they select to work in, make it possible to determine the type of injury, and in most instances to identify the specific insect responsible for the damage, since each species makes a different pattern while scoring the wood under the bark or gallery of a different shape or size while working in the wood.

FLAT-HEADED BORERS

The beetles of the group known as flat-headed borers are more uniform in general appearance than are those of the roundheaded borers. They are slightly flattened, metallic-colored, boat-shaped beetles which range in length from one-fourth inch to nearly 1 1/4 inches. (Fig. 21.)

⁷ *Neoclytus* sp., *Cyllene caryae* Gahan, etc.

⁸ *Monochamus* sp.

⁹ *Xylotrechus colonus* Fab. and *Callidium antennatum* Newm.

¹⁰ *Neoclytus* sp., *Monochamus* sp., *Cyllene caryae* Gahan.

¹¹ *Hylotrupes bajulus* L.

FIGURE 11.—Cross section of persimmon log, showing the wood destroyed by grubs of the powder-post beetle *Xylobiops basillare*. Natural size

The eggs are laid singly or in a mass on the bark or in crevices in the bark or wood. The larva is an elongate, depressed grub having a characteristic flat head often wider than the rest of the body. (Fig. 22.) The young borer mines the inner bark or the wood, making a flattened, oval, more or less tortuous mine or wormhole which, when completed, widens into a large pupal or resting cell. This connects with the outer surface by a short, oval exit hole through which the new beetle emerges after the larva has transformed and matured in the pupal cell. The larval mines may be in either the bark or wood alone, or, as in many instances, extend throughout the entire piece,¹² and are filled with tightly packed sawdustlike material. (Fig. 23.)

FIGURE 12.—Cross section of hickory handle, showing damage caused by grubs of the powder-post beetle *Xylobiops basillare*. Some of the tightly packed powderlike borings were removed from near the surface to show more clearly the extent of the damage. About actual size

¹² For further and more detailed information, see BURKE, H. E. FLAT-HEADED BORERS AFFECTING FOREST TREES IN THE UNITED STATES. U. S. Dept. Agr. Bull. 437, 8 p., illus. 1917.

FIGURE 13.—Adult beetle of southern pine sawyer, *Monochamus titillator* Fab.
Enlarged $2\frac{1}{2}$ diameters

FIGURE 14.—Adult female beetle and egg of *Chion cinctus* on a section of hickory.
About natural size

SEASONAL ACTIVITY OF THE INSECTS

Some, at least, of the insects of the kinds previously mentioned are at work during the entire growing season, which lasts from April

FIGURE 15.—Piece of log with bark removed to show eggs of white painted hickory borer, *Cyllene caryac*, which were inserted through crevices. Enlarged $3\frac{1}{2}$ diameters

to October in the vicinity of Washington, D. C., from February to December in the extreme Southern States, and from May to September in the Northeastern States. During this time beetles are active, looking for favorable woods on which to lay their eggs. During the rest of the year, however, certain species of these insects are relatively inactive and only a very few or no adult beetles are present.

The bark beetles and ambrosia beetles are active throughout nearly the entire period within the months specified although their numbers vary a great deal within the season of activity. Insects of both the bark-beetle and ambrosia-beetle types mature very rapidly and can develop from the egg to the adult stage in from four to six weeks. They may have as many as three and often five generations a year in certain parts of the South.

FIGURE 16.—Egg scan made by southern pine sawyer, *Monochamus titillator*, on bark of pine. The eggs are inserted through this opening. About natural size

The wood borers are not active for so long a period during the season as are the other two classes of beetles and do not multiply so rapidly. Many of them have one generation a year, and others have only one every two years.

FIGURE 17.—Larva or grub of a roundheaded borer, *Megacyllene antennatus* White. Enlarged 3 diameters

The powder-post beetle that is most destructive flies in the vicinity of Washington, D. C., mainly from the early part of May to the middle of June, the maximum emergence taking place between May 30 and June 8. Occasionally a partial second generation occurs, and

FIGURE 18.—Larval mining of a roundheaded borer, *Calidius antennatus*, which causes the bark to fall off: A, outside of the piece; B, inner mines of the borer. One-half natural size

a few beetles emerge during the warm days of late summer and fall and may be seen in flight from August 15 until cold weather comes. The main brood survive the winter in the mature larval stage and do not emerge as beetles until the following May. Seasonal differences affect the earliness or lateness of the flight period. During the hottest part of the summer it is

possible for the beetles to develop from eggs in from 60 to 70 days.

The beetles of the roundheaded borers usually fly only a few weeks during the spring and summer months, in the vicinity of Washington, D. C., where they have only one generation a year. Farther

FIGURE 19.—Boring dust exuded by the grub of *Cyllene caryae* as it mined in the sapwood of hickory. About one-sixth natural size

FIGURE 20.—Coarse shredded frass made by the grubs of the southern pine sawyer, *Monochamus titillator*, a roundheaded borer. Enlarged about 2 diameters

some species have two generations and sometimes a partial third.

The flat-headed borers, as beetles, fly about the same time that those of the roundheaded borers appear. Some species mature in a year; others take as long as two years.

CONDITIONS FAVORABLE AND UNFAVORABLE FOR ATTACK

The kind and condition of wood attacked are largely dependent upon the species of insect. Some species prefer freshly cut wood that is in a moist condition, others partly seasoned wood, and still others require wood that is dry and well seasoned.

FIGURE 21.—Adult beetle of a flat-headed borer, *Chrysobothris octocola* Lec. Enlarged 4 diameters

CONDITIONS FAVORABLE FOR BARK BEETLES AND AMBROSIA BEETLES

Two types of insects, the bark beetles and the ambrosia beetles, attack freshly cut logs on which the bark remains and in which the inner bark is still white and active and the sap is still present. The ambrosia beetles, however, like equally well logs, with or without bark, that have been submerged in water and which after being removed from the water remain in a moist condition, as when they are placed on the ground in the shade. Such logs may continue to be attacked until they dry out. Wood that is cut during the spring and summer months, when the weather is warm and damp, may be subject to severe injury, especially when placed in close piles on the ground so that it does not receive adequate ventilation. Wood which is cut in the early fall and dried out, or seasoned sufficiently during the winter months, is in an unfavourable condition for attack by bark beetles and ambrosia beetles when their active season arrives.

CONDITIONS FAVORABLE FOR POWDER-POST BEETLES

The powder-post beetles prefer wood that has been cut several months. They show a decided preference for wood that is cut either in the fall and slowly seasoned over winter, or that which is cut during the active season and dried rapidly. Occasionally they attack recently cut wood, but as a rule they are unsuccessful in establishing themselves in it.

CONDITIONS FAVORABLE FOR ROUNDHEADED BORERS AND FLAT-HEADED BORERS

Conditions that invite attack by roundheaded and flat-headed borers vary considerably. Many borers prefer recently cut logs, whereas other borers attack logs which have been seasoned for several months. The manner of handling the logs after they are cut has a decided influence upon whether they will ultimately be attacked. As each species of these beetles flies and lays its eggs during only a short period, and as each kind attacks only one or two kinds of wood, which must be in just the right condition to attract it, the danger of attack by any particular kind is relatively small.

PREVENTIVE AND CONTROL MEASURES

Methods of preventing insect attack and the checking of subsequent injury, once the log is infested, depend largely upon the seasonal history and habits of the insects involved. By taking advantage of what is known of these, the desired protection can be obtained.

PREVENTION OF ATTACK

SEASONAL CUTTING FOR RUSTIC STRUCTURES

The logs with the bark remaining on them and poles used in the construction of rustic cabins, summer houses, fences, etc., should be cut in October or November and piled

FIGURE 22.—Grubs of the flat-headed borer *Chalcophora angulicollis* Lec. Enlarged 2 diameters

at once,¹³ either off the ground or under cover, in such a manner as to offer the best facilities for the rapid and thorough drying of the inner bark before the beetles begin to fly in the spring.¹⁴ This will in almost every case prevent their being damaged by the insects that prefer the freshly cut wood.

FOR MANUFACTURED PRODUCTS

To prevent injury to poles used in the manufacture of rustic furniture, shuttle blocks, mallets, and mauls, the wood should be handled in the following manner:

Restrict the cutting, as previously recommended, to the late fall and winter months, and either utilize the wood before the first flight of beetles occurs in the spring or place it under a closed cover where it will be protected from insect attack. In some cases screening may have to be resorted to, screen cloth having 18 meshes to the inch being used.

If cutting is necessary during the spring and summer months, remove the poles from the forest as fast as they are cut and utilize them at once. A few days' exposure in the forest at this time is sufficient

FIGURE 23.—Work of the flat-headed eastern hemlock bark borer, *Melanophila fulvoguttata* Harr. A, section of hemlock bark showing larval mines in the inner portion, one-half natural size; B, larva; C, adult, natural size. (Burke)

for infestation to occur which may develop after the wood has been manufactured into the finished product. (Fig. 12.) If the poles can not be moved at once, they can be protected for several days by being sprayed with a solution of creosote and kerosene as hereafter described.

Avoid leaving about the factory yards, during the period of insect activity, waste material on which the bark remains, as it is likely to serve as a breeding place for the insects.

¹³ Juniper, or cedar, trees can be felled during August with very little danger of attack if the tops are removed and the poles are laid singly on the ground and turned over once a week for about three weeks to expose a fresh surface to the sun.

¹⁴ HOPKINS, A. D. INSECT INJURIES TO FOREST PRODUCTS. U. S. Dept. Agr., Bur. Ent. Circ. 128, 9 p. 1910.

TREATING WITH CHEMICALS

Certain powder-post¹⁵ and roundheaded¹⁶ borers can not always be readily controlled by such measures, and to obtain the maximum protection the wood used in the construction of rustic cabins, summer houses, etc., should be treated in the spring, before the first flight of the beetles occurs, with either crude pyridine or coal-tar creosote. These liquids are effective when diluted with kerosene in the proportion of one part of pyridine or coal-tar creosote to three parts of kerosene. Methods of applying these liquids are discussed on pages 18 and 19.

Whenever it is necessary to cut trees for these purposes during the spring and summer months, while the insects are active, and it is desired to retain the bark on the logs, the logs should be treated, as soon as cut, with one of the chemicals just mentioned. Where a slight staining of the wood is objectionable, the same protection can be obtained by carefully removing the bark in sections, applying the solutions to the sapwood, and then replacing the bark, using large-headed nails to fasten the bark in place. One nail to each square foot is usually sufficient for this purpose.

Pyridine.—Crude pyridine is a colorless, nearly stainless liquid, and possesses a strong disagreeable odor. Because of this odor, logs or furniture should be treated in the open some distance from any dwelling. If it is necessary to apply this chemical inside a building, the doors and windows should be opened so as to allow ventilation. A windy day would be preferable. Pyridine is inflammable, and reasonable care should be taken in handling it, especially where the wood treated is overhead (rafters, etc.), since the liquid dripping down might burn the skin slightly and be especially painful if it came in contact with the eyes.

Creosote.—Coal-tar creosote (grade 1, liquid oil) is a dark-brown liquid which stains the bark deeply when applied full strength. When diluted with three parts of kerosene, however, the bark is stained only slightly, and this gives a rather pleasing effect. The odor is not very strong when the creosote is diluted with kerosene. Before it is used it should be strained through burlap, and only a good grade of creosote should be used, the kind recommended above; otherwise the wood may be stained darker than is desirable.

PEELING THE POLES

Where it is not especially desirable to retain the bark, a very pleasing effect can be obtained by peeling the poles and treating them with creosote and kerosene in the proportions suggested above. The slight stain thus given is just deep enough to give to a cabin a rustic effect which usually is agreeable. The advantage of this method is that the wood can be cut at any time of the year and yet will not be injured by insects. Furthermore, the creosote aids in preserving the wood. In certain localities, peeling the poles and logs is quite

¹⁵ *Xylobiops basillare*.

¹⁶ *Callidium antennatum* in pine, *Callidium janthinum* Lec. in cedar, and *Chion cinctus* in hickory and oak.

popular. The peeled poles, however, should be kept off the ground for several days or until they have had a chance partially to season so that they will not be attractive to pinhole borers. Peeled poles thus seasoned will not need treatment with chemicals to protect them from insects.

TREATMENT AFTER ATTACK

KILLING THE INSECTS WITH CHEMICALS

Since it is not always possible to prevent injury by taking the precautions just described and because the greater part of the damage is often done before its discovery, investigations were undertaken to find a substance which will serve as an effective remedy. The results of this investigation proved that both liquid orthodichlorobenzene¹⁷ and crystalline paradichlorobenzene are effective.

Crude orthodichlorobenzene.—The crude orthodichlorobenzene product is a colorless, stainless liquid and possesses a slight odor. It is noninflammable, but it is slightly poisonous, and the odor might cause a headache if the liquid were handled for an hour or so at a time. The same precautions should be taken in handling it that are recommended in connection with handling pyridine. It should be applied full strength.

Paradichlorobenzene.—Paradichlorobenzene, which has recently proved effective in the control of the peach-tree borer,¹⁸ was found most effective when prepared, as for the preventive treatments, by being dissolved in three parts, by weight, of kerosene. Care should be taken to dissolve all of the crystals. It is noninflammable and only very slightly poisonous. Both of these solutions slowly liberate a gas which kills the insects. This chemical, like orthodichlorobenzene, may destroy the finish on furniture and necessitate revarnishing. If the infested wood is not treated it will eventually be destroyed.

SUGGESTIONS FOR APPLYING THE CHEMICALS

Wood, before being utilized, can be treated easily and efficiently by being dipped into the solution so that it will penetrate all cracks, crevices, and bark sufficiently to reach the insects and kill them. A trough can be made by bending up the sides and ends of a sheet of some metal, such as galvanized iron, to conform roughly to the shape of the log to be treated. The solution should be poured into the trough and the log immersed. A piece of cord or rope placed under the log will aid in rolling it so that all surfaces will be covered in the course of treatment. After being treated, the wood should be laid in a sunny place to dry.

¹⁷ The chemical purchased under this name (also called orthodichlorobenzol) and tested as such against insect attack, as well as most other products on the market to-day which are being sold for this purpose in the place of orthodichlorobenzene, is mainly a liquid which might be called a crude chlorinated benzol product and is not the pure orthodichlorobenzene. The crude product is manufactured by several concerns, each using its own method, and is being sold under different trade names. It is usually composed largely of monochlorobenzene and ortho-, meta-, and paradichlorobenzene products with the greater part of ortho in it. The pure orthodichlorobenzene is manufactured in limited quantities at the present time and is very expensive; in fact, too expensive to be used for this purpose.

¹⁸ BLAKESLEE, E. B. USE OF TOXIC GASES AS A POSSIBLE MEANS OF CONTROL OF THE PEACH-TREE BORER. U. S. Dept. Agr. Bul. 796, 23 p., illus. 1919.

SNAPP, O. L., and ALDEN, C. H. PARADICHLOROBENZENE EXPERIMENTS IN THE SOUTH FOR PEACH-BORER CONTROL. U. S. Dept. Agr. Tech. Bul. 58, 40 p., illus. 1928.

Wood in use can be conveniently treated by applying a liberal quantity of one of the foregoing chemicals,¹⁹ either with a good spraying apparatus or with a brush. To insure the best results it is important to treat thoroughly all parts of the wood. One gallon of the solution of paradichlorobenzene in kerosene or of the crude orthodichlorobenzene is sufficient to treat five logs, each 10 feet long and 4 inches in diameter, or approximately 50 square feet of bark surface. Two gallons will saturate approximately 100 square feet of wood surface. A pint is usually sufficient to treat a rustic chair of ordinary size.

KILLING THE INSECTS WITH HEAT

In the manufacture of rustic furniture it is customary to steam the wood prior to bending it, especially the older pieces of small-dimension hickory. Quite often the material is found to be infested by the previously mentioned types of borers, and it is usually supposed that by leaving the wood in water, through which steam is being passed, for a period of 15 to 30 minutes, the borers will be killed. An examination of material in a factory where the foregoing method was used in the manufacture of rustic furniture from hickory demonstrated that only a very small percentage of the grubs are killed in this short period of time.

Results of tests²⁰ have shown that it is necessary to subject infested hickory and ash to kiln temperatures of 125° to 130° F. for a minimum period of 1½ to 2 hours after the wood has been brought up to the initial temperature in order to kill all of the grubs in wood 1 inch thick.

Fisher,²¹ in verifying the foregoing experiments for the control of *Lyctus* beetles, determined the period of time necessary for wood of various thicknesses to reach this initial temperature, when infested ash and oak are placed in a cold kiln and raised to the lethal temperature of 130° F. In addition to this period a safety factor of one-half hour is added. Following this the wood is subjected to live steam for 1½ to 2 hours in a saturated atmosphere. The schedule is as follows:

Thickness of timber	Temperature required	Time required to overcome lag after kiln has attained 130° F.	Additional safety factor	Time then held at 130° F.	Total period in kiln at 130° F.
	° F.	Hours	Hours	Hours	Hours
1 inch.....	130	½	½	1½	2½
2 inches.....	130	2	½	1½	4
2½ inches.....	130	3¼	½	1½	5¼
3 inches.....	130	4½	½	1½	6½

¹⁹ Pyridine, paradichlorobenzene, and the crude orthodichlorobenzene product are manufactured by only a few chemical firms. Names and addresses of these firms will be furnished on application to the Bureau of Entomology, U. S. Department of Agriculture.

²⁰ CRAIGHEAD, F. C., and LOUGHBOROUGH, W. K. TEMPERATURES FATAL TO LARVAE OF THE RED-HEADED ASH BORER AS APPLICABLE TO COMMERCIAL KILN DRYING. Jour. Forestry 19: 250-254, 1921.

SNYDER, T. E., and ST. GEORGE, R. A. DETERMINATION OF TEMPERATURES FATAL TO THE POWDER-POST BEETLES, *LYCTUS PLANICOLLIS* LE CONTE, BY STEAMING INFESTED ASH AND OAK LUMBER IN A KILN. Jour. Agr. Research 28: 1033-1038, illus. 1924.

²¹ FISHER, R. C. *LYCTUS* POWDER-POST BEETLES. Dept. Sci. and Indus Res. [England], For. Prod. Res. Bul. 2, 46 p., illus. 1928.

ORGANIZATION OF THE UNITED STATES DEPARTMENT OF AGRICULTURE

June 21, 1929

<i>Secretary of Agriculture</i> -----	ARTHUR M. HYDE.
<i>Assistant Secretary</i> -----	R. W. DUNLAP.
<i>Director of Scientific Work</i> -----	A. F. WOODS.
<i>Director of Regulatory Work</i> -----	WALTER G. CAMPBELL.
<i>Director of Extension Work</i> -----	C. W. WARBURTON.
<i>Director of Personnel and Business Administration.</i>	W. W. STOCKBERGER.
<i>Director of Information</i> -----	M. S. EISENHOWER.
<i>Solicitor</i> -----	R. W. WILLIAMS.
<i>Weather Bureau</i> -----	CHARLES F. MARVIN, <i>Chief.</i>
<i>Bureau of Animal Industry</i> -----	JOHN R. MOHLER, <i>Chief.</i>
<i>Bureau of Dairy Industry</i> -----	O. E. REED, <i>Chief.</i>
<i>Bureau of Plant Industry</i> -----	WILLIAM A. TAYLOR, <i>Chief.</i>
<i>Forest Service</i> -----	R. Y. STUART, <i>Chief.</i>
<i>Bureau of Chemistry and Soils</i> -----	H. G. KNIGHT, <i>Chief.</i>
<i>Bureau of Entomology</i> -----	C. L. MARLATT, <i>Chief.</i>
<i>Bureau of Biological Survey</i> -----	PAUL G. REDINGTON, <i>Chief.</i>
<i>Bureau of Public Roads</i> -----	THOMAS H. MACDONALD, <i>Chief.</i>
<i>Bureau of Agricultural Economics</i> -----	NILS A. OLSEN, <i>Chief.</i>
<i>Bureau of Home Economics</i> -----	LOUISE STANLEY, <i>Chief.</i>
<i>Plant Quarantine and Control Administration</i> -----	C. L. MARLATT, <i>Chief.</i>
<i>Grain Futures Administration</i> -----	J. W. T. DUVEL, <i>Chief.</i>
<i>Food, Drug, and Insecticide Administration</i> -----	WALTER G. CAMPBELL, <i>Director of Regulatory Work, in Charge.</i>
<i>Office of Experiment Stations</i> -----	E. W. ALLEN, <i>Chief.</i>
<i>Office of Cooperative Extension Work</i> -----	C. B. SMITH, <i>Chief.</i>
<i>Library</i> -----	CLARIBEL R. BARNETT, <i>Librarian.</i>

This bulletin is a contribution from

<i>Bureau of Entomology</i> -----	C. L. MARLATT, <i>Chief.</i>
<i>Division of Forest Insects</i> -----	F. C. CRAIGHEAD, <i>in Charge.</i>

20

ADDITIONAL COPIES
OF THIS PUBLICATION MAY BE PROCURED FROM
THE SUPERINTENDENT OF DOCUMENTS
U. S. GOVERNMENT PRINTING OFFICE
WASHINGTON, D. C.

AT
5 CENTS PER COPY

UNITED STATES DEPARTMENT OF AGRICULTURE

DEPARTMENT CIRCULAR 411

Washington, D. C.

September, 1927

THE RELATION OF INSECTS TO SLASH DISPOSAL

Prepared by the Division of Forest Insect Investigations, Bureau of Entomology¹

CONTENTS

	Page		Page
Slash a factor in forest protection.....	1	Pine slash—Continued.....	
Few species of trees involved.....	3	Monterey pine in California.....	9
Pine slash.....	4	The southern pines.....	9
White pine in the Northeast.....	4	Spruce slash.....	10
Red pine and white pine in the Great Lakes region.....	4	Spruce in the Northeast.....	10
Western white pine in the northern Rocky Mountain region.....	5	Spruce in the Great Lakes region.....	10
Sugar pine in the Pacific coast region.....	6	Engelmann spruce in the Rocky Mountain region.....	10
Western yellow pine.....	6	Douglas fir slash.....	10
Jeffrey pine in California.....	8	Conclusions.....	11
Lodgepole pine.....	8	Literature cited.....	12

SLASH A FACTOR IN FOREST PROTECTION

Does slash from cutting operations increase the insect menace to living timber? Many times this question has been asked by foresters and timbermen, but there has been little uniformity in the answers. Considerable confusion has arisen as a result, and a great diversity of opinion is found concerning the influence of slash upon insect outbreaks.

Two widely divergent opinions are held by professional forest entomologists in North America. In Canada, Swaine (10, p. 18; 11, p. 762; 12, p. 25; 13, p. 47; 14, p. 231; 15, p. 346)² and Hopping (6, 7) advocate on entomological grounds the necessity of slash disposal as a phase of forest protection. Workers in this country, Graham (2, p. 447), for example, almost unanimously take the opposite viewpoint, that under present economic conditions slash disposal can not be justified on entomological grounds. Chamberlin (1, p. 23), however,

¹ This circular was prepared by entomologists and collaborators of the Division of Forest Insect Investigations. Those primarily responsible for the information contained in the regional statements are as follows: F. C. Craighead, northeastern and southeastern forest regions; S. A. Graham, Lake States region; J. C. Evenden, northern Rocky Mountain region; W. D. Edmonston and F. C. Craighead, southern Rocky Mountain region; J. E. Patterson, north Pacific region; J. M. Miller, H. E. Burke, and H. L. Person, California region. F. P. Keen and F. C. Craighead are largely responsible for assembling the paper as a whole. In making reference to this circular it is to be understood that it is a product of joint authorship, as explained in the foregoing.

² Figures in *italic* in parentheses refer to "Literature cited," p. 12.

recommends burning slash of western yellow pine for the control of bark beetles. In England, Munro (8, p. 27), entomologist of the forestry commission, also recommends burning slash for control of bark beetles. It is fully realized, however, that broad statements applying indiscriminately to all forested regions are out of the question; the problem varies with each forest, region, and type. Not only must the species of tree and the character of the slash be considered but the various species of insects that may be involved must be taken into account.

It is recognized that, under special conditions, slash disposal or seasonal timing of cutting operations is advantageous. Such cases are referred to in the general text.

Hopkins discusses this question in many of his papers. His general conclusions are adequately supported by all the detailed investigations so far conducted by the Division of Forest Insect Investigations. As early as 1899 (3, p. 23) he called attention to the wasteful methods of lumbering in the Northwest and the attractive breeding places for forest insects provided by the scarred trunks and the débris left on the ground. Ten years later, in his monograph, (4, p. 25-26, 48), he suggests that the cutting of living timber for commercial purposes may offer favorable conditions for the multiplication of some of the species, although in some cases it may serve as a protection to living timber, and suggests burning the slash after it has attracted beetles away from the living trees. In one of his later papers on forest entomology (5, p. 8) he broadly summarizes his opinions as follows:

Since the problem of slash disposal is on the program of this meeting and, since it is pretty generally considered by foresters and some entomologists that slash is an insect hazard to the living timber, I want to say that the results of more than thirty years of observations and some detailed study of the problem indicate that as a rule, it is not. As in all rules, there are, of course, exceptions to this one. There are a few cases as related to certain types of forests, time of year, and cases of sporadic cutting, where the slash is dangerous, not so much from the insects that breed in the tops, logs, and stumps, as that it serves to attract the tree-killing insects to the locality and from thus being concentrated they attack and kill the living timber. Continued logging operations, after they are once started within a given area, provide continuous breeding places for the insects and their natural enemies and thus the slash serves as a protection to the living timber.

The entomologists of the Division of Forest Insect Investigations feel that a paper setting forth what is now known about the entomological effect of slash would help to clear the situation and stimulate more careful observation and compilation of further and needed data. In this, as in most scientific problems, the last word can never be said, and much investigative work is still needed to clear up many phases of the slash problem. However, considerable work has already been done, and certain tentative conclusions are presented in this circular.

There are hundreds of species of insects which breed in the slash of forest trees, but few of these are capable of doing much injury to living trees. They attack the base, the trunk, the tops, the limbs, or the twigs of mature standing trees; others emerge from slash and attack seedlings, saplings, and poles.

In addition to this, slash has a very strong attractive influence which may result in bringing thousands of insects into the general slash area with a resultant varying damage to living timber.

The direct result of insect attack upon the slash itself is beneficial because the insects aid in bringing about decomposition. In this circular the direct influence of insects upon twigs, limbs, and tops can be ignored and only the role of slash need be considered in (1) supplying breeding material for insects which emerge and kill mature standing timber or seedlings, saplings, and poles, and (2) attracting insects from the surrounding forest and concentrating them in the vicinity of the slash, where they kill living trees.

In general, the insects with which this discussion is concerned breed in the same parts of a fallen tree as of a living tree. Insects which attack the base of standing trees breed in stumps, those attacking the main trunk breed in cull logs, and those attacking limbs and twigs breed in the limbs and twigs composing the smaller portions of slash. There are, however, some important exceptions to this rule, such as certain species of Ips.

Probably the heaviest losses of all are attributable to the effect of slash in attracting beetles from the surrounding territory and concentrating them in the vicinity, where they are able to do greater damage to the standing timber than if their attacks had been scattered over a wider area.

This attractive influence has been noted many times. The felling of trap trees³ has often concentrated attacks on adjacent healthy trees, even in some cases in which the felled trap tree has not been attacked. The beetle population is usually attracted to fire-scorched areas, even though the beetles are often unable to breed successfully in the trees weakened by fire. It is also usually attracted to places where heavy windfalls have occurred or to trees that have been broken down by snow.

It is evident that this class of damage can not be minimized through slash disposal. In fact, burning the slash in the summer, which is the flight period of the beetles, may even increase the attractive influence and bring about a still greater concentration.

FEW SPECIES OF TREES INVOLVED

The problem can be further simplified by limiting the discussion to such trees as are subject to primary damage from slash-breeding insects. Relatively few trees are thus involved.

All hardwoods can be eliminated from the discussion, as practically no insects which breed in hardwood slash are of importance as tree killers. Among the conifers, the slash of fir (*Abies*), larch (*Larix*), redwood (*Sequoia*), hemlock (*Tsuga*), cypress (*Cupressus*, *Taxodium*), cedar (*Thuja*, *Libocedrus*, *Chamaecyparis*), and juniper (*Juniperus*) either breeds insects of very little significance as tree killers, or the trees killed are so few or of so little value as to be of small economic importance.

As a result of these eliminations the problem in the United States, viewed from an economic standpoint under present forestry practices, resolves itself into a consideration of slash from pine (*Pinus*), spruce (*Picea*), and Douglas fir (*Pseudotsuga*). Each of these three groups will be taken up in sequence.

³ Living trees girdled or felled at the proper time to attract the flying beetles to them and away from healthy trees.

PINE SLASH

The pines suffer more than any other group from the attack of bark beetles. It is not surprising, therefore, that pine slash should be viewed with the greatest suspicion as a forest menace. Upon study, however, the danger loses much of its alarming aspect.

WHITE PINE IN THE NORTHEAST

The only really serious bark-beetle enemy of the white pine (*Pinus strobus*) which breeds in slash in the Northeast is the Pales weevil (*Hyllobius pales* Herbst). Studies by Peirson (9) indicate not only that this beetle breeds in stumps and cull logs but that the adult is also strongly attracted into slash areas for the purpose of feeding. It kills great quantities of small reproduction by gnawing off the bark until girdling results, and is frequently the limiting factor in obtaining white-pine regeneration.

No satisfactory method of control has yet been devised. Burning the bark of cull logs and thoroughly charring the stumps, or peeling the bark of logs and stumps (both of which are difficult and expensive processes), may lessen the damage to some extent, but will have no effect in lessening the attractive influence of the slash, which is, after all, probably the greatest source of danger. R. T. Fisher, in his experimental silviculture on the Harvard forest, has found that the only practical methods of preventing serious losses are (1) obtaining such an abundance of reproduction through reproduction cuttings that there are still enough seedlings left to produce a stand after the destruction wrought by the weevil, and (2) clear cutting, and subsequent planting after an interval of three years.

RED PINE AND WHITE PINE IN THE GREAT LAKES REGION

Graham (2) reports that insect species that normally breed in the slash of the red pine (*Pinus resinosa*) and white pine (*P. strobus*) in the Great Lakes region have seldom been observed attacking or killing healthy trees. The insects responsible for the few exceptions to this rule are all found breeding in the stumps and larger parts of the slash.

The bark beetle *Dendroctonus valens* Lec. May occasionally kill trees, and is one of the most common of the species that breed in fresh pine stumps. It is almost never found breeding in parts of the tree more than 6 or 8 feet above the ground. All recorded cases of injury caused by this beetle to a standing tree have been associated with the attractive influence of logging operations, windfalls, fires, or logs in storage.

Several species of *Ips* occasionally kill trees and these, like the *Dendroctonus* beetles, usually breed most abundantly in the larger parts of the slash. The most common offenders of this group are *Ips pini* Say, a primary pest of red and white pines, and *I. calligraphus* Germar, which breeds in white pine. In the light of present knowledge it is safe to say that in the Lake States only the larger parts of the trees constituting pine slash have any entomological significance so far as living standing trees are concerned. Just how important an entomological factor these parts are can not be said, but it is certain that the stumps provide the most favorable breeding places.

The ordinary method of slash disposal, by burning, destroys only the small, entomologically innocuous parts of the slash, and leaves the parts most favorable for insect breeding. Any method of slash disposal that will reduce the number of insects must take care of the larger material. Even burning slash piled over stumps does not as a rule sufficiently char the stump to prevent subsequent insect infestation. Thorough charring can be accomplished only by keeping the fire close to the sides of the stump throughout the process of burning. Thorough charring of stumps is too expensive to be practical. Bark-ing is cheaper, but must be carried down to the mineral soil if satisfactory results are to be expected.

Unpublished results of experiments in Minnesota indicate that heavy shade is unfavorable to most potentially injurious insect species. In one experiment, piling the small parts of the slash over the stumps very materially reduced the infestation. This experiment was on a small scale and only about 20 stumps were used, but the results were very clear cut. Pairs of stumps close together were selected; one of each pair was covered with the smaller parts of the slash, the other stump being left uncovered. The uncovered stumps were found to contain an average of 40 mother tunnels of *Dendroctonus valens* per stump, whereas the covered stumps averaged only 4 each.

From the data at hand the writers are forced to conclude that slash burning in the Lake States is unnecessary from the entomological point of view. Covering the stumps and larger pieces of the slash with the smaller pieces promises satisfactory results at a minimum cost.

WESTERN WHITE PINE IN THE NORTHERN ROCKY MOUNTAIN REGION

The most serious insect enemy of the western white pine (*Pinus monticola*) is the bark beetle *Dendroctonus monticolae* Hopk. This species breeds successfully in cull logs from logging operations, and also kills living trees. General observations by Evenden indicate that during continuous logging operations the progeny of such beetles as breed in the cull logs are absorbed in the new material made available, so that little damage results to standing trees. Because of the value of this timber species very close utilization is practiced, and ordinarily only a very small quantity of slash over 6 inches in diameter is left in the woods. Occasionally, however, where a certain quantity of such material is left in the woods at the close of a logging operation, and no fresh material is provided, the insects which emerge attack and kill standing trees in the vicinity, or seed trees left on the area occupied by the slash.

As these beetles do not breed in the smaller tops and limbs the present methods of brush disposal by burning can have no possible effect upon their number. And because of the close utilization of the larger material other measures for preventing injury by insects do not ordinarily appear to be justified. If, however, for any reason, any quantity of large slash material is left in the woods and no fresh material is provided to absorb the emerging broods, then as a protective measure the bark should be peeled from the infested cull logs before the beetles emerge; otherwise some damage to standing trees may result.

Two smaller species of insects (*Pityogenes carinulatus* Lec. and *P. knechteli* Sw.) breed in the smaller material from the tops. Occasionally they breed in sufficient numbers to attack and kill small

white pine reproduction. These cases are of such sporadic occurrence and of such minor importance that they are negligible.

SUGAR PINE IN THE PACIFIC COAST REGION

The bark beetle *Dendroctonus monticolae* Hopk. is also the principal enemy of the sugar pine (*Pinus lambertiana*) in the forests of the Pacific coast. As in white pine, it breeds both in cull logs and in standing trees. This species of pine is not, however, subject to heavy epidemics of infestation such as occur in yellow pine, and no heavy killings have yet been traced to the slash of logging operations. On the Yawkey tract, near Fort Klamath, Oreg., in 1922 and 1923, Patterson, Keen, and Hauge noted a considerable increase in the numbers of standing trees killed shortly after many trees were blown down by a high wind. In general, the considerations applying to utilization and slash disposal in the case of white pine apply to this species as well.

WESTERN YELLOW PINE

In California and Oregon the western pine beetle, *Dendroctonus brevicornis* Lec., is the principal insect enemy of the western yellow pine (*Pinus ponderosa*). Besides attacking living trees this insect breeds in the trunks of fallen trees which are more than 6 inches in diameter.

In some cases it has been noted that heavy losses by insects have followed logging operations, whereas in other cases logging areas have been singularly exempt from such injury. Keen reports that on the southern Oregon-northern California project two of the cleanest areas were adjacent to areas where continuous logging had been in progress for several years; that one heavily infested area was adjacent to a logged area where operations had ceased; and that in one unit a reduction of 60 per cent in infestation, equal to the best obtained by artificial control work, followed the beginning of logging operations on adjacent ground.

To determine if possible the reason for this condition, Person carried out some investigations in southern Oregon and central California. Eleven logging slash areas, two road slash areas, and one group of windfalls were studied. By bark counts made on 459 pieces of slash it was found that the average number of attacking beetles per square foot was 9.8, and the number of beetles emerging, 16. Similar studies were made by Patterson in southern Oregon. Five logging areas, one windfall area, and 24 miles of road slash were studied; the last named involved the examination of 1,425 pieces of slash, including the examination of 1 square foot of bark on each of 1,075 trees. Bark counts in this material gave 15.6 beetles per square foot attacking and 19 beetles per square foot emerging. Bark counts made on hundreds of standing trees near by showed an average attack of 23 *Dendroctonus* beetles per square foot, with 54 beetles emerging. It is evident that the attack and emergence on slash material is far below that for standing trees. In all of these studies checks on the surrounding infestation were made before, during, and for several years after the laying down of the slash.

Usually the first effect of the slash was to increase the infestation in standing trees in the vicinity. In every case, however, this increase occurred simultaneously with the attack on the slash, instead of following the emergence of the beetles breeding in the slash.

Numerous observations made by forest entomologists in the Western States point conclusively to the fact that the western pine beetle is strongly attracted to the vicinity of slash material, trees scorched by fire, trap trees, and windfalls. As long ago as 1906, Webb (17, p. 30) pointed out that storm-felled and lightning-struck western yellow pines became centers of infestation by this insect. If a concentration of the beetles is brought about through the presence of some such material, they not only attack this material but go into the standing living trees in the immediate vicinity.

From these investigations the conclusion is reached that the destruction of standing trees by insects, consequent upon logging operations in adjacent territory, is not caused by the breeding up of insects in the slash, but by the concentration in the slash of beetles from the surrounding territory, and the failure of the slash to absorb them all. This strong attractive influence can not well be avoided.

Since the western pine beetle breeds only in the slash material more than 6 inches in diameter, such as cull logs and butts, the present methods of slash disposal can be of little avail in reducing the numbers of beetles. Neither can the very strong attractive influence of the slash be avoided. The best way, therefore, to avoid injury to neighboring standing trees is to supply continuously a quantity of logs to absorb the broods of beetles, and, if necessary, to reduce, through artificial control, the total population of beetles on the area.

Several species of *Ips* (*I. emarginatus* Lec., *I. oregoni* Eich., *I. confusus* Lec., etc.) breed in the smaller pieces of the slash material as well as in some of the larger pieces. When they become sufficiently numerous, if there is a shortage of slash, they may kill young trees, and the tops and limbs of larger trees. This damage is very sporadic, and usually follows the interruption of logging operations in the summer months. Keen has reported three small areas on the North Warner division of the Modoc National Forest, in California, where species of *Ips* have caused heavy losses of reproduction after the cessation of logging operations in the summer months. The conditions were complicated by extremely dry weather. Usually a supply of fresh slash, continuously laid down during the summer months, will absorb all of these beetles and prevent injury to neighboring living trees.

In general it may be said that, in the light of present information, no special methods of slash disposal need be recommended to avoid insect damage to western yellow pine, except in the case of sporadic cutting or when logging operations cease. In such cases infested slash material should be destroyed before the broods of insects emerge. The same general considerations apply to western yellow pine in the northern Rocky Mountain region.

In the southern Rocky Mountain and Colorado Plateau regions the Black Hills beetle (*Dendroctonus ponderosae* Hopk.) is the principal insect enemy of the western yellow pine (*Pinus ponderosa* var. *scopulorum*). Edmonston has reported that in Colorado the forest areas showing least infestation are always those adjacent to places where logging operations are in progress, particularly where the operations have been continuous for several years. Several officials of the Forest Service have reported similar observations. Recent studies by Craighead, Blackman, and Keen in the Kaibab National Forest in Arizona indicate that the felled logs and culls act as traps for the

beetles during the flight period. When the material is sawn, practically all the beetles are destroyed, and in the logs that are left in the woods during the winter a very high mortality of beetles occurs, so that the emerging beetles are much fewer than those attacking. At the same time the cull logs and tops furnish good breeding material for certain borers (*Acanthocinus*) and predatory species which are detrimental to the development of the *Dendroctonus* broods. As these enemies increase with this abundance of breeding material they tend to concentrate on any standing trees which are attacked, and thus hold the infestation at a minimum. Under these conditions it is doubtful whether disposal of the larger slash would not in fact be more injurious than beneficial to neighboring standing timber.

In both the Rocky Mountain and the Pacific regions *Dendroctonus valens* Lec. breeds in stumps and butt logs, and to a limited extent it attacks the base of live trees. This injury is not often severe enough to warrant the expense of special control methods, such as piling brush over stumps, prompt removal of logs, and barking of cull logs.

In Arizona, New Mexico, and south of the Colorado Plateau, some outstanding cases of losses to reproduction resulting from insects breeding in slash have been called to our attention. Craighead reports that two species of *Ips* were found breeding in cull logs and larger tops of western yellow pine on timber sales. The spring generation of beetles is confined to this felled material, whereas the summer generations attack and kill pine reproduction in clumps of from one-tenth to 1 acre. At some places these repeated attacks have materially thinned out the stands of second growth. Modifications of logging operations will be necessary to prevent these losses.

JEFFREY PINE IN CALIFORNIA

The Jeffrey pine beetle (*Dendroctonus jeffreyi* Hopk.), although often a very destructive insect in living Jeffrey pine (*Pinus jeffreyi*) in California, is only weakly attracted to the slash of this tree. This species and other bark beetles, such as *Ips*, breed very freely in windfalls and large pieces of slash material. On large windfall areas they may occasionally breed up to the point where they become very destructive to adjacent full-grown timber, saplings, poles, and the top of mature trees. Such epidemics are, however, characteristically sporadic, and their control is not economically feasible at the present time except in cases where high timber values are involved. As far as injury by insects is concerned Jeffrey pine slash is usually of but slight importance.

LODGEPOLE PINE

The mountain pine beetle (*Dendroctonus monticolae* Hopk.) is a very serious enemy of lodgepole pine (*Pinus murrayana*), often killing hundreds and even thousands of acres of the standing timber. Here is another case, and a very striking one, of an insect which is strongly attracted to the vicinity of logging or clearing operations, yet rarely attacks the felled trees. Clearings for summer homes or hotels have often brought about such a slaughter of lodgepole pines in the vicinity as to mar the beauty of the site. Even control operations are not immune, and in one instance such a heavy concentration of beetles followed control cutting as entirely to wipe out the effect of the work.

Nevertheless, a felled tree is rarely attacked. Here, again, it is obvious that slash disposal does not reduce the number of beetles, and the concentration caused by the slash appears to be unavoidable.

Another lodgepole pine insect (*Dendroctonus murrayanae* Hopk.) has habits similar to those of *D. valens* Lec., discussed under yellow pine. Evenden has found this species of economic importance in crosstie operations in Wyoming. Breeding readily in the stumps, it subsequently attacks the bases of the larger standing trees not included in the sales. Repeated attacks of this kind weaken the tree to such an extent that it succumbs to the attack of other bark beetles on the bole.

MONTEREY PINE IN CALIFORNIA

Several serious outbreaks of *Ips* on the Monterey pine (*Pinus radiata*) near Monterey, Calif., have been accounted for as caused by wood cutting and clearing, where the wood has been left piled in the forest. Other factors, however, enter into this problem, since similar cutting in other years has produced no epidemics. The turpentine beetle (*Dendroctonus valens* Lec.), which breeds usually in stumps, often becomes primary in its attack upon standing trees of this species.

Because of the value of the Monterey pine as an ornamental and shade tree, special precautions in slash disposal may well be taken to help minimize insect damage to it, which occasionally becomes serious.

THE SOUTHERN PINES

The southern pine beetle (*Dendroctonus frontalis* Zimm.) and three species of *Ips* (*I. calligraphus* Germ., *I. avulsus* Eichh., and *I. grandicollis* Eichh.) are the most important pine insects of the South. The first species mentioned rarely attacks anything but living trees, whereas the three species of *Ips* are more commonly found in tops, cull logs, and stumps than in standing pines. All these species are so intimately associated in the destruction of living trees that it is impossible to consider them separately. On the other hand, the infrequency with which the southern pine beetle attacks slash, cull logs, or wind-blown trees immediately eliminates it as an argument for the disposal of such material.

Very few reports of dying pine timber in the South that can be associated with slash as a factor have been called to the writers' attention, and extensive observations by several entomologists have not revealed any striking examples. In fact, experimental cuttings of timber, conducted throughout the year with the express purpose of furnishing the breeding material for these insects, have utterly failed to produce attacks in the surrounding standing trees. Nevertheless, there are many examples which show that these species are all strongly attracted to areas where cutting is being carried on in the summer. If sufficient material is on the ground they invariably attack it, and occasionally small quantities of neighboring living timber are killed. For example, a freshly sawn pile of lumber, brought to a home site for building purposes, attracted to that vicinity a sufficient number of these insects to kill a number of trees adjacent to the pile.

It is therefore reasonable to conclude that slash serving as breeding material for these insects is not a menace, but that because of the attraction which it exerts on them local logging or clearing operations in the summer should be avoided.

SPRUCE SLASH

SPRUCE IN THE NORTHEAST

Several species of bark beetles breed in the stumps, logs, or tops of spruce (*Picea canadensis*, *P. rubens* and *P. mariana*), and attack and kill living trees.

The eastern spruce beetle (*Dendroctonus piceaperda* Hopk.) is a particularly destructive species which breeds in stumps and cull logs. It has destroyed large areas of mature spruce forests in the past, and active epidemics are now in progress in Canada, even in second-growth stands. Recent observations by Craighead indicate that it is most destructive only when accompanied by one or more species of Ips, which combine with it and attack the upper stems.

Several species of Ips (*I. borealis* Sw., *I. perturbatus* Eichh., and *I. chagnoni* Sw.) have in two cases been observed by Craighead breeding in slash, and in the following year attacking and killing near-by living trees. From these, however, scanty broods developed, and in the third year the outbreaks completely died out.

Special methods of slash disposal do not appear to be justified from the entomological standpoint, but cull logs and larger débris should not be left in the woods (16, p. 85). The increasingly closer utilization practiced in this region with the development of the pulp industry will undoubtedly tend to minimize such losses.

SPRUCE IN THE GREAT LAKES REGION

There is no specific evidence to prove that spruce slash in the Lake States has any entomological significance, although there is justification for suspecting that the larger pieces may be of some importance. Most of the spruce (*Picea canadensis* and *P. mariana*) cut in this region finds its way to pulp mills, a fact which makes for close utilization. A comparatively small quantity of material suitable for the breeding of potentially injurious insects is left in the woods after logging operations.

ENGELMANN SPRUCE IN THE ROCKY MOUNTAIN REGION

The Engelmann spruce beetle (*Dendroctonus engelmanni* Hopk.) breeds in the culls and stumps of the Engelmann spruce (*Picea engelmanni*) and also attacks and kills living trees. The small quantity of timber of this species which is now being cut makes of small importance the danger from this source. No opportunity has yet presented itself for the study of the effect of extensive logging on this insect.

DOUGLAS FIR SLASH

There are several insects which breed in the large pieces of slash of Douglas fir (*Pseudotsuga taxifolia*) and are capable of killing living trees, though they rarely do so. The more important of these are *Dendroctonus pseudotsugae* Hopk., *Scolytus unispinosus* Lec., and *Melanophila drummondi* Kirby.

In the region along the Pacific coast reports on observations of the death of Douglas fir from the attack of insects that breed in slash are so infrequent that they can be disregarded. This fact may be due to

the extensive and continuous character of the logging operations. Recently, however, reports from the British Columbia Forest Service indicate heavy killing of Douglas fir adjacent to certain logging operations on the coast of British Columbia.

In the northern Rocky Mountain region conditions are somewhat different. The Douglas fir beetle (*Dendroctonus pseudotsugae* Hopk.) is apparently much more aggressive in these less humid sections. In recent years it has been causing considerable damage, and coming more and more to be considered a serious insect. Evenden believes that several local outbreaks under his observation are clearly attributable to slash which has resulted from logging operations, construction of rustic buildings, spruce-budworm defoliation (resulting in the death of many trees), or windfalls.

Two small bark beetles, *Scolytus unispinosus* Lec., and *Pseudosynaldis nebulosus* (Lec.) occasionally multiply in slash, and when no more material is available they cause damage to reproduction and to young trees.

The question of the destruction of Douglas fir by insects needs much more thorough study and more systematic observations before any conclusions can be ventured.

CONCLUSIONS

The outstanding conclusions which have been reached to date in regard to the entomological aspect of slash disposal are given here.

As far as is now known, pine, spruce, and Douglas fir are the only species of timber to which insects associated with slash are of any important economic significance.

Comparatively few species of insects are capable of adapting themselves to breeding in slash and killing living trees.

The great majority of the important tree-killing insects which attack slash breed largely in the cull logs and butts. It follows that the present methods of brush burning can have little influence on their control.

Where logging operations are continuous, and a constant supply of slash is provided, the potential danger from insects need cause no concern.

Slash attracts from the surrounding forests insects which often concentrate in standing timber in the vicinity of the cutting. No method of slash disposal will avoid this effect.

In special cases where the value of the surrounding timber is greatly enhanced, either from an æsthetic standpoint as a building site or for scientific purposes such as experimental thinnings in sample plots, extreme caution should be exercised in felling green timber. In such cases the attraction of the felled timber for insects is more important than its service as a breeding ground, and all such work should be done in the fall or winter, at least two months before the growing season.

A large number of insects of secondary importance breed in the twigs, limbs, and trunks of fallen trees, and sometimes become so numerous as to kill reproduction, pole stands, and occasionally even mature trees in the vicinity. Such outbreaks are, however, sporadic in nature and of short duration, and, under present economic conditions, do not warrant special methods of slash disposal.

LITERATURE CITED

- (1) CHAMBERLIN, W. J.
1920. THE WESTERN PINE BARK BEETLE. Oreg. Agr. Expt. Sta. Bul. 172, 30 p., illus.
- (2) GRAHAM, S. A.
1922. SOME ENTOMOLOGICAL ASPECTS OF THE SLASH DISPOSAL PROBLEM. Jour. Forestry 20: 437-447.
- (3) HOPKINS, A. D.
1899. PRELIMINARY REPORT ON THE INSECT ENEMIES OF FORESTS IN THE NORTHWEST. U. S. Dept. Agr., Div. Ent. Bul. (n. s.) 21, 27 p.
- (4) ———
1909. PRACTICAL INFORMATION ON THE SCOLYTID BEETLES OF NORTH AMERICAN FORESTS. I. BARKBEETLES OF THE GENUS DENDROCTONUS. U. S. Dept. Agr., Bur. Ent. Bul. 83, pt. 1, 169 p., illus.
- (5) ———
1922. BUDWORM INFESTATION VS. PULPWOOD PRODUCTION. Amer. Paper and Pulp Assoc., Woodlands Sect. Proc. 2: 7-8.
- (6) HOPPING, R.
1915. THE ENTOMOLOGICAL ASPECT OF SLASH DISPOSAL. Soc. Amer. Foresters Proc. 10: 183-185.
- (7) ———
1921. THE CONTROL OF BARK-BEETLE OUTBREAKS IN BRITISH COLUMBIA. Canada Dept. Agr. Ent. Branch Circ. 15, 15 p., illus.
- (8) MUNRO, J. W.
[1920]. SURVEY OF FOREST INSECT CONDITIONS IN THE BRITISH ISLES, 1919. [Gt. Brit.] Forestry Comm. Bul. 2, 35 p., illus.
- (9) PEIRSON, H. B.
1921. THE LIFE HISTORY AND CONTROL OF THE PALES WEEVIL (HYLOBIUS PALES). Harvard Forest Bul. 3, 33 p., illus.
- (10) SWAINE, J. M.
1914. FOREST INSECT CONDITIONS IN BRITISH COLUMBIA. Canada Expt. Farms, Div. Ent. Bul. 7, 43 p., illus.
- (11) ———
1917. SHADE TREE AND FOREST INSECTS IN MANITOBA. Agr. Gaz. Canada 4: 755-763.
- (12) ———
1918. CANADIAN BARK-BEETLES. PART II. Canada Dept. Agr. Ent. Branch Bul. 14, 143 p., illus.
- (13) ———
1919. SOME INSECT INJURIES IN WOODLOTS. Quebec Soc. Protect. Plants Ann. Rpt. (1918/19) 11: 46-48, illus.
- (14) ———
1919. THE BALSAM INJURY IN QUEBEC AND ITS CONTROL. Agr. Gaz. Canada 6: 227-233, illus.
- (15) ———
1921. SPRUCE BUDWORM INJURIES IN EASTERN CANADA. Illus. Canad. Forestry Mag. 17: 345-346, illus.
- (16) ———CRAIGHEAD, F. C., and BAILEY, I. W.
1924. STUDIES ON THE SPRUCE BUDWORM [CACOECIA FUMIFERANA CLEM.] Canada Dept. Agr. Bul. (n. s.) 37, 91 p., illus.
- (17) WEBB, J. L.
1906. SOME INSECTS INJURIOUS TO FORESTS. II. THE WESTERN PINE-DESTROYING BARKBEETLE. U. S. Dept. Agr., Bur. Ent. Bul. 58, pt. 2, p. 17-30, illus.

ADDITIONAL COPIES

OF THIS PUBLICATION MAY BE PROCURED FROM
THE SUPERINTENDENT OF DOCUMENTS
GOVERNMENT PRINTING OFFICE
WASHINGTON, D. C.

AT

5 CENTS PER COPY

▽

U. S. DEPARTMENT OF AGRICULTURE

FARMERS' BULLETIN No. 1472

PREVENTING DAMAGE BY TERMITES OR WHITE ANTS

THROUGHOUT the United States native termites, or white ants, cause serious damage to the foundations and woodwork of buildings and articles in the buildings, as well as to living fruit and other trees, crops, and other vegetation. Such damage can be prevented by the proper construction of buildings and by chemical treatments of wood, and injury to growing vegetation by clean-culture and the use of insecticides. This bulletin describes the habits and activities of subterranean and nonsubterranean termites and sets forth in detail the precautions to be taken against them and the remedies available for damage which they have inflicted.

This bulletin supersedes Farmers' Bulletin 1037, White Ants as Pests in the United States and Methods of Preventing Their Damage.

Washington, D. C.

Issued April, 1926; revised June, 1930

PREVENTING DAMAGE BY TERMITES OR WHITE ANTS

By T. E. SNYDER, *Senior Entomologist, Division of Forest Insects, Bureau of Entomology*

CONTENTS

	Page		Page
Termites and their habits-----	1	Preventing and remedying damage to	
Distribution and forms-----	1	woodwork of buildings—Contd.	
Location of colonies or nests-----	2	Killing the winged adults will not	
Termites which are subterranean in		stop the damage-----	16
habit-----	2	Disconnecting wood from the	
Termites which are nonsubterranean		ground-----	17
in habit-----	3	Replacing with concrete-----	17
The colonizing swarm-----	3	Replacing with metal-----	17
The reproductive forms-----	5	Spraying and poisoning-----	17
Preventing and remedying damage to		Preventing and remedying injury to	
woodwork of buildings-----	6	living vegetation-----	18
Destruction of breeding places about		Fruit, nut, shade, and forest trees--	18
the building site-----	6	Young plantations or nursery stock--	18
Proper construction of buildings-----	6	Vineyards-----	19
Modifications of city building codes--	15	Field and truck crops-----	19
Locating and temporarily arresting		Flowers and greenhouse stock-----	20
the damage-----	15	Summary -----	21
Indications of infestation-----	16		

TERMITES AND THEIR HABITS

DISTRIBUTION AND FORMS

TERMITES, or "white ants," are destructive native insects of which 44 species occur in the United States. They are distributed throughout the country, although in the southern, southwestern, and Pacific coast regions, where both the subterranean and nonsubterranean kinds occur (fig. 1), they are more numerous and injurious than elsewhere.

These so-called white ants are not true ants, although they are superficially antlike and live in colonies made up of different forms or castes. In these nests or colonies both wingless and winged mature individuals are produced. The brownish or blackish, elongate, slender, antlike, colonizing, sexual adults (fig. 2) with long white wings, unlike the other forms, have functional eyes and their bodies are able to endure full sunlight. These migratory males and females appear normally once a year during a short period. There are three stages in the life of white ants: The egg, the immature form (nymph), and the mature individual (including sterile workers (fig. 12, *b*), soldiers (fig. 12, *a*), and the various fertile reproductive forms).

LOCATION OF COLONIES OR NESTS

The nests of some species of termites¹ are in the earth and in dead and decaying wood. These species are of subterranean habit, timber and trees being attacked by the workers only through the ground. The nests of some other species,² however, are excavated in wood and trees by the winged forms, there being no workers and no underground life.

With the clearing of land and the consequent destruction of their natural breeding places in the dead trees, decaying stumps, and logs of the forests, termites become increasingly destructive to the woodwork and contents of buildings (figs. 3-9), telephone poles, fences, or any timber in contact with the ground, as well as to living vegetation, including not only fruit and shade trees, shrubs, and flowers, but also truck (fig. 10) and field crops and, in California,

FIG. 1.—Map showing (line AA) the northern limit of damage by subterranean termites in the United States; BB, the northern limit of damage by dry-wood or nonsubterranean termites

grapevines. The principal food of termites is cellulose, which they obtain from either dead or living vegetation.

Termites in the United States are mainly species of subterranean or wood-boring habit and are not so spectacular or common as the mound-making or tree-nesting termites of the Tropics. Very few termites in this country have habits which make them conspicuous, or come above ground into the sunlight, except during the annual colonizing swarm; hence they largely escape notice until they become injurious.

TERMITES WHICH ARE SUBTERRANEAN IN HABIT

Subterranean termites live in forests, building their nests in the wood of standing timber, logs, or stumps, in cleared land, any wood

¹ Genera *Reticulitermes* Holmgren, *Leucotermes* Silvestri, *Amitermes* Silvestri, etc.

² Genera *Termopsis* Heer, *Kaloterms* Hagen, *Neoterms* Holmgren, *Cryptoterms* Banks, etc.

in contact with the ground or, in the plains, in a labyrinth of underground passages in the earth, usually underneath wood or vegetation.

Termites are soft-bodied and always conceal themselves within wood, in the earth, or within their earthlike carton shelter tubes (fig. 11). The grayish-white, soft-bodied, wingless, sterile "workers" (fig. 12, below) are in reality the destructive form. These workers make the excavations occupied by the colony and enlarge and extend them as the colony increases. They live underground or within the wood, are blind, and shun the light; as a result they are rarely seen. In burrowing through wood the workers often completely honeycomb it, usually following the grain and eating out the softer, thin-walled, larger-celled spring or new wood. They are able to penetrate the hardest of woods, provided they have access to moisture in the ground. In extending their galleries in wood and vegetation, subterranean species carry moisture with them by means of moist excrement mixed with earth.

**TERMITES WHICH ARE
NONSUBTERRANEAN IN
HABIT**

The nonsubterranean termites which are injurious attack wood directly; but, instead of following the grain continuously, they excavate through it longitudinal chambers of limited length. The sexual adults, after they have lost their wings, and the young or nymphs, are the destructive forms. Their pellets of excrement (fig. 13) are regularly impressed, and sometimes completely fill or block up the burrows in a compact mass; they are often expelled as dry droppings from the infested wood. These termites are destructive to the woodwork and furniture in buildings (figs. 14, 15), as well as to living trees. Apparently they can exist without the great amount of moisture necessary to the life of termites which are subterranean in habit.

FIG. 2.—Winged sexual adults of eastern subterranean termites; upper, *Reticulitermes virginicus*, nearly $3\frac{1}{2}$ times natural size; lower, *R. flavipes*, enlarged 3 times

THE COLONIZING SWARM

At certain seasons, usually spring or fall, but varying with the species and the locality, the winged, sexual individuals migrate in large numbers from the parent nests. They then lose their wings and breed new colonies. In the case of the subterranean termites, wood and trees are usually entered indirectly through the ground,

although sometimes these insects enter trees under bark loosened by sunburn, etc., or through scars or borer holes, provided sufficient moisture is present. Nonsubterranean termites enter the wood directly, or, in the case of trees, through wounds or borer holes or under loose bark; moisture is not necessary.

In the new quarters eggs are laid, the young develop, and in a few years the colony increases in numbers, and the "workers" are able to feed and care for the reproductive forms—the "king" (title page, left) and "queen" (title page, right)—as well as the soldiers (fig. 12, *a*), which are sterile forms adapted to protect the colony from insect enemies, notably the true ants. In recently formed young colonies the rate of egg laying is slow, but mating is repeated, and, although there is at first a gradual increase, later the increase in the numbers of the broods is rapid. In old colonies there are thousands and tens of thousands of individuals.

Egg laying occurs over a considerable period during the warm months in colonies out of doors. In infested buildings

FIG. 3.—Damage to oak flooring by the common eastern subterranean termite *Reticulitermes flavipes*. Note that damage is not apparent on upper surface

artificially heated, where an even temperature is maintained, the insects are active and may lay eggs every month of the year. The number of eggs laid depends on age.

THE REPRODUCTIVE FORMS

Owing to the increasing number of eggs that develop within her, the queen becomes enlarged, but never loses the power of locomotion. It was once believed that, since the queen mother was the source of the colony life, the termite colony would be exterminated if she were destroyed, but this has been disproved by more recent studies. In addition to the forms that have shed their wings, several different types of reproductive forms occur—forms with wing pads (the undeveloped wings of the nymphs) and wingless reproductive adults. The winged forms, however, are the normal type and occur as a single pair, whereas there may be hundreds of the other forms heading colonies. The reproductive forms with wing pads and the wholly wingless type usually have little color to the body, and the eyes are small. These forms rarely come above ground or leave the burrows in wood.

FIG. 4.—Quartered-oak flooring damaged by the subterranean termite *Reticulitermes flavipes* in an infested building, Washington, D. C., 1915

The location of the queens in the colony depends upon the season of the year. During periods of intense heat or drought in the plains or in arid or prairie regions the subterranean termites burrow deeply below ground or to a less depth under stones, cow chips, etc. During winter in the colder climates they burrow below the frost line.

Whole colonies of subterranean termites migrate when conditions become unfavorable. The nonsubterranean termites, on the other hand, are not able to leave the wood in which they have excavated their nests.

PREVENTING AND REMEDYING DAMAGE TO WOODWORK OF BUILDINGS

DESTRUCTION OF BREEDING PLACES ABOUT THE BUILDING SITE

If buildings are to be constructed on recently cleared woodland, decaying logs and stumps should be removed from the soil in the vicinity and burned. If, because of the presence of decaying wood

FIG. 5.—Tube of coarse yarn from bale of cotton on floor of building infested by subterranean termites, Greenville, S. C.

and humus, the subterranean termites are numerous in the earth, the soil should be deeply plowed or otherwise broken up and treated with chemicals to kill the insects. Effective poisons for this purpose are sodium cyanide;³ a 10 per cent solution of sodium arsenite;^a kerosene oil; 1 part coal-tar creosote and 3 parts kerosene oil (this mixture should be strained

through burlap before use); carbon-disulphide emulsion, which is on the market ready for use; orthodichlorobenzene; lye; or other contact poisons or gases. Live steam forced into the soil will serve the same purpose as the gases. Decaying fence posts, sidewalks, etc., should be removed and replaced with treated wood, concrete, stone, or other resistant substances; such decaying material would facilitate the formation or perpetuation of the termite colonies.

PROPER CONSTRUCTION OF BUILDINGS

Termites will infest not only old buildings but also improperly constructed new buildings, and these are often badly infested. It is not the age of the building but the manner in which it has been constructed that renders it liable to attack.

Insulation.—Complete insulation from the ground of all untreated woodwork of buildings is the only effective permanent remedy

FIG. 6.—Cotton jacket of rubber-lined fire hose from infested building in Missouri, damaged by subterranean termites (*Reticulitermes* sp.)

³ For each acre to be treated, dissolve 160 pounds of granular sodium cyanide in 12,000 gallons of water.

^a This poison is caustic and should not be used near living vegetation.

FIG. 7.—Revenue stamps damaged by the subterranean termite *Reticulitermes flavipes* on infested flooring in the Bureau of Engraving and Printing, Washington, D. C., 1921

FIG. 8.—Correspondence and advertisement regarding issue of county bonds; damaged by subterranean termites infesting the building in Virginia in which they were stored

against attack by subterranean termites, and the only relief from their presence. These insects must maintain contact with the ground to obtain the moisture necessary for their existence. When contact with their moisture supply in the earth is cut off, the subterranean insects in the damaged wood, no matter how numerous, soon dry up and die.

Foundations, supports, etc.—To prevent subterranean termites from reaching the woodwork of buildings from their nests in the ground, the foundations of buildings should be constructed, if possible, entirely of stone, brick, concrete, or concrete and steel,⁴ including the pillars in the basement or cellar. The walls, partitions, and flooring in the ground floor, basement, or cellar should also be of concrete. Wooden flooring can be laid over this concrete floor if desired. If the flooring is to be of concrete, the concrete should be laid on a gravel base.

In buildings where stone, brick, or concrete foundations are impracticable, timber impregnated with coal-tar creosote should be employed, and no untreated wood should come in contact with ground

which may be infested with termites. Wood to be protected from termites should be impregnated with coal-tar creosote by either the cylinder-pressure or the "open-tank" process. Full details regarding these processes can be obtained from the United States Forest Products Laboratory, Madison, Wis. The open-tank process is simpler, can be made to give satisfactory results where properly used, and can be operated by unskilled labor. If for any reason neither of the foregoing processes can be used, three coats of hot coal-tar creosote brushed on the wood, with sufficient intervals between brushings to permit each coat to dry, will be fairly effective. This

FIG. 9.—Damage by the subterranean termite *Reticulitermes flavipes* to shoe stored on infested woodwork in a building in New York City

FIG. 10.—Injury to carrots by the subterranean termite *Reticulitermes claripennis*, at Dallas, Tex.

⁴ In the Southern States, especially in the subtropics, the more valuable permanent buildings should be constructed entirely of steel and concrete.

method, however, is not generally recommended, since it will preserve the wood for only a few years.

The supports of porches or steps should never be laid directly on the ground, but should rest on rock or concrete. Window sills and frames in the basement or cellar should be laid over concrete and the woodwork should not come in contact with the ground. The supports of the woodwork of coal bins in basements or cellars should not be set in the ground, but should rest on concrete, not extending through the concrete into the soil.

Concrete flooring.—Even in permanent stone or concrete and steel buildings the concrete flooring is often constructed improperly.

FIG. 11.—Carton, earthlike, shelter tubes constructed by the eastern subterranean termite *Reticulitermes flavipes* over brick wall in dark, heated, damp basement: these tubes are used in passing over impenetrable substances. Inset, near view of tubes to show texture

Usually the base of the concrete floor is a loose combination of coarse gravel or cinders and cement grout. This very rough conglomerate, even if several inches thick, has many cracks and large holes running through it. Over this is laid a layer of solid concrete about 2 or 3 inches thick, of fine texture, in which untreated wooden sleepers are laid while it is still moist. These sleepers nearly or quite reach the coarse, open conglomerate in contact with the earth, and to them is nailed the wooden flooring. (Fig. 17, A.)

It will be seen that this is faulty construction. Termites have free access from the earth, in which they have galleries, through the porous conglomerate, to the untreated beams and flooring. *There should be a layer of solid concrete at least 1 inch thick between the grout and the wood.* (Fig. 17, B.)

Among the commonest means of infestation are wooden sleepers or stringers laid in concrete while the latter is still soft, or placed

FIG. 12.—Mature soldiers (a) and mature workers (b) of the eastern subterranean termite *Reticulitermes flavipes*. Slightly more than six times natural size

simply inserting a sheet of noncorroding metal to serve as a mechanical barrier or "termite shield" into the masonry and turning the projecting edges downward at an angle, communication of termites with the earth, where they obtain moisture, can be cut off. In less pretentious frame buildings, metal caps are placed over the tops of construction stone piling or pillars, or wooden supports.

This is a similar method to that used in rat-proofing corn cribs. It is effective and practicable where

before the concrete has been poured in. Termites enter through these wooden beams. Other common means of infestation are supports of coal bins extending through concrete and through the disintegrated lime mortar of brick walls.

Termite shields, guards, or metal caps.—The most injurious termites in this country are subterranean in habit and require constant access to the earth in order to attack wood which is either in contact with the earth or which they reach through covered, earthlike shelter tubes constructed over the face of stone, concrete, or brick foundations. In consequence they can be kept out of buildings by means of metal barriers. By

FIG. 13.—Impressed pellets of excrement of non-subterranean termites (*Kaloterмес* sp.), which drop from infested wood. Greatly enlarged

untreated timber is placed over masonry foundations.

Again, in the case of stone and steel buildings, the steel or stone pillars or piping extending down through the concrete floor to foundations in the earth often make infestation possible, owing to the fact that the concrete floor does not fit tightly about them. Overlapping strips of metal extending horizontally several inches from the pillars or pipes should be imbedded in the concrete floor in order to obtain a tight joint (fig. 17, B); or liquid (adhesive) asphalt should be poured in to plug up the crevice or frame.

FIG. 14.—Bottom of bureau drawer damaged by the nonsubterranean wood-boring termite *Cryptotermes brevis*, which occurs at Key West, Miami, and Palm Beach, Fla.

Avoiding dampness.—Complete dryness of foundation timbers and basement walls and flooring is an important aid in rendering buildings safe from attack by termites. Dampness, warmth, and darkness attract these insects. A deep air space should be left between the ground and wooden flooring, unless, as in factories, heavy trucking is to be used over the floor, in which case this is impracticable.

An air space should be left between the concrete floor and the wooden floor laid over it. Concrete floors should be laid on a gravel

base, which will prevent dampness and cracking. The points of juncture between concrete walls and wooden flooring should be filled in by rounding off the concrete at these places, since cracks often occur where the wall and floor join at right angles. (Fig. 17,

FIG. 15.—Furniture damaged by the nonsubterranean, wood-boring termite *Cryptotermes brevis*, Key West, Fla.

B.) Termites often come up through cracks between walls and flooring.

By mixing a heavy mineral residual oil with Portland cement, a material is formed almost perfectly nonabsorbent of water and therefore excellent for use in damp-proof construction. Where the

various patent or noiseless floorings are used on the ground floor they should always be laid over a concrete base, especially if they contain wood fiber as a constituent.

In no case should untreated beams be completely surrounded with mortar or brick; there should be a space around them sufficient to permit air circulation. Beams should not be set in earth or in moist concrete, but on rock or dry concrete, or in grooves in the latter.

Bungalows or frame buildings which have no cellar should be raised from the ground on stone, concrete, or brick foundations to a height which will allow light and air to penetrate beneath.

Lime mortar.—In brick walls, where lime mortar is used, this sometimes disintegrates after a few years and termites penetrate through the spaces thus left between the bricks, especially where the brickwork is below the ground level. (Fig. 16.) Such walls should be faced with Portland cement 1 inch thick, especially if untreated wood is to come in contact with the bricks.

FIG. 16.—Interior view of portion of whitewashed brick foundation wall of building, below ground level, showing shelter tubes of our common subterranean termite (*Reticulitermes* sp.) penetrating the disintegrated lime mortar. These termites came through the earth banked up against the exterior wall. To remedy this condition the exterior wall will have to be faced with concrete for some distance below the ground level.

Use of chemically treated wood for interior woodwork, furniture, etc.—In the Southern States, especially in the subtropics, in the more valuable permanent buildings the interior woodwork, furniture, etc., should be impregnated with preservatives, since nonsubterranean termites that attack wood directly are common in those regions. Zinc chloride, bichloride of mercury,⁵ sodium fluoride, and chlorinated naphthalene⁶ are effective preservatives.

⁵ The extremely poisonous character of mercuric chloride renders its use dangerous. It is slowly volatile, and there is a possibility that it will be given off continuously in small quantities from the treated wood. However, it has been extensively used in Germany and apparently without serious consequences.

⁶ This preservative is usually referred to as trichloronaphthalene and, as compared with other chemicals, it is really comparable to a technical product having naphthalene, monochloronaphthalene, dichloronaphthalene, and probably some of the higher chlorinations as impurities. Its melting point ranges between 190 and 210° F.—the specification under which it is sold.

Wood-pulp products, such as the various wood-fiber processed or composition boards, for interior finish and substitutes for lath (fig. 18), or for exterior use, can be protected from attack by termites by adding certain poisons, such as crude carbolic acid, to the pulp or laminated boards in the course of manufacture. Available poisons

FIG. 17.—A, Improperly constructed concrete flooring: *a*, Gravel or cinders loosely cemented with coarse concrete, 3 inches thick, but with many crevices and holes; *b*, solid, dense concrete, 2 inches thick; *c*, 2 by 4 inch untreated wood sleeper set in moist concrete over the grout; *d*, $\frac{3}{8}$ -inch pine flooring nailed to sleepers. B, Properly constructed concrete flooring: *a*, Gravel or cinders loosely cemented with coarse concrete, but with many crevices and holes; *b*, asphalt waterproofing $\frac{1}{8}$ inch thick; *c*, dense concrete, 3 inches thick; *d*, 2 by 4 inch treated wood sleeper set in a groove in concrete which insulates it from termites in the earth; *e*, $\frac{3}{8}$ -inch flooring nailed on sleepers; *f*, metal collar around pipe which runs down through the concrete (this collar should be soldered to the pipe and embedded in the concrete); *g*, shoulder of concrete at point of wall and concrete floor to avoid a right-angle connection and consequent cracking.

for this purpose are crude carbolic acid at the rate of 1 gallon to 1,000 square feet; bichloride of mercury at the rate of 49 ounces per 1,000 square feet; or copper sulphate at the rate of 113 ounces per 1,000 square feet. Chlorinated naphthalene is a suitable poison for cane-fiber boards.

MODIFICATIONS OF CITY BUILDING CODES

One of the simplest and most effective means of prevention of attack would be to modify the building regulations or codes of various cities so as to include a few simple rules to protect houses from damage by termites. As the principal object in view is to keep all untreated wood from contact with the ground, where the termites live and from which they get their moisture, the regulations should stipulate that no floors, sills, beams, clapboard, etc., of untreated wood may be laid on or in the earth and that untreated beams may not be laid in concrete without at least 1 inch of concrete underneath and separating it from the earth; that in foundations or in cellar walls in contact with the earth a special grade of hard mortar should be used, since lime mortar⁷ after some years' service disintegrates; that all brickwork extending below the surface of the ground shall be faced and capped with concrete at least 1 inch thick; and, where nonsubterranean as well as subterranean termites occur, that only woodwork impregnated with preservatives be used for exterior and interior construction, unless it is impracticable to obtain such treated wood.

FIG. 18.—Pressed wood-pulp tile mined by subterranean termites infesting the building, Biloxi, Miss.

LOCATING AND TEMPORARILY ARRESTING THE DAMAGE

Although it may be difficult to eliminate termites and stop further damage by them, when once these insects have become established in the woodwork of a building, the approximate point of entrance should be sought at once by careful examination of all woodwork in contact

⁷ The Bureau of Standards of the United States Department of Commerce recommends a mortar composed of 1 part Portland cement to 3 parts of sand graded from fine to coarse, with no grains larger than will pass through the No. 10 sieve, to which may be added 10 per cent by weight of the cement of some workability agent, such as hydrated lime, for use in locations where termites abound. Such a mortar, it is believed, will have the desirable properties of both cement and lime mortars and, furthermore, will contain a sufficient quantity of cement to prevent the penetration of it by the termites. (See fig. 16.)

with the ground. To do this, it may be necessary to tear up the foundations, flooring, and some other woodwork. The foundation timbers and interior woodwork found damaged should be removed, and the ground where they were set should be broken up and drenched with some liquid which will kill or at least temporarily prevent the further activity of termites at that point. Any of the insecticides³ mentioned under the heading "Destruction of breeding places about the building site" (p. 6) will meet the need.

INDICATIONS OF INFESTATION

The annual emergence of large numbers of the flying termites is an indication as well as a warning that the woodwork is infested, and the point of emergence indicates the approximate location of the infested timbers. Even if the insects are not observed "swarming," large numbers of the dead winged adults or of the discarded wings usually will be found near by. Frass and earth thrown out of crevices through which the insects emerge are also evidences of their presence. Another warning is the presence of branching shelter tubes of small diameter, made of earth mixed with finely powdered wood, on foundation timbers or other woodwork, or over the surface of stone, brick, or other impenetrable foundation material (fig. 11), through which the insects travel from the ground to the woodwork. These tubes should be broken off and the ground where they originate broken up and drenched with one of the liquids named in the paragraph under the heading, "Destruction of breeding places about the building site" (p. 6).

In the case of the nonsubterranean termites, which infest wood directly, evidences that they are damaging wood are the impressed pellets of excrement (fig. 13) which are expelled from the wood. Other evidences are the holes, similar in size to BB shot, where the insects entered the wood. (Fig. 19.)

KILLING THE WINGED ADULTS WILL NOT STOP THE DAMAGE

When efforts are made to prevent further damage by termites in buildings, it should be realized that the numbers of these insects may be constantly recruited from some undiscovered, outside, central colony. The destruction of the winged colonizing adults with house-

FIG. 19.—Entrance holes of sexual, colonizing adults of a non-subterranean termite (*Kaloterms hubbardi*) in wood of dead cottonwood tree in Arizona.

³ If sodium cyanide is used, a strong solution, 1 ounce to a gallon of water, is desirable.

hold ammonia, etc., at the time of emergence, although beneficial in preventing the establishment of new colonies, will not eradicate the insects infesting the woodwork. The most destructive forms are the white, wingless workers, which remain within the wood.

DISCONNECTING WOOD FROM THE GROUND

Subterranean termites infesting beams or other wood will die out if the wood is disconnected from the ground. Knowledge of this fact will save time and expense, especially in the case of old frame buildings, where extensive repairs would be unwarranted. Disconnecting untreated foundation timbers from contact with the soil will also cause the death of subterranean termites in the other woodwork, furniture, and stored material in the building, even if they have penetrated to the second or third floors. These timbers need not be removed or replaced unless seriously weakened structurally. However, if the wood is kept moist by some other means, such as water leakage, the termites will continue to work and thrive. This applies especially to damp corners of basements near outside water pipes, bathrooms, kitchens, and the like.

Wooden floors laid directly on the ground or on stringers on the ground, or set in concrete, should be removed. There should be a layer of concrete between the earth and the wooden floor (fig. 17, B). Wooden baseboards should be removed.

REPLACING WITH CONCRETE

Concrete floors and concrete baseboards should be substituted for wood. Untreated beams penetrating through concrete floors into the earth and the lower parts of door jambs and casings should be cut off at least 6 inches above the ground or floor and replaced with cement plinths, which should project one-quarter inch beyond the jambs and casings. Metal strips should be sunk down from the woodwork into the concrete. Wooden thresholds, wainscoting, window sills, subsills, and stools in the basement or ground floor should be removed and placed on concrete (a layer of concrete between the earth and wood) or replaced with concrete.

REPLACING WITH METAL

In basements and cellars steel rails or other structural metal work can sometimes be economically used to replace weakened timbers.

SPRAYING AND POISONING

Fumigation, poisons, and spraying are of no permanent value against subterranean termites in buildings, since, while they may kill the flying insects and some of those in the wood, others will continue to come up from the ground.

In the case of termites which do not live in the ground, however, but attack wood directly, it is advisable to remove and replace the wood if the damage is slight and localized. Where the wood has been seriously damaged, but not structurally weakened, saturate the infested wood with orthodichlorobenzene. The wood should be thoroughly saturated with this chemical, a rag or mop being used,

dripping wet, or the liquid can be applied as a spray. Several applications may be necessary to kill the insects.

If orthodichlorobenzene is used as a spray, it is advised that the house be opened up before and for some time after treatment, since there is quite an odor to the chemical which may prove disagreeable in a closed room. Also, in spraying timbers overhead care should be taken not to let the liquid drip down, since it might slightly burn the face and hands and would be painful if it got into the eyes.

If the orthodichlorobenzene treatment is not practicable and the wood is too thick for the solution to penetrate, blow dry Paris green by means of a bellows into holes bored with an auger into the infested wood; these holes should penetrate to the galleries of the termites. If, because of moisture, the Paris green becomes caked, the treatment should be repeated. Wood structurally weakened should be removed and replaced with treated wood, discussed in Department Bulletin 1231, or strengthened with structural steel.

By far the greater and more serious damage to timber in the United States is caused by termites of the subterranean type. Department of Agriculture Leaflet 31, on the prevention of damage by termites in buildings, discusses in detail the control of subterranean termites.

PREVENTING AND REMEDYING INJURY TO LIVING VEGETATION

FRUIT, NUT, SHADE, AND FOREST TREES

Termites cause occasional but serious injury to living trees, shrubs, nursery stock, and grapevines. A wise preventive measure is the removal of all loose wood which may afford them shelter, such as prunings, dead and dying trees and the like, and untreated fence posts and similar material. Cleanliness in orchard and forest management is important. Since termites render unmerchantable the forest trees which have been killed by insects, fire, or disease, all timber from such trees should be utilized or removed as promptly as possible.

Covering scars and pruned areas.—To prevent infestation, care should be taken that the trees do not become scarred, especially near the base. Scars and all pruned areas should be treated with a mixture of one-fourth creosote and three-fourths coal tar. This mixture should not be allowed to come in contact with the living tissues at the edges of the bark; to protect them a shellac should be applied.

Commercial fertilizers.—Soil heavily manured will attract subterranean termites, since they can obtain food from the animal manure; commercial fertilizers should be used in preference in regions where termites are common in the soil.

YOUNG PLANTATIONS OR NURSERY STOCK

Injury to nursery stock will be more serious on recently cleared land, to stock from 1 to 3 years old, and where decaying wood is abundant. Any debris in which the insects breed should be removed. In general the use of recently cleared land should be avoided in planting nursery stock. Earth used in banking should be free from rotten wood. Care should be taken not to allow the roots to dry out before planting, as weakened stock is specially susceptible to attack; particular care should be given grafted stock. In the case

of the pecan it is recommended that two or three cereal crops be grown on newly cleared land before the young trees are set out.

Tree surgery.—Properly executed tree surgery¹¹ sometimes is effective in repairing injury by termites to valuable old fruit and shade trees.

Insecticides.—When valuable trees are infested but not yet dying, subterranean termites can be killed in the soil, if it is moist and not too compact, by breaking it up near the tree and pouring carbon disulphide on the earth at a distance of about a foot from the trunk, then covering the liquid over with earth. The gas from this liquid will penetrate the subterranean galleries of the termites. It is somewhat dangerous to plant life and very large doses should not be used. *Care should be taken in handling this volatile fluid, as the gas or vapor from it is highly inflammable and explosive when mixed with air in certain proportions; no flame should be brought near it, and the fumes should not be inhaled, as they are poisonous.* There are indications that an emulsion of carbon disulphide, which is on the market ready for use, may prove more effective than carbon disulphide alone; as in the case of the disulphide itself, the earth is loosened up and the emulsion poured on, at least a foot from the tree. Carbon tetrachloride,¹² the gas from which is neither explosive nor inflammable, can be similarly used but is apparently not so effective as carbon disulphide. The gases from carbon disulphide and carbon tetrachloride, being heavier than air, will not readily rise.

If tree trunks or branches are infested, they must be treated with poisonous liquids, such as kerosene emulsion poisoned with sodium arsenite.¹³ Orthodichlorobenzene is probably useful in the same way. When treating the infested portions, which are usually in the dead heartwood, these liquids must not be permitted to come in contact with living plant tissue.

Where nonsubterranean termites are infesting living trees, they can be poisoned with dry Paris green in the manner discussed on page 18.

VINEYARDS

In vineyards all dead or diseased vines should be removed. All exposed areas left by pruning should be painted with preservative coatings and the prunings should be burned promptly. Near-by stands of tree windbreaks should be carefully cared for and kept free from infestation by termites. Trellis posts should be creosoted.

FIELD AND TRUCK CROPS

Deep fall plowing.—Deep, late, fall plowing will be of value in breaking up the galleries and nests of subterranean termites on ground planted to field or truck crops. Irrigation of the land, where practicable, will be effective; this can be done before planting the crop.

¹¹ For information on this subject the reader is referred to Farmers' Bulletin 1178, "Tree Surgery," which may be obtained on application to the United States Department of Agriculture.

¹² Carbon tetrachloride is a thin, transparent, colorless, volatile liquid, which forms a gas with a pungent, aromatic odor. Like carbon disulphide it is heavier than air. Although it is only about one-half or one-third as effective as carbon disulphide when used at the same dosage rate, it has the great advantage that its gas is neither explosive nor inflammable; there will be no fire risk attending its use in buildings.

¹³ In each gallon of the water used for diluting kerosene emulsion or miscible oil, dissolve 1 ounce of sodium arsenite.

Burning stubble.—Care should be taken not to plow under stubble which will serve as food for termites; it should be burned. The use of commercial fertilizers instead of animal manure is also recommended where subterranean termites are common in the soil.

Insecticides.—Injury to corn in the prairie region of Kansas has resulted from the presence of termites in enormous numbers in the heavily sodded soil, where they feed on the roots of the vegetation. Such soil should be plowed up and treated with insecticides before crops are planted; effective insecticides are sodium arsenite and sodium cyanide. (See p. 6.)

Rotation of crops.—As a result of the frequent stirring of the soil, rotation of crops will aid in preventing termites from injuring them.

FLOWERS AND GREENHOUSE STOCK

Termites injure a great variety of flowers in gardens, as well as many plants grown under glass in greenhouses, where the warm, moist atmosphere maintained throughout the year promotes greatly the activity of the insects. Injury from termites is especially common where the plants are perennial and have woody stalks.

Sources of infestation.—Heavily manured flower beds are a source of infestation to the stems of the flowers, as well as to the woodwork of buildings near by, if suitable protection is lacking. Untreated wooden stakes used as supports of plants often become infested, and in time the insects attack the plants. In greenhouses termites often attack old label sticks, the wooden uprights supporting wooden benches set on or in the ground, and the wooden bench bottoms and plant pots, and later attack the growing plants. The insects come up through the ground and form dirt galleries over the supports, or burrow up through the wooden bench legs and run galleries the full length of the benches. They enter the soil in the pots through the drainage holes and eat out the main stalk of the root, killing the plant very quickly.

Proper construction of greenhouses.—Proper construction of greenhouses will practically safeguard plants growing in them from all injury by termites. Iron frames and concrete work should, wherever possible, replace bricks or woodwork. In cases of infestation of old greenhouses all wooden uprights supporting wooden plant benches should be sawed off, if set on or in the ground, and rested on stone, bricks, or concrete, above the surface of the ground. Proper repairing, including the removal of all infested wood, will prevent the plants from becoming infested in turn. Where woodwork is necessary the wood should be impregnated with zinc chloride or bichloride of mercury; it can be painted after treatment. Wood impregnated with coal-tar creosote can not be used with safety in greenhouses.

Avoiding stable manure.—In flower gardens, especially those located near the woodwork of buildings, commercial fertilizers should be substituted for stable manure in order to protect not only the buildings but also the growing plants.

Insecticides.—Either carbon disulphide or carbon tetrachloride¹⁴ can be used to kill termites in the soil if it is moist and not compact. Small holes should be made near the infested plants and a small

¹⁴ See footnote 12, (p. 19).

quantity of the liquid chosen poured in and the holes immediately closed tightly with earth. Calcium cyanide¹⁵ has also been found effective, but it should not be placed near living plants; it mixes readily with and enriches the soil and gives off an insecticidal gas which *should not be inhaled, as it is poisonous*.

An effective control may be found in the use of kerosene nicotine oleate or a 5 per cent kerosene emulsion.¹⁶ If the greenhouse benches are infested, but for any reason can not be replaced, they should be soaked thoroughly with this emulsion, as should also the ashes and sand under the pots on the benches. This may be done by removing the potted plants from a section of the bench, spraying that section, and moving the pots on the bench to cover the treated area, thus giving access to another section. Potted heliotropes and geraniums have been treated directly with the 5 per cent kerosene emulsion without injury to the plants, and the white ants in the soil of the pots were all killed. The soil should be wet down before this spray is used. This treatment should be given late in the afternoon and be followed early the next morning with a thorough syringing of the soil with water to wash out the surplus oil. It is important to remove all infested pots from the bench as soon as the infestation is noticed and to destroy the termites with kerosene emulsion.

SUMMARY

Damage by termites or "white ants" is serious to many classes of crude and finished forest products and is occasionally serious to living trees and other plants. These insects are especially injurious to foundation timbers, the woodwork of buildings, and material stored in buildings to which they have gained entrance. Damage to timber in contact with the ground is especially serious in the Southern, Central, and Pacific Coast States, and the tropical possessions of the United States.

The woodwork of buildings can be protected from the attack of termites by proper construction and by the use of wood treated with preservatives. These insects can be eliminated where already established in buildings by removing wood in contact with the ground and replacing it with wood chemically treated.

To construct buildings so that they will be white-ant proof, make their foundations, where possible, entirely of stone, brick, or concrete, including stone or metal columns or pillars in the basement to support the floor above; make concrete walls and flooring in basement or cellar, and lay concrete floors on a gravel base. Where stone or concrete foundations are impracticable, use timber impregnated with coal-tar creosote.

Lay basement-window sills and frames over concrete, and do not allow woodwork to come in contact with the ground. Never sink untreated timber in the ground or in moist concrete.

Complete dryness of the foundation and of the basement walls and flooring is an important means of rendering buildings safe from

¹⁵ Two ounces of calcium cyanide to 1 square yard of ground is recommended.

¹⁶ Kerosene emulsion is made as follows: Kerosene, 2 gallons; fish-oil soap, one-half pound; water, 1 gallon. Dissolve the soap in hot water and pour in the oil slowly, with constant stirring to emulsify.

Dilution: If 37 gallons of water be added to the above stock emulsion it will give 40 gallons of 5 per cent kerosene emulsion.

attack. Provide for air spaces between the ground and wooden flooring and lay concrete floors on a gravel base.

In regions where nonsubterranean termites are common (see map, fig. 1) woodwork should be treated with preservatives.

To eliminate termites already established in buildings, examine the foundation timbers and other woodwork in the basement to determine the approximate point of entrance and the extent of damage already accomplished. After removing the damaged wood, drench the ground with insecticides or poisonous solutions. Then replace the damaged timber with rock, brick, concrete, or metal work, or substitute, for the foundation, timbers impregnated with coal-tar creosote.

Since subterranean termites always require access to damp earth, shut off this source of moisture. The insects will then be unable to extend their galleries farther and will perish.

Nonsubterranean termites can be killed in infested wood by the use of insecticides.

Injury to living vegetation is occasionally serious, especially in the Southern States, the Southwest, and the Pacific Coast States. It can be prevented by clean cultural methods, deep fall plowing, and the use of insecticides.

Reprinted from JOURNAL OF FORESTRY
Vol. XXVIII, No. 6, October, 1930

TREE DAMAGE BY THE RED SPIDER

BY GEORGE S. PERRY

TREE DAMAGE BY THE RED SPIDER¹

By GEORGE S. PERRY

Senior Research Forester, Pennsylvania Forest Research Institute

Fine webs spun over leaves or leaves appearing gray and dusty or yellowish, may indicate the presence of the Red Spider, whose life cycle, habits, distribution, detrimental activities and control are here discussed.

HARVEST mites and red spiders, at times serious pests to trees and plant life, are minute Arachnid creatures with a world-wide distribution. They are not really insects, although commonly so considered because of the similarity of their life histories and relations to the plant and animal world. Probably the species most frequently met in the eastern United States are *Tetranychus telarius* with general host habits and often a serious green-house pest, and *Bryobia pratensis*, the clover mite, which also at times attacks fruit trees. There are, however, numerous species of harvest mites and allied forms with food habits and general structure like these two. All attack vegetation in about the same way and may be combatted by similar measures. Relatively few of the species are commonly recognized as pests, chiefly because their very small size enables them to escape notice.

DISTRIBUTION AND ABUNDANCE

The red spider (1) and all of its relatives are contained in the Order *Acarina* or *Acarida*, which included 133 genera with 450 species listed in the United States in 1907. In the Order are found species which are parasitic on

both plants and animals. Birds are commonly attacked by mites. A human trouble commonly termed "itch" is caused by a minute Acarid mite. Although the red spider is now generally recognized as a nursery pest, it is probable that most foresters are even more familiar with "red bugs" or "chiggers," which are the larvae of a small red spider form. The ticks that attach themselves to dogs, cattle, and sometimes to man are among the largest representatives of the group, which feed by sucking sustenance from their host. Some very serious diseases of both men and animals are now known to be spread by these relatives of the red spider. Warm and sunny climatic conditions seem most favorable to the order.

Among the trees recorded as attacked by red spider in Pennsylvania State Forest nurseries are Colorado blue spruce, Norway spruce, white spruce, Engelmann spruce, arborvitae, catalpa, honey locust, persimmon, tulip poplar, chestnut, and American elm. The spruces and elm probably suffer the worst. Citrus fruit trees in the South are often severely injured by red spiders and mites.

Since this report is written in the interests of forestry in Pennsylvania, the

¹ Presented at Conference of Pennsylvania Foresters, Mont Alto, Pennsylvania, January 29-30, 1930.

term red spider will be used hereinafter to designate those species that may feed on the leaves and succulent parts of trees and plants, while the term *Acarid* will refer to the general group.

European forest entomologists for some years past have made accurate determinations of the number and kind of insects hibernating in the forest soil and litter during the winter. By comparative studies over a period of years, it is possible to know which pest forms are on the increase, and to predict which ones may develop in hordes and become a serious plague. The remarkable fact emphasized by all such census work has been the high representation of red spiders and related life forms (Acarids). In practically every instance these made up from 44 to 92 per cent of the total counts on all sorts of sites, from sphagnum and lichen covered tundras of the North to the desert sands of the Sahara.

It is reasonable to believe that red spiders and their relatives are just as universally distributed in America as in Europe and Africa, yet only under exceptionally favorable conditions do they become a pest demanding active control measures for the protection of trees and other crops.

Almost every year forest nursery inspectors in Pennsylvania have reported the occurrence of red spiders in the nurseries of the State. In practically every forest tree nursery it is possible to find on needles and leaves the minute pale brown spots or dots that show the work of these creatures, even if they themselves have apparently disappeared. True red spiders (4) spin a dense fine web on the twigs and foliage on which they feed, if present in damaging num-

bers. The spiders are so small, however, that even under such conditions, a magnifying lens is required to see them.

DAMAGE BY RED SPIDERS

The life history of *T. telarius* is well understood, as this species has been known since before the time of Linnaeus. The female of the species over-winters on the lower part of various plants such as moss and grass, or hides in the litter on the ground. In spring it ascends growing plants and lays eggs at the rate of five to ten per day for a period of eight to twelve days. During hot, dry weather these eggs hatch in about four days and the young are fully developed ten to fourteen days later. This life cycle permits about six generations during a summer season, and if weather and all other conditions were favorable, a single mother might have offspring to the number of 750,000,000,000 by the end of the growing period, if the sexes were equally distributed. This explains why the red spider often becomes very injurious during a protracted drought, when plants are least able to resist the pest. Because of the limited water supply at such times, they cannot replace the sap withdrawn by countless minute but extremely greedy mouths. Under such conditions entire plants and even trees may be defoliated or killed, especially in nurseries where clean culture forces the pest to concentrate on single specimens. Red spiders apparently prefer to live and feed in colonies, in which condition they are afforded maximum protection from the fine mesh of webs they spin.

Although red spiders are not mentioned by Dr. E. P. Felt in his "Manual

of Tree and Shrub Insects," and are either not listed or only casually discussed in other publications on tree pests, they certainly deserve more consideration. Pennsylvania has had seven large forest tree nurseries within its borders. In four of these it has been necessary at times to take active control measures against spiders. The necessity would probably been more frequent and acute if good irrigation and shading facilities were not usually provided.

The dependence of red spiders on hot, dry weather, is helpful to the nurseryman, since he need not fear trouble under other conditions. This characteristic, however, has probably caused these small pests to be seriously underestimated and generally overlooked, since trees that suffer more or less severely from spider attacks at such times are commonly assumed to be suffering from drought. Closer examination would reveal the red spider as an important contributory factor in their failure.

District Forester H. M. Nicholas reported a very definite case of plantation loss from the red spider during the dry summer of 1929. This occurred in a plantation of 15,000 white pine trees planted in Roaring Brook Township, Lackawanna County, in 1915. In 1929 the trees averaged 10 feet in height, and some of them were 20 feet tall. Eight per cent of all the trees were killed outright by the red spider attack, and 7 per cent were injured to the extent that their foliage turned yellow and sickly in appearance.

When trees, especially evergreens, are planted, either for forestry or ornamental purposes, there is often injury and loss from spider damage. The

danger is greatest when trees are set on warm, dry south slopes, or on the sunny, south and southwest sides of buildings. Many trees die during their first year on such sites because red spider colonies develop and work serious injury before the trees recover from the shock of transplanting. Neglected lawns and abandoned fields often harbor the red spider, where it is ready and waiting to attack any new and favorite host that may be introduced. Sometimes the pest is present on nursery stock when shipped. In the tree nursery operated by the Pennsylvania Department of Highways near Lewisburg there has been considerable trouble with the red spider. It is possible that the infestation originated in the Mont Alto Nursery, as it was first noticed on stock that was transferred from the latter nursery, where the pest recurred over a period of five successive years, whenever weather favored it.

CONTROL MEASURES

This emphasizes the necessity of keeping nursery stock free of these mites. They can be combatted easily in the nursery or on individual ornamentals, but the forester can do little in a practical way to protect extensive plantations, if the pest is generally established, because of the excessive cost. A hard, driving summer rain is most effective in controlling infestations of red spiders. Planted trees and shrubs on lawns can be often satisfactorily treated with water, applied at high pressure with a hose, so as to destroy the spiders or wash them away. However, water treatment should not be considered sufficient in a nursery. All infested nursery stock should be thoroughly sprayed to guar-

antee that the pest is eliminated and will not be scattered far and wide to places where it may do much damage—often before its presence is known.

Any contact insecticide is effective against red spiders. Kerosene emulsion, other miscible oils, lime-sulphur and nicotine sulphate are all good. They should be applied twice, with an interval of about ten days between applications.

Powdered lime, wood ashes, flowers of sulphur, and soot sifted over trees when covered with dew, have been some of the measures used in the State Forest nurseries of Pennsylvania. The results have been beneficial, but careful spraying with insecticides is undoubtedly better.

The extent to which birds eat the red spider and mites is not known definitely. The pest is always worst on low growth in the open and on such trees as spruce, whose prickly needles discourage bird visitation. The United States Biological Survey (5) has made some very careful studies of the foods of birds based on stomach contents, yet only rarely are these life forms listed. This is explained by their extremely small size and fragile structure which renders identification impossible. There is little doubt that wrens, warblers, kinglets, chickadees, creepers, nuthatches, and vireos eat these pests. Such birds deserve protection for they are of the greatest importance in keeping under control nearly all kinds of injurious insects, as well as red spiders.

SUMMARY

It is probable that the red spider is doing more injury to trees in Pennsylvania than is generally realized. It is ad-

visable that foresters, nurserymen, and gardeners be informed about this minute pest and learn to recognize its presence. Sometimes the naked eye can detect the fine webs spun over the leaves or needles of host species, or they may look gray and dusty or yellowish, particularly after a period of dry weather. Such trees or shrubs should be carefully inspected with a good hand lens or reading glass, and in a surprising number of cases the pale pin-point feeding dots will be found. If the pests are still present, their pearly eggs or their remnants will be noticed, together with the busy eight-legged reddish mites. A twig or leaf which appears to be attacked by the red spider, if sent to the Department of Forests and Waters at Harrisburg, or to the Pennsylvania Forest Research Institute at Mont Alto, Pennsylvania, will permit of verification and will bring specific recommendations for helpful control measures, when requested.

REFERENCES

1. Comstock, John H. *The Spider Book*, Doubleday, Page & Co., New York City, 1920, pp. 85-88.
2. Russell, John. *Microörganism of the Soil*, London, 1923.
3. Tragårdh, Ivar. *Svenska Skogsvårds-föreningens Tidskrift*, 1928, pp. 795-809.
4. *Insects Injurious to Deciduous Shade Trees and Their Control*. United States Department of Agriculture Farmers' Bulletin 1169, p. 93.
5. *Food of our Important Flycatchers*. United States Department of Agriculture Biological Survey Bulletin 44, p. 63. Chapin, E. A., *Food of Vireos*, United States Department of Agriculture Bulletin 1355, p. 28.