ENVIRONMENTAL RESEARCH

RESEARCH AND TECHNOLOGY BRANCH

SEDIMENT BIOASSAY RESEARCH
AND DEVELOPMENT

R. A. C. PROJECT NO. PDF 03

Environment
Ontario

1285

ISBN 0-7729-7147-1

SEDIMENT BIOASSAY RESEARCH
AND DEVELOPMENT

R. A. C. PROJECT NO. PDF 03

Prepared for Environment Ontario by:

Gail Krantzberg
Sediment Specialist
Water Resources Branch

OCTOBER 1990

Copyright: Queen's Printer for Ontario, 1990
This publication may be reproduced
for non-commercial purposes with
appropriate attribution.

PIBS 1285

ACKNOWLEDGEMENT AND DISCLAIMER

This research was based on a sediment bioassay initiated by A. Hayton. I gratefully acknowledge W. Scheider, D. Persaud, and T. Lomas for their constructive advice during the course of this work and thank S. Petro and R. Pope for assistance in conducting the experimental bioassays. I thank W. Scheider and D. Bedard for their helpful comments in review of this document.

This report was prepared for the Ontario Ministry of the Environment as part of a ministry funded project. The views and ideas expressed in this report are those of the author and do not necessarily reflect the views and the policies of the Ministry of the Environment, nor does mention of trade names or commercial products constitute endorsement or recommendation for use.

c Her Majesty the Queen in Right of Ontario as
Represented by the Minister of the Environment

EXECUTIVE SUMMARY

Under the 1987 revision of the 1978 Great Lakes Water Quality Agreement between Canada and the United States, the parties, in cooperation with state and provincial governments were assigned the task of cooperatively developing and implementing strategies for mapping, assessing and managing contaminated sediment. Adequate assessment of contaminated sediment requires information beyond bulk sediment chemistry. The biological significance of in-place pollutants can be measured on the basis of structural or functional modifications of benthic invertebrate communities, and by demonstrating the bioavailability of contaminants through a variety of toxicity tests.

Laboratory sediment bioassays are an important component of biological assessment. Bioassays range from acute lethality tests to chronic, sublethal tests. Chronic exposures provide information unachievable from acute toxicity studies. Growth, reproduction, and other physiological parameters have been used as endpoints in chronic tests. Since benthic organisms can be an important vector in the transfer of materials from sediment to other compartments of the ecosystem, the sediment bioassay should also provide information on the extent to which contaminants may be mobilized into the foodweb.

Biologically based sediment quality guidelines are under development by the Ontario Ministry of the Environment. These guidelines will, in part, provide the basis for making decisions on remedial actions. When bulk chemistry exceeds specified concentrations, the draft guidelines recommend that biological testing, including sediment bioassays, be conducted to identify whether contaminants are biologically available.

Sediment bioassays measure the effects of contaminated sediments on the biota. By far the most frequently described approach is solid phase testing with either benthic or water column organisms. For the purpose of evaluating the impacts of in-place pollutants on the biota, as opposed to the consequences arising from dredging operations, the focus of this study was on the solid phase bioassay.

The principle objective of this study was to contribute to the development of a methodology for assessing the chronic and acute toxicity of sediments to biota. This included an examination of the effects of bioassay assembly and sediment manipulation techniques to the response of the test organisms, and the sensitivity of growth as a chronic endpoint.

It is reasonable to expect that the exposure of an organism to contaminants will vary with the state to which the sediment-water system is in equilibrium. I therefore examined whether an organism's response varied with the length of settling time of the bioassay assembly proceeding the introduction of the organism. The duration of exposure required for the response of organisms in test sediments to differ significantly from the controls was also not known. As a result, the experiment was designed so that replicates could be harvested day 10 and day 21. Analysis of the growth response of Hexagenia suggested that biomass changes were influenced both by sediment type and by the duration of the period of equilibration. Growth in both test sediments was inversely proportional to the duration of the equilibration period.

In accordance with the biomass changes noted for mayflies, growth inhibition was least when the fathead minnows were added 5 days after chamber assembly. There appeared to be no notable difference between the 6 hr. (5 hr. settling plus 1 hr. aeration) and 1 day equilibration periods with respect to biomass changes. Metal accumulation was inversely proportional to bioassay settling time. The effects of fish density were variable. Growth inhibition was greater with 15 as compared to 10 fish in some, but not all cases, and density apparently exerted no influence on biomass changes in the controls. This last finding is of interest, since it may indicate that the stress of possible overcrowding was exacerbated by the contaminated sediments.

Exposing fathead minnows to test sediment for 21 days without food introduces additional stress which could exaggerate the adverse effects of contaminants. Feeding, however, could alter contaminant accumulation, and gut clearance may be important for estimating true tissue concentrations of contaminants. A feeding experiment was conducted which revealed that feeding had little effect

on accumulation of trace metals and organic contaminants, however, significant decreases in concentrations of Mn, Fe, Al, Pb, and Ni resulted when fish were held for 24 h to purge their guts. Only for control organisms was growth improved by feeding.

Current methods for assembly of sediment bioassays often involve sieving and homogenizing the sediment. This effectively exposes the organisms to a uniform dose of contaminants that is in reality a mean dose of the heterogeneously distributed contaminants. In some cases, the extensive aeration of the sediment also results in a transformation of chemical species to forms that are of greater or lesser bioavailability. I examined the question of sediment homogenization by using diver-collected cores.

Intact sediment from one station resulted in higher mortality and poorer growth than homogenized sediments for mayfly nymphs, but did not significantly influence mortality or growth in fathead minnows. Intact sediment from a second station resulted in the reverse, better growth for mayfly nymphs and substantial mortality for fathead minnows. In a third station (sandy sediment), homogenization resulted in higher mortality than in the intact cores for Hexagenia. This was most likely caused by the elimination of the surface layer of fine-grained material (present in intact cores) and therefore, the elimination of suitable substrate for burial and feeding. Homogenization did not effect growth of fathead minnows, and may have ameliorated toxicity as measured by mortality.

Sediment manipulation, bioassay assembly, organism density, and bioassay duration were important determinants of the final endpoints measured. Further efforts devoted at refining substrate and feeding requirements of test organisms would assist in calibration of the bioassays.

The results of this research support the use of a 21 day sediment bioassay with organisms introduced no more than 24 h after bioassay assembly. Several test organisms should be included in a comprehensive assessment. This work focused upon the mayfly nymph Hexagenia limbata and the fathead minnow Pimephales

promelas. Growth inhibition was demonstrated to be a sensitive indicator of sediment toxicity.

Further research into the development of full or partial life cycle tests that include reproduction as an endpoint is warranted, and establishment of cultures for all test species, along with the use of reference toxicants, would be of great value for ensuring repeatability of bioassay results.

Integrated strategies for sediment management must include a consideration of biological effects observed both in the laboratory and *in situ*. An array of biological test methods can provide an integrated approach to the determination of the toxicological qualities of sediment contaminated with a variety and sometimes unknown chemicals. Sediment bioassays are a valuable vehicle for the assessment of sediment and provide information that reflects the biologically relevant forms of mixtures of contaminants.

TABLE OF CONTENTS

	<u>Page</u>
ACKNOWLEDGEMENTS	i
EXECUTIVE SUMMARY	ii
1.0 INTRODUCTION	1
2.0 OVERVIEW OF SEDIMENT BIOASSAY METHODS	2
2.1 Matrices Used for Biological Assessment of Sediments	3
2.2 Endpoints	4
2.2.1 Acute toxicity bioassay	4
2.2.2 Sublethal chronic sediment bioassays	4
2.2.3 Bioaccumulation	5
2.3 Sediment Bioassay Design	5
3.0 EXPERIMENTAL DESIGN, RESULTS, AND DISCUSSION	6
3.1 Experiment 1 - March 1 to March 25, 1988	7
3.2 Experiment 2 - March 31 to April 25, 1988	8
3.3 Experiment 3 - April 26 to May 18, 1988	10
3.4 Experiment 4 - June 30 to July 22, 1988	11
4.0 RECOMMENDATIONS FOR RESEARCH PRIORITIES	14
4.1 Chronic Bioassays	14
4.2 Reference Toxicants	15
4.3 Sediment Manipulation	15
4.4 Invertebrate Cultures	16
4.5 Organism Age and Size	17
4.6 Substrate Properties and Colonization Potential	18
4.7 Sediment to Water Ratio	18
4.8 Flow-through, Static-Renewal and Static Water Regimes	19
4.9 Field Comparison	19

4.10 Bioaccumulation	20
5.0 RECOMMENDED PROTOCOL FOR LABORATORY SEDIMENT BIOASSESSMENT	21
6.0 OTHER ACTIVITIES RELATING TO SEDIMENT BIOASSAY DEVELOPMENT	25
7.0 REFERENCES	26
TABLES	
Table 1 - Marine and freshwater organisms used in acute toxicity tests with sediments	34
Table 2 - Chronic endpoints employed for various organisms used in sediment bioassays	37
Table 3 - Sample location and sediment composition of substrates used in sediment bioassays	39
Table 4 - Contaminant concentrations in sediment used for bioassays	40
Table 5 - Effects of settling time on growth of <u>Hexagenia limbata</u>	41
Table 6 - Trace organics in oligochaetes in relation to bioassay settling time and exposure duration	42
Table 7 - Metals in oligochaetes in relation to bioassay settling time and exposure duration	45

Table 8 -	Effect of settling time on growth of <u>Pimphales promelas</u>	46
Table 9 -	Metals in fathead minnows in relation to organism density, bioassay duration and gut clearance	53
Table 10 -	Trace organics in fathead minnows in relation to organism density, bioassay duration and gut clearance	62
Table 11 -	Effect of feeding on growth of <u>Pimphales promelas</u>	62
Table 12 -	Effect of feeding and gut clearance on metal concentrations in fathead minnows	63
Table 13 -	Effect of feeding and gut clearance on trace organic concentrations in fathead minnows	64
Table 14 -	Comparison of intact cores and homogenized sediments with reference to growth and mortality of <u>Hexagenia limbata</u> and <u>Pimephales promelas</u>	67

FIGURES

Figure 1 -	Biomass changes in <u>Hexagenia limbata</u> nymphs in 10 and 21 day bioassays	68
Figure 2 -	Biomass changes in <u>Pimephales promelas</u> in 10 and 21 day bioassays	69

Figure 3 - Cr, Cu and Pb concentrations in mayflies and fathead minnows in beaker and intact core bioassay assemblies	70
Figure 4 - As and Ni concentrations in mayflies and fathead minnows in beaker and intact core bioassay assemblies	71
Figure 5 - Cd and Hg concentrations in mayflies and fathead minnows in beaker and intact core bioassay assemblies	72
Figure 6 - Zn and Mn concentrations in mayflies and fathead minnows in beaker and intact core bioassay assemblies	73
Figure 7 - Al and Fe concentrations in mayflies and fathead minnows in beaker and intact core bioassay assemblies	74

1.0 INTRODUCTION

The principle objective of the research conducted under the Research Advisory Committee, Ministry of the Environment Post-Doctoral Grant PDF03 was to develop a methodology for assessing the chronic and acute toxicity of sediment to biota and to determine the bioavailability of sediment bound contaminants by measuring tissue retention of polar and nonpolar substances.

To achieve this goal, the following studies have been conducted:

- Examination of the effects of bioassay assembly and sediment manipulation techniques upon the response of the test organisms.
- Assessment of the relative sensitivities of a series of endpoints, including:
 - growth inhibition
 - bioaccumulation
 - mortality
- Comparison of the responses of organisms occupying different ecological niches including:
 - benthic infauna e.g. mayfly
 - water column organisms e.g. fathead minnow
- Establishment of an interim recommended protocol for multiple organism/endpoint bioassays to be conducted in concert with field collections of benthic macroinvertebrates for the assessment of community structure and physiochemical analyses of in situ contaminants.

In addition, the following studies are recommended:

- Estimation of the comparative sensitivity of the amphipod Hyalalella

azteca and Chironomus as alternate bioassay organisms, and the establishment of invertebrate cultures for these organisms and for Hexagenia limbata.

- Development of full or partial life cycle tests with reproductive success as an additional endpoint
- Introduction of the routine use of reference toxicants as a means of ensuring comparable sensitivity of test organisms for each bioassay conducted
- Calibration of growth on the basis of substrate conditions
- Further investigations into sediment manipulation with consideration of the sensitivity of endpoints to static, static-renewal, and flow through systems

2.0 OVERVIEW OF SEDIMENT BIOASSAY METHODS

Currently, the literature contains several approaches to sediment bioassessment with reference to a range of benthic and water column organisms and various lethal and nonlethal endpoints. It is useful to summarize ongoing research in the field, to better understand the context within which this report applies.

Sediment bioassays measure the effects of contaminated sediment on biota. Examining this broad statement more closely reveals the potential for a myriad of test designs. The substrates which have been explored include sediment elutriates, pore waters, and whole, sieved or suspended sediments. Chamber construction ranges from petri dishes and test tubes to 40L aquaria with static, static/renewal, flowthrough, and recycling water regimes. Test organisms are benthic infauna or epifauna, macrophytes, fish or plankton. The bioassay responses or endpoints vary from acute lethality to nonlethal impacts during acute or chronic exposures. These range from the molecular and cellular levels such as biochemical deviations and induction of carcinomas, to the organismal

level such as growth and reproductive inhibition. To avoid confusion, it is worthwhile noting that the terms acute and chronic refer to exposure duration while the endpoints may be lethal or nonlethal. The most sensitive assays are those that utilize chronic exposure intervals and monitor for nonlethal toxicity responses of critical life stages of sensitive organisms. The bioassay assembly can also be applied to monitor bioaccumulation of pollutants by the test organism in order to estimate contaminant bioavailability and potential for foodweb mobilization of potential toxicants.

2.1 Matrices used for biological assessment of sediments

Sediment elutriates have been prepared as liquid phase matrices, principally to assess the impacts of dredging activities on water column organisms (Gannon and Beaton 1969, Lee et al. 1975, Shuba et al., Munawar et al. 1983). For example, the USEPA/US Army Corps of Engineers (1977) describe a 48h toxicity test that exposes Daphnia to the elutriate.

Pore waters have been considered as an alternate liquid phase to examine the effects of contaminated sediments on the burrowing infauna and to identify the route of exposure of different organisms to different pollutants (Bahnick et al. 1980, Rodgers, J.H., Jr., N. Texas State Univ., pers. comm.). The interstitial waters may be acquired by squeezing sediment, centrifugation followed by filtration, or through the use of dialysis membranes.

By far the most frequently described approach is solid phase testing with either benthic or water column organisms (e.g., Swartz et al. 1985a, Malueg et al. 1984, Cairns et al. 1984, Ingersoll and Nelson, 1987). For the purpose of evaluating the impacts of in-place pollutants on the biota, as opposed to the consequences arising from dredging operations, this research program centred on the solid phase bioassay. Studies based on pore waters are scant, although of potential significance particularly with respect to understanding the mechanisms involved in the bioaccumulation of contaminants. Nevertheless, the remainder of this review will be dedicated to solid phase testing with a variety of organisms primarily due to its conceptual simplicity and relatively greater compatibility with field conditions.

2.2 Endpoints

2.2.1 Acute Toxicity Bioassay

Sediment bioassay reviews by Nebeker (1984), Buikeman (1982), Munawar et al. (1984), Craig (1984) and Lamberson and Swartz (1985) all demonstrate the preponderance in the literature of relatively short-term exposures with mortality as an endpoint. Acute tests typically measure the lethality of the test sediment relative to a reference substrate. Acute tests are frequently conducted by placing a specified volume or depth of sediment in a beaker, jar or aquarium. This is followed by the addition of a volume of water, often at a ratio of 4:1 (v:v) of water to sediment (Nebeker et al. 1984). The assembly is either static, flowthrough or recycling, and aeration is normally indicated. Several organisms are added to replicate units and mortality is tabulated by the end of the exposure. Exposure times vary among organisms and for the same organisms being employed by different authors. Table 1 lists a number of acute bioassays that have been described for freshwater and marine fauna.

2.2.2 Sublethal Chronic Sediment Bioassays

Chronic tests provide critical information which cannot be secured from acute toxicity studies, especially where the contaminants are materials with delayed action and of bioaccumulative potential. When an organism's life stage influences its degree of sensitivity to pollutants, chronic studies are capable of detecting significant adverse biological impacts of polluted sediment that would not be observed in an acute test. In some instances, behavioral modification may result from exposure to contaminated substrates (Swartz 1985a) and this can have consequences for an organism's ability to compete for resources, reproduce, and/or avoid predation in nature.

Growth, reproduction, and other physiological parameters have been used as endpoints in chronic bioassays (Nebeker et al. 1984) as have behavioral activity such as burrowing (Swartz 1985a) and preference/avoidance (Gagnon

and Beeton, 1969, 1971). Table 2 summarizes some of the endpoints which have been considered for a number of different taxa.

In addition to the endpoints listed in Table 2, biochemical, enzymatic, histopathological and morphological changes have also been measured. These have been reviewed by Beak (1987) and will not be considered in detail here. While these latter options can provide sensitive early indications of organism stress, the approaches currently require a degree of technical skill, expertise, laboratory specialization, and financial resources that are sufficiently great as to preclude their integration into a routine protocol.

2.2.3 Bioaccumulation

Measurements of tissue concentrations of polar and nonpolar substances can be used to demonstrate that contaminants in sediment are biologically available and have the potential to enter the food web. Nebeker et al. (1984) include tissue analysis of chironomids, mayflies and amphipods in their bioassay approach. Since benthic invertebrates can be an important vector in the transfer of materials from sediment to other compartments of the ecosystem (Krantzberg 1987) the sediment bioassay should provide information on the extent to which contaminants may become mobile.

Currently, the toxicological significance of tissue residues in invertebrates and many vertebrates is virtually unknown. Unfortunately, assessing this linkage falls beyond the scope of this research, at present. It should be stated, however, that for an organism to metabolize or detoxify contaminants, there is undoubtedly a physiological cost in the production of enzymes and other proteins. Energy diverted from normal metabolic pathways for these purposes could be quantified and the ramifications at the level of the organism should be established.

2.3 Sediment Bioassay Design

Having outlined the variety of test organisms and endpoints of potential use, it is instructive to explicitly consider the many decisions that modify the end

result of the bioassay. From a biological perspective, factors such as organism size, age, sex, reproductive status, and history of exposure to pollutants can alter the response of an organism to a given sediment (Luoma 1983, Krantzberg and Stokes 1989). Clearly, interspecific variability should also be expected and is often observed. This has led to recommendations that several taxa be used for each bioassay (e.g. Nebeker et al. 1984). From a population standpoint, density dependent effects may also be postulated and should be confirmed or refuted. From a physical and chemical perspective, different methods of sediment storage, manipulation, chamber construction and assembly, could effect the outcome of the bioassay. Sediment handling necessarily disrupts sediment physicochemistry which has direct implications for contaminant activity.

These biological, physical and chemical processes were recognized in the development of a research program directed at on the importance of bioassay design. Recommendations for the application of the results are presented in the form of a detailed protocol (Section 5.0).

3.0 EXPERIMENTAL DESIGN, RESULTS, AND DISCUSSION

The first set of experiments examined the effects of equilibration time of the bioassay assembly and bioassay duration on mortality, growth and bioaccumulation by the mayfly nymph, Hexagenia limbata and the oligochaete Tubifex tubifex. The second experiment considered the effects of organism density, chamber equilibration time and bioassay duration on mortality, growth, and bioaccumulation of contaminants by juveniles of the fathead minnow, Pimephales promelas. The third series of test bioassays examined the consequences of feeding with respect to mortality, growth and bioaccumulation of contaminants by P. promelas. The fourth bioassay assessment compared the toxicity of homogenized sediments with that of sediment collected by diver and maintained as intact cores. Test organisms for the latter experiment were mayfly nymphs and fathead minnows. For all investigations, pH and dissolved oxygen were routinely measured.

3.1 Experiment 1 - March 1 to March 25, 1988

Purpose: To determine the effects of settling time, following the addition of sediments and water to the bioassay container, on sediment toxicity to oligochaetes and mayflies. To evaluate the importance of bioassay duration on identification of toxic sediment. To determine trace metal and trace organic bioaccumulation by oligochaetes as a function of bioassay design.

It is reasonable to expect that the exposure of an organism to contaminants will vary with the state to which the sediment-water system is in equilibrium. I therefore examined whether an organism's response varied with the length of time for which the bioassay assembly was left to settle prior to the introduction of the test species. The duration of exposure required for the response of organisms in test sediments to differ significantly from that of the controls was also not known. The experiment was therefore designed so that half of the replicates could be harvested at day 10 and half could be harvested at day 21.

The two test sediments were from a silty site in the vicinity of the Toronto Main STP outfall and a sandy site at Rice Lake. The control or reference sediment was a silty substrate from Honey Harbour. Two litre widemouth glass jars were filled to a depth of 3 cm with sediment (surface area = 100 cm²) and dechlorinated tap water was gently added. Organisms were introduced at 3 time intervals; 5 hours settling plus 1 hour of aeration, 1 day settling plus 1 hour aeration and 5 days settling plus 1 hour aeration. At each time interval, either 8 mayflies weighing approximately 25 mg/individual (wet weight) or approximately 1.5 gm (wet weight) of oligochaetes (c.a. 150 individuals) were introduced into the chambers. Each treatment had 4 replicates. Preliminary tests indicated no density dependent effects for mayflies at this density.

Sediment bulk chemistry and physical properties were measured at the onset of the experiment (Tables 3 and 4). Oligochaetes were held in dechlorinated water overnight to evacuate their gut contents and were frozen and submitted for metal and trace organic analysis. Mayflies were weighed and counted, but were of

insufficient biomass for analysis.

Analysis of the growth response of Hexagenia indicated that biomass changes were influenced both by sediment type and by the duration of the period of equilibration (Table 5). Growth in both test sediments was greatest when the mayflies were added when 5 days had elapsed following chamber assembly. Growth was poorest when organisms were added when 6 hours had elapsed following assembly (1 hour after aeration).

Mayfly biomass increased most in nymphs exposed to control sediments, followed by organisms exposed to Rice Lake sediment, and increased least in mayflies subjected to sediment from the vicinity of the Toronto Main STP. Growth inhibition relative to controls was more pronounced by day 21 ($p < 0.05$), as compared to day 10 (Fig 1).

Due to difficulties in assuring retrieval of all oligochaetes, bioamass changes were not considered as an endpoint in the bioassay. Bioaccumulation results were interpreted by analysis of variance. Trace organic concentrations in oligochaetes (Table 6) did not vary significantly with experimental settling time and bioassay duration. Manganese and Al concentrations in oligochaetes (Table 7) were significantly different from time zero, with Al increasing with exposure time and Mn decreasing with exposure time. One hypothesis is that with sufficient aeration of the bioassay chamber, Mn became less bioavailable with time due to the formation of insoluble oxyhydroxides. The increase in Al concentrations in oligochaete tissues with exposure time may reflect the effect of sediment ingestion which was not mirrored by other trace metals either due to their relatively low concentrations in sediment, poor bioavailability, or active metabolic regulation by oligochaetes.

3.2 Experiment 2 - March 31 to April 25, 1988

Purpose: To determine the effects of settling time and bioassay exposure interval on sediment toxicity to fathead minnows and to establish whether toxicity is dependent on organism density.

The test design and bioassay assembly paralleled Experiment 1. The two test sediments were from a sandy site in the vicinity of the Toronto Main STP outfall and a silty site in St. Mary's River. Three to four month old juvenile fathead minnows weighing approximately 0.5 gm per individual were added to each bioassay chamber at a rate of 10 or 15 individuals per replicate. Four replicates of each treatment were harvested after 10 or 21 days exposure. Fish were reweighed and half were immediately frozen for metal and trace organic analyses. The remainder were held for one day in dechlorinated water to clear their guts and to illuminate the significance of short term depuration.

In accordance with the biomass changes noted for mayflies, growth inhibition was least if the fathead minnows were added when 5 days had elapsed following chamber assembly (Table 8). There was no notable difference between the 6 hr. (5 hr. settling plus 1 hr. aeration) and 1 day equilibration periods with respect to biomass changes, and the effects of fish density were variable. Growth inhibition was greater with 15 as compared to 10 fish in some, but not all cases, and density apparently exerted no influence on biomass changes in the controls. This last finding is of interest, since it may indicate that the stress of the exposure to contaminated sediments could be exacerbated by possible overcrowding .

By day 21, all fish had decreased in weight (Fig 2). Minnows from the test sediment lost more weight than did those from the control sediment and analysis of variance revealed that this outcome was significant by day 21 ($p < .05$).

Trace metal concentrations in fathead minnows were responsive to several bioassay treatments (Table 9). Less Pb was accumulated in the presence of 15 individuals than in the presence of 10 individuals, while the reverse was true for Al and Cd. The explanation for this remains unclear may warrant further examination. More gamma-chlordane was accumulated by fish at the higher density as compared with the lower density (Table 10). This phenomenon remained true when tissue concentrations were corrected for lipid content.

With the exception of Cd, metal accumulation was significantly greater ($p < 0.01$) when organisms were added to the bioassay chamber after 5 hrs of settling

plus one hour of aeration, as compared with longer settling intervals. In addition, As continued to increase with exposure duration while Cr decreased significantly with bioassay duration. No other significant differences in tissue residues were observed as a consequence of bioassay duration. Concentrations of trace organics in fathead minnows did not vary with settling time or exposure duration (Table 10).

The importance of permitting an interval for gut clearance in the final measurement of contaminant residues in fathead minnows is discussed under section 3.3.

3.3 Experiment 3 - April 26 to May 18, 1988

Purpose: To determine the effects of feeding on growth, bioaccumulation and mortality of fathead minnows in sediment bioassays.

Based on the results of Experiment 2, it was apparent that even control organisms were experiencing stress due to starvation. In addition, there was evidence that the expression of toxicity was density dependent. Therefore, a feeding experiment was initiated employing 10 fish in each chamber, to be introduced to the chambers when 6 hours had elapsed after assembly. Contaminated sediments were from a silty site at Canagagigue Creek, a coarse sandy site from the Algoma slip (St. Mary's River), a silty site from Rice Lake and sandy site from St. Mary's River. Silty Honey Harbour sediment was used as a reference substrate. Bioassay design was as above, with a subset of the test organisms retained for 24 hours to depurate gut contents.

For each sediment, 4 replicates were fed ad libitum every second day and 4 replicates were not fed for the duration of the 3 week bioassay. Average biomass was c.a. 0.3 - 0.4 gms per individual (wet weight) at the onset of the experiment.

Both the Canagagigue Creek and Algoma slip sediments were lethal with mortality occurring from day 1 to day 14. No relationship between time until mortality and feeding was evident.

Since fathead minnows ingest sediment and consume detritus, biomass changes may be linked to substrate properties that are independent of contaminant loads. For chronic studies, then, standardizing food availability may be necessary. Introducing food, however, may complicate the interpretation of results, particularly with respect to bioaccumulation. The only organisms to show an increase in biomass were the minnows receiving food in the Honey Harbour treatment, while biomass was lost in the Honey Harbour, unfed minnows (Table 11). Interestingly, for the other contaminated sediments, feeding did not affect the extent of biomass lost, perhaps indicating that the presence of the pollutants significantly influenced fish metabolism. This finding is in accordance with the hypothesis that the additional stress imposed upon an organism by contaminants can amplify the importance of naturally encountered stresses such as food or habitat limitation.

Highly significant ($p < 0.001$) concentrations of Mn, Fe, Al and Pb, and Ni ($p < 0.02$) were lost when fish were allowed to purge their gut contents. The only effect of feeding was an increase in Ni concentrations in fathead minnows (Table 12). Trace organic concentrations did not differ among treatments (Table 13). The loss of Mn, Fe and Al may well reflect the importance of these elements in sediment composition and their correspondingly high concentrations relative to tissue residues in fish. The loss of Al agrees with the findings for depuration effects on oligochaete tissue residues. While metal physicochemistry was not measured, Pb and Ni are known to associate with Fe and Mn oxyhydroxides. Lead and Ni may have followed pattern of Fe and Mn for this reason.

3.4 Experiment 4 - June 30-July 22, 1988

Purpose: To compare the toxicity of intact sediment cores and homogenized sediment to mayfly nymphs and fathead minnows.

The current MOE method for assembly of the sediment bioassay involves sieving and homogenizing the sediment. This effectively exposes organisms to a uniform dose of contaminants that is in reality a mean dose of the heterogeneously

distributed contaminants. A positive ramification of homogenization is that it most likely results in less variability among replicates than would be observed for intact sediment. In contrast, there are probably many circumstances where the extensive aeration of the sediment also results in a transformation of chemical species to forms that are of greater or lesser bioavailability. One approach to examining the question of how toxicity is influenced by sediment homogenization involves a comparison of endpoints achieved when organisms are exposed to diver-collected cores and to homogenized sediment.

The cores used were acrylic tubes of surface area comparable to the 2L glass jars. Organisms were introduced into the cores and into homogenized sediments from the same site as those where cores were collected.

Eight Hexagenia nymphs (c.a. 40 mg/individual net weight) or 10 juvenile fathead minnows (c.a. 400 mg/individual net weight) were the test organisms. Four replicate diver-collected cores and triplicate jars of homogenized sediment were used for each organism for each of three test substrates. Mortality, biomass changes, and bioaccumulation over three weeks were the endpoints examined. pH and dissolved oxygen were monitored in all chambers.

One sandy and two fine-grained sites in the vicinity of the Toronto Main STP outfall were sampled. Honey Harbour sediments were used for controls. In Site A (fine), intact sediment resulted in higher mortality and poorer growth than homogenized sediment for mayfly nymphs, but did not significantly influence mortality or growth in fathead minnows (Table 14). Intact sediment from Site B (fine) resulted in better growth for mayfly nymphs than homogenized sediment. Mortality was <10% in both treatments. Homogenization resulted in substantial mortality for fathead minnows (87% vs 20% in intact cores). In Site C (sandy) homogenization resulted in higher mortality than in the intact cores for Hexagenia. This was most likely caused by the elimination of the surface layer of fine-grained material (present in intact cores) and therefore, the elimination of suitable substrate for burial and feeding. Homogenization did not effect growth of fathead minnows, and may have ameliorated toxicity as measured by mortality. A point of interest is the variability in response between the two

test species, which suggests differential modes of action of contaminants upon organisms with vastly different metabolic pathways and ecological niche requirements.

The results the chemical analyses of tissue residues for trace metals are illustrated in figures 3 through 7. While there was a tendency for some metals to bioaccumulate to a greater extent in beakers as compared to core exposures, the results were not statistically significant. Considering the high cost of diver collected cores, this result is encouraging, in the sense that the beaker assay may be fairly representative, if not more conservative than intact core bioassays. The finding that intact cores were more toxic to mayflies at station A but not at station B should be perceived as justification to pursue this line of investigation and to broaden the range of sediment tested.

4.1 Chronic BioassaysRecommendation

I strongly urge that the development of chronic, nonlethal bioassays be the central focus of future research. This has begun with a consideration of a 21 day growth-inhibition test with mayflies and fathead minnows. Clearly, mortality is also documented and many other potential endpoints could be adopted.

Since sediment organic content will effect growth, independent of the level of contamination, additional studies on feeding must be performed for fathead minnows in particular. Food availability has been shown to modify the bioaccumulation of pollutants and studies designed to further assess various feeding regimes for chronic bioassays are strongly recommended.

The chironomid full or partial life cycle test is worth pursuing. Survival, changes in biomass, emergence, and reproductive success can all be measured in 10 to 28 days. Several organisms can be introduced into a single chamber (with replication) or individual organisms can be held in test-tube assembly. These two approaches could be compared.

Hyalalela azteca is also amenable to a full or partial life cycle test, but ensuring that individual sensitivity remains constant during culturing may be somewhat more difficult for the amphipod than for the chironomid.

A sensitive and more rapid chronic bioassay measures growth in larval or egg-sac stage fathead minnows. This test has been considered for effluent testing and could be applied to sediment bioassessment.

Another interesting class of endpoints to consider is behavioral in nature. The two most often cited are preference-avoidance and inability (or refusal) to burrow. I have observed oligochaetes to remain on the surface of noxious

sediment. Mayflies also elicit this response, although less frequently. Others have demonstrated reburial failure in amphipods transferred to clean substrates following exposure to contaminated sediments. These endpoints, however, may be more difficult to interpret in terms of their ecological significance than the partial life cycle tests and should be investigated once the former tests have been established.

4.2 Reference Toxicants

Recommendation

At least 1 polar (e.g. Cd) and 1 nonpolar (e.g. PCP) compound should be tested and effect concentrations established. These pretests, or positive controls, can be conducted as EC₁₀s or LC₁₀s in aqueous phase bioassays or developed as a spiked sediment bioassay.

Reference toxicants have been of value in examining seasonal changes in an organism's sensitivity, and changes related to age, reproductive status, and history of exposure to contaminants. In order to ensure that the test organism's response to contaminants is uniform and does not vary among bioassays (or among laboratories), reference toxicants should be incorporated as a pretest in all bioassays. Loss of vigour in cultured organisms has been demonstrated and could confound interpretation of sediment bioassay results. When an organism response does vary, one might justify the application of a correction factor.

4.3 Sediment Manipulation

Recommendation

Further comparisons between intact sediments and homogenized sediments should be initiated to assess the effect of sediment manipulation on acute and chronic endpoints. The effects of sediment storage on toxicity should also be investigated.

Many existing protocols, including the current MOE method for assembly of the sediment bioassay involve sieving and homogenizing the sediment. For many sediments extensive aeration will result in a transformation of chemical species to forms that are of greater or lesser bioavailability. One approach to examining the question of sediment homogenization involves comparisons of toxicity with diver-collected cores. The cores should be of comparable surface area as the glass jars being used for routine testing. Organisms can be introduced into the cores and into sediment which will be sieved and homogenized and collected from the same site as those where cores were retrieved. Since the objective of the bioassay is to assess the extent of contamination of the sediment directly, without the complicating factor of potentially contaminated site water, dechlorinated laboratory water should replace the water in the cores and should be used for the jar test as well. I conducted one such experiment (Section 3.4) and the results should be verified for other sediment types and additional bioassay organisms.

As an alternative to toxicity testing using diver collected cores, the contents of the ponar grab can be "cored" on board with minimal mixing of the test sediment. These pseudo-cores can be returned to the laboratory and subjected to the same bioassessment as the diver-collected cores and the homogenized sediment. Again, it would be useful to conduct the above experiments using a variety of organisms and a broader range of contaminants.

Due to logistical constraints sediment has in the past been stored at 4°C for several months prior to conducting the bioassay. To establish a maximum acceptable storage interval bioassays could be repeated using the same sediment after short and long-term storage (e.g. 2 weeks, 1 month, 2 months, 6 months storage).

4.4 Invertebrate Cultures

Recommendation

To facilitate the development of a multi-species approach to sediment bioassessment, invertebrate cultures should be initiated. The construction of

facilities at MOE for culturing Hyalella azteca, Hexagenia, Chironomus, and oligochaetes should be expanded and actively maintained. Protocols for culturing these invertebrates have been published by, for example, S.G. Lawrence (ed) 1981, as a Canadian Special Publication of Fisheries and Aquatic Sciences #54. It would be worthwhile for a full time technician and/or scientist to adopt responsibility for the maintenance of the cultures.

Since sensitivity to contaminants varies among species and one species is not necessarily the most sensitive to all classes of pollutants, MOE should adopt an approach that incorporates the responses of several taxa. The test organisms should represent different ecological functional groups such as filter-feeders, burrowing infauna, and benthic foraging fish, for example. Chironomus riparius can be easily cultured and used for a partial life cycle test. Collection of sufficient quantities of Hexagenia that are of a prescribed age and size is problematic and labour intensive. Establishing mayfly cultures, then, would be of great value. Hyalella has been successfully maintained in stock cultures and reproduction can be induced easily by controlling the photoperiod. Partial life cycle tests with the Hyalella would be facilitated by establishing laboratory cultures although maintaining consistency in culture vigour must be verified. Oligochaetes can be used to measure reproductive success and contaminant bioavailability as reflected by bioaccumulation. Tubificids are easily maintained in the laboratory.

4.5 Organism Age and Size

Recommendation

I have compared the response of several size classes of mayflies and fathead minnows exposed to contaminated sediments. These experiments should be conducted for all test species.

Sensitivity to contaminants and contaminant bioaccumulation has been shown to vary with age and body size for several taxa. For purposes of tissue analysis, larger organisms provide greater biomass. However, sensitive life stages are

generally early life forms. Increased replication with smaller organisms may satisfy both needs.

4.6 Substrate Properties and Colonization Potential

Recommendation

The range of substrates that the test species can potentially colonize, in the absence of contamination, must be determined. Growth rates in sediment of different organic content should be established.

Organisms such as chironomids and burrowing mayflies exhibit substrate preferences. The absence of Hexagenia from a coarse-sandy sediment, for example, may be due to its inability to thrive in such a substrate, regardless of how pristine it may be. Extremely soft oozy mud may be too unstable a substrate for chironomids. The organic content of a given sediment can be critical for growth and survival. In order to formulate decision criteria for the selection of test organisms based on substrate properties (independent of contamination), it is important to establish the range of substrates that the test species can potentially colonize.

For growth to be used as an endpoint in sediment bioassays, biomass changes should be calibrated against the same range of sediment types for uncontaminated substrates. One approach is to collect clean, fine sediments and perform a serial dilution with fine and coarse sand. These experiments should apply to all test species.

4.7 Sediment to Water Ratio

Recommendation

The effect of increasing or decreasing the currently adopted 1:4 ratio (volume sediment:volume water) should be measured to ascertain that this ratio is adequate.

The most frequently cited sediment to water ratio in a static test is 1:4 (v:v). Experimental manipulation of this ratio may reveal a graded response to the test organism. That is, toxic effects may increase when there is proportionally more sediment (or less water) in the system but may be independent of the ratio at relatively high volumes of water.

4.8 Flow-through, Static-Renewal and Static Water Regimes

Recommendation

A comparison of these three approaches, particularly in the case of chronic tests, would assist in verifying that the observed endpoints best reflect the biological significance of sediment contamination.

Static beaker tests are generally considered to represent a "worst case" or "conservative" scenario. If contaminants increase in the water column during the exposure interval, it is intuitively logical that some organisms will demonstrate toxicity responses that would not be observed in a flow-through system. It is possible, however, that under certain circumstances, flow through conditions may in fact underestimate the potential impacts of the test sediment on some organisms as would be observed in nature.

The build-up of metabolites during a chronic bioassay may complicate the interpretation of the results. A consideration of the static-renewal approach would therefore be of great utility.

4.9 Field Comparison

Recommendation

Following laboratory investigations on the ramifications of various bioassay design options, a thorough program of field testing should be instituted.

Field experiments should be designed to substantiate that laboratory tests are a meaningful representation of field conditions. In situ chambers containing the same species used in the laboratory can be positioned in the sediment. The endpoints measured would be those evaluated in the laboratory sediment bioassay. Careful consideration of chamber design, its introduction into the sediment, water flow and/or exclusion, controls for chamber effects, and other potential sources of artifact is integral to the success of the field validation program.

4.10 Bioaccumulation

Recommendation

It would be extremely beneficial to devote research efforts towards illuminating the relationship between the tissue retention of contaminants and their physiological significance to the exposed organism.

The relationship between the rate and extent of bioaccumulation of contaminants and their physiological effects are poorly understood. As a consequence, the ecological significance of tissue retention of pollutants cannot be established for many contaminants. Such information would be of great value in the establishment and implementation of sediment and biota guidelines.

The following recommendations for establishing a MOE protocol for sediment bioassays should be considered as a framework which will be further modified based upon ongoing research efforts in the development of the assays. It is anticipated that these recommendations will be complemented by recent findings of D. Bedard and A. Hayton, and that an MOE sediment bioassay protocol document will be prepared by these individuals and myself, based on available information. Further, the sediment bioassays should be recognized as an important component of a sediment management strategy that includes extensive biological assessment as well as chemical measurements.

Sample Collection

Sediment is collected by Ponar grab from test and control sites and the surficial 3 cm are placed in polyethylene-lined plastic buckets. These are returned to the laboratory and may be stored at 4°C for no more than 2 weeks.

Preparation of the Bioassay Chamber

In the laboratory, sediment is air-sieved through a 2 mm mesh to remove large particles, stones and other debris, and is then thoroughly homogenized. For the experimental bioassays, 2 L glass widemouth jars, acid washed and hexane rinsed, are filled to a depth of 3 cm with sediment (surface area = 100 cm²). Dechlorinated water is gently added at a ratio of 4:1 water to sediment (v:v). Care must be taken to avoid having sediments adhere to the glass above the 3 cm level. Resuspended sediment is allowed to settle for 5 hours (more conservative) or 24 hours (logistically more practical), after which the overlying water is aerated for 1 hour by inserting airlines through the lids of the glass jars and securing the lids in place. Water loss due to evaporation is replaced as necessary in order to maintain a water to sediment ratio of 4:1. pH and dissolved oxygen in the overlying water are monitored routinely for the duration of the experiment.

Test Organisms

Presently, organisms for the experimental bioassays are three to four month old juvenile fathead minnows (Pimephales promelas) (acquired from OMOE Rexdale laboratory cultures) weighing approximately 0.5 gm wet weight and first year mayfly nymphs (Hexagenia limbata) weighing about 30 mg wet weight, collected at a clean reference site (Honey Harbour, Georgian Bay, Ontario) or cultured in the laboratory. Nymphs collected from the field can be maintained in the laboratory in glass aquaria containing reference site sediments with gentle aeration of the overlying water. The aquaria may be kept at <10°C, and gradually brought to 20°C as mayflies are required. Alternatively, mayflies can be cultured in the laboratory and harvested upon demand. Depending upon culture availability, second instar Chironomus tentans may also be used.

Several hours prior to experimental exposure, mayflies are removed from their aquaria by sieving small volumes of sediment through a 500 µm mesh. Nymphs are randomly allocated to beakers containing dechlorinated water until each beaker has 10 individuals of similar size (c.a. 25 mg/ individual, wet weight). The nymphs are weighed after blotting them on several layers of "Kim Wipes" or acid rinsed filter papers to remove adhering water. Alternately, when a large number of assays are run concurrently, 5 to 7 aliquots of 10 individuals each can be weighed, and the mean biomass used as an estimate of initial biomass.

Similarly, juvenile fathead minnows are randomly allocated to beakers containing dechlorinated water until each beaker has 10 individuals of similar size (c.a. 0.5 gms/individual, wet weight). Wet weight of each group of 10 individuals is measured, or 5 to 7 representative samples of ten individuals each can be weighed, as above.

Chironomid length, weight, and head capsule widths can be measured for a representative subsample of individuals at the onset of the experiment.

Conducting the Bioassay

A minimum of 3 replicate jars of each bioassay organism. Organisms are added when 5 to 24 hours of settling followed by 1 hour of aeration have elapsed. Aeration persists for the duration of the 21 day exposure interval. All control and test jars are maintained at 20°C (a water bath may be required) at ambient light.

During the 21 day bioassay period, mortality is noted and dead organisms are removed. The presence of mayfly exuvia should also be recorded. As stated above, pH and dissolved oxygen are monitored and water lost due to evaporation is replaced. At day 21, organisms are removed from the bioassay jars by passing the entire contents of each jar through a 500 µm sieve and retrieving the biota. Recoveries are noted and the remaining organisms are reweighed and measured using the procedure described above.

Chemical Analyses

Mayflies and fathead minnows are placed in hexane-rinsed aluminum-foil for analysis of organic residues, or in plastic (e.g. "whirlpak" bags) for metal analysis. Depuration is currently not recommended for trace organics, however, the importance of gut clearance prior to sample analyses may be of relevance for some trace metals. If feasible, a subsample of test organisms are held in dechlorinated tap water for 24 hours to permit the depuration of gut contents. Samples are frozen until chemical treatment can be conducted. Biota must be handled gently, using teflon-coated or nylon forceps.

Concurrent with biota analyses, measurement of sediment physical and chemical characteristics should be conducted. Aliquots of the homogenized sediment can be analyzed for trace metals, a range of trace organics (e.g. PCBs, pesticides), the extent of oil and grease contamination, particle size distribution, and organic carbon and other measures of nutrient status. The selection of the parameters of interest will necessarily be site specific and depend upon available information on the pollutant source(s).

Data Interpretation

Mortality in test sediment is compared with controls. Control mortality should not exceed 10%. Growth inhibition, indicative of chronic exposure to unfavourable sediment, can be identified by comparing biomass changes of test organisms with those of control organisms. Analysis of variance should be applied to appropriately transformed data (if necessary to achieve a normal distribution), particularly if data are expressed in the form of ratios (e.g. percent reduction in growth of test organisms relative to controls). Similar statistical analyses are used to examine differences in tissue retention of contaminants following the 21 day exposure to sediment that vary in their chemical composition. Significance at $p < 0.05$ is proposed as an indicator of sediment eliciting unacceptable biological responses.

In conjunction with developing a protocol for the MOE, it has been an advantage to have the opportunity to discuss many of the growing concerns over sediment bioassays with others doing research in this area. The conference on Environmental Risk: Recognition, Assessment and Management, coordinated by the Society of Environmental Toxicology and Chemistry (SETAC) in Pensacola, Florida November 9 to 12, 1987 was of great benefit. Members of the American Society for Testing Materials also held a special session on sediment bioassay protocols. The meetings were of assistance in setting research priorities.

As a consequence of attending SETAC, I was made aware of the research efforts being conducted at North Texas State University (NTSU). After speaking with the coordinator of the NTSU research and the Liaison Officer for this project, it was agreed that visiting the NTSU facilities and gaining access to their voluminous "grey" literature would be a worthwhile investment.

As a direct result of that visit, a strategy for culturing and testing chironomids was prepared, and plans for future research on lab/field comparisons were formulated. Many of the documents acquired for MOE are an excellent source of information that would otherwise have been virtually inaccessible.

Attendance at the first International Conference on Environmental Bioassay Techniques and Their Application, Lancaster England, was most worthwhile. The title alone demonstrates the direct relevance of this meeting to my own and MOE research objectives. In becoming familiar not only with the ongoing research but with the individuals sharing common research goals, the sediment bioassay techniques that are currently recommended for development were substantiated.

Subsequent to the completion of the experiments under the RAP PDF03, additional studies were conducted on bioassay duration and the effects of chemical treatment on sediment toxicity. This work was done in 1989 in support of the Hamilton Harbour Remedial Action Plan and in conjunction with the National Water Research Institute. A paper has been submitted to a peer review journal for consideration.

Anderson, R.L., C.T. Walbridge and J.T. Fiandt, 1980. Survival and growth of Tanytarsus dissimilis (Chironomidae) exposed to copper, cadmium, zinc and lead. Arch. Environ. Contam. Toxicol. 9:329-335.

Anderson, R.V., 1978. The effects of lead on oxygen uptake in the crayfish Orconectes virilis (Hagen). Bull Environ. Contam. Toxicol. 20:394-400.

Bahnick, D.A., Swenson, W.A., Markee, T.P., Call, D.J. Anderson, C.A., and Morris, R.T., 1980. Development of bioassay procedures for defining pollution of harbour sediments. Final Report of U.S. EPA Project No. R804918-01. Center for Lake Superior Environmental Studies, University of Wisconsin, Superior, Wisconsin, 189pp.

BEAK Consultants, 1987. Development of Sediment Quality Objectives. Phase 1 - Options, Final Report. Report to the Ontario Ministry of the Environment.

Biesinger, K.E. and Christensen, G.M., 1972. Effects of various metals on survival, growth, reproduction, and metabolism of Daphnia magna. J. Fish. Res. Bd. Can., Volume 29, pp. 1691-1700.

Birge, W.J., J.A. Black, A.G. Westerman and P.C. Francis, 1987. Toxicity of sediment-associated metals and freshwater organisms: Biomonitoring Procedures. In: K.L. Dickson, A.W. Maki, and W.A. Brungs (Eds), Fate and Effects of Sediment-Bound Chemicals in Aquatic Systems Perg. Press, N.Y. pp. 199-218.

Brkovic-Popovic and Popovic, 1977. Affects of heavy metals on survival and respiration rate of tubificid worms: Part II - Effects on respiration rate. Environ. Pollut. 13:65-72.

Brungs, W.A., J.R. Geckler and M. Gast, 1976. Acute and chronic toxicity of copper to the fathead minnow in a surface water of variable quality. Water Research 10:37-43.

Buikema, A.L., Jr., Niederlehner, B.R., and Cairns, J., Jr., 1982. Biological monitoring. Part IV - toxicity testing. Water Research. 16:239-262.

Cairns, M.A., Nebeker, A.V., Gakstatter, J.H., and Griffis, W.L. (1984), Toxicity of copper-spiked sediments to fresh-water invertebrates. Environ. Toxicol. Chem. 3:435-445.

Craig, G.R., 1984. Bioassessment of sediments: Review of previous methods and recommendations for future test protocols. Report to the Ontario Ministry of the Environment, August 1984 2252.1.

deMarch, B.G.E., 1979. Reproduction and growth impairment tests with the amphipod Hyalella azteca. Toxicity Tests for Freshwater Organisms. Canadian Special Publication of Fisheries and Aquatic Sciences Volume 44, pp.1-10.

Francis, P.C., W.J. Birge, and J.A. Black, 1984. Effects of cadmium-enriched sediment on fish and amphibian embryo-larval stages. Ecotoxicol. Environ. Safety 8:378-387.

Gagnon, J.E. and Beaton, A.M., 1969. Studies on the effects of dredged materials from selected Great Lakes harbours on plankton and benthos. Special report No. 8, Center for Great Lakes, University of Wisconsin-Milwaukee, pp.80.

Gagnon, J.E. and Beaton, A.M., 1971. Procedures for determining the effects of dredged sediments on biota-benthos viability and sediment selectivity tests. J. Water Pollut. Control Fed. 43:392-398.

Friesen, M.K., T.D. Galloway, J.F. Flannagan, 1983. Toxicity of the insecticide permethrin in water and sediment to nymphs of the burrowing mayfly Hexagenia rigida (Ephemeroptera: Ephemeridae), Can. Ent. 115:1007-1014.

Hansen, D.J., G.R. Gardner and P.F. Rogerson, 1987. Effects of sediments from new Bedford Harbour, MA, on survival and reproduction of sheephead

minnows, Cyprinodon variegatus. Environmental Risk: Recognition, assessment and management. SETAC, abstracts p. 150.

Hatakeyama, S. and M. Yasuno, 1981. A method for assessing chronic effects of toxic substances on the midge, Paratanytarsus parthenogeneticus - Effects of copper. Arch. Environ. Contam. Toxicol. 10:705-713.

Ingersoll, C.G. and M.K. Nelson, 1987. Methods for conducting chronic sediment toxicity tests with Chironomus riparius. National Fisheries Contaminant Research Center SOP B5.43. USFWS, Columbia, MO.

Ingersoll, C.G., M.K. Nelson, E.K. Henry and K.S. Srigley, 1987. Effects of long-term exposure to contaminated sediment on midge and amphipods. Environmental Risk: Recognition, assessment, and management, SETAC, abstracts, p.151.

Keilly, T., D. White and P. Landrum, 1988. Responses of freshwater oligochaetes to sediment contamination. IAGLR, Abstracts of the 31st Conference of Great Lakes Research, May 17-20, 1988.

Krantzberg, G., 1987. A study of the role of biota and a biotic factors in modifying metal accumulation by Chironomus (Diptera: chironomidae). Ph.D. Thesis, University of Toronto, Ontario, Canada. 258 pp.

Krantzberg, G., and P.M. Stokes, 1989. Metal regulation, tolerance, and body burdens in the larvae of the genus Chironomus. Can. J. Fish Aquat. Sci. 46: 389-398.

Kosalwat, P. and A.W. Knight, 1987. Chronic toxicity of copper to a partial life cycle of the midge Chironomus decorus. Arch. Environ. Contam. Toxicol. 16:283-290.

Lamberson, J.O. and Swartz, R.C., 1985. Use of bioassays in determining the toxicity of sediment to benthic organisms. U.S. Corvallis Environmental Research Lab. Newport, EPA/600/D-85/221. PB86-110681 pp.19. .

LeBlanc, G.A., and D.C. Surprenant, 1985. A method of assessing the toxicity of contaminated freshwater sediments. In: R.D. Cardwell, R. Purdy, and R.C. Bahner (Eds), Aquatic Toxicology and Hazard Assessment: 7th Symp., ASTM STP 854, pp. 269-283.

Lee, G.F., M.D. Piwoni, J.M. Lopez, G.M. Mariani, J.S. Richardson, D.H. Homer, and F. Saleh, 1975. Research study for the development of dredged material disposal criteria. Dredged Material Research Program, Contract Report D-75-4.

Leonhard, S.L. (1979). Tests for the crayfish Orconectes virilis. Toxicity Test for Freshwater Organisms. Canadian Special Publication of Fisheries and Aquatic Sciences. Volume 44, pp.82-90.

Lomas, T. and G. Krantzberg, 1988. Contaminated Sediments Assessment Volume II - Sediment Bioassays. Prepared for the Canada-Ontario Polluted Sediment Committee, Feb. 1988. 9p.

Luoma, S.N., 1983. Bioavailability of trace metals to aquatic organisms - a review. Sci Total Environ. 28:1-22.

Mac, J. Michael, Edsall, C. Carol, Hesselberg, J. Robert, Sayers, E. Richard Jr., 1984. Flow-through bioassay for measuring bioaccumulation of toxic substances from sediment. U.S. Fish and Wildlife Service. Great Lakes Fishery Laboratory. EPA-905/3 - 84-007 pp.17.

Macdonald, W.A., 1979. Testing embryonic and larval life stages of fish. Toxicity Tests for Freshwater Organisms. Canadian Special Publication of Fisheries and Aquatic Sciences. Volume 44, pp.131-138.

Malueg, K.W., Schuytema, G.S., Gakstatter, J.H., and Krawczyk, D.F., 1983. Effect of Hexagenia on Daphnia response in sediment toxicity tests. Environ. Tox. Chem. 2:73-82.

Malueg, K.W., Schuytema, G.S., Gakstatter, J.H., and Krawczyk, D.F., 1984. Toxicity of sediments from three metal-contaminated areas. Environ. Tox. Chem. 3:279-291.

Marking, L.L., V.K. Dawson, S.L. Allen, T.D. Bills and J.J. Rach, 1981. Biological activity and chemical characteristics of dredge material from ten sites on the upper Mississippi River, Summary Report. U.S. Fish and Wildlife Service, LaCross, WI.

McGreer, E.R., 1979. Sublethal effects of heavy metal contaminated sediments on the bivalve Macoma balthica (L). Mar. Poll. Bull. 10:259-262.

Mcleese, D.W., L.E. Burridge, and J. Van Dinter, 1982. Toxicities of fine organochlorine compounds in water and sediment to niereis virens. Bull. Environ. Contam. Toxicol. 28:216-220.

Mcleese, D.W. and Metcalfe, 1980. Toxicities of eight organochlorine compounds in sediment and seawater to Crangon septemspinosa. Bull. Environ. Contam. Toxicol. 25:921-928.

McMurtry, M.J., 1982. Substrate selection by tubificid oligochaetes. University of Toronto, M.Sc. Thesis 115pp.

Munawar, M., A. Mudroch, I.F. Munawar and R.L. Thomas, 1983. The impact of sediment-associated contaminants from the Niagara River and mouth on various size assemblages of phytoplankton. J. Great Lakes Res. 9:303-313.

Munawar, M., R.L. Thomas, H. Shear, P. McKee, and A. Mudroch, 1984. An Overview of Sediment - Associated Contaminants and their Bioassessment. Can. Tech. Report Fish Aquatic Sci. No. 1253.

Nebeker, A.V., M.A. Cairns, J.H. Gakstater, K.W. Malueg, G.S. Schuytema and D.F. Krawczyk, 1984. Biological methods for determining toxicity of contaminated freshwater sediments to invertebrates. Environ. Toxicol. Chem. 3:617-630.

Nebeker, N.V., S.T. Onjukka, M.A. Cairns and D.F. Krawczyk, 1986. Survival of Daphnia magna and Hyalella azteca in cadmium-spiked water and sediment. Environ. Toxicol. Chem. 5:933-935.

Nebeker, A.V. and F.A. Puglisi, 1974. Effects of polychlorinated biphenyls (PCB's) on survival and reproduction of Daphnia, Gammarus, and Tanytarsus. Trans. Amer. Fish Soc. 103:722-728.

Nebeker, A.V., C. Savonen, R.J. Baker and J.K. McCrady, 1986. Effects of copper, nickel and zinc on the life cycle of the caddisfly Clistoronia magnifica. Environ. Toxicol. Chem. 3:645-649.

Phelps, H.C., J.T. Marey, W.H. Pearson and C.W. Apts, 1983. Clam burrowing behaviour: Inhibition by copper-enriched sediment. Mar. Poll. Bull. 14:452-455.

Powlesland C. and J. George, 1986. Acute and chronic toxicity of nickel to larvae of Chironomus riparis. Environ. Pollution 42:47-64.

Prater, B.L. and Anderson, M.A., 1977a. A 96-hour bioassay of Duluth and Superior harbour basins (Minnesota) using Hexagenia limbata, Asellus communis, Daphnia magna, and Pimephales promelas as test organisms. Bull. Environ. Contam. Toxicol. 18:159-168.

Prater, B.L. and Anderson, M.A., 1977b. A 96-hour bioassay of Otter Creek, Ohio. J. Water Poll. Control Fed. 49:2099-2106.

Prater, B. and Hoke, R.A., 1980. A method for the biological and chemical evaluation of sediment toxicity. Contaminants and Sediments Volume 1, pp.483-499. Ann Arbor Science Publishers.

Scherer, E., 1979. Avoidance testing for fish and invertebrates. Toxicity Tests for Freshwater Organisms. Canadian Special Publication of Fisheries and Aquatic Sciences. Volume 44, pp.160-170.

Shuba, P.J., H.E. Tatem, and J.H. Carroll, 1978. Biological assessment methods to predict the impact of open-water disposal of dredged material. Dredged Material Research Program Tech. Rep. D-78-50.

Swartz, R.C., DeBen, W.A., and Cole, F.A., 1979. A bioassay for the toxicity of sediment to marine macrobenthos. J. Water Pollut. Control Fed. 51, pp.944-950.

Swartz, R.C., DeBen, W.A., Jones, J.K. P., Lamberson, J.O., and Cole, F.A., 1985a. Phoxocephalid amphipod bioassay for marine sediment toxicity. Aquatic toxicology and hazard assessment. Seventh symposium, ASTM STP 854. pp.284-307.

Swartz, R.C., Ditsworth, G.R., Schults, D.W., and Lamberson, J.O., 1985b. Sediment toxicity to a marine infaunal amphipod: Cadmium and its interaction with sewage sludge. Marine Environ. Res. 18:133-153.

Swartz, R.C., D.W. Schults, E.R. Ditsworth and W.A. DeBen, 1984. Toxicity of sewage sludge to Rhepoxynius abronius, a marine benthic amphipod. Arch. Environ. Contam. Toxic. 13:207-216.

Tsai, L., J. Welch, K. Chang, J. Schaefer, and L.E. Cronin, 1979. Bioassay of Baltimore Harbour Sediments. Estuaries 2:141-153.

USEPA, 1988. Preliminary report on the toxicity and chemistry of sediments from Toronto and Toledo Harbours. Preliminary draft, Sediment Criteria Team, Freshwater Division, Corvallis Environmental Research Lab.

U.S. Environmental Protection Agency/U.S. Army Corps of Engineers, 1977. Ecological evaluation of proposed discharge of dredged materials into ocean waters. Environmental Effects Laboratory, U.S. Army Corps of Engineers, Waterways Experimental Station, Vicksburg, MS.

Wentzel, R.A., McIntosh, A., and Atchison, G., 1977. Sublethal effects of

heavy metal contaminated sediment on midge larvae (Chironomus tentans).
Hydrobiol. 56:153-156.

Wentzel, R., A. McIntosh, and W.P. McCafferty, 1978. Emergence of the midge Chironomus tentans when exposed to heavy metal contaminated sediment. Hydrobiol 57:195-196.

White, D.S., 1984. Redistribution of sediment-bound toxic organics by benthic invertebrates. In: The cycling of toxic organic substances in the Great Lakes Ecosystem. University of Michigan, University of Minnesota, Michigan State University, Argonne National Lab and Oak Ridge National Lab. Annual Report to NOAA pp.76-78.

Wiederholm, T., A. Wiederholm, and G. Milbrink, 1987. Bulk sediment bioassays with five species of freshwater oligochaetes. Water Air Soil Poll. 36:131-154.

TABLE 1: Marine and Freshwater Organisms Used
in Acute Toxicity Tests with Sediments

SPECIES		DURATION (d=days)	APPARATUS	REFERENCE
<u>MARINE</u>				
Amphipods	<u>Rhepoxynius abronius</u>	10d	Static, flowthrough	Swartz et al. 1979
	<u>Rhepoxynius abronius</u>		Static	Swartz et al. 1984
	<u>Parahausterius sp.</u>		Static	Swartz et al. 1985b
	<u>Parahausterius sp.</u>	10d	Static	Shuba et al. 1978
Copepods	<u>Acartia tonsa</u>	4d	Static	Shuba et al. 1978
	<u>Tigriopus californicus</u>			Shuba et al. 1978
Isopods	<u>Sphaeroma</u> <u>quadridentatum</u>	10d	Static	Shuba et al. 1978
Shrimp	<u>Palaemonetes pugio</u>	4-15d	Static	Shuba et al. 1978
	<u>P. vulgaris</u>			
	<u>Crangon septemspinosa</u>	4-12d	Static	Mcleese & Metcalfe 1980
Polychaetes	<u>Nereis virens</u>	4-12d	Static	Mcleese et al. 1982
	<u>Glycinde picta</u>	10d	Flowthrough	Swartz et al. 1979
Bivalves	<u>Macoma inquinata</u>	10d	Flowthrough	Swartz et al. 1979
	<u>Protothaca staminea</u>	10d	Flowthrough	Swartz et al. 1979
	<u>Mya arenaria</u>	2d	Static	Tsai et al. 1979
Fishes	<u>Fundulus heteroclitus</u>	2d	Static	Tsai et al. 1979
	<u>Leiostomus xanthurus</u>			

TABLE 1: Marine and Freshwater Organisms Used
in Acute Toxicity Tests with Sediments (Cont'd)

SPECIES	DURATION (d=days)	APPARATUS	REFERENCE	
FRESHWATER				
Water Fleas <u>Daphnia magna</u>	2d	Recycling	Malueg et al. 1983	
	<u>D. magna</u>	2d	Static	Cairns et al. 1984
	<u>D. magna</u>	3d	Recycling	Prater & Hake 1980
	<u>D. magna</u>	3d	Recycling	Prater & Anderson 1977
	<u>D. magna</u>	2d	Recycling Static	USEPA 1988
Amphipods <u>Hyallela azteca</u>	10d	Static	Nebeker et al. 1984	
	<u>Pontoporeia affinis</u>	3d	Static	Nebeker et al. 1986
	<u>Gammarus lacustris</u>	10d	Static	Nebeker et al. 1984
Oligochaetes unspecified	3d	Static	Keilly et al. 1988	
Mayflies <u>Hexagenia limbata</u>	10d	Static	Lomas & Krantzberg 1988	
	<u>Hexagenia limbata</u>	5d	Recycling	Malueg et al. 1983
	<u>Hexagenia limbata</u>	3d	Recycling	Prater & Anderson 1972
	<u>Hexagenia rigida</u>	7d	Static	Friesen et al. 1983
Midges <u>Chironomus</u>	10d	Static	Nebeker et al. 1984	
	<u>Chironomus</u>	10d	Static	Cairns et al. 1984
	<u>Chironomus</u>		Static	Marking et al. 1981
	<u>Chironomus</u>		Static	Gagnon & Beaton 1971
Crayfish <u>Orconectes virilis</u>	10d	Static	Leonhard 1979	
		Flowthrough		
Oligochaetes <u>Tubifex tubifex</u>				
	<u>Limnodrilus</u>			
	<u>hoffmeisteri</u>	3d	Static	McMurtry 1982

TABLE 1: Marine and Freshwater Organisms Used
in Acute Toxicity Tests with Sediments (Cont'd)

SPECIES	DURATION (d=days)	APPARATUS	REFERENCE
<u>FRESHWATER</u> (Cont'd)			
Fish <u>Micropterus salmoides</u> <u>Carassius auratus</u> <u>Salmo gairdneri</u> <u>Pimephales promelas</u> <u>P. promelas</u> <u>P. promelas</u> <u>C. auratus</u> <u>M. salmoides</u>	4d 10d 3d 10d 6-7d	Static Static Recycling Flowthrough Static	Birge et al. 1987 Lomas & Krantzberg 1988 Prater & Anderson 1977b Mac et al. 1984 Francis et al. 1984

TABLE 2: Chronic Endpoints Employed for Various
Organisms Used in Sediment Bioassays

ENDPOINT	ORGANISM	AUTHOR
Growth Inhibition	<p>Midges, <u>Chironomus decorus</u></p> <p><u>C. tentans</u></p> <p><u>C. tentans</u></p> <p><u>C. riparis</u></p> <p><u>Paratanytarsus parthenogeneticus</u></p> <p><u>Tanyarsus dissimilis</u></p> <p>Oligochaetes, <u>Tubifex tubifex</u>,</p> <p><u>Potamothenix hammoniensis</u></p> <p><u>Limnodrilus hoffmeisteri</u></p> <p><u>L. udekemianus</u>, <u>L. clapparedianus</u></p> <p>Mayflies, <u>Hexagenia limbata</u></p> <p>Amphipods, <u>Hyalella azteca</u></p> <p>Fish, <u>Pimephales promelas</u></p> <p><u>Salmo gairdneri</u></p> <p><u>Coregonus clupeaformis</u></p> <p><u>P. promela</u></p>	<p>Kosalwat & Knight 1987</p> <p>Wentzel et al. 1977</p> <p>Nebeker et al. 1984</p> <p>Powlesland & George 1986</p> <p>Hatakeyama & Yusuno 1981</p> <p>Anderson et al. 1980</p> <p>Wiederholm et al. 1987</p> <p>Krantzberg (unpubl. data)</p> <p>de March 1979</p> <p>Brungs et al. 1976</p> <p>Macdonald 1979</p> <p>LeBlanc & Suprenant 1985</p>
Full or Partial Life Cycle Completion	<p>Waterfleas <u>Daphnia magna</u></p> <p><u>D. magna</u></p> <p>Midges <u>C. tentans</u></p> <p><u>P. parthenogeneticus</u></p> <p><u>P. parthenogeneticus</u></p> <p><u>C. riparis</u></p> <p><u>C. tentans</u></p> <p><u>C. riparis</u></p>	<p>Nebeker et al. 1986</p> <p>Biesinger & Christensen 1972</p> <p>Wentzel et al. 1978</p> <p>Hatakeyama & Yasuno 1981</p> <p>LeBlanc & Suprenant 1985</p> <p>Powlesland & George 1986</p> <p>Nebeker et al. 1988</p> <p>Ingersoll et al. 1987</p>

TABLE 2: Chronic Endpoints Employed for Various
Organisms Used in Sediment Bioassays (Cont'd)

ENDPOINT	ORGANISM	AUTHOR
Full or Partial Life Cycle (Cont'd)	Amphipods <u>H. azteca</u> <u>Gammarus pseudolimnaeus</u> Oligochaetes, <u>Niad</u> Caddisflies, <u>Clistoronia magnifica</u> Fish <u>Cyprinodon variegatus</u>	Ingersoll et al. 1987 Nebeker & Puglisi 1974 Rodgers (pers. comm.) Nebeker et al. 1984b Hansen et al. 1987
Burrowing	Oligochaetes, <u>Stylodrilus</u> <u>heringianus</u> , <u>Potamothrix</u> <u>vej dovski</u> , <u>Limnodrilus</u> <u>hoffmeisteri</u> Clams, <u>Protothaca staminea</u> Amphipods, <u>Rhepoxynius abronius</u>	White 1984 Phelps et al. 1983 Swartz et al 1985
Avoidance	Amphipods, <u>Pontoporeia hoyi</u> <u>Gammarus</u> Midges, <u>Chironomus</u> <u>C. tentans</u> Oligochaetes <u>Tubifex tubifex</u> <u>L. hoffmeisteri</u> Bivalves <u>Macoma balthica</u> Fish <u>Coregonus clupeaformis</u> <u>Salmo gairdneri</u>	Gagnon & Beeton 1971 Gagnon & Beeton 1971 Wentzel et al. 1977 McMurtry 1982 McGreer 1979 Scherer 1979
Respiration	Oligochaetes, <u>T. tubifex</u> Crayfish <u>Orconectes virilis</u>	Brkovic-Popovic and Popovic 1977 Anderson et al. 1978

Table 3. Sample locations and composition of sediment used in bioassays. Numbers refer to sampling Environment Ontario Sampling stations.

SITE	LATITUDE	LONGITUDE	DEPTH (m)	% SAND ($>44 \mu\text{m}$)	% SILT ($42.27-3.73$)	% CLAY ($3.73-0.17 \mu\text{m}$)	pH	% LOI ¹	TOC ²
TORONTO STP 1	43° 39.26'	79° 18.60'	6.0	79.5	16.2	2.0	7.05	2.2	16.0
TORONTO STP 6	43° 38.76'	79° 18.99'	7.0	58.0	33.2	3.9	6.75	2.2	13.0
ST. MARYS RIVER 3	46° 30.85'	84° 14.90'	1.3	71.6	18.3	1.5	6.66	3.5	19.0
ST. MARYS RIVER 4	46° 31.70'	84° 14.18'	2.8	39.0	49.4	4.2	6.60	12.0	84.0
RICE LAKE 4	44° 16.45'	78° 19.34'	1.4	87.6	9.2	1.4	7.17	3.5	19.0
RICE LAKE 14	44° 08.90'	78° 12.39'	4.0	56.5	33.7	6.0	7.37	23.0	110.0
HONEY HARBOUR (Southern Georgian Bay)			4.0	15.8	73.3	9.9		7.5	35.0

¹percent loss on ignition;

²total organic carbon ($\text{mg}\cdot\text{g}^{-1}$)

TABLE 4
Contaminant Concentrations in Sediments Used For Bioassays
(All values are in µg/g (dry weight) unless otherwise noted)

SITE	CU	CR	HG	CD	FE	PB	ZN	AS	MN	AL	NI	SOLV. EXTRACT.	PCB (ng/g)
TORONTO STP 1	38	34	0.09	0.76	12000	82	77	2.0	200	4700	9.7	3059	nd ¹
TORONTO STP 6	32	44	0.46	1.10	12000	18	65	1.6	220	5900	11.0	2387	nd
ST. MARYS RIVER 3	21	27	0.07	0.30	13000	25	80	2.6	140	3900	8.0	n/a ²	n/a
ST. MARYS RIVER 4	86	140	0.30	nd	76000	100	370	15.0	770	8320	28.0	n/a	n/a
RICE LAKE 4	22	21	0.11	nd	10000	16	61	1.1	490	4000	8.9	1900	6140
RICE LAKE 14	63	61	0.32	3.10	24000	91	180	3.9	710	15000	27.0	5770	1160
HONEY HARBOUR	21	44	0.08	1.10	31000	37	110	4.3	900	19000	25.0	n/a	n/a

¹not detectable;

²not analyzed

TABLE 5 Effect of Settling Time on Growth of Hexagenia limbata
Values in parentheses are standard deviations

MEASUREMENT	SEDIMENT							
	TORONTO STP			RICE LAKE			CONTROL	
	SETTLING TIME							
	5h	24h	120h	5h	24h	120h	5h	24h
Initial Biomass (mg)	22 (3)	19 (4)	19 (1)	16 (2)	20 (4)	17 (2)	22 (2)	23 (3)
Percent Biomass Change (Day 10)	-3 (0.9)	-2 (0.1)	25 (7)	17 (2)	30 (6)	25 (10)	103 (8)	113 (3)
Growth Inhibition (Day 10) ¹	103	102	78	93	73	32	n.a. ²	n.a.
Percent Mortality (Day 10)	4 (5)	6 (7)	12 (10)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Percent Biomass Change (Day 21)	4 (9)	18 (10)	69 (3)	42 (9)	77 (25)	129 (50)	163 (10)	159 (15)
Growth Inhibition (Day 21)	97	89	57	74	52	19	n.a.	n.a.
Percent Mortality (Day 21)	7 (9)	12 (10)	12 (10)	8 (11)	8 (11)	0 (0)	0 (0)	0 (0)

¹ relative to controls

² not applicable

and exposure duration. All values are ug/g* unless otherwise noted.

STATION	SETTLING TIME	BIOASSAY DURATION	% LIPID	HCB	HEPTA- CHLOR	HEPTA- CHLOR EPOXIDE	ALDRIN	α-BHC	β-BHC	ALPHA CHLORDANE
RICE LAKE	5	10	0.650	0.013	0.000*	0.000	0.000	0.000	0.000	0.000
TORONTO	5	10	0.750	0.005	0.000	0.000	0.009	0.004	0.180	0.100
CONTROL	5	10	1.280	0.100	0.000	0.000	0.003	0.004	0.000	0.000
RICE LAKE	24	10	1.000	0.003	0.000	0.000	0.000	0.000	0.000	0.000
TORONTO	24	10	1.230	0.007	0.011	0.000	0.030	0.007	0.000	0.000
RICE LAKE	120	10	0.480	0.004	0.000	0.000	0.009	0.005	0.000	0.000
TORONTO	120	10	0.420	0.000	0.101	0.000	0.000	0.000	0.000	0.000
RICE LAKE	5	21	0.540	0.003	0.000	0.000	0.000	0.000	0.000	0.000
TORONTO	5	21	1.020	0.002	0.000	0.000	0.000	0.003	0.000	0.001
RICE LAKE	24	21	0.890	0.002	0.000	0.000	0.000	0.000	0.000	0.000
TORONTO	24	21	1.490	0.000	0.000	0.000	0.000	0.015	0.000	0.012
RICE LAKE	120	21	1.080	0.005	0.000	0.000	0.000	0.005	0.000	0.021
TORONTO	120	21	1.580	0.008	0.000	0.000	0.009	0.008	0.000	0.000

* Values listed as 0.000 are below detection.

STATION	SETTLING TIME	BIOASSAY DURATION	GAMMA CHLORIDANE	pp-DDE	op-DDT	PCB	LHC9/ LIPID	HEPTACHLOR/ LIPID	HCE/ LIPID	ALDRIN/ LIPID	a-BHC/ LIPID	b-BHC/ LIPID
RICE LAKE	5	10	0.000	0.000	0.000	0.26	0.020	0.000	0.000	0.000	0.000	0.000
TORONTO	5	10	0.000	0.007	0.000	0.00	0.007	0.000	0.000	0.012	0.005	0.240
CONTROL	5	10	0.000	0.000	0.000	0.00	0.078	0.000	0.000	0.002	0.003	0.000
RICE LAKE	24	10	0.000	0.000	0.000	0.66	0.003	0.000	0.000	0.000	0.000	0.000
TORONTO	24	10	0.000	0.000	0.000	0.02	0.006	0.006	0.000	0.024	0.006	0.000
RICE LAKE	120	10	0.000	0.000	0.000	0.00	0.006	0.000	0.000	0.019	0.010	0.000
TORONTO	120	10	0.000	0.000	0.000	0.09	0.000	0.240	0.000	0.000	0.000	0.000
RICE LAKE	5	21	0.000	0.000	0.000	0.52	0.006	0.000	0.000	0.000	0.000	0.000
TORONTO	5	21	0.002	0.000	0.000	0.02	0.002	0.000	0.000	0.000	0.003	0.000
RICE LAKE	24	21	0.000	0.000	0.000	0.32	0.002	0.000	0.000	0.000	0.000	0.000
TORONTO	24	21	0.000	0.000	0.057	0.00	0.000	0.000	0.000	0.000	0.010	0.000
RICE LAKE	120	21	0.000	0.000	0.000	0.55	0.005	0.000	0.000	0.000	0.005	0.000
TORONTO	120	21	0.000	0.000	0.000	0.09	0.004	0.000	0.000	0.006	0.004	0.000

STATION	SETTLING TIME	BIOASSAY DURATION	ALPHA CHLORDANE/ LIPID	GAMMA CHLORDANE/ LIPID	PP-DDE LIPID	OP-DDT/ LIPID	PCB/ LIPID
RICE LAKE	5	10	0.000	0.000	0.000	0.000	0.394
TORONTO	5	10	0.133	0.000	0.008	0.000	0.000
CONTROL	5	10	0.000	0.000	0.000	0.000	0.000
RICE LAKE	24	10	0.000	0.000	0.000	0.000	0.880
TORONTO	24	10	0.000	0.000	0.000	0.000	0.016
RICE LAKE	120	10	0.000	0.000	0.000	0.000	0.000
TORONTO	120	10	0.000	0.000	0.000	0.000	0.214
RICE LAKE	5	21	0.000	0.000	0.000	0.000	0.863
TORONTO	5	21	0.001	0.002	0.000	0.000	0.020
RICE LAKE	24	21	0.000	0.000	0.000	0.000	0.386
TORONTO	24	21	0.008	0.000	0.000	0.038	0.000
RICE LAKE	120	21	0.020	0.000	0.000	0.000	0.518
TORONTO	120	21	0.000	0.000	0.000	0.000	0.057

Table 7. Trace metals in oligochaetes as a function of bioassay settling time and exposure duration. All values are ug/g

STATION	SETTLING TIME	BIOASSAY DURATION	Al	As	Cd	Cr	Cu	Fe	Pb	Mn	Hg	Ni	Zn
RICE LAKE	5	10	102	3.60	1.31	13.3	73.9	3473	7.5	66.5	0.27	7.8	388
TORONTO	5	10	25	6.20	1.46	11.5	81.1	3846	3.8	55.8	0.13	10.6	475
CONTROL	5	0	56	7.72	1.76	12.2	88.6	4340	3.9	63.7	0.41	4.4	571
RICE LAKE	5	10	112	7.17	1.55	10.0	94.3	4344	6.3	64.1	0.36	13.1	522
TORONTO	24	10	81	7.97	2.13	21.9	111.5	5187	4.7	62.3	0.34	21.4	620
RICE LAKE	24	10	317	4.58	1.09	9.9	74.1	2968	9.8	51.9	0.14	9.7	290
TORONTO	24	10	114	2.78	0.86	8.2	45.9	2060	5.2	26.2	0.15	5.8	205
RICE LAKE	5	21	286	4.58	1.25	7.9	69.2	3253	12.4	54.5	0.17	9.6	342
TORONTO	5	21	39	0.39	0.19	1.7	9.1	506	1.5	5.8	0.00	1.8	53
RICE LAKE	24	21	90	3.21	0.73	5.2	44.1	2165	6.3	32.8	0.15	5.9	252
TORONTO	24	21	167	4.37	1.49	7.0	58.9	2968	7.8	34.2	0.15	5.4	361
RICE LAKE	21	21	175	3.09	0.66	6.4	40.5	1832	7.2	31.9	0.09	5.3	201
TORONTO	120	21	1114	5.08	1.19	19.1	60.3	4576	26.4	77.9	0.11	8.5	259
CONTROL	5	21	2530	6.86	1.25	15.8	62.2	6859	11.5	172.0	0.10	13.8	346

TABLE 8: Effect of Settling Time on Growth of Pimephales promelas at two densities.

SEDIMENT															
TORONTO STP				ST. MARY'S RIVER				CONTROL							
				SETTLING TIME											
				5h	24h	120h	5h	24h	120h	5h	24h				
DENSITY															
10 ¹	15 ²	10	15	10	15	10	15	10	15	10	15				
Initial 449	480	541	512	541	550	473	482	528	530	517	499	497	499	503	499
Biomass (18)	(15)	(35)	(18)	(30)	(10)	(23)	(43)	(58)	(44)	(40)	(23)	(34)	(22)	(26)	(9)
(mg)															
Percent -14	-21	-24	-16	-4	-15	-19	-16	-10	-17	-7	-5	-4	-3	-3	-6
Biomass (2)	(0.6)	(0.1)	(0.4)	(0.2)	(0.1)	(0.5)	(0.1)	(0.8)	(0.7)	(0.1)	(0.2)	(0)	(0)	(0)	(0)
Change (Day 10)															
Growth 250	600	700	167	0	230	375	433	233	278	100	11	n.a. ¹	n.a.	n.a.	n.a.
Inhibition ¹															
Percent -23	-30	-35	-32	-13	-20	-26	-37	-18	-23	-10	-12	-8	-6	-9	-9
Biomass (1)	(2)	(3)	(4)	(0.7)	(0.4)	(1)	(0.9)	(1.1)	(0.1)	(0.3)	(0)	(0.1)	(0.1)	(0.1)	(0.1)
Change (Day 21)															
Growth 187	400	289	256	53	167	225	512	100	156	10	33	n.a.	n.a.	n.a.	n.a.
Inhibition															

¹ 10 individuals; ² 15 individuals; ¹ relative to controls; n.a. not applicable

Table 9. Trace metals in fathead minnows as a function of bioassay settling time, exposure duration, organism density and gut clearance. All values are ug/g unless otherwise noted.

STATION	SETTLING TIME (hr)	BIOASSAY DURATION (day)	NUMBER OF FISH	GUT CLEARANCE	N	As	Cd	Cr	Cu	Fe	Pb
TORONTO	5	10	10	no	876	7.94	0.55	34.7	245	2205	8.93
TORONTO	5	10	10	no	1871	8.28	1.19	53.7	300	5636	28.32
TORONTO	5	10	10	no	2045	8.98	1.14	54.0	303	5189	47.84
TORONTO	5	10	15	no	413	5.89	0.22	22.0	216	1189	4.49
TORONTO	5	10	15	no	2087	5.37	1.03	41.2	273	6077	28.28
ST. MARY'S	5	10	10	no	1358	22.76	0.50	33.8	255	5889	30.20
ST. MARY'S	5	10	10	no	1601	14.50	0.53	30.4	255	6184	35.31
ST. MARY'S	5	10	15	no	1204	24.58	0.25	24.6	188	5413	25.83
ST. MARY'S	5	10	15	no	873	14.39	0.25	26.7	200	3741	18.56
ST. MARY'S	5	10	15	no	1059	7.15	0.43	29.7	225	4312	22.42
TORONTO	5	10	15	yes	1817	19.20	0.27	59.2	630	8569	25.32
TORONTO	5	10	15	yes	4284	14.90	0.97	75.8	429	9860	36.83
TORONTO	24	10	10	no	2859	3.53	1.76	75.3	996	8318	37.73
TORONTO	24	10	15	no	1323	15.19	0.46	31.1	269	3782	24.78
TORONTO	24	10	15	no	257	15.94	0.13	13.8	182	819	5.97
TORONTO	24	10	15	no	429	17.86	0.17	22.9	285	1104	3.25
ST. MARY'S	5	10	15	yes	1851	17.79	0.82	56.4	346	7788	37.63
ST. MARY'S	5	10	15	yes	2798	22.94	1.15	62.1	488	12806	56.71
ST. MARY'S	24	10	10	no	980	12.77	0.39	23.9	225	4156	21.01
ST. MARY'S	24	10	10	no	1310	7.83	0.49	31.8	238	5242	26.36
ST. MARY'S	24	10	10	no	1425	9.08	0.49	28.4	244	5876	29.20
ST. MARY'S	24	10	15	no	2071	11.21	0.64	43.2	214	9118	38.85
TORONTO	24	10	15	yes	732	20.69	0.25	21.6	214	2914	14.09
TORONTO	24	10	15	yes	233	34.89	0.13	15.6	163	617	2.80
ST. MARY'S	24	10	10	yes	408	21.89	0.17	18.8	256	1053	4.23
ST. MARY'S	24	10	10	yes	589	17.02	0.24	20.5	175	2532	11.20
ST. MARY'S	24	10	15	yes	363	28.08	0.22	16.8	163	1143	5.47
TORONTO	120	10	10	no	3870	11.85	1.56	63.0	263	11102	45.57
TORONTO	120	10	10	no	2434	6.88	1.70	49.4	195	7022	37.09

STATION	SETTLING TIME (hr)	BIOASSAY DURATION (day)	NUMBER OF FISH	GUT CLEARANCE	A ₁	A ₀	Cd	Cr	Cu	Fe	Pb
TORONTO	120	10	15	no	1148	3.94	0.51	24.7	182	2907	18.54
TORONTO	120	10	15	no	3433	7.14	2.17	70.9	255	10826	68.58
ST. MARYS	120	10	10	no	1124	9.06	0.56	24.9	195	4061	31.75
ST. MARYS	120	10	10	no	863	10.87	0.54	18.8	208	3788	28.39
ST. MARYS	120	10	15	no	1623	7.47	0.82	27.1	163	6630	41.33
ST. MARYS	120	10	15	no	1367	12.43	0.53	24.1	162	5743	38.39
CONTROL	24	10	10	no	1538	9.61	0.94	17.2	180	2847	3.51
CONTROL	24	10	15	no	1541	8.16	0.24	17.9	201	2500	5.85
TORONTO	120	10	15	yes	668	6.83	0.64	21.7	175	2221	12.13
TORONTO	120	10	15	yes	594	10.18	0.28	18.7	161	1601	7.18
TORONTO	120	10	15	yes	741	16.12	0.51	16.3	170	1982	9.28
ST. MARYS	120	10	15	yes	528	28.58	0.29	14.5	143	1883	13.67
ST. MARYS	120	10	15	yes	665	32.86	0.32	14.5	130	2741	15.57
CONTROL	24	10	15	yes	863	36.01	0.31	14.4	160	1713	3.85
TORONTO	24	21	10	no	2635	30.01	2.23	59.8	328	7854	50.91
TORONTO	24	21	10	no	2863	21.62	2.20	55.3	311	8305	43.85
TORONTO	24	21	15	no	973	27.89	0.63	24.6	237	2574	13.72
TORONTO	5	21	15	no	2012	12.56	1.43	48.5	235	5771	36.82
ST. MARYS	5	21	10	no	1060	32.98	0.71	37.8	254	7721	48.46
ST. MARYS	5	21	10	no	2087	22.04	0.68	32.8	248	8287	47.30
ST. MARYS	5	21	15	no	1266	17.22	0.56	31.1	204	4622	31.42
ST. MARYS	5	21	15	no	1839	17.66	0.88	30.9	226	7252	52.38
ST. MARYS	5	21	15	no	1381	15.79	0.54	24.3	208	5566	32.05
ST. MARYS	5	21	15	no	1365	39.28	0.80	24.8	202	3403	24.23
TORONTO	24	21	10	no	1104	20.23	0.52	20.7	242	2851	18.27
TORONTO	24	21	15	no	1332	36.28	0.74	30.3	278	3152	51.78
ST. MARYS	24	21	10	no	1157	27.22	0.41	17.2	161	5768	10.91
ST. MARYS	24	21	10	no	1477	28.98	0.47	28.6	247	6875	37.40
ST. MARYS	24	21	15	no	1613	27.28	0.45	28.3	222	7718	44.25
ST. MARYS	5	21	15	no	1371	15.16	0.42	26.8	341	5832	32.50
TORONTO	120	21	15	no	1016	29.17	0.36	16.2	137	2789	13.45
TORONTO	120	21	15	no	629	20.83	0.39	16.8	151	1694	18.71
TORONTO	120	21	10	no	2344	28.56	0.91	32.1	148	6047	35.95

STATION	SETTLING TIME (hr)	BIOASSAY DURATION (day)	NUMBER OF FISH	GUT CLEARANCE	Al	As	Cd	Cr	Cu	Fe	Pb
TORONTO	120	21	10	no	1274	24.68	0.42	19.9	118	3414	25.07
ST. MARY'S	120	21	10	no	557	26.65	0.18	12.8	111	2810	12.89
ST. MARY'S	120	21	10	no	587	10.40	0.21	15.4	210	2235	12.78
ST. MARY'S	120	21	10	no	1360	29.28	0.43	23.4	200	6444	32.74
ST. MARY'S	120	21	15	no	1565	22.44	0.49	27.4	206	7155	32.66
ST. MARY'S	120	21	15	no	1422	31.99	0.44	22.5	168	6852	31.18
ST. MARY'S	24	21	15	yes	429	29.54	0.24	6.3	189	1946	6.61
ST. MARY'S	5	21	10	yes	751	17.40	0.59	21.1	236	3082	14.40
CONTROL	120	21	10	no	1822	16.55	0.46	19.4	108	3232	8.26
CONTROL	120	21	15	no	1668	23.03	0.45	15.8	125	2820	4.48
TORONTO	120	21	15	yes	329	5.58	0.19	12.0	110	772	3.85
TORONTO	120	21	15	yes	1172	25.24	0.68	22.3	156	3063	15.17
ST. MARY'S	120	21	15	yes	516	19.56	0.51	15.6	181	2087	8.26
ST. MARY'S	120	21	15	yes	512	32.62	0.25	13.1	128	2532	8.72
CONTROL	120	21	15	yes	1747	9.96	0.59	16.9	178	2834	5.80

STATION	SETTLING TIME (hr)	BIOASSAY DURATION (day)	NUMBER OF FISH	GUT CLEARANCE	Mn	Hg	Ni	Zn
TORONTO	5	10	10	no	57.95	422	0.86	2154
TORONTO	5	10	10	no	101.25	480	11.82	2272
TORONTO	5	10	10	no	106.30	568	19.95	2539
TORONTO	5	10	15	no	36.17	478	9.97	2085
TORONTO	5	10	15	no	111.08	404	14.73	2176
ST. MARY'S	5	10	10	no	125.85	420	18.35	2143
ST. MARY'S	5	10	10	no	129.11	3.63	20.34	1973
ST. MARY'S	5	10	15	no	108.36	3.58	14.56	1899
ST. MARY'S	5	10	15	no	82.29	3.66	9.89	1410
ST. MARY'S	5	10	15	no	82.77	3.76	16.25	1884
TORONTO	5	10	15	yes	170.34	5.75	19.74	4811
TORONTO	5	10	15	yes	233.54	7.07	48.78	3075
TORONTO	24	10	10	no	152.20	5.71	28.36	3083
TORONTO	24	10	10	no	79.88	6.34	9.86	2843
TORONTO	24	10	15	no	34.69	4.79	3.67	1701
TORONTO	24	10	15	no	43.08	7.84	10.89	2405
ST. MARY'S	5	10	15	yes	163.38	7.07	19.00	2973
ST. MARY'S	5	10	15	yes	255.55	8.18	22.24	4844
ST. MARY'S	24	10	10	no	97.10	4.34	12.75	2144
ST. MARY'S	24	10	10	no	114.71	4.33	12.69	1881
ST. MARY'S	24	10	10	no	125.83	4.41	15.23	1927
ST. MARY'S	24	10	15	no	190.44	5.51	19.78	2568
TORONTO	24	10	15	no	70.80	5.07	11.46	1838
TORONTO	24	10	15	yes	26.69	4.79	4.74	1870
ST. MARY'S	24	10	10	yes	34.48	7.32	5.84	2223
ST. MARY'S	24	10	10	yes	59.31	5.08	7.27	1562
TORONTO	120	10	10	no	33.39	5.04	6.11	1525
TORONTO	120	10	10	no	171.88	4.94	21.56	1803
TORONTO	120	10	10	no	101.58	6.08	22.30	1317

STATION	SETTLING TIME (hr)	BIOASSAY DURATION (day)	NUMBER OF FISH	GUT CLEARANCE	Mn	Hg	Ni	Zn
TORONTO	120	10	15	no	57.83	5.48	6.56	145C
TORONTO	120	10	15	no	151.75	4.55	18.06	1853
ST. MARY'S	120	10	10	no	102.66	3.88	8.79	1408
ST. MARY'S	120	10	10	no	88.09	3.79	10.50	1650
ST. MARY'S	120	10	15	no	142.97	3.25	18.59	1586
ST. MARY'S	120	10	15	no	129.85	2.80	11.58	1481
CONTROL	24	10	10	no	112.66	4.68	10.26	1583
CONTROL	24	10	15	no	119.67	4.86	8.75	1592
TORONTO	120	10	15	yes	31.98	3.01	11.13	1503
TORONTO	120	10	15	yes	27.52	2.85	6.58	1285
TORONTO	120	10	15	yes	37.94	3.31	4.78	1451
ST. MARY'S	120	10	15	yes	44.57	2.84	5.80	1534
ST. MARY'S	120	10	15	yes	61.86	3.24	5.36	1428
CONTROL	24	10	15	yes	62.86	5.44	7.77	1632
TORONTO	5	21	10	no	131.00	7.04	19.09	2421
TORONTO	24	21	10	no	155.25	5.99	22.89	2536
TORONTO	5	21	15	no	53.36	5.70	9.10	2102
TORONTO	5	21	15	no	98.30	5.31	17.85	2417
ST. MARY'S	5	21	10	no	171.50	5.35	16.29	2124
ST. MARY'S	5	21	15	no	177.14	5.41	17.99	2236
ST. MARY'S	5	21	15	no	106.26	4.82	9.72	1862
ST. MARY'S	5	21	15	no	159.29	6.16	22.31	2108
ST. MARY'S	5	21	15	no	121.36	5.59	12.52	1813
ST. MARY'S	5	21	10	no	84.87	3.46	8.88	1810
TORONTO	24	21	10	no	57.35	4.28	8.49	2036
TORONTO	24	21	15	no	88.04	1.85	18.94	2126
ST. MARY'S	24	21	10	no	112.00	2.83	12.04	1572
ST. MARY'S	24	21	10	no	136.17	4.07	18.16	2381
ST. MARY'S	24	21	15	no	158.34	2.87	16.30	1638
ST. MARY'S	5	21	10	no	124.13	4.12	13.52	1828
TORONTO	120	21	15	no	51.86	2.51	8.68	1230
TORONTO	120	21	15	no	31.88	2.96	5.08	1420
TORONTO	120	21	10	no	116.23	2.04	13.56	1424

STATION	SETTLING TIME (hr)	BIOASSAY DURATION (day)	NUMBER OF FISH	GUT CLEARANCE	Mn	Hg	Ni	Zn
TORONTO	120	21	10	no	60.57	3.04	8.11	1327
ST. MARY'S	120	21	10	no	59.64	2.78	5.29	1210
ST. MARY'S	120	21	10	no	56.06	2.59	8.85	1364
ST. MARY'S	120	21	10	no	127.36	4.15	17.10	1619
ST. MARY'S	120	21	15	no	144.10	3.23	13.10	1526
ST. MARY'S	120	21	15	yes	133.97	1.88	14.57	1421
ST. MARY'S	24	21	15	yes	52.30	4.14	-13.34	1545
ST. MARY'S	5	21	10	yes	71.50	3.50	10.73	1841
CONTROL	120	21	10	no	130.63	2.06	8.95	1150
CONTROL	120	21	15	no	118.64	3.08	7.50	1928
TORONTO	120	21	15	yes	14.85	1.49	3.82	858
TORONTO	120	21	15	yes	70.02	3.13	13.80	1514
ST. MARY'S	120	21	15	yes	47.63	3.09	7.85	1666
ST. MARY'S	120	21	15	yes	51.34	2.56	4.84	1538
CONTROL	120	21	15	yes	112.14	2.28	12.30	1496

Table 10. Trace organics in fathead minnows as a function of bioassay settling time, exposure duration, organism density and gut clearance. All values are ug.g⁻¹ unless otherwise noted.

STATION	SETTLING TIME (hr)	BIOASSAY DURATION (day)	NUMBER OF FISH	GUT CLEARANCE	% LIPID	HCB	HEPTA-CHLOR	HEPTA-CHLOR EPOXIDE	ALDRIN	a-BHC	B-BHC
TORONTO	5	10	10	NO	8.590	0.071	0.000 ⁺	0.021	0.000	0.000	0.000
TORONTO	5	10	10	NO	8.860	0.031	0.000	0.010	0.000	0.000	0.000
TORONTO	5	10	15	NO	8.570	0.006	0.000	0.000	0.000	0.006	0.000
TORONTO	5	10	15	NO	8.940	0.008	0.000	0.001	0.000	0.000	0.000
ST. MARYS	5	10	10	NO	8.300	0.000	0.000	0.000	0.000	0.000	0.000
ST. MARYS	5	10	10	NO	8.060	0.001	0.000	0.000	0.000	0.000	0.025
ST. MARYS	5	10	15	NO	8.290	0.009	0.000	0.005	0.000	0.000	0.110
ST. MARYS	5	10	15	NO	8.560	0.000	0.000	0.002	0.000	0.001	0.000
TORONTO	5	10	15	YES	8.180	0.000	0.005	0.000	0.000	0.000	0.010
TORONTO	5	10	15	YES	8.540	0.000	0.007	0.000	0.000	0.000	0.000
TORONTO	5	10	10	NO	8.470	0.000	0.005	0.000	0.001	0.000	0.000
TORONTO	24	10	10	NO	10.370	0.000	0.000	0.004	0.000	0.000	0.000
TORONTO	24	10	15	NO	10.890	0.002	0.009	0.003	0.000	0.000	0.011
TORONTO	24	10	15	NO	9.310	0.000	0.000	0.000	0.000	0.000	0.000
ST. MARYS	5	10	15	YES	7.760	0.000	0.000	0.000	0.000	0.000	0.000
ST. MARYS	5	10	15	YES	7.920	0.002	0.010	0.003	0.000	0.010	0.000
ST. MARYS	24	10	10	NO	10.260	0.000	0.007	0.000	0.000	0.000	0.000
ST. MARYS	24	10	10	NO	10.020	0.001	0.001	0.000	0.002	0.002	0.000
ST. MARYS	24	10	15	NO	8.290	0.003	0.000	0.000	0.000	0.002	0.000
ST. MARYS	24	10	15	NO	8.950	0.004	0.003	0.004	0.000	0.003	0.000
TORONTO	24	10	15	NO	11.450	0.000	0.003	0.000	0.000	0.005	0.000
TORONTO	24	10	15	YES	11.120	0.000	0.003	0.000	0.000	0.000	0.000
ST. MARYS	24	10	10	YES	9.530	0.004	0.003	0.000	0.000	0.003	0.000
ST. MARYS	24	10	15	YES	8.660	0.003	0.000	0.000	0.000	0.003	0.000
TORONTO	120	10	10	NO	8.560	0.000	0.000	0.000	0.002	0.005	0.000
TORONTO	120	10	10	NO	7.570	0.005	0.000	0.000	0.000	0.000	0.000
TORONTO	120	10	15	NO	8.930	0.004	0.002	0.007	0.000	0.001	0.000
TORONTO	120	10	15	NO	8.030	0.005	0.000	0.000	0.000	0.002	0.000
ST. MARYS	120	10	10	NO	7.750	0.003	0.000	0.000	0.003	0.010	0.000

* Values listed as 0.000 are below detection

STATION	SETTLING TIME (hr)	BIOASSAY DURATION (day)	NUMBER OF FISH	GUT CLEARANCE	% LIPID	HCB	HEPTA- CHLOR	HEPTA- CHLOR EPOXIDE	ALDRIN	a-BHC	B-BHC
ST. MARYS	120	10	10	NO	5.450	0.003	0.004	0.001	0.000	0.000	0.000
ST. MARYS	120	10	15	NO	7.510	0.003	0.004	0.000	0.000	0.003	0.000
ST. MARYS	120	10	15	NO	14.430	0.000	0.006	0.000	0.000	0.000	0.000
CONTROL	24	10	10	NO	4.870	0.003	0.004	0.002	0.000	0.000	0.000
CONTROL	24	10	15	YES	8.270	0.005	0.009	0.000	0.000	0.010	0.000
TORONTO	120	10	15	YES	8.790	0.007	0.000	0.000	0.002	0.000	0.014
TORONTO	120	10	15	YES	7.820	0.005	0.003	0.003	0.000	0.000	0.000
ST. MARYS	120	10	15	YES	5.810	0.000	0.000	0.000	0.000	0.000	0.000
ST. MARYS	120	10	15	YES	7.870	0.002	0.000	0.000	0.000	0.000	0.000
CONTROL	24	10	15	YES	4.430	0.001	0.002	0.000	0.001	0.000	0.000
TORONTO	5	21	10	NO	4.840	0.004	0.000	0.000	0.002	0.002	0.000
TORONTO	24	21	10	NO	7.180	0.000	0.000	0.000	0.000	0.000	0.000
TORONTO	5	21	15	NO	7.340	0.000	0.000	0.000	0.004	0.000	0.000
TORONTO	5	21	15	NO	8.080	0.000	0.000	0.001	0.000	0.000	0.000
ST. MARYS	5	21	10	NO	7.010	0.004	0.003	0.000	0.000	0.004	0.000
ST. MARYS	5	21	10	NO	6.080	0.000	0.000	0.000	0.000	0.005	0.000
ST. MARYS	5	21	15	NO	6.850	0.010	0.000	0.000	0.005	0.004	0.000
ST. MARYS	5	21	15	NO	5.560	0.000	0.000	0.000	0.000	0.005	0.000
TORONTO	24	21	10	NO	6.180	0.005	0.000	0.000	0.006	0.003	0.000
TORONTO	24	21	10	NO	8.180	0.004	0.012	0.000	0.000	0.000	0.032
TORONTO	24	21	15	NO	6.170	0.008	0.011	0.000	0.000	0.006	0.000
ST. MARYS	24	21	10	NO	7.980	0.000	0.000	0.000	0.000	0.000	0.000
ST. MARYS	24	21	10	NO	6.950	0.000	0.010	0.000	0.000	0.000	0.000
ST. MARYS	5	21	10	NO	13.120	0.000	0.000	0.000	0.001	0.001	0.000
TORONTO	120	21	15	NO	7.960	0.007	0.000	0.000	0.000	0.003	0.000
TORONTO	120	21	15	NO	5.640	0.007	0.000	0.000	0.000	0.002	0.000
TORONTO	120	21	10	NO	5.140	0.004	0.004	0.000	0.000	0.002	0.000
TORONTO	120	21	10	NO	7.300	0.000	0.000	0.000	0.000	0.004	0.000
ST. MARYS	120	21	10	NO	8.330	0.004	0.005	0.000	0.000	0.000	0.000
ST. MARYS	120	21	10	NO	8.080	0.000	0.000	0.000	0.000	0.000	0.000
ST. MARYS	120	21	15	NO	6.520	0.000	0.000	0.000	0.000	0.000	0.000
ST. MARYS	120	21	15	NO	7.880	0.000	0.000	0.004	0.002	0.000	0.000
ST. MARYS	24	21	15	YES	8.490	0.000	0.000	0.000	0.004	0.008	0.000

STATION	SETTLING TIME (hr)	BIOASSAY DURATION (day)	NUMBER OF FISH	GUT CLEARANCE	% LIPID	HCB	HEPTA- CHLOR	HEPTA- CHLOR EPOXIDE	ALDRIN	a-BHC	B-BHC
ST. MARY'S CONTROL	5	21	10	YES	8.090	0.000	0.018	0.000	0.000	0.011	0.000
CONTROL	120	21	10	NO	11.120	0.004	0.000	0.000	0.000	0.004	0.000
TORONTO	120	21	15	YES	10.130	0.004	0.000	0.000	0.000	0.009	0.000
TORONTO	120	21	15	YES	6.000	0.009	0.014	0.000	0.000	0.000	0.000
ST. MARY'S	120	21	15	YES	9.250	0.000	0.000	0.000	0.000	0.007	0.000
ST. MARY'S	120	21	15	YES	3.080	0.005	0.004	0.000	0.000	0.002	0.000
ST. MARY'S	120	21	15	YES	7.970	0.004	0.004	0.000	0.000	0.003	0.000
CONTROL	120	21	15	YES	6.700	0.000	0.000	0.000	0.000	0.000	0.000

STATION	SETTLING TIME (hr)	BIOASSAY DURATION (day)	NUMBER OF FISH	GUT CLEARANCE	HCBI LIPID	HEFTA CHLOR/ LIPID	HCOI LIPID	ALDRIN/ LIPID	a-BHC/ LIPID	b-BHC/ LIPID	ALPHA CHLORDANE/ LIPID
TORONTO	5	10	10	NO	0.006	0.000	0.002	0.000	0.000	0.000	0.001
TORONTO	5	10	10	NO	0.003	0.000	0.001	0.000	0.000	0.000	0.001
TORONTO	5	10	15	NO	0.001	0.000	0.000	0.000	0.001	0.000	0.000
TORONTO	5	10	15	NO	0.001	0.000	0.000	0.000	0.000	0.000	0.000
ST. MARY'S	5	10	10	NO	0.000	0.000	0.000	0.000	0.000	0.003	0.000
ST. MARY'S	5	10	15	NO	0.001	0.000	0.001	0.000	0.000	0.013	0.001
ST. MARY'S	5	10	15	NO	0.000	0.000	0.000	0.000	0.000	0.000	0.000
TORONTO	5	10	15	YES	0.000	0.001	0.000	0.000	0.000	0.001	0.000
TORONTO	5	10	15	YES	0.000	0.001	0.000	0.000	0.000	0.000	0.000
TORONTO	24	10	10	NO	0.000	0.001	0.000	0.000	0.000	0.000	0.000
TORONTO	24	10	10	NO	0.000	0.000	0.000	0.000	0.000	0.000	0.000
TORONTO	24	10	15	NO	0.000	0.001	0.000	0.000	0.000	0.001	0.000
ST. MARY'S	5	10	15	YES	0.000	0.000	0.000	0.000	0.000	0.000	0.000
ST. MARY'S	5	10	15	YES	0.000	0.001	0.000	0.000	0.001	0.000	0.000
ST. MARY'S	24	10	10	NO	0.000	0.001	0.000	0.000	0.000	0.000	0.000
ST. MARY'S	24	10	10	NO	0.000	0.000	0.000	0.000	0.000	0.000	0.000
ST. MARY'S	24	10	15	NO	0.000	0.000	0.000	0.000	0.000	0.000	0.000
TORONTO	24	10	15	NO	0.000	0.000	0.000	0.000	0.000	0.000	0.000
TORONTO	24	10	15	YES	0.000	0.000	0.000	0.000	0.000	0.000	0.000
ST. MARY'S	24	10	15	YES	0.000	0.000	0.000	0.000	0.000	0.000	0.000
TORONTO	24	10	10	YES	0.000	0.000	0.000	0.000	0.000	0.000	0.000
ST. MARY'S	24	10	15	YES	0.000	0.000	0.000	0.000	0.000	0.000	0.000
TORONTO	120	10	10	NO	0.000	0.000	0.000	0.000	0.001	0.000	0.001
TORONTO	120	10	10	NO	0.001	0.000	0.001	0.000	0.000	0.000	0.001
TORONTO	120	10	15	NO	0.001	0.000	0.000	0.000	0.000	0.000	0.000
TORONTO	120	10	15	NO	0.001	0.000	0.000	0.000	0.000	0.000	0.000
ST. MARY'S	120	10	10	NO	0.000	0.000	0.000	0.000	0.001	0.000	0.000

STATION	SETTLING TIME (hr)	BIOASSAY DURATION (day)	NUMBER OF FISH	GUT CLEARANCE	HCB/ LIPID	HEPTA CHLOR/ LIPID	HCB/ LIPID	ALDRIN/ LIPID	a-BHC/ LIPID	b-BHC/ LIPID	ALPHA CHLORDANE/ LIPID
ST. MARYS	120	10	10	NO	0.001	0.001	0.000	0.000	0.000	0.000	0.000
ST. MARYS	120	10	15	NO	0.000	0.001	0.000	0.000	0.000	0.000	0.000
ST. MARYS	120	10	15	NO	0.000	0.001	0.000	0.000	0.000	0.000	0.000
CONTROL	24	10	10	NO	0.001	0.001	0.000	0.000	0.000	0.000	0.001
CONTROL	24	10	15	NO	0.001	0.001	0.000	0.000	0.001	0.000	0.000
TORONTO	120	10	15	YES	0.001	0.000	0.000	0.000	0.000	0.001	0.001
TORONTO	120	10	15	YES	0.001	0.000	0.000	0.000	0.000	0.000	0.001
ST. MARYS	120	10	15	YES	0.000	0.000	0.000	0.000	0.000	0.000	0.000
ST. MARYS	120	10	15	YES	0.000	0.000	0.000	0.000	0.000	0.000	0.001
ST. MARYS	120	10	15	YES	0.000	0.000	0.000	0.000	0.000	0.000	0.000
CONTROL	24	10	15	YES	0.000	0.000	0.000	0.000	0.000	0.000	0.000
TORONTO	5	21	10	NO	0.001	0.000	0.000	0.000	0.000	0.000	0.001
TORONTO	24	21	10	NO	0.000	0.000	0.000	0.000	0.000	0.000	0.001
TORONTO	5	21	15	NO	0.000	0.000	0.000	0.000	0.000	0.000	0.001
TORONTO	5	21	15	NO	0.000	0.000	0.000	0.001	0.000	0.000	0.000
ST. MARYS	5	21	10	NO	0.001	0.000	0.000	0.000	0.001	0.000	0.001
ST. MARYS	5	21	10	NO	0.000	0.000	0.000	0.000	0.001	0.000	0.001
ST. MARYS	5	21	15	NO	0.001	0.000	0.000	0.001	0.001	0.000	0.000
ST. MARYS	5	21	15	NO	0.000	0.000	0.000	0.000	0.001	0.000	0.000
TORONTO	24	21	10	NO	0.001	0.000	0.000	0.001	0.000	0.000	0.000
TORONTO	24	21	10	NO	0.000	0.000	0.000	0.000	0.000	0.004	0.000
TORONTO	24	21	15	NO	0.001	0.002	0.000	0.000	0.001	0.000	0.000
ST. MARYS	24	21	10	NO	0.000	0.000	0.000	0.000	0.000	0.000	0.000
ST. MARYS	24	21	10	NO	0.000	0.001	0.000	0.000	0.000	0.000	0.001
ST. MARYS	24	21	10	NO	0.000	0.000	0.000	0.000	0.000	0.000	0.000
TORONTO	120	21	15	NO	0.001	0.000	0.000	0.000	0.000	0.000	0.001
TORONTO	120	21	15	NO	0.001	0.000	0.000	0.000	0.000	0.000	0.001
TORONTO	120	21	10	NO	0.001	0.000	0.000	0.000	0.000	0.000	0.000
TORONTO	120	21	10	NO	0.000	0.001	0.000	0.000	0.001	0.000	0.000
ST. MARYS	120	21	10	NO	0.000	0.001	0.000	0.000	0.000	0.000	0.000
ST. MARYS	120	21	10	NO	0.000	0.000	0.000	0.000	0.000	0.000	0.000
ST. MARYS	120	21	15	NO	0.000	0.000	0.000	0.000	0.000	0.000	0.001
ST. MARYS	120	21	15	NO	0.000	0.000	0.000	0.000	0.000	0.000	0.001
ST. MARYS	120	21	15	NO	0.000	0.000	0.001	0.000	0.000	0.000	0.000
ST. MARYS	24	21	15	YES	0.000	0.000	0.000	0.000	0.001	0.000	0.000

STATION	SETTLING TIME (hr)	BIOASSAY DURATION (day)	NUMBER OF FISH	GUT CLEARANCE	HCB/ LIPID	HEPTA CHLOR/ LIPID	HCB/ LIPID	ALDRIN/ LIPID	a-BHC/ LIPID	b-BHC/ LIPID	ALPHA CHLORDANE/ LIPID
ST. MARY'S CONTROL	5	21	10	YES	0.000	0.002	0.000	0.000	0.001	0.000	0.000
CONTROL	120	21	10	NO	0.000	0.000	0.000	0.000	0.000	0.000	0.000
TORONTO	120	21	15	YES	0.000	0.000	0.000	0.000	0.001	0.000	0.000
TORONTO	120	21	15	YES	0.002	0.002	0.000	0.000	0.000	0.000	0.000
ST. MARY'S	120	21	15	YES	0.000	0.000	0.000	0.000	0.001	0.000	0.001
ST. MARY'S	120	21	15	YES	0.002	0.001	0.000	0.000	0.001	0.000	0.001
CONTROL	120	21	15	YES	0.001	0.001	0.000	0.000	0.000	0.000	0.001
CONTROL	120	21	15	YES	0.000	0.000	0.000	0.000	0.000	0.000	0.000

STATION	SETTLING TIME (hr)	BIOASSAY DURATION (day)	NUMBER OF FISH	GUT CLEARANCE	GAMMA CHLORDANE/ LIPID	PP-DDE/ LIPID	OP-DDT/ LIPID	PCB/ LIPID
TORONTO	5	10	10	NO	0.003	0.002	0.000	0.051
TORONTO	5	10	10	NO	0.001	0.002	0.000	0.000
TORONTO	5	10	15	NO	0.000	0.000	0.000	0.005
TORONTO	5	10	15	NO	0.000	0.000	0.000	0.003
ST. MARYS	5	10	10	NO	0.001	0.000	0.000	0.008
ST. MARYS	5	10	10	NO	0.000	0.000	0.000	0.000
ST. MARYS	5	10	15	NO	0.000	0.002	0.000	0.000
ST. MARYS	5	10	15	NO	0.000	0.000	0.000	0.002
TORONTO	5	10	15	YES	0.000	0.000	0.000	0.000
TORONTO	5	10	15	YES	0.000	0.000	0.000	0.000
TORONTO	24	10	10	NO	0.001	0.000	0.000	0.008
TORONTO	24	10	10	NO	0.000	0.000	0.000	0.008
TORONTO	24	10	10	NO	0.000	0.000	0.000	0.008
TORONTO	24	10	15	NO	0.000	0.000	0.000	0.011
TORONTO	24	10	15	NO	0.000	0.000	0.000	0.006
ST. MARYS	5	10	15	YES	0.000	0.000	0.000	0.006
ST. MARYS	5	10	15	YES	0.000	0.001	0.000	0.000
ST. MARYS	24	10	10	NO	0.000	0.000	0.000	0.000
ST. MARYS	24	10	10	NO	0.000	0.000	0.000	0.000
ST. MARYS	24	10	15	NO	0.000	0.000	0.006	0.000
ST. MARYS	24	10	15	NO	0.000	0.002	0.000	0.000
TORONTO	24	10	15	NO	0.000	0.000	0.000	0.000
TORONTO	24	10	15	YES	0.000	0.000	0.000	0.000
ST. MARYS	24	10	10	YES	0.000	0.000	0.000	0.000
ST. MARYS	24	10	15	YES	0.000	0.002	0.000	0.000
TORONTO	120	10	10	NO	0.000	0.002	0.000	0.001
TORONTO	120	10	10	NO	0.000	0.002	0.000	0.000
TORONTO	120	10	15	NO	0.000	0.002	0.000	0.000
ST. MARYS	120	10	10	NO	0.000	0.000	0.000	0.000

STATION	SETTLING TIME (hr)	BIOASSAY DURATION (day)	NUMBER OF FISH	CUT CLEARANCE	GAMMA CHLORIDANE/ LIPID	PP-DDE/ LIPID	OP-DDT/ LIPID	PCB/ LIPID
ST. MARYS	120	10	10	NO	0.000	0.001	0.001	0.000
ST. MARYS	120	10	15	NO	0.000	0.000	0.000	0.000
ST. MARYS	120	10	15	NO	0.000	0.000	0.000	0.000
CONTROL	24	10	10	NO	0.000	0.001	0.000	0.000
CONTROL	24	10	15	NO	0.000	0.002	0.000	0.000
TORONTO	120	10	15	YES	0.000	0.002	0.000	0.000
TORONTO	120	10	15	YES	0.000	0.002	0.000	0.000
ST. MARYS	120	10	15	YES	0.000	0.000	0.000	0.000
ST. MARYS	120	10	15	YES	0.000	0.002	0.000	0.000
CONTROL	24	10	15	YES	0.001	0.002	0.000	0.000
TORONTO	5	21	10	NO	0.001	0.000	0.027	0.000
TORONTO	24	21	10	NO	0.001	0.003	0.000	0.000
TORONTO	5	21	15	NO	0.000	0.000	0.000	0.000
TORONTO	5	21	15	NO	0.000	0.000	0.000	0.000
ST. MARYS	5	21	10	NO	0.001	0.000	0.000	0.000
ST. MARYS	5	21	10	NO	0.000	0.002	0.000	0.000
ST. MARYS	5	21	15	NO	0.000	0.000	0.000	0.016
ST. MARYS	5	21	15	NO	0.000	0.000	0.000	0.002
TORONTO	24	21	10	NO	0.000	0.000	0.000	0.000
TORONTO	24	21	10	NO	0.000	0.000	0.000	0.000
TORONTO	24	21	15	NO	0.000	0.000	0.000	0.000
ST. MARYS	24	21	10	NO	0.000	0.000	0.000	0.000
ST. MARYS	24	21	10	NO	0.000	0.003	0.000	0.000
ST. MARYS	5	21	10	NO	0.000	0.002	0.000	0.000
TORONTO	120	21	15	NO	0.001	0.003	0.000	0.000
TORONTO	120	21	15	NO	0.001	0.000	0.000	0.000
TORONTO	120	21	10	NO	0.001	0.003	0.000	0.000
ST. MARYS	120	21	10	NO	0.001	0.000	0.001	0.000
ST. MARYS	120	21	10	NO	0.001	0.003	0.002	0.000
ST. MARYS	120	21	15	NO	0.001	0.002	0.000	0.000
ST. MARYS	120	21	15	NO	0.000	0.000	0.000	0.000
ST. MARYS	24	21	15	YES	0.001	0.003	0.000	0.000

STATION	SETTLING TIME (hr)	BIOASSAY DURATION (day)	NUMBER OF FISH	GUT CLEARANCE	GAMMA CHLORIDANE/ LIPID	PP-DDE/ LIPID	OP-DDT/ LIPID	PCB/ LIPID
ST. MARY'S	5	21	10	YES	0.000	0.000	0.000	0.001
CONTROL	120	21	10	NO	0.000	0.000	0.000	0.001
CONTROL	120	21	15	NO	0.000	0.000	0.000	0.000
TORONTO	120	21	15	YES	0.000	0.000	0.000	0.000
TORONTO	120	21	15	YES	0.001	0.002	0.000	0.000
ST. MARY'S	120	21	15	YES	0.001	0.005	0.000	0.000
ST. MARY'S	120	21	15	YES	0.000	0.001	0.000	0.000
CONTROL	120	21	15	YES	0.000	0.000	0.000	0.000

TABLE 11: Effect of feeding on growth of Pimephales promelas
exposed to clean and contaminated sediments.
Values in parentheses are standard deviations

MEASUREMENT	SEDIMENT					
	RICE LAKE		ST. MARYS		CONTROL	
	TREATMENT					
	FED	UNFED	FED	UNFED	FED	UNFED
Initial Biomass (mg)	392 (47)	366 (79)	353 (58)	332 (32)	344 (23)	368 (21)
Percent Biomass Change	-14 (2)	-11 (2)	-18 (2)	-21 (3)	5 (0.2)	-9 (0.4)
Growth ¹ Inhibition	380	22	460	133	n.a. ²	n.a.

¹ relative to controls

² not applicable

Table 12. Trace metals in farhead mannows as a function of feeding and gut clearance. All values are ug.g⁻¹ unless otherwise noted.

STATION	FOOD	PURGE	Al	As	Cd	Cr	Cu	Fe	Pb	Mn	Hg	Ni	Zn
ALGOMA	NO	NO	2834	23.19	0.38	31.0	246	4754	13.90	208.49	2.10	14.63	1873
RICE LAKE	NO	NO	3224	10.19	1.99	32.2	225	8063	30.18	250.47	2.45	14.86	1758
RICE LAKE	NO	NO	2496	15.24	1.59	29.7	204	4688	26.57	198.86	2.11	9.06	1700
ST. MARY'S	NO	NO	1252	20.10	0.70	22.5	210	5387	21.19	61.96	2.32	8.69	1802
ST. MARY'S	NO	NO	968	17.01	0.48	22.2	178	3283	10.35	52.86	1.40	11.45	1780
TORONTO	NO	NO	4184	19.21	0.76	20.9	203	7544	10.39	285.29	2.88	9.46	1698
TORONTO	NO	NO	3875	15.72	0.89	21.7	176	6862	11.32	285.42	2.59	12.71	1481
ALGOMA	NO	NO	438	3.24	0.06	4.6	45	882	1.99	75.40	0.58	1.58	355
RICE LAKE	NO	YES	1466	11.59	1.02	21.5	164	2716	16.80	107.55	2.04	11.60	1539
RICE LAKE	NO	YES	1945	17.07	1.40	30.3	240	3628	20.57	130.65	3.20	13.58	1855
ST. MARY'S	NO	YES	947	11.04	0.46	21.2	205	3732	12.64	52.81	2.35	6.50	1867
ST. MARY'S	NO	YES	839	19.99	0.47	18.7	206	3369	12.44	51.45	2.17	3.74	1878
TORONTO	NO	YES	1981	8.35	1.05	21.5	200	3525	4.25	140.19	2.22	13.60	1651
TORONTO	NO	YES	1556	8.95	0.56	17.2	169	2987	4.99	106.35	2.13	5.92	1598
RICE LAKE	YES	NO	3556	14.33	2.02	35.8	209	6765	37.02	290.06	2.23	15.59	1544
RICE LAKE	YES	NO	3360	12.10	2.10	36.0	233	6452	33.50	294.45	2.26	25.92	1822
ST. MARY'S	YES	NO	1237	26.32	0.52	30.7	198	4811	22.67	78.62	1.31	12.06	1766
TORONTO	YES	NO	5728	11.65	1.30	35.2	257	10462	13.46	424.70	2.21	21.95	1821
TORONTO	YES	NO	2887	14.54	0.62	23.0	172	4787	5.56	165.49	1.60	17.91	1554
CONTROL	YES	NO	682	20.51	0.21	30.1	170	1184	2.00	69.84	1.84	13.07	1514
RICE LAKE	YES	YES	689	7.70	0.48	12.7	112	1228	5.73	50.75	1.05	3.79	876
RICE LAKE	YES	YES	2247	25.80	2.01	45.8	340	4404	18.63	183.22	2.88	14.22	3511
RICE LAKE	YES	YES	1678	12.22	1.25	20.7	209	3164	17.82	141.67	2.85	10.41	1802
ST. MARY'S	YES	YES	572	14.55	0.84	17.2	217	1752	7.98	30.90	2.72	7.72	1766
TORONTO	YES	YES	2386	15.25	0.37	20.3	204	4363	6.54	152.40	2.44	10.95	1653
TORONTO	YES	YES	2747	14.82	0.61	22.5	224	5251	8.15	177.84	2.24	12.82	1469

STATION	FOOD	PURGE	% LIPID	HCB	HEPTA- CHLOR	HEPTACHLOR EPOXIDE	ALDRIN	a-BHC	b-BHC	ALPHA CHLORDANE	GAMMA CHLORANE
ALGOMA	NO	NO	4.270	0.000*	0.000	0.000	0.000	0.000	0.000	0.000	0.007
RICE LAKE	NO	NO	6.550	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
RICE LAKE	NO	NO	8.150	0.000	0.012	0.000	0.000	0.000	0.000	0.017	0.000
ST. MARYS	NO	NO	6.890	0.000	0.000	0.026	0.000	0.011	0.000	0.000	0.010
ST. MARYS	NO	NO	2.470	0.000	0.012	0.000	0.000	0.006	0.000	0.000	0.000
TORONTO	NO	NO	4.140	0.004	0.000	0.000	0.000	0.004	0.000	0.000	0.000
TORONTO	NO	NO	4.300	0.000	0.000	0.000	0.007	0.000	0.000	0.010	0.000
CONTROL	NO	NO	5.810	0.000	0.000	0.000	0.000	0.000	0.050	0.000	0.000
ALGOMA	NO	NO	8.870	0.011	0.000	0.000	0.002	0.011	0.000	0.007	0.000
RICE LAKE	NO	YES	5.990	0.000	0.011	0.000	0.000	0.000	0.000	0.000	0.000
RICE LAKE	NO	YES	4.240	0.007	0.015	0.006	0.000	0.009	0.000	0.016	0.000
ST. MARYS	NO	YES	5.550	0.006	0.000	0.000	0.005	0.007	0.000	0.004	0.003
ST. MARYS	NO	YES	4.780	0.017	0.000	0.000	0.007	0.000	0.000	0.000	0.000
TORONTO	NO	YES	4.730	0.000	0.000	0.000	0.000	0.000	0.000	0.002	0.000
TORONTO	NO	YES	5.730	0.004	0.005	0.000	0.001	0.004	0.000	0.004	0.000
RICE LAKE	YES	NO	5.910	0.004	0.000	0.000	0.000	0.005	0.000	0.000	0.000
RICE LAKE	YES	NO	7.740	0.007	0.000	0.008	0.004	0.000	0.000	0.000	0.000
ST. MARYS	YES	NO	5.340	0.000	0.000	0.000	0.000	0.000	0.000	0.010	0.000
TORONTO	YES	NO	6.250	0.007	0.000	0.000	0.003	0.000	0.000	0.004	0.000
TORONTO	YES	NO	4.190	0.000	0.011	0.000	0.002	0.009	0.000	0.006	0.000
ST. MARYS	YES	NO	6.190	0.009	0.000	0.000	0.005	0.005	0.000	0.010	0.000
ALGOMA	YES	NO	6.240	0.000	0.000	0.000	0.000	0.030	0.130	0.000	0.000
RICE LAKE	YES	YES	4.600	0.000	0.009	0.000	0.000	0.000	0.000	0.000	0.000
RICE LAKE	YES	YES	7.690	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
ST. MARYS	YES	YES	6.860	0.000	0.000	0.000	0.000	0.000	0.000	0.014	0.000
TORONTO	YES	YES	4.970	0.000	0.022	0.000	0.000	0.000	0.000	0.004	0.000
TORONTO	YES	YES	6.690	0.014	0.007	0.000	0.000	0.012	0.000	0.003	0.000

* Values listed as 0.000 are below detection

pp-DDE	op-DDT	PCB	HCB/ LIPID	HC/ LIPID	HCE/ LIPID	ALDRIN/ LIPID	a-BHC LIPID	b-BHC LIPID	A-CHLORDANE/ LIPID	G-CHLORDANE/ LIPID	pp-DDE LIPID
0.000	0.000	0.00	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.002	0.000
0.000	0.000	0.00	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
0.017	0.000	0.00	0.000	0.001	0.000	0.000	0.000	0.000	0.002	0.000	0.002
0.000	0.000	0.00	0.000	0.000	0.004	0.000	0.002	0.000	0.000	0.001	0.000
0.000	0.000	0.00	0.000	0.005	0.000	0.000	0.002	0.000	0.000	0.000	0.000
0.004	0.000	0.01	0.001	0.000	0.000	0.000	0.001	0.000	0.000	0.000	0.001
0.036	0.000	0.01	0.000	0.000	0.000	0.002	0.000	0.000	0.002	0.000	0.008
0.007	0.000	0.00	0.000	0.000	0.000	0.000	0.000	0.009	0.000	0.000	0.001
0.025	0.000	0.00	0.001	0.000	0.000	0.000	0.001	0.000	0.001	0.000	0.003
0.013	0.000	0.00	0.002	0.002	0.000	0.000	0.000	0.000	0.000	0.004	0.002
0.015	0.000	0.00	0.000	0.004	0.002	0.000	0.002	0.000	0.004	0.001	0.004
0.000	0.000	0.00	0.001	0.000	0.000	0.001	0.001	0.000	0.001	0.000	0.000
0.000	0.000	0.00	0.004	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
0.000	0.000	0.00	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
0.009	0.000	0.00	0.001	0.001	0.000	0.000	0.001	0.000	0.001	0.000	0.002
0.000	0.077	0.00	0.001	0.000	0.000	0.000	0.001	0.000	0.000	0.000	0.000
0.000	0.000	0.00	0.001	0.000	0.001	0.001	0.000	0.000	0.002	0.000	0.000
0.000	0.029	0.00	0.000	0.000	0.000	0.000	0.000	0.000	0.001	0.000	0.005
0.031	0.000	0.00	0.001	0.000	0.000	0.000	0.000	0.000	0.001	0.000	0.000
0.016	0.000	0.00	0.000	0.003	0.000	0.000	0.002	0.000	0.001	0.000	0.004
0.045	0.000	0.02	0.001	0.000	0.000	0.001	0.001	0.000	0.002	0.000	0.007
0.011	0.000	0.00	0.000	0.000	0.000	0.000	0.005	0.021	0.000	0.000	0.002
0.020	0.000	0.00	0.000	0.002	0.000	0.000	0.000	0.000	0.000	0.000	0.004
0.000	0.000	0.00	0.000	0.000	0.000	0.000	0.000	0.000	0.002	0.000	0.000
0.030	0.000	0.00	0.000	0.000	0.000	0.000	0.000	0.000	0.002	0.000	0.004
0.020	0.000	0.00	0.000	0.004	0.000	0.000	0.000	0.000	0.001	0.000	0.004
0.012	0.000	0.01	0.002	0.001	0.000	0.000	0.002	0.000	0.000	0.000	0.002

TABLE 1. Comparison of intact cores and homogenized sediments, with reference to growth and mortality of *Hexagenia limba* and *Pilophaea promelas*. Values in parentheses are standard deviations

	SITE A ¹			SITE B ¹			SITE C ²			CONTROL ¹		
	Percent Biomass Mortality	Relative Growth Change	Inhibition	Percent Biomass Mortality	Relative Growth Change	Inhibition	Percent Biomass Mortality	Relative Growth Change	Inhibition	Percent Biomass Mortality	Relative Growth Change	Inhibition
HEXAGENIA												
Core	47 (5)	-37 (10)	186	0 (0)	129 (35)	20 (5)	45 (33)	22 (15)	86	0 (0)	162 (36)	n.s. ⁴
Homogenized	8 (11)	4 (1)	96	8 (11)	47 (16)	54 (19)	71 (12)	-80 (12)	175	0 (0)	107 (17)	n.s.
PILOPHAEA												
Core	32 (8)	-43 (6)	514	20 (14)	-36 (9)	414	25 (9)	-30 (14)	328	0 (0)	-7 (5)	n.s.
Homogenized	30 (36)	-39 (19)	680	87 (19)	-868 (187)		13 (5)	-31 (16)	520	7 (5)	-5 (1)	n.s.

¹ fine-grained sediment

² sandy sediment

³ relative to controls

⁴ not applicable

Figure 1. Biomass changes in *Hexagenia limbata* nymphs for 10 and 21 day exposure durations

Figure 2. Biomass changes in *Pimephales promelas* for 10 and 21 day exposure durations

Figure 3

METAL CONCENTRATION IN MAYFLIES AND FATHEAD MINNOWS BEAKER VERSUS CORE EXPOSURES

Figure 4

METAL CONCENTRATION IN MAYFLIES AND FATHEAD MINNOWS BEAKER VERSUS CORE EXPOSURES

METAL CONCENTRATION IN MAYFLIES AND FATHEAD MINNOWS BEAKER VERSUS CORE EXPOSURES

METAL CONCENTRATION IN MAYFLIES AND FATHEAD MINNOWS BEAKER VERSUS CORE EXPOSURES

METAL CONCENTRATION IN MAYFLIES AND FATHEAD MINNOWS BEAKER VERSUS CORE EXPOSURES