QE
747
T4
W4

UC-NRLF

QB 12 179

BERKELEY
LIBRARY
UNIVERSITY OF
CALIFORNIA

EARTH
SCIENCES
LIBRARY

Matthew L.

BULLETIN
OF
THE UNIVERSITY OF TEXAS

No. 184.

FOUR TIMES A MONTH

SCIENTIFIC SERIES NO. 18

MAY 22, 1911

Success of Economic Geology!
Contributions from the School of Geology

FAUNA OF THE BUDA LIMESTONE

BY

FRANCIS LUTHER WHITNEY

INSTRUCTOR IN GEOLOGY AND PALEONTOLOGY IN THE
UNIVERSITY OF TEXAS

By Smith
1827-1828
1827-1828

EARTH
SCIENCES
LIBRARY

PUBLISHED BY
THE UNIVERSITY OF TEXAS
AUSTIN, TEXAS

*Entered as second-class mail matter at the postoffice at
Austin, Texas*

PUBLICATIONS

OF THE

UNIVERSITY OF TEXAS

Board of Editors—Killis Campbell, Editor-in-Chief; Eugene C. Bauman, Manager; Robert A. Law, N. L. Goodrich, F. W. Simonds, A. C. Scott, A. Lomax, James J. Terrill, C. S. Potts, E. E. Rall.

The publications of the University of Texas are issued four times a month. They are arranged in the following series: RECORD, MINERAL SURVEY, AGRICULTURAL, HUMANISTIC, MEDICAL, SCIENTIFIC, REPRINT, UNIVERSITY EXTENSION, OFFICIAL, PRESS. For postal purposes they are numbered consecutively. Bulletins without regard to the arrangement in series. With the exception of the special numbers any Bulletin will be sent to citizens of Texas free of charge. Communications from other institutions in reference to exchange of publications should be addressed to the University of Texas Library.

THE UNIVERSITY OF TEXAS RECORD has been issued from two to four times a year since December, 1898, and is now in its 10th volume. Its purpose is to preserve a record of the life and progress of the University, and for that reason it is of special interest to alumni, ex-students, and friends of the University. Upon request it will be regularly mailed, free, to any citizen of Texas. Address THE UNIVERSITY OF TEXAS RECORD, Austin, Texas.

The bulletins of the UNIVERSITY EXTENSION AND OFFICIAL series, containing Announcements of Courses, Catalogues, Regents' Reports, and other administrative matter, will be mailed free to any one upon request.

The University still has for distribution copies of the following bulletins:

GENERAL SERIES

4. *Courses of Study in Law Pursued in the University of Texas*, by J. H. Townes, 16 pp. March, 1904.
7. *The Consolidation of Rural Schools*, by Una Bedichek and G. T. Bascom. New edition, enlarged by A. C. Ellis. 85 p., illus. November, 1905. 25 cents.
11. *What Should be Done by Universities to Foster the Professional Education of Teachers?* by W. S. Sutton. 24 p. 1905. 15 cents.
16. *A Study in School Supervision*, by Carl Hartman. 180 p. 1907. 50 cents.
17. *Religious Activities at the University of Texas*. 53 p., illus. August, 1907. 25 cents.

HUMANISTIC SERIES

5. *The Grotesque in the Poetry of Robert Browning*, by Lily B. Campbell. 41 p. April, 1907. 25 cents.
6. *The Beginnings of Texas*, by R. C. Clark. 94 p., map. December, 1907. 75 cents.

(Continued on inside back cover)

BULLETIN
OF
THE UNIVERSITY OF TEXAS

NO. 184

FOUR TIMES A MONTH

SCIENTIFIC SERIES NO. 18

MAY 22, 1911

Contributions from the School of Geology

FAUNA OF THE BUDA LIMESTONE

BY

FRANCIS LUTHER WHITNEY

INSTRUCTOR IN GEOLOGY AND PALEONTOLOGY IN THE
UNIVERSITY OF TEXAS

BRUCE L. CLARK

PUBLISHED BY
THE UNIVERSITY OF TEXAS
AUSTIN, TEXAS

*Entered as second-class mail matter at the postoffice at
Austin, Texas*

EARTH
SCIENCES

ERRATA

For presentation, read preparation, page 9.

For Stearnsia robinsi White, read Remondia robinsi
(White), pages 17 and 42.

Index

CONTENTS.

	Page.
.....	9
.....	7
.....	11
.....	11
.....	11
.....	11
.....	11
.....	12
.....	12
.....	13
.....	13
.....	14
.....	15
e	15
.....	16
.....	17
.....	17
.....	18
.....	19
.....	20
.....	20
.....	20
.....	22
.....	22
.....	23
.....	23
.....	24
.....	24
.....	27
.....	27
.....	27
.....	29
Index	55

QE747
T4W4

BRUCE L. CLARK

Earth
Sci. Lib.

EARTH
SCIENCES

THE UNIVERSITY OF CHICAGO
LIBRARY
540 EAST 57TH STREET
CHICAGO, ILL. 60637

QE747
T4W4

BRUCE L. CLARK

Earth
Sci. Lib.

CONTENTS.

	Page.
Prefatory note.....	9
Bibliography	7
Description of species.....	11
Mollusca	11
Pelecypoda	11
Arcidae	11
Pinnidae	11
Pernidae	12
Ostreidae	12
Pectinidae	13
Spondylidae	13
Anomidae	14
Mytilidae	15
Pholodomyacidae	15
Pachymyidae	16
Pleurophoridae	17
Crassatellitidae	17
Veneridae	18
Gastrochaenidae	19
Gasteropoda	20
Naticidae	20
Cerithiidae	20
Fusidae	22
Volutidae	22
Acteonidae	23
Ringiculidae	23
Cephalopoda	24
Turrilitidae	24
Arthropoda	27
Crustacea	27
Brachyura	27
Plates	29
Index	55

ILLUSTRATIONS

- Plate I. Arcidae, Pinnidae, Pernidae, Ostreidae, Pectinidae.
Plate II. Spondylidae, Anomidae, Mytilidae.
Plate III. Pholodomyacidae.
Plate IV. Pholodomyacidae.
Plate V. Pachymyidae.
Plate VI. Pachymyidae.
Plate VII. Pleurophoridae, Veneridae, Crassatellitidae, Gastrochaenidae.
Plate VIII. Naticidae.
Plate IX. Naticidae.
Plate X. Cerithiidae, Naticidae, Acteonidae, Ringiculidae.
Plate XI. Fusidae, Volutidae.
Plate XII. Turrilitidae.
Plate XIII. Brachyura.

BIBLIOGRAPHY.

- Bösc, E.*, Bol. Instituto Geologico de México, Nums. 24, 25.
1906, 1910.
- Choffat, P.*, Recueil D'Etudés Paléontol. Faune Cret., du Portugal. 1886.
- Conrad, T. A.*, Rept. U. S. and Mexican Bound. Surv., Vol. I,
Pt. 2, 1857.
- Cragin, F. W.*, Fourth Ann. Rept. Texas Geol. Surv., 1893.
- d'Orbigny, A.*, Paleont. francaise, Terr. Cret., 1840-1860.
- Gabb, W. M.*, Paleont. Cal., Vols. I and II, 1864-1869.
- Goldfuss, A.*, Petfr. Germaniae, 1862.
- Hill, R. T.*, Texas Geol. Surv. Bull. No. 4, 1889.
Paleont. of the Cret. Form. of Texas, 1889.
- Hill, R. T.*, and *Vaughan, T. W.*, U. S. Geol. Surv., Bull. 151,
1898.
- Lasswitz, Rudolf*, Geol. und Paleont. Abh. (E. Koken), 1904.
- Roemer, F.*, Die Kreidebildungen von Texas, 1852.
Neues Jahrbuch Min., Geol., und Paleont., 1887.
- Shattuck, G. B.*, U. S. Geol. Surv., Bull. 205, 1903.
- Shumard, B. F.*, Proc. St. Louis Acad. Sci., Vol. I, 1856-1860.
Proc. Boston Soc. Nat. Hist., Vol. VIII, 1862.
- Stolicka, F.*, Paleont. Indica, 1866-73.
- White, C. A.*, Smithson. Miss. Coll., Vol. XIX, 1880.
Eleventh Ann. Rept. U. S. Geol. and Geog. Surv. Terr.,
1877.
Twelfth Ann. Rept. U. S. Geol. and Geog. Surv. Terr.,
1882.
Archivos Museu Nacional Rio de Janeiro, 1887.
- Zekeli, F.*, Die Gasterop. der Gosaugebilde, 1852.

FAUNA OF THE BUDA LIMESTONE.

BY F. L. WHITNEY.

PREFATORY NOTE.

The species described in this paper were first figured and described in a thesis presented to the faculty of Cornell University for the A. M. degree.

A close examination of material collected since the completion of the above thesis has made necessary a few slight changes. Some species have been withheld from publication for further study with more satisfactory material, and a few forms have been added.

I am deeply indebted to Dr. F. W. Simonds and Prof. G. D. Harris for many helpful suggestions and to my wife, who, by careful collection and presentation of specimens, has added greatly to the number of species which I hope to figure and describe in the near future.

DESCRIPTION OF SPECIES.

MOLLUSCA.

PELECYPODA.

ARCIDAE.

Genus BARBATIA Gray.

BARBATIA SIMONDSI n. sp.

Pl. I, fig. 6.

Dimensions.—Length, 55mm.; breadth, 32mm.

Description.—Shell medium, elongate, inequilateral; beaks near anterior end, inflated, directed forward; anterior margin short, rounded; ventral margin nearly straight; posterior margin rounded, slightly truncate above; dorsal margin subparallel to the ventral; side of valve with a depression extending from the beak obliquely backward and downward to the ventral margin; surface marked with concentric lines of growth crossed by fine, well pronounced, radiating striae, which cover the entire surface of the shell.

This shell resembles *Arca galliennei* d'Orbigny, but it is slightly different in form, and the striae do not alternate in size. It also resembles *Barbatia micronema* Meek and *Arca galliennei* var. *tramitensis* Cragin, but comparison with Cragin's type, the equivalent of Meek's species, shows that the form and markings are somewhat different.

Localities.—Barton Creek and Shoal Creek, Austin, Texas.

PINNIDAE.

Genus PINNA Linnaeus.

PINNA sp.

Pl. I, fig. 3.

Description.—There are, at hand, two specimens of this species, but both are too fragmentary for description. The specimen figured was found near the contact of the Buda with the Del Rio.

Localities.—Shoal Creek and Bouldin Creek, Austin, Texas.

PERNIDAE.

Genus INOCERAMOUS Sowerby.

INOCERAMOUS sp.

Pl. I, fig. 5.

Dimensions.—Length, 50mm.; breadth, 35mm.*Description.*—Shell subovate, ventricose, inequilateral; hinge oblique to long axis; umbo narrow, prominent, incurved. Shell ornamented externally and internally with rounded, concentric plications, separated by grooves of similar width. The plications are very fine at the beak, but gradually increase in size toward the base.

There is only one specimen of this species in this collection, and it is too fragmentary for determination. The outlines shown in the figure may not be accurate, but are the result of following down the parts of shell and cast while removing the rock covering them.

Locality.—Barton Creek, Austin, Texas.**OSTREIDAE.**

Genus ALECTRYONIA Fischer.

ALECTRONIA CARINATA Lamarck.

Pl. I, figs. 1, 2.

Ostrea carinata Roemer, 1849, Texas, p. 394.*Ostrea carinata* Roemer, 1852, Die Kreide. von Texas, p. 75, Taf. 9, fig. 5.*Ostrea carinata* Conrad, 1857, Rept. U. S. and Mex. Bound. Surv., Vol. I, Pt. 2, p. 156, Pl. 10, fig. 6.*Ostrea carinata* White, 1884, Fourth Ann. Rept. U. S. Geol. Surv., p. 293, Pl. 43, figs. 1-4.*Ostrea carinata* Hill, 1889, Texas Geol. Surv., Bull. No. 4, p. 6.*Alectryonia carinata* Hill, 1898, U. S. Geol. Surv., Bull. No. 151, p. 26.*Dimensions.*—Length, 42mm.; breadth, 10mm.*Description.*—Shell narrow, curved, moderately auriculate; beaks depressed, slightly twisted; surface ornamented with twelve or more strong, narrow, sharp ribs, which, in some speci-

mens, are ornamented with two or three rows of tubercles; ribs more elevated on the convex side of the shell; ventral margin strongly dentate.

The specimens collected from the Buda have not attained such vigorous development as those from the Georgetown.

Locality.—Shoal Creek, Austin, Texas.

PECTINIDAE.

Genus PECTEN Mueller.

PECTEN WRIGHTII (Shumard).

Pl. I, fig. 4.

Janira wrightii Shumard, 1860, Trans. Acad. Sci., St. Louis, Vol. I, 1856-1860, p. 607.

Vola (Janira) wrightii Hill, 1889, Geol. Surv. Texas, Bull. No. 4, p. 8.

Vola wrightii Cragin, 1892, Fourth Ann. Rept. Geol. Surv. Texas, p. 217, Pl. 32, figs. 2, 3.

Dimensions.—Length, 22mm.; height, 28mm.

Original description.—Shell ovate, subtrigonal, longest diameter from beak to base; superior valve flat or slightly concave, marked with three strong, rounded, plications, which are prolonged at the base into prominent angles; inferior valve strongly convex; anterior and posterior slopes abrupt, margins straight and diverging from the beak at an angle of about 40°; surface ornamented with four very prominent, simple radiating ribs or folds on the body of the shell, and one much less developed within the margin on either side. Both valves are also elegantly marked with crowded, strongly waved concentric, filiform striæ; beaks elongated, slender, incurved; wings unknown.

Locality.—Shoal Creek, Austin, Texas.

SPONDYLIDAE.

Genus SPONDYLUS Linnæus.

SPONDYLUS CRAGINI n. sp.

Pl. II, figs. 7, 8.

Dimensions.—Greatest length, 35mm.; greatest breadth, 27mm.

Description.—Shell variable, inequilateral, inequivalve, oblique. Inferior valve the larger, ventricose, produced, orna-

mented exteriorly with faint costæ and thin lamellæ; interior with fine, flat, wavy, radiating costæ, separated by grooves of equal width. Superior valve convex, ornamented with many, irregular, wavy, radiating costæ, eight of which are stronger than the others, and separated by three strong costæ, often supplemented by weaker ones. The strong ribs sometimes bear lamellar spines, which, in the adult, are confined to the umbonal region. In the young specimens, all the ribs are spine bearing, and the spines extend throughout the length of the shell. Ears transversely ribbed.

This species resembles *S. hystrix* Goldf., both in young and adult stages. However, there are eight heavy costæ on this species, whereas *S. hystrix* has only six.

Locality.—Shoal Creek, Austin, Texas.

SPONDYLUS TEXANUS n. sp.

Pl. II, figs. 1, 2, 3.

Dimensions.—Length, 25mm.; breadth, 23mm.

Description.—Shell small, subtrigonal, inequilateral, auriculate; anterior straight; posterior slightly produced; base regularly rounded; surface marked by fifty or more fine, uneven, irregular, sometimes dichotomous costæ, about equal in width to the grooves separating them. Ears ornamented with fine, oblique, pronounced striæ; interior marked with costæ, corresponding to those on the outside, strongest at base of shell, thereby giving it a crenulate-serrate appearance; muscle impression with a raised border in thickened portion of shell. Right valve unknown.

Locality.—This and the above species were found in the chalky beds, about ten feet below the contact with the upper beds at Thirtieth Street, Shoal Creek, Austin, Texas.

ANOMIIDAE.

Genus ANOMIA Müller.

ANOMIA GENICULATA n. sp.

Pl. II, figs. 4, 5.

Dimensions.—Length, 10mm.; breadth, 11mm.

Description.—Shell small, sub-ovate, acuminate; left valve

convex, geniculate, ornamented with eight strong, radiating costæ, terminating in hollow, lamellar spines, at the geniculation. Below the geniculation, the surface is marked by heavy lines of growth, with only a trace of the costæ. Right valve ornamented with five or more radiating costæ extending to the margin and bearing four concentric rows of spines.

The hinge and interior were unknown when, in my thesis, I described this shell as *Plicatula geniculata* n. sp. Recently, however, material has been gathered, indicating that it should be referred to the genus *Anomia*.

Locality.—Shoal Creek, Austin, Texas.

MYTILIDAE.

Genus MODIOLA Lamarck.

MODIOLA AUSTINENSIS n. sp.

Pl. II, fig. 6.

Dimensions.—Length, about 40mm.; breadth, 18mm.

Description.—Shell elongate, subcuneate, arcuate, inflated anteriorly, compressed posteriorly; beaks subterminal, twisted forward; umbonal ridge prominent, sloping abruptly to the anterior; surface ornamented with concentric lines of growth, crossed by numerous fine striæ radiating from the beak, and covering the entire shell.

Locality.—Shoal Creek, Austin, Texas.

PHOLODOMYACIDAE.

Genus HOMOMYA Agassiz.

HOMOMYA BUDAENSIS n. sp.

Pl. III, figs. 1, 2. Pl. IV, figs. 1, 2.

Dimensions.—Length about 110 mm.; breadth, 60mm.; thickness 61mm.

Description.—Shell large, globose, equivalved, inequilateral, greatly produced and broadened posteriorly; anterior margin short, curved; ventral margin deeply curved; posterior margin broadly rounded; dorsal margin curved, subparallel with ventral; umbos globose, approximate, elevated, curved inward and

slightly forward, situated anteriorly; shell closed anteriorly, gaping posteriorly. Surface marked by lines of growth, which vary considerably in size.

This shell resembles very closely, *H. vulgaris* Shattuck, but differs from it in the curve of the dorsal and ventral margins, the expansion of the posterior, the greater proportional thickness of the shell, and the ornamentation.

Locality.—Shoal Creek, Austin, Texas. From the top of the formation downward, about six feet.

PACHYMYIDAE.

Genus PACHYMYA Sowerby.

PACHYMYA AUSTINENSIS VAR. BUDAENSIS n. var.

Pl. V; Pl. VI.

Pachymya austinensis Shumard, 1859, Trans. Acad. Sci. St. Louis, Vol. I, pp. 604-605.

Pachymya austinensis White, 1879, Eleventh Ann. Rept. U. S. Geol. and Geog. Surv. Terr., p. 298, Pl. 8, figs. 1, a, b; Pl. 5, figs. 7, a, b.

Pachymya austinensis Hill, 1889, Geol. Surv. Texas, Bull. No. 4, p. 15.

Pachymya austinensis Shattuck, 1903, U. S. Geol. Surv., Bull. 205, p. 26.

Dimensions.—Length, about 15cm.; breadth, about 9cm.; thickness, about 5.5cm.

Description.—Shell large, heavy, gibbose; beaks anterior, incurved, approximate; umbonal slope angulated from the beaks to the posterior extremity; anterior margin short, narrowly rounded; posterior margin truncate, narrowly rounded to angulate below; ventral margin concave in the middle, more rounded anteriorly than posteriorly; dorsal margin nearly straight, subparallel to the ventral; sides, with a depression extending from below the beaks, obliquely backward and downward to the ventral margin; surface ornamented with strong, concentric lines of growth, crossed by granulate lines radiating from the beaks.

This shell is very closely related to *P. austinensis* Shum. and may be identical with it, but the Buda type is much higher posteriorly, and not so greatly angulated on the umbonal slope.

Locality.—Shoal Creek, Austin, Texas.

PLEUROPHORIDAE.

Genus ARTICA Schumacher.

ARTICA COMPACTA (White).

Pl. VII, figs. 1, 2, 3.

Pachymya ? compacta White, 1880, Proc. U. S. Nat. Mus., Vol. II, p. 297, Pl. 6, figs. 3, 4. (Smithson. Misc. Coll., Vol. 19.)

Pachymya ? compacta White, 1883, Twelfth Ann. Rept. U. S. Geol. and Geog. Surv. Terr., Pt. I, pp. 22, 23, Pl. 17, figs. 4a, b.

Dimensions.—Length, 30mm.; breadth, 19mm.; thickness, 16mm.

Original description.—Shell small, narrower posteriorly than anteriorly, slightly gaping behind; beaks depressed, approximate, incurved, directed forward, their position being very near the front; basal margin broadly convex; posterior margin narrowly rounded; postero-dorsal margin forming an oblique downward and backward truncation of that part of the shell; cardinal margin nearly straight, subparallel with the basal margin, much shorter than the full length of the shell; ligament short, its area depressed and sharply defined; front very short, depressed beneath the beaks and narrowly rounded below; umbonal ridges prominent and angular or subangular; the space above and behind them, moderately broad and flattened; the remainder of each valve somewhat regularly convex. Hinge and interior markings unknown. Surface marked by the ordinary concentric lines of growth.

A study of the material found in the Buda, shows that this shell is not a *Pachymya*, but either a *Trapezium* (*Cypriocardia*), as White suspected, or an *Artica*. The material that I have been able to study indicates, by its hinge structure, that it belongs to the latter genus.

Locality.—Shoal Creek, Austin, Texas.

CRASSATELLITIDAE.

Genus STEARNSIA White.

STEARNSIA ROBINSI White.

Pl. VII, fig. 7.

Stearnsia robinsi White, 1887, Proc. Acad. Nat. Sci. Phila., Pt. 1, for 1887, pp. 33, 34, Pl. 2, figs. 7-9.

Astarta (Stearnsia) robinsi Hill, 1889, Texas Geol. Surv., Bull. No. 4, p. 11.

Dimensions.—Length, 22mm.; breadth, 16mm.

Original description.—Shell much compressed, trihedral in marginal outline; lunule long and narrow, nearly straight from end to end, concave transversely; escutcheon similar in shape and character to the lunule, but longer; beaks small, appressed, angular; ligament slightly exposed, and it appears to have been divided into an outer and inner portion by a calcareous septum; umbonal furrows distinct, producing an emargination at the posterior part of the convex basal border and considerable prominence of the posterior extremity; hinge strong; the lateral teeth slender and extending the full length of the lunule and escutcheon respectively; surface marked by strong concentric furrows and ridges, which end abruptly at the margin of the lunule and escutcheon respectively; the surface of both lunule and escutcheon plain.

The material studied seems to be a little different from that figured by White, but a comparison with specimens from the Del Rio and a probable Georgetown rock, leads me to believe that the differences are not sufficient to be specific.

Locality.—Shoal Creek, Austin, Texas.

VENERIDAE.

Genus MERETRIX Lamarck.

MERETRIX LEONENSIS ? (Conrad).

Pl. VII, fig. 4.

Cytherea leonensis Conrad, 1857, Rep. U. S. and Mex. Bound. Surv., Vol. I, Pt. 2, p. 153, Pl. 6, fig. 1.

Cytherea leonensis Hill, 1889, Texas Geol. Surv., Bull. No. 4, p. 14.

Dimensions.—Length, 35mm.; breadth, 44mm.

Description.—Shell medium, subovate, inequilateral, moderately ventricose; beaks, about one-fourth the length of the shell from the anterior end, small, excavated anteriorly, approximate. Anterior margin slightly curved, rounded below; ventral margin broadly rounded; posterior margin convex, rounded below. Surface ornamented with concentric lines of growth.

The material found in the Buda resembles *C. leonensis*, as figured in the Mexican Boundary Report. However, the anterior margin seems to be more gently curved in my specimens, while

the posterior is more rounded. The proportions also agree closely with those of Conrad's figure. In my thesis, I referred to this as *Meretrix budaensis* n. sp., but the study of more material inclines me to refer it doubtfully to Conrad's species.

This species appears to be quite common in the upper layers of the Buda, but it is not well preserved.

Localities.—Shoal Creek, Barton Creek, and Bouldin Creek, Austin, Texas.

Genus TAPES Mergerle von Muhlfeldt.

TAPES AUSTINENSIS n. sp.

Pl. VII, figs. 8, 9.

Dimensions.—Length, 25mm.; breadth, 17mm.

Description.—Shell small, ovate-elongate, narrowly rounded anteriorly and posteriorly; posterior broader than anterior; ventral margin broadly convex; dorsal margin convex, gently sloping downward from the beaks to the posterior margin; beaks approximate, situated about three-fifths the length of the shell from the posterior end. Surface marked by fine, unequal, concentric lines of growth.

Locality.—Shoal Creek, Austin, Texas.

GASTROCHAENIDAE.

Genus FISTULANA Bruguière.

FISTULANA RUPERTI n. sp.

Pl. VII, figs. 5, 6.

Dimensions.—Length, 23mm.; breadth, 12mm.

Description.—Shell elongate, suboval, broader anteriorly than posteriorly, substance thin; beaks nearly terminal; anterior margin broadly rounded; posterior margin narrowly rounded; dorsal and ventral margins convex and diverging from the posterior; beyond the middle the ventral margin turns abruptly to meet the anterior; ventral side gaping for a considerable part of its length; surface smooth, marked by fine lines of growth.

Tube clavate and generally vertical in the rock.

This species is very common in some beds of the Buda, and

seems to range from near the base to the top. The material is quite well preserved, but difficult to remove.

Locality.—Shoal Creek, Austin, Texas.

GASTEROPODA.

NATICIDAE.

Genus TYLOSTOMA Sharpe.

TYLOSTOMA HILLI n. sp.

Pl. VIII, figs. 1, 2. Pl. IX.

Dimensions.—Height about 100 mm.; maximum breadth, 77mm.; minimum breadth, 52mm.

Description.—Shell large, globose, solid, greatly compressed; spire elevated; whorls five and one-half, convex; body whorl much larger than the others, inflated, about two-thirds the height of the shell; surface smooth; varices well developed and strong on the spire of the adult; aperture elongate; lip unknown.

Locality.—Barton Creek, Shoal Creek, Austin, Texas.

TYLOSTOMA HARRISI n. sp.

Pl. X, figs. 12, 13, 14.

Dimensions.—Height, 39mm.; breadth, 24mm.

Description.—Spire elevated, conical, whorls five, convex; suture somewhat canaliculate; lip thin, slightly reflected; inner lip callous; aperture acute posteriorly, rounded, produced anteriorly. Surface smooth, marked by fine lines of growth. The varices are not prominent on the exterior of the type specimen. They are, however, more pronounced on some individuals of this species. They are well developed on casts.

Locality.—Shoal Creek, Barton Creek, Austin, Texas.

CERITHIIDAE.

Genus CERITHIUM Adanson.

CERITHIUM STANTONI n. sp.

Pl. X, figs. 1, 2.

Dimensions.—Height, 32mm.; breadth, 8mm.

Description.—Spire elevated, angle 16° ; whorls twenty-seven or more; aperture rounded; canal short, twisted. Decorations consist of two spiral rows of tubercles to the whorl, situated at its edges; posterior tubercles much larger than anterior of same whorl. As the shell grows larger, the tubercles become more separated.

Figure 1, Pl. X, shows an adult form; figure 2, a younger form.

Locality.—Shoal Creek, Austin, Texas.

CERITHIUM SHUMARDI n. sp.

Pl. X, fig. 3.

Dimensions.—Height, 55mm.; breadth, 30mm.

Description.—Shell turreted; whorls six or more, moderately convex; spire straight, angle about 31° ; sutures well defined; aperture subquadrate; acute anteriorly-posteriorly; columella simple; lip unknown; surface ornamented with fourteen transverse costæ. These costæ and the intervening spaces are traversed by five to seven heavy, prominent, rounded, revolving lines, separated by spaces about equally as wide, and ornamented with five to seven fine, variable, prominent, rounded lines. This ornamentation extends over about one-half the body whorl, the lower portion lacking the transverse costæ but ornamented with the same revolving lines, the heavy ones being somewhat tuberculate. This species is very similar to *C. aguileræ* Böse, but the lines are all single, whereas in that species some are paired. It has the single lines similar to *C. debile* Zekeli, but the transverse costæ are more numerous. These three species appear to be closely related.

Locality.—Shoal Creek, Barton Creek, Austin, Texas.

CERITHIUM HILLI n. sp.

Pl. X, fig. 4.

Description.—Shell turreted; spiral angle about 19° ; number of whorls unknown; surface ornamented with twenty-four transverse ribs set end to end, giving the shell a fluted appearance, broken only by the suture lines. On the poorly preserved specimens at hand, there is no evidence of revolving lines.

This species appears to be quite rare, only one specimen having been obtained. It reminds one of *C. haidingeri* Zekeli, but differs from that species in the angle of the spire and the character of the transverse costæ.

Locality.—Shoal Creek, Austin, Texas.

FUSIDAE.

Genus FUSUS Lamarck.

FUSUS SIMONDSI n. sp.

Pl. XI, fig. 2.

Dimensions.—Length, 100mm.; breadth, about 30mm.

Description.—Shell elevated, turreted; whorls nine, rounded; sutures well defined; spire straight, conical, equalling about one-half the height of the shell; angle 25° ; anterior canal long, straight, deep; lip and aperture unknown; surface of whorls ornamented with eight to ten transverse costæ, which, with the intervening spaces, are crossed by four to six prominent, rounded, revolving lines. The costæ are prominent on the upper part of the body whorl, but disappear below, leaving only the revolving lines.

This species resembles *F. texanus* Shattuck, but the spire and body whorl are nearly equal, and the angle of the spire is much less.

Locality.—Shoal Creek, Austin, Texas.

VOLUTIDAE.

Genus VOLUTILITHES Swainson.

VOLUTILITHES AUSTINENSIS n. sp.

Pl. XI, fig. 1.

Description.—Shell small, fusiform; spire elevated, sutures well defined; body whorl large, constituting the greater part of the shell, shoulder rounded; aperture narrow, elongate; canal long, straight, open and deep; columella with three strong, oblique plaits near the middle of the aperture; lip thin. Surface ornamented with about eighteen prominent, transverse costæ crossed by numerous, fine rounded, revolving lines.

Locality.—Shoal Creek, Austin, Texas.

ACTEONIDAE.

Genus CYLINDRITES Lycett.

CYLINDRITES WHITEI n. sp.

Pl. X, fig. 8.

Dimensions.—Height, 4.7mm.; breadth, 2.5mm.

Description.—Shell small; surface marked by prominent lines of growth; whorls four; spire depressed, canaliculate; aperture narrow, extending posteriorly to the end of the body whorl, anteriorly extended below the body whorl and narrowly rounded; columella with one fold.

There is, in this collection, only one specimen of this species, but it is quite unlike any other that has come to my notice.

Locality.—Shoal Creek, Austin, Texas.

RINGICULIDAE.

Genus CINULIA Gray.

CINULIA CONRADI n. sp.

Pl. X, figs. 5, 6, 7.

Dimensions.—Height, 9mm.; greatest breadth, 7mm.

Description.—Shell small, imperforate, subglobose; whorls three, spire short, about one-eighth of the height of the shell; aperture elongate, acute posteriorly, rounded anteriorly; outer lip thickened, smooth, reflected; inner lip callous, with three plaits, two of which are well up on the shell; surface of body whorl ornamented with fourteen rather broad, spiral costella, separated by narrow grooves developed by a series of spirally arranged puncta. The lip is somewhat broken in the specimen studied, but it appears to have been ornamented with two or more vertical lines.

Locality.—Shoal Creek, Austin, Texas.

CINULIA PELLETI n. sp.

Pl. X, figs. 9, 10, 11.

Dimensions.—Height, 18mm.; maximum breadth, 13mm.

Description.—Shell small, solid, subglobose; whorls about three, rounded; spire about one-sixth of the height of the shell;

aperture slightly oblique, rounded anteriorly, acute posteriorly; outer lip thickened, reflected, smooth within; inner lip callous, with three plaits, the upper one slightly above the middle of the aperture, the others well down. Surface marked with about twenty-seven revolving, well defined costella, separated by grooves equally as wide, and crossed by numerous fine, transverse lines.

Locality.—Batron Creek, Austin, Texas.

CEPHALOPODA.

TURRILITIDAE.

Genus TURRILITES Lamarek.

TURRILITES BRAZOENSIS Roemer.

Pl. XII, fig. 1.

Turrilites brazoensis Roemer, 1849, Texas, p. 415.

Turrilites brazoensis Roemer, 1852, Die Kreide. von Texas, p. 37, Taf. 3, fig. 2.

Turrilites brazoensis Hill, 1889, Texas Geol. Surv., Bull. No. 4, p. 23.

Original description.—The shell large, sinistral, tuberculate; whorls almost quadrilateral, flat outside, ornamented with four rows of large tubercles; the tubercles of the uppermost row standing on the obtusely rounded margin between the broad scarcely arched lateral surface of the whorl and the narrow superior surface, and extending themselves into plain plaits covering the superior surface; both middle rows consist of large, pointed tubercles; the lowest row of tubercles, composed of smaller tubercles, standing on the margin of the inner surface of the whorls and covered by the following whorl. The inner surface, turned toward the axis of the conical spire, covered with perpendicular plain plaits, which correspond to the tubercles of the lowest row.

The specimens collected from the Buda are mere fragments. However, a fairly complete specimen from the Georgetown formation measures in height 20cm.; breadth, 9cm.

Locality.—Barton Creek, Shoal Creek, Austin, Texas.

TURRILITES ROEMERI n. sp.

Pl. XII, figs. 2, 3.

Dimensions.—Height, 36mm.; breadth, 30mm.

Description.—Shell turriculate, conical, sinistral; angle 49° ; volutions angular in front, convex behind, overlapping; suture deep; surface ornamented with four rows of unequal tubercles; the anterior row with about thirty-two tubercles, placed on the sutural angle at the outer end of ribs which converge toward the umbilicus; a second row of narrow more elongate tubercles is placed parallel to and behind the first row. It has the same number of tubercles as the first. Below the second, is a third row of elongate tubercles, of the same number as those in the other rows. The fourth row consists of large, rounded tubercles placed in the middle of the volution, and numbering about eleven.

This species differs from *T. tuberculatus* Bosc, by the spiral angle and the shape of the tubercles; from *T. gravesianus* d'Orbiny, by the more elongate form of tubercle and the angle. In this species, the three rows of smaller tubercles are covered by the overlapping of the adjacent whorl, whereas in *T. tuberculatus* and *T. gravesianus* only one row is overlapped. It appears to be rare in the Buda, only one specimen having been found.

Locality.—Shoal Creek, Austin, Texas.

ARTHROPODA.

CRUSTACEA.

BRACHYURA.

Indeterminate Crustacean.

Pl. XIII, fig. 1.

There is, in my collection, a specimen of the great chela of a crustacean, possibly *Paramithrax*, but it is so poorly preserved that its determination is very doubtful.

Locality.—Shoal Creek, Austin, Texas.

Genus *GRAPTOCARCINUS*.

GRAPTOCARCINUS TEXANUS Roemer.

Pl. XIII, figs. 2, 3.

Graptocarcinus texanus Roemer, 1887, Neues Jahrbuch f. Mineral. und Paleont. Band 1, Heft. 2, pp. 173-176, figs *a* and *b*, text.

Dimensions.—Length, 28mm.; breadth, 36mm.

Description, original somewhat modified.—Cephalothorax transverse oval, rounded, five-sided in outline; enclosed by a fine, raised, sharp border separating the dorsal and ventral surfaces; dorsal surface moderately convex, flattened centrally; frontal bent downwards, ending in a broad, obtuse-angled, pointed rostrum, divided medially by a fine, longitudinal furrow; outer margin of rostrum raised; orbits deeply cut. Surface coarsely and thickly granulate; between the grains or tubercles are scattered separate smaller ones. Where the tubercles are not completely preserved and crumbled on the summits, they appear ring-like and deepened in the centre. On the central surface are two pairs of converging furrows. The furrows of the anterior pair are longer and stronger than those of the posterior. They are bent slightly inwards and converge at an angle of about 130° . They are deepest and broadest at their outer ends. At their inner ends, they are so weak that their junction is scarcely visible. The furrows of the posterior

pair are at first parallel to the anterior pair, then bent abruptly backwards and weakened till scarcely traceable. No other sculpture is present on the dorsal surface. The surface of the sloping and inwardly bent ventral side is almost smooth. It is finely granulate only opposite and just behind the frontal. Opposite the front pair of furrows is a small notch, in the sharp margin separating the dorsal and ventral surfaces, from which a fine, line-like furrow passes forward in a moderate curve.

Localities.—Shoal Creek, Barton Creek, Austin, Texas.

PLATE I.

PLATE 1.

	Page.
Figs. 1, 2. <i>Alectryonia carinata</i> (Lam.).....	12
Fig. 3. <i>Pinna</i> sp.....	11
Fig. 4. <i>Pecten wrightii</i> Shum.....	13
Fig. 5. <i>Inoceramous</i> sp.....	12
Fig. 6. <i>Barbatia simondsi</i> n. sp.....	11

1

2

3

4

5

6

PLATE II.

PLATE II.

	Page.
Figs. 1-3. <i>Spondylus texanus</i> n. sp.....	14
Figs. 4, 5. <i>Anomia geniculata</i> n. sp.....	14
Fig. 6. <i>Modiola austinensis</i> n. sp.....	15
Figs. 7, 8. <i>Spondylus eragini</i> n. sp.....	13

DEAR SIR: I take pleasure in sending to you a copy of Bulletin No. 9 of The University of Texas Mineral Survey, entitled "Report of a Reconnaissance of Trans-Pecos Texas, North of the Texas & Pacific Railway," by George Burr Richardson.

This Bulletin is the ninth in the series which was begun in July, 1901. The other Bulletins are as follows:

1. Texas Petroleum, July, 1901.
2. Sulphur, Oil and Quicksilver in Trans-Pecos Texas, February, 1902.
3. Coal, Lignite and Asphalt Rocks, May, 1902.
4. The Terlingua Quicksilver Deposits, Brewster County, October, 1902.
5. The Minerals and Mineral Localities of Texas, January, 1903.
6. The Mining Laws of Texas and Tables of Magnetic Declination, July, 1903.
7. Report of Progress for 1903, and Topographic Map of Terlingua Quadrangle in Brewster and Presidio Counties, January, 1904.
8. The Geology of the Shafter Silver Mine District, Presidio County, June, 1904.

These Bulletins are for gratuitous distribution among the citizens of the State and to others upon application.

The editions of Bulletins Nos. 1 to 5, inclusive, have been exhausted.

Very truly,

WM. B. PHILLIPS,

Director.

Austin, Texas, November, 1904.

1

2

3

4

5

6

7

8

PLATE III.

PLATE III.

	Page.
Figs. 1, 2. <i>Homomya budaensis</i> n. sp.....	15

1

2

PLATE IV.

PLATE IV.

	Page.
Figs. 1, 2. <i>Homomya budaensis</i> n. sp.	15

1

2

PLATE V.

PLATE V.

	Page.
<i>Pachymya austinensis</i> var. <i>budaensis</i> n. var.....	16

1

PLATE VI.

PLATE VI.

	Page.
<i>Pachymya austinensis</i> var. <i>budaensis</i> n. var. x .8.....	16
Showing surface ornamentation.

1

PLATE VII.

PLATE VII.

	Page.
Figs. 1-3. <i>Artica compacta</i> (White).....	17
Fig. 4. <i>Meretrix leonensis?</i> (Con.).....	18
Fig. 5. <i>Fistulana rupertii</i> n. sp.....	19
Fig. 6. Tubes of <i>Fistulana rupertii</i>	19
Fig. 7. <i>Stearnsia robinsi</i> White.....	17
Figs. 8, 9. <i>Tapes austinensis</i> n. sp.....	19

1

2

3

4

5

6

7

8

9

PLATE VIII.

PLATE VIII.

	Page.
<i>Tylostoma hilli</i> n. sp.	20

1

2

PLATE IX.

PLATE IX.

	Page.
<i>Tylostoma hilli</i> n. sp.	20

PLATE X.

PLATE X.

	Page.
Figs. 1, 2. <i>Cerithium stantoni</i> n. sp.....	20
Fig. 3. <i>Cerithium shumardi</i> n. sp.....	21
Fig. 4. <i>Cerithium hilli</i> n. sp.....	21
Figs. 5-7. <i>Cinulia conradi</i> n. sp.....	23
Fig. 8. <i>Cylindrites whitei</i> n. sp. x3.....	23
Figs. 9-11. <i>Cinulia pelleti</i> n. sp.....	23
Figs. 12-14. <i>Tylostoma harrisi</i> n. sp.....	20

PLATE XI.

PLATE XI.

	Page.
Fig. 1. <i>Volutilithes austinensis</i> n. sp.....	22
Fig. 2. <i>Fusus simondsi</i> n. sp.....	22

1

PLATE XII.

PLATE XII.

	Page.
Fig. 1. <i>Turrilites brazoensis</i> Roemer.....	24
Figs. 2, 3. <i>Turrilites roemeri</i> n. sp.....	24

1

2

3

PLATE XIII.

PLATE XIII.

	Page.
Figs. 1, 2. <i>Graptocarcinus texanus</i> Roemer	27
Fig. 3. Crustacean indet x .75.....	27

1

2

3

JOSE L. CLARK

INDEX.

[Names in *italic* are synonyms; figures in black-face type are numbers of page on which detailed descriptions appear; figures in *italic* denote illustrations.]

- Acteonidae23
 Alectryonia Fischer12
 Alectryonia carinata
 (Lamarck)12, 30
 Alectryonia carinata Hill.....12
 Anomia Müller14
 Anomia geniculata n. sp.14, 32
 Anomidae14
 Arca galliennei d'Orbigny.....11
 galliennei var. tramitensis
 Cragin11
 Arcidae11
 Arthropoda27-28
 Artica Schumacher17
 compacta (White)17, 42
 Barbata Gray11
 micronema Meek11
 simondsi n. sp.11, 30
 Brachyura27-28
 Cephalopoda24-26
 Cerithiidae20
 Cerithium Adanson20
 aguilerae Böse21
 debile Zekeli21
 haidingeri Zekeli22
 hilli n. sp.21, 48
 schumardi n. sp.21, 48
 stantoni n. sp.20, 48
 Cinulia Gray23
 conradi n. sp.23, 48
 pelleti n. sp.23, 48
 Crassatellitidae17
 Crustacea27
 Crustacean indet27, 54
 Cylindrites Lycett23
 whitei n. sp.23, 48
 Cytherea leonensis Conrad.....18
 leonensis Hill18
 Fistulana Bruguière19
 ruperti n. sp.19, 42
 Fusidae22
 Fusus Lamarck22
 simondsi n. sp.22, 50
 texanus Shattuck.....22
 Gasteropoda20-24
 Gastrochaenidae19
 Graptocarcinus texanus Roe-
 mer27, 54
 texanus Roemer27
 Homomya Agassiz15
 budaensis n. sp.15, 34, 36
 vulgaris Shattuck16
 Inoceramous Sowerby12
 sp.12, 30
 Janira wrightii Shumard.....13
 Meretrix Lamarck18
 budaensis n. sp.19
 leonensis? (Conrad)18, 42
 Modiola Lamarck15
 austinensis n. sp.15, 32
 Mollusca11-26
 Mytilidae15
 Ostreidae12
 Ostrea carinata Roemer et al....12
 Pachymyidae16
 Pachymya Sowerby16
 austinensis var. budaensis
 n. var.16, 38, 40
 austinensis Shumard et al....16
 compacta White17
 Pecten Müller13
 wrightii (Shumard)13, 30
 Pectinidae13
 Pelecypoda11-20
 Pernidae12
 Pholodomyacidae15
 Pinna Linnaeus11
 sp.11, 30
 Pinnidae11
 Pleurophoridae17
 Plicatula geniculata n. sp.15
 Ringiculidae23
 Spondylidae13
 Spondylus Linnaeus13
 cragini n. sp.13, 32
 hystrix Goldfuss14
 texanus n. sp.14, 32
 Stearnsia White17
 robinsi White17, 42
 robinsi White17
 Tapes Mergerle von Muhl-
 feldt19
 austinensis n. sp.19, 42
 Turrillites Lamarck24
 brazoensis Roemer24, 52
 brazoensis Roemer24
 gravesianus d'Orbigny.....25
 roemerian n. sp.24, 52
 tuberculatus Bosc25
 Turrilitidae24
 Tylostoma Sharpe20
 harrisi n. sp.20, 48
 hilli n. sp.20, 44, 46
 Veneridae18
 Vola (Janira) wrightii Hill.....13
 Vola wrightii Cragin.....13
 Volutidae22
 Volutilithes Swainson22
 austinensis n. sp.22, 50

ERUCE L. CLARK

BRUCE L. CLARK

BRUCE L. CLARK

THE UNIVERSITY OF TEXAS BULLETIN.

(Continued from inside front cover)

- Railroad Transportation in Texas*, by C. S. Potts. 214 p., 6 maps, charts. March, 1909. \$1.50.
- Crime and the Treatment of the Criminal*, by C. S. Potts. 86 p. May 1, 1910. 25 cents.
- Problems of Prison Reform*, by C. S. Potts. 40 p. December 8, 1910. 20 cents.
- Recognition of the Republic of Texas by the United States*, by Ethel Zivley Rather. 101 p. January 1, 1911. 75 cents.
- English Elements in Jonson's Early Comedy*, by Charles Read Easker-vill, 328 p. April 8, 1911. \$2.50.

MEDICAL SERIES

- Yellow Fever: a Popular Lecture*, by James Carroll. 32 p. June, 1905. 15 cents.
- The Care of the Insane*, by Dr. M. L. Graves. 16 p. 1905. 15 cents.
- The 1903 Epidemic of Yellow Fever in Texas and the Lessons to Be Learned from It*, by Dr. G. R. Tabor. 22 p. June, 1905. 15 cents.
- Further Experiments in the Use of Drugs as Stimulants in Accidents Occurring During Anaesthesia*, by O. H. Plant, 31 p. February 8, 1911. 25 cents.

SCIENTIFIC SERIES

- Vegetation of the Sotol Country in Texas*, by W. L. Bray. 24 p., pl. June, 1905. 25 cents.
- Observations on the Habits of Some Solitary Wasps of Texas*, by Carl Hartman. 72 p., pl. July, 1905. 25 cents.
- Distribution and Adaptation of the Vegetation of Texas*, by W. L. Bray, 108 p., pl. map. November, 1906. 35 cents.
- A Sketch of the Geology of the Chisos Country*, by J. A. Udden. 101 p. April, 1907. 50 cents.
- The Clays of Texas*, by Heinrich Ries, 316 p., illus. pl. 1908. \$2.00.
- The American Mistletoe*, by H. H. York. 31 p., pl. 1909. 50 cents.
- Symptoms of Disease in Plants*, by F. D. Heald. Illus. November, 1909. \$1.00.
- Field Studies of the Behavior of the Lizard *Sceloporus Floridanus**, by H. H. Newman and J. Thomas Patterson. 23 p., illus. December, 1909. 25 cents.
- The Austin Dam*, by T. U. Taylor, 85 p., illus. December 22, 1911. 75 cents.
- On the Electrostatic Effect of a Changing Magnetic Field*, by J. M. Kuehne. 15 p. January 15, 1911. 25 cents.
- Fauna of the Buda Limestone*, by Francis Luther Whitney. 54 p., illus. May 22, 1911. \$1.00.

to those who desire it a complete list of the publications of the University of Texas will be furnished. Requests for this or for Bulletins should be addressed to the University of Texas Bulletin, Austin, Texas. Exchanges should be addressed to the University of Texas Library.

THE UNIVERSITY OF TEXAS

Co-educational.

Tuition Free.

ANNUAL EXPENSES \$180 AND UPWARDS

MAIN UNIVERSITY AT AUSTIN.

COLLEGE OF ARTS: Courses leading to the Degrees of Bachelor and Master of Arts and Doctor of Philosophy.

DEPARTMENT OF EDUCATION: Professional courses for teachers, leading to elementary and permanent certificates.

ENGINEERING DEPARTMENT: Degree courses in civil, electrical and mining engineering.

LAW DEPARTMENT (in its new building): Three-year course, leading to Degree of Bachelor of Laws, with State license; course leading to Degree of Master of Laws.

SUMMER SCHOOL: Regular University and Normal courses; seven weeks.

Session of 1911 begins June 18.

For catalogue, address

THE REGISTRAR,
University Station, Austin.

DEPARTMENT OF EXTENSION: I. Correspondence Division, offering courses in various University schools, for which registration may take place at any time. II. Public Discussion and Information Division, through bibliographies and traveling libraries supplying information on current problems. III. Lecture Division, presenting members of the University Faculty in popular lectures, singly or in series.
For catalogue, address

THE DIRECTOR OF EXTENSION,
University Station, Austin.

MEDICAL DEPARTMENT AT GALVESTON

Four-year course in medicine; two-year course in pharmacy; three-year course in nursing. Thorough laboratory training. Exceptional clinical facilities in John Sealy Hospital. University Hall, a dormitory for women students of medicine.
For catalogue, address

THE DEAN, Medical College,
Galveston.

14 DAY USE

RETURN TO DESK FROM WHICH BORROWED

EARTH SCIENCES LIBRARY

**This book is due on the last date stamped below, or
on the date to which renewed.**

Renewed books are subject to immediate recall.

[illegible]

LD 21-40m-4,'64
(E4555s10)476

General Library
University of California
Berkeley

293