UC-NRLF

8 2 888 051

GIFT OF

AGRIC.
LIBRARY

Digitized by the Internet Archive
in 2007 with funding from
Microsoft Corporation

MINISTRY OF AGRICULTURE, FISHERIES

TECHNICAL AND SCIENTIFIC SERVICE

Bulletin No. 82

(FORTHY-SIXTH YEAR)

WORK CONDUCTED AT THE EXPERIMENTAL FARM AT WEAVER

(1925-1926)

GREAT MILLET

Echinochloa polystachya - *Eleusine indica*

BEUSSI

Echinochloa polystachya - *Eleusine indica*

E. J. HALL, B.S., M.S.

ASSISTANT DIRECTOR OF AGRICULTURE

U.S. DEPARTMENT OF AGRICULTURE

1926

Printed by the Government Printing Office, Washington, D.C.

MINISTRY OF AGRICULTURE, EGYPT.

TECHNICAL AND SCIENTIFIC SERVICE

Bulletin No. 8.

(BOTANICAL SECTION.)

WORK CONDUCTED AT THE EXPERIMENTAL FARM AT GIZA

IN CONNECTION WITH

GREAT MILLET

(*SORGHUM VULGARE - DHURA RAFIA*)

AND

BERSIM

(*TRIFOLIUM ALEXANDRINUM - BERSIM*),

BY

B. G. C. BOLLAND, B.A.,

BOTANIST TO THE MINISTRY OF AGRICULTURE.

(Submitted for Printing on April 11, 1916.)

CAIRO.

GOVERNMENT PRESS.

To be obtained, either directly or through any Bookseller,
from the GOVERNMENT PRESS, Bulâq, and from the SALE-ROOM,
Old Ismailia Palace, Sharia Qasr el Aini.

1916.

PRICE: P.T. 1.

MINISTRY OF AGRICULTURE, EGYPT.

(Bulletin No. 8.)

(BOTANICAL SECTION.)

WORK IN CONNECTION WITH MILLET IN 1915.

Work has been started on the different varieties of millet growing in the country with a twofold object :—

(1) To ascertain how many of the names given to millet denote separate varieties, and to write detailed descriptions of them.

(2) To improve the yield and quality of the crop by selection work and to isolate and breed pure varieties.

The following notes give the result of the work done in 1915.

The following varieties were sown between May 6 and 12 :—

Variety.	Area.			Locality from where Seed was obtained.
	P.	Q.	S.	
<i>Asfar</i>	1	8	23	Qena.
„	1	8	17	Minia.
<i>Qamhi</i>	0	15	21	Qena.
„	0	15	21	Minia.
<i>Sab'ini</i>	1	20	3	Qena.
<i>Tiss'ini</i>	1	8	7	„

The cultivation was normal and was the same for all the varieties.

For manure, nitrate of soda was applied at the rate of about ninety-five kilogrammes per feddân.

Sab'ini was the first to mature and was cut on August 24, while the *Asfar* from Minia was not ready for cutting till October 9.

The following table shows the date of cutting and yield of grain in ardebs per feddân of the different varieties :—

Variety.	Date of Cutting.	Yield.	Number of Days taken in maturing.
<i>Asfar</i> from Qena	September 4	13·45	117
<i>Qamhi</i> " "	" 1	12·80	117
<i>Tiss'ini</i> " "	August 26	10·10	109
<i>Sab'ini</i> " "	" 24	8·40	110
<i>Qamhi</i> " Minia	September 30	5·26	141
<i>Asfar</i> " "	October 9	1·19	150

1 ardeb = 5·44 bushels = 155 kilogrammes.

The sparrows ate a large quantity of grain before it was ripe enough for cutting and so reduced the yield to a considerable extent; especially was that the case with the varieties which were the last to mature, namely *Asfar* and *Qamhi* from Minia.

Detailed observations were taken, but with the exception of the *Asfar* variety from Minia no distinct differences were noticeable.

The *Asfar* from Minia, besides taking a long time to mature, was a much bigger plant in every way than the other varieties.

It was four and a half metres in height, while the next highest variety was *Tiss'ini*, which grew to three metres. The number and length and breadth of leaves were also much greater in that variety.

The following are short descriptions of the different varieties sown :—

Variety.	Maturity.	Colour of Grain.	Colour of Glumes when Ripe.	Head.
<i>Sab'ini</i> ...	Early ...	White	Cream	Compact with a truncate top.
<i>Tiss'ini</i>	" ...	"	Cream, yellow, or black.	Loosely compact with a truncate top.
<i>Qamhi</i> ...	Medium-Late.	White, dark-cream, or reddish	Light brown ...	Compact to loosely compact with a truncate top.
<i>Asfar</i> ...	Medium-Late.	Yellow	Brown "	Loosely compact with rather an indistinct truncate top.

As will be seen, the different kinds are so mixed that no accurate description can be written, and further, in different localities different names are given to the same variety.

As with maize so with millet, a fresh start will be made by writing descriptions of and growing separately as many distinct "heads" as possible.

Millets from the Sudan.—Small samples of twenty-four distinct varieties were obtained from the Sudan and grown at Gîza last year, but no comparisons of yield were made, owing to the smallness of the samples.

The following particulars were recorded, and it is hoped that in 1917 a full description will be written of each variety and the yield determined.

Variety.	Maturity.	Colour of Grain.	Colour of Glumes when Ripe.	Weight of 100 Seeds in Grammes.	Other Notes taken when in Flower or before Harvesting.
1	Medium	Cream	Black	5.18	{ Panicle loose; pedicels and spikelets ciliate. Glumes reddish, hairy; nerves not very prominent except near the tip.
2	Late ...	Orange	Deep claret ...	2.05	
3	Medium	Light orange	Red brown ...	5.08	
4	Late ...	Light brick red.	Black	5.50	{ Panicle loose. Pedicels and spikelets slightly ciliate. Glumes yellowish green, nearly glabrous; prominently nerved towards the apex.
5	" ...	Orange red ...	"	4.28	
6	Early ...	White	Light creamy brown.	2.72	{ Panicle loose. Pedicels and spikelets ciliate. Glumes very large, hairy; greenish white; prominently nerved towards the apex.
7	Late ...	"	Black	5.50	{ Panicle loosely compact. Pedicels and spikelets ciliate. Glumes dull green, hairy; prominently nerved towards the apex.
8	Medium	Light cream	Deep claret ...	3.89	Glumes covering the seed.
9	"	Cream	Claret	3.73	
10	"	Light cream	Light cream...	4.20	
11	Late ...	Light brick red.	Yellow with a tinge of pink.	2.19	
12	Medium	White	Deep claret ...	3.96	Grain small.
13	"	Light orange brown.	Nearly black	4.11	
14	"	Light cream	Cream	1.89	
15	"	Cream	Creamy brown	4.97	
16	Early ...	White	Nearly black	2.55	
17	"	Brown red ...	Deep claret ...	3.46	
18	Medium	Light cream	Red brown ...	3.52	
19	Early ...	"	Light red ...	2.29	
20	Medium	Creamy brown	Black	4.90	
21	"	Dark cream	Deep claret ...	4.00	
22	Early ...	Light cream	Dark brown...	5.21	
23	"	Yellow	Deep claret ...	5.08	
24	"	Light claret	Brown red ...	1.74	

WORK IN CONNECTION WITH *BERSIM* IN 1915.

Bersim is the general term applied to four varieties of clover included under the specified name of *Trifolium alexandrinum*, a plant apparently of Egyptian origin and, as far as is at present known, peculiar in the richness of its crops, which are taken in the winter and throughout the spring.

Small plots of the four varieties, *Ba'li*, *Fahl*, *Khadrawi*, and *Misqawi*, were sown in November 1914 in order to ascertain the characteristic differences of each.

Fahl was found to exhibit a remarkably different appearance from the three others when planted with them and is readily recognized on account of its quicker and more luxuriant growth. *Misqawi* and *Khadrawi* have many common characteristics, and are rather better in growth than *Ba'li*, but they do not approach the same standard as the *Fahl* variety.

For determining the respective yields an area of four square metres was measured off in the centre of each of the twelve plots of the four varieties. The *bersim* from the known area was cut by hand and weighed, the yields being as follows :—

<i>Fahl</i>	...	...	...	6.46	rotls per square metre.
<i>Misqawi</i>	...	...	...	4.53	" " "
<i>Khadrawi</i>	...	...	...	3.60	" " "
<i>Ba'li</i>	...	...	...	3.03	" " "

The forty-eight plots were arranged in chess-board fashion in order to eliminate any irregularities in the soil.

The following table shows the results of the observations which were taken on the height and leaves of the plant.

Variety.	Height. in Centimetres.	Number of Leaves per Plant.	Length of Leaves in Centimetres.	Breadth of Leaves in Centimetres.
<i>Fahl</i>	76.6	42	5.0	1.8
<i>Misqawi</i> ...	44.5	23	4.8	1.9
<i>Khadrawi</i>	56.8	26	4.3	1.7
<i>Ba'li</i>	48.3	23	4.1	1.6

After the *bersâm* had been cut it was noticed that the *Fahl* variety had no fresh growth, while the other varieties had several new stems growing from the base of the main stem, thus enabling them to yield further cuttings.

From the above observations it seems obvious that, if one cutting only is required to be taken, *Fahl* is the variety to grow, the disadvantage of the latter being that the land does not obtain the benefit of a green manuring from the ploughing in of the source of a supply of humus.

To get over that difficulty and at the same time to get a higher yield than if *Misqâwi* alone were sown, a mixture could be made consisting of equal parts of *Fahl* and *Misqâwi*.

Fahl, besides having by far the largest number of leaves, has also the biggest leaves, while *Ba'li* is the least luxuriant of the four varieties. The seeds of the *Fahl* variety are larger than any of the others, so that it can also be distinguished by that character.

PUBLICATIONS OF THE MINISTRY OF AGRICULTURE.

“Agricultural Journal of Egypt.” (English and Arabic.)

Vol. I, Part I. ” ” ” II. ” II, ” I. ” ” ” II.		Vol. III, Part I. ” ” ” II. ” IV, ” I. ” ” ” II. ” V, Parts I and II.
---	--	---

EGYPTIAN AGRICULTURAL PRODUCTS.

By Gerald C. Dudgeon, F.E.S., etc., Consulting Agriculturist.

- No. 1 A. “The Great Millet” (*Durra Baladi* or *Durra Rafī'a*) in Egypt. (English.)
 No. 2 A. “The Ground Nut or Earth Pea (*Ful Sudāni*) in Egypt.” (English.)
-

TECHNICAL AND SCIENTIFIC BULLETINS.

- No. 1. “Report on the First Two Years’ Working of the Plant Protection Law.” (Law No. 5 of 1913), by G. Storey, B.A., F.E.S. (English and French.)
 No. 2. “The Nature of the Damage done by the Pink Boll Worm (*Gelechia Gossypiella* Saund.),” by Lewis Gough, Ph.D., F.E.S., etc. (English and French.)
 No. 3. “Note on the Alkaloids of some Egyptian Solanaceæ,” by Frank Hughes, F.C.S., Chemist. (English and French.)
 No. 4. “The Life History of *Gelechia Gossypiella* from the Time of the Cotton Harvest to the Time of Cotton Sowing,” by Lewis Gough, Ph.D., F.E.S., etc. (English.)
 No. 5. “List of Egyptian Insects in the Collection of the Ministry of Agriculture,” by G. Storey, B.A., F.E.S. (English.)
 No. 6. “Note on a Machine to kill *Gelechia* Larvæ by Hot Air, and the Effects of Heat on *Gelechia* Larvæ and Cotton Seed,” by Lewis Gough, Ph.D., F.E.S., etc. (English.)
 No. 7. “Work in Connection with Egyptian Wheat,” by G. C. Dudgeon, F.E.S. and B. G. C. Bolland, B.A., Botanist. (English.)
 No. 8. “Work conducted at the Experimental Farm at Giza in Connection with Great Millet and *Bersim*,” by B. G. C. Bolland, B.A., Botanist. (English.)
 No. 9. “Work in Connection with Egyptian Maize,” by Gerald C. Dudgeon, F.E.S., Consulting Agriculturist, etc., and B. G. C. Bolland, B.A., Botanist. (English.)
 No. 10. “Some Tests of Flour made from Egyptian Wheat,” by Frank Hughes F.C.S., Membre de l’Institut Egyptien, Chemist. (English.)
-

LEAFLETS OF THE HORTICULTURAL SECTION. (English and Arabic.)

By T. W. Brown, Director, Horticultural Section.

- (No number) "Citrus Trees."
No. 1. "French beans (*Fasoolia*)."
" 2. "Asparagus (*Kishk Almaz*)."
" 3. "Preservation of Dates."
" 4. "Egyptian Arum (*Qolqas*)."
" 5. "Cultivation of Potatoes."
" 6. "Preservation of Apricots."
" 7. "Custard Apple."
-

COLLECTIONS OF AGRICULTURAL CIRCULARS.

From No. 1-18. (English and Arabic.)

- " " 19-35. " "
" " 36-60. " "
-

LAWS AND REGULATIONS.

- "Regulations of the Boll Worm and Seed Worm Campaign, 1915." (English and Arabic.)
"Regulations of Cotton Worm Campaign, 1915." (English and Arabic.)
"Regulations of Cotton Worm Campaign, 1916." (English and Arabic.)
"Collection of the Agricultural Laws." (French and Arabic.)
"Collection of Veterinary Laws." (French and Arabic.)
"Syllabus of the School of Veterinary Medicine." (Arabic and French.)
"Syllabus of the Intermediate School of Agriculture, Moshtohor." (English and Arabic.)
-

ADMINISTRATIVE REPORTS.

- "Annual Report of the Veterinary Service, 1913." (English and Arabic.)
"Annual Report of the Veterinary Service, 1914." (English and Arabic.)
"Report on the Great Invasion of Locusts in Egypt in 1915." (English and Arabic.)
-

EGYPTIAN AGRICULTURAL AND VETERINARY NOTES.

- "Wheat Rust." (English and Arabic.)
"Wheat Smut." (English and Arabic.)
"Pamphlet on Ticks and their Relation with Agricultural Animal Diseases." (Arabic.)
"Pamphlet on Scale Insects." (Arabic.)
-

BOOKS.

- "Agricultural Text Book," Part I. (English and Arabic.)
"Agricultural Text Book," Part II. (English and Arabic.)
"Percival's Botany." (Translation.) (Arabic.)
"Inorganic Chemistry." (Translation.) (Arabic.)
"Agricultural Chemistry," by Ingle. (Translation.) (Arabic.)
"Veterinary Medicine for Intermediate Schools of Agriculture." (Arabic.)
-

Govt. Press 1056-1916-1030 ex.

7 DAY USE
RETURN TO DESK FROM WHICH BORROWED

AGRICULTURE LIBRARY
40 Giannini Hall - Tel. No. 642-4493
This publication is due on the LAST DATE
and HOUR stamped below.

JUN - 5 1971	
NOV 14 1985	
received in Natural Resources Library	
NOV 20 1985	
PHOTOCOPY JUL 6 '88	
NOV 30 1989	
DEC 27 1989	

RB 17-50m-3.'70
(N5381s10)4188-A-32

General Library
University of California
Berkeley

...