THE EISENHOWER LIBRARY

3 1151 02676 8329

54 338

Library

Johns Hopkins University

THE
EMBRYOLOGY

—OF—

CREPIPULA

—BY—

EDWIN GRANT CONKLIN

—•—

A THESIS FOR THE DEGREE OF DOCTOR OF PHILOSOPHY
IN THE JOHNS HOPKINS UNIVERSITY

—

1891

Preliminary Note on the Embryology of *Crepidula fornicata* and of *Urosalpinx cinerea*. By E. G. CONKLIN.

Crepidula fornicata.

The time and manner in which *Crepidula* lays its eggs has been described by Dr. McMurich.¹

The cleavage follows the type found in *Fusus*, *Planorbis*, *Neritina*, etc. The first furrow is, with regard to the future embryo, a median longitudinal one, and divides the ovum into exactly equal right and left halves. (The method of determining the relation of the first furrow to the embryo cannot be explained without several figures). The second furrow is at right angles to the first and transverse to the long axis of the embryo, and it divides the egg into four equal macromeres. Of these, two meet in the centre in a line which Rabl² has called the "cross furrow;" the other two are acute toward the centre and do not meet each other. By the position of the macromeres with regard to the "cross furrow," the first and second cleavage furrows may easily be distinguished, *e.g.*, if the egg be viewed from the formative pole and so that one of the cleavage furrows is in the line of vision, the macromere to the right of this furrow and farthest from the observer will be acute at its centre if the furrow in the line of vision be the second cleavage furrow; it will be obtuse, *i.e.*, will meet the opposite macromeres in the "cross furrow" if the furrow in the line of vision be the first cleavage furrow. Of course the reverse would hold true if the egg were viewed from the vegetative pole. The examination of many hundreds of eggs has shown that the position of the macromeres in relation to the "cross furrow" and to the first and second cleavage planes is a constant one, and that the first and second furrows may always be distinguished in the way mentioned. The macromeres as distinguished from the micromeres do not again divide until late in the course of segmentation, and as they do not change their relative position it becomes very easy to orient all the future furrows and cells with reference to the first two furrows. The polar bodies which are at the centre of the ectodermal area also mark the centre of the dorsal surface of the embryo, while the blastopore closes and the definitive mouth appears almost directly opposite them. Until after the formation of the blastopore the chief axis is the one drawn from the centre of the ectodermal to the centre of the endodermal area; with regard to this axis the egg is for some time radially symmetrical; the chief axis of the embryo (the antero-posterior) is at right angles to the chief axis of the egg. At an early period there is a trace of a segmentation cavity, which however is soon obliterated. The formation of micromeres proceeds with wonderful regularity, following the law formulated by Rabl,³ new micromeres always being formed in fours either by being cut off from the four macromeres or by the division of four micromeres already formed. When twenty-four micromeres have been formed one of the macromeres, which later developments show to be on the posterior side of the ovum and to the left of the median line, divides into a larger ventral and a smaller dorsal moiety. The smaller cell, which, unlike the ectoderm cells, contains yolk, moves to the right until it comes to lie at the posterior end of the median furrow between the two posterior macromeres; it very soon divides into a right and a left half, and at a later stage these two cells give rise to two mesoblastic bands which extend forward over the right and left sides of the dorsal surface.

Save for these mesoblasts the radial symmetry at the stage with thirty-six micromeres is perfect. At this stage twelve of the micromeres

¹ A contribution to the Embryology of the Marine Prosobranchs. Studies from Biol. Lab., J. H. U., Vol. 3.

² Carl Rabl, Ueber die Entwicklung der Tellerschnecke. Morph. Jahrbuch, Bd. 5.

³ *Loc. cit.*

form a cross with three cells in each arm, the crossing of the arms being exactly at the centre of the ectodermal area. Two of the arms lie in the antero posterior axis, two in the transverse axis; one arm is anterior, one posterior, one right and one left. In the next stage the radial symmetry is no longer perfect, since three arms of the cross lengthen so as to contain four cells each, while the posterior arm does not lengthen until later. However, with this exception, the radial symmetry is preserved until fifty-two micromeres are formed; at this stage the posterior arm lengthens by one cell and the three other arms split longitudinally. At the same time each of the macromeres, except the one mentioned above as giving rise to the mesoblast, divides as did the later into a larger and a smaller moiety. The three smaller cells take positions on the periphery of the egg in the furrows separating the macromeres, one at each end of the transverse furrow and one at the anterior end of the longitudinal furrow.

In normal eggs there is not a trace of an invagination at the ectodermal pole, such as has been described for *Neritina* and *Fulga*, though eggs which were developing abnormally often showed such an invagination together with other irregularities, e. g., the spreading apart of the macromeres and the formation of large yolk containing cells at the periphery of the ectodermal area instead of the small protoplasmic ectoderm cells which are usually formed.

The gastrula is formed by typical epibole. On all sides the ectoderm cells grow around toward the ventral side at an equal rate; thus the blastopore is formed in the middle of the ventral side. The mesoblasts are carried around with the ectoderm to the ventral side, where they lie at the posterior edge of the blastopore. The mesoblastic bands are soon separated from the mesoblasts; the latter continue to proliferate mesoderm, which extends forward in the lips of the blastopore.

The cross above mentioned resembles very much an arrangement of cells figured by Blochmann¹ for *Neritina*, the terminal cells in the transverse arms of which are called by him velar cells. In *Crepidula* it seems that no part of the transverse arms forms the velum. However, the cells of the posterior arm grow very large, the nuclei are vesicular and stain lightly, and the cells become covered by fine cilia, which protrude through a thin cuticula; though at first these ciliated cells lie wholly on the posterior side of the ovum, they move forward in the course of development until they come to lie on the anterior side, and as they increase enormously in size, though they seldom divide, it comes about that they cover the anterior part of the dorsal area, extend around over the anterior end of the embryo and down over its sides. These cells finally form the walls of a large head vesicle.

The velum appears first on the ventral side, just anterior to the mouth, and consists at first of a single row of cells. Later it is composed of several rows, some of which are aloral, and at least a single row runs posterior to the mouth. It is not completed dorsally until much later, though soon after the definitive mouth is formed the velum splits on each side of the embryo, and about half way between the ventral and dorsal surfaces, into an anterior and posterior branch; the latter continues up over the dorsal surface just posterior to the large ciliated cells; the anterior branch, which is the chief one, turns forward over the sides of the head vesicle, and quite late in development the two arms of the anterior branch meet and fuse on the mid line just in front of the ventral part of the preoral velum. Thus two large velar lobes are formed, one on each side. The posterior branch of the velum appears to be the postoral ciliated band, the anterior branch the preoral; from the corners of the mouth to the middle of the sides of the embryo the two are fused, while ventrally they are separated by the mouth and dorsally by the whole diameter of the head vesicle. A postoral band of cilia has been described as present in the veligers of several gastropods, and among these *Crepidula*,² but I am not aware that any one has hitherto

¹ F. Blochmann, Ueber die Entwicklung der *Neritina fluviatilis*. Zeit. wiss. Zool. Bd. 36.

² McMurich. J. E. U. Circulars, No. 44, 1885.

found the two separated dorsally. The velum does not become ciliated until quite late in development, though the embryo swims about in the pouch by means of the cilia of the large ciliated cells which form the head vesicle.

The shell gland appears on the dorsal surface immediately posterior to the second or transverse furrow as a prominence of ectoderm cells. In the place of this prominence an invagination afterward appears; the margin of the invagination extends rapidly and a thin cuticle, the first indication of the shell, is secreted by the invaginated cells. As development proceeds the shell becomes asymmetrical, developing more rapidly on the left side than on the right.

The foot arises as a single median protuberance just posterior to the mouth. While it shows no trace of a double origin, it occupies a region along which the blastopore closed, so that really the foot may be considered as having arisen on both sides of the blastopore, though the lips of the latter have fused before the former appears. Running from the mouth backward over the median surface of the foot is a row of large ciliated cells resembling those on the dorsal area.

At the posterior end of the embryo three or four large ciliated anal cells appear, and just ventral to these the distal end of the intestine is pressed against the ectoderm. The proctodan invagination does not occur until late in development. The intestine is a tube with a distinct lumen, its walls being formed of small cells free from yolk. In the course of development its central end, where it opens into the cavity between the yolk spheres is carried anteriorly and to the right. Throughout its whole length the intestine is pressed closely against the ectoderm.

The supra-oesophageal ganglia appear as proliferation of the ectoderm on each side of and dorsal to the mouth; the eyes are formed in connection with these ganglia as involutions of ectoderm. The ganglia of the two sides are connected by a commissure, and from the centre of the latter a nerve runs forward to the centre of the apical plate, where there is a ciliated depression in the ectoderm, which I believe is a sense organ. A commissure connects the supra-oesophageal ganglion of each side with the otocysts. The latter are formed by involution of the ectoderm of the foot, and the pedal ganglion is formed by delamination from the ectoderm at the sides of the foot.

Urosalpinx cinerea.

The breeding habits of *Urosalpinx* have been fully described by Professor Brooks.³ The segmentation is almost identical with that described by Professor Brook⁴ for the oyster, and closely resembles the segmentation of *Nassa*, as described by Bobretzky.⁵ The chief difference between the segmentation in *Urosalpinx* and *Crepidula* consists in the fact that while the four macromeres of *Crepidula* are equal in size, the four macromeres of *Urosalpinx* are very unequal, one being very much larger than the other three. Two furrows appear simultaneously and seem to divide the ovum into one large sphere and two smaller ones. Really, however, one of the smaller spheres is not completely separated from the larger one, and soon after fuses with it. This smaller sphere is merely a constricted portion of the larger sphere and contains the nucleus. Thus it is seen that of the two furrows mentioned, but one is a true cleavage furrow and it divides the egg into a larger and a smaller moiety. At the next stage the smaller moiety divides into two equal parts, and at the same time two protuberances, each containing a nucleus, are pushed out from the larger moiety. One of these

³ Preliminary observations on the development of the Marine Prosobranchs. Studies from the Biological Laboratory, J. H. U., Vol. 1.

⁴ The Embryology of the Oyster. Studies from the Biological Laboratory, J. H. U., Vol. 1.

⁵ N. Bobretzky. Studien über die Embryonale Entwicklung des Gasteropoden. Archiv. f. Mik. Anat., Bd. 13.

III

1890

1891

1892

1893

1894

1895

1896

1897

1898

1899

1900

1901

1902

1903

1904

1905

1906

1907

1908

1909

1910

1911

1912

1913

1914

1915

1916

1917

1918

1919

1920

1921

1922

1923

1924

1925

1926

1927

1928

1929

1930

1931

1932

1933

1934

1935

1936

1937

1938

1939

1940

1941

1942

1943

1944

1945

1946

1947

1948

1949

1950

1951

1952

1953

1954

1955

1956

1957

1958

1959

1960

1961

1962

1963

1964

1965

1966

1967

1968

1969

1970

1971

1972

1973

1974

1975

1976

1977

1978

1979

1980

1981

1982

1983

1984

1985

1986

1987

1988

1989

1990

1991

1992

1993

1994

1995

1996

1997

1998

1999

2000

2001

2002

2003

2004

2005

2006

2007

2008

2009

2010

2011

2012

2013

2014

2015

2016

2017

2018

2019

2020

2021

2022

2023

2024

2025

2026

2027

2028

2029

2030

2031

2032

2033

2034

2035

2036

2037

2038

2039

2040

2041

2042

2043

2044

2045

2046

2047

2048

2049

2050

2051

2052

2053

2054

2055

2056

2057

2058

2059

2060

2061

2062

2063

2064

2065

2066

2067

2068

2069

2070

2071

2072

2073

2074

2075

2076

2077

2078

2079

2080

2081

2082

2083

2084

2085

2086

2087

2088

2089

2090

2091

2092

2093

2094

2095

2096

2097

2098

2099

2100

2101

2102

2103

2104

2105

2106

2107

2108

2109

2110

2111

2112

2113

2114

2115

2116

2117

2118

2119

2120

2121

2122

2123

2124

2125

2126

2127

2128

2129

2130

2131

2132

2133

2134

2135

2136

2137

2138

2139

2140

2141

2142

2143

2144

2145

2146

2147

2148

2149

2150

2151

2152

2153

2154

2155

2156

2157

2158

2159

2160

2161

2162

2163

2164

2165

2166

2167

2168

2169

2170

2171

2172

2173

2174

2175

2176

2177

2178

2179

2180

2181

2182

2183

2184

2185

2186

2187

2188

2189

2190

2191

2192

2193

2194

2195

2196

2197

2198

2199

2200

2201

2202

2203

2204

2205

2206

2207

2208

2209

2210

2211

2212

2213

2214

2215

2216

2217

2218

2219

2220

2221

2222

2223

2224

2225

2226

2227

2228

2229

2230

2231

2232

2233

2234

2235

2236

2237

2238

2239

2240

2241

2242

2243

2244

2245

2246

2247

2248

2249

2250

2251

2252

2253

2254

2255

2256

2257

2258

2259

2260

2261

2262

2263

2264

2265

2266

2267

2268

2269

2270

2271

2272

2273

2274

2275

2276

2277

2278

2279

2280

2281

2282

2283

2284

2285

2286

2287

2288

2289

2290

2291

2292

2293

2294

2295

2296

2297

2298

2299

2300

2301

2302

2303

2304

2305

2306

2307

2308

2309

2310

2311

2312

2313

2314

2315

2316

2317

2318

2319

2320

2321

2322

2323

2324

2325

2326

2327

2328

2329

2330

2331

2332

2333

2334

2335

2336

2337

2338

2339

2340

2341

2342

2343

23

I

1

VZ.

1)
 2)
 3)
 4)
 5)
 6)
 7)
 8)
 9)
 10)
 11)
 12)
 13)
 14)
 15)
 16)
 17)
 18)
 19)
 20)
 21)
 22)
 23)
 24)
 25)
 26)
 27)
 28)
 29)
 30)
 31)
 32)
 33)
 34)
 35)
 36)
 37)
 38)
 39)
 40)
 41)
 42)
 43)
 44)
 45)
 46)
 47)
 48)
 49)
 50)
 51)
 52)
 53)
 54)
 55)
 56)
 57)
 58)
 59)
 60)
 61)
 62)
 63)
 64)
 65)
 66)
 67)
 68)
 69)
 70)
 71)
 72)
 73)
 74)
 75)
 76)
 77)
 78)
 79)
 80)
 81)
 82)
 83)
 84)
 85)
 86)
 87)
 88)
 89)
 90)
 91)
 92)
 93)
 94)
 95)
 96)
 97)
 98)
 99)
 100)

) / /

extended on the 1st of May, and at 10
 o'clock the morning wind had fallen, and
 the 1st of May was a day of 1st
 rain. I had a good view of the
 rain, and the rain was not the same
 but I was not up the egg

I and I went to the 1st of May
 following the 1st of May, and I was
 not. I was not up the egg
 The first of May was at the 1st of May
 1st of May, and I was not up the egg
 The 1st of May was at the 1st of May
 1st of May, and I was not up the egg
I and I went to the 1st of May
 at the 1st of May, and I was not up the egg
 The 1st of May was at the 1st of May
 1st of May, and I was not up the egg

It would be very hard to find a
man who would follow the same path as
the one who has been followed by
the majority of men. The man who
follows the majority is not the man
who is the most useful to the world.
The man who is the most useful to
the world is the man who is the
most different from the majority.
The man who is the most different
from the majority is the man who
is the most useful to the world.
The man who is the most useful to
the world is the man who is the
most different from the majority.
The man who is the most different
from the majority is the man who
is the most useful to the world.
The man who is the most useful to
the world is the man who is the
most different from the majority.
The man who is the most different
from the majority is the man who
is the most useful to the world.

heads of the fish found were numerous large
 eggs. I did not observe any fish and
 a number of dead young fish. The eggs
 were all the size of a pea and found in the
 water and the water was very dark
 and turbid. There was a number
 of small fish and a few crabs and
 the eggs were all the same size. The water
 in the water was dark and the fish
 were all the same size and the water

The Wentworth gave these things
 on 1st October 1891. The eggs
 were found. See after the Wentworth
 were the same size and after the water was
 found the fish were found and the water
 was found. The water was found and the
 water was found. The water was found and
 the water was found. The water was found
 and the water was found. The water was
 found and the water was found. The water
 was found and the water was found. The
 water was found and the water was found.

back to the house and the
 children were all in the
 room and the mother was
 sitting on the bed and the
 father was sitting on the
 floor and the children were
 all around them.

I went back to the house and
 the mother was sitting on the
 bed and the father was sitting
 on the floor and the children
 were all around them. The
 mother was sitting on the bed
 and the father was sitting on
 the floor and the children were
 all around them. The mother
 was sitting on the bed and the
 father was sitting on the floor
 and the children were all
 around them. The mother was
 sitting on the bed and the
 father was sitting on the floor
 and the children were all
 around them. The mother was
 sitting on the bed and the
 father was sitting on the floor
 and the children were all
 around them.

The mother was sitting on the
 bed and the father was sitting
 on the floor and the children
 were all around them.

for me to be able to live the life of the heart
I have said in my life the whole of the
extended period of time to me of the
heart and I have felt that I am
the whole of the

also written for the same end of things
 The first is the most important and the
 second is the least important. The first is
 the most important because it is the most
 important of the two. The second is the
 least important because it is the least
 important of the two. The first is the most
 important because it is the most important
 of the two. The second is the least
 important because it is the least important
 of the two. The first is the most important
 because it is the most important of the two.
 The second is the least important because it
 is the least important of the two. The first
 is the most important because it is the most
 important of the two. The second is the
 least important because it is the least
 important of the two. The first is the most
 important because it is the most important
 of the two. The second is the least important
 because it is the least important of the two.

The first is the most important of the two
 because it is the most important of the two.

as the definitive mouth. With the closure of the
blastopore we may consider the first period of
development accomplished.

The foundation of the country is a
massive granite, the surface of which
is covered with a thin layer of soil
and a few scattered rocks. The soil is
generally poor, but in some places it is
fertile and produces good crops. The
climate is temperate, with a long
summer and a short winter. The
population is small, and the people
are generally poor.

There is at first a small expanse of
water, which is very shallow.

The water is very shallow, and the
bottom is composed of sand and mud.
The water is very shallow, and the
bottom is composed of sand and mud.
The water is very shallow, and the
bottom is composed of sand and mud.
The water is very shallow, and the
bottom is composed of sand and mud.
The water is very shallow, and the
bottom is composed of sand and mud.

works around - what they are in the world
124 The great power - what it is
and how it is used
in the world - what it is
to the world - what it is
125 The great power - what it is
and how it is used
in the world - what it is
to the world - what it is

The world - what it is
and how it is used
in the world - what it is
to the world - what it is
126 The world - what it is
and how it is used
in the world - what it is
to the world - what it is
127 The world - what it is
and how it is used
in the world - what it is
to the world - what it is

The first time I ever saw a man
 who was a doctor & a lawyer
 in the same person. I was
 in the city of New York
 in a building which was
 a combination of a law office
 & a doctor's office.

The man was a tall
 thin man with a long
 face & a long nose.

He was a very
 kind man & a very
 good doctor.

I was very
 much interested in
 him & his work.

He was a very
 good man & a very
 good doctor.

I was very
 much interested in
 him & his work.

Feb 1 1881

at least one hundred years ago
at least one hundred years ago

The first of these is the fact that
the first of these is the fact that

the first of these is the fact that
the first of these is the fact that

the first of these is the fact that
the first of these is the fact that

the first of these is the fact that
the first of these is the fact that

the first of these is the fact that
the first of these is the fact that

the first of these is the fact that
the first of these is the fact that

the first of these is the fact that
the first of these is the fact that

the first of these is the fact that
the first of these is the fact that

the first of these is the fact that
the first of these is the fact that

separated by a deep constriction near the
center of each of the lobes. The latter contracts
at the yolk and at about becomes a constriction,
the heart muscle now being and forth with
into the interior of this constriction, where it
rests permanently.

As the period of constriction passes on, the
muscle now increases in length and the
heart is in position to be pushed into the
constriction.

On the next week of the month, the
heart is the constriction is deepened
and the heart is pushed into the
constriction.

The position of the heart is the next of the
week is the heart is pushed into the
constriction.

The position of the heart is the next of the
week is the heart is pushed into the
constriction.

part of the
... ..
... ..

... ..
... ..
... ..
... ..
... ..
... ..

... ..
... ..
... ..
... ..
... ..
... ..

... ..
... ..
... ..
... ..
... ..

I have very often seen the old men
 to get up with the same old
 and the country there for a long time
 at a new time. I have seen the
 to be the same. I have seen the
 to be the same. I have seen the

and in the same way the
the 1st of the month is
the 2nd of the month is
the 3rd of the month is
the 4th of the month is
the 5th of the month is
the 6th of the month is
the 7th of the month is
the 8th of the month is
the 9th of the month is
the 10th of the month is
the 11th of the month is
the 12th of the month is
the 13th of the month is
the 14th of the month is
the 15th of the month is
the 16th of the month is
the 17th of the month is
the 18th of the month is
the 19th of the month is
the 20th of the month is
the 21st of the month is
the 22nd of the month is
the 23rd of the month is
the 24th of the month is
the 25th of the month is
the 26th of the month is
the 27th of the month is
the 28th of the month is
the 29th of the month is
the 30th of the month is
the 31st of the month is

... ..
... ..
... ..
... ..
... ..

... ..
... ..
... ..
... ..
... ..

... ..
... ..
... ..
... ..
... ..

... ..
... ..
... ..
... ..
... ..

at first, and I only felt a little
uneasy when I saw

the first of the things I had
promised I would do. I was not
at all sure I could do it. I had
seen the things I had to do, and I
had seen the things I had to do.

I had seen the things I had to do, and I
had seen the things I had to do. I
had seen the things I had to do, and I

had seen the things I had to do, and I
had seen the things I had to do. I

had seen the things I had to do, and I
had seen the things I had to do. I

had seen the things I had to do, and I
had seen the things I had to do. I

had seen the things I had to do, and I
had seen the things I had to do. I

 
 t .

I have been thinking of you
 and of the things you have
 done for me. I am sure
 that you will be very
 kind to me.

I am sure that you will
 be very kind to me. I am
 sure that you will be very
 kind to me. I am sure that
 you will be very kind to me.

I am sure that you will
 be very kind to me. I am
 sure that you will be very
 kind to me. I am sure that
 you will be very kind to me.

I am sure that you will
 be very kind to me. I am
 sure that you will be very
 kind to me. I am sure that
 you will be very kind to me.

I am sure that you will
 be very kind to me. I am
 sure that you will be very
 kind to me. I am sure that
 you will be very kind to me.

I am sure that you will
 be very kind to me. I am
 sure that you will be very
 kind to me. I am sure that
 you will be very kind to me.

I am sure that you will
 be very kind to me. I am
 sure that you will be very
 kind to me. I am sure that
 you will be very kind to me.

forest surface with a few more but the
 living layer is very small. The forest has
 fallen in some places. It has a
 dead layer which is about 10 feet high
 and is made of dead leaves and branches.
 The forest is very old and the trees are
 very large. The forest is very old and the
 trees are very large. The forest is very old
 and the trees are very large. The forest is
 very old and the trees are very large. The
 forest is very old and the trees are very
 large. The forest is very old and the trees
 are very large. The forest is very old and
 the trees are very large. The forest is very
 old and the trees are very large. The forest
 is very old and the trees are very large.

John moved to the southern end.

V. *Salix herbacea* L.

15. Patten, W. D. ...
...
16. Patten, W. D. ...
...
17. Patten, W. D. ...
...
18. Patten, W. D. ...
...
19. Patten, W. D. ...
...
20. Patten, W. D. ...
...
21. Patten, W. D. ...
...

27. 18. 13. 14. 15. 16. 17. 18. 19. 20. 21. 22. 23. 24. 25. 26. 27. 28. 29. 30. 31. 32. 33. 34. 35. 36. 37. 38. 39. 40. 41. 42. 43. 44. 45. 46. 47. 48. 49. 50. 51. 52. 53. 54. 55. 56. 57. 58. 59. 60. 61. 62. 63. 64. 65. 66. 67. 68. 69. 70. 71. 72. 73. 74. 75. 76. 77. 78. 79. 80. 81. 82. 83. 84. 85. 86. 87. 88. 89. 90. 91. 92. 93. 94. 95. 96. 97. 98. 99. 100.

VI. Explanation of Figures

All plates of figures are of Cephalopoda in a to and are shown with the same scale as of the plates of the same

Table of the

N. D. C. R. The first figure is a cephalopod in a to and is shown with the same scale as of the plates of the same

N. D. C. R. The second figure is a cephalopod in a to and is shown with the same scale as of the plates of the same

N. D. C. R. The third figure is a cephalopod in a to and is shown with the same scale as of the plates of the same

N. D. C. R. The fourth figure is a cephalopod in a to and is shown with the same scale as of the plates of the same

1. The first figure is a cephalopod in a to and is shown with the same scale as of the plates of the same

2. The second figure is a cephalopod in a to and is shown with the same scale as of the plates of the same

3. The third figure is a cephalopod in a to and is shown with the same scale as of the plates of the same

4. The fourth figure is a cephalopod in a to and is shown with the same scale as of the plates of the same

5. The fifth figure is a cephalopod in a to and is shown with the same scale as of the plates of the same

- 10 - 1st set of measurements given by the ... 10
- 20 - 1st set of measurements given by the ... 10
- 30 - 1st set of measurements given by the ... 10
- 40 - 1st set of measurements given by the ... 10
- 50 - 1st set of measurements given by the ... 10
- 60 - 1st set of measurements given by the ... 10
- 70 - 1st set of measurements given by the ... 10
- 80 - 1st set of measurements given by the ... 10
- 90 - 1st set of measurements given by the ... 10
- 100 - 1st set of measurements given by the ... 10

Figures

- Fig. 1 - Diagram of the ...
- Fig. 2 - Diagram of the ...
- Fig. 3 - Diagram of the ...
- Fig. 4 - Diagram of the ...
- Fig. 5 - Diagram of the ...
- Fig. 6 - Diagram of the ...
- Fig. 7 - Diagram of the ...
- Fig. 8 - Diagram of the ...
- Fig. 9 - Diagram of the ...
- Fig. 10 - Diagram of the ...

1861. 1. 2. 3. 4. 5. 6. 7. 8. 9. 10. 11. 12.

- July 21. After a long and successful season in
the field, I have returned to the
city of New York. C
- July 22. After a long and successful season in
the field, I have returned to the
city of New York. C
- July 23. After a long and successful season in
the field, I have returned to the
city of New York. C
- July 24. After a long and successful season in
the field, I have returned to the
city of New York. C
- July 25. After a long and successful season in
the field, I have returned to the
city of New York. C
- July 26. After a long and successful season in
the field, I have returned to the
city of New York. C
- July 27. After a long and successful season in
the field, I have returned to the
city of New York. C
- July 28. After a long and successful season in
the field, I have returned to the
city of New York. C
- July 29. After a long and successful season in
the field, I have returned to the
city of New York. C
- July 30. After a long and successful season in
the field, I have returned to the
city of New York. C

July 2. Dried at 100° for 24 hours, in a desiccator containing
 sulphuric acid. The residue is a
 white solid.

July 3. The residue is a white solid, melting at 100°.

July 4. The residue is a white solid, melting at 100°.

July 5. The residue is a white solid, melting at 100°.

July 6. The residue is a white solid, melting at 100°.

July 7. The residue is a white solid, melting at 100°.

July 8. The residue is a white solid, melting at 100°.

at the 6' distance from the ground the center
 of the circle is at the center of the circle
 of the circle of the circle of the circle

Figure 1. The circle of the circle of the circle
 of the circle of the circle of the circle
 of the circle of the circle of the circle

Figure 2. The circle of the circle of the circle
 of the circle of the circle of the circle
 of the circle of the circle of the circle

Figure 3. The circle of the circle of the circle
 of the circle of the circle of the circle
 of the circle of the circle of the circle

Figure 4. The circle of the circle of the circle
 of the circle of the circle of the circle
 of the circle of the circle of the circle

Figure 5. The circle of the circle of the circle
 of the circle of the circle of the circle
 of the circle of the circle of the circle

Figure 6. The circle of the circle of the circle
 of the circle of the circle of the circle
 of the circle of the circle of the circle

It is a very common mistake to suppose
that the only way to get a good
result is to use a large quantity of
the material.

Page 1. I will now discuss the question of the
best way to get a good result. It is
very important to know the
doubtful point in the matter.
The first point is the quality of the
material.

Page 2. The second point is the quantity of the
material. It is very important to know
the best way to get a good result. It is
very important to know the
doubtful point in the matter.
The first point is the quality of the
material.

Page 3. The third point is the time of the
operation. It is very important to know
the best way to get a good result. It is
very important to know the
doubtful point in the matter.
The first point is the quality of the
material.

Page 4. The fourth point is the cost of the
operation. It is very important to know
the best way to get a good result. It is
very important to know the
doubtful point in the matter.
The first point is the quality of the
material.

first portion to be a continuation of the
2nd, and the 3rd a continuation of the 1st
in general. The 1st is a continuation of the
2nd, and the 3rd a continuation of the 1st
in general. The 1st is a continuation of the
2nd, and the 3rd a continuation of the 1st
in general.

Fig 52.

The 1st is a continuation of the 2nd
in general. The 2nd is a continuation of the 1st
in general. The 3rd is a continuation of the 1st
in general. The 4th is a continuation of the 1st
in general. The 5th is a continuation of the 1st
in general.

Fig 53.

The 1st is a continuation of the 2nd
in general. The 2nd is a continuation of the 1st
in general. The 3rd is a continuation of the 1st
in general. The 4th is a continuation of the 1st
in general. The 5th is a continuation of the 1st
in general. The 6th is a continuation of the 1st
in general.

Fig 54.

The 1st is a continuation of the 2nd
in general. The 2nd is a continuation of the 1st
in general. The 3rd is a continuation of the 1st
in general. The 4th is a continuation of the 1st
in general. The 5th is a continuation of the 1st
in general.

Aug 11 - Received from the ...
...

Aug 12 - ...
...

Aug 13 - ...
...

Aug 14 - ...
...

Aug 15 - ...
...

Aug 16 - ...
...

Aug 17 - ...
...

the first of the year
the first of the year

the first of the year
the first of the year

the first of the year
the first of the year

the first of the year
the first of the year

the first of the year
the first of the year

the first of the year
the first of the year

the first of the year
the first of the year