QK
431
M93

UC-NRLF

D 2 518 500

SECOND SYSTEMATIC CENSUS
OF
AUSTRALIAN PLANTS,

WITH

Chronologic, Literary and Geographic Annotations;

BY

BARON FERDINAND VON MUELLER,

K.C.M.G., M.D., Ph.D., F.R.S.

F.L.S., F.G.S., F.C.S., F.R.G.S., CM.Z.S., H.F.R.H.S., &c.

GOVERNMENT BOTANIST FOR THE COLONY OF VICTORIA.

PART I.—VASCULARES.

"LEVATE OCULOS IN EXCELSUM ET VIDETE QUI CREAVIT EA OMNIA. QUI NUNQUIT MORTUO INVENIENS EAM. —Isaiah xl

MELBOURNE:

PRINTED FOR THE VICTORIAN GOVERNMENT

By McCARRON, BIRD & CO., 479 COLLINS STREET.

1889.

THE LIBRARY
OF
THE UNIVERSITY
OF CALIFORNIA
DAVIS

Genoss J. Dörfler & v.
zur freundlichen Erinnerung
und zur gelegentlichen Gebrauch
von Ferd. von Mueller.

Handwritten text, likely a signature or name, is visible at the top of the page. The text is written in cursive and appears to be "James M. Smith".

SECOND SYSTEMATIC CENSUS
OF
AUSTRALIAN PLANTS,

WITH
Chronologic, Literary and Geographic Annotations;

BY
BARON FERDINAND VON MUELLER

K.C.M.G., M.D., Ph.D., F.R.S.

F.L.S., F.G.S., F.C.S., F.R.G.S., C.M.Z.S., H.F.R.H.S., &c.

GOVERNMENT BOTANIST FOR THE COLONY OF VICTORIA.

PART I.—VASCULARES.

LEVATE OCULOS IN EXCELSUM ET VIDETE QUIS CREAVIT HAEC OMNIA, QUIS EDUCIT MULTITUDINEM EORUM."—*Isaiah* xl.

MELBOURNE:
PRINTED FOR THE VICTORIAN GOVERNMENT
By McCARRON, BIRD & Co., 479 COLLINS STREET.

1889.

LIBRARY
UNIVERSITY OF CALIFORNIA

Digitized by the Internet Archive
in 2007 with funding from
Microsoft Corporation

<http://www.archive.org/details/secondsystematic00muelrich>

To

SIR JOSEPH HOOKER, K.C.S.I., C.B., F.R.S., V.P.L.S.

AND TO

M. ALPHONSE DE CANDOLLE, F.M.R.S., F.M.L.S.,

WHO AS HEIRS OF GREAT NAMES WORTHILY SUSTAIN WORLD-WIDE ANCESTRAL FAME,

AND WHO AS LEADERS IN PHYTOGRAPHY DURING HALF A CENTURY

WILL BE PRE-EMINENT IN BIOMORPHIC SCIENCE THROUGH ALL AGES,

THESE PAGES

ARE OFFERED AS AN APPRECIATIVE TRIBUTE

FROM A YOUNG COLONY IN ANTIPODAL REMOTENESS.

DICOTYLEDONEAE.

Ray, Method. Plant. Nova 2 (1682).

CHORIPETALEAE HYPOGYNAE.

F. v. MUELLER, native plants of Victoria I, 1 (1879).

DILLENIACEAE.

SALISBURY, Parad. Lond. I, 73 (1801).—Genera (here) 4, Species 95.

WORMIA; Rottboell, Nye Saml. Vid. Selsk. Skrivt. II, 532 (1783). (Dilleniaceae subgenus).

W. alata; R. Brown in De Cand. syst. veg. I, 434 (1818) ...	...	...	...	...	...	...	...	...	...	Q.	—	B.fl.I,16	M.fr.VIII,124.	...	P.
HIBBERTIA; Andrews, Bot. Reposit. t. 126 (1800). (Candollea, Pleurandra, Hemistemma, Adrastaea, Ochrolasia, Hemistephus, Huttia, Warburtonia.)															
H. Banksii; F. v. Mueller, pl. of Vict. I, 14, implied (1860) ...	...	...	...	...	...	...	...	...	...	Q.	—	B.fl.I,21	M.fr.II,1; XI,92.		
H. Holtzei; F. v. M. in Wing's S. Sc. Rec. new ser. I, Nov. (1885) ...	...	...	...	...	...	...	...	...	...	—	N.A.	—			
H. Brownii; Benth. Fl. Austr. I, 21 (1863) ...	...	...	...	...	...	...	...	...	...	Q.	—	B.fl.I,21	M.fr.II,1.		
H. dealbata; F. v. M., pl. of Vict. I, 14, implied (1860) ...	...	...	...	...	...	...	...	...	...	—	N.A.	B.fl.I,21	M.fr.II,1.		
H. candicans; Benth. Fl. Austr. I, 21 (1863) ...	...	...	...	...	...	...	...	...	...	Q.	N.A.	B.fl.I,21			
H. Benthami; F. v. M., syst. Cens. of Austr. pl. 1 (1882)..	...	...	...	...	...	...	...	...	...	—	N.A.	B.fl.I,21	M.fr.II,1.		
H. Muellieri; Benth. Fl. Austr. I, 21 (1863) ...	...	...	...	...	...	...	...	...	...	—	N.A.	B.fl.I,21			
H. ledifolia; Benth. Fl. Austr. I, 22 (1863) ...	...	...	...	...	...	...	...	...	...	—	N.A.	B.fl.I,22			
H. verrucosa; F. v. M., pl. of Vict. I, 14, implied (1860) ...	...	...	...	...	...	...	...	...	...	—	—	B.fl.I,22	M.fr.XI,92.		
H. spicata; F. v. M., fragm. phytogr. Austral. II, 1 (1860) ...	...	...	...	...	...	...	...	...	...	—	—	B.fl.I,22	M.fr.XI,20,92.		
H. polystachya; Benth. Fl. Austr. I, 22 (1863) ...	...	...	...	...	...	...	...	...	...	—	—	B.fl.I,22	M.fr.XI,92.		
H. rhadinopoda; F. v. M., fragm. XI, 91 (1880) ...	...	...	...	...	...	...	...	...	...	—	—	—	M.fr.XI,91.		
H. furfuracea; Benth. Fl. Austr. I, 23 (1863) ...	...	...	...	...	...	...	...	...	...	—	—	B.fl.I,23	M.fr.I,161; XI,92.		
H. hypericoides; F. v. M., pl. of Vict. I, 13, implied (1860) ...	...	...	...	...	...	...	...	...	...	—	—	B.fl.I,23			
H. microphylla; Steudel in Lehmann, pl. Preiss. I, 273 (1844) ...	...	...	...	...	...	...	...	...	...	—	—	B.fl.I,23	M.fr.I,217.		
H. recurvifolia; Benth. Fl. Austr. I, 24 (1863) ...	...	...	...	...	...	...	...	...	...	—	—	B.fl.I,24			
H. lineata; Steudel in Lehmann, pl. Preiss. I, 272 (1844) ...	...	...	...	...	...	...	...	...	...	—	—	B.fl.I,24	M.fr.XI,92.		
H. acerosa; Benth. Fl. Austr. I, 74 (1863) ...	...	...	...	...	...	...	...	...	...	—	—	B.fl.I,24	M.fr.XI,92.		
H. aurea; Steudel in Lehmann, pl. Preiss. I, 272 (1844) ..	...	...	...	...	...	...	...	...	...	—	—	B.fl.I,25	M.fr.XI,92.		
H. crassifolia; Benth. Fl. Austr. I, 25 (1863) ...	...	...	...	...	...	...	...	...	...	—	—	B.fl.I,25			
H. nitida; F. v. M., pl. of Vict. I, 15 (1860) ...	...	...	...	...	...	...	...	...	...	N.S.W.	—	B.fl.I,25			
H. bracteata; Benth. Fl. Austr. I, 26 (1863) ...	...	...	...	...	...	...	...	...	...	N.S.W.	—	B.fl.I,25			
H. densiflora; F. v. M., pl. of Vict. I, 15 (1860) ...	...	...	...	...	...	...	...	...	...	S.A. T.	V.	N.S.W.	M.fr.VII,125; XI,92.		
H. hirsuta; Benth. Fl. Austr. I, 21 (1863) ...	...	...	...	...	...	...	...	...	...	S.A. T.	—	—	B.fl.I,26		
H. synandra; F. v. M., fragm. IV, 151 (1864) ...	...	...	...	...	...	...	...	...	...	—	—	Q.	M.fr.IV,151.		
H. stricta; R. Brown in De Candolle, syst. veg. I, 422 (1818) ...	...	...	...	...	...	...	...	...	...	W.A.	S.A. T.	V.	N.S.W.	Q.	B.fl.I,27
H. humifusa; F. v. M., pl. of Vict. I, 16, t. Suppl. I (1860) ...	...	...	...	...	...	...	...	...	...	—	—	V.	—	—	B.fl.I,27
H. Billardieri; F. v. M., pl. of Vict. I, 14 (1860) ...	...	...	...	...	...	...	...	...	...	—	S.A. T.	V.	N.S.W.	Q.	M.fr.IV,116.
H. gracilipes; Benth. Fl. Austr. I, 28 (1863) ...	...	...	...	...	...	...	...	...	...	W.A.	—	—	—	—	B.fl.I,28
H. acicularis; F. v. M., pl. of Vict. I, 17 (1860) ...	...	...	...	...	...	...	...	...	...	—	S.A. T.	V.	N.S.W.	Q.	M.fr.XI,92.
H. mucronata; F. v. M., pl. of Vict. I, 17 (1860) ...	...	...	...	...	...	...	...	...	...	W.A.	—	—	—	—	B.fl.I,29

H. hermannifolia ; De Candolle, syst. veg. I, 431 (1818) ...	...	...	...	...	...	N.S.W.	...	B.f.I,30	
H. velutina ; R. Brown in Bentham, Fl. Austr. I, 30 (1863) ...	...	...	...	...	...	Q.	...	B.f.I,30	
H. oblongata ; R. Brown in De Candolle, syst. veg. I, 431 (1818)...	...	...	...	...	...	N.A.	B.f.I,30		M.fr.XI,94.
H. melhanoides ; F. v. M., fragm. IV, 116 (1864) ...	...	...	...	...	...	Q.	...	...	M.fr.IV,116.
H. tomentosa ; R. Brown in De Candolle, syst. veg. I, 432 (1818)...	...	...	...	...	...	N.A.	B.f.I,30		
H. cistifolia ; R. Brown in De Candolle, syst. veg. I, 431 (1818) ...	...	...	...	...	...	N.A.	B.f.I,30		
H. echiifolia ; R. Brown in Bentham, Fl. Austr. I, 31 (1863) ...	...	...	...	...	...	N.A.	B.f.I,31		
H. scabra ; R. Brown in Bentham, Fl. Austr. I, 31 (1863)...	...	...	...	...	...	N.A.	B.f.I,31		
H. lepidota ; R. Brown in De Candolle, syst. veg. I, 432 (1818) ...	...	...	...	...	...	Q.	N.A.	B.f.I,31	
H. vestita ; Cunningham in Bentham, Fl. Austr. I, 31 (1863) ...	...	...	...	...	...	N.S.W.	Q.	B.f.I,31	
H. serpyllifolia ; R. Brown in De Cand. syst. veg. I, 430 (1818) ...	...	...	...	...	T.	V.	N.S.W.	B.f.I,32	
H. pedunculata ; R. Brown in De Cand. syst. veg. I, 430 (1818) ...	...	...	...	...	...	V.	N.S.W.	B.f.I,32	
H. ochrolasia ; Bentham, Fl. Austr. I, 32 (1863) ...	...	...	...	...	W.A.	...	...	B.f.I,32	M.fr.VII,125 ; XI,93.
H. angustifolia ; Salisbury, parad. Lond. 73 (1807) ...	...	...	...	...	...	T.	V.	B.f.I,33	
H. fasciculata ; R. Brown in De Candolle, syst. veg. I, 428 (1818) ...	...	...	...	...	S.A.	T.	V.	N.S.W.	Q.
H. virgata ; R. Brown in De Candolle, syst. veg. I, 428 (1818) ...	...	...	...	...	...	S.A.	T.	V.	N.S.W.
H. inclusa ; Bentham, Fl. Austr. I, 34 (1863) ...	...	...	...	...	W.A.	...	...	B.f.I,3	
H. rostellata ; Turczaninow in Bull. Soc. Mosc. XXII, part III., 8 (1849)	...	...	...	...	W.A.	...	...	B.f.I,35	M.fr.XI,93.
H. glomerata ; Bentham, Fl. Austr. I, 34 (1863) ...	...	...	...	...	W.A.	...	...	B.f.I,35	
H. argentea ; Steudel in Lehmann, pl. Preiss. I, 268 (1844) ...	...	...	...	...	W.A.	...	...	B.f.I,35	
H. pilosa ; Steudel in Lehmann, pl. Preiss. I, 272 (1844) ...	...	...	...	...	W.A.	...	...	B.f.I,35	
H. montana ; Steudel in Lehmann, pl. Preiss. I, 270 (1844) ...	...	...	...	...	W.A.	...	...	B.f.I,35	M.fr.XI,93.
H. obtusifolia ; De Candolle, syst. veg. I, 428 (1818) ...	...	...	...	...	...	...	V.	N.S.W.	Q.
H. diffusa ; R. Brown in De Candolle, syst. veg. I, 424 (1818) ...	...	...	...	...	...	...	V.	N.S.W.	Q.
H. saligna ; R. Brown in De Candolle, syst. veg. I, 427 (1818) ...	...	...	...	...	...	...	N.S.W.	B.f.I,37	M.fr.XI,94.
H. volubilis ; Andrews in Bot. Reposit. t. 126 (1800) ...	...	...	...	...	...	...	N.S.W.	Q.	B.f.I,37
H. grossularifolia ; Salisbury, parad. Lond. t. 73 (1806) ...	...	...	...	...	W.A.	...	...	B.f.I,37	M.fr.IV,116 ; VII,125 ; XI,94.
H. dentata ; R. Brown in De Candolle, syst. veg. I, 426 (1818) ...	...	...	...	...	...	...	V.	N.S.W.	Q.
H. Cunninghamii ; Aiton in Hook. Bot. Mag. t. 318 (1832)...	...	...	...	...	W.A.	...	...	B.f.I,39	M.fr.XI,93.
H. perfoliata ; Huegel, enum. pl. Nov. Holl. austr. occ. 3 (1837) ...	...	...	...	...	W.A.	...	...	B.f.I,38	M.fr.XI,93.
H. bracteosa ; Turczaninow, in Bull. Soc. Mosc. XXV, 140 (1852)	...	...	...	...	W.A.	...	...	B.f.I,38	M.fr.XI,94.
H. longifolia ; F. v. M., fragm. IV, 115 (1814) ...	...	...	...	...	...	...	Q.	...	M.fr.IV,115.
H. oenotheroides ; F. v. M., fragm. VII, 37 (1870) ...	...	...	...	...	...	...	Q.	...	M.fr.VII,37.
H. glaberrima ; F. v. M., fragm. III, 1 (1862) ...	...	...	...	...	...	S.A.	...	Q.	B.f.I,39
H. Milnei ; Bentham, Fl. Austr. I, 39 (1863) ...	...	...	...	...	W.A.	...	...	B.f.I,39	
H. lasiopus ; Bentham, Fl. Austr. I, 40 (1863) ...	...	...	...	...	W.A.	...	...	B.f.I,40	M.fr.XI,94.
H. potentilliflora ; F. v. M. in Bentham, Fl. Austr. I, 40 (1863) ...	...	...	...	...	W.A.	...	...	B.f.I,40	M.fr.XI,94.
H. pungens ; Bentham, Fl. Austr. I, 40 (1863) ...	...	...	...	...	W.A.	...	...	B.f.I,40	
H. nutans ; Bentham, Fl. Austr. I, 40 (1863)...	...	...	...	...	W.A.	...	...	B.f.I,40	
H. leptopus ; Bentham, Fl. Austr. I, 41 (1863) ...	...	...	...	...	W.A.	...	...	B.f.I,41	
H. stellaris ; Endlicher in Hueg. enum. 3 (1837) ...	...	...	...	...	W.A.	...	...	B.f.I,41	M.fr.XI,93.
H. glomerosa ; F. v. M., syst. Cens. of Austr. plants 2 (1832) ...	...	...	...	...	W.A.	...	...	B.f.I,43	M.fr.XI,94.
H. obcuneata ; Salisbury, parad. Londin. t. 73 (1807) ...	...	...	...	...	W.A.	...	...	B.f.I,42	M.fr.XI,94.
H. teretifolia ; F. v. M., fragm. IV, 117 (1864) ...	...	...	...	...	W.A.	...	...	B.f.I,43	M.fr.XI,95.
H. desmophylla ; F. v. M., fragm. XI, 95 (1880) ...	...	...	...	...	W.A.	...	...	B.f.I,43	M.fr.XI,95.
H. helianthemoides ; F. v. M., syst. Cens. of Austr. plants 2 (1882)	...	...	...	...	W.A.	...	...	B.f.I,43	
H. depressa ; Steudel in Lehm. pl. Preiss. I, 268 (1844) ...	...	...	...	...	W.A.	...	...	B.f.I,44	M.fr.XI,95.
H. Huegelii ; F. v. M., fragm. XI, 95 (1880) ...	...	...	...	...	W.A.	...	...	B.f.I,44	M.fr.XI,95.
H. polygonoides ; F. v. M., fragm. IV, 116 (1864) ...	...	...	...	...	W.A.	...	...	B.f.I,45	M.fr.XI,95.

H. vaginata; F. v. M., fragm. XI, 96 (1880)	W.A.	—	—	—	—	—	—	B.fl.I,45	M.fr.XI,96.
H. subexcisa; Steudel in Lehmann, pl. Preiss. I, 269 (1844)	W.A.	—	—	—	—	—	—	B.fl.I,46	M.fr.XI,95.
H. squamosa; Turczaninow in Bull. Soc. Mosc. XXII, part III, 9 (1844)	W.A.	—	—	—	—	—	—	B.fl.I,46	
H. uncinata; F. v. M., syst. Cens. of Austr. pl. 2 (1882)	W.A.	—	—	—	—	—	—	B.fl.I,46	
H. salicifolia; F. v. M., fragm. I, 161 (1859)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,46	M.fr.VII,124; XI,94.
H. Huttii; F. v. M., fragm. VII, 123 (1870)	W.A.	—	—	—	—	—	—	B.fl.I,47	M.fr.XI,20.
H. Goyderi; F. v. M., fragm. VII, 123 (1870)	—	—	—	—	—	—	N.A.	—	M.fr.VII,123.
TETRACERA; Linné, gen. pl. 345 (1737).									
T. Nordtiana; F. v. M., fragm. V, 1 (1865)	—	—	—	—	—	Q.	—	—	M.fr.V,1 .. P.
T. Daemeliana; F. v. M., fragm. V, 191 (1866)	—	—	—	—	—	Q.	—	—	M.fr.V,191.
T. Wuthiana; F. v. M., fragm. X, 49 (1876)	—	—	—	—	—	Q.	—	—	M.fr.X,49.
PACHYNEMA; R. Brown in De Candolle, syst. veg. I, 412 (1818).									
P. junceum; Bentham, Fl. Austr. I, 47 (1863)	—	—	—	—	—	—	N.A.	B.fl.I,47	M.fr.X,124.
P. complanatum; R. Brown in De Candolle, syst. veg. I, 412 (1818)	—	—	—	—	—	—	N.A.	B.fl.I,48	M.fr.X,124.
P. dilatatum; Bentham, Fl. Austr. I, 48 (1863)	—	—	—	—	—	—	N.A.	B.fl.I,48	M.fr.X,124.
P. sphendrum; F. v. M. & Tate in Trans. R. S. of S.A. V, 79 (1892)	—	—	—	—	—	—	N.A.	—	

RANUNCULACEAE.

A. L. DE JUSSIEU, Rec. de l'Acad. des sc. (1773) from B. de Jussieu (1759).—Genera 5, Species 17.

CLEMATIS; Linné, gen. pl. 163 (1737) from l'Ecluse (1576).

C. aristata; R. Brown in De Candolle, syst. veg. I, 142 (1818)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,6	M.fr.X,2.
C. pubescens; Huegel, enum. pl. Nov. Holl. austr. occ. 1 (1837)	W.A.	—	—	—	—	—	—	B.fl.I,6	
C. glycinoides; De Candolle, syst. veg. I, 145 (1818)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.I,7	M.fr.IX,76; X,2. P.
C. microphylla; De Candolle, syst. veg. I, 142, (1818)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,7	M.fr.X,2; XI,20,27
C. Fawcetti; F. v. M., fragm. X, 1 (1876)	—	—	—	—	N.S.W.	Q.	—	—	M. fr. X, 1.

ANEMONE; Tournefort, inst. rei herb. 275, t. 174 (1700) from Hippocrates, Theophrastus and Dioscorides.

A. crassifolia; Hooker, icones pl. t. 257 (1840)	—	—	T.	—	—	—	—	B.fl.I,8	M.fr.X,1.
---	---	---	----	---	---	---	---	----------	-----------

MYOSURUS; Dillenius, catal. pl. Giess. 106, t. 4 (1749) from L'Obel (1576). (Myosuros.)

M. minimus; Linné, sp. pl. 284 (1753)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.I,8	M.fr.X,2; XI,27 E., As., Afr., Am.
--	---	------	---	----	--------	----	---	----------	------------------------------------

RANUNCULUS; Tournefort, inst. 285, t. 149 (1700) from Bock (1552) from Plinius.

R. parviflorus; Linné, sp. pl. sec. ed. 780 (1762)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,14	M.fr.X,2; XI,27 E., As., Afr., N.Z.
R. hirtus; Banks & Solander in G. Forst. fl. ins. austr. prodr. 90 (1786)	W.A.	—	—	V.	N.S.W.	Q.	—	B.fl.I,13	M.fr.X,2 ... N.Z.
R. rivularis; Banks & Solander in G. Forst. fl. ins. austr. prodr. 90 (1786)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,13	M.fr.X,2 ... N.Z.
R. Muelleri; Bentham, Fl. Austr. I, 13 (1863)	—	—	—	V.	N.S.W.	—	—	B.fl.I,13	
R. lappaceus; Smith in Rees' Cyclopædia, XXIX (1815)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,12	M.fr.X,2 ... N.Z.
R. Gunnianus; Hooker, Jour. of Bot. I, 245, t. 233 (1834)	—	—	T.	V.	N.S.W.	—	—	B.fl.I,11	
R. anemoneus; F. v. M. in Trans. Phil. Soc. Vict. I, 97 (1855)	—	—	—	V.	N.S.W.	—	—	B.fl.I,11	
R. Millani; F. v. M. in Hook. Kew misc. VII, 358 (1855)	—	—	—	V.	N.S.W.	—	—	B.fl.I,10	M.fr.X,2.
R. acquatilis; Dodonæus, stirp. hist. pempt. 387 (1583)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.I,10	 E., As., Afr., Am.

CALTHA; Ruppins, flora Jenensis 119 (1718) from Bock (1552). (Trollius.)

C. ntroloba; F. v. M. in Trans. Phil. Soc. Vic. I, 98 (1855)	—	—	T.	V.	N.S.W.	—	—	B.fl.I,15	M.fr.X,2; XI,136 N.Z.
---	---	---	----	----	--------	---	---	-----------	-----------------------

CERATOPHYLLEAE.

J. E. GRAY; Arrangm. of Brit. pl. II, 395 (1821).—Genus 1, Species 1.

CERATOPHYLLUM, Linné, syst. nat. 9 (1735).

C. demersum; Linné, spec. plant. 992 (1753)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.II,491	M.fr.VIII,103 E., As., Afr., Am., P.
--	---	------	---	----	--------	----	------	-------------	--------------------------------------

NYMPHAEACEAE.

SALISBURY in Koenig & Sims' Ann. of Bot. II, 70 (1805).—Genera 3, Species 5.

CABOMBA ; Aublet, Hist. pl. Guian. I, 321, t. 124 (1775). (Brasenia, Hydropeltis.)

C. peltata ; F. v. M., Introd. to Bot. Teachings, 89 (1877) ... — — — V. N.S.W. Q — B.fl I,60 M.fr.X,77 ... As., Am.

NYMPHAEA ; Tournefort, inst. 260, t. 137 8 (1700) from Theophrastos and Dioscorides.

N. tetragona ; Georgi, Reise im Russ. Reich. I, 220 (1775) ... — — — — Q. — — ... As.

N. stellata ; Willdenow, spec. pl. II, 1153 (1799) ... — — — — Q. N.A. — M.fr.II,142 ; X,77 As., Afr.

N. gigantea, Hooker, Bot. Mag. t. 4647 (1852) ... — — — — N.S.W. Q. N.A. B.fl.I,61 M.fr.II,141 ; X,78 P.

NELUMBO ; Tournefort, inst. 261 (1700). (Nelumbium.)

N. nucifera ; Gaertner, de fruct. I, 73 (1788)... — — — — — Q. N.A. B.fl.I,62 M.fr.X,77 ... E., As., Afr.

PIPERACEAE.

L. CL. RICHARD in Humb. Bonpl. & Kunth, Nov. gen. I, 46 (1815).—Genera 2, Species 10.

PIPER ; Linné, gen. pl. 333 (1737), from C. Bauhin (1623). (Macropiper.)

P. subpeltatum ; Willdenow, spec. plant. V. 166 (1810) ... — — — — Q. — B.fl.VI,204 M.fr.V,198 ... As., Afr., Am., P.

P. excelsum ; G. Forster, florul. ins. Austr. prodr. 5 (1786) ... — — — — N.S.W. — B.fl.VI,204 M.fr.IX,77 ... N.Z., P.

P. Novae Hollandiae ; Miquel in Medd. Akad. Vetensk. Amsterd. sec. ser. II, 5 (1866) ... — — — — N.S.W. Q. — B.fl.VI,204 M.fr.V,198.

P. Banksii ; Miquel in Medd. Akad. Amst. sec. ser. II, 9 (1866) ... — — — — Q. — B.fl.VI,205

P. triandrum ; F. v. M., fragm. V, 197 (1866) ... — — — — Q. — B.fl.VI,205 M.fr.V,197.

P. hederaceum ; Cunningham in De Cand. prodr. XVI, 366 (1869) ... — — — — N.S.W. — B.fl.VI,205

PEPEROMIA ; Ruiz & Pavon, fl. Peru. & Chil. prodr. 8, t. 2 (1794).

P. leptostachya ; Hooker & Arnott, Bot. Beech. 96 (1841)... — — — — N.S.W. Q. — B.fl.VI,206 M.fr.V,198; IX,76 P.

P. reflexa ; A. Dietrich, spec. plant. I, 180 (1831) ... — — — — N.S.W. Q. — B.fl.VI,207 M.fr.V,198; IX,77 As., Afr., Am., P.

P. Urvilleana ; Ach. Richard in d'Urville, voy. Bot. 356 (1832) ... — — — — N.S.W. — M.fr.IX.76 ... N.Z.

P. Baneriana ; Miquel, syst. piperac. 120 (1843) ... — — — — N.S.W. — —

MAGNOLIACEAE.

J. DE ST. HILAIRE, Expos. Fam. nat. II, 74, t. 83-84 (1805).—Genus 1, Species 4.

DRIMYS ; R. & G. Forster, char. gen. 83, t. 42 (1776). (Tasmannia.)

D. aromatica ; F. v. M., pl. of Vict. I, 20 (1860) ... — — T. V. N.S.W. — B.fl.I,49 M.fr.VII,18.

D. membranacea ; F. v. M., fragm. V, 175 (1866) ... — — — — Q. — M.fr.VII,18.

D. dipetala ; F. v. M., pl. of Vict. I, 21 (1860) ... — — — — N.S.W. Q. — B.fl.I,49 M.fr.VII,81.

D. Howeana ; F. v. M., fragm. VII, 17 (1870) ... — — — — N.S.W. — M.fr.IX,76.

ANONACEAE.

A. L. DE JUSSIEU, Gen. 283 (1789) from B. de Jussieu (1759).—Genera 11, Species 18.

UVARIA ; Linné, Amoen. acad. 404 (1747).

U. membranacea ; Bentham, Fl. Austr. I, 51 (1863) ... — — — — Q. — B.fl.I,51

U. Goezeana ; F. v. M., fragm. VII, 125 (1871) ... — — — — Q. — M.fr.VII,125.

U. Holtzei ; F. v. M. in Wing's S. Sc. Rec., July 1883 ... — — — — — N.A. —

FITZALANIA; F. v. M., fragm. IV, 33 (1863).												
F. heteropetala; F. v. M., fragm. IV, 33 (1863)	...	...	...	...	...	—	—	—	—	Q.	—	B.fl.I, 51 M.fr.III, 1; VII, 126.
CANANGA; Rumphius, herb. Amboin. II, 195, t. 65 (1741). (Fitzgeraldia.)												
C. odorata; J. Hooker & Thomson, fl. Indic. 130 (1855)	...	...	...	...	...	—	—	—	—	Q.	—	M.fr.VI, 1; VIII, 279.
ANCANA; F. v. M., fragm. V, 27 (1865).												
A. stenopetala; F. v. M., fragm. V, 27 (1865)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	M.fr.V, 189; VII, 121.
POLYALTHIA; Blume, Fl. Jav. Anon. 68 (1829).												
P. nitidissima; Benth. Fl. Austr. I, 51 (1863)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	B.fl.I, 51 M.fr.III, 2, 167; VII, 126. P.
P. Holtzeana; F. v. M. in Wing's S. Sc. Rec. II, 230 (1882)	...	...	...	...	...	—	—	—	—	—	N.A.	—
P. Armitiana; F. v. M. in Austr. Journal of Pharmacy, Jan. 1887	...	...	...	...	...	—	—	—	—	Q.	—	—
POPOWIA; Endlicher, Gen. 831 (1839).												
P. Australis; Benth. Fl. Austr. I, 52 (1863)	...	...	...	...	...	—	—	—	—	—	N.A.	B.fl.I, 52
MELODORUM; Loureiro, Fl. Cochinch. I, 351 (1790).												
M. Leichhardtii; Benth. Fl. Austr. I, 52 (1863)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	B.fl.I, 52 M.fr.III, 41, VI, 2, 176; VII, 126.
M. Uhrii; F. v. M., fragm. VI, 2 (1867)	...	...	...	...	...	—	—	—	—	Q.	—	M.fr.VI, 2.
M. Maccraei; F. v. M., fragm. VI, 176 (1868)	...	...	...	...	...	—	—	—	—	Q.	—	M.fr.VI, 250.
MITREPHORA; Spach, Veget. Phanerog. VII, 497 (1839).												
M. Froggattii; F. v. M. in Austr. Journ. of Pharmacy II, 3 (1837)	...	...	...	...	...	—	—	—	—	Q.	—	—
MILIUSIA; Leschenault in Mém. Genève. V, 213 (1832).												
Species undetermined	...	...	...	...	...	—	—	—	—	Q.	—	—
SACCOPETALUM; Bennett in Horsf. pl. Jav. rar. 115, t. 35 (1838).												
S. Bidwilli; Benth. Fl. Austr. I, 53 (1863)	...	...	...	...	...	—	—	—	—	Q.	—	B.fl.I, 53
S. Brahei; F. v. M., fragm. VIII, 159 (1874)	...	...	...	...	...	—	—	—	—	Q.	—	M.fr.VIII, 159.
EUPOMATIA; R. Brown in App. Flind. voy. II, 597, t. 2 (1814).												
E. Bennettii; F. v. M., fragm. I, 45 (1859)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	B.fl.I, 54 M.fr.VI, 177.
E. lanrina; R. Brown in App. Flind. voy. II, 597, t. 2 (1814)	...	...	...	...	...	—	—	—	—	V. N.S.W.	Q.	B.fl.I, 54 M.fr.I, 45; VI, 2; VII, 126.

MONIMIACEAE.

A. L. DE JUSSIEU, in Ann. du Mus. XIV, 30 (1809).—Genera 7, Species 16.

DORYPHORA; Endlicher, Gen. 3015 (1837). (Learosa.)												
D. fastifera; Endlicher, iconogr. gen. pl., t. 10 (1838)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	B.fl.V, 283
ATHEROSPERMA; Labillardière, Nov. Holl. pl. spec. II, 74, t. 224 (1806).												
A. moschatum; Labillardière, Nov. Holl. pl. spec. II, 74, t. 224 (1806)...	...	...	...	...	...	—	—	—	—	T. V. N.S.W.	—	B.fl.V, 284 M.fr.VIII, 142; X, 106.
DAPHNANDRA; Benth. Fl. Austr. V., 285 (1870).												
D. micrantha; Benth. Fl. Austr. V, 285 (1870)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	B.fl.V, 285
D. repandula; F. v. M., fragm. X, 105 (1877)	...	...	...	...	...	—	—	—	—	Q.	—	M.fr.X, 105.
D. aromatica; Bailey, first suppl. Q.L. fl. 46 (1886)	...	...	...	...	...	—	—	—	—	Q.	—	—
MOLLINEDIA; Ruiz & Pavon, prodr. 83, t. 15 (1794). (Kibara, Wilkiea.)												
M. Huegeliana; Tulasne in Ann. sc. nat. sér. quatr. III, 45 (1855)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	B.fl.V, 286
M. Wardellii; F. v. M., fragm. V, 155 (1866)	...	...	...	...	...	—	—	—	—	Q.	—	B.fl.V, 287 M.fr.VI, 252.
M. loxocarya; Benth. Fl. Austr. V, 287 (1870)	...	...	...	...	...	—	—	—	—	Q.	—	B.fl.V, 287
M. acuminata; F. v. M., fragm. V, 155 (1866)	...	...	...	...	...	—	—	—	—	Q.	—	B.fl.V, 287 M.fr.V, 155.
M. macrophylla; Tulasne in Ann. sc. nat. sér. quatr. III, 45 (1855)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	B.fl.V, 288 M.fr.V, 3.

M. longipes ; Bentham, Fl. Austr. V, 289 (1870) — — — — — Q. — B.fl.V,289
M. laxiflora ; Bentham, Fl. Austr. V, 289 (1870) — — — — — Q. — B.fl.V,289
M. pubescens ; Bentham, Fl. Austr. V, 290 (1870) — — — — — N.S.W. Q. — B.fl.V,290

HEDYCARIA ; R. & G. Forster, char. gen. 127, t. 64 (1776).

H. Cunninghamii ; Tulasne in Archiv. du Mus. d'hist. nat. VIII, 408 (1855) ... — — — — V. N.S.W. Q. — B.fl.V,291
--

PALMERIA ; F. v. M., fragm. IV, 151 (1864).

P. scandens ; F. v. M., fragm. IV, 152 (1864) — — — — — N.S.W. Q. — B.fl.V,291 M.fr.V,2.

PIPTOCALYX ; Oliver in Bentham's Fl. Austr. V, 292 (1870).

P. Moorei ; Oliver in Bentham's Fl. Austr. V, 292 (1870) — — — — — N.S.W. — — B.fl.V,292 M.fr.X,106.

MYRISTICÆÆ.

R. BROWN, prodr. 399 (1810) from A. L. de Jussieu (1806).—Genus 1, Species 1.

MYRISTICA ; Linné, gen. pl. ed. II, 524 (1742).

M. insipida ; R. Brown, prodr. 400 (1810) — — — — — Q. N.A. B.fl.V,281

LAURACEÆ.

VENTENAT ; Tabl. I, 245 (1799).—Genera 7, Species 37.

CRYPTOCARYA ; R. Brown, prodr. 402 (1810). (Caryodaphne.)

C. Murrayi ; F. v. M., fragm. V, 170 (1866) — — — — — Q. — B.fl.V,295 M.fr.X,253.
C. Mackinnoniana ; F. v. M., fragm. V, 169 (1866) — — — — — Q. — B.fl.V,296 M.fr.V,169.
C. patentinervis ; F. v. M. in De Candolle, prodr. XV, 508 (1864) — — — — — N.S.W. Q. — B.fl.V,296 M.fr.V,196.
C. obovata ; R. Brown, prodr. 402 (1810) — — — — — N.S.W. Q. — B.fl.V,296 M.fr.V,166,170.
C. glaucescens ; R. Brown, prodr. 402 (1810)... .. — — — — — N.S.W. Q. N.A. B.fl.V,297 M.fr.V,165,166.
C. triplinervis ; R. Brown, prodr. 402 (1810)... .. — — — — — N.S.W. Q. — B.fl.V,297 M.fr.V,166.
C. cinnamomifolia ; Bentham, Fl. Austr. V, 298 (1870) — — — — — Q. — B.fl.V,298
C. Meissneri ; F. v. M., fragm. V, 170 (1866) — — — — — N.S.W. Q. — B.fl.V,298 M.fr.V,170.
C. Australis ; Bentham, Fl. Austr. V, 299 (1870) — — — — — N.S.W. Q. — B.fl.V,299 M.fr.V,167.

BEILSCHMIEDIA ; Nees in Wallich, pl. Asiat. rarior. II, 69 (1831). (Nesodaphne.)

B. obtusifolia ; Bentham in B. & H. gen. III, 1521 (1880) — — — — — N.S.W. Q. — B.fl.V,299 M.fr.V,166.

ENDIANDRA ; R. Brown, prodr. 402 (1810).

E. glauca ; R. Brown, prodr. 402 (1810) — — — — — Q. — B.fl.V,300 M.fr.V,166.
E. hypotephra ; F. v. M., fragm. V, 166 (1866) — — — — — Q. — B.fl.V,301 M.fr.V,166.
E. discolor ; Bentham, Fl. Austr. V, 301 (1870) — — — — — N.S.W. Q. — B.fl.V,301
E. Sieberi ; Nees, Syst. Laurin. 194 (1836) — — — — — N.S.W. — — B.fl.V,301 M.fr.V,166.
E. virens ; F. v. M., fragm. II, 90 (1860) — — — — — N.S.W. Q. — B.fl.V,302 M.fr.V,166.
E. Muellieri ; Meissner in De Candolle, prodr. XV, 509 (1864) — — — — — N.S.W. Q. — B.fl.V,302
E. pubens ; Meissner in De Candolle, prodr. XV, 509 (1864) — — — — — N.S.W. Q. — B.fl.V,302

CINNAMOMUM ; Burmann, fl. Zeil. 62 (1737).

C. Tamala ; C. G. & T. F. Nees, Syst. Laurin. 56 (1836) — — — — — Q. — B.fl.V,303 M.fr.V,165 ... As.

LITSEA ; Lamarck, Diction. III, 574 (1789). (Glabraria 1771, Tetranthera, Cylicodaphne, Litsaea).

L. Chinensis ; Lamarck, Diction. III, 574 (1789) — — — — — Q. N.A. B.fl.V,305 ... As.
L. Bindoniana ; F. v. M., syst. Census of Austr. plants 4 (1882) — — — — — Q. — B.fl.V,305 M.fr.V,167.
L. Hexanthus ; A. L. de Jussieu in Annal. du Mus. VI, 212 (1805) — — — — — Q. — B.fl.V,305

L. reticulata ; Bentham in B. & J. H. gen. pl. III, part I, 161 (1880) ...	...	...	—	—	—	—	—	Q.	—	B.fl.V,306			
L. Zeylanica ; C. G. & T. F. Nees, in Amoén. Bonn. I, 58, t. 5 (1823) ...	...	...	—	—	—	—	—	Q.	—	B.fl.V,307	...	...	As.
L. dealbata ; Nees, Syst. Laurin. 630 (1836) ...	...	...	...	...	...	...	N.S.W.	Q.	—	B.fl.V,307	M.fr.V,166.		
CASSYTHA ; Osbeck in Linné, sp. pl. 35 (1753).													
C. nodiflora ; Meissner in De Candolle, prodr. XV, 252 (1864) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,309			
C. glabella ; R. Brown, prodr. 404 (1810) ...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.V,309	M.fr.V,167.		
C. flava ; Nees in Lehmann, pl. Preiss. I, 620 (1845) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,310			
C. pubescens ; R. Brown, prodr. 404 (1810) ...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.V,310	M.fr.V,167.		
C. phaeolasia ; F. v. M., fragm. V. 167 (1866) ...	...	...	—	—	—	V.	N.S.W.	—	—	B.fl.V,310			
C. paniculata ; R. Brown, prodr. 404 (1810) ...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.V,311			
C. filiformis ; Linné, sp. pl. 35 (1753) ...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.V,311	M.fr.V,167	...	As., Afr., Am.
C. melantha ; R. Brown, prodr. 404 (1810) ...	...	...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.V,311	M.fr.V,167.		
C. micrantha ; Meissner in De Candolle, prodr. XV, 256 (1864) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,312			
C. racemosa ; Nees in Lehmann, pl. Preiss. I, 621 (1845) ...	...	...	W.A.	—	—	—	—	Q.	—	B.fl.V,312			
C. pomiformis ; Nees in Lehmann, pl. Preiss. I, 620 (1845) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,313			
HERNANDIA ; Plumier, Gen. 6, t. 40 (1703).													
H. peltata ; Meissner in De Candolle, prodr. XV, 263 (1864) ...	...	...	—	—	—	—	—	Q.	—	B.fl.V,314	...	...	As., Afr., P.
H. bivalvis ; Bentham, Fl. Austr. V, 314 (1870) ...	...	...	—	—	—	—	—	Q.	—	B.fl.V,314			

MENISPERMEAE.

A. L. DE JUSSIEU, Gen. 284 (1789).—Genera 15, Species 17.

TINOSPORA: Miers in Ann. and Mag. of nat. hist. sec. ser. VII, 35 (1851).

T. <i>smilacina</i> ; Bentham in Journ. Linn. Soc. V, suppl. II, 2 (1861)	...	..	—	—	—	—	—	Q.	N.A. B.fl. I, 55	M.fr. IX, 82.
T. <i>Walcottii</i> ; F. v. M. in Bentham's Fl. Austr. I, 56 (1863)	...	...	W.A.	—	—	—	—	—	N.A. B.fl. I, 56	M.fr. IX, 83.

FAWCETTIA; F. v. M., fragm. X, 93 (1877).

F. tinosporoides; F. v. M., fragm. X, 93 (1877)	...	...	...	...	...	—	—	—	—	N.S.W.	—	—	—	M.fr.XI, 136.
---	-----	-----	-----	-----	-----	---	---	---	---	--------	---	---	---	---------------

COCCULUS; De Candolle, *sys. veg.* I, 515 (1818) from C. Bauhin (1623). (*Pericampylus*, *Legnephora*.)

C. Moorei: F. v. M., fragm. I, 162 (1859) — — — — N.S.W. Q. — B.f.I, 56 M.fr. IX, 81, 197.

HYPSERPA; Miers in Ann. and Mag. of nat. hist. sec. ser. VII, 36 (1851). (Selwynia.)

H. Selwyni; F. v. M., fragm. IX, 82 (1875) ...	...	...	...	...	...	—	—	—	—	—	Q.	—	—	M.fr.IV, 153.
--	-----	-----	-----	-----	-----	---	---	---	---	---	----	---	---	---------------

ADELIOPSIS ; Bentham & J. Hooker, Gen. pl. I, 436 (1862).

A. decumbens; Benth. Fl. Austr. I, 59 (1863) — — — — — Q. — B.f.I, 59 M.fr.IX, 83

TRISTICHOCALYX; F. v. M., fragm. IV, 27 (1863).

T. pubescens; F. v. M., fragm. IV, 27 (1863)	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.I,58	M.fr.IV,27.
T. diffusus; Miers, contrib. to Bot. III, 286 (1871)	...	...	...	...	...	—	—	—	—	—	Q.	—	—	—

SARCOPETALUM; F. v. M., pl. of Vict. I, 26 (1860).

S. *Harveyanum*; F. v. M., pl. of Vict. 27 and 221 (1860) ... — — — V. N.S.W. Q. — B.f.I, 57 M.fr. IX, 83.

LEICHHARDTIA; F. v. M., fragm. X, 67 (1876).

L. clamboïdes; F. v. M., fragm. X, 68 (1876)	...	...	...	...	...	—	—	—	—	—	Q.	—	—	M.fr. X, 68.
--	-----	-----	-----	-----	-----	---	---	---	---	---	----	---	---	--------------

STEPHANIA; Loureiro, Fl. Cochinch. II, 608 (1790).

S. hernandifolia; Walpers, rept. bot. syst. I, 96 (1842) ... — — — V. N.S.W. Q. N.A. B. fl. 1, 57 M. fr. IX, 76; X, 94 As., Afr.

CISSAMPELOS; Linné, gen. plant. 368 (1737).

C. Pereira; Linné, spec. plant. 1031 (1753)	...	...	...	...	...	...	—	—	—	—	—	Q.	—	—	As., Afr., Am.
---	-----	-----	-----	-----	-----	-----	---	---	---	---	---	----	---	---	----------------

PACHYGONE; Miers in Ann. and Mag. nat. hist. sec. ser. VII, 37 (1851).												
P. Hullsii; F. v. M., fragm. IX, 81 (1875)	..	..	...	...	...	—	—	—	—	—	Q. N.A.	— M.fr.V,147.
PYCNARRHENA; Miers in Ann. and Mag. of nat. hist. sec. ser. VII, 37 (1851).												
P. Australiana; F. v. M. in Vict. Naturalist III, 61 (1886)...	...	...	...	—	—	—	—	—	—	Q.	—	—
PLEOGYNE; Miers in Taylor's Ann. and Mag. of nat. hist. sec. ser. VII, 37 (1851). (Microclisia.)												
P. Cunninghami; Miers in Ann. and Mag. of nat. hist. sec. ser. VII, 37 (1851)...	—	—	—	—	—	—	—	—	—	Q.	— B.fl.I,59	M.fr.IX,82.
HUSEMANNIA; F. v. M. in Wing's S. Sc. Rec. III, 187 (1883).												
H. protensa; F. v. M. in Wing's S. Sc. Rec. III, 127 (1883)	...	...	...	—	—	—	—	—	—	Q.	—	—
CARRONIA; F. v. M., fragm. IX, 84 (1875).												
C. multisepeala; F. v. M., fragm. IX, 171 (1875)	...	...	...	...	...	—	—	—	—	N.S.W.	—	— M.fr.IX,84.

PAPAVERACEAE.

A. L. DE JUSSIEU; Gen. 235 (1789) from B. de Jussieu (1759).—Genus 1, Species 1.

PAPAVER; Tournefort, inst. 237, t. 119 (1700) from Plinius.

P. aculeatum; Thunberg, Fl. Capensis 431 (1813)	...	...	...	...	...	—	S.A.	T. V.	N.S.W.	Q.	— B.fl.I,63	... Afr.
---	-----	-----	-----	-----	-----	---	------	-------	--------	----	-------------	----------

CAPPARIDEAE.

VENTENAT; Tabl. III, 118 (799).—Genera 6, Species 29.

CLEOME; Linné, syst. nat. 9 (1735); Linné, gen. 200 (1737). (Polanisia.)

C. oxalidea; F. v. M., fragm. I, 69 (1858)	...	...	...	...	...	—	—	—	—	—	Q. N.A.	B.fl.I,90 M.fr.I,69
C. tetrandra; Banks in De Candolle, prodr. I, 240 (1824)	...	...	...	...	W.A.	—	—	—	—	—	Q. N.A.	B.fl.I,90 M.fr.I,70, VII,40; IX,174.
C. viscosa; Linné, sp. pl. 672 (1753)	...	...	...	...	...	W.A.	S.A.	—	—	N.S.W.	Q. N.A.	B.fl.I,90 M.fr.I,70, IX,174. As., Afr.
C. grandiflora; F. v. M., ined.	...	...	...	...	...	...	...	—	—	—	Q. N.A.	—

ROEPERIA; F. v. M. in Hooker's Kew Misc. IX, 15 (1857). (Tetratelia.)

R. cleomoides; F. v. M. in Hooker's Kew Misc. IX, 15 (1857)	...	...	...	...	...	—	—	—	—	—	Q. N.A.	B.fl.I,91 M.fr.V,105; IX,174.
---	-----	-----	-----	-----	-----	---	---	---	---	---	---------	-------------------------------

EMBLINGIA; F. v. M., fragm. II, 2 (1860).

E. calceoliflora; F. v. M., fragm. II, 3 (1860)...	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.I,92 M.fr.; XI,20.
--	-----	-----	-----	-----	-----	------	---	---	---	---	---	-------------------------

CADABA; Forskal, Fl. Aeg. Arab. 67 (1775).

C. capparoides; De Candolle, prodr. I, 244 (1824)	...	...	...	...	...	—	—	—	—	—	N.A.	B.fl.I,92 ... As.
---	-----	-----	-----	-----	-----	---	---	---	---	---	------	-------------------

CAPPARIS; Tournefort, inst. 261, t. 139 (1700) from Theophrastus, Dioscorides and Plinius. (Busbeckea, Busbecquea.)

C. umbellata; R. Brown in De Candolle, prodr. I, 247 (1824)	...	...	...	...	...	—	—	—	—	—	Q. N.A.	B.fl.I,93 M.fr.V,104; IX,172.
C. lasiantha; R. Brown in De Candolle, prodr. I, 247 (1824)	...	...	...	...	W.A.	S.A.	—	—	—	N.S.W.	Q. N.A.	B.fl.I,94 M.fr.IX,173.
C. quiniflora; De Candolle, prodr. I, 247 (1824)	...	...	...	...	...	—	—	—	—	—	Q. N.A.	B.fl.I,94 M.fr.IX,173. ... P.
C. spinosa; Linné, sp. pl. 503 (1753)	...	...	...	...	...	W.A.	S.A.	—	—	—	Q. N.A.	B.fl.I,94 M.fr.I,143; V,104; IX,173. E., As., Afr., P.
C. sarmentosa; Cunningham in Benthams Fl. Austr. I, 95 (1863)	...	...	...	...	...	—	—	—	—	—	Q.	— B.fl.I,95 M.fr.V,104; IX,173.
C. ornans; F. v. M. in Benthams Fl. Austr. I, 95 (1863)	...	...	...	...	...	—	—	—	—	—	Q.	— B.fl.I,95 M.fr.V,104; IX,173.
C. nobilis; F. v. M. in Benthams Fl. Austr. I, 95 (1863)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	— B.fl.I,95 M.fr.I,163; V,104.
C. canescens; Banks in De Candolle, prodr. I, 246 (1824)	...	...	...	...	...	—	—	—	—	—	Q.	— B.fl.I,96 M.fr.V,104; IX,173.
C. Shanesiana; F. v. M., fragm. X, 94 (1877)	...	...	...	...	...	—	—	—	—	—	Q.	— M.fr.X,94.
C. lucida; R. Brown in Benthams Fl. Austr. I, 96 (1863)	...	...	...	...	...	—	—	—	—	—	Q.	— B.fl.I,96 M.fr.I,163; V,104; IX,173.
C. Mitchellii; Lindley in Mitchell's Three Exped. I, 315 (1838)	...	...	...	...	...	—	S.A.	—	V.	N.S.W.	Q. N.A.	B.fl.I,96 M.fr.IX,173.
C. Ioranthifolia; Lindley in Mitchell's Trop. Austr. 220 (1848)	...	...	...	...	...	—	—	—	—	N.S.W.	Q. N.A.	B.fl.I,97

C. umbonata; Lindley in Mitchell's Trop. Austr. 257 (1848)	...	...	...	—	—	—	—	—	Q. N.A. B.fl.I,97	M.fr.IX,173
C. uberiflora; F. v. M., fragm. IX, 172 (1875)	...	...	...	...	...	—	—	—	Q. —	M.fr.IX,172.
C. humistrata; F. v. M., fragm. V, 156 (1866)	...	...	...	...	...	—	—	—	Q. —	M.fr.V,156.
C. Thozetiana; F. v. M., fragm. V., 104 (1866)	...	...	...	...	...	—	—	—	Q. —	M.fr.V,156.
APOPHYLLUM; F. v. M. in Hooker's Kew Misc. IX, 307 (1857).										
A. anomalum; F. v. M. in Hooker's Kew Misc. IX. 307 (1857)	...	...	...	—	—	—	—	N.S.W.	Q. N.A. B.fl.I,97	M.fr.IX,173; XI,27.

CRUCIFERE.

A. L. DE JUSSIEU, Gen. 237 (1789) from B. de Jussieu (1759).—Genera 14, Species 54.

NASTURTIIUM; R. Brown in W. T. Aiton's hortus Kewensis IV, 110 (1812) from Linné, syst. nat. 9 (1735).

N. terrestre; R. Brown in Aiton's hort. Kewens. IX, 110 (1812) ... S.A. T. V. N.S.W. Q. — B.fl.I,66 M.fr.XI,60. ... E., As., Afr., Am., P., N.Z.

BARBARAEA; Beckmann, Lex. botan. 33 (1801). (Barbarea, from L'Obel 1576.)

B. vulgaris; R. Brown in De Candolle, prodr. I, 140 (1824) ... T. V. N.S.W. — — B.fl.I,66 M.fr.VIII,142; XI,6,60 E., As., Am., N.Z.

ARABIS; Linné, gen. pl. 198 (1737) from Dalechamps (1554). (Turritis.)

A. glabra; Crantz, stirp. Austrisc. 36 (1767) ... V. N.S.W. — — B.fl.I,67 ... E., As., Am.

CARDAMINE; Tournefort, inst. 224, t. 109 (1700) from l'Ecluse (1576).

C. stylosa; De Candolle, regn. veg. syst. nat. II, 24 (1821) ... T. V. N.S.W. Q. — B.fl.I,68 M.fr.VII,19; XI,60 ... N.Z.

C. dictyosperma; Hooker, Journ. of Bot. I, 246 (1834) ... T. V. N.S.W. — — B.fl.I,68 M.fr.VII,19; X,6,60.

C. radicata; J. Hooker in icones pl. t. 882 (1852) ... T. — — — B.fl.I,69 M.fr.XI,6.

C. laciniata; F. v. M. in Transact. Phil. Soc. Vict. I, 34 (1854) ... S.A. — V. N.S.W. — — B.fl.I,69 M.fr.XI,27,60.

C. hirsuta; Linné, sp. pl. 655 (1753) ... W.A. S.A. T. V. N.S.W. Q. — B.fl.I,70 M.fr.VII,19; XI,60 E., As., Am., N.Z., P.

C. eustylis; F. v. M. in Transact. Vict. Inst. I, 114 (1854) ... S.A. — V. N.S.W. Q. N.A. B.fl.I,71 M.fr.VII,19; XI,60.

WILCKIA; Scopoli, introd. hist. nat. 317 (1777). (Malcolmia.)

W. Africana; F. v. M., native plants of Vict. I, 33 (1879) ... V. N.S.W. — — — M.fr.VII,18 ... E., As., Afr.

SISYMBRIUM; Tournefort, inst. 225, t. 109 (1700). (Blennodia.)

S. filifolium; F. v. M. in Transact. Phil. Soc. Vict. I, 34 (1854) ... S.A. — — N.S.W. — — B.fl.I,73

S. trisetum; F. v. M. in Linnaea XXV, 368 (1852) ... W.A. S.A. — V. N.S.W. Q. — B.fl.I,74 M.fr.VII,20; XI,60.

S. nasturtioides; F. v. M. in Transact. Vict. Inst. I, 115 (1854) ... S.A. — V. N.S.W. Q. — B.fl.I,74 M.fr.X,53.

S. Richardsii; F. v. M., fragm. X, 105 (1877) ... W.A. S.A. — — — — M.fr.X,105.

S. eremigenum; F. v. M., fragm. II, 143 (1861) ... — — — — N.S.W. Q. — B.fl.I,74 M.fr.II,143.

S. Lucæ; F. v. M., syst. Cens. of Austr. pl. 5 (1882) ... — — — — V. N.S.W. — — — M.fr.XI,59.

S. cardaminoides; F. v. M. in Transact. Phil. Soc. Vict. I, 34 (1854) ... S.A. — V. N.S.W. Q. — B.fl.I,75 M.fr.VII,19; VII,223; X,53; XI,6,27,59.

ERYSIMUM; Linné, gen. pl. 198 (1737). (Blennodia, Alyssopsis.)

E. curvipes; F. v. M. in Linnaea XXV, 368 (1852) ... S.A. — V. N.S.W. — — B.fl.I,75 M.fr.VII,20.

E. brevipes; F. v. M. in Linnaea XXV, 367 (1852) ... W.A. S.A. — V. N.S.W. — — B.fl.I,75 M.fr.VII,20; X,53,119; XI,6.

E. blennodioides; F. v. M. in Linnaea XXV, 367 (1852) ... S.A. — V. N.S.W. Q. — B.fl.I,76 M.fr.VII,20; XI, 27,60.

E. Cunninghamii; F. v. M. Cens. 5 (1882) from Benthams (1863) ... — — — — N.S.W. Q. — B.fl.I,76

E. Blennodia; F. v. M., fragm. X, 78 (1877) ... S.A. — — N.S.W. Q. N.A. B.fl.I,76 M.fr.VII,20.

E. capsellinum; F. v. M., native pl. of Vict. I, 35 (1879) ... — — — — V. N.S.W. — — B.fl.I,77

ALYSSUM; Tournefort, inst. 217 (1700) from l'Ecluse (1576). (Meniocus, Alysson.)

A. minimum; Pallas, Reise Prov. Russ. Reich. III, 746 (1776) ... W.A. S.A. — V. N.S.W. Q. N.A. B.fl.I,71 M.fr.XI,27,60 ... E., As., Afr.

STENOPETALUM; R. Brown in De Candolle, Mém. du Mus. VII, 239 (1821.)

S. velutinum; F. v. M., pl. Vict. I, 49 (1860) ... S.A. — V. N.S.W. Q. — B.fl.I,78 M.fr.III,27; VII,20; XI,27.

S. lineare; R. Brown in De Candolle, syst. veg. I, 513 (1821) ... W.A. S.A. T. V. N.S.W. Q. — B.fl.I,78 M.fr.VII,20; XI,6.

<i>S. sphaerocarpum</i> ; F. v. M. in Transact. Phil. Soc. Vict. I, 35 (1854) ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.I,78	M.fr.VII,20.
<i>S. nutans</i> ; F. v. M., fragm. III, 27 (1862) ...	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.I,79	M.fr.VII,20; XI,6.
<i>S. robustum</i> ; Endlicher in Huegel enum. 4 (1837) ...	W.A.	S.A.	—	—	—	—	—	B.fl.I,79	M.fr.VII,20; XI,5,60.
<i>S. pedicellare</i> ; F. v. M. in Bentham, Fl. Austr. I, 79 (1863) ...	W.A.	—	—	—	—	—	—	B.fl.I,79	M.fr.VII,20.
<i>S. croceum</i> ; Bunge in Lehmann, pl. Preiss. I, 258 (1844) ...	W.A.	—	—	—	—	—	—	—	M.fr.XI,6.
MENKEA; Lehmann, Ind. sem. hort. bot. Hamb. 8 (1843).									
<i>M. Australis</i> ; Lehmann in Ind. sem. hort. bot. Hamb. 8 (1843) ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.I,80	M.fr.II,142.
<i>M. draboides</i> ; J. Hooker in Bentham Fl. Austr. I, 80 (1863) ...	W.A.	—	—	—	—	—	—	B.fl.I,80	M.fr.II,142; VIII,223; XII,19.
<i>M. sphaerocarpa</i> ; F. v. M., fragm. VIII, 223 (1874) ...	—	S.A.	—	—	—	—	—	—	M.fr.VIII,223.
CAPSELLA; Medicus, Pflanzen-Gattungen 85 (1792). (Hutchinsia partly, Thlaspi partly, Microlepidium).									
<i>C. elliptica</i> ; C. A. Meyer, Verz. der. Pfl. v. Caucas. 194 (1831) ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.I,81	M.fr.VII,19; XI,60 E., As., Afr., Am.
<i>C. antipoda</i> ; F. v. M., pl. of Vict. I, 44 (1860) ...	—	—	T.	V.	—	—	—	B.fl.I,81	
<i>C. pilosula</i> ; F. v. M., pl. of Vict. I, 44 (1860) ...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.I,82	M.fr.XI,60.
<i>C. Tasmanica</i> ; F. v. M., fragm. XI, 26 (1878) ...	—	—	T.	—	—	—	—	B.fl.I,87	
<i>C. cochlearina</i> ; F. v. M., pl. of Vict. I, 51 (1860) ...	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.I,88	N.fr.XI,26.
<i>C. Drummondii</i> ; F. v. M., fragm. XI, 21 (1878) ...	W.A.	S.A.	—	—	—	—	—	B.fl.I,88	M.fr.VII,90; XI,26.
<i>C. humistrata</i> ; F. v. M., fragm. XI, 25 (1878) ...	—	S.A.	—	—	N.S.W.	Q.	—	—	M.fr.XI,25.
<i>C. Andraeana</i> ; F. v. M. in Wing's S. So. Rec. March p. 49 (1885) ...	—	—	—	—	N.S.W.	—	—	—	
SENEBIERA; De Candolle in Mém. soc. d'hist. nat. Par. I, 140 (1799). Carara, Medicus (1792) from Caesalpini (1583).									
<i>S. integrifolia</i> ; De Candolle in Mém. soc. d'hist. nat. Par. I, 144 (1799) ...	—	—	—	—	—	Q.	—	B.fl.I,82	... As., Afr.
LEPIDIDIUM; Tournefort, inst. 215, t. 103 (1700) from Dioscorides and Plinius. (Iberis partly, Lepia, Monoploca).									
<i>L. strongylophyllum</i> ; F. v. M. in Benth. Fl. Austr. I, 84 (1863) ...	W.A.	S.A.	—	—	—	Q.	N.A.	B.fl.I,84	M.fr.XI,6,60.
<i>L. linifolium</i> ; Bentham, Fl. Austr. I, 84 (1863) ...	W.A.	—	—	—	—	—	—	B.fl.I,84	M.fr.VII,20; XI,6.
<i>L. leptopetalum</i> ; F. v. M., pl. of Vict. I, 48 (1860) ...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.I,84	
<i>L. rotundum</i> ; De Candolle, syst. veg. II, 537 (1818) ...	W.A.	S.A.	—	—	—	—	—	B.fl.I,85	M.fr.VII,20; XI,6.
<i>L. phlebotopetalum</i> ; F. v. M., pl. of Vict. I, 47 (1860) ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.I,85	M.fr.VII,19; XI,60.
<i>L. pedicellosum</i> ; F. v. M., fragm. XI, 27 (1878) ...	—	—	—	—	—	—	N.A.	—	M.fr.XI,27.
<i>L. monoplocoides</i> ; F. v. M. in Trans. Phil. Soc. Vict. 35 (1854) ...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.I,85	M.fr.XI,27.
<i>L. papillosum</i> ; F. v. M. in Linnaea XXV, 370 (1852) ...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.I,86	M.fr.VII,19; XI,6,28.
<i>L. foliosum</i> ; Desvaux, Journ. de Bot. III, 164 et 180 (1813) ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.I,86	M.fr.VII,19; IX,76.
<i>L. ruderales</i> ; Linné, sp. pl. 645 (1753) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.I,86	M.fr.VII,19; XI,27,61.
CAKILE; Tournefort, coroll. 49, t. 483 (1700) from L'Obel (1576).									
<i>C. maritima</i> ; Scopoli, Fl. Carniolica, 344 (1760) ...	—	S.A.	T.	V.	N.S.W.	—	—	—	M.fr.VII,19; XI,6,28,60 E., As., Afr.

VIOLACEAE.

DE CANDOLLE; Flor. Franç. IV, 801 (1805).—Genera 4, Species 13.

VIOLA; Tournefort, inst. 419, t. 236 (1700) from Plinius. (Erpetion.)									
<i>V. betonicifolia</i> ; Smith in Rees's Cyclopaedia XXXVII (1817) ...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,99	M.fr.X,82.
<i>V. hederacea</i> ; Labillardière, Nov. Holl. pl. spec. I, 66, t. 91 (1804) ...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,99	M.fr.X,82.
<i>V. Cunninghamii</i> ; J. Hooker, Fl. N. Zel. I, 16 (1853) ...	—	—	T.	—	—	—	—	B.fl.I,100	... N.Z.
<i>V. Caleyana</i> ; G. Don, gen. syst. dichlam. pl. I, 329 (1831) ...	—	—	T.	V.	N.S.W.	—	—	B.fl.I,100	M.fr.X,82.
HYBANTHUS; Jacquin, Stirp. Amer. hist. 77, t. 75 (1763). (Ionidium, Pigea, Vlamingia).									
<i>H. floribundus</i> ; F. v. M., nat. pl. of Vict. I, 45 (1879) ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.I,112	M.fr.X,81.
<i>H. Vernonii</i> ; F. v. M., nat. pl. of Vict. I, 45 (1879) ...	—	—	—	V.	N.S.W.	—	—	B.fl.I,103	M.fr.X,81.
<i>H. filiformis</i> ; F. v. M., nat. pl. of Vict. I, 44 (1879) ...	—	—	—	V.	N.S.W.	Q.	—	B.fl.I,103	M.fr.X,81.

H. debilissimus; F. v. M., fragm. XI, 4 (1878)	W.A.	—	—	—	—	—	—	—	M.fr.X,4.
H. Tatei; F. v. M. in Trans. R. Soc. of S. Austr. IX, 102 (1882)	—	S.A.	—	—	—	—	—	—	
H. calycinus; F. v. M., fragm. X, 81 (1877)	W.A.	—	—	—	—	—	—	B.fl.I,104	M.fr.XI,4.
H. enneaspermus; F. v. M., fragm. X, 81 (1877)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.I,101	M.fr.X,81 ... As., Afr.
HYMENANTHERA; R. Brown in Tuck. Cong. 442 (1818).									
H. Banksii; F. v. M., first gen. report 9 (1853)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.I,104	M.fr.IX,77; X,82 N.Z.
MELICYTUS; R. & G. Forster, char. gen. 123, t. 62 (1776).									
M. ramiflorus; R. & G. Forst., char. gen. 123, t. 2 (1776)	—	—	—	—	N.S.W.	—	—	—	 N.Z.

FLACOURTIEAE.

CL. RICHARD in Mém. du Mus. I, 366 (1815).—Genera 4, Species 7.

COCHLOSPERMUM; Kunth, Malvac. 6 (1822).									
C. Fraseri; Planchon in Hook. Lond. Journ. VI, 307 (1847)	—	—	—	—	—	—	N.A.	B.fl.I,106	M.fr.IX,83.
C. Gillivraei; Benth. Fl. Austr. I, 106 (1863)	—	—	—	—	—	Q.	—	B.fl.I,106	
C. Gregorii; F. v. M., fragm. I, 71 (1853)	—	—	—	—	—	Q.	N.A.	B.fl.I,106	M.fr.IX,61.
SCOLOPIA; Schreber, Gen. 355 (1789). (Phoberos).									
S. Brownii; F. v. M., fragm. III, 11 (1862)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,107	M.fr.IV,174; IX,61.
XYLOSMA; G. Forster, prodr. 380 (1786).									
X. ovata; Benth. Fl. Austr. I, 108 (1863)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,108	M.fr.IX,60.
STREPTOTHAMNUS; F. v. M., fragm. III, 28 (1862).									
S. Moorei; F. v. M., fragm. III, 28 (1862)	—	—	—	—	N.S.W.	—	—	B.fl.I,108	M.fr.III,28.
S. Beckleri; F. v. M., fragm. III, 28 (1862)	—	—	—	—	N.S.W.	—	—	B.fl.I,108	M.fr.IX,61.

SAMYDACEAE.

J. GAERTNER; de Fruct. III, 328 (1805).—Genera 2, Species 4.

CASEARIA; Jacquin, Stirp. Amer. hist. 132, t. 85 (1763).									
C. esculenta; Roxburgh. Fl. Ind. II, 422 (1832)	—	—	—	—	—	Q.	—	B.fl.VI,309	 As.
C. Dallachii; F. v. M., fragm. V, 107 (1866)	—	—	—	—	—	Q.	—	B.fl.III,309	M.fr.V,214; VI,252.
HOMALIMUM; Jacquin, Stirp. Amer. hist. 170, t. 183 (1763). (Blackwellia.)									
H. Vitiense; Benth. in Journ. Linn. Soc. IV, 36 (1860)	—	—	—	—	—	Q.	—	B.fl.III,310	 P.
H. brachybotrys; F. v. M., fragm. II, 127 (1861)	—	—	—	—	—	Q.	—	B.fl.III,310	M.fr.II,127.

PITTOSPOREAE.

R. BROWN in Flinders' voy. 542 (1814).—Genera 8, Species 40.

PITTOSPORUM; Banks in Gaertner, de Fruct. I, 266, t. 59 (1788).									
P. rhombifolium; Cunningham in Hook. icon. pl. t. 621 (1844)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,110	M.fr.XII,4.
P. melanosperrum; F. v. M., fragm. I, 70 (1853)	—	—	—	—	—	—	N.A.	B.fl.I,111	M.fr.VI,167; XII,4.
P. undulatum; Andrews, Bot. reposit. 383 (1800)	—	—	—	—	T.	V.	N.S.W.	Q.	— B.fl.I,111 M.fr.VI,167; XII,4.
P. revolutum; Aiton, Hort. Kew, sec. ed. II, 27 (1811)	—	—	—	—	V.	N.S.W.	Q.	— B.fl.I,111	M.fr.VI,167,186; VII,140; XII,4.
P. venulosum; F. v. M., fragm. VI, 186 (1868)	—	—	—	—	—	Q.	—	—	M.fr.VII,140; XII,4.
P. ferrugineum; Aiton, Hort. Kew, sec. ed. II, 27 (1811)	—	—	—	—	—	Q.	—	B.fl.I,112	M.fr.II,78; III,165; VI,167; XII,4.
P. bracteolatum; Endlicher, prodr. fl. ias. Norf. 78 (1833)	—	—	—	—	N.S.W.	—	—	—	 As., P.

P. Wingii; F. v. M. in Wing's S. Sc. Rec. March (1885), p. 49	...	...	...	...	...	...	...	...	...	Q.	—			
P. rubiginosum; Cunningham in Ann. of Nat. Hist. IV, 108 (1842)	...	...	...	...	...	...	...	...	...	Q.	—	B. fl. I, 112	M. fr. VI, 167; VII, 140; IX, 190.	
P. phillyroides; De Candolle, prodr. I, 347 (1824)	...	...	...	...	...	W. A.	S. A.	—	V.	N. S. W.	Q.	N. A.	B. fl. I, 112	M. fr. VI, 167; IX, 75.
P. bicolor; Hooker, Journ. Bot. I, 249 (1834)	...	...	...	...	...	—	—	T.	V.	N. S. W.	—	—	B. fl. I, 113	M. fr. VIII, 142; XII, 4.
P. erioloma; C. Moore et F. v. M., fragm. VII, 139 (1871)	...	...	...	...	...	—	—	—	—	N. S. W.	—	—	—	M. fr. VII, 139.
HYMENOSPORUM; R. Brown in Hook. Bot. Magaz. 4799 (1854).														
H. flavum; F. v. M., fragm. II, 77 (1860)	...	...	...	...	...	...	—	—	—	N. S. W.	Q.	—	B. fl. I, 114	M. fr. V, 210; VI, 168; VII, 140; XII, 4.
BURSARIA; Cavanilles, icon. et descr. pl. IV, 30, t. 350 (1797).														
B. spinosa; Cavanilles, icon. pl. IV, 30, t. 350 (1797)	...	...	...	...	...	W. A.	S. A.	T.	V.	N. S. W.	Q.	N. A.	B. fl. I, 115	M. fr. XII, 4.
MARIANTHUS; Huegel, enum. pl. Nov. Holl. 8 (1837). (Oncosporum, Rhytidosporum, Calopetalon, Spiranthera.)														
M. procumbens; Benth. Fl. Austr. I, 117 (1863)	...	...	...	...	...	—	—	T.	V.	N. S. W.	Q.	—	B. fl. I, 117	M. fr. XII, 2.
M. rhytidosporeus; F. v. M., fragm. II, 145 (1861)	...	...	...	...	...	W. A.	—	—	—	—	—	—	B. fl. I, 117	M. fr. II, 145.
M. villosus; Benth. Fl. Austr. I, 117 (1863)	...	...	...	...	...	W. A.	—	—	—	—	—	—	B. fl. I, 117	
M. parviflorus; F. v. M., fragm. II, 144 (1861)	...	...	...	...	...	W. A.	—	—	—	—	—	—	B. fl. I, 118	M. fr. II, 144; XII, 2.
M. bignoniaceus; F. v. M. in Traas. Phil. Soc. Vict. 6 (1854)	...	...	...	...	...	—	S. A.	—	V.	—	—	—	B. fl. I, 118	M. fr. XII, 2.
M. Drummondianus; Benth. Fl. Austr. I, 119 (1863)	...	...	...	...	...	W. A.	—	—	—	—	—	—	B. fl. I, 119	M. fr. XII, 2.
M. coeruleo-punctatus; Klotzsch in Link and Otto, icon. pl. 28, t. 12 (1841)	...	...	...	...	...	W. A.	—	—	—	—	—	—	B. fl. I, 119	M. fr. XII, 2.
M. candidus; Huegel, enum. pl. Nov. Holl. 8 (1837)	...	...	...	...	...	W. A.	—	—	—	—	—	—	B. fl. I, 120	M. fr. XII, 2.
M. erubescens; Putterlick in Endl. nov. stirp. dec. 60 (1839)	...	...	...	...	...	W. A.	—	—	—	—	—	—	B. fl. I, 120	M. fr. I, 219; XII, 2.
M. ringens; F. v. M., fragm. I, 218 (1859)	...	...	...	...	...	W. A.	—	—	—	—	—	—	B. fl. I, 120	M. fr. I, 218; XII, 2.
M. pictus; Lindley in Bot. Regist. XXV, App. XXV (1839)	...	...	...	...	...	W. A.	—	—	—	—	—	—	B. fl. I, 121	M. fr. I, 217; II, 78, 182; XII, 2.
CITRIBATUS; Cunningham in Loudon's hort. Brit. Suppl. I, 585 (1832). (Ixiosporum.)														
C. multiflorus; Cunningham in Loud. hort. Brit. I, 585 (1832)	...	...	...	...	...	—	—	—	—	N. S. W.	Q.	—	B. fl. I, 121	M. fr. II, 77; VI, 168; XII, 4.
C. pauciflorus; Cunningham in Loud. hort. Brit. I, 585 (1832)	...	...	...	...	...	—	—	—	—	N. S. W.	Q.	N. A.	B. fl. I, 122	M. fr. II, 76; VI, 168; VII, 180.
BILLARDIERA; Smith, Sp. Bot. New. Holl. I, t. 1 (1793). (Pronaya, Campylanthera.)														
B. longiflora; Labillardiere, Nov. Holl. plant. spec. I, 64, t. 89 (1804)	...	...	...	...	...	—	—	T.	V.	N. S. W.	—	—	B. fl. I, 123	
B. scandens; Smith, Spec. bot. New Holl. I, t. 1 (1793)	...	...	...	...	...	—	S. A.	T.	V.	N. S. W.	Q.	—	B. fl. I, 123	M. fr. XI, 123.
B. coriacea; Benth. Fl. Austr. I, 124 (1863)	...	...	...	...	...	W. A.	—	—	—	—	—	—	B. fl. I, 124	M. fr. XII, 1.
B. cymosa; F. v. M. in Transact. Vict. Inst. I, 29 (1855)	...	...	...	...	...	—	S. A.	—	V.	N. S. W.	—	—	B. fl. I, 124	M. fr. VI, 168; VII, 140.
B. elegans; F. v. M., pl. of Vict. I, 78 (1860)	...	...	...	...	...	W. A.	—	—	—	—	—	—	B. fl. I, 125	M. fr. XII, 2.
B. varifolia; De Candolle, prodr. I, 346 (1824)	...	...	...	...	...	W. A.	—	—	—	—	—	—	B. fl. I, 125	M. fr. II, 145.
B. Lehmanniana; F. v. M., pl. of Vict. I, 78 (1860)	...	...	...	...	...	W. A.	—	—	—	—	—	—	B. fl. I, 125	M. fr. XII, 2.
B. floribunda; F. v. M. in South. Sc. Record II, 1 (1882)	...	...	...	...	...	W. A.	—	—	—	—	—	—	B. fl. I, 120	M. fr. XII, 1.
SOLLYA; Lindley, Bot. Regist. XVII, t. 1460 (1831). (Xerosollya.)														
S. heterophylla; Lindley, Bot. Regist. XVII, t. 1466 (1831)	...	...	...	...	...	W. A.	—	—	—	—	—	—	B. fl. I, 126	M. fr. XII, 3.
S. parviflora; Turcz. in Bull. Soc. Mosc. XXVII, 361 (1854)	...	...	...	...	...	W. A.	—	—	—	—	—	—	B. fl. I, 126	M. fr. XII, 3.
CHEIRANTHERA; Cunningham in Brongn. Bot. voy. Coquille, t. 77 (1829).														
C. linearis; Cunningham in Bot. Regist. 1719 (1834)	...	...	...	...	...	—	S. A.	—	V.	N. S. W.	Q.	—	B. fl. I, 127	M. fr. I, 97; II, 180; VI, 168; IX, 168; XII, 4.
C. filifolia; Turczaninow in Bull. Soc. Mosc. XXVII, 364 (1854)	...	...	...	...	...	W. A.	—	—	—	—	—	—	B. fl. I, 127	M. fr. I, 97; II, 79; VI, 168; IX, 168; XII, 4.
C. parviflora; Benth. Fl. Austr. I, 128 (1863)	...	...	...	...	...	W. A.	—	—	—	—	—	—	B. fl. I, 128	M. fr. XII, 3.

DROSERACEAE.

SALISBURY; Parad. Lond. 95 (1809).—Genera 3, Species 46.

ALDROVANDA; Monti in Act. Bonon. II, 404, t. 12 (1747).

A. vesiculosa; Linné, sp. pl. 281 (1753)	...	...	...	...	...	...	—	—	—	—	Q.	—	—	M.fr.VI,104;X,79	...	E., As.
--	-----	-----	-----	-----	-----	-----	---	---	---	---	----	---	---	------------------	-----	---------

DROSERA ; Linné, gen. pl. 89 (1773). (Sondera).

D. Indica ; Linné, sp. pl. 232 (1753)	W.A. S.A. — V. N.S.W. Q. N.A. B.fl.II,456 M.fr.X,79 ... As., Afr.
D. Adelaë ; F. v. M., fragm. IV, 154, t. XXXIII (1864)	— — — — — Q. — — M.fr.X,80.
D. Arcturi ; Hooker, Journ. of Bot. I, 247 (1834)	— — — — — T. V. N.S.W. — — B.fl.II,456 ... N.Z.
D. glanduligera ; Lehmann, pugill. VIII, 37 (1844)	W.A. S.A. — V. N.S.W. — — B.fl.II,457 M.fr.X,80.
D. pygmaea ; De Candolle, prodr. I, 317 (1824)	— S.A. T. V. N.S.W. — — B.fl.II,457 M.fr.X,80 ... N.Z.
D. platystigma ; Lebmann, pugill. VIII, 37 (1844)	W.A. — — — — — B.fl.II,457 M.fr.X,80.
D. pulchella ; Lehmann, pugill. VIII, 38 (1844)	W.A. — — — — — B.fl.II,458 M.fr.X,80.
D. leucoblasta ; Bentham, Fl. Austr. II, 458 (1864)	W.A. — — — — — B.fl.II,458 M.fr.X,80.
D. nitidula ; Planchon in Ann. des sc. nat. IX, 285 (1848)	W.A. — — — — — B.fl.II,458
D. paleacea ; R. Brown in De Candolle, prodr. I, 318 (1824)	W.A. — — — — — B.fl.II,458
D. parvula ; Planchon in Ann. des sc. nat. IX, 287 (1848)	W.A. — — — — — B.fl.II,459
D. Burmanni ; Vahl, Symbol. III, 50 (1794)	— S.A. — — N.S.W. Q. N.A. B.fl.II,459 M.fr.X,80. ... As.
D. spatbulata ; Labillardière, Nov. Holl. pl. spec. I, 79, t. 106 (1804)	— S.A. T. V. N.S.W. Q. — B.fl.II,459 ... As., N.Z.
D. Drummondii ; Lehmann, pl. Preiss. II, 235 (1847)	W.A. — — — — — B.fl.II,460
D. scorpoides ; Planchon in Ann. des sc. nat. IX, 288 (1848)	W.A. — — — — — B.fl.II,460 M.fr.VIII,185 ; X,85.
D. petiolaris ; R. Brown in De Candolle, prodr. I, 318 (1824)	— — — — — Q. N.A. B.fl.II,460 M.fr.X,80.
D. binata ; Labillardière, Nov. Holl. pl. spec. I, 78, t. 105 (1804)	— S.A. T. V. N.S.W. Q. — B.fl.II,461 M.fr.X,80 ... N.Z.
D. zonaria ; Planchon in Ann. des sc. nat. IX, 303 (1848)	W.A. — — — — — B.fl.II,462
D. bulbosa ; Hooker, icon. pl. t. 375 (1841)	W.A. — — — — — B.fl.II,462
D. rosulata ; Lehmann, pugill. VIII, 36 (1844)	W.A. — — — — — B.fl.II,462 M.fr.X,80.
D. Whittakerii ; Planchon in Ann. des sc. nat. IX, 302 (1848)	— S.A. — V. — — — B.fl.II,462
D. macrophylla ; Lindley, Bot. Regist. XXV, App. XX (1839)	W.A. — — — — — B.fl.II,463
D. squamosa ; Bentham, Fl. Austr. II, 463 (1864)	W.A. — — — — — B.fl.II,463
D. erythrorrhiza ; Lindley, Bot. Regist. XXV, App. XX (1839)	W.A. — — — — — B.fl.II,463
D. stolonifera ; Endlicher in Hueg. enum. 5 (1837)	W.A. — — — — — B.fl.II,463
D. humilis ; Planchon in Ann. des sc. nat. IX, 300 (1848)	W.A. — — — — — B.fl.II,464
D. ramellosa ; Lehmann, pugill. VIII, 40 (1844)	W.A. — — — — — B.fl.II,464
D. platypoda ; Turczaninow in Bull. de l'Ac. de St. Petersb. 343 (1852)	W.A. — — — — — B.fl.II,464 M.fr.VIII,185 ; X,80.
D. auriculata ; Backhouse in Ann. des sc. nat. IX, 295 (1848)	— S.A. T. V. N.S.W. Q. — B.fl.II,465 M.fr.X,80 ... N.Z.
D. lunata ; Buchanan in De Candolle, prodr. I, 319 (1824)	— — — — — Q. — — ... As.
D. peltata ; Smith in Willd. spec. pl. I, 1546 (1797)	— S.A. T. V. N.S.W. Q. — B.fl.II,465 M.fr.X,80.
D. Neesii ; Lehmann, pugill. VIII, 42 (1844)	W.A. — — — — — B.fl.II,466
D. gigantea ; Lindley, Bot. Regist. XXV, App. XX (1839)	W.A. — — — — — B.fl.II,466
D. myriantha ; Planchon in Ann. des sc. nat. IX, 291 (1848)	W.A. — — — — — B.fl.II,466
D. pallida ; Lindley, Bot. Regist. XXV, App. XX (1839)	W.A. — — — — — B.fl.II,467
D. penicillaris ; Bentham, Fl. Austr. II, 467 (1864)	W.A. — — — — — B.fl.II,467
D. filicaulis ; Endlicher in Hueg. enum. 6 (1837)	W.A. — — — — — B.fl.II,467
D. Huegelii ; Endlicher in Hueg. enum. 6 (1837)	W.A. — — — — — B.fl.II,467 M.fr.VIII,185 ; X,81.
D. macrantha ; Endlicher in Hueg. enum. 6 (1837)	W.A. — — — — — B.fl.II,468
D. Menziesii ; R. Brown in De Candolle, prodr. I, 319 (1824)	W.A. S.A. T. V. N.S.W. — — B.fl.II,468
D. calycina ; Planchon in Ann. des sc. nat. IX, 299 (1848)	W.A. — — — — — B.fl.II,468
D. heterophylla ; Lindley, Bot. Regist. XXV, App. XX (1839)	W.A. — — — — — B.fl.II,469
D. Banksii ; R. Brown in De Candolle, prodr. I, 319 (1824)	— — — — — Q. — B.fl.II,469

BYBLIS ; Salisbury Parad. Lond. t. 95 (1808).

B. liniflora ; Salisbury, Parad. Lond. t. 95 (1808)	W.A. — — — — — Q. N.A. B.fl.II,470 M.fr.X,81.
B. gigantea ; Lindley Bot. Regist. XXV, App. XXI (1839)	W.A. — — — — — ... B.fl.II,470

ELATINEAE.

CAMBESSEDES in Mém. du Mus. XVIII 225 (1829).—Genera 2, Species 4.

ELATINE; Linné, gen. pl., 18 (1737).

E. Americana; Arnott in Edinb. journ. nat. sc. I, 431 (1830) ... W.A. S.A. T. V. N.S.W. Q. ... B.fl.I,178 M.fr.XI,27 ... Am., P., N.Z.

BERGIA; Linné, mantiss. II, 152 (1771).

B. ammaonioides; Roxburgh, hort. Bengh. 34 (1814) ... — S.A. — V. N.S.W. Q. N.A. B.fl.I,180 M.fr.II,147; IV,173 As., Afr

B. pedicellaris; F. v. M. in Benth. Fl. Austr. I, 180 (1863) ... — — — — — N.A. B.fl.I,180 M.fr.II,145; IV,173.

B. perennis; F. v. M. in Benth. Fl. Austr. I, 180 (1863) ... — S.A. — — — Q. N.A. B.fl.I,181 M.fr.II,146.

HYPERICINAE.

J. DE ST. HILAIRE, Expos. Fam. II, 23, t. 75 (1805).—Genus 1, Species 1.

HYPERICUM; Tournefort, inst. 254, t. 131 (1700), from Dioscorides and Plinius. (Ascyrum, Brathys).

H. Japonicum; Thunberg, Fl. Jap. 295, t. 31 (1784) ... W.A. S.A. T. V. N.S.W. Q. N.A. B.fl.I,182 ... As., N.Z.

TERNSTROEMIACEAE.

MIRBEL in Bullet. Soc. Philomat. Dec. p. 381 (1813).—Genus 1, Species 1.

SARAUIA; Willdenow in Schrift. nat. Freunde, Berlin III, 406, t. 4 (1801).

S. Andraeana; Dyer & Oliver in Bailey, synops. Queensl. flora, suppl. (1885) ... — — — — — Q. — — M.fr.V,175.

GUTTIFERAE.

A. L. DE JUSSIEU; Gen. 243 (1789).—Genera 2, Species 3.

CALOPHYLLUM; Linné, gen. 154 (1737).

C. inophyllum; Linné, sp. pl. 513 (1753) ... — — — — — Q. N.A. B.fl.I,183 M.fr.IX,175 ... As., Afr., P.

C. tomentosum; Wight, Illustrat. Pl. Ind. Orient. I, 128 (1840) ... — — — — — Q. — — M.fr.IX,174 ... As.

KAYEA; Wallich, pl. As. rar. III, 4 (1832).

K. Larnachiana; F. v. M. in Viet. Nat. III, 126 (1887) ... — — — — — Q. —

POLYGALEAE.

A. L. DE JUSSIEU, in Ann. c. Mus. XIV, 386 (1809).—Genera 4, Species 32.

SALOMONIA; Loureiro, Fl. Cochinch. I, 14 (1790)

S. oblongifolia; De Candolle, prodr. I, 334 (1824) ... — — — — N.S.W. Q. N.A. B.fl.I,138 M.fr.XI,4 ... As., P.

POLYGALA; Tournefort, inst. 174, t. 79 (1700), from Dioscorides and Plinius.

P. Sibirica; Linné, sp. pl. 702 (1753) ... — — — — V. N.S.W. Q. — B.fl.I,139 M.fr.XI,4 ... As.

P. leptalea; De Candolle, prodr. I, 325 (1824) ... — — — — — Q. N.A. B.fl.I,139 M.fr.XI,4 ... As.

P. eriocephala; F. v. M. in Benth. Fl. Austr. I, 139 (1863) ... — — — — — N.A. B.fl.I,139

P. persicarifolia; De Candolle, prodr. I, 326 (1818) ... — — — — — Q. N.A. — M.fr.XI,4 ... As., Afr.

P. Chinensis; Linné, sp. pl. 704 (1753) ... — S.A. — — N.S.W. Q. N.A. B.fl.I,140 M.fr.XI,4 ... As.

P. stenoclada; Benth, Fl. Austr. I, 141 (1863) ... — — — — — N.A. B.fl.I,141 M.fr.XI,4

COMESPERMA; Labillardière, Nov. Holl. pl. spec. II, 21 (1806).

C. sphaerocarpum; Steetz in Lehm. pl. Preiss. II, 304 (1847)	...	...	...	...	...	...	...	N.S.W.	Q.	—	B.fl.I, 143	M.fr.I, 145; XI, 3.			
C. scoparium; Steetz in Lehm. pl. Preiss. II, 314 (1847)	...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	—	B.fl.I, 143	M.fr.I, 145.		
C. aphyllum; R. Brown in Benth. Fl. Austr. I, 143 (1863)...	...	...	...	...	...	...	—	—	—	—	N.A.	B.fl.I, 143.			
C. spinosum; F. v. M., fragm. I, 144 (1859) ...	...	...	...	...	...	...	W.A.	—	—	—	—	B.fl.I, 144	M.fr.I, 144.		
C. volubile; Labillardière, Nov. Holl. pl. spec. II, 24, t. 163 (1806)	...	...	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	B.fl.I, 144	M.fr.XI, 3.	
C. ciliatum; Steetz in Lehm. pl. Preiss. II, 304 (1847)	...	...	...	...	...	...	W.A.	—	—	—	—	B.fl.I, 144	M.fr.X, 3.		
C. integerimum; Endlicher in Hueg. enum. (1837) ...	...	...	...	...	...	...	W.A.	—	—	—	—	B.fl.I, 145	M.fr.XI, 3.		
C. secundum; Banks in De Candolle, prodr. I, 334 (1824) ...	...	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.I, 145.	
C. Drummondii; Steetz in Lehm. pl. Preiss. II, 301 (1847)...	...	...	...	...	...	...	W.A.	—	—	—	—	B.fl.I, 145	M.fr.X, 5; XI, 2.		
C. praelsum; F. v. M., fragm. XI, 2 (1878) ...	...	...	...	...	...	...	...	—	—	—	—	Q.	—	M.fr.XI, 2.	
C. rectusum; Labillardière, Nov. Holl. pl. spec. II, 22, t. 160 (1806)	...	...	...	...	...	...	—	—	T.	V.	N.S.W.	Q.	B.fl.I, 145	M.fr.XI, 2.	
C. sylvestre; Lindley in Mitch. Trop. Austr. 342 (1848) ...	...	...	...	...	...	...	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.I, 146	M.fr.I, 49; X, 5; XI, 2.
C. viscidulum; F. v. M., fragm. X, 4 (1876) ...	...	...	...	...	...	...	W.A.	S.A.	—	—	—	—	—	M.fr.XI, 2.	
C. accrosum; Steetz in Lehm. pl. Preiss. II, 299 (1847) ...	...	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.I, 146	M.fr.XI, 3.	
C. ericium; De Candolle, prodr. I, 334 (1824) ...	...	...	...	...	...	...	—	—	T.	V.	N.S.W.	Q.	—	B.fl.I, 146	M.fr.I, 48; XI, 2.
C. confertum; Labillardière, Nov. Holl. pl. spec. II, 23, t. 161 (1806)	...	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.I, 147	M.fr.XI, 3.	
C. flavum; De Candolle, prodr. I, 334 (1824) ...	...	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.I, 147	M.fr.XI, 3.	
C. calymega; Labillardière, Nov. Holl. pl. spec. II, 23, t. 162 (1806)	...	...	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	—	B.fl.I, 147	M.fr.XI, 3.	
C. rhadinocarpum; F. v. M., fragm. XI, 1 (1878) ...	...	...	...	...	...	...	W.A.	—	—	—	—	—	—	M.fr.XI, 1.	
C. lanceolatum; R. Brown in Benth. Fl. Austr. I, 148 (1863) ...	...	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.I, 148	M.fr.XI, 1.	
C. defoliatum; F. v. M., pl. of Vict. I, 189 (1862) ...	...	...	...	...	...	...	—	—	T.	V.	N.S.W.	Q.	—	B.fl.I, 148	M.fr.XI, 3.
C. nudiusculum; De Candolle, prodr. I, 334 (1824) ...	...	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.I, 148	M.fr.XI, 4.	
C. virgatum; Labillardière, Nov. Holl. pl. spec. II, 21, t. 159 (1806)	...	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.I, 149	M.fr.XI, 2.	
C. polygaloides; F. v. M. in Trans. Phil. Soc. Vict. I, 7 (1854) ...	...	...	...	...	...	...	W.A.	S.A.	—	V.	—	—	B.fl.I, 149	M.fr.XI, 4.	

XANTHOPHYLLUM; Roxburgh, Pl. Corom. III, 81, t. 284 (1819). (Macintyria.)

X. Macintyrii; F. v. M., fragm. V, 57 (1865)...	...	...	...	...	...	—	—	—	—	—	—	—	—	M.fr.V, 8; XI, 4.
---	-----	-----	-----	-----	-----	---	---	---	---	---	---	---	---	-------------------

TREMANDREAE.

R. BROWN in Flind. voy. II, 554 App. (1814).—Genera 3, Species 17.

PLATYTHECA; Steetz in Lehmann, pl. Preiss. I, 220 (1845).

P. galioides; Steetz in Lehm. pl. Preiss. I, 220 (1845) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.I, 136	M.fr.XII.
---	-----	-----	-----	-----	------	---	---	---	---	---	---	-------------	-----------

TETRATHECA; Smith, Spec. Bot. New Holl. I, t. 2 (1793).

T. Harperi; F. v. M., fragm. V, 49 (1865) ...	...	...	...	...	...	W.A.	—	—	—	—	—	—	—	M.fr.V, 189.
T. ciliata; Lindley in Mitch. three Exped. II, 206 (1838) ...	...	...	...	...	...	—	S.A.	T.	V.	—	—	—	B.fl.I, 130	M.fr.XII, 7.
T. efoliata; F. v. M., fragm. X, 3 (1876) ...	...	...	...	...	...	W.A.	—	—	—	—	—	—	—	M.fr.X, 3.
T. aphylla; F. v. M. in South. Science Record, vol. II, May (1882) ...	...	...	...	...	...	W.A.	—	—	—	—	—	—	—	M.fr.XII, 5.
T. ericifolia; Smith, Exot. Bot. I, 37, t. 20 (1804) ...	...	...	...	...	...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I, 131	M.fr.XII, 6.
T. juncea; Smith, specim. of Bot. of New Holl. 5, t. 2 (1793) ...	...	...	...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.I, 132	M.fr.XII, 6.
T. affinis; Endlicher in Hueg. enum. 7 (1837) ...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.I, 132	M.fr.XII, 5.
T. nuda; Lindley, Bot. Regist. XXV, App. XXXVIII (1839) ...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.I, 133	M.fr.XII, 5.
T. virgata; Steetz in Lehm. pl. Preiss. I, 212 (1845)...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.I, 133	M.fr.XII, 5.
T. confertifolia; Steetz in Lehm. pl. Preiss. I, 214 (1845) ...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.I, 133	M.fr.XII, 6.
T. setigera; Endlicher in Hueg. enum. 8 (1837) ...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.I, 133	M.fr.XII, 8.
T. hirsuta; Lindley, Bot. Regist. XXV, App. XXXVIII (1839) ...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.I, 134	M.fr.XII, 7.
T. pilifera; Lindley, Bot. Regist. XXV, App. XXXVIII (1839) ...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.I, 135	M.fr.XII, 8.
T. filiformis; Bentham, Fl. Austr. I, 135 (1863) ...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.I, 135	M.fr.XII, 8.

TREMANDRA; R. Brown in App. Flind. voy. II, 554 (1814).

T. stelligera; R. Brown in De Candolle, prodr. I, 344 (1824)	W.A.	—	—	—	—	—	—	B.fl.I,136	M.fr.XII,8.
T. diffusa; R. Brown in De Candolle, prodr. I, 344 (1824)	W.A.	—	—	—	—	—	—	B.fl.I,137	M.fr.XII,9.

MELIACEAE.

VENTENAT; Tabl. III, 159 (1799).—Genera 11, Species 36.

TURRAEA; Linné mantiss. II, 150 (1771).

T. pubescens; Hellenius in K. Sv. Vet. Acad. Handl. 26, t. 10 (1788)	N.S.W.	Q.	N.A.	B.fl.I,379	M.fr.V,144	...	As.
---	--------	----	------	------------	------------	-----	-----

MELIA; Linné, gen. pl. 127 (1737).

M. Azedarach; Linné, sp. pl. 384 (1753)	N.S.W.	Q.	N.A.	B.fl.I,380	M.fr.V,144 ; IX,134 ; XI,135	As., Afr.
--	--------	----	------	------------	------------------------------	-----------

DYSOXYLUM; Blume, Bijdrag. 172 (1825). (Dysoxylon, Epicharis, Hartigsheea.)

D. oppositifolium; F. v. M., fragm. V, 144 (1866)	Q.	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
--	----	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---

AMOORA; Roxburgh, pl. Corom. III, 54, t. 258 (1819).

A. nitidula; Benth. Fl. Austr. I, 383 (1863)	N.S.W.	Q.	—	B.fl.I,383	M.fr.V,145	...	P.
---	--------	----	---	------------	------------	-----	----

SYNOUM; Adr. de Jussieu in Mém. du Mus. XIX, 226, t. 15 (1830).

A. glandulosum; A. de Jussieu in Mém. du Mus. XIX, 227, t. 15 (1830)	N.S.W.	Q.	—	B.fl.I,384	M.fr.V,145 ; IX,134.
A. Muellieri; C. de Candolle in Monogr. phanerog. I, 593 (1878)	Q.	—	—	—	M.fr.V,145 ; XI,138.

OWENIA; F. v. M. in Hook. Kew Misc. IX, 304 (1857).

<i>O. acidula</i> ; F. v. M. in Hook. Kew Misc. IX, 303 (1857)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.I,385	M.fr.III,14 ; IX,134.
<i>O. vernicosa</i> ; F. v. M., fragm. III, 15 (1862)... ..	—	—	—	—	—	—	N.A.	B.fl.I,385	M.fr.IX,134.
<i>O. venosa</i> ; F. v. M. in Hook. Kew Misc. IX, 304 (1857)	—	—	—	—	—	Q.	—	B.fl.I,386	M.fr.III,14 ; IX,134 ; XI,82.
<i>O. cepiodora</i> ; F. v. M., fragm. XI, 81 (1890)... ..	—	—	—	—	N.S.W.	—	—	—	M.fr.XI,87
<i>O. reticulata</i> ; F. v. M. in Hook. Kew Misc. IX, 305 (1857)	—	W.A.	—	—	—	—	N.A.	B.fl.I,386	M.fr.III,13 ; IX,134.

AGLAIA; Loureiro, Fl. Cochinch. I, 173 (1790), non Allemand (1770). (Milnea, Nemedra.)

A. elaeagnoides; Benth. Fl. Austr. I, 383 (1863)	Q.	N.A.	B.fl.I,383	M.fr.V,145 ; IX,134.
---	----	------	------------	----------------------

HEARNIA; F. v. M., fragm. V, 55 (1865). (Aglaiopsis).

H. sapindina; F. v. M., fragm. V, 55 (1865)	Q.	—	B.fl.I,387	M.fr.V,55.
--	----	---	------------	------------

CARAPA; Aublet, Hist. pl. Guian. II, Suppl. 32, t. 387 (1775). (Xylocarpus.)

C. Moluccensis; Lamarck, Encycl. meth. Bot. I, 621 (1783)	Q.	N.A.	B.fl.I,387	M.f.V,145 ; IX,134.	As., Afr.
--	----	------	------------	---------------------	-----------

CEDRELA; P. Browne, Nat. hist. of Jamaica 153 (1756).

C. Australis; F. v. M., fragm. I, 4 (1858)	N.S.W.	Q.	—	B.fl.I,387	M.fr.IX,133 ; XI,82.
---	--------	----	---	------------	----------------------

FLINDERSIA; R. Brown in Flind. voy. II, 595 (1814). (Oxleya, Strzeleckia.)

F. Australis; R. Brown in Flind. voy. II, 595, t. 1 (1814)	...	...	...	...	...	...	...	N.S.W.	Q.	—	B.fl.I, 388	M.fr.I, 65; II, 174; III, 26; IX, 133.
F. Bourjotiana; F. v. M., fragm. IX, 133 (1875)	...	...	...	...	...	...	...	—	Q.	—	—	M.fr.IX, 133.
F. Schottiana; F. v. M., fragm. III, 25 (1862)	...	...	...	...	...	...	...	N.S.W.	Q.	—	B.fl.I, 388	M.fr.V, 143; VI, 143; IX, 133.
F. Oxleyana; F. v. M., fragm. I, 65 (1858)	...	...	...	...	...	...	...	N.S.W.	Q.	—	B.fl.I, 389	M.fr.III, 25; IX, 133.
F. Pimenteliana; F. v. M., fragm. IX, 132 (1875)	...	...	...	...	...	...	...	—	Q.	—	—	M.fr.IX, 132.
F. Bennettiana; F. v. M. in Benth. Fl. Austr. I, 389 (1863)	...	...	...	...	...	...	...	N.S.W.	Q.	—	B.fl.I, 389	M.fr.IX, 131; XI, 135.
F. Iflaiana; F. v. M., fragm. X, 94 (1877)	...	...	...	...	...	...	...	—	Q.	—	—	M.fr.X, 94.
F. Brayleyana; F. v. M., fragm. V, 143 (1866)	...	...	...	...	...	...	...	—	Q.	—	—	M.fr.VI, 252.
F. Strzeleckiana; F. v. M., fragm. I, 65 (1858)	...	...	...	...	...	...	...	N.S.W.	Q.	—	B.fl.I, 389	M.fr.IX, 133.

OCHNACEAE.

DE CANDOLLE in Ann. du Mus. XVII, 398 (1811), Ochnae subgenus.—Genus 1, Species 1.

BRACKENRIDGEA; Asa Gray, Bot. Wilk. Expl. Exped. 361, t. 42 (1854).

B. Australiana; F. v. M., fragm. V, 29 (1865)	...	...	...	...	...	...	...	—	Q.	—	—	M.fr.V, 213.
---	-----	-----	-----	-----	-----	-----	-----	---	----	---	---	--------------

RUTACEAE.

A. L. DE JUSSIEU, Gen. 296 (1789).—Genera 26, Species 190.

ZIERIA; Smith in Transact. Linn. Soc. IV, 216 (1798).

Z. laevigata; Smith in Rees's Cyclopaedia XXXIX (1819)	...	...	...	—	—	—	V.	N.S.W.	Q.	—	B.fl.I,304	M.fr.I,101; IX,116.	
Z. pilosa; Rudge in Transact. Linn. Soc. X, 293, t. 17 (1809)	...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.I,305	M.fr.I,101; IX,116.	
Z. obcordata; Cunningham in Field, N.S.Wales, 330 (1625)	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,305	M.fr.100; IX,116.	
Z. cytisoides; Smith in Rees's Cyclopaedia XXXIX (1819)	..	..	...	—	—	—	T.	V.	N.S.W.	Q.	—	B.fl.I,306	M.fr.IX,116.
Z. Smithii; Andrews, Bot. Rep. t. 606 (1810)	...	...	...	—	—	—	T.	V.	N.S.W.	Q.	—	B.fl.I,306	M.fr.I,100; IX,116.
Z. granulata; C. Moore in Benth. Fl. Austr. I, 307 (1863)...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,307	M.fr.I,101; IX,116.	
Z. veronicea; F. v. M. in Trans. phil. Soc. Vict. I, 11 (1854)	...	...	...	—	—	—	S.A.	T.	V.	—	—	B.fl.I,305	M.fr.IX,116.

BORONIA; Smith, Tracts relat. nat. hist. 285 (1798). (Cyanothamnus.)

B. grandisepala; F. v. M., fragm. I, 66 (1858)	...	...	...	...	...	...	...	—	—	—	—	—	N.A.	B.fl.I,311	M.fr.II,179.
B. artemisifolia; F. v. M., fragm. I, 66 (1858)	...	...	...	...	...	...	...	—	—	—	—	—	N.A.	B.fl.I,311	M.fr.IX,111.
B. affinis; R. Brown in Benth. Fl. Austr. I, 311 (1863)	...	...	...	...	...	...	...	—	—	—	—	—	N.A.	B.fl.I,311	
B. filicifolia; Cunningham in Benth. Fl. Austr. I, 311 (1863)	...	...	...	...	...	...	...	—	—	—	—	—	N.A.	B.fl.I,311	
B. alata; Smith in Transact. Linn. Soc. VIII, 283 (1807)	...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.I,312	M.fr.IX,111.
B. algida; F. v. M. in Transact. phil. Soc. Vict. I, 100 (1855)	...	...	...	...	...	...	...	—	—	—	V.	N.S.W.	—	B.fl.I,312	M.fr.IX,111.
B. Ewardsii; Bentham, Fl. Austr. I, 312 (1863)	...	...	...	...	...	...	...	S.A.	—	—	—	—	—	B.fl.I,312	
B. ternata; Endlicher, nov. stirp. dec. 6 (1839)	...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.I,312	M.fr.IX,111.
B. ericifolia; Bentham, Fl. Austr. I, 313 (1863)	...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.I,313	M.fr.IX,111.
B. inconspicua; Bentham, Fl. Austr. I, 313 (1863)	...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.I,313	
B. eriantha; Lindley in Mitch. Trop. Austr. 298 (1848)	...	...	...	...	...	...	...	—	—	—	—	—	Q.	B.fl.I,313	
B. alulata; Solander in Benth. Fl. Austr. I, 313 (1863)	...	...	...	...	...	...	...	—	—	—	—	—	Q.	B.fl.I,313	
B. ledifolia; J. Gay, monogr. des Lasiopetalées 20 (1821)	...	...	...	...	...	...	...	—	—	—	N.S.W.	Q.	—	B.fl.I,314	M.fr.I,67; IX,111.
B. lanceolata; F. v. M., fragm. I, 66 (1858)	...	...	...	...	...	...	...	—	—	—	—	Q.	N.A.	B.fl.I,314	M.fr.VII,38; IX,112; XI,97.
B. Bowmanii; F. v. M., fragm. IV, 135 (1864)	...	...	...	...	...	...	...	—	—	—	—	Q.	—	—	M.fr.VII,38.
B. Fraseri; Hooker, Bot. Mag. t. 4052 (1844)	...	...	...	...	...	...	...	—	—	—	N.S.W.	—	—	B.fl.I,315	M.fr.IX,111.
B. mollis; Cunningham in Lindl. Bot. Reg. 47 (1841)	...	...	...	...	...	...	...	—	—	—	N.S.W.	—	—	B.fl.I,315	M.fr.IX,112.

B. megastigma; Nees in Lehm. Pl. Preiss. II, 227 (1847)	W.A.	—	—	—	—	—	—	B.fl.I,315	M.fr.II,97; IV,172; IX,112.
B. heterophylla; F. v. M., fragm. II, 98 (1861)	W.A.	—	—	—	—	—	—	B.fl.I,315	M.fr.IX,112.
B. elatior; Bartling in Lehm. pl. Preiss. I, 170 (1844)	W.A.	—	—	—	—	—	—	B.fl.I,316	M.fr.II,98; IX,122; XI,97.
B. tetrandra; Labillardière, Nov. Holl. pl. spec. I, 98, t. 125 (1804)	W.A.	—	—	—	—	—	—	B.fl.I,316	
B. crassifolia; Bartling in Lehm. pl. Preiss. I, 169 (1844)	W.A.	—	—	—	—	—	—	B.fl.I,316	M.fr.IX,112.
B. albiflora; R. Brown in Benth. Fl. Austr. I, 317 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,317	M.fr.IX,114.
B. lanuginosa; Endlicher in Hueg. enum. 16 (1837)	W.A.	—	—	—	—	—	—	B.fl.I,317	M.fr.IX,114.
B. pulchella; Turczaninow in Bull. Soc. Mosc. XXV, 162 (1852) ..	W.A.	—	—	—	—	—	—	B.fl.I,318	M.fr.IX,114.
B. gracilipes; F. v. M., fragm. II, 99 (1861)	W.A.	—	—	—	—	—	—	B.fl.I,318	M.fr.IX,114.
B. microphylla; Sieber in Spreng. syst. cur. post. 148 (1827)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,318	
B. floribunda; Sieber in Spreng. syst. veg. cur. post. 148 (1827)	—	—	—	—	N.S.W.	—	—	—	
B. pinnata; Smith, Tracts relat. nat. hist. 290, t. 4 (1798)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,318	M.fr.IX,114.
B. pilosa; Labillardière, Nov. Holl. pl. spec. I, 97, t. 124 (1804)	—	—	T.	V.	N.S.W.	—	—	B.fl.I,319	
B. spinescens; Benth. Fl. Austr. I, 319 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,319	
B. baeckeacea; F. v. M., fragm. IV, 28 (1863)	W.A.	—	—	—	—	—	—	—	M.fr.IX,112.
B. coerulescens; F. v. M. in Trans. phil. Soc. Vict. II (1854)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.I,320	M.fr.VII,38; IX,112.
B. defoliata; F. v. M., fragm. IX, 113 (1875)	W.A.	—	—	—	—	—	—	—	M.fr.IX,113.
B. Busselliana; F. v. M., fragm. IX, 113 (1875)	W.A.	—	—	—	—	—	—	—	M.fr.IX,113.
B. subcoerulea; F. v. M., fragm. II, 100 (1860)	W.A.	—	—	—	—	—	—	B.fl.I,320	M.fr.IX,112&198.
B. polygalifolia; Smith, Tracts relat. nat. hist. 297, t. 7 (1798)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,320	M.fr.IX,114.
B. falcifolia; Cunningham in Lindl. Bot. Reg. 47 (1841)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,322	M.fr.IX,114.
B. penicillata; Benth. Fl. Austr. I, 322 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,322	
B. crassipes; Bartling in Lehm. pl. Preiss. I, 168 (1844)	W.A.	—	—	—	—	—	—	B.fl.I,322	M.fr.IX,116.
B. subsessilis; Benth. Fl. Austr. I, 322 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,322	
B. capitata; Benth. Fl. Austr. I, 323 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,323	M.fr.IX,116.
B. nematophylla; F. v. M., fragm. II, 100 (1861)	W.A.	—	—	—	—	—	—	B.fl.I,323	M.fr.II,100.
B. crenulata; Smith in Transact. Linn. Soc. VIII, 284 (1807)	W.A.	—	—	—	—	—	—	B.fl.I,323	M.fr.XI,97,115&198.
B. serrulata; Smith, Tracts relat. nat. hist. 292 (1798)	—	—	—	—	N.S.W.	—	—	B.fl.I,323	M.fr.IX,115.
B. rhomboidea; Hooker icon. pl. t. 722 (1848)	—	—	T.	—	N.S.W.	—	—	B.fl.I,324	
B. parviflora; Smith, Tracts relat. nat. hist. 295, t. 5 (1798)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,324	M.fr.IX,115.
B. viminea; Lindley, Bot. Regist. XXV, App. XVII (1839)	W.A.	—	—	—	—	—	—	B.fl.I,324	
B. filifolia; F. v. M., fragm. I, 3 (1858)	—	S.A.	—	V.	—	—	—	B.fl.I,325	M.fr.IX,114.
B. clavellifolia; F. v. M. in Trans. phil. Soc. Vict. I, 12 (1854)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.I,325	M.fr.I,99; VII,38; IX,114&198.
B. scabra; Lindley, Bot. Regist. XXV, App. XVII (1839)	W.A.	—	—	—	—	—	—	B.fl.I,326	
B. thymifolia; Turczaninow in Bull. Soc. Mosc. XXV, part II, 165 (1852)	W.A.	—	—	—	—	—	—	B.fl.I,326	M.fr.I,99; II,99; IX,116.
B. ovata; Lindley, Bot. Regist. 47 (1841)	W.A.	—	—	—	—	—	—	B.fl.I,326	M.fr.IX,116.
B. Barkeriana; F. v. M., fragm. XI, 96 (1880)	—	—	—	—	N.S.W.	—	—	—	M.fr.XI,96.
B. denticulata; Smith in Transact. Linn. Soc. VIII, 284 (1807)	W.A.	—	—	—	—	—	—	B.fl.I,327	M.fr.IX,115&198.
B. Machardiana; F. v. M., fragm. IX, 115 (1875)	W.A.	—	—	—	—	—	—	—	M.fr.IX,115.
B. spatulata; Lindley, Bot. Regist. XXV, App. XVII (1839)	W.A.	—	—	—	—	—	—	B.fl.I,327	M.fr.IX,115.
B. juncea; Bartling in Lehm. pl. Preiss. I, 166 (1844)	W.A.	—	—	—	—	—	—	B.fl.I,327	M.fr.IX,115.
B. cymosa; Endlicher in Hueg. enum. 16 (1837)	W.A.	—	—	—	—	—	—	B.fl.I,328	M.fr.II,101; IX,116.

ERIOSTEMON; Smith in Transact. Linn. Soc. IX, 221 (1798). (Crowea, Phebalium, Asterolasia, Microcybe, Pleurandropsis, Geleznovia, Urocarpus, Sandfordia, Actinostigma.)

E. virgatus; Cunningham in Hooker's Journ. of Bot. II, 417 (1840)	—	—	T.	—	—	—	—	B.fl.I,332	M.fr.IX,111.
E. nodiflorus; Lindley, Bot. Regist. XXV, App. XVII (1839)	W.A.	—	—	—	—	—	—	B.fl.I,336	M.fr.IX,110; X,53.
E. spicatus; A. Richard, sert. Astrolab. 76, t. 27 (1833)	W.A.	—	—	—	—	—	—	B.fl.I,336	M.fr.IX,110.
E. pungens; Lindley in Mitch. Three Exped. II, 156 (1838)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.I,338	M.fr.IX,110.

<i>E. montanus</i> ; F. v. M., pl. of Vict. I, 129 (1860)	—	—	T.	—	—	—	—	B.fl.I,338	M.fr.IX,109.
<i>E. phyllicifolius</i> ; F. v. M., fragm. I, 105 (1859)	—	—	—	V.	N.S.W.	—	—	B.fl.I,339	M.fr.I,105.
<i>E. umbellatus</i> ; Turczaninow in Bull. Soc. Mosc. XXII, part III, 15 (1849)	—	—	—	V.	N.S.W.	—	—	B.fl.I,339	M.fr.I,104; IX,109.
<i>E. Ralstoni</i> ; F. v. M., fragm. II, 101, t. 14 (1861)	—	—	—	V.	N.S.W.	—	—	B.fl.I,339	M.fr.III,165; IX,109.
<i>E. Hillebrandi</i> ; F. v. M. in Transact. phil. Soc. Vict. I, 10 (1854)	—	—	S.A. T.	V.	—	—	—	B.fl.I,340	M.fr.I,4; IX,109.
<i>E. Coxii</i> ; F. v. M. in Melb. Chem., Dec. (1884)	—	—	—	—	N.S.W.	—	—	—	—
<i>E. lamprophyllus</i> ; F. v. M. in Journ. pharm. Soc. Vict. II, 43 (1858)	—	—	—	V.	N.S.W.	—	—	B.fl.I,340	M.fr.IX,109.
<i>E. elatior</i> ; F. v. M., fragm. I, 181 (1859)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,340	M.fr.IX,109.
<i>E. amplifolius</i> ; F. v. M. in Melb. Chem., Dec. (1884)	—	—	—	V.	N.S.W.	—	—	—	—
<i>E. ambiens</i> ; F. v. M., fragm. VI, 166 (1868)	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,110.
<i>E. Oldfieldii</i> ; F. v. M., fragm. I, 3 (1858)	—	—	T.	—	—	—	—	B.fl.I,340	M.fr.I,181; IX,109.
<i>E. rotundifolius</i> ; Cunningham in Hueg. enum. 15 (1837)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,341	—
<i>E. microphyllus</i> ; F. v. M. in Transact. phil. Soc. Vict. I, 99 (1854)	—	—	S.A.	—	—	—	—	B.fl.I,341	M.fr.IX,109.
<i>E. phyllicoides</i> ; F. v. M., fragm. I, 107 (1859)	—	—	—	V.	N.S.W.	—	—	B.fl.I,341	M.fr.IX,107.
<i>E. ozothamnoides</i> ; F. v. M., fragm. I, 103 (1859)	—	—	—	V.	N.S.W.	—	—	B.fl.I,342	M.fr.IX,107.
<i>E. Mortonii</i> ; F. v. M., fragm. IX, 108 (1875)	—	—	—	—	N.S.W.	—	—	B.fl.I,342	M.fr.IX,108.
<i>E. sediflorus</i> ; F. v. M., fragm. I, 102 (1859)	—	—	S.A. T.	V.	N.S.W.	Q.	—	B.fl.I,342	M.fr.I,102.
<i>E. Nottii</i> ; F. v. M., fragm. VI, 22 (1867)	—	—	—	—	—	Q.	—	—	M.fr.VI,22.
<i>E. lepidotus</i> ; Sprengel, syst. II, 322 (1825)	—	—	S.A.	V.	N.S.W.	Q.	—	B.fl.I,343	M.fr.I,104; IX,107.
<i>E. stenophyllus</i> ; F. v. M., from Benth. Fl. Austr. I, 343 (1863)	—	—	S.A.	V.	N.S.W.	—	—	—	—
<i>E. alpinus</i> ; F. v. M., fragm. I, 103 (1859)	—	—	—	V.	N.S.W.	—	—	—	M.fr.I,244.
<i>E. tuberculosus</i> ; F. v. M., pl. of Vict. I, 130 (1862)	W.A.	—	—	—	—	—	—	B.fl.I,343	M.fr.IX,108.
<i>E. Benthami</i> ; F. v. M., fragm. IX, 108 (1875)	W.A.	—	—	—	—	—	—	B.fl.I,343	M.fr.IX,108.
<i>E. Maxwellii</i> ; F. v. M., fragm. IX, 108 (1875)	W.A.	—	—	—	—	—	—	—	M.fr.IX,103.
<i>E. filifolius</i> ; F. v. M., fragm. IX, 108 (1875)	W.A.	—	—	—	—	—	—	B.fl.I,344	M.fr.IX,108.
<i>E. squameus</i> ; Labillardière, Nov. Holl. pl. spec. I, 3, t. 141 (1804)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.I,344	M.fr.I,104; IX,107.
<i>E. anceps</i> ; Sprengel, syst. II, 322 (1825)	W.A.	—	—	—	—	—	—	B.fl.I,345	M.fr.I,103; IX,108.
<i>E. ovatifolius</i> ; F. v. M., fragm. I, 103 (1859)	—	—	—	V.	N.S.W.	—	—	B.fl.I,345	M.fr.I,103.
<i>E. bilobus</i> ; F. v. M., fragm. I, 102 (1859)	W.A.	—	—	—	—	—	—	B.fl.I,345	M.fr.IX,108.
<i>E. Beckleri</i> ; F. v. M., fragm. IX, 109 (1875)	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,108.
<i>E. capitatus</i> ; F. v. M., fragm. I, 106 (1859)	W.A.	S.A.	—	V.	—	—	—	B.fl.I,346	M.fr.IX,107.
<i>E. Geleznowii</i> ; F. v. M., fragm. I, 107 (1859)	W.A.	—	—	—	—	—	—	B.fl.I,347	M.fr.IX,107.
<i>E. correifolius</i> ; F. v. M., fragm. I, 105 (1859)	—	—	—	V.	N.S.W.	—	—	B.fl.I,350	M.fr.IX,107.
<i>E. Cunninghamii</i> ; F. v. M., fragm. IX, 107 (1875)	—	—	—	—	N.S.W.	—	—	B.fl.I,351	M.fr.IX,107.
<i>E. mollis</i> ; F. v. M., syst. cens. of Austr. pl. II (1882), from Benth. (1863)	—	—	—	—	N.S.W.	—	—	B.fl.I,349	—
<i>E. pleurandroides</i> ; F. v. M., fragm. I, 106 (1859)	—	—	—	V.	—	—	—	B.fl.I,349	M.fr.IX,107.
<i>E. trymalioides</i> ; F. v. M., fragm. I, 106 (1859)	—	—	—	V.	N.S.W.	—	—	B.fl.I,351	M.fr.I,106.
<i>E. Hookeri</i> ; F. v. M., fragm. I, 104 (1859)	W.A.	—	—	—	—	—	—	B.fl.I,352	M.fr.VII,22; IX,107.
<i>E. pallidus</i> ; F. v. M., syst. cens. of Austr. pl. II (1882), from Benth. (1863)	W.A.	—	—	—	—	—	—	B.fl.I,352	—
<i>E. Drummondii</i> ; F. v. M., fragm. I, 105 (1859)	W.A.	—	—	—	—	—	—	B.fl.I,352	M.fr.VII,22.
<i>E. grandiflorus</i> ; F. v. M., fragm. I, 105 (1859)	W.A.	—	—	—	—	—	—	B.fl.I,353	M.fr.VI,21.
<i>E. Crowei</i> ; F. v. M., pl. of Vict. I, 119 (1860)	—	—	—	V.	N.S.W.	Q.	—	B.fl.I,329	M.fr.IX,106.
<i>E. Turczaninowii</i> ; F. v. M., pl. of Vict. I, 120 (1860)	W.A.	—	—	—	—	—	—	B.fl.I,330	M.fr.IX,106.
<i>E. lanceolatus</i> ; K. F. Gaertner, de fruct. III, 154, t. 210 (1807)	—	—	—	V.	N.S.W.	—	—	B.fl.I,331	M.fr.IX,110.
<i>E. Banksii</i> ; Cunningham in Hueg. enum. 15 (1837)	—	—	—	—	—	Q.	—	B.fl.I,332	—
<i>E. trachyphyllus</i> ; F. v. M. in Transact. phil. Soc. Vict. I, 99 (1854)	—	—	—	V.	N.S.W.	—	—	B.fl.I,332	M.fr.IX,110.
<i>E. myoporoides</i> ; De Candolle, prodr. I, 720 (1824)	—	—	—	V.	N.S.W.	Q.	—	B.fl.I,333	M.fr.XI,110.
<i>E. hispidulus</i> ; Sieber in Spreng. syst. cur. post. 164 (1827)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,333	M.fr.XI,110.

<i>E. buxifolius</i> ; Smith in Rees' Cyclop. XIII, 2 (1809)	...	...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.I,333	M.fr.IX,111.	
<i>E. obovalis</i> ; Cunningham in Field's N.S. Wales, 331 (1825)	...	...	...	...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.I,334	M.fr.IX,110.	
<i>E. scaber</i> ; Paxton, Mag. of Bot. XIII, 127 (1846)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,334	M.fr.XI,111.
<i>E. Brucei</i> ; F. v. M., fragm. VII, 38 (1869)	...	...	...	...	...	W.A.	—	—	—	—	—	—	—	M.fr.IX,110.
<i>E. linearis</i> ; Cunningham in Hueg. enum. 16 (1837)	...	...	...	...	...	—	S.A.	—	—	N.S.W.	—	—	B.fl.I,334	
<i>E. difformis</i> ; Cunningham in Hueg. enum. 15 (1837)	...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.I,335	M.fr.IX,110.
<i>E. parvifolius</i> ; R. Brown in Benth. Fl. Austr. I, 335 (1863)	...	...	...	...	...	...	—	—	—	—	Q.	—	B.fl.I,335	
<i>E. ericifolius</i> ; Cunningham in Benth. Fl. Austr. I, 335 (1863)	...	...	...	...	...	...	—	—	—	N.S.W.	—	—	B.fl.I,335	

PHILOTHECA; Rudge in Transact. Linn. Soc. XI, 298, t. 21 (1815) (Drummondia.)

<i>P. Australis</i> ; Rudge in Transact. Linn. Soc. XI, 298, t. 21 (1815)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,348	M.fr.VII,20,141.
<i>P. ericoides</i> ; F. v. M., fragm. I, 107 (1859) implied	W.A.	—	—	—	—	—	—	B.fl.I,349	M.fr.VII,21.
<i>P. Hasselli</i> ; F. v. M. in Wing's S. Sc. Rec. III, 3 (1883)	W.A.	—	—	—	—	—	—	—	
<i>P. calida</i> ; F. v. M., fragm. VII, 21 (1869)	—	—	—	—	—	Q.	—	—	M.fr.VII,38.

CORREA; Smith in Transact. Linn. Soc. IV, 219 (1798). (Didymeria, Mazeutoxeron, Automachia.)

<i>C. aemula</i> ; F. v. M., fragm. I, 3 (1859)	—	S.A.	—	V.	—	—	—	B.fl.I,353	M.fr.IX,117.
<i>C. alba</i> ; Andrews, Bot. Reposit. t. 18 (1797)... ..	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.I,354	M.fr.IX,117.
<i>C. speciosa</i> ; Andrews, Bot. Reposit. t. 653 (1811)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,354	M.fr.IX,117.
<i>C. Laurenciana</i> ; Hooker, Journ. of Bot. I, 254 (1834)	—	—	—	T.	V.	N.S.W.	—	B.fl.I,355	M.fr.VIII,142; IX,117.
<i>C. decumbens</i> ; F. v. M. in Transact. phil. Soc. Vict. I, 30 (1854)	—	S.A.	—	—	—	—	—	B.fl.I,356	
<i>C. Bacnerlenii</i> ; F. v. M. in proc. L. S. of N.S.W. IX, 960 (1884)	—	—	—	—	N.S.W.	—	—	—	

NEMATOLEPIS; Turczaninow in Bull. Soc. Mosc. XXV, part III, 158 (1852). (Symphyopetalum.)

<i>N. phlebaloides</i> ; Turczaninow in Bull. Soc. Mosc. part III, 158 (1852)... ..	W.A.	—	—	—	—	—	—	B.fl.I,351	M.fr.II,102; IX,106.
<i>N. Euphemiae</i> ; F. v. M., fragm. III, 149, t. 25 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,345	M.fr.IX,166.

CHORILAENA; Endlicher in Hueg. enum. 17 (1837).

<i>C. quercifolia</i> ; Endlicher in Hueg. enum. 17 (1837)	W.A.	—	—	—	—	—	—	B.fl.I,357	M.fr.IX,106.
<i>C. hirsuta</i> ; Benth. Fl. Austr. I, 357 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,357	M.fr.IX,106.

DIPLOLAENA; R. Brown in Append. Flind. voy. II, 546 (1814).

<i>D. Dampieri</i> ; Desfontaines in Mém. du Mus. 452, t. 20 (1817)	W.A.	—	—	—	—	—	—	B.fl.I,358	M.fr.IX,106.
--	------	---	---	---	---	---	---	------------	--------------

BOSISTOA; F. v. M. in Benth. Fl. Austr. I, 359 (1863).

<i>B. sapindiformis</i> ; F. v. M. in Benth. Fl. Austr. I, 359 (1863)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,359	M.fr.IX,103; III,41,168; IV,175.
<i>B. euodiformis</i> ; F. v. M., fragm. IX, 174 (1875)	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,174.

ACRADENIA; Kippist in Transact. Linn. Soc. XXI, 207 (1855).

<i>A. Frankliniae</i> ; Kippist in Transact. Linn. Soc. XXI, 207 (1855)	—	—	—	T.	—	—	—	B.fl.I,323	M.fr.VI,167; IX,103.
--	---	---	---	----	---	---	---	------------	----------------------

EUODIA; R. et G. Forster, char. gen. 13, t. 7 (1776). (Melicope, Medicosma.)

<i>E. Cunninghami</i> ; F. v. M., fragm. III, 2 (1862)	—	—	—	—	N.S.W.	—	—	B.fl.I,362	M.fr.I,27; II,178; IX,102.
<i>E. Fareana</i> ; F. v. M., fragm. IX, 101 (1875)... ..	—	—	—	—	—	Q.	—	—	M.fr.IX,101.
<i>E. erythrocoeca</i> ; F. v. M., fragm. I, 26 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,360	M.fr.II,103,178; IX,102.
<i>E. contermina</i> ; Moore & F. v. M., fragm. VII, 144 (1871)	—	—	—	—	N.S.W.	—	—	—	M.fr.VII,144.
<i>E. octandra</i> ; F. v. M., fragm. II, 102 (1861)	—	—	—	—	N.S.W.	—	—	B.fl.I,360	M.fr.IV,117; IX,102.
<i>E. microcoeca</i> ; F. v. M., fragm. I, 144 (1859)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,361	M.fr.II,103; III,3,180; IX,102.
<i>E. xanthoxyloides</i> ; F. v. M., fragm. IV, 155 (1864)... ..	—	—	—	—	—	Q.	—	—	M.fr.IV,155.
<i>E. Bonwickii</i> ; F. v. M., fragm. V, 56 (1865)	—	—	—	—	—	Q.	—	—	M.fr.V,179.
<i>E. alata</i> ; F. v. M., fragm. VII, 142 (1871)	—	—	—	—	—	Q.	—	—	M.fr.VII,142
<i>E. accedens</i> ; Blume, Bijdr. 246 (1825)... ..	—	—	—	—	N.S.W.	Q.	—	—	M.fr.V,4,56,179; VII,22; IX,102. As.
<i>E. litoralis</i> ; Endlicher, prodr. Fl. Norfolk. 86 (1833)	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,102.

E. polybotrya; Moore & F. v. M., fragm. VII, 143 (1871) ...	...	...	...	...	...	...	...	...	N.S.W.	—	—	—	M.fr.IX,193.
E. vitiflora; F. v. M., fragm. VII, 144 (1871) ...	...	...	...	...	...	...	...	...	—	Q.	—	—	M.fr.VII,144.
E. haplophylla; F. v. M., fragm. V, 179 (1866) ...	...	...	...	...	...	...	...	...	—	Q.	—	—	M.fr.V,179.
BROMBYA; F. v. M., fragm. V, 4 (1865).													
B. platynema; F. v. M., fragm. V, 4 (1865) ...	...	...	...	...	...	...	...	...	—	Q.	—	—	M.fr.V,56; IX,102.
PAGETIA; F. v. M., fragm. V, 178 (1866).													
P. medicinalis; F. v. M., fragm. V, 178 (1866) ...	...	...	...	...	...	...	...	...	—	Q.	—	—	M.fr.V,215; VI,167; IX,103.
BOUCHARDATIA; Baillon in Adans. VII, 350 t. 10 (1867).													
B. neurococca; Baillon in Adans. IX, 110 (1869) ...	...	...	...	...	...	...	...	...	N.S.W.	Q.	—	B.fl.I,360	M.fr.I,28; II,103,178; IX,103.
XANTHOXYLUM; Catesby Nat. Hist. of Carolina I, 26 (1731). (Zanthoxylum, Blackburnia.)													
X. brachyacanthum; F. v. M., pl. of Vict. I, 108 (1862) ...	...	...	...	...	...	...	...	...	N.S.W.	Q.	—	B.fl.I,363	M.fr.VII,141; IX,104.
X. Blackburnia; Bentham, Fl. Austr. I, 363 (1863) ...	...	...	...	...	...	...	...	...	N.S.W.	—	—	B.fl.I,363	M.fr.VII,141.
X. parviflorum; Bentham, Fl. Austr. I, 363 (1863) ...	...	...	...	...	...	...	...	...	—	N.A.	B.fl.I,363	M.fr.	IX,104.
X. torvum; F. v. M., fragm. VII, 140 (1871) ...	...	...	...	...	...	...	...	...	—	Q.	—	—	M.fr.VII,140; IX,104.
X. venenificum; Bailey, catal. of Ql. woods, 8 (1886) ...	...	...	...	...	...	...	...	...	—	Q.	—	—	
GEIJERA; Schott, Rutac. 7, t. 4 (1834). (Coatesia.)													
G. Muellieri; Bentham, Fl. Austr. I, 364 (1863) ...	...	...	...	...	...	...	...	...	—	Q.	—	B.fl.I,364	M.fr.I,26; IX,104.
G. salicifolia; Schott, Rutac. 7, t. 4 (1834) ...	...	...	...	...	...	...	...	...	N.S.W.	Q.	—	B.fl.I,364	M.fr.VII,22; IX,105; XI,134.
G. parviflora; Lindley in Mitch. Trop. Austr. 102 (1848) ...	...	...	...	...	...	...	...	W.A. S.A.	V. N.S.W.	Q.	—	B.fl.I,364	M.fr.VII,22; IX,105; X,53; XI,139.
PLEIOCOCCA; F. v. M., fragm. IX, 117 (1875).													
P. Wilcoxiana; F. v. M., fragm. IX, 117 (1875) ...	...	...	...	...	...	...	...	...	N.S.W.	Q.	—	—	M.fr.IX,117.
ACRONYCHIA; R. & G. Forster, char. gen. 53, t. 27 (1776). (Cyminosma.)													
A. Baueri; Schott, Rutac. 5, t. 3 (1834) ...	...	...	...	...	...	...	...	...	N.S.W.	Q.	—	B.fl.I,366	M.fr.I,26; II,178; V,179; IX,103.
A. laevis; R. & G. Forster, char. gen. 53, t. 27 (1776) ...	...	...	...	...	...	...	...	...	V. N.S.W.	Q.	—	B.fl.I,366	M.fr.I,27; V,179; IX,104. P.
A. imperforata; F. v. M., fragm. I, 26 (1858) ...	...	...	...	...	...	...	...	...	—	Q.	—	B.fl.I,367	M.fr.IX,104.
A. melicopoides; F. v. M., fragm. V, 3 (1865) ...	...	...	...	...	...	...	...	...	N.S.W.	Q.	—	—	M.fr.IV,117; V,179; VII,145.
A. acidula; F. v. M., fragm. IV, 154 (1864) ...	...	...	...	...	...	...	...	...	—	Q.	—	—	M.fr.IV,154.
A. vestita; F. v. M., fragm. IV, 155 (1864) ...	...	...	...	...	...	...	...	...	—	Q.	—	—	M.fr.IX,104.
A. Endlicheri; Schott, fragm. bot. III, t. 2 ...	...	...	...	...	...	...	...	...	N.S.W.	—	—	—	M.fr.IX,103.
A. tetrandra; F. v. M., fragm. IX, 104 (1875) ...	...	...	...	...	...	...	...	...	—	Q.	—	—	M.fr.V,179.
HALFORDIA; F. v. M., fragm. V, 43 (1865).													
H. drupifera; F. v. M., fragm. V, 43 (1865) ...	...	...	...	...	...	...	...	...	N.S.W.	Q.	—	—	M.fr.V,5,189; VII,142; IX,103.
H. scleroxyla; F. v. M., fragm. VII, 142 (1871) ...	...	...	...	...	...	...	...	...	—	Q.	—	—	M.fr.IX,103.
GLYCOSMIS; Correa in Ann. du Mus. VI, 384 (1805).													
G. pentaphylla; Correa in Annal. du Mus. VI, 384 (1805) ...	...	...	...	...	...	...	...	...	—	Q.	—	B.fl.I,367	M.fr.IX,105. ... As., P.
MICROMELUM; Blume, Bijdrag. 137 (1825).													
M. pubescens; Blume, Bijdrag. 137 (1825) ...	...	...	...	...	...	...	...	...	—	Q.	N.A.	B.fl.I,368	M.fr.I,25; IX,105. ... As., P.
MURRAYA; Koenig in Linné, mant. alt. 563 (1771). (Chaslas, 1767.)													
M. exotica; Koenig in Linné, mant. alt. 563 (1771) ...	...	...	...	...	...	...	...	...	—	Q.	—	B.fl.I,369	M.fr.IX,106. ... As., P.
M. crenulata; Oliver in Journ. Linn. Soc. V, suppl. II, 29 (1861)...	...	...	...	...	...	...	...	...	—	Q.	—	B.fl.I,369	... As.
CLAUSENA; Burmann, Fl. Indica index & t. 29 (1768).													
C. brevistyla; Oliver in Journ. Linn. Soc. V, suppl. II, 31 (1861) ...	...	...	...	...	...	...	...	...	—	Q.	—	B.fl.I,369	M.fr.IX,106.
ATALANTIA; Correa in Ann. du Mus. VI, 383 (1805).													
A. glauca; J. Hooker in B. & H. Gen. pl. I, 305 (1862) ...	...	...	...	...	...	...	...	...	N.S.W.	Q.	—	B.fl.I,370	M.fr.XI,105.
A. recurva; Bentham, Fl. Austr. I, 370 (1863) ...	...	...	...	...	...	...	...	...	—	N.A.	B.fl.I,370	M.fr.VII,142.	

CITRUS; Lioné, gen. pl. 230 (1737), from Plinius.												
C. Planchonii; F. v. M., Report Intercol. Exhib. 23 (1867)	...	...	...	...	...	...	...	N.S.W.	Q.	—	B.fl.I,371	M.fr.IX,105.
C. Anstralasica; F. v. M., fragm. I, 26 (1858)	...	...	...	...	...	...	...	N.S.W.	Q.	—	B.fl.I,371	M.fr.II,178; III,167; IX,105.
PENTACERAS; J. Hooker in B. & H. Gen. pl. I, 298 (1862).												
P. Australis; J. Hooker in B. & H. Gen. pl. I, 298 (1862)	...	...	...	...	...	...	...	N.S.W.	Q.	—	B.fl.I,365	M.fr.I,25; IX,104.

SIMARUBEAE.

DE CANDOLLE in Ann. du Mus. XVII, 323 (1811).—Genera 6, Species 7.

AILANTUS; Desfontaines, in Act. Acad. Paris 265 (1786).												
A. imberbiflora; F. v. M., fragm. III, 42 (1862)	...	...	...	...	...	...	...	N.S.W.	Q.	—	B.fl.I,373	M.fr.VI,166.
BRUCEA; J. S. Miller, icon. plantar. t. 25 (1780).												
B. Sumatrana; Roxburgh, Fl. Indic. I, 469 (1820)	...	...	...	...	...	...	...	—	Q.	N.A.	B.fl.I,373	M.fr.VI,166 ... As.
HYPTIANDRA; J. Hooker in B. & H. Gen. pl. 294 (1862).												
H. Bidwilli; J. Hooker in B. & H. Gen. pl. 294 (1862)	...	...	...	...	...	...	...	N.S.W.	Q.	—	B.fl.I,374	M.fr.VI,165.
CADELLIA; F. v. M., fragm. II, 25 (1860).												
C. pentastylis; F. v. M., fragm. II, 26, t. 12 (1860)	...	...	...	...	...	...	...	N.S.W.	—	—	B.fl.I,374	M.fr.II,171.
C. monostylis; Benth. Fl. Austr. I, 375 (1863)	...	...	...	...	...	...	...	N.S.W.	Q.	—	B.fl.I,375	M.fr.VI,166; VIII,34.
SURIANA; Plumier, nov. plant. Amer. gen. 37, t. 40 (1703).												
S. maritima; Linné, sp. pl. 284 (1753)	...	...	...	...	...	...	...	—	Q.	—	B.fl.I,375	... As., Afr., Am., P.
HARRISONIA; R. Brown in Mém. du Mus. XII, 517 (1825). (Ebelingia.)												
H. Brownii; A. L. de Jussieu in Mém. du Mus. Par. XII, 530, t. 28 (1825)	...	...	...	...	...	...	...	—	Q.	N.A.	B.fl.I,376	... As., P.

ZYGOPHYLLEAE.

R. BROWN in App. Flind. voy. II, 545 (1814)—Genera 3, Species 22.

NITRARIA; Linné, syst. nat. X, 1044 (1759).												
N. Schoberi; Linné, sp. pl. ed. sec. 639 (1762)	...	...	...	...	...	...	...	W.A. S.A.	—	V.	N.S.W.	Q. N.A. B.fl.I,291 M.fr.IX,75; XI,30 ... As., Afr.
ZYGOPHYLLUM; Linné syst. nat. 8 (1735), Linné, gen. 126 (1737). (Sarcozygium, Roepera.)												
Z. apiculatum; F. v. M. in Linnaea XXV, 373 (1852)	...	...	...	...	...	...	...	W.A. S.A.	T.	V.	N.S.W.	Q. — B.fl.I,292 M.fr.XI,29.
Z. glaucescens; F. v. M., pl. of Vict. I, 228 (1862)	...	...	...	...	...	...	...	W.A. S.A.	—	V.	N.S.W.	Q. N.A. B.fl.I,293 M.fr.XI,29.
Z. crenatum; F. v. M. in Linnaea XXV, 374 (1852)	...	...	...	...	...	...	...	W.A. S.A.	—	V.	N.S.W.	Q. — —
Z. iodocarpum; F. v. M. in Linnaea XXV, 372 (1852)	...	...	...	...	...	...	...	W.A. S.A.	—	V.	N.S.W.	Q. N.A. B.fl.I,293 M.fr.XI,29.
Z. prismatothecum; F. v. M. in Linnaea XXV, 375 (1852)	...	...	...	...	...	...	...	—	S.A.	—	—	N.S.W. Q. N.A. B.fl.I,293
Z. ammophilum; F. v. M. in Linnaea XXV, 376 (1852)	...	...	...	...	...	...	...	W.A. S.A.	—	V.	N.S.W.	Q. N.A. — M.fr.XI,28.
Z. Billardieri; De Candolle, prodr. I, 705 (1824)	...	...	...	...	...	...	...	W.A. S.A.	T.	V.	N.S.W.	— — B.fl.I,293 M.fr.XI,29.
Z. fruticosum; De Candolle, prodr. I, 705 (1824)	...	...	...	...	...	...	...	W.A. S.A.	—	V.	N.S.W.	Q. N.A. B.fl.I,594 M.fr.IX,75; XI,29.
Z. Howittii; F. v. M., fragm. III, 150 (1863)	...	...	...	...	...	...	...	—	S.A.	—	—	— — — M.fr.XI,29 ... P.
TRIBULUS; L'Obel, Icon. II, 84 (1581), from Theophrastos, Dioscorides & Plinius). (Tribulopsis.)												
T. terrestris; L'Obel, stirp. icones II, 84 (1581)	...	...	...	...	...	...	...	W.A. S.A.	—	V.	N.S.W.	Q. N.A. B.fl.I,288 M.fr.IX,75; XI,30; XII,4. E., As., Afr., Am.,
T. ranunculiflorus; F. v. M., fragm. I, 48 (1858)	...	...	...	...	...	...	...	—	—	—	—	N.A. B.fl.I,288 M.fr.I,48. [P.]
T. hystrix; R. Brown in App. Sturt. Exped. 6 (1849)	...	...	...	...	...	...	...	W.A. S.A.	—	—	—	N.A. B.fl.I,289 M.fr.XI,30.
T. macrocarpus; F. v. M. in Benth. Fl. Austr. I, 289 (1863)	...	...	...	...	...	...	...	W.A. S.A.	—	—	—	N.A. B.fl.I,289
T. Forrestii; F. v. M. in Wing's S. Sc. rec. Nov. (1885)	...	...	...	...	...	...	...	W.A. S.A.	—	—	—	— — —

T. platypterus ; Bentham, Fl. Austr. I, 289 (1863)	W.A.	—	—	—	—	—	N.A.	B.fl.I,289	
T. hirsutus ; Bentham, Fl. Austr. I, 289 (1863)	W.A.	—	—	—	—	—	N.A.	B.fl.I,289	
T. minutus ; Leichhardt in Benth. Fl. Austr. I, 291 (1863)... ..	—	—	—	—	—	N.S.W.	Q.	B.fl.I,241	
T. Brownii ; F. v. M., pl. of Vict. I, 99 (1860)	—	—	—	—	—	—	Q.	N.A. B.fl.I,290	M.fr.I,48.
T. bicolor ; F. v. M., pl. of Vict. I, 99 (1860)	—	—	—	—	—	—	N.A.	B.fl.I,290	M.fr.I,47.
T. Solandri ; F. v. M., pl. of Vict. I, 99 (1860)	—	—	—	—	—	—	Q.	N.A. B.fl.I,290	M.fr.I,47 ; XI,30.
T. astrocarpus ; F. v. M., fragm. XII, 4 (1882)	W.A.	S.A.	—	—	—	—	—	—	

LINEAE.

DE CANDOLLE, Théor. élém. bot. 214 (1813).—Genera 3, Species 4.

HUGONIA ; Linné, gen. pl. 134 (1737).

H. Jenkinsii ; F. v. M., fragm. V, 7 (1865) Q. — M.fr.IX,190.

ERYTHROXYLUM ; P. Brown, Hist. pl. of Jamaica 278 (1756), from Theophrastos, Dioscorides & Plinius. (Erythroxyylon.)

E. Australe ; F. v. M. in Transact. Vict. Inst. III, 22 (1855) Q. — B.fl.I,284

E. ellipticum ; R. Brown in Benth. Fl. Austr. I, 284 (1863) — N.A. B.fl.I,284

LINUM ; Tournefort, inst. 339, t. 176 (1700), from Theophrastos, Dioscorides & Plinius.

L. marginale ; Cunningham in Hook. Lond. Jour. Bot. VII, 169 (1848) ... W.A. S.A. T. V. N.S.W. Q. — B.fl.I,283 M.fr.XI,27.

GERANIACEAE.

A. L. DE JUSSIEU, Gen. 268 (1789), from B. de Jussieu (1759).—Genera 5, Species 8.

GERANIUM ; Tournefort, inst. 266, t. 142 (1700), from Dioscorides and Plinius.

G. pilosum ; Solander in G. Forst., Fl. Ins. Austr. prodr. 91 (1786) W.A. S.A. T. V. N.S.W. Q. — B.fl.I,296 M.fr.XI,27 ... Am., N.Z.

G. sessiliflorum ; Cavanilles, Diss. 198, t. 77 (1790) — T. V. N.S.W. — — B.fl.I,297 ... Am., N.Z.

ERODIUM ; L. Héritier, Geraniologia, t. 7 (1787).

E. cygnorum ; Nees in Lehm. pl. Preiss. I, 162 (1844) W.A. S.A. — V. N.S.W. Q. N.A. B.fl.I,297 M.fr.IX,75 ; XI,20.

PELARGONIUM ; L. Héritier, Geraniologia, t. 1 (1787).

P. Australe ; Willdenow, sp. pl. III, 675 (1800) W.A. S.A. T. V. N.S.W. Q. — B.fl.I,298 M.fr.IX,77 ... Afr., N.Z.

P. Rodueyanum ; Mitchell, Three Exped. II, 144 (1838) W.A. S.A. — V. N.S.W. — — B.fl.I,299

OXALIS ; Linné, gen. pl. 134 (1737), from Plinius.

O. Magellanica ; G. Forster in Comm. Goetting. IX, 33 (1789) — T. V. — — — B.fl.I,300 ... Am., N.Z.

O. corniculata ; Linné, sp. pl. 435 (1753) W.A. S.A. T. V. N.S.W. Q. — B.fl.I,301 M.fr.IX,75,77 E., As., Afr., Am., P., N.Z.

BIOPHYTUM ; De Candolle, prodr. I, 689 (1824).

B. apodoscias ; Turczaninow in Bullet. des Nat. de Mosc. XXXVI, 595 (1863) — — — — — Q. — —

MALVACEAE.

ADANSON in Mém. Acad. fr. 224 (1761).—Genera 15, Species 110.

LAVATERA ; Tournefort in Act. Acad. Par. 86, t. 3 (1706).

L. plebeja ; Sinus in Bot. Mag. t. 2269 (1821) W.A. S.A. T. V. N.S.W. Q. N.A. B.fl.I,185 M.fr.IX,75,130 ; X,53.

MALVASTRUM; Asa Gray in Mem. Amer. Acad. IV, 21 (1849). (Malva partly.)

M. spicatum; A. Gray in Mem. Amer. Acad. IV, 21 (1849)	W.A. S.A. — —	N.S.W. Q. N.A. B.fl.I, 187	M.fr.VI, 170; IX, 130	As., Afr., Am., P.
M. tricuspidatum; A. Gray in Mem. Amer. Acad. IV, 21 (1849)	— — — —	N.S.W. Q. — B.fl.I, 187		As., Afr., Am., P.

PLAGIANTHUS; R. & G. Forster, char. gen. 85, t. 43 (1776). (Asterotrichon, Blepharanthemum, Lawrenceia, Halothamnus, Gynothrix.)

P. sidoides; Hooker, Bot. Mag. t. 3396 (1835)	— — T. — —	— — B.fl.I, 188	M.fr.VI, 169.	
P. pulchellus; A. Gray, Bot. Amer. Expl. Exped. I, 181 (1854)	— — T. V. N.S.W. — —	B.fl.I, 189	M.fr.IX, 130.	
P. spicatus; Benth in proceed. Linn. Soc. VI, 103 (1862)	W.A. S.A. T. V. N.S.W. — —	B.fl.I, 189	M.fr.VI, 170; IX, 130.	
P. glomeratus; Benth in proceed. Linn. Soc. VI, 103 (1862)	W.A. S.A. — —	N.S.W. Q. — B.fl.I, 190	M.fr.IX, 130.	
P. diffusus; Benth, Fl. Austr. I, 190 (1863)	W.A. — — — —	N.A. B.fl.I, 190		
P. microphyllus; F. v. M., fragm. I, 29 (1858)	W.A. S.A. — V. N.S.W. — —	B.fl.I, 190	M.fr.VI, 170; IX, 13.	
P. Berthae; F. v. M., fragm. V, 103 (1866)	W.A. S.A. — — — —	— — — —	M.fr.V. 114.	

SIDA; Linné, gen. pl. 205 (1737) from Theophrastos.

S. corrugata; Lindley in Mitch. Three Exped. II, 13 (1838)	W.A. S.A. — V. N.S.W. Q. N.A. B.fl.I, 192	M.fr.IX, 75.		
S. Spenceriana; F. v. M. in Wing's S. Sc. Record, April 1885	— — — — N.S.W. Q. —			
S. intricata; F. v. M. in Transact. phil. Soc. Vict. I, 19 (1860)	W.A. S.A. — V. N.S.W. Q. N.A. B.fl.I, 193			
S. macropoda; F. v. M. in Benth. Fl. Austr. I, 193 (1863)	— — — — —	N.A. B.fl.I, 193		
S. virgata; Hooker in Mitch. Trop. Austr. 361 (1848)	W.A. S.A. — — N.S.W. Q. N.A. B.fl.I, 194			
S. echinocarpa; F. v. M., fragm. XI, 62 (1879)	— — — — —	N.A. —	M.fr.XI, 62.	
S. cryphiopetala; F. v. M., fragm. III, 4 (1862)	W.A. S.A. — — — —	Q. N.A. B.fl.I, 194	M.fr.III, 4.	
S. petrophila; F. v. M. in Linnaea XXV, 381 (1852)	W.A. S.A. — — N.S.W. Q. N.A. B.fl.I, 194			
S. Kiogii; F. v. M. in Vict. Natur. III, 138 (1887)	W.A. — — — — —	— — — —		
S. calyxhymentia; J. Gay in De Candolle, prodr. I, 462 (1824)	W.A. S.A. — — — —	N.A. B.fl.I, 194	M.fr.IX, 75.	
S. physocalyx; F. v. M., fragm. III, 3 (1862)	W.A. — — — — —	N.A. B.fl.I, 195	M.fr.X, 73.	
S. subspicata; F. v. M. in Benth. Fl. Austr. I, 195 (1863)	— — — — N.S.W. Q. N.A. B.fl.I, 195			
S. pleiantha; F. v. M. in Benth. Fl. Austr. I, 195 (1863)	— — — — —	Q. — B.fl.I, 195	M.fr.VI, 169.	
S. spinosa; Linné, sp. pl. 683 (1753)	— — — — —	Q. N.A. B.fl.I, 196		As., Afr., Am., P.
S. rhombifolia; Linné, sp. pl. 684 (1753)	— S.A. — — N.S.W. Q. N.A. B.fl.I, 196			As., Afr., Am., P.
S. cordifolia; Linné, sp. pl. 684 (1753)	— — — — —	Q. N.A. B.fl.I, 196		As., Afr., Am., P.
S. platycalyx; F. v. M. in Benth. Fl. Austr. I, 197 (1863)	— S.A. — — N.S.W. Q. N.A. B.fl.I, 197	M.fr.X, 73.		
S. cleisocalyx; F. v. M., fragm. X, 73 (1876)	— — — — —	N.A. —	M.fr.X, 73.	
S. inclusa; Benth, Fl. Austr. I, 197 (1863)	— S.A. — — — —	N.A. B.fl.I, 197	M.fr.IX, 181; XI, 32.	
S. cardiophylla; F. v. M., fragm. VIII, 242 (1874)	W.A. S.A. — — — —	N.A. —	M.fr.VIII, 242.	
S. Hookeriana; Miquel in Lehm. pl. Preiss. I, 242 (1844)	W.A. — — — — —	B.fl.I, 197		
S. lepida; F. v. M., fragm. VI, 168 (1868)	W.A. S.A. — — — —	N.A. —	M.fr.XI, 32.	

ABUTILON; Tournefort, inst. 99, t. 25 (1700) from Camerarius (1586). (Fleischera, Abutilaea.)

A. tubulosum; Hooker in Mitch. Trop. Austr. 300 (1848)	— S.A. — — N.S.W. Q. N.A. B.fl.I, 200	M.fr.II, 12; IX, 131.		
A. amplum; Benth, Fl. Austr. I, 200 (1863)	— — — — —	N.A. B.fl.I, 200		
A. leucopetalum; F. v. M. in Benth. Fl. Austr. I, 200 (1863)	— S.A. — — N.S.W. Q. N.A. B.fl.I, 200	M.fr.II, 12.		
A. Mitchellii; Benth, Fl. Austr. I, 201 (1863)	— S.A. — — — —	Q. — B.fl.I, 201		
A. micropetalum; F. v. M. in Benth. Fl. Austr. I, 201 (1863)	— — — — —	Q. — B.fl.I, 201	M.fr.VI, 169.; IX, 131; XI, 32; M.fr.IX, 131.	
A. cryptopetalum; F. v. M. in Benth. Fl. Austr. I, 201 (1863)	W.A. S.A. — — N.S.W. — N.A. B.fl.I, 201	M.fr.II, 11; IX, 13.		
A. geranioides; Benth, Fl. Austr. I, 202 (1863)	— — — — —	N.A. B.fl.I, 202	M.fr.VI, 168.	
A. otocarpum; F. v. M. in Transact. phil. Soc. Vic. I, 13 (1854)	W.A. S.A. — — N.S.W. Q. N.A. B.fl.I, 202	M.fr.XI, 32.		
A. longilobum; F. v. M., fragm. IX, 130 (1875)	W.A. — — — — —	N.A. —	M.fr.IX, 130.	
A. subviscosum; Benth, Fl. Austr. I, 202 (1863)	— — — — —	Q. — B.fl.I, 202		
A. Indicum; J. Bauhin, hist. plant. univ. II, 959 (1651)	— — — — —	Q. N.A. B.fl.I, 202		As., Afr.
A. auritum; G. Don, gen. syst. I, 500 (1831)	— — — — N.S.W. Q. — B.fl.I, 203	M.fr.XI, 32		As., Afr., P.

A. Avicennae; Gerard, gen. hist. pl. 790 (1597)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.I,203	...	...	...	E., As., Am.
A. graveolens; Wight & Arnott, Prodr. Fl. Pen. Ind. or. I, 56 (1840)	—	—	—	—	—	—	—	Q. — B.fl.I,204	...	...	...	As., Afr.
A. oxycarpum; F. v. M. in Benth. Fl. Austr. I, 204 (1863)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.I,204	M.fr.II,12; IX,131; XI,32.			
A. mnticum; G. Don, gen. syst. I, 502 (1831)	—	—	—	—	—	Q.	—	B.fl.I,204	...	...	...	As., Afr.
A. Cunninghamii; Beatham, Fl. Austr. I, 205 (1863)	—	—	—	—	—	Q.	N.A.	B.fl.I,205				
A. exonemum; F. v. M., fragm. XI, 63 (1879)	—	—	—	—	—	—	N.A.	—	M.fr.XI,63.			
A. Fraseri; Hooker in Walpera, annal. II, 158 (1851)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.I,205	M.fr.II,11.			
A. halophilum; F. v. M. in Linnaea XXV, 381 (1852)	W.A.	S.A.	—	—	N.S.W.	Q.	—	B.fl.I,206	M.fr.XI,27.			
A. macrum; F. v. M., fragm. IX, 59 (1875)	—	S.A.	—	—	—	—	—	—	M.fr.IX,59.			
A. crispum; G. Don, gen. syst. I, 502 (1831)	—	—	—	—	—	—	N.A.	B.fl.I,206	M.fr.VI,169	...	...	As., Afr., Am., P.
A. Julianae; Endl. prodr. fl. ina. Norf. 75 (1833)	—	—	—	—	N.S.W.	—	—	—				
URENA; Dillenius, Hort. Eltham. 430, t. 319 (1732).												
U. lobata; Linné, sp. pl. 692 (1753)	—	—	—	—	—	Q.	N.A.	B.fl.I,206	M.fr.VII,69; IX,130	...	...	As., Afr., Am., P.
U. Armitiana; F. v. M., fragm. X, 78 (1877)	—	—	—	—	—	Q.	N.A.	—	M.fr.X,78.			
PAVONIA; Cavanilles, monadelph. disaert. II, App. II (1786). (Greevesia.)												
P. hastata; Cavanilles, dissert. III, 138, t. 47 (1786)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,207	M.fr.IX,130; X,73	...	...	As., Afr., Am.
HOWITTIA; F. v. M. in Transact. Vict. Inst. I, 116 (1855).												
H. trilocularis; F. v. M. in Transact. Vict. Inst. I, 116 (1855)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.I,198	M.fr.VI,169.			
HIBISCUS; Linné, syst. nat. (1735); gen. pl. 207 (1737). (Fugosia partly, Abelmoschus, Paritium.)												
H. ficulnea; Linné, sp. pl. 605 (1753)	—	—	—	—	—	Q.	N.A.	B.fl.I,209	M.fr.I,67.	...	...	As.
H. Abelmoschua; Linné, sp. pl. 696 (1753)	—	—	—	—	—	—	N.A.	—	M.fr.VI,169	...	...	As., Afr., P.
H. rhodopetalus; F. v. M. in Benth. Fl. Austr. I, 209 (1863)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.I,209	M.fr.II,113; V,210; IX,229; XI,31.			
H. Manihot; Linné, sp. pl. 696 (1753)	—	—	—	—	—	Q.	—	B.fl.I,210	...	...	...	As.
H. Notho-Manihot; F. v. M., fragm. V, 57 (1865)	—	—	—	—	—	Q.	—	—	M.fr.V,213; IX,130.			
H. Trionum; Linné, sp. pl. 697 (1753)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.I,210	M.fr.II,115; IX,129; XI,31.	E., As., Afr., N.Z.		
H. insularia; Endlicher, prodr. fl. ina. Norf. 74 (1833)	—	—	—	—	N.S.W.	—	—	—				
H. brachysiphonia; F. v. M., fragm. I, 67 (1858)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.I,210	M.fr.I,220,243; VI,169.			
H. Drummondii; Turczaninow in Bull. Soc. Mosc. XXXI, 195 (1858)	W.A.	—	—	—	—	—	—	B.fl.I,211	M.fr.I,220; IX,129; XI,31.			
H. microlaenus; F. v. M., fragm. II, 116 (1861)	W.A.	S.A.	—	—	—	—	N.A.	B.fl.I,211	M.fr.III,5; IX,129.			
H. Pinonians; Gaudichaud in Freyc. voy. Bot. 476, t. 100 (1826)	W.A.	S.A.	—	—	—	—	N.A.	B.fl.I,211	M.fr.II,116.			[As., Afr.]
H. cannabinus; Linné, sp. pl. ed. sec., 979 (1763)	—	—	—	—	—	Q.	N.A.	B.fl.I,212	M.fr.II,117; III,166; VI,169; IX,129; XI,31			
H. divaricata; Graham in Edinb. phil. Journ. (1830)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,212	M.fr.II,118; IX,129; XI,31.			
H. Fitzgeraldii; F. v. M., fragm. VIII, 242 (1874)	—	—	—	—	—	Q.	—	—	M.fr.VIII,242.			
H. Elaworthii; F. v. M., fragm. VIII, 241 (1874)	—	—	—	—	—	Q.	—	—	M.fr.VIII,241.			
H. Haynaldi; F. v. M. in Wing's S. Sc. Rec. March (1883)	W.A.	—	—	—	—	—	—	—				
H. heterophyllus; Ventenat, jard. de la Malm., t. 103 (1804)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,212	M.fr.VI,170; XI,31.			
H. diversifolia; N. Jacquin, icon. pl. rar. 551 (1789)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,213	M.fr.II,117; VI,169	...	...	As., Afr., P.
H. splendens; Fraser in Edinb. phil. Journ. (1830)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,213	M.fr.VI,170; XI,31&123.			
H. zonatus; F. v. M., fragm. I, 221 (1859)	—	—	—	—	—	—	N.A.	B.fl.I,213	M.fr.VI,170.			
H. Coatesii; F. v. M., fragm. III, 5 (1862)	W.A.	—	—	—	—	—	N.A.	B.fl.I,214	M.fr.III,5.			
H. leptocladus; Benth. Fl. Austr. I, 214 (1863)	—	—	—	—	—	—	N.A.	B.fl.I,214	M.fr.VI,169; IX,129.			
H. setulosus; F. v. M., fragm. I, 221 (1859)	—	—	—	—	—	—	N.A.	B.fl.I,214	M.fr.IX,31.			
H. Goldaworthii; F. v. M., fragm. XI, 30 (1878)	—	—	—	—	—	—	N.A.	—	M.fr.IX,30.			
H. pentaphyllus; F. v. M., fragm. II, 13 (1860)	—	—	—	—	—	—	N.A.	B.fl.I,214	M.fr.VI,169; XI,32.	...	...	As., Afr.
H. geranioides; Cunningham in Benth. Fl. Austr. I, 215 (1863)	—	—	—	—	—	—	N.A.	B.fl.I,215				
H. vitifolia; Linné, sp. pl. 696 (1753)	—	—	—	—	—	Q.	—	B.fl.I,215	M.fr.I,220; II,114; VI,169.			As., Afr.
H. panduriformis; N. Burmann, Fl. Ind. 151, t. 47 (1768)	W.A.	—	—	—	—	Q.	N.A.	B.fl.I,215	M.fr.II,115; VI,169; IX,129.			As., Afr.

H. Normani; F. v. M., fragm. III, 4 (1862)	...	...	...	...	...	—	—	—	—	Q.	—	B.fl.I,216	M.fr.V,44; VI,169; IX,128.		
H. Krichauffii; F. v. M., Rep. Babb. Exped. 7 (1858)	...	...	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.I,216	M.fr.VI,169; IX,75,129.	
H. Farragci; F. v. M., fragm. VIII, 241 (1874)	...	...	...	...	...	...	W.A.	S.A.	—	N.S.W.	—	N.A.	—	M.fr.VIII,284; IX,129&196; XI,32,134.	
H. Sturtii; Hooker in Mitch. trop. Austr. 363 (1848)	...	...	...	...	...	...	W.A.	S.A.	—	N.S.W.	Q.	N.A.	B.fl.I,216	M.fr.II,13; IX,128.	
H. phyllochaenus; F. v. M., fragm. IX, 128 (1875)	...	...	...	...	...	...	—	—	—	—	Q.	—	—	M.fr.IX,129.	
H. Huegelii; Endlicher in Hueg. enum. 10 (1837)	...	...	...	...	...	...	W.A.	S.A.	—	—	—	—	B.fl.I,217	M.fr.IX,128.	
H. Wrayae; Lindley, bot. Regist. t. 69 (1840)	...	...	...	...	...	...	W.A.	S.A.	—	—	—	—	—	M.fr.IX,128.	
H. tiliaceus; Linné, sp. pl. 694 (1753)	...	...	...	...	...	...	—	—	—	N.S.W.	Q.	N.A.	B.fl.I,218	M.fr.VI,170; IX,129; XI,32.	As., Afr.,
H. cuneiformis; De Candolle, prodr. I, 454 (1824)	...	...	...	...	...	...	W.A.	—	—	—	—	N.A.	B.fl.I,219	M.fr.XI,32.	[Am., P.
H. hakeaefolius; Giordano, Mém. di nuov. sp. d'Ibisco (1833)	...	...	...	...	...	...	W.A.	S.A.	—	—	—	—	B.fl.I,220	M.fr.IX,128.	

LAGUNARIA; G. Don, gen. syst. dichlam. pl. I, 485 (1831).

L. Patersoni; G. Don, gen. syst. I, 485 (1831) ...	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,218	M.fr.IX,130; XI,32.
--	-----	-----	-----	-----	-----	---	---	---	---	--------	----	---	------------	---------------------

THESPESIA; Solander in Ann. du. Mus. IX, 290 (1807).

T. populnea; Solander in Ann. du. Mus. IX, 290, t. 8 (1807) ...	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.I,221	M.fr.VI,170; IX,130; XI,32. As., Afr., P.
---	-----	-----	-----	-----	-----	---	---	---	---	---	----	------	------------	---

GOSSYPIUM; Linné, gen. pl. 206 (1737) from Camerarius (1586). (Fugosia partly, Sturtia.)

G. Sturtii; F. v. M., fragm. III, 6 (1862) ...	...	...	...	...	...	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.I,222	M.fr.IV,174; VI,169&251; IX,127; XI,63.
G. Robinsoni; F. v. M., fragm. IX, 126 (1875) ...	...	...	...	...	...	W.A.	—	—	—	—	—	N.A.	—	M.fr.XI,63&135.
G. Australe; F. v. M., fragm. I, 46 (1858) ...	...	...	...	...	...	W.A.	S.A.	—	—	—	Q.	N.A.	B.fl.I,220	M.fr.III,6; V,44; IX,127; XI,32&63.
G. thespesioides; F. v. M., fragm. IX, 127 (1875) ...	...	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.I,220	M.fr.V,44&177; VI,169; XI,32&63.
G. Cunninghamii; Todaro, Monogr. Gen. Gossyp. 7 (1878) ...	...	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.I,220	M.fr.VI,169; IX,129; XI,32&63.
G. costulatum; Todaro, Monogr. Gen. Gossyp. 7 (1878) ...	...	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.I,221	M.fr.IX,127; XI,32&63.
G. flaviflorum; F. v. M., fragm. IX, 127 (1875) ...	...	...	...	...	...	—	—	—	—	—	—	N.A.	—	M.fr.V,44; XI,63.
G. populifolium; F. v. M., fragm. IX, 127 (1875) ...	...	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.I,221	M.fr.XI,32&63.

ADANSONIA; Linné, sp. pl. I, 1190 (1753).

A. Gregorii; F. v. M. in Hook. Kew misc. IX, 14 (1857) ...	...	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.I,223	M.fr.VII,40; XI,32.
--	-----	-----	-----	-----	-----	---	---	---	---	---	---	------	------------	---------------------

CAMPTOSTEMON; Masters in J. Hook. icon. pl. XII, 18, t. 1119 (1876).

C. Schultzii; Masters in J. Hook. icon. pl. XII, 18, t. 1119 (1876) ...	...	...	...	...	...	—	—	—	—	—	—	N.A.	—	M.fr.IX,43; XI,32.
---	-----	-----	-----	-----	-----	---	---	---	---	---	---	------	---	--------------------

BOMBAX; Linné, sp. pl. I, 511 (1753).

B. Malabarica; De Candolle, prodr. I, 479 (1824) ...	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.I,223	M.fr.XI,32 ... As.
--	-----	-----	-----	-----	-----	---	---	---	---	---	----	------	------------	--------------------

STERCULIACEAE.

VENTENAT; Jard. de la Malm. II, 91 (1804).—Genera 20, Species 125.

STERCULIA; Linné, Fl. Zeyl. 166 (1747).

S. quadrifida; R. Brown in Benn. pl. Jav. rar. 233 (1852) ...	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.I,227	M.fr.VI,173.
S. laurifolia; F. v. M., fragm. VI, 172 (1868) ...	...	...	...	...	...	—	—	—	—	—	Q.	—	—	M.fr.VI,172.

BRACHYCHITON; Schott & Endlicher, Melet. bot. 34 (1832). (Trichosiphon, Poeciloderms, Delabechea.)

B. paradoxus; Schott, Meletem. 34 (1832) ...	...	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.I,227	
B. Bidwilli; Hooker, Bot. Mag. t. 5133 (1859) ...	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl.I,228	
B. discolor; F. v. M., fragm. I, 1 (1858) ...	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.I,228	M.fr.I,242; II,177.
B. incanus; R. Brown in Benn. pl. Jav. rar. 234 (1852) ...	...	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.I,228	
B. luridus; C. Moore in F. v. M., fragm. I, 1 (1858) ...	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,228	M.fr.II,177.
B. platanoides; R. Brown in Benn. pl. Jav. rar. 234 (1852) ...	...	...	...	...	...	W.A.	—	—	—	—	Q.	N.A.	B.fl.I,229	M.fr.VI,173.
B. acerifolius; F. v. M., fragm. I, 1 (1858) ...	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,229	M.fr.II,177; V,209; VI,173.

B. diversifolius; R. Brown in Benn. pl. Jav. rar. 234 (1852)	...	...	...	W.A.	—	—	—	—	Q. N.A.	B.fl.I,230	
B. populneus; B. Brown in Benn. pl. Jav. rar. 234 (1852)	...	...	...	—	—	—	—	V. N.S.W.	Q. —	B.fl.I,229	M.fr.IV,137.
B. populneo-acerifolius; F. v. M. in proceed. L.S. of N.S.W. IX, 380 (1884)	...	...	...	—	—	—	—	N.S.W.	—	—	
B. Delabechii; F. v. M., pl. of Vict. I, 157 (1860)	...	...	...	...	—	—	—	—	Q. —	B.fl.I,230	M.fr.IV,137.
B. Gregorii; F. v. M. in Hook. Kew misc. IX, 199 (1857)...	...	...	...	W.A.	S.A.	—	—	—	—	—	M.fr.IV,137; V,212; X,53.
TARRIETIA; Blume, Bijdr. 227 (1825). (Argyrodendron.)											
T. Argyrodendron; Benth. Fl. Austr. I, 230 (1863)	...	...	...	...	—	—	—	—	N.S.W.	Q. —	B.fl.I,230 M.fr.VI,173; IX,42.
T. trifoliolata; F. v. M., fragm. IX, 43 (1875)	...	...	...	...	—	—	—	—	N.S.W.	Q. —	M.fr.I,2; II,177.
HERITIERA; Dryander in Aiton, Hort. Kew III, 546 (1789).											
H. litoralis; Dryander in Aiton, Hort. Kew III, 546 (1789)	...	...	...	...	—	—	—	—	Q. N.A.	B.fl.I,231	M.fr.VI,173 ... As., Afr., P.
UNGHERIA; Schott & Endlicher, Melet. 27—31 (1832).											
U. floribunda; Schott & Endlicher, Melet. 27, t. 4 (1832)	...	...	...	...	—	—	—	—	N.S.W.	—	M.fr.IX,169.
HELICTERES; Plukenet, Phytogr. 181, t. 245 (1692).											
H. Isora; Linné, sp. pl. 963 (1753)	...	...	...	...	—	—	—	—	—	N.A.	B.fl.I,232 M.fr.VI,173 ... As.
METHORIUM; Schott & Endlicher, Melet. 29—30 (1832).											
M. spicatum; F. v. M., syst. Cens. of Austr. pl. 15 (1882) from Colebr. (1831)	...	...	...	...	—	—	—	—	Q. —	—	M.fr.VII,39 ... As.
M. canum; Schott, Meletem. 29, t. 5 (1832)	...	...	...	...	—	—	—	—	—	N.A.	B.fl.I,232
M. dentatum; F. v. M. in Benth. Fl. Austr. I, 232 (1863)...	...	...	...	...	—	—	—	—	—	N.A.	B.fl.I,232
M. semiglabrum; F. v. M., fragm. V, 43 (1865)	...	...	...	...	—	—	—	—	Q. —	—	M.fr.V,43.
MELHANIA; Forskael, Fl. Aeg. Arab. 64 (1775).											
M. incana; Heyne in Wall. numerit. list 1200 (1828)	...	...	...	...	—	S.A.	—	—	N.S.W.	Q. N.A.	B.fl.I,234 M.fr.I,69; VI,174 As., Afr., P.
MELOCHIA; Dillenius, Hort. Eltham. 221, t. 176 (1732). (Riedleya.)											
M. pyramidata; Linné, sp. pl. 674 (1753)	...	...	...	...	—	—	—	—	Q. N.A.	B.fl.I,234	M.fr.VI,173. ... As., Afr., Am., P.
M. corchorifolia; Linné, sp. pl. 675 (1753)	...	...	...	...	—	—	—	—	Q. N.A.	B.fl.I,235	M.fr.VI,173. ... As., Afr.
HERMANNIA; Tournefort, inst. 656, t. 432 (1700). (Gilesia, Mahernia partly.)											
H. Gilesii; F. v. M., fragm. IX, 42 (1875)	...	...	...	...	—	S.A.	—	—	—	—	M.fr.IX,42.
DICARPIDIUM; F. v. M. in Hook. Kew Misc. IX, 302 (1857).											
D. monoicum; F. v. M. in Hook. Kew misc. IX, 302 (1857)	...	...	...	...	—	—	—	—	—	N.A.	B.fl.I,235
WALTHERIA; Linné, gen. pl. 203 (1737).											
W. Indica; Linné, sp. pl. 637 (1753)	...	...	...	...	...	W.A.	S.A.	—	—	Q. N.A.	B.fl.I,236 M.fr.VI,173. ... As., Afr. Am., P.
ABROMA; N. Jacquin, Hort. Vindob. III, t. 1 (1776).											
A. fastuosa; Salisbury, Parad. Lond. 102 (1807)	...	...	...	...	...	—	—	—	—	Q. —	B.fl.I,236 M.fr.VI,173 ... As.
COMMERCONIA; R. & G. Forster, char. gen. 43, t. 22 (1776). (Ruelingia, Achilleopsis.)											
C. salvifolia; Benth. Fl. Austr. I, 238 (1863)	...	...	...	...	...	—	—	—	—	Q. —	B.fl.I,238
C. dasyphylla; Andrews, Bot. Reposit. t. 603 (1804)	...	...	...	...	...	—	—	—	V. N.S.W.	Q. —	B.fl.I,238 M.fr.X,22.
C. rugosa; F. v. M., syst. Cens. of Austr. pl. 16 (1882)	...	...	...	...	...	—	—	—	N.S.W.	—	B.fl.I,238
C. Preissii; Steudel in Lehm. pl. Preiss. I, 237 (1844)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.I,339
C. cinerea; Steudel in Lehm. pl. Preiss. I, 237 (1844)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.I,239
C. cygnorum; Steudel in Lehm. pl. in Preiss. I, 238 (1844)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.I,239
C. platycalyx; F. v. M., syst. Cens. of Austr. pl. 16 (1882) from Benth. (1863)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.I,240
C. parviflora; F. v. M., syst. Cens. of Austr. pl. 16 (1882) from Endl. (1837)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.I,240
C. hermanniaefolia; F. v. M., syst. Cens. of Austr. pl. 16 (1882) from Steetz (1847)	...	...	...	...	—	—	—	—	N.S.W.	—	B.fl.I,240
C. Kempeana; F. v. M., fragm. IX, 113 (1881)	...	...	...	...	...	—	S.A.	—	—	—	M.fr.XI,113.
C. loxophylla; F. v. M., fragm. X., 22 (1876)	...	...	...	...	...	—	S.A.	—	—	—	N.A. B.fl.I,240 M.fr.I,68

C. magniflora; F. v. M., fragm. VIII, 223 (1874) from Turcz. (1852)	...	...	...	...	S.A.	—	—	—	—	—	—	M.fr.X,22; XI,114.
C. cuneata; F. v. M., syst. Cens. of Austr. pl. 16 (1882) from Turcz. (1852)	...	W.A.	—	—	—	—	—	—	—	—	B.fl.I,241	
C. rotundifolia; F. v. M., syst. Cens. of Austr. pl. 16 (1882)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.I,241	
C. densiflora; F. v. M., fragm. X, 21 (1876)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.I,241	M.fr.VI,175; X,22; XI,114.
C. Fraseri; J. Gay in Mém. du Mus. Par. X, 215, t. 15 (1823)	...	...	...	...	—	—	—	V.	N.S.W.	Q.	—	B.fl.I,242
C. Leichhardtii; Benthām, Fl. Austr. I, 242 (1863)	...	...	...	...	—	—	—	—	—	Q.	—	B.fl.I,242
C. echinata; R. & G. Forster, char. gen. 43, t. 22 (1776)	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,243
C. Gaudichaudi; J. Gay in Mém. du Mus. Par. X, 213, t. 14 (1823)	...	...	W.A.	—	—	—	—	—	—	—	—	B.fl.I,243
C. crispa; Turczaninow in Bull. Soc. Mosc. XIX, 501 (1846)	...	...	W.A.	—	—	—	—	—	—	—	—	B.fl.I,243
C. melanopetala; F. v. M., fragm. X, 21 (1876)	...	...	W.A.	—	—	—	—	—	—	—	—	M.fr.X,21.
C. pulchella; Turczaninow in Bull. Soc. Mosc. XIX, 502 (1846)	...	...	W.A.	—	—	—	—	—	—	—	—	B.fl.I,244
C. microphylla; Benthām, Fl. Austr. I, 244 (1863)	...	...	W.A.	—	—	—	—	—	—	—	—	B.fl.I,244
C. cratrophylla; F. v. M., fragm. IX, 59 (1875)	...	...	W.A.	—	—	—	—	—	—	—	—	M.fr.IX,59.
C. Tatei; F. v. M. in Trans. Roy. Soc. of S. Austr. X (1888)	...	...	—	S.A.	—	—	—	—	—	—	—	
HANNAFORDIA; F. v. M., fragm. II, 9 (1860).												
H. quadrivalvis; F. v. M., fragm. II, 9 (1860)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.I,248
H. Shanesii; F. v. M., fragm. VI, 175 (1868)	...	...	...	...	—	—	—	—	—	Q.	—	M.fr.X,96.
H. Bissillii; F. v. M., fragm. X, 95 (1877)	...	...	...	...	W.A.	S.A.	—	—	N.S.W.	—	—	M.fr.XI,115.
SERINGEA; Sprengel, Anleit. II, 649 (1818). (Keraudrenia.)												
S. platyphylla; J. Gay in Mém. du Mus. Par. VII, 443 (1821)	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,244
S. lanceolata; Steetz in Lehm. pl. Preiss. II, 349 (1847)	...	...	...	...	—	—	—	—	—	Q.	—	B.fl.I,245
S. corollata; Steetz in Lehm. pl. Preiss. II, 330 (1847)	...	...	...	...	—	S.A.	—	—	—	Q.	N.A.	B.fl.I,246
S. Hillii; F. v. M., syst. Cens. of Austr. pl. 16 (1882)	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,246
S. nephrosperma; F. v. M. in Hook. Kew misc. IX, 15 (1857)	...	...	...	...	—	S.A.	—	—	—	N.A.	B.fl.I,246	M.fr.I,142; II,5; X,97.
S. microphylla; F. v. M., fragm. II, 5 (1860)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.I,247
S. adenolasia; F. v. M., fragm. X, 96 (1877)	...	...	...	...	—	—	—	—	—	Q.	N.A.	—
S. integrifolia; F. v. M., fragm. II, 5 (1860)	...	...	...	...	W.A.	S.A.	—	—	—	—	N.A.	B.fl.I,247
THOMASIA; J. Gay in Mém. du Mus. VII, t. 6—7 (1821). (Leucothamnus, Rhynchostemon.)												
T. macrocarpa; Huegel in Endlicher, nov. stirp. dec. 32 (1839)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.I,250
T. rugosa; Turczaninow in Bull. Soc. Mosc. XIX, 501 (1846)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.I,250
T. montana; Steudel in Lehm. pl. Preiss. I, 230 (1844)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.I,250
T. tenuivestita; F. v. M., fragm. II, 7 (1860)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.I,251
T. solanacea; J. Gay in Mém. du Mus. VII, 456, t. 21 (1821)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.I,251
T. brachystachys; Turczaninow in Bull. Soc. Mosc. XXV, 143 (1852)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.I,251
T. discolor; Steudel in Lehm. pl. Preiss. I, 233 (1844)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.I,251
T. quercifolia; J. Gay in Mém. du Mus. VII, 459, t. 21 (1821)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.I,252
T. foliosa; J. Gay in Mém. du Mus. VII, 454, t. 22 (1821)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.I,252
T. triloba; Turczaninow in Bull. Soc. Mosc. XIX, 500 (1846)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.I,252
T. triphylla; J. Gay in Mém. du Mus. VII, 458 (1821)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.I,252
T. purpurea; J. Gay in Mém. du Mus. VII, 452, t. 21 (1821)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.I,253
T. macrocalyx; Steudel in Lehm. pl. Preiss. I, 230 (1844)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.I,253
T. pauciflora; Lindley Bot. Regist. XXV, App. XVII (1839)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.I,253
T. rhynchocharpa; Turczaninow in Bull. Soc. Mosc. XXV, 142 (1852)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.I,254
T. grandiflora; Lindley, Bot. Regist. XXV, App. XVIII (1839)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.I,254
T. cognata; Steudel in Lehm. pl. Preiss. I, 232 (1844)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.I,254
T. rulingioides; Steudel in Lehm. pl. Preiss. I, 232 (1844)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.I,255

<i>T. angustifolia</i> ; Steudel in Lehm. pl. Preiss. I, 232 (1844)...	...	...	...	W.A.	—	—	—	—	—	B.fl.I, 255
<i>T. petalocalyx</i> ; F. v. M. in Transact. phil. Soc. Vict. I, 35 (1854)	...	...	...	W.A. S.A.	—	V.	—	—	—	B.fl.I, 255
<i>T. Sarotes</i> ; Turczaninow in Bull. Soc. Mosc. XXV, 145 (1852)	...	...	...	W.A.	—	—	—	—	—	B.fl.I, 255

GUICHENOTIA; J. Gay in Mém. du Mus. VII, 448, t. 5 (1821). (Sarotes, Ditomostrophe.)

<i>G. ledifolia</i> ; J. Gay in Mém. du Mus. VII, 449, t. 5 (1821)	...	...	...	W.A.	—	—	—	—	—	B.fl.I, 258	M.fr. VII, 174; XI, 114.
<i>G. macrantha</i> ; Turczaninow in Bull. Soc. Mosc. XIX, 500 (1846)	...	...	...	W.A.	—	—	—	—	—	B.fl.I, 258	
<i>G. semihastata</i> ; Benth. Fl. Austr. I, 258 (1863)	...	...	...	W.A.	—	—	—	—	—	B.fl.I, 258	M.fr. II, 4, 7.
<i>G. Sarotes</i> ; Benth. Fl. Austr. I, 258 (1863)	...	...	...	W.A.	—	—	—	—	—	B.fl.I, 258	
<i>G. micrantha</i> ; Benth. Fl. Austr. I, 258 (1863)	...	...	...	W.A.	—	—	—	—	—	B.fl.I, 258	M.fr. II, 7.

LASIOPEPALUM; Smith in Transact. Linn. Soc. IV, 216 (1789). (Corethrostylis, Asterochiton.)

<i>L. laxiflorum</i> ; F. v. M., fragm. XI, 112 (1881)	...	...	...	W.A.	—	—	—	—	—	B.fl.I, 256	M.fr. XI, 112.
<i>L. pygmalum</i> ; Benth. Fl. Austr. I, 256 (1863) implied...	...	...	...	W.A.	—	—	—	—	—	B.fl.I, 257	M.fr. XI, 113.
<i>L. stelligerum</i> ; Benth. Fl. Austr. I, 256 (1863) implied...	...	...	...	W.A.	—	—	—	—	—	B.fl.I, 256	M.fr. XI, 113.
<i>L. glutinosum</i> ; F. v. M., fragm. XI, 113 (1881)	...	...	...	W.A.	—	—	—	—	—	B.fl.I, 256	M.fr. XI, 113.
<i>L. discolor</i> ; Hooker, Comp. to Bot. Mag. I, 276 (1835)	...	...	...	W.A. S.A. T.	—	—	—	—	—	B.fl.I, 260	M.fr. XI, 110.
<i>L. dasyphyllum</i> ; Sieber in Hook. Journ. of bot. II, 414 (1840)	...	...	...	—	—	T.	V.	N.S.W.	—	B.fl.I, 261	M.fr. VI, 174; XI, 110.
<i>L. indutum</i> ; Steudel in Lehm. pl. Preiss. I, 235 (1844)	...	...	...	W.A.	—	—	—	—	—	B.fl.I, 261	M.fr. XI, 110.
<i>L. Maxwellii</i> ; F. v. M., fragm. XI, 107 (1881)	...	...	...	W.A.	—	—	—	—	—	—	M.fr. XI, 107.
<i>L. Ogilvieanum</i> ; F. v. M., fragm. XI, 107 (1881)	...	...	...	W.A.	—	—	—	—	—	—	M.fr. XI, 107.
<i>L. Behrii</i> ; F. v. M. in Transact. phil. Soc. Vict. I, 36 (1854)	...	...	...	—	S.A.	—	V.	N.S.W.	—	B.fl.I, 261	
<i>L. parviflorum</i> ; Rudge in Transact. Linn. Soc. X, 297, t. 19 (1811)	...	...	...	—	—	—	V.	N.S.W.	—	B.fl.I, 262	M.fr. II, 7; VI, 174.
<i>L. parvuliflorum</i> ; F. v. M., fragm. VI, 174 (1868)	...	...	...	W.A.	—	—	—	—	—	—	M.fr. VI, 174.
<i>L. Fitzgibbonii</i> ; F. v. M. in South. Sc. Record II, Apr. (1882)	...	...	...	W.A.	—	—	—	—	—	—	M.fr. XII.
<i>L. oppositifolium</i> ; F. v. M., fragm. II, 5 (1860)	...	...	...	W.A.	—	—	—	—	—	B.fl.I, 262	M.fr. XI, 112.
<i>L. micranthum</i> ; J. Hooker, Fl. Tasman. I, 51 (1860)	...	...	...	—	—	T.	—	—	—	B.fl.I, 262	M.fr. XI, 110.
<i>L. macrophyllum</i> ; Graham in Bot. Mag. t. 3098 (1844)	...	...	...	—	—	—	—	N.S.W.	—	B.fl.I, 262	
<i>L. Baueri</i> ; Steetz in Lehm. pl. Preiss. II, 339 (1847)	...	...	...	—	S.A.	—	V.	N.S.W.	—	B.fl.I, 263	M.fr. VI, 174; XI, 110.
<i>L. Tepperi</i> ; F. v. M., fragm. XI, 109 (1881)	...	...	...	—	S.A.	—	—	—	—	—	M.fr. IX, 109.
<i>L. rufum</i> ; R. Brown in Benth. Fl. Austr. I, 263 (1863)	...	...	...	—	—	—	—	N.S.W.	—	B.fl.I, 263	
<i>L. ferrugineum</i> ; Smith in Bot. Reposit. t. 308 (1799)	...	...	...	—	—	—	V.	N.S.W.	—	B.fl.I, 263	
<i>L. acutiflorum</i> ; Turczaninow in Bull. Soc. Mosc. XXV, 145 (1852)	...	...	...	W.A.	—	—	—	—	—	B.fl.I, 264	M.fr. XI, 109.
<i>L. Oldfieldii</i> ; F. v. M., fragm. II, 6 (1860)	...	...	...	W.A.	—	—	—	—	—	—	M.fr. II, 6.
<i>L. quinquenervium</i> ; Turczaninow in Bull. Soc. Mosc. XXV, 146 (1852)	...	...	...	W.A.	—	—	—	—	—	—	M.fr. XI, 109.
<i>L. Drummondii</i> ; Benth. Fl. Austr. I, 264 (1863)...	...	...	...	W.A.	—	—	—	—	—	B.fl.I, 264	M.fr. VI, 174; XI, 112.
<i>L. rosmarinifolium</i> ; Benth. Fl. Austr. I, 264 (1863)	...	...	...	W.A.	—	—	—	—	—	B.fl.I, 264	M.fr. XI, 112.
<i>L. cordifolium</i> ; Endlicher in Hueg. enum. 10 (1837)	...	...	...	W.A.	—	—	—	—	—	B.fl.I, 265	M.fr. VI, 174; XI, 112.
<i>L. Schulzenii</i> ; F. v. M., pl. of Vict. I, 145 (1861) implied...	...	...	...	—	S.A.	—	V.	—	—	B.fl.I, 265	
<i>L. floribundum</i> ; Benth. Fl. Austr. I, 265 (1863)	...	...	...	W.A.	—	—	—	—	—	B.fl.I, 265	M.fr. XI, 112.
<i>L. molle</i> ; Benth. Fl. Austr. I, 265 (1863)	...	...	...	W.A.	—	—	—	—	—	B.fl.I, 265	M.fr. XI, 111.
<i>L. membranaceum</i> ; Benth. Fl. Austr. I, 266 (1863)	...	...	...	W.A.	—	—	—	—	—	B.fl.I, 266	M.fr. XI, 112.
<i>L. bracteatum</i> ; Benth. Fl. Austr. I, 266 (1863)	...	...	...	W.A.	—	—	—	—	—	B.fl.I, 266	M.fr. XI, 111.

LYSIOSEPALUM; F. v. M., fragm. I, 143 (1859). (Lasiopetali subgenus.)

<i>L. Barryanum</i> ; F. v. M., fragm. I, 143 (1859)	...	...	...	W.A.	—	—	—	—	—	B.fl.I, 267	M.fr. VI, 174.
<i>L. rugosum</i> ; Benth. Fl. Austr. I, 267 (1863)	...	...	...	W.A.	—	—	—	—	—	B.fl.I, 267	

ARISTOTELIA; L'Héritier, stirp. II, 21, t. 16 (1784). (Friesia.)

<i>A. peduncularis</i> ; J. Hooker, Fl. Nov. Zel. I, 33 (1853)	...	...	...	...	—	—	T.	—	—	—	—	B.fl.I,280	M.fr.II,80; VIII,2.
<i>A. Australasica</i> ; F. v. M., fragm. II, 79 (1860)	...	...	...	...	—	—	—	N.S.W.	Q.	—	—	B.fl.I,280	M.fr.VIII,2.
<i>A. megalosperma</i> ; F. v. M., fragm. IX, 84 (1875)	...	...	...	...	—	—	—	—	—	Q.	—	—	M.fr.IX,84.

E. Bancroftii; F. v. M. & Bailey in proceed. Roy. Soc. of Q. II, pt. II (1885)...	—	—	—	—	—	Q.	—	—
E. holopetalus; F. v. M., fragm. II, 143 (1861)	—	—	—	V.	N.S.W.	—	—	B.fl.I, 281 M.fr.IV, 173; V, 211; VI, 172; VIII, 2.
E. Arnheimicus; F. v. M., Decum. intercol. Exhib. 24 (1867)	—	—	—	—	—	—	N.A.	— M.fr.X, 4; VI, 172.
E. obovatus; G. Don, gen. syst. dichl. pl. I, 599 (1831)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.I, 281 M.fr.II, 80; VI, 172; IX, 85.
E. reticulatus; Smith in Rees Cycl. XII (1809)	—	—	—	T.	V.	N.S.W.	Q.	— B.fl.I, 281 M.fr.VI, 172; VIII, 2.
E. grandis; F. v. M., fragm. II, 81 (1860)	—	—	—	—	N.S.W.	Q.	—	B.fl.I, 281 M.fr.II, 175; VI, 172; VIII, 12.
E. foveolatus; F. v. M., fragm. V, 157 (1866)	—	—	—	—	—	Q.	—	— M.fr.VI, 172; VIII, 2.
E. ruminatus; F. v. M., fragm. VIII, 1 (1872)	—	—	—	—	—	Q.	—	— M.fr.X, 4.
E. Grahami; F. v. M., fragm. X, 3 (1876)	—	—	—	—	—	Q.	—	— M.fr.X, 3.
E. sericopetalus; F. v. M., fragm. VI, 171 (1868)	—	—	—	—	—	Q.	—	— M.fr.VI, 171.

S. <i>Australis</i> ; F. v. M., fragm. IV, 91 (1864) ...	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.f.I, 279	M.fr.V, 29; VI, 172; VIII, 2.
S. <i>Langii</i> ; F. v. M., fragm. V, 28 (1865) ...	...	...	...	...	...	—	—	—	—	—	Q.	—	—	M.fr.VIII, 2.
S. <i>Macbrydei</i> ; F. v. M., fragm. VI, 170 (1868) ...	...	...	...	...	...	—	—	—	—	—	Q.	—	—	M.fr.VI, 170.
S. <i>Woolsi</i> ; F. v. M., fragm. VI, 171 (1868) ...	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	—	M.fr.VIII, 2.

B. ammonilla; Roxburgh, pl. Coromand. III, 60, t. 264 (1819)	...	...	...	—	—	—	—	—	Q.	—	B.f.I, 268	...	...	...	As.
--	-----	-----	-----	---	---	---	---	---	----	---	------------	-----	-----	-----	-----

C. Rothii; Triumfetta pilosa, Roth, Nov. pl. sp. 223 (1821)	...	...	...	—	—	—	—	Q.	—	—	M.fr.IV,28; VIII,5	As., Afr.
C. Elderi; F. v. M. in transact., R. S. of S.A. IX, 58 (1886)	...	...	...	—	S.A.	—	—	—	N.A.	—		
C. echinatus; Bentham, Fl. Austr. I, 276 (1863)	...	...	...	...	—	—	—	—	N.A.	B.fl.I,276	M.fr.III,8; VIII,5.	
C. hygrophilus; Cunningham in Fl. Austr. I, 276 (1863)	...	...	...	...	—	—	—	—	Q.	N.A. B.fl.I,276	M.fr.VIII,5.	
C. Cunninghamii; F. v. M., fragm. III, 8 (1862)	...	...	...	...	—	—	—	N.S.W.	Q.	— B.fl.I,276	M.fr.III,167; VI,172; VIII,3.	
C. olitorius; Linné, sp. pl. 529 (1753)	...	...	...	...	—	—	—	—	—	N.A. B.fl.I,276	M.fr.VIII,5	... As., Afr.
C. tridens; Linné, mantiss. alter. 566 (1771)	...	...	...	...	—	—	—	—	—	N.A. B.fl.I,276	...	... As., Afr.
C. acutangulus; Lamarck, encycl. meth. II, 104 (1786)	...	...	...	...	—	—	—	—	Q.	N.A. B.fl.I,277	M.fr.VIII,6	... As., Afr.
C. fascicularis; Lamarck, encycl. meth. II, 104 (1786)	...	...	...	...	—	—	—	—	—	N.A. B.fl.I,277	...	... As., Afr.
C. pumilio; Bentham, Fl. Austr. I, 277 (1863)	...	...	...	...	—	—	—	—	—	N.A. B.fl.I,277	...	... As., Afr.
C. vermicularis; F. v. M., fragm. III, 10 (1862)	...	...	...	...	—	—	—	—	—	N.A. B.fl.I,277	M.fr.VIII,5.	
C. tomentellus; F. v. M., fragm. III, 10 (1862)	...	...	...	...	—	—	—	—	Q.	— B.fl.I,278	M.fr.III,10.	
C. sidoides; F. v. M., fragm. III, 9 (1862)	...	...	...	...	...	W.A.	S.A.	—	Q.	N.A. B.fl.I,278	M.fr.VIII,6; XI,62.	
C. leptocarpus; Cunningham in Benth. Fl. Austr. I, 278 (1863)	...	...	...	...	—	—	—	—	—	N.A. B.fl.I,278		
C. elachocarpus; F. v. M., fragm. VIII, 6 (1872)	...	...	...	...	—	—	—	—	—	N.A.	M.fr.III,6.	
C. Walcottii; F. v. M., fragm. III, 9 (1862)	...	...	...	...	—	—	—	—	—	N.A. B.fl.I,278	M.fr.VI,172; VIII,6; XI,62.	
C. trilocularis; Linné, mantiss. 77 (1771)	...	...	...	...	...	—	—	—	Q.	—	M.fr.VIII,5.	... As., Afr.

T. procumbens; G. Forster, prodr. fl. insul. Austr. 35 (1786)	...	...	...	—	—	—	—	—	Q.	—	B.fl.I, 273	M.fr.III, 8; VI, 172.	As., P.
T. appendiculata; F. v. M., fragm. III, 7 (1862)	...	...	...	...	—	—	—	—	—	N.A.	B.fl.I, 273	M.fr.VIII, 5.	

T. chaetocarpa; F. v. M., fragm. XI, 61 (1879)	...	...	...	...	W.A.	—	—	—	—	N.A.	—	M.fr.XI,61.
T. glaucescens; R. Brown in Benth. Fl. Austr. I, 273 (1863)	...	...	...	...	...	—	—	—	—	N.A.	B.fl.I,273	
T. denticulata; R. Brown in Benth. Fl. Austr. I, 273 (1863)	...	...	...	...	...	—	—	—	—	N.A.	B.fl.I,274	
T. micracantha; F. v. M., fragm. III, 7 (1862)	...	...	...	...	...	—	—	—	—	N.A.	B.fl.I,274	M.fr.VIII,5; XI,62.
T. leptacantha; F. v. M., fragm. XI, 62 (1879)	...	...	...	...	...	—	—	—	—	N.A.	—	M.fr.XI,62.
T. plumigera; F. v. M., fragm. I, 69 (1858)	...	...	...	...	...	—	—	—	—	N.A.	B.fl.I,274	M.fr.III,8; VI,172.
T. parviflora; Benth. Fl. Austr. I, 274 (1863)	...	...	...	...	...	—	—	—	—	N.A.	B.fl.I,274	
T. Winneckeana; F. v. M. in Winnecke's Report (1885)	...	...	...	...	...	—	S.A.	—	—	Q.	N.A.	—
T. rhomboidea; N. Jacquin, select. stirp. Arn. hist. 147, t. 90 (1763)	...	...	...	...	...	—	—	—	—	Q.	—	...
T. Johnstonii; F. v. M. in Melb. Chem. Feb. (1885)	...	...	...	...	...	W.A.	—	—	—	—	—	As., Afr., Am., P.

GREWIA; Linné, syst. nat. 9 (1735); gen. pl. 276 (1737). (Nettoa.)

G. orientalis; Linné, sp. pl. 964 (1753)	...	...	...	...	...	—	—	—	—	Q.	N.A.	B.fl.I,270	...	...	As.
G. laevigata; Vahl, symbol. Bot. I, 34 (1790)	...	...	...	...	...	—	—	—	—	Q.	N.A.	B.fl.I,270	M.fr.VI,172	...	As., Afr., P.
G. breviflora; Benth. Fl. Austr. I, 270 (1863)	...	...	...	...	...	—	—	—	—	—	N.A.	B.fl.I,270			
G. latifolia; F. v. M. in Benth. Fl. Austr. I, 271 (1863)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,271	M.fr.VI,172; VIII,4.	As.
G. polygama; Roxburgh, Fl. Ind. II, 588 (1832)	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.I,271	...	As.
G. xanthopetala; F. v. M. in Benth. Fl. Austr. I, 271 (1863)	...	...	...	...	...	—	—	—	—	—	N.A.	B.fl.I,271			
G. scabrella; Benth. Fl. Austr. I, 272 (1863)	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl.I,272		
G. orbifolia; F. v. M. in Benth. Fl. Austr. I, 272 (1863)	...	...	...	...	...	—	—	—	—	—	N.A.	B.fl.I,272			
G. pleiostigma; F. v. M., fragm. VIII, 4 (1872)	...	...	...	...	...	—	—	—	—	—	Q.	—	—	M.fr.VIII,4.	

EUPHORBIACEAE.

A. L. DE JUSSIEU, Gen. 384 (1789) from B. de Jussieu (1759).—Genera 39, Species 226.

CALYCOPEPLUS; Planchon in Bull. de la Soc. Bot. de Fr. VIII, 31 (1861).

C. ephedroides; Planchon in Bull. Soc. Bot. Fr. VIII, 31 (1861)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.VI,53	
C. marginatus; Benth. Fl. Austr. VI, 53 (1873)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.VI,53	

EUPHORBIA; Linné, gen. pl. 152 (1737) from Plinius.

E. Atoto; G. Forster, florul. ins. Austr. prodr. 36 (1786)	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.VI,46	...	...	As., P.
E. obliqua; Bauer in Endl. fl. Norf. 85 (1833)	...	...	...	...	...	—	—	—	—	N.S.W.	—	—	—			
E. Sparmanni; Boissier, Cent. Euphorb. 5 (1860)	...	...	...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.VI,46	M.fr.IX,77	...	P.
E. Mitchelliana; Boissier in De Candolle, prodr. XV, 25 (1866)	...	...	...	...	...	...	W.A.	—	—	—	—	Q.	N.A.	B.fl.VI,47	M.fr.VI,182.	
E. schizolepis; F. v. M. in De Candolle, prodr. XV, pt. II, 20 (1862)	...	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.VI,47		
E. Schultzii; Benth. in Fl. Austr. VI, 47 (1873)	...	...	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.VI,47		
E. Armstrongiana; Boissier in De Candolle, prodr. XV, 47 (1866)	...	...	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.VI,48		
E. erythrantha; F. v. M., fragm. II, 152 (1861)	...	...	...	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VI,48	M.fr.XI,64.	
E. Muelleri; Boissier in De Candolle, prodr. XV, 27 (1866)	...	...	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.VI,48		
E. Drummondii; Boissier, Cent. Euphorb. 14 (1860)	...	...	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.VI,49	M.fr.VI,182; XI,64.	
E. alsiniflora; Baillon, Adans. VI, 288 (1866)	...	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.VI,49		
E. Wheeleri; Baillon, Adans. VI, 286 (1866)	...	...	...	...	...	...	—	S.A.	—	—	—	—	N.A.	B.fl.VI,49		
E. myrtoides; Boissier in De Candolle, prodr. XV, 15 (1866)	...	...	...	...	...	...	W.A.	—	—	—	—	—	N.A.	B.fl.VI,50		
E. micradenia; Boissier in De Candolle, prodr. XV, 27 (1866)	...	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.VI,50		
E. Macgillivrayi; Boissier in De Candolle, prodr. XV, 26 (1866)	...	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,50		
E. serrulata; Reinwardt in Blum. Bijdrag. 635 (1826)	...	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.VI,51	...	As.
E. filipes; Benth. Fl. Austr. VI, 51 (1873)	...	...	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.VI,51		
E. Carey; F. v. M., fragm. XI, 64 (1879)	...	...	...	...	...	...	W.A.	—	—	—	—	N.A.	—	M.fr.XI,64		

<i>E. pilulifera</i> ; Linné, dissert. de Euphorb. 19 (1752)...	...	...	...	...	—	—	—	—	Q. N.A.	B.fl.VI,51	M.fr.IV,35.	...	As., Afr., Am., P.
<i>E. Tannensis</i> ; Sprengel, syst. veg. III, 791 (1826) ...	...	...	...	...	—	—	—	N.S.W.	—	—	M.fr.IX,169.	...	P.
<i>E. eremophila</i> ; Cunningham in Mitch. Trop. Austr. 348 (1848) ...	...	...	...	...	W.A.	S.A.	V.	N.S.W.	Q. N.A.	B.fl.VI,52			
<i>E. corynoclada</i> ; F. v. M. in Wing's S. Sc. Rec., new ser., vol. II, Apr. (1886)...	...	...	...	...	—	—	—	—	Q.	—	—		

MONOTAXIS; Brogniart in Duperr. Voy. Bot. 223, t. 49 (1829). (Hippocrepandra.)

<i>M. macrophylla</i> ; Benth. Fl. Austr. VI, 79 (1873) ...	...	...	...	...	—	—	—	N.S.W.	Q.	—	B.fl.VI,79		
<i>M. linifolia</i> ; Brongniart in Duperr. Voy. Bot. 223, t. 49 (1829) ...	...	...	...	...	—	—	—	N.S.W.	—	—	B.fl.VI,79		
<i>M. occidentalis</i> ; Endlicher in Hueg. enum. 19 (1837) ...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.VI,79		
<i>M. lurida</i> ; Benth. Fl. Austr. VI, 80 (1873) ...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.VI,80		
<i>M. megacarpa</i> ; F. v. M., fragm. IV, 143 (1865) ...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.VI,80	M.fr.IV,143.	
<i>M. gracilis</i> ; Baillon, Adans. VI, 293 (1866) ...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.VI,80		
<i>M. grandiflora</i> ; Endlicher in Hueg. enum. 19 (1837) ...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.VI,81	M.fr.X,53.	
<i>M. luteiflora</i> ; F. v. M., fragm. X, 51 (1877) ...	...	...	...	...	W.A.	—	—	—	—	—	—	M.fr.X,51.	

PORANTHERA; Rudge in Transact. Linn. Soc. X, 302 t. 22 (1811).

<i>P. ericifolia</i> ; Rudge in Transact. Linn. Soc. X, 302, t. 22 (1811) ...	...	...	...	...	—	—	—	N.S.W.	—	—	B.fl.VI,55		
<i>P. ericoides</i> ; Klotzsch in Lehm. pl. Preiss. II, 239 (1847) ...	...	...	...	...	W.A.	S.A.	—	V.	—	—	B.fl.VI,55		
<i>P. Huegelii</i> ; Klotzsch in Lehm. pl. Preiss. II, 231 (1847) ...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.VI,55		
<i>P. corymbosa</i> ; Brongniart in Duperr. Voy. Coq. Bot. 219, t. 56 (1826) ...	...	...	...	...	—	—	—	V.	N.S.W.	—	B.fl.VI,56	M.fr.VI,183.	
<i>P. microphylla</i> ; Brongniart in Duperr. Voy. Coq. Bot. 218, t. 50 (1826) ...	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q. N.A.	B.fl.VI,56	M.fr.XI,20; XII,11	

MICRANTHEUM; Desfontaines in Mém. du Mus. hist. nat. IV, 253 (1818). (Caletia.)

<i>M. ericoides</i> ; Desfontaines in Mém. du Mus. IV, 253, t. 14 (1818) ..	...	...	...	...	—	—	—	N.S.W.	Q.	—	B.fl.VI,57	M.fr.I,32.	
<i>M. hexandrum</i> ; J. Hooker in Lond. Journ. VI, 283 (1847) ...	...	...	...	...	—	S.A.	T.	V.	N.S.W.	—	B.fl.VI,57	M.fr.XII,11.	

PSEUDANTHUS; Sieber in Sprengel, syst. veg. cur. post. 25 (1827). (Stachystemon, Chrysostemon, Choirizotheca.)

<i>P. pimeloides</i> ; Sieber in Sprengel, syst. veg. cur. post. 25 (1827) ...	...	...	...	...	—	—	—	N.S.W.	Q.	—	B.fl.VI,59	M.fr.I,107.	
<i>P. ovalifolius</i> ; F. v. M. in Transact. phil. Inst. Vict. II, 66 (1857) ...	...	...	...	...	—	—	T.	V.	N.S.W.	Q.	—	B.fl.VI,59	M.fr.I,107.
<i>P. micranthus</i> ; Benth. Fl. Austr. VI, 59 (1873) ...	...	...	...	...	—	S.A.	—	—	—	—	B.fl.VI,59	M.fr.XII,11.	
<i>P. divaricatissimus</i> ; Benth. Fl. Austr. VI, 60 (1873) ...	...	...	...	...	—	—	—	V.	N.S.W.	—	B.fl.VI,60		
<i>P. orientalis</i> ; F. v. M., fragm. II, 14 (1860) ...	...	...	...	...	—	—	—	N.S.W.	Q.	—	B.fl.VI,60	M.fr.II,14.	
<i>P. occidentalis</i> ; F. v. M., fragm. I, 107 (1859) ...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.VI,60	M.fr.I,107; IV,35.	
<i>P. nematophorus</i> ; F. v. M., fragm. II, 14 (1860) ...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.VI,61	M.II,14.	
<i>P. polyandrus</i> ; F. v. M., fragm. II, 153 (1861) ...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.VI,62	M.fr.II,153; IV,35,173.	
<i>P. brachyphyllus</i> ; F. v. M. in Transact. R. Soc. N.S.W. 11 (1881) ...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.VI,62		
<i>P. vermicularis</i> ; F. v. M. in Transact. R. Soc. N.S.W. 11 (1881)...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.VI,62		

BEYERIA; Miquel in Ann. sc. nat. sér. trois. I, 350 (1844). (Calyptrastigma, Beyeropsis.)

<i>B. viscosa</i> ; Miquel in Ann. des sc. nat. sér. trois. I, 350, t. 15 (1844) ...	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,64	M.fr.I,230; II,181; XII,10.
<i>B. lasiocarpa</i> ; F. v. M. in Benth. Fl. Austr. VI, 65 (1873)...	...	...	...	...	—	—	—	V.	N.S.W.	—	—	B.fl.VI,65	M.fr.VI,182.
<i>B. opaca</i> ; F. v. M. in Transact. phil. Soc. Vict. I, 16 (1854) ...	...	...	...	...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,65	M.fr.VI,182; X,53.
<i>B. uncinata</i> ; F. v. M. in Benth. Fl. Austr. VI, 65 (1873) ...	...	...	...	...	—	S.A.	—	—	—	—	—	B.fl.VI,65	
<i>B. latifolia</i> ; Baillon, Adans. VI, 304 (1866) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.VI,66	
<i>B. cygnorum</i> ; Baillon, Adans. VI, 309 (1866) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.VI,66	
<i>B. cinerea</i> ; Baillon, Adans. VI, 309 (1866) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.VI,66	
<i>B. cyanescens</i> ; Benth. Fl. Austr. VI, 66 (1873) ..	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.VI,66	
<i>B. lepidopetala</i> ; F. v. M., fragm. I, 230 (1859) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.VI,67	M.fr.I,230; III,164.
<i>B. similis</i> ; Baillon, Adans. VI, 309 (1866) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.VI,67	
<i>B. brevifolia</i> ; Baillon, Adans. VI, 309 (1866) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.VI,67	
<i>B. Drummondii</i> ; J. Mueller in Linnaea XXXIV, 58 (1865) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.VI,68	
<i>B. tristigma</i> ; F. v. M., fragm. VI, 181 (1868) ...	...	...	...	...	—	—	—	—	Q.	—	B.fl.VI,68	M.fr.VI,181.	

RICINOCARPUS; Desfontaines in Mém. du Mus. III, 459, t. 22 (1817). (Roeperia.)

R. pinifolius; Desfontaines in Mém. du Mus. III, 459, t. 22 (1817)	...	...	...	—	T.	V.	N.S.W.	Q.	—	B.fl.VI,70	M.fr.VI,182.
R. tuberculatus; J. Mueller in Linnaea XXXIV, 60 (1865)	...	...	...	W.A.	—	—	—	—	—	B.fl.VI,70	
R. cyanescens; J. Mueller in Linnaea XXXIV, 60 (1865)	...	...	...	W.A.	—	—	—	—	—	B.fl.VI,70	
R. psilocladus; Benth. Fl. Austr. VI, 71 (1873)	...	...	...	W.A.	—	—	—	—	—	B.fl.VI,70	
R. glaucus; Eudlicher in Hueg. enum. 18 (1873)	...	...	...	W.A.	—	—	—	—	—	B.fl.VI,71	M.fr.VI,182.
R. major; J. Mueller in Linnaea XXXIV, 59 (1865)	...	...	...	—	T.	—	—	—	—	B.fl.VI,72	
R. Bowmanii; F. v. M., fragm. I, 181 (1859)	...	...	...	—	—	—	N.S.W.	Q.	—	B.fl.VI,72	M.fr.III,164; IV,35.
R. ledifolius; F. v. M., fragm. I, 76 (1858)	...	...	...	—	—	—	—	Q.	—	B.fl.VI,72	M.fr.IV,35.
R. rosmarinifolius; Benth. Fl. Austr. VI, 72 (1873)	...	...	...	—	—	—	—	—	N.A.	B.fl.VI,72	
R. marginatus; Benth. Fl. Austr. VI, 73 (1873)	...	...	...	—	—	—	—	—	N.A.	B.fl.VI,73	
R. speciosus; J. Mueller in De Candolle, prodr. XV, 204 (1863)	...	...	...	—	—	—	N.S.W.	—	—	B.fl.VI,73	
R. trichoporus; J. Mueller in Linnaea XXXIV, 60 (1865)	...	...	...	W.A.	—	—	—	—	—	B.fl.VI,73	
R. muricatus; J. Mueller in Linnaea XXXIV, 61 (1865)	...	...	...	W.A.	—	—	—	—	—	B.fl.VI,73	
R. velutinus; F. v. M., fragm. IX, 2 (1875)	...	...	...	W.A.	—	—	—	—	—	—	M.fr.IX,2.

BERTYA; Planchon in Hook. Lond. Journ. Bot. IV, 472, t. 16 (1845). (Lambertya.)

B. gummifera; Planchon in Hook. Lond. Journ. IV, 473 (1845)	...	...	...	—	—	—	N.S.W.	—	—	B.fl.VI,75	M.fr.IV,35.
B. pinifolia; Planchon in Hook. Lond. Journ. IV, 473 (1845)	...	...	...	—	—	—	N.S.W.	Q.	—	B.fl.VI,75	
B. quadriepala; F. v. M., fragm. X, 52 (1876)	...	...	...	W.A.	—	—	—	—	—	—	M.fr.X,52.
B. Cunninghamii; Planchon in Hook. Lond. Journ. IV, 473 (1845)	...	...	...	—	—	—	V.	N.S.W.	—	B.fl.VI,75	M.fr.IV,35; X,52.
B. dimerostigma; F. v. M. in Wing's Science Journ. II (1882)	...	...	...	W.A.	—	—	—	—	—	—	M.fr.XII,9.
B. rosmarinifolia; Planchon in Hook. Lond. Journ. IV, 473 (1845)	...	...	...	—	—	T.	—	N.S.W.	—	B.fl.VI,76	M.fr.IV,35; XII,10.
B. Mitchelli; J. Mueller in Linnaea XXXIV, 63 (1865)	...	...	...	—	S.A.	—	V.	N.S.W.	—	B.fl.VI,76	
B. oppositifolia; F. v. M. & O'Shanesey in S. Sc. Rec. II, May (1882)	...	...	...	—	—	—	—	Q.	—	—	M.fr.XII,9.
B. oleaefolia; Planchon in Hook. Lond. Journ. IV, 473 (1845)	...	...	...	—	—	—	N.S.W.	Q.	—	B.fl.VI,76	
B. rotundifolia; F. v. M., fragm. IV, 34 (1864)	...	...	...	—	S.A.	—	—	—	—	B.fl.VI,77	M.fr.XII,10.
B. pedicellata; F. v. M., fragm. IV, 143 (1865)	...	...	...	—	—	—	—	Q.	—	B.fl.VI,77	M.fr.IV,143.
B. pomaderrioides; F. v. M., fragm. IV, 34 (1864)	...	...	...	—	—	—	N.S.W.	—	—	B.fl.VI,77	M.fr.VIII,141.
B. Findlayi; F. v. M., fragm. VIII, 141 (1874)	...	...	...	—	—	—	V.	N.S.W.	—	—	M.fr.VIII,141.

AMPPEREA; Adr. de Jussieu, Euphorb. gen. 35, t. 10 (1824).

A. protensa; Nees in Lehm. pl. Preiss. II, 229 (1847)	...	...	...	W.A.	—	—	—	—	—	B.fl.VI,82	
A. volubilis; F. v. M. in Benth. Fl. Austr. VI, 82 (1873)	...	...	...	W.A.	—	—	—	—	—	B.fl.VI,82	M.fr.VI,182.
A. micrantha; Benth. Fl. Austr. VI, 83 (1873)	...	...	...	W.A.	—	—	—	—	—	B.fl.VI,83	
A. conferta; Benth. Fl. Austr. VI, 83 (1873)	...	...	...	W.A.	—	—	—	—	—	B.fl.VI,83	
A. ericoides; Adr. de Jussieu, Tent. Euphorb. 112, t. 10 (1824)	...	...	...	W.A.	—	—	—	—	—	B.fl.VI,83	
A. spartioides; Brongniart in Duperr. Voy. Coq. 221, t. 49 (1826)	...	...	...	—	S.A.	T.	V.	N.S.W.	Q.	B.fl.VI,84	

ANTIDESMA; Linné, Fl. Zeyl. 169 (1747).

A. Ghaesembilla; Gaertner, de Fruct. I, 189, t. 39 (1758)	...	...	...	—	—	—	—	—	N.A.	B.fl.VI,85	M.fr.IV,86; VI,183.	...	As., P.
A. Dallachyanum; Baillon, Adans. VI, 337 (1866)	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.VI,85	M.fr.VI,183.	
A. Bunius; Sprengel, syst. veg. I, 826 (1825)	...	...	...	—	—	—	—	—	Q.	—	B.fl.VI,86		...
A. parvifolium; Thwaites & F. v. M., fragm. IV, 86 (1864)	...	...	...	—	—	—	—	—	Q.	—	B.fl.VI,86	M.fr.VI,183.	...
A. Schultzii; Benth. Fl. Austr. VI, 86 (1873)	...	...	...	—	—	—	—	—	—	N.A.	B.fl.VI,86		...
A. erostre; F. v. M. in Benth. Fl. Austr. VI, 87 (1873)	...	...	...	—	—	—	—	—	Q.	—	B.fl.VI,87		
A. sinuatum; Benth. Fl. Austr. VI, 87 (1873)	...	...	...	—	—	—	—	—	Q.	—	B.fl.VI,87		

ANDRACHNE; Linné, syst. nat. 9 (1735); gen. pl. 287 (1737).

A. Decaisnei; Benth. Fl. Austr. VI, 88 (1873)	...	...	...	W.A.	—	—	—	—	Q.	N.A.	B.fl.VI,88	M.fr.VI,183	...	...	As.
---	-----	-----	-----	------	---	---	---	---	----	------	------------	-------------	-----	-----	-----

ACTEPHILA; Blume, Bijdr. XII, 581 (1825).

A. grandifolia; Baillon, Adans. VI, 330 & 360, t. 10 (1866)	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl. VI, 89	
A. Mooreana; Baillon, Adans. VI, 330 & 366 (1866)...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl. VI, 89	
A. latifolia; Benth. Fl. Austr. VI, 89 (1873)	...	...	...	—	—	—	—	—	Q.	—	B.fl. VI, 89	
A. petiolaris; Benth. Fl. Austr. VI, 89 (1873)	...	...	...	—	—	—	—	—	Q.	—	B.fl. VI, 89	
A. sessilifolia; Benth. Fl. Austr. VI, 90 (1873)	...	...	...	—	—	—	—	—	Q.	—	B.fl. VI, 90	

DISSILIARIA; F. v. M. in Baillon, Adans. VII, 356, pl. 1 (1867). (Choriceras.)

D. baloghioides; F. v. M. in Baillon, Adans. VII, 356 (1867)	...	...	...	—	—	—	—	—	Q.	—	B.fl. VI, 90	
D. Muellieri; Baillon, Adans. VII, 359, t. 1 (1867)	...	...	...	—	—	—	—	—	Q.	—	B.fl. VI, 91	
D. tricornis; Benth. Fl. Austr. VI, 91 (1873)	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl. VI, 91	M.fr. XII, 11.

PETALOSTIGMA; F. v. M. in Hook. Kew Misc. IX, 17 (1857). (Hylococcus.)

P. quadriloculare; F. v. M. in Hook. Kew Misc. IX, 17 (1857)	...	...	...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl. VI, 92	M.fr. VI, 182.
--	-----	-----	-----	---	---	---	---	--------	----	------	--------------	----------------

PHYLLANTHUS; Linné, gen. pl. 282 (1737), from J. Commelin. (Glochidion, Bradleia, Synostemon.)

P. Ferdinandi; J. Mueller in Regensb. Flora, 379 (1865)	...	...	...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl. VI, 96	M.fr. VI, 183.
P. lobocarpus; Benth. Fl. Austr. VI, 97 (1873)	...	...	...	—	—	—	—	—	Q.	—	B.fl. VI, 97	
P. ditassoides; J. Mueller in Regensb. Flora, 487 (1864)	...	...	...	—	—	—	—	—	—	N.A.	B.fl. VI, 97	
P. Adami; J. Mueller in De Candolle, prodr. XV, 327 (1866)	...	...	...	—	—	—	—	—	—	N.A.	B.fl. VI, 97	M.fr. I, 33.
P. thesioides; Benth. Fl. Austr. VI, 98 (1873)	...	...	...	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl. VI, 98	
P. hirtellus; J. Mueller in De Candolle, prodr. XV, 326 (1866)	...	...	...	—	—	—	—	—	Q.	—	B.fl. VI, 98	M.fr. III, 89.
P. rigens; J. Mueller in Regensb. Flora, 513 (1864)...	...	...	...	—	S.A.	—	—	N.S.W.	Q.	—	B.fl. VI, 99	M.fr. II, 153; XII, 11.
P. ochrophyllus; Benth. Fl. Austr. VI, 99 (1873)	...	...	...	—	—	—	—	—	—	N.A.	B.fl. VI, 99	
P. rigidulus; F. v. M. in Linnaea XXXIV, 72 (1865)	...	...	...	—	—	—	—	—	—	N.A.	B.fl. VI, 99	
P. ramosissimus; J. Mueller in Linnaea XXXIV, 70 (1865)	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl. VI, 100	M.fr. I, 33; III, 163.
P. Tatei; F. v. M. in Wing's South. Science Record II, 55 (1882)	...	...	...	—	S.A.	—	—	N.S.W.	—	—	—	M.fr. XII, 10.
P. rhytidospermus; F. v. M. in Linnaea XXXIV, 70 (1865)	...	...	...	—	S.A.	—	—	—	—	N.A.	B.fl. VI, 100	M.fr. XII, 11.
P. albiflorus; F. v. M. in Linnaea XXXIV, 70 (1865)	...	...	...	—	—	—	—	—	Q.	—	B.fl. VI, 100	
P. crassifolius; J. Mueller in Regensb. Flora, 513 (1864)	...	...	...	...	W.A.	—	—	—	—	—	B.fl. VI, 100	
P. elachophyllus; F. v. M. in Benth. Fl. Austr. VI, 101 (1873)	...	...	...	...	—	—	—	—	Q.	—	B.fl. VI, 101	
P. uberiflorus; F. v. M. in Adansonia, 343 (1866)	...	...	...	...	—	—	—	—	Q.	—	B.fl. VI, 101	
P. reticulatus; Poir. encycl. meth. V, 298 (1804)	...	...	...	...	—	—	—	—	—	N.A.	B.fl. VI, 101	...
P. baccatus; F. v. M. in Benth. Fl. Austr. VI, 102 (1873)...	...	...	...	...	—	—	—	—	—	N.A.	B.fl. VI, 102	
P. Urinaria; Linné, sp. pl. 982 (1753)	...	...	...	...	—	—	—	—	—	N.A.	B.fl. VI, 102	...
P. trachygyna; Benth. Fl. Austr. VI, 103 (1873)	...	...	...	...	—	—	—	—	—	N.A.	B.fl. VI, 103	As., Afr.
P. Maderaspatanus; Linné, sp. pl. 982 (1753)	...	...	...	...	—	—	—	—	Q.	N.A.	B.fl. VI, 103	
P. Mitchelli; Benth. Fl. Austr. VI, 103 (1873)	...	...	...	...	—	—	—	—	Q.	—	B.fl. VI, 103	...
P. Gastroemii; J. Mueller in De Cand., prodr. XV, 358 (1866)	...	...	...	...	—	—	—	N.S.W.	Q.	—	B.fl. VI, 104	M.fr. XII, 11.
P. Dallachyanus; Benth. Fl. Austr. VI, 104 (1873)	...	...	...	...	—	—	—	—	Q.	—	B.fl. VI, 104	
P. subcrenulatus; F. v. M., fragm. I, 108 (1859)	...	...	...	...	—	—	—	N.S.W.	Q.	—	B.fl. VI, 104	M.fr. I, 108.
P. calycinus; Labillardiere, Nov. Holl. pl. spec. II, 75, t. 225 (1806)	...	...	...	...	W.A.	S.A.	—	—	—	—	B.fl. VI, 105	
P. flagellaris; Benth. Fl. Austr. VI, 106 (1873)	...	...	...	...	—	—	—	—	—	N.A.	B.fl. VI, 106	
P. similis; J. Mueller in Linnaea XXXIV, 71 (1865)	...	...	...	...	—	—	—	—	Q.	—	B.fl. VI, 106	
P. microcladus; J. Mueller in Linnaea XXXIV, 71 (1865)...	...	...	...	...	—	—	—	N.S.W.	Q.	—	B.fl. VI, 106	
P. grandisepalus; F. v. M. in Linnaea XXXIV, 72 (1865)...	...	...	...	...	—	—	—	—	—	N.A.	B.fl. VI, 106	
P. Carpentariae; J. Mueller in Linnaea XXXIV, 72 (1865)	...	...	...	...	—	—	—	—	—	N.A.	B.fl. VI, 107	
P. Fuernrohrrii; F. v. M. in Transact. phil. Soc. Vict. I, 15 (1854)	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl. VI, 107
P. hebecarpus; Benth. Fl. Austr. VI, 108 (1873)...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl. VI, 108

<i>P. lacunarius</i> ; F. v. M. in Transact. phil. Soc. Vict. I, 14 (1854) ...	...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VI,108			
<i>P. trachyspermus</i> ; F. v. M. in Transact. phil. Soc. Vict. I, 14 (1854) ...	...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VI,108			
<i>P. Australis</i> ; J. Hooker in Lond. Journ. VI, 284 (1847) ...	...	...	—	S.A.	T.	—	N.S.W.	—	—	B.fl.VI,108	M.fr.XII,11.		
<i>P. thymoides</i> ; Sieber in Linnaea XXVIII, 566 (1856) ...	...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VI,109			
<i>P. scaber</i> ; Klotzsch in Lehm. pl. Preiss. I, 179 (1844) ...	...	...	—	W.A.	—	—	—	—	—	B.fl.VI,109			
<i>P. indigoferoidea</i> ; Benth. Fl. Austr. VI, 110 (1873) ...	...	...	—	—	—	—	—	—	N.A.	B.fl.VI,110			
<i>P. aridus</i> ; Benth. Fl. Austr. VI, 110 (1873) ...	...	...	—	—	—	—	—	—	N.A.	B.fl.VI,110			
<i>P. Gunnii</i> ; J. Hooker in Lond. Journ. VI, 284 (1847) ...	...	...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,110			
<i>P. simplex</i> ; Retzius, observ. V, 29 (1789) ...	...	...	—	—	—	—	—	Q.	—	B.fl.VI,111	M.fr.VI,183.	...	As., P.
<i>P. filicaulis</i> ; Benth. Fl. Austr. VI, 111 (1873) ...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.VI,111			
<i>P. minutiflorus</i> ; F. v. M. in Linnaea XXXIV, 75 (1865) ...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.VI,112			
<i>P. Armstrongii</i> ; Benth. Fl. Austr. VI, 112 (1873) ...	...	...	—	—	—	—	—	—	N.A.	B.fl.VI,112			
<i>P. buxifolius</i> ; Reinwardt in Blume, catal. plant. Buitenz. 106 (1823) ...	...	...	—	—	—	—	—	Q.	—	—	M.fr.X,121.	...	As.
BREYNIA ; R. & G. Forster, char. gen. 145, t. 73 (1776). (Melanthesa, Melanthesiopais.)													
<i>B. cernua</i> ; J. Mueller in De Candolle, prodr. XV, 439 (1866) ...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.VI,113			
<i>B. oblongifolia</i> ; J. Mueller in De Candolle, prodr. XV, 110 (1866) ...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,114			
<i>B. stipitata</i> ; J. Mueller in De Candolle, prodr. XV, 442 (1866) ...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.VI,114			
<i>B. rhynchocarpa</i> ; Benth. Fl. Austr. VI, 114 (1873) ...	...	...	—	—	—	—	—	—	N.A.	B.fl.VI,114			
SECURINEGA ; A. L. de Jussieu, Gen. 388 (1789). (Flueggea.)													
<i>S. Abyssinica</i> ; A. Richard, tentam. Fl. Abyss. II, 256 (1851) ...	...	...	...	W.A.	—	—	—	Q.	N.A.	B.fl.VI,116	...	...	As., Afr., P.
<i>S. Leucopyrus</i> ; J. Mueller in De Candolle, prodr. XV, 451 (1866) ...	...	...	—	S.A.	—	—	—	Q.	N.A.	B.fl.VI,116	...	...	As.
NEOROEPERA ; F. & J. Mueller in De Candolle, prodr. XV, 489 (1866).													
<i>N. buxifolia</i> ; F. & J. Mueller in De Candolle, prodr. XV, 489 (1866) ...	...	...	—	—	—	—	—	Q.	—	B.fl.VI,116			
<i>N. Banksii</i> ; Benth. Fl. Austr. VI, 117 (1873) ...	...	...	—	—	—	—	—	Q.	—	B.fl.VI,117			
BISCHOFFIA ; Blume, Bijdr. 1168 (1825).													
<i>B. Javanica</i> ; Blume, Bijdr. 1168 (1825) ...	...	...	—	—	—	—	—	Q.	N.A.	—	M.fr.VIII,141	...	As., P.
HEMICYCLIA ; Wight & Arnott in Edinb. phil. Journ. XIV, 297 (1833).													
<i>H. sepiaria</i> ; Wight & Arnott in Edinb. phil. Journ. XIV, 297 (1833) ...	...	...	—	—	—	—	—	—	N.A.	B.fl.VI,117	M.fr.IV,119 ; VI,182 ; IX,77.	...	As.
<i>H. Australasica</i> ; J. Mueller in De Candolle, prodr. XV, 487 (1866) ...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,118			
<i>H. lasiogyne</i> ; F. v. M., fragm. IV, 119 (1864) ...	...	...	—	—	—	—	—	—	N.A.	B.fl.VI,118	M.fr.IV,119.		
APOROSA ; Blume, Bijdrag. 514 (1824).													
<i>A. Australiana</i> ; F. v. M., syst. cons. of Austr. pl. 20 (1882) ...	...	...	—	—	—	—	—	Q.	—	—	M.fr.VIII,141.		
BRIDELIA ; Willdenow, Spec. pl. IV, 978 (1805).													
<i>B. exaltata</i> ; F. v. M., fragm. III, 32 (1862) ...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,119	M.fr.III,32.		
<i>B. ovata</i> ; Decaisne in nouv. ann. du Mus. V, 484 (1835) ...	...	...	—	—	—	—	—	—	N.A.	B.fl.VI,120	...	...	As.
<i>B. tomentosa</i> ; Blume, Bijdrag. 597 (1825) ...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.VI,120	M.fr.III,32 ; VI,182	...	As., P.
<i>B. faginea</i> ; F. v. M. in Benth. Fl. Austr. VI, 120 (1873) ...	...	...	—	—	—	—	—	Q.	—	B.fl.VI,120			
CLEISTANTHUS ; J. Hooker, icon. pl. t. 779 (1847). (Lebediera, Amanoa.)													
<i>C. Cunninghamii</i> ; J. Mueller in De Candolle, prodr. XV, 506 (1866) ...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,122			
<i>C. apodus</i> ; Benth. Fl. Austr. VI, 122 (1873) ...	...	...	—	—	—	—	—	Q.	—	B.fl.VI,122			
<i>C. Dallachyanus</i> ; Baillon in Benth. Fl. Austr. VI, 122 (1873) ...	...	...	—	—	—	—	—	Q.	—	B.fl.VI,122	M.fr.VI,182.		
<i>C. semiopacus</i> ; F. v. M. in Benth. Fl. Austr. VI, 123 (1873) ...	...	...	—	—	—	—	—	Q.	—	B.fl.VI,123			

CROTON; Linné, gen. pl. 288 (1737).

C. Schultzii; Benth. Fl. Austr. VI, 124 (1873)	...	...	...	...	...	—	—	—	—	—	N.A.	B.fl. VI, 124			
C. insularis; Baillon, Adans. II, 217 (1862)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl. VI, 124		
C. phebaloides; F. v. M. in Benth. Fl. Austr. VI, 125 (1873)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl. VI, 125 M.fr. IV, 140	...	P.
C. opposens; F. v. M. in Benth. Fl. Austr. VI, 125 (1873)	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl. VI, 125		
C. tomentellus; F. v. M., fragm. IV, 141 (1864)	...	...	...	...	...	—	—	—	—	—	N.A.	B.fl. VI, 126 M.fr. IV, 141.			
C. Verreauxii; Baillon, Etud. Euph. 357 (1858)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl. VI, 126 M.fr. IV, 141 ; VI, 186.		
C. acronychioides; F. v. M., fragm. IV, 142 (1864)	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl. VI, 127 M.fr. VI, 186.		
C. triacros; F. v. M., fragm. VI, 185 (1867)	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl. VI, 127 M.fr. VI, 185.		
C. Arnheimicus; F. v. M. in Linnaea XXXIV, 112 (1865)	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl. VI, 127 M.fr. VI, 186.		

ALEURITES; R. & G. Forster, char. gen. 111, t. 56 (1776).

A. triloba; R. & G. Forster, char. gen. 111, t. 56 (1776)	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl. VI, 128 M.fr. V, 30; VI, 182. As., P.
---	-----	-----	-----	-----	-----	---	---	---	---	---	----	---	---

CLAOXYLON; Adr. de Jussieu, Euphorb. gen. 43, t. 14 (1824).

C. angustifolium; J. Mueller in Linnaea XXXIV, 165 (1865)	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl. VI, 129
C. tenerifolium; F. v. M. in Baillon, Adans. VI, 323 (1866)	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl. VI, 130 M.fr. VI, 183.
C. Australe; Baillon, Etud. Euphorb. 493 (1858)	...	...	...	...	...	—	—	—	—	V. N.S.W.	Q.	—	B.fl. VI, 130 M.fr. IV, 142.
C. Hillii; Benth. Fl. Austr. VI, 131 (1873)	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl. VI, 131

ACALYPHA; Royen in Linné, hort. Cliffort. 495 (1737).

A. nemorum; F. v. M. in Linnaea XXXIV, 38 (1865)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl. VI, 132
A. eremorum; F. v. M. in Regensb. Flora 440 (1864)	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl. VI, 132
A. capillipes; F. v. M. in Linnaea XXXIV, 40 (1865)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl. VI, 133

ADRIANA; Gaudichaud in Ann. sc. nat. V. 223 (1825). (Trachycaryon.)

A. tomentosa; Gaudichaud in Ann. sc. nat. V, 223 (1825)	...	...	...	...	...	W.A. S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl. VI, 134 M.fr. IX, 2; XII, 11.
A. quadripartita; Gaudichaud in Freyc. voy. Bot. 489 (1826)	...	...	...	...	...	W.A. S.A.	—	V.	—	—	—	B.fl. VI, 135 M.fr. XII, 11.

ALCHORNEA; Solander in Swartz, prodr. 698 (1788). (Coelebogynne, Cladodes.)

A. ilicifolia; J. Mueller in Linnaea XXXIV, 170 (1865)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl. VI, 136
A. Thozetiana; F. v. M. in Benth. Fl. Austr. VI, 137 (1873)	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl. VI, 137

TRAGIA; Plumier, gen. 14, t. 12 (1703).

T. Novae Hollandiae; J. Mueller in Linnaea XXXIV, 180 (1865)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl. VI, 137 M.fr. VI, 183; XII, 11.
--	-----	-----	-----	-----	-----	---	---	---	---	--------	----	---	---------------------------------------

MALLOTUS; Loureiro, Fl. Cochinch. II, 635 (1790). (Echinus, Rottlera, Echinocroton.)

M. ricinoides; J. Mueller in Linnaea XXXIV, 187 (1865)	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl. VI, 140 M.fr. IV, 138; VI, 184 As., P.
M. Chinensis; Loureiro, Fl. Cochinchin. II, 635 (1770)	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl. VI, 140 M.fr. VI, 184 ... As., P.
M. claoxyloides; J. Mueller in Linnaea XXXIV, 192 (1865)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl. VI, 140 M.fr. I, 32; VI, 184.
M. Philippinensis; J. Mueller in Linnaea XXXIV, 196 (1865)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl. VI, 141 M.fr. VI, 183 ... As., P.
M. angustifolius; Benth. in Fl. Austr. VI, 141 (1873)	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl. VI, 142
M. polyadenos; F. v. M., fragm. VI, 184 (1867)	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl. VI, 142 M.fr. VI, 184.
M. repandus; J. Mueller in Linnaea XXXIV, 197 (1865)	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl. VI, 142 M.fr. VI, 185.
M. mesophilus; F. v. M. in Linnaea XXXIV, 196 (1865)	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl. VI, 143 M.fr. VI, 184.
M. discolor; F. v. M. in Benth. Fl. Austr. VI, 143 (1873)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl. VI, 143 M.fr. IV, 139.

MACARANGA; Petit-Thouars, Gen. Madag. n. 88 (1809). (Mappa.)

M. Dallachyi; F. v. M. in Benth. Fl. Austr. VI, 144 (1873)	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl. VI, 144 M.fr. VI, 185.
M. subdentata; Benth. Fl. Austr. VI, 145 (1873)	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl. VI, 145
M. inamoena; F. v. M. in Benth. Fl. Austr. VI, 145 (1873)	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl. VI, 145
M. involucreta; Baillon, Etud. Euphorb. 432 (1858)	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl. VI, 146 M.fr. IV, 139, 177; VI, 183 As.
M. Tanaria; J. Mueller in De Candolle, prodr. XV, 997 (1866)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl. VI, 146 M.fr. IV, 140; VI, 183 ... As., P.

CODIAEUM; Rumphius, herbar. Amboin. IV, 65, t. 25 (1743).

C. variegatum; Blume, Bijdrag. 605 (1825) ... — — — — — Q. — B.fl. VI, 147 M.fr. VI, 182 ... As., P.

BALOGHIA; Endlicher, prodr. fl. Norfolk, 84 (1833).

B. lucida; Endlicher, prodr. fl. Norfolk, 84 (1833) ... — — — — — N.S.W. Q. — B.fl. VI, 148 M.fr. VI, 182; IX, 77 ... P.

FONTAINEA; Heckel, thèse inaugurale, Montpell. (1870).

F. Pancheri; Heckel, thèse inaugurale (1870) ... — — — — — Q. — B.fl. VI, 149 M.fr. XII, 11 ... P.

OMALANTHUS; Adr. de Jussieu, Euphorb. gen. 50, t. 16 (1824). (Carumbium, Wartmannia, Homalanthus.)

O. Leschenaultianus; A. de Jussieu, Euph. gen. 50, t. 16, f. 53 (1824) ... — — — — — V. N.S.W. Q. N.A. B.fl. VI, 150 M.fr. I, 32; VI, 183; IX, 77; XII, 11 As., P.

O. stillingiifolius; F. v. M., fragm. I, 32 (1853) ... — — — — — N.S.W. Q. — B.fl. VI, 150 M.fr. I, 33; III, 163

SEBASTIANIA; Sprengel, Neue Entd. II, 118 (1821). (Microstachys, Elachocroton.)

S. chamaelaea; J. Mueller in De Candolle, prodr. XV, 1175 (1866) ... — — — — — Q. N.A. B.fl. VI, 151 ... As., Afr., P.

EXCAECARIA; Linné, syst. veg. ed. decim. 1288 (1759).

E. Agallocha; Linné, syst. ed. decim. 1288 (1759) ... — — — — — N.S.W. Q. N.A. B.fl. VI, 152 M.fr. XI, 64 ... As., P.

E. Dallachyana; Baillon, Adans. VI, 324 (1866) ... — — — — — Q. — B.fl. VI, 153

E. parvifolia; J. Mueller in Flora, 433 (1864) ... — — — — — Q. N.A. B.fl. VI, 153

URTICACEAE.

VENTENAT; Tabl. du règn. vég. III, 524 (1799).—Genera 19, Species 65.

CELTIS; Tournefort, iust. 612, t. 383 (1700) from Camerarius (1586). (Solenostigma.)

C. Philippinensis; Blanco, Fl. de Filipinas 197 (1837) ... — — — — — Q. N.A. B.fl. VI, 156 M.fr. IV, 88 ... As., P.

C. paniculata; Planchon in Ann. sc. nat. sér. trois. X, 305 (1848) ... — — — — — N.S.W. Q. N.A. B.fl. VI, 156 M.fr. IV, 88 ... As.

C. amblyphylla; F. v. M., fragm. IX, 76 (1875) ... — — — — — N.S.W. — — — M.fr. IX, 76.

TREMA; Loureiro, Fl. Cochinch. II, 565 (1790). (Sponia.)

T. cannabina; Loureiro, Fl. Cochinch. II, 689 (1790) ... — — — — — V. N.S.W. Q. N.A. B.fl. VI, 158 M.fr. VI, 194; VIII, 247 As., P.

APHANANTHE; Planchon in Ann. sc. nat. X, 265 (1848).

A. Philippinensis; Planchon in Ann. sc. nat. X, 337 (1848) ... — — — — — N.S.W. Q. — B.fl. VI, 160 M.fr. VI, 192 ... As.

FICUS; Tournefort, inst. 662 (1700) from Plinius. (Urostigma, Covellia.)

F. colossea; F. v. M. in Benth. Fl. Austr. VI, 163 (1873) ... — — — — — Q. — B.fl. VI, 163

F. pilosa; Reinwardt in Blume, Bijdr. 446 (1825) ... — — — — — Q. — B.fl. VI, 164 M.fr. VI, 195 ... As.

F. glabella; Blume, Bijdr. 452 (1825) ... — — — — — Q. N.A. B.fl. VI, 164 M.fr. VI, 195 ... As.

F. Cunninghamii; Miquel in Ann. Mus. Lugd. Bat. III, 286 (1867) ... — — — — — Q. — B.fl. VI, 165 M.fr. VI, 195; VIII, 246.

F. Henneana; Miquel in Ann. Mus. Lugd. Bat. III, 216 (1867) ... — — — — — Q. N.A. B.fl. VI, 165

F. validinervis; F. v. M. in Benth. Fl. Austr. VI, 166 (1873) ... — — — — — Q. — B.fl. VI, 166

F. retusa; Linné, mantissa plant. 129 (1767) ... — — — — — Q. N.A. B.fl. VI, 166 M.fr. VI, 195 ... As., P.

F. eugenioides; F. v. M., Docum. Interc. Exhib. 26 (1866) ... — — — — — N.S.W. Q. N.A. B.fl. VI, 166 M.fr. VI, 195

F. Benjaminea; Linné, mantissa plant. 129 (1767) ... — — — — — Q. — B.fl. VI, 167 M.fr. VI, 195 ... As.

F. Muellieri; Miquel in Ann. Mus. Lugd. Bat. III, 287 (1867) ... — — — — — N.S.W. — — B.fl. VI, 167 M.fr. VI, 195; VIII, 246.

F. leucotricha; Miquel in Ann. Mus. Lugd. Bat. III, 285 (1867) ... — — — — — Q. N.A. B.fl. VI, 167 M.fr. VI, 195.

F. rubiginosa; Desfontaines, catal. pl. hort. Paris. 209 (1804) ... — — — — — N.S.W. Q. — B.fl. VI, 168 M.fr. IV, 194; VIII, 247.

F. columnaris; F. v. M. & Moore in proc. Accl. Soc. Vict. III, 71 (1874) ... — — — — — N.S.W. — — — M.fr. VIII, 247.

F. macrophylla; Desfontaines, catal. pl. hort. Paris. 209 (1804) ... — — — — — N.S.W. Q. — B.fl. VI, 170 M.fr. VII, 194; VIII, 247.

F. paberula; Cunningham in Hook. Lond. Journ. VI, 562 (1847) ... — — — — — — N.A. B.fl. VI, 169

F. platypoda; Cunningham in Hook. Lond. Journ. VI, 561 (1847)	...	...	W.A.	S.A.	—	—	—	Q.	N.A.	B.fl. VI, 169	M.fr. VI, 194; VIII, 247.			
F. dictyophleba; F. v. M. in Benth. Fl. Austr. VI, 170 (1873)	...	...	—	—	—	—	—	Q.	—	B.fl. VI, 170				
F. magnifolia; F. v. M., fragm. IV, 50 (1864)	...	...	—	—	—	—	—	Q.	—	B.fl. VI, 171	M.fr. IV, 177; VI, 196.			
F. ehretioides; F. v. M. in Benth. Fl. Austr. VI, 171 (1873)	...	...	—	—	—	—	—	Q.	—	B.fl. VI, 171				
F. pumila; Linné, spec. plant. 1060 (1753)	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl. VI, 171	M.fr. VI, 196	...	As.	
F. coronulata; F. v. M. in Journ. Bot. Neerland. 241 (1861)	...	...	—	—	—	—	—	—	N.A.	B.fl. VI, 172	M.fr. IV, 49; VI, 196; VIII, 246.			
F. leptoclada; Benth. Fl. Austr. VI, 172 (1873)	...	...	—	—	—	—	—	Q.	—	B.fl. VI, 172				
F. depressa; Benth. Fl. Austr. VI, 172 (1873)	...	...	—	—	—	—	—	Q.	—	B.fl. VI, 172				
F. subulata; Blume Bijdr., 461 (1825)	...	...	—	—	—	—	—	Q.	—	—	...	...	As.	
F. infectoria; Roxburgh. flor. Indic. III, 550 (1832)	...	...	—	—	—	—	—	Q.	—	—	...	...	As.	
F. Philippinensis; Miquel in Hooker, Lond. Journ. VII, 435 (1848)	...	...	—	—	—	—	—	Q.	—	B.fl. VI, 173	...	...	As., P.	
F. mollior; F. v. M. in Benth. Fl. Austr. VI, 173 (1873)	...	...	—	—	—	—	—	Q.	—	B.fl. VI, 173				
F. stenocarpa; F. v. M. in Benth. Fl. Austr. VI, 174 (1873)	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl. VI, 174	M.fr. IX, 152.			
F. scabra; G. Forster, florul. ins. Austr. prodr. 76 (1786)	...	...	—	—	—	—	V. N.S.W.	Q.	N.A.	B.fl. VI, 174	M.fr. IV, 85; VI, 196; IX, 152; X, 114.	As., P.		
F. subglabra; F. v. M., fragm. IX, 152 (1875)	...	...	—	—	—	—	—	Q.	—	—				
F. Pinkiana; F. v. M. in Wing's S. Sec. Rec. II, 273 (1882)	...	...	—	—	—	—	—	Q.	—	—				
F. orbicularis; Cunningham in Hook. Lond. Journ. VII, 426 (1848)	...	...	—	S.A.	—	—	—	—	N.A.	B.fl. VI, 175	M.fr. VI, 196.			
F. aculeata; Cunningham in Hook. Lond. Journ. VII, 426 (1848)	...	...	—	—	—	—	—	—	N.A.	B.fl. VI, 175				
F. opposita; Miquel in Hook. Lond. Journ. VII, 426 (1848)	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl. VI, 176	M.fr. VI, 196; VIII, 246.			
F. scobina; Benth. Fl. Austr. VI, 176 (1873)	...	...	—	—	—	—	—	—	N.A.	B.fl. VI, 176				
F. hispida; Linné, fil. suppl. 442 (1781)	...	...	—	—	—	—	—	Q.	N.A.	B.fl. VI, 176	...	...	As.	
F. fasciculata; F. v. M. in Benth. Fl. Austr. VI, 177 (1873)	...	...	—	—	—	—	—	Q.	—	B.fl. VI, 177				
F. casearia; F. v. M. in Benth. Fl. Austr. VI, 177 (1873)	...	...	—	—	—	—	—	Q.	—	B.fl. VI, 177	...	...	As.	
F. glomerata; Willdenow, spec. pl. IV, 1148 (1806)	...	...	—	—	—	—	—	Q.	N.A.	B.fl. VI, 178	M.fr. VI, 195	...	As.	
F. pleurocarpa; F. v. M., fragm. VIII, 246 (1874)	...	...	—	—	—	—	—	Q.	—	—	M.fr. VIII, 246.			
CUDRANIA; Trécul in Ann. des sc. nat. VIII, 122 (1847).														
C. Javanensis; Trécul in Ann. des sc. nat. VIII, 123 (1847)	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl. VI, 179	M.fr. IV, 88	...	As., Afr., P.	
ANTIARIS; Leschenault in Ann. du Mus. XVI, 476 (1810).														
A. macrophylla; R. Brown in Flind. voy. II, 602, t. 5 (1814)	...	...	—	—	—	—	—	—	N.A.	B.fl. VI, 179	M.fr. VI, 194.			
MALAISIA; Blanco, Fl. Filip. 789 (1837). (Caturus.)														
M. tortuosa; Blanco, Fl. Filip. 789 (1837)	...	...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl. VI, 180	M.fr. IV, 88; VI, 193&255; VIII, 247; IX, 77.			[As., P.]
PSEUDOMORUS; Bureau in Ann. des sc. nat. sér. cinq. XI, 372 (1872).														
P. Brunoniana; Bureau in Ann. des sc. nat. sér. cinq. XI, 372 (1872)	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl. VI, 181	M.fr. IV, 88; VI, 192.	P.		
FATOUA; Gaudichaud in Freyc. voy. Bot. 509 (1826).														
F. pilosa; Gaudichaud in Freyc. voy. Bot. 509 (1826)	...	...	—	—	—	—	—	—	N.A.	B.fl. VI, 182	...	...	As., P.	
ELATOSTEMMA; R. & G. Forster, char. gen. 105, t. 53 (1776).														
E. reticulatum; Weddell in Ann. des sc. nat. sér. 4, I, 188 (1854)	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl. VI, 183	M.fr. IX, 77.			
E. stipitatum; Weddell in Ann. des sc. nat. sér. 4, I, 190 (1854)	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl. VI, 184				
PROCRIS; Commerçon in A. L. de Jussieu, gen. 403 (1789).														
P. montana; Steudel, nomencl. bot. II, 398 (1841)	...	...	—	—	—	—	N.S.W.	—	—	—	M.fr. IX, 169.			
BOEHMERIA; N. J. Jacquin, stirp. Amer. hist. 246, t. 157 (1763).														
B. calophleba; Moore & Mueller, fragm. VIII, 11 (1870)	...	...	—	—	—	—	N.S.W.	—	—	B.fl. VI, 184	M.fr. VIII, 11.			
B. Australis; Endlicher, prodr. Fl. Norfolk. 38 (1833)	...	...	—	—	—	—	N.S.W.	—	—	—	M.fr. VIII, 12; IX, 169.			

PIPTURUS; Weddell in Ann. des sc. nat. sér. quatr. I, 196 (1854).

P. propinquus; Weddell in Archiv. du Mus. 447, t. 15 (1856) ... — — — — N.S.W. Q. — B.fl. VI, 185 M.fr. IV, 87; VI, 197; VIII, 12. As., P.

POUZOLZIA; Gaudichaud in Freyc. voy. Bot. 503 (1826). (Memoralis, Hyrtanandra.)

P. Indica; Gaudichaud in Freyc. voy. Bot. 503 (1826) ... — — — — Q. N.A. B.fl. VI, 186 M.fr. IV, 87 ... As.

P. quinquenervis; Bennett, Pl. Jav. rar. 66 (1852) ... — — — — Q. N.A. B.fl. VI, 187 M.fr. V, 194 ... As.

PARIETARIA; Tournefort, inst. 509, t. 289 (1700) from C. Bauhin (1623). Freirea, from Brunfels (1532).

[N.Z.]

P. debilis; G. Forster, florul. ins. Austr. prodr. 73 (1786) ... W.A. S.A. T. V. N.S.W. Q. N.A. B.fl. VI, 188 M.fr. IV, 88; IX, 77; XI, 20. As., Afr., Am., P.,

AUSTRALINA; Gaudichaud in Freyc. voy. Bot. 497 (1826).

A. pusilla; Gaudichaud in Freyc. voy. Bot. 505 (1826) ... — — T. V. N.S.W. Q. — B.fl. VI, 189 M.fr. IV, 88; VIII, 247 ... N.Z.

FLEURYA; Gaudichaud in Freyc. voy. Bot. 497 (1826).

F. interrupta; Gaudichaud in Freyc. voy. Bot. 497 (1826) ... — — — — Q. — — M.fr. X, 114 ... As., Afr., P.

URTICA; Tournefort, inst. 534, t. 308 (1700) from Plinius.

U. incisa; Poiret, Encycl. meth. suppl. IV, 223 (1817) ... — S.A. T. V. N.S.W. Q. — B.fl. VI, 190 M.fr. IV, 87 ... N.Z.

LAPORTEA; Gaudichaud in Freyc. voy. Bot. 498 (1826).

L. gigas; Weddell in Arch. du Mus. IX, 129, t. 3 & 4 (1856) ... — — — — N.S.W. Q. — B.fl. VI, 192 M.fr. IV, 87.

L. photiniphylla; Weddell in Arch. du Mus. IX, 138 (1856) ... — — — — N.S.W. Q. — B.fl. VI, 192 M.fr. IV, 88, 194.

L. moroides; Weddell in Arch. du Mus. IX, 142 (1856) ... — — — — Q. — B.fl. VI, 192 M.fr. VIII, 248.

CUPULIFERAE.

L. CL. RICHARD; Anal. de fr. 32 & 92 (1808).—Genera 2, Species 4.

FAGUS; Tournefort, inst. 584, t. 351 (1700) from Camerarius (1586).

F. Gunnii; J. Hooker, icon. plant., t. 881 (1852) ... — — T. — — — B.fl. VI, 210 M.fr. V, 110.

F. Cunninghamii; Hooker, Journ. of Bot. II, 152, t. 7 (1840) ... — — T. V. — — — B.fl. VI, 210 M.fr. V, 110.

F. Moorei; F. v. M., fragm. V, 109 (1866) ... — — — — N.S.W. — — B.fl. VI, 211 M.fr. V, 109.

BALANOPS; Baillon, Hist. des pl. VI, 237 & 258 (1876).

B. Australiana; F. v. M., fragm. X, 114 (1877) ... — — — — Q. — — M.fr. X, 114.

CASUARINEAE.

MIRBEL in Ann. du Mus. XVI, 451 (1810).—Genus 1, Species 24.

CASUARINA; Rumphius, herb. Amboin. III, 87, t. 58 (1743).

C. quadrivalvis; Labillardière, nov. Holl. pl. spec. II, 67, t. 218 (1806) ... W.A. S.A. T. V. N.S.W. — — B.fl. VI, 195 M.fr. VI, 18 & 19; X, 62.

C. trichodon; Miquel in Lehm. pl. Preiss. I, 641 (1845) ... W.A. — — — — B.fl. VI, 196 M.fr. VI, 17.

C. lepidophloia; F. v. M., fragm. X, 115 (1877) ... — S.A. — — N.S.W. — — M.fr. X, 115.

C. glauca; Sieber in Sprengel, syst. III, 803 (1826) ... — S.A. — V. N.S.W. Q. — B.fl. VI, 196 M.fr. VI, 18.

C. Huegeliana; Miquel in Lehm. pl. Preiss. I, 640 (1845) ... W.A. — — — — B.fl. VI, 196 M.fr. VI, 19; X, 62.

C. equisetifolia; R. & G. Forster, char. gen. 103, t. 52 (1776) ... — — — — N.S.W. Q. N.A. B.fl. VI, 197 M.fr. VI, 17 ... As., Afr., P.

C. suberosa; Otto & Dietrich, allgem. Gartenzeit. 155 (1841) ... — S.A. T. V. N.S.W. Q. — B.fl. VI, 197 M.fr. VI, 20 & 254.

C. Cunninghamiana; Miquel, Revis. Casuarin. 56, t. 6 (1848) ... — — — — N.S.W. Q. — B.fl. VI, 198

C. paludosa; Sieber in Spreng. syst. veg. III, 803 (1826) ... — — — — V. N.S.W. — — —

<i>C. distyla</i> ; Ventenat, pl. dans le jard. de Cels. t. 62 (1800)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl. VI, 198	M.fr. VI, 19.
<i>C. inophloia</i> ; F. v. M. & Bailey in Melb. Chemist, April (1882)	—	—	—	—	N.S.W.	Q.	—	—	M.fr. XII.
<i>C. Fraseriana</i> ; Miquel, Revis. Casuarin. 50, t. 6 (1848)	W.A.	—	—	—	—	—	—	B.fl. VI, 199	M.fr. VI, 19.
<i>C. nana</i> ; Sieber in Sprengel, syst. III, 804 (1826)	—	—	—	V.	N.S.W.	—	—	B.fl. VI, 199	
<i>C. humilis</i> ; Otto & Dietrich, allgem. Gartenzeit. 163 (1841)	W.A.	S.A.	—	—	—	—	—	B.fl. VI, 200	M.fr. VI, 18.
<i>C. torulosa</i> ; Aiton, hort. Kew. III, 320 (1789)	—	—	—	—	N.S.W.	Q.	—	B.fl. VI, 200	M.fr. VI, 17.
<i>C. decussata</i> ; Bentham, Fl. Austr. VI, 200 (1873)	W.A.	—	—	—	—	—	—	B.fl. VI, 200	
<i>C. Decaisneana</i> ; F. v. M., fragm. I, 61 (1858)	W.A.	S.A.	—	—	—	Q.	N.A.	B.fl. VI, 201	M.fr. I, 61; X, 62.
<i>C. Drummondiana</i> ; Miquel, Revis. Casuar. 26, t. 1 (1848)	W.A.	—	—	—	—	—	—	B.fl. VI, 201	
<i>C. microstachya</i> ; Miquel in Lehm. pl. Preiss. I, 642 (1845)	W.A.	—	—	—	—	—	—	B.fl. VI, 201	M.fr. VI, 17; X, 62.
<i>C. acutivalvis</i> ; F. v. M., fragm. X, 61 (1876)	W.A.	—	—	—	—	—	—	—	M.fr. X, 61.
<i>C. bicuspidata</i> ; Bentham, Fl. Austr. VI, 202 (1873)	W.A.	S.A.	T.	—	—	—	—	B.fl. VI, 202	
<i>C. thuyoides</i> ; Miquel in Lehm. pl. Preiss. I, 641 (1845)	W.A.	—	—	—	—	—	—	B.fl. VI, 202	M.fr. VI, 17.
<i>C. corniculata</i> ; F. v. M., fragm. X, 62 (1876)	W.A.	—	—	—	—	—	—	—	M.fr. X, 62.
<i>C. acuaria</i> ; F. v. M., fragm. VI, 16 (1868)	W.A.	—	—	—	—	—	—	B.fl. VI, 202	M.fr. VI, 250.

CELASTRINAE.

R. BROWN in App. Flind. voy. 554 (1814).—Genera 10, Species 17.

EUONYMUS; Tournefort, inst. 617, t. 388 (1700) from Theophrastos & Plinius. (Evonymus.)

E. Australianus; F. v. M., fragm. IV, 118 (1864) Q. — — M.fr. VI, 202.

CELASTRUS; Linné, gen. pl. 59 (1737).

C. Australis; Harvey & F. v. M. in Trans. phil. Soc. Vict. I, 41 (1854) V. N.S.W. Q. — B.fl. I, 398 M.fr. III, 93.
C. Muelleri; Bentham, Fl. Austr. I, 399 (1863) — — — — — N.A. B.fl. I, 399
C. dispermus; F. v. M. in Transact. phil. Inst. Vict. III, 31 (1859) Q. — B.fl. I, 399 M.fr. VI, 167, 203.
C. bilocularis; F. v. M. in Transact. phil. Inst. Vict. III, 31 (1859) N.S.W. Q. — B.fl. I, 399 M.fr. VI, 204.
C. Cunninghamii; F. v. M. in Transact. phil. Inst. Vict. III, 30 (1859) N.S.W. Q. N.A. B.fl. I, 399 M.fr. VI, 203.

GYMNOSPORA; Wight & Arnott, prodr. Fl. penins. Ind. I, 159 (1834).

G. montana; Wight & Arnott, prodr. I, 159 (1834) Q. — B.fl. I, 400 As.

HEDRAIANTHERA; F. v. M., fragm. V, 58 (1865).

H. porphyropetala; F. v. M., fragm. V, 59 (1865) N.S.W. Q. — — M.fr. V, 214.

HYP SOPHILA; F. v. M. in Vict. Nat. III, 169 (1887).

H. Halleyana; F. v. M. in Vict. Nat. III, 169 (1887) Q. — —

LEUCOCARPUM; A. Richard, sert. Astrolabe 46 (1834). (Denhamia, Leucocarpon.)

L. Oleaster; F. v. M. in Transact. Roy. Soc. of N.S.W. 15 (1881) Q. — B.fl. I, 401 M.fr. VI, 203.
L. obscurum; A. Richard, sert. Astrol. 46, t. 18 (1834) Q. N.A. B.fl. I, 401 M.fr. VI, 203.
L. pittosporoides; F. v. M. in Transact. Roy. Soc. of N.S.W. 15 (1881) N.S.W. Q. — B.fl. I, 402 M.fr. VI, 203.

ELAEODENDRON; J. F. Jacquin in Nov. Act. Helv. I, 36 (1787). (Portenschlagia.)

E. Australe; Ventenat, Jard. Malm. t. 117 (1804) N.S.W. Q. N.A. B.fl. I, 402 M.fr. III, 61, 168; IV, 175; VI, 204; IX, 77.
E. curtispiculum; Endlicher, prodr. fl. ins. Norf. 81 (1833) N.S.W. — — —

CARYOSPERMUM; Blume, Mns. Bot. Lugd. I, 175 (1850). (Perrottetiae subgenus.)

C. arborescens; F. v. M., fragm. VI, 202 (1868) Q. — — M.fr. VIII, 279.

SIPHONODON; Griffith in Calc. Journ. of Nat. Hist. IV, 150 (1843).

S. Australe; Benth. Fl. Austr. I, 403 (1863) ... — — — — N.S.W. Q. — B.fl.I,403 M.fr.VI,204.

HIPPOCRATEA; Linné, gen. pl. 363 (1737).

H. obtusifolia; Roxburgh, flor. Ind. edit. Carey, I, 170 (1820) ... — — — — N.S.W. Q. — B.fl.I,404 ... As.

SALACIA; Linné, mantiss. II, 159 (1771).

S. prinoidea; De Candolle, prodr. I, 571 (1824) ... — — — — Q. — — M.fr.VI,202.

SAPINDACEAE.

A. L. DE JUSSIEU, Gen. plant. 246 (1789).—Genera 14, Species 100.

CARDIOSPERMUM; Linné, syst. nat. (1735); gen. pl. 117 (1737).

C. Halicacabum; Linné, sp. pl. 366 (1753) ... — — — — Q. N.A. B.fl.I,453 M.fr.IX,89 ... As., Afr., Am., P.

ATALAYA; Blume, Rumphia III, 186 (1847). (Pseudatalaya, Sapindus partly.)

A. coriacea; Radlkofer in Sitzungsber. der Akad. zu Münch. 326 (1878) ... — — — — N.S.W. — — —
 A. salicifolia; Blume, Rumphia III, 186 (1847) ... — — — — — N.A. B.fl.I,463 ... As.
 A. Australis; Radlkofer in Sitzungsber. der Akad. zu Münch. 327 (1878) ... — — — — Q. — B.fl.I,464
 A. multiflora; Benth. Fl. Austr. I, 463 (1863) ... — — — — N.S.W. Q. — B.fl.I,463 M.fr.IX,90.
 A. hemiglauc; F. v. M. in Benth. Fl. Austr. I, 463 (1863) ... — S.A. — — N.S.W. Q. N.A. B.fl.I,463 M.fr.IX,90.
 A. variifolia; F. v. M. in Benth. Fl. Austr. I, 463 (1863) ... — — — — Q. N.A. B.fl.I,463 M.fr.I,45; IX,90.

DIPLOGLOTTIS; J. Hooker in H. & B. gen. pl. 395 (1862). (Stadmannia partly.)

D. Cunninghamii; J. Hooker in H. & B. gen. pl. 395 (1862) ... — — — — N.S.W. Q. — B.fl.I,454 M.fr.V,145; IX,90.

ERIOGLOSSUM; Blume, Bijdr. 229 (1829). (Pancovia partly.)

E. edule; Blume, Bijdr. 229 (1829) ... — — — — — N.A. B.fl.I,454 M.fr.IX,100 ... As.

CASTANOSPORA; F. v. M., fragm. IX, 92 (1875).

C. Alphandi; F. v. M., fragm. IX, 92 (1875) ... — — — — N.S.W. Q. — — M.fr.IV,158; IX,197.

ALLOPHYLUS; Linné, amoen. acad. 124 (1747). (Schmidelia.)

A. ternatus; Loureiro, Fl. Cochinch. ed. Willd. 286 (1790) ... — — — — Q. N.A. B.fl.I,455 M.fr.IX,90 ... As., P.

CUPANIA; Plumier, Gen. 45, t. 19 (1703). (Guioa, Ratonía, Aryteria, Elattostachys, Mischocarpus, Euphoria partly.)

C. Wadsworthii; F. v. M., syst. Cens. of Austr. pl. 24 (1882) ... — — — — N.S.W. Q. — — M.fr.IV,1; V,147; IX,89,197.
 C. anacardioides; A. Richard, sert. Astrol. 33, t. 13 (1833) ... — — — — N.S.W. Q. N.A. B.fl.I,458 M.fr.IX,91.
 C. Robertsonii; F. v. M., fragm. V, 146 (1866) ... — — — — Q. — — M.fr.IX,94.
 C. serrata; F. v. M., fragm. III, 43 (1862) ... — — — — N.S.W. Q. — B.fl.I,453 M.fr.IX,94,198.
 C. foveolata; F. v. M., fragm. IX, 95 (1875) ... — — — — Q. — — M.fr.IX,95.
 C. tomentella; F. v. M. in Benth. Fl. Austr. I, 458 (1863) ... — — — — Q. — B.fl.I,458 M.fr.IX,94.
 C. Pseudorhus; A. Richard, sert. Astrol. 34, t. 14 (1833) ... — — — — N.S.W. Q. — B.fl.I,459 M.fr.V,146; IX,92.
 C. pleurophylla; F. v. M. in Vict. Natural. III, 170 (1887) ... — — — — Q. — —
 C. xylocarpa; Cunningham in Benth. Fl. Austr. I, 459 (1863) ... — — — — N.S.W. Q. — B.fl.I,459 M.fr.V,7; IX,95.
 C. erythrocarpa; F. v. M., fragm. V, 7 (1865) ... — — — — Q. — — M.fr.IX,91.
 C. Mortoniana; F. v. M., fragm. V, 177 (1866) ... — — — — Q. — — M.fr.IX,94.
 C. lachnocarpa; F. v. M., fragm. IV, 157 (1864) ... — — — — Q. — — M.fr.V,6; IX,91.
 C. grandissima; F. v. M., fragm. IV, 156 (1864) ... — — — — Q. — — M.fr.IX,91.
 C. pyriformis; F. v. M., App. Rep. Intercol. Exhib. 25 (1867) ... — — — — N.S.W. Q. — B.fl.I,461 M.fr.I,2; II,76; IX,90.
 C. anodonta; F. v. M., fragm. II, 76 (1860) ... — — — — Q. — B.fl.I,461 M.fr.I,2; IX,90.
 C. stipitata; F. v. M., fragm. II, 75 (1860) ... — — — — N.S.W. — — B.fl.I,461 M.fr.III,12; IV,157; V,6,147,210; IX,91.
 C. punctulata; F. v. M., fragm. III, 12 (1862) ... — — — — Q. — B.fl.I,458 M.fr.IX,91.

C. exangulata; F. v. M., fragm. IV, 156 (1864)	—	—	—	—	—	Q.	—	—	M.fr.IX,91.
C. O'Shanesiana; F. v. M., fragm. IX, 96 (1875)	—	—	—	—	—	Q.	—	—	M.fr.IX,96.
C. Daemeliana; F. v. M., fragm. IX, 96 (1875)	—	—	—	—	—	Q.	—	—	M.fr.IX,96.
C. Martiana; F. v. M., fragm. V, 6 (1865)	—	—	—	—	—	Q.	—	—	M.fr.IX,94.
C. Cordieri; F. v. M., fragm. IX, 93 (1875)	—	—	—	—	—	Q.	—	—	M.fr.IX,93.
C. tenax; Cunningham in Benth. Fl. Austr. I, 461 (1863)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,461	M.fr.IX,94.

NEPHELIUM; Linné, mantiss. 18 (1767). (Euphoria partly, Alectryon, Spanoghea.)

N. semiglaucum; F. v. M. in Transact. phil. Soc. Vict. III, 25 (1857)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,457	M.fr.IV,158; V,147; IX,97.
N. connatum; F. v. M. in Benth. Fl. Austr. I, 465 (1863)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,465	M.fr.IV,158; IX,98.
N. semicinerum; F. v. M., fragm. IV, 158 (1864)	—	—	—	—	—	Q.	—	—	M.fr.IV,158.
N. subdentatum; F. v. M. in Benth. Fl. Austr. I, 465 (1863)	—	—	—	—	N.S.W.	—	—	B.fl.I,465	M.fr.IX,99.
N. tomentosum; F. v. M. in Transact. Vict. Inst. II, 64 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,466	M.fr.IX,99.
N. coriaceum; Benth. Fl. Austr. I, 466 (1863)	—	—	—	—	—	Q.	—	B.fl.I,466	—
N. foveolatum; F. v. M. in Benth. Fl. Austr. I, 466 (1863)	—	—	—	—	—	Q.	—	B.fl.I,466	M.fr.IX,99.
N. leiocarpum; F. v. M. in Transact. phil. Inst. Vict. III, 25 (1858)	—	—	—	—	V. N.S.W.	Q.	—	B.fl.I,467	M.fr.IX,98.
N. Beckleri; Benth. Fl. Austr. I, 467 (1863)	—	—	—	—	N.S.W.	—	—	B.fl.I,467	—
N. divaricatum; F. v. M. in Benth. Fl. Austr. 467 (1863)	—	—	—	—	—	Q.	—	B.fl.I,467	M.fr.IX,98.
N. microphyllum; Benth. Fl. Austr. I, 468 (1863)	—	—	—	—	—	Q.	—	B.fl.I,468	—
N. Leichhardtii; F. v. M., App. Rep. Intercol. Exhib. 25 (1867)	—	—	—	—	—	Q.	—	B.fl.I,468	M.fr.IX,99.
N. distyle; F. v. M., fragm. IX, 99 (1875)	—	—	—	—	—	Q.	—	B.fl.I,462	M.fr.IX,99.

HETERODENDRON; Desfontaines in Mém. du Mus. IV, 8, t. 3 (1818).

H. oleaeifolium; Desfontaines in Mém. du Mus. IV, 8, t. 3 (1818)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.I,469	M.fr.IX,75,90; X,82.
H. diversifolium; F. v. M., fragm. I, 46 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,469	M.fr.IX,90; X,82.

HARPULLIA; Roxburgh, Fl. Ind. ed. Carey II, 44 (1826).

H. alata; F. v. M., fragm. II, 103 (1861)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,470	M.fr.IX,89,198.
H. Hillii; F. v. M. in Transact. Vict. Inst. III, 26 (1859)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,470	M.fr.II,104; IX,100,197.
H. Leichhardtii; F. v. M. in Benth. Fl. Austr. I, 470 (1863)	—	—	—	—	—	—	N.A.	B.fl.I,470	—
H. pendula; Planchon in Transact. Vict. Inst. III, 26 (1859)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,471	M.fr.II,104; IX,89,100&187.

AKANIA; J. Hooker in B. & H. Gen. pl. 409 (1862).

A. Hillii; J. Hooker in B. & H. Gen. pl. 409 (1862)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,471	M.fr.III,44; IX,89&197.
--	---	---	---	---	--------	----	---	------------	-------------------------

DIPLOPELTIS; Endlicher in Huegel, enum. 13 (1837).

D. Huegelii; Endlicher in Huegel, enum. 13 (1837)	W.A.	—	—	—	—	—	N.A.	B.fl.I,456	M.fr.III,12; IX,89.
D. Stuartii; F. v. M., fragm. III, 12 (1862)	—	S.A.	—	—	—	—	N.A.	B.fl.I,456	M.fr.III,12&167; IX,89.

DODONAEA; Linné, gen. pl. 341 (1737). (Empleurosma.)

D. triquetra; Wendland, Bot. Beobacht. p. 44 (1798)	—	—	—	V.	N.S.W.	Q.	—	B.fl.I,474	M.fr.I,73; IX,88.
D. lanceolata; F. v. M., fragm. I, 73 (1858)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.I,475	M.fr.I,73.
D. petiolaris; F. v. M., fragm. III, 13 (1862)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.I,475	M.fr.IX,89.
D. viscosa; Linné, mantissa altera 149 & 228 (1771)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.I,475	M.fr.IX,81 ... As., Afr., Am., P., N.Z.
D. peduncularis; Lindley in Mitch. Trop. Austr. 361 (1848)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,478	M.fr.IX,87.
D. procumbens; F. v. M. in Transact. Vict. Inst. I, 3 (1854)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.I,478	—
D. ericifolia; G. Don, gen. syst. I, 674 (1831)	—	—	T.	—	—	—	—	B.fl.I,478	M.fr.IX,87.
D. filifolia; Hooker in Mitch. Trop. Austr. 241 (1848)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,478	M.fr.I,71.
D. lobulata; F. v. M. in Linnæa XXV, 372 (1852)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.I,479	M.fr.IX,87,197.
D. ptarmicifolia; Turczaninow in Bull. Soc. Mosc. XXV, part III, 165 (1852)	W.A.	—	—	—	—	—	—	B.fl.I,479	M.fr.I,97; IX,87.
D. calycina; Cunningham in A. Gray, Bot. of Wilk. Expl. Exped. I, 262 (1854)	—	—	—	V.	N.S.W.	Q.	—	B.fl.I,479	M.fr.II,143; IX,89.
D. platyptera; F. v. M., fragm. I, 73 (1858)	W.A.	—	—	—	—	—	N.A.	B.fl.I,480	M.fr.IX,87.
D. pachyneura; F. v. M. in Transact. R. S. of Vict. XXIII, 51 (1886)	W.A.	—	—	—	—	—	N.A.	—	—
D. stenophylla; F. v. M., fragm. I, 72 (1858)	—	—	—	—	—	Q.	—	B.fl.I,480	M.fr.IX,87.

D. pinifolia; Miquel in Lehm. pl. Preiss. I, 227 (1844)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.I,480	M.fr.IX,88.	
D. ceratocarpa; Endlicher in Hueg. enum. 13 (1837)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.I,481	M.fr.X,88.	
D. divaricata; Benth. Fl. Austr. I, 481 (1863)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.I,481	M.fr.IX,87.	
D. triangularis; Lindley in Mitch. Trop. Austr. 219 (1848)	...	...	...	...	—	—	—	—	N.S.W.	Q.	B.fl.I,481	M.fr.IX,87.	
D. aptera; Miquel in Lehm. pl. Preiss. I, 225 (1844)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.I,481	M.fr.IX,88.	
D. bursarifolia; Behr & F. v. M. in Transact. Vict. Inst. I, 8 (1855)	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	—	B.fl.I,482	M.fr.IX,87.	
D. Baueri; Endlicher in Hueg. enum. 13 (1837)	...	...	...	...	—	S.A.	—	V.	N.S.W.	—	B.fl.I,482	M.fr.IX,87.	
D. trifida; F. v. M., fragm. IX, 88 (1875)	...	...	...	...	W.A.	—	—	—	—	—	—	M.fr.IX,88.	
D. humifusa; Miquel in Lehm. pl. Preiss. I, 206 (1844)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.I,482	M.fr.IX,88.	
D. hexandra; F. v. M. in Transact. Vict. Inst. I, 117 (1855)	...	...	...	...	—	S.A.	—	—	—	—	B.fl.I,483		
D. ericoides; Miquel in Lehm. pl. Preiss. I, 227 (1844)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.I,483	M.fr.IX,88.	
D. polyzyga; F. v. M., fragm. I, 74 (1858)	...	...	...	...	—	—	—	—	—	N.A.	B.fl.I,483	M.fr.I,219.	
D. megazyga; F. v. M. in Benth. Fl. Austr. I, 483 (1863)	...	...	...	...	—	—	—	—	N.S.W.	—	B.fl.I,483	M.fr.IX,86.	
D. macrozyga; F. v. M., fragm. IV, 135 (1864)	...	...	...	...	—	S.A.	—	—	—	Q.	—	M.fr.IX,86.	
D. physocarpa; F. v. M., fragm. I, 74 (1858)	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.I,484	M.fr.IX,86.
D. vestita; Hooker in Mitch. Trop. Austr. 265 (1848)	...	...	...	...	—	—	—	—	—	Q.	—	B.fl.I,484	M.fr.IX,86.
D. pinnata; Smith in Rees' Cycl. XII (1809)	...	...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.I,484	
D. oxyptera; F. v. M., fragm. I, 74 (1858)	...	...	...	...	—	—	—	—	—	N.A.	B.fl.I,484	M.fr.I,74.	
D. humilis; Endlicher, nov. stirp. dec. 26 (1839)	...	...	...	...	—	S.A.	—	—	—	N.A.	B.fl.I,485	M.fr.IX,86.	
D. Macrossani; F. v. M. & Scortechini in Melb. Chemist Jan. 1882	...	...	...	...	—	—	—	—	—	Q.	—	—	M.fr.XII.
D. microzyga; F. v. M., annual Rep. Bot. Gard. 12 (1863)	...	...	...	...	—	S.A.	—	—	—	—	—	—	M.fr.IX,86.
D. boronifolia; G. Don, gen. syst. I, 674 (1831)	...	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.I,485	M.fr.IX,86.
D. multijuga; G. Don, gen. syst. I, 674 (1831)	...	...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.I,485	M.fr.IX,86.
D. larraeoides; Turczaninow in Bull. Soc. Mosc. XXXI, 408 (1858)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.I,486	M.fr.I,210; II,182.
D. inaequifolia; Turczaninow in Bull. Soc. Mosc. XXXI, 408 (1858)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.I,486	M.fr.I,219; IV,135; IX,89.
D. adenophora; Miquel in Linnaea XVIII, 95 (1845)	...	...	...	...	W.A.	—	—	—	N.S.W.	Q.	—	B.fl.I,486	M.fr.I,98.
D. tenuifolia; Lindley in Mitch. Trop. Austr. 248 (1848)	...	...	...	...	—	S.A.	—	—	N.S.W.	Q.	N.A.	—	M.fr.IX,85.
D. stenozyga; F. v. M., fragm. I, 98 (1859)	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.I,486	M.fr.IX,87.
D. concinna; Benth. Fl. Austr. I, 487 (1863)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.I,487	M.fr.IX,87.

DISTICHOSTEMON; F. v. M. in Hook. Kew Misc. IX, 306 (1857). (Pleistichostemon.)

D. phyllopterus; F. v. M. in Hook. Kew Misc. IX, 306 (1857)	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.I,487	M.fr.IX,89.
---	-----	-----	-----	-----	---	---	---	---	---	----	------	------------	-------------

MALPIGHIACEAE.

VENTENAT; Tabl. III, 131 (1799).—Genera 2, Species 2.

RYSSOPTERYS; Blume in Delessert, ic. sel. III, 21, t. 35 (1837).

R. Timorensis; Blume in Deless. ic. sel. III, 21, t. 35 (1837)	...	...	...	...	—	—	—	—	—	Q.	—	B.fl.I,285	...	...	As.
--	-----	-----	-----	-----	---	---	---	---	---	----	---	------------	-----	-----	-----

TRISTELLATEIA; Petit-Thouars in Roemer, Collect. bot. 206 (1809).

T. Australasica; A. Richard, Sert. Astrol. 38, t. 15 (1833)	...	...	...	...	—	—	—	—	—	Q.	—	B.fl.I,286	...	...	As., P.
---	-----	-----	-----	-----	---	---	---	---	---	----	---	------------	-----	-----	---------

BURSERACEAE.

KUNTH in Ann. des sc. nat. II, 346 (1824).—Genera 3, Species 3.

GARUGA; Roxburgh Pl. of Coromandel III, 5, t. 208 (1819).

G. floribunda; Descaisne, in nouv. annal. du Mus. III, 477 (1834)	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.I,377	...	...	As.
---	-----	-----	-----	-----	---	---	---	---	---	----	------	------------	-----	-----	-----

- CANARIUM; Rumphius, herb. Amboin. II, 145, t. 47, 49, 54 (1741). (Sonzaya.)
 C. Australianum; F. v. M., fragm. III, 15 (1860) Q. N.A. B.fl.I, 378 M.fr.V, 211; VII, 25.
 GANOPHYLLUM; Blume, mus. bot. Lugd. I, 230 (1850).
 G. falcatum; Blume, mus. bot. Lugd. I, 230 (1850) Q. — — M.fr.VII, 24; IX, 193; XI, 16. As.

ANACARDIACEAE.

R. BROWN in Tuckey's Expedit. Congo 431 (1818).—Genera 6, Species 9.

- EUROSCHINUS; J. Hooker in B. & H. Gen. pl. I, 422 (1862).
 E. falcatus; J. Hooker in B. & H. Gen. pl. I, 422 (1862) N.S.W. Q. — B.fl.I, 490 M.fr.VII, 24.
 RHUS; Tournefort, Inst. 611, t. 381 (1700), from Hippocrates, Theophrastus, Dioscorides & Plinius. (Rhodosphaera.)
 R. rhodanthema; F. v. M. in Journ. pharm. soc. Vict. pl. I, 43 (1858) N.S.W. Q. — B.fl.I, 489 M.fr.VII, 23.
 R. rufa; Teyssman & Binnendyk in Naturk. Tijd. XXVII, 52 (1863) Q. — — M.fr.VII, 22.
 BLEPHAROCARYA; F. v. M., fragm. XI, 15 (1878).
 B. involucrigera; F. v. M., fr. XI, 16 (1878) Q. — — M.fr.XI, 137.
 BUCHANANIA; Sprengel in Schrader's Journ. II, 234 (1800). (Coniogeton.)
 B. mangoides; F. v. M., fragm. VII, 23 (1869) Q. — — M.fr.VII, 23.
 B. obovata; Engler in A. & C. de Candolle, monogr. phaner. IV, 187 (1883) — N.A. —
 B. Muelleri; Engler in A. & C. de Candolle, monogr. phaner. IV, 190 (1883) Q. N.A. B.fl.I, 490 M.fr.VII, 23.
 SEMECARPUS; Linné fil., suppl. 25 (1781).
 S. Anacardium; Linné, fil., suppl. 182 (1781) Q. N.A. B.fl.I, 491 M.fr.VII, 23 ... As., P.
 SPONDIAS; Linné, gen. pl. 365 (1737). (Pleiogynium.)
 S. Solandri; Benth, Fl. Austr. I, 492 (1863) Q. — B.fl.I, 492 M.fr.III, 15; V, 177.

STACKHOUSIEAE.

R. BROWN in App. to Flind. Voy. 555 (1814).—Genera 2, Species 13.

- STACKHOUSIA; Smith in Transact. Linn. Soc. IV, 218 (1798). (Tripterococcus, Plokiostigma.)
 S. pulvaris; F. v. M. in Transact. phil. Soc. Vict. I, 101 (1855) — — T. V. N.S.W. — — B.fl.I, 405 M.fr.III, 88.
 S. linearifolia; Cunningham in Field's N. S. Wales 365 (1825) — — S.A. T. V. N.S.W. Q. — B.fl.I, 406 M.fr.III, 87; XI, 27.
 S. pubescens; A. Richard, sert. Astrol. 89, t. 33 (1834) W.A. — — — — B.fl.I, 407
 S. Huegelii; Endlicher in Hueg. enum. 17 (1837) W.A. — — — — B.fl.I, 407
 S. flava; Hooker, icon. pl. t. 269 (1840) W.A. S.A. T. V. — — — B.fl.I, 407
 S. muricata; Lindley, Bot. Reg. XXII, 1917 (1837) — S.A. — — N.S.W. Q. N.A. B.fl.I, 408 M.fr.VIII, 36.
 S. viminea; Smith in Rees' Cycl. XXXIII (1816) W.A. S.A. — V. N.S.W. Q. N.A. B.fl.I, 408 M.fr.III, 86; VIII, 36.
 S. elata; F. v. M., fragm. III, 86 (1862) W.A. — — — — B.fl.I, 408 M.fr.III, 86.
 S. scoparia; Benth, Fl. Austr. I, 409 (1863) W.A. — — — — B.fl.I, 409
 S. spathulata; Sieber in Spreng. syst. cur. post. 124 (1827) — S.A. T. V. N.S.W. Q. — B.fl.I, 406 M.fr.III, 86.
 S. megaloptera; F. v. M., fragm. VIII, 35 (1873) W.A. S.A. — — — — M.fr.X, 119.
 S. Brunonis; Benth, Fl. Austr. I, 409 (1863) W.A. — — — — N.A. B.fl.I, 409 M.fr.VIII, 36.
 MACGREGORIA; F. v. M. in Caruel, Giorn. Bot. Italian. 128 (1873).
 M. racemigera; F. v. M. in Caruel, Giorn. Bot. Ital. 128 (1873) — S.A. — — N.S.W. Q. — — M.fr.VIII, 161; XI, 134.

FRANKENIACEAE.

A. DE ST. HILAIRE in Bull. de la Soc. Philom. 22 (1815).—Genus 1, Species 7.

FRANKENIA; Linné, gen. pl. 92 (1737).

F. laevis; Linné, spec. plant. 331 (1753)	...	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.I,151	M.fr.IX,75	...	E., As., Afr.
F. bracteata; Turczaninow in Bull. Soc. Mosc. XXVII, 367 (1854)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.I,150			
F. glomerata; Turczaninow in Bull. Soc. Mosc. XXVII, 368 (1854)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.I,151			
F. parvula; Turczaninow in Bull. Soc. Mosc. XXVII, 368 (1854)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.I,152			
F. Drummondii; Benth, Fl. Austr. I, 152 (1863)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.I,152			
F. tetrapetala; Labillardière, nov. Holl. pl. spec. I, 83, t. 114 (1804)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.I,152			
F. punctata; Turczaninow in Bull. Soc. Mosc. XXVII, 367 (1854)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.I,153			

PLUMBAGINEAE.

A. L. DE JUSSIEU, gen. plant. 92 (1789).—Genera 3, Species 4.

PLUMBAGO; Tournefort, inst. 140, t. 58 (1700).

P. Zeilanica; Linné, sp. pl. 151 (1753)	...	...	...	...	...	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.IV,267	M.fr.XI,7	...	As., Afr., P.
---	-----	-----	-----	-----	-----	---	------	---	---	--------	----	------	-------------	-----------	-----	---------------

STATICE; Linné, gen. pl. 88 (1737). (Taxanthema.)

S. Taxanthema; Roemer & Schultes, syst. veg. VI, 798 (1820)	...	...	...	...	...	—	—	T.	V.	N.S.W.	Q.	—	B.fl.IV,267	M.fr.XI,7	...	As., P.
S. salicorniacea; F. v. M., fragm. XI, 7 (1878)	...	...	...	...	...	W.A.	—	—	—	—	—	N.A.	—	M.fr.XI,7.		

AEGIALITIS; R. Brown, prodr. 426 (1810). (Aegialinites.)

A. annulata; R. Brown, prodr. 426 (1810)	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.IV,266	M.fr.XI,7	...	P.
--	-----	-----	-----	-----	-----	---	---	---	---	---	----	------	-------------	-----------	-----	----

PORTULACEAE.

A. L. DE JUSSIEU, gen. plant. 312 (1789).—Genera 3, Species 32.

PORTULACA; Tournefort, inst. 236, t. 118 (1700) from W. Turner (1538).

P. oleracea; Linné, spec. plant. 445 (1753)	...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.I,169	M.fr.X,71	...	E., As., Afr., Am., P.
P. napiformis; F. v. M. in Benth. Fl. Austr. I, 169 (1863)	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.I,169	M.fr.X,72	...	As.
P. Australis; Endlicher, Atakta 7, t. 6 (1833)	...	...	...	...	...	—	S.A.	—	—	—	Q.	N.A.	B.fl.I,169	M.fr.X,72.		
P. filifolia; F. v. M., fragm. I, 169 (1859)	...	...	...	...	...	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.I,169	M.fr.III,164; X,72.		
P. digyna; F. v. M., fragm. I, 170 (1859)	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.I,170	M.fr.I,170.		
P. oligosperma; F. v. M., fragm. I, 170 (1859)	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.I,170	M.fr.III,164; X,72.	...	As.
P. bicolor; F. v. M., fragm. I, 171 (1859)	...	...	...	...	...	—	S.A.	—	—	—	Q.	N.A.	B.fl.I,170	M.fr.III,164; X,72.		
P. Armitii; F. v. M., fragm. X, 97 (1877)	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	—	M.fr.X,97.		

CLAYTONIA; Gronovius in Linné, gen. pl. 339 (1737). (Calandrinia, Talinum, Montiae subgenus.)

C. spargularina; F. v. M., Cens. 27 (1882)	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.I,176	M.fr.I,175; IV,137; X,71.		
C. ptychosperma; F. v. M., Cens. 27 (1882)	...	...	...	...	...	W.A.	S.A.	—	—	—	Q.	N.A.	—	M.fr.IV,137; X,70.		
C. strophiolata; F. v. M., fragm. XI, 82 (1880)	...	...	...	...	...	W.A.	—	—	—	—	—	N.A.	—	M.fr.XI,82.		
C. Lehmanni; F. v. M., Cens. 27 (1882) from Endl. (1847)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.I,172	M.fr.X,71.		
C. uniflora; F. v. M., Cens. 27 (1882)	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.I,172	M.fr.X,71.		
C. Balonnensis; F. v. M., Cens. 27 (1882) from Lindl. (1848)	...	...	...	...	...	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.I,172	M.fr.I,177; X,71.		
C. polyandra; F. v. M., Cens. 27 (1882) from Hook. (1855)	...	...	...	...	...	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.I,172	M.fr.I,177; X,11.		
C. pleiopetala; F. v. M., Cens. 27 (1882)	...	...	...	...	...	—	S.A.	—	—	—	Q.	—	—	M.fr.X,70.		
C. quadrivalvis; F. v. M., Cens. 27 (1882)	...	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.I,173	M.fr.I,176.		

<i>C. liniflora</i> ; F. v. M., Cens. 27 (1882) from Fenzl (1837)	W.A.	—	—	—	—	—	B.fl.I,173	M.fr.X,71.
<i>C. gracilis</i> ; F. v. M., Cens. 27 (1882) from Benth. (1863)	—	—	—	—	—	N.A.	B.fl.I,173	
<i>C. polypetala</i> ; F. v. M., Cens. 27 (1882) from Fenzl (1837)	W.A.	—	—	—	—	—	B.fl.I,174	
<i>C. Pickeringi</i> ; F. v. M., Cens. 27 (1882) from A. Gray (1854)	—	—	—	—	N.S.W.	—	—	M.fr.X,69; XI,136.
<i>C. volubilis</i> ; F. v. M., Cens. 27 (1882) from Benth. (1863)... ..	W.A.	S.A.	—	V.	N.S.W.	Q.	B.fl.I,174	M.fr.X,68; XI,27.
<i>C. calyptrata</i> ; F. v. M., fragm. III, 89 (1862)	W.A.	S.A.	T.	V.	N.S.W.	Q.	B.fl.I,174	M.fr.I,175,177; IV,175; X,68.
<i>C. pumila</i> ; F. v. M., Cens. 27 (1882)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	— M.fr.X,68.
<i>C. composita</i> ; F. v. M., Cens. 27 (1882) from Nees (1844)... ..	W.A.	—	—	—	—	—	B.fl.I,175	M.fr.X,69.
<i>C. corrigioloides</i> ; F. v. M., Cens. 27 (1882)	W.A.	S.A.	—	V.	N.S.W.	—	B.fl.I,175	M.fr.X,69; XI,83.
<i>C. brevipedata</i> ; F. v. M., Cens. 27 (1882)	W.A.	S.A.	—	V.	—	—	—	M.fr.X,69; IX,136.
<i>C. granulifera</i> ; F. v. M., Cens. 27 (1882) from Benth. (1863)	W.A.	—	—	—	—	—	B.fl.I,176	
<i>C. pogonophora</i> ; F. v. M., Cens. 27 (1882)	—	—	—	—	—	Q.	—	M.fr.X,69.
<i>C. pygmaea</i> ; F. v. M., fragm. III, 89 (1862)... ..	W.A.	S.A.	—	V.	N.S.W.	—	B.fl.I,176	M.fr.I,175; X,71; XI,83.
<i>C. Australasica</i> ; J. Hooker, icon. plant. t. 293 (1840)	W.A.	S.A.	T.	V.	N.S.W.	—	B.fl.I,177	M.fr.III,89; X,71 ... N.Z.

MONTIA; Micheli, nov. pl. gen. 17, t. 13 (1729).

<i>M. fontana</i> ; Linné, spec. plant. 82 (1753)	—	—	T.	V.	N.S.W.	—	B.fl.I,177	 E., As., Am., N.Z.
--	---	---	----	----	--------	---	------------	---------------------------

CARYOPHYLLEAE.

LINNE, Philosoph. Bot. 31 (1751).—Genera 10, Species 26.

STELLARIA; Linné, syst. nat. ed. VI, (1748).

<i>S. pungens</i> ; Brongniart, Voy. sur la Coquille, t. 78 (1826)	—	S.A.	T.	V.	N.S.W.	—	B.fl.I,157	
<i>S. palustris</i> ; Retzius, Fl. Scandinav., prodr. sec. ed. n. 548 (1779)	—	S.A.	T.	V.	N.S.W.	Q.	B.fl.I,158	 E., As., Am.
<i>S. flaccida</i> ; Hooker, Comp. Bot. Mag. I, 275 (1835)	—	—	T.	V.	N.S.W.	Q.	B.fl.I,158	
<i>S. multiflora</i> ; Hooker, Comp. Bot. Mag. I, 275 (1835)	W.A.	S.A.	T.	V.	N.S.W.	—	B.fl.I,159	M.fr.IX,27.

SAGINA; Linné, syst. nat. 8 (1735); gen. pl. 118 (1737).

<i>S. procumbens</i> ; Linné, spec. plant. 128 (1753)	—	—	—	V.	N.S.W.	—	B.fl.I,160	 E., As., Afr. Am.
<i>S. apetala</i> ; Arduino, animadv. Bot. spec. alt. 22, t. 8 (1764)	W.A.	S.A.	T.	V.	—	—	B.fl.I,160	 E., As., Afr.

COLOBANTHUS; Bartling, Reliq. Haenk. II, 13, t. 49 (1830).

<i>C. Benthamianus</i> ; Fenzl in Ann. des Wien. Mus. I, 49 (1838)	—	—	—	V.	N.S.W.	—	B.fl.I,160	 Am., N.Z.
<i>C. Billardieri</i> ; Fenzl in Ann. des Wien. Mus. I, 48 (1838)... ..	—	—	T.	V.	—	—	B.fl.I,161	

SCLERANTHUS; Linné, gen. plant. 130 (1737). (Mniarum, Ditoca.)

<i>S. pungens</i> ; R. Brown, prodr. 412 (1810)	—	S.A.	—	V.	N.S.W.	—	B.fl.V,260	
<i>S. diander</i> ; R. Brown, prodr. 412 (1810)	—	S.A.	T.	V.	N.S.W.	—	B.fl.V,260	
<i>S. biflorus</i> ; J. Hooker, Fl. N. Zeal. I, 74 (1853)	—	—	T.	V.	N.S.W.	Q.	B.fl.V,259	 N.Z.
<i>S. mniaroides</i> ; F. v. M., pl. of Vict. I, 215 (1862)	—	—	—	V.	N.S.W.	—	B.fl.V,259	

SAPONARIA; Linné, gen. plant. 130 (1737) from Bock (1529). (Gypsophila, Dichoglottis.)

<i>S. tubulosa</i> ; F. v. M., native pl. of Vict. I, 136 (1879)	—	S.A.	—	V.	N.S.W.	Q.	B.fl.I,155	
---	---	------	---	----	--------	----	------------	--

SPERGULARIA; Persoon, synops. pl. I, 504 (1805). (Tissa (1763), Lepigonum, Arenaria partly.)

<i>S. rubra</i> ; Cambessèdes in St. Hilaire, Fl. Brazil. II, 179 (1829)	W.A.	S.A.	T.	V.	N.S.W.	Q.	B.fl.I,162	M.fr.XI,27... .. E., As., Am.
---	------	------	----	----	--------	----	------------	-------------------------------

DRYMARIA; Willdenow in Roemer & Schultes, syst. V, 406 (1819).

<i>D. filiformis</i> ; Bentham, Fl. Austr. I, 162 (1863)	W.A.	S.A.	—	—	—	—	B.fl.I,162	
<i>D. diandra</i> ; Blume, Bijdr. 63 (1823)	—	—	—	—	—	Q.	—	 As.

HERNIARIA; Tournefort, instit. rei. herb. 507, t. 288 (1700).

<i>H. incana</i> ; Lamarck, encyl. meth. III, 124 (1789)	—	S.A.	—	—	—	—	—	 E., As., Afr.
---	---	------	---	---	---	---	---	----------------------

POLYCARPON; Loefling, iter Hispanic. 7 (1758).

P. tetraphyllum; Loefling in Linné, syst. nat. ed. X, 881 (1759) W.A. S.A. T. V. N.S.W. Q. — B.fl.I, 163 E., As., Afr.

POLYCARPAEA; Lamarck in Journ. d'hist. nat. II, 8, t. 25 (1792). (Aylmeyera.)

P. longiflora; F. v. M., Rep. of Babb. Exped. 8 (1858) W.A. — — — — — N.A. B.fl.I, 164
P. spirostylis; F. v. M., Rep. of Babb. Exped. 8 (1858) — — — — — Q. N.A. B.fl.I, 165
P. synandra; F. v. M., Rep. of Babb. Exped. 8 (1858) — S.A. — — — — — Q. N.A. B.fl.I, 165
P. violacea; Benth. Fl. Austr. I, 165 (1863) — — — — — — N.A. B.fl.I, 165
P. Indica; Lamarck in Journ. d'hist. nat. II, 8 (1792) — S.A. — — — N.S.W. Q. N.A. B.fl.I, 166 As., Afr., Am.
P. breviflora; F. v. M., Rep. of Babb. Exped. 9 (1858) — — — — — Q. N.A. B.fl.I, 166
P. spicata; Arnott in Ann. of nat. hist. III, 91 (1839) — — — — — — N.A. B.fl.I, 167
P. involucrata; F. v. M., Rep. of Babb. Exped. 9 (1858) — — — — — — N.A. B.fl.I, 167

AMARANTACEAE.

A. L. DE JUSSIEU, gen. plant. 87 (1789).—Genera 12, Species 100.

GOMPHRENA; Linné, gen. plant. 69 (1737). (Philoxerus.)

G. canescens; R. Brown, prodr. 416 (1810) W.A. — — — — — Q. N.A. B.fl.V, 253
G. flaccida; R. Brown, prodr. 416 (1810) — — — — — Q. N.A. B.fl.V, 254 M.fr.III, 123.
G. affinis; F. v. M. in Benth. Fl. Austr. V, 254 (1870) — — — — — — N.A. B.fl.V, 254
G. humilis; R. Brown, prodr. 416 (1810) — — — — — Q. N.A. B.fl.V, 254
G. platandra; F. v. M. in Trans. R. S. Vict. XXIII, 52 (1886) W.A. — — — — — — N.A. —
G. lanata; R. Brown, prodr. 416 (1810) — S.A. — — — — — Q. N.A. B.fl.V, 255
G. brachystylis; F. v. M., fragm. III, 124 (1862) — — — — — — N.A. B.fl.V, 255 M.fr.III, 124.
G. leptoclada; Benth. Fl. Austr. V, 255 (1870) — — — — — — N.A. B.fl.V, 255
G. Maitlandi; F. v. M., fragm. III, 124, t. 23 (1862) W.A. — — — — — — N.A. B.fl.V, 256 M.fr.VI, 234.
G. pusilla; Benth. Fl. Austr. V, 256 (1870) — — — — — — N.A. B.fl.V, 256
G. tenella; Benth. Fl. Austr. V, 256 (1870) — — — — — — N.A. B.fl.V, 256
G. conica; Sprengel, syst. I, 824 (1825) — — — — — Q. N.A. B.fl.V, 256 M.fr.III, 125.
G. conferta; Benth. Fl. Austr. V, 257 (1870) W.A. — — — — — Q. N.A. B.fl.V, 257
G. diffusa; Sprengel, syst. I, 824 (1825) — — — — — Q. N.A. B.fl.V, 257
G. parviflora; Benth. Fl. Austr. V, 257 (1870) — — — — — — N.A. B.fl.V, 257

ALTERNANTHERA; Forskael, flor. Aegypt. Arab. 28 (1775). (Telanthera.)

A. triandra; Lamarck, encycl. meth. I, 95 (1783) W.A. S.A. T. V. N.S.W. Q. N.A. B.fl.V, 249 As., Afr.
A. nana; R. Brown, prodr. 417 (1810) — S.A. — — — N.S.W. Q. N.A. B.fl.V, 250
A. angustifolia; R. Brown, prodr. 417 (1810) — — — — — Q. N.A. B.fl.V, 250
A. decipiens; Benth. Fl. Austr. V, 251 (1870) — — — — — Q. — B.fl.V, 251
A. polycephala; Benth. Fl. Austr. V, 251 (1870) — — — — — — N.A. B.fl.V, 251
A. leptophylla; Benth. Fl. Austr. V, 251 (1870) — — — — — — N.A. B.fl.V, 251
A. longipes; Benth. Fl. Austr. V, 252 (1870) — — — — — — N.A. B.fl.V, 252

ACHYRANTHES; Linné, gen. plant. 34 (1737).

A. aspera; Linné, spec. plant. 204 (1753) W.A. S.A. — — — N.S.W. Q. N.A. B.fl.V, 246 M.fr.IX, 77.
A. arborescens; R. Brown, prodr. 417 (1810) — — — — — N.S.W. — — — M.fr.IX, 169.

NYSSANTHES; R. Brown, prodr. 418 (1810).

N. erecta; R. Brown, prodr. 418 (1810) — — — — — N.S.W. Q. — B.fl.V, 247
N. diffusa; R. Brown, prodr. 418 (1810) — — — — — N.S.W. Q. — B.fl.V, 247

DIPTERANTHEMUM; F. v. M. in Wing's S. Sc. Rec. III, 281 (1884).

D. Crosslandi; F. v. M. in Wing's S. Sc. Rec. III, 281 (1884) ... W.A. — — — — —

PTILOIOTUS; R. Brown, prodr. 415 (1810). (Trichinium, Psilotrichum, Goniotriche, Hemisteirus, Arthrotrichum.)

P. conicus; R. Brown, prodr. 415 (1810) ...	—	—	—	—	—	N.A.	B.fl. V, 242	
P. Macleayi; F. v. M. in proceed. L. S. of N.S.W. sec. ser. III, 162 (1888) ...	—	—	—	—	—	N.A.	—	
P. corymbosus; R. Brown, prodr. 415 (1810) ...	—	—	—	—	—	Q. N.A.	B.fl. V, 243	M.fr. III, 125.
P. grandiflorus; F. v. M., fragm. I, 237 (1859) ...	W.A.	—	—	—	—	—	B.fl. V, 243	M.fr. I, 237.
P. spicatus; F. v. M. in Benth. Fl. Austr. V, 243 (1870) ...	—	—	—	—	—	Q. N.A.	B.fl. V, 243	
P. Murrai; F. v. M., fragm. III, 145 (1863) ...	W.A.	S.A.	—	—	—	N.A.	B.fl. V, 243	M.fr. III, 145.
P. gomphrenoides; F. v. M., fragm. VI, 233 (1868) ...	W.A.	—	—	—	—	N.A.	B.fl. V, 244	
P. latifolius; R. Brown in App. Sturt's Exped. 25 (1849) ...	W.A.	S.A.	—	—	—	Q. N.A.	B.fl. V, 244	M.fr. III, 145; VI, 232.
P. macrotrichus; F. v. M., fragm. IV, 90 (1864) ...	W.A.	—	—	—	—	N.A.	B.fl. V, 244	M.fr. VI, 232.
P. villosiflorus; F. v. M., fragm. III, 125 (1863) ...	—	—	—	—	—	N.A.	B.fl. V, 245	M.fr. III, 125.
P. humilis; F. v. M., fragm. VI, 229 (1868) ...	W.A.	—	—	—	—	—	B.fl. V, 245	M.fr. III, 161; VI, 229.
P. obovatus; F. v. M., fragm. VI, 228 (1868) ...	W.A.	S.A.	—	V.	N.S.W.	Q. N.A.	B.fl. V, 220	M.fr. XI, 27.
P. incanus; Poiret, Encycl. méth. suppl. IV, 620 (1817) ...	—	S.A.	—	—	—	N.A.	B.fl. V, 221	M.fr. VI, 228.
P. parviflorus; F. v. M., syst. Cens. of Austr. pl. 28 (1882) from Lindley (1838) ...	—	—	—	—	N.S.W.	Q. N.A.	B.fl. V, 222	
P. astrolasius; F. v. M., fragm. VI, 233 (1868) ...	—	—	—	—	—	N.A.	B.fl. V, 222	M.fr. VI, 233.
P. rotundifolius; F. v. M., fragm. VI, 230 (1868) ...	W.A.	—	—	—	—	—	B.fl. V, 223	M.fr. III, 122.
P. dissitiflorus; F. v. M., syst. Cens. of Austr. pl. 28 (1882) ...	—	—	—	—	—	N.A.	B.fl. V, 223	M.fr. IV, 89.
P. distans; Poiret, Encycl. suppl. IV, 620 (1817) ...	—	—	—	—	—	Q. N.A.	B.fl. V, 223	M.fr. VI, 228.
P. alopecuroides; F. v. M., fragm. VI, 227 (1868) ...	W.A.	S.A.	—	V.	N.S.W.	Q. N.A.	B.fl. V, 224	M.fr. VI, 227.
P. nobilis; F. v. M., fragm. VI, 227 (1868) ...	—	S.A.	—	V.	N.S.W.	—	B.fl. V, 224	M.fr. VI, 227.
P. macrocephalus; Poiret, Encycl. suppl. IV, 620 (1817) ...	—	S.A.	—	V.	N.S.W.	Q. —	B.fl. V, 225	M.fr. VI, 228; XI, 27.
P. hemisteirus; F. v. M., fragm. IV, 90 (1864) ...	W.A.	S.A.	—	—	N.S.W.	N.A.	B.fl. V, 226	M.fr. VI, 231.
P. parvifolius; F. v. M., fragm. VI, 229 (1868) ...	—	S.A.	—	—	N.S.W.	—	B.fl. V, 241	M.fr. VI, 229.
P. Polakii; F. v. M. in S. Sc. Rec. II, 12 (1882) ...	W.A.	—	—	—	—	—	—	
P. exaltatus; Nees in Lehm. pl. Preiss. I, 630 (1845) ...	W.A.	S.A.	—	V.	N.S.W.	Q. N.A.	B.fl. V, 227	
P. Manglesii; F. v. M., fragm. VI, 230 (1868) ...	W.A.	—	—	—	—	N.A.	B.fl. V, 228	M.fr. VI, 230.
P. Beckeri; F. v. M., fragm. VI, 233 (1868) ...	—	S.A.	—	—	—	—	B.fl. V, 228	
P. gomphrenoides; Moquin in De Cand., prodr. XIII, 2, 287 (1849) ...	W.A.	S.A.	—	—	—	—	B.fl. V, 229	
P. esquamatus; F. v. M., syst. Cens. of Austr. pl. 28 (1882) from Benth (1838) ...	W.A.	—	—	—	—	—	B.fl. V, 229	
P. declinatus; Nees in Lehm. pl. Preiss. I, 631 (1845) ...	W.A.	—	—	—	—	—	B.fl. V, 230	
P. erubescens; Schlechtendal in Linnaea XX, 575 (1847) ...	—	S.A.	—	V.	N.S.W.	—	B.fl. V, 230	M.fr. VI, 229.
P. divaricatus; F. v. M., fragm. VI, 229 (1868) ...	W.A.	—	—	—	—	N.A.	B.fl. V, 231	M.fr. VI, 229.
P. helipteroides; F. v. M., fragm. VI, 231 (1868) ...	W.A.	S.A.	—	—	—	N.A.	B.fl. V, 231	M.fr. III, 122 & 161.
P. Stirlingi; F. v. M., syst. Cens. of Austr. pl. 28 (1882) from Lindley (1839) ...	W.A.	—	—	—	—	N.A.	B.fl. V, 232	
P. laxus; F. v. M., syst. Cens. of Austr. pl. 28 (1882) from Benth (1870) ...	W.A.	—	—	—	—	—	B.fl. V, 232	
P. axillaris; F. v. M., syst. Cens. of Austr. pl. 28 (1882) ...	W.A.	—	—	—	—	N.A.	B.fl. V, 232	
P. striatus; F. v. M., syst. Cens. of Austr. pl. 28 (1882) from Moquin (1870) ...	W.A.	—	—	—	—	N.A.	B.fl. V, 233	
P. auriculifolius; F. v. M., Cens. of Austr. pl. 28 (1882) from Cunningham (1849) ...	—	—	—	—	—	N.A.	B.fl. V, 233	
P. sericostachyus; F. v. M., fragm. VI, 231 (1868) ...	W.A.	—	—	—	—	—	B.fl. V, 234	M.fr. VI, 230.
P. roseus; F. v. M., syst. Cens. of Austr., pl. 28 (1882) from Moquin (1849) ...	W.A.	—	—	—	—	—	B.fl. V, 234	
P. fusiformis; Poiret, Encycl. suppl. IV, 619 (1817) ...	—	—	—	—	—	Q. N.A.	B.fl. V, 234	
P. gracilis; Poiret, Encycl. suppl. IV, 620 (1817) ...	W.A.	—	—	—	—	Q. N.A.	B.fl. V, 235	M.fr. VI, 228.
P. Drummondii; F. v. M., fragm. VI, 229 (1868) ...	W.A.	—	—	—	—	—	B.fl. V, 235	M.fr. VI, 229.
P. calostachyus; F. v. M., fragm. VI, 231 (1868) ...	—	—	—	—	—	Q. N.A.	B.fl. V, 236	M.fr. VI, 231.
P. Fraseri; F. v. M., Cens. 29 (1882) from Cunningham (1849) ...	—	—	—	—	N.S.W.	—	B.fl. V, 236	
P. Carlsoni; F. v. M. in Vict. Nat. V, 74 (1888) ...	W.A.	—	—	—	—	—	—	
P. Sewelliae; F. v. M. in Vict. Nat. Jan. (1885) ...	W.A.	—	—	—	—	—	—	

A. Muelleri; Benth. Fl. Austr. V, 175 (1870)	...	...	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V, 175	M.fr.VII, 9.
A. semibaccatum; R. Brown, prodr. 406 (1810)	...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.V, 175	M.fr.VI, 19.
A. humile; F. v. M., fragm. IV, 48 (1866)	...	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.V, 174	M.fr.IV, 48.
A. prostratum; R. Brown, prodr. 406 (1810)	...	...	...	...	...	W.A.	S.A.	—	—	N.S.W.	—	—	B.fl.V, 176	M.fr.VII, 9; XII, 17.
A. angulatum; Benth. Fl. Austr. V, 174 (1870)	...	...	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.V, 174	
A. Quioii; F. v. M. in Vict. Nat. V, 96 (1888)	...	...	...	...	...	—	S.A.	—	—	N.S.W.	Q.	—		
A. stipitatum; Benth. Fl. Austr. V, 168 (1870)	...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.V, 168	M.fr.VII, 9.
A. paludosum; R. Brown, prodr. 406 (1810)	...	...	...	...	...	W.A.	S.A.	T.	V.	—	—	—	B.fl.V, 169	M.fr.VII, 9; XII, 16.
A. Moquinianum; Webb in De Candolle, prodr. XIII, pars II, 97 (1849)	...	...	...	...	...	W.A.	—	—	—	—	—	N.A.	B.fl.V, 169	
A. Drummondii; Moquin in De Candolle, prodr. XIII, pars II, 102 (1849)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V, 170	
A. cinereum; Poiret, Encycl. suppl. I, 471 (1810)	...	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.V, 171	M.fr.VII, 9; IX, 77.
A. rhagodioides; F. v. M. in Transact. phil. Inst. Vict. II, 74 (1857)	...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.V, 172	
A. incrassatum; F. v. M., Rep. Babb. Exp. 20 (1858)	...	...	...	...	...	—	[S.A.	—	—	—	—	—	B.fl.V, 172	
A. nummularium; Lindley in Mitch. Trop. Austr. 64 (1848)	...	...	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.V, 170	M.fr.VII, 10; XI, 27; XII, 17.
A. hymenothecum; Moquin in De Candolle, prodr. XIII, pars II, 101 (1849)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V, 173	M.fr.VII, 10.
A. vesicarium; Heward in Benth. Fl. Austr. V, 172 (1870)	...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V, 172	
A. balimoides; Lindley in Mitch. Three Exped. I, 282 (1838)	...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V, 179	M.fr.VII, 10; XI, 27; XII, 17.
A. spongiosum; F. v. M. in Transact. phil. Inst. Vict. II, 74 (1857)	...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V, 179	M.fr.XII, 17.

RHAGODIA; R. Brown, prodr. 408 (1810).

R. Billardieri; R. Brown, prodr. 408 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.V, 152	M.fr.VII, 10; IX, 77; XII, 16.
R. parabolica; R. Brown, prodr. 408 (1810)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.V, 153	M.fr.VII, 10; XII, 15.
R. dioica; Nees in Lehm. pl. Preiss. I, 636 (1845)	W.A.	—	—	—	—	—	—	B.fl.V, 154	
R. Gaudichaudiana; Moquin, Chenop. monogr. enum. 11 (1840)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.V, 154	M.fr.VII, 10; XII, 15.
R. crassifolia; R. Brown, prodr. 408 (1810)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V, 154	M.fr.XII, 15.
R. Pressii; Moquin in De Candolle, prodr. XIII, part II, 49 (1849)	W.A.	S.A.	—	—	—	—	—	B.fl.V, 155	M.fr.VII, 10; XII, 15.
R. obovata; Moquin, Chenop. monogr. enum. 10 (1840)	W.A.	—	—	—	—	—	—	B.fl.V, 155	
R. spinescens; R. Brown, prodr. 408 (1810)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V, 155	M.fr.VII, 10; XI, 20, 75; XII, 15.
R. hastata; R. Brown, prodr. 408 (1810)	—	—	—	V.	N.S.W.	Q.	—	B.fl.V, 156	M.fr.VII, 10.
R. nutans; R. Brown, prodr. 408 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.V, 156	M.fr.VII, 10; XII, 15.
R. linifolia; R. Brown, prodr. 408 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.V, 157	M.fr.VII, 10; XII, 15.

CHENOPODIUM; Tournefort, inst. 506, t. 288 (1700). (Blitum, Ambrina.)

C. nitrariaceum; F. v. M. in Benth. Fl. Austr. V, 158 (1870)	...	...	...	...	...	W.A.	S.A.	— V.	N.S.W.	Q.	—	B.fl.V, 158	M.fr.XI, 20&27; XII, 16.
C. auricomum; Lindley in Mitch. Trop. Austr. 94 (1848)	...	...	...	...	...	—	S.A.	— V.	N.S.W.	Q.	N.A.	B.fl.V, 159	M.fr.VII, 11.
C. triangulare; R. Brown, prodr. 407 (1810)	...	...	...	...	...	—	—	—	N.S.W.	Q.	—	B.fl.V, 160	M.fr.VII, 10.
C. microphyllum; F. v. M. in Transact. phil. Inst. Vict. II, 74 (1857)	...	...	...	...	...	—	S.A.	— V.	N.S.W.	—	—	B.fl.V, 161	M.fr.VII, 11; XII, 16.
C. atriplicinum; F. v. M., fragm. VII, 11 (1869)	...	...	...	...	...	—	S.A.	— V.	N.S.W.	—	N.A.	B.fl.V, 163	M.fr.I, 140; XI, 27.
C. cristatum; F. v. M., fragm. VII, 11 (1869)	...	...	...	...	...	W.A.	S.A.	— V.	N.S.W.	Q.	—	B.fl.V, 163	M.fr.XI, 27; IX, 75; XII, 17.
C. carinatum; R. Brown, prodr. 407 (1810)	...	...	...	...	...	W.A.	S.A.	— V.	N.S.W.	Q.	N.A.	B.fl.V, 162	M.fr.VII, 11 ... N.Z., P.
C. rhadinostachyum; F. v. M. in Wing's S. Sc. Rec. May (1880)	...	...	...	...	...	—	S.A.	—	—	—	—	—	M.fr.XII.
C. simulans, F. v. M. & Tate; Dysphania, F. v. M. & T. in proc. Roy. Soc. S.A. VIII, 71 (1885)	...	...	...	...	...	—	S.A.	—	—	—	—	—	

DYSPHANIA; R. Brown, prodr. VI, 411 (1810).

D. plantaginella; F. v. M., fragm. I, 61 (1858)	...	...	...	...	...	W.A.	S.A.	—	—	—	—	N.A.	B.fl.V, 164	M.fr.IX, 75
D. litoralis; R. Brown, prodr. 411 (1810)	...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V, 164	M.fr.VII, 11; XII, 17.

KOCHIA; Roth in Schrader's Journ. Bot. I, 303 (1799). (Maireana, Sclerochlamys.)

K. fimbriolata; F. v. M., fragm. IX, 75 (1875)	...	...	...	...	...	—	S.A.	—	—	—	—	—	M.fr.IX, 75.
K. lobiflora; F. v. M. in Benth. Fl. Austr. V, 184 (1870)	...	...	...	...	...	—	S.A.	—	N.S.W.	—	—	B.fl.V, 184	M.fr.IX, 75.
K. lanosa; Lindley in Mitch. Trop. Austr. 88 (1848)	...	...	...	...	...	W.A.	S.A.	— V.	N.S.W.	Q.	—	B.fl.V, 184	M.fr.VII, 13; XII, 15
K. prosthocochaeta; F. v. M., fragm. XII, 14 (1882)	...	...	...	...	...	W.A.	—	—	—	—	—	—	M.fr.XII, 14.
K. melanocoma; F. v. M., fragm. XII, 14 (1882)	...	...	...	...	...	W.A.	—	—	—	—	—	—	M.fr.XII, 14.

K. triptera; Benth, Fl. Austr. V, 185 (1870)	W.A. S.A. — V. N.S.W. Q. N.A. B.fl.V, 185	
K. oppositifolia; F. v. M. in Transact. Vict. Inst. 134 (1855)	W.A. S.A. — V. N.S.W. — — B.fl.V, 185	M.fr.VII, 12; XII, 15.
K. brevifolia; R. Brown, prodr. 409 (1810)	W.A. S.A. — V. N.S.W. Q. — — B.fl.V, 185	M.fr.VII, 13.
K. pyramidata; Benth, Fl. Austr. V, 186 (1870)	— S.A. — V. N.S.W. — — B.fl.V, 186	M.fr.XII, 15.
K. spongiocarpa; F. v. M. in Vict. Nat. III, 92, (1886)	W.A. S.A. — — N.S.W. Q. — —	
K. eriantha; F. v. M., Rep. Babb. Exped. 20 (1858)	— S.A. — — N.S.W. — — B.fl.V, 186	M.fr.VII, 13.
K. villosa; Lindley in Mitch. Trop. Austr. 91 (1848)	W.A. S.A. — V. N.S.W. Q. N.A. B.fl.V, 186	M.fr.VII, 12; IX, 75; XII, 15.
K. planifolia; F. v. M., fragm. I, 213 (1859)	W.A. — — — — — B.fl.V, 187	M.fr.VII, 12.
K. sedifolia; F. v. M. in Transact. Vict. Inst. I, 134 (1855)	W.A. S.A. — V. N.S.W. Q. N.A. B.fl.V, 187	M.fr.VII, 12; VIII, 37; XII, 15.
K. humillima; F. v. M., fragm. IX, 168 (1875)	— S.A. — V. N.S.W. — — —	M.fr.XII, 15.
K. microphylla; F. v. M., fragm. VIII, 148 (1874)	— — — V. N.S.W. Q. — — B.fl.V, 181	M.fr.VIII, 37; XII, 15.
K. dichoptera; F. v. M., fragm. VIII, 37 (1873)	— — — — — Q. — — —	M.fr.VIII, 163.
K. decaptera; F. v. M., fragm. IX, 75 (1875)	— S.A. — — — — —	M.fr.IX, 75.
K. ciliata; F. v. M., Rep. Babb. Exped. 20 (1858)	— S.A. — V. N.S.W. — — B.fl.V, 188	M.fr.VII, 12; XII, 15.
K. brachyptera; F. v. M., second gen. Rep. 15 (1854)	— S.A. — V. N.S.W. Q. — — B.fl.V, 189	M.fr.VII, 13; XII, 15.
K. stelligera; F. v. M., fragm. VII, 13 (1869)	— S.A. — V. N.S.W. Q. — — B.fl.V, 189	M.fr.I, 139; XII, 15.

DIDYMANTHUS; Endlicher, nov. stirp. decad. 7 (1839).

D. Roei; Endlicher, nov. stirp. decad. 8 (1839)	W.A. — — — — — N.A. B.fl.V, 193	M.fr.VII, 11.
--	---------------------------------	---------------

BABBAGIA; F. v. M., Rep. on pl. of Babb. Exped. 21 (1858).

B. dipterocarpa; F. v. M., Rep. of Babb. Exped. 21 (1858)	— S.A. — — N.S.W. Q. N.A. B.fl.V, 192	M.fr.VII, 11.
B. scleroptera; F. v. M. in Wing's S. Sc. Rec. Nov. (1855)	— — — — — N.S.W. Q. — —	
B. acroptera; F. v. M. & Tate in Transact. Roy. Soc. of S. Austr. VI, 108 (1883)	— S.A. — — — — — Q. N.A. —	
B. pentaptera; Tate in Transact. Roy. Soc. of S. Austr. VI, 108 (1883)	— S.A. — — — — —	

BASSIA; Allioni in Misc. Taurin. III, 117, t. 4 (1766). (Chenolea, Sclerolaena, Anisacantha, Echinopsilon, Kentropsia, Dissocarpus, Eriochiton, Osteocarpum, Coilocarpus.)

B. carnosa; F. v. M., Cens. of Austr. pl. 30 (1882) from Benth (1870)	W.A. — — — — — B.fl.V, 190	M.fr.VIII, 13; XII, 13.
B. Dallachyana; F. v. M., Cens. of Austr. pl. 30 (1882) from Benth (1870)	— — — V. N.S.W. — — B.fl.V, 191	
B. tricornis; F. v. M., Cens. of Austr. pl. 30 (1882) from Benth (1870)	W.A. S.A. — — — — — B.fl.V, 191	
B. tridens; F. v. M., fragm. XII, 12 (1882)	W.A. — — — — — —	M.fr.XII, 12.
B. enrotioides; F. v. M., Cens. of Austr. pl. 30 (1882)	W.A. — — — — — B.fl.V, 191	
B. Muelleri; F. v. M., Cens. of Austr. pl. 30 (1882) from Benth (1870)	— — — — — N.A. B.fl.V, 191	M.fr.XII, 13.
B. astrocarpa; F. v. M., fragm. XII, 12 (1882)	W.A. — — — — — —	M.fr.XII, 12.
B. sclerolaenoides; F. v. M., Cens. of Austr. pl. 30 (1882)	W.A. S.A. — V. N.S.W. Q. — B.fl.V, 192	M.fr.VII, 13; XII, 13.
B. Cornishiana; F. v. M. in Austr. Chemist (1885)	— S.A. — — — — —	
B. uniflora; F. v. M., Cens. of Austr. pl. 30 (1882) from R. Brown (1810)	W.A. S.A. — — — — — B.fl.V, 194	M.fr.VIII, 38.
B. diancantha; F. v. M., Cens. of Austr. pl. 30 (1882) from Nees (1844)	W.A. S.A. — V. N.S.W. Q. N.A. B.fl.V, 194	M.fr.VII, 14; IX, 75; X, 91.
B. lanicuspis; F. v. M., Cens. of Austr. pl. 30 (1882)	W.A. S.A. — — N.S.W. Q. N.A. B.fl.V, 195	M.fr.II, 140, 170; VII, 14; IX, 27; XII, 13.
B. bicornis; F. v. M., Cens. of Austr. pl. 30 (1882) from Lindley (1838)	— S.A. — V. N.S.W. Q. N.A. B.fl.V, 195	M.fr.I, 139; VII, 14; XII, 13.
B. biflora; F. v. M., Cens. of Austr. pl. 30 (1882) from R. Brown (1838)	— S.A. — V. N.S.W. Q. — B.fl.V, 196	M.fr.VII, 11; VIII, 38; IX, 27; X, 91; XII, 13.
B. paradoxa; F. v. M., Cens. of Austr. pl. 30 (1882) from R. Brown (1810)	W.A. S.A. — V. N.S.W. Q. — B.fl.V, 196	M.fr.VII, 11; VIII, 38, 149; X, 91; XII, 13.
B. quinquecuspidis; F. v. M., Cens. of Austr. pl. 30 (1882)	— S.A. — V. N.S.W. Q. — B.fl.V, 199	M.fr.VII, 14; X, 91.
B. Forrestiana; F. v. M., fragm. XII, 12 (1882)	W.A. — — — — — —	M.fr.XII, 12.
B. Drummondii; F. v. M., Cens. of Austr. pl. 30 (1882) from Benth (1870)	W.A. — — — — — B.fl.V, 199	
B. Birchii; F. v. M., Cens. of Austr. pl. 30 (1882)	— — — — — Q. — —	M.fr.VIII, 163.
B. divaricata; F. v. M., Cens. of Austr. pl. 30 (1882) from R. Brown (1810)	— S.A. — V. N.S.W. Q. — B.fl.V, 200	M.fr.VII, 14; X, 92; XII, 13.
B. bicuspidis; F. v. M., Cens. of Austr. pl. 30 (1882)	— S.A. — — N.S.W. Q. — B.fl.V, 200	M.fr.VII, 13.
B. glabra; F. v. M., Cens. of Austr. pl. 30 (1882)	— — — — — N.A. B.fl.V, 200	M.fr.I, 139.
B. echinopsila; F. v. M., Cens. of Austr. pl. 30 (1882)	W.A. S.A. — V. N.S.W. Q. N.A. B.fl.V, 201	M.fr.I, 139; VII, 14; X, 92; XII, 13.
B. cnchylaenoides; F. v. M., Cens. of Austr. pl. 30 (1882)	— S.A. — V. N.S.W. Q. — B.fl.V, 192	M.fr.X, 92; XII, 13.

- B. micrantha; F. v. M., Cens. of Austr. pl. 30 (1882) from Benth (1870) ... W.A. — — — — — B.fl.V,181
 B. brevicuapis; F. v. M., Cens. of Austr. pl. 30 (1882) ... — — — — N.S.W. Q. — B.fl.V,198 M.fr.IV,150; VII,14; VII,38.
 B. salsuginosa; F. v. M., Cens. of Austr. pl. 30 (1882) ... — S.A. — V. N.S.W. — — B.fl.V,197 M.fr.VII,12; X,92; XI,27; XII,13.

THRELKELDIA; R. Brown, prodr. 409 (1810).

- T. proceriflora; F. v. M., fragm. VIII, 38 (1873) ... — — — — N.S.W. Q. — — M.fr.VIII,38.
 T. diffusa; R. Brown, prodr. 410 (1810) ... W.A. S.A. T. V. N.S.W. — — B.fl.V,197 M.fr.VII,12.

ENCHYLAENA; R. Brown, prodr. 408 (1810).

- E. tomentosa; R. Brown, prodr. 408 (1810) ... W.A. S.A. — V. N.S.W. Q. N.A. B.fl.V,181 M.fr.VII,12; IX,75; XI,27; XII,15.

SALICORNIA; Tournefort, coroll. 51 (1703). (Halocnemum, Arthrocnemum, Tecticornia, Pachycornia) from Gerard (1597).

- S. robusta; F. v. M., fragm. VI, 251 (1868) ... W.A. S.A. — V. N.S.W. — N.A. B.fl.V,202 M.fr.I,139; VII,14; XII,18.
 S. arbuscula; R. Brown, prodr. 411 (1810) ... W.A. S.A. T. V. N.S.W. Q. N.A. B.fl.V,203 M.fr.VII,15; XII,18.
 S. cinerea; F. v. M., fragm. VI, 251 (1869) ... — — — — Q. N.A. B.fl.V,203 M.fr.I,140; VII,15; XII,18.
 S. leiostachya; Benth, Fl. Austr. V, 203 (1870) ... W.A. S.A. — — N.S.W. Q. N.A. B.fl.V,203
 S. bidens; Benth, Fl. Austr. V, 204 (1870) ... W.A. — — — — — B.fl.V,204
 S. tenuis; Benth, Fl. Austr. V, 204 (1870) ... — S.A. — — N.S.W. Q. — B.fl.V,204
 S. Australis; Solander in G. Forster, prodr. 88 (1786) ... W.A. S.A. T. V. N.S.W. Q. N.A. B.fl.V,205 M.fr.VII,15; IX,75; XI,20,27; XII,18. N.Z.

SUAEDA; Forskael, Fl. Aegypt. Arab. 69 (1779). (Chenopodium parly, Schoberia, Chenopodina.)

- S. maritima; Dumortier, Fl. Belg. 22 (1827) ... W.A. S.A. T. V. N.S.W. Q. N.A. B.fl.V,206 M.fr.XI,20. ... E., As., Afr., Am., P., N.Z.

SALSOLA; Linné, gen. pl. 67 (1737), from Cissalpinii.

- S. Kali; Linné, spec. plant. 222 (1753) ... W.A. S.A. — V. N.S.W. Q. N.A. B.fl.V,207 M.fr.IX,75; XI,20,27; XII,18. E., As., Afr., [Am., P.]

FICOIDEAE.

A. L. DE JUSSIEU, Gen. plant. 315 (1789).—Genera 9, Species 28.

MESEMBRIANTHEMUM; Breyne, prodr. fasc. rar. pl. II, 67 (1689).

- M. aequilaterale; Haworth, misc. nat. 77 (1803) ... W.A. S.A. T. V. N.S.W. Q. N.A. B.fl.III,324 M.fr.V,158; XI,20,27. ... Am.
 M. Australe; Solander in G. Forster, prodr. 90 (1786) ... W.A. S.A. T. V. N.S.W. Q. N.A. B.fl.III,324 M.fr.V,158; X,83; XI,20,27. N.Z.

TETRAGONIA; Linné, syst. nat. 9 (1735); gen. pl. 144 (1737). (Tetragonella.)

- T. expansa; Murray in Comm. Goetting. VI, 13, t. 5 (1783) ... W.A. S.A. T. V. N.S.W. Q. — B.fl.III,325 M.fr.VII,129; X,84; XI,27. As., Am., P., N.Z.
 T. diptera; F. v. M., fragm. XI, 8 (1878) ... W.A. — — — — N.A. — M.fr.XI,8.
 T. implexicoma; J. Hooker, Fl. Tasman. I, 148 (1859) ... W.A. S.A. T. V. N.S.W. — N.A. B.fl.III,326 M.fr.X,84; XI,8 ... N.Z.

AIZOON; Linné, gen. pl. 161 (1737). (Aizoon, Gunniopsis.)

- A. quadrifidum; F. v. M., fragm. II, 148 (1861) ... W.A. S.A. — — N.S.W. Q. — B.fl.III,327 M.fr.VII,129; X,83.
 A. zygophylloides; F. v. M., fragm. VII, 129 (1871) ... W.A. S.A. — — N.S.W. Q. — — M.fr.X,83.

GUNNIA; F. v. M., Rep. on pl. of Babb. Exped. 9 (1858).

- G. septifraga; F. v. M., Rep. of Babb. Exped. 9 (1858) ... — S.A. — — N.S.W. Q. — B.fl.III,327
 G. Drummondii; Benth, Fl. Austr. III, 327 (1866) ... W.A. — — — — — B.fl.III,327

SESUVIUM; Linné, syst. ed. decim. 1058 (1759).

- S. portulacastrum; Linné, syst. ed. dec. 1058 (1759) ... — — — — N.S.W. Q. N.A. B.fl.III,328 M.fr.III,164; VII,129; IX,77; X,83. As., [Afr., Am]

ZALEYA; N. L. Burmann, Fl. Indic. 110, t. 31 (1768).

- Z. decandra; N. L. Burmann, Fl. Ind. 110, t. 31 (1768) ... W.A. S.A. — — N.S.W. Q. N.A. B.fl.III,329 M.fr.I,172; III,164; X,83. As.

TRIANTHEMA; Sauvage, Méth. fol. 127 (1751). (Ancistrostigma.)

- T. turgidifolia; F. v. M., fragm. X, 83 (1876) ... — S.A. — — — Q. N.A. — M.fr.X,83.
 T. crystallina; Vahl, symbol. I, 32 (1790) ... W.A. S.A. — — N.S.W. Q. N.A. B.fl.III,330 M.fr.I,171-2. As., Afr.
 T. glaucifolia; F. v. M., fragm. I, 172 (1859) ... W.A. — — — — Q. — — M.fr.I,172.

<i>T. pilosa</i> ; F. v. M., fragm. I, 174 (1859)	W.A. S.A. — — —	Q. N.A. B.fl.III,330 M.fr.X,83.
<i>T. oxycalyptra</i> ; F. v. M., fragm. I, 173 (1859)	— — — — —	— N.A. B.fl.III,330 M.fr.I,173.
<i>T. rhynchocalyptra</i> ; F. v. M., fragm. I, 174 (1859)	— — — — —	Q. N.A. B.fl.III,330 M.fr.I,174.
<i>T. glossostigma</i> ; F. v. M. in Wing's S. Sc. Rec. III, 282 (1884)	W.A. — — — — —	— — —
<i>T. cypseloides</i> ; Benth., Fl. Austr. III, 331 (1866)	— — — — — N.S.W.	Q. N.A. B.fl.III,331 M.fr.X,83.

MACARTHURIA; Huegel, enum. pl. Nov. Holl. Austr. occ. II (1837).

<i>M. apetal</i> ; Harvey in Hook. Kew. Misc. VII, 55 (1855)	W.A. — — — — —	— N.A. B.fl.III,332 M.fr.XI,20.
<i>M. Australis</i> ; Huegel, enum. pl. II (1837)	W.A. — — — — —	— B.fl.III,332 M.fr.II,11; V,28.
<i>M. Neo-Cambrica</i> ; F. v. M., fragm. V, 28 (1866)	— — — — — N.S.W.	Q. — B.fl.III,332 M.fr.V,28.

MOLLUGO; Linné, gen. plant. 336 (1737). (Glinus, Trigastrotheca.)

<i>M. hirta</i> ; Thunberg, prodr. pl. cap. I, 24 (1794)	W.A. S.A. T. V. N.S.W.	Q. N.A. B.fl.III,333	E., As., Afr., Am.
<i>M. orygioides</i> ; F. v. M. in Benth. Fl. Austr. III, 333 (1866)	— S.A. — — N.S.W.	Q. — B.fl.III,333	
<i>M. Spergula</i> ; Linné, spec. pl. ed. alt. 131 (1762)	W.A. S.A. — V. N.S.W.	Q. N.A. B.fl.III,334 M.fr.II,148	As., Afr.
<i>M. stricta</i> ; Linné, spec. pl. ed. alt. 31 (1762)	— — — — —	Q. — —	As., P.
<i>M. trigastrotheca</i> ; F. v. M., pl. of Vict. I, 201 (1860)	W.A. — — — — —	— N.A. B.fl.III,334	
<i>M. Cerviana</i> ; Séringe in De Candolle, prodr. I, 392 (1824)	W.A. S.A. — V. N.S.W.	Q. N.A. B.fl.III,334 M.fr.II,148	E., As., Afr.

POLYGONACEAE.

A. L. DE JUSSIEU, Gen. pl. 82 (1789), from B. de Jussieu (1759).—Genera 4, Species 25.

EMEX; Necker, elem. bot. II, 214 (1790). (Perhaps immigrated.)

<i>E. Australis</i> ; Steinheil in Ann. des sc. nat. ser. 2, IX, 195, t. 7 (1839)	W.A. S.A. — — — — —	B.fl.V,262	Afr.
--	---------------------	-------------------	------

RUMEX; Linné, gen. pl. 105 (1737), from Plinius.

<i>R. Brownii</i> ; Campdera, Monogr. Rum. 81 (1819)	— S.A. T. V. N.S.W.	Q. — B.fl.V,263	
<i>R. flexuosus</i> ; Solander in G. Forst., fl. ins. Austr. prodr. 90 (1786)	— S.A. — — N.S.W.	— — B.fl.V,264	N.Z.
<i>R. crystallinus</i> ; Lange, ind. sem. hort. Hafn. 28 (1861)	W.A. S.A. — V. N.S.W.	Q. N.A. B.fl.V,265 M.fr.IV,48.	
<i>R. bidens</i> ; R. Brown, prodr. 421 (1810)	— S.A. T. V. N.S.W.	Q. — B.fl.V,265	

POLYGONUM; l'Obel, stirp. hist. 228 (1576), from Dioscorides and Plinius.

<i>P. plebejum</i> ; R. Brown, prodr. 420 (1810)	— S.A. — V. N.S.W.	Q. N.A. B.fl.V,267	E., As., Afr.
<i>P. strigosum</i> ; R. Brown, prodr. 420 (1810)	— — T. V. N.S.W.	Q. — B.fl.V,268	As., N.Z.
<i>P. prostratum</i> ; R. Brown, prodr. 419 (1810)	W.A. S.A. T. V. N.S.W.	Q. — B.fl.V,268	
<i>P. hydropiper</i> ; Linné, spec. plant. 361 (1753)	— S.A. — V. N.S.W.	Q. — B.fl.V,269	E., As., Afr., Am.
<i>P. minus</i> ; Hudson, Fl. Anglica I, 148 (1762)	W.A. S.A. T. V. N.S.W.	Q. — B.fl.V,269	E., As., Afr.
<i>P. subscissile</i> ; R. Brown, prodr. 419 (1810)	— — T. V. N.S.W.	Q. — B.fl.V,269	P.
<i>P. barbatum</i> ; Linné, spec. plant. 362 (1753)	— — — — N.S.W.	Q. N.A. B.fl.V,270	As., Afr.
<i>P. articulatum</i> ; R. Brown, prodr. 420 (1810)	— — — — —	Q. — B.fl.V,270	
<i>P. lapathifolium</i> ; Linné, spec. plant. 360 (1753)	— S.A. T. V. N.S.W.	Q. — B.fl.V,270	E., As., Afr., Am.
<i>P. orientale</i> ; Linné, spec. plant. 362 (1753)	— — — — N.S.W.	Q. — B.fl.V,271	As., P.
<i>P. attenuatum</i> ; R. Brown, prodr. 420 (1810)	— S.A. — — N.S.W.	Q. N.A. B.fl.V,272	Am.

MUEHLENBECKIA; Meissner, gen. pl. vasc. 316 (1840). (Sarcogonum, Coccoloba partly.)

<i>M. adpressa</i> ; Meissner, gen. pl. comm. 227 (1843)	W.A. S.A. T. V. N.S.W.	— — B.fl.V,273 M.fr.XI,20.	N.Z., Am.
<i>M. Australis</i> ; Meissner, gen. pl. comm. 227 (1843)	— — — — N.S.W.	— — — M.fr.XI,169.	
<i>M. gracillima</i> ; Meissner in De Candolle, prodr. XIV, 145 (1856)	— — — — N.S.W.	Q. — B.fl.V,274	
<i>M. rhyticarya</i> ; F. v. M., fragm. V, 92 (1866)	— — — — N.S.W.	Q. — B.fl.V,274	
<i>M. axillaris</i> ; J. Hooker in Loud. Journ. VI, 278 (1847)	— — — T. V. N.S.W.	— — B.fl.V,275 M.fr.IX,77	N.Z.
<i>M. polybotrya</i> ; Meissner in Lehm. pl. Preiss. I, 623 (1845)	W.A. — — — — —	B.fl.V,475 M.fr.IV,130.	

M. polygonoides; F. v. M., fragm. V, 73 (1865)	...	...	...	...	...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.V,275	M.fr.V,73.
M. stenophylla; F. v. M., fragm. I, 138 (1859)	...	...	...	...	...	—	—	—	V.	N.S.W.	—	—	B.fl.V,275	M.fr.I,138.
M. Cunninghamii; F. v. M., fragm. V, 91 (1865)	...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V,275	M.fr.IV,131.

PHYTOLACCEAE.

R. BROWN in Tuck. Exped. Cong. App. V, 454 (1818).—Genera 6, Species 11.

MONOCOCCUS; F. v. M., fragm. I, 47 (1858).

M. echinophorus; F. v. M., fragm. I, 47 (1858)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.V,144	M.fr.I,47	...	P.
--	-----	-----	-----	-----	-----	---	---	---	---	--------	----	---	------------	-----------	-----	----

DIDYMOTHECA; J. Hooker in Lond. Journ. Bot. VI, 278 (1847). (Cyclotheca.)

D. thesioides; J. Hooker in Lond. Journ. V, 279 (1847)	...	...	...	...	...	W.A.	S.A.	T.	—	—	—	—	B.fl.V,145			
D. pleiococca; F. v. M., fragm. I, 202 (1859)...	...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	—	N.A.	B.fl.V,146	M.fr.I,202.		

GYROSTEMON; Desfontaines in Mém. du Mus. VI, 16, t. 6 (1820).

G. brachystigma; F. v. M. in Benth. Fl. Austr. V, 146 (1870)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,146			
G. ramulosus; Desfontaines in Mém. du Mus. V, 17, t. 6 (1820)	...	...	...	...	...	W.A.	S.A.	—	—	—	—	N.A.	B.fl.V,147			

CODONOCARPUS; Cunningham in Hook. Bot. Misc. I, 244 (1830). (Hymenotheca.)

C. pyramidalia; F. v. M., plants of Vict. I, 201 (1862)	...	...	...	...	...	—	S.A.	—	—	—	—	—	B.fl.V,148	M.fr.I,202.		
C. Australis; Cunningham in Hook. bot. misc. I, 244 (1830)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.V,148			
C. cotinifolius; F. v. M., pl. of Vict. I, 200 (1862)	...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V,149	M.fr.II,181; X,53.		

TERSONIA; Moquin in De Candolle, prodr. XIII, pars sec. 40 (1849).

T. brevipes; Moquin in De Candolle, prodr. XIII, pars II, 40 (1849)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,149			
T. subvolubilis; Benth. Fl. Austr. V, 150 (1870)...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,150			

CYPSELOCARPUS; F. v. M., fragm. VIII, 36 (1873).

C. haloragoides; F. v. M., fragm. VIII, 36 (1873)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,198	M.fr.VIII,36.		
---	-----	-----	-----	-----	-----	------	---	---	---	---	---	---	------------	---------------	--	--

NYCTAGINEAE.

A. L. DE JUSSIEU, Gen. plant. 90 (1789).—Genera 2, Species 6.

BOERHAAVIA; Vaillant, serm. de struct. fl. 50 (1718).

B. diffusa; Linné, fl. Zeylan. 4 (1747)	...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V,277	M.fr.IX,75,77	As., Afr., P.
B. elegans; Choisy in De Candolle, prodr. XIII, pars II, 453 (1849)	...	...	...	...	...	—	—	—	—	—	—	N.A.	—	...	As., Afr.
B. plantaginea; Cavanilles, icon. et descr. pl. II, 7, t. 112 (1793)	...	...	...	...	...	W.A.	S.A.	—	—	—	—	Q.	N.A.	B.fl.V,278	E., As., Afr.

PISONIA; Plumier, nov. pl. Amer. gen. 7, t. 11 (1703).

<i>P. aculeata</i> ; Linné, sp. pl. 1026 (1753)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.V,279	M.fr.VI,197	...	As., Afr., Am., P.
<i>P. inermis</i> ; G. Forster, prodr. 75 (1786)	...	..	...	...	...	—	—	—	—	—	—	Q.	N.A.	B.fl.V,280	M.fr.VI,197	P.
<i>P. Brunoniana</i> ; Endlicher, prodr. fl. Norf. 43 (1833)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.V,280	M.fr.I,20; IV,169; VI,198; IX,77.	N.Z.	

CHORIPETALEAE PERIGYNAE.

F. v. M. in Woolls, pl. of the neighb. of Sydney 18 (1880).

THYMELEAE.

A. L. DE JUSSIEU, Gen. plant. 76 (1789).—Genera 4, Species 75.

PIMELEA; Banks & Solander in Gaert. de fruct. I, 186 (1788). (Thecanthes, Gymnococca, Heterolaena, Calyptrostegia, Macrostegia;—Banksia, Forst., 1776.)

P. punicea; R. Brown, prodr. 359 (1810)	...	...	...	...	...	—	—	—	—	—	N.A.	B.fl. VI, 6	M.fr. V, 74; VII, 3.	
P. concreta; F. v. M., fragm. V, 73 (1865)	...	...	...	...	...	—	—	—	—	—	N.A.	B.fl. VI, 6	M.fr. V, 73.	
P. cornucopiae; Vahl, enum. plant. I, 305 (1804)	...	...	...	...	...	—	—	—	—	Q.	—	B.fl. VI, 6	M.fr. VII, 3.	
P. sanguinea; F. v. M., fragm. I, 84 (1858)	...	...	...	...	...	—	—	—	—	Q.	N.A.	B.fl. VI, 7	M.fr. VII, 3; XI, 47.	
P. alpina; F. v. M., second general report, 14 (1854)	...	...	...	...	...	—	—	—	V.	N.S.W.	—	B.fl. VI, 7	M.fr. VII, 4.	
P. longifolia; Banks & Solander in Rees, Cycl. XXVI (1814)	...	...	...	...	...	—	—	—	—	N.S.W.	—	B.fl. VI, 7	M.fr. VIII, 9. ... N.Z.	
P. cinerea; R. Brown, prodr. 361 (1810)	...	...	...	...	...	—	—	T.	—	—	—	B.fl. VI, 8		
P. Milligani; Meissner in De Candolle, prodr. XIV, 509 (1856)	...	...	...	...	...	—	—	T.	—	—	—	B.fl. VI, 8		
P. spectabilis; Lindley, Bot. Regist. t. 33 (1841)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl. VI, 9	M.fr. VII, 4.	
P. Lehmanniana; Meissner in Lehm. pl. Preiss. I, 603 (1845)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl. VI, 9	M.fr. VII, 4.	
P. hispida; R. Brown, prodr. 360 (1810)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl. VI, 10		
P. rosea; R. Brown, prodr. 360 (1810)...	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl. VI, 10		
P. ferruginea; Labillardière, Nov. Holl. pl. spec. I, 10, t. 5 (1804)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl. VI, 10	M.fr. VII, 4.	
P. brachyphylla; Bentham, Fl. Austr. VI, 11 (1873)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl. VI, 11		
P. sylvestris; R. Brown, prodr. 361 (1810)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl. VI, 11	M.fr. VII, 2&5.	
P. brevifolia; R. Brown, prodr. 359 (1810)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl. VI, 12	M.fr. VII, 4.	
P. Maxwelli; F. v. M. in Benth. Fl. Austr. VI, 12 (1873)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl. VI, 12		
P. angustifolia; R. Brown, prodr. 360 (1810)...	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl. VI, 13	M.fr. VII, 3.	
P. nervosa; Meissner in Lehm. pl. Preiss. II, 269 (1847)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl. VI, 13	M.fr. VII, 3.	
P. sulphurea; Meissner in Mohl & Schlecht Bot. Zeitung, 396 (1848)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl. VI, 14	M.fr. VII, 5.	
P. floribunda; Meissner in De Candolle, prodr. XIV, 505 (1856)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl. VI, 14	M.fr. VII, 5.	
P. suaveolens; Meissner in Lehm. pl. Preiss. I, 603 (1845)...	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl. VI, 14	M.fr. VI, 2, 4, 5.	
P. physodes; Hooker, icon. plant. t. 865 (1852)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl. VI, 15	M.fr. VII, 4.	
P. glauca; R. Brown, prodr. 360 (1810)	...	...	...	...	...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl. VI, 15	M.fr. VII, 4; XI, 123.
P. colorans; Cunningham in De Candolle, prodr. XIV, 499 (1856)	...	...	...	...	...	—	—	—	—	N.S.W.	—	—	B.fl. VI, 16	
P. collina; R. Brown, prodr. 359 (1810)	...	...	...	...	...	—	—	—	V.	N.S.W.	Q.	—	B.fl. VI, 16	
P. spathulata; Labillardière, Nov. Holl. pl. spec. I, 9, t. 4 (1804)	...	...	...	...	...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl. VI, 17	
P. linifolia; Smith, Specim. of Bot. of N. Holl. 31, t. 11 (1793)	...	...	...	...	...	—	—	T.	V.	N.S.W.	Q.	—	B.fl. VI, 17	M.fr. VII, 3.
P. ligustrina; Labillardière, Nov. Holl. pl. spec. I, 9, t. 3 (1804)...	...	...	...	...	...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl. VI, 18	M.fr. I, 87; IV, 163; VII, 5.
P. humilis; R. Brown, prodr. 361 (1810)	...	...	...	...	...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl. VI, 19	M.fr. VII, 4; XI, 47.
P. sericea; R. Brown, prodr. 361 (1810)	...	...	...	...	...	—	—	T.	—	—	—	—	B.fl. VI, 19	M.fr. VII, 5.
P. nivca; Labillardière, Nov. Holl. pl. spec. I, 10, t. 6 (1804)	...	...	...	...	...	—	—	T.	—	—	—	—	B.fl. VI, 20	M.fr. VII, 5.
P. imbricata; R. Brown, prodr. 361 (1810)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 20	M.fr. VII, 5.
P. villifera; Meissner in Lehm. pl. Preiss. II, 271 (1847)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 21	
P. drupacea; Labillardière, Nov. Holl. pl. spec. I, 10, t. 7 (1804)	...	...	...	...	...	—	—	T.	V.	—	—	—	B.fl. VI, 21	M.fr. VII, 8.
P. haematostachya; F. v. M., fragm. I, 84 (1858)	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl. VI, 21	M.fr. I, 84; VII, 8; XI, 47.
P. spicata; R. Brown, prodr. 362 (1810)	...	...	...	...	...	—	—	—	—	N.S.W.	—	—	B.fl. VI, 22	M.fr. VII, 7; VIII, 9.
P. spiculigera; F. v. M., in Bentham Fl. Austr. VI, 23 (1873)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 23	M.fr. XI, 46.
P. Forrestiana; F. v. M., fragm. XI, 46 (1879)	...	...	...	...	...	—	—	—	—	—	N.A.	—	—	M.fr. XI, 46.
P. filiformis; J. Hooker in Lond. Journ. VI, 280 (1847)	...	...	...	...	...	—	—	T.	—	—	—	—	B.fl. VI, 23	M.fr. VII, 7.
P. latifolia; R. Brown, prodr. 362 (1810)	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl. VI, 23	M.fr. IV, 49; VII, 7.
P. simplex; F. v. M. in Linnaea XXV, 443 (1852)	...	...	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl. VI, 23	

P. sericostachya; F. v. M., fragm. IV, 162 (1864)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,24	M.fr.XI,47.
P. trichostachya; Lindley in Mitch. Trop. Austr. 355 (1848)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VI,24	M.fr.VII,7.
P. leptostachya; Benth. Fl. Austr. VI, 24 (1873)	—	—	—	—	—	Q.	—	B.fl.VI,24	
P. argentea; R. Brown, prodr. 362 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,25	M.fr.VII,7.
P. clavata; Labillardière, Nov. Holl. pl. spec. I, 11 (1804)... ..	W.A.	—	—	—	—	—	—	B.fl.VI,25	M.fr.VI,160; VII,5.
P. axiflora; F. v. M., First general Report 17 (1853)	—	—	T.	V.	N.S.W.	—	—	B.fl.VI,26	M.fr.VII,7.
P. leptospermoides; F. v. M., fragm. VII, 2 (1870)	—	—	—	—	—	Q.	—	B.fl.VI,27	M.fr.VII,2.
P. microcephala; R. Brown, prodr. 361 (1810)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VI,27	M.fr.XI,27,47.
P. pauciflora; R. Brown, prodr. 360 (1810)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.VI,27	M.fr.VII,8; XI,47.
P. elachantha; F. v. M., First general Report 17 (1853)	—	S.A.	—	V.	—	—	—	B.fl.VI,28	M.fr.VII,6.
P. pygmaea; F. v. M., in Schlecht. Linnaea XXVI, 346 (1853)	—	—	T.	—	—	—	—	B.fl.VI,28	
P. serpyllifolia; R. Brown, prodr. 360 (1810)... ..	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,29	M.fr.VII,6.
P. flava; R. Brown, prodr. 361 (1810)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,29	M.fr.VII,6.
P. petrophila; F. v. M., in Schlecht. Linnaea XXV, 442 (1852)	—	S.A.	—	N.S.W.	—	—	—	B.fl.VI,30	
P. Bowmanni; F. v. M. in Benth. Fl. Austr. VI, 30 (1873)	—	—	—	—	—	Q.	—	B.fl.VI,30	M.fr.VII,6.
P. ammodendris; F. v. M. in Hook. Kew Misc. IX, 24 (1857)	W.A.	S.A.	—	—	—	—	N.A.	B.fl.VI,30	M.fr.VII,5; XI,47.
P. penicillaris; F. v. M. in Melb. Chemist Oct. (1883)	—	—	—	—	N.S.W.	Q.	—	—	
P. curviflora; R. Brown, prodr. 362 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,31	M.fr.VII,6; VIII,9.
P. hirsuta; Meissner in De Candolle, prodr. XIV, 513 (1856)	—	—	—	—	N.S.W.	—	—	B.fl.VI,32	
P. altior; F. v. M., fragm. I, 84 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,32	M.fr.VII,7.
P. octophylla; R. Brown, prodr. 361 (1810)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.VI,32	M.fr.VII,6.
P. petraea; Meissner in Schlecht. Linnaea XXVI, 347 (1853)	—	S.A.	—	—	N.S.W.	—	—	B.fl.VI,33	
P. phylloides; Meissner in Lehm. pl. Preiss. II, 271 (1847)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.VI,33	M.fr.VII,6.
P. Eyrei; F. v. M., fragm. V, 109 (1865)	W.A.	—	—	—	—	—	—	B.fl.VI,34	M.fr.V,109.
P. longiflora; R. Brown, prodr. 361 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,34	M.fr.VII,5.
P. stricta; Meissner in Schlecht. Linnaea XXVI, 348 (1853)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,34	M.fr.VII,4.
P. Preissi; Meissner in Lehm. pl. Preiss. I, 601 (1845)	W.A.	—	—	—	—	—	—	B.fl.VI,35	M.fr.VII,3.
P. Holroydii; F. v. M., fragm. VI, 159, t. 59 (1868)	—	—	—	—	—	—	N.A.	B.fl.VI,35	M.fr.250.

DRAPETES; Lamarck, Journ. d'hist. nat. I, 119, t. 10 (1792). (Kelleria, Daphnobryon.)

D. Tasmanica; J. Hooker in Kew Misc. V, 299, t. 7 (1853) T. V. N.S.W. — — B.fl.VI,36 M.fr.VII,1.

WICKSTROEMIA; Endlicher, prodr. fl. insul. Norfolk. 47 (1833). (Subgenus Stelleriae (1747) Daphne partly.)

W. Indica; C. A. Meyer in Bull. de l'Acad. de Petersb. I, 357 (1843) N.S.W. Q. N.A. B.fl.VI,37 M.fr.VII,1 ... As., P.

PHALERIA; Jack in Hook. Comp. to Bot. Mag. I, 156 (1835). (Drymispermum.)

P. Blumei; Benth. Fl. Austr. VI, 38 (1873) — — — — — Q. — B.fl.VI,38 As.

P. Neumannii; F. v. M., fragm. VIII, 9 (1873) — — — — — Q. — B.fl.VI,38 M.fr.V,26; VII,1.

P. Clerodendron; F. v. M., fragm. VIII, 9 (1873) — — — — — Q. — B.fl.VI,38 M.fr.VII,1.

LEGUMINOSAE.

HALLER, Enum. stirp. Helv. II, 565 (1742).—Genera 95, Species 1065.

JANSONIA; Kippist in Gard. Chron. 19 (1847). (Cryptosema.)

J. formosa; Kippist in Transact. Linn. Soc. XX, 384, t. 16 (1851) W.A. — — — — — B.fl.II,8

BRACHYSEMA; R. Brown in Ait. hort. Kew. sec. edit. III, 10 (1811). (Leptosema, Kaleniczenkia, Burgesia.)

B. praemorsum; Meissner in Lehm. pl. Preiss. I, 25 (1844) W.A. — — — — — B.fl.II,10

B. lanceolatum; Meissner in Lehm. pl. Preiss. I, 24 (1844) W.A. — — — — — B.fl.II,10

B. latifolium; R. Brown in Ait. hort. Kew. sec. ed. III, 10 (1811) W.A. — — — — — B.fl.II,10

B. undulatum; Ker in Bot. Regist. t. 642 (1822) W.A. — — — — — B.fl.II,11 M.fr.IV,11.

B. subcordatum ; Benth. Fl. Austr. II, 11 (1864) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,11	
B. bracteolosum ; F. v. M., fragm. IV, 10 (1864) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,11	M.fr.IV,10.
B. bossiaeioides ; Benth. Fl. Austr. II, 12 (1864) ...	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.II,12	
B. oxylobioides ; Benth. Fl. Austr. II, 12 (1864) ...	...	...	...	...	—	—	—	—	—	—	Q. N.A.	B.fl.II,12	
B. uniflorum ; R. Brown in Benth., Fl. Austr. II, 12 (1864) ...	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.II,12	
B. aphyllum ; Hooker, Bot. Mag. t. 4481 (1849) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,12	M.fr.I,222; II,182.
B. macrocarpum ; Benth. Fl. Austr. II, 13 (1864) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,13	
B. tomentosum ; Benth. Fl. Austr. II, 13 (1864) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,13	
B. Chambersii ; F. v. M. in Benth. Fl. Austr. II, 13 (1864) ...	...	...	...	...	—	S.A.	—	—	—	—	N.A.	B.fl.II,13	
B. daviesioides ; Benth. Fl. Austr. II, 13 (1864) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,13	

OXYLOBIUM ; Andrews, Bot. Reposit. t. 492 (1809). (Chorizematis subgenus, Callistachys, Podolobium.)

O. Callistachys ; Benth. Fl. Austr. II, 16 (1864) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,16	M.fr.IV,18.
O. ellipticum ; R. Brown in Ait. hort. Kew. sec. ed. III, 10 (1811) ...	...	...	...	...	—	—	T.	V.	N.S.W.	Q.	—	B.fl.II,16	
O. alpestre ; F. v. M. in Transact. phil. Soc. Vict. I, 38 (1854) ...	...	...	...	...	—	—	—	V.	N.S.W.	—	—	B.fl.II,17	
O. lineare ; Benth. Fl. Austr. II, 17 (1864) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,17	M.fr.IV,17.
O. carinatum ; Benth. Fl. Austr. II, 18 (1864) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,18	
O. spathulatum ; Benth. Fl. Austr. II, 18 (1864) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,18	
O. obtusifolium ; Sweet, Fl. Austr. t. 5 (1827) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,18	
O. cordifolium ; Andrews, Bot. Reposit. t. 492 (1809) ...	...	...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.II,19	M.fr.IV,17.
O. microphyllum ; Benth. Fl. Austr. II, 19 (1864) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,19	
O. Pulteneae ; De Candolle, prodr. II, 104 (1825) ...	...	...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.II,19	M.fr.IV,19.
O. hamulosum ; Benth. in A. Gray, Bot. Wilk. Exped. t. 379 ...	...	...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.II,20	
O. scandens ; Benth. in Ann. Wien. Mus. II, 70 (1839) ...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,20	
O. procumbens ; F. v. M. in Transact. phil. Soc. Vict. I, 37 (1854) ...	...	...	...	...	—	—	—	—	V.	N.S.W.	—	B.fl.II,20	
O. trienspidatum ; Meissner in Lehm. pl. Preiss. I, 30 (1844) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,21	
O. spectabile ; Endlicher, nov. stirp. dec. 2 (1839) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,21	
O. atropurpureum ; Turczaninow in Bull. Soc. Mosc. XXVI, 250 (1853) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,22	
O. retusum ; R. Brown in Bot. Regist. t. 913 (1825) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,22	
O. virgatum ; Benth. Fl. Austr. II, 22 (1864) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,22	
O. reticulatum ; Meissner in Lehm. in Preiss. I, 29 (1844) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,23	
O. capitatum ; Benth. in Hueg. enum. 28 (1837) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,23	
O. cuneatum ; Benth. in Bot. Regist. XXV, App. XII (1839) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,23	
O. acutum ; Benth. Fl. Austr. II, 24 (1864) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,24	
O. parviflorum ; Benth. in Bot. Regist. XXV, App. XII (1839) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,24	
O. heterophyllum ; Benth. Fl. Austr. II, 25 (1864) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,25	
O. aciculiferum ; F. v. M., fragm. I, 75 implied (1859) ...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl.II,25	M.fr.I,75.
O. trilobatum ; F. v. M., annual report 23 (1858) ...	...	...	...	...	—	—	—	—	V.	N.S.W.	Q.	B.fl.II,25	M.fr.IV,19.

CHORIZEMA ; Labillardière, Voy. I, 405 (1799). (Orthotropis.)

C. Dicksonii ; Graham in Maund's Botanist, t. 106 (1841) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,27	
C. nervosum ; Th. Moore in Gard. Companion (1852) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,27	
C. varium ; Benth. in Bot. Regist. XXV, t. 49 (1839) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,28	
C. cordatum ; Lindley, Bot. Regist. t. 10 (1838) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,28	
C. ilicifolium ; Labillardière, Voy. I, 405, t. 21 (1799) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,28	
C. rhombeum ; R. Brown in Ait. hort. Kew. sec. ed. III, 9 (1811) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,28	
C. diversifolium ; A. de Candolle, Pl. rar. jard. Genève, 44, t. 8 (1836) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,29	
C. angustifolium ; Benth. in Hueg. enum. 28 (1837) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,29	
C. reticulatum ; Meissner in Lehm. pl. Preiss. I, 34 (1844) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,30	
C. trigonum ; Turczaninow in Bull. de la Soc. Mosc. XXVI, 254 (1853) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,30	
C. humile ; Turczaninow in Bull. de la Soc. Mosc. XXVI, 254 (1853) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,30	

C. parviflorum ; Bentham in Ann. des Wien. Mus. II, 71 (1838) ...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,30
C. cytiseoides ; Turczaninow in Bull. Soc. Mosc. XXVI, 256 (1853) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,31
C. Henchmanni ; R. Brown in Bot. Regist. t. 986 (1826) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,31
C. ericifolium ; Meissner in Lehm. pl. Preiss. II, 209 (1847) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,32

GASTROLOBIIUM ; R. Brown in Ait. hort. Kew. sec. edit. III, 16 (1811). (Chorizematis subgenus.)

G. pyramidale ; Th. Moore in Gard. Comp. I, 81 (1852) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,98
G. Lehmanni ; Meissner in Lehm. pl. Preiss. I, 70 (1844) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,98 M.fr.,X,35.
G. pulchellum ; Turczaninow in Bull. de la Soc. Mosc. XXVI, 274 (1853) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,98
G. stipulare ; Meissner in Lehm. pl. Preiss. II, 218 (1847) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,99
G. Brownii ; Meissner in Lehm. pl. Preiss. I, 71 (1844) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,99
G. reticulatum ; Bentham, Fl. Austr. II, 99 (1864) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,99 M.fr.X,35.
G. truncatum ; Bentham, Fl. Austr. II, 99 (1864) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,99
G. spathulatum ; Bentham in Bot. Regist. XX, App. XIV (1839) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,100
G. plicatum ; Turczaninow in Bull. Soc. Mosc. XXVI, 274 (1853) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,100
G. tricuspidatum ; Meissner in Lehm. pl. Preiss. I, 66 (1844) ..	...	...	W.A.	—	—	—	—	—	—	B.fl.II,100
G. obovatum ; Bentham in Bot. Regist. XXV, App. XIV (1839) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,101
G. epacridioides ; Meissner in Lehm. Preiss. I, 72 (1844) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,102
G. trilobum ; Bentham in Bot. Regist. XXV, App. XIII (1839) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,101 M.fr.X,35.
G. ilicifolium ; Meissner in Lehm. pl. Preiss. I, 67 (1844) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,101
G. villosum ; Bentham in Bot. Regist. XXV, App. XIII (1839) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,102
G. polystachyum ; Meissner in Lehm. pl. Preiss. II, 217 (1847) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,102
G. ovalifolium ; Henfrey in Gard. Comp. I, 41 (1852) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,102
G. grandiflorum ; F. v. M., fragm. III, 17 (1862) ...	...	...	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.II,103 M.fr.IV,174 ; X,35.
G. pycnostachyum ; Bentham, Fl. Austr. II, 103 (1864) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,103
G. spinosum ; Bentham in Bot. Regist. XXV, App. XIII (1839) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,103 M.fr.X,35.
G. rotundifolium ; Meissner in Lehm. pl. Preiss. II, 216 (1847) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,104
G. microcarpum ; Meissner in Lehm. pl. Preiss. I, 70 (1844) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,104
G. oxylobioides ; Bentham in Bot. Regist. XXV, App. XIII (1839) ..	...	...	W.A.	—	—	—	—	—	—	B.fl.II,104
G. calycinum ; Bentham in Bot. Regist. XXV, App. XIII (1839)...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,104 M.fr.X,35.
G. Callistachys ; Meissner in Lehm. pl. Preiss. II, 216 (1846) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,105
G. stenophyllum ; Turczaninow in Bull. Soc. Mosc. XXVI, 275 (1853) ..	...	...	W.A.	—	—	—	—	—	—	B.fl.II,105
G. crassifolium ; Bentham, Fl. Austr. II, 105 (1864) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,105
G. parvifolium ; Bentham, Fl. Austr. II, 106 (1864)...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,106
G. hamulosum ; Meissner in Lehm. pl. Preiss. II, 218 (1847) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,106
G. velutinum ; Lindley & Paxton, Flower-Gard. III, 76 (1853) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,106 M.fr.X,35.
G. bidens ; Meissner in Regensb. Bot. Zeit. 29 (1855) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,106
G. bilobum ; R. Brown in Ait. hort. Kew. sec. ed. III, 16 (1811) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,107 M.fr.X,35.
G. seorsifolium ; F. v. M., fragm. X, 45 (1876) ...	...	...	W.A.	—	—	—	—	—	—	M.fr.X,35.

ISOTROPIS ; Bentham in Hueg. enum. 28 (1837).

I. striata ; Bentham in Hueg. enum. 28 (1837)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,39	M.fr.III,16.
I. Drummondii ; Meissner in Lehm. pl. Preiss, I, 31 (1844)	...	..	...	...	W.A.	—	—	—	—	—	B.fl.II,39	
I. juncea ; Turczaninow in Bull. Soc. Mosc. XXVI, 251 (1853)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,39	M.fr.X,51.
I. canescens ; F. v. M., fragm. X, 51 (1876)	...	...	...	...	W.A.	—	—	—	—	—	—	M.fr.X,51.
I. atropurpurea ; F. v. M., fragm. III, 16 (1862)	...	...	...	...	W.A.	S.A.	—	—	—	N.A.	B.fl.II,40	M.fr.X,51.
I. filicaulis ; Bentham in Ann. des Wien. Mus. II, 71 (1838)	...	...	...	...	—	—	—	—	—	Q.	B.fl.II,40	M.fr.IV,20.
I. Forrestii ; F. v. M. in Wing's S. Sc. Rec. II, 252 (1882)	...	...	...	...	W.A.	—	—	—	—	—	—	
I. parviflora ; Bentham in Ann. des Wien. Mus. II, 71 (1838)	...	...	...	...	—	—	—	—	—	N.A.	B.fl.II,40	
I. Wheeleri ; F. v. M. in Benth. Fl. Austr. II, 40 (1864)	...	...	...	...	—	S.A.	—	—	N.S.W.	—	B.fl.II,40	
I. Winneckiana ; F. v. M. in Austr. Chemist (1884)	...	...	...	...	—	S.A.	—	—	—	—	—	

MIRBELIA; Smith in Koenig & Sims, Ann. of Bot. I, 511 (1805). (Dichosema, Oxycladium.)

<i>M. dilatata</i> ; R. Brown in Ait. hort. Kew. sec. ed. III, 21 (1811) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,33	
<i>M. racemosa</i> ; Turczaninow in Bull. Soc. Mosc. XXVI, 282 (1853) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,33	M.fr.IV,18.
<i>M. grandiflora</i> ; Aiton in Bot. Mag. t. 2771 (1827) ...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.II,34	
<i>M. subcordata</i> ; Turczaninow in Bull. Soc. Mosc. XXVI, 282 (1853) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,34	
<i>M. ovata</i> ; Meissner in Lehm. pl. Preiss. I, 77 (1844) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,34	M.fr.II,154.
<i>M. oxylobioides</i> ; F. v. M., fragm. II, 154 (1861) ...	...	...	—	—	—	—	V. N.S.W.	—	—	B.fl.II,34	M.fr.III,167; IV,12.
<i>M. reticulata</i> ; Smith in Koen. & Sims, Ann. of Bot. I, 511 (1805) ...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.II,35	M.fr.IV,12.
<i>M. aotoides</i> ; F. v. M. in Transact. phil. Inst. Vict. III, 53 (1859) ...	...	...	—	—	—	—	N.S.W. Q.	—	—	B.fl.II,35	M.fr.IV,12.
<i>M. pungens</i> ; Cunningham in G. Don, gen. syst. II, 126 (1832) ...	...	...	—	—	—	—	N.S.W. Q.	—	—	B.fl.II,35	
<i>M. speciosa</i> ; Sieber in De Candolle, prodr. II, 115 (1824) ...	...	...	—	—	—	—	N.S.W. Q.	—	—	B.fl.II,36	M.fr.IV,12.
<i>M. floribunda</i> ; Benthām in Bot. Regist. XXV, App. XII (1839)...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,36	
<i>M. spinosa</i> ; Benthām, Fl. Austr. II, 36 (1864) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,36	
<i>M. multicaulis</i> ; Benthām, Fl. Austr. II, 37 (1864) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,37	
<i>M. microphylla</i> ; Benthām, Fl. Austr. II, 37 (1864)...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,37	
<i>M. aphylla</i> ; F. v. M., fragm. IV, 11 (1863) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,37	M.fr.IV,11.
<i>M. oxyclada</i> ; F. v. M., fragm. IV, 12 (1864)...	...	...	—	—	—	—	Q. N.A.	—	—	B.fl.II,38	M.fr.I,168.

GOMPHOLOBIUM; Smith in Transact. Linn. Soc. IV, 220 (1798).

<i>G. amplexicaule</i> ; Meissner in Lehm. pl. Preiss. I, 36 (1844) ...	...	...	W.A.	—	—	—	—	—	B.fl.II,42
<i>G. latifolium</i> ; Smith in Koenig & Sims, Ann. of Bot. I, 505 (1805) ...	...	...	—	—	—	V.	N.S.W.	Q.	B.fl.II,42
<i>G. pedunculare</i> ; Loddiges, Bot. Cabinet, t. 1639 (1830) ...	...	...	—	—	T.	V.	N.S.W.	—	B.fl.II,43
<i>G. polymorphum</i> ; R. Brown in Ait. hort. Kew sec. ed. III, 11 (1811) ...	...	...	W.A.	—	—	—	—	—	B.fl.II,43
<i>G. obcordatum</i> ; Turczaninow in Bull. Soc. Mosc. XXVI, 258 (1853) ...	...	...	W.A.	—	—	—	—	—	B.fl.II,44
<i>G. marginatum</i> ; R. Brown in Ait. hort. Kew. sec. ed. III, 11 (1811) ...	...	...	W.A.	—	—	—	—	—	B.fl.II,44
<i>G. grandiflorum</i> ; Smith, Exot. Bot. t. 5 (1804) ...	...	...	—	—	—	—	N.S.W.	—	B.fl.II,44
<i>G. virgatum</i> ; Sieber in De Candolle, prodr. II, 105 (1824)...	...	...	—	—	—	—	N.S.W.	Q.	B.fl.II,45
<i>G. minus</i> ; Smith in Transact. Linn. Soc. IX, 251 (1808) ...	...	...	—	S.A.	—	V.	N.S.W.	—	B.fl.II,45
<i>G. uncinatum</i> ; Cunningham in Annal. des Wien. Mus. II, 72 (1838) ...	...	...	—	—	—	—	N.S.W.	—	B.fl.II,46
<i>G. Eatoniae</i> ; F. v. M. in Vict. Nat. VI (1889) ...	...	...	W.A.	—	—	—	—	—	—
<i>G. Baxteri</i> ; Benth. Fl. Austr. II, 46 (1864) ...	...	...	W.A.	—	—	—	—	—	B.fl.II,46
<i>G. aristatum</i> ; Benth. in Ann. des Wien. Mus. II, 72 (1838) ...	...	...	W.A.	—	—	—	—	—	B.fl.II,46
<i>G. burtonioides</i> ; Meissner in Lehm. pl. Preiss. I, 37 (1844) ...	...	...	W.A.	—	—	—	—	—	B.fl.II,46
<i>G. capitatum</i> ; Cunningham in Bot. Regist. t. 1563 (1839) ...	...	...	W.A.	—	—	—	—	—	B.fl.II,47
<i>G. tomentosum</i> ; Labillardière, Nov. Holl. pl. sp. 1, 106, t. 134 (1804) ...	...	...	W.A.	—	—	—	—	—	B.fl.II,47
<i>G. Preissii</i> ; Meissner in Lehm. pl. Preiss. I, 40 (1844) ...	...	...	W.A.	—	—	—	—	—	B.fl.II,47
<i>G. viscidulum</i> ; Meissner in Lehm. pl. Preiss. I, 39 (1844) ...	...	...	W.A.	—	—	—	—	—	B.fl.II,48
<i>G. glabratum</i> ; De Candolle, prodr. II, 106 (1824) ...	...	...	—	—	—	—	N.S.W.	—	B.fl.II,48
<i>G. nitidum</i> ; Solander in Benth. Fl. Austr. II, 48 (1864) ...	...	...	—	—	—	—	—	Q.	B.fl.II,48
<i>G. pinnatum</i> ; Smith in Transact. Linn. Soc. IX, 251 (1808) ...	...	...	—	—	—	—	N.S.W.	Q.	B.fl.II,48
<i>G. Shuttleworthii</i> ; Meissner in Lehm. pl. Preiss. I, 39 (1844) ...	...	...	W.A.	—	—	—	—	—	B.fl.II,49
<i>G. venustum</i> ; R. Brown in Ait. hort. Kew. sec. ed. III, 12 (1811) ...	...	...	W.A.	—	—	—	—	—	B.fl.II,49
<i>G. Knightiauum</i> ; Lindley, Bot. Regist. t. 1468 (1837) ...	...	...	W.A.	—	—	—	—	—	B.fl.II,49

BURTONIA; R. Brown in Ait. hort. Kew. ed. sec. III, 13 (1811). (Gompholobii subgenus.)

<i>B. subulata</i> ; Bentham, Fl. Austr. II, 50 (1864)	...	...	...	...	...	—	—	—	—	—	N.A.	B.fl.II,50	M.fr.III,30; X,35.
<i>B. foliolosa</i> ; Bentham, Fl. Austr. II, 50 (1864)	..	..,	...	...	..	—	—	—	—	—	Q.	— B.fl.II,50	
<i>B. polyzyga</i> ; Bentham, Fl. Austr. II, 51 (1864)	...	...	...	...	...	—	S.A.	—	—	—	—	N.A. B.fl.II,51	M.fr.III,29; X,34.
<i>B. villosa</i> ; Meissner in Lehmann, pl. Preiss. I, 41 (1844)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,51	M.fr.X,35.
<i>B. Hendersonii</i> ; Bentham, Fl. Austr. II, 51 (1864)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,51	
<i>B. gompholobioides</i> ; F. v. M., fragm. X, 34 (1876)	...	...	...	...	...	W.A.	—	—	—	—	—	—	M.fr.X,34.

B. scabra; R. Brown in Ait. hort. Kew. sec. ed. III, 12 (1811)	...	...	...	W.A.	—	—	—	—	—	B.fl.II,51	M.fr.X,35.
B. conferta; De Candolle, prodr. II, 106 (1825)	...	...	...	..	W.A.	—	—	—	—	B.fl.II,52	M.fr.X,35.

JACKSONIA; R. Brown in Ait. hort. Kew. sec. ed. III, 12 (1811). (Piptomeris.)

J. dilatata ; Bentham in Ann. des Wien. Mus. II, 74 (1838)	W.A.	—	—	—	—	—	—	N.A.	B.fl.II,54	M.fr.X,37.
J. densiflora ; Bentham in Bot. Regist. XXV, App. XIII (1839)	W.A.	—	—	—	—	—	—	—	B.fl.II,54	M.fr.X,38.
J. carduacea ; Meissner in Bot. Zeit. 25 (1855)	W.A.	—	—	—	—	—	—	—	B.fl.II,55	
J. floribunda ; Endlicher in Ann. des Wien. Mus. II, 197 (1838) ..	W.A.	—	—	—	—	—	—	—	B.fl.II,55	
J. odontoclada ; F. v. M. in Benth. Fl. Austr. II, 55 (1864)	—	—	—	—	—	—	—	N.A.	B.fl.II,55	M.fr.X,37.
J. ramosissima ; Bentham in Mitch. Trop. Austr. 258 (1848)	—	—	—	—	—	—	—	Q.	B.fl.II,56	M.fr.X,37.
J. foliosa ; Turczaninow in Bull. de la Soc. Imp. de Mosc. XXVI, 260 (1853) ...	W.A.	—	—	—	—	—	—	—	B.fl.II,56	
J. spinosa ; R. Brown in Ait. hort. Kew. sec. ed. III, 13 (1811)	W.A.	—	—	—	—	—	—	—	B.fl.II,56	M.fr.X,38&39.
J. stricta ; Meissner in Regensb. Bot. Zeit. 27 (1855)	W.A.	—	—	—	—	—	—	—	B.fl.II,56	
J. hakeoides ; Meissner in Lehm. pl. Preiss. I, 45 (1844)	W.A.	—	—	—	—	—	—	—	B.fl.II,57	M.fr.X,39.
J. furcellata ; De Candolle, prodr. II, 107 (1824)	W.A.	—	—	—	—	—	—	—	B.fl.II,57	M.fr.X,39.
J. horrida ; De Candolle, prodr. II, 107 (1824)	W.A.	—	—	—	—	—	—	—	B.fl.II,57	
J. sericea ; Bentham in Hueg. enum. 31 (1837)	W.A.	—	—	—	—	—	—	—	B.fl.II,58	
J. Sternbergiana ; Huegel, Bot. Archiv. t. 3 (1837)	W.A.	—	—	—	—	—	—	—	B.fl.II,58	M.fr.X,39.
J. vernicosa ; F. v. M. in Benth. Fl. Austr. II, 58 (1864)	—	—	—	—	—	—	—	N.A.	B.fl.II,58	M.fr.X,37.
J. rhadinoclona ; F. v. M., fragm. X, 37 (1876)	—	—	—	—	—	N.S.W.	Q.	—	—	M.fr.X,37.
J. Forrestii ; F. v. M. in proceed. L. S. of N.S.W. sec. ser. II, 194 (1887)	W.A.	—	—	—	—	—	—	—	—	
J. thesioides ; Cunningham in den Ann. des Wien. Mus. II, 74 (1838)	—	—	—	—	—	—	—	Q.	N.A. B.fl.II,59	M.fr.IV,161 ; X,37.
J. compressa ; Turczaninow in Bull. de la Soc. de Mosc. XXVI, 260 (1853)	W.A.	—	—	—	—	—	—	—	B.fl.II,59	M.fr.X,38.
J. Clarkei ; F. v. M. in proceed. L. S. of N.S.W. sec. ser. II, 193 (1887)	—	—	—	—	—	V. N.S.W.	—	—	—	
J. scoparia ; R. Bröwn in Ait. hort. Kew. sec. ed. III, 13 (1811)	—	—	—	—	—	N.S.W.	Q.	—	B.fl.II,59	M.fr.X,37.
J. cupulifera ; Meissner in der Bot. Zeit. 27 (1855)	W.A.	—	—	—	—	—	—	—	—	M.fr.X,36.
J. restioides ; Meissner in Lehm. pl. Preiss. I, 46 (1844)	W.A.	—	—	—	—	—	—	—	B.fl.II,60	
J. velutina ; Bentham, Fl. Austr. II, 60 (1864)	W.A.	—	—	—	—	—	—	—	B.fl.II,60	
J. Lehmanni ; Meissner in Lehm. pl. Preiss. I, 46 (1844)	W.A.	—	—	—	—	—	—	—	B.fl.II,60	
J. racemosa ; Meissner in Lehm. pl. Preiss. II, 212 (1847)	W.A.	—	—	—	—	—	—	—	B.fl.II,60	
J. umbellata ; Turczaninow in Bull. Soc. Mosc. XXVI, 261 (1853)	W.A.	—	—	—	—	—	—	—	B.fl.II,61	M.fr.X,38.
J. nematoclada ; F. v. M., fragm. X, 50 (1876)	W.A.	—	—	—	—	—	—	—	—	M.fr.X,50.
J. rhadinoclada ; F. v. M., fragm. X, 38 (1876)	W.A.	—	—	—	—	—	—	—	—	M.fr.X,38.
J. capitata ; Meissner in Lehm. pl. Preiss. I, 45 (1844)	W.A.	—	—	—	—	—	—	—	B.fl.II,61	M.fr.X,38.
J. alata ; Bentham in Hueg. enum. 30 (1837)	W.A.	—	—	—	—	—	—	—	B.fl.II,61	M.fr.X,38.
J. angulata ; Bentham, Fl. Austr. II, 62 (1864)	W.A.	—	—	—	—	—	—	—	B.fl.II,62	
J. Stackhousii ; F. v. M. in proc. Linn. Soc. N.S.W., VI, 791 (1881)	—	—	—	—	—	N.S.W.	Q.	—	—	M.fr.XI,138.
J. pteroclada ; F. v. M., fragm. X, 37 (1876)	W.A.	—	—	—	—	—	—	—	—	M.fr.X,37.
J. macrocalyx ; Meissner in der Bot. Zeit. 26 (1855)	W.A.	—	—	—	—	—	—	—	B.fl.II,62	
J. purpurascens ; F. v. M., fragm. IV, 161 (1864)	—	—	—	—	—	—	—	Q.	—	M.fr.IV,161.
J. piptomeris ; Bentham, Fl. Austr. II, 62 (1864)	W.A.	—	—	—	—	—	—	—	B.fl.II,62	

SPHAEROLOBIUM; Smith in Koenig & Sims, Ann. of Bot. I, 509 (1805). (Roea.)

<i>S. linophyllum</i> ; F. v. M., fragm. I, 167 (1859)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,64	M.fr.I,167.
<i>S. foliosum</i> ; F. v. M., fragm. I, 166 (1859)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,64	M.fr.I,166.
<i>S. gracile</i> ; Bentham, Fl. Austr. II, 64 (1864)...	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,64	
<i>S. racemosum</i> ; Bentham, Fl. Austr. II, 64 (1864)...	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,65	
<i>S. alatum</i> ; Bentham in Hueg. enum. 32 (1837)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,65	
<i>S. vimineum</i> ; Smith in Koen. & Sims, Ann. of Bot. I, 509 (1805)	...	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	B.fl.II,65	
<i>S. grandiflorum</i> ; R. Brown in Benth. Fl. Austr. II, 66 (1864)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,66	
<i>S. fornicatum</i> ; Bentham in Hueg. enum. 32 (1837)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,66	

S. medium; R. Brown in Ait. hort. Kew. sec. ed. III, 14 (1811) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,66
S. scabriusculum; Meissner in Lehm. pl. Preiss. II, 214 (1849) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,66
S. macranthum; Meissner in Lehm. pl. Preiss. II, 213 (1847) ..	...	...	W.A.	—	—	—	—	—	—	B.fl.II,67
S. davisoides; Turczaninow in Bull. Soc. Mosc. XXVI, 266 (1853) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,67

VIMINARIA; Smith, Exotic. Bot. 51, t. 27 (1804).

V. denudata; Smith, Exot. Bot. 51, t. 27 (1804) ...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,68
---	-----	-----	------	------	----	----	--------	----	---	------------

DAVIESIA; Smith in Transact. Linn. Soc. IV, 222 (1798).

D. cordata; Smith in Transact. Linn. Soc. IX, 259 (1808) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,72
D. ovata; Benth. Fl. Austr. II, 72 (1864) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,72
D. crenulata; Turczaninow in Bull. Soc. de Mosc. XXVI, 265 (1853) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,72 M.fr.IV,16.
D. oppositifolia; Endlicher in den Ann. des Wien. Mus. II, 199 (1838) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,73
D. alternifolia; Endlicher in Ann. des Wien. Mus. II, 199 (1838)...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,73 M.fr.II,105.
D. elongata; Benth. Fl. Austr. II, 74 (1864) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,74
D. pedunculata; Benth. in Bot. Regist. XXV, App. XIV (1839) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,74
D. arthropoda; F. v. M., fragm. VIII, 225 (1874) ...	...	...	W.A.	S.A.	—	—	—	—	—	M.fr. VIII, 225.
D. mollis; Turczaninow in Bull. Soc. Mosc. XXVI, 263 (1853) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,74
D. concinna; R. Brown in Benth. Fl. Austr. II, 75 (1864)...	...	...	—	—	—	N.S.W.	Q.	—	—	B.fl.II,75
D. umbellulata; Smith in Koen. & Sims, Ann. of Bot. I, 507 (1805) ...	...	...	—	—	—	N.S.W.	Q.	—	—	B.fl.II,75
D. Wyattii; Bailey in pap. Garden. Assoc. of S. Austr. (1880) ...	...	...	—	—	—	V.	N.S.W.	Q.	—	M.fr.X,138.
D. latifolia; R. Brown in Ait. hort. Kew. sec. ed. III, 20 (1811) ...	...	...	—	—	—	T.	V.	N.S.W.	—	B.fl.II,76
D. corymbosa; Smith in Koen. & Sims, Ann. of Bot. I, 507 (1805) ...	...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.II,76
D. horrida; Meissner in Lehm. pl. Preiss. I, 54 (1844) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,77
D. reclinata; Cunningham in Benth. Fl. Austr. II, 77 (1864) ...	...	...	—	—	—	—	—	N.A.	—	B.fl.II,77
D. obtusifolia; F. v. M., fragm. II, 104 (1861) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,77 M.fr.II,104.
D. obovata; Turczaninow in Bull. Soc. Mosc. XXVI, 261 (1853) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,78
D. longifolia; Benth. in Bot. Regist. XXV, App. XIV (1839) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,78
D. chordophylla; Meissner in Lehm. plant. Preiss. I, 48 (1844) ..	...	...	W.A.	—	—	—	—	—	—	B.fl.II,78
D. nematophylla; F. v. M. in Benth. Fl. Austr. II, 78 (1864) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,78
D. daphnoides; Meissner in Lehm. plant. Preiss. I, 54 (1844) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,79
D. nudiflora; Meissner in Lehm. plant. Preiss. I, 53 (1844) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,79
D. rhombifolia; Meissner in Lehm. plant. Preiss. I, 56 (1844) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,79
D. Drummondii; Meissner in Lehm. plant. Preiss. I, 53 (1844) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,80 M.fr.II,106.
D. filipes; Benth. in Mitch. Trop. Austr. 363 (1848) ...	...	...	—	—	—	N.S.W.	Q.	—	—	B.fl.II,80
D. cardiophylla; F. v. M., fragm. II, 105 (1861) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,79 M.fr.II,105.
D. squarrosa; Smith in Koen. & Sims, Ann. of Bot. I, 507 (1805) ...	...	...	—	—	—	N.S.W.	Q.	—	—	B.fl.II,80
D. ulicina; Smith in Koen. & Sims, Ann. of Bot. I, 506 (1805) ...	...	...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,81 M.fr.II,106.
D. acicularis; Smith in Koen. & Sims, Ann. of Bot. I, 506 (1805) ...	...	...	—	—	—	N.S.W.	Q.	—	—	B.fl.II,81
D. pachyphylla; F. v. M., fragm. IV, 15 (1864) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,82 M.fr.IV,15.
D. teretifolia; R. Brown in Benth. Fl. Austr. II, 82 (1864) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,82
D. genistifolia; Cunningham in Annal. Wien. Mus. II, 75 (1838) ...	...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.II,82
D. hakeoides; Meissner in Lehm. plant. Preiss. I, 47 (1844) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,82
D. colletioides; Meissner in Lehm. plant. Preiss. I, 48 (1844) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,83
D. reversifolia; F. v. M., fragm. I, 145 (1859) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,83 M.fr.I,145.
D. incrassata; Smith in Transact. Linn. Soc. IX, 255 (1808) ...	...	...	W.A.	S.A.	—	—	—	—	—	B.fl.II,83 M.fr.I,145.
D. brevifolia; Lindley in Mitch. Three Exped. II, 201 (1838) ...	...	...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.II,84
D. acanthoclona; F. v. M., fragm. X, 32 (1876) ...	...	...	W.A.	—	—	—	—	—	—	M.fr.X,32.
D. Preissii; Meissner in Lehm. plant. Preiss. I, 50 (1844) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,84
D. spinosissima; Meissner in Lehm. plant. Preiss. I, 51 (1844) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,85
D. pachylina; Turczaninow in Bull. Soc. Mosc. XXVI, 263 (1853) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,85

D. quadrilatera; Bentham in Bot. Regist. XXV, App. XIV (1839) ...	...	W.A.	—	—	—	—	—	B.fl.II,85	
D. striata; Turczaninow in Bull. Soc. Mosc. XXVI, 264 (1853) ...	...	W.A.	—	—	—	—	—	B.fl.II,85	M.fr.II,105.
D. polyphylla; Bentham, in Bot. Regist. XXV, App. XIV (1839) ...	...	W.A.	—	—	—	—	—	B.fl.II,86	
D. microphylla; Bentham, Fl. Austr. II, 86 (1864) ...	...	W.A.	—	—	—	—	—	B.fl.II,86	
D. flexuosa; Bentham in Hueg. enum. 32 (1837) ...	...	W.A.	—	—	—	—	—	B.fl.II,86	M.fr.I,145.
D. pectinata; Liodley in Mitch. Three Exped. II, 151 (1838) ...	...	W.A. S.A.	—	V.	N.S.W.	—	—	B.fl.II,87	
D. trigonophylla; Meissner in Lehm. plant. Preiss. II, 213 (1847) ...	...	W.A.	—	—	—	—	—	B.fl.II,87	
D. epiphylla; Meissner in Regensb. Bot. Zeit. 27 (1855) ...	...	W.A.	—	—	—	—	—	B.fl.II,87	
D. euphorbioides; Bentham, Fl. Austr. II, 88 (1864) ...	...	W.A.	—	—	—	—	—	B.fl.II,88	
D. divaricata; Bentham in Hueg. enum. 31 (1837) ...	...	W.A.	—	—	—	—	—	B.fl.II,88	
D. paniculata; Bentham in Hueg. enum. 31 (1837) ...	...	W.A.	—	—	—	—	—	B.fl.II,88	
D. aphylla; F. v. M. in Benth. Fl. Austr. 88 (1864) ...	...	W.A.	—	—	—	—	—	B.fl.II,88	
D. juncea; Smith in Transact. Linn. Soc. IX, 260 (1808) ...	...	W.A.	—	—	—	—	—	B.fl.II,88	
D. alata; Smith in Transact. Linn. Soc. IX, 259 (1808) ...	...	—	—	—	N.S.W.	—	—	B.fl.II,89	
D. anceps; Turczaninow in Bull. Soc. Mosc. XXVI, 266 (1853) ...	...	W.A.	—	—	—	—	—	B.fl.II,89	

AOTUS; Smith in Koenig & Sims Ann. of Bot. I, 504 (1805).

A. gracillima; Meissner in Lehm. pl. Preiss. I, 59 (1844) ...	...	W.A.	—	—	—	—	—	B.fl.II,90	
A. villosa; Smith in Koen. & Sims, Ann. of Bot. I, 504 (1805) ..	...	—	T.	V.	N.S.W.	Q.	—	B.fl.II,90	
A. mollis; Bentham in Mitch. Trop. Austr. 236 (1848) ...	...	—	—	—	N.S.W.	Q.	—	B.fl.II,91	
A. Preissii; Meissner in Lehm. plant. Preiss. II, 214 (1847) ...	...	W.A.	—	—	—	—	—	B.fl.II,91	
A. phyllicoides; F. v. M. in Benth. Fl. Austr. II, 72 (1864) ...	...	W.A.	—	—	—	—	—	B.fl.II,92	
A. lanigera; Cunningham in Annal. des Wien. Mus. II, 78 (1838) ...	...	—	—	—	N.S.W.	Q.	—	B.fl.II,92	
A. genistoides; Turczaninow in Bull. Soc. Mosc. XXVI, 268 (1853) ...	...	W.A.	—	—	—	—	—	B.fl.II,92	
A. carinata; Meissner in Lehm. plant. Preiss. II, 215 (1847) ...	...	W.A.	—	—	—	—	—	B.fl.II,93	
A. passerinoides; Meissner in Lehm. plant. Preiss. I, 61 (1844) ...	...	W.A.	—	—	—	—	—	B.fl.II,93	
A. Tietkensii; F. v. M., fragm. X, 33 (1876) ...	...	W.A.	—	—	—	—	—	—	M.fr.X,33.
A. cordifolia; Bentham in Hueg. enum. 33 (1837) ...	...	W.A.	—	—	—	—	—	B.fl.II,93	

PHYLLOTA; De Candolle, prodr. II, 113 (1825). (Urodon.)

P. barbata; Bentham in Hueg. enum. 33 (1837) ...	...	W.A.	—	—	—	—	—	B.fl.II,94	
P. gracilis; Turczaninow in Bull. Soc. Mosc. XXVI, 267 (1853) ...	...	W.A.	—	—	—	—	—	B.fl.II,94	
P. Sturtii; Bentham, Fl. Austr. II, 95 (1864) ...	...	S.A.	—	—	—	—	—	B.fl.II,95	
P. phyllicoides; Bentham in Ann. des Wien. Mus. II, 77 (1838) ..	...	—	—	—	N.S.W.	Q.	—	B.fl.II,95	M.fr.X,33.
P. humifusa; Bentham, Fl. Austr. II, 95 (1864) ...	...	—	—	—	N.S.W.	—	—	B.fl.II,95	
P. diffusa; F. v. M., fragm. I, 8 (1858) ...	...	—	T.	—	—	—	—	B.fl.II,119	M.fr.I,8.
P. pleurandroides; F. v. M. in Transact. phil. Inst. Vict. I, 38 (1854) ...	...	S.A.	—	V.	—	—	—	B.fl.II,96	M.fr.X,33.
P. Luehmanni; F. v. M., fragm. X, 33 (1876) ...	...	W.A.	—	—	—	—	—	—	M.fr.X,33.
P. Urodon; F. v. M. in papers of the Roy. Soc. of Tasm. 120 (1877) ...	...	W.A.	—	—	—	—	—	B.fl.II,124	

PULTENAEA; Smith, Specim. of the Bot. of New Holl. 35, t. 12 (1793). (Euchilus, Spadostyles, Bartlingia, Pulteneya.)

P. daphnoides; Wendland, Bot. Beob. 49 (1798) ...	...	S.A. T.	V.	N.S.W.	—	—	—	B.fl.II,112	
P. stricta; Sims, Bot. Mag. 1588 (1813) ...	...	S.A. T.	V.	N.S.W.	—	—	—	B.fl.II,112	
P. retusa; Smith in Koen. & Sims, Ann. of Bot. I, 502 (1805) ...	...	—	—	V.	N.S.W.	Q.	—	B.fl.II,113	
P. Benthamii; F. v. M. in Transact. phil. Inst. Vict. I, 38 (1854) ...	...	—	—	V.	—	—	—	B.fl.II,113	
P. pycnocephala; F. v. M. in Benth. Fl. Austr. II, 114 (1864) ...	...	—	—	—	N.S.W.	—	—	B.fl.II,114	
P. myrtoides; Cunningham in Annal. des Wien. Mus. II, 81 (1838) ...	...	—	—	—	—	Q.	—	B.fl.II,114	
P. mucronata; F. v. M., fragm. I, 8 (1858) ...	...	S.A.	—	V.	N.S.W.	—	—	B.fl.II,114	M.fr.I,8.
P. polifolia; Cunningham in Field's N. S. Wales 346 (1825) ...	...	—	—	—	N.S.W.	—	—	B.fl.II,115	
P. petiolaris; Cunningham in Ann. des Wien. Mus. II, 82 (1838) ...	...	—	—	—	—	Q.	—	B.fl.II,115	
P. paleacea; Willdenow, spec. plant. II, 506 (1799) ...	...	—	—	V.	N.S.W.	—	—	B.fl.II,115	

P. Gunnii; Benth. in Ann. des Wien. Mus. II, 82 (1839)	...	...	...	...	...	T.	V.	N.S.W.	—	—	B.fl.II, 116	
P. scabra; R. Brown in Ait. hort. Kew. sec. ed. III, 18 (1811)	...	...	...	...	...	S.A.	—	V.	N.S.W.	—	—	B.fl.II, 116
P. Hartmanni; F. v. M., fragm. VIII, 166 (1874)	...	...	...	...	...	—	—	—	N.S.W.	Q.	—	M.fr. VIII, 166.
P. microphylla; Sieber in De Candolle prodr. II, 11 (1825)	...	...	...	...	...	—	—	—	N.S.W.	Q.	—	B.fl.II, 117
P. Skinneri; F. v. M., fragm. VIII, 166 (1874)	...	...	...	...	...	W.A.	—	—	—	—	—	M.fr. VIII, 166.
P. Drummondii; Meissner in Lehm. plant. Preiss. II, 219 (1847)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II, 117
P. pinifolia; Meissner in Lehm. plant. Preiss. II, 220 (1847)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II, 118
P. pedunculata; Hooker, Bot. Magaz. t. 2859 (1828)	...	...	...	...	...	—	S.A.	T.	V.	N.S.W.	—	B.fl.II, 118
P. conferta; Benth. Fl. Austr. II, 118 (1864)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II, 118
P. aciphylla; Benth. in Hueg. enum. 35 (1837)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II, 119
P. ochreate; Meissner in Lehm. plant. Preiss. I, 73 (1844)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II, 120
P. aspalathoides; Meissner in Lehm. plant. Preiss. I, 73 (1844)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II, 120
P. obcordata; Benth. Fl. Austr. II, 120 (1864)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II, 120
P. rotundifolia; Benth. Fl. Austr. II, 121 (1864)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II, 121 M.fr. I, 145.
P. calycina; Benth. Fl. Austr. II, 121 (1864)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II, 121
P. spinulosa; Benth. Fl. Austr. II, 121 (1864)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II, 121
P. tenella; Benth. Fl. Austr. II, 122 (1864)	...	...	...	...	...	—	—	—	V.	—	—	B.fl.II, 122
P. ternata; F. v. M., fragm. I, 8 (1858)	...	...	...	...	...	—	—	—	V.	N.S.W.	Q.	B.fl.II, 122 M.fr. I, 242; III, 21; IV, 10.
P. styphelioides; Cunningham in G. Don, gen. ayst. II, 124 (1832)	...	...	...	...	...	—	—	—	V.	N.S.W.	—	B.fl.II, 122 M.fr. III, 21.
P. obovata; Benth. Fl. Austr. II, 123 (1864)	...	...	...	...	...	—	—	—	N.S.W.	—	—	B.fl.II, 123
P. incurvata; Cunningham in Field's N. S. Wales 346 (1825)	...	...	...	...	...	—	—	—	N.S.W.	—	—	B.fl.II, 124
P. subumbellata; Hooker, Bot. Mag. t. 3254 (1833)	...	...	...	...	...	—	—	T.	V.	N.S.W.	—	B.fl.II, 124
P. stipularis; Smith, Specim. of the Bot. of New Holl. 35, t. 12 (1793)	...	...	...	...	...	—	—	—	N.S.W.	—	—	B.fl.II, 125
P. glabra; Benth. Fl. Austr. II, 124 (1864)	...	...	...	...	...	—	—	—	N.S.W.	—	—	B.fl.II, 125
P. dentata; Labillardière, Nov. Holl. pl. spec. I, 103, t. 131 (1804)	...	...	...	...	...	—	—	T.	V.	N.S.W.	—	B.fl.II, 125
P. aristata; Sieber in De Candolle, prodr. II, 112 (1825)	...	...	...	...	...	—	—	—	N.S.W.	—	—	B.fl.II, 126
P. Baeuerlenii; F. v. M. in proceed. Linn. Soc. N.S.W. sec. ser. I, 1109 (1886)	...	...	...	...	...	—	—	—	N.S.W.	—	—	—
P. plumosa; Sieber in De Candolle, prodr. II, 111 (1825)	...	...	...	...	...	—	—	—	N.S.W.	—	—	B.fl.II, 126
P. viscosa; R. Brown in Benth. Fl. Austr. II, 127 (1864)	...	...	...	...	...	—	—	—	V.	N.S.W.	—	B.fl.II, 127
P. echinula; Sieber in De Candolle, prodr. II, 112 (1825)	...	...	...	...	...	—	—	—	N.S.W.	Q.	—	B.fl.II, 127
P. hibernioides; J. Hooker, Fl. Tasm. I, 89 (1860)	...	...	...	...	...	—	—	T.	V.	N.S.W.	—	B.fl.II, 127
P. rosea; F. v. M., fragm. II, 15 (1860)	...	...	...	...	...	—	—	—	V.	—	—	B.fl.II, 128 M.fr. II, 15.
P. mollis; Lindley in Mitch. Three Exped. II, 260 (1838)	...	...	...	...	...	—	S.A.	—	V.	—	—	B.fl.II, 128
P. strobilifera; Meissner in Lehm. pl. Preiss. I, 75 (1844)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II, 128
P. ericifolia; Benth. in Bot. Regist. XXV, App. XIII (1839)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II, 129
P. verruculosa; Turczaninow in Bull. Soc. Mosc. XXVI, 278 (1853)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II, 129
P. empetrifolia; Meissner in Lehm. plant. Preiss. I, 76 (1844)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II, 129
P. adunca; Turczaninow in Bull. Soc. Mosc. XXVI, 279 (1853)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II, 130
P. neurocalyx; Turczaninow in Bull. Soc. Mosc. XXVI, 297 (1853)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II, 130
P. rigida; R. Brown in Benth. Fl. Austr. II, 130 (1864)	...	...	...	...	...	—	S.A.	—	—	—	—	B.fl.II, 130
P. juniperina; Labillardière, Nov. Holl. pl. spec. I, 102, t. 130 (1804)	...	...	...	...	...	—	—	T.	V.	N.S.W.	—	B.fl.II, 131
P. acerosa; R. Brown in Benth. Fl. Austr. II, 131 (1864)	...	...	...	...	...	—	S.A.	—	—	—	—	B.fl.II, 131
P. humilis; Benth. in J. Hook. Fl. Tasm. I, 91 (1860)	...	...	...	...	...	—	S.A.	T.	V.	N.S.W.	—	B.fl.II, 131
P. parviflora; Sieber in De Candolle, prodr. II, 111 (1825)	...	...	...	...	...	—	—	—	V.	N.S.W.	—	B.fl.II, 132
P. setulosa; Benth. Fl. Austr. II, 132 (1864)	...	...	...	...	...	—	—	—	—	Q.	—	B.fl.II, 132
P. vestita; R. Brown in Ait. hort. Kew. sec. ed. III, 19 (1811)	...	...	...	...	...	W.A.	S.A.	—	—	—	—	B.fl.II, 132
P. procumbens; Cunningham in Field's N. S. Wales 347 (1838)	...	...	...	...	...	—	—	—	N.S.W.	—	—	B.fl.II, 133
P. hispidula; R. Brown in Benth. Fl. Austr. II, 133 (1864)	...	...	...	...	...	—	—	—	N.S.W.	—	—	B.fl.II, 133
P. laxiflora; Benth. Fl. Austr. I, 133 (1864)	...	...	...	...	...	—	S.A.	—	V.	—	—	B.fl.II, 133
P. largiflorens; F. v. M. in Benth. Fl. Austr. II, 134 (1864)	...	...	...	...	...	—	S.A.	—	V.	—	—	B.fl.II, 134

<i>P. villosa</i> ; Willdenow, spec. plant. II, 507 (1799)	—	—	—	V.	N.S.W.	Q.	—	B.fl.II,134	
<i>P. foliolosa</i> ; Cunningham in Ann. des Wien. Mus. II, 83 (1838)	—	—	—	V.	N.S.W.	—	—	B.fl.II,135	
<i>P. flexilis</i> ; Smith in Koen. & Sims, Ann. of Bot. I, 502 (1805) ..	—	—	—	V.	N.S.W.	—	—	B.fl.II,135	
<i>P. euchila</i> ; De Candolle, prodr. II, 112 (1825)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,135	
<i>P. selaginoides</i> ; J. Hooker, Fl. Tasm. I, 87 (1860) ..	—	—	—	T.	—	—	—	B.fl.II,136	
<i>P. densifolia</i> ; F. v. M. in Transact. Vict. Inst. 119 (1854)...	—	S.A.	—	V.	—	—	—	B.fl.II,136	
<i>P. elliptica</i> ; Smith in Transact. Linn. Soc. IX, 246 (1808)...	—	—	—	—	N.S.W.	—	—	B.fl.II,136	
<i>P. subspicata</i> ; Benth. Fl. Austr. II, 137 (1864)	—	—	—	—	N.S.W.	—	—	B.fl.II,137	
<i>P. villifera</i> ; Sieber in De Candolle, prodr. II, 111 (1825) ..	—	S.A.	V.	—	N.S.W.	—	—	B.fl.II,137	
<i>P. involuerata</i> ; Benth. Fl. Austr. II, 138 (1864)	—	S.A.	—	—	—	—	—	B.fl.II,138	
<i>P. Muelleri</i> ; Benth. Fl. Austr. II, 138 (1864)	—	—	—	V.	—	—	—	B.fl.II,138	
<i>P. polyota</i> ; F. v. M. in J. Hooker, Fl. Tasm. I, 89 (1860)...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.II,138	
<i>P. canaliculata</i> ; F. v. M. in Transact. Vict. Inst. 119 (1854) ..	—	S.A.	—	V.	—	—	—	B.fl.II,139	
<i>P. fasciculata</i> ; Benth. in Ann. des Wien. Mus. II, 82 (1838)	—	—	—	T.	V.	N.S.W.	—	B.fl.II,139	
<i>P. tenuifolia</i> ; R. Brown in Bot. Mag. t. 2086 (1819)	W.A.	S.A.	T.	V.	—	—	—	B.fl.II,139	M.fr.I,9.
<i>P. viscidula</i> ; Tate in Transact. Roy. Soc. of S. Austr. VI, 140 (1883)	—	S.A.	—	—	—	—	—	—	

LATROBEA ; Meissner in Lehm. plant. Preiss. II, 219 (1847). (Leptocytisus.)

<i>L. pungens</i> ; Benth. Fl. Austr. II, 140 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,140	M.fr.106.
<i>L. genistoides</i> ; Meissner in Lehm. plant. Preiss. II, 219 (1847)	W.A.	—	—	—	—	—	—	B.fl.II,141	
<i>L. Brunonis</i> ; Meissner in Lehm. plant. Preiss. II, 219 (1847)	W.A.	—	—	—	—	—	—	B.fl.II,141	
<i>L. tenella</i> ; Benth. Fl. Austr. II, 141 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,141	
<i>L. hirtella</i> ; Benth. Fl. Austr. II, 142 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,142	
<i>L. diosmifolia</i> ; Benth. Fl. Austr. II, 142 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,142	

EUTAXIA ; R. Brown in Ait. hort. Kew. sec. ed. III, 16 (1811). (Sclerothamnus.)

<i>E. cuneata</i> ; Meissner in Lehm. plant. Preiss. I, 65 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,143	
<i>E. myrtifolia</i> ; R. Brown in Ait. hort. Kew. sec. ed. III, 16 (1811)	W.A.	—	—	—	—	—	—	B.fl.II,144	
<i>E. epacridoides</i> ; Meissner in Lehm. plant. Preiss. I, 64 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,144	
<i>E. virgata</i> ; Benth. in Hueg. enum. 34 (1837)	W.A.	—	—	—	—	—	—	B.fl.II,144	
<i>E. densifolia</i> ; Turczaninow in Bull. Soc. Mosc. XXVI, 271 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,145	
<i>E. dillwynioides</i> ; Meissner in Lehm. plant. Preiss. I, 63 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,145	
<i>E. parvifolia</i> ; Benth. in Hueg. enum. 34 (1837)	W.A.	—	—	—	—	—	—	B.fl.II,145	
<i>E. empetrifolia</i> ; Schlechtendal, Linnæa XX, 667 (1847)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.II,145	M.fr.I,7.

DILLWYNIA ; Smith in Koenig & Sims, Ann. of Bot. I, 510 (1805).

<i>D. hispida</i> ; Lindley in Mitch. Three Exped. II, 251 (1838)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.II,147	
<i>D. ericifolia</i> ; Smith in Koen. & Sims, Ann. of Bot. I, 510 (1805)...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,147	
<i>D. floribunda</i> ; Smith in Koen. & Sims, Ann. of Bot. I, 510 (1805)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,149	
<i>D. Preissii</i> ; Benth. Fl. Austr. II, 149 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,149	
<i>D. brunoides</i> ; Meissner in Lehm. plant. Preiss. I, 62 (1844)	—	—	—	—	N.S.W.	—	—	B.fl.II,150	
<i>D. juniperina</i> ; Sieber in Hueg. enum. plant. N. Holl. occ. 33 (1837)	—	—	—	V.	N.S.W.	Q.	—	B.fl.II,150	
<i>D. pungens</i> ; Mackay in den Annual. des Wien. Mus. II, 79 (1833)	W.A.	—	—	—	—	—	—	B.fl.II,150	
<i>D. cinerascens</i> ; R. Brown in Bot. Mag. t. 2247 (1821)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.II,151	
<i>D. divaricata</i> ; Benth. Fl. Austr. II, 151 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,151	
<i>D. patula</i> ; F. v. M. fragm. IV, 16 (1864)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.II,151	M.fr.IV,16.

EUCHILOPSIS ; F. v. M. in Melbourne Chemist, June (1882).

<i>E. linearis</i> ; F. v. M. in Melb. Chem., June (1882)	W.A.	—	—	—	—	—	—	B.fl.II,67	
--	------	---	---	---	---	---	---	------------	--

PLATYLOBIUM ; Smith in Transact. Linn. Soc. II, 350 (1794).

<i>P. formosum</i> ; Smith in Transact. Linn. Soc. II, 350 (1794)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.II,153	M.fr.IX,157.
<i>P. obtusangulum</i> ; Hooker, Bot. Mag. t. 3258 (1833)	—	S.A.	T.	V.	—	—	—	B.fl.II,153	

<i>P. triangulare</i> ; R. Brown in Ait. hort. Kew. sec. ed. IV, 266 (1812) ...	...	...	—	S.A.	T.	V.	—	—	—	B.fl.II,152	M.fr.IX,158.
<i>P. alternifolium</i> ; F. v. M. in Wing's S. Sc. Rec. III, 99 (1883) ...	...	...	—	—	—	V.	—	—	—	—	—

BOSSIAEA; Ventenat, Jardin de Cels. I, 7, t. 7 (1800). (Lalage, Scottia.)

<i>B. dentata</i> ; Bentham, Fl. Austr. II, 156 (1864) ...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,156
<i>B. Aquifolium</i> ; Bentham, Fl. Austr. II, 157 (1864) ...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,157
<i>B. Webbii</i> ; F. v. M. in Melb. Chemist No. 56, p. 65 (1882) ...	...	...	...	...	W.A.	—	—	—	—	—	—
<i>B. strigillosa</i> ; Bentham, Fl. Austr. II, 157 (1864) ...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,157
<i>B. cordigera</i> ; Bentham in J. Hook. Fl. Tasm. I, 95, t. 16 (1860)...	...	...	...	...	—	—	T.	V.	—	—	B.fl.II,157 M.fr.III,100; IX,45.
<i>B. lenticularis</i> ; Sieber in De Candolle, prodr. II, 117 (1825) ...	...	...	...	...	—	—	—	—	N.S.W.	—	B.fl.II,157
<i>B. Kiamensis</i> ; Bentham, Fl. Austr. II, 158 (1864) ...	...	...	...	...	—	—	—	—	N.S.W.	—	B.fl.II,158
<i>B. ornata</i> ; F. v. M., fragm. IV, 13 (1863) ...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,158 M.fr.IX,45.
<i>B. eriocarpa</i> ; Bentham in Hueg. enum. 36 (1837) ...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,159 M.fr.IX,45.
<i>B. divaricata</i> ; Turczaninow in Bull. Soc. Mosc. XXVI, 285 (1853) ...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,159 M.fr.IV,12.
<i>B. calycina</i> ; Bentham, Fl. Austr. II, 159 (1864) ...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,159
<i>B. foliosa</i> ; Cunningham in Field's N. S. Wales, 347 (1825)...	...	...	...	...	—	—	—	V.	N.S.W.	—	B.fl.II,160 M.fr.III,100; IX,45.
<i>B. cinerea</i> ; R. Brown in Ait. hort. Kew. sec. ed. IV, 268 (1812) ...	...	...	...	...	—	—	S.A.	T.	V.	N.S.W.	B.fl.II,160 M.fr.II,98.
<i>B. biloba</i> ; Bentham in Hueg. enum. 36 (1837) ...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,160 M.fr.IX,45.
<i>B. Preissii</i> ; Meissner in Lehm. pl. Preiss. I, 82 (1844) ...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,161 M.fr.IX,45.
<i>B. concinna</i> ; Bentham, Fl. Austr. II, 161 (1864) ...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,161
<i>B. carinalis</i> ; Bentham in Mitch. Trop. Austr. 290 (1848) ...	...	...	...	...	—	—	—	—	—	Q.	B.fl.II,161 M.fr.IX,45.
<i>B. rupicola</i> ; Cunningham in Benth. Fl. Austr. II, 162 (1864) ...	...	...	...	...	—	—	—	—	—	Q.	B.fl.II,162
<i>B. linophylla</i> ; R. Brown in Ait. hort. Kew. sec. ed. IV, 268 (1812) ...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,162 M.fr.IX,45.
<i>B. disticha</i> ; Lindley, Bot. Regist. t. 55 (1841) ...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,162
<i>B. prostrata</i> ; R. Brown in Ait. hort. Kew. sec. ed. IV, 268 (1812) ...	...	...	...	...	—	—	S.A.	T.	V.	N.S.W.	Q. — B.fl.II,162 M.fr.I,9; III,97.
<i>B. Scortechinii</i> ; F. v. M. in Wing's S. Sc. Record III, 1 (1883) ...	...	...	...	...	—	—	—	—	—	N.S.W.	—
<i>B. Neo-Anglica</i> ; F. v. M., fragm. V, 106 (1866) ...	...	...	...	...	—	—	—	—	—	N.S.W.	— M.fr.V,106.
<i>B. buxifolia</i> ; Cunningham in Field's N. S. Wales, 348 (1825) ...	...	...	...	...	—	—	—	V.	N.S.W.	Q.	B.fl.II,163 M.fr.I,9; III,98; IX,45.
<i>B. Brownii</i> ; Bentham, Fl. Austr. II, 163 (1864) ...	...	...	...	...	—	—	—	—	—	Q.	B.fl.II,163
<i>B. rhombifolia</i> ; Sieber in De Candolle, prodr. II, 117 (1825) ...	...	...	...	...	—	—	—	—	N.S.W.	Q.	B.fl.II,164 M.fr.III,99; IX,45.
<i>B. pulchella</i> ; Meissner in Lehm. pl. Preiss. I, 84 (1844) ...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,164
<i>B. microphylla</i> ; Smith in Transact. Linn. Soc. IX, 303 (1808) ...	...	...	...	...	—	—	—	V.	N.S.W.	—	B.fl.II,164 M.fr.III,99.
<i>B. peduncularis</i> ; Turczaninow in Bull. Soc. Mosc. XXVI, 287 (1853) ...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,165
<i>B. Stephensonii</i> ; F. v. M. in proceed. L. S. of N.S.W. sec. ser. I, 1107 (1886) ...	...	...	...	...	—	—	—	—	N.S.W.	—	—
<i>B. heterophylla</i> ; Ventenat, Jardin de Cels. t. 7 (1800) ...	...	...	...	...	—	—	—	V.	N.S.W.	Q.	B.fl.II,165 M.fr.I,10; III,97.
<i>B. rufa</i> ; R. Brown in Ait. hort. Kew. sec. ed. IV, 267 (1812) ...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,165 M.fr.III,97; IX,45.
<i>B. bracteosa</i> ; F. v. M. in Benth. Fl. Austr. II, 166 (1864) ...	...	...	...	...	—	—	—	V.	—	—	B.fl.II,166
<i>B. riparia</i> ; Cunningham in Benth. Fl. Austr. II, 166 (1864) ...	...	...	...	...	—	—	S.A.	T.	V.	N.S.W.	— B.fl.II,166
<i>B. ensata</i> ; Sieber in De Candolle, prodr. II, 117 (1825) ...	...	...	...	...	—	—	—	V.	N.S.W.	Q.	B.fl.II,167 M.fr.II,120; XI,27&123.
<i>B. scolopendria</i> ; Smith in Transact. Linn. Soc. IX, 307 (1808) ...	...	...	...	...	—	—	—	—	N.S.W.	—	B.fl.II,167 M.fr.II,120; III,95.
<i>B. Walkeri</i> ; F. v. M., fragm. II, 120 (1861) ...	...	...	...	...	—	—	S.A.	—	N.S.W.	—	B.fl.II,167 M.fr.III,97,166; IX,45; XI,27.
<i>B. Armitii</i> ; F. v. M., fragm. IX, 44 (1875) ...	...	...	...	...	—	—	—	—	—	Q.	— M.fr.IX,44.
<i>B. phylloclada</i> ; F. v. M. in Transact. phil. Inst. Viet. III, 52 (1859) ...	...	...	...	...	—	—	—	—	—	N.A.	B.fl.II,168 M.fr.II,120.

TEMPLETONIA; R. Brown in Ait. hort. Kew. sec. ed. 269 (1812).

<i>T. retusa</i> ; R. Brown in Ait. hort. Kew. sec. ed. IV, 269 (1812) ...	...	...	...	...	W.A.	S.A.	—	—	—	—	B.fl.II,169 M.fr.IX,158.
<i>T. Drummondii</i> ; Bentham, Fl. Austr. II, 169 (1864) ...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,169
<i>T. Muelleri</i> ; Bentham, Fl. Austr. II, 169 (1864) ...	...	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	B.fl.II,169 M.fr.I,9; II,178; IX,158.
<i>T. aculeata</i> ; Bentham, Fl. Austr. II, 170 (1864) ...	...	...	...	...	W.A.	S.A.	—	—	N.S.W.	—	B.fl.II,170 M.fr.II,120.
<i>T. egena</i> ; Bentham, Fl. Austr. II, 170 (1864) ...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A. B.fl.II,170 M.fr.III,94; IX,158.
<i>T. Battii</i> ; F. v. M. in Melb. Chemist new ser. II, 31 (1887) ...	...	...	...	...	W.A.	S.A.	—	—	—	—	—
<i>T. sulcata</i> ; Bentham, Fl. Austr. II, 171 (1864) ...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	—	B.fl.II,171 M.fr.III,94,163.

HOVEA; R. Brown in Ait. hort. Kew. sec. ed. IV, 275 (1812). (Poirotia, Plagiolobium, Platychilum.)

H. linearis; R. Brown in Ait. hort. Kew. sec. ed. IV, 275 (1812)...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,172	
H. heterophylla; Cunningham in J. Hook. Fl. Tasm. I, 93, t. 15 (1860)	...	...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,172	M.fr.IX,158.
H. longifolia; R. Brown in Ait. hort. Kew. sec. ed. IV, 275 (1812)	...	...	—	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.II,172	M.fr.IX,158.
H. acutifolia; Cunningham in G. Don, gen. syst. II, 126 (1832)	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,174	M.fr.IX,158.
H. longipes; Benth in Hueg. enum. 37 (1837)	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,174	M.fr.IX,158.
H. acanthoclada; F. v. M., fragm. IV, 15 (1864)	...	...	...	...	...	...	W.A.	—	—	—	B.fl.II,174 M.fr.IV,15.
H. chorizemifolia; De Caudolle, prodr. II, 116 (1825)	...	...	...	...	...	...	W.A.	—	—	—	B.fl.II,174 M.fr.IX,158.
H. Celsi; Bonpland, plant. cult. à Malmaison. t. 51 (1813)	...	...	...	...	...	...	W.A.	—	—	—	B.fl.II,175 M.fr.IX,158.
H. trisperma; Benth in Hueg. enum. 37 (1837)	...	...	...	...	...	...	W.A.	—	—	—	B.fl.II,176 M.fr.IX,158.
H. stricta; Meissner in Lehm. pl. Preiss. I, 79 (1844)	...	...	...	...	...	...	W.A.	—	—	—	B.fl.II,176 M.fr.IX,158.
H. pungens; Benth in Hueg. enum. 37 (1837)	...	...	...	...	...	...	W.A.	—	—	—	B.fl.II,176

NEMATOPHYLLUM; F. v. M. in Hook. Kew. miscell. IX, 20 (1857).

N. Hookeri; F. v. M. in Hook. Kew. miscell. IX, 20 (1857)	...	...	...	...	—	S.A.	—	—	—	Q. N.A.	B.fl.II,170 M.fr.I,169; IV,170; IX,157.
---	-----	-----	-----	-----	---	------	---	---	---	---------	---

GOODIA; Salisbury, paradis. Londin. t. 41 (1806).

G. lotifolia; Salisbury, parad. Lond. t. 41 (1806)	...	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,177	M.fr.I,10; IX,157.
G. medicaginea; F. v. M., Second Gen. Rep. 11 (1854)	...	...	...	...	...	W.A.	S.A.	—	V.	—	—	—	—	M.fr.I,10,242; IX,157.

CROTALARIA; Hermann, hort. acad. Lugd. Bat. 196—202, t. 197—203 (1687). (Pentadynamis.)

C. verrucosa; Linné, spec. plant. 715 (1753)	—	—	—	—	Q. N.A. B.fl.II,179	M.fr.III,54; IX,156.	As., Afr., Am., P.	
C. crispata; F. v. M. in Benth. Fl. Austr. II, 179 (1864)	—	—	—	—	Q. N.A. B.fl.II,179	M.fr.III,55.		
C. juncea; Linné, spec. plant. 714 (1753)	—	—	—	—	Q. N.A. B.fl.II,179	M.fr.III,51; IX,156.	As., P.	
C. linifolia; Linné, fl. suppl. 322 (1781)	—	—	—	N.S.W.	Q. N.A. B.fl.II,180	M.fr.III,55; IX,156.	As., P.	
C. humifusa; Graham in Wall. numeric. list 5421 (1828)	—	—	—	—	Q. N.A. —	M.fr.VI,225; IX,156.	As.	
C. calycina; Schranck, pl. rar. hort. Monac. t. 12 (1819)	—	—	—	—	Q. N.A. B.fl.II,180	M.fr.III,56; IX,156.	As., P.	
C. retusa; Linné, spec. plant. 715 (1753)	—	—	—	N.S.W.	Q. N.A. B.fl.II,181	M.fr.III,51; IX,156.	As., Afr., Am.	
C. Mitchellii; Benth in Mitch. Trop. Austr. 120 (1848)	—	S.A.	—	N.S.W.	Q. N.A. B.fl.II,181	M.fr.III,50; IX,156.		
C. Novae-Hollandiae; De Caudolle, prodr. II, 127 (1825)	W.A.	—	—	—	Q. N.A. B.fl.II,181	M.fr.III,55; IX,156.		
C. crassipes; Hooker, icon. plant. t. 830 (1852)	—	—	—	—	—	N.A. B.fl.II,182		
C. Cunninghamii; B. Brown, App. to Sturt's Centr. Austr. 8 (1849)	W.A.	S.A.	—	—	Q. N.A. B.fl.II,182	M.fr.III,52; IX,156.		
C. medicaginea; Lamarck, encycl. meth. II, 201 (1786)	W.A.	—	—	N.S.W.	Q. N.A. B.fl.II,183	M.fr.III,56; IX,157.	As., Afr., Am., P.	
C. incana; Linné, spec. plant. 716 (1753)	W.A.	—	—	N.S.W.	Q. N.A. B.fl.II,183	M.fr.III,53; IX,157.	As., Afr., Am., P.	
C. dissitiflora; Benth in Mitch. Trop. Austr. 386 (1848)	W.A.	S.A.	—	N.S.W.	Q. N.A. B.fl.II,184	M.fr.III,56; IX,156.		
C. striata; De Caudolle, prodr. II, 137 (1825)	—	—	—	—	Q. —	—	As., Afr.	
C. laburnifolia; Linné, spec. plant. 715 (1753)	—	—	—	—	Q. —	B.fl.II,184	M.fr.III,52; IX,157.	As., P.
C. quinquefolia; Linné, spec. plant. 716 (1753)	—	—	—	—	Q. —	B.fl.II,184	M.fr.IX,157	As., P.
C. alata; Hamilton in D. Don, prodr. Fl. Nepal. 241 (1825)	—	—	—	—	Q. N.A. —	M.fr.XI,138	As.	

WESTONIA; Sprengel, syst. veg. III, 152 (1826).

W. trifoliata; Sprengel, syst. veg. III, 230 (1826)	...	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.II,185	...	...	...	As.
---	-----	-----	-----	-----	-----	---	---	---	---	---	---	------	-------------	-----	-----	-----	-----

TRIGONELLA; Linné, gen. pl. 351 (1737).

T. suavissima; Lindley in Mitch. Three Exped. I, 255 (1838)	...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.II,187	
---	-----	-----	-----	-----	-----	------	------	---	----	--------	----	---	-------------	--

LOTUS; Tournefort, inst. 402, t. 227 (1700) from C. Bauhin (1623).

L. corniculatus; Linné, spec. plant. 775 (1753)	...	...	...	...	...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.II,188	...	...	...	E., As., Afr.
L. australis; Andrews, Botanist's Reposit. t. 624 (1810)	...	...	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.II,188	M.fr.XI,20&27	...	P.

PSORALEA; Linné, gen. ed. sec. 358 (1742).

P. badocana; Benth in Fl. Austr. II, 190 (1864)	...	...	...	...	...	—	—	—	—	—	Q. N.A.	B.fl.II,190	M.fr.IX,156	...	...	...	As., P.
P. cephalantha; F. v. M., fragm. IV, 35 (1863)	...	...	...	...	...	—	—	—	—	—	Q. N.A.	—	M.fr.IV,35.				

P. Martini; F. v. M., fragm. V, 11 (1865)	—	—	—	—	—	N.A.	—	M.fr.V,11.
P. Archeri; F. v. M., fragm. IV, 21 (1863)	—	—	—	—	—	N.A.	B.fl.II,190	M.fr.IV,21.
P. balsamica; F. v. M. in Transact. Vict. Inst. III, 55 (1858)	—	S.A.	—	—	—	N.A.	B.fl.II,191	
P. plumosa; F. v. M., fragm. IV, 22 (1864)	—	—	—	—	—	Q. N.A.	B.fl.II,191	M.fr.IV,22.
P. pustulata; F. v. M. in Transact. Vict. Inst. III, 54 (1858)	W.A.	—	—	—	—	N.A.	B.fl.II,191	
P. lachnostachys; F. v. M., fragm. III, 105 (1863)	—	—	—	—	—	N.A.	B.fl.II,191	M.fr.III,195.
P. eriantha; Benth in Mitch. Trop. Austr. 131 (1848)	W.A.	S.A.	—	V.	N.S.W.	Q. N.A.	B.fl.II,192	M.fr.III,45.
P. patens; Lindley in Mitch. Three Exped. II, 9 (1838)	W.A.	S.A.	—	V.	N.S.W.	Q. N.A.	B.fl.II,192	M.fr.IX,155; III,45; IV,22.
P. cinerea; Lindley in Mitch. Three Exped. II, 65 (1838)	W.A.	—	—	N.S.W.	—	Q. N.A.	B.fl.II,192	M.fr.III,45; IV,22.
P. leucantha; F. v. M. in Transact. Vict. Inst. III, 54 (1859)	W.A.	S.A.	—	—	—	Q. N.A.	B.fl.II,193	M.fr.IX,156.
P. parva; F. v. M. in Transact. Vict. Inst. I, 40 (1854)	—	S.A.	—	V.	N.S.W.	—	B.fl.II,194	M.fr.IX,155.
P. tenax; Lindley in Mitch. Three Exped. II, 10 (1838)	—	—	—	V.	N.S.W.	Q. —	B.fl.II,193	M.fr.III,45; IX,155.
P. adscendens; F. v. M. in Transact. Vict. Inst. I, 40 (1854)	—	S.A.	T.	V.	N.S.W.	—	B.fl.II,193	M.fr.III,45; IX,155.
P. Testariae; F. v. M., fragm. V, 45 (1865)	—	—	—	—	—	Q. —	—	M.fr.V,45,189.

INDIGOFERA; Royen, pl. hort. Lugd. 372 (1740), from Isnard (1724).

I. linifolia; Retzius, observ. bot. IV, 29 (1786)	W.A.	S.A.	—	N.S.W.	Q. N.A.	B.fl.II,195	M.fr.III,101; IX,44	... As., Afr., P.
I. cordifolia; Heyne in Roth, nov. pl. spec. 357 (1821)	—	—	—	—	—	N.A.	B.fl.II,196	... As., Afr.
I. eorneaphylla; Linné, mantissa altera 272 (1771)	—	S.A.	—	N.S.W.	Q. N.A.	B.fl.II,196	M.fr.III,102; IX,44	... As., Afr. P.
I. glandulosa; Willdenow, spec. plant. III, 1227 (1800)	—	—	—	—	Q. —	B.fl.II,196	M.fr.IV,22; VII,106; IX,44.	As.
I. haplophylla; F. v. M., fragm. III, 102 (1863)	—	—	—	—	Q. N.A.	B.fl.II,196	M.fr.IX,44.	
I. trifoliata; Linné, amoen. acad. IV, 327 (1759)	—	—	—	—	Q. —	B.fl.II,197	M.fr.III,164	... As., P.
I. trita; Linné fil., suppl. plant. 335 (1781)	—	—	—	N.S.W.	Q. N.A.	B.fl.II,197	M.fr.III,103	... As., Afr., P.
I. parviflora; Heyne in Wall. numeric. list 5457 (1828)	—	—	—	—	Q. N.A.	B.fl.II,197	M.fr.III,103; IX,44	... As., Afr.
I. viscosa; Lamarck, Encycl. méth. III, 247 (1789)	—	S.A.	—	—	Q. N.A.	B.fl.II,198	M.fr.III,104; IX,44	... As., Afr.
I. hirsuta; Linné, spec. plant. 751 (1753)	—	S.A.	—	N.S.W.	Q. N.A.	B.fl.II,198	M.fr.III,105; IX,44	... As., Afr., Am., P.
I. pratensis; F. v. M., essay on plants of Burdek. Exped. 10 (1860)	—	—	—	—	Q. —	B.fl.II,198		
I. Schultzeana; F. v. M., fragm. VII, 105 (1870)	—	—	—	—	N.A.	—	M.fr.VII,105.	
I. monophylla; De Candolle, prodr. II, 222 (1825)	W.A.	S.A.	—	—	N.A.	B.fl.II,199	M.fr.III,44; VII,106; IX,44.	
I. rugosa; Benth in Mitch. Fl. Austr. II, 199 (1864)	—	—	—	—	N.A.	B.fl.II,199		
I. saxicola; F. v. M. in Benth. Fl. Austr. II, 199 (1864)	—	—	—	—	N.A.	B.fl.II,199		
I. Australis; Willdenow, spec. plant. III, 1235 (1800)	W.A.	S.A.	T.	V.	N.S.W.	Q. —	B.fl.II,199	
I. brevidens; Benth in Mitch. Trop. Austr. 385 (1848)	W.A.	S.A.	—	N.S.W.	Q. N.A.	B.fl.II,200		
I. efoliata; F. v. M. in Melb. Chemist. Nov. (1883)	—	—	—	N.S.W.	—	—		
I. Baileyi; F. v. M., fragm. IX, 43 (1875)	—	—	—	—	Q. —	—	M.fr.IX,43.	
I. coronillifolia; Cunningham in Benth. Fl. Austr. II, 201 (1864)	—	S.A.	—	N.S.W.	—	B.fl.II,201		

PTYCHOSEMA; Benth in Lindl. Bot. Regist. XXV. App. XVI (1839).

P. pusillum; Benth in Lindl. Bot. Regist. XXV, App. XVI (1839)	W.A.	—	—	—	—	B.fl.II,201		
P. anomalum; F. v. M., fragm. IX, 62 (1875)	—	S.A.	—	—	—	—	M.fr.IX,62.	
P. trifoliolatum; F. v. M. in Wing's S. Science Record II, 72 (1882)	W.A.	S.A.	—	N.S.W.	—	—		

LAMPROLOBIUM; Benth in Mitch. Fl. Austr. II, 202 (1864).

L. fruticosum; Benth in Mitch. Fl. Austr. II, 202 (1864)	—	—	—	—	Q. —	B.fl.II,202	M.fr.IX,68.	
L. megalophyllum; F. v. M., fragm. IX, 67 (1875)	—	—	—	—	N.A.	—	M.fr.IX,67.	

TEPHROSIA; Persoon, synops. pl. II, 328 (1807).

T. coriacea; Benth in Mitch. Fl. Austr. II, 204 (1864)	—	—	—	—	N.A.	B.fl.II,204	M.fr.IX,65&66.	
T. lamprolobioides; F. v. M., fragm. IX, 64 (1875)	—	—	—	—	N.A.	—	M.fr.IX,64.	
T. Forrestiana; F. v. M., fragm. XI, 98 (1880)	—	—	—	—	N.A.	—	M.fr.XI,98.	
T. flammea; F. v. M. in Benth in Mitch. Fl. Austr. II, 204 (1864)	—	—	—	—	Q. N.A.	B.fl.II,204		
T. reticulata; Benth in Mitch. Fl. Austr. II, 205 (1864)	—	—	—	—	Q. N.A.	B.fl.II,205	M.fr.IX,64.	

T. crocea; Benth. Fl. Austr. II, 205 (1864)	...	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.II,205					
T. oblongata; Benth. Fl. Austr. II, 205 (1864)	...	...	...	...	...	—	—	—	—	—	—	—	N.A.	B.fl.II,205				
T. porrecta; Benth. Fl. Austr. II, 206 (1864)	...	...	...	...	...	—	—	—	—	—	—	—	N.A.	B.fl.II,206				
T. polyzyga; F. v. M. in Benth. Fl. Austr. II, 206 (1864)	...	...	...	...	...	—	—	—	—	—	—	—	N.A.	B.fl.II,206				
T. graminifolia; F. v. M. in Benth. Fl. Austr. II, 206 (1864)	...	...	...	...	...	—	—	—	—	—	—	—	N.A.	B.fl.II,206	M.fr.IX,6			
T. simplicifolia; F. v. M. in Benth. Fl. Austr. II, 206 (1864)	...	...	...	...	...	—	—	—	—	—	—	—	N.A.	B.fl.II,206				
T. nematophylla; F. v. M., fragm. IX, 63 (1879)	...	...	...	...	...	—	—	—	—	—	—	—	N.A.	—	M.fr.IX,63.			
T. leptoclada; Benth. Fl. Austr. II, 207 (1864)	...	...	...	...	...	—	—	—	—	—	—	—	Q.	N.A.	B.fl.II,207			
T. brachycarpa; F. v. M. in Benth. Fl. Austr. II, 207 (1864)	...	...	...	...	...	—	—	—	—	—	—	—	—	N.A.	B.fl.II,207			
T. uniovulata; F. v. M., fragm. XI, 70 (1879)	...	...	...	...	...	—	—	—	—	—	—	—	—	N.A.	—	M.fr.IX,70.		
T. Stuartii; Benth. Fl. Austr. II, 207 (1864)	...	...	...	...	...	—	—	—	—	—	—	—	—	N.A.	B.fl.II,207			
T. conferta; F. v. M., fragm. IX, 65 (1879)	...	...	...	...	...	—	—	—	—	—	—	—	—	N.A.	—	M.fr.IX,75.		
T. eriocarpa; Benth. Fl. Austr. II, 207 (1864)	...	...	...	...	...	—	—	—	—	—	—	—	—	N.A.	B.fl.II,207			
T. phaeosperma; F. v. M. in Benth. Fl. Austr. II, 207 (1864)	...	...	...	...	...	—	—	—	—	—	—	—	—	N.A.	B.fl.II,208			
T. astragaloides; Benth. Fl. Austr. II, 208 (1864)	...	...	...	...	...	—	—	—	—	—	—	—	Q.	—	B.fl.II,208			
T. juncea; Benth. Fl. Austr. II, 208 (1864)	...	...	...	...	...	—	—	—	—	—	—	—	Q.	N.A.	B.fl.II,208	M.fr.IX,64		
T. filipes; Benth. Fl. Austr. II, 208 (1864)	...	...	...	...	...	—	—	—	—	—	—	—	Q.	N.A.	B.fl.II,208			
T. sphaerospora; F. v. M. in Wing's S. Sc. Record III, 128 (1883)	...	...	...	...	...	—	S.A.	—	—	—	—	—	—	N.A.	—			
T. remotiflora; F. v. M. in Benth. Fl. Austr. II, 209 (1864)	...	...	...	...	...	—	—	—	—	—	—	—	—	N.A.	B.fl.II,209			
T. oligophylla; Benth. Fl. Austr. II, 209 (1864)	...	...	...	...	...	—	—	—	—	—	—	—	Q.	N.A.	B.fl.II,209			
T. macrocarpa; Benth. Fl. Austr. II, 209 (1864)	...	...	...	...	...	—	—	—	—	—	—	—	—	N.A.	B.fl.II,209			
T. purpurea; Persoon, synops. plant. II, 329 (1807)	...	...	...	...	...	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.II,209	...	..	As, Afr., P.		
T. Bidwilli; Benth. Fl. Austr. II, 210 (1864)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,210					
T. rosea; F. v. M. in Benth. Fl. Austr. II, 211 (1864)	...	...	...	...	...	—	—	—	—	—	—	—	N.A.	B.fl.II,211				
WISTARIA; Nuttall, Gen. of N. Amer. pl. II, 115 (1818). (Milletia.)																		
W. megasperma; F. v. M., fragm. I, 10 (1858)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,211	M.fr.I,10.				
W. australis; F. v. M. in Britten's Journ. of Bot. XXII, 290 (1884)	...	...	...	...	...	—	—	—	—	N.S.W.	—	—	—					
SESBANIA; Scopoli, Introd. 308 (1777). (Agati from Alpino 1592.)																		
S. grandiflora; Persoon, synops. pl. II, 316 (1807)	...	...	...	...	...	W.A.	—	—	—	—	Q.	N.A.	B.fl.II,212	M.fr.II,88; XI,33	...	As.		
S. Aegyptiaca; Persoon, synops. pl. II, 316 (1807)	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.II,212	M.fr.XI,33	...	As, Afr		
S. aculeata; Persoon, synops. pl. II, 316 (1807)	...	...	...	...	...	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.II,213	M.fr.XI,33	...	As, Afr., P.		
S. simpliciuscula; F. v. M. in Benth. Fl. Austr. II, 213 (1864)	...	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.II,213					
S. brachycarpa; F. v. M., fragm. XI, 32 (1878)	...	...	...	...	...	—	—	—	—	—	—	N.A.	—	M.fr.IX,32.				
CARMICHAELIA; R. Brown in Bot. Regist. XI, 912 (1825).																		
C. exsul; F. v. M., fragm. VII, 126 (1871)	...	...	...	...	...	—	—	—	—	N.S.W.	—	—	—	M.fr.VII,121,193.				
CLIANTHUS; Banks & Solander in G. Don, dichlam. pl. II, 468; indicative (1832). (Donia.)																		
C. Dampieri; Cunningham in Trans. hort. soc. Lond. sec. ser. I, 522 (1835)	...	...	...	...	...	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.II,214					
STREBLORRHIZA; Endlicher, prodr. fl. Norfolk. 92 (1833).																		
S. speciosa; Endlicher, prodr. fl. Norf. 93 (1833)	...	...	...	...	...	—	—	—	—	N.S.W.	—	—	—					
SWAINSONIA; Salisbury, parad. Londin. t. 28 (1806). (Cyclogyne, Diplobium.)																		
S. Greyana; Lindley, Bot. Regist. t. 66 (1846)	...	...	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.II,216	M.fr.IX,154; X,7.				
S. galiegifolia; R. Brown in Ait. hort. Kew. sec. ed. III, 327 (1811)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,217					
S. coronillifolia; Salisbury, paradis. Lond. t. 28 (1806)	...	...	...	...	...	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.II,217	M.fr.XI,70.				
S. brachycarpa; Benth. Fl. Austr. II, 217 (1864)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,217	M.fr.IX,155.				
S. phacoides; Benth. in Mitch. Trop. Austr. 363 (1848)	...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.II,217					
S. Burkittii; F. v. M. in Benth. Fl. Austr. II, 218 (1864)	...	...	...	...	...	—	S.A.	—	—	N.S.W.	—	—	B.fl.II,218					
S. Burkei; F. v. M. in Benth. Fl. Austr. II, 218 (1864)	...	...	...	...	...	W.A.	S.A.	—	—	—	—	N.A.	B.fl.II,218	M.fr.IX,155.				

S. oligophylla; F. v. M. in Benth. Fl. Austr. II, 219 (1864)	W.A.	S.A.	—	—	N.S.W.	—	N.A.	B.fl.II,219			
S. campylantha; F. v. M. in Gregory's pap. rel. to Leichh. Exp. 6 (1858) ...	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.II,219	M.fr.III,46; IX,154.		
S. occidentalis; F. v. M., fragm. III, 46 (1862)	W.A.	—	—	—	—	—	N.A.	B.fl.II,219	M.fr.III,46.		
S. stenodonta; F. v. M., fragm. XI, 70 (1879)	W.A.	—	—	—	—	—	N.A.	—	M.fr.XI,70.		
S. oncinotropis; F. v. M. in Melb. Chem. Oct. (1884)	—	—	—	V.	N.S.W.	—	—	—			
S. gracilis; Benth. Fl. Austr. II, 220 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,220			
S. plagiotropis; F. v. M., fragm. IX, 153 (1875)	—	—	—	V.	N.S.W.	—	—	—	M.fr.IX,143.		
S. procumbens; F. v. M., fragm. III, 46 (1862)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.II,220	M.fr.I,76; III,46; IX,154; XI,27.		
S. Drummondii; Benth. Fl. Austr. II, 220 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,220			
S. Oliveri; F. v. M. in Wing's S. Sc. Rec. II, 152 (1882)	W.A.	S.A.	—	—	N.S.W.	—	—	—			
S. canescens; F. v. M., fragm. III, 46 (1862)... ..	W.A.	—	—	—	—	—	—	B.fl.II,221	M.fr.III,46; IX,154; XI,70.		
S. Beasleyana; F. v. M. in Melb. Chemist, new ser. II, 84 (1882) ...	W.A.	—	—	—	—	—	—	—			
S. stipularis; F. v. M. in Linnaea XXV, 393 (1852)... ..	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.II,221	M.fr.I,76; IX,154.		
S. oroboides; F. v. M. in Benth. Fl. Austr. II, 222 (1864)... ..	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.II,222			
S. Kingii; F. v. M. in Transact. Roy. Soc. Vict. XXIII, 53 (1886) ...	W.A.	—	—	—	—	—	N.A.	—			
S. unifoliolata; F. v. M., fragm. VIII, 226 (1874)	W.A.	S.A.	—	—	—	—	—	—	M.fr.VIII,226.		
S. lessertiifolia; De Candolle, prodr. II, 271 (1825)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.II,222	M.fr.IX,154.		
S. luteola; F. v. M., fragm. I, 75 (1858)	—	—	—	—	—	Q.	—	B.fl.II,223	M.fr.XI,70.		
S. parviflora; Benth. Fl. Austr. II, 223 (1864)	—	—	—	—	—	Q.	—	B.fl.II,223			
S. monticola; Cunningham in A. Gray's Bot. Wilk. Exp. I, 411 (1854) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,223	M.fr.IX,155.		
S. microphylla; A. Gray, Bot. Wilk. Exped. I, 410 (1854)... ..	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.II,223	M.fr.IX,155.		
S. laxa; R. Brown in Sturt's Centr. Austr. App. 18 (1849)... ..	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.II,224			
S. colutoides; F. v. M., fragm. X, 6 (1876)	W.A.	S.A.	—	—	—	—	—	—	M.fr.X,6.		
S. Fraseri; Benth. Fl. Austr. II, 224 (1864)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,224			
S. Macullochiana; F. v. M., fragm. VII, 25 (1869)	W.A.	—	—	—	—	—	N.A.	—	M.fr.IX,154.		

GLYCYRRHIZA; Tournefort, inst. 389, t. 210 (1700) from Dioscorides and Plinius. (Clidanthera.)

G. psoraleoides; Benth. Fl. Austr. II, 225 (1864)... ..	—	S.A.	—	V.	N.S.W.	—	—	B.fl.II,225	M.fr.III,45.		
---	---	------	---	----	--------	---	---	-------------	--------------	--	--

ORMOCARPUM; Palisot, Fl. d'Ovar. I. 95, t. 58 (1805).

O. scnnoides; De Candolle, prodr. II, 315 (1825)	—	—	—	—	—	Q.	—	B.fl.II,226	...	...	As.
---	---	---	---	---	---	----	---	-------------	-----	-----	-----

AESCHYNOMENE; Linné, gen. pl. 350 (1737).

A. Indica; Linné, spec. plant. 713 (1753)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.II,227	M.fr.XII,18	...	As., Afr., P.
A. falcata; De Candolle, prodr. II, 322 (1825)	—	—	—	—	—	Q.	—	B.fl.II,227	M.fr.XII,18	...	Afr., Am.
A. Americana; Linné, spec. plant. 713 (1753)	—	—	—	—	—	Q.	—	—	M.fr.XII,18	...	As., Am.

SMITHIA; Aiton, hort. Kew. III, 496 (1789).

S. capitata; Desvaux, Journ. de Bot. I, (1813)	—	—	—	—	—	Q.	N.A.	B.fl.II,228	...	...	As., P.
S. sensitiva; Aiton, hort. Kew. III, 496, t. 13 (1789)	—	—	—	—	—	Q.	—	—	M.fr.VII,27; IX,193.	...	As., Afr.

ZORNIA; Gmelin, Syst. Nat. 1076 (1791).

Z. diphylla; Persoon, synops. II, 318 (1807)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.II,228	...	...	As., Afr., Am., P.
Z. chaetophora; F. v. M. in Transact. phil. Inst. Vict. III, 56 (1857) ...	—	—	—	—	—	—	N.A.	B.fl.II,229			

DESMODIUM; Desvaux, Journ. de Bot. III, 122 (1813). (Dicerna, Phyllodium, Pteroloma.)

D. umbellatum; De Candolle, prodr. II, 325 (1825)	—	—	—	—	—	Q.	—	B.fl.II,230	M.fr.IX,66	...	As., Afr., Am.
D. pulchellum; Benth. Fl. Hongk. 83 (1861)	—	—	—	—	—	—	N.A.	B.fl.II,231	...	...	As., P.
D. biarticulatum; F. v. M., fragm. II, 121 (1861)	—	—	—	—	—	Q.	N.A.	B.fl.II,231	M.fr.IX,66	...	As.
D. acanthocladum; F. v. M., fragm. II, 122 (1861)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,231	M.fr.IX,66.		
D. triquetrum; De Candolle, prodr. II, 326 (1825)	—	—	—	—	—	Q.	—	—	M.fr.IX,66	...	As.
D. Gangeticum; De Candolle, prodr. II, 327 (1825)	—	—	—	—	—	Q.	N.A.	B.fl.II,232	...	...	As., As., P.
D. dependens; Blume in Miq. Fl. Ind. Batav. I, 249 (1855)	—	—	—	—	—	Q.	—	—	M.fr.VIII,225,284; IX,66&195; X,120.	...	As., P.

D. brachypodium; A. Gray, Bot. Wilk. Exped. 434 (1854)	—	—	—	V.	N.S.W.	Q.	—	B.fl.II,232				
D. varians; Endlicher in Ann. des Wien. Mus. I, 185 (1839)	—	—	—	T.	V.	N.S.W.	Q.	—	B.fl.II,232	...	...	P.
D. flagellare; Benth. Fl. Austr. II, 233 (1864)	—	—	—	—	—	—	—	N.A.	B.fl.II,233			
D. rhytidophyllum; F. v. M. in Benth. Fl. Austr. II, 223 (1864)	—	—	—	—	—	N.S.W.	Q.	—	B.fl.II,233			
D. campylocaulon; F. v. M. in Benth. Fl. Austr. II, 233 (1864)	—	—	—	—	—	—	—	Q.	N.A. B.fl.II,233			
D. nemorosum; F. v. M. in Benth. Fl. Austr. II, 234 (1864)	—	—	—	—	—	N.S.W.	Q.	—	B.fl.II,234			
D. neurocarpum; Benth. Fl. Austr. II, 234 (1864)	—	—	—	—	—	—	—	N.A.	B.fl.II,234			
D. trichostachyum; Benth. Fl. Austr. II, 234 (1864)	—	—	—	—	—	—	—	Q.	N.A. B.fl.II,234			
D. polycarpum; De Candolle, prodr. II, 334 (1825)	—	—	—	—	—	N.S.W.	Q.	N.A. B.fl.II,235	M.fr.IX,66	...	...	Aa., Afr., P.
D. trichocaulon; De Candolle, prodr. II, 335 (1825)	—	—	—	—	—	N.S.W.	Q.	—	B.fl.II,235			
D. Muelleri; Benth. Fl. Austr. II, 235 (1864)	—	—	—	—	—	—	—	Q.	N.A. B.fl.II,235			
D. reniforme; De Candolle, prodr. II, 327 (1825)	—	—	—	—	—	—	—	Q.	N.A. —	M.fr.XII	...	As.
D. parvifolium; De Candolle, prodr. II, 334 (1825)	—	—	—	—	—	N.S.W.	Q.	N.A. B.fl.II,235	M.fr.IX,66	...	...	As., P.
PYCNOSPORA; R. Brown in Wight & Arnott, prodr. I, 197 (1834).												
P. hedyaroides; R. Brown in Wight & Arnott, prodr. 197 (1834)	—	—	—	—	—	—	—	Q.	N.A. B.fl.II,236	M.fr.V,30	...	As., P.
URARIA; Desvaux, Journ. de Bot. III, 122 (1813).												
U. picta; Desvaux, Journ. de Bot. III, 122 (1813)	—	—	—	—	—	N.S.W.	Q.	—	B.fl.II,237	...	...	As., Afr., P.
U. cylindracea; Benth. Fl. Austr. II, 237 (1864)	—	—	—	—	—	—	—	Q.	N.A. B.fl.II,237			
U. lagopoides; De Candolle, prodr. II, 324 (1825)	—	—	—	—	—	—	—	Q.	N.A. B.fl.II,237	...	...	As., P.
LOUREA; Necker, elem. Bot. III, 17 (1790).												
L. obcordata; Desvaux, Journ. de Bot. III, 122 (1813)	—	—	—	—	—	—	—	—	N.A. B.fl.II,238	...	...	As., P.
ALYSICARPUS; Necker, elem. Bot. III, 15 (1790).												
A. vaginalia; De Candolle, prodr. II, 353 (1825)	—	—	—	—	—	—	—	Q.	N.A. B.fl.II,239	...	...	As., Afr., P.
A. longifolia; Wight & Arnott, prodr. 223 (1834)	—	—	—	—	—	—	—	—	N.A. B.fl.II,239	...	...	As.
A. cylindricus; Desvaux in Annal. Soc. Linn. Par. 301 (1825)	—	—	—	—	—	—	—	Q.	N.A. B.fl.II,239	...	...	As., Afr.
LESPEDEZA; Cl. Richard in Michaux, fl. bor. am. II, 70 (1803).												
L. cuneata; G. Don, gen. syst. II, 307 (1832)	—	—	—	—	—	V.	N.S.W.	Q.	—	B.fl.II,240	...	As.
L. lanata; Benth. Fl. Austr. II, 241 (1864)	—	S.A.	—	—	—	—	—	—	N.A. B.fl.II,241			
CLITORIA; Linné, gen. plant. 216 (1737), from Petiver.												
C. Ternatea; Linné, sp. pl. 753 (1753)	—	—	—	—	—	—	—	Q.	—	—	...	As., Afr., P.
C. Australis; Benth. Fl. Austr. II, 242 (1864)	—	—	—	—	—	—	—	—	N.A. B.fl.II,242			
GLYCINE; Linné, gen. pl. 349 (1737). (Leptolobium, Leptocyanus.)												
G. falcata; Benth. Fl. Austr. II, 243 (1864)	—	S.A.	—	—	—	N.S.W.	Q.	N.A. B.fl.II,243				
G. clandestina; Wendland, Bot. Beob. 54 (1798)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,243				
G. Latrobeana; Benth. Fl. Austr. II, 224 (1864)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.II,244				
G. tabacina; Benth. Fl. Austr. II, 244 (1864)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.II,244	...	...	...	P.
G. sericea; Benth. Fl. Austr. II, 245 (1864)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A. B.fl.II,245					
G. tomentosa; Benth. Fl. Austr. II, 245 (1864)	—	S.A.	—	—	N.S.W.	Q.	N.A. B.fl.II,245	...	...	...	As., P.	
KENNEDYA; Ventenat, Jardin de la Malmaison II, t. 104—106 (1804). (Caulina, Moench 1802, Hardenbergia, Physolobium, Zichya, Amphodus.)												
K. nigricans; Lindley, Bot. Regist. t. 1715 (1835)	W.A.	—	—	—	—	—	—	B.fl.II,249	M.fr.VII,127.			
K. Beckxiana; F. v. M., fragm. XI, 98 (1880)	W.A.	—	—	—	—	—	—	—	M.fr.XI,98.			
K. rubicunda; Ventenat, Jard. de la Malm. t. 104 (1804)	—	—	—	V.	N.S.W.	Q.	—	B.fl.II,249	M.fr.VII,127.			
K. procurrens; Benth. in Mitch. trop. Austr. 365 (1848)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,249				
K. prorepens; F. v. M., fragm. VIII, 225 (1874)	—	S.A.	—	—	—	—	—	—	M.fr.VIII,225.			
K. prostrata; R. Brown in Ait. hort. Kew. sec. ed. IV, 299 (1812)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.II,250	M.fr.VII,128.			
K. bracteata; Gaudichaud in Freyc. voy. 286, t. 113 (1826)	W.A.	—	—	—	—	—	—	B.fl.II,250	M.fr.IV,79.			

K. eximia; Lindley in Paxt. Mag. XVI, 35 (1849)	W.A.	—	—	—	—	—	B.fl.II,250				
K. microphylla; Meissner in Lehm. pl. Preiss. I, 91 (1844)	W.A.	—	—	—	—	—	B.fl.II,251				
K. parviflora; Meissner in Lehm. pl. Preiss. I, 91 (1844)	W.A.	—	—	—	—	—	B.fl.II,251				
K. Stirlingii; Lindley, Bot. Regist. t. 1845 (1837)	W.A.	—	—	—	—	—	B.fl.II,252				
K. glabrata; Lindley Bot. Regist. t. 1838 (1837)	W.A.	—	—	—	—	—	B.fl.II,252	M.fr.VII,128.			
K. macrophylla; F. v. M., fragm. IV, 79 (1864)	W.A.	—	—	—	—	—	B.fl.II,252	M.fr.IV,79.			
K. coccinea; Vntenat, Jard. de la Malm. t. 105 (1804)	W.A.	—	—	—	—	—	B.fl.II,250	M.fr.VII,128.			
K. monophylla; Ventenat, Jard. de la Malm. t. 106 (1804)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,246	M.fr.VII,128.		
K. Comptoniana; Link, enum. pl. hort. bot. Berol. II, 235 (1822)	W.A.	—	—	—	—	—	—	B.fl.II,247	M.fr.VII,128.		
K. retusa; F. v. M., fragm. V, 106 (1866)	—	—	—	—	—	—	Q.	—	B.fl.II,247	M.fr.VII,128	... P.
ERYTHRINA; Linné, gen. pl. 216 (1737).											
E. vesperilio; Benth in Mitch. Trop. Austr. 218 (1848)	W.A.	S.A.	—	—	—	Q.	N.A.	B.fl.II,253			
E. Indica; Lamarck, Encyclop. méthod. II, 391 (1786)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.II,253	...	...	As., P.
STRONGYLODON; T. Vogel in Linnaea X, 585 (1836).											
S. ruber; T. Vogel in Linnaea X, 585 (1836)... ..	—	—	—	—	—	Q.	—	—	...	...	As., P.
MUCUNA; Maregraf, hist. nat. Brazil. 18 (1648).											
M. gigantea; De Candolle, prodr. II, 405 (1825)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.II,254	...	...	As., P.
GALACTIA; P. Browne, Civil & Nat. Hist. of Jamaica 298 (1756).											
G. tenuiflora; Wight & Arnott, prodr. fl. penins. Ind. 206 (1834)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.II,255	...	...	As., Afr., Am., P.
G. Muellieri; Benth in Fl. Austr. II, 255 (1864)	—	—	—	—	—	—	N.A.	B.fl.II,255			
CANAVALIA; De Candolle, prodr. II, 403 (1825). (Canavali.)											
C. obtusifolia; De Candolle, prodr. II, 404 (1825)	W.A.	—	—	—	N.S.W.	Q.	N.A.	B.fl.II,256	...	...	As., Afr., Am., P.
PHASEOLUS; Tournefort, inst. 412, t. 232 (1700), from Dioscorides & Columella.)											
P. Truxillensis; Humb. Bonpl. & Kunth, nov. gen. Amer. VI, 451 (1823)	—	—	—	—	—	Q.	N.A.	B.fl.II,257	...	...	As., Afr., Am., P.
P. Max; Linné, sp. pl. 725 (1753)	—	—	—	—	—	Q.	N.A.	B.fl.II,257	...	...	As., Afr., P.
VIGNA; Savi, Dissert. 6 (1824). (Callicystus.)											
V. vexillata; Benth in Mart. Fl. Bras. Papil. 193, t. 50 (1862)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.II,258	...	...	As., Afr., Am., P.
V. lutea; A. Gray, Bot. Wilk. Exped. I, 454 (1854)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.II,259	...	...	As., Afr., Am., P.
V. luteola; Benth in Mart. Fl. Bras. Papil. 194, t. 50 (1862)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,260	...	...	As., Afr., Am.
V. lanceolata; Benth in Mitch. trop. Austr. 350 (1848)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.II,260			
DOLICHOS; Linné, gen. plant. 222 (1737).											
D. biflorus; Linné, spec. plant. 727 (1753)	—	—	—	—	—	Q.	N.A.	B.fl.II,261	...	...	As., Afr.
DUNBARIA; Wight & Arnott, prodr. I, 258 (1834).											
D. conspersa; Benth in plant. Junghuhn. (1851)	—	—	—	—	—	Q.	—	B.fl.II,262	...	...	As.
CAJANUS; De Candolle, Catal. hort. Monsp. 85 (1813). (Cajan, Atylosia.)											
C. marmoratus; F. v. M., Cens. 41 (1882) from Benth. (1864)	—	—	—	—	—	Q.	N.A.	B.fl.II,263			
C. scaraboides; F. v. M., Cens. 41 (1882) from Benth. (1851)	—	—	—	—	—	Q.	—	B.fl.II,263	...	...	As., Afr.
C. reticulatus; F. v. M., Cens. 41 (1882) from Benth. (1864)	—	—	—	—	—	Q.	N.A.	B.fl.II,263			
C. cinereus; F. v. M., Cens. 41 (1882)... ..	W.A.	—	—	—	—	—	N.A.	B.fl.II,264			
C. grandifolius; F. v. M., Essay on pl. Burdekin Exped. 9 (1860)	—	—	—	—	—	Q.	N.A.	B.fl.II,264			
C. confertiflorus; F. v. M., Essay on pl. Burdekin Exped. 9 (1860)	—	—	—	—	—	Q.	—	B.fl.II,264			
RHYNCHOSIA; Loureiro, Fl. Cochinch. II, 460 (1790).											
R. rhomboidea; F. v. M. in Benth. Fl. Austr. II, 265 (1864)	—	—	—	—	—	—	N.A.	B.fl.II,265			
R. acutifolia; Benth in Fl. Austr. II, 266 (1864)	—	—	—	—	—	Q.	N.A.	B.fl.II,266			

R. rostrata ; Benth. Fl. Austr. II, 266 (1864)	...	...	...	...	...	—	—	—	—	—	N.A.	B.fl.II,266				
R. Cunninghamii ; Benth. Fl. Austr. II, 266 (1864)	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl.II,266	..	...	P.
R. minima ; De Candolle, prodr. II, 385 (1825)	...	...	...	...	...	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.II,267	...	..	As., Afr., Am., P.
ERIOSEMA ; De Candolle, prodr. II, 388 (1825).																
E. Chinense ; T. Vogel in Meyen's Beitr. zur Bot. 31 (1843)	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.II,268	...	...	As., P.
FLEMINGIA ; Roxburgh in Ait. hort. Kew. sec. ed. IV, 349 (1812).																
F. lineata ; Roxburgh in Ait. hort. Kew. sec. ed. IV, 350 (1812)	...	...	..	—	—	—	—	—	—	—	Q.	N.A.	B.fl.II,268	...	...	As., P.
F. pauciflora ; Benth. Fl. Austr. II, 269 (1864)	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.II,269			
F. parviflora ; Benth. Fl. Austr. II, 269 (1864)	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl.II,269			
F. involucrata ; Benth. in pl. Jungh. I, 246 (1851)	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl.II,269	...	...	As.
ABRUS ; Dalechamps, hist. general. plant. append. 193 (1587), from Rauwolf (1582).																
A. precatorius ; Linné, syst. natur. ed. XII, 533 (1767)	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.II,270	...	...	As., Afr., Am., P.
DALBERGIA ; Linné, fil. suppl. 52 (1781).																
D. densa ; Benth. in Hook. Lond. Journ. II, 217 (1843)	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl.II,271	...	...	P.
LONCHOCARPUS ; Humboldt, Bonpland & Kunth, nov. gen. VI, 383 (1823).																
L. Blackii ; Benth. Fl. Austr. II, 271 (1864)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,271	M.fr.II,123.		
DERRIS ; Loureiro, Fl. Cochinch. II, 432 (1790). (Brachypterum.)																
D. scandens ; Benth. in Journ. Linn. Soc. IV, suppl. 103 (1860)	...	...	—	—	—	—	—	—	—	N.S.W.	Q.	—	B.fl.II,272	...	...	As.
D. uliginosa ; Benth. in pl. Jungh. I, 252 (1851)	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.II,272	...	...	As., Afr., P.
PONGAMIA ; Lamarck in Ventenat, Jardin de la Malmaison t. 28 (1803) from Adanson (1763).																
P. glabra ; Ventenat, Jard. Malm. t. 28 (1803)	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.II,273	...	..	As., P.
SOPHORA ; Linné, gen. plant. 125 (1737). (Calpurnia partim.)																
S. tomentosa ; Linné, spec. plant. 373 (1753)	..	...	...	...	...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.II,274	M.fr.VII,27	...	As., Afr., Am., P.
S. Fraseri ; Benth. Fl. Austr. II, 274 (1864)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,274	M.fr.V,31 ; VII,27.		
S. tetraptera ; J. Miller, icon. plant. t. 1 (1780)	...	...	...	...	...	—	—	—	—	N.S.W.	—	—	—	M.fr.VII,26	...	N.Z.
PODOPETALEM ; F. v. M., in Melbourne Chemist, June (1882).																
P. Ormondi ; F. v. M. in Melb. Chem. June (1882)	...	...	...	...	...	—	—	—	—	—	Q.	—	—			
CASTANOSPERMUM ; Cunningham & Fraser in Hook. bot. misc. I, 241, t. 51 (1830).																
C. Australe ; Cunningham & Fraser in Hook. bot. misc. I, 241, t. 51 (1830)	..	—	—	—	—	—	—	—	—	N.S.W.	Q.	—	B.fl.II,275	M.fr.VII,27.		
BARKLYA ; F. v. M., fragm. I, 109 (1859).																
B. syringifolia ; F. v. M. in Journ. Linn. Soc. III, 158 (1859)	...	...	...	—	—	—	—	—	—	N.S.W.	Q.	—	B.fl.II,275	M.fr.I,109 ; III,163 ; VIII,27.		
CAESALPINA ; Plumier, nov. pl. Amer. gen. 28, t. 9 (1703) from Ammand. (Guilandina, Caesalpinia 1810.)																
C. Bonducella ; Fleming in Asiat. Research XI, 159 (1801)	...	...	...	—	—	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.II,276	...	...	As., Afr., Am., P.
C. Nuga ; W. T. Aiton, hort. Kew. sec. ed. III, 32 (1811)	...	...	...	—	—	—	—	—	—	—	Q.	—	B.fl.II,277	M.fr.X,7	...	As., P.
C. sepiaria ; Roxburgh, Fl. Indic. II, 360 (1832)	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl.II,277	M.fr.X,7	...	As.
MEZONEURON ; Desfontaines in Mém. du Mus. IV, 245, t. 10 et 11 (1818).																
M. brachycarpum ; Benth. Fl. Austr. II, 278 (1864)	...	...	...	—	—	—	—	—	—	N.S.W.	Q.	—	B.fl.II,278	M.fr.X,7.		
M. Scortechinii ; F. v. M. in Wing's South. Sc. Record II, 73 (1882)	...	...	—	—	—	—	—	—	—	N.S.W.	Q.	—	—			
PTEROLOBIUM ; R. Brown in Salt. Abyssin. 65 (1814).																
P. nitens ; F. v. M. in Benth. Fl. Austr. II, 279 (1864)	...	...	...	—	—	—	—	—	—	—	Q.	—	B.fl.II,279	M.fr.X,7.		
PELTOPHORUM ; Vogel in Linnaea XI, 406 (1837).																
P. ferrugineum ; Benth. Fl. Austr. II, 279 (1864)	...	...	...	—	—	—	—	—	—	—	Q.	N.A.	B.fl.II,279	M.fr.X,7	...	As.

CASSIA ; Tournefort, inst. 619, t. 392 (1700), from Plumier (1693). (Cathartocarpus.)

C. Brewsteri ; F. v. M., fourth ann. Rep. 17 (1858)...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,282	M.fr.I,110; X,8.				
C. laevigata ; Willdenow, enum. pl. hort. bot. Berol. I, 441 (1809)	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,282	M.fr.IV,14; X,8	...	Am.		
C. Sophera ; Linné, spec. plant. 379 (1753)	...	...	...	...	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.II,283	M.fr.III,49; IV,14; X,9.		As.,	P.	
C. laxiflora ; Benth, Fl. Austr. II, 283 (1864)	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.II,283					
C. magnifolia ; F. v. M., fragm. I, 166 (1859)	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.II,283	M.fr.X,9.				
C. venusta ; F. v. M., fragm. I, 165 (1859)	...	...	...	...	...	W.A.	S.A.	—	—	Q.	N.A.	B.fl.II,284	M.fr.X,9.				
C. notabilis ; F. v. M., fragm. III, 28 (1862)...	...	...	...	...	...	W.A.	S.A.	—	—	—	N.A.	B.fl.II,284	M.fr.X,9.				
C. pleurocarpa ; F. v. M., fragm. I, 223 (1859)	...	...	...	...	...	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.II,284	M.fr.X,9.			
C. glauca ; Lamarck, Encycl. méth. I, 647 (1783)	...	...	...	...	...	—	—	—	—	S.W.	Q.	N.A.	B.fl.II,285	M.fr.IV,13; X,9	...	As.,	P.
C. retusa ; Solander in Linnaea XI, 72 (1842)	...	...	...	...	...	—	—	—	—	Q.	—	B.fl.II,285					
C. Australis ; Sims in Bot. Mag. t. 2676 (1826)	...	...	...	...	...	—	—	—	V.	N.S.W.	Q.	N.A.	B.fl.II,285	M.fr.X,9.			
C. Chatelainiana ; Gaudichaud in Freyc. voy. 485, t. 111 (1826)	...	...	...	...	...	W.A.	—	—	—	—	N.A.	B.fl.II,286					
C. glutinosa ; De Candolle, prodr. II, 495 (1825)	...	...	...	...	...	—	S.A.	—	—	—	N.A.	B.fl.II,286					
C. Cuthbertsoni ; F. v. M. in Vict. Nat. V, 75 (1888)	...	...	...	...	...	W.A.	—	—	—	—	—	—					
C. pruinosa ; F. v. M., fragm. III, 48 (1862)...	...	...	...	...	...	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.II,286	M.fr.III,48.			
C. circinata ; Benth in Mitch. trop. Austr. 384 (1848)	...	...	...	...	...	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.II,286	M.fr.I,165.			
C. phyllodinea ; R. Brown, App. to Sturt's Centr. Austr. 15 (1849)	...	...	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.II,287				
C. eremophila ; Cunningham in T. Vogel, gen. Cass. synops. 47 (1837)	...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.II,287				
C. artemisioides ; Gaudichaud in De Candolle, prodr. II, 495 (1825)	...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.II,288				
C. Sturtii ; R. Brown, App. to Sturt's Centr. Austr. 14 (1849)	...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.II,288				
C. desolata ; F. v. M. in Linnaea XXV, 387 (1852)	...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.II,289				
C. oligophylla ; F. v. M., fragm. III, 49 (1862)	...	...	...	...	...	—	—	—	—	—	N.A.	B.fl.II,289	M.fr.III,49.				
C. cardiosperma ; F. v. M., fragm. X, 50 (1877)	...	...	...	...	...	W.A.	—	—	—	—	—	—	M.fr.X,50.				
C. oligoclada ; F. v. M., fragm. III, 49 (1862)	...	...	...	...	...	—	—	—	—	—	N.A.	B.fl.II,289	M.fr.X,9.				
C. heptanthera ; F. v. M., fragm. X, 8 (1876)	...	...	...	...	...	—	—	—	—	—	N.A.	—	M.fr.X,8.				
C. leptoclada ; Benth, Fl. Austr. II, 290 (1864)	...	...	...	...	...	—	—	—	—	Q.	N.A.	B.fl.II,290					
C. Absus ; Linné, spec. plant. 376 (1753)	...	...	...	...	...	—	—	—	—	Q.	N.A.	B.fl.II,290	M.fr.III,50; X,9	...	As.,	Afr.	
C. pumila ; Lamarck, Encycl. méth. I, 620 (1783)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,290	M.fr.III,47; X,10	...	As.	
C. concinna ; Benth, Fl. Austr. II, 291 (1864)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.II,291	M.fr.X,10.			
C. mimosoides ; Linné, spec. plant. 379 (1753)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.II,291	M.fr.III,48 ; X,10	...	As.,	Afr., Am., P.

PETALOSTYLIS ; R. Brown, App. to Sturt's Centr. Austr. 79 (1849). (Petalostylos, Petalogyne.)

P. labichoides ; R. Brown, App. to Sturt's Centr. Austr. 17 (1849)	...	...	...	...	...	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.II,292	M.fr.X,7.			
--	-----	-----	-----	-----	-----	------	------	---	---	--------	----	------	-------------	-----------	--	--	--

LABICHEA ; Gaudichaud in De Candolle, prodr. II, 507 (1825).

L. cassioides ; Gaudichaud in De Candolle, prodr. II, 507 (1825)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,292					
L. nitida ; Benth, Fl. Austr. II, 293 (1864)	...	...	...	...	...	—	—	—	—	Q.	N.A.	B.fl.II,293	M.fr.X,7.				
L. lanceolata ; Benth in Hueg. enum. 41 (1837)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,293	M.fr.X,7.				
L. rupestris ; Benth in Mitch. Trop. Austr. 342 (1848)	...	...	...	...	...	—	—	—	—	Q.	—	B.fl.II,293	M.fr.X,7; XII,19.				
L. punctata ; Benth in Bot. Regist. XXV, App. XV (1839)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,294	M.fr.X,8; XII,19.				
L. Buettneriana ; F. v. M. in Melb. Chemist, June (1882)	...	...	...	...	...	—	—	—	—	Q.	—	—	M.fr.XII,18.				

TAMARINDUS ; Tournefort, inst. 660, t. 445, App. (1700) from C. Bauhin (1623).

T. Indicus ; Linné, spec. plant. 34 (1753)	...	...	...	...	...	—	—	—	—	—	N.A.	B.fl.II,294	...	...	...	As.,	Afr.
--	-----	-----	-----	-----	-----	---	---	---	---	---	------	-------------	-----	-----	-----	------	------

BAUHINIA ; Plumier, nov. gen. pl. Amer. 22, t. 13 (1703).

B. acuminata ; Linné, spec. plant. 375 (1753)...	...	...	...	...	...	—	—	—	—	Q.	—	—	...	...	...	As.	
B. Leichhardtii ; F. v. M. in Transact. Vict. Inst. III, 50 (1858)...	...	...	...	...	...	—	S.A.	—	—	Q.	N.A.	B.fl.II,295	M.fr.X,8.				
B. Carronii ; F. v. M. in Transact. Vict. Inst. III, 49 (1858)	...	...	...	...	...	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.II,295	M.fr.X,8.			
B. Hookeri ; F. v. M. in Transact. Vict. Inst. III, 51 (1858)	...	...	...	...	...	—	—	—	—	Q.	N.A.	B.fl.II,295	M.fr.X,8.				

B. Gilesii; F. v. M. & Bailey in Wing's S. Soc. Record, II, 151 (1882) ...	...	—	—	—	—	—	—	N.A.	—										
B. monandra; Kurz, Forest Flora of Burma I, 39 (1877) ...	...	...	...	—	—	—	—	—	Q	—	—	...	...	...	...	...	...	...	As.
AFZELIA; Smith in Transact. Linn. Soc. IV, 221 (1798).																			
A. bijuga; A. Gray, Bot. Wilk. Expl. Exped. 467, t. 51 (1854) ...	...	...	...	—	—	—	—	—	Q.	—	—	...	...	...	...	...	...	...	As., P.
CYNOMETRA; Linné in Act. Soc. Ups. 78 (1741).																			
C. ramiflora; Linné, spec. plant. 382 (1753) ...	...	...	...	—	—	—	—	—	Q.	—	B.fl.II,296	...	...	...	...	...	...	...	As., P.
ERYTHROPHLAEUM; Afzelius in Tuckey's Congo 430 (1818). (Laboucheria.)																			
E. Laboucherii; F. v. M. in Benth. Fl. Austr. II, 297 (1864) ...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.II,297								
ENTADA; Adanson, Familles des plantes II, 318 (1763).																			
E. Pursaetha; De Candolle, memoir. sur la fam. des Légum. 12 (1825) ...	...	...	...	—	—	—	—	—	Q.	—	B.fl.II,298	M.fr.V,179				...	...	As., Afr., Am., P.	
ADENANTHERA; Royen in Linné, coroll. 7 (1737).																			
A. pavonina; Linné, spec. plant. 384 (1753) ...	...	...	...	—	—	—	—	—	Q.	N.A.	—	M.fr.V,30				...	...	...	As.
A. abrosperma; F. v. M., fragm. V, 30 (1865) ...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.II,298	M.fr.V,31.							
NEPTUNIA; Loureiro, Fl. Cochinch. II, 653 (1790). (Dichrostachys.)																			
N. cinerea; F. v. M., Cens. 43 (1882) from Wight & Arnott (1834) ...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.II,299	...	...	...	...	...	...	...	As.
N. spicata; F. v. M., fragm. III, 151 (1862) ...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.II,299	M.fr.III,151.							
N. gracilis; Benth. in Hook. Journ. of Bot. IV, 355 (1842) ...	...	...	...	—	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.II,300							
N. monosperma; F. v. M. in Benth. Fl. Austr. II, 299 (1864) ...	...	...	...	—	S.A.	—	—	—	N.S.W.	Q.	N.A.	B.fl.II,300							
ACACIA; Tournefort, inst. 605, t. 375 (1700) from Dioscorides & Plinius. (Vachellia, Chithonanthus, Tetracheilus, Mimosa partly.)																			
I. ALATAE.																			
A. glaucoptera; Benth. in Schlecht. Linnaea XXVI, 604 (1853) ...	...	...	...	W.A.	—	—	—	—	—	—	—	B.fl.II,320							
A. alata; R. Brown in Ait. hort. Kew. sec. ed. V, 464 (1813) ...	...	...	...	W.A.	—	—	—	—	—	—	—	B.fl.II,300							
A. diptera; Lindley, Bot. Regist. XXV, App. XV (1839) ...	...	...	...	W.A.	—	—	—	—	—	—	—	B.fl.II,321							
A. stenoptera; Benth. in Hook. Lond. Journ. I, 325 (1842) ...	...	...	...	W.A.	—	—	—	—	—	—	—	B.fl.II,321							
II. CONTINUAE.																			
A. incurva; Benth. in Hook. Lond. Journ. I, 325 (1842) ...	...	...	...	W.A.	—	—	—	—	—	—	—	B.fl.II,322							
A. trigonophylla; Meissner in Lehm. pl. Preiss. II, 199 (1846) ...	...	...	...	W.A.	—	—	—	—	—	—	—	B.fl.II,322	M.fr.IV,3.						
A. continua; Benth. in Fl. Austr. II, 322 (1864) ...	...	...	...	—	S.A.	—	V.	N.S.W.	—	—	—	B.fl.II,322							
A. Peuce; F. v. M., fragm. III, 151 (1863) ...	...	...	...	—	S.A.	—	—	—	Q.	—	—	B.fl.II,323	M.fr.III,151.						
III. PUNGENTES.																			
A. spinescens; Benth. in Hook. Lond. Journ. I, 323 (1842) ...	...	...	...	—	S.A.	—	V.	N.S.W.	—	—	—	B.fl.II,323							
A. cochlearis; H. L. Wendland, comm. de Acac. aphyll. 15 (1820) ...	...	...	...	W.A.	S.A.	—	—	—	—	—	—	B.fl.II,324							
A. lanigera; Cunningham in Field's N.S.W. 345 (1825) ...	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	—	—	B.fl.II,324							
A. phlebocarpa; F. v. M. in Journ. Linn. Soc. III, 119 (1858) ...	...	...	...	—	—	—	—	—	—	Q.	N.A.	B.fl.II,325							
A. trinervata; Sieber in De Candolle, prodr. II, 451 (1825) ...	...	...	...	—	—	—	—	—	N.S.W.	—	—	B.fl.II,325							
A. colletioides; Cunningham in Hook. Lond. Journ. I, 336 (1842) ...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	—	—	—	B.fl.II,325	M.fr.IV,4.						
A. striatula; Benth. in Hook. Lond. Journ. I, 336 (1842) ...	...	...	...	W.A.	—	—	—	—	—	—	—	B.fl.II,326							
A. campylophylla; Benth. in Schlecht. Linnaea XXVI, 605 (1853) ...	...	...	...	W.A.	—	—	—	—	—	—	—	B.fl.II,326							
A. teretifolia; Benth. in Hook. Lond. Journ. I, 326 (1842) ...	...	...	...	W.A.	—	—	—	—	—	—	—	B.fl.II,326							
A. sulcata; R. Brown in Ait. hort. Kew. sec. ed. V, 460 (1813) ...	...	...	...	W.A.	—	—	—	—	—	—	—	B.fl.II,327							
A. costata; Benth. in Hook. Lond. Journ. I, 339 (1842) ...	...	...	...	W.A.	—	—	—	—	—	—	—	B.fl.II,327							
A. barbinervis; Benth. in Hook. Lond. Journ. I, 326 (1842) ...	...	...	...	W.A.	—	—	—	—	—	—	—	B.fl.II,327							
A. ataxiphylla; Benth. in Schlecht. Linnaea XXVI, 605 (1853) ...	...	...	...	W.A.	—	—	—	—	—	—	—	B.fl.II,328							
A. Baxteri; Benth. in Hook. Lond. Journ. I, 327 (1842) ...	...	...	...	W.A.	—	—	—	—	—	—	—	B.fl.II,328							
A. anreo-nitens; Lindley, Bot. Regist. XXV, App. XV (1839) ...	...	...	...	W.A.	—	—	—	—	—	—	—	B.fl.II,328							
A. quadrisulcata; F. v. M., fragm. III, 127 (1863) ...	...	...	...	W.A.	—	—	—	—	—	—	—	B.fl.II,328	M.fr.III,127.						

A. erioclada; Benth. in Schlecht. Linnaea XXVI, 606 (1853) ...	...	...	W.A.	—	—	—	—	—	—	B.f.II,329
A. sicutiformis; Cunningham in Hook. Lond. Journ. I, 337 (1842) ...	...	...	—	—	T.	V.	N.S.W.	—	—	B.f.II,329
A. patens; F. v. M. in Journ. Linn. Soc. III, 120 (1858) ...	...	...	—	—	—	—	—	N.A.	—	B.f.II,329 M.fr.III,46.
A. laricina; Meissner in Lehm. pl. Preiss. I, 6 (1844) ...	...	...	W.A.	—	—	—	—	—	—	B.f.II,330
A. tetragonophylla; F. v. M. in Journ. Linn. Soc. III, 121 (1858) ...	...	...	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.f.II,330 M.fr.IV,3.
A. genistoides; Cunningham in Benth. Fl. Austr. II, 330 (1864) ...	...	...	W.A.	S.A.	—	—	—	—	N.A.	B.f.II,330
A. sphacelata; Benth. in Hook. Lond. Journ. I, 338 (1842) ...	...	...	W.A.	—	—	—	—	—	—	B.f.II,331
A. ingrata; Benth. Fl. Austr. II, 331 (1864) ...	...	...	W.A.	—	—	—	—	—	—	B.f.II,331
A. juniperina; Willdenow, spec. plant. IV, 1049 (1805) ...	...	...	—	—	T.	V.	N.S.W.	Q.	—	B.f.II,331
A. asparagoides; Cunningham in Field's N. S. Wales 343 (1825) ...	...	...	—	—	—	—	N.S.W.	—	—	B.f.II,332
A. tenuifolia; F. v. M. in Transact. phil. Soc. Vict. I, 37 (1854) ...	...	...	—	—	—	V.	—	—	—	B.f.II,332
A. diffusa; Edwards, Bot. Regist. t. 634 (1822) ...	...	...	—	—	T.	V.	N.S.W.	—	—	B.f.II,332
A. rupicola; F. v. M. in Schlecht. Linnaea XXVI, 610 (1853) ...	...	...	—	S.A.	—	V.	—	—	—	B.f.II,333

IV. CALAMIFORMES.

A. volubilis; F. v. M., fragm. X, 98 (1877) ...	...	...	W.A.	—	—	—	—	—	—	M.fr.X,98.
A. tetragonocarpa; Meissner in Lehm. pl. Preiss. I, 4 (1844) ...	...	...	W.A.	—	—	—	—	—	—	B.f.II,336
A. restiacea; Benth. in Hook. Lond. Journ. I, 3 (1842) ...	...	...	W.A.	—	—	—	—	—	—	B.f.II,336
A. squamata; Lindley, Bot. Regist. XXV, App. XV (1839) ...	...	...	W.A.	—	—	—	—	—	—	B.f.II,336
A. brachyphylla; Benth. in Schlecht. Linnaea XXVI, 615 (1853) ...	...	...	W.A.	—	—	—	—	—	—	B.f.II,337
A. Bynoeana; Benth. in Schlecht. Linnaea XXVI, 614 (1853) ...	...	...	—	—	—	—	—	N.A.	—	B.f.II,337 M.fr.IV,9.
A. triptycha; F. v. M. in Benth. Fl. Austr. II, 337 (1864) ...	...	...	W.A.	—	—	—	—	—	—	B.f.II,337
A. leptoneura; Benth. in Hook. Lond. Journ. I, 341 (1842) ...	...	...	W.A.	—	—	—	—	—	—	B.f.II,337
A. rigens; Cunningham in Loudon, hort. Brit. II, 406 (1830) ...	...	...	—	S.A.	—	V.	N.S.W.	—	—	B.f.II,337
A. papyrocarpa; Benth. Fl. Austr. II, 338 (1864) ...	...	...	W.A.	S.A.	—	—	—	—	—	B.f.II,338
A. Gilesiana; F. v. M. in Melb. Chemist, July (1882) ...	...	...	—	S.A.	—	—	—	—	—	—
A. sessiliceps; F. v. M. in Melb. Chemist, July (1882) ...	...	...	—	S.A.	—	—	—	—	—	—
A. pugioniformis; Wendland, comm. de Acac. aphyll. 38, t. 9 (1820) ...	...	...	—	—	—	—	N.S.W.	Q.	—	B.f.II,338
A. juncifolia; Benth. in Hook. Lond. Journ. I, 341 (1842) ...	...	...	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.f.II,339
A. calamifolia; Sweet in Bot. Regist. t. 839 (1824) ...	...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.f.II,339
A. spodioperma; F. v. M. in proc. L. Soc. of N.S.W. sec. ser. III, 164 (1888) ...	...	...	W.A.	—	—	—	—	—	—	—
A. scirpifolia; Meissner in Bot. Zeitung 10 (1855) ...	...	...	W.A.	S.A.	—	—	N.S.W.	—	—	B.f.II,339
A. extensa; Lindley, Bot. Regist. XXV, App. XV (1839) ...	...	...	W.A.	—	—	—	—	—	—	B.f.II,340
A. gonophylla; Benth. in Schlecht. Linnaea XXVI, 613 (1853) ...	...	...	W.A.	S.A.	—	—	—	—	—	B.f.II,340
A. ericifolia; Benth. in Hook. Lond. Journ. I, 345 (1842) ...	...	...	W.A.	—	—	—	—	—	—	B.f.II,340
A. uncinella; Benth. in Schlecht. Linnaea XXVI, 613 (1853) ...	...	...	W.A.	—	—	—	—	—	—	B.f.II,341

V. BRUNIOIDAE.

A. cedroides; Benth. in Schlecht. Linnaea XXVI, 615 (1853) ...	...	...	W.A.	—	—	—	—	—	—	B.f.II,341
A. lycopodiifolia; Cunningham in Hook. icon. plant. t. 172 (1837) ...	...	...	—	S.A.	—	—	—	Q.	N.A.	B.f.II,342
A. spondylophylla; F. v. M., fragm. VIII, 243 (1874) ...	...	...	—	S.A.	—	—	—	—	N.A.	— M.fr.243.
A. galioides; Benth. in Hook. Lond. Journ. I, 344 (1842) ...	...	...	—	—	—	—	—	Q.	N.A.	B.f.II,342
A. Baueri; Benth. in Hook. Lond. Journ. I, 344 (1842) ...	...	...	—	—	—	—	N.S.W.	Q.	—	B.f.II,342 M.fr.XI,33.
A. subternata; F. v. M. in Journ. Linn. Soc. III, 124 (1858) ...	...	...	—	—	—	—	—	—	N.A.	B.f.II,343
A. minutifolia; F. v. M., fragm. VIII, 243 (1874) ...	...	...	—	S.A.	—	—	—	—	—	— M.fr.X,243.
A. brunioides; Cunningham in G. Don, gen. syst. II, 404 (1832) ...	...	...	—	—	—	—	—	Q.	—	B.f.II,343
A. conferta; Cunningham in Hook. Lond. Journ. I, 345 (1842) ...	...	...	—	—	—	—	N.S.W.	Q.	—	B.f.II,343 M.fr.I,6.

VI. UNINERVES.

A. scabra; Benth. in Schlecht. Linnaea XXVI, 605 (1853) ...	...	...	W.A.	—	—	—	—	—	—	B.f.II,344
A. nodiflora; Benth. in Schlecht. Linnaea XXVI, 621 (1853) ...	...	...	W.A.	—	—	—	—	—	—	B.f.II,344
A. spinosissima; Benth. in Schlecht. Linnaea XXVI, 621 (1853) ...	...	...	W.A.	—	—	—	—	—	—	B.f.II,344
A. ulicina; Meissner in Lehm. pl. Preiss. II, 202 (1846) ...	...	...	W.A.	—	—	—	—	—	—	B.f.II,345

<i>A. erinacea</i> ; Benth in Hook. Lond. Journ. I, 360 (1842)	...	...	...	W.A.	S.A.	—	—	—	—	—	B.fl.II,345	
<i>A. Huegelii</i> ; Benth in Hueg. enum. 42 (1837)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,345	
<i>A. nervosa</i> ; De Candolle, Mém. sur la fam. des Légum. 444 (1825)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,346	
<i>A. obovata</i> ; Benth in Hook. Lond. Journ. I, 329 (1842)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,346	
<i>A. congesta</i> ; Benth in Hook. Lond. Journ. I, 327 (1842)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,346	
<i>A. dermatophylla</i> ; Benth in Fl. Austr. II, 346 (1864)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,346	
<i>A. aspera</i> ; Lindley in Mitch. Three Exped. II, 139 (1838)	...	...	...	...	—	S.A.	—	V.	N.S.W.	—	B.fl.II,347	
<i>A. armata</i> ; R. Brown in Ait. hort. Kew. sec. ed. V, 463 (1813)	...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	B.fl.II,347
<i>A. idiomorpha</i> ; Cunningham in Hook. Lond. Journ. I, 329 (1842)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,348
<i>A. Shuttleworthii</i> ; Meissner in Lehm. pl. Preiss. I, 7 (1844)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,348
<i>A. Gregorii</i> ; F. v. M., fragm. III, 47 (1862)	...	...	...	...	...	—	—	—	—	—	N.A.	B.fl.II,348 M.fr.III,47.
<i>A. pilosa</i> ; Benth in Schlecht. Linnaea XXVI, 607 (1853)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,348
<i>A. crispula</i> ; Benth in Schlecht. Linnaea XXVI, 607 (1853)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,349
<i>A. crassistipulea</i> ; Benth in Hook. Lond. Journ. I, 326 (1842)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,349
<i>A. hastulata</i> ; Smith in Rees' Cyclopaed. XXXIX, Suppl. (1819)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,349
<i>A. horridula</i> ; Meissner in Lehm. pl. Preiss. I, 9 (1844)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,350
<i>A. divergens</i> ; Benth in Hook. Lond. Journ. I, 331 (1842)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,350
<i>A. vomeriformis</i> ; Cunningham in Hook. Lond. Journ. I, 332 (1842)	...	...	...	...	...	—	S.A.	T.	V.	N.S.W.	—	B.fl.II,350
<i>A. plagiophylla</i> ; F. v. M. in Journ. Linn. Soc. III, 131 (1858)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—
<i>A. biflora</i> ; R. Brown in Ait. hort. Kew. sec. ed. V, 463 (1813)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,351
<i>A. decipiens</i> ; R. Brown in Ait. hort. Kew. sec. ed. V, 463 (1813)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,351
<i>A. cuneata</i> ; Benth in Hueg. enumer. 42 (1837)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,351
<i>A. dilatata</i> ; Benth in Schlecht. Linnaea XXVI, 608 (1853)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,352
<i>A. bidentata</i> ; Benth in Hook. Lond. Journ. I, 333 (1842)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,352
<i>A. acanthoclada</i> ; F. v. M., fragm. III, 127 (1863)	...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	—	B.fl.II,352 M.fr.III,127.
<i>A. obliqua</i> ; Cunningham in Hook. Lond. Journ. I, 334 (1842)	...	...	...	...	...	—	S.A.	—	V.	N.S.W.	—	B.fl.II,353
<i>A. acinacea</i> ; Lindley in Mitch. Three Exped. II, 267 (1838)	...	...	...	...	...	—	S.A.	—	V.	N.S.W.	—	B.fl.II,353
<i>A. runciformis</i> ; Cunningham in G. Don, gen. syst. II, 404 (1832)	...	...	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	B.fl.II,353 M.fr.I,5; II,177.
<i>A. lachnophylla</i> ; F. v. M. in Wing's S. Sc. Record II, 150 (1882)	...	...	...	...	...	W.A.	—	—	—	—	—	—
<i>A. triquetra</i> ; Benth in Hook. Lond. Journ. I, 358 (1842)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,354
<i>A. ligustrina</i> ; Meissner in Lehm. pl. Preiss. II, 203 (1846)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,354
<i>A. Meissneri</i> ; Lehmann, delect. sem. hort. bot. Hamb. (1842)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,354
<i>A. anceps</i> ; De Candolle, Mém. sur la fam. des Légum. 446 (1825)	...	...	...	...	...	W.A.	S.A.	—	—	—	—	B.fl.II,355
<i>A. hispidula</i> ; Willdenow, spec. plant. IV, 1054 (1805)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	B.fl.II,355
<i>A. undulifolia</i> ; Fraser in Loddiges, Bot. Cabinet XVI, 1544 (1829)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	B.fl.II,355
<i>A. flexifolia</i> ; Cunningham in Hook. Lond. Journ. I, 359 (1842)	...	...	...	...	...	—	—	—	—	N.S.W.	—	B.fl.II,356
<i>A. dura</i> ; Benth in Schlecht. Linnaea XXVI, 622 (1853)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,356
<i>A. spathulata</i> ; F. v. M. in Benth. Fl. Austr. II, 356 (1864)	...	...	...	...	...	W.A.	—	—	—	—	N.A.	B.fl.II,356
<i>A. microcarpa</i> ; F. v. M., second gen. Report 7 (1854)	...	...	...	...	...	—	S.A.	—	V.	N.S.W.	—	B.fl.II,357 M.fr.I,6.
<i>A. montana</i> ; Benth in Hook. Lond. Journ. I, 360 (1842)	...	...	...	...	...	—	S.A.	—	V.	N.S.W.	—	B.fl.II,357
<i>A. verniciflua</i> ; Cunningham in Field's N. S. Wales 344 (1825)	...	...	...	...	...	—	S.A.	T.	V.	N.S.W.	—	B.fl.II,358
<i>A. leprosa</i> ; Sieber in De Candolle, prodr. II, 450 (1825)	...	...	...	...	...	—	—	—	V.	N.S.W.	—	B.fl.II,358
<i>A. stricta</i> ; Willdenow, spec. plant. IV, 1052 (1805)	...	...	...	...	...	—	—	T.	V.	N.S.W.	—	B.fl.II,358
<i>A. dodonaeifolia</i> ; Willdenow, enum. pl. hort. Berol. suppl. 68 (1813)	...	...	...	...	...	—	S.A.	—	—	—	—	B.fl.II,359
<i>A. Gnidium</i> ; Benth in Fl. Austr. II, 359 (1864)	...	...	...	...	...	—	—	—	—	—	Q.	B.fl.II,359
<i>A. ramosissima</i> ; Benth in Hook. Lond. Journ. I, 356 (1842)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,360
<i>A. Dempsteri</i> ; F. v. M., fragm. XI, 65 (1879)	...	...	...	...	...	W.A.	—	—	—	—	—	M.fr.XI,65.
<i>A. Sentis</i> ; F. v. M., second gen. report 11 (1854)	...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A. B.fl.II,360 M.fr.XI,65.
<i>A. dentifera</i> ; Benth in Maund, Botanist IV, t. 179 (1839)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,361
<i>A. sclerosperma</i> ; F. v. M. in Wing's South. Sc. Rec. II, 150 (1882)	...	...	...	...	...	W.A.	—	—	—	—	N.A.	—

A. fasciculifera; F. v. M. in Benth. Fl. Austr. II, 361 (1864)	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,361	
A. falcata; Willdenow, spec. plant. IV, 1053 (1805)	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,361	
A. macradenia; Benth. in Mitch. Trop. Austr. 360 (1848)	...	...	...	—	—	—	—	—	Q.	—	B.fl.II,362	
A. penninervis; Sieber in De Candolle, prodr. II, 452 (1825)	...	...	...	—	—	T.	V.	N.S.W.	Q.	—	B.fl.II,362	
A. amblyphylla; F. v. M. in Wing's S. Sc. Record, II, 149 (1882)	...	...	W.A.	—	—	—	—	—	—	—	—	
A. retinodes; Schlechtendal, Linnaea XX, 664 (1847)	...	...	—	S.A.	—	V.	—	—	—	—	B.fl.II,362	
A. nerifolia; Cunningham in Hook. Lond. Journ. I, 357 (1842)	...	...	—	—	—	—	—	N.S.W.	Q.	—	B.fl.II,363	
A. microbotrya; Benth. in Hook. Lond. Journ. I, 353 (1842)	...	...	W.A.	—	—	—	—	—	—	—	B.fl.II,363	
A. leiophylla; Benth. in Hook. Lond. Journ. I, 351 (1842)	...	...	W.A.	—	—	—	—	—	—	—	B.fl.II,364	
A. cyanophylla; Lindley, Bot. Regist. Misc. 49 (1835)	...	...	W.A.	—	—	—	—	—	—	—	B.fl.II,364	
A. pycnantha; Benth. in Hook. Lond. Journ. I, 351 (1842)	...	...	—	S.A.	—	V.	N.S.W.	—	—	—	B.fl.II,365	
A. notabilis; F. v. M., fragm. I, 6 (1858)	...	...	W.A.	S.A.	—	—	N.S.W.	—	N.A.	B.fl.II,365	M.fr.I,6.	
A. gladiiformis; Cunningham in Hook. Lond. Journ. I, 354 (1842)	...	...	—	—	—	—	—	N.S.W.	—	—	B.fl.II,365	
A. obtusata; Sieber in De Candolle, prodr. II, 453 (1825)	...	...	—	—	—	V.	N.S.W.	—	—	—	B.fl.II,366	
A. rubida; Cunningham in Field's N.S.W. 344 (1825)	...	...	—	—	—	—	—	N.S.W.	—	—	B.fl.II,366	
A. amoena; H. L. Wendland, comm. de Acac. aphyll. 10 t. 4 (1820)	...	...	—	—	—	V.	N.S.W.	—	—	—	B.fl.II,366	
A. hakeoides; Cunningham in Hook. Lond. Journ. I, 354 (1842)	...	...	—	S.A.	—	V.	N.S.W.	—	—	—	B.fl.II,367	
A. salicina; Lindley in Mitch. Three Exped. II, 20 (1838)	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.II,367		
A. rostellifera; Benth. in Hook. Lond. Journ. I, 356 (1842)	...	...	W.A.	—	—	—	—	—	—	—	B.fl.II,368	
A. crassiuscula; H. L. Wendland, comm. de Acac. 31, t. 8 (1820)	...	...	W.A.	—	—	—	—	—	—	—	B.fl.II,368	
A. Harveyi; Benth. Fl. Austr. II, 363 (1864)	...	...	W.A.	—	—	—	—	—	—	—	B.fl.II,368	
A. iteaphylla; F. v. M. in Schlecht. Linnaea XXVI, 617 (1853)	...	...	—	S.A.	—	—	—	—	—	—	—	
A. suaveolens; Willdenow, spec. plant. IV, 1050 (1805)	...	...	—	S.A.	T.	V.	N.S.W.	Q.	—	—	B.fl.II,369	
A. subcoerulea; Lindley, Bot. Regist. t. 1075 (1827)	...	...	W.A.	—	—	—	—	—	—	—	B.fl.II,369	
A. Graffiana; F. v. M. in Melb. Chemist, new ser. II, 118 (1887)	...	...	W.A.	—	—	—	—	—	—	—	—	
A. Lindleyi; Meissner in Lehm. pl. Preiss. I, 14 (1844)	...	...	W.A.	—	—	—	—	—	—	—	B.fl.II,370	
A. leptopetala; Benth. in Schlecht. Linnaea XXVI, 619 (1853)	...	...	W.A.	—	—	—	—	—	—	—	B.fl.II,370	
A. Dietrichiana; F. v. M. in Wing's S. Sc. Rec. II, 149 (1882)	...	...	—	—	—	—	—	—	Q.	—	—	
A. Murrayana; F. v. M. in Benth. Fl. Austr. II, 370 (1864)	...	...	—	S.A.	—	—	N.S.W.	Q.	—	—	B.fl.II,370	
A. linifolia; Willdenow, spec. plant. IV, 1051 (1805)	...	...	—	—	—	V.	N.S.W.	Q.	—	—	B.fl.II,371	
A. Leichhardtii; Benth. Fl. Austr. II, 372 (1864)	...	...	—	—	—	—	—	—	Q.	—	B.fl.II,372	
A. buxifolia; Cunningham in Field's N.S.W. 344 (1825)	...	...	—	—	—	—	N.S.W.	Q.	—	—	B.fl.II,372	
A. lunata; Sieber in De Candolle, prodr. II, 452 (1825)	...	...	—	—	—	V.	N.S.W.	Q.	—	—	B.fl.II,373	
A. brachybotrya; Benth. in Hook. Lond. Journ. I, 347 (1842)	...	...	—	S.A.	—	V.	N.S.W.	—	—	—	B.fl.II,373	
A. Wattsiana; F. v. M. in Benth. Fl. Austr. II, 374 (1864)	...	...	—	S.A.	—	—	—	—	—	—	B.fl.II,374	
A. podalyrifolia; Cunningham in Loudon, hort. Brit. 707 (1830)	...	...	—	—	—	—	N.S.W.	Q.	—	—	B.fl.II,374	
A. uncifera; Benth. in Mitch. Trop. Austr. 341 (1848)	...	...	—	—	—	—	—	Q.	—	—	B.fl.II,374	
A. vestita; Edwards in Bot. Regist. t. 698 (1823)	...	...	—	—	—	V.	N.S.W.	—	—	—	B.fl.II,375	
A. cultriformis; Cunningham in G. Don. gen. syst. II, 406 (1832)	...	...	—	—	—	—	N.S.W.	—	—	—	B.fl.II,375	M.fr.XI,123.
A. pravissima; F. v. M., first gen. report 12 (1853)	...	...	—	—	—	V.	N.S.W.	—	—	—	B.fl.II,375	M.fr.I,5,242.
A. stronglylphylla; F. v. M., fragm. VIII, 226 (1874)	...	...	—	S.A.	—	—	—	—	—	—	—	M.fr.VIII,226.
A. pyrifolia; De Candolle, Mém. sur la fam. des Légum. 447 (1825)	...	...	W.A.	—	—	—	—	—	N.A.	B.fl.II,376	M.fr.III,17.	
A. myrtifolia; Willdenow, spec. plant. IV, 1054 (1805)	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	—	B.fl.II,376	
VII. PLURINERVES.												
A. scalpelliformis; Meissner in Lehm. pl. Preiss. II, 200 (1847)	...	...	W.A.	—	—	—	—	—	—	—	B.fl.II,377	
A. urophylla; Benth. in Bot. Reg. Misc. 24 (1841)	...	...	W.A.	—	—	—	—	—	—	—	B.fl.II,377	
A. Luehmanni; F. v. M., fragm. XI, 116 (1881)	...	...	—	—	—	—	—	—	N.A.	—	—	M.fr.XI,116.
A. sublanata; Benth. in Hueg. enum. 42 (1837)	...	...	—	S.A.	—	—	—	—	—	—	B.fl.II,378	M.fr.XI,117.
A. pravifolia; F. v. M., fragm. I, 4 (1858)	...	...	—	S.A.	—	—	N.S.W.	Q.	—	—	—	M.fr.XI,117.
A. amblygona; Cunningham in Hook. Lond. Journ. I, 332 (1842)	...	...	—	—	—	—	N.S.W.	Q.	—	—	B.fl.II,378	M.fr.IV,3; XI,117.

<i>A. amentifera</i> ; F. v. M. in Journ. Linn. Soc. III, 141 (1858)	...	...	...	...	...	...	...	...	...	N.A. B.fl.II,392	
<i>A. Wickhami</i> ; Bentham in Hook. Lond. Journ. I, 379 (1842)	...	...	...	...	...	...	...	...	...	N.A. B.fl.II,392,	
<i>A. lysiphloia</i> ; F. v. M. in Journ. Linn. Soc. III, 137 (1858)	...	...	...	...	S.A.	...	...	...	...	Q. N.A. B.fl.II,393	
<i>A. craspedocarpa</i> ; F. v. M. in Melb. Chem. new ser. II, 73 (1887)	...	...	W.A.	...	...	...	...	...	...	...	...
<i>A. linarioides</i> ; Bentham in Hook. Lond. Journ. I, 371 (1842)	...	...	...	...	...	...	...	...	...	N.A. B.fl.II,393	
<i>A. stipuligera</i> ; F. v. M. in Journ. Linn. Soc. III, 144 (1858)	...	...	...	...	S.A.	...	...	...	...	Q. N.A. B.fl.II,393	
<i>A. ptychophylla</i> ; F. v. M. in Journ. Linn. Soc. III, 142 (1858)	...	...	...	...	...	...	...	...	...	N.A. B.fl.II,394	
<i>A. stigmatophylla</i> ; Cunningham in Hook. Lond. Journ. I, 377 (1842)	...	...	...	...	...	...	...	...	...	N.A. B.fl.II,394	
<i>A. acradenia</i> ; F. v. M. in Journ. Linn. Soc. III (1858)	...	...	...	...	...	...	...	...	...	Q. N.A. B.fl.II,394	
<i>A. leptophleba</i> ; F. v. M. in Journ. Linn. Soc. III, 143 (1858)	...	...	...	...	...	...	...	...	...	N.A. B.fl.II,395	
<i>A. limbata</i> ; F. v. M. in Journ. Linn. Soc. III, 145 (1858)	...	...	...	...	...	...	...	...	...	N.A. B.fl.II,395	
<i>A. brevifolia</i> ; Bentham, Fl. Austr. II, 395 (1864)	...	...	...	...	...	...	...	...	...	Q. N.A. B.fl.II,395	
<i>A. megalantha</i> ; F. v. M. in Journ. Linn. Soc. III, 143 (1858)	...	...	...	...	...	...	...	...	...	N.A. B.fl.II,395	
<i>A. gonoclada</i> ; F. v. M. in Journ. Linn. Soc. III, 140 (1858)	...	...	...	...	...	...	...	...	...	Q. N.A. B.fl.II,396	
<i>A. pycnostachya</i> ; F. v. M., pl. of Vict. II, 33 (1860)	...	...	...	...	...	...	...	N.S.W.	...	B.fl.II,396	
<i>A. subtilinervis</i> ; F. v. M., pl. of Vict. II, 32 (1860)... ..	...	...	...	...	...	...	...	V. N.S.W.	...	B.fl.II,396	M.fr.IV,8.
<i>A. cochliocarpa</i> ; Meissner in Bot. Zeitung 10 (1855)... ..	...	...	...	W.A.	...	...	...	...	...	B.fl.II,397	
<i>A. Dallachiana</i> ; F. v. M., fragm. I, 7 (1858)	...	...	...	...	...	...	...	V.	...	B.fl.II,397	M.fr.I,7.
<i>A. alpina</i> ; F. v. M., fragm. III, 129 (1863)	...	...	...	...	...	...	...	V.	...	B.fl.II,397	M.fr.III,129.
<i>A. longifolia</i> ; Willdenow, spec. pl. IV, 1052 (1805)... ..	...	...	...	...	S.A.	T.	V.	N.S.W.	Q.	B.fl.II,397	
<i>A. linearis</i> ; Sims, Bot. Mag. t. 2156 (1820)	...	...	...	...	...	T.	V.	N.S.W.	...	B.fl.II,399	
<i>A. aciphylla</i> ; Bentham in Schlecht. Linnaea, XXVI, 627 (1853)	...	...	...	W.A.	...	...	...	...	...	B.fl.II,399	
<i>A. ephedroides</i> ; Bentham in Hook. Lond. Journ. I, 370 (1842)	...	...	...	W.A.	...	...	...	...	...	B.fl.II,399	
<i>A. Burkittii</i> ; F. v. M. in Benth. Fl. Austr. II, 400 (1864)... ..	...	...	...	...	S.A.	...	...	N.S.W.	...	B.fl.II,400	
<i>A. microneura</i> ; Meissner in Lehm. pl. Preiss. I, 19 (1844)... ..	...	...	...	W.A.	...	...	...	...	...	B.fl.II,400	
<i>A. cyperophylla</i> ; F. v. M. in Benth. Fl. Austr. II, 400 (1864)	...	...	...	...	S.A.	...	...	N.S.W.	Q.	B.fl.II,400	
<i>A. multispicata</i> ; Bentham, Fl. Austr. II, 400 (1864)	...	...	...	W.A.	...	...	...	...	...	B.fl.II,400	
<i>A. pityoides</i> ; F. v. M. in Journ. Linn. Soc. III, 135 (1858)	...	...	...	...	...	...	...	...	Q. N.A.	B.fl.II,400	
<i>A. xylocarpa</i> ; Cunningham in Hook. Lond. Journ. I, 370 (1842)	...	...	...	...	...	...	...	...	...	N.A. B.fl.II,401	
<i>A. gonocarpa</i> ; F. v. M. in Journ. Linn. Soc. III, 136 (1858)	...	...	...	...	...	...	...	...	...	N.A. B.fl.II,401	
<i>A. oncinophylla</i> ; Lindley, Bot. Regist. XXV, App. XV (1839)	...	...	...	W.A.	...	...	...	...	...	B.fl.II,401	
<i>A. conjunctifolia</i> ; F. v. M., fragm. XI, 68 (1879)	...	...	...	...	...	...	...	...	N.A.	...	M.fr.XI,68.
<i>A. drepanocarpa</i> ; F. v. M. in Journ. Linn. Soc. III, 137 (1858)	...	...	...	...	...	...	...	...	...	N.A. B.fl.II,402	
<i>A. arida</i> ; Bentham in Hook. Lond. Journ. I, 370 (1842)	...	...	...	...	...	...	...	...	...	N.A. B.fl.II,402	
<i>A. aneura</i> ; F. v. M. in Schlecht. Linnaea XXVI, 627 (1853)	...	...	...	W.A.	S.A.	...	V.	N.S.W.	Q.	B.fl.II,402	M.fr.IV,8.
<i>A. cibaria</i> ; F. v. M. in Melb. Chemist, July (1882)	...	...	...	W.A.	S.A.	...	...	N.S.W.	Q.	...	
<i>A. Kempeana</i> ; F. v. M. in Melb. Chemist, July (1882)	...	...	...	...	S.A.	...	...	N.S.W.	Q.	...	
<i>A. conspersa</i> ; F. v. M. in Journ. Linn. Soc. III, 140 (1858)	...	...	...	...	...	...	...	...	...	N.A. B.fl.II,403	
<i>A. Doratoxylon</i> ; Cunningham in Field's N.S.W. 345 (1825)	...	...	...	W.A.	S.A.	...	V.	N.S.W.	Q.	N.A. B.fl.II,403	
<i>A. acuminata</i> ; Bentham in Hook. Lond. Journ. I, 373 (1862)	...	...	...	W.A.	...	...	...	...	...	B.fl.II,404	M.fr.IV,7.
<i>A. stereophylla</i> ; Meissner in Lehm. pl. Preiss. II, 203 (1847)	...	...	...	W.A.	...	...	...	...	...	B.fl.II,404	
<i>A. signata</i> ; F. v. M., fragm. IV, 7 (1864)	...	...	...	W.A.	...	...	...	...	...	B.fl.II,404	M.fr.IV,7.
<i>A. delibrata</i> ; Cunningham in Hook. Lond. Journ. I, 374 (1842)	...	...	...	...	...	...	...	...	Q. N.A.	B.fl.II,404	
<i>A. oligoneura</i> ; F. v. M. in Journ. Linn. Soc. III, 139 (1858)	...	...	...	...	...	...	...	...	...	N.A. B.fl.II,405	
<i>A. torulosa</i> ; Bentham in Journ. Linn. Soc. III, 139 (1858)	...	...	...	...	...	...	...	...	Q. N.A.	B.fl.II,405	
<i>A. julifera</i> ; Bentham in Hook. Lond. Journ. I, 374 (1842)	...	...	...	...	...	...	...	...	Q.	B.fl.II,405	
<i>A. Solandri</i> ; Bentham, Fl. Austr. II, 406 (1864)	...	...	...	...	...	...	...	...	Q.	B.fl.II,406	
<i>A. leptostachya</i> ; Bentham, Fl. Austr. II, 406 (1864)	...	...	...	...	...	...	...	...	Q. N.A.	B.fl.II,406	
<i>A. glaucescens</i> ; Willdenow, spec. plant. IV, 1052 (1805)	...	...	...	...	...	...	V.	N.S.W.	Q.	B.fl.II,406	
<i>A. Cunninghamii</i> ; Hooker, icon. plant. t. 165 (1837)	...	...	...	...	...	...	...	N.S.W.	Q.	B.fl.II,407	

A. leptocarpa ; Cunningham in Hook. Lond. Journ. I, 376 (1842)	...	...	—	—	—	—	Q.	—	B.fl.II,407				
A. polystachya ; Cunningham in Hook. Lond. Journ. I, 376 (1842)	...	...	—	—	—	—	Q.	N.A.	B.fl.II,407				
A. holcocarpa ; Benth. Fl. Austr. II, 408 (1864) ...	...	...	—	—	—	—	Q.	—	B.fl.II,408	M.fr.XI,69.			
A. plectocarpa ; Cunningham in Hook. Lond. Journ. I, 375 (1842)	...	...	—	—	—	—	Q.	N.A.	B.fl.II,408				
A. pachycarpa ; F. v. M. in Journ. Linn. Soc. III, 139 (1858) ...	...	...	—	—	—	—	—	N.A.	B.fl.II,408				
A. tumida ; F. v. M. in Journ. Linn. Soc. III, 144 (1858) ...	...	...	—	—	—	—	—	N.A.	B.fl.II,409				
A. loxocarpa ; Benth. in Hook. Lond. Journ. I, 377 (1842) ...	...	...	—	—	—	—	—	N.A.	B.fl.II,409				
A. oncinocarpa ; Benth. in Hook. Lond. Journ. I, 378 (1842) ...	...	...	—	—	—	—	—	N.A.	B.fl.II,409				
A. retinervis ; Benth. in Hook. Lond. Journ. I, 379 (1842) ...	...	...	—	—	—	—	—	N.A.	B.fl.II,410				
A. aulacocarpa ; Cunningham in Hook. Lond. Journ. I, 378 (1842)	...	...	—	—	—	—	Q.	N.A.	B.fl.II,410	M.fr.XI,69.			
A. calyculata ; Cunningham in Hook. Lond. Journ. I, 379 (1842)...	...	...	—	—	—	—	Q.	—	B.fl.II,410				
A. auriculiformis ; Cunningham in Hook. Lond. Journ. I, 377 (1842)	...	...	—	—	—	—	Q.	N.A.	B.fl.II,411				
A. latifolia ; Benth. in Hook. Lond. Journ. I, 382 (1842) ...	...	...	—	—	—	—	Q.	N.A.	B.fl.II,411				
A. Mangium ; Willdenow, spec. plant. IV, 1053 (1805) ...	...	...	—	—	—	—	Q.	—	—	M.fr.XI,36	...	As	
A. holosericea ; Cunningham in G. Don, gen. syst. II, 407 (1832)...	...	...	—	—	—	—	Q.	N.A.	B.fl.II,411	...	...	P.	
A. cincinnata ; F. v. M., fragm. XI, 35 (1878) ...	...	...	—	—	—	—	Q.	—	—	M.fr.XI,35.			
A. dimidiata ; Benth. in Hook. Lond. Journ. I, 381 (1842) ...	...	...	—	—	—	—	—	N.A.	B.fl.II,412				
A. humifusa ; A. Cunningham in Lond. Journ. I, 382 (1842) ...	...	...	—	—	—	—	Q.	N.A.	B.fl.II,412				
A. denticulosa ; F. v. M., fragm. X, 32 (1876) ...	...	...	W.A.	—	—	—	—	—	—	M.fr.X,32.			

IX. BIPINNATAE.

A. Drummondii ; Lindley, Bot. Regist. XXV, App. XV (1839) ...	...	...	W.A.	—	—	—	—	—	B.fl.II,419				
A. pulchella ; R. Brown in Ait. hort. Kew. sec. ed. V, 464 (1813) ...	...	...	W.A.	—	—	—	—	—	B.fl.II,416				
A. Mitchellii ; Benth. in Hook. Lond. Journ. I, 387 (1842) ...	...	...	—	S.A.	—	V.	—	—	B.fl.II,417				
A. pentadenia ; Lindley, Bot. Regist. t. 1521 (1832)...	...	...	W.A.	—	—	—	—	—	B.fl.II,417				
A. Gilbertii ; Meissner in Lehm. pl. Preiss. II, 204 (1847) ...	...	...	W.A.	—	—	—	—	—	B.fl.II,417				
A. nigricans ; R. Brown in Ait. hort. Kew. sec. ed. V, 465 (1813) ...	...	...	W.A.	—	—	—	—	—	B.fl.II,418				
A. Tayloriana ; F. v. M. in Wing's S. Sc. Record, July (1882) ...	...	...	W.A.	—	—	—	—	—	—				
A. strigosa ; Link, enum. pl. hort. Berol. 444 (1822) ...	...	...	W.A.	—	—	—	—	—	B.fl.II,418				
A. elata ; Cunningham in Hook. Lond. Journ. I, 383 (1842) ...	...	...	—	—	—	—	N.S.W.	—	B.fl.II,413				
A. pruinosa ; Cunningham in Hook. Lond. Journ. I, 383 (1842) ...	...	...	—	—	—	—	N.S.W.	—	B.fl.II,413				
A. spectabilis ; Cunningham in Hook. Lond. Journ. I, 383 (1842)...	...	...	—	—	—	—	N.S.W.	Q.	B.fl.II,413				
A. Baileyana ; F. v. M. in Transact. Roy. Soc. of Vict. XXIV, 168 (1887)	...	...	—	—	—	—	N.S.W.	—	—				
A. polybotrya ; Benth. in Hook. Lond. Journ. I, 384 (1842) ...	...	...	—	—	—	—	N.S.W.	Q.	B.fl.II,414				
A. discolor ; Willdenow, spec. plant. IV, 1068 (1805) ...	...	...	—	—	T.	V.	N.S.W.	—	B.fl.II,414				
A. decurrens ; Willdenow, spec. plant. IV, 1872 (1805) ...	...	...	—	—	—	V.	N.S.W.	Q.	B.fl.II,414				
A. mollissima ; Willdenow, enum. pl. hort. Berol. 1053 (1809) ...	...	...	—	S.A.	T.	V.	N.S.W.	—	—				
A. dealbata ; Link, enum. pl. hort. Berol. 445 (1822) ...	...	...	—	—	—	V.	N.S.W.	Q.	B.fl.II,415				
A. cardiophylla ; Cunningham in Hook. Lond. Journ. I, 385 (1842) ...	...	...	—	—	—	—	N.S.W.	—	B.fl.II,415				
A. leptoclada ; Cunningham in Hook. Lond. Journ. I, 385 (1842) ...	...	...	—	—	—	—	N.S.W.	Q.	B.fl.II,416				
A. pubescens ; R. Brown in Ait. hort. Kew. sec. ed. V, 464 (1813) ...	...	...	—	—	—	—	N.S.W.	—	B.fl.II,416				
A. Farnesiana ; Willdenow, spec. plant. IV, 1083 (1805) ...	...	...	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.II,419	...	E., As., Afr., Am., P.	
A. suberosa ; Cunningham in Hook. Lond. Journ. I, 499 (1842) ...	...	...	—	—	—	—	—	—	N.A.	B.fl.II,420	M.fr.XI,36.		
A. Bidwilli ; Benth. in Schlecht. Linnaea XXVI, 629 (1853) ...	...	...	—	—	—	—	—	—	Q.	N.A.	B.fl.II,420		
A. pallida ; F. v. M. in Journ. Linn. Soc. III, 14 (1858) ...	...	...	—	—	—	—	—	—	—	N.A.	B.fl.II,421		
A. Sutherlandii ; F. v. M., inconogr. of Austr. Acac. twelfth decade (1888)	...	...	—	—	—	—	—	—	Q.	—	M.fr.VI,22; IX,179.		

ALBIZZIA ; Durazzini in Magazzino Toscano III, part IV, 13 (1772). (Zygia 1756, Pithecolobium, Cathormion, Calliandra, Enterolobium, Serianthes, Archidendron.)

A. lophantha ; Benth. in Hook. Lond. Journ. III, 86 (1844) ...	...	...	W.A.	—	—	—	—	—	B.fl.II,421				
A. basaltica ; Benth. Fl. Austr. II, 422 (1864) ...	...	...	—	—	—	—	—	—	Q.	—	B.fl.II,422		
A. Thozetiana ; F. v. M. in Benth. Fl. Austr. II, 422 (1864) ...	...	...	—	—	—	—	—	—	Q.	—	B.fl.II,422	M.fr.IV,9&176.	

A. procera; Benth. in Hook. Journ. III, 88 (1844)	...	...	...	—	—	—	—	—	—	N.A.	B.fl.II,422	...	...	As.	
A. pruinosa; F. v. M. in Trimen's Journ. of Bot. 9 (1872)...	...	...	...	—	—	—	—	—	N.S.W.	Q.	—	B.fl.II,423	M.fr.IX,179	...	As.
A. Lebbek; Durazzini in Magaz. Tosc. III, pt. IV, 13 (1772)	...	...	...	—	—	—	—	—	—	Q.	—	—	...	...	As., Afr.
A. canescens; Benth. Fl. Austr. II, 423 (1864)	...	...	...	—	—	—	—	—	—	Q.	—	B.fl.II,423	M.fr.IX,179.		
A. Tozeri; F. v. M. in Trimen's Journ. of Bot. 10 (1872)	...	...	...	—	—	—	—	—	N.S.W.	Q.	—	B.fl.II,424	M.fr.V,106; IX,179.		
A. Hendersoni; F. v. M. in Trimen's Journ. of Bot. 10 (1872)	...	...	...	—	—	—	—	—	N.S.W.	Q.	—	—	M.fr.V,191; VI,202; IX,178.		
A. monilifera; F. v. M. in Trimen's Journ. of Bot. 10 (1872)	...	...	...	—	—	—	—	—	—	N.A.	B.fl.II,424	M.fr.V,191	...	..	As.
A. ramiflora; F. v. M., fragm. IX, 178 (1875)	...	...	...	—	—	—	—	—	—	Q.	—	—	M.fr.V,10; IX,178.		
A. Vaillantii; F. v. M., fragm. V, 9 (1865)	...	...	...	—	—	—	—	—	—	Q.	—	—	M.fr.V,60; IX,178.		
A. Lucyi; F. v. M., iconogr. of Austr. Acai. 13, Dec. 6 (1888)	...	...	...	—	—	—	—	—	—	Q.	—	—	M.fr.V,201; IX,178.		

CONNARACEAE.

R. BROWN, Narr. Exped. Cong. App. V, 431 (1818).—Genera 2, Species 2.

ROUREA; Aublet, Hist. des pl. de la Guian. I, 467, t. 187 (1775).

R. brachyandra; F. v. M., fragm. VIII, 6 (1872) ... — — — — — Q. — — M.fr.X,119.

TRICHOLOBUS; Blume, Mus. Bot. Lugd. I, 237 (1850).

T. connaroides; F. v. M., fragm. VIII, 224 (1874) .. — — — — — Q. — — M.fr.V,105; X,120.

ROSACEAE.

A. L. DE JUSSIEU, Gen. pl. 334 (1789), from B. de Jussieu (1759).—Genera 8, Species 17.

PARINARIUM; A. L. de Jussieu, gen. pl. 342 (1789).

P. Nonda; F. v. M. in Benth. Fl. Austr. II, 426 (1864) ... — — — — — Q. N.A. B.fl.II,426
P. Griffithianum; Benth. in Hook. Niger-Flora 334 (1849) ... — — — — — Q. N.A. B.fl.II,426 M.fr.VIII,34 ... As., P.

STYLOBASIUM; Desfontaines in Mém. du Mus. V, 37, t. 2 (1819). (Macrostigma.)

S. spatulatum; Desfontaines in Mém. du Mus. V, 37, t. 2 (1819) ... W.A. S.A. — — — — — N.A. B.fl.II,427
S. lineare; Nees in Lehm. pl. Preiss. I, 95 (1844) ... W.A. — — — — — — — B.fl.II,427

GEUM; Linné, gen. plant. 148 (1737) from Plinius. (Sieversia.)

G. urbanum; Linné, spec. plant. 501 (1753) ... — S.A. T. V. N.S.W. — — B.fl.II,428 ... E., As.
G. renifolium; F. v. M. in Transact. phil. Inst. Vict. II, 66 (1857) ... — — T. — — — — B.fl.II,428

POTENTILLA; Linné, gen. plant. 147 (1737) from Camerarius (1586).

P. anserina; Linné, spec. plant. 495 (1753) ... — S.A. T. V. — — — — B.fl.II,429 ... E., As., Af., Am., N.Z.

RUBUS; Tournefort, Inst. 614, t. 385 (1700) from Plinius.

R. Gunnianus; Hooker, icon. plant. t. 291 (1840) ... — — T. — — — — B.fl.II,430 M.fr.IV,32
R. parvifolius; Linné, spec. plant. 1197 (1753) ... — S.A. T. V. N.S.W. Q. — — B.fl.II,430 M.fr.IV,30 ... As.
R. rosifolius; Smith, plantar. icon. ex herb. Linn. t. 60 (1790) ... — — — — V. N.S.W. Q. — — B.fl.II,431 M.fr.IV,32; VIII,34. As., Afr.
R. Moluccanus; Linné, spec. plant. 1197 (1753) ... — — — — V. N.S.W. Q. N.A. B.fl.II,430 M.fr.IV,31; VIII,34. As., P.
R. Moorei; F. v. M. in Transact. phil. Inst. Vict. II, 67 (1857) ... — — — — N.S.W. Q. — — B.fl.II,431 M.fr.IV,29&176; VIII,34.

AGRIMONIA; Tournefort, inst. 301, t. 155 (1700) from C. Bauhin (1623).

A. Eupatoria; Linné, spec. plant. 448 (1753) ... — — — — N.S.W. — — — — M.fr.VIII,34 ... As., Afr., Am.

ALCHEMILLA; Linné, gen. pl. 30 (1737) from C. Bauhin (1623). (Alchimilla.)

A. vulgaris; C. Bauhin, pinax 319 (1623) ... — — — — V. N.S.W. — — — — B.fl.II,432 ... E., As., Am.

ACAENA ; Mutis in Linné, mantiss. II, 200 (1771). (Ancistrum.)

A. ovina ; Cunningham in Field's N. S. Wales, 358 (1825)...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,433	...	...	...	Am., N.Z.
A. sanguisorbae ; Vahl, enum. plant. I, 294 (1804) ...	...	...	...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,434	...	...	...	N.Z.
A. montana ; J. Hooker in Hook. Lond. Journ. VI, 276 (1847) ...	...	...	...	—	—	T.	—	—	—	—	B.fl.II,434	...	...	...	

SAXIFRAGEAE.

VENTENAT, Tabl. III, 277 (1799).—Genera 22, Species 36.

ARGOPHYLLUM ; R. et G. Forster, charact. gen. 29, t. 15 (1776).

A. Lejournandii ; F. v. M., fragm. IV, 33 (1864) ...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,437	M.fr.IV,177 ; VI,188.
--	-----	-----	-----	---	---	---	---	--------	----	---	-------------	-----------------------

ABROPHYLLUM ; J. Hooker in Benth. Fl. Aust. II, 437 (1864). (Brachynema.)

A. ornans ; J. Hooker in Benth. Fl. Austr. II, 437 (1864) ...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,437	M.fr.III,90 ; VI,189 ; VII,150.
---	-----	-----	-----	---	---	---	---	--------	----	---	-------------	---------------------------------

CUTTSIA ; F. v. M., fragm. V, 47 (1865).

C. viburnea ; F. v. M., fragm. V, 47 (1865) ...	...	...	...	...	—	—	—	N.S.W.	Q.	—	—	M.fr.V,47&189 ; VI,189.
---	-----	-----	-----	-----	---	---	---	--------	----	---	---	-------------------------

COLMEIROA ; F. v. M., fragm. VII, 149 (1871).

C. carpodetoides ; F. v. M., fragm. VII, 149 (1871) ...	...	...	...	...	—	—	—	N.S.W.	—	—	—	M.fr.VII,149.
---	-----	-----	-----	-----	---	---	---	--------	---	---	---	---------------

QUINTINIA ; A. de Candolle, Mon. Campan. 92 (1830).

Q. Sieberi ; A. De Candolle, Monogr. Camp. 92 (1830) ...	...	...	...	...	—	—	—	N.S.W.	Q.	—	B.fl.II,438	M.fr.II,126 ; III,165 ; VI,189.
Q. Verdonii ; F. v. M., fragm. II, 125 (1861)...	...	...	...	...	—	—	—	N.S.W.	Q.	—	B.fl.II,438	M.fr.VI,189.
Q. Fawkneri ; F. v. M., fragm. VI, 92 (1867)...	...	...	...	...	—	—	—	—	Q.	—	—	M.fr.VII,150.

POLYOSMA ; Blume, Bijdr. 658 (1825).

P. Cunninghamii ; J. J. Bennett, pl. Jav. rar. 196 (1840) ...	...	...	...	...	—	—	—	N.S.W.	Q.	—	B.fl.II,438	M.fr.VI,189.
P. alangiacea ; F. v. M., fragm. VIII, 8 (1872) ...	...	...	...	...	—	—	—	—	Q.	—	—	M.fr.VIII,8.

ANOPTERUS ; Labillardière, Nov. Holl. pl. spec. I, 86, t. 112 (1804).

A. glandulosus ; Labillardière, Nov. Holl. pl. spec. I, 86, t. 112 (1804) ...	...	...	...	...	—	—	T.	—	—	—	B.fl.II,439	M.fr.II,126.
A. Macleayanus ; F. v. M. in Journ. Pharm. Soc. Vict. 43 (1858)...	...	...	...	...	—	—	—	N.S.W.	Q.	—	B.fl.II,439	M.fr.VI,188 ; VII,150.

CALLICOMA ; Andrews, Bot. Reposit. t. 566 (1809). (Calycomis partly.)

C. serratifolia ; Andrews, Bot. Reposit. t. 566 (1809) ...	...	...	...	...	—	—	—	N.S.W.	Q.	—	B.fl.II,440	M.fr.V,32.
C. Stutzeri ; F. v. M., fragm. V, 34 (1865) ...	...	...	...	...	...	—	—	—	—	Q.	—	M.fr.VI,188&252 ; VII,150.

ANODOPETALUM ; Cunningham in Endl. gen. 818 (1839).

A. biglandulosum ; Cunningham in J. Hook. Fl. Tasm. I, 148 (1860) ...	...	...	...	...	—	—	T.	—	—	—	B.fl.II,440	M.fr.VI,189.
---	-----	-----	-----	-----	---	---	----	---	---	---	-------------	--------------

APHANOPETALUM ; Endlicher, nov. stirp. decad. I, 34 (1839). (Platyptelea.)

A. resinosum ; Endlicher, nov. stirp. dec. I, 35 (1839) ...	...	...	...	...	—	—	—	V.	N.S.W.	Q.	—	B.fl.II,441	M.fr.I,228.
A. occidentale ; F. v. M. fragm. I, 228 (1858) ...	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,441	M.fr.XI,20.

CERATOPETALUM ; Smith, specim. of the Bot. of New Holland, 9 (1793).

C. gummiferum ; Smith, specim. of the Bot. of New Holl. t. 3 (1793) ...	...	...	...	...	—	—	—	—	N.S.W.	—	B.fl.II,442	M.fr.II,126.
C. apetalum ; D. Don in Edinb. new phil. journ. (1830) ...	...	...	...	...	...	—	—	—	N.S.W.	—	B.fl.II,442	M.fr.II,126 ; VI,189 ; VII,150.

SCHIZOMERIA ; D. Don in Edinb. new phil. journ. IX, 94 (1830).

S. ovata ; D. Don in Edinb. new phil. journ. IX, 94 (1830)...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,443	M.fr.VI,189.
---	-----	-----	-----	-----	---	---	---	---	--------	----	---	-------------	--------------

ACROPHYLLUM ; Bentham in Maund & Henslow, Botanist II, 95 (1840). (Calycomis partly.)

A. venosum ; Bentham in Maund & Henslow, Bot. II, 95 (1840) ...	...	...	...	...	—	—	—	—	N.S.W.	—	B.fl.II,443	M.fr.VI,189 ; VII,150.
---	-----	-----	-----	-----	---	---	---	---	--------	---	-------------	------------------------

WEINMANNIA ; Linné, syst. ed. X, 1005 (1759). (Ackama, Geissois.)

W. paniculosa ; F. v. M., fragm. II, 126 (1861) ...	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,444	M.fr.II,83,175 ; VI,188 ; VII,150.
---	-----	-----	-----	-----	-----	---	---	---	---	--------	----	---	-------------	------------------------------------

W. Biagiana; F. v. M., fragm. V, 16 (1865) ...	—	—	—	—	—	Q.	—	—	M.fr. VII, 150.
W. lachnocarpa; F. v. M., fragm. VIII, 7 (1782) ...	—	—	—	—	N.S.W.	Q.	—	—	M.fr. VIII, 7, 281.
W. rubifolia; Benthams, Fl. Aust. II, 445 (1864) ...	—	—	—	—	N.S.W.	Q.	—	B.fl. II, 445	M.fr. II, 82 & 175; V, 180; VI, 181.
W. Benthams; F. v. M., Docum. intercol. Exhibit. 31 (1866) ...	—	—	—	—	N.S.W.	Q.	—	B.fl. II, 446	M.fr. V, 180; VI, 188.

SPIRAEANTHEMUM; Asa Gray in proceed. Amer. Acad. of Science III (1853).

S. Davidsoni; F. v. M. in Austr. Journ. of Pharm. new ser. II, 85 (1887) ...	—	—	—	—	—	Q.	—	—	
--	---	---	---	---	---	----	---	---	--

GILLBEEA; F. v. M., fragm. V, 17 (1865).

G. adenopetala; F. v. M., fragm. V, 17 (1865) ...	—	—	—	—	—	Q.	—	—	M.fr. VI, 188.
---	---	---	---	---	---	----	---	---	----------------

DAVIDSONIA; F. v. M., fragm. VI, 4 (1867).

D. pruriens; F. v. M., fragm. VI, 4 (1867) ...	—	—	—	—	N.S.W.	Q.	—	—	M.fr. VI, 249.
--	---	---	---	---	--------	----	---	---	----------------

TETRACARPAEA; J. Hooker, icon. plant. t. 264 (1840).

T. Tasmanica; J. Hooker, icon. plant. t. 264 (1840) ...	—	—	T.	—	—	—	—	B.fl. II, 445	
---	---	---	----	---	---	---	---	---------------	--

EUCRYPTIA; Cavanilles, icon. IV, 48, t. 372 (1797). (Carpodontos.)

E. Billardieri; Spach, veg. phan. V, 344 (1836) ...	—	—	T.	—	—	—	—	B.fl. II, 446	M.fr. IV, 2; VI, 189; VII, 150.
E. Moorei; F. v. M., fragm. IV, 2 (1864) ...	—	—	—	—	V.	N.S.W.	—	B.fl. II, 447	M.fr. IV, 176; VI, 189; VII, 150.

BAUERA; Banks & Kennedy in Andrews, Bot. Rep. t. 198 (1793).

B. rubioides; Andrews, Bot. Reposit. t. 198 (1793) ...	—	—	S.A. T.	V.	N.S.W.	Q.	—	B.fl. II, 447	M.fr. IV, 23.
B. capitata; Séringe in De Candolle, prodr. IV, 13 (1830) ...	—	—	—	—	N.S.W.	Q.	—	B.fl. II, 448	M.fr. IV, 24.
B. sessiliflora; F. v. M. in Transact. phil. Soc. Vict. I, 41 (1854) ...	—	—	—	—	V.	—	—	B.fl. II, 448	M.fr. IV, 24; VII, 150.

EREMOSYNE; Endlicher in Hueg. enum. pl. Nov. Holl. Austr. occ. 53 (1837).

E. pectinata; Endlicher in Hueg. enum. pl. 53 (1837) ...	—	—	—	—	—	—	—	B.fl. II, 449	M.fr. VI, 189.
--	---	---	---	---	---	---	---	---------------	----------------

CEPHALOTUS; Labillardière, Nov. Holl. pl. spec. II, 7, t. 145 (1806).

C. follicularis; Labillardière, Nov. Holl. pl. spec. II, 7, t. 145 (1806) ...	—	—	—	—	—	—	—	B.fl. II, 449	M.fr. VI, 189.
---	---	---	---	---	---	---	---	---------------	----------------

NEPENTHACEAE.

REICHENBACH, Conspectus regni vegetabilis 45 (1828).—Genus 1, Species 2.

NEPENTHES; Linné, syst. nat. 9 (1735); Linné, gen. pl. 273 (1737).

N. Kennedyana; F. v. M., fragm. V, 154 (1866) ...	—	—	—	—	—	Q.	—	B.fl. VI, 40	M.fr. V, 154.
N. phyllamphora; Willdenow, sp. plant. IV, 874 (1805) ...	—	—	—	—	—	Q.	—	...	... As., P.

ARISTOLOCHIEAE.

A. L. DE JUSSIEU in Ann. du Mus. V, 221 (1804).—Genus 1, Species 5.

ARISTOLOCHIA; Tournefort, inst. 162, t. 71 (1700), from Hippocrates, Theophrastos & Dioscorides.

A. deltantha; F. v. M., fragm. VI, 179 (1869) ...	—	—	—	—	—	Q.	—	B.fl. VI, 208	M.fr. VI, 179 & 255.
A. praevenosa; F. v. M., fragm. II, 166 (1861) ...	—	—	—	—	N.S.W.	—	—	B.fl. VI, 208	M.fr. VI, 180.
A. pubera; R. Brown, prodr. 349 (1810) ...	—	—	—	—	N.S.W.	Q.	—	B.fl. VI, 208	
A. Thozetii; F. v. M., fragm. II, 167 (1861) ...	—	—	—	—	—	Q.	—	B.fl. VI, 208	M.fr. II, 167.
A. Indica; Linné, spec. plant. 960 (1753) ...	—	—	—	—	—	Q.	—	B.fl. VI, 209	... As., P.

CRASSULACEAE.

DE CANDOLLE, Bull. de la Soc. Philom. n. 49 (1801).—Genus 1, Species 6.

TILLAEA; Micheli, nov. gen. 22, t. 20 (1729). (Bulliardia, Crassulae subgenus.)

T. verticillaris; De Candolle, prodr. III, 382 (1828) ...	—	—	—	—	W.A. S.A. T.	V.	N.S.W.	Q.	—	B.fl. II, 451	M.fr. XI, 118	... Am., N.Z.
T. purpurata; J. Hooker in Lond. Journ. of Bot. VI, 472 (1847) ...	—	—	—	—	W.A. S.A. T.	V.	N.S.W.	—	—	B.fl. II, 451	M.fr. XI, 118.	

T. macrantha; J. Hooker in Hook. icon. plant. t. 310 (1841)	...	...	...	...	S.A. T. V. N.S.W.	—	—	B.fl.II,452	M.fr.XI,117.
T. pedicellosa; F. v. M., fragm. XI, 118 (1881)	...	...	...	...	W.A.	—	—	—	M.fr.XI,118.
T. intricata; Nees in Lehm. pl. Preiss. I, 278 (1844)	...	...	...	...	W.A.	—	—	—	M.fr.XI,117.
T. recurva; J. Hooker, Fl. Tasm. I, 146 (1860)	...	...	...	...	W.A. S.A. T. V. N.S.W.	Q.	—	B.fl.II,452	M.fr.XI,118.

HAMAMELIDEAE.

R. BROWN in Abel, Narr. journ. Chin. 374 (1818).

Genus and species undertermined	...	...	...	...	...	...	—	—	—	—	Q.	—	—
---------------------------------	-----	-----	-----	-----	-----	-----	---	---	---	---	----	---	---

ONAGREAE.

ADANSON, Familles des plantes II, 81 (1763), from B. de Jussieu (1759).—Genera 4, Species 5.

OENOTHERA; Linné, syst. nat. 8 (1735); Linné, gen. plant. 112 (1737).

O. Tasmanica; J. Hooker, Fl. Tasm. I, 119 (1860)	...	...	...	...	...	—	—	T.	—	—	—	—	B.fl.III,303
--	-----	-----	-----	-----	-----	---	---	----	---	---	---	---	--------------

EPILOBIUM; Dillenius in Linné, syst. nat. 8 (1735), from Gesner (1542).

E. glabellum; G. Forster, florul. ins. Austr. prodr. 27 (1786)	...	...	...	...	W.A. S.A. T. V. N.S.W.	Q.	—	B.fl.III,305	...	...	...	N.Z.
--	-----	-----	-----	-----	------------------------	----	---	--------------	-----	-----	-----	------

JUSSIEUA; Linné, gen. plant. 126 (1737). (Jussiaea.)

J. diffusa; Forskael, fl. Aeg. Arab. 210 (1775)	...	...	...	...	...	—	S.A.	—	V. N.S.W.	Q.	—	B.fl.III,306	...	...	...	As., Afr.
J. suffruticosa; Linné, spec. plant. 388 (1753)	...	...	...	...	...	—	—	—	N.S.W.	Q. N.A.	B.fl.III,307	M.fr.III,13	...	...	...	As., Afr., Am., P.

LUDWIGIA; Linné, coroll. 3 (1737).

L. parviflora; Roxburgh, Fl. Ind. edit. Carey. I, 440 (1820)	...	...	...	...	...	—	—	—	—	—	Q. N.A.	B.fl.III,307	M.fr.III,129; IV,175	...	...	As., Afr.
--	-----	-----	-----	-----	-----	---	---	---	---	---	---------	--------------	----------------------	-----	-----	-----------

SALICARIEAE.

ADANSON, Familles des plantes II, 232 (1763), from B. de Jussieu (1759).—Genera 7, Species 19.

ROOTALA; Linné, mantissa altera 143 (1771).

R. verticillaris; Linné, mantissa altera 143 (1771)	...	...	...	...	...	—	—	—	—	—	N.A.	B.fl.III,295	M.fr.III,108	...	...	As.
R. Roxburghiana; Wight, Icon. t. 260 (1840)	...	...	...	...	...	—	—	—	—	—	Q. N.A.	B.fl.III,296	M.fr.XI,118	...	...	As.
R. Mexicana; Chamisso & Schlechtendal in Linnaea V, 567 (1830)	...	...	...	...	...	—	—	—	—	—	Q. N.A.	—	...	...	...	As., Am.
R. occultiflora; Koehne in Engler's Jahrb. I, 152 (1880)	...	...	...	...	...	—	—	—	—	—	Q. N.A.	—	...	...	...	As.
R. diandra; F. v. M. in Transact. Roy. Soc. of N.S.W. 89 (1880)	...	...	...	...	...	—	S.A.	—	—	—	Q. N.A.	B.fl.III,296	M.fr.III,108,168.	...	...	As.

AMMANNIA; Houston in Linné, gen. plant. 337 (1737). (Koehnea, Crinopodium.)

A. crinipes; F. v. M. in Transact. phil. Soc. Vict. III, 49 (1858)	...	...	...	...	...	—	—	—	—	—	N.A.	B.fl.III,296	...	...	...	As.
A. triflora; R. Brown in Benth. Fl. Austr. III, 297 (1866)	...	...	...	...	...	—	—	—	—	—	Q. N.A.	B.fl.III,297	...	...	...	As.
A. baccifera; Linné, spec. plant. ed. sec. 175 (1762)	...	...	...	...	...	—	S.A.	—	—	—	Q. N.A.	B.fl.III,297	...	...	...	As., Afr.
A. auriculata; Willdenow, hort. Berol. t. 7 (1806)	...	...	...	...	...	—	—	—	—	—	Q.	B.fl.III,297	...	...	...	As., Afr.
A. multiflora; Roxburgh, Fl. Ind. edit. Carey. I, 447 (1820)	...	...	...	...	...	W.A. S.A.	—	V. N.S.W.	Q. N.A.	B.fl.III,298	...	...	...	...	...	As., Afr.

LAGERSTROEMIA; Linné, syst. ed. X, 1076 (1759).

L. Indica; Linné, spec. plant. ed. sec. 734 (1762)	...	...	...	...	...	—	—	—	—	—	Q.	—	—	M.fr.VIII,35	...	As.
L. Flos Reginae; Retzius, observ. botan. V, 25 (1789)	...	...	...	...	...	—	—	—	—	—	Q.	—	—	M.fr.XII,20	...	As.
L. Archeriana; Bailey, Synop. of Queensland Flora 196 (1883)	...	...	...	...	...	—	—	—	—	—	Q.	—	—	...	...	As.

LYTHRUM; Linné, syst. nat. 8 (1735); Linné, gen. plant. 136 (1737).

L. Salicaria; Linné, spec. plant. 446 (1753)	...	...	...	...	...	—	S.A. T. V. N.S.W.	Q.	—	B.fl.III,298	M.fr.III,110; VII,147	E., As., Afr., Am.
L. Hyssopifolia; Linné, spec. plant. 447 (1753)	...	...	...	...	...	—	S.A. T. V. N.S.W.	Q.	—	B.fl.III,299	M.fr.III,109; VII,147	E., As., Afr., Am.

NESAEA; Commerçon in A. L. de Jussieu, gen. plant. 332 (1789). (Decadon.)

N. Robertsii; F. v. M., fragm. VII, 145 (1871)	...	...	...	...	...	—	—	—	—	—	Q.	—	—	M.fr. VII, 145.
N. Arnhemica; F. v. M., fragm. VII, 146 (1871)	..	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl. III, 299	M.fr. III, 109, 168.

PEMPHIS ; R. et G. Forster, charact. gen. 67, t. 34 (1776).

P. acidula; R. & G. Forster, charact. gen. 67, t. 34 (1776)... .. — — — — — Q. N.A. B.f.III,300 M.fr.III,109; VII,147 As., Afr., P.

LAWSONIA; Linné, gen. plant. 111 (1737).

L. alba ; Lamarck, Encycl. méth. III, 106 (1789)	...	...	...	...	...	—	—	—	—	—	—	N. A. B.f.III,301	...	...	...	As., Afr.
--	-----	-----	-----	-----	-----	---	---	---	---	---	---	-------------------	-----	-----	-----	-----------

HALORAGEAE.

R. BROWN in Flind. Voy. II, 549 (1814).—Genera 4, Species 58.

GUNNERA; Linné, mantiss. plant. 16 (1767). (Milligania.)

<i>G. cordifolia</i> ; J. Hooker, Fl. Tas. I, 125 (1860)	...	...	...	...	...	—	—	T.	—	—	—	—	B.fl.II, 491
--	-----	-----	-----	-----	-----	---	---	----	---	---	---	---	--------------

LOUDONIA; Lindley, Bot. Regist. XXV, App. XLII (1839). (Glischrocaryon.)

L. aurea; Lindley, Bot. Regist. XXV, App. XLII (1839) ...	...	...	...	...	W.A.	S.A.	—	—	—	—	—	B.f. II, 472 M.fr. VIII, 162; XI, 20.
L. Behrii; Schlechtendal, Linnaea XX, 648 (1847) ...	...	...	...	...	—	S.A.	—	V.	N.S.W.	—	—	B.f. II, 472 M.fr. VIII, 162.
L. Roei; Schlechtendal, Linnaea XX, 648 (1847) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.f. II, 472 M.fr. VIII, 162.

HALORAGIS; R. et G. Forster, charact. gen. 61, t. 31 (1776). (Cercodia, Goniocarpus, Meionectes.)

H. Gossei ; F. v. M., fragm. VIII, 161 (1874)	...	...	...	...	...	W.A. S.A.	—	—	—	Q. N.A.	—	M.fr.XI,134.
H. trigonocarpa ; F. v. M., fragm. X, 84 (1876)	...	...	...	...	...	W.A. S.A.	—	—	—	—	—	M.fr.X,84.
H. digyna ; Labillardière, Nov. Holl. pl. spec. I, 101, t. 129 (1804)	...	...	...	...	...	W.A. S.A.	—	—	—	—	B.fl.II,475	
H. mucronata ; Bentham, Fl. Austr. II, 475 (1864)	...	...	...	...	...	W.A. S.A.	—	V.	N.S.W.	—	B.fl.II,475	
H. pityoides ; Bentham, Fl. Austr. II, 476 (1864)	...	...	...	...	...	W.A.	—	—	—	—	B.fl.II,476	
H. cordigera ; Fenzl in Hueg. enum. 45 (1837)	...	...	...	...	...	W.A.	—	—	—	—	B.fl.II,476	
H. elata ; Cunningham in Hueg. enum. 45 (1837)	...	...	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	B.fl.II,476 M.fr.VIII,162.
H. tenuifolia ; Bentham, Fl. Austr. II, 477 (1864)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,477
H. scoparia ; Fenzl in Hueg. enum. 45 (1837)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,477 M.fr.X,54.
H. aculeolata ; Bentham, Fl. Austr. II, 477 (1864)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,477 M.fr.X,54.
H. foliosa ; Bentham, Fl. Austr. II, 477 (1864)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,477
H. pycnostachya ; F. v. M. in Transact. R. S. Vict. XXIV, p. 135 (1887)	...	...	...	...	...	W.A.	—	—	—	—	—	
H. confertifolia ; F. v. M., fragm. X, 53 (1876)	...	...	...	...	...	W.A.	—	—	—	—	—	M.fr.X,54.
H. platycarpa ; Bentham, Fl. Austr. II, 478 (1864)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,478
H. acutangula ; F. v. M. in Transact. Vict. Inst. 125 (1854)	...	...	...	...	...	—	S.A.	—	—	—	—	B.fl.II,478
H. hexandra ; F. v. M., fragm. III, 31 (1862)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,478 M.fr.III,31.
H. odontocarpa ; F. v. M., fragm. I, 108 (1859)	...	...	...	...	...	—	S.A.	—	V.	N.S.W.	—	B.fl.II,479 M.fr.I,108; X,54.
H. serra ; Brongniart in Duperr. Voy. sur la Coquille, t. 69 (1829)	...	...	...	...	...	—	—	—	—	N.S.W.	—	B.fl.II,479
H. aspera ; Lindley in Mitch. Trop. Austr. 306 (1848)	...	...	...	...	...	W.A. S.A.	—	V.	N.S.W.	Q. N.A.	B.fl.II,479	
H. exalata ; F. v. M. in Transact. R. Soc. of Vict. XXIV, 132 (1887)	...	...	...	...	...	—	—	—	—	N.S.W.	—	B.fl.II,479
H. racemosa ; Labillardière, Nov. Holl. pl. spec. I, 100, t. 128 (1804)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,480 M.fr.VIII,162.
H. Bauerlenii ; F. v. M. in Transact. R. Soc. Vict. XXIV, p. 132 (1887)	...	...	...	...	...	—	—	—	V.	N.S.W.	—	
H. rotundifolia ; Bentham, Fl. Austr. II, 480 (1864)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,480 M.fr.VIII,162.
H. rudis ; Bentham, Fl. Austr. II, 480 (1864)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,480
H. paniculata ; R. Brown in Benth. Fl. Austr. II, 481 (1864)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,481
H. pusilla ; R. Brown in Benth. Fl. Austr. II, 481 (1864)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,481
H. intricata ; Bentham, Fl. Austr. II, 481 (1864)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,481
H. nodulosa ; Walpers, rept. V, 671 (1846)...	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,481
H. trichostachya ; Bentham, Fl. Austr. II, 481 (1864)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,481

H. lanceolata ; R. Brown in Benth. Fl. Austr. II, 481 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,482					
H. micrantha ; R. Brown in Flind. Voy. II, 550 (1814) implied	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,482	M.fr.VIII,162; X,54	...	As., N.Z.		
H. stricta ; R. Brown in Benth. Fl. Austr. II, 482 (1864)	—	—	—	—	—	—	Q.	—	B.fl.II,482				
H. heterophylla ; Brongniart in Duperr. Voy. t. 68 (1829)	—	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.II,483					
H. leptotheca ; F. v. M., fragm. III, 32 (1862)	—	—	—	—	—	—	N.A.	B.fl.II,483	M.fr.VIII,162.				
H. tetragyna ; R. Brown in Flind. Voy. II, 550 (1814) implied	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,484	M.fr.III,32; IV,26	...	N.Z.		
H. teucroides ; Schlechtendal, Linnaea XX, 648 (1847)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,484					
H. scordioides ; Benth. Fl. Austr. II, 485 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,485					
H. depressa ; Walpers, repertor. II, 99 (1843)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.II,485	M.fr.IV,26	...	N.Z.		
H. salsoloides ; Benth. Fl. Austr. II, 485 (1864)	—	—	—	—	N.S.W.	—	—	B.fl.II,485					
H. monosperma ; F. v. M. in proceed. Linn. Soc. N.S.W., X, 197 (1885)	—	—	—	V.	N.S.W.	—	—	—					
H. Meionectes ; F. v. M., Key to the syst. of Vict. pl. I, 261 (1888)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.II,486	M.fr.VIII,163.				

MYRIOPHYLLUM ; P'Ecluse, rar. stirp. hist. II, 252 (1853) from Dioscorides. (Pelonastes).

M. variifolium ; J. Hooker in Hook. icon. plant. t. 239 (1840)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,487	...	...	N.Z.		
M. elatinooides ; Gaudichaud in Ann. des sc. nat. V, 105 (1825)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.II,487	...	...	Am., N.Z.		
M. verrucosum ; Lindley in Mitch. Trop. Austr. 384 (1848)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.II,488	M.fr.VIII,163.				
M. latifolium ; F. v. M., fragm. II, 87 (1860)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,488	M.fr.II,87; IV,172; X,54.				
M. Muellieri ; Sonder in Linnaea XXVIII, 233 (1855)	W.A.	S.A.	—	V.	—	—	—	B.fl.II,488	M.fr.VII,162.				
M. amphibium ; Labillardière, Nov. Holl. pl. spec. II, 70, t. 220 (1806)	—	S.A.	T.	V.	—	—	—	B.fl.II,489	M.fr.VIII,163.				
M. pedunculatum ; J. Hooker in Lond. Journ. VI, 474 (1847)	W.A.	S.A.	T.	V.	—	—	—	B.fl.II,489	M.fr.VIII,162	...	N.Z.		
M. dicoccum ; F. v. M. in Transact. phil. Inst. Vict. III, 41 (1858)	—	—	—	—	—	—	N.A.	B.fl.II,489	M.fr.VIII,162.				
M. trachycarpum ; F. v. M., fragm. II, 87 (1860)	—	—	—	—	—	—	N.A.	B.fl.II,489	M.fr.II,87.				
M. gracile ; Benth. Fl. Austr. II, 489 (1864)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,489					
M. filiforme ; Benth. Fl. Austr. II, 489 (1864)	—	—	—	—	—	—	N.A.	B.fl.II,489					
M. integrifolium ; J. Hooker, Fl. Tasm. I, 123, t. 23 (1860)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.II,490	M.fr.VIII,162.				
M. Drummondii ; J. Hooker, Fl. Tasm. I, 123 implied (1860)	W.A.	—	—	—	—	—	—	B.fl.II,490					

CALLITRICHINAE.

LINK, Enumer. plant. hort. reg. Berol. I, 7 (1821).—Genus 1, Species 2.

CALLITRICHE ; Linné, syst. nat. ed. sext. 82 (1748) from Colonna (1616).

C. verna ; Linné, flor. Suec. edit. sec. 32 (1755)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,492	...	...	E., As., Afr., Am., P., N.Z.		
C. Muellieri ; Sonder in Linnaea XXVIII, 229 (1855)	—	—	—	V.	N.S.W.	—	—	—	M.fr.VIII,163.				

RHIZOPHOREAE.

R. BROWN in Flind. Voy. II, 549 (1814).—Genera 4, Species 7.

RHIZOPHORA ; Linné, gen. plant. 137 (1737).

R. mucronata ; Lamarck, Encycl. méth. VI, 169 (1804)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.II,493	M.fr.IX,159	...	As., Afr., P.		
---	---	---	---	---	--------	----	------	-------------	-------------	-----	---------------	--	--

CERIOPS ; Arnott in Jardine, Ann. of nat. hist. I, 363 (1838).

C. Candolleana ; Arnott in Ann. of nat. hist. I, 364 (1838)	—	—	—	—	—	Q.	N.A.	B.fl.II,494	M.fr.IX,159	...	As., Afr.		
--	---	---	---	---	---	----	------	-------------	-------------	-----	-----------	--	--

BRUGUIERA ; Lamarck, Encycl. méth. IV, 696, t. 397 (1796).

B. Rhœdii ; Blume, enum. pl. Jav. 92 (1827)	—	—	—	—	—	Q.	N.A.	B.fl.II,494	...	...	As., P.		
B. caryophylloides ; Blume, enum. pl. Jav. 93 (1827)	—	—	—	—	—	Q.	—	—	M.fr.IX,159	...	As.		
B. parviflora ; Wight & Arnott, prodr. fl. penins. Ind. 311 (1834)	—	—	—	—	—	Q.	—	—	...	...	As.		
B. gymnorhiza ; Lamarck, Encycl. méth. IV, 696, t. 397 (1796)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.II,495	M.fr.IX,159.	...	As., Afr., P.		

CARALLIA ; Roxburgh in Flind. Voy. II, 549 (1814).

C. integerrima ; De Candolle, prodr. III, 33 (1828)	—	—	—	—	—	Q.	N.A.	B.fl.II,495	M.fr.IX,159.	...	As.		
--	---	---	---	---	---	----	------	-------------	--------------	-----	-----	--	--

COMBRETACEAE.

R. BROWN, prodr. 351 (1810).—Genera 4, Species 27.

TERMINALIA; Linné, mantiss. plant. 21 (1767). (Chuncoa.)

T. platyptera; F. v. M., fragm. II, 151 (1861)	...	...	...	...	...	—	—	—	—	Q. N.A. B.fl.II, 498	M.fr.II, 151; IX, 160.	
T. volucris; R. Brown in Benth. Fl. Austr. II, 498 (1864)...	...	...	...	...	...	—	—	—	—	—	N.A. B.fl.II, 498	
T. oblongata; F. v. M., fragm. II, 152 (1861)	...	...	...	...	...	—	—	—	—	Q. —	B.fl.II, 499	M.fr.III, 92, 167; IX, 160.
T. bursarina; F. v. M., fragm. II, 149 (1861)	...	...	...	...	...	—	—	—	—	Q. N.A. B.fl.II, 499	M.fr.II, 149.	
T. circumalata; F. v. M., fragm. III, 91 (1862)	...	...	...	...	...	W.A.	—	—	—	—	N.A. B.fl.II, 499	M.fr.III, 91.
T. pterocarya; F. v. M., fragm. II, 152 (1861)	...	...	...	...	...	—	—	—	—	—	N.A. B.fl.II, 500	M.fr.II, 152.
T. Thozetii; Benth. Fl. Austr. II, 500 (1864)	...	...	...	...	...	—	—	—	—	Q. —	B.fl.II, 500	M.fr.IX, 160.
T. Catappa; Linné, mantiss. pl. 128 (1767)	...	...	...	...	...	—	—	—	—	Q. —	—	M.fr.IX, 160. ... As., P.
T. melanocarpa; F. v. M., fragm. III, 92 (1862)	...	...	...	...	...	—	—	—	—	Q. N.A. B.fl.II, 500	M.fr.III, 92; IX, 160.	
T. Muelleri; Benth. Fl. Austr. II, 500 (1864)	...	...	...	...	...	—	—	—	—	Q. —	B.fl.II, 500	M.fr.IX, 160.
T. latipes; Benth. Fl. Austr. II, 501 (1864)	...	...	...	...	...	—	—	—	—	—	N.A. B.fl.II, 501	
T. edulis; F. v. M., fragm. II, 151 (1861)	...	...	...	...	...	—	—	—	—	—	N.A. B.fl.II, 501	M.fr.II, 151; IX, 160.
T. discolor; F. v. M., fragm. III, 92 (1862)	...	...	...	...	...	—	—	—	—	—	N.A. B.fl.II, 501	M.fr.III, 92.
T. porphyrocarpa; F. v. M. in Benth. Fl. Austr. II, 501 (1864)	...	...	...	...	...	—	—	—	—	Q. —	B.fl.II, 501	
T. platyphylla; F. v. M., fragm. II, 150 (1861)	...	...	...	...	...	—	—	—	—	Q. N.A. B.fl.II, 502	M.fr.II, 150.	
T. sericocarpa; F. v. M., fragm. IX, 159 (1875)	...	...	...	...	...	—	—	—	—	Q. —	—	M.fr.IX, 159.
T. microcarpa; Decaisne in Ann. du Mus. III (1834)	...	...	...	...	...	—	—	—	—	—	N.A. B.fl.II, 502	M.fr.II, 151; III, 92. As.
T. petiolaris; Cunningham in Benth. Fl. Austr. II, 502 (1864)	...	...	...	...	...	—	—	—	—	—	N.A. B.fl.II, 502	
T. erythrocarpa; F. v. M., fragm. II, 150 (1861)	...	...	...	...	...	—	—	—	—	—	N.A. B.fl.II, 503	M.fr.II, 150.
T. grandiflora; Benth. Fl. Austr. II, 503 (1864)	...	...	...	...	...	—	—	—	—	Q. N.A. B.fl.II, 503	M.fr.IX, 160.	

LUMNITZERA; Willdenow in den Verh. der Ges. Nat. Freunde zu Berl. IV, 186 (1803).

L. coccinea; Wight & Arnott, prodr. 316 (1834)	...	...	...	...	...	—	—	—	—	Q. —	B.fl.II, 503	M.fr.IX, 160 ... As., P.
L. racemosa; Willdenow in Verh. der Nat. Freunde zu Berl. IV, 186 (1803)	...	...	...	...	...	—	—	—	—	Q. N.A. B.fl.II, 504	M.fr.III, 92; IX, 160.	As., Afr., P.

MACROPTERANTHES; F. v. M., fragm. III, 91 (1862).

M. montana; F. v. M., fragm. III, 91 (1862)...	...	...	...	...	...	—	—	—	—	Q. —	B.fl.II, 504	M.fr.II, 149; III, 91 & 166.
M. Kekwickii; F. v. M., fragm. III, 151 (1863)	...	...	...	...	...	—	—	—	—	—	N.A. B.fl.II, 504	M.fr.II, 151.
M. Leichhardtii; F. v. M., fragm. III, 91 (1862)	...	...	...	...	...	—	—	—	—	Q. —	B.fl.II, 505	M.fr.III, 91.
M. Fitzalani; F. v. M., fragm. VIII, 160 (1874)	...	...	...	...	...	—	—	—	—	Q. —	—	M.fr. VIII, 160.

GYROCARPUS; N. Jacquin, select. stirp. Amer. hist. 282 (1763).

G. Americanus; N. Jacquin, select. stirp. Amer. hist. 282, t. 178 (1763)...	...	—	—	—	—	—	—	—	—	Q. N.A. B.fl.II, 505	M.fr.IX, 160...	... As., Am., P.
---	-----	---	---	---	---	---	---	---	---	----------------------	-----------------	------------------

MYRTACEAE.

ADANSON, Familles des plantes II, 86 (1763), from B. de Jussieu (1759).—Genera 40, Species 663.

ACTINODIUM; Schauer in Schlecht. Linnaea X, 311 (1835). (Triphelia.)

A. Cunninghamii; Schauer in Lindl. introd. to the nat. syst., sec. ed. 440 (1835)	W.A.	—	—	—	—	—	—	—	—	—	B.fl.III, 5	
---	------	---	---	---	---	---	---	---	---	---	-------------	--

DARWINIA; Rudge in Transact. Linn. Soc. XI, 299 (1813). (Genetyllis, Chamaelaucium, Homoranthus, Hedaroma, Polyzone, Francissia, Decalophium, Schuermannia, Cryptostemon.)

D. macrostegia; Benth. in Journ. Linn. Soc. IX, 179 (1865)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III, 8	
D. Hookeriana; Benth. in Journ. Linn. Soc. IX, 179 (1865)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III, 9	M.fr.VIII, 182.
D. fimbriata; Benth. in Journ. Linn. Soc. IX, 179 (1865)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III, 9	M.fr.VIII, 182; IX, 177.
D. speciosa; Benth. in Journ. Linn. Soc. IX, 179 (1865)...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III, 9	M.fr.XI, 10.
D. Meissneri; Benth. in Journ. Linn. Soc. IX, 179 (1865)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III, 9	M.fr.VIII, 182; IX, 177.

D. helichrysoides; Bentham in Journ. Linn. Soc. IX, 179 (1865) ...	...	W.A.	—	—	—	—	—	B.fl.III,10.
D. Neildiana; F. v. M., fragm. IX, 177 (1875) ...	...	W.A.	—	—	—	—	—	M.fr.IX,177.
D. oederoides; Bentham in Journ. Linn. Soc. IX, 179 (1865) ...	...	W.A.	—	—	—	—	—	B.fl.III,10 M.fr.IX,177; XI,9.
D. virescens; Bentham in Journ. Linn. Soc. IX, 179 (1865) ...	...	W.A.	—	—	—	—	—	B.fl.III,10 M.fr.XI,10.
D. Oldfieldii; Bentham in Journ. Linn. Soc. IX, 180 (1865) ...	...	W.A.	—	—	—	—	—	B.fl.III,10
D. purpurea; Bentham in Journ. Linn. Soc. IX, 180 (1865) ...	...	W.A.	—	—	—	—	—	B.fl.III,11 M.fr.XI,9.
D. citriodora; Bentham in Journ. Linn. Soc. IX, 180 (1865) ...	...	W.A.	—	—	—	—	—	B.fl.III,11 M.fr.I,169; IV,174; IX,177; XI,9.
D. thymoides; Bentham in Journ. Linn. Soc. IX, 180 (1865) ...	...	W.A.	—	—	—	—	—	B.fl.III,11 M.fr.VIII,182; IX,176.
D. taxifolia; Cunningham in Field's N. S. Wales 352 (1825) ...	...	—	—	—	V.	N.S.W.	—	B.fl.III,12 M.fr.IX,176.
D. vestita; Bentham in Journ. Linn. Soc. IX, 180 (1865) ...	...	W.A.	—	—	—	—	—	B.fl.III,12 M.fr.VIII,182; IX,76.
D. pauciflora; Bentham in Journ. Linn. Soc. IX, 180 (1865) ...	...	W.A.	—	—	—	—	—	B.fl.III,12 M.fr.XI,9.
D. diosmoides; Bentham in Journ. Linn. Soc. IX, 180 (1865) ...	...	W.A.	—	—	—	—	—	B.fl.III,13 M.fr.IX,176.
D. fascicularis; Rudge in Transact. Linn. Soc. XI, 299, t. 22 (1813) ...	...	—	—	—	N.S.W.	—	—	B.fl.III,13 M.fr.IX,176.
D. pinifolia; Bentham in Journ. Linn. Soc. IX, 181 (1865) ...	...	W.A.	—	—	—	—	—	B.fl.III,14
D. rhadinophylla; F. v. M., fragm. IX, 175 (1875) ...	...	W.A.	—	—	—	—	—	M.fr.IX,175.
D. sanguinea; Bentham in Journ. Linn. Soc. IX, 181 (1865) ...	...	W.A.	—	—	—	—	—	B.fl.III,14 M.fr.IX,176.
D. micropetala; Bentham in Journ. Linn. Soc. IX, 181 (1865) ...	...	—	S.A.	—	V.	—	—	B.fl.III,14 M.fr.I,12; IV,58; IX,176.
D. Schuermanni; Bentham in Journ. Linn. Soc. IX, 181 (1865) ...	...	—	S.A.	—	—	—	—	B.fl.III,14 M.fr.I,12; IV,58.
D. verticordina; Bentham in Journ. Linn. Soc. IX, 181 (1865) ...	...	W.A.	—	—	—	—	—	B.fl.III,15 M.fr.IV,57.
D. Thomasii; Bentham in Journ. Linn. Soc. IX, 181 (1865) ...	...	—	—	—	—	Q.	—	B.fl.III,15 M.fr.IV,137.
D. virgata; F. v. M., fragm. IX, 176 (1875) from Cunningham (1835) ...	...	—	—	—	V.	N.S.W.	Q.	B.fl.III,16 M.fr.IX,176.
D. ciliata; F. v. M., Cens. 51 (1882) from Desf. (1821) ...	...	W.A.	—	—	—	—	—	B.fl.III,36 M.fr.VIII,182.
D. gracilis; F. v. M., Cens. 51 (1882) ...	...	W.A.	—	—	—	—	—	B.fl.III,36 M.fr.IV,62; VIII,182.
D. Forrestii; F. v. M., fragm. XI, 9 (1878) ...	...	W.A.	—	—	—	—	—	M.fr.XI,9.
D. heterandra; F. v. M., Cens. 51 (1882) from Benth. (1866) ...	...	W.A.	—	—	—	—	—	B.fl.III,36
D. Drummondii; F. v. M., Cens. 51 (1882) from Meissn. (1857) ...	...	W.A.	—	—	—	—	—	B.fl.III,37
D. Endlicheri; F. v. M., Cens. 51 (1882) ...	...	W.A.	—	—	—	—	—	B.fl.III,37
D. brevifolia; F. v. M., Cens. 51 (1882) from Benth. (1866) ...	...	W.A.	—	—	—	—	—	B.fl.III,37
D. uncinata; F. v. M., Cens. 51 (1882) from Schauer (1844) ...	...	W.A.	—	—	—	—	—	B.fl.III,37
D. megalopetala; F. v. M., Cens. 51 (1882) ...	...	W.A.	—	—	—	—	—	B.fl.III,38 M.fr.VIII,182.
D. Turczaninowii; F. v. M., Cens. 51 (1882) ...	...	W.A.	—	—	—	—	—	B.fl.III,38 M.fr.IX,176; XI,9.
D. axillaris; F. v. M., Cens. 51 (1882) ...	...	W.A.	—	—	—	—	—	B.fl.III,38

VERTICORDIA; De Candolle in Dict. class XI, 400 (1826). (Chrysorrhoea, perhaps also Huegelia.)

V. Wilhelmii; F. v. M. in Transact. Vict. Inst. 122 (1855) ...	...	—	S.A.	—	—	—	—	B.fl.III,19 M.fr.X,28.
V. densiflora; Lindley in Bot. Regist. XXV, App. VI (1839) ...	...	W.A.	—	—	—	—	—	B.fl.III,20
V. stelluligera; Meissner in Journ. Linn. Soc. I, 38 (1857) ...	...	W.A.	—	—	—	—	—	B.fl.III,20
V. minutiflora; F. v. M., fragm. IV, 58 (1864) ...	...	W.A.	—	—	—	—	—	B.fl.III,20 M.fr.IV,38.
V. Fontainesii; De Candolle, prodr. III, 209 (1828) ...	...	W.A.	—	—	—	—	—	B.fl.III,21 M.fr.VIII,182; X,28.
V. helichrysantha; F. v. M. in Benth. Fl. Austr. III, 21 (1866) ...	...	W.A.	—	—	—	—	—	B.fl.III,21
V. Brownii; De Candolle, prodr. III, 209 (1828) ...	...	W.A.	—	—	—	—	—	B.fl.III,21
V. fastigiata; Turczaninow in Bullet. de l'Ac. de St. Petersb. 402 (1852) ...	...	W.A.	—	—	—	—	—	B.fl.III,22 M.fr.VIII,122; X,28.
V. Harveyi; Bentham, Fl. Austr. III, 22 (1864) ...	...	W.A.	—	—	—	—	—	B.fl.III,22
V. fimbrilepis; Turczaninow in Bull. Mosc. XX, 158 (1847) ...	...	W.A.	—	—	—	—	—	B.fl.III,22 M.fr.X,28.
V. serrata; Schauer in nov. act. acad. Caes. XIX, suppl. II, 70 (1841) ...	...	W.A.	—	—	—	—	—	B.fl.III,23
V. nitens; Schauer in nov. act. acad. Caes. XIX, suppl. II, 71, t. 4 (1841) ...	...	W.A.	—	—	—	—	—	B.fl.III,23 M.fr.X,28.
V. grandiflora; Endlicher in Ann. Wien. Mus. II, 195 (1838) ...	...	W.A.	—	—	—	—	—	B.fl.III,23
V. chrysanthia; Endlicher in Ann. Wien. Mus. II, 195 (1838) ...	...	W.A.	—	—	—	—	—	B.fl.III,24 M.fr.X,28.
V. Preissii; Schauer in Lehm. pl. Preiss. I, 101 (1844) ...	...	W.A.	—	—	—	—	—	B.fl.III,24 M.fr.VIII,182.
V. acerosa; Lindley in Bot. Regist. XXV, App. VI (1839) ...	...	W.A.	—	—	—	—	—	B.fl.III,25
V. polytricha; Bentham, Fl. Austr. III, 25 (1866) ...	...	W.A.	—	—	—	—	—	B.fl.III,25

V. oxylepis; Turczaninow in Bulet. de l'Ac. de St. Petersb. 402 (1852) ...	...	W.A.	—	—	—	—	—	—	B.fl.III,25	M.fr.VIII,182.
V. humilis; Benth. Fl. Austr. III, 26 (1866) ...	...	W.A.	—	—	—	—	—	—	B.fl.III,26	
V. penicillaris; F. v. M., fragm. I, 226 (1859) ...	...	W.A.	—	—	—	—	—	—	B.fl.III,26	M.fr.I,226.
V. multiflora; Turczaninow in Bull. Mosc. XX, 159 (1847) ...	...	W.A.	—	—	—	—	—	—	B.fl.III,26	M.fr.X,28.
V. Huegelii; Endlicher in Hueg. enum. 16 (1837) ...	...	W.A.	—	—	—	—	—	—	B.fl.III,27	
V. insignis; Endlicher in Hueg. enum. 47 (1837) ...	...	W.A.	—	—	—	—	—	—	B.fl.III,27	M.fr.X,28.
V. habrantha; Schauer in Lehm. pl. Preiss. I, 100 (1844) ...	...	W.A.	—	—	—	—	—	—	B.fl.III,28	M.fr.I,164.
V. monadelpha; Turczaninow in Bull. Mosc. XX, 158 (1847) ...	...	W.A.	—	—	—	—	—	—	B.fl.III,28	M.fr.I,227; X,28.
V. Lehmanni; Schauer in Lehm. pl. Preiss. I, 99 (1844) ...	...	W.A.	—	—	—	—	—	—	B.fl.III,29	
V. Cunninghamii; Schauer in nov. act. acad. Caes. XIX, suppl. II, 55 (1841) ...	...	—	—	—	—	—	—	Q. N.A.	B.fl.III,29	M.fr.X,28.
V. Jamiesonii; F. v. M. in Wieg's S. Sc. Rec. III, 68 (1883) ...	...	W.A.	—	—	—	—	—	—	—	
V. picta; Endlicher in Ann. des Wien. Mus. II, 194 (1838) ...	...	W.A.	—	—	—	—	—	—	B.fl.III,30	M.fr.X,28.
V. pennigera; Endlicher in Hueg. enum. 46 (1837) ...	...	W.A.	—	—	—	—	—	—	B.fl.III,30	M.fr.X,28.
V. Drummondii; Schauer in nov. act. acad. Caes. XIX, suppl. II, 56 (1841) ...	...	W.A.	—	—	—	—	—	—	B.fl.III,31	
V. Hughani; F. v. M., fragm. XI, 10 (1878) ...	...	W.A.	—	—	—	—	—	—	—	M.fr.XI,10.
V. pholidophylla; F. v. M., fragm. I, 227 (1859) ...	...	W.A.	—	—	—	—	—	—	B.fl.III,31	M.fr.I,227.
V. spicata; F. v. M., fragm. I, 226 (1859) ...	...	W.A.	—	—	—	—	—	—	B.fl.III,32	M.fr.I,226.
V. lepidophylla; F. v. M., fragm. I, 228 (1859) ...	...	W.A.	—	—	—	—	—	—	B.fl.III,32	M.fr.I,228.
V. Forrestii; F. v. M. in Wing's S. Sc. Rec. III, 67 (1883) ...	...	W.A.	—	—	—	—	—	—	—	
V. ovalifolia; Meissner in Journ. Linn. Soc. I, 40 (1857) ...	...	W.A.	—	—	—	—	—	—	B.fl.III,32	M.fr.I,227; X,28.
V. chrysostachya; Meissner in Journ. Linn. Soc. I, 41 (1857) ...	...	W.A.	—	—	—	—	—	—	B.fl.III,33	M.fr.I,227; X,28.
V. oculata; Meissner in Journ. Linn. Soc. I, 41 (1857) ...	...	W.A.	—	—	—	—	—	—	B.fl.III,33	M.fr.I,227.
V. grandis; Drummond in Hook. Kew Misc. V, 119 (1853) ...	...	W.A.	—	—	—	—	—	—	B.fl.III,33	M.fr.V,14; X,28; XI,10.

PILEANTHUS; Labillardière, Nov. Holl. pl. specim. II, 11 (1806).

P. peduncularis; Endlicher in Ann. des Wien. Mus. II, 196 (1838) ...	...	W.A.	—	—	—	—	—	—	B.fl.III,34	M.fr.I,225.
P. Limacis; Labillardière, Nov. Holl. pl. specim. II, 11, t. 149 (1806) ...	...	W.A.	—	—	—	—	—	—	B.fl.III,34	
P. filifolius; Meissner in Journ. Linn. Soc. I, 45 (1857) ...	...	W.A.	—	—	—	—	—	—	B.fl.III,35	

CALYCOTHRIX; Labillardière, Nov. Holl. pl. specim. II, 8, t. 146 (1806). (Calytrix, Calythrux.)

C. aurea; Lindley, Bot. Regist. XXV, App. 5, t. 3 (1839) ...	...	W.A.	—	—	—	—	—	—	B.fl.III,41	
C. puberula; Meissner in Journ. Linn. Soc. I, 48 (1857) ...	...	W.A.	—	—	—	—	—	—	B.fl.III,42	
C. flavesces; Cunningham in Bot. Mag. 3323 (1834) ...	...	W.A.	—	—	—	—	—	—	B.fl.III,42	
C. asperula; Schauer in Lehm. pl. Preiss. I, 106 (1844) ...	...	W.A.	—	—	—	—	—	—	B.fl.III,42	
C. sapphirina; Lindley, Bot. Regist. XXV, App. 5 (1839) ...	...	W.A.	—	—	—	—	—	—	B.fl.III,43	M.fr.I,224.
C. breviseta; Lindley, Bot. Regist. XXV, App. 5 (1839) ...	...	W.A.	—	—	—	—	—	—	B.fl.III,43	M.fr.VIII,182.
C. simplex; Lindley, Bot. Regist. XXV, App. 5 (1839) ...	...	W.A.	—	—	—	—	—	—	B.fl.III,43	
C. empetroides; Schauer in nov. act. acad. Caes. XIX, suppl. II, 102 (1841) ...	...	W.A.	—	—	—	—	—	—	B.fl.III,44	
C. variabilis; Lindley, Bot. Regist. XXV, App. 5 (1839) ...	...	W.A.	—	—	—	—	—	—	B.fl.III,44	
C. muricata; F. v. M., fragm. I, 224 (1859) ...	...	W.A.	—	—	—	—	—	—	B.fl.III,44	M.fr.I,224.
C. gracilis; Benth. Fl. Austr. III, 45 (1866) ...	...	W.A.	—	—	—	—	—	—	B.fl.III,45	
C. brevifolia; Meissner in Journ. Linn. Soc. I, 46 (1857) ...	...	W.A.	—	—	—	—	—	—	B.fl.III,45	
C. brachyphylla; Turczaninow in Bull. Mosc. XX, 161 (1847) ...	...	W.A.	—	—	—	—	—	—	B.fl.III,45	
C. Leschenaultii; Schauer in Lehm. pl. Preiss. I, 104 (1844) ...	...	W.A.	—	—	—	—	—	—	B.fl.III,46	
C. Oldfieldii; Benth. Fl. Austr. III, 46 (1866) ...	...	W.A.	—	—	—	—	—	—	B.fl.III,46	
C. glutinosa; Lindley, Bot. Regist. XXV, App. 5 (1839) ...	...	W.A.	—	—	—	—	—	—	B.fl.III,46	
C. angulata; Lindley, Bot. Regist. XXV, App. 6 (1839) ...	...	W.A.	—	—	—	—	—	—	B.fl.III,47	
C. depressa; Turczaninow in Bull. Soc. Mosc. XX, 162 (1847) ...	...	W.A.	—	—	—	—	—	—	B.fl.III,47	
C. tenuifolia; Meissner in Journ. Linn. Soc. I, 46 (1857) ...	...	W.A.	—	—	—	—	—	—	B.fl.III,47	
C. strigosa; Cunningham in Bot. Mag. 3323 (1834) ...	...	W.A.	—	—	—	—	—	—	B.fl.III,47	
C. plumulosa; F. v. M., fragm. X, 27 (1876) ..	...	W.A.	—	—	—	—	—	—	—	M.fr.X,27.

C. decandra; R. Brown in De Candolle, prodr. III, 208 (1828)	W.A.	—	—	—	—	—	—	B.f.III,48	M.fr.I,146.
C. Birdii; F. v. M., fragm. X, 26 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,26.
C. tenuiramea; Turczaninow in Bull. Mosc. XXII, 20 (1849)	W.A.	—	—	—	—	—	—	B.f.III,48	
C. Fraseri; Cunningham in Bot. Mag. 3323 (1834)	W.A.	—	—	—	—	—	—	B.f.III,48	M.fr.I,13.
C. granulosa; Benth. Fl. Austr. III, 49 (1866)	W.A.	—	—	—	—	—	—	B.f.III,49	
C. Creswelli; F. v. M., fragm. X, 27 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,27.
C. microphylla; Cunningham in Bot. Mag. 3323 (1834)	—	—	—	—	—	—	—	Q. N.A. B.f.III,49	
C. longiflora; F. v. M., fragm. I, 12 (1858)	—	S.A.	—	—	N.S.W.	Q.	—	B.f.III,49	M.fr.I,12.
C. leptophylla; Benth. Fl. Austr. III, 50 (1866)	—	—	—	—	—	Q.	—	B.f.III,50	
C. megaphylla; F. v. M., fragm. I, 146 (1859)	—	—	—	—	—	—	N.A.	B.f.III,50	M.fr.I,13,146.
C. tetragona; Labillardière, Nov. Holl. pl. specim. II, S, t. 146 (1806) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.f.III,50	M.fr.IV,36; VIII,183.
C. Sullivanii; F. v. M., fragm. IX, 1 (1875)	—	—	—	—	V.	—	—	—	M.fr.IX,1.
C. conferta; Cunningham in Bot. Mag. 3323 (1834)	—	—	—	—	—	—	N.A.	B.f.III,51	
C. arborescens; F. v. M. in Transact. phil. Inst. Vict. III, 42 (1858) ...	—	—	—	—	—	—	N.A.	B.f.III,51	
C. brachychaeta; F. v. M. in Transact. phil. Inst. Vict. III, 43 (1858) ...	—	—	—	—	—	—	N.A.	B.f.III,52	
C. achaeta; F. v. M. in Transact. phil. Inst. Vict. III, 43 (1858)	—	—	—	—	—	—	N.A.	B.f.III,52	
C. loricata; R. Brown in Benth. Fl. Austr. III, 52 (1866)	—	—	—	—	—	Q. N.A.	B.f.III,52		

LHOTZKYA; Schauer in Linnaea X, 309 (1835).

L. glaberrima; F. v. M., fragm. I, 13 (1858)	—	—	S.A.	—	—	—	—	B.f.III,53	M.fr.I,13.
L. genetyloides; F. v. M. in Transact. phil. Soc. Vict. I, 16 (1854) ...	—	—	S.A.	—	V.	—	—	B.f.III,54	
L. Smeatoniana; F. v. M. in Austral. Journ. of Pharm. II, 5 (1887) ...	—	—	S.A.	—	—	—	—	—	
L. violacea; Lindley, Bot. Regist. XXV, App. VII (1839)	W.A.	—	—	—	—	—	—	B.f.III,54	M.fr.XI,8.
L. ciliata; F. v. M. in Benth. Fl. Austr. III, 54 (1866)	W.A.	—	—	—	—	—	—	B.f.III,54	
L. brevifolia; Schauer in Lehm. pl. Preiss. I, 103 (1844)	W.A.	—	—	—	—	—	—	B.f.III,54	
L. purpurea; F. v. M., fragm. I, 224 (1859)	W.A.	—	—	—	—	—	—	B.f.III,55	M.fr.I,224.
L. ericoides; Schauer in Lindl. Introd. sec. ed. 439 (1835)	W.A.	—	—	—	—	—	—	B.f.III,55	M.fr.XI,9.
L. acutifolia; Lindley, Bot. Regist. XXV, App. VII (1839)	W.A.	—	—	—	—	—	—	B.f.III,55	M.fr.XI,8.
L. Harvestiana; F. v. M., fragm. XI, 8 (1878)	W.A.	—	—	—	—	—	—	—	M.fr.XI,8.

THRYPTOMENE; Endlicher in den Ann. des Wien. Mus. II, 192 (1838). (Homalocalyx, Paryphantha, Astraea, Eremopyxis, Micromyrtus.)

T. mucronulata; Turczaninow in Bull. Mosc. XX, 156 (1847)	W.A.	—	—	—	—	—	—	B.f.III,58	
T. Australis; Endlicher in den Ann. des Wien. Mus. II, 192 (1838) ...	W.A.	—	—	—	—	—	—	B.f.III,58	
T. tenella; Benth. Fl. Austr. III, 59 (1866)	W.A.	—	—	—	—	—	—	B.f.III,59	M.fr.VIII,182; X,24.
T. prolifera; Turczaninow in Bull. Mosc. XXXV, 324 (1862)	W.A.	—	—	—	—	—	—	B.f.III,59	
T. saxicola; Schauer in Lehm. pl. Preiss. I, 102 (1844)	W.A.	—	—	—	—	—	—	B.f.III,59	M.fr.IV,75.
T. Johnsonii; F. v. M., fragm. IV, 77 (1864)	W.A.	—	—	—	—	—	—	B.f.III,60	M.fr.IV,77.
T. racemulosa; Turczaninow in Bull. Mosc. XX, 156 (1847)	W.A.	—	—	—	—	—	—	B.f.III,60	
T. denticulata; Benth. in Fl. Austr. III, 60 (1866)	W.A.	—	—	—	—	—	—	B.f.III,60	M.fr.IV,75.
T. baeckeacea; F. v. M., fragm. IV, 65 (1864)	W.A.	—	—	—	—	—	—	B.f.III,60	M.fr.IV,65.
T. stronglyphylla; F. v. M. in Benth. Fl. Austr. III, 61 (1866) ..	W.A.	—	—	—	—	—	—	B.f.III,61	
T. hyporhytis; Turczaninow in Bull. Mosc. XXXV, 324 (1862)	W.A.	—	—	—	—	—	—	B.f.III,61	
T. Maisonneuvii; F. v. M., fragm. IV, 64 (1864)	—	S.A.	—	—	—	—	N.A.	B.f.III,61	M.fr.IV,64.
T. urceolaris; F. v. M., fragm. X, 25 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,25.
T. Mitchelliana; F. v. M., fragm. I, 11 (1858)	—	S.A.	—	V.	—	—	—	B.f.III,61	M.fr.XI,62.
T. Miqueliana; F. v. M., fragm. I, 11 (1858)	—	S.A.	—	—	N.S.W.	—	—	B.f.III,62	M.fr.I,11.
T. micrantha; J. Hooker in Hook. Kew. Misc. V, 299, t. 8 (1853) ...	—	—	T.	—	—	—	—	B.f.III,62	
T. stenocalyx; F. v. M., fragm. X, 23 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,23.
T. ericaea; F. v. M., fragm. I, 12 (1858)	—	S.A.	—	V.	—	—	—	B.f.III,62	M.fr.I,12.
T. oligandra; F. v. M., fragm. I, 11 (1858)	W.A.	—	—	—	—	Q. N.A.	B.f.III,63	M.fr.IV,169.	
T. elobata; F. v. M., fragm. IV, 63 (1864)	W.A.	—	—	—	—	—	—	B.f.III,64	M.fr.IV,63.

T. racemosa; F. v. M. in Benth. Fl. Austr. III, 64 (1866) ...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,64	
T. trachycalyx; F. v. M., fragm. X, 25 (1876) ...	...	...	...	W.A.	—	—	—	—	—	—	—	M.fr.X,25.
T. imbricata; F. v. M., Cens. 53 (1882) from Benth. (1866) ...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,64	
T. flaviflora; F. v. M., fragm. VIII, 13 (1873) ...	...	...	...	W.A.	S.A.	—	—	—	—	—	—	M.fr.X,25.
T. Drummondii; F. v. M., Cens. 53 (1882) from Benth. (1866) ...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,64	M.fr.VIII,183.
T. Elliottii; F. v. M., fragm. IX, 62 (1875) ...	...	...	...	...	S.A.	—	—	—	—	—	—	M.fr.IX,62.
T. auriculata; F. v. M., fragm. X, 24 (1876) ...	...	...	...	...	S.A.	—	—	—	—	—	—	M.fr.X,24.
T. hymenonema; F. v. M., fragm. X, 26 (1876) ...	...	...	...	W.A.	—	—	—	—	—	—	—	M.fr.X,26.
T. ciliata; F. v. M. in Woolls, pl. of the neighb. of Sydney 23 (1880) ...	...	...	...	...	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,65	M.fr.I,30&242; IV,63.
T. minutiflora; F. v. M. in Woolls, pl. of the neighb. of Sydney 23 (1880) ...	...	...	...	...	—	—	—	—	N.S.W.	—	B.fl.III,65	
T. leptocalyx; F. v. M., Cens. 53 (1882) from Benth. (1866) ...	...	...	...	...	—	—	—	—	—	Q.	B.fl.III,65	M.fr.I,30.
T. homalocalyx; F. v. M., fragm. IV, 63 (1864) ...	...	...	...	...	—	—	—	—	—	Q. N.A.	B.fl.III,56	M.fr.IV,63.
T. polyandra; F. v. M., fragm. IV, 77 (1864) ...	...	...	...	...	—	—	—	—	—	Q. N.A.	B.fl.III,56	M.fr.IV,77.

WEHLIA; F. v. M., fragm. X, 22 (1876).

W. thryptomenoides; F. v. M., fragm. X, 22 (1876) ...	...	...	...	W.A.	—	—	—	—	—	—	—	M.fr.X,22.
W. coarctata; F. v. M., fragm. X, 23 (1876) ...	...	...	...	W.A.	—	—	—	—	—	—	—	M.fr.X,23.
W. staminosa; F. v. M. in Wing's S. Sc. Rec. III, 282 (1884) ...	...	...	...	W.A.	—	—	—	—	—	—	—	

BAECKEA; Linné, spec. plant. 358 (1753). (Jungia, Imbricaria, Scholtzia, Schidiomyrtus, Rinzia, Earyomyrtus, Camphoromyrtus, Tetrapora, Harmogia, Oxymyrrhine, Babingtonia, Ericomyrtus, Piptandra, Anticoryne.)

B. platystemonea; Benth. Fl. Austr. III, 74 (1866) ...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,74	M.fr.VIII,183.
B. Funmana; F. v. M., fragm. IV, 68 (1864) ...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,74	M.fr.IV,68.
B. dimorphandra; F. v. M. in Benth. Fl. Austr. III, 74 (1866) ...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,74	
B. schollerifolia; Lehmann, pl. Preiss. II, 369 (1847) ...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,75	M.fr.IV,68; VIII,183.
B. oxycoccoides; Benth. Fl. Austr. III, 75 (1866) ...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,75	M.fr.VIII,183.
B. Drummondii; Benth. Fl. Austr. III, 75 (1866) ...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,75	M.fr.VIII,183.
B. diffusa; Sieber in De Candolle, prodr. III, 230 (1828) ...	...	...	...	...	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,76	M.fr.IV,67.
B. crassifolia; Lindley in Mitch. Three Exped. II, 115 (1838) ...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,76	M.fr.IV,66.
B. cryptandroides; F. v. M., fragm. X, 29 (1876) ...	...	...	...	W.A.	—	—	—	—	—	—	—	M.fr.X,29.
B. tetragona; F. v. M. in Benth. Fl. Austr. III, 77 (1866) ...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,77	
B. ericaea; F. v. M., fragm. I, 31 (1858) ...	...	...	...	...	S.A.	—	V.	N.S.W.	—	—	B.fl.III,77	
B. polystemonea; F. v. M., fragm. II, 124 (1861) ...	...	...	...	...	S.A.	—	—	—	—	—	B.fl.III,77	M.fr.III,24.
B. crenulata; R. Brown in Flind. voy. 548 (1814) ...	...	...	...	...	—	—	—	N.S.W.	Q.	—	B.fl.III,78	M.fr.IV,65.
B. brevifolia; De Candolle, prodr. III, 230 (1828) ...	...	...	...	...	—	—	—	N.S.W.	—	—	B.fl.III,78	
B. Gunniana; Shauer in Walp. Rep. II, 920 (1843) ...	...	...	...	...	—	—	T.	V.	N.S.W.	—	B.fl.III,78	M.fr.IV,66.
B. diosmifolia; Rudge in Transact. Linn. Soc. VIII, 298, t. 13 (1807) ...	...	...	...	...	—	—	—	N.S.W.	Q.	—	B.fl.III,79	M.fr.I,29.
B. leptocaulis; J. Hooker, icon. pl. t. 298 (1840) ...	...	...	...	...	—	—	T.	—	—	—	B.fl.III,79	
B. arbuscula; R. Brown in Benth. Fl. Austr. III, 79 (1866) ...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,79	
B. astarteoides; Benth. Fl. Austr. III, 80 (1866) ...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,80	
B. linifolia; Rudge in Transact. Linn. Soc. VIII, 297, t. 12 (1807) ...	...	...	...	...	—	—	—	V.	N.S.W.	—	B.fl.III,80	M.fr.IV,71.
B. stenophylla; F. v. M., fragm. I, 13 (1858) ...	...	...	...	...	—	—	—	N.S.W.	Q.	—	B.fl.III,80	M.fr.I,13.
B. camphorata; R. Brown in Bot. Mag. t. 2694 (1826) ...	...	...	...	...	—	—	—	V.	N.S.W.	—	B.fl.III,81	M.fr.IV,70; IV,177.
B. virgata; Andrews, Bot. Reposit. t. 598 (1810) ...	...	...	...	...	—	—	—	V.	N.S.W.	Q. N.A.	B.fl.III,81	M.fr.I,69; II,31 ... P.
B. crenatifolia; F. v. M., fragm. IV, 70 (1864) ...	...	...	...	...	—	—	—	V.	—	—	B.fl.III,82	M.fr.IV,70.
B. Cunninghamii; Benth. Fl. Austr. III, 82 (1866) ...	...	...	...	...	—	—	—	N.S.W.	—	—	B.fl.III,82	
B. densifolia; Smith in Transact. Linn. Soc. III, 260 (1797) ...	...	...	...	...	—	—	—	N.S.W.	Q.	—	B.fl.III,82	M.fr.IV,71.
B. Behrii; F. v. M., fragm. IV, 68 (1864) ...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,83	M.fr.IV,68.
B. uncinella; Benth. Fl. Austr. III, 84 (1866) ...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,84	
B. polyandra; F. v. M., fragm. IV, 12 (1864) ...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,84	M.fr.II,30; IV,72.
B. corynophylla; F. v. M., fragm. IV, 72 (1864) ...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,84	M.fr.IV,72.

B. pachyphylla ; Benth. Fl. Austr. III, 85 (1866)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,85
B. crispiflora ; F. v. M., fragm. IV, 72 (1864)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,85 M.fr.II,31.
B. Blackettii ; F. v. M., fragm. VIII, 181 (1874)	...	...	...	...	W.A.	—	—	—	—	—	—	M.fr.VIII,181.
B. camphoresmae ; Endlicher in Hueg. enum. 51 (1837)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,86
B. pulchella ; De Candolle, prodr. III, 230 (1828)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,86 M.fr.IV,67.
B. ochropetala ; F. v. M., fragm. X, 29 (1876)	...	...	...	...	W.A.	—	—	—	—	—	—	M.fr.X,29.
B. pygmaea ; R. Brown in Benth. Fl. Austr. III, 86 (1866)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,86
B. corymbulosa ; Benth. Fl. Austr. III, 87 (1866)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,87 M.fr.VIII,183.
B. floribunda ; Benth. Fl. Austr. III, 87 (1866)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,87
B. pentandra ; F. v. M., fragm. II, 31 ; IV, 72 (1864)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,87 M.fr.II,31 ; IV,72 ; VIII,183.
B. pentagonantha ; F. v. M., fragm. IV, 73 (1864)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,87 M.fr.IV,73.
B. robusta ; F. v. M., fragm. IV, 72 (1864)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,88 M.fr.IV,72.
B. ovalifolia ; F. v. M., fragm. IV, 72 (1864)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,88 M.fr.II,32 ; IV,72 ; VIII,183.
B. subcuneata ; F. v. M., fragm. IV, 73 (1864)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,88 M.fr.IV,73.
B. grandiflora ; Benth. Fl. Austr. III, 89 (1866)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,89
B. uberiflora ; F. v. M., Cens. 53 (1882)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,67 M.fr.IV,74.
B. involucrata ; Endlicher in Hueg. enum. 51 (1837)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,68 M.fr.IV,74.
B. spathulata ; F. v. M., Cens. 54 (1882) from Benth. (1866)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,68
B. ciliata ; F. v. M., Cens. 54 (1882)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,68 M.fr.IV,76.
B. capitata ; F. v. M., Cens. 54 (1882)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,69
B. umbellifera ; F. v. M., Cens. 54 (1882)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,69 M.fr.IV,75.
B. laxiflora ; F. v. M., Cens. 54 (1882) from Benth. (1866)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,69
B. leptantha ; F. v. M., Cens. 54 (1882) from Benth. (1866)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,69
B. parviflora ; F. v. M., Cens. 54 (1882)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,70 M.fr.IV,76.
B. oligandra ; F. v. M., Cens. 54 (1882)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,70
B. serpillifolia ; F. v. M., fragm. X, 30 (1876)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,70 M.fr.VIII,183.
B. teretifolia ; F. v. M., Cens. 54 (1882) from Benth. (1866)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,70

ASTARTEA ; De Candolle in Dict. class. XI, 400 (1826).

A. ambigua ; F. v. M., fragm. II, 32 (1860)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,89 M.fr.II,32.
A. fascicularis ; De Candolle, prodr. III, 210 (1828)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,90 M.fr.VIII,183.
A. intratropica ; F. v. M., fragm. I, 83 (1858)	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.III,90 M.fr.I,83.

HYPOCALYMMMA ; Endlicher in Hueg. enum. pl. Nov. Holl. Austr. occ. 51 (1837), implied.

H. xanthopetalum ; F. v. M., fragm. II, 29 (1860)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,92 M.fr.II,29.
H. robustum ; Endlicher in Hueg. enum. 50 (1837)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,92
H. longifolium ; F. v. M., fragm. II, 28 (1860)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,92 M.fr.II,28.
H. strictum ; Schauer in Lehm. pl. Preiss. I, 111 (1844)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,93
H. tetrapterum ; Turczaninow in Bull. Mosc. XXXV, 325 (1862)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,93
H. linifolium ; Turczaninow in Bull. Mosc. XXXV, 325 (1862)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,93
H. angustifolium ; Endlicher in Hueg. enum. 50 (1837)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,94 M.fr.II,28.
H. ericifolium ; Benth. Fl. Austr. III, 94 (1866)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,94
H. cordifolium ; Lehmann, plant. Preiss. I, 112 (1844)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,94
H. speciosum ; Turczaninow in Bull. de l'Acad. de St. Pet. 409 (1852)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,95 M.fr.VIII,183.
H. Phillipsii ; Harvey in nat. hist. Rev. V, 296, t. 22 (1865)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,95
H. myrtifolium ; Turczaninow in Bull. de St. Pet. 410 (1852)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,95 M.fr.VIII,183.

BALAUSTION ; Hooker, icones plantarum 852 (1852). (Punicella, Cheynia.)

B. pulcherrimum ; Hooker, icon. plant. 852 (1852)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,96 M.fr.VIII,183.
---	-----	-----	-----	-----	------	---	---	---	---	---	---	----------------------------

AGONIS ; De Candolle, prodr. III, 226 (1828), implied. (Billiotia, Billetia, Billottia.)

A. spathulata ; Schauer in Lehm. pl. Preiss. I, 117 (1844)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,97 M.fr.XI,120.
A. floribunda ; Turczaninow in Bull. Mosc. II, 20 (1849)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,97

A. marginata; De Candolle, prodr. III, 226 (1828)	W.A.	—	—	—	—	—	—	B.fl.III,98	M.fr.XI,120.
A. lysicephala; F. v. M. & Bailey in synops. Queensl. Flora sec. suppl. 24 (1888)	—	—	—	—	—	—	Q.	—	—
A. linearifolia; De Candolle, prodr. III, 227 (1828)	W.A.	—	—	—	—	—	—	B.fl.III,98	M.fr.VIII,183; XI,120.
A. juniperina; Schauer in Lehm. pl. Preiss. I, 118 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,98	M.fr.XI,120.
A. parviceps; Schauer in Lehm. pl. Preiss. I, 119 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,99	M.fr.XI,120.
A. obtusissima; F. v. M., fragm. XI, 119 (1881)	W.A.	—	—	—	—	—	—	—	M.fr.XI,119.
A. flexuosa; De Candolle, prodr. III, 226 (1828)	W.A.	—	—	—	—	—	—	B.fl.III,99	M.fr.XI,120.
A. undulata; Bentham, Fl. Austr. III, 100 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,100	—
A. hypericifolia; Schauer in Lehm. pl. Preiss. I, 117 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,100	M.fr.XI,120.
A. Scortechiniana; F. v. M., fragm. XI, 118 (1881)	—	—	—	—	—	N.S.W.	Q.	—	M.fr.XI,118.
A. grandiflora; Bentham, Fl. Austr. III, 100 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,100	M.fr.XI,120.

LEPTOSPERMUM; R. et G. Forster, char. gen. 71, t. 36 (1776). (Fabricia, Homalospermum, Pericalymma.)

L. Fabricia; Bentham, Fl. Austr. III, 102 (1866)	—	—	—	—	—	—	Q.	—	B.fl.III,102
L. laevigatum; F. v. M., Annual Report 22 (1858)	—	S.A.	T.	V.	N.S.W.	—	—	—	B.fl.III,103 M.fr.IV,60.
L. firmum; Bentham, Fl. Austr. III, 104 (1866)	W.A.	—	—	—	—	—	—	—	B.fl.III,104
L. flavescens; Smith in Transact. Linn. Soc. III, 262 (1797)	—	—	T.	V.	N.S.W.	Q.	—	—	B.fl.III,104 M.fr.IV,61 ... As.
L. scoparium; R. & G. Forster, char. gen. 48 (1776)	—	S.A.	T.	V.	N.S.W.	Q.	—	—	B.fl.III,105 M.fr.IV,61 ... N.Z.
L. arachnoideum; Smith in Transact. Linn. Soc. III, 263 (1797)	—	—	—	—	N.S.W.	—	—	—	B.fl.III,105
L. spinescens; Endlicher in Hueg. enum. 51 (1837)	W.A.	—	—	—	—	—	—	—	B.fl.III,106
L. lanigerum; Smith in Transact. Linn. Soc. III, 263 (1797)	—	S.A.	T.	V.	N.S.W.	Q.	—	—	B.fl.III,106
L. parvifolium; Smith in Transact. Linn. Soc. III, 263 (1797)	—	—	—	—	N.S.W.	—	—	—	B.fl.III,107
L. stellatum; Cavanilles, icon. pl. IV, 16, t. 330	—	—	—	—	N.S.W.	Q.	—	—	B.fl.III,107
L. attenuatum; Smith in Transact. Linn. Soc. III, 262 (1797)	—	—	—	—	V.	N.S.W.	Q.	—	B.fl.III,108 M.fr.II,28; IV,62.
L. myrtifolium; Sieber in De Candolle, prodr. III, 238 (1828)	—	—	T.	V.	N.S.W.	Q.	—	—	B.fl.III,108
L. rupestre; J. Hooker in Hook. icon. pl. t. 308 (1841)	—	—	T.	—	—	—	—	—	B.fl.III,108
L. myrsinoides; Schlechtendal in Linnaea XX, 653 (1847)	—	S.A.	—	V.	N.S.W.	—	—	—	B.fl.III,109 M.fr.VI,61.
L. erubescens; Schauer in Lehm. pl. Preiss. I, 121 (1844)	W.A.	—	—	—	—	—	—	—	B.fl.III,109 M.fr.II,28; IV,61; VIII,183.
L. abnorme; F. v. M. in Benth. Fl. Austr. III, 109 (1866)	—	—	—	—	N.S.W.	Q.	N.A.	—	B.fl.III,109 M.fr.II,27.
L. Roei; Bentham, Fl. Austr. III, 110 (1866)	W.A.	—	—	—	—	—	—	—	B.fl.III,110
L. floridum; Bentham, Fl. Austr. III, 110 (1866)	W.A.	—	—	—	—	—	—	—	B.fl.III,110
L. ellipticum; Endlicher in Hueg. enum. 51 (1837)	W.A.	—	—	—	—	—	—	—	B.fl.III,110
L. crassipes; Lehmann, ind. sem. hort. Hamb. (1842)	W.A.	—	—	—	—	—	—	—	B.fl.III,110

KUNZEA; Reichenbach, conspect. regn. veget. 175 (1828). (Salisia, Pentagonaster.)

K. micrantha; Schauer in Lehm. pl. Preiss. I, 125 (1844)	W.A.	—	—	—	—	—	—	—	B.fl.III,112 M.fr.VIII,183.
K. eriocalyx; F. v. M., fragm. II, 28 (1860)	W.A.	—	—	—	—	—	—	—	B.fl.III,112 M.fr.II,28.
K. Muelleri; Bentham, Fl. Austr. III, 113 (1866)	—	—	—	V.	N.S.W.	—	—	—	B.fl.III,113
K. ericifolia; Reichenbach, consp. 175 (1828)	W.A.	—	—	—	—	—	—	—	B.fl.III,113
K. Preissiana; Schauer in Lehm. pl. Preiss. I, 125 (1844)	W.A.	—	—	—	—	—	—	—	B.fl.III,113
K. recurva; Schauer in Lehm. pl. Preiss. I, 125 (1844)	W.A.	—	—	—	—	—	—	—	B.fl.III,114 M.fr.VIII,183.
K. micromera; Schauer in Lehm. pl. Preiss. III, 223 (1847)	W.A.	—	—	—	—	—	—	—	B.fl.III,114
K. pauciflora; Schauer in Lehm. pl. Preiss. I, 124 (1844)	W.A.	—	—	—	—	—	—	—	B.fl.III,114 M.fr.II,28; VIII,183.
K. calida; F. v. M., fragm. VI, 23 (1867)	—	—	—	—	—	—	Q.	—	M.fr.VI,23.
K. parvifolia; Schauer in Lehm. pl. Preiss. I, 124 (1844)	—	—	—	—	N.S.W.	—	—	—	B.fl.III,115 M.fr.II,28; VIII,13.
K. peduncularis; F. v. M. in Transact. Vict. Inst. 144 (1855)	—	—	—	V.	N.S.W.	Q.	—	—	B.fl.III,115 M.fr.II,28; X,55.
K. corifolia; Reichenbach, consp. 175 (1828)	—	—	—	T.	V.	N.S.W.	—	—	B.fl.III,115
K. capitata; Reichenbach, consp. 175 (1828)	—	—	—	—	N.S.W.	—	—	—	B.fl.III,116 M.fr.II,28.
K. pomifera; F. v. M. in Transact. Vict. Inst. 124 (1855)	—	S.A.	—	V.	—	—	—	—	B.fl.III,116 M.fr.IV,62.
K. sericea; Turczaninow in Bull. Mosc. XX, 162 (1847)	W.A.	—	—	—	—	—	—	—	B.fl.III,117
K. Baxteri; Schauer in Lehm. pl. Preiss. I, 123 (1844)	W.A.	—	—	—	—	—	—	—	B.fl.III,117 M.fr.III,153.
K. opposita; F. v. M., fragm. VI, 24 (1867)	—	—	—	—	N.S.W.	—	—	—	M.fr.VI,24.

CALLISTEMON; R. Brown in Flind. voy. II, 547 (1814).

<i>C. speciosus</i> ; De Candolle, prodr. III, 224 (1828)	W.A.	—	—	—	—	—	B.f.III, 119	M.fr.I, 14.
<i>C. phoeniceus</i> ; Lindley, Bot. Regist. XXV, App. 10 (1839)	W.A.	—	—	—	—	—	B.f.III, 119	M.fr.IV, 53.
<i>C. lanceolatus</i> ; De Candolle, prodr. III, 224 (1828)	—	—	—	V.	N.S.W.	Q.	B.f.III, 120	M.fr.I, 14; IV, 53.
<i>C. coccineus</i> ; F. v. M., fragm. I, 13 (1858)	—	S.A.	—	V.	N.S.W.	—	B.f.III, 120	M.fr.IV, 54.
<i>C. salignus</i> ; De Candolle, prodr. III, 223 (1828)	—	S.A.	T.	V.	N.S.W.	Q.	B.f.III, 120	M.fr.I, 14; IV, 54.
<i>C. pityoides</i> ; Miquel in Nederl. Kruidk. Arch. IV, 140 (1859)	—	—	—	V.	N.S.W.	Q.	—	—
<i>C. rigidus</i> ; R. Brown in Bot. Regist. t. 393 (1819)	—	—	—	—	N.S.W.	—	B.f.III, 121	M.fr.IV, 54.
<i>C. linearis</i> ; De Candolle, prodr. III, 223 (1828)	—	—	—	V.	N.S.W.	—	B.f.III, 122	M.fr.IV, 54.
<i>C. pinifolius</i> ; De Candolle, prodr. III, 223 (1828)	—	—	—	—	N.S.W.	—	B.f.III, 122	M.fr.IV, 53.
<i>C. teretifolius</i> ; F. v. M. in Schlecht. Linnæa XXV, 387 (1852)	—	S.A.	—	—	—	—	B.f.III, 122	M.fr.IV, 53.
<i>C. brachyandrus</i> ; Lindley in Journ. Hort. Soc. IV, 112 (1849)	—	S.A.	—	V.	N.S.W.	—	B.f.III, 122	M.fr.IV, 52.

MELALEUCA; Linné, mantiss. plant. 14 (1767). (*Gymnagathis*, *Asteromyrtus*.)

<i>M. macronychia</i> ; Turczaninow in Bullet. de St. Petersburg. 420 (1852)	W.A.	—	—	—	—	—	B.f.III, 129	M.fr.VIII, 183.
<i>M. lateritia</i> ; Otto in allgem. Gart. Zeit. II, 257 (1834)	W.A.	—	—	—	—	—	B.f.III, 130	—
<i>M. calothamnoides</i> ; F. v. M., fragm. III, 114 (1862)	W.A.	—	—	—	—	—	B.f.III, 130	M.fr.III, 114.
<i>M. blaeriaefolia</i> ; Turczaninow in Bull. Mosc. XX, 165 (1847)	W.A.	—	—	—	—	—	B.f.III, 130	—
<i>M. diosmifolia</i> ; Andrews, Bot. Reposit. t. 476 (1808)	W.A.	—	—	—	—	—	B.f.III, 130	—
<i>M. elliptica</i> ; Labillardière, Nov. Holl. pl. spec. II, 31, t. 173 (1806)	W.A.	—	—	—	—	—	B.f.III, 131	M.fr.VIII, 183.
<i>M. hypericifolia</i> ; Smith in Transact. Linn. Soc. III, 279 (1797)	—	—	—	V.	N.S.W.	—	B.f.III, 131	—
<i>M. fulgens</i> ; R. Brown in Ait. hort. Kew. sec. ed. IV, 415 (1812)	W.A.	—	—	—	—	—	B.f.III, 131	—
<i>M. acuminata</i> ; F. v. M., fragm. I, 15 (1858)	W.A.	S.A.	—	V.	N.S.W.	—	B.f.III, 132	M.fr.I, 15.
<i>M. leptoclada</i> ; Benth. Fl. Austr. III, 132 (1866)	W.A.	—	—	—	—	—	B.f.III, 132	—
<i>M. basiccephala</i> ; Benth. Fl. Austr. III, 133 (1866)	W.A.	—	—	—	—	—	B.f.III, 133	—
<i>M. gibbosa</i> ; Labillardière, Nov. Holl. pl. spec. II, 30, t. 172 (1806)	—	S.A.	T.	V.	—	—	B.f.III, 133	—
<i>M. decussata</i> ; R. Brown in Ait. hort. Kew. sec. ed. IV, 415 (1812)	—	S.A.	—	V.	—	—	B.f.III, 133	—
<i>M. Wilsonii</i> ; F. v. M., fragm. II, 124, t. 15 (1861)	—	S.A.	—	V.	—	—	B.f.III, 134	M.fr.II, 124.
<i>M. thymifolia</i> ; Smith in Transact. Linn. Soc. III, 278 (1797)	—	—	—	—	N.S.W.	Q.	B.f.III, 134	M.fr.X, 55.
<i>M. violacea</i> ; Lindley, Bot. Regist. XXV, App. 8 (1839)	W.A.	—	—	—	—	—	B.f.III, 135	—
<i>M. cardiophylla</i> ; F. v. M., fragm. I, 225 (1859)	W.A.	—	—	—	—	—	B.f.III, 135	M.fr.III, 164.
<i>M. undulata</i> ; Benth. Fl. Austr. III, 135 (1866)	W.A.	—	—	—	—	—	B.f.III, 135	—
<i>M. depanperata</i> ; Turczaninow in Bullet. de St. Petersburg. 424 (1852)	W.A.	—	—	—	—	—	B.f.III, 136	M.fr.III, 120; VIII, 183.
<i>M. lateriflora</i> ; Benth. Fl. Austr. III, 136 (1866)	W.A.	—	—	—	—	—	B.f.III, 136	—
<i>M. exarata</i> ; F. v. M., fragm. III, 114 (1862)	W.A.	—	—	—	—	—	B.f.III, 136	M.fr.VIII, 184.
<i>M. teretifolia</i> ; Endlicher in Hueg. ennm. 49 (1837)	W.A.	—	—	—	—	—	B.f.III, 137	M.fr.III, 117.
<i>M. alsophila</i> ; Cunningham in Benth. Fl. Austr. III, 137 (1866)	—	—	—	—	—	N.A.	B.f.III, 137	—
<i>M. acacioides</i> ; F. v. M., fragm. III, 116 (1862)	—	—	—	—	—	N.A.	B.f.III, 138	M.fr.III, 116.
<i>M. Baxteri</i> ; Benth. Fl. Austr. III, 138 (1866)	W.A.	—	—	—	—	—	B.f.III, 138	—
<i>M. symphyocarpa</i> ; F. v. M. in Transact. phil. Inst. Viet. III, 44 (1858)	—	—	—	—	—	Q. N.A.	B.f.III, 138	—
<i>M. angustifolia</i> ; Gaertner, de fruct. I, 172, t. 35 (1788)	—	—	—	—	—	Q.	B.f.III, 139	—
<i>M. pauciflora</i> ; Turczaninow in Bull. Mosc. XX, 166 (1847)	—	—	—	—	N.S.W.	—	B.f.III, 139	—
<i>M. squarrosa</i> ; Donn, Hort. Cantabr. ed. sec. 101 (1800)	—	S.A.	T.	V.	N.S.W.	—	B.f.III, 139	M.fr.X, 55.
<i>M. adnata</i> ; Turczaninow in Bullet. de St. Petersburg. 425 (1852)	W.A.	—	—	—	—	—	B.f.III, 140	M.fr.III, 117; VIII, 184.
<i>M. linariifolia</i> ; Smith in Transact. Linn. Soc. III, 278 (1797)	—	—	—	—	N.S.W.	Q.	B.f.III, 140	M.fr.X, 55.
<i>M. trichostachya</i> ; Lindley in Mitch. Trop. Austr. 277 (1848)	—	S.A.	—	—	—	Q. N.A.	—	—
<i>M. radula</i> ; Lindley, Bot. Regist. XXV, App. 8 (1839)	W.A.	—	—	—	—	—	B.f.III, 141	M.fr.X, 55.
<i>M. pulchella</i> ; R. Brown in Ait. hort. Kew. sec. ed. IV, 414 (1812)	W.A.	—	—	—	—	—	B.f.III, 141	M.fr.III, 120.
<i>M. conferta</i> ; Benth. Fl. Austr. III, 142 (1866)	W.A.	—	—	—	—	—	B.f.III, 142	—
<i>M. Leucadendron</i> ; Linné, mantiss. 105 (1767)	W.A.	—	—	—	N.S.W.	Q. N.A.	B.f.III, 142	M.fr.IV, 55 ... As., P.

<i>M. lasiantha</i> ; F. v. M., fragm. III, 115 (1862)	—	—	—	—	—	—	—	—	N.S.W.	Q.	N.S.W.	R. d. III, 143	M. d. III, 115.	
<i>M. gemistifolia</i> ; Smith in Transact. Linn. Soc. III, 277 (1797)	—	—	—	—	—	—	—	—	N.S.W.	Q.	N.S.W.	R. d. III, 143	M. d. I, 15; X, 53.	
<i>M. stypheloides</i> ; Smith in Transact. Linn. Soc. III, 276 (1797)	—	—	—	—	—	—	—	—	N.S.W.	Q.	—	R. d. III, 144	M. d. I, 15.	
<i>M. Hoegghii</i> ; Endlicher in Hæg. enum. 48 (1837)	—	—	—	—	—	—	W.A.	—	—	—	—	R. d. III, 144	—	
<i>M. acroclora</i> ; F. v. M. in Austr. Jour. of Phasn. I, 276 (1896)	—	—	—	—	—	—	W.A.	—	—	—	—	—	—	
<i>M. distichocla</i> ; F. v. M., fragm. III, 153 (1863)	—	—	—	—	—	—	—	—	—	—	N.S.W.	R. d. III, 144	M. d. III, 153.	
<i>M. leucophylla</i> ; F. v. M., fragm. III, 115 (1863)	—	—	—	—	—	—	—	—	—	—	N.S.W.	R. d. III, 145	M. d. III, 115.	
<i>M. parviflora</i> ; Lindley, Bot. Regist. XXV, App. 8 (1839)	—	—	—	—	—	—	W.A.	S.A.	—	V.	N.S.W.	Q.	R. d. III, 145	M. d. III, 115.
<i>M. Deanei</i> ; F. v. M. in proceed. L. Soc. of N.S.W. sec. sec. I, 1106 (1896)	—	—	—	—	—	—	—	—	—	—	N.S.W.	—	—	—
<i>M. laxiflora</i> ; Turczaninow in Bull. de St. Petersburg. 421 (1892)	—	—	—	—	—	—	W.A.	—	—	—	—	R. d. III, 145	M. d. VIII, 184; X, 53.	
<i>M. armillaris</i> ; Smith in Transact. Linn. Soc. III, 277 (1797)	—	—	—	—	—	—	—	—	—	V.	N.S.W.	Q.	R. d. III, 146	—
<i>M. cylindrica</i> ; R. Brown in Benth. Fl. Austr. III, 146 (1866)	—	—	—	—	—	—	—	S.A.	—	—	—	—	—	—
<i>M. hantziæ</i> ; Turczaninow in Bull. Mosc. XX, 165 (1847)	—	—	—	—	—	—	W.A.	—	—	—	—	R. d. III, 146	M. d. III, 116.	
<i>M. subulata</i> ; Turczaninow in Bull. de St. Petersburg. 422 (1892)	—	—	—	—	—	—	W.A.	—	—	—	—	R. d. III, 146	M. d. III, 119; VIII, 184.	
<i>M. glaberrima</i> ; F. v. M., fragm. III, 119 (1862)	—	—	—	—	—	—	W.A.	—	—	—	—	R. d. III, 147	M. d. III, 119.	
<i>M. raphiophylla</i> ; Schauer in Lehm. pl. Preiss. I, 143 (1844)	—	—	—	—	—	—	W.A.	—	—	—	—	R. d. III, 147	—	
<i>M. cymbifolia</i> ; Benth. Fl. Austr. III, 145 (1866)	—	—	—	—	—	—	W.A.	—	—	—	—	R. d. III, 148	—	
<i>M. cuticularia</i> ; Labillardière, Nov. Holl. pl. spec. II, 30, t. 171 (1806)	—	—	—	—	—	—	W.A.	—	—	—	—	R. d. III, 148	—	
<i>M. sparsiflora</i> ; Turczaninow in Bull. Mosc. XX, 167 (1847)	—	—	—	—	—	—	W.A.	—	—	—	—	R. d. III, 148	—	
<i>M. calycina</i> ; R. Brown in Ait. hort. Kew. sec. ed. IV, 416 (1812)	—	—	—	—	—	—	W.A.	—	—	—	—	R. d. III, 148	M. d. VIII, 184.	
<i>M. cordata</i> ; Turczaninow in Bull. de St. Petersburg. 418 (1892)	—	—	—	—	—	—	W.A.	—	—	—	—	R. d. III, 149	M. d. VIII, 184.	
<i>M. globifera</i> ; R. Brown in Ait. hort. Kew. sec. ed. IV, 411 (1812)	—	—	—	—	—	—	W.A.	—	—	—	—	R. d. III, 149	—	
<i>M. megacephala</i> ; F. v. M., fragm. III, 117 (1862)	—	—	—	—	—	—	W.A.	—	—	—	—	R. d. III, 149	M. d. III, 117.	
<i>M. necephala</i> ; F. v. M., fragm. III, 113 (1862)	—	—	—	—	—	—	W.A.	—	—	—	—	R. d. III, 150	M. d. III, 113.	
<i>M. Oldfieldii</i> ; F. v. M., fragm. III, 118 (1862)	—	—	—	—	—	—	W.A.	—	—	—	—	R. d. III, 150	—	
<i>M. uncinata</i> ; R. Brown in Ait. hort. Kew. sec. ed. IV, 414 (1812)	—	—	—	—	—	—	W.A.	S.A.	—	V.	N.S.W.	Q.	R. d. III, 150	M. d. III, 118.
<i>M. concreta</i> ; F. v. M., fragm. III, 116 (1862)	—	—	—	—	—	—	W.A.	—	—	—	—	R. d. III, 151	M. d. III, 118.	
<i>M. filifolia</i> ; F. v. M., fragm. III, 119 (1862)	—	—	—	—	—	—	W.A.	—	—	—	—	R. d. III, 151	M. d. III, 119.	
<i>M. glomerata</i> ; F. v. M., Rep. Babb. Exped. 5 (1858)	—	—	—	—	—	—	W.A.	S.A.	—	N.S.W.	Q.	N.S.W.	R. d. III, 151	—
<i>M. pentagona</i> ; Labillardière, Nov. Holl. pl. spec. II, 27, t. 166 (1806)	—	—	—	—	—	—	W.A.	—	—	—	—	R. d. III, 152	M. d. VIII, 184.	
<i>M. ciliosa</i> ; Turczaninow in Bull. Soc. Mosc. XXXV, 326 (1892)	—	—	—	—	—	—	W.A.	—	—	—	—	R. d. III, 152	—	
<i>M. polycephala</i> ; Benth. Fl. Austr. III, 152 (1866)	—	—	—	—	—	—	W.A.	—	—	—	—	R. d. III, 152	M. d. VIII, 184.	
<i>M. spiritalata</i> ; Schauer in Lehm. pl. Preiss. I, 134 (1844)	—	—	—	—	—	—	W.A.	—	—	—	—	R. d. III, 153	M. d. VIII, 184.	
<i>M. eriantha</i> ; Benth. Fl. Austr. III, 153 (1866)	—	—	—	—	—	—	W.A.	—	—	—	—	R. d. III, 153	—	
<i>M. subtrigona</i> ; Schauer in Lehm. pl. Preiss. I, 139 (1844)	—	—	—	—	—	—	W.A.	—	—	—	—	R. d. III, 153	—	
<i>M. seriata</i> ; Lindley, Bot. Regist. XXV, App. 8 (1839)	—	—	—	—	—	—	W.A.	—	—	—	—	R. d. III, 154	—	
<i>M. scabra</i> ; R. Brown in Ait. hort. Kew. sec. ed. IV, 414 (1812)	—	—	—	—	—	—	W.A.	—	—	—	—	R. d. III, 154	M. d. VIII, 184.	
<i>M. urceolaris</i> ; F. v. M. in Benth. Fl. Austr. III, 154 (1866)	—	—	—	—	—	—	W.A.	—	—	—	—	R. d. III, 154	—	
<i>M. trichophylla</i> ; Lindley, Bot. Regist. XXV, App. 8 (1839)	—	—	—	—	—	—	W.A.	—	—	—	—	R. d. III, 155	M. d. III, 114.	
<i>M. holosericea</i> ; Schauer in Lehm. pl. Preiss. I, 139 (1844)	—	—	—	—	—	—	W.A.	—	—	—	—	R. d. III, 155	—	
<i>M. squamea</i> ; Labillardière, Nov. Holl. pl. spec. II, 28, t. 168 (1806)	—	—	—	—	—	—	—	S.A.	T.	V.	N.S.W.	—	R. d. III, 155	—
<i>M. densa</i> ; R. Brown in Ait. hort. Kew. sec. ed. IV, 415 (1812)	—	—	—	—	—	—	W.A.	—	—	—	—	R. d. III, 156	M. d. VIII, 184.	
<i>M. thymoides</i> ; Labillardière, Nov. Holl. pl. spec. II, 27, t. 167 (1806)	—	—	—	—	—	—	W.A.	—	—	—	—	R. d. III, 156	—	
<i>M. striata</i> ; Labillardière, Nov. Holl. pl. spec. II, 28, t. 165 (1806)	—	—	—	—	—	—	W.A.	—	—	—	—	R. d. III, 157	—	
<i>M. polygaloides</i> ; Schauer in Lehm. pl. Preiss. I, 142 (1844)	—	—	—	—	—	—	W.A.	—	—	—	—	R. d. III, 157	—	
<i>M. incana</i> ; R. Brown in Bot. Regist. t. 410 (1819)	—	—	—	—	—	—	W.A.	—	—	—	—	R. d. III, 157	—	
<i>M. nodosa</i> ; Smith in Transact. Linn. Soc. III, 276 (1797)	—	—	—	—	—	—	—	—	—	—	N.S.W.	Q.	R. d. III, 158	—
<i>M. pungens</i> ; Schauer in Lehm. pl. Preiss. I, 135 (1844)	—	—	—	—	—	—	W.A.	—	—	—	—	R. d. III, 158	M. d. VIII, 184.	
<i>M. ericifolia</i> ; Smith in Transact. Linn. Soc. III, 276 (1797)	—	—	—	—	—	—	—	S.A.	T.	V.	N.S.W.	Q.	R. d. III, 159	—
<i>M. pustulata</i> ; J. Hooker in Lond. Journ. VI, 476 (1844)	—	—	—	—	—	—	—	S.A.	T.	V.	N.S.W.	—	R. d. III, 160	—

M. viminea; Lindley, Bot. Regist. XXV, App. 8 (1839)	W.A.	—	—	—	—	—	—	B.fl.III,159
M. microphylla; Smith in Rees, Cyclop. XXIII (1812)	W.A.	—	—	—	—	—	—	B.fl.III,159 M.fr.III,116.
M. tenella; Benth. Fl. Austr. III, 160 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,160
M. leiocarpa; F. v. M., fragm. X, 55 (1876)	W.A.	—	—	—	—	—	—	M.fr.X,55.
M. leiopyxis; F. v. M. in Benth. Fl. Austr. III, 160 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,160
M. acerosa; Schauer in Lehm. pl. Preiss. I, 137 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,160 M.fr.III,118.
M. pauperiflora; F. v. M., fragm. III, 116 (1862)	W.A.	—	—	—	—	—	—	B.fl.III,161 M.fr.III,116.
M. aspalathoides; Schauer in Lehm. pl. Preiss. I, 140 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,161 M.fr.VIII,184.
M. cucullata; Turczaninow in Bullet. de St. Petersb. 424 (1852)	W.A.	—	—	—	—	—	—	B.fl.III,161 M.fr.VIII,184.
M. minutifolia; F. v. M. in Transact. phil. Soc. Vict. III, 45 (1858)	—	—	—	—	—	—	Q. N.A.	B.fl.III,162
M. foliolosa; Cunningham in Benth. Fl. Austr. III, 162 (1866)	—	—	—	—	—	—	Q.	B.fl.III,162
M. micromera; Schauer in Lehm. pl. Preiss. I, 146 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,162
M. thuyoides; Turczaninow in Bull. Soc. Mosc. XX, 167 (1847)	W.A.	—	—	—	—	—	—	B.fl.III,162 M.fr.III,114.
M. tamariscina; Hooker in Mitch. Trop. Austr. 262 (1848)	—	—	—	—	—	—	Q.	B.fl.III,163 M.fr.III,115.
M. quadrifaria; F. v. M. in Wing's S. Sc. Rec. April (1886)	W.A. S.A.	—	—	—	—	—	—	—

CONOTHAMNUS; Lindley, Bot. Regist. XXV, App. 11 (1839). (Trichobasis.)

C. trinervis; Lindley, Bot. Regist. XXV, App. IX (1839)	W.A.	—	—	—	—	—	—	B.fl.III,163
C. divaricatus; Benth. Fl. Austr. III, 164 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,164 M.fr.VIII,184.

BEAUFORTIA; R. Brown in Ait. hort. Kew. sec. ed. IV, 418 (1812). (Schizopleura.)

B. sparsa; R. Brown in Ait. hort. Kew. sec. ed. IV, 419 (1812)	W.A.	—	—	—	—	—	—	B.fl.III,165
B. decussata; R. Brown in Ait. hort. Kew. sec. ed. IV, 418 (1812)	W.A.	—	—	—	—	—	—	B.fl.III,166
B. squarrosa; Schauer in nov. act. acad. Caes. XXI, 11 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,166
B. orbifolia; F. v. M., fragm. III, 110 (1863)	W.A.	—	—	—	—	—	—	B.fl.III,166 M.fr.III,110.
B. anisandra; Schauer in nov. act. acad. Caes. XXI, 13, t. I (1844)	W.A.	—	—	—	—	—	—	B.fl.III,167
B. macrostemon; Lindley, Bot. Regist. XXV, App. 10 (1839)	W.A.	—	—	—	—	—	—	B.fl.III,167
B. heterophylla; Turczaninow in Bullet. de St. Petersb. 427 (1852)	W.A.	—	—	—	—	—	—	B.fl.III,167 M.fr.VIII,184.
B. Schaueri; Preiss in nov. act. acad. Caes. XXI, 18 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,168
B. purpurea; Lindley, Bot. Regist. XXV, App. X, t. 3 (1839)	W.A.	—	—	—	—	—	—	B.fl.III,168
B. Dampieri; Cunningham in Bot. Mag. t. 3272 (1833)	W.A.	—	—	—	—	—	—	B.fl.III,168 M.fr.III,111.
B. elegans; Schauer in nov. act. acad. Caes. XXI, 20 (1844)	W.A.	—	—	—	—	—	N.A.	B.fl.III,169
B. interstans; F. v. M., fragm. X, 30 (1876)	W.A.	—	—	—	—	—	—	M.fr.X,30.
B. micrantha; Schauer in nov. act. acad. Caes. XXI, 22 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,169 M.fr.VIII,184.

REGELIA; Schauer in Linnaea, XVII, 243 (1843).

R. grandiflora; Benth. Fl. Austr. III, 170 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,170 M.fr.VIII,184; IX,46; XI,11.
R. ciliata; Schauer in nov. act. acad. Caes. XXI, 11 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,171 M.fr.I,225; XI,11.
R. inops; Schauer in Lehm. pl. Preiss. II, 224 (1847)	W.A.	—	—	—	—	—	—	B.fl.III,171 M.fr.XI,11.

PHYMATOCARPUS; F. v. M., fragm. III, 120 (1862).

P. porphyrocephalus; F. v. M., fragm. III, 121 (1862)	W.A.	—	—	—	—	—	—	B.fl.III,172 M.fr.III,121.
P. Maxwelli; F. v. M., fragm. IX, 45 (1875)	W.A.	—	—	—	—	—	—	M.fr.IX,45.

CALOTHAMNUS; Labillardière, Nov. Holl. pl. specim. II, 25 (1806). (Billotia.)

C. pachystachyus; Benth. Fl. Austr. III, 173 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,173
C. longissimus; F. v. M., fragm. III, 112 (1862)	W.A.	—	—	—	—	—	—	B.fl.III,174 M.fr.III,112.
C. blepharanthus; F. v. M., fragm. III, 112 (1862)	W.A.	—	—	—	—	—	—	B.fl.III,174 M.fr.X,31&55.
C. sanguineus; Labillardière, Nov. Holl. pl. spec. II, 25, t. 164 (1806)	W.A.	—	—	—	—	—	—	B.fl.III,174 M.fr.X,31.
C. torulosus; Schauer in nov. act. acad. Caes. XXI, 25 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,175 M.fr.X,31.
C. gibbosus; Benth. Fl. Austr. III, 175 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,175
C. gracilis; R. Brown in Ait. hort. Kew. sec. ed. IV, 418 (1812)	W.A.	—	—	—	—	—	—	B.fl.III,175

C. blepharospermus; F. v. M., fragm. III, 111 (1862)	W.A.	—	—	—	—	—	B.fl.III,175	M.fr.III,111.
C. Gilesii; F. v. M., fragm. X, 31 (1876)	W.A.	—	—	—	—	—	—	M.fr.X,31.
C. chrysanthus; F. v. M., fragm. III, 112 (1862)	W.A.	—	—	—	—	—	B.fl.III,176	M.fr.III,112.
C. Oldfieldii; F. v. M., fragm. III, 113 (1862)	W.A.	—	—	—	—	—	B.fl.III,176	M.fr.III,113.
C. planifolius; Lehmann, delect. sem. hort. bot. Hamb. 7 (1842)	W.A.	—	—	—	—	—	B.fl.III,176	M.fr.X,31.
C. lateralis; Lindley, Bot. Regist. XXV, App. 9 (1839)	W.A.	—	—	—	—	—	B.fl.III,177	—
C. affinis; Turczaninow in Bullet. de St. Petersb. 428 (1852)	W.A.	—	—	—	—	—	B.fl.III,177	M.fr.III,111; VIII,184.
C. Preissii; Schauer in nov. act. acad. Caes. XXI, 31 (1844)	W.A.	—	—	—	—	—	B.fl.III,177	—
C. Schaueri; Lehmann, delect. sem. hort. bot. Hamb. 7 (1842)	W.A.	—	—	—	—	—	B.fl.III,178	—
C. Lehmanni; Schauer in nov. act. acad. Caes. XXI, 31 (1844)	W.A.	—	—	—	—	—	B.fl.III,178	—
C. villosus; R. Brown in Ait. hort. Kew. sec. ed. IV, 418 (1812)	W.A.	—	—	—	—	—	B.fl.III,178	—
C. pinifolius; F. v. M., fragm. III, 153 (1863)	W.A.	—	—	—	—	—	B.fl.III,178	M.fr.III,153.
C. rupestris; Schauer in nov. act. acad. Caes. XXI, 26 (1844)	W.A.	—	—	—	—	—	B.fl.III,179	—
C. quadrifidus; R. Brown in Ait. hort. Kew. sec. ed. IV, 418 (1812)	W.A.	—	—	—	—	—	B.fl.III,179	M.fr.X,31.
C. asper; Turczaninow in Bull. Mosc. XXII, 25 (1849)	W.A.	—	—	—	—	—	B.fl.III,180	M.fr.X,31.
C. homalophyllus; F. v. M., fragm. III, 111 (1863)	W.A.	—	—	—	—	—	B.fl.III,180	M.fr.X,31.

LAMARCHEA; Gaudichaud in Freyc. Voy. Bot. 483, t. 110 (1826).

L. hakeaeifolia; Gaudichaud in Freyc. Voy. 484, t. 110 (1826)	W.A.	—	—	—	—	—	B.fl.III,123	—
--	------	---	---	---	---	---	--------------	---

EREMAEA; Lindley, Bot. Regist. XXV, App. 11 (1839).

E. fimbriata; Lindley, Bot. Regist. XXV, App. 11 (1839)	W.A.	—	—	—	—	—	B.fl.III,181	M.fr.II,30; XI,11.
E. acutifolia; F. v. M., fragm. II, 30 (1860)	W.A.	—	—	—	—	—	B.fl.III,181	M.fr.II,30.
E. violacea; F. v. M., fragm. XI, 10 (1878)	W.A.	—	—	—	—	—	—	M.fr.XI,10.
E. pilosa; Lindley, Bot. Regist. XXV, App. 11 (1839)	W.A.	—	—	—	—	—	B.fl.III,182	M.fr.II,30; XI,11.
E. ebracteata; F. v. M., fragm. II, 29 (1860)	W.A.	—	—	—	—	—	B.fl.III,182	M.fr.XI,11.
E. beaufortioides; Benth, Fl. Austr. III, 182 (1866)	W.A.	—	—	—	—	—	B.fl.III,182	M.fr.XI,11.

ANGOPHORA; Cavanilles, icon. IV, 21, t. 331 (1797).

A. cordifolia; Cavanilles, icon. IV, 21, t. 338 (1797)	—	—	—	—	N.S.W.	—	B.fl.III,183	—
A. subvelutina; F. v. M., fragm. I, 31 (1858)	—	—	—	—	N.S.W.	Q.	B.fl.III,184	M.fr.I,31.
A. intermedia; De Candolle, prodr. III, 222 (1828)	—	—	—	—	V.	N.S.W.	Q.	B.fl.III,184
A. lanceolata; Cavanilles, icon. IV, 22, t. 339 (1797)	—	—	—	—	N.S.W.	Q.	B.fl.III,184	—

EUCALYPTUS; L'Heritier, sertum Anglicum 18 (1788). (Eudesmia, Symphyomyrtus.)

I. RENANTHERAE.

E. stellulata; Sieber in De Candolle, prodr. III, 217 (1828)	—	—	—	V.	N.S.W.	—	B.fl.III,200	M.fr.II,45; IX,172.
E. paniciflora; Sieber in Sprengel, cur. poster. 195 (1827)	—	S.A.	T.	V.	N.S.W.	—	B.fl.III,201	M.fr.II,52.
E. regnans; F. v. M. in Report Acclim. Soc. Vict. 20 (1870)	—	—	T.	V.	N.S.W.	—	—	—
E. amygdalina; Labillardiere, Nov. Holl. pl. spec. II, 14, t. 154 (1806)	—	S.A.	T.	V.	N.S.W.	—	B.fl.III,202	M.fr.II,53; VII,42&44.
E. coccifera; J. Hooker in Lond. Journ. of Bot. VI, 477 (1847)	—	—	T.	—	—	—	B.fl.III,204	—
E. obliqua; L'Heritier, sertum Anglicum 18, t. 20 (1788)	—	S.A.	T.	V.	N.S.W.	—	B.fl.III,204	M.fr.II,44&171; XI,33.
E. stricta; Sieber in Spreng. syst. veg. cur. post. 195 (1827)	—	—	—	—	N.S.W.	—	B.fl.III,217	M.fr.XI,44.
E. virgata; Sieber in Spreng. syst. veg. cur. post. 195 (1827)	—	—	—	—	N.S.W.	—	B.fl.III,202	M.fr.XI,38.
E. Planchoniana; F. v. M., fragm. XI, 43 (1878)	—	—	—	—	N.S.W.	Q.	—	M.fr.XI,43.
E. Baileyana; F. v. M., fragm. XI, 37 (1878)	—	—	—	—	N.S.W.	Q.	—	M.fr.XI,37.
E. sepulcralis; F. v. M., Eucalyptographia eighth dec. (1882)	W.A.	—	—	—	—	—	—	—
E. buprestium; F. v. M., fragm. III, 57 (1862)	W.A.	—	—	—	—	—	B.fl.III,205	M.fr.III,57.
E. marginata; Donn, hort. Cantabr. sec. edit. 101 (1800)	W.A.	—	—	—	—	—	B.fl.III,209	M.fr.II,40&174; VII,43; XI,38.
E. santalifolia; F. v. M. in Transact. Vict. Inst. I, 35 (1854)	W.A.	S.A.	—	—	—	—	B.fl.III,206	—
E. macrorrhyncha; F. v. M., First general Report, 12 (1853)	—	S.A.	—	V.	N.S.W.	—	B.fl.III,207	M.fr.XI,45.
E. capitellata; Smith in White, Journ. Voy. N. S. Wales 216 (1790)	—	S.A.	—	V.	N.S.W.	—	B.fl.III,206	M.fr.II,173; XI,38.

E. eugenioides; Sieber in Spreng. cur. poster. 195 (1827)	—	—	—	V.	N.S.W.	Q.	—	—	M.fr.II,52,64.
E. piperita; Smith in White, Journ. Voy. N. S. Wales 226 (1790)	—	—	—	V.	N.S.W.	Q.	—	B.fl.III,207	M.fr.II,173; VII,42,44; XI,33.
E. pilularis; Smith in Transact. Linn. Soc. III, 284 (1797)	—	—	—	V.	N.S.W.	Q.	—	B.fl.III,208	M.fr.II,61&172; VII,44; XI,33.
E. triantha; Link, enum. pl. hort. bot. Berol. II, 20 (1822)	—	—	—	—	N.S.W.	Q.	—	—	M.fr.VII,42.
E. haemastoma; Smith in Transact. Linn. Soc. III, 285 (1797)	—	—	—	T.	V.	N.S.W.	Q.	B.fl.III,212	M.fr.II,51&175; VII,43; XI,38.
E. Sieberiana; F. v. M., Eucalyptographia sec. decade (1879)	—	—	—	S.A.	T.	V.	N.S.W.	B.fl.III,202	M.fr.XI,37.
E. microcorys; F. v. M., fragm. II, 50 (1860)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,212	M.fr.II,50.

II. PORANTHERAE.

E. paniculata; Smith in Transact. Linn. Soc. III, 287 (1797)	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,211	M.fr.II,52,174; VII,43.	
E. Leucoxydon; F. v. M. in Transact. Vict. Inst. I, 33 (1854)	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,209	M.fr.II,60&175.	
E. melliodora; Cunningham in Walp. rep. bot. syst. II, 924 (1843)	...	...	...	—	—	—	V.	N.S.W.	Q.	—	B.fl.III,210	M.fr.VII,43.	
E. polyanthema; Schauer in Walpers, rep. bot. syst. II, 924 (1843)	...	...	...	—	—	—	V.	N.S.W.	Q.	—	B.fl.III,213	M.fr.VII,44.	
E. populifolia; Hooker, icon. plant. 279 (1852)	...	...	...	...	—	—	—	N.S.W.	Q.	N.A.	B.fl.III,214&243	M.fr.II,67.	
E. ochrophloia; F. v. M., fragm. XI, 36 (1878)	...	...	...	...	—	—	—	N.S.W.	Q.	—	—	M.fr.XI,36.	
E. gracilis; F. v. M. in Transact. Vict. Inst. I, 35 (1854)	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,222	M.fr.II,55,57; VII,43; VIII,184; XI,36.
E. uncinata; Turczaninow in Bull. de Mosc. XXII, 23 (1849)	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,216	
E. odorata; Behr in Schlecht. Linnæa XX, 657 (1847)	...	...	...	...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,215	M.fr.II,66.
E. largiflorens; F. v. M. in Transact. Vict. Inst. I, 34 (1854)	...	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.III,215	M.fr.VII,43.
E. Behriana; F. v. M. in Transact. Vict. Inst. I, 34 (1854)	...	...	...	...	—	S.A.	—	V.	—	—	—	B.fl.III,214	
E. hemiphloia; F. v. M., fragm. II, 62 (1860)	...	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,216	M.fr.II,62.
E. pruinosa; Schauer in Walp. rep. bot. syst. II, 926 (1843)	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.III,213	M.fr.III,132.
E. melanophloia; F. v. M. in Journ. Linn. Soc. III, 93 (1858)	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.III,220	M.fr.VII,43.

III. PARALLELANATHERAE.

E. Cloeziana; F. v. M., fragm. XI, 44 (1878)...	...	...	...	...	...	—	—	—	Q.	—	—	M.fr.XI,44.
E. Howittiana; F. v. M. in Wing's S. Sc. Record, Aug. (1882)	...	...	...	...	...	—	—	—	Q.	—	—	
E. drepanophylla; F. v. M. in Benth. Fl. Austr. III, 221 (1866)	...	...	...	...	...	—	—	—	Q.	—	B.fl.III,221	
E. crebra; F. v. M. in Journ. Linn. Soc. III, 87 (1858)	...	...	...	...	...	—	—	—	N.S.W.	Q.	N.A.	B.fl.III,221 M.fr.XI,44.
E. Staigeriana; F. v. M. in Bailey's synops. Q. L. Flora 176 (1883)	...	...	...	...	...	—	—	—	—	Q.	—	
E. decipiens; Endlicher in Hueg. enum. 49 (1837)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.III,218	
E. concolor; Schauer in Lehm. pl. Preiss. I, 129 (1844)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.III,247	M.fr.II,42.
E. siderophloia; Benth. Fl. Austr. III, 220 (1866)	...	...	...	...	...	—	—	—	N.S.W.	Q.	—	B.fl.III,220 M.fr.II,61; VII,44.
E. microtheca; F. v. M. in Journ. Linn. Soc. III, 87 (1858)	...	...	...	...	...	—	S.A.	—	N.S.W.	Q.	N.A.	B.fl.III,223 M.fr.II,40; VII,42; X,100; XI,14.
E. Raveretiana; F. v. M., fragm. X, 99 (1877)	...	...	...	...	...	—	—	—	—	Q.	—	M.fr.X,99.
E. brachyandra; F. v. M. in Journ. Linn. Soc. III, 97 (1858)	...	...	...	...	...	—	—	—	—	N.A.	B.fl.III,223	M.fr.X,100.
E. pachyphylla; F. v. M. in Journ. Linn. Soc. III, 98 (1858)	...	...	...	...	...	—	S.A.	—	—	Q.	N.A.	B.fl.III,237
E. pyriformis; Turczaninow in Bull. Mosc. XXII, 22 (1849)	...	...	...	...	W.A.	S.A.	—	—	—	—	B.fl.III,226	M.fr.II,32; X,5&84.
E. macrocarpa; Hooker, icones plant. t. 405-407 (1842)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.III,224	M.fr.II,41; XI,38.
E. Preissiana; Schauer in Lehm. pl. Preiss. I, 131 (1844)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.III,232	M.fr.II,38; VII,42.
E. alpina; Lindley in Mitch. Three Exped. II, 175 (1838)	...	...	...	...	...	—	—	—	V.	—	B.fl.III,225	M.fr.II,68; VII,42.
E. Globulus; Labillardière, Voy. I, 153, t. 13 (1799)	...	...	...	...	...	—	—	—	T.	V.	N.S.W.	B.fl.III,225 M.fr.II,68.
E. megacarpa; F. v. M., fragm. II, 70 (1860)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.III,232	M.fr.VII,43.
E. cosmophylla; F. v. M. in Transact. Vict. Inst. 32 (1854)	...	...	...	...	...	—	S.A.	—	—	—	B.fl.III,225	M.fr.VII,43.
E. cordata; Labillardière, Nov. Holl. pl. spec. II, 13, t. 152 (1806)	...	...	...	...	...	—	—	—	T.	—	—	B.fl.III,224
E. urnigera; J. Hooker in Lond. Journ. of Bot. VI, 477 (1847)	...	...	...	...	...	—	—	—	T.	—	—	B.fl.III,227 M.fr.II,43.
E. longifolia; Link, enum. pl. hort. bot. Berol. II, 29 (1822)	...	...	...	...	...	—	—	—	V.	N.S.W.	—	B.fl.III,226 M.fr.II,50,147; VII,43.
E. erythronema; Turczaninow in Bull. de l'acad. de St. Petersb. 415 (1852)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.III,227	M.fr.VIII,184.
E. caesia; Benth. Fl. Austr. III, 227 (1866)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.III,227	M.fr.XI,40.
E. tetraptera; Turczaninow in Bull. de Mosc. XXII, 22 (1849)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.III,228	M.fr.II,34; IV,172; VIII,184
E. miniata; A. Cunningham in Walp. rep. bot. syst. II, 925 (1843)	...	...	...	...	...	—	—	—	—	Q.	N.A.	B.fl.III,228 M.fr.XI,38&42.
E. phoenicea; F. v. M. in Journ. Linn. Soc. III, 91 (1858)...	...	...	...	...	...	—	—	—	—	—	N.A.	B.fl.III,251

<i>E. robusta</i> ; Smith, Specim. Bot. New Holl. 40, t. 13 (1793) ...	...	...	—	—	—	N.S.W.	Q.	—	B.fl.III,228	M.fr.II,43&171; VII,44.
<i>E. botryoides</i> ; Smith in Transact. Linn. Soc. III, 286 (1797) ...	...	...	—	—	—	V. N.S.W.	Q.	—	B.fl.III,229	M.fr.II,48&175; VII,43.
<i>E. gonicalyx</i> ; F. v. M. in Nederl. Kruidk. Arch. IV, 134 (1859) ...	...	...	—	S.A.	—	V. N.S.W.	—	—	B.fl.III,229	M.fr.II,48; IV,52.
<i>E. pallidifolia</i> ; F. v. M., fragm. III, 131 (1863) ...	...	...	—	—	—	—	—	N.A.	B.fl.III,236	M.fr.III,131.
<i>E. iucrassata</i> ; Labillardière, Nov. Holl. pl. spec. II, 12, t. 150 (1806) ...	...	...	W.A.	S.A.	—	V. N.S.W.	—	—	B.fl.III,231	M.fr.II,59.
<i>E. oleosa</i> ; F. v. M. in Nederl. Kruidk. Arch. IV, 127 (1859) ...	...	...	W.A.	S.A.	—	V. N.S.W.	—	—	B.fl.III,248	M.fr.II,56; XI,14.
<i>E. goniantha</i> ; Turczaninow in Bullet. de Mosc. XX, 163 (1847) ...	...	...	W.A.	—	—	—	—	—	B.fl.III,248	
<i>E. falcata</i> ; Turczaninow in Bullet. de Mosc. XX, 163 (1847) ...	...	...	W.A.	—	—	—	—	—	B.fl.III,248	
<i>E. salmonophloia</i> ; F. v. M., fragm. XI, 11 (1878) ...	...	...	W.A.	—	—	—	—	—	—	M.fr.XI,11.
<i>E. leptopoda</i> ; Benth. Fl. Austr. III, 238 (1866) ...	...	...	W.A.	—	—	—	—	—	B.fl.III,238	M.fr.XI,11&13.
<i>E. salubris</i> ; F. v. M., fragm. X, 54 (1876) ...	...	...	W.A.	—	—	—	—	—	—	M.fr.XI,12.
<i>E. angustissima</i> ; F. v. M., fragm. IV, 25 (1863) ...	...	...	W.A.	S.A.	—	—	—	—	B.fl.III,238	M.fr.IV,25.
<i>E. Doratoxylon</i> ; F. v. M., fragm. II, 55 (1860) ...	...	...	W.A.	—	—	—	—	—	B.fl.III,249	M.fr.II,55.
<i>E. decurva</i> ; F. v. M., fragm. III, 130 (1863) ...	...	...	W.A.	—	—	—	—	—	B.fl.III,249	M.fr.III,130.
<i>E. Cooperiana</i> ; F. v. M., fragm. XI, 83 (1880) ...	...	...	W.A.	—	—	—	—	—	—	M.fr.XI,83.
<i>E. corynocalyx</i> ; F. v. M., fragm. II, 43 (1860) ...	...	...	—	S.A.	—	V.	—	—	B.fl.III,218	M.fr.VII,43.
<i>E. gomphocephala</i> ; De Candolle, prodr. III, 220 (1828) ...	...	...	W.A.	—	—	—	—	—	B.fl.III,231	M.fr.II,36; V,14.
<i>E. Oldfieldii</i> ; F. v. M., fragm. II, 37 (1860) ...	...	...	W.A.	S.A.	—	—	—	—	B.fl.III,237	M.fr.II,37.
<i>E. orbifolia</i> ; F. v. M., fragm. V, 50 (1865) ...	...	...	W.A.	—	—	—	—	—	B.fl.III,238	M.fr.V,50.
<i>E. diversicolor</i> ; F. v. M., fragm. III, 131 (1863) ...	...	...	W.A.	—	—	—	—	—	B.fl.III,251	M.fr.III,131; V,14; VII,42.
<i>E. patellaris</i> ; F. v. M. in Journ. Linn. Soc. III, 84 (1858) ...	...	...	—	—	—	—	—	N.A.	B.fl.III,244	
<i>E. tessellaris</i> ; F. v. M. in Journ. Linn. Soc. III, 88 (1858) ...	...	...	—	S.A.	—	—	N.S.W.	Q. N.A.	B.fl.III,251	
<i>E. clavigera</i> ; A. Cunningham in Walp. rep. bot. syst. II, 926 (1843) ...	...	...	—	—	—	—	—	Q. N.A.	B.fl.III,250	... P.
<i>E. ferruginea</i> ; Schauer in Walp. repert. II, 926 (1843) ...	...	...	—	—	—	—	—	—	N.A.	B.fl.III,254
<i>E. grandifolia</i> ; R. Brown in Benth. Fl. Austr. III, 250 (1866) ...	...	...	—	—	—	—	—	Q. N.A.	B.fl.III,250	
<i>E. aspera</i> ; F. v. M. in Journ. Linn. Soc. III, 95 (1858) ...	...	...	—	—	—	—	—	—	N.A.	B.fl.III,250
<i>E. patens</i> ; Benth. Fl. Austr. III, 247 (1866) ...	...	...	W.A.	—	—	—	—	—	B.fl.III,247	
<i>E. Todtiana</i> ; F. v. M. in Wing's S. Sc. Record II, 171 (1882) ...	...	...	W.A.	—	—	—	—	—	—	
<i>E. vernicosa</i> ; J. Hooker in Lond. Journ. of Bot. VI, 478 (1847) ...	...	...	—	—	T.	—	—	—	B.fl.III,232	
<i>E. Gunnii</i> ; J. Hooker in Lond. Journ. Bot. III, 499 (1844) ...	...	...	—	S.A.	T.	V.	N.S.W.	—	B.fl.III,246	M.fr.II,62; VII,43; XI,38.
<i>E. resinifera</i> ; Smith in White's Journ. Voy. N.S. Wales 231 (1790) ...	...	...	—	—	—	—	N.S.W.	Q.	B.fl.III,245	M.fr.II,172; VII,43.
<i>E. saligna</i> ; Smith in Transact. Linn. Soc. III, 285 (1797) ...	...	...	—	—	—	—	N.S.W.	Q.	B.fl.III,245	M.fr.II,173; VII,44; II,52.
<i>E. punctata</i> ; De Candolle, prodr. III, 217 (1828) ...	...	...	—	—	—	—	N.S.W.	—	B.fl.III,242	M.fr.VII,43.
<i>E. oligantha</i> ; Schauer in Walp. rep. II, 926 (1843) ...	...	...	—	—	—	—	—	—	N.A.	B.fl.III,213
<i>E. alba</i> ; Reinwardt in Blume, Bijdr. 1101 (1826) ...	...	...	—	—	—	—	—	Q. N.A.	B.fl.III,243	... As., P.
<i>E. pulverulenta</i> ; Sims, Bot. Mag. t. 2089 (1819) ...	...	...	—	—	—	V.	N.S.W.	—	B.fl.III,224	M.fr.II,70.
<i>E. Stuartiana</i> ; F. v. M. in Benth. Fl. Austr. III, 243 (1866) ...	...	...	—	S.A.	T.	V.	N.S.W.	Q.	B.fl.III,243	M.fr.II,62.
<i>E. viminalis</i> ; Labillardière, Nov. Holl. pl. spec. II, 12, t. 151 (1806) ...	...	...	—	S.A.	T.	V.	N.S.W.	—	B.fl.III,239	M.fr.II,64; VII,44.
<i>E. rostrata</i> ; Schlechtendal, Linnaea XX, 655 (1847) ...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q. N.A.	B.fl.III,240	M.fr.II,40,65; VII,44.
<i>E. tereticornis</i> ; Smith, Spec. Bot. New Holl. 41 (1793) ...	...	...	—	—	—	V.	N.S.W.	Q.	B.fl.III,241	
<i>E. rudis</i> ; Endlicher in Hueg. enum. 49 (1837) ...	...	...	W.A.	—	—	—	—	—	B.fl.III,244	M.fr.XI,14.
<i>E. foecunda</i> ; Schauer in Lehm. pl. Preiss. I, 130 (1844) ...	...	...	W.A.	—	—	—	—	—	B.fl.III,252	
<i>E. redunda</i> ; Schauer in Lehm. pl. Preiss. I, 127 (1844) ...	...	...	W.A.	—	—	—	—	—	B.fl.III,253	M.fr.XI,15.
<i>E. grossa</i> ; F. v. M. in Benth. Fl. Austr. III, 232 (1866) ...	...	...	W.A.	—	—	—	—	—	B.fl.III,232	M.fr.VII,41.
<i>E. obcordata</i> ; Turczaninow in Bull. Acad. Petersb. 416 (1852) ...	...	...	W.A.	—	—	—	—	—	B.fl.III,234	M.fr.VII,44; VIII,184.
<i>E. occidentalis</i> ; Endlicher in Hueg. enum. 49 (1837) ...	...	...	W.A.	—	—	—	—	—	B.fl.III,235	M.fr.II,39; VII,43.
<i>E. cornuta</i> ; Labillardière, Voy. I, 403, t. 20 (1799) ..	...	...	W.A.	—	—	—	—	—	B.fl.III,234	M.fr.II,39; V,210; XI,38.
<i>E. gamophylla</i> ; F. v. M., fragm. XI, 40 (1878) ...	...	...	—	S.A.	—	—	—	N.A.	—	M.fr.XI,40.
<i>E. perfoliata</i> ; R. Brown in Benth. Fl. Austr. III, 253 (1866) ...	...	...	—	—	—	—	—	—	N.A.	B.fl.III,253
<i>E. setosa</i> ; Schauer in Walpers, rep. bot. syst. II, 926 (1843) ..	...	...	—	—	—	—	—	Q. N.A.	B.fl.III,254	M.fr.III,123; X,106.

E. Torelliana; F. v. M., fragm. X, 106 (1877)	...	...	...	...	...	—	—	—	—	—	Q.	—	—	M.fr.XI,42.
E. peltata; Benth. Fl. Austr. III, 254 (1866)	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl.III,254	
E. latifolia; F. v. M. in Journ. Linn. Soc. III, 94 (1858)	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.III,255	
E. ptychocarpa; F. v. M. in Journ. Linn. Soc. III, 90 (1858)	...	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.III,255	M.fr.VII,44.
E. Abergiana; F. v. M., fragm. XI, 41 (1878)	...	...	...	...	...	—	—	—	—	—	Q.	—	—	M.fr.XI,41.
E. calophylla; R. Brown in Journ. Geogr. Soc. 20 (1831)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,255	M.fr.II,35&171.
E. ficifolia; F. v. M., fragm. II, 85 (1860)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,256	M.fr.II,85&176; VI,25; XI,15.
E. corymbosa; Smith, Specim. Bot. New Holl. 43 (1793)	...	...	...	...	...	—	—	—	V.	N.S.W.	Q.	N.A.	B.fl.III,256	M.fr.II,46&174; VII,43; XI,42.
E. terminalis; F. v. M. in Journ. Linn. Soc. III, 89 (1858)	...	...	...	...	...	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.III,257	M.fr.XI,42.
E. Foelscheana; F. v. M. in Melb. Chemist, No. 55, p. 56 (1882)	...	...	...	...	...	—	—	—	—	—	—	N.A.	—	
E. dichromophloia; F. v. M. in Journ. Linn. Soc. III, 89 (1858)	...	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.III,257	
E. trachyphloia; F. v. M. in Journ. Linn. Soc. III, 90 (1858)	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl.III,221	M.fr.XI,42.
E. eximia; Schauer in Walpers, rep. bot. syst. II, 925 (1843)	...	...	...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.III,258	
E. maculata; Hooker, icones plant. t. 619 (1844)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.III,258	M.fr.II,47,147; VII,43; XI,45.
E. Watsoniana; F. v. M., fragm. X, 98 (1877)	...	...	...	...	...	—	—	—	—	—	Q.	—	—	M.fr.X,98.
E. odontocarpa; F. v. M. in Journ. Linn. Soc. III, 98 (1858)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,260	
E. tetrodonta; F. v. M. in Journ. Linn. Soc. III, 97 (1858)	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.III,260	
E. erythrocorys; F. v. M., fragm. II, 33 (1860)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,258	M.fr.II,33.
E. tetragona; F. v. M., fragm. VI, 51 (1864)...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,259	M.fr.II,37.
E. endesmioides; F. v. M., fragm. II, 35 (1860)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,260	M.fr.II,35.

TRISTANIA; R. Brown in Ait. hort. Kew. sec. ed. IV, 417 (1812). (Lophostemon, Tristaniopsis.)

T. neriifolia; R. Brown in Ait. hort. Kew. sec. ed. IV, 417 (1812)	...	...	—	—	—	—	N.S.W.	—	—	B.fl.III,261	M.fr.IV,56.
T. suaveolens; Smith in Rees, Cycl. XXXVI (1817)...	...	...	—	—	—	—	N.S.W.	Q. N.A.	B.fl.III,261	M.fr.I,81; IV,57.	
T. conferta; R. Brown in Ait. hort. Kew. sec. ed. IV, 417 (1812)	...	...	—	—	—	—	N.S.W.	Q. N.A.	B.fl.III,263	M.fr.I,82; IV,57.	
T. lactiflora; F. v. M., fragm. I, 82 (1858)	...	...	—	—	—	—	—	N.A.	B.fl.III,263	M.fr.I,82.	
T. exiliflora; F. v. M., fragm. V, 11 (1865)	...	...	—	—	—	—	—	Q.	B.fl.III,264	M.fr.IX,190.	
T. laurina; R. Brown in Ait. hort. Kew. sec. ed. IV, 417 (1812)...	...	...	—	—	—	—	V. N.S.W.	Q.	B.fl.III,264	M.fr.I,81; IV,56.	
T. psidioides; Cunningham in Bot. Regist. XXII, n. 1839 (1837)...	...	...	—	—	—	—	—	N.A.	B.fl.III,264		
T. umbrosa; Cunningham in Bot. Regist. XXII, n. 1839 (1837)	...	...	—	—	—	—	—	N.A.	B.fl.III,265		
T. longivalvis; F. v. M. in Wing's S. Sc. Record, new ser. vol. II, April (1886)	—	—	—	—	—	—	—	Q. N.A.	—		

METROSIDEROS; Banks in Gaertner, de fruct. I, 170, t. 34 (1788). (Syncarpia, Kamptzia, Lysicarpus, Xanthostemon, Nania, Fremya, Cloezia, Spermalepis.)

M. glomulifera; Smith in Transact. Linn. Soc. III, 269 (1797)	...	...	...	...	...	...	...	...	...	—	—	—	N.S.W.	Q.	—	B.fl.III,265	M.fr.I,79.
M. leptopetala; F. v. M., Docum. Intercol. Exhib. 30 (1867)	...	...	...	...	...	...	...	...	...	—	—	—	N.S.W.	Q.	—	B.fl.III,266	M.fr.I,79.
M. ternifolia; F. v. M., Docum. Intercol. Exhib. 30 (1867)	...	...	...	...	...	...	...	...	...	—	—	—	—	Q.	—	B.fl.III,267	M.fr.I,82.
M. encalyptoides; F. v. M., fragm. I, 243 (1858)	...	...	...	...	...	...	...	...	...	—	—	—	—	—	N.A.	B.fl.III,267	M.fr.I,81.
M. polymorpha; Gaudichaud, Freyc. Voy. Bot. 99, t. 108 (1826)	...	...	...	...	...	...	...	...	...	—	—	—	N.S.W.	—	—	—	M.fr.VIII,14 ... P.
M. nervulosa; C. Moore & F. v. M., fragm. VIII, 15 (1873)	...	...	...	...	...	...	...	...	...	—	—	—	N.S.W.	—	—	—	M.fr.I,80.
M. chrysantha; F. v. M., fragm. IV, 159 (1864)	...	...	...	...	...	...	...	...	...	—	—	—	—	Q.	—	B.fl.III,268	M.fr.IV,159.
M. paradoxa; F. v. M., fragm. I, 243 (1859)	...	...	...	...	...	...	...	...	...	—	—	—	—	—	N.A.	B.fl.III,269	M.fr.I,80.
M. tetrapetala; F. v. M., fragm. VII, 41 (1870)	...	...	...	...	...	...	...	...	...	—	—	—	—	Q. N.A.	—	—	M.fr.VII,41.
M. pachysperma; F. v. M. & Bailey in Bailey's occas. papers on Q. Flora I, 4 (1886)	...	...	...	...	...	...	...	...	...	—	—	—	—	Q.	—	—	

BACKHOUSIA; Hooker and Harvey in Bot. Mag. t. 4133 (1845).

B. myrtifolia; Hooker & Harvey in Bot. Mag. t. 4133 (1845)	...	...	...	...	...	...	...	...	...	—	—	—	V. N.S.W.	Q.	—	B.fl.III,269	M.fr.I,78; IX,43.
B. angustifolia; F. v. M., fragm. I, 79 (1858)	...	...	...	...	...	...	...	...	...	—	—	—	—	Q.	—	B.fl.III,270	M.fr.IX,43.
B. sciadophora; F. v. M., fragm. II, 26 (1860)	...	...	...	...	...	...	...	...	...	—	—	—	N.S.W.	Q.	—	B.fl.III,270	M.fr.II,171.
B. citriodora; F. v. M., fragm. I, 78 (1858)	...	...	...	...	...	...	...	...	...	—	—	—	—	Q.	—	B.fl.III,270	M.fr.I,78.
B. Bancroftii; F. v. M. & Bailey in catal. of Queensland Woods 46 (1886)	...	...	...	...	...	...	...	...	...	—	—	—	—	Q.	—	—	

OSBORNIA; F. v. M., fragm. III, 30 (1862).

O. octodonta; F. v. M., fragm. III, 31 (1862)	...	...	...	...	...	...	...	...	...	—	—	—	—	Q. N.A.	B.fl.III,271	M.fr.IX,143	... As., P.
---	-----	-----	-----	-----	-----	-----	-----	-----	-----	---	---	---	---	---------	--------------	-------------	-------------

RHODOMYRTUS; De Candolle, prodr. III, 240 (1828).

R. psidioides; Benth. Fl. Austr. III, 272 (1866) ...	...	...	...	...	...	...	...	N.S.W.	Q.	—	B.fl.III,272	M.fr.II,86; III,165; IX,142.
R. trineura; F. v. M. in Benth. Fl. Austr. III, 272 (1866)...	...	...	...	...	...	...	...	—	Q.	—	B.fl.III,272	M.fr.IV,117; IX,142.
R. cymiflora; F. v. M. in Benth. Fl. Austr. III, 273 (1866) ...	...	...	...	...	...	...	...	—	Q.	—	B.fl.III,273	M.fr.V,12; IX,149.
R. macrocarpa; Benth. Fl. Austr. III, 273 (1866) ...	...	...	...	...	...	...	...	—	Q.	—	B.fl.III,273	M.fr.IX,142.

MYRTUS; Tournefort, inst. 640, t. 409 (1700), from Plinius.

M. rhytisperma; F. v. M., fragm. I, 77 (1858) ...	...	...	...	...	...	...	...	N.S.W.	Q.	—	B.fl.III,274	M.fr.IX,147.
M. tenuifolia; Smith in Transact. Linn. Soc. III, 280 (1797) ...	...	...	...	...	...	...	...	N.S.W.	Q.	—	B.fl.III,274	M.fr.IX,147.
M. gonoclada; F. v. M. in Benth. Fl. Austr. III, 275 (1866) ...	...	...	...	...	...	...	...	—	Q.	—	B.fl.III,275	M.fr.IX,144.
M. lasioclada; F. v. M., fragm. IX, 148 (1875) ...	...	...	...	...	...	...	...	—	Q.	—	—	M.fr.IX,148.
M. Hillii; Benth. Fl. Austr. III, 275 (1866) ...	...	...	...	...	...	...	...	—	Q.	—	B.fl.III,275	M.fr.IX,149.
M. Becklerii; F. v. M., fragm. II, 85 (1861) ...	...	...	...	...	...	...	...	N.S.W.	Q.	—	B.fl.III,272	M.fr.IX,147.
M. Bidwillii; Benth. Fl. Austr. III, 275 (1866) ...	...	...	...	...	...	...	...	—	Q.	—	B.fl.III,275	—
M. racemulosa; Benth. Fl. Austr. III, 276 (1866) ...	...	...	...	...	...	...	...	—	Q.	—	B.fl.III,276	M.fr.IX,147.
M. acmenioides; F. v. M., Fragm. I, 77 (1858) ...	...	...	...	...	...	...	...	N.S.W.	Q.	—	B.fl.III,276	M.fr.IX,147.
M. fragrantissima; F. v. M. in Benth. Fl. Austr. III, 277 (1866)...	...	...	...	...	...	...	...	N.S.W.	Q.	—	B.fl.III,277	—
M. Shepherdi; F. v. M., fragm. IX, 148 (1875) ...	...	...	...	...	...	...	...	—	Q.	—	—	M.fr.IX,148.

RHODAMNIA; Jack in Hook. Comp. to the Bot. Mag. I, 153 (1835).

R. sessiliflora; Benth. Fl. Austr. III, 275 (1866) ...	...	...	...	...	...	...	...	—	Q.	—	B.fl.III,277	M.fr.IX,142.
R. trinervia; Blume, Mus. Bot. Lugd. I, 79 (1849) ...	...	...	...	...	...	...	...	N.S.W.	Q.	—	B.fl.III,278	M.fr.I,76; IX,142.
R. argentea; Benth. Fl. Austr. III, 276 (1866) ...	...	...	...	...	...	...	...	N.S.W.	Q.	—	B.fl.III,278	M.fr.IX,142.
R. Blairiana; F. v. M., fragm. IX, 141 (1875) ...	...	...	...	...	...	...	...	—	Q.	—	—	M.fr.IX,141.

FENZLIA; Endlicher in Atacta 19, t. 17 & 18 (1833).

F. obtusa; Endlicher, Atacta 19, t. 17 (1833) ...	...	...	...	...	...	...	...	—	Q.	—	B.fl.III,279	M.fr.IX,142.
F. retusa; Endlicher, Atacta 20, t. 18 (1833)...	...	...	...	...	...	...	...	—	Q.	N.A.	B.fl.III,279	—

DECASPERMUM; R. & G. Forster, char. gen. 73, t. 37 (1776). (Nelitris.)

D. paniculatum; Baillon, hist. des plant. 355 (1871) ...	...	...	...	...	...	...	...	—	Q.	—	B.fl.III,279	M.fr.IV,56; IX,149 ... As.
--	-----	-----	-----	-----	-----	-----	-----	---	----	---	--------------	----------------------------

EUGENIA; Micheli, nov. pl. gen. 226, t. 108 (1729). (Plinia, Plumier 1703, Acmene, Jambosa, Syzygium, Cleistocalyx partly.)

E. rariflora; Benth. in Hook. Lond. Journ. II, 221 (1843) ...	...	...	...	...	...	...	...	—	Q.	—	B.fl.III,282	M.fr.III,130; V,15; IX,143. P.
E. Smithii; Poir. Encycl. méth. Suppl. III, 126 (1813) ...	...	...	...	...	...	...	...	V. N.S.W.	Q.	N.A.	B.fl.III,282	M.fr.IV,59; IX,145.
E. hemilampra; F. v. M., fragm. IX, 145 (1875) ...	...	...	...	...	...	...	...	N.S.W.	Q.	—	—	M.fr.IX,145.
E. Ventenatii; Benth. Fl. Austr. III, 283 (1866) ..	...	...	...	...	...	...	...	N.S.W.	Q.	—	B.fl.III,283	M.fr.IV,58; IX,145.
E. leptantha; Wight, Illustr. II, 15 (1850) ...	...	...	...	...	...	...	...	—	Q.	—	B.fl.III,283	M.fr.IX,143 ... As.
E. Moorei; F. v. M., fragm. V, 33 (1865) ...	...	...	...	...	...	...	...	N.S.W.	Q.	—	B.fl.III,283	M.fr.IX,143.
E. corynantha; F. v. M., fragm. IX, 144 (1875) ...	...	...	...	...	...	...	...	N.S.W.	Q.	—	—	M.fr.IX,144.
E. corniflora; F. v. M., fragm. V, 32 (1865)...	...	...	...	...	...	...	...	—	Q.	—	B.fl.III,284	M.fr.V,32.
E. Hodgkinsoniae; F. v. M., fragm. IX, 145 (1875)...	...	...	...	...	...	...	...	N.S.W.	Q.	—	—	M.fr.IX,145.
E. Tierneyana; F. v. M., fragm. V, 14 (1865) ...	...	...	...	...	...	...	...	—	Q.	—	B.fl.III,284	M.fr.IX,143.
E. grandis; Wight, Illustr. II, 17 (1850) ...	...	...	...	...	...	...	...	—	Q.	—	B.fl.III,285	M.fr.V,13; IX,143 ... As.
E. suborbicularis; Benth. Fl. Austr. III, 285 (1866) ...	...	...	...	...	...	...	...	—	Q.	—	B.fl.III,285	M.fr.IX,143.
E. Wilsonii; F. v. M., fragm. V, 12 (1865) ...	...	...	...	...	...	...	...	—	Q.	—	B.fl.III,285	M.fr.IX,143.
E. eucalyptoides; F. v. M., fragm. IV, 55 (1863) ...	...	...	...	...	...	...	...	—	Q.	N.A.	B.fl.III,285	M.fr.I,226; IX,143.
E. Australis; Wendland in Link, enum. pl. hort. bot. II, 28 (1822) ...	...	...	...	...	...	...	...	N.S.W.	Q.	—	B.fl.III,286	M.fr.I,225; V,33; IX,143.
E. angophoroides; F. v. M., fragm. V, 33 (1865) ...	...	...	...	...	...	...	...	—	Q.	—	B.fl.III,286	M.fr.IX,145.
E. Armstrongii; Benth. Fl. Austr. III, 286 (1866) ...	...	...	...	...	...	...	...	—	N.A.	—	B.fl.III,286	—
E. oleosa; F. v. M., fragm. V, 15 (1865) ...	...	...	...	...	...	...	...	—	Q.	—	B.fl.III,287	M.fr.V,33; IX,146.
E. Dallachiana; F. v. M. in Benth. Fl. Austr. III, 287 (1866) ...	...	...	...	...	...	...	...	—	Q.	—	B.fl.III,287	M.fr.IX,145.
E. cryptophlebia; F. v. M., fragm. IX, 144 (1875) ...	...	...	...	...	...	...	...	—	Q.	—	—	M.fr.IX,144.
E. Holtziana; F. v. M. in Austr. Journ. of Pharm. I, 199 (1886)...	...	...	...	...	...	...	...	—	N.A.	—	—	—

ACICALYPTUS; A. Gray, New Gen. Un. St. Exped. 9 (1853). (Calyptanthus partly, Cleistocalycis subgenus.)

A. Fullagari; F. v. M., fragm. VIII, 15 (1873) ... — — — — N.S.W. — — — M.fr. VIII, 281.

BARRINGTONIA; R. & G. Forster, char. gen. 75, t. 38 (1776). (Stravadium.)

B. speciosa; R. & G. Forster, char. gen. 76, t. 38 (1776) ... — — — — — Q. — B.fl. III, 288 M.fr. IX, 118 ... As., P.
 B. racemosa; Gaudichaud in Freyc. Voy. Bot. 483, t. 107 (1826) ... — — — — — Q. — — M.fr. IX, 118 ... As., P.
 B. acutangula; Gaertner, de fruct. II, 97, t. 101 (1791) ... — — — — — Q. N.A. B.fl. III, 288 M.fr. IX, 119 ... As.

CAREYA; Roxburgh, pl. Corom. III, 13, t. 217 (1816).

C. Australis; F. v. M., fragm. V, 183 (1866) ... — — — — — Q. N.A. B.fl. III, 289 M.fr. V, 183.

SONNERATIA; Linné, fil. suppl. plant. 38 (1781).

S. acida; Linné, fil. suppl. plant. 38 (1781) ... — — — — — Q. N.A. B.fl. III, 301 M.fr. III, 109 ... As., Afr.

MELASTOMACEAE.

A. L. DE JUSSIEU, Gen. plant. 328 (1789).—Genera 5, Species 7.

MEMECYLON; Linné, Fl. Zeyl. app. 9 (1747).

M. umbellatum; N. L. Burmann, Fl. Ind. 87 (1768) ... — — — — — Q. N.A. B.fl. III, 293 ... As.

OSBECKIA; Linné, spec. plant. I, 345 (1753).

O. Chinensis; Linné, spec. plant. 345 (1753) ... — — — — — Q. — B.fl. III, 291 M.fr. IV, 160 ... As.
 O. Australiana; Naudin in Ann. sc. Nat. trois. sér. XIV, 59 (1855) ... — — — — — — N.A. B.fl. III, 291 M.fr. IV, 161.
 O. perangusta; F. v. M., fragm. V, 181 (1866) ... — — — — — — N.A. — M.fr. V, 181.

OTANTHERA; Blume in Flora, Regensb. IV, 488 (1831).

O. bracteata; Korthals in Verh. Nat. Gesch. der Ned. Bes. 235, t. 51 (1841) ... — — — — — Q. — B.fl. III, 292 ... P.

MELASTOMA; J. Burmann, thesaur. Zeyl. 72 (1737).

M. Malabathricum; Linné, spec. plant. 390 (1753) ... — — — — — N.S.W. Q. N.A. B.fl. III, 292 M.fr. IV, 161 ... As., P.

MEDINILLA; Gaudichaud in Freycin. voy. Bot. 484, t. 106 (1826).

Balls-Headleyi; F. v. M. in Vict. Naturalist new ser. II, 125 (1887) ... — — — — — Q. — —

RHAMNACEAE.

A. L. DE JUSSIEU, Gen. plant. 376 (1789), from B. de Jussieu (1759).—Genera 10, Species 90.

VENTILAGO; Gaertner, de fruct. I, 223, t. 49 (1788).

V. viminalis; Hooker in Mitch. Trop. Austr. 369 (1848) ... — S.A. — — N.S.W. Q. N.A. B.fl. I, 411 M.fr. III, 63; IX, 140.
 V. ecorollata; F. v. M. in Wing's S. Sc. Rec. III, 176 (1883) ... — — — — — Q. — — M.fr. IX, 141.

ZIZYPHUS; Tournefort, inst. 627, t. 403 (1700), from Plinius. (Ziziphus.)

Z. Oenopia; Miller, Gardeners' Dictionary n. 3 (1731) ... — — — — — Q. N.A. B.fl. I, 412
 Z. Jujuba; Lamarck, Encycl. méth. III, 318 (1789) ... — — — — — Q. — B.fl. I, 412 ... As., Afr.
 Z. quadrilocularis; F. v. M., fragm. III, 57 (1862) ... — — — — — — N.A. B.fl. I, 412 M.fr. III, 57.

DALLACHYA; F. v. M., fragm. IX, 140 (1875). (Rhamnus partly.)

D. Vitiensis; F. v. M., fragm. IX, 140 (1875) ... — — — — — Q. — — M.fr. XI, 134 ... P.

COLUBRINA; L. C. Richard in Ann. des sc. nat. X, 368 (1827).

C. Asiatica; Brogniart in Ann. des sc. nat. X, 369 (1827) ... — — — — — Q. N.A. B.fl. I, 413 M.fr. III, 63; IX, 139 ... As., Afr., P.

ALPHITONIA; Reisseck in Endl. gen. plant. 1098 (1840).

A. excelsa; Reisseck in Endl. gen. plant. 1098 (1840) ... — — — — — N.S.W. Q. N.A. B.fl. I, 414 M.fr. III, 63; IX, 140 ... As., P.

EMMENOSPERMA; F. v. M., fragm. III, 63 (1862).

E. alphonoides; F. v. M., fragm. III, 63 (1862)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,415	M.fr.IX,140.
E. Cunninghamii; Benth., Fl. Austr. I, 415 (1863)	—	—	—	—	—	—	N.A.	B.fl.I,415	

POMADERIS; Labillardière, Nov. Holl. pl. specim. I, 61, t. 86 (1804). (Pomatoderris.)

P. lanigera; Sims, Bot. Mag. t. 1823 (1816)	—	—	T.	V.	N.S.W.	—	—	B.fl.I,416	
P. elliptica; Labillardière, Nov. Holl. pl. spec. I, 61, t. 86 (1804) ..	—	—	T.	V.	N.S.W.	Q.	—	B.fl.I,417	M.fr.III,69; IX,139.
P. phillyroides; Sieber in De Candolle, prodr. II, 33 (1825) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,418	
P. vacciniifolia; Reisseck & F. v. M. in Linnæa XXIX, 266 (1857) ...	—	—	—	V.	N.S.W.	—	—	B.fl.I,418	M.fr.III,71.
P. myrtilloides; Fenzl in Hueg. enum. 22 (1837)	W.A.	S.A.	—	—	—	—	—	B.fl.I,419	M.fr.IX,138.
P. grandis; F. v. M., fragm. III, 68 (1862)	W.A.	—	—	—	—	—	—	B.fl.I,417	M.fr.IX,139.
P. Calvertiana; F. v. M., fragm. IX, 138 (1875)	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,138.
P. ledifolia; Cunningham in Field, N.S.W. 351 (1825)	—	—	—	V.	N.S.W.	—	—	B.fl.I,419	M.fr.III,69.
P. apetala; Labillardière, Nov. Holl. pl. spec. I, 62, t. 87 (1804) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.I,419	M.fr.III,69&73; IX,139.
P. cinerea; Benth., Fl. Austr. I, 420 (1863)	—	—	—	V.	N.S.W.	—	—	B.fl.I,420	
P. prunifolia; Cunningham in Hueg. enum. 22 (1837)	—	—	—	V.	N.S.W.	Q.	—	B.fl.I,420	M.fr.III,75; IX,139.
P. ligustrina; Sieber in De Candolle, prodr. II, 34 (1825)	—	—	—	V.	N.S.W.	—	—	B.fl.I,420	M.fr.III,71; IX,139.
P. betulina; Cunningham in Bot. Mag. t. 3212 (1833)	—	—	—	V.	N.S.W.	—	—	B.fl.I,421	M.fr.III,76; IX,139.
P. Forrestiana; F. v. M., fragm. IX, 139 (1875)	W.A.	—	—	—	—	—	—	—	M.fr.IX,139.
P. obcordata; Fenzl in Hueg. enum. 23 (1837)	W.A.	S.A.	—	V.	—	—	—	B.fl.I,421	M.fr.III,73.
P. racemosa; Hooker, Journ. of Bot. I, 256 (1834)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.I,421	M.fr.III,75&168; IX,139.
P. subrepanda; Reisseck & F. v. M. in Linnæa XXIX, 267 (1857) ...	—	—	—	V.	—	—	—	B.fl.I,422	M.fr.III,74.
P. elachophylla; F. v. M., fragm. II, 131 (1861)	—	—	—	V.	—	—	—	B.fl.I,422	M.fr.III,72,166; IX,139 ... N.Z.
P. phyllifolia; Loddiges, Bot. Cabinet t. 120 (1818)	—	—	T.	V.	N.S.W.	—	—	B.fl.I,422	M.fr.III,73.
P. intangenda; F. v. M., fragm. X, 52 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,53.

CRYPTANDRA; Smith in Transact. Linn. Soc. IV, 217 (1798). (Trymalium, Spyridium, Stenanthemum, Wichurea, Stenodiscus.)

C. albicans; F. v. M., Cens. 60 (1882) from Reisseck (1847)	W.A.	—	—	—	—	—	—	B.fl.I,423	
C. Billardieri; F. v. M., Cens. 60 (1882) from Fenzl (1837)	W.A.	—	—	—	—	—	—	B.fl.I,423	
C. ledifolia; F. v. M., Cens. 60 (1882) from Fenzl (1837)	W.A.	—	—	—	—	—	—	B.fl.I,424	
C. Daltoni; F. v. M., Cens. 60 (1882)	—	—	—	V.	—	—	—	—	M.fr.IX,135,198.
C. angustifolia; F. v. M., Cens. 60 (1882) from Reisseck (1847)	W.A.	—	—	—	—	—	—	B.fl.I,424	M.fr.IX,135.
C. Wichureae; F. v. M., Cens. 60 (1882) from Nees (1847)	W.A.	—	—	—	—	—	—	B.fl.I,425	M.fr.IX,135.
C. ericifolia; Smith in Rees, Cycloped. (1808)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,438	M.fr.III,66; IX,137.
C. propinqua; Cunningham in Hueg. enum. 23 (1837)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.I,442	M.fr.III,67; IX,137.
C. hispidula; Reisseck & F. v. M. in Linnæa XXIX, 294 (1857)	—	S.A.	—	—	—	—	—	B.fl.I,439	
C. spyridioides; F. v. M., fragm. III, 68 (1862)	W.A.	—	—	—	—	—	—	B.fl.I,439	M.fr.IX,138.
C. scoparia; Reisseck in Lehm. pl. Preiss. II, 283 (1847)	W.A.	—	—	—	—	—	—	B.fl.I,439	
C. spinescens; Sieber in De Candolle, prodr. II, 38 (1825)	—	—	—	—	N.S.W.	—	—	B.fl.I,439	
C. amara; Smith in Transact. Linn. Soc. X, 245, t. 18 (1811)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,440	M.fr.III,66&67; IX,137.
C. lanosiflora; F. v. M., fragm. III, 65 (1862)	—	—	—	—	N.S.W.	—	—	B.fl.I,440	M.fr.III,65.
C. tomentosa; Lindley in Mitch. Three Exped. II, 178 (1838)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.I,441	M.fr.III,67; IX,138.
C. nutans; Steudel in Lehm. pl. Preiss. I, 186 (1844)	W.A.	—	—	—	—	—	—	B.fl.I,441	M.fr.IX,138.
C. glabriflora; Benth., Fl. Austr. I, 441 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,441	
C. alpina; J. Hooker, Fl. Tasm. I, 75, t. 12 (1860)	—	—	T.	—	—	—	—	B.fl.I,441	
C. Leucopogon; Meissner in Lehm. pl. Preiss. II, 287 (1847)	W.A.	—	—	—	—	—	—	B.fl.I,442	
C. parvifolia; Turczaninow in Bull. de Mose. I, 459 (1858)	W.A.	—	—	—	—	—	—	B.fl.I,442	
C. buxifolia; Fenzl in Hueg. enum. 23 (1837)	—	—	—	—	N.S.W.	—	—	B.fl.I,442	M.fr.IX,138.
C. pungens; Steudel in Lehm. pl. Preiss. I, 187 (1844)	W.A.	—	—	—	—	—	—	B.fl.I,443	M.fr.IX,138.
C. mutila; Nees in Lehm. pl. Preiss. II, 289 (1847)	W.A.	—	—	—	—	—	—	B.fl.I,443	
C. longistaminea; F. v. M., fragm. III, 64 (1862)	—	—	—	—	N.S.W.	—	—	B.fl.I,443	M.fr.IX,138.

<i>C. arbutiflora</i> ; Fenzl in Hueg. enum. 26 (1837)	W.A.	—	—	—	—	—	—	B.fl.I, 444	M.fr.III, 64; IX, 138.
<i>C. miliaris</i> ; Reisseck in Lehm. pl. Preiss. II, 288 (1847)	W.A.	—	—	—	—	—	—	B.fl.I, 444	
<i>C. nudiflora</i> ; F. v. M., fragm. III, 64 (1862)	W.A.	—	—	—	—	—	—	B.fl.I, 444	M.fr.III, 64.
<i>C. ulicina</i> ; Hooker, Journ. of Bot. I, 257 (1834)	—	—	T.	—	—	—	—	B.fl.I, 434	M.fr.IX, 137.
<i>C. pomaderoides</i> ; Reisseck in Endl. nov. stirp. dec. 29 (1839)	W.A.	—	—	—	—	—	—	B.fl.I, 435	M.fr.IX, 137.
<i>C. pumila</i> ; F. v. M., Cens. 61 (1882)	W.A.	—	—	—	—	—	—	—	M.fr.IX, 137.
<i>C. leucophracta</i> ; Schlechtendal, Linnaea XX, 640 (1847)	—	S.A.	T.	V.	—	—	—	B.fl.I, 435	M.fr.III, 77; IX, 136.
<i>C. coronata</i> ; Reisseck in Lehm. pl. Preiss. II, 288 (1847)	W.A.	—	—	—	—	—	—	B.fl.I, 436	M.fr.IX, 137.
<i>C. Scortechinii</i> ; F. v. M. in Melb. Chem. VI, 71 (1884)	—	—	—	V.	N.S.W.	Q.	—	—	
<i>C. humilis</i> ; F. v. M., Cens. 61 (1882) from Bentham (1863)	W.A.	—	—	—	—	—	—	B.fl.I, 436	
<i>C. Waterhousii</i> ; F. v. M., Cens. 61 (1882)	—	S.A.	—	—	—	—	—	B.fl.I, 436	M.fr.III, 83.
<i>C. tridentata</i> ; Steudel in Lehm. pl. Preiss. I, 186 (1844)	W.A.	—	—	—	—	—	—	B.fl.I, 427	M.fr.IX, 137.
<i>C. divaricata</i> ; F. v. M., Cens. 61 (1882) from Bentham (1863)	W.A.	—	—	—	—	—	—	B.fl.I, 427	M.fr.III, 68.
<i>C. obcordata</i> ; J. Hooker, Fl. Tasm. I, 71 (1860)	—	—	T.	V.	—	—	—	B.fl.I, 427	M.fr.III, 80.
<i>C. Lawrencii</i> ; J. Hooker, Fl. Tasm. I, 72 (1860)	—	—	T.	—	—	—	—	B.fl.I, 430	M.fr.IX, 136.
<i>C. Hookeri</i> ; F. v. M., Cens. 61 (1882)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.I, 428	M.fr.III, 79; IX, 136.
<i>C. spadicea</i> ; F. v. M., Cens. 61 (1882) from Fenzl (1837)	W.A.	—	—	—	—	—	—	B.fl.I, 428	M.fr.IX, 137.
<i>C. globulosa</i> ; F. v. M., Cens. 61 (1882) from G. Don (1830)	W.A.	—	—	—	—	—	—	B.fl.I, 429	M.fr.IX, 137.
<i>C. obovata</i> ; J. Hooker, Fl. Tasm. I, 74 (1860)	—	S.A.	T.	—	—	—	—	B.fl.I, 429	M.fr.IX, 136.
<i>C. spathulata</i> ; F. v. M., Cens. 61 (1882)	—	S.A.	—	—	—	—	—	B.fl.I, 430	M.fr.IX, 136.
<i>C. cordata</i> ; Turczaninow in Bull. de Mosc. XXXI, 459 (1858)	W.A.	—	—	—	—	—	—	B.fl.I, 430	M.fr.IV, 25; IX, 137.
<i>C. rotundifolia</i> ; F. v. M., Cens. 61 (1882)	W.A.	—	—	—	—	—	—	—	M.fr.IV, 25.
<i>C. phlebophylla</i> ; F. v. M., Cens. 61 (1882)	—	S.A.	—	—	—	—	—	B.fl.I, 430	
<i>C. coactilifolia</i> ; F. v. M., Cens. 61 (1882) from Reiss. (1856)	—	S.A.	—	—	—	—	—	B.fl.I, 431	M.fr.IX, 136.
<i>C. complicata</i> ; F. v. M., Cens. 61 (1882)	W.A.	—	—	—	—	—	—	B.fl.I, 431	M.fr.III, 78.
<i>C. westringifolia</i> ; F. v. M., Cens. 61 (1882) from Reiss. (1847)	W.A.	—	—	—	—	—	—	B.fl.I, 431	
<i>C. villosa</i> ; Turczaninow in Bull. de Mosc. XXXI, 458 (1858)	W.A.	—	—	—	—	—	—	B.fl.I, 432	
<i>C. pauciflora</i> ; Turczaninow in Bull. de Mosc. XXXI, 458 (1858)	W.A.	—	—	—	—	—	—	B.fl.I, 432	
<i>C. halmaturina</i> ; F. v. M., Cens. 61 (1882)	—	S.A.	—	—	—	—	—	B.fl.I, 432	
<i>C. bifida</i> ; F. v. M., Cens. 61 (1882)	—	S.A.	—	—	—	—	—	B.fl.I, 432	M.fr.IX, 136.
<i>C. subochreatea</i> ; F. v. M., Cens. 61 (1882)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.I, 432	M.fr.III, 82; IX, 136.
<i>C. vexillifera</i> ; Hooker, Journ. Bot. I, 257 (1834)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.I, 433	M.fr.III, 81; IX, 136.
<i>C. Wayii</i> ; F. v. M. and Tate in Transact. R. Soc. S.A. V, 89 (1882)	—	S.A.	—	—	—	—	—	—	

COLLETIA; Commerçon in A. L. de Jussieu, gen. 380 (1789). (Discaria, Tetrapasma.)

<i>C. pubescens</i> ; Brogniart in Annal. des sc. nat. X, 366 (1827)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.I, 445	M.fr.III, 83; IV, 175; IX, 140.
---	---	---	----	----	--------	----	---	-------------	---------------------------------

GOUANIA; N. J. Jacquin, stirp. Amer. hist. 263, t. 179 (1763).

<i>G. Hillii</i> ; F. v. M., fragm. IX, 163 (1875)	—	—	—	—	—	Q.	—	—	M.fr.IX, 163.
<i>G. Australiana</i> ; F. v. M., fragm. IV, 144 (1864)	—	—	—	—	—	Q.	—	—	M.fr.IV, 144.

VINIFERAE.

J. DE ST. HILAIRE, Exp. fam. II, 48, t. 79 (1805).—Genus 1, Species 17.

VITIS; Tournefort, inst. 613, t. 384 (1700), e Latinis. (Cissus.)

<i>V. Baudiniana</i> ; F. v. M., fragm. IV, 136 (1864)	—	—	—	V.	N.S.W.	Q.	—	B.fl.I, 447	M.fr.V, 212; VI, 178.
<i>V. oblonga</i> ; Bentham, Fl. Austr. I, 447 (1863)	—	—	—	—	—	Q.	—	B.fl.I, 447	M.fr.IV, 136; VI, 178; IX, 125.
<i>V. repens</i> ; Wight & Arnott, prodr. fl. penins. Ind. I, 186 (1834)	—	—	—	—	—	Q.	N.A.	B.fl.I, 447	M.fr.II, 73; V, 210; VI, 178; IX, 126. As.
<i>V. adnata</i> ; Wallich in Wight & Arnott, prodr. 126 (1834)	—	—	—	—	—	—	N.A.	B.fl.I, 448	M.fr.VI, 178. As, P
<i>V. nitens</i> ; F. v. M., fragm. II, 73 (1860)	—	—	—	—	N.S.W.	Q.	—	B.fl.I, 448	M.fr.VI, 178.
<i>V. brachypoda</i> ; F. v. M., fragm. IX, 125 (1875)	—	—	—	—	—	Q.	—	—	M.fr.IX, 125.

V. saponaria ; Seemann, Bonplandia, 254 (1859)	— — — — —	Q. —	B.fl.I,448	M.fr.VI,178.	
V. acris ; F. v. M., fragm. II, 75 (1860)	— — — — —	N.S.W. Q. —	B.fl.I,449	M.fr.VI,178.	
V. trifolia ; Linné, spec. plant. 203 (1753)	— — — — —	Q. N.A. B.fl.I,449	M.fr.II,75; VI,177; IX,126	...	As., P.
V. clematidea ; F. v. M., fragm. II, 74 (1860)	— — — — —	N.S.W. Q. —	B.fl.I,449	M.fr.V,210; VI,177; IX,126.	
V. Japonica ; Thunberg, Fl. Jap. 104 (1775)	— — — — —	Q. —	—	...	As., P.
V. acetosa ; F. v. M., plants of Vict. I, 44 (1860)	— — — — —	Q. N.A. B.fl.I,449	M.fr.VI,177; IX,126.		
V. hypoglaucia ; F. v. M., plants of Vict. I, 95 (1860)	— — — — —	V. N.S.W. Q. —	B.fl.I,450	M.fr.VI,177.	
V. sterculiifolia ; F. v. M. in Benth. Fl. Austr. I, 450 (1863)	— — — — —	N.S.W. Q. —	B.fl.I,450	M.fr.IX,126.	
V. penninervis ; F. v. M., fragm. VI, 177 (1868)	— — — — —	Q. —	—	M.fr.VI,177.	
V. opaca ; F. v. M. in Benth. Fl. Austr. I, 450 (1863)	— — — — —	N.S.W. Q. —	B.fl.I,450	M.fr.VI,177; IX,125.	
V. augustissima ; F. v. M., fragm. I, 141 (1859)	W.A. — — — — —	— — — — —	B.fl.I,450	M.fr.I,244; II,180; IX,126.	

LEEACEAE.

F. v. M. from De Candolle, prodr. I, 635 (1824).—Genus 1, Species 2.

LEEAE ; Linné, mantiss. 17 (1767).

L. staphylea ; Roxburgh, hort. Beng. 18 (1814)	— — — — —	Q. —	B.fl.I,451	M.fr.VI,178	... As.
F. Brunouiana ; Clarke in Britten's journ. of Bot. X, 105 (1881)	— — — — —	Q. N.A. —	—	M.fr.VI,178.	

ARALIACEAE.

VENTENAT, Tabl. III, (1799). (Umbelliferarum subordo.)—Genera 10, Species 22.

ASTROTTRICHA ; De Candolle, Mémoire sur la fam. des Ombellifères 25 (1829).

A. pterocarpa ; Bentham, Fl. Austr. III, 379 (1866)	— — — — —	Q. —	B.fl.III,379	M.fr.VII,148.	
A. floccosa ; De Candolle, Mém. des Ombell. 30, t. 5 (1829)	— — — — —	N.S.W. Q. —	B.fl.III,380		
A. longifolia ; Bentham in Hueg. enum. 55 (1837)	— — — — —	N.S.W. Q. —	B.fl.III,380	M.fr.VII,148.	
A. ledifolia ; De Candolle, Mém. des Ombell. 30, t. 6 (1829)	— — — — —	V. N.S.W. —	B.fl.III,380	M.fr.VII,148.	
A. Hamptoni ; F. v. M., fragm. VI, 125 (1868)	W.A. — — — — —	— N.A. —	—	M.fr.VI,125&250.	

DELARBREA ; Vieillard in Bulet. de la Soc. Linn. Normand. IX, 342 (1865). (Porospermum.)

D. Michiana ; F. v. M., fragm. VII, 95 (1870)	— — — — —	Q. —	—	M.fr.VII,95.	
--	-----------	------	---	--------------	--

PANAX ; Linné, spec. plant. edit. sec. 1513 (1763). (Nothopanax.)

P. Gunnii ; J. Hooker in Lond. Journ. of Bot. VI, 466 (1847)	— — — — —	T. — — — —	B.fl.III,381		
P. Murrayi ; F. v. M., fragm. II, 106 (1861)	— — — — —	V. N.S.W. Q. —	B.fl.III,381	M.fr.VII,96.	
P. mollis ; Bentham, Fl. Austr. III, 382 (1866)	— — — — —	Q. —	B.fl.III,382	M.fr.VII,96.	
P. Macgillivraei ; Bentham, Fl. Austr. III, 382 (1866)	— — — — —	Q. —	B.fl.III,382		
P. sambucifolius ; Sieber in De Candolle, prodr. III, 255 (1825)	— — — — —	T. V. N.S.W. Q. —	B.fl.III,382	M.fr.II,107; VII,95.	
P. cephalobotrys ; F. v. M., fragm. II, 83 (1861)	— — — — —	N.S.W. Q. —	B.fl.III,382	M.fr.II,83,107,176; VII,95.	
P. elegans ; C. Moore and F. v. M. in Trans. phil. Inst. Vict. II, 68 (1858)	— — — — —	N.S.W. Q. —	B.fl.III,383	M.fr.II,107.	
P. Cissodendron ; C. Moore and F. v. M., fragm. VII, 96 (1870)	— — — — —	N.S.W. — — — —	—	M.fr.VIII,280; IX,96.	

PENTAPANAX ; Seemann, Journal of Botany II, 294 (1864).

P. Willmottii ; F. v. M. in Vict. Naturalist, new ser. II, 125 (1887)	— — — — —	Q. —	—		
--	-----------	------	---	--	--

MOTHERWELLIA ; F. v. M., fragm. VII, 107 (1871).

M. haplosciadea ; F. v. M., fragm. VII, 107 (1871)	— — — — —	Q. —	—	M.fr.VII,107.	
---	-----------	------	---	---------------	--

CISSODENDRON ; Seemann, Journ. of Bot. II, 303 (1864). (Hedera partly, Irvingia, Kissodendron.)

C. Australianum ; Seemann, Journ. of Bot. II, 303 (1864)	— — — — —	Q. —	B.fl.III,384	M.fr.IV,120; V,18.	
---	-----------	------	--------------	--------------------	--

HEPTAPLEURUM ; Gaertner, de fruct. II, 472, t. 178 (1791). (Paratropia.)

H. venulosum ; Seemann, Journ. of Bot. III, 80 (1865)	— — — — —	Q. —	B.fl.III,384	M.fr.II,108...	.. As.
--	-----------	------	--------------	----------------	--------

BRASSAIA; Eodlicher, nov. stirp. decad. I, 89 (1839).

B. actinophylla; Eodlicher, nov. stirp. dec. I, 89 (1839) ... Q. — B.fl.III,385 M.fr.II,108; IV,121.

MERYTA; R. & G. Forster, char. gen. 119, t. 60 (1776). (Bothryodendron.)

M. latifolia; Seemann in Bonplandia, 295 (1862) ... N.S.W. — — M.fr.IX,169.

M. angustifolia; Seemann in Bonplandia, 296 (1862) ... N.S.W. — — M.fr.IX,169.

MACKINLAYA; F. v. M., fragm. IV, 120 (1864).

M. macrosciadea; F. v. M., fragm. IV, 120 (1864) ... Q. — B.fl.III,383 M.fr.II,108&176.

UMBELLIFERAE.

MORISON, hist. pl. II, lib. 3, sect. 9 (1680).—Genera 16, Species 106.

HYDROCOTYLE; Tournefort, inst. 328, t. 173 (1700).

H. vulgaris; Linné, spec. plant. 234 (1753) ... S.A. — V. N.S.W. Q. — B.fl.III,339 M.fr.IV,180 ... E., As., Afr., Am.

H. hirta; R. Brown in annal. des sc. phys. VI, 64 (1820) ... W.A. S.A. T. V. N.S.W. Q. — B.fl.III,339 M.fr.IV,181.

H. Candollei; F. v. M., key to the syst. of Vict. pl. I, 267 (1888) ... S.A. — V. N.S.W. Q. — B.fl.III,340 M.fr.IV,179&180.

H. rhombifolia; F. v. M., fragm. VII, 147 (1871) ... N.S.W. — — M.fr.VII,147.

H. pedicellosa; F. v. M., fragm. IV, 182 (1864) ... N.S.W. Q. — B.fl.III,341 M.fr.VII,147.

H. tripartita; R. Brown in annal. des sc. phys. VI, 69, t. 61 (1820) ... S.A. T. V. N.S.W. Q. — B.fl.III,341 M.fr.IV,181; VII,147.

H. comocarpa; F. v. M. in Vict. Nat. III, 126 (1887) ... S.A. — — — —

H. crassiuscula; Tate in Transact. R. S. of S.A. VI, 140 ... S.A. — — — —

H. plebeja; R. Brown in annal. des sc. phys. VI, 46, t. 60 (1820) ... W.A. — — — — B.fl.III,341

H. pterocarpa; F. v. M. in Transact. Vict. Inst. 126 (1855) ... S.A. T. V. — — — — B.fl.III,342 ... N.Z.

H. geraniifolia; F. v. M. in Transact. Vict. Inst. 126 (1855) ... V. N.S.W. — — B.fl.III,342 M.fr.IV,181.

H. medicaginoides; Turczaninow in Bull. Mosc. XXII, 27 (1849) ... W.A. S.A. — V. — — — B.fl.III,342 M.fr.IV,178.

H. muriculata; Turczaninow in Bull. Mosc. 28 (1849) ... W.A. — — — — B.fl.III,342

H. callicarpa; Bunge in Lehm. pl. Preiss. I, 283 (1844) ... W.A. S.A. T. V. N.S.W. — — B.fl.III,343 M.fr.IV,181.

H. scutellifera; Benth, Fl. Austr. III, 343 (1866) ... W.A. — — — — B.fl.III,343

H. hispidula; Bunge in Lehm. pl. Preiss. I, 283 (1844) ... W.A. — — — — B.fl.III,343

H. trachycarpa; F. v. M. in Schlecht. Linnæa XXV, 394 (1852) ... S.A. — — N.S.W. — — B.fl.III,343 M.fr.IV,181.

H. pilifera; Turczaninow in Bull. Mosc. XXII, 26 (1849) ... W.A. — — — — B.fl.III,344

H. capillaris; F. v. M., First Gen. Report 13 (1853) ... W.A. S.A. — V. N.S.W. — — B.fl.III,344 M.fr.IV,178.

H. rugulosa; Turczaninow in Bull. Soc. Mosc. XXII, 27 (1849) ... W.A. — — — — B.fl.III,344

H. grammatocarpa; F. v. M., fragm. II, 128 (1861) ... — — — — N.A. B.fl.III,344 M.fr.II,128.

H. diantha; De Candolle, prodr. IV, 63 (1830) ... W.A. S.A. — — — — B.fl.III,345 M.fr.IV,179.

H. lemnoides; Benth, Fl. Austr. III, 345 (1866) ... W.A. — — — — B.fl.III,345

H. alata; R. Brown in annal. des sc. phys. VI, 75, t. 61 (1820) ... W.A. — — — — B.fl.III,345

H. tetragonocarpa; Bunge in Lehm. pl. Preiss. I, 284 (1844) ... W.A. — — — — B.fl.III,345

H. blepharocarpa; F. v. M. in Wing's S. Sc. Rec. III, 175 (1883) ... W.A. — — — —

H. glochidiata; Benth, Fl. Austr. III, 346 (1866) ... W.A. — — — — B.fl.III,346

H. homalocarpa; F. v. M., fragm. II, 129 (1861) ... W.A. — — — — B.fl.III,346 M.fr.IV,181; VII,147.

H. Asiatica; Linné, spec. plant. 234 (1753) ... W.A. S.A. T. V. N.S.W. Q. N.A. B.fl.III,346 M.fr.VII,147 ... As., Afr., Am., P., N.Z.

DIDISCUS; De Candolle in Bot. Mag. t. 2857 (1828). (Dimetopia, Pritzelia, Huegelia, Cesatia, Hemicarpus.)

D. pusillus; F. v. M., fragm. IX, 47 (1875) ... W.A. S.A. — V. N.S.W. — — B.fl.III,348 M.fr.I,231.

D. cyanopetalus; F. v. M., fragm. IX, 46 (1875) ... W.A. S.A. — V. N.S.W. Q. — B.fl.III,348 M.fr.I,231.

D. eriocarpus; F. v. M., fragm. IX, 46 (1875) ... W.A. S.A. — — — — B.fl.III,348 M.fr.I,231; IX,46.

D. villosus; F. v. M. in Proc. Roy. Soc. Tasm. III, 238 (1857) ... — — — — N.A. B.fl.III,349

D. coeruleus; De Candolle in Hook. Bot. Mag. 2875 (1828) ... W.A. — — — — B.fl.III,349 M.fr.IX,47.

D. pilosus; Benth in Hueg. enum. 54 (1837) ... W.A. S.A. T. V. N.S.W. Q. — B.fl.III,349

D. glaucifolius; F. v. M. in Schlecht. Linnaea XXV, 395 (1852) ...	...	...	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.III,350	M.fr.IX,47.
D. glandulosus; F. v. M. in Proc. Roy. Soc. Tasm. III, 238 (1857) ...	...	...	—	—	—	—	—	—	Q.	N.A.	B.fl.III,350
D. incisus; Hooker, Bot. Magaz. 2875 (1828) ...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.III,350	
D. procumbens; F. v. M. in Proc. Roy. Soc. Tasm. III, 237 (1857) ...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.III,350	M.fr.IX,47.
D. hemicarpus; F. v. M. in Transact. Bot. Soc. Edinb. VII, 491 (1857) ...	...	...	—	—	—	—	—	—	N.A.	B.fl.III,351	M.fr.IX,47.
D. humilis; J. Hooker in icon. pl. t. 304 (1841) ...	...	...	—	—	—	T.	V.	N.S.W.	—	—	B.fl.III,351

TRACHYMENE; Rudge in Transact. Linn. Soc. X, 300 (1810). (Siebera, Fischera, Platysace, Platycarpidium.)

T. compressa; Rudge in Transact. Linn. Soc. X, 300 (1810) implied ...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,352	
T. juncea; Bunge in Lehm. pl. Preiss. I, 236 (1844) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,353	
T. haplosciadea; F. v. M., Cens. 62 (1882) from Benthams (1866) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,353	
T. cirrosa; F. v. M., Cens. 62 (1882) from Bunge (1844) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,354	M.fr.I,231.
T. heterophylla; F. v. M., First Gen. Rep. 13 (1853) ...	...	...	—	S.A.	—	V.	—	—	—	B.fl.III,354	
T. tenuissima; F. v. M., Cens. 62 (1866) from Benthams (1866) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,354	
T. dissecta; F. v. M., Cens. 62 (1882) from Benthams (1866) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,354	
T. commutata; Turczaninow in Bull. Mosc. XXII, 30 (1849) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,355	
T. effusa; Turczaninow in Bull. Mosc. XXII, 31 (1849) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,355	M.fr.I,231.
T. deflexa; Turczaninow in Bull. Mosc. XXII, 31 (1849) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,355	
T. valida; F. v. M., Cens. 62 (1882) ...	...	...	—	—	—	—	—	Q.	—	B.fl.III,355	M.fr.I,232.
T. ericoides; Sieber in Spreng. syst. veg. eur. post. 116 (1827) ...	...	...	—	—	—	V.	N.S.W.	Q.	—	B.fl.III,356	
T. linearis; Sprengel, spec. umbellif. 7 (1818) ...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.III,356	
T. Billardieri; F. v. M., Cens. 62 (1882) from Benthams (1866) ...	...	...	—	—	—	V.	N.S.W.	Q.	—	B.fl.III,356	
T. Stephensonii; Turczaninow in Bull. de Mosc. XX, 170 (1847) ...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.III,357	

XANTHOSIA; Rudge in Transact. Linn. Soc. X, 361 (1810). (Leucolaena, Schoenolaena, Pentapeltis.)

X. juncea; Benthams, Fl. Austr. III, 359 (1866) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,359	
X. tenuior; Benthams, Fl. Austr. III, 359 (1866) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,359	
X. peltigera; Steudel, nomencl. bot. 791 (1841) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,359	
X. hederifolia; Benthams, Fl. Austr. III, 359 (1866) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,359	
X. candida; Steudel, nomencl. bot. 790 (1841) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,259	
X. tridentata; De Candolle, prodr. IV, 75 (1830) ...	...	...	—	—	—	V.	N.S.W.	—	—	B.fl.III,260	
X. singuliflora; F. v. M., fragm. IV, 184 (1864) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,260	M.fr.IV
X. pilosa; Rudge in Transact. Linn. Soc. X, 301, t. 22 (1811) ...	...	...	—	—	T.	V.	N.S.W.	Q.	—	B.fl.III,260	
X. ciliata; Hooker, icon. pl. t. 726 (1848) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,261	
X. pusilla; Bunge in Lehm. pl. Preiss. I, 291 (1844) ...	...	...	W.A.	S.A.	T.	V.	—	—	—	B.fl.III,261	
X. fruticulosa; Benthams, Fl. Austr. III, 361 (1866) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,261	
X. Huegelii; Steudel, nomencl. bot. 791 (1841) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,262	
X. dissecta; J. Hooker in icon. plant. t. 302 (1841) ...	...	...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,362	
X. peduncularis; Benthams, Fl. Austr. III, 362 (1866) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,362	
X. vestita; Benthams, Fl. Austr. III, 363 (1866) ...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.III,363	
X. Atkinsoniana; F. v. M., fragm. II, 127 (1861) ...	...	...	W.A.	—	—	V.	N.S.W.	—	—	B.fl.III,363	M.fr.II,127.
X. rotundifolia; De Candolle, prodr. IV, 75 (1830) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,633	M.fr.II,128.

AZORELLA; Lamarek, Encycl. Méthod. I, 344, t. 189 (1783). (Fragosa, Pozoa, Microsciadium, Schizeilema, Oschatzia, Centella partly, Dichopetalum.)

A. Muelleri; Benthams, Fl. Austr. III, 364 (1866) ...	...	...	—	—	—	V.	—	—	—	B.fl.III,364	
A. cuneifolia; F. v. M. in Transact. phil. Inst. Vict. I, 103 (1855) ...	...	...	—	—	—	V.	—	—	—	B.fl.III,365	
A. saxifraga; Benthams, Fl. Austr. III, 365 (1866) ...	...	...	—	—	T.	—	—	—	—	B.fl.III,365	
A. dichopetala; Benthams, Fl. Austr. III, 365 (1866) ...	...	...	—	—	T.	V.	—	—	—	B.fl.III,365	

HUANACA; Cavanilles, icon. VI, 18, t. 528 (1801). (Diapaspis, Pozoopsis.)

H. hydrocotylea; Benthams & J. Hooker, gen. pl. I, 877 (1867) ...	...	...	—	—	T.	V.	—	—	—	B.fl.III,366	
H. cordifolia; Benthams & J. Hooker, gen. pl. I, 877 (1867) ...	...	...	—	—	T.	—	—	—	—	B.fl.III,366	

ACTINOTUS ; Labillardière, Nov. Holl. pl. spec. I, 67, t. 92 (1804). (Hemiphues, Eriocalia, Holotome.)

A. Helianthi ; Labillardière, Nov. Holl. pl. spec. I, 67, t. 92 (1804)	...	...	—	—	—	V.	N.S.W.	Q.	—	B.f.III,367			
A. Schwarzii ; F. v. M. in proceed. R. Soc. of S.A. IX, 7 & 200 (1886)	...	...	—	S.A.	—	—	—	—	—	—			
A. leucocephalus ; Benth in Hueg. enum. 56 (1837)	...	...	...	W.A.	—	—	—	—	—	B.f.III,368	M.fr.II,130.		
A. minor ; De Candolle, prodr. IV, 83 (1830)...	...	...	...	—	—	—	—	N.S.W.	—	B.f.III,368	M.fr.VI,23.		
A. Gibbonsii ; F. v. M., fragm. VI, 23 (1867)	...	...	...	—	—	—	V.	N.S.W.	—	—	M.fr.VI,23.		
A. omnifertilis ; Benth, Fl. Austr. III, 368 (1866)	...	...	...	W.A.	—	—	—	—	—	B.f.III,368	M.fr.II,129.		
A. rhomboideus ; Benth, Fl. Austr. III, 368 (1866)	...	...	...	W.A.	—	—	—	—	—	B.f.III,368			
A. bellidioides ; Benth, Fl. Austr. III, 369 (1866)	...	...	...	—	—	T.	—	—	—	B.f.III,369			
A. glomeratus ; Benth, Fl. Austr. III, 369 (1866)	...	...	...	W.A.	—	—	—	—	—	B.f.III,369	M.fr.VI,23.		

ERYNGIUM ; Tournefort, inst. 327, t. 173 (1700) from Theophrastos and Dioscorides.

E. rostratum ; Cavanilles, icon. pl. VI, 35, t. 552 (1801)	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.f.III,370	...	...	Am.
E. vesiculosum ; Labillardière, Nov. Holl. pl. spec. I, 73, t. 98 (1804)	...	...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.f.III,370	M.fr.VII,148	...	Am., N.Z.	
E. plantagineum ; F. v. M. in Pap. Roy. Soc. Tasm. III, 235 (1857)	...	...	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.f.III,371	M.fr.VII,148.			
E. expansum ; F. v. M. in Pap. Roy. Soc. Tasm. III, 236 (1857)	..	...	...	—	—	—	N.S.W.	Q.	—	B.f.III,371	M.fr.VII,148.			

APIUM ; Tournefort, inst. 305, t. 160 (1700).

A. prostratum ; Labillardière, relat. du voy. I, 141 (1799)	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.f.III,372	M.fr.VII,148	...	As., Afr., N.Z., P.
A. leptophyllum ; F. v. M. in Benth. Fl. Austr. III, 372 (1866)	...	...	...	—	—	—	V.	N.S.W.	Q.	—	B.f.III,372	M.fr.IV,184; VII,148.	...	Am., P.

SIUM ; Tournefort, inst. 308, t. 162 (1700) from C. Bannin (1623). (Helosciadium).

S. latifolium ; C. Bauhin, pinax 154 (1623)	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.f.III,336	...	...	E., As.
S. erectum ; Hudson, fl. Angl. p. 103 (1762)	...	...	...	—	—	—	N.S.W.	—	—	B.f.III,336	...	...	...	E., As., Am.

SESELI ; Rivinus in Rupp. Fl. Jen. 267 (1718).

S. Harveyanum ; F. v. M. in Transact. phil. Soc. Vict. I, 104 (1855)	...	...	—	—	—	V.	N.S.W.	—	—	B.f.III,373				
S. algens ; F. v. M. in Transact. phil. Soc. Vict. I, 104 (1855)	...	...	—	—	—	V.	N.S.W.	—	—	B.f.III,373				

CRANTZIA ; Nuttall, Gen. of North Amer. plants I, 177 (1818). (Non Scopoli, 1777, Crantziola, F. v. M.)

C. lineata ; Nuttall, Gen. of North Amer. pl. I, 178 (1818)	..	...	...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.f.III,374	...	...	Am., N.Z.
---	----	-----	-----	---	------	----	----	--------	----	---	-------------	-----	-----	-----------

OENANTHE ; Tournefort, inst. 312, t. 166 (1700) from l'Obel (1576).

O. stolonifera ; Wallich, numerical list 585 (1825)	...	...	...	—	—	—	—	—	Q.	—	M.fr.V,182	...	...	As.
---	-----	-----	-----	---	---	---	---	---	----	---	------------	-----	-----	-----

ACIPHYLLA ; R. & G. Forster, char. gen. 135, t. 68 (1776). (Anisotome, Calosciadium, Gingidium partly, Ligustici subgenus.)

A. simplicifolia ; F. v. M. in Benth. Fl. Austr. III, 375 (1866)	...	...	...	—	—	—	V.	N.S.W.	—	—	B.f.III,375			
A. glacialis ; F. v. M. in Benth. Fl. Austr. III, 375 (1866)...	...	...	...	—	—	—	V.	N.S.W.	—	—	B.f.III,375			
A. procumbens ; F. v. M. in Benth. Fl. Austr. III, 375 (1866)	...	...	...	—	—	T.	—	—	—	—	B.f.III,375			

DAUCUS ; Tournefort, inst. 307, t. 161 (1700) from l'Ecluse (1576).

D. brachiatus ; Sieber in De Candolle, prodr. IV, 214 (1830)	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.f.III,376	...	..	Am., N.Z.
--	-----	-----	-----	------	------	----	----	--------	----	------	-------------	-----	----	-----------

OREOMYRRHIS ; Endlicher, gen. plantar. 787 (1839). (Caldasia.)

O. andicola ; Endlicher, gen. plantar. 787 (1839)	...	...	...	—	S.A.	T.	V.	N.S.W.	—	—	B.f.III,377	M.fr.VIII,185	...	Am., N.Z.
O. pulvinifera ; F. v. M., fragm. VIII, 185 (1874)	...	...	...	—	—	—	V.	N.S.W.	—	—	—	M.fr.VIII,185.		

SYNPETALEAE PERIGYNAE.

F. v. M. in Woolls's plants of the neighb. of Sydney, 27 (1880).

ELAEAGNEAE.

R. BROWN, prodr. I, 350 (1810).—Genus 1, Species 1.

ELAEAGNUS; Tournefort, coroll. 53, t. 459 (1703) from Camerarius (1586).

E. latifolia; Linné, spec. plant. 121 (1753) — — — — Q. — B.fl.VI,39 M.fr.IV,146; IX,190 .. As.

OLACINEAE.

MIRBEL in Bull. de la Soc. philomat. 377 (1813).—Genera 9, Species 15.

XIMENIA; Plumier, nova plant. Amer. genera 6, t. 21 (1703).

X. Americana; Linné, spec. plant. 1193 (1753) — — — — — Q. N.A. B.fl.I,391 M.fr.VI,3; IX,150 ... As., Afr., Am., P.

OLAX; Linné, amoen. acad. 387 (1747). (Spermaxyrum, Lopadocalyx.)

O. phyllanthi; R. Brown, prodr. 358 (1810) W.A. — — — — — B.fl.I,392 M.fr.IX,151.

O. retusa; F. v. M. in Benth. Fl. Austr. I, 392 (1863) — — — — N.S.W. Q. — B.fl.I,392 M.fr.IX,151.

O. stricta; R. Brown, prodr. 358 (1810) — — — — V. N.S.W. Q. — B.fl.I,392 M.fr.IX,151.

O. Benthamiana; Miquel in Lehm. pl. Preiss. I, 228 (1844) W.A. S.A. — — — — N.A. B.fl.I,393 M.fr.IX,151.

O. aphylla; R. Brown, prodr. 358 (1810) — — — — — Q. N.A. B.fl.I,393 M.fr.IX,151.

CANSJERA; A. L. de Jussieu, gen. pl. 448 (1789).

C. leptostachya; Benth in Hook. Lond. Journ. of Bot. II, 231 (1843) — — — — — Q. N.A. B.fl.I,394 M.fr.VI,3 P.

OPILIA; Roxburgh, pl. Corom. II, 31, t. 158 (1799).

O. amentacea; Roxburgh, pl. Corom. II, 31, t. 158 (1799) — — — — — Q. N.A. B.fl.I,394 M.fr.VI,4; IX,150 ... As., Afr.

PHLEBOCALYMMA; Griffith in Benth. & J. Hook. gen. pl. I, 353 (1862). (Phlebocalymna.)

P. lobospora; F. v. M., fragm. IX, 151 (1875) — — — — — Q. — — M.fr.IX,151.

PENNANTIA; R. & G. Forster, char. gen. 133, t. 67 (1776).

P. Cunninghamii; Miers in Ann. of nat. hist. sec. ser. IX, 491 (1852) — — — — N.S.W. Q. — B.fl.I,395 M.fr.VI,4; IX,150.

P. Endlicheri; Reisseck in Schlecht. Linnaea XVI, 341, t. 13 (1843) — — — — N.S.W. — — — M.fr.IX,150.

APODYTES; E. Meyer in Hook. Journ. of Bot. III, 155 (1841).

A. brachystylis; F. v. M., fragm. IX, 149 (1875) — — — — — Q. — — M.fr.IX,149.

VILLARESIA; Ruiz & Pavon, flor. Peruv. et Chil. prodr. 35 (1794).

V. Moorei; F. v. in Benth. Fl. Austr. I, 396 (1863)... .. — — — — N.S.W. — — B.fl.I,395

V. Smythii; F. v. M., fragm. V, 156 (1866) — — — — — Q. — — M.fr.IX,150.

GOMPHANDRA; Wallich, Catal. n. 3718 & 7204 (1832).

G. Australiana; F. v. M., fragm. VI, 3 (1877) — — — — — Q. — — M.fr.VI,3&253; IX,150.

BALANOPHOREAE.

L. CL. RICHARD in Mém. du Mus. VIII, 404 & 429 (1822).—Genus 1, Species 1.

BALANOPHORA; R. & G. Forster, char. gen. 99, t. 50 (1776).

B. fungosa; R. & G. Forster, char. gen. 99, t. 50 (1776) — — — — — Q. — B.fl.VI,232 P.

SANTALACEAE.

R. BROWN, prodr. I, 350 (1810).—Genera 7, Species 43.

EXOCARPOS; Labillardière, Relat. du voy. à la rech. de La Pérouse I, 155 (1789). (Exocarpus.)

<i>E. latifolia</i> ; R. Brown, prodr. 356 (1810)	...	...	...	...	...	—	—	—	—	N.S.W.	Q. N.A.	B.f. VI, 228	M.fr. VIII, 10	...	...	As., P.
<i>E. odorata</i> ; A. de Candolle, prodr. XIV, 689 (1856)	...	...	...	...	...	W.A.	—	—	—	—	—	B.f. VI, 228	M.fr. VIII, 10.			
<i>E. cupressiformis</i> ; Labillardière, voy. I, 155, t. 14 (1798)	...	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.f. VI, 229			
<i>E. spartea</i> ; R. Brown, prodr. 356 (1810)	...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q. N.A.	B.f. VI, 229	M.fr. VIII, 10; IX, 20.			
<i>E. apbylla</i> ; R. Brown, prodr. 357 (1810)	...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.f. VI, 230	M.fr. VIII, 10; IX, 169; XI, 20.		
<i>E. homaloclada</i> ; C. Moore & F. v. M., fragm. VIII. 9 (1871)	...	...	...	...	...	—	—	—	—	N.S.W.	—	—	B.f. VI, 230	M.fr. VIII, 9.		
<i>E. phyllanthoides</i> ; Endlicher, prodr. pl. ins. Norf. 46 (1833)	...	...	...	...	...	—	—	—	—	N.S.W.	—	—	—	M.fr. IX, 169.		
<i>E. stricta</i> ; R. Brown, prodr. 357 (1810)	...	...	...	...	...	—	S.A.	T.	V.	N.S.W.	—	—	B.f. VI, 230	M.fr. IX, 169.		
<i>E. humifusa</i> ; R. Brown, prodr. 356 (1810)	...	...	...	...	...	—	—	T.	—	—	—	—	B.f. VI, 231			
<i>E. nana</i> ; J. Hooker in Lond. Journ. of Bot. VI, 281 (1847)	...	...	...	...	...	—	—	T.	V.	—	—	—	B.f. VI, 230			

ANTHOBOLUS; R. Brown, prodr. 357 (1810).

A. filifolius; R. Brown, prodr. 357 (1810)	...	...	...	...	...	...	—	—	—	—	Q. N.A.	B.fl. VI, 226	M, fr. I, 213; IX, 3.	
A. triqueter; R. Brown, prodr. 357 (1810)	...	...	...	...	...	...	—	—	—	—	Q.	—	B.fl. VI, 226	
A. foveolatus; F. v. M., fragm. I, 212 (1859)	...	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl. VI, 226	M. fr. II, 181; IX, 3; XI, 20.
A. leptomerioides; F. v. M., fragm. I, 21 (1858)	...	...	...	...	...	...	—	—	—	—	Q.	—	B.fl. VI, 227	M. fr. IX, 3.
A. exocarpoides; F. v. M., fragm. IX, 3 (1875)	...	...	...	...	...	...	—	S.A.	—	—	—	—	—	M. fr. IX, 3.

OMPHACOMERIA; Endlicher, gen. pl. 326 (1838).

O. acerba; A. de Candolle, prodr. XIV, 681 (1856)... — — — V. N.S.W. — — B.fl.VI,224

LEPTOMERIA ; R. Brown, prodr. 353 (1810).

<i>L. spinosa</i> ; A. de Candolle, prodr. XIV, 678 (1856)	...	...	...	...	W.A.	—	—	—	—	—	—	B.f. VI, 220	M.fr. VIII, 10.
<i>L. Preissiana</i> ; A. de Candolle, prodr. XIV, 678 (1856)	...	...	...	...	W.A.	—	—	—	—	—	—	B.f. VI, 221	M.fr. VIII, 10.
<i>L. pauciflora</i> ; R. Brown, prodr. 354 (1810)	...	...	...	...	W.A.	—	—	—	—	—	—	B.f. VI, 221	M.fr. VIII, 10.
<i>L. scrobiculata</i> ; R. Brown, prodr. 354 (1810)	...	...	...	...	W.A.	—	—	—	—	—	—	B.f. VI, 221	M.fr. VIII, 10.
<i>L. acida</i> ; R. Brown, prodr. 353 (1810)	...	...	...	...	—	—	—	V.	N.S.W.	Q.	—	B.f. VI, 222	M.fr. VIII, 10.
<i>L. Billardieri</i> ; R. Brown, prodr. 354 (1810)	...	...	...	...	—	—	T.	—	N.S.W.	Q.	—	B.f. VI, 222	M.fr. VIII, 10.
<i>L. aphylla</i> ; R. Brown, prodr. 354 (1810)	...	...	...	...	—	S.A.	—	V.	N.S.W.	—	—	B.f. VI, 222	M.fr. VIII, 10.
<i>L. glomerata</i> ; F. v. M. in J. Hook. Fl. Tasm. II, 370 (1860)	...	...	...	...	—	—	T.	—	—	—	—	B.f. VI, 223	M.fr. VIII, 10.
<i>L. squarrosa</i> ; R. Brown, prodr. 354 (1810)	...	...	...	...	W.A.	—	—	—	—	—	—	B.f. VI, 223	M.fr. VIII, 11.
<i>L. Cunninghamii</i> ; Miquel in Lehm. pl. Preiss. I, 611 (1845)	...	...	...	...	W.A.	—	—	—	—	—	—	B.f. VI, 223	M.fr. VIII, 10.
<i>L. empetriformis</i> ; Miquel in Lehm. pl. Preiss. I, 610 (1845)	...	...	...	...	W.A.	—	—	—	—	—	—	B.f. VI, 224	
<i>L. axillaris</i> ; R. Brown, prodr. 354 (1810)	...	...	...	...	W.A.	—	—	—	—	—	—	B.f. VI, 224	
<i>L. laxa</i> ; Miquel in Lehm. pl. Preiss. I, 612 (1845)	...	...	...	...	W.A.	—	—	—	—	—	—	B.f. VI, 224	
<i>L. obovata</i> ; Miquel in Lehm. pl. Preiss. I, 613 (1845)	...	...	...	...	W.A.	—	—	—	—	—	—	B.f. VI, 224	

CHORETRUM ; R. Brown, prodr. 354 (1810).

<i>C. glomeratum</i> ; R. Brown, prodr. 354 (1810)	...	...	...	...	W.A. S.A.	—	V. N.S.W.	—	—	B.fl.VI,218	M.fr.I,21.	
<i>C. chrysanthum</i> ; F. v. M. in Transact. phil. Soc. Vict. I, 23 (1854)	...	...	...	...	—	S.A.	—	V. N.S.W.	—	—	B.fl.VI,218 M.fr.I,21; VIII,11; XI,138.	
<i>C. spicatum</i> ; F. v. M., fragm. I, 21 (1858)	...	...	...	...	—	S.A.	—	V. N.S.W.	—	—	B.fl.VI,218 M.fr.I,21; VIII,11.	
<i>C. lateriflorum</i> ; R. Brown, prodr. 354 (1810)...	...	...	...	...	—	—	—	V. N.S.W.	Q.	—	B.fl.VI.219 M.fr.I,21.	
<i>C. Candollei</i> ; F. v. M. in Benthani's Fl. Austr. VI, 219 (1873)	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.VI.219

SANTALUM; Linné, gen. ed. sec. 165 (1742) from C. Bauhin (1623). (Fusanus, Eucarya.)

<i>S. lanceolatum</i> ; R. Brown, prodr. 356 (1810)	...	...	...	...	...	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl. VI, 214	M.fr. I, 85; VIII, 11; XI, 3.
<i>S. ovatum</i> ; R. Brown, prodr. 355 (1810)	...	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl. VI, 214	
<i>S. obtusifolium</i> ; R. Brown, prodr. 356 (1810)	...	...	...	...	...	—	—	—	V.	N.S.W.	—	—	B.fl. VI, 215	M.fr. VIII, 11.

<i>S. acuminatum</i> ; A. de Candolle, prodr. XIV, 684 (1856)	W.A. S.A. — V. N.S.W. Q. — B.fl.VI,216 M.fr.I,85; IX,169; XI,20.
<i>S. persicarium</i> ; F. v. M. in Transact. Vict. Inst. 41 (1854)	W.A. S.A. — V. N.S.W. Q. — B.fl.VI,216 M.fr.I,86; IX,3; XI,20.
<i>S. cygnorum</i> ; Miquel in Lehm. pl. Preiss. I, 615 (1845)	W.A. — — — — — B.fl.VI,217 M.fr.VIII,11.
<i>S. crassifolium</i> ; A. de Candolle, prodr. XIV, 685 (1856)	— — — — — N.S.W. — — B.fl.VI,217

THESIUM; Linné, gen. pl. 60 (1737).

<i>T. Australe</i> ; R. Brown, prodr. 353 (1810)	— — T. V. N.S.W. Q. — B.fl.VI,212 M.fr.IX,3 As.
---	--

LORANTHACEAE.

A. L. DE JUSSIEU in Ann. du Mus. XII, 292 (1808).—Genera 5, Species 27.

VISCUM; Tournefort, inst. 609, t. 380 (1700), from *Camerarius* (1586), from W. Turner (1538).

<i>V. orientale</i> ; Willdenow, spec. plant. IV, 737 (1805)	— — — — — Q. — B.fl.III,396 As., P.
<i>V. angulatum</i> ; Heyne in De Candolle, prodr. IV, 283 (1830)	— — — — — N.S.W. Q. N.A. B.fl.III,396 As.
<i>V. articulatum</i> ; N. L. Burmann, Fl. Ind. 311 (1768)	— S.A. — — — N.S.W. Q. — B.fl.III,396 As., P.

NOTOTHIXOS; Oliver in Journ. Linn. Soc. VII, 104 (1865).

<i>N. incaus</i> ; Oliver in Journ. Linn. Soc. VII, 104 (1865)	— — — — — V. N.S.W. Q. — B.fl.III,397 M.fr.II,109,176; IV,173.
---	--

LORANTHUS; Linné, syst. nat. ed. sec. 22 (1740).

<i>L. celastroides</i> ; Sieber in Roem. and Schult. syst. veg. VII, 163 (1829)	— — — — — V. N.S.W. Q. — B.fl.III,389 M.fr.VI,252.
<i>L. Bidwillii</i> ; Benth. Fl. Austr. III, 390 (1866)	— — — — — Q. — B.fl.III,390
<i>L. myrtifolius</i> ; Cunningham in Benth. Fl. Austr. III, 390 (1866)... ..	— — — — — N.S.W. Q. — B.fl.III,390
<i>L. longiflorus</i> ; Desrousseaux in Lam. Encycl. méth. IV, 598 (1796)	— — — — — N.S.W. Q. N.A. B.fl.III,390 As.
<i>L. angustifolius</i> ; R. Brown in Benth. Fl. Austr. III, 390 (1866)	— — S.A. — — — — — B.fl.III,390
<i>L. dictyophlebus</i> ; F. v. M., Rep. Burdek. Exped. 14 (1858)	— — — — — N.S.W. Q. — B.fl.III,391
<i>L. maytenifolius</i> ; A. Gray, Bot. Wilk. Expl. Exped. I, 739, t. 99 (1854)	— — — — — N.S.W. — — B.fl.III,393
<i>L. alyxifolius</i> ; F. v. M. in Benth. Fl. Austr. III, 391 (1866)	— — — — — N.S.W. Q. — B.fl.III,391
<i>L. odontocalyx</i> ; F. v. M. in Benth. Fl. Austr. III, 391 (1866)	— — — — — Q. N.A. B.fl.III,391
<i>L. linearifolius</i> ; Hooker in Mitchell, Trop. Austr. 102 (1848)	W.A. S.A. — — — — — Q. — B.fl.III,391
<i>L. Murrayi</i> ; Tate in Transact. R. Soc. of S.A. VI, 109 (1883)	— — S.A. — — — — — —
<i>L. Exocarpi</i> ; Behr in Schlecht. Linnæa XX, 624 (1847)	W.A. S.A. — V. N.S.W. Q. N.A. B.fl.III,392
<i>L. acacioides</i> ; Cunningham in Benth. Fl. Austr. III, 392 (1866)	— — — — — — — N.A. B.fl.III,392
<i>L. signatus</i> ; F. v. M. in Benth. Fl. Austr. III, 392 (1866)	— — — — — Q. N.A. B.fl.III,392 As.
<i>L. sanguineus</i> ; F. v. M., fragm. I, 177 (1859)	— — — — — Q. N.A. B.fl.III,393 M.fr.IV,171.
<i>L. bifurcatus</i> ; Benth. Fl. Austr. III, 393 (1866)	— — — — — Q. N.A. B.fl.III,393
<i>L. linophyllus</i> ; Fenzl in Huegel, enum. 56 (1837)	W.A. S.A. — V. N.S.W. Q. N.A. B.fl.III,393
<i>L. gibberulosus</i> ; Tate in Transact. R. S. of S.A. VIII, 71 Oct. (1885)	— — S.A. — — — — — —
<i>L. pendulus</i> ; Sieber in Sprengel, cur. poster. 139 (1827)	W.A. S.A. — V. N.S.W. Q. N.A. B.fl.III,394
<i>L. Quandang</i> ; Lindley in Mitch. Three Exped. II, 69 (1838)	W.A. S.A. — V. N.S.W. Q. N.A. B.fl.III,395
<i>L. grandibracteus</i> ; F. v. M., Rep. Burdek. Exped. 14 (1858)	— — S.A. — — — — — Q. N.A. B.fl.III,395

ATKINSONIA; F. v. M., fragm. V, 34 (1865). (*Gaiadendron* partly.)

<i>A. ligustrina</i> ; F. v. M., fragm. V, 34 (1865)	— — — — — N.S.W. — — B.fl.III,388 M.fr.II,130; VI,252.
---	--

NUYTSIA; R. Brown in Journ. of the geogr. Soc. I, 17 (1831).

<i>N. floribunda</i> ; R. Brown in Journ. geogr. Soc. I, 17 (1831)	W.A. — — — — — B.fl.III,387 M.fr.VI,252.
---	--

PROTEACEAE.

A. L. DE JUSSIEU, gen. plant. 78 (1789).—Genera 33, Species 597.

PETROPHILA; R. Brown in Transact. Linn. Soc. X, 67 (1809).

<i>P. teretifolia</i> ; R. Brown in Transact. Linn. Soc. X, 68 (1809)	W.A. — — — — — B.fl.V,321 M.fr.VI,245.
<i>P. longifolia</i> ; R. Brown, prot. nov. 5 (1830)	W.A. — — — — — B.fl.V,322 M.fr.VI,255.

<i>P. media</i> ; R. Brown, prot. nov. 5 (1830)	W.A.	—	—	—	—	—	—	B.fl. V, 322	M.fr. VI, 255.
<i>P. acicularis</i> ; R. Brown in Transact. Linn. Soc. X, 69 (1809)	W.A.	—	—	—	—	—	—	B.fl. V, 323	M.fr. VI, 245.
<i>P. megalostegia</i> ; F. v. M., fragm. X, 61 (1876)	W.A.	—	—	—	—	—	—	—	M.fr. X, 61.
<i>P. linearis</i> ; R. Brown, prot. nov. 6 (1830)	W.A.	—	—	—	—	—	—	B.fl. V, 323	M.fr. VI, 243.
<i>P. anceps</i> ; R. Brown, prot. nov. 5 (1830)	W.A.	—	—	—	—	—	—	—	M.fr. VI, 243; X, 47.
<i>P. heterophylla</i> ; Lindley, Bot. Regist. XXV, App. 35 (1839)	W.A.	—	—	—	—	—	—	B.fl. V, 324	M.fr. VI, 244.
<i>P. biloba</i> ; R. Brown, prot. nov. 7 (1830)	W.A.	—	—	—	—	—	—	B.fl. V, 324	M.fr. VI, 255.
<i>P. propinqua</i> ; R. Brown, prot. nov. 7 (1830)	W.A.	—	—	—	—	—	—	B.fl. V, 325	—
<i>P. squamata</i> ; R. Brown in Transact. Linn. Soc. X, 70 (1809)	W.A.	—	—	—	—	—	—	B.fl. V, 325	M.fr. VI, 244.
<i>P. colorata</i> ; Meissner in Lehm. pl. Preiss. II, 246 (1847)	W.A.	—	—	—	—	—	—	B.fl. V, 326	—
<i>P. striata</i> ; R. Brown, prot. nov. 6 (1830)	W.A.	—	—	—	—	—	—	B.fl. V, 326	M.fr. VI, 244.
<i>P. divaricata</i> ; R. Brown, prot. nov. 7 (1830)	W.A.	—	—	—	—	—	—	B.fl. V, 326	M.fr. VI, 344.
<i>P. Serruriae</i> ; R. Brown, prot. nov. 6 (1830)	W.A.	—	—	—	—	—	—	B.fl. V, 327	M.fr. VI, 243.
<i>P. inconspicua</i> ; Meissner in Hook. Kew Misc. VII, 68 (1855)	W.A.	—	—	—	—	—	—	B.fl. V, 327	M.fr. VI, 243.
<i>P. trifida</i> ; R. Brown in Transact. Linn. Soc. X, 70 (1809)	W.A.	—	—	—	—	—	—	B.fl. V, 328	M.fr. VI, 244.
<i>P. carduacea</i> ; Meissner in Hook. Kew Misc. IV, 182 (1852)	W.A.	—	—	—	—	—	—	B.fl. V, 328	M.fr. VI, 244.
<i>P. Shuttleworthiana</i> ; Meissner in Lehm. pl. Preiss. II, 246 (1847)	W.A.	—	—	—	—	—	—	B.fl. V, 329	M.fr. VI, 245.
<i>P. macrostachya</i> ; R. Brown, prot. nov. 7 (1830)	W.A.	—	—	—	—	—	—	B.fl. V, 329	M.fr. VI, 245.
<i>P. diversifolia</i> ; R. Brown in Transact. Linn. Soc. X, 70 (1809)	W.A.	—	—	—	—	—	—	B.fl. V, 329	M.fr. VI, 244.
<i>P. biternata</i> ; Meissner in Hook. Kew Misc. VII, 69 (1855)	W.A.	—	—	—	—	—	—	B.fl. V, 330	M.fr. VI, 245.
<i>P. plumosa</i> ; Meissner in Hook. Kew Misc. VII, 69 (1855)	W.A.	—	—	—	—	—	—	B.fl. V, 330	M.fr. VI, 244.
<i>P. ericifolia</i> ; R. Brown, prot. nov. 5 (1830)	W.A.	—	—	—	—	—	—	B.fl. V, 331	M.fr. VI, 243.
<i>P. chrysantha</i> ; Meissner in Hook. Kew Misc. VII, 68 (1855)	W.A.	—	—	—	—	—	—	B.fl. V, 331	M.fr. VI, 243.
<i>P. pedunculata</i> ; R. Brown in Transact. Linn. Soc. X, 70 (1809)	—	—	—	—	N.S.W.	Q.	—	B.fl. V, 332	M.fr. VI, 242.
<i>P. pulchella</i> ; R. Brown in Transact. Linn. Soc. X, 69 (1809)	—	—	—	—	N.S.W.	—	—	B.fl. V, 332	M.fr. VI, 242.
<i>P. sessilis</i> ; Sieber in Roem. and Schult. mant. 262 (1822)	—	—	—	—	N.S.W.	Q.	—	B.fl. V, 332	—
<i>P. fastigiata</i> ; R. Brown in Transact. Linn. Soc. X, 70 (1809)	W.A.	—	—	—	—	—	—	B.fl. V, 333	M.fr. VI, 245.
<i>P. seminuda</i> ; Lindley, Bot. Regist. XXV, App. 34 (1839)	W.A.	—	—	—	—	—	—	B.fl. V, 333	M.fr. VI, 243.
<i>P. circinata</i> ; Kippist in Hook. Kew Misc. VII, 67 (1855)	W.A.	—	—	—	—	—	—	B.fl. V, 333	M.fr. VI, 244.
<i>P. Drummondii</i> ; Meissner in Lehm. pl. Preiss. I, 496 (1845)	W.A.	—	—	—	—	—	—	B.fl. V, 334	M.fr. VI, 243.
<i>P. crispata</i> ; R. Brown, prot. nov. 6 (1830)	W.A.	—	—	—	—	—	—	B.fl. V, 334	—
<i>P. rigida</i> ; R. Brown in Transact. Linn. Soc. X, 69 (1809)	W.A.	—	—	—	—	—	—	B.fl. V, 334	M.fr. VI, 245.
<i>P. multisecta</i> ; F. v. M., fragm. VI, 242 (1868)	—	S.A.	—	—	—	—	—	B.fl. V, 335	M.fr. VI, 242.
<i>P. conifera</i> ; Meissner in Hook. Kew Misc. VII, 67 (1855)	W.A.	—	—	—	—	—	—	B.fl. V, 335	M.fr. VI, 245.
<i>P. semifurcata</i> ; F. v. M. in Bentham Fl. Austr. V, 335 (1870)	W.A.	—	—	—	—	—	—	B.fl. V, 335	M.fr. X, 47.

ISOPOGON; R. Brown in Transact. Linn. Soc. IX, 70 (1809).

<i>I. latifolius</i> ; R. Brown, prot. nov. 8 (1830)	W.A.	—	—	—	—	—	—	B.fl. V, 338	M.fr. VI, 237.
<i>I. cuneatus</i> ; R. Brown in Transact. Linn. Soc. X, 73 (1809)	W.A.	—	—	—	—	—	—	B.fl. V, 339	M.fr. VI, 238.
<i>I. linearis</i> ; Meissner in Hook. Kew Misc. VII, 69 (1855)	W.A.	—	—	—	—	—	—	B.fl. V, 339	M.fr. VI, 236.
<i>I. polycephalus</i> ; R. Brown in Transact. Linn. Soc. X, 73 (1809)	W.A.	—	—	—	—	—	—	B.fl. V, 339	M.fr. VI, 236.
<i>I. attenuatus</i> ; R. Brown in Transact. Linn. Soc. X, 73 (1809)	W.A.	—	—	—	—	—	—	B.fl. V, 340	M.fr. VI, 237.
<i>I. sphaerocephalus</i> ; Lindley, Bot. Regist. XXV, App. 34 (1839)	W.A.	—	—	—	—	—	—	B.fl. V, 340	M.fr. VI, 236.
<i>I. uncinatus</i> ; R. Brown, prot. nov. 8 (1830)	W.A.	—	—	—	—	—	—	B.fl. V, 341	M.fr. VI, 237.
<i>I. buxifolius</i> ; R. Brown in Transact. Linn. Soc. X, 74 (1809)	W.A.	—	—	—	—	—	—	B.fl. V, 341	M.fr. VI, 237.
<i>I. axillaris</i> ; R. Brown in Transact. Linn. Soc. X, 74 (1809)	W.A.	—	—	—	—	—	—	B.fl. V, 351	M.fr. VI, 237.
<i>I. tridens</i> ; F. v. M., fragm. VI, 239 (1868)	W.A.	—	—	—	—	—	—	B.fl. V, 342	M.fr. VI, 239.
<i>I. Baxteri</i> ; R. Brown, prot. nov. 9 (1830)	W.A.	—	—	—	—	—	—	B.fl. V, 342	M.fr. VI, 240.
<i>I. roseus</i> ; Lindley, Bot. Regist. Misc. n. 37 (1842)	W.A.	—	—	—	—	—	—	B.fl. V, 343	M.fr. VI, 240.
<i>I. adenanthoides</i> ; Meissner in Hook. Kew Misc. VII, 69 (1855)	W.A.	—	—	—	—	—	—	B.fl. V, 343	M.fr. VI, 241.
<i>I. trilobus</i> ; R. Brown in Transact. Linn. Soc. X, 72 (1809)	W.A.	—	—	—	—	—	—	B.fl. V, 343	M.fr. VI, 239.

I. tripartitus; R. Brown, prot. nov. 8 (1830)...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,344	
I. longifolius; R. Brown in Transact. Linn. Soc. X, 73 (1809)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,344	M.fr.VI,237.
I. Drummondii; Benth. Fl. Austr. V, 344 (1870)...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,344	M.fr.VI,241.
I. heterophyllus; Meissner in Lehm. pl. Preiss. I, 504 (1845)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,345	M.fr.VI,241.
I. villosus; Meissner in De Candolle, prodr. XIV, 277 (1856)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,345	M.fr.VI,241.
I. teretifolius; R. Brown in Transact. Linn. Soc. X, 71 (1809)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,345	M.fr.VI,241.
I. anethifolius; R. Brown in J. Knight, proteac. 94 (1809)	...	...	...	...	W.A.	—	—	—	N.S.W.	—	B.fl.V,346	M.fr.VI,238.
I. petiolaris; Cunningham in R. Brown, prot. nov. 8 (1830)	...	...	...	...	—	—	—	—	N.S.W.	Q.	B.fl.V,346	
I. anemonifolius; R. Brown in J. Knight, prot. 93 (1809)	...	...	...	...	—	—	—	V.	N.S.W.	Q.	B.fl.V,347	M.fr.VI,238; X,90.
I. ceratophyllus; R. Brown in Transact. Linn. Soc. X, 72 (1809)	...	...	...	...	—	S.A.	T.	V.	N.S.W.	—	B.fl.V,347	M.fr.VI,238.
I. asper; R. Brown, prot. nov. 8 (1830)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,348	M.fr.VI,240.
I. crithmifolius; F. v. M., fragm. VI, 239 (1868)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,348	M.fr.VI,239.
I. formosus; R. Brown in Transact. Linn. Soc. X, 72 (1809)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,349	M.fr.VI,240.
I. divergens; R. Brown, prot. nov. 7 (1830)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,349	M.fr.VI,241.
I. scabriusculus; Meissner in Hook. Kew Misc. IV, 182 (1852)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,349	M.fr.VI,240.

ADENANTHOS; Labillardière, Nov. Holl. pl. specim. I, 28, t. 36 (1804).

A. Detmoldi; F. v. M., fragm. VIII, 149 (1874)	...	...	...	...	W.A.	—	—	—	—	—	—	M.fr.VIII,149.
A. barbigerus; Lindley, Bot. Regist. XXV, App. 36 (1839)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,351	M.fr.VI,205; VIII,149.
A. obovatus; Labillardière, Nov. Holl. pl. spec. I, 29, t. 37 (1804)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,352	
A. cuneatus; Labillardière, Nov. Holl. pl. spec. I, 28, t. 36 (1804)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,352	
A. Cunninghami; Meissner in Lehm. pl. Preiss. I, 513 (1845)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,352	
A. pungens; Meissner in Lehm. pl. Preiss. I, 515 (1845)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,353	
A. venosus; Meissner in Hook. Kew Misc. IV, 183 (1852)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,353	
A. Dobsoni; F. v. M., fragm. VI, 204 (1868)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,353	M.fr.VI,204.
A. Forrestii; F. v. M. in Wing's S. Sc. Rec. II, 230 (1882)	...	...	...	...	W.A.	—	—	—	—	—	—	
A. linearis; Meissner in Hook. Kew Misc. IV, 183 (1852)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,354	
A. sericeus; Labillardière, Nov. Holl. pl. specim. I, 29, t. 38 (1804)	...	...	...	...	W.A.	S.A.	—	—	—	—	B.fl.V,354	
A. Meissneri; Lehmann, pl. Preiss. I, 515 (1845)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,354	
A. filifolius; Benth. Fl. Austr. V, 355 (1873)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,355	
A. terminalis; R. Brown in Transact. Linn. Soc. X, 152 (1809)	...	...	...	...	—	S.A.	—	V.	—	—	B.fl.V,355	M.fr.I,158.
A. flavidiflorus; F. v. M., fragm. I, 157 (1859)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,355	M.fr.I,157.
A. apiculatus; R. Brown, prot. nov. 9 (1830)...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,356	

SIMSIA; R. Brown in Transact. Linn. Soc. X, 152 (1809). (Stirlingia.)

S. simplex; F. v. M., Cens. of Austr. pl. 66 (1882) from Lindley (1839)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,357	M.fr.VI,248.
S. abrotanoides; F. v. M., Cens. of Austr. pl. 66 (1882) from Meissner (1845)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,357	
S. teretifolia; F. v. M., Cens. of Austr. pl. 66 (1882) from Meissner (1845)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,357	
S. tenuifolia; R. Brown in Transact. Linn. Soc. X, 152 (1809)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,358	
S. latifolia; R. Brown, prot. nov. 9 (1830)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,358	M.fr.VI,248.

SYNAPHEA; R. Brown in Transact. Linn. Soc. X, 155 (1809).

S. polymorpha; R. Brown in Transact. Linn. Soc. X, 156 (1809)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,360	
S. dilatata; R. Brown in Transact. Linn. Soc. X, 156 (1809)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,360	
S. favosa; R. Brown in Transact. Linn. Soc. X, 156 (1809)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,360	
S. Preissii; Meissner in Lehm. pl. Preiss. I, 529 (1845)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,361	
S. acutifolia; Meissner in Lehm. pl. Preiss. I, 528 (1845)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,361	
S. petiolaris; R. Brown in Transact. Linn. Soc. X, 156 (1809)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,361	
S. decorticans; Lindley, Bot. Regist. XXV, App. 32 (1839)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,362	
S. pinuata; Lindley, Bot. Regist. XXV, App. 32 (1839)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,362	

CONOSPERMUM; Smith in Transact. Linn. Soc. IV, 213 (1798).

C. capitatum; R. Brown in Transact. Linn. Soc. X, 155 (1809)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,365	
C. petiolare; R. Brown, prot. nov. 11 (1830)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,365	

C. teretifolium; R. Brown in Transact. Linn. Soc. X, 155 (1809) ...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,365	M.fr.VI,224.
C. flexuosum; R. Brown, prot. nov. 11 (1830) ...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,365	M.fr.VI,224.
C. acerosum; Lindley, Bot. Regist. XXV, App. 30 (1839) ...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,366	
C. amoenum; Meissner in Lehm. pl. Preiss. I, 522 (1845) ...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,366	M.fr.I,157; VI,224.
C. nervosum; Meissner in Hook. Kew Misc. VII, 71 (1855) ...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,367	M.fr.VI,224.
C. diffusum; Bentham, Fl. Austr. V, 367 (1870) ...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,367	
C. glumaceum; Lindley, Bot. Regist. XXV, App. 30 (1839) ...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,367	
C. ephedroides; Kippist in Hook. Kew Misc. VII, 70 (1855) ...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,368	
C. Toddii; F. v. M., fragm. X, 20 (1876) ...	...	...	...	W.A.	—	—	—	—	—	—	M.fr.X,20,90.
C. polycephalum; Meissner in Lehm. pl. Preiss. II, 249 (1847) ...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,368	
C. coeruleum; R. Brown in Transact. Linn. Soc. X, 154 (1809) ...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,368	M.fr.VI,224.
C. debile; Kippist in Hook. Kew Misc. VII, 70 (1855) ...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,369	
C. scaposum; Bentham, Fl. Austr. V, 369 (1870) ...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,369	
C. Huegelii; R. Brown in Endlicher, nov. stirp. dec. 58 (1839) ...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,369	
C. densiflorum; Lindley, Bot. Regist. XXV, App. 32 (1839) ...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,369	
C. Brownii; Meissner in Lehm. pl. Preiss. II, 249 (1847) ...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,370	
C. longifolium; Smith, Exot. Bot. II, 45, t. 82 (1805) ...	...	...	...	—	—	—	—	N.S.W.	—	B.fl.V,370	
C. tenuifolium; R. Brown in Transact. Linn. Soc. X, 154 (1809) ...	...	...	...	—	—	—	—	N.S.W.	—	B.fl.V,370	M.fr.VI,223.
C. Mitchellii; Meissner in De Candolle, prodr. XIV, 320 (1856) ...	...	...	...	—	S.A.	—	V.	—	—	B.fl.V,371	
C. sphacelatum; Hooker in Mitchell, Trop. Austr. 342 (1848) ...	...	...	...	—	—	—	—	—	Q. N.A.	B.fl.V,371	M.fr.VI,223.
C. patens; Schlechtendal, Linnaea XX, 587 (1847) ...	...	...	...	—	S.A.	—	V.	N.S.W.	—	B.fl.V,371	M.fr.VI,223.
C. taxifolium; Smith in Rees's Cycl. IX (1808) ...	...	...	...	—	—	T.	V.	N.S.W.	Q.	B.fl.V,372	M.fr.VI,224.
C. ericifolium; Smith in Rees's Cycl. IX (1808) ...	...	...	...	—	—	—	—	N.S.W.	—	B.fl.V,372	
C. ellipticum; Smith in Rees's Cycl. IX (1808) ...	...	...	...	—	—	—	—	N.S.W.	—	B.fl.V,372	
C. distichum; R. Brown in Transact. Linn. Soc. X, 155 (1809) ...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,373	M.fr.VI,224.
C. florihundum; Bentham, Fl. Austr. V, 373 (1870) ...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,373	
C. incurvum; Lindley, Bot. Regist. XXV, App. 30 (1839) ...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,373	
C. brachyphyllum; Lindley, Bot. Regist. XXV, App. 31 (1839) ...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,374	
C. stoechadis; Endlicher in Ann. Wien. Mus. II, 208 (1838) ...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,374	M.fr.XI,20.
C. triplinervium; R. Brown, prot. nov. 11 (1830) ...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,375	
C. bracteosum; Meissner in Lehm. pl. Preiss. I, 518 (1845) ...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,375	M.fr.VI,224.
C. crassinervium; Meissner in Hook. Kew Misc. IV, 184 (1852) ...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,375	
FRANKLANDIA; R. Brown in Transact. Linn. Soc. X, 157 (1809).											
F. fucifolia; R. Brown in Transact. Linn. Soc. X, 157 (1809) ...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,376	M.fr.VI,223; VII,133.
F. triaristata; Bentham, Fl. Austr. V, 377 (1870) ...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,377	
SYMPHYONEMA; R. Brown in Transact. Linn. Soc. X, 157 (1809). (Symphonema.)											
S. montanum; R. Brown in Transact. Linn. Soc. X, 158 (1809) ...	...	...	...	—	—	—	—	N.S.W.	—	B.fl.V,377	M.fr.VI,223.
S. paludosum; R. Brown in Transact. Linn. Soc. X, 158 (1809) ...	...	...	...	—	—	—	—	N.S.W.	—	B.fl.V,378	M.fr.VI,223.
BELLENDENA; R. Brown in Transact. Linn. Soc. X, 156 (1809).											
B. montana; R. Brown in Transact. Linn. Soc. X, 166 (1809) ...	...	...	...	—	—	T.	—	—	—	B.fl.V,378	
AGASTACHYS; R. Brown in Transact. Linn. Soc. X, 158 (1809).											
A. odorata; R. Brown in Transact. Linn. Soc. X, 158 (1809) ...	...	...	...	—	—	T.	—	—	—	B.fl.V,379	M.fr.X,90.
CENARRHENES; Labillardière, Nov. Holl. pl. spec. I, 36, t. 50 (1804).											
C. nitida; Labillardière, Nov. Holl. pl. spec. I, 36, t. 50 (1804) ...	...	...	...	—	—	T.	—	—	—	B.fl.V,380	
PERSOONIA; Smith in Transact. Linn. Soc. IV, 215 (1798). (Linkia 1797, Pentadactylon.)											
P. hakeiformis; Meissner in Hook. Kew Misc. IV, 185 (1852) ...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,383	M.fr.VI,223.
P. teretifolia; R. Brown in Transact. Linn. Soc. X, 160 (1809) ...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,383	M.fr.VI,223.
P. saccata; R. Brown, prot. nov. 12 (1830) ...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,384	M.fr.VI,223.

<i>P. Saundersiana</i> ; Kippist in Hook. Kew Misc. VII, 72 (1855) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 384	
<i>P. diadema</i> ; F. v. M., fragm. X, 46 (1876) ...	...	...	W.A.	—	—	—	—	—	—	—	M.fr. X, 46.
<i>P. comata</i> ; Meissner in Hooker's Kew Misc. VII, 71 (1855) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 385	
<i>P. brachystylis</i> ; F. v. M., fragm. VI, 221 (1868) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 385	
<i>P. falcata</i> ; R. Brown in Transact. Linn. Soc. X, 162 (1809) ...	...	...	—	—	—	—	—	—	Q. N.A.	B.fl. V, 385	M.fr. VI, 222.
<i>P. trinervis</i> ; Meissner in Hook. Kew Misc. IV, 185 (1852) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 386	
<i>P. tortifolia</i> ; Meissner in Hook. Kew Misc. IV, 185 (1852) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 386	
<i>P. angustiflora</i> ; Benth. Fl. Austr. V, 386 (1870) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 386	
<i>P. rudis</i> ; Meissner in Hook. Kew Misc. IV, 185 (1852) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 387	
<i>P. microcarpa</i> ; R. Brown in Transact. Linn. Soc. X, 160 (1809) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 387	M.fr. VI, 222.
<i>P. sulcata</i> ; Meissner in Hook. Kew Misc. IV, 185 (1852) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 387	
<i>P. acicularius</i> ; F. v. M., fragm. VI, 220 (1868) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 388	M.fr. VI, 220.
<i>P. scabrella</i> ; Meissner in Hooker's Kew Misc. VII, 72 (1855) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 388	
<i>P. dillwynioides</i> ; Meissner in Hook. Kew Misc. IV, 185 (1852) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 388	M.fr. VI, 222.
<i>P. striata</i> ; R. Brown, prot. nov. 13 (1830) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 388	M.fr. VI, 223.
<i>P. quinquenervis</i> ; Hooker, icon. plant. t. 425 (1842) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 389	
<i>P. rufiflora</i> ; Meissner in Hooker's Kew Misc. VII, 72 (1855) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 389	
<i>P. scabra</i> ; R. Brown in Transact. Linn. Soc. X, 162 (1809) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 389	
<i>P. graminea</i> ; R. Brown in Transact. Linn. Soc. X, 164 (1809) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 390	M.fr. VI, 222.
<i>P. longifolia</i> ; R. Brown in Transact. Linn. Soc. X, 164 (1809) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 390	M.fr. VI, 222.
<i>P. articulata</i> ; R. Brown in Transact. Linn. Soc. X, 164 (1809) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 390	
<i>P. elliptica</i> ; R. Brown in Transact. Linn. Soc. X, 164 (1809) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 391	M.fr. VI, 222.
<i>P. ferruginea</i> ; Smith, Exot. Bot. II, 47, t. 83 (1805) ...	...	...	—	—	—	—	N.S.W.	—	—	B.fl. V, 391	
<i>P. media</i> ; R. Brown, prot. nov. 16 (1830) ...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl. V, 391	
<i>P. cornifolia</i> ; Cunningham in R. Brown, prot. nov. 16 (1830) ...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl. V, 392	
<i>P. marginata</i> ; R. Brown, prot. nov. 16 (1830) ...	...	...	—	—	—	—	N.S.W.	—	—	B.fl. V, 492	
<i>P. sericea</i> ; Cunningham in R. Brown, prot. nov. 14 (1830) ...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl. V, 393	
<i>P. Mitchellii</i> ; Meissner in Hook. Kew Misc. VII, 73 (1809) ...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl. V, 393	
<i>P. fastigiata</i> ; R. Brown, prot. nov. 13 (1830) ...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl. V, 393	
<i>P. hirsuta</i> ; Persoon, synops. plant. I, 118 (1805) ...	...	...	—	—	—	—	N.S.W.	—	—	B.fl. V, 394	M.fr. VI, 222.
<i>P. Chamaepitys</i> ; Cunningham in Field, N.S.W. 329 (1825) ...	...	...	—	—	—	—	N.S.W.	—	—	B.fl. V, 394	M.fr. VI, 223.
<i>P. arborea</i> ; F. v. M., gen. rep. 18 (1860) ...	...	...	—	—	—	V.	—	—	—	B.fl. V, 394	M.fr. V, 37; VI, 221.
<i>P. salicina</i> ; Persoon, synops. plant. I, 118 (1805) ...	...	...	—	—	—	V.	N.S.W.	—	—	B.fl. V, 395	
<i>P. prostrata</i> ; R. Brown in Transact. Linn. Soc. X, 163 (1809) ...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl. V, 395	
<i>P. lanceolata</i> ; Andrews, Bot. Reposit. t. 74 (1800) ...	...	...	—	—	—	V.	N.S.W.	Q.	—	B.fl. V, 395	M.fr. VI, 222.
<i>P. confertiflora</i> ; Benth. Fl. Austr. V, 396 (1870) ...	...	...	—	—	—	V.	—	—	—	B.fl. V, 396	
<i>P. lucida</i> ; R. Brown in Transact. Linn. Soc. X, 161 (1809) ...	...	...	—	—	—	—	N.S.W.	—	—	B.fl. V, 396	
<i>P. linearis</i> ; Andrews, Bot. Reposit. t. 77 (1800) ...	...	...	—	—	—	V.	N.S.W.	Q.	—	B.fl. V, 397	M.fr. VI, 222.
<i>P. pinifolia</i> ; R. Brown in Transact. Linn. Soc. X, 160 (1809) ...	...	...	—	—	—	—	N.S.W.	—	—	B.fl. V, 397	M.fr. VI, 222; X, 47.
<i>P. Caley</i> ; R. Brown, prot. nov. 13 (1830) ...	...	...	—	—	—	—	N.S.W.	—	—	B.fl. V, 398	
<i>P. ledifolia</i> ; Cunningham in De Candolle, prodr. XIV, 339 (1856) ...	...	...	—	—	—	—	N.S.W.	—	—	B.fl. V, 398	
<i>P. revoluta</i> ; Sieber in Roem. & Schult. mant. 272 (1822) ...	...	...	—	—	—	V.	N.S.W.	—	—	B.fl. V, 398	M.fr. VI, 221.
<i>P. Gunnii</i> ; J. Hooker in Lond. Journ. Bot. VI, 283 (1843) ...	...	...	—	—	—	T.	—	—	—	B.fl. V, 399	
<i>P. mollis</i> ; R. Brown in Transact. Linn. Soc. X, 161 (1809) ...	...	...	—	—	—	—	N.S.W.	—	—	B.fl. V, 399	M.fr. VI, 221.
<i>P. rigida</i> ; R. Brown, prot. nov. 14 (1830) ...	...	...	—	—	—	V.	N.S.W.	—	—	B.fl. V, 399	M.fr. VI, 221; VIII, 149; X, 47.
<i>P. carvifolia</i> ; R. Brown, prot. nov. 13 (1830) ...	...	...	—	—	—	—	N.S.W.	—	—	B.fl. V, 400	
<i>P. oblongata</i> ; Cunningham in R. Brown, prot. nov. 14 (1830) ...	...	...	—	—	—	—	N.S.W.	—	—	B.fl. V, 400	
<i>P. Cunninghamii</i> ; R. Brown, prot. nov. 15, (1830) ...	...	...	—	—	—	—	N.S.W.	—	—	B.fl. V, 400	
<i>P. myrtilloides</i> ; Sieber in Roem. & Schult. mant. 272 (1822) ...	...	...	—	—	—	V.	N.S.W.	—	—	B.fl. V, 401	M.fr. VI, 221.
<i>P. oxycoccoides</i> ; Sieber in Roem. & Schult. mant. 270 (1822) ...	...	...	—	—	—	V.	N.S.W.	—	—	B.fl. V, 401	M.fr. VI, 221.

<i>P. nutans</i> ; R. Brown in Transact. Linn. Soc. X, 162 (1809) — — — — N.S.W. — — B.fl.V,401 M.fr.VI,222.
<i>P. angulata</i> ; R. Brown, prot. nov. 14 (1830) — — — — N.S.W. — — B.fl.V,402
<i>P. virgata</i> ; R. Brown in Transact. Linn. Soc. X, 161 (1809) — — — — N.S.W. Q. — B.fl.V,402
<i>P. Chamaepeuce</i> ; Lhotsky in De Candolle, prodr. XIV, 336 (1856) — — — — V. N.S.W. — — B.fl.V,402 M.fr.VIII,149; X,47.
<i>P. juniperina</i> ; Labillardière, Nov. Holl. pl. spec. I, 33, t. 45 (1804) — — S.A. T. V. N.S.W. — — B.fl.V,403
<i>P. tenuifolia</i> ; R. Brown, prot. nov. 12 (1830) — — — — N.S.W. Q. — B.fl.V,403
<i>P. acerosa</i> ; Sieber in Roem. & Schult. mant. 269 (1822) — — — — N.S.W. — — B.fl.V,403 M.fr.VI,222.

MACADAMIA ; F. v. M. in Transact. phil. Inst. Vict. II, 72 (1857). (Panopsis partly, Andriapetalum partly.)

<i>M. ternifolia</i> ; F. v. M. in Transact. phil. Inst. Vict. II, 72 (1857) — — — — N.S.W. Q. — B.fl.V,406 M.fr.II,91; III,38; VI,191,224; VII,59.
--

HELICIA ; Loureiro, Fl. Cochinch. I, 83 (1790).

<i>H. Youngiana</i> ; C. Moore & F. v. M., fragm. IV, 84 (1864) — — — — N.S.W. Q. — B.fl.V,406 M.fr.V,186.
<i>H. praealta</i> ; F. v. M., fragm. III, 37 (1862) — — — — N.S.W. Q. — B.fl.V,404 M.fr.IV,174; VI,224.
<i>H. Australasica</i> ; F. v. M. in Hook. Kew Misc. IX, 27 (1857) — — — — — N.A. B.fl.V,405 M.fr.II,91.
<i>H. glabriflora</i> ; F. v. M., fragm. II, 91 (1861) — — — — N.S.W. Q. — B.fl.V,405 M.fr.III,38; V,38,186.
<i>H. ferruginea</i> ; F. v. M., fragm. III, 37 (1862) — — — — N.S.W. Q. — B.fl.V,405 M.fr.IV,174; V,186; VI,191,224.

ROUPALA ; Aublet, Hist. des pl. de la Guian. I, 83, t. 32 (1775). (Rhopala, Bleasdalea, Adenostephanus partly, Kermadecia.)

<i>R. Bleasdalei</i> ; F. v. M. in Journ. Roy. Soc. N.S.W. XV, 212 (1881) implied — — — — — Q. — B.fl.V,417 M.fr.V,90.

HICKSBEACHIA ; F. v. M. in Wing's South. Science Record III, 32 (1883).

<i>H. pinnatifolia</i> ; F. v. M. in S. Sc. Rec. III, 33 (1883) — — — — — N.S.W. Q. — —
--

XYLOMELUM ; Smith in Transact. Linn. Soc. IV, 214 (1798).

<i>X. pyriforme</i> ; Smith in J. Knight, prot. 105 (1809) — — — — N.S.W. Q. — B.fl.V,408 M.fr.VI,220.
<i>X. occidentale</i> ; R. Brown, prot. nov. 31 (1830) — — — — W.A. — — — — B.fl.V,408
<i>X. salicinum</i> ; Cunningham in R. Brown, prot. 31 (1830) — — — — — Q. — B.fl.V,408 M.fr.VI,220.
<i>X. Scottianum</i> ; F. v. M., fragm. V, 174 (1866) — — — — — Q. — — M.fr.IV,107; V,215.
<i>X. angustifolium</i> ; Kippist in Hook. Kew Misc. IV, 209 (1852) — — — — W.A. — — — — B.fl.V,409 M.fr.VI,220.

LAMBERTIA ; Smith in Transact. Linn. Soc. IV, 214 (1798).

<i>L. uniflora</i> ; R. Brown in Transact. Linn. Soc. X, 188 (1809) — — — — W.A. — — — — B.fl.V,413 M.fr.VI,248.
<i>L. rariflora</i> ; Meissner in Lehm. pl. Preiss. II, 263 (1847) — — — — W.A. — — — — B.fl.V,414
<i>L. inermis</i> ; R. Brown in Transact. Linn. Soc. X, 188 (1809) — — — — W.A. — — — — B.fl.V,414 M.fr.VI,248.
<i>L. ericifolia</i> ; R. Brown, prot. nov. 30 (1830) — — — — W.A. — — — — B.fl.V,414 M.fr.VI,248.
<i>L. multiflora</i> ; Lindley, Bot. Regist. XXV, App. 32 (1839) — — — — W.A. — — — — B.fl.V,415
<i>L. formosa</i> ; Smith in Transact. Linn. Soc. IV, 214, t. 20 (1798) — — — — — N.S.W. — — B.fl.V,415 M.fr.VII,133.
<i>L. echinata</i> ; R. Brown in Transact. Linn. Soc. X, 189 (1809) — — — — W.A. — — — — B.fl.V,416 M.fr.VI,248.
<i>L. ilicifolia</i> ; Hooker, icon. plant. t. 553 (1843) — — — — W.A. — — — — B.fl.V,416

ORITES ; R. Brown in Transact. Linn. Soc. X, 189 (1809). (Oritina.)

<i>O. excelsa</i> ; R. Brown, prot. nov. 32 (1832) — — — — N.S.W. Q. — B.fl.V,411 M.fr.V,153; VI,223.
<i>O. diversifolia</i> ; R. Brown in Transact. Linn. Soc. X, 190 (1809) — — — — T. — — — — B.fl.V,411 M.fr.V,153.
<i>O. Milligani</i> ; Meissner in Hook. Kew Misc. IV, 209 (1852) — — — — T. — — — — B.fl.V,411
<i>O. lancifolia</i> ; F. v. M. in Transact. phil. Soc. Vict. I, 108 (1855) — — — — V. N.S.W. — — B.fl.V,412 M.fr.V,153; VI,223; VII,133.
<i>O. revoluta</i> ; R. Brown in Transact. Linn. Soc. X, 19 (1809) — — — — T. — — — — B.fl.V,412
<i>O. acicularis</i> ; R. Brown, prot. nov. 32 (1830) — — — — T. — — — — B.fl.V,412

STRANGAEA ; Meissner in Hooker's Kew Misc. VII, 66 (1855). (Molloya.)

<i>S. linearis</i> ; Meissner in Hook. Kew Misc. VII, 66 (1855) — — — — N.S.W. Q. — B.fl.V,453 M.fr.VI,223; VII,132.
<i>S. cynanchocarpa</i> ; F. v. M., fragm. VII, 132 (1871) — — — — W.A. — — — — B.fl.V,453 M.fr.VII,132.

GREVILLIA; R. Brown in Transact. Linn. Soc. X, 168 (1809). (Anadenia, Lyssanthe, Stylurus, Manglesia.)

G. Pinaster; Meissner in Hook. Kew Misc. VII, 76 (1855)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V, 427	M.fr.VI, 212.
G. obtusifolia; Meissner in Hook. Kew Misc. IV, 187 (1852)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V, 427	
G. sparsiflora; F. v. M., fragm. VI, 206 (1868)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V, 428	M.fr.VI, 206.
G. macrostylis; F. v. M., fragm. I, 137 (1859)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V, 428	M.fr.VI, 211.
G. tripartita; Meissner in Hook. Kew Misc. IV, 186 (1852)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V, 428	M.fr.VI, 213.
G. platypoda; F. v. M., fragm. VI, 205 (1868)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V, 428	M.fr.VI, 205.
G. patentiloba; F. v. M., fragm. I, 137 (1859)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V, 429	M.fr.I, 137.
G. pectinata; R. Brown, prot. nov. 23 (1830)...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V, 429	M.fr.VI, 212.
G. plurijuga; F. v. M., fragm. IV, 84 (1864)...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V, 430	M.fr.VI, 212.
G. nudiflora; Meissner in Hook. Kew Misc. IV, 186 (1852)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V, 430	M.fr.I, 35; VI, 212.
G. stenomera; F. v. M., fragm. IV, 85 (1864)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V, 430	M.fr.IV, 85.
G. Thelemanniana; Endlicher, nov. stirp. dec. 6 (1839)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V, 431	M.fr.VI, 212.
G. concinna; R. Brown in Transact. Linn. Soc. X, 172 (1809)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V, 431	M.fr.VI, 212.
G. Hookeriana; Meissner in Lehm. pl. Preiss. I, 546 (1845)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V, 432	M.fr.VI, 212.
G. Baxteri; R. Brown, prot. nov. 22 (1830)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V, 432	
G. eriobotrya; F. v. M., fragm. X, 44 (1876)...	...	...	...	W.A.	—	—	—	—	—	—	—	B.fl.X, 44.
G. pterosperma; F. v. M. in Transact. phil. Soc. Vict. I, 22 (1854)	...	...	...	W.A. S.A.	—	V.	N.S.W.	—	—	—	B.fl.V, 432	M.fr.VI, 210; X, 45.
G. stenobotrya; F. v. M., fragm. IX, 3 (1875)	...	...	...	W.A. S.A.	—	—	—	Q. N.A.	—	—	—	M.fr.IX, 3.
G. eriostachya; Lindley, Bot. Regist. XXV, App. 36 (1839)	...	...	...	—	—	—	—	—	N.A.	B.fl.V, 433	—	M.fr.VI, 208&213; X, 45.
G. thyrsoides; Meissner in Hooker's Kew Misc. VII, 77 (1855)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V, 433	
G. Chrysodendron; R. Brown in Transact. Linn. Soc. X, 176 (1809)	...	...	...	—	—	—	—	Q. N.A.	B.fl.V, 434	—	—	M.fr.VI, 210.
G. Banksii; R. Brown in Transact. Linn. Soc. X, 176 (1809)	...	...	...	—	—	—	—	Q. —	B.fl.V, 434	—	—	M.fr.VI, 210; VIII, 150; X, 46.
G. Caley; R. Brown, prot. nov. 22 (1830)	...	...	...	—	—	—	N.S.W.	—	—	—	B.fl.V, 435	
G. asplenifolia; R. Brown in J. Knight, prot. 120 (1869)	...	...	...	—	—	—	N.S.W.	—	—	—	B.fl.V, 435	M.fr.VI, 212.
G. cirsiifolia; Meissner in Lehm. pl. Preiss. II, 253 (1847)...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V, 436	
G. laurifolia; Sieber in Roem. & Schult. mant. 279 (1822)...	...	...	...	—	—	—	N.S.W.	—	—	—	B.fl.V, 436	
G. Barklyana; F. v. M., gen. report 14 & 18 (1861)...	...	...	...	—	—	—	V.	—	—	—	B.fl.V, 436	M.fr.VII, 133; VIII, 150; X, 46.
G. repens; F. v. M. in Schlecht. Linnaea XXVI, 355 (1853)	...	...	...	—	—	—	V.	—	—	—	B.fl.V, 437	
G. Aquifolium; Lindley in Mitchell, Three Exped. II, 178 (1838)	...	...	...	—	S.A.	—	V.	—	—	—	B.fl.V, 437	
G. ilicifolia; R. Brown, prot. nov. 21 (1830)	...	...	...	—	S.A.	—	V.	—	Q.	—	B.fl.V, 437	M.fr.VI, 212.
G. Scortechinii; F. v. M. in Bailey's synop. of Queensland Flora 435 (1883)	...	...	...	—	—	—	N.S.W.	Q.	—	—	—	
G. Gandichandii; R. Brown in Freyc. Voy. Bot. 443, t. 46 (1826)	...	...	...	—	—	—	N.S.W.	—	—	—	B.fl.V, 438	
G. acanthifolia; Cunningham in Field, N.S.W. 328 (1825)...	...	...	...	—	—	—	N.S.W.	—	—	—	B.fl.V, 438	M.fr.VI, 212; VIII, 150.
G. bipinnatifida; R. Brown, prot. nov. 23 (1830)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V, 439	M.fr.VI, 212.
G. armigera; Meissner in Hook. Kew Misc. IV, 186 (1852)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V, 439	
G. asparagoides; Meissner in Hook. Kew Misc. IV, 186 (1852)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V, 439	M.fr.IX, 123.
G. Treueriana; F. v. M., fragm. IX, 123 (1875)	...	...	...	W.A. S.A.	—	—	—	—	—	—	—	M.fr.IX, 123.
G. floribunda; R. Brown, prot. nov. 19 (1830)	...	...	...	—	—	—	V.	N.S.W.	Q.	—	B.fl.V, 440	
G. cinerea; R. Brown in Transact. Linn. Soc. X, 173 (1809)	...	...	...	—	—	—	N.S.W.	—	—	—	B.fl.V, 440	
G. alpina; Lindley in Mitchell, Three Exped. II, 179 (1838)	...	...	...	—	—	—	V.	—	—	—	B.fl.V, 441	
G. montana; R. Brown in Transact. Linn. Soc. X, 172 (1809)	...	...	...	—	—	—	N.S.W.	—	—	—	B.fl.V, 441	
G. obtusiflora; R. Brown, prot. nov. 19 (1830)	...	...	...	—	—	—	N.S.W.	—	—	—	B.fl.V, 442	
G. arenaria; R. Brown in Transact. Linn. Soc. X, 172 (1809)	...	...	...	—	—	—	N.S.W.	—	—	—	B.fl.V, 442	
G. mucronulata; R. Brown in Transact. Linn. Soc. X, 173 (1809)	...	...	...	—	—	—	N.S.W.	—	—	—	B.fl.V, 443	
G. Baueri; R. Brown in Transact. Linn. Soc. X, 173 (1809)	...	...	...	—	—	—	N.S.W.	—	—	—	B.fl.V, 443	
G. lanigera; Cunningham in R. Brown, prot. nov. 20 (1830)	...	...	...	—	—	—	V.	N.S.W.	—	—	B.fl.V, 444	
G. rosmarinifolia; Cunningham in Field, N. S. Wales, 328 (1825)	...	...	...	—	—	—	V.	N.S.W.	—	—	B.fl.V, 445	
G. Goodii; R. Brown in Transact. Linn. Soc. X, 174 (1809)	...	...	...	—	—	—	—	Q. N.A.	B.fl.V, 446	—	—	M.fr.VI, 211.
G. vennsta; R. Brown in Transact. Linn. Soc. X, 175 (1809)	...	...	...	—	—	—	—	Q.	—	—	B.fl.V, 446	

G. longistyla; Hooker in Mitchell, Trop. Austr. 343 (1848)	...	...	...	—	—	—	N.S.W.	Q. N.A.	B.fl.V,446	M.fr.VI,211.
G. juncifolia; Hooker in Mitchell, Trop. Austr. 341 (1848)	...	...	...	—	S.A.	—	N.S.W.	Q. N.A.	B.fl.V,447	M.fr.VI,209; VIII,150.
G. Wilsoni; Cunningham in Wilson's Voy. 273 (1835)	...	...	...	W.A.	—	—	—	—	B.fl.V,447	M.fr.VI,212; X,45.
G. erectiloba; F. v. M., fragm. X, 44 (1876)	...	...	...	W.A.	—	—	—	—	—	M.fr.X,44.
G. lavandulacea; Schlechtendal, Linnaea XX, 586 (1847)	...	...	...	—	S.A.	—	V. N.S.W.	—	B.fl.V,448	
G. insignis; Kippist in De Candolle, prodr. XIV, 379 (1856)	...	...	...	W.A.	—	—	—	—	B.fl.V,448	M.fr.VIII,150; X,45.
G. Brownii; Meissner in Lehm. pl. Preiss. I, 537 (1845)	...	...	...	W.A.	—	—	—	—	B.fl.V,449	M.fr.VI,213.
G. fasciculata; R. Brown, prot. nov. 20 (1830)	...	...	...	W.A.	—	—	—	—	B.fl.V,449	
G. aspera; R. Brown in Transact. Linn. Soc. X, 172 (1809)	...	...	...	W.A.	S.A.	—	—	—	B.fl.V,450	M.fr.VII,211.
G. brachystylis; Meissner in Lehm. pl. Preiss. I, 538 (1845)	...	...	...	W.A.	—	—	—	—	B.fl.V,450	
G. faccata; Bentham, Fl. Austr. V, 450 (1870)	...	...	...	W.A.	—	—	—	—	B.fl.V,450	
G. Drummondii; Meissner in Lehm. pl. Preiss. I, 536 (1845)	...	...	...	W.A.	—	—	—	—	B.fl.V,451	
G. deflexa, F. v. M. in Melb. Chemist 57, p. 72 (1883)	...	...	...	W.A.	—	—	—	—	—	
G. disjuncta; F. v. M., fragm. VI, 206 (1868)	...	...	...	W.A.	—	—	—	—	B.fl.V,451	M.fr.VI,206.
G. haplantha; F. v. M. in Bentham, Fl. Austr. V, 451 (1870)	...	...	...	W.A.	—	—	—	—	B.fl.V,452	
G. pinifolia; Meissner in Hook. Kew Misc. 186 (1852)	...	...	...	W.A.	—	—	—	—	B.fl.V,452	
G. Kennediana; F. v. M. in proceed. R.S. Vict. XXIV, p. 172 (1887)	...	...	...	—	—	—	N.S.W.	Q.	—	
G. acuaria; F. v. M. in Bentham, Fl. Austr. V, 452 (1870)	...	...	...	W.A.	—	—	—	—	B.fl.V,452	
G. singuliflora; F. v. M., fragm. VI, 92 (1868)	...	...	...	—	—	—	—	Q.	B.fl.V,452	M.fr.VI,92.
G. pauciflora; R. Brown in Transact. Linn. Soc. X, 171 (1809)	...	...	...	W.A.	S.A.	—	—	—	B.fl.V,453	M.fr.I,135; VI,206,208.
G. quercifolia; R. Brown, prot. nov. 23 (1830)	...	...	...	W.A.	—	—	—	—	B.fl.V,454	M.fr.VI,213.
G. angulata; R. Brown, prot. nov. 24 (1830)	...	...	...	—	S.A.	—	—	N.A.	B.fl.V,455	
G. Wickhami; Meissner in Hook. Kew Misc. 187 (1852)	...	...	...	W.A.	S.A.	—	—	Q. N.A.	B.fl.V,455	
G. agrifolia; Cunningham in R. Brown, prot. nov. 24 (1830)	...	...	...	—	—	—	—	N.A.	B.fl.V,455	
G. Cunninghamii; R. Brown, prot. nov. 23 (1830)	...	...	...	—	—	—	—	N.A.	B.fl.V,456	
G. pungens; R. Brown in Transact. Linn. Soc. X, 175 (1809)	...	...	...	—	—	—	—	N.A.	B.fl.V,456	
G. Huegelii; Meissner in Lehm. pl. Preiss. I, 543 (1845)	...	...	...	W.A.	S.A.	—	V. N.S.W.	—	B.fl.V,457	M.fr.VI,212; X,46.
G. dimidiata; F. v. M., fragm. III, 146 (1863)	...	...	...	—	—	—	—	N.A.	B.fl.V,457	M.fr.III,146.
G. heliosperma; R. Brown in Transact. Linn. Soc. X, 176 (1809)	...	...	...	—	—	—	—	N.A.	B.fl.V,457	M.fr.VI,210.
G. refracta; R. Brown in Transact. Linn. Soc. X, 176 (1809)	...	...	...	—	—	—	—	Q. N.A.	B.fl.V,458	
G. Dryandri; R. Brown in Transact. Linn. Soc. X, 175 (1809)	...	...	...	—	—	—	—	Q. N.A.	B.fl.V,458	M.fr.VI,209.
G. polystachya; R. Brown in Transact. Linn. Soc. X, 177 (1809)	...	...	...	—	—	—	—	Q. N.A.	B.fl.V,459	M.fr.VI,210.
G. robusta; Cunningham in R. Brown, prot. nov. 24 (1830)	...	...	...	—	—	—	N.S.W.	Q.	B.fl.V,459	M.fr.VI,210; VII,133.
G. annulifera; F. v. M., fragm. IV., 85 (1864)	...	...	...	W.A.	—	—	—	—	B.fl.V,460	M.fr.IV,85; VI,212; VII,132; XI,20.
G. leucopteris, Meissner in Hook. Kew Misc. VII, 76 (1855)	...	...	...	W.A.	—	—	—	—	B.fl.V,460	M.fr.III,145; IV,85,176; XI,20.
G. Leucadendron; Cunningham in R. Brown, prot. nov. 25 (1830)	...	...	...	—	—	—	—	N.A.	B.fl.V,461	M.fr.I,136-7.
G. pyramidalis; Cunningham in R. Brown, prot. nov. 25 (1830)	...	...	...	—	—	—	—	N.A.	B.fl.V,462	
G. striata; R. Brown in Transact. Linn. Soc. X, 177 (1809)	...	...	...	W.A.	S.A.	—	N.S.W.	Q. N.A.	B.fl.V,462	
G. mimosoides; R. Brown in Transact. Linn. Soc. X, 177 (1809)	...	...	...	—	—	—	—	N.A.	B.fl.V,462	
G. Hilliana; F. v. M. in Transact. phil. Inst. Vict. II, 72 (1857)	...	...	...	—	—	—	N.S.W.	Q.	B.fl.V,463	
G. pinnatifida; Bailey, papers on Q. plants I, 6 (1896)	...	...	...	—	—	—	—	Q.	—	
G. gibbosa; R. Brown in Transact. Linn. Soc. X, 177 (1809)	...	...	...	—	—	—	—	Q. N.A.	B.fl.V,463	M.fr.VI,210
G. buxifolia; R. Brown in Transact. Linn. Soc. X, 174 (1809)	...	...	...	—	—	—	N.S.W.	—	B.fl.V,464	P.
G. phyllicoides; R. Brown in Transact. Linn. Soc. X, 174 (1809)	...	...	...	—	—	—	N.S.W.	—	B.fl.V,464	M.fr.VI,213.
G. sphacelata; R. Brown in Transact. Linn. Soc. X, 174 (1809)	...	...	...	—	—	—	N.S.W.	—	B.fl.V,464	
G. occidentalis; R. Brown in Transact. Linn. Soc. X, 173 (1809)	...	...	...	W.A.	—	—	—	—	B.fl.V,465	M.fr.I,136.
G. accrosa; F. v. M., fragm. I, 136 (1859)	...	...	...	W.A.	—	—	—	—	B.fl.V,465	M.fr.I,136.
G. umbellulata; Meissner in Lehm. pl. Preiss. II, 252 (1847)	...	...	...	W.A.	—	—	—	—	B.fl.V,465	
G. oxystigma; Meissner in Lehm. pl. Preiss. I, 540 (1845)	...	...	...	W.A.	—	—	—	—	B.fl.V,466	
G. Candolleana; Meissner in Lehm. pl. Preiss. I, 541 (1845)	...	...	...	W.A.	—	—	—	—	B.fl.V,466	M.fr.VI,213.

G. scabra; Meissner in Lehm. pl. Preiss. I, 541 (1845)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,466	
G. Miqueliana; F. v. M. in Transact. Vict. Inst. 132 (1855)	...	...	...	...	—	—	—	V.	N.S.W.	—	B.fl.V,467	
G. Victoriae; F. v. M. in Transact. phil. Soc. Vict. I, 107 (1855)	...	...	...	...	—	—	—	V.	N.S.W.	—	B.fl.V,468	M.fr. VIII,142,150.
G. punicea; R. Brown in Transact. Linn. Soc. X, 169 (1809)	...	...	...	...	—	—	—	—	N.S.W.	—	B.fl.V,468	M.fr. XI,123.
G. oleoides; Sieber in Roem. & Schult. mant. 277 (1822)	...	...	...	...	—	—	—	V.	N.S.W.	—	B.fl.V,468	
G. trinervis; R. Brown, prot. nov. 18 (1830)	...	...	...	...	—	—	—	—	N.S.W.	—	B.fl.V,469	M.fr. VI,211.
G. juniperina; R. Brown in Transact. Linn. Soc. X, 171 (1809)	...	...	...	...	—	—	—	—	N.S.W.	—	B.fl.V,469	
G. sericea; R. Brown in Transact. Linn. Soc. X, 170 (1809)	...	...	...	...	—	—	—	—	N.S.W.	—	B.fl.V,470	
G. capitellata; Meissner in Hook. Kew Misc. IV, 187 (1852)	...	...	...	...	—	—	—	—	N.S.W.	—	B.fl.V,470	
G. leiophylla; F. v. M. in Benth. Fl. Austr. V, 471 (1870)	...	...	...	...	—	—	—	—	N.S.W.	Q.	B.fl.V,471	
G. linearis; R. Brown in Transact. Linn. Soc. X, 170 (1809)	...	...	...	...	—	—	—	—	N.S.W.	—	B.fl.V,471	
G. confertifolia; F. v. M. in Transact. phil. Soc. Vict. I, 22 (1854)	...	...	...	...	—	—	—	V.	—	—	B.fl.V,471	M.fr. VI,211.
G. parviflora; R. Brown in Transact. Linn. Soc. X, 171 (1809)	...	...	...	...	—	—	—	S.A.	V.	N.S.W.	B.fl.V,472	
G. halmaturina; Tate in Transact. R.S. of S.A. VI, 141 (1883)	...	...	...	...	—	—	—	S.A.	—	—	—	
G. Australis; R. Brown in Transact. Linn. Soc. X, 171 (1809)	...	...	...	...	—	—	—	T.	V.	N.S.W.	B.fl.V,472	M.fr. VII,133.
G. commutata; F. v. M., fragm. VI, 207 (1868)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,473	M.fr. VII,207.
G. pinnatisecta; F. v. M. in Benth. Fl. Austr. V, 473 (1870)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,473	
G. argyrophylla; Meissner in Hook. Kew Misc. VII, 75 (1855)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,474	M.fr. VI,411.
G. brachystachys; Meissner in Lehm. pl. Preiss. II, 254 (1847)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,474	M.fr. VI,211.
G. Endlicheriana; Meissner in Lehm. pl. Preiss. I, 546 (1845)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,474	M.fr. VI,211.
G. manglesoides; Meissner in Lehm. pl. Preiss. I, 547 (1845)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,475	
G. diversifolia; Meissner in Lehm. pl. Preiss. I, 547 (1845)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,475	
G. filifolia; Meissner in Lehm. pl. Preiss. I, 547 (1845)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,475	
G. hakeoides; Meissner in Lehm. pl. Preiss. II, 252 (1847)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,476	
G. teretifolia; Meissner in Lehm. pl. Preiss. II, 255 (1847)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,476	
G. eryngioides; Benth. Fl. Austr. V, 476 (1870)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,476	M.fr. X,46.
G. bracteosa; Meissner in Lehm. pl. Preiss. II, 254 (1847)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,477	
G. crithmifolia; R. Brown, prot. nov. 23 (1830)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,477	
G. trachytheca; F. v. M., fragm. VI, 207 (1868)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,477	M.fr. VI,207.
G. triternata; R. Brown, prot. nov. 21 (1830)	...	...	...	...	—	—	—	V.	N.S.W.	—	B.fl.V,478	
G. ramosissima; Meissner in De Candolle, prodr. XIV, 388	...	...	...	...	—	—	—	V.	N.S.W.	—	B.fl.V,478	M.fr. VIII,150; X,46.
G. monticola; Meissner in Lehm. pl. Preiss. II, 259 (1847)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,478	M.fr. VI,209
G. Muellieri; Benth. Fl. Austr. V, 479 (1870)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,479	
G. trifida; Meissner in Lehm. pl. Preiss. I, 553 (1845)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,479	M.fr. VI,209.
G. Synapheae; R. Brown, prot. nov. 23 (1830)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,480	
G. flexuosa; Meissner in Lehm. pl. Preiss. I, 553 (1845)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,480	
G. leptobotrya; Meissner in Lehm. pl. Preiss. II, 256 (1847)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,480	M.fr. VI,209.
G. brevicuspis; Meissner in Lehm. pl. Preiss. II, 256 (1847)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,481	
G. intricata; Meissner in Hook. Kew Misc. VII, 74 (1855)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,481	
G. didymobotrya; Meissner in Hook. Kew Misc. IV, 186 (1852)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,481	M.fr. I,136; VI,208; X,45.
G. polybotrya; Meissner in Hook. Kew Misc. IV, 185 (1852)	...	...	...	...	W.A.	—	—	—	—	N.A.	B.fl.V,482	M.fr. IV,129; VI,208.
G. nematophylla; F. v. M., fragm. I, 136 (1859)	...	...	...	...	—	S.A.	—	—	N.S.W.	—	B.fl.V,482	M.fr. I,136; VI,208.
G. anethifolia; R. Brown, prot. nov. 21 (1830)	...	...	...	...	—	—	—	—	N.S.W.	—	B.fl.V,482	M.fr. VI,209.
G. paradoxa; F. v. M., fragm. VI, 246 (1868)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,483	M.fr. VI,246.
G. petrophiloides; Meissner in Lehm. pl. Preiss. II, 257 (1847)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,483	M.fr. VI,209; X,46.
G. tenuiflora; Meissner in Lehm. pl. Preiss. I, 554 (1845)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,483	
G. pulchella; Meissner in Lehm. pl. Preiss. I, 553 (1845)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,484	M.fr. VI,209.
G. rudis; Meissner in Hook. Kew Misc. VII, 73 (1855)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,484	
G. apiculoba; F. v. M., fragm. X, 45 (1876)	...	...	...	...	W.A.	—	—	—	—	—	—	M.fr. X,45.
G. Shuttleworthiana; Meissner in Lehm. pl. Preiss. II, 258 (1847)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.V,484	

<i>G. integrifolia</i> ; Meissner in De Candolle, prodr. XIV, 385 (1856) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.V, 485	
<i>G. stenocarpa</i> ; F. v. M. in Benth. Fl. Austr. V, 485 (1870) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.V, 485	
<i>G. acrobotrya</i> ; Meissner in Hooker's Kew Misc. VII, 74 (1855) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.V, 486	M.fr.VI, 213.
<i>G. glabrata</i> ; Meissner in Lehm. pl. Preiss. I, 549 (1845) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.V, 486	
<i>G. ornithopoda</i> ; Meissner in Lehm. pl. Preiss. II, 256 (1847) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.V, 486	
<i>G. paniculata</i> ; Meissner in Lehm. pl. Preiss. I, 550 (1845) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.V, 487	
<i>G. biternata</i> ; Meissner in Lehm. pl. Preiss. I, 549 (1845) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.V, 487	M.fr.VI, 213.
<i>G. triloba</i> ; Meissner in Hooker's Kew Misc. VII, 74 (1855) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.V, 487	M.fr.VI, 213.
<i>G. amplexans</i> ; F. v. M. in Benth. Fl. Austr. V, 488 (1870) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.V, 488	
<i>G. vestita</i> ; Meissner in Lehm. pl. Preiss. I, 548 (1845) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.V, 488	M.fr.VI, 213.
<i>G. tridentifera</i> ; Meissner in Lehm. pl. Preiss. I, 547 (1845) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.V, 488	
<i>G. erinacea</i> ; Meissner in Hooker's Kew Misc. VII, 74 (1855) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.V, 489	M.fr.X, 45.

HAKEA; Schrader, sert. Hannov. I, fasc. 3, 27, t. 17 (1797). (Conchium.)

<i>H. chordophylla</i> ; F. v. M. in Hooker's Kew Misc. IX, 23 (1857) ...	...	...	—	—	—	—	—	Q. N.A.	B.fl.V, 495	M.fr.V, 190.
<i>H. Cunninghamii</i> ; R. Brown, prot. nov. 26 (1830) ...	...	...	W.A.	—	—	—	—	—	N.A. B.fl.V, 495	M.fr.VI, 190.
<i>H. lorea</i> ; R. Brown, prot. nov. 25 (1830) ...	...	...	W.A.	S.A.	—	—	N.S.W.	Q. N.A.	B.fl.V, 495	M.fr.VI, 189; VII, 133.
<i>H. Fraseri</i> ; R. Brown, prot. nov. 26 (1830) ...	...	...	—	—	—	—	—	—	B.fl.V, 496	
<i>H. macrocarpa</i> ; Cunningham in R. Brown, prot. nov. 30 (1830) ...	...	...	W.A.	S.A.	—	—	—	Q. N.A.	B.fl.V, 496	M.fr.VI, 190, 210.
<i>H. arborescens</i> ; R. Brown in Transact. Linn. Soc. X, 187 (1809) ...	...	...	—	—	—	—	—	Q. N.A.	B.fl.V, 497	M.fr.VI, 190.
<i>H. Ednieana</i> ; Tate in Transact. R. Soc. of S.A. VII, 70 (1884) ...	...	...	—	S.A.	—	—	—	—	—	
<i>H. Persiehana</i> ; F. v. M. in Austr. Journ. of Pharm. I, 430 (1886) ...	...	...	—	—	—	—	—	Q. —	—	
<i>H. stenophylla</i> ; Cunningham in De Candolle, prodr. XIV, 417 (1856) ...	...	...	—	—	—	—	—	—	N.A. B.fl.V, 497	
<i>H. trineura</i> ; F. v. M., fragm. III, 146 (1863) ...	...	...	—	—	—	—	—	Q. —	B.fl.V, 497	M.fr.VI, 216.
<i>H. cyclocarpa</i> ; Lindley, Bot. Regist. XXV, App. 36 (1839) ...	...	...	W.A.	—	—	—	—	—	B.fl.V, 498	
<i>H. crassifolia</i> ; Meissner in Lehm. pl. Preiss. I, 570 (1845) ...	...	...	W.A.	—	—	—	—	—	B.fl.V, 498	M.fr.VI, 216.
<i>H. Brooksiana</i> ; F. v. M. in Austr. Journ. of Pharm. I, 430 (1886) ...	...	...	W.A.	—	—	—	—	—	—	
<i>H. pandanocarpa</i> ; R. Brown, prot. nov. 29 (1830) ...	...	...	W.A.	—	—	—	—	—	B.fl.V, 499	M.fr.VI, 216.
<i>H. Roei</i> ; Benth. Fl. Austr. V, 499 (1870) ...	...	...	W.A.	—	—	—	—	—	B.fl.V, 499	
<i>H. adnata</i> ; R. Brown, prot. nov. 26 (1830) ...	...	...	W.A.	—	—	—	—	—	B.fl.V, 499	
<i>H. obliqua</i> ; R. Brown in Transact. Linn. Soc. X, 180 (1809) ...	...	...	W.A.	—	—	—	—	—	B.fl.V, 500	M.fr.VI, 218.
<i>H. Hookeriana</i> ; Meissner in Hook. Kew Misc. IV, 208 (1852) ...	...	...	W.A.	—	—	—	—	—	B.fl.V, 500	M.fr.VI, 216.
<i>H. incrassata</i> ; R. Brown, prot. nov. 29 (1839) ...	...	...	W.A.	—	—	—	—	—	B.fl.V, 500	
<i>H. flabellifolia</i> ; Meissner in Hook. Kew Misc. VII, 116 (1855) ...	...	...	W.A.	—	—	—	—	—	B.fl.V, 501	
<i>H. Brownii</i> ; Meissner in Lehm. pl. Preiss. I, 569 (1845) ...	...	...	W.A.	—	—	—	—	—	B.fl.V, 501	
<i>H. Baxteri</i> ; R. Brown, prot. nov. 28 (1830) ...	...	...	W.A.	S.A.	—	—	—	—	B.fl.V, 501	
<i>H. ceratophylla</i> ; R. Brown in Transact. Linn. Soc. X, 184 (1809) ...	...	...	W.A.	—	—	—	—	—	B.fl.V, 501	M.fr.IV, 49; VI, 217.
<i>H. lasiantha</i> ; R. Brown, prot. nov. 29 (1830) ...	...	...	W.A.	—	—	—	—	—	B.fl.V, 502	M.fr.VI, 215.
<i>H. eriantha</i> ; R. Brown, prot. nov. 29 (1830) ...	...	...	—	—	—	—	V. N.S.W.	Q. —	B.fl.V, 502	M.fr.VI, 217.
<i>H. megalosperma</i> ; Meissner in Hook. Kew Misc. VII, 117 (1855) ...	...	...	W.A.	—	—	—	—	—	B.fl.V, 503	
<i>H. clavata</i> ; Labillardière, Nov. Holl. pl. spec. I, 31, t. 41 (1804) ...	...	...	W.A.	—	—	—	—	—	B.fl.V, 503	M.fr.VI, 216.
<i>H. orthorrhyncha</i> ; F. v. M., fragm. V, 214 (1868) ...	...	...	W.A.	—	—	—	—	—	B.fl.V, 503	M.fr.VI, 214.
<i>H. Candolleana</i> ; Meissner in Lehm. pl. Preiss. II, 262 (1847) ...	...	...	W.A.	—	—	—	—	—	B.fl.V, 504	M.fr.VI, 215.
<i>H. trifurcata</i> ; R. Brown in Transact. Linn. Soc. X, 183 (1809) ...	...	...	W.A.	—	—	—	—	—	B.fl.V, 504	M.fr.VI, 220.
<i>H. erinacea</i> ; Meissner in Lehm. pl. Preiss. I, 559 (1845) ...	...	...	W.A.	—	—	—	—	—	B.fl.V, 505	M.fr.VI, 220.
<i>H. platysperma</i> ; Hooker, icon. pl. t. 433 (1842) ...	...	...	W.A.	—	—	—	—	—	B.fl.V, 505	
<i>H. brachyptera</i> ; Meissner in Hook. Kew Misc. IV, 208 (1852) ...	...	...	W.A.	—	—	—	—	—	B.fl.V, 505	M.fr.VI, 217.
<i>H. Kippistiana</i> ; Meissner in Hooker's Kew Misc. VII, 115 (1855) ...	...	...	W.A.	—	—	—	—	—	B.fl.V, 506	
<i>H. Preissii</i> ; Meissner in Lehm. pl. Preiss. I, 557 (1845) ...	...	...	W.A.	—	—	—	—	—	B.fl.V, 506	M.fr.VI, 219.
<i>H. pugioniformis</i> ; Cavanilles, Annal. hist. nat. I, 213, t. 11 (1800) ...	...	...	—	—	—	T. V.	N.S.W.	Q. —	B.fl.V, 506	M.fr.VI, 217.
<i>H. vittata</i> ; R. Brown in Transact. Linn. Soc. X, 182 (1809) ...	...	...	—	S.A.	—	V. N.S.W.	Q. —	—	B.fl.V, 507	

H. rostrata; F. v. M. in Schlecht. Linnaea XXVI, 259 (1853) ...	...	...	...	...	...	—	S.A.	—	V.	—	—	—	—	B.fl. V, 508	M.fr. VI, 218; VII, 133.
H. rugosa; R. Brown in Transact. Linn. Soc. X, 179 (1809) ...	...	...	...	...	...	—	S.A.	—	V.	—	—	—	—	B.fl. V, 508	M.fr. VI, 218.
II. Epiglottis; Labillardière, Nov. Holl. pl. spec. I, 30, t. 40 (1804) ...	...	...	...	...	...	—	—	—	T.	—	—	—	—	B.fl. V, 508	
H. amplexicaulis; R. Brown in Transact. Linn. Soc. X, 184 (1809) ...	...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 509	M.fr. VI, 217; VII, 133.
H. glabella; R. Brown, prot. nov. 28 (1830) ...	...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 509	
H. auriculata; Meissner in Hook. Kew Misc. VII, 116 (1855) ...	...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 510	M.fr. VI, 217.
H. cristata; R. Brown, prot. nov. 28 (1830) ...	...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 510	
H. linearis; R. Brown in Transact. Linn. Soc. X, 183 (1809) ...	...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 511	
H. stenocarpoides; F. v. M. in Benth. Fl. V, 511 (1870) ...	...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 511	
H. pedunculata; F. v. M. in Melb. Chem. July (1883) ...	...	...	...	...	...	...	—	—	—	—	—	Q.	—	—	
H. ruscifolia; Labillardière, Nov. Holl. pl. spec. I, 30, t. 39 (1804) ...	...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 511	
H. saligna; R. Brown in J. Knight, prot. 108 (1809) ...	...	...	...	...	...	...	—	—	—	V.	N.S.W.	Q.	—	B.fl. V, 512	M.fr. VI, 217.
H. verrucosa; F. v. M., fragm. V, 25 (1865) ..	...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 512	M.fr. V, 25; VI, 218.
H. rhombales; F. v. M., fragm. X, 90 (1876) ..	...	...	...	...	...	...	—	—	—	—	—	—	N.A.	—	M.fr. X, 90.
H. purpurea; Hooker in Mitch. Trop. Austr. 348 (1848) ...	...	...	...	...	...	...	—	S.A.	—	—	N.S.W.	Q.	—	B.fl. V, 513	
H. pubescens; Schrader. sert. Hannover. I, 25 (1795), ...	...	...	...	...	...	...	—	—	—	—	N.S.W.	—	—	B.fl. V, 513	M.fr. VI, 218.
H. propinqua; Cunningham in Field's N.S. Wales, 327 (1825) ...	...	...	...	...	...	...	—	—	—	—	N.S.W.	—	—	B.fl. V, 513	M.fr. V, 26.
H. Macraeana; F. v. M. in Austr. Journ. of Phar. I, 430 (1886) ...	...	...	...	...	...	...	—	—	—	—	N.S.W.	—	—	—	
H. nodosa; R. Brown in Transact. Linn. Soc. X, 179 (1809) ...	...	...	...	...	...	...	—	S.A.	T.	V.	—	—	—	B.fl. V, 514	M.fr. V, 26; VI, 217.
H. sericea; Schrader sert. Hannover. I, 27 (1795) ...	...	...	...	...	...	...	—	—	T.	V.	N.S.W.	—	—	B.fl. V, 514	M.fr. VI, 218.
H. leucoptera; R. Brown in Transact. Linn. Soc. X, 180 (1809) ...	...	...	...	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl. V, 515	M.fr. VI, 219; VII, 133.
H. cycloptera; R. Brown in Transact. Linn. Soc. X, 182 (1809) ...	...	...	...	...	...	...	—	S.A.	—	—	—	—	—	B.fl. V, 515	M.fr. VI, 217.
H. microcarpa; R. Brown in Transact. Linn. Soc. X, 182 (1809) ...	...	...	...	...	...	...	—	—	T.	V.	N.S.W.	—	—	B.fl. V, 516	M.fr. VI, 217.
H. recurva; Meissner in Hook. Kew Misc. IV, 207 (1852) ...	...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 516	M.fr. VI, 218.
H. circumalata; Meissner in Hook. Kew Misc. VII, 114 (1855) ...	...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 516	M.fr. VI, 219.
H. commutata; F. v. M., fragm. V, 26 (1865) ...	...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 517	M.fr. VI, 219.
H. strumosa; Meissner in Hook. Kew Misc. IV, 208 (1852) ...	...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 517	M.fr. VI, 219.
H. multilineata; Meissner in Lehm. pl. Preiss. II, 261 (1847) ...	...	...	...	...	...	...	W.A.	S.A.	—	—	—	—	N.A.	B.fl. V, 518	M.fr. V, 25; VI, 215.
H. laurina; R. Brown, prot. nov. 29 (1830) ...	...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 518	M.fr. IV, 130, 216.
H. obtusa; Meissner in Hook. Kew Misc. IV, 209 (1852) ...	...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 519	
H. cinerea; R. Brown in Transact. Linn. Soc. X, 186 (1809) ...	...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 519	
H. corymbosa; R. Brown, prot. nov. 28 (1830) ...	...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 519	M.fr. VI, 216.
H. undulata; R. Brown in Transact. Linn. Soc. X, 185 (1809) ...	...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 520	M.fr. VI, 216.
H. petiolaris; Meissner in Lehm. pl. Preiss. I, 577 (1845) ...	...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 520	
H. neurophylla; Meissner in Hook. Kew Misc. VII, 117 (1809) ...	...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 521	
H. loranthifolia; Meissner in Lehm. pl. Preiss. I, 574 (1845) ...	...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 521	
H. cucullata; R. Brown, prot. nov. 30 (1830) ...	...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 521	M.fr. VI, 216.
H. ferruginea; Sweet, Fl. Austral. t. 45 (1828) ...	...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 522	M.fr. VI, 216.
H. smilacifolia; Meissner in Lehm. pl. Preiss. I, 567 (1845) ...	...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 522	M.fr. VI, 217.
H. elliptica; R. Brown in Transact. Linn. Soc. X, 187 (1809) ...	...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 523	M.fr. VI, 216.
H. ambigua; Meissner in Lehm. pl. Preiss. II, 260 (1847) ...	...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 523	M.fr. VI, 216.
H. plurinervia; F. v. M. in Benth. Fl. Austr. V, 523 (1870) ...	...	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl. V, 523	
H. dactyloides; Cavanilles, Anal. hist. nat. I, 215, t. 12 (1803) ...	...	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl. V, 524	M.fr. VI, 215.
H. ulicina; R. Brown, prot. nov. 29 (1830) ...	...	...	...	...	...	...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl. V, 524	
H. falcata; R. Brown, prot. nov. 29 (1830) ...	...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 524	
H. pycnoneura; Meissner in Hook. Kew Misc. VII, 117 (1855) ...	...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 525	M.fr. VI, 215.
II. stenocarpa; R. Brown, prot. nov. 29 (1830) ...	...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 525	
H. marginata; R. Brown in Transact. Linn. Soc. X, 185 (1809) ...	...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 526	M.fr. VI, 216.
H. myrtoides; Meissner in Lehm. pl. Preiss. I, 577 (1845) ...	...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 526	
H. costata; Meissner in Lehm. pl. Preiss. I, 575 (1845) ...	...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 526	

<i>B. nutans</i> ; R. Brown in Transact. Linn. Soc. X, 203 (1809) ...	...	...	W.A.	—	—	—	—	—	B.fl.V,545	M.fr.IV,108; VII,54.
<i>B. sphaerocarpa</i> ; R. Brown in Transact. Linn. Soc. X, 203 (1809) ...	...	...	W.A.	—	—	—	—	—	B.fl.V,546	M.fr.VII,54.
<i>B. tricuspis</i> ; Meissner in Hook. Kew Misc. VII, 118 (1855) ...	...	...	W.A.	—	—	—	—	—	B.fl.V,546	M.fr.VII,54.
<i>B. occidentalis</i> ; R. Brown in Transact. Linn. Soc. X, 204 (1809) ...	...	...	W.A.	—	—	—	—	—	B.fl.V,546	M.fr.VII,54.
<i>B. litoralis</i> ; R. Brown in Transact. Linn. Soc. X, 204 (1809) ...	...	...	W.A.	—	—	—	—	—	B.fl.V,547	M.fr.VII,54.
<i>B. ericifolia</i> ; Linné fil., suppl. 127 (1781) ...	...	...	—	—	—	—	N.S.W.	—	B.fl.V,547	M.fr.VII,54.
<i>B. spinulosa</i> ; Smith, Specim. Bot. New Holl. 13, t. 4 (1793) ...	...	...	—	—	—	—	N.S.W.	—	B.fl.V,547	
<i>B. collina</i> ; R. Brown in Transact. Linn. Soc. X, 204 (1809) ...	...	...	—	—	—	V.	N.S.W.	Q.	B.fl.V,548	M.fr.VII,54.
<i>B. verticillata</i> ; R. Brown in Transact. Linn. Soc. X, 207 (1809) ...	...	...	W.A.	—	—	—	—	—	B.fl.V,548	
<i>B. dryandroides</i> ; Baxter in Sweet, hort. Brit., 350 (1827) ...	...	...	W.A.	—	—	—	—	—	B.fl.V,549	M.fr.VII,58.
<i>B. Brownii</i> ; Baxter in R. Brown prot. nov. 37 (1830) ...	...	...	W.A.	—	—	—	—	—	B.fl.V,549	M.fr.VII,58.
<i>B. attenuata</i> ; R. Brown in Transact. Linn. Soc. X, 209 (1809) ...	...	...	W.A.	—	—	—	—	—	B.fl.V,549	M.fr.VII,55.
<i>B. media</i> ; R. Brown, prot. nov. 35 (1830) ...	...	...	W.A.	—	—	—	—	—	B.fl.V,550	M.fr.IV,109.
<i>B. Solandri</i> ; R. Brown, prot. nov. 35 (1830) ...	...	...	W.A.	—	—	—	—	—	B.fl.V,550	M.fr.VII,58.
<i>B. Goodii</i> ; R. Brown, prot. nov. 36 (1830) ...	...	...	W.A.	—	—	—	—	—	B.fl.V,550	M.fr.VII,58.
<i>B. petiolaris</i> ; F. v. M., fragm. IV, 109 (1864) ...	...	...	W.A.	—	—	—	—	—	B.fl.V,551	M.fr.IV,109.
<i>B. repens</i> ; Labillardière, Voy. I, 411, t. 23 (1798) ...	...	...	W.A.	—	—	—	—	—	B.fl.V,551	M.fr.IV,108,177; VII,58.
<i>B. prostrata</i> ; R. Brown, prot. nov. 36 (1830) ...	...	...	W.A.	—	—	—	—	—	B.fl.V,551	M.fr.VII,57.
<i>B. grandis</i> ; Willdenow, spec. plant. I, 535 (1797) ...	...	...	W.A.	—	—	—	—	—	B.fl.V,552	M.fr.VII,57.
<i>B. quercifolia</i> ; R. Brown in Transact. Linn. Soc. X, 210 (1809) ...	...	...	W.A.	—	—	—	—	—	B.fl.V,552	M.fr.VII,57.
<i>B. Baueri</i> ; R. Brown, prot. nov. 35 (1830) ...	...	...	W.A.	—	—	—	—	—	B.fl.V,552	M.fr.IV,107; VII,56.
<i>B. marginata</i> ; Cavanilles, Anal. hist. nat. I, 227, t. 13 (1800) ...	...	...	—	S.A.	T.	V.	N.S.W.	—	B.fl.V,553	M.fr.VII,55,133; X,90.
<i>B. integrifolia</i> ; Linné fil., suppl. 127 (1781) ...	...	...	—	—	—	V.	N.S.W.	Q.	B.fl.V,554	M.fr.VII,55.
<i>B. dentata</i> ; Linné fil., suppl. 127 (1781) ...	...	...	—	—	—	—	Q.	N.A.	B.fl.V,555	M.fr.VII,57.
<i>B. Robur</i> ; Cavanilles in Ansl. hist. nat. I, 226 (1800) ...	...	...	—	—	—	—	N.S.W.	Q.	B.fl.V,555	M.fr.VII,56.
<i>B. serrata</i> ; Linné fil., suppl. 126 (1781) ...	...	...	—	—	—	T.	V.	N.S.W.	B.fl.V,556	M.fr.VII,56.
<i>B. serratifolia</i> ; Salisbury, prodr. 51 (1796) ...	...	...	—	—	—	—	N.S.W.	Q.	B.fl.V,556	M.fr.VII,56.
<i>B. ornata</i> ; F. v. M. in Schlecht. Linnaea XXVI, 352 (1853) ...	...	...	—	S.A.	—	V.	—	—	B.fl.V,557	M.fr.VII,56.
<i>B. coccinea</i> ; R. Brown in Transact. Linn. Soc. X, 207 (1809) ...	...	...	W.A.	—	—	—	—	—	B.fl.V,557	M.fr.VII,56.
<i>B. Sceptrum</i> ; Meissner in Hook. Kew Misc. VII, 120 (1855) ...	...	...	W.A.	—	—	—	—	—	B.fl.V,557	M.fr.VII,56.
<i>B. Menziesii</i> ; R. Brown, prot. nov. 36 (1830) ...	...	...	W.A.	—	—	—	—	—	B.fl.V,558	
<i>B. laevigata</i> ; Meissner in Hook. Kew Misc. IV, 210 (1852) ...	...	...	W.A.	—	—	—	—	—	B.fl.V,558	M.fr.VII,56.
<i>B. Hookeriana</i> ; Meissner in Hook. Kew Misc. VII, 119 (1855) ...	...	...	W.A.	—	—	—	—	—	B.fl.V,558	
<i>B. prionotes</i> ; Lindley, Bot. Regist. XXV, App. 34 (1839) ...	...	...	W.A.	—	—	—	—	—	B.fl.V,558	M.fr.VII,56.
<i>B. Victoriae</i> ; Meissner in Hook. Kew Misc. VII, 119 (1855) ...	...	...	W.A.	—	—	—	—	—	B.fl.V,559	M.fr.VII,58.
<i>B. speciosa</i> ; R. Brown in Transact. Linn. Soc. X, 210 (1809) ...	...	...	W.A.	—	—	—	—	—	B.fl.V,559	M.fr.VII,58.
<i>B. Baxteri</i> ; R. Brown, prot. nov. 36 (1830) ...	...	...	W.A.	—	—	—	—	—	B.fl.V,559	M.fr.VII,58.
<i>B. praemorsa</i> ; Andrews, Bot. repos. t. 258 (1802) ...	...	...	W.A.	—	—	—	—	—	B.fl.V,560	M.fr.VII,56.
<i>B. Lehmanniana</i> ; Meissner in Hook. Kew Misc. 210 (1852) ...	...	...	W.A.	—	—	—	—	—	B.fl.V,560	M.fr.VII,57.
<i>B. Caleyii</i> ; R. Brown, prot. nov. 35 (1830) ...	...	...	W.A.	—	—	—	—	—	B.fl.V,560	M.fr.VII,57.
<i>B. Lindleyana</i> ; Meissner in Hook. Kew Misc. VII, 120 (1855) ...	...	...	W.A.	—	—	—	—	—	B.fl.V,561	M.fr.VII,55.
<i>B. elegans</i> ; Meissner in Hook. Kew Misc. VII, 119 (1855) ...	...	...	W.A.	—	—	—	—	—	B.fl.V,561	
<i>B. Candolleana</i> ; Meissner in Hook. Kew Misc. VII, 118 (1855) ...	...	...	W.A.	—	—	—	—	—	B.fl.V,561	M.fr.VII,58.
<i>B. ilicifolia</i> ; R. Brown in Transact. Linn. Soc. X, 211 (1809) ...	...	...	W.A.	—	—	—	—	—	B.fl.V,561	M.fr.VII,58.

DRYANDRA; R. Brown in Transact. Linn. Soc. X, 211 (1809). (Josephia, Hemiclidia.)

<i>D. quercifolia</i> ; Meissner in Hook. Kew Misc. IV, 210 (1852) ...	...	...	W.A.	—	—	—	—	—	B.fl.V,566	M.fr.VII,50.
<i>D. praemorsa</i> ; Meissner in Lehm. pl. Preiss. II, 265 (1847) ...	...	...	W.A.	—	—	—	—	—	B.fl.V,566	M.fr.VII,50.
<i>D. cuneata</i> ; R. Brown in Transact. Linn. Soc. X, 212 (1809) ...	...	...	W.A.	—	—	—	—	—	B.fl.V,566	M.fr.VII,50.
<i>D. falcata</i> ; R. Brown in Transact. Linn. Soc. X, 213 (1809) ...	...	...	W.A.	—	—	—	—	—	B.fl.V,567	
<i>D. armata</i> ; R. Brown in Transact. Linn. Soc. X, 212 (1809) ...	...	...	W.A.	—	—	—	—	—	B.fl.V,567	M.fr.VII,50.

D. longifolia ; R. Brown in Transact. Linn. Soc. X, 215 (1809) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,568	
D. Fraseri ; R. Brown, prot. nov. 39 (1830) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,568	M.fr.VII,52.
D. floribunda ; R. Brown in Transact. Linn. Soc. X, 212 (1809) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,569	M.fr.VI,92; VII,50.
D. carduacea ; Lindley, Bot. Regist. XXV, App. 33 (1839)...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,569	M.fr.VII,51.
D. carlinoides ; Meissner in Lehm. pl. Preiss. II, 267 (1847) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,569	M.fr.VII,53.
D. polycephala ; Bentham, Fl. Austr. V, 570 (1870)...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,570	
D. Kippistiana ; Meissner in Hook. Kew Misc. VII, 122 (1855) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,570	M.fr.VII,52.
D. squarrosa ; R. Brown, prot. nov. 38 (1830) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,571	M.fr.VII,52.
D. Serra ; R. Brown, prot. nov. 38 (1830) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,571	M.fr.VII,51.
D. concinna ; R. Brown, prot. nov. 38 (1830) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,571	M.fr.VII,52.
D. foliolata ; R. Brown, prot. nov. 38 (1830) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,572	M.fr.VII,51.
D. stupposa ; Lindley, Bot. Regist. XXV, App. 33 (1839) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,572	M.fr.VII,51.
D. nobilis ; Lindley, Bot. Regist. XXV, App. 33 (1839) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,572	
D. mucronulata ; R. Brown in Transact. Linn. Soc. X, 213 (1809)...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,573	
D. formosa ; R. Brown in Transact. Linn. Soc. X, 213, t. 3 (1809)...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,573	M.fr.VII,51.
D. Baxteri ; R. Brown, prot. nov. 38 (1830) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,573	M.fr.VII,51.
D. nivea ; R. Brown in Transact. Linn. Soc. X, 214 (1809)...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,574	M.fr.VII,52.
D. Arctotidis ; R. Brown, prot. nov. 39 (1830) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,574	
D. nana ; Meissner in Hook. Kew Misc. VII, 121 (1855) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,575	M.fr.VII,52.
D. Preissii ; Meissner in Lehm. pl. Preiss. I, 599 (1845) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,575	M.fr.VII,54.
D. sclerophylla ; Meissner in Hook. Kew Misc. VII, 123 (1855) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,576	M.fr.VII,52.
D. pulchella ; Meissner in Hook. Kew Misc. IV, 211 (1852) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,576	
D. plumosa ; R. Brown in Transact. Linn. Soc. X, 214 (1809) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,576	M.fr.VII,51.
D. senecionifolia ; R. Brown, prot. nov. 39 (1830) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,577	M.fr.VII,53.
D. vestita ; Kippist in Hook. Kew Misc. VII, 121 (1855) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,577	M.fr.VII,53.
D. cirsioides ; Meissner in Hook. Kew Misc. IV, 211 (1852) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,577	M.fr.VII,52.
D. Hewardiana ; Meissner in Hook. Kew Misc. IV, 210 (1852) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,577	M.fr.VII,53.
D. patens ; Bentham, Fl. Austr. V, 578 (1870) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,578	
D. conferta ; Bentham, Fl. Austr. V, 578 (1870) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,578	
D. horrida ; Meissner in Hook. Kew Misc. IV, 211 (1852) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,579	M.fr.VII,52.
D. serratuloides ; Meissner in Hook. Kew Misc. VII, 123 (1855) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,579	M.fr.VII,52.
D. comosa ; Meissner in Hook. Kew Misc. IV, 211 (1852) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,579	M.fr.VII,53.
D. Shuttleworthiana ; Meissner in Hook. Kew Misc. VII, 122 (1855) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,580	M.fr.VII,52.
D. speciosa ; Meissner in Hook. Kew Misc. IV, 211 (1852)...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,580	
D. tridentata ; Meissner in Hook. Kew Misc. VII, 120 (1855) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,580	M.fr.VII,53.
D. tenuifolia ; R. Brown in Transact. Linn. Soc. X, 215 (1809) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,581	M.fr.VII,53.
D. proteoides ; Lindley, Bot. Regist. XXV, App. 33 (1839) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,582	
D. runcinata ; Meissner in Hook. Kew Misc. IV, 210 (1852) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,582	
D. obtusa ; R. Brown in Transact. Linn. Soc. X, 214 (1809) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,582	M.fr.V,185.
D. bipinnatifida ; R. Brown, prot. nov. 39 (1830) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,583	
D. pteridifolia ; R. Brown in Transact. Linn. Soc. X, 215 (1809) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,583	M.fr.VII,54.
D. calophylla ; R. Brown, prot. nov. 40 (1830) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,583	M.fr.VII,54.

CORNACEAE.

HUMBOLDT, Bonpland & Kunth, nov. gen. Americ. III, 430 (1818).—Genus 1, Species 1.

STYLIDIUM ; Loureiro, Fl. Cochinch. I, 220 (1790). (Marlea, Rhytidandra, Pseudalangium.)

S. Vitiense ; F. v. M., Cens. 74 (1832) from Asa Gray (1854) N.S.W. Q. — B.fl.III,386 M.fr.II,84,176 P.

RUBIACEAE.

A. L. DE JUSSIEU, gen. plant. 196 (1789), from B. de Jussieu (1759).—Genera 30, Species 127.

SARCOCEPHALUS; Afzelius in Transact. hort. soc. Lond. V, 422, t. 18 (1824).

S. cordatus; Miquel, Fl. Ind. Batav. II, 133 (1856)... Q. N.A. B.fl.III,402 ... As., P.

OLDENLANDIA; Plumier, nov. plant. Americ. gen. 42, t. 36 (1703). (Hedyotis, Synaptantha, Anotis.)

O. eorymbosa; Linné, sp. pl. 119 (1753) ... M.fr.VII,45 ... As., Afr.
 O. paniculata; Linné, spec. plant. edit. sec. 1667 (1763) ... M.fr.VII,45; VIII,47 ... As., P.
 O. psychotrioides; F. v. M. in Viet. Naturalist VI, 54 (1889) ... As., P.
 O. auricularia; F. v. M., Cens. of Austr. pl. 74 (1882) from Linné (1753) ... B.fl.III,404
 O. polyclada; F. v. M., Cens. of Austr. pl. 74 (1882) ... M.fr.VIII,146.
 O. coerulescens; F. v. M., Cens. of Austr. pl. 74 (1882) ... B.fl.III,404 M.fr.IV,38.
 O. spermacocoides; F. v. M., Cens. of Austr. pl. 74 (1882) ... N.A. — M.fr.VIII,146.
 O. Cronchiana; F. v. M., Cens. of Austr. pl. 74 (1882) ... N.A. — M.fr.X,85.
 O. mitrasaemoides; F. v. M., Cens. of Austr. pl. 74 (1882)... N.A. B.fl.III,404 M.fr.IV,37.
 O. galioides; F. v. M., Cens. of Austr. pl. 74 (1882) ... W.A. — Q. N.A. B.fl.III,404 M.fr.IV,38.
 O. scleranthoides; Benth. & J. Hooker, gen. pl. II, 88 (1873) ... N.A. B.fl.III,405 M.fr.IV,39.
 O. elatinoides; F. v. M., Cens. of Austr. pl. 74 (1882) from Benth. (1866) ... W.A. — B.fl.III,405 M.fr.VIII,147.
 O. tillaeacea; F. v. M., Cens. of Austr. pl. 74 (1882) ... W.A. S.A. — N.S.W. Q. N.A. B.fl.III,405 M.fr.IV,39.
 O. trachymenioides; F. v. M., Cens. of Austr. pl. 74 (1882) ... Q. — B.fl.III,405 M.fr.IV,40.
 O. pterospora; F. v. M., Cens. of Austr. pl. 74 (1882) ... N.A. B.fl.III,406 M.fr.IV,40,177.

DENTELLA; R. et G. Forster, char. gen. 25, t. 13 (1776). (Lippaya.)

D. repens; R. & G. Forster, char. gen. 25, t. 13 (1776) ... S.A. — N.S.W. Q. N.A. B.fl.III,406 M.fr.IX,187 ... As., P.

OPHIORRHIZA; Linné, Fl. Zeylan. 190 et 239 (1747).

O. Australiana; Benth. Fl. Austr. III, 407 (1866) ... B.fl.III,407 M.fr.IX,187.

ABBOTTIA; F. v. M., fragm. IX, 181 (1875).

A. siogularis; F. v. M., fragm. IX, 181 (1875) ... M.fr.IX,181.

GARDENIA; Ellis in Phil. Transact. LI, 935, t. 23 (1761).

G. edulis; F. v. M., fragm. I, 54 (1858) ... Q. N.A. B.fl.III,408 M.fr.I,54.
 G. resinosa; F. v. M., fragm. I, 54 (1858) ... N.A. B.fl.III,408 M.fr.I,54.
 G. pyriformis; Cunningham in Benth. Fl. Austr. III, 409 (1866) ... N.A. B.fl.III,409
 G. megasperma; F. v. M., fragm. I, 54 (1858) ... N.A. B.fl.III,409 M.fr.I,54.
 G. Macgillivraei; Benth. Fl. Austr. III, 409 (1866) ... Q. — B.fl.III,409 M.fr.VII,46.
 G. ochreatea; F. v. M., fragm. I, 55 (1858) ... Q. — B.fl.III,409 M.fr.155;VII,46.
 G. anfruticosa; R. Brown in Benth. Fl. Austr. III, 410 (1866) ... Q. N.A. B.fl.III,410
 G. fucata; R. Brown in Benth. Fl. Austr. III, 410 (1866) ... Q. N.A. B.fl.III,410
 G. Jardinei; F. v. M. in Benth. Fl. Austr. III, 410 (1866) ... Q. — B.fl.III,410 M.fr.VII,46.

RANDIA; Houston in Linné, hort. Cliffort. 485 (1737).

R. hirta; F. v. M., fragm. VII, 46 (1869) ... M.fr.IX,181.
 R. echartacea; F. v. M., fragm. IX, 180 (1875) ... N.S.W. Q. — B.fl.III,410 M.fr.VII,47.
 R. Benthamiana; F. v. M., fragm. IX, 180 (1875) ... N.S.W. Q. — M.fr.IX,180.
 R. Moorei; F. v. M. in Benth. Fl. Austr. III, 411 (1866) ... N.S.W. Q. — B.fl.III,411
 R. stipularis; F. v. M., Papuan plants I, 69 (1876) ... N.S.W. — M.fr.VII,47;IX,70,181.
 R. Fitzalani; F. v. M., Rep. Burdek. Exped. 12 (1860) ... Q. — B.fl.III,411 M.fr.V,19;IX,180.
 R. sessilis; F. v. M., fragm. VII, 47 (1869) ... Q. N.A. — M.fr.VII,47.
 R. densiflora; Benth. Fl. Hongk. 155 (1861) ... N.S.W. Q. N.A. B.fl.III,412 M.fr.II,132; III,166; IX,180 As.

DIPLOSPORA; De Candolle, prodr. IV, 477 (1830). (Discospermum.)

D. Australis; Benth, Fl. Austr. III, 413 (1866)	— — — — —	Q. —	B.fl.III,413						
D. ixoroides; F. v. M., fragm. IX, 182 (1875)	— — — — —	Q. —	B.fl.III,416	M.fr.IX,182.					

IXORA; Linné, syst. nat. 8 (1735), gen. pl. 27 (1737). (Pavetta, Tarenna, Webera, Stylocoryne.)

I. Pavetta; Roxburgh, Fl. Ind. I, 395, ed. Carey (1820)	— — — — —	Q. N.A.	B.fl.III,414	M.fr.IX,182	...	...	As., P.
I. tomentosa; Roxburgh, Fl. Ind. I, 396, ed. Carey (1820)... ..	— — — — —	— N.A.	B.fl.III,414		...	...	As.
I. coccinea; Linné, sp. pl. 110 (1753)	— — — — —	— N.A.	B.fl.III,415		...	...	As.
I. Timorensis; Decaisne in Nouv. Ann. du Mus. III, 419 (1834)	— — — — —	Q. N.A.	B.fl.III,415	M.fr.V,18; IX,183		...	As.
I. Becklerii; Benth, Fl. Austr. III, 415 (1866)	— — — — — N.S.W.	Q. —	B.fl.III,415	M.fr.IX,183.			
I. pentamera; Benth, Fl. Austr. III, 416 (1866)	— — — — —	— N.A.	B.fl.III,416				
I. Dallachyana; F. v. M., Cens. of Austr. pl. 75 (1882) from Benth (1866)	— — — — —	Q. N.A.	B.fl.III,412				
I. expandens; F. v. M., Cens. of Austr. pl. 75 (1882)	— — — — —	— N.A.	—	M.fr.VI,25.			

COFFEA; Linné, syst. nat. 8 (1735); gen. pl. 55 (1737) from Ray (1691).

C. Bengalensis; Roxburgh, hort. Bengal. 15 (1814)	— — — — —	Q. —	—	M.fr.VIII,147	...	...	As.
--	-----------	------	---	---------------	-----	-----	-----

TIMONIUS; Rumphius, herb. Amboin. III, 216, t. 140 (1743). (Polyphragmon.)

T. Rumphii; De Candolle, prodr. IV, 461 (1830)	— — — — —	Q. N.A.	B.fl.III,417	M.fr.II,134; IX,187		...	As., P.
---	-----------	---------	--------------	---------------------	--	-----	---------

SCYPHIPHORA; K. F. Gaertner, de fruct. III, 91, t. 196 (1805). (Epithinia.)

S. hydrophyllacea; K. F. Gaertner, de fruct. III, 91, t. 196 (1805)	— — — — —	Q. N.A.	B.fl.III,418	M.fr.VII,46; IX,187		...	As., P.
--	-----------	---------	--------------	---------------------	--	-----	---------

GUETTARDA; Osbeck in Linné, spec. plant. 991 (1753). (Antirrhoea, Guettardella, Bohea, Haplocaryon.)

G. speciosa; Linné, spec. plant. 991 (1753)	— — — — —	Q. N.A.	B.fl.III,419	M.fr.IX,183	...	...	As., Afr., P.
G. tenuiflora; F. v. M., fragm. VII, 48 (1869)	— — — — —	Q. —	B.fl.III,418	M.fr.IX,183.			
G. putaminosa; F. v. M., fragm. IX, 183 (1875)	— — — — —	Q. —	B.fl.III,419	M.fr.IV,92; V,212.			
G. myrtoides; F. v. M., fragm. IX, 184 (1875)	— — — — —	Q. —	—	M.fr.IX,184.			

HODGKINSONIA; F. v. M., fragm. II, 132 (1861).

H. ovatiflora; F. v. M., fragm. II, 132 (1861)	— — — — — N.S.W.	Q. —	B.fl.III,420	M.fr.III,166.			
---	------------------	------	--------------	---------------	--	--	--

CANTHIUM; Lamarek, Encycl. méthod. I, 602 (1783). (Psydrax, Electronia partly.)

C. cymosum; Persoon, synops. plant. I, 200 (1805)	— — — — —	Q. —	—	M.fr.IX,185	...	...	As., P.
C. latifolium; F. v. M. in Benth. Fl. Austr. III, 421 (1866)	W.A. S.A. — — N.S.W.	Q. —	B.fl.III,421	M.fr.IX,185			
C. attenuatum; R. Brown in Benth. Fl. Austr. III, 421 (1866)	— — — — —	Q. N.A.	B.fl.III,421				
C. lucidum; Hooker & Arnott, Bot. of Beechey's voy. 65 (1841)	— — — — — N.S.W.	Q. N.A.	B.fl.III,421	M.fr.II,133; IX,185; XI,135		P.	
C. oleifolium; Hooker in Mitch. Trop. Austr. 397 (1848)	— — — — — N.S.W.	Q. —	B.fl.III,422	M.fr.II,134; IX,185.			
C. buxifolium; Benth, Fl. Austr. III, 422 (1866)... ..	— — — — — N.S.W.	Q. —	B.fl.III,422				
C. vacciniifolium; F. v. M. in Transact. phil. Inst. Vict. III, 47 (1858)	— — — — — N.S.W.	Q. —	B.fl.III,422	M.fr.II,134; IX,186.			
C. coprosmoides; F. v. M. in Trausact. phil. Inst. Vict. III, 47 (1858)	— — — — — N.S.W.	Q. N.A.	B.fl.III,422				

MORINDA; Vaillant in act. Acad. Par. 202 (1722).

M. citrifolia; Linné, spec. plant. 176 (1753)	— — — — —	Q. N.A.	B.fl.III,422		...	...	As., Afr., P.
M. acutifolia; F. v. M., fragm. IX, 179 (1875)	— — — — —	Q. —	B.fl.III,429	M.fr.IX,179.			
M. jasminoides; Cunningham in Hook. Bot. Mag. t. 3351 (1834)	— — — — — V. N.S.W.	Q. —	B.fl.III,423				
M. umbellata; Linné, spec. plant. 176 (1753)	— — — — —	Q. —	B.fl.III,423		...	...	As., P.
M. reticulata; Benth, Fl. Austr. III, 424 (1866)	— — — — —	Q. N.A.	B.fl.III,423	M.fr.IX,180.			
M. hypotephra; F. v. M. in Vict. Nat. III, 170 (1887)	— — — — —	Q. —	—				

COELOSPERMUM; Blume, Bijdr. 994 (1826). (Pogonolobus.)

C. paniculatum; F. v. M., fragm. V, 19 (1865)	— — — — — N.S.W.	Q. —	B.fl.III,425	M.fr.VII,46; IX,185.			
C. reticulatum; Benth, Fl. Austr. III, 425 (1866)	— — — — —	Q. N.A.	B.fl.III,425	M.fr.I,56; IX,185.			

HYDNOPHYTUM; Jack in Transact. Linn. Soc. XIV, 124 (1823).

H. formicarum; Jack in Transact. Linn. Soc. XIV, 124 (1823) — — — — — Q. N.A. — M.fr.VII,45 As., P.

MYRMECODIA; Jack in Transact. Linn. Soc. XIV, 122 (1823).

M. Antoinii; Beccari, Malesia II, 116, t. 14 (1884) — — — — — Q. — — M.fr.VII,45.

LASIANTHUS; Jack in Transact. Linn. Soc. XIV, 125 (1823).

L. strigosus; Wight in Calc. Journ. nat. hist. VI, 512 (1846) — — — — — Q. — B.fl.III,426 As.

PSYCHOTRIA; Linné, syst. ed. decim. 929 (1759). (Uragoga, 1737, Grumilia.)

P. nesophila; F. v. M., fragm. II, 135 (1861)... .. — — — — — Q. N.A. B.fl.III,427 M.fr.IX,184.
 P. Dallachyana; Benth., Fl. Austr. III, 427 (1866) — — — — — Q. — B.fl.III,427
 P. nematopoda; F. v. M., fragm. VII, 48 (1869) — — — — — Q. — — M.fr.IX,184.
 P. loniceroides; Sieber in De Candolle, prodr. IV, 523 (1830) — — — — — N.S.W. Q. — B.fl.III,427 M.fr.IX,184.
 P. daphnoides; Cunningham in Hook. Bot. Mag. 3228 (1833) — — — — — N.S.W. Q. — B.fl.III,428 M.fr.II,135; IX,184.
 P. polioctemma; Benth., Fl. Austr. III, 427 (1866) — — — — — Q. — B.fl.III,428
 P. Fitzalani; Benth., Fl. Austr. III, 427 (1866) — — — — — Q. — B.fl.III,428
 P. Carronis; Moore & F. v. M., fragm. VII, 49 (1869) — — — — — N.S.W. — — — M.fr.IX,70,185.

GEOPHILA; D. Don, prodr. fl. Nepal. 136 (1825). (Cephaelis partly.)

G. reniformis; D. Don, prodr. fl. Nepal. 136 (1825)... .. — — — — — Q. — — M.fr.VII,45; VIII,147; IX,187. [Am., P. As., Afr.,

COPROSMA; R. et G. Forster, char. gen. 137, t. 69 (1776). (Nertera, Marquisia.)

C. Nertera; F. v. M., fragm. IX, 186 (1875) — — — — — T. V. N.S.W. — — B.fl.III,431 M.fr.IX,186 Am., P., N.Z.
 C. reptans; F. v. M., fragm. IX, 186 (1875) — — — — — V. N.S.W. — — B.fl.III,431 M.fr.VIII,147.
 C. pumila; J. Hooker in Lond. Journ. VI, 465 (1847) — — — — — T. V. — — — B.fl.III,430 M.fr.IX,186 N.Z.
 C. nitida; J. Hooker in Lond. Journ. VI, 465 (1847) — — — — — T. V. — — — B.fl.III,430 M.fr.IX,186.
 C. Billardieri; J. Hooker in Lond. Journ. VI, 465 (1847) — — — — — T. V. N.S.W. — — B.fl.III,430
 C. hirtella; Labillardière, Nov. Holl. pl. spec. I, 70, t. 95 (1804)... .. — — — — — S.A. T. V. N.S.W. — — B.fl.III,429 M.fr.IX,186.
 C. Baueri; Endlicher, iconogr. gen. pl. XI, t. 111 (1838) — — — — — N.S.W. — — — M.fr.IX,69.
 C. putida; C. Moore & F. v. M., fragm. VII, 45 (1869) — — — — — N.S.W. — — — M.fr.IX,69.
 C. lanceolaris; F. v. M., fragm. IX, 70 (1875) — — — — — N.S.W. — — — M.fr.IX,70.
 C. pilosa; Endlicher, florul. ins. Norf. prodr. 60 (1833) — — — — — N.S.W. — — —

OPERCULARIA; J. Gaertner, de fruct. I, 111, t. 24 (1788).

O. vaginata; Labillardière, Nov. Holl. pl. spec. I, 34, t. 46 (1804) W.A. — — — — — B.fl.III,433
 O. spermacocca; Labillardière, Nov. Holl. pl. spec. I, 35, t. 47 (1804) W.A. — — — — — B.fl.III,433
 O. scabrida; Schlechtendal, Linnaea XX, 604 (1847) — — — — — S.A. — — — V. N.S.W. — — B.fl.III,433
 O. hirsuta; F. v. M. in Benth. Fl. Austr. III, 434 (1866) W.A. — — — — — — — B.fl.III,434
 O. aspera; Gaertner, de fruct. I, 112, t. 24 (1788) — — — — — V. N.S.W. Q. — B.fl.III,434
 O. hispida; Sprengel, syst. veg. I, 385 (1825)... .. — — — — — V. N.S.W. — — B.fl.III,434
 O. diphylla; Gaertner, de fruct. I, 113 (1788) — — — — — N.S.W. Q. — B.fl.III,434
 O. ovata; J. Hooker in Lond. Journ. VI, 465 (1847) — — — — — S.A. T. V. N.S.W. — — B.fl.III,435
 O. varia; J. Hooker in Lond. Journ. VI, 466 (1847) — — — — — S.A. T. V. N.S.W. Q. — B.fl.III,435
 O. rubioides; A. L. de Jussieu in Annal. du Mus. IV, 428 (1804)... .. W.A. — — — — — — — B.fl.III,435
 O. volubilis; R. Brown in Benth. Fl. Austr. III, 435 (1866) W.A. — — — — — — — B.fl.III,435
 O. hispidula; Endlicher in Hueg. enum. 58 (1837) W.A. — — — — — — — B.fl.III,436
 O. echinocephala; Benth., Fl. Austr. III, 436 (1866) W.A. — — — — — — — B.fl.III,436
 O. apiciflora; Labillardière, Nov. Holl. pl. spec. I, 35, t. 48 (1804) W.A. — — — — — — — B.fl.III,436

POMAX; Solander in Gaertner, de fruct. I, 112 (1788).

P. umbellata; Solander in Gaertner, de fruct. I, 112 (1788) W.A. S.A. — — — — — V. N.S.W. Q. — B.fl.III,437 M.fr.IX,187.

ELEUTHRANTHES; F. v. M., fragm. IV, 92 (1864).

E. opercularina; F. v. M., fragm. IV, 92 (1864) W.A. — — — — — — — B.fl.III,437 M.fr.IV,92.

KNOXIA; Linné, Fl. Zeyl. 189 (1747).

K. corymbosa; Willdenow, spec. pl. I, 582 (1798) — — — — — Q. N.A. B.f.III,438 M.fr.IV,41; IX,187 ... As., P.

SPERMACOCE; Dillenius, hort. Elth. 369, t. 277 (1732).

S. brachystoma; R. Brown in Benth. Fl. Austr. III, 439 (1866) — — — — — Q. N.A. B.f.III,439 M.fr.IV,41.
 S. pogostoma; Bentham, Fl. Austr. III, 440 (1866) — — — — — — N.A. B.f.III,440
 S. leptoloba; Bentham, Fl. Austr. III, 440 (1866) — — — — — — N.A. B.f.III,440
 S. marginata; Bentham, Fl. Austr. III, 440 (1866) — S.A. — — — — — Q. N.A. B.f.III,440
 S. multicaulis; Bentham, Fl. Austr. III, 440 (1866) — — — — — N.S.W. Q. N.A. B.f.III,440
 S. exserta; Bentham, Fl. Austr. III, 441 (1866) — — — — — Q. N.A. B.f.III,441
 S. membranacea; R. Brown in Benth. Fl. Austr. III, 441 (1866) — — — — — Q. N.A. B.f.III,441
 S. debilis; Bentham, Fl. Austr. III, 441 (1866) — — — — — Q. — B.f.III,441
 S. inaperta; F. v. M., fragm. IV, 43 (1863) — — — — — — N.A. B.f.III,441 M.fr.IV,43.
 S. stenophylla; F. v. M., fragm. IV, 43 (1863) — — — — — Q. N.A. B.f.III,442 M.fr.IV,43.
 S. laevigata; F. v. M., fragm. IV, 41 (1863) — — — — — Q. N.A. B.f.III,442 M.fr.IV,41.
 S. breviflora; F. v. M. in Benth. Fl. Austr. III, 442 (1866) — — — — — Q. N.A. B.f.III,442
 S. auriculata; F. v. M., fragm. IV, 42 (1863) — — — — — — N.A. B.f.III,442 M.fr.IV,42.
 S. suffruticosa; R. Brown in Benth. Fl. Austr. III, 443 (1866) — — — — — Q. N.A. B.f.III,443

ASPERULA; Dodoens, pemptad. 355 (1583) from Dalechamps (1554).

A. geminifolia; F. v. M., fragm. V, 147 (1865) — S.A. — V. N.S.W. Q. — B.f.III,443 M.fr.IX,188; XI,27.
 A. oligantha; F. v. M. in Neerl. Kruidk. Arch. IV, 111 et 112 (1859) — S.A. T. V. N.S.W. Q. — B.f.III,444 M.fr.IX,187; XI,27.

GALIUM; Dodoens, pemptad. 335 (1583) from Dioscorides.

G. umbrosum; Solander in G. Forster prodr. 89 (1786) — S.A. T. V. N.S.W. — — B.f.III,446 M.fr.IX,188... .. N.Z.
 G. Australe; De Candolle, prodr. IV, 608 (1830) W.A. S.A. T. V. N.S.W. Q. — B.f.III,446

CAPRIFOLIACEAE.

ADANSON, Familles des plantes II, 133 (1763).—Genus 1, Species 2.

SAMBUCUS; Tournefort, inst. 606, t. 396 (1700) from Dodoens (1583) and Plinius. (Tripetelus.)

S. xanthocarpa; F. v. M. in Transact. phil. Soc. Vict. I, 42 (1855) — — — V. N.S.W. Q. — B.f.III,398
 S. Gaudichaudiana; De Candolle, prodr. IV, 322 (1830) — S.A. T. V. N.S.W. Q. — B.f.III,398

PASSIFLOREAE.

A. L. DE JUSSIEU in Ann. du Mus. VI, 102 (1805).—Genera 2, Species 6.

PASSIFLORA; Plukenet, phytoph. 202 et 282, t. 104, 210, 211, 212 (1692). (Disemma).

P. Herbertiana; Lindley, Bot. Regist. t. 737 (1823) — — — — — N.S.W. Q. — B.f.III,311 M.fr.IX,68.
 P. cinnabarina; Lindley in Gard. Chron. 724 (1855) — — — V. N.S.W. — — — M.fr.IX,68.
 P. glabra; Wendland, collect. pl. I, 55, t. 17 (1808) — — — — — N.S.W. — — —
 P. aurantia; G. Forster, fl. ins. Austr. prodr. 62 (1786) — — — — — N.S.W. Q. — B.f.III,312 M.fr.IX,68 P.
 P. brachystephanea; F. v. M. in Benth. Fl. Austr. III, 312 (1866) — — — — — N.S.W. Q. — B.f.III,312 M.fr.I,56.

MODECCA, Lamarck, Encycl. méthod. IV, 208 (1797).

M. Australis; R. Brown in De Candolle prodr. III, 337 (1828) — — — — — Q. N.A. B.f.III,313 M.fr.IX,69.

CUCURBITACEAE.

HALLER, enum. stirp. Helv. praef. 34 (1742).—Genera 10, Species 27.

TRICHOSANTHES; Linné, gen. plant. 295 (1737).

T. Holtzei; F. v. M. in Austr. Journ. Pharm. I, 447 (1886) — — — — — — N.A. —
 T. pentaphylla; F. v. M. in Benth. Fl. Austr. III, 314 (1866) — — — — — Q. — B.f.III,314

T. cucumerina; Linné, spec. plant. 1008 (1753)	...	...	...	...	...	...	...	...	Q. N.A. B.fl.III,314	...	...	...	As.
T. subvelutina; F. v. M. in A. & C. De Candolle, mon. phan. III, 366 (1881)	...	...	...	...	...	...	...	N.S.W.	Q. — B.fl.III,315	M.fr.VI,187.			
T. Hearnii; F. v. M. in Benth. Fl. Austr. III, 315 (1866)	...	...	...	...	...	...	...	...	Q. — B.fl.III,315				
T. Muelleri; Cogniaux in F. v. M., Cens. fourth suppl. 4 (1889)	...	...	...	...	...	...	...	...	— N.A. —				
LAGENARIA; Séringé in Mém. Soc. Genév. III, 25, t. 2 (1825).													
L. vulgaris; Séringé in De Candolle, prodr. III, 279 (1828)	...	...	...	...	...	...	...	...	Q. — B.fl.III,316	M.fr.VII,62...	...	As., Afr.	
LUFFA; Tournefort in act. Acad. Paris. 84, t. 2 (1706), from Vesling (1638).													
L. Aegyptiaca; Miller, Gardener's Dictionary (1731)	...	...	...	...	...	...	...	...	Q. N.A. B.fl.III,316	M.fr.III,107	...	As., Afr., P.	
L. graveolens; Roxburgh, Fl. Ind. III, 716 (1832)	...	...	...	...	...	...	...	...	— N.A. B.fl.III,316	M.fr.III,106; IV,175	...	As., P.	
L. foetida; Cavanilles, icon. I, 7, t. 9—10 (1791)	...	...	...	...	...	...	...	...	— N.A. —	M.fr.XI,138...	...	As., Afr., P.	
ZANONIA; Linné, coroll. 19 (1737). (Alsomitra.)													
Z. Capricornica; F. v. M., fragm. VII, 61 (1870)	...	...	...	...	...	...	...	...	Q. — —	M.fr.VII,61			
Z. Hookeri; F. v. M., fragm. VI, 188 (1868)	...	...	...	...	...	...	...	...	Q. — —	M.fr.V,181; VII,62.			
Z. Stephensiana; F. v. M., fragm. VIII, 181 (1874)	...	...	...	...	...	...	...	...	Q. — —	M.fr.VII,181.			
CUCUMIS; Tournefort, inst. 104, t. 31 (1700) from Plinius.													
C. trigonus; Roxburgh, Fl. Ind. III, 722 (1832)	...	...	...	...	...	W.A. — — —	N.S.W.	Q. N.A. B.fl.III,317	M.fr.VI,187...	...	As.		
C. Chate; Linné, syst. nat. edit. decim. 5 (1759)	...	...	...	...	...	W.A. S.A. — — —	N.S.W.	Q. N.A. —	...	...	As., Afr.		
BENINCASA; Savi in Bibl. Ital. IX, 158 (1818).													
B. vacua; F. v. M., Cens. of Austr. pl. 76 (1882)	...	...	...	...	...	...	...	...	Q. — —	M.fr.VI,186.			
MOMORDICA; Tournefort, institut. rei herbar. 103, t. 29 et 30 (1700).													
M. Balsamina; Linné, sp. pl. 1009 (1753)	...	...	...	...	...	...	...	N.S.W.	Q. — B.fl.III,318	M.fr.VI,187...	...	As., Afr.	
M. Charantia; Linné, sp. pl. 1009 (1753)	...	...	...	...	...	...	S.A. — — —	N.S.W.	Q. N.A. —	...	...	As., Afr.	
BRYONOPSIS; Arnott in Hook. Journ. of Bot. III, 274 (1841). (Bryonia partly.)													
B. laciniosa; Naudin in Annal. des sc. nat. sér. cinq. VI, 30 (1869)	...	...	...	...	...	...	...	N.S.W.	Q. N.A. B.fl.III,319	...	...	As., Afr.	
B. Pancheri; Naudin in Annal. des sc. nat. sér. cinq. V, 30 (1869)	...	...	...	...	...	...	...	N.S.W.	— — —	...	...	P.	
MELOTHRIA; Linné, coroll. 1 (1737). (Zehneria, Mukia.)													
M. Cunninghamii; Benth. Fl. Austr. III, 320 (1866)	...	...	...	...	...	...	...	N.S.W.	Q. N.A. B.fl.III,320	M.fr.VI,188; VII,62.			
M. Baueriana; F. v. M., fragm. VI, 188 (1866)	...	...	...	...	...	...	...	N.S.W.	— — —	M.fr.VI,188.			
M. Muelleri; Benth. Fl. Austr. III, 320 (1866)	...	...	...	...	...	...	W.A. S.A. — V.	N.S.W.	Q. — B.fl.III,320	M.fr.I,182; II,180; III,107.			
M. Maderaspatana; Cogniaux in De Candolle mon. phan. III, 623 (1881)	...	...	...	...	...	...	W.A. S.A. — — —	N.S.W.	Q. N.A. B.fl.III,321	M.fr.VI,187	...	As., Afr., P.	
M. subpellucida; Cogniaux in F. v. M., Cens. fourth suppl. 4 (1889)	...	...	...	...	...	...	...	...	Q. N.A. —				
M. Celebica; Cogniaux in De Candolle, mon. phan. III, 625 (1881)	...	...	...	...	...	...	...	...	Q. N.A. —	...	...	As.	
SICYOS; Linné, syst. nat. 9 (1735); gen. plant. 297 (1737). (Sycios.)													
S. angulata; Linné, spec. plant. 1013 (1753)	...	...	...	...	...	...	...	T. V. N.S.W.	Q. — B.fl.III,322	M.fr.VI,187; VII,62	...	Afr., Am., N.Z., P.	

COMPOSITAE.

VAILLANT in act. Acad. Paris. 143 (1718).—Genera 90, Species 539.

ETHULIA; Linné, fil. decas. prima plant. rar. hort. Upsal. 1, t. 1 (1762).

E. conyzoides; Linné, fil. dec. prim. 1, t. 1 (1762) N.S.W. Q. — — ... As., Afr.

CENTRATHERUM; Cassini in Bull. de la Soc. philom. 31 (1817).

C. muticum; Lessing in Schlecht. Linnaea II, 320 (1829) N.S.W. Q. — B.fl.III,460 Am., P.

PLEUROCARPAEA; Benth. Fl. Austr. III, 460 (1866).

P. denticulata; Benth. Fl. Austr. III, 460 (1866) Q. N.A. B.fl.III,460

B. cheilocarpa; F. v. M. in Wing's South. Sc. Record, Aug. (1882) ...	...	W.A.	—	—	—	—	—	—	—	—	—
B. ciliaris; Lessing, syn. comp. 172 (1832) ...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	—	B.fl.III, 518	
B. calocarpa; F. v. M. in Schlecht. Linnaea XXV, 399 (1852) ...	...	—	S.A.	—	V.	N.S.W.	—	—	—	B.fl.III, 519	M.fr.XI, 27.
B. chrysoglossa; F. v. M. in Transact. phil. Soc. Vict. I, 44 (1854) ...	...	—	S.A.	—	V.	N.S.W.	—	—	—	—	—
B. marginata; Benth. in Hueg. enum. 60 (1837) ...	...	—	—	—	—	N.S.W.	Q.	—	—	B.fl.III, 519	
B. Sieberi; De Candolle, prodr. V, 306 (1836) ...	...	—	—	—	—	N.S.W.	—	—	—	B.fl.III, 520	
B. discolor; C. Stuart in Benth. Fl. Austr. III, 520 (1866) ...	...	—	—	—	—	N.S.W.	Q.	—	—	B.fl.III, 520	
B. multifida; De Candolle, prodr. V, 306 (1836) ...	...	—	—	—	V.	N.S.W.	Q.	—	—	B.fl.III, 520	M.fr.VIII, 145.
B. latisquamea; F. v. M., fragm. XI, 16 (1878) ...	...	W.A.	—	—	—	—	—	—	—	—	M.fr.XI, 16.
B. glandulosa; Benth. Fl. Austr. III, 520 (1866) ...	...	W.A.	—	—	—	—	—	—	—	B.fl.III, 520	
B. collina; Benth. Fl. Austr. III, 521 (1866) ...	...	W.A.	S.A.	—	V.	N.S.W.	—	—	—	B.fl.III, 521	

ERODIOPHYLLUM; F. v. M., fragm. IX, 119 (1875). (Subgenus of Erigeron, aect. Trimorphaea.)

E. Elderi; F. v. M., fragm. IX, 120 (1875) ...	...	—	S.A.	—	—	—	—	—	—	—	M.fr.IX, 119.
--	-----	---	------	---	---	---	---	---	---	---	---------------

MINURIA; De Candolle, prodr. V, 298 (1836). (Therogeron, Elachothamnus, Kippistia.)

M. leptophylla; De Candolle, prodr. V, 298 (1836) ...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.III, 498	M.fr.IX, 119; X, 56.	
M. Cunninghamii; Benth. Fl. Austr. III, 498 (1866) ...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III, 498		
M. Candollei; F. v. M., fragm. IX, 119 (1875) ...	...	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.III, 499	M.fr.V, 87; X, 56.	
M. suaedifolia; F. v. M. in Benth. Fl. Austr. III, 499 (1866) ...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III, 499	M.fr.X, 56.	

ACHNOPHORA; F. v. M. in Proceed. Roy. Soc. South Austr. VI, 32 (1883).

A. Tatei; F. v. M. in Proc. Roy. Soc. S. Austr. VI, 33 (1883) ...	...	—	S.A.	—	—	—	—	—	—	—	
---	-----	---	------	---	---	---	---	---	---	---	--

CALOTIS; R. Brown in Bot. Regist. t. 504 (1820). (Huenefeldia, Goniopogon, Cheiroloma.)

C. dentex; R. Brown in Bot. Regist. 504 (1820) ...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.III, 501		
C. cuneifolia; R. Brown in Bot. Regist. 504 (1820) ...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III, 501		
C. glandulosa; F. v. M. in Transact. Vict. Inst. 129 (1855) ...	...	—	—	—	V.	N.S.W.	—	—	B.fl.III, 502		
C. cymbacantha; F. v. M. in Schlecht. Linnaea XXV, 400 (1852) ...	...	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.III, 502		
C. erinacea; Steetz in Lehm. pl. Preiss. I, 424 (1845) ...	...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III, 502		
C. acabiosifolia; Sonder & F. v. M. in Linnaea XXV, 471 (1852) ...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III, 503	M.fr.XI, 27.	
C. scapigera; Hooker in Mitch. Trop. Austr. 75 (1848) ...	...	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.III, 503	M.fr.XI, 27.	
C. anthemoides; F. v. M. in Transact. phil. soc. Vict. I, 44 (1854) ...	...	—	—	—	V.	N.S.W.	—	—	B.fl.III, 504		
C. lappulacea; Benth. in Hueg. enum. 60 (1837) ...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III, 504		
C. microcephala; Benth. Fl. Austr. III, 504 (1866) ...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III, 504		
C. breviseta; Benth. in Hueg. enum. 60 (1837) ...	...	—	—	—	—	—	Q.	N.A.	B.fl.III, 505		
C. plumulifera; F. v. M. in Transact. phil. Inst. Vict. III, 57 (1859) ...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III, 505		
C. porphyroglossa; F. v. M. in Benth. Fl. Austr. III, 505 (1866) ...	...	—	S.A.	—	—	N.S.W.	—	N.A.	B.fl.III, 505		
C. pteroperma; R. Brown in Benth. Fl. Austr. III, 505 (1866) ...	...	—	—	—	—	—	—	N.A.	B.fl.III, 505		
C. hispidula; F. v. M. in Transact. Vict. Inst. 130 (1855) ...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III, 506		
C. Kempei; F. v. M. in Transact. Roy. Soc. S. Austr. IV, 112 (1881) ...	...	—	S.A.	—	—	—	—	—	—	M.fr.XI, 139.	

ASTER; Tournefort, inst. 481, t. 274 (1700), from Dioscorides. (Olcaria, Celmisia, Eurybia, Steetzia, Diplopappus, Diplostephium.)

A. megaphyllus; F. v. M., fragm. V, 70 (1865) ...	...	—	—	—	V.	N.S.W.	—	—	B.fl.III, 468	M.fr.V, 70.	
A. chrysophyllus; Cunningham in De Candolle, prodr. V, 266 (1836) ...	...	—	—	—	—	N.S.W.	—	—	B.fl.III, 468	M.fr.II, 88; V, 71.	
A. alpicola; F. v. M., fragm. V, 70 (1865) ...	...	—	—	—	V.	N.S.W.	—	—	B.fl.III, 468	M.fr.V, 70.	
A. rosmarinifolius; Cunningham in De Candolle, prodr. V, 268 (1836) ...	...	—	—	—	—	N.S.W.	—	—	B.fl.III, 468		
A. viscosus; Labillardière, Nov. Holl. pl. spec. II, 53, t. 203 (1806) ...	...	—	—	—	T.	V.	—	—	B.fl.III, 468	M.fr.V, 71.	
A. Sonderi; F. v. M., fragm. V, 83 (1865) ...	...	—	S.A.	—	—	—	—	—	B.fl.III, 469	M.fr.V, 83.	
A. pannosus; F. v. M., fragm. V, 83 (1865) ...	...	—	S.A.	—	V.	—	—	—	B.fl.III, 469	M.fr.V, 83.	
A. oliganthemus; F. v. M., Cens. of Austr. pl. 78 (1882) ...	...	—	—	—	—	N.S.W.	—	—	B.fl.III, 469		
A. argophyllus; Labillardière, Nov. Holl. pl. spec. II, 52, t. 201 (1806) ...	...	—	—	—	T.	V.	N.S.W.	—	B.fl.III, 470	M.fr.V, 68.	
A. cydonifolius; Cunningham in De Candolle, prodr. V, 267 (1836) ...	...	—	—	—	—	N.S.W.	—	—	B.fl.III, 470		

A. myrsinoides; Labillardière, Nov. Holl. pl. spec. II, 53, t. 202 (1806)...	..	—	S.A.	T.	V.	N.S.W.	—	—	B.fl. III, 471	M.fr. V, 69.
A. persoonioides; Cunningham in De Candolle, prodr. V, 267 (1836) ...	...	—	—	T.	—	—	—	—	B.fl. III, 471	M.fr. V, 69.
A. obcordatus; F. v. M., fragm. V, 69 (1865) ...	...	—	—	T.	—	—	—	—	B.fl. III, 471	M.fr. V, 69.
A. pinifolius; F. v. M., fragm. V, 71 (1865) ...	...	—	—	T.	—	—	—	—	B.fl. III, 472	M.fr. V, 71.
A. ledifolius; Cunningham in De Candolle, prodr. V, 269 (1836)...	...	—	—	T.	—	—	—	—	B.fl. III, 472	
A. dentatus; Andrews, Bot. Reposit. t. 61 (1799) ...	...	—	—	—	V.	N.S.W.	—	—	B.fl. III, 472	M.fr. V, 82.
A. stellulatus; Labillardière, Nov. Holl. pl. spec. II, 50, t. 196 (1806) ...	...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl. III, 473	M.fr. V, 81.
A. asterotrichus; F. v. M., fragm. V, 79 (1865) ...	...	—	S.A.	—	V.	N.S.W.	—	—	B.fl. III, 473	M.fr. I, 111; II, 180.
A. gravis; F. v. M., fragm. V, 82 (1865) ...	...	—	—	—	—	N.S.W.	—	—	B.fl. III, 474	M.fr. V, 82.
A. Nernstii; F. v. M., fragm. V, 81 (1865) ...	...	—	—	—	—	N.S.W.	Q.	—	B.fl. III, 474	M.fr. V, 81.
A. hygrophilus; Cunningham in De Candolle, prodr. V, 269 (1836) ...	...	—	—	—	—	—	Q.	—	B.fl. III, 474	
A. Siemsseni; F. v. M., fragm. V, 71 (1865)...	...	—	—	—	—	N.S.W.	—	—	B.fl. III, 475	M.fr. I, 50.
A. tubuliflorus; F. v. M., fragm. V, 65 (1865) ...	...	—	S.A.	—	V.	N.S.W.	—	—	B.fl. III, 475	M.fr. V, 65.
A. axillaris; F. v. M., fragm. V, 64 (1865) ...	...	W.A.	S.A.	T.	V.	N.S.W.	—	N.A.	B.fl. III, 475	M.fr. V, 64.
A. revolutus; F. v. M., Cens. of Austr. pl. 78 (1882) ...	...	W.A.	—	—	—	—	—	—	B.fl. III, 476	
A. exilifolius; F. v. M., fragm. V, 69 (1865) ...	...	W.A.	—	—	—	—	—	—	B.fl. III, 476	M.fr. V, 69.
A. ramulosus; Labillardière, Nov. Holl. plant. spec. II, 51, t. 198 (1806)	...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl. III, 476	
A. florulentus; F. v. M., fragm. V, 70 (1865) ...	...	—	—	T.	V.	N.S.W.	—	—	B.fl. III, 477	M.fr. V, 82.
A. microphyllus; Ventenat, Jard. de Malmaison. 83 (1803) ...	...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl. III, 478	M.fr. V, 70.
A. lepidophyllus; Persoon, synop. plant. II, 442 (1807) ...	...	—	S.A.	T.	V.	N.S.W.	—	—		
A. Mitchelli; F. v. M., fragm. V, 78 (1865) ...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl. III, 478	M.fr. V, 78.
A. exiguifolius; F. v. M., fragm. V, 67 (1865) ...	...	W.A.	S.A.	—	—	—	—	—	B.fl. III, 478	M.fr. V, 67.
A. Cassiniae; F. v. M., fragm. V, 68 (1865) ...	...	W.A.	—	—	—	—	—	—	B.fl. III, 479	M.fr. V, 68.
A. cyanodiscus; F. v. M., fragm. V, 82 (1865) ...	...	—	—	—	—	N.S.W.	—	—	B.fl. III, 479	M.fr. V, 82.
A. pimeleoides; Cunningham in De Candolle, prodr. V, 268 (1836) ...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl. III, 479	
A. iodochrus; F. v. M., fragm. V, 81 (1865)...	...	—	—	—	V.	N.S.W.	—	—	B.fl. III, 480	M.fr. II, 110.
A. adenolasius; F. v. M., fragm. V, 67 (1865) ...	...	W.A.	—	—	—	—	—	—	B.fl. III, 480	M.fr. V, 67.
A. calcareus; F. v. M., Cens. of Austr. pl. 78 (1882) ...	...	—	S.A.	—	V.	N.S.W.	—	—	B.fl. III, 481	
A. magniflorus; F. v. M., fragm. V, 80 (1865) ...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl. III, 481	M.fr. V, 80.
A. Muelleri; Cens. 78 (1882), from Sonder (1852) ...	...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl. III, 481	
A. Stuartii; F. v. M., fragm. V, 76 (1865) ...	...	—	S.A.	—	—	—	—	—	B.fl. III, 481	M.fr. I, 202.
A. decurrens; Cunningham in De Candolle, prodr. V, 269 (1836) ...	...	—	S.A.	—	V.	N.S.W.	—	—	B.fl. III, 481	
A. glutescens; F. v. M., fragm. V, 77 (1865) ...	...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl. III, 482	M.fr. V, 77.
A. orarius; F. v. M., fragm. V, 78 (1865) ...	...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl. III, 482	M.fr. V, 78.
A. vernicosus; F. v. M., fragm. V, 67 (1865) ...	...	W.A.	—	—	—	—	—	—	B.fl. III, 482	M.fr. V, 67.
A. teretifolius; F. v. M., fragm. V, 77 (1865) ...	...	—	S.A.	—	V.	N.S.W.	—	—	B.fl. III, 483	M.fr. V, 77.
A. Turczaninowii; F. v. M., fragm. V, 67 (1865) ...	...	W.A.	—	—	—	—	—	—	B.fl. III, 483	M.fr. V, 67.
A. ellipticus; Cunningham in De Candolle, prodr. V, 271 (1836) ...	...	—	—	—	—	N.S.W.	Q.	—	B.fl. III, 483	M.fr. I, 16, 242.
A. glandulosus; Labillardière, Nov. Holl. pl. spec. II, 50, t. 197 (1806)...	...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl. III, 484	
A. heleophilus; F. v. M., fragm. V, 66 (1865) ...	...	W.A.	—	—	—	—	—	—	B.fl. III, 484	M.fr. V, 66.
A. muricatus; F. v. M., fragm. V, 66 (1865) ...	...	W.A.	—	—	—	—	—	—	B.fl. III, 484	M.fr. V, 66.
A. Benthami; F. v. M., Cens. of Austr. pl. 78 (1882) ...	...	—	—	—	V.	—	—	—	B.fl. III, 485	
A. Steetzii; F. v. M., fragm. V, 66 (1865) ...	...	W.A.	—	—	—	—	—	—	B.fl. III, 485	M.fr. V, 66.
A. Cunninghami; F. v. M., Cens. 78 (1882), from De Candolle (1836) ..	...	—	—	—	—	N.S.W.	—	—	B.fl. III, 486	
A. paucidentatus; F. v. M., fragm. V, 66 (1865) ...	...	W.A.	—	—	—	—	—	—	B.fl. III, 485	M.fr. V, 66.
A. megalodontus; F. v. M., fragm. VIII, 244 (1874) ...	...	W.A.	S.A.	—	—	—	—	—		M.fr. VIII, 244.
A. adenophorus; F. v. M., fragm. V, 78 (1865) ...	...	—	—	—	V.	N.S.W.	—	—	B.fl. III, 486	M.fr. I, 111.
A. homolepis; F. v. M., fragm. V, 65 (1865)...	...	W.A.	—	—	—	—	—	—	B.fl. III, 486	M.fr. V, 65.
A. xerophilus; F. v. M., fragm. V, 76 (1865) ...	...	—	—	—	—	—	Q.	—	B.fl. III, 486	M.fr. I, 51, 76 & 86.
A. Ferresii; F. v. M., fragm. V, 75 (1865) ...	...	—	S.A.	—	—	—	—	N.A.	B.fl. III, 487	M.fr. III, 18.

A. exul ; Lindley, Bot. Regist. XXV, App. 24 (1839)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,487	M.fr.I,51; V,75.		
A. argutus ; R. Brown in Benth. Fl. Austr. III, 488 (1866)	—	—	—	—	—	—	N.A.	B.fl.III,488			
A. Huegelii ; F. v. M., fragm. V, 79 (1865)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,488	M.fr.V,79.		
A. Balli ; F. v. M., fragm. VIII, 143 (1874)	—	—	—	—	N.S.W.	—	—	—	M.fr.VIII,143.		
A. Mooneyi ; F. v. M., fragm. VIII, 144 (1874)	—	—	—	—	N.S.W.	—	—	—	M.fr.VIII,144.		
A. Celmisia ; F. v. M., fragm. V, 84 (1865)	—	—	T.	V.	N.S.W.	—	—	B.fl.III,489	M.fr.V,84	...	N.Z.

VITTADINIA ; Ach. Richard, Voy. de l'Astrol. bot. 250 (1832). (Eurybiopsis, Microgyne.)

V. brachycomoides ; F. v. M., fragm. V, 86 (1865)	—	—	—	—	—	Q.	N.A.	B.fl.III,490	M.fr.V,86.		
V. Australis ; A. Richard, Voy. de l'Astrol. 251 (1832)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,490	M.fr.V,87; XI,20		N.Z.
V. scabra ; De Candolle, prodr. V, 281 (1836)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,491			
V. macrorrhiza ; A. Gray in Proc. Am. Acad. V, 118 (1863)	—	—	—	—	—	—	N.A.	B.fl.III,492			

PODOCOMA ; Cassini in Bull. de la Soc. philom. 137 (1817). (Ixiochlamys.)

P. cuneifolia ; R. Brown, App. to Sturt, Centr. Austr. 17 (1849)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.III,493			
---	------	------	---	---	--------	----	------	--------------	--	--	--

ERIGERON ; Linné, hort. Cliffort. 407 (1737). (Haplopappus partly.)

E. pappochromus ; Labillardière, Nov. Holl. pl. spec. II, 47, t. 193 (1806) ..	—	—	T.	V.	N.S.W.	—	—	B.fl.III,494	M.fr.V,87.		
E. ambiguus ; F. v. M. in Transact. phil. Inst. Vict. III, 58 (1858)	—	—	—	—	—	Q.	N.A.	B.fl.III,494			
E. sessilifolius ; F. v. M., fragm. X, 100 (1881)	—	—	—	—	—	—	N.A.	—	M.fr.X,137.		
E. minurioides ; Benth. Fl. Austr. III, 495 (1866)	—	—	—	—	V.	—	—	B.fl.III,495			
E. conyzoides ; F. v. M. in Transact. phil. Soc. Vict. I, 105 (1855)	—	—	—	—	V.	N.S.W.	—	B.fl.III,495	M.fr.V,87.		

CONYZA ; Lamarck, Encycl. méthod. II, 87, 89 (1786).

C. viscidula ; Wallich, numer. list 3006 (1828)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,496...	...	...	As., P.
C. Aegyptiaca ; Aiton, hort. Kew. III, 183 (1789)	—	—	—	—	—	Q.	N.A.	B.fl.III,497...	...	...	As., Afr.

BLUMEA ; De Candolle in Guillemin, Arch. de Bot. II, 514 (1833).

B. glandulosa ; De Candolle in Wight, contrib. 14 (1834)	—	—	—	—	—	Q.	N.A.	B.fl.III,525...	...	...	As., Afr.
B. lacera ; De Candolle in Wight, contrib. 14 (1834)	—	—	—	—	—	Q.	N.A.	B.fl.III,526...	...	...	As., Afr.
B. amplexans ; De Candolle in Wight, contrib. 13 (1834)	—	—	—	—	—	Q.	—	B.fl.III,525...	...	...	As.
B. integrifolia ; De Candolle, prodr. V, 433 (1836)	—	—	—	—	—	Q.	N.A.	B.fl.III,525			
B. diffusa ; R. Brown in Benth. Fl. Austr. III, 525 (1866)	—	—	—	—	—	Q.	N.A.	B.fl.III,525			
B. hieracifolia ; De Candolle in Wight, contrib. 15 (1834)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.II,526...	...	...	As.
B. Cunninghamii ; De Candolle, prodr. V, 435 (1836)	—	—	—	—	—	Q.	N.A.	B.fl.III,526			
B. acutata ; De Candolle in Ann. du Mus. d'hist. nat. Par. III, 409 (1834)	—	—	—	—	—	Q.	—	—	M.fr.IX,160.	...	As.

PLUCHEA ; Cassini in Bull. de la Soc. philom. 31 (1817). (Spiropodium, Eyrea.)

P. Indica ; Lessing in Schlecht. Linnaea VI, 150 (1831)	—	—	—	—	—	Q.	N.A.	B.fl.III,527	M.fr.X,101	...	As., P.
P. Dioscoridis ; De Candolle, prodr. V, 450 (1836)	—	—	—	—	—	Q.	—	—	...	...	As., Afr.
P. tetranthera ; F. v. M. Rep. Babb. Exped. 12 (1860)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.III,528			
P. concephala ; F. v. M. in Transact. Roy. Soc. Vict. 24, 138 (1887)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,480	M.fr.V,79.		
P. baccharoides ; F. v. M. in Benth. Fl. Austr. III, 528 (1866)	—	—	—	—	—	Q.	N.A.	B.fl.III,528	M.fr.I,34.		
P. Eyrea ; F. v. M. Rep. Babb. Exped. 11, 12 (1860)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.III,528			
P. squarrosa ; Benth. Fl. Austr. III, 529 (1866)	W.A.	—	—	—	—	—	N.A.	B.fl.III,529			
P. dentex ; R. Brown in Benth. Fl. Austr. III, 529 (1866)	—	—	—	—	—	Q.	—	B.fl.III,529			

PTERIGERON ; De Candolle, prodr. V, 293 (1836). (Streptoglossa, Oliganthemum.)

P. decurrens ; De Candolle, prodr. V, 293 (1836)	W.A.	—	—	—	—	—	N.A.	B.fl.III,531			
P. liatroides ; Benth. Fl. Austr. III, 532 (1866)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.III,532			
P. odoratus ; Benth. Fl. Austr. III, 532 (1866)	—	—	—	—	—	Q.	N.A.	B.fl.III,532			
P. macrocephalus ; Benth. Fl. Austr. 532 (1866)	—	—	—	—	—	Q.	N.A.	B.fl.III,532			
P. microglossus ; Benth. Fl. Austr. III, 532 (1866)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.III,532			

- P. dentatifolius*; F. v. M., fragm. IX, 119 (1875) ... — S.A. — — — — — M.fr.X,119.
P. adscendens; Benth. Fl. Austr. III, 533 (1866) ... — S.A. — — — — — Q. N.A. B.f.III,533
P. filifolius; Benth. Fl. Austr. III, 533 (1866) ... — — — — — — N.A. B.f.III,533

THESPIDIUM; F. v. M. in Journ. of Landsborough's Exped. app. (1862).

- T. basiflorum*; F. v. M. in Journ. of Landsb. Exped. App. (1862) ... — — — — — Q. N.A. B.f.III,534 M.fr.XI,101.

COLEOCOMA; F. v. M. in Hook. Kew Misc. IX, 19 (1857).

- C. Centaurea*; F. v. M. in Hook. Kew Misc. IX, 19 (1857) ... — — — — — — N.A. B.f.III,533 M.fr.X,101.

EPALTES; Cassini in Bull. de la Soc. philom. 139 (1818). (Sphaeromorphaea partly, Ethuliopsis, Gymnaphanes.)

- E. Cunninghamii*; Benth. Fl. Austr. III, 530 (1866) ... — S.A. — V. N.S.W. Q. — B.f.III,530 M.fr.II,155.
E. Australis; Lessing in Linnaea V, 148 (1831) ... — S.A. — V. N.S.W. Q. N.A. B.f.III,530 M.fr.XI,101 ... As.
E. pleiochaeta; F. v. M., fragm. X, 100 (1877) ... — — — — N.S.W. — — — M.fr.X,100.
E. Harrisii; F. v. M., fragm. XI, 101 (1881) ... — — — — — Q. — — — M.fr.XI,101.
E. Tatei; F. v. M. in Transact. Roy. Soc. of S.A. VI, 31 (1883) ... — — — — — — — —

SPHAERANTHOS; Vaillant in Act. Acad. Par. 289 (1719). (Sphaeranthus.)

- S. Indicus*; Linné, sp. pl., 927 (1753) ... — — — — — — Q. N.A. B.f.III,521 M.fr.III,138 ... As., Afr.
S. Africanus; Linné, sp. pl. ed. sec. 13, 14 (1762) ... — — — — — — — N.A. B.f.III,522 M.fr.III,138 ... As., Afr.

PTEROCAULON; Elliot, Sketch of the Bot. of S. Carolina and Georgia II, 323 (1824). (Monenteles.)

- P. verbascifolius*; Benth. & J. Hooker, gen. pl. II, 294 (1873) ... — — — — — Q. N.A. B.f.III,523
P. Billardieri; F. v. M., Papuan plants, 43 (1876) ... — S.A. — — N.S.W. Q. N.A. B.f.III,523 M.fr.XI,101 ... As., P.
P. sphacelatus; Benth. & J. Hooker, gen. plant. II, 94 (1873) ... — — — — — W.A. S.A. — — N.S.W. Q. N.A. B.f.III,523 ... P.
P. glandulosus; Benth. & J. Hooker, gen. plant. II, 94 (1873) ... — — — — — — Q. N.A. B.f.III,523
P. sphaeranthoides; Benth. & J. Hooker, gen. plant. II, 94 (1873) ... — — — — — — — N.A. B.f.III,524

STUARTINA; Sonder in Schlechtendal's Linnaea XXV, 521 (1852).

- S. Mnelleri*; Sonder in Schlecht. Linnaea XXV. 522 (1852) ... — S.A. — V. N.S.W. Q. — B.f.III,657 M.fr.VIII,145; XI,27.

GNAPHALIUM; Linné, gen. plant. 250 (1737), from J. & C. Bauhin (1619). (Euchiton.)

- G. luteo-album*; Linné, spec. plant. 851 (1753) ... — — — — — W.A. S.A. T. V. N.S.W. Q. N.A. B.f.III,653 M.fr.V,150 E., As., Afr., Am., P., N.Z.
G. Japonicum; Thunberg, Fl. Japon. 311 (1784) ... — — — — — W.A. S.A. T. V. N.S.W. Q. — B.f.III,653 M.fr.V,150; XI,27. As., N.Z.
G. alpinum; F. v. M., second gen. Report 12 (1854) ... — — — — — T. V. — — — — B.f.III,654 M.fr.V,150.
G. purpureum; Linné, spec. plant. 854 (1753) ... — — — — — — — — N.S.W. Q. — B.f.III,655... ... Am.
G. Indicum; Linné, spec. plant. 852 (1753) ... — — — — — S.A. — — N.S.W. Q. N.A. B.f.III,655 M.fr.V,149 ... As., Afr., P.
G. indutum; J. Hooker in Lond. Journ. of Bot. VI, 121 (1847) ... — — — — — W.A. S.A. T. V. N.S.W. Q. — B.f.III,655 M.fr.V,150.
G. Traversii; J. Hooker, Handb. N. Zeal. Fl. 154 ... — — — — — V. N.S.W. — — — B.f.III,656... ... N.Z.

ANTENNARIA; Gaertner, de fruct. II, 410, t. 167 (1791). (Raoulia partly.)

- A. Planchoni*; F. v. M., Cens. 80 (1882), from J. Hook. (1860) ... — — — — — T. — — — — B.f.III,651
A. uniceps; F. v. M. in Transact. phil. Soc. Vict. t. 105 (1854) ... — — — — — V. N.S.W. — — — B.f.III,652 M.fr.VIII,145.

LEONTOPODIUM; R. Brown in Transact. Linn. Soc. XII, 124 (1817). (Raoulia partly.)

- L. Catipes*; F. v. M. in Papers of the Roy. Soc. of Tasm. 44 (1882) ... — — — — — T. V. N.S.W. — — — B.f.III,651
L. Meredithae; F. v. M., Cens. of Austr. pl. 80 (1882) ... — — — — — T. — — — — —

PTERYGOPAPPUS; J. Hooker in Lond. Journ. of Bot. VII, 120 (1847).

- P. Lawrencii*; J. Hooker in Lond. Journ. VII, 120 (1847) ... — — — — — T. — — — — — B.f.III,656

PODOSPERMA; Labillardière, Nov. Holl. plant. specim. II, 35, t. 177 (1806). (Podospermum, Podotheca, Phaenopoda, Lophoclinium.)

- P. gnaphalioides*; F. v. M., fragm. XII, 22 (1882) from Graham (1842) ... — — — — — W.A. — — — — — B.f.III,601 M.fr.XII,21.
P. angustifolium; Labillardière, Nov. Holl. pl. specim. II, 35, t.177 (1806) ... — — — — — W.A. S.A. T. V. N.S.W. — — — B.f.III,601 M.fr.XI,20; XII,21.
P. pygmaeum; F. v. M., fragm. XII, 22 (1882) implied (1852) ... — — — — — W.A. — — — — — B.f.III,602

P. chrysanthia; F. v. M., fragm. XII, 22 (1882)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.III,602	M.fr.XII,22.
P. Polakii; F. v. M., fragm. XII, 21 (1882) ...	...	...	...	...	...	W.A.	—	—	—	—	—	—	M.fr.XII,21.

IXIOLAENA; Bentham in Hueg. enum. 66 (1837).

I. brevicompta; F. v. M., fragm. I, 53 (1858)	...	...	...	...	...	—	—	—	N.S.W.	Q.	—	B.fl.III,597	M.fr.I,53.
I. leptolepis; Benthani, Fl. Austr. III, 597 (1866)	...	...	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.III,597
I. supina; F. v. M. in Transact. Vict. Inst. 37 (1855)	...	...	...	...	...	—	S.A.	T.	—	—	—	—	B.fl.III,598
I. tomentosa; Sonder & F. v. M. in Linnaea XXV, 504 (1852)	...	...	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,598 M.fr.I,53; XI,27.
I. viscosa; Benthani in Hueg. enum. 66 (1837)	...	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.III,598

PODOLEPIS; Labillardière, Nov. Holl. pl. specim. II, 57, t. 28 (1806). (Scalia, Panaetia, Scaliopsis, Siemssenia, Stylolepis, Rutidochlamys.)

<i>P. rhytidochlamys</i> ; F. v. M., fragm. IV, 79 (1864) ...	...	...	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl. III, 603	M.fr. IV, 79.
<i>P. longipedata</i> ; Cunningham in De Candolle, prodr. VI, 163 (1837) ...	...	...	...	...	...	—	—	—	V.	N.S.W.	Q.	—	B.fl. III, 604	M.fr. I, 112.
<i>P. acuminata</i> ; R. Brown in Ait. hort. Kew. ed. sec. V, 82 (1813) ...	...	...	...	...	...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl. III, 604	M.fr. XI, 27.
<i>P. canescens</i> ; Cunningham in De Candolle, prodr. VI, 163 (1837) ...	...	...	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl. III, 605	
<i>P. aristata</i> ; Bentham in Hueg. enum. 64 (1837) ...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. III, 605	
<i>P. pallida</i> ; Turczaninow in Bull. de Mosc. XXIV, 78 (1851) ...	...	...	...	...	...	W.A.	—	—	—	—	—	N.A.	B.fl. III, 605	
<i>P. nutans</i> ; Steetz in Lehm. pl. Preiss. I, 464 (1845) ...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. III, 605	
<i>P. gracilis</i> ; Graham in Edinb. New phil. journ. V, 379 (1828) ...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. III, 606	
<i>P. rugata</i> ; Labillardière, Nov. Holl. pl. spec. II, 57, t. 208 (1806) ...	...	...	...	...	...	W.A.	S.A.	—	V.	—	—	N.A.	B.fl. III, 606	
<i>P. Lessoni</i> ; Beutham, Fl. Austr. III, 606 (1866) ...	...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl. III, 606	M.fr. XI, 27.
<i>P. Siemssenii</i> ; F. v. M., fragm. V, 200 (1866) ...	...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl. III, 607	M.fr. V, 200.
<i>P. microcephala</i> ; Bentham, Fl. Austr. 607 (1866) ...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. III, 607	
<i>P. Kendallii</i> ; F. v. M. in Wing's S. Sc. Rec. III, 68 (1883) ...	...	...	...	...	...	W.A.	—	—	—	—	—	—	—	M.fr. VIII, 168; X, 108.

ATHRIXIA ; Ker in Bot. Regist. VIII, 681 (1823). (Asteridea, Chrysodiscus, Trichostegia.)

A. Australis; Steetz in Lehm. pl. Preiss. I, 482 (1845)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,599
A. gracilis; Bentham, Fl. Austr. III, 599 (1866)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,599
A. multiceps; Bentham, Fl. Austr. III, 599 (1866)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,599
A. stricta; Bentham, Fl. Austr. III, 600 (1866)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,600
A. tenella; Bentham, Fl. Austr. III, 600 (1866)	...	...	...	...	W.A.	S.A.	—	V.	N.S.V.	—	—	B.fl.III,600
A. Croniniana; F. v. M. in Vict. Nat. V, 54 (1888)	...	...	...	...	W.A.	—	—	—	—	—	—	—
A. chaetopoda; F. v. M., fragm. X, 56 (1876)	...	...	...	...	W.A.	—	—	—	—	—	—	M.fr.X,56.

LEPTORRHYNCHOS; Lessing, syn. composit. 273 (1832). (Rhytidanthe, Doratolepis, Waitziae subgenus.)

<i>L. squamatus</i> ; Lessing, syn. comp. 273 (1832)	...	...	...	...	...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,608	M.fr.I,52.
<i>L. panaetioides</i> ; Bentham, Fl. Austr. III, 609 (1866)	...	...	...	...	...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,609	
<i>L. tenuifolius</i> ; F. v. M., fragm. I, 52 (1858)	...	...	...	...	...	—	S.A.	—	V.	—	—	—	B.fl.III,609	M.fr.I,52.
<i>L. pulchellus</i> ; F. v. M. in Linnaea XXV, 500 (1852)	...	...	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,610	M.fr.I,53; XI,27.
<i>L. Baileyi</i> ; F. v. M., fragm. X, 101 (1877)	...	...	...	...	...	—	—	—	—	—	Q.	—	M.fr.X,101.	
<i>L. elongatus</i> ; De Candolle, prodr. VI, 160 (1837)	...	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,610	M.fr.XI,27,28.
<i>L. medius</i> ; Cunningham in De Candolle, prodr. VI, 160 (1837)	...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	—	—	—	M.fr.XI,85.
<i>L. Waitzia</i> ; Sonder in Schlecht. Linnaea XXV, 501 (1852)	...	...	...	...	...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,610	
<i>L. nitidulus</i> ; De Candolle, prodr. VI, 160 (1837)	...	...	...	...	...	—	—	T.	V.	N.S.W.	—	—	B.fl.III,611	

WAITZIA; Wendland, Collect. pl. II, 13, t. 42 (1808). (Viraya, Morna, Pterochaete.)

<i>W. corymbosa</i> ; Wendland, Coll. pl. II, 13, t. 42 (1808) ...	...	...	...	W.A. S.A. —	V. N.S.W. Q. —	B.A.III, 635
<i>W. aurea</i> ; Steetz in Lehm. pl. Preiss. I, 452 (1845)...	...	...	...	W.A. — — —	— — —	B.A.III, 636
<i>W. nivea</i> ; Bentham, Fl. Austr. III, 636 (1866) ...	...	...	...	W.A. — — —	— — —	B.A.III, 636
<i>W. Steetziana</i> ; Lehmann, pl. Preiss. I, 454 (1845) ...	...	...	...	W.A. — — —	— — —	B.A.III, 636
<i>W. podolepis</i> ; Steetz in Lehm. pl. Preiss. I, 450 (1845) ...	...	...	...	W.A. — — —	— — —	B.A.III, 637
<i>W. paniculata</i> ; F. v. M. in Benth. Fl. Austr. III, 637 (1866) ...	...	...	...	W.A. — — —	— — —	B.A.III, 637

HELIPTERUM; De Candolle, prodr. VI, 211 (1837). (Argyrocome partly, Pteropogon, Rodanthe, Xyridanthe, Anisolepis, Hyalosperma, Triptilodiscus, Acroclinium, Monenyanthes, Dimorpholepis, Duttonia, Cassiniola.)

H. Manglesii; F. v. M., fragm. V, 200 (1866)	...	...	...	...	W.A.	—	—	—	—	—	B.f.III, 640	M.fr.X, 108.	
H. Margaritae; F. v. M., fragm. XI, 48 (1878)	...	...	...	...	—	—	—	—	—	N.A.	—	M.fr.XI, 48.	
H. roseum; Benth. Fl. Austr. III, 640 (1866)	...	...	...	...	W.A.	S.A.	—	—	—	N.A.	B.f.III, 640	M.fr.X, 108.	
H. fuscescens; Turczaninow in Bull. de Mosc., 80 (1851)	...	...	...	...	W.A.	—	—	—	—	—	B.f.III, 602	M.fr.XII, 22.	
H. rubellum; Benth. Fl. Austr. III, 641 (1866)	...	...	...	...	W.A.	—	—	—	—	—	B.f.III, 641	M.fr.X, 108.	
H. chlorocephalum; Benth. Fl. Austr. III, 641 (1866)	...	...	...	...	W.A.	—	—	—	—	—	B.f.III, 641		
H. anthemoides; De Candolle, prodr. VI, 216 (1837)	...	...	...	...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.f.III, 641	M.fr.X, 108.
H. polygalifolium; De Candolle, prodr. VI, 216 (1837)	...	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.f.III, 641	M.fr.X, 107, 108.
H. floribundum; De Candolle, prodr. VI, 217 (1837)	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.f.III, 642	M.fr.X, 108.
H. Pyrethrum; Benth. Fl. Austr. III, 642 (1866)	...	...	...	...	W.A.	—	—	—	—	—	B.f.III, 642	M.fr.X, 110.	
H. heteranthum; Turczaninow in Bull. de Mosc. XXIV, 198 (1851)	...	...	...	...	W.A.	S.A.	—	—	—	—	B.f.III, 642	M.fr.III, 137; X, 108.	
H. stipitatum; F. v. M. in Benth. Fl. Austr. III, 643 (1866)	...	...	...	...	—	S.A.	—	—	—	—	B.f.III, 643	M.fr.III, 133; X, 109.	
H. incanum; De Candolle, prodr. VI, 215 (1837)	...	...	...	...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.f.III, 643	
H. Cotula; De Candolle, prodr. VI, 215 (1837)	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.f.III, 644	M.fr.II, 157; III, 134; X, 110.
H. hyalospermum; F. v. M. in Benth. Fl. Austr. III, 644 (1866)	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.f.III, 644	M.fr.X, 109.
H. condensatum; F. v. M., fragm. III, 136 (1863)	...	...	...	...	W.A.	—	—	—	—	N.A.	B.f.III, 645	M.fr.III, 136.	
H. polyphyllum; F. v. M., fragm. I, 35 (1858)	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.f.III, 645	M.fr.I, 35.
H. Humboldtianum; De Candolle, prodr. VI, 216 (1866)	...	...	...	...	W.A.	—	—	—	—	N.A.	B.f.III, 645	M.fr.III, 135; X, 109.	
H. Haigii; F. v. M., fragm. X, 107 (1877)	...	...	...	...	W.A.	S.A.	—	—	—	—	—	—	M.fr.X, 107.
H. tenellum; Turczaninow in Bull. de Mosc. XXIV, 198 (1851)	...	...	...	...	W.A.	S.A.	—	—	—	—	B.f.III, 646	M.fr.III, 135; X, 109; XI, 20.	
H. strictum; Benth. Fl. Austr. III, 646 (1866)	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.f.III, 646	M.fr.X, 109.
H. corymbiflorum; Schlechtendal, Linnaea XXI, 448 (1848)	...	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.f.III, 647	M.fr.X, 109.
H. sterileseens; F. v. M. in Wing's S. Sc. Rec. II, 273 (1882)	...	...	...	...	W.A.	—	—	—	—	—	—	—	
H. pygmaeum; Benth. Fl. Austr. III, 647 (1866)	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.f.III, 647	M.fr.X, 109.
H. spicatum; F. v. M. in Benth. Fl. Austr. III, 647 (1866)	...	...	...	...	W.A.	—	—	—	—	—	B.f.III, 647	M.fr.X, 109.	
H. Charsleyae; F. v. M., fragm. VIII, 168 (1874)	...	...	...	...	W.A.	S.A.	—	—	—	—	—	—	M.fr.XI, 134.
H. moschatum; Benth. Fl. Austr. III, 648 (1866)	...	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.f.III, 648	M.fr.X, 107.
H. Tietkensii; F. v. M., fragm. VIII, 227 (1874)	...	...	...	...	—	S.A.	—	—	N.S.W.	—	—	—	M.fr.VIII, 227.
H. pterochaetum; Benth. Fl. Austr. III, 648 (1866)	...	...	...	...	—	S.A.	—	—	N.S.W.	—	—	B.f.III, 648	M.fr.X, 109.
H. Frenchii; F. v. M. in Wing's S. Sc. Rec. III, 2 (1883)	...	...	...	...	W.A.	—	—	—	—	—	—	—	
H. Forrestii; F. v. M. in Wing's S. Sc. Rec. II, 273 (1882)	...	...	...	...	W.A.	—	—	—	—	—	—	—	
H. polyccephalum; Benth. Fl. Austr. III, 649 (1866)	...	...	...	...	W.A.	—	—	—	—	—	B.f.III, 649	M.fr.III, 139; X, 109.	
H. corymbosum; Benth. Fl. Austr. III, 649 (1866)	...	...	...	...	W.A.	—	—	—	—	—	B.f.III, 649	M.fr.V, 200.	
H. laeve; Benth. Fl. Austr. III, 649 (1866)	...	...	...	...	W.A.	S.A.	—	—	N.S.W.	—	—	B.f.III, 649	M.fr.V, 200.
H. exiguum; F. v. M. in Transact. Vict. Inst. 39 (1855)	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.f.III, 644	M.fr.X, 109.
H. Dimorpholepis; Benth. Fl. Austr. III, 650 (1866)	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.f.III, 650	M.fr.X, 109; XI, 27.

HELICHRYSUM; Vaillant in Act. Acad. Par. 290 (1719), from Theophrastos and Dioscorides. (Elichrysium, Schoenia, Petalolepis, Faustula, Ozothamnus, Swammerdamia, Lawrenceella, ArgYROPHANES, Chrysocephalum, Conanthodium, Xanthochrysium, ArgYROGLOTTIS, Acanthocladium, Raoulia partly.)

H. Cassinianum; Gaudichaud in Freye. Voy. Bot. 466, t. 87 (1826)	...	...	...	...	W.A.	S.A.	—	—	—	—	—	B.fl.III,616
H. Lawrenceella; F. v. M. in Benth. Fl. Austr. III, 616 (1866)	...	...	...	...	W.A.	S.A.	—	—	—	—	—	B.fl.III,616
H. subnifolium; F. v. M., fragm. III, 134 (1863)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,616 M.fr.III,134.
H. filifolium; F. v. M., fragm. III, 134 (1863)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,617 M.fr.III,134.
H. MacIvorii; F. v. M. in Wing's S. Sc. Rec. III, 99 (1883)	...	...	...	...	W.A.	—	—	—	—	—	—	—
H. semifertile; F. v. M., Rep. Babb. Exped. 14 (1858)	...	...	...	...	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.III,617
H. Baxteri; Cunningham in De Candolle, prodr. VI, 193 (1837)	...	...	...	...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,617
H. Ayersii; F. v. M., fragm. VIII, 167 (1874)	...	...	...	...	—	S.A.	—	—	—	—	—	— M.fr.IX,195.
H. rutidolepis; De Candolle, prodr. VI, 194 (1837)	...	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,618
H. scorpioides; Labillardière, Nov. Holl. pl. spec. II, 45, t. 191 (1806)	...	...	...	...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,618 M.fr.VIII,167.

H. oligochsetum; F. v. M., fragm. VI, 235 (1868) ...	...	W.A.	—	—	—	—	N.A.	—	M.fr. VI, 235.
H. Tepperi; F. v. M. in Wing's South. Science Record II, 1 (1882) ...	...	—	S.A.	—	—	—	—	—	
H. obtusifolium; Sonder & F. v. M. in Linnaea XXV, 515 (1852) ...	...	W.A.	S.A.	—	V.	N.S.W.	—	B.fl. III, 619	
H. Calvertianum; F. v. M., fragm. X, 108 (1877) ...	...	—	—	—	—	N.S.W.	—	—	M.fr. XI, 136.
H. Spicerii; F. v. M., fragm. XI, 47 (1878) ...	...	—	—	T.	—	—	—	—	M.fr. XI, 47.
H. dealbatum; Labillardière, Nov. Holl. pl. spec. II, 45, t. 190 (1806) ...	...	—	—	T.	V.	—	—	B.fl. III, 619	
H. pumilum; J. Hooker, Fl. Tasman. I, 213, t. 60 (1860) ...	...	—	—	T.	—	—	—	B.fl. III, 619	
H. lucidum; Henckel, adumbr. pl. hort. Hal. 5 (1806) ...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl. III, 620 M.fr. XI, 48.
H. elatum; Cunningham in De Candolle, prodr. VI, 193 (1837) ...	...	—	—	—	V.	N.S.W.	Q.	—	B.fl. III, 621
H. adenophorum; F. v. M. in Transact. Vict. Inst. 38 (1855) ...	...	—	S.A.	—	V.	—	—	—	B.fl. III, 622
H. leucopsidium; De Candolle, prodr. VI, 193 (1837) ...	...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl. III, 622 M.fr. VI, 255.
H. Blandowskianum; Steetz in Schlecht. Linnaea XXV, 512 (1852) ...	...	—	S.A.	—	V.	N.S.W.	—	—	B.fl. III, 622
H. oxylepis; F. v. M., fragm. I, 35 (1858) ...	...	—	—	—	—	N.S.W.	Q.	—	B.fl. III, 623 M.fr. I, 35.
H. collinum; De Candolle, prodr. VI, 190 (1837) ...	...	—	—	—	—	N.S.W.	Q.	—	B.fl. III, 623
H. rupicola; De Candolle, prodr. VI, 190 (1837) ...	...	—	—	—	—	—	Q.	—	B.fl. III, 623
H. podolepideum; F. v. M., Rep. Babb. Exped. 13 (1858) ...	...	—	S.A.	—	—	N.S.W.	Q.	—	B.fl. III, 624 M.fr. III, 134.
H. Gilesii; F. v. M., fragm. X, 85 (1876) ...	...	W.A.	—	—	—	—	N.A.	—	M.fr. X, 86.
H. ambiguum; Turczaninow in Bull. de Mosc. 195 (1851) ...	...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl. III, 609
H. apiculatum; De Candolle, prodr. VI, 195 (1837) ...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl. III, 624
H. senipapposum; De Candolle, prodr. VI, 195 (1837) ...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl. III, 625
H. Dockerii; F. v. M. in Benth. Fl. Austr. III, 626 (1866) ...	...	—	S.A.	—	—	N.S.W.	—	—	B.fl. III, 626 M.fr. II, 156.
H. Agryroglottis; Benth. Fl. Austr. III, 626 (1866) ...	...	W.A.	—	—	—	—	—	—	B.fl. III, 626
H. ramosum; De Candolle, prodr. VI, 181 (1837) ...	...	W.A.	—	—	—	—	—	—	B.fl. III, 626
H. cordatum; De Candolle, prodr. VI, 180 (1837) ...	...	W.A.	—	—	—	—	—	—	B.fl. III, 627
H. obovatum; De Candolle, prodr. VI, 180 (1837) ...	...	—	—	—	—	N.S.W.	—	—	B.fl. III, 627 M.fr. II, 89.
H. Bidwillii; Benth. Fl. Austr. III, 627 (1866) ...	...	—	—	—	—	N.S.W.	Q.	—	B.fl. III, 627
H. Becklerii; F. v. M. in Benth. Fl. Austr. III, 627 (1866) ...	...	—	—	—	—	N.S.W.	Q.	—	B.fl. III, 627 M.fr. I, 183, 244.
H. diotophyllum; F. v. M., fragm. V, 150 (1866) ...	...	—	—	—	—	N.S.W.	Q.	—	B.fl. III, 628 M.fr. VIII, 46.
H. Thomsoni; F. v. M., fragm. VIII, 45 (1873) ...	...	—	S.A.	—	—	—	—	—	M.fr. VIII, 45.
H. diosmifolium; Don in Steud. Nom. Bot. ed. sec. 738 (1841) ...	...	—	—	—	—	N.S.W.	Q.	—	B.fl. III, 628
H. retusum; F. v. M., fragm. VIII, 46 (1873) ...	...	—	S.A.	—	V.	N.S.W.	—	—	B.fl. III, 628 M.fr. VIII, 46.
H. decurrens; F. v. M., fragm. VIII, 46 (1873) ...	...	—	S.A.	—	V.	N.S.W.	—	—	M.fr. VIII, 46.
H. Cunninghamii; Benth. Fl. Austr. III, 629 (1866) ...	...	—	—	—	—	N.S.W.	—	—	B.fl. III, 629
H. reticulatum; Lessing in Steud. Nom. Bot. ed. sec. 740 (1841) ...	...	—	—	T.	—	—	—	—	B.fl. III, 629
H. cinereum; F. v. M. in Benth. Fl. Austr. III, 629 (1866) ...	...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl. III, 629 M.fr. V, 199.
H. hracteolatum; Benth. Fl. Austr. III, 630 (1866) ...	...	—	—	T.	—	—	—	—	B.fl. III, 630
H. Kempei; F. v. M. in Melb. Chemist, Jan. (1882) ...	...	—	S.A.	—	—	—	—	—	—
H. cassinioides; Benth. Fl. Austr. III, 630 (1866) ...	...	—	—	—	—	—	Q.	—	B.fl. III, 630
H. Gunnii; F. v. M. in Benth. Fl. Austr. III, 630 (1866) ...	...	—	—	T.	—	—	—	—	B.fl. III, 630 M.fr. V, 199.
H. rosmarinifolium; Lessing, synops. gen. comp. 274 (1832) ...	...	—	—	T.	V.	N.S.W.	—	—	B.fl. III, 631
H. ledifolium; F. v. M. in Benth. Fl. Austr. III, 630 (1866) ...	...	—	—	T.	—	—	—	—	B.fl. III, 630 M.fr. V, 199.
H. ferrugineum; Lessing, synops. gen. comp. 274 (1832) ...	...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl. III, 631
H. antennarium; F. v. M. in Benth. Fl. Austr. III, 632 (1866) ...	...	—	—	T.	—	—	—	—	B.fl. III, 632 M.fr. V, 199.
H. obo cordatum; F. v. M. in Benth. Fl. Austr. III, 632 (1866) ...	...	—	—	T.	V.	N.S.W.	—	—	B.fl. III, 632 M.fr. V, 199.
H. Backhausii; F. v. M. in Benth. Fl. Austr. III, 632 (1866) ...	...	—	—	T.	V.	—	—	—	B.fl. III, 632 M.fr. V, 199.
H. cuneifolium; F. v. M. in Benth. Fl. Austr. III, 633 (1866) ...	...	—	—	—	V.	N.S.W.	—	—	B.fl. III, 633
H. baccharoides; F. v. M. in Benth. Fl. Austr. III, 633 (1866) ...	...	—	—	T.	V.	N.S.W.	—	—	B.fl. III, 633 M.fr. V, 199.
H. lepidophyllum; F. v. M. in Benth. Fl. Austr. III, 633 (1866) ...	...	W.A.	—	—	—	—	—	—	B.fl. III, 633 M.fr. V, 199.
H. scutellifolium; Benth. Fl. Austr. III, 633 (1866) ...	...	—	—	T.	—	—	—	—	B.fl. III, 633 M.fr. V, 199.
H. lycopodioides; Benth. Fl. Austr. III, 634 (1866) ...	...	—	—	T.	—	—	—	—	B.fl. III, 634 M.fr. V, 199.
H. selaginoides; F. v. M. in Benth. Fl. Austr. III, 634 (1866) ...	...	—	—	T.	—	—	—	—	B.fl. III, 634 M.fr. V, 199.

CASSINIA; R. Brown in Transact. Linn. Soc. XII, 126 (1817). (Non zoologorum, Apolochlamys, Achromolaena, Chromochiton.)

C. leptcephala; F. v. M., fragm. III, 138 (1863)	—	—	—	—	N.S.W.	—	—	B.fl.III,585	M.fr.III,138.
C. compacta; F. v. M., fragm. I, 18 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,585	M.fr.I,18.
C. denticulata; R. Brown in Transact. Linn. Soc. XII, 127 (1817)	—	—	—	—	N.S.W.	—	—	B.fl.III,586	
C. longifolia; R. Brown in Transact. Linn. Soc. XII, 127 (1817)	—	—	—	T.	V.	N.S.W.	Q.	—	B.fl.III,586
C. aurea; R. Brown in Transact. Linn. Soc. XII, 127 (1817)	—	—	—	—	—	N.S.W.	—	—	B.fl.III,586
C. aculeata; R. Brown in Transact. Linn. Soc. XII, 127	—	—	—	S.A.	T.	V.	N.S.W.	—	B.fl.III,586
C. laevis; R. Brown in Transact. Linn. Soc. XII, 128 (1817)	—	—	—	S.A.	—	—	N.S.W.	Q.	—
C. tenuifolia; Bentham, Fl. Austr. III, 585 (1866)	—	—	—	—	—	N.S.W.	—	—	B.fl.III,585
C. quinquefaria; R. Brown in Transact. Linn. Soc. XII, 128 (1817)	—	—	—	—	—	V.	N.S.W.	Q.	—
C. arcuata; R. Brown in Transact. Linn. Soc. XII, 128 (1817)	—	—	—	W.A.	S.A.	—	V.	N.S.W.	—
C. subtropica; F. v. M., fragm. I, 17 (1858)	—	—	—	—	—	—	N.S.W.	Q.	—
C. Theodori; F. v. M., fragm. V, 148 (1866)	—	—	—	—	—	—	N.S.W.	—	—
C. spectabilis; R. Brown in Transact. Linn. Soc. XII, 128 (1817)	—	—	—	S.A.	T.	V.	—	—	—

HUMEA; Smith, Exot. bot. I, t. 1 (1804). (Haeckeria, Calomeria.)

H. elegans; Smith, Exot. bot. I, t. 1 (1804)	—	—	—	V.	N.S.W.	—	—	B.fl.III,589	M.fr.I,17.
H. cassiniacea; F. v. M., fragm. I, 17 (1858)... ..	—	—	—	S.A.	—	—	—	—	B.fl.III,589
H. ozothamnoides; F. v. M., fragm. I, 17 (1858)	—	—	—	V.	N.S.W.	—	—	B.fl.III,590	M.fr.VIII,142.
H. squamata; F. v. M., fragm. XI, 86 (1880)	—	—	—	S.A.	—	V.	N.S.W.	—	—

ACOMIS; F. v. M., fragm. II, 89 (1860).

A. Rutidosis; F. v. M., fragm. II, 89 (1860)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,591	M.fr.II,89.
A. macra; F. v. M., fragm. IV, 145 (1864)	—	—	—	—	—	—	Q.	—	B.fl.III,591

RUTIDOSIS; De Candolle, prodr. VI, 158 (1837). (Pumilo, Actinopappus, Lepidocoma.)

R. leioplepis; F. v. M. in Transact. Vict. Inst. 131 (1855)	—	—	—	V.	—	—	—	B.fl.III,593	M.fr.I,34.
R. leptorrhynchoides; F. v. M., fragm. V, 148 (1866)	—	—	—	V.	N.S.W.	—	—	B.fl.III,593	M.fr.V,148.
R. helichrysoides; De Candolle, prodr. VI, 159 (1837)	—	—	—	S.A.	—	V.	N.S.W.	Q.	N.A.
R. leucantha; F. v. M., fragm. I, 35 (1858)	—	—	—	—	—	—	—	Q.	—
R. Murchisonii; F. v. M., fragm. I, 34 (1858)	—	—	—	—	—	—	—	Q.	—
R. Pumilo; Bentham, Fl. Austr. III, 595 (1866)	—	—	—	W.A.	S.A.	T.	V.	N.S.W.	—

PITHOCARPA; Lindley, Bot. Regist. XXV, App. 23 (1839).

P. corymbosa; Lindley, Bot. Regist. XXV, App. 23 (1839)	—	—	—	W.A.	—	—	—	—	B.fl.III,590
--	---	---	---	------	---	---	---	---	--------------

AMMOBIUM; R. Brown in Bot. Magaz. t. 2459 (1824).

A. alatum; R. Brown in Bot. Magaz. t. 2459 (1824)... ..	—	—	—	V.	N.S.W.	Q.	—	B.fl.III,583	
A. craspedioides; Bentham, Fl. Austr. III, 584 (1866)	—	—	—	—	N.S.W.	—	—	B.fl.III,584	

IXODIA; R. Brown in Ait. hort. Kew. sec. ed. IV, 17 (1812).

I. achilleoides; R. Brown in Ait. hort. Kew. sec. ed. IV, 17 (1812)	—	—	—	S.A.	—	V.	—	—	B.fl.III,583
--	---	---	---	------	---	----	---	---	--------------

MILLOTIA; Cassini in Ann. sc. nat. XVII, 416 (1829).

M. tenuifolia; Cassini in Ann. sc. nat. XVII, 416 (1829)	—	—	—	W.A.	S.A.	T.	V.	N.S.W.	—
M. Greevesii; F. v. M., fragm. III, 18, t. 19 (1863)	—	—	—	S.A.	—	—	—	N.S.W.	Q.
M. Kempei; F. v. M. in Wing's South. Sc. Record II, 1 (1882)	—	—	—	S.A.	—	—	—	—	—

TOXANTHUS; Turczaninow in Bull. de Mosc. XXIV, 177 (1851). (Anthocerastes, Scyphocoronis.)

T. perpusillus; Turczaninow in Bull. Mosc. XXIV, 177 (1851)	—	—	—	W.A.	S.A.	—	V.	N.S.W.	—
T. Muelleri; Bentham, Fl. Austr. III, 592 (1866)	—	—	—	W.A.	S.A.	—	V.	—	—
T. major; Turczaninow in Bull. de Mosc. XXIV, 64 (1851)	—	—	—	W.A.	—	—	—	—	—

QUINETIA; Cassini in Diction. sc. nat. LX, 579 (1830).

Q. Urvillei; Cassini in Diction. LX, 579 (1830) ... W.A. S.A. — — — — B.fl.III,595

PHACELOTHRIX; F. v. M., fragm. XI, 49 (1878).

P. cladochaeta; F. v. M., fragm. XI, 49 (1878) ... Q. N.A. B.fl.III,594 M.fr.V,199.

DECAZESIA; F. v. M., fragm. XI, 71 (1879).

D. hecatocephala; F. v. M., fragm. XI, 72 (1879) ... W.A. — — — — N.A. — M.fr.XI,72.

MYRIOCEPHALUS; Bentham in Hueg. enum. 61 (1837). (Hyalolepis, Anthediosorus, Gilberta, Lamprochlaena, Elachopappus, Polycalymma; some readily to be restituted.)

M. rhizocephalus; Bentham, Fl. Austr. III, 557 (1866) ... W.A. S.A. — V. N.S.W. Q. — B.fl.III,557 M.fr.III,155.
 M. nudus; A. Gray in Hook. Kew Misc. III, 174 (1851) ... W.A. — — — — — B.fl.III,558 M.fr.III,157.
 M. appendiculatus; Bentham in Hueg. enum. 61 (1837) ... W.A. — — — — — B.fl.III,558
 M. Rudallii; F. v. M., fragm. III, 157 (1863) ... W.A. S.A. — — — — N.A. B.fl.III,559 M.fr.VI,235; VII,145.
 M. gracilis; Bentham, Fl. Austr. III, 559 (1866) ... W.A. — — — — — B.fl.III,559 M.fr.VIII,169.
 M. helichrysoides; A. Gray in Hook. Kew Misc. III, 175 (1851) ... W.A. — — — — — B.fl.III,559 M.fr.III,159.
 M. suffruticosus; Bentham, Fl. Austr. III, 559 (1866) ... W.A. — — — — — B.fl.III,559
 M. Stuartii; Bentham, Fl. Austr. III, 560 (1866) ... — S.A. — V. N.S.W. Q. N.A. B.fl.III,560
 M. Guerinae; F. v. M., fragm. VIII, 169 (1874) ... W.A. — — — — — M.fr.VIII,169.

ANGIANTHUS; Wendland, Collect. pl. II, 31 (1809). (Siloxerus, Styloncerus, Ogcerostylus, Cylindrosorus, Phyllocalymma, Skirrophorus, Chrysocoryne, Eriocladium, Pogonolepis, Piptostemma, Epitriche, Gamozygis, Cephalosorus partly, Hyalochlamys, Dithyrostegia, Pleuropappus; some readily to be restituted.)

A. humifusus; Bentham, Fl. Austr. III, 563 (1866) ... W.A. — — — — — B.fl.III,563
 A. tomentosus; Wendland, Collect. pl. II, 31, t. 48 (1809) ... W.A. S.A. — V. N.S.W. — N.A. B.fl.III,562
 A. pleuropappus; Bentham, Fl. Austr. III, 563 (1866) ... — S.A. — — — — B.fl.III,563
 A. brachypappus; F. v. M. in Transact. phil. Soc. Vict. I, 44 (1854) ... — S.A. — V. N.S.W. Q. — B.fl.III,563
 A. myosuroides; Bentham, Fl. Austr. III, 563 (1866) ... W.A. — — — — — B.fl.III,563
 A. tenellus; Bentham, Fl. Austr. III, 564 (1866) ... W.A. S.A. — — — — B.fl.III,564
 A. pusillus; Bentham, Fl. Austr. III, 564 (1866) ... W.A. S.A. — V. N.S.W. Q. — B.fl.III,564
 A. Milnei; Bentham, Fl. Austr. III, 564 (1866) ... W.A. — — — — — B.fl.III,564
 A. Cunninghamii; Bentham, Fl. Austr. III, 565 (1866) ... W.A. — — — — — B.fl.III,565
 A. phyllocephalus; Bentham, Fl. Austr. III, 565 (1866) ... W.A. — — — — — B.fl.III,565 M.fr.III,159.
 A. micropoides; Bentham, Fl. Austr. III, 565 (1866) ... W.A. — — — — — B.fl.III,565
 A. microcephalus; Bentham, Fl. Austr. III, 566 (1866) ... W.A. — — — — — B.fl.III,566 M.fr.III,158.
 A. platycephalus; Bentham, Fl. Austr. III, 566 (1866) ... W.A. — — — — — B.fl.III,566
 A. Drummondii; Bentham, Fl. Austr. III, 566 (1866) ... W.A. — — — — — B.fl.III,566
 A. Preissianus; Bentham, Fl. Austr. III, 566 (1866) ... W.A. S.A. T. V. N.S.W. — — B.fl.III,566
 A. pygmaeus; Bentham, Fl. Austr. III, 567 (1863) ... W.A. — — — — — B.fl.III,567
 A. globifer; Bentham, Fl. Austr. 567 (1866) .. W.A. — — — — — B.fl.III,567
 A. demissus; Bentham, Fl. Austr. III, 567 (1866) ... W.A. — — — — — B.fl.III,567
 A. strictus; Bentham, Fl. Austr. III, 568 (1866) ... W.A. S.A. — V. N.S.W. — — B.fl.III,568
 A. plumiger; Bentham, Fl. Austr. III, 568 (1866) ... W.A. — — — — — B.fl.III,568
 A. amplexicaulis; Bentham, Fl. Austr. III, 568 (1866) ... W.A. — — — — — B.fl.III,568 M.fr.III,159.

GNEPHOSIS; Cassini in Bull. de la Soc. philom. 45 (1820). (Cephalosorus partly, Nematopus, Crossolepis, Leptotriche, Trichanthodium, Cyathopappus; some readily to be restituted.)

G. Burkittii; Bentham, Fl. Austr. III, 570 (1866) ... — S.A. — — N.S.W. — — B.fl.III,570
 G. eriocarpa; Bentham, Fl. Austr. III, 570 (1866) ... — S.A. — — N.S.W. Q. — B.fl.III,570 M.fr.III,156.
 G. macrocephala; Turczaninow in Bull. Soc. Mosc. XXIV, 190 (1851) ... W.A. — — — — — B.fl.III,570 M.fr.II,158.
 G. skirphora; Bentham, Fl. Austr. III, 570 (1866) ... W.A. S.A. — V. N.S.W. — — B.fl.III,570

G. cyathopappa ; Bentham, Fl. Austr. III, 571 (1866)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.III,571	M.fr.II,158.
G. codonopappa ; F. v. M. in Giles, geogr. trav. 217 (1875)	—	S.A.	—	—	—	—	—	—	M.fr.IX,2.
G. leptoclada ; Bentham, Fl. Austr. III, 571 (1866)... ..	...	W.A.	—	—	—	—	—	B.fl.III,571	M.fr.III,158.
G. arachnoidea ; Turczaninow in Bull. Soc. Mosc. XXIV, 189 (1851)	...	W.A.	S.A.	—	—	N.S.W.	Q.	—	B.fl.III,571 M.fr.II,158.
G. tenuissima ; Cassini, in Bull. de la Soc. philom. 43 (1820)	...	W.A.	—	—	—	—	—	B.fl.III,572	
G. acicularis ; Bentham, Fl. Austr. III, 572 (1866)	...	W.A.	—	—	—	—	—	B.fl.III,572	
G. pygmaea ; Bentham, Fl. Austr. III, 572 (1866)	...	W.A.	—	—	—	—	—	B.fl.III,572	
G. brevifolia ; Bentham, Fl. Austr. III, 572 (1866)	...	W.A.	—	—	—	—	—	B.fl.III,572	
G. eriocephala ; Bentham, Fl. Austr. III, 573 (1866)	...	W.A.	—	—	—	—	—	B.fl.III,573	

ERIOCHLAMYS ; Sonder & F. v. M. in Schlecht. Linnaea XXV, 488 (1852).

E. Behrii ; Sonder & F. v. M. in Linnaea XXV, 488 (1852)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,591	
E. Knappii ; F. v. M. in Melb. Chem. VI, 61, p. 4 (1883)	...	W.A.	S.A.	—	—	—	—	N.A.	—

CALOCEPHALUS ; R. Brown in Transact. Linn. Soc. XII, 106 (1817). (Leucophyta, Pachysura, Blennospora, Achrysum.)

C. Drummondii ; Bentham, Fl. Austr. III, 574 (1866)	...	W.A.	S.A.	—	—	—	—	B.fl.III,574	
C. Brownii ; F. v. M., Rep. Babb. Exped. 13 (1858)	...	W.A.	S.A.	T.	V.	N.S.W.	—	B.fl.III,574	
C. Sonderi ; F. v. M., sec. gen. Report 12 (1854)	...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,775
C. lacteus ; Lessing, syn. gen. comp. 271 (1832)	...	W.A.	S.A.	T.	V.	N.S.W.	—	B.fl.III,575	
C. citreus ; Lessing, syn. gen. comp. 271 (1832)	...	—	S.A.	T.	V.	N.S.W.	—	B.fl.III,575	
C. angianthoides ; Bentham, Fl. Austr. III, 575 (1866)	...	W.A.	—	T.	—	—	—	B.fl.III,575	
C. Francisii ; Bentham, Fl. Austr. III, 576 (1866)	...	W.A.	—	—	—	—	—	B.fl.III,576	M.fr.III,155.
C. Dittrichii ; F. v. M. in Uhlworm's bot. Centralbl. XXVII, 300 (1886)	...	—	S.A.	—	—	N.S.W.	Q.	N.A.	—
C. platycephalus ; Bentham, Fl. Austr. III, 576 (1866)	...	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.III,576 M.fr.III,154.
C. multiflorus ; Bentham, Fl. Austr. III, 576 (1866)	...	W.A.	—	—	—	—	—	N.A.	B.fl.III,576
C. aeruoides ; Bentham, Fl. Austr. III, 576 (1866)	...	W.A.	—	—	—	—	—	—	B.fl.III,576 M.fr.III,154.

CEPHALIPTERUM ; A. Gray in Hook. Kew Misc. IV, 271 (1852).

C. Drummondii ; A. Gray in Hook. Kew Misc. IV, 272 (1852)	...	W.A.	S.A.	—	—	—	—	—	B.fl.III,577
--	-----	------	------	---	---	---	---	---	--------------

GNAPHALODES ; A. Gray in Hook. Kew Misc. IV, 228 (1852).

G. uliginosum ; A. Gray in Hook. Kew Misc. IV, 228 (1852)	...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,578 M.fr.XI,27.
G. condensatum ; A. Gray in Hook. Kew Misc. IV, 228 (1852)	...	W.A.	—	—	—	—	—	—	B.fl.III,578
G. filifolium ; Bentham, Fl. Austr. III, 578 (1866)	...	W.A.	—	—	—	—	—	—	B.fl.III,578

CRASPEDIA ; G. Forster, florul. insul. Austr. prodr. 306 (1786). (Richea, Pycnosorus.)

C. Richea ; Cassini, in Diction. XI, 353 (1818)	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,579 N.Z.
C. pleiocephala ; F. v. M. in Schlecht. Linnaea XXV, 404 (1852)... ..	...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,580 M.fr.XI,27.
C. chrysantha ; Bentham, Fl. Austr. III, 580 (1866)	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,580
C. globosa ; Bentham, Fl. Austr. III, 580 (1866)	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,580

CHTHONOCEPHALUS ; Steetz in Lehm. pl. Preiss. I, 444 (1845). (Chamaesphaerion, Gyrostephium, Lachnothalamus.)

C. tomentellus ; Bentham, Fl. Austr. III, 581 (1866)	...	W.A.	—	—	—	—	—	—	B.fl.III,581 M.fr.III,156; IV,176.
C. Pseudevax ; Steetz in Lehm. pl. Preiss. I, 445 (1845)	...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,582
C. pygmaeus ; Bentham, Fl. Austr. III, 582 (1866)	...	W.A.	—	—	—	—	—	—	B.fl.III,582

NABLONIUM ; Cassini in Diction. XXXIV, 101 (1825).

N. calyceroides ; Cassini, Dict. XXXIV, 101 (1825)... ..	...	—	—	T.	—	—	—	—	B.fl.III,545
--	-----	---	---	----	---	---	---	---	--------------

CHRYSOGONUM ; Linné, hort. Cliffort. 424 (1737). (Moonia, Pentalepis).

C. trichodesmoides ; Bentham & J. Hooker, gen. pl. II, 350 (1873)	...	—	—	—	—	—	—	N.A.	B.fl.III,540 M.fr.VIII,145.
C. eclipoides ; Bentham & J. Hooker, gen. pl. II, 350 (1873)	...	—	—	—	—	—	—	N.A.	B.fl.III,540 M.fr.VIII,145.
C. procumbens ; Bentham & J. Hooker, gen. pl. II, 350 (1873)	...	—	—	—	—	—	—	N.A.	B.fl.III,540 M.fr.VIII,145.

SIEGESBECKIA ; Linné, hort. Cliffort. 412 (1737).

S. orientalis ; Linné, spec. plant. 900 (1753) W.A. S.A. — V. N.S.W. Q. — B.f.III,535 As., Afr., Am., P.

ENHYDRA ; Loureiro, Flor. Cochinch. II, 510 (1790). (Euydra.)

E. fluctuans ; Loureiro, fl. Cochinch. II, 511 (1799) — — — — N.S.W. Q. — B.f.III,546 M.fr.III,139 As., Afr., Am.

ECLIPTA ; Linné, mantissa altera, 157 (1771). (Blainvillea.)

E. alba ; Hasskarl, plant. Javan. rar. 528 (1848) — — — — N.S.W. Q. N.A. B.f.III,536 As., Afr., Am.

E. latifolia ; Linné, fil. suppl. 378 (1781) — — — — — Q. N.A. — M.fr.VIII,145 As., Afr., Am.

E. platyglossa ; F. v. M., fragm. II, 135 (1861) — S.A. — V. N.S.W. Q. N.A. B.f.III,536 M.fr.XI,27.

WEDELIA ; Jacquin, stirp. Amer. hist. 217, t. 130 (1763). (Wollastonia.)

W. calendulacea ; Lessing, synops. gen. comp. 222 (1832) — — — — — Q. — B.f.III,537 As.

W. urticifolia ; De Candolle in Wight, Contrib. Bot. Ind. 18 (1834) — — — — — Q. N.A. B.f.III,538 As.

W. spilanthis ; F. v. M., fragm. V, 64 (1865) — — — — N.S.W. Q. — B.f.III,538 M.fr.V,64.

W. verbesinoides ; F. v. M. in Benth. Fl. Austr. III, 538 (1866) — S.A. — — — N.A. B.f.III,538

W. biflora ; De Candolle in Wight, Contrib. Bot. Ind. 18 (1834) — — — — N.S.W. Q. N.A. B.f.III,539 As., Afr., P.

W. asperima ; Benth, Fl. Austr. III, 539 (1866) — — — — — Q. N.A. B.f.III,539

SPILANTHES ; Jacquin, stirp. Amer. hist. 214, t. 126 (1763).

S. grandiflora ; Turczaninow in Bull. de Mosc. XXIV, 185 (1851) — — — — N.S.W. Q. N.A. B.f.III,541 M.fr.V,63.

S. anactina ; F. v. M., fragm. V, 63 (1865) — — — — — Q. N.A. B.f.III,541 M.fr.V,63 As.

BIDENS ; Dillenius, hort. Eltham. 51 et 52, t. 43 et 44 (1732), from Cesalpini (1583). (Possibly immigrated.)

B. tripartitus ; Linné, spec. plant. 831 (1753) — — — — V. N.S.W. Q. — B.f.III,543 E., As.

B. pilosus ; Linné, spec. plant. 832 (1753) — — — — N.S.W. Q. — B.f.III,543 As., Afr., Am., P.

B. bipinnatus ; Linné, spec. plant. 832 (1753) — S.A. — — N.S.W. Q. N.A. B.f.III,543 As., Afr., Am.

GLOSSOGYNE ; Cassini in Diction. LI, 475 (1827). (Diodontium.)

G. tenuifolia ; Cassini in Dict. LI, 475 (1827) — S.A. — V. N.S.W. Q. N.A. B.f.III,544 M.fr.I,51 As., P.

G. retroflexa ; F. v. M., fragm. I, 51 (1858) — — — — — Q. — B.f.III,544

G. filifolia ; F. v. M. in Benth. Fl. Austr. III, 544 (1866) — — — — — N.A. B.f.III,544

FLAVERIA ; A. L. de Jussieu, gen. plant. 186 (1789).

F. Australasica ; Hooker in Mitch. Trop. Austr. 118 (1848) W.A. S.A. — — N.S.W. Q. N.A. B.f.III,546 M.fr.I,183.

SOLIVA ; Ruiz et Pavon, prodr. 113, t. 24 (1794). (Gymnostyles ; possibly immigrated.)

S. anthemifolia ; R. Brown in Transact. Linn. Soc. XII, 102 (1817) — — — — N.S.W. Q. — B.f.III,552 Am.

COTULA ; Linné, syst. nat. 9 (1735) ; Linné, gen. pl. 256 (1737). (Gymnogyne, Strongylospermum, Pleiogyne, Symphyomera, Ctenosperma, Leptinella.)

C. filifolia ; Thunberg, prodr. pl. Capens. 161 (1800) W.A. S.A. T. V. N.S.W. — — B.f.III,548 M.fr.VI,114 Afr.

C. coronopifolia ; Linné, sp. pl. 892 (1753) W.A. S.A. T. V. N.S.W. — — B.f.III,549 M.fr.VI,114 E., Afr., Am., N.Z.

C. Gymnogyne ; F. v. M. in Benth. Fl. Austr. III, 549 (1866) W.A. — — — — — B.f.III,549

C. Australis ; J. Hooker, Fl. N. Zeal. I, 128 (1853) W.A. S.A. T. V. N.S.W. Q. — B.f.III,550 N.Z.

C. alpina ; J. Hooker, Fl. Tasman. I, 192, t. 51 (1860) — — T. V. N.S.W. Q. — B.f.III,550

C. Drummondii ; Benth, Fl. Austr. III, 550 (1866) W.A. — — — — — B.f.III,550

C. reptans ; Benth, Fl. Austr. III, 551 (1866) — S.A. T. V. N.S.W. — — B.f.III,551

C. filicula ; J. Hooker in Benth. Fl. Austr. III, 551 (1866) — — T. V. N.S.W. — — B.f.III,551

CENTIPEDA ; Loureiro, Fl. Cochinch. II, 492 (1790). (Myriogyne, Sphaeromorphaea partly.)

C. racemosa ; F. v. M., Cens. 84 (1882) from Hook. (1848) — — — — N.S.W. Q. N.A. B.f.III,553

C. orbicularis ; Loureiro, Fl. Cochinch. II, 492 (1790) W.A. S.A. T. V. N.S.W. Q. N.A. B.f.III,553 M.fr.VIII,142 As., Am., P., N.Z.

C. Cunninghamii ; F. v. M., fragm. VIII, 143 (1874) W.A. S.A. — V. N.S.W. Q. N.A. —

C. thespidioides ; F. v. M., fragm. VIII, 143 (1874) — S.A. — V. N.S.W. Q. — —

ABROTANELLA; Cassini in Dict. XXXVI, 27 (1825). (Scleroleima, Trineuron.)

A. fosterioides; J. Hooker, Hand. N. Zeal. Fl. 139 (1864)...	...	...	...	—	—	T.	—	—	—	—	B.fl.III,554
A. nivigena; F. v. M. in Benth. Fl. Austr. III, 554 (1866)...	...	...	...	—	—	V.	—	—	—	—	B.fl.III,554
A. scapigera; F. v. M. in Benth. Fl. Austr. III, 554 (1866)	...	...	...	—	—	T.	—	—	—	—	B.fl.III,554

ELACHANTHUS; F. v. M. in Schlecht. Linnaea XXV, 410 (1852).

E. pusillus; F. v. M. in Schlecht. Linnaea XXV, 411 (1852)	...	...	...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,555
--	-----	-----	-----	---	------	---	----	--------	---	---	--------------

DIMORPHOCOMA; F. v. M. & Tate in Trans. Roy. Soc. of S. Austr. VI, 107 (1883).

D. minutula; F. v. M. & Tate in Transact. Roy. Soc. of S.A. VI, 107 (1883)	...	—	S.A.	—	—	—	—	—	—	—	—
--	-----	---	------	---	---	---	---	---	---	---	---

CERATOGYNE; Turczaninow in Bull. de Mosc. XXIV, 68 (1851). (Diotosperma.)

C. obionoides; Turczaninow in Bull. de Mosc. XXIV, 68 (1851)	...	...	...	W.A.	S.A.	—	—	N.S.W.	Q.	—	B.fl.III,555 M.fr.III,159.
--	-----	-----	-----	------	------	---	---	--------	----	---	----------------------------

ISOETOPSIS; Turczaninow in Bull. de Mosc. XXIV, 175 (1851).

I. graminifolia; Turczaninow in Bull. de Mosc. XXIV, 175, t. 3 (1851)	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,556
---	-----	-----	-----	------	------	---	----	--------	----	---	--------------

EMILIA; Cassini in Bulletin de la Soc. philomat. 68 (1817).

E. purpurea; Cassini in Diction. des sc. nat. XXXIV, 393 (1825)	...	...	—	—	—	—	—	Q.	—	—	M.fr.XII ...	...	As., Afr.
---	-----	-----	---	---	---	---	---	----	---	---	--------------	-----	-----------

GYNURA; Cassini in Diction. des sc. nat. XXXIV, 391 (1825).

G. Pseudo-China; De Candolle, prodr. VI, 299 (1837)	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.III,661	...	...	...	As.
---	-----	-----	-----	---	---	---	---	--------	----	---	--------------	-----	-----	-----	-----

SENECIO; Tournefort, inst. 456, t. 260 (1700), from Plinius. (Bedfordia, Centropappus.)

S. Gregorii; F. v. M., pl. of Greg. Search. Exped. for Leichh. 7 (1858)	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.III,663					
S. platylepis; De Candolle, prodr. VI, 371 (1837)	...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,664	M.fr.XI,27.				
S. papillosus; F. v. M. in Transact. phil. Inst. Vict. II, 69 (1857)	...	...	—	—	T.	—	—	—	—	B.fl.III,664					
S. primulifolius; F. v. M. in Transact. phil. Inst. Vict. II, 69 (1857)	...	...	—	—	T.	—	—	—	—	B.fl.III,664					
S. pectinatus; De Candolle, prodr. VI, 372 (1837)	...	...	—	—	T.	V.	N.S.W.	—	—	B.fl.III,664					
S. spathulatus; A. Richard, sert. Astrol. 125 (1833)	...	...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,665					
S. megaglossus; F. v. M. in Linnaea XXV, 419 (1852)	...	...	—	S.A.	—	—	—	—	—	B.fl.III,665	M.fr.VI,255.				
S. magnificus; F. v. M. in Linnaea XXV, 418 (1852)	...	...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,665					
S. insularis; Benth. Fl. Austr. III, 666 (1866)	...	...	—	—	—	—	N.S.W.	—	—	B.fl.III,666					
S. Centropappus; F. v. M., annual report 26 (1858)	...	...	—	—	T.	—	—	—	—	B.fl.III,666					
S. macranthus; A. Richard, sert. Astrol. 126 (1833)	...	...	—	—	—	—	N.S.W.	—	—	B.fl.III,666					
S. Daltoni; F. v. M., fragm. VI, 27 (1867)	...	...	—	—	—	—	N.S.W.	Q.	—	—	M.fr.VI,27.				
S. lautus; Solander in G. Forst. prodr. 91 (1786)	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.III,667	M.fr.XI,20	...	N.Z.		
S. capillifolius; J. Hooker in Lond. Journ. VI, 123 (1847)...	...	...	—	—	T.	—	—	—	—	B.fl.III,667					
S. vagus; F. v. M. in Transact. phil. Soc. Vict. I, 46 (1854)	...	...	—	—	—	V.	N.S.W.	—	—	B.fl.III,667					
S. amygdalifolius; F. v. M., fragm. I, 232 (1859)	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.III,668	M.fr.I,232.				
S. velleioides; Cunningham in De Candolle, prodr. VI, 374 (1837)	...	...	—	—	T.	V.	N.S.W.	Q.	—	B.fl.III,668					
S. dryadens; Sieber in Spreng. syst. veg. III, 562 (1826)	...	...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,668					
S. Behrianus; Sonder & F. v. M. in Linnaea XXV, 527 (1852)	...	...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,669					
S. leucoglossus; F. v. M., fragm. II, 15 (1861)	...	...	W.A.	—	—	—	—	—	—	B.fl.III,669	M.fr.II,15.				
S. brachyglossus; F. v. M. in Schlecht. Linnaea XXV, 525 (1852)	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,669	M.fr.XI,27.				
S. Georgianus; De Candolle, prodr. VI, 371 (1837)	...	...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,670					
S. Gilherti; Turczaninow in Bull. Soc. Mosc. XXIV, 208 (1851)	...	...	W.A.	—	—	—	—	—	—	B.fl.III,670					
S. ramosissimus; De Candolle, prodr. VI, 371 (1837)	...	...	W.A.	—	—	—	—	—	—	B.fl.III,671					
S. odoratus; Hornemann, hort. reg. bot. Hafn. II, 809 (1815)	...	...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,671					
S. Cunninghamii; De Candolle, prodr. VI, 371 (1837)	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,671					
S. anethifolius; Cunningham in De Candolle, prodr. VI, 371 (1837)	...	...	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.III,672					
S. Bedfordii; F. v. M., annual report 26 (1858)	...	...	—	—	T.	V.	N.S.W.	—	—	B.fl.III,673					
S. Billardieri; F. v. M., annual report 26 (1858)	...	...	—	—	T.	—	—	—	—	B.fl.III,673					

ERECHTITES ; Rafinesque, Fl. Ludov. 65 (1817).

E. prenanthoides ; De Candolle, prodr. VI, 296 (1837)	...	...	...	...	—	S.A. T.	V.	N.S.W.	—	—	B.fl.III,658	...	...	N.Z.
E. Atkinsoniae ; F. v. M., fragm. V, 88 (1866)	...	...	...	...	—	—	—	N.S.W.	Q.	—	B.fl.III,658	M.fr.V,88.		
E. arguta ; De Candolle, prodr. VI, 296 (1837)	...	...	...	...	...	W.A.	S.A. T.	V.	N.S.W.	Q.	—	B.fl.III,659	...	N.Z.
E. picridioides ; Turczaninow in Bullet. de Mosc. XXIV, 200 (1851)	...	...	...	...	...	W.A.	S.A.	—	V.	—	—	—		
R. mixta ; De Candolle, prodr. VI, 297 (1837)	...	...	...	...	...	—	S.A.	—	V.	N.S.W.	—	B.fl.III,659		
E. quadridentata ; De Candolle, prodr. VI, 295 (1837)	...	...	...	...	...	W.A.	S.A. T.	V.	N.S.W.	Q.	—	B.fl.III,660	...	N.Z.
E. hispidula ; De Candolle, prodr. VI, 296 (1837)	...	...	...	...	...	W.A.	S.A. T.	V.	N.S.W.	—	—	B.fl.III,660		

CYMBONOTUS ; Cassini in Diction. XXXV, 397 (1825).

C. Lawsonianus ; Gaudichaud in Freyc. Voy. Bot. 462, t. 86 (1826)	...	...	...	...	...	W.A.	S.A. T.	V.	N.S.W.	Q.	—	B.fl.III,674	M.fr.XI,27.	
---	-----	-----	-----	-----	-----	------	---------	----	--------	----	---	--------------	-------------	--

SAUSSUREA ; De Candolle in Ann. du Mus. XVI, 156 (1810). (Aplotaxis, Haplotaxis.)

S. affinis ; Sprengel in De Candolle, prodr. VI, 540 (1837)	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.III,456	M.fr.I,36	As.
---	-----	-----	-----	-----	---	---	---	---	--------	----	---	--------------	-----------	-----

CENTAUREA ; Linné, hort. Cliffort. 420 (1737). (Leuzea partly.)

C. Australis ; Benth. & J. Hooker, gen. plant. II, 479 (1873)	...	...	...	...	—	—	—	V.	N.S.W.	Q.	—	B.fl.III,457		
---	-----	-----	-----	-----	---	---	---	----	--------	----	---	--------------	--	--

TRICHOCLINE ; Cassini in Bull. de la Soc. philom. 13 (1817). (Amblysperma, Gerberae subgenus.)

T. scapigera ; Benth. & J. Hooker, gen. plant. II, 479 (1873)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,676		
---	-----	-----	-----	-----	------	---	---	---	---	---	---	--------------	--	--

MICROSERIS ; D. Don in Edinb. phil. Mag. XI, 388 (1832). (Phyllopappus, Scozonera partly.)

M. Forsteri ; J. Hooker, Fl. N. Zeal. I, 151 (1853)	...	...	...	...	...	W.A.	S.A. T.	V.	N.S.W.	Q.	—	B.fl.III,676	M.fr.XI,27	N.Z.
---	-----	-----	-----	-----	-----	------	---------	----	--------	----	---	--------------	------------	------

CREPIS ; Linné, gen. pl. 240 (1737). (Youngia.)

C. Japonica ; Benth. & J. Hongk. 194 (1861)	...	...	...	...	...	—	—	—	N.S.W.	Q.	—	B.fl.III,679	...	As., Afr., P.
---	-----	-----	-----	-----	-----	---	---	---	--------	----	---	--------------	-----	---------------

CAMPANULACEAE.

A. L. DE JUSSIEU, gen. plant. 163 (1789), from B. de Jussieu (1759).—Genera 4, Species 35.

LOBELIA ; Linné, Fl. Lappon. 227 (1737). (Rapuntium, Pratia, Grammatotheca, Holostigma, Isolobus.)

L. heterophylla ; Labillardière, Nov. Holl. pl. spec. I, 52, t. 74 (1804)	...	...	...	...	...	W.A.	S.A.	—	—	—	N.A.	B.fl.IV,124	M.fr.IV,184; X,41.	
L. Browniana ; Roemer & Schultes, syst. veg. V, 71 (1819)	...	...	...	...	...	—	S.A. T.	V.	N.S.W.	Q.	—	—	M.fr.X,41.	
L. simplicicaulis ; R. Brown, prodr. 564 (1810)	...	...	...	...	...	—	S.A. T.	V.	N.S.W.	—	—	B.fl.IV,124	M.fr.IV,184; X,41.	
L. microsperma ; F. v. M., fragm. X, 41 (1876)	...	...	...	...	...	W.A.	S.A. T.	V.	N.S.W.	Q.	—	—	M.fr.X,41.	
L. dentata ; Cavanilles, icon. pl. VI, 14, t. 522 (1801)	...	...	...	...	...	—	—	—	N.S.W.	Q.	—	B.fl.IV,125	M.fr.IV,183; X,42.	
L. gracilis ; Andrews, Bot. Reposit. t. 340 (1805)	...	...	...	...	...	—	—	—	N.S.W.	Q.	—	B.fl.IV,125	M.fr.IV,183; X,42.	
L. rhytidosperra ; Benth. & J. Austr. IV, 126 (1869)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV,126	M.fr.X,40.	
L. tenuior ; R. Brown, prodr. 564 (1810)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV,126	M.fr.IV,183; X,42.	
L. rhombifolia ; De Vriese in Lehm. pl. Preiss. I, 397 (1845)	...	...	...	...	...	W.A.	S.A.	—	V.	—	—	B.fl.IV,126	M.fr.X,42.	
L. parvifolia ; R. Brown, prodr. 564 (1810)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV,127	M.fr.IV,183.	
L. trigonocaulis ; F. v. M., fragm. I, 18 (1858)	...	...	...	...	...	—	—	—	N.S.W.	Q.	—	B.fl.IV,127	M.fr.I,18,242.	
L. Bergiana ; Chamisso in Schlecht. Linnaea VIII, 217 (1835)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV,127	M.fr.II,20; IV,171; X,43.	Afr.
L. anceps ; Thunberg, prodr. plant. Cap. 40 (1794)	...	...	...	...	...	W.A.	S.A. T.	V.	N.S.W.	Q.	—	B.fl.IV,128	M.fr.IV,184; VI,114; X,43	Afr., Am., N.Z.
L. surrepens ; J. Hooker, Fl. Tasman. I, 237, t. 69 (1860)	...	...	...	...	...	—	—	T.	—	—	—	B.fl.IV,129	M.fr.IV,183.	
L. membranacea ; R. Brown, prodr. 563 (1810)	...	...	...	...	...	—	—	—	—	—	Q.	B.fl.IV,129	M.fr.IV,171; X,42.	
L. stenophylla ; Benth. & J. Austr. III, 130 (1869)	...	...	...	...	...	—	—	—	—	—	Q. N.A.	B.fl.IV,130	M.fr.X,42.	
L. humistrata ; F. v. M. in Benth. Fl. Austr. III, 130 (1869)	...	...	...	...	...	W.A.	—	—	—	—	Q. N.A.	B.fl.IV,130	M.fr.IV,171; X,42.	
L. dioica ; R. Brown, prodr. 565 (1810)	...	...	...	...	...	—	—	—	—	—	Q. N.A.	B.fl.IV,130	M.fr.IV,183; X,43.	
L. purpurascens ; R. Brown, prodr. 563 (1810)	...	...	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	B.fl.IV,131	M.fr.IV,183; X,42.	
L. pratioides ; Benth. & J. Austr. IV, 131 (1869)	...	...	...	...	...	—	S.A. T.	V.	N.S.W.	—	—	B.fl.IV,131	M.fr.X,43.	
L. irrigua ; R. Brown, prodr. 563 (1810)	...	...	...	...	...	—	—	T.	—	—	—	B.fl.IV,132		

L. gelida; F. v. M., fragm. IV, 183 (1864) ..	...	...	...	...	...	—	—	—	V.	N.S.W.	—	—	B.fl.IV,132	M.fr.IV,183.
L. platycalyx; F. v. M., fragm. IV, 183 (1864) ..	...	...	...	...	...	—	S.A.	T.	V.	—	—	—	B.fl.IV,133	M.fr.X,43.
L. concolor; R. Brown, prodr. 563 (1810) ..	...	...	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.IV,133	M.fr.X,43.
L. Benthani; F. v. M., Cens. of Austr. pl. 85 (1882) ..	...	...	...	...	...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.IV,133	
L. pedunculata; R. Brown, prodr. 563 (1810) ..	...	...	...	...	...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,133	M.fr.X,42.

ISOTOMA; R. Brown, prodr. 565 (1810). (Lobelia partly, Enchysia partly, Laurencia partly.)

I. Brownii; G. Don, gen. syst. III, 716 (1834) ..	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,134	M.fr.X,40.
I. axillaris; Lindley, Bot. Regist. t. 964 (1826) ..	...	...	...	...	...	—	—	—	V.	N.S.W.	Q.	—	B.fl.IV,135	M.fr.X,39.
I. petraea; F. v. M. in Linnaea XXV, 420 (1852) ..	...	...	...	...	...	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.IV,135	M.fr.X,39.
I. pusilla; Bentham in Hueg. enum. 75 (1837) ..	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,135	
I. Gulliverii; F. v. M., fragm. X, 39 (1876) ..	...	...	...	...	...	—	—	—	—	—	—	Q. N.A.	—	M.fr.X,39.
I. scapigera; G. Don, gen. syst. III, 716 (1834) ..	...	...	...	...	...	W.A.	S.A.	—	—	—	—	—	B.fl.IV,136	M.fr.IV,183; X,39.
I. fluviatilis; F. v. M. in Bentham, Fl. Austr. IV, 136 (1869) ..	...	...	...	...	...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,136	M.fr.X,39.

WAHLENBERGIA; Schrader, cat. hort. bot. Goetting. (1814). (Strzeleckia.)

W. gracilis; A. de Candolle, Monogr. comp. 142 (1830) ..	...	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.IV,137	...	...	...	As., N.Z.
--	-----	-----	-----	-----	-----	------	------	----	----	--------	----	------	-------------	-----	-----	-----	-----------

SPHENOCLEA; Gaertner, de fructibus I, 113 (1788).

S. Zeilanica; Gaertner de fruct. I, 113, t. 24 (1788)...	...	...	...	...	...	—	—	—	—	—	—	N.A.	—	...	...	...	As., Afr., Am.
--	-----	-----	-----	-----	-----	---	---	---	---	---	---	------	---	-----	-----	-----	----------------

CANDOLLEACEAE.

F. v. M., fragm. VIII, 41 (1873). (Stylideae; R. Brown, prodr. 565 (1810).—Genera 4, Species 97.

CANDOLLEA; Labillardière in Ann. du Mus. VI, 451 (1805.) (Stylidium 1805, Ventenata, Forsteropsis.)

C. carnosa; F. v. M., Cens. 85 (1882) from Benth. (1837) ..	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,6	
C. pilosa; Labillardière in. Annal. du Mus. VI, 453, t. 63 (1805)...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,7	
C. reduplicata; F. v. M., Cens. 85 (1882) from R. Brown (1810) ..	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,7	
C. scabrida; F. v. M., Cens. 85 (1882) from Lindley (1839)...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,8	
C. hirsuta; F. v. M., Cens. 85 (1882) from R. Brown (1810) ..	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,8	
C. crossoccephala; F. v. M., Cens. 85 (1882) ..	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,8	M.fr.VI,5,254; X,58.
C. juncea; F. v. M., Cens. 85 (1882) from R. Brown (1810) ..	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,9	M.fr.X,58.
C. guttata; F. v. M., Cens. 85 (1882) from R. Brown (1810) ..	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,9	
C. repens; F. v. M., Cens. 85 (1882) from R. Brown (1810) ..	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,9	M.fr.X,58.
C. serrulata; Labillardière in Annal. du Mus. VI, 454, t. 63 (1805) ..	...	...	...	...	...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,10	M.fr.IV,94; VI,6; X,58.
C. linearis; F. v. M., Cens. 85 (1882) from Swartz (1807) ..	...	...	...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.IV,11	M.fr.IV,95.
C. elongata; F. v. M., Cens. 85 (1882) from Benth. (1869)...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,583	M.fr.X,86.
C. spinulosa; F. v. M., Cens. 85 (1882) from R. Brown (1810) ..	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,11	
C. Tepperiana; F. v. M., in Melb. Chem. and Drugg. II, 15 (1887) ..	...	...	...	...	...	—	S.A.	—	—	—	—	—	—	
C. limbata; F. v. M., Cens. 86 (1882) ..	...	...	...	...	...	W.A.	—	—	—	—	—	—	—	M.fr.X,57.
C. caespitosa; F. v. M., Cens. 86 (1882) from R. Brown (1810) ..	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,11	M.fr.X,57.
C. squamellosa; F. v. M., Cens. 86 (1882) from De Candolle (1838) ..	...	...	...	...	...	W.A.	—	—	—	—	—	—	—	M.fr.X,56.
C. violacea; F. v. M., Cens. 86 (1882) from R. Brown (1810) ..	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,12	
C. lutea; F. v. M., Cens. 86 (1882) from R. Brown (1810) ..	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,12	
C. filifera; F. v. M., Cens. 86 (1882) from R. Brown (1810) ..	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,12	M.fr.X,58,86.
C. ciliata; F. v. M., Cens. 86 (1882) from Lindley (1839) ..	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,13	
C. sobolifera; F. v. M., Cens. 86 (1882) ..	...	...	...	...	...	—	—	—	V.	—	—	—	B.fl.IV,13	M.fr.IV,94.
C. Flondii; F. v. M., Cens. 86 (1882) ..	...	...	...	...	...	—	—	—	—	—	Q. N.A.	B.fl.IV,13	M.fr.I,149; VI,6; X,58.	
C. disperma; F. v. M., Cens. 86 (1882) ..	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,14	M.fr.IV,93.
C. calcarata; F. v. M., Cens. 86 (1882) from R. Brown (1810) ..	...	...	...	...	...	W.A.	S.A.	—	V.	—	—	N.A.	B.fl.IV,14	M.fr.IV,94; VI,6.
C. perpusilla; F. v. M., Cens. 86 (1882) from J. Hooker (1847) ..	...	...	...	...	...	W.A.	S.A.	T.	V.	—	—	—	B.fl.IV,15	M.fr.VI,78; X,58.

C. eriorrhiza; F. v. M., Cens. 86 (1882) from R. Brown (1810) ...	...	...	—	—	—	—	Q.	—	B.fl.IV,15	M.fr.I,147; VI,6; X,58.	
C. debilis; F. v. M., Cens. 86 (1882) ...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,15	M.fr.I,149; VI,6; X,58.
C. floribunda; F. v. M., Cens. 86 (1882) from R. Brown (1810) ...	...	...	—	S.A.	—	—	Q.	N.A.	B.fl.IV,15	M.fr.I,148; VI,6; X,58.	
C. leptorrhiza; F. v. M., Cens. 86 (1882) ...	...	...	—	—	—	—	—	N.A.	B.fl.IV,16	M.fr.I,148.	
C. assimilis; F. v. M., Cens. 86 (1882) from R. Brown (1810) ...	...	...	W.A.	—	—	—	—	—	B.fl.IV,16	M.fr.X,57.	
C. rupestris; F. v. M., Cens. 86 (1882) from Sonder (1845)...	...	...	W.A.	—	—	—	—	—	B.fl.IV,16		
C. spatulata; F. v. M., Cens. 86 (1882) from R. Brown (1810) ...	...	...	W.A.	—	—	—	—	—	B.fl.IV,17		
C. Barleei; F. v. M., Cens. 86 (1882) ...	...	...	W.A.	—	—	—	—	—	B.fl.IV,17	M.fr.VI,5,249.	
C. lineata; F. v. M., Cens. 86 (1882) from Sonder (1845) ...	...	...	W.A.	—	—	—	—	—	B.fl.IV,17		
C. glauca; Labillardière in Annal. du Mus. VI, 454, t. 64 (1805) ...	...	...	W.A.	—	—	—	—	—	B.fl.IV,17		
C. amoena; F. v. M., Cens. 86 (1882) from R. Brown (1810) ...	...	...	W.A.	—	—	—	—	—	B.fl.IV,18		
C. striata; F. v. M., Cens. 86 (1882) from Lindley (1839) ...	...	...	W.A.	—	—	—	—	—	B.fl.IV,18		
C. diversifolia; F. v. M., Cens. 86 (1882) from R. Brown (1810) ...	...	...	W.A.	—	—	—	—	—	B.fl.IV,18	M.fr.X,58.	
C. articulata; F. v. M., Cens. 86 (1882) from R. Brown (1810) ...	...	...	W.A.	—	—	—	—	—	B.fl.IV,19		
C. Brunonia; F. v. M., Cens. 86 (1882) from Benthams (1837) ...	...	...	W.A.	—	—	—	—	—	B.fl.IV,19		
C. diuroides; F. v. M., Cens. 86 (1882) from Lindley (1839) ...	...	...	W.A.	—	—	—	—	—	B.fl.IV,20		
C. scandens; F. v. M., Cens. 86 (1882) from R. Brown (1810) ...	...	...	W.A.	—	—	—	—	—	B.fl.IV,20	M.fr.VI,78.	
C. verticillata; F. v. M., Cens. 86 (1882) ...	...	...	W.A.	—	—	—	—	—	B.fl.IV,20	M.fr.IV,94.	
C. glandulosa; F. v. M., Cens. 86 (1882) from Salisbury (1807) ...	...	...	W.A.	—	—	—	—	—	B.fl.IV,20		
C. laricifolia; F. v. M., Cens. 86 (1882) from Richard (1807) ...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,21	
C. Preissii; F. v. M., Cens. 86 (1882) ...	...	...	W.A.	—	—	—	—	—	B.fl.IV,21	M.fr.III,122.	
C. imbricata; F. v. M., Cens. 86 (1882) from Benthams (1837) ...	...	...	W.A.	—	—	—	—	—	B.fl.IV,21		
C. adpressa; F. v. M., Cens. 86 (1882) from Benthams (1869) ...	...	...	W.A.	—	—	—	—	—	B.fl.IV,22		
C. despecta; F. v. M., Cens. 86 (1882) from R. Brown (1810) ...	...	...	W.A.	S.A.	T.	V.	N.S.W.	—	B.fl.IV,22	M.fr.IV,94; VI,58.	
C. utricularioides; F. v. M., Cens. 86 (1882) from Benthams (1839) ...	...	...	W.A.	—	—	—	—	—	B.fl.IV,23		
C. pygmaea; F. v. M., Cens. 86 (1882) from R. Brown (1810) ...	...	...	W.A.	—	—	—	—	—	B.fl.IV,23		
C. longitubea; F. v. M., Cens. 86 (1882) from Benthams (1837) ...	...	...	W.A.	—	—	—	—	—	B.fl.IV,23		
C. diffusa; F. v. M., Cens. 86 (1882) from R. Brown (1810) ...	...	...	—	—	—	—	Q.	—	B.fl.IV,23		
C. fissiloba; F. v. M., Cens. 86 (1882) ...	...	...	—	—	—	—	—	N.A.	B.fl.IV,24	M.fr.I,154.	
C. trichopoda; F. v. M., Cens. 86 (1882) ...	...	...	—	—	—	—	Q.	—	—	M.fr.X,86.	
C. alsinoides; F. v. M., Cens. 86 (1882) from R. Brown (1810) ...	...	...	—	—	—	—	Q.	N.A.	B.fl.IV,24	M.fr.I,151; VI,6; X,58.	
C. tencrura; F. v. M., Cens. 86 (1882) ...	...	...	—	—	—	—	—	N.A.	B.fl.IV,24	M.fr.I,150.	
C. brachyphylla; F. v. M., Cens. 86 (1882) from Sonder (1845) ...	...	...	W.A.	—	—	—	—	—	B.fl.IV,24		
C. capillaris; F. v. M., Cens. 86 (1882) from R. Brown (1810) ...	...	...	—	—	—	—	Q.	—	B.fl.IV,25		
C. rotundifolia; F. v. M., Cens. 86 (1882) from R. Brown (1810) ...	...	...	—	—	—	—	Q.	N.A.	B.fl.IV,25	M.fr.I,151.	
C. schizantha; F. v. M., Cens. 86 (1882) ...	...	...	—	—	—	—	—	N.A.	B.fl.IV,25	M.fr.I,152.	
C. lobuliflora; F. v. M., Cens. 86 (1882) ...	...	...	—	—	—	—	—	N.A.	B.fl.IV,25	M.fr.I,153.	
C. uliginosa; F. v. M., Cens. 86 (1882) from Swartz (1807) ...	...	...	—	—	—	—	Q.	—	B.fl.IV,26	... As.	
C. pulchella; F. v. M., Cens. 86 (1882) from Sonder (1845) ...	...	...	W.A.	—	—	—	—	—	B.fl.IV,26		
C. petiolaris; F. v. M., Cens. 86 (1882) from Sonder (1845) ...	...	...	W.A.	—	—	—	—	—	B.fl.IV,26		
C. emarginata; F. v. M., Cens. 86 (1882) from Sonder (1845) ...	...	...	W.A.	—	—	—	—	—	B.fl.IV,27		
C. corymbosa; F. v. M., Cens. 86 (1882) from R. Brown (1810) ...	...	...	W.A.	—	—	—	—	—	B.fl.IV,27		
C. lepida; F. v. M., Cens. 86 (1882) from Benthams (1869) ...	...	...	W.A.	—	—	—	—	—	B.fl.IV,27		
C. streptocarpa; F. v. M., Cens. 86 (1882) from Sonder (1845) ...	...	...	W.A.	—	—	—	—	—	B.fl.IV,27		
C. uniflora; F. v. M., Cens. 86 (1882) from Sonder (1845) ...	...	...	W.A.	—	—	—	—	—	B.fl.IV,28	M.fr.VI,6.	
C. pedunculata; F. v. M., Cens. 86 (1882) from R. Brown (1810) ...	...	...	—	—	—	—	Q.	N.A.	B.fl.IV,28	M.fr.VI,91.	
C. pachyrrhiza; F. v. M., Cens. 86 (1882) ...	...	...	—	—	—	—	—	N.A.	B.fl.IV,28	M.fr.I,152.	
C. muscicola; F. v. M., Cens. 86 (1882) ...	...	...	—	—	—	—	—	N.A.	B.fl.IV,29	M.fr.I,153.	
C. crassifolia; F. v. M., Cens. 86 (1882) from R. Brown (1810) ...	...	...	W.A.	—	—	—	—	—	B.fl.IV,29		
C. pycnostachya; F. v. M., Cens. 86 (1882) from Lindley (1839) ...	...	...	W.A.	—	—	—	—	—	B.fl.IV,29		

<i>C. pubigera</i> ; F. v. M., Cens. 86 (1882) from Sonder (1845) ...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,30	
<i>C. canaliculata</i> ; F. v. M., Cens. 86 (1882) from Lindley (1839) ...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,30	
<i>C. leptophylla</i> ; F. v. M., Cens. 86 (1882) from De Candolle (1838) ...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,30	
<i>C. dichotoma</i> ; F. v. M., Cens. 86 (1882) from De Candolle (1838)...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,30	
<i>C. bulbifera</i> ; F. v. M., Cens. 86 (1882) from Bentham (1837) ...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,31	
<i>C. Merralliana</i> ; F. v. M. in Vict. Naturalist V, 76 (1888)...	...	...	...	W.A.	—	—	—	—	—	—	—	
<i>C. breviscapa</i> ; F. v. M., Cens. 86 (1882) from R. Brown (1810) ...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,31	
<i>C. eglandulosa</i> ; F. v. M., Cens. 86 (1882) ...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,31	M.fr.I,150; VI,6; X,58.
<i>C. fasciculata</i> ; F. v. M., Cens. 86 (1882) from R. Brown (1810) ...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,32	
<i>C. falcata</i> ; F. v. M., Cens. 86 (1882) from R. Brown (1810) ...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,32	
<i>C. rhynchocharpa</i> ; F. v. M., Cens. 86 (1882) from Sonder (1845) ...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,33	
<i>C. adnata</i> ; F. v. M., Cens. 86 (1882) from R. Brown (1810) ...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,33	M.fr.X,58.

LEEWENHOEKIA; R. Brown, prodr. 572 (1810). (Levenhookia, Coleostylis.)

L. pusilla; R. Brown, prodr. 573 (1810)	...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,34	M.fr.X,58.
L. dubia; Sonder in Lehm. pl. Preiss. I, 392 (1845)	...	...	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.IV,34	M.fr.III,121; IV,94; X,58.
L. Sonderi; F. v. M., fragm. I, 18 (1858)	...	...	...	...	...	...	—	—	—	V.	—	—	—	B.fl.IV,35	
L. pauciflora; Benth in Hueg. enum. 74 (1837)	...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,35	M.fr.VI,77.
L. leptantha; Benth, Fl. Austr. IV, 35 (1869)	...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,35	
L. stipitata; F. v. M., fragm. IV, 94 (1864)	...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,36	M.fr.IV,94.
L. Preissii; F. v. M., fragm. IV, 94 (1864)	...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,36	M.fr.IV,94.

PHYLLACHNE; R. et G. Forster, char. gen. 115, t. 58 (1776). (Forstera, Helophyllum, Oreostylidium.)

P. bellidifolia; F. v. M., fragm. VIII, 39 (1873) ... — — T. — — — — B.fl.IV,37 M.fr.X,58.

DONATIA ; R. et G. Forster, char. gen. 9, t. 5 (1776).

D. Novae Zealandiae; J. Hooker, Fl. N. Zeal. I, 81, t. 20 (1853)... — — T. — — — — B.fl.II,450 M.fr.IV,3; VII,150; VIII,41; X,58. N.Z.

GOODENIACEAE.

R. BROWN, prodr. 573 (1810).—Genera 11, Species 220.

BRUNONIA ; Smith in Transact. Linn. Soc. X, 366 (1809).

B. Australis; Smith in Transact. Linn. Soc. X, 367, t. 28 (1809)... .. W.A. S.A. T. V. N.S.W. Q. N.A. B.f.IV, 121 M.fr.X, 111.

DAMPIERA ; R. Brown, prodr. 587 (1810). (Linschottenia.)

D. luteiflora ; F. v. M., fragm. X, 11 (1876) ...	...	...	...	...	...	W.A.	—	—	—	—	—	—	—	M.fr.X, 11.
D. Linschotenii ; F. v. M., fragm. VI, 28 (1866) ...	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl.IV, 108	M.fr.VI, 28.
D. spicigera ; Benth., Fl. Austr. IV, 199 (1869) ...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV, 109	
D. candicans ; F. v. M., fragm. X, 86 (1876) ...	...	...	...	...	...	W.A.	S.A.	—	—	—	—	—	—	M.fr.X, 86.
D. teres ; Lindley, Bot. Regist. XXV, App. 27 (1839) ...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV, 109	
D. trigona ; De Vriese in Lehm. pl. Preiss. I, 401 (1845) ...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV, 109	M.fr.II, 17; VIII, 245; X, 11.
D. prostrata ; De Vriese, Goodenov. 83 (1854) ...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV, 110	
D. alata ; Lindley, Bot. Regist. XXV, App. 27 (1839) ...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV, 110	M.fr.X, 11.
D. coronata ; Lindley, Bot. Regist. XXV, App. 27 (1839) ...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV, 110	
D. carinata ; Benth., Fl. Austr. IV, 111 (1869) ...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV, 111	
D. sacculata ; F. v. M. in Benth. Fl. Austr. IV, 111 (1869) ...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV, 111	
D. incana ; R. Brown, prodr. 588 (1810) ...	...	...	...	...	...	W.A.	—	—	—	—	N.A.	B.fl.IV, 111	M.fr.II, 17.	
D. hederacea ; R. Brown, prodr. 588 (1810) ...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV, 112	M.fr.X, 12.
D. ferruginea ; R. Brown, prodr. 588 (1810) ...	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl.IV, 112	M.fr.VI, 29.
D. Brownii ; F. v. M., fragm. VI, 29 (1866) ...	...	...	...	...	...	—	—	—	V.	N.S.W.	Q.	—	B.fl.IV, 112	M.fr.VI, 29.
D. lanceolata ; Cunningham in De Candolle, prodr. VII, 503 (1839) ...	...	...	...	...	...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.IV, 113	M.fr.VI, 29.
D. altissima ; F. v. M. in Benth. Fl. Austr. IV, 113 (1869) ...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV, 113	

SCAEVOLOA; Linné, mantiss. II, 145 (1771). (Pogonetes, Crossotoma, Temminckia, Kamphusia, Molkenboeria, Merkusia, Verrcauxia.)

S. Koenigii; Vahl, symb. bot. III, 36 (1794) ...	...	...	...	...	...	...	...	...	...	Q. N.A.	B.fl.IV,86	M.fr.XI,76	...	...	As., Afr.
S. enantophylla; F. v. M., fragm. VIII, 58 (1873) ...	...	...	...	...	...	...	...	...	...	Q.	—	M.fr.VI,225.			
S. spinescens; R. Brown, prodr. 586 (1810) ...	...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q. N.A.	B.fl.IV,87	M.fr.XI,20,27,76.		
S. Groeneri; F. v. M., fragm. VI, 15 (1866) ...	...	...	...	...	...	W.A.	S.A.	—	—	—	—	B.fl.IV,88	M.fr.VIII,58.		
S. tomentosa; Gaudichaud in Freyc. Voy. Bot. 460, t. 81 (1826) ...	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV,88			
S. atriplicina; F. v. M., fragm. II, 18 (1860) ...	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV,88	M.fr.III,165,181.		
S. striata; R. Brown, prodr. 586 (1810) ...	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV,89			
S. phlebotopetala; F. v. M., fragm. II, 18 (1860) ...	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV,89	M.fr.II,VI,16; XI,76.		
S. pilosa; Benth. in Hueg. enum. 69 (1837) ...	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV,89			
S. hispida; Cavanilles, icon. et descr. pl. VI, 7, t. 510 (1801) ...	...	...	...	...	...	—	—	—	V.	N.S.W.	Q.	—	B.fl.IV,90	M.fr.I,121,207; XI,76.	
S. apterantha; F. v. M., fragm. I, 121 (1859) ...	...	...	...	...	...	—	—	—	V.	—	—	—	B.fl.IV,90	M.fr.I,121.	
S. Hookeri; F. v. M., gen. report, 15 (1853) ...	...	...	...	...	...	—	—	—	T.	V.	N.S.W.	—	—	B.fl.IV,90	M.fr.XI,76.
S. parvifolia; F. v. M. in Benth. Fl. Austr. IV, 91 (1869) ...	...	...	...	...	...	W.A.	S.A.	—	—	—	Q. N.A.	B.fl.IV,91	M.fr.VIII,58; X,59; XI,76.		
S. oxyclona; F. v. M., fragm. X, 58 (1876) ...	...	...	...	...	...	W.A.	—	—	—	—	—	—	M.fr.X,58.		
S. restiacea; Benth. Fl. Austr. IV, 91 (1869) ...	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV,91			
S. depauperata; R. Brown, App. Sturt's Exped. 20 (1849) ...	...	...	...	...	...	—	S.A.	—	—	—	N.A.	B.fl.IV,91	M.fr.III,33; VIII,58; XI,76.		
S. tortuosa; Benth. Fl. Austr. IV, 91 (1869) ...	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV,91			
S. Cunninghamii; De Candolle, prodr. VII, 508 (1838) ...	...	...	...	...	...	—	—	—	—	—	N.A.	B.fl.IV,92			
S. collaris; F. v. M., Rep. Babb. Exped. 15 (1858) ...	...	...	...	...	...	—	S.A.	—	—	—	—	B.fl.IV,92	M.fr.II,20; VIII,58; XI,76.		
S. angulata; R. Brown, prodr. 586 (1810) ...	...	...	...	...	...	—	—	—	—	—	Q. N.A.	B.fl.IV,92			
S. Brooksiana; F. v. M. in Vict. Naturalist I, 122 (1884) ...	...	...	...	...	...	W.A.	—	—	—	—	—	—			
S. nitida; R. Brown, prodr. 584 (1810) ...	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV,93			
S. globuliflora; Labillardière, Nov. Holl. pl. spec. I, 55, t. 78 (1804) ...	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV,93			
S. porocarya; F. v. M., fragm. II, 19 (1860) ...	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV,94	M.fr.II,19; XI,76.		
S. attenuata; R. Brown, prodr. 583 (1810) ...	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV,94			
S. glandulifera; De Candolle, prodr. VII, 510 (1838) ...	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV,94			
S. anchusifolia; Benth. in Hueg. enum. 68 (1837) ...	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV,94			
S. holosericea; De Vriese in Lehm. pl. Preiss. I, 408 (1845) ...	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV,95			
S. snaveolens; R. Brown, prodr. 585 (1810) ...	...	...	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.IV,95		
S. revoluta; R. Brown, prodr. 586 (1810) ...	...	...	...	...	...	—	—	—	—	—	Q. N.A.	B.fl.IV,96	M.fr.I,121.		
S. crassifolia; Labillardière, Nov. Holl. pl. spec. I, 56, t. 79 (1804) ...	...	...	...	...	...	W.A.	S.A.	—	V.	—	—	B.fl.IV,96	M.fr.VIII,245; XI,20,76.		
S. macrostachya; Benth. Fl. Austr. IV, 97 (1869) ...	...	...	...	...	...	—	—	—	—	—	N.A.	B.fl.IV,97	M.fr.VIII,58; XI,76.		
S. longifolia; De Vriese in Lehm. pl. Preiss. I, 410 (1845) ...	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV,97			
S. lanceolata; Benth. in Hueg. enum. 69 (1837) ...	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV,97	M.fr.I,207; XI,76.		
S. thesioides; Benth. in Hueg. enum. 68 (1837) ...	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV,98	M.fr.I,207; XI,77.		
S. macrophylla; Benth. Fl. Austr. IV, 98 (1869) ...	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV,98			
S. platyphylla; Lindley, Bot. Regist. XXV, App. 26 (1839) ...	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV,98	M.fr.XI,77.		
S. auriculata; Benth. Fl. Austr. IV, 99 (1869) ...	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV,99	M.fr.XI,77.		
S. ovalifolia; R. Brown, prodr. 584 (1810) ...	...	...	...	...	...	W.A.	S.A.	—	N.S.W.	Q. N.A.	B.fl.IV,96	M.fr.VIII,58; XI,76.			
S. aemula; R. Brown, prodr. 584 (1810) ...	...	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,99	M.fr.XI,77.	
S. humilis; R. Brown, prodr. 585 (1810) ...	...	...	...	...	...	—	S.A.	—	—	—	—	B.fl.IV,100			
S. amblyanthera; F. v. M., fragm. I, 121 (1859) ...	...	...	...	...	...	—	—	—	—	—	Q. N.A.	B.fl.IV,100	M.fr.I,121.		
S. microphylla; Benth. Fl. Austr. IV, 100 (1869) ...	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV,100			
S. cuneiformis; Labillardière, Nov. Holl. pl. spec. I, 56, t. 80 (1804) ...	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV,101	M.fr.XI,77.		
S. microcarpa; Cavanilles, icon. et descr. pl. VI, 6, t. 509 (1801) ...	...	...	...	...	...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,101	M.fr.VI,16; XI,77.	
S. linearis; R. Brown, prodr. 586 (1810) ...	...	...	...	...	...	—	S.A.	—	—	—	—	B.fl.IV,102			
S. Oldfieldii; F. v. M., fragm. II, 19 (1860) ...	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV,102	M.fr.II,19; XI,77.		
S. paludosa; R. Brown, prodr. 586 (1810) ...	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV,102	M.fr.XI,77.		
S. sericophylla; F. v. M. in Benth. Fl. Austr. IV, 102 (1869) ...	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV,102	M.fr.XI,77.		

S. caesescens; Benth in Hueg. enum. 69 (1837)	W.A.	—	—	—	—	—	—	B.fl.IV,103	
S. humifusa; De Vriese in Lehm. pl. Preiss. I, 410 (1845)	W.A.	—	—	—	—	—	—	B.fl.IV,103	M.fr.XI,77.
S. fasciculata; Benth in Hueg. enum. 68 (1837)	W.A.	—	—	—	—	—	—	B.fl.IV,104	M.fr.XI,77.
S. stenophylla; Benth in Austr. IV, 104 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,104	M.fr.I,113; XI,77.
S. Reinwardtii; De Vriese in Lehm. pl. Preiss. I, 409 (1845)	W.A.	—	—	—	—	—	—	B.fl.IV,106	M.fr.II,17.
S. Verreauxii; F. v. M., Bot. Teachings 65 (1877)	W.A.	—	—	—	—	—	—	B.fl.IV,105	M.fr.II,17.

SELLIERA; Cavanilles in Anal. Cien. Nat. I, 41, t. 5 (1799).

S. radicans; Cavanilles, icon. et descr. pl. V, 49, t. 474 (1799)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.IV,82	M.fr.VIII,58	 Am., N.Z.
S. exigua; F. v. M., fragm. III, 142 (1862)	W.A.	—	—	—	—	—	—	B.fl.IV,82	M.fr.III,142.	

CALOGYNE; R. Brown, prodr. 579 (1810). (Distylis.)

C. pilosa; R. Brown, prodr. 579 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,81	M.fr.VI,7,28.
C. Berardiana; F. v. M., fragm. VI, 7 (1867)	W.A.	S.A.	—	—	—	—	N.A.	B.fl.IV,81	M.fr.VI,7; VIII,58.
C. heteroptera; F. v. M., fragm. X, 43 (1876)	—	—	—	—	—	Q.	N.A.	—	M.fr.X,43.
C. purpurea; F. v. M., fragm. VIII, 57 (1873)	—	—	—	—	—	—	N.A.	—	M.fr.VIII,57.

GOODENIA; Smith in Transact. Linn. Soc. II, 347 (1794). (Goodenoughia, Tetraphylax, Picrophyta, Stekhovia, Aillya.)

G. phyllicoides; F. v. M., fragm. I, 206 (1859)	W.A.	—	—	—	—	—	—	B.fl.IV,55	M.fr.I,206.
G. viscida; R. Brown, prodr. 578 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,55	M.fr.I,120; III,35,163.
G. xanthotricha; De Vriese, Gooden. 155 (1854)	W.A.	—	—	—	—	—	—	B.fl.IV,56	M.fr.VI,13.
G. scapigera; R. Brown, prodr. 578 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,56	M.fr.I,114; III,163; XI,50.
G. quadrilocularis; R. Brown, prodr. 578 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,57	M.fr.III,141; IV,13.
G. Ramelii; F. v. M., fragm. III, 20, t. 17 (1862)	W.A.	S.A.	—	—	—	—	N.A.	B.fl.IV,57	M.fr.VIII,245; X,13.
G. pinifolia; De Vriese, Gooden. 157, t. 30 (1854)	W.A.	—	—	—	—	—	—	B.fl.IV,58	
G. decurrens; R. Brown, prodr. 575 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.IV,58	
G. racemosa; F. v. M., fragm. I, 114 (1859)	—	—	—	—	—	Q.	—	B.fl.IV,58	M.fr.I,114.
G. bellidifolia; Smith in Transact. Linn. Soc. II, 349 (1794)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,58	M.fr.I,117; XI,50.
G. stelligera; R. Brown, prodr. 575 (1810)	—	—	—	—	V.	N.S.W.	Q.	B.fl.IV,59	M.fr.VI,14.
G. Stevensonii; F. v. M. in Vict. Naturalist III, 138 (1887)	—	—	—	—	—	N.S.W.	—	—	
G. ovata; Smith in Transact. Linn. Soc. II, 347 (1794)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,59	M.fr.I,205; X,110.
G. varia; R. Brown, prodr. 576 (1810)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.IV,61	M.fr.I,205.
G. amplexans; F. v. M. in Transact. phil. Inst. Vict. II, 70 (1859)	—	S.A.	—	V.	—	—	—	B.fl.IV,60	M.fr.VI,15.
G. strophilata; F. v. M., fragm. I, 119 (1859)	W.A.	—	—	—	—	—	—	B.fl.IV,60	M.fr.I,119; II,111.
G. laevis; Benth in Austr. IV, 61 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,61	
G. barbata; R. Brown, prodr. 576 (1810)	—	—	T.	V.	N.S.W.	—	—	B.fl.IV,61	M.fr.II,110,176; VI,15.
G. arthrotricha; F. v. M. in Benth. Fl. Austr. IV, 62 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,62	
G. disperma; F. v. M., fragm. I, 113 (1859)	—	—	—	—	—	Q.	—	B.fl.IV,62	M.fr.IV,145; VI,15.
G. geniculata; R. Brown, prodr. 577 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,62	M.fr.VI,14.
G. hederacea; Smith in Transact. Linn. Soc. II, 349 (1794)	—	—	—	V.	N.S.W.	Q.	—	B.fl.IV,63	M.fr.VI,15,110.
G. xanthosperma; F. v. M., fragm. X, 12 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,12.
G. hirsuta; F. v. M., fragm. III, 35 (1862)	—	S.A.	—	—	—	—	N.A.	B.fl.IV,64	M.fr.III,35.
G. heterophylla; Smith in Transact. Linn. Soc. II, 349 (1794)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,64	M.fr.I,117.
G. deflexa; F. v. M. in Melb. Chem. Jan. (1883)	—	—	—	—	—	—	N.A.	—	
G. glabra; R. Brown, prodr. 577 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,64	M.fr.VI,14.
G. stronglylphylla; F. v. M., fragm. VI, 12 (1867)	—	—	—	—	—	Q.	—	B.fl.IV,65	M.fr.VI,12.
G. rotundifolia; R. Brown, prodr. 576 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,65	M.fr.VI,15.
G. azurea; F. v. M., fragm. I, 117 (1859)	—	—	—	—	—	—	N.A.	B.fl.IV,66	M.fr.X,87.
G. scaevolina; F. v. M., fragm. I, 118 (1859)	—	—	—	—	—	—	N.A.	B.fl.IV,66	M.fr.XI,50.
G. Stobbsiana; F. v. M., fragm. XI, 49 (1879)	—	—	—	—	—	—	N.A.	—	M.fr.XI,49,137.
G. incana; R. Brown, prodr. 578 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,66	M.fr.I,155; III,141.
G. leptoclada; Benth in Austr. IV, 67 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,67	M.fr.VI,227; VIII,186; XI,50.

G. Eatoniana; F. v. M., fragm. VIII, 186 (1874)	W.A.	—	—	—	—	—	—	M.fr.XI,51.	
G. coerulea; R. Brown, prodr. 578 (1810)	W.A.	—	—	—	—	—	B.fl.IV,67	M.fr.III,140; XI,20.	
G. trichophylla; De Vriese in Benth. Fl. Austr. IV, 67 (1869)	W.A.	—	—	—	—	—	B.fl.IV,67		
G. Hassallii; F. v. M., fragm. VI, 10, t. 51 (1867)	W.A.	—	—	—	—	—	B.fl.IV,68	M.fr.XI,50.	
G. pterygosperma; R. Brown, prodr. 578 (1810)	W.A.	—	—	—	—	—	B.fl.IV,68	M.fr.I,155; III,163.	
G. Vilmoriniae; F. v. M., fragm. III, 19, t. 16 (1862)	—	S.A.	—	—	—	N.A.	B.fl.IV,68	M.fr.VIII,245.	
G. Bonneyana; F. v. M. in Benth. Fl. Austr. IV, 69 (1869)	W.A.	—	—	—	—	—	B.fl.IV,69	M.fr.VI,226,250.	
G. calcarata; F. v. M., fragm. VI, 14 (1867)	—	S.A.	—	—	N.S.W.	Q.	B.fl.IV,69	M.fr.I,119; VIII,245.	
G. albiflora; Schlechtendal, Linnaea XX, 599 (1847)	—	S.A.	—	—	—	—	B.fl.IV,70	M.fr.I,119.	
G. Nicholsoni; F. v. M., fragm. I, 203, t. 4 (1859)	—	S.A.	—	—	—	—	B.fl.IV,69	M.fr.I,203.	
G. Macmillani; F. v. M., fragm. I, 119, t. 5 (1859)	—	—	—	V.	—	—	B.fl.IV,69	M.fr.I,119.	
G. grandiflora; Sims, Bot. Mag. t. 890 (1806)	W.A.	S.A.	—	—	N.S.W.	Q.	B.fl.IV,69	M.fr.I,204; VI,14.	
G. Chambersii; F. v. M., fragm. I, 204 (1859)	—	S.A.	—	—	—	N.A.	B.fl.IV,70	M.fr.I,204.	
G. Strangfordii; F. v. M., fragm. VI, 11, t. 52 (1867)	—	S.A.	—	—	—	Q.	N.A.	B.fl.IV,70	
G. Mitchellii; Benth. Fl. Austr. IV, 71 (1869)	—	S.A.	—	—	—	Q.	—	B.fl.IV,71	M.fr.XI,75.
G. Mueckeana; F. v. M., fragm. VIII, 56 (1873)	W.A.	S.A.	—	—	—	—	—	—	M.fr.IX,194; XI,133.
G. melanoptera; F. v. M., fragm. I, 115 (1859)	—	—	—	—	—	N.A.	—	—	M.fr.I,115.
G. heterochila; F. v. M., fragm. III, 142 (1863)	—	S.A.	—	—	—	N.A.	B.fl.IV,71	—	M.fr.VI,12; VIII,245.
G. sepulosa; F. v. M. in Benth. Fl. Austr. IV, 72 (1869)	—	—	—	—	—	—	N.A.	B.fl.IV,72	
G. hispida; R. Brown, prodr. 577 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,72	M.fr.VI,12.
G. auriculata; Benth. Fl. Austr. IV, 72 (1869)	—	—	—	—	—	—	N.A.	B.fl.IV,72	
G. Armstrongiana; De Vriese, Goodenov. 138, t. 24 (1854)	—	—	—	—	—	—	N.A.	B.fl.IV,73	M.fr.I,205; VII,40.
G. corynocarpa; F. v. M., fragm. II, 16 (1860)	W.A.	—	—	—	—	—	—	B.fl.IV,73	M.fr.VI,15.
G. mollissima; F. v. M. in Benth. Fl. Austr. IV, 73 (1869)	—	—	—	—	—	Q.	—	B.fl.IV,73	
G. O'Donnelli; F. v. M. in Austr. Journ. of Pharm. I, 278 (1886)	—	—	—	—	—	—	N.A.	—	
G. cycloptera; R. Brown in App. Sturt's Exped. 83 (1849)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.IV,74	M.fr.VI,14; XI,51.
G. tenella; R. Brown, prodr. 577 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,74	M.fr.XI,51.
G. elongata; Labillardière, Nov. Holl. pl. spec. I, 52, t. 75 (1804)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.IV,74	M.fr.VI,14; VIII,142,245; X,110.
G. pinnatifida; Schlechtendal, Linnaea XXI, 440 (1848)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.IV,75	M.fr.VI,14; XI,51.
G. pusilliflora; F. v. M., Key to the Syst. of Vict. pl. I, 354 (1888)	—	S.A.	—	V.	N.S.W.	Q.	—	—	
G. tenuiloba; F. v. M. in Wing's S. Sc. Rec. new ser. I, 25 (1885)	W.A.	—	—	—	—	—	—	—	
G. coronopifolia; R. Brown, prodr. 577 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,75	M.fr.X,110.
G. heteromera; F. v. M., Sec. Rep. 13 (1854)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.IV,76	M.fr.I,115; VI,14; XI,27.
G. concinna; Benth. Fl. Austr. IV, 76 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,76	
G. glauca; F. v. M. in Transact. Vict. Inst. 40 (1855)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.IV,76	M.fr.VI,14; XI,27.
G. filiformis; R. Brown, prodr. 578 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,77	M.fr.I,116; VI,14; VIII,245.
G. Armitiana; F. v. M., fragm. X, 110 (1877)	—	—	—	—	—	Q.	N.A.	—	M.fr.X,110.
G. cirrifica; F. v. M. in Austr. Journ. of Pharm. I, 81 (1886)	—	—	—	—	—	—	N.A.	—	
G. microptera; F. v. M., fragm. III, 34 (1862)	—	S.A.	—	—	—	—	N.A.	B.fl.IV,77	M.fr.VI,227; XI,75.
G. bicolor; F. v. M. in Benth. Fl. Austr. IV, 80 (1869)	—	—	—	—	—	—	N.A.	B.fl.IV,80	
G. humilis; R. Brown, prodr. 575 (1810)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.IV,79	M.fr.VII,227; VIII,142,245; X,110.
G. paniculata; Smith in Transact. Linn. Soc. II, 348 (1794)	—	—	—	V.	N.S.W.	Q.	—	B.fl.IV,78	M.fr.I,116.
G. purpurascens; R. Brown, prodr. 578 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,78	M.fr.I,117.
G. gracilis; R. Brown, prodr. 575 (1810)	—	—	—	V.	N.S.W.	Q.	N.A.	B.fl.IV,79	M.fr.XI,51.
G. lamprosperma; F. v. M., fragm. I, 116 (1859)	W.A.	—	—	—	—	—	N.A.	B.fl.IV,79	M.fr.III,34.
G. minutiflora; F. v. M., fragm. VIII, 244 (1874)	—	—	—	—	—	—	N.A.	—	M.fr.VIII,244.
G. claytoniacea; F. v. M. in Benth. Fl. Austr. IV, 79 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,79	M.fr.II,111.
G. Pumilio; R. Brown, prodr. 579 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,80	M.fr.X,111.

VELLEYA; Smith in Transact. Linn. Soc. IV, 217 (1798). (Velleia, Euthales.)

V. Daviesii; F. v. M., fragm. X, 10 (1876) W.A. — — — — — M.fr.X,10,,54.

V. panduriformis; Cunningham in Benth. Fl. Austr. IV, 46 (1869) ...	...	...	—	—	—	—	—	N.A.	B.fl.IV,46	M.fr.XI,73.
V. connata; F. v. M. in Transact. phil. Soc. Vict. I, 18 (1855) ...	...	...	W.A.	S.A.	—	V.	N.S.W.	—	N.A.	B.fl.IV,46 M.fr.VI,227; VIII,246.
V. perfoliata; R. Brown, prodr. 581 (1810) ...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.IV,46
V. discophora; F. v. M., fragm. X, 10 (1876) ...	...	...	W.A.	—	—	—	—	—	—	M.fr.X,10.
V. trinervis; Labillardière, Nov. Holl. pl. spec. I, 54, t. 77 (1804) ...	...	...	W.A.	—	—	—	—	—	B.fl.IV,47	M.fr.XI,75.
V. macrophylla; Benth. Fl. Austr. IV, 47 (1869) ...	...	...	W.A.	—	—	—	—	—	B.fl.IV,47	M.fr.XI,75.
V. montana; J. Hooker in Lond. Journ. of Bot. VI, 265 (1847) ...	...	...	—	—	T.	V.	N.S.W.	—	—	B.fl.IV,50
V. pubescens; R. Brown, prodr. 581 (1810) ...	...	...	—	—	—	—	—	Q.	—	B.fl.IV,49 M.fr.VI,11; XI,75.
V. spathulata; R. Brown, prodr. 580 (1810) ...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,50 M.fr.XI,75.
V. macrocalyx; De Vriese, Goodenov. 176, t. 34 (1854) ...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,49 M.fr.X,75.
V. lyrata; R. Brown, prodr. 580 (1810) ...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.IV,49
V. cynopotamica; F. v. M., fragm. VI, 7 (1867) ...	...	...	W.A.	—	—	—	—	—	B.fl.IV,48	M.fr.VI,249; X,10.
V. paradoxa; R. Brown, prodr. 580 (1810) ...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,48 M.fr.VI,11; XI,75; XII,23.
V. macropectra; F. v. M., fragm. XII, 22 (1882) ...	...	...	W.A.	—	—	—	—	—	—	—

SYNPETALEAE HYPOGYNAE.

F. v. M. in Woolls' pl. of the neighb. of Sydney, 34 (1880).

GENTIANEAEE.

NECKER in Act. Acad. Theod. Pal. II, 477 (1770), from B. de Jussieu (1759).—Genera 5, Species 24.

LIMNANTHEMUM; Gmelin, Nov. Comm. Acad. Petrop. XIV, 257 (1770). (Villarsia, Liparophyllum.)

L. exaltatum; F. v. M., fragm. IX, 165 (1875) ...	...	...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,376 M.fr.II,21; VI,139,254.
L. parnassifolium; F. v. M., fragm. IX, 165 (1875) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,376 M.fr.VI,140.
L. violifolium; F. v. M., Cens. 90 (1882) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,377 M.fr.VI,138.
L. lasiospermum; F. v. M., fragm. IX, 165 (1875) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,377 M.fr.VI,137.
L. albiflorum; F. v. M., Cens. 90 (1882) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,377 M.fr.VI,138.
L. latifolium; F. v. M., fragm. IX, 164 (1875) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,375 M.fr.VI,141.
L. calthifolium; F. v. M., fragm. IX, 164 (1875) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,374 M.fr.VI,140.
L. congestiflorum; F. v. M., Cens. 90 (1882) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,375 M.fr.VI,141.
L. capitatum; F. v. M., fragm. IX, 164 (1875) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,375 M.fr.VI,142.
L. Gunnii; J. Hooker, Fl. Tasman. II, 368 (1860) implied ...	...	...	—	—	T.	—	—	—	—	B.fl.IV,382 M.fr.VI,137; IX,164.
L. exiguum; F. v. M., fragm. I, 40 (1858) ...	...	...	—	—	T.	—	—	—	—	B.fl.IV,381 M.fr.VI,137.
L. exiliflorum; F. v. M., fragm. V, 46 (1865) ...	...	...	—	—	—	—	—	Q.	—	B.fl.IV,381 M.fr.VI,137.
L. hydrocharoides; F. v. M. in Benth. Fl. Austr. IV, 380 (1869) ...	...	...	—	—	—	—	—	Q.	—	B.fl.IV,380 M.fr.VI,139; IX,165.
L. geminatum; Grisebach, gen. et spec. Gent. 346 (1839) ...	...	...	—	—	—	V.	N.S.W.	Q.	N.A.	B.fl.IV,380 M.fr.VI,137,254.
L. crenatum; F. v. M. in Transact. phil. Soc. Vict. I, 17 (1854) ...	...	...	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.IV,379 M.fr.IV,127; VI,137; IX,165.
L. minimum; F. v. M., fragm. I, 40 (1858) ...	...	...	—	—	—	—	—	—	N.A.	B.fl.IV,379 M.fr.VI,137; IX,164.
L. Moonii; Thwaites, enum. pl. Ceyl. 205 (1862) ...	...	...	—	—	—	—	—	Q.	—	M.fr.IX,163 ... As., P.
L. Indicum; Thwaites, enum. pl. Ceyl. 205 (1862) ...	...	...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.IV,378 M.fr.IV,128; VII,136; IX,164.

SEBAEA; Solander in R. Brown, prodr. 451 (1810).

S. ovata; R. Brown, prodr. 452 (1810) ...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,371 M.fr.VI,136; IX,165 ... N.Z.
S. albidiflora; F. v. M. in Transact. phil. Soc. Vict. I, 46 (1854) ...	...	...	—	S.A.	T.	V.	—	—	—	B.fl.IV,371 M.fr.VI,136.

ERYTHRAEA; Reneaulme, specim. hist. plant. 77, t. 76 (1611).

E. spicata; Persoon, syn. plant. I, 283 (1805) ...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.IV,371 M.fr.VI,136; IX,165; XI,20. E., As., Af., P.
--	-----	-----	------	------	----	----	--------	----	------	--

CANSORA; Lamarck, Encycl. method. I, 602 (1783). (Orthostemon.)														
C. diffusa; R. Brown, prodr. 451 (1810)	...	...	...	...	...	—	—	—	—	—	Q. N.A.	B.fl.IV,372	M.fr.VI,136; IX,165	... As., Afr., P.
GENTIANA; Tournefort, inst. 80, t. 40 (1700), from Dioscorides.														
G. quadrifaria; Blume, Bijdr. 804 (1825)	...	...	...	...	...	—	—	—	—	V. N.S.W.	—	—	...	As.
G. saxosa; Forster in Svensk. Kongl. Vet. Ac. Handl. 183, t. 5 (1777)	...	...	—	S.A. T.	V. N.S.W.	—	—	B.fl.IV,373	M.fr.VI,136,255; IX,165	..	Am, N.Z.			

LOGANIACEAE.

R. BROWN in App. Flind. Voy. 564 (1814).—Genera 7, Species 52.

STRYCHNOS; Linné, syst. nat. 8 (1735); spec. plant. 189 (1753).														
S. lucida; R. Brown, prodr. 469 (1810)	...	...	...	...	...	—	—	—	—	—	Q. N.A.	B.fl.IV,369	M.fr.IV,44	... As.
S. pilosperma; F. v. M., fragm. IV, 44 (1863)	...	...	...	...	...	—	—	—	—	—	Q. —	B.fl.IV,369	M.fr.VI,131	
FAGRAEA; Thunberg in Kongl. Svensk. Vetensk. Acad. Handl. 125 (1782).														
F. racemosa; Jack in Roxb. Fl. Ind. ed. Carey. II, 35 (1825)	...	...	...	...	...	—	—	—	—	—	Q. N.A.	B.fl.IV,367	M.fr.II,137	... As.
GARDNERIA; Wallich in Roxb. Fl. Indic. I, 400 (1820). (Loranthophyllum.)														
G. fagraeacea; F. v. M., fragm. VI, 130 (1868)	...	...	...	...	...	—	—	—	—	—	Q. —	B.fl.IV,368	M.fr.VI,130.	
MITREOLA; Linné, hort. Cliffort. 492 (1737).														
M. oldenlandioides; Wallich, numeric list. 4350 (1828)	...	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.IV,349	M.fr.I,130; VI,131. As.
MITRASACME; Labillardière, Nov. Holl. pl. spec. I, 36, t. 49 (1804). (Mitragyne.)														
M. Archeri; J. Hooker, Fl. Tasman. II, 368 (1860)	...	...	...	...	...	—	—	T. —	—	—	—	B.fl.IV,351		
M. montana; J. Hooker, Fl. Tasman. I, 274, t. 88 (1859)	...	...	...	...	...	—	—	T. V.	—	—	—	B.fl.IV,351	M.fr.VI,131.	
M. serpillifolia; R. Brown, prodr. 454 (1810)	...	...	...	...	...	—	—	T. V.	N.S.W.	—	—	B.fl.IV,352	M.fr.I,130.	
M. paludosa; R. Brown, prodr. 453 (1810)	...	...	...	...	...	—	—	—	N.S.W.	Q. —	—	B.fl.IV,352	M.fr.I,134.	
M. pilosa; Labillardière, Nov. Holl. pl. spec. I, 36, t. 49 (1804)	...	...	...	...	...	—	S.A. T.	V. N.S.W.	—	—	—	B.fl.IV,352	M.fr.I,130; VI,131.	
M. alsinoides; R. Brown, prodr. 453 (1810)	...	...	...	...	...	—	—	—	N.S.W.	Q. —	—	B.fl.IV,353		
M. polymorpha; R. Brown, prodr. 452 (1810)	...	...	...	...	...	—	—	—	V. N.S.W.	Q. —	—	B.fl.IV,353	M.fr.VI,131.	
M. longiflora; F. v. M. in Benth. Fl. Austr. IV, 353 (1869)	...	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.IV,353	
M. elata; R. Brown, prodr. 453 (1810)	...	...	...	...	...	—	—	—	—	—	—	Q. N.A.	B.fl.IV,354	M.fr.I,132.
M. exserta; F. v. M., fragm. I, 131 (1859)	...	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.IV,354	M.fr.I,131.
M. tenuiflora; Benth. Fl. Austr. IV, 354 (1869)	...	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.IV,354	
M. ambigua; R. Brown, prodr. 454 (1810)	...	...	...	...	...	—	—	—	—	—	—	Q. —	B.fl.IV,354	
M. nudicaulis; Reinwardt in Blume, Bijdr. 849 (1826)	...	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.IV,355	M.fr.I,132. As.
M. connata; R. Brown, prodr. 454 (1810)	...	...	...	...	...	—	—	—	—	—	—	Q. N.A.	B.fl.IV,355	M.fr.I,131.
M. laevis; Benth. in Jour. Linn. Soc. I, 93 (1857)	...	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.IV,355	M.fr.I,133.
M. Indica; Wight, icon. pl. Ind. or. IV, t. 1601 (1850)	...	...	...	...	...	—	—	—	—	N.S.W.	Q. N.A.	B.fl.IV,356	...	As.
M. stellata; R. Brown, prodr. 454 (1810)	...	...	...	...	...	—	—	—	—	—	—	Q. N.A.	B.fl.IV,356	
M. Cunninghamii; Benth. Fl. Austr. IV, 357 (1869)	...	...	...	...	...	—	—	—	—	—	—	Q. —	B.fl.IV,357	
M. pygmaea; R. Brown, prodr. 453 (1810)	...	...	...	...	...	—	—	—	—	—	—	Q. —	B.fl.IV,357	
M. lutea; F. v. M., fragm. I, 133 (1859)	...	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.IV,357	M.fr.IV,170.
M. multicaulis; R. Brown, prodr. 453 (1810)	...	...	...	...	...	—	—	—	—	—	—	Q. N.A.	B.fl.IV,357	
M. laricifolia; R. Brown, prodr. 453 (1810)	...	...	...	...	...	—	—	—	—	—	—	Q. N.A.	B.fl.IV,358	M.fr.VI,131.
M. prolifera; R. Brown, prodr. 453 (1810)	...	...	...	...	...	—	—	—	—	—	—	Q. N.A.	B.fl.IV,358	
M. gentianea; F. v. M., fragm. I, 130 (1859)	...	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.IV,358	M.fr.I,130.
M. phascoides; R. Brown, prodr. 454 (1810)	...	...	...	...	...	—	—	—	—	—	—	Q. N.A.	B.fl.IV,359	
M. paradoxa; R. Brown, prodr. 454 (1810)	...	...	...	...	...	...	W.A. S.A. T.	V. N.S.W.	—	—	—	B.fl.IV,359	M.fr.I,130; VI,131.	
M. distylis; F. v. M. in Transact. phil. Soc. Vict. I, 20 (1854)	...	...	...	...	...	...	S.A. T.	V.	—	—	—	B.fl.IV,359	M.fr.VI,131.	

GENIOSTOMA ; R. & G. Forster, char. gen. 23, t. 12 (1776).

G. Australianum ; F. v. M., fragm. V, 19 (1865)	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl.IV,367	M.fr.V,19.
G. petiolosum ; Moore & F. v. M., fragm. VII, 28 (1869)	...	...	...	...	...	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,193.

LOGANIA ; R. Brown, prodr. 454 (1810). (Euosma 1808, restorable.)

L. longifolia ; R. Brown, prodr. 456 (1810)	...	...	...	...	...	W.A.	S.A.	—	—	—	—	—	B.fl.IV,361	M.fr.VI,132.
L. latifolia ; R. Brown, prodr. 455 (1810)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,361	M.fr.VI,132.
L. crassifolia ; R. Brown, prodr. 455 (1810)	...	...	...	...	...	W.A.	S.A.	—	—	—	—	—	B.fl.IV,362	M.fr.VI,132.
L. ovata ; R. Brown, prodr. 455 (1810)	...	...	...	...	...	W.A.	S.A.	—	V.	—	—	—	B.fl.IV,362	M.fr.VI,132.
L. buxifolia ; F. v. M., fragm. VI, 132 (1868)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,362	M.fr.VI,132.
L. stenophylla ; F. v. M., fragm. I, 128 (1859)	...	...	...	...	...	W.A.	S.A.	—	—	—	—	—	B.fl.IV,362	M.fr.I,128.
L. micrantha ; Bentham in Journ. Linn. Soc. I, 94 (1857)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,363	M.fr.I,129; VI,132.
L. linifolia ; Schlechtendal, Linnaea XX, 605 (1847)...	...	...	...	...	...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.IV,363	M.fr.I,129; VI,131.
L. fasciculata ; R. Brown, prodr. 456 (1810)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,363	M.fr.I,129; VI,131.
L. floribunda ; R. Brown, prodr. 456 (1810)	...	...	...	...	...	—	—	—	V.	N.S.W.	Q.	—	B.fl.IV,364	M.fr.VI,131.
L. cordifolia ; Hooker in Mitch. Trop. Austr. 341 (1848)	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl.IV,364	M.fr.VI,132.
L. nuda ; F. v. M., fragm. I, 129 (1859)	...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.IV,365	M.fr.II,163; VI,131.
L. spermacoea ; F. v. M., fragm. VI, 134 (1868)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,365	M.fr.VI,134.
L. flaviflora ; F. v. M. in Vict. Naturalist V, 165 (1889)	...	...	...	...	...	W.A.	—	—	—	—	—	—	—	—
L. callosa ; F. v. M., fragm. VI, 134 (1868)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,365	M.fr.VI,134.
L. campanulata ; R. Brown, prodr. 456 (1810)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,365	M.fr.I,130; VI,133.
L. serpillifolia ; R. Brown, prodr. 456 (1810)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,366	M.fr.I,130; VI,133.
L. pusilla ; R. Brown, prodr. 456 (1810)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,366	M.fr.VI,131.

PLANTAGINEAE.

A. L. DE JUSSIEU, Gen. plant. 89 (1789).—Genus 1, Species 4.

PLANTAGO ; Tournefort, inst. 126, t. 48 (1700), from l'Ecluse (1576), from W. Turner (1538).

P. varia ; R. Brown, prodr. 424 (1810)...	...	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.V,139	M.fr.XI,20.
P. stellaris ; F. v. M., fragm. II, 23 (1860)	...	...	...	...	...	—	—	—	V.	N.S.W.	—	—	—	M.fr.VIII,148.
P. Brownii ; Rapin in Mém. de la Soc. Linn. Par. VI, 484 (1827)	...	...	...	...	...	—	—	T.	—	—	—	—	B.fl.V,141	M.fr.II,23 ... N.Z.
P. Gunnii ; J. Hooker in Lond. Journ. V. 446, t. 13 (1846)	...	...	...	...	...	—	—	T.	V.	N.S.W.	—	—	B.fl.V,142	M.fr.VIII,148; XI,134.

PRIMULACEAE.

VENTENAT, Tabl. du régn. végét. II, 285 (1799).—Genera 3, Species 6.

ANAGALLIS ; Tournefort, inst. 142, t. 59 (1700) from Hippocrates and Dioscorides. (Centunculus, Micropyxis.)

A. pumila ; Swartz, Fl. Ind. occ. I, 345 (1797)	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.IV,270	...	...	...	As., Afr., Am.
A. Centunculus ; Afzelius, de Vegetab. Suecan. (1785)	...	...	...	...	...	—	S.A.	—	V.	—	—	—	—	...	...	...	E., As., Afr., Am.

LYSIMACHIA ; Tournefort, inst. 141, t. 59 (1700) from Dioscorides.

L. salicifolia ; F. v. M. in Benth. Fl. Austr. IV, 269 (1869)	...	...	...	...	...	—	—	—	V.	N.S.W.	—	—	B.fl.IV,269	...	...	...	...
L. Japonica ; Thunberg, Fl. Japon. 83 (1784)...	...	...	...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.IV,269	...	...	...	As.

SAMOLUS ; Tournefort, inst. 143, t. 60 (1700). (Sheffieldia.)

S. Valerandi ; Linné, spec. plant. 171 (1753)	...	...	...	...	...	—	—	—	V.	N.S.W.	Q.	—	B.fl.IV,271	...	...	...	E., As., Afr., Am.
S. repens ; Persoon, synopsis. plant. I, 171 (1805)	...	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.IV,271	M.fr.XI,20	...	...	Afr., Am., P., N.Z.

MYRSINACEAE.

R. BROWN, prodr. 532 (1810). (Primulacearum subordo).—Genera 5, Species 12.

MAESA; Forskael, Fl. Aegypt. Arab. 66 (1775). (Baeobotrys.)

M. dependens; F. v. M., fragm. V, 107 (1865)	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl. IV, 273	M.fr. V, 107.
M. haplobotrys; F. v. M., fragm. V, 161 (1866)	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl. IV, 273	M.fr. V, 161.

SAMARA; Linné, mantiss. II, 144 (1771). (Choripetalum.)

S. Australiana; F. v. M., fragm. VI, 164 (1868)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl. IV, 274	M.fr. III, 36; IV, 82, 174.
---	-----	-----	-----	-----	-----	---	---	---	---	--------	----	---	---------------	-----------------------------

MYRSINE; Linné, syst. nat. 8 (1735); Linné, gen. plant. 54 (1737).

M. urceolata; R. Brown, prodr. 534 (1810)	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl. IV, 274	
M. campanulata; F. v. M., fragm. VI, 235 (1868)	...	...	...	...	...	—	—	—	—	—	Q.	—	—	M.fr. VI, 235.
M. crassifolia; R. Brown, prodr. 534 (1810)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl. IV, 275	M.fr. IV, 81; V, 108; VI, 164; VIII, 48.
M. platystigma; F. v. M., fragm. VIII, 48 (1873)	...	...	...	...	...	—	—	—	—	N.S.W.	—	—	—	M.fr. VIII, 48.
M. variabilis; R. Brown, prodr. 534 (1810)	...	...	...	...	...	—	—	—	—	V. N.S.W.	Q.	—	B.fl. IV, 275	M.fr. IV, 81.
M. achradifolia; F. v. M., fragm. VI, 164 (1868)	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl. IV, 275	M.fr. VI, 164.

ARDISIA; Swartz, nov. gen. et spec. 3 et 48 (1788). (Labisia partly.)

A. Pseudo-Jambosa; F. v. M., fragm. IV, 81 (1864)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl. IV, 276	M.fr. IV, 81.
A. brevipedata; F. v. M., fragm. VI, 163 (1868)	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl. IV, 276	M.fr. VI, 163.

AEGICERAS; Gaertner, de fruct. I, 216, t. 46 (1788).

A. majus; Gaertner, de fruct. I, 216, t. 46 (1788)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl. IV, 277	 As., P.
--	-----	-----	-----	-----	-----	---	---	---	---	--------	----	------	---------------	----------------

SAPOTACEAE.

A. L. DE JUSSIEU, Gen. plant. 151 (1789), from B. de Jussieu (1759).—Genera 6, Species 19.

NIEMEYERA; F. v. M., fragm. VII, 114 (1870). (Chrysophyllum partly.)

N. prunifera; F. v. M., fragm. VII, 114 (1870)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl. IV, 278	M.fr. VI, 26.
--	-----	-----	-----	-----	-----	---	---	---	---	--------	----	---	---------------	---------------

AMORPHOSPERMUM; F. v. M., fragm. VII, 112 (1870). (Sersalisia partly.)

A. antilogum; F. v. M., fragm. VII, 113 (1870)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	—	M.fr. X, 119.
--	-----	-----	-----	-----	-----	---	---	---	---	--------	----	---	---	---------------

LUCUMA; Molina, Saggio 186 (1782). (Sersalisia partly.)

L. sericea; Benth & J. Hooker, gen. pl. II, 654 (1876)	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl. IV, 279	M.fr. V, 162; VII, 112.
L. Galactoxylon; Benth & J. Hooker, gen. pl. II, 654 (1876)	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl. IV, 279	

SIDEROXYLON; Dillenius, hort. Eltham. 357, t. 265 (1732). (Achras partly, Sapota partly, Sersalisia partly.)

S. Arnemicum; Benth & J. Hooker, gen. pl. II, 655 (1876)	...	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl. IV, 280	
S. Pohlmanianum; Benth & J. Hooker, gen. pl. II, 655 (1876)	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl. IV, 281	M.fr. V, 184.
S. xerocarpum; Benth & J. Hooker, gen. pl. II, 655 (1876)	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl. IV, 281	
S. chartaceum; Benth & J. Hooker, gen. pl. II, 655 (1876)	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl. IV, 281	
S. Richardi; F. v. M., Cens. of Austr. pl. 92 (1882)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl. IV, 282	
S. Australe; Benth & J. Hooker, gen. pl. II, 655 (1876)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl. IV, 282	M.fr. V, 162; IX, 73.
S. Brownii; F. v. M., Cens. of Austr. pl. 92 (1882)	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl. IV, 283	
S. myrsinoides; Benth & J. Hooker, gen. pl. II, 655 (1876)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl. IV, 283	M.fr. IX, 73.
S. euphlebiun; F. v. M., Cens. of Austr. pl. 92 (1882)	...	...	...	...	...	—	—	—	—	—	Q.	—	—	M.fr. VII, 110.
S. Brownlessianum; F. v. M., fragm. VIII, 111 (1870)	...	...	...	...	...	—	—	—	—	—	Q.	—	—	M.fr. VII, 111.
S. Howeanum; F. v. M., Cens. of Austr. pl. 92 (1882)	...	...	...	...	...	—	—	—	—	N.S.W.	—	—	—	M.fr. IX, 72.
S. costatum; F. v. M., Cens. 92 (1882) from Endlicher (1833)	...	...	...	...	...	—	—	—	—	N.S.W.	—	—	—	M.fr. IX, 72.

HORMOGYNE; A. de Candolle, prodr. VIII, 176 (1844).

H. cotinifolia; A. De Candolle, prodr. VIII, 176 (1844)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl. IV, 284	M.fr. V, 161; VII, 111.
---	-----	-----	-----	-----	-----	---	---	---	---	--------	----	---	---------------	-------------------------

MIMUSOPS; Linné, Fl. Zeil. 57 (1747).

M. parvifolia; R. Brown, prodr. 531 (1810)	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl. IV, 284	M.fr. V, 162; VI, 178; VII, 114.
M. Kauki; Linné, spec. plant. 349 (1753)	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl. IV, 285	M.fr. V, 162 As.

EBENACEAE.

VENTENAT, Tabl. II, 443 (1799).—Genera 2, Species 15.

DIOSPYROS; Linné, gen. plant. 143 (1737). (Cargillia.)

D. cordifolia; Roxburgh, pl. Corom. I, 38, t. 50 (1795)	—	—	—	—	—	—	N.A.	B.fl.IV, 286	...	...	...	As.
D. hebecarpa; Cunningham in Benth. Fl. Austr. IV, 286 (1869)	—	—	—	—	—	—	Q.	N.A.	B.fl.IV, 286			
D. maritima; Blume, Bijdrag. 669 (1825)	—	—	—	—	—	—	Q.	N.A.	B.fl.IV, 287	M.fr.V, 163; X, 75	...	As.
D. mabacea; F. v. M., Docum. intercol. Exhib. 35 (1866)	—	—	—	—	—	N.S.W.	Q.	—	B.fl.IV, 288	M.fr.V, 162; X, 75.		
D. Cargillia; F. v. M., Docum. intercol. Exhib. 35 (1866)	—	—	—	—	—	N.S.W.	Q.	—	B.fl.IV, 288	M.fr.IV, 82; V, 162; X, 74.		
D. pentanera; F. v. M. & Woolls, Doc. intercol. Exhib. 35 (1866)	—	—	—	—	—	N.S.W.	Q.	—	B.fl.IV, 288	M.fr.IV, 82; V, 163; X, 75.		

MABA; R. & G. Forster, char. gen. 121, t. 61 (1776).

M. laurina; R. Brown, prodr. 527 (1810)	—	—	—	—	—	—	Q.	—	B.fl.IV, 289			
M. rufa; Labillardière, sert. Austr. Caled. 33, t. 36 (1824)	—	—	—	—	—	—	Q.	—	B.fl.IV, 289	M.fr.V, 164; VI, 253; X, 74	...	P.
M. hemicycloides; F. v. M. in Benth. Fl. Austr. IV, 290 (1869)	—	—	—	—	—	—	Q.	—	B.fl.IV, 290			
M. fasciculosa; F. v. M., fragm. V, 163 (1866)	—	—	—	—	—	N.S.W.	Q.	—	B.fl.IV, 290	M.fr.X, 74.		
M. compacta; R. Brown, prodr. 528 (1810)	—	—	—	—	—	—	Q.	N.A.	B.fl.IV, 290	M.fr.V, 165.		
M. reticulata; R. Brown, prodr. 528 (1810)	—	—	—	—	—	—	Q.	N.A.	B.fl.IV, 291			
M. geminata; R. Brown, prodr. 527 (1810)	—	—	—	—	—	—	Q.	N.A.	B.fl.IV, 291	M.fr.V, 164.		
M. humilis; R. Brown, prodr. 527 (1810)	—	—	—	—	—	—	Q.	N.A.	B.fl.IV, 291	M.fr.V, 164; X, 74.		
M. buxifolia; Persoon, synops. pl. II, 602 (1807)	—	—	—	—	—	—	—	N.A.	—	M.fr.X, 74	...	As., Afr.

AQUIFOLIACEAE.

DE CANDOLLE, Théor. élém. I, 217 (1813).—Genera 2, Species 2.

BRYONIA; Endlicher in den Ana. des Wien. Mus. I, 184 (1836).

B. Arnhemensis; F. v. M., fragm. II, 119 (1861)	—	—	—	—	—	—	—	N.A.	B.fl.I, 397	M.fr.II, 119.		
--	---	---	---	---	---	---	---	------	-------------	---------------	--	--

ILEX; Linné, gen. plant. 33 (1737) from C. Bauhin (1623).

I. peduncularis; F. v. M., fragm. VII, 105 (1871)	—	—	—	—	—	—	Q.	—	—	M.fr.VII, 105.		
--	---	---	---	---	---	---	----	---	---	----------------	--	--

STYRACEAE.

L. C. RICHARD, Analyse du fruit, 48 (1808).—Genus 1, Species 2.

SYMPLOCOS; N. J. Jacquin, ennm. plant. Carib. 24 (1760).

S. Stawellii; F. v. M., fragm. V, 60 (1865)	—	—	—	—	—	N.S.W.	Q.	—	B.fl.IV, 292	M.fr.V, 60.		
S. Thwaitesii; F. v. M., fragm. III, 22 (1862)	—	—	—	—	—	N.S.W.	Q.	—	B.fl.IV, 293	M.fr.V, 211.		

JASMINEAE.

A. L. DE JUSSIEU, Gen. plant. 104 (1789) from Necker (1770).—Genera 5, Species 21.

JASMINUM; Tournefort, inst. 597, t. 368 (1700), from l'Ecluse (1611), from W. Turner (1538).

J. didymum; G. Forster, prodr. 3 (1786)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.IV, 294	M.fr.IV, 150; VI, 87; VIII, 43.	As., P.
J. racemosum; F. v. M., fragm. I, 19 (1858)	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl.IV, 295	M.fr.VI, 87; VIII, 43	
J. lineare; R. Brown, prodr. 521 (1810)	...	...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.IV, 295	M.fr.VI, 87; VIII, 43.
J. simplicifolium; G. Forster, prodr. 3 (1786)	...	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.IV, 296	M.fr.VI, 87; VIII, 43 ... As., P.
J. aemulum; R. Brown, prodr. 521 (1810)	...	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.IV, 296	M.fr.VI, 86.
J. calcareum; F. v. M., fragm. I, 212 (1859)	...	...	...	...	...	...	W.A.	S.A.	—	—	—	—	N.A.	B.fl.IV, 297	M.fr.VI, 87; VIII, 43.
J. suavisimum; Lindley in Mitch. Trop. Austr. 355 (1848)	...	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV, 297	M.fr.I, 183; II, 181; VI, 87.
J. singuliflorum; F. v. M. & Bailey in suppl. to Queensl. Fl. 38 (1886)	...	...	...	...	...	...	—	—	—	—	—	Q.	—	—	

OLEA; Tournefort, inst. 598, t. 370 (1700) from Plinius.

O. paniculata; R. Brown, prodr. 523 (1810)	—	—	—	—	—	N.S.W.	Q.	—	B.fl.IV, 297	M.fr.IV, 83; VIII, 43	...	P.
O. Endlicheri; F. v. M., fragm. VIII, 43 (1872)	—	—	—	—	—	N.S.W.	—	—	—	M.fr.VIII, 43; IX, 169.		

LIGUSTRUM ; Tournefort. inst. 596, t. 367 (1700) from C. Bauhin (1623).

L. Australianum ; F. v. M., fragm. V, 20 (1865) — — — — — Q. — B.fl.IV, 298 M.fr. V, 20.

NOTELAEA ; Ventenat, Choix des pl. dans le jard. de Cels. t. 25 (1803).

N. ovata ; R. Brown, prodr. 524 (1810) — — — — — N.S.W. Q. — B.fl.IV, 299 M.fr. VIII, 43.
 N. longifolia ; Ventenat, Choix t. 25 (1803) — — — — — V. N.S.W. Q. — B.fl.IV, 299 M.fr. VIII, 43.
 N. punctata ; R. Brown, prodr. 524 (1810) — — — — — Q. — B.fl.IV, 300
 N. microcarpa ; R. Brown, prodr. 524 (1810) — — — — — N.S.W. Q. — B.fl.IV, 300 M.fr. VIII, 43.
 N. ligustrina ; Ventenat, Choix t. 25 (1803) — — — — — T. V. N.S.W. — — B.fl.IV, 300 M.fr. VIII, 43, 142.
 N. linearis ; Benth. Fl. Austr. IV, 300 (1869) — — — — — N.S.W. Q. — B.fl.IV, 300

MAYEPEA ; Aublet, Hist. des pl. de la Guian. I, 81, t. 31 (1775). (Chionanthus partly, Linociera partly, Ceranthes.)

M. ramiflora ; F. v. M., Cens. 92 (1882) from Roxburgh (1820) — — — — — Q. — B.fl.IV, 301 M.fr. IV, 83; VIII, 42.
 M. picophloia ; F. v. M., Cens. of Austr. plant. 92 (1882) — — — — — Q. — B.fl.IV, 301 M.fr. III, 139, 24; IV, 176.
 M. axillaris ; F. v. M., Cens. 92 (1882) from R. Brown (1810) — — — — — Q. N.A. B.fl.IV, 301 M.fr. VIII, 42.
 M. quadristaminea ; F. v. M., fragm. X, 89 (1876) — — — — — N.S.W. — — — M.fr. VIII, 41.

APOCYNEAE.

A. L. DE JUSSIEU, Gen. plant. 143 (1789).—Genera 13, Species 47.

CHILOCARPUS ; Blume, catal. hort. Buitenz. (1823).

C. Australis ; F. v. M., fragm. II, 90 (1861) — — — — — N.S.W. Q. — B.fl.IV, 303 M.fr. VI, 118.

MELODINUS ; R. & G. Forster, char. gen. 37, t. 19 (1776).

M. acutiflorus ; F. v. M. in Transact. phil. Soc. Vict. II, 71 (1857) — — — — — Q. — B.fl.IV, 304
 M. Guilfoylei ; F. v. M., fragm. VI, 118 (1868) — — — — — Q. — B.fl.IV, 304 M.fr. VI, 118.
 M. Baueri ; Endlicher, prodr. pl. Norfolk. 59 (1833) — — — — — N.S.W. — — — M.fr. IX, 169.

CARISSA ; Linné, mantiss. pl. 7 (1767).

C. laxiflora ; Benth. Fl. Austr. IV, 305 (1869) — — — — — Q. — B.fl.IV, 305
 C. Brownii ; F. v. M., fragm. IV, 45 (1863) — — — — — S.A. — — N.S.W. Q. N.A. B.fl.IV, 305 M.fr. IV, 45.

CERBERA ; Linné, gen. plant. 62 (1737).

C. Odollam ; Gaertner, de fruct. 193, t. 124 (1791) — — — — — Q. N.A. B.fl.IV, 306 M.fr. VI, 117 ... As., P.

ALYXIA ; Banks in R. Brown, prodr. 469 (1810). (Gynopogon, 1776.)

A. buxifolia ; R. Brown, prodr. 470 (1810) W.A. S.A. T. V. N.S.W. — — B.fl.IV, 307 M.fr. VI, 117; VII, 131.
 A. ruscifolia ; R. Brown, prodr. 470 (1810) — — — — — N.S.W. Q. — B.fl.IV, 308 M.fr. VI, 117; VII, 131. P.
 A. ilicifolia ; F. v. M., fragm. IV, 149 (1864) — — — — — Q. — B.fl.IV, 308 M.fr. IV, 149.
 A. squamulosa ; C. Moore & F. v. M., fragm. VIII, 47 (1873) — — — — — N.S.W. — — — M.fr. VI, 117; VIII, 47.
 A. obtusifolia ; R. Brown, prodr. 470 (1810) — — — — — Q. — B.fl.IV, 308
 A. Gynopogon ; Roemer & Schultes, syst. veg. IV, 440 (1819) — — — — — N.S.W. — — — M.fr. VIII, 47.
 A. spicata ; R. Brown, prodr. 470 (1810) — — — — — Q. N.A. B.fl.IV, 308 M.fr. VI, 117.
 A. Lindii ; F. v. M., fragm. VIII, 46 (1873) — — — — — N.S.W. — — — M.fr. VIII, 46.
 A. thyrsoiflora ; Benth. Fl. Austr. IV, 309 (1869) — — — — — Q. — B.fl.IV, 309
 A. Thozetii ; F. v. M., fragm. X, 103 (1877) — — — — — Q. — — M.fr. X, 103.

OCHROSIA ; A. L. de Jussieu, gen. plant. 143 (1789). (Lactaria.)

O. elliptica ; Labillardière, eert. Austr. Caled. 27, t. 30 (1824) — — — — — Q. — B.fl.IV, 310 M.fr. III, 110; VI, 118. As., P.
 O. Moorei ; F. v. M. in Benth. Fl. Austr. IV, 310 (1869) — — — — — N.S.W. Q. — B.fl.IV, 310 M.fr. III, 110; VI, 118; VII, 131.
 O. Kilneri ; F. v. M., fragm. VII, 129 (1871) — — — — — Q. — — M.fr. VIII, 129.

NOTONERIUM ; Bentham in B. & J. H. gen. II, 698 (1876).

N. Gossei ; Bentham in B. & J. H. gen. pl. 698 (1876) ... — S.A. — — — — — M.fr.X,103.

TABERNAEMONTANA ; Plumier, nov. pl. Amer. gen. 18, t. 30 (1703).

T. orientalis ; R. Brown, prodr. 468 (1810) ... — — — — — N.S.W. Q. N.A. B.fl.IV,311 M.fr.VI,117 ... P.
T. pubescens ; R. Brown, prodr. 468 (1810) ... — — — — — Q. N.A. B.fl.IV,311 M.fr.VI,118 ... P.

ALSTONIA ; R. Brown in Mem. Wern. Soc. I, 75 (1809).

A. scholaris ; R. Brown in Mem. Wern. Soc. I, 75 (1809) ... — — — — — Q. — B.fl.IV,312 M.fr.VI,117 ... As., Afr.
A. verticillata ; F. v. M., fragm. VI, 116 (1868) ... — — — — — Q. N.A. B.fl.IV,313 M.fr.VI,116.
A. ophioxylodes ; F. v. M., fragm. I, 57 (1858) ... — — — — — — N.A. B.fl.IV,313 M.fr.VI,117.
A. villosa ; Blume, Bijdr. 1038 (1826) ... — — — — — Q. — B.fl.IV,313 M.fr.VI,117 ... As.
A. linearis ; Bentham, Fl. Austr. IV, 314 (1869) ... — — — — — — N.A. B.fl.IV,314
A. constricta ; F. v. M., fragm. I, 57 (1858) ... — — — — — N.S.W. Q. — B.fl.IV,314 M.fr.III,163; IV,170; VI,117; VII,131.

ICHNOCARPUS ; R. Brown in Mem. Wern. Soc. I, 75 (1809).

I. frutescens ; R. Brown in Mem. Wern. Soc. I, 75 (1809) ... — — — — — Q. N.A. B.fl.IV,315 M.fr.VI,118 ... As.

WRIGHTIA ; R. Brown in Mem. Wern. Soc. I, 75 (1809). (Balfouria.)

W. pubescens ; R. Brown, prodr. 467 (1810) ... — — — — — — N.A. B.fl.IV,316 ... As., P.
W. saligna ; F. v. M. in Benth. Fl. Austr. IV, 316 (1869) ... — — — — — Q. N.A. B.fl.IV,316 M.fr.V,161; VI,118.
W. Cunninghamii ; Bentham, Fl. Austr. IV, 317 (1869) ... — — — — — W.A. — — — — — B.fl.IV,317

PARSONIA ; R. Brown in Mem. Wern. Soc. I, 64 (1809). (Lyonsiae subgenus.)

P. lanceolata ; R. Brown, prodr. 466 (1810) ... — — — — — N.S.W. Q. — B.fl.IV,318 M.fr.VI,126.
P. velutina ; R. Brown, prodr. 466 (1810) ... — — — — — N.S.W. Q. N.A. B.fl.IV,318 M.fr.VI,130.
P. Leichhardtii ; F. v. M., fragm. VI, 128 (1868) ... — — — — — Q. — B.fl.IV,319 M.fr.VI,128.
P. ventricosa ; F. v. M. in Transact. phil. Inst. Vict. II, 71 (1857) ... — — — — — N.S.W. Q. — B.fl.IV,319 M.fr.VI,130.

LYONSIA ; R. Brown in Mem. Wern. Soc. I, 66 (1809).

L. lilacina ; F. v. M. in Bentham, Fl. Austr. IV, 321 (1869) ... — — — — — N.S.W. Q. — B.fl.IV,321 M.fr.VI,127.
L. induplicata ; F. v. M. in Bentham, Fl. Austr. IV, 321 (1869) ... — — — — — N.S.W. — — B.fl.IV,321 M.fr.VI,129.
L. straminea ; R. Brown, prodr. 466 (1810) ... — — — — — T. V. N.S.W. — — B.fl.IV,321 M.fr.VI,129.
L. reticulata ; F. v. M., Rep. Burdek. Exped. 16 (1860) ... — — — — — N.S.W. Q. — B.fl.IV,321 M.fr.VI,129.
L. diaphanophleba ; F. v. M. in Benth. Fl. Austr. IV, 322 (1869) ... — — — — — W.A. — — — — — B.fl.IV,322 M.fr.II,158; VI,130.
L. Langiana ; F. v. M., fragm. VI, 128 (1868) ... — — — — — Q. — B.fl.IV,322 M.fr.VI,128.
L. largiflorens ; F. v. M. in Benth. Fl. Austr. IV, 322 (1869) ... — — — — — N.S.W. Q. — B.fl.IV,322
L. latifolia ; Bentham, Fl. Austr. IV, 323 (1869) ... — — — — — Q. — B.fl.IV,323
L. oblongifolia ; Bentham, Fl. Austr. IV, 323 (1869) ... — — — — — Q. — B.fl.IV,323
L. eucalyptifolia ; F. v. M. in Benth. Fl. Austr. IV, 323 (1869) ... — — — — — N.S.W. Q. — B.fl.IV,323 M.fr.II,159; VI,130.

ASOLEPIADEAE.

N. J. JACQUIN, Misc. Austr. I, 35 (1778).—Genera 15, Species 61.

GYMNANTHERA ; R. Brown in Mem. Wern. Soc. I, 58 (1809).

G. nitida ; R. Brown, prodr. 464 (1810) ... — — — — — Q. N.A. B.fl.IV,326 M.fr.V,160 ... P.

SECAMONE ; R. Brown in Mem. Wern. Soc. I, 55 (1809).

S. elliptica ; R. Brown, prodr. 464 (1810) ... — — — — — N.S.W. Q. N.A. B.fl.IV,327 M.fr.V,161 ... As.
S. ovata ; R. Brown, prodr. 464 (1810) ... — — — — — Q. N.A. B.fl.IV,328

VINCETOXICUM ; Dodoens, stirp. hist. pemptades, 704 (1583). (Oxystelma partly, Cynoctonum, Cynanchi subgenus.)

V. ovatum ; Bentham, Fl. Austr. IV, 330 (1869) ... — — — — — Q. — B.fl.IV,330
V. elegans ; Bentham, Fl. Austr. IV, 330 (1869) ... — — — — — N.S.W. Q. N.A. B.fl.IV,330 M.fr.V,160.

- V. carnosum*; Benth. Fl. Austr. IV, 331 (1869) ... — — — — N.S.W. Q. N.A. B.f.IV, 331
V. leptolepis; Benth. Fl. Austr. IV, 331 (1869) ... — — — — — Q. — B.f.IV, 331

CYNANCHUM; Linné, gen. plant. 63 (1737). (Endotropis.)

- C. erubescens*; R. Brown, prodr. 463 (1810) ... — — — — — Q. — B.f.IV, 332
C. floribundum; R. Brown, prodr. 463 (1810) ... W.A. S.A. — — — — Q. N.A. B.f.IV, 332 M.fr.V, 160.
C. pedunculatum; R. Brown, prodr. 463 (1810) ... — — — — — Q. N.A. B.f.IV, 333 M.fr.V, 160.
C. puberulum; F. v. M. in Benth. Fl. Austr. IV, 333 (1869) ... — — — — — — N.A. B.f.IV, 333

SARCOSTEMMA; R. Brown in Mem. Wern. Soc. I, 50 (1809).

- S. Australe*; R. Brown, prodr. 463 (1810) ... — — — — — W.A. S.A. — V. N.S.W. Q. N.A. B.f.IV, 328 M.fr.V, 160; XI, 80.

DAEMIA; R. Brown in Mem. Wern. Soc. I, 50 (1809). (Pentatropis, Rhyncharrhena.)

- D. linearis*; F. v. M., Cens. 94 (1882) from Decaisne (1833) ... — — — — — W.A. — — — — B.f.IV, 329
D. atropurpurea; F. v. M., Cens. 94 (1882) ... — — — — — Q. — B.f.IV, 329 M.fr.I, 128.
D. quinquepartita; F. v. M., Cens. 94 (1882) ... — — — — — V. N.S.W. — — B.f.IV, 329 M.fr.I, 128; XI, 80.
D. Kempeana; F. v. M. in Wing's S. Sc. Record, Ang. (1882) ... — — — — — S.A. — — — — —

GYMNEMA; R. Brown in Mem. Wern. Soc. I, 33 (1809). (Bidaria.)

- G. sylvestre*; R. Brown in Mem. Wern. Soc. I, 33 (1809) ... — — — — — Q. N.A. B.f.IV, 343 ... As.
G. pleiadenium; F. v. M., fragm. XI, 77 (1879) ... — — — — — Q. — — M.fr.XI, 77.
G. Muelleri; Benth. Fl. Austr. IV, 343 (1869) ... — — — — — — N.A. B.f.IV, 343
G. brevifolium; Benth. Fl. Austr. IV, 343 (1809) ... — — — — — Q. — B.f.IV, 343
G. trinerve; R. Brown, prodr. 462 (1810) ... — — — — — Q. N.A. B.f.IV, 343
G. stenophyllum; A. Gray in Proc. Am. Ac. of Sc. V, 335 (1864) ... — — — — — — N.A. B.f.IV, 344 ... P.

GONGRONEMA; Decaisne in De Candolle, prodr. VIII, 624 (1844).

- G. micradenia*; Benth. & J. Hooker, gen. plant. II, 770 (1876) ... — — — — — Q. — B.f.IV, 344 M.fr.XI, 78.

TYLOPHORA; R. Brown in Mem. Wern. Soc. I, 2S (1809). (Hybanthera.)

- T. erecta*; F. v. M. in Benth. Fl. Austr. IV, 334 (1869) ... — — — — — Q. N.A. B.f.IV, 334 M.fr.IX, 71; XI, 79.
T. macrophylla; Benth. Fl. Austr. IV, 334 (1869) ... — — — — — — N.A. B.f.IV, 334
T. grandiflora; R. Brown, prodr. 460 (1810) ... — — — — — N.S.W. Q. — B.f.IV, 334 M.fr.IX, 71; XI, 80.
T. floribunda; Benth. Fl. Austr. IV, 335 (1869) ... — — — — — N.S.W. Q. — B.f.IV, 335
T. calcarata; Benth. Fl. Austr. IV, 335 (1869) ... — — — — — Q. — B.f.IV, 335
T. barbata; R. Brown, prodr. 460 (1810) ... — — — — — V. N.S.W. — — B.f.IV, 335 M.fr.V, 160; XI, 79.
T. Woollsii; Benth. Fl. Austr. IV, 335 (1869) ... — — — — — N.S.W. — — B.f.IV, 335
T. paniculata; R. Brown, prodr. 460 (1810) ... — — — — — N.S.W. Q. — B.f.IV, 336 M.fr.IX, 71.
T. enervis; F. v. M., fragm. IX, 70 (1875) ... — — — — — N.S.W. — — M.fr.IX, 70.
T. flexuosa; R. Brown, prodr. 460 (1810) ... — — — — — — N.A. B.f.IV, 336 M.fr.XI, 79.
T. biglandulosa; A. Gray in Proc. of Am. acad. sc. vol. V (1864) ... — — — — — N.S.W. — — — M.fr.IX, 169.

MARSDENIA; R. Brown in Mem. Wern. Soc. I, 2S (1809). (Leichhardtia, Stephanotis.)

- M. cinerascens*; R. Brown, prodr. 461 (1810) ... — — — — — Q. N.A. B.f.IV, 337
M. rhyncholepis; F. v. M., fragm. XI, 78 (1879) ... — — — — — Q. — — M.fr.XI, 78.
M. flavescens; Cunningham in Bot. Mag. t. 3289 (1833) ... — — — — — V. N.S.W. Q. — B.f.IV, 337 M.fr.V, 160.
M. cymulosa; Benth. Fl. Austr. IV, 338 (1869) ... — — — — — Q. — B.f.IV, 338
M. velutina; R. Brown, prodr. 461 (1810) ... — — — — — Q. N.A. B.f.IV, 338
M. Hullsii; F. v. M. in Benth. Fl. Austr. IV, 338 (1869) ... — — — — — — N.A. B.f.IV, 338
M. araujacea; F. v. M., fragm. VI, 135 (1868) ... — — — — — Q. — B.f.IV, 339 M.fr.VI, 135.
M. tubulosa; F. v. M., fragm. IX, 71 (1875) ... — — — — — N.S.W. — — M.fr.IX, 71.
M. rostrata; R. Brown, prodr. 461 (1810) ... — — — — — V. N.S.W. Q. — B.f.IV, 339 M.fr.V, 160; VI, 135; IX, 71.
M. Fraseri; Benth. Fl. Austr. IV, 339 (1869) ... — — — — — Q. — B.f.IV, 339
M. longiloba; Benth. Fl. Austr. IV, 340 (1869) ... — — — — — N.S.W. Q. — B.f.IV, 340 M.fr.XI, 79.

M. suaveolens ; R. Brown, prodr. 461 (1810) ...	...	...	...	...	...	N.S.W.	—	—	B.fl.IV, 340		
M. leptophylla ; F. v. M. in Benth. Fl. Austr. IV, 340 (1869) ...	...	...	...	...	...	—	—	—	Q. —	B.fl.IV, 340	
M. Leichhardtiana ; F. v. M., fragm. V, 160 (1866) ...	...	...	...	...	...	W.A. S.A.	—	V.	N.S.W.	Q. N.A. B.fl.IV, 341	M.fr.V, 160.
M. viridiflora ; R. Brown, prodr. 461 (1810) ...	...	...	...	...	...	—	—	—	N.S.W.	Q. N.A. B.fl.IV, 341	M.fr.V, 160.
M. coronata ; Benth., Fl. Austr. IV, 341 (1810) ...	...	...	...	...	...	—	—	—	—	Q. —	B.fl.IV, 341
M. microlepis ; Benth., Fl. Austr. IV, 342 (1869) ...	...	...	...	...	...	—	—	—	—	Q. —	B.fl.IV, 342
THOZETIA ; F. v. M. in Benth. Fl. Austr. IV, 347 (1869).											
T. racemosa ; F. v. M. in Benth. Fl. Austr. IV, 347 (1869) ...	...	...	...	...	...	—	—	—	—	Q. —	B.fl.IV, 347
HOYA ; R. Brown in Mem. Wern. Soc. I, 26 (1809).											
H. carnosa ; R. Brown in Mem. Wern. Soc. I, 26 (1809) ...	...	...	...	...	...	—	—	—	—	Q. —	B.fl.IV, 346 ... As.
H. Australis ; R. Brown in Transact. hort. Soc. VII, 28 (1826) ...	...	...	...	...	...	—	—	—	N.S.W.	Q. —	B.fl.IV, 346 M.fr.V, 159; XI, 80 ... P.
H. Nicholsoniae ; F. v. M., fragm. V, 159 (1886) ...	...	...	...	...	...	—	—	—	—	Q. —	B.fl.IV, 347 M.fr.IX, 71, 190.
DISCHIDIA ; R. Brown in Mem. Wern. Soc. I, 32 (1809).											
D. nummularia ; R. Brown, prodr. 461 (1810) ...	...	...	...	...	...	—	—	—	—	Q. —	B.fl.IV, 345 M.fr.V, 161 ... As., P.
D. Timorensis ; Decaisne in nouv. annal. du Mus. III, 377, t. 17 (1835) ...	...	...	...	...	...	—	—	—	—	Q. N.A.	... As.
D. Rafflesiana ; Wallich, pl. Asiat. rar. II, 35, t. 142 (1831) ...	...	...	...	...	...	—	—	—	—	Q. N.A.	... As.
MICROSTEMMA ; R. Brown in Mem. Wern. Soc. I, 25 (1809).											
M. tuberosum ; R. Brown, prodr. 459 (1810) ...	...	...	...	...	...	—	—	—	—	Q. N.A. B.fl.IV, 345	M.fr.I, 58.
M. glabriflorum ; F. v. M., fragm. I, 58 (1858) ...	...	...	...	...	...	—	—	—	—	—	N.A. B.fl.IV, 345 M.fr.I, 58.
CEROPEGIA ; Linné, gen. plant. 65 (1737).											
C. Cumingiana ; Decaisne in De Candolle, prodr. VIII, 642 (1844) ...	...	...	...	...	...	—	—	—	—	Q. —	B.fl.IV, 348 M.fr.V, 159 ... As.

CONVOLVULACEAE.

A. L. DE JUSSIEU, Gen. plant. 132 (1789), from B. de Jussieu (1759).—Genera 13, Species 70.

ERYCIBE ; Roxburgh, pl. Corom. III, 31, t. 159 (1798).

E. paniculata ; Roxburgh, pl. Corom. III, 31, t. 159 (1798) ...	...	...	...	...	...	—	—	—	—	Q. —	B.fl.IV, 411 M.fr.VI, 100 ... As.
---	-----	-----	-----	-----	-----	---	---	---	---	------	-----------------------------------

LEPISTEMON ; Blume, Bijdrag. 722 (1825).

L. urceolatus ; F. v. M. ; L. Fitzalani, F. v. M., fragm. X, III (1877) ...	...	...	...	...	...	—	—	—	—	Q. —	B.fl.IV, 427 M.fr.X, 111 ... P.
L. Lucae ; F. v. M. in Vict. Naturalist, Oct. (1885) ...	...	...	...	...	...	—	—	—	—	Q. —	

IPOMOEA ; Linné, syst. nat. 8 (1735); Linné, gen. plant. 47 (1737). (Pharbitis, Batatas, Calonyction, Aniseia, Skinnera.)

I. digitata ; Linné, syst. nat. edit. decim. n. 5 (1758) ...	...	...	...	...	...	—	—	—	—	Q. N.A. B.fl.IV, 414	M.fr.VI, 98; X, 111 ... As., Afr., Am., P.
I. Davenporti ; F. v. M., fragm. VI, 97 (1868) ...	...	...	...	...	...	—	S.A.	—	—	—	N.A. B.fl.IV, 415 M.fr.VII, 254; VIII, 17; X, 117.
I. bona nox ; Linné, spec. plant. edit. secund. 228 (1762) ...	...	...	...	...	...	—	—	—	N.S.W.	Q. —	M.fr.IX, 74; X, 73, 113. As., Am., P.
I. palmata ; Forskael, Fl. Aegypt. Arabic. 43 (1775) ...	...	...	...	...	...	—	—	—	N.S.W.	Q. —	B.fl.IV, 415 M.fr.VI, 98; X, 111 ... As., Afr., Am., P.
I. quinata ; R. Brown, prodr. 486 (1810) ...	...	...	...	...	...	—	—	—	—	Q. N.A. B.fl.IV, 415	M.fr.VI, 98; X, 112 ... As., P. [N.Z.]
I. diversifolia ; R. Brown, prodr. 487 (1810) ...	...	...	...	...	...	—	—	—	—	Q. N.A. B.fl.IV, 416	M.fr.X, 112.
I. dissecta ; Willdenow, phytogr. 5, t. 2 (1794) ...	...	...	...	...	...	—	—	—	—	Q. N.A. B.fl.IV, 416	M.fr.VI, 98; X, 112 ... As., Afr., P.
I. sinuata ; Ortega, decad. nov. pl. hort. Matrit. VII, 84 (1789) ...	...	...	...	...	...	—	—	—	—	Q. —	M.fr.VIII, 17; X, 113 ... As., Afr., Am., P.
I. hederacea ; N. J. Jacquin, Collectan. I, 124 (1776) ...	...	...	...	...	...	—	—	—	—	Q. N.A. B.fl.IV, 416	M.fr.VI, 99; X, 112 ... As., Afr., Am., P.
I. congesta ; R. Brown, prodr. 485 (1810) ...	...	...	...	...	...	—	—	—	—	Q. —	B.fl.IV, 417 M.fr.VI, 99; X, 112 ... P.
I. purpurea ; Roth, Catalect. bot. I, 36 (1797) ...	...	...	...	...	...	—	—	—	—	Q. —	B.fl.IV, 417 M.fr.VI, 99; X, 112 ... Am.
I. peltata ; Choisy, Convolvulac. orient. 70 (1834) ...	...	...	...	...	...	—	—	—	N.S.W.	Q. —	B.fl.IV, 418 ... As., Afr., P.
I. Calobra ; Hill & F. v. M., fragm. XI, 73 (1879) ...	...	...	...	...	...	—	—	—	—	Q. —	M.fr.XI, 73, 137.
I. alata ; R. Brown, prodr. 484 (1810) ...	...	...	...	...	...	—	—	—	—	—	N.A. B.fl.IV, 418 M.fr.X, 112 ... Am.
I. Turpethum ; R. Brown, prodr. 485 (1810) ...	...	...	...	...	...	—	—	—	—	Q. N.A. B.fl.IV, 418	M.fr.VI, 98 ... As., Afr., Am., P.

I. grandiflora; Lamareck, illustr. des. genr. I, 407 (1791)	...	...	...	...	...	...	...	...	...	Q. N.A. B.fl.IV,418 M.fr.VI,98; IX,74; X,112. As., Afr., P.
I. costata; F. v. M. in Benth. Fl. Austr. IV, 419 (1869)	...	...	...	...	W.A. S.A.	—	—	—	—	— N.A. B.fl.IV,419 M.fr.X,112.
I. Pes Caprae; Roth. nov. spec. plant. 109 (1821)	...	...	...	...	W.A.	—	—	—	N.S.W.	Q. N.A. B.fl.IV,419 M.fr.VI,97; X,112 ... As., Afr., Am., P.
I. carnosa; R. Brown, prodr. 485 (1810)	...	...	...	...	...	—	—	—	—	Q. N.A. B.fl.IV,420 ... As., Afr., Am., P.
I. aquatica; Forskael, fl. Aeg. Arab. 44 (1775)	...	...	...	...	W.A.	—	—	—	N.S.W.	Q. N.A. B.fl.IV,420 M.fr.VI,97 ... As., Afr.
I. graminea; R. Brown, prodr. 485 (1810)	...	...	...	...	...	—	—	—	—	Q. N.A. B.fl.IV,421
I. velutina; R. Brown, prodr. 485 (1810)	...	...	...	...	...	—	—	—	—	Q. N.A. B.fl.IV,421
I. abrupta; R. Brown, prodr. 485 (1810)	...	...	...	...	...	—	—	—	—	Q. N.A. B.fl.IV,421 M.fr.X,112.
I. denticulata; Choisy in De Candolle, prodr. IX, 379 (1845)	...	...	...	...	...	—	—	—	—	Q. N.A. B.fl.IV,421 M.fr.I,185; X,112 ... As., P.
I. gracilis; R. Brown, prodr. 484 (1810)	...	...	...	...	...	—	—	—	—	Q. N.A. B.fl.IV,422 M.fr.I,185; X,112.
I. sepriaria; Koenig in Roxb. Fl. Indic. ed. Carey II, 90 (1824)	...	...	...	...	...	—	—	—	N.S.W.	Q. N.A. B.fl.IV,422 ... As., P.
I. Muelleri; Benth. Fl. Austr. IV, 423 (1869)	...	...	...	...	W.A. S.A.	—	—	—	—	— N.A. B.fl.IV,423 M.fr.X,112.
I. cymosa; Roemer & Schultes, syst. veg. IV, 241 (1819)	...	...	...	...	...	—	—	—	—	Q. — B.fl.IV,423 ... As., Afr., P.
I. linifolia; Blume, Bijdrag. 721 (1825)	...	...	...	...	...	—	—	—	—	Q. — B.fl.IV,423 ... As.
I. chryseides; Ker in Bot. Regist. t. 270 (1818)	...	...	...	...	...	—	—	—	—	Q. — B.fl.IV,423 M.fr.VI,98 ... As., Afr.
I. flava; F. v. M. in Benth. Fl. Austr. IV, 424 (1869)	...	...	...	...	...	—	—	—	—	Q. N.A. B.fl.IV,424
I. obscura; Ker in Bot. Regist. t. 239 (1817)... ..	...	...	...	...	...	—	—	—	—	Q. — B.fl.IV,424 ... As., Afr.
I. incisa; R. Brown, prodr. 486 (1810)	...	...	...	...	...	—	—	—	—	Q. N.A. B.fl.IV,424 M.fr.I,186.
I. uniflora; Roemer & Schultes, syst. IV, 247 (1819)	...	...	...	...	...	—	—	—	—	Q. — B.fl.IV,425 M.fr.VI,99 ... As., Afr., P.
I. angustiflora; N. J. Jacquin, Collect. II, 367 (1788)	...	...	...	...	...	—	—	—	—	Q. N.A. B.fl.IV,425 M.fr.VI,97; X,112 ... As., Afr., P.
I. plebeja; R. Brown, prodr. 484 (1810)	...	...	...	...	...	—	—	—	N.S.W.	Q. — B.fl.IV,426 M.fr.VI,99; X,112.
I. eriocarpa; R. Brown, prodr. 484 (1810)	...	...	...	...	...	—	—	—	—	Q. N.A. B.fl.IV,426 M.fr.VI,98; X,112 ... As., Afr., P.
I. heterophylla; R. Brown, prodr. 487 (1810)	...	...	...	...	...	—	S.A.	—	—	Q. N.A. B.fl.IV,426 M.fr.VI,98; X,112.
I. erecta; R. Brown, prodr. 487 (1810)	...	...	...	...	...	—	—	—	—	Q. N.A. B.fl.IV,427 M.fr.VI,98.
I. cataractae; Endl. prod. fl. ins. Norf. 53 (1833)	...	...	...	...	...	—	—	—	N.S.W.	Q. — —

ARGYREIA; Loureiro, Fl. Cochinch. I, 134 (1790).

A. speciosa; Sweet, hort. Brit. sec. ed. 373 (1830)	...	...	...	...	...	—	—	—	—	Q. — — ... As.
--	-----	-----	-----	-----	-----	---	---	---	---	----------------

CONVOLVULUS; Tournefort, inst. 82, t. 17 (1700), from W. Turner (1538). (Calystegia.)

C. erubescens; Sims, Bot. Magaz. t. 1067 (1808)	...	...	...	...	W.A. S.A. T. V. N.S.W.	Q. — B.fl.IV,429 M.fr.VI,99; X,113; XI,73.
C. parviflorus; Vahl, symbol. bot. III, 29 (1794)	...	...	...	...	...	Q. N.A. B.fl.IV,430 M.fr.VI,99; X,113 ... As., P.
C. marginatus; Poiret, Encycl. méth. suppl. II (1811)	...	...	...	...	— — — V. N.S.W.	Q. — B.fl.IV,430 M.fr.VI,99; X,113 ... N.Z.
C. sepium; Linné, spec. plant. 153 (1753)	...	...	...	...	W.A. S.A. T. V. N.S.W.	— — B.fl.IV,430 M.fr.VI,99; X,113. E., As., Afr., Am., N.Z.

POLYMERIA; R. Brown, prodr. 488 (1810).

P. marginata; Benth. Fl. Austr. IV, 432 (1869)	...	...	...	...	...	Q. — B.fl.IV,432
P. longifolia; Lindley in Mitch. Trop. Austr. 398 (1848)	...	...	...	...	— S.A. — — N.S.W.	Q. — B.fl.IV,432 M.fr.X,114.
P. angusta; F. v. M., fragm. VI, 100 (1868)	...	...	...	...	— — — S.A. — — —	— N.A. B.fl.IV,432 M.fr.VI,100.
P. distigma; Benth. Fl. Austr. IV, 433 (1869)	...	...	...	...	...	— N.A. B.fl.IV,433
P. occidentalis; F. v. M., inedit.	...	...	...	...	W.A. — — — —	— — B.fl.IV,433
P. calycina; R. Brown, prodr. 488 (1810)	...	...	...	...	...	N.S.W. Q. — B.fl.IV,433 M.fr.VI,100.
P. ambigua; R. Brown, prodr. 488 (1810)	...	...	...	...	...	Q. N.A. B.fl.IV,433

PORANA; Burmann, Fl. Ind. 51, t. 21 (1768). (Duperreya.)

P. sericea; F. v. M., fragm. VI, 100 (1868)	...	...	...	...	W.A. — — — —	— N.A. B.fl.IV,434 M.fr.II,22.
--	-----	-----	-----	-----	--------------	--------------------------------

BREWERIA; R. Brown, prodr. 487 (1810).

B. linearis; R. Brown, prodr. 488 (1810)	...	...	...	...	...	Q. N.A. B.fl.IV,435
B. media; R. Brown, prodr. 488 (1810)	...	...	...	...	— S.A. — — N.S.W.	Q. N.A. B.fl.IV,436 M.fr.X,113.
B. brevifolia; Benth. Fl. Austr. IV, 436 (1869)	...	...	...	...	...	— N.A. B.fl.IV,436
B. rosea; F. v. M., fragm. I, 233 (1858)	...	...	...	...	W.A. S.A. — — — —	— N.A. B.fl.IV,436 M.fr.VI,100; X,113.
B. pannosa; R. Brown, prodr. 488 (1810)	...	...	...	...	...	Q. N.A. B.fl.IV,436 M.fr.I,233; VI,100; X,113.

EVOLVULUS ; Linné, spec. plant. ed. sec. 391 (1762).

E. linifolius ; Linné, spec. plant. ed. sec. 392 (1762) ... W.A. S.A. — — N.S.W. Q. N.A. B.fl.IV,437 M.fr.VI,100; X,113. As., Afr., Am., P.

DICHONDRA ; R. & G. Forster, char. gen. 39, t. 20 (1776).

D. repens ; R. & G. Forster, char. gen. 39, t. 20 (1776) ... W.A. S.A. T. V. N.S.W. Q. N.A. B.fl.IV,438 M.fr.VI,100; X,113. E., As., Afr., Am.

CRESSA ; Linné, Amoen. Acad. I, 121 (1747).

C. Cretica ; Linné, spec. plant. 223 (1753) ... W.A. S.A. — V. N.S.W. Q. N.A. B.fl.IV,437 M.fr.VI,100 ... E., As., Afr., Am.

WILSONIA ; R. Brown, prodr. 490 (1810).

W. humilis ; R. Brown, prodr. 490 (1810) ... W.A. S.A. T. V. N.S.W. — — B.fl.IV,439 M.fr.VI,101; X,113.

W. rotundifolia ; Hooker, icon. plant. t. 410 (1842) ... W.A. S.A. — V. N.S.W. — — B.fl.IV,440 M.fr.VI,101.

W. Backhousi ; J. Hooker in Lond. Journ. VI, 275 (1847) ... W.A. S.A. T. V. N.S.W. — — B.fl.IV,440 M.fr.VI,101; VIII,17.

CUSCUTA ; Tournefort, inst. app. 652, t. 422 (1700), from C. Bahin (1623), from Bock (1539).

C. Chinensis ; Lamarck, Encycl. méth. II, 229 (1786) ... — — — — — Q. — B.fl.IV,441 M.fr.X,113 ... As., Afr.

C. Australis ; R. Brown, prodr. 491 (1810) ... W.A. S.A. — V. N.S.W. Q. — B.fl.IV,441 M.fr.I,100; X,113... As., Afr., Am.

C. Tasmanica ; Engelmann in Transact. Acad. St. Louis I, 512 (1860) ... — S.A. T. V. — — — B.fl.IV,441

HYDROPHYLLEAE.

R. BROWN in Bot. Regist. t. 242 (1817).—Genus 1, Species 2.

HYDROLEA ; Linné, spec. plant. ed. sec. 328 (1762).

H. Zeilanica ; Vahl, sym. bot. II, 46 (1791) ... — — — — — Q. N.A. B.fl.IV,382 ... As., Afr., Am.

H. spinosa ; Linné, spec. plant. ed. sec. 328 (1762) ... — — — — — Q. — B.fl.IV,383 ... Am.

SOLANACEAE.

HALLER, enum. stirp. Helv. praef. 34 (1742).—Genera 9, Species 79.

PHYSALIS ; Linné, syst. nat. 8 (1735); gen. plant. 51 (1737).

P. minima ; Linné, spec. plant. 183 (1753) ... — — — — — N.S.W. Q. N.A. B.fl.IV,466 M.fr.VI,144 ... As., Afr., Am., P.

SOLANUM ; Tournefort, inst. 148, t. 62 (1700) from Celsus.

S. nigrum ; Linné, spec. plant. 186 (1753) ... W.A. S.A. T. V. N.S.W. Q. N.A. B.fl.IV,446 M.fr.VI,145 ... E., As., Afr., Am., P., [N.Z.]

S. vescum ; F. v. M. in Transact. Vict. Inst. 69 (1855) ... — — T. V. N.S.W. — — — M.fr.I,123 ... N.Z.

S. aviculare ; G. Forster, prodr. 18 (1786) ... — S.A. T. V. N.S.W. Q. — B.fl.IV,447 M.fr.I,123; VI,144 ... N.Z.

S. simile ; F. v. M. in Transact. phil. Soc. Vict. I, 19 (1854) ... W.A. S.A. — V. N.S.W. Q. — B.fl.IV,448 M.fr.I,123; VI,144.

S. fasciculatum ; F. v. M., fragm. 1, 123 (1859) ... W.A. S.A. — — — — — M.fr.VI,144.

S. Bauerianum ; Endlicher, prodr. fl. Norfolk, 54 (1833) ... — — — — — N.S.W. — — — M.fr.IX,74.

S. viride ; Solander in G. Forster, prodr. 89 (1786) ... — — — — — Q. — B.fl.IV,449 M.fr.VI,145; IX,75 P.

S. tetrandrum ; R. Brown, prodr. 445 (1810) ... — — — — — Q. N.A. B.fl.IV,449

S. Shanesii ; F. v. M., fragm. VI, 144 (1868) ... — — — — — Q. — B.fl.IV,448 M.fr.VI,254.

S. verbascifolium ; Linné, spec. plant. 184 (1753) ... — — — — — N.S.W. Q. — B.fl.IV,449 M.fr.VI,145 ... As., Am., P.

S. discolor ; R. Brown, prodr. 445 (1810) ... — — — — — N.S.W. Q. N.A. B.fl.IV,450 M.fr.II,166; VI,147.

S. stelligerum ; Smith, Exot. Bot. II, 57, t. 88 (1805) ... — — — — — N.S.W. Q. — B.fl.IV,450 M.fr.VI,147.

S. parvifolium ; R. Brown, prodr. 446 (1810) ... — — — — — N.S.W. Q. — B.fl.IV,451 M.fr.II,164; VI,147.

S. ferocissimum ; Lindley in Mitch. Three Exped. II, 58 (1838) ... — S.A. — — N.S.W. Q. — B.fl.IV,451

S. sporadotrichum ; F. v. M. in Melb. Chemist, Oct. (1882) ... — — — — — N.S.W. Q. — —

S. defensum ; F. v. M., fragm. V, 193 (1866) ... — — — — — Q. — B.fl.IV,451 M.fr.V,193.

S. violaceum ; R. Brown, prodr. 445 (1810) ... — — — — — N.S.W. Q. — B.fl.IV,452

S. tethrathecum ; F. v. M., fragm. II, 165 (1861) ... — — — — — N.S.W. Q. — B.fl.IV,453 M.fr.II,165.

S. elachophyllum ; F. v. M., fragm. II, 164 (1861) ... — — — — — Q. — B.fl.IV,453 M.fr.II,164.

S. orbiculatum ; Dunal in Poir. Encycl. méth. XIII, 762 (1817) ... W.A. S.A. — — — — — B.fl.IV,453 M.fr.VI,145.

S. oligacanthum; F. v. M. in Transact. phil. Soc. Vict. I, 19 (1854)	...	...	—	S.A.	—	—	N.S.W.	—	—	B.fl.IV,454	M.fr.VI,145.	
S. esuriale; Lindley in Mitch. Three Exped. II, 43 (1838)...	...	...	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.IV,454		
S. chenopodinum; F. v. M., fragm. II, 165 (1861)	...	...	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.IV,454	M.fr.VI,146.	
S. Sturtianum; F. v. M. in Transact. phil. Soc. Vict. I, 19 (1854)	...	...	—	S.A.	—	—	N.S.W.	—	N.A.	B.fl.IV,454	M.fr.VI,146.	
S. furfuraceum; R. Brown, prodr. 446 (1810)	...	...	—	—	—	—	—	Q.	—	B.fl.IV,455		
S. dianthophorum; Dunal, hist. des Solan. 183 (1813)	...	...	—	—	—	—	—	Q.	—	B.fl.IV,455	M.fr.VI,145.	
S. Dallachyi; Benth. Fl. Austr. IV, 456 (1869)	...	...	—	—	—	—	—	Q.	—	B.fl.IV,456	M.fr.VI,145.	
S. densivestitum; F. v. M. in Benth. Fl. Austr. IV, 456 (1869)	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,456		
S. nemophilum; F. v. M., fragm. II, 161 (1861)	...	...	—	—	—	—	—	Q.	—	B.fl.IV,456	M.fr.II,161.	
S. Oldfieldii; F. v. M., fragm. II, 161 (1861)...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV,457	M.fr.VI,145.	
S. semiarmatum; F. v. M., fragm. II, 163 (1861)	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,457	M.fr.VI,146.	
S. xanthocarpum; Schrader in Wendl. sert. Hannov. 8, t. 2 (1795)	...	...	—	—	—	V.	N.S.W.	Q.	—	B.fl.IV,458	...	...
S. Hystrix; R. Brown, prodr. 446 (1810)	...	...	...	W.A.	S.A.	—	—	—	—	B.fl.IV,458		...
S. cataphractum; Cunningham in Benth. Fl. Austr. IV, 459 (1869)	...	...	—	—	—	—	—	—	N.A.	B.fl.IV,459		As., P.
S. pungetium; R. Brown, prodr. 446 (1810)	...	...	—	—	—	V.	N.S.W.	Q.	—	B.fl.IV,459	M.fr.VI,146.	
S. eremophilum; F. v. M. in Schlecht. Linnaea XXV, 432 (1852)	...	...	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.IV,459	M.fr.VI,146.	
S. campanulatum; R. Brown, prodr. 446 (1810)	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,460	M.fr.VI,146.	
S. adenophorum; F. v. M., fragm. II, 162 (1861)	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,460	M.fr.II,162.	
S. cinereum; R. Brown, prodr. 446 (1810)	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,460		
S. lacunarium; F. v. M. in Transact. phil. Soc. Vict. I, 18 (1854)	...	...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.IV,461	M.fr.VI,146.	
S. petrophilum; F. v. M. in Linnaea XXV, 433 (1852)	...	...	—	S.A.	—	—	N.S.W.	—	—	B.fl.IV,461	M.fr.VI,146.	
S. diversiflorum; F. v. M., fragm. VI, 146 (1868)	...	...	—	—	—	—	—	—	N.A.	B.fl.IV,461	M.fr.VI,146.	
S. carduiforme; F. v. M., fragm. II, 163 (1861)	...	...	—	—	—	—	—	Q.	N.A.	B.fl.IV,462	M.fr.II,163.	
S. melanospermum; F. v. M., fragm. II, 163 (1861)	...	...	—	—	—	—	—	—	N.A.	B.fl.IV,462	M.fr.II,163.	
S. horridum; Dunal, Solan. synops. 28 (1816)	...	...	—	—	—	—	—	—	N.A.	B.fl.IV,463		
S. echinatum; R. Brown, prodr. 447 (1810)	...	...	—	—	—	—	—	Q.	N.A.	B.fl.IV,463	M.fr.VI,145.	
S. lasiophyllum; Dunal in Poir. Encycl. méth. XIII, 764 (1817)	...	...	...	W.A.	—	—	—	—	N.A.	B.fl.IV,463	M.fr.VI,145.	
S. ellipticum; R. Brown, prodr. 446 (1810)	...	...	...	W.A.	S.A.	—	N.S.W.	Q.	N.A.	B.fl.IV,464	M.fr.VI,145.	
S. quadriloculatum; F. v. M., fragm. II, 161 (1861)	...	...	—	—	—	—	—	Q.	N.A.	B.fl.IV,464	M.fr.II,161.	
S. phlomoides; Cunningham in Benth. Fl. Austr. IV, 464 (1869)...	...	...	...	W.A.	—	—	—	—	N.A.	B.fl.IV,464		
S. Cunninghamii; Benth. Fl. Austr. IV, 465 (1869)	...	...	—	—	—	—	—	—	N.A.	B.fl.IV,465		

LYCIUM; Linné, gen. plant. 57 (1737), from Celsus.

L. Australe; F. v. M. in Transact. phil. Soc. Vict. I, 20 (1854) ... W.A. S.A. — V. N.S.W. Q. — B.fl.IV,467 M.fr.I,83,243; II,179.

DATURA; Linné, gen. plant. 48 (1737).

D. Leichhardtii; F. v. M. in Transact. phil. Soc. Vict. I, 20 (1854) ... W.A. S.A. — — — Q. N.A. B.fl.IV,468 M.fr.VI,144.

NICOTIANA; Tournefort, inst. 117, t. 41 (1700), from C. Bauhin (1623).

N. suaveolens; Lehmann, gener. Nicot. hist. 43 (1818) ... W.A. S.A. — V. N.S.W. Q. N.A. B.fl.IV,469 M.fr.VI,144.

ANTHOTROCHE; Endlicher, nov. stirp. mus. Vindob. dec. 6 (1839).

A. pannosa; Endlicher, nov. stirp. mus. Vind. dec. 7 (1839) ... W.A. — — — — — B.fl.IV,467 M.fr.VI,142.
A. Blackii; F. v. M., fragm. VIII, 232 (1874) ... W.A. S.A. — — — — — M.fr.VIII,232.
A. Walcottii; F. v. M., fragm. I, 123 (1859) ... W.A. — — — — — B.fl.IV,468 M.fr.I,123.

ANTHOCERCIS; Labillardière, Nov. Holl. pl. spec. II, 19 (1806). (Cyphanthera, Eadesia.)

A. viscosa; R. Brown, prodr. 448 (1810) ... W.A. — — — — — B.fl.IV,475 M.fr.VI,143.
A. litorea; Labillardière, Nov. Holl. pl. spec. II, 19, t. 158 (1806) ... W.A. — — — — — B.fl.IV,476 M.fr.VI,143; XI,123.
A. gracilis; Benth. in De Candolle, prodr. X, 192 (1846) ... W.A. — — — — — B.fl.IV,476
A. genistoides; Miers, Illustr. of plants II, App. 27, t. 83 (1857) ... W.A. — — — — — B.fl.IV,476 M.fr.I,122; VI,143.
A. anisantha; Endlicher in Ann. Wien. Mus. II, 201 (1838) ... W.A. S.A. — — — — — B.fl.IV,477 M.fr.I,122; VI,143.

<i>A. intricata</i> ; F. v. M., fragm. I, 211 (1859)	W.A.	—	—	—	—	—	—	B.fl.IV, 477	M.fr.I, 211.
<i>A. arborea</i> ; F. v. M., fragm. I, 212 (1859)	W.A.	—	—	—	—	—	—	B.fl.IV, 477	M.fr.I, 212; VI, 143.
<i>A. angustifolia</i> ; F. v. M. in Transact. phil. Soc. Vict. I, 21 (1854) ..	—	S.A.	—	—	—	—	—	B.fl.IV, 478	M.fr.II, 139.
<i>A. fasciculata</i> ; F. v. M., fragm. I, 122 (1859)	W.A.	—	—	—	—	—	—	B.fl.IV, 478	M.fr.I, 122; VI, 143.
<i>A. microphylla</i> ; F. v. M., fragm. I, 179 (1859)	W.A.	—	—	—	—	—	—	B.fl.IV, 478	M.fr.I, 179; VI, 143.
<i>A. myosotidea</i> ; F. v. M. in Transact. phil. Soc. Vict. I, 20 (1854) ...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.IV, 478	M.fr.I, 179; VI, 143.
<i>A. scabrella</i> ; Benth in De Candolle, prodr. X, 192 (1846)	—	—	—	—	N.S.W.	—	—	B.fl.IV, 479	
<i>A. albicans</i> ; Cunningham in Field's N.S.W. 335, t. 2 (1825)	—	—	—	V.	N.S.W.	—	—	B.fl.IV, 479	M.fr.VI, 143.
<i>A. Odgersii</i> ; F. v. M., fragm. X, 19 (1876)	—	—	—	—	—	—	N.A.	—	M.fr.X, 19.
<i>A. Tasmanica</i> ; J. Hooker, Fl. Tasman. I, 289, t. 92 (1860)	—	—	—	T.	—	—	—	B.fl.IV, 479	M.fr.VI, 143.
<i>A. Eadeii</i> ; F. v. M., fragm. II, 139 (1861)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.IV, 480	M.fr.VI, 142.
<i>A. racemosa</i> ; F. v. M., fragm. I, 211 (1859)	W.A.	—	—	—	—	—	—	B.fl.IV, 480	M.fr.I, 211.

ISANDRA: F. v. M. in Wing's S. Sc. Rec. III, 2 (1883).

I. Bancroftii; F. v. M. in Wing's S. Sc. Rec. III, 2 (1883)	...	...	...	W.A.	-	-	-	-	-	-	-	-
---	-----	-----	-----	------	---	---	---	---	---	---	---	---

DUBOISIA : R. Brown, prodr. fl. Nov. Holl. 448 (1810).

D. myoporoides; R. Brown, prodr. 448 (1810)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,474	M.fr.VI,144	...	P.
D. Hopwoodi; F. v. M., fragm. VIII, 232 (1874)	...	...	...	...	...	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.IV,480	M.fr.II,138; VI,143; X,20,121.		
D. Leichhardtii; F. v. M. in Wing's S. Sc. Rec. II, 222 (1882)	...	...	...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.IV,480	M.fr.VI,142.		

SCROPHULARINAE.

MIRBEL, Elém. de Botan. II, 879 (1815).—Genera 22, Species 80.

MIMULUS : Linné in Act. Upsal. 82 (1741). (Uvedalia.)

M. Uvedaliae; Benth in De Candolle, prodr. X, 369 (1846)	...	...	...	—	—	—	—	—	Q. N.A. B.fl.IV,482	M.fr.VI,103; IX,167.	
M. debilis; F. v. M. in Transact. phil. Inst. Vict. I, 62 (1857)	...	..	...	—	—	—	—	—	N.A. —	M.fr.VI,103.	
M. gracilis; R. Brown, prodr. 439 (1810)	...	...	...	...	W.A. S.A.	—	V. N.S.W.	Q. N.A. B.fl.IV,482	M.fr.VI,103; IX,167...	As., Afr.	
M. repens; R. Brown, prodr. 439 (1810)	...	...	...	...	W.A. S.A. T.	V. N.S.W.	Q. —	B.fl.IV,482	M.fr.VI,103; IX,166...	N.Z.	
M. prostratus; Benth in De Candolle, prodr. X, 373 (1846)	...	...	...	...	—	S.A.	—	V. N.S.W.	Q. —	B.fl.IV,483	M.fr.IX,167.

MAZUS ; Loureiro, Fl. Cochinch. II, 385 (1790).

M. Pumilio; R. Brown, prodr. 439 (1810) — S.A. T. V. N.S.W. Q. — B.fl.IV,484 M.fr.VI,104; IX,167... N.Z.

ADENOSMA; R. Brown, prodr. fl. Nov. Holl. 442 (1810). (Pterostigma.)

A. coerulea; R. Brown, prodr. 443 (1810)	...	...	...	...	...	...	—	—	—	—	Q.	—	B.f.IV,484	...	...	...	Aa.
A. Muelleri; Bentham, Fl. Austr. IV, 485 (1869)	...	...	...	...	...	...	—	—	—	—	—	N.A.	B.f.IV,485				

STEMODIA ; Linné, syst. ed. dec. 1118 (1759). (Morgania, Limnophila.)

<i>S. lythrifolia</i> ; F. v. M. in Benth. Fl. Austr. IV, 486 (1869)	...	...	...	—	—	—	—	—	N.A.	B.fl.IV, 486				
<i>S. grossa</i> ; Bentham, Fl. Austr. IV, 486 (1869)	...	...	...	...	W.A.	—	—	—	—	N.A.	B.fl.IV, 486			
<i>S. viscosa</i> ; Roxburgh, pl. of Coromand. II, 33, t. 163 (1798)	...	...	...	...	W.A.	S.A.	—	—	Q.	N.A.	B.fl.IV, 486	...	...	As.
<i>S. linophylla</i> ; F. v. M., fragm. X, 88 (1876) ...	...	...	...	...	W.A.	—	—	—	—	—	M.fr.X, 88.			
<i>S. debilis</i> ; Bentham, Fl. Austr. IV, 487 (1869)	...	...	...	...	—	—	—	—	—	N.A.	B.fl.IV, 487			
<i>S. pedicellaris</i> ; F. v. M., fragm. VIII, 231 (1874)	...	...	...	...	W.A.	S.A.	—	—	—	—	M.fr.VIII, 231.			
<i>S. Morgania</i> ; F. v. M., fragm. X, 89 (1876) ...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.IV, 488	M.fr.VI, 104.	
<i>S. sessiliflora</i> ; F. v. M., Cens. 97 (1882) from Blume (1826)	...	...	...	...	—	—	—	—	—	N.A.	—	...	...	As.
<i>S. gratioloides</i> ; F. v. M., fragm. X, 89 (1876) from R. Brown (1810)	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.IV, 489	M.fr.X, 89	As., Afr.
<i>S. punctata</i> ; F. v. M., Cens. 97 (1882) from Blume (1826)...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl.IV, 490	...	As.
<i>S. hirsuta</i> ; Heyne in Wallich, numer. list. 3930 (1828)	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.IV, 490	...	As.
<i>S. tenuiflora</i> ; Bentham, Scrophul. Indic. 23 (1835) ...	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.IV, 490	...	As., P.
<i>S. Kingii</i> ; F. v. M. in Transact. R. S. of Vict. XXIII, 57 (1886)	...	...	...	...	W.A.	—	—	—	—	—	N.A.	—		

BRAMIA ; Lamarck, Encycl. méthod. I, 459 (1783). (Herpestis.)

B. floribunda ; F. v. M., fragm. IX, 167 (1875) from R. Brown (1810) — — — — — Q. N.A. B.fl.IV,491 M.fr.VI,104 ... As., Afr.
B. Indica ; Lamarck, Encycl. méth. I, 459 (1783) — — — — — N.S.W. Q. N.A. B.fl.IV,491 M.fr.IX,167 ... As., Afr., Am.

GRATIOLA ; Ruppis, Fl. Jenens. 241 (1718), from Dodoens (1556).

G. pedunculata ; R. Brown, prodr. 435 (1810) W.A. S.A. — V. N.S.W. Q. — B.fl.IV,492 M.fr.VI,108; IX,167.
G. Peruviana ; Linné, spec. plant. 17 (1753) W.A. S.A. T. V. N.S.W. Q. N.A. B.fl.IV,493 M.fr.VI,103; IX,167... Am., N.Z.
G. nana ; Benth in De Candolle, prodr. X, 404 (1846) — — T. V. N.S.W. — — B.fl.IV,493 M.fr.IX,167 ... N.Z.

DOPATRIUM ; Hamilton in Benth. Scroph. Ind. 4 (1835).

D. junceum ; Hamilton in Benth. Scroph. Ind. 4 (1835) — — — — — Q. — B.fl.IV,494 As.
--

ARTANEMA ; D. Don in Sweet, Fl. gard. t. 234 (1829).

A. fimbriatum ; D. Don in Sweet, Fl. Gard. t. 234 (1829) — — — — — N.S.W. Q. — B.fl.IV,495 M.fr.VI,104; IX,167.
--

LINDERNIA ; Allioni, stirp. agr. Nicaeensis enum. 178, t. 5 (1757). (Pyxidaria, Vandellia, Bonnaya, Ilysanthes, Torenia partly, Tittmannia partly, Ilyogeton.)

L. crustacea ; F. v. M., Cens. 97 (1882) from Linné (1767) — — — — — N.S.W. Q. N.A. B.fl.IV,496 As., Afr., Am.
L. pubescens ; F. v. M., Cens. 97 (1882) from Benth (1846) — — — — — — N.A. B.fl.IV,496 M.fr.VI,102.
L. pyxidaria ; Linné, mant. alter. 252 (1771) W.A. — — — — — N.S.W. Q. N.A. B.fl.IV,496 E., As., P.
L. scapigera ; R. Brown, prodr. 441 (1810) — — — — — Q. N.A. B.fl.IV,497 M.fr.VI,102.
L. subulata ; R. Brown, prodr. 441 (1810) — — — — — Q. N.A. B.fl.IV,497 M.fr.VI,102.
L. lobelioides ; F. v. M., Cens. 97 (1882) — — — — — — N.A. B.fl.IV,497 M.fr.VI,102.
L. clausa ; F. v. M., fragm. VI, 102 (1868) — — — — — — N.A. B.fl.IV,498
L. veronicifolia ; F. v. M., fragm. VI, 101 (1868) — — — — — Q. — B.fl.IV,498 M.fr.VI,101 ... As., P.
L. serrata ; F. v. M., Cens. 97 (1882) from Roxburgh (1820) — — — — — Q. — — — — — As.

HEMIARRHENA ; Benth, Fl. Austr. IV, 518 (1869).

H. plantaginea ; Benth in Hook. icon. pl. t. 1059 (1869) — — — — — — N.A. B.fl.IV,518 M.fr.VI,102.

PEPLIDIUM ; Delile, Fl. Aegypt. 148 (1813).

P. humifusum ; Delile, Fl. Aegypt. 148 (1813) — S.A. — — N.S.W. Q. N.A. B.fl.IV,500 M.fr.VI,105 ... As., Afr.
P. Muellieri ; Benth, Fl. Austr. IV, 500 (1869) W.A. S.A. — — — Q. N.A. B.fl.IV,500

MICROCARPAEA ; R. Brown, prodr. 435 (1810).

M. muscosa ; R. Brown, prodr. 435 (1810) — — — — — Q. N.A. B.fl.IV,501 M.fr.VI,104 ... As.

GLOSSOSTIGMA ; Arnott in nov. act. acad. Caes. Leop. Carol. XVIII, 355 (1836). (Tricholoma.)

G. spathulatum ; Arnott in nov. act. Leop. XVIII, 355 (1836) — — — — — Q. — B.fl.IV,501 As., Afr.
G. Drummondii ; Benth in De Candolle, prodr. X, 426 (1846) W.A. S.A. — V. N.S.W. Q. N.A. B.fl.IV,502 M.fr.VI,104.
G. elatinoide ; Benth in J. Hook. Fl. New Zeal. I, 189 (1853) — S.A. T. V. N.S.W. — — B.fl.IV,502 N.Z.

LIMOSELLA ; Lindern, opusc. plant. Argentorat. 156, t. 5 (1728).

L. aquatica ; Linné, spec. plant. 631 (1753) W.A. S.A. T. V. N.S.W. — — B.fl.IV,502 M.fr.VI,105; IX,165... F., As., Afr., Am.,
L. Curdieana ; F. v. M., fragm. IX, 166 (1875) — S.A. — V. N.S.W. Q. — — M.fr.IX,166; XI,135.

SCOPARIA ; Linné, syst. nat. ed. sext. 87 (1748).

S. dulcis ; Linné, spec. plant. 116 (1753) — — — — — Q. N.A. B.fl.IV,504 M.fr.VI,104 ... Am.

OURISIA ; Commerçon in A. L. de Jussieu, gen. pl. 100 (1789).

O. integrifolia ; R. Brown, prodr. 439 (1810) — — — — — T. — — — — B.fl.IV,512 M.fr.VI,105; IX,168.
--

VERONICA ; Tournefort, inst. 143, t. 60 (1700) from Fuchs (1543). (Pygmaea.)

V. densifolia ; F. v. M., fragm. II, 137, t. 63 (1861) — — — — — V. N.S.W. — — B.fl.IV,505 M.fr.VI,103.
V. formosa ; R. Brown, prodr. 434 (1810) — — — — — T. — — — — B.fl.IV,506 M.fr.VI,102; IX,167.
V. decorosa ; F. v. M. in Schlechtend. Linnaea XXV, 430 (1852) — S.A. — — — — B.fl.IV,506
V. perfoliata ; R. Brown, prodr. 434 (1810) — — — — — V. N.S.W. — — B.fl.IV,507 M.fr.VI,102; IX,167

V. Derwentia ; Littlejohn in Andr. Bot. Reposit. t. 531 (1808) — S.A. T. V. N.S.W. Q. — B.fl.IV,507 M.fr.VI,102; IX,168.
V. arenaria ; Cunningham in De Candolle, prodr. X, 463 (1846) — — — — N.S.W. — — B.fl.IV,507 M.fr.VI,102; IX,167.
V. nivea ; Lindley, Bot. Regist. Misc. 42 (1842) — — — — T. V. N.S.W. — — B.fl.IV,508 M.fr.VI,102; IX,167.
V. gracilis ; R. Brown, prodr. 435 (1810) — — — — S.A. T. V. N.S.W. — — B.fl.IV,508 M.fr.VI,103; IX,168.
V. distans ; R. Brown, prodr. 435 (1810) — — — — W.A. S.A. T. V. — — B.fl.IV,509 M.fr.VI,102.
V. calycina ; R. Brown, prodr. 435 (1810) — — — — W.A. S.A. T. V. N.S.W. Q. — B.fl.IV,509 M.fr.VI,103.
V. plebeja ; R. Brown, prodr. 435 (1810) — — — — T. V. N.S.W. Q. — B.fl.IV,510 M.fr.VI,102. ... N.Z.
V. notabilis ; F. v. M. in Benth. Fl. Austr. IV, 511 (1869) — — — — T. V. N.S.W. Q. — B.fl.IV,511 M.fr.IX,168.
V. serpillifolia ; Linné, spec. plant. 12 (1753)... .. — — — — V. N.S.W. — — B.fl.IV,511 M.fr.VI,103 ... E., As., Afr., Am.
V. peregrina ; Linné, spec. plant. 14 (1753) — — — — W.A. S.A. T. V. N.S.W. Q. — B.fl.IV,511 M.fr.VI,103; IX,168. Am.

RAMPHICARPA ; Benth in Hook. Comp. to the Bot. Mag. I, 368 (1835).

R. longiflora ; Benth in Hook. Comp. Bot. Mag. I, 368 (1835) — — — — — Q. N.A. B.fl.IV,518 M.fr.VI,104; IX,168. As.
--

CENTRANTHERA ; R. Brown, prodr. fl. Nov. Holl. 438 (1810).

C. hispida ; R. Brown, prodr. 438 (1810) — — — — — N.S.W. Q. N.A. B.fl.IV,513 M.fr.VI,104; IX,168. As.

SOPUBIA ; Hamilton in D. Don, prodrom. flor. Nepalens. 88 (1825).

S. trifida ; Hamilton in D. Don, prodr. fl. Nep. 88 (1825) — — — — — Q. — B.fl.IV,513
--

BUCHNERA ; Linné, hort. Cliffort. 501 (1737). (Buchnera, Striga, Campuleia.)

B. tetragona ; R. Brown, prodr. 437 (1810) — — — — — Q. N.A. B.fl.IV,514 M.fr.IX,168.
B. urticifolia ; R. Brown, prodr. 437 (1810) — — — — — N.S.W. Q. N.A. B.fl.IV,514 M.fr.VI,104.
B. linearis ; R. Brown, prodr. 437 (1810) — — — — — S.A. — — — Q. N.A. B.fl.IV,515
B. tenella ; R. Brown, prodr. 437 (1810) — — — — — — Q. N.A. B.fl.IV,515
B. gracilis ; R. Brown, prodr. 437 (1810) — — — — — N.S.W. — — B.fl.IV,515 M.fr.VI,104.
B. ramosissima ; R. Brown, prodr. 438 (1810) — — — — — Q. N.A. B.fl.IV,515
B. coccinea ; Benth in Scrophul. Ind. 40 (1835) — — — — — Q. — B.fl.IV,516
B. parviflora ; R. Brown, prodr. 438 (1810) — — — — — W.A. — — — Q. N.A. B.fl.IV,517 M.fr.VI,104 ... As., Afr., P.
B. multiflora ; F. v. M., Cens. 98 (1882) from Benth in (1836) — — — — — — N.A. B.fl.IV,517 As.
B. curviflora ; R. Brown, prodr. 438 (1810) — — — — — Q. N.A. B.fl.IV,517 M.fr.VI,104.

EUPHRASIA ; Tournefort, inst. 174, t. 78 (1700) from Matthaeus (1478).

E. Brownii ; F. v. M., fragm. V, 88 (1865) — — — — — W.A. S.A. T. V. N.S.W. Q. — B.fl.IV,521 M.fr.IX,168 ... P.
E. scabra ; R. Brown, prodr. 437 (1810) — — — — — W.A. S.A. T. V. N.S.W. Q. — B.fl.IV,521 M.fr.V,89; IX,168.
E. arguta ; R. Brown, prodr. 437 (1810) — — — — — N.S.W. — — B.fl.IV,522
E. cuspidata ; J. Hooker, Fl. Tasm. I, 298 (1860) — — — — — T. — — — B.fl.IV,522 M.fr.V,90.
E. antarctica ; Benth in De Candolle, prodr. X, 555 (1846) — — — — — V. N.S.W. — — B.fl.IV,522 M.fr.V,90 ... Am., N.Z.

OROBANCHEAE.

A. L. DE JUSSIEU in Ann. Mus. V, 25 (1804).—Genus 1, Species 1.

OROBANCHE ; Tournefort, inst. 175, t. 81 (1700), from l'Ecluse (1583).

O. Australiana ; F. v. M., Cens. 98 (1882) — — — — — W.A. S.A. — — V. N.S.W. — — B.fl.IV,533

LENTIBULARINAE.

L. C. RICHARD in Bulliard, Fl. Paris. ed. sec. I, 26 (1796).—Genera 2, Species 25.

UTRICULARIA ; Linné, syst. nat. 8 (1735); Linné, gen. plant. 5 (1737).

U. stellaris ; Linné, fl. snupl. plant. 86 (1781) — — — — — Q. — B.fl.IV,525 M.fr.VI,161; IX,48... As., Afr.
U. flexuosa ; Vahl, enum. plant. I, 198 (1804) — — — — — W.A. S.A. T. V. N.S.W. Q. N.A. B.fl.IV,525 As.
U. exoleta ; R. Brown, prodr. 430 (1810) — — — — — N.S.W. Q. N.A. B.fl.IV,526 M.fr.VI,161 ... As.

U. tubulata; F. v. M., fragm. IX, 48 (1875) ...	...	...	...	...	...	...	...	...	...	Q.	—	—	M.fr.IX, 48.
U. albiflora; R. Brown, prodr. 431 (1810) ...	...	...	...	...	...	...	...	...	...	Q.	—	B.fl.IV, 526	
U. pygmaea; R. Brown, prodr. 432 (1810) ...	...	...	...	...	...	...	...	...	...	Q.	—	B.fl.IV, 526	
U. fulva; F. v. M. in Transact. phil. Inst. Vict. III, 63 (1858) ...	...	...	...	...	...	...	...	...	...	—	N.A.	B.fl.IV, 527	M.fr.VI, 160.
U. chrysantha; R. Brown, prodr. 432 (1810) ...	...	...	...	...	...	...	...	...	...	Q.	N.A.	B.fl.IV, 527	M.fr.VI, 161; IX, 48.
U. bifida; Linné, spec. plant. 18 (1753) ...	...	...	...	...	...	...	...	...	...	Q.	—	B.fl.IV, 527	...
U. cyanea; R. Brown, prodr. 431 (1810) ...	...	...	...	...	...	...	...	...	N.S.W.	Q.	N.A.	B.fl.IV, 527	M.fr.VI, 161. ... As.
U. lateriflora; R. Brown, prodr. 431 (1810) ...	...	...	...	...	...	...	...	S.A. T. V.	N.S.W.	Q.	—	B.fl.IV, 528	M.fr.VI, 162; IX, 48.
U. simplex; R. Brown, prodr. 431 (1810) ...	...	...	...	...	...	...	...	W.A.	—	—	—	B.fl.IV, 528	M.fr.VI, 162.
U. monantha; J. Hooker, Fl. Tasm. I, 299 (1860) ...	...	...	...	...	...	...	...	—	T.	—	—	B.fl.IV, 528	
U. dichotoma; Labillardière, Nov. Holl. pl. spec. I, 11, t. 8 (1804) ...	...	...	...	...	...	...	...	S.A. T. V.	N.S.W.	Q.	—	B.fl.IV, 529	M.fr.VI, 160.
U. volubilis; R. Brown, prodr. 430 (1810) ...	...	...	...	...	...	...	...	W.A.	—	—	—	B.fl.IV, 529	M.fr.VI, 161.
U. Hookeri; Lehmann, nov. stirp. pugill. VIII, 47 (1844) ...	...	...	...	...	...	...	...	W.A.	—	—	—	B.fl.IV, 530	M.fr.VI, 162.
U. violacea; R. Brown, prodr. 431 (1810) ...	...	...	...	...	...	...	...	W.A.	—	—	—	B.fl.IV, 530	M.fr.VI, 160.
U. Menziesii; R. Brown, prodr. 431 (1810) ...	...	...	...	...	...	...	...	W.A.	—	—	—	B.fl.IV, 530	M.fr.VI, 160.
U. biloba; R. Brown, prodr. 432 (1810) ...	...	...	...	...	...	...	...	—	—	N.S.W.	Q.	—	B.fl.IV, 531 ... As.
U. leptoplectra; F. v. M. in Austr. Chem. VIII, 50 (1885) ...	...	...	...	...	...	...	...	—	—	—	—	N.A.	—
U. limosa; R. Brown, prodr. 432 (1810) ...	...	...	...	...	...	...	...	—	—	—	—	Q.	— B.fl.IV, 531
U. Baueri; R. Brown, prodr. 431 (1810) ...	...	...	...	...	...	...	...	—	—	—	—	Q.	— B.fl.IV, 531 ... As.
U. lasiocaulis; F. v. M. in Austr. Chem. VIII, 55 (1885) ...	...	...	...	...	...	...	...	—	—	—	—	N.A.	—

POLYPOMPHOLYX; Lehmann in der Bot. Zeitung, Halle, 109 (1844). (Tetralobus.)

P. multifida; F. v. M., fragm. VI, 162 (1868) ...	...	...	...	...	...	...	...	W.A.	—	—	—	—	B.fl.IV, 532	M.fr.VI, 162.
P. tenella; Lehmann, nov. stirp. pugill. VIII, 50 (1844) ...	...	...	...	...	...	...	...	W.A.	S.A. T. V.	—	—	—	B.fl.IV, 532	M.fr.VI, 162.

PODOSTEMONACEAE.

L. CL. RICHARD in Humb. Bonpl. & Kunth, Nov. gen. I, 246 (1815).—Genus 1, Species 1.

Genus and species not yet determined ...	...	...	...	...	...	...	...	...	...	Q.	—	—		
--	-----	-----	-----	-----	-----	-----	-----	-----	-----	----	---	---	--	--

GESNERIACEAE.

HUMBOLDT, Bonpland & Kunth, nov. gen. et spec. pl. II, 392 (1817).—Genera 4, Species 4.

FIELDIA; Cunningham in Field's New South Wales 363, t. 2 (1825).

F. Australis; Cunningham in Field's N.S.W. 364, t. 2 (1825) ...	...	...	...	...	...	...	...	...	V.	N.S.W.	—	—	B.fl.IV, 535	M.fr.IV, 147; IX, 191.
---	-----	-----	-----	-----	-----	-----	-----	-----	----	--------	---	---	--------------	------------------------

NEGRIA; F. v. M., fragm. VII, 151 (1871). (Rhabdothamnus partly.)

N. rhabdothamnoides; F. v. M., fragm. VII, 152 (1871) ...	...	...	...	...	...	...	...	...	...	N.S.W.	—	—	—	M.fr.VII, 152; VIII, 281; XI, 133.
---	-----	-----	-----	-----	-----	-----	-----	-----	-----	--------	---	---	---	------------------------------------

BAEA; Commerçon in Lamarck, Encycl. méth. I, 401 (1783). (Streptocarpus, Boea, Bea.)

B. hygroscopica; F. v. M., fragm. IV, 146 (1864) ...	...	...	...	...	...	...	...	...	...	Q.	—	—	B.fl.IV, 535	M.fr.IV, 146.
--	-----	-----	-----	-----	-----	-----	-----	-----	-----	----	---	---	--------------	---------------

ROETTLERA; Vahl. enum. plant. I, 87 (1805). (Didymocarpus.)

R. Kinnearii; F. v. M. in Vict. Naturalist III, 159 (1887) ...	...	...	...	...	...	...	...	...	...	Q.	—	—		
--	-----	-----	-----	-----	-----	-----	-----	-----	-----	----	---	---	--	--

BIGNONIACEAE.

VENTENAT; Tabl. du règn. végét. II, 402 (1799).—Genera 4, Species 7.

TECOMA; A. L. de Jussieu, gen. plant. 139 (1789).

T. Australis; R. Brown, prodr. 471 (1810) ...	...	...	...	...	...	...	...	...	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.IV, 537	
T. Hillii; F. v. M., fragm. X, 101 (1877) ...	...	...	...	...	...	...	...	...	—	—	—	—	Q.	—	—	M.fr.X, 101; XI, 136.
T. jasminoides; Lindley, Bot. Regist. t. 2002 (1838) ...	...	...	...	...	...	...	...	...	—	—	—	N.S.W.	Q.	—	—	B.fl.IV, 537

DOLICHANDRONE; Fenzl in Ann. nat. hist. third ser. X, 31 (1862). (Spathodea partly, Dolichandra partly.)

- D. heterophylla; F. v. M., fragm. IV, 149 (1864) ... — — — — — Q. N.A. B.fl.IV,539 M.fr.IV,149.
D. filiformis; Seemann, Jour. of Bot. I, 226 (1863) ... — — — — — — N.A. B.fl.IV,539

HAUSSMANNIA; F. v. M., fragm. IV, 148 (1864).

- H. juncunda; F. v. M., fragm. IV, 148 (1864)... — — — — — Q. — B.fl.IV,540 M.fr.IV,148.

DEPLANCHEA; Vieillard in Bullet. de la Soc. Linn. de Norm. VII, 11 (1862). (Diplanthera, Bulweria.)

- D. tetraphylla; F. v. M., from R. Brown (1810) ... — — — — — Q. — B.fl.IV,556 M.fr.IV,147;VI,72.

PEDALINAE.

R. BROWN, prodr. Fl. Nov. Holl. 519 (1810).—Genus 1, Species 3.

JOSEPHINA; Ventenat, Jardin de la Malmaison 103, t. 67 (1804).

- J. grandiflora; R. Brown, prodr. 520 (1810) ... — — — — — Q. — B.fl.IV,540 ... P.
J. Imperatricis; Ventenat, Jard. de la Malm. t. 67 (1804)... — — — — — — N.A. B.fl.IV,557 M.fr.VI,163... As.
J. Eugeniae; F. v. M. in Hook. Kew Misc. IX, 370, t. 11 (1857)... — — — — — W.A. S.A. — — Q. N.A. B.fl.IV,557 M.fr.VI,163.

ACANTHACEAE.

A. L. DE JUSSIEU, gen. plant. 102 (1789), from B. de Jussieu (1759).—Genera 13, Species 30.

THUNBERGIA; Retzius, Physiogr. Saellsk. Handl. I, 163 (1776).

- T. Arnhemica; F. v. M., fragm. IX, 73 (1875) ... — — — — — — N.A. — M.fr.IX,73 ... As.
T. Powellii; F. v. M. in Wing's South. Sc. Rec. II, 34 (1882) ... — — — — — — N.A. —
T. fragrans; J. Koenig in Roxb. pl. Coromand. I & IV (1795) ... — — — — — — N.A. —

NELSONIA; R. Brown, prodr. fl. Nov. Holl. 485 (1810).

- N. campestris; R. Brown, prodr. 481 (1810) ... — — — — — Q. N.A. B.fl.IV,543 M.fr.VI,88; XI,18 ... As., Afr., Am.

EBERMAYERA; Nees in Wallich, pl. Asiat. rar. III, 75 (1832).

- E. glauca; Nees in De Candolle, prodr. XI, 73 (1847) ... — — — — — Q. N.A. B.fl.IV,544 M.fr.VI,88; XI,18 ... As.

HYGROPHILA; R. Brown, prodr. fl. Nov. Holl. 479 (1810).

- H. angustifolia; R. Brown, prodr. 479 (1810) ... — — — — — Q. N.A. B.fl.IV,544 M.fr.VI,88 ... As.

STROBILANTHES; Blume, Bijdr. tot de fl. van Nederl. Ind. 796 (1826).

- S. Tatei; F. v. M. in Transact. Roy. Soc. S. Austr. V, Aug. (1882) ... — — — — — — N.A. —

RUELLIA; Plumier, nov. gen. 12, t. 2 (1703). (Dipteracanthus.)

- R. bracteata; R. Brown, prodr. 479 (1810) ... — — — — — Q. N.A. B.fl.IV,546 M.fr.VI,91 ... As.
R. primulaea; F. v. M. in Benth. Fl. Austr. IV, 546 (1869) ... — S.A. — — — Q. — B.fl.IV,546 M.fr.XI,18.
R. corynotheca; F. v. M. in Benth. Fl. Austr. IV, 546 (1869) ... — — — — — N.S.W. Q. — B.fl.IV,546 M.fr.XI,18.
R. Australis; R. Brown, prodr. 479 (1810) ... — S.A. — — — N.S.W. Q. — B.fl.IV,547 M.fr.VI,91.
R. spiciflora; F. v. M. in Benth. Fl. Austr. IV, 547 (1869) ... — — — — — Q. — B.fl.IV,547
R. acaulis; R. Brown, prodr. 479 (1810) ... — — — — — Q. — B.fl.IV,547

ACANTHUS; Tournefort, inst. 176, t. 81 (1700) from Theophrastos. (Dilivaria.)

- A. ilicifolius; Linné, spec. plant. 939 (1753) ... — — — — — Q. N.A. B.fl.IV,548 M.fr.X,18 ... As., P.

JUSTICIA; Houston in Linné, gen. plant. 4 (1737). (Rostellaria, Rostellularia.)

- J. procumbens; Linné, spec. plant. 15 (1753) ... — — — — — W.A. S.A. — — N.S.W. Q. N.A. B.fl.IV,549 M.fr.VI,91; XI,18; XII,24 As., Afr., P.
J. Bonneyana; F. v. M. in Wing's S. Sc. Record II, 74 (1882) ... — — — — — S.A. — — N.S.W. — — M.fr.X,73.
J. hygrophiloides; F. v. M., fragm. VI, 89 (1868) ... — — — — — — Q. — B.fl.IV,550 M.fr.VI,89.
J. Kempeana; F. v. M., fragm. XI, 101 (1881) ... — — — — — W.A. S.A. — — — N.A. — M.fr.XI,101.

- J. cavernarum*; F. v. M., fragm. VI, 91 (1868) ... — — — — — Q. — B.fl.IV,550 M.fr.VI,91.
J. eranthemoides; F. v. M., fragm. VI, 90 (1868) ... — — — — — N.S.W. Q. — B.fl.IV,551 M.fr.VI,90.

GRAPTOPHYLLUM; Nees in Wallich, pl. Asiat. rar. III, 102 (1832). (Earlia, Thyracanthus.)

- G. Earlii*; F. v. M., fragm. VI, 87 (1868) ... — — — — — Q. — B.fl.IV,551 M.fr.III,160.
G. ilicifolium; F. v. M. in Benth. Fl. Austr. IV, 552 (1869) ... — — — — — Q. — B.fl.IV,552
G. spinigerum; F. v. M., fragm. XI, 17 (1878) ... — — — — — Q. — — M.fr.XI,17.

DICLIPTERA; A. L. de Jussieu in Ann. du Mus. IX, 267 (1807). (Brochosiphon.)

- D. glabra*; Decaisne in Nouv. Annal. du Mus. III, 388 (1834) ... — — — — — — N.A. B.fl.IV,552 M.fr.VI,88; XI,18 ... As.
D. spicata; Decaisne in Nouv. Annal. du Mus. III, 389 (1834) ... — — — — — Q. N.A. B.fl.IV,553 M.fr.VI,89 ... As., P.
D. Burmanni; Nees in Wallich, pl. Asiat. rar. III, 112 (1832) ... — — — — — Q. — — M.fr.VII,62 ... As.

HYPOESTES; Solander in R. Brown, prodr. 474 (1810).

- H. floribunda*; R. Brown, prodr. 474 (1810) ... — — — — — N.S.W. Q. N.A. B.fl.IV,553 M.fr.XI,18.

ERANTHEMUM; Linné, Fl. Zeyl. 6 (1747).

- E. variabile*; R. Brown, prodr. 477 (1810) ... — — — — — N.S.W. Q. — B.fl.IV,555 M.fr.VI,91; XI,18 ... P.
E. tenellum; Benth. Fl. Austr. IV, 555 (1869) ... — — — — — Q. — B.fl.IV,555

DICLADANTHERA; F. v. M., fragm. XII, 23 (1882).

- D. Forrestii*; F. v. M., fragm. XII, 23 (1882) ... W.A. — — — — — — — M.fr.XII,23

LABIATAE.

ADANSON, Famill. des plant. II, 180 (1763), from B. de Jussieu (1759).—Genera 20, Species 125.

OCIMUM; Tournefort, inst. 203, t. 96 (1700), from Theophrastos and Dioscorides. (Ocymum.)

- O. sanctum*; Linné, mantiss. plant. 85 (1767) ... — — — — — Q. N.A. B.fl.V,74 M.fr.IV,46 ... As., Afr., P.

MOSCHOSMA; Reichenbach, conspect. regn. vegetab. 115 (1828). (Basilicum 1802.)

- M. polystachya*; Benth. in Wall. pl. As. rar. II, 13 (1831) ... — — — — — N.S.W. Q. N.A. B.fl.V,75 M.fr.VI,108 ... As., Afr., P.

ORTHOSIPHON; Benth. in Wall. pl. As. rar. II, 15 (1831).

- O. stamineus*; Benth. in Wall. pl. As. rar. II, 15 (1831) ... — — — — — Q. N.A. B.fl.V,76 M.fr.IV,47; VI,108; IX,161. As., P.

COLEUS; Loureiro, Fl. Cochinch. II, 372 (1790).

- C. scutellarioides*; Benth. in Wall. pl. Asiat. rar. II, 16 (1831) ... — — — — — Q. N.A. B.fl.V,79 M.fr.IV,47; VI,108 ... As., P.

PLECTRANTHUS; L'Héritier, stirp. nov. I, 85 (1785).

- P. foetidus*; Benth. labiat. 35 (1832) ... — — — — — Q. — — M.fr.IX,161; XI,135.
P. congestus; R. Brown, prodr. 506 (1810) ... — — — — — Q. — B.fl.V,78 M.fr.VI,108.
P. parviflorus; Henckel, adumbrat. pl. hort. Halens. (1806) ... — S.A. — V. N.S.W. Q. N.A. B.fl.V,78 M.fr.IV,47; VI,108; IX,161. P.
P. longicornis; F. v. M., fragm. V, 51 (1865) ... — — — — — Q. — B.fl.V,77 M.fr.V,51.

POGOSTEMON; Desfontaines in Mém. du Mus. II, 154, t. 6 (1815). (Dysophylla.)

- P. verticillatus*; Hasskarl, catal. hort. Bogor. 131 (1844) ... — — — — — Q. N.A. B.fl.V,81 M.fr.V,200; IX,161 ... As.

MENTHA; Tournefort, inst. 188, t. 89 (1700), from Hippocrates and Theophrastos.

- M. laxiflora*; Benth. in De Candolle, prodr. XII, 174 (1848) ... — — — — — V. N.S.W. — — B.fl.V,82 M.fr.VI,109; IX,162.
M. grandiflora; Benth. in Mitch. Trop. Austr. 362 (1848) ... — — — — — N.S.W. Q. — B.fl.V,83
M. Australis; R. Brown, prodr. 505 (1810) ... — S.A. T. V. N.S.W. Q. N.A. B.fl.V,83 M.fr.VI,109; IX,162.
M. gracilis; R. Brown, prodr. 505 (1810) ... — S.A. T. V. N.S.W. — — B.fl.V,83 M.fr.VI,109.
M. satrejoideis; R. Brown, prodr. 505 (1810) ... W.A. S.A. T. V. N.S.W. Q. — B.fl.V,84 M.fr.VI,109.

LYCOPUS; Tournefort, inst. 180, t. 83 (1700), from Plinius.

- L. Australis*; R. Brown, prodr. 500 (1810) ... — S.A. T. V. N.S.W. Q. — B.fl.V,85 M.fr.VI,109.

SALVIA; Linné, gen. plant. 6 (1737), from Plinius.														
S. plebeja; R. Brown, prodr. 501 (1810)	...	...	...	...	...	—	—	—	V. N.S.W.	Q.	—	B.fl.V,85	M.fr.VI,109; IX,161	As., P.
BRUNELLA; L'Ecluse, rar. stirp. hist. II, 42, 43 (1576), from Brunfels (1536). (Prunella.)														
B. vulgaris; A. De Candolle, prodr. XII, 410 (1848)	...	...	...	...	...	—	—	—	S.A. T.	V. N.S.W.	Q.	—	B.fl.V,87	M.fr.VI,109 ... E., As., Afr. Am.
SCUTTELLARIA; Hermann, hort. Lugd. bot. cat. (1687), from Cortusi (1591).														
S. mollis; R. Brown, prodr. 507 (1810)	...	...	...	...	...	—	—	—	V. N.S.W.	—	—	B.fl.V,88	M.fr.VI,109.	
S. humilis; R. Brown, prodr. 507 (1810)	...	...	...	...	...	—	—	—	S.A. T.	V. N.S.W.	Q.	—	B.fl.V,88	M.fr.VI,109.
ANISOMELES; R. Brown, prodr. 503 (1810).														
A. salvifolia; R. Brown, prodr. 503 (1810)	...	...	...	...	...	—	—	—	—	—	—	Q. N.A.	B.fl.V,89	M.fr.VI,109; IX,162 ... P.
LEUCAS; J. Burmann, thesaur. Zeyl. 140 (1737).														
L. flaccida; R. Brown, prodr. 505 (1810)	...	...	...	...	...	—	—	—	—	—	—	Q. N.A.	B.fl.V,90	M.fr.VI,109 ... As.
PROSTANTHERA; Labillardière, Nov. Holl. pl. spec. 18, t. 157 (1806). (Chilodia, Cryphia, Klanderia.)														
P. lasiantha; Labillardière, Nov. Holl. pl. spec. II, 18, t. 157 (1806)	...	...	...	...	...	—	—	—	S.A. T.	V. N.S.W.	Q.	—	B.fl.V,93	M.fr.VI,105.
P. prunelloides; R. Brown, prodr. 508 (1810)	...	...	...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.V,94	M.fr.VI,106.
P. coerulea; R. Brown, prodr. 508 (1810)	...	...	...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.V,94	M.fr.VI,106.
P. ovalifolia; R. Brown, prodr. 509 (1810)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.V,95	M.fr.VI,106.
P. melissifolia; F. v. M., fragm. I, 19 (1858)...	...	...	...	...	...	—	—	—	V.	—	—	—	B.fl.V,95	M.fr.I,19,242.
P. incisa; R. Brown, prodr. 509 (1810)	...	...	...	...	...	—	—	—	V. N.S.W.	Q.	—	—	B.fl.V,95	M.fr.VI,107; IX,162.
P. rotundifolia; R. Brown, prodr. 509 (1810)...	...	...	...	...	...	—	—	—	S.A. T.	V. N.S.W.	—	—	B.fl.V,96	M.fr.VI,107; IX,162.
P. Schultzii; F. v. M., cens. fourth suppl. 4 (1889)	...	...	...	...	...	—	—	—	S.A.	—	—	—	N.A.	—
P. violacea; R. Brown, prodr. 509 (1810)	...	...	...	...	...	—	—	—	V. N.S.W.	—	—	—	B.fl.V,96	M.fr.VI,107; IX,162.
P. incana; Cunningham in Benth. lab. 455 (1836)	...	...	...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.V,97	M.fr.IX,162.
P. hirtula; F. v. M., first gen. Report 16 (1853)	...	...	...	...	...	—	—	—	V. N.S.W.	—	—	—	B.fl.V,97	
P. denticulata; R. Brown, prodr. 509 (1810)	...	...	...	...	...	—	—	—	V. N.S.W.	—	—	—	B.fl.V,97	M.fr.VI,105.
P. rugosa; Cunningham in Benth. lab. 456 (1836)	...	...	...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.V,98	
P. marifolia; R. Brown, prodr. 509 (1810)	...	...	...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.V,98	M.fr.VI,105.
P. rhombea; R. Brown, prodr. 509 (1810)	...	...	...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.V,99	
P. spinosa; F. v. M. in Transact. phil. Soc. Vict. I, 48 (1854)	...	...	...	...	...	—	—	—	S.A.	—	V.	—	B.fl.V,99	M.fr.VI,108; IX,162.
P. cuneata; Benth in De Candolle, prodr. XII, 560 (1848)	...	...	...	...	...	—	—	—	T.	V. N.S.W.	—	—	B.fl.V,99	M.fr.VI,108.
P. linearis; R. Brown, prodr. 509 (1810)	...	...	...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.V,100	M.fr.VI,106.
P. phyllifolia; F. v. M., fragm. I, 19 (1858)...	...	...	...	...	...	—	—	—	V. N.S.W.	Q.	—	—	B.fl.V,100	M.fr.I,19; IX,162.
P. decussata; F. v. M., fragm. I, 126 (1858)	...	...	...	...	...	—	—	—	V.	—	—	—	B.fl.V,100	M.fr.I,126.
P. empetrifolia; Sieber in Spreng. syst. cur. post. 226 (1827)	...	...	...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.V,101	
P. lithospermoides; F. v. M., fragm. VI, 107 (1868)	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl.V,101	M.fr.VI,107.
P. Behriana; Schlechtendal, Linnaea XX, 610 (1847)	...	...	...	...	...	—	—	—	S.A.	—	V. N.S.W.	—	B.fl.V,102	M.fr.VI,106.
P. Wilkieana; F. v. M., fragm. VIII, 230 (1874)	...	...	...	...	...	—	—	—	S.A.	—	—	—	—	M.fr.VIII,230.
P. Baxteri; Cunningham in Benth. lab. 452 (1836)	...	...	...	...	...	...	...	...	W.A.	S.A.	—	—	B.fl.V,102	M.fr.VI,106.
P. Campbelli; F. v. M. in Wing's S. Sc. Record II, 25 (1882)	...	...	...	...	...	...	...	...	W.A.	—	—	—	—	
P. canaliculata; F. v. M., fragm. VI, 105 (1868)	...	...	...	...	...	...	...	...	W.A.	—	—	—	B.fl.V,102	M.fr.VI,105.
P. nivea; Cunningham in Benth. lab. 452 (1836)	...	...	...	...	...	...	...	...	—	—	V. N.S.W.	Q.	B.fl.V,103	M.fr.VI,106.
P. striatiflora; F. v. M. in Schlechtend. Linnaea XXV, 425 (1852)	...	...	...	...	...	...	...	...	W.A.	S.A.	—	—	N.S.W. Q. N.A.	B.fl.V,103 M.fr.VI,105; IX,162.
P. saxicola; R. Brown, prodr. 509 (1810)	...	...	...	...	...	...	...	...	—	—	V. N.S.W.	Q.	B.fl.V,104	M.fr.VI,107; IX,162
P. debilis; F. v. M., fragm. VIII, 147 (1874)	...	...	...	...	...	...	...	...	—	—	V.	—	—	M.fr.VIII,147.
P. odoratissima; Benth in Mitch. Trop. Austr. 291 (1848)	...	...	...	...	...	...	...	...	—	—	—	—	Q.	B.fl.V,104
P. Eckersleyana; F. v. M., fragm. X, 17 (1876)	...	...	...	...	...	...	...	...	W.A.	—	—	—	—	M.fr.X,17.
P. euphrasioides; Benth in Mitch. Trop. Austr. 360 (1848)	...	...	...	...	...	...	...	...	—	—	—	N.S.W. Q.	B.fl.V,104	M.fr.VI,108.
P. staurophylla; F. v. M., fragm. IX, 73 (1875)	...	...	...	...	...	...	...	...	—	—	—	N.S.W.	—	M.fr.IX,73.

<i>P. cryptandroides</i> ; Cunningham in Benth. lab. 453 (1836)	—	—	—	—	N.S.W.	—	—	B.fl.V,105	
<i>P. eurybioides</i> ; F. v. M. in Transact. phil. Soc. Vict. I, 48 (1854)	—	—	S.A.	—	—	—	—	B.fl.V,105	M.fr.VI,105.
<i>P. ringens</i> ; Benth. in Mitch. Trop. Austr. 363 (1848)	—	—	S.A.	—	—	N.S.W.	Q.	B.fl.V,106	
<i>P. Leichhardtii</i> ; Benth. in Fl. Austr. V, 106 (1870)	—	—	—	—	—	—	Q.	B.fl.V,106	M.fr.VII,108.
<i>P. Walteri</i> ; F. v. M., fragm. VII, 108 (1870)	—	—	—	—	V.	N.S.W.	—	—	M.fr.VII,108.
<i>P. coccinea</i> ; F. v. M. in Transact. phil. Soc. Vict. I, 48 (1854)	—	—	W.A.	S.A.	—	V.	N.S.W.	Q.	B.fl.V,106
<i>P. chlorantha</i> ; F. v. M. in Benth. Fl. Austr. V, 108 (1870)	—	—	—	S.A.	—	V.	N.S.W.	—	B.fl.V,108
<i>P. calycina</i> ; F. v. M. in Benth. Fl. Austr. V, 107 (1870)	—	—	—	S.A.	—	—	—	—	B.fl.V,107
<i>P. Grylloana</i> ; F. v. M., fragm. X, 17 (1876)	—	—	W.A.	—	—	—	—	—	M.fr.X,17.

WRIXONIA; F. v. M., fragm. X, 18 (1876).

<i>W. prostantheroides</i> ; F. v. M., fragm. X, 18 (1876)	—	—	W.A.	—	—	—	—	—	M.fr.X,18.
---	---	---	------	---	---	---	---	---	------------

HEMIGENIA; R. Brown, prodr. 502 (1810). (Hemiandra, Colobandra, Atelandra.)

<i>H. macrantha</i> ; F. v. M., fragm. I, 210 (1859)	—	—	W.A.	—	—	—	—	B.fl.V,112	M.fr.I,210; XI,19.
<i>H. rigida</i> ; Benth. in De Candolle, prodr. XII, 563 (1848)	—	—	W.A.	—	—	—	—	B.fl.V,112	
<i>H. ramosissima</i> ; Benth. in De Candolle, prodr. XII, 565 (1848)	—	—	W.A.	—	—	—	—	B.fl.V,113	
<i>H. microphylla</i> ; Benth. in De Candolle, prodr. XII, 565 (1848)	—	—	W.A.	—	—	—	—	B.fl.V,113	M.fr.VI,112; IX,162.
<i>H. incana</i> ; Benth. in De Candolle, prodr. XII, 566 (1848)	—	—	W.A.	—	—	—	—	B.fl.V,113	
<i>H. canescens</i> ; Benth. in De Candolle, prodr. XII, 566 (1848)	—	—	W.A.	—	—	—	—	B.fl.V,114	
<i>H. podalyrina</i> ; F. v. M., fragm. VI, 112 (1868)	—	—	W.A.	—	—	—	—	B.fl.V,114	M.fr.VI,112.
<i>H. platyphylla</i> ; Benth. in De Candolle, prodr. XII, 566 (1848)	—	—	W.A.	—	—	—	—	B.fl.V,115	
<i>H. obovata</i> ; F. v. M., fragm. XI, 19 (1878)	—	—	W.A.	—	—	—	—	—	B.fl.XI,19.
<i>H. glabrescens</i> ; Benth. in De Candolle, prodr. XII, 566 (1848)	—	—	W.A.	—	—	—	—	B.fl.V,115	
<i>H. obtusa</i> ; Benth. in De Candolle, prodr. XII, 567 (1848)	—	—	W.A.	—	—	—	—	B.fl.V,115	
<i>H. sericea</i> ; Benth. in Hueg. enum. 80 (1837)	—	—	W.A.	—	—	—	—	B.fl.V,116	M.fr.VI,112.
<i>H. barbata</i> ; Bartling in Lehm. pl. Preiss. I, 360 (1845)	—	—	W.A.	—	—	—	—	B.fl.V,116	
<i>H. curvifolia</i> ; F. v. M., fragm. I, 210 (1859)	—	—	W.A.	—	—	—	—	B.fl.V,117	M.fr.I,210.
<i>H. scabra</i> ; Benth. in Fl. Austr. V, 117 (1870)	—	—	W.A.	—	—	—	—	B.fl.V,117	
<i>H. humilis</i> ; Benth. in De Candolle, prodr. XII, 567 (1848)	—	—	W.A.	—	—	—	—	B.fl.V,117	
<i>H. westringioides</i> ; Benth. in De Candolle, prodr. XII, 568 (1848)	—	—	W.A.	—	—	—	—	B.fl.V,117	
<i>H. teretiuscula</i> ; F. v. M., fragm. VI, 111 (1868)	—	—	W.A.	—	—	—	—	B.fl.V,118	M.fr.VI,111.
<i>H. brachyphylla</i> ; F. v. M., fragm. X, 19 (1876)	—	—	W.A.	—	—	—	—	—	M.fr.X,19.
<i>H. purpurea</i> ; R. Brown, prodr. 502 (1810)	—	—	—	—	—	N.S.W.	—	B.fl.V,118	M.fr.VI,112.
<i>H. cuneifolia</i> ; Benth. in Fl. Austr. V, 118 (1870)	—	—	—	—	—	N.S.W.	—	B.fl.V,118	
<i>H. Drummondii</i> ; Benth. in Fl. Austr. V, 119 (1870)	—	—	W.A.	—	—	—	—	B.fl.V,119	
<i>H. pimelifolia</i> ; F. v. M., fragm. VI, 112 (1868)	—	—	W.A.	—	—	—	—	B.fl.V,119	M.fr.VI,112.
<i>H. diplanthera</i> ; F. v. M., fragm. VI, 111 (1868)	—	—	W.A.	—	—	—	—	B.fl.V,119	M.fr.VI,111.
<i>H. pungens</i> ; F. v. M., fragm. XI, 20 (1878)	—	—	W.A.	—	—	—	—	B.fl.V,109	M.fr.XI,20.
<i>H. leiantha</i> ; F. v. M., fragm. XI, 20 (1878)	—	—	W.A.	—	—	—	—	B.fl.V,110	M.fr.XI,20.
<i>H. loganiacea</i> ; F. v. M., fragm. XI, 19 (1878)	—	—	W.A.	—	—	—	—	B.fl.V,112	M.fr.VI,113.

MICROCORYS; R. Brown, prodr. 502 (1810). (Anisandra.)

<i>M. longifolia</i> ; Benth. in De Candolle, prodr. XII, 568 (1848)	—	—	W.A.	—	—	—	—	B.fl.V,121	
<i>M. longiflora</i> ; F. v. M., fragm. VI, 113 (1868)	—	—	W.A.	—	—	—	—	B.fl.V,121	M.fr.VI,113.
<i>M. tenuifolia</i> ; Benth. in Fl. Austr. V, 122 (1870)	—	—	W.A.	—	—	—	—	B.fl.V,122	
<i>M. capitata</i> ; Benth. in De Candolle, prodr. XII, 568 (1848)	—	—	W.A.	—	—	—	—	B.fl.V,122	
<i>M. pimeloides</i> ; F. v. M., fragm. I, 156 (1859)	—	—	W.A.	—	—	—	—	B.fl.V,123	M.fr.I,156.
<i>M. subcanescens</i> ; Benth. in Fl. Austr. V, 123 (1870)	—	—	W.A.	—	—	—	—	B.fl.V,123	
<i>M. ericifolia</i> ; Benth. in De Candolle, prodr. XII, 569 (1848)	—	—	W.A.	—	—	—	—	B.fl.V,123	
<i>M. glabra</i> ; Benth. in De Candolle, prodr. XII, 569 (1848)	—	—	W.A.	—	—	—	—	B.fl.V,124	

<i>M. exserta</i> ; Benth. Fl. Austr. V, 124 (1870)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 124	
<i>M. virgata</i> ; R. Brown, prodr. 502 (1810)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 125	M.fr. VI, 112; VIII, 231.
<i>M. Macrediana</i> ; F. v. M., fragm. VIII, 231 (1874)	...	...	...	...	...	W.A.	S.A.	—	—	—	—	—	—	M.fr. VIII, 231.
<i>M. barbata</i> ; R. Brown, prodr. 502 (1810)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 125	
<i>M. lenticularis</i> ; F. v. M., fragm. VI, 113 (1868)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 126	M.fr. VI, 113.
<i>M. obovata</i> ; Benth. in De Candolle, prodr. XII, 569 (1848)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 126	M.fr. XI, 19.
<i>M. purpurea</i> ; R. Brown, prodr. 502 (1810)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. V, 126	M.fr. VI, 112.

WESTRINGIA ; Smith in Kongl. Svenska Vetenskaps Akad. Handlingar 171 (1797).

<i>W. cephalantha</i> ; F. v. M., fragm. VI, 110 (1868)	W.A.	—	—	—	—	—	—	B.fl. V, 127	M.fr. VI, 110.
<i>W. rosmariniformis</i> ; Smith in Vet. Acad. Handl. 171 (1797)	—	—	T.	V.	N.S.W.	Q.	—	B.fl. V, 128	M.fr. VI, 110; IX, 162.
<i>W. Dampieri</i> ; R. Brown, prodr. 501 (1810)	W.A.	S.A.	—	—	—	—	—	B.fl. V, 129	M.fr. VI, 110; IX, 163.
<i>W. rigida</i> ; R. Brown, prodr. 501 (1810)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl. V, 129	M.fr. IX, 163.
<i>W. angustifolia</i> ; R. Brown, prodr. 501 (1810)	—	—	T.	—	—	—	—	—	M.fr. VI, 110; IX, 163.
<i>W. senifolia</i> ; F. v. M. in Transact. phil. Soc. Vict. I, 49 (1854)	—	—	—	V.	—	—	—	B.fl. V, 130	M.fr. VI, 110; IX, 163.
<i>W. longifolia</i> ; R. Brown, prodr. 501 (1810)	—	—	—	V.	N.S.W.	Q.	—	B.fl. V, 131	M.fr. VI, 110; IX, 163.
<i>W. glabra</i> ; R. Brown, prodr. 501 (1810)	—	—	—	V.	N.S.W.	Q.	—	B.fl. V, 131	M.fr. VI, 110; IX, 163.
<i>W. rubrifolia</i> ; R. Brown, prodr. 501 (1810)	—	—	T.	—	—	—	—	B.fl. V, 131	M.fr. VI, 110.

AJUGA; Linné, gen. plant. 167 (1737), from Scribonius (1529).

A. Australis; R. Brown, prodr. 503 (1810) — S.A. T. V. N.S.W. Q. — B.f.V, 136 M.fr.VI, 109.

TEUCRIUM; Tournefort, inst. 207, t. 98 (1700), from Dioscorides.

<i>T. ajugaceum</i> ; F. v. M. & Bailey in syn. Queensland Flora, sec. suppl. 43 (1888)	—	—	—	—	—	Q.	—	—
<i>T. argutum</i> ; R. Brown, prodr. 504 (1810)	...	...	...	...	...	N.S.W.	Q.	B.fl.V, 135 M.fr.VI, 110; IX, 162.
<i>T. sessiliflorum</i> ; Benth in De Candolle, prodr. XI, 580 (1848)	...	...	W.A.	S.A.	—	V.	N.S.W.	— B.fl.V, 134 M.fr.VI, 110.
<i>T. corymbosum</i> ; R. Brown, prodr. 504 (1810)	...	...	...	...	...	—	S.A. T.	V. N.S.W. Q. — B.fl.V, 133 M.fr.VI, 110.
<i>T. fililobum</i> ; F. v. M. in Benth. Fl. Austr. V, 134 (1870)	...	...	...	...	...	W.A.	—	— — B.fl.V, 134 M.fr.IX, 162.
<i>T. integrifolium</i> ; F. v. M. in Benth. Fl. Austr. V, 133 (1870)	...	...	...	...	...	—	S.A.	— — Q. N.A. B.fl.V, 133 M.fr.IX, 162.
<i>T. racemosum</i> ; R. Brown, prodr. 504 (1810)	...	...	...	...	...	W.A.	S.A.	— V. N.S.W. Q. N.A. B.fl.V, 133 M.fr.VI, 109; IX, 162.

VERBENACEAE.

ADANSON, *Familles des plantes* II, 195 (1763), from B. de Jussieu (1759).—Genera 22, Species 82.

SPARTOTHAMNUS; Cunningham in Loudon's hort. Brit. 600 (1830).

<i>S. teucriflorus</i> ; F. v. M. in Wing's S. Sc. Rec. II, 55 (1882)	...	...	...	W.A.	S.A.	—	—	—	—	N.A.	—		
<i>S. junceus</i> ; Cunningham in Loud. hort. Brit. 600 (1830)	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.V,55	M.fr.VI,153; IX,5	... N.Z.
<i>S. puberulus</i> ; F. v. M., in Wing's S. Sc. Rec. II, 55 (1882)	...	...	...	—	S.A.	—	—	N.S.W.	Q.	—	—		

ONCINOCALYX; F. v. M. in Wing's S. Sc. Rec. III, 70 (1883).

O. Betchei; F. v. M. in Wing's S. Sc. Rec. III, 70 (1883) ...	...	...	...	...	—	—	—	—	N.S.W.	—	—	—
---	-----	-----	-----	-----	---	---	---	---	--------	---	---	---

LIPPIA ; Houston in Linné, syst. nat. 8 (1735); Linné, gen. plant. 347 (1737). (Zapania.)

L. nodiflora; Cl. Richard in Mich. fl. bor. Amer. II, 15 (1803)	...	...	...	W.A.	—	—	—	N.S.W.	Q.	N.A.	B.fl.V,35	M.fr.VI,151; IX,4	...	As., Afr., Am., P.
L. geminata; Humboldt, Bonpland et Kunth, nov. gen. et spec. pl. II, 215 (1817)	—	—	—	—	—	—	—	Q.	—	B.fl.V,35	M.fr.VI,151	...	Am.	

VERBENA; Tournefort, inst. 200, t. 94 (1700), from l'Ecluse (1576).

V. <i>officinalis</i> ; Linné, spec. plant. 11 (1753) ...	...	...	...	...	...	—	S.A. T.	V.	N.S.W.	Q.	—	B.fl. V, 36	M.fr. VI, 153; IX, 5	...	E., As., Afr.
V. <i>macrostachya</i> ; F. v. M., fragm. I, 60 (1858) ...	...	...	...	...	...	—	S.A. —	—	—	Q. N.A.	B.fl. V, 36	M.fr. I, 60; IX, 5.			

LACHNOSTACHYS; Hooker, icon. plant. t. 414 (1842). (Walcottia, Pycnolachne.)

<i>L. albicans</i> ; Hooker, icon. plant. 414 (1842) W.A.	— — — — —	B.fl. V, 38	M.fr. V, 159.
<i>L. Cliftoni</i> ; F. v. M., fragm. IX, 3 (1875) W.A.	— — — — —	—	M.fr. IX, 3.

L. verbascifolia; F. v. M., fragm. VI, 158 (1868)	W.A.	—	—	—	—	—	—	B.fl.V,38	M.fr.VI,158; IX,4.
L. ferruginea; Hooker, icon. plant. 415 (1842)	W.A.	—	—	—	—	—	—	B.fl.V,39	M.fr.VI,159.
L. Walcottii; F. v. M., fragm. II, 140 (1861)	W.A.	—	—	—	—	—	—	B.fl.V,39	M.fr.I,241; VI,159.

NEWCASTLIA; F. v. M. in Hook. Kew Misc. IX, 22 (1857).

N. cladotricha; F. v. M. in Hook. Kew Misc. IX, 22 (1857)	W.A.	S.A.	—	—	—	—	N.A.	B.fl.V,40	M.fr.I,184; III,21; VI,155; VIII,50.
N. bracteosa; F. v. M., fragm. VIII, 49 (1873)	—	S.A.	—	—	—	—	—	—	M.fr.VIII,230; IX,4.
N. chrysotricha; F. v. M., fragm. X, 15 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,15.
N. hexarrhena; F. v. M., fragm. X, 16 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,16.
N. spodiostriata; F. v. M., fragm. III, 21, t. 21 (1862)	W.A.	S.A.	—	—	—	—	N.A.	B.fl.V,40	M.fr.VIII,230; IX,4.
N. Dixoni; F. v. M. & Tate in Proceed. R. S. of S.A. X, 81 (1887)	—	S.A.	—	V.	—	—	—	—	—
N. cephalantha; F. v. M., fragm. IX, 4 (1875)	—	S.A.	—	—	—	—	—	—	M.fr.IX,4.

PHYSOPSIS; Turczaninow in Bullet. de la Soc. Imper. des Natur. de Mosc. XXII, 34 (1849).

P. spicata; Turczaninow in Bull. de Mosc. XXII, 35 (1849)	W.A.	—	—	—	—	—	—	B.fl.V,41	M.fr.VI,154.
--	------	---	---	---	---	---	---	-----------	--------------

MALLOPHORA; Endlicher in Annal. des Wien. Mus. II, 206 (1838). (Lachnocephalus.)

M. globiflora; Endlicher in Ann. des Wien. Mus. II, 206 (1838)	W.A.	—	—	—	—	—	—	B.fl.V,41	M.fr.VI,154; X,15.
---	------	---	---	---	---	---	---	-----------	--------------------

DICRASTYLIS; Drummond & Harvey in Hook. Kew Misc. VII, 56 (1855). (Dicrastyles.)

D. Doranii; F. v. M., fragm. VIII, 230 (1874)	—	S.A.	—	—	—	—	—	—	M.fr.VIII,230.
D. ochrotricha; F. v. M., fragm. IV, 161 (1864)	—	S.A.	—	—	—	—	N.A.	B.fl.V,42	M.fr.I,60; VI,155; VIII,230; IX,4.
D. Gilesii; F. v. M., fragm. VIII, 229 (1874)	—	S.A.	—	—	—	—	—	—	M.fr.X,15.
D. fulva; Drummond & Harvey in Kew Misc. VII, 56 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,43	M.fr.I,236; II,160; VI,155.
D. reticulata; Drummond & Harvey in Hook. Kew Misc. VII, 57 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,43	M.fr.VI,155; IX,4.
D. Nicholasii; F. v. M., fragm. X, 15 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,15.
D. parvifolia; F. v. M., fragm. II, 16 (1861)	W.A.	—	—	—	—	—	—	B.fl.V,43	M.fr.VI,155; X,15.
D. Beveridgei; F. v. M., fragm. VIII, 50 (1873)	—	S.A.	—	—	—	—	—	—	M.fr.IX,4.
D. Stoechas; Drummond & Harvey in Kew Misc. VII, 57 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,44	M.fr.VI,155.
D. Lewellini; F. v. M., fragm. XI, 86 (1880)	—	S.A.	—	—	—	—	—	—	M.fr.VIII,50; XI,137.

CHLOANTHES; R. Brown, prodr. Flor. Nov. Holl. 513 (1810). (Pityrodia, Quoya, Dasymalla, Depremesnilia.)

C. Stoechadis; R. Brown, prodr. 514 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.V,45	M.fr.VI,156.
C. parviflora; Walpers, repertor. bot. syst. IV, 58 (1848)	—	—	—	—	V. N.S.W.	Q.	—	B.fl.V,46	M.fr.IX,5.
C. coccinea; Bartling in Lehm. pl. Preiss. I, 352 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,48	M.fr.VI,156; IX,5.
C. salvifolia; F. v. M., Pityrodia, R. Brown, prodr. 513 (1810)	—	—	—	—	—	Q.	—	B.fl.V,48	M.fr.VI,155,255; IX,5.
C. hemigenioides; F. v. M., fragm. VI, 156 (1868)	W.A.	—	—	—	—	—	N.A.	B.fl.V,48	M.fr.VI,156.
C. uncinata; Turczaninow in Bull. de Mosc. XXXVI, 194 (1863)	W.A.	—	—	—	—	—	—	B.fl.V,48	M.fr.VI,156.
C. Bartlingii; Lehmann, Ind. sem. hort. Hamb. (1844)	W.A.	—	—	—	—	—	—	B.fl.V,49	M.fr.VI,156.
C. verbascina; F. v. M., fragm. I, 233 (1859)	W.A.	—	—	—	—	—	—	B.fl.V,50	M.fr.IV,80; VI,157; IX,5.
C. stachyodes; F. v. M., fragm. V, 50 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,50	M.fr.VI,158.
C. loxocarpa; F. v. M., fragm. II, 22 (1861)	W.A.	—	—	—	—	—	N.A.	B.fl.V,51	M.fr.IV,80; VI,156.
C. dilatata; F. v. M., fragm. VI, 157 (1868)	W.A.	—	—	—	—	—	—	B.fl.V,51	M.fr.VI,157.
C. cnneata; F. v. M., Cens. 103 (1882)	W.A.	—	—	—	—	—	—	B.fl.V,51	—
C. Oldfieldii; F. v. M., fragm. I, 234 (1859)	W.A.	—	—	—	—	—	—	B.fl.V,52	M.fr.II,182; IV,80.
C. atriplicina; F. v. M., fragm. I, 235 (1859)	W.A.	—	—	—	—	—	N.A.	B.fl.V,52	M.fr.VI,157; IX,5.
C. paniculata; F. v. M., fragm. IV, 80 (1864)	W.A.	—	—	—	—	—	N.A.	B.fl.V,53	M.fr.IV,80.
C. lepidota; F. v. M. in Wing's S. Sc. Rec. III, 3 (1883)	W.A.	—	—	—	—	—	—	—	—
C. loricata; F. v. M., fragm. X, 14 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,14.
C. halganiacea; F. v. M., fragm. X, 14 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,14.
C. Depremesnilii; Depremesnilia chrysocalyx, F. v. M. X, 59 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,59.

NESOGENES; A. de Candolle, prodr. XI, 703 (1847).

N. euphrasioides; A. de Candolle, prodr. XI, 703 (1847)	—	—	—	—	—	Q.	—	—	—
--	---	---	---	---	---	----	---	---	---

HEMIPHORA ; F. v. M., fragm. X, 13 (1876).

H. Elderi ; F. v. M., fragm. X, 13 (1876) ... W.A. — — — — — M.fr.X,13.

DENISONIA ; F. v. M. in Journ. Linn. Soc. II, 159 (1858).

D. ternifolia ; F. v. M., fragm. I, 124, t. 2 (1859) ... — — — — — N.A. B.fl.V,54 M.fr.I,124.

CYANOSTEGIA ; Turczaninow in Bull. Soc. Mosc. XXII, 35 (1849). (Bunnya.)

C. Turczaninowii ; F. v. M., fragm. VI, 154 (1868) ... W.A. — — — — — B.fl.V,54 M.fr.IX,5.
C. Bunnya ; F. v. M., fragm. V, 36 (1865) ... — — — — — N.A. B.fl.V,54 M.fr.VI,153.

CALLICARPA ; Linné in act. litter. et scien. Upsal. 80 (1741).

C. cana ; Linné, mantisa. plant. alt. 198 (1771) ... — — — — — Q. N.A. B.fl.V,56 ... Aa., P.
C. pedunculata ; R. Brown, prodr. 513 (1810) ... — — — — — N.S.W. Q. — B.fl.V,57
C. longifolia ; Lamarek, Encycl. méth. I, 562 (1783) ... — — — — — Q. — B.fl.V,57 ... As.

TATEA ; F. v. M., in Transact. Roy. Soc. of S. Austr. VI, 33 (1883).

T. acaulis ; F. v. M. in Transact. R. Soc. of S. Austr. VI, 34 (1883) ... — — — — — N.A. —

PREMNA ; Linné, mantiss. plant. alter. 154 (1771).

P. obtusifolia ; R. Brown, prodr. 512 (1810) ... — — — — — Q. N.A. B.fl.V,58 M.fr.III,36.
P. integrifolia ; Linné, mant. plant. alt. 154 (1771) ... — — — — — Q. N.A. B.fl.V,59 M.fr.VI,153 ... As., P.
P. limbata ; Benth, Fl. Austr. V, 59 (1870) ... — — — — — Q. — B.fl.V,59
P. Dallachyana ; Benth, Fl. Austr. V, 59 (1870) ... — — — — — Q. — B.fl.V,59
P. acuminata ; R. Brown, prodr. 512 (1810) ... — — — — — Q. N.A. B.fl.V,60 M.fr.III,36.

CLERODENDRUM ; Burmann, thesaur. Zeylan. 66, t. 29 (1737). (Clerodendron, Glossocarya.)

C. hemiderma ; F. v. M. in Benth. Fl. Austr. V, 61 (1870) ... — — — — — Q. — B.fl.V,61 M.fr.VI,151; IX,5.
C. inerme ; Gaertner, de fruct. I, 271, t. 57 (1788) ... — — — — — N.S.W. Q. N.A. B.fl.V,61 M.fr.VI,152 ... As., P.
C. Tracynum ; F. v. M. in Benth. Fl. Austr. V, 62 (1870) ... — — — — — Q. — B.fl.V,62 M.fr.V,61.
C. tomentosum ; R. Brown, prodr. 510 (1810) ... — — — — — N.S.W. Q. — B.fl.V,62 M.fr.VI,152.
C. lanceolatum ; F. v. M., fragm. III, 145 (1863) ... W.A. — — — — — N.A. B.fl.V,63 M.fr.III,145.
C. floribundum ; R. Brown, prodr. 511 (1810) ... — S.A. — — — — — Q. N.A. B.fl.V,63 M.fr.III,145; IX,5. P.
C. Cunninghamii ; Benth, Fl. Austr. V, 64 (1870) ... — — — — — Q. N.A. B.fl.V,64
C. costatum ; R. Brown, prodr. 511 (1810) ... — — — — — Q. — B.fl.V,64

GMELINA ; Linné, gen. plant. ed. sec. 526 (1742).

G. macrophylla ; Benth, Fl. Austr. V, 65 (1870) ... — — — — — Q. N.A. B.fl.V,65 M.fr.IV,128.
G. fasciculiflora ; Benth, Fl. Austr. V, 65 (1870) ... — — — — — Q. — B.fl.V,65 M.fr.V,35.
G. Leichhardtii ; F. v. M. in Benth. Fl. Austr. V, 66 (1870) ... — — — — — N.S.W. Q. — B.fl.V,66 M.fr.III,58,158; VI,153.

VITEX ; Tournefort, inst. 603, t. 373 (1700), from Plinius.

V. trifolia ; Linné, spec. plant. 638 (1753) ... — — — — — N.S.W. Q. N.A. B.fl.V,66 M.fr.III,59; VI,152; IX,5. Aa., P.
V. Lignum vitae ; Cunningham in De Candolle, prodr. XI, 692 (1847) ... — — — — — N.S.W. Q. — B.fl.V,67 M.fr.III,58; IX,5.
V. acuminata ; R. Brown, prodr. 512 (1810) ... — — — — — Q. N.A. B.fl.V,67 M.fr.V,34; VI,152; IX,5.
V. glabrata ; R. Brown, prodr. 512 (1810) ... — — — — — N.S.W. Q. N.A. B.fl.V,68 M.fr.III,59; V,35; VI,153.

FARADAYA ; F. v. M., fragm. phytogr. Austr. V, 21 (1865).

F. splendida ; F. v. M., fragm. V, 21 (1865) ... — — — — — Q. — B.fl.V,69 M.fr.VI,153.

AVICENNIA ; Linné, syst. nat. 8 (1735); Linné, gen. plant. 27 (1737).

A. officinalis ; Linné, spec. plant. 110 (1753) ... W.A. S.A. — V. N.S.W. Q. N.A. B.fl.V,69 ... As., Afr., Am., P., N.Z.

MYOPORINAE.

R. BROWN, prodr. 514 (1810).—Genera 2, Species 76.

EREMOPHILA; R. Brown, prodr. 518 (1810). (Pholidia, Stenochilus, Eremodendron, Pseudopholidia, Sentis, Duttonia, Pholidiposis.)

E. Mackinlayi; F. v. M., fragm. IV, 80 (1864)	...	...	...	...	W.A.	—	—	—	—	—	B.fl. V, 17	M. fr. V, 109.
E. Bowmani; F. v. M., fragm. II, 139 (1863)	...	...	...	...	—	S.A.	—	—	N.S.W.	Q.	B.fl. V, 18	M. fr. II, 139; III, 166.
E. strongylophylla; F. v. M., fragm. X, 87 (1876)	...	...	...	...	W.A.	S.A.	—	—	N.S.W.	Q.	—	M. fr. X, 87.
E. leucophylla; Benth. Fl. Austr. V, 18 (1870)	...	...	...	...	W.A.	S.A.	—	—	—	—	B.fl. V, 18	M. fr. X, 87.
E. Forrestii; F. v. M., fragm. VII, 49 (1869)	...	...	...	...	W.A.	—	—	—	—	—	B.fl. V, 18	M. fr. VII, 49.
E. eriocalyx; F. v. M., fragm. I, 236 (1859)	...	...	...	...	W.A.	—	—	—	—	—	B.fl. V, 19	M. fr. I, 236.
E. Maitlandi; F. v. M. in Benth. Fl. Austr. V, 19 (1870)	...	...	...	...	W.A.	—	—	—	—	—	B.fl. V, 19	—
E. rotundifolia; F. v. M., fragm. I, 207 (1859)	...	...	...	...	—	S.A.	—	—	—	—	B.fl. V, 19	M. fr. I, 207.
E. oppositifolia; R. Brown, prodr. 518 (1810)	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	—	B.fl. V, 20	—
E. Mitchellii; Benth. in Mitch. Trop. Austr. 31 (1848)	...	...	...	...	—	S.A.	—	—	N.S.W.	Q.	B.fl. V, 21	M. fr. VI, 148.
E. Paisleyi; F. v. M., Rep. Babb. Exped. 17 (1858)	...	...	...	...	—	S.A.	—	—	—	—	B.fl. V, 20	—
E. Sturtii; R. Brown, App. to Sturt's Exped. 22 (1849)	...	...	...	...	—	S.A.	—	—	N.S.W.	Q.	B.fl. V, 21	M. fr. VI, 148.
E. exilifolia; F. v. M., fragm. X, 88 (1876)	...	...	...	...	W.A.	S.A.	—	—	—	—	—	M. fr. X, 88.
E. Dempsteri; F. v. M., fragm. X, 60 (1876)	...	...	...	...	W.A.	—	—	—	—	—	—	M. fr. X, 60.
E. Gibsoni; F. v. M., fragm. VIII, 227 (1874)	...	...	...	...	—	S.A.	—	—	—	—	—	M. fr. VIII, 227.
E. Berryi; F. v. M., fragm. VIII, 228 (1874)	...	...	...	...	W.A.	S.A.	—	—	—	—	—	M. fr. VIII, 228; X, 88.
E. Clarkei; F. v. M., fragm. I, 208 (1859)	...	...	...	...	W.A.	—	—	—	—	—	B.fl. V, 21	M. fr. I, 208.
E. Gilesii; F. v. M., fragm. VIII, 49 (1873)	...	...	...	...	—	S.A.	—	—	—	—	—	M. fr. VIII, 49.
E. Latrobei; F. v. M. in Proceed. Roy. Soc. Tasm. III, 294 (1858)	...	...	...	...	W.A.	S.A.	—	—	N.S.W.	Q.	N.A. B.fl. V, 22	M. fr. I, 125.
E. graciliflora; F. v. M., fragm. I, 208 (1858)	...	...	...	...	W.A.	—	—	—	—	—	B.fl. V, 23	M. fr. II, 181.
E. longifolia; F. v. M. in Proceed. Roy. Soc. Tasm. III, 295 (1858)	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A. B.fl. V, 23	M. fr. VI, 148.
E. Laanii; F. v. M. in Melb. Chem. Jan. (1885)	...	...	...	...	W.A.	—	—	—	—	—	—	—
E. Drummondii; F. v. M., fragm. VI, 147 (1868)	...	...	...	...	W.A.	—	—	—	—	—	B.fl. V, 24	M. fr. VI, 147.
E. Hughesii; F. v. M., fragm. VIII, 228 (1874)	...	...	...	...	W.A.	S.A.	—	—	—	—	—	M. fr. VIII, 228.
E. polyclada; F. v. M. in Proceed. Roy. Soc. Tasm. III, 294 (1858)	...	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	B.fl. V, 24	—
E. bignoniflora; F. v. M. in Proceed. Roy. Soc. Tasm. III, 294 (1858)	...	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	N.A. B.fl. V, 25	M. fr. VI, 148.
E. Freelingii; F. v. M. in Proceed. Roy. Soc. Tasm. III, 295 (1858)	...	...	...	...	W.A.	S.A.	—	—	N.S.W.	—	B.fl. V, 25	M. fr. VI, 148.
E. Fraseri; F. v. M., fragm. XI, 51 (1878)	...	...	...	...	W.A.	—	—	—	—	N.A.	—	M. fr. XI, 51.
E. Macdonnellii; F. v. M., Rep. Babb. Exped. 18 (1858)	...	...	...	...	—	S.A.	—	—	N.S.W.	Q.	N.A. B.fl. V, 25	M. fr. V, 23; VI, 150.
E. Goodwinii; F. v. M., Rep. Babb. Exped. 17 (1858)	...	...	...	...	—	S.A.	—	—	N.S.W.	Q.	N.A. B.fl. V, 25	M. fr. VI, 148.
E. Elderi; F. v. M., fragm. VIII, 228 (1874)	...	...	...	...	W.A.	S.A.	—	—	—	—	—	M. fr. VIII, 228.
E. Willsii; F. v. M., fragm. III, 21, t. 20 (1862)	...	...	...	...	W.A.	S.A.	—	—	—	Q.	B.fl. V, 26	M. fr. III, 21.
E. platycalyx; F. v. M., fragm. V, 109 (1866)	...	...	...	...	W.A.	—	—	—	—	—	B.fl. V, 26	M. fr. V, 109.
E. viscida; Endlicher, nov. stirp. dec. 51 (1839)	...	...	...	...	W.A.	—	—	—	—	—	B.fl. V, 26	—
E. Brownii; F. v. M. in Proceed. Roy. Soc. Tasm. III, 297 (1858)	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	B.fl. V, 27	M. fr. VI, 148.
E. subfloccosa; Benth. Fl. Austr. V, 28 (1870)	...	...	...	...	W.A.	—	—	—	—	—	B.fl. V, 28	—
E. Oldfieldii; F. v. M., fragm. I, 208 (1859)	...	...	...	...	W.A.	—	—	—	—	—	B.fl. V, 28	M. fr. I, 208.
E. Duttonii; F. v. M., Rep. Babb. Exped. 16 (1858)	...	...	...	...	—	S.A.	—	—	N.S.W.	—	B.fl. V, 28	M. fr. VI, 148.
E. maculata; F. v. M. in Proceed. Roy. Soc. Tasm. III, 297 (1858)	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A. B.fl. V, 29	M. fr. I, 125.
E. denticulata; F. v. M., fragm. I, 125 (1859)	...	...	...	...	W.A.	S.A.	—	—	—	—	B.fl. V, 29	M. fr. IV, 170.
E. latifolia; F. v. M. in Schlechtend. Linnaea XXV, 428 (1852)	...	...	...	...	W.A.	S.A.	—	—	N.S.W.	—	B.fl. V, 30	M. fr. V, 109; VI, 148.
E. alternifolia; R. Brown, prodr. 518 (1810)	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	—	B.fl. V, 30	M. fr. V, 109; VI, 148.
E. Dalyana; F. v. M., fragm. V, 22 (1865)	...	...	...	...	—	S.A.	—	—	N.S.W.	—	B.fl. V, 10	M. fr. V, 22.
E. Pantoni; F. v. M. in Wing's S. Sc. Rec. II, 25 (1882)	...	...	...	...	—	—	—	—	—	N.A.	—	—
E. scoparia; F. v. M. in Proceed. Roy. Soc. Tasm. III, 296 (1858)	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	—	B.fl. V, 10	M. fr. V, 22; VI, 148.
E. Delisserii; F. v. M., fragm. V, 108, t. 42 (1865)	...	...	...	...	W.A.	S.A.	—	—	—	—	B.fl. V, 11	M. fr. V, 189.

<i>E. crassifolia</i> ; F. v. M. in Proceed. Roy. Soc. Tasm. III, 297 (1858) ...	...	—	S.A.	—	—	—	—	—	B.fl.V,11	M.fr.VI,148.
<i>E. resinosa</i> ; F. v. M. in Proceed. Roy. Soc. Tasm. III, 296 (1858) ...	...	W.A.	—	—	—	—	—	—	B.fl.V,11	
<i>E. Behrii</i> ; F. v. M. in Proceed. Roy. Soc. Tasm. III, 296 (1858)...	...	—	S.A.	—	—	—	—	—	B.fl.V,12	
<i>E. Woollsiana</i> ; F. v. M., fragm. I, 125, t. 7 (1859) ...	...	W.A.	—	—	—	—	—	—	B.fl.V,12	M.fr.VI,151.
<i>E. Christophori</i> ; F. v. M., fragm. IX, 120 (1875) ..	...	—	S.A.	—	—	—	—	—	—	M.fr.IX,198.
<i>E. brevifolia</i> ; F. v. M., fragm. I, 126 (1859) ...	...	W.A.	—	—	—	—	—	—	B.fl.V,12	M.fr.VI,150.
<i>E. Weldii</i> ; F. v. M., fragm. VII, 109 (1870) ...	...	W.A.	S.A.	—	—	—	—	—	—	M.fr.IX,120.
<i>E. densifolia</i> ; F. v. M., fragm. II, 160 (1861) ...	...	W.A.	—	—	—	—	—	—	B.fl.V,13	M.fr.II,160.
<i>E. gibbosifolia</i> ; F. v. M., Rep. Babb. Exped. 18 (1858) ...	...	—	S.A.	—	V.	—	—	—	B.fl.V,13	M.fr.VI,150.
<i>E. divaricata</i> ; F. v. M. in Proceed. Roy. Soc. Tasm. III, 293 (1858) ...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.V,14	M.fr.IV,48;VI,150.
<i>E. microtheca</i> ; F. v. M. in Benth. Fl. Austr. V, 14 (1870) ...	...	W.A.	—	—	—	—	—	—	B.fl.V,14	M.fr.II,160.
<i>E. adenotricha</i> ; F. v. M. in Benth. Fl. Austr. V, 15 (1870) ...	...	W.A.	—	—	—	—	—	—	B.fl.V,14	
<i>E. santalina</i> ; F. v. M. in Proceed. Roy. Soc. Tasm. III, 295 (1858) ...	...	—	S.A.	—	—	—	—	—	B.fl.V,15	
<i>E. Youngii</i> ; F. v. M., fragm. X, 16 (1876) ...	...	W.A.	—	—	—	—	—	—	—	M.fr.X,16.

MYOPORUM; Banks & Solander in G. Forster, prodr. 44 (1786). (Bertolonia, Andrewsia, Pogonia, Disoon.)

<i>M. tenuifolium</i> ; G. Forster, fl. ins. Austr. prodr. 44 (1786) ...	...	—	—	—	V.	N.S.W.	Q.	N.A.	B.fl.V,4	M.fr.VI,149	... P.
<i>M. obscurum</i> ; Endlicher, prodr. fl. Norf. 54 (1833)...	...	—	—	—	—	N.S.W.	—	—	—		
<i>M. Dampieri</i> ; Cunningham in De Candolle, prodr. XI, 708 (1847) ...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V,4	M.fr.VI,149.	
<i>M. deserti</i> ; Cunningham in Hueg. enum. 78 (1837) ...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.V,5	M.fr.VI,148; VII,110.	
<i>M. laxiflorum</i> ; Benth. Fl. Austr. V, 6 (1870) ...	...	—	—	—	—	—	Q.	—	B.fl.V,6	M.fr.V,23.	
<i>M. insulare</i> ; R. Brown, prodr. 515 (1810) ...	...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.V,6	M.fr.VI,149; VII,110.	
<i>M. viscosum</i> ; R. Brown, prodr. 516 (1810) ...	...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.V,5	M.fr.VII,109.	
<i>M. serratum</i> ; R. Brown, prodr. 516 (1810) ...	...	W.A.	—	—	—	—	—	—	B.fl.V,4	M.fr.VI,149; VII,110.	
<i>M. oppositifolium</i> ; R. Brown, prodr. 516 (1810) ...	...	W.A.	—	—	—	—	—	—	B.fl.V,7	M.fr.VI,149.	
<i>M. brevipes</i> ; Benth. Fl. Austr. V, 6 (1870) ...	...	—	S.A.	—	—	—	—	—	B.fl.V,6		
<i>M. humile</i> ; R. Brown, prodr. 516 (1810) ...	...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.V,6	M.fr.VI,150.	
<i>M. debile</i> ; R. Brown, prodr. 516 (1810) ...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.V,6	M.fr.VI,149.	
<i>M. platycarpum</i> ; R. Brown, prodr. 516 (1810) ...	...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.V,7	M.fr.VI,150.	
<i>M. floribundum</i> ; Cunningham in Hueg. enum. 78 (1837) ...	...	—	—	—	—	V.	N.S.W.	—	B.fl.V,8	M.fr.I,126; VI,150.	
<i>M. Beckeri</i> ; F. v. M. in Benth. Fl. Austr. V, 7 (1870) ...	...	W.A.	—	—	—	—	—	—	B.fl.V,7	M.fr.I,156,244; VI,150.	
<i>M. salsoloides</i> ; Turczaninow in Bull. de Mosc. XXXVI, 226 (1863) ...	...	W.A.	—	—	—	—	—	—	B.fl.V,8	M.fr.I,126; VI,150.	

ASPERIFOLIAE.

HALLER, enum. stirp. Helv. Praef. 34 (1742).—Genera 12, Species 52.

CYNOGLOSSUM; Tournefort, inst. 139, t. 57 (1700) from Dioscorides and Nicandros.

<i>C. latifolium</i> ; R. Brown, prodr. 495 (1810) ...	...	—	—	T.	V.	N.S.W.	Q.	—	B.fl.IV,408	M.fr.VI,115; IX,122; XI,125.
<i>C. suaveolens</i> ; R. Brown, prodr. 495 (1810) ...	...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,409	M.fr.VI,115; IX,122.
<i>C. Australe</i> ; R. Brown, prodr. 495 (1810) ...	...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,409	M.fr.VI,115; IX,122.
<i>C. Drummondii</i> ; Benth. Fl. Austr. IV, 409 (1869) ...	...	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.IV,409	M.fr.IX,122.

ROCHELIA; Reichenbach in Fl. Regensb. bot. Zeit. I, 243 (1824). (Maccoya.)

<i>R. Maccoya</i> ; F. v. M. in Benth. Fl. Austr. IV, 408 (1869) ...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.IV,408	M.fr.I,127; IX,121.
--	-----	---	------	---	----	--------	----	---	-------------	---------------------

LAPPULA; Rivinus in Rupp. Fl. Jenens. (1718), from Dalechamp. (Echinosperrum.)

<i>L. concava</i> ; F. v. M., Cens. 100 (1882) ...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.IV,407	M.fr.VI,116; IX,121.
--	-----	------	------	---	----	--------	----	---	-------------	----------------------

ERITRICHUM; Schrader in commentat. Goetting. IV, 186 (1820). (Eritricium, Lappulae subgenus.)

<i>E. Australasicum</i> ; A. de Candolle, prodr. X, 134 (1846) ...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.IV,406	M.fr.XI,125.
--	-----	------	------	---	----	--------	----	---	-------------	--------------

MYOSOTIS; Ruppius, Fl. Jenens. 9 (1718), from Dioscorides and Plinius. (Exarrhena.)

M. Australis; R. Brown, prodr. 495 (1810)	...	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.IV,405	M.fr.VI,116; IX,122	...	N.Z.
M. exarrhena; F. v. M., inedit.	...	...	...	...	...	—	—	T.	V.	N.S.W.	—	—	B.fl.IV,406	M.fr.VI,115; IX,122.		

POLLICHIA; Medikus, Bot. Beobach. 247 (1783). (Trichodesma.)

P. Zeylanica; F. v. M., Cens. 100 (1882) from N. Burmann (1768)	...	...	...	...	...	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.IV,404	M.fr.VI,115; IX,122; XI,125.	As., Afr., P.
P. latiseppalea; F. v. M., Cens. 100 (1882)	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	—	M.fr.X,102,121.	

HALGANIA; Gaudichaud in Freyc. voy. Bot. 448 (1826).

H. solanacea; F. v. M. in Hook. Kew Misc. IX, 21 (1857)	...	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.IV,401	M.fr.VI,115.	
H. litoralis; Gaudichaud in Freyc. voy. Bot. 449, t. 59 (1826)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,401	M.fr.VI,115.	
H. corymbosa; Lindley, Bot. Regist. XXV, App. 40 (1839)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,402		
H. sericiflora; Benth. Fl. Austr. IV, 402 (1869)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,402		
H. cyanea; Lindley, Bot. Regist. XXV, App. 40 (1839)	...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.IV,402	M.fr.VI,115; IX,122.	
H. lavandulacea; Endlicher in Ann. Wien. Mus. II, 205 (1838)	...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.IV,402	M.fr.I,209; VI,115; XI,125.	
H. integerrima; Endlicher in Ann. Wien. Mus. II, 205 (1838)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,402		

HELIOTROPIUM; Tournefort, inst. 138, t. 57 (1700) from Theophrastos, Dioscorides & Plinius. (Tiaridium, Heliophytum, Schleidenia restorable.)

H. Indicum; Linné, spec. plant. 130 (1753)	...	...	...	...	...	—	—	—	—	—	Q. N.A.	—	M.fr.XI,124	...	...	As., Afr., Am.
H. pleiopterum; F. v. M., fragm. IX, 121 (1875)	...	...	...	...	...	—	S.A.	—	—	—	—	—	M.fr.IX,121.			
H. Curassavicum; Linné, spec. plant. 130 (1753)	...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q. N.A.	B.fl.IV,393	M.fr.VI,116; IX,121	...	Afr., Am., P.	
H. Europaeum; Linné, spec. plant. 130 (1753)	...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.IV,394	M.fr.VI,116	...	E., As., Afr.
H. undulatum; Vahl, symbol. bot. I, 13 (1795)	...	...	...	...	...	W.A.	S.A.	—	—	—	—	N.A.	B.fl.IV,394	M.fr.VI,116; IX,121; XI,124.	As., Afr.	
H. asperinum; R. Brown, prodr. 493 (1810)	...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q. N.A.	B.fl.IV,394	M.fr.VI,116; IX,121.			
H. crispatum; F. v. M. in Benth. Fl. Austr. IV, 395 (1869)	...	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.IV,395			
H. fasciculatum; R. Brown, prodr. 494 (1810)	...	...	...	...	...	—	—	—	—	—	Q. N.A.	B.fl.IV,395				
H. vestitum; Benth. Fl. Austr. IV, 395 (1869)	...	...	...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.IV,395			
H. brachygyne; Benth. Fl. Austr. IV, 396 (1869)	...	...	...	...	...	—	—	—	—	—	Q. —	B.fl.IV,396				
H. epacrideum; F. v. M. in Benth. Fl. Austr. IV, 396 (1869)	...	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.IV,396			
H. ovalifolium; Forskael, Fl. Aegypt. Arab. 47 (1775)	...	...	...	...	...	—	S.A.	—	—	—	Q. N.A.	B.fl.IV,396	M.fr.VI,115	...	...	As., Afr.
H. strigosum; Willdenow, sp. pl. I, 743 (1797)	...	...	...	...	...	—	—	—	—	—	Q. N.A.	B.fl.IV,397	...	...	...	As., Afr.
H. prostratum; R. Brown, prodr. 494 (1810)...	...	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.IV,397			
H. bracteatum; R. Brown, prodr. 493 (1810)	...	...	...	...	...	—	—	—	—	—	Q. N.A.	B.fl.IV,397				
H. pauciflorum; R. Brown, prodr. 493 (1810)	...	...	...	...	...	—	—	—	—	—	Q. N.A.	B.fl.IV,398				
H. conocarpum; F. v. M. in Benth. Fl. Austr. 398 (1869)	...	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.IV,398			
H. filaginoides; Benth. Fl. Austr. IV, 398 (1869)	...	...	...	...	...	—	S.A.	—	—	—	—	—	B.fl.IV,398			
H. ventricosum; R. Brown, prodr. 494 (1810)	...	...	...	...	...	—	—	—	—	—	Q. N.A.	B.fl.IV,399				
H. paniculatum; R. Brown, prodr. 494 (1810)	...	...	...	...	...	W.A.	S.A.	—	—	—	Q. N.A.	B.fl.IV,399				
H. Cunninghamii; Benth. Fl. Austr. IV, 400 (1869)	...	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.IV,400			
H. diversifolium; F. v. M. in Benth. Fl. Austr. IV, 400 (1869)	...	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.IV,400			

COLDENIA; Linné, amoen. acad. ed. prim. 119 (1747). (Lobophyllum.)

C. procumbens; Linné, spec. plant. 125 (1753)	...	...	...	...	...	—	S.A.	—	—	—	Q.	N.A.	B.fl.IV,391	M.fr.VI,115	...	...	As., Afr.
---	-----	-----	-----	-----	-----	---	------	---	---	---	----	------	-------------	-------------	-----	-----	-----------

TOURNEFORTIA; Linné, syst. nat. 8 (1735); Linné, gen. plant. 55 (1737).

T. argentea; Linné, fil. suppl. plant. 133 (1781)	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.IV,389	M.fr.IV,95; VI,116	...	As., Afr., P.
T. mollis; F. v. M., fragm. I, 59 (1858)	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.IV,390	M.fr.VI,116.		
T. sarmentosa; Lamarck, Illustr. des genr. I, 416 (1791)	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl.IV,390	M.fr.IV,95; VI,116	...	P.

EHRETIA; P. Browne, Civ. and Nat. Hist. of Jamaic. 168, t. 16 (1756).

E. acuminata; R. Brown, prodr. 497 (1810)	...	...	...	...	...	—	—	—	V.	N.S.W.	Q.	—	B.fl.IV,387	M.fr.V,21; IX,122; XI,124.	As.
E. pilosula; F. v. M., fragm. V, 20 (1865)	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl.IV,388	M.fr.V,20.	
E. saligna; R. Brown, prodr. 497 (1810)	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.IV,388	M.fr.V,21; IX,122.	

E. membranifolia; R. Brown, prodr. 497 (1810)	— — — — N.S.W.	Q. —	B.fl.IV,388	M.fr.V,21; VI,115; IX,122.	
E. laevis; Roxburgh, Pl. Corom. I, 42, t. 56 (1795)	— — — — —	Q. —	B.fl.IV,389		As.
CORDIA; Plumier, gen. plant. 13, t. 14 (1703).					
C. subcordata; Lamarck, Illustr. des geur. I, 421 (1791)	— — — — —	Q. N.A.	B.fl.IV,385	M.fr.VI,114; IX,122	.. As., Afr., P.
C. aspera; G. Forster, florul. ins. Austr. prodr. 18 (1786)	— — — — —	Q. —	B.fl.IV,386	M.fr.V,193; VI,114; IX,122.	P.
C. Myxa; Linné, spec. plant. 190 (1753)	— — — — —	Q. —	B.fl.IV,386	M.fr.I,59; VI,114...	.. As., Afr., P.

ERICACEAE.

A. L. DE JUSSIEU, Gen. plant. 159 (1789)—Genera 5, Species 7.

RHODODENDRON; Linné, spec. plant. I, 392 (1753).					
R. Lochar; F. v. M. in Vict. Naturalist III, 157 (1887)	— — — — —	Q. —	—	—	
PERNETTYA; Gaudichaud in Ann. des sc. nat. V, 102 (1825).					
P. Tasmanica; J. Hooker in Lond. Journ. of Bot. VI, 268 (1847)	— — — — T.	— — — —	B.fl.IV,140		N.Z.
GAULTIERA; Kalm in Linné, amoen. acad. III, 14, fig. 6 (1751). (Gaultheria.)					
G. hispida; R. Brown, prodr. 559 (1810)	— — — — T.	V. N.S.W.	— —	B.fl.IV,141	
G. lanceolata; J. Hooker in Lond. Journ. of Bot. VI, 267 (1847)... ..	— — — — T.	— — — —	— —	B.fl.IV,141	
G. antipoda; G. Forster, florul. ins. Austr. pr. 34 (1786)	— — — — T.	— — — —	— —	B.fl.IV,142	 N.Z.
AGAPETES; D. Don in G. Don, gen. syst. of dich. plants III, 862 (1834). (Vaccinii subgenus.)					
A. Meiniana; F. v. M. in Vict. Naturalist III, 158 (1887)... ..	— — — — —	Q. —	—	—	
WITTSTEINIA; F. v. M., fragm. II, 136 (1861).					
W. vacciniacea; F. v. M., fragm. II, 136 (1861)	— — — — V.	— — — —	B.fl.IV,139	M.fr.III,166.	

EPACRIDAE.

R. BROWN, prodr. 535 (1810).—Genera 18, Species 275.

STYPHELIA; Solander in G. Forster, fl. ins. Austr. prodr. 13 (1786). (Epacris, Forst. (1776) partly, Stiphelia, Ardisia, Peroa, Perojoa, Ventenatia, Cyathodes, Stenanthera, Astroloma, Leucopogon, Melichrus, Acrotiche, Monotoca, Soleniscia, Stomarrhena, Pentataphrus, Mesotriche, Phanerandra, Froebelina, Pentaptelion, Androstoma, Cyathopsis, Lissanthe partly.)					
S. adscendens; R. Brown, prodr. 537 (1810)	— S.A.	T. V. N.S.W.	— —	B.fl.IV,146	M.fr.VI,36; XI,122.
S. longifolia; R. Brown, prodr. 537 (1810)	— — — —	N.S.W.	— —	B.fl.IV,147	
S. laeta; R. Brown, prodr. 537 (1810)	— — — —	N.S.W.	— —	B.fl.IV,147	
S. triflora; Andrews, Bot. Reposit. t. 72 (1799)	— — — —	N.S.W.	Q. —	B.fl.IV,147	M.fr.XI,122.
S. viridis; Andrews, Bot. Reposit. t. 312 (1806)	— — — —	N.S.W.	Q. —	B.fl.IV,148	
S. Hainesii; F. v. M., fragm. IV, 96, t. 28 (1864)	W.A.	— — — —	— —	B.fl.IV,148	M.fr.VIII,55.
S. tubiflora; Smith, Specim. of the Bot. of New Holl. 45, t. 14 (1793)	— — — —	N.S.W.	— —	B.fl.IV,148	
S. tenuiflora; Lindley, Bot. Regist. XXV, App. 25 (1839)... ..	W.A.	— — — —	— —	B.fl.IV,148	M.fr.IV,97; VI,46; VIII,54.
S. melaleucoides; F. v. M., fragm. IV, 97 (1864)	W.A.	— — — —	— —	B.fl.IV,149	M.fr.VI,30.
S. pusilliflora; F. v. M., fragm. IV, 105 (1864)	— S.A.	— — — —	— —	B.fl.IV,149	M.fr.I,178; VI,31.
S. Leucopogon; F. v. M., fragm. IV, 97 (1864)	W.A.	— — — —	— —	B.fl.IV,149	M.fr.III,143; VI,31.
S. lasionema; F. v. M., fragm. VI, 40 (1867)... ..	W.A.	— — — —	— —	B.fl.IV,153	M.fr.VIII,54.
S. macrocalyx; F. v. M., fragm. VI, 37 (1867)	W.A.	— — — —	— —	B.fl.IV,153	M.fr.VI,37.
S. xerophylla; F. v. M., fragm. VI, 38 (1867)	W.A.	— — — —	— —	B.fl.IV,153	M.fr.VI,38.
S. pentapogonea; F. v. M., fragm. VI, 36 (1867)	W.A.	— — — —	— —	B.fl.IV,153	M.fr.VI,36.
S. prostrata; F. v. M., Cens. 105 (1882) from R. Brown (1810)	W.A.	— — — —	— —	B.fl.IV,154	
S. tecta; Sprengel, syst. I, 657 (1825)... ..	W.A.	— — — —	— —	B.fl.IV,154	M.fr.VI,37.

S. Candolleana; F. v. M., fragm. VI, 38 (1867)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV, 154	M.fr. VIII, 54.
S. microdonta; F. v. M., Cens. 105 (1882)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV, 155	
S. pallida; Sprengel, syst. I, 658 (1825)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV, 155	M.fr. VI, 37.
S. compacta; Sprengel, syst. I, 657 (1825)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV, 155	M.fr. VI, 38.
S. humifusa; Persoon, synops. plant. I, 174 (1805)	...	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	—	B.fl.IV, 156	M.fr. VI, 37; VIII, 54.
S. Epacridis; F. v. M., fragm. VI, 38 (1867)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV, 156	M.fr. VI, 38.
S. Drummondii; F. v. M., fragm. VI, 37 (1867)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV, 157	M.fr. VI, 37.
S. microcalyx; F. v. M., fragm. VI, 37 (1867)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV, 157	M.fr. VI, 37.
S. Baxteri; F. v. M., fragm. VI, 37 (1867)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV, 158	M.fr. IV, 97.
S. Sonderi; F. v. M., fragm. VI, 36 (1867)	...	...	...	...	...	—	S.A.	—	V.	N.S.W.	—	B.fl.IV, 158	M.fr. VIII, 54; IX, 48; XI, 122.
S. longiflora; F. v. M., fragm. VIII, 54 (1873)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV, 158	M.fr. VIII, 54.
S. pinifolia; Sprengel, syst. veg. I, 659 (1825)	...	...	...	...	...	—	—	T.	V.	N.S.W.	—	B.fl.IV, 159	M.fr. IV, 99; VI, 36.
S. procumbens; Persoon, synops. I, 174 (1805)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	B.fl.IV, 162	M.fr. VI, 38; VIII, 55.
S. urceolata; F. v. M., fragm. VI, 38 (1867)	...	...	...	...	...	—	—	—	V.	N.S.W.	Q.	B.fl.IV, 162	M.fr. VIII, 54.
S. Billardieri; F. v. M., fragm. VI, 43 (1867)	...	...	...	...	...	—	—	T.	—	—	—	B.fl.IV, 169	M.fr. VIII, 54.
S. Milligani; F. v. M., Pentschondra verticillata, J. Hook. Fl. Tas. I, 256 (1860)	...	...	...	...	...	—	—	T.	—	—	—	B.fl.IV, 164	
S. straminea; Sprengel, syst. I, 656 (1825)	...	...	...	...	...	—	—	T.	—	—	—	B.fl.IV, 169	M.fr. VI, 43.
S. Hookeri; F. v. M., fragm. VI, 44 (1867)	...	...	...	...	...	—	—	T.	—	—	—	B.fl.IV, 169	M.fr. VI, 44; VIII, 54.
S. dealbata; Sprengel, syst. I, 659 (1825)	...	...	...	...	...	—	—	T.	—	—	—	B.fl.IV, 170	M.fr. VI, 43; VIII, 54.
S. abietina; Labillardière, Nov. Holl. pl. spec. I, 48, t. 68 (1804)	...	...	...	...	...	—	—	T.	—	—	—	B.fl.IV, 170	M.fr. VI, 43.
S. Oxycedrus; Labillardière, Nov. Holl. pl. spec. I, 49, t. 69 (1804)	...	...	...	...	...	—	—	T.	V.	—	—	B.fl.IV, 170	M.fr. IV, 100; VI, 43 ... N.Z.
S. parvifolia; F. v. M. in papers of R. S. Tasm. 86 (1874)	...	...	...	...	...	—	—	T.	—	—	—	B.fl.IV, 171	
S. sapida; F. v. M., fragm. VI, 42 (1867)	...	...	...	...	...	—	—	—	—	N.S.W.	—	B.fl.IV, 175	M.fr. VI, 42; VIII, 54.
S. strigosa; Smith, Specim. of the Bot. of New Holl. 49 (1793)	...	...	...	...	...	—	S.A.	T.	V.	N.S.W.	Q.	B.fl.IV, 176	M.fr. VI, 42; IX, 48; XI, 122.
S. verticillata; Sprengel, syst. I, 656 (1825)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV, 184	M.fr. IV, 122; VI, 43; VIII, 55.
S. interrupta; Sprengel, syst. I, 656 (1825)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV, 184	
S. amplexicaulis; Rudge in Transact. Linn. Soc. VIII, 292, t. 8 (1805)	...	...	...	...	...	—	—	—	—	N.S.W.	—	B.fl.IV, 185	M.fr. VI, 44.
S. alternifolia; Sprengel, syst. I, 655 (1825)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV, 185	
S. lanceolata; Smith, Specim. of the Bot. of New Holl. 49 (1793)	...	...	...	...	...	—	—	T.	V.	N.S.W.	Q.	B.fl.IV, 185	M.fr. IV, 124; VI, 43.
S. Richei; Labillardière, Nov. Holl. pl. spec. I, 44, t. 60 (1804)	...	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	B.fl.IV, 186	M.fr. IV, 123; VI, 42; VIII, 54. N.Z.
S. Australis; F. v. M., fragm. VI, 43 (1867)	...	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	—	B.fl.IV, 186	M.fr. IV, 123.
S. capitellata; F. v. M., fragm. VI, 31 (1867)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV, 187	M.fr. VI, 31.
S. revoluta; Sprengel, syst. I, 657 (1825)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV, 187	M.fr. VI, 31.
S. grandiuscula; F. v. M., fragm. VI, 47 (1867)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV, 188	M.fr. VI, 47.
S. reflexa; Sprengel, syst. I, 655 (1825)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV, 188	M.fr. VI, 32.
S. corifolia; F. v. M., Cens. 105 (1882) from Endlicher (1839)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV, 188	
S. distans; Sprengel, syst. I, 655 (1825)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV, 189	M.fr. VI, 32.
S. gibbosa; F. v. M., Cens. 106 (1882) from Stschegleew (1859)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV, 189	
S. thymifolia; F. v. M., Cens. 106 (1882) from Lindley (1869)	...	...	...	...	...	—	—	—	V.	—	—	B.fl.IV, 189	
S. cordata; F. v. M., Cens. 106 (1882) from Sonder (1845)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV, 190	
S. Bossiaea; F. v. M., fragm. VI, 47 (1867)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV, 190	M.fr. VI, 47.
S. hirsuta; F. v. M., fragm. VI, 31 (1867)	...	...	...	...	...	W.A.	S.A.	—	—	—	—	B.fl.IV, 191	M.fr. VI, 31.
S. collina; Labillardière, Nov. Holl. pl. spec. I, 47, t. 65 (1804)	...	...	...	...	...	—	S.A.	T.	V.	N.S.W.	—	B.fl.IV, 191	M.fr. VI, 45; VIII, 54.
S. glacialis; F. v. M., Cens. 106 (1882) from Lindley (1838)	...	...	...	...	...	—	—	—	V.	—	—	B.fl.IV, 191	
S. compacta; F. v. M., Cens. 106 (1882) from Stschegl. (1859)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV, 192	
S. squarrosa; F. v. M., fragm. VI, 31 (1867)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV, 192	M.fr. VI, 31.
S. microphylla; Sprengel, syst. I, 656 (1825)	...	...	...	...	...	—	—	—	V.	N.S.W.	—	B.fl.IV, 192	M.fr. VI, 31.
S. tetragona; F. v. M., fragm. VI, 31 (1867)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV, 193	M.fr. VI, 31.
S. phyllostachys; F. v. M., Cens. 106 (1882) from Bentham (1869)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV, 193	M.fr. VI, 32.
S. glabella; Sprengel, syst. I, 655 (1825)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV, 194	M.fr. VI, 32.
S. semiopposita; F. v. M., fragm. VI, 49 (1867)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.IV, 194	M.fr. VI, 49.

<i>S. florulenta</i> ; F. v. M., Cens. 106 (1882) from Benth. (1869) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,194
<i>S. striata</i> ; Sprengel, syst. I, 656 (1825) ...	...	...	W.A.	S.A.	—	—	—	—	—	B.fl.IV,194
<i>S. costata</i> ; F. v. M. in Wing's S. Sc. Rec. I, new ser. (1885) ...	...	...	—	S.A.	—	V.	—	—	—	—
<i>S. lasiostachya</i> ; F. v. M., Cens. 106 (1882) from Stschegl. (1859)...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,195
<i>S. carinata</i> ; Sprengel, syst. I, 658 (1825) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,195
<i>S. opponens</i> ; F. v. M., fragm. VI, 48 (1867) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,196 M.fr.VI,48.
<i>S. oppositifolia</i> ; F. v. M., fragm. VI, 32 (1867) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,196 M.fr.VI,32.
<i>S. tamariscina</i> ; Sprengel, syst. I, 656 (1825)...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,197
<i>S. bracteolaris</i> ; F. v. M., Cens. 106 (1882) from Benth. (1869)...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,197
<i>S. blepharophylla</i> ; F. v. M., fragm. VI, 34 (1867) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,197 M.fr.VI,34.
<i>S. tenuis</i> ; F. v. M., Cens. 106 (1882) from De Candolle (1833) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,197
<i>S. gnaphalioides</i> ; F. v. M., Cens. 106 (1882) from Stschegl. (1859) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,198
<i>S. concurva</i> ; F. v. M., fragm. VI, 36 (1867) ...	...	...	—	S.A.	—	—	—	—	—	B.fl.IV,198 M.fr.III,144.
<i>S. Gilbertii</i> ; F. v. M., Cens. 106 (1882) from Stschegl. (1859) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,198
<i>S. gracilis</i> ; Sprengel, syst. I, 658 (1825) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,199
<i>S. acicularis</i> ; F. v. M., Cens. 106 (1882) from Benth. (1869) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,199
<i>S. cryptantha</i> ; F. v. M., Cens. 106 (1882) from Benth. (1869) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,199
<i>S. gracillima</i> ; F. v. M., fragm. VI, 34 (1867) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,200 M.fr.VI,34.
<i>S. cymbiformis</i> ; F. v. M., fragm. VI, 34 (1867) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,200 M.fr.VI,34.
<i>S. apiculata</i> ; Sprengel, syst. I, 656 (1825) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,201 M.fr.III,144; IV,105; VI,31.
<i>S. polystachya</i> ; Sprengel, syst. I, 659 (1825) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,201 M.fr.VI,31.
<i>S. virgata</i> ; Labillardiere, Nov. Holl. pl. spec. I, 46, t. 64 (1804)...	...	...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,201 M.fr.VI,42.
<i>S. pulchella</i> ; F. v. M., fragm. VI, 34 (1867) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,202 M.fr.VI,34.
<i>S. polymorpha</i> ; F. v. M., fragm. VI, 31 (1867) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,202 M.fr.VI,31.
<i>S. assimilis</i> ; F. v. M., Cens. 106 (1882) from R. Brown (1810) ..	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,202 M.fr.IV,106; VI,32.
<i>S. Oldfieldii</i> ; F. v. M., Cens. 106 (1882) from Benth. (1869) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,203
<i>S. cucullata</i> ; Sprengel, syst. I, 656 (1825) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,203 M.fr.VI,32.
<i>S. sprengelioides</i> ; F. v. M., Cens. 106 (1882) from Sonder (1845)...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,204 M.fr.VI,34.
<i>S. obtusata</i> ; F. v. M., Cens. 106 (1882) from Sonder (1845) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,204
<i>S. fimbriata</i> ; F. v. M., Cens. 106 (1882) from Stschegl. (1859) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,204
<i>S. ozothamnoides</i> ; F. v. M., Cens. 106 (1882) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,204 M.fr.VI,155.
<i>S. plumiflora</i> ; F. v. M., fragm. VI, 29 (1867) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,205 M.fr.XI,122.
<i>S. unilateralis</i> ; F. v. M., Cens. 106 (1882) from Stschegl. (1859) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,205
<i>S. montana</i> ; F. v. M., fragm. VI, 45 (1867) ...	...	...	—	—	T.	V.	N.S.W.	—	—	B.fl.IV,206 M.fr.VI,45.
<i>S. Macraei</i> ; F. v. M., fragm. VI, 46 (1867) ...	...	...	—	—	—	V.	N.S.W.	—	—	B.fl.IV,206 M.fr.VI,46; VIII,54.
<i>S. pleurandroides</i> ; F. v. M., fragm. VI, 32 (1867) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,206 M.fr.III,143.
<i>S. linifolia</i> ; F. v. M., fragm. VI, 36 (1867) ...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,207 M.fr.VI,36.
<i>S. pluriloculata</i> ; F. v. M., fragm. VI, 32 (1867) ...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,207 M.fr.I,37.
<i>S. pleiosperma</i> ; F. v. M., fragm. VI, 41 (1867) ...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,207 M.fr.VI,41.
<i>S. rubicunda</i> ; F. v. M., fragm. VI, 31 (1867) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,208 M.fr.IV,99.
<i>S. attenuata</i> ; F. v. M., Cens. 106 (1882) from Cunningham (1825) ...	...	...	—	—	—	V.	N.S.W.	—	—	B.fl.IV,208
<i>S. conferta</i> ; F. v. M., Cens. 106 (1882) from Benth. (1869) ...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.IV,208
<i>S. mutica</i> ; F. v. M., fragm. VI, 45 (1867) ...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.IV,209 M.fr.VI,45.
<i>S. ericoides</i> ; Smith, Specim. of the Bot. of New Holl. 48 (1793) ...	...	...	—	—	S.A.	T.	V.	N.S.W.	Q.	B.fl.IV,209 M.fr.VI,45.
<i>S. brevicuspidis</i> ; F. v. M., Cens. 106 (1882) from Benth. (1869) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,210
<i>S. propinqua</i> ; Sprengel, syst. I, 658 (1825) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,210 M.fr.IV,103; VI,34.
<i>S. subulifolia</i> ; F. v. M., fragm. VI, 33 (1867) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,211 M.fr.IV,103.
<i>S. Allittii</i> ; F. v. M., fragm. VI, 34 (1867) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,211 M.fr.IV,103.
<i>S. racemulosa</i> ; F. v. M., fragm. VI, 33 (1867) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,211 M.fr.IV,103.
<i>S. pendula</i> ; Sprengel, syst. I, 657 (1825) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,212 M.fr.VI,33.
<i>S. concinna</i> ; F. v. M., Cens. 106 (1882) from Benth. (1869) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,212

<i>S. margarodes</i> ; Sprengel, syst. I, 657 (1825) ...	...	...	...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.IV, 213	M.fr.VI, 36.
<i>S. flavescens</i> ; F. v. M., fragm. VI, 33 (1867) ...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV, 213	M.fr.IV, 100.
<i>S. blepharolepis</i> ; F. v. M., fragm. VI, 48 (1867) ...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV, 213	M.fr.VI, 48.
<i>S. esquamata</i> ; Sprengel, syst. I, 658 (1825) ...	...	...	...	...	...	—	—	—	—	V.	N.S.W.	—	B.fl.IV, 214	
<i>S. rotundifolia</i> ; Sprengel, syst. I, 655 (1825) ...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV, 214	
<i>S. cordifolia</i> ; F. v. M., fragm. VIII, 54 (1873) ...	...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.IV, 214	M.fr.IV, 105; VI, 45.
<i>S. megacarpa</i> ; F. v. M., fragm. VI, 32 (1867) ...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV, 215	M.fr.IV, 102.
<i>S. ruscifolia</i> ; Sprengel, syst. I, 656 (1825) ...	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl.IV, 215	
<i>S. imbricata</i> ; Sprengel, syst. I, 656 (1825) ...	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl.IV, 215	
<i>S. cuspidata</i> ; Sprengel, syst. I, 657 (1825) ...	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl.IV, 216	
<i>S. leptospermoides</i> ; Sprengel, syst. I, 659 (1825) ...	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl.IV, 216	
<i>S. acuminata</i> ; Sprengel, syst. I, 659 (1825) ...	...	...	...	...	...	—	—	—	—	—	N.A.	—	B.fl.IV, 216	
<i>S. flexifolia</i> ; Sprengel, syst. I, 659 (1825) ...	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl.IV, 217	
<i>S. biflora</i> ; Sprengel, syst. I, 659 (1825) ...	...	...	...	...	...	—	—	—	V.	N.S.W.	Q.	—	B.fl.IV, 217	M.fr.IV, 104.
<i>S. setigera</i> ; Sprengel, syst. I, 659 (1825) ...	...	...	...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.IV, 217	M.fr.VI, 45.
<i>S. exolasia</i> ; F. v. M., fragm. VI, 34 (1867) ...	...	...	...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.IV, 217	M.fr.VI, 34.
<i>S. Fraseri</i> ; F. v. M., Cens. 107 (1882) from Cunningham (1840) ...	...	...	...	...	...	—	—	T.	V.	N.S.W.	—	—	B.fl.IV, 218	M.fr.IV, 105; VI, 46 ... N.Z.
<i>S. hirtella</i> ; F. v. M., Cens. 107 (1882) ...	...	...	...	...	...	—	S.A.	—	—	—	—	—	B.fl.IV, 218	
<i>S. Oldfieldii</i> ; F. v. M., Cens. 107 (1882) ...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV, 218	
<i>S. erubescens</i> ; F. v. M., fragm. VI, 33 (1867) ...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV, 219	M.fr.IV, 102.
<i>S. lissanthoides</i> ; F. v. M., fragm. VI, 33 (1867) ...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV, 219	M.fr.IV, 101.
<i>S. stricta</i> ; F. v. M., Cens. 107 (1882) from Benthams (1869) ...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV, 219	
<i>S. Mitchellii</i> ; F. v. M., Cens. 107 (1882) from Benthams (1869) ...	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl.IV, 220	
<i>S. juniperina</i> ; Sprengel, syst. I, 658 (1825) ...	...	...	...	...	...	—	—	—	V.	N.S.W.	Q.	—	B.fl.IV, 220	M.fr.IV, 104; VI, 46.
<i>S. rufa</i> ; F. v. M., fragm. VI, 46 (1867) ...	...	...	...	...	...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.IV, 221	M.fr.VIII, 54; IX, 48.
<i>S. conostephioides</i> ; F. v. M., fragm. VI, 34 (1867) ...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV, 221	M.fr.IV, 98.
<i>S. deformis</i> ; Sprengel, syst. I, 658 (1825) ...	...	...	...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.IV, 221	
<i>S. pogonocalyx</i> ; F. v. M., Cens. 107 (1882) ...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV, 222	
<i>S. breviflora</i> ; F. v. M., Cens. 107 (1882) ...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV, 222	M.fr.IV, 102.
<i>S. dura</i> ; F. v. M., Cens. 107 (1882) from Benthams (1869) ...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV, 222	
<i>S. multiflora</i> ; Sprengel, syst. I, 658 (1825) ...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV, 222	M.fr.VI, 32.
<i>S. appressa</i> ; Sprengel, syst. I, 658 (1825) ...	...	...	...	...	...	—	—	—	V.	N.S.W.	—	—	B.fl.IV, 223	
<i>S. Neo-Anglica</i> ; F. v. M., Cens. 107 (1882) ...	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV, 223	
<i>S. obtecta</i> ; F. v. M., Cens. 107 (1882) from Benthams (1869) ...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV, 223	
<i>S. crassifolia</i> ; F. v. M., fragm. VI, 40 (1867) ...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV, 224	M.fr.VI, 40.
<i>S. stronglylophylla</i> ; F. v. M., fragm. VI, 33 (1867) ...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV, 224	M.fr.IV, 101.
<i>S. crassiflora</i> ; F. v. M., fragm. VI, 33 (1867) ...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV, 224	M.fr.VI, 33.
<i>S. corynocarpa</i> ; F. v. M., Cens. 107 (1882) from Sonder (1845) ...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV, 224	
<i>S. Woodsii</i> ; F. v. M., Cens. 107 (1882) ...	...	...	...	...	...	W.A.	S.A.	—	V.	—	—	—	B.fl.IV, 225	M.fr.VI, 33.
<i>S. leptantha</i> ; F. v. M., Cens. 107 (1882) from Benthams (1869) ...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV, 225	
<i>S. ericifolia</i> ; F. v. M.; Pentachondra; J. Hook. in Lond. Journ. VI, 271 (1847) ...	...	...	...	...	...	—	—	T.	—	—	—	—	B.fl.IV, 164	
<i>S. divaricata</i> ; Sprengel, syst. I, 658 (1825) ...	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV, 226	M.fr.VI, 44.
<i>S. aggregata</i> ; Sprengel, syst. I, 657 (1825) ...	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV, 227	M.fr.VI, 44; VIII, 55.
<i>S. serrulata</i> ; Labillardière, Nov. Holl. pl. spec. I, 45, t. 62 (1804) ...	...	...	...	...	...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.IV, 227	M.fr.VI, 44.
<i>S. patula</i> ; Sprengel, syst. I, 657 (1825) ...	...	...	...	...	...	W.A.	S.A.	—	—	—	—	—	B.fl.IV, 227	M.fr.VI, 44; XI, 122.
<i>S. ovalifolia</i> ; Sprengel, syst. I, 656 (1825) ...	...	...	...	...	...	W.A.	S.A.	—	V.	—	—	—	B.fl.IV, 228	M.fr.VI, 44.
<i>S. ramiflora</i> ; Sprengel, syst. I, 659 (1825) ...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV, 228	M.fr.
<i>S. depressa</i> ; Sprengel, syst. I, 655 (1825) ...	...	...	...	...	...	W.A.	S.A.	—	—	—	—	—	B.fl.IV, 228	M.fr.VI, 44.
<i>S. fasciculiflora</i> ; F. v. M., fragm. VIII, 55 (1873) ...	...	...	...	...	...	—	S.A.	—	—	—	—	—	B.fl.IV, 229	M.fr.VI, 44.
<i>S. elliptica</i> ; Smith, Specim. of the Bot. of New Holl. 49 (1793) ...	...	...	...	...	...	—	—	T.	V.	N.S.W.	Q.	—	B.fl.IV, 230	M.fr.VI, 58.
<i>S. glauca</i> ; Labillardière, Nov. Holl. pl. spec. I, 45, t. 61 (1804) ...	...	...	...	...	...	—	—	T.	—	—	—	—	B.fl.IV, 230	

<i>S. scoparia</i> ; Smith, Specim. of the Bot. of New Holl. 49 (1793) ...	...	...	—	—	T.	V.	N.S.W.	Q.	—	B.fl.IV, 231	M.fr.VI, 58; XI, 122.
<i>S. ledifolia</i> ; F. v. M., Cens. 107 (1882) from Cunningham (1839) ...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.IV, 231	
<i>S. empetrifolia</i> ; F. v. M. in Papers of R. S. Tasm. 86 (1874) ...	...	...	—	—	T.	—	—	—	—	B.fl.IV, 231	
<i>S. oligarrhenoides</i> ; F. v. M., fragm. IX, 47 (1875) ...	...	...	...	...	W.A.	—	—	—	—	—	M.fr.IX, 47.
<i>S. minutiflora</i> ; F. v. M., fragm. XI, 122 (1881) ...	...	...	...	...	W.A.	—	—	—	—	B.fl.IV, 231	M.fr.XI, 122.

OLIGARRHENA; R. Brown, prodr. Fl. Nov. Holl. 549 (1810).

<i>O. micrantha</i> ; R. Brown, prodr. 459 (1810) ...	...	...	...	...	W.A.	—	—	—	—	B.fl.IV, 232	
---	-----	-----	-----	-----	------	---	---	---	---	--------------	--

NEEDHAMIA; R. Brown, prodr. Fl. Nov. Holl. 549 (1810).

<i>N. pumilio</i> ; R. Brown, prodr. 549 (1810) ...	...	...	...	...	W.A.	—	—	—	—	B.fl.IV, 174	
---	-----	-----	-----	-----	------	---	---	---	---	--------------	--

BRACHYLOMA; Sonder in Lehm. pl. Preiss. I, 304 (1845). (Lobopogon, Lissanthe partly.)

<i>B. daphnoides</i> ; Bentham, Fl. Austr. IV, 173 (1869) ...	...	...	...	...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV, 173	M.fr.VI, 42; VIII, 55.
<i>B. ciliatum</i> ; Bentham, Fl. Austr. IV, 173 (1869) ...	...	...	...	...	—	S.A.	T.	V.	—	—	—	B.fl.IV, 173	M.fr.VI, 42.
<i>B. depressum</i> ; Bentham, Fl. Austr. IV, 173 (1869) ...	...	...	...	...	—	—	T.	V.	—	—	—	B.fl.IV, 173	M.fr.I, 36; VI, 42.
<i>B. Scortechinii</i> ; F. v. M., fragm. XI, 121 (1881) ...	...	...	...	...	—	—	—	—	—	Q.	—	—	M.fr.XI, 121.
<i>B. Preissii</i> ; Sonder in Lehm. pl. Preiss. I, 305 (1845) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV, 172	M.fr.VI, 39; VIII, 55.
<i>B. concolor</i> ; F. v. M., fragm. VI, 39 (1867) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV, 172	M.fr.IV, 98.
<i>B. ericoides</i> ; Sonder in Schlecht. Linnaea XXVI, 247 (1853) ...	...	...	...	...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.IV, 172	M.fr.IV, 98; VI, 39; XI, 46.

CONOSTEPHIUM; Bentham in Hueg. enum. 76 (1837). (Conostephiopsis.)

<i>C. pendulum</i> ; Bentham in Hueg. enum. 76 (1837) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV, 160	M.fr.VI, 40.
<i>C. minus</i> ; Lindley, Bot. Regist. XXV, App. 25 (1839) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV, 160	M.fr.VI, 40.
<i>C. Roei</i> ; Bentham, Fl. Austr. IV, 160 (1869) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV, 160	M.fr.VIII, 55.
<i>C. Preissii</i> ; Sonder in Lehm. pl. Preiss. I, 304 (1845) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV, 161	M.fr.VI, 40.
<i>C. planifolium</i> ; F. v. M., fragm. VI, 30 (1867) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV, 161	M.fr.VI, 30.

COLEANTHERA; Stschegleew in Bull. Soc. Mosc. XXXII, 4 (1859). (Michiea.)

<i>C. coelophylla</i> ; Bentham, Fl. Austr. IV, 150 (1869) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV, 150	
<i>C. myrtoides</i> ; Stschegleew in Bull. de Mosc. XXXII, 4 (1859) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV, 150	M.fr.IV, 96; VI, 80.
<i>C. virgata</i> ; Stschegleew in Bull. de Mosc. XXXII, 5 (1859) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV, 151	

TROCHOCARPA; R. Brown, prodr. 548 (1810). (Decaspora, Pentachondra.)

<i>T. laurina</i> ; R. Brown, prodr. 548 (1810) ...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV, 166	M.fr.VI, 57; IX, 48.
<i>T. disticha</i> ; Sprengel, syst. I, 660 (1825) ...	...	...	...	...	—	—	T.	—	—	—	—	B.fl.IV, 166	M.fr.VI, 57; VIII, 55.
<i>T. thymifolia</i> ; Sprengel, syst. I, 660 (1825) ...	...	...	...	...	—	—	T.	—	—	—	—	B.fl.IV, 166	M.fr.VI, 57; VIII, 55.
<i>T. Clarkei</i> ; F. v. M., fragm. VI, 57 (1867) ...	...	...	...	...	—	—	—	V.	—	—	—	B.fl.IV, 167	M.fr.VI, 57.
<i>T. Gunnii</i> ; Bentham, Fl. Austr. IV, 167 (1869) ...	...	...	...	...	—	—	T.	—	—	—	—	B.fl.IV, 167	M.fr.VIII, 55.
<i>T. parviflora</i> ; Bentham, Fl. Austr. IV, 167 (1869) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV, 167.	
<i>T. involucrata</i> ; F. v. M., fragm. VI, 57 (1867) ...	...	...	...	...	—	—	T.	—	—	—	—	B.fl.IV, 164	M.fr.VIII, 55.
<i>T. pumila</i> ; F. v. M., fragm. VI, 57 (1867) ...	...	...	...	...	—	—	T.	V.	N.S.W.	—	—	B.fl.IV, 164	M.fr.VI, 57 ... N.Z.

EPACRIS; Cavanilles, icon. et descr. pl. IV, 25, t. 344 et 345 (1797). (Archeria.)

<i>E. micranthera</i> ; F. v. M., fragm. VI, 72 (1867) ...	...	...	...	...	—	—	T.	—	—	—	—	B.fl.IV, 246	M.fr.VIII, 56.
<i>E. hirtella</i> ; J. Hooker in Lond. Journ. of Bot. VI, 271 (1847) ...	...	...	...	...	—	—	T.	—	—	—	—	B.fl.IV, 246	M.fr.VI, 71.
<i>E. longiflora</i> ; Cavanilles, icon. et descr. pl. IV, 25, t. 344 (1797) ...	...	...	...	...	—	—	—	V.	N.S.W.	—	—	B.fl.IV, 234	M.fr.VI, 70; VIII, 56.
<i>E. reclinata</i> ; Cunningham in Benth. Fl. Austr. IV, 234 (1869) ...	...	...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.IV, 234	M.fr.VIII, 56.
<i>E. impressa</i> ; Labillardière, Nov. Holl. pl. spec. I, 43, t. 58 (1804) ...	...	...	...	...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.IV, 235	M.fr.VI, 70; VIII, 56; XI, 122.
<i>E. sparsa</i> ; R. Brown, prodr. 551 (1810) ...	...	...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.IV, 235	M.fr.VIII, 53.
<i>E. petrophila</i> ; J. Hooker, Fl. Tasman. I, 261 (1860) ...	...	...	...	...	—	—	T.	V.	N.S.W.	—	—	B.fl.IV, 236	
<i>E. rigida</i> ; Sieber in Spreng. eur. poster. 64 (1827) ...	...	...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.IV, 236	
<i>E. coriacea</i> ; Cunningham in De Candolle, prodr. VII, 763 (1839) ...	...	...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.IV, 236	
<i>E. crassifolia</i> ; R. Brown, prodr. 551 (1810) ...	...	...	...	...	—	—	—	V.	N.S.W.	—	—	B.fl.IV, 237	M.fr.IV, 127.

<i>E. robusta</i> ; Bentham, Fl. Austr. IV, 237 (1869)	—	—	—	V.	N.S.W.	—	—	B.fl.IV, 237	
<i>E. obtusifolia</i> ; Smith, Exot. Bot. I, 77, t. 40 (1804)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV, 237	M.fr.VI, 71; VIII, 56.
<i>E. myrtifolia</i> ; Labillardière, Nov. Holl. pl. spec. I, 41, t. 55 (1804)	—	—	T.	—	—	—	—	B.fl.IV, 238	
<i>E. exserta</i> ; R. Brown, prodr. 551 (1810)	—	—	T.	—	—	—	—	B.fl.IV, 238	
<i>E. mucronulata</i> ; R. Brown, prodr. 552 (1810)	—	—	T.	—	—	—	—	B.fl.IV, 238	
<i>E. lanuginosa</i> ; Labillardière, Nov. Holl. pl. spec. I, 42, t. 57 (1804)	—	S.A.	T.	V.	—	—	—	B.fl.IV, 238	M.fr.VIII, 56; XI, 122.
<i>E. paludosa</i> ; R. Brown, prodr. 551 (1810)	—	—	—	V.	N.S.W.	—	—	B.fl.IV, 239	M.fr.VIII, 56.
<i>E. Calvertiana</i> ; F. v. M., fragm. VIII, 53 (1874)	—	—	—	—	N.S.W.	—	—	—	M.fr.VIII, 53.
<i>E. heteronema</i> ; Labillardière, Nov. Holl. pl. spec. I, 42, t. 56 (1804)	—	—	—	T.	V.	N.S.W.	—	B.fl.IV, 239	M.fr.IV, 127; VI, 71; VIII, 56. N.Z.
<i>E. serpillifolia</i> ; R. Brown, prodr. 240 (1810)	—	—	—	T.	V.	N.S.W.	—	B.fl.IV, 240	
<i>E. microphylla</i> ; R. Brown, prodr. 550 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV, 240	
<i>E. acuminata</i> ; Bentham, Fl. Austr. IV, 240 (1869)	—	—	T.	—	—	—	—	B.fl.IV, 240	
<i>E. apiculata</i> ; Cunningham in Field, N. S. Wales, 340 (1825)	—	—	—	—	N.S.W.	—	—	B.fl.IV, 241	
<i>E. pulchella</i> ; Cavanilles, icon. IV, 26, t. 345 (1797)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV, 241	M.fr.VI, 71; XI, 122.
<i>E. pnrpurascens</i> ; R. Brown, prodr. 550 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.IV, 241	M.fr.IV, 127; VI, 254; VIII, 56.
WOOLLSIA; F. v. M., fragm. VIII, 52 (1872). (Lysinema partly.) [122.									
<i>W. pungens</i> ; F. v. M., fragm. VIII, 52 (1872)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV, 243	M.fr.IV, 126; VI, 70, 88; VIII, 55; IX, 48; XI,
LYSINEMA; R. Brown, prodr. 552 (1810).									
<i>L. lasianthum</i> ; R. Brown, prodr. 552 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV, 243	M.fr.III, 142; VI, 70; VIII, 55.
<i>L. conspicuum</i> ; R. Brown, prodr. 552 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV, 243	M.fr.VI, 70; VIII, 55; XI, 122.
<i>L. ciliatum</i> ; R. Brown, prodr. 552 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV, 243	M.fr.VI, 69, 70; VIII, 55.
<i>L. fimbriatum</i> ; F. v. M., fragm. IV, 125 (1864)	W.A.	—	—	—	—	—	—	B.fl.IV, 244	M.fr.VI, 70.
<i>L. elegans</i> ; Sonder in Lehm. pl. Preiss. I, 327 (1845)	W.A.	—	—	—	—	—	—	B.fl.IV, 244	M.fr.VI, 70.
PRIONOTES; R. Brown, prodr. fl. Nov. Holl. 553 (1810).									
<i>P. cerinthoides</i> ; R. Brown, prodr. 553 (1810)	—	—	T.	—	—	—	—	B.fl.IV, 246	M.fr.VI, 69; XI, 122.
COSMELIA; R. Brown, prodr. fl. Nov. Holl. 553 (1810).									
<i>C. rubra</i> ; R. Brown, prodr. 553 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV, 247	M.fr.VI, 64.
PONCELETIA; R. Brown, prodr. fl. Nov. Holl. 554 (1810).									
<i>P. sprengelioides</i> ; R. Brown, prodr. 554 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV, 248	M.fr.I, 39; VI, 60.
<i>P. monticola</i> ; Cunningham in De Candolle, prodr. VII, 768 (1843)	—	—	—	—	N.S.W.	—	—	B.fl.IV, 249	M.fr.VI, 60.
SPRENGELIA; Smith in Svensk. Vetens. Acad. Handling. 260 (1794). (Poirotia.)									
<i>S. incarnata</i> ; Smith in Vetens. Ac. Handl. 260, t. 8 (1794)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.IV, 249	M.fr.I, 39; VI, 54; IX, 48; VIII, 56; XI, 122.
ANDERSONIA; R. Brown, prodr. 553 (1810). (Atherocephala, Homalostoma, Sphincterostoma.)									
<i>A. colossea</i> ; F. v. M., fragm. VI, 63 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV, 251	M.fr.VI, 63.
<i>A. patricia</i> ; F. v. M., fragm. VI, 79 (1868)	W.A.	—	—	—	—	—	—	B.fl.IV, 251	M.fr.VI, 79.
<i>A. grandiflora</i> ; Stachegleew in Bull. de Mosc. XXXII, 21 (1859)	W.A.	—	—	—	—	—	—	B.fl.IV, 252	M.fr.VI, 62.
<i>A. setifolia</i> ; Bentham, Fl. Austr. IV, 252 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV, 252	M.fr.VIII, 51.
<i>A. involucreta</i> ; Sonder in Lehm. pl. Preiss. I, 331 (1845)	W.A.	—	—	—	—	—	—	B.fl.IV, 252	M.fr.VI, 62.
<i>A. homalostoma</i> ; Bentham, Fl. Austr. IV, 253 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV, 253	M.fr.VIII, 63.
<i>A. sprengelioides</i> ; R. Brown, prodr. 554 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV, 253	M.fr.VI, 64.
<i>A. latiflora</i> ; F. v. M., fragm. VI, 61 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV, 253	M.fr.VI, 61.
<i>A. gracilis</i> ; De Candolle, prodr. VII, 767 (1843)	W.A.	—	—	—	—	—	—	B.fl.IV, 254	
<i>A. aristata</i> ; Lindley, Bot. Regist. XXV, App. 25 (1839)	W.A.	—	—	—	—	—	—	B.fl.IV, 254	M.fr.VI, 56.
<i>A. macronema</i> ; F. v. M., fragm. VIII, 51 (1873)	W.A.	—	—	—	—	—	—	—	M.fr.VIII, 51.
<i>A. parvifolia</i> ; R. Brown, prodr. 554 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV, 254	M.fr.IV, 125; VI, 62.
<i>A. depressa</i> ; R. Brown, prodr. 554 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV, 255	
<i>A. coerules</i> ; R. Brown, prodr. 554 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV, 255	M.fr.VI, 64.

A. subulata; Benth. Fl. Austr. IV, 256 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV, 256	
A. heterophylla; Sonder in Lehm. pl. Preiss. I, 333 (1845)	W.A.	—	—	—	—	—	—	B.fl.IV, 256	M.fr.IV, 124.
A. brachyanthera; F. v. M., fragm. VI, 61 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV, 256	M.fr.VI, 61.
A. brevifolia; Sonder in Lehm. pl. Preiss. I, 332 (1845)	W.A.	—	—	—	—	—	—	B.fl.IV, 256	M.fr.VI, 61.
A. variegata; Sonder in Lehm. pl. Preiss. I, 334 (1845)	W.A.	—	—	—	—	—	—	B.fl.IV, 257	
A. micrantha; R. Brown, prodr. 554 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV, 257	M.fr.VI, 61.

RICHEA; R. Brown, prodr. 555 (1810). (Cystanthe, Pilitis.)

R. sprengelioides; F. v. M., fragm. VI, 68 (1867)	—	—	T.	—	—	—	—	B.fl.IV, 258	M.fr.I, 38; VI, 68.
R. procera; F. v. M., fragm. VI, 68 (1867)	—	—	T.	—	—	—	—	B.fl.IV, 259	M.fr.I, 38; VI, 68.
R. acerosa; F. v. M., fragm. VI, 69 (1867)	—	—	T.	—	—	—	—	B.fl.IV, 259	M.fr.I, 38; VI, 69.
R. Milligani; F. v. M., fragm. VI, 69 (1867)	—	—	T.	—	—	—	—	B.fl.IV, 259	M.fr.I, 38; VIII, 56.
R. Gunnii; J. Hooker in Lond. Journ. VI, 273 (1847)	—	—	T.	V.	—	—	—	B.fl.IV, 260	M.fr.VI, 67; XI, 122.
R. scoparia; J. Hooker in Lond. Journ. VI, 273 (1847)	—	—	T.	—	—	—	—	B.fl.IV, 260	M.fr.VI, 68.
R. dracophylla; R. Brown, prodr. 555 (1810)	—	—	T.	—	—	—	—	B.fl.IV, 260	M.fr.VI, 67.
R. paudanifolia; J. Hooker, Fl. Antarct. I, 50 (1844)	—	—	T.	—	—	—	—	B.fl.IV, 261	M.fr.VI, 67; IX, 48.

DRACOPHYLLUM; Labillardière, Voy. II, 211, t. 40 (1798). (Epacris, Forst. (1776) partly.)

D. Milligani; Hooker, icon. t. 845 (1852)	—	—	T.	—	—	—	—	B.fl.IV, 262	M.fr.VI, 65.
D. Fitzgeraldi; F. v. M., fragm. VII, 27 (1869)	—	—	—	—	—	N.S.W.	—	—	M.fr.VIII, 280; X, 119.
D. Sayeri; F. v. M. in Aust. Journ. of Pharm. new ser. II, 85 (1887)	—	—	—	—	—	—	Q.	—	—
D. secundum; R. Brown, prodr. 556 (1810)	—	—	—	—	—	N.S.W.	—	B.fl.IV, 262	M.fr.VI, 65; VIII, 56.
D. micromum; F. v. M., fragm. I, 39 (1858)	—	—	T.	—	—	—	—	B.fl.IV, 265	M.fr.VI, 65; VII, 56 ... N.Z.

SPHENOTOMA; Sweet, Flora Australasica, t. 44 (1828), from R. Brown (1810).

S. squarrosus; G. Don, gen. syst. III, 785 (1834)	W.A.	—	—	—	—	—	—	B.fl.IV, 263	M.fr.VI, 66.
S. Drummondii; F. v. M., Cens. 109 (1882) from Benth. (1869)	W.A.	—	—	—	—	—	—	B.fl.IV, 263	
S. dracophylloides; Sonder in Lehm. pl. Preiss. I, 33 (1844)	W.A.	—	—	—	—	—	—	B.fl.IV, 264	M.fr.VI, 65.
S. capitatum; Lindley, Bot. Regist. t. 1515 (1832)	W.A.	—	—	—	—	—	—	B.fl.IV, 264	
S. gracile; Sweet, Fl. Austral. t. 44 (1828)	W.A.	—	—	—	—	—	—	B.fl.IV, 264	M.fr.VI, 66.
S. parviflorum; F. v. M., Cens. of Aust. pl. 109 (1882)	W.A.	—	—	—	—	—	—	B.fl.IV, 265	M.fr.VI, 66.

APETALEAE GYMNOSPERMAE.

F. v. M. in Woolls's plants of the neighb. of Sydney, 40 (1880).

CONIFERAE.

HALLER, enum. stirp. Helv. I, 145 (1742).—Genera 10, Species 29.

DAMMARA; Rumphius, herb. Amboin. II, 174, t. 54 (1741) from Sherard (1704). (Agathis.)

D. robusta; C. Moore in Transact. pharm. Soc. Vict. II, 174 (1860)	—	—	—	—	—	—	Q.	—	B.fl.VI, 244	M.fr.XI, 104.
---	---	---	---	---	---	---	----	---	--------------	---------------

ARAUCARIA; A. L. de Jussieu, gen. plant. 413 (1789). (Colymbea, Columbea, Eutassa, Eutacta, Altingia.)

A. Cunninghamii; Aiton in Sweet. hort. Brit. 475 (1827)	—	—	—	—	—	N.S.W.	Q.	—	B.fl.VI, 243	 P.
A. Bidwilli; Hooker, Lond. Journ. of Bot. II, 503, t. 18 (1850)	—	—	—	—	—	—	Q.	—	B.fl.VI, 243	
A. excelsa; R. Brown in Ait. hort. Kew, sec. ed. V, 412 (1813)	—	—	—	—	—	N.S.W.	—	—	—	M.fr.IX, 169.

ATHROTAXIS; D. Don in Transact. Linn. Soc. XVIII, 171 (1839). (Arthrotaxis, Sequoia.)

A. cupressoides; D. Don in Trans. Linn. Soc. XVIII, 173, t. 13 (1839)	—	—	T.	—	—	—	—	B.fl.VI, 242	
A. laxifolia; Hooker, icon. plant. t. 573 (1843)	—	—	T.	—	—	—	—	B.fl.VI, 242	
A. selaginoides; D. Don in Transact. Linn. Soc. XVIII, 172, t. 14 (1839)	—	—	T.	—	—	—	—	B.fl.VI, 242	M.fr.XI, 104.

CALLITRIS; Ventenat, decas generum novorum 10 (1808). (Frenela, Actinostrobus, Leichhardtia, Octoclinis.)											
C. Macleayana; F. v. M. in Rep. Burdek. Exped. 17 (1860)	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl. VI, 235 M.fr. XI, 104.
C. Parlatores; F. v. M. in Seemann, Journ. of Bot. 267 (1866)	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl. VI, 235 M.fr. V, 186; XI, 104.
C. Roei; Benthams & J. Hooker, gen. plant. III, 424 (1880) implied	...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 236
C. Drummondii; Benthams & J. Hooker, gen. plant. III, 424 (1880)	...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 236
C. verrucosa; R. Brown in Mém. du Mus. Par. XIII, 74 (1826)	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl. VI, 236 M.fr. V, 237.
C. columellaris; F. v. M., fragm. V, 198 (1866)	...	...	...	...	—	—	—	N.S.W.	Q.	—	M.fr. V, 198.
C. Muelleri; Benthams & J. Hooker, gen. plant. III, 424 (1880)	...	...	...	...	—	—	—	N.S.W.	Q.	—	B.fl. VI, 237
C. cupressiformis; Ventenat, decas. gen. nov. 10 (1808)	...	...	...	...	—	S.A.	T.	V.	N.S.W.	Q.	— B.fl. VI, 237
C. calcarata; R. Brown in Mém. du Mus. Par. XIII, 74 (1826)	...	...	...	...	—	—	—	V.	N.S.W.	Q.	— B.fl. VI, 238
C. oblonga; L. Cl. Richard, comment. de conf. 49, t. 18 (1826)	...	...	...	...	—	—	T.	—	—	—	B.fl. VI, 238
C. Actinostrobus; F. v. M., Rep. Burdek. Exped. 19 (1860)	...	...	...	...	W.A.	—	—	—	—	—	B.fl. VI, 240
C. acuminata; F. v. M., Cens. 109 (1882) from Parlatores (1862)	...	...	...	...	W.A.	—	—	—	—	—	B.fl. VI, 240 M.fr. XI, 104.
FITZROYA; J. Hooker in Journ. of the hort. Soc. VI, 264 (1851). (Diselma.)											
F. Archeri; Benthams & J. Hooker, gen. pl. III, 426 (1880)	...	...	...	—	—	T.	—	—	—	—	B.fl. VI, 240
PHEROSPHERA; Archer in Hooker's Kew Misc. II, 52 (1850). (Dacrydium partly.)											
P. Hookeriana; Archer in Hooker, Kew Misc. II, 52 (1850)	...	...	...	—	—	T.	—	—	—	—	B.fl. VI, 245 M.fr. XI, 103.
P. Fitzgeraldi; F. v. M. in J. Hooker, icon. pl. XIV, 64, t. 1383 (1882)	...	...	...	—	—	—	—	N.S.W.	—	—	M.fr. XI, 102, 138.
MICROCACHRYIS; J. Hooker in Lond. Journ. of Bot. IV, 149 (1845). (Dacrydium partly.)											
M. tetragona; J. Hooker, Fl. Tasman. I, 358, t. 100 (1860)	...	...	...	—	—	T.	—	—	—	—	B.fl. VI, 241
DACRYDIUM; Solander in G. Forster, pl. escul. insul. ocean. austr. 80 (1786).											
D. Franklinii; J. Hooker in Lond. Journ. of Bot. IV, 152, t. 6 (1845)	...	...	...	—	—	T.	—	—	—	—	B.fl. VI, 245
NAGEIA; Gaertner, de fructib. t. 39 (1788). (Podocarpus.)											
N. elata; F. v. M., Cens. 109 (1882) from R. Brown (1826)	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl. VI, 247 M.fr. IV, 87.
N. spinulosa; F. v. M., Cens. 109 (1882) from R. Brown (1826)	...	...	...	—	—	—	—	N.S.W.	—	—	B.fl. VI, 247
N. Drouyniana; F. v. M., Cens. of Aust. pl. 109 (1882)	...	...	...	...	W.A.	—	—	—	—	—	B.fl. VI, 247 M.fr. IV, 86; VI, 251; XI, 104.
N. alpina; F. v. M. in papers Roy. Soc. of Tasm. 23 (1879)	...	...	...	...	—	—	T.	V.	N.S.W.	—	B.fl. VI, 248 M.fr. IV, 87; XI, 104.
THALAMIA; Sprengel, Anleit. zur Kenntn. der Gew. zweite Ausg. II, 218 (1817). (Phyllocladus.)											
T. asplenifolia; Sprengel, Anl. zur Kenntn. der. Gew. II, 218 (1817)	...	...	...	—	—	T.	—	—	—	—	B.fl. VI, 246 M.fr. XI, 104.

CYCADEAE.

L. C. RICHARD in Persoon, synops. 630 (1807).—Genera 3, Species 14.

CYCAS; Linné, hort. Cliffort. 482 (1737).											
C. media; R. Brown, prodr. 348 (1810)	...	...	...	...	...	—	—	—	—	Q. N.A.	B.fl.VI,249 M.fr.VIII,171.
C. Normanbyana; F. v. M., fragm. VIII, 169 (1874)	...	...	...	...	...	—	—	—	—	Q. —	M.fr.VIII,169,283.
C. Kennedyana; F. v. M. in Melb. Chemist, March (1882)	...	...	...	...	...	—	—	—	—	Q. —	—
C. Cairnsiana; F. v. M., fragm. X, 63 (1876)...	...	...	...	...	...	—	—	—	—	Q. —	M.fr.X,63,121.
ENCEPHALARTOS; Lehmann, pugill. VI, 9, t. I, et 3 (1834). (Macrozamia, Lepidozamia, Catakidozamia, Zamia partly.)											
E. Pauli Guilielmi; F. v. M. in Trans. Pharm. Soc. Vict. II, 91 (1859) ...	...	...	...	...	...	—	—	—	—	N.S.W. Q. —	B.fl.VI,251 M.fr.I,86,243; II,179; IX,124.
E. tridentatus; Lehmann, pugill. VI, 13 (1834)	...	...	...	...	...	—	—	—	—	N.S.W. —	B.fl.VI,253 M.fr.III,38; XI,125.
E. spiralis; Lehmann, pugill. VI, 13 (1834)	...	...	...	...	...	—	—	—	—	N.S.W. Q. —	B.fl.VI,251 M.fr.I,41; II,179; VIII,173.
E. Douglasii; F. v. M., Cens. 110 (1882)	...	...	...	...	...	—	—	—	—	Q. —	—
E. Fraseri; Miquel in Versl. K. Akad. Wet. Amst. XV, 368 (1863)	...	...	...	...	...	W.A.	—	—	—	—	B.fl.VI,252 M.fr.I,41,243; II,179; III,167; VIII,173; XI,
E. Moorei; F. v. M., fragm. XI, 125 (1881)	...	...	...	...	...	—	—	—	—	Q. —	M.fr.I,243; XI,125.
E. Macdonnelli; F. v. M. in Versl. K. Akad. Wet. Amst. XV, 376 (1863)	...	...	...	...	...	—	S.A.	—	—	—	M.fr.IX,124; XI,126.
E. Dyeri; F. v. M. in Melb. Chemist, June (1885)	...	...	...	...	...	W.A.	—	—	—	—	—
E. Denisonii; F. v. M. in Journ. Pharm. Soc. Vict. II, 90 (1859)...	...	...	...	...	...	—	—	—	—	N.S.W. Q. —	B.fl.VI,253 M.fr.I,41; VIII,173; IX,124.
BOWENIA; Hooker, Bot. Magaz. t. 5398 (1863).											
B. spectabilis; Hooker, Bot. Mag. t. 5398 (1863)	...	...	...	...	...	—	—	—	—	Q. —	B.fl.VI,254 M.fr.V,171; VIII,279.

MONOCOTYLEDONEAE.

RAY, meth. plant. nov. (1682).

EUCALYCEAE PERIGYNAE.

F. v. M. in Woolls's plants of the neighb. of Sydney 41 (1880).

SCITAMINEAE.

LINNE, philos. botan. 27 (1751).—Genera 7, Species 11.

MUSA; l'Ecluse, exoticorum libri decem 229 (1605).

M. Banksii; F. v. M., fragm. IV, 132 (1864)...	...	...	...	...	...	...	...	...	Q.	—	B.fl. VI, 261	M.fr. IX, 190.
M. Fitzalani; F. v. M., fragm. IX, 188 (1875)	...	...	...	...	...	...	...	...	Q.	—	—	M.fr. IX, 188.
M. Hillii; F. v. M., fragm. IX, 169 (1875)	...	...	...	...	...	...	...	...	Q.	—	—	M.fr. IX, 169.

CURCUMA; Linné, gen. plant. 332 (1737) from Hermann (1687).

C. Australasica; J. Hooker, Bot. Mag. t. 5620 (1867)	...	...	...	...	...	...	...	...	Q.	—	B.fl. VI, 263	M.fr. VIII, 26.
--	-----	-----	-----	-----	-----	-----	-----	-----	----	---	---------------	-----------------

AMOMUM; Linné, gen. plant. 330 (1737).

A. Dallachyi; F. v. M., fragm. VIII, 25 (1872)	...	...	...	...	...	...	...	...	Q.	—	B.fl. VI, 263	M.fr. VIII, 25.
--	-----	-----	-----	-----	-----	-----	-----	-----	----	---	---------------	-----------------

ELETTARIA; White & Matton in Transact. Linn. Soc. X, 249 (1809) from Rheede (1692).

E. Scottiana; F. v. M., fragm. VIII, 24 (1872)	...	...	...	...	...	...	...	...	Q.	—	B.fl. VI, 264	M.fr. VIII, 24.
--	-----	-----	-----	-----	-----	-----	-----	-----	----	---	---------------	-----------------

ALPINIA; Linné, gen. plant. 332 (1737). (Hellenia.)

A. racemigera; F. v. M., fragm. VIII, 27 (1872)	...	...	...	...	...	...	...	...	Q.	—	B.fl. VI, 265	M.fr. VIII, 27.
A. coerulea; Benthams, Fl. Austr. VI, 265 (1873)	...	...	...	...	...	...	...	N.S.W.	Q.	—	B.fl. VI, 265	M.fr. VIII, 26.
A. arctiflora; F. v. M., fragm. VIII, 25 (1872)	...	...	...	...	...	...	...	...	Q.	—	B.fl. VI, 266	M.fr. VIII, 25.

COSTUS; C. Bauhin, pinax 36 (1623).

C. Potierae; F. v. M., fragm. IV, 164 (1864)...	...	...	...	...	...	...	...	...	Q.	—	B.fl. VI, 266	M.fr. VIII, 27.
---	-----	-----	-----	-----	-----	-----	-----	-----	----	---	---------------	-----------------

TAPEINOCHEILOUS; Miquel in annal. mus. bot. Lugdun. IV, 101, t. 4 (1869).

T. pungens; Miquel in ann. mus. Lugd. IV, 101, t. 4 (1869)	...	...	...	...	...	...	...	...	Q.	—	B.fl. VI, 267	M.fr. VIII, 26	...	As.
--	-----	-----	-----	-----	-----	-----	-----	-----	----	---	---------------	----------------	-----	-----

ORCHIDEAE.

HALLER, enum. stirp. Helv. praefat. 33 (1742).—Genera 49, Species 271.

PHOLIDOTA; Lindley in Hooker, Exotic Flora II, t. 138 (1825).

P. imbricata; Lindley in Hook. Exot. Fl. II, t. 138 (1825)	...	...	...	...	...	...	...	...	Q.	—	B.fl. VI, 290	M.fr. VI, 163; V, 105	...	As.
--	-----	-----	-----	-----	-----	-----	-----	-----	----	---	---------------	-----------------------	-----	-----

STURMIA; Reichenbach in Moessler's Handb. II, 1552 (1828). (Liparis, Richard 1818 not of Zoologists 1738.)

S. reflexa; F. v. M., fragm. II, 72 (1861)	...	...	...	...	...	...	...	N.S.W.	Q.	—	B.fl. VI, 273	M.fr. III, 165; V, 102.
S. angustilabris; F. v. M., fragm. IV, 164 (1864)	...	...	...	...	...	...	...	—	Q.	—	B.fl. VI, 273	M.fr. IV, 164.
S. coelogynoides; F. v. M., fragm. II, 71 (1861)	...	...	...	...	...	...	...	N.S.W.	Q.	—	B.fl. VI, 273	M.fr. IX, 50.
S. habenarina; F. v. M., fragm. IV, 131 (1864)	...	...	...	...	...	...	...	—	Q.	—	B.fl. VI, 273	M.fr. V, 102; XI, 53.

OBERONIA; Lindley, gen. and spec. of orchid. pl. 15 (1830). (Titania.)

O. iridifolia; Lindley, gen. and spec. of orchid. pl. 15 (1830)	...	...	...	...	...	...	...	N.S.W.	Q.	—	B.fl. VI, 274	M.fr. II, 24.
O. Titania; Lindley, fol. orchidac. VIII, 8 (1859)	...	...	...	...	...	...	...	N.S.W.	Q.	—	B.fl. VI, 274	M.fr. II, 24; VII, 30; XI, 53, 88.

PHREATIA; Lindley, gen. and spec. of orchid. pl. 63 (1830). (Plexaure.)

P. Limenophylax; G. Reichenbach in Seem. Bonpl. 54 (1857) ... Q. — B.fl. VI, 290 M.fr. X, 64.

DENDROBIUM; Swartz in nov. act. Upsal. VI, 82 (1799). (Thelychiton partly, Pedilonum, Coelandria.)

D. bigibbum; Lindley in Paxt. Flower Gard. III, 25 (1853) ... Q. N.A. B.fl. VI, 277 M.fr. VI, 119; XI, 87.
 D. dicuphum; F. v. M., fragm. VIII, 28 (1871) ... — N.A. B.fl. VI, 277 M.fr. VI, 119.
 D. Sumneri; F. v. M., fragm. VI, 94 (1868) ... Q. — B.fl. VI, 278 M.fr. VI, 94.
 D. superbiens; G. Reichenbach in Garden. Chron. 516 (1876) ... Q. — M.fr. XI, 22.
 D. undulatum; R. Brown, prodr. 332 (1810) ... Q. — B.fl. VI, 278 M.fr. I, 87; III, 39; XI, 87. As.
 D. Johannis; G. Reichenbach in Garden. Chron. 890 (1865) ... Q. — B.fl. VI, 279
 D. speciosum; Smith, Exot. Botany I, 17, t. 10 (1804) ... V. N.S.W. Q. — B.fl. VI, 279 M.fr. I, 180; VIII, 249.
 D. falconirostre; Fitzgerald, Austr. Orch. I, pt. V (1879) ... N.S.W. — M.fr. XI, 87.
 D. tetragonum; Cunningham in Bot. Regiat. XXV, Misc. 33 (1839) ... N.S.W. Q. — B.fl. VI, 279 M.fr. I, 87; VI, 119.
 D. aemulum; R. Brown, prodr. 333 (1810) ... N.S.W. Q. — B.fl. VI, 280 M.fr. I, 213; XI, 87.
 D. Adae; Bailey in Proceed. Roy. Soc. of Q. I, pt. 3 (1884) ... Q. —
 D. Eriae; D. eriaeoides, Bailey, papers on the Queenal. Flora. II, Sept. (1886) ... Q. —
 D. Fitzgeraldi; F. v. M. in Melb. Chemist, June (1884) ... Q. —
 D. macropus; Benthams & J. Hooker, gen. pl. III, 500 (1883) implied ... N.S.W. —
 D. Bairdianum; Bailey, suppl. to syn. of Q. Fl. Nov. (1884) ... Q. —
 D. Stuartii; Bailey in Proc. Roy. Soc. of Q. I, pt. 2 (1884) ... Q. —
 D. Kingianum; Bidwill in Bot. Regist. XXV, Misc. II (1839) ... N.S.W. Q. — B.fl. VI, 280 M.fr. III, 60.
 D. brachypus; G. Reichenbach in Garcke, Linnaea XLI, 42 (1876) ... N.S.W. —
 D. gracilicaule; F. v. M., fragm. I, 179 (1859) ... N.S.W. Q. — B.fl. VI, 281 M.fr. III, 164; IV, 171.
 D. Moorei; F. v. M., fragm. VII, 29 (1870) ... N.S.W. — B.fl. VI, 281 M.fr. XI, 54.
 D. antennatum; Lindley in Hook. Lond. Jour. of Bot. II, 236 (1843) ... Q. —
 D. agrostophyllum; F. v. M., fragm. VIII, 28 (1871) ... Q. — B.fl. VI, 282 M.fr. VIII, 28.
 D. Baileyi; F. v. M., fragm. VIII, 173 (1874) ... Q. — M.fr. VIII, 173.
 D. Smilliae; F. v. M., fragm. VI, 94 (1868) ... Q. — B.fl. VI, 282 M.fr. VII, 63; XI, 87.
 D. ophioglossum; G. Reichenbach in Journ. L. Soc. XV, 113 (1876) ... Q. — M.fr. XI, 54.
 D. canaliculatum; R. Brown, prodr. 333 (1810) ... Q. — B.fl. VI, 282 M.fr. III, 126.
 D. Foelschei; F. v. M. in Wing's S. Sc. Record, Oct. (1882) ... — N.A. —
 D. monophyllum; F. v. M., fragm. I, 189 (1859) ... N.S.W. Q. — B.fl. VI, 283 M.fr. III, 164; VII, 64.
 D. hispidum; A. Richard, sert. Astrol. 13, t. 5 (1833) ... Q. — B.fl. VI, 283 M.fr. VII, 30; VIII, 150.
 D. cucumerinum; MacLeay in Bot. Reg., new ser. V, Misc. 58 (1842) ... N.S.W. Q. — B.fl. VI, 283 M.fr. III, 59; VIII, 151.
 D. pugioniforme; Cunningham in Bot. Reg. XXV, Misc. 33 (1839) ... N.S.W. Q. — B.fl. VI, 284 M.fr. I, 189; III, 59.
 D. rigidum; R. Brown, prodr. 333 (1810) ... Q. — B.fl. VI, 284 M.fr. VIII, 248; XI, 87.
 D. linguiforme; Swartz in Svensk. Vetensk. Ac. Handl. 247 (1800) ... N.S.W. Q. — B.fl. VI, 284 M.fr. I, 214.
 D. teretifolium; R. Brown, prodr. 333 (1810) ... N.S.W. Q. — B.fl. VI, 285 M.fr. I, 89; III, 60.
 D. Fairfaxii; Fitzgerald & F. v. M. in Sydney Mail 360, Nov. (1872) ... N.S.W. Q. — B.fl. VI, 285 M.fr. X, 122.
 D. striolatum; G. Reichenbach in Hamburg Gartenz. 313 (1857) ... T. V. N.S.W. — B.fl. VI, 285 M.fr. I, 88; III, 39; VII, 30.
 D. Mortii; F. v. M., fragm. I, 214 (1859) ... N.S.W. Q. — B.fl. VI, 286 M.fr. II, 93; III, 60; IX, 50.
 D. Beckleri; F. v. M., fragm. V, 95 (1865) ... N.S.W. Q. — B.fl. VI, 296 M.fr. VII, 59.

BULBOPHYLLUM; Petit-Thouars, orchid. Afric. t. 95 et 108 (1822). (Bolbophyllum, Thelychiton partly, Cirrhopetalum.)

B. Lichenastrum; F. v. M., fragm. VII, 60 (1869) ... Q. — B.fl. VI, 287 M.fr. X, 119.
 B. Baileyi; F. v. M., fragm. IX, 5 (1875) ... Q. — M.fr. IX, 5.
 B. nematopodium; F. v. M., fragm. VIII, 30 (1873) ... Q. — B.fl. VI, 287 M.fr. VIII, 130.
 B. Shepherdii; F. v. M., fragm. III, 40 (1862) ... N.S.W. Q. — B.fl. VI, 288 M.fr. I, 190.
 B. aurantiacum; F. v. M., fragm. III, 39 (1862) ... N.S.W. Q. — B.fl. VI, 288 M.fr. VII, 98.
 B. argyropus; G. Reichenbach in Garcke, Linnaea XLI, 42 (1876) ... N.S.W. —
 B. exiguum; F. v. M., fragm. II, 72 (1861) ... N.S.W. Q. — B.fl. VI, 288 M.fr. II, 175; XI, 22.

B. minutissimum; F. v. M., fragm. XI, 53 (1878)	—	—	—	—	N.S.W.	Q.	—	—	M.fr.III,60; V,95; XI,88.
B. Prenticei; F. v. M. in Wing's South. Sc. Record I, 173 (1881)	—	—	—	—	—	Q.	—	—	B.fl.X,139.
B. Elisae; F. v. M., fragm. VI, 120 (1868)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,289	M.fr.VI,254; VIII,28.
B. punctatum; Fitzgerald in Britten's Journ. Bot. XXI, 205 (1883)	—	—	—	—	—	Q.	—	—	
B. clavigerum; Cirrhopetalum; Fitzgerald in Brit. Journ. Bot. XXI, 204 (1883)	—	—	—	—	—	Q.	—	—	
B. purpurascens; Bailey in Proc. Roy. Soc. Queensl. I, 88 (1884)	—	—	—	—	—	Q.	—	—	
B. Bowkettiae; Bailey in Proc. Roy. Soc. Queensl. I, 88 (1884)	—	—	—	—	—	Q.	—	—	

ERIA; Lindley in Edw. Bot. Regist. t. 904 (1825).

E. Fitzalani; F. v. M. in Wing's South. Sc. Rec. II, 251 (1882)	—	—	—	—	—	Q.	—	—	
--	---	---	---	---	---	----	---	---	--

LUISIA; Gaudichaud in Freyc. voy. bot. 426, t. 37 (1826).

L. teretifolia; Gaudichaud in Freyc. voy. bot. 426, t. 37 (1826)	—	—	—	—	—	Q.	—	—	 P.
---	---	---	---	---	---	----	---	---	-----------

SARCOCHILUS; R. Brown, prodr. 332 (1810). (Thrixspermum, Gunnia, Cleisostoma, Saccolabium partly.)

S. erectus; F. v. M., Cens. 111 (1882) from Fitzgerald (1877)	—	—	—	—	N.S.W.	Q.	—	—	M.fr.XI,52,139.
S. tridentatus; G. Reichenbach in Walp. Annal. VI, 500 (1861)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,296	M.fr.I,192; II,181; VII,98.
S. Armitii; F. v. M., fragm. IX, 49 (1875)	—	—	—	—	—	Q.	—	—	M.fr.XI,49.
S. Beckleri; F. v. M., Cens. of Austr. Pl. 111 (1882)	—	—	—	—	N.S.W.	—	—	B.fl.VI,296	M.fr.VIII,248.
S. brevibras; F. v. M., Cens. of Austr. Pl. 111 (1882)	—	—	—	—	—	Q.	—	—	M.fr.VIII,248; XI,139.
S. Macphersoni; F. v. M., fragm. VIII, 248 (1874)	—	—	—	—	—	Q.	—	B.fl.VI,297	M.fr.VII,96.
S. Hartmanni; F. v. M., fragm. VIII, 248 (1874)	—	—	—	—	—	Q.	—	—	M.fr.VIII,248.
S. divitiflorus; F. v. M. in Benth. Fl. Austr. VI, 292 (1870)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,292	M.fr.IX,50; XI,22.
S. falcatus; R. Brown, prodr. 332 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,293	M.fr.I,89,243; VII,87; X,50.
S. Fitzgeraldi; F. v. M., fragm. VII, 115 (1870)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,293	M.fr.VII,115.
S. olivaceus; Lindley, Bot. Regist. Misc. XXV, 32 (1839)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,293	M.fr.I,90,191; VII,97.
S. parviflorus; Lindley, Bot. Regist. Misc. XXIV, 34 (1838)	—	—	—	—	T. V.	N.S.W.	Q.	B.fl.VI,294	M.fr.I,89,243; VII,98.
S. Ceciliae; F. v. M., fragm. V, 42, t. 42 (1865)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,294	M.fr.VII,98; XI,22.
S. Hillii; F. v. M., fragm. II, 94 (1861)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,295	M.fr.I,88,243; III,39; VII,98.
S. phyllorrhizus; F. v. M., fragm. V, 201 (1866)	—	—	—	—	—	Q.	—	B.fl.VI,295	M.fr.VII,30.
S. Keffordii; F. v. M., Cens. sec. suppl. (1885) from Bailey (1884)	—	—	—	—	—	Q.	—	—	
S. platystachys; Bailey, syn. Q. Fl., first suppl. 56 (1886)	—	—	—	—	—	Q.	—	—	

TAENIOPHYLLUM; Blume, Bijdr. 355 (1825).

T. Muelleri; Lindley in Benth. Fl. Austr. VI, 291 (1870)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,291	
---	---	---	---	---	--------	----	---	-------------	--

ORNITHOCHILUS; Wallich in Lindley, gen and spec. of orchid. plants 242 (1833). (Saccolabium partly.)

O. Hillii; Bentham in Journ. Linn. Soc. XVIII, 334 (1880)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,298	M.fr.VII,97.
--	---	---	---	---	--------	----	---	-------------	--------------

GEODORUM; Jackson in Andrews, Bot. Reposit. t. 626 (1810).

G. pictum; Lindley, gen. and spec. of orchid. pl. 175 (1833)	—	—	—	—	—	Q.	N.A.	B.fl.VI,299	M.fr.III,24; V,102; VIII,31. As., P.
---	---	---	---	---	---	----	------	-------------	--------------------------------------

EULOPHIA; R. Brown in Edwards, Bot. Regist. 686 (1822). (Eulophus 1822.)

E. venosa; G. Reichenbach in Benth. Fl. Austr. VI, 300 (1870)	—	—	—	—	—	Q.	N.A.	B.fl.VI,300	M.fr.I,61.
E. Fitzalani; F. v. M., fragm. VIII, 30 (1872)	—	—	—	—	—	Q.	—	B.fl.VI,300	M.fr.VII,30.
E. Holtzei; F. v. M. in Vict. Nat. VI, 55 (1889)	—	—	—	—	—	—	N.A.	—	

DIPODIUM; R. Brown, prodr. 336 (1810). (Waillesia.)

D. punctatum; R. Brown, prodr. 331 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.VI,301	M.fr.X,64.
D. ensifolium; F. v. M., fragm. V, 42 (1865)	—	—	—	—	—	Q.	—	B.fl.VI,301	M.fr.V,42.

CYMBIDIUM; Swartz in nov. act. Upsal. VI, 70 (1799).

C. Hillii; F. v. M. in Regel's Gartenflora, 138 (1879)	—	—	—	—	—	Q.	—	—	M.fr.XI,88.
C. canaliculatum; R. Brown, prodr. 331 (1810)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.VI,202	M.fr.XI,88.

C. albuciflorum; F. v. M., fragm. I, 188 (1859)	...	...	...	...	...	...	...	...	...	N.S.W.	Q.	—	B.fl. VI,303	M.fr.I,188			
C. suave; R. Brown, prodr. 331 (1810)	...	...	...	...	...	...	...	...	...	N.S.W.	Q.	—	B.fl. VI,303	M.fr.I,187.			
SPATHOGLOTTIS; Blume, Bijdr. 400 (1825).																	
S. Paulinæ; F. v. M., fragm. VI, 95 (1868)	...	...	...	...	...	...	...	...	...	—	Q.	—	B.fl. VI,304	M.fr.X,64.			
PHAJUS; Loureiro, Fl. Cochinch. II, 529 (1790).																	
P. grandifolius; Loureiro, Fl. Cochinch. II, 529 (1790)	...	...	...	...	...	...	...	...	...	N.S.W.	Q.	—	B.fl. VI,304	M.fr.I,42; IV,163; X,6	As., P.		
CALANTHE; R. Brown in Edwards, Bot. Regist. 573 (1821).																	
C. veratrifolia; R. Brown in Bot. Regist. 573 (1821)	...	...	...	...	...	...	...	...	...	N.S.W.	Q.	—	B.fl. VI,305	M.fr. VIII,30; IX,50	As., P.		
DIDYMOPLEXIS; Griffith in Calcutta Journ. of Nat. Hist. IV, 383 (1844). (Leucorchis.)																	
D. pallens; Griffith in Calc. Journ. IV, 383, t. 17 (1844)	...	...	...	...	...	...	...	...	...	—	N.A.	—	...	...	As.		
GALEOLA; Loureiro, Fl. Cochinch. II, 520 (1790). (Erythrorchis, Ledgeria.)																	
G. cassythoides; G. Reichenbach, Xen. Orchid. II, 77 (1869)	...	...	...	...	...	...	...	...	...	N.S.W.	Q.	—	B.fl. VI,307	M.fr.II,167; V,102; VIII,31; XI,105.			
G. Ledgeriana; F. v. M., fragm. XI, 127 (1872)	...	...	...	...	...	...	...	...	...	N.S.W.	Q.	—	B.fl. VI,307	M.fr.II,167; III,40; V,102; VIII,31.			
EPIPOGUM; Gmelin, Fl. Sibir. I, 11, t. 2 (1747).																	
E. nutans; Lindley in Journ. Linn. Soc. I, 177 (1856)	...	...	...	...	...	...	...	...	...	N.S.W.	Q.	—	B.fl. VI,308	M.fr. VIII,30; IX,50; XI,105.	As., Afr.		
GASTRODIA; R. Brown, prodr. 330 (1810).																	
G. sesamoides; R. Brown, prodr. 330 (1810)	...	...	...	...	...	...	...	W.A.	—	T.	V.	N.S.W.	Q.	—	B.fl. VI,309	M.fr. VIII,151; X,64.	
POGONIA; A. L. de Jussieu, gen. plant. 65 (1789).																	
P. uniflora; F. v. M., fragm. V, 201 (1866)	...	...	...	...	...	...	...	...	...	—	Q.	—	B.fl. VI,310	M.fr. V,201.			
P. holochila; F. v. M., fragm. V, 200 (1866)...	...	...	...	...	...	...	...	...	...	—	Q.	—	B.fl. VI,310	M.fr. V,200.			
P. flabelliformis; Lindley in Wallich's numer. list. 7400 (1828)	...	...	...	...	...	...	...	...	...	—	N.A.	—	...	...	As.		
P. Dallachyana; F. v. M. in Benth. Fl. Austr. VI, 310 (1870)	...	...	...	...	...	...	...	...	...	—	Q.	—	B.fl. VI,310				
P. pachystomoides; F. v. M., fragm. VIII, 174 (1874)	...	...	...	...	...	...	...	...	...	—	Q.	—		M.fr. VIII,174.			
CORYMBORCHIS; Petit-Thouars, hist. des plant. Orchid. recueil. sur les isles d'Afrique. t. 37-38 (1822). (Corymbis.)																	
C. veratrifolia; Blume, Collect. des Orchidées 125, t. 42 & 43 (1858)	...	...	...	...	...	...	...	...	...	—	Q.	—	B.fl. VI,311	M.fr. VIII,30; X,64	As., P.		
ETAERIA; Blume, Bijdr. 409 (1825). (Hetaeria, Rhamphidia.)																	
E. tenuis; Bentham in Journ. Linn. Soc. XVIII, 345 (1880)	...	...	...	...	...	...	...	...	...	—	Q.	—	B.fl. VI,312	...	As., P.		
MICRÖSTYLIS; Nuttall, Genera of North American plants II, 196 (1818).																	
M. congesta; G. Reichenbach in Walp. Annal. bot. syst. VI, 206 (1861)	...	...	...	...	...	...	...	...	...	—	Q.	—		M.fr. XI,21.	As.		
GOODYERA; R. Brown in Aiton. hort. Kew sec. ed. V, 197 (1813).																	
G. viridiflora; Blume, Coll. des Orchid. 41, t. 9 (1858)	...	...	...	...	...	...	...	...	...	—	Q.	—	B.fl. VI,313	M.fr. VIII,29.	As.		
G. polygonoides; F. v. M., fragm. VIII, 29 (1872)	...	...	...	...	...	...	...	...	...	—	Q.	—	B.fl. VI,313	M.fr. VII,30; VIII,29; X,64.			
SPIRANTHES; L. C. Richard in Mém. du Mus. IV, 40 (1818). (Gyrostachis, 1807.)																	
S. Australis; Lindley, Bot. Regist. 823 (1824)	...	...	...	...	...	...	...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl. VI,314	M.fr. X,65	E., As., N.Z.
ANOECTOCHILUS; Blume, flor. Jav. prae fat. 6 (1828).																	
A. species unascertained	...	...	...	...	...	...	...	...	...	—	Q.	—					
THELYMITRA; R. & G. Forster, char. gen. 97, t. 49 (1776). (Macdonaldia.)																	
T. ixioides; Swartz in Kong. Svensk. Acad. Handl. 228, t. 3 (1800)	...	...	...	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl. VI,317		
T. circumsepta; Fitzgerald, Austral. Orchids I, part 4 (1877)	...	...	...	...	...	...	...	...	...	—	—	N.S.W.	—	—			
T. erinita; Lindley, Bot. Regist. XXV, App. 49 (1839)	...	...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI,319	M.fr. VI,84.	
T. aristata; Lindley, gen. and spec. of orchid. pl. 521 (1840)	...	...	...	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl. VI,319		
T. epipactoides; F. v. M., fragm. V, 174 (1866)	...	...	...	...	...	...	...	...	...	—	V.	—	—	—		M.fr. V,174.	

T. longifolia; R. & G. Forster, char. gen. 98, t. 49 (1776) ...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,319	...	...	...	P., N.Z.
T. villosa; Lindley, Bot. Regist. XXV, App. 49 (1839) ...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.VI,320	M.fr.V,94,214.			
T. tigrina; R. Brown, prodr. 315 (1810) ...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.VI,320				
T. fusco-lutea; R. Brown, prodr. 315 (1810) ...	...	...	...	W.A.	S.A.	—	V.	—	—	—	B.fl.VI,321	M.fr.V,94.			
T. stellata; Lindley, Bot. Regist. XXV, App. 49 (1839) ...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.VI,321				
T. carnea; R. Brown, prodr. 314 (1810) ...	...	...	...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,321	M.fr.V,98.			
T. flexuosa; Endlicher, nov. stirp. decad. 23 (1839)...	...	...	...	W.A.	S.A.	T.	V.	—	—	—	B.fl.VI,322				
T. antennifera; J. Hooker, Fl. Tasman. II, 4, t. 101 (1860) ...	...	...	...	W.A.	S.A.	T.	V.	—	—	—	B.fl.VI,322				
T. Macmillani; F. v. M., fragm. V, 93 (1865) ...	...	...	...	—	—	—	V.	—	—	—	B.fl.VI,323	M.fr.V,93.			
T. Mackibbinii; F. v. M. in Melb. Chemist 44 (1881) ...	...	...	...	W.A.	S.A.	—	V.	—	—	—	—	M.fr.XI,139.			
T. variegata; Lindley in Benth. Fl. Austr. VI, 323 (1870)...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.VI,323	M.fr.V,97,98.			
T. venosa; R. Brown, prodr. 314 (1870) ...	...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.VI,323	...	...	...	N.Z.
T. cyanea; Lindley in Benth. Fl. Austr. VI, 323 (1870) ...	...	...	...	—	—	T.	—	—	—	—	B.fl.VI,323				
T. mucida; Fitzgerald in Gard. Chron. new ser. XVII, 495 (1882) ...	...	...	...	W.A.	—	—	—	—	—	—	—				
T. luteo-ciliata; Fitzgerald in Gard. Chron. new ser. XVII, 433 (1882) ...	...	...	...	—	S.A.	—	—	—	—	—	—				

EPIBLEMA; R. Brown, prodr. fl. Nov. Holl. 315 (1810).

E. grandiflorum; R. Brown, prodr. 315 (1810) ...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.VI,324	M.fr.XI,21.			
--	-----	-----	-----	------	---	---	---	---	---	---	-------------	-------------	--	--	--

DIURIS; Smith in Transact. Linn. Soc. IV, 222 (1798).

D. alba; R. Brown, prodr. 316 (1810) ...	...	...	...	—	—	—	V.	N.S.W.	Q.	—	B.fl.VI,325	M.fr.V,102.			
D. punctata; Smith, Exot. Bot. I, 13, t. 8 (1804) ...	...	...	...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.VI,326	M.fr.V,102.			
D. secundiflora; Fitzgerald, Austral. Orchids I, part 4 (1877) ...	...	...	...	—	—	—	—	N.S.W.	Q.	—	—				
D. aurea; Smith, Exot. Bot. 15, t. 9 (1804) ...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,327	M.fr.V,172.			
D. palustris; Lindley, gen. and spec. of orchid. pl. 507 (1840) ...	...	...	...	—	S.A.	T.	V.	—	—	—	B.fl.VI,327	M.fr.V,102.			
D. maculata; Smith, Exot. Bot. I, 57, t. 30 (1804) ...	...	...	...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,327	M.fr.V,102.			
D. pedunculata; R. Brown, prodr. 316 (1810) ...	...	...	...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,328	M.fr.V,102,173.			
D. pallens; Benth. Fl. Austr. VI, 329 (1870)...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,329				
D. abbreviata; F. v. M. in Benth. Fl. Austr. VI, 329 (1870) ...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,329				
D. setacea; R. Brown, prodr. 316 (1810) ...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.VI,329	M.fr.V,101.			
D. emarginata; R. Brown, prodr. 316 (1810)...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.VI,330				
D. sulphurea; R. Brown, prodr. 316 (1810) ...	...	...	...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,330	M.fr.V,102,173.			
D. aequalis; F. v. M. in Fitzgerald, Austral. Orchids I, part 2 (1873) ...	...	...	...	—	—	—	—	N.S.W.	—	—	—	M.fr.X,65.			
D. dendrobioides; Fitzgerald, Austral. Orchids I, part 7 (1882) ...	...	...	...	—	—	—	—	N.S.W.	—	—	—				
D. longifolia; R. Brown, prodr. 316 (1810) ...	...	...	...	W.A.	S.A.	T.	V.	—	—	—	B.fl.VI,331	M.fr.V,101,172.			
D. pauciflora; R. Brown, prodr. 316 (1810) ...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.VI,331				

ORTHOCERAS; R. Brown, prodr. fl. Nov. Holl. 317 (1810).

O. strictum; R. Brown, prodr. 317 (1810) ...	...	...	...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.VI,332	...	...	...	N.Z.
--	-----	-----	-----	---	------	---	----	--------	---	---	-------------	-----	-----	-----	------

CALOCHILUS; R. Brown, prodr. fl. Nov. Holl. 320 (1810).

C. campester; R. Brown, prodr. 320 (1810) ...	...	...	...	—	—	T.	V.	N.S.W.	Q.	—	B.fl.VI,315	M.fr.VIII,151.			
C. Robertsoni; Benth. Fl. Austr. VI, 315 (1870)...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.VI,315				
C. paludosus; R. Brown, prodr. 320 (1810) ...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,316				

CRYPTOSTYLIS; R. Brown, prodr. 317 (1810). (Zosterostylis.)

C. longifolia; R. Brown, prodr. 317 (1810) ...	...	...	...	—	—	T.	V.	N.S.W.	Q.	—	B.fl.VI,333	M.fr.VII,116; XI,21.			
C. ovata; R. Brown, prodr. 317 (1810) ...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.VI,334	M.fr.VII,116.			
C. erecta; R. Brown, prodr. 317 (1810) ...	...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.VI,334	M.fr.VII,115.			
C. leptochila; F. v. M. in Benth. Fl. Austr. VI, 324 (1870) ...	...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.VI,334				

PRASOPHYLLUM; R. Brown, prodr. 317 (1810). (Genoplesium, Carunastylis).

P. Australe; R. Brown, prodr. 318 (1810) ...	...	...	...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,337				
P. flavum; R. Brown, prodr. 318 (1810) ...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,337				

<i>P. elatum</i> ; R. Brown, prodr. 318 (1810)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,337	M.fr.V,100.	
<i>P. brevilabre</i> ; J. Hooker, Fl. Tasman. II, 11, t. 110 (1860)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.VI,338	M.fr.V,101.	
<i>P. hians</i> ; G. Reichenbach, Beitr. zur syst. Pflanzenk. 61 (1871)	W.A.	—	—	—	—	—	—	B.fl.VI,338		
<i>P. patens</i> ; R. Brown, prodr. 318 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,339	M.fr.V,101.	
<i>P. Deaneanum</i> ; Fitzgerald in proc. L. S. of N.S.W. sec. ser. II, 142 (1887)	—	—	—	—	N.S.W.	—	—	—		
<i>P. fuscum</i> ; R. Brown, prodr. 318 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,339	M.fr.V,101.	
<i>P. cyphochilum</i> ; Benth. Fl. Austr. VI, 340 (1870)	W.A.	—	—	—	—	—	—	B.fl.VI,340		
<i>P. ovale</i> ; Lindley, Bot. Regist. XXV, App. 54 (1839)	W.A.	—	—	—	—	—	—	B.fl.VI,341		
<i>P. macrostachyum</i> ; R. Brown, prodr. 318 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,341		
<i>P. Reichenbachii</i> ; F. v. M., Fimbria; G. Reichenbach, Beitr. 60 (1871)	W.A.	—	—	V.	—	—	—	B.fl.VI,341		
<i>P. striatum</i> ; R. Brown, prodr. 318 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.VI,342	M.fr.V,100.	
<i>P. parvifolium</i> ; Lindley, Bot. Regist. XXV, App. 54 (1839)	W.A.	—	—	—	—	—	—	B.fl.VI,342	M.fr.V,101.	
<i>P. gibbosum</i> ; R. Brown, prodr. 381 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,342	M.fr.V,100.	
<i>P. cucullatum</i> ; G. Reichenbach, Beitr. zur syst. Pflanzenk. 59 (1871)	W.A.	—	—	—	—	—	—	B.fl.VI,343		
<i>P. nigricans</i> ; R. Brown, prodr. 319 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,343	M.fr.IX,50.	
<i>P. rufum</i> ; R. Brown, prodr. 319 (1810)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.VI,344	M.fr.V,100	... N.Z.
<i>P. brachystachyum</i> ; Lindley, gen. and spec. of orchid. pl. 513 (1840)	—	—	T.	—	—	—	—	B.fl.VI,344	M.fr.V,100.	
<i>P. despectans</i> ; J. Hooker, Fl. Tasman. II, 13, t. 113 (1860)	—	S.A.	T.	V.	—	—	—	B.fl.VI,345	M.fr.V,100.	
<i>P. fimbriatum</i> ; R. Brown, prodr. 319 (1810)	—	—	—	V.	N.S.W.	—	—	B.fl.VI,345	M.fr.V,100.	
<i>P. Archeri</i> ; J. Hooker, Fl. Tasman. II, 14, t. 113 (1860)	—	—	T.	V.	—	—	—	B.fl.VI,345	M.fr.V,100.	
<i>P. iotricatum</i> ; C. Stuart in Benth. Fl. Austr. VI, 346 (1870)	—	—	T.	V.	—	—	—	B.fl.VI,346		
<i>P. Woollsi</i> ; F. v. M., fragm. V, 100 (1866)	—	—	—	—	N.S.W.	—	—	B.fl.VI,346	M.fr.V,100.	
<i>P. apostasioides</i> ; Carunastylis, Fitzgerald, Austr. Orchids II, part 3 (1888)	—	—	—	—	N.S.W.	—	—	—		

MICROTIS; R. Brown, prodr. fl. Nov. Holl. 320 (1810).

<i>M. porrifolia</i> ; R. Brown, prodr. 320 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,347	M.fr.X,65; XI,21	... N.Z., P.
<i>M. media</i> ; R. Brown, prodr. 321 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,348		
<i>M. alba</i> ; R. Brown, prodr. 321 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,348		
<i>M. pulchella</i> ; R. Brown, prodr. 321 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,349		
<i>M. atrata</i> ; Lindley, Bot. Regist. XXV, App. 54 (1839)	W.A.	S.A.	—	V.	—	—	—	B.fl.VI,349	M.fr.I,90,244; X,65.	

CORYSANTHES; R. Brown, prodr. 328 (1810). (Corybas, Nematoceras.)

<i>C. unguiculata</i> ; R. Brown, prodr. 328 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.VI,350		
<i>C. pruinosa</i> ; R. Cunningham in N.S. Wales Magaz. n. 1 (1833)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	—	M.fr.IX,49.	
<i>C. fimbriata</i> ; R. Brown, prodr. 328 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.VI,351	M.fr.X,65.	
<i>C. bicalcarata</i> ; R. Brown, prodr. 328 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,351		

PTEROSTYLIS; R. Brown, prodr. fl. Nov. Holl. 326 (1810).

<i>P. ophioglossa</i> ; R. Brown, prodr. 326 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,354		
<i>P. concinna</i> ; R. Brown, prodr. 326 (1810)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VI,355	M.fr.V,99; XI,105.	
<i>P. curta</i> ; R. Brown, prodr. 326 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,355	M.fr.V,99.	
<i>P. acuminata</i> ; R. Brown, prodr. 326 (1810)	—	—	—	V.	N.S.W.	Q.	—	B.fl.VI,355		
<i>P. nutans</i> ; R. Brown, prodr. 327 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,356	M.fr.V,99.	
<i>P. pedaloglossa</i> ; Fitzgerald, Austral. Orchids II, part 3 (1887)	—	—	—	V.	N.S.W.	—	—	—		
<i>P. pedunculata</i> ; R. Brown, prodr. 327 (1810)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,356	M.fr.VIII,249; XI,105.	
<i>P. nana</i> ; R. Brown, prodr. 327 (1810)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,357	M.fr.V,99.	
<i>P. cucullata</i> ; R. Brown, prodr. 327 (1810)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,357	M.fr.V,99.	
<i>P. grandiflora</i> ; R. Brown, prodr. 327 (1810)	—	—	—	V.	N.S.W.	Q.	—	B.fl.VI,358	M.fr.V,99; XI,105.	
<i>P. truncata</i> ; Fitzgerald, Austral. Orchids I, part 4 (1878)	—	—	—	—	N.S.W.	—	—	—		
<i>P. reflexa</i> ; R. Brown, prodr. 327 (1810)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.VI,359	M.fr.XI,105.	
<i>P. praecox</i> ; Lindley, gen. and spec. of Orchid. pl. 388 (1840)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,359		

<i>P. obtusa</i> ; R. Brown, prodr. 327 (1810)	—	S.A. T.	V.	N.S.W.	—	—	B.fl. VI, 360	M.fr. IX, 50; XI, 53.
<i>P. recurva</i> ; Benth. Fl. Austr. VI, 360 (1870)	W.A.	—	—	—	—	—	B.fl. VI, 360	
<i>P. parviflora</i> ; R. Brown, prodr. 327 (1810)	—	—	T.	V.	N.S.W.	Q.	B.fl. VI, 361	M.fr. V, 100.
<i>P. barbata</i> ; Lindley, Bot. Regist. XXV, App. 53 (1839)	W.A.	S.A. T.	V.	N.S.W.	—	—	B.fl. VI, 362	M.fr. VIII, 151; XI, 22, 105. N.Z.
<i>P. mutica</i> ; R. Brown, prodr. 328 (1810)	—	S.A. T.	V.	N.S.W.	Q.	—	B.fl. VI, 362	M.fr. V, 100.
<i>P. rufa</i> ; R. Brown, prodr. 327 (1810)	W.A.	S.A. T.	V.	N.S.W.	Q.	—	B.fl. VI, 363	M.fr. V, 100.
<i>P. Daintreyana</i> ; F. v. M. in Benth. Fl. Austr. VI, 360 (1870)	—	—	—	N.S.W.	—	—	B.fl. VI, 360	
<i>P. longifolia</i> ; R. Brown, prodr. 327 (1810)	—	S.A. T.	V.	N.S.W.	Q.	—	B.fl. VI, 364	M.fr. V, 100.
<i>P. vittata</i> ; Lindley, Bot. Regist. XXV, App. 53 (1839)	W.A.	S.A. T.	V.	—	—	—	B.fl. VI, 364	M.fr. XI, 126.

CALEYA; R. Brown in Aiton, Hort. Kew sec. ed. V, 204 (1813). (Caleana 1810).

<i>C. major</i> ; R. Brown, prodr. 329 (1810)	—	—	T.	V.	N.S.W.	Q.	—	B.fl. VI, 366	M.fr. XI, 105.
<i>C. minor</i> ; R. Brown, prodr. 329 (1810)	—	—	T.	—	N.S.W.	—	—	B.fl. VI, 366	
<i>C. nigrita</i> ; Lindley, Bot. Regist. XXV, App. 54 (1839)	W.A.	—	—	—	—	—	—	B.fl. VI, 366	
<i>C. Sullivani</i> ; F. v. M. in Melb. Chemist 44 (1882)	—	—	—	V.	—	—	—	—	

DRAKAEA; Lindley, Bot. Regist. XXV, App. 55 (1834). (Spiculaea, Arthrochilus.)

<i>D. ciliata</i> ; G. Reichenbach, Beitr. zur. syst. Pflanzenk. 68 (1871)	W.A.	—	—	—	—	—	—	B.fl. VI, 367	
<i>D. irritabilis</i> ; G. Reichenbach, Beitr. zur. syst. Pflanzenk. 68 (1871)	—	—	—	V.	N.S.W.	Q.	—	B.fl. VI, 368	M.fr. I, 43; VIII, 151; XI, 53.
<i>D. elastica</i> ; Lindley, Bot. Regist. XXV, App. 55 (1839)	W.A.	—	—	—	—	—	—	B.fl. VI, 368	

ACIANTHUS; R. Brown, prodr. fl. Nov. Holl. 321 (1810).

<i>A. caudatus</i> ; R. Brown, prodr. 321 (1810)	—	S.A. T.	V.	N.S.W.	—	—	B.fl. VI, 369	
<i>A. fornicatus</i> ; R. Brown, prodr. 321 (1810)	—	—	—	N.S.W.	Q.	—	B.fl. VI, 370	
<i>A. exsertus</i> ; R. Brown, prodr. 321 (1810)	W.A.	S.A. T.	V.	N.S.W.	Q.	—	B.fl. VI, 370	M.fr. XI, 53, 105.
<i>A. viridis</i> ; J. Hooker, Fl. Tasman. II, 372 (1860)	—	—	T.	—	—	—	B.fl. IV, 371	M.fr. IX, 50.

CYRTOSTYLIS; R. Brown, prodr. fl. Nov. Holl. 322 (1810).

<i>C. reniformis</i> ; R. Brown, prodr. 322 (1810)	W.A.	S.A. T.	V.	N.S.W.	Q.	—	B.fl. VI, 376	
---	------	---------	----	--------	----	---	---------------	--

LYPERANTHUS; R. Brown, prodr. 325 (1810). (Burnettia, Fitzgeraldia.)

<i>L. nigricans</i> ; R. Brown, prodr. 325 (1810)	W.A.	S.A. T.	V.	N.S.W.	—	—	B.fl. VI, 374	M.fr. VII, 134; X, 65.
<i>L. Burnettii</i> ; F. v. M., fragm. V, 96 (1865)	—	—	T.	V.	—	—	B.fl. VI, 375	M.fr. V, 96; VII, 134.
<i>L. Forrestii</i> ; F. v. M. in Wing's S. Sc. Rec. II, 55 (1882)	W.A.	—	—	—	—	—	—	
<i>L. ellipticus</i> ; R. Brown, prodr. 325 (1810)	—	—	—	N.S.W.	—	—	B.fl. VI, 374	M.fr. VII, 133.

ERIOCHILUS; R. Brown, prodr. fl. Nov. Holl. 323 (1810).

<i>E. autumnalis</i> ; R. Brown, prodr. 323 (1810)	—	S.A. T.	V.	N.S.W.	Q.	—	B.fl. VI, 372	M.fr. XI, 22, 105.
<i>E. scaber</i> ; Lindley, Bot. Regist. XXV, App. 53 (1839)	W.A.	—	—	—	—	—	B.fl. VI, 372	
<i>E. tenuis</i> ; Lindley, Bot. Regist. XXV, App. 53 (1839)	W.A.	—	—	—	—	—	B.fl. VI, 372	
<i>E. dilatatus</i> ; Lindley, Bot. Regist. XXV, App. 53 (1839)	W.A.	—	—	—	—	—	B.fl. VI, 372	
<i>E. multiflorus</i> ; Lindley, Bot. Regist. XXV, App. 53 (1839)	W.A.	—	—	—	—	—	B.fl. VI, 373	M.fr. XI, 105.
<i>E. fimbriatus</i> ; F. v. M. in Wing's S. Sc. Record II, 152 (1882)	W.A.	S.A.	—	V.	—	—	B.fl. VI, 377	M.fr. XI, 104.

CALADENIA; R. Brown, prodr. 325 (1810). (Leptoceras, Adenochilus.)

<i>C. Menziesii</i> ; R. Brown, prodr. 325 (1810)	W.A.	S.A. T.	V.	—	—	—	B.fl. VI, 377	M.fr. V, 101; XI, 105.
<i>C. Nortoni</i> ; F. v. M., Cens. 113 (1882) from Fitzgerald (1876)	—	—	—	N.S.W.	—	—	—	M.fr. X, 64.
<i>C. Cairnsiana</i> ; F. v. M., fragm. VII, 31 (1869)	W.A.	S.A.	—	—	—	—	B.fl. VI, 378	M.fr. VII, 31.
<i>C. multiclavia</i> ; G. Reichenbach, Beitr. zur. syst. Pflanzenk. 64 (1871)	W.A.	—	—	—	—	—	B.fl. VI, 378	
<i>C. discoidea</i> ; Lindley, Bot. Regist. XXV, App. 52 (1839)	W.A.	—	—	—	—	—	B.fl. VI, 382	
<i>C. Patersoni</i> ; R. Brown, prodr. 324 (1810)	W.A.	S.A. T.	V.	N.S.W.	Q.	—	B.fl. VI, 382	M.fr. V, 101.
<i>C. Drummondii</i> ; Benth. Fl. Austr. VI, 383 (1873)	W.A.	—	—	—	—	—	B.fl. VI, 383	
<i>C. hirta</i> ; Lindley, Bot. Regist. XXV, App. 52 (1839)	W.A.	—	—	—	—	—	B.fl. VI, 383	

C. Roei; Bentham, Fl. Austr. VI, 383 (1873)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 383
C. Barbarossae; G. Reichenbach, Beitr. zur syst. Pflanzenk. 64 (1871)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 383
C. flava; R. Brown, prodr. 324 (1810)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 384 M.fr. V, 101; VI, 83; XI, 105.
C. latifolia; R. Brown, prodr. 324 (1810)	...	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl. VI, 384 M.fr. V, 101; XI, 105.
C. reptans; Lindley, Bot. Regist. XXV, App. 52 (1839)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 385
C. suaveolens; G. Reichenbach, Beitr. zur syst. Pflanzenk. 67 (1871)	...	...	...	...	...	—	—	T.	V.	N.S.W.	—	—	B.fl. VI, 385 M.fr. V, 93.
C. serrata; G. Reichenbach, Beitr. zur syst. Pflanzenk. 67 (1871)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 386 M.fr. V, 99.
C. carnea; R. Brown, prodr. 324 (1810)	...	...	...	...	...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl. VI, 386 M.fr. V, 101.
C. congesta; R. Brown, prodr. 324 (1810)	...	...	...	...	...	—	—	T.	V.	N.S.W.	—	—	B.fl. VI, 387
C. aphylla; Bentham, Fl. Austr. VI, 387 (1873)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 387
C. coerulea; R. Brown, prodr. 324 (1810)	...	...	...	...	...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl. VI, 388 M.fr. V, 101; XI, 22.
C. deformis; R. Brown, prodr. 324 (1810)	...	...	...	...	...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl. VI, 388 M.fr. V, 101; XI, 105.
C. sericea; Lindley, Bot. Regist. XXV, App. 52 (1839)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 389
C. gemmata; Lindley, Bot. Regist. XXV, App. 52 (1839)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 389 M.fr. V, 101.
C. ixioides; Lindley, Bot. Regist. XXV, App. 52 (1839)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 389
C. unita; Fitzgerald in Gardeners' Chronicle 461 (1882)	...	...	...	...	...	W.A.	—	—	—	—	—	—	—

CHILOGLOTTIS; R. Brown, prodr. 322 (1810).

C. diphylla; R. Brown, prodr. 323 (1810)	...	...	...	...	...	—	—	T.	V.	N.S.W.	Q.	—	B.fl. VI, 390 M.fr. X, 116.
C. Gunnii; Lindley, gen and spec. orchid. 387 (1840)	...	...	...	...	...	—	—	T.	V.	N.S.W.	—	—	B.fl. VI, 391 M.fr. VIII, 151; IX, 50; X, 116; XI, 22.
C. trilabra; Fitzgerald in Britten's Journ. of Bot. XXI, 304 (1883)	...	...	...	...	...	—	—	—	—	—	Q.	—	—

GLOSSODIA; R. Brown, prodr. 325 (1810).

G. major; R. Brown, prodr. 326 (1810)	...	...	...	...	...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl. VI, 392
G. minor; R. Brown, prodr. 326 (1810)	...	...	...	...	...	—	—	—	V.	N.S.W.	Q.	—	B.fl. VI, 392
G. Brunonis; Endlicher, nov. stirp. dec. 16 (1839)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 393 M.fr. VI, 83.
G. intermedia; Fitzgerald in Gardeners' Chronicle 462 (1882)	...	...	...	...	...	W.A.	—	—	—	—	—	—	—
G. emarginata; Lindley, gen. and spec. of orchid. pl. 424 (1840)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 393 M.fr. XI, 22.

HABENARIA; Willdenow, spec. plant. IV, 44 (1805).

H. trinervis; Wight, icon. plant. Ind. or. t. 1701 (1852)	...	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl. VI, 394	...	...	...	As.
H. elongata; R. Brown, prodr. 313 (1810)	...	...	...	...	...	—	—	—	—	—	—	Q.	N.A.	B.fl. VI, 394	M.fr. VII, 15.		
H. graminea; Lindley, gen. and spec. of orchid. pl. 318 (1840)	...	...	...	...	...	—	—	—	—	—	—	Q.	N.A.	B.fl. IV, 394	M.fr. VII, 16.		
H. ochroleuca; R. Brown, prodr. 313 (1810)	...	...	...	...	...	—	—	—	—	—	—	—	N.A.	B.fl. IV, 394	M.fr. V, 102.		
H. xanthantha; F. v. M., fragm. VII, 16 (1869)	...	...	...	...	...	—	—	—	—	—	—	Q.	—	B.fl. VI, 395	M.fr. VII, 16.		

APOSTASIACEAE.

ACH. RICHARD, in Ch. d'Orbigny Dictionn. univ. d'hist. nat. II, 34 (1842), from R. Brown (1830).

APOSTASIA; Blume, Bijdr. 423 (1825). (Niemeyera.)

A. stylidioides; G. Reichenbach in Flora Regens. C. B. Z. 278 (1872)	...	...	—	—	—	—	—	Q.	—	B.fl. VI, 396	M.fr. VI, 96; VII, 114.				
--	-----	-----	---	---	---	---	---	----	---	---------------	-------------------------	--	--	--	--

BURMANNIACEAE.

SPRENGEL, syst. veget. I, 125 (1825).—Genus 1, Species 2.

BURMANNIA; Linné, syst. nat. 8 (1735); gen. plant. 100 (1737).

B. disticha; Linné, spec. plant. 287 (1753)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl. VI, 397	...	...	...	As.
B. juncea; Solander in R. Brown, prodr. 265 (1810)	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl. VI, 397	...	...	...	As.

IRIDEAE.

VENTENAT, Tableau du règne végétal II, 188 (1799).—Genera 4, Species 24.

DIPLARRHENA ; Labillardière, Relation du voy. à la rech. de La Per. I, 275, t. 15 (1799)

D. Moraea ; Labillardière, Voy. I, 157, t. 15 (1799) ... — — T. V. N.S.W. — — B.fl.VI,400 M.fr.VII,94.

PATERSONIA ; R. Brown in Bot. Magaz. t. 1041 (1807). (Genosiris, 1804.)

P. glauca ; R. Brown, prodr. 304 (1810) ... — S.A. T. V. N.S.W. — — B.fl.VI,402 M.fr.VII,36.
 P. longiscapa ; Sweet, Fl. Austr. t. 39 (1828) ... — S.A. T. V. — — B.fl.VI,402 M.fr.VII,36.
 P. occidentalis ; R. Brown, prodr. 304 (1810) ... W.A. — — — — B.fl.VI,403 M.fr.VII,31.
 P. umbrosa ; Endlicher in Lehm. pl. Preiss. II, 31 (1846) ... W.A. — — — — B.fl.VI,404 M.fr.VII,32.
 P. xanthina ; F. v. M., fragm. I, 214 (1859) ... W.A. — — — — B.fl.VI,404 M.fr.I,214.
 P. limbata ; Endlicher in Lehm. pl. Preiss. II, 29 (1846) ... W.A. — — — — B.fl.VI,404 M.fr.VII,32.
 P. juncea ; Lindley, Bot. Regist. XXV, App. 58 (1839) ... W.A. — — — — B.fl.VI,404 M.fr.VII,33.
 P. Maxwelli ; F. v. M. in Benth. Fl. Austr. VI, 405 (1873) ... W.A. — — — — B.fl.VI,405 M.fr.VII,34.
 P. pygmaea ; Lindley, Bot. Regist. XXV, App. 58 (1839) ... W.A. — — — — B.fl.VI,405 M.fr.VII,33.
 P. sericea ; R. Brown in Bot. Mag. t. 1041 (1807) ... — — — V. N.S.W. Q. — B.fl.VI,406 M.fr.VII,35.
 P. lanata ; R. Brown, prodr. 303 (1810) ... W.A. — — — — B.fl.VI,406 M.fr.VII,35.
 P. rudis ; Endlicher in Lehm. pl. Preiss. II, 29 (1846) ... W.A. — — — — B.fl.VI,406 M.fr.VII,35.
 P. macrantha ; Benth. Fl. Austr. VI, 407 (1873) ... W.A. — — — — B.fl.VI,407
 P. glabrata ; R. Brown, prodr. 304 (1810) ... — — — V. N.S.W. Q. — B.fl.VI,407 M.fr.VII,35.
 P. Drummondii ; F. v. M. in Benth. Fl. Austr. VI, 407 (1873) ... W.A. — — — — B.fl.VI,407
 P. inaequalis ; Benth. Fl. Austr. VI, 408 (1873) ... W.A. — — — — B.fl.VI,408
 P. graminea ; Benth. Fl. Austr. VI, 408 (1873) ... W.A. — — — — B.fl.VI,408 M.fr.XI,23.
 P. habianoides ; Benth. Fl. Austr. VI, 408 (1873) ... W.A. — — — — B.fl.VI,408

IRIS ; Tournefort, inst. 358, t. 186, 188 (1700), from Hippocrates, Dioscorides and Plinius. (Moraea partly.)

I. Robinsoniana ; F. v. M., fragm. VII, 153 (1870) ... — — — — N.S.W. — — B.fl.VI,409 M.fr.VII,153; VIII,281.

SISYRINCHIUM ; Linné, gen. plant. 273 (1737). (Libertia, Orthrosanthus, Renealmia, Nematostigma.)

S. paniculatum ; R. Brown, prodr. 305 (1810) ... — — — V. N.S.W. Q. — B.fl.VI,413 M.fr.VII,91.
 S. pulchellum ; R. Brown, prodr. 305 (1810) ... — — — T. V. N.S.W. — — B.fl.VI,413 M.fr.VII,92 ... P., N.Z.
 S. cyaneum ; Lindley, Bot. Regist. t. 1090 (1827) ... W.A. S.A. — V. — — B.fl.VI,410 M.fr.VII,92.
 S. polystachyum ; F. v. M., Orthrosanthus, Benth. Fl. Austr. VI, 411 (1873) W.A. — — — — B.fl.VI,411

HYDROCHARIDEAE.

LAMARCK & DE CANDOLLE, Flore Française, III, 265 (1805).—Genera 7, Species 9.

HYDROCHARIS ; Linné, syst. nat. 8 (1735); gen. plant. 308 (1737).

H. morsus ranae ; Linné, spec. plant. 1036 (1753) ... — — — — N.S.W. Q. — B.fl.VI,256 M.fr.VI,199 ... E., As.

ENHALUS ; L. C. Richard in Mém. de l'Inst. II, 64 (1811).

E. Koenigii ; Richard in Mém. de l'Inst. II, 64 (1811) ... — — — — Q. — — M.fr.VIII,219 ... As.

HALOPHILA ; Du Petit-Thouars, gen. nov. Madag. n. 6 (1806).

H. ovata ; Gaudichaud, Bot. Voy. Freyc. 470, t. 40 (1826) ... — S.A. T. V. N.S.W. Q. — B.fl.VII,182 M.fr.VIII,219 ... As.
 H. spinulosa ; Ascherson in Neumayer, Anleit. zu wiss. Beoh. 368 (1875) ... — — — — Q. — B.fl.VII,183 M.fr.VIII,219,283.

OTTELIA ; Persoon, synops. plant. I, 400 (1805). (Damasonium partly, Hymenotheca.)

O. alismoides ; Persoon, synops. plant. I, 400 (1805) ... — — — — Q. N.A. B.fl.VI,257 M.fr.I,94; VI,199. As.
 O. ovalifolia ; L. C. Richard in Mém. de l'Inst. II, 78 (1811) ... W.A. S.A. — V. N.S.W. Q. N.A. B.fl.VI,257

BLYXA; Noronha in Du Petit-Thouars, gen. nov. Madag. n. 14 (1806).

B. Roxburghii; L. C. Richard, Mém. de l'Inst. II, 77, t. 5 (1811) ... — S.A. — — — Q. N.A. B.fl. VI, 258 As.

VALLISNERIA; Micheli, nov. plant. gen. 12, t. 10 (1729).

V. spiralis; Linné, spec. plant. 1015 (1753) ... — S.A. T. V. N.S.W. Q. N.A. B.fl. VI, 259 E., As., Afr., Am.

HYDRILLA; L. C. Richard in Mém. de l'Inst. II, 61 (1811). (Elodeae subgenus.)

E. verticillata; F. v. M., Key to the System of Vict. plants I, 423 (1888) ... — S.A. — V. N.S.W. Q. N.A. B.fl. VI, 259 M.fr. I, 94 As., Afr.

TACCACEAE.

PRESL, reliq. Haenk. I, 149 (1828).—Genus 1, Species 1.

TACCA, R. et G. Forster, char. gen. 65 (1776).

T. pinnatifida; R. et G. Forster, char. gen. 65 (1776) ... — — — — — Q. N.A. B.fl. VI, 453 As., P.

HAEMODORACEAE.

R. BROWN, prodr. Fl. Nov. Holl. 299 (1810).—Genera 5, Species 66.

HAEMODORUM, Smith in Transact. Linn. Soc. 213 (1798).

H. distichophyllum; Hooker, icon. plant. t. 866 (1852) ... — — — — — T. — — — — — B.fl. VI, 419 M.fr. VII, 118; IX, 51.
H. brevicaule; F. v. M., fragm. I, 64 (1858) ... — — — — — — — — — — — N.A. B.fl. VI, 419 M.fr. VII, 118.
H. sparsiflorum; F. v. M., fragm. VII, 117 (1870) ... — — — — — W.A. — — — — — — — — — — — B.fl. VI, 420 M.fr. VII, 117.
H. spicatum; R. Brown, prodr. 300 (1810) ... — — — — — W.A. — — — — — — — — — — — B.fl. VI, 420 M.fr. VII, 118.
H. brevisepaleum; Benth. Fl. Austr. VI, 420 (1873) ... — — — — — W.A. — — — — — — — — — — — B.fl. VI, 420
H. paniculatum; Lindley, Bot. Regist. XXV, App. 44 (1839) ... — — — — — W.A. — — — — — — — — — — — B.fl. VI, 420
H. laxum; R. Brown, prodr. 300 (1810) ... — — — — — W.A. — — — — — — — — — — — B.fl. VI, 421 M.fr. VII, 118.
H. simplex; Lindley, Bot. Regist. XXV, App. 44 (1839) ... — — — — — W.A. — — — — — — — — — — — B.fl. VI, 421
H. simulans; F. v. M., fragm. VII, 117 (1870) ... — — — — — W.A. — — — — — — — — — — — B.fl. VI, 421 M.fr. VII, 117.
H. planifolium; R. Brown, prodr. 300 (1810) ... — — — — — — — — — — — N.S.W. Q. — — — — — B.fl. VI, 422 M.fr. VII, 118.
H. teretifolium; R. Brown, prodr. 300 (1810) ... — — — — — — — — — — — N.S.W. — — — — — B.fl. VI, 422 M.fr. VII, 117.
H. coccineum; R. Brown, prodr. 300 (1810) ... — — — — — — — — — — — Q. N.A. B.fl. VI, 422 M.fr. VII, 117.
H. subvirens; F. v. M., fragm. I, 63 (1858) ... — — — — — — — — — — — — — — — — — N.A. B.fl. VI, 423 M.fr. I, 63.
H. ensifolium; F. v. M., fragm. I, 64 (1858) ... — — — — — — — — — — — Q. N.A. B.fl. VI, 423 M.fr. VII, 118.
H. parviflorum; Benth. Fl. Austr. VI, 423 (1873) ... — — — — — — — — — — — — — — — — — N.A. B.fl. VI, 423
H. leptostachyum; Benth. Fl. Austr. VI, 423 (1873) ... — — — — — — — — — — — — — — — — — N.A. B.fl. VI, 423
H. tenuifolium; Cunningham in Benth. Fl. Austr. VI, 425 (1873) ... — — — — — — — — — — — Q. — — — — — B.fl. VI, 423

PHLEBOCARYA; R. Brown, prodr. 301 (1810).

P. ciliata; R. Brown, prodr. 301 (1810) ... — — — — — W.A. — — — — — — — — — — — B.fl. VI, 424 M.fr. VIII, 23.
P. pilosissima; F. v. M. in Benth. Fl. Austr. VI, 425 (1873) ... — — — — — W.A. — — — — — — — — — — — B.fl. VI, 425 M.fr. VIII, 23.
P. filifolia; F. v. M. in Benth. Fl. Austr. VI, 425 (1873) ... — — — — — W.A. — — — — — — — — — — — B.fl. VI, 425 M.fr. VIII, 23.

TRIBONANTHES; Endlicher, nov. stirp. dec. I, 27 (1839).

T. brachypetala; Lindley, Bot. Regist. XXV, App. 44 (1839) ... — — — — — W.A. — — — — — — — — — — — B.fl. VI, 426 M.fr. VIII, 23.
T. uniflora; Lindley, Bot. Regist. XXV, App. 44 (1839) ... — — — — — W.A. — — — — — — — — — — — B.fl. VI, 427 M.fr. VIII, 23.
T. Australis; Endlicher, nov. stirp. dec. I, 27 (1839) ... — — — — — W.A. — — — — — — — — — — — B.fl. VI, 427 M.fr. VIII, 23.
T. variabilis; Lindley, Bot. Regist. XXV, App. 44 (1839) ... — — — — — W.A. — — — — — — — — — — — B.fl. VI, 427
T. longipetala; Lindley, Bot. Regist. XXV, App. 44 (1839) ... — — — — — W.A. — — — — — — — — — — — B.fl. VI, 427 M.fr. VIII, 23.

CONOSTYLIS; R. Brown, prodr. 301 (1810). (Blancoa, Androstemma.)

C. breviscapa; R. Brown, prodr. 301 (1810) ... — — — — — W.A. — — — — — — — — — — — B.fl. VI, 430
C. vaginata; Endlicher in Lehm. pl. Preiss. II, 23 (1846) ... — — — — — W.A. — — — — — — — — — — — B.fl. VI, 431 M.fr. VIII, 19.

<i>C. petrophiloides</i> ; F. v. M. in Benth. Fl. Austr. VI, 431 (1873) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 431	
<i>C. setosa</i> ; Lindley, Bot. Regist. XXV, App. 44, t. 6 (1839) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 431	M.fr. VIII, 19.
<i>C. aurea</i> ; Lindley, Bot. Regist. XXV, App. 44 (1839) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 432	M.fr. VIII, 19.
<i>C. melanopogon</i> ; Endlicher in Lehm. pl. Preiss. II, 18 (1846) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 432	M.fr. VIII, 20.
<i>C. setigera</i> ; R. Brown, prodr. 300 (1810) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 432	M.fr. VIII, 20.
<i>C. Psyllium</i> ; Endlicher in Lehm. pl. Preiss. II, 21 (1846) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 433	
<i>C. villosa</i> ; Benth. Fl. Austr. VI, 433 (1873) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 433	
<i>C. Drummondii</i> ; Benth. Fl. Austr. VI, 433 (1873) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 433	
<i>C. involucrata</i> ; Endlicher in Lehm. pl. Preiss. II, 23 (1846) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 434	
<i>C. gladiata</i> ; Benth. Fl. Austr. VI, 434 (1873) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 434	
<i>C. seorsiflora</i> ; F. v. M., fragm. I, 158 (1859) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 435	M.fr. VIII, 19.
<i>C. teretiuscula</i> ; F. v. M., fragm. VIII, 18 (1872) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 435	M.fr. VIII, 18.
<i>C. stylidioides</i> ; F. v. M., fragm. VIII, 17 (1872) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 435	M.fr. VIII, 17.
<i>C. prolifera</i> ; Benth. Fl. Austr. VI, 436 (1873) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 436	
<i>C. racemosa</i> ; Benth. Fl. Austr. VI, 436 (1873) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 436	
<i>C. candicans</i> ; Endlicher, nov. stirp. dec. 20 (1839) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 436	M.fr. VI, 255.
<i>C. dealbata</i> ; Lindley, Bot. Regist. XXV, App. 45 (1839) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 436	
<i>C. Preissii</i> ; Endlicher in Lehm. pl. Preiss. II, 18 (1846) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 437	M.fr. VIII, 20.
<i>C. bracteata</i> ; Endlicher in Lehm. pl. Preiss. II, 16 (1846) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 437	M.fr. VIII, 20.
<i>C. filifolia</i> ; F. v. M., fragm. VIII, 18 (1873) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 438	M.fr. VIII, 18.
<i>C. spinuligera</i> ; F. v. M. in Benth. Fl. Austr. VI, 438 (1873) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 438	
<i>C. bromelioides</i> ; Endlicher in Lehm. pl. Preiss. II, 18 (1846) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 438	
<i>C. aculeata</i> ; R. Brown, prodr. 300 (1810) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 438	M.fr. VIII, 19.
<i>C. laxiflora</i> ; Benth. Fl. Austr. VI, 439 (1873) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 439	
<i>C. cymosa</i> ; F. v. M. in Benth. Fl. Austr. VI, 439 (1873) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 439	
<i>C. serrulata</i> ; R. Brown, prodr. 300 (1810) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 439	M.fr. VIII, 19.
<i>C. caricina</i> ; Lindley, Bot. Regist. XXV, App. 45 (1839) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 440	M.fr. VIII, 20.
<i>C. Androstemma</i> ; F. v. M., fragm. VIII, 19 (1873) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 440	M.fr. VIII, 19; IX, 51.
<i>C. Bealiana</i> ; F. v. M., fragm. IX, 50 (1875) ...	...	...	W.A.	—	—	—	—	—	—	—	M.fr. IX, 50.
<i>C. canescens</i> ; F. v. M., fragm. VIII, 19 (1873) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 441	M.fr. VIII, 19; IX, 51.

ANIGOZANTHOS; Labillardière, Relat. du voy. à la rech. de La Perouse I, 411 (1798). (Anigosanthos, Macropidia, Macropodia, Schwaegrichenia.)

<i>A. rufa</i> ; Labillardière, Voy. I, 411, t. 22 (1798) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 442	M.fr. VIII, 21.
<i>A. pulcherrima</i> ; Hooker, Bot. Mag. t. 4180 (1845) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 443	M.fr. VIII, 21.
<i>A. flavida</i> ; Redouté et De Candolle, les Liliacées t. 176 (1807) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 443	M.fr. VIII, 21.
<i>A. Preissii</i> ; Endlicher in Lehm. pl. Preiss. II, 26 (1846) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 444	M.fr. VIII, 23.
<i>A. humilis</i> ; Lindley, Bot. Regist. XXV, App. 46 (1839) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 444	M.fr. VIII, 21.
<i>A. viridis</i> ; Endlicher in Lehm. pl. Preiss. II, 25 (1846) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 445	M.fr. VIII, 22.
<i>A. Manglesii</i> ; D. Don in Sweet, Brit. Fl. Gard. sec. ser. t. 265 (1836) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 445	M.fr. VIII, 22.
<i>A. bicolor</i> ; Endlicher in Lehm. pl. Preiss. II, 26 (1846) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 446	M.fr. VIII, 22.
<i>A. fuliginosus</i> ; Hooker, Bot. Mag. t. 4291 (1847) ...	...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 447	M.fr. VIII, 20.

AMARYLLIDEEAE.

J. ST. HILAIRE, Exposit. des famill. natur. des plantes I, 134, t. 21 (1805).—Genera 7, Species 21.

CAMPYNEMA, Labillardière, Nov. Holl. plant. spec. I, 93 (1804).

<i>C. lineare</i> ; Labillardière, Nov. Holl. pl. spec. I, 93, t. 121 (1804) ...	...	...	—	—	T.	—	—	—	—	B.fl. VI, 415	M.fr. VI, 96.
--	-----	-----	---	---	----	---	---	---	---	---------------	---------------

HYPOXIS; Linné, syst. ed. decim. 986 (1759).

<i>H. hygrometrica</i> ; Labillardière, Nov. Holl. pl. spec. I, 82, t. 108 (1804) ...	...	—	S.A.	T.	V.	N.S.W.	Q.	—	—	B.fl. VI, 449	M.fr. VI, 96.
<i>H. glabella</i> ; R. Brown, prodr. 289 (1810) ...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl. VI, 450	M.fr. VI, 96.

- H. occidentalis*; Benth. Fl. Austr. VI, 451 (1873) ... W.A. — — — — — B.fl. VI, 451
H. marginata; R. Brown, prodr. 289 (1810) ... — — — — — N.A. B.fl. VI, 451

CURCULIGO; Gaertner, de fruct. I, 63, t. 16 (1788).

- C. recurvata*; Aiton, hort. Kew. sec. ed. II, 253 (1811) ... — — — — — Q. — B.fl. VI, 448 M.fr. VIII, 27 ... As.
C. ensifolia; R. Brown, prodr. 290 (1810) ... — — — — — N.S.W. Q. N.A. B.fl. VI, 448 M.fr. VI, 96... ... Aa., P.

DORYANTHES; Correa in Transact. Linn. Soc. VI, 211 (1802).

- D. excelsa*; Correa in Transact. Linn. Soc. VI, 211, t. 23 (1802) ... — — — — — N.S.W. Q. — B.fl. VI, 452 M.fr. IX, 51.
D. Palmeri; W. Hill in Benth. Fl. Austr. VI, 453 (1873) ... — — — — — Q. — B.fl. VI, 452 M.fr. IX, 51.

CRINUM; Linné, gen. plant. 97 (1737).

- C. pedunculatum*; R. Brown, prodr. 297 (1810) ... — — — — — N.S.W. Q. — B.fl. VI, 455 M.fr. III, 23.
C. Asiaticum; Linné, spec. plant. 292 (1753)... — — — — — Q. N.A. B.fl. VI, 454 M.fr. III, 23 ... As.
C. venosum; R. Brown, prodr. 297 (1810) ... — — — — — Q. N.A. B.fl. VI, 453 M.fr. III, 22.
C. brachyandrum; Herbert in Bot. Mag. t. 2121 (1820) ... — — — — — Q. — — M.fr. XI, 139.
C. flaccidum; Herbert in Bot. Mag. t. 2133 (1820) ... W.A. S.A. — V. N.S.W. Q. N.A. B.fl. VI, 454 M.fr. III, 23.
C. angustifolium; R. Brown, prodr. 297 (1810) ... — — — — — Q. — — M.fr. III, 23; XI, 139.
C. uniflorum; F. v. M., fragm. III, 23 (1862) ... — — — — — Q. N.A. B.fl. VI, 454 M.fr. VI, 251.

CALOSTEMMA; R. Brown, prodr. fl. Nov. Holl. 297 (1810).

- C. purpureum*; R. Brown, prodr. 298 (1810) ... — S.A. — V. N.S.W. — — B.fl. VI, 457
C. luteum; Sims, Bot. Mag. t. 2101 (1819) ... — S.A. — — N.S.W. Q. — B.fl. VI, 457

EURYCYLES; Salisbury in Transact. of the Hort. Soc. I, 337 (1812).

- E. silvestris*; Salisbury in Transact. Hort. Soc. I, 337 (1812) ... — — — — — Q. — B.fl. VI, 456 As.
E. Cunninghamii; Aiton in Loud. Hort. Brit. Suppl. 588 (1832) ... — — — — — N.S.W. Q. — B.fl. VI, 456
E. alba; F. v. M., Calostemma; R. Brown, prodr. 298 (1810) ... — — — — — Q. N.A. B.fl. VI, 458

DIOSCORIDEAE.

DU MORTIER, Analyse des familles des plantes 57 (1829), from R. Brown (1810).—Genera 2, Species 4.

DIOSCOREA; Plumier, nov. plant. Americ. gen. 9, t. 26 (1703). (Dioscorideae, Helmia.)

- D. hastifolia*; Nees in Lehm. pl. Preisa. II, 33 (1846) ... W.A. — — — — — B.fl. VI, 461 M.fr. VII, 80; XI, 20.
D. transversa; R. Brown, prodr. 295 (1810) ... — — — — — N.S.W. Q. N.A. B.fl. VI, 460 M.fr. VII, 80.
D. sativa; Linné, spet. plant. 1033 (1753) ... — — — — — Q. N.A. B.fl. VI, 461 M.fr. VII, 81 ... Aa., P.

PETERMANNIA; F. v. M., fragm. II, 92 (1860).

- P. cirrosa*; F. v. M., fragm. II, 93 (1860) ... — — — — — N.S.W. — — B.fl. VI, 462 M.fr. II, 176.

EUCALYCEAE HYPOGYNAE.

F. v. M. in Woolla's plants of the neighb. of Sydney 44 (1880).

ROXBURGHIAEAE.

LINDLEY in Wallich, pl. Asiat. rar. III, 50 (1832).—Genus 1, Species 1.

ROXBURGHIA; Jones in Roxburgh, pl. Corom. I, 29, t. 32 (1795). (Stemona, 1790.)

- R. Javanica*; Kunth, enum. V, 288 (1850) ... W.A. — — — — — Q. N.A. B.fl. VII, 1 M.fr. VII, 81 ... As.

LILIACEAE.

HALLEK, enum. stirp. Helv. I, 279 (1742).—Genera 45, Species 161.

SMILAX; Tournefort, inst. 654, t. 421 (1700), from Theophrastos and Dioscorides.

S. glycyphylla; Smith in White, Voy. to N.S. Wales 230 (1790) ...	...	...	...	...	...	N.S.W.	Q.	—	B.fl. VII, 7	M.fr. VII, 77.	
S. Australis; R. Brown, prodr. 293 (1810) ...	...	...	...	...	...	V. N.S.W.	Q.	N.A.	B.fl. VII, 7	M.fr. VII, 78.	
S. purpurata; G. Forster, fl. ins. Austr. prodr. 70 (1786) ...	...	...	...	...	...	N.S.W.	—	—	—	—	

RHIPOGONUM; R. & G. Forster, char. gen. 49, t. 25 (1776). (Ripogonum.)

R. album; R. Brown, prodr. 293 (1810) ...	...	...	...	...	...	V. N.S.W.	Q.	—	B.fl. VII, 9	M.fr. I, 43, 243; VII, 79; VIII, 280.	
R. discolor; F. v. M., fragm. VII, 78 (1870) ...	...	...	...	...	...	N.S.W.	Q.	—	B.fl. VII, 9	M.fr. VII, 78.	
R. Fawcettianum; F. v. M. in Benth. Fl. Austr. VII, 9 (1878) ...	...	...	...	...	...	N.S.W.	Q.	—	B.fl. VII, 9	M.fr. VII, 80.	
R. Elseyanum; F. v. M., fragm. I, 44 (1858) ...	...	...	...	...	...	N.S.W.	Q.	—	B.fl. VII, 10	M.fr. VII, 80.	

DRYMOPHILA; R. Brown, prodr. fl. Nov. Holl. 292 (1810).

D. cyanocarpa; R. Brown, prodr. 292 (1810) ...	...	...	...	...	...	T. V. N.S.W.	—	—	B.fl. VII, 12	M.fr. VII, 73; VIII, 280; IX, 191.	
D. Moorei; Baker in Journ. Linn. Soc. XIV, 571 (1875) ...	...	...	...	...	...	N.S.W.	—	—	B.fl. VII, 13	M.fr. IX, 190.	

DIANELLA; Lamarck, Encyclop. method. II, t. 250 (1786).

D. Tasmanica; J. Hooker, Fl. Tasman. II, 57, t. 133 (1860) ...	...	...	...	...	...	T. V. N.S.W.	—	—	B.fl. VII, 14	M.fr. VI, 121; VIII, 279.	
D. longifolia; R. Brown, prodr. 280 (1810) ..	...	...	...	...	...	S.A. T. V. N.S.W.	Q.	—	B.fl. VII, 14	M.fr. VI, 122.	
D. revoluta; R. Brown, prodr. 280 (1810) ...	...	...	...	...	...	W.A. S.A. T. V. N.S.W.	Q.	—	B.fl. VII, 15	M.fr. VI, 121; XI, 20.	
D. caerulea; Sims, Bot. Mag. t. 505 (1799) ...	...	...	...	...	...	N.S.W.	Q.	—	B.fl. VII, 16	M.fr. VI, 123.	
D. ensifolia; De Candolle & Redouté, Liliac. t. 1 (1802) ...	...	...	...	...	...	—	Q. N.A.	B.fl. VII, 16	M.fr. VI, 123	...	As., Afr., P.

ASPARAGUS; Tournefort, inst. 300, t. 154 (1700), from Theophrastos, Dioscorides, and Plinius. (Asparagopsis.)

A. racemosus; Willdenow, spec. plant. II, 152 (1799) ...	...	...	...	...	...	—	Q. N.A.	B.fl. VII, 17	M.fr. VII, 73	...	As., Afr.
--	-----	-----	-----	-----	-----	---	---------	---------------	---------------	-----	-----------

EUSTREPHUS; R. Brown, prodr. 281 (1810).

E. Brownii; F. v. M., fragm. VII, 73 (1870) ...	...	...	...	...	...	V. N.S.W.	Q.	—	B.fl. VII, 18	M.fr. VII, 73; VIII, 280.	P.
---	-----	-----	-----	-----	-----	-----------	----	---	---------------	---------------------------	----

GEITONOPLESIMUM; Cunningham in Bot. Magaz. t. 3131 (1832). (Luzuriago partly, Culcoa.)

G. cymosum; Cunningham in Bot. Mag. t. 3131 (1832) ...	...	...	...	...	...	V. N.S.W.	Q.	—	B.fl. VII, 19	M.fr. VII, 74	...	As., P.
--	-----	-----	-----	-----	-----	-----------	----	---	---------------	---------------	-----	---------

ELACHANTHERA; F. v. M., in Vict. Naturalist III, 108 (1886).

E. Sewelliae; F. v. M. in Vict. Natural. III, 108 (1886) ...	...	...	...	...	...	—	N.A.	—	—	—	
--	-----	-----	-----	-----	-----	---	------	---	---	---	--

DRACAENA; Vandelli, dissertatio de Dracaena t. 1 (1762).

D. angustifolia; Roxburgh, Fl. Indic. II, 155 (1832) ...	...	...	...	...	...	—	Q. N.A.	B.fl. VII, 20	M.fr. V, 194; VI, 120; VIII, 279; IX, 190.	As.,	[P.]
--	-----	-----	-----	-----	-----	---	---------	---------------	--	------	------

CORDYLINE; Commerçon in R. Hedwig, gen. plant. 243 (1806). (Charlwoodia.)

C. terminalis; Kunth in act. acad. Berol. 30 (1820) ...	...	...	...	...	...	N.S.W.	Q.	—	B.fl. VII, 21	M.fr. V, 196	...	As., P.
C. Murchisoniae; F. v. M., fragm. V, 195 (1866) ...	...	...	...	...	...	—	Q.	—	B.fl. VII, 22	M.fr. V, 195		
C. stricta; Endlicher in Ann. Wien. Mus. I, 162 (1838) ...	...	...	...	...	...	N.S.W.	Q.	—	B.fl. VII, 22	M.fr. V, 195; VI, 121.		
C. Baueri; J. Hooker in Gardeners' Chronicle 792 (1860) ...	...	...	...	...	...	N.S.W.	—	—	—	—		

PHORMIUM; R. & G. Forster, charact. gener. 47, t. 24 (1776). (Chlamydia.)

P. tenax; R. & G. Forster, char. gen. 47, t. 24 (1776) ...	...	...	...	...	...	N.S.W.	—	—	—	...	...	N.Z.
--	-----	-----	-----	-----	-----	--------	---	---	---	-----	-----	------

BLANDFORDIA; Smith, Exot. Bot. I, 5, t. 4 (1804).

B. marginata; Herbert in Bot. Regist. Misc. 84 (1842) ...	...	...	...	...	...	T.	—	—	—	B.fl. VII, 23	M.fr. VII, 71.	
B. grandiflora; R. Brown, prodr. 296 (1810) ...	...	...	...	...	...	N.S.W.	—	—	—	B.fl. VII, 23		
B. nobilis; Smith, Exot. Bot. I, 5, t. 4 (1804) ...	...	...	...	...	...	N.S.W.	—	—	—	B.fl. VII, 24	M.fr. VII, 70.	
B. flammea; Hooker, Bot. Mag. t. 4819 (1854) ...	...	...	...	...	...	N.S.W.	Q.	—	—	B.fl. VII, 24		

HEWARDIA; Hooker, icon. plant. 858 (1852).

H. Tasmanica; Hooker, icon. plant. t. 858 (1852) ...	...	...	...	...	...	T.	—	—	—	B.fl. VII, 25	M.fr. VII, 72.	
--	-----	-----	-----	-----	-----	----	---	---	---	---------------	----------------	--

ASTELIA ; Banks & Solander in R. Brown, prodr. 291 (1810). (Hamelinia, Funkia, 1805).

A. alpina ; R. Brown, prodr. 291 (1810) — — T. V. N.S.W. — — B.fl.VII,11

MILLIGANIA ; J. Hooker in Kew misc. V, 296, t. 9 (1853).

M. densiflora ; J. Hooker, Fl. Tasman. II, 62 (1860) — — T. — — — — B.fl.VII,26
 M. longifolia ; J. Hooker in Kew misc. V, 296, t. 9 (1853) — — T. — — — — B.fl.VII,26
 M. Johnstoni ; F. v. M. in Benth. Fl. Austr. VII, 26 (1878) — — T. — — — — B.fl.VII,26
 M. stylosa ; F. v. M. in papers of the R. S. of Tasm. 11 (1876) — — T. — — — — B.fl.VII,27

WURMBESEA ; Thunberg, nov. gen. I, 18 (1781). (Anguillaria, Wurmbesa, Melanthium partly, Pleea partly.)

W. tubulosa ; Benth. Fl. Austr. VII, 28 (1878) W.A. — — — — — — B.fl.VII,28 M.fr.VII,75.
 W. dioica ; F. v. M., fragm. X, 119 (1877) W.A. S.A. T. V. N.S.W. Q. N.A. B.fl.VII,30 M.fr.VII,74.

IPHIGENIA ; Kunth, enum. plant. omn. hucusq. cognit. IV, 212 (1843)

I. Indica ; Kunth, enum. plant. IV, 213 (1843) — — — — — — Q. N.A. B.fl.VII,31 M.fr.VII,74; IX,193 ... As.

SCHELHAMMERA ; R. Brown, prodr. 273 (1810). (Pardunya, 1806).

S. undulata ; R. Brown, prodr. 274 (1810) — — — — V. N.S.W. — — B.fl.VII,31 M.fr.VII,71.
 S. multiflora ; R. Brown, prodr. 274 (1810) — — — — — — Q. — — B.fl.VII,32 M.fr.VII,72.

KREYSSIGIA ; Reichenbach, iconogr. bot. exotic. III, 11, t. 229 (1829). (Kreysigia, Tripladenia.)

K. Cunninghami ; F. v. M., fragm. VII, 71 (1870) — — — — — — N.S.W. Q. — — B.fl.VII,32 M.fr.VII,71; VIII,280.

BURCHARDIA ; R. Brown, prodr. 272 (1810).

B. umbellata ; R. Brown, prodr. 272 (1810) W.A. S.A. T. V. N.S.W. Q. — — B.fl.VII,33 M.fr.VII,77; VIII,280.

BULBINE ; Linné, hort. Cliffort. 122 (1737). (Anthericum partly, Chrysobactron.)

B. bulbosa ; Haworth, revis. pl. succul. 33 (1821) — — S.A. T. V. N.S.W. Q. — — B.fl.VII,34 M.fr.VII,70; XI,27.
 B. semibarbata ; Haworth, rev. pl. succ. 33 (1821) W.A. S.A. T. V. N.S.W. Q. — — B.fl.VII,35 M.fr.VII,70.

AGROSTOCRINUM ; F. v. M., fragm. phytogr. Austr. II, 95 (1860).

A. stypanoides ; F. v. M., fragm. II, 95 (1860) W.A. — — — — — — B.fl.VII,36 M.fr.III,165; VII,65.

THYSANOTUS ; R. Brown, prodr. 282 (1810). (Thysanella 1806, Chlamysporum 1809, Isandra.)

T. multiflorus ; R. Brown, prodr. 285 (1810) W.A. — — — — — — B.fl.VII,38 M.fr.VII,69.
 T. triandrus ; R. Brown, prodr. 284 (1810) W.A. — — — — — — B.fl.VII,39 M.fr.VII,69.
 T. glaucus ; Endlicher in Lehm. pl. Preiss. II, 38 (1846) W.A. — — — — — — B.fl.VII,39
 T. Drummondii ; Baker in Journ. Linn. Soc. XV, 341 (1876) W.A. — — — — — — B.fl.VII,39
 T. pauciflorus ; R. Brown, prodr. 285 (1810) W.A. — — — — — — B.fl.VII,39
 T. asper ; Lindley, Bot. Regist. XXV, App. 58 (1839) W.A. — — — — — — B.fl.VII,39
 T. chrysantherus ; F. v. M. in Benth. flor. Hongk. 372 (1861) — — — — — — Q. N.A. B.fl.VII,40 M.fr.V,202 As.
 T. exiliflorus ; F. v. M. in Transact. R. S. of S. Austr. IV, 112 (1881) — — S.A. — — — — — — M.fr.XI,139.
 T. isantherus ; R. Brown, prodr. 283 (1810) W.A. — — — — — — B.fl.VII,40
 T. tenellus ; Endlicher in Lehm. pl. Preiss. II, 37 (1846) W.A. — — — — — — B.fl.VII,40
 T. scaber ; Endlicher in Lehm. pl. Preiss. II, 37 (1846) W.A. — — — — — — B.fl.VII,41
 T. tuberosus ; R. Brown, prodr. 282 (1810) — — S.A. — — V. N.S.W. Q. N.A. B.fl.VII,41 M.fr.VII,69.
 T. exasperatus ; F. v. M., fragm. I, 21 (1858) — — S.A. — — — — — — B.fl.VII,42 M.fr.I,21.
 T. thyrsoides ; Baker in Journ. Linn. Soc. XV, 336 (1876) W.A. — — — — — — B.fl.VII,42
 T. Baueri ; R. Brown, prodr. 283 (1810) — — S.A. — — V. N.S.W. Q. — — B.fl.VII,42 M.fr.I,22; VII,69.
 T. Patersoni ; R. Brown, prodr. 284 (1810) W.A. S.A. T. V. N.S.W. — — — — B.fl.VII,43 M.fr.VII,69; XI,20.
 T. junceus ; R. Brown, prodr. 283 (1810) — — — — — — N.S.W. — — B.fl.VII,43 M.fr.VII,69.
 T. dichotomus ; R. Brown, prodr. 284 (1810) W.A. S.A. — — V. — — — — B.fl.VII,44 M.fr.VII,69.
 T. arbuscula ; Baker in Journ. Linn. Soc. XV, 339 (1876) W.A. — — — — — — B.fl.VII,45
 T. anceps ; Lindley, Bot. Regist. XXV, App. 58 (1839) W.A. — — — — — — B.fl.VII,45

HODGSONIOLA; F. v. M., fragm. II, 176 (1861). (Hodgsonia).

H. junciformis; F. v. M., fragm. II, 176 (1861) W.A. — — — — — B.fl.VII,46 M.fr.II,96,176.

CAESIA; R. Brown, prodr. fl. Nov. Holl. 277 (1810).

C. vittata; R. Brown, prodr. 277 (1810) — S.A. T. V. N.S.W. Q. — B.fl.VII,47 M.fr.I,63; VII,68.
 C. chlorantha; F. v. M., fragm. I, 63 (1858) — — — — — Q. N.A. — M.fr.I,63.
 C. parviflora; R. Brown, prodr. 277 (1810) — S.A. T. V. N.S.W. Q. — B.fl.VII,47 M.fr.VII,67.
 C. rigidifolia; F. v. M., fragm. X, 48 (1876) W.A. — — — — — B.fl.VII,47 M.fr.X,48.
 C. setifera; Baker in Journ. Linn. Soc. XV, 359 (1876) — — — — — N.A. B.fl.VII,47

CHAMAESCILLA; F. v. M., fragm. VII, 68 (1870). (Caesia partly.)

C. corymbosa; F. v. M., fragm. VII, 68 (1870) W.A. S.A. T. V. N.S.W. — — B.fl.VII,48 M.fr.VII,68.
 C. spiralis; F. v. M., fragm. VII, 68 (1870) W.A. — — — — — B.fl.VII,48 M.fr.VII,68.

CORYNOTHECA; F. v. M., fragm. VII, 68 (1870). (Caesia partly.)

C. lateriflora; F. v. M., fragm. VII, 68 (1870) W.A. S.A. — V. N.S.W. Q. N.A. B.fl.VII,49 M.fr.I,215.
 C. dichotoma; F. v. M., fragm. VII, 68 (1870) W.A. — — — — — B.fl.VII,50 M.fr.I,215; XI,20.
 C. acanthoclada; F. v. M., fragm. VII, 68 (1870) W.A. — — — — — B.fl.VII,50 M.fr.I,215; XI,20.

TRICORYNE; R. Brown, prodr. fl. Nov. Holl. 278 (1810).

T. platyptera; G. Reichenbach, Beitr. zur syst. Pflanzenk. 72 (1871) — — — — — Q. — B.fl.VII,51
 T. anceps; R. Brown, prodr. 278 (1810) — — — — — Q. — B.fl.VII,51 M.fr.VII,67.
 T. muricata; Baker in Journ. Linn. Soc. XV, 363 (1876) — — — — — Q. — B.fl.VII,52
 T. elatior; R. Brown, prodr. 278 (1810) W.A. S.A. T. V. N.S.W. Q. — B.fl.VII,52 M.fr.VII,67; XI,20.
 T. simplex; R. Brown, prodr. 278 (1810) — — — V. N.S.W. Q. — B.fl.VII,52 M.fr.VII,67.
 T. humilis; Endlicher in Lehm. pl. Preiss. II, 36 (1846) W.A. — — — — — B.fl.VII,53 M.fr.VII,67.

STYPANDRA; R. Brown, prodr. flor. Nov. Holl. 278 (1810).

S. glauca; R. Brown, prodr. 278 (1810) W.A. — — V. N.S.W. Q. — B.fl.VII,53 M.fr.VII,64.
 S. caespitosa; R. Brown, prodr. 279 (1810) — — T. V. N.S.W. Q. — B.fl.VII,54 M.fr.VII,65.

ARTHROPODIUM; R. Brown, prodr. 276 (1810). (Dichopogon, Siona.)

A. paniculatum; R. Brown, prodr. 276 (1810) — S.A. T. V. N.S.W. Q. — B.fl.VII,56 M.fr.VII,66.
 A. minus; R. Brown, prodr. 276 (1810) — S.A. T. V. N.S.W. — — B.fl.VII,56 M.fr.VII,67; XI,27.
 A. capillipes; Endlicher in Lehm. pl. Preiss. II, 34 (1846) W.A. — — — — — B.fl.VII,57 M.fr.VII,67.
 A. Preissii; Endlicher in Lehm. pl. Preiss. II, 35 (1846) W.A. — — — — — B.fl.VII,57 M.fr.VII,66.
 A. dianellaceum; F. v. M., fragm. X, 65 (1876) — — — — — Q. — B.fl.VII,57 M.fr.X,65.
 A. strictum; R. Brown, prodr. 276 (1810) — S.A. T. V. N.S.W. Q. — B.fl.VII,58 M.fr.VII,66.
 A. fimbriatum; R. Brown, prodr. 276 (1810) — S.A. — V. N.S.W. Q. — B.fl.VII,59

CHLOROPHYTUM; Ker in Bot. Mag. t. 1071 (1808). (Chlorophyton, Phalangium partly.)

C. laxum; R. Brown, prodr. 277 (1810) — — — — — N.A. B.fl.VII,60 M.fr.I,63; VII,71 ... As., Afr.
 C. alpinum; Baker in Journ. Linn. Soc. XV, 329 (1876) — — T. — — — — B.fl.VII,60

HERPOLIRION; J. Hooker, Fl. of New Zealand I, 258 (1853).

H. Novae Zelandiae; J. Hooker, Fl. of New Zeal. I, 258 (1853) — — T. V. N.S.W. — — B.fl.VII,61 M.fr.VII,68; VIII,280. N.Z.

SOWERBAEA; Smith in Transact. Linn. Soc. V, 159 (1800). (Sowerbya.)

S. juncea; Smith in Transact. Linn. Soc. V, 160, t. 6 (1800) — — — V. N.S.W. Q. — B.fl.VII,61 M.fr.VI,180.
 S. laxiflora; Lindley, Botan. Regist. new ser. IV, t. 10 (1841) W.A. — — — — — B.fl.VII,62 M.fr.VI,180; VII,87.
 S. alliacea; F. v. M., fragm. VI, 180 (1868) — — — — — N.A. B.fl.VII,62 M.fr.VII,87.

ALLANIA; Endlicher, gen. plant. 151 (1837). (Alania.)

A. Endlicheri; Kunth, enum. IV, 644 (1843) — — — — — N.S.W. — — B.fl.VII,63 M.fr.VI,180.

BARTLINGIA; F. v. M., fragm. VII, 88 (1870). (Laxmannia.)

B. grandiflora; F. v. M., Cens. 118 (1882) from Lindley (1839)	W.A.	—	—	—	—	—	B.fl.VII,64	M.fr.I,159; VII,88.
B. squarrosa; F. v. M., Cens. 118 (1882) from Lindley (1839)	W.A.	—	—	—	—	—	B.fl.VII,64	
B. minor; F. v. M., Cens. 118 (1882) from R. Brown (1810)	W.A.	—	—	—	—	—	B.fl.VII,65	M.fr.VII,88,89.
B. gracilis; F. v. M., fragm. VII, 88 (1870)	—	—	—	V.	N.S.W.	Q.	B.fl.VII,65	M.fr.I,159.
B. ramosa; F. v. M., fragm. VII, 88 (1870)	W.A.	—	—	—	—	—	B.fl.VII,66	M.fr.VI,255.
B. sessiliflora; F. v. M. in pap. R. S. of Tasm. 116 (1877)	W.A.	S.A.	T.	V.	N.S.W.	—	B.fl.VII,66	
B. brachyphylla; F. v. M., Cens. of Austr. pl. 119 (1882)	W.A.	—	—	—	—	—	B.fl.VII,66	M.fr.I,158.
B. sessilis; F. v. M., Cens. 119 (1882) from Lindley (1839)	W.A.	—	—	—	—	—	B.fl.VII,67	

STAWELLIA; F. v. M., fragm. phytogr. Austr. VII, 85 (1870).

S. dimorphantha; F. v. M., fragm. VII, 85 (1870)	W.A.	—	—	—	—	—	B.fl.VII,67	M.fr.VII,85.
---	------	---	---	---	---	---	-------------	--------------

JOHNSONIA; R. Brown, prodr. fl. Nov. Holl. 287 (1810).

J. lupulina; R. Brown, prodr. 287 (1810)	W.A.	—	—	—	—	—	B.fl.VII,68	M.fr.VII,86.
J. pubescens; Lindley, Bot. Regist. XXV, App. 57 (1839)	W.A.	—	—	—	—	—	B.fl.VII,68	M.fr.VI,255; VII,87.

ARNOCRINUM; Endlicher & Lehmann in pl. Preiss. II, 41 (1846).

A. Drummondii; Endlicher in Lehm. pl. Preiss. II, 41 (1846)	W.A.	—	—	—	—	—	B.fl.VII,69	M.fr.VII,73.
A. Preissii; Lehmann, pl. Preiss. II, 42 (1846)	W.A.	—	—	—	—	—	B.fl.VII,70	

BORYA; Labillardière, Nov. Holl. pl. specim. 81, t. 107 (1804).

B. nitida; Labillardière, Nov. Holl. pl. spec. 81, t. 107 (1804)	W.A.	—	—	—	—	—	B.fl.VII,71	M.fr.VII,87.
B. septentrionalis; F. v. M., fragm. V, 41 (1865)	—	—	—	—	—	Q.	B.fl.VII,71	M.fr.V,41.

CALECTASIA; R. Brown, prodr. fl. Nov. Holl. 263 (1810).

C. cyanea; R. Brown, prodr. 264 (1810)	W.A.	S.A.	—	V.	—	—	B.fl.VII,121	
---	------	------	---	----	---	---	--------------	--

BAXTERIA; R. Brown in Hook. Lond. Journ. of Bot. II, 492, t. 13-15 (1843).

B. Australis; R. Brown in Hook. Lond. Journ. II, 492 (1843)	W.A.	—	—	—	—	—	B.fl.VII,120	
--	------	---	---	---	---	---	--------------	--

XEROTES; R. Brown, prodr. 257 (1810). (Lomandra 1804, Acanthocarpus, Chamaexeros.)

X. echinata; Cunningham in Benth. fl. Austr. VII, 112 (1878)	...	...	...	W.A.	—	—	—	—	—	B.fl.VII,111				
X. Serra; Endlicher in Lehm. pl. Preiss. II, 49 (1846)	...	...	...	W.A.	—	—	—	—	—	B.fl.VII,111	M.fr.VIII,211; XI,24.			
X. fimbriatus; F. v. M., fragm. VIII, 211 (1874)	...	...	...	W.A.	—	—	—	—	—	B.fl.VII,111	M.fr.VIII,211.			
X. Banksii; R. Brown, prodr. 263 (1810)	...	...	...	...	—	—	—	—	Q.	B.fl.VII,96	M.fr.VIII,213			
X. dura; F. v. M. in Transact. Vict. Inst. 42 (1854)	...	...	...	..	—	S.A.	—	N.S.W.	—	B.fl.VII,97	M.fr.VIII,207.			
X. longifolia; R. Brown, prodr. 262 (1810)	...	...	...	...	—	S.A.	T.	V.	N.S.W.	Q	N.A.	B.fl.VII,98	M.fr.VIII,210.	
X. rigida; R. Brown, prodr. 262 (1810)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.VII,99	M.fr.VIII,209.		
X. Sonderi; F. v. M., fragm. VIII, 206 (1874)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.VII,99	M.fr.		
X. odora; Endlicher in Lehm. pl. Preiss. II, 50 (1846)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.VII,99			
X. Brownii; F. v. M., fragm. VIII, 206 (1874)	...	...	...	...	...	—	S.A.	—	V.	N.S.W.	Q	N.A.	B.fl.VII,100	M.fr.VIII,206.
X. Ordii; F. v. M., fragm. XI, 23 (1878)	...	...	...	...	...	W.A.	—	—	—	—	—		M.fr.XI,137.	
X. sororia; F. v. M., first gen. Report 15 (1854)	...	...	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VII,100	
X. Endlicheri; F. v. M., fragm. VIII, 205 (1874)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.VII,101	M.fr.VIII,205.	
X. sericea; Endlicher in Lehm. pl. Preiss. II, 51 (1846)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.VII,101	M.fr.VIII,207.	
X. purpurea; Endlicher in Lehm. pl. Preiss. II, 49 (1846)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.VII,101	M.fr.VIII,213.	
X. Preissii; Endlicher in Lehm. pl. Preiss. II, 50 (1846)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl.VII,102		
X. effusa; Lindley in Mitch. Three Exped. II, 151 (1838)	...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VII,102	M.fr.VIII,209.
X. micrantha; Endlicher in Lehm. pl. Preiss. II, 49 (1846)	...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.VII,103	M.fr.VIII,212.
X. Thunbergii; F. v. M., fragm. VIII, 208 (1874)	...	...	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VII,103	M.fr.VIII,208.
X. caespitosa; Benth. Fl. Austr. VII, 104 (1878)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.VII,104	
X. pauciflora; R. Brown, prodr. 261 (1810)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.VII,104	M.fr.212.
X. flexifolia; R. Brown, prodr. 260 (1810)	...	...	...	...	...	—	—	—	N.S.W.	—	—	—	B.fl.VII,105	M.fr.VIII,207.

X. glauca; R. Brown, prodr. 260 (1810)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII, 105	M.fr.VIII, 208.
X. elongata; Benth. Fl. Austr. VII, 106 (1878)	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.VII, 106	
X. rupestris; Endlicher in Lehm. pl. Preiss. II, 50 (1846)	W.A.	—	—	—	—	—	—	B.fl.VII, 106	
X. collina; R. Brown, prodr. 260 (1810)	W.A.	—	—	—	—	—	—	B.fl.VII, 107	
X. suaveolens; Endlicher in Lehm. pl. Preiss. II, 50 (1846)	W.A.	—	—	—	—	—	—	B.fl.VII, 107	M.fr.VIII, 207.
X. turbinata; Endlicher in Lehm. pl. Preiss. II, 51 (1846)	W.A.	—	—	—	—	—	—	B.fl.VII, 107	M.fr.VIII, 211.
X. apartea; Endlicher in Lehm. pl. Preiss. II, 51 (1846)	W.A.	—	—	—	—	—	—	B.fl.VII, 108	M.fr.VIII, 211.
X. juncea; F. v. M. in Transact. Vict. Inst. 135 (1855)	—	S.A.	—	V.	—	—	—	B.fl.VII, 108	M.fr.VIII, 210.
X. leucocephala; R. Brown, prodr. 260 (1810)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VII, 109	M.fr.VIII, 210; IX, 195.
X. hastilis; R. Brown, prodr. 263 (1810)	W.A.	—	—	—	—	—	—	B.fl.VII, 110	M.fr.VIII, 210; XI, 54.

DASYPOGON; R. Brown, prodr. fl. Nov. Holl. 263 (1810).

D. bromelifolius; R. Brown, prodr. 263 (1810)	W.A.	—	—	—	—	—	—	B.fl.VII, 118	M.fr.II, 112.
D. Hookeri; Drummond in Hook. Lond. Journ. II, 168 (1843)	W.A.	—	—	—	—	—	—	B.fl.VII, 119	M.fr.II, 112; IV, 173.

XANTHORRHOEA; Smith in Transact. Linn. Soc. IV, 219 (1798).

X. gracilis; Endlicher in Lehm. pl. Preiss. II, 39 (1846)	W.A.	—	—	—	—	—	—	B.fl.VII, 114	M.fr.IV, 112.
X. pumilio; R. Brown, prodr. 288 (1810)	—	—	—	—	—	—	Q.	B.fl.VII, 114	
X. macronema; F. v. M., fragm. IV, 112 (1864)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII, 113	
X. hastilis; R. Brown, prodr. 288 (1810)	—	—	—	V.	N.S.W.	Q.	—	B.fl.VII, 115	M.fr.IV, 113.
X. resinosa; Persoon, syn. pl. I, 370 (1805)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII, 115	M.fr.IV, 113.
X. semiplana; F. v. M., fragm. IV, 111 (1864)	—	S.A.	—	—	—	—	—	B.fl.VII, 115	M.fr.IV, 111.
X. bracteata; R. Brown, prodr. 288 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.VII, 116	
X. minor; R. Brown, prodr. 288 (1810)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII, 114	M.fr.IV, 112.
X. Tateana; F. v. M. in Zeitschr. Oesterr. Apoth. Ver. XXIII, 294 (1885)	—	S.A.	—	—	—	—	—	—	
X. Australis; R. Brown, prodr. 288 (1810)	—	—	T.	V.	—	—	—	B.fl.VII, 116	M.fr.IV, 110.
X. quadrangulata; F. v. M., fragm. IV, 111 (1864)	—	S.A.	—	—	—	—	—	B.fl.VII, 117	M.fr.IV, 111.
X. Preissii; Endlicher in Lehm. pl. Preiss. II, 39 (1846)	W.A.	—	—	—	—	—	—	B.fl.VII, 117	M.fr.IV, 110.

KINGIA; R. Brown in King's Narrat. of survey of Austr. coasts II, 529 (1827).

K. Australis; R. Brown in King's Narr. of surv. II, 535 (1827)	W.A.	—	—	—	—	—	—	B.fl.VII, 119	
---	------	---	---	---	---	---	---	---------------	--

PALMAE.

RAY, method. pl. emend. 135 (1703).—Genera 10, Species 25.

CALAMUS; Linné, spec. plant. 325 (1753).

C. Australis; Martius, hist. nat. palm. III, 213 (1839)	—	—	—	—	—	Q.	—	B.fl.VII, 134	M.fr.V, 48.
C. Muelleri; Wendland & Drude in Linnaea XXXIX, 193 (1871)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII, 135	
C. caryotoides; A. Cunningham in Martius, hist. nat. palm. III, 212 (1839)	—	—	—	—	—	Q.	—	B.fl.VII, 135	

BACULARIA; F. v. M., fragm. VII, 103 (1870). (Linospadix.)

B. monostachya; F. v. M., fragm. VII, 103 (1870)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII, 136	M.fr.VII, 82, 193; XI, 58.
B. minor; F. v. M., fragm. XI, 58 (1878)	—	—	—	—	—	Q.	—	B.fl.VII, 137	M.fr.VIII, 235; IX, 195; XI, 137.

KENTIA; Blume in Bull. Neerl. 64 (1838). (Grisebachia, Hydriastele, Hedysepe, Kentiopsis, Veitchia partly, Howea, Howiea.)

K. Wendlandiana; F. v. M., fragm. VII, 102 (1870)	—	—	—	—	—	Q.	N.A.	B.fl.VII, 137	M.fr.VII, 102.
K. Belmoreana; C. Moore & F. v. M., fragm. VII, 99 (1870)	—	—	—	—	N.S.W.	—	—	B.fl.VII, 137	M.fr.VIII, 234.
K. Forsteriana; C. Moore & F. v. M., fragm. VII, 100 (1870)	—	—	—	—	N.S.W.	—	—	—	
K. Baueri; Seemann, fl. Vitiens. 269 (1869)	—	—	—	—	N.S.W.	—	—	—	M.fr.VII, 102; VIII, 233.
K. Canterburyana; C. Moore & F. v. M., fragm. VII, 101 (1870)	—	—	—	—	N.S.W.	—	—	B.fl.VII, 138	M.fr.VIII, 234; X, 119.
K. acuminata; Wendland & Drude in Garcke, Linnaea XXXIX, 207 (1871)	—	—	—	—	—	—	N.A.	B.fl.VII, 138	

CLINOSTIGMA ; Herm. Wendland in Seemann's Bonplandia X, 196 (1862). (Cyphokentia.)

C. Mooreanum ; F. v. M., fragm. VIII, 235 (1874) — — — — N.S.W. — — B.fl.VII,139 M.fr.VII,101; IX,195; X,120.

PTYCHOSPERMA ; Labillardière in Mém. de l'Inst. ann. 1808, I, 251, pl. X (1809). (Seaforthia, Laccospadix, Archontophoenix, Drymophlaeus, Calyptrocalyx).

P. Laccospadix ; Bentham, Fl. Austr. VII, 140 (1878) — — — — Q. — B.fl.VII,140 M.fr.XI,57.
 P. Beatricae ; F. v. M. in Melb. Chemist. Febr. (1882) — — — — Q. — —
 P. Alexandrae ; F. v. M., fragm. V, 47, t. 43 (1865) — — — — Q. — B.fl.VII,140 M.fr.V,47,196,213.
 P. Cunninghamii ; Herm. Wendland in Bot. Zeit. 346 (1858) — — — — N.S.W. Q. — B.fl.VII,141 M.fr.VIII,222.
 P. elegans ; Blume, Rumphia II, 118 (1836) — — — — Q. — B.fl.VII,141 M.fr.VIII,222.
 P. Normanbyi ; F. v. M., fragm. XI, 56 (1878) — — — — Q. — B.fl.VII,142 M.fr.VIII,235.

ARECA ; Linné, spec. plant. 1189 (1753) from Ray (1688).

A. Alicae ; F. v. M. in Regel's Gartenflora, 199-201 (1879) — — — — Q. — — M.fr.XI,88.

COCOS ; Linné, spec. plant. 1188 (1753).

C. nucifera ; Linné, spec. plant. 1188 (1753) — — — — Q. — B.fl.VII,143 As., Afr., Am., P.

CARYOTA ; Linné, gen. plant. 355 (1737).

C. Rumphiana ; Martius, hist. nat. palm. III, 195 (1839) — — — — Q. — B.fl.VII,144 As., P.

LICUALA ; Rumphius, herb. Amboin. I, 44, t. 9 (1741).

L. Muelleri ; Wendland & Drude in Garcke, Linnaea XXXIX, 223 (1871) — — — — Q. — B.fl.VII,145 M.fr.VIII,221; X,120. As.

LIVISTONA ; R. Brown, prodr. fl. Nov. Holl. 267 (1810). (Corypha partly.)

L. Leichhardtii ; F. v. M., fragm. V, 49 (1865) — — — — — N.A. B.fl.VII,146 M.fr.VIII,221; XI,55.
 L. Mariae ; F. v. M. in Giles's Geog. Trav. in Centr. Austr. 222 (1875) — S.A. — — — N.A. — M.fr.VIII,283; XI,54,55.
 L. Australis ; Martius, hist. nat. palm. III, 241 (1839) — — — — V. N.S.W. Q. — B.fl.VII,146 M.fr.XI,55.

NIPACEAE.

BROGNIART, enum. des genr. de plantes 15 (1843).—Genus 1, Species 1.

NIPA ; Wurm in Verhandl. Batav. Genootsch. I, 349 (1779) from Camellus, Ray & Rumphius. (Nypa.)

N. fruticans ; Wurm in Verhandl. Batav. Genootsch. I, 349 (1779) — — — — Q. — — M.fr.XI,128. ... As., P.

PANDANEAE.

R. BROWN, prodr. fl. Nov. Holl. (1810).—Genera 2, Species 11.

PANDANUS ; Rumphius, herb. Amboin. IV, 139, t. 74 (1750). (Athrodactylis, Fiquetia.)

P. odoratissimus ; Linné, fil. suppl. 424 (1781) — — — — Q. N.A. B.fl.VII,148 M.fr.IV,40; XI,55 ... As., P.
 P. pedunculatus ; R. Brown, prodr. 341 (1810) — — — — N.S.W. Q. — B.fl.VII,149 M.fr.IV,40; VIII,220; IX,193.
 P. Moorei ; F. v. M., Cens. 140 (1882) — — — — N.S.W. — — —
 P. Forsteri ; Moore & F. v. M., fragm. VIII, 220 (1874) — — — — N.S.W. — — B.fl.VII,149 M.fr.VIII,220.
 P. Solms-Laubachii ; F. v. M. in der bot. Zeitung (1887) — — — — Q. — —
 P. aquaticus ; F. v. M., fragm. V, 40 (1865) — — — — Q. N.A. B.fl.VII,148 M.fr.VIII,220.
 P. monticola ; F. v. M., fragm. V, 40 (1865) — — — — Q. — B.fl.VII,150 M.fr.VII,63; VIII,220; IX,193.

FREYCINETIA ; Gaudichaud in Annal. des scienc. natur. III, 509 (1824).

F. insignis ; Blume, Rumphia I, 158, t. 42 (1835) — — — — Q. — — ... As.
 F. Gaudichaudii ; Brown & Bennett, pl. Jav. rar. 31, t. 9 (1838) — — — — Q. — B.fl.VII,151 ... As.
 F. excelsa ; F. v. M., fragm. V, 39 (1865) — — — — Q. — B.fl.VII,151 M.fr.V,39.
 F. Baueriana ; Endlicher, prodr. fl. ins. Norf. 25 (1833) — — — — N.S.W. — — —

AROIDEAE.

A. L. DE JUSSIEU, gen. plant. 23 (1789), from B. de Jussieu (1759).—Genera 6, Species 10.

TYPHONIUM; Schott in Wien. Zeitschr. III, 72 (1829). (Arum partly.)

T. liliifolium; F. v. M. in Schott, prodr. syst. aroid. 107 (1860) ...	...	...	—	—	—	—	—	N.A.	B.f. VII, 153	M.fr. VIII, 187.	
T. alismifolium; F. v. M., fragm. VIII, 186 (1874) ...	...	...	—	—	—	—	—	Q.	B.f. VII, 153	M.fr. VIII, 186.	
T. Brownii; Schott, synops. Aroid. I, 11 (1855) ...	...	...	—	—	—	—	N.S.W.	Q.	N.A.	B.f. VII, 154	M.fr. VIII, 187.
T. angustilobum; F. v. M., fragm. X, 66 (1876) ...	...	...	—	—	—	—	—	Q.	—	B.f. VII, 154	M.fr. X, 66.

AMORPHOPHALLUS; Blume in Nouv. Ann. du Mns. III, 366 (1834). (Brachyspatha.)

A. variabilis; Blume, Rumphia I, 146, t. 35 (1835) ...	...	...	—	—	—	—	—	N.A.	B.f. VII, 155	M.fr. VIII, 87.	As.
--	-----	-----	---	---	---	---	---	------	---------------	-----------------	-----

COLOCASIA; Ray, Method. 157 (1682). (Caladium partly, Alocasia.)

C. antiquorum; Schott, Meletemat. botan. 18 (1832) ...	...	...	—	—	—	—	—	Q.	N.A.	B.f. VII, 155	M.fr. VIII, 87.	As., Afr., P.
C. macrorrhiza; Schott, Meletemat. botan. 18 (1832) ...	...	...	—	—	—	—	N.S.W.	Q.	—	B.f. VII, 155	M.fr. VIII, 87.	As., P.

RHAPHIDOPHORA; Hasskarl in Fl. Regensb. Beibl. II, 11 (1842). (Epipremnum partly.)

R. pertusa; Schott in Seem. Bonplandia V, 45 (1857) ...	...	...	—	—	—	—	—	Q.	—	B.f. VII, 156	M.fr. VIII, 87.	As., P.
---	-----	-----	---	---	---	---	---	----	---	---------------	-----------------	---------

GYMNOSTACHYS; R. Brown, prodr. fl. Nov. Holl. 337 (1810).

G. anceps; R. Brown, prodr. 337 (1810) ...	...	...	—	—	—	—	N.S.W.	Q.	—	B.f. VII, 157	M.fr. VIII, 87.	
--	-----	-----	---	---	---	---	--------	----	---	---------------	-----------------	--

POTHOS; Linné, amœn. acad. I, 410 (1747).

P. Loureiri; Hooker & Arnott, Bot. of Beech. Voy. 220 (1841) ...	...	...	—	—	—	—	N.S.W.	Q.	—	B.f. VII, 158	M.fr. I, 62; III, 163; VIII, 87.	As., P.
--	-----	-----	---	---	---	---	--------	----	---	---------------	----------------------------------	---------

TYPHACEAE.

A. L. DE JUSSIEU, Gen. plant. 25 (1789).—Genera 2, Species 2.

TYPHA; Tournefort, inst. 530, t. 301 (1700) from Theophrastos, Dioscorides and Plinius.

T. angustifolia; Linné, spec. plant. 971 (1753) ...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.f. VII, 159	M.fr. VII, 116.	E., As., Afr., Am., P., N.Z.
---	-----	-----	-----	------	------	----	----	--------	----	------	---------------	-----------------	------------------------------

SPARGANIUM; Tournefort, inst. 530, t. 302 (1700).

S. angustifolium; R. Brown, prodr. 338 (1810) ...	...	...	...	...	—	—	—	V.	N.S.W.	Q.	—	B.f. VII, 160	M.fr. VII, 117.
---	-----	-----	-----	-----	---	---	---	----	--------	----	---	---------------	-----------------

LEMNACEAE.

J. E. GRAY, Nat. arrang. of Brit. pl. II, 729 (1821).—Genera 2, Species 6.

LEMNA; Linné, syst. nat. 9 (1735); Fl. Lappon. 351 (1737). (Telmatophace, Spirodela.)

L. trisulca; Linné, spec. plant. 970 (1753)	...	...	...	...	...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.f. VII, 162	M.fr. VIII, 188.	E., As., Afr., Am.
L. minor; Linné, spec. plant. 970 (1753)	...	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.f. VII, 163	M.fr. VIII, 188.	E., As., Afr., Am., P., N.Z.
L. oligorrhiza; Kurz in Journ. Linn. Soc. IX, 267 (1867)	...	...	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.f. VII, 163	M.fr. VIII, 188.	As., P.
L. polyrrhiza; Linné, spec. plant. 970 (1753)	...	...	...	...	...	—	—	—	V.	N.S.W.	—	—	B.f. VII, 164	M.fr. VIII, 188.	E., As., Afr., Am.
L. gibba; Linné, spec. plant. 970 (1753)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.f. VII, 163	M.fr. VIII, 188.	E., As., Afr., Am., N.Z.

WOLFFIA; Horkel & Schleiden in Schlecht. Linnaea XIII, 389 (1839). (Wolfia.)

W. Michelii; Schleiden, Beitr. zur Botan. 233 (1844) ...	...	...	...	...	—	S.A.	—	V.	N.S.W.	—	—	B.f. VII, 162	M.fr. VIII, 187.	E., As., Afr.
--	-----	-----	-----	-----	---	------	---	----	--------	---	---	---------------	------------------	---------------

FLUVIALES.

VENTENAT, Tabl. du règn. végét. II, 80 (1799).—Genera 10, Species 36.

NAJAS; Linné, syst. nat. 9 (1735); gen. plant. 278 (1737). (Caulinia partly.)

N. major; Allioni, Fl. Pedem. II, 221 (1785) ...	...	...	...	...	...	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.f. VII, 181	M.fr. VIII, 218.	E., As., Afr., Am.
N. tenuifolia; R. Brown, prodr. 345 (1810) ...	...	...	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.f. VII, 181	M.fr. I, 94; VIII, 219.	As., Afr.

LEPILAENA; Drummond & Harvey in Hook. Kew misc. VII, 57 (1855). (Zannichellia partly, Hexatheca, Altheniae subgenus.)

L. Australis; J. Drummond in Hook. Kew misc. VII, 53 (1855) ...	...	...	W.A. S.A. — V. — — —	B.fl. VII, 179 M.fr. VIII, 217.
L. Preissii; F. v. M., fragm. VIII, 217 (1874) ...	...	...	W.A. S.A. T. V. — — —	B.fl. VII, 180 M.fr. VIII, 217.

DIPLANTHERA; Thouars, gen. nov. Madag. 3 (1806). (Halodule).

D. tridentata; Thouars, gen. nov. Madag. 3 (1806) ...	...	...	— — — — —	N.A. — M.fr. VIII, 218. As., Afr.
---	-----	-----	-----------	-----------------------------------

CYMODOCEA; C. Koenig, Annals of Botany II, 96, t. 7 (1806). (Amphibolis, Graumuelleria, Kernera partly.)

C. zosterifolia; F. v. M., Cens. 121 (1882) from Agardh. (1822) ...	...	...	W.A. S.A. T. V. — — —	B.fl. VII, 177 M.fr. IV, 113; VIII, 218; IX, 196.
C. ciliata; Ehrenberg in Garcke, Linnaea XXXV, 162 (1867) ..	...	...	— — — — —	Q. N.A. B.fl. VII, 178 M.fr. VIII, 218. As., Afr.
C. rotundata; Ascherson & Schweinfurth in Neum. Anl. zu wiss. Beob. 361 (1875)	...	...	— — — — —	N.A. — M.fr. X, 162. As.
C. serrulata; Ascherson & Magnus in Neum. Anleit. zu wiss. Beob. 362 (1875)	...	...	— — — — —	Q. — B.fl. VII, 178 M.fr. VIII, 218.
C. isoetifolia; Ascherson in Garcke, Linnaea XXXV, 163 (1867) ...	...	...	— — — — —	Q. N.A. B.fl. VII, 178 ... As.

ZOSTERA; Linné, Waesgoeta-Ressa, 166-168 et fig. (1747).

Z. nana; F. K. Mertens in Roth, enum. pl. phaner. Germ. 8 (1827) ...	...	...	— S.A. T. V. N.S.W. Q. —	B.fl. VII, 176 M.fr. VIII, 218. E., As., Afr., Am.
Z. Tasmanica; G. v. Martens in Garcke, Linnaea XXXV, 68 (1867) ...	...	...	— S.A. T. V. — — —	B.fl. VII, 176 M.fr. VIII, 218.

POSIDONIA; Koenig, Annals of Botany II, 95, t. 6 (1806). (Caulinia partly, Kernera partly.)

P. Australis; J. Hooker, Fl. Tasman. II, 43 (1860) ...	...	...	W.A. S.A. T. V. — — —	B.fl. VII, 175 M.fr. VI, 198; VIII, 218.
--	-----	-----	-----------------------	--

RUPPIA; Linné, syst. nat. 9 (1735); gen. plant. 277 (1737).

R. maritima; Linné, spec. plant. 127 (1753) ...	...	...	W.A. S.A. T. V. N.S.W. Q. —	B.fl. VII, 174 M.fr. VIII, 217. E., As., Afr., Am., N.Z.
---	-----	-----	-----------------------------	--

POTAMOGETON; Fuchs, histor. stirp. comment. 651 (1542).

P. tricarínatus; F. v. M. & A. Bennett in Brit. Journ. XXV, 178 (1887)	...	...	— — — — —	Q. N.A. —
P. natans; Linné, spec. plant. 126 (1753) ...	...	...	W.A. S.A. T. V. N.S.W. Q. N.A.	B.fl. VII, 170 M.fr. VIII, 217. E., As., Afr., Am., P., N.Z.
P. coloratus; Hornemann, fl. Danic. t. 1449 (1813) ...	...	...	— — — V. — — —	... E., As., Afr.
P. polygonifolia; Pourret in Mém. de Toulouse (1788) ...	...	...	— — T. V. N.S.W. — — —	... E., As., Afr.
P. lucens; Linné, spec. plant. 126 (1753) ...	...	...	— — — V. — — —	... E., As., Afr., Am.
P. Tepperi; A. Bennett in Brit. Journ. XXV, 177 (1887) ...	...	...	— S.A. — — — Q. — —	... N.Z.
P. Cheesemani; A. Bennett in Brit. Journ. of Bot. XXI, 66 (1883)	...	...	— — — T. — — —	... N.Z.
P. Javanicus; Hasskarl in Act. Soc. Ind. Neerl. I, 26 (1857) ...	...	...	— — — — N.S.W. Q. N.A.	B.fl. VII, 171 M.fr. VIII, 217. As.
P. Drummondii; Benth. Fl. Austr. VII, 171 (1878) ...	...	...	W.A. — — — — —	B.fl. VII, 171
P. perfoliatus; Linné, spec. plant. 126 (1753) ...	...	...	— — T. V. N.S.W. Q. —	B.fl. VII, 172 M.fr. VIII, 217. E., As., Afr., Am.
P. crispus; Linné, spec. plant. 126 (1753) ...	...	...	— S.A. — V. N.S.W. Q. N.A.	B.fl. VII, 172 M.fr. VIII, 217. E., As., Afr., Am.
P. obtusifolius; Mertens & Koch, Deutschl. Fl. I, 855 (1823) ...	...	...	W.A. S.A. T. V. N.S.W. Q. —	B.fl. VII, 172 M.fr. VIII, 216. E., As., Afr., Am., N.Z.
P. acutifolius; Link in Roem. & Schult. syst. veg. III, 513 (1818)	...	...	— S.A. — V. N.S.W. — —	B.fl. VII, 173 M.fr. VIII, 216. E., As.
P. pectinatus; Linné, spec. plant. 127 (1753) ...	...	...	— S.A. T. V. — — —	B.fl. VII, 173 M.fr. VIII, 217. E., As., Afr., Am., N.Z.

APONOGETON; Linné, fil. suppl. plant. 32 (1781).

A. monostachyus; Linné fil., suppl. 214 (1781) ...	...	...	— — — — N.S.W. Q. N.A.	B.fl. VII, 188 M.fr. VIII, 215. As.
A. elongatus; F. v. M. in Benth. Fl. Austr. VII, 188 (1878) ...	...	...	— — — — N.S.W. Q. N.A.	B.fl. VII, 188 M.fr. VIII, 216.

TRIGLOCHIN; Rivinus in Rupp. fl. Jenens. 54 et 366 (1718) from Dalechamps (1586). (Cycnogeton, Maundia.)

T. calcitrapa; Hooker, icon. pl. t. 731 (1848) ...	...	...	W.A. S.A. — — N.S.W. Q. —	M.fr. VI, 82.
T. centrocarpa; Hooker, icon. plant. t. 728 (1848) ...	...	...	W.A. S.A. T. V. N.S.W. Q. —	B.fl. VII, 167 M.fr. VI, 82.
T. striata; Ruiz & Pavon, Fl. Per. et Chil. III, 72 (1802) ...	...	...	W.A. S.A. T. V. N.S.W. Q. —	B.fl. VII, 166 M.fr. VI, 83. Am., N.Z.
T. mucronata; R. Brown, prodr. 343 (1810) ...	...	...	W.A. S.A. — V. — —	B.fl. VII, 168 M.fr. VI, 81.
T. procera; R. Brown, prodr. 343 (1810) ...	...	...	W.A. S.A. T. V. N.S.W. Q. N.A.	B.fl. VII, 168 M.fr. VI, 83.
T. Maundii; F. v. M., fragm. VI, 83 (1867) ...	...	...	— — — — N.S.W. Q. —	B.fl. VII, 169 M.fr. I, 23.

ALISMACEAE.

VENTENAT, Tabl. du règn. végét. II, 157 (1799).—Genera 3, Species 6.

ALISMA; Rivinus in Ruppia, fl. Jen. 54 (1718), from Dioscorides and Plinius. (Caldesia; Echinodorus partly.)

<i>A. Plantago</i> ; Linné, spec. plant. 842 (1753) ...	...	...	...	...	...	—	—	—	V. N.S.W.	—	—	B.f.VII,185	M.fr.VIII,215.	E., As., Afr., Am.
<i>A. acanthocarpum</i> ; F. v. M., fragm. I, 23 (1858) ...	...	...	...	...	...	—	—	—	—	—	—	Q. N.A.	B.f.VII,185	M.fr.VIII,214.
<i>A. oligococcum</i> ; F. v. M., fragm. I, 23 (1858) ...	...	...	...	...	...	—	—	—	—	—	—	Q. N.A.	B.f.VII,185	M.fr.VI,169; VIII,214.
<i>A. parnassifolium</i> ; Bassi in Comment. Scienc. inst. Bonon. VI, 13, t. 1 (1768) ...	...	...	...	...	...	—	—	—	—	—	—	Q. —	B.f.VII,186	M.fr.VIII,214.

DAMASONIUM ; Tournefort, inst. 256 (1700). (Actinocarpus.)

D. Australe; Salisbury in Transact. hort. Soc. Lond. I, 268 (1812) ... W.A. S.A. — V. N.S.W. Q. — B.f.VII,186 M.fr.VIII,215.

TENAGOCHARIS; Hochstetter in Regensb. bot. Zeit. June (1841). (Butomopaisa.)

T. Cordofana; Hochstetter in Regenab. bot. Zeit. June (1841)	...	...	...	—	—	—	—	—	Q. N.A. B.fl.VII,187 M.fr.X,103.	As., Afr.
--	-----	-----	-----	---	---	---	---	---	----------------------------------	-----------

PONTEDERIACEAE.

HUMBOLDT, Bonpland & Kunth, nov. gen. et spec. plant. I, 265 (1815).—Genus 1, Species 1.

MONOCHORIA; Preal, reliq. Haenk. I, 127 (1827). (Limnostachya.)

M. cyanea; F. v. M., fragm. VIII, 44 (1872)... — — — — — Q. N.A. B.fl. VII, 74 M.fr. VIII, 44.

PHILHYDREAE.

R. BROWN in Flinders' Voy. II, 57 (1814).—Genera 3, Species 4.

PHILHYDRUM; Banks in Gaertner, de fruct. I, 62, t. 16 (1788). . (Philydrum.)

P. lanuginosum; Banks in Gaertn. de fruct. I. 62, t. 16 (1788) ... — — — V. N.S.W. Q. N.A. B.fl.VII,73 M.fr.V,203. Aa.

PRITZELIA; F. v. M., Papuan plants 13 (1875). (Hetaeria, Philydrella.)

P. pygmaea; F. v. M., Papuan plants 13 (1875) W.A. — — — — — B.fl.VII,74 M.fr.V,203.

HELMHOLTZIA ; F. v. M., fragm. V, 202 (1866).

H. acorifolia; F. v. M., fragm. V, 203 (1866) — — — — N.S.W. Q. — B.fl.VII,75 M.fr.V,203.

H. glaberrima; Caruel in A. & C. de Candolle, monogr. phanerog. III, 6 (1881) — — — — — Q. — —

COMMELINEAE.

R. BROWN, prodr. fl. Nov. Holl. 268 (1810).—Genera 6, Species 19.

ZYGOMENES; Salisbury in Transact. hort. Soc. I, 271 (1812). (Cyanotis).

Z. axillaria; Salisbury in Transact. hort. Soc. I, 271 (1812) Q. N.A. B.fl.VII,82 M.fr.VIII,62. A8.

COMMELINA; Plumier, nov. pl. Amer. gen. 48, t. 38 (1703). (Commelynia.)

C. ensifolia; R. Brown, prodr. 269 (1810) W.A. S.A. — — N.S.W. Q. N.A. B.f.VII,83 M.fr.VIII,60; IX,191. As., P.

C. cyanea; R. Brown, prodr. 269 (1810) — — — — N.S.W. Q. N.A. B.fl.VII,84 M.fr.VIII,59; IX,191. P.

C. lanceolata; R. Brown, prodr. 269 (1810) — — — — — Q. N.A. B.fl.VII,84 M.fr.VIII,59.

C. agrostophylla; F. v. M. in J. Hooker, Fl. Tasman. p. 47 (1860)	...	...	—	—	—	—	—	Q. N.A.	—	M.fr.VIII,59.
---	-----	-----	---	---	---	---	---	---------	---	---------------

ANEILEMA; R. Brown, prodr. fl. Nov. Holl. 270 (1810).

A. acuminatum; R. Brown, prodr. 270 (1810) — — — — N.S.W. Q. — B.fl.VII,85 M.fr.VIII,61.

A. biflorum; R. Brown, prodr. 270 (1810)	...	..	...	...	...	...	—	—	—	—	N.S. W.	Q.	—	B.fl.VII,86
--	-----	----	-----	-----	-----	-----	---	---	---	---	---------	----	---	-------------

A. sclerocarpum; F. v. M., fragm. VIII, 61 (1873) ... — — — — — Q. — B.f. VII, 86 M.fr. VIII, 61.

A. siliculosum; R. Brown, prodr. 270 (1810)...	...	...	...	...	...	—	—	—	—	Q. N.A. B.fl. VII, 87	M.fr. VIII, 61; IX, 191.
A. gramineum; R. Brown, prodr. 270 (1810)...	...	...	...	...	...	—	—	—	N.S.W.	Q. N.A. B.fl. VII, 87	M.fr. VIII, 62.
A. giganteum; R. Brown, prodr. 271 (1810)...	...	...	...	...	...	—	—	—	—	Q. N.A. B.fl. VII, 88	M.fr. VII, 62. As., Afr.

FLORISCOPA; Loureiro, Fl. Cochinch. I, 192 (1790). (Floscopa, Dithyrocarpus.)

F. scandens; Loureiro, fl. Cochinch. I, 193 (1790)...	...	...	...	...	...	—	—	—	—	Q. —	B.fl. VII, 89	M.fr. VIII, 63. As., P.
---	-----	-----	-----	-----	-----	---	---	---	---	------	---------------	-------------------------

POLLIA; Thunberg, nov. gen. plant. I, 11 (1781).

P. cyanococca; F. v. M., fragm. V, 40 (1865)...	...	...	...	...	...	—	—	—	N.S.W.	Q. —	B.fl. VII, 90	M.fr. VIII, 63.
---	-----	-----	-----	-----	-----	---	---	---	--------	------	---------------	-----------------

CARTONEMA; R. Brown, prodr. fl. Nov. Holl. 271 (1810).

C. philhydroides; F. v. M., fragm. I, 62 (1858)...	...	...	...	...	...	W.A.	—	—	—	—	B.fl. VII, 91	M.fr. VII, 64.
C. spicatum; R. Brown, prodr. 271 (1810)...	...	...	...	...	...	—	—	—	—	—	N.A. B.fl. VII, 91	M.fr. I, 62; VIII, 64.
C. parviflorum; Hasskarl in Flora Regensb. 365 (1869)...	...	...	...	...	...	—	—	—	—	—	Q. N.A. B.fl. VII, 91	
C. trigonospermum; Clarke in De Candolle, mon. phaner. III, 24 (1881)...	...	...	...	...	...	—	—	—	—	—	N.A. —	M.fr. XI, 139.
C. brachyantherum; Benth, Fl. Austr. VII, 92 (1878)...	...	...	...	...	...	—	—	—	—	Q. —	B.fl. VII, 92	
C. tenue; Benth in De Candolle, monogr. phaner. III, 264 (1881)...	...	...	...	...	...	—	—	—	—	—	N.A. —	M.fr. XI, 139.

XYRIDEAE.

SALISBURY in Transact. of the Hort. Soc. I, 326 (1812).—Genus I, Species 9.

XYRIS; Gronovius in Linné, genera plantar. 11 (1737).

X. elongata; Rudge in Trans. Linn. Soc. X, 289, t. 15 (1809)	...	...	...	...	...	—	—	—	—	N.S.W.	Q. N.A.	B.fl. VII, 77	M.fr. VIII, 205.	
X. pauciflora; Willdenow, phytogr. 2, t. 1 (1794)	...	...	...	...	...	—	—	—	—	—	Q. N.A.	B.fl. VII, 77	M.fr. VIII, 205.	As.
X. lacera; R. Brown, prodr. 257 (1810)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl. VII, 78	M.fr. VIII, 204.	
X. flexifolia; R. Brown, prodr. 256 (1810)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl. VII, 78	M.fr. VIII, 204.	
X. gracilis; R. Brown, prodr. 256 (1810)	...	...	...	...	...	—	S.A. T.	V.	N.S.W.	Q.	—	B.fl. VII, 79	M.fr. VIII, 204.	
X. operculata; Labillardière, Nov. Holl. pl. spec. I, 14, t. 10 (1804)	...	...	...	...	...	—	S.A. T.	V.	N.S.W.	Q.	—	B.fl. VII, 79	M.fr. VIII, 204.	
X. lanata; R. Brown, prodr. 257 (1810)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl. VII, 80	M.fr. VIII, 204.	
X. laxiflora; F. v. M., fragm. VIII, 203 (1874)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl. VII, 80	M.fr. VIII, 203.	
X. gracillima; F. v. M., fragm. VIII, 203 (1874)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl. VII, 80	M.fr. VIII, 203.	

FLAGELLARIACEAE.

ENDLICHER, gener. plantar. 131 (1836).—Genus 1, Species 1.

FLAGELLARIA; Linné, Amoenit. academ. 396 (1747).

E. Indica; Linné, spec. plant. 333 (1753)...	...	...	...	...	...	—	—	—	N.S.W.	Q. N.A. B.fl. VII, 10	...	...	As., Afr., P.
--	-----	-----	-----	-----	-----	---	---	---	--------	-----------------------	-----	-----	---------------

JUNCEAE.

R. BROWN, prodr. fl. Nov. Holl. 257 (1810).—Genera 2, Species 16.

LUZULA; De Candolle, Fl. franc. III, 158 (1805).

L. campestris; De Candolle, Fl. franc. III, 158 (1805)...	...	...	...	...	...	W.A. S.A. T.	V.	N.S.W.	Q. —	B.fl. VII, 123	—	...	E., As., Afr., Am., P., N.Z.
L. longiflora; Benth, Fl. Austr. VII, 123 (1878)...	...	...	...	...	...	—	—	—	N.S.W.	—	B.fl. VII, 123		

JUNCUS; Tournefort, inst. 246, t. 127 (1700), from Camerarius, J. & C. Bauhin, Ray & Morison.

J. gracilis; R. Brown, prodr. 259 (1810)	...	...	...	...	...	W.A.	—	—	—	—	—	B.fl. VII, 125		
J. planifolius; R. Brown, prodr. 259 (1810)	...	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl. VII, 125	... N.Z.
J. caespititius; E. Meyer in Lehm. pl. Preiss. II, 47 (1846)	...	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl. VII, 126	
J. falcatus; E. Meyer, synops. luzular. 34 (1823)	...	...	...	...	...	—	—	T.	V.	N.S.W.	—	—	B.fl. VII, 126	... Am.

<i>J. bufonius</i> ; Linné, spec. plant. 328 (1753)	W.A. S.A. T. V. N.S.W. Q. —	B.fl.VII,127	...	...	E., As., Afr., Am., P., N.Z.
<i>J. homalocaulis</i> ; F. v. M., first gen. Rep. 19 (1853)	W.A. S.A. — V. N.S.W. — —	B.fl.VII,128			
<i>J. Brownii</i> ; F. v. M., first gen. Rep. 19 (1853)	— — T. V. N.S.W. — —	B.fl.VII,128	M.fr.IX,78.		
<i>J. communis</i> ; E. Meyer, synops. Juncor. 12 (1822)	W.A. S.A. T. V. N.S.W. Q. —	B.fl.VII,128	...	...	E., As., Afr., Am., P., N.Z.
<i>J. vaginatus</i> ; R. Brown, prodr. 258 (1810)	— — — — N.S.W. Q. —	B.fl.VII,129			
<i>J. pauciflorus</i> ; R. Brown, prodr. 259 (1810)	— S.A. T. V. N.S.W. Q. —	B.fl.VII,129			
<i>J. pallidus</i> ; R. Brown, prodr. 258 (1810)	W.A. S.A. T. V. N.S.W. Q. —	B.fl.VII,130	...	...	N.Z.
<i>J. maritimus</i> ; Lamarek, Encycl. méth. III, 264 (1789)	W.A. S.A. T. V. N.S.W. Q. —	B.fl.VII,130	...	...	Aa., Afr., N.Z.
<i>J. prismatocarpus</i> ; R. Brown, prodr. 259 (1810)	W.A. S.A. T. V. N.S.W. Q. —	B.fl.VII,131			
<i>J. stipulatus</i> ; Meyen & Nees in Act. Acad. Caes. L.C. XVIII, suppl. 127 (1843) — —	T. V. N.S.W. — —	B.fl.VII,132	...	...	N.Z.

ERIOCAULEAE.

HUMBOLDT, Bonpland & Kunth, Nov. gen. et spec. plant. I, 251 (1815).—Genus 1, Species 18.

ERIOCAULON; Linné, gen. plant. ed. sec. 35 (1742). (Electrosperma)

<i>E. setaceum</i> ; Linné, spec. plant 87 (1753)	— — — — —	Q. N.A. B.fl.VII,191	...	...	As.
<i>E. Australe</i> ; R. Brown, prodr. 254 (1810)	— — — — N.S.W. Q. —	B.fl.VII,192			
<i>E. quinquangulare</i> ; Linné, spec. plant. 87 (1753)	— — — — —	— N.A. B.fl.VII,192	...	...	As.
<i>E. Smithii</i> ; R. Brown, prodr. 254 (1810)	— — — — V. N.S.W. Q. —	B.fl.VII,192			
<i>E. nanum</i> ; R. Brown, prodr. 254 (1810)	— — — — —	Q. — B.fl.VII,193			
<i>E. cinereum</i> ; R. Brown, prodr. 254 (1810)	— — — — —	Q. N.A. B.fl.VII,193	M.fr.I,95.	...	As.
<i>E. pusillum</i> ; R. Brown, prodr. 254 (1810)	— — — — —	Q. — B.fl.VII,194			
<i>E. pallidum</i> ; R. Brown, prodr. 254 (1810)	— — — — —	Q. — B.fl.VII,194			
<i>E. nigricans</i> ; R. Brown, prodr. 254 (1810)	— — — — —	Q. N.A. B.fl.VII,194			
<i>E. electrospermum</i> ; F. v. M., Cens. 123 (1832)	— — — — V. N.S.W. — —	B.fl.VII,195			
<i>E. lividum</i> ; F. v. M., fragm. I, 92 (1858)	— — — — —	— N.A. B.fl.VII,195	M.fr.I,92.		
<i>E. concretum</i> ; F. v. M., fragm. I, 92 (1858)	— — — — —	— N.A. B.fl.VII,195	M.fr.I,92.		
<i>E. Schultzei</i> ; Benth. Fl. Austr. VII, 195 (1878)	— — — — —	— N.A. B.fl.VII,195			
<i>E. tortuosum</i> ; F. v. M., fragm. I, 94 (1858)	— — — — —	— N.A. B.fl.VII,196	M.fr.I,91.		
<i>E. monoscapum</i> ; F. v. M., fragm. I, 94 (1858)	— — — — —	— N.A. B.fl.VII,196	M.fr.I,94.		
<i>E. spectabile</i> ; F. v. M., fragm. I, 95 (1858)	— — — — —	Q. N.A. B.fl.VII,196	M.fr.I,95.		
<i>E. scariosum</i> ; R. Brown, prodr. 255 (1810)	— — — — —	Q. N.A. B.fl.VII,197			
<i>E. depressum</i> ; R. Brown, prodr. 255 (1810)	— — — — —	Q. N.A. B.fl.VII,197	M.fr.I,92.		

RESTIACEAE.

R. BROWN, prodr. flor. Nov. Holl. 243 (1810).—Genera 14, Species 93.

TRITHURIA; J. Hooker, Fl. Tasman. II, 79, t. 138 (1860). (Juncella, 1854.)

<i>T. submersa</i> ; J. Hooker, Fl. Tasman. II, 79, t. 138 (1860)	W.A. S.A. T. V. N.S.W. — —	B.fl.VII,199	M.fr.I,96; VIII,237.		
--	----------------------------	--------------	----------------------	--	--

APHELIA; R. Brown, prodr. fl. Nov. Holl. 251 (1810). (Brizula.)

<i>A. cyperoides</i> ; R. Brown, prodr. 252 (1810)	W.A. — — — — —	B.fl.VII,200	M.fr.VIII,236.		
<i>A. nutans</i> ; J. Hooker in Benth. Fl. Austr. VII, 200 (1878)	W.A. — — — — —	B.fl.VII,200			
<i>A. gracilis</i> ; Sonder in Schlecht. Linnaea XXVIII, 227 (1855)	— S.A. T. V. N.S.W. — —	B.fl.VII,201	M.fr.VIII,237.		
<i>A. Drummondii</i> ; Benth. Fl. Austr. VII, 201 (1878)	W.A. — — — — —	B.fl.VII,201			
<i>A. Pumilio</i> ; F. v. M., first gen. Report 20 (1853)	— S.A. T. V. N.S.W. — —	B.fl.VII,201	M.fr.VIII,237.		
<i>A. Brizula</i> ; F. v. M., fragm. V, 203 (1866)	W.A. — — — — —	B.fl.VII,202	M.fr.VIII,237.		

CENTROLEPIS; Labillardière, Nov. Holl. pl. specim. I, 7 (1804). (Desvauxia, Devauxia, Alepyrum.)

<i>C. humillima</i> ; F. v. M. in Benth. Fl. Austr. VII, 203 (1878)	W.A. — — — — —	B.fl.VII,203			
<i>C. polygyna</i> ; Hieronymus in Abh. naturf. Ges. Halle XII, 96 (1873)	W.A. S.A. T. V. N.S.W. Q. —	B.fl.VII,203	M.fr.VIII,237.		

R. deformis ; R. Brown, prodr. 245 (1810)	W.A.	—	—	—	—	—	B.fl. VII, 225	M.fr. VIII, 65.
R. crispatus ; R. Brown, prodr. 246 (1810)	W.A.	—	—	—	—	—	B.fl. VII, 225	
R. nitens ; Nees in Lehm. pl. Preiss. II, 59 (1846)	W.A.	—	—	—	—	—	B.fl. VII, 226	M.fr. VIII, 70.
R. gracilior ; F. v. M. in Benth. Fl. Austr. VII, 226 (1878)	W.A.	—	—	—	—	—	B.fl. VII, 226	
R. chauuocoleus ; F. v. M., fragm. VIII, 64 (1873)	W.A.	—	—	—	—	—	B.fl. VII, 226	M.fr.
R. Australis ; R. Brown, prodr. 245 (1810)	—	—	T. V.	N.S.W.	—	—	B.fl. VII, 227	M.fr. VIII, 69.
R. gracilis ; R. Brown, prodr. 245 (1810)	—	—	T. V.	N.S.W.	Q.	—	B.fl. VII, 227	M.fr. VIII, 69.
R. complanatus ; R. Brown, prodr. 245 (1810)	—	—	S.A. T. V.	N.S.W.	Q.	—	B.fl. VII, 228	M.fr. VIII, 67.
R. tetraphyllus ; Labillardière, Nov. Holl. pl. spec. II, 77 (1806)	—	—	S.A. T. V.	N.S.W.	Q.	—	B.fl. VII, 228	M.fr. VIII, 66.
R. laxus ; R. Brown, prodr. 245 (1810)	W.A.	—	—	—	—	—	B.fl. VII, 228	M.fr. VIII, 71.
R. ornatus ; Steudel, syn. pl. glum. II, 256 (1855)	W.A.	—	—	—	—	—	B.fl. VII, 229	M.fr. VIII, 70.
R. leptocarpoides ; Benth. Fl. Austr. VII, 229 (1878)	W.A.	—	—	—	—	—	B.fl. VII, 229	M.fr. VIII, 65.
R. amblycoleus ; F. v. M., fragm. VIII, 65 (1873)	W.A.	—	—	—	—	—	B.fl. VII, 230	M.fr. VIII, 65.
R. tremulus ; R. Brown, prodr. 230 (1810)	W.A.	—	—	—	—	—	B.fl. VII, 230	M.fr. VIII, 70.

LOXOCARYA ; R. Brown, prodr. fl. Nov. Holl. 249 (1810). (Desmocladus.)

L. densa ; Benth. Fl. Austr. VII, 241 (1878)	W.A.	—	—	—	—	—	B.fl. VII, 241	
L. Benthami ; F. v. M., Cens. 124 (1882) from Masters (1878)	W.A.	—	—	—	—	—	B.fl. VII, 242	
L. pubescens ; Benth. Fl. Austr. VII, 242 (1878)	W.A.	—	—	—	—	—	B.fl. VII, 242	M.fr. VIII, 100.
L. fasciculata ; Benth. Fl. Austr. VII, 242 (1878)	W.A.	—	—	—	—	—	B.fl. VII, 242	M.fr. VIII, 98.
L. flexuosa ; Benth. Fl. Austr. VII, 243 (1878)	W.A.	—	—	—	—	—	B.fl. VII, 243	M.fr. VIII, 100.
L. cinerea ; R. Brown, prodr. 249 (1810)	W.A.	—	—	—	—	—	B.fl. VII, 243	

CALOSTROPHUS ; Labillardière, Nov. Holl. pl. spec. II, 78 (1806). (Calorophus, Hypolaena.)

C. elongatus ; Labillardière, Nov. Holl. pl. spec. II, 78 (1806)	—	—	T.	—	—	—	B.fl. VII, 238	M.fr. VIII, 86; IX, 94; X, 120.
C. lateriflorus ; F. v. M., fragm. VIII, 87 (1873)	—	—	S.A. T. V.	N.S.W.	Q.	—	B.fl. VII, 238	M.fr. VIII, 87.
C. gracillimus ; F. v. M., fragm. VIII, 88 (1873)	W.A.	—	—	—	—	—	B.fl. VII, 239	M.fr. VIII, 88.
C. fastigiatus ; F. v. M. in pap. Roy. Soc. Tasm. 117 (1878)	—	—	S.A. T. V.	N.S.W.	Q.	—	B.fl. VII, 239	M.fr. VIII, 84.
C. exsulcus ; F. v. M., Cens. 124 (1882) from R. Brown (1810)	W.A.	—	—	—	—	—	B.fl. VII, 240	M.fr. VIII, 86.

LEPTOCARPUS ; R. Brown, prodr. 250 (1810). (Schoenodum partly.)

L. scariosus ; R. Brown, prodr. 250 (1810)	W.A.	—	—	—	—	—	B.fl. VII, 232	M.fr. VIII, 94.
L. tenax ; R. Brown, prodr. 250 (1810)	W.A. S.A. T. V.	N.S.W.	—	—	—	—	B.fl. VII, 232	M.fr. VIII, 96.
L. Brownii ; J. Hooker, Fl. Tasman. II, 73, t. 136 (1860)	—	—	S.A. T. V.	N.S.W.	—	—	B.fl. VII, 233	M.fr. VIII, 91.
L. caeus ; Nees in Ann. of nat. hist. VI, 50 (1843)	W.A.	—	—	—	—	—	B.fl. VII, 234	M.fr. VIII, 95.
L. coangustatus ; Nees in Lehm. pl. Preiss. II, 65 (1846)	W.A.	—	—	—	—	—	B.fl. VII, 234	M.fr. VIII, 94.
L. aristatus ; R. Brown, prodr. 250 (1810)	W.A.	—	—	—	—	—	B.fl. VII, 235	M.fr. VIII, 90.
L. erianthus ; Benth. Fl. Austr. VII, 235 (1878)	W.A.	—	—	—	—	—	B.fl. VII, 235	
L. ramosus ; R. Brown, prodr. 250 (1810)	—	—	—	—	—	Q.	B.fl. VII, 236	M.fr. VIII, 92.
L. elatior ; R. Brown, prodr. 250 (1810)	—	—	—	—	—	Q. N.A.	B.fl. VII, 236	
L. spathaceus ; R. Brown, prodr. 250 (1810)	—	—	—	—	—	N.A.	B.fl. VII, 236	M.fr. VIII, 93.
L. Schultzii ; Benth. Fl. Austr. VII, 237 (1878)	—	—	—	—	—	N.A.	B.fl. VII, 237	M.fr. VIII, 93.

LEPIDOBOLUS ; Nees in Lehm. pl. Preiss. II, 66 (1846).

L. drapetocoleus ; F. v. M., fragm. VIII, 84 (1873)	—	—	S.A.	—	V.	—	B.fl. VII, 244	M.fr. VIII, 84.
L. Preissianus ; Nees in Lehm. pl. Preiss. II, 66 (1846)	W.A.	—	—	—	—	—	B.fl. VII, 245	M.fr. VIII, 83.
L. chaetocephalus ; F. v. M., fragm. VIII, 84 (1873)	W.A.	—	—	—	—	—	B.fl. VII, 245	

CHAETANTHUS ; R. Brown, prodr. 251 (1810). (Prionosepalum.)

C. leptocarpoides ; R. Brown, prodr. 251 (1810)	W.A.	—	—	—	—	—	B.fl. VII, 246	M.fr. VIII, 97.
--	------	---	---	---	---	---	----------------	-----------------

ONYCHOSEPALUM ; Steudel, syn. pl. glum. II, 249 (1855).

O. laxiflorum ; Steudel, syn. pl. glum. II, 249 (1855)	W.A.	—	—	—	—	—	B.fl. VII, 246	M.fr. IX, 51.
---	------	---	---	---	---	---	----------------	---------------

ACALYCEAE HYPOGYNAE.

F. v. M. in Woolls's plants of the neighb. of Sydney 48 (1880).

CYPERACEAE.

HALLER, enum. stirp. Helvet. I, 234 (1742).—Genera 29, Species 380.

KYLINGIA; Rottboell, descr. et icon. rar. et nov. plant. 13, t. 4 (1773). (Killiogia.)

K. intermedia; R. Brown, prodr. 219 (1810) ...	...	...	...	...	...	S.A.	—	V.	N.S.W.	Q.	—	B.f. VII, 251	M.fr. VIII, 27	...	As., P.
K. monocephala; Rottboell, descr. et icon. pl. 13, t. 4 (1773) ...	...	...	...	...	...	S.A.	—	—	N.S.W.	Q.	—	B.f. VII, 251	M.fr. VIII, 271; IX, 54.	...	As., Afr., Am., P.
K. cylindrica; Nees in Wight's Contr. to Bot. of Ind. 91 (1834) ...	...	...	...	...	...	—	—	—	N.S.W.	Q.	—	B.f. VII, 252	M.fr. VIII, 271	...	As.
K. triceps; Rottboell, descr. et icon. pl. 14, t. 4 (1773) ...	...	...	...	...	...	—	—	—	—	Q.	—	B.f. VII, 252	...	...	As., Afr.

CYPERUS; Tournefort, inst. 527, t. 299 (1700) from Hippocrates, Theophrastos & Plinius. (Pycrus, Mariscus, Anosporum, Sorostachys.)

C. pumilus; Linné, amoen. acad. IV, 302 (1759) ...	...	...	...	...	...	—	—	—	—	Q.	—	B.f. VII, 258	M.fr. VIII, 267	...	As., Afr.
C. Eragrostis; Vahl, enum. plant. II, 322 (1806) ...	...	...	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.f. VII, 258	M.fr. VIII, 260; IX, 52.	As., Afr., P.
C. flavescens; Linné, spec. plant. 46 (1753) ...	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.f. VII, 259	...	E., As., Afr.
C. globosus; Allioni, actuar. ad fl. Pedem. 49 (1789) ...	...	...	...	...	...	—	—	—	V.	N.S.W.	Q.	—	B.f. VII, 260	M.fr. VIII, 260; IX, 52.	E., As., Afr.
C. unioloides; R. Brown, prodr. 216 (1810) ...	...	...	...	...	...	—	—	—	V.	N.S.W.	Q.	—	B.f. VII, 260	M.fr. VIII, 259; IX, 52.	As., Afr., Am.
C. polystachyus; Rottboell, descr. et icon. pl. 39, t. 11 (1773) ...	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	N.A.	B.f. VII, 261	M.fr. VIII, 265, 270; XI, 53; XII, 25	E., As.
C. Hochstetteri; Nees in Regensb. Flora 755 (1845) ...	...	...	...	...	...	—	—	—	—	—	—	N.A.	B.f. VII, 261	M.fr. VIII, 265; IX, 53.	As., Afr., Am.
C. lucidulus; Klein in Link, Jahrb. III, 86 (1820) ...	...	...	...	...	...	—	—	—	—	—	Q.	—	...	...	As.
C. pygmaeus; Rottboell, descr. et icon. pl. 20, t. 14 (1773) ...	...	...	...	...	...	—	—	—	V.	N.S.W.	Q.	N.A.	B.f. VII, 262	M.fr. VIII, 268; IX, 53.	E., As., Afr., Am.
C. cephalotes; Vahl, enum. II, 311 (1806) ...	...	...	...	...	...	—	—	—	—	—	Q.	—	B.f. VII, 263	M.fr. VIII, 272; IX, 54.	As.
C. laevigatus; Linné, mantissa alt. 179 (1771) ...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.f. VII, 263	M.fr. VIII, 266; IX, 53.	E., As., Afr., Am.
C. platystylis; R. Brown, prodr. 241 (1810) ...	...	...	...	...	...	—	—	—	—	N.S.W.	—	—	B.f. VII, 264	...	...
C. alopecuroides; Rottboell, descr. et icon. pl. 38, t. 8 (1773) ...	...	...	...	...	...	—	—	—	—	—	Q.	—	B.f. VII, 264	M.fr. VIII, 263; IX, 52.	As., Afr.
C. leucocephalus; Retzius, observ. V, 11 (1788) ...	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.f. VII, 265	M.fr. VIII, 271; XII, 25.	As.
C. tenellus; Linné, fil. suppl. 103 (1781) ...	...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.f. VII, 265	M.fr. VIII, 261; IX, 53; XII, 25.	Afr., N.Z.
C. gracilis; R. Brown, prodr. 213 (1810) ...	...	...	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.f. VII, 265	M.fr. VIII, 264; IX, 53; XII, 25.	P.
C. enervis; R. Brown, prodr. 213 (1810) ...	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.f. VII, 266	...	...
C. debilis; R. Brown, prodr. 213 (1810) ...	...	...	...	...	...	—	—	—	—	N.S.W.	—	—	B.f. VII, 266	...	...
C. castaneus; Willdenow, spec. plant. I, 278 (1797) ...	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.f. VII, 267	...	As.
C. cuspidatus; Humboldt, Bonpland & Kunth, nov. gen. et sp. pl. I, 204 (1815) ...	...	...	...	...	...	—	—	—	—	—	Q.	—	B.f. VII, 267	...	As., Afr., Am.
C. squarrosus; Linné, amoen. acad. IV, 303 (1759) ...	...	...	...	...	...	—	—	—	—	—	—	N.A.	—	...	...
C. ixiocarpus; F. v. M. in proc. R. S. Vict. XXIII, 53 (1885) ...	...	...	...	...	...	W.A.	S.A.	—	—	—	Q.	N.A.	B.f. VII, 268	M.fr. VIII, 262; IX, 53; XII, 25.	As., Afr., Am.
C. difformis; Lioné, amoen. acad. IV, 302 (1759) ...	...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.f. VII, 268	M.fr. VIII, 262; IX, 52.	E., As., Afr.
C. tetraphyllus; R. Brown, prodr. 214 (1810) ...	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.f. VII, 269	M.fr. VIII, 264.	...
C. trinervis; R. Brown, prodr. 213 (1810) ...	...	...	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.f. VII, 269	M.fr. VIII, 267; IX, 53.	...
C. Haspan; Linné, spec. plant. 45 (1753) ...	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	N.A.	B.f. VII, 270	M.fr. VIII, 260; IX, 52.	As., Afr., Am., P.
C. concinnus; R. Brown, prodr. 214 (1810) ...	...	...	...	...	...	—	—	—	V.	N.S.W.	Q.	—	B.f. VII, 270	M.fr. VIII, 261.	...
C. filipes; Benth, Fl. Austr. VII, 271 (1878) ...	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.f. VII, 271	...	...
C. pedunculatus; F. v. M., fragm. VIII, 266 (1874) ...	...	...	...	...	...	—	—	—	—	—	Q.	—	B.f. VII, 272	M.fr.	...
C. vaginatus; R. Brown, prodr. 213 (1810) ...	...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.f. VII, 272	M.fr. VIII, 261; IX, 53.	...
C. Holoschoenus; R. Brown, prodr. 215 (1810) ...	...	...	...	...	...	—	S.A.	—	—	—	Q.	N.A.	B.f. VII, 273	M.fr. I, 200; VIII, 262.	...
C. dactyloides; Benth, Fl. Austr. VII, 273 (1878) ...	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	N.A.	B.f. VII, 273	...	...
C. Gilesii; Benth, Fl. Austr. VII, 274 (1878) ...	...	...	...	...	...	—	S.A.	—	—	N.S.W.	Q.	—	B.f. VII, 274	...	...
C. fulvus; R. Brown, prodr. 215 (1810) ...	...	...	...	...	...	W.A.	S.A.	—	—	N.S.W.	Q.	—	B.f. VII, 274	M.fr. VIII, 268.	...
C. carinatus; R. Brown, prodr. 216 (1810) ...	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	N.A.	B.f. VII, 274	...	...
C. alterniflorus; R. Brown, prodr. 216 (1810) ...	...	...	...	...	...	W.A.	S.A.	—	—	—	Q.	—	B.f. VII, 275	...	...
C. pilosus; Vahl, enum. II, 354 (1806) ...	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.f. VII, 275	M.fr. VIII, 260; IX, 52.	As.

<i>C. procerus</i> ; Rottboell, descr. et. icon. 29, t. 5 (1773)	— — — — —	N.S.W.	Q. —	B.fl.VII,276				
<i>C. Iria</i> ; Linné, spec. plant. 45 (1753)	— S.A. — — —	N.S.W.	Q. N.A.	B.fl.VII,276	M.fr.VIII,266; IX,53; XII,25.	As., Afr.		
<i>C. eleusinoides</i> ; Wallich in Kunth. enum. II, 39 (1837)	— — — — —	—	Q. N.A.	B.fl.VII,277	M.fr.VIII,264; IX,53.	As., Afr.		
<i>C. distans</i> ; Linné, fil. suppl. 103 (1781)	— — — — —	—	Q. —	B.fl.VII,277	M.fr.VIII,266; IX,53.	As., Afr., Am., P.		
<i>C. Malaccensis</i> ; Lamarck, ill. des gen. I, 146 (1791)	— — — — —	—	—	N.A.	B.fl.VII,278	M.fr.VIII,260; IX,52.	As.	
<i>C. articulatus</i> ; Linné, spec. plant. 44 (1753)	— — — — —	—	—	N.A.	B.fl.VII,278	...	As., Afr., Am.	
<i>C. corymbosus</i> ; Rottboell, descr. et icon. pl. 42, t. 7 (1773)	— S.A. — — —	—	Q. N.A.	B.fl.VII,279			[Am., P.	
<i>C. rotundus</i> ; Linné, spec. plant. 45 (1755)	W.A. S.A. — V.	N.S.W.	Q. N.A.	B.fl.VII,279	M.fr.VIII,269; IX,53; XII,25.	As., Afr.,		
<i>C. stenostachyus</i> ; Benth. Fl. Austr. VII, 280 (1878)	W.A. — — —	—	—	B.fl.VII,280				
<i>C. Lhotzkyanus</i> ; Boeckeler in Engler's bot. Jahrb. 498 (1884)	— — — — —	N.S.W.	—	—				
<i>C. flavidus</i> ; Retzius, obs. bot. V, 13 (1788)	— — — — —	—	Q. —	—				
<i>C. tenuiflorus</i> ; Rottboell, descr. et icon. pl. 30, t. 14, f. I (1773)	— — — — —	—	Q. N.A.	—				
<i>C. congestus</i> ; Vahl, enum. II, 350 (1806)	W.A. — — —	N.S.W.	Q. —	B.fl.VII,280	M.fr.VIII,269; IX,53.	Afr.		
<i>C. subulatus</i> ; R. Brown, prodr. 217 (1810)	— S.A. — — —	N.S.W.	Q. —	B.fl.VII,281				
<i>C. sporobolus</i> ; R. Brown, prodr. 215 (1810)	— — — — —	—	Q. N.A.	B.fl.VII,281				
<i>C. angustatus</i> ; R. Brown, prodr. 214 (1810)	— — — — —	—	Q. N.A.	B.fl.VII,282	M.fr.IX,54.			
<i>C. inornatus</i> ; Boeckeler in Fl. Regensb. 86 (1875)	— — — — —	—	Q. —	B.fl.VII,282	M.fr.IX,54.			
<i>C. lucidus</i> ; R. Brown, prodr. 218 (1810)	— S.A. T. V.	N.S.W.	Q. N.A.	B.fl.VII,283	M.fr.VIII,270; IX,53.			
<i>C. pennatus</i> ; Lamarck, illustr. des genr. I, 144 (1799)	— — — — —	—	Q. N.A.	B.fl.VII,284	M.fr.VIII,263; IX,53.	As., P.		
<i>C. exaltatus</i> ; Retzius, observ. bot. V, 11 (1789)	— S.A. — V.	N.S.W.	Q. N.A.	B.fl.VII,285	M.fr.VIII,263; IX,53.	As., P.		
<i>C. haematodes</i> ; Endlicher, prodr. fl. Norf. 22 (1839)	— — — — —	N.S.W.	—	B.fl.VII,285	M.fr.IX,54.			
<i>C. digitatus</i> ; Roxburgh, fl. Ind. I, 209 (1820)	— — — — —	—	Q. —	B.fl.VII,286	M.fr.VIII,263	...	As., Afr., P.	
<i>C. ferax</i> ; Cl. Richard in act. soc. hist. nat. Par. I, 106 (1792)	— — — — —	—	Q. —	B.fl.VII,286	...	...	As., Afr., Am.	
<i>C. stoloniferus</i> ; Retzius, observ. IV, 10 (1786)	— — — — —	—	—	N.A.	—	...	As.	
<i>C. esculentus</i> ; Linné, sp. pl. 45 (1753)	— — — — —	—	Q. —	—	...	...	E., As., Afr.	
<i>C. Bowmani</i> ; F. v. M. in Benth. Fl. Austr. VII, 287 (1878)	— — — — —	N.S.W.	Q. —	B.fl.VII,287				
<i>C. trichostachys</i> ; Benth. Fl. Austr. VII, 287 (1878)	— — — — —	—	Q. —	B.fl.VII,287				
<i>C. leiocaulis</i> ; Benth. Fl. Austr. VII, 287 (1878)	— — — — —	N.S.W.	Q. —	B.fl.VII,287				
<i>C. scaber</i> ; Benth. Fl. Austr. VII, 288 (1878)	— — — — —	—	Q. N.A.	B.fl.VII,288				
<i>C. decompositus</i> ; F. v. M., fragm. VIII, 267 (1874)	— — — — —	—	Q. N.A.	B.fl.VII,288	M.fr.XII,25.			
<i>C. Armstrongii</i> ; Benth. Fl. Austr. VII, 289 (1878)	— — — — —	—	Q. N.A.	B.fl.VII,289				
<i>C. umbellatus</i> ; Benth. Fl. Hongk. 386 (1861)	— — — — —	—	Q. —	B.fl.VII,289	M.fr.VII,267; IX,53.	As., Afr., P.		
<i>C. conicus</i> ; Boeckeler in Schlecht. Liliaceae XXXVIII, 371 (1874)	W.A. — — —	—	Q. N.A.	B.fl.VII,290	M.fr.VIII,268; IX,53.			

HELEOCHARIS; R. Brown, prodr. 224 (1810). (Eleocharis, Eleochariton, Scirpidium.)

<i>H. sphacelata</i> ; R. Brown, prodr. 224 (1810)	— S.A. T. V.	N.S.W.	Q. N.A.	B.fl.VII,292	M.fr.VI,94; VIII,239.	N.Z., P.		
<i>H. spiralis</i> ; R. Brown, prodr. 224 (1810)	— — — — —	—	Q. N.A.	B.fl.VII,292	...	As., Afr., Am.		
<i>H. compacta</i> ; R. Brown, prodr. 224 (1810)	— — — — —	N.S.W.	Q. N.A.	B.fl.VII,293	M.fr.VIII,239	As., Afr., Am.		
<i>H. fistulosa</i> ; J. A. & J. H. Schultes, mantiss. II, 89 (1824)	— — — — —	—	Q. —	B.fl.VII,293	M.fr.VI,93; VIII,239.	As., Afr., Am.		
<i>H. tetraquetra</i> ; Nees in Wight's Contrib. Bot. of Ind. 113 (1834)	— — — — —	N.S.W.	Q. —	B.fl.VII,294	M.fr.VIII,239	As.		
<i>H. cylindrostachys</i> ; Boeckeler in Regensb. Flora, 108 (1875)	— — — — —	N.S.W.	Q. —	B.fl.VII,294	M.fr.VIII,240.			
<i>H. acuta</i> ; R. Brown, prodr. 224 (1810)	W.A. S.A. T. V.	N.S.W.	Q. N.A.	B.fl.VII,294	M.fr.VIII,240.	...	Am., N.Z.	
<i>H. multicaulis</i> ; Smith, Engl. Flor. II, 64 (1824)	W.A. S.A. — V.	N.S.W.	—	B.fl.VII,296	...	...	E., Am., N.Z.	
<i>H. atricha</i> ; R. Brown, prodr. 225 (1810)	— — — — —	N.S.W.	Q. N.A.	B.fl.VII,295	M.fr.VII,252; IX,100.			
<i>H. capitata</i> ; R. Brown, prodr. 225 (1810)	W.A. — — —	—	Q. N.A.	B.fl.VII,296	M.fr.VI,94; VIII,240.	As., Afr., Am.		
<i>H. atropurpurea</i> ; Kunth, enum. II, 151 (1837)	— — — — —	N.S.W.	Q. N.A.	B.fl.VII,296	M.fr.VIII,240	...	E., As., Afr., Am.	
<i>H. acicularis</i> ; R. Brown, prodr. 224 (1810)	— S.A. T. V.	N.S.W.	—	B.fl.VII,297	M.fr.VIII,240	...	E., As., Afr., Am.	

FIMBRISTYLIS; Vahl, enum. plant. II, 285 (1806). (Abildgaardia, Trichelostylis, Oncostylis.)

<i>F. acicularis</i> ; R. Brown, prodr. 226 (1810)	— — — — —	—	Q. N.A.	B.fl.VII,301	M.fr.IX,54.			
<i>F. acuminata</i> ; Vahl, enum. II, 285 (1806)	— — — — —	N.S.W.	Q. N.A.	B.fl.VII,301	M.fr.VIII,274; IX,54.	As.		

F. punctata ; R. Brown, prodr. 226 (1810)	Q. — B.f.VII,302		
F. rhyticarya ; F. v. M., fragm. I, 215 (1859)	Q. N.A. B.f.VII,302 M.fr.I,215.		
F. leucostachya ; Boeckeler in Garcke, Linnaea XXXVIII, 385 (1874)	— N.A. B.f.VII,302 M.fr.IX,54.		
F. nutans ; Vahl, enum. II, 285 (1806)	Q. N.A. B.f.VII,303 M.fr.VIII,274; IX,54.	As. P.	
F. pauciflora ; R. Brown, prodr. 225 (1810)	Q. N.A. B.f.VII,303	As.	
F. cardiocarpa ; F. v. M., fragm. I, 194 (1859)	— N.A. B.f.VII,303 M.fr.I,194.		
F. leucocolea ; Bentham, Fl. Austr. VII, 304 (1878)	— N.A. B.f.VII,304		
F. polytrichoides ; R. Brown, prodr. 226 (1810)	Q. N.A. B.f.VII,304	As.	
F. androgyna ; R. Brown, prodr. 226 (1810)	— N.A. B.f.VII,304		
F. subbulbosa ; Bentham, Fl. Austr. VII, 305 (1878)	Q. — B.f.VII,305		
F. tetragona ; R. Brown, prodr. 226 (1810)	Q. N.A. B.f.VII,305 M.fr.I,194; VIII,274.	As.	
F. trigastrocarya ; F. v. M., fragm. I, 194 (1859)	— N.A. B.f.VII,305 M.fr.I,194.		
F. monandra ; F. v. M., fragm. I, 195 (1859)	— N.A. B.f.VII,306 M.fr.I,195.		
F. pterygosperma ; R. Brown, prodr. 226 (1810)	Q. N.A. B.f.VII,306 M.fr.I,193; IX,55.		
F. sphaerocephala ; Bentham, Fl. Austr. VII, 306 (1878)	Q. N.A. B.f.VII,306		
F. xyridis ; R. Brown, prodr. 226 (1810)	Q. N.A. B.f.VII,307 M.fr.VIII,274; IX,54.		
F. oxystachya ; F. v. M., fragm. I, 195 (1859)	— N.A. B.f.VII,307 M.fr.VIII,272.		
F. macrantha ; Boeckeler in Garcke, Linnaea XXXVIII, 388 (1874)	— N.A. B.f.VII,307 M.fr.IX,55.		
F. aquarrulosa ; F. v. M., fragm. I, 216 (1859)	Q. N.A. B.f.VII,308 M.fr.I,216.		
F. monostachya ; Hasskarl, pl. Jav. rar. 61-64 (1848)	Q. N.A. B.f.VII,308 M.fr.VIII,272; IX,54.	As., Afr., Am.	
F. Brownii ; Bentham, Fl. Austr. VII, 308 (1878)	— N.A. B.f.VII,308 M.fr.VIII,273.		
F. Dallachyi ; F. v. M. in Benth. Fl. Austr. VII, 309 (1878)	Q. — B.f.VII,309 M.fr.VIII,273.		
F. velata ; R. Brown, prodr. 227 (1810)	Q. N.A. B.f.VII,309 M.fr.IX,11,54	N.Z.	
F. aestivalis ; Vahl, enum. II, 288 (1806)	Q. N.A. B.f.VII,310 M.fr.IX,11,54	As., Am., P.	
F. dichotoma ; Vahl, enum. II, 287 (1806)	Q. N.A. B.f.VII,310 M.fr.IX,10,54	As., Afr., Am., P.	
F. depauperata ; R. Brown, prodr. 227 (1810)	— N.A. B.f.VII,311		
F. spirostachya ; F. v. M. in Benth. Fl. Austr. VII, 311 (1878)	— N.A. B.f.VII,311		
F. communis ; Kunth, enum. II, 234 (1837)	Q. N.A. B.f.VII,311 M.fr.I,196; IX,10,54.	As., Afr., Am., P.	
F. ferruginea ; Vahl, enum. 291 (1806)	Q. N.A. B.f.VII,312 M.fr.I,197; IX,10,54.	As., Afr., Am., P.	
F. denudata ; R. Brown, prodr. 227 (1810)	Q. N.A. B.f.VII,313 M.fr.IX,9.		
F. elata ; R. Brown, prodr. 227 (1810)	— N.A. B.f.VII,313		
F. caespitosa ; R. Brown, prodr. 228 (1810)	Q. N.A. B.f.VII,313 M.fr.I,199.		
F. spiralis ; R. Brown, prodr. 226 (1810)	— N.A. B.f.VII,314		
F. subaristata ; Bentham, Fl. Austr. VII, 314 (1878)	— N.A. B.f.VII,314		
F. leptoclada ; Bentham, Fl. Austr. VII, 314 (1878)	Q. — B.f.VII,314		
F. debilis ; F. v. M., fragm. I, 198 (1859)	— N.A. B.f.VII,315 M.fr.I,198.		
F. corynocarya ; F. v. M., fragm. I, 197 (1859)	— N.A. B.f.VII,315 M.fr.I,187.		
F. solidifolia ; F. v. M., fragm. I, 198 (1859)	— N.A. B.f.VII,315 M.fr.I,198.		
F. obtusangula ; F. v. M., fragm. I, 198 (1859)	Q. N.A. B.f.VII,315 M.fr.I,198.		
F. miliacea ; Vahl, enum. II, 287 (1806)	Q. N.A. B.f.VII,316 M.fr.I,199; IX,12,54.	As., P.	
F. rara ; R. Brown, prodr. 227 (1810)	— N.A. B.f.VII,316		
F. microcarya ; F. v. M., fragm. I, 200 (1859)	Q. N.A. B.f.VII,316 M.fr.I,200.		
F. quinquangularis ; Kunth, enum. II, 229 (1837)	— N.A. B.f.VII,317	As., Afr.	
F. cyperoides ; R. Brown, prodr. 228 (1810)	Q. N.A. B.f.VII,317 M.fr.VIII,273; IX,11,54.	As.	
F. furva ; R. Brown, prodr. 228 (1810)	Q. — B.f.VII,318		
F. cymosa ; R. Brown, prodr. 228 (1810)	Q. N.A. B.f.VII,318		
F. multifolia ; Boeckeler in Garcke, Linnaea XXXVIII, 397 (1874)	— N.A. B.f.VII,319 M.fr.IX,55.		
F. aericea ; R. Brown, prodr. 228 (1810)	— N.A. B.f.VII,319		
F. macrostachya ; Boeckeler in Garcke, Linnaea XXXVIII, 386 (1874)	— N.A. B.f.VII,319 M.fr.IX,54.		
F. Neilsoni ; F. v. M., fragm. IX, 79 (1875)	Q. — B.f.VII,320 M.fr.IX,79.		
F. capitata ; R. Brown, prodr. 228 (1810)	Q. N.A. B.f.VII,320 M.fr.I,196.		

F. Schultzii ; Boeckeler in Gareke, Linnaea XXXVIII, 391 (1874)	...	...	—	—	—	—	—	N.A.	B.fl.VII,321	M.fr.IX,55.		
F. barbata ; Benth. Fl. Austr. VII, 321 (1878)	...	...	...	...	—	S.A.	—	N.S.W.	Q. N.A.	B.fl.VII,321	M.fr.IX,55	...
F. capillaris ; A. Gray, Man. of Bot. N.U.S. fifth ed. 567 (1867)	...	...	...	...	W.A.	—	—	—	Q. N.A.	B.fl.VII,322	M.fr.IX,55	...

SCIRPUS ; Tournefort, inst. 528, t. 360 (1700) from Terentius. (Isoplepis, Malacochaete.)

S. humillimus ; Benth. Fl. Austr. VII, 324 (1878)	...	...	...	...	—	—	—	—	N.A.	B.fl.VII,324	M.fr.IX,7.	
S. fluitans ; Linné, spec. plant. 48 (1753)	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	—	B.fl.VII,325	M.fr.IX,55
S. arenarius ; Benth. Fl. Austr. VII, 325 (1878)	...	...	...	...	W.A.	—	—	V.	—	—	B.fl.VII,325	...
S. lenticularis ; Sprengel, syst. I, 208 (1825)	...	...	...	...	...	—	—	—	N.S.W.	—	B.fl.VII,326	M.fr.IX,55
S. crassiusculus ; J. Hooker in Benth. Fl. Austr. VII, 326 (1878)	...	...	...	...	...	—	—	T.	V.	N.S.W.	—	B.fl.VII,326
S. cyperoides ; Sprengel, syst. I, 208 (1825)	...	...	...	...	W.A.	—	—	—	—	—	B.fl.VII,326	...
S. setaceus ; Linné, spec. plant. 49 (1753)	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	B.fl.VII,327	M.fr.IX,55
S. riparius ; Sprengel, syst. I, 208 (1825)	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	—	B.fl.VII,327	...
S. cartilagineus ; Sprengel, syst. I, 208 (1825)	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	—	B.fl.VII,328	M.fr.IX,55
S. squarrosus ; Linné, mant. plant. alt. 181 (1771)	...	...	...	...	...	—	—	—	—	Q. N.A.	B.fl.VII,329	...
S. inundatus ; Sprengel, syst. I, 207 (1825)	...	...	...	...	...	—	S.A.	T.	V.	N.S.W.	Q.	B.fl.VII,329
S. prolifer ; Rottboell, descr. et icon. pl. 55, t. 17 (1773)	...	...	...	...	...	—	—	—	N.S.W.	—	B.fl.VII,330	M.fr.IX,55
S. articulatus ; Linné, spec. plant. 47 (1753)	...	...	...	...	...	—	—	—	—	Q. N.A.	B.fl.VII,331	M.fr.IX,55
S. nodosus ; Rottboell, descr. et icon. pl. 52, t. 8 (1773)	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	B.fl.VII,331	M.fr.IX,55
S. supinus ; Linné, spec. plant. 49 (1753)	...	...	...	...	...	—	S.A.	—	V.	N.S.W.	Q. N.A.	B.fl.VII,330
S. debilis ; Pursch, flor. Amer. Sept. I, 55 (1814)	...	...	...	...	...	—	—	—	—	Q. N.A.	B.fl.VII,332	...
S. mucronatus ; Linné, spec. plant. 50 (1753)	...	...	...	...	...	—	—	—	N.S.W.	Q.	B.fl.VII,332	M.fr.IX,8,56...
S. pungens ; Vahl, enum. II, 255 (1806)	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	—	B.fl.VII,333	M.fr.IX,8,56...
S. lacustris ; Linné, spec. plant. 48 (1753)	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	B.fl.VII,333	M.fr.IX,7,56...
S. litoralis ; Schrader, Fl. Germ. I, 142, t. 5 (1806)	...	...	...	...	...	—	S.A.	—	N.S.W.	Q. N.A.	B.fl.VII,334	M.fr.IX,7,56; XII,25
S. maritimus ; Linné, spec. plant. 51 (1753)	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q. N.A.	B.fl.VII,335	M.fr.IX,8,56...
S. grossus ; Linné, fil. suppl. plant. 104 (1781)	...	...	...	...	...	—	—	—	—	Q.	—	M.fr.XII,24
S. polystachyus ; F. v. M. in Transact. phil. Soc. Vict. I, 108 (1854)	...	...	...	...	...	—	—	—	V.	N.S.W.	—	B.fl.VII,335

LIPOCARPHA ; R. Brown in Tuckey's narrat. of an Exped. to Congo, 459 (1818). (Hypaeptyum.)

L. argentea ; R. Brown in Tuck. Congo, 459 (1818)	...	...	...	...	—	—	—	—	Q.	—	B.fl.VII,336	M.fr.VIII,238; IX,57.	As., Afr.
L. microcephala ; R. Brown in Tuck. Congo, 459 (1818)	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q. N.A.	B.fl.VII,337	M.fr.VIII,238; IX,57.	As., P.

FUIRENA ; Rottboell, descr. et icon. rar. pl. illustr. 70 (1773).

F. umbellata ; Rottboell, descr. et icon. pl. 70, t. 19 (1773)	...	...	...	—	—	—	—	Q. N.A.	B.fl.VII,337	M.fr.VIII,238; IX,57.	As., Afr., Am., P.
F. glomerata ; Lamarck, Illustr. des genr. I, 150 (1791)	...	...	...	—	—	—	N.S.W.	Q. N.A.	B.fl.VII,338	M.fr.VIII,238; IX,57.	As., Afr., P.

HYPELYTRUM ; L. C. Richard in Persoon, synops. I, 70 (1805). (Hypolytrum.)

H. latifolium ; L. C. Richard in Pers. syn. I, 70 (1805)	...	...	...	...	—	—	—	—	Q.	—	B.fl.VII,339	M.fr.VIII,238; IX,57.	As., Afr., Am., P.
--	-----	-----	-----	-----	---	---	---	---	----	---	--------------	-----------------------	--------------------

EXOCARYA ; Benth. in J. Hooker, icon. plant. t. 1206 (1877).

E. scleroides ; Benth. in Hook. icon. pl. t. 1206 (1877)	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,339	M.fr.IX,12.
--	-----	-----	-----	-----	---	---	---	---	--------	----	---	--------------	-------------

MAPANIA ; Aublet, Hist. pl. Guian. 47, t. 17 (1775). (Pandaphyllum.)

M. hypelytroides ; F. v. M., fragm. IX, 16 (1875)	...	...	...	...	—	—	—	—	Q.	—	B.fl.VII,341	M.fr.IX,16	...
---	-----	-----	-----	-----	---	---	---	---	----	---	--------------	------------	-----

SCIRPODENDRON ; Zippelius in Journ. Asiat. Soc. Beng. XXXVIII, 85 (1869).

S. costatum ; Kurz in Journ. Asiat. Soc. Beng. XXXVIII, 85 (1869)	...	...	...	...	—	—	—	—	Q.	—	B.fl.VII,341	M.fr.X,104	...
---	-----	-----	-----	-----	---	---	---	---	----	---	--------------	------------	-----

LEPIRONIA ; L. C. Richard in Persoon, synops. I, 70 (1805). (Chondrachne.)

L. mucronata ; L. C. Richard in Persoon, syn. I, 70 (1805)	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,342	M.fr.IX,17,58
--	-----	-----	-----	-----	---	---	---	---	--------	----	---	--------------	---------------

CHORIZANDRA ; R. Brown, prodr. 221 (1810). (Chorisandra.)

C. spaerocephala ; R. Brown, prodr. 221 (1810)	...	...	...	...	...	—	—	—	N.S.W.	Q.	—	B.fl.VII,344	M.fr.IX,18,58.
C. enodis ; Nees in Lehm. pl. Preiss. II, 73 (1846)	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,344	M.fr.IX,18.

C. multiarticulata; Nees in Ann. and Mag. of nat. Hist. VI, 48 (1842) ...	W.A.	—	—	—	—	—	—	B.fl.VII,345	M.fr.IX,18.		
C. cymbaria; R. Brown, prodr. 221 (1810) ...	W.A.	—	—	V.	N.S.W.	Q.	—	B.fl.VII,345	M.fr.IX,18,58	...	P.
OREOBOLUS; R. Brown, prodr. fl. Nov. Holl. 235 (1810).											
O. Pumilio; R. Brown, prodr. 226 (1810) ...	—	—	T.	V.	N.S.W.	—	—	B.fl.VII,346	M.fr.IX,20,58	...	Am., P., N.Z.
REMIREA; Aublet, Hist. des pl. de la Guian. franc. I, 44, t. 16 (1775).											
R. maritima; Aublet, Hist. pl. Guian. I, 45, t. 16 (1775) ...	—	—	—	—	—	—	Q.	—	B.fl.VII,347	M.fr.V,92; IX,20,58.	As., Afr., Am., P.
RHYNCHOSPORA; Vahl, enum. plant. II, 229 (1806). (Morisia.)											
R. aurea; Vahl, enum. II, 229 (1806) ...	—	—	—	—	—	—	Q.	—	B.fl.VII,348	M.fr.IX,17,57	As., Afr., Am., P.
R. glauca; Vahl, enum. II, 229 (1806) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,349	M.fr.IX,17,57	As., Afr., Am.	
R. Wallichiana; Kunth, enum. II, 289 (1837) ...	—	—	—	—	—	Q.	N.A.	B.fl.VII,349	M.fr.IX,17,52	As., Afr., Am.	
R. longisetis; R. Brown, prodr. 230 (1810) ...	—	—	—	—	—	Q.	N.A.	B.fl.VII,350	M.fr.IX,17,58.		
R. tenuifolia; Benth, Fl. Austr. VII, 350 (1878) ...	—	—	—	—	—	Q.	N.A.	B.fl.VII,350			
CYATHOCHAETE; Nees in Lehm. pl. Preiss. II, 86 (1846). (Tetralepis.)											
C. clandestina; Benth, Fl. Austr. VII, 351 (1878) ...	W.A.	—	—	—	—	—	—	B.fl.VII,351	M.fr.IX,40; XII,25.		
C. avenacea; Benth, Fl. Austr. VII, 351 (1878) ...	W.A.	—	—	—	—	—	—	B.fl.VII,351	M.fr.IX,40.		
C. diandra; Nees in Lehm. pl. Preiss. II, 86 (1846) ...	—	—	—	—	V.	N.S.W.	—	—	B.fl.VII,352	M.fr.IX,39; XII,25.	
CARPHA; Banks & Solander in R. Brown, prodr. 230 (1810).											
C. alpina; R. Brown, prodr. 230 (1810) ...	—	—	T.	V.	N.S.W.	—	—	B.fl.VII,381	M.fr.IX,39 ...	...	P., N.Z.
SCHOENUS; Linné, coroll. gen. 2 (1737). (Chaetospora, Elynanthus, Tricostularia, Helothrix, Isoschoenus, Gymnochaete, Gymnoschoenus, Mesomelaena, Discopodium.)											
S. cruentus; F. v. M., fragm. IX, 36 (1875) ...	W.A.	—	—	—	—	—	—	B.fl.VII,357	M.fr.IX,37.		
S. Benthami; F. v. M., syst. Cens. of Austr. pl. 127 (1882) ...	W.A.	—	—	—	—	—	—	B.fl.VII,357	M.fr.IX,37.		
S. lanatus; Labillardière, Nov. Holl. pl. spec. I, 19, t. 20 (1804) ...	W.A.	—	—	—	—	—	—	B.fl.VII,357	M.fr.IX,37.		
S. curvifolius; Poir, Encycl. méth. suppl. II, 251 (1811) ...	W.A.	—	—	—	—	—	—	B.fl.VII,358	M.fr.IX,36.		
S. capitatus; F. v. M., fragm. IX, 58 (1875) ...	W.A.	—	—	—	—	—	—	B.fl.VII,358	M.fr.IX,37.		
S. setifolius; Benth, Fl. Austr. VII, 359 (1878) ...	W.A.	—	—	—	—	—	—	B.fl.VII,359			
S. Drummondii; Benth, Fl. Austr. VII, 359, non. Steud. (1870) ...	W.A.	—	—	—	—	—	—	B.fl.VII,359	M.fr.IX,37.		
S. turbinatus; Poir, Encycl. méth. suppl. II, 251 (1811) ...	—	—	—	—	N.S.W.	—	—	B.fl.VII,359	M.fr.IX,33.		
S. barbatus; Boeckeler in Garcke, Linnaea XXXVIII, 277 (1874) ...	W.A.	—	—	—	—	—	—	B.fl.VII,360	M.fr.IX,30.		
S. flavus; Boeckeler in Garcke, Linnaea XXXVIII, 278 (1874) ...	W.A.	—	—	—	—	—	—	B.fl.VII,360	M.fr.IX,30.		
S. brevisetis; Poir, Encycl. méth. suppl. II, 251 (1811) ...	W.A.	—	—	—	—	—	—	B.fl.VII,360	M.fr.IX,30.		
S. Armeria; Boeckeler in Garcke, Linnaea XXXVIII, 279 (1874) ...	W.A.	—	—	—	—	—	—	B.fl.VII,361	M.fr.IX,30.		
S. aphyllus; Boeckeler in Garcke, Linnaea XXXVIII, 280 (1874) ...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.VII,361	M.fr.IX,28; XII,26.		
S. imberbis; R. Brown, prodr. 231 (1810) ...	—	—	—	V.	N.S.W.	—	—	B.fl.VII,361	M.fr.IX,28.		
S. ericetorum; R. Brown, prodr. 231 (1810) ...	—	—	—	V.	N.S.W.	Q.	—	B.fl.VII,362	M.fr.IX,58; XII,26.		
S. nitens; Poir, Encycl. méth. suppl. II, 251 (1811) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,362	M.fr.IX,35	...	Am., N.Z.
S. cygneus; Nees in Lehmann, pl. Preiss. II, 81 (1846) ...	W.A.	—	—	—	—	—	—	B.fl.VII,363	M.fr.IX,80.		
S. minutulus; F. v. M., fragm. IX, 32 (1875) ...	W.A.	—	—	—	—	—	—	B.fl.VII,363	M.fr.IX,32.		
S. trachycarpus; F. v. M., fragm. IX, 33 (1875) ...	W.A.	—	—	—	—	—	—	B.fl.VII,363	M.fr.IX,33.		
S. Tepperi; F. v. M., fragm. XI, 106 (1880) ...	—	S.A.	T.	—	—	—	—	—	M.fr.XI,106.		
S. nanus; F. v. M., fragm. IX, 36 (1875) ...	W.A.	—	—	—	—	—	—	B.fl.VII,364	M.fr.IX,28.		
S. pleiostemoneus; F. v. M., fragm. IX, 52 (1875) ...	W.A.	—	—	—	—	—	—	B.fl.VII,364	M.fr.IX,52.		
S. breviculmis; Benth, Fl. Austr. VII, 364 (1878) ...	W.A.	—	—	—	—	—	—	B.fl.VII,364			
S. deformis; Poir, Encycl. méth. suppl. II, 251 (1811) ...	—	S.A.	—	—	—	—	—	B.fl.VII,364	M.fr.IX,39.		
S. unispiculatus; F. v. M. in Benth. Fl. Austr. VII, 365 (1878) ...	W.A.	—	—	—	—	—	—	B.fl.VII,365			
S. obtusifolius; Boeckeler in Garcke, Linnaea XXXVIII, 281 (1874) ...	W.A.	—	—	—	—	—	—	B.fl.VII,366	M.fr.IX,31.		
S. grammatophyllus; F. v. M., fragm. IX, 31 (1875) ...	W.A.	—	—	—	—	—	—	B.fl.VII,366	M.fr.IX,31.		
S. asperocarpus; F. v. M., fragm. IX, 29 (1875) ...	W.A.	—	—	—	—	—	—	B.fl.VII,366	M.fr.IX,29.		

S. Moorei; Benth. Fl. Austr. VII, 367 (1878)	...	...	...	...	...	...	...	...	N.S.W.	—	—	B.fl. VII, 367				
S. villosus; R. Brown, prodr. 231 (1810)	...	...	...	...	...	...	...	...	N.S.W.	—	—	B.fl. VII, 367	M.fr. IX, 28.			
S. grandiflorus; F. v. M., fragm. IX, 30 (1875)	...	...	...	...	...	...	...	W.A.	—	—	—	B.fl. VII, 367	M.fr. IX, 30.			
S. distans; F. v. M., syst. Cens. of Austr. pl. 127 (1882)	...	...	...	...	...	...	...	W.A.	—	—	—	—	M.fr. IX, 35.			
S. calostachyus; Poir. Encycl. méth. suppl. II, 251 (1811)	...	...	...	...	...	...	...	...	N.S.W.	Q.	—	B.fl. VII, 368	M.fr. IX, 35.			
S. scabripes; Benth. Fl. Austr. VII, 368 (1878)	...	...	...	...	...	...	...	...	—	—	Q.	B.fl. VII, 368				
S. multiglumis; Benth. Fl. Austr. VII, 368 (1878)	...	...	...	...	...	...	...	W.A.	—	—	—	B.fl. VII, 368				
S. efoliatus; F. v. M., fragm. IX, 32 (1875)	...	...	...	...	...	...	...	W.A.	—	—	—	B.fl. VII, 369	M.fr. IX, 32.			
S. acuminatus; R. Brown, prodr. 231 (1810)	...	...	...	...	...	...	...	W.A.	—	—	—	B.fl. VII, 369	M.fr. IX, 80.			
S. pedicellatus; Poir. Encycl. méth. suppl. II, 251 (1811)	...	...	...	...	...	...	...	W.A.	—	—	—	B.fl. VII, 369				
S. fascicularis; Nees in Ann. and Mag. of nat. Hist. VI, 48 (1842)	...	...	...	...	...	...	...	W.A.	—	—	—	B.fl. VII, 370				
S. brevifolius; R. Brown, prodr. 231 (1810)	...	...	...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	B.fl. VII, 370	M.fr. IX, 29, 80.	
S. melanostachys; R. Brown, prodr. 231 (1810)	...	...	...	...	...	...	...	...	—	—	V.	N.S.W.	Q.	B.fl. VII, 370	M.fr. IX, 29.	
S. sparteus; R. Brown, prodr. 231 (1810)	...	...	...	...	...	...	...	...	—	—	—	Q. N.A.	B.fl. VII, 371			
S. vaginatus; F. v. M. in Benth. Fl. Austr. VII, 371 (1878)	...	...	...	...	...	...	...	...	—	—	—	N.S.W.	Q.	B.fl. VII, 371		
S. falcatus; R. Brown, prodr. 232 (1810)	...	...	...	...	...	...	...	...	—	—	—	Q. N.A.	B.fl. VII, 372	M.fr. IX, 29.		
S. punctatus; R. Brown, prodr. 232 (1810)	...	...	...	...	...	...	...	...	—	—	—	—	N.A.	B.fl. VII, 372		
S. indutus; F. v. M. in Benth. Fl. Austr. VII, 372 (1878)	...	...	...	...	...	...	...	W.A.	—	—	—	Q.	B.fl. VII, 372			
S. bifida; Boeckeler in Garcke, Linnaea XXXVIII, 282 (1874)	...	...	...	...	...	...	...	W.A.	—	—	—	—	B.fl. VII, 373	M.fr. IX, 38, 80.		
S. apogon; Roemer et Schultes, syst. veg. II, 77 (1817)	...	...	...	...	...	...	...	—	S.A.	T.	V.	N.S.W.	Q.	B.fl. VII, 373	M.fr. IX, 56; X, 120. N.Z.	
S. odontocarpus; F. v. M., fragm. IX, 32 (1875)	...	...	...	...	...	...	...	W.A.	—	—	—	—	B.fl. VII, 374	M.fr. IX, 32.		
S. humilis; Benth. Fl. Austr. VII, 374 (1878)	...	...	...	...	...	...	...	W.A.	—	—	—	—	B.fl. VII, 374			
S. sculptus; Boeckeler in Garcke, Linnaea XXXVIII, 286 (1874)	...	...	...	...	...	...	...	W.A.	S.A.	—	—	—	B.fl. VII, 374	M.fr. IX, 30.		
S. axillaris; Poir. Encycl. méth. suppl. II, 251 (1811)	...	...	...	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	—	B.fl. VII, 375	M.fr. IX, 34 ... N.Z.	
S. tenellus; Benth. Fl. Austr. VII, 375 (1878)	...	...	...	...	...	...	...	W.A.	—	—	—	—	B.fl. VII, 375			
S. natans; F. v. M., fragm. IX, 36 (1875)	...	...	...	...	...	...	...	W.A.	—	—	—	—	B.fl. VII, 375	M.fr. IX, 38.		
S. fluitans; J. Hooker, Fl. Tasman. II, 81, t. 141 (1860)	...	...	...	...	...	...	...	...	—	—	T.	—	B.fl. VII, 376	M.fr. IX, 28.		
S. octandrus; F. v. M., fragm. IX, 31 (1875)	...	...	...	...	...	...	...	W.A.	—	—	—	—	B.fl. VII, 377	M.fr. IX, 31.		
S. capillaris; F. v. M., syst. Cens. of Austr. pl. 128 (1882)	...	...	...	...	...	...	...	—	S.A.	T.	V.	—	B.fl. VII, 377	M.fr. IX, 34, 80.		
S. stygius; Poir. Encycl. méth. suppl. II, 251 (1811)	...	...	...	...	...	...	...	W.A.	—	—	—	—	B.fl. VII, 378	M.fr. IX, 36, 34, 80.		
S. deustus; F. v. M., syst. Cens. of Austr. pl. 128 (1882) from R. Brown (1810)	...	...	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	B.fl. VII, 379	M.fr. IX, 39.	
S. tetragonus; Poir. Encycl. méth. suppl. II, 251 (1811)	...	...	...	...	...	...	...	W.A.	—	—	—	—	B.fl. VII, 380	M.fr. IX, 36.		
S. sphaerocephalus; Poir. Encycl. méth. suppl. II, 251 (1811)	...	...	...	...	...	...	...	...	—	—	T.	V.	N.S.W.	—	B.fl. VII, 380	M.fr. IX, 33.
S. anceps; Poir. Encycl. méth. suppl. II, 251 (1811)	...	...	...	...	...	...	...	W.A.	—	—	—	—	B.fl. VII, 380	M.fr. IX, 30.		
S. paludosus; Poir. Encycl. méth. suppl. II, 251 (1811)	...	...	...	...	...	...	...	...	—	—	—	N.S.W.	Q.	B.fl. VII, 382	M.fr. IX, 35.	
S. pauciflorus; syst. Cens. of Austr. pl. 128 (1882)	...	...	...	...	...	...	...	...	—	—	—	V.	—	B.fl. VII, 383	M.fr. IX, 23.	
S. Tricostularia; F. v. M., syst. Cens. of Austr. pl. 128 (1882)	...	...	...	...	...	...	...	W.A.	—	—	—	—	B.fl. VII, 383	M.fr. IX, 37.		
S. Neesii; F. v. M., syst. Cens. of Austr. pl. 128 (1882) from Lehmann (1846)	...	...	...	...	...	...	...	W.A.	—	—	—	—	B.fl. VII, 383	M.fr. IX, 37.		
S. fimbriatylodes; F. v. M., syst. Cens. of Austr. pl. 128 (1882)	...	...	...	...	...	...	...	...	—	—	—	—	N.A.	B.fl. VII, 384	M.fr. IX, 34.	

LEPIDOSPOREA; F. v. M., fragm. phytograph. Austral. IX, 34 (1875).

L. tenuissima; F. v. M., fragm. IX, 34 (1875)	...	...	...	...	...	...	...	...	S.A.	T.	V.	N.S.W.	—	—	B.fl. VII, 365	M.fr. IX, 34; XII, 26.
--	-----	-----	-----	-----	-----	-----	-----	-----	------	----	----	--------	---	---	----------------	------------------------

LEPIDOSPERMA; Labillardière, Nov. Holl. pl. spec. I, 14 (1804).

L. gladiatum; Labillardière, Nov. Holl. pl. spec. I, 15, t. 12 (1804)	...	...	...	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl. VII, 387	M.fr. XII, 26.
L. effusum; Benth. Fl. Austr. VII, 387 (1878)	...	...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. VII, 387	
L. rupestre; Benth. Fl. Austr. VII, 388 (1878)	...	...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. VII, 388	
L. elatius; Labillardière, Nov. Holl. pl. spec. I, 15, t. 11 (1804)	...	...	...	...	...	...	...	—	S.A.	T.	V.	—	—	—	B.fl. VII, 388	M.fr. IX, 25.
L. tetraquetrum; Nees in Lehm. pl. Preisa. II, 90 (1846)	...	...	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. VII, 388	M.fr. IX, 24.
L. Oldfieldii; J. Hooker, Fl. Tasman. II, 91, t. 146 (1860)	...	...	...	...	...	...	...	—	—	T.	—	—	—	—	B.fl. VII, 389	
L. exaltatum; R. Brown, prodr. 234 (1810)	...	...	...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl. VII, 389	

L. longitudinale; Labillardière, Nov. Holl. pl. spec. I, 16, t. 13 (1804) ...	... W.A. S.A. T. V. N.S.W. — — B.fl.VII,389 M.fr.IX,25.	
L. concavum; R. Brown, prodr. 234 (1810) ...	... — S.A. T. V. N.S.W. Q. — B.fl.VII,390 M.fr.IX,24 ...	... N.Z.
L. angustatum; R. Brown, prodr. 235 (1810) ...	... W.A. — — — — — B.fl.VII,391 M.fr.IX,26,80.	
L. Drummondii; Benth, Fl. Austr. VII, 391 (1878) ...	... W.A. — — — — — B.fl.VII,391	
L. Brunonianum; Nees in Lehm. pl. Preiss. II, 92 (1846) ...	... W.A. — — — — — B.fl.VII,392	
L. tuberculatum; Nees in Lehm. pl. Preiss. II, 90 (1846) ...	... W.A. — — — — — B.fl.VII,392 M.fr.IX,26.	
L. resinosum; F. v. M., fragm. IX, 26 (1875) implied ...	... W.A. — — — — — B.fl.VII,392 M.fr.IX,26.	
L. viscidum; R. Brown, prodr. 234 (1810) ...	... — S.A. — V. N.S.W. — — B.fl.VII,393	
L. costale; Nees in Lehm. pl. Preiss. II, 92 (1846) ...	... W.A. — — — — — B.fl.VII,393	
L. laterale; R. Brown, prodr. 234 (1810) ...	... — S.A. T. V. N.S.W. Q. — B.fl.VII,393	
L. congestum; R. Brown, prodr. 234 (1810) ...	... — S.A. — — — — — B.fl.VII,394	
L. globosum; Labillardière, Nov. Holl. pl. spec. I, 16, t. 14 (1804) ...	... — S.A. T. V. — — — B.fl.VII,394	
L. lineare; R. Brown, prodr. 235 (1810) ...	... — S.A. T. V. N.S.W. — — B.fl.VII,395 M.fr.IX,26.	
L. aphyllum; R. Brown, prodr. 235 (1810) ...	... W.A. — — — — — B.fl.VII,395	
L. gracile; R. Brown, prodr. 235 (1810) ...	... W.A. — — — — — B.fl.VII,395	
L. semiteres; F. v. M. in Garcke, Linnaea XXXVIII, 327 (1874) ...	... — S.A. — V. — — — B.fl.VII,396	
L. canescens; Boeckeler in Garcke, Linnaea XXXVIII, 330 (1874) ...	... — S.A. — V. N.S.W. — — B.fl.VII,396 M.fr.IX,24.	
L. pubisquameum; Steudel, syn. pl. glum. II, 158 (1855) ...	... W.A. — — — — — B.fl.VII,396 M.fr.IX,27.	
L. scabrum; Nees in Lehm. pl. Preiss. II, 92 (1846) ...	... W.A. — — — — — B.fl.VII,397 M.fr.IX,27.	
L. tenue; Benth, Fl. Austr. VII, 397 (1878) ...	... W.A. — — — — — B.fl.VII,397	
L. leptostachyum; Benth, Fl. Austr. VII, 397 (1878) ...	... W.A. — — — — — B.fl.VII,397	
L. leptophyllum; Benth, Fl. Austr. VII, 398 (1878) ...	... W.A. — — — — — B.fl.VII,398	
L. tortuosum; F. v. M., general Report 9 (1855) ...	... — — — V. — — — B.fl.VII,398 M.fr.IX,23.	
L. flexuosum; R. Brown, prodr. 235 (1810) ...	... — — — — N.S.W. — — B.fl.VII,398	
L. filiforme; Labillardière, Nov. Holl. pl. spec. I, 17, t. 15 (1804) ...	... — S.A. T. V. N.S.W. — — B.fl.VII,398 M.fr.IX,27; XII,26.	
L. striatum; R. Brown, prodr. 235 (1810) ...	... W.A. — — — — — B.fl.VII,399 M.fr.IX,26,27.	
L. Neesii; Kunth, enum. II, 319 (1837) ...	... — — — V. N.S.W. — — B.fl.VII,399 M.fr.IX,24,27.	
L. carphoides; F. v. M. in Benth. Fl. Austr. VII, 400 (1878) ...	... W.A. S.A. — V. — — — B.fl.VII,400	

GAHNIA; R. et G. Forster, char. gen. 51 (1776). (Cladium, Lampocarya, Melachne, Epiandra, Didymonema, Baumea, Chapelliera, Morelotia.)

G. Mariscus; F. v. M., Key to Vict. pl. I, 456 (1888) from Linné (1753) ...	... — S.A. — V. N.S.W. Q. N.A. B.fl.VII,402 M.fr.IX,14,57; XII,25.	E.,As.,Afr.,Am.,P.
G. insularis; Cladium; Benth, Fl. Austr. 403 (1878) ...	... — — — — N.S.W. — — B.fl.VII,403	
G. articulata; F. v. M., Key to Vict. pl. I, 453 (1888) from R. Brown (1810) ...	... W.A. S.A. — V. N.S.W. Q. — B.fl.VII,403 M.fr.IX,14,57 ...	N.Z., P.
G. arthropphylla; Cladium; F. v. M., fragm. IX, 14 (1875) ...	... W.A. — — — — — B.fl.VII,403 M.fr.IX,14.	
G. glomerata; F. v. M., Key to Vict. pl. I, 455 (1888) from R. Brown (1810) ...	... W.A. S.A. T. V. N.S.W. Q. N.A. B.fl.VII,404 M.fr.IX,15,57 ...	N.Z.
G. Preissii; Baumea; Nees in Lehm. pl. Preiss. II, 75 (1846) ...	... W.A. — — — — — B.fl.VII,405 M.fr.IX,13,15,56.	
G. laxa; Chapelliera; Nees in Lehm. pl. Preiss. II, 76 (1846) ...	... W.A. — — — — — B.fl.VII,405	
G. riparia; Cladium; Benth, Fl. Austr. VII, 405 (1878) ...	... W.A. — — — — — B.fl.VII,405	
G. teretifolia; Cladium; R. Brown, prodr. 237 (1810) ...	... — — — — N.S.W. Q. — B.fl.VII,406 M.fr.IX,15,56; XII,25.	N.Z.
G. tetraquetra; F. v. M., Key to Vict. pl. I, 454 (1888) from J. Hooker (1860) ...	... — S.A. T. V. N.S.W. — — B.fl.VII,406 ...	N.Z.
G. schoenoides; F. v. M., Key to Vict. pl. I, 454 (1888) from R. Brown (1810) ...	... W.A. S.A. T. V. N.S.W. Q. — B.fl.VII,407 M.fr.XII,26.	
G. Gunnii; F. v. M., Key to Vict. pl. I, 455 (1888) from J. Hooker (1860) ...	... — S.A. T. V. N.S.W. — — B.fl.VII,407 M.fr.IX,15,57.	
G. juncea; F. v. M., Key to Vict. pl. I, 453 (1888) from R. Brown (1810) ...	... W.A. S.A. T. V. N.S.W. Q. N.A. B.fl.VII,408 M.fr.IX,16,57 ...	N.Z.
G. vaginalis; Cladium; Benth, Fl. Austr. VII, 409 (1878) ...	... W.A. — — — — — B.fl.VII,408	
G. elynanthoides; Cladium; F. v. M., fragm. IX, 31 (1875) ...	... W.A. — — — — — B.fl.VII,409 M.fr.IX,31.	
G. Filum; F. v. M., Key to Vict. pl. I, 456 (1888) from Labillardière (1804) ...	... — S.A. T. V. — — — B.fl.VII,409 M.fr.IX,14,57; XII,26.	
G. trifida; Labillardière, Nov. Holl. pl. spec. I, 89, t. 116 (1804) ...	... W.A. S.A. T. V. N.S.W. — — B.fl.VII,413 M.fr.XII,26.	
G. decomposita; Benth, Fl. Austr. VII, 417 (1876) from R. Brown (1810) ...	... W.A. — — — — — B.fl.VII,417 M.fr.IX,18.	
G. microstachya; Benth, Fl. Austr. VII, 414 (1878) ...	... — — — V. N.S.W. — — B.fl.VII,414	
G. polyphylla; Benth, Fl. Austr. VII, 415 (1878) ...	... W.A. — — — — — B.fl.VII,415	
G. ancistrophylla; F. v. M. in Benth. Fl. Austr. VII, 415 (1878) ...	... W.A. — — — — — B.fl.VII,415	

G. lanigera; Benth. Fl. Austr. VII, 415 (1878) from R. Brown (1810)	...	W.A.	S.A.	—	—	—	—	—	B.f. VII, 415	M.fr. IX, 14, 57.	
G. aristata; F. v. M. in Benth. Fl. Austr. 416 (1878)	...	W.A.	—	—	—	—	—	—	B.f. VII, 416		
G. deusta; Benth. Fl. Austr. VII, 416 (1878) from R. Brown (1810)	...	W.A.	S.A.	—	—	—	—	—	B.f. VII, 416	M.fr. IX, 14.	
G. Radula; Benth. Fl. Austr. VII, 417 (1878) from R. Brown (1810)	...	—	S.A.	T.	V.	—	—	—	B.f. VII, 414	M.fr. IX, 13, 56.	
G. Sieberi; Boeckeler in Garcke, Linn. XXXVIII, 343 (1874) from Schrader (1830)	...	—	—	—	—	N.S.W.	—	—	B.f. VII, 414	M.fr. IX, 14, 57.	
G. tetragonocarpa; Boeckeler in Linnaea XXXVIII, 347 (1874)	...	—	—	—	V.	N.S.W.	—	—	B.f. VII, 418		
G. melanocarpa; R. Brown, prodr. 239 (1810)	...	—	—	T.	V.	N.S.W.	—	—	B.f. VII, 414	M.fr. IX, 13, 56; XII, 26.	
G. psittacorum; Labillardiere, Nov. Holl. pl. spec. I, 89, t. 115 (1804)	...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.f. VII, 418	M.fr. IX, 13, 56; XII, 25.	
G. aspera; Sprengel, syst. veg. II, 114 (1825)	...	—	—	—	—	N.S.W.	Q.	—	B.f. VII, 412	M.fr. IX, 12, 56	P.
G. xanthocarpa; J. Hooker, Handb. of the N.Z. Flor. 306 (1864)	...	—	—	—	—	N.S.W.	—	—	B.f. VII, 418	M.fr. IX, 13, 56	N.Z.

CAUSTIS; R. Brown, prodr. 239 (1810). (Eurostorrhiza.)

C. pentandra; R. Brown, prodr. 240 (1810)	...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.f. VII, 420	M.fr. IX, 19, 58; XII, 26.	
C. flexuosa; R. Brown, prodr. 239 (1810)	...	—	—	—	V.	N.S.W.	Q.	N.A.	B.f. VII, 421	M.fr. IX, 58.	
C. recurvata; Sprengel, syst. cur. post. 26 (1827)	...	—	—	—	—	N.S.W.	—	—	B.f. VII, 421	M.fr. IX, 19.	
C. restiacea; F. v. M. in Benth. Fl. Austr. VII, 421 (1878)	...	—	—	—	V.	N.S.W.	—	—	B.f. VII, 421	M.fr. IX, 19.	
C. dioica; R. Brown, prodr. 239 (1810)	...	W.A.	—	—	—	—	—	—	B.f. VII, 422	M.fr. IX, 19, 58.	

ARTHROSTYLIS; R. Brown, prodr. 229 (1810). (Arthrostyles.)

A. aphylla; R. Brown, prodr. 229 (1810)	...	—	—	—	—	—	—	Q. N.A.	B.f. VII, 423	M.fr. IX, 9, 55; XII, 25.	
---	-----	---	---	---	---	---	---	---------	---------------	---------------------------	--

REEDIA; F. v. M., fragm. I, 240, t. 10 (1859).

R. spathacea; F. v. M., fragm. I, 240, t. 10 (1859)	...	W.A.	—	—	—	—	—	—	B.f. VII, 423	M.fr. IX, 15.	
---	-----	------	---	---	---	---	---	---	---------------	---------------	--

EUANDRA; R. Brown, prodr. 239 (1810). (Evandra.)

E. aristata; R. Brown, prodr. 239 (1810)	...	W.A.	—	—	—	—	—	—	B.f. VII, 424	M.fr. IX, 18.	
E. pauciflora; R. Brown, prodr. 239 (1810)	...	W.A.	—	—	—	—	—	—	B.f. VII, 425	M.fr. IX, 18.	

SCLERIA; Bergius in Svensk. Vet. Acad. Handl. 149, t. 4 (1765). (Diplacrum, Hypoporum, Sphaeropus.)

S. caricina; Benth. Fl. Austr. VII (1878)	...	—	—	—	—	—	Q.	—	B.f. VII, 426	M.fr. IX, 58...	As.
S. pygmaea; R. Brown, prodr. 240 (1810)	...	—	—	—	—	—	Q. N.A.	B.f. VII, 427	M.fr. IX, 22, 58.		
S. rugosa; R. Brown, prodr. 240 (1810)	...	—	—	—	—	—	Q. N.A.	B.f. VII, 428	M.fr. IX, 22, 58.		
S. laxa; R. Brown, prodr. 240 (1810)	...	—	—	—	—	—	Q. N.A.	B.f. VII, 428	M.fr. IX, 21.		
S. Brownii; Kunth, enum. II, 349 (1837)	...	—	—	—	—	—	Q. N.A.	B.f. VII, 429	M.fr. IX, 21, 58.		
S. lithosperma; Willdenow, spec. plant. IV, 316 (1805)	...	—	—	—	—	—	Q. N.A.	B.f. VII, 429	M.fr. IX, 21, 58	As., P.	
S. tessellata; Willdenow, spec. plant. IV, 315 (1805)	...	—	—	—	—	—	Q.	—	B.f. VII, 430	...	As.
S. margaritifera; Willdenow, spec. plant. IV, 312 (1805)	...	—	—	—	—	—	Q.	—	B.f. VII, 430	M.fr. IX, 21, 58	As., P.
S. Graeffeana; Boeckeler in Regensb. Flora 121 (1875)	...	—	—	—	—	—	Q. N.A.	B.f. VII, 431			
S. hebecarpa; Nees in Wight, Contrib. to Bot. of Ind. 117 (1834)	...	—	—	—	—	N.S.W.	Q. N.A.	B.f. VII, 431	M.fr. IX, 21, 58	As.	
S. Chinensis; Kunth, enum. II, 357 (1837)	...	—	—	—	—	—	Q.	—	B.f. VII, 431	M.fr. IX, 20, 58	As.
S. oryzoides; Presl, reliq. Haenk. I, 201 (1830)	...	—	—	—	—	—	—	N.A.	B.f. VII, 432	...	As., P.
S. sphacelata; F. v. M., fragm. IX, 20 (1875)	...	—	—	—	—	N.S.W.	Q.	—	B.f. VII, 432	M.fr. IX, 20.	

UNCINIA; Persoon, synops. plant. I, 534 (1807).

U. tenella; R. Brown, prodr. 241 (1810)	...	—	—	T.	V.	N.S.W.	—	—	B.f. VII, 433	M.fr. VIII, 151.	
U. compacta; R. Brown, prodr. 241 (1810)	...	—	—	T.	V.	—	—	—	B.f. VII, 434	M.fr. VIII, 152.	
U. riparia; R. Brown, prodr. 241 (1810)	...	—	—	T.	V.	N.S.W.	—	—	B.f. VII, 434	M.fr. VIII, 152.	
U. debilior; F. v. M., fragm. VIII, 151 (1874)	...	—	—	—	—	N.S.W.	—	—	B.f. VII, 435	M.fr. VIII, 151.	

CAREX; Ruppius, Fl. Jencns. 306 (1718).

C. cephalotes; F. v. M. in Transact. phil. Soc. Vict. I, 110 (1854)	...	—	—	—	V.	N.S.W.	—	—	B.f. VII, 437	M.fr. VIII, 251.	
C. acicularis; Boott in J. Hook. Fl. N. Zel. I, 280, t. 63 (1853)	...	—	—	T.	V.	N.S.W.	—	—	B.f. VII, 437	M.fr. VIII, 251	N.Z.
C. capillacea; Boott, Illustr. of the gen. Carex I, 44, t. 110 (1858)	...	—	—	—	—	N.S.W.	—	—	B.f. VII, 438	M.fr. IX, 191	As.
C. inversa; R. Brown, prodr. 242 (1810)	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.f. VII, 438	M.fr. VIII, 252; XII, 26.	N.Z.

<i>P. gibbosum</i> ; R. Brown, prodr. 193 (1810)	— — — — —	Q. N.A. B.fl. VII, 471	M.fr. VIII, 155; XI, 129.	
<i>P. leucophaeum</i> ; Humboldt & Kuuth, nov. gen. et spec. I, 97 (1815)	— S.A. — V. N.S.W.	Q. N.A. B.fl. VII, 472	M.fr. VIII, 155	... Afr., Am.
<i>P. semialatum</i> ; R. Brown, prodr. 192 (1810)	— — — — —	N.S.W. Q. N.A. B.fl. VII, 472	M.fr. VIII, 196	... As.
<i>P. rarum</i> ; R. Brown, prodr. 189 (1810)	— — — — —	— N.A. B.fl. VII, 473		
<i>P. argenteum</i> ; R. Brown, prodr. 190 (1810)	— S.A. — — —	— N.A. B.fl. VII, 473		
<i>P. holosericeum</i> ; R. Brown, prodr. 190 (1810)	— — — — —	Q. N.A. B.fl. VII, 473	M.fr. VII, 155.	
<i>P. flavidum</i> ; Retzius, observ. IV, 15 (1787)	— — — — —	N.S.W. Q. N.A. B.fl. VII, 474	M.fr. VIII, 189	... As.
<i>P. gracile</i> ; R. Brown, prodr. 190 (1810)	W.A. S.A. — V. N.S.W.	Q. N.A. B.fl. VII, 475	M.fr. VIII, 189.	
<i>P. tubulatum</i> ; Haeckel in Engler's Jahrb. VI, 234 (1885)	— — — — —	— N.A. —		
<i>P. prostratum</i> ; Lamarck, illustr. I, 171 (1791)	— — — — —	Q. N.A. B.fl. VII, 476	M.fr. VIII, 191	... As., Afr., Am.
<i>P. helopus</i> ; Trinius in Spreng. Neue Entd. II, 84 (1821)	— S.A. — — N.S.W.	Q. N.A. B.fl. VII, 476		... As., Afr.
<i>P. Gilesii</i> ; Benth. Fl. Austr. VII, 477 (1878)	— — S.A. — — —	Q. N.A. B.fl. VII, 477	M.fr. XI, 129.	
<i>P. piligerum</i> ; F. v. M. in Benth. Fl. Austr. VII, 477 (1878)	— — — — —	— N.A. B.fl. VII, 477		
<i>P. polyphyllum</i> ; R. Brown, prodr. 190 (1810)	— — — — —	— N.A. B.fl. VII, 477	M.fr. VIII, 194.	
<i>P. distachyon</i> ; Linné, mantiss. alt. pl. 138 (1771)	— — S.A. — — N.S.W.	Q. N.A. B.fl. VII, 478	M.fr. VIII, 194	... As.
<i>P. reversum</i> ; F. v. M., fragm. VIII, 152 (1874)	W.A. S.A. — — N.S.W.	Q. — B.fl. VII, 478	M.fr. X, 76.	
<i>P. colonum</i> ; Linné, syst. nat. ed. decim. 5 (1758)	— — — — —	— N.A. B.fl. VII, 478		... E., As., Afr., Am.
<i>P. Crus Galli</i> ; Linné, spec. plant. 56 (1753)	W.A. S.A. — V. N.S.W.	Q. N.A. B.fl. VII, 479	M.fr. VIII, 198	... As., Afr., Am.
<i>P. myosuroides</i> ; R. Brown, prodr. 189 (1810)	— — — — —	Q. N.A. B.fl. VII, 480		... As., Afr.
<i>P. Indicum</i> ; Linné, mantiss. pl. alt. 184 (1771)	— — — — —	N.S.W. Q. N.A. B.fl. VII, 480	M.fr. VIII, 197; XI, 129.	As., Afr.
<i>P. Myurus</i> ; Lamarck, illustr. I, 172 (1791)	— — — — —	Q. — B.fl. VII, 480	M.fr. X, 114	... As., Afr., Am.
<i>P. foliosum</i> ; R. Brown, prodr. 191 (1810)	— — — — —	N.S.W. Q. — B.fl. VII, 481	M.fr. VIII, 194.	
<i>P. adpersum</i> ; Trinius, spec. gram. t. 169 (1830)	— — S.A. — — N.S.W.	Q. — B.fl. VII, 481		... Am.
<i>P. inaequale</i> ; F. v. M., fragm. VIII, 189 (1874)	— — — — —	Q. — B.fl. VII, 482	M.fr. XI, 129.	
<i>P. uncinulatum</i> ; R. Brown, prodr. 191 (1810)	— — — — —	N.S.W. Q. — B.fl. VII, 482	M.fr. VIII, 195; X, 76; XI, 129.	
<i>P. majusculum</i> ; F. v. M. in Benth. Fl. Austr. VII, 482 (1878)	— — — — —	— N.A. B.fl. VII, 482		
<i>P. pauciflorum</i> ; R. Brown, prodr. 483 (1810)	W.A. S.A. — — —	Q. N.A. B.fl. VII, 483	M.fr. VIII, 193.	
<i>P. semitonsum</i> ; F. v. M. in Benth. Fl. Austr. VII, 483 (1878)	— — — — —	— N.A. B.fl. VII, 483		
<i>P. antidotale</i> ; Retzius, observ. bot. IV, 17 (1787)	— — — — —	— N.A. B.fl. VII, 483		
<i>P. repens</i> ; Linné, spec. pl. ed. sec. 87 (1762)	— — — — —	V. N.S.W. Q. N.A. B.fl. VII, 484		... E., As., Afr., Am., P.
<i>P. Buncei</i> ; F. v. M. in Benth. Fl. Austr. VII, 487 (1878)	— — — — —	Q. — B.fl. VII, 487		
<i>P. capillipes</i> ; Benth. Fl. Austr. VII, 484 (1878)	— — — — —	— N.A. B.fl. VII, 484	M.fr. VIII, 193.	
<i>P. pygmaeum</i> ; R. Brown, prodr. 191 (1810)	— — — — —	N.S.W. Q. — B.fl. VII, 484	M.fr. VIII, 193.	
<i>P. brevifolium</i> ; Linné, spec. plant. 59 (1753)	— — — — —	— Q. N.A. B.fl. VII, 485	M.fr. VIII, 193; XI, 129.	
<i>P. hermaphroditum</i> ; Stendel, syn. glum. I, 67 (1855)	— — — — —	Q. — B.fl. VII, 485		... As.
<i>P. marginatum</i> ; R. Brown, prodr. 190 (1810)	— — — — —	V. N.S.W. Q. — B.fl. VII, 485	M.fr. VIII, 190.	
<i>P. lachnophyllum</i> ; Benth. Fl. Austr. VII, 486 (1878)	— — — — —	Q. — B.fl. VII, 486		
<i>P. obseptum</i> ; Trinius, gram. panic. diss. alt. 149 (1826)	— — — — —	N.S.W. — B.fl. VII, 486		
<i>P. bicolor</i> ; R. Brown, prodr. 191 (1810)	— — — — —	N.S.W. Q. — B.fl. VII, 487		
<i>P. melanthum</i> ; F. v. M. in Transact. Vict. Inst. 47 (1854)	— — — — —	V. N.S.W. Q. — B.fl. VII, 488	M.fr. VIII, 192.	
<i>P. effusum</i> ; R. Brown, prodr. 191 (1810)	W.A. S.A. — V. N.S.W.	Q. N.A. B.fl. VII, 488	M.fr. VIII, 191.	
<i>P. Mitchellii</i> ; Benth. Fl. Austr. VII, 489 (1878)	— S.A. — V. N.S.W.	Q. N.A. B.fl. VII, 489	M.fr. X, 76.	
<i>P. decompositum</i> ; R. Brown, prodr. 191 (1810)	W.A. S.A. — V. N.S.W.	Q. N.A. B.fl. VII, 489	M.fr. VI, 86; VIII, 191.	As.
<i>P. trachyrhachis</i> ; Benth. Fl. Austr. VII, 490 (1878)	— — — — —	N.S.W. Q. N.A. B.fl. VII, 490	M.fr. VIII, 192; XI, 129.	
<i>P. prolutum</i> ; F. v. M. in Transact. Vict. Inst. 46 (1856)	— S.A. — V. N.S.W.	Q. — B.fl. VII, 490	M.fr. VIII, 192.	
<i>P. spinescens</i> ; R. Brown, prodr. 193 (1810)	W.A. S.A. — V. N.S.W.	Q. N.A. B.fl. VII, 499	M.fr. VIII, 197; XI, 129.	As.
<i>P. paradoxum</i> ; R. Brown, prodr. 193 (1810)	— — — — —	V. N.S.W. Q. — B.fl. VII, 499	M.fr. VIII, 197.	
<i>P. Chamaeraphis</i> ; Trinius in Mem. acad. Petersb. ser. 6, III, 217 (1835)	— — — — —	— N.A. B.fl. VII, 500		
<i>P. atro-virens</i> ; Trinius in Spreng. neue Entdeckung. II, 89 (1821)	— — — — —	V. N.S.W. Q. — B.fl. VII, 625	M.fr. VIII, 193; XI, 129.	As.
<i>P. Myosotis</i> ; Steudel, syn. glum. I, 96 (1855)	— — — — —	Q. — B.fl. VII, 625	M.fr. VIII, 193	... As.

OPLISMENUS; Palisot, Flore d'Oware et de Benin. II, 14 (1807). (Orthopogon.)

O. compositus; Palisot, Agrostogr. 54 (1812) ... — — — V. N.S.W. Q. — B.fl.VII,491 M.fr.VIII,199 ... As., P., N.Z.

SETARIA; Palisot, Essai d'une nouv. Agrostogr. 51, t. 13 (1812).

S. glauca; Palisot, Agrostogr. 51, t. 13 (1812) ... W.A. S.A. — V. N.S.W. Q. N.A. B.fl.VII,492 M.fr.VIII,110. E., As., Afr., Am., P.
S. macrostachya; Humboldt & Kunth, nov. gen. et spec. I, 110 (1815) ... W.A. S.A. — — N.S.W. Q. N.A. B.fl.VII,493 M.fr.VIII,110; XI,129. As., Am.
S. viridis; Palisot, Agrostogr. 51 (1812) ... W.A. S.A. — — — N.A. B.fl.VII,494 ... E., As., Afr.
S. verticillata; Palisot, Agrostogr. 51 (1812) ... W.A. — — — — N.A. B.fl.VII,494 M.fr.VIII,110 ... E., As., Afr.

PENNISETUM; L. C. Richard in Persoon, synops. plant. I, 72 (1805). (Gymnothrix, Plagiosetum.)

P. compressum; R. Brown, prodr. 195 (1810) ... — — — — N.S.W. Q. — B.fl.VII,495 M.fr.VIII,109. ... As., P.
P. Arnemicum; F. v. M., fragm. VIII, 109 (1874) ... — — — — — N.A. B.fl.VII,496 M.fr.VIII,109.
P. refractum; F. v. M., fragm. VIII, 109 (1874) ... — S.A. — — — Q. N.A. B.fl.VII,495 M.fr.III,147; IX,194.

CENCHRUS; Linné, coroll. gen. 20 (1737).

C. Australis; R. Brown, prodr. 196 (1810) ... — — — — N.S.W. Q. — B.fl.VII,497 M.fr.VIII,107; XI,129.
C. inflexus; R. Brown, prodr. 195 (1810) ... — — — — — N.A. B.fl.VII,497 M.fr.VIII,107.
C. elymoides; F. v. M., fragm. VIII, 107 (1874) ... — — — — — Q. N.A. B.fl.VII,497 M.fr.VIII,107.

XEROCHLOA; R. Brown, prodr. fl. Nov. Holl. 196 (1810).

X. imberbis; R. Brown, prodr. 197 (1810) ... — — — — — Q. N.A. B.fl.VII,501 M.fr.VIII,117.
X. barbata; R. Brown, prodr. 197 (1810) ... — — — — — Q. N.A. B.fl.VII,501 M.fr.VIII,117.
X. laniflora; Benth, Fl. Austr. VII, 502 (1878) ... — — — — — N.A. B.fl.VII,502

STENOTAPHRUM; Trinius, fundam. Agrostogr. 175 (1820). (Diastemanthe.)

S. dimidiatum; Trinius, fund. Agrost. 175 (1820) ... — — — — — Q. N.A. B.fl.VII,500 M.fr.VIII,156 ... As., Afr., Am.

THUAREA; Persoon, synops. plant. I, 110 (1805). (Thouarsea.)

T. sarmentosa; Persoon, synops. I, 110 (1805) ... — — — — — Q. N.A. B.fl.VII,502 M.fr.VI,86 ... As., Afr., P.

SPINIFEX; Linné, syst. nat. ed. XII, 757 (1767).

S. hirsutus; Labillardière, Nov. Holl. pl. spec. II, 81, t. 230 (1806) ... W.A. S.A. T. V. N.S.W. Q. — B.fl.VII,503 M.fr.VIII,133 ... N.Z., P.
S. longifolius; R. Brown, prodr. 198 (1810) ... W.A. — — — — Q. N.A. B.fl.VII,504 M.fr.VIII,139.
S. paradoxus; Benth, in Hook. icon. plant. t. 1243 (1877) ... — S.A. — V. N.S.W. Q. N.A. B.fl.VII,504 M.fr.VI,86; VIII,117,199; IX,195.

LEPTASPIS; R. Brown, prodr. fl. Nov. Holl. 211 (1810).

L. Banksii; R. Brown, prodr. 211 (1810) ... — — — — — Q. — B.fl.VII,548 M.fr.VIII,116. ... P.

PEROTIS; Aiton, hort. Kew. I, 85 (1789). (Xystidium.)

P. rara; R. Brown, prodr. 172 (1810) ... — S.A. — — N.S.W. Q. N.A. B.fl.VII,509 M.fr.VIII,115 ... As., P.

POLYPOGON; Desfontaines, flor. Atlantic. I, 66 (1798).

P. fugax; Nees in Steudel, syn. glum. I, 184 (1855) ... W.A. — — — — — B.fl.VII,547 ... As.
P. tenellus; R. Brown, prodr. 173 (1810) ... W.A. — — — — — B.fl.VII,547 M.fr.VIII,114.

ARUNDINELLA; Raddi, Agrostogr. Brasil. 37 (1823).

A. Nepalensis; Trinius, spec. gram. t. 268 (1833) ... — — — — N.S.W. Q. N.A. B.fl.VII,545 M.fr.VIII,139 ... As., Afr.
A. Schultzii; Benth, Fl. Austr. VII, 545 (1878) ... — — — — — N.A. B.fl.VII,545

TRAGUS; Haller, hist. stirp. Helvet. n. 1413 (1768). (Lappago.)

T. racemosus; Haller, stirp. Helv. n. 1413 (1768) ... — S.A. — V. N.S.W. Q. N.A. B.fl.VII,507 M.fr.VIII,107 ... E., As., Afr., Am.

NEURACHNE; R. Brown, prodr. fl. Nov. Holl. 196 (1810).

N. alopecuroides; R. Brown, prodr. 196 (1810) ... W.A. S.A. — V. N.S.W. — — B.fl.VII,507 M.fr.VI,86; VIII,200; XI,129.
N. Mitchelliana; Nees in Hook. Lond. Journ. of Bot. II, 410 (1843) ... — S.A. — V. N.S.W. Q. — B.fl.VII,508 M.fr.VI,86; VIII,200.
N. Munroi; F. v. M., fragm. VIII, 200 (1874) ... — S.A. — V. N.S.W. — — B.fl.VII,508 M.fr.V,204; XI,129.

ZOYSIA; Willdenow, Neue Schrift. nat. Freunde zu Berl. III, 440 (1801).

Z. pungens; Willdenow, Neue Schrift. nat. Fr. Berl. III, 440 (1801) ... — — T. V. N.S.W. Q. — B.fl.VII,506 M.fr.VIII,116 ... As., N.Z.

IMPERATA; Cyrillo, plant. rar. Neap. II, t. 11 (1792).

I. arundinacea; Cyrillo, plant. rar. Neap. II, t. 11 (1792)... ... W.A. S.A. T. V. N.S.W. Q. N.A. B.fl.VII,536 M.fr.VIII,126 ... E., As., Afr., Am., P.

ERIANTHUS; L. C. Richard in Michaux, fl. bor. Amer. I, 54 (1803). (Pollinia, Enlalia, Pogonatherum partly, Leptatherum, Saccharum partly.)

E. articulatus; F. v. M., fragm. VIII, 118 (1874) ... — — — — — Q. N.A. B.fl.VII,525 M.fr.X,76 ... As.

E. irritans; Kunth, Graminées I, 160 (1832) ... — — — — — Q. N.A. B.fl.VII,526 M.fr.VIII,118.

E. fulvus; Kunth, Graminées I, 160 (1832) ... W.A. S.A. — V. N.S.W. Q. N.A. B.fl.VII,526 M.fr.VIII,118.

E. Roxburghii; F. v. M., fragm. VIII, 117 (1874) ... — — — — — Q. — B.fl.VII,527 M.fr.VIII,117 ... As.

E. Mackinlayi; F. v. M., Cens. 131 (1882) ... — — — — — — N.A. B.fl.VII,527 M.fr.VIII,118.

POGONATHERUM; Palisot, Agrostogr. 176 (1812).

P. saccharoideum; Palisot, Agrostogr. 56, t. 11 (1812) ... — — — — — Q. — — — — — As.

DIMERIA; R. Brown, prodr. fl. Nov. Holl. 204 (1810).

D. acinaciformis; R. Brown, prodr. 204 (1810) ... — — — — — Q. — B.fl.VII,523

D. ornithopoda; Trinius, fundam. agrostogr. 167 (1820) ... — — — — — Q. N.A. B.fl.VII,523 M.fr.VII,104; XI,121. As.

ARTHRAON; Palisot, Agrostogr. III, t. 11, f. 6 (1812). (Batratherum.)

A. ciliaris; Palisot, Agrostogr. III, t. 11 (1812) ... — — — — — N.S.W. Q. — B.fl.VII,524 M.fr.VIII,119; X,76. As.

ELIONURUS; Humboldt & Bonpland in Willd. sp. pl. IV, 941 (1805).

E. citreus; Munro in Benth. Fl. Austr. VII, 510 (1878) ... — — — — — Q. — B.fl.VII,510 M.fr.XI,129.

LEPTURUS; R. Brown, prodr. fl. Nov. Holl. 207 (1810). (Monerma.)

L. incurvatus; Trinius, fundam. Agrostogr. 123 (1820) ... — S.A. — V. N.S.W. — — B.fl.VII,668 M.fr.VIII,117 ... E., As., Afr., N.Z.

L. cylindricus; Trinius, fundam. Agrostogr. 123 (1820) ... W.A. S.A. — V. N.S.W. — — B.fl.VII,668 ... E., As., Afr.

L. repens; R. Brown, prodr. 207 (1810) ... — — — — — Q. — B.fl.VII,668 M.fr.VI,86 ... As., P.

HEMARTHRIA; R. Brown, prodr. fl. Nov. Holl. 207 (1810). (Rottboelliae subgenus.)

H. compressa; R. Brown, prodr. 207 (1810) ... W.A. S.A. T. V. N.S.W. Q. — B.fl.VII,511 M.fr.XI,129.

ROTTBOELLIA; Linné, fil. gram. gen. 22 (1779). (Coelorrhachis.)

R. formosa; R. Brown, prodr. 206 (1810) ... — — — — — Q. N.A. B.fl.VII,513 M.fr.VI,85; XI,129.

R. exaltata; Linné, fil. suppl. 114 (1781) ... — — — — — Q. N.A. B.fl.VII,513 ... As.

R. ophiuroides; Benth. Fl. Austr. VII, 514 (1878) ... — — — — — Q. N.A. B.fl.VII,514 M.fr.VIII,123 ... As., P.

OPHIUROS; K. F. Gaertner, de fruct. III, 3, t. 181 (1805). (Ophiurus.)

O. corymbosus; K. F. Gaertner, de fruct. III, 3, t. 181 (1805) ... — — — — — Q. N.A. B.fl.VII,512 M.fr.VI,85 ... As., Afr.

MANISURIS; Linné, syst. nat. ed. duodec. 762 (1767).

M. granularis; Swartz, Fl. Ind. occid. I, 186 (1797) ... — — — — — Q. N.A. B.fl.VII,514 M.fr.VII,116; XI,129. As., Afr., Am.

ISCHAEMUM; Linné, syst. nat. 8 (1735); Linné, gen. pl. ed. sec. 525 (1742). (Hologamium, Spodiopogon partly.)

I. truncatiglume; F. v. M. in Benth. Fl. Austr. VII, 518 (1878) ... — — — — — — N.A. B.fl.VII,518

I. arundinaceum; F. v. M. in Benth. Fl. Austr. VII, 519 (1878) ... — — — — — — N.A. B.fl.VII,519

I. triticeum; R. Brown, prodr. 205 (1810) ... — — — — — N.S.W. Q. — B.fl.VII,519

I. Australe; R. Brown, prodr. 205 (1810) ... — — — — — N.S.W. Q. N.A. B.fl.VII,519

I. muticum; Linné, spec. plant. 1049 (1753) ... — — — — — Q. — B.fl.VII,520 M.fr.VIII,120 ... As., P.

I. ciliare; Retzius, observ. bot. VI, 36 (1791) ... — — — — — N.S.W. — B.fl.VII,520 M.fr.VIII,119 ... As.

I. decumbens; Benth. Fl. Austr. VII, 521 (1878) ... — — — — — — N.A. B.fl.VII,521

I. pectinatum; Trinius in Mem. de l'Acad. de Petersb. six. sér. II, 296 (1833)... — — — — — N.S.W. Q. N.A. B.fl.VII,521 M.fr.VIII,118 ... As.

I. fragile; R. Brown, prodr. 205 (1810) ... — — — — — Q. — B.fl.VII,522 M.fr.VIII,123; XI,130.

I. laxum; R. Brown, prodr. 205 (1810) ... — — — — — Q. N.A. B.fl.VII,522 M.fr.VIII,119 ... As., Afr.

ANDROPOGON ; Royen. prodr. exhib. pl. hort. acad. Lugd. 52 (1740). (Sorghum, Chrysopogon, Heteropogon, Schizachyrium, Trachypogon, Holcus partly, Spodiopogon partly.)

A. erianthoides ; F. v. M., fragm. X, 75 (1876)	...	...	...	...	...	—	—	—	N.S.W.	Q.	—	B.fl. VII, 529	M.fr. X, 75.			
A. sericens ; R. Brown, prodr. 201 (1810)	...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl. VII, 529	...	...	As., P.
A. affinis ; R. Brown, prodr. 201 (1810)	...	...	...	...	...	—	—	—	V.	N.S.W.	Q.	—	B.fl. VII, 530			
A. pertusus ; Willdenow, spec. plant. IV, 922 (1805)	...	...	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl. VII, 530	M.fr. VIII, 122; X, 76.		As.
A. annulatus ; Forskael, Fl. Aegypt. Arab. 173 (1775)	...	...	...	...	...	—	S.A.	—	V.	—	Q.	N.A.	B.fl. VII, 531	M.fr. VIII, 123	...	As., Afr.
A. Ischaemum ; Linné, spec. plant. 1047 (1753)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl. VII, 531	M.fr. VIII, 122	...	E., As., Afr.
A. punctatus ; Roxburgh, fl. Indic. I, 268 (1820)	...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl. VII, 531	..	...	As.
A. procerus ; R. Brown, prodr. 202 (1810)	...	...	...	...	...	W.A.	—	—	—	—	—	N.A.	B.fl. VII, 532	M.fr. VIII, 124.		
A. exaltatus ; R. Brown, prodr. 202 (1810)	...	...	...	...	...	W.A.	S.A.	—	—	—	—	N.A.	B.fl. VII, 532	M.fr. VIII, 124.		
A. bombycinus ; R. Brown, prodr. 202 (1810)	...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl. VII, 533	M.fr. XI, 20.		
A. Schoenanthus ; Linné, mater. med. 31 (1749)	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl. VII, 534	M.fr. VIII, 124	...	As., P.
A. refractus ; R. Brown, prodr. 202 (1810)	...	...	...	...	...	—	—	—	V.	N.S.W.	Q.	N.A.	B.fl. VII, 534	M.fr. VIII, 124; X, 76.		P.
A. filipendulinus ; Hochstetter in Regensb. Fl. 175 (1846)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl. VII, 534	...	...	Afr.
A. brevifolius ; Swartz, fl. Ind. occid. I, 209 (1797)	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl. VII, 535			
A. contortus ; Linné, spec. plant. 1045 (1753)	...	...	...	...	...	W.A.	—	—	—	N.S.W.	Q.	N.A.	B.fl. VII, 517	M.fr. VIII, 120	...	As., Afr., Am., P.
A. triticeus ; R. Brown, prodr. 201 (1810)	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl. VII, 517	M.fr. VIII, 120	...	As.
A. Gryllus ; Linné, amoen. acad. IV, 332 (1759)	...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl. VII, 537	M.fr. VIII, 121; XI, 130.		As., Afr., P.
A. micranthus ; Kunth, gram. I, 165 (1835)	...	...	...	...	...	—	—	—	V.	N.S.W.	Q.	N.A.	B.fl. VII, 538	M.fr. VIII, 122; X, 76; XI, 130.		As., P.
A. aciculatus ; Retzius, observ. bot. V, 22 (1789)	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl. VII, 538	M.fr. VIII, 122	...	As., P.
A. elongatus ; Sprengel, syst. I, 289 (1825)	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl. VII, 539	M.fr. VIII, 121.		
A. Halpensis ; Sibthorp & Smith, Fl. Graeca I, 52, t. 68 (1806)	...	...	...	...	...	W.A.	—	—	—	N.S.W.	Q.	—	B.fl. VII, 540	M.fr. VIII, 119	...	E., As., P.
A. Australis ; Sprengel, syst. I, 237 (1825)	...	...	...	...	...	W.A.	—	—	V.	N.S.W.	Q.	N.A.	B.fl. VII, 540			
A. serratus ; Thunberg, fl. Japon. 41 (1784)	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl. VII, 541	...	...	As.
A. intrans ; F. v. M., Cens. of Austr. pl. 132 (1882)	...	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl. VII, 541			

ANTHISTIRIA ; Linné, fil. gram. gen. 35 (1779). (Themeda 1775, Iscilema.)

A. ciliata ; Linné, fil. gram. gen. 35 (1779)	...	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl. VII, 542	M.fr. V, 207; VIII, 139.		As., Afr.
A. frondosa ; R. Brown, prodr. 200 (1810)	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl. VII, 542	M.fr. VIII, 139; XI, 130.		
A. avenacea ; F. v. M., fragm. V, 206 (1866)	...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl. VII, 543	M.fr. VI, 253; VIII, 139.		
A. membranacea ; Lindley in Mitch. Trop. Austr. 88 (1848)	...	...	...	...	...	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl. VII, 543	M.fr. V, 207; VIII, 139.		

APLUDA ; Linné, spec. plant. 82 (1753).

A. mutica ; Linné, spec. plant. 82 (1753)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl. VII, 544	M.fr. VIII, 140	...	As., Afr., P.
---	-----	-----	-----	-----	-----	---	---	---	---	--------	----	---	----------------	-----------------	-----	---------------

CHIONACHNE ; R. Brown in Horsfield's pl. Jav. rar. 15 (1838).

C. barbata ; R. Brown in Bennett, pl. Javan. rar. 18 (1838)	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl. VII, 515	...	...	As.
C. cyathopoda ; F. v. M. in Benth. Fl. Austr. VII, 516 (1878)	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl. VII, 515	M.fr. VIII, 116.		

ALOPECURUS ; Linné, syst. nat. 8 (1735); gen. plant. 18 (1737). (Perhaps immigrated.)

A. geniculatus ; Linné, spec. plant. 60 (1753)	...	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl. VII, 555	M.fr. VIII, 138	...	As., Am., N.Z.
--	-----	-----	-----	-----	-----	------	------	----	----	--------	----	---	----------------	-----------------	-----	----------------

LEERSIA ; Solander in Swartz, nov. gen. et spec. plant. 1 et 21 (1788). (Asprella, Asperella.)

L. hexandra ; Swartz, nov. gen. et spec. pl. 21 (1788)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl. VII, 549	M.fr. VIII, 115	...	E., As., Afr., Am.
--	-----	-----	-----	-----	-----	---	---	---	---	--------	----	---	----------------	-----------------	-----	--------------------

ORYZA ; Tournefort, inst. 513, t. 296 (1700), from Theophrastos & Dioscorides.

O. sativa ; Linné, spec. plant. 333 (1753)	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl. VII, 550	M.fr. VIII, 115; XI, 130.		As.
--	-----	-----	-----	-----	-----	---	---	---	---	---	----	------	----------------	---------------------------	--	-----

POTAMOPHILA ; R. Brown, prodr. fl. Nov. Holl. 211 (1810).

P. parviflora ; R. Brown, prodr. 211 (1810)	...	...	...	...	...	—	—	—	—	N.S.W.	—	—	B.fl. VII, 550	M.fr. VIII, 126.		
---	-----	-----	-----	-----	-----	---	---	---	---	--------	---	---	----------------	------------------	--	--

EHRHARTA ; Thunberg in Svensk. Vetensk. Acad. Handl. 216, t. 8 (1779). (Tetrarrhena, Microlaena, Diplax.)

E. distichophylla ; Labillardière, Nov. Holl. pl. spec. I, 90, t. 117 (1804)	...	...	...	...	...	—	—	T.	V.	—	—	—	B.fl. VII, 554	M.fr. VII, 90.		
E. juncea ; Sprengel, syst. II, 114 (1825)	...	...	...	...	...	—	—	T.	V.	N.S.W.	—	—	B.fl. VII, 554	M.fr. VII, 90.		

E. laevis ; Sprengel, syst. II, 115 (1825)	W.A.	—	—	—	—	—	B.fl. VII, 554	M.fr. VII, 90.	
E. acuminata ; Sprengel, syst. II, 114 (1825)	—	—	T.	V.	—	—	B.fl. VII, 555	M.fr. VII, 90; XI, 130.	
E. stipoides ; Labillardière, Nov. Holl. pl. spec. I, 91, t. 118 (1804)	W.A.	S.A.	T.	V.	N.S.W.	Q.	B.fl. VII, 552	M.fr. VII, 90	... N.Z.
E. Diarrhena ; F. v. M., fragm. VII, 89 (1870)	—	—	T.	—	—	—	B.fl. VII, 553	M.fr. VII, 89.	

HIEROCHLOE ; J. G. Gmelin, Fl. Sibiric. I, 100 (1747). (Hierochloa, Disarrhenum.)

H. redolens ; R. Brown, prodr. 209 (1810)	—	—	T.	V.	N.S.W.	—	B.fl. VII, 558	M.fr. VIII, 137	... Am., N.Z.
H. rariflora ; J. Hooker, Fl. Antarct. I, 93 (1844)	—	—	T.	V.	N.S.W.	—	B.fl. VII, 559	M.fr. VIII, 138.	

ARISTIDA ; Linuë, spec. plant. 82 (1753). (Chaetaria, Arthratherum.)

A. hygrometrica ; R. Brown, prodr. 174 (1810)	—	—	—	—	—	Q. N.A.	B.fl. VII, 561		
A. stipoides ; R. Brown, prodr. 174 (1810)	W.A.	S.A.	—	—	N.S.W.	Q. N.A.	B.fl. VII, 561	M.fr. VIII, 111.	
A. arenaria ; Gaudichaud in Freyc. voy. Bot. 407 (1826)	W.A.	S.A.	—	V.	N.S.W.	Q. N.A.	B.fl. VII, 561		
A. Behriana ; F. v. M. in Transact. Vict. Inst. 44 (1854)	—	S.A.	—	V.	N.S.W.	Q.	B.fl. VII, 562	M.fr. XI, 130.	
A. leptopoda ; Benthams, Fl. Austr. VII, 562 (1878)	—	S.A.	—	V.	N.S.W.	Q.	B.fl. VII, 562	M.fr. XI, 130.	
A. vagans ; Cavanilles, icon. et descr. pl. V, 45, t. 471 (1799)	—	—	—	V.	N.S.W.	Q.	B.fl. VII, 562	M.fr. VIII, 111.	
A. ramosa ; R. Brown, prodr. 173 (1810)	—	S.A.	—	—	N.S.W.	Q.	B.fl. VII, 563		
A. calycina ; R. Brown, prodr. 173 (1810)	—	S.A.	—	V.	N.S.W.	Q. N.A.	B.fl. VII, 563	M.fr. XI, 130.	
A. depressa ; Retzius, observ. bot. IV, 22 (1787)	—	—	—	—	N.S.W.	Q.	B.fl. VII, 563		... As., Afr.

STIPA ; Linné, spec. plant. 78 (1737). (Streptachne, Dichelachne, Urachne partly, Muehlenbergia partly.)

S. elegantissima ; Labillardière, Nov. Holl. pl. spec. I, 23, t. 29 (1804)	W.A.	S.A.	—	V.	N.S.W.	Q. N.A.	B.fl. VII, 565	M.fr. VIII, 103.	
S. Tuckeri ; F. v. M., fragm. XI, 128 (1881)	—	—	—	V.	N.S.W.	—	—	M.fr. XI, 129.	
S. flavescens ; Labillardière, Nov. Holl. pl. spec. I, 24, t. 30 (1804)	W.A.	S.A.	T.	V.	N.S.W.	—	B.fl. VII, 566	M.fr. VIII, 104.	
S. teretifolia ; Steudel, syn. glum. I, 128 (1855)	W.A.	S.A.	T.	V.	—	—	B.fl. VII, 567	M.fr. VIII, 104	... N.Z.
S. compressa ; R. Brown, prodr. 175 (1810)	W.A.	—	—	—	—	—	B.fl. VII, 567		
S. Drummondii ; Steudel, syn. glum. I, 128 (1855)	W.A.	—	—	—	—	—	B.fl. VII, 567		
S. pyenostachya ; Benthams, Fl. Austr. VII, 568 (1878)	W.A.	—	—	—	—	—	B.fl. VII, 568		
S. setacea ; R. Brown, prodr. 174 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	B.fl. VII, 568		
S. semibarbata ; R. Brown, prodr. 174 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	B.fl. VII, 568	M.fr. VIII, 104; XI, 129.	
S. hemipogon ; Benthams, Fl. Austr. VII, 569 (1878)	W.A.	—	—	—	—	—	B.fl. VII, 569	M.fr. VIII, 104.	
S. pubescens ; R. Brown, prodr. 174 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	B.fl. VII, 569	M.fr. VIII, 104.	
S. aristiglumis ; F. v. M. in Transact. Vict. Inst. 43 (1854)	—	S.A.	—	V.	N.S.W.	Q.	B.fl. VII, 570	M.fr. VIII, 103.	
S. Eriopus ; Benthams, Fl. Austr. VII, 570 (1878)	W.A.	—	—	—	—	—	B.fl. VII, 570		
S. trichophylla ; Benthams, Fl. Austr. VII, 570 (1878)	W.A.	—	—	—	—	—	B.fl. VII, 570		
S. scabra ; Lindley in Mitch. Trop. Austr. 31 (1848)	W.A.	S.A.	—	V.	N.S.W.	Q.	B.fl. VII, 570		
S. Streptachne ; F. v. M. in Journ. R.S. of N.S.W. 237 (1881)	—	—	—	—	—	Q.	B.fl. VII, 572		
S. micrantha ; Cavanilles, icon. et descr. pl. V, 42, t. 467 (1799)	W.A.	S.A.	T.	V.	N.S.W.	Q.	B.fl. VII, 574	M.fr. VIII, 105	... N.Z.
S. verticillata ; Nees in Spreng. syst. veg. cur. poster. 30 (1827)	—	—	—	—	N.S.W.	Q.	B.fl. VII, 566	M.fr. VIII, 105.	
S. Dichelachne ; Steudel, nomencl. botan. edit. sec. I, 502 (1841)	W.A.	S.A.	T.	V.	N.S.W.	Q.	B.fl. VII, 574	M.fr. VIII, 105; XI, 129.	N.Z.

PENTAPOGON ; R. Brown, prodr. fl. Nov. Holl. 173 (1810).

P. Billardieri ; R. Brown, prodr. 173 (1810)	—	S.A.	T.	V.	N.S.W.	—	B.fl. VII, 572	M.fr. VIII, 106.	
---	---	------	----	----	--------	---	----------------	------------------	--

ECHINOPOGON ; Palisot, Essai d'une nouv. Agrostogr. 42, t. 9, f. 5 (1812). (Cinna partly.)

E. ovatus ; Palisot, Agrostogr. 42, t. 9 (1812)	W.A.	S.A.	T.	V.	N.S.W.	Q.	B.fl. VII, 599	M.fr. VIII, 106	... N.Z.
--	------	------	----	----	--------	----	----------------	-----------------	----------

DIPLOPOGON ; R. Brown, prodr. fl. Nov. Holl. 176 (1810). (Dipogonia.)

D. setaceus ; R. Brown, prodr. 176 (1810)	W.A.	—	—	—	—	—	B.fl. VII, 573	M.fr. VI, 86; VIII, 106.	
--	------	---	---	---	---	---	----------------	--------------------------	--

AMPHIPOGON ; R. Brown, prodr. 175 (1810). (Aegopogon partly, Pentacraspedion, Gamelythrum.)

A. debilis ; R. Brown, prodr. 175 (1810)	W.A.	—	—	—	—	—	B.fl. VII, 597	M.fr. VIII, 201.	
A. strictus ; R. Brown, prodr. 175 (1810)	W.A.	S.A.	—	V.	N.S.W.	Q. N.A.	B.fl. VII, 597	M.fr. VIII, 201.	

A. laguroides ; R. Brown, prodr. 175 (1810)	W.A.	—	—	—	—	—	—	B.fl.VII,598	M.fr.VI,86; VIII,201.
A. cygnornm ; Nees in Lehm. pl. Preiss. II, 100 (1846)	W.A.	—	—	—	—	—	—	B.fl.VII,599	
A. turbinatus ; R. Brown, prodr. 175 (1810)	W.A.	—	—	—	—	—	—	B.fl.VII,599	M.fr.VI,86; VIII,201.

PAPPOPHORUM ; Schreber, gen. pl. II, 787 (1791). (Enneapogon.)

P. commune ; F. v. M., plants of Gregory's Exp. 10 (1859)...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,600	M.fr.VIII,200	... P.
--	------	------	---	----	--------	----	------	--------------	---------------	--------

SPOROBOLUS ; R. Brown, prodr. fl. Nov. Holl. 169 (1810). (Vilfa partly.)

S. Virginicus ; Humboldt & Kunth, gram. I, 67 (1832)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.VII,621	M.fr.VI,84	... As., Afr., Am.
S. Indicus ; R. Brown, prodr. 170 (1810)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VII,622	M.fr.VI,84; X,76.	As., Afr., Am., N.Z.
S. diander ; Palisot, Agrostogr. 26 (1812)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,622	M.fr.VI,84; VIII,140.	As.
S. pulchellus ; R. Brown, prodr. 170 (1810)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.VII,623	M.fr.VIII,140.	
S. Lindleyi ; Benth, Fl. Austr. VII, 623 (1878)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,623	M.fr.XI,130.	
S. actinocladus ; F. v. M., fragm. VIII, 140 (1874)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.VII,623	M.fr.VI,84; IX,194; XI,130.	

AGROSTIS ; Linné, syst. nat. 8 (1135) ; gen. plant. 19 (1737). (Deyeuxia, Didymochaeta, Trichodium, Bromidium, Lachnagrostis partly, Apera partly.)

A. Muelleri ; Benth, Fl. Austr. VII, 576 (1878)	—	—	—	V.	N.S.W.	—	—	B.fl.VII,576	M.fr.VI,86	... N.Z.
A. scabra ; Willdenow, spec. plant. I, 378 (1797)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,576		... Am.
A. venusta ; Trinius in Mém. de l'Acad. de Petersh. VI, 340 (1850)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,576		
A. aequata ; Nees in Hook. Lond. Journ. of Bot. II, 412 (1843)	—	—	T.	—	—	—	—	B.fl.VII,579		
A. Solandri ; F. v. M., Veget. of the Chath. Isl. 60 (1864)...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,579	M.fr.XI,27,130	... N.Z.
A. montana ; R. Brown, prodr. 171 (1810)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,581		
A. quadriseta ; R. Brown, prodr. 171 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,581		... N.Z.
A. cylindrica ; R. Brown, prodr. 171 (1810)	W.A.	—	—	—	—	—	—	B.fl.VII,582		
A. densa ; F. v. M., Cens. 133 (1882) from Benth, (1878)	—	S.A.	—	V.	—	—	—	B.fl.VII,582		
A. frigida ; F. v. M., first gen. rep. 20 (1853)...	—	—	T.	V.	N.S.W.	—	—	B.fl.VII,583		
A. rudis ; Roemer & Schultes, syst. veg. II, 360 (1817)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.VII,583	M.fr.VI,85.	
A. nivalis ; F. v. M. in Transact. Vict. Inst. 43 (1854)	—	—	—	V.	N.S.W.	—	—	B.fl.VII,584		
A. Gunniana ; F. v. M., Cens. 133 (1882) from Benth, (1878)	—	—	T.	—	—	—	—	B.fl.VII,584		
A. breviglumis ; F. v. M., Cens. 133 (1882) from Benth, (1878)...	—	—	—	—	N.S.W.	—	—	B.fl.VII,584		

COELACHNE ; R. Brown, prodr. fl. Nov. Holl. 187 (1810).

C. pulchella ; R. Brown, prodr. 187 (1810)	—	—	—	—	—	Q.	—	B.fl.VII,626		... As.
---	---	---	---	---	---	----	---	--------------	--------	---------

MICRAIRA ; F. v. M., fragm. V, 208 (1866).

M. subulifolia ; F. v. M., fragm. V, 208 (1866)	—	—	—	—	—	Q.	—	B.fl.VII,624	M.fr.V,208.	
--	---	---	---	---	---	----	---	--------------	-------------	--

AIRA ; Linné, gen. plant. 335 (1737). (Aera, Deschampsia.)

A. caespitosa ; Linné, spec. plant. 64 (1753)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,587	M.fr.VIII,138	... E., As., Am., N.Z.
--	---	------	----	----	--------	---	---	--------------	---------------	------------------------

TRisetum ; Persoon, synops. plant. I, 97 (1805). (Airae subgenus.)

T. subspicatum ; Palisot, Agrostogr. 88 (1812)	—	—	T.	V.	N.S.W.	—	—	B.fl.VII,588	M.fr.VIII,136	... E., As., Am., N.Z.
---	---	---	----	----	--------	---	---	--------------	---------------	------------------------

ERIACHNE ; R. Brown, prodr. fl. Nov. Holl. 183 (1810.)

E. stipacea ; F. v. M., fragm. V, 206 (1866)	—	—	—	—	—	Q.	N.A.	B.fl.VII,626	M.fr.VIII,137; X,76.	
E. Armitii ; F. v. M. in Benth. Fl. Austr. VII, 627 (1878)	—	—	—	—	—	Q.	N.A.	B.fl.VII,627		... As.
E. squarrosa ; R. Brown, prodr. 183 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.VII,628		
E. glauca ; R. Brown, prodr. 184 (1810)	W.A.	—	—	—	—	Q.	N.A.	B.fl.VII,628		
E. rara ; R. Brown, prodr. 183 (1810)	—	—	—	—	—	Q.	—	B.fl.VII,628		
E. agrostidea ; F. v. M., fragm. VII, 82 (1870)	—	—	—	—	—	—	N.A.	B.fl.VII,628	M.fr.VIII,137.	
E. ciliata ; R. Brown, prodr. 184 (1810)	—	—	—	—	—	—	N.A.	B.fl.VII,629		
E. setacea ; Benth, Fl. Austr. VII, 629 (1878)	—	—	—	—	—	—	N.A.	B.fl.VII,629		
E. avenacea ; R. Brown, prodr. 184 (1810)	—	—	—	—	—	—	N.A.	B.fl.VII,629		

E. aristidea; F. v. M., fragm. V, 205 (1866) ...	...	...	...	...	...	W.A. S.A. — —	N.S.W. Q. N.A. B.fl.VII,629	M.fr.V,205.		
E. pallescens; R. Brown, prodr. 184 (1810) ...	...	...	...	...	...	— — — —	Q. — B.fl.VII,630	...	...	As.
E. festucacea; F. v. M., fragm. V, 205 (1866) ...	...	...	...	...	...	— — — —	— N.A. B.fl.VII,630	M.fr.V,205.		
E. ovata; Nees in Hook. Lond. Journ. II, 418 (1843) ...	...	...	...	...	...	W.A. S.A. — —	— N.A. B.fl.VII,630	M.fr.VIII,137.		
E. meliacea; F. v. M., fragm. V, 205 (1866) ...	...	...	...	...	...	— — — —	— N.A. B.fl.VII,631	M.fr.V,205.		
E. pallida; F. v. M. in Benth. Fl. Austr. VII, 631 (1878) ...	...	...	...	...	...	— S.A. — —	N.S.W. Q. N.A. B.fl.VII,631			
E. scleranthoides; F. v. M., fragm. VIII, 233 (1874) ...	...	...	...	...	...	— S.A. — —	— — B.fl.VII,631			
E. mucronata; R. Brown, prodr. 184 (1810) ...	...	...	...	...	...	— S.A. — —	N.S.W. Q. — B.fl.VII,632			
E. obtusa; R. Brown, prodr. 184 (1810) ...	...	...	...	...	...	W.A. S.A. — —	N.S.W. Q. N.A. B.fl.VII,632	M.fr.VIII,137; XI,130.		
E. capillaris; R. Brown, prodr. 184 (1810) ...	...	...	...	...	...	— — — —	— — N.A. B.fl.VII,632			

ANISOPOGON; R. Brown, prodr. fl. Nov. Holl. 176 (1810).

A. avenaceus; R. Brown, prodr. 176 (1810) ...	...	...	...	...	...	— — — —	V. N.S.W. Q. — B.fl.VII,590	M.fr.VII,106.		
---	-----	-----	-----	-----	-----	---------	-----------------------------	---------------	--	--

DANTHONIA; De Candolle, Fl. Franç. III, 32 (1805). (Amphibromus, Monacather, Plinthacanthus, Danthoinia.)

D. paradoxa; R. Brown, prodr. 177 (1810) ...	...	...	...	...	...	— — — —	N.S.W. — — B.fl.VII,591			
D. bipartita; F. v. M., fragm. I, 160 (1859) ...	...	...	...	...	...	W.A. S.A. — V.	N.S.W. Q. — B.fl.VII,592	M.fr.VIII,136.		
D. carphoides; F. v. M. in Benth. Fl. Austr. VII, 592 (1878) ...	...	...	...	...	...	— S.A. — V.	N.S.W. — — B.fl.VII,592	M.fr.XI,130.		
D. penicillata; F. v. M., fragm. VIII, 135 (1873) ...	...	...	...	...	...	W.A. S.A. T. V.	N.S.W. Q. — B.fl.VII,592	M.fr.VIII,135	...	P., N.Z.
D. robusta; F. v. M. in Transact. Vict. Inst. 44 (1854) ...	...	...	...	...	...	— — — —	V. N.S.W. — — B.fl.VII,593	M.fr.VIII,136.		
D. pauciflora; R. Brown, prodr. 177 (1810) ...	...	...	...	...	...	— — — —	T. V. N.S.W. — — B.fl.VII,596	...	...	N.Z.
D. nervosa; J. Hooker, Fl. Tasman. II, 121, t. 163 (1860) ...	...	...	...	...	...	W.A. S.A. T. V.	N.S.W. — — B.fl.VII,589	M.fr.VIII,135.		

ASTREBLA; F. v. M. fragm. phytogr. Austr. X, 76 (1876).

A. pectinata; F. v. M., fragm. X, 76 (1876) ...	...	...	...	...	...	W.A. S.A. — —	N.S.W. Q. N.A. B.fl.VII,602	M.fr.VIII,134.		
A. triticoides; F. v. M., fragm. X, 76 (1876) ...	...	...	...	...	...	— S.A. — —	N.S.W. Q. N.A. B.fl.VII,602	M.fr.VIII,134.		

MICROCHLOA; R. Brown, prodr. fl. Nov. Holl. 208 (1810).

M. setacea; R. Brown, prodr. 208 (1810) ...	...	...	...	...	...	— — — —	— — N.A. B.fl.VII,608	...	...	As., Afr., Am.
---	-----	-----	-----	-----	-----	---------	-----------------------	-----	-----	----------------

CYNODON; L. C. Richard in Persoon, synops. I, 85 (1805). (Fibrichia 1802.)

C. Dactylon; L. C. Richard in Persoon, synops. I, 85 (1805) ...	...	...	...	...	...	W.A. S.A. — V.	N.S.W. Q. — B.fl.VII,609	M.fr.VIII,113	...	E., As., Afr.
C. tenellus; R. Brown, prodr. 187 (1810) ...	...	...	...	...	...	— — — —	— Q. N.A. B.fl.VII,609	M.fr.VIII,113	...	As.
C. convergens; F. v. M., fragm. VIII, 113 (1874) ...	...	...	...	...	...	— — — —	— Q. N.A. B.fl.VII,610	M.fr.X,76.		
C. ciliaris; Benth. Fl. Austr. VII, 610 (1878) ...	...	...	...	...	...	— S.A. — —	— — B.fl.VII,610			

CHLORIS; Swartz, nov. gen. et spec. plant. 25 (1788).

C. unispicea; F. v. M., fragm. VII, 118 (1871) ...	...	...	...	...	...	— — — —	— Q. — B.fl.VII,611	M.fr.VII,118.		
C. pumilio; R. Brown, prodr. 186 (1810) ...	...	...	...	...	...	— — — —	— Q. N.A. B.fl.VII,611			
C. pectinata; Benth. Fl. Austr. VII, 612 (1878) ...	...	...	...	...	...	— S.A. — —	— Q. N.A. B.fl.VII,612			
C. divaricata; R. Brown, prodr. 186 (1810) ...	...	...	...	...	...	— — — —	— Q. N.A. B.fl.VII,612	M.fr.VI,85.		
C. acicularis; Lindley in Mitch. Trop. Austr. 33 (1848) ...	...	...	...	...	...	W.A. S.A. — V.	N.S.W. Q. — B.fl.VII,612	M.fr.VI,85.		
C. truncata; R. Brown, prodr. 186 (1810) ...	...	...	...	...	...	— S.A. — V.	N.S.W. Q. — B.fl.VII,612	M.fr.VI,85; X,130.		
C. ventricosa; R. Brown, prodr. 186 (1810) ...	...	...	...	...	...	— — — —	N.S.W. Q. — B.fl.VII,613	M.fr.VI,85.		
C. barbata; Swartz, nov. gen. et spec. plant. 25 (1788) ...	...	...	...	...	...	— S.A. — —	— N.A. B.fl.VII,613	M.fr.VI,85	...	As., Afr.
C. pallida; Hackel in Engler's bot. Jahrb. VI, 248 (1885) ...	...	...	...	...	...	W.A. — — — —	— N.A. — —			
C. scariosa; F. v. M., fragm. VI, 85 (1868) ...	...	...	...	...	...	W.A. S.A. — —	— Q. N.A. B.fl.VII,614	M.fr.XI,130.		

ELEUSINE; J. Gaertner, de fruct. I, 7 (1788). (Leptochloa, Dactyloctenium, Acrachne.)

E. cruciata; Lamarck, tabl. enclop. méth. I, 203, t. 48 (1791) ...	...	...	...	...	...	W.A. S.A. — V.	N.S.W. Q. N.A. B.fl.VII,615	M.fr.VIII,111	...	As., Afr., Am., P.
E. Indica; J. Gaertner, de fruct. I, 7 (1788) ...	...	...	...	...	...	— — — —	N.S.W. Q. — B.fl.VII,615	M.fr.VIII,112	...	As., Afr., Am., P.
E. verticillata; Roxburgh, Fl. Ind. ed. Car. & Wall. I, 346 (1820) ...	...	...	...	...	...	— — — —	— Q. N.A. B.fl.VII,616	M.fr.VIII,112	...	As., Afr.
E. digitata; Sprengel, syst. cur. post. 36 (1827) ...	...	...	...	...	...	W.A. S.A. — —	N.S.W. Q. N.A. B.fl.VII,617	M.fr.I,216; VIII,112; XI,130.		
E. Chinensis; F. v. M., Cens. 134 (1882) from Koenig (1821) ...	...	...	...	...	...	— — — —	N.S.W. Q. — B.fl.VII,617	M.fr.VIII,132	...	As.
E. polystachya; F. v. M.; Cynodon; R. Brown, prodr. 187 (1810) ...	...	...	...	...	...	— — — —	— Q. N.A. B.fl.VII,617			

CENTOTHECA ; Desvaux in Palisot, Agrost. 69, t. 14 (1812).

C. lappacea ; Desvaux in Palisot, Agrost. 69, t. 14 (1812) Q. — B.f.VII,640 M.fr.VI,85; VIII,116; X,76. As., Afr., P.

POA ; Linné, gen. plant. 20 (1737). (Glyceria partly.)

P. Labillardieri ; Steudel, syn. glum. I, 262 (1855) W.A. S.A. T. V. — — N.A. B.f.VII,651
P. homomalla ; Nees in Lehm. pl. Preiss. II, 104 (1846) W.A. — — — — — B.f.VII,651
P. caespitosa ; G. Forster, fl. ins. Austr. prodr. 89 (1786) W.A. S.A. T. V. N.S.W. Q. — B.f.VII,651 M.fr.VIII,133; XI,13. N.Z.
P. Maxwelli ; Benth. Fl. Austr. VII, 653 (1878) W.A. — — — — — B.f.VII,652
P. nodosa ; Nees in Lehm. pl. Preiss. II, 105 (1846) W.A. S.A. — V. N.S.W. Q. — B.f.VII,653 M.fr.VIII,132.
P. saxicola ; R. Brown, prodr. 654 (1810) — — T. — — — — B.f.VII,654
P. lepida ; F. v. M., fragm. VIII, 130 (1874) W.A. S.A. — V. N.S.W. — — B.f.VII,654 M.fr.XI,130.
P. Fordeana ; F. v. M., fragm. VIII, 130 (1874) — S.A. — V. N.S.W. Q. — B.f.VII,657 M.fr.VIII,130.
P. fluitans ; Scopoli, flor. Carniol. 106 (1772) W.A. S.A. T. V. N.S.W. — — B.f.VII,657 M.fr.VIII,129 ... E., As., Afr., Am.
P. latispicea ; F. v. M., Cens. of Austr. pl. 134 (1882) — — — — — N.S.W. Q. — B.f.VII,658 M.fr.VIII,127.
P. syrtica ; F. v. M. in Transact. Vict. Inst. 45 (1854) W.A. S.A. T. V. — — — — B.f.VII,658 M.fr.VIII,130 ... N.Z.
P. dives ; F. v. M., Cens. of Austr. pl. 134 (1882) — — — V. — — — B.f.VII,659 M.fr.VI,147; VIII,129; XI,130; III,147.
P. ramigera ; F. v. M. in Transact. Vict. Inst. 45 (1854) W.A. S.A. — V. N.S.W. Q. N.A. B.f.VII,659 M.fr.VI,85; VIII,131; XI,130.

FESTUCA ; Dillenius, nov. gen. 90, t. 3 (1719). (Glyceria partly, Vulpia partly, Schedonorus, Schenodorus, Brizopyrum.)

F. duriuscula ; Linné, spec. plant. 74 (1753) — S.A. T. V. N.S.W. — — B.f.VII,663 M.fr.VI,85 ... E., As., Am., P.
F. scirpoidea ; F. v. M., fragm. VIII, 129 (1874) W.A. — — — — — B.f.VII,665 M.fr.XI,130.
F. litoralis ; Labillardière, Nov. Holl. plant. spec. I, 22, t. 27 (1804) W.A. S.A. T. V. N.S.W. Q. — B.f.VII,666 M.fr.VIII,129 ... N.Z.
F. Hookeriana ; F. v. M. in J. Hooker, Fl. Tasman. II, 122, t. 115 (1860) — — T. V. N.S.W. — — B.f.VII,666 M.fr.VIII,131.

DIPLACHNE ; Palisot, Essai d'une nouv. Agrostogr. 9, t. 16 (1812). (Uralepis.)

D. loliiformis ; F. v. M. in Benth. Fl. Austr. VII, 618 (1878) — S.A. — V. N.S.W. Q. — B.f.VII,616 M.fr.VIII,128; XI,130. As.
D. Muelleri ; Benth. Fl. Austr. VII, 619 (1878) W.A. S.A. — — — — N.A. B.f.VII,619
D. fusca ; Palisot, Agrost. 163 (1812) W.A. S.A. — V. N.S.W. Q. — B.f.VII,619 M.fr.VIII,127 ... As., Afr.
D. parviflora ; Benth. Fl. Austr. VII, 620 (1878) — — — — — N.A. B.f.VII,620 M.fr.VIII,129.

TRIODIA ; R. Brown, prodr. fl. Nov. Holl. 182 (1810). (Sieglingia 1799.)

T. Mitchelli ; Benth. Fl. Austr. VII, 606 (1878) W.A. S.A. — — N.S.W. Q. N.A. B.f.VII,606
T. pungens ; R. Brown, prodr. 182 (1810) — S.A. — — — — Q. N.A. B.f.VII,606 M.fr.VIII,129.
T. Cunninghamii ; Benth. Fl. Austr. VII, 606 (1878) — — — — — Q. N.A. B.f.VII,606
T. irritans ; R. Brown, prodr. 182 (1810) W.A. S.A. — V. N.S.W. Q. — B.f.VII,607 M.fr.VIII,129.
T. proceera ; R. Brown, prodr. 182 (1810) — — — — — N.A. B.f.VII,607
T. microstachya ; R. Brown, prodr. 182 (1810) — — — — — N.A. B.f.VII,607
T. microdon ; F. v. M., syst. Cens. of Austr. pl. 135 (1882) from Benth. (1878) — — — — N.S.W. — — B.f.VII,605

DISTICHLIS ; Rafinesque in Journ. de Phys. LXXXIX, 104 (1819). (Uniola partly, Brizopyrum partly.)

D. maritima ; Rafinesque in Journ. de Phys. LXXXIX, 104 (1819) — S.A. T. V. N.S.W. — — B.f.VII,637 M.fr.VIII,129 ... Am.

BROMUS ; Dillenius in Linné, syst. nat. 8 (1735) ; Linné, gen. plant. 15 (1737).

B. arenarius ; Labillardière, Nov. Holl. pl. spec. I, 23, t. 28 (1804) W.A. S.A. — V. N.S.W. Q. — B.f.VII,660 M.fr.VI,126; XI,20 ... N.Z.

ERAGROSTIS ; Palisot, Essai d'une nouv. Agrostogr. 70, t. 14 (1812). (Poa partly.)

E. tenella ; Palisot in Roem. et Schult. syst. veg. II, 576 (1817) W.A. S.A. — V. N.S.W. Q. N.A. B.f.VII,643 M.fr.VIII,132; XI,130 As.
E. nigra ; Nees in Steud. nomencl. bot. I, 563 (1841) — — — — — N.S.W. Q. — B.f.VII,643 As.
E. imbecilla ; Benth. Fl. Austr. VII, 643 (1878) — — — — — Q. — B.f.VII,643 N.Z.
E. trichophylla ; Benth. Fl. Austr. VII, 644 (1878) — S.A. — — — — — B.f.VII,644
E. leptocarpa ; Benth. Fl. Austr. VII, 644 (1878) — S.A. — — — — Q. N.A. B.f.VII,644
E. megalosperma ; F. v. M. in Benth. Fl. Austr. VII, 644 (1878) — — — — N.S.W. Q. — B.f.VII,644 [P.
E. pilosa ; Palisot, Agrost. 71 (1812) W.A. S.A. — V. N.S.W. Q. — B.f.VII,645 M.fr.VIII,133; XI,130. E., As., Afr., Am.,

E. leptostachya; Steudel, syn. glum. I, 279 (1855) ...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,645	
E. Schultzii; Benth. Fl. Austr. VII, 646 (1878) ...	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.VII,646	
E. diandra; Steudel, syn. glum. I, 279 (1855) ...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,646	M.fr.XI,130.
E. Brownii; Nees in Steud. nomencl. bot. I, 562 (1841) ...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,646	... As.
E. concinna; Steudel, syn. glum. I, 279 (1855) ...	...	...	...	...	—	S.A.	—	—	—	Q.	N.A.	B.fl.VII,647	
E. speciosa; Steudel, syn. glum. I, 279 (1855) ...	...	...	...	...	W.A.	S.A.	—	—	—	Q.	N.A.	B.fl.VII,648	
E. laniflora; Benth. Fl. Austr. VII, 648 (1878) ...	...	...	...	...	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.VII,648	
E. eriopoda; Benth. Fl. Austr. VII, 648 (1878) ...	...	...	...	...	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.VII,648	
E. setifolia; Nees in Hook. Lond. Journ. II, 419 (1843) ...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,648	
E. lacunaria; F. v. M., first gen. Rep. 20 (1853) ...	...	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VII,649	
E. falcata; Gaudichaud in Voy. Freyc. Bot. 408, t. 25 (1826) ...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,649	
E. stenostachya; Steudel, syn. glum. I, 279 (1855) ...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.VII,650	

HETERACHNE; Benth. in J. Hook. icon. plant. XIII, 39 (1877).

H. Brownii; Benth. in J. Hook. icon. XIII, 40 (1877) ...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.VII,635	M.fr.VIII,132.
H. Gulliveri; Benth. in J. Hook. icon. XIII, 39, t. 1250 (1877)...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.VII,635	

ECTROSIA; R. Brown, prodr. fl. Nov. Holl. 185 (1810).

E. Schultzii; Benth. Fl. Austr. VII, 633 (1878) ...	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.VII,633	
E. leporina; R. Brown, prodr. 186 (1810) ...	...	...	...	...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.VII,633	M.fr.VIII,109.
E. agrostoides; Benth. Fl. Austr. VII, 634 (1878) ...	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.VII,634	
E. Gulliveri; F. v. M., fragm. VIII, 201 (1874) ...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.VII,634	M.fr.VIII,201.

ELYTHROPHORUS; Palisot, Essai d'une nouv. Agrostogr. 67, t. 14 (1812).

E. articulatus; Palisot, Agrost. 67, t. 14 (1812) ...	...	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,638	M.fr.VII,109 ... As., Afr.
---	-----	-----	-----	-----	---	------	---	----	--------	----	------	--------------	----------------------------

TRIRHAPHIS; R. Brown, prodr. fl. Nov. Holl. 185 (1810.) (Triraphis.)

T. mollis; R. Brown, prodr. 185 (1810) ...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,603	M.fr.VIII,108.
T. pungens; R. Brown, prodr. 185 (1810) ...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.VII,604	M.fr.VIII,108,125.
T. bromoides; F. v. M., fragm. VIII, 108 (1874) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.VII,604	M.fr.VIII,108.
T. danthonioides; F. v. M., fragm. VIII, 125 (1874) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.VII,604	M.fr.VIII,108.

AGROPYRON; J. Gaertner, nov. comm. Petrop. XIV, 539 (1770). (Agropyrum, Triticum partly, Anthosachne, Vulpia partly.)

A. scabrum; Palisot, Agrost. 102 (1812) ...	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,665	M.fr.VI,85; XI,27,130. N.Z.
A. velutinum; Nees in Hook. Lond. Journ. of Bot. II, 417 (1843) ...	...	...	...	...	—	—	T.	V.	N.S.W.	—	—	B.fl.VII,665	
A. pectinatum; Palisot, Agrost. 102 (1812) ...	...	...	...	...	—	—	T.	V.	N.S.W.	—	—	B.fl.VII,666	M.fr.VI,85.

ARUNDO; Tournefort, inst. 526 (1700), from Varro. (Phragmites.)

A. Phragmites; Dodoens, stirp. hist. pemptad. 602 (1583) ...	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,637	... E., As., Afr., Am., P., N.Z.
A. Roxburghii; F. v. M., syst. Cens. of Austr. pl. 135 (1882) from Kunth (1843) ...	...	...	...	...	—	—	—	—	—	Q.	N.A.	—	M.fr.XI,139. As.

BAMBUSA; Schreber, gen. plantar. I, 236 (1789).

B. Arnhemica; F. v. M. in Austral. journ. of Pharm. new ser. I, 447 (1886) ...	...	...	...	...	—	—	—	—	—	—	N.A.	—	M.fr.VI,85; XI,129.
--	-----	-----	-----	-----	---	---	---	---	---	---	------	---	---------------------

ACOTYLEDONEAE.

A. L. DE JUSSIEU, genera plantarum 1 (1789).

ACOTYLEDONEAE VASCULARES.

MEISSNER, plantarum vascularium genera, 429 (1843).

RHIZOSPERMAE.

G. WEBER, primit. Fl. Holsat. 74 (1780).—Genera 5, Species 11.

AZOLLA; Lamarek, Encycl. méthod. I, 343 (1783).

A. pinnata; R. Brown, prodr. 167 (1810) ... S.A. — V. N.S.W. Q. — B.fl.VII,679 M.fr.V,140 ... As., Afr.
 A. filiculoides; Lamarek, Encycl. méth. I, 343 (1783) ... — S.A. T. V. N.S.W. Q. — B.fl.VII,680 M.fr.V,140; XI,127. Am., N.Z.

SALVINIA; Micheli, nov. plant. gen. 107, t. 58 (1729).

S. cucullata; Roxburgh, fl. Indic. IV, 547 (1843) ... W.A. — — — — — — — — — — As.

MARSILEA; Linné, syst. nat. 8 (1735); gen. plant. 326 (1737). (Marsiglia.)

M. quadrifolia; Linné, spec. plant. 1099 (1753) ... W.A. S.A. — V. N.S.W. Q. N.A. B.fl.VII,683 M.fr.V,140 ... E., As., Afr.

PILULARIA; Vaillant, Bot. Paris. 159, t. 15 (1727).

P. globulifera; Linné, spec. plant. 1100 (1753) ... W.A. S.A. T. V. N.S.W. — — B.fl.VII,684 M.fr.V,140 ... E., As., Afr., N.Z.

ISOETES; Linné, skanska resa 420 (1751).

I. tripus; A. Braun in Berl. Monatsb. 559 (1863) ... W.A. — — — — — — — — — —
 I. Drummondii; A. Braun in Berl. Monatsb. 593 (1863) ... W.A. — — V. N.S.W. — — B.fl.VII,672 M.fr.V,140.
 I. elatior; F. v. M. & A. Braun in Linnaea XXV, 722 (1852) ... — — T. — — — — — M.fr.V,140; XI,139.
 I. humilior; F. v. M. & A. Braun in Linnaea XXV, 722 (1852) ... — — T. — — — — — M.fr.V,140; XI,139.
 I. Gunnii; A. Braun in Berl. Monatsb. 535 (1868) ... — — T. — — — — — M.fr.XI,139.
 I. Muelleri; A. Braun in Berl. Monatsb. 541 (1868) ... — — — — — Q. — — M.fr.XI,139.

LYCOPODINAE.

SWARTZ, synops. filic. XV (1806).—Genera 5, Species 21.

PSILOTUM; Swartz in Schrader, Journ. fuer die Bot. II, 109 (1800).

P. triquetrum; Swartz, synops. filic. 187 (1806) ... — — — V. N.S.W. Q. — B.fl.VII,681 M.fr.V,112; X,118. As., Afr., Am.
 P. complanatum; Swartz, synops. filic. 187 (1806) ... — — — — — Q. — B.fl.VII,682 M.fr.V,112 ... As., Afr., Am., P.

TMESIPTERIS; Bernhardt in Schrader, Journ. fuer die Bot. II, 131 (1800).

T. Tannensis; Bernhardt in Schrad. Journ. II, 131, t. 2 (1800) ... — — — T. V. N.S.W. Q. — B.fl.VII,680 M.fr.V,112; X,118. P., N.Z.

LYCOPODIUM; Ruppis, Fl. Jenens. 32 (1718), from Tabernaemontanus (1588).

L. Selago; Linné, spec. plant. 1102 (1753) ... — — — T. V. N.S.W. — — B.fl.VII,674 M.fr.V,111; X,118. E., As., Afr., Am., N.Z.
 L. Phlegmaria; Linné, spec. plant. 1101 (1753) ... — — — — — Q. — B.fl.VII,674 M.fr.V,111; X,118. As., Afr. [N.Z.]
 L. clavatum; Linné, spec. plant. 1101 (1753) ... — — — T. V. N.S.W. — — B.fl.VII,675 M.fr.V,111 ... E., As., Afr., Am., P.,
 L. Carolinianum; Linné, spec. plant. 1104 (1753) ... W.A. S.A. T. V. N.S.W. Q. — B.fl.VII,675 M.fr.V,111; X,118. As., Afr., Am., P., N.Z.
 L. laterale; R. Brown, prodr. 165 (1810) ... — S.A. T. V. N.S.W. Q. — B.fl.VII,675 M.fr.V,111; VI,124; X,118. P., N.Z.
 L. cernuum; Linné, spec. plant. 1103 (1753) ... — — — — — Q. N.A. B.fl.VII,676 M.fr.V,111; X,118. As., Afr., Am., N.Z.
 L. densum; Labillardière, Nov. Holl. plant. spec. II, 104, t. 251 (1806) ... — S.A. T. V. N.S.W. Q. — B.fl.VII,676 M.fr.V,111; X,118. P., N.Z.
 L. scarosum; G. Forster, florul. ins. Austr. prodr. 87 (1786) ... — — — T. V. — — B.fl.VII,676 M.fr.V,112. ... Am., N.Z.
 L. volubile; G. Forster, florul. ins. Austr. prodr. 86 (1786) ... — — — — — — — N.A. B.fl.VII,677 M.fr.V,112. ... As., Am., P., N.Z.

SELAGINELLA; Palisot, prodr. des fam. de l'Aethéogamie 101 (1805). (Lycopodium partly.)

S. Preissiana; Spring in Mém. de l'Ac. de Brux. 61 (1842) ...	...	...	...	W.A. S.A. T. V. N.S.W. Q. —	B.fl. VII, 677 M.fr. V, 112; VIII, 275.
S. uliginosa; Spring in Mém. de l'Ac. de Brux. 60 (1842) ...	...	...	...	W.A. S.A. T. V. N.S.W. Q. —	B.fl. VII, 678 M.fr. V, 112; VIII, 275.
S. flabellata; Spring in Mém. de l'Ac. de Brux. 174 (1849) ...	...	...	...	— — — — — Q. —	B.fl. VII, 678 M.fr. VIII, 175, 275. As., Am., P.
S. Australiensis; Baker in Britten's Journ. of Bot. 144 (1883) ...	...	...	...	— — — — — Q. —	B.fl. VII, 678 M.fr. V, 112; VIII, 274.
S. Belangeri; Spring in Mém. de l'Ac. de Brux. 242 (1849) ...	...	...	...	— — — — — N.S.W. Q. N.A.	B.fl. VII, 679 ... As., P.
S. Bakeriana; Bailey, suppl. Queensl. Fl. 62 (1886) ...	...	...	...	— — — — — Q. —	—
S. Brisbaneensis; Bailey, suppl. Queensl. Fl. 62 (1886) ...	...	...	...	— — — — — Q. —	—
S. Pumilio; Spring in Mém. de l'Ac. de Brux. 241 (1849) ...	...	...	...	— — — — — Q. —	—

PHYLOGLOSSUM; Kunze in Muhl & Schlechtendal, Bot. Zeitung, 721 (1843).

P. Drummondii; Kunze in der Bot. Zeit. 721 (1843) ...	...	...	...	W.A. — T. V. N.S.W. — —	B.fl. VII, 672 M.fr. V, 112; X, 118.
---	-----	-----	-----	-------------------------	--------------------------------------

FILICES.

Linné, gen. plant. 322 (1737).—Genera 35, Species 212.

OPHIOGLOSSUM; Tournefort, inst. rei herb. 548, t. 325 (1700), from Bock (1539). (Ophioderma.)

O. vulgatum; C. Bauhin, pinax. theatr. bot. 354 (1623) ...	...	...	...	— S.A. T. V. N.S.W. Q. N.A.	B.fl. VII, 688 M.fr. V, 112; VI, 124. E., As., Afr., Am., P.,
O. pendulum; Linné, sp. pl. edit. sec. 1518 (1763) ...	...	...	...	— — — — — N.S.W. Q. —	B.fl. VII, 689 M.fr. V, 113; VI, 124. As., Afr. [N.Z.]

BOTRYCHIUM; Swartz in Schrader, Journ. fuer die Bot. II, 110 (1800).

B. Lunaria; Swartz in Schrad. Journ. II, 110 (1800) ...	...	...	...	— — T. V. N.S.W. — —	B.fl. VII, 690 M.fr. V, 113 ... E., As., Am.
B. ternatum; Swartz, synops. filic. 172 (1806) ...	...	...	...	— — T. V. N.S.W. Q. —	B.fl. VII, 690 M.fr. V, 113 ... As., Am., N.Z.

HELMINTOSTACHYS; Kaulfuss, enum. filic. Chamiss. 28 (1824).

H. Zeylanica; Kaulfuss, enum. fil. 28 (1824) ...	...	...	...	— — — — — Q. —	B.fl. VII, 690 M.fr. V, 113; IX, 190. As., P.
--	-----	-----	-----	----------------	---

LYGODIUM; Swartz in Schrader, Journ. fuer die Bot. II, 7 & 106 (1800). (Hydroglossum, Lygodictyon.)

L. scandens; Swartz in Schrad. Journ. II, 106 (1800) ...	...	...	...	— — — — — N.S.W. Q. N.A.	B.fl. VII, 691 M.fr. V, 113; VII, 84; X, 118. As., Afr., P.,
L. reticulatum; Schkuhr, kryptog. Gewaechse 139, t. 139 (1809) ...	...	...	...	— S.A. — — — Q. —	B.fl. VII, 692 M.fr. VII, 83; VIII, 157. P.
L. Japonicum; Swartz, syn. filic. 154 (1806) ...	...	...	...	— — — — — Q. N.A.	B.fl. VII, 692 M.fr. V, 113; VII, 84, 156. As., P.
L. dichotomum; Swartz, syn. filic. 154 (1806) ...	...	...	...	— — — — — Q. —	—

SCHIZAEA; Smith in Mém. de l'Acad. Turin V, 149, t. 19 (1791). (Lopidium, Actinostachys.)

S. rupestris; R. Brown, prodr. 162 (1810) ...	...	...	...	— — — — — N.S.W. — —	B.fl. VII, 693 M.fr. V, 113.
S. fistulosa; Labillardière, Nov. Holl. pl. spec. II, 103, t. 250 (1806) ...	...	...	...	— S.A. T. V. N.S.W. — —	B.fl. VII, 693 M.fr. V, 113 ... Afr., Am., P., N.Z.
S. dichotoma; Smith in Mém. de l'Ac. Tur. V, 149 (1791) ...	...	...	...	— — — — — N.S.W. Q. N.A.	B.fl. VII, 694 M.fr. V, 113; VIII, 275; X, 118 As., Afr.,
S. bifida; Willdenow in act. Acad. Erfurt. 30, t. 3, f. 3 (1802) ...	...	...	...	— S.A. T. V. N.S.W. Q. N.A.	B.fl. VII, 693 M.fr. V, 113 ... N.Z. [Am., P., N.Z.]

ANGIOPTERIS; G. Hoffmann in Comment. Goett. XII, 29 (1796).

A. evecta; G. Hoffmann in Comment. Goett. XII, 29 (1796) ...	...	...	...	— — — — — N.S.W. Q. —	B.fl. VII, 694 M.fr. V, 114; VII, 120; X, 118. As., Afr., P.
--	-----	-----	-----	-----------------------	--

MARATTIA; Swartz, nov. gen. et spec. plant. 8 & 128 (1788).

M. fraxinea; Smith, plant. icon. hact. ined. t. 48 (1790) ...	...	...	...	— — — — — N.S.W. Q. —	B.fl. VII, 695 M.fr. V, 114; X, 118. As., Afr., Am., P., N.Z.
---	-----	-----	-----	-----------------------	---

TRICHOMANES; Linné, syst. nat. 8 (1835); hort. Cliffort. 476 (1737).

T. peltatum; Baker in Journ. Linn. Soc. IX, 336, t. 8 (1865) ...	...	...	...	— — — — — Q. —	B.fl. VII, 701 M.fr. X, 117 ... P.
T. Vitiense; Baker in Journ. Linn. Soc. IX, 338, t. 8 (1865) ...	...	...	...	— — — — — N.S.W. Q. —	B.fl. VII, 701 M.fr. VIII, 32; IX, 132. P.
T. Motleyi; Bosch, Hymenoph. Javan. (1861) ...	...	...	...	— — — — — Q. —	— ... As., P.
T. cuspidatum; Willdenow, spec. plant. V, 499 (1810) ...	...	...	...	— — — — — Q. —	M.fr. XI, 131 & 139. As., Afr.
T. parvulum; Poirer, Encycl. méth. VIII, 44 (1810) ...	...	...	...	— — — — — N.S.W. Q. —	B.fl. VII, 701 M.fr. VII, 155 ... As., Afr., P.
T. digitatum; Swartz, synops. filic. 370 & 422 (1806) ...	...	...	...	— — — — — N.S.W. — —	B.fl. VII, 702 ... As., Afr., P.
T. venosum; R. Brown, prodr. 159 (1810) ...	...	...	...	— — T... V. N.S.W. Q. —	B.fl. VII, 702 M.fr. V, 116; IX, 191. N.Z.
T. Javanicum; Blume, flor. Jav. filic. 224 (1829) ...	...	...	...	— — — — — Q. —	B.fl. VII, 702 ... As., P.

T. rigidum; Swartz, nov. gen. et spec. pl. 137 (1788)	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.f.VII,702	M.fr.V,115	...	As., Afr., Am., P., N.Z.
T. maximum; Blume, enum. fil. Javan. 228 (1826)	...	...	...	...	—	—	—	—	—	Q.	—	...	...	...	As., P.
T. bipunctatum; Poiret, Encycl. méth. VIII, 44 (1810)	...	...	...	...	—	—	—	—	—	Q.	—	B.f.VII,703	M.fr.X,117; XI,132	...	As., P.
T. humile; G. Forster, floral. ins. Austr. prodr. 84 (1786)	...	...	...	...	—	—	—	—	V.	N.S.W.	—	—	...	...	As., P., N.Z.
T. caudatum; Brackenridge, Bot. of Wilk. U.S. Exped. fil. 256, t. 36 (1854)	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.f.VII,703	M.fr.V,116; VII,122	...	P.
T. meifolium; Bory in Willd. sp. pl. V, 509 (1810)	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.f.VII,703	M.fr.VII,121; XI,132	...	As., Afr., P.
T. parviflorum; Poiret, Encycl. méth. VIII, 44 (1810)	...	...	...	...	—	—	—	—	—	Q.	—	B.f.VII,704	M.fr.VII,122	...	As., Afr.
T. Sayeri; F. v. M. & Baker in Britten's Journ. of Bot. XXV, 163 (1887)	...	...	...	...	—	—	—	—	—	Q.	—	—	...	...	

HYMENOPHYLLUM; Smith in Roemer, Archiv I, 56 (1797).

H. Malingii; J. Hooker, Handb. N.Z. Flora I, 357 (1864) implied	...	...	—	—	T.	—	—	—	—	—	—	—	...	...	N.Z.
H. marginatum; Hooker & Greville, icon. filic. I, t. 34 (1829)	...	...	—	—	T.	—	—	—	N.S.W.	—	—	B.f.VII,705	...	...	
H. nitens; R. Brown, prodr. 159 (1810)	...	...	—	—	T.	V.	N.S.W.	Q.	—	—	—	B.f.VII,705	...	...	P., N.Z.
H. Javanicum; Sprengel, syst. veg. IV, 132 (1827)	...	...	—	—	T.	V.	N.S.W.	Q.	—	—	—	B.f.VII,705	M.fr.V,116	...	As., P., N.Z.
H. Tunbridgense; Smith in Roemer, Archiv. I, 56 (1797)	...	...	—	—	T.	V.	N.S.W.	Q.	—	—	—	B.f.VII,706	M.fr.V,116	...	E., As., Afr., Am., N.Z., P.
H. multifidum; Swartz, synops. filic. 149 & 378 (1806)	...	...	—	—	—	—	—	—	N.S.W.	—	—	B.f.VII,707	...	...	P., N.Z.
H. bivalve; Swartz, synops. filic. 146 & 372 (1806)	...	...	—	—	—	—	—	—	N.S.W.	—	—	—	...	...	P., N.Z.

CERATOPTERIS; Brogniart in Bull. de la Soc. philom. 186 (1821). (Parkeria.)

C. thalictroides; Brogniart in Bull. Soc. philom. 186 (1821)	...	...	—	—	—	—	—	—	—	Q.	N.A.	B.f.VII,695	M.fr.V,122; VIII,158	...	As., Afr., Am., P.
--	-----	-----	---	---	---	---	---	---	---	----	------	-------------	----------------------	-----	--------------------

GLEICHENIA; Smith in Mém. de l'Acad. Turin V, 418 (1791). (Mertensia, Platyzoma, Stromatopteris.)

G. platyzoma; F. v. M., fragm. V, 114 (1866)	...	...	—	—	—	—	—	—	N.S.W.	Q.	N.A.	B.f.VII,696	M.fr.V,114; VIII,157; X,118	...	
G. circinata; Swartz, syn. filic. 165 & 394 (1806)	...	...	—	—	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.f.VII,697	M.fr.V,115	...	...	As., P., N.Z.
G. dicarpa; R. Brown, prodr. 161 (1810)	...	...	—	—	—	—	—	—	T.	V.	N.S.W.	Q.	—	...	As., P., N.Z.
G. flabellata; R. Brown, prodr. 161 (1810)	...	...	—	—	—	—	—	—	T.	V.	N.S.W.	Q.	N.A.	B.f.VII,698	M.fr.V,114; VIII,157. P., N.Z.
G. Hermannii; R. Brown, prodr. 161 (1810)	...	...	—	—	—	—	—	—	V.	N.S.W.	Q.	N.A.	B.f.VII,698	M.fr.V,114; X,118	As., Afr., Am., P., N.Z.

OSMUNDA; Tournefort, inst. rei herb. 547, t. 324 (1700) from l'Obel (1576). (Todea, Leptopteris.)

O. barbara; Thunberg, prodr. plant. Capens. 171 (1800)	...	...	—	—	S.A.	T.	V.	N.S.W.	Q.	—	—	B.f.VII,699	M.fr.V,114	...	Afr., N.Z.
O. Fraseri; F. v. M., Cens. 137 (1882) from Hooker & Greville (1829)	...	...	—	—	—	—	—	—	N.S.W.	—	—	B.f.VII,700	M.fr.V,114; VII,156	...	P.
O. Moorei; F. v. M., Cens. 137 (1882) from Baker (1873)	...	...	—	—	—	—	—	—	N.S.W.	—	—	B.f.VII,700	M.fr.VIII,157	...	

CYATHEA; Smith in Mém. de l'Acad. Turin V, 416 (1791). (Hemitelia, Amphicosmia.)

C. Lindsayana; Hooker, syn. filic. 25 (1865)	...	...	—	—	—	—	—	—	N.S.W.	Q.	—	B.f.VII,708	M.fr.VIII,177	...	
C. arachnoidea; Hooker, syn. filic. 24 (1866)	...	...	—	—	—	—	—	—	—	Q.	—	B.f.VII,708	M.fr.VI,200	...	As.
C. Macarthurii; F. v. M., fragm. VII, 177 (1874)	...	...	—	—	—	—	—	—	N.S.W.	—	—	B.f.VII,708	M.fr.VIII,176; X,120	...	
C. medullaris; Swartz, synops. filic. 140 (1806)	...	...	—	—	—	—	—	—	T.	V.	N.S.W.	—	B.f.VII,708	M.fr.V,116; VIII,177	As., P., N.Z.
C. Cunninghami; J. Hooker, icon. plant. 985 (1854)	...	...	—	—	—	—	—	—	T.	V.	—	—	...	...	N.Z.
C. brevipinna; Baker in Benth, fl. Austr. VII, 709 (1878)	...	...	—	—	—	—	—	—	N.S.W.	—	—	B.f.VII,709	M.fr.XI,139	...	
C. Moorei; F. v. M., Cens. 137 (1882) from Baker (1872)	...	...	—	—	—	—	—	—	N.S.W.	—	—	B.f.VII,709	M.fr.VIII,157	...	

ALSOPIHILA; R. Brown, prodr. 158 (1810).

A. Rebeccae; F. v. M., fragm. V, 53 (1865)	...	...	—	—	—	—	—	—	—	Q.	—	B.f.VII,710	M.fr.V,117,215; VIII,180	...	
A. Loddigesii; Kunze in Schlecht. Linnaea XXIII, 221 (1850)	...	...	—	—	—	—	—	—	N.S.W.	Q.	—	B.f.VII,710	...	...	
A. Australis; R. Brown, prodr. 158 (1810)	...	...	—	—	—	—	—	—	T.	V.	N.S.W.	Q.	—	B.f.VII,710	M.fr.V,52,116; VII,120; VIII,178
A. excelsa; R. Brown, prodr. 158 (1810)	...	...	—	—	—	—	—	—	N.S.W.	Q.	—	—	...	...	
A. Leichhardtiana; F. v. M., fragm. V, 53 (1865)	...	...	—	—	—	—	—	—	N.S.W.	Q.	—	B.f.VII,711	M.fr.V,117; VI,200; VII,120	...	
A. Robertiana; F. v. M., fragm. V, 54 (1865)	...	...	—	—	—	—	—	—	—	Q.	—	B.f.VII,712	M.fr.V,117,213	...	

DICKSONIA; L'Héritier, sert. Angl. 31 (1788). (Cibotium, Patania, Dennstaedtia, Deparia, Balantium.)

D. Billardieri; F. v. M., fragm. VIII, 175 (1874)	...	...	...	...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.f.VII,712	M.fr.V,117; VI,199; VII,120.	P., N.Z.
D. Youngiae; C. Moore in Hooker & Baker synops. filic. 51 (1865)	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.f.VII,713	M.fr.VI,200.	

- D. davallioides*; R. Brown, prodr. 158 (1810) ... — — — V. N.S.W. Q. — B.f. VII, 713 M.fr. VIII, 158 ... As., N.Z.
D. nephrodioides; F. v. M., Cens. 137 (1882) from Baker (1872) ... — — — — N.S.W. — — B.f. VII, 714 M.fr. VIII, 157; IX, 78; X, 104.

DAVALLIA; Smith in Mém. de l'Acad. de Turin V, 414 (1791). (Microlepia, Humata.)

- D. solida*; Swartz, synops. filic. 132 & 345 (1806) ... — — — — Q. — B.f. VII, 715 ... As., P.
D. elegans; Swartz, synops. filic. 132 & 347 (1806) ... — — — — Q. — B.f. VII, 715 M.fr. V, 118; VIII, 157. As., Afr.
D. pyxidata; Cavanilles, description de las plantas n. 694 (1802) ... — — — — V. N.S.W. Q. — B.f. VII, 716 M.fr. V, 118 ... P.
D. pedata; Smith in Mém. de l'Ac. de Turin V, 414 (1791) ... — — — — Q. — B.f. VII, 716 M.fr. V, 118 ... As., Afr.
D. dubia; R. Brown, prodr. 157 (1810) ... — — — — T. V. N.S.W. Q. — B.f. VII, 716 M.fr. V, 118.
D. flaccida; R. Brown, prodr. 157 (1810) ... — — — — Q. — B.f. VII, 717 M.fr. V, 118; VII, 156. As., Afr., P.
D. tripinnata; F. v. M. in Benth. Fl. Austr. VII, 717 (1878) ... — — — — Q. — B.f. VII, 717

VITTARIA; Smith in Mém. de l'Acad. de Turin V, 414 (1791).

- V. elongata*; Swartz, synops. filic. 109 & 302 (1806) ... — — — — N.S.W. Q. — B.f. VII, 718 M.fr. V, 122; VII, 121. As., Afr., P.
V. falcata; Kunze in Feé, gen. fil. t. 6, f. 1 (1852) ... — — — — Q. — — — — As.

LINDSAYA; Dryander in Mém. de l'Acad. de Turin V, 413 (1791). (Lindsaea, Isoloma, Synophlebium, Schizoloma.)

- L. linearis*; Swartz, synops. filic. 118 & 318, t. 3 (1806) ... W.A. S.A. T. V. N.S.W. Q. — B.f. VII, 719 M.fr. V, 119 ... P., N.Z.
L. dimorpha; Bailey, Queensland Ferns 19 (1874) ... — — — — N.S.W. Q. — B.f. VII, 719
L. cultrata; Swartz, synops. filic. 119 (1806) ... — — — — Q. — B.f. VII, 719 ... As., Afr.
L. flabellulata; Dryander in Transact. Linn. Soc. III, 41, t. 8 (1797) ... — — — — Q. N.A. B.f. VII, 720 M.fr. V, 119 ... As.
L. decomposita; Willdenow, spec. plant. V, 425 (1810) ... — — — — Q. — B.f. VII, 720 ... As., P.
L. trichomanoides; Dryander in Transact. Linn. Soc. III, 43, t. 11 (1797) ... — — — — T. — N.S.W. — — B.f. VII, 720 M.fr. V, 118 ... N.Z.
L. microphylla; Swartz, synops. filic. 120 (1806) ... — — — — N.S.W. Q. — B.f. VII, 721 M.fr. V, 119.
L. incisa; Prentice in Trim. Journ. of Bot. 295 (1873) ... — — — — N.S.W. Q. — B.f. VII, 721 ... N.Z.
L. Fraseri; Hooker, spec. filic. I, 221, t. 70 (1846) ... — — — — Q. — B.f. VII, 721 M.fr. V, 118.
L. ensifolia; Swartz, synops. filic. 118 (1806) ... — — — — Q. N.A. B.f. VII, 721 M.fr. V, 118 ... As., Afr., P.
L. lanuginosa; Wallich, numerical list n. 154 (1828) ... — — — — Q. — B.f. VII, 721 M.fr. V, 118 ... As., Afr.

ADIANTUM; Tournefort in 543, t. 317 (1700) from Hippocrates, Theophrastus, Dioscorides and Plinius.

- A. lunulatum*; N. Burman, Fl. Ind. 235 (1768) ... — — — — Q. N.A. B.f. VII, 723 M.fr. V, 119 ... As., Afr., Am.
A. Capillus veneris; Linné, spec. plant. 1096 (1753) ... — — — — Q. — B.f. VII, 723 ... E., As., Am., Afr.
A. Aethiopicum; Linné, syst. nat. edit. declin. n. 15 (1759) ... W.A. S.A. T. V. N.S.W. Q. — B.f. VII, 724 M.fr. V, 119 ... E., As., Afr., Am., P., [N.Z.]
A. formosum; R. Brown, prodr. 155 (1810) ... — — — — V. N.S.W. Q. — B.f. VII, 724 M.fr. V, 120; VIII, 157. N.Z.
A. affine; Willdenow, spec. plant. V, 448 (1810) ... — — — — N.S.W. Q. — B.f. VII, 724 M.fr. V, 119 ... N.Z.
A. diaphanum; Blume, flora Javae, filic. 215 (1829) ... — — — — V. N.S.W. Q. — B.f. VII, 725 M.fr. VII, 121 ... As., N.Z., P.
A. hispidulum; Swartz, syn. filic. 124 & 321 (1806) ... — — — — V. N.S.W. Q. — B.f. VII, 725 M.fr. V, 120 ... As., Afr., Am., P., N.Z.

CHEILANTHES; Swartz, syn. filic. 126, t. 3 (1806). (Notholaena, Nothochlaena.)

- C. Prenticei*; Luerssen in Uhlw. Bot. Centralblatt IX, 440 (1882) ... — — — — Q. N.A. — —
C. Clelandi; F. v. M. & Tate in proceed. R.S. of S.A. X, 81 (1887) ... — — — — S.A. — —
C. pumilio; F. v. M., Cens. 138 (1882) from R. Brown (1810) ... — — — — Q. — B.f. VII, 773 M.fr. VIII, 175.
C. fragillima; F. v. M., fragm. V, 123 (1866) ... — — — — Q. N.A. B.f. VII, 774 M.fr. V, 123.
C. vellea; F. v. M., fragm. V, 123 (1866) ... W.A. S.A. — V. N.S.W. Q. N.A. B.f. VII, 773 M.fr. V, 123; VIII, 176. E., As., Afr.
C. distans; A. Braun, index sem. hort. Berol. (1859) ... W.A. S.A. — V. N.S.W. Q. — B.f. VII, 774 M.fr. V, 122; VIII, 176. P., N.Z.
C. tenuifolia; Swartz, syn. filic. 126 (1806) ... W.A. S.A. T. V. N.S.W. Q. N.A. B.f. VII, 726 M.fr. V, 122 ... As., N.Z., P.

PTERIS; Linné, syst. nat. 9 (1735); gen. plant. 322 (1737). (Pellaea, Cheiloplecton, Platyloma, Litobrochia, Pteridium.)

- P. concolor*; Langsdorf & Fischer, icon. filic. t. 21 (1818) ... — — — — N.S.W. Q. — B.f. VII, 728 M.fr. V, 124 ... As., Afr., Am., P.
P. paradoxa; Baker in Benth. Fl. Austr. VII, 729 (1878) ... — — — — N.S.W. Q. — B.f. VII, 729
P. falcata; R. Brown, prodr. 154 (1810) ... — — — — T. V. N.S.W. Q. — B.f. VII, 729 M.fr. V, 123 ... As., N.Z.
P. rotundifolia; G. Forster, florul. ins. Austr. prodr. 79 (1786) ... — — — — Q. — B.f. VII, 730 ... N.Z.
P. longifolia; Linné, spec. plant. 1074 (1753) ... — — — — V. N.S.W. Q. — B.f. VII, 730 M.fr. V, 126 ... E., As., Afr., Am., P.

<i>P. ensiformis</i> ; N. Burmann, Fl. Ind. 230 (1768) — — — — — Q. — B.fl.VII,730 M.fr.V,123 As., P.
<i>P. umbrosa</i> ; R. Brown, prodr. 154 (1810) — — — — — V. N.S.W. Q. — B.fl.VII,730 M.fr.V,126.
<i>P. baurita</i> ; Linné, spec. plant. 1076 (1753) — — — — — Q. — B.fl.VII,731 M.fr.V,125 As., Afr., Am., P., N.Z.
<i>P. arguta</i> ; Aiton, hort. Kew. III, 458 (1789) — — — — — V. N.S.W. Q. — B.fl.VII,731 M.fr.V,126 As., Afr., Am., P., N.Z.
<i>P. aquilina</i> ; Linné, spec. plant. 1075 (1753) W.A. S.A. T. V. N.S.W. Q. — B.fl.VII,731 M.fr.V,126 E., As., Afr., Am., P., N.Z.
<i>P. incisa</i> ; Thunberg, prodr. pl. Capens. 171 (1800) — — — — — S.A. T. V. N.S.W. Q. — B.fl.VII,732 M.fr.V,124; VIII,158. As., Afr., Am., P.,
<i>P. marginata</i> ; Bory. voy. dans les isles d'Afr. II, 192 (1804) — — — — — N.S.W. Q. — B.fl.VII,733 M.fr.V,125 As., Afr., N.Z. [N.Z.]
<i>P. comans</i> ; G. Forster, florul. ins. Austr. prodr. 79 (1786) — — — — — T. V. N.S.W. Q. — B.fl.VII,733 M.fr.V,125; VII,121. Am., P., N.Z.

LOMARIA; Willdenow in Magaz. der Ges. naturf. Freunde zu Berlin, III, 160 (1809). (Stegania, Plagiogyria.)

<i>L. Patersoni</i> ; Sprengel, sys. veg. IV, 62 (1827) — — — — — T. V. N.S.W. Q. — B.fl.VII,734 M.fr.V,122; VII,158. As., P., N.Z.
<i>L. vulcanica</i> ; Blume, fl. Javae, filic. 202 (1836) — — — — — T. — — — — — Q. — B.fl.VII,734 M.fr.V,121 ... As., P., N.Z.
<i>L. discolor</i> ; Willdenow, spec. plant. V, 293 (1810) — — — — — S.A. T. V. N.S.W. Q. — B.fl.VII,735 M.fr.V,121 ... N.Z.
<i>L. lanceolata</i> ; Sprengel, syst. veg. IV, 62 (1827) — — — — — S.A. T. V. N.S.W. — — — — — B.fl.VII,735 M.fr.V,121 ... P., N.Z.
<i>L. attenuata</i> ; Willdenow, spec. plant. V, 290 (1810) — — — — — — — — — — — B.fl.VII,736 ... Afr., Am., P.
<i>L. alpina</i> ; Sprengel, syst. veg. IV, 62 (1827) — — — — — T. V. N.S.W. — — — — — B.fl.VII,736 M.fr.V,121 ... Am., N.Z.
<i>L. fluviatilis</i> ; Sprengel, syst. veg. IV, 65 (1827) — — — — — T. V. N.S.W. — — — — — B.fl.VII,736 M.fr.V,121 ... N.Z.
<i>L. Fullagari</i> ; F. v. M., fragm. VIII, 157 (1874) — — — — — — — — — — — B.fl.VII,737 M.fr.VIII,157.
<i>L. Capensis</i> ; Willdenow, spec. plant. V, 291 (1810) — — — — — S.A. T. V. N.S.W. Q. — B.fl.VII,737 M.fr.V,121 ... As., Afr., Am., P., N.Z.
<i>L. euphlebia</i> ; Kunze in Mohl & Schlecht. Bot. Zeit. VI, 521 (1849) — — — — — — — — — — — Q. — B.fl.VII,738 M.fr.VI,124 ... As.

BLECHNUM; Linné, spec. plant. 1077 (1753).

<i>B. cartilagineum</i> ; Swartz, syn. filic. 114 & 312 (1806) — — — — — V. N.S.W. Q. — B.fl.VII,738 M.fr.V,120.
<i>B. laevigatum</i> ; Cavanilles, descripcion de las plantas 650 (1802) — — — — — — — — — — — B.fl.VII,739 M.fr.V,120.
<i>B. serrulatum</i> ; C. Richard in Annal. du Mus. d'Hist. nat. Par. I, 114 (1802) — — — — — — — — — — — Q. N.A. B.fl.VII,739 M.fr.V,120 ... As., Am., P
<i>B. orientale</i> ; Linné, syst. nat. ed. decim. (1759) — — — — — — — — — — — Q. N.A. B.fl.VII,740 — ... As., P.

MONOGRAMMA; Commerçon in Schkuhr's Kryptog. Gew. 82, t. 87 (1809). (Monogramme, Vaginularia, Pleurogramme, Diclidopteris.)

<i>M. Junghuhnii</i> ; Hooker, spec. filic. V, 123, t. 289 (1864) — — — — — — — — — — — Q. — B.fl.VII,740 M.fr.VII,119 ... As., P.

WOODWARDIA; Smith in mém. de l'Acad. de Turin V, 411 (1791). (Doodia.)

<i>W. aspera</i> ; Mettenius, fil. hort. bot. Lips. 65 (1856) — — — — — V. N.S.W. Q. — B.fl.VII,740 M.fr.V,130.
<i>W. caudata</i> ; Cavanilles, descripcion de las pl. n. 653 (1802) — — — — — T. V. N.S.W. Q. N.A. B.fl.VII,742 M.fr.V,129; VIII,158. P., N.Z.

ASPLENIUM; Linné, gen. plant. 322 (1737) from C. Bauhin (1651). (Scolopendrium, Allantodea, Diplazium, Callipteris, Anisogonium, Thamnopteris, Neottopteris,

Darea, Caenopteris, Athyrium, Diplora, Asplenium.)

<i>A. Nidus</i> ; Linné, spec. plant. 1079 (1753) — — — — — V. N.S.W. Q. — B.fl.VII,744 M.fr.V,130 ... As., Afr., P.
<i>A. Robinsonii</i> ; F. v. M. in Britten's Journ. Bot. XXII, 289 (1884) — — — — — — — — — — — N.S.W. — — — — —
<i>A. simplicifrons</i> ; F. v. M., fragm. V, 74 (1865) — — — — — — — — — — — Q. — B.fl.VII,744 M.fr.V,74,130; VIII,158.
<i>A. attenuatum</i> ; R. Brown, prodr. 150 (1810) — — — — — — — — — — — N.S.W. Q. — B.fl.VII,745 M.fr.V,130. [N.Z.]
<i>A. Trichomanes</i> ; Linné, spec. plant. 1080 (1753) — — — — — T. V. N.S.W. — — — — — B.fl.VII,745 M.fr.V,131 ... E., As., Afr., Am., P.,
<i>A. flabellifolium</i> ; Cavanilles, descripcion de las pl. n. 636 (1802) W.A. S.A. T. V. N.S.W. Q. — B.fl.VII,745 M.fr.V,131 ... N.Z.
<i>A. paleaceum</i> ; R. Brown, prodr. 150 (1810) — — — — — — — — — — — Q. — B.fl.VII,746 M.fr.V,131; VIII,158.
<i>A. falcatum</i> ; Lamarck, Encycl. méth. I, 303 (1783) — — — — — — — — — — — N.S.W. Q. — B.fl.VII,746 M.fr.V,131; VII,156. As., Afr., P., N.Z.
<i>A. Hookerianum</i> ; Colenso in Journ. of Nat. Sc. Tasm. II, 169 (1842) — — — — — V. N.S.W. — — — — — B.fl.VII,747 ... N.Z.
<i>A. furcatum</i> ; Thunberg, prodr. pl. Cap. 172 (1800) W.A. S.A. — — — — — V. N.S.W. — — — — — B.fl.VII,748 M.fr.V,131 ... As., Afr., Am., P.
<i>A. laserpitifolium</i> ; Lamarck, Encycl. méth. I, 310 (1783) — — — — — — — — — — — Q. — B.fl.VII,749 M.fr.V,131 ... As., P. [N.Z.]
<i>A. marinum</i> ; Linné, spec. plant. 1081 (1753) — — — — — T. V. N.S.W. Q. — B.fl.VII,747 M.fr.V,132 ... E., As., Afr., Am., P.,
<i>A. bulbiferum</i> ; G. Forster, floral. ins. Austr. prodr. 433 (1786) — — — — — S.A. T. V. N.S.W. Q. — B.fl.VII,749 M.fr.V,132 ... As., Afr., Am., P., N.Z.
<i>A. pteridoides</i> ; Baker, syn. filic. sec. edit. 488 (1874) — — — — — — — — — — — N.S.W. — — — — — B.fl.VII,749
<i>A. umbrosum</i> ; J. Smith in Hook. Lond. Journ. IV, 174 (1842) — — — — — T. V. N.S.W. Q. — B.fl.VII,749 M.fr.V,132 ... As., Afr., P., N.Z.
<i>A. pallidum</i> ; Blume, fl. Javae, filic. 776 (1836) — — — — — — — — — — — Q. — — — — — M.fr.XI,131 ... As., P.
<i>A. sylvaticum</i> ; Presl, reliq. Haenk. I, 42 (1830) — — — — — — — — — — — Q. — B.fl.VII,750 M.fr.V,133 ... As., Afr., P.

A. maximum ; D. Don, prodr. fl. Nepal. 8 (1825)	— — — — N.S.W.	Q. —	B.fl. VII, 751	...	...	As.
A. polypodoides ; Mettenius, fl. hort. Lips. 78 (1856)	— — — — —	Q. —	B.fl. VII, 751	M.fr. V, 132	...	As.
A. melanochlamsy ; Hooker, spec. filic. III, 259 (1860)	— — — — — N.S.W.	— —	B.fl. VII, 751			
A. decussatum ; Swartz, spec. fil. 76 & 260 (1806)	— — — — —	Q. —	B.fl. VII, 751	...	...	As., Afr., P.
A. diversifolium ; Cunningham in Lond. hort. Brit. suppl. 581 (1832)	— — — — — N.S.W.	— —	—			

CYSTOPTERIS ; Bernhardt in Schrader's neuem Journ. II, 26 (1806).

C. fragilis ; Bernhardt in Schrad. Journ. I, 26 (1806)	— — T. — — — —	— —	B.fl. VII, 752	M.fr. V, 118	...	E., As., Afr., Am., P. N.Z.
---	----------------	-----	----------------	--------------	-----	-----------------------------

ASPIDIUM ; Swartz in Schrader's Journ. II, 429 (1800). (Nephrodium, Nephrolepis, Polystichum partly, 1797, Lastraea, Sagenia, Oleandra.)

A. cordifolium ; Swartz, syn. filic. 45 (1806)	— — — — — N.S.W.	Q. —	B.fl. VII, 754	M.fr. V, 136	...	As., Afr., Am., P., N.Z.
A. exaltatum ; Swartz, syn. filic. 45 (1806)	— — — — —	Q. N.A.	B.fl. VII, 754	M.fr. V, 136	...	As., Afr., Am.
A. ramosum ; Palisot, Fl. d'Oware et de Benin II, 53, t. 91 (1807)	— — — — — N.S.W.	Q. —	B.fl. VII, 754	M.fr. V, 135	...	As., Afr., P.
A. unitum ; Swartz, syn. filic. 47 (1806)	W.A. — — — — N.S.W.	Q. N.A.	B.fl. VII, 755	M.fr. V, 137; VIII, 158.	...	As., Afr.
A. pteroides ; Swartz, syn. filic. 49 (1806)	— — — — —	Q. —	B.fl. VII, 755	...	...	As., P.
A. molle ; Swartz, syn. filic. 49 (1806)	— S.A. — V. N.S.W.	Q. N.A.	B.fl. VII, 756	M.fr. V, 135	...	As., Afr., P., N.Z.
A. truncatum ; Gaudichaud in Voy. Freyc. Bot. 332, t. 10 (1826)	— — — — — N.S.W.	Q. —	B.fl. VII, 756	M.fr. V, 135	...	As., P.
A. confluent ; Mettenius in Schlecht. Linnaea XXXVI, 125 (1863)	— — — — —	Q. —	B.fl. VII, 757	M.fr. V, 133; VIII, 158.	...	
A. aculeatum ; Swartz in Schrad. Journ. II, 37 (1800)	— S.A. T. V. N.S.W.	Q. —	B.fl. VII, 757	M.fr. V, 134	...	E., As., Afr., Am., P., N.Z.
A. aristatum ; Swartz, syn. filic. 53 & 253 (1806)	— — — — — N.S.W.	Q. —	B.fl. VII, 757	M.fr. V, 134	...	As., Afr., P., N.Z.
A. coriaceum ; Swartz, syn. filic. 57 (1806)	— — T. V. N.S.W.	— —	B.fl. VII, 758	M.fr. V, 124	...	As., Afr., Am., N.Z.
A. decompositum ; Sprengel, syst. IV, 109 (1827)	— S.A. T. V. N.S.W.	Q. —	B.fl. VII, 758	M.fr. V, 136	...	P., N.Z.
A. tenerum ; Sprengel, syst. IV, 109 (1827)	— — — — V. N.S.W.	Q. —	B.fl. VII, 759			
A. uliginosum ; Kunze in Schlecht. Linnaea XX, 6 (1847)	— — — — — N.S.W.	Q. —	B.fl. VII, 759	M.fr. V, 133	...	As., P.
A. hispidum ; Swartz, syn. filic. 56 (1806)	— — T. V. N.S.W.	— —	B.fl. VII, 760	M.fr. V, 133	...	N.Z.

MENISCIUM ; Schreber, gen. plant. II, 757 (1791).

M. triphyllum ; Swartz, syn. filic. 19 & 206 (1806)	— — — — —	Q. —	—	...	...	As.
--	-----------	------	---	-----	-----	-----

POLYPODIUM ; Tournefort, inst. 540, t. 316 (1700), from Theophrastos, Dioscorides and Plinius. (Nipholobolus, Goniophlebium, Goniopteris, Phlebodium, Phegopteris, Phymatodes, Pleopeltis, Drynaria, Dictyopteris, Arthropteris, Xiphopteris, Meniscium partly ; from Dodoens, l'Ecluse, Bauhin, Morison, Ray, Plumier and particularly Petiver as Polypodium.)

P. Australe ; Mettenius, filic. hort. bot. Lips. 36 (1856)	— — T. V. N.S.W.	Q. —	B.fl. VII, 762	M.fr. V, 127	...	Am., N.Z.
P. Hookeri ; Brackenridge, Bot. Wilk. Exped. Filic. 4 (1854)	— — — — — N.S.W.	Q. —	B.fl. VII, 763	...	...	As., P.
P. contiguum ; Brackenridge, Bot. Wilk. Exped. Filic. 6, t. 2 (1854)	— — — — —	Q. —	B.fl. VII, 763	M.fr. V, 127	...	P.
P. grammitidis ; R. Brown, prodr. 147 (1810)	— — T. V. N.S.W.	— —	B.fl. VII, 764	M.fr. V, 127	...	N.Z.
P. tenellum ; G. Forster, florul. ins. Austr. prodr. 440 (1786)	— — — — — N.S.W.	Q. —	B.fl. VII, 764	M.fr. V, 126; VII, 121.	...	P., N.Z.
P. proliferum ; Roxburgh in Wall. num. list 11 (1828)	— — — — — N.S.W.	Q. N.A.	B.fl. VII, 765	M.fr. V, 128	...	As., Afr., P.
P. urophyllum ; Wallich, numerical list 11 (1828)	— — — — —	Q. —	B.fl. VII, 765	M.fr. IV, 165; V, 137; VIII, 121.	...	As., P.
P. Hillii ; Baker in Hooker, syn. filic. sec. edit. 505 (1874)	— — — — —	Q] —	B.fl. VII, 766			
P. poecilophlebium ; Hooker, spec. filic. V, 14 (1863)	— — — — —	Q. —	B.fl. VII, 766	M.fr. V, 127.		
P. serpens ; G. Forster, florul. insul. Austr. prodr. 81 (1786)	— — — — V. N.S.W.	Q. —	B.fl. VII, 767	M.fr. V, 129	...	P., N.Z.
P. confluent ; R. Brown, prodr. 146 (1810)	— — — — — N.S.W.	Q. —	B.fl. VII, 767	...	...	P.
P. acrostichoides ; G. Forster, florul. insul. Austr. prodr. 81 (1786)	— — — — —	Q. —	B.fl. VII, 767	M.fr. V, 129	...	As., P.
P. attenuatum ; R. Brown, prodr. 146 (1810)	— — — — — N.S.W.	Q. —	B.fl. VII, 768	M.fr. V, 128; V, 121.	...	P.
P. simplicissimum ; F. v. M. in Hooker & Baker, syn. filic. sec. edit. 513 (1874)	— — — — —	Q. —	B.fl. VII, 768	M.fr. VII, 126, 156.		
P. fusco-pilosum ; F. v. M. & Baker in Britt. journ. bot. XXV, 163 (1887)	— — — — —	Q. —	—			
P. nigrescens ; Blume, Fl. Javae, filic. 161, t. 70 (1829)	— — — — —	Q. —	B.fl. VII, 769	...	...	As., Afr., Am., P.
P. phymatodes ; Linné, mant. pl. alt. 306 (1771)	— — — — —	Q. N.A.	B.fl. VII, 769	...	...	As., Afr., P., N.Z.
P. pustulatum ; G. Forster, florul. ins. Austr. pr. 81 (1786)	— — T. V. N.S.W.	Q. —	B.fl. VII, 769	M.fr. V, 127	...	N.Z.
P. scandens ; G. Forster, florul. ins. Austr. pr. 81 (1786)	— — — — V. N.S.W.	Q. —	B.fl. VII, 770	M.fr. V, 128	...	P., N.Z.
P. verrucosum ; Wallich, numerical list II (1828)	— — — — —	Q. —	B.fl. VII, 770	...	...	As., P., N.Z.

<i>P. subauriculatum</i> ; Blume, Fl. Javae, filic. 177, t. 83 (1829)	Q.	B.fl.VII,770	M.fr.V,128	...	As., P.
<i>P. rigidulum</i> ; Swartz, synops. filic. 38 & 320 (1806)	Q.	B.fl.VII,771	...	...	As., P.
<i>P. quercifolium</i> ; Linné, spec. plant. 1087 (1753)	Q. N.A.	B.fl.VII,772	M.fr.V,127	...	As., P.
<i>P. irioides</i> ; Poiret, Encycl. méth. V, 513 (1804)	Q.	B.fl.VII,771	M.fr.V,127	...	As., Afr., P.
<i>P. punctatum</i> ; Thunberg, fl. Japon. 337 (1784)	Q.	B.fl.VII,764	M.fr.V,129	...	As., Afr., N.Z.

HYPOLEPIS; Bernhardt in Schrader, neu. Journ. I, 34 (1806).

<i>H. tenuifolia</i> ; Bernhardt in Schrad. Journ. II, 24 (1800)	Q.	B.fl.VII,726	M.fr.V,129	...	As., P., N.Z.
---	----	--------------	------------	-----	---------------

GRAMMITIS; Swartz in Schrader, Journ. II, 3 & 17 (1800). (Gymnogramme, Selligaea.)

<i>G. Reynoldsii</i> ; F. v. M. in Benth. Fl. Austr. VII, 775 (1878)	Q.	B.fl.VII,775	M.fr.VIII,175.		
<i>G. Muelleri</i> ; Hooker in F. v. M., fragm. V, 138 (1866)	Q.	B.fl.VII,775	M.fr.VI,124; VII,121.		
<i>G. rutifolia</i> ; R. Brown, prodr. 146 (1810)	Q.	B.fl.VII,775	M.fr.V,137	...	E., Am., N.Z.
<i>G. leptophylla</i> ; Swartz, syn. filic. 23 & 218, t. 1 (1806)	Q.	B.fl.VII,776	M.fr.V,137	...	E., As., Afr., Am., P., N.Z.
<i>G. pinnata</i> ; F. v. M., fragm. VI, 124 (1868)	Q.	B.fl.VII,776	M.fr.VI,124...		As., P.
<i>G. ampla</i> ; F. v. M., fragm. V, 188 (1866)	Q.	B.fl.VII,777	M.fr.V,188.		
<i>G. Sayeri</i> ; F. v. M. & Baker in Britt. journ. bot. XXV, 163 (1887)	Q.	—	—		

ANTROPHYUM; Kaulfuss, enum. filic. 197 (1824).

<i>A. reticulatum</i> ; Kaulfuss, enum. filic. 197 (1824)	Q.	B.fl.VII,777	M.fr.V,138	...	As., P.
--	----	--------------	------------	-----	---------

ACROSTICHUM; Linné, gen. plant. 322 (1737), indicative. (Elaphoglossum, Stenochlaena, Lomariopsis, Hymenolepis, Gymnopteris, Chrysodium.)

<i>A. conforme</i> ; Swartz, syn. filic. 10 & 192, t. 1 (1806)	Q.	B.fl.VII,778	M.fr.V,138	...	As., Afr., Am.
<i>A. scandens</i> ; J. Smith in Hook. spec. fil. V, 249 (1864)	Q. N.A.	B.fl.VII,778	M.fr.VI,124...		As., P.
<i>A. sorbifolium</i> ; Linné, spec. plant. 1069 (1753)	Q.	B.fl.VII,779	M.fr.VII,119.		As., Afr., Am.
<i>A. neglectum</i> ; Bailey in proc. Linn. Soc. of N.S.W. V, 32 (1880)	Q.	—	M.fr.XI,131.		
<i>A. repandum</i> ; Blume, Fl. Javae 39, t. 14 & 15 (1829)	Q.	B.fl.VII,779	M.fr.V,138	...	As., P.
<i>A. aureum</i> ; Linné, spec. plant. 1069 (1753)	Q. N.A.	B.fl.VII,779	M.fr.V,138; VII,120.		As., Afr., Am., P.
<i>A. spicatum</i> ; Linné, fil. suppl. pl. 444 (1781)... ..	Q.	B.fl.VII,780	M.fr.V,138	...	As., Afr., P.
<i>A. pteroides</i> ; R. Brown, prodr. 145 (1810)	Q. N.A.	B.fl.VII,780	M.fr.X,76.		
<i>A. Taylori</i> ; Bailey in trans. Accl. Soc. of Queensl. (1884)	Q.	—	—		

PLATYCERIUM; Desvaux in Mém. de la Soc. Linn. de Paris VI, 213 (1827).

<i>P. alcornu</i> ; Desvaux in Mém. Soc. Linn. Par. VI, 213 (1827)	Q.	B.fl.VII,780	M.fr.V,139	...	Afr.
<i>P. grande</i> ; J. Smith in Hooker, Lond. Journ. III, 402 (1841)	Q.	B.fl.VII,781	M.fr.V,139	...	As.

SEQUENCE OF ORDERS ACCORDING TO SYSTEMATIC AFFINITY, WITH INDICATION OF THEIR NUMBERS OF SPECIES.

Dilleniaceae ...	95	Rutaceae ...	190	Phytolaceae ...	11	Olacineae ...	15	Convolvulaceae...	70	Haemodoraceae	66
Ranunculaceae ...	17	Simarubeae ...	7	Nyctagineae ...	6	Balanophoreae ...	1	Hydrophyllaeae ...	2	Amaryllideae ...	21
Ceratophylleae ...	1	Zygophylleae ...	22	Thymeleae ...	75	Santalaceae ...	43	Solanaceae ...	79	Dioscorideae ...	4
Nymphaeaceae ...	5	Lineae ...	4	Leguminosae ..	1065	Loranthaceae ...	27	Scrophularinae...	80	Roxburghiaceae	1
Piperaceae ...	10	Geraniaceae ...	8	Connaraceae ...	2	Proteaceae ...	597	Orobanchaeae ...	1	Liliaceae ...	161
Magnoliaceae ...	4	Malvaceae ...	110	Rosaceae ...	17	Cornaceae ...	1	Lentibularinae ...	25	Palmae ...	25
Anonaceae ...	18	Sterculiaceae ...	125	Saxifrageae ...	36	Rubiaceae ...	127	Podostemoneae...	1	Nipaceae ...	1
Monimiaceae ...	16	Tiliaceae ...	56	Nepenthaceae ...	2	Caprifoliaceae ...	2	Gesneriaceae ...	4	Pandaneae ...	11
Myristiceae ...	1	Euphorbiaceae ...	224	Aristolochieae ...	5	Passifloreae ...	6	Bignoniaceae ...	7	Aroideae ...	10
Lauraceae ...	37	Urticaceae ...	65	Crassulaceae ...	6	Cucurbitaceae ...	27	Pedaliinae ...	3	Typhaceae ...	2
Menispermaceae ...	17	Cupuliferae ...	4	Hamamelideae ...	1	Compositae ...	539	Acanthaceae ...	30	Lemnaceae ...	6
Papaveraceae ...	1	Casuarineae ...	24	Onagreae ...	5	Campanulaceae	35	Labiatae ...	125	Fluviales ...	36
Capparideae ...	24	Celastrinae ...	18	Salicarieae ...	19	Candolleaceae ...	97	Verbenaceae ...	82	Alismaceae ...	6
Cruciferae ...	54	Meliaceae ...	36	Stackhousieae ...	13	Goodeniaceae ...	220	Myoporinae ...	76	Pontederiaceae ...	1
Violaceae ...	13	Sapindaceae ...	100	Halorageae ...	58	Gentianeae ...	24	Asperifoliae ...	52	Phillydreae ...	4
Flacourtiaceae ...	7	Malpighiaceae ...	2	Callitrichinae ...	2	Loganiaceae ...	52	Ericaceae ...	7	Commelineae ...	19
Samydaceae ...	4	Burseraceae ...	3	Rhizophoreae ...	7	Plantagineae ...	4	Epacrideae ...	275	Xyrideae ...	9
Pittosporae ...	40	Anacardiaceae ...	9	Combretaceae ...	27	Primulaceae ...	6	Coniferae ...	29	Flagellariaceae...	1
Droseraceae ...	46	Frankeniaceae ...	7	Myrtaceae ...	663	Myrsinaceae ...	12	Cycadeae ...	14	Junceae...	16
Elatineae ...	4	Plumbagineae ...	4	Melastomaceae ...	7	Sapotaceae ...	19	Scitamineae ...	11	Eriocaulaeae ...	18
Hypericinae ...	1	Portulacaeae ...	32	Rhamnaceae ...	90	Ebenaceae ...	15	Orchideae ...	271	Restiaceae ...	93
Ternstroemiaceae	1	Caryophylleae ...	26	Viniferae ...	17	Aquifoliaceae ...	2	Apostasiaceae ...	1	Cyperaceae ...	380
Guttiferae ...	3	Amarantaceae ...	102	Leeceaeae ...	2	Styraceae ...	2	Burmanniaceae	2	Gramineae ...	345
Polygaleae ...	32	Salsolaceae ...	111	Araliaceae ...	22	Jasmineae ...	21	Irideae ...	24	Rhizospermae ...	11
Tremandreae ...	17	Ficoideae ...	28	Umbelliferae ...	106	Apocyneae ...	47	Hydrocharideae	9	Lycopodinae ...	21
Ochnaceae ...	1	Polygonaceae ...	25	Elaeagneae ...	1	Asclepiadeae ...	61	Taccaceae ...	1	Filices ...	212

SEQUENCE OF ORDERS ACCORDING TO PREDOMINANCE, WITH INDICATION OF THEIR NUMBERS OF SPECIES.

Leguminosae ... 1065	Scrophularinae ... 80	Ficoideae ... 28	Viniferae ... 17	Ericaceae ... 7	Callitrichinae ... 2
Myrtaceae ... 663	Solanaceae ... 79	Combretaceae ... 27	Rosaceae ... 17	Nyctagineae ... 6	Leeaceae ... 2
Proteaceae ... 597	Myoporinae ... 76	Cucurbitaceae ... 27	Monimiaceae ... 16	Crassulaceae ... 6	Caprifoliaceae ... 2
Compositae ... 539	Thymeleae ... 75	Loranthaceae ... 27	Junceae ... 16	Passifloreae ... 6	Aquifoliaceae ... 2
Cyperaceae ... 380	Convolvulaceae ... 70	Caryophylleae ... 26	Olacineae ... 15	Primulaceae ... 6	Styraceae ... 2
Gramineae ... 345	Haemodoraceae ... 66	Polygonaceae ... 25	Ebenaceae ... 15	Lemnaceae ... 6	Hydrophyllaeae ... 2
Epacrideae ... 275	Urticaceae ... 65	Lentibularinae ... 25	Cycadeae ... 14	Alismaceae ... 6	Burmanniaceae ... 2
Orchideae ... 271	Asclepiadeae ... 61	Palmae ... 25	Violaceae ... 13	Nymphaeaceae ... 5	Typhaceae ... 2
Euphorbiaceae ... 224	Halorageae ... 58	Capparideae ... 24	Stackhousieae ... 13	Aristolochieae ... 5	Ceratophylleae ... 1
Goodeniaceae ... 220	Tiliaceae ... 56	Casuarineae ... 24	Myrsinaceae ... 12	Onagreae ... 5	Myristiceae ... 1
Filices ... 212	Cruciferae ... 54	Gentianeae ... 24	Phytolacceae ... 11	Magnoliaceae ... 4	Papaveraceae ... 1
Rutaceae ... 190	Loganiaceae ... 52	Irideae ... 24	Scitamineae ... 11	Samydaceae ... 4	Ternstroemiaceae ... 1
Liliaceae ... 161	Asperifoliae ... 52	Zygophylleae ... 22	Pandaneae ... 11	Elatineae ... 4	Hypericinae ... 1
Rubiaceae ... 127	Apocynaeae ... 47	Araliaceae ... 22	Rhizospermae ... 11	Lineae ... 4	Ochnaceae ... 1
Sterculiaceae ... 125	Droseraceae ... 46	Jasmineae ... 21	Piperaceae ... 10	Cupuliferae ... 4	Podostemoneae ... 1
Labiatae ... 125	Santalaceae ... 43	Amaryllideae ... 21	Aroideae ... 10	Plumbagineae ... 4	Balanophoreae ... 1
Salsolaceae ... 111	Pittosporaeae ... 40	Lycopodinae ... 21	Anacardiaceae ... 9	Plantagineae ... 4	Hamamelideae ... 1
Malvaceae ... 110	Lauraceae ... 37	Salicariaceae ... 19	Hydrocharideae ... 9	Gesneriaceae ... 4	Elaeagneae ... 1
Umbelliferae ... 106	Meliaceae ... 36	Sapotaceae ... 19	Xyrideae ... 9	Dioscorideae ... 4	Cornaceae ... 1
Amarantaceae ... 102	Saxifrageae ... 36	Commelineae ... 19	Geraniaceae ... 8	Philhydroae ... 4	Orobanchaeae ... 1
Sapindaceae ... 100	Fluviales ... 36	Anonaceae ... 18	Flacourtiaeae ... 7	Guttiferae ... 3	Apostasiaceae ... 1
Candolleaceae ... 97	Campanulaceae ... 35	Celastrinae ... 18	Simarubeae ... 7	Burseraceae ... 3	Taccaceae ... 1
Dilleniaceae ... 95	Polygaleae ... 32	Ericoaulaeae ... 18	Frankeniaceae ... 7	Pedalinae ... 3	Roxburghiaceae ... 1
Restiaceae ... 93	Portulaceae ... 32	Ranunculaceae ... 17	Rhizophoreae ... 7	Malpighiaceae ... 2	Nipaceae ... 1
Rhamnaceae ... 90	Acanthaceae ... 30	Menispermaceae ... 17	Melastomaceae ... 7	Connaraceae ... 2	Pontederiaceae ... 1
Verbenaceae ... 82	Coniferae ... 29	Tremandreae ... 17	Bignoniaceae ... 7	Nepenthaeae ... 2	Flagellariaceae ... 1

Total Number of Orders of Vasculares, 156; of Genera, 1409; of Species, 8839. Of these occur in W.A., 3560: in S.A., 1892; in T., 1029; in V., 1894; in N.S.W., 3251; in Q., 3753; in N.A., 1956. Admitted from Flora Australiensis, 7814; additional species recorded in Census, 1025. Proportion of distribution through the respective Colonial territories:—In W.A., 40.3 per cent.; in S.A., 21.4 per cent.; in T., 11.6 per cent.; in V., 21.4 per cent.; in N.S.W., 36.8 per cent.; in Q., 42.5 per cent.; in N.A., 22.1 per cent. Of the total number 7501 are endemic in Continental Australia and Tasmania; therefore 1338 or 15.1 per cent. extend also to other countries. Of these 160 are found in Europe; 1032 in Asia; 515 in Africa; 315 in America; 558 in Polynesia; 291 in New Zealand.

Species added since the publication of the first Census 193, irrespective of some systematic changes. Geographic entries added since 1150.

Chronologic quotations of original authors of such species, as are transferred to other genera, will be given in the next supplement for which also many species additional for Australia have already been recorded in manuscript, while these pages were going through the press.

In all instances immigrated plants are excluded.

INDEX OF ORDERS AND GENERA.

Abbottia ...	125	Adiantum ...	231	Altingia ...	183	Anisotome ...	108	Argyrophanes ...	136	Athrotaxis ...	183
Abelmoschus ...	25	Adraستا ...	1	Alysicarpus ...	70	Anodopetalum ...	82	Argyroglossis ...	136	Athyrium ...	232
Abildgaardia ...	211	Adriana ...	36	Alyssopsis ...	9	Anoetochilus ...	188	Aristida ...	223	Atkinsonia ...	111
Abroma ...	27	Aegialinites ...	45	Alyssum ...	9	Anonaceae ...	4	Aristolochia ...	83	Atriplex ...	49
Abrophyllum ...	82	Aegialitis ...	45	Alyxia ...	156	Anopterus ...	82	Aristolochiaceae ...	83	Atylosia ...	71
Abrotanella ...	142	Aegiceras ...	154	Amanoa ...	35	Anosporum ...	210	Aristotelia ...	30	Australina ...	39
Abrus ...	72	Aegopogon ...	223	Amarantaceae ...	47	Anotis ...	125	Arnocrinum ...	200	Automachia ...	20
Abutilaea ...	24	Aera ...	224	Amarantus ...	49	Antennaria ...	134	Aroideae ...	203	Avicennia ...	173
Abutilon ...	24	Aeschynomene ...	69	Amaryllidaceae ...	195	Antheidosorus ...	139	Artanema ...	164	Aylmeyeria ...	47
Acacia ...	74	Afzelia ...	74	Amblyosperma ...	143	Anthericum ...	198	Arthratherum ...	223	Azolla ...	228
Acacna ...	82	Agapetes ...	177	Ambrina ...	50	Anthistiria ...	222	Arthaxon ...	221	Azorella ...	107
Acalypha ...	36	Agastachys ...	114	Ammannia ...	84	Anthobolus ...	110	Arthrochilus ...	191		
Acanthaceae ...	167	Agathis ...	183	Ammobium ...	138	Anthocerastes ...	138	Arthrocnemum ...	52	Babbagia ...	51
Acanthocarpus ...	200	Agati ...	68	Amomum ...	185	Anthocercis ...	162	Arthropodium ...	199	Babingtonia ...	91
Acanthocladium ...	136	Ageratum ...	130	Amoora ...	16	Anthosachne ...	227	Arthropteris ...	233	Backhousia ...	100
Acanthus ...	167	Aglia ...	16	Amorphophallus ...	203	Anthotium ...	147	Arthrostylis ...	217	Bacularia ...	201
Achilleopsis ...	27	Aglaiopsis ...	16	Amorphospermum ...	154	Anthotroche ...	162	Arthrotaxis ...	183	Baea ...	166
Achnophora ...	131	Agnostus ...	122	Amperea ...	33	Antiaris ...	38	Arthrotichum ...	48	Baeckea ...	91
Achras ...	154	Agonis ...	92	Amphibolus ...	204	Anticoryne ...	91	Arum ...	203	Baeobotrys ...	154
Achromolaena ...	138	Agromonia ...	81	Amphibromus ...	225	Antidesma ...	33	Arundinella ...	220	Balanophora ...	109
Achrysium ...	140	Agropyron ...	227	Amphicosmia ...	230	Antirrhoea ...	126	Arundo ...	227	Balanophoreae ...	109
Achyranthes ...	47	Agrostis ...	224	Amphipogon ...	223	Anthophyllum ...	234	Aryteria ...	41	Balanops ...	39
Acianthus ...	191	Agrostocrinum ...	192	Amphodus ...	70	Aotus ...	62	Asclepiadeae ...	157	Balantium ...	230
Acicalyptus ...	102	Ailantus ...	22	Anacardiaceae ...	44	Apera ...	224	Ascyrum ...	14	Balaustion ...	92
Aciphylla ...	108	Aillya ...	149	Anadenia ...	117	Aphananthe ...	37	Asparagopsis ...	197	Balfouria ...	157
Ackama ...	82	Aira ...	224	Anagallis ...	153	Aphanopetalum ...	82	Asparagus ...	197	Baloghia ...	37
Acmena ...	101	Aizoon ...	52	Anarthria ...	208	Aphelia ...	207	Asperella ...	222	Bambusa ...	227
Acomis ...	138	Ajuga ...	171	Ancana ...	5	Apium ...	108	Asperifoliae ...	175	Banksia ...	55, 122
Acrachne ...	225	Akania ...	42	Ancistrostigma ...	52	Aplotaxis ...	143	Asperula ...	128	Barbarea ...	9
Acradenia ...	20	Alania ...	199	Ancistrum ...	82	Apluda ...	222	Aspidium ...	233	Barklya ...	72
Acroclinium ...	136	Albizzia ...	80	Andersonia ...	182	Apocynae ...	156	Asplenium ...	232	Barringtonia ...	102
Acronychia ...	21	Alchemilla ...	81	Andrachne ...	33	Apodytes ...	109	Asprella ...	222	Bartlingia ...	62, 200
Acrophylum ...	82	Alchornea ...	36	Andrewia ...	175	Apolochlamys ...	138	Astartea ...	92	Basilicum ...	168
Acrostichum ...	234	Aldrovanda ...	12	Andriapetalum ...	116	Aponogeton ...	204	Astelia ...	198	Bassia ...	51
Acrotiche ...	177	Alectryon ...	42	Andropogon ...	222	Apophyllum ...	9	Aster ...	131	Batatas ...	159
Actephila ...	34	Alepyrum ...	207	Androstemma ...	194	Aporosa ...	35	Asteridea ...	135	Batrachium ...	221
Actinocarpus ...	205	Aleurites ...	36	Androstoma ...	177	Apostasia ...	192	Asterochiton ...	29	Bauera ...	83
Actinodium ...	87	Alisma ...	205	Anailema ...	205	Apostasiaceae ...	192	Asterolasia ...	18	Bauhinia ...	73
Actinopappus ...	138	Alismaceae ...	205	Anemone ...	3	Asterofoliaceae ...	155	Asteromyrtus ...	94	Baumea ...	216
Actinostachys ...	229	Allania ...	199	Angianthus ...	139	Arabis ...	9	Asterotrichon ...	24	Baxteria ...	200
Actinostigma ...	18	Allantodea ...	232	Angiopteris ...	229	Araliaceae ...	105	Astraea ...	90	Bea ...	166
Actinostrobilus ...	184	Allophylus ...	41	Angophora ...	97	Araucaria ...	183	Astrebla ...	225	Beaufortia ...	96
Actinotus ...	108	Alocasia ...	203	Anguillaria ...	198	Archeria ...	181	Astroloma ...	177	Bedfordia ...	142
Adansonia ...	26	Alopecurus ...	222	Anigozanthos ...	195	Archidendron ...	80	Astroticha ...	105	Beilschmiedia ...	6
Adeliopsis ...	7	Alphitonia ...	102	Anisacantha ...	51	Archontophoenix ...	202	Atalantia ...	21	Bellendenia ...	114
Adenanthra ...	74	Alpinia ...	185	Anisandra ...	170	Ardisia ...	154, 177	Atalaya ...	41	Bellis ...	130
Adenanthos ...	113	Alsomitra ...	129	Aniseia ...	159	Areca ...	202	Atelandra ...	170	Benincasa ...	129
Adenochilus ...	191	Alsophila ...	230	Anisogonium ...	232	Arenaria ...	46	Atherocephala ...	182	Bergia ...	14
Adenosma ...	163	Alstonia ...	157	Anisolepis ...	136	Argophyllum ...	82	Atherosperma ...	5	Berrya ...	30
Adenostemma ...	130	Alternanthera ...	47	Anisomeles ...	169	Argyrocome ...	136	Athrixia ...	135	Bertolonia ...	175
Adenostephanus ...	116	Althenia ...	204	Anisopogon ...	225	Argyrodendron ...	27	Athrodactylis ...	202	Bertya ...	33

Beyeria 32	Brunonia 146	Calostrophus... .. 209	Cathartocarpus 73	Chlamysporum 198	Coldenia 176
Beyeropsis 32	Bryonia 129	Calothamnus... .. 96	Cathormion 80	Chloanthes 172	Coleanthera 181
Bidaria 158	Bryonopsis 129	Calotis 131	Catosperma 147	Chloris 225	Coleocoma 134
Bidens 141	Buchanania 44	Calpurnia 72	Caturus 38	Chlorophytum 199	Coleostylis 146
Bignoniaceae... .. 166	Buchnera 165	Calthae... .. 3	Caulinia ... 70, 203, 204	Chlorizotheca 32	Coleus 168
Billardiera 12	Buckinghamia 122	Calycomis 82	Caustis 217	Chondrachne... .. 213	Colletia 104
Billotia 92, 96	Buechnera 165	Calycopeplus... .. 31	Cedrela 16	Choretrum 110	Colmeiroa 82
Biophytum 23	Bulbine 198	Calycothrix 89	Celastrinae 40	Choriceras 34	Colobandra 170
Bischofia 35	Bulbophyllum 186	Calyptanthaea 102	Celastrus 40	Chorilaena 20	Colobanthus 46
Blackburnia 21	Bulliarda 83	Calyptocalyx 202	Celmisia 131	Choripetalum 154	Colocasia 203
Blackwellia 11	Bulweria 167	Calyptrostegia 55	Celosia 49	Chorizandra 213	Colubrina 102
Blainvillea 141	Bunnya 173	Calyptrostigma 32	Celtis 37	Chorizema 57, 58	Columbea 183
Blancoa 194	Burchardia 198	Calystegia 160	Cenarrhenes 114	Chromochiton 138	Colymbea 183
Blandfordia 197	Burgesia 56	Calythrix 89	Cenchrus 220	Chrysobactran 198	Combretaceae 87
Bleasdalea 116	Burmannaia 192	Campanulaceae 143	Centaurea 143	Chrysocephalum 136	Comesperma 15
Blechnum 232	Burmanniaceae 192	Camphoromyrtus 91	Centella 107	Chrysocoryne 139	Commelina 205
Blennodia 9	Burnettia 191	Camptostemon 26	Centipeda 141	Chrysodiscus... .. 135	Commelineae 205
Blennospora 140	Bursaria 12	Campuleia 165	Centotheca 226	Chrysodium 234	Commerconia 27
Blepharanthemum 24	Burseraceae 43	Campylanthera 12	Centranthera 165	Chrysogonium 140	Compositae 129
Blepharocarya 44	Burtonia 59	Campynema 195	Centratherum 129	Chrysophyllum 154	Conanthodium 136
Blitum 50	Busbeckea 8	Cananga 5	Centrolepis 207	Chrysopogon 222	Conchium 120
Blumea 133	Butomopsis 205	Canarium 44	Centropappus 142	Chrysostemon 32	Couiferae 183
Blyxa 194	Byblis 13	Canavalia 71	Centunculus 153	Chrysorrhoea 88	Coniogeton 44
Bobea 126	Byronia 155	Candollea 1, 144	Cephaelis 127	Chthonocephalus 140	Connaraceae 81
Boca 166		Candolleaceae 144	Cephalopterum 140	Chuncoa 87	Conospermum 113
Boehmeria 38	Cabomba 4	Canscora 152	Cephalosorus... .. 139	Cibotium 230	Conostephiopsis 181
Boerhaavia 54	Cadaba 8	Cansjera 109	Cephalotus 83	Cinna 223	Conostephium 181
Bolbophyllum 186	Cadellia 22	Canthium 126	Ceranthia 156	Cinnamomum 6	Conostylis 194
Bombax 26	Caenopteris 232	Capparidaceae 8	Ceratogyne 142	Cirrhopterium 186	Conothamnus 96
Bonnaya 164	Caesalpinia 72	Capparis 8	Ceratopetalum 82	Cissampelos 7	Convolvulaceae 159
Boronia 17	Caesia... .. 199	Caprifoliaceae 128	Ceratophylleae 3	Cissodendron 105	Convolvulus... .. 160
Borya 200	Cajanus 71	Capsella 10	Ceratophyllum 3	Cissus 104	Conyza 133
Bosistoa 20	Cakile... .. 10	Carallia 86	Ceratopteris 230	Citriobatus 12	Coprosma 127
Bossiaea 65	Caladenia 191	Carapa 16	Cerbera 156	Citrus 22	Corchorus 30
Bothryodendron 106	Caladium 203	Carara 10	Cercodia 85	Cladium 216	Cordia 177
Botrychium 229	Calamus 201	Cardamine 9	Ceriops 86	Cladodes 36	Corlyline 197
Bouchardatia 21	Calandrinia 45	Cardiospermum 41	Ceropogia 159	Claoxylon 36	Corethrostylis 29
Bowenia 184	Calanthe 188	Cardwellia 122	Cesatia 106	Clausena 21	Coridochloa 218
Brachychiton 26	Caldasie 108	Carex 217	Chaetanthera 209	Claytonia 45	Cornaceae 124
Brachycome 130	Caldesia 205	Careya 102	Chaetaria 223	Cleisostoma 187	Correa 20
Brachyloa 181	Calceana 191	Cargillia 155	Chaetospora 214	Cleistanthus 35	Corybas 190
Brachynema... .. 82	Calectasia 200	Carissa 156	Chalcas 21	Cleistocalyx 101, 102	Corymbis 188
Brachypterum 72	Caletia 32	Carmichaelia... .. 68	Chamaelancium 87	Clematis 3	Corymborchis 188
Brachysema 56	Calaya 191	Carnarvonina 122	Chamaeraphis 218	Cleome 8	Corynotheca 199
Brachyspatha 203	Calliandra 80	Carpha 214	Chamaescilla... .. 199	Clerodendrum 173	Coryospermum 40
Brachystephium 130	Callicarpa 173	Carpodontoa 83	Chamaespherion 140	Clianthus 68	Corypha 202
Brackenridgea 17	Callicoma 82	Carronia 8	Chamaexeros... .. 200	Clidanthra 69	Corysanthes 190
Bradleya 34	Callicystus 71	Cartonema 206	Chapeliera 216	Clinostigma 202	Cosmelia 182
Bramia 164	Callipteris 232	Carumbium 37	Charlwoodia 197	Clitoria 70	Costus 185
Brasenia 4	Callistachya 57	Carunostylis... .. 189	Cheilanthes 231	Cloeia 100	Cotula 141
Brassaia 106	Callistemon 94	Caryodaphne... .. 6	Cheiloplecton 231	Coatesia 21	Covellia 37
Brathys 14	Callitriche 86	Caryophylleae 46	Cheiranthera... .. 12	Coccoloba 53	Crantzia 108
Breweria 160	Callitrichinae 86	Caryospermum 40	Cheioloma 131	Cocculus 7	Crantzioia 108
Breynia 35	Callitris 184	Caryota 202	Chenolea 51	Cochlospermum 11	Craspedia 140
Bridelia 35	Calocephalus... .. 140	Casearia 11	Chenopodina... .. 52	Cocos 202	Crassula 83
Brizopyrum 226	Calochilus 189	Cassia 73	Chenopodium 50, 52	Codiaeum 37	Crassulaceae 83
Brizula 207	Calogyne 149	Cassinia 138	Chenynia 92	Codonocarpus 54	Crepis... .. 143
Brochosiphon 168	Calomeria 138	Cassiniola 136	Chillocarpus 156	Coelachne 224	Cressa... .. 161
Brombya 21	Calonyction 159	Cassytha 7	Chilodia 169	Coelandria 186	Crinopodium... .. 84
Bromidium 224	Calopetalon 12	Castanospermum 72	Chiloglottis 192	Coelebogyne 36	Crinum 186
Bromus 226	Calophyllum... .. 14	Castanospora 41	Chionachne 222	Coelorrhachis 221	Crossolepis 139
Brucea 22	Calorophyllum 209	Casuarina 39	Chionanthus 156	Coelospermum 126	Crossotoma 143
Bruguiera 86	Calosciadium 108	Casuarineae 39	Chithonanthus 74	Coffea 126	Crotalaria 66
Brunella 169	Calostemma 196	Catakidozamia 184	Chlamydia 197	Coilocarpus 51	Croton 36

Crowea	18	Daphne	56	Dilleniaceae	1	Drosera	13	Emilia	142	Ethulopsis	134
Cruciferae	9	Daphnobyron	56	Dillwynia	64	Droseraceae	12	Emmenospermum	103	Euandra	217
Cryphia	169	Darea	232	Dimeria	221	Drummondia	20	Emphysopus	130	Eucalyptus	97
Cryptandra	103	Darlingia	122	Dimetopia	106	Dryandra	123	Empleurosma	42	Eucarya	110
Cryptocarya	6	Darwinia	87	Dimorphocoma	142	Drymaria	46	Encephalartos	184	Euchilopsis	64
Cryptosema	56	Dasmalla	172	Dimorpholepis	136	Drymisperrum	56	Enchylaena	52	Euchilus	62
Cryptostemon	87	Dasypogon	201	Diodontium	141	Drymophila	197	Enchysia	144	Euchiton	134
Cryptostylia	189	Datura	162	Dioscorea	196	Drymophlaeus	202	Endiandra	6	Eucryphia	83
Ctenosperma	141	Daucus	108	Dioscorideae	196	Drynaria	233	Enhalus	193	Eudeamia	97
Cucumia	129	Davallia	231	Diospyros	155	Duboisia	163	Enhydra	141	Eugenia	101
Cucurbitaceae	128	Davidsonia	83	Diotosperma	142	Dunbaria	71	Enneapogon	224	Eulalia	22
Cudrania	38	Daviesia	61	Diplachne	226	Duperreya	160	Entada	74	Eulophia	18
Culcoa	197	Decadon	85	Diplacrum	217	Duttonia	136, 174	Enterolobium	80	Eulophus	187
Cupania	41	Decalophium	87	Diplanthera	167, 204	Dysophylla	168	Nydra	141	Euodia	20
Cupuliferae	39	Decaspermum	101	Diplarrhena	193	Dysoxylum	16	Epacridaeae	177	Euonymus	40
Curculigo	196	Decaspora	181	Diplaspis	107	Dyaphania	50	Epacris	177, 181, 183	Euosma	153
Curcuma	185	Decazesia	139	Diplax	222	Eadesia	162	Epaltes	134	Eupatorinm	130
Cuscuta	161	Deeringia	49	Diplazium	232	Earlia	168	Epiandra	216	Euphorbia	31
Cuttisia	82	Delabechea	26	Diploglottis	41	Ebelingia	22	Epiblema	189	Euphorbiaceae	31
Cyanostegia	173	Delarbrea	105	Diplolaena	20	Ebenaceae	155	Epicharis	16	Euphoria	41, 42
Cyanothamnus	17	Dendrobium	186	Diplobolium	68	Ebermayera	167	Epilobium	84	Euphrasia	165
Cyanotis	205	Denhamia	40	Diplopappus	131	Eedeiocolea	208	Epipogum	188	Eupomatia	5
Cyathae	230	Denisonia	173	Diplopeltia	42	Echinocarpa	30	Epipremnum	203	Euroschinus	44
Cyathochaete	214	Dennstaedtia	230	Diplopogon	223	Echinocroton	36	Epithinia	126	Eurostorrhiza	217
Cyathodes	177	Dentella	125	Diplospora	126	Echinopogon	218	Epitriche	139	Eurybia	131
Cyathopappus	139	Deparia	230	Diplostephium	131	Echinopsilon	51	Eragrostis	226	Eurybiopsis	133
Cyathopsis	177	Deplanchea	167	Diplora	232	Echinodorus	205	Eranthemum	168	Eurycles	196
Cyathula	49	Depremea	172	Dipodium	187	Echinopogon	223	Erechtites	143	Euryomyrtus	91
Cybele	122	Derris	72	Dipogonia	223	Echinopsilon	51	Eremaea	97	Eustrephus	197
Cycadeae	184	Deschampsia	224	Dipteracanthus	167	Echinopsilum	175	Eremodendron	174	Eutacta	183
Cycas	184	Desmodcladus	209	Dipteranthemum	48	Echinua	36	Eremophila	174	Eutassa	183
Cyclogyne	68	Desmodium	69	Disarrhenum	223	Eclipta	141	Eremopyxis	90	Eutaxia	64
Cyclotheca	54	Desvauxia	207	Discaria	104	Ectrosia	227	Eremosyne	83	Euthalea	150
Cyenogeton	204	Deyeuxia	224	Dischidia	159	Ehretia	176	Eria	187	Euxolus	49
Cylicodaphne	6	Dianella	197	Diseopodium	214	Ehrharta	222	Eriachne	224	Evandra	217
Cylindrosorus	139	Diaspasis	147	Discospermum	126	Elachanthera	197	Erianthus	221	Evolvulus	160
Cymbidium	187	Diastemanthe	220	Diselma	184	Elachanthus	142	Ericaceae	177	Evonymus	40
Cymbonotus	143	Dicarpidium	27	Disemma	128	Elachocroton	37	Ericomyrtus	91	Exarrhena	176
Cyminosma	21	Dicera	30	Disoon	175	Elachopappus	139	Erigeron	133	Excaecaria	37
Cymodocea	204	Dicerna	69	Disiliaria	34	Elachothamnus	131	Eriocalia	108	Exocarpoea	110
Cynanchum	157, 158	Dichelachne	223	Dissocarpus	51	Elaeagneae	109	Eriocaulaeae	207	Exocarya	213
Cynoetionum	157	Diehoglottis	46	Distichlis	226	Elaeagnus	109	Eriocaulon	207	Eyrea	133
Cynodon	225	Dichondra	161	Distichostemon	43	Elaeocarpus	30	Eriochilus	191		
Cynoglossum	175	Dichopetalum	107	Distylis	149	Elaeodendron	40	Eriochiton	51	Fabricia	93
Cynometra	74	Dichopogon	199	Dithyrocarpus	206	Elaphoglossum	234	Eriochlamys	140	Fagraea	152
Cyperaceae	210	Dichosema	59	Dithyrostegia	139	Elatine	14	Eriochloa	218	Fagus	39
Cyperus	210	Dichrostachys	74	Ditoca	46	Elatineae	14	Eriocladium	139	Faradaya	173
Cyphanthera	162	Dicksonia	230	Ditomostrophe	29	Elatostemma	38	Erioglossum	41	Fatona	38
Cyphokentia	202	Di cladantha	168	Diuiris	189	Elatostachys	41	Eriosema	72	Faustula	136
Cypselocarpus	54	Diclidopteris	232	Dodonaea	42	Electrosperma	207	Eriotatemon	18	Fawcettia	7
Cyrtostylis	191	Dicliptera	168	Dolichandra	167	Eleocharis	211	Eritrichum	175	Fenzlia	101
Cystanthe	183	Dierastylis	172	Dolichandrone	167	Eleoichiton	211	Erodiphyllum	131	Festuca	226
Cystopteris	233	Dietyopteris	233	Dolichos	71	Elephantopus	130	Erodium	23	Fibichia	225
		Didiscus	106	Donatia	146	Elettaria	185	Erpetion	10	Ficoideae	52
Dacrydium	184	Didymanthus	51	Donia	68	Eleusine	225	Erycihe	159	Ficus	37
Dactyloctenium	225	Didymeria	20	Doodia	232	Eleuthanthas	127	Eryngium	108	Fieldia	166
Daemia	158	Didymocarpus	166	Dopatrium	164	Elichrysaum	136	Eryaimum	9	Filices	229
Dalbergia	72	Didymochaeta	224	Doryanthes	196	Elionurua	221	Erythraea	151	Fimbristylis	211
Dallachya	102	Didymonema	216	Doryphora	5	Elodea	194	Erythrina	71	Figueta	202
Damasonium	193, 205	Didymoplexis	188	Dracaena	197	Elynanthus	214	Erythrophlaeum	74	Fischera	107
Dammara	183	Didymotheca	54	Dracophyllum	183	Elythrophorus	227	Erythrorrhis	188	Fitzalania	5
Dampiera	146	Digitaria	218	Drakaea	191	Emblingia	8	Erythroxylum	23	Fitzgeraldia	5, 191
Danthonia	225	Dilivaria	167	Drapetes	56	Embothrium	122	Etaeria	188	Fitzroya	184
Daphnandra	5	Dillenia	1	Drinya	4	Emex	53	Ethulia	129	Flacourtieae	11

Flagellaria ...	206	Gingidium ...	108	Gymnosporia ...	40	Hemichroa ...	49	Hydriastele ...	201	Ixodia ...	138
Flagellariaceae ...	206	Glabraria ...	6	Gymnostachya ...	203	Hemiclidia ...	123	Hydrilla ...	194	Ixora [...	126
Flaveria ...	141	Gleichenia ...	230	Gymnostyles ...	141	Hemicyclia ...	35	Hydrocharideae ...	193	Jacksonia ...	60
Fleischera ...	24	Glinus ...	53	Gymnothrix ...	220	Hemigenia ...	170	Hydrocharia ...	193	Jambosa ...	101
Flemingia ...	72	Glischrocaryon ...	85	Gynopogon ...	156	Hemiphora ...	173	Hydrocotyle ...	106	Jansonia ...	56
Fleurya ...	39	Glochidion ...	34	Gynothrix ...	24	Hemiphysa ...	108	Hydroglossum ...	229	Jasminum ...	155
Flindersia ...	17	Glossocarya ...	173	Gynura ...	142	Hemisteirus ...	48	Hydroleia ...	161	Johnsonia ...	200
Floriscopa ...	206	Glossodia ...	192	Gysophila ...	46	Hemistemma ...	1	Hydropeltis ...	4	Josephia ...	123
Floscopa ...	206	Glossogyne ...	141	Gyrocarpus ...	87	Hemistephus ...	1	Hydrophyllaeae ...	161	Josephina ...	167
Flueggea ...	35	Glossostigma ...	164	Gyrostachis ...	188	Hemitelia ...	230	Hylococoua ...	34	Juncea ...	206
Fluvialea ...	203	Glyceria ...	226	Gyrostemon ...	54	Hemitelia ...	105	Hylogyne ...	122	Juncella ...	207
Fontainea ...	37	Glycine ...	70	Gyrostephium ...	140	Heritiera ...	27	Hymenanthra ...	11	Juncua ...	206
Forstera ...	146	Glycosmis ...	21			Hermannia ...	27	Hymenolepis ...	234	Jungia ...	91
Forsteropsis ...	144	Glycyrrhiza ...	69	Habenaria ...	192	Hernandia ...	7	Hymenophyllum ...	230	Jusieua ...	84
Fragosa ...	107	Gmelina ...	173	Haecckeria ...	138	Herniaria ...	46	Hymenoporum ...	12	Justicia ...	167
Franciisia ...	87	Gnaphalium ...	134	Haemodoraceae ...	194	Herpestis ...	164	Hypocaulymma ...	92		
Frankenia ...	45	Gnaphalodes ...	140	Haemodorum ...	194	Herpolirium ...	199	Hypoestes ...	168		
Frankeniaceae ...	45	Gnephosia ...	139	Hakea ...	120	Hetaeria ...	188, 205	Hypocaulymma ...	213		
Franklandia ...	114	Gomphandra ...	109	Halfordia ...	21	Heterachne ...	227	Hypolepis ...	234		
Freirea ...	39	Gompholobium ...	59	Halgania ...	176	Heterodendron ...	42	Hypolytrum ...	213		
Fremya ...	100	Gomphrena ...	47	Halocnemum ...	52	Heterolaena ...	55	Hypericinae ...	14		
Frenela ...	184	Gongronema ...	158	Halodule ...	204	Heteropogon ...	222	Hypericum ...	14		
Freycinetia ...	202	Goniocarpus ...	85	Halophila ...	193	Hewardia ...	197	Hypocaulymma ...	92		
Friesia ...	30	Goniophlebium ...	233	Halorageae ...	85	Hexatheca ...	204	Hypoestes ...	168		
Froebelina ...	177	Goniopogon ...	131	Haloragis ...	85	Hibbertia ...	1	Hypolaena ...	209		
Fugosia ...	25, 26	Goniopteris ...	233	Halothamnus ...	24	Hibiscus ...	25	Hypolepis ...	234		
Fuirena ...	213	Goniotriche ...	48	Hamamelideae ...	84	Hicksbeachia ...	116	Hypolytrum ...	213		
Funkia ...	198	Goodenia ...	149	Hamelina ...	198	Hierochloa ...	223	Hypoporum ...	217		
Fusanus ...	110	Goodeniaceae ...	146	Hannafordia ...	28	Hippocatea ...	41	Hypoxia ...	195		
		Goodenoughia ...	140	Haplocaryon ...	126	Hippocrepandra ...	32	Hyperpa ...	7		
Gahnia ...	216	Goodia ...	66	Haplopappus ...	133	Hodgkinsonia ...	126	Hypophila ...	40		
Gaiadendron ...	111	Goodyera ...	188	Haplotaxia ...	143	Hodgsonia ...	199	Hyptiandra ...	22		
Galactia ...	71	Gossypium ...	26	Haplobergia ...	70	Hodgsonioli ...	199	Hyrtanandra ...	39		
Galeola ...	188	Gouania ...	104	Harmogia ...	91	Holcus ...	222				
Galium ...	128	Gramineae ...	218	Harpullia ...	42	Hollandaea ...	122	Iberis ...	10		
Gamelythrum ...	223	Grammatotheca ...	143	Harrisonia ...	22	Hologamium ...	221	Ichnocarpus ...	157		
Gamozygis ...	139	Grammitis ...	234	Hartigsheia ...	16	Holostigma ...	143	Ilex ...	155		
Ganophyllum ...	44	Graptophyllum ...	168	Haussmannia ...	167	Holotome ...	108	Ilyogeton ...	164		
Gardenia ...	125	Gratiola ...	164	Hearnia ...	16	Holomanthus ...	37	Ilysanthes ...	164		
Gardneria ...	152	Granmuellera ...	204	Hedera ...	87	Homalium ...	11	Imbricaria ...	91		
Garuga ...	43	Greevesia ...	25	Hedera ...	105	Homalocalyx ...	90	Imperata ...	221		
Gastrodia ...	188	Grevillea ...	117	Hedraianthera ...	40	Homalospermum ...	93	Indigofera ...	67		
Gastrolobium ...	58	Grewia ...	31	Hedycarya ...	6	Homalostoma ...	182	Ionidium ...	10		
Gaultiera ...	177	Grisebachia ...	201	Hedyotis ...	125	Homoranthus ...	87	Iphigenia ...	198		
Geijera ...	21	Grumilia ...	127	Hedyscepe ...	201	Homoranthus ...	154	Ipidocaea ...	159		
Geiasois ...	82	Guettarda ...	126	Heleocharis ...	211	Hormogyne ...	66	Iridaeae ...	193		
Geitonoplesium ...	197	Guettardella ...	126	Helichrysum ...	136	Hovea ...	201	Iris ...	193		
Geleznovia ...	18	Guichenotia ...	29	Helicia ...	116	Howea ...	201	Iringia ...	105		
Genetyllis ...	87	Guilandina ...	72	Helicteres ...	27	Howittia ...	25	Isachne ...	218		
Geniostoma ...	153	Guioa ...	41	Heliotropium ...	176	Hoya ...	159	Isandra ...	163, 198		
Genoplesium ...	189	Gunnera ...	85	Heliotropium ...	176	Huanaca ...	107	Ischaemum ...	221		
Genosiris ...	193	Gunnia ...	52, 187	Heliotropium ...	136	Huegelia ...	88, 106	Iseilema ...	222		
Gentiana ...	152	Gunnopsia ...	52	Heliopsis ...	185	Huenefeldia ...	131	Isoetes ...	228		
Gentianeae ...	151	Guttiferae ...	14	Hellera ...	205	Hugonia ...	23	Isoetopsis ...	142		
Geodorum ...	187	Gymnagathis ...	94	Helmholtzia ...	196	Humata ...	231	Isolepis ...	213		
Geophila ...	127	Gymnanthera ...	157	Helmia ...	196	Humea ...	138	Isolobus ...	143		
Geraniaceae ...	123	Gymnaphanes ...	134	Helminthostachys ...	229	Husemannia ...	8	Isoloma ...	231		
Geranium ...	23	Gymnema ...	158	Helophyllum ...	146	Hutchinsia ...	10	Isopogon ...	112		
Gerbera ...	143	Gymnochaete ...	214	Helopha ...	218	Huttia ...	1	Isoschoenus ...	214		
Gesneriaceae ...	166	Gymnococca ...	55	Helosciadium ...	108	Hyalochlamys ...	139	Isotoma ...	144		
Geum ...	81	Gymnogramme ...	234	Hemolthrix ...	214	Hyalolepis ...	138	Isotropis ...	58		
Gilberta ...	139	Gymnogyne ...	141	Hemithria ...	220	Hyalosperma ...	136	Ixauchenus ...	130		
Gilesia ...	27	Gymnopteris ...	234	Hemiandra ...	171	Hybanthera ...	158	Ixioclamys ...	133		
Gillbeea ...	83	Gymnoschoenus ...	214	Hemiarrhena ...	164	Hybanthus ...	10	Ixiolaena ...	135		
				Hemicarpus ...	106	Hydnophytum ...	127	Ixioporum ...	12		

Laportea ...	39	Leucopogon ...	177	Lyginia ...	208	Melastoma ...	102	Mitrephora ...	5	Nematostigma ...	193
Lappago ...	220	Leucorchis ...	188	Lygodictyon ...	229	Melastomaceae ...	102	Mniarum ...	46	Nemeda ...	16
Lappula ...	175	Leucothamnus ...	28	Lygodium ...	229	Melhania ...	27	Modecca ...	128	Neoroepa ...	35
Lasianthus ...	127	Leuzea ...	143	Lyonsia ...	157	Melia ...	16	Molkenboeria ...	148	Neottopteris ...	232
Lasiopetalum ...	29	Levenhookia ...	146	Lyperanthus ...	191	Meliaceae ...	16	Mollinedia ...	5	Nepenthaceae ...	83
Lasraea ...	233	Lietzkya ...	90	Lysicarpus ...	100	Melichrus ...	177	Molloya ...	116	Nepenthe ...	83
Latourea ...	147	Libertia ...	193	Lysimachia ...	153	Melicope ...	20	Mollugo ...	53	Nephelium ...	42
Latrobea ...	64	Licuala ...	202	Lysinema ...	182	Melicytus ...	11	Momordica ...	129	Nephrodium ...	233
Lauraceae ...	6	Ligusticum ...	108	Lysiosepalum ...	29	Melochia ...	27	Monacather ...	225	Nephrolepis ...	233
Laurencia ...	144	Ligustrum ...	156	Lyssanthe ...	117	Melodinus ...	156	Monenyanthes ...	136	Neptunia ...	74
Lavatera ...	23	Liliaceae ...	197	Lythrum ...	84	Melodorum ...	5	Monenteles ...	134	Nertera ...	127
Lawrencella ...	136	Limnanthemum ...	151	Maba ...	155	Melothria ...	129	Monimiaceae ...	5	Nesaea ...	85
Lawrencia ...	24	Limnophila ...	163	Macadamia ...	116	Memecylon ...	102	Monochoria ...	205	Nesodaphne ...	6
Lawsonia ...	85	Limnrostachya ...	205	Macaranga ...	36	Memorialis ...	39	Monococcus ...	54	Nesogenes ...	172
Laxmannia ...	200	Limosella ...	164	Macarthuria ...	53	Meniocus ...	9	Monogramma ...	232	Nettoa ...	31
Learosa ...	5	Lindernia ...	164	Maccoya ...	175	Meniacium ...	233	Monoploca ...	10	Neurachne ...	220
Lebediera ...	35	Lindsaya ...	231	Macdonaldia ...	188	Menispermaceae ...	7	Monotaxis ...	32	Newcastlia ...	172
Ledgeria ...	188	Lineae ...	23	Macgregoria ...	44	Menkea ...	10	Monotoca ...	177	Nicotiana ...	162
Leea ...	105	Linkia ...	114	Mackinlaya ...	106	Mentha ...	168	Montia ...	45, 46	Niemeyera ...	154, 192
Leeaceae ...	105	Linociera ...	156	Macintyria ...	15	Merkusia ...	148	Moonia ...	140	Nipa ...	202
Leersia ...	222	Linospadix ...	201	Macropidia ...	195	Mertensia ...	230	Moraea ...	193	Nipaceae ...	202
Leewenhockia ...	146	Linschottenia ...	146	Macropiper ...	4	Meryta ...	106	Morelotia ...	216	Nipholobus ...	233
Legnephora ...	7	Linum ...	23	Macropodia ...	195	Mesembrianthemum ...	52	Morgania ...	163	Nitraria ...	22
Leguminosae ...	56	Liparis ...	185	Macropteranthus ...	87	Mesomelaena ...	214	Moriuda ...	126	Notelaea ...	156
Leichhardtia ...	7, 158, 184	Liparophyllum ...	151	Macrostegia ...	55	Mesotriche ...	177	Morisia ...	214	Nothochlaena ...	231
Lemna ...	203	Lipocarpha ...	213	Macrostigma ...	81	Methorium ...	27	Morna ...	135	Notholaena ...	231
Lemnaceae ...	203	Lippaya ...	125	Macrozamia ...	184	Metrosideros ...	100	Moachosma ...	168	Nothopanax ...	105
Lentibularinae ...	165	Lippia ...	171	Maesa ...	154	Mezoneuron ...	72	Motherwellia ...	105	Notomerium ...	157
Leontopodium ...	134	Lissanthe ...	177, 181	Magnoliaceae ...	4	Michiea ...	181	Mucuna ...	71	Notothixos ...	111
Lepia ...	10	Litobrochia ...	231	Mahernia ...	27	Micraira ...	224	Muehlenbeckia ...	53	Nuytsia ...	111
Lepidium ...	10	Litsea ...	6	Maireana ...	50	Micranthemum ...	32	Muehlenbergia ...	223	Nyctagineae ...	54
Lepidobolus ...	209	Livistona ...	202	Malacochaete ...	213	Microcachrys ...	184	Mukia ...	129	Nymphaea ...	4
Lepidocoma ...	138	Lobelia ...	143, 144	Malaisia ...	38	Microcarpaea ...	164	Murraya ...	21	Nymphaeaceae ...	4
Lepidosperma ...	215	Lobophyllum ...	176	Malcolmia ...	9	Microchloa ...	225	Musa ...	185	Nyssanthus ...	47
Lepidospora ...	215	Lobopogon ...	181	Mallophora ...	172	Microclisia ...	8	Myoporinae ...	174		
Lepidozamia ...	184	Logania ...	153	Mallotus ...	36	Microcorys ...	170	Myoporium ...	175	Oberonia ...	185
Lepigonum ...	46	Loganiaceae ...	152	Malpighiaceae ...	43	Microcybe ...	18	Myosotis ...	176	Obione ...	49
Lepilaena ...	204	Lomandra ...	200	Malva ...	24	Microgyne ...	133	Myosurus ...	3	Ochna ...	17
Lepironia ...	213	Lomaria ...	232	Malvaceae ...	23	Microlaena ...	222	Myrioccephalus ...	139	Ochnaceae ...	17
Lepistemon ...	159	Lomariopsis ...	234	Malvastrum ...	24	Microlepidium ...	10	Myriogyne ...	141	Ochrolasia ...	1
Leptaspis ...	220	Lomatia ...	122	Manglesia ...	117	Microlepis ...	231	Myriophyllum ...	86	Ochrosia ...	156
Leptatherum ...	221	Lonchocarpus ...	72	Manisauris ...	221	Micromelum ...	21	Myristica ...	6	Ocimum ...	168
Leptinella ...	141	Lopadocalyx ...	109	Mapania ...	213	Micromyrtus ...	90	Myristiceae ...	6	Octoclinis ...	184
Leptocarpus ...	209	Lophoclinium ...	134	Mappa ...	36	Micropyxis ...	153	Myrmecodia ...	127	Ocymum ...	168
Leptoceras ...	191	Lophostemon ...	100	Marattia ...	229	Microsciadium ...	107	Myrsinaceae ...	154	Oenanthe ...	108
Leptochloa ...	225	Lopidium ...	229	Marianthus ...	12	Microseris ...	143	Myrsine ...	154	Oenothera ...	84
Leptocymna ...	70	Loranthaceae ...	111	Mariacus ...	210	Microstachya ...	37	Myrtaceae ...	87	Ogerostylus ...	139
Leptocytisus ...	64	Loranthophyllum ...	152	Marlea ...	124	Microstemma ...	159	Myrtus ...	101	Olacineae ...	109
Leptolobium ...	70	Loranthus ...	111	Marquisia ...	127	Microstylis ...	188			Olae ...	109
Leptomeria ...	110	Lotus ...	66	Marsdenia ...	158	Microtis ...	190	Nablonium ...	140	Oldenlandia ...	125
Leptopteris ...	230	Loudonia ...	85	Marsilea ...	228	Milusia ...	5	Nageia ...	184	Olea ...	155
Leptorrhynchus ...	135	Lourea ...	70	Maundia ...	204	Milletia ...	68	Najas ...	203	Oleandra ...	233
Leptosema ...	56	Loxocarya ...	209	Mayepa ...	156	Milligania ...	85, 198	Nania ...	100	Olearia ...	131
Leptospermum ...	93	Lucuma ...	154	Mazentoxeron ...	20	Millotia ...	138	Nasturtium ...	9	Oliganthemum ...	133
Leptotriche ...	139	Ludwigia ...	84	Mazus ...	163	Milnea ...	16	Needhamia ...	181	Oligarrhena ...	181
Lepturus ...	221	Luffa ...	129	Medicosma ...	20	Mimosa ...	74	Negria ...	166	Omalanthus ...	37
Lepyrodia ...	208	Luisia ...	187	Medinilla ...	102	Mimulus ...	163	Nelitris ...	101	Omphacomeria ...	110
Leschenaultia ...	147	Lumnitzera ...	87	Megalotheca ...	208	Mimulus ...	154	Nelsonia ...	167	Onagreae ...	84
Lespedeza ...	70	Luzula ...	206	Meionectea ...	85	Minurys ...	131	Nelumbium ...	4	Onchospalum ...	209
Lesibudesia ...	49	Luzuriago ...	197	Melachne ...	216	Mirbelia ...	59	Nelumbo ...	4	Oncinocalyx ...	171
Leucas ...	169	Lycium ...	162	Melaleuca ...	94	Mischocarpus ...	41	Nematoceras ...	190	Oncosporum ...	12
Leucocarpum ...	40	Lycopodinae ...	228	Melanthesa ...	35	Mitragyne ...	152	Nematolepis ...	20	Oncostylia ...	211
Leucolaena ...	107	Lycopodium ...	228, 229	Melanthesiopsis ...	35	Mitrasacme ...	152	Nematophyllum ...	66	Opercularia ...	127
Leucophyta ...	140	Lycopus ...	168	Melanthium ...	198	Mitreola ...	152	Nematopus ...	139	Ophioderma ...	229

Ophioglossum ...	229	Paritium ...	25	Philoxerus ...	47	Pleiococca ...	21	Poranthera ...	32	Randia ...	125
Ophiorrhiza ...	125	Parkeria ...	230	Philydrella ...	205	Pleioogyne ...	141	Porospermum ...	105	Ranunculaceae ...	3
Ophiuros ...	221	Parsonsia ...	157	Philydram ...	205	Pleio gynium ...	44	Portenschlagia ...	40	Ranunculus ...	3
Opilia ...	109	Paryphantha ...	90	Phlebocallymma ...	109	Pleistichostemon ...	43	Portulaca ...	45	Raulia ...	134, 136
Oplismenus ...	220	Paspalum ...	218	Phlebocarya ...	194	Pleogyne ...	8	Portulacaceae ...	45	Rapuntium ...	143
Orchideae ...	185	Passiflora ...	128	Phlebodium ...	233	Pleopeltis ...	233	Posidonia ...	204	Ratonia ...	41
Oreobolus ...	214	Passifloraeae ...	128	Phoberos ...	11	Pleurandra ...	1	Potamogeton ...	204	Reedia ...	217
Oreocallis ...	122	Patania ...	230	Pholidia ...	174	Pleurandropsis ...	18	Potamophila ...	222	Regelia ...	96
Oreomyrrhis ...	108	Patersonia ...	193	Pholidopsis ...	174	Pleurocarpaea ...	129	Potentilla ...	81	Remirea ...	214
Oreostylichium ...	146	Pavetta ...	126	Pholidota ...	185	Pleurogramme ...	232	Pothos ...	203	Renalmia ...	193
Orites ...	116	Pavonia ...	25	Phormium ...	197	Pleuropappus ...	139	Pouzolzia ...	39	Restiaceae ...	207
Oritina ...	116	Pedalinæ ...	167	Phragmites ...	227	Plexaure ...	186	Pozoa ...	107	Restio ...	208
Ormocarpum ...	69	Pedilonum ...	186	Phreatia ...	186	Pinia ...	101	Pozoopsis ...	107	Rhabdotheramnus ...	166
Ornithochilus ...	187	Pelargonium ...	23	Phyllachne ...	146	Plinthacanthus ...	225	Prasophyllum ...	189	Rhagodia ...	50
Orobanche ...	165	Pellsea ...	231	Phyllanthus ...	34	Plokistigma ...	44	Pratia ...	143	Rhamnaceae ...	102
Orobanchaeae ...	165	Pelonastes ...	86	Phyllocallymma ...	139	Pluchea ...	133	Premna ...	173	Rhamnus ...	102
Orthoceras ...	189	Peltophorum ...	72	Phyllocladus ...	184	Plumbagineae ...	45	Primulaceae ...	153, 154	Rhamphicarpa ...	165
Orthopogon ...	220	Pemphis ...	85	Phyllodium ...	69	Plumbago ...	45	Prionosepalum ...	209	Rhamphidophora ...	188
Orthosiphon ...	168	Pennantia ...	109	Phylloglossum ...	229	Poa ...	226	Prionotes ...	182	Rhaphidophora ...	203
Orthostemon ...	152	Pennisetum ...	220	Phyllopappus ...	143	Podocarpus ...	184	Pritzelia ...	106, 205	Rhipogonum ...	197
Orthotropis ...	57	Pentaceras ...	22	Phyllota ...	62	Podocoma ...	133	Procris ...	38	Rhizophora ...	86
Orthrosanthus ...	193	Pentachondra ...	181	Phymatocarpus ...	96	Podolepis ...	135	Pronaya ...	12	Rhizophoreae ...	86
Oryza ...	222	Pentactropedion ...	223	Phymatodes ...	233	Podolobium ...	57	Prostanthera ...	169	Rhizospermae ...	228
Osbeckia ...	102	Pentadactylon ...	114	Physalis ...	161	Podopetalum ...	72	Proteaaceae ...	111	Rhodamnia ...	101
Osbornia ...	100	Pentadynamis ...	66	Physolobium ...	70	Podosperma ...	134	Prunella ...	169	Rhodanthe ...	136
Oschstzja ...	107	Pentagonaster ...	93	Physopsis ...	172	Podostemoneae ...	166	Pseudalangium ...	124	Rhododendron ...	177
Osmunda ...	230	Pentalepis ...	140	Phytolaccaeae ...	54	Podothea ...	134	Pseudanthus ...	32	Rhodomlyrtus ...	101
Osteocarpum ...	51	Pentapanax ...	105	Picrophyta ...	149	Poecilodermis ...	26	Pseudatalaya ...	41	Rhodosphaera ...	44
Otaothera ...	102	Pentapeltis ...	107	Pigea ...	10	Pogonatherum ...	221	Pseudomorus ...	38	Rhopala ...	116
Otella ...	193	Pentapogon ...	223	Pileanthus ...	89	Pogonetes ...	148	Pseudophilodia ...	174	Rhus ...	44
Ourisia ...	164	Pentaptelion ...	177	Pilitis ...	183	Pogonia ...	175, 188	Psilotrichum ...	48	Rhyncharrhena ...	158
Owenia ...	16	Pentataphrus ...	177	Pilularia ...	228	Pogonolepis ...	139	Psilotum ...	228	Rhynchosia ...	71
Oxalis ...	23	Pentatropis ...	158	Pimelea ...	55	Pogonolobus ...	126	Psoralea ...	66	Rhynchospora ...	214
Oxleya ...	17	Peperomia ...	4	Piper ...	4	Pogostemon ...	168	Psychotria ...	127	Rhynchosostemon ...	28
Oxycyladium ...	59	Peplidium ...	164	Piperaceae ...	4	Poiretia ...	66, 182	Psydrax ...	126	Rhytidandra ...	124
Oxylobium ...	57	Pericalymma ...	93	Piptandra ...	91	Polanisia ...	8	Pteridium ...	231	Rhytidanthe ...	135
Oxymyrrhine ...	91	Pericampylus ...	7	Piptocalyx ...	6	Polia ...	206	Pterigeron ...	133	Rhytidospurum ...	12
Oxystelma ...	157	Pernettya ...	177	Piptomeria ...	60	Pollichia ...	176	Pteris ...	231	Richea ...	140, 183
Ozothamnus ...	136	Peroa ...	177	Piptostemma ...	139	Pollioia ...	221	Pterocaulon ...	134	Ricinocarpus ...	33
Pachycornia ...	52	Perojoa ...	177	Pipturus ...	39	Polyalthia ...	5	Pterochaete ...	135	Riedleya ...	27
Pachygone ...	8	Perotis ...	220	Pisonia ...	54	Polycalymma ...	139	Pterolobium ...	72	Rinzia ...	91
Pachynema ...	3	Perrottetia ...	40	Pithecolobium ...	80	Polycarpaea ...	47	Pteroloma ...	69	Ripogonum ...	197
Pachysura ...	140	Persoonia ...	114	Pithocarpa ...	138	Polycarpon ...	47	Pteropogon ...	136	Rochelia ...	175
Pacquerina ...	130	Petalogyne ...	73	Pithyrodia ...	172	Polygonemon ...	49	Pterostigma ...	163	Roea ...	60
Pagetia ...	21	Petalolepis ...	136	Pittosporaeae ...	11	Polygala ...	14	Pterostylis ...	190	Roettlera ...	166
Palmae ...	201	Petalostigma ...	34	Pittosporum ...	11	Polygaleae ...	14	Pterygopappus ...	134	Roepera ...	22
Palmeria ...	6	Petalostylis ...	73	Plagianthus ...	24	Polygonaceae ...	53	Ptilotus ...	48	Roeperia ...	8, 33
Panaetia ...	135	Petermannia ...	196	Plagiogyria ...	232	Polygonum ...	53	Ptychosema ...	67	Rosaceae ...	81
Panax ...	105	Petrolobium ...	72	Plagiolobium ...	66	Polymeria ...	160	Ptychosperma ...	202	Rostellaria ...	167
Pancovia ...	41	Petrophila ...	111	Plagiosetum ...	220	Polyosma ...	82	Pultenaea ...	62	Rostellularia ...	67
Pandanæae ...	202	Phacelothrix ...	139	Plantagineae ...	153	Polyphragmon ...	126	Pumilo ...	138	Rotala ...	84
Pandanophyllum ...	213	Phaenopoda ...	134	Plantago ...	153	Polypodium ...	233	Punicella ...	92	Rottboellia ...	221
Pandanus ...	202	Phajus ...	188	Platycarpidium ...	107	Polypogon ...	220	Pycnarrhena ...	8	Rottlera ...	136
Panicum ...	218	Phalangium ...	199	Platycerium ...	234	Polypompholyx ...	166	Pycnolachne ...	171	Roupaia ...	116
Panopsis ...	116	Phaleria ...	56	Platycheilum ...	66	Polystichum ...	233	Pycnosorus ...	140	Rourea ...	81
Papaver ...	8	Phanerandra ...	177	Platylobium ...	64	Polyzone ...	87	Pycnospora ...	70	Roxburghia ...	196
Papaveraceae ...	8	Pharbitis ...	159	Platyloma ...	231	Pomaderris ...	103	Pycnus ...	210	Roxburghiaceae ...	196
Pappophorum ...	224	Phaseolus ...	71	Platytelea ...	82	Pomatoderris ...	103	Pygmaea ...	164	Rubiaceae ...	125
Paractaenium ...	218	Phebalium ...	18	Platysace ...	107	Pomax ...	127	Pyxidaria ...	164	Rubus ...	81
Paratrophia ...	105	Phegopteris ...	238	Platytheca ...	15	Ponceletia ...	182	Quinetia ...	139	Ruelingia ...	27
Parduyna ...	198	Pherosphaera ...	184	Platythoma ...	230	Pongamia ...	72	Quintinia ...	82	Ruellia ...	167
Parietaria ...	39	Philbydreae ...	205	Plectranthus ...	168	Pontederiaceae ...	205	Quoya ...	172	Rumex ...	53
Parinarium ...	81	Philhydrium ...	205	Pleconia ...	126	Popowia ...	5			Rupia ...	204
		Philotheca ...	20	Pleea ...	198	Porana ...	160			Rutaceae ...	17

Rutidochlamys ...	135	Scirpodendron ...	213	Soleniscia ...	177	Stephania ...	7	Taxanthema ...	45	Trachycaryon ...	36
Rutidosia ...	138	Scirpus ...	213	Solenogyne ...	130	Sterculia ...	26	Tecoma ...	166	Trachymene ...	107
Ryssopterys ...	43	Scitamineae ...	185	Solenostigma ...	37	Sterculiaceae ...	26	Tecticornia ...	52	Trachypogon ...	222
Saccharum ...	221	Scleranthus ...	46	Soliva ...	141	Stipa ...	223	Telanthera ...	47	Tragia ...	36
Saccolabium ...	187	Scleria ...	217	Sollia ...	12	Stipelia ...	177	Telmatophace ...	203	Tragus ...	220
Saccopetalum ...	5	Sclerochlamys ...	50	Sondera ...	13	Stirlingia ...	113	Telesia ...	122	Trema ...	37
Sagena ...	233	Sclerolaena ...	51	Sonneratia ...	102	Stomarrhena ...	177	Temminckia ...	148	Tremandra ...	16
Sagina ...	46	Scleroleima ...	142	Sonzaya ...	44	Strangea ...	116	Templetonia ...	65	Tremandreae ...	15
Salacia ...	41	Sclerothamnus ...	64	Sophora ...	72	Stravadium ...	102	Tenagocharis ...	205	Trianthema ...	52
Salicariaceae ...	84	Scolopendrium ...	232	Sopubia ...	165	Streblorrhiza ...	68	Tephrosia ...	67	Tribonanthus ...	194
Salicornia ...	52	Scopia ...	11	Sorghum ...	222	Streptachne ...	223	Terminalia ...	87	Tribulopsis ...	22
Salisia ...	93	Scoparia ...	164	Sorostachys ...	210	Streptocarpus ...	166	Ternstroemiaceae ...	14	Tribulus ...	22
Salomonia ...	14	Scorzenera ...	143	Sowerbaea ...	199	Streptoglossa ...	133	Tersonia ...	54	Trichanthodium ...	139
Salsola ...	52	Scottia ...	65	Spadostyles ...	62	Streptothamnus ...	11	Tetracarpaea ...	83	Trichelostylis ...	211
Salsolaceae ...	49	Scrophularinae ...	163	Spanoghea ...	42	Striga ...	165	Tetracera ...	3	Trichinium ...	48
Salvia ...	169	Scutellaria ...	169	Sparganium ...	203	Strobilanthes ...	167	Tetracheilus ...	74	Trichobasis ...	96
Salvinia ...	228	Scyphiphora ...	126	Spartothamnus ...	171	Stromatopteris ...	230	Tetragonella ...	52	Trichobolus ...	81
Samara ...	154	Scyphocoronis ...	138	Spathodea ...	167	Strongylodon ...	71	Tetragonia ...	52	Trichocline ...	143
Sambucus ...	128	Seaforthia ...	202	Spathoglottis ...	188	Strongylopermum ...	141	Tetralepis ...	214	Trichodesma ...	176
Samolus ...	153	Sebaea ...	151	Spergularia ...	46	Strychnos ...	152	Tetralobus ...	166	Trichodium ...	224
Samydaceae ...	11	Sebastiania ...	37	Spermaceae ...	128	Strzeleckia ...	17, 144	Tetranthera ...	6	Tricholoma ...	164
Sandfordia ...	18	Secamone ...	157	Spermalepis ...	100	Sturtia ...	26	Tetrapasma ...	104	Trichomanes ...	229
Santalaceae ...	110	Securinea ...	35	Spermaxyrum ...	109	Sturmia ...	185	Tetraphylax ...	149	Trichosanthes ...	128
Santalum ...	110	Selaginella ...	229	Sphaeranthos ...	134	Sturtia ...	26	Tetrapora ...	91	Trichosiphon ...	26
Sapindaceae ...	41	Selliera ...	149	Sphaerolobium ...	60	Stylideae ...	144	Tetrarrhena ...	222	Trichostegia ...	135
Sapindus ...	41	Selligera ...	234	Sphaeromorphaca ...	134, 141	Stylidium ...	124, 144	Tetralelia ...	8	Tricondylus ...	122
Saponaria ...	46	Selwynia ...	7	Sphaeropus ...	217	Stylobasium ...	81	Tetratheca ...	15	Tricoryne ...	199
Sapota ...	154	Semecarpus ...	44	Sphenoclea ...	144	Stylodoryne ...	126	Teucrium ...	171	Tricostularia ...	214
Sapotaceae ...	154	Senebiera ...	10	Sphenotoma ...	183	Stylolopsis ...	135	Thalamia ...	184	Trigastrotecha ...	53
Sarauia ...	14	Senecio ...	142	Sphincterostoma ...	182	Styloncerus ...	139	Thamnopteris ...	232	Triglochis ...	204
Sarcocephalus ...	125	Sentis ...	174	Spiculaea ...	191	Stylurus ...	117	Thecanthus ...	55	Trigonella ...	66
Sarcochilus ...	187	Sequoia ...	183	Spilanthus ...	141	Stypandra ...	199	Theleophyton ...	49	Trineuron ...	142
Sarcogonum ...	53	Serianthes ...	80	Spinifex ...	220	Styphelia ...	177	Thelychiton ...	186	Triodia ...	226
Sarcopetalum ...	7	Seringea ...	28	Spiraeanthemum ...	83	Styraceae ...	155	Thelymitra ...	188	Tripetelus ...	128
Sarcostemma ...	158	Sersalisia ...	154	Spiranthera ...	12	Suaeda ...	52	Themeda ...	222	Triphelia ...	87
Sarcozygium ...	22	Sesbania ...	68	Spiranthes ...	188	Suriana ...	22	Therogeron ...	131	Tripladenia ...	198
Sarotes ...	29	Seseli ...	108	Spirodela ...	203	Swainsona ...	68	Thesium ...	111	Tripterococcus ...	44
Saussurea ...	143	Sesuvium ...	52	Spiropodium ...	133	Swammerdamia ...	136	Thespesia ...	26	Triptilodiscus ...	136
Saxifragaceae ...	82	Setaria ...	220	Spodiopogon ...	221, 222	Sycios ...	129	Thespidium ...	134	Trirhaphis ...	227
Scaevola ...	148	Sheffieldia ...	153	Spondias ...	44	Symphionema ...	141	Thlaspi ...	10	Trisetum ...	224
Scalia ...	135	Sicyos ...	129	Sponia ...	37	Symphionema ...	114	Thomasia ...	28	Tristania ...	100
Scallioptis ...	135	Sida ...	24	Sporobolus ...	224	Symphomyrtus ...	97	Thouarsea ...	220	Tristaniopsis ...	100
Schelhammra ...	198	Sideroxylon ...	154	Sprengelia ...	182	Symphyonema ...	114	Thozetia ...	159	Tristellateia ...	43
Schedonorus ...	226	Sieberta ...	107	Spyridium ...	103	Symphopetalum ...	20	Threlkeldia ...	52	Tristichocalyx ...	7
Schenodorus ...	226	Siegesbeckia ...	141	Stachystemon ...	32	Symplocos ...	155	Thrixpermum ...	187	Trithuria ...	207
Schidiomyrtus ...	91	Sieglingia ...	226	Stackhousia ...	44	Synaphea ...	113	Thryptomene ...	90	Triticum ...	227
Schizachyrium ...	222	Siemsenia ...	135	Stackhousiacea ...	44	Synaptantha ...	125	Thuarea ...	220	Triumfetta ...	30
Schizaea ...	229	Sieversia ...	81	Stadmanidia ...	41	Synonym ...	100	Thunbergia ...	167	Trochocarpa ...	181
Schizeilema ...	107	Siloxerus ...	139	Statice ...	45	Synophlebium ...	231	Thymeleae ...	55	Trollius ...	3
Schizoloma ...	231	Silphiospermum ...	130	Stawellia ...	200	Synostemon ...	34	Thysacanthus ...	168	Trymalium ...	103
Schizomeria ...	82	Simarubeae ...	22	Stetzia ...	131	Synoum ...	16	Thysanella ...	198	Turraea ...	16
Schizopleura ...	96	Simsia ...	113	Stegania ...	232	Syzygium ...	101	Thysanotus ...	198	Turritia ...	9
Schleidenia ...	176	Siona ...	199	Steiroglossa ...	130	Tabernaemontana ...	157	Tiaridium ...	176	Tylophora ...	158
Schmidelia ...	41	Siphonodon ...	41	Stekhovia ...	149	Tacca ...	194	Tiliaceae ...	30	Typha ...	203
Schoberia ...	52	Sisymbrium ...	9	Stellaria ...	46	Taccaceae ...	194	Tillaea ...	83	Typhaceae ...	203
Schoenia ...	136	Sisyrrinchium ...	193	Stemodia ...	163	Taeniophyllum ...	187	Timonius ...	126	Typhonium ...	203
Schoenodum ...	208, 209	Sium ...	108	Stenanthemum ...	103	Talinum ...	45	Tinospora ...	7	Umbelliferae ...	106
Schoenolaena ...	107	Skinnera ...	159	Stenanthus ...	177	Tamarindus ...	73	Titania ...	185	Uncinia ...	217
Schoenus ...	214	Skirophorus ...	139	Stenocarpus ...	122	Tapeinocheilos ...	185	Tittmannsia ...	164	Ungeria ...	27
Scholtzia ...	91	Sloanea ...	30	Stenochilus ...	174	Tarensa ...	126	Tmesipteris ...	228	Uniola ...	226
Schuermannia ...	87	Smilax ...	197	Stenochlaena ...	234	Tarrietia ...	27	Todea ...	230	Urachne ...	223
Schwaegrichenia ...	195	Smithia ...	69	Stenodiscus ...	103	Tasmania ...	4	Tournefortia ...	176	Uragoga ...	127
Scirpidium ...	211	Solanaceae ...	161	Stenopetalum ...	9	Tatea ...	173	Toxanthus ...	138	Uralespis ...	226
		Solanum ...	161	Stenotaphrum ...	220						

Uria	...	70	Ventenatia	...	177	Vitex	...	173	Wichurea	...	103	Xanthorrhoea	...	201	Zaleya	...	52
Urena...	...	25	Ventilago	...	102	Vitis	...	104	Wickstroemia	...	56	Xanthosia	...	107	Zamia...	...	184
Urocarpus	...	18	Verbena	...	171	Vittadinia	...	133	Wilckia	...	9	Xanthostemon	...	100	Zannichellia	...	204
Urodon	...	62	Verbenaceae	...	171	Vittaria	...	231	Wilkiea	...	5	Xanthoxylum	...	21	Zanonia	...	129
Urostigma	...	37	Vernonia	...	130	Vlamingia	...	10	Wilsonia	...	161	Xerochloa	...	220	Zanthoxylum	...	21
Urtica	...	39	Veronica	...	164	Vulpia	...	226, 227	Wistaria	...	68	Xerosollya	...	12	Zapania	...	171
Urticaceae	...	37	Verreauxia	...	148				Wittsteinia	...	177	Xerotes	...	200	Zehneria	...	129
Utricularia	...	165	Verticordia	...	88	Wahlenbergia	...	144	Woffia	...	203	Ximcna	...	109	Zichya	...	70
Uvaria	...	4	Vigna	...	71	Waitzia	...	135	Wollastonia	...	141	Xiphopteris	...	233	Zieria	...	17
Uvedalia	...	163	Vilfa	...	224	Walcottia	...	171	Woodwardia	...	232	Xylocarpus	...	16	Zizyphus	...	102
			Villaresia	...	109	Waltheria	...	27	Woollsia	...	182	Xylomelum	...	116	Zornia...	...	69
Vaccinium	...	177	Villarsia	...	151	Warburtonia	...	1	Wormia	...	1	Xylosma	...	11	Zoatera	...	204
Vachellia	...	74	Viminaria	...	61	Wartmannia	...	37	Wrightia	...	157	Xyridanthe	...	136	Zosterostylis	...	189
Vaginularia	...	232	Vincetoxicum	...	157	Webera	...	126	Wrixonia	...	170	Xyrideae	...	206	Zoysia	...	221
Vallisneria	...	194	Viniferae	...	104	Wedelia	...	141	Wurmbsea	...	198	Xyris	...	256	Zygia	...	80
Vandellia	...	164	Viola	...	10	Wehlia	...	91				Xystidium	...	220	Zygomenes	...	205
Veitchia	...	201	Violaceae	...	10	Weinmannia	...	82	Xanthochrysum	...	136				Zygophylleae	...	22
Velleya	...	150	Viraya	...	135	Westonia	...	66	Xanthophyllum	...	15	Youngia	...	143	Zygophyllum	...	22
Ventenata	...	144	Viscum	...	111	Westringia	...	171									

OMISSIONS.

Senebiera (Coronopus)...	...	...	...	...	...	10
Sisymbrium (Geococcus)	...	...	...	...	...	9
Ventilago (Berchemia)...	...	...	...	...	...	102

THIS BOOK IS DUE ON THE LAST DATE
STAMPED BELOW

AN INITIAL FINE OF 25 CENTS

WILL BE ASSESSED FOR FAILURE TO RETURN THIS BOOK
ON THE DATE DUE. THE PENALTY WILL INCREASE TO
50 CENTS ON THE FOURTH DAY AND TO \$1.00 ON THE
SEVENTH DAY OVERDUE.

JUN 6 1960

JAN 24 1963

FEB 13 1963

LIBRARY USE

NOV 21 1963

DEC 12 1968

Call Number:

174669

Mueller, F.

Second systematic
census of Australian

QK431

M93

Mueller

QK431

M93

174669